

Computer Algebra Independent Integration Tests

Summer 2024

7-Inverse-hyperbolic-functions/7.3-Inverse-hyperbolic-
tangent/337-7.3.2

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 8:02am

Contents

1	Introduction	10
1.1	Listing of CAS systems tested	11
1.2	Results	12
1.3	Time and leaf size Performance	16
1.4	Performance based on number of rules Rubi used	18
1.5	Performance based on number of steps Rubi used	19
1.6	Solved integrals histogram based on leaf size of result	20
1.7	Solved integrals histogram based on CPU time used	21
1.8	Leaf size vs. CPU time used	22
1.9	list of integrals with no known antiderivative	23
1.10	List of integrals solved by CAS but has no known antiderivative	23
1.11	list of integrals solved by CAS but failed verification	23
1.12	Timing	24
1.13	Verification	24
1.14	Important notes about some of the results	25
1.15	Current tree layout of integration tests	28
1.16	Design of the test system	29
2	detailed summary tables of results	30
2.1	List of integrals sorted by grade for each CAS	31
2.2	Detailed conclusion table per each integral for all CAS systems	37
2.3	Detailed conclusion table specific for Rubi results	98
3	Listing of integrals	106
3.1	$\int x^5(a + \operatorname{barctanh}(cx)) dx$	114
3.2	$\int x^4(a + \operatorname{barctanh}(cx)) dx$	120
3.3	$\int x^3(a + \operatorname{barctanh}(cx)) dx$	126
3.4	$\int x^2(a + \operatorname{barctanh}(cx)) dx$	132
3.5	$\int x(a + \operatorname{barctanh}(cx)) dx$	138
3.6	$\int (a + \operatorname{barctanh}(cx)) dx$	144
3.7	$\int \frac{a + \operatorname{barctanh}(cx)}{x} dx$	149

3.8	$\int \frac{a+\operatorname{barctanh}(cx)}{x^2} dx$	154
3.9	$\int \frac{a+\operatorname{barctanh}(cx)}{x^3} dx$	160
3.10	$\int \frac{a+\operatorname{barctanh}(cx)}{x^4} dx$	166
3.11	$\int \frac{a+\operatorname{barctanh}(cx)}{x^5} dx$	172
3.12	$\int \frac{a+\operatorname{barctanh}(cx)}{x^6} dx$	178
3.13	$\int x^5(a + \operatorname{barctanh}(cx))^2 dx$	184
3.14	$\int x^4(a + \operatorname{barctanh}(cx))^2 dx$	194
3.15	$\int x^3(a + \operatorname{barctanh}(cx))^2 dx$	203
3.16	$\int x^2(a + \operatorname{barctanh}(cx))^2 dx$	211
3.17	$\int x(a + \operatorname{barctanh}(cx))^2 dx$	219
3.18	$\int (a + \operatorname{barctanh}(cx))^2 dx$	226
3.19	$\int \frac{(a+\operatorname{barctanh}(cx))^2}{x} dx$	232
3.20	$\int \frac{(a+\operatorname{barctanh}(cx))^2}{x^2} dx$	239
3.21	$\int \frac{(a+\operatorname{barctanh}(cx))^2}{x^3} dx$	245
3.22	$\int \frac{(a+\operatorname{barctanh}(cx))^2}{x^4} dx$	253
3.23	$\int \frac{(a+\operatorname{barctanh}(cx))^2}{x^5} dx$	260
3.24	$\int x^5(a + \operatorname{barctanh}(cx))^3 dx$	270
3.25	$\int x^4(a + \operatorname{barctanh}(cx))^3 dx$	283
3.26	$\int x^3(a + \operatorname{barctanh}(cx))^3 dx$	294
3.27	$\int x^2(a + \operatorname{barctanh}(cx))^3 dx$	304
3.28	$\int x(a + \operatorname{barctanh}(cx))^3 dx$	313
3.29	$\int (a + \operatorname{barctanh}(cx))^3 dx$	321
3.30	$\int \frac{(a+\operatorname{barctanh}(cx))^3}{x} dx$	328
3.31	$\int \frac{(a+\operatorname{barctanh}(cx))^3}{x^2} dx$	336
3.32	$\int \frac{(a+\operatorname{barctanh}(cx))^3}{x^3} dx$	343
3.33	$\int \frac{(a+\operatorname{barctanh}(cx))^3}{x^4} dx$	351
3.34	$\int \frac{(a+\operatorname{barctanh}(cx))^3}{x^5} dx$	361
3.35	$\int (dx)^{5/2}(a + \operatorname{barctanh}(cx)) dx$	370
3.36	$\int (dx)^{3/2}(a + \operatorname{barctanh}(cx)) dx$	378
3.37	$\int \sqrt{dx}(a + \operatorname{barctanh}(cx)) dx$	385
3.38	$\int \frac{a+\operatorname{barctanh}(cx)}{\sqrt{dx}} dx$	392
3.39	$\int \frac{a+\operatorname{barctanh}(cx)}{(dx)^{3/2}} dx$	399
3.40	$\int \frac{a+\operatorname{barctanh}(cx)}{(dx)^{5/2}} dx$	406
3.41	$\int \frac{a+\operatorname{barctanh}(cx)}{(dx)^{7/2}} dx$	413

3.42	$\int \frac{a+\operatorname{barctanh}(cx)}{(dx)^{9/2}} dx$	420
3.43	$\int (dx)^m (a + \operatorname{barctanh}(cx))^3 dx$	428
3.44	$\int (dx)^m (a + \operatorname{barctanh}(cx))^2 dx$	433
3.45	$\int (dx)^m (a + \operatorname{barctanh}(cx)) dx$	438
3.46	$\int \frac{(dx)^m}{a+\operatorname{barctanh}(cx)} dx$	443
3.47	$\int \frac{(dx)^m}{(a+\operatorname{barctanh}(cx))^2} dx$	448
3.48	$\int (a + \operatorname{barctanh}(cx))^p dx$	453
3.49	$\int (dx)^m (a + \operatorname{barctanh}(cx))^p dx$	458
3.50	$\int x^7 (a + \operatorname{barctanh}(cx^2)) dx$	463
3.51	$\int x^5 (a + \operatorname{barctanh}(cx^2)) dx$	469
3.52	$\int x^3 (a + \operatorname{barctanh}(cx^2)) dx$	475
3.53	$\int x (a + \operatorname{barctanh}(cx^2)) dx$	481
3.54	$\int \frac{a+\operatorname{barctanh}(cx^2)}{x} dx$	487
3.55	$\int \frac{a+\operatorname{barctanh}(cx^2)}{x^3} dx$	492
3.56	$\int \frac{a+\operatorname{barctanh}(cx^2)}{x^5} dx$	498
3.57	$\int \frac{a+\operatorname{barctanh}(cx^2)}{x^7} dx$	504
3.58	$\int x^4 (a + \operatorname{barctanh}(cx^2)) dx$	510
3.59	$\int x^2 (a + \operatorname{barctanh}(cx^2)) dx$	517
3.60	$\int (a + \operatorname{barctanh}(cx^2)) dx$	524
3.61	$\int \frac{a+\operatorname{barctanh}(cx^2)}{x^2} dx$	530
3.62	$\int \frac{a+\operatorname{barctanh}(cx^2)}{x^4} dx$	537
3.63	$\int \frac{a+\operatorname{barctanh}(cx^2)}{x^6} dx$	544
3.64	$\int x^7 (a + \operatorname{barctanh}(cx^2))^2 dx$	551
3.65	$\int x^5 (a + \operatorname{barctanh}(cx^2))^2 dx$	560
3.66	$\int x^3 (a + \operatorname{barctanh}(cx^2))^2 dx$	568
3.67	$\int x (a + \operatorname{barctanh}(cx^2))^2 dx$	576
3.68	$\int \frac{(a+\operatorname{barctanh}(cx^2))^2}{x} dx$	583
3.69	$\int \frac{(a+\operatorname{barctanh}(cx^2))^2}{x^3} dx$	590
3.70	$\int \frac{(a+\operatorname{barctanh}(cx^2))^2}{x^5} dx$	597
3.71	$\int x^4 (a + \operatorname{barctanh}(cx^2))^2 dx$	606
3.72	$\int x^2 (a + \operatorname{barctanh}(cx^2))^2 dx$	614
3.73	$\int (a + \operatorname{barctanh}(cx^2))^2 dx$	622
3.74	$\int \frac{(a+\operatorname{barctanh}(cx^2))^2}{x^2} dx$	629
3.75	$\int \frac{(a+\operatorname{barctanh}(cx^2))^2}{x^4} dx$	637

3.76	$\int \frac{(a+b\operatorname{arctanh}(cx^2))^2}{x^6} dx$	645
3.77	$\int x^3(a+b\operatorname{arctanh}(cx^2))^3 dx$	653
3.78	$\int x(a+b\operatorname{arctanh}(cx^2))^3 dx$	662
3.79	$\int \frac{(a+b\operatorname{arctanh}(cx^2))^3}{x} dx$	669
3.80	$\int \frac{(a+b\operatorname{arctanh}(cx^2))^3}{x^3} dx$	676
3.81	$\int \frac{(a+b\operatorname{arctanh}(cx^2))^3}{x^5} dx$	683
3.82	$\int (dx)^{5/2}(a+b\operatorname{arctanh}(cx^2)) dx$	690
3.83	$\int (dx)^{3/2}(a+b\operatorname{arctanh}(cx^2)) dx$	705
3.84	$\int \sqrt{dx}(a+b\operatorname{arctanh}(cx^2)) dx$	719
3.85	$\int \frac{a+b\operatorname{arctanh}(cx^2)}{\sqrt{dx}} dx$	733
3.86	$\int \frac{a+b\operatorname{arctanh}(cx^2)}{(dx)^{3/2}} dx$	745
3.87	$\int \frac{a+b\operatorname{arctanh}(cx^2)}{(dx)^{5/2}} dx$	758
3.88	$\int \frac{a+b\operatorname{arctanh}(cx^2)}{(dx)^{7/2}} dx$	771
3.89	$\int \frac{a+b\operatorname{arctanh}(cx^2)}{(dx)^{9/2}} dx$	787
3.90	$\int \sqrt{dx}(a+b\operatorname{arctanh}(cx^2))^2 dx$	803
3.91	$\int \frac{(a+b\operatorname{arctanh}(cx^2))^2}{\sqrt{dx}} dx$	810
3.92	$\int \frac{(a+b\operatorname{arctanh}(cx^2))^2}{(dx)^{3/2}} dx$	817
3.93	$\int \frac{(a+b\operatorname{arctanh}(cx^2))^2}{(dx)^{5/2}} dx$	824
3.94	$\int (dx)^m(a+b\operatorname{arctanh}(cx^2))^3 dx$	831
3.95	$\int (dx)^m(a+b\operatorname{arctanh}(cx^2))^2 dx$	836
3.96	$\int (dx)^m(a+b\operatorname{arctanh}(cx^2)) dx$	841
3.97	$\int \frac{(dx)^m}{a+b\operatorname{arctanh}(cx^2)} dx$	846
3.98	$\int \frac{(dx)^m}{(a+b\operatorname{arctanh}(cx^2))^2} dx$	851
3.99	$\int x^{11}(a+b\operatorname{arctanh}(cx^3)) dx$	856
3.100	$\int x^8(a+b\operatorname{arctanh}(cx^3)) dx$	862
3.101	$\int x^5(a+b\operatorname{arctanh}(cx^3)) dx$	868
3.102	$\int x^2(a+b\operatorname{arctanh}(cx^3)) dx$	874
3.103	$\int \frac{a+b\operatorname{arctanh}(cx^3)}{x} dx$	879
3.104	$\int \frac{a+b\operatorname{arctanh}(cx^3)}{x^4} dx$	884
3.105	$\int \frac{a+b\operatorname{arctanh}(cx^3)}{x^7} dx$	890
3.106	$\int \frac{a+b\operatorname{arctanh}(cx^3)}{x^{10}} dx$	896

3.107	$\int x^3(a + \operatorname{barctanh}(cx^3)) dx$	902
3.108	$\int (a + \operatorname{barctanh}(cx^3)) dx$	913
3.109	$\int \frac{a+\operatorname{barctanh}(cx^3)}{x^3} dx$	919
3.110	$\int \frac{a+\operatorname{barctanh}(cx^3)}{x^6} dx$	929
3.111	$\int x^7(a + \operatorname{barctanh}(cx^3)) dx$	938
3.112	$\int x^4(a + \operatorname{barctanh}(cx^3)) dx$	950
3.113	$\int x(a + \operatorname{barctanh}(cx^3)) dx$	959
3.114	$\int \frac{a+\operatorname{barctanh}(cx^3)}{x^2} dx$	970
3.115	$\int \frac{a+\operatorname{barctanh}(cx^3)}{x^5} dx$	979
3.116	$\int x^{11}(a + \operatorname{barctanh}(cx^3))^2 dx$	990
3.117	$\int x^8(a + \operatorname{barctanh}(cx^3))^2 dx$	999
3.118	$\int x^5(a + \operatorname{barctanh}(cx^3))^2 dx$	1008
3.119	$\int x^2(a + \operatorname{barctanh}(cx^3))^2 dx$	1016
3.120	$\int \frac{(a+\operatorname{barctanh}(cx^3))^2}{x} dx$	1023
3.121	$\int \frac{(a+\operatorname{barctanh}(cx^3))^2}{x^4} dx$	1030
3.122	$\int \frac{(a+\operatorname{barctanh}(cx^3))^2}{x^7} dx$	1036
3.123	$\int \frac{(a+\operatorname{barctanh}(cx^3))^2}{x^{10}} dx$	1044
3.124	$\int x^8(a + \operatorname{barctanh}(cx^3))^3 dx$	1052
3.125	$\int x^5(a + \operatorname{barctanh}(cx^3))^3 dx$	1060
3.126	$\int x^2(a + \operatorname{barctanh}(cx^3))^3 dx$	1069
3.127	$\int \frac{(a+\operatorname{barctanh}(cx^3))^3}{x} dx$	1076
3.128	$\int \frac{(a+\operatorname{barctanh}(cx^3))^3}{x^4} dx$	1083
3.129	$\int \frac{(a+\operatorname{barctanh}(cx^3))^3}{x^7} dx$	1090
3.130	$\int (dx)^m (a + \operatorname{barctanh}(cx^3))^3 dx$	1097
3.131	$\int (dx)^m (a + \operatorname{barctanh}(cx^3))^2 dx$	1102
3.132	$\int (dx)^m (a + \operatorname{barctanh}(cx^3)) dx$	1107
3.133	$\int \frac{(dx)^m}{a+\operatorname{barctanh}(cx^3)} dx$	1112
3.134	$\int \frac{(dx)^m}{(a+\operatorname{barctanh}(cx^3))^2} dx$	1117
3.135	$\int x^3(a + \operatorname{barctanh}(\frac{c}{x})) dx$	1122
3.136	$\int x^2(a + \operatorname{barctanh}(\frac{c}{x})) dx$	1128
3.137	$\int x(a + \operatorname{barctanh}(\frac{c}{x})) dx$	1134
3.138	$\int (a + \operatorname{barctanh}(\frac{c}{x})) dx$	1140
3.139	$\int \frac{a+\operatorname{barctanh}(\frac{c}{x})}{x} dx$	1145

3.140	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x})}{x^2} dx$	1150
3.141	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x})}{x^3} dx$	1155
3.142	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x})}{x^4} dx$	1161
3.143	$\int x^3 (a + b \operatorname{arctanh}(\frac{c}{x}))^2 dx$	1168
3.144	$\int x^2 (a + b \operatorname{arctanh}(\frac{c}{x}))^2 dx$	1178
3.145	$\int x (a + b \operatorname{arctanh}(\frac{c}{x}))^2 dx$	1186
3.146	$\int (a + b \operatorname{arctanh}(\frac{c}{x}))^2 dx$	1194
3.147	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x} dx$	1201
3.148	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^2} dx$	1209
3.149	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^3} dx$	1216
3.150	$\int x^3 (a + b \operatorname{arctanh}(\frac{c}{x}))^3 dx$	1224
3.151	$\int x^2 (a + b \operatorname{arctanh}(\frac{c}{x}))^3 dx$	1234
3.152	$\int x (a + b \operatorname{arctanh}(\frac{c}{x}))^3 dx$	1245
3.153	$\int (a + b \operatorname{arctanh}(\frac{c}{x}))^3 dx$	1254
3.154	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x} dx$	1262
3.155	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^2} dx$	1271
3.156	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^3} dx$	1279
3.157	$\int x^7 (a + b \operatorname{arctanh}(\frac{c}{x^2})) dx$	1288
3.158	$\int x^5 (a + b \operatorname{arctanh}(\frac{c}{x^2})) dx$	1295
3.159	$\int x^3 (a + b \operatorname{arctanh}(\frac{c}{x^2})) dx$	1302
3.160	$\int x (a + b \operatorname{arctanh}(\frac{c}{x^2})) dx$	1308
3.161	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{x} dx$	1314
3.162	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{x^3} dx$	1319
3.163	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{x^5} dx$	1325
3.164	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{x^7} dx$	1332
3.165	$\int x^4 (a + b \operatorname{arctanh}(\frac{c}{x^2})) dx$	1339
3.166	$\int x^2 (a + b \operatorname{arctanh}(\frac{c}{x^2})) dx$	1347
3.167	$\int (a + b \operatorname{arctanh}(\frac{c}{x^2})) dx$	1355
3.168	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{x^2} dx$	1361
3.169	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{x^4} dx$	1368
3.170	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{x^6} dx$	1376

3.171	$\int x^3 \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2 dx$	1384
3.172	$\int x \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2 dx$	1394
3.173	$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x} dx$	1401
3.174	$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^3} dx$	1408
3.175	$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^5} dx$	1415
3.176	$\int x^4 \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2 dx$	1422
3.177	$\int x^2 \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2 dx$	1430
3.178	$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2 dx$	1438
3.179	$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^2} dx$	1446
3.180	$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^4} dx$	1454
3.181	$\int (dx)^m \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^3 dx$	1462
3.182	$\int (dx)^m \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2 dx$	1467
3.183	$\int (dx)^m \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right) dx$	1472
3.184	$\int \frac{(dx)^m}{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)} dx$	1477
3.185	$\int \frac{(dx)^m}{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2} dx$	1482
3.186	$\int x^3 \left(a + b \operatorname{arctanh}(c\sqrt{x})\right) dx$	1487
3.187	$\int x^2 \left(a + b \operatorname{arctanh}(c\sqrt{x})\right) dx$	1494
3.188	$\int x \left(a + b \operatorname{arctanh}(c\sqrt{x})\right) dx$	1501
3.189	$\int \left(a + b \operatorname{arctanh}(c\sqrt{x})\right) dx$	1508
3.190	$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x} dx$	1513
3.191	$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2} dx$	1518
3.192	$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3} dx$	1524
3.193	$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4} dx$	1531
3.194	$\int x^3 \left(a + b \operatorname{arctanh}(c\sqrt{x})\right)^2 dx$	1538
3.195	$\int x^2 \left(a + b \operatorname{arctanh}(c\sqrt{x})\right)^2 dx$	1550
3.196	$\int x \left(a + b \operatorname{arctanh}(c\sqrt{x})\right)^2 dx$	1559
3.197	$\int \left(a + b \operatorname{arctanh}(c\sqrt{x})\right)^2 dx$	1567
3.198	$\int \frac{\left(a + b \operatorname{arctanh}(c\sqrt{x})\right)^2}{x} dx$	1574
3.199	$\int \frac{\left(a + b \operatorname{arctanh}(c\sqrt{x})\right)^2}{x^2} dx$	1582
3.200	$\int \frac{\left(a + b \operatorname{arctanh}(c\sqrt{x})\right)^2}{x^3} dx$	1591
3.201	$\int x^3 \left(a + b \operatorname{arctanh}(c\sqrt{x})\right)^3 dx$	1602

3.202	$\int x^2 (a + \operatorname{barctanh}(c\sqrt{x}))^3 dx$	1617
3.203	$\int x (a + \operatorname{barctanh}(c\sqrt{x}))^3 dx$	1631
3.204	$\int (a + \operatorname{barctanh}(c\sqrt{x}))^3 dx$	1642
3.205	$\int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{x} dx$	1651
3.206	$\int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{x^2} dx$	1661
3.207	$\int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{x^3} dx$	1670
3.208	$\int x^{3/2} \operatorname{arctanh}(\sqrt{x}) dx$	1680
3.209	$\int \sqrt{x} \operatorname{arctanh}(\sqrt{x}) dx$	1686
3.210	$\int \frac{\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} dx$	1691
3.211	$\int \frac{\operatorname{arctanh}(\sqrt{x})}{x^{3/2}} dx$	1696
3.212	$\int x^3 (a + \operatorname{barctanh}(cx^{3/2})) dx$	1702
3.213	$\int x^2 (a + \operatorname{barctanh}(cx^{3/2})) dx$	1714
3.214	$\int x (a + \operatorname{barctanh}(cx^{3/2})) dx$	1720
3.215	$\int (a + \operatorname{barctanh}(cx^{3/2})) dx$	1731
3.216	$\int \frac{a + \operatorname{barctanh}(cx^{3/2})}{x} dx$	1738
3.217	$\int \frac{a + \operatorname{barctanh}(cx^{3/2})}{x^2} dx$	1743
3.218	$\int \frac{a + \operatorname{barctanh}(cx^{3/2})}{x^3} dx$	1753
3.219	$\int \frac{a + \operatorname{barctanh}(cx^{3/2})}{x^4} dx$	1764
3.220	$\int x^2 (a + \operatorname{barctanh}(cx^{3/2}))^2 dx$	1770
3.221	$\int \frac{(a + \operatorname{barctanh}(cx^{3/2}))^2}{x} dx$	1777
3.222	$\int \frac{(a + \operatorname{barctanh}(cx^{3/2}))^2}{x^4} dx$	1785
3.223	$\int x^2 (a + \operatorname{barctanh}(cx^n)) dx$	1794
3.224	$\int x (a + \operatorname{barctanh}(cx^n)) dx$	1799
3.225	$\int (a + \operatorname{barctanh}(cx^n)) dx$	1804
3.226	$\int \frac{a + \operatorname{barctanh}(cx^n)}{x} dx$	1809
3.227	$\int \frac{a + \operatorname{barctanh}(cx^n)}{x^2} dx$	1814
3.228	$\int \frac{a + \operatorname{barctanh}(cx^n)}{x^3} dx$	1819
3.229	$\int \frac{a + \operatorname{barctanh}(cx^n)}{x^4} dx$	1824
3.230	$\int x (a + \operatorname{barctanh}(cx^n))^2 dx$	1829
3.231	$\int (a + \operatorname{barctanh}(cx^n))^2 dx$	1834
3.232	$\int \frac{(a + \operatorname{barctanh}(cx^n))^2}{x} dx$	1839
3.233	$\int \frac{(a + \operatorname{barctanh}(cx^n))^2}{x^2} dx$	1846

3.234	$\int \frac{(a+b\operatorname{arctanh}(cx^n))^2}{x^3} dx$	1851
3.235	$\int \frac{\operatorname{arctanh}(ax^n)}{x} dx$	1856
3.236	$\int \frac{\operatorname{arctanh}(ax^5)}{x} dx$	1861
3.237	$\int \operatorname{arctanh}\left(\frac{1}{x}\right) dx$	1866
3.238	$\int (dx)^m (a+b\operatorname{arctanh}(cx^n))^3 dx$	1871
3.239	$\int (dx)^m (a+b\operatorname{arctanh}(cx^n))^2 dx$	1876
3.240	$\int (dx)^m (a+b\operatorname{arctanh}(cx^n)) dx$	1881
3.241	$\int \frac{(dx)^m}{a+b\operatorname{arctanh}(cx^n)} dx$	1886
3.242	$\int \frac{(dx)^m}{(a+b\operatorname{arctanh}(cx^n))^2} dx$	1891
4	Appendix	1896
4.1	Listing of Grading functions	1896
4.2	Links to plain text integration problems used in this report for each CAS	1914

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	11
1.2	Results	12
1.3	Time and leaf size Performance	16
1.4	Performance based on number of rules Rubi used	18
1.5	Performance based on number of steps Rubi used	19
1.6	Solved integrals histogram based on leaf size of result	20
1.7	Solved integrals histogram based on CPU time used	21
1.8	Leaf size vs. CPU time used	22
1.9	list of integrals with no known antiderivative	23
1.10	List of integrals solved by CAS but has no known antiderivative	23
1.11	list of integrals solved by CAS but failed verification	23
1.12	Timing	24
1.13	Verification	24
1.14	Important notes about some of the results	25
1.15	Current tree layout of integration tests	28
1.16	Design of the test system	29

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [242]. This is test number [337].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (242)	0.00 (0)
Mathematica	95.45 (231)	4.55 (11)
Maple	84.71 (205)	15.29 (37)
Maxima	63.64 (154)	36.36 (88)
Fricas	60.74 (147)	39.26 (95)
Reduce	59.92 (145)	40.08 (97)
Mupad	52.89 (128)	47.11 (114)
Giac	52.48 (127)	47.52 (115)
Sympy	33.88 (82)	66.12 (160)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	88.017	1.240	0.000	10.744
Mathematica	75.207	0.826	8.678	15.289
Maple	50.000	9.504	14.876	25.620
Maxima	42.975	9.917	0.000	47.107
Fricas	40.496	4.959	4.545	50.000
Giac	18.182	23.140	0.413	58.264
Sympy	16.116	11.157	0.000	72.727
Mupad	0.000	42.149	0.000	57.851
Reduce	0.000	49.174	0.000	50.826

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	11	100.00	0.00	0.00
Maple	37	97.30	2.70	0.00
Maxima	88	100.00	0.00	0.00
Fricas	95	100.00	0.00	0.00
Reduce	97	100.00	0.00	0.00
Mupad	114	0.00	100.00	0.00
Giac	115	100.00	0.00	0.00
Sympy	160	64.38	35.62	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Fricas	0.11
Reduce	0.18
Giac	0.21
Maxima	0.21
Mathematica	0.47
Rubi	0.91
Maple	3.66
Mupad	4.19
Sympy	7.63

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	91.72	1.33	59.00	1.12
Reduce	92.19	1.65	75.00	1.21
Mathematica	129.17	1.25	99.00	1.17
Maxima	157.56	2.96	97.00	1.19
Fricas	161.18	1.95	69.00	1.47
Giac	163.75	2.50	109.00	1.44
Sympy	232.59	4.06	70.50	1.29
Rubi	243.16	1.09	89.50	1.04
Maple	409.34	3.35	93.00	1.14

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{43, 44, 46, 47, 48, 49, 94, 95, 97, 98, 130, 131, 133, 134, 181, 182, 184, 185, 230, 231, 233, 234, 238, 239, 241, 242}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {90, 93, 143, 194, 195, 200}

Mathematica {}

Maple {19, 25, 27, 30, 31, 32, 33, 34, 69, 77, 117, 121, 123, 125, 147, 150, 151, 152, 153, 154, 156, 171, 172, 198, 201, 202, 203, 204, 205, 206, 207, 221, 222, 232}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals.

These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	31
2.2	Detailed conclusion table per each integral for all CAS systems	37
2.3	Detailed conclusion table specific for Rubi results	98

2.1 List of integrals sorted by grade for each CAS

Rubi	31
Mma	32
Maple	32
Fricas	33
Maxima	33
Giac	34
Mupad	34
Sympy	35
Reduce	35

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 45, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 96, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 132, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 183, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 232, 235, 236, 237, 240 }

B grade { 24, 201, 202 }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 32, 34, 35, 36, 37, 38, 39, 40, 41, 42, 45, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 69, 70, 73, 74, 77, 78, 81, 82, 83, 84, 85, 86, 87, 88, 89, 96, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 121, 122, 123, 124, 125, 126, 129, 132, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 148, 149, 150, 152, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 174, 175, 178, 179, 183, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 199, 200, 201, 202, 203, 204, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 220, 223, 224, 225, 227, 228, 229, 236, 237, 240 }

B grade { 219, 222 }

C grade { 19, 30, 31, 33, 68, 79, 80, 120, 127, 128, 147, 151, 153, 154, 173, 198, 205, 221, 226, 232, 235 }

F normal fail { 71, 72, 75, 76, 90, 91, 92, 93, 176, 177, 180 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 21, 23, 35, 36, 37, 38, 39, 40, 41, 42, 50, 51, 52, 53, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 66, 67, 70, 82, 83, 84, 85, 86, 87, 88, 89, 99, 100, 101, 102, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 118, 119, 122, 135, 136, 137, 138, 140, 141, 142, 143, 145, 148, 149, 157, 158, 159, 160, 162, 163, 164, 165, 166, 167, 168, 169, 170, 174, 175, 186, 187, 188, 189, 191, 192, 193, 194, 208, 209, 210, 211, 212, 213, 214, 215, 217, 218, 219, 226, 235, 237 }

B grade { 20, 22, 24, 26, 28, 29, 54, 65, 78, 126, 139, 144, 146, 155, 161, 190, 195, 196, 197, 199, 200, 216, 220 }

C grade { 19, 25, 27, 30, 31, 32, 33, 34, 69, 77, 103, 117, 121, 123, 125, 147, 150, 151, 152, 153, 154, 156, 171, 172, 198, 201, 202, 203, 204, 205, 206, 207, 221, 222, 232, 236 }

F normal fail { 45, 68, 71, 72, 73, 74, 75, 76, 79, 80, 81, 90, 91, 92, 93, 96, 120, 124, 127, 128, 129, 132, 173, 176, 177, 178, 179, 180, 183, 223, 224, 225, 227, 228, 229, 240 }

F(-1) timedout fail { 242 }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13, 15, 17, 21, 23, 35, 36, 37, 38, 39, 40, 41, 42, 50, 51, 52, 53, 55, 56, 57, 58, 59, 61, 62, 63, 64, 66, 70, 99, 100, 101, 102, 104, 105, 106, 108, 110, 112, 114, 115, 116, 118, 122, 135, 136, 137, 138, 140, 141, 142, 143, 145, 149, 157, 158, 159, 160, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 175, 186, 187, 188, 189, 191, 192, 193, 194, 195, 196, 200, 208, 209, 210, 211, 213, 218, 219, 237 }

B grade { 60, 107, 109, 111, 113, 197, 199, 217, 220, 222, 226, 235 }

C grade { 82, 83, 84, 85, 86, 87, 88, 89, 212, 214, 215 }

F normal fail { 7, 14, 16, 18, 19, 20, 22, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 45, 54, 65, 67, 68, 69, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 90, 91, 92, 93, 96, 103, 117, 119, 120, 121, 123, 124, 125, 126, 127, 128, 129, 132, 139, 144, 146, 147, 148, 150, 151, 152, 153, 154, 155, 156, 161, 172, 173, 174, 176, 177, 178, 179, 180, 183, 190, 198, 201, 202, 203, 204, 205, 206, 207, 216, 221, 223, 224, 225, 227, 228, 229, 232, 236, 240 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maxima

A grade { 1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13, 15, 35, 36, 37, 38, 39, 40, 41, 42, 50, 51, 52, 53, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 82, 83, 84, 85, 86, 87, 88, 89, 99, 100, 101, 102, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 135, 136, 137, 138, 140, 141, 142, 143, 145, 157, 158, 159, 160, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 186, 187, 188, 189, 191, 192, 193, 194, 195, 208, 209, 210, 211, 212, 213, 214, 215, 217, 218, 219, 237 }

B grade { 17, 21, 23, 66, 70, 118, 122, 149, 175, 190, 196, 197, 199, 200, 201, 202, 203, 206, 207, 216, 220, 222, 235, 236 }

C grade { }

F normal fail { 7, 14, 16, 18, 19, 20, 22, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 45, 54, 65, 67, 68, 69, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 90, 91, 92, 93, 96, 103, 117, 119, 120, 121, 123, 124, 125, 126, 127, 128, 129, 132, 139, 144, 146, 147, 148, 150, 151, 152, 153, 154, 155, 156, 161, 172, 173, 174, 176, 177, 178, 179, 180, 183, 198, 204, 205, 221, 223, 224, 225, 226, 227, 228, 229, 232, 240 }

F(-1) timedout fail { }

F(-2) exception fail { }

Giac

A grade { 38, 39, 40, 41, 42, 50, 51, 55, 56, 57, 58, 59, 63, 64, 99, 100, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 157, 158, 162, 163, 164, 165, 166, 167, 168, 169, 170, 215, 217, 218, 219 }

B grade { 1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13, 15, 17, 21, 23, 52, 53, 60, 61, 62, 66, 85, 86, 87, 88, 89, 101, 102, 118, 135, 136, 137, 138, 140, 141, 142, 143, 145, 149, 159, 160, 171, 186, 187, 188, 189, 191, 192, 193, 208, 209, 210, 211, 213, 237 }

C grade { 212 }

F normal fail { 7, 14, 16, 18, 19, 20, 22, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 45, 54, 65, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 90, 91, 92, 93, 96, 103, 117, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 132, 139, 144, 146, 147, 148, 150, 151, 152, 153, 154, 155, 156, 161, 172, 173, 174, 175, 176, 177, 178, 179, 180, 183, 190, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 214, 216, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 232, 235, 236, 240 }

F(-1) timeout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13, 15, 17, 21, 23, 50, 51, 52, 53, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 66, 70, 99, 100, 101, 102, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 118, 122, 135, 136, 137, 138, 140, 141, 142, 143, 145, 149, 157, 158, 159, 160, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 175, 186, 187, 188, 189, 191, 192, 193, 194, 195, 196, 197, 199, 200, 208, 210, 211, 212, 213, 214, 215, 217, 218, 219, 220, 222, 237 }

C grade { }

F normal fail { }

F(-1) timeout fail { 7, 14, 16, 18, 19, 20, 22, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 45, 54, 65, 67, 68, 69, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 96, 103, 117, 119, 120, 121, 123, 124, 125, 126, 127, 128, 129, 132, 139, 144, 146, 147, 148, 150, 151, 152, 153, 154, 155, 156, 161, 172, 173, 174, 176, 177, 178, 179, 180, 183, 190, 198, 201, 202, 203, 204, 205, 206, 207, 209, 216, 221, 223, 224, 225, 226, 227, 228, 229, 232, 235, 236, 240 }

F(-2) exception fail { }

Sympy

A grade { 1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13, 15, 17, 21, 23, 50, 52, 56, 57, 60, 64, 135, 136, 137, 138, 140, 141, 142, 143, 145, 149, 157, 159, 160, 163, 167, 171, 237 }

B grade { 51, 53, 55, 58, 59, 61, 62, 63, 66, 70, 158, 162, 164, 165, 166, 168, 169, 170, 175, 191, 192, 193, 199, 200, 208, 210, 211 }

C grade { }

F normal fail { 7, 14, 16, 18, 19, 20, 22, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 45, 54, 65, 67, 68, 69, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 83, 84, 85, 86, 87, 90, 91, 92, 93, 96, 103, 120, 139, 144, 146, 147, 148, 150, 151, 152, 153, 154, 155, 156, 161, 172, 173, 174, 176, 177, 178, 179, 180, 183, 186, 187, 188, 189, 190, 194, 195, 196, 197, 198, 201, 202, 203, 204, 205, 206, 207, 209, 223, 224, 225, 226, 227, 229, 232, 235, 240 }

F(-1) timedout fail { 42, 49, 82, 88, 89, 98, 99, 100, 101, 102, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 185, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 228, 236, 238, 239, 242 }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13, 15, 17, 21, 23, 35, 36, 37, 38, 39, 40, 41, 42, 50, 51, 52, 53, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 66, 70, 82, 83, 84, 85, 86, 87, 88, 89, 99, 100, 101, 102, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 118, 122, 135, 136, 137, 138, 140, 141, 142, 143, 145, 149, 157, 158, 159, 160, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 175, 186, 187, 188, 189, 191, 192, 193, 194, 195, 196, 197, 199, 200, 208, 209, 210, 211, 212, 213, 214, 215, 217, 218, 219, 220, 222, 237 }

C grade { }

F normal fail { 7, 14, 16, 18, 19, 20, 22, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 45, 54, 65, 67, 68, 69, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 90, 91, 92, 93, 96, 103, 117, 119, 120, 121, 123, 124, 125, 126, 127, 128, 129, 132, 139, 144, 146, 147, 148, 150, 151, 152, 153, 154, 155, 156, 161, 172, 173, 174, 176, 177, 178, 179, 180, 183, 190, 198, 201, 202, 203, 204, 205, 206, 207, 216, 221, 223, 224, 225, 226, 227, 228, 229, 232, 235, 236, 240 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	57	81	59	70	67	63	442	58	52
N.S.	1	0.97	1.37	1.00	1.19	1.14	1.07	7.49	0.98	0.88
time (sec)	N/A	0.339	0.017	0.228	0.029	0.088	0.417	0.121	0.178	3.779

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	62	58	55	69	68	403	65	53
N.S.	1	1.00	1.09	1.02	0.96	1.21	1.19	7.07	1.14	0.93
time (sec)	N/A	0.348	0.016	0.213	0.030	0.080	0.365	0.122	0.169	3.752

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	47	70	50	61	58	53	296	48	43
N.S.	1	0.98	1.46	1.04	1.27	1.21	1.10	6.17	1.00	0.90
time (sec)	N/A	0.332	0.016	0.200	0.024	0.072	0.306	0.122	0.164	3.754

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	47	51	50	44	58	58	258	56	44
N.S.	1	1.02	1.11	1.09	0.96	1.26	1.26	5.61	1.22	0.96
time (sec)	N/A	0.364	0.015	0.190	0.031	0.087	0.278	0.121	0.159	3.666

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	59	40	50	48	42	148	37	35
N.S.	1	1.00	1.59	1.08	1.35	1.30	1.14	4.00	1.00	0.95
time (sec)	N/A	0.331	0.014	0.203	0.030	0.076	0.236	0.118	0.156	3.636

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	30	29	30	42	27	156	35	27
N.S.	1	1.00	1.00	0.97	1.00	1.40	0.90	5.20	1.17	0.90
time (sec)	N/A	0.245	0.003	0.174	0.024	0.079	0.128	0.116	0.168	3.672

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	24	28	0	0	0	0	17	0
N.S.	1	1.00	0.92	1.08	0.00	0.00	0.00	0.00	0.65	0.00
time (sec)	N/A	0.315	0.016	0.217	0.000	0.000	0.000	0.000	0.168	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	38	39	42	39	47	41	94	45	33
N.S.	1	1.06	1.08	1.17	1.08	1.31	1.14	2.61	1.25	0.92
time (sec)	N/A	0.269	0.015	0.154	0.031	0.084	0.283	0.120	0.172	3.674

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	34	59	39	45	43	36	135	33	46
N.S.	1	0.92	1.59	1.05	1.22	1.16	0.97	3.65	0.89	1.24
time (sec)	N/A	0.337	0.015	0.187	0.029	0.071	0.233	0.118	0.170	3.748

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	52	59	56	49	59	70	251	64	46
N.S.	1	0.96	1.09	1.04	0.91	1.09	1.30	4.65	1.19	0.85
time (sec)	N/A	0.245	0.016	0.175	0.024	0.092	0.377	0.117	0.166	3.731

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	46	70	43	60	52	46	292	43	59
N.S.	1	0.96	1.46	0.90	1.25	1.08	0.96	6.08	0.90	1.23
time (sec)	N/A	0.258	0.016	0.190	0.025	0.080	0.293	0.132	0.172	3.992

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	62	70	64	61	70	80	397	73	71
N.S.	1	0.95	1.08	0.98	0.94	1.08	1.23	6.11	1.12	1.09
time (sec)	N/A	0.250	0.016	0.210	0.026	0.092	0.526	0.126	0.172	3.819

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	145	204	164	182	215	193	211	889	180	171
N.S.	1	1.41	1.13	1.26	1.48	1.33	1.46	6.13	1.24	1.18
time (sec)	N/A	1.319	0.042	0.588	0.034	0.087	0.520	0.139	0.177	3.971

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	162	199	161	229	0	0	0	0	173	0
N.S.	1	1.23	0.99	1.41	0.00	0.00	0.00	0.00	1.07	0.00
time (sec)	N/A	1.308	0.314	0.783	0.000	0.000	0.000	0.000	0.182	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	137	132	143	189	160	168	603	143	134
N.S.	1	1.21	1.17	1.27	1.67	1.42	1.49	5.34	1.27	1.19
time (sec)	N/A	0.964	0.040	0.520	0.036	0.085	0.418	0.140	0.170	3.937

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	130	142	122	201	0	0	0	0	132	0
N.S.	1	1.09	0.94	1.55	0.00	0.00	0.00	0.00	1.02	0.00
time (sec)	N/A	1.395	0.184	0.710	0.000	0.000	0.000	0.000	0.174	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	78	90	106	158	122	114	301	106	89
N.S.	1	1.04	1.20	1.41	2.11	1.63	1.52	4.01	1.41	1.19
time (sec)	N/A	0.694	0.082	0.464	0.032	0.087	0.308	0.128	0.174	3.846

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	88	82	115	0	0	0	0	56	0
N.S.	1	1.19	1.11	1.55	0.00	0.00	0.00	0.00	0.76	0.00
time (sec)	N/A	0.724	0.131	0.694	0.000	0.000	0.000	0.000	0.178	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	117	143	151	630	0	0	0	0	37	0
N.S.	1	1.22	1.29	5.38	0.00	0.00	0.00	0.00	0.32	0.00
time (sec)	N/A	0.801	0.396	5.319	0.000	0.000	0.000	0.000	0.179	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	77	94	195	0	0	0	0	90	0
N.S.	1	1.08	1.32	2.75	0.00	0.00	0.00	0.00	1.27	0.00
time (sec)	N/A	0.497	0.101	0.533	0.000	0.000	0.000	0.000	0.183	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	77	101	134	151	135	126	278	126	246
N.S.	1	0.96	1.26	1.68	1.89	1.69	1.58	3.48	1.58	3.08
time (sec)	N/A	0.526	0.048	0.332	0.033	0.118	0.358	0.127	0.176	4.211

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	130	119	145	247	0	0	0	0	149	0
N.S.	1	0.92	1.12	1.90	0.00	0.00	0.00	0.00	1.15	0.00
time (sec)	N/A	0.803	0.242	0.550	0.000	0.000	0.000	0.000	0.177	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	136	164	171	224	173	184	612	163	303
N.S.	1	1.16	1.40	1.46	1.91	1.48	1.57	5.23	1.39	2.59
time (sec)	N/A	0.929	0.045	0.346	0.036	0.103	0.472	0.126	0.174	4.499

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	247	499	305	803	0	0	0	0	357	0
N.S.	1	2.02	1.23	3.25	0.00	0.00	0.00	0.00	1.45	0.00
time (sec)	N/A	3.527	0.476	9.291	0.000	0.000	0.000	0.000	0.176	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	262	365	383	1070	0	0	0	0	341	0
N.S.	1	1.39	1.46	4.08	0.00	0.00	0.00	0.00	1.30	0.00
time (sec)	N/A	2.898	0.507	12.007	0.000	0.000	0.000	0.000	0.170	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	185	290	245	667	0	0	0	0	268	0
N.S.	1	1.57	1.32	3.61	0.00	0.00	0.00	0.00	1.45	0.00
time (sec)	N/A	3.065	0.334	8.063	0.000	0.000	0.000	0.000	0.170	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	197	216	250	973	0	0	0	0	259	0
N.S.	1	1.10	1.27	4.94	0.00	0.00	0.00	0.00	1.31	0.00
time (sec)	N/A	1.964	0.368	10.256	0.000	0.000	0.000	0.000	0.179	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	138	161	488	0	0	0	0	185	0
N.S.	1	1.12	1.31	3.97	0.00	0.00	0.00	0.00	1.50	0.00
time (sec)	N/A	0.965	0.240	6.592	0.000	0.000	0.000	0.000	0.187	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	108	123	161	230	0	0	0	0	77	0
N.S.	1	1.14	1.49	2.13	0.00	0.00	0.00	0.00	0.71	0.00
time (sec)	N/A	0.735	0.207	1.191	0.000	0.000	0.000	0.000	0.196	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	184	212	315	1304	0	0	0	0	57	0
N.S.	1	1.15	1.71	7.09	0.00	0.00	0.00	0.00	0.31	0.00
time (sec)	N/A	1.772	0.472	6.395	0.000	0.000	0.000	0.000	0.172	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	102	115	196	1329	0	0	0	0	140	0
N.S.	1	1.13	1.92	13.03	0.00	0.00	0.00	0.00	1.37	0.00
time (sec)	N/A	1.307	0.238	9.424	0.000	0.000	0.000	0.000	0.172	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	123	119	192	4599	0	0	0	0	240	0
N.S.	1	0.97	1.56	37.39	0.00	0.00	0.00	0.00	1.95	0.00
time (sec)	N/A	1.099	0.186	8.773	0.000	0.000	0.000	0.000	0.188	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	200	197	323	1581	0	0	0	0	303	0
N.S.	1	0.98	1.62	7.90	0.00	0.00	0.00	0.00	1.52	0.00
time (sec)	N/A	1.926	0.644	12.881	0.000	0.000	0.000	0.000	0.194	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	187	242	295	1136	0	0	0	0	307	0
N.S.	1	1.29	1.58	6.07	0.00	0.00	0.00	0.00	1.64	0.00
time (sec)	N/A	1.838	0.437	9.128	0.000	0.000	0.000	0.000	0.176	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	142	128	107	134	296	0	0	101	0
N.S.	1	1.15	1.03	0.86	1.08	2.39	0.00	0.00	0.81	0.00
time (sec)	N/A	0.521	0.072	1.166	0.112	0.116	0.000	0.000	0.161	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	123	115	93	118	255	0	0	91	0
N.S.	1	1.16	1.08	0.88	1.11	2.41	0.00	0.00	0.86	0.00
time (sec)	N/A	0.479	0.058	0.372	0.112	0.105	0.000	0.000	0.171	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	119	114	89	119	223	0	0	83	0
N.S.	1	1.12	1.08	0.84	1.12	2.10	0.00	0.00	0.78	0.00
time (sec)	N/A	0.422	0.052	0.394	0.106	0.107	0.000	0.000	0.179	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	92	98	68	103	211	0	88	73	0
N.S.	1	1.08	1.15	0.80	1.21	2.48	0.00	1.04	0.86	0.00
time (sec)	N/A	0.411	0.031	0.416	0.111	0.097	0.000	0.113	0.190	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	95	99	69	94	208	0	93	75	0
N.S.	1	1.12	1.16	0.81	1.11	2.45	0.00	1.09	0.88	0.00
time (sec)	N/A	0.335	0.050	0.507	0.107	0.120	0.000	0.152	0.211	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	120	107	93	101	231	0	117	92	0
N.S.	1	1.12	1.00	0.87	0.94	2.16	0.00	1.09	0.86	0.00
time (sec)	N/A	0.390	0.061	0.462	0.109	0.103	0.000	0.140	0.184	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	122	108	93	112	242	0	117	109	0
N.S.	1	1.14	1.01	0.87	1.05	2.26	0.00	1.09	1.02	0.00
time (sec)	N/A	0.317	0.053	0.389	0.108	0.113	0.000	0.153	0.174	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	125	142	122	107	130	259	0	135	118	0
N.S.	1	1.14	0.98	0.86	1.04	2.07	0.00	1.08	0.94	0.00
time (sec)	N/A	0.302	0.064	0.394	0.107	0.109	0.000	0.170	0.182	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	379	44	15	18	209	18
N.S.	1	1.00	1.12	1.00	23.69	2.75	0.94	1.12	13.06	1.12
time (sec)	N/A	0.198	2.890	0.336	2.222	0.094	5.263	0.169	0.219	3.988

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	218	30	15	18	159	18
N.S.	1	1.00	1.12	1.00	13.62	1.88	0.94	1.12	9.94	1.12
time (sec)	N/A	0.198	1.763	0.288	1.148	0.099	2.844	0.146	0.197	4.086

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	59	0	0	0	0	0	111	0
N.S.	1	1.00	0.82	0.00	0.00	0.00	0.00	0.00	1.54	0.00
time (sec)	N/A	0.233	0.060	0.000	0.000	0.000	0.000	0.000	0.167	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	18	18	14	18	20	18
N.S.	1	1.00	1.12	1.00	1.12	1.12	0.88	1.12	1.25	1.12
time (sec)	N/A	0.215	0.201	0.602	0.080	0.074	1.313	0.132	0.167	3.740

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	116	32	15	18	34	18
N.S.	1	1.00	1.12	1.00	7.25	2.00	0.94	1.12	2.12	1.12
time (sec)	N/A	0.203	0.428	0.569	0.129	0.080	9.552	0.139	0.169	3.648

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	10	12	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	1.00	1.20	1.20	1.20
time (sec)	N/A	0.176	0.316	0.358	0.153	0.095	1.723	0.144	0.174	3.623

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	18	18	0	18	20	18
N.S.	1	1.00	1.12	1.00	1.12	1.12	0.00	1.12	1.25	1.12
time (sec)	N/A	0.197	0.249	0.553	0.181	0.093	0.000	0.169	0.185	3.633

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	53	78	56	69	64	58	78	54	69
N.S.	1	0.98	1.44	1.04	1.28	1.19	1.07	1.44	1.00	1.28
time (sec)	N/A	0.252	0.030	0.303	0.025	0.071	6.084	0.131	0.173	3.878

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	49	53	45	46	62	85	57	58	61
N.S.	1	1.02	1.10	0.94	0.96	1.29	1.77	1.19	1.21	1.27
time (sec)	N/A	0.241	0.025	0.325	0.027	0.077	4.398	0.120	0.168	3.643

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	67	46	58	54	48	181	43	60
N.S.	1	1.00	1.56	1.07	1.35	1.26	1.12	4.21	1.00	1.40
time (sec)	N/A	0.218	0.025	0.275	0.030	0.085	3.067	0.131	0.164	3.749

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	37	42	37	37	50	71	188	43	52
N.S.	1	0.88	1.00	0.88	0.88	1.19	1.69	4.48	1.02	1.24
time (sec)	N/A	0.202	0.015	0.217	0.029	0.088	2.644	0.126	0.167	3.539

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	36	28	126	0	0	0	0	19	0
N.S.	1	1.20	0.93	4.20	0.00	0.00	0.00	0.00	0.63	0.00
time (sec)	N/A	0.246	0.022	0.162	0.000	0.000	0.000	0.000	0.175	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	42	45	47	41	55	80	51	54	55
N.S.	1	1.05	1.12	1.18	1.02	1.38	2.00	1.28	1.35	1.38
time (sec)	N/A	0.221	0.023	0.135	0.026	0.085	4.302	0.118	0.169	3.677

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	38	65	45	51	49	41	67	39	52
N.S.	1	0.93	1.59	1.10	1.24	1.20	1.00	1.63	0.95	1.27
time (sec)	N/A	0.284	0.023	0.192	0.030	0.074	2.932	0.119	0.168	3.889

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	54	61	63	51	65	97	65	68	67
N.S.	1	0.96	1.09	1.12	0.91	1.16	1.73	1.16	1.21	1.20
time (sec)	N/A	0.401	0.022	0.225	0.028	0.091	7.712	0.122	0.171	3.768

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	74	93	53	69	197	185	73	88	72
N.S.	1	1.14	1.43	0.82	1.06	3.03	2.85	1.12	1.35	1.11
time (sec)	N/A	0.381	0.028	0.441	0.103	0.097	4.471	0.192	0.165	3.848

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	70	91	51	66	186	670	75	84	70
N.S.	1	1.11	1.44	0.81	1.05	2.95	10.63	1.19	1.33	1.11
time (sec)	N/A	0.387	0.025	0.349	0.103	0.094	2.928	0.157	0.174	3.832

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	44	57	37	55	160	702	83	71	55
N.S.	1	1.00	1.30	0.84	1.25	3.64	15.95	1.89	1.61	1.25
time (sec)	N/A	0.290	0.016	0.209	0.105	0.093	2.211	0.118	0.177	3.540

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	55	75	42	61	157	1374	79	70	62
N.S.	1	1.20	1.63	0.91	1.33	3.41	29.87	1.72	1.52	1.35
time (sec)	N/A	0.304	0.025	0.198	0.110	0.099	3.944	0.143	0.165	3.744

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	69	91	51	65	181	1904	95	89	71
N.S.	1	1.10	1.44	0.81	1.03	2.87	30.22	1.51	1.41	1.13
time (sec)	N/A	0.313	0.030	0.296	0.105	0.091	5.297	0.142	0.167	3.845

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	71	91	51	66	187	1948	91	98	71
N.S.	1	1.13	1.44	0.81	1.05	2.97	30.92	1.44	1.56	1.13
time (sec)	N/A	0.359	0.031	0.353	0.110	0.087	6.924	0.168	0.169	3.884

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	125	153	146	163	217	176	206	175	159	335
N.S.	1	1.22	1.17	1.30	1.74	1.41	1.65	1.40	1.27	2.68
time (sec)	N/A	1.025	0.053	0.732	0.035	0.080	7.684	0.175	0.165	4.408

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	146	162	132	380	0	0	0	0	151	0
N.S.	1	1.11	0.90	2.60	0.00	0.00	0.00	0.00	1.03	0.00
time (sec)	N/A	0.957	0.199	1.010	0.000	0.000	0.000	0.000	0.168	0.000

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	92	106	121	186	138	163	361	120	275
N.S.	1	1.01	1.16	1.33	2.04	1.52	1.79	3.97	1.32	3.02
time (sec)	N/A	0.549	0.038	0.664	0.034	0.075	4.514	0.154	0.156	4.147

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	104	99	134	0	0	0	0	68	0
N.S.	1	1.11	1.05	1.43	0.00	0.00	0.00	0.00	0.72	0.00
time (sec)	N/A	0.566	0.120	1.056	0.000	0.000	0.000	0.000	0.167	0.000

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	137	163	181	0	0	0	0	0	41	0
N.S.	1	1.19	1.32	0.00	0.00	0.00	0.00	0.00	0.30	0.00
time (sec)	N/A	0.852	0.299	0.000	0.000	0.000	0.000	0.000	0.169	0.000

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	87	91	119	820	0	0	0	0	105	0
N.S.	1	1.05	1.37	9.43	0.00	0.00	0.00	0.00	1.21	0.00
time (sec)	N/A	0.579	0.112	0.856	0.000	0.000	0.000	0.000	0.169	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	87	111	152	175	151	175	0	140	278
N.S.	1	0.99	1.26	1.73	1.99	1.72	1.99	0.00	1.59	3.16
time (sec)	N/A	0.637	0.060	0.526	0.035	0.087	6.564	0.000	0.167	3.964

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	1173	1173	0	0	0	0	0	0	227	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.19	0.00
time (sec)	N/A	2.624	0.000	0.000	0.000	0.000	0.000	0.000	0.173	0.000

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	1129	1129	0	0	0	0	0	0	205	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.18	0.00
time (sec)	N/A	2.497	0.000	0.000	0.000	0.000	0.000	0.000	0.157	0.000

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	958	958	566	0	0	0	0	0	91	0
N.S.	1	1.00	0.59	0.00	0.00	0.00	0.00	0.00	0.09	0.00
time (sec)	N/A	2.665	1.813	0.000	0.000	0.000	0.000	0.000	0.164	0.000

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	942	942	566	0	0	0	0	0	95	0
N.S.	1	1.00	0.60	0.00	0.00	0.00	0.00	0.00	0.10	0.00
time (sec)	N/A	1.724	2.317	0.000	0.000	0.000	0.000	0.000	0.164	0.000

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	1102	1102	0	0	0	0	0	0	119	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.11	0.00
time (sec)	N/A	2.102	0.000	0.000	0.000	0.000	0.000	0.000	0.167	0.000

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	1176	1176	0	0	0	0	0	0	128	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.11	0.00
time (sec)	N/A	2.362	0.000	0.000	0.000	0.000	0.000	0.000	0.176	0.000

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	141	158	185	798	0	0	0	0	211	0
N.S.	1	1.12	1.31	5.66	0.00	0.00	0.00	0.00	1.50	0.00
time (sec)	N/A	1.085	0.332	1.177	0.000	0.000	0.000	0.000	0.169	0.000

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	134	143	213	265	0	0	0	0	93	0
N.S.	1	1.07	1.59	1.98	0.00	0.00	0.00	0.00	0.69	0.00
time (sec)	N/A	0.827	0.201	1.674	0.000	0.000	0.000	0.000	0.175	0.000

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	207	240	371	0	0	0	0	0	63	0
N.S.	1	1.16	1.79	0.00	0.00	0.00	0.00	0.00	0.30	0.00
time (sec)	N/A	1.316	0.326	0.000	0.000	0.000	0.000	0.000	0.175	0.000

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	125	133	222	0	0	0	0	0	161	0
N.S.	1	1.06	1.78	0.00	0.00	0.00	0.00	0.00	1.29	0.00
time (sec)	N/A	1.412	0.302	0.000	0.000	0.000	0.000	0.000	0.170	0.000

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	139	137	218	0	0	0	0	0	270	0
N.S.	1	0.99	1.57	0.00	0.00	0.00	0.00	0.00	1.94	0.00
time (sec)	N/A	1.690	0.202	0.000	0.000	0.000	0.000	0.000	0.173	0.000

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F(-1)	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	257	356	241	256	316	434	0	0	246	0
N.S.	1	1.39	0.94	1.00	1.23	1.69	0.00	0.00	0.96	0.00
time (sec)	N/A	0.722	0.122	4.388	0.122	0.115	0.000	0.000	0.179	0.000

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	257	350	240	248	310	376	0	0	243	0
N.S.	1	1.36	0.93	0.96	1.21	1.46	0.00	0.00	0.95	0.00
time (sec)	N/A	0.634	0.083	0.385	0.114	0.108	0.000	0.000	0.181	0.000

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	241	328	227	240	301	385	0	0	228	0
N.S.	1	1.36	0.94	1.00	1.25	1.60	0.00	0.00	0.95	0.00
time (sec)	N/A	0.610	0.068	0.408	0.122	0.108	0.000	0.000	0.173	0.000

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	227	320	227	242	296	354	0	493	226	0
N.S.	1	1.41	1.00	1.07	1.30	1.56	0.00	2.17	1.00	0.00
time (sec)	N/A	0.586	0.051	0.417	0.111	0.105	0.000	0.134	0.173	0.000

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	228	324	268	246	296	388	0	505	244	0
N.S.	1	1.42	1.18	1.08	1.30	1.70	0.00	2.21	1.07	0.00
time (sec)	N/A	0.586	0.092	0.395	0.112	0.102	0.000	0.295	0.172	0.000

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	241	326	268	238	277	421	0	516	254	0
N.S.	1	1.35	1.11	0.99	1.15	1.75	0.00	2.14	1.05	0.00
time (sec)	N/A	0.575	0.101	0.391	0.112	0.100	0.000	0.468	0.177	0.000

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F(-1)	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	257	351	275	250	298	460	0	532	281	0
N.S.	1	1.37	1.07	0.97	1.16	1.79	0.00	2.07	1.09	0.00
time (sec)	N/A	0.629	0.083	0.428	0.112	0.107	0.000	1.587	0.178	0.000

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F(-1)	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	257	349	281	250	297	447	0	519	284	0
N.S.	1	1.36	1.09	0.97	1.16	1.74	0.00	2.02	1.11	0.00
time (sec)	N/A	0.616	0.092	0.407	0.117	0.120	0.000	5.070	0.176	0.000

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	6274	5360	0	0	0	0	0	0	260	0
N.S.	1	0.85	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.00
time (sec)	N/A	12.418	0.000	0.000	0.000	0.000	0.000	0.000	0.172	0.000

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	6127	5216	0	0	0	0	0	0	257	0
N.S.	1	0.85	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.00
time (sec)	N/A	11.694	0.000	0.000	0.000	0.000	0.000	0.000	0.179	0.000

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	6281	5171	0	0	0	0	0	0	282	0
N.S.	1	0.82	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.00
time (sec)	N/A	11.441	0.000	0.000	0.000	0.000	0.000	0.000	0.179	0.000

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	6464	5305	0	0	0	0	0	0	295	0
N.S.	1	0.82	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.00
time (sec)	N/A	10.211	0.000	0.000	0.000	0.000	0.000	0.000	0.189	0.000

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	413	50	17	20	133	20
N.S.	1	1.00	1.11	1.00	22.94	2.78	0.94	1.11	7.39	1.11
time (sec)	N/A	0.189	1.476	0.085	2.477	0.082	52.964	0.190	0.169	3.289

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	236	34	17	20	88	20
N.S.	1	1.00	1.11	1.00	13.11	1.89	0.94	1.11	4.89	1.11
time (sec)	N/A	0.203	0.942	0.079	1.296	0.084	38.196	0.166	0.167	3.336

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	64	0	0	0	0	0	45	0
N.S.	1	1.00	0.86	0.00	0.00	0.00	0.00	0.00	0.61	0.00
time (sec)	N/A	0.244	0.062	0.000	0.000	0.000	0.000	0.000	0.173	0.000

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	20	20	15	20	22	20
N.S.	1	1.00	1.11	1.00	1.11	1.11	0.83	1.11	1.22	1.11
time (sec)	N/A	0.205	0.316	0.073	0.085	0.076	45.029	0.143	0.174	3.469

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	134	36	0	20	38	20
N.S.	1	1.00	1.11	1.00	7.44	2.00	0.00	1.11	2.11	1.11
time (sec)	N/A	0.200	0.332	0.073	0.157	0.079	0.000	0.158	0.166	3.489

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	53	78	56	69	64	0	78	54	69
N.S.	1	0.98	1.44	1.04	1.28	1.19	0.00	1.44	1.00	1.28
time (sec)	N/A	0.243	0.027	0.612	0.029	0.088	0.000	0.132	0.161	3.817

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	49	53	45	46	62	0	57	79	61
N.S.	1	1.02	1.10	0.94	0.96	1.29	0.00	1.19	1.65	1.27
time (sec)	N/A	0.237	0.029	0.477	0.029	0.082	0.000	0.123	0.162	3.390

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	67	46	58	54	0	181	43	60
N.S.	1	1.00	1.56	1.07	1.35	1.26	0.00	4.21	1.00	1.40
time (sec)	N/A	0.221	0.022	0.397	0.025	0.093	0.000	0.126	0.164	3.490

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	37	42	37	37	50	0	188	63	52
N.S.	1	0.88	1.00	0.88	0.88	1.19	0.00	4.48	1.50	1.24
time (sec)	N/A	0.204	0.014	0.270	0.024	0.095	0.000	0.128	0.165	3.317

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	36	28	92	0	0	0	0	19	0
N.S.	1	1.20	0.93	3.07	0.00	0.00	0.00	0.00	0.63	0.00
time (sec)	N/A	0.244	0.029	0.204	0.000	0.000	0.000	0.000	0.175	0.000

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	42	45	47	41	55	0	51	79	55
N.S.	1	1.05	1.12	1.18	1.02	1.38	0.00	1.28	1.98	1.38
time (sec)	N/A	0.225	0.023	0.210	0.032	0.090	0.000	0.118	0.175	3.398

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	38	65	45	51	49	0	67	39	52
N.S.	1	0.93	1.59	1.10	1.24	1.20	0.00	1.63	0.95	1.27
time (sec)	N/A	0.220	0.023	0.300	0.031	0.070	0.000	0.126	0.176	3.571

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	54	61	63	51	65	0	65	95	67
N.S.	1	0.96	1.09	1.12	0.91	1.16	0.00	1.16	1.70	1.20
time (sec)	N/A	0.246	0.024	0.426	0.026	0.082	0.000	0.117	0.167	3.441

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	177	196	184	162	981	0	207	112	125
N.S.	1	1.26	1.40	1.31	1.16	7.01	0.00	1.48	0.80	0.89
time (sec)	N/A	0.411	0.051	0.362	0.112	0.110	0.000	0.163	0.164	4.004

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	101	136	99	90	260	0	109	119	107
N.S.	1	1.00	1.35	0.98	0.89	2.57	0.00	1.08	1.18	1.06
time (sec)	N/A	0.271	0.030	0.189	0.107	0.092	0.000	0.114	0.165	5.537

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	166	187	159	155	228	0	165	122	118
N.S.	1	1.27	1.43	1.21	1.18	1.74	0.00	1.26	0.93	0.90
time (sec)	N/A	0.374	0.053	0.292	0.115	0.106	0.000	0.168	0.174	4.076

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	127	196	172	100	151	0	125	166	135
N.S.	1	1.10	1.70	1.50	0.87	1.31	0.00	1.09	1.44	1.17
time (sec)	N/A	0.462	0.045	0.405	0.105	0.085	0.000	0.122	0.175	6.000

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	142	191	198	186	164	248	0	208	114	127
N.S.	1	1.35	1.39	1.31	1.15	1.75	0.00	1.46	0.80	0.89
time (sec)	N/A	0.656	0.036	0.641	0.108	0.099	0.000	0.204	0.169	4.008

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	125	198	114	103	149	0	126	132	124
N.S.	1	1.07	1.69	0.97	0.88	1.27	0.00	1.08	1.13	1.06
time (sec)	N/A	0.536	0.034	0.392	0.111	0.085	0.000	0.120	0.164	5.216

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	176	187	177	155	238	0	173	105	118
N.S.	1	1.34	1.43	1.35	1.18	1.82	0.00	1.32	0.80	0.90
time (sec)	N/A	0.552	0.028	0.283	0.126	0.083	0.000	0.202	0.163	4.006

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	104	110	183	105	94	117	0	106	133	117
N.S.	1	1.06	1.76	1.01	0.90	1.12	0.00	1.02	1.28	1.12
time (sec)	N/A	0.315	0.030	0.284	0.108	0.086	0.000	0.122	0.172	5.310

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	186	196	172	160	196	0	193	141	125
N.S.	1	1.33	1.40	1.23	1.14	1.40	0.00	1.38	1.01	0.89
time (sec)	N/A	0.407	0.046	0.362	0.114	0.082	0.000	0.335	0.174	4.151

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	125	153	146	163	217	176	0	175	182	335
N.S.	1	1.22	1.17	1.30	1.74	1.41	0.00	1.40	1.46	2.68
time (sec)	N/A	0.914	0.054	1.534	0.040	0.088	0.000	0.179	0.198	4.339

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	146	162	132	2906	0	0	0	0	175	0
N.S.	1	1.11	0.90	19.90	0.00	0.00	0.00	0.00	1.20	0.00
time (sec)	N/A	0.893	0.206	1.361	0.000	0.000	0.000	0.000	0.206	0.000

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	92	106	121	186	138	0	361	143	275
N.S.	1	1.01	1.16	1.33	2.04	1.52	0.00	3.97	1.57	3.02
time (sec)	N/A	0.516	0.040	1.030	0.036	0.086	0.000	0.159	0.201	4.175

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	96	104	99	134	0	0	0	0	92	0
N.S.	1	1.08	1.03	1.40	0.00	0.00	0.00	0.00	0.96	0.00
time (sec)	N/A	0.561	0.108	1.100	0.000	0.000	0.000	0.000	0.219	0.000

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	140	163	181	0	0	0	0	0	41	0
N.S.	1	1.16	1.29	0.00	0.00	0.00	0.00	0.00	0.29	0.00
time (sec)	N/A	0.838	0.307	0.000	0.000	0.000	0.000	0.000	0.187	0.000

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	90	91	117	2993	0	0	0	0	131	0
N.S.	1	1.01	1.30	33.26	0.00	0.00	0.00	0.00	1.46	0.00
time (sec)	N/A	0.556	0.118	0.866	0.000	0.000	0.000	0.000	0.178	0.000

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	87	111	152	175	151	0	0	169	278
N.S.	1	0.99	1.26	1.73	1.99	1.72	0.00	0.00	1.92	3.16
time (sec)	N/A	0.657	0.067	0.665	0.035	0.090	0.000	0.000	0.180	4.267

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	144	137	159	3062	0	0	0	0	191	0
N.S.	1	0.95	1.10	21.26	0.00	0.00	0.00	0.00	1.33	0.00
time (sec)	N/A	1.222	0.268	1.242	0.000	0.000	0.000	0.000	0.177	0.000

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	231	244	334	0	0	0	0	0	346	0
N.S.	1	1.06	1.45	0.00	0.00	0.00	0.00	0.00	1.50	0.00
time (sec)	N/A	2.242	0.350	0.000	0.000	0.000	0.000	0.000	0.196	0.000

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	139	158	185	800	0	0	0	0	235	0
N.S.	1	1.14	1.33	5.76	0.00	0.00	0.00	0.00	1.69	0.00
time (sec)	N/A	1.062	0.086	1.608	0.000	0.000	0.000	0.000	0.181	0.000

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	130	143	191	265	0	0	0	0	121	0
N.S.	1	1.10	1.47	2.04	0.00	0.00	0.00	0.00	0.93	0.00
time (sec)	N/A	0.806	0.176	1.937	0.000	0.000	0.000	0.000	0.193	0.000

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	210	240	368	0	0	0	0	0	63	0
N.S.	1	1.14	1.75	0.00	0.00	0.00	0.00	0.00	0.30	0.00
time (sec)	N/A	1.180	0.323	0.000	0.000	0.000	0.000	0.000	0.186	0.000

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	120	133	223	0	0	0	0	0	189	0
N.S.	1	1.11	1.86	0.00	0.00	0.00	0.00	0.00	1.58	0.00
time (sec)	N/A	0.852	0.287	0.000	0.000	0.000	0.000	0.000	0.195	0.000

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	136	137	218	0	0	0	0	0	300	0
N.S.	1	1.01	1.60	0.00	0.00	0.00	0.00	0.00	2.21	0.00
time (sec)	N/A	1.059	0.200	0.000	0.000	0.000	0.000	0.000	0.190	0.000

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	416	50	0	20	133	20
N.S.	1	1.00	1.11	1.00	23.11	2.78	0.00	1.11	7.39	1.11
time (sec)	N/A	0.253	1.588	0.098	2.491	0.084	0.000	0.182	0.197	3.561

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	239	34	0	20	88	20
N.S.	1	1.00	1.11	1.00	13.28	1.89	0.00	1.11	4.89	1.11
time (sec)	N/A	0.255	0.991	0.091	1.293	0.083	0.000	0.159	0.190	3.552

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	64	0	0	0	0	0	45	0
N.S.	1	1.00	0.86	0.00	0.00	0.00	0.00	0.00	0.61	0.00
time (sec)	N/A	0.390	0.063	0.000	0.000	0.000	0.000	0.000	0.184	0.000

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	20	20	0	20	22	20
N.S.	1	1.00	1.11	1.00	1.11	1.11	0.00	1.11	1.22	1.11
time (sec)	N/A	0.335	0.315	0.102	0.083	0.071	0.000	0.143	0.206	3.335

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	141	36	0	20	38	20
N.S.	1	1.00	1.11	1.00	7.83	2.00	0.00	1.11	2.11	1.11
time (sec)	N/A	0.331	0.330	0.076	0.158	0.076	0.000	0.156	0.220	3.603

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	48	67	46	57	48	46	262	45	45
N.S.	1	0.96	1.34	0.92	1.14	0.96	0.92	5.24	0.90	0.90
time (sec)	N/A	0.377	0.018	0.555	0.030	0.086	0.163	0.119	0.174	3.621

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	42	50	57	42	49	49	227	52	42
N.S.	1	0.93	1.11	1.27	0.93	1.09	1.09	5.04	1.16	0.93
time (sec)	N/A	0.345	0.017	0.428	0.026	0.078	0.144	0.120	0.169	3.443

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	35	56	43	44	39	36	130	36	36
N.S.	1	0.90	1.44	1.10	1.13	1.00	0.92	3.33	0.92	0.92
time (sec)	N/A	0.259	0.015	0.473	0.032	0.072	0.130	0.115	0.174	3.580

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	29	32	29	35	24	150	34	27
N.S.	1	1.00	1.00	1.10	1.00	1.21	0.83	5.17	1.17	0.93
time (sec)	N/A	0.254	0.003	0.359	0.027	0.081	0.123	0.122	0.174	3.433

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	33	28	57	0	0	0	0	19	0
N.S.	1	1.10	0.93	1.90	0.00	0.00	0.00	0.00	0.63	0.00
time (sec)	N/A	0.246	0.024	0.405	0.000	0.000	0.000	0.000	0.181	0.000

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	38	37	37	48	39	87	48	43
N.S.	1	1.00	1.09	1.06	1.06	1.37	1.11	2.49	1.37	1.23
time (sec)	N/A	0.255	0.015	0.260	0.029	0.086	0.362	0.119	0.173	3.500

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	60	42	52	46	44	123	43	49
N.S.	1	1.00	1.40	0.98	1.21	1.07	1.02	2.86	1.00	1.14
time (sec)	N/A	0.219	0.016	0.427	0.031	0.070	0.312	0.123	0.167	3.517

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	55	62	57	55	62	68	234	68	59
N.S.	1	1.15	1.29	1.19	1.15	1.29	1.42	4.88	1.42	1.23
time (sec)	N/A	0.249	0.016	0.422	0.031	0.080	0.381	0.126	0.176	3.487

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	140	131	161	189	149	158	552	154	142
N.S.	1	1.14	1.07	1.31	1.54	1.21	1.28	4.49	1.25	1.15
time (sec)	N/A	1.015	0.043	1.833	0.035	0.082	0.237	0.129	0.171	3.607

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	142	131	145	298	0	0	0	0	147	0
N.S.	1	0.92	1.02	2.10	0.00	0.00	0.00	0.00	1.04	0.00
time (sec)	N/A	0.880	0.214	1.483	0.000	0.000	0.000	0.000	0.179	0.000

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	82	92	118	136	111	104	268	112	101
N.S.	1	0.99	1.11	1.42	1.64	1.34	1.25	3.23	1.35	1.22
time (sec)	N/A	0.615	0.029	1.511	0.032	0.081	0.162	0.122	0.169	3.474

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	80	97	235	0	0	0	0	55	0
N.S.	1	1.08	1.31	3.18	0.00	0.00	0.00	0.00	0.74	0.00
time (sec)	N/A	0.524	0.086	0.810	0.000	0.000	0.000	0.000	0.181	0.000

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	133	159	177	704	0	0	0	0	41	0
N.S.	1	1.20	1.33	5.29	0.00	0.00	0.00	0.00	0.31	0.00
time (sec)	N/A	0.849	0.286	13.598	0.000	0.000	0.000	0.000	0.170	0.000

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	101	101	134	0	0	0	0	101	0
N.S.	1	1.16	1.16	1.54	0.00	0.00	0.00	0.00	1.16	0.00
time (sec)	N/A	0.882	0.107	1.355	0.000	0.000	0.000	0.000	0.184	0.000

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	119	132	165	130	124	255	133	235
N.S.	1	1.00	1.37	1.52	1.90	1.49	1.43	2.93	1.53	2.70
time (sec)	N/A	0.859	0.048	2.035	0.037	0.092	0.365	0.126	0.177	4.298

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	203	266	286	1267	0	0	0	0	304	0
N.S.	1	1.31	1.41	6.24	0.00	0.00	0.00	0.00	1.50	0.00
time (sec)	N/A	2.546	0.423	20.358	0.000	0.000	0.000	0.000	0.170	0.000

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	217	216	316	1780	0	0	0	0	282	0
N.S.	1	1.00	1.46	8.20	0.00	0.00	0.00	0.00	1.30	0.00
time (sec)	N/A	2.035	0.563	36.307	0.000	0.000	0.000	0.000	0.189	0.000

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	135	131	193	5036	0	0	0	0	203	0
N.S.	1	0.97	1.43	37.30	0.00	0.00	0.00	0.00	1.50	0.00
time (sec)	N/A	1.046	0.213	22.586	0.000	0.000	0.000	0.000	0.177	0.000

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	108	121	198	1475	0	0	0	0	77	0
N.S.	1	1.12	1.83	13.66	0.00	0.00	0.00	0.00	0.71	0.00
time (sec)	N/A	1.067	0.201	27.389	0.000	0.000	0.000	0.000	0.189	0.000

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	208	236	373	1452	0	0	0	0	63	0
N.S.	1	1.13	1.79	6.98	0.00	0.00	0.00	0.00	0.30	0.00
time (sec)	N/A	1.723	0.322	15.621	0.000	0.000	0.000	0.000	0.188	0.000

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	140	205	265	0	0	0	0	158	0
N.S.	1	1.11	1.63	2.10	0.00	0.00	0.00	0.00	1.25	0.00
time (sec)	N/A	0.890	0.133	2.178	0.000	0.000	0.000	0.000	0.190	0.000

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	139	154	195	6081	0	0	0	0	262	0
N.S.	1	1.11	1.40	43.75	0.00	0.00	0.00	0.00	1.88	0.00
time (sec)	N/A	1.128	0.226	22.868	0.000	0.000	0.000	0.000	0.174	0.000

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	52	73	48	62	53	51	71	47	66
N.S.	1	0.96	1.35	0.89	1.15	0.98	0.94	1.31	0.87	1.22
time (sec)	N/A	0.250	0.025	0.658	0.033	0.082	3.164	0.118	0.189	4.346

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	42	50	59	42	52	75	52	50	56
N.S.	1	0.93	1.11	1.31	0.93	1.16	1.67	1.16	1.11	1.24
time (sec)	N/A	0.244	0.023	0.727	0.025	0.072	2.340	0.121	0.178	3.997

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	39	62	45	49	44	41	162	38	57
N.S.	1	0.91	1.44	1.05	1.14	1.02	0.95	3.77	0.88	1.33
time (sec)	N/A	0.230	0.022	0.550	0.030	0.077	1.687	0.118	0.185	3.894

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	34	39	42	34	43	61	184	39	47
N.S.	1	0.87	1.00	1.08	0.87	1.10	1.56	4.72	1.00	1.21
time (sec)	N/A	0.209	0.015	0.391	0.026	0.072	1.234	0.146	0.182	3.697

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	37	28	149	0	0	0	0	19	0
N.S.	1	1.23	0.93	4.97	0.00	0.00	0.00	0.00	0.63	0.00
time (sec)	N/A	0.244	0.024	0.366	0.000	0.000	0.000	0.000	0.176	0.000

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	42	37	37	55	76	52	54	56
N.S.	1	1.00	1.14	1.00	1.00	1.49	2.05	1.41	1.46	1.51
time (sec)	N/A	0.208	0.015	0.333	0.029	0.080	4.178	0.115	0.176	3.823

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	64	44	56	52	49	66	45	59
N.S.	1	1.00	1.42	0.98	1.24	1.16	1.09	1.47	1.00	1.31
time (sec)	N/A	0.233	0.022	0.377	0.025	0.081	5.207	0.121	0.181	3.806

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	55	62	45	55	67	94	65	68	66
N.S.	1	1.15	1.29	0.94	1.15	1.40	1.96	1.35	1.42	1.38
time (sec)	N/A	0.248	0.022	0.404	0.024	0.086	7.968	0.128	0.172	3.793

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	72	88	53	62	170	845	67	84	67
N.S.	1	1.14	1.40	0.84	0.98	2.70	13.41	1.06	1.33	1.06
time (sec)	N/A	0.245	0.027	0.705	0.103	0.090	3.851	0.151	0.194	3.875

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	65	86	51	61	162	830	69	74	65
N.S.	1	1.07	1.41	0.84	1.00	2.66	13.61	1.13	1.21	1.07
time (sec)	N/A	0.234	0.025	0.542	0.102	0.117	2.798	0.130	0.168	3.734

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	44	54	39	51	138	632	57	56	52
N.S.	1	1.00	1.23	0.89	1.16	3.14	14.36	1.30	1.27	1.18
time (sec)	N/A	0.173	0.014	0.409	0.104	0.095	2.550	0.118	0.172	3.613

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	55	72	44	57	159	886	62	72	59
N.S.	1	1.20	1.57	0.96	1.24	3.46	19.26	1.35	1.57	1.28
time (sec)	N/A	0.223	0.027	0.394	0.106	0.092	3.530	0.140	0.172	3.816

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	72	90	55	64	189	1046	72	91	69
N.S.	1	1.11	1.38	0.85	0.98	2.91	16.09	1.11	1.40	1.06
time (sec)	N/A	0.252	0.031	0.426	0.106	0.109	4.827	0.154	0.174	3.981

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	74	90	55	65	196	994	74	94	69
N.S.	1	1.14	1.38	0.85	1.00	3.02	15.29	1.14	1.45	1.06
time (sec)	N/A	0.235	0.029	0.476	0.107	0.093	6.723	0.175	0.173	3.916

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	94	88	104	805	157	126	151	327	118	247
N.S.	1	0.94	1.11	8.56	1.67	1.34	1.61	3.48	1.26	2.63
time (sec)	N/A	0.596	0.043	0.267	0.032	0.080	2.096	0.143	0.174	4.065

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	94	90	107	889	0	0	0	0	58	0
N.S.	1	0.96	1.14	9.46	0.00	0.00	0.00	0.00	0.62	0.00
time (sec)	N/A	0.571	0.098	1.805	0.000	0.000	0.000	0.000	0.188	0.000

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	144	163	183	0	0	0	0	0	41	0
N.S.	1	1.13	1.27	0.00	0.00	0.00	0.00	0.00	0.28	0.00
time (sec)	N/A	0.834	0.303	0.000	0.000	0.000	0.000	0.000	0.181	0.000

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	105	114	134	0	0	0	0	108	0
N.S.	1	1.06	1.15	1.35	0.00	0.00	0.00	0.00	1.09	0.00
time (sec)	N/A	0.575	0.045	1.598	0.000	0.000	0.000	0.000	0.185	0.000

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	91	131	138	183	143	172	0	135	262
N.S.	1	0.94	1.35	1.42	1.89	1.47	1.77	0.00	1.39	2.70
time (sec)	N/A	0.536	0.059	244.924	0.033	0.089	5.877	0.000	0.177	4.449

Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	1214	1214	0	0	0	0	0	0	233	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.19	0.00
time (sec)	N/A	3.157	0.000	0.000	0.000	0.000	0.000	0.000	0.182	0.000

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	1172	1172	0	0	0	0	0	0	197	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.17	0.00
time (sec)	N/A	2.629	0.000	0.000	0.000	0.000	0.000	0.000	0.173	0.000

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	1050	1050	565	0	0	0	0	0	81	0
N.S.	1	1.00	0.54	0.00	0.00	0.00	0.00	0.00	0.08	0.00
time (sec)	N/A	1.752	2.249	0.000	0.000	0.000	0.000	0.000	0.193	0.000

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	1080	1117	568	0	0	0	0	0	98	0
N.S.	1	1.03	0.53	0.00	0.00	0.00	0.00	0.00	0.09	0.00
time (sec)	N/A	2.428	1.970	0.000	0.000	0.000	0.000	0.000	0.182	0.000

Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	1263	1263	0	0	0	0	0	0	124	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.10	0.00
time (sec)	N/A	2.525	0.000	0.000	0.000	0.000	0.000	0.000	0.186	0.000

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	398	50	17	20	133	20
N.S.	1	1.00	1.11	1.00	22.11	2.78	0.94	1.11	7.39	1.11
time (sec)	N/A	0.200	1.980	0.117	2.541	0.085	54.854	0.168	0.193	3.905

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	228	34	17	20	88	20
N.S.	1	1.00	1.11	1.00	12.67	1.89	0.94	1.11	4.89	1.11
time (sec)	N/A	0.201	1.268	0.119	1.402	0.087	35.633	0.145	0.194	3.968

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	75	78	68	0	0	0	0	0	45	0
N.S.	1	1.04	0.91	0.00	0.00	0.00	0.00	0.00	0.60	0.00
time (sec)	N/A	0.278	0.070	0.000	0.000	0.000	0.000	0.000	0.177	0.000

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	20	20	15	20	22	20
N.S.	1	1.00	1.11	1.00	1.11	1.11	0.83	1.11	1.22	1.11
time (sec)	N/A	0.206	0.304	0.112	0.080	0.075	45.615	0.135	0.186	3.421

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	128	36	0	20	38	20
N.S.	1	1.00	1.11	1.00	7.11	2.00	0.00	1.11	2.11	1.11
time (sec)	N/A	0.204	0.303	0.120	0.153	0.084	0.000	0.133	0.187	3.769

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	101	114	79	86	89	0	359	74	86
N.S.	1	1.15	1.30	0.90	0.98	1.01	0.00	4.08	0.84	0.98
time (sec)	N/A	0.253	0.032	0.296	0.026	0.091	0.000	0.141	0.178	4.616

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	84	101	71	78	80	0	301	63	58
N.S.	1	1.12	1.35	0.95	1.04	1.07	0.00	4.01	0.84	0.77
time (sec)	N/A	0.240	0.026	0.293	0.030	0.084	0.000	0.137	0.172	4.251

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	67	88	63	69	70	0	239	51	49
N.S.	1	1.08	1.42	1.02	1.11	1.13	0.00	3.85	0.82	0.79
time (sec)	N/A	0.222	0.023	0.289	0.032	0.080	0.000	0.134	0.178	4.226

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	42	50	53	56	0	174	35	32
N.S.	1	1.00	1.08	1.28	1.36	1.44	0.00	4.46	0.90	0.82
time (sec)	N/A	0.169	0.037	0.286	0.032	0.090	0.000	0.133	0.170	4.050

Problem 190	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	40	29	57	61	0	0	0	18	0
N.S.	1	1.38	1.00	1.97	2.10	0.00	0.00	0.00	0.62	0.00
time (sec)	N/A	0.247	0.018	0.355	0.125	0.000	0.000	0.000	0.168	0.000

Problem 191	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	43	67	57	51	53	231	168	33	52
N.S.	1	1.08	1.68	1.42	1.28	1.32	5.78	4.20	0.82	1.30
time (sec)	N/A	0.216	0.025	0.305	0.031	0.081	2.555	0.131	0.175	4.348

Problem 192	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	59	86	65	64	64	342	356	46	61
N.S.	1	0.98	1.43	1.08	1.07	1.07	5.70	5.93	0.77	1.02
time (sec)	N/A	0.218	0.028	0.307	0.029	0.087	7.342	0.136	0.173	4.296

Problem 193	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	99	73	72	74	371	534	57	69
N.S.	1	1.00	1.36	1.00	0.99	1.01	5.08	7.32	0.78	0.95
time (sec)	N/A	0.228	0.030	0.305	0.024	0.088	21.144	0.139	0.169	4.065

Problem 194	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	211	305	224	309	265	273	0	0	230	453
N.S.	1	1.45	1.06	1.46	1.26	1.29	0.00	0.00	1.09	2.15
time (sec)	N/A	2.040	0.078	1.033	0.035	0.105	0.000	0.000	0.175	6.488

Problem 195	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	F	F	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	173	226	194	279	241	241	0	0	189	185
N.S.	1	1.31	1.12	1.61	1.39	1.39	0.00	0.00	1.09	1.07
time (sec)	N/A	2.353	0.073	1.014	0.035	0.116	0.000	0.000	0.183	4.259

Problem 196	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	A	F	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	129	159	160	249	215	207	0	0	147	143
N.S.	1	1.23	1.24	1.93	1.67	1.60	0.00	0.00	1.14	1.11
time (sec)	N/A	1.064	0.062	1.035	0.034	0.107	0.000	0.000	0.174	3.977

Problem 197	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	96	115	213	175	165	0	0	104	94
N.S.	1	1.13	1.35	2.51	2.06	1.94	0.00	0.00	1.22	1.11
time (sec)	N/A	0.515	0.038	0.999	0.036	0.102	0.000	0.000	0.182	3.768

Problem 198	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	145	177	203	662	0	0	0	0	39	0
N.S.	1	1.22	1.40	4.57	0.00	0.00	0.00	0.00	0.27	0.00
time (sec)	N/A	0.848	0.328	12.204	0.000	0.000	0.000	0.000	0.173	0.000

Problem 199	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	89	129	231	174	157	680	0	121	278
N.S.	1	1.05	1.52	2.72	2.05	1.85	8.00	0.00	1.42	3.27
time (sec)	N/A	0.626	0.080	0.457	0.038	0.096	2.622	0.000	0.183	4.426

Problem 200	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	A	B	F	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	133	152	178	263	234	201	972	0	168	341
N.S.	1	1.14	1.34	1.98	1.76	1.51	7.31	0.00	1.26	2.56
time (sec)	N/A	1.039	0.092	0.452	0.035	0.097	7.444	0.000	0.175	5.050

Problem 201	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	C	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	374	949	418	1341	1972	0	0	0	469	0
N.S.	1	2.54	1.12	3.59	5.27	0.00	0.00	0.00	1.25	0.00
time (sec)	N/A	5.926	0.748	19.236	0.665	0.000	0.000	0.000	0.187	0.000

Problem 202	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	C	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	304	611	351	1264	1579	0	0	0	374	0
N.S.	1	2.01	1.15	4.16	5.19	0.00	0.00	0.00	1.23	0.00
time (sec)	N/A	4.133	0.535	16.195	0.595	0.000	0.000	0.000	0.175	0.000

Problem 203	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	234	356	285	1145	1184	0	0	0	276	0
N.S.	1	1.52	1.22	4.89	5.06	0.00	0.00	0.00	1.18	0.00
time (sec)	N/A	2.182	0.367	15.424	0.515	0.000	0.000	0.000	0.179	0.000

Problem 204	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	142	170	201	5673	0	0	0	0	183	0
N.S.	1	1.20	1.42	39.95	0.00	0.00	0.00	0.00	1.29	0.00
time (sec)	N/A	1.052	0.188	14.079	0.000	0.000	0.000	0.000	0.173	0.000

Problem 205	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	224	262	423	1363	0	0	0	0	60	0
N.S.	1	1.17	1.89	6.08	0.00	0.00	0.00	0.00	0.27	0.00
time (sec)	N/A	1.203	0.325	14.581	0.000	0.000	0.000	0.000	0.190	0.000

Problem 206	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	142	151	230	4700	528	0	0	0	237	0
N.S.	1	1.06	1.62	33.10	3.72	0.00	0.00	0.00	1.67	0.00
time (sec)	N/A	1.059	0.218	15.121	1.018	0.000	0.000	0.000	0.174	0.000

Problem 207	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	234	306	333	1219	703	0	0	0	318	0
N.S.	1	1.31	1.42	5.21	3.00	0.00	0.00	0.00	1.36	0.00
time (sec)	N/A	2.009	0.488	16.687	1.168	0.000	0.000	0.000	0.182	0.000

Problem 208	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	35	31	35	24	36	121	170	31	24
N.S.	1	0.92	0.82	0.92	0.63	0.95	3.18	4.47	0.82	0.63
time (sec)	N/A	0.216	0.013	0.112	0.030	0.087	0.793	0.118	0.174	3.237

Problem 209	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	28	25	30	19	31	0	121	24	0
N.S.	1	0.90	0.81	0.97	0.61	1.00	0.00	3.90	0.77	0.00
time (sec)	N/A	0.207	0.010	0.076	0.043	0.087	0.000	0.123	0.200	0.000

Problem 210	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	17	16	25	87	72	20	14
N.S.	1	1.00	1.00	0.85	0.80	1.25	4.35	3.60	1.00	0.70
time (sec)	N/A	0.180	0.006	0.097	0.025	0.090	0.205	0.111	0.175	3.268

Problem 211	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	29	18	37	126	72	34	22
N.S.	1	1.00	1.00	1.21	0.75	1.54	5.25	3.00	1.42	0.92
time (sec)	N/A	0.188	0.015	0.138	0.026	0.091	0.375	0.117	0.179	3.250

Problem 212	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F(-1)	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	160	212	222	194	172	2039	0	227	120	231
N.S.	1	1.32	1.39	1.21	1.08	12.74	0.00	1.42	0.75	1.44
time (sec)	N/A	0.456	0.053	0.178	0.109	0.880	0.000	0.195	0.178	16.649

Problem 213	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	54	75	55	58	64	0	97	43	110
N.S.	1	1.10	1.53	1.12	1.18	1.31	0.00	1.98	0.88	2.24
time (sec)	N/A	0.252	0.028	0.356	0.027	0.090	0.000	0.124	0.183	4.416

Problem 214	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F(-1)	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	160	200	222	194	172	1488	0	0	117	247
N.S.	1	1.25	1.39	1.21	1.08	9.30	0.00	0.00	0.73	1.54
time (sec)	N/A	0.513	0.041	0.187	0.114	0.785	0.000	0.000	0.174	15.596

Problem 215	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	170	114	179	158	1914	0	186	105	107
N.S.	1	1.21	0.81	1.28	1.13	13.67	0.00	1.33	0.75	0.76
time (sec)	N/A	0.663	0.108	0.197	0.108	0.854	0.000	0.139	0.185	8.189

Problem 216	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	42	32	57	62	0	0	0	19	0
N.S.	1	1.24	0.94	1.68	1.82	0.00	0.00	0.00	0.56	0.00
time (sec)	N/A	0.412	0.026	0.444	0.169	0.000	0.000	0.000	0.183	0.000

Problem 217	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	142	180	205	167	163	234	0	172	116	220
N.S.	1	1.27	1.44	1.18	1.15	1.65	0.00	1.21	0.82	1.55
time (sec)	N/A	0.579	0.044	0.162	0.115	0.104	0.000	0.212	0.189	10.402

Problem 218	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	158	207	220	180	168	214	0	200	145	228
N.S.	1	1.31	1.39	1.14	1.06	1.35	0.00	1.27	0.92	1.44
time (sec)	N/A	0.704	0.055	0.184	0.108	0.110	0.000	0.220	0.182	10.473

Problem 219	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	49	140	55	51	59	0	67	39	114
N.S.	1	1.04	2.98	1.17	1.09	1.26	0.00	1.43	0.83	2.43
time (sec)	N/A	0.307	0.035	0.176	0.036	0.091	0.000	0.146	0.185	3.922

Problem 220	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	102	122	220	186	179	0	0	143	105
N.S.	1	1.01	1.21	2.18	1.84	1.77	0.00	0.00	1.42	1.04
time (sec)	N/A	0.533	0.072	1.157	0.037	0.114	0.000	0.000	0.180	3.890

Problem 221	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	156	179	207	706	0	0	0	0	41	0
N.S.	1	1.15	1.33	4.53	0.00	0.00	0.00	0.00	0.26	0.00
time (sec)	N/A	0.860	0.327	18.046	0.000	0.000	0.000	0.000	0.191	0.000

Problem 222	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	B	B	F(-1)	F	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	96	95	210	3062	175	173	0	0	170	281
N.S.	1	0.99	2.19	31.90	1.82	1.80	0.00	0.00	1.77	2.93
time (sec)	N/A	0.622	0.094	2.239	0.042	0.105	0.000	0.000	0.197	4.687

Problem 223	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	73	0	0	0	0	0	21	0
N.S.	1	1.00	1.11	0.00	0.00	0.00	0.00	0.00	0.32	0.00
time (sec)	N/A	0.235	0.048	0.000	0.000	0.000	0.000	0.000	0.184	0.000

Problem 224	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	73	0	0	0	0	0	19	0
N.S.	1	1.00	1.09	0.00	0.00	0.00	0.00	0.00	0.28	0.00
time (sec)	N/A	0.224	0.042	0.000	0.000	0.000	0.000	0.000	0.186	0.000

Problem 225	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	14	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.24	0.00
time (sec)	N/A	0.188	0.088	0.000	0.000	0.000	0.000	0.000	0.180	0.000

Problem 226	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	39	35	0	141	0	0	19	0
N.S.	1	1.00	1.08	0.97	0.00	3.92	0.00	0.00	0.53	0.00
time (sec)	N/A	0.248	0.075	0.676	0.000	0.091	0.000	0.000	0.183	0.000

Problem 227	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	66	0	0	0	0	0	23	0
N.S.	1	1.00	0.99	0.00	0.00	0.00	0.00	0.00	0.34	0.00
time (sec)	N/A	0.240	0.056	0.000	0.000	0.000	0.000	0.000	0.178	0.000

Problem 228	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	70	71	73	0	0	0	0	0	27	0
N.S.	1	1.01	1.04	0.00	0.00	0.00	0.00	0.00	0.39	0.00
time (sec)	N/A	0.232	0.039	0.000	0.000	0.000	0.000	0.000	0.176	0.000

Problem 229	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	73	0	0	0	0	0	27	0
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	0.38	0.00
time (sec)	N/A	0.232	0.037	0.000	0.000	0.000	0.000	0.000	0.177	0.000

Problem 230	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	16	14	155	32	14	16	39	16
N.S.	1	1.00	1.14	1.00	11.07	2.29	1.00	1.14	2.79	1.14
time (sec)	N/A	0.194	12.172	0.037	1.292	0.088	30.205	0.408	0.193	3.460

Problem 231	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	143	28	12	14	32	14
N.S.	1	1.00	1.17	1.00	11.92	2.33	1.00	1.17	2.67	1.17
time (sec)	N/A	0.175	1.545	0.033	0.399	0.078	18.779	0.243	0.187	3.310

Problem 232	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	148	163	181	750	0	0	0	0	41	0
N.S.	1	1.10	1.22	5.07	0.00	0.00	0.00	0.00	0.28	0.00
time (sec)	N/A	0.847	0.318	13.485	0.000	0.000	0.000	0.000	0.179	0.000

Problem 233	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	154	32	15	18	46	18
N.S.	1	1.00	1.12	1.00	9.62	2.00	0.94	1.12	2.88	1.12
time (sec)	N/A	0.193	12.206	0.041	0.506	0.077	19.734	0.405	0.183	3.387

Problem 234	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	153	32	15	18	52	18
N.S.	1	1.00	1.12	1.00	9.56	2.00	0.94	1.12	3.25	1.12
time (sec)	N/A	0.200	12.120	0.034	0.498	0.079	43.166	0.387	0.180	3.379

Problem 235	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	B	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	28	33	29	147	129	0	0	12	0
N.S.	1	0.93	1.10	0.97	4.90	4.30	0.00	0.00	0.40	0.00
time (sec)	N/A	0.231	0.030	0.612	0.098	0.090	0.000	0.000	0.177	0.000

Problem 236	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	28	22	61	104	0	0	0	12	0
N.S.	1	1.17	0.92	2.54	4.33	0.00	0.00	0.00	0.50	0.00
time (sec)	N/A	0.228	0.011	0.345	0.036	0.000	0.000	0.000	0.179	0.000

Problem 237	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	17	16	15	22	15	101	15	15
N.S.	1	1.00	0.89	0.84	0.79	1.16	0.79	5.32	0.79	0.79
time (sec)	N/A	0.178	0.002	0.327	0.033	0.081	0.093	0.114	0.184	0.070

Problem 238	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	468	65	0	20	133	20
N.S.	1	1.00	1.11	1.00	26.00	3.61	0.00	1.11	7.39	1.11
time (sec)	N/A	0.205	6.758	0.049	1.489	0.102	0.000	0.774	0.200	3.616

Problem 239	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	265	44	0	20	88	20
N.S.	1	1.00	1.11	1.00	14.72	2.44	0.00	1.11	4.89	1.11
time (sec)	N/A	0.204	14.298	0.037	0.876	0.095	0.000	0.454	0.191	3.608

Problem 240	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	84	90	77	0	0	0	0	0	45	0
N.S.	1	1.07	0.92	0.00	0.00	0.00	0.00	0.00	0.54	0.00
time (sec)	N/A	0.308	0.077	0.000	0.000	0.000	0.000	0.000	0.179	0.000

Problem 241	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	20	20	15	20	22	20
N.S.	1	1.00	1.11	1.00	1.11	1.11	0.83	1.11	1.22	1.11
time (sec)	N/A	0.200	0.343	0.036	0.090	0.075	36.479	0.241	0.177	3.362

Problem 242	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	F(-1)	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	0	169	36	0	20	38	20
N.S.	1	1.00	1.11	0.00	9.39	2.00	0.00	1.11	2.11	1.11
time (sec)	N/A	0.204	1.757	180.000	0.238	0.078	0.000	0.389	0.194	3.470

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [201] had the largest ratio of [1.2222200000000008]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	3	3	0.97	12	0.250
2	A	5	4	1.00	12	0.333
3	A	3	3	0.98	12	0.250
4	A	5	4	1.02	12	0.333
5	A	3	3	1.00	10	0.300
6	A	1	1	1.00	8	0.125
7	A	1	1	1.00	12	0.083
8	A	6	5	1.06	12	0.417
9	A	3	3	0.92	12	0.250
10	A	5	4	0.96	12	0.333
11	A	4	4	0.96	12	0.333
12	A	5	4	0.95	12	0.333
13	A	15	14	1.41	14	1.000
14	A	14	13	1.23	14	0.929
15	A	10	9	1.21	14	0.643
16	A	10	9	1.09	14	0.643
17	A	4	4	1.04	12	0.333
18	A	6	5	1.19	10	0.500
19	A	4	4	1.22	14	0.286
20	A	4	4	1.08	14	0.286
21	A	9	8	0.96	14	0.571

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	8	8	0.92	14	0.571
23	A	14	13	1.16	14	0.929
24	B	18	17	2.02	14	1.214
25	A	16	15	1.39	14	1.071
26	A	14	13	1.57	14	0.929
27	A	10	10	1.10	14	0.714
28	A	9	8	1.12	12	0.667
29	A	5	5	1.14	10	0.500
30	A	5	5	1.15	14	0.357
31	A	5	5	1.13	14	0.357
32	A	7	7	0.97	14	0.500
33	A	15	14	0.98	14	1.000
34	A	11	11	1.29	14	0.786
35	A	8	7	1.15	16	0.438
36	A	8	7	1.16	16	0.438
37	A	7	6	1.12	16	0.375
38	A	7	6	1.08	16	0.375
39	A	6	5	1.12	16	0.312
40	A	8	7	1.12	16	0.438
41	A	7	6	1.14	16	0.375
42	A	9	8	1.14	16	0.500
43	N/A	1	0	1.00	16	0.000
44	N/A	1	0	1.00	16	0.000
45	A	2	2	1.00	14	0.143
46	N/A	1	0	1.00	16	0.000
47	N/A	1	0	1.00	16	0.000
48	N/A	1	0	1.00	10	0.000
49	N/A	1	0	1.00	16	0.000
50	A	5	4	0.98	14	0.286
51	A	5	4	1.02	14	0.286
52	A	5	4	1.00	14	0.286
53	A	2	2	0.88	12	0.167

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	3	2	1.20	14	0.143
55	A	6	5	1.05	14	0.357
56	A	5	4	0.93	14	0.286
57	A	5	4	0.96	14	0.286
58	A	5	5	1.14	14	0.357
59	A	5	5	1.11	14	0.357
60	A	1	1	1.00	10	0.100
61	A	4	4	1.20	14	0.286
62	A	5	5	1.10	14	0.357
63	A	5	5	1.13	14	0.357
64	A	11	10	1.22	16	0.625
65	A	11	10	1.11	16	0.625
66	A	6	5	1.01	16	0.312
67	A	7	6	1.11	14	0.429
68	A	6	5	1.19	16	0.312
69	A	6	5	1.05	16	0.312
70	A	10	9	0.99	16	0.562
71	A	2	2	1.00	16	0.125
72	A	2	2	1.00	16	0.125
73	A	2	2	1.00	12	0.167
74	A	2	2	1.00	16	0.125
75	A	2	2	1.00	16	0.125
76	A	2	2	1.00	16	0.125
77	A	10	9	1.12	16	0.562
78	A	7	6	1.07	14	0.429
79	A	7	6	1.16	16	0.375
80	A	7	6	1.06	16	0.375
81	A	9	8	0.99	16	0.500
82	A	17	16	1.39	18	0.889
83	A	16	15	1.36	18	0.833
84	A	16	15	1.36	18	0.833
85	A	16	15	1.41	18	0.833

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
86	A	16	15	1.42	18	0.833
87	A	15	14	1.35	18	0.778
88	A	17	16	1.37	18	0.889
89	A	17	16	1.36	18	0.889
90	A	5	4	0.85	20	0.200
91	A	5	4	0.85	20	0.200
92	A	5	4	0.82	20	0.200
93	A	5	4	0.82	20	0.200
94	N/A	1	0	1.00	18	0.000
95	N/A	1	0	1.00	18	0.000
96	A	2	2	1.00	16	0.125
97	N/A	1	0	1.00	18	0.000
98	N/A	1	0	1.00	18	0.000
99	A	5	4	0.98	14	0.286
100	A	5	4	1.02	14	0.286
101	A	5	4	1.00	14	0.286
102	A	2	2	0.88	14	0.143
103	A	3	2	1.20	14	0.143
104	A	6	5	1.05	14	0.357
105	A	5	4	0.93	14	0.286
106	A	5	4	0.96	14	0.286
107	A	12	11	1.26	14	0.786
108	A	1	1	1.00	10	0.100
109	A	11	10	1.27	14	0.714
110	A	11	10	1.10	14	0.714
111	A	12	11	1.35	14	0.786
112	A	11	10	1.07	14	0.714
113	A	11	10	1.34	12	0.833
114	A	10	9	1.06	14	0.643
115	A	12	11	1.33	14	0.786
116	A	11	10	1.22	16	0.625
117	A	11	10	1.11	16	0.625

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
118	A	6	5	1.01	16	0.312
119	A	7	6	1.08	16	0.375
120	A	6	5	1.16	16	0.312
121	A	6	5	1.01	16	0.312
122	A	10	9	0.99	16	0.562
123	A	10	9	0.95	16	0.562
124	A	12	11	1.06	16	0.688
125	A	10	9	1.14	16	0.562
126	A	7	6	1.10	16	0.375
127	A	7	6	1.14	16	0.375
128	A	7	6	1.11	16	0.375
129	A	9	8	1.01	16	0.500
130	N/A	1	0	1.00	18	0.000
131	N/A	1	0	1.00	18	0.000
132	A	2	2	1.00	16	0.125
133	N/A	1	0	1.00	18	0.000
134	N/A	1	0	1.00	18	0.000
135	A	4	4	0.96	14	0.286
136	A	7	6	0.93	14	0.429
137	A	4	4	0.90	12	0.333
138	A	1	1	1.00	10	0.100
139	A	3	2	1.10	14	0.143
140	A	2	2	1.00	14	0.143
141	A	4	4	1.00	14	0.286
142	A	7	6	1.15	14	0.429
143	A	15	14	1.14	16	0.875
144	A	10	9	0.92	16	0.562
145	A	10	9	0.99	14	0.643
146	A	10	9	1.08	12	0.750
147	A	6	5	1.20	16	0.312
148	A	7	6	1.16	16	0.375
149	A	6	5	1.00	16	0.312

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
150	A	13	12	1.31	16	0.750
151	A	16	15	1.00	16	0.938
152	A	9	8	0.97	14	0.571
153	A	9	9	1.12	12	0.750
154	A	7	6	1.13	16	0.375
155	A	7	6	1.11	16	0.375
156	A	10	9	1.11	16	0.562
157	A	7	6	0.96	14	0.429
158	A	7	6	0.93	14	0.429
159	A	7	6	0.91	14	0.429
160	A	3	3	0.87	12	0.250
161	A	3	2	1.23	14	0.143
162	A	2	2	1.00	14	0.143
163	A	7	6	1.00	14	0.429
164	A	7	6	1.15	14	0.429
165	A	7	7	1.14	14	0.500
166	A	6	6	1.07	14	0.429
167	A	1	1	1.00	10	0.100
168	A	5	5	1.20	14	0.357
169	A	7	7	1.11	14	0.500
170	A	6	6	1.14	14	0.429
171	A	10	9	0.94	16	0.562
172	A	6	5	0.96	14	0.357
173	A	6	5	1.13	16	0.312
174	A	7	6	1.06	16	0.375
175	A	6	5	0.94	16	0.312
176	A	3	3	1.00	16	0.188
177	A	3	3	1.00	16	0.188
178	A	3	3	1.00	12	0.250
179	A	3	3	1.03	16	0.188
180	A	3	3	1.00	16	0.188
181	N/A	1	0	1.00	18	0.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
182	N/A	1	0	1.00	18	0.000
183	A	4	3	1.04	16	0.188
184	N/A	1	0	1.00	18	0.000
185	N/A	1	0	1.00	18	0.000
186	A	8	7	1.15	16	0.438
187	A	7	6	1.12	16	0.375
188	A	6	5	1.08	14	0.357
189	A	1	1	1.00	12	0.083
190	A	3	2	1.38	16	0.125
191	A	5	4	1.08	16	0.250
192	A	6	5	0.98	16	0.312
193	A	7	6	1.00	16	0.375
194	A	21	20	1.45	18	1.111
195	A	16	15	1.31	18	0.833
196	A	11	10	1.23	16	0.625
197	A	6	5	1.13	14	0.357
198	A	6	5	1.22	18	0.278
199	A	10	9	1.05	18	0.500
200	A	15	14	1.14	18	0.778
201	B	23	22	2.54	18	1.222
202	B	19	18	2.01	18	1.000
203	A	15	14	1.52	16	0.875
204	A	10	9	1.20	14	0.643
205	A	7	6	1.17	18	0.333
206	A	9	8	1.06	18	0.444
207	A	13	12	1.31	18	0.667
208	A	3	3	0.92	12	0.250
209	A	3	3	0.90	12	0.250
210	A	2	2	1.00	12	0.167
211	A	4	4	1.00	12	0.333
212	A	13	12	1.32	16	0.750
213	A	6	5	1.10	16	0.312

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
214	A	13	12	1.25	14	0.857
215	A	1	1	1.21	12	0.083
216	A	3	2	1.24	16	0.125
217	A	12	11	1.27	16	0.688
218	A	13	12	1.31	16	0.750
219	A	6	5	1.04	16	0.312
220	A	6	5	1.01	18	0.278
221	A	6	5	1.15	18	0.278
222	A	10	9	0.99	18	0.500
223	A	2	2	1.00	14	0.143
224	A	2	2	1.00	12	0.167
225	A	1	1	1.00	10	0.100
226	A	3	2	1.00	14	0.143
227	A	2	2	1.00	14	0.143
228	A	2	2	1.01	14	0.143
229	A	2	2	1.00	14	0.143
230	N/A	1	0	1.00	14	0.000
231	N/A	1	0	1.00	12	0.000
232	A	6	5	1.10	16	0.312
233	N/A	1	0	1.00	16	0.000
234	N/A	1	0	1.00	16	0.000
235	A	3	2	0.93	10	0.200
236	A	3	2	1.17	10	0.200
237	A	3	3	1.00	4	0.750
238	N/A	1	0	1.00	18	0.000
239	N/A	1	0	1.00	18	0.000
240	A	3	3	1.07	16	0.188
241	N/A	1	0	1.00	18	0.000
242	N/A	1	0	1.00	18	0.000

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^5(a + \operatorname{barctanh}(cx)) dx$	114
3.2	$\int x^4(a + \operatorname{barctanh}(cx)) dx$	120
3.3	$\int x^3(a + \operatorname{barctanh}(cx)) dx$	126
3.4	$\int x^2(a + \operatorname{barctanh}(cx)) dx$	132
3.5	$\int x(a + \operatorname{barctanh}(cx)) dx$	138
3.6	$\int (a + \operatorname{barctanh}(cx)) dx$	144
3.7	$\int \frac{a+\operatorname{barctanh}(cx)}{x} dx$	149
3.8	$\int \frac{a+\operatorname{barctanh}(cx)}{x^2} dx$	154
3.9	$\int \frac{a+\operatorname{barctanh}(cx)}{x^3} dx$	160
3.10	$\int \frac{a+\operatorname{barctanh}(cx)}{x^4} dx$	166
3.11	$\int \frac{a+\operatorname{barctanh}(cx)}{x^5} dx$	172
3.12	$\int \frac{a+\operatorname{barctanh}(cx)}{x^6} dx$	178
3.13	$\int x^5(a + \operatorname{barctanh}(cx))^2 dx$	184
3.14	$\int x^4(a + \operatorname{barctanh}(cx))^2 dx$	194
3.15	$\int x^3(a + \operatorname{barctanh}(cx))^2 dx$	203
3.16	$\int x^2(a + \operatorname{barctanh}(cx))^2 dx$	211
3.17	$\int x(a + \operatorname{barctanh}(cx))^2 dx$	219
3.18	$\int (a + \operatorname{barctanh}(cx))^2 dx$	226
3.19	$\int \frac{(a+\operatorname{barctanh}(cx))^2}{x} dx$	232
3.20	$\int \frac{(a+\operatorname{barctanh}(cx))^2}{x^2} dx$	239
3.21	$\int \frac{(a+\operatorname{barctanh}(cx))^2}{x^3} dx$	245
3.22	$\int \frac{(a+\operatorname{barctanh}(cx))^2}{x^4} dx$	253
3.23	$\int \frac{(a+\operatorname{barctanh}(cx))^2}{x^5} dx$	260
3.24	$\int x^5(a + \operatorname{barctanh}(cx))^3 dx$	270

3.25	$\int x^4(a + \operatorname{barctanh}(cx))^3 dx$	283
3.26	$\int x^3(a + \operatorname{barctanh}(cx))^3 dx$	294
3.27	$\int x^2(a + \operatorname{barctanh}(cx))^3 dx$	304
3.28	$\int x(a + \operatorname{barctanh}(cx))^3 dx$	313
3.29	$\int (a + \operatorname{barctanh}(cx))^3 dx$	321
3.30	$\int \frac{(a+\operatorname{barctanh}(cx))^3}{x} dx$	328
3.31	$\int \frac{(a+\operatorname{barctanh}(cx))^3}{x^2} dx$	336
3.32	$\int \frac{(a+\operatorname{barctanh}(cx))^3}{x^3} dx$	343
3.33	$\int \frac{(a+\operatorname{barctanh}(cx))^3}{x^4} dx$	351
3.34	$\int \frac{(a+\operatorname{barctanh}(cx))^3}{x^5} dx$	361
3.35	$\int (dx)^{5/2}(a + \operatorname{barctanh}(cx)) dx$	370
3.36	$\int (dx)^{3/2}(a + \operatorname{barctanh}(cx)) dx$	378
3.37	$\int \sqrt{dx}(a + \operatorname{barctanh}(cx)) dx$	385
3.38	$\int \frac{a+\operatorname{barctanh}(cx)}{\sqrt{dx}} dx$	392
3.39	$\int \frac{a+\operatorname{barctanh}(cx)}{(dx)^{3/2}} dx$	399
3.40	$\int \frac{a+\operatorname{barctanh}(cx)}{(dx)^{5/2}} dx$	406
3.41	$\int \frac{a+\operatorname{barctanh}(cx)}{(dx)^{7/2}} dx$	413
3.42	$\int \frac{a+\operatorname{barctanh}(cx)}{(dx)^{9/2}} dx$	420
3.43	$\int (dx)^m(a + \operatorname{barctanh}(cx))^3 dx$	428
3.44	$\int (dx)^m(a + \operatorname{barctanh}(cx))^2 dx$	433
3.45	$\int (dx)^m(a + \operatorname{barctanh}(cx)) dx$	438
3.46	$\int \frac{(dx)^m}{a+\operatorname{barctanh}(cx)} dx$	443
3.47	$\int \frac{(dx)^m}{(a+\operatorname{barctanh}(cx))^2} dx$	448
3.48	$\int (a + \operatorname{barctanh}(cx))^p dx$	453
3.49	$\int (dx)^m(a + \operatorname{barctanh}(cx))^p dx$	458
3.50	$\int x^7(a + \operatorname{barctanh}(cx^2)) dx$	463
3.51	$\int x^5(a + \operatorname{barctanh}(cx^2)) dx$	469
3.52	$\int x^3(a + \operatorname{barctanh}(cx^2)) dx$	475
3.53	$\int x(a + \operatorname{barctanh}(cx^2)) dx$	481
3.54	$\int \frac{a+\operatorname{barctanh}(cx^2)}{x} dx$	487
3.55	$\int \frac{a+\operatorname{barctanh}(cx^2)}{x^3} dx$	492
3.56	$\int \frac{a+\operatorname{barctanh}(cx^2)}{x^5} dx$	498
3.57	$\int \frac{a+\operatorname{barctanh}(cx^2)}{x^7} dx$	504
3.58	$\int x^4(a + \operatorname{barctanh}(cx^2)) dx$	510
3.59	$\int x^2(a + \operatorname{barctanh}(cx^2)) dx$	517

3.60	$\int (a + \operatorname{barctanh}(cx^2)) dx$	524
3.61	$\int \frac{a + \operatorname{barctanh}(cx^2)}{x^2} dx$	530
3.62	$\int \frac{a + \operatorname{barctanh}(cx^2)}{x^4} dx$	537
3.63	$\int \frac{a + \operatorname{barctanh}(cx^2)}{x^6} dx$	544
3.64	$\int x^7 (a + \operatorname{barctanh}(cx^2))^2 dx$	551
3.65	$\int x^5 (a + \operatorname{barctanh}(cx^2))^2 dx$	560
3.66	$\int x^3 (a + \operatorname{barctanh}(cx^2))^2 dx$	568
3.67	$\int x (a + \operatorname{barctanh}(cx^2))^2 dx$	576
3.68	$\int \frac{(a + \operatorname{barctanh}(cx^2))^2}{x} dx$	583
3.69	$\int \frac{(a + \operatorname{barctanh}(cx^2))^2}{x^3} dx$	590
3.70	$\int \frac{(a + \operatorname{barctanh}(cx^2))^2}{x^5} dx$	597
3.71	$\int x^4 (a + \operatorname{barctanh}(cx^2))^2 dx$	606
3.72	$\int x^2 (a + \operatorname{barctanh}(cx^2))^2 dx$	614
3.73	$\int (a + \operatorname{barctanh}(cx^2))^2 dx$	622
3.74	$\int \frac{(a + \operatorname{barctanh}(cx^2))^2}{x^2} dx$	629
3.75	$\int \frac{(a + \operatorname{barctanh}(cx^2))^2}{x^4} dx$	637
3.76	$\int \frac{(a + \operatorname{barctanh}(cx^2))^2}{x^6} dx$	645
3.77	$\int x^3 (a + \operatorname{barctanh}(cx^2))^3 dx$	653
3.78	$\int x (a + \operatorname{barctanh}(cx^2))^3 dx$	662
3.79	$\int \frac{(a + \operatorname{barctanh}(cx^2))^3}{x} dx$	669
3.80	$\int \frac{(a + \operatorname{barctanh}(cx^2))^3}{x^3} dx$	676
3.81	$\int \frac{(a + \operatorname{barctanh}(cx^2))^3}{x^5} dx$	683
3.82	$\int (dx)^{5/2} (a + \operatorname{barctanh}(cx^2)) dx$	690
3.83	$\int (dx)^{3/2} (a + \operatorname{barctanh}(cx^2)) dx$	705
3.84	$\int \sqrt{dx} (a + \operatorname{barctanh}(cx^2)) dx$	719
3.85	$\int \frac{a + \operatorname{barctanh}(cx^2)}{\sqrt{dx}} dx$	733
3.86	$\int \frac{a + \operatorname{barctanh}(cx^2)}{(dx)^{3/2}} dx$	745
3.87	$\int \frac{a + \operatorname{barctanh}(cx^2)}{(dx)^{5/2}} dx$	758
3.88	$\int \frac{a + \operatorname{barctanh}(cx^2)}{(dx)^{7/2}} dx$	771
3.89	$\int \frac{a + \operatorname{barctanh}(cx^2)}{(dx)^{9/2}} dx$	787
3.90	$\int \sqrt{dx} (a + \operatorname{barctanh}(cx^2))^2 dx$	803

3.91	$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{\sqrt{dx}} dx$	810
3.92	$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{3/2}} dx$	817
3.93	$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{5/2}} dx$	824
3.94	$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^3 dx$	831
3.95	$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^2 dx$	836
3.96	$\int (dx)^m (a + b \operatorname{arctanh}(cx^2)) dx$	841
3.97	$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^2)} dx$	846
3.98	$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^2))^2} dx$	851
3.99	$\int x^{11} (a + b \operatorname{arctanh}(cx^3)) dx$	856
3.100	$\int x^8 (a + b \operatorname{arctanh}(cx^3)) dx$	862
3.101	$\int x^5 (a + b \operatorname{arctanh}(cx^3)) dx$	868
3.102	$\int x^2 (a + b \operatorname{arctanh}(cx^3)) dx$	874
3.103	$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x} dx$	879
3.104	$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^4} dx$	884
3.105	$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^7} dx$	890
3.106	$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^{10}} dx$	896
3.107	$\int x^3 (a + b \operatorname{arctanh}(cx^3)) dx$	902
3.108	$\int (a + b \operatorname{arctanh}(cx^3)) dx$	913
3.109	$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^3} dx$	919
3.110	$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^6} dx$	929
3.111	$\int x^7 (a + b \operatorname{arctanh}(cx^3)) dx$	938
3.112	$\int x^4 (a + b \operatorname{arctanh}(cx^3)) dx$	950
3.113	$\int x (a + b \operatorname{arctanh}(cx^3)) dx$	959
3.114	$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^2} dx$	970
3.115	$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^5} dx$	979
3.116	$\int x^{11} (a + b \operatorname{arctanh}(cx^3))^2 dx$	990
3.117	$\int x^8 (a + b \operatorname{arctanh}(cx^3))^2 dx$	999
3.118	$\int x^5 (a + b \operatorname{arctanh}(cx^3))^2 dx$	1008
3.119	$\int x^2 (a + b \operatorname{arctanh}(cx^3))^2 dx$	1016
3.120	$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x} dx$	1023
3.121	$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^4} dx$	1030
3.122	$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^7} dx$	1036

3.123	$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^{10}} dx$	1044
3.124	$\int x^8 (a + b \operatorname{arctanh}(cx^3))^3 dx$	1052
3.125	$\int x^5 (a + b \operatorname{arctanh}(cx^3))^3 dx$	1060
3.126	$\int x^2 (a + b \operatorname{arctanh}(cx^3))^3 dx$	1069
3.127	$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x} dx$	1076
3.128	$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^4} dx$	1083
3.129	$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^7} dx$	1090
3.130	$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^3 dx$	1097
3.131	$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^2 dx$	1102
3.132	$\int (dx)^m (a + b \operatorname{arctanh}(cx^3)) dx$	1107
3.133	$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^3)} dx$	1112
3.134	$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^3))^2} dx$	1117
3.135	$\int x^3 (a + b \operatorname{arctanh}(\frac{c}{x})) dx$	1122
3.136	$\int x^2 (a + b \operatorname{arctanh}(\frac{c}{x})) dx$	1128
3.137	$\int x (a + b \operatorname{arctanh}(\frac{c}{x})) dx$	1134
3.138	$\int (a + b \operatorname{arctanh}(\frac{c}{x})) dx$	1140
3.139	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x})}{x} dx$	1145
3.140	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x})}{x^2} dx$	1150
3.141	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x})}{x^3} dx$	1155
3.142	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x})}{x^4} dx$	1161
3.143	$\int x^3 (a + b \operatorname{arctanh}(\frac{c}{x}))^2 dx$	1168
3.144	$\int x^2 (a + b \operatorname{arctanh}(\frac{c}{x}))^2 dx$	1178
3.145	$\int x (a + b \operatorname{arctanh}(\frac{c}{x}))^2 dx$	1186
3.146	$\int (a + b \operatorname{arctanh}(\frac{c}{x}))^2 dx$	1194
3.147	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x} dx$	1201
3.148	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^2} dx$	1209
3.149	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^3} dx$	1216
3.150	$\int x^3 (a + b \operatorname{arctanh}(\frac{c}{x}))^3 dx$	1224
3.151	$\int x^2 (a + b \operatorname{arctanh}(\frac{c}{x}))^3 dx$	1234
3.152	$\int x (a + b \operatorname{arctanh}(\frac{c}{x}))^3 dx$	1245
3.153	$\int (a + b \operatorname{arctanh}(\frac{c}{x}))^3 dx$	1254
3.154	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x} dx$	1262

3.155	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^2} dx$	1271
3.156	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^3} dx$	1279
3.157	$\int x^7 (a + b \operatorname{arctanh}(\frac{c}{x^2})) dx$	1288
3.158	$\int x^5 (a + b \operatorname{arctanh}(\frac{c}{x^2})) dx$	1295
3.159	$\int x^3 (a + b \operatorname{arctanh}(\frac{c}{x^2})) dx$	1302
3.160	$\int x (a + b \operatorname{arctanh}(\frac{c}{x^2})) dx$	1308
3.161	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{x} dx$	1314
3.162	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{x^3} dx$	1319
3.163	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{x^5} dx$	1325
3.164	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{x^7} dx$	1332
3.165	$\int x^4 (a + b \operatorname{arctanh}(\frac{c}{x^2})) dx$	1339
3.166	$\int x^2 (a + b \operatorname{arctanh}(\frac{c}{x^2})) dx$	1347
3.167	$\int (a + b \operatorname{arctanh}(\frac{c}{x^2})) dx$	1355
3.168	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{x^2} dx$	1361
3.169	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{x^4} dx$	1368
3.170	$\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{x^6} dx$	1376
3.171	$\int x^3 (a + b \operatorname{arctanh}(\frac{c}{x^2}))^2 dx$	1384
3.172	$\int x (a + b \operatorname{arctanh}(\frac{c}{x^2}))^2 dx$	1394
3.173	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x} dx$	1401
3.174	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^3} dx$	1408
3.175	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^5} dx$	1415
3.176	$\int x^4 (a + b \operatorname{arctanh}(\frac{c}{x^2}))^2 dx$	1422
3.177	$\int x^2 (a + b \operatorname{arctanh}(\frac{c}{x^2}))^2 dx$	1430
3.178	$\int (a + b \operatorname{arctanh}(\frac{c}{x^2}))^2 dx$	1438
3.179	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^2} dx$	1446
3.180	$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^4} dx$	1454
3.181	$\int (dx)^m (a + b \operatorname{arctanh}(\frac{c}{x^2}))^3 dx$	1462
3.182	$\int (dx)^m (a + b \operatorname{arctanh}(\frac{c}{x^2}))^2 dx$	1467
3.183	$\int (dx)^m (a + b \operatorname{arctanh}(\frac{c}{x^2})) dx$	1472
3.184	$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(\frac{c}{x^2})} dx$	1477

3.185	$\int \frac{(dx)^m}{(a+b\operatorname{arctanh}(\frac{c}{x^2}))^2} dx$	1482
3.186	$\int x^3(a+b\operatorname{arctanh}(c\sqrt{x})) dx$	1487
3.187	$\int x^2(a+b\operatorname{arctanh}(c\sqrt{x})) dx$	1494
3.188	$\int x(a+b\operatorname{arctanh}(c\sqrt{x})) dx$	1501
3.189	$\int (a+b\operatorname{arctanh}(c\sqrt{x})) dx$	1508
3.190	$\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x} dx$	1513
3.191	$\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x^2} dx$	1518
3.192	$\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x^3} dx$	1524
3.193	$\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x^4} dx$	1531
3.194	$\int x^3(a+b\operatorname{arctanh}(c\sqrt{x}))^2 dx$	1538
3.195	$\int x^2(a+b\operatorname{arctanh}(c\sqrt{x}))^2 dx$	1550
3.196	$\int x(a+b\operatorname{arctanh}(c\sqrt{x}))^2 dx$	1559
3.197	$\int (a+b\operatorname{arctanh}(c\sqrt{x}))^2 dx$	1567
3.198	$\int \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{x} dx$	1574
3.199	$\int \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{x^2} dx$	1582
3.200	$\int \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{x^3} dx$	1591
3.201	$\int x^3(a+b\operatorname{arctanh}(c\sqrt{x}))^3 dx$	1602
3.202	$\int x^2(a+b\operatorname{arctanh}(c\sqrt{x}))^3 dx$	1617
3.203	$\int x(a+b\operatorname{arctanh}(c\sqrt{x}))^3 dx$	1631
3.204	$\int (a+b\operatorname{arctanh}(c\sqrt{x}))^3 dx$	1642
3.205	$\int \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^3}{x} dx$	1651
3.206	$\int \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^3}{x^2} dx$	1661
3.207	$\int \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^3}{x^3} dx$	1670
3.208	$\int x^{3/2}\operatorname{arctanh}(\sqrt{x}) dx$	1680
3.209	$\int \sqrt{x}\operatorname{arctanh}(\sqrt{x}) dx$	1686
3.210	$\int \frac{\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} dx$	1691
3.211	$\int \frac{\operatorname{arctanh}(\sqrt{x})}{x^{3/2}} dx$	1696
3.212	$\int x^3(a+b\operatorname{arctanh}(cx^{3/2})) dx$	1702
3.213	$\int x^2(a+b\operatorname{arctanh}(cx^{3/2})) dx$	1714
3.214	$\int x(a+b\operatorname{arctanh}(cx^{3/2})) dx$	1720
3.215	$\int (a+b\operatorname{arctanh}(cx^{3/2})) dx$	1731
3.216	$\int \frac{a+b\operatorname{arctanh}(cx^{3/2})}{x} dx$	1738

3.217	$\int \frac{a + \operatorname{barctanh}(cx^{3/2})}{x^2} dx$	1743
3.218	$\int \frac{a + \operatorname{barctanh}(cx^{3/2})}{x^3} dx$	1753
3.219	$\int \frac{a + \operatorname{barctanh}(cx^{3/2})}{x^4} dx$	1764
3.220	$\int x^2 (a + \operatorname{barctanh}(cx^{3/2}))^2 dx$	1770
3.221	$\int \frac{(a + \operatorname{barctanh}(cx^{3/2}))^2}{x} dx$	1777
3.222	$\int \frac{(a + \operatorname{barctanh}(cx^{3/2}))^2}{x^4} dx$	1785
3.223	$\int x^2 (a + \operatorname{barctanh}(cx^n)) dx$	1794
3.224	$\int x (a + \operatorname{barctanh}(cx^n)) dx$	1799
3.225	$\int (a + \operatorname{barctanh}(cx^n)) dx$	1804
3.226	$\int \frac{a + \operatorname{barctanh}(cx^n)}{x} dx$	1809
3.227	$\int \frac{a + \operatorname{barctanh}(cx^n)}{x^2} dx$	1814
3.228	$\int \frac{a + \operatorname{barctanh}(cx^n)}{x^3} dx$	1819
3.229	$\int \frac{a + \operatorname{barctanh}(cx^n)}{x^4} dx$	1824
3.230	$\int x (a + \operatorname{barctanh}(cx^n))^2 dx$	1829
3.231	$\int (a + \operatorname{barctanh}(cx^n))^2 dx$	1834
3.232	$\int \frac{(a + \operatorname{barctanh}(cx^n))^2}{x} dx$	1839
3.233	$\int \frac{(a + \operatorname{barctanh}(cx^n))^2}{x^2} dx$	1846
3.234	$\int \frac{(a + \operatorname{barctanh}(cx^n))^2}{x^3} dx$	1851
3.235	$\int \frac{\operatorname{arctanh}(ax^n)}{x} dx$	1856
3.236	$\int \frac{\operatorname{arctanh}(ax^5)}{x} dx$	1861
3.237	$\int \operatorname{arctanh}\left(\frac{1}{x}\right) dx$	1866
3.238	$\int (dx)^m (a + \operatorname{barctanh}(cx^n))^3 dx$	1871
3.239	$\int (dx)^m (a + \operatorname{barctanh}(cx^n))^2 dx$	1876
3.240	$\int (dx)^m (a + \operatorname{barctanh}(cx^n)) dx$	1881
3.241	$\int \frac{(dx)^m}{a + \operatorname{barctanh}(cx^n)} dx$	1886
3.242	$\int \frac{(dx)^m}{(a + \operatorname{barctanh}(cx^n))^2} dx$	1891

3.1 $\int x^5(a + \operatorname{barctanh}(cx)) dx$

Optimal result	114
Mathematica [A] (verified)	114
Rubi [A] (verified)	115
Maple [A] (verified)	116
Fricas [A] (verification not implemented)	116
Sympy [A] (verification not implemented)	117
Maxima [A] (verification not implemented)	117
Giac [B] (verification not implemented)	118
Mupad [B] (verification not implemented)	118
Reduce [B] (verification not implemented)	119

Optimal result

Integrand size = 12, antiderivative size = 59

$$\int x^5(a + \operatorname{barctanh}(cx)) dx = \frac{bx}{6c^5} + \frac{bx^3}{18c^3} + \frac{bx^5}{30c} - \frac{\operatorname{barctanh}(cx)}{6c^6} + \frac{1}{6}x^6(a + \operatorname{barctanh}(cx))$$

output

```
1/6*b*x/c^5+1/18*b*x^3/c^3+1/30*b*x^5/c-1/6*b*arctanh(c*x)/c^6+1/6*x^6*(a+
b*arctanh(c*x))
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.37

$$\int x^5(a + \operatorname{barctanh}(cx)) dx = \frac{bx}{6c^5} + \frac{bx^3}{18c^3} + \frac{bx^5}{30c} + \frac{ax^6}{6} + \frac{1}{6}bx^6\operatorname{arctanh}(cx) + \frac{b \log(1 - cx)}{12c^6} - \frac{b \log(1 + cx)}{12c^6}$$

input

```
Integrate[x^5*(a + b*ArcTanh[c*x]),x]
```

output

```
(b*x)/(6*c^5) + (b*x^3)/(18*c^3) + (b*x^5)/(30*c) + (a*x^6)/6 + (b*x^6*Arc
Tanh[c*x])/6 + (b*Log[1 - c*x])/(12*c^6) - (b*Log[1 + c*x])/(12*c^6)
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.97, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6452, 254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^5(a + b \operatorname{arctanh}(cx)) dx$$

$$\downarrow 6452$$

$$\frac{1}{6}x^6(a + b \operatorname{arctanh}(cx)) - \frac{1}{6}bc \int \frac{x^6}{1 - c^2x^2} dx$$

$$\downarrow 254$$

$$\frac{1}{6}x^6(a + b \operatorname{arctanh}(cx)) - \frac{1}{6}bc \int \left(-\frac{x^4}{c^2} - \frac{x^2}{c^4} + \frac{1}{c^6(1 - c^2x^2)} - \frac{1}{c^6} \right) dx$$

$$\downarrow 2009$$

$$\frac{1}{6}x^6(a + b \operatorname{arctanh}(cx)) - \frac{1}{6}bc \left(\frac{\operatorname{arctanh}(cx)}{c^7} - \frac{x}{c^6} - \frac{x^3}{3c^4} - \frac{x^5}{5c^2} \right)$$

input `Int[x^5*(a + b*ArcTanh[c*x]),x]`

output `(x^6*(a + b*ArcTanh[c*x]))/6 - (b*c*(-(x/c^6) - x^3/(3*c^4) - x^5/(5*c^2) + ArcTanh[c*x]/c^7))/6`

Defintions of rubi rules used

rule 254 `Int[(x_)^(m_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^2, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 3]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00

method	result
parallelrisc	$-\frac{-15b \operatorname{arctanh}(cx)x^6c^6 - 15ac^6x^6 - 3bc^5x^5 - 5b^2c^3x^3 - 15bcx + 15b \operatorname{arctanh}(cx)}{90c^6}$
parts	$\frac{ax^6}{6} + \frac{b \left(\frac{c^6x^6 \operatorname{arctanh}(cx)}{6} + \frac{c^5x^5}{30} + \frac{c^3x^3}{18} + \frac{cx}{6} + \frac{\ln(cx-1)}{12} - \frac{\ln(cx+1)}{12} \right)}{c^6}$
derivativdivides	$\frac{\frac{ac^6x^6}{6} + b \left(\frac{c^6x^6 \operatorname{arctanh}(cx)}{6} + \frac{c^5x^5}{30} + \frac{c^3x^3}{18} + \frac{cx}{6} + \frac{\ln(cx-1)}{12} - \frac{\ln(cx+1)}{12} \right)}{c^6}$
default	$\frac{\frac{ac^6x^6}{6} + b \left(\frac{c^6x^6 \operatorname{arctanh}(cx)}{6} + \frac{c^5x^5}{30} + \frac{c^3x^3}{18} + \frac{cx}{6} + \frac{\ln(cx-1)}{12} - \frac{\ln(cx+1)}{12} \right)}{c^6}$
risc	$\frac{bx^6 \ln(cx+1)}{12} - \frac{bx^6 \ln(-cx+1)}{12} + \frac{ax^6}{6} + \frac{bx^5}{30c} + \frac{bx^3}{18c^3} + \frac{bx}{6c^5} + \frac{b \ln(-cx+1)}{12c^6} - \frac{b \ln(cx+1)}{12c^6}$
orering	$\frac{(3c^6x^6 + c^4x^4 + 5c^2x^2 - 9)(a + b \operatorname{arctanh}(cx))}{9c^6} - \frac{(3c^4x^4 + 5c^2x^2 + 15)(cx-1)(cx+1) \left(5x^4(a + b \operatorname{arctanh}(cx)) + \frac{x^5bc}{-c^2x^2+1} \right)}{90x^4c^6}$

input

```
int(x^5*(a+b*arctanh(c*x)),x,method=_RETURNVERBOSE)
```

output

```
-1/90*(-15*b*arctanh(c*x)*x^6*c^6-15*a*c^6*x^6-3*b*c^5*x^5-5*b*c^3*x^3-15*
b*c*x+15*b*arctanh(c*x))/c^6
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.14

$$\int x^5(a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{30ac^6x^6 + 6bc^5x^5 + 10bc^3x^3 + 30bcx + 15(bc^6x^6 - b) \log\left(-\frac{cx+1}{cx-1}\right)}{180c^6}$$

input `integrate(x^5*(a+b*arctanh(c*x)),x, algorithm="fricas")`

output `1/180*(30*a*c^6*x^6 + 6*b*c^5*x^5 + 10*b*c^3*x^3 + 30*b*c*x + 15*(b*c^6*x^6 - b)*log(-(c*x + 1)/(c*x - 1)))/c^6`

Sympy [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.07

$$\int x^5(a + b \operatorname{arctanh}(cx)) dx$$

$$= \begin{cases} \frac{ax^6}{6} + \frac{bx^6 \operatorname{atanh}(cx)}{6} + \frac{bx^5}{30c} + \frac{bx^3}{18c^3} + \frac{bx}{6c^5} - \frac{b \operatorname{atanh}(cx)}{6c^6} & \text{for } c \neq 0 \\ \frac{ax^6}{6} & \text{otherwise} \end{cases}$$

input `integrate(x**5*(a+b*atanh(c*x)),x)`

output `Piecewise((a*x**6/6 + b*x**6*atanh(c*x)/6 + b*x**5/(30*c) + b*x**3/(18*c**3) + b*x/(6*c**5) - b*atanh(c*x)/(6*c**6), Ne(c, 0)), (a*x**6/6, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.19

$$\int x^5(a + b \operatorname{arctanh}(cx)) dx = \frac{1}{6} ax^6 + \frac{1}{180} \left(30 x^6 \operatorname{artanh}(cx) + c \left(\frac{2(3c^4x^5 + 5c^2x^3 + 15x)}{c^6} - \frac{15 \log(cx + 1)}{c^7} + \frac{15 \log(cx - 1)}{c^7} \right) \right) b$$

input `integrate(x^5*(a+b*arctanh(c*x)),x, algorithm="maxima")`

output `1/6*a*x^6 + 1/180*(30*x^6*arctanh(c*x) + c*(2*(3*c^4*x^5 + 5*c^2*x^3 + 15*x)/c^6 - 15*log(c*x + 1)/c^7 + 15*log(c*x - 1)/c^7))*b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 442 vs. $2(49) = 98$.

Time = 0.12 (sec) , antiderivative size = 442, normalized size of antiderivative = 7.49

$$\int x^5(a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{1}{45} c \left(\frac{15 \left(\frac{3(cx+1)^5 b}{(cx-1)^5} + \frac{10(cx+1)^3 b}{(cx-1)^3} + \frac{3(cx+1)b}{cx-1} \right) \log\left(-\frac{cx+1}{cx-1}\right)}{\frac{(cx+1)^6 c^7}{(cx-1)^6} - \frac{6(cx+1)^5 c^7}{(cx-1)^5} + \frac{15(cx+1)^4 c^7}{(cx-1)^4} - \frac{20(cx+1)^3 c^7}{(cx-1)^3} + \frac{15(cx+1)^2 c^7}{(cx-1)^2} - \frac{6(cx+1)c^7}{cx-1} + c^7} + \frac{90(cx+1)^5 a}{(cx-1)^5} + \frac{300(cx+1)^3 a}{(cx-1)^3} + \frac{90(cx+1)^5 a}{(cx-1)^5} + \frac{300(cx+1)^3 a}{(cx-1)^3} \right)$$

input `integrate(x^5*(a+b*arctanh(c*x)),x, algorithm="giac")`

output

```
1/45*c*(15*(3*(c*x + 1)^5*b/(c*x - 1)^5 + 10*(c*x + 1)^3*b/(c*x - 1)^3 + 3*(c*x + 1)*b/(c*x - 1))*log(-(c*x + 1)/(c*x - 1))/((c*x + 1)^6*c^7/(c*x - 1)^6 - 6*(c*x + 1)^5*c^7/(c*x - 1)^5 + 15*(c*x + 1)^4*c^7/(c*x - 1)^4 - 20*(c*x + 1)^3*c^7/(c*x - 1)^3 + 15*(c*x + 1)^2*c^7/(c*x - 1)^2 - 6*(c*x + 1)*c^7/(c*x - 1) + c^7) + (90*(c*x + 1)^5*a/(c*x - 1)^5 + 300*(c*x + 1)^3*a/(c*x - 1)^3 + 90*(c*x + 1)*a/(c*x - 1) + 45*(c*x + 1)^5*b/(c*x - 1)^5 - 135*(c*x + 1)^4*b/(c*x - 1)^4 + 230*(c*x + 1)^3*b/(c*x - 1)^3 - 210*(c*x + 1)^2*b/(c*x - 1)^2 + 93*(c*x + 1)*b/(c*x - 1) - 23*b)/((c*x + 1)^6*c^7/(c*x - 1)^6 - 6*(c*x + 1)^5*c^7/(c*x - 1)^5 + 15*(c*x + 1)^4*c^7/(c*x - 1)^4 - 20*(c*x + 1)^3*c^7/(c*x - 1)^3 + 15*(c*x + 1)^2*c^7/(c*x - 1)^2 - 6*(c*x + 1)*c^7/(c*x - 1) + c^7))
```

Mupad [B] (verification not implemented)

Time = 3.78 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.88

$$\int x^5(a + b \operatorname{arctanh}(cx)) dx = \frac{bc^3 x^3}{18} - \frac{b \operatorname{atanh}(cx)}{6} + \frac{bc^5 x^5}{30} + \frac{bcx}{6} + \frac{ax^6}{6} + \frac{bx^6 \operatorname{atanh}(cx)}{6}$$

input `int(x^5*(a + b*atanh(c*x)),x)`

output

```
((b*c^3*x^3)/18 - (b*atanh(c*x))/6 + (b*c^5*x^5)/30 + (b*c*x)/6)/c^6 + (a*x^6)/6 + (b*x^6*atanh(c*x))/6
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.98

$$\int x^5(a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{15 \operatorname{atanh}(cx) b c^6 x^6 - 15 \operatorname{atanh}(cx) b + 15 a c^6 x^6 + 3 b c^5 x^5 + 5 b c^3 x^3 + 15 b c x}{90 c^6}$$

input `int(x^5*(a+b*atanh(c*x)),x)`

output `(15*atanh(c*x)*b*c**6*x**6 - 15*atanh(c*x)*b + 15*a*c**6*x**6 + 3*b*c**5*x**5 + 5*b*c**3*x**3 + 15*b*c*x)/(90*c**6)`

3.2 $\int x^4(a + b \operatorname{arctanh}(cx)) dx$

Optimal result	120
Mathematica [A] (verified)	120
Rubi [A] (verified)	121
Maple [A] (verified)	122
Fricas [A] (verification not implemented)	123
Sympy [A] (verification not implemented)	123
Maxima [A] (verification not implemented)	124
Giac [B] (verification not implemented)	124
Mupad [B] (verification not implemented)	125
Reduce [B] (verification not implemented)	125

Optimal result

Integrand size = 12, antiderivative size = 57

$$\int x^4(a + b \operatorname{arctanh}(cx)) dx = \frac{bx^2}{10c^3} + \frac{bx^4}{20c} + \frac{1}{5}x^5(a + b \operatorname{arctanh}(cx)) + \frac{b \log(1 - c^2x^2)}{10c^5}$$

output

```
1/10*b*x^2/c^3+1/20*b*x^4/c+1/5*x^5*(a+b*arctanh(c*x))+1/10*b*ln(-c^2*x^2+1)/c^5
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.09

$$\int x^4(a + b \operatorname{arctanh}(cx)) dx = \frac{bx^2}{10c^3} + \frac{bx^4}{20c} + \frac{ax^5}{5} + \frac{1}{5}bx^5 \operatorname{arctanh}(cx) + \frac{b \log(1 - c^2x^2)}{10c^5}$$

input

```
Integrate[x^4*(a + b*ArcTanh[c*x]),x]
```

output

```
(b*x^2)/(10*c^3) + (b*x^4)/(20*c) + (a*x^5)/5 + (b*x^5*ArcTanh[c*x])/5 + (b*Log[1 - c^2*x^2])/(10*c^5)
```

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6452, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^4(a + b \operatorname{arctanh}(cx)) dx \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{5}x^5(a + b \operatorname{arctanh}(cx)) - \frac{1}{5}bc \int \frac{x^5}{1 - c^2x^2} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{5}x^5(a + b \operatorname{arctanh}(cx)) - \frac{1}{10}bc \int \frac{x^4}{1 - c^2x^2} dx^2 \\
 & \quad \downarrow \text{49} \\
 & \frac{1}{5}x^5(a + b \operatorname{arctanh}(cx)) - \frac{1}{10}bc \int \left(-\frac{x^2}{c^2} - \frac{1}{c^4(c^2x^2 - 1)} - \frac{1}{c^4} \right) dx^2 \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{5}x^5(a + b \operatorname{arctanh}(cx)) - \frac{1}{10}bc \left(-\frac{x^2}{c^4} - \frac{x^4}{2c^2} - \frac{\log(1 - c^2x^2)}{c^6} \right)
 \end{aligned}$$

input `Int [x^4*(a + b*ArcTanh[c*x]),x]`

output `(x^5*(a + b*ArcTanh[c*x]))/5 - (b*c*(-(x^2/c^4) - x^4/(2*c^2) - Log[1 - c^2*x^2]/c^6))/10`

Defintions of rubi rules used

rule 49 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}*((c_.) + (d_.)(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{IGtQ}[m, 0] \&\& \text{IGtQ}[m + n + 2, 0]$

rule 243 $\text{Int}[(x_)^{(m_.)}*((a_) + (b_.)(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{((m - 1)/2)*(a + b*x)^p}, x], x, x^2], x] /; \text{FreeQ}[\{a, b, m, p\}, x] \&\& \text{IntegerQ}[(m - 1)/2]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 6452 $\text{Int}[(a_.) + \text{ArcTanh}[(c_.)(x_)^{(n_.)}]*(b_.)^{(p_.)}(x_)^{(m_.)}], x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m + 1)), x] - \text{Simp}[b*c*n*(p/(m + 1)) \text{ Int}[x^{(m + n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p - 1)/(1 - c^2*x^{(2*n)})}), x], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \&\& \text{IGtQ}[p, 0] \&\& (\text{EqQ}[p, 1] \|\| (\text{EqQ}[n, 1] \&\& \text{IntegerQ}[m])) \&\& \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.02

method	result	size
parts	$\frac{ax^5}{5} + \frac{b \left(\frac{c^5 x^5 \operatorname{arctanh}(cx)}{5} + \frac{c^4 x^4}{20} + \frac{c^2 x^2}{10} + \frac{\ln(cx-1)}{10} + \frac{\ln(cx+1)}{10} \right)}{c^5}$	58
derivativedivides	$\frac{\frac{ac^5x^5}{5} + b \left(\frac{c^5 x^5 \operatorname{arctanh}(cx)}{5} + \frac{c^4 x^4}{20} + \frac{c^2 x^2}{10} + \frac{\ln(cx-1)}{10} + \frac{\ln(cx+1)}{10} \right)}{c^5}$	62
default	$\frac{\frac{ac^5x^5}{5} + b \left(\frac{c^5 x^5 \operatorname{arctanh}(cx)}{5} + \frac{c^4 x^4}{20} + \frac{c^2 x^2}{10} + \frac{\ln(cx-1)}{10} + \frac{\ln(cx+1)}{10} \right)}{c^5}$	62
parallelrisch	$\frac{4b \operatorname{arctanh}(cx)x^5 c^5 + 4a c^5 x^5 + b c^4 x^4 + 2b c^2 x^2 + 4 \ln(cx-1)b + 4b \operatorname{arctanh}(cx) + 2b}{20c^5}$	65
risch	$\frac{bx^5 \ln(cx+1)}{10} - \frac{bx^5 \ln(-cx+1)}{10} + \frac{ax^5}{5} + \frac{bx^4}{20c} + \frac{bx^2}{10c^3} + \frac{b \ln(c^2 x^2 - 1)}{10c^5}$	67

input $\text{int}(x^4*(a+b*\operatorname{arctanh}(c*x)), x, \text{method}=_RETURNVERBOSE)$

output `1/5*a*x^5+b/c^5*(1/5*c^5*x^5*arctanh(c*x)+1/20*c^4*x^4+1/10*c^2*x^2+1/10*ln(c*x-1)+1/10*ln(c*x+1))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.21

$$\int x^4(a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{2bc^5x^5 \log\left(-\frac{cx+1}{cx-1}\right) + 4ac^5x^5 + bc^4x^4 + 2bc^2x^2 + 2b \log(c^2x^2 - 1)}{20c^5}$$

input `integrate(x^4*(a+b*arctanh(c*x)),x, algorithm="fricas")`

output `1/20*(2*b*c^5*x^5*log(-(c*x + 1)/(c*x - 1)) + 4*a*c^5*x^5 + b*c^4*x^4 + 2*b*c^2*x^2 + 2*b*log(c^2*x^2 - 1))/c^5`

Sympy [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.19

$$\int x^4(a + b \operatorname{atanh}(cx)) dx$$

$$= \begin{cases} \frac{ax^5}{5} + \frac{bx^5 \operatorname{atanh}(cx)}{5} + \frac{bx^4}{20c} + \frac{bx^2}{10c^3} + \frac{b \log\left(x - \frac{1}{c}\right)}{5c^5} + \frac{b \operatorname{atanh}(cx)}{5c^5} & \text{for } c \neq 0 \\ \frac{ax^5}{5} & \text{otherwise} \end{cases}$$

input `integrate(x**4*(a+b*atanh(c*x)),x)`

output `Piecewise((a*x**5/5 + b*x**5*atanh(c*x)/5 + b*x**4/(20*c) + b*x**2/(10*c**3) + b*log(x - 1/c)/(5*c**5) + b*atanh(c*x)/(5*c**5), Ne(c, 0)), (a*x**5/5, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.96

$$\int x^4(a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{1}{5} ax^5 + \frac{1}{20} \left(4x^5 \operatorname{arctanh}(cx) + c \left(\frac{c^2 x^4 + 2x^2}{c^4} + \frac{2 \log(c^2 x^2 - 1)}{c^6} \right) \right) b$$

input `integrate(x^4*(a+b*arctanh(c*x)),x, algorithm="maxima")`

output `1/5*a*x^5 + 1/20*(4*x^5*arctanh(c*x) + c*((c^2*x^4 + 2*x^2)/c^4 + 2*log(c^2*x^2 - 1)/c^6))*b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 403 vs. 2(49) = 98.

Time = 0.12 (sec) , antiderivative size = 403, normalized size of antiderivative = 7.07

$$\int x^4(a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{1}{5} c \left(\frac{\left(\frac{5(cx+1)^4 b}{(cx-1)^4} + \frac{10(cx+1)^2 b}{(cx-1)^2} + b \right) \log\left(-\frac{cx+1}{cx-1}\right)}{\frac{(cx+1)^5 c^6}{(cx-1)^5} - \frac{5(cx+1)^4 c^6}{(cx-1)^4} + \frac{10(cx+1)^3 c^6}{(cx-1)^3} - \frac{10(cx+1)^2 c^6}{(cx-1)^2} + \frac{5(cx+1)c^6}{cx-1} - c^6} + \frac{2 \left(\frac{5(cx+1)^4 a}{(cx-1)^4} + \frac{10(cx+1)^2 a}{(cx-1)^2} + a + \frac{2}{c} \right)}{\frac{(cx+1)^5 c^6}{(cx-1)^5} - \frac{5(cx+1)^4 c^6}{(cx-1)^4} + \frac{10(cx+1)^3 c^6}{(cx-1)^3} - \frac{10(cx+1)^2 c^6}{(cx-1)^2} + \frac{5(cx+1)c^6}{cx-1} - c^6} \right)$$

input `integrate(x^4*(a+b*arctanh(c*x)),x, algorithm="giac")`

output `1/5*c*((5*(c*x + 1)^4*b/(c*x - 1)^4 + 10*(c*x + 1)^2*b/(c*x - 1)^2 + b)*log(-(c*x + 1)/(c*x - 1))/((c*x + 1)^5*c^6/(c*x - 1)^5 - 5*(c*x + 1)^4*c^6/(c*x - 1)^4 + 10*(c*x + 1)^3*c^6/(c*x - 1)^3 - 10*(c*x + 1)^2*c^6/(c*x - 1)^2 + 5*(c*x + 1)*c^6/(c*x - 1) - c^6) + 2*(5*(c*x + 1)^4*a/(c*x - 1)^4 + 10*(c*x + 1)^2*a/(c*x - 1)^2 + a + 2*(c*x + 1)^4*b/(c*x - 1)^4 - 4*(c*x + 1)^3*b/(c*x - 1)^3 + 4*(c*x + 1)^2*b/(c*x - 1)^2 - 2*(c*x + 1)*b/(c*x - 1))/((c*x + 1)^5*c^6/(c*x - 1)^5 - 5*(c*x + 1)^4*c^6/(c*x - 1)^4 + 10*(c*x + 1)^3*c^6/(c*x - 1)^3 - 10*(c*x + 1)^2*c^6/(c*x - 1)^2 + 5*(c*x + 1)*c^6/(c*x - 1) - c^6) - b*log(-(c*x + 1)/(c*x - 1) + 1)/c^6 + b*log(-(c*x + 1)/(c*x - 1))/c^6)`

Mupad [B] (verification not implemented)

Time = 3.75 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.93

$$\int x^4(a + b \operatorname{arctanh}(cx)) dx = \frac{a x^5}{5} + \frac{b \ln(c^2 x^2 - 1)}{10} + \frac{b c^2 x^2}{10} + \frac{b c^4 x^4}{20} + \frac{b x^5 \operatorname{atanh}(cx)}{5}$$

input `int(x^4*(a + b*atanh(c*x)),x)`output `(a*x^5)/5 + ((b*log(c^2*x^2 - 1))/10 + (b*c^2*x^2)/10 + (b*c^4*x^4)/20)/c^5 + (b*x^5*atanh(c*x))/5`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.14

$$\int x^4(a + b \operatorname{arctanh}(cx)) dx = \frac{4 \operatorname{atanh}(cx) b c^5 x^5 + 4 \operatorname{atanh}(cx) b + 4 \log(c^2 x - c) b + 4 a c^5 x^5 + b c^4 x^4 + 2 b c^2 x^2}{20 c^5}$$

input `int(x^4*(a+b*atanh(c*x)),x)`output `(4*atanh(c*x)*b*c**5*x**5 + 4*atanh(c*x)*b + 4*log(c**2*x - c)*b + 4*a*c**5*x**5 + b*c**4*x**4 + 2*b*c**2*x**2)/(20*c**5)`

3.3 $\int x^3(a + \operatorname{barctanh}(cx)) dx$

Optimal result	126
Mathematica [A] (verified)	126
Rubi [A] (verified)	127
Maple [A] (verified)	128
Fricas [A] (verification not implemented)	128
Sympy [A] (verification not implemented)	129
Maxima [A] (verification not implemented)	129
Giac [B] (verification not implemented)	130
Mupad [B] (verification not implemented)	130
Reduce [B] (verification not implemented)	131

Optimal result

Integrand size = 12, antiderivative size = 48

$$\int x^3(a + \operatorname{barctanh}(cx)) dx = \frac{bx}{4c^3} + \frac{bx^3}{12c} - \frac{\operatorname{barctanh}(cx)}{4c^4} + \frac{1}{4}x^4(a + \operatorname{barctanh}(cx))$$

output `1/4*b*x/c^3+1/12*b*x^3/c-1/4*b*arctanh(c*x)/c^4+1/4*x^4*(a+b*arctanh(c*x))`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.46

$$\int x^3(a + \operatorname{barctanh}(cx)) dx = \frac{bx}{4c^3} + \frac{bx^3}{12c} + \frac{ax^4}{4} + \frac{1}{4}bx^4\operatorname{arctanh}(cx) + \frac{b \log(1 - cx)}{8c^4} - \frac{b \log(1 + cx)}{8c^4}$$

input `Integrate[x^3*(a + b*ArcTanh[c*x]),x]`

output `(b*x)/(4*c^3) + (b*x^3)/(12*c) + (a*x^4)/4 + (b*x^4*ArcTanh[c*x])/4 + (b*Log[1 - c*x])/(8*c^4) - (b*Log[1 + c*x])/(8*c^4)`

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.98, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6452, 254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3(a + b \operatorname{arctanh}(cx)) dx$$

$$\downarrow 6452$$

$$\frac{1}{4}x^4(a + b \operatorname{arctanh}(cx)) - \frac{1}{4}bc \int \frac{x^4}{1 - c^2x^2} dx$$

$$\downarrow 254$$

$$\frac{1}{4}x^4(a + b \operatorname{arctanh}(cx)) - \frac{1}{4}bc \int \left(-\frac{x^2}{c^2} + \frac{1}{c^4(1 - c^2x^2)} - \frac{1}{c^4} \right) dx$$

$$\downarrow 2009$$

$$\frac{1}{4}x^4(a + b \operatorname{arctanh}(cx)) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(cx)}{c^5} - \frac{x}{c^4} - \frac{x^3}{3c^2} \right)$$

input `Int[x^3*(a + b*ArcTanh[c*x]),x]`

output `(x^4*(a + b*ArcTanh[c*x]))/4 - (b*c*(-(x/c^4) - x^3/(3*c^2) + ArcTanh[c*x]/c^5))/4`

Defintions of rubi rules used

rule 254 `Int[(x_)^(m_)/((a_) + (b_.)*(x_)^2), x_Symbol] :> Int[PolynomialDivide[x^m, a + b*x^2, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 3]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.04

method	result	size
parallelrisc	$-\frac{-3b \operatorname{arctanh}(cx)x^4c^4 - 3ac^4x^4 - bc^3x^3 - 3bcx + 3b \operatorname{arctanh}(cx)}{12c^4}$	50
parts	$\frac{x^4a}{4} + \frac{b\left(\frac{c^4x^4 \operatorname{arctanh}(cx)}{4} + \frac{c^3x^3}{12} + \frac{cx}{4} + \frac{\ln(cx-1)}{8} - \frac{\ln(cx+1)}{8}\right)}{c^4}$	54
derivativedivides	$\frac{\frac{ac^4x^4}{4} + b\left(\frac{c^4x^4 \operatorname{arctanh}(cx)}{4} + \frac{c^3x^3}{12} + \frac{cx}{4} + \frac{\ln(cx-1)}{8} - \frac{\ln(cx+1)}{8}\right)}{c^4}$	58
default	$\frac{\frac{ac^4x^4}{4} + b\left(\frac{c^4x^4 \operatorname{arctanh}(cx)}{4} + \frac{c^3x^3}{12} + \frac{cx}{4} + \frac{\ln(cx-1)}{8} - \frac{\ln(cx+1)}{8}\right)}{c^4}$	58
risc	$\frac{bx^4 \ln(cx+1)}{8} - \frac{bx^4 \ln(-cx+1)}{8} + \frac{x^4a}{4} + \frac{bx^3}{12c} + \frac{bx}{4c^3} + \frac{b \ln(-cx+1)}{8c^4} - \frac{b \ln(cx+1)}{8c^4}$	74
orering	$\frac{(c^4x^4 + c^2x^2 - 2)(a + b \operatorname{arctanh}(cx))}{2c^4} - \frac{(c^2x^2 + 3)(cx - 1)(cx + 1)\left(3x^2(a + b \operatorname{arctanh}(cx)) + \frac{x^3bc}{-c^2x^2 + 1}\right)}{12x^2c^4}$	90

input

```
int(x^3*(a+b*arctanh(c*x)),x,method=_RETURNVERBOSE)
```

output

```
-1/12*(-3*b*arctanh(c*x)*x^4*c^4-3*a*c^4*x^4-b*c^3*x^3-3*b*c*x+3*b*arctanh
(c*x))/c^4
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.21

$$\int x^3(a + b \operatorname{arctanh}(cx)) dx = \frac{6ac^4x^4 + 2bc^3x^3 + 6bcx + 3(bc^4x^4 - b) \log\left(-\frac{cx+1}{cx-1}\right)}{24c^4}$$

input

```
integrate(x^3*(a+b*arctanh(c*x)),x, algorithm="fricas")
```

output $\frac{1}{24}(6ac^4x^4 + 2b^2c^3x^3 + 6b^2cx + 3(b^2c^4x^4 - b^2)\log(-(cx + 1)/(cx - 1)))/c^4$

Sympy [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.10

$$\int x^3(a + b \operatorname{arctanh}(cx)) dx = \begin{cases} \frac{ax^4}{4} + \frac{bx^4 \operatorname{atanh}(cx)}{4} + \frac{bx^3}{12c} + \frac{bx}{4c^3} - \frac{b \operatorname{atanh}(cx)}{4c^4} & \text{for } c \neq 0 \\ \frac{ax^4}{4} & \text{otherwise} \end{cases}$$

input `integrate(x**3*(a+b*atanh(c*x)),x)`

output `Piecewise((a*x**4/4 + b*x**4*atanh(c*x)/4 + b*x**3/(12*c) + b*x/(4*c**3) - b*atanh(c*x)/(4*c**4), Ne(c, 0)), (a*x**4/4, True))`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.27

$$\begin{aligned} & \int x^3(a + b \operatorname{arctanh}(cx)) dx \\ &= \frac{1}{4} ax^4 \\ &+ \frac{1}{24} \left(6x^4 \operatorname{artanh}(cx) + c \left(\frac{2(c^2x^3 + 3x)}{c^4} - \frac{3 \log(cx + 1)}{c^5} + \frac{3 \log(cx - 1)}{c^5} \right) \right) b \end{aligned}$$

input `integrate(x^3*(a+b*arctanh(c*x)),x, algorithm="maxima")`

output $\frac{1}{4}ax^4 + \frac{1}{24}(6x^4 \operatorname{arctanh}(cx) + c(2(c^2x^3 + 3x)/c^4 - 3 \log(cx + 1)/c^5 + 3 \log(cx - 1)/c^5))b$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 296 vs. $2(40) = 80$.

Time = 0.12 (sec) , antiderivative size = 296, normalized size of antiderivative = 6.17

$$\int x^3(a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{1}{3} c \left(\frac{3 \left(\frac{(cx+1)^3 b}{(cx-1)^3} + \frac{(cx+1)b}{cx-1} \right) \log\left(-\frac{cx+1}{cx-1}\right)}{\frac{(cx+1)^4 c^5}{(cx-1)^4} - \frac{4(cx+1)^3 c^5}{(cx-1)^3} + \frac{6(cx+1)^2 c^5}{(cx-1)^2} - \frac{4(cx+1)c^5}{cx-1} + c^5} + \frac{\frac{6(cx+1)^3 a}{(cx-1)^3} + \frac{6(cx+1)a}{cx-1} + \frac{3(cx+1)^3 b}{(cx-1)^3} - \frac{6(cx+1)^2 b}{(cx-1)^2} + \frac{5}{cx-1}}{\frac{(cx+1)^4 c^5}{(cx-1)^4} - \frac{4(cx+1)^3 c^5}{(cx-1)^3} + \frac{6(cx+1)^2 c^5}{(cx-1)^2} - \frac{4(cx+1)c^5}{cx-1} + c^5} \right)$$

input `integrate(x^3*(a+b*arctanh(c*x)),x, algorithm="giac")`

output `1/3*c*(3*((c*x + 1)^3*b/(c*x - 1)^3 + (c*x + 1)*b/(c*x - 1))*log(-(c*x + 1)/(c*x - 1))/((c*x + 1)^4*c^5/(c*x - 1)^4 - 4*(c*x + 1)^3*c^5/(c*x - 1)^3 + 6*(c*x + 1)^2*c^5/(c*x - 1)^2 - 4*(c*x + 1)*c^5/(c*x - 1) + c^5) + (6*(c*x + 1)^3*a/(c*x - 1)^3 + 6*(c*x + 1)*a/(c*x - 1) + 3*(c*x + 1)^3*b/(c*x - 1)^3 - 6*(c*x + 1)^2*b/(c*x - 1)^2 + 5*(c*x + 1)*b/(c*x - 1) - 2*b)/((c*x + 1)^4*c^5/(c*x - 1)^4 - 4*(c*x + 1)^3*c^5/(c*x - 1)^3 + 6*(c*x + 1)^2*c^5/(c*x - 1)^2 - 4*(c*x + 1)*c^5/(c*x - 1) + c^5))`

Mupad [B] (verification not implemented)

Time = 3.75 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.90

$$\int x^3(a + b \operatorname{arctanh}(cx)) dx = \frac{a x^4}{4} + \frac{b c^3 x^3}{12} - \frac{b \operatorname{atanh}(cx)}{4} + \frac{b c x}{4} + \frac{b x^4 \operatorname{atanh}(cx)}{4}$$

input `int(x^3*(a + b*atanh(c*x)),x)`

output `(a*x^4)/4 + ((b*c^3*x^3)/12 - (b*atanh(c*x))/4 + (b*c*x)/4)/c^4 + (b*x^4*a tanh(c*x))/4`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00

$$\int x^3(a + b \operatorname{arctanh}(cx)) dx = \frac{3 \operatorname{atanh}(cx) b c^4 x^4 - 3 \operatorname{atanh}(cx) b + 3 a c^4 x^4 + b c^3 x^3 + 3 b c x}{12 c^4}$$

input `int(x^3*(a+b*atanh(c*x)),x)`

output `(3*atanh(c*x)*b*c**4*x**4 - 3*atanh(c*x)*b + 3*a*c**4*x**4 + b*c**3*x**3 + 3*b*c*x)/(12*c**4)`

3.4 $\int x^2(a + b \operatorname{arctanh}(cx)) dx$

Optimal result	132
Mathematica [A] (verified)	132
Rubi [A] (verified)	133
Maple [A] (verified)	134
Fricas [A] (verification not implemented)	135
Sympy [A] (verification not implemented)	135
Maxima [A] (verification not implemented)	136
Giac [B] (verification not implemented)	136
Mupad [B] (verification not implemented)	137
Reduce [B] (verification not implemented)	137

Optimal result

Integrand size = 12, antiderivative size = 46

$$\int x^2(a + b \operatorname{arctanh}(cx)) dx = \frac{bx^2}{6c} + \frac{1}{3}x^3(a + b \operatorname{arctanh}(cx)) + \frac{b \log(1 - c^2x^2)}{6c^3}$$

output $1/6*b*x^2/c+1/3*x^3*(a+b*\operatorname{arctanh}(c*x))+1/6*b*\ln(-c^2*x^2+1)/c^3$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.11

$$\int x^2(a + b \operatorname{arctanh}(cx)) dx = \frac{bx^2}{6c} + \frac{ax^3}{3} + \frac{1}{3}bx^3 \operatorname{arctanh}(cx) + \frac{b \log(1 - c^2x^2)}{6c^3}$$

input $\operatorname{Integrate}[x^2*(a + b*\operatorname{ArcTanh}[c*x]),x]$

output $(b*x^2)/(6*c) + (a*x^3)/3 + (b*x^3*\operatorname{ArcTanh}[c*x])/3 + (b*\operatorname{Log}[1 - c^2*x^2])/(6*c^3)$

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6452, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2(a + b \operatorname{arctanh}(cx)) dx \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{3}x^3(a + b \operatorname{arctanh}(cx)) - \frac{1}{3}bc \int \frac{x^3}{1 - c^2x^2} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{3}x^3(a + b \operatorname{arctanh}(cx)) - \frac{1}{6}bc \int \frac{x^2}{1 - c^2x^2} dx^2 \\
 & \quad \downarrow \text{49} \\
 & \frac{1}{3}x^3(a + b \operatorname{arctanh}(cx)) - \frac{1}{6}bc \int \left(-\frac{1}{c^2} - \frac{1}{c^2(c^2x^2 - 1)} \right) dx^2 \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{3}x^3(a + b \operatorname{arctanh}(cx)) - \frac{1}{6}bc \left(-\frac{x^2}{c^2} - \frac{\log(1 - c^2x^2)}{c^4} \right)
 \end{aligned}$$

input `Int [x^2*(a + b*ArcTanh[c*x]),x]`

output `(x^3*(a + b*ArcTanh[c*x]))/3 - (b*c*(-(x^2/c^2) - Log[1 - c^2*x^2]/c^4))/6`

Definitions of rubi rules used

rule 49 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}*((c_.) + (d_.)(x_)^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{IGtQ}[m, 0] \&\& \text{IGtQ}[m + n + 2, 0]$

rule 243 $\text{Int}[(x_)^{(m_.)}*((a_) + (b_.)(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] /; \text{FreeQ}[\{a, b, m, p\}, x] \&\& \text{IntegerQ}[(m-1)/2]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 6452 $\text{Int}[(a_.) + \text{ArcTanh}[(c_.)(x_)^{(n_.)}]*(b_.)]^{(p_.)}(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{2*n}))}, x], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \&\& \text{IGtQ}[p, 0] \&\& (\text{EqQ}[p, 1] \|\| (\text{EqQ}[n, 1] \&\& \text{IntegerQ}[m])) \&\& \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.09

method	result	size
parts	$\frac{ax^3}{3} + \frac{b\left(\frac{c^3x^3 \operatorname{arctanh}(cx)}{3} + \frac{c^2x^2}{6} + \frac{\ln(cx-1)}{6} + \frac{\ln(cx+1)}{6}\right)}{c^3}$	50
parallelrisch	$\frac{2b \operatorname{arctanh}(cx)x^3c^3 + 2c^3x^3a + bc^2x^2 + 2\ln(cx-1)b + 2b \operatorname{arctanh}(cx)}{6c^3}$	53
derivativdivides	$\frac{\frac{c^3x^3a}{3} + b\left(\frac{c^3x^3 \operatorname{arctanh}(cx)}{3} + \frac{c^2x^2}{6} + \frac{\ln(cx-1)}{6} + \frac{\ln(cx+1)}{6}\right)}{c^3}$	54
default	$\frac{\frac{c^3x^3a}{3} + b\left(\frac{c^3x^3 \operatorname{arctanh}(cx)}{3} + \frac{c^2x^2}{6} + \frac{\ln(cx-1)}{6} + \frac{\ln(cx+1)}{6}\right)}{c^3}$	54
risch	$\frac{bx^3 \ln(cx+1)}{6} - \frac{bx^3 \ln(-cx+1)}{6} + \frac{ax^3}{3} + \frac{bx^2}{6c} + \frac{b \ln(c^2x^2-1)}{6c^3}$	58

input $\text{int}(x^2*(a+b*\operatorname{arctanh}(c*x)), x, \text{method}=_RETURNVERBOSE)$

output

```
1/3*a*x^3+b/c^3*(1/3*c^3*x^3*arctanh(c*x)+1/6*c^2*x^2+1/6*ln(c*x-1)+1/6*ln
(c*x+1))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.26

$$\int x^2(a + b \operatorname{arctanh}(cx)) dx = \frac{bc^3 x^3 \log\left(-\frac{cx+1}{cx-1}\right) + 2ac^3 x^3 + bc^2 x^2 + b \log(c^2 x^2 - 1)}{6c^3}$$

input

```
integrate(x^2*(a+b*arctanh(c*x)),x, algorithm="fricas")
```

output

```
1/6*(b*c^3*x^3*log(-(c*x + 1)/(c*x - 1)) + 2*a*c^3*x^3 + b*c^2*x^2 + b*log
(c^2*x^2 - 1))/c^3
```

Sympy [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.26

$$\int x^2(a + b \operatorname{arctanh}(cx)) dx = \begin{cases} \frac{ax^3}{3} + \frac{bx^3 \operatorname{atanh}(cx)}{3} + \frac{bx^2}{6c} + \frac{b \log\left(x - \frac{1}{c}\right)}{3c^3} + \frac{b \operatorname{atanh}(cx)}{3c^3} & \text{for } c \neq 0 \\ \frac{ax^3}{3} & \text{otherwise} \end{cases}$$

input

```
integrate(x**2*(a+b*atanh(c*x)),x)
```

output

```
Piecewise((a*x**3/3 + b*x**3*atanh(c*x)/3 + b*x**2/(6*c) + b*log(x - 1/c)/
(3*c**3) + b*atanh(c*x)/(3*c**3), Ne(c, 0)), (a*x**3/3, True))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.96

$$\int x^2(a + b \operatorname{arctanh}(cx)) dx = \frac{1}{3} ax^3 + \frac{1}{6} \left(2x^3 \operatorname{arctanh}(cx) + c \left(\frac{x^2}{c^2} + \frac{\log(c^2 x^2 - 1)}{c^4} \right) \right) b$$

input `integrate(x^2*(a+b*arctanh(c*x)),x, algorithm="maxima")`

output `1/3*a*x^3 + 1/6*(2*x^3*arctanh(c*x) + c*(x^2/c^2 + log(c^2*x^2 - 1)/c^4))*
b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 258 vs. 2(40) = 80.

Time = 0.12 (sec) , antiderivative size = 258, normalized size of antiderivative = 5.61

$$\int x^2(a + b \operatorname{arctanh}(cx)) dx = \frac{1}{3} c \left(\frac{\left(\frac{3(cx+1)^2 b}{(cx-1)^2} + b \right) \log\left(-\frac{cx+1}{cx-1}\right)}{\frac{(cx+1)^3 c^4}{(cx-1)^3} - \frac{3(cx+1)^2 c^4}{(cx-1)^2} + \frac{3(cx+1)c^4}{cx-1} - c^4} + \frac{2 \left(\frac{3(cx+1)^2 a}{(cx-1)^2} + a + \frac{(cx+1)^2 b}{(cx-1)^2} - \frac{(cx+1)b}{cx-1} \right)}{\frac{(cx+1)^3 c^4}{(cx-1)^3} - \frac{3(cx+1)^2 c^4}{(cx-1)^2} + \frac{3(cx+1)c^4}{cx-1} - c^4} - \frac{b \log\left(-\frac{cx+1}{cx-1} + 1\right)}{c^4} \right)$$

input `integrate(x^2*(a+b*arctanh(c*x)),x, algorithm="giac")`

output `1/3*c*((3*(c*x + 1)^2*b/(c*x - 1)^2 + b)*log(-(c*x + 1)/(c*x - 1))/((c*x + 1)^3*c^4/(c*x - 1)^3 - 3*(c*x + 1)^2*c^4/(c*x - 1)^2 + 3*(c*x + 1)*c^4/(c*x - 1) - c^4) + 2*(3*(c*x + 1)^2*a/(c*x - 1)^2 + a + (c*x + 1)^2*b/(c*x - 1)^2 - (c*x + 1)*b/(c*x - 1))/((c*x + 1)^3*c^4/(c*x - 1)^3 - 3*(c*x + 1)^2*c^4/(c*x - 1)^2 + 3*(c*x + 1)*c^4/(c*x - 1) - c^4) - b*log(-(c*x + 1)/(c*x - 1) + 1)/c^4 + b*log(-(c*x + 1)/(c*x - 1))/c^4`

Mupad [B] (verification not implemented)

Time = 3.67 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.96

$$\int x^2(a + b \operatorname{arctanh}(cx)) dx = \frac{b \ln(c^2 x^2 - 1)}{6} + \frac{b c^2 x^2}{6} + \frac{a x^3}{3} + \frac{b x^3 \operatorname{atanh}(cx)}{3}$$

input `int(x^2*(a + b*atanh(c*x)),x)`

output `((b*log(c^2*x^2 - 1))/6 + (b*c^2*x^2)/6)/c^3 + (a*x^3)/3 + (b*x^3*atanh(c*x))/3`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.22

$$\int x^2(a + b \operatorname{arctanh}(cx)) dx = \frac{2 \operatorname{atanh}(cx) b c^3 x^3 + 2 \operatorname{atanh}(cx) b + 2 \log(c^2 x - c) b + 2 a c^3 x^3 + b c^2 x^2}{6 c^3}$$

input `int(x^2*(a+b*atanh(c*x)),x)`

output `(2*atanh(c*x)*b*c**3*x**3 + 2*atanh(c*x)*b + 2*log(c**2*x - c)*b + 2*a*c**3*x**3 + b*c**2*x**2)/(6*c**3)`

3.5 $\int x(a + b \operatorname{arctanh}(cx)) dx$

Optimal result	138
Mathematica [A] (verified)	138
Rubi [A] (verified)	139
Maple [A] (verified)	140
Fricas [A] (verification not implemented)	141
Sympy [A] (verification not implemented)	141
Maxima [A] (verification not implemented)	141
Giac [B] (verification not implemented)	142
Mupad [B] (verification not implemented)	142
Reduce [B] (verification not implemented)	143

Optimal result

Integrand size = 10, antiderivative size = 37

$$\int x(a + b \operatorname{arctanh}(cx)) dx = \frac{bx}{2c} - \frac{b \operatorname{arctanh}(cx)}{2c^2} + \frac{1}{2}x^2(a + b \operatorname{arctanh}(cx))$$

output `1/2*b*x/c-1/2*b*arctanh(c*x)/c^2+1/2*x^2*(a+b*arctanh(c*x))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.59

$$\int x(a + b \operatorname{arctanh}(cx)) dx = \frac{bx}{2c} + \frac{ax^2}{2} + \frac{1}{2}bx^2 \operatorname{arctanh}(cx) + \frac{b \log(1 - cx)}{4c^2} - \frac{b \log(1 + cx)}{4c^2}$$

input `Integrate[x*(a + b*ArcTanh[c*x]),x]`

output `(b*x)/(2*c) + (a*x^2)/2 + (b*x^2*ArcTanh[c*x])/2 + (b*Log[1 - c*x])/(4*c^2) - (b*Log[1 + c*x])/(4*c^2)`

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {6452, 262, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(a + b \operatorname{arctanh}(cx)) dx$$

$$\downarrow 6452$$

$$\frac{1}{2}x^2(a + b \operatorname{arctanh}(cx)) - \frac{1}{2}bc \int \frac{x^2}{1 - c^2x^2} dx$$

$$\downarrow 262$$

$$\frac{1}{2}x^2(a + b \operatorname{arctanh}(cx)) - \frac{1}{2}bc \left(\frac{\int \frac{1}{1 - c^2x^2} dx}{c^2} - \frac{x}{c^2} \right)$$

$$\downarrow 219$$

$$\frac{1}{2}x^2(a + b \operatorname{arctanh}(cx)) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(cx)}{c^3} - \frac{x}{c^2} \right)$$

input `Int[x*(a + b*ArcTanh[c*x]),x]`

output `(x^2*(a + b*ArcTanh[c*x]))/2 - (b*c*(-(x/c^2) + ArcTanh[c*x]/c^3))/2`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 262

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.08

method	result	size
parallelrisch	$\frac{-\operatorname{arctanh}(cx)b c^2 x^2 - a c^2 x^2 - b c x + b \operatorname{arctanh}(cx)}{2c^2}$	40
parts	$\frac{a x^2}{2} + \frac{b \left(\frac{c^2 x^2 \operatorname{arctanh}(cx)}{2} + \frac{cx}{2} + \frac{\ln(cx-1)}{4} - \frac{\ln(cx+1)}{4} \right)}{c^2}$	46
derivativedivides	$\frac{\frac{a c^2 x^2}{2} + b \left(\frac{c^2 x^2 \operatorname{arctanh}(cx)}{2} + \frac{cx}{2} + \frac{\ln(cx-1)}{4} - \frac{\ln(cx+1)}{4} \right)}{c^2}$	50
default	$\frac{\frac{a c^2 x^2}{2} + b \left(\frac{c^2 x^2 \operatorname{arctanh}(cx)}{2} + \frac{cx}{2} + \frac{\ln(cx-1)}{4} - \frac{\ln(cx+1)}{4} \right)}{c^2}$	50
orering	$\frac{(c^2 x^2 - 1)(a + b \operatorname{arctanh}(cx))}{c^2} - \frac{(cx-1)(cx+1) \left(a + b \operatorname{arctanh}(cx) + \frac{xbc}{-c^2 x^2 + 1} \right)}{2c^2}$	62
risch	$\frac{x^2 b \ln(cx+1)}{4} - \frac{x^2 b \ln(-cx+1)}{4} + \frac{a x^2}{2} + \frac{bx}{2c} + \frac{b \ln(-cx+1)}{4c^2} - \frac{b \ln(cx+1)}{4c^2}$	65

input

```
int(x*(a+b*arctanh(c*x)),x,method=_RETURNVERBOSE)
```

output

```
-1/2*(-arctanh(c*x)*b*c^2*x^2-a*c^2*x^2-b*c*x+b*arctanh(c*x))/c^2
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.30

$$\int x(a + b \operatorname{arctanh}(cx)) dx = \frac{2ac^2x^2 + 2bcx + (bc^2x^2 - b) \log\left(-\frac{cx+1}{cx-1}\right)}{4c^2}$$

input `integrate(x*(a+b*arctanh(c*x)),x, algorithm="fricas")`

output `1/4*(2*a*c^2*x^2 + 2*b*c*x + (b*c^2*x^2 - b)*log(-(c*x + 1)/(c*x - 1)))/c^2`

Sympy [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.14

$$\int x(a + b \operatorname{arctanh}(cx)) dx = \begin{cases} \frac{ax^2}{2} + \frac{bx^2 \operatorname{atanh}(cx)}{2} + \frac{bx}{2c} - \frac{b \operatorname{atanh}(cx)}{2c^2} & \text{for } c \neq 0 \\ \frac{ax^2}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*(a+b*atanh(c*x)),x)`

output `Piecewise((a*x**2/2 + b*x**2*atanh(c*x)/2 + b*x/(2*c) - b*atanh(c*x)/(2*c**2), Ne(c, 0)), (a*x**2/2, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.35

$$\int x(a + b \operatorname{arctanh}(cx)) dx = \frac{1}{2} ax^2 + \frac{1}{4} \left(2x^2 \operatorname{artanh}(cx) + c \left(\frac{2x}{c^2} - \frac{\log(cx+1)}{c^3} + \frac{\log(cx-1)}{c^3} \right) \right) b$$

input `integrate(x*(a+b*arctanh(c*x)),x, algorithm="maxima")`

output $1/2*a*x^2 + 1/4*(2*x^2*\operatorname{arctanh}(c*x) + c*(2*x/c^2 - \log(c*x + 1)/c^3 + \log(c*x - 1)/c^3))*b$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 148 vs. $2(31) = 62$.

Time = 0.12 (sec) , antiderivative size = 148, normalized size of antiderivative = 4.00

$$\int x(a + b \operatorname{arctanh}(cx)) dx$$

$$= c \left(\frac{(cx+1)b \log\left(-\frac{cx+1}{cx-1}\right)}{\left(\frac{(cx+1)^2 c^3}{(cx-1)^2} - \frac{2(cx+1)c^3}{cx-1} + c^3\right)(cx-1)} + \frac{\frac{2(cx+1)a}{cx-1} + \frac{(cx+1)b}{cx-1} - b}{\frac{(cx+1)^2 c^3}{(cx-1)^2} - \frac{2(cx+1)c^3}{cx-1} + c^3} \right)$$

input `integrate(x*(a+b*arctanh(c*x)),x, algorithm="giac")`

output $c*((c*x + 1)*b*\log(-(c*x + 1)/(c*x - 1))/(((c*x + 1)^2*c^3/(c*x - 1)^2 - 2*(c*x + 1)*c^3/(c*x - 1) + c^3)*(c*x - 1)) + (2*(c*x + 1)*a/(c*x - 1) + (c*x + 1)*b/(c*x - 1) - b)/((c*x + 1)^2*c^3/(c*x - 1)^2 - 2*(c*x + 1)*c^3/(c*x - 1) + c^3))$

Mupad [B] (verification not implemented)

Time = 3.64 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.95

$$\int x(a + b \operatorname{arctanh}(cx)) dx = \frac{a x^2}{2} - \frac{b \operatorname{atanh}(cx)}{2} - \frac{b c x}{2} + \frac{b x^2 \operatorname{atanh}(cx)}{2}$$

input `int(x*(a + b*atanh(c*x)),x)`

output $(a*x^2)/2 - ((b*\operatorname{atanh}(c*x))/2 - (b*c*x)/2)/c^2 + (b*x^2*\operatorname{atanh}(c*x))/2$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

$$\int x(a + b \operatorname{arctanh}(cx)) dx = \frac{\operatorname{atanh}(cx) b c^2 x^2 - \operatorname{atanh}(cx) b + a c^2 x^2 + bcx}{2c^2}$$

input `int(x*(a+b*atanh(c*x)),x)`

output `(atanh(c*x)*b*c**2*x**2 - atanh(c*x)*b + a*c**2*x**2 + b*c*x)/(2*c**2)`

3.6 $\int (a + b \operatorname{arctanh}(cx)) dx$

Optimal result	144
Mathematica [A] (verified)	144
Rubi [A] (verified)	145
Maple [A] (verified)	146
Fricas [A] (verification not implemented)	146
Sympy [A] (verification not implemented)	147
Maxima [A] (verification not implemented)	147
Giac [B] (verification not implemented)	147
Mupad [B] (verification not implemented)	148
Reduce [B] (verification not implemented)	148

Optimal result

Integrand size = 8, antiderivative size = 30

$$\int (a + b \operatorname{arctanh}(cx)) dx = ax + b \operatorname{arctanh}(cx) + \frac{b \log(1 - c^2 x^2)}{2c}$$

output `a*x+b*x*arctanh(c*x)+1/2*b*ln(-c^2*x^2+1)/c`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int (a + b \operatorname{arctanh}(cx)) dx = ax + b \operatorname{arctanh}(cx) + \frac{b \log(1 - c^2 x^2)}{2c}$$

input `Integrate[a + b*ArcTanh[c*x],x]`

output `a*x + b*x*ArcTanh[c*x] + (b*Log[1 - c^2*x^2])/(2*c)`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{arctanh}(cx)) dx$$

$$\downarrow \text{2009}$$

$$ax + b \operatorname{arctanh}(cx) + \frac{b \log(1 - c^2 x^2)}{2c}$$

input `Int[a + b*ArcTanh[c*x],x]`

output `a*x + b*x*ArcTanh[c*x] + (b*Log[1 - c^2*x^2])/(2*c)`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.97

method	result	size
default	$ax + bx \operatorname{arctanh}(cx) + \frac{b \ln(-c^2x^2+1)}{2c}$	29
parts	$ax + bx \operatorname{arctanh}(cx) + \frac{b \ln(-c^2x^2+1)}{2c}$	29
derivativedivides	$\frac{cxa+b \left(cx \operatorname{arctanh}(cx) + \frac{\ln(-c^2x^2+1)}{2} \right)}{c}$	33
parallelrisch	$-\frac{b(-cx \operatorname{arctanh}(cx) - \ln(cx-1) - \operatorname{arctanh}(cx))}{c} + ax$	34
risch	$ax + \frac{bx \ln(cx+1)}{2} - \frac{b \ln(-cx+1)x}{2} + \frac{b \ln(c^2x^2-1)}{2c}$	42

input `int(a+b*arctanh(c*x),x,method=_RETURNVERBOSE)`output `a*x+b*x*arctanh(c*x)+1/2*b*ln(-c^2*x^2+1)/c`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.40

$$\int (a + b \operatorname{arctanh}(cx)) dx = \frac{bcx \log\left(-\frac{cx+1}{cx-1}\right) + 2acx + b \log(c^2x^2 - 1)}{2c}$$

input `integrate(a+b*arctanh(c*x),x, algorithm="fricas")`output `1/2*(b*c*x*log(-(c*x + 1)/(c*x - 1)) + 2*a*c*x + b*log(c^2*x^2 - 1))/c`

Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.90

$$\int (a + \operatorname{barctanh}(cx)) dx = ax + b \begin{cases} x \operatorname{atanh}(cx) + \frac{\log(cx+1)}{c} - \frac{\operatorname{atanh}(cx)}{c} & \text{for } c \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

input `integrate(a+b*atanh(c*x),x)`

output `a*x + b*Piecewise((x*atanh(c*x) + log(c*x + 1)/c - atanh(c*x)/c, Ne(c, 0)), (0, True))`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int (a + \operatorname{barctanh}(cx)) dx = ax + \frac{(2cx \operatorname{artanh}(cx) + \log(-c^2x^2 + 1))b}{2c}$$

input `integrate(a+b*arctanh(c*x),x, algorithm="maxima")`

output `a*x + 1/2*(2*c*x*arctanh(c*x) + log(-c^2*x^2 + 1))*b/c`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 156 vs. 2(28) = 56.

Time = 0.12 (sec) , antiderivative size = 156, normalized size of antiderivative = 5.20

$$\int (a + \operatorname{barctanh}(cx)) dx = bc \left(\frac{\log\left(\frac{|-cx-1|}{|cx-1|}\right)}{c^2} - \frac{\log\left(\left|-\frac{cx+1}{cx-1} + 1\right|\right)}{c^2} + \frac{\log\left(\frac{\frac{c\left(\frac{cx+1}{cx-1}+1\right)}{\frac{(cx+1)c}{cx-1}-c}+1}{-\frac{c\left(\frac{cx+1}{cx-1}+1\right)}{\frac{(cx+1)c}{cx-1}-c}-1}\right)}{c^2\left(\frac{cx+1}{cx-1}-1\right)} \right) + ax$$

input `integrate(a+b*arctanh(c*x),x, algorithm="giac")`

output `b*c*(log(abs(-c*x - 1)/abs(c*x - 1))/c^2 - log(abs(-(c*x + 1)/(c*x - 1) + 1))/c^2 + log(-(c*((c*x + 1)/(c*x - 1) + 1)/((c*x + 1)*c/(c*x - 1) - c) + 1)/(c*((c*x + 1)/(c*x - 1) + 1)/((c*x + 1)*c/(c*x - 1) - c) - 1))/c^2*((c*x + 1)/(c*x - 1) - 1)) + a*x`

Mupad [B] (verification not implemented)

Time = 3.67 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.90

$$\int (a + b \operatorname{arctanh}(cx)) dx = ax + \frac{b \ln(c^2 x^2 - 1)}{2c} + bx \operatorname{atanh}(cx)$$

input `int(a + b*atanh(c*x),x)`

output `a*x + (b*log(c^2*x^2 - 1))/(2*c) + b*x*atanh(c*x)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.17

$$\int (a + b \operatorname{arctanh}(cx)) dx = \frac{\operatorname{atanh}(cx) bcx + \operatorname{atanh}(cx) b + \log(c^2 x - c) b + acx}{c}$$

input `int(a+b*atanh(c*x),x)`

output `(atanh(c*x)*b*c*x + atanh(c*x)*b + log(c**2*x - c)*b + a*c*x)/c`

3.7 $\int \frac{a+b\operatorname{arctanh}(cx)}{x} dx$

Optimal result	149
Mathematica [A] (verified)	149
Rubi [A] (verified)	150
Maple [A] (verified)	151
Fricas [F]	151
Sympy [F]	151
Maxima [F]	152
Giac [F]	152
Mupad [F(-1)]	152
Reduce [F]	153

Optimal result

Integrand size = 12, antiderivative size = 26

$$\int \frac{a + b\operatorname{arctanh}(cx)}{x} dx = a \log(x) - \frac{1}{2}b \operatorname{PolyLog}(2, -cx) + \frac{1}{2}b \operatorname{PolyLog}(2, cx)$$

output `a*ln(x)-1/2*b*polylog(2,-c*x)+1/2*b*polylog(2,c*x)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \frac{a + b\operatorname{arctanh}(cx)}{x} dx = a \log(x) + \frac{1}{2}b(-\operatorname{PolyLog}(2, -cx) + \operatorname{PolyLog}(2, cx))$$

input `Integrate[(a + b*ArcTanh[c*x])/x,x]`

output `a*Log[x] + (b*(-PolyLog[2, -(c*x)] + PolyLog[2, c*x]))/2`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x} dx$$

↓ 6446

$$a \log(x) - \frac{1}{2}b \operatorname{PolyLog}(2, -cx) + \frac{1}{2}b \operatorname{PolyLog}(2, cx)$$

input `Int[(a + b*ArcTanh[c*x])/x,x]`

output `a*Log[x] - (b*PolyLog[2, -(c*x)])/2 + (b*PolyLog[2, c*x])/2`

Defintions of rubi rules used

rule 6446 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))/(x_), x_Symbol] := Simp[a*Log[x], x] + (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) / ; FreeQ[{a, b, c}, x]`

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

method	result	size
risch	$a \ln(-cx) + \frac{b \operatorname{dilog}(-cx+1)}{2} - \frac{b \operatorname{dilog}(cx+1)}{2}$	28
parts	$a \ln(x) + b \left(\ln(cx) \operatorname{arctanh}(cx) - \frac{\operatorname{dilog}(cx+1)}{2} - \frac{\ln(cx) \ln(cx+1)}{2} - \frac{\operatorname{dilog}(cx)}{2} \right)$	44
derivativedivides	$a \ln(cx) + b \left(\ln(cx) \operatorname{arctanh}(cx) - \frac{\operatorname{dilog}(cx+1)}{2} - \frac{\ln(cx) \ln(cx+1)}{2} - \frac{\operatorname{dilog}(cx)}{2} \right)$	46
default	$a \ln(cx) + b \left(\ln(cx) \operatorname{arctanh}(cx) - \frac{\operatorname{dilog}(cx+1)}{2} - \frac{\ln(cx) \ln(cx+1)}{2} - \frac{\operatorname{dilog}(cx)}{2} \right)$	46

input `int((a+b*arctanh(c*x))/x,x,method=_RETURNVERBOSE)`

output `a*ln(-c*x)+1/2*b*dilog(-c*x+1)-1/2*b*dilog(c*x+1)`

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x} dx = \int \frac{b \operatorname{artanh}(cx) + a}{x} dx$$

input `integrate((a+b*arctanh(c*x))/x,x, algorithm="fricas")`

output `integral((b*arctanh(c*x) + a)/x, x)`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x} dx = \int \frac{a + b \operatorname{atanh}(cx)}{x} dx$$

input `integrate((a+b*atanh(c*x))/x,x)`

output `Integral((a + b*atanh(c*x))/x, x)`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x} dx = \int \frac{b \operatorname{artanh}(cx) + a}{x} dx$$

input `integrate((a+b*arctanh(c*x))/x,x, algorithm="maxima")`

output `1/2*b*integrate((log(c*x + 1) - log(-c*x + 1))/x, x) + a*log(x)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x} dx = \int \frac{b \operatorname{artanh}(cx) + a}{x} dx$$

input `integrate((a+b*arctanh(c*x))/x,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x} dx = \int \frac{a + b \operatorname{atanh}(cx)}{x} dx$$

input `int((a + b*atanh(c*x))/x,x)`

output `int((a + b*atanh(c*x))/x, x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x} dx = \left(\int \frac{\operatorname{atanh}(cx)}{x} dx \right) b + \log(x) a$$

input `int((a+b*atanh(c*x))/x,x)`

output `int(atanh(c*x)/x,x)*b + log(x)*a`

3.8 $\int \frac{a+b\operatorname{arctanh}(cx)}{x^2} dx$

Optimal result	154
Mathematica [A] (verified)	154
Rubi [A] (verified)	155
Maple [A] (verified)	156
Fricas [A] (verification not implemented)	157
Sympy [A] (verification not implemented)	157
Maxima [A] (verification not implemented)	158
Giac [B] (verification not implemented)	158
Mupad [B] (verification not implemented)	159
Reduce [B] (verification not implemented)	159

Optimal result

Integrand size = 12, antiderivative size = 36

$$\int \frac{a + b\operatorname{arctanh}(cx)}{x^2} dx = -\frac{a + b\operatorname{arctanh}(cx)}{x} + bc \log(x) - \frac{1}{2}bc \log(1 - c^2x^2)$$

output `-(a+b*arctanh(c*x))/x+b*c*ln(x)-1/2*b*c*ln(-c^2*x^2+1)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.08

$$\int \frac{a + b\operatorname{arctanh}(cx)}{x^2} dx = -\frac{a}{x} - \frac{b\operatorname{arctanh}(cx)}{x} + bc \log(x) - \frac{1}{2}bc \log(1 - c^2x^2)$$

input `Integrate[(a + b*ArcTanh[c*x])/x^2,x]`

output `-(a/x) - (b*ArcTanh[c*x])/x + b*c*Log[x] - (b*c*Log[1 - c^2*x^2])/2`

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.06, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {6452, 243, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}(cx)}{x^2} dx \\
 & \quad \downarrow \text{6452} \\
 & bc \int \frac{1}{x(1-c^2x^2)} dx - \frac{a + b \operatorname{arctanh}(cx)}{x} \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2}bc \int \frac{1}{x^2(1-c^2x^2)} dx^2 - \frac{a + b \operatorname{arctanh}(cx)}{x} \\
 & \quad \downarrow \text{47} \\
 & \frac{1}{2}bc \left(c^2 \int \frac{1}{1-c^2x^2} dx^2 + \int \frac{1}{x^2} dx^2 \right) - \frac{a + b \operatorname{arctanh}(cx)}{x} \\
 & \quad \downarrow \text{14} \\
 & \frac{1}{2}bc \left(c^2 \int \frac{1}{1-c^2x^2} dx^2 + \log(x^2) \right) - \frac{a + b \operatorname{arctanh}(cx)}{x} \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{2}bc (\log(x^2) - \log(1-c^2x^2)) - \frac{a + b \operatorname{arctanh}(cx)}{x}
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c*x])/x^2,x]`

output `-((a + b*ArcTanh[c*x])/x) + (b*c*(Log[x^2] - Log[1 - c^2*x^2]))/2`

Definitions of rubi rules used

rule 14 $\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ ; FreeQ}[a, x]$

rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$

rule 47 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x]$

rule 243 $\text{Int}[(x_)^{(m_)*((a_)+(b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] \text{ ; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$

rule 6452 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)]*(b_)]^{(p_)*(x_)^{(m_)}, x_Symbol] : > \text{Simp}[x^{(m+1)*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{2*n}))}, x], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.17

method	result	size
parallelrisch	$\frac{bc \ln(x)x - \ln(cx-1)xbc - x \operatorname{arctanh}(cx)bc - b \operatorname{arctanh}(cx) - a}{x}$	42
parts	$-\frac{a}{x} + bc \left(-\frac{\operatorname{arctanh}(cx)}{cx} + \ln(cx) - \frac{\ln(cx-1)}{2} - \frac{\ln(cx+1)}{2} \right)$	44
derivativedivides	$c \left(-\frac{a}{cx} + b \left(-\frac{\operatorname{arctanh}(cx)}{cx} + \ln(cx) - \frac{\ln(cx-1)}{2} - \frac{\ln(cx+1)}{2} \right) \right)$	48
default	$c \left(-\frac{a}{cx} + b \left(-\frac{\operatorname{arctanh}(cx)}{cx} + \ln(cx) - \frac{\ln(cx-1)}{2} - \frac{\ln(cx+1)}{2} \right) \right)$	48
risch	$-\frac{b \ln(cx+1)}{2x} + \frac{2bc \ln(x)x - bc \ln(c^2x^2-1)x + b \ln(-cx+1) - 2a}{2x}$	54

input `int((a+b*arctanh(c*x))/x^2,x,method=_RETURNVERBOSE)`

output `(b*c*ln(x)*x-ln(c*x-1)*x*b*c-x*arctanh(c*x)*b*c-b*arctanh(c*x)-a)/x`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.31

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^2} dx = -\frac{bcx \log(c^2x^2 - 1) - 2bcx \log(x) + b \log\left(-\frac{cx+1}{cx-1}\right) + 2a}{2x}$$

input `integrate((a+b*arctanh(c*x))/x^2,x, algorithm="fricas")`

output `-1/2*(b*c*x*log(c^2*x^2 - 1) - 2*b*c*x*log(x) + b*log(-(c*x + 1)/(c*x - 1)) + 2*a)/x`

Sympy [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.14

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^2} dx = \begin{cases} -\frac{a}{x} + bc \log(x) - bc \log\left(x - \frac{1}{c}\right) - bc \operatorname{atanh}(cx) - \frac{b \operatorname{atanh}(cx)}{x} & \text{for } c \neq 0 \\ -\frac{a}{x} & \text{otherwise} \end{cases}$$

input `integrate((a+b*atanh(c*x))/x**2,x)`

output `Piecewise((-a/x + b*c*log(x) - b*c*log(x - 1/c) - b*c*atanh(c*x) - b*atanh(c*x)/x, Ne(c, 0)), (-a/x, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.08

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^2} dx = -\frac{1}{2} \left(c(\log(c^2 x^2 - 1) - \log(x^2)) + \frac{2 \operatorname{arctanh}(cx)}{x} \right) b - \frac{a}{x}$$

input `integrate((a+b*arctanh(c*x))/x^2,x, algorithm="maxima")`

output `-1/2*(c*(log(c^2*x^2 - 1) - log(x^2)) + 2*arctanh(c*x)/x)*b - a/x`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 94 vs. 2(34) = 68.

Time = 0.12 (sec) , antiderivative size = 94, normalized size of antiderivative = 2.61

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^2} dx = \left(b \log \left(-\frac{cx+1}{cx-1} - 1 \right) - b \log \left(-\frac{cx+1}{cx-1} \right) + \frac{b \log \left(-\frac{cx+1}{cx-1} \right)}{\frac{cx+1}{cx-1} + 1} + \frac{2a}{\frac{cx+1}{cx-1} + 1} \right) c$$

input `integrate((a+b*arctanh(c*x))/x^2,x, algorithm="giac")`

output `(b*log(-(c*x + 1)/(c*x - 1) - 1) - b*log(-(c*x + 1)/(c*x - 1)) + b*log(-(c*x + 1)/(c*x - 1))/((c*x + 1)/(c*x - 1) + 1) + 2*a/((c*x + 1)/(c*x - 1) + 1))*c`

Mupad [B] (verification not implemented)

Time = 3.67 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.92

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^2} dx = bc \ln(x) - \frac{a + b \operatorname{atanh}(cx)}{x} - \frac{bc \ln(c^2 x^2 - 1)}{2}$$

input `int((a + b*atanh(c*x))/x^2,x)`output `b*c*log(x) - (a + b*atanh(c*x))/x - (b*c*log(c^2*x^2 - 1))/2`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.25

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^2} dx = \frac{-\operatorname{atanh}(cx)bcx - \operatorname{atanh}(cx)b - \log(c^2x - c)bcx + \log(x)bcx - a}{x}$$

input `int((a+b*atanh(c*x))/x^2,x)`output `(- atanh(c*x)*b*c*x - atanh(c*x)*b - log(c**2*x - c)*b*c*x + log(x)*b*c*x - a)/x`

3.9 $\int \frac{a+b\operatorname{arctanh}(cx)}{x^3} dx$

Optimal result	160
Mathematica [A] (verified)	160
Rubi [A] (verified)	161
Maple [A] (verified)	162
Fricas [A] (verification not implemented)	163
Sympy [A] (verification not implemented)	163
Maxima [A] (verification not implemented)	163
Giac [B] (verification not implemented)	164
Mupad [B] (verification not implemented)	164
Reduce [B] (verification not implemented)	165

Optimal result

Integrand size = 12, antiderivative size = 37

$$\int \frac{a + b\operatorname{arctanh}(cx)}{x^3} dx = -\frac{bc}{2x} + \frac{1}{2}bc^2\operatorname{arctanh}(cx) - \frac{a + b\operatorname{arctanh}(cx)}{2x^2}$$

output

```
-1/2*b*c/x+1/2*b*c^2*arctanh(c*x)-1/2*(a+b*arctanh(c*x))/x^2
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.59

$$\int \frac{a + b\operatorname{arctanh}(cx)}{x^3} dx = -\frac{a}{2x^2} - \frac{bc}{2x} - \frac{b\operatorname{arctanh}(cx)}{2x^2} - \frac{1}{4}bc^2 \log(1-cx) + \frac{1}{4}bc^2 \log(1+cx)$$

input

```
Integrate[(a + b*ArcTanh[c*x])/x^3,x]
```

output

```
-1/2*a/x^2 - (b*c)/(2*x) - (b*ArcTanh[c*x])/(2*x^2) - (b*c^2*Log[1 - c*x])/4 + (b*c^2*Log[1 + c*x])/4
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.92, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6452, 264, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{arctanh}(cx)}{x^3} dx$$

$$\downarrow 6452$$

$$\frac{1}{2}bc \int \frac{1}{x^2(1-c^2x^2)} dx - \frac{a + \operatorname{arctanh}(cx)}{2x^2}$$

$$\downarrow 264$$

$$\frac{1}{2}bc \left(c^2 \int \frac{1}{1-c^2x^2} dx - \frac{1}{x} \right) - \frac{a + \operatorname{arctanh}(cx)}{2x^2}$$

$$\downarrow 219$$

$$\frac{1}{2}bc \left(\operatorname{arctanh}(cx) - \frac{1}{x} \right) - \frac{a + \operatorname{arctanh}(cx)}{2x^2}$$

input `Int[(a + b*ArcTanh[c*x])/x^3,x]`

output `-1/2*(a + b*ArcTanh[c*x])/x^2 + (b*c*(-x^(-1) + c*ArcTanh[c*x]))/2`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 264

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]
```

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.05

method	result	size
parallelrisch	$-\frac{\operatorname{arctanh}(cx)bc^2x^2+ac^2x^2+bcx+b\operatorname{arctanh}(cx)+a}{2x^2}$	39
parts	$-\frac{a}{2x^2} + bc^2\left(-\frac{\operatorname{arctanh}(cx)}{2c^2x^2} - \frac{1}{2cx} - \frac{\ln(cx-1)}{4} + \frac{\ln(cx+1)}{4}\right)$	50
derivativedivides	$c^2\left(-\frac{a}{2c^2x^2} + b\left(-\frac{\operatorname{arctanh}(cx)}{2c^2x^2} - \frac{1}{2cx} - \frac{\ln(cx-1)}{4} + \frac{\ln(cx+1)}{4}\right)\right)$	54
default	$c^2\left(-\frac{a}{2c^2x^2} + b\left(-\frac{\operatorname{arctanh}(cx)}{2c^2x^2} - \frac{1}{2cx} - \frac{\ln(cx-1)}{4} + \frac{\ln(cx+1)}{4}\right)\right)$	54
risch	$-\frac{b\ln(cx+1)}{4x^2} + \frac{bc^2\ln(-cx-1)x^2-x^2b\ln(-cx+1)c^2-2bcx+b\ln(-cx+1)-2a}{4x^2}$	68
orering	$\frac{(2c^2x^3-2x)(a+b\operatorname{arctanh}(cx))}{x^3} + \frac{(cx-1)(cx+1)x^2\left(\frac{bc}{(-c^2x^2+1)x^3} - \frac{3(a+b\operatorname{arctanh}(cx))}{x^4}\right)}{2}$	73

input

```
int((a+b*arctanh(c*x))/x^3,x,method=_RETURNVERBOSE)
```

output

```
-1/2*(-arctanh(c*x)*b*c^2*x^2+a*c^2*x^2+b*c*x+b*arctanh(c*x)+a)/x^2
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.16

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^3} dx = -\frac{2bcx - (bc^2x^2 - b) \log\left(-\frac{cx+1}{cx-1}\right) + 2a}{4x^2}$$

input `integrate((a+b*arctanh(c*x))/x^3,x, algorithm="fricas")`output `-1/4*(2*b*c*x - (b*c^2*x^2 - b)*log(-(c*x + 1)/(c*x - 1)) + 2*a)/x^2`**Sympy [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.97

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^3} dx = -\frac{a}{2x^2} + \frac{bc^2 \operatorname{atanh}(cx)}{2} - \frac{bc}{2x} - \frac{b \operatorname{atanh}(cx)}{2x^2}$$

input `integrate((a+b*atanh(c*x))/x**3,x)`output `-a/(2*x**2) + b*c**2*atanh(c*x)/2 - b*c/(2*x) - b*atanh(c*x)/(2*x**2)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.22

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^3} dx = \frac{1}{4} \left(\left(c \log(cx + 1) - c \log(cx - 1) - \frac{2}{x} \right) c - \frac{2 \operatorname{artanh}(cx)}{x^2} \right) b - \frac{a}{2x^2}$$

input `integrate((a+b*arctanh(c*x))/x^3,x, algorithm="maxima")`output `1/4*((c*log(c*x + 1) - c*log(c*x - 1) - 2/x)*c - 2*arctanh(c*x)/x^2)*b - 1/2*a/x^2`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 135 vs. $2(31) = 62$.

Time = 0.12 (sec) , antiderivative size = 135, normalized size of antiderivative = 3.65

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^3} dx = \left(\frac{(cx+1)bc \log\left(-\frac{cx+1}{cx-1}\right)}{(cx-1)\left(\frac{(cx+1)^2}{(cx-1)^2} + \frac{2(cx+1)}{cx-1} + 1\right)} + \frac{\frac{2(cx+1)ac}{cx-1} + \frac{(cx+1)bc}{cx-1} + bc}{\frac{(cx+1)^2}{(cx-1)^2} + \frac{2(cx+1)}{cx-1} + 1} \right) c$$

input `integrate((a+b*arctanh(c*x))/x^3,x, algorithm="giac")`

output `((c*x + 1)*b*c*log(-(c*x + 1)/(c*x - 1))/((c*x - 1)*((c*x + 1)^2/(c*x - 1)^2 + 2*(c*x + 1)/(c*x - 1) + 1)) + (2*(c*x + 1)*a*c/(c*x - 1) + (c*x + 1)*b*c/(c*x - 1) + b*c)/((c*x + 1)^2/(c*x - 1)^2 + 2*(c*x + 1)/(c*x - 1) + 1)*c`

Mupad [B] (verification not implemented)

Time = 3.75 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.24

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^3} dx = \frac{bc \operatorname{atan}\left(\frac{c^2 x}{\sqrt{-c^2}}\right) \sqrt{-c^2}}{2} - \frac{a}{2} + \frac{b \operatorname{atanh}(cx)}{2} + \frac{bcx}{2x^2}$$

input `int((a + b*atanh(c*x))/x^3,x)`

output `(b*c*atan((c^2*x)/(-c^2)^(1/2))*(-c^2)^(1/2))/2 - (a/2 + (b*atanh(c*x))/2 + (b*c*x)/2)/x^2`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.89

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^3} dx = \frac{\operatorname{atanh}(cx) b c^2 x^2 - \operatorname{atanh}(cx) b - a - bcx}{2x^2}$$

input `int((a+b*atanh(c*x))/x^3,x)`

output `(atanh(c*x)*b*c**2*x**2 - atanh(c*x)*b - a - b*c*x)/(2*x**2)`

3.10 $\int \frac{a+b\operatorname{arctanh}(cx)}{x^4} dx$

Optimal result	166
Mathematica [A] (verified)	166
Rubi [A] (verified)	167
Maple [A] (verified)	168
Fricas [A] (verification not implemented)	169
Sympy [A] (verification not implemented)	169
Maxima [A] (verification not implemented)	170
Giac [B] (verification not implemented)	170
Mupad [B] (verification not implemented)	171
Reduce [B] (verification not implemented)	171

Optimal result

Integrand size = 12, antiderivative size = 54

$$\int \frac{a + b\operatorname{arctanh}(cx)}{x^4} dx = -\frac{bc}{6x^2} - \frac{a + b\operatorname{arctanh}(cx)}{3x^3} + \frac{1}{3}bc^3 \log(x) - \frac{1}{6}bc^3 \log(1 - c^2x^2)$$

output
$$-1/6*b*c/x^2-1/3*(a+b*\operatorname{arctanh}(c*x))/x^3+1/3*b*c^3*\ln(x)-1/6*b*c^3*\ln(-c^2*x^2+1)$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.09

$$\int \frac{a + b\operatorname{arctanh}(cx)}{x^4} dx = -\frac{a}{3x^3} - \frac{bc}{6x^2} - \frac{b\operatorname{arctanh}(cx)}{3x^3} + \frac{1}{3}bc^3 \log(x) - \frac{1}{6}bc^3 \log(1 - c^2x^2)$$

input
$$\operatorname{Integrate}[(a + b*\operatorname{ArcTanh}[c*x])/x^4, x]$$

output
$$-1/3*a/x^3 - (b*c)/(6*x^2) - (b*\operatorname{ArcTanh}[c*x])/(3*x^3) + (b*c^3*\operatorname{Log}[x])/3 - (b*c^3*\operatorname{Log}[1 - c^2*x^2])/6$$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6452, 243, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{arctanh}(cx)}{x^4} dx$$

$$\downarrow 6452$$

$$\frac{1}{3}bc \int \frac{1}{x^3(1-c^2x^2)} dx - \frac{a + \operatorname{arctanh}(cx)}{3x^3}$$

$$\downarrow 243$$

$$\frac{1}{6}bc \int \frac{1}{x^4(1-c^2x^2)} dx^2 - \frac{a + \operatorname{arctanh}(cx)}{3x^3}$$

$$\downarrow 54$$

$$\frac{1}{6}bc \int \left(-\frac{c^4}{c^2x^2-1} + \frac{c^2}{x^2} + \frac{1}{x^4} \right) dx^2 - \frac{a + \operatorname{arctanh}(cx)}{3x^3}$$

$$\downarrow 2009$$

$$\frac{1}{6}bc \left(c^2 \log(x^2) - c^2 \log(1-c^2x^2) - \frac{1}{x^2} \right) - \frac{a + \operatorname{arctanh}(cx)}{3x^3}$$

input `Int[(a + b*ArcTanh[c*x])/x^4,x]`

output `-1/3*(a + b*ArcTanh[c*x])/x^3 + (b*c*(-x^(-2) + c^2*Log[x^2] - c^2*Log[1 - c^2*x^2]))/6`

Defintions of rubi rules used

rule 54 $\text{Int}[(a + b \cdot x)^m \cdot (c + d \cdot x)^n, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b \cdot x)^m \cdot (c + d \cdot x)^n, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{ILtQ}[m, 0] \&\& \text{IntegerQ}[n] \&\& \text{!(IGtQ}[n, 0] \&\& \text{LtQ}[m + n + 2, 0])$

rule 243 $\text{Int}[x^{(m-1)/2} \cdot (a + b \cdot x)^p, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2} \cdot (a + b \cdot x)^p, x], x, x^2], x] /; \text{FreeQ}[\{a, b, m, p\}, x] \&\& \text{IntegerQ}[(m-1)/2]$

rule 2009 $\text{Int}[u, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 6452 $\text{Int}[(a + \text{ArcTanh}[c \cdot x]^n) \cdot (b \cdot x)^m, x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x]^n)^{p/(m+1)}, x] - \text{Simp}[b \cdot c \cdot n \cdot (p/(m+1)) \text{ Int}[x^{m+n} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x]^n)^{p-1} / (1 - c^2 \cdot x^{2n}), x], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \&\& \text{IGtQ}[p, 0] \&\& (\text{EqQ}[p, 1] \|\| (\text{EqQ}[n, 1] \&\& \text{IntegerQ}[m])) \&\& \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.04

method	result	size
parts	$-\frac{a}{3x^3} + b c^3 \left(-\frac{\text{arctanh}(cx)}{3c^3 x^3} - \frac{\ln(cx+1)}{6} - \frac{1}{6c^2 x^2} + \frac{\ln(cx)}{3} - \frac{\ln(cx-1)}{6} \right)$	56
derivativedivides	$c^3 \left(-\frac{a}{3c^3 x^3} + b \left(-\frac{\text{arctanh}(cx)}{3c^3 x^3} - \frac{\ln(cx+1)}{6} - \frac{1}{6c^2 x^2} + \frac{\ln(cx)}{3} - \frac{\ln(cx-1)}{6} \right) \right)$	60
default	$c^3 \left(-\frac{a}{3c^3 x^3} + b \left(-\frac{\text{arctanh}(cx)}{3c^3 x^3} - \frac{\ln(cx+1)}{6} - \frac{1}{6c^2 x^2} + \frac{\ln(cx)}{3} - \frac{\ln(cx-1)}{6} \right) \right)$	60
risch	$-\frac{b \ln(cx+1)}{6x^3} + \frac{2b c^3 \ln(x)x^3 - b c^3 \ln(c^2 x^2 - 1)x^3 - bcx + b \ln(-cx+1) - 2a}{6x^3}$	67
parallelrisc	$\frac{2b c^3 \ln(x)x^3 - 2 \ln(cx-1)x^3 b c^3 - 2b \text{arctanh}(cx)x^3 c^3 - b c^3 x^3 - bcx - 2b \text{arctanh}(cx) - 2a}{6x^3}$	70

input $\text{int}((a+b \cdot \text{arctanh}(c \cdot x))/x^4, x, \text{method}=_RETURNVERBOSE)$

output

```
-1/3*a/x^3+b*c^3*(-1/3/c^3/x^3*arctanh(c*x)-1/6*ln(c*x+1)-1/6/c^2/x^2+1/3*
ln(c*x)-1/6*ln(c*x-1))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.09

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^4} dx$$

$$= -\frac{bc^3 x^3 \log(c^2 x^2 - 1) - 2bc^3 x^3 \log(x) + bcx + b \log\left(-\frac{cx+1}{cx-1}\right) + 2a}{6x^3}$$

input

```
integrate((a+b*arctanh(c*x))/x^4,x, algorithm="fricas")
```

output

```
-1/6*(b*c^3*x^3*log(c^2*x^2 - 1) - 2*b*c^3*x^3*log(x) + b*c*x + b*log(-(c*
x + 1)/(c*x - 1)) + 2*a)/x^3
```

Sympy [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.30

$$\int \frac{a + b \operatorname{atanh}(cx)}{x^4} dx$$

$$= \begin{cases} -\frac{a}{3x^3} + \frac{bc^3 \log(x)}{3} - \frac{bc^3 \log\left(\frac{x-1}{c}\right)}{3} - \frac{bc^3 \operatorname{atanh}(cx)}{3} - \frac{bc}{6x^2} - \frac{b \operatorname{atanh}(cx)}{3x^3} & \text{for } c \neq 0 \\ -\frac{a}{3x^3} & \text{otherwise} \end{cases}$$

input

```
integrate((a+b*atanh(c*x))/x**4,x)
```

output

```
Piecewise((-a/(3*x**3) + b*c**3*log(x)/3 - b*c**3*log(x - 1/c)/3 - b*c**3*
atanh(c*x)/3 - b*c/(6*x**2) - b*atanh(c*x)/(3*x**3), Ne(c, 0)), (-a/(3*x**
3), True))
```

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.91

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^4} dx$$

$$= -\frac{1}{6} \left(\left(c^2 \log(c^2 x^2 - 1) - c^2 \log(x^2) + \frac{1}{x^2} \right) c + \frac{2 \operatorname{artanh}(cx)}{x^3} \right) b - \frac{a}{3x^3}$$

input `integrate((a+b*arctanh(c*x))/x^4,x, algorithm="maxima")`

output `-1/6*((c^2*log(c^2*x^2 - 1) - c^2*log(x^2) + 1/x^2)*c + 2*arctanh(c*x)/x^3)*b - 1/3*a/x^3`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 251 vs. $2(46) = 92$.

Time = 0.12 (sec) , antiderivative size = 251, normalized size of antiderivative = 4.65

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^4} dx$$

$$= \frac{1}{3} \left(bc^2 \log\left(-\frac{cx+1}{cx-1} - 1\right) - bc^2 \log\left(-\frac{cx+1}{cx-1}\right) + \frac{\left(\frac{3(cx+1)^2 bc^2}{(cx-1)^2} + bc^2\right) \log\left(-\frac{cx+1}{cx-1}\right)}{\frac{(cx+1)^3}{(cx-1)^3} + \frac{3(cx+1)^2}{(cx-1)^2} + \frac{3(cx+1)}{cx-1} + 1} + \frac{2 \left(\frac{3(cx+1)^2 ac^2}{(cx-1)^2} + a\right)}{\frac{(cx+1)^3}{(cx-1)^3} + \frac{3(cx+1)^2}{(cx-1)^2} + \frac{3(cx+1)}{cx-1} + 1} \right)$$

input `integrate((a+b*arctanh(c*x))/x^4,x, algorithm="giac")`

output `1/3*(b*c^2*log(-(c*x + 1)/(c*x - 1) - 1) - b*c^2*log(-(c*x + 1)/(c*x - 1)) + (3*(c*x + 1)^2*b*c^2/(c*x - 1)^2 + b*c^2)*log(-(c*x + 1)/(c*x - 1))/((c*x + 1)^3/(c*x - 1)^3 + 3*(c*x + 1)^2/(c*x - 1)^2 + 3*(c*x + 1)/(c*x - 1) + 1) + 2*(3*(c*x + 1)^2*a*c^2/(c*x - 1)^2 + a*c^2 + (c*x + 1)^2*b*c^2/(c*x - 1)^2 + (c*x + 1)*b*c^2/(c*x - 1))/((c*x + 1)^3/(c*x - 1)^3 + 3*(c*x + 1)^2/(c*x - 1)^2 + 3*(c*x + 1)/(c*x - 1) + 1))*c`

Mupad [B] (verification not implemented)

Time = 3.73 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.85

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^4} dx = \frac{b c^3 \ln(x)}{3} - \frac{b c^3 \ln(c^2 x^2 - 1)}{6} - \frac{\frac{a}{3} + \frac{b \operatorname{atanh}(cx)}{3} + \frac{bcx}{6}}{x^3}$$

input `int((a + b*atanh(c*x))/x^4,x)`output `(b*c^3*log(x))/3 - (b*c^3*log(c^2*x^2 - 1))/6 - (a/3 + (b*atanh(c*x))/3 + (b*c*x)/6)/x^3`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.19

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^4} dx = \frac{-2 \operatorname{atanh}(cx) b c^3 x^3 - 2 \operatorname{atanh}(cx) b - 2 \log(c^2 x - c) b c^3 x^3 + 2 \log(x) b c^3 x^3 - 2a - bcx}{6x^3}$$

input `int((a+b*atanh(c*x))/x^4,x)`output `(- 2*atanh(c*x)*b*c**3*x**3 - 2*atanh(c*x)*b - 2*log(c**2*x - c)*b*c**3*x**3 + 2*log(x)*b*c**3*x**3 - 2*a - b*c*x)/(6*x**3)`

3.11 $\int \frac{a+b\operatorname{arctanh}(cx)}{x^5} dx$

Optimal result	172
Mathematica [A] (verified)	172
Rubi [A] (verified)	173
Maple [A] (verified)	174
Fricas [A] (verification not implemented)	175
Sympy [A] (verification not implemented)	175
Maxima [A] (verification not implemented)	176
Giac [B] (verification not implemented)	176
Mupad [B] (verification not implemented)	177
Reduce [B] (verification not implemented)	177

Optimal result

Integrand size = 12, antiderivative size = 48

$$\int \frac{a + b\operatorname{arctanh}(cx)}{x^5} dx = -\frac{bc}{12x^3} - \frac{bc^3}{4x} + \frac{1}{4}bc^4\operatorname{arctanh}(cx) - \frac{a + b\operatorname{arctanh}(cx)}{4x^4}$$

output `-1/12*b*c/x^3-1/4*b*c^3/x+1/4*b*c^4*arctanh(c*x)-1/4*(a+b*arctanh(c*x))/x^4`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.46

$$\int \frac{a + b\operatorname{arctanh}(cx)}{x^5} dx = -\frac{a}{4x^4} - \frac{bc}{12x^3} - \frac{bc^3}{4x} - \frac{b\operatorname{arctanh}(cx)}{4x^4} - \frac{1}{8}bc^4 \log(1 - cx) + \frac{1}{8}bc^4 \log(1 + cx)$$

input `Integrate[(a + b*ArcTanh[c*x])/x^5,x]`

output `-1/4*a/x^4 - (b*c)/(12*x^3) - (b*c^3)/(4*x) - (b*ArcTanh[c*x])/(4*x^4) - (b*c^4*Log[1 - c*x])/8 + (b*c^4*Log[1 + c*x])/8`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.96, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6452, 264, 264, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{arctanh}(cx)}{x^5} dx \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{4}bc \int \frac{1}{x^4(1-c^2x^2)} dx - \frac{a + \operatorname{arctanh}(cx)}{4x^4} \\
 & \quad \downarrow \text{264} \\
 & \frac{1}{4}bc \left(c^2 \int \frac{1}{x^2(1-c^2x^2)} dx - \frac{1}{3x^3} \right) - \frac{a + \operatorname{arctanh}(cx)}{4x^4} \\
 & \quad \downarrow \text{264} \\
 & \frac{1}{4}bc \left(c^2 \left(c^2 \int \frac{1}{1-c^2x^2} dx - \frac{1}{x} \right) - \frac{1}{3x^3} \right) - \frac{a + \operatorname{arctanh}(cx)}{4x^4} \\
 & \quad \downarrow \text{219} \\
 & \frac{1}{4}bc \left(c^2 \left(\operatorname{arctanh}(cx) - \frac{1}{x} \right) - \frac{1}{3x^3} \right) - \frac{a + \operatorname{arctanh}(cx)}{4x^4}
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c*x])/x^5,x]`

output `-1/4*(a + b*ArcTanh[c*x])/x^4 + (b*c*(-1/3*1/x^3 + c^2*(-x^(-1) + c*ArcTanh[c*x])))/4`

Defintions of rubi rules used

```
rule 219 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 264 Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(
m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + 2*p + 3)/(a*c
^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p
}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]
```

```
rule 6452 Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.90

method	result	size
parallelrisch	$-\frac{-3b \operatorname{arctanh}(cx)x^4c^4+3bc^3x^3+bcx+3b \operatorname{arctanh}(cx)+3a}{12x^4}$	43
parts	$-\frac{a}{4x^4} + bc^4 \left(-\frac{\operatorname{arctanh}(cx)}{4c^4x^4} + \frac{\ln(cx+1)}{8} - \frac{\ln(cx-1)}{8} - \frac{1}{12c^3x^3} - \frac{1}{4cx} \right)$	58
derivativedivides	$c^4 \left(-\frac{a}{4c^4x^4} + b \left(-\frac{\operatorname{arctanh}(cx)}{4c^4x^4} + \frac{\ln(cx+1)}{8} - \frac{\ln(cx-1)}{8} - \frac{1}{12c^3x^3} - \frac{1}{4cx} \right) \right)$	62
default	$c^4 \left(-\frac{a}{4c^4x^4} + b \left(-\frac{\operatorname{arctanh}(cx)}{4c^4x^4} + \frac{\ln(cx+1)}{8} - \frac{\ln(cx-1)}{8} - \frac{1}{12c^3x^3} - \frac{1}{4cx} \right) \right)$	62
risch	$-\frac{b \ln(cx+1)}{8x^4} + \frac{3bc^4 \ln(-cx-1)x^4 - 3bx^4 \ln(-cx+1)c^4 - 6bc^3x^3 - 2bcx + 3b \ln(-cx+1) - 6a}{24x^4}$	79
orering	$\frac{\left(\frac{3}{2}c^4x^5 - \frac{5}{6}c^2x^3 - \frac{2}{3}x\right)(a+b \operatorname{arctanh}(cx))}{x^5} + \frac{(3c^2x^2+1)(cx-1)(cx+1)x^2 \left(\frac{bc}{(-c^2x^2+1)x^5} - \frac{5(a+b \operatorname{arctanh}(cx))}{x^6} \right)}{12}$	91

```
input int((a+b*arctanh(c*x))/x^5,x,method=_RETURNVERBOSE)
```

output

$$-1/12*(-3*b*\operatorname{arctanh}(c*x)*x^4*c^4+3*b*c^3*x^3+b*c*x+3*b*\operatorname{arctanh}(c*x)+3*a)/x^4$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.08

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^5} dx = -\frac{6bc^3x^3 + 2bcx - 3(bc^4x^4 - b) \log\left(-\frac{cx+1}{cx-1}\right) + 6a}{24x^4}$$

input

```
integrate((a+b*arctanh(c*x))/x^5,x, algorithm="fricas")
```

output

$$-1/24*(6*b*c^3*x^3 + 2*b*c*x - 3*(b*c^4*x^4 - b)*\log(-(c*x + 1)/(c*x - 1)) + 6*a)/x^4$$

Sympy [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.96

$$\int \frac{a + b \operatorname{atanh}(cx)}{x^5} dx = -\frac{a}{4x^4} + \frac{bc^4 \operatorname{atanh}(cx)}{4} - \frac{bc^3}{4x} - \frac{bc}{12x^3} - \frac{b \operatorname{atanh}(cx)}{4x^4}$$

input

```
integrate((a+b*atanh(c*x))/x**5,x)
```

output

$$-a/(4*x**4) + b*c**4*atanh(c*x)/4 - b*c**3/(4*x) - b*c/(12*x**3) - b*atanh(c*x)/(4*x**4)$$

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.25

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^5} dx$$

$$= \frac{1}{24} \left(\left(3c^3 \log(cx+1) - 3c^3 \log(cx-1) - \frac{2(3c^2x^2+1)}{x^3} \right) c - \frac{6 \operatorname{arctanh}(cx)}{x^4} \right) b - \frac{a}{4x^4}$$

input `integrate((a+b*arctanh(c*x))/x^5,x, algorithm="maxima")`

output `1/24*((3*c^3*log(c*x + 1) - 3*c^3*log(c*x - 1) - 2*(3*c^2*x^2 + 1)/x^3)*c - 6*arctanh(c*x)/x^4)*b - 1/4*a/x^4`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 292 vs. 2(40) = 80.

Time = 0.13 (sec) , antiderivative size = 292, normalized size of antiderivative = 6.08

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^5} dx$$

$$= \frac{1}{3} c \left(\frac{3 \left(\frac{(cx+1)^3 bc^3}{(cx-1)^3} + \frac{(cx+1)bc^3}{cx-1} \right) \log\left(-\frac{cx+1}{cx-1}\right)}{\frac{(cx+1)^4}{(cx-1)^4} + \frac{4(cx+1)^3}{(cx-1)^3} + \frac{6(cx+1)^2}{(cx-1)^2} + \frac{4(cx+1)}{cx-1} + 1} + \frac{\frac{6(cx+1)^3 ac^3}{(cx-1)^3} + \frac{6(cx+1)ac^3}{cx-1} + \frac{3(cx+1)^3 bc^3}{(cx-1)^3} + \frac{6(cx+1)^2 bc^3}{(cx-1)^2} + \frac{5(cx+1)bc^3}{cx-1}}{\frac{(cx+1)^4}{(cx-1)^4} + \frac{4(cx+1)^3}{(cx-1)^3} + \frac{6(cx+1)^2}{(cx-1)^2} + \frac{4(cx+1)}{cx-1} + 1} \right)$$

input `integrate((a+b*arctanh(c*x))/x^5,x, algorithm="giac")`

output `1/3*c*(3*((c*x + 1)^3*b*c^3/(c*x - 1)^3 + (c*x + 1)*b*c^3/(c*x - 1))*log(-(c*x + 1)/(c*x - 1))/((c*x + 1)^4/(c*x - 1)^4 + 4*(c*x + 1)^3/(c*x - 1)^3 + 6*(c*x + 1)^2/(c*x - 1)^2 + 4*(c*x + 1)/(c*x - 1) + 1) + (6*(c*x + 1)^3*a*c^3/(c*x - 1)^3 + 6*(c*x + 1)*a*c^3/(c*x - 1) + 3*(c*x + 1)^3*b*c^3/(c*x - 1)^3 + 6*(c*x + 1)^2*b*c^3/(c*x - 1)^2 + 5*(c*x + 1)*b*c^3/(c*x - 1) + 2*b*c^3)/((c*x + 1)^4/(c*x - 1)^4 + 4*(c*x + 1)^3/(c*x - 1)^3 + 6*(c*x + 1)^2/(c*x - 1)^2 + 4*(c*x + 1)/(c*x - 1) + 1))`

Mupad [B] (verification not implemented)

Time = 3.99 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.23

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^5} dx = \frac{b \ln(1 - cx)}{8x^4} - \frac{b \ln(cx + 1)}{8x^4} - \frac{bc^3 x^3 + \frac{bcx}{3} + a}{4x^4} - \frac{bc^4 \operatorname{atan}(cx) \operatorname{li}}{4}$$

input `int((a + b*atanh(c*x))/x^5,x)`output `(b*log(1 - c*x))/(8*x^4) - (b*c^4*atan(c*x*1i)*1i)/4 - (b*log(c*x + 1))/(8*x^4) - (a + b*c^3*x^3 + (b*c*x)/3)/(4*x^4)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.90

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^5} dx = \frac{3 \operatorname{atanh}(cx) b c^4 x^4 - 3 \operatorname{atanh}(cx) b - 3a - 3b c^3 x^3 - bcx}{12x^4}$$

input `int((a+b*atanh(c*x))/x^5,x)`output `(3*atanh(c*x)*b*c**4*x**4 - 3*atanh(c*x)*b - 3*a - 3*b*c**3*x**3 - b*c*x)/(12*x**4)`

3.12 $\int \frac{a+b\operatorname{arctanh}(cx)}{x^6} dx$

Optimal result	178
Mathematica [A] (verified)	178
Rubi [A] (verified)	179
Maple [A] (verified)	180
Fricas [A] (verification not implemented)	181
Sympy [A] (verification not implemented)	181
Maxima [A] (verification not implemented)	182
Giac [B] (verification not implemented)	182
Mupad [B] (verification not implemented)	183
Reduce [B] (verification not implemented)	183

Optimal result

Integrand size = 12, antiderivative size = 65

$$\int \frac{a + b\operatorname{arctanh}(cx)}{x^6} dx = -\frac{bc}{20x^4} - \frac{bc^3}{10x^2} - \frac{a + b\operatorname{arctanh}(cx)}{5x^5} + \frac{1}{5}bc^5 \log(x) - \frac{1}{10}bc^5 \log(1 - c^2x^2)$$

output

```
-1/20*b*c/x^4-1/10*b*c^3/x^2-1/5*(a+b*arctanh(c*x))/x^5+1/5*b*c^5*ln(x)-1/10*b*c^5*ln(-c^2*x^2+1)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.08

$$\int \frac{a + b\operatorname{arctanh}(cx)}{x^6} dx = -\frac{a}{5x^5} - \frac{bc}{20x^4} - \frac{bc^3}{10x^2} - \frac{b\operatorname{arctanh}(cx)}{5x^5} + \frac{1}{5}bc^5 \log(x) - \frac{1}{10}bc^5 \log(1 - c^2x^2)$$

input

```
Integrate[(a + b*ArcTanh[c*x])/x^6,x]
```

output

$$-1/5*a/x^5 - (b*c)/(20*x^4) - (b*c^3)/(10*x^2) - (b*ArcTanh[c*x])/(5*x^5) + (b*c^5*Log[x])/5 - (b*c^5*Log[1 - c^2*x^2])/10$$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.95, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6452, 243, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}(cx)}{x^6} dx \\ & \quad \downarrow \text{6452} \\ & \frac{1}{5}bc \int \frac{1}{x^5(1-c^2x^2)} dx - \frac{a + b \operatorname{arctanh}(cx)}{5x^5} \\ & \quad \downarrow \text{243} \\ & \frac{1}{10}bc \int \frac{1}{x^6(1-c^2x^2)} dx^2 - \frac{a + b \operatorname{arctanh}(cx)}{5x^5} \\ & \quad \downarrow \text{54} \\ & \frac{1}{10}bc \int \left(-\frac{c^6}{c^2x^2-1} + \frac{c^4}{x^2} + \frac{c^2}{x^4} + \frac{1}{x^6} \right) dx^2 - \frac{a + b \operatorname{arctanh}(cx)}{5x^5} \\ & \quad \downarrow \text{2009} \\ & \frac{1}{10}bc \left(c^4 \log(x^2) - \frac{c^2}{x^2} - c^4 \log(1-c^2x^2) - \frac{1}{2x^4} \right) - \frac{a + b \operatorname{arctanh}(cx)}{5x^5} \end{aligned}$$

input

$$\text{Int}[(a + b*ArcTanh[c*x])/x^6, x]$$

output

$$-1/5*(a + b*ArcTanh[c*x])/x^5 + (b*c*(-1/2*1/x^4 - c^2/x^2 + c^4*Log[x^2] - c^4*Log[1 - c^2*x^2]))/10$$

Defintions of rubi rules used

rule 54 $\text{Int}[(a + b \cdot x)^m \cdot (c + d \cdot x)^n, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b \cdot x)^m \cdot (c + d \cdot x)^n, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{ILtQ}[m, 0] \&\& \text{IntegerQ}[n] \&\& \text{!(IGtQ}[n, 0] \&\& \text{LtQ}[m + n + 2, 0])$

rule 243 $\text{Int}[x^{(m-1)/2} \cdot (a + b \cdot x)^p, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2} \cdot (a + b \cdot x)^p, x], x, x^2], x] /; \text{FreeQ}[\{a, b, m, p\}, x] \&\& \text{IntegerQ}[(m-1)/2]$

rule 2009 $\text{Int}[u, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 6452 $\text{Int}[(a + \text{ArcTanh}[c \cdot x]^n) \cdot (b \cdot x)^m, x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x]^n)^{p/(m+1)}, x] - \text{Simp}[b \cdot c \cdot n \cdot (p/(m+1)) \text{ Int}[x^{m+n} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x]^n)^{p-1} / (1 - c^2 \cdot x^{2n}), x], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \&\& \text{IGtQ}[p, 0] \&\& (\text{EqQ}[p, 1] \|\| (\text{EqQ}[n, 1] \&\& \text{IntegerQ}[m])) \&\& \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.98

method	result	size
parts	$-\frac{a}{5x^5} + b c^5 \left(-\frac{\text{arctanh}(cx)}{5c^5 x^5} - \frac{1}{20c^4 x^4} - \frac{1}{10c^2 x^2} + \frac{\ln(cx)}{5} - \frac{\ln(cx+1)}{10} - \frac{\ln(cx-1)}{10} \right)$	64
derivativedivides	$c^5 \left(-\frac{a}{5c^5 x^5} + b \left(-\frac{\text{arctanh}(cx)}{5c^5 x^5} - \frac{1}{20c^4 x^4} - \frac{1}{10c^2 x^2} + \frac{\ln(cx)}{5} - \frac{\ln(cx+1)}{10} - \frac{\ln(cx-1)}{10} \right) \right)$	68
default	$c^5 \left(-\frac{a}{5c^5 x^5} + b \left(-\frac{\text{arctanh}(cx)}{5c^5 x^5} - \frac{1}{20c^4 x^4} - \frac{1}{10c^2 x^2} + \frac{\ln(cx)}{5} - \frac{\ln(cx+1)}{10} - \frac{\ln(cx-1)}{10} \right) \right)$	68
risch	$-\frac{b \ln(cx+1)}{10x^5} + \frac{4b c^5 \ln(x)x^5 - 2b c^5 \ln(c^2 x^2 - 1)x^5 - 2b c^3 x^3 - bcx + 2b \ln(-cx+1) - 4a}{20x^5}$	77
parallelrisc	$\frac{4b c^5 \ln(x)x^5 - 4 \ln(cx-1)x^5 b c^5 - 4b \text{arctanh}(cx)x^5 c^5 - 2b c^5 x^5 - 2b c^3 x^3 - bcx - 4b \text{arctanh}(cx) - 4a}{20x^5}$	79

input $\text{int}((a+b \cdot \text{arctanh}(c \cdot x))/x^6, x, \text{method}=_RETURNVERBOSE)$

output

```
-1/5*a/x^5+b*c^5*(-1/5/c^5/x^5*arctanh(c*x)-1/20/c^4/x^4-1/10/c^2/x^2+1/5*
ln(c*x)-1/10*ln(c*x+1)-1/10*ln(c*x-1))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.08

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^6} dx$$

$$= -\frac{2bc^5x^5 \log(c^2x^2 - 1) - 4bc^5x^5 \log(x) + 2bc^3x^3 + bcx + 2b \log\left(-\frac{cx+1}{cx-1}\right) + 4a}{20x^5}$$

input

```
integrate((a+b*arctanh(c*x))/x^6,x, algorithm="fricas")
```

output

```
-1/20*(2*b*c^5*x^5*log(c^2*x^2 - 1) - 4*b*c^5*x^5*log(x) + 2*b*c^3*x^3 + b
*c*x + 2*b*log(-(c*x + 1)/(c*x - 1)) + 4*a)/x^5
```

Sympy [A] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.23

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^6} dx$$

$$= \begin{cases} -\frac{a}{5x^5} + \frac{bc^5 \log(x)}{5} - \frac{bc^5 \log\left(x - \frac{1}{c}\right)}{5} - \frac{bc^5 \operatorname{atanh}(cx)}{5} - \frac{bc^3}{10x^2} - \frac{bc}{20x^4} - \frac{b \operatorname{atanh}(cx)}{5x^5} & \text{for } c \neq 0 \\ -\frac{a}{5x^5} & \text{otherwise} \end{cases}$$

input

```
integrate((a+b*atanh(c*x))/x**6,x)
```

output

```
Piecewise((-a/(5*x**5) + b*c**5*log(x)/5 - b*c**5*log(x - 1/c)/5 - b*c**5*
atanh(c*x)/5 - b*c**3/(10*x**2) - b*c/(20*x**4) - b*atanh(c*x)/(5*x**5), N
e(c, 0)), (-a/(5*x**5), True))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.94

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^6} dx$$

$$= -\frac{1}{20} \left(\left(2c^4 \log(c^2x^2 - 1) - 2c^4 \log(x^2) + \frac{2c^2x^2 + 1}{x^4} \right) c + \frac{4 \operatorname{artanh}(cx)}{x^5} \right) b - \frac{a}{5x^5}$$

input `integrate((a+b*arctanh(c*x))/x^6,x, algorithm="maxima")`

output `-1/20*((2*c^4*log(c^2*x^2 - 1) - 2*c^4*log(x^2) + (2*c^2*x^2 + 1)/x^4)*c + 4*arctanh(c*x)/x^5)*b - 1/5*a/x^5`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 397 vs. 2(55) = 110.

Time = 0.13 (sec) , antiderivative size = 397, normalized size of antiderivative = 6.11

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^6} dx$$

$$= \frac{1}{5} \left(bc^4 \log \left(-\frac{cx+1}{cx-1} - 1 \right) - bc^4 \log \left(-\frac{cx+1}{cx-1} \right) + \frac{\left(\frac{5(cx+1)^4 bc^4}{(cx-1)^4} + \frac{10(cx+1)^2 bc^4}{(cx-1)^2} + bc^4 \right) \log \left(-\frac{cx+1}{cx-1} \right)}{\frac{(cx+1)^5}{(cx-1)^5} + \frac{5(cx+1)^4}{(cx-1)^4} + \frac{10(cx+1)^3}{(cx-1)^3} + \frac{10(cx+1)^2}{(cx-1)^2} + \frac{5(cx+1)}{cx-1}} \right) - \frac{a}{5x^5}$$

input `integrate((a+b*arctanh(c*x))/x^6,x, algorithm="giac")`

output `1/5*(b*c^4*log(-(c*x + 1)/(c*x - 1) - 1) - b*c^4*log(-(c*x + 1)/(c*x - 1)) + (5*(c*x + 1)^4*b*c^4/(c*x - 1)^4 + 10*(c*x + 1)^2*b*c^4/(c*x - 1)^2 + b*c^4)*log(-(c*x + 1)/(c*x - 1))/(c*x + 1)^5/(c*x - 1)^5 + 5*(c*x + 1)^4/(c*x - 1)^4 + 10*(c*x + 1)^3/(c*x - 1)^3 + 10*(c*x + 1)^2/(c*x - 1)^2 + 5*(c*x + 1)/(c*x - 1) + 1) + 2*(5*(c*x + 1)^4*a*c^4/(c*x - 1)^4 + 10*(c*x + 1)^2*a*c^4/(c*x - 1)^2 + a*c^4 + 2*(c*x + 1)^4*b*c^4/(c*x - 1)^4 + 4*(c*x + 1)^3*b*c^4/(c*x - 1)^3 + 4*(c*x + 1)^2*b*c^4/(c*x - 1)^2 + 2*(c*x + 1)*b*c^4/(c*x - 1))/(c*x + 1)^5/(c*x - 1)^5 + 5*(c*x + 1)^4/(c*x - 1)^4 + 10*(c*x + 1)^3/(c*x - 1)^3 + 10*(c*x + 1)^2/(c*x - 1)^2 + 5*(c*x + 1)/(c*x - 1) + 1))*c`

Mupad [B] (verification not implemented)

Time = 3.82 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.09

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^6} dx = \frac{b c^5 \ln(x)}{5} - \frac{b c^5 \ln(c^2 x^2 - 1)}{10} - \frac{\frac{b c^3 x^3}{2} + \frac{b c x}{4} + a}{5 x^5} - \frac{b \ln(cx + 1)}{10 x^5} + \frac{b \ln(1 - cx)}{10 x^5}$$

input `int((a + b*atanh(c*x))/x^6,x)`output `(b*c^5*log(x))/5 - (b*c^5*log(c^2*x^2 - 1))/10 - (a + (b*c^3*x^3)/2 + (b*c*x)/4)/(5*x^5) - (b*log(c*x + 1))/(10*x^5) + (b*log(1 - c*x))/(10*x^5)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.12

$$\int \frac{a + b \operatorname{arctanh}(cx)}{x^6} dx = \frac{-4 \operatorname{atanh}(cx) b c^5 x^5 - 4 \operatorname{atanh}(cx) b - 4 \log(c^2 x - c) b c^5 x^5 + 4 \log(x) b c^5 x^5 - 4a - 2b c^3 x^3 - bcx}{20x^5}$$

input `int((a+b*atanh(c*x))/x^6,x)`output `(- 4*atanh(c*x)*b*c**5*x**5 - 4*atanh(c*x)*b - 4*log(c**2*x - c)*b*c**5*x**5 + 4*log(x)*b*c**5*x**5 - 4*a - 2*b*c**3*x**3 - b*c*x)/(20*x**5)`

3.13 $\int x^5(a + b \operatorname{arctanh}(cx))^2 dx$

Optimal result	184
Mathematica [A] (verified)	185
Rubi [A] (verified)	185
Maple [A] (verified)	189
Fricas [A] (verification not implemented)	190
Sympy [A] (verification not implemented)	190
Maxima [A] (verification not implemented)	191
Giac [B] (verification not implemented)	191
Mupad [B] (verification not implemented)	192
Reduce [B] (verification not implemented)	193

Optimal result

Integrand size = 14, antiderivative size = 145

$$\int x^5(a + b \operatorname{arctanh}(cx))^2 dx = \frac{abx}{3c^5} + \frac{4b^2x^2}{45c^4} + \frac{b^2x^4}{60c^2} + \frac{b^2x \operatorname{arctanh}(cx)}{3c^5} + \frac{bx^3(a + b \operatorname{arctanh}(cx))}{9c^3} + \frac{bx^5(a + b \operatorname{arctanh}(cx))}{15c} - \frac{(a + b \operatorname{arctanh}(cx))^2}{6c^6} + \frac{1}{6}x^6(a + b \operatorname{arctanh}(cx))^2 + \frac{23b^2 \log(1 - c^2x^2)}{90c^6}$$

output

```
1/3*a*b*x/c^5+4/45*b^2*x^2/c^4+1/60*b^2*x^4/c^2+1/3*b^2*x*arctanh(c*x)/c^5
+1/9*b*x^3*(a+b*arctanh(c*x))/c^3+1/15*b*x^5*(a+b*arctanh(c*x))/c-1/6*(a+b
*arctanh(c*x))^2/c^6+1/6*x^6*(a+b*arctanh(c*x))^2+23/90*b^2*ln(-c^2*x^2+1)
/c^6
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.13

$$\int x^5(a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{60abcx + 16b^2c^2x^2 + 20abc^3x^3 + 3b^2c^4x^4 + 12abc^5x^5 + 30a^2c^6x^6 + 4bcx(15ac^5x^5 + b(15 + 5c^2x^2 + 3c^4x^4)) \operatorname{ArcTanh}[cx] + 30b^2(-1 + c^6x^6) \operatorname{ArcTanh}[cx]^2 + 2b(15a + 23b) \operatorname{Log}[1 - cx] - 30ab \operatorname{Log}[1 + cx] + 46b^2 \operatorname{Log}[1 + cx]}{(180c^6)}$$

input `Integrate[x^5*(a + b*ArcTanh[c*x])^2,x]`

output `(60*a*b*c*x + 16*b^2*c^2*x^2 + 20*a*b*c^3*x^3 + 3*b^2*c^4*x^4 + 12*a*b*c^5*x^5 + 30*a^2*c^6*x^6 + 4*b*c*x*(15*a*c^5*x^5 + b*(15 + 5*c^2*x^2 + 3*c^4*x^4))*ArcTanh[c*x] + 30*b^2*(-1 + c^6*x^6)*ArcTanh[c*x]^2 + 2*b*(15*a + 23*b)*Log[1 - c*x] - 30*a*b*Log[1 + c*x] + 46*b^2*Log[1 + c*x])/(180*c^6)`

Rubi [A] (verified)

Time = 1.32 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.41, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {6452, 6542, 6452, 243, 49, 2009, 6542, 6452, 243, 49, 2009, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^5(a + b \operatorname{arctanh}(cx))^2 dx$$

$$\downarrow 6452$$

$$\frac{1}{6}x^6(a + b \operatorname{arctanh}(cx))^2 - \frac{1}{3}bc \int \frac{x^6(a + b \operatorname{arctanh}(cx))}{1 - c^2x^2} dx$$

$$\downarrow 6542$$

$$\frac{1}{6}x^6(a + b \operatorname{arctanh}(cx))^2 - \frac{1}{3}bc \left(\frac{\int \frac{x^4(a + b \operatorname{arctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\int x^4(a + b \operatorname{arctanh}(cx)) dx}{c^2} \right)$$

$$\downarrow 6452$$

$$\begin{aligned}
& \frac{1}{6}x^6(a + \operatorname{barctanh}(cx))^2 - \\
& \frac{1}{3}bc \left(\frac{\int \frac{x^4(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{5}x^5(a + \operatorname{barctanh}(cx)) - \frac{1}{5}bc \int \frac{x^5}{1 - c^2x^2} dx}{c^2} \right) \\
& \quad \downarrow 243 \\
& \frac{1}{6}x^6(a + \operatorname{barctanh}(cx))^2 - \\
& \frac{1}{3}bc \left(\frac{\int \frac{x^4(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{5}x^5(a + \operatorname{barctanh}(cx)) - \frac{1}{10}bc \int \frac{x^4}{1 - c^2x^2} dx^2}{c^2} \right) \\
& \quad \downarrow 49 \\
& \frac{1}{6}x^6(a + \operatorname{barctanh}(cx))^2 - \\
& \frac{1}{3}bc \left(\frac{\int \frac{x^4(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{5}x^5(a + \operatorname{barctanh}(cx)) - \frac{1}{10}bc \int \left(-\frac{x^2}{c^2} - \frac{1}{c^4(c^2x^2 - 1)} - \frac{1}{c^4} \right) dx^2}{c^2} \right) \\
& \quad \downarrow 2009 \\
& \frac{1}{6}x^6(a + \operatorname{barctanh}(cx))^2 - \\
& \frac{1}{3}bc \left(\frac{\int \frac{x^4(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{5}x^5(a + \operatorname{barctanh}(cx)) - \frac{1}{10}bc \left(-\frac{x^2}{c^4} - \frac{x^4}{2c^2} - \frac{\log(1 - c^2x^2)}{c^6} \right)}{c^2} \right) \\
& \quad \downarrow 6542 \\
& \frac{1}{6}x^6(a + \operatorname{barctanh}(cx))^2 - \\
& \frac{1}{3}bc \left(\frac{\frac{\int \frac{x^2(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\int \frac{x^2(a + \operatorname{barctanh}(cx))}{c^2} dx}{c^2} - \frac{\frac{1}{5}x^5(a + \operatorname{barctanh}(cx)) - \frac{1}{10}bc \left(-\frac{x^2}{c^4} - \frac{x^4}{2c^2} - \frac{\log(1 - c^2x^2)}{c^6} \right)}{c^2}}{c^2} \right) \\
& \quad \downarrow 6452 \\
& \frac{1}{6}x^6(a + \operatorname{barctanh}(cx))^2 - \\
& \frac{1}{3}bc \left(\frac{\frac{\int \frac{x^2(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a + \operatorname{barctanh}(cx)) - \frac{1}{3}bc \int \frac{x^3}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{5}x^5(a + \operatorname{barctanh}(cx)) - \frac{1}{10}bc \left(-\frac{x^2}{c^4} - \frac{x^4}{2c^2} - \frac{\log(1 - c^2x^2)}{c^6} \right)}{c^2}}{c^2} \right) \\
& \quad \downarrow 243
\end{aligned}$$

$$\frac{1}{3}bc \left(\frac{\frac{\int \frac{x^2(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx)) - \frac{1}{6}bc \int \frac{x^2}{1-c^2x^2} dx^2}{c^2}}{c^2} - \frac{\frac{1}{5}x^5(a+b\operatorname{arctanh}(cx)) - \frac{1}{10}bc \left(-\frac{x^2}{c^4} - \frac{x^4}{2c^2} \right)}{c^2} \right)$$

↓ 49

$$\frac{1}{3}bc \left(\frac{\frac{\int \frac{x^2(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx)) - \frac{1}{6}bc \int \left(-\frac{1}{c^2} - \frac{1}{c^2(c^2x^2-1)} \right) dx^2}{c^2}}{c^2} - \frac{\frac{1}{5}x^5(a+b\operatorname{arctanh}(cx)) - \frac{1}{10}bc \left(-\frac{x^2}{c^4} - \frac{x^4}{2c^2} \right)}{c^2} \right)$$

↓ 2009

$$\frac{1}{3}bc \left(\frac{\frac{\int \frac{x^2(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx)) - \frac{1}{6}bc \left(-\frac{x^2}{c^2} - \frac{\log(1-c^2x^2)}{c^4} \right)}{c^2}}{c^2} - \frac{\frac{1}{5}x^5(a+b\operatorname{arctanh}(cx)) - \frac{1}{10}bc \left(-\frac{x^2}{c^4} - \frac{x^4}{2c^2} \right)}{c^2} \right)$$

↓ 6542

$$\frac{1}{3}bc \left(\frac{\frac{\frac{\int \frac{a+b\operatorname{arctanh}(cx)}{1-c^2x^2} dx}{c^2} - \frac{\int (a+b\operatorname{arctanh}(cx)) dx}{c^2}}{c^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx)) - \frac{1}{6}bc \left(-\frac{x^2}{c^2} - \frac{\log(1-c^2x^2)}{c^4} \right)}{c^2}}{c^2} - \frac{\frac{1}{5}x^5(a+b\operatorname{arctanh}(cx)) - \frac{1}{10}bc \left(-\frac{x^2}{c^4} - \frac{x^4}{2c^2} \right)}{c^2} \right)$$

↓ 2009

$$\frac{1}{3}bc \left(\frac{\frac{\frac{\int \frac{a+b\operatorname{arctanh}(cx)}{1-c^2x^2} dx}{c^2} - \frac{ax+b\operatorname{arctanh}(cx) + \frac{b \log(1-c^2x^2)}{2c}}{c^2}}{c^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx)) - \frac{1}{6}bc \left(-\frac{x^2}{c^2} - \frac{\log(1-c^2x^2)}{c^4} \right)}{c^2}}{c^2} - \frac{\frac{1}{5}x^5(a+b\operatorname{arctanh}(cx)) - \frac{1}{10}bc \left(-\frac{x^2}{c^4} - \frac{x^4}{2c^2} \right)}{c^2} \right)$$

↓ 6510

$$\frac{1}{6}x^6(a + b\operatorname{arctanh}(cx))^2 - \frac{1}{3}bc \left(\frac{\frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^3} - \frac{ax+b\operatorname{arctanh}(cx) + \frac{b\log(1-c^2x^2)}{2c}}{c^2}}{c^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx)) - \frac{1}{6}bc \left(-\frac{x^2}{c^2} - \frac{\log(1-c^2x^2)}{c^4} \right)}{c^2} \right) - \frac{1}{5}x^5(a + b\operatorname{arctanh}(cx))$$

input `Int[x^5*(a + b*ArcTanh[c*x])^2,x]`

output `(x^6*(a + b*ArcTanh[c*x])^2)/6 - (b*c*(-(((x^5*(a + b*ArcTanh[c*x])))/5 - (b*c*(-(x^2/c^4) - x^4/(2*c^2) - Log[1 - c^2*x^2]/c^6))/10)/c^2) + (-(((x^3*(a + b*ArcTanh[c*x])))/3 - (b*c*(-(x^2/c^2) - Log[1 - c^2*x^2]/c^4))/6)/c^2) + ((a + b*ArcTanh[c*x])^2/(2*b*c^3) - (a*x + b*x*ArcTanh[c*x] + (b*Log[1 - c^2*x^2])/(2*c))/c^2)/c^2)/c^2)/3`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

```
rule 6510 Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol]
:> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x]
&& EqQ[c^2*d + e, 0] && NeQ[p, -1]
```

```
rule 6542 Int[(((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)*((f_.)*(x_)^(m_)))/((d_) + (e_.)*(x_)^2), x_Symbol]
:> Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]
```

Maple [A] (verified)

Time = 0.59 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.26

method	result
parallelrisch	$30b^2 \operatorname{arctanh}(cx)^2 x^6 c^6 + 60ab \operatorname{arctanh}(cx) x^6 c^6 + 30a^2 c^6 x^6 + 12b^2 \operatorname{arctanh}(cx) x^5 c^5 + 12ab c^5 x^5 + 3b^2 c^4 x^4 + 20b^2 \operatorname{arctanh}(cx) x^3 c^3 + 20ab c^3 x^3 + 10a^2 c^2 x^2 + 10b^2 \operatorname{arctanh}(cx) x c + 10ab c \operatorname{arctanh}(cx) + 10a^2 \operatorname{arctanh}(cx) + 10b^2 \operatorname{arctanh}(cx) \ln(cx-1) - \operatorname{arctanh}(cx) \ln(cx+1)$
parts	$\frac{a^2 x^6}{6} + \frac{b^2 \left(\frac{c^6 x^6 \operatorname{arctanh}(cx)^2}{6} + \frac{c^5 x^5 \operatorname{arctanh}(cx)}{15} + \frac{c^3 x^3 \operatorname{arctanh}(cx)}{9} + \frac{cx \operatorname{arctanh}(cx)}{3} + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{6} - \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{6} \right)}{6}$
derivativedivides	$\frac{a^2 c^6 x^6}{6} + b^2 \left(\frac{c^6 x^6 \operatorname{arctanh}(cx)^2}{6} + \frac{c^5 x^5 \operatorname{arctanh}(cx)}{15} + \frac{c^3 x^3 \operatorname{arctanh}(cx)}{9} + \frac{cx \operatorname{arctanh}(cx)}{3} + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{6} - \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{6} \right)$
default	$\frac{a^2 c^6 x^6}{6} + b^2 \left(\frac{c^6 x^6 \operatorname{arctanh}(cx)^2}{6} + \frac{c^5 x^5 \operatorname{arctanh}(cx)}{15} + \frac{c^3 x^3 \operatorname{arctanh}(cx)}{9} + \frac{cx \operatorname{arctanh}(cx)}{3} + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{6} - \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{6} \right)$
risch	$\frac{b^2 (c^6 x^6 - 1) \ln(cx+1)^2}{24c^6} + \frac{b(-15b x^6 \ln(-cx+1) c^6 + 30a c^6 x^6 + 6b c^5 x^5 + 10b c^3 x^3 + 30bcx + 15b \ln(-cx+1)) \ln(cx+1)}{180c^6}$

```
input int(x^5*(a+b*arctanh(c*x))^2,x,method=_RETURNVERBOSE)
```

```
output 1/180*(30*b^2*arctanh(c*x)^2*x^6*c^6+60*a*b*arctanh(c*x)*x^6*c^6+30*a^2*c^6*x^6+12*b^2*arctanh(c*x)*x^5*c^5+12*a*b*c^5*x^5+3*b^2*c^4*x^4+20*b^2*arctanh(c*x)*x^3*c^3+20*a*b*c^3*x^3+16*b^2*c^2*x^2+60*b^2*arctanh(c*x)*x*c+60*a*b*c*x-30*b^2*arctanh(c*x)^2+92*ln(c*x-1)*b^2-60*arctanh(c*x)*a*b+92*arctanh(c*x)*b^2+16*b^2)/c^6
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.33

$$\int x^5(a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{60 a^2 c^6 x^6 + 24 a b c^5 x^5 + 6 b^2 c^4 x^4 + 40 a b c^3 x^3 + 32 b^2 c^2 x^2 + 120 a b c x + 15 (b^2 c^6 x^6 - b^2) \log\left(-\frac{cx+1}{cx-1}\right)^2 - 4}{c^6}$$

input `integrate(x^5*(a+b*arctanh(c*x))^2,x, algorithm="fricas")`output
$$\frac{1}{360} \cdot (60 a^2 c^6 x^6 + 24 a b c^5 x^5 + 6 b^2 c^4 x^4 + 40 a b c^3 x^3 + 32 b^2 c^2 x^2 + 120 a b c x + 15 (b^2 c^6 x^6 - b^2) \log(-\frac{c x + 1}{c x - 1})^2 - 4 (15 a b - 23 b^2) \log(c x + 1) + 4 (15 a b + 23 b^2) \log(c x - 1) + 4 (15 a b c^6 x^6 + 3 b^2 c^5 x^5 + 5 b^2 c^3 x^3 + 15 b^2 c x) \log(-\frac{c x + 1}{c x - 1})) / c^6$$
Sympy [A] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.46

$$\int x^5(a + b \operatorname{atanh}(cx))^2 dx$$

$$= \begin{cases} \frac{a^2 x^6}{6} + \frac{a b x^6 \operatorname{atanh}(cx)}{3} + \frac{a b x^5}{15 c} + \frac{a b x^3}{9 c^3} + \frac{a b x}{3 c^5} - \frac{a b \operatorname{atanh}(cx)}{3 c^6} + \frac{b^2 x^6 \operatorname{atanh}^2(cx)}{6} + \frac{b^2 x^5 \operatorname{atanh}(cx)}{15 c} + \frac{b^2 x^4}{60 c^2} + \frac{b^2 x^3 \operatorname{atanh}(cx)}{9 c^3} \\ \frac{a^2 x^6}{6} \end{cases}$$

input `integrate(x**5*(a+b*atanh(c*x))**2,x)`output `Piecewise((a**2*x**6/6 + a*b*x**6*atanh(c*x)/3 + a*b*x**5/(15*c) + a*b*x**3/(9*c**3) + a*b*x/(3*c**5) - a*b*atanh(c*x)/(3*c**6) + b**2*x**6*atanh(c*x)**2/6 + b**2*x**5*atanh(c*x)/(15*c) + b**2*x**4/(60*c**2) + b**2*x**3*atanh(c*x)/(9*c**3) + 4*b**2*x**2/(45*c**4) + b**2*x*atanh(c*x)/(3*c**5) + 23*b**2*log(x - 1/c)/(45*c**6) - b**2*atanh(c*x)**2/(6*c**6) + 23*b**2*atanh(c*x)/(45*c**6), Ne(c, 0)), (a**2*x**6/6, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.48

$$\int x^5(a + b \operatorname{arctanh}(cx))^2 dx = \frac{1}{6} b^2 x^6 \operatorname{arctanh}(cx)^2 + \frac{1}{6} a^2 x^6 + \frac{1}{90} \left(30 x^6 \operatorname{arctanh}(cx) + c \left(\frac{2(3c^4 x^5 + 5c^2 x^3 + 15x)}{c^6} - \frac{15 \log(cx+1)}{c^7} + \frac{15 \log(cx-1)}{c^7} \right) \right) ab + \frac{1}{360} \left(4c \left(\frac{2(3c^4 x^5 + 5c^2 x^3 + 15x)}{c^6} - \frac{15 \log(cx+1)}{c^7} + \frac{15 \log(cx-1)}{c^7} \right) \operatorname{arctanh}(cx) + \frac{6c^4 x^4 + 32c^2 x^2 - 2(15 \log(cx-1) - 46) \log(cx+1) + 15 \log(cx+1)^2 + 15 \log(cx-1)^2 + 92 \log(cx-1)}{c^6} \right) b^2$$

input `integrate(x^5*(a+b*arctanh(c*x))^2,x, algorithm="maxima")`

output `1/6*b^2*x^6*arctanh(c*x)^2 + 1/6*a^2*x^6 + 1/90*(30*x^6*arctanh(c*x) + c*(2*(3*c^4*x^5 + 5*c^2*x^3 + 15*x)/c^6 - 15*log(c*x + 1)/c^7 + 15*log(c*x - 1)/c^7))*a*b + 1/360*(4*c*(2*(3*c^4*x^5 + 5*c^2*x^3 + 15*x)/c^6 - 15*log(c*x + 1)/c^7 + 15*log(c*x - 1)/c^7)*arctanh(c*x) + (6*c^4*x^4 + 32*c^2*x^2 - 2*(15*log(c*x - 1) - 46)*log(c*x + 1) + 15*log(c*x + 1)^2 + 15*log(c*x - 1)^2 + 92*log(c*x - 1))/c^6)*b^2`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 889 vs. $2(127) = 254$.

Time = 0.14 (sec) , antiderivative size = 889, normalized size of antiderivative = 6.13

$$\int x^5(a + b \operatorname{arctanh}(cx))^2 dx = \text{Too large to display}$$

input `integrate(x^5*(a+b*arctanh(c*x))^2,x, algorithm="giac")`

output

```

1/90*(15*(3*(c*x + 1)^5*b^2/(c*x - 1)^5 + 10*(c*x + 1)^3*b^2/(c*x - 1)^3 +
  3*(c*x + 1)*b^2/(c*x - 1))*log(-(c*x + 1)/(c*x - 1))^2/((c*x + 1)^6*c^7/(
c*x - 1)^6 - 6*(c*x + 1)^5*c^7/(c*x - 1)^5 + 15*(c*x + 1)^4*c^7/(c*x - 1)^
4 - 20*(c*x + 1)^3*c^7/(c*x - 1)^3 + 15*(c*x + 1)^2*c^7/(c*x - 1)^2 - 6*(c
*x + 1)*c^7/(c*x - 1) + c^7) + 2*(90*(c*x + 1)^5*a*b/(c*x - 1)^5 + 300*(c*
x + 1)^3*a*b/(c*x - 1)^3 + 90*(c*x + 1)*a*b/(c*x - 1) + 45*(c*x + 1)^5*b^2
/(c*x - 1)^5 - 135*(c*x + 1)^4*b^2/(c*x - 1)^4 + 230*(c*x + 1)^3*b^2/(c*x
- 1)^3 - 210*(c*x + 1)^2*b^2/(c*x - 1)^2 + 93*(c*x + 1)*b^2/(c*x - 1) - 23
*b^2)*log(-(c*x + 1)/(c*x - 1))/((c*x + 1)^6*c^7/(c*x - 1)^6 - 6*(c*x + 1)
^5*c^7/(c*x - 1)^5 + 15*(c*x + 1)^4*c^7/(c*x - 1)^4 - 20*(c*x + 1)^3*c^7/(
c*x - 1)^3 + 15*(c*x + 1)^2*c^7/(c*x - 1)^2 - 6*(c*x + 1)*c^7/(c*x - 1) +
c^7) + 4*(45*(c*x + 1)^5*a^2/(c*x - 1)^5 + 150*(c*x + 1)^3*a^2/(c*x - 1)^3
+ 45*(c*x + 1)*a^2/(c*x - 1) + 45*(c*x + 1)^5*a*b/(c*x - 1)^5 - 135*(c*x
+ 1)^4*a*b/(c*x - 1)^4 + 230*(c*x + 1)^3*a*b/(c*x - 1)^3 - 210*(c*x + 1)^2
*a*b/(c*x - 1)^2 + 93*(c*x + 1)*a*b/(c*x - 1) - 23*a*b + 11*(c*x + 1)^5*b^
2/(c*x - 1)^5 - 38*(c*x + 1)^4*b^2/(c*x - 1)^4 + 54*(c*x + 1)^3*b^2/(c*x -
1)^3 - 38*(c*x + 1)^2*b^2/(c*x - 1)^2 + 11*(c*x + 1)*b^2/(c*x - 1))/((c*x
+ 1)^6*c^7/(c*x - 1)^6 - 6*(c*x + 1)^5*c^7/(c*x - 1)^5 + 15*(c*x + 1)^4*c
^7/(c*x - 1)^4 - 20*(c*x + 1)^3*c^7/(c*x - 1)^3 + 15*(c*x + 1)^2*c^7/(c*x
- 1)^2 - 6*(c*x + 1)*c^7/(c*x - 1) + c^7) - 46*b^2*log(-(c*x + 1)/(c*x ...

```

Mupad [B] (verification not implemented)

Time = 3.97 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.18

$$\int x^5 (a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{46 b^2 \ln(c^2 x^2 - 1) - 30 b^2 \operatorname{atanh}(cx)^2 + 30 a^2 c^6 x^6 + 16 b^2 c^2 x^2 + 3 b^2 c^4 x^4 - 60 a b \operatorname{atanh}(cx) + 20 b^2 c^2 x^2}{180 c^6}$$

input

```
int(x^5*(a + b*atanh(c*x))^2,x)
```

output

```

(46*b^2*log(c^2*x^2 - 1) - 30*b^2*atanh(c*x)^2 + 30*a^2*c^6*x^6 + 16*b^2*c
^2*x^2 + 3*b^2*c^4*x^4 - 60*a*b*atanh(c*x) + 20*b^2*c^3*x^3*atanh(c*x) + 1
2*b^2*c^5*x^5*atanh(c*x) + 60*b^2*c*x*atanh(c*x) + 30*b^2*c^6*x^6*atanh(c*
x)^2 + 20*a*b*c^3*x^3 + 12*a*b*c^5*x^5 + 60*a*b*c*x + 60*a*b*c^6*x^6*atanh
(c*x))/(180*c^6)

```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.24

$$\int x^5 (a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{30 \operatorname{atanh}(cx)^2 b^2 c^6 x^6 - 30 \operatorname{atanh}(cx)^2 b^2 + 60 \operatorname{atanh}(cx) ab c^6 x^6 - 60 \operatorname{atanh}(cx) ab + 12 \operatorname{atanh}(cx) b^2 c^5 x^5 + \dots}{180 c^6}$$

input `int(x^5*(a+b*atanh(c*x))^2,x)`output `(30*atanh(c*x)**2*b**2*c**6*x**6 - 30*atanh(c*x)**2*b**2 + 60*atanh(c*x)*a*b*c**6*x**6 - 60*atanh(c*x)*a*b + 12*atanh(c*x)*b**2*c**5*x**5 + 20*atanh(c*x)*b**2*c**3*x**3 + 60*atanh(c*x)*b**2*c*x + 92*atanh(c*x)*b**2 + 92*log(c**2*x - c)*b**2 + 30*a**2*c**6*x**6 + 12*a*b*c**5*x**5 + 20*a*b*c**3*x**3 + 60*a*b*c*x + 3*b**2*c**4*x**4 + 16*b**2*c**2*x**2)/(180*c**6)`

3.14 $\int x^4(a + \operatorname{arctanh}(cx))^2 dx$

Optimal result	194
Mathematica [A] (verified)	195
Rubi [A] (verified)	195
Maple [A] (verified)	199
Fricas [F]	200
Sympy [F]	201
Maxima [F]	201
Giac [F]	202
Mupad [F(-1)]	202
Reduce [F]	202

Optimal result

Integrand size = 14, antiderivative size = 162

$$\begin{aligned} \int x^4(a + \operatorname{arctanh}(cx))^2 dx = & \frac{3b^2x}{10c^4} + \frac{b^2x^3}{30c^2} - \frac{3b^2\operatorname{arctanh}(cx)}{10c^5} \\ & + \frac{bx^2(a + \operatorname{arctanh}(cx))}{5c^3} + \frac{bx^4(a + \operatorname{arctanh}(cx))}{10c} \\ & + \frac{(a + \operatorname{arctanh}(cx))^2}{5c^5} + \frac{1}{5}x^5(a + \operatorname{arctanh}(cx))^2 \\ & - \frac{2b(a + \operatorname{arctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{5c^5} \\ & - \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{5c^5} \end{aligned}$$

output

```
3/10*b^2*x/c^4+1/30*b^2*x^3/c^2-3/10*b^2*arctanh(c*x)/c^5+1/5*b*x^2*(a+b*arctanh(c*x))/c^3+1/10*b*x^4*(a+b*arctanh(c*x))/c+1/5*(a+b*arctanh(c*x))^2/c^5+1/5*x^5*(a+b*arctanh(c*x))^2-2/5*b*(a+b*arctanh(c*x))*ln(2/(-c*x+1))/c^5-1/5*b^2*polylog(2,1-2/(-c*x+1))/c^5
```

Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.99

$$\int x^4(a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{-9ab + 9b^2cx + 6abc^2x^2 + b^2c^3x^3 + 3abc^4x^4 + 6a^2c^5x^5 + 6b^2(-1 + c^5x^5) \operatorname{arctanh}(cx)^2 + 3b \operatorname{arctanh}(cx)}{30c^5}$$

input `Integrate[x^4*(a + b*ArcTanh[c*x])^2,x]`

output `(-9*a*b + 9*b^2*c*x + 6*a*b*c^2*x^2 + b^2*c^3*x^3 + 3*a*b*c^4*x^4 + 6*a^2*c^5*x^5 + 6*b^2*(-1 + c^5*x^5)*ArcTanh[c*x]^2 + 3*b*ArcTanh[c*x]*(4*a*c^5*x^5 + b*(-3 + 2*c^2*x^2 + c^4*x^4) - 4*b*Log[1 + E^(-2*ArcTanh[c*x])])) + 6*a*b*Log[-1 + c^2*x^2] + 6*b^2*PolyLog[2, -E^(-2*ArcTanh[c*x])])/(30*c^5)`

Rubi [A] (verified)

Time = 1.31 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.23, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.929$, Rules used = {6452, 6542, 6452, 254, 2009, 6542, 6452, 262, 219, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4(a + b \operatorname{arctanh}(cx))^2 dx$$

$$\downarrow 6452$$

$$\frac{1}{5}x^5(a + b \operatorname{arctanh}(cx))^2 - \frac{2}{5}bc \int \frac{x^5(a + b \operatorname{arctanh}(cx))}{1 - c^2x^2} dx$$

$$\downarrow 6542$$

$$\frac{1}{5}x^5(a + b \operatorname{arctanh}(cx))^2 - \frac{2}{5}bc \left(\frac{\int \frac{x^3(a + b \operatorname{arctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\int x^3(a + b \operatorname{arctanh}(cx)) dx}{c^2} \right)$$

$$\downarrow 6452$$

$$\begin{aligned}
& \frac{1}{5}x^5(a + \operatorname{barctanh}(cx))^2 - \\
& \frac{2}{5}bc \left(\frac{\int \frac{x^3(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx)) - \frac{1}{4}bc \int \frac{x^4}{1 - c^2x^2} dx}{c^2} \right) \\
& \quad \downarrow 254 \\
& \frac{2}{5}bc \left(\frac{\int \frac{x^3(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx)) - \frac{1}{4}bc \int \left(-\frac{x^2}{c^2} + \frac{1}{c^4(1 - c^2x^2)} - \frac{1}{c^4} \right) dx}{c^2} \right) \\
& \quad \downarrow 2009 \\
& \frac{2}{5}bc \left(\frac{\int \frac{x^3(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx)) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(cx)}{c^5} - \frac{x}{c^4} - \frac{x^3}{3c^2} \right)}{c^2} \right) \\
& \quad \downarrow 6542 \\
& \frac{2}{5}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\int x(a + \operatorname{barctanh}(cx)) dx}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx)) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(cx)}{c^5} - \frac{x}{c^4} - \frac{x^3}{3c^2} \right)}{c^2} \right) \\
& \quad \downarrow 6452 \\
& \frac{2}{5}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx)) - \frac{1}{2}bc \int \frac{x^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx)) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(cx)}{c^5} - \frac{x}{c^4} - \frac{x^3}{3c^2} \right)}{c^2} \right) \\
& \quad \downarrow 262 \\
& \frac{2}{5}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx)) - \frac{1}{2}bc \left(\frac{\int \frac{1}{1 - c^2x^2} dx}{c^2} - \frac{x}{c^2} \right)}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx)) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(cx)}{c^5} - \frac{x}{c^4} - \frac{x^3}{3c^2} \right)}{c^2} \right) \\
& \quad \downarrow 219
\end{aligned}$$

$$\frac{2}{5}bc \left(\frac{\int \frac{x(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a+b\operatorname{arctanh}(cx)) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(cx)}{c^3} - \frac{x}{c^2} \right)}{c^2}}{c^2} - \frac{\frac{1}{4}x^4(a+b\operatorname{arctanh}(cx)) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(cx)}{c^5} - \frac{x}{c^2} \right)}{c^2} \right)$$

6546

$$\frac{2}{5}bc \left(\frac{\int \frac{a+b\operatorname{arctanh}(cx)}{c} dx}{c^2} - \frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^2} - \frac{\frac{1}{2}x^2(a+b\operatorname{arctanh}(cx)) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(cx)}{c^3} - \frac{x}{c^2} \right)}{c^2}}{c^2} - \frac{\frac{1}{4}x^4(a+b\operatorname{arctanh}(cx)) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(cx)}{c^5} - \frac{x}{c^2} \right)}{c^2} \right)$$

6470

$$\frac{2}{5}bc \left(\frac{\frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))}{c} - b \int \frac{\log\left(\frac{2}{1-cx}\right)}{1-c^2x^2} dx}{c^2} - \frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^2} - \frac{\frac{1}{2}x^2(a+b\operatorname{arctanh}(cx)) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(cx)}{c^3} - \frac{x}{c^2} \right)}{c^2}}{c^2} - \frac{\frac{1}{4}x^4(a+b\operatorname{arctanh}(cx)) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(cx)}{c^5} - \frac{x}{c^2} \right)}{c^2} \right)$$

2849

$$\frac{2}{5}bc \left(\frac{\frac{b \int \frac{\log\left(\frac{2}{1-cx}\right)}{1-cx} dx + \frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))}{c}}{c^2} - \frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^2} - \frac{\frac{1}{2}x^2(a+b\operatorname{arctanh}(cx)) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(cx)}{c^3} - \frac{x}{c^2} \right)}{c^2}}{c^2} - \frac{\frac{1}{4}x^4(a+b\operatorname{arctanh}(cx)) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(cx)}{c^5} - \frac{x}{c^2} \right)}{c^2} \right)$$

2752

$$\frac{2}{5}bc \left(\frac{\frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))}{c} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{2c}}{c^2} - \frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^2} - \frac{\frac{1}{2}x^2(a+b\operatorname{arctanh}(cx)) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(cx)}{c^3} - \frac{x}{c^2} \right)}{c^2}}{c^2} - \frac{\frac{1}{4}x^4(a+b\operatorname{arctanh}(cx)) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(cx)}{c^5} - \frac{x}{c^2} \right)}{c^2} \right)$$

input

`Int [x^4*(a + b*ArcTanh [c*x])^2, x]`

output

$$\begin{aligned} & (x^5(a + b\text{ArcTanh}[c*x])^2)/5 - (2*b*c*(-(((x^4*(a + b\text{ArcTanh}[c*x]))/4 - \\ & (b*c*(-(x/c^4) - x^3/(3*c^2) + \text{ArcTanh}[c*x]/c^5))/4)/c^2) + (-(((x^2*(a + \\ & b*\text{ArcTanh}[c*x]))/2 - (b*c*(-(x/c^2) + \text{ArcTanh}[c*x]/c^3))/2)/c^2) + (-1/2* \\ & (a + b*\text{ArcTanh}[c*x])^2/(b*c^2) + (((a + b*\text{ArcTanh}[c*x])*\text{Log}[2/(1 - c*x)])/ \\ & c + (b*\text{PolyLog}[2, 1 - 2/(1 - c*x)])/(2*c))/c/c^2)/c^2)/5 \end{aligned}$$
Defintions of rubi rules used

rule 219

$$\text{Int}[\{(a_) + (b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))* \\ \text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] \text{ ; FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{Gt} \\ \text{Q}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 254

$$\text{Int}[(x_)^m/\{(a_) + (b_)*(x_)^2\}, x_Symbol] \rightarrow \text{Int}[\text{PolynomialDivide}[x^m, \\ a + b*x^2, x], x] \text{ ; FreeQ}\{a, b\}, x \ \&\& \ \text{IGtQ}[m, 3]$$

rule 262

$$\text{Int}[\{(c_)*(x_)^m\}*\{(a_) + (b_)*(x_)^2\}^p, x_Symbol] \rightarrow \text{Simp}[c*(c*x) \\ ^{m-1}*\{(a + b*x^2)^{p+1}/(b*(m+2*p+1))\}, x] - \text{Simp}[a*c^{2*(m-1)}/ \\ (b*(m+2*p+1)) \ \text{Int}[(c*x)^{m-2}*(a + b*x^2)^p, x], x] \text{ ; FreeQ}\{a, b \\ , c, p\}, x \ \&\& \ \text{GtQ}[m, 2-1] \ \&\& \ \text{NeQ}[m+2*p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c \\ , 2, m, p, x]$$

rule 2009

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ ; SumQ}[u]$$

rule 2752

$$\text{Int}[\text{Log}[(c_)*(x_)]/\{(d_) + (e_)*(x_)\}, x_Symbol] \rightarrow \text{Simp}[(-e^{-1})*\text{PolyLo} \\ \text{g}[2, 1 - c*x], x] \text{ ; FreeQ}\{c, d, e\}, x \ \&\& \ \text{EqQ}[e + c*d, 0]$$

rule 2849

$$\text{Int}[\text{Log}[(c_)]/\{(d_) + (e_)*(x_)\}]/\{(f_) + (g_)*(x_)^2\}, x_Symbol] \rightarrow \text{Simp} \\ [-e/g \ \text{Subst}[\text{Int}[\text{Log}[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] \text{ ; FreeQ} \\ \{c, d, e, f, g\}, x \ \&\& \ \text{EqQ}[c, 2*d] \ \&\& \ \text{EqQ}[e^2*f + d^2*g, 0]$$

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6470 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol] :> Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6542 `Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_))/((d_) + (e_.)*(x_)^2), x_Symbol] :> Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

rule 6546 `Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*(x_)/((d_) + (e_.)*(x_)^2), x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 0.78 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.41

method	result
parts	$\frac{a^2 x^5}{5} + \frac{b^2 \left(\frac{c^5 x^5 \operatorname{arctanh}(cx)^2}{5} + \frac{c^4 x^4 \operatorname{arctanh}(cx)}{10} + \frac{c^2 x^2 \operatorname{arctanh}(cx)}{5} + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{5} + \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{5} + \frac{c^3 x^3}{30} \right)}{5}$
derivativedivides	$\frac{a^2 c^5 x^5}{5} + b^2 \left(\frac{c^5 x^5 \operatorname{arctanh}(cx)^2}{5} + \frac{c^4 x^4 \operatorname{arctanh}(cx)}{10} + \frac{c^2 x^2 \operatorname{arctanh}(cx)}{5} + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{5} + \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{5} + \frac{c^3 x^3}{30} \right)$
default	$\frac{a^2 c^5 x^5}{5} + b^2 \left(\frac{c^5 x^5 \operatorname{arctanh}(cx)^2}{5} + \frac{c^4 x^4 \operatorname{arctanh}(cx)}{10} + \frac{c^2 x^2 \operatorname{arctanh}(cx)}{5} + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{5} + \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{5} + \frac{c^3 x^3}{30} \right)$
risch	$-\frac{ab \ln(-cx+1)x^5}{5} - \frac{b^2 \ln(-cx+1)x^4}{20c} - \frac{b^2 \ln(-cx+1)x^2}{10c^3} + \frac{ab \ln(-cx+1)}{5c^5} + \frac{b^2 \ln(cx+1)^2 x^5}{20} + \frac{ab x^4}{10c} + \frac{b^2 \ln(cx+1)}{10c}$

input `int(x^4*(a+b*arctanh(c*x))^2,x,method=_RETURNVERBOSE)`

output `1/5*a^2*x^5+b^2/c^5*(1/5*c^5*x^5*arctanh(c*x)^2+1/10*c^4*x^4*arctanh(c*x)+1/5*c^2*x^2*arctanh(c*x)+1/5*arctanh(c*x)*ln(c*x-1)+1/5*arctanh(c*x)*ln(c*x+1)+1/30*c^3*x^3+3/10*c*x+3/20*ln(c*x-1)-3/20*ln(c*x+1)-1/5*dilog(1/2*c*x+1/2)-1/10*ln(c*x-1)*ln(1/2*c*x+1/2)+1/20*ln(c*x-1)^2+1/10*(ln(c*x+1)-ln(1/2*c*x+1/2))*ln(-1/2*c*x+1/2)-1/20*ln(c*x+1)^2)+2*a*b/c^5*(1/5*c^5*x^5*arctanh(c*x)+1/20*c^4*x^4+1/10*c^2*x^2+1/10*ln(c*x-1)+1/10*ln(c*x+1))`

Fricas [F]

$$\int x^4 (a + b \operatorname{arctanh}(cx))^2 dx = \int (b \operatorname{arctanh}(cx) + a)^2 x^4 dx$$

input `integrate(x^4*(a+b*arctanh(c*x))^2,x, algorithm="fricas")`

output `integral(b^2*x^4*arctanh(c*x)^2 + 2*a*b*x^4*arctanh(c*x) + a^2*x^4, x)`

Sympy [F]

$$\int x^4(a + \operatorname{barctanh}(cx))^2 dx = \int x^4(a + b \operatorname{atanh}(cx))^2 dx$$

input `integrate(x**4*(a+b*atanh(c*x))**2,x)`

output `Integral(x**4*(a + b*atanh(c*x))**2, x)`

Maxima [F]

$$\int x^4(a + \operatorname{barctanh}(cx))^2 dx = \int (b \operatorname{artanh}(cx) + a)^2 x^4 dx$$

input `integrate(x^4*(a+b*arctanh(c*x))^2,x, algorithm="maxima")`

output `1/5*a^2*x^5 + 1/10*(4*x^5*arctanh(c*x) + c*((c^2*x^4 + 2*x^2)/c^4 + 2*log(c^2*x^2 - 1)/c^6))*a*b - 1/36000*(24*c^6*(2*(3*c^4*x^5 + 5*c^2*x^3 + 15*x)/c^10 - 15*log(c*x + 1)/c^11 + 15*log(c*x - 1)/c^11) - 45*c^5*((c^2*x^4 + 2*x^2)/c^8 + 2*log(c^2*x^2 - 1)/c^10) - 1080000*c^5*integrate(1/150*x^5*log(c*x + 1)/(c^6*x^2 - c^4), x) + 50*c^4*(2*(c^2*x^3 + 3*x)/c^8 - 3*log(c*x + 1)/c^9 + 3*log(c*x - 1)/c^9) - 300*c^3*(x^2/c^6 + log(c^2*x^2 - 1)/c^8) + 900*c^2*(2*x/c^6 - log(c*x + 1)/c^7 + log(c*x - 1)/c^7) - 540000*c*integrate(1/150*x*log(c*x + 1)/(c^6*x^2 - c^4), x) - 60*(30*c^5*x^5*log(c*x + 1)^2 + (12*c^5*x^5 - 15*c^4*x^4 + 20*c^3*x^3 - 30*c^2*x^2 + 60*c*x - 60*(c^5*x^5 + 1)*log(c*x + 1))*log(-c*x + 1))/c^5 - (72*(c*x - 1)^5*(25*log(-c*x + 1)^2 - 10*log(-c*x + 1) + 2) + 1125*(c*x - 1)^4*(8*log(-c*x + 1)^2 - 4*log(-c*x + 1) + 1) + 2000*(c*x - 1)^3*(9*log(-c*x + 1)^2 - 6*log(-c*x + 1) + 2) + 9000*(c*x - 1)^2*(2*log(-c*x + 1)^2 - 2*log(-c*x + 1) + 1) + 9000*(c*x - 1)*(log(-c*x + 1)^2 - 2*log(-c*x + 1) + 2))/c^5 + 1800*log(150*c^6*x^2 - 150*c^4)/c^5 - 540000*integrate(1/150*log(c*x + 1)/(c^6*x^2 - c^4), x))*b^2`

Giac [F]

$$\int x^4(a + b \operatorname{arctanh}(cx))^2 dx = \int (b \operatorname{artanh}(cx) + a)^2 x^4 dx$$

input `integrate(x^4*(a+b*arctanh(c*x))^2,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^2*x^4, x)`

Mupad [F(-1)]

Timed out.

$$\int x^4(a + b \operatorname{arctanh}(cx))^2 dx = \int x^4(a + b \operatorname{atanh}(cx))^2 dx$$

input `int(x^4*(a + b*atanh(c*x))^2,x)`

output `int(x^4*(a + b*atanh(c*x))^2, x)`

Reduce [F]

$$\int x^4(a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{6 \operatorname{atanh}(cx)^2 b^2 c^5 x^5 - 6 \operatorname{atanh}(cx)^2 b^2 cx + 12 \operatorname{atanh}(cx) ab c^5 x^5 + 12 \operatorname{atanh}(cx) ab + 3 \operatorname{atanh}(cx) b^2 c^4 x^4 + 6 \operatorname{atanh}(cx) b^2 c^3 x^3 + 9 b^2 c^2 x^2 - 9 \operatorname{atanh}(cx) b^2 c^2 x + 6 \operatorname{int}(\operatorname{atanh}(cx))^2, x) b^2 c^2 + 12 \log(c^2 x - c) a b + 6 a^2 c^5 x^5 + 3 a b c^4 x^4 + 6 a b c^3 x^3 + 9 b^2 c^2 x}{(30 c^5)}$$

input `int(x^4*(a+b*atanh(c*x))^2,x)`

output `(6*atanh(c*x)**2*b**2*c**5*x**5 - 6*atanh(c*x)**2*b**2*c*x + 12*atanh(c*x)*a*b*c**5*x**5 + 12*atanh(c*x)*a*b + 3*atanh(c*x)*b**2*c**4*x**4 + 6*atanh(c*x)*b**2*c**3*x**3 - 9*atanh(c*x)*b**2*c**2*x**2 - 9*atanh(c*x)*b**2*c**2*x + 6*int(atanh(c*x)**2,x)*b**2*c**2 + 12*log(c**2*x - c)*a*b + 6*a**2*c**5*x**5 + 3*a*b*c**4*x**4 + 6*a*b*c**3*x**3 + b**2*c**3*x**3 + 9*b**2*c*x)/(30*c**5)`

3.15 $\int x^3(a + b \operatorname{arctanh}(cx))^2 dx$

Optimal result	203
Mathematica [A] (verified)	203
Rubi [A] (verified)	204
Maple [A] (verified)	207
Fricas [A] (verification not implemented)	207
Sympy [A] (verification not implemented)	208
Maxima [A] (verification not implemented)	208
Giac [B] (verification not implemented)	209
Mupad [B] (verification not implemented)	210
Reduce [B] (verification not implemented)	210

Optimal result

Integrand size = 14, antiderivative size = 113

$$\int x^3(a + b \operatorname{arctanh}(cx))^2 dx = \frac{abx}{2c^3} + \frac{b^2x^2}{12c^2} + \frac{b^2x \operatorname{arctanh}(cx)}{2c^3} + \frac{bx^3(a + b \operatorname{arctanh}(cx))}{6c} - \frac{(a + b \operatorname{arctanh}(cx))^2}{4c^4} + \frac{1}{4}x^4(a + b \operatorname{arctanh}(cx))^2 + \frac{b^2 \log(1 - c^2x^2)}{3c^4}$$

output

$$\frac{1}{2}abx/c^3 + \frac{1}{12}b^2x^2/c^2 + \frac{1}{2}b^2x \operatorname{arctanh}(cx)/c^3 + \frac{1}{6}b^2x^3(a + b \operatorname{arctanh}(cx))/c - \frac{1}{4}(a + b \operatorname{arctanh}(cx))^2/c^4 + \frac{1}{4}x^4(a + b \operatorname{arctanh}(cx))^2 + \frac{1}{3}b^2 \ln(-c^2x^2 + 1)/c^4$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.17

$$\int x^3(a + b \operatorname{arctanh}(cx))^2 dx = \frac{6abcx + b^2c^2x^2 + 2abc^3x^3 + 3a^2c^4x^4 + 2bcx(3ac^3x^3 + b(3 + c^2x^2)) \operatorname{arctanh}(cx) + 3b^2(-1 + c^4x^4) \operatorname{arctan}}{12c^4}$$

input

```
Integrate[x^3*(a + b*ArcTanh[c*x])^2,x]
```


output

```
(6*a*b*c*x + b^2*c^2*x^2 + 2*a*b*c^3*x^3 + 3*a^2*c^4*x^4 + 2*b*c*x*(3*a*c^3*x^3 + b*(3 + c^2*x^2))*ArcTanh[c*x] + 3*b^2*(-1 + c^4*x^4)*ArcTanh[c*x]^2 + b*(3*a + 4*b)*Log[1 - c*x] - 3*a*b*Log[1 + c*x] + 4*b^2*Log[1 + c*x])/(12*c^4)
```

Rubi [A] (verified)

Time = 0.96 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.21, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {6452, 6542, 6452, 243, 49, 2009, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3(a + b \operatorname{arctanh}(cx))^2 dx \\
 & \quad \downarrow 6452 \\
 & \frac{1}{4}x^4(a + b \operatorname{arctanh}(cx))^2 - \frac{1}{2}bc \int \frac{x^4(a + b \operatorname{arctanh}(cx))}{1 - c^2x^2} dx \\
 & \quad \downarrow 6542 \\
 & \frac{1}{4}x^4(a + b \operatorname{arctanh}(cx))^2 - \frac{1}{2}bc \left(\frac{\int \frac{x^2(a + b \operatorname{arctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\int x^2(a + b \operatorname{arctanh}(cx)) dx}{c^2} \right) \\
 & \quad \downarrow 6452 \\
 & \frac{1}{4}x^4(a + b \operatorname{arctanh}(cx))^2 - \\
 & \frac{1}{2}bc \left(\frac{\int \frac{x^2(a + b \operatorname{arctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a + b \operatorname{arctanh}(cx)) - \frac{1}{3}bc \int \frac{x^3}{1 - c^2x^2} dx}{c^2} \right) \\
 & \quad \downarrow 243 \\
 & \frac{1}{4}x^4(a + b \operatorname{arctanh}(cx))^2 - \\
 & \frac{1}{2}bc \left(\frac{\int \frac{x^2(a + b \operatorname{arctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a + b \operatorname{arctanh}(cx)) - \frac{1}{6}bc \int \frac{x^2}{1 - c^2x^2} dx^2}{c^2} \right) \\
 & \quad \downarrow 49
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{2}bc \left(\frac{\int \frac{x^2(a + \operatorname{arctanh}(cx))}{1-c^2x^2} dx}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{arctanh}(cx))^2 - \frac{1}{3}x^3(a + \operatorname{arctanh}(cx)) - \frac{1}{6}bc \int \left(-\frac{1}{c^2} - \frac{1}{c^2(c^2x^2-1)}\right) dx^2}{c^2} \right) \\
& \quad \downarrow \text{2009} \\
& \frac{1}{2}bc \left(\frac{\int \frac{x^2(a + \operatorname{arctanh}(cx))}{1-c^2x^2} dx}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{arctanh}(cx))^2 - \frac{1}{3}x^3(a + \operatorname{arctanh}(cx)) - \frac{1}{6}bc \left(-\frac{x^2}{c^2} - \frac{\log(1-c^2x^2)}{c^4}\right)}{c^2} \right) \\
& \quad \downarrow \text{6542} \\
& \frac{1}{2}bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(cx)}{1-c^2x^2} dx}{c^2} - \frac{\int (a + \operatorname{arctanh}(cx)) dx}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{arctanh}(cx))^2 - \frac{1}{3}x^3(a + \operatorname{arctanh}(cx)) - \frac{1}{6}bc \left(-\frac{x^2}{c^2} - \frac{\log(1-c^2x^2)}{c^4}\right)}{c^2} \right) \\
& \quad \downarrow \text{2009} \\
& \frac{1}{2}bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(cx)}{1-c^2x^2} dx}{c^2} - \frac{ax + bx \operatorname{arctanh}(cx) + \frac{b \log(1-c^2x^2)}{2c}}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{arctanh}(cx))^2 - \frac{1}{3}x^3(a + \operatorname{arctanh}(cx)) - \frac{1}{6}bc \left(-\frac{x^2}{c^2} - \frac{\log(1-c^2x^2)}{c^4}\right)}{c^2} \right) \\
& \quad \downarrow \text{6510} \\
& \frac{1}{2}bc \left(\frac{(a + \operatorname{arctanh}(cx))^2}{2bc^3} - \frac{ax + bx \operatorname{arctanh}(cx) + \frac{b \log(1-c^2x^2)}{2c}}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{arctanh}(cx))^2 - \frac{1}{3}x^3(a + \operatorname{arctanh}(cx)) - \frac{1}{6}bc \left(-\frac{x^2}{c^2} - \frac{\log(1-c^2x^2)}{c^4}\right)}{c^2} \right)
\end{aligned}$$

input `Int [x^3*(a + b*ArcTanh[c*x])^2,x]`

output $(x^4*(a + b*ArcTanh[c*x])^2)/4 - (b*c*(-(((x^3*(a + b*ArcTanh[c*x])))/3 - (b*c*(-(x^2/c^2) - \operatorname{Log}[1 - c^2*x^2]/c^4))/6)/c^2) + ((a + b*ArcTanh[c*x])^2)/(2*b*c^3) - (a*x + b*x*ArcTanh[c*x] + (b*\operatorname{Log}[1 - c^2*x^2])/(2*c))/c^2/c^2)/2$

Definitions of rubi rules used

- rule 49 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}((c_.) + (d_.)(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[m + n + 2, 0]$
- rule 243 $\text{Int}(x_)^{(m_.)}((a_) + (b_.)(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/2 \ \text{Subst}[\text{Int}[x^{((m - 1)/2)*(a + b*x)^p}, x], x, x^2], x] /; \text{FreeQ}\{a, b, m, p\}, x \ \&\& \ \text{IntegerQ}[(m - 1)/2]$
- rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$
- rule 6452 $\text{Int}[(a_.) + \text{ArcTanh}[(c_.)(x_)^{(n_.)}](b_.)]^{(p_.)}(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}((a + b*\text{ArcTanh}[c*x^n])^p/(m + 1)), x] - \text{Simp}[b*c*n*(p/(m + 1)) \ \text{Int}[x^{(m + n)}((a + b*\text{ArcTanh}[c*x^n])^{(p - 1)/(1 - c^2*x^{(2*n)})}), x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$
- rule 6510 $\text{Int}[(a_.) + \text{ArcTanh}[(c_.)(x_)](b_.)]^{(p_.)}/((d_) + (e_.)(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^{(p + 1)}/(b*c*d*(p + 1)), x] /; \text{FreeQ}\{a, b, c, d, e, p\}, x \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{NeQ}[p, -1]$
- rule 6542 $\text{Int}[(a_.) + \text{ArcTanh}[(c_.)(x_)](b_.)]^{(p_.)}((f_.)(x_)^{(m_.)}), x_Symbol] \rightarrow \text{Simp}[f^2/e \ \text{Int}[(f*x)^{(m - 2)}(a + b*\text{ArcTanh}[c*x])^p, x], x] - \text{Simp}[d*(f^2/e) \ \text{Int}[(f*x)^{(m - 2)}(a + b*\text{ArcTanh}[c*x])^p/(d + e*x^2), x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{GtQ}[m, 1]$

Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.27

method	result
parallelsch	$\frac{3b^2 \operatorname{arctanh}(cx)^2 x^4 c^4 + 6ab \operatorname{arctanh}(cx) x^4 c^4 + 3a^2 c^4 x^4 + 2b^2 \operatorname{arctanh}(cx) x^3 c^3 + 2ab c^3 x^3 + b^2 c^2 x^2 + 6b^2 \operatorname{arctanh}(cx) x c + 6b^2 \operatorname{arctanh}(cx)}{12c^4}$
parts	$\frac{a^2 x^4}{4} + \frac{b^2 \left(\frac{c^4 x^4 \operatorname{arctanh}(cx)^2}{4} + \frac{c^3 x^3 \operatorname{arctanh}(cx)}{6} + \frac{cx \operatorname{arctanh}(cx)}{2} + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{4} - \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{4} - \frac{\ln(cx-1)}{c^4} \right)}{c^4}$
derivativdivides	$\frac{a^2 c^4 x^4}{4} + b^2 \left(\frac{c^4 x^4 \operatorname{arctanh}(cx)^2}{4} + \frac{c^3 x^3 \operatorname{arctanh}(cx)}{6} + \frac{cx \operatorname{arctanh}(cx)}{2} + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{4} - \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{4} - \frac{\ln(cx-1)}{c^4} \right)$
default	$\frac{a^2 c^4 x^4}{4} + b^2 \left(\frac{c^4 x^4 \operatorname{arctanh}(cx)^2}{4} + \frac{c^3 x^3 \operatorname{arctanh}(cx)}{6} + \frac{cx \operatorname{arctanh}(cx)}{2} + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{4} - \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{4} - \frac{\ln(cx-1)}{c^4} \right)$
risch	$\frac{b^2 (c^4 x^4 - 1) \ln(cx+1)^2}{16c^4} + \frac{b(-3bx^4 \ln(-cx+1)c^4 + 6ac^4 x^4 + 2bc^3 x^3 + 6bcx + 3b \ln(-cx+1)) \ln(cx+1)}{24c^4} + \frac{\ln(-cx+1)^2}{16}$

input `int(x^3*(a+b*arctanh(c*x))^2,x,method=_RETURNVERBOSE)`

output $\frac{1}{12} * (3 * b^2 * \operatorname{arctanh}(c * x)^2 * x^4 * c^4 + 6 * a * b * \operatorname{arctanh}(c * x) * x^4 * c^4 + 3 * a^2 * c^4 * x^4 + 2 * b^2 * \operatorname{arctanh}(c * x) * x^3 * c^3 + 2 * a * b * c^3 * x^3 + b^2 * c^2 * x^2 + 6 * b^2 * \operatorname{arctanh}(c * x) * x * c + 6 * a * b * c * x - 3 * b^2 * \operatorname{arctanh}(c * x)^2 + 8 * \ln(c * x - 1) * b^2 - 6 * \operatorname{arctanh}(c * x) * a * b + 8 * \operatorname{arctanh}(c * x) * b^2 + b^2) / c^4$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.42

$$\int x^3 (a + b \operatorname{arctanh}(cx))^2 dx = \frac{12 a^2 c^4 x^4 + 8 abc^3 x^3 + 4 b^2 c^2 x^2 + 24 abcx + 3 (b^2 c^4 x^4 - b^2) \log\left(-\frac{cx+1}{cx-1}\right)^2 - 4 (3 ab - 4 b^2) \log(cx+1) + 48 c^4}{48 c^4}$$

input `integrate(x^3*(a+b*arctanh(c*x))^2,x, algorithm="fricas")`

output

```
1/48*(12*a^2*c^4*x^4 + 8*a*b*c^3*x^3 + 4*b^2*c^2*x^2 + 24*a*b*c*x + 3*(b^2*c^4*x^4 - b^2)*log(-(c*x + 1)/(c*x - 1))^2 - 4*(3*a*b - 4*b^2)*log(c*x + 1) + 4*(3*a*b + 4*b^2)*log(c*x - 1) + 4*(3*a*b*c^4*x^4 + b^2*c^3*x^3 + 3*b^2*c*x)*log(-(c*x + 1)/(c*x - 1)))/c^4
```

Sympy [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.49

$$\int x^3(a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \begin{cases} \frac{a^2 x^4}{4} + \frac{abx^4 \operatorname{atanh}(cx)}{2} + \frac{abx^3}{6c} + \frac{abx}{2c^3} - \frac{ab \operatorname{atanh}(cx)}{2c^4} + \frac{b^2 x^4 \operatorname{atanh}^2(cx)}{4} + \frac{b^2 x^3 \operatorname{atanh}(cx)}{6c} + \frac{b^2 x^2}{12c^2} + \frac{b^2 x \operatorname{atanh}(cx)}{2c^3} + \frac{2b^2}{12c^2} \\ \frac{a^2 x^4}{4} \end{cases}$$

input

```
integrate(x**3*(a+b*atanh(c*x))**2,x)
```

output

```
Piecewise((a**2*x**4/4 + a*b*x**4*atanh(c*x)/2 + a*b*x**3/(6*c) + a*b*x/(2*c**3) - a*b*atanh(c*x)/(2*c**4) + b**2*x**4*atanh(c*x)**2/4 + b**2*x**3*a*tanh(c*x)/(6*c) + b**2*x**2/(12*c**2) + b**2*x*atanh(c*x)/(2*c**3) + 2*b**2*log(x - 1/c)/(3*c**4) - b**2*atanh(c*x)**2/(4*c**4) + 2*b**2*atanh(c*x)/(3*c**4), Ne(c, 0)), (a**2*x**4/4, True))
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.67

$$\int x^3(a + b \operatorname{arctanh}(cx))^2 dx = \frac{1}{4} b^2 x^4 \operatorname{artanh}(cx)^2 + \frac{1}{4} a^2 x^4$$

$$+ \frac{1}{12} \left(6x^4 \operatorname{artanh}(cx) + c \left(\frac{2(c^2 x^3 + 3x)}{c^4} - \frac{3 \log(cx + 1)}{c^5} + \frac{3 \log(cx - 1)}{c^5} \right) \right) ab$$

$$+ \frac{1}{48} \left(4c \left(\frac{2(c^2 x^3 + 3x)}{c^4} - \frac{3 \log(cx + 1)}{c^5} + \frac{3 \log(cx - 1)}{c^5} \right) \operatorname{artanh}(cx) + \frac{4c^2 x^2 - 2(3 \log(cx - 1) - 3 \log(cx + 1))}{c^5} \right)$$

input

```
integrate(x^3*(a+b*arctanh(c*x))^2,x, algorithm="maxima")
```

output

```
1/4*b^2*x^4*arctanh(c*x)^2 + 1/4*a^2*x^4 + 1/12*(6*x^4*arctanh(c*x) + c*(2
*(c^2*x^3 + 3*x)/c^4 - 3*log(c*x + 1)/c^5 + 3*log(c*x - 1)/c^5))*a*b + 1/4
8*(4*c*(2*(c^2*x^3 + 3*x)/c^4 - 3*log(c*x + 1)/c^5 + 3*log(c*x - 1)/c^5)*a
rctanh(c*x) + (4*c^2*x^2 - 2*(3*log(c*x - 1) - 8)*log(c*x + 1) + 3*log(c*x
+ 1)^2 + 3*log(c*x - 1)^2 + 16*log(c*x - 1))/c^4)*b^2
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 603 vs. 2(99) = 198.

Time = 0.14 (sec) , antiderivative size = 603, normalized size of antiderivative = 5.34

$$\int x^3(a + b \operatorname{arctanh}(cx))^2 dx = \text{Too large to display}$$

input

```
integrate(x^3*(a+b*arctanh(c*x))^2,x, algorithm="giac")
```

output

```
1/6*(3*((c*x + 1)^3*b^2/(c*x - 1)^3 + (c*x + 1)*b^2/(c*x - 1))*log(-(c*x +
1)/(c*x - 1))^2/((c*x + 1)^4*c^5/(c*x - 1)^4 - 4*(c*x + 1)^3*c^5/(c*x - 1
)^3 + 6*(c*x + 1)^2*c^5/(c*x - 1)^2 - 4*(c*x + 1)*c^5/(c*x - 1) + c^5) + 2
*(6*(c*x + 1)^3*a*b/(c*x - 1)^3 + 6*(c*x + 1)*a*b/(c*x - 1) + 3*(c*x + 1)^
3*b^2/(c*x - 1)^3 - 6*(c*x + 1)^2*b^2/(c*x - 1)^2 + 5*(c*x + 1)*b^2/(c*x -
1) - 2*b^2)*log(-(c*x + 1)/(c*x - 1))/((c*x + 1)^4*c^5/(c*x - 1)^4 - 4*(c
*x + 1)^3*c^5/(c*x - 1)^3 + 6*(c*x + 1)^2*c^5/(c*x - 1)^2 - 4*(c*x + 1)*c^
5/(c*x - 1) + c^5) + 2*(6*(c*x + 1)^3*a^2/(c*x - 1)^3 + 6*(c*x + 1)*a^2/(c
*x - 1) + 6*(c*x + 1)^3*a*b/(c*x - 1)^3 - 12*(c*x + 1)^2*a*b/(c*x - 1)^2 +
10*(c*x + 1)*a*b/(c*x - 1) - 4*a*b + (c*x + 1)^3*b^2/(c*x - 1)^3 - 2*(c*x
+ 1)^2*b^2/(c*x - 1)^2 + (c*x + 1)*b^2/(c*x - 1))/((c*x + 1)^4*c^5/(c*x -
1)^4 - 4*(c*x + 1)^3*c^5/(c*x - 1)^3 + 6*(c*x + 1)^2*c^5/(c*x - 1)^2 - 4*
(c*x + 1)*c^5/(c*x - 1) + c^5) - 4*b^2*log(-(c*x + 1)/(c*x - 1) + 1)/c^5 +
4*b^2*log(-(c*x + 1)/(c*x - 1))/c^5)*c
```

Mupad [B] (verification not implemented)

Time = 3.94 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.19

$$\int x^3(a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{4b^2 \ln(c^2 x^2 - 1) - 3b^2 \operatorname{atanh}(cx)^2 + 3a^2 c^4 x^4 + b^2 c^2 x^2 - 6ab \operatorname{atanh}(cx) + 2b^2 c^3 x^3 \operatorname{atanh}(cx) + 6b^2 c^4 x^4 \operatorname{atanh}(cx)}{12c^4}$$

input `int(x^3*(a + b*atanh(c*x))^2,x)`output `(4*b^2*log(c^2*x^2 - 1) - 3*b^2*atanh(c*x)^2 + 3*a^2*c^4*x^4 + b^2*c^2*x^2 - 6*a*b*atanh(c*x) + 2*b^2*c^3*x^3*atanh(c*x) + 6*b^2*c*x*atanh(c*x) + 3*b^2*c^4*x^4*atanh(c*x)^2 + 2*a*b*c^3*x^3 + 6*a*b*c*x + 6*a*b*c^4*x^4*atanh(c*x))/(12*c^4)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.27

$$\int x^3(a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{3 \operatorname{atanh}(cx)^2 b^2 c^4 x^4 - 3 \operatorname{atanh}(cx)^2 b^2 + 6 \operatorname{atanh}(cx) ab c^4 x^4 - 6 \operatorname{atanh}(cx) ab + 2 \operatorname{atanh}(cx) b^2 c^3 x^3 + 6 \operatorname{atanh}(cx) b^2 c^4 x^4 \operatorname{atanh}(cx) + 2 a^2 c^4 x^4 + b^2 c^2 x^2 - 6 a b \operatorname{atanh}(cx) + 2 b^2 c^3 x^3 \operatorname{atanh}(cx) + 6 b^2 c^4 x^4 \operatorname{atanh}(cx)}{12c^4}$$

input `int(x^3*(a+b*atanh(c*x))^2,x)`output `(3*atanh(c*x)**2*b**2*c**4*x**4 - 3*atanh(c*x)**2*b**2 + 6*atanh(c*x)*a*b*c**4*x**4 - 6*atanh(c*x)*a*b + 2*atanh(c*x)*b**2*c**3*x**3 + 6*atanh(c*x)*b**2*c*x + 8*atanh(c*x)*b**2 + 8*log(c**2*x - c)*b**2 + 3*a**2*c**4*x**4 + 2*a*b*c**3*x**3 + 6*a*b*c*x + b**2*c**2*x**2)/(12*c**4)`

3.16 $\int x^2(a + b \operatorname{arctanh}(cx))^2 dx$

Optimal result	211
Mathematica [A] (verified)	212
Rubi [A] (verified)	212
Maple [A] (verified)	215
Fricas [F]	216
Sympy [F]	217
Maxima [F]	217
Giac [F]	218
Mupad [F(-1)]	218
Reduce [F]	218

Optimal result

Integrand size = 14, antiderivative size = 130

$$\int x^2(a + b \operatorname{arctanh}(cx))^2 dx = \frac{b^2 x}{3c^2} - \frac{b^2 \operatorname{arctanh}(cx)}{3c^3} + \frac{bx^2(a + b \operatorname{arctanh}(cx))}{3c} + \frac{(a + b \operatorname{arctanh}(cx))^2}{3c^3} + \frac{1}{3}x^3(a + b \operatorname{arctanh}(cx))^2 - \frac{2b(a + b \operatorname{arctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{3c^3} - \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{3c^3}$$

output

```
1/3*b^2*x/c^2-1/3*b^2*arctanh(c*x)/c^3+1/3*b*x^2*(a+b*arctanh(c*x))/c+1/3*(a+b*arctanh(c*x))^2/c^3+1/3*x^3*(a+b*arctanh(c*x))^2-2/3*b*(a+b*arctanh(c*x))*ln(2/(-c*x+1))/c^3-1/3*b^2*polylog(2,1-2/(-c*x+1))/c^3
```


Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.94

$$\int x^2(a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{b^2 cx + abc^2 x^2 + a^2 c^3 x^3 + b^2(-1 + c^3 x^3) \operatorname{arctanh}(cx)^2 + b \operatorname{arctanh}(cx) (-b + bc^2 x^2 + 2ac^3 x^3 - 2b \log(1 + c^2 x^2))}{3c^3}$$

input `Integrate[x^2*(a + b*ArcTanh[c*x])^2,x]`

output `(b^2*c*x + a*b*c^2*x^2 + a^2*c^3*x^3 + b^2*(-1 + c^3*x^3)*ArcTanh[c*x]^2 + b*ArcTanh[c*x]*(-b + b*c^2*x^2 + 2*a*c^3*x^3 - 2*b*Log[1 + E^(-2*ArcTanh[c*x])])) + a*b*Log[-1 + c^2*x^2] + b^2*PolyLog[2, -E^(-2*ArcTanh[c*x])])/(3*c^3)`

Rubi [A] (verified)

Time = 1.39 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.09, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {6452, 6542, 6452, 262, 219, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2(a + b \operatorname{arctanh}(cx))^2 dx$$

$$\downarrow 6452$$

$$\frac{1}{3}x^3(a + b \operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \int \frac{x^3(a + b \operatorname{arctanh}(cx))}{1 - c^2 x^2} dx$$

$$\downarrow 6542$$

$$\frac{1}{3}x^3(a + b \operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\int \frac{x(a + b \operatorname{arctanh}(cx))}{1 - c^2 x^2} dx}{c^2} - \frac{\int x(a + b \operatorname{arctanh}(cx)) dx}{c^2} \right)$$

$$\downarrow 6452$$

$$\frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx)) - \frac{1}{2}bc \int \frac{x^2}{1 - c^2x^2} dx}{c^2} \right)$$

↓ 262

$$\frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx)) - \frac{1}{2}bc \left(\frac{\int \frac{1}{1 - c^2x^2} dx}{c^2} - \frac{x}{c^2} \right)}{c^2} \right)$$

↓ 219

$$\frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx)) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(cx)}{c^3} - \frac{x}{c^2} \right)}{c^2} \right)$$

↓ 6546

$$\frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(cx)}{1 - cx} dx}{c} - \frac{(a + \operatorname{barctanh}(cx))^2}{2bc^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx)) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(cx)}{c^3} - \frac{x}{c^2} \right)}{c^2} \right)$$

↓ 6470

$$\frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\frac{\log\left(\frac{2}{1 - cx}\right)(a + b \operatorname{arctanh}(cx))}{c} - b \int \frac{\log\left(\frac{2}{1 - cx}\right)}{1 - c^2x^2} dx}{c} - \frac{(a + \operatorname{barctanh}(cx))^2}{2bc^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx)) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(cx)}{c^3} - \frac{x}{c^2} \right)}{c^2} \right)$$

↓ 2849

$$\frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\frac{b \int \frac{\log\left(\frac{2}{1 - cx}\right)}{1 - \frac{2}{1 - cx}} d \frac{1}{1 - cx} + \log\left(\frac{2}{1 - cx}\right)(a + b \operatorname{arctanh}(cx))}{c} - \frac{(a + \operatorname{barctanh}(cx))^2}{2bc^2}}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx)) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(cx)}{c^3} - \frac{x}{c^2} \right)}{c^2} \right)$$

$$\begin{array}{c} \downarrow 2752 \\ \frac{1}{3}x^3(a + b\operatorname{arctanh}(cx))^2 - \\ \frac{2}{3}bc \left(\frac{\frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))}{c} + \frac{b\operatorname{PolyLog}\left(2,1-\frac{2}{1-cx}\right)}{2c}}{c^2} - \frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^2} - \frac{\frac{1}{2}x^2(a + b\operatorname{arctanh}(cx)) - \frac{1}{2}bc\left(\frac{\operatorname{arctanh}(cx)}{c^3}\right)}{c^2} \right) \end{array}$$

input `Int[x^2*(a + b*ArcTanh[c*x])^2,x]`

output `(x^3*(a + b*ArcTanh[c*x])^2)/3 - (2*b*c*(-((x^2*(a + b*ArcTanh[c*x]))/2 - (b*c*(-(x/c^2) + ArcTanh[c*x]/c^3))/2)/c^2) + (-1/2*(a + b*ArcTanh[c*x])^2/(b*c^2) + (((a + b*ArcTanh[c*x])*Log[2/(1 - c*x)])/c + (b*PolyLog[2, 1 - 2/(1 - c*x)])/(2*c))/c)/c^2)/3`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 262 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2752 `Int[Log[(c_)*(x_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_)]/((d_) + (e_)*(x_)]/((f_) + (g_)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6470 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol] :> Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6542 `Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_))/((d_) + (e_.)*(x_)^2), x_Symbol] :> Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

rule 6546 `Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*(x_)/((d_) + (e_.)*(x_)^2), x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 0.71 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.55

method	result
parts	$\frac{a^2 x^3}{3} + \frac{b^2 \left(\frac{c^3 x^3 \operatorname{arctanh}(cx)^2}{3} + \frac{c^2 x^2 \operatorname{arctanh}(cx)}{3} + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{3} + \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{3} + \frac{cx}{3} + \frac{\ln(cx-1)}{6} - \frac{\ln(cx+1)}{6} \right)}{c^3}$
derivativedivides	$\frac{a^2 c^3 x^3}{3} + b^2 \left(\frac{c^3 x^3 \operatorname{arctanh}(cx)^2}{3} + \frac{c^2 x^2 \operatorname{arctanh}(cx)}{3} + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{3} + \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{3} + \frac{cx}{3} + \frac{\ln(cx-1)}{6} - \frac{\ln(cx+1)}{6} \right)$
default	$\frac{a^2 c^3 x^3}{3} + b^2 \left(\frac{c^3 x^3 \operatorname{arctanh}(cx)^2}{3} + \frac{c^2 x^2 \operatorname{arctanh}(cx)}{3} + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{3} + \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{3} + \frac{cx}{3} + \frac{\ln(cx-1)}{6} - \frac{\ln(cx+1)}{6} \right)$
risch	$-\frac{b^2 \ln(-cx+1)^2}{12c^3} + \frac{b^2 \ln(cx+1)x^2}{6c} + \frac{11b^2 \ln(-cx+1)}{18c^3} + \frac{b^2 \ln(-cx+1)^2 x^3}{12} + \frac{a^2 x^3}{3} + \frac{ba \ln(cx+1)}{3c^3} - \frac{ab \ln(-cx+1)}{3c^3}$

```
input int(x^2*(a+b*arctanh(c*x))^2,x,method=_RETURNVERBOSE)
```

```
output 1/3*a^2*x^3+b^2/c^3*(1/3*c^3*x^3*arctanh(c*x)^2+1/3*c^2*x^2*arctanh(c*x)+1/3*arctanh(c*x)*ln(c*x-1)+1/3*arctanh(c*x)*ln(c*x+1)+1/3*c*x+1/6*ln(c*x-1)-1/6*ln(c*x+1)+1/12*ln(c*x-1)^2-1/3*dilog(1/2*c*x+1/2)-1/6*ln(c*x-1)*ln(1/2*c*x+1/2)-1/12*ln(c*x+1)^2+1/6*(ln(c*x+1)-ln(1/2*c*x+1/2))*ln(-1/2*c*x+1/2))+2*a*b/c^3*(1/3*c^3*x^3*arctanh(c*x)+1/6*c^2*x^2+1/6*ln(c*x-1)+1/6*ln(c*x+1))
```

Fricas [F]

$$\int x^2(a + b \operatorname{arctanh}(cx))^2 dx = \int (b \operatorname{arctanh}(cx) + a)^2 x^2 dx$$

```
input integrate(x^2*(a+b*arctanh(c*x))^2,x, algorithm="fricas")
```

```
output integral(b^2*x^2*arctanh(c*x)^2 + 2*a*b*x^2*arctanh(c*x) + a^2*x^2, x)
```

Sympy [F]

$$\int x^2(a + \operatorname{arctanh}(cx))^2 dx = \int x^2(a + b \operatorname{atanh}(cx))^2 dx$$

input `integrate(x**2*(a+b*atanh(c*x))**2,x)`

output `Integral(x**2*(a + b*atanh(c*x))**2, x)`

Maxima [F]

$$\int x^2(a + \operatorname{arctanh}(cx))^2 dx = \int (b \operatorname{artanh}(cx) + a)^2 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c*x))^2,x, algorithm="maxima")`

output `1/3*a^2*x^3 + 1/3*(2*x^3*arctanh(c*x) + c*(x^2/c^2 + log(c^2*x^2 - 1)/c^4)
)*a*b - 1/216*(2*c^4*(2*(c^2*x^3 + 3*x)/c^6 - 3*log(c*x + 1)/c^7 + 3*log(c
*x - 1)/c^7) - 3*c^3*(x^2/c^4 + log(c^2*x^2 - 1)/c^6) - 648*c^3*integrate(
1/9*x^3*log(c*x + 1)/(c^4*x^2 - c^2), x) + 9*c^2*(2*x/c^4 - log(c*x + 1)/c
^5 + log(c*x - 1)/c^5) - 324*c*integrate(1/9*x*log(c*x + 1)/(c^4*x^2 - c^2
), x) - 6*(3*c^3*x^3*log(c*x + 1)^2 + (2*c^3*x^3 - 3*c^2*x^2 + 6*c*x - 6*(
c^3*x^3 + 1)*log(c*x + 1))*log(-c*x + 1))/c^3 - (2*(c*x - 1)^3*(9*log(-c*x
+ 1)^2 - 6*log(-c*x + 1) + 2) + 27*(c*x - 1)^2*(2*log(-c*x + 1)^2 - 2*log
(-c*x + 1) + 1) + 54*(c*x - 1)*(log(-c*x + 1)^2 - 2*log(-c*x + 1) + 2))/c^
3 + 18*log(9*c^4*x^2 - 9*c^2)/c^3 - 324*integrate(1/9*log(c*x + 1)/(c^4*x^
2 - c^2), x))*b^2`

Giac [F]

$$\int x^2(a + b \operatorname{arctanh}(cx))^2 dx = \int (b \operatorname{artanh}(cx) + a)^2 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c*x))^2,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^2*x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^2(a + b \operatorname{arctanh}(cx))^2 dx = \int x^2(a + b \operatorname{atanh}(cx))^2 dx$$

input `int(x^2*(a + b*atanh(c*x))^2,x)`

output `int(x^2*(a + b*atanh(c*x))^2, x)`

Reduce [F]

$$\int x^2(a + b \operatorname{arctanh}(cx))^2 dx = \frac{\operatorname{atanh}(cx)^2 b^2 c^3 x^3 - \operatorname{atanh}(cx)^2 b^2 cx + 2 \operatorname{atanh}(cx) ab c^3 x^3 + 2 \operatorname{atanh}(cx) ab + \operatorname{atanh}(cx) b^2 c^2 x^2 - \operatorname{atanh}(cx)}{3c^3}$$

input `int(x^2*(a+b*atanh(c*x))^2,x)`

output `(atanh(c*x)**2*b**2*c**3*x**3 - atanh(c*x)**2*b**2*c*x + 2*atanh(c*x)*a*b*c**3*x**3 + 2*atanh(c*x)*a*b + atanh(c*x)*b**2*c**2*x**2 - atanh(c*x)*b**2 + int(atanh(c*x)**2,x)*b**2*c + 2*log(c**2*x - c)*a*b + a**2*c**3*x**3 + a*b*c**2*x**2 + b**2*c*x)/(3*c**3)`

3.17 $\int x(a + b \operatorname{arctanh}(cx))^2 dx$

Optimal result	219
Mathematica [A] (verified)	219
Rubi [A] (verified)	220
Maple [A] (verified)	222
Fricas [A] (verification not implemented)	222
Sympy [A] (verification not implemented)	223
Maxima [B] (verification not implemented)	223
Giac [B] (verification not implemented)	224
Mupad [B] (verification not implemented)	224
Reduce [B] (verification not implemented)	225

Optimal result

Integrand size = 12, antiderivative size = 75

$$\int x(a + b \operatorname{arctanh}(cx))^2 dx = \frac{abx}{c} + \frac{b^2 x \operatorname{arctanh}(cx)}{c} - \frac{(a + b \operatorname{arctanh}(cx))^2}{2c^2} + \frac{1}{2} x^2 (a + b \operatorname{arctanh}(cx))^2 + \frac{b^2 \log(1 - c^2 x^2)}{2c^2}$$

output

$a*b*x/c + b^2*x*\operatorname{arctanh}(c*x)/c - 1/2*(a+b*\operatorname{arctanh}(c*x))^2/c^2 + 1/2*x^2*(a+b*\operatorname{arctanh}(c*x))^2 + 1/2*b^2*\ln(-c^2*x^2+1)/c^2$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.20

$$\int x(a + b \operatorname{arctanh}(cx))^2 dx = \frac{2abcx + a^2c^2x^2 + 2bcx(b + acx)\operatorname{arctanh}(cx) + b^2(-1 + c^2x^2)\operatorname{arctanh}(cx)^2 + b(a + b)\log(1 - cx) - ab \log(1 - c^2x^2)}{2c^2}$$

input

$\operatorname{Integrate}[x*(a + b*\operatorname{ArcTanh}[c*x])^2, x]$

output

$$(2*a*b*c*x + a^2*c^2*x^2 + 2*b*c*x*(b + a*c*x)*\text{ArcTanh}[c*x] + b^2*(-1 + c^2*x^2)*\text{ArcTanh}[c*x]^2 + b*(a + b)*\text{Log}[1 - c*x] - a*b*\text{Log}[1 + c*x] + b^2*\text{Log}[1 + c*x])/(2*c^2)$$
Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.04, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6452, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(a + \text{barctanh}(cx))^2 dx$$

$$\downarrow 6452$$

$$\frac{1}{2}x^2(a + \text{barctanh}(cx))^2 - bc \int \frac{x^2(a + \text{barctanh}(cx))}{1 - c^2x^2} dx$$

$$\downarrow 6542$$

$$\frac{1}{2}x^2(a + \text{barctanh}(cx))^2 - bc \left(\frac{\int \frac{a + \text{barctanh}(cx)}{1 - c^2x^2} dx}{c^2} - \frac{\int (a + \text{barctanh}(cx)) dx}{c^2} \right)$$

$$\downarrow 2009$$

$$\frac{1}{2}x^2(a + \text{barctanh}(cx))^2 - bc \left(\frac{\int \frac{a + \text{barctanh}(cx)}{1 - c^2x^2} dx}{c^2} - \frac{ax + b\text{barctanh}(cx) + \frac{b \log(1 - c^2x^2)}{2c}}{c^2} \right)$$

$$\downarrow 6510$$

$$\frac{1}{2}x^2(a + \text{barctanh}(cx))^2 - bc \left(\frac{(a + \text{barctanh}(cx))^2}{2bc^3} - \frac{ax + b\text{barctanh}(cx) + \frac{b \log(1 - c^2x^2)}{2c}}{c^2} \right)$$

input

$$\text{Int}[x*(a + b*\text{ArcTanh}[c*x])^2, x]$$

output $(x^2(a + b \operatorname{ArcTanh}[c x])^2)/2 - b c ((a + b \operatorname{ArcTanh}[c x])^2/(2 b c^3) - (a x + b x \operatorname{ArcTanh}[c x] + (b \operatorname{Log}[1 - c^2 x^2])/(2 c))/c^2)$

Defintions of rubi rules used

rule 2009 $\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] \text{ /; SumQ}[u]$

rule 6452 $\operatorname{Int}[(a + \operatorname{ArcTanh}[c x]^n] \cdot b)^p \cdot x^m, x_Symbol] \rightarrow \operatorname{Simp}[x^{m+1} \cdot (a + b \operatorname{ArcTanh}[c x^n])^{p/(m+1)}, x] - \operatorname{Simp}[b c^n \cdot (p/(m+1)) \operatorname{Int}[x^{m+n} \cdot (a + b \operatorname{ArcTanh}[c x^n])^{p-1} / (1 - c^2 x^{2n})], x], x] \text{ /; FreeQ}\{a, b, c, m, n\}, x \ \&\& \operatorname{IGtQ}[p, 0] \ \&\& (\operatorname{EqQ}[p, 1] \ || \ (\operatorname{EqQ}[n, 1] \ \&\& \operatorname{IntegerQ}[m])) \ \&\& \operatorname{NeQ}[m, -1]$

rule 6510 $\operatorname{Int}[(a + \operatorname{ArcTanh}[c x]) \cdot b]^p / (d + e x^2), x_Symbol] \rightarrow \operatorname{Simp}[(a + b \operatorname{ArcTanh}[c x])^{p+1} / (b c d (p+1)), x] \text{ /; FreeQ}\{a, b, c, d, e, p\}, x \ \&\& \operatorname{EqQ}[c^2 d + e, 0] \ \&\& \operatorname{NeQ}[p, -1]$

rule 6542 $\operatorname{Int}[(a + \operatorname{ArcTanh}[c x]) \cdot b]^p \cdot (f x)^m / (d + e x^2), x_Symbol] \rightarrow \operatorname{Simp}[f^2/e \operatorname{Int}[(f x)^{m-2} \cdot (a + b \operatorname{ArcTanh}[c x])^p, x], x] - \operatorname{Simp}[d \cdot (f^2/e) \operatorname{Int}[(f x)^{m-2} \cdot (a + b \operatorname{ArcTanh}[c x])^p / (d + e x^2)], x], x] \text{ /; FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \operatorname{GtQ}[p, 0] \ \&\& \operatorname{GtQ}[m, 1]$

Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.41

method	result
parallelrisc	$\frac{b^2 \operatorname{arctanh}(cx)^2 x^2 c^2 + 2x^2 \operatorname{arctanh}(cx) ab c^2 + a^2 c^2 x^2 + 2b^2 \operatorname{arctanh}(cx) xc + 2abcx - b^2 \operatorname{arctanh}(cx)^2 + 2 \ln(cx-1) b^2 - 2 a}{2c^2}$
parts	$\frac{a^2 x^2}{2} + \frac{b^2 \left(\frac{c^2 x^2 \operatorname{arctanh}(cx)^2}{2} + cx \operatorname{arctanh}(cx) + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{2} - \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{2} + \frac{\ln(cx-1)^2}{8} - \frac{\ln(cx-1) \ln\left(\frac{cx}{2}\right)}{4} \right)}{c^2}$
derivativedivides	$\frac{a^2 c^2 x^2}{2} + b^2 \left(\frac{c^2 x^2 \operatorname{arctanh}(cx)^2}{2} + cx \operatorname{arctanh}(cx) + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{2} - \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{2} + \frac{\ln(cx-1)^2}{8} - \frac{\ln(cx-1) \ln\left(\frac{cx}{2}\right)}{4} \right)$
default	$\frac{a^2 c^2 x^2}{2} + b^2 \left(\frac{c^2 x^2 \operatorname{arctanh}(cx)^2}{2} + cx \operatorname{arctanh}(cx) + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{2} - \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{2} + \frac{\ln(cx-1)^2}{8} - \frac{\ln(cx-1) \ln\left(\frac{cx}{2}\right)}{4} \right)$
risc	$\frac{b^2 (c^2 x^2 - 1) \ln(cx+1)^2}{8c^2} + \frac{b(-x^2 b \ln(-cx+1) c^2 + 2a c^2 x^2 + 2bcx + b \ln(-cx+1)) \ln(cx+1)}{4c^2} + \frac{\ln(-cx+1)^2 b^2 x^2}{8} - \frac{\ln(-cx+1) \ln\left(\frac{cx}{2}\right) b^2}{4}$

input `int(x*(a+b*arctanh(c*x))^2,x,method=_RETURNVERBOSE)`

output $\frac{1}{2} * (b^2 * \operatorname{arctanh}(c * x)^2 * x^2 * c^2 + 2 * x^2 * \operatorname{arctanh}(c * x) * a * b * c^2 + a^2 * c^2 * x^2 + 2 * b^2 * \operatorname{arctanh}(c * x) * x * c + 2 * a * b * c * x - b^2 * \operatorname{arctanh}(c * x)^2 + 2 * \ln(c * x - 1) * b^2 - 2 * \operatorname{arctanh}(c * x) * a * b + 2 * \operatorname{arctanh}(c * x) * b^2 + a^2) / c^2$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.63

$$\int x(a + b \operatorname{arctanh}(cx))^2 dx = \frac{4a^2c^2x^2 + 8abcx + (b^2c^2x^2 - b^2) \log\left(-\frac{cx+1}{cx-1}\right)^2 - 4(ab - b^2) \log(cx + 1) + 4(ab + b^2) \log(cx - 1) + 4(a^2c^2x^2 + b^2c^2x) \log\left(-\frac{cx+1}{cx-1}\right)}{8c^2}$$

input `integrate(x*(a+b*arctanh(c*x))^2,x, algorithm="fricas")`

output $\frac{1}{8} * (4 * a^2 * c^2 * x^2 + 8 * a * b * c * x + (b^2 * c^2 * x^2 - b^2) * \log\left(-\frac{c * x + 1}{c * x - 1}\right)^2 - 4 * (a * b - b^2) * \log(c * x + 1) + 4 * (a * b + b^2) * \log(c * x - 1) + 4 * (a * b * c^2 * x^2 + b^2 * c * x) * \log\left(-\frac{c * x + 1}{c * x - 1}\right)) / c^2$

Sympy [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.52

$$\int x(a + \operatorname{arctanh}(cx))^2 dx$$

$$= \begin{cases} \frac{a^2 x^2}{2} + abx^2 \operatorname{atanh}(cx) + \frac{abx}{c} - \frac{ab \operatorname{atanh}(cx)}{c^2} + \frac{b^2 x^2 \operatorname{atanh}^2(cx)}{2} + \frac{b^2 x \operatorname{atanh}(cx)}{c} + \frac{b^2 \log(x - \frac{1}{c})}{c^2} - \frac{b^2 \operatorname{atanh}^2(cx)}{2c^2} + \frac{b^2}{2c^2} \\ \frac{a^2 x^2}{2} \end{cases}$$

input `integrate(x*(a+b*atanh(c*x))**2,x)`

output `Piecewise((a**2*x**2/2 + a*b*x**2*atanh(c*x) + a*b*x/c - a*b*atanh(c*x)/c**2 + b**2*x**2*atanh(c*x)**2/2 + b**2*x*atanh(c*x)/c + b**2*log(x - 1/c)/c**2 - b**2*atanh(c*x)**2/(2*c**2) + b**2*atanh(c*x)/c**2, Ne(c, 0)), (a**2*x**2/2, True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 158 vs. 2(69) = 138.

Time = 0.03 (sec) , antiderivative size = 158, normalized size of antiderivative = 2.11

$$\int x(a + \operatorname{arctanh}(cx))^2 dx = \frac{1}{2} b^2 x^2 \operatorname{artanh}(cx)^2 + \frac{1}{2} a^2 x^2$$

$$+ \frac{1}{2} \left(2x^2 \operatorname{artanh}(cx) + c \left(\frac{2x}{c^2} - \frac{\log(cx+1)}{c^3} + \frac{\log(cx-1)}{c^3} \right) \right) ab$$

$$+ \frac{1}{8} \left(4c \left(\frac{2x}{c^2} - \frac{\log(cx+1)}{c^3} + \frac{\log(cx-1)}{c^3} \right) \operatorname{artanh}(cx) - \frac{2(\log(cx-1) - 2)\log(cx+1) - \log(cx+1)^2 - \log(cx-1)^2 - 4\log(cx-1)}{c^2} \right) b^2$$

input `integrate(x*(a+b*arctanh(c*x))^2,x, algorithm="maxima")`

output `1/2*b^2*x^2*arctanh(c*x)^2 + 1/2*a^2*x^2 + 1/2*(2*x^2*arctanh(c*x) + c*(2*x/c^2 - log(c*x + 1)/c^3 + log(c*x - 1)/c^3))*a*b + 1/8*(4*c*(2*x/c^2 - log(c*x + 1)/c^3 + log(c*x - 1)/c^3)*arctanh(c*x) - (2*(log(c*x - 1) - 2)*log(c*x + 1) - log(c*x + 1)^2 - log(c*x - 1)^2 - 4*log(c*x - 1))/c^2)*b^2`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 301 vs. $2(69) = 138$.

Time = 0.13 (sec) , antiderivative size = 301, normalized size of antiderivative = 4.01

$$\int x(a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{1}{2} \left(\frac{(cx+1)b^2 \log\left(-\frac{cx+1}{cx-1}\right)^2}{\left(\frac{(cx+1)^2 c^3}{(cx-1)^2} - \frac{2(cx+1)c^3}{cx-1} + c^3\right)(cx-1)} + \frac{2\left(\frac{2(cx+1)ab}{cx-1} + \frac{(cx+1)b^2}{cx-1} - b^2\right) \log\left(-\frac{cx+1}{cx-1}\right)}{\frac{(cx+1)^2 c^3}{(cx-1)^2} - \frac{2(cx+1)c^3}{cx-1} + c^3} + \frac{4\left(\frac{(cx+1)a^2}{cx-1} + \frac{(cx+1)ab}{cx-1} - a^2\right) \log\left(-\frac{cx+1}{cx-1}\right)}{\frac{(cx+1)^2 c^3}{(cx-1)^2} - \frac{2(cx+1)c^3}{cx-1} + c^3} \right)$$

input `integrate(x*(a+b*arctanh(c*x))^2,x, algorithm="giac")`

output `1/2*((c*x + 1)*b^2*log(-(c*x + 1)/(c*x - 1))^2/(((c*x + 1)^2*c^3/(c*x - 1)^2 - 2*(c*x + 1)*c^3/(c*x - 1) + c^3)*(c*x - 1)) + 2*(2*(c*x + 1)*a*b/(c*x - 1) + (c*x + 1)*b^2/(c*x - 1) - b^2)*log(-(c*x + 1)/(c*x - 1))/((c*x + 1)^2*c^3/(c*x - 1)^2 - 2*(c*x + 1)*c^3/(c*x - 1) + c^3) + 4*((c*x + 1)*a^2/(c*x - 1) + (c*x + 1)*a*b/(c*x - 1) - a^2)/((c*x + 1)^2*c^3/(c*x - 1)^2 - 2*(c*x + 1)*c^3/(c*x - 1) + c^3) - 2*b^2*log(-(c*x + 1)/(c*x - 1) + 1)/c^3 + 2*b^2*log(-(c*x + 1)/(c*x - 1))/c^3)*c`

Mupad [B] (verification not implemented)

Time = 3.85 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.19

$$\int x(a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{a^2 x^2}{2} - \frac{b^2 \operatorname{atanh}(cx)^2}{2} - \frac{b^2 \ln(c^2 x^2 - 1)}{2} - \frac{c(x \operatorname{atanh}(cx) b^2 + a x b) + a b \operatorname{atanh}(cx)}{c^2} + \frac{b^2 x^2 \operatorname{atanh}(cx)^2}{2} + a b x^2 \operatorname{atanh}(cx)$$

input `int(x*(a + b*atanh(c*x))^2,x)`

output `(a^2*x^2)/2 - ((b^2*atanh(c*x)^2)/2 - (b^2*log(c^2*x^2 - 1))/2 - c*(b^2*x*atanh(c*x) + a*b*x) + a*b*atanh(c*x))/c^2 + (b^2*x^2*atanh(c*x)^2)/2 + a*b*x^2*atanh(c*x)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.41

$$\int x(a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{\operatorname{atanh}(cx)^2 b^2 c^2 x^2 - \operatorname{atanh}(cx)^2 b^2 + 2 \operatorname{atanh}(cx) ab c^2 x^2 - 2 \operatorname{atanh}(cx) ab + 2 \operatorname{atanh}(cx) b^2 cx + 2 \operatorname{atanh}(cx) a^2}{2c^2}$$

input

```
int(x*(a+b*atanh(c*x))^2,x)
```

output

```
(atanh(c*x)**2*b**2*c**2*x**2 - atanh(c*x)**2*b**2 + 2*atanh(c*x)*a*b*c**2*x**2 - 2*atanh(c*x)*a*b + 2*atanh(c*x)*b**2*c*x + 2*atanh(c*x)*b**2 + 2*log(c**2*x - c)*b**2 + a**2*c**2*x**2 + 2*a*b*c*x)/(2*c**2)
```

3.18 $\int (a + b \operatorname{arctanh}(cx))^2 dx$

Optimal result	226
Mathematica [A] (verified)	226
Rubi [A] (verified)	227
Maple [A] (verified)	229
Fricas [F]	229
Sympy [F]	230
Maxima [F]	230
Giac [F]	230
Mupad [F(-1)]	231
Reduce [F]	231

Optimal result

Integrand size = 10, antiderivative size = 74

$$\int (a + b \operatorname{arctanh}(cx))^2 dx = \frac{(a + b \operatorname{arctanh}(cx))^2}{c} + x(a + b \operatorname{arctanh}(cx))^2 - \frac{2b(a + b \operatorname{arctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{c} - \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{c}$$

output

```
(a+b*arctanh(c*x))^2/c+x*(a+b*arctanh(c*x))^2-2*b*(a+b*arctanh(c*x))*ln(2/(-c*x+1))/c-b^2*polylog(2,1-2/(-c*x+1))/c
```

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.11

$$\int (a + b \operatorname{arctanh}(cx))^2 dx = \frac{b^2(-1 + cx) \operatorname{arctanh}(cx)^2 + 2b \operatorname{arctanh}(cx) (acx - b \log(1 + e^{-2 \operatorname{arctanh}(cx)})) + a(acx + b \log(1 - c^2 x^2))}{c}$$

input

```
Integrate[(a + b*ArcTanh[c*x])^2,x]
```

output

```
(b^2*(-1 + c*x)*ArcTanh[c*x]^2 + 2*b*ArcTanh[c*x]*(a*c*x - b*Log[1 + E^(-2
*ArcTanh[c*x]])) + a*(a*c*x + b*Log[1 - c^2*x^2]) + b^2*PolyLog[2, -E^(-2*
ArcTanh[c*x]]))/c
```

Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.19, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6436, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + b \operatorname{arctanh}(cx))^2 dx \\
 & \quad \downarrow \text{6436} \\
 & x(a + b \operatorname{arctanh}(cx))^2 - 2bc \int \frac{x(a + b \operatorname{arctanh}(cx))}{1 - c^2 x^2} dx \\
 & \quad \downarrow \text{6546} \\
 & x(a + b \operatorname{arctanh}(cx))^2 - 2bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(cx)}{1 - cx} dx}{c} - \frac{(a + b \operatorname{arctanh}(cx))^2}{2bc^2} \right) \\
 & \quad \downarrow \text{6470} \\
 & 2bc \left(\frac{x(a + b \operatorname{arctanh}(cx))^2 - \frac{\log\left(\frac{2}{1 - cx}\right)(a + b \operatorname{arctanh}(cx))}{c} - b \int \frac{\log\left(\frac{2}{1 - cx}\right)}{1 - c^2 x^2} dx}{c} - \frac{(a + b \operatorname{arctanh}(cx))^2}{2bc^2} \right) \\
 & \quad \downarrow \text{2849} \\
 & 2bc \left(\frac{x(a + b \operatorname{arctanh}(cx))^2 - \frac{b \int \frac{\log\left(\frac{2}{1 - cx}\right)}{1 - c^2 x^2} dx \frac{1}{1 - cx} + \frac{\log\left(\frac{2}{1 - cx}\right)(a + b \operatorname{arctanh}(cx))}{c}}{c} - \frac{(a + b \operatorname{arctanh}(cx))^2}{2bc^2} \right) \\
 & \quad \downarrow \text{2752}
 \end{aligned}$$

$$2bc \left(\frac{\frac{\log\left(\frac{2}{1-cx}\right)(a+\operatorname{barctanh}(cx))}{c} + \frac{b \operatorname{PolyLog}\left(2, 1-\frac{2}{1-cx}\right)}{2c}}{c} - \frac{(a + \operatorname{barctanh}(cx))^2}{2bc^2} \right)$$

input `Int[(a + b*ArcTanh[c*x])^2,x]`

output `x*(a + b*ArcTanh[c*x])^2 - 2*b*c*(-1/2*(a + b*ArcTanh[c*x])^2/(b*c^2) + ((a + b*ArcTanh[c*x])*Log[2/(1 - c*x)]/c + (b*PolyLog[2, 1 - 2/(1 - c*x)])/(2*c))/c)`

Definitions of rubi rules used

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 6436 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.), x_Symbol] := Simp[x*(a + b*ArcTanh[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`

rule 6470 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6546

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)*(x_.)/((d_.) + (e_.)*(x_.)^2),
 x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/
 (c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.55

method	result
derivativedivides	$\frac{cx a^2 + b^2 \left(\operatorname{arctanh}(cx)^2 (cx-1) + 2 \operatorname{arctanh}(cx)^2 - 2 \operatorname{arctanh}(cx) \ln \left(1 + \frac{(cx+1)^2}{-c^2 x^2 + 1} \right) - \operatorname{polylog} \left(2, -\frac{(cx+1)^2}{-c^2 x^2 + 1} \right) \right) + 2abcx}{c}$
default	$\frac{cx a^2 + b^2 \left(\operatorname{arctanh}(cx)^2 (cx-1) + 2 \operatorname{arctanh}(cx)^2 - 2 \operatorname{arctanh}(cx) \ln \left(1 + \frac{(cx+1)^2}{-c^2 x^2 + 1} \right) - \operatorname{polylog} \left(2, -\frac{(cx+1)^2}{-c^2 x^2 + 1} \right) \right) + 2abcx}{c}$
parts	$x a^2 + \frac{b^2 \left(\operatorname{arctanh}(cx)^2 (cx-1) + 2 \operatorname{arctanh}(cx)^2 - 2 \operatorname{arctanh}(cx) \ln \left(1 + \frac{(cx+1)^2}{-c^2 x^2 + 1} \right) - \operatorname{polylog} \left(2, -\frac{(cx+1)^2}{-c^2 x^2 + 1} \right) \right)}{c} + 2$
risch	$\frac{b^2 \ln(cx+1)^2 x}{4} + x a^2 - \ln(-cx+1) abx + \frac{\ln(-cx+1) ab}{c} - \frac{b^2 \ln(-cx+1) \ln(cx+1)}{2c} + \frac{b^2 \ln(-\frac{cx}{2} + \frac{1}{2})}{c}$

input

```
int((a+b*arctanh(c*x))^2,x,method=_RETURNVERBOSE)
```

output

```
1/c*(c*x*a^2+b^2*(arctanh(c*x)^2*(c*x-1)+2*arctanh(c*x)^2-2*arctanh(c*x)*
ln(1+(c*x+1)^2/(-c^2*x^2+1))-polylog(2,-(c*x+1)^2/(-c^2*x^2+1)))+2*a*b*c*x*
arctanh(c*x)+a*b*ln(-c^2*x^2+1))
```

Fricas [F]

$$\int (a + b \operatorname{arctanh}(cx))^2 dx = \int (b \operatorname{arctanh}(cx) + a)^2 dx$$

input

```
integrate((a+b*arctanh(c*x))^2,x, algorithm="fricas")
```

output

```
integral(b^2*arctanh(c*x)^2 + 2*a*b*arctanh(c*x) + a^2, x)
```

Sympy [F]

$$\int (a + b \operatorname{arctanh}(cx))^2 dx = \int (a + b \operatorname{atanh}(cx))^2 dx$$

input `integrate((a+b*atanh(c*x))**2,x)`

output `Integral((a + b*atanh(c*x))**2, x)`

Maxima [F]

$$\int (a + b \operatorname{arctanh}(cx))^2 dx = \int (b \operatorname{artanh}(cx) + a)^2 dx$$

input `integrate((a+b*arctanh(c*x))^2,x, algorithm="maxima")`

output `-1/4*(c^2*(2*x/c^2 - log(c*x + 1)/c^3 + log(c*x - 1)/c^3) - 6*c*integrate(x*log(c*x + 1)/(c^2*x^2 - 1), x) - (c*x - 1)*(log(-c*x + 1)^2 - 2*log(-c*x + 1) + 2)/c - (c*x*log(c*x + 1)^2 + 2*(c*x - (c*x + 1)*log(c*x + 1))*log(-c*x + 1))/c + log(c^2*x^2 - 1)/c - 2*integrate(log(c*x + 1)/(c^2*x^2 - 1), x))*b^2 + a^2*x + (2*c*x*arctanh(c*x) + log(-c^2*x^2 + 1))*a*b/c`

Giac [F]

$$\int (a + b \operatorname{arctanh}(cx))^2 dx = \int (b \operatorname{artanh}(cx) + a)^2 dx$$

input `integrate((a+b*arctanh(c*x))^2,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + b \operatorname{arctanh}(cx))^2 dx = \int (a + b \operatorname{atanh}(cx))^2 dx$$

input `int((a + b*atanh(c*x))^2,x)`output `int((a + b*atanh(c*x))^2, x)`**Reduce [F]**

$$\int (a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{2 \operatorname{atanh}(cx) abcx + 2 \operatorname{atanh}(cx) ab + \left(\int \operatorname{atanh}(cx)^2 dx \right) b^2 c + 2 \log(c^2 x - c) ab + a^2 cx}{c}$$

input `int((a+b*atanh(c*x))^2,x)`output `(2*atanh(c*x)*a*b*c*x + 2*atanh(c*x)*a*b + int(atanh(c*x)**2,x)*b**2*c + 2*log(c**2*x - c)*a*b + a**2*c*x)/c`

3.19 $\int \frac{(a+b\operatorname{arctanh}(cx))^2}{x} dx$

Optimal result	232
Mathematica [C] (verified)	233
Rubi [A] (verified)	233
Maple [C] (warning: unable to verify)	235
Fricas [F]	237
Sympy [F]	237
Maxima [F]	237
Giac [F]	238
Mupad [F(-1)]	238
Reduce [F]	238

Optimal result

Integrand size = 14, antiderivative size = 117

$$\int \frac{(a + b\operatorname{arctanh}(cx))^2}{x} dx = 2(a + b\operatorname{arctanh}(cx))^2 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx}\right) - b(a + b\operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx}\right) + b(a + b\operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - cx}\right) + \frac{1}{2}b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - cx}\right) - \frac{1}{2}b^2 \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 - cx}\right)$$

output

```
-2*(a+b*arctanh(c*x))^2*arctanh(-1+2/(-c*x+1))-b*(a+b*arctanh(c*x))*polylog(2,1-2/(-c*x+1))+b*(a+b*arctanh(c*x))*polylog(2,-1+2/(-c*x+1))+1/2*b^2*polylog(3,1-2/(-c*x+1))-1/2*b^2*polylog(3,-1+2/(-c*x+1))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.40 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.29

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x} dx = a^2 \log(cx) + ab(-\operatorname{PolyLog}(2, -cx) + \operatorname{PolyLog}(2, cx))$$

$$+ b^2 \left(\frac{i\pi^3}{24} - \frac{2}{3} \operatorname{arctanh}(cx)^3 \right.$$

$$\quad - \operatorname{arctanh}(cx)^2 \log(1 + e^{-2\operatorname{arctanh}(cx)})$$

$$\quad + \operatorname{arctanh}(cx)^2 \log(1 - e^{2\operatorname{arctanh}(cx)})$$

$$+ \operatorname{arctanh}(cx) \operatorname{PolyLog}(2, -e^{-2\operatorname{arctanh}(cx)})$$

$$+ \operatorname{arctanh}(cx) \operatorname{PolyLog}(2, e^{2\operatorname{arctanh}(cx)})$$

$$\quad + \frac{1}{2} \operatorname{PolyLog}(3, -e^{-2\operatorname{arctanh}(cx)})$$

$$\quad \left. - \frac{1}{2} \operatorname{PolyLog}(3, e^{2\operatorname{arctanh}(cx)}) \right)$$

input

```
Integrate[(a + b*ArcTanh[c*x])^2/x,x]
```

output

```
a^2*Log[c*x] + a*b*(-PolyLog[2, -(c*x)] + PolyLog[2, c*x]) + b^2*((I/24)*P
i^3 - (2*ArcTanh[c*x]^3)/3 - ArcTanh[c*x]^2*Log[1 + E^(-2*ArcTanh[c*x])] +
ArcTanh[c*x]^2*Log[1 - E^(2*ArcTanh[c*x])] + ArcTanh[c*x]*PolyLog[2, -E^(-
2*ArcTanh[c*x])] + ArcTanh[c*x]*PolyLog[2, E^(2*ArcTanh[c*x])] + PolyLog[
3, -E^(-2*ArcTanh[c*x])]/2 - PolyLog[3, E^(2*ArcTanh[c*x])]/2)
```

Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.22,
 number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules
 used = {6448, 6614, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{(a + b \operatorname{arctanh}(cx))^2}{x} dx \\
& \quad \downarrow \text{6448} \\
& 2 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx}\right) (a + b \operatorname{arctanh}(cx))^2 - 4bc \int \frac{(a + b \operatorname{arctanh}(cx)) \operatorname{arctanh}\left(1 - \frac{2}{1 - cx}\right)}{1 - c^2 x^2} dx \\
& \quad \downarrow \text{6614} \\
& 2 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx}\right) (a + b \operatorname{arctanh}(cx))^2 - \\
& 4bc \left(\frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(cx)) \log\left(2 - \frac{2}{1 - cx}\right)}{1 - c^2 x^2} dx - \frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(cx)) \log\left(\frac{2}{1 - cx}\right)}{1 - c^2 x^2} dx \right) \\
& \quad \downarrow \text{6620} \\
& 2 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx}\right) (a + b \operatorname{arctanh}(cx))^2 - \\
& 4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx}\right) (a + b \operatorname{arctanh}(cx))}{2c} - \frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx}\right)}{1 - c^2 x^2} dx \right) + \frac{1}{2} \left(\frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, \frac{2}{1 - cx}\right)}{1 - c^2 x^2} dx \right) \right) \\
& \quad \downarrow \text{7164} \\
& 2 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx}\right) (a + b \operatorname{arctanh}(cx))^2 - \\
& 4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx}\right) (a + b \operatorname{arctanh}(cx))}{2c} - \frac{b \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - cx}\right)}{4c} \right) + \frac{1}{2} \left(\frac{b \operatorname{PolyLog}\left(3, \frac{2}{1 - cx} - 1\right)}{4c} \right) \right)
\end{aligned}$$

input `Int[(a + b*ArcTanh[c*x])^2/x,x]`

output `2*(a + b*ArcTanh[c*x])^2*ArcTanh[1 - 2/(1 - c*x)] - 4*b*c*(((a + b*ArcTanh[c*x])*PolyLog[2, 1 - 2/(1 - c*x)])/(2*c) - (b*PolyLog[3, 1 - 2/(1 - c*x)])/(4*c))/2 + (-1/2*((a + b*ArcTanh[c*x])*PolyLog[2, -1 + 2/(1 - c*x)]))/c + (b*PolyLog[3, -1 + 2/(1 - c*x)])/(4*c)/2`

Definitions of rubi rules used

rule 6448

```
Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_)/(x_), x_Symbol] := Simp[2*(a +
b*ArcTanh[c*x])^p*ArcTanh[1 - 2/(1 - c*x)], x] - Simp[2*b*c*p Int[(a + b
*ArcTanh[c*x])^(p - 1)*(ArcTanh[1 - 2/(1 - c*x)]/(1 - c^2*x^2)), x], x] /;
FreeQ[{a, b, c}, x] && IGtQ[p, 1]
```

rule 6614

```
Int[(ArcTanh[u_] * ((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.))/((d_) + (e_.)*(
x_)^2), x_Symbol] := Simp[1/2 Int[Log[1 + u]*((a + b*ArcTanh[c*x])^p/(d +
e*x^2)), x], x] - Simp[1/2 Int[Log[1 - u]*((a + b*ArcTanh[c*x])^p/(d + e
*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e,
0] && EqQ[u^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 6620

```
Int[(Log[u_] * ((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.))/((d_) + (e_.)*(x_)^
2), x_Symbol] := Simp[(- (a + b*ArcTanh[c*x])^p) * (PolyLog[2, 1 - u]/(2*c*d))
, x] + Simp[b*(p/2 Int[(a + b*ArcTanh[c*x])^(p - 1) * (PolyLog[2, 1 - u]/(
d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d
+ e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 5.32 (sec) , antiderivative size = 630, normalized size of antiderivative = 5.38

method	result
parts	$a^2 \ln(x) + b^2 \left(\ln(cx) \operatorname{arctanh}(cx)^2 - \operatorname{arctanh}(cx) \operatorname{polylog}\left(2, -\frac{(cx+1)^2}{-c^2x^2+1}\right) + \frac{\operatorname{polylog}(3, \dots)}{\dots} \right)$
derivativedivides	$a^2 \ln(cx) + b^2 \left(\ln(cx) \operatorname{arctanh}(cx)^2 - \operatorname{arctanh}(cx) \operatorname{polylog}\left(2, -\frac{(cx+1)^2}{-c^2x^2+1}\right) + \frac{\operatorname{polylog}(3, \dots)}{\dots} \right)$
default	$a^2 \ln(cx) + b^2 \left(\ln(cx) \operatorname{arctanh}(cx)^2 - \operatorname{arctanh}(cx) \operatorname{polylog}\left(2, -\frac{(cx+1)^2}{-c^2x^2+1}\right) + \frac{\operatorname{polylog}(3, \dots)}{\dots} \right)$

```
input int((a+b*arctanh(c*x))^2/x,x,method=_RETURNVERBOSE)
```

```
output a^2*ln(x)+b^2*(ln(c*x)*arctanh(c*x)^2-arctanh(c*x)*polylog(2,-(c*x+1)^2/(-c^2*x^2+1))+1/2*polylog(3,-(c*x+1)^2/(-c^2*x^2+1))-arctanh(c*x)^2*ln((c*x+1)^2/(-c^2*x^2+1)-1)+arctanh(c*x)^2*ln(1+(c*x+1)/(-c^2*x^2+1)^(1/2))+2*arctanh(c*x)*polylog(2,-(c*x+1)/(-c^2*x^2+1)^(1/2))-2*polylog(3,-(c*x+1)/(-c^2*x^2+1)^(1/2))+arctanh(c*x)^2*ln(1-(c*x+1)/(-c^2*x^2+1)^(1/2))+2*arctanh(c*x)*polylog(2,(c*x+1)/(-c^2*x^2+1)^(1/2))-2*polylog(3,(c*x+1)/(-c^2*x^2+1)^(1/2))+1/2*I*Pi*csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1)/(1-(c*x+1)^2/(c^2*x^2-1)))*(csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1))*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1))))-csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1))*csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1)/(1-(c*x+1)^2/(c^2*x^2-1)))-csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1)/(1-(c*x+1)^2/(c^2*x^2-1)))+csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))+csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^2)*arctanh(c*x)+2*a*b*(ln(c*x)*arctanh(c*x)-1/2*dilog(c*x+1)-1/2*ln(c*x)*ln(c*x+1)-1/2*dilog(c*x))
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x))^2/x,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x)^2 + 2*a*b*arctanh(c*x) + a^2)/x, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x} dx = \int \frac{(a + b \operatorname{atanh}(cx))^2}{x} dx$$

input `integrate((a+b*atanh(c*x))**2/x,x)`

output `Integral((a + b*atanh(c*x))**2/x, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x))^2/x,x, algorithm="maxima")`

output `a^2*log(x) + integrate(1/4*b^2*(log(c*x + 1) - log(-c*x + 1))^2/x + a*b*(log(c*x + 1) - log(-c*x + 1))/x, x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x))^2/x,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^2/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x} dx = \int \frac{(a + b \operatorname{atanh}(cx))^2}{x} dx$$

input `int((a + b*atanh(c*x))^2/x,x)`

output `int((a + b*atanh(c*x))^2/x, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x} dx = 2 \left(\int \frac{\operatorname{atanh}(cx)}{x} dx \right) ab + \left(\int \frac{\operatorname{atanh}(cx)^2}{x} dx \right) b^2 + \log(x) a^2$$

input `int((a+b*atanh(c*x))^2/x,x)`

output `2*int(atanh(c*x)/x,x)*a*b + int(atanh(c*x)**2/x,x)*b**2 + log(x)*a**2`

3.20 $\int \frac{(a+b\operatorname{arctanh}(cx))^2}{x^2} dx$

Optimal result	239
Mathematica [A] (verified)	239
Rubi [A] (verified)	240
Maple [B] (verified)	242
Fricas [F]	242
Sympy [F]	243
Maxima [F]	243
Giac [F]	243
Mupad [F(-1)]	244
Reduce [F]	244

Optimal result

Integrand size = 14, antiderivative size = 71

$$\int \frac{(a + b\operatorname{arctanh}(cx))^2}{x^2} dx = c(a + b\operatorname{arctanh}(cx))^2 - \frac{(a + b\operatorname{arctanh}(cx))^2}{x} + 2bc(a + b\operatorname{arctanh}(cx)) \log\left(2 - \frac{2}{1 + cx}\right) - b^2c \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + cx}\right)$$

output

```
c*(a+b*arctanh(c*x))^2-(a+b*arctanh(c*x))^2/x+2*b*c*(a+b*arctanh(c*x))*ln(2-2/(c*x+1))-b^2*c*polylog(2,-1+2/(c*x+1))
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.32

$$\int \frac{(a + b\operatorname{arctanh}(cx))^2}{x^2} dx = \frac{b^2(-1 + cx)\operatorname{arctanh}(cx)^2 + 2b\operatorname{arctanh}(cx) (-a + bcx \log(1 - e^{-2\operatorname{arctanh}(cx)})) - a(a - 2bcx \log(cx) + bcx^2)}{x}$$

input

```
Integrate[(a + b*ArcTanh[c*x])^2/x^2,x]
```

output

```
(b^2*(-1 + c*x)*ArcTanh[c*x]^2 + 2*b*ArcTanh[c*x]*(-a + b*c*x*Log[1 - E^(-2*ArcTanh[c*x])]) - a*(a - 2*b*c*x*Log[c*x] + b*c*x*Log[1 - c^2*x^2]) - b^2*c*x*PolyLog[2, E^(-2*ArcTanh[c*x])])/x
```

Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.08, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6452, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^2} dx$$

$$\downarrow 6452$$

$$2bc \int \frac{a + b \operatorname{arctanh}(cx)}{x(1 - c^2x^2)} dx - \frac{(a + b \operatorname{arctanh}(cx))^2}{x}$$

$$\downarrow 6550$$

$$2bc \left(\int \frac{a + b \operatorname{arctanh}(cx)}{x(cx + 1)} dx + \frac{(a + b \operatorname{arctanh}(cx))^2}{2b} \right) - \frac{(a + b \operatorname{arctanh}(cx))^2}{x}$$

$$\downarrow 6494$$

$$2bc \left(-bc \int \frac{\log \left(2 - \frac{2}{cx+1} \right)}{1 - c^2x^2} dx + \frac{(a + b \operatorname{arctanh}(cx))^2}{2b} + \log \left(2 - \frac{2}{cx+1} \right) (a + b \operatorname{arctanh}(cx)) \right) - \frac{(a + b \operatorname{arctanh}(cx))^2}{x}$$

$$\downarrow 2897$$

$$2bc \left(\frac{(a + b \operatorname{arctanh}(cx))^2}{2b} + \log \left(2 - \frac{2}{cx+1} \right) (a + b \operatorname{arctanh}(cx)) - \frac{1}{2} b \operatorname{PolyLog} \left(2, \frac{2}{cx+1} - 1 \right) \right) - \frac{(a + b \operatorname{arctanh}(cx))^2}{x}$$

input `Int[(a + b*ArcTanh[c*x])^2/x^2,x]`

output `-((a + b*ArcTanh[c*x])^2/x) + 2*b*c*((a + b*ArcTanh[c*x])^2/(2*b) + (a + b*ArcTanh[c*x])*Log[2 - 2/(1 + c*x)] - (b*PolyLog[2, -1 + 2/(1 + c*x)]))/2)`

Defintions of rubi rules used

rule 2897 `Int[Log[u_]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6494 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((x_)*((d_) + (e_)*(x_))), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6550 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((x_)*((d_) + (e_)*(x_)^2)), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 194 vs. $2(71) = 142$.

Time = 0.53 (sec) , antiderivative size = 195, normalized size of antiderivative = 2.75

method	result
parts	$-\frac{a^2}{x} + b^2 c \left(-\frac{\operatorname{arctanh}(cx)^2}{cx} + 2 \ln(cx) \operatorname{arctanh}(cx) - \operatorname{arctanh}(cx) \ln(cx+1) - \operatorname{arctanh}(cx) \ln(cx-1) \right)$
derivativedivides	$c \left(-\frac{a^2}{cx} + b^2 \left(-\frac{\operatorname{arctanh}(cx)^2}{cx} + 2 \ln(cx) \operatorname{arctanh}(cx) - \operatorname{arctanh}(cx) \ln(cx+1) - \operatorname{arctanh}(cx) \ln(cx-1) \right) \right)$
default	$c \left(-\frac{a^2}{cx} + b^2 \left(-\frac{\operatorname{arctanh}(cx)^2}{cx} + 2 \ln(cx) \operatorname{arctanh}(cx) - \operatorname{arctanh}(cx) \ln(cx+1) - \operatorname{arctanh}(cx) \ln(cx-1) \right) \right)$

input `int((a+b*arctanh(c*x))^2/x^2,x,method=_RETURNVERBOSE)`

output `-1/x*a^2+b^2*c*(-1/c/x*arctanh(c*x)^2+2*ln(c*x)*arctanh(c*x)-arctanh(c*x)*ln(c*x+1)-arctanh(c*x)*ln(c*x-1)-1/4*ln(c*x-1)^2+dilog(1/2*c*x+1/2)+1/2*ln(c*x-1)*ln(1/2*c*x+1/2)+1/4*ln(c*x+1)^2-1/2*(ln(c*x+1)-ln(1/2*c*x+1/2))*ln(-1/2*c*x+1/2)-dilog(c*x+1)-ln(c*x)*ln(c*x+1)-dilog(c*x))+2*a*b*c*(-1/c/x*arctanh(c*x)+ln(c*x)-1/2*ln(c*x+1)-1/2*ln(c*x-1))`

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^2} dx = \int \frac{(b \operatorname{arctanh}(cx) + a)^2}{x^2} dx$$

input `integrate((a+b*arctanh(c*x))^2/x^2,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x)^2 + 2*a*b*arctanh(c*x) + a^2)/x^2, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^2} dx = \int \frac{(a + b \operatorname{atanh}(cx))^2}{x^2} dx$$

input `integrate((a+b*atanh(c*x))**2/x**2,x)`

output `Integral((a + b*atanh(c*x))**2/x**2, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^2} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^2}{x^2} dx$$

input `integrate((a+b*arctanh(c*x))^2/x^2,x, algorithm="maxima")`

output `-(c*(log(c^2*x^2 - 1) - log(x^2)) + 2*arctanh(c*x)/x)*a*b - 1/4*b^2*(log(-c*x + 1)^2/x + integrate(-((c*x - 1)*log(c*x + 1)^2 + 2*(c*x - (c*x - 1)*log(c*x + 1))*log(-c*x + 1))/(c*x^3 - x^2), x)) - a^2/x`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^2} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^2}{x^2} dx$$

input `integrate((a+b*arctanh(c*x))^2/x^2,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^2/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^2} dx = \int \frac{(a + b \operatorname{atanh}(cx))^2}{x^2} dx$$

input `int((a + b*atanh(c*x))^2/x^2,x)`output `int((a + b*atanh(c*x))^2/x^2, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^2} dx$$

$$= \frac{-\operatorname{atanh}(cx)^2 b^2 - 2 \operatorname{atanh}(cx) abcx - 2 \operatorname{atanh}(cx) ab - 2 \left(\int \frac{\operatorname{atanh}(cx)}{c^2 x^3 - x} dx \right) b^2 cx - 2 \log(c^2 x - c) abcx + 2 \log(c^2 x - c) ab}{x}$$

input `int((a+b*atanh(c*x))^2/x^2,x)`output `(- atanh(c*x)**2*b**2 - 2*atanh(c*x)*a*b*c*x - 2*atanh(c*x)*a*b - 2*int(a
tanh(c*x)/(c**2*x**3 - x),x)*b**2*c*x - 2*log(c**2*x - c)*a*b*c*x + 2*log(
x)*a*b*c*x - a**2)/x`

3.21 $\int \frac{(a+b\operatorname{arctanh}(cx))^2}{x^3} dx$

Optimal result	245
Mathematica [A] (verified)	245
Rubi [A] (verified)	246
Maple [A] (verified)	248
Fricas [A] (verification not implemented)	249
Sympy [A] (verification not implemented)	249
Maxima [B] (verification not implemented)	250
Giac [B] (verification not implemented)	250
Mupad [B] (verification not implemented)	251
Reduce [B] (verification not implemented)	252

Optimal result

Integrand size = 14, antiderivative size = 80

$$\int \frac{(a + b\operatorname{arctanh}(cx))^2}{x^3} dx = -\frac{bc(a + b\operatorname{arctanh}(cx))}{x} + \frac{1}{2}c^2(a + b\operatorname{arctanh}(cx))^2 - \frac{(a + b\operatorname{arctanh}(cx))^2}{2x^2} + b^2c^2 \log(x) - \frac{1}{2}b^2c^2 \log(1 - c^2x^2)$$

output

```
-b*c*(a+b*arctanh(c*x))/x+1/2*c^2*(a+b*arctanh(c*x))^2-1/2*(a+b*arctanh(c*x))^2/x^2+b^2*c^2*ln(x)-1/2*b^2*c^2*ln(-c^2*x^2+1)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.26

$$\int \frac{(a + b\operatorname{arctanh}(cx))^2}{x^3} dx = \frac{a^2 + 2abcx + 2b(a + bcx)\operatorname{arctanh}(cx) - b^2(-1 + c^2x^2)\operatorname{arctanh}(cx)^2 - 2b^2c^2x^2 \log(x) + b(a + b)c^2x^2 \log(1 - c^2x^2)}{2x^2}$$

input

```
Integrate[(a + b*ArcTanh[c*x])^2/x^3,x]
```

output

$$-1/2*(a^2 + 2*a*b*c*x + 2*b*(a + b*c*x)*\text{ArcTanh}[c*x] - b^2*(-1 + c^2*x^2)*\text{ArcTanh}[c*x]^2 - 2*b^2*c^2*x^2*\text{Log}[x] + b*(a + b)*c^2*x^2*\text{Log}[1 - c*x] - (a - b)*b*c^2*x^2*\text{Log}[1 + c*x])/x^2$$
Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.96, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {6452, 6544, 6452, 243, 47, 14, 16, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^3} dx \\
 & \quad \downarrow \text{6452} \\
 & bc \int \frac{a + b \operatorname{arctanh}(cx)}{x^2(1 - c^2x^2)} dx - \frac{(a + b \operatorname{arctanh}(cx))^2}{2x^2} \\
 & \quad \downarrow \text{6544} \\
 & bc \left(c^2 \int \frac{a + b \operatorname{arctanh}(cx)}{1 - c^2x^2} dx + \int \frac{a + b \operatorname{arctanh}(cx)}{x^2} dx \right) - \frac{(a + b \operatorname{arctanh}(cx))^2}{2x^2} \\
 & \quad \downarrow \text{6452} \\
 & bc \left(c^2 \int \frac{a + b \operatorname{arctanh}(cx)}{1 - c^2x^2} dx + bc \int \frac{1}{x(1 - c^2x^2)} dx - \frac{a + b \operatorname{arctanh}(cx)}{x} \right) - \\
 & \quad \frac{(a + b \operatorname{arctanh}(cx))^2}{2x^2} \\
 & \quad \downarrow \text{243} \\
 & bc \left(c^2 \int \frac{a + b \operatorname{arctanh}(cx)}{1 - c^2x^2} dx + \frac{1}{2} bc \int \frac{1}{x^2(1 - c^2x^2)} dx^2 - \frac{a + b \operatorname{arctanh}(cx)}{x} \right) - \\
 & \quad \frac{(a + b \operatorname{arctanh}(cx))^2}{2x^2} \\
 & \quad \downarrow \text{47} \\
 & bc \left(c^2 \int \frac{a + b \operatorname{arctanh}(cx)}{1 - c^2x^2} dx + \frac{1}{2} bc \left(c^2 \int \frac{1}{1 - c^2x^2} dx^2 + \int \frac{1}{x^2} dx^2 \right) - \frac{a + b \operatorname{arctanh}(cx)}{x} \right) - \\
 & \quad \frac{(a + b \operatorname{arctanh}(cx))^2}{2x^2}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 14 \\
bc \left(c^2 \int \frac{a + \operatorname{arctanh}(cx)}{1 - c^2 x^2} dx + \frac{1}{2} bc \left(c^2 \int \frac{1}{1 - c^2 x^2} dx^2 + \log(x^2) \right) - \frac{a + \operatorname{arctanh}(cx)}{x} \right) - \\
& \frac{(a + \operatorname{arctanh}(cx))^2}{2x^2} \\
& \downarrow 16 \\
bc \left(c^2 \int \frac{a + \operatorname{arctanh}(cx)}{1 - c^2 x^2} dx - \frac{a + \operatorname{arctanh}(cx)}{x} + \frac{1}{2} bc (\log(x^2) - \log(1 - c^2 x^2)) \right) - \\
& \frac{(a + \operatorname{arctanh}(cx))^2}{2x^2} \\
& \downarrow 6510 \\
bc \left(\frac{c(a + \operatorname{arctanh}(cx))^2}{2b} - \frac{a + \operatorname{arctanh}(cx)}{x} + \frac{1}{2} bc (\log(x^2) - \log(1 - c^2 x^2)) \right) - \\
& \frac{(a + \operatorname{arctanh}(cx))^2}{2x^2}
\end{aligned}$$

input `Int[(a + b*ArcTanh[c*x])^2/x^3,x]`

output `-1/2*(a + b*ArcTanh[c*x])^2/x^2 + b*c*(-((a + b*ArcTanh[c*x])/x) + (c*(a + b*ArcTanh[c*x])^2)/(2*b) + (b*c*(Log[x^2] - Log[1 - c^2*x^2]))/2)`

Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

```
rule 243 Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]
```

```
rule 6452 Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]
```

```
rule 6510 Int[((a_.) + ArcTanh[(c_.)*(x_)*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]
```

```
rule 6544 Int((((a_.) + ArcTanh[(c_.)*(x_)*(b_.))^(p_.)*((f_.)*(x_)^(m_)))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.68

method	result
paralelrisch	$\frac{b^2 \operatorname{arctanh}(cx)^2 x^2 c^2 + 2b^2 c^2 \ln(x) x^2 - 2 \ln(cx-1) x^2 b^2 c^2 + 2x^2 \operatorname{arctanh}(cx) ab c^2 - 2x^2 \operatorname{arctanh}(cx) b^2 c^2 - a^2 c^2 x^2 - 2b^2 \operatorname{arctanh}(cx) c^2}{2x^2}$
parts	$-\frac{a^2}{2x^2} + b^2 c^2 \left(-\frac{\operatorname{arctanh}(cx)^2}{2c^2 x^2} + \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{2} - \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{2} - \frac{\operatorname{arctanh}(cx)}{cx} - \frac{\ln(cx-1)}{8} \right)$
derivativedivides	$c^2 \left(-\frac{a^2}{2c^2 x^2} + b^2 \left(-\frac{\operatorname{arctanh}(cx)^2}{2c^2 x^2} + \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{2} - \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{2} - \frac{\operatorname{arctanh}(cx)}{cx} - \frac{\ln(cx-1)}{8} \right) \right)$
default	$c^2 \left(-\frac{a^2}{2c^2 x^2} + b^2 \left(-\frac{\operatorname{arctanh}(cx)^2}{2c^2 x^2} + \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{2} - \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{2} - \frac{\operatorname{arctanh}(cx)}{cx} - \frac{\ln(cx-1)}{8} \right) \right)$
risch	$\frac{b^2 (c^2 x^2 - 1) \ln(cx+1)^2}{8x^2} - \frac{b(x^2 b \ln(-cx+1) c^2 + 2bcx - b \ln(-cx+1) + 2a) \ln(cx+1)}{4x^2} + \frac{b^2 c^2 x^2 \ln(-cx+1)^2 + 4b c^2 \ln(-cx+1)}{4x^2}$

```
input int((a+b*arctanh(c*x))^2/x^3,x,method=_RETURNVERBOSE)
```

output

```
1/2*(b^2*arctanh(c*x)^2*x^2*c^2+2*b^2*c^2*ln(x)*x^2-2*ln(c*x-1)*x^2*b^2*c^2+2*x^2*arctanh(c*x)*a*b*c^2-2*x^2*arctanh(c*x)*b^2*c^2-a^2*c^2*x^2-2*b^2*arctanh(c*x)*x*c-2*a*b*c*x-b^2*arctanh(c*x)^2-2*arctanh(c*x)*a*b-a^2)/x^2
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.69

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^3} dx$$

$$= \frac{8b^2c^2x^2 \log(x) + 4(ab - b^2)c^2x^2 \log(cx + 1) - 4(ab + b^2)c^2x^2 \log(cx - 1) - 8abcx + (b^2c^2x^2 - b^2) \log}{8x^2}$$

input

```
integrate((a+b*arctanh(c*x))^2/x^3,x, algorithm="fricas")
```

output

```
1/8*(8*b^2*c^2*x^2*log(x) + 4*(a*b - b^2)*c^2*x^2*log(c*x + 1) - 4*(a*b + b^2)*c^2*x^2*log(c*x - 1) - 8*a*b*c*x + (b^2*c^2*x^2 - b^2)*log(-(c*x + 1)/(c*x - 1))^2 - 4*a^2 - 4*(b^2*c*x + a*b)*log(-(c*x + 1)/(c*x - 1)))/x^2
```

Sympy [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.58

$$\int \frac{(a + b \operatorname{atanh}(cx))^2}{x^3} dx$$

$$= \begin{cases} -\frac{a^2}{2x^2} + abc^2 \operatorname{atanh}(cx) - \frac{abc}{x} - \frac{ab \operatorname{atanh}(cx)}{x^2} + b^2c^2 \log(x) - b^2c^2 \log\left(x - \frac{1}{c}\right) + \frac{b^2c^2 \operatorname{atanh}^2(cx)}{2} - b^2c^2 \operatorname{atanh} \\ -\frac{a^2}{2x^2} \end{cases}$$

input

```
integrate((a+b*atanh(c*x))**2/x**3,x)
```

output

```
Piecewise((-a**2/(2*x**2) + a*b*c**2*atanh(c*x) - a*b*c/x - a*b*atanh(c*x)/x**2 + b**2*c**2*log(x) - b**2*c**2*log(x - 1/c) + b**2*c**2*atanh(c*x)**2/2 - b**2*c**2*atanh(c*x) - b**2*c*atanh(c*x)/x - b**2*atanh(c*x)**2/(2*x**2), Ne(c, 0)), (-a**2/(2*x**2), True))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 151 vs. $2(74) = 148$.

Time = 0.03 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.89

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^3} dx$$

$$= \frac{1}{2} \left(\left(c \log(cx + 1) - c \log(cx - 1) - \frac{2}{x} \right) c - \frac{2 \operatorname{artanh}(cx)}{x^2} \right) ab$$

$$+ \frac{1}{8} \left((2 \log(cx - 1) - 2) \log(cx + 1) - \log(cx + 1)^2 - \log(cx - 1)^2 - 4 \log(cx - 1) + 8 \log(x) \right) c^2 +$$

$$- \frac{b^2 \operatorname{artanh}(cx)^2}{2x^2} - \frac{a^2}{2x^2}$$

input `integrate((a+b*arctanh(c*x))^2/x^3,x, algorithm="maxima")`

output `1/2*((c*log(c*x + 1) - c*log(c*x - 1) - 2/x)*c - 2*arctanh(c*x)/x^2)*a*b + 1/8*((2*(log(c*x - 1) - 2)*log(c*x + 1) - log(c*x + 1)^2 - log(c*x - 1)^2 - 4*log(c*x - 1) + 8*log(x))*c^2 + 4*(c*log(c*x + 1) - c*log(c*x - 1) - 2/x)*c*arctanh(c*x))*b^2 - 1/2*b^2*arctanh(c*x)^2/x^2 - 1/2*a^2/x^2`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 278 vs. $2(74) = 148$.

Time = 0.13 (sec) , antiderivative size = 278, normalized size of antiderivative = 3.48

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^3} dx$$

$$= \frac{1}{2} \left(2b^2c \log \left(-\frac{cx + 1}{cx - 1} - 1 \right) - 2b^2c \log \left(-\frac{cx + 1}{cx - 1} \right) + \frac{(cx + 1)b^2c \log \left(-\frac{cx + 1}{cx - 1} \right)^2}{(cx - 1) \left(\frac{(cx + 1)^2}{(cx - 1)^2} + \frac{2(cx + 1)}{cx - 1} + 1 \right)} + \frac{2 \left(\frac{2(cx + 1)abc}{cx - 1} \right)}{\left(\frac{cx + 1}{cx - 1} \right)^2} \right)$$

input `integrate((a+b*arctanh(c*x))^2/x^3,x, algorithm="giac")`

output

```

1/2*(2*b^2*c*log(-(c*x + 1)/(c*x - 1) - 1) - 2*b^2*c*log(-(c*x + 1)/(c*x -
1)) + (c*x + 1)*b^2*c*log(-(c*x + 1)/(c*x - 1))^2/((c*x - 1)*((c*x + 1)^2
/(c*x - 1)^2 + 2*(c*x + 1)/(c*x - 1) + 1)) + 2*(2*(c*x + 1)*a*b*c/(c*x - 1
) + (c*x + 1)*b^2*c/(c*x - 1) + b^2*c)*log(-(c*x + 1)/(c*x - 1))/((c*x + 1
)^2/(c*x - 1)^2 + 2*(c*x + 1)/(c*x - 1) + 1) + 4*((c*x + 1)*a^2*c/(c*x - 1
) + (c*x + 1)*a*b*c/(c*x - 1) + a*b*c)/((c*x + 1)^2/(c*x - 1)^2 + 2*(c*x +
1)/(c*x - 1) + 1))*c

```

Mupad [B] (verification not implemented)

Time = 4.21 (sec) , antiderivative size = 246, normalized size of antiderivative = 3.08

$$\begin{aligned}
\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^3} dx = & \frac{b^2 c^2 \ln(cx + 1)^2}{8} - \frac{a^2}{2x^2} + \frac{b^2 c^2 \ln(1 - cx)^2}{8} \\
& - \frac{b^2 \ln(cx + 1)^2}{8x^2} - \frac{b^2 \ln(1 - cx)^2}{8x^2} + b^2 c^2 \ln(x) \\
& - \frac{b^2 c^2 \ln(cx - 1)}{2} - \frac{b^2 c^2 \ln(cx + 1)}{2} - \frac{ab \ln(cx + 1)}{2x^2} \\
& + \frac{ab \ln(1 - cx)}{2x^2} + \frac{b^2 \ln(cx + 1) \ln(1 - cx)}{4x^2} - \frac{abc}{x} \\
& - \frac{b^2 c \ln(cx + 1)}{2x} + \frac{b^2 c \ln(1 - cx)}{2x} - \frac{abc^2 \ln(cx - 1)}{2} \\
& + \frac{abc^2 \ln(cx + 1)}{2} - \frac{b^2 c^2 \ln(cx + 1) \ln(1 - cx)}{4}
\end{aligned}$$

input

```
int((a + b*atanh(c*x))^2/x^3,x)
```

output

```

(b^2*c^2*log(c*x + 1)^2)/8 - a^2/(2*x^2) + (b^2*c^2*log(1 - c*x)^2)/8 - (b
^2*log(c*x + 1)^2)/(8*x^2) - (b^2*log(1 - c*x)^2)/(8*x^2) + b^2*c^2*log(x)
- (b^2*c^2*log(c*x - 1))/2 - (b^2*c^2*log(c*x + 1))/2 - (a*b*log(c*x + 1)
)/(2*x^2) + (a*b*log(1 - c*x))/(2*x^2) + (b^2*log(c*x + 1)*log(1 - c*x))/(
4*x^2) - (a*b*c)/x - (b^2*c*log(c*x + 1))/(2*x) + (b^2*c*log(1 - c*x))/(2*
x) - (a*b*c^2*log(c*x - 1))/2 + (a*b*c^2*log(c*x + 1))/2 - (b^2*c^2*log(c*
x + 1)*log(1 - c*x))/4

```


Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.58

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^3} dx$$

$$= \frac{\operatorname{atanh}(cx)^2 b^2 c^2 x^2 - \operatorname{atanh}(cx)^2 b^2 + 2 \operatorname{atanh}(cx) ab c^2 x^2 - 2 \operatorname{atanh}(cx) ab - 2 \operatorname{atanh}(cx) b^2 c^2 x^2 - 2 \operatorname{atanh}(cx) b^2 c^2}{2x^2}$$

input `int((a+b*atanh(c*x))^2/x^3,x)`output `(atanh(c*x)**2*b**2*c**2*x**2 - atanh(c*x)**2*b**2 + 2*atanh(c*x)*a*b*c**2*x**2 - 2*atanh(c*x)*a*b - 2*atanh(c*x)*b**2*c**2*x**2 - 2*atanh(c*x)*b**2*c*x - 2*log(c**2*x - c)*b**2*c**2*x**2 + 2*log(x)*b**2*c**2*x**2 - a**2 - 2*a*b*c*x)/(2*x**2)`

3.22 $\int \frac{(a+b\operatorname{arctanh}(cx))^2}{x^4} dx$

Optimal result	253
Mathematica [A] (verified)	254
Rubi [A] (verified)	254
Maple [B] (verified)	257
Fricas [F]	258
Sympy [F]	258
Maxima [F]	258
Giac [F]	259
Mupad [F(-1)]	259
Reduce [F]	259

Optimal result

Integrand size = 14, antiderivative size = 130

$$\begin{aligned} \int \frac{(a + b\operatorname{arctanh}(cx))^2}{x^4} dx = & -\frac{b^2c^2}{3x} + \frac{1}{3}b^2c^3\operatorname{arctanh}(cx) - \frac{bc(a + b\operatorname{arctanh}(cx))}{3x^2} \\ & + \frac{1}{3}c^3(a + b\operatorname{arctanh}(cx))^2 - \frac{(a + b\operatorname{arctanh}(cx))^2}{3x^3} \\ & + \frac{2}{3}bc^3(a + b\operatorname{arctanh}(cx)) \log\left(2 - \frac{2}{1 + cx}\right) \\ & - \frac{1}{3}b^2c^3 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + cx}\right) \end{aligned}$$

output

```
-1/3*b^2*c^2/x+1/3*b^2*c^3*arctanh(c*x)-1/3*b*c*(a+b*arctanh(c*x))/x^2+1/3
*c^3*(a+b*arctanh(c*x))^2-1/3*(a+b*arctanh(c*x))^2/x^3+2/3*b*c^3*(a+b*arct
anh(c*x))*ln(2-2/(c*x+1))-1/3*b^2*c^3*polylog(2,-1+2/(c*x+1))
```

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.12

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^4} dx = \frac{a^2 + abcx + b^2c^2x^2 + b^2(1 - c^3x^3) \operatorname{arctanh}(cx)^2 + b \operatorname{arctanh}(cx) (2a + bcx - bc^3x^3 - 2bc^3x^3 \log(1 - e^{-2 \operatorname{arctanh}(cx)}))}{3x^3}$$

input

```
Integrate[(a + b*ArcTanh[c*x])^2/x^4,x]
```

output

```
-1/3*(a^2 + a*b*c*x + b^2*c^2*x^2 + b^2*(1 - c^3*x^3)*ArcTanh[c*x]^2 + b*ArcTanh[c*x]*(2*a + b*c*x - b*c^3*x^3 - 2*b*c^3*x^3*Log[1 - E^(-2*ArcTanh[c*x])])) - 2*a*b*c^3*x^3*Log[c*x] + a*b*c^3*x^3*Log[1 - c^2*x^2] + b^2*c^3*x^3*PolyLog[2, E^(-2*ArcTanh[c*x])])/x^3
```

Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.92, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {6452, 6544, 6452, 264, 219, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^4} dx \\ & \quad \downarrow \text{6452} \\ & \frac{2}{3}bc \int \frac{a + b \operatorname{arctanh}(cx)}{x^3(1 - c^2x^2)} dx - \frac{(a + b \operatorname{arctanh}(cx))^2}{3x^3} \\ & \quad \downarrow \text{6544} \\ & \frac{2}{3}bc \left(c^2 \int \frac{a + b \operatorname{arctanh}(cx)}{x(1 - c^2x^2)} dx + \int \frac{a + b \operatorname{arctanh}(cx)}{x^3} dx \right) - \frac{(a + b \operatorname{arctanh}(cx))^2}{3x^3} \\ & \quad \downarrow \text{6452} \end{aligned}$$

$$\begin{aligned}
& \frac{2}{3}bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{x(1-c^2x^2)} dx + \frac{1}{2}bc \int \frac{1}{x^2(1-c^2x^2)} dx - \frac{a + \operatorname{barctanh}(cx)}{2x^2} \right) - \\
& \qquad \qquad \qquad \frac{(a + \operatorname{barctanh}(cx))^2}{3x^3} \\
& \qquad \qquad \qquad \downarrow \text{264} \\
& \frac{2}{3}bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{x(1-c^2x^2)} dx + \frac{1}{2}bc \left(c^2 \int \frac{1}{1-c^2x^2} dx - \frac{1}{x} \right) - \frac{a + \operatorname{barctanh}(cx)}{2x^2} \right) - \\
& \qquad \qquad \qquad \frac{(a + \operatorname{barctanh}(cx))^2}{3x^3} \\
& \qquad \qquad \qquad \downarrow \text{219} \\
& \frac{2}{3}bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{x(1-c^2x^2)} dx - \frac{a + \operatorname{barctanh}(cx)}{2x^2} + \frac{1}{2}bc \left(\operatorname{carctanh}(cx) - \frac{1}{x} \right) \right) - \\
& \qquad \qquad \qquad \frac{(a + \operatorname{barctanh}(cx))^2}{3x^3} \\
& \qquad \qquad \qquad \downarrow \text{6550} \\
& \frac{2}{3}bc \left(c^2 \left(\int \frac{a + \operatorname{barctanh}(cx)}{x(cx+1)} dx + \frac{(a + \operatorname{barctanh}(cx))^2}{2b} \right) - \frac{a + \operatorname{barctanh}(cx)}{2x^2} + \frac{1}{2}bc \left(\operatorname{carctanh}(cx) - \frac{1}{x} \right) \right) - \\
& \qquad \qquad \qquad \frac{(a + \operatorname{barctanh}(cx))^2}{3x^3} \\
& \qquad \qquad \qquad \downarrow \text{6494} \\
& \frac{2}{3}bc \left(c^2 \left(-bc \int \frac{\log \left(2 - \frac{2}{cx+1} \right)}{1-c^2x^2} dx + \frac{(a + \operatorname{barctanh}(cx))^2}{2b} + \log \left(2 - \frac{2}{cx+1} \right) (a + \operatorname{barctanh}(cx)) \right) - \frac{a + \operatorname{barctanh}(cx)}{2x^2} \right) - \\
& \qquad \qquad \qquad \frac{(a + \operatorname{barctanh}(cx))^2}{3x^3} \\
& \qquad \qquad \qquad \downarrow \text{2897} \\
& \frac{2}{3}bc \left(c^2 \left(\frac{(a + \operatorname{barctanh}(cx))^2}{2b} + \log \left(2 - \frac{2}{cx+1} \right) (a + \operatorname{barctanh}(cx)) - \frac{1}{2}b \operatorname{PolyLog} \left(2, \frac{2}{cx+1} - 1 \right) \right) - \frac{a + \operatorname{barctanh}(cx)}{2x^2} \right) - \\
& \qquad \qquad \qquad \frac{(a + \operatorname{barctanh}(cx))^2}{3x^3}
\end{aligned}$$

input

Int[(a + b*ArcTanh[c*x])^2/x^4, x]

output

```
-1/3*(a + b*ArcTanh[c*x])^2/x^3 + (2*b*c*(-1/2*(a + b*ArcTanh[c*x])/x^2 +
(b*c*(-x^(-1) + c*ArcTanh[c*x]))/2 + c^2*((a + b*ArcTanh[c*x])^2/(2*b) + (
a + b*ArcTanh[c*x])*Log[2 - 2/(1 + c*x)] - (b*PolyLog[2, -1 + 2/(1 + c*x)]
)/2))/3
```

Defintions of rubi rules used

rule 219

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

rule 264

```
Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(
m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*((m + 2*p + 3)/(a*c
^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p
}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]
```

rule 2897

```
Int[Log[u_]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/
D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] &&
PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u,
x][[2]], Expon[Pq, x]]
```

rule 6452

```
Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

rule 6494

```
Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((x_)*((d_) + (e_)*(x_))), x
_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] -
Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]]
/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c
^2*d^2 - e^2, 0]
```

rule 6544

```
Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_))/((d_) + (
e_.)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x
], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x
^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

rule 6550

```
Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_)^2)),
x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/
d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 246 vs. 2(116) = 232.

Time = 0.55 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.90

method	result
parts	$-\frac{a^2}{3x^3} + b^2c^3 \left(-\frac{\operatorname{arctanh}(cx)^2}{3c^3x^3} - \frac{\operatorname{arctanh}(cx)}{3c^2x^2} + \frac{2\ln(cx)\operatorname{arctanh}(cx)}{3} - \frac{\operatorname{arctanh}(cx)\ln(cx-1)}{3} - \frac{\operatorname{arctanh}(cx)\ln(cx+1)}{3} \right)$
derivativedivides	$c^3 \left(-\frac{a^2}{3c^3x^3} + b^2 \left(-\frac{\operatorname{arctanh}(cx)^2}{3c^3x^3} - \frac{\operatorname{arctanh}(cx)}{3c^2x^2} + \frac{2\ln(cx)\operatorname{arctanh}(cx)}{3} - \frac{\operatorname{arctanh}(cx)\ln(cx-1)}{3} - \frac{\operatorname{arctanh}(cx)\ln(cx+1)}{3} \right) \right)$
default	$c^3 \left(-\frac{a^2}{3c^3x^3} + b^2 \left(-\frac{\operatorname{arctanh}(cx)^2}{3c^3x^3} - \frac{\operatorname{arctanh}(cx)}{3c^2x^2} + \frac{2\ln(cx)\operatorname{arctanh}(cx)}{3} - \frac{\operatorname{arctanh}(cx)\ln(cx-1)}{3} - \frac{\operatorname{arctanh}(cx)\ln(cx+1)}{3} \right) \right)$

input

```
int((a+b*arctanh(c*x))^2/x^4,x,method=_RETURNVERBOSE)
```

output

```
-1/3*a^2/x^3+b^2*c^3*(-1/3/c^3/x^3*arctanh(c*x)^2-1/3/c^2/x^2*arctanh(c*x)
+2/3*ln(c*x)*arctanh(c*x)-1/3*arctanh(c*x)*ln(c*x-1)-1/3*arctanh(c*x)*ln(c
*x+1)-1/3/c/x-1/6*ln(c*x-1)+1/6*ln(c*x+1)-1/12*ln(c*x-1)^2+1/3*dilog(1/2*c
*x+1/2)+1/6*ln(c*x-1)*ln(1/2*c*x+1/2)+1/12*ln(c*x+1)^2-1/6*(ln(c*x+1)-ln(1
/2*c*x+1/2))*ln(-1/2*c*x+1/2)-1/3*dilog(c*x+1)-1/3*ln(c*x)*ln(c*x+1)-1/3*d
ilog(c*x))+2*a*b*c^3*(-1/3/c^3/x^3*arctanh(c*x)-1/6/c^2/x^2+1/3*ln(c*x)-1/
6*ln(c*x-1)-1/6*ln(c*x+1))
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^4} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^2}{x^4} dx$$

input `integrate((a+b*arctanh(c*x))^2/x^4,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x)^2 + 2*a*b*arctanh(c*x) + a^2)/x^4, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^4} dx = \int \frac{(a + b \operatorname{atanh}(cx))^2}{x^4} dx$$

input `integrate((a+b*atanh(c*x))**2/x**4,x)`

output `Integral((a + b*atanh(c*x))**2/x**4, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^4} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^2}{x^4} dx$$

input `integrate((a+b*arctanh(c*x))^2/x^4,x, algorithm="maxima")`

output `-1/3*((c^2*log(c^2*x^2 - 1) - c^2*log(x^2) + 1/x^2)*c + 2*arctanh(c*x)/x^3)*a*b - 1/12*b^2*(log(-c*x + 1)^2/x^3 + 3*integrate(-1/3*(3*(c*x - 1)*log(c*x + 1)^2 + 2*(c*x - 3*(c*x - 1)*log(c*x + 1))*log(-c*x + 1))/(c*x^5 - x^4), x)) - 1/3*a^2/x^3`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^4} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^2}{x^4} dx$$

input `integrate((a+b*arctanh(c*x))^2/x^4,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^2/x^4, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^4} dx = \int \frac{(a + b \operatorname{atanh}(cx))^2}{x^4} dx$$

input `int((a + b*atanh(c*x))^2/x^4,x)`

output `int((a + b*atanh(c*x))^2/x^4, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^4} dx$$

$$= \frac{-\operatorname{atanh}(cx)^2 b^2 - 2 \operatorname{atanh}(cx) ab c^3 x^3 - 2 \operatorname{atanh}(cx) ab + \operatorname{atanh}(cx) b^2 c^3 x^3 - \operatorname{atanh}(cx) b^2 cx - 2 \left(\int \frac{\operatorname{atanh}(cx)}{c^2 x^3} dx \right)}{3x^3}$$

input `int((a+b*atanh(c*x))^2/x^4,x)`

output `(- atanh(c*x)**2*b**2 - 2*atanh(c*x)*a*b*c**3*x**3 - 2*atanh(c*x)*a*b + atanh(c*x)*b**2*c**3*x**3 - atanh(c*x)*b**2*c*x - 2*int(atanh(c*x)/(c**2*x**3 - x),x)*b**2*c**3*x**3 - 2*log(c**2*x - c)*a*b*c**3*x**3 + 2*log(x)*a*b*c**3*x**3 - a**2 - a*b*c*x - b**2*c**2*x**2)/(3*x**3)`

3.23 $\int \frac{(a+b\operatorname{arctanh}(cx))^2}{x^5} dx$

Optimal result	260
Mathematica [A] (verified)	260
Rubi [A] (verified)	261
Maple [A] (verified)	264
Fricas [A] (verification not implemented)	265
Sympy [A] (verification not implemented)	265
Maxima [B] (verification not implemented)	266
Giac [B] (verification not implemented)	267
Mupad [B] (verification not implemented)	268
Reduce [B] (verification not implemented)	268

Optimal result

Integrand size = 14, antiderivative size = 117

$$\int \frac{(a + b\operatorname{arctanh}(cx))^2}{x^5} dx = -\frac{b^2c^2}{12x^2} - \frac{bc(a + b\operatorname{arctanh}(cx))}{6x^3} - \frac{bc^3(a + b\operatorname{arctanh}(cx))}{2x} + \frac{1}{4}c^4(a + b\operatorname{arctanh}(cx))^2 - \frac{(a + b\operatorname{arctanh}(cx))^2}{4x^4} + \frac{2}{3}b^2c^4 \log(x) - \frac{1}{3}b^2c^4 \log(1 - c^2x^2)$$

output

```
-1/12*b^2*c^2/x^2-1/6*b*c*(a+b*arctanh(c*x))/x^3-1/2*b*c^3*(a+b*arctanh(c*x))/x+1/4*c^4*(a+b*arctanh(c*x))^2-1/4*(a+b*arctanh(c*x))^2/x^4+2/3*b^2*c^4*ln(x)-1/3*b^2*c^4*ln(-c^2*x^2+1)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.40

$$\int \frac{(a + b\operatorname{arctanh}(cx))^2}{x^5} dx = \frac{3a^2 + 2abcx + b^2c^2x^2 + 6abc^3x^3 + 2b(3a + bcx + 3bc^3x^3) \operatorname{arctanh}(cx) - 3b^2(-1 + c^4x^4) \operatorname{arctanh}(cx)^2}{x^4}$$

input `Integrate[(a + b*ArcTanh[c*x])^2/x^5,x]`

output
$$-1/12*(3*a^2 + 2*a*b*c*x + b^2*c^2*x^2 + 6*a*b*c^3*x^3 + 2*b*(3*a + b*c*x + 3*b*c^3*x^3)*ArcTanh[c*x] - 3*b^2*(-1 + c^4*x^4)*ArcTanh[c*x]^2 - 8*b^2*c^4*x^4*Log[x] + 3*a*b*c^4*x^4*Log[1 - c*x] + 4*b^2*c^4*x^4*Log[1 - c*x] - 3*a*b*c^4*x^4*Log[1 + c*x] + 4*b^2*c^4*x^4*Log[1 + c*x])/x^4$$

Rubi [A] (verified)

Time = 0.93 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.16, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.929$, Rules used = {6452, 6544, 6452, 243, 54, 2009, 6544, 6452, 243, 47, 14, 16, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^5} dx \\ & \quad \downarrow 6452 \\ & \frac{1}{2}bc \int \frac{a + b \operatorname{arctanh}(cx)}{x^4(1 - c^2x^2)} dx - \frac{(a + b \operatorname{arctanh}(cx))^2}{4x^4} \\ & \quad \downarrow 6544 \\ & \frac{1}{2}bc \left(c^2 \int \frac{a + b \operatorname{arctanh}(cx)}{x^2(1 - c^2x^2)} dx + \int \frac{a + b \operatorname{arctanh}(cx)}{x^4} dx \right) - \frac{(a + b \operatorname{arctanh}(cx))^2}{4x^4} \\ & \quad \downarrow 6452 \\ & \frac{1}{2}bc \left(c^2 \int \frac{a + b \operatorname{arctanh}(cx)}{x^2(1 - c^2x^2)} dx + \frac{1}{3}bc \int \frac{1}{x^3(1 - c^2x^2)} dx - \frac{a + b \operatorname{arctanh}(cx)}{3x^3} \right) - \\ & \quad \frac{(a + b \operatorname{arctanh}(cx))^2}{4x^4} \\ & \quad \downarrow 243 \\ & \frac{1}{2}bc \left(c^2 \int \frac{a + b \operatorname{arctanh}(cx)}{x^2(1 - c^2x^2)} dx + \frac{1}{6}bc \int \frac{1}{x^4(1 - c^2x^2)} dx^2 - \frac{a + b \operatorname{arctanh}(cx)}{3x^3} \right) - \\ & \quad \frac{(a + b \operatorname{arctanh}(cx))^2}{4x^4} \\ & \quad \downarrow 54 \end{aligned}$$

$$\frac{1}{2}bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{x^2(1-c^2x^2)} dx + \frac{1}{6}bc \int \left(-\frac{c^4}{c^2x^2-1} + \frac{c^2}{x^2} + \frac{1}{x^4} \right) dx^2 - \frac{a + \operatorname{barctanh}(cx)}{3x^3} \right) - \frac{(a + \operatorname{barctanh}(cx))^2}{4x^4}$$

↓ 2009

$$\frac{1}{2}bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{x^2(1-c^2x^2)} dx - \frac{a + \operatorname{barctanh}(cx)}{3x^3} + \frac{1}{6}bc \left(c^2 \log(x^2) - c^2 \log(1-c^2x^2) - \frac{1}{x^2} \right) \right) - \frac{(a + \operatorname{barctanh}(cx))^2}{4x^4}$$

↓ 6544

$$\frac{1}{2}bc \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{1-c^2x^2} dx + \int \frac{a + \operatorname{barctanh}(cx)}{x^2} dx \right) - \frac{a + \operatorname{barctanh}(cx)}{3x^3} + \frac{1}{6}bc \left(c^2 \log(x^2) - c^2 \log(1-c^2x^2) - \frac{1}{x^2} \right) \right) - \frac{(a + \operatorname{barctanh}(cx))^2}{4x^4}$$

↓ 6452

$$\frac{1}{2}bc \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{1-c^2x^2} dx + bc \int \frac{1}{x(1-c^2x^2)} dx - \frac{a + \operatorname{barctanh}(cx)}{x} \right) - \frac{a + \operatorname{barctanh}(cx)}{3x^3} + \frac{1}{6}bc \left(c^2 \log(x^2) - c^2 \log(1-c^2x^2) - \frac{1}{x^2} \right) \right) - \frac{(a + \operatorname{barctanh}(cx))^2}{4x^4}$$

↓ 243

$$\frac{1}{2}bc \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{1-c^2x^2} dx + \frac{1}{2}bc \int \frac{1}{x^2(1-c^2x^2)} dx^2 - \frac{a + \operatorname{barctanh}(cx)}{x} \right) - \frac{a + \operatorname{barctanh}(cx)}{3x^3} + \frac{1}{6}bc \left(c^2 \log(x^2) - c^2 \log(1-c^2x^2) - \frac{1}{x^2} \right) \right) - \frac{(a + \operatorname{barctanh}(cx))^2}{4x^4}$$

↓ 47

$$\frac{1}{2}bc \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{1-c^2x^2} dx + \frac{1}{2}bc \left(c^2 \int \frac{1}{1-c^2x^2} dx^2 + \int \frac{1}{x^2} dx^2 \right) - \frac{a + \operatorname{barctanh}(cx)}{x} \right) - \frac{a + \operatorname{barctanh}(cx)}{3x^3} + \frac{1}{6}bc \left(c^2 \log(x^2) - c^2 \log(1-c^2x^2) - \frac{1}{x^2} \right) \right) - \frac{(a + \operatorname{barctanh}(cx))^2}{4x^4}$$

↓ 14

$$\frac{1}{2}bc \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{1-c^2x^2} dx + \frac{1}{2}bc \left(c^2 \int \frac{1}{1-c^2x^2} dx^2 + \log(x^2) \right) - \frac{a + \operatorname{barctanh}(cx)}{x} \right) - \frac{a + \operatorname{barctanh}(cx)}{3x^3} + \frac{1}{6}bc \left(c^2 \log(x^2) - c^2 \log(1-c^2x^2) - \frac{1}{x^2} \right) \right) - \frac{(a + \operatorname{barctanh}(cx))^2}{4x^4}$$

↓ 16

$$\frac{1}{2}bc \left(c^2 \int \frac{a + \operatorname{arctanh}(cx)}{1 - c^2x^2} dx - \frac{a + \operatorname{arctanh}(cx)}{x} + \frac{1}{2}bc(\log(x^2) - \log(1 - c^2x^2)) \right) - \frac{a + \operatorname{arctanh}(cx)}{3x^3} - \frac{(a + \operatorname{arctanh}(cx))^2}{4x^4}$$

↓ 6510

$$\frac{1}{2}bc \left(c^2 \left(\frac{c(a + \operatorname{arctanh}(cx))^2}{2b} - \frac{a + \operatorname{arctanh}(cx)}{x} + \frac{1}{2}bc(\log(x^2) - \log(1 - c^2x^2)) \right) - \frac{a + \operatorname{arctanh}(cx)}{3x^3} + \frac{1}{6} \right) - \frac{(a + \operatorname{arctanh}(cx))^2}{4x^4}$$

input `Int[(a + b*ArcTanh[c*x])^2/x^5,x]`

output `-1/4*(a + b*ArcTanh[c*x])^2/x^4 + (b*c*(-1/3*(a + b*ArcTanh[c*x])/x^3 + c^2*(-((a + b*ArcTanh[c*x])/x) + (c*(a + b*ArcTanh[c*x])^2)/(2*b) + (b*c*(Log[x^2] - Log[1 - c^2*x^2]))/2) + (b*c*(-x^(-2) + c^2*Log[x^2] - c^2*Log[1 - c^2*x^2]))/6)/2`

Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && !LtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 243 $\text{Int}[(x_)^{(m_)}*((a_) + (b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{((m - 1)/2)*(a + b*x)^p}, x], x, x^2], x] /; \text{FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m - 1)/2]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 6452 $\text{Int}[((a_) + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m + 1)), x] - \text{Simp}[b*c*n*(p/(m + 1)) \text{ Int}[x^{(m + n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p - 1)/(1 - c^2*x^{(2*n)})}), x], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 6510 $\text{Int}[((a_) + \text{ArcTanh}[(c_)*(x_)]*(b_))^{(p_)} / ((d_) + (e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^{(p + 1)} / (b*c*d*(p + 1)), x] /; \text{FreeQ}[\{a, b, c, d, e, p\}, x] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{NeQ}[p, -1]$

rule 6544 $\text{Int}[(((a_) + \text{ArcTanh}[(c_)*(x_)]*(b_))^{(p_)}*((f_)*(x_))^{(m_)} / ((d_) + (e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[1/d \text{ Int}[(f*x)^m*(a + b*\text{ArcTanh}[c*x])^p, x], x] - \text{Simp}[e/(d*f^2) \text{ Int}[(f*x)^{(m + 2)}*((a + b*\text{ArcTanh}[c*x])^p / (d + e*x^2)), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{LtQ}[m, -1]$

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.46

method	result
parallelrisch	$\frac{3b^2 \operatorname{arctanh}(cx)^2 x^4 c^4 + 8b^2 c^4 \ln(x) x^4 - 8 \ln(cx-1) x^4 b^2 c^4 + 6ab \operatorname{arctanh}(cx) x^4 c^4 - 8 \operatorname{arctanh}(cx) x^4 b^2 c^4 - b^2 c^4 x^4 - 6b^2 a}{12x^4}$
parts	$-\frac{a^2}{4x^4} + b^2 c^4 \left(-\frac{\operatorname{arctanh}(cx)^2}{4c^4 x^4} + \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{4} - \frac{\operatorname{arctanh}(cx)}{6c^3 x^3} - \frac{\operatorname{arctanh}(cx)}{2cx} - \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{4} \right)$
derivativedivides	$c^4 \left(-\frac{a^2}{4c^4 x^4} + b^2 \left(-\frac{\operatorname{arctanh}(cx)^2}{4c^4 x^4} + \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{4} - \frac{\operatorname{arctanh}(cx)}{6c^3 x^3} - \frac{\operatorname{arctanh}(cx)}{2cx} - \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{4} \right) \right)$
default	$c^4 \left(-\frac{a^2}{4c^4 x^4} + b^2 \left(-\frac{\operatorname{arctanh}(cx)^2}{4c^4 x^4} + \frac{\operatorname{arctanh}(cx) \ln(cx+1)}{4} - \frac{\operatorname{arctanh}(cx)}{6c^3 x^3} - \frac{\operatorname{arctanh}(cx)}{2cx} - \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{4} \right) \right)$
risch	$\frac{b^2(c^4 x^4 - 1) \ln(cx+1)^2}{16x^4} - \frac{b(3b x^4 \ln(-cx+1) c^4 + 6b c^3 x^3 + 2bcx - 3b \ln(-cx+1) + 6a) \ln(cx+1)}{24x^4} - \frac{-3b^2 c^4 x^4 \ln(-cx+1)}{24x^4}$

input `int((a+b*arctanh(c*x))^2/x^5,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{12}(3b^2\operatorname{arctanh}(cx)^2x^4c^4+8b^2c^4\ln(x)x^4-8\ln(cx-1)x^4b^2c^4+6ab\operatorname{arctanh}(cx)x^4c^4-8\operatorname{arctanh}(cx)x^4b^2c^4-b^2c^4x^4-6b^2\operatorname{arctanh}(cx)x^3c^3-6ab^2c^3x^3-b^2c^2x^2-2b^2\operatorname{arctanh}(cx)x^2c^2+ab^2cx-3b^2\operatorname{arctanh}(cx)^2-6\operatorname{arctanh}(cx)ab-3a^2)/x^4$$

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.48

$$\int \frac{(a + b\operatorname{arctanh}(cx))^2}{x^5} dx$$

$$= \frac{32b^2c^4x^4\log(x) + 4(3ab - 4b^2)c^4x^4\log(cx + 1) - 4(3ab + 4b^2)c^4x^4\log(cx - 1) - 24abc^3x^3 - 4b^2c^2x^2}{48x^4}$$

input `integrate((a+b*arctanh(c*x))^2/x^5,x, algorithm="fricas")`

output
$$\frac{1}{48}(32b^2c^4x^4\log(x) + 4(3ab - 4b^2)c^4x^4\log(cx + 1) - 4(3ab + 4b^2)c^4x^4\log(cx - 1) - 24abc^3x^3 - 4b^2c^2x^2 - 8ab^2cx + 3(b^2c^4x^4 - b^2)\log(-(cx + 1)/(cx - 1))^2 - 12a^2 - 4(3b^2c^3x^3 + b^2cx + 3ab)\log(-(cx + 1)/(cx - 1)))/x^4$$

Sympy [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.57

$$\int \frac{(a + b\operatorname{atanh}(cx))^2}{x^5} dx$$

$$= \begin{cases} -\frac{a^2}{4x^4} + \frac{abc^4\operatorname{atanh}(cx)}{2} - \frac{abc^3}{2x} - \frac{abc}{6x^3} - \frac{ab\operatorname{atanh}(cx)}{2x^4} + \frac{2b^2c^4\log(x)}{3} - \frac{2b^2c^4\log(x-\frac{1}{c})}{3} + \frac{b^2c^4\operatorname{atanh}^2(cx)}{4} - \frac{2b^2c^4\operatorname{atanh}(cx)}{3} \\ -\frac{a^2}{4x^4} \end{cases}$$

input `integrate((a+b*atanh(c*x))**2/x**5,x)`

output

```
Piecewise((-a**2/(4*x**4) + a*b*c**4*atanh(c*x)/2 - a*b*c**3/(2*x) - a*b*c
/(6*x**3) - a*b*atanh(c*x)/(2*x**4) + 2*b**2*c**4*log(x)/3 - 2*b**2*c**4*log(x - 1/c)/3 + b**2*c**4*atanh(c*x)**2/4 - 2*b**2*c**4*atanh(c*x)/3 - b**2*c**3*atanh(c*x)/(2*x) - b**2*c**2/(12*x**2) - b**2*c*atanh(c*x)/(6*x**3) - b**2*atanh(c*x)**2/(4*x**4), Ne(c, 0)), (-a**2/(4*x**4), True))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 224 vs. $2(103) = 206$.

Time = 0.04 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.91

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^5} dx$$

$$= \frac{1}{12} \left(\left(3c^3 \log(cx + 1) - 3c^3 \log(cx - 1) - \frac{2(3c^2x^2 + 1)}{x^3} \right) c - \frac{6 \operatorname{arctanh}(cx)}{x^4} \right) ab$$

$$+ \frac{1}{48} \left(\left(32c^2 \log(x) - \frac{3c^2x^2 \log(cx + 1)^2 + 3c^2x^2 \log(cx - 1)^2 + 16c^2x^2 \log(cx - 1) - 2(3c^2x^2 \log(cx - 1) - 8c^2x^2) \log(cx + 1) + 4}{x^2} \right) c^2 + 4(3c^3 \log(cx + 1) - 3c^3 \log(cx - 1) - 2(3c^2x^2 + 1)/x^3) c \operatorname{arctanh}(cx) \right) b^2 - \frac{1}{4} b^2 \operatorname{arctanh}(cx)^2/x^4 - \frac{1}{4} a^2/x^4$$

input

```
integrate((a+b*arctanh(c*x))^2/x^5,x, algorithm="maxima")
```

output

```
1/12*((3*c^3*log(c*x + 1) - 3*c^3*log(c*x - 1) - 2*(3*c^2*x^2 + 1)/x^3)*c
- 6*arctanh(c*x)/x^4)*a*b + 1/48*((32*c^2*log(x) - (3*c^2*x^2*log(c*x + 1)
^2 + 3*c^2*x^2*log(c*x - 1)^2 + 16*c^2*x^2*log(c*x - 1) - 2*(3*c^2*x^2*log
(c*x - 1) - 8*c^2*x^2)*log(c*x + 1) + 4)/x^2)*c^2 + 4*(3*c^3*log(c*x + 1)
- 3*c^3*log(c*x - 1) - 2*(3*c^2*x^2 + 1)/x^3)*c*arctanh(c*x))*b^2 - 1/4*b^
2*arctanh(c*x)^2/x^4 - 1/4*a^2/x^4
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 612 vs. $2(103) = 206$.

Time = 0.13 (sec) , antiderivative size = 612, normalized size of antiderivative = 5.23

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^5} dx = \text{Too large to display}$$

input `integrate((a+b*arctanh(c*x))^2/x^5,x, algorithm="giac")`

output

```
1/6*(4*b^2*c^3*log(-(c*x + 1)/(c*x - 1) - 1) - 4*b^2*c^3*log(-(c*x + 1)/(c*x - 1)) + 3*((c*x + 1)^3*b^2*c^3/(c*x - 1)^3 + (c*x + 1)*b^2*c^3/(c*x - 1)))*log(-(c*x + 1)/(c*x - 1))^2/((c*x + 1)^4/(c*x - 1)^4 + 4*(c*x + 1)^3/(c*x - 1)^3 + 6*(c*x + 1)^2/(c*x - 1)^2 + 4*(c*x + 1)/(c*x - 1) + 1) + 2*(6*(c*x + 1)^3*a*b*c^3/(c*x - 1)^3 + 6*(c*x + 1)*a*b*c^3/(c*x - 1) + 3*(c*x + 1)^3*b^2*c^3/(c*x - 1)^3 + 6*(c*x + 1)^2*b^2*c^3/(c*x - 1)^2 + 5*(c*x + 1)*b^2*c^3/(c*x - 1) + 2*b^2*c^3)*log(-(c*x + 1)/(c*x - 1))/((c*x + 1)^4/(c*x - 1)^4 + 4*(c*x + 1)^3/(c*x - 1)^3 + 6*(c*x + 1)^2/(c*x - 1)^2 + 4*(c*x + 1)/(c*x - 1) + 1) + 2*(6*(c*x + 1)^3*a^2*c^3/(c*x - 1)^3 + 6*(c*x + 1)*a^2*c^3/(c*x - 1) + 6*(c*x + 1)^3*a*b*c^3/(c*x - 1)^3 + 12*(c*x + 1)^2*a*b*c^3/(c*x - 1)^2 + 10*(c*x + 1)*a*b*c^3/(c*x - 1) + 4*a*b*c^3 + (c*x + 1)^3*b^2*c^3/(c*x - 1)^3 + 2*(c*x + 1)^2*b^2*c^3/(c*x - 1)^2 + (c*x + 1)*b^2*c^3/(c*x - 1))/((c*x + 1)^4/(c*x - 1)^4 + 4*(c*x + 1)^3/(c*x - 1)^3 + 6*(c*x + 1)^2/(c*x - 1)^2 + 4*(c*x + 1)/(c*x - 1) + 1))*c
```


input `int((a+b*atanh(c*x))^2/x^5,x)`

output `(3*atanh(c*x)**2*b**2*c**4*x**4 - 3*atanh(c*x)**2*b**2 + 6*atanh(c*x)*a*b*c**4*x**4 - 6*atanh(c*x)*a*b - 8*atanh(c*x)*b**2*c**4*x**4 - 6*atanh(c*x)*b**2*c**3*x**3 - 2*atanh(c*x)*b**2*c*x - 8*log(c**2*x - c)*b**2*c**4*x**4 + 8*log(x)*b**2*c**4*x**4 - 3*a**2 - 6*a*b*c**3*x**3 - 2*a*b*c*x - b**2*c**2*x**2)/(12*x**4)`

3.24 $\int x^5(a + \operatorname{arctanh}(cx))^3 dx$

Optimal result	270
Mathematica [A] (verified)	271
Rubi [B] (verified)	271
Maple [B] (verified)	278
Fricas [F]	279
Sympy [F]	280
Maxima [F]	280
Giac [F]	281
Mupad [F(-1)]	281
Reduce [F]	281

Optimal result

Integrand size = 14, antiderivative size = 247

$$\int x^5(a + \operatorname{arctanh}(cx))^3 dx = \frac{19b^3x}{60c^5} + \frac{b^3x^3}{60c^3} - \frac{19b^3\operatorname{arctanh}(cx)}{60c^6} + \frac{4b^2x^2(a + \operatorname{arctanh}(cx))}{15c^4} + \frac{b^2x^4(a + \operatorname{arctanh}(cx))}{20c^2} + \frac{23b(a + \operatorname{arctanh}(cx))^2}{30c^6} + \frac{bx(a + \operatorname{arctanh}(cx))^2}{2c^5} + \frac{bx^3(a + \operatorname{arctanh}(cx))^2}{6c^3} + \frac{bx^5(a + \operatorname{arctanh}(cx))^2}{10c} - \frac{(a + \operatorname{arctanh}(cx))^3}{6c^6} + \frac{1}{6}x^6(a + \operatorname{arctanh}(cx))^3 - \frac{23b^2(a + \operatorname{arctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{15c^6} - \frac{23b^3 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{30c^6}$$

output

```
19/60*b^3*x/c^5+1/60*b^3*x^3/c^3-19/60*b^3*arctanh(c*x)/c^6+4/15*b^2*x^2*(a+b*arctanh(c*x))/c^4+1/20*b^2*x^4*(a+b*arctanh(c*x))/c^2+23/30*b*(a+b*arctanh(c*x))^2/c^6+1/2*b*x*(a+b*arctanh(c*x))^2/c^5+1/6*b*x^3*(a+b*arctanh(c*x))^2/c^3+1/10*b*x^5*(a+b*arctanh(c*x))^2/c-1/6*(a+b*arctanh(c*x))^3/c^6+1/6*x^6*(a+b*arctanh(c*x))^3-23/15*b^2*(a+b*arctanh(c*x))*ln(2/(-c*x+1))/c^6-23/30*b^3*polylog(2,1-2/(-c*x+1))/c^6
```

Mathematica [A] (verified)

Time = 0.48 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.23

$$\int x^5(a + b \operatorname{arctanh}(cx))^3 dx$$

$$= \frac{-19ab^2 + 30a^2bcx + 19b^3cx + 16ab^2c^2x^2 + 10a^2bc^3x^3 + b^3c^3x^3 + 3ab^2c^4x^4 + 6a^2bc^5x^5 + 10a^3c^6x^6 + 2b^2c^6x^6}{60c^6}$$

input

```
Integrate[x^5*(a + b*ArcTanh[c*x])^3,x]
```

output

```
(-19*a*b^2 + 30*a^2*b*c*x + 19*b^3*c*x + 16*a*b^2*c^2*x^2 + 10*a^2*b*c^3*x^3 + b^3*c^3*x^3 + 3*a*b^2*c^4*x^4 + 6*a^2*b*c^5*x^5 + 10*a^3*c^6*x^6 + 2*b^2*(b*(-23 + 15*c*x + 5*c^3*x^3 + 3*c^5*x^5) + 15*a*(-1 + c^6*x^6))*ArcTanh[c*x]^2 + 10*b^3*(-1 + c^6*x^6)*ArcTanh[c*x]^3 + b*ArcTanh[c*x]*(30*a^2*c^6*x^6 + 4*a*b*c*x*(15 + 5*c^2*x^2 + 3*c^4*x^4) + b^2*(-19 + 16*c^2*x^2 + 3*c^4*x^4) - 92*b^2*Log[1 + E^(-2*ArcTanh[c*x])]) + 15*a^2*b*Log[1 - c*x] - 15*a^2*b*Log[1 + c*x] + 46*a*b^2*Log[1 - c^2*x^2] + 46*b^3*PolyLog[2, -E^(-2*ArcTanh[c*x])])/(60*c^6)
```

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 499 vs. 2(247) = 494.

Time = 3.53 (sec) , antiderivative size = 499, normalized size of antiderivative = 2.02, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 1.214$, Rules used = {6452, 6542, 6452, 6542, 6452, 254, 2009, 6542, 6436, 6452, 262, 219, 6510, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^5(a + b \operatorname{arctanh}(cx))^3 dx$$

$$\downarrow 6452$$

$$\frac{1}{6}x^6(a + b \operatorname{arctanh}(cx))^3 - \frac{1}{2}bc \int \frac{x^6(a + b \operatorname{arctanh}(cx))^2}{1 - c^2x^2} dx$$

$$\begin{aligned}
 & \downarrow 6542 \\
 & \frac{1}{6}x^6(a + \operatorname{barctanh}(cx))^3 - \frac{1}{2}bc \left(\frac{\int \frac{x^4(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\int x^4(a + \operatorname{barctanh}(cx))^2 dx}{c^2} \right) \\
 & \downarrow 6452 \\
 & \frac{1}{6}x^6(a + \operatorname{barctanh}(cx))^3 - \\
 & \frac{1}{2}bc \left(\frac{\int \frac{x^4(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{5}x^5(a + \operatorname{barctanh}(cx))^2 - \frac{2}{5}bc \int \frac{x^5(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} \right) \\
 & \downarrow 6542 \\
 & \frac{1}{6}x^6(a + \operatorname{barctanh}(cx))^3 - \\
 & \frac{1}{2}bc \left(\frac{\frac{\int \frac{x^2(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\int x^2(a + \operatorname{barctanh}(cx))^2 dx}{c^2}}{c^2} - \frac{\frac{1}{5}x^5(a + \operatorname{barctanh}(cx))^2 - \frac{2}{5}bc \left(\frac{\int \frac{x^3(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} \right)}{c^2} \right) \\
 & \downarrow 6452 \\
 & \frac{1}{6}x^6(a + \operatorname{barctanh}(cx))^3 - \\
 & \frac{1}{2}bc \left(\frac{\frac{\int \frac{x^2(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^2 - \frac{2}{3}bc \int \frac{x^3(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2}}{c^2} - \frac{\frac{1}{5}x^5(a + \operatorname{barctanh}(cx))^2 - \frac{2}{5}bc \left(\right)}{c^2} \right) \\
 & \downarrow 254 \\
 & \frac{1}{6}x^6(a + \operatorname{barctanh}(cx))^3 - \\
 & \frac{1}{2}bc \left(\frac{\frac{\int \frac{x^2(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^2 - \frac{2}{3}bc \int \frac{x^3(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2}}{c^2} - \frac{\frac{1}{5}x^5(a + \operatorname{barctanh}(cx))^2 - \frac{2}{5}bc \left(\right)}{c^2} \right) \\
 & \downarrow 2009
 \end{aligned}$$

$$\frac{1}{6}x^6(a + b\operatorname{arctanh}(cx))^3 - \frac{1}{2}bc \left(\frac{\int \frac{x^2(a+b\operatorname{arctanh}(cx))^2 dx}{1-c^2x^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \int \frac{x^3(a+b\operatorname{arctanh}(cx)) dx}{1-c^2x^2}}{c^2}}{c^2} - \frac{\frac{1}{5}x^5(a + b\operatorname{arctanh}(cx))^2 - \frac{2}{5}bc}{c^2} \right)$$

↓ 6542

$$\frac{1}{6}x^6(a + b\operatorname{arctanh}(cx))^3 - \frac{1}{2}bc \left(\frac{\frac{\int \frac{(a+b\operatorname{arctanh}(cx))^2 dx}{1-c^2x^2}}{c^2} - \frac{\int (a+b\operatorname{arctanh}(cx))^2 dx}{c^2}}{c^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\int \frac{x(a+b\operatorname{arctanh}(cx)) dx}{1-c^2x^2}}{c^2} - \int \frac{x(a+b\operatorname{arctanh}(cx))}{c^2} \right)}{c^2}}{c^2} \right)$$

↓ 6436

$$\frac{1}{6}x^6(a + b\operatorname{arctanh}(cx))^3 - \frac{1}{2}bc \left(\frac{\frac{\int \frac{(a+b\operatorname{arctanh}(cx))^2 dx}{1-c^2x^2}}{c^2} - \frac{x(a+b\operatorname{arctanh}(cx))^2 - 2bc \int \frac{x(a+b\operatorname{arctanh}(cx)) dx}{1-c^2x^2}}{c^2}}{c^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\int \frac{x(a+b\operatorname{arctanh}(cx))}{1-c^2x^2}}{c^2} \right)}{c^2}}{c^2} \right)$$

↓ 6452

$$\frac{1}{6}x^6(a + b\operatorname{arctanh}(cx))^3 - \frac{1}{2}bc \left(\frac{\frac{\int \frac{(a+b\operatorname{arctanh}(cx))^2 dx}{1-c^2x^2}}{c^2} - \frac{x(a+b\operatorname{arctanh}(cx))^2 - 2bc \int \frac{x(a+b\operatorname{arctanh}(cx)) dx}{1-c^2x^2}}{c^2}}{c^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\int \frac{x(a+b\operatorname{arctanh}(cx))}{1-c^2x^2}}{c^2} \right)}{c^2}}{c^2} \right)$$

↓ 262

$$\frac{1}{6}x^6(a + b\operatorname{arctanh}(cx))^3 - \frac{1}{2}bc \left(\frac{\int \frac{(a+b\operatorname{arctanh}(cx))^2}{1-c^2x^2} dx}{c^2} - \frac{x(a+b\operatorname{arctanh}(cx))^2 - 2bc \int \frac{x(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \left(\int \frac{x(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx \right)}{c^2} \right)$$

↓ 219

$$\frac{1}{6}x^6(a + b\operatorname{arctanh}(cx))^3 - \frac{1}{2}bc \left(\frac{\int \frac{(a+b\operatorname{arctanh}(cx))^2}{1-c^2x^2} dx}{c^2} - \frac{x(a+b\operatorname{arctanh}(cx))^2 - 2bc \int \frac{x(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \left(\int \frac{x(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx \right)}{c^2} \right)$$

↓ 6510

$$\frac{1}{6}x^6(a + b\operatorname{arctanh}(cx))^3 - \frac{1}{2}bc \left(\frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^3} - \frac{x(a+b\operatorname{arctanh}(cx))^2 - 2bc \int \frac{x(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \left(\int \frac{x(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx \right)}{c^2} \right)$$

↓ 6546

$$\frac{1}{6}x^6(a + b\operatorname{arctanh}(cx))^3 -$$

$$\frac{1}{2}bc \left(\frac{\frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^3} - \frac{x(a+b\operatorname{arctanh}(cx))^2 - 2bc \left(\frac{\int \frac{a+b\operatorname{arctanh}(cx)}{1-cx} dx}{c} - \frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^2} \right)}{c^2}}{c^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\int \frac{a+b\operatorname{arctanh}(cx)}{1-cx} dx}{c} - \frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^2} \right)}{c^2}}{c^2} \right)$$

6470

$$\frac{1}{6}x^6(a + b\operatorname{arctanh}(cx))^3 -$$

$$\frac{1}{2}bc \left(\frac{\frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^3} - \frac{x(a+b\operatorname{arctanh}(cx))^2 - 2bc \left(\frac{\frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))}{c}}{c} - b \int \frac{\log\left(\frac{2}{1-cx}\right)}{1-c^2x^2} dx - \frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^2} \right)}{c^2}}{c^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))}{c} - b \int \frac{\log\left(\frac{2}{1-cx}\right)}{1-c^2x^2} dx - \frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^2} \right)}{c^2}}{c^2} \right)$$

2849

$$\frac{1}{6}x^6(a + b\operatorname{arctanh}(cx))^3 -$$

$$\frac{1}{2}bc \left(\frac{\frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^3} - \frac{x(a+b\operatorname{arctanh}(cx))^2 - 2bc \left(\frac{b \int \frac{\log\left(\frac{2}{1-cx}\right) d\frac{1}{1-cx}}{1-\frac{2}{1-cx}}}{c} + \frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))}{c} - \frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^2} \right)}{c^2}}{c^2} - \frac{\frac{1}{3}x^3(a+b\operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{b \int \frac{\log\left(\frac{2}{1-cx}\right) d\frac{1}{1-cx}}{1-\frac{2}{1-cx}}}{c} + \frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))}{c} - \frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^2} \right)}{c^2}}{c^2} \right)$$

2752

$$\frac{1}{6}x^6(a + b\operatorname{arctanh}(cx))^3 - \frac{1}{2}bc \left(\frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^3} - \frac{x^{(a+b\operatorname{arctanh}(cx))^2-2bc} \left(\frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))}{c} + \frac{b\operatorname{PolyLog}\left(2,1-\frac{2}{1-cx}\right)}{2c} - \frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^2} \right)}{c^2} \right) - \frac{1}{3}x^3$$

```
input Int [x^5*(a + b*ArcTanh[c*x])^3,x]
```

```
output (x^6*(a + b*ArcTanh[c*x])^3)/6 - (b*c*(-((x^5*(a + b*ArcTanh[c*x])^2)/5 - (2*b*c*(-((x^4*(a + b*ArcTanh[c*x]))/4 - (b*c*(-(x/c^4) - x^3/(3*c^2) + ArcTanh[c*x]/c^5))/4)/c^2) + (-((x^2*(a + b*ArcTanh[c*x]))/2 - (b*c*(-(x/c^2) + ArcTanh[c*x]/c^3))/2)/c^2) + (-1/2*(a + b*ArcTanh[c*x])^2/(b*c^2) + (((a + b*ArcTanh[c*x])*Log[2/(1 - c*x)])/c + (b*PolyLog[2, 1 - 2/(1 - c*x)])/((2*c))/c)/c^2)/5)/c^2) + (-((x^3*(a + b*ArcTanh[c*x])^2)/3 - (2*b*c*(-((x^2*(a + b*ArcTanh[c*x]))/2 - (b*c*(-(x/c^2) + ArcTanh[c*x]/c^3))/2)/c^2) + (-1/2*(a + b*ArcTanh[c*x])^2/(b*c^2) + ((a + b*ArcTanh[c*x])*Log[2/(1 - c*x)])/c + (b*PolyLog[2, 1 - 2/(1 - c*x)])/((2*c))/c)/c^2))/3)/c^2) + ((a + b*ArcTanh[c*x])^3/(3*b*c^3) - (x*(a + b*ArcTanh[c*x])^2 - 2*b*c*(-1/2*(a + b*ArcTanh[c*x])^2/(b*c^2) + (((a + b*ArcTanh[c*x])*Log[2/(1 - c*x)])/c + (b*PolyLog[2, 1 - 2/(1 - c*x)])/((2*c))/c))/c^2)/c^2)/2
```

Defintions of rubi rules used

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))* ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt Q[a, 0] || LtQ[b, 0])
```

```
rule 254 Int[(x_)^(m_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^2, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 3]
```

rule 262 $\text{Int}[(c \cdot x)^m \cdot (a + b \cdot x^2)^p, x_Symbol] \rightarrow \text{Simp}[c \cdot (c \cdot x)^{m-1} \cdot (a + b \cdot x^2)^{p+1} / (b \cdot (m + 2 \cdot p + 1)), x] - \text{Simp}[a \cdot c^2 \cdot (m-1) / (b \cdot (m + 2 \cdot p + 1)) \text{Int}[(c \cdot x)^{m-2} \cdot (a + b \cdot x^2)^p, x], x] /;$ $\text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{GtQ}[m, 2 - 1] \ \&\& \ \text{NeQ}[m + 2 \cdot p + 1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 2009 $\text{Int}[u, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /;$ $\text{SumQ}[u]$

rule 2752 $\text{Int}[\text{Log}[(c \cdot x) / ((d) + (e \cdot x))], x_Symbol] \rightarrow \text{Simp}[(-e^{-1}) \cdot \text{PolyLog}[2, 1 - c \cdot x], x] /;$ $\text{FreeQ}\{c, d, e\}, x \ \&\& \ \text{EqQ}[e + c \cdot d, 0]$

rule 2849 $\text{Int}[\text{Log}[(c \cdot x) / ((d) + (e \cdot x))] / ((f) + (g \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[-e/g \text{Subst}[\text{Int}[\text{Log}[2 \cdot d \cdot x] / (1 - 2 \cdot d \cdot x), x], x, 1 / (d + e \cdot x)], x] /;$ $\text{FreeQ}\{c, d, e, f, g\}, x \ \&\& \ \text{EqQ}[c, 2 \cdot d] \ \&\& \ \text{EqQ}[e^2 \cdot f + d^2 \cdot g, 0]$

rule 6436 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n]) \cdot (b \cdot x)^p, x_Symbol] \rightarrow \text{Simp}[x \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^p, x] - \text{Simp}[b \cdot c \cdot n \cdot p \text{Int}[x^n \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2 \cdot n})], x], x] /;$ $\text{FreeQ}\{a, b, c, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[n, 1] \ || \ \text{EqQ}[p, 1])$

rule 6452 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n]) \cdot (b \cdot x)^p \cdot (x)^m, x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^p / (m+1), x] - \text{Simp}[b \cdot c \cdot n \cdot (p / (m+1)) \text{Int}[x^{m+n} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2 \cdot n})], x], x] /;$ $\text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 6470 $\text{Int}[(a + \text{ArcTanh}[c \cdot x]) \cdot (b \cdot x)^p / ((d) + (e \cdot x)), x_Symbol] \rightarrow \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{Log}[2 / (1 + e \cdot (x/d))] / e), x] + \text{Simp}[b \cdot c \cdot (p/e) \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{Log}[2 / (1 + e \cdot (x/d))] / (1 - c^2 \cdot x^2)), x], x] /;$ $\text{FreeQ}\{a, b, c, d, e\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d^2 - e^2, 0]$

rule 6510 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6542 `Int((((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_)))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

rule 6546 `Int((((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*(x_))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 802 vs. $2(221) = 442$.

Time = 9.29 (sec) , antiderivative size = 803, normalized size of antiderivative = 3.25

method	result
risch	$-\frac{19b^3 \ln(-cx-1)}{120c^6} + \frac{a^3 x^6}{6} + \left(\frac{b^3(c^6 x^6 - 1) \ln(-cx+1)^2}{16c^6} - \frac{b^2 x(15a c^5 x^5 + 3b c^4 x^4 + 5b c^2 x^2 + 15b) \ln(-cx+1)}{60c^5} \right) -$
derivativedivides	Expression too large to display
default	Expression too large to display
parts	Expression too large to display

input `int(x^5*(a+b*arctanh(c*x))^3,x,method=_RETURNVERBOSE)`

output

```

-19/120/c^6*b^3*ln(-c*x-1)+1/6*a^3*x^6+(1/16*b^3*(c^6*x^6-1)/c^6*ln(-c*x+1)
)^2-1/60*b^2*x*(15*a*c^5*x^5+3*b*c^4*x^4+5*b*c^2*x^2+15*b)/c^5*ln(-c*x+1)-
1/120*b*(-30*a^2*c^6*x^6-12*a*b*c^5*x^5-3*b^2*c^4*x^4-20*a*b*c^3*x^3-16*b^
2*c^2*x^2-60*a*b*c*x-30*b*ln(-c*x+1)*a-46*b^2*ln(-c*x+1))/c^6)*ln(c*x+1)-1
/10/c*a*b^2*ln(-c*x+1)*x^5-1/6/c^6*a^3+19/60*b^3*x/c^5+1/60*b^3*x^3/c^3+1/
20/c^2*a*b^2*x^4+4/15/c^4*a*b^2*x^2+1/10/c*a^2*b*x^5+1/6/c^3*a^2*b*x^3+1/2
/c^5*a^2*b*x-1/8/c^6*a*b^2*ln(-c*x+1)^2-19/60/c^6*a*b^2-23/30/c^6*a^2*b-1/
3*b^3/c^6-1/48*b^3*ln(-c*x+1)^3*x^6+1/48/c^6*b^3*ln(-c*x+1)^3-23/120/c^6*b
^3*ln(-c*x+1)^2+19/120/c^6*b^3*ln(-c*x+1)-1/6/c^3*a*b^2*ln(-c*x+1)*x^3-1/2
/c^5*a*b^2*ln(-c*x+1)*x+23/30/c^6*a*b^2*ln(-c*x+1)+1/4/c^6*a^2*b*ln(-c*x+1
)+1/40/c*b^3*ln(-c*x+1)^2*x^5+1/24/c^3*b^3*ln(-c*x+1)^2*x^3+1/8/c^5*b^3*ln
(-c*x+1)^2*x-1/40/c^2*b^3*ln(-c*x+1)*x^4-2/15/c^4*b^3*ln(-c*x+1)*x^2+1/8*a
*b^2*ln(-c*x+1)^2*x^6-1/4*a^2*b*ln(-c*x+1)*x^6+23/30*b^3/c^6*dilog(-1/2*c*
x+1/2)+23/30*b^2/c^6*ln(-c*x-1)*a-1/4*b/c^6*ln(-c*x-1)*a^2-23/30*b^3/c^6*ln
(-c*x+1)*ln(1/2*c*x+1/2)+23/30*b^3/c^6*ln(-1/2*c*x+1/2)*ln(1/2*c*x+1/2)+1
/48*b^3*(c^6*x^6-1)/c^6*ln(c*x+1)^3+1/240*b^2*(-15*b*x^6*ln(-c*x+1)*c^6+30
*a*c^6*x^6+6*b*c^5*x^5+10*b*c^3*x^3+30*b*c*x+15*b*ln(-c*x+1)-30*a+46*b)/c^
6*ln(c*x+1)^2

```

Fricas [F]

$$\int x^5(a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{arctanh}(cx) + a)^3 x^5 dx$$

input

```
integrate(x^5*(a+b*arctanh(c*x))^3,x, algorithm="fricas")
```

output

```
integral(b^3*x^5*arctanh(c*x)^3 + 3*a*b^2*x^5*arctanh(c*x)^2 + 3*a^2*b*x^5
*arctanh(c*x) + a^3*x^5, x)
```

Sympy [F]

$$\int x^5(a + \operatorname{arctanh}(cx))^3 dx = \int x^5(a + b \operatorname{atanh}(cx))^3 dx$$

input `integrate(x**5*(a+b*atanh(c*x))**3,x)`

output `Integral(x**5*(a + b*atanh(c*x))**3, x)`

Maxima [F]

$$\int x^5(a + \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{artanh}(cx) + a)^3 x^5 dx$$

input `integrate(x^5*(a+b*arctanh(c*x))^3,x, algorithm="maxima")`

output `1/2*a*b^2*x^6*arctanh(c*x)^2 + 1/6*a^3*x^6 + 1/60*(30*x^6*arctanh(c*x) + c*(2*(3*c^4*x^5 + 5*c^2*x^3 + 15*x)/c^6 - 15*log(c*x + 1)/c^7 + 15*log(c*x - 1)/c^7))*a^2*b + 1/120*(4*c*(2*(3*c^4*x^5 + 5*c^2*x^3 + 15*x)/c^6 - 15*log(c*x + 1)/c^7 + 15*log(c*x - 1)/c^7)*arctanh(c*x) + (6*c^4*x^4 + 32*c^2*x^2 - 2*(15*log(c*x - 1) - 46)*log(c*x + 1) + 15*log(c*x + 1)^2 + 15*log(c*x - 1)^2 + 92*log(c*x - 1))/c^6)*a*b^2 - 1/1728000*(500*c^7*((2*c^4*x^6 + 3*c^2*x^4 + 6*x^2)/c^11 + 6*log(c^2*x^2 - 1)/c^13) + 728*c^6*(2*(3*c^4*x^5 + 5*c^2*x^3 + 15*x)/c^11 - 15*log(c*x + 1)/c^12 + 15*log(c*x - 1)/c^12) + 1485*c^5*((c^2*x^4 + 2*x^2)/c^9 + 2*log(c^2*x^2 - 1)/c^11) - 622080000*c^5*integrate(1/3600*x^5*log(c*x + 1)/(c^7*x^2 - c^5), x) + 9750*c^4*(2*(c^2*x^3 + 3*x)/c^9 - 3*log(c*x + 1)/c^10 + 3*log(c*x - 1)/c^10) - 2700*c^3*(x^2/c^7 + log(c^2*x^2 - 1)/c^9) - 1036800000*c^3*integrate(1/3600*x^3*log(c*x + 1)/(c^7*x^2 - c^5), x) + 227700*c^2*(2*x/c^7 - log(c*x + 1)/c^8 + log(c*x - 1)/c^8) - 5495040000*c*integrate(1/3600*x*log(c*x + 1)/(c^7*x^2 - c^5), x) + (1000*(36*log(-c*x + 1)^3 - 18*log(-c*x + 1)^2 + 6*log(-c*x + 1) - 1)*(c*x - 1)^6 + 1728*(125*log(-c*x + 1)^3 - 75*log(-c*x + 1)^2 + 30*log(-c*x + 1) - 6)*(c*x - 1)^5 + 16875*(32*log(-c*x + 1)^3 - 24*log(-c*x + 1)^2 + 12*log(-c*x + 1) - 3)*(c*x - 1)^4 + 80000*(9*log(-c*x + 1)^3 - 9*log(-c*x + 1)^2 + 6*log(-c*x + 1) - 2)*(c*x - 1)^3 + 135000*(4*log(-c*x + 1)^3 - 6*log(-c*x + 1)^2 + 6*log(-c*x + 1) - 3)*(c*x - 1)^2 + 216000*(log...`

Giac [F]

$$\int x^5(a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{artanh}(cx) + a)^3 x^5 dx$$

input `integrate(x^5*(a+b*arctanh(c*x))^3,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^3*x^5, x)`

Mupad [F(-1)]

Timed out.

$$\int x^5(a + b \operatorname{arctanh}(cx))^3 dx = \int x^5(a + b \operatorname{atanh}(cx))^3 dx$$

input `int(x^5*(a + b*atanh(c*x))^3,x)`

output `int(x^5*(a + b*atanh(c*x))^3, x)`

Reduce [F]

$$\int x^5(a + b \operatorname{arctanh}(cx))^3 dx$$

$$= \frac{-10 \operatorname{atanh}(cx)^3 b^3 - 19 \operatorname{atanh}(cx) b^3 + 30 \operatorname{atanh}(cx)^2 a b^2 c^6 x^6 + 30 \operatorname{atanh}(cx) a^2 b c^6 x^6 + 12 \operatorname{atanh}(cx) a b^2 c^6 x^6}{c^6}$$

input `int(x^5*(a+b*atanh(c*x))^3,x)`

output

```
(10*atanh(c*x)**3*b**3*c**6*x**6 - 10*atanh(c*x)**3*b**3 + 30*atanh(c*x)**2*a*b**2*c**6*x**6 - 30*atanh(c*x)**2*a*b**2 + 6*atanh(c*x)**2*b**3*c**5*x**5 + 10*atanh(c*x)**2*b**3*c**3*x**3 + 30*atanh(c*x)**2*b**3*c*x + 30*atanh(c*x)*a**2*b*c**6*x**6 - 30*atanh(c*x)*a**2*b + 12*atanh(c*x)*a*b**2*c**5*x**5 + 20*atanh(c*x)*a*b**2*c**3*x**3 + 60*atanh(c*x)*a*b**2*c*x + 92*atanh(c*x)*a*b**2 + 3*atanh(c*x)*b**3*c**4*x**4 + 16*atanh(c*x)*b**3*c**2*x**2 - 19*atanh(c*x)*b**3 + 92*int((atanh(c*x)*x)/(c**2*x**2 - 1),x)*b**3*c**2 + 92*log(c**2*x - c)*a*b**2 + 10*a**3*c**6*x**6 + 6*a**2*b*c**5*x**5 + 10*a**2*b*c**3*x**3 + 30*a**2*b*c*x + 3*a*b**2*c**4*x**4 + 16*a*b**2*c**2*x**2 + b**3*c**3*x**3 + 19*b**3*c*x)/(60*c**6)
```

3.25 $\int x^4(a + b \operatorname{arctanh}(cx))^3 dx$

Optimal result	283
Mathematica [A] (verified)	284
Rubi [A] (verified)	284
Maple [C] (warning: unable to verify)	290
Fricas [F]	291
Sympy [F]	291
Maxima [F]	291
Giac [F]	292
Mupad [F(-1)]	292
Reduce [F]	293

Optimal result

Integrand size = 14, antiderivative size = 262

$$\begin{aligned} \int x^4(a + b \operatorname{arctanh}(cx))^3 dx = & \frac{9ab^2x}{10c^4} + \frac{b^3x^2}{20c^3} + \frac{9b^3x \operatorname{arctanh}(cx)}{10c^4} \\ & + \frac{b^2x^3(a + b \operatorname{arctanh}(cx))}{10c^2} - \frac{9b(a + b \operatorname{arctanh}(cx))^2}{20c^5} \\ & + \frac{3bx^2(a + b \operatorname{arctanh}(cx))^2}{10c^3} + \frac{3bx^4(a + b \operatorname{arctanh}(cx))^2}{20c} \\ & + \frac{(a + b \operatorname{arctanh}(cx))^3}{5c^5} + \frac{1}{5}x^5(a + b \operatorname{arctanh}(cx))^3 \\ & - \frac{3b(a + b \operatorname{arctanh}(cx))^2 \log\left(\frac{2}{1-cx}\right)}{5c^5} + \frac{b^3 \log(1 - c^2x^2)}{2c^5} \\ & - \frac{3b^2(a + b \operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{5c^5} \\ & + \frac{3b^3 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx}\right)}{10c^5} \end{aligned}$$

output

```
9/10*a*b^2*x/c^4+1/20*b^3*x^2/c^3+9/10*b^3*x*arctanh(c*x)/c^4+1/10*b^2*x^3
*(a+b*arctanh(c*x))/c^2-9/20*b*(a+b*arctanh(c*x))^2/c^5+3/10*b*x^2*(a+b*ar
ctanh(c*x))^2/c^3+3/20*b*x^4*(a+b*arctanh(c*x))^2/c+1/5*(a+b*arctanh(c*x))
^3/c^5+1/5*x^5*(a+b*arctanh(c*x))^3-3/5*b*(a+b*arctanh(c*x))^2*ln(2/(-c*x+
1))/c^5+1/2*b^3*ln(-c^2*x^2+1)/c^5-3/5*b^2*(a+b*arctanh(c*x))*polylog(2,1-
2/(-c*x+1))/c^5+3/10*b^3*polylog(3,1-2/(-c*x+1))/c^5
```


Mathematica [A] (verified)

Time = 0.51 (sec) , antiderivative size = 383, normalized size of antiderivative = 1.46

$$\int x^4(a + b \operatorname{arctanh}(cx))^3 dx$$

$$= \frac{-b^3 + 18ab^2cx + 6a^2bc^2x^2 + b^3c^2x^2 + 2ab^2c^3x^3 + 3a^2bc^4x^4 + 4a^3c^5x^5 - 18ab^2 \operatorname{arctanh}(cx) + 18b^3cx \operatorname{arctanh}(cx) + 6a^2b^2c^2x^2 \operatorname{arctanh}(cx) + 12ab^2c^3x^3 \operatorname{arctanh}(cx) + 6a^2b^2c^4x^4 \operatorname{arctanh}(cx) - 12ab^2c^5x^5 \operatorname{arctanh}(cx) - 12ab^2c^2x^2 \operatorname{arctanh}(cx)^2 - 9b^3c^2x^2 \operatorname{arctanh}(cx)^2 + 6b^3c^2x^2 \operatorname{arctanh}(cx)^2 + 3b^3c^4x^4 \operatorname{arctanh}(cx)^2 + 12a^2b^2c^5x^5 \operatorname{arctanh}(cx)^2 - 4b^3c^3 \operatorname{arctanh}(cx)^3 + 4b^3c^5x^5 \operatorname{arctanh}(cx)^3 - 24a^2b^2 \operatorname{arctanh}(cx) \operatorname{Log}[1 + E^{-2 \operatorname{ArcTanh}[c*x]}] - 12b^3 \operatorname{arctanh}(cx)^2 \operatorname{Log}[1 + E^{-2 \operatorname{ArcTanh}[c*x]}] + 6a^2b \operatorname{Log}[1 - c^2x^2] + 10b^3 \operatorname{Log}[1 - c^2x^2] + 12b^2(a + b \operatorname{ArcTanh}[c*x]) \operatorname{PolyLog}[2, -E^{-2 \operatorname{ArcTanh}[c*x]}] + 6b^3 \operatorname{PolyLog}[3, -E^{-2 \operatorname{ArcTanh}[c*x]}]}{20c^5}$$

input

```
Integrate[x^4*(a + b*ArcTanh[c*x])^3,x]
```

output

```
(-b^3 + 18*a*b^2*c*x + 6*a^2*b*c^2*x^2 + b^3*c^2*x^2 + 2*a*b^2*c^3*x^3 + 3*a^2*b*c^4*x^4 + 4*a^3*c^5*x^5 - 18*a*b^2*ArcTanh[c*x] + 18*b^3*c*x*ArcTanh[c*x] + 12*a*b^2*c^2*x^2*ArcTanh[c*x] + 2*b^3*c^3*x^3*ArcTanh[c*x] + 6*a*b^2*c^4*x^4*ArcTanh[c*x] + 12*a^2*b*c^5*x^5*ArcTanh[c*x] - 12*a*b^2*ArcTanh[c*x]^2 - 9*b^3*ArcTanh[c*x]^2 + 6*b^3*c^2*x^2*ArcTanh[c*x]^2 + 3*b^3*c^4*x^4*ArcTanh[c*x]^2 + 12*a*b^2*c^5*x^5*ArcTanh[c*x]^2 - 4*b^3*ArcTanh[c*x]^3 + 4*b^3*c^5*x^5*ArcTanh[c*x]^3 - 24*a*b^2*ArcTanh[c*x]*Log[1 + E^(-2*ArcTanh[c*x])] - 12*b^3*ArcTanh[c*x]^2*Log[1 + E^(-2*ArcTanh[c*x])] + 6*a^2*b*Log[1 - c^2*x^2] + 10*b^3*Log[1 - c^2*x^2] + 12*b^2*(a + b*ArcTanh[c*x])*PolyLog[2, -E^(-2*ArcTanh[c*x])] + 6*b^3*PolyLog[3, -E^(-2*ArcTanh[c*x])])/(20*c^5)
```

Rubi [A] (verified)

Time = 2.90 (sec) , antiderivative size = 365, normalized size of antiderivative = 1.39, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 1.071$, Rules used = {6452, 6542, 6452, 6542, 6452, 243, 49, 2009, 6542, 2009, 6510, 6546, 6470, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4(a + b \operatorname{arctanh}(cx))^3 dx$$

↓ 6452

$$\begin{aligned}
& \frac{1}{5}x^5(a + \operatorname{barctanh}(cx))^3 - \frac{3}{5}bc \int \frac{x^5(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx \\
& \quad \downarrow 6542 \\
& \frac{1}{5}x^5(a + \operatorname{barctanh}(cx))^3 - \frac{3}{5}bc \left(\frac{\int \frac{x^3(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\int x^3(a + \operatorname{barctanh}(cx))^2 dx}{c^2} \right) \\
& \quad \downarrow 6452 \\
& \frac{1}{5}x^5(a + \operatorname{barctanh}(cx))^3 - \\
& \frac{3}{5}bc \left(\frac{\int \frac{x^3(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx))^2 - \frac{1}{2}bc \int \frac{x^4(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} \right) \\
& \quad \downarrow 6542 \\
& \frac{1}{5}x^5(a + \operatorname{barctanh}(cx))^3 - \\
& \frac{3}{5}bc \left(\frac{\frac{\int \frac{x(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\int x(a + \operatorname{barctanh}(cx))^2 dx}{c^2}}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx))^2 - \frac{1}{2}bc \left(\frac{\int \frac{x^2(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \int \frac{x^2}{1 - c^2x^2} dx \right)}{c^2}}{c^2} \right) \\
& \quad \downarrow 6452 \\
& \frac{1}{5}x^5(a + \operatorname{barctanh}(cx))^3 - \\
& \frac{3}{5}bc \left(\frac{\frac{\int \frac{x(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx))^2 - bc \int \frac{x^2(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2}}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx))^2 - \frac{1}{2}bc \left(\frac{\int \frac{x^2}{1 - c^2x^2} dx}{c^2} - \int \frac{x^2}{1 - c^2x^2} dx \right)}{c^2}}{c^2} \right) \\
& \quad \downarrow 243 \\
& \frac{1}{5}x^5(a + \operatorname{barctanh}(cx))^3 - \\
& \frac{3}{5}bc \left(\frac{\frac{\int \frac{x(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx))^2 - bc \int \frac{x^2(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2}}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx))^2 - \frac{1}{2}bc \left(\frac{\int \frac{x^2}{1 - c^2x^2} dx}{c^2} - \int \frac{x^2}{1 - c^2x^2} dx \right)}{c^2}}{c^2} \right) \\
& \quad \downarrow 49
\end{aligned}$$

$$\frac{1}{5}x^5(a + \operatorname{barctanh}(cx))^3 - \frac{3}{5}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx))^2 - bc \int \frac{x^2(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx))^2 - \frac{1}{2}bc \left(\int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx \right)}{c^2} \right)$$

2009

$$\frac{1}{5}x^5(a + \operatorname{barctanh}(cx))^3 - \frac{3}{5}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx))^2 - bc \int \frac{x^2(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx))^2 - \frac{1}{2}bc \left(\int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx \right)}{c^2} \right)$$

6542

$$\frac{1}{5}x^5(a + \operatorname{barctanh}(cx))^3 - \frac{3}{5}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx))^2 - bc \left(\frac{\int \frac{a + \operatorname{barctanh}(cx)}{1 - c^2x^2} dx}{c^2} - \frac{\int (a + \operatorname{barctanh}(cx)) dx}{c^2} \right)}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx))^2 - \frac{1}{2}bc \left(\int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx \right)}{c^2} \right)$$

2009

$$\frac{1}{5}x^5(a + \operatorname{barctanh}(cx))^3 - \frac{3}{5}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx))^2 - bc \left(\frac{\int \frac{a + \operatorname{barctanh}(cx)}{1 - c^2x^2} dx}{c^2} - \frac{ax + b \operatorname{arctanh}(cx) + \frac{b \log(1 - c^2x^2)}{2c}}{c^2} \right)}{c^2} - \frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx))^2 - \frac{1}{2}bc \left(\int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx \right)}{c^2} \right)$$

6510

$$\frac{1}{5}x^5(a + b\operatorname{arctanh}(cx))^3 - \frac{3}{5}bc \left(\frac{\int \frac{x(a+b\operatorname{arctanh}(cx))^2}{1-c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a+b\operatorname{arctanh}(cx))^2 - bc \left(\frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^3} - \frac{ax+b\operatorname{arctanh}(cx)}{c^2} + \frac{b \log(1-c^2x^2)}{2c} \right)}{c^2}}{c^2} - \frac{1}{4}x^4(a + b\operatorname{arctanh}(cx))^3 \right)$$

6546

$$\frac{1}{5}x^5(a + b\operatorname{arctanh}(cx))^3 - \frac{3}{5}bc \left(\frac{\int \frac{(a+b\operatorname{arctanh}(cx))^2}{1-cx} dx}{c} - \frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^2}}{c^2} - \frac{\frac{1}{2}x^2(a+b\operatorname{arctanh}(cx))^2 - bc \left(\frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^3} - \frac{ax+b\operatorname{arctanh}(cx)}{c^2} + \frac{b \log(1-c^2x^2)}{2c} \right)}{c^2}}{c^2} \right)$$

6470

$$\frac{1}{5}x^5(a + b\operatorname{arctanh}(cx))^3 - \frac{3}{5}bc \left(\frac{\frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))^2}{c} - 2b \int \frac{(a+b\operatorname{arctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{1-c^2x^2} dx}{c} - \frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^2}}{c^2} - \frac{\frac{1}{2}x^2(a+b\operatorname{arctanh}(cx))^2 - bc \left(\frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^3} - \frac{ax+b\operatorname{arctanh}(cx)}{c^2} + \frac{b \log(1-c^2x^2)}{2c} \right)}{c^2}}{c^2} \right)$$

6620

$$\frac{1}{5}x^5(a + b\operatorname{arctanh}(cx))^3 - \frac{3}{5}bc \left(\frac{\frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))^2}{c} - 2b \left(\frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{1-c^2x^2} dx - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))}{2c} \right)}{c} - \frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^2}}{c^2} - \frac{\frac{1}{2}x^2(a+b\operatorname{arctanh}(cx))^2 - bc \left(\frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^3} - \frac{ax+b\operatorname{arctanh}(cx)}{c^2} + \frac{b \log(1-c^2x^2)}{2c} \right)}{c^2}}{c^2} \right)$$

7164

$$\frac{1}{5}x^5(a + \operatorname{arctanh}(cx))^3 - \frac{3}{5}bc \left(\frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))^2}{c} - 2b \left(\frac{b \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx}\right)}{4c} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))}{2c} \right) - \frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^2} - \frac{\frac{1}{2}x^2(a+b\operatorname{arctanh}(cx))}{c^2} \right)$$

input `Int[x^4*(a + b*ArcTanh[c*x])^3,x]`

output $(x^5*(a + b*ArcTanh[c*x])^3)/5 - (3*b*c*(-((x^4*(a + b*ArcTanh[c*x])^2)/4 - (b*c*(-((x^3*(a + b*ArcTanh[c*x])))/3 - (b*c*(-(x^2/c^2) - \operatorname{Log}[1 - c^2*x^2]/c^4))/6)/c^2) + ((a + b*ArcTanh[c*x])^2/(2*b*c^3) - (a*x + b*x*ArcTanh[c*x] + (b*\operatorname{Log}[1 - c^2*x^2])/(2*c))/c^2)/2)/c^2) + (-((x^2*(a + b*ArcTanh[c*x])^2)/2 - b*c*((a + b*ArcTanh[c*x])^2/(2*b*c^3) - (a*x + b*x*ArcTanh[c*x] + (b*\operatorname{Log}[1 - c^2*x^2])/(2*c))/c^2))/c^2) + (-1/3*(a + b*ArcTanh[c*x])^3/(b*c^2) + (((a + b*ArcTanh[c*x])^2*\operatorname{Log}[2/(1 - c*x)])/c - 2*b*(-1/2*((a + b*ArcTanh[c*x])*PolyLog[2, 1 - 2/(1 - c*x)])/c + (b*PolyLog[3, 1 - 2/(1 - c*x)]/(4*c))/c)/c^2)/c^2)/5$

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 $\text{Int}[(a + \text{ArcTanh}[c \cdot x]^n] \cdot (b \cdot x)^m, x_Symbol] :> \text{Simp}[x^{m+1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x]^n)^p / (m+1), x] - \text{Simp}[b \cdot c \cdot n \cdot (p/(m+1)) \text{Int}[x^{m+n} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x]^n)^{p-1} / (1 - c^2 \cdot x^{2n})], x] /; \text{FreeQ}\{a, b, c, m, n\}, x\} \&\& \text{IGtQ}[p, 0] \&\& (\text{EqQ}[p, 1] \|\| (\text{EqQ}[n, 1] \&\& \text{IntegerQ}[m])) \&\& \text{NeQ}[m, -1]$

rule 6470 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x))^p / ((d + e \cdot x)), x_Symbol] :> \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{Log}[2/(1 + e \cdot (x/d))]) / e, x] + \text{Simp}[b \cdot c \cdot (p/e) \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{Log}[2/(1 + e \cdot (x/d))]) / (1 - c^2 \cdot x^2)], x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x\} \&\& \text{IGtQ}[p, 0] \&\& \text{EqQ}[c^2 \cdot d^2 - e^2, 0]$

rule 6510 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x))^p / ((d + e \cdot x)^2), x_Symbol] :> \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot c \cdot d \cdot (p+1)), x] /; \text{FreeQ}\{a, b, c, d, e, p\}, x\} \&\& \text{EqQ}[c^2 \cdot d + e, 0] \&\& \text{NeQ}[p, -1]$

rule 6542 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x))^p \cdot (f \cdot x)^m / ((d + e \cdot x)^2), x_Symbol] :> \text{Simp}[f^2/e \text{Int}[(f \cdot x)^{m-2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x] - \text{Simp}[d \cdot (f^2/e) \text{Int}[(f \cdot x)^{m-2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2)], x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x\} \&\& \text{GtQ}[p, 0] \&\& \text{GtQ}[m, 1]$

rule 6546 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x))^p \cdot (x) / ((d + e \cdot x)^2), x_Symbol] :> \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot e \cdot (p+1)), x] + \text{Simp}[1/(c \cdot d) \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p / (1 - c \cdot x)], x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x\} \&\& \text{EqQ}[c^2 \cdot d + e, 0] \&\& \text{IGtQ}[p, 0]$

rule 6620 $\text{Int}[(\text{Log}[u] \cdot (a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x))^p) / ((d + e \cdot x)^2), x_Symbol] :> \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{PolyLog}[2, 1 - u] / (2 \cdot c \cdot d)), x] + \text{Simp}[b \cdot (p/2) \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{PolyLog}[2, 1 - u]) / (d + e \cdot x^2)], x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x\} \&\& \text{IGtQ}[p, 0] \&\& \text{EqQ}[c^2 \cdot d + e, 0] \&\& \text{EqQ}[(1 - u)^2 - (1 - 2/(1 - c \cdot x))^2, 0]$

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 12.01 (sec) , antiderivative size = 1070, normalized size of antiderivative = 4.08

method	result	size
derivativedivides	Expression too large to display	1070
default	Expression too large to display	1070
parts	Expression too large to display	1072

input

```
int(x^4*(a+b*arctanh(c*x))^3,x,method=_RETURNVERBOSE)
```

output

```
1/c^5*(1/5*a^3*c^5*x^5+b^3*(3/20*c^4*x^4*arctanh(c*x)^2-1/10-9/20*arctanh(
c*x)^2+1/10*(c^2*x^2-4*c*x+7)*(c*x+1)*arctanh(c*x)+3/20*I*Pi*csgn(I/(1-(c*
x+1)^2/(c^2*x^2-1)))*csgn(I*(c*x+1)^2/(c^2*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x
^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))*arctanh(c*x)^2+1/10*c*x+1/20*(c*x-1)^2-ln
(1+(c*x+1)^2/(-c^2*x^2+1))+3/10*polylog(3,-(c*x+1)^2/(-c^2*x^2+1))-3/5*arc
tanh(c*x)*polylog(2,-(c*x+1)^2/(-c^2*x^2+1))+3/10*c^2*x^2*arctanh(c*x)^2-3
/20*I*Pi*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^3*arctanh
(c*x)^2+3/10*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))^2*arctanh(c*x)^2-3/10*
I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))^3*arctanh(c*x)^2-3/10*I*Pi*csgn(I*(
c*x+1)/(-c^2*x^2+1)^(1/2))*csgn(I*(c*x+1)^2/(c^2*x^2-1))^2*arctanh(c*x)^2-
3/20*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(
1-(c*x+1)^2/(c^2*x^2-1)))^2*arctanh(c*x)^2+3/20*I*Pi*csgn(I*(c*x+1)^2/(c^2
*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^2*arctanh
(c*x)^2-3/20*I*Pi*csgn(I*(c*x+1)/(-c^2*x^2+1)^(1/2))^2*csgn(I*(c*x+1)^2/(c
^2*x^2-1))*arctanh(c*x)^2+3/10*arctanh(c*x)^2*ln(c*x-1)+3/10*arctanh(c*x)^
2*ln(c*x+1)-3/5*arctanh(c*x)^2*ln((c*x+1)/(-c^2*x^2+1)^(1/2))-3/5*ln(2)*ar
ctanh(c*x)^2+6/5*(c*x+1)*arctanh(c*x)+1/5*c^5*x^5*arctanh(c*x)^3-3/10*I*Pi
*arctanh(c*x)^2+1/5*arctanh(c*x)^3-3/20*I*Pi*csgn(I*(c*x+1)^2/(c^2*x^2-1))
^3*arctanh(c*x)^2+3/10*(c*x-3)*(c*x+1)*arctanh(c*x))+3*a*b^2*(1/5*c^5*x^5*
arctanh(c*x)^2+1/10*c^4*x^4*arctanh(c*x)+1/5*c^2*x^2*arctanh(c*x)+1/5*a...
```

Fricas [F]

$$\int x^4(a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{artanh}(cx) + a)^3 x^4 dx$$

input `integrate(x^4*(a+b*arctanh(c*x))^3,x, algorithm="fricas")`

output `integral(b^3*x^4*arctanh(c*x)^3 + 3*a*b^2*x^4*arctanh(c*x)^2 + 3*a^2*b*x^4*arctanh(c*x) + a^3*x^4, x)`

Sympy [F]

$$\int x^4(a + b \operatorname{arctanh}(cx))^3 dx = \int x^4(a + b \operatorname{atanh}(cx))^3 dx$$

input `integrate(x**4*(a+b*atanh(c*x))**3,x)`

output `Integral(x**4*(a + b*atanh(c*x))**3, x)`

Maxima [F]

$$\int x^4(a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{artanh}(cx) + a)^3 x^4 dx$$

input `integrate(x^4*(a+b*arctanh(c*x))^3,x, algorithm="maxima")`

output

```
1/5*a^3*x^5 + 3/20*(4*x^5*arctanh(c*x) + c*((c^2*x^4 + 2*x^2)/c^4 + 2*log(
c^2*x^2 - 1)/c^6))*a^2*b - 1/80*(2*(b^3*c^5*x^5 - b^3)*log(-c*x + 1)^3 - 3
*(4*a*b^2*c^5*x^5 + b^3*c^4*x^4 + 2*b^3*c^2*x^2 + 2*(b^3*c^5*x^5 + b^3)*lo
g(c*x + 1))*log(-c*x + 1)^2)/c^5 - integrate(-1/40*(5*(b^3*c^5*x^5 - b^3*c
^4*x^4)*log(c*x + 1)^3 + 30*(a*b^2*c^5*x^5 - a*b^2*c^4*x^4)*log(c*x + 1)^2
- 3*(4*a*b^2*c^5*x^5 + b^3*c^4*x^4 + 2*b^3*c^2*x^2 + 5*(b^3*c^5*x^5 - b^3
*c^4*x^4)*log(c*x + 1)^2 - 2*(10*a*b^2*c^4*x^4 - (10*a*b^2*c^5 + b^3*c^5)*
x^5 - b^3)*log(c*x + 1))*log(-c*x + 1))/(c^5*x - c^4), x)
```

Giac [F]

$$\int x^4(a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{artanh}(cx) + a)^3 x^4 dx$$

input

```
integrate(x^4*(a+b*arctanh(c*x))^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x) + a)^3*x^4, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^4(a + b \operatorname{arctanh}(cx))^3 dx = \int x^4(a + b \operatorname{atanh}(cx))^3 dx$$

input

```
int(x^4*(a + b*atanh(c*x))^3,x)
```

output

```
int(x^4*(a + b*atanh(c*x))^3, x)
```

Reduce [F]

$$\int x^4 (a + b \operatorname{arctanh}(cx))^3 dx$$

$$= \frac{12 \left(\int \operatorname{atanh}(cx)^2 dx \right) a b^2 c + 20 \operatorname{atanh}(cx) b^3 + b^3 c^2 x^2 + 4 \operatorname{atanh}(cx)^3 b^3 c^5 x^5 - 4 \operatorname{atanh}(cx)^3 b^3 c x + 3 \operatorname{atanh}(cx)^3 b^3 c^2 x^2}{20 c^5}$$

input `int(x^4*(a+b*atanh(c*x))^3,x)`

output

```
(4*atanh(c*x)**3*b**3*c**5*x**5 - 4*atanh(c*x)**3*b**3*c*x + 12*atanh(c*x)
**2*a*b**2*c**5*x**5 - 12*atanh(c*x)**2*a*b**2*c*x + 3*atanh(c*x)**2*b**3*
c**4*x**4 + 6*atanh(c*x)**2*b**3*c**2*x**2 - 9*atanh(c*x)**2*b**3 + 12*ata
nh(c*x)*a**2*b*c**5*x**5 + 12*atanh(c*x)*a**2*b + 6*atanh(c*x)*a*b**2*c**4
*x**4 + 12*atanh(c*x)*a*b**2*c**2*x**2 - 18*atanh(c*x)*a*b**2 + 2*atanh(c*
x)*b**3*c**3*x**3 + 18*atanh(c*x)*b**3*c*x + 20*atanh(c*x)*b**3 + 4*int(at
anh(c*x)**3,x)*b**3*c + 12*int(atanh(c*x)**2,x)*a*b**2*c + 12*log(c**2*x -
c)*a**2*b + 20*log(c**2*x - c)*b**3 + 4*a**3*c**5*x**5 + 3*a**2*b*c**4*x*
*4 + 6*a**2*b*c**2*x**2 + 2*a*b**2*c**3*x**3 + 18*a*b**2*c*x + b**3*c**2*x
**2)/(20*c**5)
```

3.26 $\int x^3(a + \operatorname{arctanh}(cx))^3 dx$

Optimal result	294
Mathematica [A] (verified)	295
Rubi [A] (verified)	295
Maple [B] (verified)	300
Fricas [F]	301
Sympy [F]	301
Maxima [F]	302
Giac [F]	302
Mupad [F(-1)]	303
Reduce [F]	303

Optimal result

Integrand size = 14, antiderivative size = 185

$$\begin{aligned} \int x^3(a + \operatorname{arctanh}(cx))^3 dx = & \frac{b^3x}{4c^3} - \frac{b^3\operatorname{arctanh}(cx)}{4c^4} + \frac{b^2x^2(a + \operatorname{arctanh}(cx))}{4c^2} \\ & + \frac{b(a + \operatorname{arctanh}(cx))^2}{c^4} + \frac{3bx(a + \operatorname{arctanh}(cx))^2}{4c^3} \\ & + \frac{bx^3(a + \operatorname{arctanh}(cx))^2}{4c} - \frac{(a + \operatorname{arctanh}(cx))^3}{4c^4} \\ & + \frac{1}{4}x^4(a + \operatorname{arctanh}(cx))^3 \\ & - \frac{2b^2(a + \operatorname{arctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{c^4} \\ & - \frac{b^3 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{c^4} \end{aligned}$$

output

```
1/4*b^3*x/c^3-1/4*b^3*arctanh(c*x)/c^4+1/4*b^2*x^2*(a+b*arctanh(c*x))/c^2+
b*(a+b*arctanh(c*x))^2/c^4+3/4*b*x*(a+b*arctanh(c*x))^2/c^3+1/4*b*x^3*(a+b
*arctanh(c*x))^2/c-1/4*(a+b*arctanh(c*x))^3/c^4+1/4*x^4*(a+b*arctanh(c*x))
^3-2*b^2*(a+b*arctanh(c*x))*ln(2/(-c*x+1))/c^4-b^3*polylog(2,1-2/(-c*x+1))
/c^4
```

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.32

$$\int x^3(a + b \operatorname{arctanh}(cx))^3 dx$$

$$= \frac{-2ab^2 + 6a^2bcx + 2b^3cx + 2ab^2c^2x^2 + 2a^2bc^3x^3 + 2a^3c^4x^4 + 2b^2(b(-4 + 3cx + c^3x^3) + 3a(-1 + c^4x^4))}{8c^4}$$

input

```
Integrate[x^3*(a + b*ArcTanh[c*x])^3,x]
```

output

```
(-2*a*b^2 + 6*a^2*b*c*x + 2*b^3*c*x + 2*a*b^2*c^2*x^2 + 2*a^2*b*c^3*x^3 +
2*a^3*c^4*x^4 + 2*b^2*(b*(-4 + 3*c*x + c^3*x^3) + 3*a*(-1 + c^4*x^4))*ArcT
anh[c*x]^2 + 2*b^3*(-1 + c^4*x^4)*ArcTanh[c*x]^3 + 2*b*ArcTanh[c*x]*(3*a^2
*c^4*x^4 + b^2*(-1 + c^2*x^2) + 2*a*b*c*x*(3 + c^2*x^2) - 8*b^2*Log[1 + E^
(-2*ArcTanh[c*x])]) + 3*a^2*b*Log[1 - c*x] - 3*a^2*b*Log[1 + c*x] + 8*a*b^
2*Log[1 - c^2*x^2] + 8*b^3*PolyLog[2, -E^(-2*ArcTanh[c*x])])/(8*c^4)
```

Rubi [A] (verified)

Time = 3.06 (sec) , antiderivative size = 290, normalized size of antiderivative = 1.57, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.929$, Rules used = {6452, 6542, 6452, 6542, 6436, 6452, 262, 219, 6510, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3(a + b \operatorname{arctanh}(cx))^3 dx$$

$$\downarrow 6452$$

$$\frac{1}{4}x^4(a + b \operatorname{arctanh}(cx))^3 - \frac{3}{4}bc \int \frac{x^4(a + b \operatorname{arctanh}(cx))^2}{1 - c^2x^2} dx$$

$$\downarrow 6542$$

$$\frac{1}{4}x^4(a + b \operatorname{arctanh}(cx))^3 - \frac{3}{4}bc \left(\frac{\int \frac{x^2(a + b \operatorname{arctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\int x^2(a + b \operatorname{arctanh}(cx))^2 dx}{c^2} \right)$$

↓ 6452

$$\frac{1}{4}x^4(a + \operatorname{barctanh}(cx))^3 - \frac{3}{4}bc \left(\frac{\int \frac{x^2(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^2 - \frac{2}{3}bc \int \frac{x^3(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} \right)$$

↓ 6542

$$\frac{1}{4}x^4(a + \operatorname{barctanh}(cx))^3 - \frac{3}{4}bc \left(\frac{\int \frac{(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\int (a + \operatorname{barctanh}(cx))^2 dx}{c^2} - \frac{\frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \int x(a + \operatorname{barctanh}(cx)) dx \right)}{c^2} \right)$$

↓ 6436

$$\frac{1}{4}x^4(a + \operatorname{barctanh}(cx))^3 - \frac{3}{4}bc \left(\frac{\int \frac{(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{x(a + \operatorname{barctanh}(cx))^2 - 2bc \int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \int x(a + \operatorname{barctanh}(cx)) dx \right)}{c^2} \right)$$

↓ 6452

$$\frac{1}{4}x^4(a + \operatorname{barctanh}(cx))^3 - \frac{3}{4}bc \left(\frac{\int \frac{(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{x(a + \operatorname{barctanh}(cx))^2 - 2bc \int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \int x(a + \operatorname{barctanh}(cx)) dx \right)}{c^2} \right)$$

↓ 262

$$\frac{1}{4}x^4(a + \operatorname{barctanh}(cx))^3 - \frac{3}{4}bc \left(\frac{\int \frac{(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{x(a + \operatorname{barctanh}(cx))^2 - 2bc \int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^2 - \frac{2}{3}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} - \int x(a + \operatorname{barctanh}(cx)) dx \right)}{c^2} \right)$$

$$\begin{aligned} & \downarrow 219 \\ & \frac{1}{4}x^4(a + \operatorname{arctanh}(cx))^3 - \\ \frac{3}{4}bc & \left(\frac{\int \frac{(a+b\operatorname{arctanh}(cx))^2}{1-c^2x^2} dx}{c^2} - \frac{x(a+b\operatorname{arctanh}(cx))^2 - 2bc \int \frac{x(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a + \operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \left(\int \frac{x(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx \right)}{c^2} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 6510 \\ & \frac{1}{4}x^4(a + \operatorname{arctanh}(cx))^3 - \\ \frac{3}{4}bc & \left(\frac{\frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^3}}{c^2} - \frac{x(a+b\operatorname{arctanh}(cx))^2 - 2bc \int \frac{x(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx}{c^2} - \frac{\frac{1}{3}x^3(a + \operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \left(\int \frac{x(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx \right)}{c^2} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 6546 \\ & \frac{1}{4}x^4(a + \operatorname{arctanh}(cx))^3 - \\ \frac{3}{4}bc & \left(\frac{\frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^3}}{c^2} - \frac{x(a+b\operatorname{arctanh}(cx))^2 - 2bc \left(\frac{\int \frac{a+b\operatorname{arctanh}(cx)}{1-cx} dx}{c} - \frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^2} \right)}{c^2} - \frac{\frac{1}{3}x^3(a + \operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \left(\int \frac{x(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx \right)}{c^2} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 6470 \\ & \frac{1}{4}x^4(a + \operatorname{arctanh}(cx))^3 - \\ \frac{3}{4}bc & \left(\frac{\frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^3}}{c^2} - \frac{x(a+b\operatorname{arctanh}(cx))^2 - 2bc \left(\frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))}{c} - b \int \frac{\log\left(\frac{2}{1-cx}\right)}{1-c^2x^2} dx - \frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^2} \right)}{c^2} - \frac{\frac{1}{3}x^3(a + \operatorname{arctanh}(cx))^2 - \frac{2}{3}bc \left(\int \frac{x(a+b\operatorname{arctanh}(cx))}{1-c^2x^2} dx \right)}{c^2} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 2849 \end{aligned}$$

$$\frac{1}{4}x^4(a + \operatorname{arctanh}(cx))^3 - \frac{3}{4}bc \frac{\frac{(a + \operatorname{arctanh}(cx))^3}{3bc^3} - \frac{x(a + \operatorname{arctanh}(cx))^2 - 2bc \left(\frac{b \int \frac{\log\left(\frac{2}{1-cx}\right) dx}{1-cx} + \frac{\log\left(\frac{2}{1-cx}\right)(a + \operatorname{arctanh}(cx))}{c} - \frac{(a + \operatorname{arctanh}(cx))^2}{2bc^2} \right)}{c^2}}{c^2}}$$

↓ 2752

$$\frac{1}{4}x^4(a + \operatorname{arctanh}(cx))^3 - \frac{3}{4}bc \frac{\frac{(a + \operatorname{arctanh}(cx))^3}{3bc^3} - \frac{x(a + \operatorname{arctanh}(cx))^2 - 2bc \left(\frac{\log\left(\frac{2}{1-cx}\right)(a + \operatorname{arctanh}(cx))}{c} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{2c} - \frac{(a + \operatorname{arctanh}(cx))^2}{2bc^2} \right)}{c^2}}{c^2}}$$

input `Int [x^3*(a + b*ArcTanh[c*x])^3,x]`

output `(x^4*(a + b*ArcTanh[c*x])^3)/4 - (3*b*c*(-(((x^3*(a + b*ArcTanh[c*x])^2)/3 - (2*b*c*(-(((x^2*(a + b*ArcTanh[c*x]))/2 - (b*c*(-(x/c^2) + ArcTanh[c*x]/c^3))/2)/c^2) + (-1/2*(a + b*ArcTanh[c*x])^2/(b*c^2) + (((a + b*ArcTanh[c*x])*Log[2/(1 - c*x)])/c + (b*PolyLog[2, 1 - 2/(1 - c*x)]/(2*c))/c)/c^2))/3)/c^2) + ((a + b*ArcTanh[c*x])^3/(3*b*c^3) - (x*(a + b*ArcTanh[c*x])^2 - 2*b*c*(-1/2*(a + b*ArcTanh[c*x])^2/(b*c^2) + (((a + b*ArcTanh[c*x])*Log[2/(1 - c*x)])/c + (b*PolyLog[2, 1 - 2/(1 - c*x)]/(2*c))/c))/c^2)/c^2)/4`

Defintions of rubi rules used

rule 219 $\text{Int}[\{(a_)+(b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /;$ $\text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 262 $\text{Int}[\{(c_)*(x_)\}^{(m_)}*\{(a_)+(b_)*(x_)^2\}^{(p_)}, x_Symbol] \rightarrow \text{Simp}[c*(c*x)^{(m-1)}*((a+b*x^2)^{(p+1)}/(b*(m+2*p+1))), x] - \text{Simp}[a*c^{2*((m-1)/(b*(m+2*p+1)))} \ \text{Int}[(c*x)^{(m-2)}*(a+b*x^2)^p, x], x] /;$ $\text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{GtQ}[m, 2-1] \ \&\& \ \text{NeQ}[m+2*p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 2752 $\text{Int}[\text{Log}[(c_)*(x_)]/((d_)+(e_)*(x_)), x_Symbol] \rightarrow \text{Simp}[(-e^{-1})*\text{PolyLog}[2, 1-c*x], x] /;$ $\text{FreeQ}\{c, d, e\}, x \ \&\& \ \text{EqQ}[e+c*d, 0]$

rule 2849 $\text{Int}[\text{Log}[(c_)/((d_)+(e_)*(x_))]/((f_)+(g_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[-e/g \ \text{Subst}[\text{Int}[\text{Log}[2*d*x]/(1-2*d*x), x], x, 1/(d+e*x)], x] /;$ $\text{FreeQ}\{c, d, e, f, g\}, x \ \&\& \ \text{EqQ}[c, 2*d] \ \&\& \ \text{EqQ}[e^2*f+d^2*g, 0]$

rule 6436 $\text{Int}[\{(a_)+\text{ArcTanh}[(c_)*(x_)^{(n_)}]\}*(b_)\}^{(p_)}, x_Symbol] \rightarrow \text{Simp}[x*(a+b*\text{ArcTanh}[c*x^n])^p, x] - \text{Simp}[b*c^n*p \ \text{Int}[x^n*((a+b*\text{ArcTanh}[c*x^n])^{(p-1)}/(1-c^2*x^{(2*n)})), x], x] /;$ $\text{FreeQ}\{a, b, c, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[n, 1] \ || \ \text{EqQ}[p, 1])$

rule 6452 $\text{Int}[\{(a_)+\text{ArcTanh}[(c_)*(x_)^{(n_)}]\}*(b_)\}^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a+b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c^n*(p/(m+1)) \ \text{Int}[x^{(m+n)}*((a+b*\text{ArcTanh}[c*x^n])^{(p-1)}/(1-c^2*x^{(2*n)})), x], x] /;$ $\text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 6470

```
Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol
] := Simp[(-a + b*ArcTanh[c*x])^p*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c
*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^
2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2
, 0]
```

rule 6510

```
Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symb
ol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b
, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]
```

rule 6542

```
Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_))/((d_) + (
e_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*
x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/
(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m,
1]
```

rule 6546

```
Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*(x_)/((d_) + (e_.)*(x_)^2),
x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/
(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 666 vs. $2(171) = 342$.

Time = 8.06 (sec) , antiderivative size = 667, normalized size of antiderivative = 3.61

method	result
risch	$\frac{b^3x}{4c^3} + \frac{3\ln(-cx+1)^2ab^2x^4}{16} - \frac{3\ln(-cx+1)a^2bx^4}{8} - \frac{b^3}{4c^4} - \frac{3ab^2\ln(-cx+1)x}{4c^3} + \frac{ab^2x^2}{4c^2} + \frac{a^2bx^3}{4c} + \frac{3a^2bx}{4c^3}$
derivativdivides	Expression too large to display
default	Expression too large to display
parts	Expression too large to display

input

```
int(x^3*(a+b*arctanh(c*x))^3,x,method=_RETURNVERBOSE)
```

output

```
1/4*b^3*x/c^3+3/16*ln(-c*x+1)^2*a*b^2*x^4-3/8*ln(-c*x+1)*a^2*b*x^4-1/4*b^3
/c^4-3/4/c^3*a*b^2*ln(-c*x+1)*x+1/4/c^2*a*b^2*x^2+1/4/c*a^2*b*x^3+3/4/c^3*
a^2*b*x+3/8/c^4*a^2*b*ln(-c*x+1)-3/16/c^4*a*b^2*ln(-c*x+1)^2+1/c^4*a*b^2*ln
(-c*x+1)+1/16/c*b^3*ln(-c*x+1)^2*x^3+3/16/c^3*b^3*ln(-c*x+1)^2*x-1/8/c^2*
b^3*ln(-c*x+1)*x^2+1/4*a^3*x^4-1/4/c^4*a*b^2-1/c^4*a^2*b-1/4/c^4*a^3-1/8/c
^4*b^3*ln(-c*x-1)+b^3/c^4*dilog(-1/2*c*x+1/2)+1/32/c^4*b^3*ln(-c*x+1)^3-1/
4/c^4*b^3*ln(-c*x+1)^2+1/8/c^4*b^3*ln(-c*x+1)+(3/32*b^3*(c^4*x^4-1)/c^4*ln
(-c*x+1)^2-1/8*b^2*x*(3*a*c^3*x^3+b*c^2*x^2+3*b)/c^3*ln(-c*x+1)-1/8*b*(-3*
a^2*c^4*x^4-2*a*b*c^3*x^3-b^2*c^2*x^2-6*a*b*c*x-3*b*ln(-c*x+1)*a-4*b^2*ln(
-c*x+1))/c^4)*ln(c*x+1)-1/4/c*a*b^2*ln(-c*x+1)*x^3+1/32*b^3*(c^4*x^4-1)/c^
4*ln(c*x+1)^3+1/32*b^2*(-3*b*x^4*ln(-c*x+1)*c^4+6*a*c^4*x^4+2*b*c^3*x^3+6*
b*c*x+3*b*ln(-c*x+1)-6*a+8*b)/c^4*ln(c*x+1)^2-1/32*ln(-c*x+1)^3*b^3*x^4+b^
2/c^4*ln(-c*x-1)*a-3/8*b/c^4*ln(-c*x-1)*a^2+b^3/c^4*ln(-1/2*c*x+1/2)*ln(1/
2*c*x+1/2)-b^3/c^4*ln(1/2*c*x+1/2)*ln(-c*x+1)
```

Fricas [F]

$$\int x^3(a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{artanh}(cx) + a)^3 x^3 dx$$

input

```
integrate(x^3*(a+b*arctanh(c*x))^3,x, algorithm="fricas")
```

output

```
integral(b^3*x^3*arctanh(c*x)^3 + 3*a*b^2*x^3*arctanh(c*x)^2 + 3*a^2*b*x^3
*arctanh(c*x) + a^3*x^3, x)
```

Sympy [F]

$$\int x^3(a + b \operatorname{arctanh}(cx))^3 dx = \int x^3(a + b \operatorname{atanh}(cx))^3 dx$$

input

```
integrate(x**3*(a+b*atanh(c*x))**3,x)
```

output

```
Integral(x**3*(a + b*atanh(c*x))**3, x)
```

Maxima [F]

$$\int x^3(a + \operatorname{arctanh}(cx))^3 dx = \int (\operatorname{arctanh}(cx) + a)^3 x^3 dx$$

input `integrate(x^3*(a+b*arctanh(c*x))^3,x, algorithm="maxima")`

output

```

3/4*a*b^2*x^4*arctanh(c*x)^2 + 1/4*a^3*x^4 + 1/8*(6*x^4*arctanh(c*x) + c*(
2*(c^2*x^3 + 3*x)/c^4 - 3*log(c*x + 1)/c^5 + 3*log(c*x - 1)/c^5))*a^2*b +
1/16*(4*c*(2*(c^2*x^3 + 3*x)/c^4 - 3*log(c*x + 1)/c^5 + 3*log(c*x - 1)/c^5
)*arctanh(c*x) + (4*c^2*x^2 - 2*(3*log(c*x - 1) - 8)*log(c*x + 1) + 3*log(
c*x + 1)^2 + 3*log(c*x - 1)^2 + 16*log(c*x - 1))/c^4)*a*b^2 - 1/9216*(27*c
^5*((c^2*x^4 + 2*x^2)/c^7 + 2*log(c^2*x^2 - 1)/c^9) + 74*c^4*(2*(c^2*x^3 +
3*x)/c^7 - 3*log(c*x + 1)/c^8 + 3*log(c*x - 1)/c^8) + 60*c^3*(x^2/c^5 + l
og(c^2*x^2 - 1)/c^7) - 221184*c^3*integrate(1/96*x^3*log(c*x + 1)/(c^5*x^2
- c^3), x) + 1692*c^2*(2*x/c^5 - log(c*x + 1)/c^6 + log(c*x - 1)/c^6) - 1
105920*c*integrate(1/96*x*log(c*x + 1)/(c^5*x^2 - c^3), x) + (9*(32*log(-c
*x + 1)^3 - 24*log(-c*x + 1)^2 + 12*log(-c*x + 1) - 3)*(c*x - 1)^4 + 128*(
9*log(-c*x + 1)^3 - 9*log(-c*x + 1)^2 + 6*log(-c*x + 1) - 2)*(c*x - 1)^3 +
432*(4*log(-c*x + 1)^3 - 6*log(-c*x + 1)^2 + 6*log(-c*x + 1) - 3)*(c*x -
1)^2 + 1152*(log(-c*x + 1)^3 - 3*log(-c*x + 1)^2 + 6*log(-c*x + 1) - 6)*(c
*x - 1))/c^4 - 12*(24*(c^4*x^4 - 1)*log(c*x + 1)^3 + 48*(c^3*x^3 + 3*c*x)*
log(c*x + 1)^2 - 6*(3*c^4*x^4 - 4*c^3*x^3 + 6*c^2*x^2 - 12*c*x - 12*(c^4*x
^4 - 1)*log(c*x + 1) + 7)*log(-c*x + 1)^2 + (9*c^4*x^4 + 28*c^3*x^3 - 18*c
^2*x^2 - 72*(c^4*x^4 - 1)*log(c*x + 1)^2 + 300*c*x - 96*(c^3*x^3 + 3*c*x +
4)*log(c*x + 1))*log(-c*x + 1))/c^4 + 1800*log(96*c^5*x^2 - 96*c^3)/c^4 -
442368*integrate(1/96*log(c*x + 1)/(c^5*x^2 - c^3), x))*b^3

```

Giac [F]

$$\int x^3(a + \operatorname{arctanh}(cx))^3 dx = \int (\operatorname{arctanh}(cx) + a)^3 x^3 dx$$

input `integrate(x^3*(a+b*arctanh(c*x))^3,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^3*x^3, x)`

3.27 $\int x^2(a + b \operatorname{arctanh}(cx))^3 dx$

Optimal result	304
Mathematica [A] (verified)	305
Rubi [A] (verified)	305
Maple [C] (warning: unable to verify)	309
Fricas [F]	310
Sympy [F]	311
Maxima [F]	311
Giac [F]	311
Mupad [F(-1)]	312
Reduce [F]	312

Optimal result

Integrand size = 14, antiderivative size = 197

$$\int x^2(a + b \operatorname{arctanh}(cx))^3 dx = \frac{ab^2x}{c^2} + \frac{b^3x \operatorname{arctanh}(cx)}{c^2} - \frac{b(a + b \operatorname{arctanh}(cx))^2}{2c^3} + \frac{bx^2(a + b \operatorname{arctanh}(cx))^2}{2c} + \frac{(a + b \operatorname{arctanh}(cx))^3}{3c^3} + \frac{1}{3}x^3(a + b \operatorname{arctanh}(cx))^3 - \frac{b(a + b \operatorname{arctanh}(cx))^2 \log\left(\frac{2}{1-cx}\right)}{c^3} + \frac{b^3 \log(1 - c^2x^2)}{2c^3} - \frac{b^2(a + b \operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{c^3} + \frac{b^3 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx}\right)}{2c^3}$$

output

```
a*b^2*x/c^2+b^3*x*arctanh(c*x)/c^2-1/2*b*(a+b*arctanh(c*x))^2/c^3+1/2*b*x^2*(a+b*arctanh(c*x))^2/c+1/3*(a+b*arctanh(c*x))^3/c^3+1/3*x^3*(a+b*arctanh(c*x))^3-b*(a+b*arctanh(c*x))^2*ln(2/(-c*x+1))/c^3+1/2*b^3*ln(-c^2*x^2+1)/c^3-b^2*(a+b*arctanh(c*x))*polylog(2,1-2/(-c*x+1))/c^3+1/2*b^3*polylog(3,1-2/(-c*x+1))/c^3
```

Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.27

$$\int x^2(a + b \operatorname{arctanh}(cx))^3 dx$$

$$= \frac{3a^2bc^2x^2 + 2a^3c^3x^3 + 6a^2bc^3x^3 \operatorname{arctanh}(cx) + 3a^2b \log(1 - c^2x^2) + 6ab^2(cx + (-1 + c^3x^3) \operatorname{arctanh}(cx))^2}{6c^3}$$

input `Integrate[x^2*(a + b*ArcTanh[c*x])^3,x]`

output $(3a^2bc^2x^2 + 2a^3c^3x^3 + 6a^2bc^3x^3 \operatorname{ArcTanh}[cx] + 3a^2b \operatorname{Log}[1 - c^2x^2] + 6ab^2(cx + (-1 + c^3x^3) \operatorname{ArcTanh}[cx])^2 + \operatorname{ArcTanh}[cx](-1 + c^2x^2 - 2 \operatorname{Log}[1 + E^{-2 \operatorname{ArcTanh}[cx]}])) + \operatorname{PolyLog}[2, -E^{-2 \operatorname{ArcTanh}[cx]}]) + b^3(6cx \operatorname{ArcTanh}[cx] - 3 \operatorname{ArcTanh}[cx]^2 + 3c^2x^2 \operatorname{ArcTanh}[cx]^2 - 2 \operatorname{ArcTanh}[cx]^3 + 2c^3x^3 \operatorname{ArcTanh}[cx]^3 - 6 \operatorname{ArcTanh}[cx]^2 \operatorname{Log}[1 + E^{-2 \operatorname{ArcTanh}[cx]}] + 3 \operatorname{Log}[1 - c^2x^2] + 6 \operatorname{ArcTanh}[cx] \operatorname{PolyLog}[2, -E^{-2 \operatorname{ArcTanh}[cx]}] + 3 \operatorname{PolyLog}[3, -E^{-2 \operatorname{ArcTanh}[cx]}])))/(6c^3)$

Rubi [A] (verified)

Time = 1.96 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.10, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {6452, 6542, 6452, 6542, 2009, 6510, 6546, 6470, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2(a + b \operatorname{arctanh}(cx))^3 dx$$

$$\downarrow \text{6452}$$

$$\frac{1}{3}x^3(a + b \operatorname{arctanh}(cx))^3 - bc \int \frac{x^3(a + b \operatorname{arctanh}(cx))^2}{1 - c^2x^2} dx$$

$$\downarrow \text{6542}$$

$$\begin{aligned}
& \frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^3 - bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\int x(a + \operatorname{barctanh}(cx))^2 dx}{c^2} \right) \\
& \quad \downarrow 6452 \\
& \frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^3 - \\
& bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx))^2 - bc \int \frac{x^2(a + \operatorname{barctanh}(cx))}{1 - c^2x^2} dx}{c^2} \right) \\
& \quad \downarrow 6542 \\
& \frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^3 - \\
& bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx))^2 - bc \left(\frac{\int \frac{a + \operatorname{barctanh}(cx)}{1 - c^2x^2} dx}{c^2} - \frac{\int (a + \operatorname{barctanh}(cx)) dx}{c^2} \right)}{c^2} \right) \\
& \quad \downarrow 2009 \\
& \frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^3 - \\
& bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx))^2 - bc \left(\frac{\int \frac{a + \operatorname{barctanh}(cx)}{1 - c^2x^2} dx}{c^2} - \frac{ax + bx \operatorname{arctanh}(cx) + \frac{b \log(1 - c^2x^2)}{2c}}{c^2} \right)}{c^2} \right) \\
& \quad \downarrow 6510 \\
& \frac{1}{3}x^3(a + \operatorname{barctanh}(cx))^3 - \\
& bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{barctanh}(cx))^2 - bc \left(\frac{(a + \operatorname{barctanh}(cx))^2}{2bc^3} - \frac{ax + bx \operatorname{arctanh}(cx) + \frac{b \log(1 - c^2x^2)}{2c}}{c^2} \right)}{c^2} \right) \\
& \quad \downarrow 6546
\end{aligned}$$

$$bc \left(\frac{\frac{1}{3}x^3(a + \operatorname{arctanh}(cx))^3 - \int \frac{(a+b\operatorname{arctanh}(cx))^2 dx}{c} - \frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^2}}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{arctanh}(cx))^2 - bc \left(\frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^3} - \frac{ax+b\operatorname{arctanh}(cx)}{c} \right)}{c^2} \right)$$

6470

$$bc \left(\frac{\frac{1}{3}x^3(a + \operatorname{arctanh}(cx))^3 - \frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))^2}{c} - 2b \int \frac{(a+b\operatorname{arctanh}(cx)) \log\left(\frac{2}{1-cx}\right) dx}{1-c^2x^2} - \frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^2}}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{arctanh}(cx))^2 - bc \left(\frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^3} - \frac{ax+b\operatorname{arctanh}(cx)}{c} \right)}{c^2} \right)$$

6620

$$bc \left(\frac{\frac{1}{3}x^3(a + \operatorname{arctanh}(cx))^3 - \frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))^2}{c} - 2b \left(\frac{\frac{1}{2}b \int \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right) dx}{1-c^2x^2} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))}{2c} \right)}{c} - \frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^2}}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{arctanh}(cx))^2 - bc \left(\frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^3} - \frac{ax+b\operatorname{arctanh}(cx)}{c} \right)}{c^2} \right)$$

7164

$$bc \left(\frac{\frac{1}{3}x^3(a + \operatorname{arctanh}(cx))^3 - \frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))^2}{c} - 2b \left(\frac{b \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx}\right)}{4c} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))}{2c} \right)}{c} - \frac{(a+b\operatorname{arctanh}(cx))^3}{3bc^2}}{c^2} - \frac{\frac{1}{2}x^2(a + \operatorname{arctanh}(cx))^2 - bc \left(\frac{(a+b\operatorname{arctanh}(cx))^2}{2bc^3} - \frac{ax+b\operatorname{arctanh}(cx)}{c} \right)}{c^2} \right)$$

input `Int [x^2*(a + b*ArcTanh [c*x])^3, x]`

output

$$\begin{aligned} & (x^3(a + b\operatorname{ArcTanh}[c*x])^3)/3 - b*c*(-((x^2*(a + b\operatorname{ArcTanh}[c*x])^2)/2 - \\ & b*c*((a + b\operatorname{ArcTanh}[c*x])^2/(2*b*c^3) - (a*x + b*x*\operatorname{ArcTanh}[c*x] + (b*\operatorname{Log}[1 \\ & - c^2*x^2])/(2*c))/c^2))/c^2) + (-1/3*(a + b\operatorname{ArcTanh}[c*x])^3/(b*c^2) + ((\\ & (a + b\operatorname{ArcTanh}[c*x])^2*\operatorname{Log}[2/(1 - c*x)]/c - 2*b*(-1/2*((a + b\operatorname{ArcTanh}[c*x] \\ &])*\operatorname{PolyLog}[2, 1 - 2/(1 - c*x)])/c + (b*\operatorname{PolyLog}[3, 1 - 2/(1 - c*x)]/(4*c)) \\ &)/c)/c^2) \end{aligned}$$

Defintions of rubi rules used

rule 2009

$$\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] \;/; \operatorname{SumQ}[u]$$

rule 6452

$$\begin{aligned} & \operatorname{Int}[(a + \operatorname{ArcTanh}[c*x]^n] * (b*x)^m, x_Symbol] : \\ & > \operatorname{Simp}[x^{m+1} * ((a + b\operatorname{ArcTanh}[c*x]^n)^p / (m+1)), x] - \operatorname{Simp}[b*c*n*(p/(m \\ & + 1)) \operatorname{Int}[x^{m+n} * ((a + b\operatorname{ArcTanh}[c*x]^n)^{p-1} / (1 - c^2*x^{2*n})), x \\ &], x] \;/; \operatorname{FreeQ}\{a, b, c, m, n\}, x \ \&\& \operatorname{IGtQ}[p, 0] \ \&\& (\operatorname{EqQ}[p, 1] \ || \ (\operatorname{EqQ}[n, 1] \\ &] \ \&\& \operatorname{IntegerQ}[m])) \ \&\& \operatorname{NeQ}[m, -1] \end{aligned}$$

rule 6470

$$\begin{aligned} & \operatorname{Int}[(a + \operatorname{ArcTanh}[c*x])^p * (b*x)^m / ((d) + (e)*x), x_Symbol] \\ &] \rightarrow \operatorname{Simp}[(-a + b\operatorname{ArcTanh}[c*x])^p * (\operatorname{Log}[2/(1 + e*(x/d))]/e), x] + \operatorname{Simp}[b*c \\ & *(p/e) \operatorname{Int}[(a + b\operatorname{ArcTanh}[c*x])^{p-1} * (\operatorname{Log}[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] \;/; \operatorname{FreeQ}\{a, b, c, d, e\}, x \ \&\& \operatorname{IGtQ}[p, 0] \ \&\& \operatorname{EqQ}[c^2*d^2 - e^2, 0] \end{aligned}$$

rule 6510

$$\begin{aligned} & \operatorname{Int}[(a + \operatorname{ArcTanh}[c*x])^p * (b*x)^m / ((d) + (e)*x^2), x_Symbol] \\ &] \rightarrow \operatorname{Simp}[(a + b\operatorname{ArcTanh}[c*x])^{p+1} / (b*c*d*(p+1)), x] \;/; \operatorname{FreeQ}\{a, b, c, d, e, p\}, x \ \&\& \operatorname{EqQ}[c^2*d + e, 0] \ \&\& \operatorname{NeQ}[p, -1] \end{aligned}$$

rule 6542

$$\begin{aligned} & \operatorname{Int}[(a + \operatorname{ArcTanh}[c*x])^p * (b*x)^m * (f*x)^n / ((d) + (e)*x^2), x_Symbol] \\ &] \rightarrow \operatorname{Simp}[f^2/e \operatorname{Int}[(f*x)^{m-2} * (a + b\operatorname{ArcTanh}[c*x])^p, x], x] - \operatorname{Simp}[d*(f^2/e) \operatorname{Int}[(f*x)^{m-2} * (a + b\operatorname{ArcTanh}[c*x])^p / \\ & (d + e*x^2)), x], x] \;/; \operatorname{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \operatorname{GtQ}[p, 0] \ \&\& \operatorname{GtQ}[m, 1] \end{aligned}$$

rule 6546

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*(x_)/((d_) + (e_.)*(x_)^2),
 x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/
 (c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

rule 6620

```
Int[(Log[u_] * ((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.))/((d_) + (e_.)*(x_)^
 2), x_Symbol] := Simp[(-(a + b*ArcTanh[c*x])^p)*(PolyLog[2, 1 - u]/(2*c*d))
 , x] + Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(
 d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d
 + e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
 x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 10.26 (sec) , antiderivative size = 973, normalized size of antiderivative = 4.94

method	result	size
derivativedivides	Expression too large to display	973
default	Expression too large to display	973
parts	Expression too large to display	975

input

```
int(x^2*(a+b*arctanh(c*x))^3,x,method=_RETURNVERBOSE)
```

output

```

1/c^3*(1/3*a^3*c^3*x^3+b^3*(1/3*c^3*x^3*arctanh(c*x)^3+1/2*c^2*x^2*arctanh
(c*x)^2+1/2*arctanh(c*x)^2*ln(c*x-1)+1/2*arctanh(c*x)^2*ln(c*x+1)-arctanh(
c*x)*polylog(2,-(c*x+1)^2/(-c^2*x^2+1))+1/2*polylog(3,-(c*x+1)^2/(-c^2*x^2
+1))-arctanh(c*x)^2*ln((c*x+1)/(-c^2*x^2+1)^(1/2))+1/3*arctanh(c*x)^3+1/4*
I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(c*x+1)^2/(c^2*x^2-1))*csgn(
I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))*arctanh(c*x)^2-ln(1+(c*
x+1)^2/(-c^2*x^2+1))-1/2*arctanh(c*x)^2-1/4*I*Pi*csgn(I*(c*x+1)^2/(c^2*x^2
-1))^3*arctanh(c*x)^2-1/2*I*Pi*arctanh(c*x)^2+1/4*I*Pi*csgn(I*(c*x+1)^2/(c
^2*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^2*arcta
nh(c*x)^2+(c*x+1)*arctanh(c*x)-1/4*I*Pi*csgn(I*(c*x+1)/(-c^2*x^2+1)^(1/2))
^2*csgn(I*(c*x+1)^2/(c^2*x^2-1))*arctanh(c*x)^2-1/4*I*Pi*csgn(I*(c*x+1)^2/
(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^3*arctanh(c*x)^2-1/2*I*Pi*csgn(I*(c
*x+1)/(-c^2*x^2+1)^(1/2))*csgn(I*(c*x+1)^2/(c^2*x^2-1))^2*arctanh(c*x)^2+1
/2*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))^2*arctanh(c*x)^2-1/4*I*Pi*csgn(I
/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2
*x^2-1)))^2*arctanh(c*x)^2-ln(2)*arctanh(c*x)^2-1/2*I*Pi*csgn(I/(1-(c*x+1)
^2/(c^2*x^2-1)))^3*arctanh(c*x)^2)+3*a*b^2*(1/3*c^3*x^3*arctanh(c*x)^2+1/3
*c^2*x^2*arctanh(c*x)+1/3*arctanh(c*x)*ln(c*x-1)+1/3*arctanh(c*x)*ln(c*x+1
)+1/3*c*x+1/6*ln(c*x-1)-1/6*ln(c*x+1)+1/12*ln(c*x-1)^2-1/3*dilog(1/2*c*x+1
/2)-1/6*ln(c*x-1)*ln(1/2*c*x+1/2)-1/12*ln(c*x+1)^2+1/6*(ln(c*x+1)-ln(1/...

```

Fricas [F]

$$\int x^2(a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{arctanh}(cx) + a)^3 x^2 dx$$

input

```
integrate(x^2*(a+b*arctanh(c*x))^3,x, algorithm="fricas")
```

output

```
integral(b^3*x^2*arctanh(c*x)^3 + 3*a*b^2*x^2*arctanh(c*x)^2 + 3*a^2*b*x^2
*arctanh(c*x) + a^3*x^2, x)
```

Sympy [F]

$$\int x^2(a + b \operatorname{arctanh}(cx))^3 dx = \int x^2(a + b \operatorname{atanh}(cx))^3 dx$$

input `integrate(x**2*(a+b*atanh(c*x))**3,x)`

output `Integral(x**2*(a + b*atanh(c*x))**3, x)`

Maxima [F]

$$\int x^2(a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{artanh}(cx) + a)^3 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c*x))^3,x, algorithm="maxima")`

output `1/3*a^3*x^3 + 1/2*(2*x^3*arctanh(c*x) + c*(x^2/c^2 + log(c^2*x^2 - 1)/c^4)
)*a^2*b - 1/24*((b^3*c^3*x^3 - b^3)*log(-c*x + 1)^3 - 3*(2*a*b^2*c^3*x^3 +
b^3*c^2*x^2 + (b^3*c^3*x^3 + b^3)*log(c*x + 1))*log(-c*x + 1)^2)/c^3 - in
tegrate(-1/8*((b^3*c^3*x^3 - b^3*c^2*x^2)*log(c*x + 1)^3 + 6*(a*b^2*c^3*x^
3 - a*b^2*c^2*x^2)*log(c*x + 1)^2 - (4*a*b^2*c^3*x^3 + 2*b^3*c^2*x^2 + 3*(
b^3*c^3*x^3 - b^3*c^2*x^2)*log(c*x + 1)^2 - 2*(6*a*b^2*c^2*x^2 - (6*a*b^2*
c^3 + b^3*c^3)*x^3 - b^3)*log(c*x + 1))*log(-c*x + 1))/(c^3*x - c^2), x)`

Giac [F]

$$\int x^2(a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{artanh}(cx) + a)^3 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c*x))^3,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^3*x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^2(a + b \operatorname{arctanh}(cx))^3 dx = \int x^2(a + b \operatorname{atanh}(cx))^3 dx$$

input `int(x^2*(a + b*atanh(c*x))^3,x)`output `int(x^2*(a + b*atanh(c*x))^3, x)`**Reduce [F]**

$$\int x^2(a + b \operatorname{arctanh}(cx))^3 dx$$

$$= \frac{2 \operatorname{atanh}(cx)^3 b^3 c^3 x^3 - 2 \operatorname{atanh}(cx)^3 b^3 cx + 6 \operatorname{atanh}(cx)^2 a b^2 c^3 x^3 - 6 \operatorname{atanh}(cx)^2 a b^2 cx + 3 \operatorname{atanh}(cx)^2 b^3 c^2 x^2}{6c^3}$$

input `int(x^2*(a+b*atanh(c*x))^3,x)`output `(2*atanh(c*x)**3*b**3*c**3*x**3 - 2*atanh(c*x)**3*b**3*c*x + 6*atanh(c*x)*
*2*a*b**2*c**3*x**3 - 6*atanh(c*x)**2*a*b**2*c*x + 3*atanh(c*x)**2*b**3*c*
*2*x**2 - 3*atanh(c*x)**2*b**3 + 6*atanh(c*x)*a**2*b*c**3*x**3 + 6*atanh(c
*x)*a**2*b + 6*atanh(c*x)*a*b**2*c**2*x**2 - 6*atanh(c*x)*a*b**2 + 6*atanh
(c*x)*b**3*c*x + 6*atanh(c*x)*b**3 + 2*int(atanh(c*x)**3,x)*b**3*c + 6*int
(atanh(c*x)**2,x)*a*b**2*c + 6*log(c**2*x - c)*a**2*b + 6*log(c**2*x - c)*
b**3 + 2*a**3*c**3*x**3 + 3*a**2*b*c**2*x**2 + 6*a*b**2*c*x)/(6*c**3)`

3.28 $\int x(a + \operatorname{barctanh}(cx))^3 dx$

Optimal result	313
Mathematica [A] (verified)	314
Rubi [A] (verified)	314
Maple [B] (verified)	317
Fricas [F]	318
Sympy [F]	318
Maxima [F]	319
Giac [F]	319
Mupad [F(-1)]	320
Reduce [F]	320

Optimal result

Integrand size = 12, antiderivative size = 123

$$\int x(a + \operatorname{barctanh}(cx))^3 dx = \frac{3b(a + \operatorname{barctanh}(cx))^2}{2c^2} + \frac{3bx(a + \operatorname{barctanh}(cx))^2}{2c} - \frac{(a + \operatorname{barctanh}(cx))^3}{2c^2} + \frac{1}{2}x^2(a + \operatorname{barctanh}(cx))^3 - \frac{3b^2(a + \operatorname{barctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{c^2} - \frac{3b^3 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{2c^2}$$

output

```
3/2*b*(a+b*arctanh(c*x))^2/c^2+3/2*b*x*(a+b*arctanh(c*x))^2/c-1/2*(a+b*arctanh(c*x))^3/c^2+1/2*x^2*(a+b*arctanh(c*x))^3-3*b^2*(a+b*arctanh(c*x))*ln(2/(-c*x+1))/c^2-3/2*b^3*polylog(2,1-2/(-c*x+1))/c^2
```

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.31

$$\int x(a + b \operatorname{arctanh}(cx))^3 dx$$

$$= \frac{6b^2(-1 + cx)(a + b + acx) \operatorname{arctanh}(cx)^2 + 2b^3(-1 + c^2x^2) \operatorname{arctanh}(cx)^3 + 6b \operatorname{arctanh}(cx) (acx(2b + acx))}{4c^2}$$

input

```
Integrate[x*(a + b*ArcTanh[c*x])^3,x]
```

output

```
(6*b^2*(-1 + c*x)*(a + b + a*c*x)*ArcTanh[c*x]^2 + 2*b^3*(-1 + c^2*x^2)*ArcTanh[c*x]^3 + 6*b*ArcTanh[c*x]*(a*c*x*(2*b + a*c*x) - 2*b^2*Log[1 + E^(-2*ArcTanh[c*x])]) + a*(6*a*b*c*x + 2*a^2*c^2*x^2 + 3*a*b*Log[1 - c*x] - 3*a*b*Log[1 + c*x] + 6*b^2*Log[1 - c^2*x^2]) + 6*b^3*PolyLog[2, -E^(-2*ArcTanh[c*x])])/(4*c^2)
```

Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.12, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {6452, 6542, 6436, 6510, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(a + b \operatorname{arctanh}(cx))^3 dx$$

$$\downarrow 6452$$

$$\frac{1}{2}x^2(a + b \operatorname{arctanh}(cx))^3 - \frac{3}{2}bc \int \frac{x^2(a + b \operatorname{arctanh}(cx))^2}{1 - c^2x^2} dx$$

$$\downarrow 6542$$

$$\frac{1}{2}x^2(a + b \operatorname{arctanh}(cx))^3 - \frac{3}{2}bc \left(\frac{\int \frac{(a + b \operatorname{arctanh}(cx))^2}{1 - c^2x^2} dx}{c^2} - \frac{\int (a + b \operatorname{arctanh}(cx))^2 dx}{c^2} \right)$$

$$\downarrow 6436$$

$$\frac{3}{2}bc \left(\frac{\int \frac{(a+\operatorname{barctanh}(cx))^2}{1-c^2x^2} dx}{c^2} - \frac{\frac{1}{2}x^2(a+\operatorname{barctanh}(cx))^3 - x(a+\operatorname{barctanh}(cx))^2 - 2bc \int \frac{x(a+\operatorname{barctanh}(cx))}{1-c^2x^2} dx}{c^2} \right)$$

↓ 6510

$$\frac{3}{2}bc \left(\frac{(a+\operatorname{barctanh}(cx))^3}{3bc^3} - \frac{\frac{1}{2}x^2(a+\operatorname{barctanh}(cx))^3 - x(a+\operatorname{barctanh}(cx))^2 - 2bc \int \frac{x(a+\operatorname{barctanh}(cx))}{1-c^2x^2} dx}{c^2} \right)$$

↓ 6546

$$\frac{3}{2}bc \left(\frac{(a+\operatorname{barctanh}(cx))^3}{3bc^3} - \frac{x(a+\operatorname{barctanh}(cx))^2 - 2bc \left(\frac{\int \frac{a+\operatorname{barctanh}(cx)}{1-cx} dx}{c} - \frac{(a+\operatorname{barctanh}(cx))^2}{2bc^2} \right)}{c^2} \right)$$

↓ 6470

$$\frac{3}{2}bc \left(\frac{(a+\operatorname{barctanh}(cx))^3}{3bc^3} - \frac{x(a+\operatorname{barctanh}(cx))^2 - 2bc \left(\frac{\log\left(\frac{2}{1-cx}\right)(a+\operatorname{barctanh}(cx))}{c} - b \int \frac{\log\left(\frac{2}{1-cx}\right)}{1-c^2x^2} dx \right) - \frac{(a+\operatorname{barctanh}(cx))^2}{2bc^2}}{c^2} \right)$$

↓ 2849

$$\frac{3}{2}bc \left(\frac{(a+\operatorname{barctanh}(cx))^3}{3bc^3} - \frac{x(a+\operatorname{barctanh}(cx))^2 - 2bc \left(\frac{b \int \frac{\log\left(\frac{2}{1-cx}\right)}{1-\frac{2}{1-cx}} d\frac{1}{1-cx}}{c} + \frac{\log\left(\frac{2}{1-cx}\right)(a+\operatorname{barctanh}(cx))}{c} \right) - \frac{(a+\operatorname{barctanh}(cx))^2}{2bc^2}}{c^2} \right)$$

↓ 2752

$$\frac{3}{2}bc \left(\frac{(a + \operatorname{arctanh}(cx))^3}{3bc^3} - \frac{x(a + \operatorname{arctanh}(cx))^2 - 2bc \left(\frac{\log\left(\frac{2}{1-cx}\right)(a+b\operatorname{arctanh}(cx))}{c} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{2c} \right)}{c^2} - \frac{(a+b\operatorname{arctanh}(cx))^3}{2bc^3} \right)$$

input `Int [x*(a + b*ArcTanh [c*x])^3,x]`

output `(x^2*(a + b*ArcTanh [c*x])^3)/2 - (3*b*c*((a + b*ArcTanh [c*x])^3/(3*b*c^3) - (x*(a + b*ArcTanh [c*x])^2 - 2*b*c*(-1/2*(a + b*ArcTanh [c*x])^2/(b*c^2) + ((a + b*ArcTanh [c*x])*Log [2/(1 - c*x)]/c + (b*PolyLog [2, 1 - 2/(1 - c*x)])/(2*c))/c))/c^2)/2`

Defintions of rubi rules used

rule 2752 `Int [Log [(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp [(-e^(-1))*PolyLog [2, 1 - c*x], x] /; FreeQ [{c, d, e}, x] && EqQ [e + c*d, 0]`

rule 2849 `Int [Log [(c_.)/(d_) + (e_.)*(x_)]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp [-e/g Subst [Int [Log [2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ [{c, d, e, f, g}, x] && EqQ [c, 2*d] && EqQ [e^2*f + d^2*g, 0]`

rule 6436 `Int [((a_.) + ArcTanh [(c_.)*(x_)^(n_.)]*(b_.))^ (p_.), x_Symbol] := Simp [x*(a + b*ArcTanh [c*x^n])^p, x] - Simp [b*c*n*p Int [x^n*((a + b*ArcTanh [c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ [{a, b, c, n}, x] && IGtQ [p, 0] && (EqQ [n, 1] || EqQ [p, 1])`

rule 6452 `Int [((a_.) + ArcTanh [(c_.)*(x_)^(n_.)]*(b_.))^ (p_.)*(x_)^(m_.), x_Symbol] := Simp [x^(m + 1)*((a + b*ArcTanh [c*x^n])^p/(m + 1)), x] - Simp [b*c*n*(p/(m + 1)) Int [x^(m + n)*((a + b*ArcTanh [c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ [{a, b, c, m, n}, x] && IGtQ [p, 0] && (EqQ [p, 1] || (EqQ [n, 1] && IntegerQ [m])) && NeQ [m, -1]`

rule 6470

```
Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol
] := Simp[(-a + b*ArcTanh[c*x])^p*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c
*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^
2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2
, 0]
```

rule 6510

```
Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symb
ol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b
, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]
```

rule 6542

```
Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_))/((d_) + (
e_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*
x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/
(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m,
1]
```

rule 6546

```
Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*(x_)/((d_) + (e_.)*(x_)^2),
x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/
(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 487 vs. $2(113) = 226$.

Time = 6.59 (sec) , antiderivative size = 488, normalized size of antiderivative = 3.97

method	result
risch	$\frac{a^3 x^2}{2} + \left(\frac{3(c^2 x^2 - 1)b^3 \ln(-cx + 1)^2}{16c^2} - \frac{3b^2 x(cx + b) \ln(-cx + 1)}{4c} - \frac{3b(-a^2 c^2 x^2 - 2abcx - b \ln(-cx + 1)a - b^2 \ln(-cx + 1)^2)}{4c^2} \right)$
derivativdivides	Expression too large to display
default	Expression too large to display
parts	Expression too large to display

input

```
int(x*(a+b*arctanh(c*x))^3,x,method=_RETURNVERBOSE)
```

output

```

1/2*a^3*x^2+(3/16*(c^2*x^2-1)*b^3/c^2*ln(-c*x+1)^2-3/4*b^2*x*(a*c*x+b)/c*ln(-c*x+1)-3/4*b*(-a^2*c^2*x^2-2*a*b*c*x-b*ln(-c*x+1)*a-b^2*ln(-c*x+1))/c^2)*ln(c*x+1)-1/2/c^2*a^3+3/2*b^3/c^2*dilog(-1/2*c*x+1/2)-3/2/c^2*a^2*b+1/16/c^2*b^3*ln(-c*x+1)^3-3/8/c^2*b^3*ln(-c*x+1)^2+3/2/c*a^2*b*x-3/8/c^2*a*b^2*ln(-c*x+1)^2+3/2/c^2*a*b^2*ln(-c*x+1)+3/4/c^2*a^2*b*ln(-c*x+1)+3/8/c*b^3*ln(-c*x+1)^2*x+3/8*ln(-c*x+1)^2*a*b^2*x^2-3/4*ln(-c*x+1)*a^2*b*x^2-3/4*a^2*b/c^2*ln(-c*x-1)+3/2*a*b^2/c^2*ln(-c*x-1)-3/2/c*a*b^2*ln(-c*x+1)*x-1/16*ln(-c*x+1)^3*b^3*x^2+3/2*b^3/c^2*ln(-1/2*c*x+1/2)*ln(1/2*c*x+1/2)-3/2*b^3/c^2*ln(1/2*c*x+1/2)*ln(-c*x+1)+1/16*(c^2*x^2-1)*b^3/c^2*ln(c*x+1)^3+3/16*b^2*(-x^2*b*ln(-c*x+1)*c^2+2*a*c^2*x^2+2*b*c*x+b*ln(-c*x+1)-2*a+2*b)/c^2*ln(c*x+1)^2

```

Fricas [F]

$$\int x(a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{artanh}(cx) + a)^3 x dx$$

input

```
integrate(x*(a+b*arctanh(c*x))^3,x, algorithm="fricas")
```

output

```
integral(b^3*x*arctanh(c*x)^3 + 3*a*b^2*x*arctanh(c*x)^2 + 3*a^2*b*x*arctanh(c*x) + a^3*x, x)
```

Sympy [F]

$$\int x(a + b \operatorname{arctanh}(cx))^3 dx = \int x(a + b \operatorname{atanh}(cx))^3 dx$$

input

```
integrate(x*(a+b*atanh(c*x))**3,x)
```

output

```
Integral(x*(a + b*atanh(c*x))**3, x)
```

Maxima [F]

$$\int x(a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{arctanh}(cx) + a)^3 x dx$$

input `integrate(x*(a+b*arctanh(c*x))^3,x, algorithm="maxima")`

output `3/2*a*b^2*x^2*arctanh(c*x)^2 + 1/2*a^3*x^2 + 3/4*(2*x^2*arctanh(c*x) + c*(2*x/c^2 - log(c*x + 1)/c^3 + log(c*x - 1)/c^3))*a^2*b + 3/8*(4*c*(2*x/c^2 - log(c*x + 1)/c^3 + log(c*x - 1)/c^3)*arctanh(c*x) - (2*(log(c*x - 1) - 2)*log(c*x + 1) - log(c*x + 1)^2 - log(c*x - 1)^2 - 4*log(c*x - 1))/c^2)*a*b^2 - 1/64*(3*c^3*(x^2/c^3 + log(c^2*x^2 - 1)/c^5) + 21*c^2*(2*x/c^3 - log(c*x + 1)/c^4 + log(c*x - 1)/c^4) - 576*c*integrate(1/4*x*log(c*x + 1)/(c^3*x^2 - c), x) - 2*(12*c*x*log(c*x + 1)^2 + 2*(c^2*x^2 - 1)*log(c*x + 1)^3 - 3*(c^2*x^2 - 2*c*x - 2*(c^2*x^2 - 1)*log(c*x + 1) + 1)*log(-c*x + 1)^2 + 3*(c^2*x^2 - 2*(c^2*x^2 - 1)*log(c*x + 1)^2 + 6*c*x - 8*(c*x + 1)*log(c*x + 1))*log(-c*x + 1)/c^2 + ((4*log(-c*x + 1)^3 - 6*log(-c*x + 1)^2 + 6*log(-c*x + 1) - 3)*(c*x - 1)^2 + 8*(log(-c*x + 1)^3 - 3*log(-c*x + 1)^2 + 6*log(-c*x + 1) - 6)*(c*x - 1))/c^2 + 18*log(4*c^3*x^2 - 4*c)/c^2 - 192*integrate(1/4*log(c*x + 1)/(c^3*x^2 - c), x))*b^3`

Giac [F]

$$\int x(a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{arctanh}(cx) + a)^3 x dx$$

input `integrate(x*(a+b*arctanh(c*x))^3,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^3*x, x)`

3.29 $\int (a + b \operatorname{arctanh}(cx))^3 dx$

Optimal result	321
Mathematica [A] (verified)	322
Rubi [A] (verified)	322
Maple [B] (verified)	324
Fricas [F]	325
Sympy [F]	325
Maxima [F]	326
Giac [F]	326
Mupad [F(-1)]	326
Reduce [F]	327

Optimal result

Integrand size = 10, antiderivative size = 108

$$\int (a + b \operatorname{arctanh}(cx))^3 dx = \frac{(a + b \operatorname{arctanh}(cx))^3}{c} + x(a + b \operatorname{arctanh}(cx))^3 - \frac{3b(a + b \operatorname{arctanh}(cx))^2 \log\left(\frac{2}{1-cx}\right)}{c} - \frac{3b^2(a + b \operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{c} + \frac{3b^3 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx}\right)}{2c}$$

output

```
(a+b*arctanh(c*x))^3/c+x*(a+b*arctanh(c*x))^3-3*b*(a+b*arctanh(c*x))^2*ln(
2/(-c*x+1))/c-3*b^2*(a+b*arctanh(c*x))*polylog(2,1-2/(-c*x+1))/c+3/2*b^3*p
olylog(3,1-2/(-c*x+1))/c
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.49

$$\int (a + b \operatorname{arctanh}(cx))^3 dx$$

$$= \frac{2a^3 cx + 6a^2 bcx \operatorname{arctanh}(cx) + 3a^2 b \log(1 - c^2 x^2) + 6ab^2 (\operatorname{arctanh}(cx) ((-1 + cx) \operatorname{arctanh}(cx) - 2 \log(1 - c^2 x^2)))}{2c}$$

input `Integrate[(a + b*ArcTanh[c*x])^3,x]`

output $(2a^3 cx + 6a^2 bcx \operatorname{ArcTanh}[c*x] + 3a^2 b \operatorname{Log}[1 - c^2 x^2] + 6a^2 b^2 (\operatorname{ArcTanh}[c*x] * ((-1 + c*x) \operatorname{ArcTanh}[c*x] - 2 \operatorname{Log}[1 + E^{(-2 \operatorname{ArcTanh}[c*x])}]])) + \operatorname{PolyLog}[2, -E^{(-2 \operatorname{ArcTanh}[c*x])}] + b^3 (2 \operatorname{ArcTanh}[c*x]^2 * ((-1 + c*x) \operatorname{ArcTanh}[c*x] - 3 \operatorname{Log}[1 + E^{(-2 \operatorname{ArcTanh}[c*x])}])) + 6 \operatorname{ArcTanh}[c*x] * \operatorname{PolyLog}[2, -E^{(-2 \operatorname{ArcTanh}[c*x])}] + 3 \operatorname{PolyLog}[3, -E^{(-2 \operatorname{ArcTanh}[c*x])}])) / (2*c)$

Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.14, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6436, 6546, 6470, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{arctanh}(cx))^3 dx$$

$$\downarrow 6436$$

$$x(a + b \operatorname{arctanh}(cx))^3 - 3bc \int \frac{x(a + b \operatorname{arctanh}(cx))^2}{1 - c^2 x^2} dx$$

$$\downarrow 6546$$

$$x(a + b \operatorname{arctanh}(cx))^3 - 3bc \left(\frac{\int \frac{(a + b \operatorname{arctanh}(cx))^2}{1 - cx} dx}{c} - \frac{(a + b \operatorname{arctanh}(cx))^3}{3bc^2} \right)$$

$$\downarrow 6470$$

$$\begin{aligned}
& 3bc \left(\frac{\log\left(\frac{2}{1-cx}\right)(a+\operatorname{barctanh}(cx))^2}{c} - \frac{2b \int \frac{(a+\operatorname{barctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{1-c^2x^2} dx}{c} - \frac{(a+\operatorname{barctanh}(cx))^3}{3bc^2} \right) \\
& \quad \downarrow 6620 \\
& 3bc \left(\frac{\log\left(\frac{2}{1-cx}\right)(a+\operatorname{barctanh}(cx))^2}{c} - \frac{2b \left(\frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, 1-\frac{2}{1-cx}\right)}{1-c^2x^2} dx - \frac{\operatorname{PolyLog}\left(2, 1-\frac{2}{1-cx}\right)(a+\operatorname{barctanh}(cx))}{2c} \right)}{c} \right) - \frac{(a+\operatorname{barctanh}(cx))^3}{3bc^2} \\
& \quad \downarrow 7164 \\
& 3bc \left(\frac{\log\left(\frac{2}{1-cx}\right)(a+\operatorname{barctanh}(cx))^2}{c} - \frac{2b \left(\frac{b \operatorname{PolyLog}\left(3, 1-\frac{2}{1-cx}\right)}{4c} - \frac{\operatorname{PolyLog}\left(2, 1-\frac{2}{1-cx}\right)(a+\operatorname{barctanh}(cx))}{2c} \right)}{c} \right) - \frac{(a+\operatorname{barctanh}(cx))^3}{3bc^2}
\end{aligned}$$

input `Int[(a + b*ArcTanh[c*x])^3,x]`

output `x*(a + b*ArcTanh[c*x])^3 - 3*b*c*(-1/3*(a + b*ArcTanh[c*x])^3/(b*c^2) + ((a + b*ArcTanh[c*x])^2*Log[2/(1 - c*x)]/c - 2*b*(-1/2*((a + b*ArcTanh[c*x])*PolyLog[2, 1 - 2/(1 - c*x)]/c + (b*PolyLog[3, 1 - 2/(1 - c*x)]/(4*c)))/c)`

Defintions of rubi rules used

rule 6436

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^p, x_Symbol] := Simp[x*(a + b*ArcTanh[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])
```



```
rule 6470 Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol]
  := Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c
  *(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^
  2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]
```

```
rule 6546 Int[(((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*(x_))/((d_) + (e_.)*(x_)^2),
  x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/
  (c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d,
  e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

```
rule 6620 Int[(Log[u_]*((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.))/((d_) + (e_.)*(x_)^
  2), x_Symbol] := Simp[(-(a + b*ArcTanh[c*x])^p)*(PolyLog[2, 1 - u]/(2*c*d))
  , x] + Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(
  d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d
  + e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]
```

```
rule 7164 Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
  x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 229 vs. 2(106) = 212.

Time = 1.19 (sec) , antiderivative size = 230, normalized size of antiderivative = 2.13

method	result
derivativedivides	$a^3cx + b^3 \left(\operatorname{arctanh}(cx)^3(cx-1) + 2 \operatorname{arctanh}(cx)^3 - 3 \operatorname{arctanh}(cx)^2 \ln \left(1 + \frac{(cx+1)^2}{-c^2x^2+1} \right) - 3 \operatorname{arctanh}(cx) \operatorname{polylog} \left(2, -\frac{cx+1}{-c^2x} \right) \right)$
default	$a^3cx + b^3 \left(\operatorname{arctanh}(cx)^3(cx-1) + 2 \operatorname{arctanh}(cx)^3 - 3 \operatorname{arctanh}(cx)^2 \ln \left(1 + \frac{(cx+1)^2}{-c^2x^2+1} \right) - 3 \operatorname{arctanh}(cx) \operatorname{polylog} \left(2, -\frac{cx+1}{-c^2x} \right) \right)$
parts	$x a^3 + \frac{b^3 \left(\operatorname{arctanh}(cx)^3(cx-1) + 2 \operatorname{arctanh}(cx)^3 - 3 \operatorname{arctanh}(cx)^2 \ln \left(1 + \frac{(cx+1)^2}{-c^2x^2+1} \right) - 3 \operatorname{arctanh}(cx) \operatorname{polylog} \left(2, -\frac{cx+1}{-c^2x} \right) \right)}{c}$

input `int((a+b*arctanh(c*x))^3,x,method=_RETURNVERBOSE)`

output `1/c*(a^3*c*x+b^3*(arctanh(c*x)^3*(c*x-1)+2*arctanh(c*x)^3-3*arctanh(c*x)^2*ln(1+(c*x+1)^2/(-c^2*x^2+1))-3*arctanh(c*x)*polylog(2,-(c*x+1)^2/(-c^2*x^2+1))+3/2*polylog(3,-(c*x+1)^2/(-c^2*x^2+1)))+3*a*b^2*(arctanh(c*x)^2*(c*x-1)+2*arctanh(c*x)^2-2*arctanh(c*x)*ln(1+(c*x+1)^2/(-c^2*x^2+1))-polylog(2,-(c*x+1)^2/(-c^2*x^2+1)))+3*a^2*b*(c*x*arctanh(c*x)+1/2*ln(-c^2*x^2+1))`

Fricas [F]

$$\int (a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{artanh}(cx) + a)^3 dx$$

input `integrate((a+b*arctanh(c*x))^3,x, algorithm="fricas")`

output `integral(b^3*arctanh(c*x)^3 + 3*a*b^2*arctanh(c*x)^2 + 3*a^2*b*arctanh(c*x) + a^3, x)`

Sympy [F]

$$\int (a + b \operatorname{arctanh}(cx))^3 dx = \int (a + b \operatorname{atanh}(cx))^3 dx$$

input `integrate((a+b*atanh(c*x))**3,x)`

output `Integral((a + b*atanh(c*x))**3, x)`

Maxima [F]

$$\int (a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{artanh}(cx) + a)^3 dx$$

input `integrate((a+b*arctanh(c*x))^3,x, algorithm="maxima")`

output `a^3*x + 3/2*(2*c*x*arctanh(c*x) + log(-c^2*x^2 + 1))*a^2*b/c - 1/8*((b^3*c*x - b^3)*log(-c*x + 1)^3 - 3*(2*a*b^2*c*x + (b^3*c*x + b^3)*log(c*x + 1))*log(-c*x + 1)^2)/c - integrate(-1/8*((b^3*c*x - b^3)*log(c*x + 1)^3 + 6*(a*b^2*c*x - a*b^2)*log(c*x + 1)^2 - 3*(4*a*b^2*c*x + (b^3*c*x - b^3)*log(c*x + 1)^2 - 2*(2*a*b^2 - b^3 - (2*a*b^2*c + b^3*c)*x)*log(c*x + 1))*log(-c*x + 1))/(c*x - 1), x)`

Giac [F]

$$\int (a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{artanh}(cx) + a)^3 dx$$

input `integrate((a+b*arctanh(c*x))^3,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + b \operatorname{arctanh}(cx))^3 dx = \int (a + b \operatorname{atanh}(cx))^3 dx$$

input `int((a + b*atanh(c*x))^3,x)`

output `int((a + b*atanh(c*x))^3, x)`

Reduce [F]

$$\int (a + b \operatorname{arctanh}(cx))^3 dx$$

$$= \frac{3 \operatorname{atanh}(cx) a^2 b c x + 3 \operatorname{atanh}(cx) a^2 b + \left(\int \operatorname{atanh}(cx)^3 dx \right) b^3 c + 3 \left(\int \operatorname{atanh}(cx)^2 dx \right) a b^2 c + 3 \log(c^2 x - c)}{c}$$

input `int((a+b*atanh(c*x))^3,x)`

output `(3*atanh(c*x)*a**2*b*c*x + 3*atanh(c*x)*a**2*b + int(atanh(c*x)**3,x)*b**3*c + 3*int(atanh(c*x)**2,x)*a*b**2*c + 3*log(c**2*x - c)*a**2*b + a**3*c*x)/c`

3.30 $\int \frac{(a+b\operatorname{arctanh}(cx))^3}{x} dx$

Optimal result	328
Mathematica [C] (verified)	329
Rubi [A] (verified)	330
Maple [C] (warning: unable to verify)	332
Fricas [F]	333
Sympy [F]	334
Maxima [F]	334
Giac [F]	334
Mupad [F(-1)]	335
Reduce [F]	335

Optimal result

Integrand size = 14, antiderivative size = 184

$$\int \frac{(a + b\operatorname{arctanh}(cx))^3}{x} dx = 2(a + b\operatorname{arctanh}(cx))^3 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx}\right) - \frac{3}{2}b(a + b\operatorname{arctanh}(cx))^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx}\right) + \frac{3}{2}b(a + b\operatorname{arctanh}(cx))^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - cx}\right) + \frac{3}{2}b^2(a + b\operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - cx}\right) - \frac{3}{2}b^2(a + b\operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 - cx}\right) - \frac{3}{4}b^3 \operatorname{PolyLog}\left(4, 1 - \frac{2}{1 - cx}\right) + \frac{3}{4}b^3 \operatorname{PolyLog}\left(4, -1 + \frac{2}{1 - cx}\right)$$

output

```
-2*(a+b*arctanh(c*x))^3*arctanh(-1+2/(-c*x+1))-3/2*b*(a+b*arctanh(c*x))^2*
polylog(2,1-2/(-c*x+1))+3/2*b*(a+b*arctanh(c*x))^2*polylog(2,-1+2/(-c*x+1))
)+3/2*b^2*(a+b*arctanh(c*x))*polylog(3,1-2/(-c*x+1))-3/2*b^2*(a+b*arctanh(
c*x))*polylog(3,-1+2/(-c*x+1))-3/4*b^3*polylog(4,1-2/(-c*x+1))+3/4*b^3*pol
ylog(4,-1+2/(-c*x+1))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.47 (sec) , antiderivative size = 315, normalized size of antiderivative = 1.71

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x} dx = a^3 \log(cx) + \frac{3}{2} a^2 b (-\operatorname{PolyLog}(2, -cx) + \operatorname{PolyLog}(2, cx))$$

$$+ 3ab^2 \left(\frac{i\pi^3}{24} - \frac{2}{3} \operatorname{arctanh}(cx)^3 \right.$$

$$\quad - \operatorname{arctanh}(cx)^2 \log(1 + e^{-2\operatorname{arctanh}(cx)})$$

$$\quad + \operatorname{arctanh}(cx)^2 \log(1 - e^{2\operatorname{arctanh}(cx)})$$

$$+ \operatorname{arctanh}(cx) \operatorname{PolyLog}(2, -e^{-2\operatorname{arctanh}(cx)})$$

$$+ \operatorname{arctanh}(cx) \operatorname{PolyLog}(2, e^{2\operatorname{arctanh}(cx)})$$

$$\quad + \frac{1}{2} \operatorname{PolyLog}(3, -e^{-2\operatorname{arctanh}(cx)})$$

$$\quad \left. - \frac{1}{2} \operatorname{PolyLog}(3, e^{2\operatorname{arctanh}(cx)}) \right) + \frac{1}{64} b^3 (\pi^4$$

$$- 32 \operatorname{arctanh}(cx)^4 - 64 \operatorname{arctanh}(cx)^3 \log(1 + e^{-2\operatorname{arctanh}(cx)})$$

$$+ 64 \operatorname{arctanh}(cx)^3 \log(1 - e^{2\operatorname{arctanh}(cx)})$$

$$+ 96 \operatorname{arctanh}(cx)^2 \operatorname{PolyLog}(2, -e^{-2\operatorname{arctanh}(cx)})$$

$$+ 96 \operatorname{arctanh}(cx)^2 \operatorname{PolyLog}(2, e^{2\operatorname{arctanh}(cx)})$$

$$+ 96 \operatorname{arctanh}(cx) \operatorname{PolyLog}(3, -e^{-2\operatorname{arctanh}(cx)})$$

$$- 96 \operatorname{arctanh}(cx) \operatorname{PolyLog}(3, e^{2\operatorname{arctanh}(cx)})$$

$$+ 48 \operatorname{PolyLog}(4, -e^{-2\operatorname{arctanh}(cx)})$$

$$+ 48 \operatorname{PolyLog}(4, e^{2\operatorname{arctanh}(cx)})$$

input `Integrate[(a + b*ArcTanh[c*x])^3/x,x]`

output

```

a^3*Log[c*x] + (3*a^2*b*(-PolyLog[2, -(c*x)] + PolyLog[2, c*x])/2 + 3*a*b
^2*((I/24)*Pi^3 - (2*ArcTanh[c*x]^3)/3 - ArcTanh[c*x]^2*Log[1 + E^(-2*ArcT
anh[c*x])) + ArcTanh[c*x]^2*Log[1 - E^(2*ArcTanh[c*x])] + ArcTanh[c*x]*Pol
yLog[2, -E^(-2*ArcTanh[c*x])] + ArcTanh[c*x]*PolyLog[2, E^(2*ArcTanh[c*x])
] + PolyLog[3, -E^(-2*ArcTanh[c*x])]/2 - PolyLog[3, E^(2*ArcTanh[c*x])]/2)
+ (b^3*(Pi^4 - 32*ArcTanh[c*x]^4 - 64*ArcTanh[c*x]^3*Log[1 + E^(-2*ArcTan
h[c*x])] + 64*ArcTanh[c*x]^3*Log[1 - E^(2*ArcTanh[c*x])] + 96*ArcTanh[c*x]
^2*PolyLog[2, -E^(-2*ArcTanh[c*x])] + 96*ArcTanh[c*x]^2*PolyLog[2, E^(2*Ar
cTanh[c*x])] + 96*ArcTanh[c*x]*PolyLog[3, -E^(-2*ArcTanh[c*x])] - 96*ArcTa
nh[c*x]*PolyLog[3, E^(2*ArcTanh[c*x])] + 48*PolyLog[4, -E^(-2*ArcTanh[c*x]
)] + 48*PolyLog[4, E^(2*ArcTanh[c*x])]))/64

```

Rubi [A] (verified)

Time = 1.77 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.15, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {6448, 6614, 6620, 6624, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{(a + b \operatorname{arctanh}(cx))^3}{x} dx \\
& \quad \downarrow \text{6448} \\
& 2 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx}\right) (a + b \operatorname{arctanh}(cx))^3 - 6bc \int \frac{(a + b \operatorname{arctanh}(cx))^2 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx}\right)}{1 - c^2 x^2} dx \\
& \quad \downarrow \text{6614} \\
& 2 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx}\right) (a + b \operatorname{arctanh}(cx))^3 - \\
& 6bc \left(\frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(cx))^2 \log\left(2 - \frac{2}{1 - cx}\right)}{1 - c^2 x^2} dx - \frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(cx))^2 \log\left(\frac{2}{1 - cx}\right)}{1 - c^2 x^2} dx \right) \\
& \quad \downarrow \text{6620}
\end{aligned}$$

$$\begin{aligned}
& 2\operatorname{arctanh}\left(1 - \frac{2}{1-cx}\right) (a + b\operatorname{arctanh}(cx))^3 - \\
6bc & \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right) (a + b\operatorname{arctanh}(cx))^2}{2c} - b \int \frac{(a + b\operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{1 - c^2x^2} dx \right) + \frac{1}{2} \right. \\
& \quad \downarrow 6624 \\
& 2\operatorname{arctanh}\left(1 - \frac{2}{1-cx}\right) (a + b\operatorname{arctanh}(cx))^3 - \\
6bc & \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right) (a + b\operatorname{arctanh}(cx))^2}{2c} - b \left(\frac{\operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx}\right) (a + b\operatorname{arctanh}(cx))}{2c} - \frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx}\right) (a + b\operatorname{arctanh}(cx))}{1 - c^2x^2} dx \right) \right. \\
& \quad \downarrow 7164 \\
& 2\operatorname{arctanh}\left(1 - \frac{2}{1-cx}\right) (a + b\operatorname{arctanh}(cx))^3 - \\
6bc & \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right) (a + b\operatorname{arctanh}(cx))^2}{2c} - b \left(\frac{\operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx}\right) (a + b\operatorname{arctanh}(cx))}{2c} - \frac{b \operatorname{PolyLog}\left(4, 1 - \frac{2}{1-cx}\right) (a + b\operatorname{arctanh}(cx))}{4c} \right) \right)
\end{aligned}$$

input `Int[(a + b*ArcTanh[c*x])^3/x, x]`

output `2*(a + b*ArcTanh[c*x])^3*ArcTanh[1 - 2/(1 - c*x)] - 6*b*c*(((a + b*ArcTanh[c*x])^2*PolyLog[2, 1 - 2/(1 - c*x)]/(2*c) - b*(((a + b*ArcTanh[c*x])*PolyLog[3, 1 - 2/(1 - c*x)]/(2*c) - (b*PolyLog[4, 1 - 2/(1 - c*x)]/(4*c)))/2 + (-1/2*((a + b*ArcTanh[c*x])^2*PolyLog[2, -1 + 2/(1 - c*x)]/c + b*(((a + b*ArcTanh[c*x])*PolyLog[3, -1 + 2/(1 - c*x)]/(2*c) - (b*PolyLog[4, -1 + 2/(1 - c*x)]/(4*c)))/2)`

Defintions of rubi rules used

rule 6448

```
Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^p/(x_), x_Symbol] := Simp[2*(a + b*ArcTanh[c*x])^p*ArcTanh[1 - 2/(1 - c*x)], x] - Simp[2*b*c*p Int[(a + b*ArcTanh[c*x])^(p - 1)*(ArcTanh[1 - 2/(1 - c*x)]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1]
```


rule 6614

```
Int[(ArcTanh[u_]*((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/2 Int[Log[1 + u]*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] - Simp[1/2 Int[Log[1 - u]*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[u^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 6620

```
Int[(Log[u_]*((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(-a + b*ArcTanh[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)), x] + Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 6624

```
Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*PolyLog[k_, u_])/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(PolyLog[k + 1, u]/(2*c*d)), x] - Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[k + 1, u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, k}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[u^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v, x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 6.40 (sec) , antiderivative size = 1304, normalized size of antiderivative = 7.09

method	result	size
parts	Expression too large to display	1304
derivativedivides	Expression too large to display	1306
default	Expression too large to display	1306

input

```
int((a+b*arctanh(c*x))^3/x,x,method=_RETURNVERBOSE)
```

output

```

a^3*ln(x)+b^3*(ln(c*x)*arctanh(c*x)^3-arctanh(c*x)^3*ln((c*x+1)^2/(-c^2*x^
2+1)-1)+arctanh(c*x)^3*ln(1+(c*x+1)/(-c^2*x^2+1)^(1/2))+3*arctanh(c*x)^2*p
olylog(2,-(c*x+1)/(-c^2*x^2+1)^(1/2))-6*arctanh(c*x)*polylog(3,-(c*x+1)/(-
c^2*x^2+1)^(1/2))+6*polylog(4,-(c*x+1)/(-c^2*x^2+1)^(1/2))+arctanh(c*x)^3*
ln(1-(c*x+1)/(-c^2*x^2+1)^(1/2))+3*arctanh(c*x)^2*polylog(2,(c*x+1)/(-c^2*
x^2+1)^(1/2))-6*arctanh(c*x)*polylog(3,(c*x+1)/(-c^2*x^2+1)^(1/2))+6*polyl
og(4,(c*x+1)/(-c^2*x^2+1)^(1/2))+1/2*I*Pi*csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1
))/(1-(c*x+1)^2/(c^2*x^2-1)))*(csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1))*csgn(I/(1
-(c*x+1)^2/(c^2*x^2-1)))-csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1))*csgn(I*(-(c*x+
1)^2/(c^2*x^2-1)-1)/(1-(c*x+1)^2/(c^2*x^2-1)))-csgn(I*(-(c*x+1)^2/(c^2*x^
2-1)-1)/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))+csgn(I
*(-(c*x+1)^2/(c^2*x^2-1)-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^2)*arctanh(c*x)^3-3
/2*arctanh(c*x)^2*polylog(2,-(c*x+1)^2/(-c^2*x^2+1))+3/2*arctanh(c*x)*poly
log(3,-(c*x+1)^2/(-c^2*x^2+1))-3/4*polylog(4,-(c*x+1)^2/(-c^2*x^2+1))+3*a
*b^2*(ln(c*x)*arctanh(c*x)^2-arctanh(c*x)*polylog(2,-(c*x+1)^2/(-c^2*x^2+1
))+1/2*polylog(3,-(c*x+1)^2/(-c^2*x^2+1))-arctanh(c*x)^2*ln((c*x+1)^2/(-c^
2*x^2+1)-1)+arctanh(c*x)^2*ln(1+(c*x+1)/(-c^2*x^2+1)^(1/2))+2*arctanh(c*x)
*polylog(2,-(c*x+1)/(-c^2*x^2+1)^(1/2))-2*polylog(3,-(c*x+1)/(-c^2*x^2+1)^
(1/2))+arctanh(c*x)^2*ln(1-(c*x+1)/(-c^2*x^2+1)^(1/2))+2*arctanh(c*x)*poly
log(2,(c*x+1)/(-c^2*x^2+1)^(1/2))-2*polylog(3,(c*x+1)/(-c^2*x^2+1)^(1/2)...

```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x} dx = \int \frac{(b \operatorname{arctanh}(cx) + a)^3}{x} dx$$

input

```
integrate((a+b*arctanh(c*x))^3/x,x, algorithm="fricas")
```

output

```
integral((b^3*arctanh(c*x)^3 + 3*a*b^2*arctanh(c*x)^2 + 3*a^2*b*arctanh(c*
x) + a^3)/x, x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x} dx = \int \frac{(a + b \operatorname{atanh}(cx))^3}{x} dx$$

input `integrate((a+b*atanh(c*x))**3/x,x)`

output `Integral((a + b*atanh(c*x))**3/x, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{x} dx$$

input `integrate((a+b*arctanh(c*x))^3/x,x, algorithm="maxima")`

output `a^3*log(x) + integrate(1/8*b^3*(log(c*x + 1) - log(-c*x + 1))^3/x + 3/4*a*b^2*(log(c*x + 1) - log(-c*x + 1))^2/x + 3/2*a^2*b*(log(c*x + 1) - log(-c*x + 1))/x, x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{x} dx$$

input `integrate((a+b*arctanh(c*x))^3/x,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^3/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x} dx = \int \frac{(a + b \operatorname{atanh}(cx))^3}{x} dx$$

input `int((a + b*atanh(c*x))^3/x,x)`output `int((a + b*atanh(c*x))^3/x, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x} dx = 3 \left(\int \frac{\operatorname{atanh}(cx)}{x} dx \right) a^2 b + \left(\int \frac{\operatorname{atanh}(cx)^3}{x} dx \right) b^3$$

$$+ 3 \left(\int \frac{\operatorname{atanh}(cx)^2}{x} dx \right) a b^2 + \log(x) a^3$$

input `int((a+b*atanh(c*x))^3/x,x)`output `3*int(atanh(c*x)/x,x)*a**2*b + int(atanh(c*x)**3/x,x)*b**3 + 3*int(atanh(c*x)**2/x,x)*a*b**2 + log(x)*a**3`

3.31 $\int \frac{(a+b\operatorname{arctanh}(cx))^3}{x^2} dx$

Optimal result	336
Mathematica [C] (verified)	337
Rubi [A] (verified)	337
Maple [C] (warning: unable to verify)	340
Fricas [F]	341
Sympy [F]	341
Maxima [F]	341
Giac [F]	342
Mupad [F(-1)]	342
Reduce [F]	342

Optimal result

Integrand size = 14, antiderivative size = 102

$$\int \frac{(a + b\operatorname{arctanh}(cx))^3}{x^2} dx = c(a + b\operatorname{arctanh}(cx))^3 - \frac{(a + b\operatorname{arctanh}(cx))^3}{x} + 3bc(a + b\operatorname{arctanh}(cx))^2 \log\left(2 - \frac{2}{1 + cx}\right) - 3b^2c(a + b\operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + cx}\right) - \frac{3}{2}b^3c \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 + cx}\right)$$

output

```
c*(a+b*arctanh(c*x))^3-(a+b*arctanh(c*x))^3/x+3*b*c*(a+b*arctanh(c*x))^2*ln(2-2/(c*x+1))-3*b^2*c*(a+b*arctanh(c*x))*polylog(2,-1+2/(c*x+1))-3/2*b^3*c*polylog(3,-1+2/(c*x+1))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.24 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.92

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^2} dx = -\frac{a^3}{x} - \frac{3a^2 b \operatorname{arctanh}(cx)}{x} + 3a^2 b c \log(x) - \frac{3}{2} a^2 b c \log(1 - c^2 x^2) \\ + 3ab^2 c \left(\operatorname{arctanh}(cx) \left(\operatorname{arctanh}(cx) - \frac{\operatorname{arctanh}(cx)}{cx} \right) \right. \\ \left. + 2 \log(1 - e^{-2 \operatorname{arctanh}(cx)}) \right) - \operatorname{PolyLog}(2, e^{-2 \operatorname{arctanh}(cx)}) \\ + b^3 c \left(\frac{i\pi^3}{8} - \operatorname{arctanh}(cx)^3 - \frac{\operatorname{arctanh}(cx)^3}{cx} \right. \\ \left. + 3 \operatorname{arctanh}(cx)^2 \log(1 - e^{2 \operatorname{arctanh}(cx)}) \right. \\ \left. + 3 \operatorname{arctanh}(cx) \operatorname{PolyLog}(2, e^{2 \operatorname{arctanh}(cx)}) \right. \\ \left. - \frac{3}{2} \operatorname{PolyLog}(3, e^{2 \operatorname{arctanh}(cx)}) \right)$$

input

```
Integrate[(a + b*ArcTanh[c*x])^3/x^2,x]
```

output

```
-(a^3/x) - (3*a^2*b*ArcTanh[c*x])/x + 3*a^2*b*c*Log[x] - (3*a^2*b*c*Log[1
- c^2*x^2])/2 + 3*a*b^2*c*(ArcTanh[c*x]*(ArcTanh[c*x] - ArcTanh[c*x]/(c*x)
+ 2*Log[1 - E^(-2*ArcTanh[c*x])]) - PolyLog[2, E^(-2*ArcTanh[c*x])]) + b^
3*c*((I/8)*Pi^3 - ArcTanh[c*x]^3 - ArcTanh[c*x]^3/(c*x) + 3*ArcTanh[c*x]^2
*Log[1 - E^(2*ArcTanh[c*x])] + 3*ArcTanh[c*x]*PolyLog[2, E^(2*ArcTanh[c*x]
)]) - (3*PolyLog[3, E^(2*ArcTanh[c*x])])/2)
```

Rubi [A] (verified)

Time = 1.31 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.13, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {6452, 6550, 6494, 6618, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{(a + \operatorname{barctanh}(cx))^3}{x^2} dx \\
& \quad \downarrow \text{6452} \\
& 3bc \int \frac{(a + \operatorname{barctanh}(cx))^2}{x(1 - c^2x^2)} dx - \frac{(a + \operatorname{barctanh}(cx))^3}{x} \\
& \quad \downarrow \text{6550} \\
& 3bc \left(\int \frac{(a + \operatorname{barctanh}(cx))^2}{x(cx + 1)} dx + \frac{(a + \operatorname{barctanh}(cx))^3}{3b} \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{x} \\
& \quad \downarrow \text{6494} \\
& 3bc \left(-2bc \int \frac{(a + \operatorname{barctanh}(cx)) \log \left(2 - \frac{2}{cx+1} \right)}{1 - c^2x^2} dx + \frac{(a + \operatorname{barctanh}(cx))^3}{3b} + \log \left(2 - \frac{2}{cx+1} \right) (a + \operatorname{barctanh}(cx)) \right) \\
& \quad \quad \quad \frac{(a + \operatorname{barctanh}(cx))^3}{x} \\
& \quad \quad \quad \downarrow \text{6618} \\
& 3bc \left(-2bc \left(\frac{\operatorname{PolyLog} \left(2, \frac{2}{cx+1} - 1 \right) (a + \operatorname{barctanh}(cx))}{2c} - \frac{1}{2} b \int \frac{\operatorname{PolyLog} \left(2, \frac{2}{cx+1} - 1 \right)}{1 - c^2x^2} dx \right) + \frac{(a + \operatorname{barctanh}(cx))^3}{3b} \right) \\
& \quad \quad \quad \frac{(a + \operatorname{barctanh}(cx))^3}{x} \\
& \quad \quad \quad \downarrow \text{7164} \\
& 3bc \left(-2bc \left(\frac{\operatorname{PolyLog} \left(2, \frac{2}{cx+1} - 1 \right) (a + \operatorname{barctanh}(cx))}{2c} + \frac{b \operatorname{PolyLog} \left(3, \frac{2}{cx+1} - 1 \right)}{4c} \right) + \frac{(a + \operatorname{barctanh}(cx))^3}{3b} + \log \left(2 - \frac{2}{cx+1} \right) (a + \operatorname{barctanh}(cx)) \right) \\
& \quad \quad \quad \frac{(a + \operatorname{barctanh}(cx))^3}{x}
\end{aligned}$$

input

```
Int[(a + b*ArcTanh[c*x])^3/x^2,x]
```

output

```
-((a + b*ArcTanh[c*x])^3/x) + 3*b*c*((a + b*ArcTanh[c*x])^3/(3*b) + (a + b*ArcTanh[c*x])^2*Log[2 - 2/(1 + c*x)] - 2*b*c*(((a + b*ArcTanh[c*x])*PolyLog[2, -1 + 2/(1 + c*x)])/(2*c) + (b*PolyLog[3, -1 + 2/(1 + c*x)]/(4*c))))
```

Definitions of rubi rules used

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6494 `Int[((a_.) + ArcTanh[(c_.)*(x_)*(b_.)]^(p_.)/((x_)*((d_) + (e_.)*(x_))), x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6550 `Int[((a_.) + ArcTanh[(c_.)*(x_)*(b_.)]^(p_.)/((x_)*((d_) + (e_.)*(x_)^2)), x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]`

rule 6618 `Int[(Log[u_]*)((a_.) + ArcTanh[(c_.)*(x_)*(b_.)]^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)), x] - Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 + c*x))^2, 0]`

rule 7164 `Int[(u_)*PolyLog[n_, v_], x_Symbol] :> With[{w = DerivativeDivides[v, u*v, x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 9.42 (sec) , antiderivative size = 1329, normalized size of antiderivative = 13.03

method	result	size
parts	Expression too large to display	1329
derivativedivides	Expression too large to display	1331
default	Expression too large to display	1331

input `int((a+b*arctanh(c*x))^3/x^2,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & -1/x*a^3+b^3*c*(-1/c/x*arctanh(c*x)^3-3/2*arctanh(c*x)^2*\ln(c*x-1)+3*\ln(c*x) \\
 & *arctanh(c*x)^2-3/2*arctanh(c*x)^2*\ln(c*x+1)+3*arctanh(c*x)^2*\ln((c*x+1) \\
 & /(-c^2*x^2+1)^{(1/2)})-arctanh(c*x)^3+3/4*(2*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x \\
 & ^2-1)))^3+I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(c*x+1)^2/(c^2*x^2 \\
 & -1)/(1-(c*x+1)^2/(c^2*x^2-1)))^2+2*I*Pi*csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1)/ \\
 & (1-(c*x+1)^2/(c^2*x^2-1)))^3+2*I*Pi-2*I*Pi*csgn(I*(-(c*x+1)^2/(c^2*x^2-1)- \\
 & 1))*csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^2-I*Pi*csgn \\
 & (I*(c*x+1)^2/(c^2*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2 \\
 & *x^2-1)))^2-2*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))^2+I*Pi*csgn(I*(c*x+1) \\
 & /(-c^2*x^2+1)^{(1/2)})^2*csgn(I*(c*x+1)^2/(c^2*x^2-1))+I*Pi*csgn(I*(c*x+1)^2 \\
 & /(-c^2*x^2+1)^{(1/2)})^3+2*I*Pi*csgn(I*(c*x+1)/(-c^2*x^2+1)^{(1/2)})*csgn(I*(c*x+1)^2 \\
 & /(-c^2*x^2+1)^{(1/2)})^2+I*Pi*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1) \\
 &))^3-2*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(-(c*x+1)^2/(c^2*x^2- \\
 & 1)-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^2-I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1))) * \\
 & csgn(I*(c*x+1)^2/(c^2*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c \\
 & ^2*x^2-1)))+2*I*Pi*csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1))*csgn(I/(1-(c*x+1)^2/ \\
 & (c^2*x^2-1)))*csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1)/(1-(c*x+1)^2/(c^2*x^2-1))) \\
 & +4*\ln(2)*arctanh(c*x)^2-3*arctanh(c*x)^2*\ln((c*x+1)^2/(-c^2*x^2+1)-1)+3*a \\
 & rctanh(c*x)^2*\ln(1+(c*x+1)/(-c^2*x^2+1)^{(1/2)})+6*arctanh(c*x)*polylog(2,-(\\
 & c*x+1)/(-c^2*x^2+1)^{(1/2)})-6*polylog(3,-(c*x+1)/(-c^2*x^2+1)^{(1/2)})+3*a...
 \end{aligned}$$

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^2} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{x^2} dx$$

input `integrate((a+b*arctanh(c*x))^3/x^2,x, algorithm="fricas")`

output `integral((b^3*arctanh(c*x)^3 + 3*a*b^2*arctanh(c*x)^2 + 3*a^2*b*arctanh(c*x) + a^3)/x^2, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^2} dx = \int \frac{(a + b \operatorname{atanh}(cx))^3}{x^2} dx$$

input `integrate((a+b*atanh(c*x))**3/x**2,x)`

output `Integral((a + b*atanh(c*x))**3/x**2, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^2} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{x^2} dx$$

input `integrate((a+b*arctanh(c*x))^3/x^2,x, algorithm="maxima")`

output `-3/2*(c*(log(c^2*x^2 - 1) - log(x^2)) + 2*arctanh(c*x)/x)*a^2*b - a^3/x - 1/8*((b^3*c*x - b^3)*log(-c*x + 1)^3 + 3*(2*a*b^2 + (b^3*c*x + b^3)*log(c*x + 1))*log(-c*x + 1)^2)/x - integrate(-1/8*((b^3*c*x - b^3)*log(c*x + 1)^3 + 6*(a*b^2*c*x - a*b^2)*log(c*x + 1)^2 + 3*(4*a*b^2*c*x - (b^3*c*x - b^3)*log(c*x + 1)^2 + 2*(b^3*c^2*x^2 + 2*a*b^2 - (2*a*b^2*c - b^3*c)*x)*log(c*x + 1))*log(-c*x + 1))/(c*x^3 - x^2), x)`

3.32 $\int \frac{(a+b\operatorname{arctanh}(cx))^3}{x^3} dx$

Optimal result	343
Mathematica [A] (verified)	344
Rubi [A] (verified)	344
Maple [C] (warning: unable to verify)	347
Fricas [F]	348
Sympy [F]	348
Maxima [F]	348
Giac [F]	349
Mupad [F(-1)]	349
Reduce [F]	350

Optimal result

Integrand size = 14, antiderivative size = 123

$$\begin{aligned} \int \frac{(a + b\operatorname{arctanh}(cx))^3}{x^3} dx &= \frac{3}{2}bc^2(a + b\operatorname{arctanh}(cx))^2 - \frac{3bc(a + b\operatorname{arctanh}(cx))^2}{2x} \\ &\quad + \frac{1}{2}c^2(a + b\operatorname{arctanh}(cx))^3 - \frac{(a + b\operatorname{arctanh}(cx))^3}{2x^2} \\ &\quad + 3b^2c^2(a + b\operatorname{arctanh}(cx)) \log\left(2 - \frac{2}{1 + cx}\right) \\ &\quad - \frac{3}{2}b^3c^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + cx}\right) \end{aligned}$$

output

```
3/2*b*c^2*(a+b*arctanh(c*x))^2-3/2*b*c*(a+b*arctanh(c*x))^2/x+1/2*c^2*(a+b
*arctanh(c*x))^3-1/2*(a+b*arctanh(c*x))^3/x^2+3*b^2*c^2*(a+b*arctanh(c*x))
*ln(2-2/(c*x+1))-3/2*b^3*c^2*polylog(2,-1+2/(c*x+1))
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.56

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^3} dx$$

$$= \frac{6b^2(-1 + cx)(a + acx + bcx)\operatorname{arctanh}(cx)^2 + 2b^3(-1 + c^2x^2)\operatorname{arctanh}(cx)^3 - 6b\operatorname{arctanh}(cx)(a^2 + 2abcx$$

input

```
Integrate[(a + b*ArcTanh[c*x])^3/x^3,x]
```

output

```
(6*b^2*(-1 + c*x)*(a + a*c*x + b*c*x)*ArcTanh[c*x]^2 + 2*b^3*(-1 + c^2*x^2)*ArcTanh[c*x]^3 - 6*b*ArcTanh[c*x]*(a^2 + 2*a*b*c*x - 2*b^2*c^2*x^2*Log[1 - E^(-2*ArcTanh[c*x])]) + a*(-2*a^2 - 6*a*b*c*x - 3*a*b*c^2*x^2*Log[1 - c*x] + 3*a*b*c^2*x^2*Log[1 + c*x] + 12*b^2*c^2*x^2*Log[(c*x)/Sqrt[1 - c^2*x^2]]) - 6*b^3*c^2*x^2*PolyLog[2, E^(-2*ArcTanh[c*x])])/(4*x^2)
```

Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.97, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6452, 6544, 6452, 6510, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^3} dx$$

$$\downarrow 6452$$

$$\frac{3}{2}bc \int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^2(1 - c^2x^2)} dx - \frac{(a + b \operatorname{arctanh}(cx))^3}{2x^2}$$

$$\downarrow 6544$$

$$\frac{3}{2}bc \left(c^2 \int \frac{(a + b \operatorname{arctanh}(cx))^2}{1 - c^2x^2} dx + \int \frac{(a + b \operatorname{arctanh}(cx))^2}{x^2} dx \right) - \frac{(a + b \operatorname{arctanh}(cx))^3}{2x^2}$$

$$\downarrow 6452$$

$$\frac{3}{2}bc \left(c^2 \int \frac{(a + \operatorname{barctanh}(cx))^2}{1 - c^2x^2} dx + 2bc \int \frac{a + \operatorname{barctanh}(cx)}{x(1 - c^2x^2)} dx - \frac{(a + \operatorname{barctanh}(cx))^2}{x} \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{2x^2}$$

↓ 6510

$$\frac{3}{2}bc \left(2bc \int \frac{a + \operatorname{barctanh}(cx)}{x(1 - c^2x^2)} dx + \frac{c(a + \operatorname{barctanh}(cx))^3}{3b} - \frac{(a + \operatorname{barctanh}(cx))^2}{x} \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{2x^2}$$

↓ 6550

$$\frac{3}{2}bc \left(2bc \left(\int \frac{a + \operatorname{barctanh}(cx)}{x(cx + 1)} dx + \frac{(a + \operatorname{barctanh}(cx))^2}{2b} \right) + \frac{c(a + \operatorname{barctanh}(cx))^3}{3b} - \frac{(a + \operatorname{barctanh}(cx))^2}{x} \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{2x^2}$$

↓ 6494

$$\frac{3}{2}bc \left(2bc \left(-bc \int \frac{\log\left(2 - \frac{2}{cx+1}\right)}{1 - c^2x^2} dx + \frac{(a + \operatorname{barctanh}(cx))^2}{2b} + \log\left(2 - \frac{2}{cx+1}\right) (a + \operatorname{barctanh}(cx)) \right) + \frac{c(a + \operatorname{barctanh}(cx))^3}{2x^2} \right)$$

↓ 2897

$$\frac{3}{2}bc \left(2bc \left(\frac{(a + \operatorname{barctanh}(cx))^2}{2b} + \log\left(2 - \frac{2}{cx+1}\right) (a + \operatorname{barctanh}(cx)) - \frac{1}{2}b \operatorname{PolyLog}\left(2, \frac{2}{cx+1} - 1\right) \right) + \frac{c(a + \operatorname{barctanh}(cx))^3}{2x^2} \right)$$

input `Int[(a + b*ArcTanh[c*x])^3/x^3,x]`

output `-1/2*(a + b*ArcTanh[c*x])^3/x^2 + (3*b*c*(-((a + b*ArcTanh[c*x])^2/x) + (c*(a + b*ArcTanh[c*x])^3)/(3*b) + 2*b*c*((a + b*ArcTanh[c*x])^2/(2*b) + (a + b*ArcTanh[c*x])*Log[2 - 2/(1 + c*x)] - (b*PolyLog[2, -1 + 2/(1 + c*x)]))/2))/2`

Definitions of rubi rules used

rule 2897 `Int[Log[u_]*(Pq_)^(m_.), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6494 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_))), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6510 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6544 `Int[(((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*((f_.)*(x_)^(m_)))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]`

rule 6550 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_)^2)), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 8.77 (sec) , antiderivative size = 4599, normalized size of antiderivative = 37.39

method	result	size
derivativedivides	Expression too large to display	4599
default	Expression too large to display	4599
parts	Expression too large to display	4601

input `int((a+b*arctanh(c*x))^3/x^3,x,method=_RETURNVERBOSE)`

output

```
c^2*(-1/2*a^3/c^2/x^2+b^3*(3/8*I*Pi*csgn(I*(c*x+1)/(-c^2*x^2+1)^(1/2))^2*csgn(I*(c*x+1)^2/(c^2*x^2-1))*arctanh(c*x)*ln(1-(c*x+1)/(-c^2*x^2+1)^(1/2))-3/8*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(c*x+1)^2/(c^2*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))*polylog(2,-(c*x+1)/(-c^2*x^2+1)^(1/2))-3/8*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(c*x+1)^2/(c^2*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))*polylog(2,(c*x+1)/(-c^2*x^2+1)^(1/2))+3/8*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(c*x+1)^2/(c^2*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))*dilog(1+(c*x+1)/(-c^2*x^2+1)^(1/2))-3/8*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(c*x+1)^2/(c^2*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))*dilog((c*x+1)/(-c^2*x^2+1)^(1/2))-3/8*I*Pi*csgn(I*(c*x+1)^2/(c^2*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^2*arctanh(c*x)*ln(1-(c*x+1)/(-c^2*x^2+1)^(1/2))+3/8*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(c*x+1)^2/(c^2*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))*arctanh(c*x)^2-3/2*arctanh(c*x)^2+3/8*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(c*x+1)^2/(c^2*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^2*arctanh(c*x)*ln(1-(c*x+1)/(-c^2*x^2+1)^(1/2))+3/2*polylog(2,-(c*x+1)/(-c^2*x^2+1)^(1/2))-3/8*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(c*x+1)^2/(c^2*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))*arctanh(c*x)*ln(1-(c*x+1)/(-c^2*x^2+1)^(...
```


Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^3} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{x^3} dx$$

input `integrate((a+b*arctanh(c*x))^3/x^3,x, algorithm="fricas")`

output `integral((b^3*arctanh(c*x)^3 + 3*a*b^2*arctanh(c*x)^2 + 3*a^2*b*arctanh(c*x) + a^3)/x^3, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^3} dx = \int \frac{(a + b \operatorname{atanh}(cx))^3}{x^3} dx$$

input `integrate((a+b*atanh(c*x))**3/x**3,x)`

output `Integral((a + b*atanh(c*x))**3/x**3, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^3} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{x^3} dx$$

input `integrate((a+b*arctanh(c*x))^3/x^3,x, algorithm="maxima")`

output

```
3/4*((c*log(c*x + 1) - c*log(c*x - 1) - 2/x)*c - 2*arctanh(c*x)/x^2)*a^2*b
+ 3/8*((2*(log(c*x - 1) - 2)*log(c*x + 1) - log(c*x + 1)^2 - log(c*x - 1)
^2 - 4*log(c*x - 1) + 8*log(x))*c^2 + 4*(c*log(c*x + 1) - c*log(c*x - 1) -
2/x)*c*arctanh(c*x))*a*b^2 - 1/16*b^3*(((c^2*x^2 - 1)*log(-c*x + 1)^3 + 3
*(2*c*x - (c^2*x^2 - 1)*log(c*x + 1))*log(-c*x + 1)^2)/x^2 + 2*integrate(-
((c*x - 1)*log(c*x + 1)^3 + 3*(2*c^2*x^2 - (c*x - 1)*log(c*x + 1)^2 - (c^3
*x^3 - c*x)*log(c*x + 1))*log(-c*x + 1))/(c*x^4 - x^3), x)) - 3/2*a*b^2*ar
ctanh(c*x)^2/x^2 - 1/2*a^3/x^2
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^3} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{x^3} dx$$

input

```
integrate((a+b*arctanh(c*x))^3/x^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x) + a)^3/x^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^3} dx = \int \frac{(a + b \operatorname{atanh}(cx))^3}{x^3} dx$$

input

```
int((a + b*atanh(c*x))^3/x^3,x)
```

output

```
int((a + b*atanh(c*x))^3/x^3, x)
```


3.33 $\int \frac{(a+b\operatorname{arctanh}(cx))^3}{x^4} dx$

Optimal result	351
Mathematica [C] (verified)	352
Rubi [A] (verified)	352
Maple [C] (warning: unable to verify)	357
Fricas [F]	358
Sympy [F]	358
Maxima [F]	358
Giac [F]	359
Mupad [F(-1)]	359
Reduce [F]	360

Optimal result

Integrand size = 14, antiderivative size = 200

$$\int \frac{(a + b\operatorname{arctanh}(cx))^3}{x^4} dx = -\frac{b^2c^2(a + b\operatorname{arctanh}(cx))}{x} + \frac{1}{2}bc^3(a + b\operatorname{arctanh}(cx))^2 - \frac{bc(a + b\operatorname{arctanh}(cx))^2}{2x^2} + \frac{1}{3}c^3(a + b\operatorname{arctanh}(cx))^3 - \frac{(a + b\operatorname{arctanh}(cx))^3}{3x^3} + b^3c^3 \log(x) - \frac{1}{2}b^3c^3 \log(1 - c^2x^2) + bc^3(a + b\operatorname{arctanh}(cx))^2 \log\left(2 - \frac{2}{1 + cx}\right) - b^2c^3(a + b\operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + cx}\right) - \frac{1}{2}b^3c^3 \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 + cx}\right)$$

output

```
-b^2*c^2*(a+b*arctanh(c*x))/x+1/2*b*c^3*(a+b*arctanh(c*x))^2-1/2*b*c*(a+b*arctanh(c*x))^2/x^2+1/3*c^3*(a+b*arctanh(c*x))^3-1/3*(a+b*arctanh(c*x))^3/x^3+b^3*c^3*ln(x)-1/2*b^3*c^3*ln(-c^2*x^2+1)+b*c^3*(a+b*arctanh(c*x))^2*ln(2-2/(c*x+1))-b^2*c^3*(a+b*arctanh(c*x))*polylog(2,-1+2/(c*x+1))-1/2*b^3*c^3*polylog(3,-1+2/(c*x+1))
```


$$\begin{aligned}
& \downarrow 6544 \\
& bc \left(c^2 \int \frac{(a + \operatorname{barctanh}(cx))^2}{x(1-c^2x^2)} dx + \int \frac{(a + \operatorname{barctanh}(cx))^2}{x^3} dx \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{3x^3} \\
& \downarrow 6452 \\
& bc \left(c^2 \int \frac{(a + \operatorname{barctanh}(cx))^2}{x(1-c^2x^2)} dx + bc \int \frac{a + \operatorname{barctanh}(cx)}{x^2(1-c^2x^2)} dx - \frac{(a + \operatorname{barctanh}(cx))^2}{2x^2} \right) - \\
& \quad \frac{(a + \operatorname{barctanh}(cx))^3}{3x^3} \\
& \downarrow 6544 \\
& bc \left(c^2 \int \frac{(a + \operatorname{barctanh}(cx))^2}{x(1-c^2x^2)} dx + bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{1-c^2x^2} dx + \int \frac{a + \operatorname{barctanh}(cx)}{x^2} dx \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{3x^3} \right) - \frac{(a + \operatorname{barctanh}(cx))^2}{2x^2} \\
& \downarrow 6452 \\
& bc \left(c^2 \int \frac{(a + \operatorname{barctanh}(cx))^2}{x(1-c^2x^2)} dx + bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{1-c^2x^2} dx + bc \int \frac{1}{x(1-c^2x^2)} dx - \frac{a + \operatorname{barctanh}(cx)}{x} \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{3x^3} \right) \\
& \downarrow 243 \\
& bc \left(c^2 \int \frac{(a + \operatorname{barctanh}(cx))^2}{x(1-c^2x^2)} dx + bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{1-c^2x^2} dx + \frac{1}{2} bc \int \frac{1}{x^2(1-c^2x^2)} dx^2 - \frac{a + \operatorname{barctanh}(cx)}{x} \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{3x^3} \right) \\
& \downarrow 47 \\
& bc \left(c^2 \int \frac{(a + \operatorname{barctanh}(cx))^2}{x(1-c^2x^2)} dx + bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{1-c^2x^2} dx + \frac{1}{2} bc \left(c^2 \int \frac{1}{1-c^2x^2} dx^2 + \int \frac{1}{x^2} dx^2 \right) - \frac{a + \operatorname{barctanh}(cx)}{x} \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{3x^3} \right) \\
& \downarrow 14 \\
& bc \left(c^2 \int \frac{(a + \operatorname{barctanh}(cx))^2}{x(1-c^2x^2)} dx + bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{1-c^2x^2} dx + \frac{1}{2} bc \left(c^2 \int \frac{1}{1-c^2x^2} dx^2 + \log(x^2) \right) - \frac{a + \operatorname{barctanh}(cx)}{x} \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{3x^3} \right)
\end{aligned}$$

↓ 16

$$bc \left(c^2 \int \frac{(a + \operatorname{barctanh}(cx))^2}{x(1 - c^2x^2)} dx + bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{1 - c^2x^2} dx - \frac{a + \operatorname{barctanh}(cx)}{x} + \frac{1}{2}bc(\log(x^2) - \log(1 - c^2x^2)) \right) \right) \frac{(a + \operatorname{barctanh}(cx))^3}{3x^3}$$

↓ 6510

$$bc \left(c^2 \int \frac{(a + \operatorname{barctanh}(cx))^2}{x(1 - c^2x^2)} dx + bc \left(\frac{c(a + \operatorname{barctanh}(cx))^2}{2b} - \frac{a + \operatorname{barctanh}(cx)}{x} + \frac{1}{2}bc(\log(x^2) - \log(1 - c^2x^2)) \right) \right) \frac{(a + \operatorname{barctanh}(cx))^3}{3x^3}$$

↓ 6550

$$bc \left(c^2 \left(\int \frac{(a + \operatorname{barctanh}(cx))^2}{x(cx + 1)} dx + \frac{(a + \operatorname{barctanh}(cx))^3}{3b} \right) + bc \left(\frac{c(a + \operatorname{barctanh}(cx))^2}{2b} - \frac{a + \operatorname{barctanh}(cx)}{x} + \frac{1}{2}bc(\log(x^2) - \log(1 - c^2x^2)) \right) \right) \frac{(a + \operatorname{barctanh}(cx))^3}{3x^3}$$

↓ 6494

$$bc \left(c^2 \left(-2bc \int \frac{(a + \operatorname{barctanh}(cx)) \log\left(2 - \frac{2}{cx+1}\right)}{1 - c^2x^2} dx + \frac{(a + \operatorname{barctanh}(cx))^3}{3b} + \log\left(2 - \frac{2}{cx+1}\right) (a + \operatorname{barctanh}(cx)) \right) \right) \frac{(a + \operatorname{barctanh}(cx))^3}{3x^3}$$

↓ 6618

$$bc \left(c^2 \left(-2bc \left(\frac{\operatorname{PolyLog}\left(2, \frac{2}{cx+1} - 1\right) (a + \operatorname{barctanh}(cx))}{2c} - \frac{1}{2}b \int \frac{\operatorname{PolyLog}\left(2, \frac{2}{cx+1} - 1\right)}{1 - c^2x^2} dx \right) + \frac{(a + \operatorname{barctanh}(cx))^3}{3b} \right) \right) \frac{(a + \operatorname{barctanh}(cx))^3}{3x^3}$$

↓ 7164

$$bc \left(c^2 \left(-2bc \left(\frac{\operatorname{PolyLog}\left(2, \frac{2}{cx+1} - 1\right) (a + \operatorname{barctanh}(cx))}{2c} + \frac{b \operatorname{PolyLog}\left(3, \frac{2}{cx+1} - 1\right)}{4c} \right) + \frac{(a + \operatorname{barctanh}(cx))^3}{3b} \right) \right) \frac{(a + \operatorname{barctanh}(cx))^3}{3x^3}$$

input `Int[(a + b*ArcTanh[c*x])^3/x^4,x]`

output `-1/3*(a + b*ArcTanh[c*x])^3/x^3 + b*c*(-1/2*(a + b*ArcTanh[c*x])^2/x^2 + b*c*(-((a + b*ArcTanh[c*x])/x) + (c*(a + b*ArcTanh[c*x])^2)/(2*b) + (b*c*(Log[x^2] - Log[1 - c^2*x^2]))/2) + c^2*((a + b*ArcTanh[c*x])^3/(3*b) + (a + b*ArcTanh[c*x])^2*Log[2 - 2/(1 + c*x)] - 2*b*c*((a + b*ArcTanh[c*x])*PolyLog[2, -1 + 2/(1 + c*x)])/(2*c) + (b*PolyLog[3, -1 + 2/(1 + c*x)])/(4*c))`

Defintions of rubi rules used

rule 14 `Int[(a_)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_)/((a_) + (b_)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/(((a_) + (b_)*(x_))*((c_) + (d_)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

rule 243 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6494 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((x) \cdot (d + e \cdot x)), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{Log}[2 - 2/(1 + e \cdot (x/d))]/d), x] - \text{Simp}[b \cdot c \cdot (p/d) \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{Log}[2 - 2/(1 + e \cdot (x/d))]) / (1 - c^2 \cdot x^2)], x], x] /;$ FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]

rule 6510 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((d + e \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot c \cdot d \cdot (p+1)), x] /;$ FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]

rule 6544 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p \cdot (f \cdot x)^m / ((d + e \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[1/d \text{Int}[(f \cdot x)^m \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x] - \text{Simp}[e/(d \cdot f^2) \text{Int}[(f \cdot x)^{m+2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2)], x], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]

rule 6550 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((x) \cdot (d + e \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot d \cdot (p+1)), x] + \text{Simp}[1/d \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p / (x \cdot (1 + c \cdot x)), x], x] /;$ FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]

rule 6618 $\text{Int}[(\text{Log}[u] \cdot (a + \text{ArcTanh}[c \cdot x] \cdot b)^p) / ((d + e \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{PolyLog}[2, 1 - u] / (2 \cdot c \cdot d)), x] - \text{Simp}[b \cdot (p/2) \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{PolyLog}[2, 1 - u] / (d + e \cdot x^2))], x], x] /;$ FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 + c \cdot x))^2, 0]

rule 7164 $\text{Int}(u \cdot \text{PolyLog}[n, v], x_Symbol) \rightarrow \text{With}\{w = \text{DerivativeDivides}[v, u \cdot v, x]\}, \text{Simp}[w \cdot \text{PolyLog}[n + 1, v], x] /;$!FalseQ[w] /; FreeQ[n, x]

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 12.88 (sec) , antiderivative size = 1581, normalized size of antiderivative = 7.90

method	result	size
derivativeldivides	Expression too large to display	1581
default	Expression too large to display	1581
parts	Expression too large to display	1583

input `int((a+b*arctanh(c*x))^3/x^4,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & c^3*(-1/3*a^3/c^3/x^3+b^3*(1/2*arctanh(c*x)^2-1/4*I*Pi*csgn(I/(1-(c*x+1)^2 \\
 & / (c^2*x^2-1))) *csgn(I*(c*x+1)^2/(c^2*x^2-1)) *csgn(I*(c*x+1)^2/(c^2*x^2-1)/ \\
 & (1-(c*x+1)^2/(c^2*x^2-1))) *arctanh(c*x)^2+\ln(1+(c*x+1)/(-c^2*x^2+1)^(1/2)) \\
 & -2*polylog(3,-(c*x+1)/(-c^2*x^2+1)^(1/2))+\ln(c*x)*arctanh(c*x)^2-arctanh(c \\
 & *x)^2*\ln((c*x+1)^2/(-c^2*x^2+1)-1)+arctanh(c*x)^2*\ln(1+(c*x+1)/(-c^2*x^2+1 \\
 &)^(1/2))+2*arctanh(c*x)*polylog(2,-(c*x+1)/(-c^2*x^2+1)^(1/2))+arctanh(c*x \\
 &)^2*\ln(1-(c*x+1)/(-c^2*x^2+1)^(1/2))+2*arctanh(c*x)*polylog(2,(c*x+1)/(-c^ \\
 & 2*x^2+1)^(1/2))-1/2/c^2/x^2*arctanh(c*x)^2-2*polylog(3,(c*x+1)/(-c^2*x^2+1 \\
 &)^(1/2))+1/2*I*Pi*csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1)) *csgn(I/(1-(c*x+1)^2/(\\
 & c^2*x^2-1))) *csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1)/(1-(c*x+1)^2/(c^2*x^2-1))) * \\
 & arctanh(c*x)^2+1/4*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1))) *csgn(I*(c*x+1)^2 \\
 & / (c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^2*arctanh(c*x)^2-1/2*arctanh(c*x)* \\
 & (c*x+(-c^2*x^2+1)^(1/2)+1)/c/x+1/2*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))^ \\
 & 3*arctanh(c*x)^2+1/2*I*Pi*csgn(I*(-(c*x+1)^2/(c^2*x^2-1)-1)/(1-(c*x+1)^2/(\\
 & c^2*x^2-1)))^3*arctanh(c*x)^2-1/2*I*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))^2 \\
 & *arctanh(c*x)^2+1/4*I*Pi*csgn(I*(c*x+1)^2/(c^2*x^2-1))^3*arctanh(c*x)^2+1/ \\
 & 4*I*Pi*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^3*arctanh(c \\
 & *x)^2-1/2*arctanh(c*x)^2*\ln(c*x-1)-1/2*arctanh(c*x)^2*\ln(c*x+1)+arctanh(c \\
 & *x)^2*\ln((c*x+1)/(-c^2*x^2+1)^(1/2))+\ln(2)*arctanh(c*x)^2-1/4*I*Pi*csgn(I*(\\
 & c*x+1)^2/(c^2*x^2-1)) *csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^...
 \end{aligned}$$

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^4} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{x^4} dx$$

input `integrate((a+b*arctanh(c*x))^3/x^4,x, algorithm="fricas")`

output `integral((b^3*arctanh(c*x)^3 + 3*a*b^2*arctanh(c*x)^2 + 3*a^2*b*arctanh(c*x) + a^3)/x^4, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^4} dx = \int \frac{(a + b \operatorname{atanh}(cx))^3}{x^4} dx$$

input `integrate((a+b*atanh(c*x))**3/x**4,x)`

output `Integral((a + b*atanh(c*x))**3/x**4, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^4} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{x^4} dx$$

input `integrate((a+b*arctanh(c*x))^3/x^4,x, algorithm="maxima")`

output

```
-1/2*((c^2*log(c^2*x^2 - 1) - c^2*log(x^2) + 1/x^2)*c + 2*arctanh(c*x)/x^3
)*a^2*b - 1/3*a^3/x^3 - 1/24*((b^3*c^3*x^3 - b^3)*log(-c*x + 1)^3 + 3*(b^3
*c*x + 2*a*b^2 + (b^3*c^3*x^3 + b^3)*log(c*x + 1))*log(-c*x + 1)^2)/x^3 -
integrate(-1/8*((b^3*c*x - b^3)*log(c*x + 1)^3 + 6*(a*b^2*c*x - a*b^2)*log
(c*x + 1)^2 + (2*b^3*c^2*x^2 + 4*a*b^2*c*x - 3*(b^3*c*x - b^3)*log(c*x + 1
)^2 + 2*(b^3*c^4*x^4 + 6*a*b^2 - (6*a*b^2*c - b^3*c)*x)*log(c*x + 1))*log(
-c*x + 1))/(c*x^5 - x^4), x)
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^4} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{x^4} dx$$

input

```
integrate((a+b*arctanh(c*x))^3/x^4,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x) + a)^3/x^4, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^4} dx = \int \frac{(a + b \operatorname{atanh}(cx))^3}{x^4} dx$$

input

```
int((a + b*atanh(c*x))^3/x^4,x)
```

output

```
int((a + b*atanh(c*x))^3/x^4, x)
```


3.34 $\int \frac{(a+b\operatorname{arctanh}(cx))^3}{x^5} dx$

Optimal result	361
Mathematica [A] (verified)	362
Rubi [A] (verified)	362
Maple [C] (warning: unable to verify)	366
Fricas [F]	367
Sympy [F]	367
Maxima [F]	367
Giac [F]	368
Mupad [F(-1)]	368
Reduce [F]	369

Optimal result

Integrand size = 14, antiderivative size = 187

$$\begin{aligned} \int \frac{(a + b\operatorname{arctanh}(cx))^3}{x^5} dx = & -\frac{b^3c^3}{4x} + \frac{1}{4}b^3c^4\operatorname{arctanh}(cx) - \frac{b^2c^2(a + b\operatorname{arctanh}(cx))}{4x^2} \\ & + bc^4(a + b\operatorname{arctanh}(cx))^2 - \frac{bc(a + b\operatorname{arctanh}(cx))^2}{4x^3} \\ & - \frac{3bc^3(a + b\operatorname{arctanh}(cx))^2}{4x} \\ & + \frac{1}{4}c^4(a + b\operatorname{arctanh}(cx))^3 - \frac{(a + b\operatorname{arctanh}(cx))^3}{4x^4} \\ & + 2b^2c^4(a + b\operatorname{arctanh}(cx)) \log\left(2 - \frac{2}{1 + cx}\right) \\ & - b^3c^4 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + cx}\right) \end{aligned}$$

output

```
-1/4*b^3*c^3/x+1/4*b^3*c^4*arctanh(c*x)-1/4*b^2*c^2*(a+b*arctanh(c*x))/x^2
+b*c^4*(a+b*arctanh(c*x))^2-1/4*b*c*(a+b*arctanh(c*x))^2/x^3-3/4*b*c^3*(a
+b*arctanh(c*x))^2/x+1/4*c^4*(a+b*arctanh(c*x))^3-1/4*(a+b*arctanh(c*x))^3/
x^4+2*b^2*c^4*(a+b*arctanh(c*x))*ln(2-2/(c*x+1))-b^3*c^4*polylog(2,-1+2/(c
*x+1))
```


$$\frac{3}{4}bc \left(c^2 \int \frac{(a + \operatorname{barctanh}(cx))^2}{x^2(1-c^2x^2)} dx + \frac{2}{3}bc \int \frac{a + \operatorname{barctanh}(cx)}{x^3(1-c^2x^2)} dx - \frac{(a + \operatorname{barctanh}(cx))^2}{3x^3} \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{4x^4}$$

↓ 6452

$$\frac{3}{4}bc \left(c^2 \left(c^2 \int \frac{(a + \operatorname{barctanh}(cx))^2}{1-c^2x^2} dx + \int \frac{(a + \operatorname{barctanh}(cx))^2}{x^2} dx \right) + \frac{2}{3}bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{x(1-c^2x^2)} dx + \int \frac{a + \operatorname{barctanh}(cx)}{x} dx \right) \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{4x^4}$$

↓ 6544

$$\frac{3}{4}bc \left(c^2 \left(c^2 \int \frac{(a + \operatorname{barctanh}(cx))^2}{1-c^2x^2} dx + 2bc \int \frac{a + \operatorname{barctanh}(cx)}{x(1-c^2x^2)} dx - \frac{(a + \operatorname{barctanh}(cx))^2}{x} \right) + \frac{2}{3}bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{x(1-c^2x^2)} dx + \int \frac{a + \operatorname{barctanh}(cx)}{x} dx \right) \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{4x^4}$$

↓ 6452

$$\frac{3}{4}bc \left(c^2 \left(c^2 \int \frac{(a + \operatorname{barctanh}(cx))^2}{1-c^2x^2} dx + 2bc \int \frac{a + \operatorname{barctanh}(cx)}{x(1-c^2x^2)} dx - \frac{(a + \operatorname{barctanh}(cx))^2}{x} \right) + \frac{2}{3}bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{x(1-c^2x^2)} dx + \int \frac{a + \operatorname{barctanh}(cx)}{x} dx \right) \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{4x^4}$$

↓ 264

$$\frac{3}{4}bc \left(c^2 \left(c^2 \int \frac{(a + \operatorname{barctanh}(cx))^2}{1-c^2x^2} dx + 2bc \int \frac{a + \operatorname{barctanh}(cx)}{x(1-c^2x^2)} dx - \frac{(a + \operatorname{barctanh}(cx))^2}{x} \right) + \frac{2}{3}bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{x(1-c^2x^2)} dx + \int \frac{a + \operatorname{barctanh}(cx)}{x} dx \right) \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{4x^4}$$

↓ 219

$$\frac{3}{4}bc \left(c^2 \left(2bc \int \frac{a + \operatorname{barctanh}(cx)}{x(1-c^2x^2)} dx + \frac{c(a + \operatorname{barctanh}(cx))^3}{3b} - \frac{(a + \operatorname{barctanh}(cx))^2}{x} \right) + \frac{2}{3}bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{x(1-c^2x^2)} dx + \int \frac{a + \operatorname{barctanh}(cx)}{x} dx \right) \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{4x^4}$$

↓ 6510

$$\frac{3}{4}bc \left(c^2 \left(2bc \int \frac{a + \operatorname{barctanh}(cx)}{x(1-c^2x^2)} dx + \frac{c(a + \operatorname{barctanh}(cx))^3}{3b} - \frac{(a + \operatorname{barctanh}(cx))^2}{x} \right) + \frac{2}{3}bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx)}{x(1-c^2x^2)} dx + \int \frac{a + \operatorname{barctanh}(cx)}{x} dx \right) \right) - \frac{(a + \operatorname{barctanh}(cx))^3}{4x^4}$$

↓ 6550

$$\frac{3}{4}bc \left(\frac{2}{3}bc \left(c^2 \left(\int \frac{a + \operatorname{barctanh}(cx)}{x(cx+1)} dx + \frac{(a + \operatorname{barctanh}(cx))^2}{2b} \right) - \frac{a + \operatorname{barctanh}(cx)}{2x^2} + \frac{1}{2}bc \left(\operatorname{carctanh}(cx) - \frac{1}{x} \right) \right) \right) \frac{(a + \operatorname{barctanh}(cx))^3}{4x^4}$$

↓ 6494

$$\frac{3}{4}bc \left(c^2 \left(2bc \left(-bc \int \frac{\log \left(2 - \frac{2}{cx+1} \right)}{1 - c^2x^2} dx + \frac{(a + \operatorname{barctanh}(cx))^2}{2b} + \log \left(2 - \frac{2}{cx+1} \right) (a + \operatorname{barctanh}(cx)) \right) \right) \right) \frac{(a + \operatorname{barctanh}(cx))^3}{4x^4} + c(a$$

↓ 2897

$$\frac{3}{4}bc \left(\frac{2}{3}bc \left(c^2 \left(\frac{(a + \operatorname{barctanh}(cx))^2}{2b} + \log \left(2 - \frac{2}{cx+1} \right) (a + \operatorname{barctanh}(cx)) - \frac{1}{2}b \operatorname{PolyLog} \left(2, \frac{2}{cx+1} - 1 \right) \right) \right) \right) \frac{(a + \operatorname{barctanh}(cx))^3}{4x^4} - c$$

input `Int[(a + b*ArcTanh[c*x])^3/x^5,x]`

output `-1/4*(a + b*ArcTanh[c*x])^3/x^4 + (3*b*c*(-1/3*(a + b*ArcTanh[c*x])^2/x^3 + c^2*(-((a + b*ArcTanh[c*x])^2/x) + (c*(a + b*ArcTanh[c*x])^3)/(3*b) + 2*b*c*((a + b*ArcTanh[c*x])^2/(2*b) + (a + b*ArcTanh[c*x])*Log[2 - 2/(1 + c*x)] - (b*PolyLog[2, -1 + 2/(1 + c*x)]/2)) + (2*b*c*(-1/2*(a + b*ArcTanh[c*x])/x^2 + (b*c*(-x^(-1) + c*ArcTanh[c*x]))/2 + c^2*((a + b*ArcTanh[c*x])^2/(2*b) + (a + b*ArcTanh[c*x])*Log[2 - 2/(1 + c*x)] - (b*PolyLog[2, -1 + 2/(1 + c*x)]/2))))/3)/4`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 264 $\text{Int}[(c \cdot x)^m \cdot (a + b \cdot x^2)^p, x_Symbol] \rightarrow \text{Simp}[(c \cdot x)^{m+1} \cdot (a + b \cdot x^2)^{p+1} / (a \cdot c \cdot (m+1)), x] - \text{Simp}[b \cdot (m+2 \cdot p+3) / (a \cdot c^2 \cdot (m+1)) \text{Int}[(c \cdot x)^{m+2} \cdot (a + b \cdot x^2)^p, x], x] /;$ FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]

rule 2897 $\text{Int}[\text{Log}[u] \cdot (Pq)^m, x_Symbol] \rightarrow \text{With}[\{C = \text{FullSimplify}[Pq^m \cdot ((1-u)/D[u, x])]\}, \text{Simp}[C \cdot \text{PolyLog}[2, 1-u], x] /;$ FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]

rule 6452 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot b)^p \cdot x^m, x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^p / (m+1), x] - \text{Simp}[b \cdot c \cdot n \cdot (p/(m+1)) \text{Int}[x^{m+n} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2n}), x], x] /;$ FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]

rule 6494 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((d + e \cdot x)), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{Log}[2 - 2/(1 + e \cdot (x/d))]/d), x] - \text{Simp}[b \cdot c \cdot (p/d) \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{Log}[2 - 2/(1 + e \cdot (x/d))]) / (1 - c^2 \cdot x^2), x], x] /;$ FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2 \cdot d^2 - e^2, 0]

rule 6510 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((d + e \cdot x^2)), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot c \cdot d \cdot (p+1)), x] /;$ FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2 \cdot d + e, 0] && NeQ[p, -1]

rule 6544 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p \cdot (f \cdot x)^m / (d + e \cdot x^2), x_Symbol] \rightarrow \text{Simp}[1/d \text{Int}[(f \cdot x)^m \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x] - \text{Simp}[e/(d \cdot f^2) \text{Int}[(f \cdot x)^{m+2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2), x], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]

rule 6550

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)/((x_.)*((d_.) + (e_.)*(x_)^2)),
 x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/
 d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 9.13 (sec) , antiderivative size = 1136, normalized size of antiderivative = 6.07

method	result	size
derivativeldivides	Expression too large to display	1136
default	Expression too large to display	1136
parts	Expression too large to display	1193

input

```
int((a+b*arctanh(c*x))^3/x^5,x,method=_RETURNVERBOSE)
```

output

```
c^4*(3/8*I*b^3*Pi*arctanh(c*x)^2-1/4*a^3/c^4/x^4+3/16*I*b^3*Pi*csgn(I*(c*x
+1)^2/(c^2*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))
^2*arctanh(c*x)^2+2*b^3*arctanh(c*x)*ln(1+(c*x+1)/(-c^2*x^2+1)^(1/2))-3/8*
I*b^3*Pi*csgn(I*(c*x+1)/(-c^2*x^2+1)^(1/2))*csgn(I*(c*x+1)^2/(c^2*x^2-1))^
2*arctanh(c*x)^2+3/16*I*b^3*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(c
*x+1)^2/(c^2*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)
))*arctanh(c*x)^2-3/16*I*b^3*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(
c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^2*arctanh(c*x)^2-3/16*I*b^
3*Pi*csgn(I*(c*x+1)/(-c^2*x^2+1)^(1/2))^2*csgn(I*(c*x+1)^2/(c^2*x^2-1))*ar
ctanh(c*x)^2-2*b^3*dilog((c*x+1)/(-c^2*x^2+1)^(1/2))+2*b^3*dilog(1+(c*x+1)
/(-c^2*x^2+1)^(1/2))-1/4*b^3/c^4/x^4*arctanh(c*x)^3-1/4*b^3*arctanh(c*x)^2
/c^3/x^3-1/4*b^3*arctanh(c*x)/c^2/x^2-3/4*b^3/c/x*arctanh(c*x)^2+1/4*b^3*a
rctanh(c*x)+1/4*b^3*arctanh(c*x)^3-b^3*arctanh(c*x)^2+3/8*I*b^3*Pi*csgn(I/
(1-(c*x+1)^2/(c^2*x^2-1)))^3*arctanh(c*x)^2-3/8*I*b^3*Pi*csgn(I/(1-(c*x+1)
^2/(c^2*x^2-1)))^2*arctanh(c*x)^2-3/16*I*b^3*Pi*csgn(I*(c*x+1)^2/(c^2*x^2-
1)/(1-(c*x+1)^2/(c^2*x^2-1)))^3*arctanh(c*x)^2-3/16*I*b^3*Pi*csgn(I*(c*x+1)
)^2/(c^2*x^2-1))^3*arctanh(c*x)^2-3/4*b^3*arctanh(c*x)^2*ln((c*x+1)/(-c^2*
x^2+1)^(1/2))-3/8*b^3*arctanh(c*x)^2*ln(c*x-1)+3/8*b^3*arctanh(c*x)^2*ln(c
*x+1)+3*a*b^2*(-1/4/c^4/x^4*arctanh(c*x)^2+1/4*arctanh(c*x)*ln(c*x+1)-1/6/
c^3/x^3*arctanh(c*x)-1/2/c/x*arctanh(c*x)-1/4*arctanh(c*x)*ln(c*x-1)+1/...
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^5} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{x^5} dx$$

input `integrate((a+b*arctanh(c*x))^3/x^5,x, algorithm="fricas")`

output `integral((b^3*arctanh(c*x)^3 + 3*a*b^2*arctanh(c*x)^2 + 3*a^2*b*arctanh(c*x) + a^3)/x^5, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^5} dx = \int \frac{(a + b \operatorname{atanh}(cx))^3}{x^5} dx$$

input `integrate((a+b*atanh(c*x))**3/x**5,x)`

output `Integral((a + b*atanh(c*x))**3/x**5, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^5} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{x^5} dx$$

input `integrate((a+b*arctanh(c*x))^3/x^5,x, algorithm="maxima")`

output

```
1/8*((3*c^3*log(c*x + 1) - 3*c^3*log(c*x - 1) - 2*(3*c^2*x^2 + 1)/x^3)*c -
6*arctanh(c*x)/x^4)*a^2*b + 1/16*((32*c^2*log(x) - (3*c^2*x^2*log(c*x + 1)
)^2 + 3*c^2*x^2*log(c*x - 1)^2 + 16*c^2*x^2*log(c*x - 1) - 2*(3*c^2*x^2*lo
g(c*x - 1) - 8*c^2*x^2)*log(c*x + 1) + 4)/x^2)*c^2 + 4*(3*c^3*log(c*x + 1)
- 3*c^3*log(c*x - 1) - 2*(3*c^2*x^2 + 1)/x^3)*c*arctanh(c*x))*a*b^2 - 1/3
2*b^3*(((c^4*x^4 - 1)*log(-c*x + 1)^3 + (6*c^3*x^3 + 2*c*x - 3*(c^4*x^4 -
1)*log(c*x + 1))*log(-c*x + 1)^2)/x^4 + 4*integrate(-1/2*(2*(c*x - 1)*log(
c*x + 1)^3 + (6*c^4*x^4 + 2*c^2*x^2 - 6*(c*x - 1)*log(c*x + 1)^2 - 3*(c^5*x
^5 - c*x)*log(c*x + 1))*log(-c*x + 1))/(c*x^6 - x^5), x)) - 3/4*a*b^2*arc
tanh(c*x)^2/x^4 - 1/4*a^3/x^4
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^5} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{x^5} dx$$

input

```
integrate((a+b*arctanh(c*x))^3/x^5,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x) + a)^3/x^5, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{x^5} dx = \int \frac{(a + b \operatorname{atanh}(cx))^3}{x^5} dx$$

input

```
int((a + b*atanh(c*x))^3/x^5,x)
```

output

```
int((a + b*atanh(c*x))^3/x^5, x)
```


3.35 $\int (dx)^{5/2}(a + \operatorname{barctanh}(cx)) dx$

Optimal result	370
Mathematica [A] (verified)	370
Rubi [A] (verified)	371
Maple [A] (verified)	374
Fricas [A] (verification not implemented)	374
Sympy [F]	375
Maxima [A] (verification not implemented)	375
Giac [F]	376
Mupad [F(-1)]	376
Reduce [B] (verification not implemented)	376

Optimal result

Integrand size = 16, antiderivative size = 124

$$\int (dx)^{5/2}(a + \operatorname{barctanh}(cx)) dx = \frac{4bd^2\sqrt{dx}}{7c^3} + \frac{4b(dx)^{5/2}}{35c} - \frac{2bd^{5/2} \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{7c^{7/2}} + \frac{2(dx)^{7/2}(a + \operatorname{barctanh}(cx))}{7d} - \frac{2bd^{5/2}\operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{7c^{7/2}}$$

output

```
4/7*b*d^2*(d*x)^(1/2)/c^3+4/35*b*(d*x)^(5/2)/c-2/7*b*d^(5/2)*arctan(c^(1/2)
)*(d*x)^(1/2)/d^(1/2))/c^(7/2)+2/7*(d*x)^(7/2)*(a+b*arctanh(c*x))/d-2/7*b*
d^(5/2)*arctanh(c^(1/2)*(d*x)^(1/2)/d^(1/2))/c^(7/2)
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.03

$$\int (dx)^{5/2}(a + \operatorname{barctanh}(cx)) dx = \frac{(dx)^{5/2} (20b\sqrt{c}\sqrt{x} + 4bc^{5/2}x^{5/2} + 10ac^{7/2}x^{7/2} - 10b \arctan(\sqrt{c}\sqrt{x}) + 10bc^{7/2}x^{7/2}\operatorname{arctanh}(\sqrt{c}\sqrt{x}))}{35c^{7/2}x^{5/2}}$$

input

```
Integrate[(d*x)^(5/2)*(a + b*ArcTanh[c*x]), x]
```

output

```
((d*x)^(5/2)*(20*b*Sqrt[c]*Sqrt[x] + 4*b*c^(5/2)*x^(5/2) + 10*a*c^(7/2)*x^(7/2) - 10*b*ArcTan[Sqrt[c]*Sqrt[x]] + 10*b*c^(7/2)*x^(7/2)*ArcTanh[c*x] + 5*b*Log[1 - Sqrt[c]*Sqrt[x]] - 5*b*Log[1 + Sqrt[c]*Sqrt[x]])/(35*c^(7/2)*x^(5/2))
```

Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.15, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {6464, 262, 262, 266, 756, 218, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (dx)^{5/2} (a + \text{barctanh}(cx)) dx \\
 & \quad \downarrow \text{6464} \\
 & \frac{2(dx)^{7/2} (a + \text{barctanh}(cx))}{7d} - \frac{2bc \int \frac{(dx)^{7/2}}{1-c^2x^2} dx}{7d} \\
 & \quad \downarrow \text{262} \\
 & \frac{2(dx)^{7/2} (a + \text{barctanh}(cx))}{7d} - \frac{2bc \left(\frac{d^2 \int \frac{(dx)^{3/2}}{1-c^2x^2} dx}{c^2} - \frac{2d(dx)^{5/2}}{5c^2} \right)}{7d} \\
 & \quad \downarrow \text{262} \\
 & \frac{2(dx)^{7/2} (a + \text{barctanh}(cx))}{7d} - \frac{2bc \left(\frac{d^2 \left(\frac{d^2 \int \frac{1}{\sqrt{dx}(1-c^2x^2)} dx}{c^2} - \frac{2d\sqrt{dx}}{c^2} \right)}{c^2} - \frac{2d(dx)^{5/2}}{5c^2} \right)}{7d} \\
 & \quad \downarrow \text{266} \\
 & \frac{2(dx)^{7/2} (a + \text{barctanh}(cx))}{7d} - \frac{2bc \left(\frac{d^2 \left(\frac{2d \int \frac{1}{1-c^2x^2} d\sqrt{dx}}{c^2} - \frac{2d\sqrt{dx}}{c^2} \right)}{c^2} - \frac{2d(dx)^{5/2}}{5c^2} \right)}{7d}
 \end{aligned}$$

$$\frac{2(dx)^{7/2}(a + \operatorname{barctanh}(cx))}{7d} - \frac{2bc \left(\frac{d^2 \left(\frac{2d \left(\frac{1}{2} d \int \frac{1}{d-cdx} d\sqrt{dx} + \frac{1}{2} d \int \frac{1}{cx+d} d\sqrt{dx} \right) - \frac{2d\sqrt{dx}}{c^2} \right)}{c^2} \right)}{7d} - \frac{2d(dx)^{5/2}}{5c^2} \right)}{7d}$$

$$\frac{2(dx)^{7/2}(a + \operatorname{barctanh}(cx))}{7d} - \frac{2bc \left(\frac{d^2 \left(\frac{2d \left(\frac{1}{2} d \int \frac{1}{d-cdx} d\sqrt{dx} + \frac{\sqrt{d} \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt{c}} \right)}{c^2} - \frac{2d\sqrt{dx}}{c^2} \right)}{c^2} \right)}{7d} - \frac{2d(dx)^{5/2}}{5c^2} \right)}{7d}$$

$$\frac{2(dx)^{7/2}(a + \operatorname{barctanh}(cx))}{7d} - \frac{2bc \left(\frac{d^2 \left(\frac{2d \left(\frac{\sqrt{d} \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt{c}} + \frac{\sqrt{d} \operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt{c}} \right)}{c^2} - \frac{2d\sqrt{dx}}{c^2} \right)}{c^2} \right)}{7d} - \frac{2d(dx)^{5/2}}{5c^2} \right)}{7d}$$

input `Int[(d*x)^(5/2)*(a + b*ArcTanh[c*x]),x]`

output `(2*(d*x)^(7/2)*(a + b*ArcTanh[c*x]))/(7*d) - (2*b*c*((-2*d*(d*x)^(5/2))/(5*c^2) + (d^2*((-2*d*sqrt[d*x])/c^2 + (2*d*((sqrt[d]*ArcTan[(sqrt[c]*sqrt[d*x])/sqrt[d]])/sqrt[d]))/(2*sqrt[c]) + (sqrt[d]*ArcTanh[(sqrt[c]*sqrt[d*x])/sqrt[d]])/(2*sqrt[c])))/c^2))/c^2)/(7*d)`

Definitions of rubi rules used

rule 218 $\text{Int}[\{(a_)+(b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

rule 221 $\text{Int}[\{(a_)+(b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$

rule 262 $\text{Int}[\{(c_)*(x_)\}^{(m_)}*\{(a_)+(b_)*(x_)^2\}^{(p_)}, x_Symbol] \rightarrow \text{Simp}[c*(c*x)^{(m-1)}*\{(a+b*x^2)^{(p+1)}/(b*(m+2*p+1))\}, x] - \text{Simp}[a*c^2*((m-1)/(b*(m+2*p+1))) \ \text{Int}[(c*x)^{(m-2)}*(a+b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{GtQ}[m, 2-1] \ \&\& \ \text{NeQ}[m+2*p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 266 $\text{Int}[\{(c_)*(x_)\}^{(m_)}*\{(a_)+(b_)*(x_)^2\}^{(p_)}, x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \ \text{Subst}[\text{Int}[x^{k*(m+1)-1}*(a+b*(x^{2*k}/c^2))^p, x], x, (c*x)^{(1/k)}], x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 756 $\text{Int}[\{(a_)+(b_)*(x_)^4\}^{-1}, x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[r/(2*a) \ \text{Int}[1/(r-s*x^2), x], x] + \text{Simp}[r/(2*a) \ \text{Int}[1/(r+s*x^2), x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ !\text{GtQ}[a/b, 0]$

rule 6464 $\text{Int}[\{(a_)+\text{ArcTanh}[(c_)*(x_)^{(n_)}]\}*(b_)*\{(d_)*(x_)\}^{(m_)}, x_Symbol] \rightarrow \text{Simp}[(d*x)^{(m+1)}*\{(a+b*\text{ArcTanh}[c*x^n])/(d*(m+1))\}, x] - \text{Simp}[b*c*(n/(d^n*(m+1))) \ \text{Int}[(d*x)^{(m+n)}/(1-c^2*x^{2*n}), x], x] /; \text{FreeQ}[\{a, b, c, d, m, n\}, x] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 1.17 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.86

method	result
derivativedivides	$\frac{\frac{2a(dx)^{\frac{7}{2}}}{7} + \frac{2b(dx)^{\frac{7}{2}} \operatorname{arctanh}(cx)}{7} + \frac{4bd(dx)^{\frac{5}{2}}}{35c} + \frac{4bd^3\sqrt{dx}}{7c^3} - \frac{2bd^4 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{7c^3\sqrt{cd}} - \frac{2bd^4 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{7c^3\sqrt{cd}}}{d}$
default	$\frac{\frac{2a(dx)^{\frac{7}{2}}}{7} + \frac{2b(dx)^{\frac{7}{2}} \operatorname{arctanh}(cx)}{7} + \frac{4bd(dx)^{\frac{5}{2}}}{35c} + \frac{4bd^3\sqrt{dx}}{7c^3} - \frac{2bd^4 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{7c^3\sqrt{cd}} - \frac{2bd^4 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{7c^3\sqrt{cd}}}{d}$
parts	$\frac{2a(dx)^{\frac{7}{2}}}{7d} + \frac{2b(dx)^{\frac{7}{2}} \operatorname{arctanh}(cx)}{7d} + \frac{4b(dx)^{\frac{5}{2}}}{35c} + \frac{4bd^2\sqrt{dx}}{7c^3} - \frac{2bd^3 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{7c^3\sqrt{cd}} - \frac{2bd^3 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{7c^3\sqrt{cd}}$

input `int((d*x)^(5/2)*(a+b*arctanh(c*x)),x,method=_RETURNVERBOSE)`

output `2/d*(1/7*a*(d*x)^(7/2)+1/7*b*(d*x)^(7/2)*arctanh(c*x)+2/35*b*d/c*(d*x)^(5/2)+2/7*b*d^3/c^3*(d*x)^(1/2)-1/7*b*d^4/c^3/(c*d)^(1/2)*arctanh(c*(d*x)^(1/2)/(c*d)^(1/2))-1/7*b*d^4/c^3/(c*d)^(1/2)*arctan(c*(d*x)^(1/2)/(c*d)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 296, normalized size of antiderivative = 2.39

$$\int (dx)^{5/2} (a + b \operatorname{arctanh}(cx)) dx = \left[\frac{10bd^2 \sqrt{\frac{d}{c}} \operatorname{arctan}\left(\frac{\sqrt{dxc}\sqrt{\frac{d}{c}}}{d}\right) - 5bd^2 \sqrt{\frac{d}{c}} \log\left(\frac{cdx - 2\sqrt{dxc}\sqrt{\frac{d}{c}} + d}{cx - 1}\right) - (5bc^3d^2x^3 \log(\dots))}{35c^3} \right]$$

input `integrate((d*x)^(5/2)*(a+b*arctanh(c*x)),x, algorithm="fricas")`

output

```
[-1/35*(10*b*d^2*sqrt(d/c)*arctan(sqrt(d*x)*c*sqrt(d/c)/d) - 5*b*d^2*sqrt(d/c)*log((c*d*x - 2*sqrt(d*x)*c*sqrt(d/c) + d)/(c*x - 1)) - (5*b*c^3*d^2*x^3*log(-(c*x + 1)/(c*x - 1)) + 10*a*c^3*d^2*x^3 + 4*b*c^2*d^2*x^2 + 20*b*d^2)*sqrt(d*x))/c^3, 1/35*(10*b*d^2*sqrt(-d/c)*arctan(sqrt(d*x)*c*sqrt(-d/c)/d) + 5*b*d^2*sqrt(-d/c)*log((c*d*x - 2*sqrt(d*x)*c*sqrt(-d/c) - d)/(c*x + 1)) + (5*b*c^3*d^2*x^3*log(-(c*x + 1)/(c*x - 1)) + 10*a*c^3*d^2*x^3 + 4*b*c^2*d^2*x^2 + 20*b*d^2)*sqrt(d*x))/c^3]
```

Sympy [F]

$$\int (dx)^{5/2} (a + b \operatorname{arctanh}(cx)) dx = \int (dx)^{5/2} (a + b \operatorname{atanh}(cx)) dx$$

input

```
integrate((d*x)**(5/2)*(a+b*atanh(c*x)),x)
```

output

```
Integral((d*x)**(5/2)*(a + b*atanh(c*x)), x)
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.08

$$\int (dx)^{5/2} (a + b \operatorname{arctanh}(cx)) dx = \frac{10 (dx)^{7/2} a + \left(10 (dx)^{7/2} \operatorname{artanh}(cx) - \frac{\left(\frac{10 d^5 \operatorname{arctan}\left(\frac{\sqrt{dxc}}{\sqrt{cd}}\right)}{\sqrt{cdc^4}} - \frac{5 d^5 \log\left(\frac{\sqrt{dxc}-\sqrt{cd}}{\sqrt{dxc}+\sqrt{cd}}\right)}{\sqrt{cdc^4}} - \frac{4 \left((dx)^{5/2} c^2 d^2 + 5 \sqrt{dxc}\right)}{c^4} \right)}{d}}{35 d}$$

input

```
integrate((d*x)^(5/2)*(a+b*arctanh(c*x)),x, algorithm="maxima")
```

output

```
1/35*(10*(d*x)^(7/2)*a + (10*(d*x)^(7/2)*arctanh(c*x) - (10*d^5*arctan(sqrt(d*x)*c/sqrt(c*d))/(sqrt(c*d)*c^4) - 5*d^5*log((sqrt(d*x)*c - sqrt(c*d))/(sqrt(d*x)*c + sqrt(c*d)))/(sqrt(c*d)*c^4) - 4*((d*x)^(5/2)*c^2*d^2 + 5*sqrt(d*x)*d^4)/c^4)*c/d)*b)/d
```

Giac [F]

$$\int (dx)^{5/2} (a + b \operatorname{arctanh}(cx)) dx = \int (dx)^{5/2} (b \operatorname{artanh}(cx) + a) dx$$

input

```
integrate((d*x)^(5/2)*(a+b*arctanh(c*x)),x, algorithm="giac")
```

output

```
integrate((d*x)^(5/2)*(b*arctanh(c*x) + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int (dx)^{5/2} (a + b \operatorname{arctanh}(cx)) dx = \int (a + b \operatorname{atanh}(cx)) (dx)^{5/2} dx$$

input

```
int((a + b*atanh(c*x))*(d*x)^(5/2),x)
```

output

```
int((a + b*atanh(c*x))*(d*x)^(5/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.81

$$\int (dx)^{5/2} (a + b \operatorname{arctanh}(cx)) dx = \frac{\sqrt{d} d^2 \left(-10\sqrt{c} \operatorname{atan}\left(\frac{\sqrt{x}c}{\sqrt{c}}\right) b + 10\sqrt{c} \operatorname{atanh}(cx) b + 10\sqrt{x} \operatorname{atanh}(cx) b c^4 x^3 + 10\sqrt{c} \log\left(\frac{\sqrt{d}(\sqrt{x}c - \sqrt{c})}{\sqrt{d}(\sqrt{x}c + \sqrt{c})}\right) b \right)}{35c^4}$$

input `int((d*x)^(5/2)*(a+b*atanh(c*x)),x)`

output `(sqrt(d)*d**2*(- 10*sqrt(c)*atan((sqrt(x)*c)/sqrt(c))*b + 10*sqrt(c)*atan
h(c*x)*b + 10*sqrt(x)*atanh(c*x)*b*c**4*x**3 + 10*sqrt(c)*log(sqrt(x)*sqrt
(c) - 1)*b - 5*sqrt(c)*log(c*x + 1)*b + 10*sqrt(x)*a*c**4*x**3 + 4*sqrt(x)
*b*c**3*x**2 + 20*sqrt(x)*b*c))/(35*c**4)`

3.36 $\int (dx)^{3/2}(a + b \operatorname{arctanh}(cx)) dx$

Optimal result	378
Mathematica [A] (verified)	378
Rubi [A] (verified)	379
Maple [A] (verified)	381
Fricas [A] (verification not implemented)	382
Sympy [F]	382
Maxima [A] (verification not implemented)	383
Giac [F]	383
Mupad [F(-1)]	384
Reduce [B] (verification not implemented)	384

Optimal result

Integrand size = 16, antiderivative size = 106

$$\int (dx)^{3/2}(a + b \operatorname{arctanh}(cx)) dx = \frac{4b(dx)^{3/2}}{15c} + \frac{2bd^{3/2} \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{5c^{5/2}} + \frac{2(dx)^{5/2}(a + b \operatorname{arctanh}(cx))}{5d} - \frac{2bd^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{5c^{5/2}}$$

output

```
4/15*b*(d*x)^(3/2)/c+2/5*b*d^(3/2)*arctan(c^(1/2)*(d*x)^(1/2)/d^(1/2))/c^(5/2)+2/5*(d*x)^(5/2)*(a+b*arctanh(c*x))/d-2/5*b*d^(3/2)*arctanh(c^(1/2)*(d*x)^(1/2)/d^(1/2))/c^(5/2)
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.08

$$\int (dx)^{3/2}(a + b \operatorname{arctanh}(cx)) dx = \frac{(dx)^{3/2} (4bc^{3/2}x^{3/2} + 6ac^{5/2}x^{5/2} + 6b \arctan(\sqrt{c}\sqrt{x}) + 6bc^{5/2}x^{5/2} \operatorname{arctanh}(cx) + 3b \log(\dots))}{15c^{5/2}x^{3/2}}$$

input

```
Integrate[(d*x)^(3/2)*(a + b*ArcTanh[c*x]),x]
```

output

$$\left((dx)^{3/2} (4bc^{3/2}x^{3/2} + 6ac^{5/2}x^{5/2} + 6b\text{ArcTan}[\text{Sqrt}[c]\text{Sqrt}[x]] + 6bc^{5/2}x^{5/2}\text{ArcTanh}[cx] + 3b\text{Log}[1 - \text{Sqrt}[c]\text{Sqrt}[x]] - 3b\text{Log}[1 + \text{Sqrt}[c]\text{Sqrt}[x]]) \right) / (15c^{5/2}x^{3/2})$$
Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.16, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {6464, 262, 266, 27, 827, 218, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (dx)^{3/2} (a + b \operatorname{arctanh}(cx)) dx \\ & \quad \downarrow \text{6464} \\ & \frac{2(dx)^{5/2} (a + b \operatorname{arctanh}(cx))}{5d} - \frac{2bc \int \frac{(dx)^{5/2}}{1-c^2x^2} dx}{5d} \\ & \quad \downarrow \text{262} \\ & \frac{2(dx)^{5/2} (a + b \operatorname{arctanh}(cx))}{5d} - \frac{2bc \left(\frac{d^2 \int \frac{\sqrt{dx}}{1-c^2x^2} dx}{c^2} - \frac{2d(dx)^{3/2}}{3c^2} \right)}{5d} \\ & \quad \downarrow \text{266} \\ & \frac{2(dx)^{5/2} (a + b \operatorname{arctanh}(cx))}{5d} - \frac{2bc \left(\frac{2d \int \frac{d^3x}{d^2-c^2d^2x^2} d\sqrt{dx}}{c^2} - \frac{2d(dx)^{3/2}}{3c^2} \right)}{5d} \\ & \quad \downarrow \text{27} \\ & \frac{2(dx)^{5/2} (a + b \operatorname{arctanh}(cx))}{5d} - \frac{2bc \left(\frac{2d^3 \int \frac{dx}{d^2-c^2d^2x^2} d\sqrt{dx}}{c^2} - \frac{2d(dx)^{3/2}}{3c^2} \right)}{5d} \\ & \quad \downarrow \text{827} \end{aligned}$$

$$\frac{2(dx)^{5/2}(a + \operatorname{barctanh}(cx))}{5d} - \frac{2bc \left(\frac{2d^3 \left(\frac{\int \frac{1}{d-cdx} d\sqrt{dx}}{2c} - \frac{\int \frac{1}{cx+d} d\sqrt{dx}}{2c} \right)}{c^2} - \frac{2d(dx)^{3/2}}{3c^2} \right)}{5d}$$

↓ 218

$$\frac{2(dx)^{5/2}(a + \operatorname{barctanh}(cx))}{5d} - \frac{2bc \left(\frac{2d^3 \left(\frac{\int \frac{1}{d-cdx} d\sqrt{dx}}{2c} - \frac{\arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/2}\sqrt{d}} \right)}{c^2} - \frac{2d(dx)^{3/2}}{3c^2} \right)}{5d}$$

↓ 221

$$\frac{2(dx)^{5/2}(a + \operatorname{barctanh}(cx))}{5d} - \frac{2bc \left(\frac{2d^3 \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/2}\sqrt{d}} - \frac{\arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/2}\sqrt{d}} \right)}{c^2} - \frac{2d(dx)^{3/2}}{3c^2} \right)}{5d}$$

input `Int[(d*x)^(3/2)*(a + b*ArcTanh[c*x]), x]`

output `(2*(d*x)^(5/2)*(a + b*ArcTanh[c*x]))/(5*d) - (2*b*c*((-2*d*(d*x)^(3/2))/(3*c^2) + (2*d^3*(-1/2*ArcTan[(Sqrt[c]*Sqrt[d*x])/Sqrt[d]]/(c^(3/2)*Sqrt[d]) + ArcTanh[(Sqrt[c]*Sqrt[d*x])/Sqrt[d]]/(2*c^(3/2)*Sqrt[d]))/c^2))/(5*d)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 262 $\text{Int}[(c \cdot x)^m \cdot (a + b \cdot x^2)^p, x_Symbol] \rightarrow \text{Simp}[c \cdot (c \cdot x)^{m-1} \cdot (a + b \cdot x^2)^{p+1} / (b \cdot (m + 2 \cdot p + 1))], x] - \text{Simp}[a \cdot c^2 \cdot (m - 1) / (b \cdot (m + 2 \cdot p + 1)) \text{Int}[(c \cdot x)^{m-2} \cdot (a + b \cdot x^2)^p, x], x] /;$ FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2 * p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]

rule 266 $\text{Int}[(c \cdot x)^m \cdot (a + b \cdot x^2)^p, x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \text{Subst}[\text{Int}[x^{k \cdot (m + 1) - 1} \cdot (a + b \cdot x^{2 \cdot k}) / c^2]^p, x], x, (c \cdot x)^{1/k}], x] /;$ FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]

rule 827 $\text{Int}[x^2 / (a + b \cdot x^4), x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[s / (2 \cdot b) \text{Int}[1 / (r + s \cdot x^2), x], x] - \text{Simp}[s / (2 \cdot b) \text{Int}[1 / (r - s \cdot x^2), x], x] /;$ FreeQ[{a, b}, x] && !GtQ[a/b, 0]

rule 6464 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n]) \cdot b \cdot (d \cdot x)^m, x_Symbol] \rightarrow \text{Simp}[(d \cdot x)^{m+1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n]) / (d \cdot (m + 1))], x] - \text{Simp}[b \cdot c \cdot n / (d^n \cdot (m + 1)) \text{Int}[(d \cdot x)^{m+n} / (1 - c^2 \cdot x^{2 \cdot n}), x], x] /;$ FreeQ[{a, b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]

Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{\frac{2(dx)^{\frac{5}{2}}a}{5} + \frac{2(dx)^{\frac{5}{2}}b \operatorname{arctanh}(cx)}{5} + \frac{4bd(dx)^{\frac{3}{2}}}{15c} - \frac{2bd^3 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{5c^2\sqrt{cd}} + \frac{2bd^3 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{5c^2\sqrt{cd}}}{d}$	93
default	$\frac{\frac{2(dx)^{\frac{5}{2}}a}{5} + \frac{2(dx)^{\frac{5}{2}}b \operatorname{arctanh}(cx)}{5} + \frac{4bd(dx)^{\frac{3}{2}}}{15c} - \frac{2bd^3 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{5c^2\sqrt{cd}} + \frac{2bd^3 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{5c^2\sqrt{cd}}}{d}$	93
parts	$\frac{2a(dx)^{\frac{5}{2}}}{5d} + \frac{2b(dx)^{\frac{5}{2}} \operatorname{arctanh}(cx)}{5d} + \frac{4b(dx)^{\frac{3}{2}}}{15c} - \frac{2bd^2 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{5c^2\sqrt{cd}} + \frac{2bd^2 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{5c^2\sqrt{cd}}$	93

input `int((d*x)^(3/2)*(a+b*arctanh(c*x)),x,method=_RETURNVERBOSE)`

output

```
2/d*(1/5*(d*x)^(5/2)*a+1/5*(d*x)^(5/2)*b*arctanh(c*x)+2/15*b*d*(d*x)^(3/2)
/c-1/5*b*d^3/c^2/(c*d)^(1/2)*arctanh(c*(d*x)^(1/2)/(c*d)^(1/2))+1/5*b*d^3/
c^2/(c*d)^(1/2)*arctan(c*(d*x)^(1/2)/(c*d)^(1/2)))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 255, normalized size of antiderivative = 2.41

$$\int (dx)^{3/2} (a + b \operatorname{arctanh}(cx)) dx = \frac{6bd\sqrt{\frac{d}{c}} \arctan\left(\frac{\sqrt{dxc}\sqrt{\frac{d}{c}}}{d}\right) + 3bd\sqrt{\frac{d}{c}} \log\left(\frac{cdx-2\sqrt{dxc}\sqrt{\frac{d}{c}}+d}{cx-1}\right) + (3bc^2 dx^2 \log\left(-\frac{cx+1}{cx-1}\right))}{15c^2}$$

input

```
integrate((d*x)^(3/2)*(a+b*arctanh(c*x)),x, algorithm="fricas")
```

output

```
[1/15*(6*b*d*sqrt(d/c)*arctan(sqrt(d*x)*c*sqrt(d/c)/d) + 3*b*d*sqrt(d/c)*1
og((c*d*x - 2*sqrt(d*x)*c*sqrt(d/c) + d)/(c*x - 1)) + (3*b*c^2*d*x^2*log(-
(c*x + 1)/(c*x - 1)) + 6*a*c^2*d*x^2 + 4*b*c*d*x)*sqrt(d*x))/c^2, 1/15*(6*
b*d*sqrt(-d/c)*arctan(sqrt(d*x)*c*sqrt(-d/c)/d) + 3*b*d*sqrt(-d/c)*log((c*
d*x + 2*sqrt(d*x)*c*sqrt(-d/c) - d)/(c*x + 1)) + (3*b*c^2*d*x^2*log(-(c*x
+ 1)/(c*x - 1)) + 6*a*c^2*d*x^2 + 4*b*c*d*x)*sqrt(d*x))/c^2]
```

Sympy [F]

$$\int (dx)^{3/2} (a + b \operatorname{arctanh}(cx)) dx = \int (dx)^{\frac{3}{2}} (a + b \operatorname{atanh}(cx)) dx$$

input

```
integrate((d*x)**(3/2)*(a+b*atanh(c*x)),x)
```

output

```
Integral((d*x)**(3/2)*(a + b*atanh(c*x)), x)
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.11

$$\int (dx)^{3/2} (a + b \operatorname{arctanh}(cx)) dx = \frac{6(dx)^{\frac{5}{2}} a + \left(6(dx)^{\frac{5}{2}} \operatorname{artanh}(cx) + \frac{\left(\frac{4(dx)^{\frac{3}{2}} d^2}{c^2} + \frac{6d^4 \arctan\left(\frac{\sqrt{dxc}}{\sqrt{cd}}\right)}{\sqrt{cd}c^3} + \frac{3d^4 \log\left(\frac{\sqrt{dxc}-\sqrt{cd}}{\sqrt{dxc}+\sqrt{cd}}\right)}{\sqrt{cd}c^3} \right) c}{d} \right) b}{15d}$$

input `integrate((d*x)^(3/2)*(a+b*arctanh(c*x)),x, algorithm="maxima")`output `1/15*(6*(d*x)^(5/2)*a + (6*(d*x)^(5/2)*arctanh(c*x) + (4*(d*x)^(3/2)*d^2/c^2 + 6*d^4*arctan(sqrt(d*x)*c/sqrt(c*d))/(sqrt(c*d)*c^3) + 3*d^4*log((sqrt(d*x)*c - sqrt(c*d))/(sqrt(d*x)*c + sqrt(c*d)))/(sqrt(c*d)*c^3))*c/d)*b/d`**Giac [F]**

$$\int (dx)^{3/2} (a + b \operatorname{arctanh}(cx)) dx = \int (dx)^{\frac{3}{2}} (b \operatorname{artanh}(cx) + a) dx$$

input `integrate((d*x)^(3/2)*(a+b*arctanh(c*x)),x, algorithm="giac")`output `integrate((d*x)^(3/2)*(b*arctanh(c*x) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int (dx)^{3/2} (a + b \operatorname{arctanh}(cx)) dx = \int (a + b \operatorname{atanh}(cx)) (dx)^{3/2} dx$$

input `int((a + b*atanh(c*x))*(d*x)^(3/2), x)`output `int((a + b*atanh(c*x))*(d*x)^(3/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.86

$$\int (dx)^{3/2} (a + b \operatorname{arctanh}(cx)) dx = \frac{\sqrt{d} d \left(6\sqrt{c} \operatorname{atan}\left(\frac{\sqrt{x}c}{\sqrt{c}}\right) b + 6\sqrt{c} \operatorname{atanh}(cx) b + 6\sqrt{x} \operatorname{atanh}(cx) b c^3 x^2 + 6\sqrt{c} \log(\sqrt{x} \sqrt{c}) b \right)}{15c^3}$$

input `int((d*x)^(3/2)*(a+b*atanh(c*x)), x)`output `(sqrt(d)*d*(6*sqrt(c)*atan((sqrt(x)*c)/sqrt(c))*b + 6*sqrt(c)*atanh(c*x)*b + 6*sqrt(x)*atanh(c*x)*b*c**3*x**2 + 6*sqrt(c)*log(sqrt(x)*sqrt(c) - 1)*b - 3*sqrt(c)*log(c*x + 1)*b + 6*sqrt(x)*a*c**3*x**2 + 4*sqrt(x)*b*c**2*x) / (15*c**3)`

3.37 $\int \sqrt{dx}(a + b \operatorname{arctanh}(cx)) dx$

Optimal result	385
Mathematica [A] (verified)	385
Rubi [A] (verified)	386
Maple [A] (verified)	388
Fricas [A] (verification not implemented)	389
Sympy [F]	389
Maxima [A] (verification not implemented)	390
Giac [F]	390
Mupad [F(-1)]	391
Reduce [B] (verification not implemented)	391

Optimal result

Integrand size = 16, antiderivative size = 106

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx)) dx = \frac{4b\sqrt{dx}}{3c} - \frac{2b\sqrt{d} \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{3c^{3/2}} + \frac{2(dx)^{3/2}(a + b \operatorname{arctanh}(cx))}{3d} - \frac{2b\sqrt{d} \operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{3c^{3/2}}$$

output

```
4/3*b*(d*x)^(1/2)/c-2/3*b*d^(1/2)*arctan(c^(1/2)*(d*x)^(1/2)/d^(1/2))/c^(3/2)+2/3*(d*x)^(3/2)*(a+b*arctanh(c*x))/d-2/3*b*d^(1/2)*arctanh(c^(1/2)*(d*x)^(1/2)/d^(1/2))/c^(3/2)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.08

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx)) dx = \frac{\sqrt{dx}(4b\sqrt{c}\sqrt{x} + 2ac^{3/2}x^{3/2} - 2b \arctan(\sqrt{c}\sqrt{x}) + 2bc^{3/2}x^{3/2} \operatorname{arctanh}(cx) + b \log(1 - \sqrt{c}\sqrt{x}) - b \log(1 + \sqrt{c}\sqrt{x}))}{3c^{3/2}\sqrt{x}}$$

input

```
Integrate[Sqrt[d*x]*(a + b*ArcTanh[c*x]),x]
```

output

$$\frac{(\text{Sqrt}[d*x]*(4*b*\text{Sqrt}[c]*\text{Sqrt}[x] + 2*a*c^{(3/2)}*x^{(3/2)} - 2*b*\text{ArcTan}[\text{Sqrt}[c]*\text{Sqrt}[x]] + 2*b*c^{(3/2)}*x^{(3/2)}*\text{ArcTanh}[c*x] + b*\text{Log}[1 - \text{Sqrt}[c]*\text{Sqrt}[x]] - b*\text{Log}[1 + \text{Sqrt}[c]*\text{Sqrt}[x]])}{(3*c^{(3/2)}*\text{Sqrt}[x])}$$
Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6464, 262, 266, 756, 218, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx)) dx$$

$$\downarrow 6464$$

$$\frac{2(dx)^{3/2}(a + b \operatorname{arctanh}(cx))}{3d} - \frac{2bc \int \frac{(dx)^{3/2}}{1-c^2x^2} dx}{3d}$$

$$\downarrow 262$$

$$\frac{2(dx)^{3/2}(a + b \operatorname{arctanh}(cx))}{3d} - \frac{2bc \left(\frac{d^2 \int \frac{1}{\sqrt{dx}(1-c^2x^2)} dx}{c^2} - \frac{2d\sqrt{dx}}{c^2} \right)}{3d}$$

$$\downarrow 266$$

$$\frac{2(dx)^{3/2}(a + b \operatorname{arctanh}(cx))}{3d} - \frac{2bc \left(\frac{2d \int \frac{1}{1-c^2x^2} d\sqrt{dx}}{c^2} - \frac{2d\sqrt{dx}}{c^2} \right)}{3d}$$

$$\downarrow 756$$

$$\frac{2(dx)^{3/2}(a + b \operatorname{arctanh}(cx))}{3d} - \frac{2bc \left(\frac{2d \left(\frac{1}{2} d \int \frac{1}{d-cdx} d\sqrt{dx} + \frac{1}{2} d \int \frac{1}{cx+d} d\sqrt{dx} \right)}{c^2} - \frac{2d\sqrt{dx}}{c^2} \right)}{3d}$$

$$\downarrow 218$$

$$\frac{2(dx)^{3/2}(a + b\operatorname{arctanh}(cx))}{3d} - \frac{2bc \left(\frac{2d \left(\frac{1}{2}d \int \frac{1}{d-cx} d\sqrt{dx} + \frac{\sqrt{d} \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt{c}} \right)}{c^2} - \frac{2d\sqrt{dx}}{c^2} \right)}{3d}$$

↓ 221

$$\frac{2(dx)^{3/2}(a + b\operatorname{arctanh}(cx))}{3d} - \frac{2bc \left(\frac{2d \left(\frac{\sqrt{d} \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt{c}} + \frac{\sqrt{d}\operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt{c}} \right)}{c^2} - \frac{2d\sqrt{dx}}{c^2} \right)}{3d}$$

input `Int[Sqrt[d*x]*(a + b*ArcTanh[c*x]), x]`

output `(2*(d*x)^(3/2)*(a + b*ArcTanh[c*x]))/(3*d) - (2*b*c*((-2*d*Sqrt[d*x])/c^2 + (2*d*((Sqrt[d]*ArcTan[(Sqrt[c]*Sqrt[d*x])/Sqrt[d]])/(2*Sqrt[c]) + (Sqrt[d]*ArcTanh[(Sqrt[c]*Sqrt[d*x])/Sqrt[d]])/(2*Sqrt[c])))/c^2)/(3*d)`

Defintions of rubi rules used

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 262 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m-1)*((a + b*x^2)^(p+1)/(b*(m+2*p+1))), x] - Simp[a*c^2*(m-1)/(b*(m+2*p+1)) Int[(c*x)^(m-2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2-1] && NeQ[m+2*p+1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2)) ^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 756 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 6464 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_)), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.84

method	result	size
parts	$\frac{2a(dx)^{\frac{3}{2}}}{3d} + \frac{2b(dx)^{\frac{3}{2}} \operatorname{arctanh}(cx)}{3d} + \frac{4bd\sqrt{dx}}{3c} - \frac{2bd \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{3c\sqrt{cd}} - \frac{2bd \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{3c\sqrt{cd}}$	89
derivativedivides	$\frac{\frac{2(dx)^{\frac{3}{2}}a}{3} + \frac{2(dx)^{\frac{3}{2}}b \operatorname{arctanh}(cx)}{3} + \frac{4bd\sqrt{dx}}{3c} - \frac{2bd^2 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{3c\sqrt{cd}} - \frac{2bd^2 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{3c\sqrt{cd}}}{d}$	93
default	$\frac{\frac{2(dx)^{\frac{3}{2}}a}{3} + \frac{2(dx)^{\frac{3}{2}}b \operatorname{arctanh}(cx)}{3} + \frac{4bd\sqrt{dx}}{3c} - \frac{2bd^2 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{3c\sqrt{cd}} - \frac{2bd^2 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{3c\sqrt{cd}}}{d}$	93

input `int((d*x)^(1/2)*(a+b*arctanh(c*x)),x,method=_RETURNVERBOSE)`

output $\frac{2}{3}a*(d*x)^{(3/2)}/d+2/3*b/d*(d*x)^{(3/2)}*arctanh(c*x)+4/3*b*(d*x)^{(1/2)}/c-2/3*b*d/c/(c*d)^{(1/2)}*arctanh(c*(d*x)^{(1/2)}/(c*d)^{(1/2)})-2/3*b*d/c/(c*d)^{(1/2)}*arctan(c*(d*x)^{(1/2)}/(c*d)^{(1/2)})$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 223, normalized size of antiderivative = 2.10

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx)) dx$$

$$= \left[\frac{2b\sqrt{\frac{d}{c}} \operatorname{arctan}\left(\frac{\sqrt{dx}c\sqrt{\frac{d}{c}}}{d}\right) - b\sqrt{\frac{d}{c}} \log\left(\frac{cdx - 2\sqrt{dx}c\sqrt{\frac{d}{c}} + d}{cx-1}\right) - (bcx \log\left(-\frac{cx+1}{cx-1}\right) + 2acx + 4b)\sqrt{dx}}{3c}, \frac{2b\sqrt{\frac{d}{c}} \operatorname{arctan}\left(\frac{\sqrt{dx}c\sqrt{\frac{d}{c}}}{d}\right) - b\sqrt{\frac{d}{c}} \log\left(\frac{cdx - 2\sqrt{dx}c\sqrt{\frac{d}{c}} + d}{cx-1}\right) - (bcx \log\left(-\frac{cx+1}{cx-1}\right) + 2acx + 4b)\sqrt{dx}}{3c} \right]$$

input `integrate((d*x)^(1/2)*(a+b*arctanh(c*x)),x, algorithm="fricas")`

output `[-1/3*(2*b*sqrt(d/c)*arctan(sqrt(d*x)*c*sqrt(d/c)/d) - b*sqrt(d/c)*log((c*d*x - 2*sqrt(d*x)*c*sqrt(d/c) + d)/(c*x - 1)) - (b*c*x*log(-(c*x + 1)/(c*x - 1)) + 2*a*c*x + 4*b)*sqrt(d*x))/c, 1/3*(2*b*sqrt(-d/c)*arctan(sqrt(d*x)*c*sqrt(-d/c)/d) + b*sqrt(-d/c)*log((c*d*x - 2*sqrt(d*x)*c*sqrt(-d/c) - d)/(c*x + 1)) + (b*c*x*log(-(c*x + 1)/(c*x - 1)) + 2*a*c*x + 4*b)*sqrt(d*x))/c]`

Sympy [F]

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx)) dx = \int \sqrt{dx}(a + b \operatorname{atanh}(cx)) dx$$

input `integrate((d*x)**(1/2)*(a+b*atanh(c*x)),x)`

output `Integral(sqrt(d*x)*(a + b*atanh(c*x)), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.12

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{2(dx)^{\frac{3}{2}} a + \left(2(dx)^{\frac{3}{2}} \operatorname{artanh}(cx) - \frac{\left(\frac{2d^3 \arctan\left(\frac{\sqrt{dx}c}{\sqrt{cd}}\right)}{\sqrt{cd}c^2} - \frac{d^3 \log\left(\frac{\sqrt{dx}c - \sqrt{cd}}{\sqrt{dx}c + \sqrt{cd}}\right)}{\sqrt{cd}c^2} - \frac{4\sqrt{dx}d^2}{c^2} \right) c}{d} \right) b}{3d}$$

input `integrate((d*x)^(1/2)*(a+b*arctanh(c*x)),x, algorithm="maxima")`output `1/3*(2*(d*x)^(3/2)*a + (2*(d*x)^(3/2)*arctanh(c*x) - (2*d^3*arctan(sqrt(d*x)*c/sqrt(c*d))/(sqrt(c*d)*c^2) - d^3*log((sqrt(d*x)*c - sqrt(c*d))/(sqrt(d*x)*c + sqrt(c*d)))/(sqrt(c*d)*c^2) - 4*sqrt(d*x)*d^2/c^2)*c/d)`**Giac [F]**

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx)) dx = \int \sqrt{dx}(b \operatorname{artanh}(cx) + a) dx$$

input `integrate((d*x)^(1/2)*(a+b*arctanh(c*x)),x, algorithm="giac")`output `integrate(sqrt(d*x)*(b*arctanh(c*x) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx)) dx = \int (a + b \operatorname{atanh}(cx)) \sqrt{dx} dx$$

input `int((a + b*atanh(c*x))*(d*x)^(1/2), x)`

output `int((a + b*atanh(c*x))*(d*x)^(1/2), x)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.78

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{\sqrt{d} \left(-2\sqrt{c} \operatorname{atan}\left(\frac{\sqrt{x}c}{\sqrt{c}}\right) b + 2\sqrt{c} \operatorname{atanh}(cx) b + 2\sqrt{x} \operatorname{atanh}(cx) b c^2 x + 2\sqrt{c} \log(\sqrt{x} \sqrt{c} - 1) b - \sqrt{c} \log(ca) \right)}{3c^2}$$

input `int((d*x)^(1/2)*(a+b*atanh(c*x)), x)`

output `(sqrt(d)*(-2*sqrt(c)*atan((sqrt(x)*c)/sqrt(c))*b + 2*sqrt(c)*atanh(c*x)*
b + 2*sqrt(x)*atanh(c*x)*b*c**2*x + 2*sqrt(c)*log(sqrt(x)*sqrt(c) - 1)*b -
sqrt(c)*log(c*x + 1)*b + 2*sqrt(x)*a*c**2*x + 4*sqrt(x)*b*c)/(3*c**2)`

3.38 $\int \frac{a+b\operatorname{arctanh}(cx)}{\sqrt{dx}} dx$

Optimal result	392
Mathematica [A] (verified)	392
Rubi [A] (verified)	393
Maple [A] (verified)	395
Fricas [A] (verification not implemented)	395
Sympy [F]	396
Maxima [A] (verification not implemented)	396
Giac [A] (verification not implemented)	397
Mupad [F(-1)]	397
Reduce [B] (verification not implemented)	397

Optimal result

Integrand size = 16, antiderivative size = 85

$$\int \frac{a + b\operatorname{arctanh}(cx)}{\sqrt{dx}} dx = \frac{2b \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{c}\sqrt{d}} + \frac{2\sqrt{dx}(a + b\operatorname{arctanh}(cx))}{d} - \frac{2b\operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{c}\sqrt{d}}$$

output

$2*b*\arctan(c^{(1/2)}*(d*x)^{(1/2)}/d^{(1/2)})/c^{(1/2)}/d^{(1/2)}+2*(d*x)^{(1/2)}*(a+b*\operatorname{arctanh}(c*x))/d-2*b*\operatorname{arctanh}(c^{(1/2)}*(d*x)^{(1/2)}/d^{(1/2)})/c^{(1/2)}/d^{(1/2)}$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.15

$$\int \frac{a + b\operatorname{arctanh}(cx)}{\sqrt{dx}} dx = \frac{\sqrt{x}(2a\sqrt{c}\sqrt{x} + 2b \arctan(\sqrt{c}\sqrt{x}) + 2b\sqrt{c}\sqrt{x}\operatorname{arctanh}(cx) + b \log(1 - \sqrt{c}\sqrt{x}) - b \log(1 + \sqrt{c}\sqrt{x}))}{\sqrt{c}\sqrt{dx}}$$

input

`Integrate[(a + b*ArcTanh[c*x])/Sqrt[d*x], x]`

output

```
(Sqrt[x]*(2*a*Sqrt[c]*Sqrt[x] + 2*b*ArcTan[Sqrt[c]*Sqrt[x]] + 2*b*Sqrt[c]*
Sqrt[x]*ArcTanh[c*x] + b*Log[1 - Sqrt[c]*Sqrt[x]] - b*Log[1 + Sqrt[c]*Sqrt
[x]]))/(Sqrt[c]*Sqrt[d*x])
```

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.08, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6464, 266, 27, 827, 218, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}(cx)}{\sqrt{dx}} dx \\
 & \quad \downarrow \text{6464} \\
 & \frac{2\sqrt{dx}(a + b \operatorname{arctanh}(cx))}{d} - \frac{2bc \int \frac{\sqrt{dx}}{1-c^2x^2} dx}{d} \\
 & \quad \downarrow \text{266} \\
 & \frac{2\sqrt{dx}(a + b \operatorname{arctanh}(cx))}{d} - \frac{4bc \int \frac{d^3x}{d^2-c^2d^2x^2} d\sqrt{dx}}{d^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{2\sqrt{dx}(a + b \operatorname{arctanh}(cx))}{d} - 4bc \int \frac{dx}{d^2 - c^2d^2x^2} d\sqrt{dx} \\
 & \quad \downarrow \text{827} \\
 & \frac{2\sqrt{dx}(a + b \operatorname{arctanh}(cx))}{d} - 4bc \left(\frac{\int \frac{1}{d-cdx} d\sqrt{dx}}{2c} - \frac{\int \frac{1}{cx d+d} d\sqrt{dx}}{2c} \right) \\
 & \quad \downarrow \text{218} \\
 & \frac{2\sqrt{dx}(a + b \operatorname{arctanh}(cx))}{d} - 4bc \left(\frac{\int \frac{1}{d-cdx} d\sqrt{dx}}{2c} - \frac{\arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/2}\sqrt{d}} \right) \\
 & \quad \downarrow \text{221}
 \end{aligned}$$

$$\frac{2\sqrt{dx}(a + b\operatorname{arctanh}(cx))}{d} - 4bc \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/2}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/2}\sqrt{d}} \right)$$

input `Int[(a + b*ArcTanh[c*x])/Sqrt[d*x], x]`

output `(2*Sqrt[d*x]*(a + b*ArcTanh[c*x])/d - 4*b*c*(-1/2*ArcTan[(Sqrt[c]*Sqrt[d*x])/Sqrt[d]]/(c^(3/2)*Sqrt[d]) + ArcTanh[(Sqrt[c]*Sqrt[d*x])/Sqrt[d]]/(2*c^(3/2)*Sqrt[d]))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 827 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x], x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 6464

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))*((d_)*(x_)^(m_), x_Symbol] :
> Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n
/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a,
b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.80

method	result	size
derivativedivides	$\frac{2\sqrt{dx} a + 2\sqrt{dx} b \operatorname{arctanh}(cx) - \frac{2bd \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{\sqrt{cd}} + \frac{2bd \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{\sqrt{cd}}}{d}$	68
default	$\frac{2\sqrt{dx} a + 2\sqrt{dx} b \operatorname{arctanh}(cx) - \frac{2bd \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{\sqrt{cd}} + \frac{2bd \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{\sqrt{cd}}}{d}$	68
parts	$\frac{2a\sqrt{dx}}{d} + \frac{2b\sqrt{dx} \operatorname{arctanh}(cx)}{d} - \frac{2b \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{\sqrt{cd}} + \frac{2b \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{\sqrt{cd}}$	70

input

```
int((a+b*arctanh(c*x))/(d*x)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
2/d*((d*x)^(1/2)*a+(d*x)^(1/2)*b*arctanh(c*x)-b*d/(c*d)^(1/2)*arctanh(c*(d
*x)^(1/2)/(c*d)^(1/2))+b*d/(c*d)^(1/2)*arctan(c*(d*x)^(1/2)/(c*d)^(1/2)))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 211, normalized size of antiderivative = 2.48

$$\int \frac{a + b \operatorname{arctanh}(cx)}{\sqrt{dx}} dx$$

$$= \left[-\frac{2\sqrt{cdb} \operatorname{arctan}\left(\frac{\sqrt{cd}\sqrt{dx}}{cdx}\right) - \sqrt{cdb} \log\left(\frac{cdx - 2\sqrt{cd}\sqrt{dx} + d}{cx - 1}\right) - (bc \log\left(-\frac{cx+1}{cx-1}\right) + 2ac)\sqrt{dx}}{cd}, \frac{2\sqrt{-cdb} \operatorname{arctan}\left(\frac{\sqrt{-cd}\sqrt{dx}}{cdx}\right) - \sqrt{-cdb} \log\left(\frac{cdx - 2\sqrt{-cd}\sqrt{dx} + d}{cx - 1}\right) - (bc \log\left(\frac{cx+1}{cx-1}\right) + 2ac)\sqrt{dx}}{cd} \right]$$

input

```
integrate((a+b*arctanh(c*x))/(d*x)^(1/2),x, algorithm="fricas")
```


output

```
[-(2*sqrt(c*d)*b*arctan(sqrt(c*d)*sqrt(d*x)/(c*d*x)) - sqrt(c*d)*b*log((c*
d*x - 2*sqrt(c*d)*sqrt(d*x) + d)/(c*x - 1)) - (b*c*log(-(c*x + 1)/(c*x - 1
)) + 2*a*c)*sqrt(d*x))/(c*d), (2*sqrt(-c*d)*b*arctan(sqrt(-c*d)*sqrt(d*x)/
(c*d*x)) - sqrt(-c*d)*b*log((c*d*x - 2*sqrt(-c*d)*sqrt(d*x) - d)/(c*x + 1
)) + (b*c*log(-(c*x + 1)/(c*x - 1)) + 2*a*c)*sqrt(d*x))/(c*d)]
```

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{\sqrt{dx}} dx = \int \frac{a + b \operatorname{atanh}(cx)}{\sqrt{dx}} dx$$

input

```
integrate((a+b*atanh(c*x))/(d*x)**(1/2),x)
```

output

```
Integral((a + b*atanh(c*x))/sqrt(d*x), x)
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.21

$$\int \frac{a + b \operatorname{arctanh}(cx)}{\sqrt{dx}} dx$$

$$= \frac{\left(2 \sqrt{dx} \operatorname{artanh}(cx) + \frac{\left(\frac{2 d^2 \arctan\left(\frac{\sqrt{dxc}}{\sqrt{cd}}\right)}{\sqrt{cdc}} + \frac{d^2 \log\left(\frac{\sqrt{dxc}-\sqrt{cd}}{\sqrt{dxc}+\sqrt{cd}}\right)}{\sqrt{cdc}} \right) c}{d} \right) b + 2 \sqrt{dxa}}{d}$$

input

```
integrate((a+b*arctanh(c*x))/(d*x)^(1/2),x, algorithm="maxima")
```

output

```
((2*sqrt(d*x)*arctanh(c*x) + (2*d^2*arctan(sqrt(d*x)*c/sqrt(c*d))/(sqrt(c*
d)*c) + d^2*log((sqrt(d*x)*c - sqrt(c*d))/(sqrt(d*x)*c + sqrt(c*d)))/(sqrt
(c*d)*c))*c/d)*b + 2*sqrt(d*x)*a)/d
```

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.04

$$\int \frac{a + b \operatorname{arctanh}(cx)}{\sqrt{dx}} dx$$

$$= \frac{\left(2cd \left(\frac{\arctan\left(\frac{\sqrt{dxc}}{\sqrt{cd}}\right)}{\sqrt{cdc}} + \frac{\arctan\left(\frac{\sqrt{dxc}}{\sqrt{-cd}}\right)}{\sqrt{-cdc}} \right) + \sqrt{dx} \log\left(-\frac{cx+1}{cx-1}\right) \right) b + 2\sqrt{dxa}}{d}$$

input `integrate((a+b*arctanh(c*x))/(d*x)^(1/2),x, algorithm="giac")`output `((2*c*d*(arctan(sqrt(d*x)*c/sqrt(c*d))/(sqrt(c*d)*c) + arctan(sqrt(d*x)*c/sqrt(-c*d))/(sqrt(-c*d)*c)) + sqrt(d*x)*log(-(c*x + 1)/(c*x - 1)))*b + 2*sqrt(d*x)*a)/d`**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx)}{\sqrt{dx}} dx = \int \frac{a + b \operatorname{atanh}(cx)}{\sqrt{dx}} dx$$

input `int((a + b*atanh(c*x))/(d*x)^(1/2),x)`output `int((a + b*atanh(c*x))/(d*x)^(1/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.86

$$\int \frac{a + b \operatorname{arctanh}(cx)}{\sqrt{dx}} dx$$

$$= \frac{\sqrt{d} \left(2\sqrt{c} \operatorname{atan}\left(\frac{\sqrt{xc}}{\sqrt{c}}\right) b + 2\sqrt{c} \operatorname{atanh}(cx) b + 2\sqrt{x} \operatorname{atanh}(cx) bc + 2\sqrt{c} \log(\sqrt{x} \sqrt{c} - 1) b - \sqrt{c} \log(cx + 1) \right)}{cd}$$

input `int((a+b*atanh(c*x))/(d*x)^(1/2),x)`

output `(sqrt(d)*(2*sqrt(c)*atan((sqrt(x)*c)/sqrt(c))*b + 2*sqrt(c)*atanh(c*x)*b +
2*sqrt(x)*atanh(c*x)*b*c + 2*sqrt(c)*log(sqrt(x)*sqrt(c) - 1)*b - sqrt(c)
*log(c*x + 1)*b + 2*sqrt(x)*a*c))/(c*d)`

3.39 $\int \frac{a+b\operatorname{arctanh}(cx)}{(dx)^{3/2}} dx$

Optimal result	399
Mathematica [A] (verified)	399
Rubi [A] (verified)	400
Maple [A] (verified)	402
Fricas [A] (verification not implemented)	402
Sympy [F]	403
Maxima [A] (verification not implemented)	403
Giac [A] (verification not implemented)	404
Mupad [F(-1)]	404
Reduce [B] (verification not implemented)	404

Optimal result

Integrand size = 16, antiderivative size = 85

$$\int \frac{a + b\operatorname{arctanh}(cx)}{(dx)^{3/2}} dx = \frac{2b\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{d^{3/2}} - \frac{2(a + b\operatorname{arctanh}(cx))}{d\sqrt{dx}} + \frac{2b\sqrt{c}\operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{d^{3/2}}$$

output

```
2*b*c^(1/2)*arctan(c^(1/2)*(d*x)^(1/2)/d^(1/2))/d^(3/2)-2*(a+b*arctanh(c*x))/d/(d*x)^(1/2)+2*b*c^(1/2)*arctanh(c^(1/2)*(d*x)^(1/2)/d^(1/2))/d^(3/2)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.16

$$\int \frac{a + b\operatorname{arctanh}(cx)}{(dx)^{3/2}} dx = \frac{x(-2a + 2b\sqrt{c}\sqrt{x} \arctan(\sqrt{c}\sqrt{x}) - 2b\operatorname{arctanh}(cx) - b\sqrt{c}\sqrt{x} \log(1 - \sqrt{c}\sqrt{x}))}{(dx)^{3/2}}$$

input

```
Integrate[(a + b*ArcTanh[c*x])/(d*x)^(3/2), x]
```

output

```
(x*(-2*a + 2*b*Sqrt[c]*Sqrt[x]*ArcTan[Sqrt[c]*Sqrt[x]] - 2*b*ArcTanh[c*x]
- b*Sqrt[c]*Sqrt[x]*Log[1 - Sqrt[c]*Sqrt[x]] + b*Sqrt[c]*Sqrt[x]*Log[1 + S
qrt[c]*Sqrt[x]]))/(d*x)^(3/2)
```

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.12, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6464, 266, 756, 218, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barctanh}(cx)}{(dx)^{3/2}} dx \\
 & \quad \downarrow \text{6464} \\
 & \frac{2bc \int \frac{1}{\sqrt{dx}(1-c^2x^2)} dx}{d} - \frac{2(a + \operatorname{barctanh}(cx))}{d\sqrt{dx}} \\
 & \quad \downarrow \text{266} \\
 & \frac{4bc \int \frac{1}{1-c^2x^2} d\sqrt{dx}}{d^2} - \frac{2(a + \operatorname{barctanh}(cx))}{d\sqrt{dx}} \\
 & \quad \downarrow \text{756} \\
 & \frac{4bc \left(\frac{1}{2} d \int \frac{1}{d-cdx} d\sqrt{dx} + \frac{1}{2} d \int \frac{1}{cx+d} d\sqrt{dx} \right)}{d^2} - \frac{2(a + \operatorname{barctanh}(cx))}{d\sqrt{dx}} \\
 & \quad \downarrow \text{218} \\
 & \frac{4bc \left(\frac{1}{2} d \int \frac{1}{d-cdx} d\sqrt{dx} + \frac{\sqrt{d} \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt{c}} \right)}{d^2} - \frac{2(a + \operatorname{barctanh}(cx))}{d\sqrt{dx}} \\
 & \quad \downarrow \text{221} \\
 & \frac{4bc \left(\frac{\sqrt{d} \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt{c}} + \frac{\sqrt{d} \operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt{c}} \right)}{d^2} - \frac{2(a + \operatorname{barctanh}(cx))}{d\sqrt{dx}}
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c*x])/(d*x)^(3/2), x]`

output `(-2*(a + b*ArcTanh[c*x]))/(d*Sqrt[d*x]) + (4*b*c*((Sqrt[d]*ArcTan[(Sqrt[c]*Sqrt[d*x])/Sqrt[d]])/(2*Sqrt[c]) + (Sqrt[d]*ArcTanh[(Sqrt[c]*Sqrt[d*x])/Sqrt[d]])/(2*Sqrt[c])))/d^2`

Defintions of rubi rules used

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 266 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntegerBinomialQ[a, b, c, 2, m, p, x]`

rule 756 `Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 6464 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)*((d_)*(x_)^(m_)), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\frac{-\frac{2a}{\sqrt{dx}} - \frac{2b \operatorname{arctanh}(cx)}{\sqrt{dx}} + \frac{2bc \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{d} + \frac{2bc \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{\sqrt{cd}}}{d}$	69
default	$\frac{-\frac{2a}{\sqrt{dx}} - \frac{2b \operatorname{arctanh}(cx)}{\sqrt{dx}} + \frac{2bc \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{d} + \frac{2bc \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{\sqrt{cd}}}{d}$	69
parts	$-\frac{2a}{\sqrt{dx}d} - \frac{2b \operatorname{arctanh}(cx)}{d\sqrt{dx}} + \frac{2bc \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{d\sqrt{cd}} + \frac{2bc \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{d\sqrt{cd}}$	78

input `int((a+b*arctanh(c*x))/(d*x)^(3/2),x,method=_RETURNVERBOSE)`

output
$$\frac{2/d*(-a/(d*x)^{(1/2)}-b/(d*x)^{(1/2)}*\operatorname{arctanh}(c*x)+b*c/(c*d)^{(1/2)}*\operatorname{arctanh}(c*(d*x)^{(1/2)}/(c*d)^{(1/2}))+b*c/(c*d)^{(1/2)}*\operatorname{arctan}(c*(d*x)^{(1/2)}/(c*d)^{(1/2}))}{d}$$

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 208, normalized size of antiderivative = 2.45

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{3/2}} dx = \left[\frac{2 b dx \sqrt{\frac{c}{d}} \operatorname{arctan}\left(\sqrt{dx} \sqrt{\frac{c}{d}}\right) + b dx \sqrt{\frac{c}{d}} \log\left(\frac{cx+2\sqrt{dx}\sqrt{\frac{c}{d}}+1}{cx-1}\right) - \sqrt{dx} \left(b \log\left(-\frac{cx+1}{cx-1}\right) + 2a\right)}{d^2 x} \right. \\ \left. - \frac{2 b dx \sqrt{-\frac{c}{d}} \operatorname{arctan}\left(\sqrt{dx} \sqrt{-\frac{c}{d}}\right) - b dx \sqrt{-\frac{c}{d}} \log\left(\frac{cx+2\sqrt{dx}\sqrt{-\frac{c}{d}}-1}{cx+1}\right) + \sqrt{dx} \left(b \log\left(-\frac{cx+1}{cx-1}\right) + 2a\right)}{d^2 x} \right]$$

input `integrate((a+b*arctanh(c*x))/(d*x)^(3/2),x, algorithm="fricas")`

output

```
[(2*b*d*x*sqrt(c/d)*arctan(sqrt(d*x)*sqrt(c/d)) + b*d*x*sqrt(c/d)*log((c*x
+ 2*sqrt(d*x)*sqrt(c/d) + 1)/(c*x - 1)) - sqrt(d*x)*(b*log(-(c*x + 1)/(c*
x - 1)) + 2*a))/(d^2*x), -(2*b*d*x*sqrt(-c/d)*arctan(sqrt(d*x)*sqrt(-c/d))
- b*d*x*sqrt(-c/d)*log((c*x + 2*sqrt(d*x)*sqrt(-c/d) - 1)/(c*x + 1)) + sq
rt(d*x)*(b*log(-(c*x + 1)/(c*x - 1)) + 2*a))/(d^2*x)]
```

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{3/2}} dx = \int \frac{a + b \operatorname{atanh}(cx)}{(dx)^{3/2}} dx$$

input

```
integrate((a+b*atanh(c*x))/(d*x)**(3/2), x)
```

output

```
Integral((a + b*atanh(c*x))/(d*x)**(3/2), x)
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.11

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{3/2}} dx = \frac{b \left(\frac{\left(\frac{2d \operatorname{arctan}\left(\frac{\sqrt{dx}c}{\sqrt{cd}}\right) - d \log\left(\frac{\sqrt{dx}c - \sqrt{cd}}{\sqrt{dx}c + \sqrt{cd}}\right)}{\sqrt{cd}} \right) c}{d} - \frac{2 \operatorname{arctanh}(cx)}{\sqrt{dx}} \right) - \frac{2a}{\sqrt{dx}}}{d}$$

input

```
integrate((a+b*arctanh(c*x))/(d*x)^(3/2), x, algorithm="maxima")
```

output

```
(b*((2*d*arctan(sqrt(d*x)*c/sqrt(c*d))/sqrt(c*d) - d*log((sqrt(d*x)*c - sq
rt(c*d))/(sqrt(d*x)*c + sqrt(c*d)))/sqrt(c*d))*c/d - 2*arctanh(c*x)/sqrt(d
*x)) - 2*a/sqrt(d*x))/d
```


Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.09

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{3/2}} dx = \frac{2bcd \left(\frac{\arctan\left(\frac{\sqrt{dxc}}{\sqrt{cd}}\right)}{\sqrt{cdd}} - \frac{\arctan\left(\frac{\sqrt{dxc}}{\sqrt{-cd}}\right)}{\sqrt{-cdd}} \right) - \frac{b \log\left(\frac{-cdx+d}{cdx-d}\right)}{\sqrt{dx}} - \frac{2a}{\sqrt{dx}}}{d}$$

input `integrate((a+b*arctanh(c*x))/(d*x)^(3/2),x, algorithm="giac")`output `(2*b*c*d*(arctan(sqrt(d*x)*c/sqrt(c*d))/(sqrt(c*d)*d) - arctan(sqrt(d*x)*c/sqrt(-c*d))/(sqrt(-c*d)*d)) - b*log(-(c*d*x + d)/(c*d*x - d))/sqrt(d*x) - 2*a/sqrt(d*x))/d`**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{3/2}} dx = \int \frac{a + b \operatorname{atanh}(cx)}{(dx)^{3/2}} dx$$

input `int((a + b*atanh(c*x))/(d*x)^(3/2),x)`output `int((a + b*atanh(c*x))/(d*x)^(3/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.88

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{3/2}} dx = \frac{\sqrt{d} \left(2\sqrt{x} \sqrt{c} \operatorname{atan}\left(\frac{\sqrt{x}c}{\sqrt{c}}\right) b - 2\sqrt{x} \sqrt{c} \operatorname{atanh}(cx) b - 2 \operatorname{atanh}(cx) b - 2\sqrt{x} \sqrt{c} \log\left(\frac{-cdx+d}{cdx-d}\right) \right)}{\sqrt{x} d^2}$$

input `int((a+b*atanh(c*x))/(d*x)^(3/2),x)`

output

```
(sqrt(d)*(2*sqrt(x)*sqrt(c)*atan((sqrt(x)*c)/sqrt(c))*b - 2*sqrt(x)*sqrt(c)
)*atanh(c*x)*b - 2*atanh(c*x)*b - 2*sqrt(x)*sqrt(c)*log(sqrt(x)*sqrt(c) -
1)*b + sqrt(x)*sqrt(c)*log(c*x + 1)*b - 2*a)/(sqrt(x)*d**2)
```

3.40 $\int \frac{a+b\operatorname{arctanh}(cx)}{(dx)^{5/2}} dx$

Optimal result	406
Mathematica [A] (verified)	406
Rubi [A] (verified)	407
Maple [A] (verified)	409
Fricas [A] (verification not implemented)	410
Sympy [F]	410
Maxima [A] (verification not implemented)	411
Giac [A] (verification not implemented)	411
Mupad [F(-1)]	412
Reduce [B] (verification not implemented)	412

Optimal result

Integrand size = 16, antiderivative size = 107

$$\int \frac{a + b\operatorname{arctanh}(cx)}{(dx)^{5/2}} dx = -\frac{4bc}{3d^2\sqrt{dx}} - \frac{2bc^{3/2} \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{3d^{5/2}} - \frac{2(a + b\operatorname{arctanh}(cx))}{3d(dx)^{3/2}} + \frac{2bc^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{3d^{5/2}}$$

output

```
-4/3*b*c/d^2/(d*x)^(1/2)-2/3*b*c^(3/2)*arctan(c^(1/2)*(d*x)^(1/2)/d^(1/2))
/d^(5/2)-2/3*(a+b*arctanh(c*x))/d/(d*x)^(3/2)+2/3*b*c^(3/2)*arctanh(c^(1/2)
)*(d*x)^(1/2)/d^(1/2))/d^(5/2)
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.00

$$\int \frac{a + b\operatorname{arctanh}(cx)}{(dx)^{5/2}} dx = \frac{x(2a + 4bcx + 2bc^{3/2}x^{3/2} \arctan(\sqrt{c}\sqrt{x}) + 2b\operatorname{arctanh}(cx) + bc^{3/2}x^{3/2} \log(1 - \sqrt{c}\sqrt{x}) - bc^{3/2}x^{3/2} \log(1 + \sqrt{c}\sqrt{x}))}{3(dx)^{5/2}}$$

input `Integrate[(a + b*ArcTanh[c*x])/(d*x)^(5/2), x]`

output `-1/3*(x*(2*a + 4*b*c*x + 2*b*c^(3/2)*x^(3/2)*ArcTan[Sqrt[c]*Sqrt[x]] + 2*b*ArcTanh[c*x] + b*c^(3/2)*x^(3/2)*Log[1 - Sqrt[c]*Sqrt[x]] - b*c^(3/2)*x^(3/2)*Log[1 + Sqrt[c]*Sqrt[x]])/(d*x)^(5/2)`

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.12, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {6464, 264, 266, 27, 827, 218, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{5/2}} dx \\
 & \quad \downarrow 6464 \\
 & \frac{2bc \int \frac{1}{(dx)^{3/2}(1-c^2x^2)} dx}{3d} - \frac{2(a + b \operatorname{arctanh}(cx))}{3d(dx)^{3/2}} \\
 & \quad \downarrow 264 \\
 & \frac{2bc \left(\frac{c^2 \int \frac{\sqrt{dx}}{1-c^2x^2} dx}{d^2} - \frac{2}{d\sqrt{dx}} \right)}{3d} - \frac{2(a + b \operatorname{arctanh}(cx))}{3d(dx)^{3/2}} \\
 & \quad \downarrow 266 \\
 & \frac{2bc \left(\frac{2c^2 \int \frac{d^3x}{d^2-c^2d^2x^2} d\sqrt{dx}}{d^3} - \frac{2}{d\sqrt{dx}} \right)}{3d} - \frac{2(a + b \operatorname{arctanh}(cx))}{3d(dx)^{3/2}} \\
 & \quad \downarrow 27 \\
 & \frac{2bc \left(\frac{2c^2 \int \frac{dx}{d^2-c^2d^2x^2} d\sqrt{dx}}{d} - \frac{2}{d\sqrt{dx}} \right)}{3d} - \frac{2(a + b \operatorname{arctanh}(cx))}{3d(dx)^{3/2}} \\
 & \quad \downarrow 827
 \end{aligned}$$

$$\begin{aligned}
& \frac{2bc \left(\frac{2c^2 \left(\frac{\int \frac{1}{d-cdx} d\sqrt{dx}}{2c} - \frac{\int \frac{1}{cxd+d} d\sqrt{dx}}{2c} \right)}{d} - \frac{2}{d\sqrt{dx}} \right)}{3d} - \frac{2(a + b \operatorname{arctanh}(cx))}{3d(dx)^{3/2}} \\
& \quad \downarrow \text{218} \\
& \frac{2bc \left(\frac{2c^2 \left(\frac{\int \frac{1}{d-cdx} d\sqrt{dx}}{2c} - \frac{\arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/2}\sqrt{d}} \right)}{d} - \frac{2}{d\sqrt{dx}} \right)}{3d} - \frac{2(a + b \operatorname{arctanh}(cx))}{3d(dx)^{3/2}} \\
& \quad \downarrow \text{221} \\
& \frac{2bc \left(\frac{2c^2 \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/2}\sqrt{d}} - \frac{\arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/2}\sqrt{d}} \right)}{d} - \frac{2}{d\sqrt{dx}} \right)}{3d} - \frac{2(a + b \operatorname{arctanh}(cx))}{3d(dx)^{3/2}}
\end{aligned}$$

input `Int[(a + b*ArcTanh[c*x])/(d*x)^(5/2), x]`

output `(-2*(a + b*ArcTanh[c*x]))/(3*d*(d*x)^(3/2)) + (2*b*c*(-2/(d*Sqrt[d*x])) + (2*c^2*(-1/2*ArcTan[(Sqrt[c]*Sqrt[d*x])/Sqrt[d]]/(c^(3/2)*Sqrt[d]) + ArcTanh[(Sqrt[c]*Sqrt[d*x])/Sqrt[d]]/(2*c^(3/2)*Sqrt[d])))/d)/(3*d)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 264 `Int[((c.)*(x.))^(m.)*((a.) + (b.)*(x.)^2)^(p.), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c.)*(x.))^(m.)*((a.) + (b.)*(x.)^2)^(p.), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 827 `Int[(x.)^2/((a.) + (b.)*(x.)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x], x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 6464 `Int[((a.) + ArcTanh[(c.)*(x.)^(n.)]*(b.))*((d.)*(x.))^(m.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.87

method	result	size
derivativedivides	$\frac{-\frac{2a}{3(dx)^{\frac{3}{2}}} - \frac{2b \operatorname{arctanh}(cx)}{3(dx)^{\frac{3}{2}}} + \frac{2b c^2 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{3d\sqrt{cd}} - \frac{2b c^2 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{3d\sqrt{cd}} - \frac{4bc}{3d\sqrt{dx}}}{d}$	93
default	$\frac{-\frac{2a}{3(dx)^{\frac{3}{2}}} - \frac{2b \operatorname{arctanh}(cx)}{3(dx)^{\frac{3}{2}}} + \frac{2b c^2 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{3d\sqrt{cd}} - \frac{2b c^2 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{3d\sqrt{cd}} - \frac{4bc}{3d\sqrt{dx}}}{d}$	93
parts	$-\frac{2a}{3(dx)^{\frac{3}{2}}d} - \frac{2b \operatorname{arctanh}(cx)}{3d(dx)^{\frac{3}{2}}} + \frac{2b c^2 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{3d^2\sqrt{cd}} - \frac{2b c^2 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{3d^2\sqrt{cd}} - \frac{4bc}{3d^2\sqrt{dx}}$	94

input `int((a+b*arctanh(c*x))/(d*x)^(5/2),x,method=_RETURNVERBOSE)`

output

```
2/d*(-1/3*a/(d*x)^(3/2)-1/3*b/(d*x)^(3/2)*arctanh(c*x)+1/3*b/d*c^2/(c*d)^(
1/2)*arctanh(c*(d*x)^(1/2)/(c*d)^(1/2))-1/3*b/d*c^2/(c*d)^(1/2)*arctan(c*(
d*x)^(1/2)/(c*d)^(1/2))-2/3*b/d*c/(d*x)^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 231, normalized size of antiderivative = 2.16

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{5/2}} dx = \left[\frac{2 b c d x^2 \sqrt{\frac{c}{d}} \arctan\left(\sqrt{dx} \sqrt{\frac{c}{d}}\right) - b c d x^2 \sqrt{\frac{c}{d}} \log\left(\frac{cx+2\sqrt{dx}\sqrt{\frac{c}{d}}+1}{cx-1}\right) + (4 b c x + b \log(-\frac{cx+1}{cx-1})) \sqrt{dx}}{3 d^3 x^2} \right. \\ \left. - \frac{2 b c d x^2 \sqrt{-\frac{c}{d}} \arctan\left(\sqrt{dx} \sqrt{-\frac{c}{d}}\right) - b c d x^2 \sqrt{-\frac{c}{d}} \log\left(\frac{cx-2\sqrt{dx}\sqrt{-\frac{c}{d}}-1}{cx+1}\right) + (4 b c x + b \log(-\frac{cx+1}{cx-1}) + 2 a) \sqrt{dx}}{3 d^3 x^2} \right]$$

input

```
integrate((a+b*arctanh(c*x))/(d*x)^(5/2),x, algorithm="fricas")
```

output

```
[-1/3*(2*b*c*d*x^2*sqrt(c/d)*arctan(sqrt(d*x)*sqrt(c/d)) - b*c*d*x^2*sqrt(
c/d)*log((c*x + 2*sqrt(d*x)*sqrt(c/d) + 1)/(c*x - 1)) + (4*b*c*x + b*log(-
(c*x + 1)/(c*x - 1)) + 2*a)*sqrt(d*x))/(d^3*x^2), -1/3*(2*b*c*d*x^2*sqrt(-
c/d)*arctan(sqrt(d*x)*sqrt(-c/d)) - b*c*d*x^2*sqrt(-c/d)*log((c*x - 2*sqrt
(d*x)*sqrt(-c/d) - 1)/(c*x + 1)) + (4*b*c*x + b*log(-(c*x + 1)/(c*x - 1))
+ 2*a)*sqrt(d*x))/(d^3*x^2)]
```

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{5/2}} dx = \int \frac{a + b \operatorname{atanh}(cx)}{(dx)^{5/2}} dx$$

input

```
integrate((a+b*atanh(c*x))/(d*x)**(5/2),x)
```

output

```
Integral((a + b*atanh(c*x))/(d*x)**(5/2), x)
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.94

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{5/2}} dx = - \frac{b \left(\frac{\left(\frac{2c \arctan\left(\frac{\sqrt{dxc}}{\sqrt{cd}}\right) + \frac{c \log\left(\frac{\sqrt{dxc}-\sqrt{cd}}{\sqrt{dxc}+\sqrt{cd}}\right) + \frac{4}{\sqrt{dx}}\right)}{d} \right) c}{d} + \frac{2 \operatorname{artanh}(cx)}{(dx)^{3/2}} \right) + \frac{2a}{(dx)^{3/2}}}{3d}$$

input `integrate((a+b*arctanh(c*x))/(d*x)^(5/2),x, algorithm="maxima")`

output `-1/3*(b*((2*c*arctan(sqrt(d*x)*c/sqrt(c*d))/sqrt(c*d) + c*log((sqrt(d*x)*c - sqrt(c*d))/(sqrt(d*x)*c + sqrt(c*d)))/sqrt(c*d) + 4/sqrt(d*x))*c/d + 2*arctanh(c*x)/(d*x)^(3/2)) + 2*a/(d*x)^(3/2))/d`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.09

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{5/2}} dx = - \frac{\frac{2bc^2 \arctan\left(\frac{\sqrt{dxc}}{\sqrt{cd}}\right)}{\sqrt{cdd}} + \frac{2bc^2 \arctan\left(\frac{\sqrt{dxc}}{\sqrt{-cd}}\right)}{\sqrt{-cdd}} + \frac{b \log\left(\frac{-cdx+d}{cdx-d}\right)}{\sqrt{dxdx}} + \frac{2(2bcdx+ad)}{\sqrt{dxd^2x}}}{3d}$$

input `integrate((a+b*arctanh(c*x))/(d*x)^(5/2),x, algorithm="giac")`

output `-1/3*(2*b*c^2*arctan(sqrt(d*x)*c/sqrt(c*d))/(sqrt(c*d)*d) + 2*b*c^2*arctan(sqrt(d*x)*c/sqrt(-c*d))/(sqrt(-c*d)*d) + b*log(-(c*d*x + d)/(c*d*x - d))/(sqrt(d*x)*d*x) + 2*(2*b*c*d*x + a*d)/(sqrt(d*x)*d^2*x))/d`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{5/2}} dx = \int \frac{a + b \operatorname{atanh}(cx)}{(dx)^{5/2}} dx$$

input `int((a + b*atanh(c*x))/(d*x)^(5/2), x)`output `int((a + b*atanh(c*x))/(d*x)^(5/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.86

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{5/2}} dx = \frac{\sqrt{d} \left(-2\sqrt{x} \sqrt{c} \operatorname{atan}\left(\frac{\sqrt{x}c}{\sqrt{c}}\right) b c x - 2\sqrt{x} \sqrt{c} \operatorname{atanh}(cx) b c x - 2 \operatorname{atanh}(cx) b - 2\sqrt{x} \sqrt{c} \right)}{3\sqrt{x} d^3 x}$$

input `int((a+b*atanh(c*x))/(d*x)^(5/2), x)`output `(sqrt(d)*(- 2*sqrt(x)*sqrt(c)*atan((sqrt(x)*c)/sqrt(c))*b*c*x - 2*sqrt(x)*sqrt(c)*atanh(c*x)*b*c*x - 2*atanh(c*x)*b - 2*sqrt(x)*sqrt(c)*log(sqrt(x)*sqrt(c) - 1)*b*c*x + sqrt(x)*sqrt(c)*log(c*x + 1)*b*c*x - 2*a - 4*b*c*x)/(3*sqrt(x)*d**3*x)`

3.41 $\int \frac{a+b\operatorname{arctanh}(cx)}{(dx)^{7/2}} dx$

Optimal result	413
Mathematica [A] (verified)	413
Rubi [A] (verified)	414
Maple [A] (verified)	416
Fricas [A] (verification not implemented)	417
Sympy [F]	417
Maxima [A] (verification not implemented)	418
Giac [A] (verification not implemented)	418
Mupad [F(-1)]	419
Reduce [B] (verification not implemented)	419

Optimal result

Integrand size = 16, antiderivative size = 107

$$\int \frac{a + b\operatorname{arctanh}(cx)}{(dx)^{7/2}} dx = -\frac{4bc}{15d^2(dx)^{3/2}} + \frac{2bc^{5/2} \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{5d^{7/2}} - \frac{2(a + b\operatorname{arctanh}(cx))}{5d(dx)^{5/2}} + \frac{2bc^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{5d^{7/2}}$$

output `-4/15*b*c/d^2/(d*x)^(3/2)+2/5*b*c^(5/2)*arctan(c^(1/2)*(d*x)^(1/2)/d^(1/2))/d^(7/2)-2/5*(a+b*arctanh(c*x))/d/(d*x)^(5/2)+2/5*b*c^(5/2)*arctanh(c^(1/2)*(d*x)^(1/2)/d^(1/2))/d^(7/2)`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.01

$$\int \frac{a + b\operatorname{arctanh}(cx)}{(dx)^{7/2}} dx = \frac{x(-6a - 4bcx + 6bc^{5/2}x^{5/2} \arctan(\sqrt{c}\sqrt{x}) - 6b\operatorname{arctanh}(cx) - 3bc^{5/2}x^{5/2} \log(1 + \sqrt{c}\sqrt{x}))}{15(dx)^{7/2}}$$

input `Integrate[(a + b*ArcTanh[c*x])/(d*x)^(7/2), x]`

output

```
(x*(-6*a - 4*b*c*x + 6*b*c^(5/2)*x^(5/2)*ArcTan[Sqrt[c]*Sqrt[x]] - 6*b*ArcTanh[c*x] - 3*b*c^(5/2)*x^(5/2)*Log[1 - Sqrt[c]*Sqrt[x]] + 3*b*c^(5/2)*x^(5/2)*Log[1 + Sqrt[c]*Sqrt[x]])/(15*(d*x)^(7/2))
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.14, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6464, 264, 266, 756, 218, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barctanh}(cx)}{(dx)^{7/2}} dx \\
 & \quad \downarrow \text{6464} \\
 & \frac{2bc \int \frac{1}{(dx)^{5/2}(1-c^2x^2)} dx}{5d} - \frac{2(a + \operatorname{barctanh}(cx))}{5d(dx)^{5/2}} \\
 & \quad \downarrow \text{264} \\
 & \frac{2bc \left(\frac{c^2 \int \frac{1}{\sqrt{dx}(1-c^2x^2)} dx}{d^2} - \frac{2}{3d(dx)^{3/2}} \right)}{5d} - \frac{2(a + \operatorname{barctanh}(cx))}{5d(dx)^{5/2}} \\
 & \quad \downarrow \text{266} \\
 & \frac{2bc \left(\frac{2c^2 \int \frac{1}{1-c^2x^2} d\sqrt{dx}}{d^3} - \frac{2}{3d(dx)^{3/2}} \right)}{5d} - \frac{2(a + \operatorname{barctanh}(cx))}{5d(dx)^{5/2}} \\
 & \quad \downarrow \text{756} \\
 & \frac{2bc \left(\frac{2c^2 \left(\frac{1}{2} d \int \frac{1}{d-cdx} d\sqrt{dx} + \frac{1}{2} d \int \frac{1}{cx+d} d\sqrt{dx} \right)}{d^3} - \frac{2}{3d(dx)^{3/2}} \right)}{5d} - \frac{2(a + \operatorname{barctanh}(cx))}{5d(dx)^{5/2}} \\
 & \quad \downarrow \text{218}
 \end{aligned}$$

$$\frac{2bc \left(\frac{2c^2 \left(\frac{1}{2} d \int \frac{1}{d-cx} d\sqrt{dx} + \frac{\sqrt{d} \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt{c}} \right)}{d^3} - \frac{2}{3d(dx)^{3/2}} \right)}{5d} - \frac{2(a + b \operatorname{arctanh}(cx))}{5d(dx)^{5/2}}$$

↓ 221

$$\frac{2bc \left(\frac{2c^2 \left(\frac{\sqrt{d} \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt{c}} + \frac{\sqrt{d} \operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt{c}} \right)}{d^3} - \frac{2}{3d(dx)^{3/2}} \right)}{5d} - \frac{2(a + b \operatorname{arctanh}(cx))}{5d(dx)^{5/2}}$$

input `Int[(a + b*ArcTanh[c*x])/(d*x)^(7/2), x]`

output `(-2*(a + b*ArcTanh[c*x]))/(5*d*(d*x)^(5/2)) + (2*b*c*(-2/(3*d*(d*x)^(3/2)) + (2*c^2*((Sqrt[d]*ArcTan[(Sqrt[c]*Sqrt[d*x])/Sqrt[d]])/(2*Sqrt[c]) + (Sqrt[d]*ArcTanh[(Sqrt[c]*Sqrt[d*x])/Sqrt[d]])/(2*Sqrt[c])))/d^3))/(5*d)`

Defintions of rubi rules used

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 264 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^(p), x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 756 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 6464 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_)), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.87

method	result	size
derivativedivides	$\frac{-\frac{2a}{5(dx)^{\frac{5}{2}}} - \frac{2b \operatorname{arctanh}(cx)}{5(dx)^{\frac{5}{2}}} + \frac{2b c^3 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{5d^2\sqrt{cd}} - \frac{4bc}{15d(dx)^{\frac{3}{2}}} + \frac{2b c^3 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{5d^2\sqrt{cd}}}{d}$	93
default	$\frac{-\frac{2a}{5(dx)^{\frac{5}{2}}} - \frac{2b \operatorname{arctanh}(cx)}{5(dx)^{\frac{5}{2}}} + \frac{2b c^3 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{5d^2\sqrt{cd}} - \frac{4bc}{15d(dx)^{\frac{3}{2}}} + \frac{2b c^3 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{5d^2\sqrt{cd}}}{d}$	93
parts	$-\frac{2a}{5(dx)^{\frac{5}{2}}d} - \frac{2b \operatorname{arctanh}(cx)}{5d(dx)^{\frac{5}{2}}} + \frac{2b c^3 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{5d^3\sqrt{cd}} + \frac{2b c^3 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{5d^3\sqrt{cd}} - \frac{4bc}{15d^2(dx)^{\frac{3}{2}}}$	94

input `int((a+b*arctanh(c*x))/(d*x)^(7/2),x,method=_RETURNVERBOSE)`

output `2/d*(-1/5*a/(d*x)^(5/2)-1/5*b/(d*x)^(5/2)*arctanh(c*x)+1/5*b/d^2*c^3/(c*d)^(1/2)*arctanh(c*(d*x)^(1/2)/(c*d)^(1/2))-2/15*b/d*c/(d*x)^(3/2)+1/5*b/d^2*c^3/(c*d)^(1/2)*arctan(c*(d*x)^(1/2)/(c*d)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 242, normalized size of antiderivative = 2.26

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{7/2}} dx = \frac{\left[6bc^2 dx^3 \sqrt{\frac{c}{d}} \arctan\left(\sqrt{dx} \sqrt{\frac{c}{d}}\right) + 3bc^2 dx^3 \sqrt{\frac{c}{d}} \log\left(\frac{cx+2\sqrt{dx}\sqrt{\frac{c}{d}}+1}{cx-1}\right) - (4bcx + 3b \log(-\frac{cx+1}{cx-1})) + 6a \sqrt{dx} \right]}{15d^4 x^3} - \frac{\left[6bc^2 dx^3 \sqrt{-\frac{c}{d}} \arctan\left(\sqrt{dx} \sqrt{-\frac{c}{d}}\right) - 3bc^2 dx^3 \sqrt{-\frac{c}{d}} \log\left(\frac{cx+2\sqrt{dx}\sqrt{-\frac{c}{d}}-1}{cx+1}\right) + (4bcx + 3b \log(-\frac{cx+1}{cx-1})) + 6a \sqrt{dx} \right]}{15d^4 x^3}$$

input `integrate((a+b*arctanh(c*x))/(d*x)^(7/2),x, algorithm="fricas")`

output `[1/15*(6*b*c^2*d*x^3*sqrt(c/d)*arctan(sqrt(d*x)*sqrt(c/d)) + 3*b*c^2*d*x^3*sqrt(c/d)*log((c*x + 2*sqrt(d*x)*sqrt(c/d) + 1)/(c*x - 1)) - (4*b*c*x + 3*b*log(-(c*x + 1)/(c*x - 1)) + 6*a)*sqrt(d*x))/(d^4*x^3), -1/15*(6*b*c^2*d*x^3*sqrt(-c/d)*arctan(sqrt(d*x)*sqrt(-c/d)) - 3*b*c^2*d*x^3*sqrt(-c/d)*log((c*x + 2*sqrt(d*x)*sqrt(-c/d) - 1)/(c*x + 1)) + (4*b*c*x + 3*b*log(-(c*x + 1)/(c*x - 1)) + 6*a)*sqrt(d*x))/(d^4*x^3)]`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{7/2}} dx = \int \frac{a + b \operatorname{atanh}(cx)}{(dx)^{7/2}} dx$$

input `integrate((a+b*atanh(c*x))/(d*x)**(7/2),x)`

output `Integral((a + b*atanh(c*x))/(d*x)**(7/2), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.05

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{7/2}} dx = \frac{b \left(\frac{6c^2 \arctan\left(\frac{\sqrt{dxc}}{\sqrt{cd}}\right) - 3c^2 \log\left(\frac{\sqrt{dxc}-\sqrt{cd}}{\sqrt{dxc}+\sqrt{cd}}\right) - \frac{4}{(dx)^{3/2}}}{\sqrt{cdd}} \right) c - \frac{6 \operatorname{arctanh}(cx)}{(dx)^{5/2}} - \frac{6a}{(dx)^{5/2}}}{15d}$$

input `integrate((a+b*arctanh(c*x))/(d*x)^(7/2),x, algorithm="maxima")`

output `1/15*(b*((6*c^2*arctan(sqrt(d*x)*c/sqrt(c*d))/(sqrt(c*d)*d) - 3*c^2*log((sqrt(d*x)*c - sqrt(c*d))/(sqrt(d*x)*c + sqrt(c*d)))/(sqrt(c*d)*d) - 4/(d*x)^(3/2))*c/d - 6*arctanh(c*x)/(d*x)^(5/2)) - 6*a/(d*x)^(5/2))/d`

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.09

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{7/2}} dx = \frac{6bc^3 \left(\frac{\arctan\left(\frac{\sqrt{dxc}}{\sqrt{cd}}\right)}{\sqrt{cdd^2}} - \frac{\arctan\left(\frac{\sqrt{dxc}}{\sqrt{-cd}}\right)}{\sqrt{-cdd^2}} \right) - \frac{3b \log\left(\frac{-cdx+d}{cdx-d}\right)}{\sqrt{dx}d^2x^2} - \frac{2(2bcdx+3ad)}{\sqrt{dx}d^3x^2}}{15d}$$

input `integrate((a+b*arctanh(c*x))/(d*x)^(7/2),x, algorithm="giac")`

output `1/15*(6*b*c^3*(arctan(sqrt(d*x)*c/sqrt(c*d))/(sqrt(c*d)*d^2) - arctan(sqrt(d*x)*c/sqrt(-c*d))/(sqrt(-c*d)*d^2)) - 3*b*log(-(c*d*x + d)/(c*d*x - d))/(sqrt(d*x)*d^2*x^2) - 2*(2*b*c*d*x + 3*a*d)/(sqrt(d*x)*d^3*x^2))/d`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{7/2}} dx = \int \frac{a + b \operatorname{atanh}(cx)}{(dx)^{7/2}} dx$$

input `int((a + b*atanh(c*x))/(d*x)^(7/2), x)`output `int((a + b*atanh(c*x))/(d*x)^(7/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.02

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{7/2}} dx = \frac{\sqrt{d} \left(6\sqrt{x} \sqrt{c} \operatorname{atan}\left(\frac{\sqrt{x}c}{\sqrt{c}}\right) b c^2 x^2 - 6\sqrt{x} \sqrt{c} \operatorname{atanh}(cx) b c^2 x^2 - 6 \operatorname{atanh}(cx) b - 6\sqrt{x} \sqrt{c} \right)}{15\sqrt{x} d^4 x}$$

input `int((a+b*atanh(c*x))/(d*x)^(7/2), x)`output `(sqrt(d)*(6*sqrt(x)*sqrt(c)*atan((sqrt(x)*c)/sqrt(c))*b*c**2*x**2 - 6*sqrt(x)*sqrt(c)*atanh(c*x)*b*c**2*x**2 - 6*atanh(c*x)*b - 6*sqrt(x)*sqrt(c)*log(sqrt(x)*sqrt(c) - 1)*b*c**2*x**2 + 3*sqrt(x)*sqrt(c)*log(c*x + 1)*b*c**2*x**2 - 6*a - 4*b*c*x))/(15*sqrt(x)*d**4*x**2)`

3.42 $\int \frac{a+b\operatorname{arctanh}(cx)}{(dx)^{9/2}} dx$

Optimal result	420
Mathematica [A] (verified)	420
Rubi [A] (verified)	421
Maple [A] (verified)	424
Fricas [A] (verification not implemented)	425
Sympy [F(-1)]	425
Maxima [A] (verification not implemented)	426
Giac [A] (verification not implemented)	426
Mupad [F(-1)]	427
Reduce [B] (verification not implemented)	427

Optimal result

Integrand size = 16, antiderivative size = 125

$$\int \frac{a + b\operatorname{arctanh}(cx)}{(dx)^{9/2}} dx = -\frac{4bc}{35d^2(dx)^{5/2}} - \frac{4bc^3}{7d^4\sqrt{dx}} - \frac{2bc^{7/2} \arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{7d^{9/2}} - \frac{2(a + b\operatorname{arctanh}(cx))}{7d(dx)^{7/2}} + \frac{2bc^{7/2}\operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{7d^{9/2}}$$

output

```
-4/35*b*c/d^2/(d*x)^(5/2)-4/7*b*c^3/d^4/(d*x)^(1/2)-2/7*b*c^(7/2)*arctan(c^(1/2)*(d*x)^(1/2)/d^(1/2))/d^(9/2)-2/7*(a+b*arctanh(c*x))/d/(d*x)^(7/2)+2/7*b*c^(7/2)*arctanh(c^(1/2)*(d*x)^(1/2)/d^(1/2))/d^(9/2)
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.98

$$\int \frac{a + b\operatorname{arctanh}(cx)}{(dx)^{9/2}} dx = \frac{\sqrt{dx}(10a + 4bcx + 20bc^3x^3 + 10bc^{7/2}x^{7/2} \arctan(\sqrt{c}\sqrt{x}) + 10b\operatorname{arctanh}(cx) + 5bc^{7/2}x^{7/2} \log(1 - \sqrt{c}\sqrt{x}))}{35d^5x^4}$$

input

```
Integrate[(a + b*ArcTanh[c*x])/(d*x)^(9/2), x]
```

output

$$-1/35*(\text{Sqrt}[d*x]*(10*a + 4*b*c*x + 20*b*c^3*x^3 + 10*b*c^{(7/2)*x^{(7/2)*\text{ArcTan}[\text{Sqrt}[c]*\text{Sqrt}[x]] + 10*b*\text{ArcTanh}[c*x] + 5*b*c^{(7/2)*x^{(7/2)*\text{Log}[1 - \text{Sqrt}[c]*\text{Sqrt}[x]] - 5*b*c^{(7/2)*x^{(7/2)*\text{Log}[1 + \text{Sqrt}[c]*\text{Sqrt}[x]]}))/d^5*x^4)$$
Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.14, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6464, 264, 264, 266, 27, 827, 218, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \text{barctanh}(cx)}{(dx)^{9/2}} dx$$

$$\downarrow 6464$$

$$\frac{2bc \int \frac{1}{(dx)^{7/2}(1-c^2x^2)} dx}{7d} - \frac{2(a + \text{barctanh}(cx))}{7d(dx)^{7/2}}$$

$$\downarrow 264$$

$$\frac{2bc \left(\frac{c^2 \int \frac{1}{(dx)^{3/2}(1-c^2x^2)} dx}{d^2} - \frac{2}{5d(dx)^{5/2}} \right)}{7d} - \frac{2(a + \text{barctanh}(cx))}{7d(dx)^{7/2}}$$

$$\downarrow 264$$

$$\frac{2bc \left(\frac{c^2 \left(\frac{c^2 \int \frac{\sqrt{dx}}{1-c^2x^2} dx}{d^2} - \frac{2}{d\sqrt{dx}} \right)}{d^2} - \frac{2}{5d(dx)^{5/2}} \right)}{7d} - \frac{2(a + \text{barctanh}(cx))}{7d(dx)^{7/2}}$$

$$\downarrow 266$$

$$\frac{2bc \left(\frac{c^2 \left(\frac{2c^2 \int \frac{d^3x}{d^2-c^2d^2x^2} d\sqrt{dx}}{d^3} - \frac{2}{d\sqrt{dx}} \right)}{d^2} - \frac{2}{5d(dx)^{5/2}} \right)}{7d} - \frac{2(a + \text{barctanh}(cx))}{7d(dx)^{7/2}}$$

$$\downarrow 27$$

$$\frac{2bc \left(\frac{c^2 \left(\frac{2c^2 \int \frac{dx}{d^2 - c^2 d^2 x^2} d\sqrt{dx} - \frac{2}{d\sqrt{dx}} \right)}{d^2} - \frac{2}{5d(dx)^{5/2}} \right)}{7d} - \frac{2(a + \operatorname{barctanh}(cx))}{7d(dx)^{7/2}} \right)}{7d}$$

$$\downarrow 827$$

$$\frac{2bc \left(\frac{c^2 \left(\frac{2c^2 \left(\frac{\int \frac{1}{d-cdx} d\sqrt{dx}}{2c} - \frac{\int \frac{1}{cx+d} d\sqrt{dx}}{2c} \right)}{d} - \frac{2}{d\sqrt{dx}} \right)}{d^2} - \frac{2}{5d(dx)^{5/2}} \right)}{7d} - \frac{2(a + \operatorname{barctanh}(cx))}{7d(dx)^{7/2}} \right)}{7d}$$

$$\downarrow 218$$

$$\frac{2bc \left(\frac{c^2 \left(\frac{2c^2 \left(\frac{\int \frac{1}{d-cdx} d\sqrt{dx}}{2c} - \frac{\arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/2}\sqrt{d}} \right)}{d} - \frac{2}{d\sqrt{dx}} \right)}{d^2} - \frac{2}{5d(dx)^{5/2}} \right)}{7d} - \frac{2(a + \operatorname{barctanh}(cx))}{7d(dx)^{7/2}} \right)}{7d}$$

$$\downarrow 221$$

$$\frac{2bc \left(\frac{c^2 \left(\frac{2c^2 \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/2}\sqrt{d}} - \frac{\arctan\left(\frac{\sqrt{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/2}\sqrt{d}} \right)}{d} - \frac{2}{d\sqrt{dx}} \right)}{d^2} - \frac{2}{5d(dx)^{5/2}} \right)}{7d} - \frac{2(a + \operatorname{barctanh}(cx))}{7d(dx)^{7/2}} \right)}{7d}$$

input `Int[(a + b*ArcTanh[c*x])/(d*x)^(9/2), x]`

output
$$\frac{(-2*(a + b*\text{ArcTanh}[c*x]))/(7*d*(d*x)^{(7/2)}) + (2*b*c*(-2/(5*d*(d*x)^{(5/2)}) + (c^2*(-2/(d*\text{Sqrt}[d*x]) + (2*c^2*(-1/2*\text{ArcTan}[(\text{Sqrt}[c]*\text{Sqrt}[d*x])/(\text{Sqrt}[d])]/(c^{(3/2)*\text{Sqrt}[d]} + \text{ArcTanh}[(\text{Sqrt}[c]*\text{Sqrt}[d*x])/(\text{Sqrt}[d])]/(2*c^{(3/2)*\text{Sqrt}[d]})))/d))/d^2))/(7*d)}$$

Defintions of rubi rules used

rule 27
$$\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x) /; \text{FreeQ}[b, x]]$$

rule 218
$$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$$

rule 221
$$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$$

rule 264
$$\text{Int}[(c_)*(x_)^m * (a_ + (b_)*(x_)^2)^p, x_Symbol] \rightarrow \text{Simp}[(c*x)^{(m+1)} * ((a + b*x^2)^{(p+1}) / (a*c*(m+1))), x] - \text{Simp}[b*(m+2*p+3) / (a*c^2*(m+1)) \text{ Int}[(c*x)^{(m+2)} * (a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 266
$$\text{Int}[(c_)*(x_)^m * (a_ + (b_)*(x_)^2)^p, x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \text{ Subst}[\text{Int}[x^{(k*(m+1)-1)} * (a + b*(x^{(2*k)/c^2})^p, x], x, (c*x)^{(1/k)}], x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 827
$$\text{Int}[(x_)^2 / ((a_ + (b_)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[s/(2*b) \text{ Int}[1/(r + s*x^2), x], x] - \text{Simp}[s/(2*b) \text{ Int}[1/(r - s*x^2), x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ !\text{GtQ}[a/b, 0]$$

rule 6464

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))*((d_)*(x_)^(m_), x_Symbol] :
> Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n
/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a,
b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.86

method	result	size
derivativedivides	$\frac{-\frac{2a}{7(dx)^{\frac{7}{2}}} - \frac{2b \operatorname{arctanh}(cx)}{7(dx)^{\frac{7}{2}}} - \frac{2b c^4 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{7d^3\sqrt{cd}} - \frac{4bc}{35d(dx)^{\frac{5}{2}}} - \frac{4b c^3}{7d^3\sqrt{dx}} + \frac{2b c^4 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{7d^3\sqrt{cd}}}{d}$	10
default	$\frac{-\frac{2a}{7(dx)^{\frac{7}{2}}} - \frac{2b \operatorname{arctanh}(cx)}{7(dx)^{\frac{7}{2}}} - \frac{2b c^4 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{7d^3\sqrt{cd}} - \frac{4bc}{35d(dx)^{\frac{5}{2}}} - \frac{4b c^3}{7d^3\sqrt{dx}} + \frac{2b c^4 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{7d^3\sqrt{cd}}}{d}$	10
parts	$-\frac{2a}{7(dx)^{\frac{7}{2}}d} - \frac{2b \operatorname{arctanh}(cx)}{7d(dx)^{\frac{7}{2}}} - \frac{4bc}{35d^2(dx)^{\frac{5}{2}}} - \frac{4b c^3}{7d^4\sqrt{dx}} + \frac{2b c^4 \operatorname{arctanh}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{7d^4\sqrt{cd}} - \frac{2b c^4 \operatorname{arctan}\left(\frac{c\sqrt{dx}}{\sqrt{cd}}\right)}{7d^4\sqrt{cd}}$	10

input

```
int((a+b*arctanh(c*x))/(d*x)^(9/2),x,method=_RETURNVERBOSE)
```

output

```
2/d*(-1/7*a/(d*x)^(7/2)-1/7*b/(d*x)^(7/2)*arctanh(c*x)-1/7*b/d^3*c^4/(c*d)
^(1/2)*arctan(c*(d*x)^(1/2)/(c*d)^(1/2))-2/35*b/d*c/(d*x)^(5/2)-2/7*b/d^3*
c^3/(d*x)^(1/2)+1/7*b/d^3*c^4/(c*d)^(1/2)*arctanh(c*(d*x)^(1/2)/(c*d)^(1/2)
)))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 259, normalized size of antiderivative = 2.07

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{9/2}} dx = \left[\frac{10bc^3 dx^4 \sqrt{\frac{c}{d}} \arctan\left(\sqrt{dx} \sqrt{\frac{c}{d}}\right) - 5bc^3 dx^4 \sqrt{\frac{c}{d}} \log\left(\frac{cx+2\sqrt{dx}\sqrt{\frac{c}{d}}+1}{cx-1}\right) + (20bc^3 x^3 + 4bcx + 5b \log\left(\frac{cx-2\sqrt{dx}\sqrt{\frac{c}{d}}-1}{cx+1}\right))}{35d^5 x^4} \right]$$

input `integrate((a+b*arctanh(c*x))/(d*x)^(9/2),x, algorithm="fricas")`

output `[-1/35*(10*b*c^3*d*x^4*sqrt(c/d)*arctan(sqrt(d*x)*sqrt(c/d)) - 5*b*c^3*d*x^4*sqrt(c/d)*log((c*x + 2*sqrt(d*x)*sqrt(c/d) + 1)/(c*x - 1)) + (20*b*c^3*x^3 + 4*b*c*x + 5*b*log(-(c*x + 1)/(c*x - 1)) + 10*a)*sqrt(d*x))/(d^5*x^4), -1/35*(10*b*c^3*d*x^4*sqrt(-c/d)*arctan(sqrt(d*x)*sqrt(-c/d)) - 5*b*c^3*d*x^4*sqrt(-c/d)*log((c*x - 2*sqrt(d*x)*sqrt(-c/d) - 1)/(c*x + 1)) + (20*b*c^3*x^3 + 4*b*c*x + 5*b*log(-(c*x + 1)/(c*x - 1)) + 10*a)*sqrt(d*x))/(d^5*x^4)]`

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{9/2}} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x))/(d*x)**(9/2),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.04

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{9/2}} dx =$$

$$\frac{b \left(\frac{\left(\frac{10 c^3 \arctan\left(\frac{\sqrt{dxc}}{\sqrt{cd}}\right) + 5 c^3 \log\left(\frac{\sqrt{dxc}-\sqrt{cd}}{\sqrt{dxc}+\sqrt{cd}}\right) + 4 (5 c^2 d^2 x^2 + d^2)}{(dx)^{5/2} d^2} \right) c}{d} + \frac{10 \operatorname{arctanh}(cx)}{(dx)^{7/2}} \right) + \frac{10 a}{(dx)^{7/2}}}{35 d}$$

input `integrate((a+b*arctanh(c*x))/(d*x)^(9/2),x, algorithm="maxima")`output `-1/35*(b*((10*c^3*arctan(sqrt(d*x)*c/sqrt(c*d))/(sqrt(c*d)*d^2) + 5*c^3*log((sqrt(d*x)*c - sqrt(c*d))/(sqrt(d*x)*c + sqrt(c*d)))/(sqrt(c*d)*d^2) + 4*(5*c^2*d^2*x^2 + d^2)/((d*x)^(5/2)*d^2))*c/d + 10*arctanh(c*x)/(d*x)^(7/2)) + 10*a/(d*x)^(7/2))/d`**Giac [A] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.08

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{9/2}} dx =$$

$$\frac{\frac{10 bc^4 \arctan\left(\frac{\sqrt{dxc}}{\sqrt{cd}}\right)}{\sqrt{cdd^3}} + \frac{10 bc^4 \arctan\left(\frac{\sqrt{dxc}}{\sqrt{-cd}}\right)}{\sqrt{-cdd^3}} + \frac{5 b \log\left(-\frac{cdx+d}{cdx-d}\right)}{\sqrt{dxd^3x^3}} + \frac{2 (10 bc^3 d^3 x^3 + 2 bcd^3 x + 5 ad^3)}{\sqrt{dxd^6x^3}}}{35 d}$$

input `integrate((a+b*arctanh(c*x))/(d*x)^(9/2),x, algorithm="giac")`output `-1/35*(10*b*c^4*arctan(sqrt(d*x)*c/sqrt(c*d))/(sqrt(c*d)*d^3) + 10*b*c^4*arctan(sqrt(d*x)*c/sqrt(-c*d))/(sqrt(-c*d)*d^3) + 5*b*log(-(c*d*x + d)/(c*d*x - d))/(sqrt(d*x)*d^3*x^3) + 2*(10*b*c^3*d^3*x^3 + 2*b*c*d^3*x + 5*a*d^3)/(sqrt(d*x)*d^6*x^3))/d`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{9/2}} dx = \int \frac{a + b \operatorname{atanh}(cx)}{(dx)^{9/2}} dx$$

input `int((a + b*atanh(c*x))/(d*x)^(9/2), x)`output `int((a + b*atanh(c*x))/(d*x)^(9/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.94

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(dx)^{9/2}} dx = \frac{\sqrt{d} \left(-10\sqrt{x} \sqrt{c} \operatorname{atan}\left(\frac{\sqrt{x}c}{\sqrt{c}}\right) b c^3 x^3 - 10\sqrt{x} \sqrt{c} \operatorname{atanh}(cx) b c^3 x^3 - 10 \operatorname{atanh}(cx) b \right)}{(dx)^{9/2}}$$

input `int((a+b*atanh(c*x))/(d*x)^(9/2), x)`output `(sqrt(d)*(- 10*sqrt(x)*sqrt(c)*atan((sqrt(x)*c)/sqrt(c))*b*c**3*x**3 - 10*sqrt(x)*sqrt(c)*atanh(c*x)*b*c**3*x**3 - 10*atanh(c*x)*b - 10*sqrt(x)*sqrt(c)*log(sqrt(x)*sqrt(c) - 1)*b*c**3*x**3 + 5*sqrt(x)*sqrt(c)*log(c*x + 1)*b*c**3*x**3 - 10*a - 20*b*c**3*x**3 - 4*b*c*x))/(35*sqrt(x)*d**5*x**3)`

3.43 $\int (dx)^m (a + \operatorname{barctanh}(cx))^3 dx$

Optimal result	428
Mathematica [N/A]	428
Rubi [N/A]	429
Maple [N/A]	429
Fricas [N/A]	430
Sympy [N/A]	430
Maxima [N/A]	430
Giac [N/A]	431
Mupad [N/A]	431
Reduce [N/A]	432

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int (dx)^m (a + \operatorname{barctanh}(cx))^3 dx = \operatorname{Int}((dx)^m (a + \operatorname{barctanh}(cx))^3, x)$$

output `Defer(Int)((d*x)^m*(a+b*arctanh(c*x))^3,x)`

Mathematica [N/A]

Not integrable

Time = 2.89 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (dx)^m (a + \operatorname{barctanh}(cx))^3 dx = \int (dx)^m (a + \operatorname{barctanh}(cx))^3 dx$$

input `Integrate[(d*x)^m*(a + b*ArcTanh[c*x])^3,x]`

output `Integrate[(d*x)^m*(a + b*ArcTanh[c*x])^3, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^3 dx$$

↓ 6468

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^3 dx$$

input `Int[(d*x)^m*(a + b*ArcTanh[c*x])^3,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^3 dx$$

input `int((d*x)^m*(a+b*arctanh(c*x))^3,x)`

output `int((d*x)^m*(a+b*arctanh(c*x))^3,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.75

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{artanh}(cx) + a)^3 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x))^3,x, algorithm="fricas")`

output `integral((b^3*arctanh(c*x)^3 + 3*a*b^2*arctanh(c*x)^2 + 3*a^2*b*arctanh(c*x) + a^3)*(d*x)^m, x)`

Sympy [N/A]

Not integrable

Time = 5.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^3 dx = \int (dx)^m (a + b \operatorname{atanh}(cx))^3 dx$$

input `integrate((d*x)**m*(a+b*atanh(c*x))**3,x)`

output `Integral((d*x)**m*(a + b*atanh(c*x))**3, x)`

Maxima [N/A]

Not integrable

Time = 2.22 (sec) , antiderivative size = 379, normalized size of antiderivative = 23.69

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{artanh}(cx) + a)^3 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x))^3,x, algorithm="maxima")`

output

```
-1/8*b^3*d^m*x*x^m*log(-c*x + 1)^3/(m + 1) + (d*x)^(m + 1)*a^3/(d*(m + 1))
+ integrate(1/8*((b^3*c*d^m*(m + 1)*x - b^3*d^m*(m + 1))*x^m*log(c*x + 1)
^3 + 6*(a*b^2*c*d^m*(m + 1)*x - a*b^2*d^m*(m + 1))*x^m*log(c*x + 1)^2 + 12
*(a^2*b*c*d^m*(m + 1)*x - a^2*b*d^m*(m + 1))*x^m*log(c*x + 1) + 3*((b^3*c*
d^m*(m + 1)*x - b^3*d^m*(m + 1))*x^m*log(c*x + 1) - (2*a*b^2*d^m*(m + 1) -
(2*a*b^2*c*d^m*(m + 1) + b^3*c*d^m)*x)*x^m*log(-c*x + 1)^2 - 3*((b^3*c*d
^m*(m + 1)*x - b^3*d^m*(m + 1))*x^m*log(c*x + 1)^2 + 4*(a*b^2*c*d^m*(m + 1)
)*x - a*b^2*d^m*(m + 1))*x^m*log(c*x + 1) + 4*(a^2*b*c*d^m*(m + 1)*x - a^2
*b*d^m*(m + 1))*x^m*log(-c*x + 1))/(c*(m + 1)*x - m - 1), x)
```

Giac [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^3 dx = \int (b \operatorname{artanh}(cx) + a)^3 (dx)^m dx$$

input

```
integrate((d*x)^m*(a+b*arctanh(c*x))^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x) + a)^3*(d*x)^m, x)
```

Mupad [N/A]

Not integrable

Time = 3.99 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^3 dx = \int (a + b \operatorname{atanh}(cx))^3 (dx)^m dx$$

input

```
int((a + b*atanh(c*x))^3*(d*x)^m,x)
```

output

```
int((a + b*atanh(c*x))^3*(d*x)^m, x)
```


3.44 $\int (dx)^m (a + \operatorname{barctanh}(cx))^2 dx$

Optimal result	433
Mathematica [N/A]	433
Rubi [N/A]	434
Maple [N/A]	434
Fricas [N/A]	435
Sympy [N/A]	435
Maxima [N/A]	435
Giac [N/A]	436
Mupad [N/A]	436
Reduce [N/A]	437

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int (dx)^m (a + \operatorname{barctanh}(cx))^2 dx = \operatorname{Int}((dx)^m (a + \operatorname{barctanh}(cx))^2, x)$$

output `Defer(Int)((d*x)^m*(a+b*arctanh(c*x))^2,x)`

Mathematica [N/A]

Not integrable

Time = 1.76 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (dx)^m (a + \operatorname{barctanh}(cx))^2 dx = \int (dx)^m (a + \operatorname{barctanh}(cx))^2 dx$$

input `Integrate[(d*x)^m*(a + b*ArcTanh[c*x])^2,x]`

output `Integrate[(d*x)^m*(a + b*ArcTanh[c*x])^2, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^2 dx$$

↓ 6468

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^2 dx$$

input `Int[(d*x)^m*(a + b*ArcTanh[c*x])^2,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^2 dx$$

input `int((d*x)^m*(a+b*arctanh(c*x))^2,x)`

output `int((d*x)^m*(a+b*arctanh(c*x))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.10 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.88

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^2 dx = \int (b \operatorname{artanh}(cx) + a)^2 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x))^2,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x)^2 + 2*a*b*arctanh(c*x) + a^2)*(d*x)^m, x)`

Sympy [N/A]

Not integrable

Time = 2.84 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^2 dx = \int (dx)^m (a + b \operatorname{atanh}(cx))^2 dx$$

input `integrate((d*x)**m*(a+b*atanh(c*x))**2,x)`

output `Integral((d*x)**m*(a + b*atanh(c*x))**2, x)`

Maxima [N/A]

Not integrable

Time = 1.15 (sec) , antiderivative size = 218, normalized size of antiderivative = 13.62

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^2 dx = \int (b \operatorname{artanh}(cx) + a)^2 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x))^2,x, algorithm="maxima")`

output

```
1/4*b^2*d^m*x^m*log(-c*x + 1)^2/(m + 1) + (d*x)^(m + 1)*a^2/(d*(m + 1))
- integrate(-1/4*((b^2*c*d^m*(m + 1)*x - b^2*d^m*(m + 1))*x^m*log(c*x + 1)
^2 + 4*(a*b*c*d^m*(m + 1)*x - a*b*d^m*(m + 1))*x^m*log(c*x + 1) - 2*((b^2*
c*d^m*(m + 1)*x - b^2*d^m*(m + 1))*x^m*log(c*x + 1) - (2*a*b*d^m*(m + 1) -
(2*a*b*c*d^m*(m + 1) + b^2*c*d^m)*x)*x^m*log(-c*x + 1))/(c*(m + 1)*x - m
- 1), x)
```

Giac [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^2 dx = \int (b \operatorname{arctanh}(cx) + a)^2 (dx)^m dx$$

input

```
integrate((d*x)^m*(a+b*arctanh(c*x))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x) + a)^2*(d*x)^m, x)
```

Mupad [N/A]

Not integrable

Time = 4.09 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^2 dx = \int (a + b \operatorname{atanh}(cx))^2 (dx)^m dx$$

input

```
int((a + b*atanh(c*x))^2*(d*x)^m,x)
```

output

```
int((a + b*atanh(c*x))^2*(d*x)^m, x)
```

Reduce [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 159, normalized size of antiderivative = 9.94

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{d^m (2x^m \operatorname{atanh}(cx) abcmx + x^m a^2 cmx + 2x^m ab + 2 \left(\int \frac{x^m}{c^2 m x^3 + c^2 x^3 - mx - x} dx \right) ab m^2 + 2 \left(\int \frac{x^m}{c^2 m x^3 + c^2 x^3 - mx - x} dx \right) cm(m+1)}{cm(m+1)}$$

input `int((d*x)^m*(a+b*atanh(c*x))^2,x)`output `(d**m*(2*x**m*atanh(c*x)*a*b*c*m*x + x**m*a**2*c*m*x + 2*x**m*a*b + 2*int(x**m/(c**2*m*x**3 + c**2*x**3 - m*x - x),x)*a*b*m**2 + 2*int(x**m/(c**2*m*x**3 + c**2*x**3 - m*x - x),x)*a*b*m + int(x**m*atanh(c*x)**2,x)*b**2*c*m**2 + int(x**m*atanh(c*x)**2,x)*b**2*c*m))/(c*m*(m + 1))`

3.45 $\int (dx)^m (a + \operatorname{barctanh}(cx)) dx$

Optimal result	438
Mathematica [A] (verified)	438
Rubi [A] (verified)	439
Maple [F]	440
Fricas [F]	440
Sympy [F]	441
Maxima [F]	441
Giac [F]	441
Mupad [F(-1)]	442
Reduce [F]	442

Optimal result

Integrand size = 14, antiderivative size = 72

$$\int (dx)^m (a + \operatorname{barctanh}(cx)) dx = \frac{(dx)^{1+m} (a + \operatorname{barctanh}(cx))}{d(1+m)} - \frac{bc(dx)^{2+m} \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, c^2x^2\right)}{d^2(1+m)(2+m)}$$

output `(d*x)^(1+m)*(a+b*arctanh(c*x))/d/(1+m)-b*c*(d*x)^(2+m)*hypergeom([1, 1+1/2*m], [2+1/2*m], c^2*x^2)/d^2/(1+m)/(2+m)`

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.82

$$\int (dx)^m (a + \operatorname{barctanh}(cx)) dx = \frac{x(dx)^m (-(2+m)(a + \operatorname{barctanh}(cx))) + bcx \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{m}{2}, 2 + \frac{m}{2}, c^2x^2\right)}{(1+m)(2+m)}$$

input `Integrate[(d*x)^m*(a + b*ArcTanh[c*x]), x]`

output

$$-\left(\frac{x(d*x)^m(-((2+m)(a+b*\text{ArcTanh}[c*x]))+b*c*x*\text{Hypergeometric2F1}[1, 1+m/2, 2+m/2, c^2*x^2])}{(1+m)(2+m)}\right)$$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6464, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx)) dx$$

$$\downarrow 6464$$

$$\frac{(dx)^{m+1} (a + b \operatorname{arctanh}(cx))}{d(m+1)} - \frac{bc \int \frac{(dx)^{m+1}}{1-c^2x^2} dx}{d(m+1)}$$

$$\downarrow 278$$

$$\frac{(dx)^{m+1} (a + b \operatorname{arctanh}(cx))}{d(m+1)} - \frac{bc(dx)^{m+2} \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{d^2(m+1)(m+2)}$$

input

$$\text{Int}[(d*x)^m*(a + b*\text{ArcTanh}[c*x]), x]$$

output

$$\frac{(d*x)^{(1+m)}*(a + b*\text{ArcTanh}[c*x])}{d*(1+m)} - \frac{(b*c*(d*x)^{(2+m)}*\text{Hypergeometric2F1}[1, (2+m)/2, (4+m)/2, c^2*x^2])}{d^2*(1+m)*(2+m)}$$

Definitions of rubi rules used

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 6464 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_.), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]`

Maple [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx)) dx$$

input `int((d*x)^m*(a+b*arctanh(c*x)),x)`

output `int((d*x)^m*(a+b*arctanh(c*x)),x)`

Fricas [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx)) dx = \int (b \operatorname{arctanh}(cx) + a)(dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x)),x, algorithm="fricas")`

output `integral((b*arctanh(c*x) + a)*(d*x)^m, x)`

Sympy [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx)) dx = \int (dx)^m (a + b \operatorname{atanh}(cx)) dx$$

input `integrate((d*x)**m*(a+b*atanh(c*x)),x)`

output `Integral((d*x)**m*(a + b*atanh(c*x)), x)`

Maxima [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx)) dx = \int (b \operatorname{artanh}(cx) + a)(dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x)),x, algorithm="maxima")`

output `1/2*(2*c*d^m*integrate(x*x^m/(c^2*(m + 1)*x^2 - m - 1), x) + (d^m*x*x^m*log(c*x + 1) - d^m*x*x^m*log(-c*x + 1))/(m + 1))*b + (d*x)^(m + 1)*a/(d*(m + 1))`

Giac [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx)) dx = \int (b \operatorname{artanh}(cx) + a)(dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x)),x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)*(d*x)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx)) dx = \int (a + b \operatorname{atanh}(cx)) (dx)^m dx$$

input `int((a + b*atanh(c*x))*(d*x)^m,x)`output `int((a + b*atanh(c*x))*(d*x)^m, x)`**Reduce [F]**

$$\int (dx)^m (a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{d^m (x^m \operatorname{atanh}(cx) b c m x + x^m a c m x + x^m b + \left(\int \frac{x^m}{c^2 m x^3 + c^2 x^3 - m x - x} dx \right) b m^2 + \left(\int \frac{x^m}{c^2 m x^3 + c^2 x^3 - m x - x} dx \right) b m)}{c m (m + 1)}$$

input `int((d*x)^m*(a+b*atanh(c*x)),x)`output `(d**m*(x**m*atanh(c*x)*b*c*m*x + x**m*a*c*m*x + x**m*b + int(x**m/(c**2*m*x**3 + c**2*x**3 - m*x - x),x)*b*m**2 + int(x**m/(c**2*m*x**3 + c**2*x**3 - m*x - x),x)*b*m))/(c*m*(m + 1))`

3.46 $\int \frac{(dx)^m}{a+b\mathbf{arctanh}(cx)} dx$

Optimal result	443
Mathematica [N/A]	443
Rubi [N/A]	444
Maple [N/A]	444
Fricas [N/A]	445
Sympy [N/A]	445
Maxima [N/A]	445
Giac [N/A]	446
Mupad [N/A]	446
Reduce [N/A]	447

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{(dx)^m}{a + b\mathbf{arctanh}(cx)} dx = \text{Int}\left(\frac{(dx)^m}{a + b\mathbf{arctanh}(cx)}, x\right)$$

output `Defer(Int)((d*x)^m/(a+b*arctanh(c*x)),x)`

Mathematica [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(dx)^m}{a + b\mathbf{arctanh}(cx)} dx = \int \frac{(dx)^m}{a + b\mathbf{arctanh}(cx)} dx$$

input `Integrate[(d*x)^m/(a + b*ArcTanh[c*x]),x]`

output `Integrate[(d*x)^m/(a + b*ArcTanh[c*x]), x]`

Rubi [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx)} dx$$

↓ 6468

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx)} dx$$

input `Int[(d*x)^m/(a + b*ArcTanh[c*x]),x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.60 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx)} dx$$

input `int((d*x)^m/(a+b*arctanh(c*x)),x)`

output `int((d*x)^m/(a+b*arctanh(c*x)),x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx)} dx = \int \frac{(dx)^m}{b \operatorname{artanh}(cx) + a} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x)),x, algorithm="fricas")`

output `integral((d*x)^m/(b*arctanh(c*x) + a), x)`

Sympy [N/A]

Not integrable

Time = 1.31 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx)} dx = \int \frac{(dx)^m}{a + b \operatorname{atanh}(cx)} dx$$

input `integrate((d*x)**m/(a+b*atanh(c*x)),x)`

output `Integral((d*x)**m/(a + b*atanh(c*x)), x)`

Maxima [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx)} dx = \int \frac{(dx)^m}{b \operatorname{artanh}(cx) + a} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x)),x, algorithm="maxima")`

output `integrate((d*x)^m/(b*arctanh(c*x) + a), x)`

Giac [N/A]

Not integrable

Time = 0.13 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx)} dx = \int \frac{(dx)^m}{b \operatorname{artanh}(cx) + a} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x)),x, algorithm="giac")`

output `integrate((d*x)^m/(b*arctanh(c*x) + a), x)`

Mupad [N/A]

Not integrable

Time = 3.74 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx)} dx = \int \frac{(dx)^m}{a + b \operatorname{atanh}(cx)} dx$$

input `int((d*x)^m/(a + b*atanh(c*x)),x)`

output `int((d*x)^m/(a + b*atanh(c*x)), x)`

Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx)} dx = d^m \left(\int \frac{x^m}{\operatorname{atanh}(cx) b + a} dx \right)$$

input `int((d*x)^m/(a+b*atanh(c*x)),x)`output `d**m*int(x**m/(atanh(c*x)*b + a),x)`

$$3.47 \quad \int \frac{(dx)^m}{(a+b\operatorname{arctanh}(cx))^2} dx$$

Optimal result	448
Mathematica [N/A]	448
Rubi [N/A]	449
Maple [N/A]	449
Fricas [N/A]	450
Sympy [N/A]	450
Maxima [N/A]	450
Giac [N/A]	451
Mupad [N/A]	451
Reduce [N/A]	452

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{(dx)^m}{(a + \operatorname{arctanh}(cx))^2} dx = \operatorname{Int}\left(\frac{(dx)^m}{(a + \operatorname{arctanh}(cx))^2}, x\right)$$

output `Defer(Int)((d*x)^m/(a+b*arctanh(c*x))^2,x)`

Mathematica [N/A]

Not integrable

Time = 0.43 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(dx)^m}{(a + \operatorname{arctanh}(cx))^2} dx = \int \frac{(dx)^m}{(a + \operatorname{arctanh}(cx))^2} dx$$

input `Integrate[(d*x)^m/(a + b*ArcTanh[c*x])^2,x]`

output `Integrate[(d*x)^m/(a + b*ArcTanh[c*x])^2, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx))^2} dx$$

↓ 6468

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx))^2} dx$$

input `Int[(d*x)^m/(a + b*ArcTanh[c*x])^2,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.57 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx))^2} dx$$

input `int((d*x)^m/(a+b*arctanh(c*x))^2,x)`

output `int((d*x)^m/(a+b*arctanh(c*x))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 32, normalized size of antiderivative = 2.00

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx))^2} dx = \int \frac{(dx)^m}{(b \operatorname{artanh}(cx) + a)^2} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x))^2,x, algorithm="fricas")`

output `integral((d*x)^m/(b^2*arctanh(c*x)^2 + 2*a*b*arctanh(c*x) + a^2), x)`

Sympy [N/A]

Not integrable

Time = 9.55 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx))^2} dx = \int \frac{(dx)^m}{(a + b \operatorname{atanh}(cx))^2} dx$$

input `integrate((d*x)**m/(a+b*atanh(c*x))**2,x)`

output `Integral((d*x)**m/(a + b*atanh(c*x))**2, x)`

Maxima [N/A]

Not integrable

Time = 0.13 (sec) , antiderivative size = 116, normalized size of antiderivative = 7.25

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx))^2} dx = \int \frac{(dx)^m}{(b \operatorname{artanh}(cx) + a)^2} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x))^2,x, algorithm="maxima")`

output

```
2*(c^2*d^m*x^2 - d^m)*x^m/(b^2*c*log(c*x + 1) - b^2*c*log(-c*x + 1) + 2*a*
b*c) + integrate(-2*(c^2*d^m*(m + 2)*x^2 - d^m*m)*x^m/(b^2*c*x*log(c*x + 1)
) - b^2*c*x*log(-c*x + 1) + 2*a*b*c*x), x)
```

Giac [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx))^2} dx = \int \frac{(dx)^m}{(b \operatorname{artanh}(cx) + a)^2} dx$$

input

```
integrate((d*x)^m/(a+b*arctanh(c*x))^2,x, algorithm="giac")
```

output

```
integrate((d*x)^m/(b*arctanh(c*x) + a)^2, x)
```

Mupad [N/A]

Not integrable

Time = 3.65 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx))^2} dx = \int \frac{(dx)^m}{(a + b \operatorname{atanh}(cx))^2} dx$$

input

```
int((d*x)^m/(a + b*atanh(c*x))^2,x)
```

output

```
int((d*x)^m/(a + b*atanh(c*x))^2, x)
```


Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.12

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx))^2} dx = d^m \left(\int \frac{x^m}{\operatorname{atanh}(cx)^2 b^2 + 2 \operatorname{atanh}(cx) ab + a^2} dx \right)$$

input `int((d*x)^m/(a+b*atanh(c*x))^2,x)`output `d**m*int(x**m/(atanh(c*x)**2*b**2 + 2*atanh(c*x)*a*b + a**2),x)`

3.48 $\int (a + b \operatorname{arctanh}(cx))^p dx$

Optimal result	453
Mathematica [N/A]	453
Rubi [N/A]	454
Maple [N/A]	454
Fricas [N/A]	455
Sympy [N/A]	455
Maxima [N/A]	455
Giac [N/A]	456
Mupad [N/A]	456
Reduce [N/A]	457

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int (a + b \operatorname{arctanh}(cx))^p dx = \operatorname{Int}((a + b \operatorname{arctanh}(cx))^p, x)$$

output `Defer(Int)((a+b*arctanh(c*x))^p,x)`

Mathematica [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int (a + b \operatorname{arctanh}(cx))^p dx = \int (a + b \operatorname{arctanh}(cx))^p dx$$

input `Integrate[(a + b*ArcTanh[c*x])^p,x]`

output `Integrate[(a + b*ArcTanh[c*x])^p, x]`

Rubi [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{arctanh}(cx))^p dx$$

↓ 6444

$$\int (a + b \operatorname{arctanh}(cx))^p dx$$

input `Int[(a + b*ArcTanh[c*x])^p,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int (a + b \operatorname{arctanh}(cx))^p dx$$

input `int((a+b*arctanh(c*x))^p,x)`

output `int((a+b*arctanh(c*x))^p,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int (a + b \operatorname{arctanh}(cx))^p dx = \int (b \operatorname{artanh}(cx) + a)^p dx$$

input `integrate((a+b*arctanh(c*x))^p,x, algorithm="fricas")`

output `integral((b*arctanh(c*x) + a)^p, x)`

Sympy [N/A]

Not integrable

Time = 1.72 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int (a + b \operatorname{arctanh}(cx))^p dx = \int (a + b \operatorname{atanh}(cx))^p dx$$

input `integrate((a+b*atanh(c*x))**p,x)`

output `Integral((a + b*atanh(c*x))**p, x)`

Maxima [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int (a + b \operatorname{arctanh}(cx))^p dx = \int (b \operatorname{artanh}(cx) + a)^p dx$$

input `integrate((a+b*arctanh(c*x))^p,x, algorithm="maxima")`

output `integrate((b*arctanh(c*x) + a)^p, x)`

Giac [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int (a + b \operatorname{arctanh}(cx))^p dx = \int (b \operatorname{artanh}(cx) + a)^p dx$$

input `integrate((a+b*arctanh(c*x))^p,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^p, x)`

Mupad [N/A]

Not integrable

Time = 3.62 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int (a + b \operatorname{arctanh}(cx))^p dx = \int (a + b \operatorname{atanh}(cx))^p dx$$

input `int((a + b*atanh(c*x))^p,x)`

output `int((a + b*atanh(c*x))^p, x)`

Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int (a + b \operatorname{arctanh}(cx))^p dx = \int (a \operatorname{tanh}(cx) b + a)^p dx$$

input `int((a+b*atanh(c*x))^p,x)`output `int((atanh(c*x)*b + a)**p,x)`

3.49 $\int (dx)^m (a + \operatorname{barctanh}(cx))^p dx$

Optimal result	458
Mathematica [N/A]	458
Rubi [N/A]	459
Maple [N/A]	459
Fricas [N/A]	460
Sympy [F(-1)]	460
Maxima [N/A]	460
Giac [N/A]	461
Mupad [N/A]	461
Reduce [N/A]	461

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int (dx)^m (a + \operatorname{barctanh}(cx))^p dx = \operatorname{Int}((dx)^m (a + \operatorname{barctanh}(cx))^p, x)$$

output `Defer(Int)((d*x)^m*(a+b*arctanh(c*x))^p,x)`

Mathematica [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (dx)^m (a + \operatorname{barctanh}(cx))^p dx = \int (dx)^m (a + \operatorname{barctanh}(cx))^p dx$$

input `Integrate[(d*x)^m*(a + b*ArcTanh[c*x])^p,x]`

output `Integrate[(d*x)^m*(a + b*ArcTanh[c*x])^p, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^p dx$$

↓ 6468

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^p dx$$

input `Int[(d*x)^m*(a + b*ArcTanh[c*x])^p,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.55 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^p dx$$

input `int((d*x)^m*(a+b*arctanh(c*x))^p,x)`

output `int((d*x)^m*(a+b*arctanh(c*x))^p,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^p dx = \int (dx)^m (b \operatorname{arctanh}(cx) + a)^p dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x))^p,x, algorithm="fricas")`

output `integral((d*x)^m*(b*arctanh(c*x) + a)^p, x)`

Sympy [F(-1)]

Timed out.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^p dx = \text{Timed out}$$

input `integrate((d*x)**m*(a+b*atanh(c*x))**p,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^p dx = \int (dx)^m (b \operatorname{arctanh}(cx) + a)^p dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x))^p,x, algorithm="maxima")`

output `integrate((d*x)^m*(b*arctanh(c*x) + a)^p, x)`

Giac [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^p dx = \int (dx)^m (b \operatorname{artanh}(cx) + a)^p dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x))^p,x, algorithm="giac")`

output `integrate((d*x)^m*(b*arctanh(c*x) + a)^p, x)`

Mupad [N/A]

Not integrable

Time = 3.63 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^p dx = \int (a + b \operatorname{atanh}(cx))^p (dx)^m dx$$

input `int((a + b*atanh(c*x))^p*(d*x)^m,x)`

output `int((a + b*atanh(c*x))^p*(d*x)^m, x)`

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int (dx)^m (a + b \operatorname{arctanh}(cx))^p dx = d^m \left(\int x^m (\operatorname{atanh}(cx) b + a)^p dx \right)$$

input `int((d*x)^m*(a+b*atanh(c*x))^p,x)`

output `d**m*int(x**m*(atanh(c*x)*b + a)**p,x)`

3.50 $\int x^7(a + \operatorname{barctanh}(cx^2)) dx$

Optimal result	463
Mathematica [A] (verified)	463
Rubi [A] (verified)	464
Maple [A] (verified)	465
Fricas [A] (verification not implemented)	466
Sympy [A] (verification not implemented)	466
Maxima [A] (verification not implemented)	467
Giac [A] (verification not implemented)	467
Mupad [B] (verification not implemented)	468
Reduce [B] (verification not implemented)	468

Optimal result

Integrand size = 14, antiderivative size = 54

$$\int x^7(a + \operatorname{barctanh}(cx^2)) dx = \frac{bx^2}{8c^3} + \frac{bx^6}{24c} - \frac{\operatorname{barctanh}(cx^2)}{8c^4} + \frac{1}{8}x^8(a + \operatorname{barctanh}(cx^2))$$

output

```
1/8*b*x^2/c^3+1/24*b*x^6/c-1/8*b*arctanh(c*x^2)/c^4+1/8*x^8*(a+b*arctanh(c*x^2))
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.44

$$\int x^7(a + \operatorname{barctanh}(cx^2)) dx = \frac{bx^2}{8c^3} + \frac{bx^6}{24c} + \frac{ax^8}{8} + \frac{1}{8}bx^8\operatorname{arctanh}(cx^2) + \frac{b \log(1 - cx^2)}{16c^4} - \frac{b \log(1 + cx^2)}{16c^4}$$

input

```
Integrate[x^7*(a + b*ArcTanh[c*x^2]),x]
```

output

```
(b*x^2)/(8*c^3) + (b*x^6)/(24*c) + (a*x^8)/8 + (b*x^8*ArcTanh[c*x^2])/8 + (b*Log[1 - c*x^2])/(16*c^4) - (b*Log[1 + c*x^2])/(16*c^4)
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.98, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6452, 807, 254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^7(a + b \operatorname{arctanh}(cx^2)) dx \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{8}x^8(a + b \operatorname{arctanh}(cx^2)) - \frac{1}{4}bc \int \frac{x^9}{1 - c^2x^4} dx \\
 & \quad \downarrow \text{807} \\
 & \frac{1}{8}x^8(a + b \operatorname{arctanh}(cx^2)) - \frac{1}{8}bc \int \frac{x^8}{1 - c^2x^4} dx^2 \\
 & \quad \downarrow \text{254} \\
 & \frac{1}{8}x^8(a + b \operatorname{arctanh}(cx^2)) - \frac{1}{8}bc \int \left(-\frac{x^4}{c^2} + \frac{1}{c^4(1 - c^2x^4)} - \frac{1}{c^4} \right) dx^2 \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{8}x^8(a + b \operatorname{arctanh}(cx^2)) - \frac{1}{8}bc \left(\frac{\operatorname{arctanh}(cx^2)}{c^5} - \frac{x^2}{c^4} - \frac{x^6}{3c^2} \right)
 \end{aligned}$$

input `Int[x^7*(a + b*ArcTanh[c*x^2]),x]`

output `(x^8*(a + b*ArcTanh[c*x^2]))/8 - (b*c*(-(x^2/c^4) - x^6/(3*c^2) + ArcTanh[c*x^2]/c^5))/8`

Definitions of rubi rules used

rule 254 $\text{Int}[(x_)^{(m_)} / ((a_) + (b_)*(x_)^2), x_Symbol] \rightarrow \text{Int}[\text{PolynomialDivide}[x^m, a + b*x^2, x], x] /;$ FreeQ[{a, b}, x] && IGtQ[m, 3]

rule 807 $\text{Int}[(x_)^{(m_)} * ((a_) + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{With}[\{k = \text{GCD}[m + 1, n]\}, \text{Simp}[1/k \text{ Subst}[\text{Int}[x^{(m+1)/k - 1} * (a + b*x^{(n/k)})^p, x], x, x^k], x] /;$ k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /;$ SumQ[u]

rule 6452 $\text{Int}[(a_) + \text{ArcTanh}[(c_)*(x_)^{(n_)}] * (b_)]^{(p_)} * (x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)} * ((a + b*\text{ArcTanh}[c*x^n])^{(p/(m+1))}), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)} * ((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{(2*n)})}), x], x] /;$ FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.04

method	result	size
parallelrisch	$-\frac{-3b \operatorname{arctanh}(cx^2)x^8c^4 - 3ac^4x^8 - b^3c^3x^6 - 3bcx^2 + 3b \operatorname{arctanh}(cx^2)}{24c^4}$	56
default	$\frac{ax^8}{8} + \frac{bx^8 \operatorname{arctanh}(cx^2)}{8} + \frac{bx^6}{24c} + \frac{bx^2}{8c^3} - \frac{b \ln(cx^2+1)}{16c^4} + \frac{b \ln(cx^2-1)}{16c^4}$	66
parts	$\frac{ax^8}{8} + \frac{bx^8 \operatorname{arctanh}(cx^2)}{8} + \frac{bx^6}{24c} + \frac{bx^2}{8c^3} - \frac{b \ln(cx^2+1)}{16c^4} + \frac{b \ln(cx^2-1)}{16c^4}$	66
risch	$\frac{x^8 b \ln(cx^2+1)}{16} - \frac{x^8 b \ln(-cx^2+1)}{16} + \frac{ax^8}{8} + \frac{bx^6}{24c} + \frac{bx^2}{8c^3} + \frac{b \ln(cx^2-1)}{16c^4} - \frac{b \ln(cx^2+1)}{16c^4}$	83
orering	$\frac{(13c^4x^8 + 14c^2x^4 - 27)(a + b \operatorname{arctanh}(cx^2))}{48c^4} - \frac{(c^2x^4 + 3)(cx^2 - 1)(cx^2 + 1)(7x^6(a + b \operatorname{arctanh}(cx^2)) + \frac{2x^8bc}{-c^2x^4 + 1})}{48x^6c^4}$	101

input $\text{int}(x^7*(a+b*\operatorname{arctanh}(c*x^2)), x, \text{method}=_RETURNVERBOSE)$

output

```
-1/24*(-3*b*arctanh(c*x^2)*x^8*c^4-3*a*c^4*x^8-b*c^3*x^6-3*b*c*x^2+3*b*arc
tanh(c*x^2))/c^4
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.19

$$\int x^7(a + b \operatorname{arctanh}(cx^2)) dx = \frac{6ac^4x^8 + 2bc^3x^6 + 6bcx^2 + 3(bc^4x^8 - b) \log\left(-\frac{cx^2+1}{cx^2-1}\right)}{48c^4}$$

input

```
integrate(x^7*(a+b*arctanh(c*x^2)),x, algorithm="fricas")
```

output

```
1/48*(6*a*c^4*x^8 + 2*b*c^3*x^6 + 6*b*c*x^2 + 3*(b*c^4*x^8 - b)*log(-(c*x^
2 + 1)/(c*x^2 - 1)))/c^4
```

Sympy [A] (verification not implemented)

Time = 6.08 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.07

$$\int x^7(a + b \operatorname{atanh}(cx^2)) dx = \begin{cases} \frac{ax^8}{8} + \frac{bx^8 \operatorname{atanh}(cx^2)}{8} + \frac{bx^6}{24c} + \frac{bx^2}{8c^3} - \frac{b \operatorname{atanh}(cx^2)}{8c^4} & \text{for } c \neq 0 \\ \frac{ax^8}{8} & \text{otherwise} \end{cases}$$

input

```
integrate(x**7*(a+b*atanh(c*x**2)),x)
```

output

```
Piecewise((a*x**8/8 + b*x**8*atanh(c*x**2)/8 + b*x**6/(24*c) + b*x**2/(8*c
**3) - b*atanh(c*x**2)/(8*c**4), Ne(c, 0)), (a*x**8/8, True))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.28

$$\int x^7 (a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{1}{8} ax^8$$

$$+ \frac{1}{48} \left(6x^8 \operatorname{artanh}(cx^2) + c \left(\frac{2(c^2x^6 + 3x^2)}{c^4} - \frac{3 \log(cx^2 + 1)}{c^5} + \frac{3 \log(cx^2 - 1)}{c^5} \right) \right) b$$

input `integrate(x^7*(a+b*arctanh(c*x^2)),x, algorithm="maxima")`

output `1/8*a*x^8 + 1/48*(6*x^8*arctanh(c*x^2) + c*(2*(c^2*x^6 + 3*x^2)/c^4 - 3*log(c*x^2 + 1)/c^5 + 3*log(c*x^2 - 1)/c^5))*b`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.44

$$\int x^7 (a + b \operatorname{arctanh}(cx^2)) dx = \frac{1}{16} bx^8 \log\left(-\frac{cx^2 + 1}{cx^2 - 1}\right) + \frac{1}{8} ax^8 + \frac{bx^6}{24c}$$

$$+ \frac{bx^2}{8c^3} - \frac{b \log(cx^2 + 1)}{16c^4} + \frac{b \log(cx^2 - 1)}{16c^4}$$

input `integrate(x^7*(a+b*arctanh(c*x^2)),x, algorithm="giac")`

output `1/16*b*x^8*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 1/8*a*x^8 + 1/24*b*x^6/c + 1/8*b*x^2/c^3 - 1/16*b*log(c*x^2 + 1)/c^4 + 1/16*b*log(c*x^2 - 1)/c^4`

Mupad [B] (verification not implemented)

Time = 3.88 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.28

$$\int x^7(a + b \operatorname{arctanh}(cx^2)) dx = \frac{ax^8}{8} + \frac{bx^2}{8c^3} + \frac{bx^6}{24c} + \frac{bx^8 \ln(cx^2 + 1)}{16} - \frac{bx^8 \ln(1 - cx^2)}{16} + \frac{b \operatorname{atan}(cx^2 \operatorname{li}) \operatorname{li}}{8c^4}$$

input `int(x^7*(a + b*atanh(c*x^2)),x)`output `(a*x^8)/8 + (b*x^2)/(8*c^3) + (b*x^6)/(24*c) + (b*atan(c*x^2*1i)*1i)/(8*c^4) + (b*x^8*log(c*x^2 + 1))/16 - (b*x^8*log(1 - c*x^2))/16`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00

$$\int x^7(a + b \operatorname{arctanh}(cx^2)) dx = \frac{3 \operatorname{atanh}(cx^2) b c^4 x^8 - 3 \operatorname{atanh}(cx^2) b + 3 a c^4 x^8 + b c^3 x^6 + 3 b c x^2}{24 c^4}$$

input `int(x^7*(a+b*atanh(c*x^2)),x)`output `(3*atanh(c*x**2)*b*c**4*x**8 - 3*atanh(c*x**2)*b + 3*a*c**4*x**8 + b*c**3*x**6 + 3*b*c*x**2)/(24*c**4)`

3.51 $\int x^5(a + \operatorname{barctanh}(cx^2)) dx$

Optimal result	469
Mathematica [A] (verified)	469
Rubi [A] (verified)	470
Maple [A] (verified)	471
Fricas [A] (verification not implemented)	472
Sympy [B] (verification not implemented)	472
Maxima [A] (verification not implemented)	473
Giac [A] (verification not implemented)	473
Mupad [B] (verification not implemented)	473
Reduce [B] (verification not implemented)	474

Optimal result

Integrand size = 14, antiderivative size = 48

$$\int x^5(a + \operatorname{barctanh}(cx^2)) dx = \frac{bx^4}{12c} + \frac{1}{6}x^6(a + \operatorname{barctanh}(cx^2)) + \frac{b \log(1 - c^2x^4)}{12c^3}$$

output

```
1/12*b*x^4/c+1/6*x^6*(a+b*arctanh(c*x^2))+1/12*b*ln(-c^2*x^4+1)/c^3
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.10

$$\int x^5(a + \operatorname{barctanh}(cx^2)) dx = \frac{bx^4}{12c} + \frac{ax^6}{6} + \frac{1}{6}bx^6\operatorname{arctanh}(cx^2) + \frac{b \log(1 - c^2x^4)}{12c^3}$$

input

```
Integrate[x^5*(a + b*ArcTanh[c*x^2]),x]
```

output

```
(b*x^4)/(12*c) + (a*x^6)/6 + (b*x^6*ArcTanh[c*x^2])/6 + (b*Log[1 - c^2*x^4])/12*c^3
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6452, 798, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^5(a + b \operatorname{arctanh}(cx^2)) dx$$

$$\downarrow 6452$$

$$\frac{1}{6}x^6(a + b \operatorname{arctanh}(cx^2)) - \frac{1}{3}bc \int \frac{x^7}{1 - c^2x^4} dx$$

$$\downarrow 798$$

$$\frac{1}{6}x^6(a + b \operatorname{arctanh}(cx^2)) - \frac{1}{12}bc \int \frac{x^4}{1 - c^2x^4} dx^4$$

$$\downarrow 49$$

$$\frac{1}{6}x^6(a + b \operatorname{arctanh}(cx^2)) - \frac{1}{12}bc \int \left(-\frac{1}{c^2} - \frac{1}{c^2(c^2x^4 - 1)} \right) dx^4$$

$$\downarrow 2009$$

$$\frac{1}{6}x^6(a + b \operatorname{arctanh}(cx^2)) - \frac{1}{12}bc \left(-\frac{x^4}{c^2} - \frac{\log(1 - c^2x^4)}{c^4} \right)$$

input `Int [x^5*(a + b*ArcTanh[c*x^2]), x]`

output `(x^6*(a + b*ArcTanh[c*x^2]))/6 - (b*c*(-(x^4/c^2) - Log[1 - c^2*x^4]/c^4)) /12`

Definitions of rubi rules used

rule 49 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}*((c_.) + (d_.)(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{IGtQ}[m, 0] \&\& \text{IGtQ}[m + n + 2, 0]$

rule 798 $\text{Int}[(x_)^{(m_.)}*((a_) + (b_.)(x_)^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n) - 1}*(a + b*x)^p, x], x, x^n], x] /; \text{FreeQ}[\{a, b, m, n, p\}, x] \&\& \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 6452 $\text{Int}[(a_.) + \text{ArcTanh}[(c_.)(x_)^{(n_.)}]*(b_.)]^{(p_.)}(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m + 1)), x] - \text{Simp}[b*c*n*(p/(m + 1)) \text{ Int}[x^{(m + n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p - 1)/(1 - c^2*x^{(2*n)})}), x], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \&\& \text{IGtQ}[p, 0] \&\& (\text{EqQ}[p, 1] \|\| (\text{EqQ}[n, 1] \&\& \text{IntegerQ}[m])) \&\& \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.94

method	result	size
default	$\frac{ax^6}{6} + \frac{bx^6 \operatorname{arctanh}(cx^2)}{6} + \frac{bx^4}{12c} + \frac{b \ln(c^2x^4 - 1)}{12c^3}$	45
parts	$\frac{ax^6}{6} + \frac{bx^6 \operatorname{arctanh}(cx^2)}{6} + \frac{bx^4}{12c} + \frac{b \ln(c^2x^4 - 1)}{12c^3}$	45
parallelrisch	$\frac{2b \operatorname{arctanh}(cx^2)x^6c^3 + 2ac^3x^6 + bc^2x^4 + 2b \ln(cx^2 - 1) + 2b \operatorname{arctanh}(cx^2)}{12c^3}$	59
risch	$\frac{bx^6 \ln(cx^2 + 1)}{12} - \frac{bx^6 \ln(-cx^2 + 1)}{12} + \frac{ax^6}{6} + \frac{bx^4}{12c} + \frac{b \ln(c^2x^4 - 1)}{12c^3}$	62

input `int(x^5*(a+b*arctanh(c*x^2)),x,method=_RETURNVERBOSE)`

output `1/6*a*x^6+1/6*b*x^6*arctanh(c*x^2)+1/12*b*x^4/c+1/12*b/c^3*ln(c^2*x^4-1)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.29

$$\int x^5(a + b \operatorname{arctanh}(cx^2)) dx = \frac{bc^3x^6 \log\left(-\frac{cx^2+1}{cx^2-1}\right) + 2ac^3x^6 + bc^2x^4 + b \log(c^2x^4 - 1)}{12c^3}$$

input `integrate(x^5*(a+b*arctanh(c*x^2)),x, algorithm="fricas")`

output `1/12*(b*c^3*x^6*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 2*a*c^3*x^6 + b*c^2*x^4 + b*log(c^2*x^4 - 1))/c^3`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 85 vs. 2(39) = 78.

Time = 4.40 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.77

$$\int x^5(a + b \operatorname{arctanh}(cx^2)) dx = \begin{cases} \frac{ax^6}{6} + \frac{bx^6 \operatorname{atanh}(cx^2)}{6} + \frac{bx^4}{12c} + \frac{b \log\left(x - \sqrt{-\frac{1}{c}}\right)}{6c^3} + \frac{b \log\left(x + \sqrt{-\frac{1}{c}}\right)}{6c^3} - \frac{b \operatorname{atanh}(cx^2)}{6c^3} & \text{for } c \neq 0 \\ \frac{ax^6}{6} & \text{otherwise} \end{cases}$$

input `integrate(x**5*(a+b*atanh(c*x**2)),x)`

output `Piecewise((a*x**6/6 + b*x**6*atanh(c*x**2)/6 + b*x**4/(12*c) + b*log(x - sqrt(-1/c))/(6*c**3) + b*log(x + sqrt(-1/c))/(6*c**3) - b*atanh(c*x**2)/(6*c**3), Ne(c, 0)), (a*x**6/6, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.96

$$\int x^5 (a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{1}{6} ax^6 + \frac{1}{12} \left(2x^6 \operatorname{arctanh}(cx^2) + \left(\frac{x^4}{c^2} + \frac{\log(c^2 x^4 - 1)}{c^4} \right) c \right) b$$

input `integrate(x^5*(a+b*arctanh(c*x^2)),x, algorithm="maxima")`output `1/6*a*x^6 + 1/12*(2*x^6*arctanh(c*x^2) + (x^4/c^2 + log(c^2*x^4 - 1)/c^4)*c)*b`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.19

$$\int x^5 (a + b \operatorname{arctanh}(cx^2)) dx = \frac{1}{12} bx^6 \log\left(-\frac{cx^2 + 1}{cx^2 - 1}\right) + \frac{1}{6} ax^6 + \frac{bx^4}{12c} + \frac{b \log(c^2 x^4 - 1)}{12c^3}$$

input `integrate(x^5*(a+b*arctanh(c*x^2)),x, algorithm="giac")`output `1/12*b*x^6*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 1/6*a*x^6 + 1/12*b*x^4/c + 1/12*b*log(c^2*x^4 - 1)/c^3`**Mupad [B] (verification not implemented)**

Time = 3.64 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.27

$$\int x^5 (a + b \operatorname{arctanh}(cx^2)) dx = \frac{ax^6}{6} + \frac{b \ln(c^2 x^4 - 1)}{12c^3} + \frac{bx^4}{12c}$$

$$+ \frac{bx^6 \ln(cx^2 + 1)}{12} - \frac{bx^6 \ln(1 - cx^2)}{12}$$

input `int(x^5*(a + b*atanh(c*x^2)),x)`

output

$$(a*x^6)/6 + (b*\log(c^2*x^4 - 1))/(12*c^3) + (b*x^4)/(12*c) + (b*x^6*\log(c*x^2 + 1))/12 - (b*x^6*\log(1 - c*x^2))/12$$
Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.21

$$\int x^5(a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{2 \operatorname{atanh}(cx^2) b c^3 x^6 - 2 \operatorname{atanh}(cx^2) b + 2 \log(cx^2 + 1) b + 2 a c^3 x^6 + b c^2 x^4}{12 c^3}$$

input

$$\operatorname{int}(x^5*(a+b*\operatorname{atanh}(c*x^2)),x)$$

output

$$(2*\operatorname{atanh}(c*x**2)*b*c**3*x**6 - 2*\operatorname{atanh}(c*x**2)*b + 2*\log(c*x**2 + 1)*b + 2*a*c**3*x**6 + b*c**2*x**4)/(12*c**3)$$

3.52 $\int x^3(a + \operatorname{barctanh}(cx^2)) dx$

Optimal result	475
Mathematica [A] (verified)	475
Rubi [A] (verified)	476
Maple [A] (verified)	477
Fricas [A] (verification not implemented)	478
Sympy [A] (verification not implemented)	478
Maxima [A] (verification not implemented)	479
Giac [B] (verification not implemented)	479
Mupad [B] (verification not implemented)	480
Reduce [B] (verification not implemented)	480

Optimal result

Integrand size = 14, antiderivative size = 43

$$\int x^3(a + \operatorname{barctanh}(cx^2)) dx = \frac{bx^2}{4c} - \frac{\operatorname{barctanh}(cx^2)}{4c^2} + \frac{1}{4}x^4(a + \operatorname{barctanh}(cx^2))$$

output `1/4*b*x^2/c-1/4*b*arctanh(c*x^2)/c^2+1/4*x^4*(a+b*arctanh(c*x^2))`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.56

$$\int x^3(a + \operatorname{barctanh}(cx^2)) dx = \frac{bx^2}{4c} + \frac{ax^4}{4} + \frac{1}{4}bx^4\operatorname{arctanh}(cx^2) + \frac{b \log(1 - cx^2)}{8c^2} - \frac{b \log(1 + cx^2)}{8c^2}$$

input `Integrate[x^3*(a + b*ArcTanh[c*x^2]),x]`

output `(b*x^2)/(4*c) + (a*x^4)/4 + (b*x^4*ArcTanh[c*x^2])/4 + (b*Log[1 - c*x^2])/(8*c^2) - (b*Log[1 + c*x^2])/(8*c^2)`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6452, 807, 262, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3(a + \operatorname{arctanh}(cx^2)) dx \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{4}x^4(a + \operatorname{arctanh}(cx^2)) - \frac{1}{2}bc \int \frac{x^5}{1 - c^2x^4} dx \\
 & \quad \downarrow \text{807} \\
 & \frac{1}{4}x^4(a + \operatorname{arctanh}(cx^2)) - \frac{1}{4}bc \int \frac{x^4}{1 - c^2x^4} dx \\
 & \quad \downarrow \text{262} \\
 & \frac{1}{4}x^4(a + \operatorname{arctanh}(cx^2)) - \frac{1}{4}bc \left(\frac{\int \frac{1}{1 - c^2x^4} dx^2}{c^2} - \frac{x^2}{c^2} \right) \\
 & \quad \downarrow \text{219} \\
 & \frac{1}{4}x^4(a + \operatorname{arctanh}(cx^2)) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(cx^2)}{c^3} - \frac{x^2}{c^2} \right)
 \end{aligned}$$

input `Int [x^3*(a + b*ArcTanh[c*x^2]), x]`

output `(x^4*(a + b*ArcTanh[c*x^2]))/4 - (b*c*(-(x^2/c^2) + ArcTanh[c*x^2]/c^3))/4`

Definitions of rubi rules used

rule 219 $\text{Int}[\{(a_)+(b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{[a, b], x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 262 $\text{Int}[\{(c_)*(x_)\}^{(m_)}*\{(a_)+(b_)*(x_)^2\}^{(p_)}, x_Symbol] \rightarrow \text{Simp}[c*(c*x)^{(m-1)}*((a+b*x^2)^{(p+1)}/(b*(m+2*p+1))), x] - \text{Simp}[a*c^2*((m-1)/(b*(m+2*p+1))) \ \text{Int}[(c*x)^{(m-2)}*(a+b*x^2)^p, x], x] /; \text{FreeQ}\{[a, b, c, p], x\} \ \&\& \ \text{GtQ}[m, 2-1] \ \&\& \ \text{NeQ}[m+2*p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 807 $\text{Int}[(x_)^{(m_)}*\{(a_)+(b_)*(x_)^n\}^{(p_)}, x_Symbol] \rightarrow \text{With}[\{k = \text{GCD}[m+1, n]\}, \text{Simp}[1/k \ \text{Subst}[\text{Int}[x^{(m+1)/k-1}*(a+b*x^{(n/k)})^p, x], x, x^k], x] /; k \neq 1] /; \text{FreeQ}\{[a, b, p], x\} \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$

rule 6452 $\text{Int}[\{(a_)+\text{ArcTanh}[(c_)*(x_)^n]\}*(b_)^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a+b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \ \text{Int}[x^{(m+n)}*((a+b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{(2*n)})}), x], x] /; \text{FreeQ}\{[a, b, c, m, n], x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.07

method	result	size
parallelrisch	$-\frac{\arctanh(cx^2)bc^2x^4 - ac^2x^4 - bcx^2 + b \arctanh(cx^2)}{4c^2}$	46
default	$\frac{x^4a}{4} + \frac{bx^4 \arctanh(cx^2)}{4} + \frac{bx^2}{4c} - \frac{b \ln(cx^2+1)}{8c^2} + \frac{b \ln(cx^2-1)}{8c^2}$	57
parts	$\frac{x^4a}{4} + \frac{bx^4 \arctanh(cx^2)}{4} + \frac{bx^2}{4c} - \frac{b \ln(cx^2+1)}{8c^2} + \frac{b \ln(cx^2-1)}{8c^2}$	57
orering	$\frac{5(c^2x^4-1)(a+b \arctanh(cx^2))}{8c^2} - \frac{(cx^2-1)(cx^2+1)(3x^2(a+b \arctanh(cx^2))+\frac{2x^4bc}{-c^2x^4+1})}{8c^2x^2}$	83
risch	$\frac{bx^4 \ln(cx^2+1)}{8} - \frac{bx^4 \ln(-cx^2+1)}{8} + \frac{x^4a}{4} + \frac{bx^2}{4c} - \frac{b \ln(cx^2+1)}{8c^2} + \frac{b \ln(cx^2-1)}{8c^2} + \frac{b^2}{16ac^2}$	85

input `int(x^3*(a+b*arctanh(c*x^2)),x,method=_RETURNVERBOSE)`

output `-1/4*(-arctanh(c*x^2)*b*c^2*x^4-a*c^2*x^4-b*c*x^2+b*arctanh(c*x^2))/c^2`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.26

$$\int x^3(a + b \operatorname{arctanh}(cx^2)) dx = \frac{2ac^2x^4 + 2bcx^2 + (bc^2x^4 - b) \log\left(-\frac{cx^2+1}{cx^2-1}\right)}{8c^2}$$

input `integrate(x^3*(a+b*arctanh(c*x^2)),x, algorithm="fricas")`

output `1/8*(2*a*c^2*x^4 + 2*b*c*x^2 + (b*c^2*x^4 - b)*log(-(c*x^2 + 1)/(c*x^2 - 1)))/c^2`

Sympy [A] (verification not implemented)

Time = 3.07 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.12

$$\int x^3(a + b \operatorname{arctanh}(cx^2)) dx = \begin{cases} \frac{ax^4}{4} + \frac{bx^4 \operatorname{atanh}(cx^2)}{4} + \frac{bx^2}{4c} - \frac{b \operatorname{atanh}(cx^2)}{4c^2} & \text{for } c \neq 0 \\ \frac{ax^4}{4} & \text{otherwise} \end{cases}$$

input `integrate(x**3*(a+b*atanh(c*x**2)),x)`

output `Piecewise((a*x**4/4 + b*x**4*atanh(c*x**2)/4 + b*x**2/(4*c) - b*atanh(c*x**2)/(4*c**2), Ne(c, 0)), (a*x**4/4, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.35

$$\int x^3(a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{1}{4} ax^4 + \frac{1}{8} \left(2x^4 \operatorname{artanh}(cx^2) + c \left(\frac{2x^2}{c^2} - \frac{\log(cx^2 + 1)}{c^3} + \frac{\log(cx^2 - 1)}{c^3} \right) \right) b$$

input `integrate(x^3*(a+b*arctanh(c*x^2)),x, algorithm="maxima")`

output `1/4*a*x^4 + 1/8*(2*x^4*arctanh(c*x^2) + c*(2*x^2/c^2 - log(c*x^2 + 1)/c^3 + log(c*x^2 - 1)/c^3))*b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 181 vs. 2(37) = 74.

Time = 0.13 (sec) , antiderivative size = 181, normalized size of antiderivative = 4.21

$$\int x^3(a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{1}{2} c \left(\frac{(cx^2 + 1)b \log\left(-\frac{cx^2+1}{cx^2-1}\right)}{\left(\frac{(cx^2+1)^2 c^3}{(cx^2-1)^2} - \frac{2(cx^2+1)c^3}{cx^2-1} + c^3\right)(cx^2 - 1)} + \frac{\frac{2(cx^2+1)a}{cx^2-1} + \frac{(cx^2+1)b}{cx^2-1} - b}{\left(\frac{(cx^2+1)^2 c^3}{(cx^2-1)^2} - \frac{2(cx^2+1)c^3}{cx^2-1} + c^3\right)} \right)$$

input `integrate(x^3*(a+b*arctanh(c*x^2)),x, algorithm="giac")`

output `1/2*c*((c*x^2 + 1)*b*log(-(c*x^2 + 1)/(c*x^2 - 1))/(((c*x^2 + 1)^2*c^3/(c*x^2 - 1)^2 - 2*(c*x^2 + 1)*c^3/(c*x^2 - 1) + c^3)*(c*x^2 - 1)) + (2*(c*x^2 + 1)*a/(c*x^2 - 1) + (c*x^2 + 1)*b/(c*x^2 - 1) - b)/((c*x^2 + 1)^2*c^3/(c*x^2 - 1)^2 - 2*(c*x^2 + 1)*c^3/(c*x^2 - 1) + c^3))`

Mupad [B] (verification not implemented)

Time = 3.75 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.40

$$\int x^3(a + b \operatorname{arctanh}(cx^2)) dx = \frac{ax^4}{4} + \frac{bx^2}{4c} + \frac{bx^4 \ln(cx^2 + 1)}{8} - \frac{bx^4 \ln(1 - cx^2)}{8} + \frac{b \operatorname{atan}(cx^2) \operatorname{li}}{4c^2}$$

input `int(x^3*(a + b*atanh(c*x^2)),x)`output `(a*x^4)/4 + (b*x^2)/(4*c) + (b*atan(c*x^2*1i)*1i)/(4*c^2) + (b*x^4*log(c*x^2 + 1))/8 - (b*x^4*log(1 - c*x^2))/8`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int x^3(a + b \operatorname{arctanh}(cx^2)) dx = \frac{\operatorname{atanh}(cx^2) b c^2 x^4 - \operatorname{atanh}(cx^2) b + a c^2 x^4 + b c x^2}{4c^2}$$

input `int(x^3*(a+b*atanh(c*x^2)),x)`output `(atanh(c*x**2)*b*c**2*x**4 - atanh(c*x**2)*b + a*c**2*x**4 + b*c*x**2)/(4*c**2)`

3.53 $\int x(a + b \operatorname{arctanh}(cx^2)) dx$

Optimal result	481
Mathematica [A] (verified)	481
Rubi [A] (verified)	482
Maple [A] (verified)	483
Fricas [A] (verification not implemented)	483
Sympy [B] (verification not implemented)	484
Maxima [A] (verification not implemented)	484
Giac [B] (verification not implemented)	485
Mupad [B] (verification not implemented)	485
Reduce [B] (verification not implemented)	486

Optimal result

Integrand size = 12, antiderivative size = 42

$$\int x(a + b \operatorname{arctanh}(cx^2)) dx = \frac{ax^2}{2} + \frac{1}{2}bx^2 \operatorname{arctanh}(cx^2) + \frac{b \log(1 - c^2x^4)}{4c}$$

output

```
1/2*a*x^2+1/2*b*x^2*arctanh(c*x^2)+1/4*b*ln(-c^2*x^4+1)/c
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int x(a + b \operatorname{arctanh}(cx^2)) dx = \frac{ax^2}{2} + \frac{1}{2}bx^2 \operatorname{arctanh}(cx^2) + \frac{b \log(1 - c^2x^4)}{4c}$$

input

```
Integrate[x*(a + b*ArcTanh[c*x^2]),x]
```

output

```
(a*x^2)/2 + (b*x^2*ArcTanh[c*x^2])/2 + (b*Log[1 - c^2*x^4])/(4*c)
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6452, 792}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(a + b \operatorname{arctanh}(cx^2)) dx$$

$$\downarrow 6452$$

$$\frac{1}{2}x^2(a + b \operatorname{arctanh}(cx^2)) - bc \int \frac{x^3}{1 - c^2x^4} dx$$

$$\downarrow 792$$

$$\frac{1}{2}x^2(a + b \operatorname{arctanh}(cx^2)) + \frac{b \log(1 - c^2x^4)}{4c}$$

input `Int[x*(a + b*ArcTanh[c*x^2]),x]`

output `(x^2*(a + b*ArcTanh[c*x^2]))/2 + (b*Log[1 - c^2*x^4])/(4*c)`

Defintions of rubi rules used

rule 792 `Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_.)), x_Symbol] := Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

method	result	size
parts	$\frac{ax^2}{2} + \frac{bx^2 \operatorname{arctanh}(cx^2)}{2} + \frac{b \ln(-c^2x^4+1)}{4c}$	37
derivativedivides	$\frac{cx^2a+b \left(cx^2 \operatorname{arctanh}(cx^2) + \frac{\ln(-c^2x^4+1)}{2} \right)}{2c}$	40
default	$\frac{cx^2a+b \left(cx^2 \operatorname{arctanh}(cx^2) + \frac{\ln(-c^2x^4+1)}{2} \right)}{2c}$	40
parallelrisc	$\frac{b \operatorname{arctanh}(cx^2)x^2c+cx^2a+b \ln(cx^2-1)+b \operatorname{arctanh}(cx^2)}{2c}$	43
risc	$\frac{x^2b \ln(cx^2+1)}{4} - \frac{bx^2 \ln(-cx^2+1)}{4} + \frac{ax^2}{2} + \frac{b \ln(c^2x^4-1)}{4c}$	53

input `int(x*(a+b*arctanh(c*x^2)),x,method=_RETURNVERBOSE)`

output `1/2*a*x^2+1/2*b*x^2*arctanh(c*x^2)+1/4*b*ln(-c^2*x^4+1)/c`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.19

$$\int x(a + b \operatorname{arctanh}(cx^2)) dx = \frac{bcx^2 \log\left(-\frac{cx^2+1}{cx^2-1}\right) + 2acx^2 + b \log(c^2x^4 - 1)}{4c}$$

input `integrate(x*(a+b*arctanh(c*x^2)),x, algorithm="fricas")`

output `1/4*(b*c*x^2*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 2*a*c*x^2 + b*log(c^2*x^4 - 1))/c`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 71 vs. $2(34) = 68$.

Time = 2.64 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.69

$$\int x(a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \begin{cases} \frac{ax^2}{2} + \frac{bx^2 \operatorname{atanh}(cx^2)}{2} + \frac{b \log(x - \sqrt{-1/c})}{2c} + \frac{b \log(x + \sqrt{-1/c})}{2c} - \frac{b \operatorname{atanh}(cx^2)}{2c} & \text{for } c \neq 0 \\ \frac{ax^2}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*(a+b*atanh(c*x**2)),x)`

output `Piecewise((a*x**2/2 + b*x**2*atanh(c*x**2)/2 + b*log(x - sqrt(-1/c))/(2*c) + b*log(x + sqrt(-1/c))/(2*c) - b*atanh(c*x**2)/(2*c), Ne(c, 0)), (a*x**2/2, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

$$\int x(a + b \operatorname{arctanh}(cx^2)) dx = \frac{1}{2} ax^2 + \frac{(2cx^2 \operatorname{artanh}(cx^2) + \log(-c^2x^4 + 1))b}{4c}$$

input `integrate(x*(a+b*arctanh(c*x^2)),x, algorithm="maxima")`

output `1/2*a*x^2 + 1/4*(2*c*x^2*arctanh(c*x^2) + log(-c^2*x^4 + 1))*b/c`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 188 vs. $2(36) = 72$.

Time = 0.13 (sec) , antiderivative size = 188, normalized size of antiderivative = 4.48

$$\int x(a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{1}{2} ax^2 + \frac{1}{2} bc \left(\frac{\log\left(\frac{|-cx^2-1|}{|cx^2-1|}\right)}{c^2} - \frac{\log\left(\left|-\frac{cx^2+1}{cx^2-1} + 1\right|\right)}{c^2} + \frac{\log\left(\frac{\frac{c\left(\frac{cx^2+1}{cx^2-1}+1\right)}{(cx^2+1)c-c}+1}{-\frac{cx^2-1}{cx^2-1}-c}\right)}{c^2\left(\frac{cx^2+1}{cx^2-1}-1\right)} \right)$$

input `integrate(x*(a+b*arctanh(c*x^2)),x, algorithm="giac")`

output `1/2*a*x^2 + 1/2*b*c*(log(abs(-c*x^2 - 1)/abs(c*x^2 - 1))/c^2 - log(abs(-(c*x^2 + 1)/(c*x^2 - 1) + 1))/c^2 + log(-(c*((c*x^2 + 1)/(c*x^2 - 1) + 1))/((c*x^2 + 1)*c/(c*x^2 - 1) - c) + 1)/(c*((c*x^2 + 1)/(c*x^2 - 1) + 1)/((c*x^2 + 1)*c/(c*x^2 - 1) - c) - 1))/(c^2*((c*x^2 + 1)/(c*x^2 - 1) - 1))`

Mupad [B] (verification not implemented)

Time = 3.54 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.24

$$\int x(a + b \operatorname{arctanh}(cx^2)) dx = \frac{ax^2}{2} + \frac{b \ln(c^2 x^4 - 1)}{4c} + \frac{bx^2 \ln(cx^2 + 1)}{4} - \frac{bx^2 \ln(1 - cx^2)}{4}$$

input `int(x*(a + b*atanh(c*x^2)),x)`

output `(a*x^2)/2 + (b*log(c^2*x^4 - 1))/(4*c) + (b*x^2*log(c*x^2 + 1))/4 - (b*x^2*log(1 - c*x^2))/4`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.02

$$\int x(a + b \operatorname{arctanh}(cx^2)) dx = \frac{\operatorname{atanh}(cx^2)bcx^2 - \operatorname{atanh}(cx^2)b + \log(cx^2 + 1)b + acx^2}{2c}$$

input `int(x*(a+b*atanh(c*x^2)),x)`

output `(atanh(c*x**2)*b*c*x**2 - atanh(c*x**2)*b + log(c*x**2 + 1)*b + a*c*x**2)/
(2*c)`

3.54 $\int \frac{a+b\operatorname{arctanh}(cx^2)}{x} dx$

Optimal result	487
Mathematica [A] (verified)	487
Rubi [A] (verified)	488
Maple [B] (verified)	489
Fricas [F]	489
Sympy [F]	490
Maxima [F]	490
Giac [F]	490
Mupad [F(-1)]	491
Reduce [F]	491

Optimal result

Integrand size = 14, antiderivative size = 30

$$\int \frac{a + b\operatorname{arctanh}(cx^2)}{x} dx = a \log(x) - \frac{1}{4}b \operatorname{PolyLog}(2, -cx^2) + \frac{1}{4}b \operatorname{PolyLog}(2, cx^2)$$

output `a*ln(x)-1/4*b*polylog(2,-c*x^2)+1/4*b*polylog(2,c*x^2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{a + b\operatorname{arctanh}(cx^2)}{x} dx = a \log(x) + \frac{1}{4}b(-\operatorname{PolyLog}(2, -cx^2) + \operatorname{PolyLog}(2, cx^2))$$

input `Integrate[(a + b*ArcTanh[c*x^2])/x,x]`

output `a*Log[x] + (b*(-PolyLog[2, -(c*x^2)] + PolyLog[2, c*x^2]))/4`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.20, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6450, 6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x} dx$$

↓ 6450

$$\frac{1}{2} \int \frac{a + b \operatorname{arctanh}(cx^2)}{x^2} dx^2$$

↓ 6446

$$\frac{1}{2} \left(a \log(x^2) - \frac{1}{2} b \operatorname{PolyLog}(2, -cx^2) + \frac{1}{2} b \operatorname{PolyLog}(2, cx^2) \right)$$

input `Int[(a + b*ArcTanh[c*x^2])/x,x]`

output `(a*Log[x^2] - (b*PolyLog[2, -(c*x^2)])/2 + (b*PolyLog[2, c*x^2])/2)/2`

Defintions of rubi rules used

rule 6446 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))/(x_), x_Symbol] := Simp[a*Log[x], x] + (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) / ; FreeQ[{a, b, c}, x]`

rule 6450 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_.)/(x_), x_Symbol] := Simp[1/n Subst[Int[(a + b*ArcTanh[c*x])^p/x, x], x, x^n], x] / ; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 125 vs. $2(26) = 52$.

Time = 0.16 (sec) , antiderivative size = 126, normalized size of antiderivative = 4.20

method	result
default	$a \ln(x) + b \left(\ln(x) \operatorname{arctanh}(cx^2) - 2c \left(\frac{\ln(x)(\ln(1+\sqrt{-c}x) + \ln(1-\sqrt{-c}x))}{4c} + \frac{\operatorname{dilog}(1+\sqrt{-c}x) + \operatorname{dilog}(1-\sqrt{-c}x)}{4c} \right) \right)$
parts	$a \ln(x) + b \left(\ln(x) \operatorname{arctanh}(cx^2) - 2c \left(\frac{\ln(x)(\ln(1+\sqrt{-c}x) + \ln(1-\sqrt{-c}x))}{4c} + \frac{\operatorname{dilog}(1+\sqrt{-c}x) + \operatorname{dilog}(1-\sqrt{-c}x)}{4c} \right) \right)$
risch	$a \ln(x) - \frac{\ln(x) \ln(-cx^2+1)b}{2} + \frac{b \ln(x) \ln(1+\sqrt{c}x)}{2} + \frac{b \ln(x) \ln(1-\sqrt{c}x)}{2} + \frac{b \operatorname{dilog}(1-\sqrt{c}x)}{2} + \frac{b \operatorname{dilog}(1+\sqrt{c}x)}{2} - \frac{\ln(x)}{2}$

input `int((a+b*arctanh(c*x^2))/x,x,method=_RETURNVERBOSE)`

output `a*ln(x)+b*(ln(x)*arctanh(c*x^2)-2*c*(1/4*ln(x)*(ln(1+(-c)^(1/2)*x)+ln(1-(-c)^(1/2)*x))/c+1/4*(dilog(1+(-c)^(1/2)*x)+dilog(1-(-c)^(1/2)*x))/c-1/4*ln(x)*(ln(1-c^(1/2)*x)+ln(1+c^(1/2)*x))/c-1/4*(dilog(1-c^(1/2)*x)+dilog(1+c^(1/2)*x))/c)`

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x} dx = \int \frac{b \operatorname{artanh}(cx^2) + a}{x} dx$$

input `integrate((a+b*arctanh(c*x^2))/x,x, algorithm="fricas")`

output `integral((b*arctanh(c*x^2) + a)/x, x)`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x} dx = \int \frac{a + b \operatorname{atanh}(cx^2)}{x} dx$$

input `integrate((a+b*atanh(c*x**2))/x,x)`

output `Integral((a + b*atanh(c*x**2))/x, x)`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x} dx = \int \frac{b \operatorname{artanh}(cx^2) + a}{x} dx$$

input `integrate((a+b*arctanh(c*x^2))/x,x, algorithm="maxima")`

output `1/2*b*integrate((log(c*x^2 + 1) - log(-c*x^2 + 1))/x, x) + a*log(x)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x} dx = \int \frac{b \operatorname{artanh}(cx^2) + a}{x} dx$$

input `integrate((a+b*arctanh(c*x^2))/x,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x} dx = \int \frac{a + b \operatorname{atanh}(cx^2)}{x} dx$$

input `int((a + b*atanh(c*x^2))/x,x)`output `int((a + b*atanh(c*x^2))/x, x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x} dx = \left(\int \frac{\operatorname{atanh}(cx^2)}{x} dx \right) b + \log(x) a$$

input `int((a+b*atanh(c*x^2))/x,x)`output `int(atanh(c*x**2)/x,x)*b + log(x)*a`

3.55 $\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^3} dx$

Optimal result	492
Mathematica [A] (verified)	492
Rubi [A] (verified)	493
Maple [A] (verified)	494
Fricas [A] (verification not implemented)	495
Sympy [B] (verification not implemented)	495
Maxima [A] (verification not implemented)	496
Giac [A] (verification not implemented)	496
Mupad [B] (verification not implemented)	496
Reduce [B] (verification not implemented)	497

Optimal result

Integrand size = 14, antiderivative size = 40

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^3} dx = -\frac{a + b \operatorname{arctanh}(cx^2)}{2x^2} + bc \log(x) - \frac{1}{4}bc \log(1 - c^2x^4)$$

output `-1/2*(a+b*arctanh(c*x^2))/x^2+b*c*ln(x)-1/4*b*c*ln(-c^2*x^4+1)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.12

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^3} dx = -\frac{a}{2x^2} - \frac{b \operatorname{arctanh}(cx^2)}{2x^2} + bc \log(x) - \frac{1}{4}bc \log(1 - c^2x^4)$$

input `Integrate[(a + b*ArcTanh[c*x^2])/x^3,x]`

output `-1/2*a/x^2 - (b*ArcTanh[c*x^2])/(2*x^2) + b*c*Log[x] - (b*c*Log[1 - c^2*x^4])/4`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {6452, 798, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{arctanh}(cx^2)}{x^3} dx \\
 & \quad \downarrow 6452 \\
 & bc \int \frac{1}{x(1-c^2x^4)} dx - \frac{a + \operatorname{arctanh}(cx^2)}{2x^2} \\
 & \quad \downarrow 798 \\
 & \frac{1}{4}bc \int \frac{1}{x^4(1-c^2x^4)} dx^4 - \frac{a + \operatorname{arctanh}(cx^2)}{2x^2} \\
 & \quad \downarrow 47 \\
 & \frac{1}{4}bc \left(c^2 \int \frac{1}{1-c^2x^4} dx^4 + \int \frac{1}{x^4} dx^4 \right) - \frac{a + \operatorname{arctanh}(cx^2)}{2x^2} \\
 & \quad \downarrow 14 \\
 & \frac{1}{4}bc \left(c^2 \int \frac{1}{1-c^2x^4} dx^4 + \log(x^4) \right) - \frac{a + \operatorname{arctanh}(cx^2)}{2x^2} \\
 & \quad \downarrow 16 \\
 & \frac{1}{4}bc (\log(x^4) - \log(1-c^2x^4)) - \frac{a + \operatorname{arctanh}(cx^2)}{2x^2}
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c*x^2])/x^3,x]`

output `-1/2*(a + b*ArcTanh[c*x^2])/x^2 + (b*c*(Log[x^4] - Log[1 - c^2*x^4]))/4`

Definitions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.18

method	result	size
default	$-\frac{a}{2x^2} + b\left(-\frac{\operatorname{arctanh}(cx^2)}{2x^2} + c\left(\ln(x) - \frac{\ln(cx^2+1)}{4} - \frac{\ln(cx^2-1)}{4}\right)\right)$	47
parts	$-\frac{a}{2x^2} + b\left(-\frac{\operatorname{arctanh}(cx^2)}{2x^2} + c\left(\ln(x) - \frac{\ln(cx^2+1)}{4} - \frac{\ln(cx^2-1)}{4}\right)\right)$	47
parallelrisch	$\frac{2bc\ln(x)x^2 - \ln(cx^2-1)x^2bc - b\operatorname{arctanh}(cx^2)x^2c - b\operatorname{arctanh}(cx^2) - a}{2x^2}$	56
risch	$-\frac{b\ln(cx^2+1)}{4x^2} + \frac{4bc\ln(x)x^2 - bc\ln(c^2x^4-1)x^2 + b\ln(-cx^2+1) - 2a}{4x^2}$	62

input `int((a+b*arctanh(c*x^2))/x^3,x,method=_RETURNVERBOSE)`

output `-1/2*a/x^2+b*(-1/2/x^2*arctanh(c*x^2)+c*(ln(x)-1/4*ln(c*x^2+1)-1/4*ln(c*x^2-1)))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.38

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^3} dx = -\frac{bcx^2 \log(c^2x^4 - 1) - 4bcx^2 \log(x) + b \log\left(-\frac{cx^2+1}{cx^2-1}\right) + 2a}{4x^2}$$

input `integrate((a+b*arctanh(c*x^2))/x^3,x, algorithm="fricas")`

output `-1/4*(b*c*x^2*log(c^2*x^4 - 1) - 4*b*c*x^2*log(x) + b*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 2*a)/x^2`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 80 vs. $2(39) = 78$.

Time = 4.30 (sec) , antiderivative size = 80, normalized size of antiderivative = 2.00

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^3} dx = \begin{cases} -\frac{a}{2x^2} + bc \log(x) - \frac{bc \log\left(x - \sqrt{-\frac{1}{c}}\right)}{2} - \frac{bc \log\left(x + \sqrt{-\frac{1}{c}}\right)}{2} + \frac{bc \operatorname{atanh}(cx^2)}{2} - \frac{b \operatorname{atanh}(cx^2)}{2x^2} & \text{for } c \neq 0 \\ -\frac{a}{2x^2} & \text{otherwise} \end{cases}$$

input `integrate((a+b*atanh(c*x**2))/x**3,x)`

output `Piecewise((-a/(2*x**2) + b*c*log(x) - b*c*log(x - sqrt(-1/c))/2 - b*c*log(x + sqrt(-1/c))/2 + b*c*atanh(c*x**2)/2 - b*atanh(c*x**2)/(2*x**2), Ne(c, 0)), (-a/(2*x**2), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.02

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^3} dx = -\frac{1}{4} \left(c(\log(c^2x^4 - 1) - \log(x^4)) + \frac{2 \operatorname{artanh}(cx^2)}{x^2} \right) b - \frac{a}{2x^2}$$

input `integrate((a+b*arctanh(c*x^2))/x^3,x, algorithm="maxima")`output `-1/4*(c*(log(c^2*x^4 - 1) - log(x^4)) + 2*arctanh(c*x^2)/x^2)*b - 1/2*a/x^2`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.28

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^3} dx = -\frac{1}{4} bc \log(c^2x^4 - 1) + bc \log(x) - \frac{b \log\left(-\frac{cx^2+1}{cx^2-1}\right)}{4x^2} - \frac{a}{2x^2}$$

input `integrate((a+b*arctanh(c*x^2))/x^3,x, algorithm="giac")`output `-1/4*b*c*log(c^2*x^4 - 1) + b*c*log(x) - 1/4*b*log(-(c*x^2 + 1)/(c*x^2 - 1))/x^2 - 1/2*a/x^2`**Mupad [B] (verification not implemented)**

Time = 3.68 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.38

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^3} dx = bc \ln(x) - \frac{a}{2x^2} - \frac{bc \ln(c^2x^4 - 1)}{4} - \frac{b \ln(cx^2 + 1)}{4x^2} + \frac{b \ln(1 - cx^2)}{4x^2}$$

input `int((a + b*atanh(c*x^2))/x^3,x)`

output $b*c*\log(x) - a/(2*x^2) - (b*c*\log(c^2*x^4 - 1))/4 - (b*\log(c*x^2 + 1))/(4*x^2) + (b*\log(1 - c*x^2))/(4*x^2)$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.35

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^3} dx$$

$$= \frac{\operatorname{atanh}(cx^2) bcx^2 - \operatorname{atanh}(cx^2) b - \log(cx^2 + 1) bcx^2 + 2 \log(x) bcx^2 - a}{2x^2}$$

input `int((a+b*atanh(c*x^2))/x^3,x)`

output $(\operatorname{atanh}(c*x**2)*b*c*x**2 - \operatorname{atanh}(c*x**2)*b - \log(c*x**2 + 1)*b*c*x**2 + 2*\log(x)*b*c*x**2 - a)/(2*x**2)$

3.56 $\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^5} dx$

Optimal result	498
Mathematica [A] (verified)	498
Rubi [A] (verified)	499
Maple [A] (verified)	500
Fricas [A] (verification not implemented)	501
Sympy [A] (verification not implemented)	501
Maxima [A] (verification not implemented)	502
Giac [A] (verification not implemented)	502
Mupad [B] (verification not implemented)	503
Reduce [B] (verification not implemented)	503

Optimal result

Integrand size = 14, antiderivative size = 41

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^5} dx = -\frac{bc}{4x^2} + \frac{1}{4}bc^2 \operatorname{arctanh}(cx^2) - \frac{a + b \operatorname{arctanh}(cx^2)}{4x^4}$$

output

```
-1/4*b*c/x^2+1/4*b*c^2*arctanh(c*x^2)-1/4*(a+b*arctanh(c*x^2))/x^4
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.59

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^5} dx = -\frac{a}{4x^4} - \frac{bc}{4x^2} - \frac{b \operatorname{arctanh}(cx^2)}{4x^4} - \frac{1}{8}bc^2 \log(1 - cx^2) + \frac{1}{8}bc^2 \log(1 + cx^2)$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])/x^5, x]
```

output

```
-1/4*a/x^4 - (b*c)/(4*x^2) - (b*ArcTanh[c*x^2])/(4*x^4) - (b*c^2*Log[1 - c*x^2])/8 + (b*c^2*Log[1 + c*x^2])/8
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.93, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6452, 807, 264, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^5} dx$$

$$\downarrow 6452$$

$$\frac{1}{2}bc \int \frac{1}{x^3(1-c^2x^4)} dx - \frac{a + b \operatorname{arctanh}(cx^2)}{4x^4}$$

$$\downarrow 807$$

$$\frac{1}{4}bc \int \frac{1}{x^4(1-c^2x^4)} dx^2 - \frac{a + b \operatorname{arctanh}(cx^2)}{4x^4}$$

$$\downarrow 264$$

$$\frac{1}{4}bc \left(c^2 \int \frac{1}{1-c^2x^4} dx^2 - \frac{1}{x^2} \right) - \frac{a + b \operatorname{arctanh}(cx^2)}{4x^4}$$

$$\downarrow 219$$

$$\frac{1}{4}bc \left(\operatorname{arctanh}(cx^2) - \frac{1}{x^2} \right) - \frac{a + b \operatorname{arctanh}(cx^2)}{4x^4}$$

input `Int[(a + b*ArcTanh[c*x^2])/x^5,x]`

output `-1/4*(a + b*ArcTanh[c*x^2])/x^4 + (b*c*(-x^(-2) + c*ArcTanh[c*x^2]))/4`

Defintions of rubi rules used

rule 219

$$\text{Int}[\{(a_)+(b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[\{1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2])\} * \text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 264

$$\text{Int}[\{(c_)*(x_)\}^{(m_)}*\{(a_)+(b_)*(x_)^2\}^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(c*x)^{(m+1)}*\{(a+b*x^2)^{(p+1)}/(a*c*(m+1))\}, x] - \text{Simp}[b*((m+2*p+3)/(a*c^{2*(m+1)})) \ \text{Int}[(c*x)^{(m+2)}*(a+b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 807

$$\text{Int}[(x_)^{(m_)}*\{(a_)+(b_)*(x_)^{(n_)}\}^{(p_)}, x_Symbol] \rightarrow \text{With}[\{k = \text{GCD}[m+1, n]\}, \text{Simp}[1/k \ \text{Subst}[\text{Int}[x^{(m+1)/k-1}*(a+b*x^{(n/k)})^p, x], x, x^k], x] /; k \neq 1] /; \text{FreeQ}\{a, b, p\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$$

rule 6452

$$\text{Int}[\{(a_)+\text{ArcTanh}[(c_)*(x_)^{(n_)}]\}*(b_)\}^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*\{(a+b*\text{ArcTanh}[c*x^n])^{p/(m+1)}\}, x] - \text{Simp}[b*c*n*(p/(m+1)) \ \text{Int}[x^{(m+n)}*\{(a+b*\text{ArcTanh}[c*x^n])^{p-1}/(1-c^2*x^{2*n})\}, x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$$

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.10

method	result	size
parallelrisch	$-\frac{-\arctanh(cx^2)bc^2x^4+ac^2x^4+bcx^2+b\arctanh(cx^2)+a}{4x^4}$	45
default	$-\frac{a}{4x^4} - \frac{b\arctanh(cx^2)}{4x^4} + \frac{bc^2\ln(cx^2+1)}{8} - \frac{bc^2\ln(cx^2-1)}{8} - \frac{bc}{4x^2}$	55
parts	$-\frac{a}{4x^4} - \frac{b\arctanh(cx^2)}{4x^4} + \frac{bc^2\ln(cx^2+1)}{8} - \frac{bc^2\ln(cx^2-1)}{8} - \frac{bc}{4x^2}$	55
risch	$-\frac{b\ln(cx^2+1)}{8x^4} - \frac{bc^2\ln(cx^2-1)x^4-bc^2\ln(cx^2+1)x^4+2bcx^2-b\ln(-cx^2+1)+2a}{8x^4}$	77
orering	$\frac{(\frac{7}{8}c^2x^5-\frac{7}{8}x)(a+b\arctanh(cx^2))}{x^5} + \frac{x^2(cx^2+1)(cx^2-1)\left(\frac{2bc}{x^4(-c^2x^4+1)}-\frac{5(a+b\arctanh(cx^2))}{x^6}\right)}{8}$	82

input `int((a+b*arctanh(c*x^2))/x^5,x,method=_RETURNVERBOSE)`

output `-1/4*(-arctanh(c*x^2)*b*c^2*x^4+a*c^2*x^4+b*c*x^2+b*arctanh(c*x^2)+a)/x^4`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.20

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^5} dx = -\frac{2bcx^2 - (bc^2x^4 - b) \log\left(-\frac{cx^2+1}{cx^2-1}\right) + 2a}{8x^4}$$

input `integrate((a+b*arctanh(c*x^2))/x^5,x, algorithm="fricas")`

output `-1/8*(2*b*c*x^2 - (b*c^2*x^4 - b)*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 2*a)/x^4`

Sympy [A] (verification not implemented)

Time = 2.93 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^5} dx = -\frac{a}{4x^4} + \frac{bc^2 \operatorname{atanh}(cx^2)}{4} - \frac{bc}{4x^2} - \frac{b \operatorname{atanh}(cx^2)}{4x^4}$$

input `integrate((a+b*atanh(c*x**2))/x**5,x)`

output `-a/(4*x**4) + b*c**2*atanh(c*x**2)/4 - b*c/(4*x**2) - b*atanh(c*x**2)/(4*x**4)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.24

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^5} dx$$

$$= \frac{1}{8} \left(\left(c \log(cx^2 + 1) - c \log(cx^2 - 1) - \frac{2}{x^2} \right) c - \frac{2 \operatorname{arctanh}(cx^2)}{x^4} \right) b - \frac{a}{4x^4}$$

input `integrate((a+b*arctanh(c*x^2))/x^5,x, algorithm="maxima")`output `1/8*((c*log(c*x^2 + 1) - c*log(c*x^2 - 1) - 2/x^2)*c - 2*arctanh(c*x^2)/x^4)*b - 1/4*a/x^4`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.63

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^5} dx = \frac{1}{8} bc^2 \log(cx^2 + 1) - \frac{1}{8} bc^2 \log(cx^2 - 1)$$

$$- \frac{b \log\left(-\frac{cx^2+1}{cx^2-1}\right)}{8x^4} - \frac{bcx^2 + a}{4x^4}$$

input `integrate((a+b*arctanh(c*x^2))/x^5,x, algorithm="giac")`output `1/8*b*c^2*log(c*x^2 + 1) - 1/8*b*c^2*log(c*x^2 - 1) - 1/8*b*log(-(c*x^2 + 1)/(c*x^2 - 1))/x^4 - 1/4*(b*c*x^2 + a)/x^4`

Mupad [B] (verification not implemented)

Time = 3.89 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.27

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^5} dx = \frac{bc^2 \operatorname{atanh}(cx^2)}{4} - \frac{a}{4} + \frac{b \ln(cx^2+1)}{8} - \frac{b \ln(1-cx^2)}{8} + \frac{bcx^2}{4}$$

input `int((a + b*atanh(c*x^2))/x^5,x)`output `(b*c^2*atanh(c*x^2))/4 - (a/4 + (b*log(c*x^2 + 1))/8 - (b*log(1 - c*x^2)))/8 + (b*c*x^2)/4/x^4`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.95

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^5} dx = \frac{\operatorname{atanh}(cx^2)bc^2x^4 - \operatorname{atanh}(cx^2)b - a - bcx^2}{4x^4}$$

input `int((a+b*atanh(c*x^2))/x^5,x)`output `(atanh(c*x**2)*b*c**2*x**4 - atanh(c*x**2)*b - a - b*c*x**2)/(4*x**4)`

3.57 $\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^7} dx$

Optimal result	504
Mathematica [A] (verified)	504
Rubi [A] (verified)	505
Maple [A] (verified)	506
Fricas [A] (verification not implemented)	507
Sympy [A] (verification not implemented)	507
Maxima [A] (verification not implemented)	508
Giac [A] (verification not implemented)	508
Mupad [B] (verification not implemented)	509
Reduce [B] (verification not implemented)	509

Optimal result

Integrand size = 14, antiderivative size = 56

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^7} dx = -\frac{bc}{12x^4} - \frac{a + b \operatorname{arctanh}(cx^2)}{6x^6} + \frac{1}{3}bc^3 \log(x) - \frac{1}{12}bc^3 \log(1 - c^2x^4)$$

output

```
-1/12*b*c/x^4-1/6*(a+b*arctanh(c*x^2))/x^6+1/3*b*c^3*ln(x)-1/12*b*c^3*ln(-c^2*x^4+1)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.09

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^7} dx = -\frac{a}{6x^6} - \frac{bc}{12x^4} - \frac{b \operatorname{arctanh}(cx^2)}{6x^6} + \frac{1}{3}bc^3 \log(x) - \frac{1}{12}bc^3 \log(1 - c^2x^4)$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])/x^7, x]
```

output

$$-1/6*a/x^6 - (b*c)/(12*x^4) - (b*ArcTanh[c*x^2])/(6*x^6) + (b*c^3*Log[x])/3 - (b*c^3*Log[1 - c^2*x^4])/12$$

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6452, 798, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}(cx^2)}{x^7} dx \\ & \quad \downarrow 6452 \\ & \frac{1}{3}bc \int \frac{1}{x^5(1-c^2x^4)} dx - \frac{a + b \operatorname{arctanh}(cx^2)}{6x^6} \\ & \quad \downarrow 798 \\ & \frac{1}{12}bc \int \frac{1}{x^8(1-c^2x^4)} dx^4 - \frac{a + b \operatorname{arctanh}(cx^2)}{6x^6} \\ & \quad \downarrow 54 \\ & \frac{1}{12}bc \int \left(-\frac{c^4}{c^2x^4-1} + \frac{c^2}{x^4} + \frac{1}{x^8} \right) dx^4 - \frac{a + b \operatorname{arctanh}(cx^2)}{6x^6} \\ & \quad \downarrow 2009 \\ & \frac{1}{12}bc \left(c^2 \log(x^4) - c^2 \log(1-c^2x^4) - \frac{1}{x^4} \right) - \frac{a + b \operatorname{arctanh}(cx^2)}{6x^6} \end{aligned}$$

input

$$\text{Int}[(a + b*ArcTanh[c*x^2])/x^7, x]$$

output

$$-1/6*(a + b*ArcTanh[c*x^2])/x^6 + (b*c*(-x^(-4) + c^2*Log[x^4] - c^2*Log[1 - c^2*x^4]))/12$$

Definitions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.12

method	result	size
default	$-\frac{a}{6x^6} + b \left(-\frac{\operatorname{arctanh}(cx^2)}{6x^6} + \frac{c \left(-\frac{1}{4x^4} + c^2 \ln(x) - \frac{c^2 \ln(cx^2+1)}{4} - \frac{c^2 \ln(cx^2-1)}{4} \right)}{3} \right)$	63
parts	$-\frac{a}{6x^6} + b \left(-\frac{\operatorname{arctanh}(cx^2)}{6x^6} + \frac{c \left(-\frac{1}{4x^4} + c^2 \ln(x) - \frac{c^2 \ln(cx^2+1)}{4} - \frac{c^2 \ln(cx^2-1)}{4} \right)}{3} \right)$	63
risch	$-\frac{b \ln(cx^2+1)}{12x^6} + \frac{4bc^3 \ln(x)x^6 - bc^3 \ln(c^2x^4-1)x^6 - bcx^2 + b \ln(-cx^2+1) - 2a}{12x^6}$	73
parallelrisch	$\frac{4bc^3 \ln(x)x^6 - 2 \ln(cx^2-1)x^6 bc^3 - 2b \operatorname{arctanh}(cx^2)x^6 c^3 - bc^3x^6 - bcx^2 - 2b \operatorname{arctanh}(cx^2) - 2a}{12x^6}$	78

input `int((a+b*arctanh(c*x^2))/x^7,x,method=_RETURNVERBOSE)`

output

```
-1/6*a/x^6+b*(-1/6/x^6*arctanh(c*x^2)+1/3*c*(-1/4/x^4+c^2*ln(x)-1/4*c^2*ln
(c*x^2+1)-1/4*c^2*ln(c*x^2-1)))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.16

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^7} dx$$

$$= -\frac{bc^3 x^6 \log(c^2 x^4 - 1) - 4bc^3 x^6 \log(x) + bcx^2 + b \log\left(-\frac{cx^2+1}{cx^2-1}\right) + 2a}{12x^6}$$

input

```
integrate((a+b*arctanh(c*x^2))/x^7,x, algorithm="fricas")
```

output

```
-1/12*(b*c^3*x^6*log(c^2*x^4 - 1) - 4*b*c^3*x^6*log(x) + b*c*x^2 + b*log(-
(c*x^2 + 1)/(c*x^2 - 1)) + 2*a)/x^6
```

Sympy [A] (verification not implemented)

Time = 7.71 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.73

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^7} dx$$

$$= \begin{cases} -\frac{a}{6x^6} + \frac{bc^3 \log(x)}{3} - \frac{bc^3 \log\left(x - \sqrt{-\frac{1}{c}}\right)}{6} - \frac{bc^3 \log\left(x + \sqrt{-\frac{1}{c}}\right)}{6} + \frac{bc^3 \operatorname{atanh}(cx^2)}{6} - \frac{bc}{12x^4} - \frac{b \operatorname{atanh}(cx^2)}{6x^6} & \text{for } c \neq 0 \\ -\frac{a}{6x^6} & \text{otherwise} \end{cases}$$

input

```
integrate((a+b*atanh(c*x**2))/x**7,x)
```

output

```
Piecewise((-a/(6*x**6) + b*c**3*log(x)/3 - b*c**3*log(x - sqrt(-1/c))/6 -
b*c**3*log(x + sqrt(-1/c))/6 + b*c**3*atanh(c*x**2)/6 - b*c/(12*x**4) - b*
atanh(c*x**2)/(6*x**6), Ne(c, 0)), (-a/(6*x**6), True))
```


Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.91

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^7} dx$$

$$= -\frac{1}{12} \left(\left(c^2 \log(c^2 x^4 - 1) - c^2 \log(x^4) + \frac{1}{x^4} \right) c + \frac{2 \operatorname{artanh}(cx^2)}{x^6} \right) b - \frac{a}{6 x^6}$$

input `integrate((a+b*arctanh(c*x^2))/x^7,x, algorithm="maxima")`output `-1/12*((c^2*log(c^2*x^4 - 1) - c^2*log(x^4) + 1/x^4)*c + 2*arctanh(c*x^2)/x^6)*b - 1/6*a/x^6`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.16

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^7} dx = -\frac{1}{12} bc^3 \log(c^2 x^4 - 1) + \frac{1}{3} bc^3 \log(x)$$

$$- \frac{b \log\left(-\frac{cx^2+1}{cx^2-1}\right)}{12 x^6} - \frac{bcx^2 + 2a}{12 x^6}$$

input `integrate((a+b*arctanh(c*x^2))/x^7,x, algorithm="giac")`output `-1/12*b*c^3*log(c^2*x^4 - 1) + 1/3*b*c^3*log(x) - 1/12*b*log(-(c*x^2 + 1)/(c*x^2 - 1))/x^6 - 1/12*(b*c*x^2 + 2*a)/x^6`

Mupad [B] (verification not implemented)

Time = 3.77 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.20

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^7} dx = \frac{bc^3 \ln(x)}{3} - \frac{bc^3 \ln(c^2 x^4 - 1)}{12} - \frac{a}{6x^6} - \frac{bc}{12x^4} - \frac{b \ln(cx^2 + 1)}{12x^6} + \frac{b \ln(1 - cx^2)}{12x^6}$$

input `int((a + b*atanh(c*x^2))/x^7,x)`output `(b*c^3*log(x))/3 - (b*c^3*log(c^2*x^4 - 1))/12 - a/(6*x^6) - (b*c)/(12*x^4) - (b*log(c*x^2 + 1))/(12*x^6) + (b*log(1 - c*x^2))/(12*x^6)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.21

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^7} dx = \frac{2 \operatorname{atanh}(cx^2) b c^3 x^6 - 2 \operatorname{atanh}(cx^2) b - 2 \log(cx^2 + 1) b c^3 x^6 + 4 \log(x) b c^3 x^6 - 2a - bcx^2}{12x^6}$$

input `int((a+b*atanh(c*x^2))/x^7,x)`output `(2*atanh(c*x**2)*b*c**3*x**6 - 2*atanh(c*x**2)*b - 2*log(c*x**2 + 1)*b*c**3*x**6 + 4*log(x)*b*c**3*x**6 - 2*a - b*c*x**2)/(12*x**6)`

3.58 $\int x^4(a + b \operatorname{arctanh}(cx^2)) dx$

Optimal result	510
Mathematica [A] (verified)	510
Rubi [A] (verified)	511
Maple [A] (verified)	513
Fricas [A] (verification not implemented)	513
Sympy [B] (verification not implemented)	514
Maxima [A] (verification not implemented)	514
Giac [A] (verification not implemented)	515
Mupad [B] (verification not implemented)	515
Reduce [B] (verification not implemented)	516

Optimal result

Integrand size = 14, antiderivative size = 65

$$\int x^4(a + b \operatorname{arctanh}(cx^2)) dx = \frac{2bx^3}{15c} + \frac{b \arctan(\sqrt{cx})}{5c^{5/2}} - \frac{b \operatorname{arctanh}(\sqrt{cx})}{5c^{5/2}} + \frac{1}{5}x^5(a + b \operatorname{arctanh}(cx^2))$$

output

```
2/15*b*x^3/c+1/5*b*arctan(c^(1/2)*x)/c^(5/2)-1/5*b*arctanh(c^(1/2)*x)/c^(5/2)+1/5*x^5*(a+b*arctanh(c*x^2))
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.43

$$\int x^4(a + b \operatorname{arctanh}(cx^2)) dx = \frac{2bx^3}{15c} + \frac{ax^5}{5} + \frac{b \arctan(\sqrt{cx})}{5c^{5/2}} + \frac{1}{5}bx^5 \operatorname{arctanh}(cx^2) + \frac{b \log(1 - \sqrt{cx})}{10c^{5/2}} - \frac{b \log(1 + \sqrt{cx})}{10c^{5/2}}$$

input

```
Integrate[x^4*(a + b*ArcTanh[c*x^2]), x]
```

output

$$(2bx^3)/(15c) + (ax^5)/5 + (b\text{ArcTan}[\text{Sqrt}[c]x])/(5c^{(5/2)}) + (bx^5\text{ArcTanh}[cx^2])/5 + (b\text{Log}[1 - \text{Sqrt}[c]x])/(10c^{(5/2)}) - (b\text{Log}[1 + \text{Sqrt}[c]x])/(10c^{(5/2)})$$
Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.14, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {6452, 843, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4(a + \text{barctanh}(cx^2)) dx$$

$$\downarrow 6452$$

$$\frac{1}{5}x^5(a + \text{barctanh}(cx^2)) - \frac{2}{5}bc \int \frac{x^6}{1 - c^2x^4} dx$$

$$\downarrow 843$$

$$\frac{1}{5}x^5(a + \text{barctanh}(cx^2)) - \frac{2}{5}bc \left(\frac{\int \frac{x^2}{1 - c^2x^4} dx}{c^2} - \frac{x^3}{3c^2} \right)$$

$$\downarrow 827$$

$$\frac{1}{5}x^5(a + \text{barctanh}(cx^2)) - \frac{2}{5}bc \left(\frac{\int \frac{1}{1 - cx^2} dx}{2c} - \frac{\int \frac{1}{cx^2 + 1} dx}{2c} - \frac{x^3}{3c^2} \right)$$

$$\downarrow 216$$

$$\frac{1}{5}x^5(a + \text{barctanh}(cx^2)) - \frac{2}{5}bc \left(\frac{\int \frac{1}{1 - cx^2} dx}{2c} - \frac{\arctan(\sqrt{cx})}{2c^{3/2}} - \frac{x^3}{3c^2} \right)$$

$$\downarrow 219$$

$$\frac{1}{5}x^5(a + \text{barctanh}(cx^2)) - \frac{2}{5}bc \left(\frac{\arctanh(\sqrt{cx})}{2c^{3/2}} - \frac{\arctan(\sqrt{cx})}{2c^{3/2}} - \frac{x^3}{3c^2} \right)$$

input `Int[x^4*(a + b*ArcTanh[c*x^2]),x]`

output `(-2*b*c*(-1/3*x^3/c^2 + (-1/2*ArcTan[Sqrt[c]*x]/c^(3/2) + ArcTanh[Sqrt[c]*x]/(2*c^(3/2)))/c^2)/5 + (x^5*(a + b*ArcTanh[c*x^2]))/5`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 827 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x], x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 843 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Simp[a*c^n*((m - n + 1)/(b*(m + n*p + 1))) Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.82

method	result
default	$\frac{ax^5}{5} + \frac{bx^5 \operatorname{arctanh}(cx^2)}{5} + \frac{2bx^3}{15c} + \frac{b \arctan(\sqrt{cx})}{5c^{\frac{5}{2}}} - \frac{b \operatorname{arctanh}(\sqrt{cx})}{5c^{\frac{5}{2}}}$
parts	$\frac{ax^5}{5} + \frac{bx^5 \operatorname{arctanh}(cx^2)}{5} + \frac{2bx^3}{15c} + \frac{b \arctan(\sqrt{cx})}{5c^{\frac{5}{2}}} - \frac{b \operatorname{arctanh}(\sqrt{cx})}{5c^{\frac{5}{2}}}$
risch	$\frac{bx^5 \ln(cx^2+1)}{10} - \frac{bx^5 \ln(-cx^2+1)}{10} + \frac{ax^5}{5} + \frac{2bx^3}{15c} + \frac{b \ln(1-\sqrt{cx})}{10c^{\frac{5}{2}}} - \frac{b \ln(1+\sqrt{cx})}{10c^{\frac{5}{2}}} + \frac{\sqrt{-c} \ln(-\sqrt{-c}c-c^2x)b}{10c^3} - \frac{\sqrt{-c}}{10c^3}$

input `int(x^4*(a+b*arctanh(c*x^2)),x,method=_RETURNVERBOSE)`output $\frac{1}{5}ax^5 + \frac{1}{5}bx^5 \operatorname{arctanh}(cx^2) + \frac{2}{15}bx^3/c + \frac{1}{5}b \arctan(c^{1/2}x)/c^{5/2} - \frac{1}{5}b \operatorname{arctanh}(c^{1/2}x)/c^{5/2}$ **Fricas [A] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 102 vs. 2(49) = 98.

Time = 0.10 (sec) , antiderivative size = 197, normalized size of antiderivative = 3.03

$$\int x^4(a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{3bc^3x^5 \log\left(-\frac{cx^2+1}{cx^2-1}\right) + 6ac^3x^5 + 4bc^2x^3 + 6b\sqrt{c} \arctan(\sqrt{cx}) + 3b\sqrt{c} \log\left(\frac{cx^2-2\sqrt{cx}+1}{cx^2-1}\right)}{30c^3}, \frac{3bc^3x^5 \log\left(\frac{cx^2-2\sqrt{cx}+1}{cx^2-1}\right)}{30c^3}$$

input `integrate(x^4*(a+b*arctanh(c*x^2)),x, algorithm="fricas")`output $[1/30*(3*b*c^3*x^5*\log(-(c*x^2 + 1)/(c*x^2 - 1)) + 6*a*c^3*x^5 + 4*b*c^2*x^3 + 6*b*\sqrt{c}*\arctan(\sqrt{c}*x) + 3*b*\sqrt{c}*\log((c*x^2 - 2*\sqrt{c}*x + 1)/(c*x^2 - 1)))/c^3, 1/30*(3*b*c^3*x^5*\log(-(c*x^2 + 1)/(c*x^2 - 1)) + 6*a*c^3*x^5 + 4*b*c^2*x^3 + 6*b*\sqrt{-c}*\arctan(\sqrt{-c}*x) - 3*b*\sqrt{-c}*\log((c*x^2 - 2*\sqrt{-c}*x - 1)/(c*x^2 + 1)))/c^3]$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 185 vs. $2(58) = 116$.

Time = 4.47 (sec) , antiderivative size = 185, normalized size of antiderivative = 2.85

$$\int x^4(a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \begin{cases} \frac{ax^5}{5} + \frac{bx^5 \operatorname{atanh}(cx^2)}{5} + \frac{2bx^3}{15c} - \frac{b\sqrt{-\frac{1}{c}} \log\left(x - \sqrt{-\frac{1}{c}}\right)}{10c^2} + \frac{b\sqrt{-\frac{1}{c}} \log\left(x + \sqrt{-\frac{1}{c}}\right)}{10c^2} - \frac{b \log\left(x - \sqrt{-\frac{1}{c}}\right)}{10c^3 \sqrt{\frac{1}{c}}} - \frac{b \log\left(x + \sqrt{-\frac{1}{c}}\right)}{10c^3 \sqrt{\frac{1}{c}}} + \dots \\ \frac{ax^5}{5} \end{cases}$$

input `integrate(x**4*(a+b*atanh(c*x**2)),x)`

output `Piecewise((a*x**5/5 + b*x**5*atanh(c*x**2)/5 + 2*b*x**3/(15*c) - b*sqrt(-1/c)*log(x - sqrt(-1/c))/(10*c**2) + b*sqrt(-1/c)*log(x + sqrt(-1/c))/(10*c**2) - b*log(x - sqrt(-1/c))/(10*c**3*sqrt(1/c)) - b*log(x + sqrt(-1/c))/(10*c**3*sqrt(1/c)) + b*log(x - sqrt(1/c))/(5*c**3*sqrt(1/c)) + b*atanh(c*x**2)/(5*c**3*sqrt(1/c)), Ne(c, 0)), (a*x**5/5, True))`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.06

$$\int x^4(a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{1}{5} ax^5 + \frac{1}{30} \left(6x^5 \operatorname{artanh}(cx^2) + c \left(\frac{4x^3}{c^2} + \frac{6 \arctan(\sqrt{cx})}{c^{\frac{7}{2}}} + \frac{3 \log\left(\frac{cx - \sqrt{c}}{cx + \sqrt{c}}\right)}{c^{\frac{7}{2}}} \right) \right) b$$

input `integrate(x^4*(a+b*arctanh(c*x^2)),x, algorithm="maxima")`

output `1/5*a*x^5 + 1/30*(6*x^5*arctanh(c*x^2) + c*(4*x^3/c^2 + 6*arctan(sqrt(c)*x)/c^(7/2) + 3*log((c*x - sqrt(c))/(c*x + sqrt(c)))/c^(7/2)))*b`

Giac [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.12

$$\int x^4(a + b \operatorname{arctanh}(cx^2)) dx = \frac{1}{10} bx^5 \log\left(-\frac{cx^2+1}{cx^2-1}\right) + \frac{1}{5} ax^5 + \frac{2bx^3}{15c} + \frac{b \arctan(\sqrt{cx})}{5c^{5/2}} + \frac{b \arctan\left(\frac{cx}{\sqrt{-c}}\right)}{5\sqrt{-cc^2}}$$

input `integrate(x^4*(a+b*arctanh(c*x^2)),x, algorithm="giac")`output `1/10*b*x^5*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 1/5*a*x^5 + 2/15*b*x^3/c + 1/5*b*arctan(sqrt(c)*x)/c^(5/2) + 1/5*b*arctan(c*x/sqrt(-c))/(sqrt(-c)*c^2)`**Mupad [B] (verification not implemented)**

Time = 3.85 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.11

$$\int x^4(a + b \operatorname{arctanh}(cx^2)) dx = \frac{ax^5}{5} + \frac{2bx^3}{15c} + \frac{b \operatorname{atan}(\sqrt{c}x)}{5c^{5/2}} + \frac{bx^5 \ln(cx^2+1)}{10} - \frac{bx^5 \ln(1-cx^2)}{10} + \frac{b \operatorname{atan}(\sqrt{c}x \operatorname{li}) \operatorname{li}}{5c^{5/2}}$$

input `int(x^4*(a + b*atanh(c*x^2)),x)`output `(a*x^5)/5 + (2*b*x^3)/(15*c) + (b*atan(c^(1/2)*x))/(5*c^(5/2)) + (b*atan(c^(1/2)*x*li)*li)/(5*c^(5/2)) + (b*x^5*log(c*x^2 + 1))/10 - (b*x^5*log(1 - c*x^2))/10`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.35

$$\int x^4 (a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{6\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) b + 6\sqrt{c} \operatorname{atanh}(cx^2) b + 6 \operatorname{atanh}(cx^2) b c^3 x^5 + 6\sqrt{c} \log(\sqrt{c}x - 1) b - 3\sqrt{c} \log(cx^2 + 1) b}{30c^3}$$

input

```
int(x^4*(a+b*atanh(c*x^2)),x)
```

output

```
(6*sqrt(c)*atan((c*x)/sqrt(c))*b + 6*sqrt(c)*atanh(c*x**2)*b + 6*atanh(c*x**2)*b*c**3*x**5 + 6*sqrt(c)*log(sqrt(c)*x - 1)*b - 3*sqrt(c)*log(c*x**2 + 1)*b + 6*a*c**3*x**5 + 4*b*c**2*x**3)/(30*c**3)
```

3.59 $\int x^2(a + b \operatorname{arctanh}(cx^2)) dx$

Optimal result	517
Mathematica [A] (verified)	517
Rubi [A] (verified)	518
Maple [A] (verified)	520
Fricas [A] (verification not implemented)	520
Sympy [B] (verification not implemented)	521
Maxima [A] (verification not implemented)	521
Giac [A] (verification not implemented)	522
Mupad [B] (verification not implemented)	522
Reduce [B] (verification not implemented)	523

Optimal result

Integrand size = 14, antiderivative size = 63

$$\int x^2(a + b \operatorname{arctanh}(cx^2)) dx = \frac{2bx}{3c} - \frac{b \arctan(\sqrt{cx})}{3c^{3/2}} - \frac{b \operatorname{arctanh}(\sqrt{cx})}{3c^{3/2}} + \frac{1}{3}x^3(a + b \operatorname{arctanh}(cx^2))$$

output

```
2/3*b*x/c-1/3*b*arctan(c^(1/2)*x)/c^(3/2)-1/3*b*arctanh(c^(1/2)*x)/c^(3/2)
+1/3*x^3*(a+b*arctanh(c*x^2))
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.44

$$\int x^2(a + b \operatorname{arctanh}(cx^2)) dx = \frac{2bx}{3c} + \frac{ax^3}{3} - \frac{b \arctan(\sqrt{cx})}{3c^{3/2}} + \frac{1}{3}bx^3 \operatorname{arctanh}(cx^2) + \frac{b \log(1 - \sqrt{cx})}{6c^{3/2}} - \frac{b \log(1 + \sqrt{cx})}{6c^{3/2}}$$

input

```
Integrate[x^2*(a + b*ArcTanh[c*x^2]), x]
```

output

$$(2bx)/(3c) + (ax^3)/3 - (b\text{ArcTan}[\text{Sqrt}[c]x])/(3c^{(3/2)}) + (bx^3\text{ArcTanh}[cx^2])/3 + (b\text{Log}[1 - \text{Sqrt}[c]x])/(6c^{(3/2)}) - (b\text{Log}[1 + \text{Sqrt}[c]x])/(6c^{(3/2)})$$
Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.11, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {6452, 843, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2(a + \text{barctanh}(cx^2)) dx$$

$$\downarrow 6452$$

$$\frac{1}{3}x^3(a + \text{barctanh}(cx^2)) - \frac{2}{3}bc \int \frac{x^4}{1 - c^2x^4} dx$$

$$\downarrow 843$$

$$\frac{1}{3}x^3(a + \text{barctanh}(cx^2)) - \frac{2}{3}bc \left(\frac{\int \frac{1}{1 - c^2x^4} dx}{c^2} - \frac{x}{c^2} \right)$$

$$\downarrow 756$$

$$\frac{1}{3}x^3(a + \text{barctanh}(cx^2)) - \frac{2}{3}bc \left(\frac{\frac{1}{2} \int \frac{1}{1 - cx^2} dx + \frac{1}{2} \int \frac{1}{cx^2 + 1} dx}{c^2} - \frac{x}{c^2} \right)$$

$$\downarrow 216$$

$$\frac{1}{3}x^3(a + \text{barctanh}(cx^2)) - \frac{2}{3}bc \left(\frac{\frac{1}{2} \int \frac{1}{1 - cx^2} dx + \frac{\arctan(\sqrt{cx})}{2\sqrt{c}}}{c^2} - \frac{x}{c^2} \right)$$

$$\downarrow 219$$

$$\frac{1}{3}x^3(a + \text{barctanh}(cx^2)) - \frac{2}{3}bc \left(\frac{\frac{\arctan(\sqrt{cx})}{2\sqrt{c}} + \frac{\text{arctanh}(\sqrt{cx})}{2\sqrt{c}}}{c^2} - \frac{x}{c^2} \right)$$

input `Int[x^2*(a + b*ArcTanh[c*x^2]),x]`

output `(-2*b*c*(-(x/c^2) + (ArcTan[Sqrt[c]*x]/(2*Sqrt[c]) + ArcTanh[Sqrt[c]*x]/(2*Sqrt[c]))/c^2))/3 + (x^3*(a + b*ArcTanh[c*x^2]))/3`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 756 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 843 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Simp[a*c^n*(m - n + 1)/(b*(m + n*p + 1)) Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.81

method	result
default	$\frac{ax^3}{3} + \frac{bx^3 \operatorname{arctanh}(cx^2)}{3} + \frac{2bx}{3c} - \frac{b \operatorname{arctan}(\sqrt{c}x)}{3c^{\frac{3}{2}}} - \frac{b \operatorname{arctanh}(\sqrt{c}x)}{3c^{\frac{3}{2}}}$
parts	$\frac{ax^3}{3} + \frac{bx^3 \operatorname{arctanh}(cx^2)}{3} + \frac{2bx}{3c} - \frac{b \operatorname{arctan}(\sqrt{c}x)}{3c^{\frac{3}{2}}} - \frac{b \operatorname{arctanh}(\sqrt{c}x)}{3c^{\frac{3}{2}}}$
risch	$\frac{bx^3 \ln(cx^2+1)}{6} - \frac{bx^3 \ln(-cx^2+1)}{6} + \frac{ax^3}{3} - \frac{b \ln(1+\sqrt{c}x)}{6c^{\frac{3}{2}}} + \frac{b \ln(\sqrt{c}x-1)}{6c^{\frac{3}{2}}} + \frac{2bx}{3c} - \frac{\sqrt{-c} \ln(\sqrt{-c}x-1)b}{6c^2} + \frac{\sqrt{-c} \ln(1-\sqrt{-c}x)b}{6c^2}$

input `int(x^2*(a+b*arctanh(c*x^2)),x,method=_RETURNVERBOSE)`

output $\frac{1}{3}ax^3 + \frac{1}{3}bx^3 \operatorname{arctanh}(cx^2) + \frac{2}{3}bx/c - \frac{1}{3}b \operatorname{arctan}(c^{1/2}x)/c^{3/2} - \frac{1}{3}b \operatorname{arctanh}(c^{1/2}x)/c^{3/2}$

Fricas [A] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 97 vs. 2(47) = 94.

Time = 0.09 (sec) , antiderivative size = 186, normalized size of antiderivative = 2.95

$$\int x^2(a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \left[\frac{bc^2 x^3 \log\left(-\frac{cx^2+1}{cx^2-1}\right) + 2ac^2 x^3 + 4bcx - 2b\sqrt{c} \operatorname{arctan}(\sqrt{c}x) + b\sqrt{c} \log\left(\frac{cx^2-2\sqrt{c}x+1}{cx^2-1}\right)}{6c^2}, \frac{bc^2 x^3 \log\left(-\frac{cx^2+1}{cx^2-1}\right)}{6c^2} \right]$$

input `integrate(x^2*(a+b*arctanh(c*x^2)),x, algorithm="fricas")`

output $\left[\frac{1}{6}(b*c^2*x^3*\log(-(c*x^2 + 1)/(c*x^2 - 1)) + 2*a*c^2*x^3 + 4*b*c*x - 2*b*\sqrt{c}*\operatorname{arctan}(\sqrt{c}*x) + b*\sqrt{c}*\log((c*x^2 - 2*\sqrt{c}*x + 1)/(c*x^2 - 1)))/c^2, \frac{1}{6}(b*c^2*x^3*\log(-(c*x^2 + 1)/(c*x^2 - 1)) + 2*a*c^2*x^3 + 4*b*c*x + 2*b*\sqrt{-c}*\operatorname{arctan}(\sqrt{-c}*x) - b*\sqrt{-c}*\log((c*x^2 + 2*\sqrt{-c}*x - 1)/(c*x^2 + 1)))/c^2 \right]$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 670 vs. $2(56) = 112$.

Time = 2.93 (sec) , antiderivative size = 670, normalized size of antiderivative = 10.63

$$\int x^2(a + \operatorname{barctanh}(cx^2)) dx$$

$$= \begin{cases} \frac{4ac^2x^3\sqrt{-\frac{1}{c}}}{12c^2\sqrt{-\frac{1}{c}}+12c^2\sqrt{\frac{1}{c}}} + \frac{4ac^2x^3\sqrt{\frac{1}{c}}}{12c^2\sqrt{-\frac{1}{c}}+12c^2\sqrt{\frac{1}{c}}} + \frac{4bc^2x^3\sqrt{-\frac{1}{c}}\operatorname{atanh}(cx^2)}{12c^2\sqrt{-\frac{1}{c}}+12c^2\sqrt{\frac{1}{c}}} + \frac{4bc^2x^3\sqrt{\frac{1}{c}}\operatorname{atanh}(cx^2)}{12c^2\sqrt{-\frac{1}{c}}+12c^2\sqrt{\frac{1}{c}}} - \frac{bc^2(-\frac{1}{c})^{\frac{3}{2}}\sqrt{\frac{1}{c}}\log\left(x+\sqrt{-\frac{1}{c}}\right)}{12c^2\sqrt{-\frac{1}{c}}+12c^2\sqrt{\frac{1}{c}}} \\ \frac{ax^3}{3} \end{cases}$$

input `integrate(x**2*(a+b*atanh(c*x**2)),x)`

output

```
Piecewise((4*a*c**2*x**3*sqrt(-1/c)/(12*c**2*sqrt(-1/c) + 12*c**2*sqrt(1/c)) + 4*a*c**2*x**3*sqrt(1/c)/(12*c**2*sqrt(-1/c) + 12*c**2*sqrt(1/c)) + 4*b*c**2*x**3*sqrt(-1/c)*atanh(c*x**2)/(12*c**2*sqrt(-1/c) + 12*c**2*sqrt(1/c)) + 4*b*c**2*x**3*sqrt(1/c)*atanh(c*x**2)/(12*c**2*sqrt(-1/c) + 12*c**2*sqrt(1/c)) - b*c**2*(-1/c)**(3/2)*sqrt(1/c)*log(x + sqrt(-1/c))/(12*c**2*sqrt(-1/c) + 12*c**2*sqrt(1/c)) + b*c**2*sqrt(-1/c)*(1/c)**(3/2)*log(x + sqrt(-1/c))/(12*c**2*sqrt(-1/c) + 12*c**2*sqrt(1/c)) + 8*b*c*x*sqrt(-1/c)/(12*c**2*sqrt(-1/c) + 12*c**2*sqrt(1/c)) + 8*b*c*x*sqrt(1/c)/(12*c**2*sqrt(-1/c) + 12*c**2*sqrt(1/c)) - 6*b*c*sqrt(-1/c)*sqrt(1/c)*log(x + sqrt(-1/c))/(12*c**2*sqrt(-1/c) + 12*c**2*sqrt(1/c)) + 4*b*c*sqrt(-1/c)*sqrt(1/c)*log(x - sqrt(1/c))/(12*c**2*sqrt(-1/c) + 12*c**2*sqrt(1/c)) + 4*b*c*sqrt(-1/c)*sqrt(1/c)*atanh(c*x**2)/(12*c**2*sqrt(-1/c) + 12*c**2*sqrt(1/c)) - 4*b*log(x - sqrt(-1/c))/(12*c**2*sqrt(-1/c) + 12*c**2*sqrt(1/c)) + 4*b*log(x - sqrt(1/c))/(12*c**2*sqrt(-1/c) + 12*c**2*sqrt(1/c)) + 4*b*atanh(c*x**2)/(12*c**2*sqrt(-1/c) + 12*c**2*sqrt(1/c)), Ne(c, 0)), (a*x**3/3, True))
```

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.05

$$\int x^2(a + \operatorname{barctanh}(cx^2)) dx$$

$$= \frac{1}{3} ax^3 + \frac{1}{6} \left(2x^3 \operatorname{artanh}(cx^2) + c \left(\frac{4x}{c^2} - \frac{2 \arctan(\sqrt{cx})}{c^{\frac{5}{2}}} + \frac{\log\left(\frac{cx-\sqrt{c}}{cx+\sqrt{c}}\right)}{c^{\frac{5}{2}}} \right) \right) b$$

input `integrate(x^2*(a+b*arctanh(c*x^2)),x, algorithm="maxima")`

output $\frac{1}{3}ax^3 + \frac{1}{6}(2x^3\operatorname{arctanh}(cx^2) + c(4x/c^2 - 2\operatorname{arctan}(\sqrt{c}x)/c^{5/2} + \log((cx - \sqrt{c})/(cx + \sqrt{c}))/c^{5/2}))*b$

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.19

$$\int x^2(a + b\operatorname{arctanh}(cx^2)) dx = -\frac{1}{3}bc^5 \left(\frac{\operatorname{arctan}(\sqrt{cx})}{c^{\frac{13}{2}}} - \frac{\operatorname{arctan}\left(\frac{cx}{\sqrt{-c}}\right)}{\sqrt{-cc^6}} \right) + \frac{1}{6}bx^3 \log\left(-\frac{cx^2+1}{cx^2-1}\right) + \frac{1}{3}ax^3 + \frac{2bx}{3c}$$

input `integrate(x^2*(a+b*arctanh(c*x^2)),x, algorithm="giac")`

output $-\frac{1}{3}b*c^5*(\operatorname{arctan}(\sqrt{c}x)/c^{13/2} - \operatorname{arctan}(cx/\sqrt{-c})/(\sqrt{-c}*c^6)) + \frac{1}{6}b*x^3*\log(-(cx^2+1)/(cx^2-1)) + \frac{1}{3}a*x^3 + \frac{2}{3}b*x/c$

Mupad [B] (verification not implemented)

Time = 3.83 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.11

$$\int x^2(a + b\operatorname{arctanh}(cx^2)) dx = \frac{ax^3}{3} - \frac{b\operatorname{atan}(\sqrt{c}x)}{3c^{3/2}} + \frac{2bx}{3c} + \frac{bx^3 \ln(cx^2+1)}{6} - \frac{bx^3 \ln(1-cx^2)}{6} + \frac{b\operatorname{atan}(\sqrt{c}x) \operatorname{li}}{3c^{3/2}}$$

input `int(x^2*(a + b*atanh(c*x^2)),x)`

output $(a*x^3)/3 - (b*\operatorname{atan}(c^{1/2}*x))/(3*c^{3/2}) + (b*\operatorname{atan}(c^{1/2}*x)*\operatorname{li})/(3*c^{3/2}) + (2*b*x)/(3*c) + (b*x^3*\log(cx^2+1))/6 - (b*x^3*\log(1-cx^2))/6$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.33

$$\int x^2 (a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{-2\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) b + 2\sqrt{c} \operatorname{atanh}(cx^2) b + 2 \operatorname{atanh}(cx^2) b c^2 x^3 + 2\sqrt{c} \log(\sqrt{c}x - 1) b - \sqrt{c} \log(cx^2 + 1) b}{6c^2}$$

input

```
int(x^2*(a+b*atanh(c*x^2)),x)
```

output

```
( - 2*sqrt(c)*atan((c*x)/sqrt(c))*b + 2*sqrt(c)*atanh(c*x**2)*b + 2*atanh(
c*x**2)*b*c**2*x**3 + 2*sqrt(c)*log(sqrt(c)*x - 1)*b - sqrt(c)*log(c*x**2
+ 1)*b + 2*a*c**2*x**3 + 4*b*c*x)/(6*c**2)
```


3.60 $\int (a + b \operatorname{arctanh}(cx^2)) dx$

Optimal result	524
Mathematica [A] (verified)	524
Rubi [A] (verified)	525
Maple [A] (verified)	525
Fricas [B] (verification not implemented)	526
Sympy [A] (verification not implemented)	526
Maxima [A] (verification not implemented)	527
Giac [B] (verification not implemented)	528
Mupad [B] (verification not implemented)	528
Reduce [B] (verification not implemented)	529

Optimal result

Integrand size = 10, antiderivative size = 44

$$\int (a + b \operatorname{arctanh}(cx^2)) dx = ax + \frac{b \arctan(\sqrt{cx})}{\sqrt{c}} - \frac{b \operatorname{arctanh}(\sqrt{cx})}{\sqrt{c}} + b \operatorname{arctanh}(cx^2)$$

output

```
a*x+b*arctan(c^(1/2)*x)/c^(1/2)-b*arctanh(c^(1/2)*x)/c^(1/2)+b*x*arctanh(c*x^2)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.30

$$\int (a + b \operatorname{arctanh}(cx^2)) dx = ax + b \operatorname{arctanh}(cx^2) + \frac{b(2 \arctan(\sqrt{cx}) + \log(1 - \sqrt{cx}) - \log(1 + \sqrt{cx}))}{2\sqrt{c}}$$

input

```
Integrate[a + b*ArcTanh[c*x^2], x]
```

output

```
a*x + b*x*ArcTanh[c*x^2] + (b*(2*ArcTan[Sqrt[c]*x] + Log[1 - Sqrt[c]*x] - Log[1 + Sqrt[c]*x]))/(2*Sqrt[c])
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{arctanh}(cx^2)) dx$$

↓ 2009

$$ax + \frac{b \arctan(\sqrt{cx})}{\sqrt{c}} + bx \operatorname{arctanh}(cx^2) - \frac{b \operatorname{arctanh}(\sqrt{cx})}{\sqrt{c}}$$

input `Int[a + b*ArcTanh[c*x^2], x]`

output `a*x + (b*ArcTan[Sqrt[c]*x])/Sqrt[c] - (b*ArcTanh[Sqrt[c]*x])/Sqrt[c] + b*x*ArcTanh[c*x^2]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.84

method	result
default	$ax + \frac{b \arctan(\sqrt{cx})}{\sqrt{c}} - \frac{b \operatorname{arctanh}(\sqrt{cx})}{\sqrt{c}} + bx \operatorname{arctanh}(cx^2)$
parts	$ax + \frac{b \arctan(\sqrt{cx})}{\sqrt{c}} - \frac{b \operatorname{arctanh}(\sqrt{cx})}{\sqrt{c}} + bx \operatorname{arctanh}(cx^2)$
risch	$ax + \frac{bx \ln(cx^2+1)}{2} - \frac{bx \ln(-cx^2+1)}{2} + \frac{b\sqrt{-c} \ln(cx+\sqrt{-c})}{2c} - \frac{b\sqrt{-c} \ln(-cx+\sqrt{-c})}{2c} + \frac{b \ln(1-\sqrt{cx})}{2\sqrt{c}} - \frac{b \ln(1+\sqrt{cx})}{2\sqrt{c}}$

input `int(a+b*arctanh(c*x^2),x,method=_RETURNVERBOSE)`

output `a*x+b*arctan(c^(1/2)*x)/c^(1/2)-b*arctanh(c^(1/2)*x)/c^(1/2)+b*x*arctanh(c*x^2)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 84 vs. $2(36) = 72$.

Time = 0.09 (sec) , antiderivative size = 160, normalized size of antiderivative = 3.64

$$\int (a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \left[\frac{bcx \log\left(-\frac{cx^2+1}{cx^2-1}\right) + 2acx + 2b\sqrt{c} \arctan(\sqrt{cx}) + b\sqrt{c} \log\left(\frac{cx^2-2\sqrt{cx}+1}{cx^2-1}\right)}{2c}, \frac{bcx \log\left(-\frac{cx^2+1}{cx^2-1}\right) + 2acx +}{2c} \right]$$

input `integrate(a+b*arctanh(c*x^2),x, algorithm="fricas")`

output `[1/2*(b*c*x*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 2*a*c*x + 2*b*sqrt(c)*arctan(sqrt(c)*x) + b*sqrt(c)*log((c*x^2 - 2*sqrt(c)*x + 1)/(c*x^2 - 1)))/c, 1/2*(b*c*x*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 2*a*c*x + 2*b*sqrt(-c)*arctan(sqrt(-c)*x) - b*sqrt(-c)*log((c*x^2 - 2*sqrt(-c)*x - 1)/(c*x^2 + 1)))/c]`

Sympy [A] (verification not implemented)

Time = 2.21 (sec) , antiderivative size = 702, normalized size of antiderivative = 15.95

$$\int (a + b \operatorname{arctanh}(cx^2)) dx = \text{Too large to display}$$

input `integrate(a+b*atanh(c*x**2),x)`

output

```
a*x + b*Piecewise((4*c*x*sqrt(-1/c)*sqrt(1/c)*atanh(c*x**2)/(c**2*(-1/c)**
(3/2)*sqrt(1/c) - c**2*sqrt(-1/c)*(1/c)**(3/2) + 6*c*sqrt(-1/c)*sqrt(1/c))
- c*(-1/c)**(3/2)*log(x + sqrt(-1/c))/(c**2*(-1/c)**(3/2)*sqrt(1/c) - c**
2*sqrt(-1/c)*(1/c)**(3/2) + 6*c*sqrt(-1/c)*sqrt(1/c)) + c*(1/c)**(3/2)*log
(x + sqrt(-1/c))/(c**2*(-1/c)**(3/2)*sqrt(1/c) - c**2*sqrt(-1/c)*(1/c)**(3
/2) + 6*c*sqrt(-1/c)*sqrt(1/c)) - 2*sqrt(-1/c)*log(x - sqrt(-1/c))/(c**2*(
-1/c)**(3/2)*sqrt(1/c) - c**2*sqrt(-1/c)*(1/c)**(3/2) + 6*c*sqrt(-1/c)*sqr
t(1/c)) - 3*sqrt(-1/c)*log(x + sqrt(-1/c))/(c**2*(-1/c)**(3/2)*sqrt(1/c) -
c**2*sqrt(-1/c)*(1/c)**(3/2) + 6*c*sqrt(-1/c)*sqrt(1/c)) + 4*sqrt(-1/c)*l
og(x - sqrt(1/c))/(c**2*(-1/c)**(3/2)*sqrt(1/c) - c**2*sqrt(-1/c)*(1/c)**(
3/2) + 6*c*sqrt(-1/c)*sqrt(1/c)) + 4*sqrt(-1/c)*atanh(c*x**2)/(c**2*(-1/c)
**3/2)*sqrt(1/c) - c**2*sqrt(-1/c)*(1/c)**(3/2) + 6*c*sqrt(-1/c)*sqrt(1/c
)) + 2*sqrt(1/c)*log(x - sqrt(-1/c))/(c**2*(-1/c)**(3/2)*sqrt(1/c) - c**2*
sqrt(-1/c)*(1/c)**(3/2) + 6*c*sqrt(-1/c)*sqrt(1/c)) - 3*sqrt(1/c)*log(x +
sqrt(-1/c))/(c**2*(-1/c)**(3/2)*sqrt(1/c) - c**2*sqrt(-1/c)*(1/c)**(3/2) +
6*c*sqrt(-1/c)*sqrt(1/c)), Ne(c, 0)), (0, True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.25

$$\int (a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{1}{2} \left(c \left(\frac{2 \arctan(\sqrt{cx})}{c^{\frac{3}{2}}} + \frac{\log\left(\frac{cx-\sqrt{c}}{cx+\sqrt{c}}\right)}{c^{\frac{3}{2}}} \right) + 2x \operatorname{arctanh}(cx^2) \right) b + ax$$

input

```
integrate(a+b*arctanh(c*x^2),x, algorithm="maxima")
```

output

```
1/2*(c*(2*arctan(sqrt(c)*x)/c^(3/2) + log((c*x - sqrt(c))/(c*x + sqrt(c)))
/c^(3/2)) + 2*x*arctanh(c*x^2))*b + a*x
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 83 vs. 2(36) = 72.

Time = 0.12 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.89

$$\int (a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{1}{2} \left(c \left(\frac{2 \sqrt{|c|} \arctan \left(x \sqrt{|c|} \right)}{c^2} - \frac{\sqrt{|c|} \log \left(\left| x + \frac{1}{\sqrt{|c|}} \right| \right)}{c^2} + \frac{\sqrt{|c|} \log \left(\left| x - \frac{1}{\sqrt{|c|}} \right| \right)}{c^2} \right) + x \log \left(-\frac{cx^2 + 1}{cx^2 - 1} \right) \right) + ax$$

input `integrate(a+b*arctanh(c*x^2),x, algorithm="giac")`

output `1/2*(c*(2*sqrt(abs(c))*arctan(x*sqrt(abs(c)))/c^2 - sqrt(abs(c))*log(abs(x + 1/sqrt(abs(c))))/c^2 + sqrt(abs(c))*log(abs(x - 1/sqrt(abs(c))))/c^2) + x*log(-(c*x^2 + 1)/(c*x^2 - 1))*b + a*x`

Mupad [B] (verification not implemented)

Time = 3.54 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.25

$$\int (a + b \operatorname{arctanh}(cx^2)) dx = ax + \frac{b \operatorname{atan}(\sqrt{c}x)}{\sqrt{c}} + \frac{bx \ln(cx^2 + 1)}{2} - \frac{bx \ln(1 - cx^2)}{2} + \frac{b \operatorname{atan}(\sqrt{c}x) \operatorname{li} \operatorname{li}}{\sqrt{c}}$$

input `int(a + b*atanh(c*x^2),x)`

output `a*x + (b*atan(c^(1/2)*x))/c^(1/2) + (b*atan(c^(1/2)*x*1i)*1i)/c^(1/2) + (b*x*log(c*x^2 + 1))/2 - (b*x*log(1 - c*x^2))/2`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.61

$$\int (a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{2\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) b + 2\sqrt{c} \operatorname{atanh}(cx^2) b + 2 \operatorname{atanh}(cx^2) bcx + 2\sqrt{c} \log(\sqrt{c}x - 1) b - \sqrt{c} \log(cx^2 + 1) b + 2a}{2c}$$

input

```
int(a+b*atanh(c*x^2),x)
```

output

```
(2*sqrt(c)*atan((c*x)/sqrt(c))*b + 2*sqrt(c)*atanh(c*x**2)*b + 2*atanh(c*x**2)*b*c*x + 2*sqrt(c)*log(sqrt(c)*x - 1)*b - sqrt(c)*log(c*x**2 + 1)*b + 2*a*c*x)/(2*c)
```

3.61 $\int \frac{a+b\operatorname{arctanh}(cx^2)}{x^2} dx$

Optimal result	530
Mathematica [A] (verified)	530
Rubi [A] (verified)	531
Maple [A] (verified)	532
Fricas [A] (verification not implemented)	533
Sympy [B] (verification not implemented)	533
Maxima [A] (verification not implemented)	534
Giac [B] (verification not implemented)	535
Mupad [B] (verification not implemented)	535
Reduce [B] (verification not implemented)	536

Optimal result

Integrand size = 14, antiderivative size = 46

$$\int \frac{a + b\operatorname{arctanh}(cx^2)}{x^2} dx = b\sqrt{c} \arctan(\sqrt{cx}) + b\sqrt{c}\operatorname{arctanh}(\sqrt{cx}) - \frac{a + b\operatorname{arctanh}(cx^2)}{x}$$

output

```
b*c^(1/2)*arctan(c^(1/2)*x)+b*c^(1/2)*arctanh(c^(1/2)*x)-(a+b*arctanh(c*x^2))/x
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.63

$$\int \frac{a + b\operatorname{arctanh}(cx^2)}{x^2} dx = -\frac{a}{x} + b\sqrt{c} \arctan(\sqrt{cx}) - \frac{b\operatorname{arctanh}(cx^2)}{x} - \frac{1}{2}b\sqrt{c} \log(1 - \sqrt{cx}) + \frac{1}{2}b\sqrt{c} \log(1 + \sqrt{cx})$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])/x^2,x]
```

output

```
-(a/x) + b*Sqrt[c]*ArcTan[Sqrt[c]*x] - (b*ArcTanh[c*x^2])/x - (b*Sqrt[c]*Log[1 - Sqrt[c]*x])/2 + (b*Sqrt[c]*Log[1 + Sqrt[c]*x])/2
```

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.20, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6452, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^2} dx$$

$$\downarrow 6452$$

$$2bc \int \frac{1}{1 - c^2 x^4} dx - \frac{a + b \operatorname{arctanh}(cx^2)}{x}$$

$$\downarrow 756$$

$$2bc \left(\frac{1}{2} \int \frac{1}{1 - cx^2} dx + \frac{1}{2} \int \frac{1}{cx^2 + 1} dx \right) - \frac{a + b \operatorname{arctanh}(cx^2)}{x}$$

$$\downarrow 216$$

$$2bc \left(\frac{1}{2} \int \frac{1}{1 - cx^2} dx + \frac{\arctan(\sqrt{cx})}{2\sqrt{c}} \right) - \frac{a + b \operatorname{arctanh}(cx^2)}{x}$$

$$\downarrow 219$$

$$2bc \left(\frac{\arctan(\sqrt{cx})}{2\sqrt{c}} + \frac{\operatorname{arctanh}(\sqrt{cx})}{2\sqrt{c}} \right) - \frac{a + b \operatorname{arctanh}(cx^2)}{x}$$

input `Int[(a + b*ArcTanh[c*x^2])/x^2,x]`

output `2*b*c*(ArcTan[Sqrt[c]*x]/(2*Sqrt[c]) + ArcTanh[Sqrt[c]*x]/(2*Sqrt[c])) - (a + b*ArcTanh[c*x^2])/x`

Definitions of rubi rules used

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 756 `Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)]*(b_))^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.91

method	result	size
default	$-\frac{a}{x} - \frac{b \operatorname{arctanh}(cx^2)}{x} + b\sqrt{c} \arctan(\sqrt{c}x) + b\sqrt{c} \operatorname{arctanh}(\sqrt{c}x)$	42
parts	$-\frac{a}{x} - \frac{b \operatorname{arctanh}(cx^2)}{x} + b\sqrt{c} \arctan(\sqrt{c}x) + b\sqrt{c} \operatorname{arctanh}(\sqrt{c}x)$	42
risch	$-\frac{b \ln(cx^2+1)}{2x} + b\sqrt{c} \arctan(\sqrt{c}x) - \frac{a}{x} + \frac{b \ln(-cx^2+1)}{2x} + b\sqrt{c} \operatorname{arctanh}(\sqrt{c}x)$	59

input `int((a+b*arctanh(c*x^2))/x^2,x,method=_RETURNVERBOSE)`

output

```
-a/x-b/x*arctanh(c*x^2)+b*c^(1/2)*arctan(c^(1/2)*x)+b*c^(1/2)*arctanh(c^(1/2)*x)
```

Fricas [A] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 82 vs. $2(38) = 76$.

Time = 0.10 (sec) , antiderivative size = 157, normalized size of antiderivative = 3.41

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^2} dx$$

$$= \left[\frac{2b\sqrt{cx} \arctan(\sqrt{cx}) + b\sqrt{cx} \log\left(\frac{cx^2 + 2\sqrt{cx} + 1}{cx^2 - 1}\right) - b \log\left(-\frac{cx^2 + 1}{cx^2 - 1}\right) - 2a}{2x}, \right.$$

$$\left. - \frac{2b\sqrt{-cx} \arctan(\sqrt{-cx}) - b\sqrt{-cx} \log\left(\frac{cx^2 + 2\sqrt{-cx} - 1}{cx^2 + 1}\right) + b \log\left(-\frac{cx^2 + 1}{cx^2 - 1}\right) + 2a}{2x} \right]$$

input

```
integrate((a+b*arctanh(c*x^2))/x^2,x, algorithm="fricas")
```

output

```
[1/2*(2*b*sqrt(c)*x*arctan(sqrt(c)*x) + b*sqrt(c)*x*log((c*x^2 + 2*sqrt(c)*x + 1)/(c*x^2 - 1)) - b*log(-(c*x^2 + 1)/(c*x^2 - 1)) - 2*a)/x, -1/2*(2*b*sqrt(-c)*x*arctan(sqrt(-c)*x) - b*sqrt(-c)*x*log((c*x^2 + 2*sqrt(-c)*x - 1)/(c*x^2 + 1)) + b*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 2*a)/x]
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1374 vs. $2(41) = 82$.

Time = 3.94 (sec) , antiderivative size = 1374, normalized size of antiderivative = 29.87

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^2} dx = \text{Too large to display}$$

input

```
integrate((a+b*atanh(c*x**2))/x**2,x)
```

output

```
Piecewise((-a/x, Eq(c, 0)), (-a - oo*b)/x, Eq(c, -1/x**2)), (-a + oo*b)/
x, Eq(c, x**(-2))), (-a*c*x**4*sqrt(-1/c)/(c*x**5*sqrt(-1/c) + c*x**5*sqrt
(1/c) - x*sqrt(-1/c)/c - x*sqrt(1/c)/c) - a*c*x**4*sqrt(1/c)/(c*x**5*sqrt(
-1/c) + c*x**5*sqrt(1/c) - x*sqrt(-1/c)/c - x*sqrt(1/c)/c) + a*sqrt(-1/c)/
(c**2*x**5*sqrt(-1/c) + c**2*x**5*sqrt(1/c) - x*sqrt(-1/c) - x*sqrt(1/c))
+ a*sqrt(1/c)/(c**2*x**5*sqrt(-1/c) + c**2*x**5*sqrt(1/c) - x*sqrt(-1/c) -
x*sqrt(1/c)) + b*c**2*x**5*sqrt(-1/c)*sqrt(1/c)*log(x + sqrt(-1/c))/(c*x*
*5*sqrt(-1/c) + c*x**5*sqrt(1/c) - x*sqrt(-1/c)/c - x*sqrt(1/c)/c) - b*c**
2*x**5*sqrt(-1/c)*sqrt(1/c)*log(x - sqrt(1/c))/(c*x**5*sqrt(-1/c) + c*x**5
*sqrt(1/c) - x*sqrt(-1/c)/c - x*sqrt(1/c)/c) - b*c**2*x**5*sqrt(-1/c)*sqrt
(1/c)*atanh(c*x**2)/(c*x**5*sqrt(-1/c) + c*x**5*sqrt(1/c) - x*sqrt(-1/c)/c
- x*sqrt(1/c)/c) + b*c*x**5*log(x - sqrt(-1/c))/(c*x**5*sqrt(-1/c) + c*x*
*5*sqrt(1/c) - x*sqrt(-1/c)/c - x*sqrt(1/c)/c) - b*c*x**5*log(x - sqrt(1/c
))/(c*x**5*sqrt(-1/c) + c*x**5*sqrt(1/c) - x*sqrt(-1/c)/c - x*sqrt(1/c)/c)
- b*c*x**5*atanh(c*x**2)/(c*x**5*sqrt(-1/c) + c*x**5*sqrt(1/c) - x*sqrt(-
1/c)/c - x*sqrt(1/c)/c) - b*c*x**4*sqrt(-1/c)*atanh(c*x**2)/(c*x**5*sqrt(-
1/c) + c*x**5*sqrt(1/c) - x*sqrt(-1/c)/c - x*sqrt(1/c)/c) - b*c*x**4*sqrt(
1/c)*atanh(c*x**2)/(c*x**5*sqrt(-1/c) + c*x**5*sqrt(1/c) - x*sqrt(-1/c)/c
- x*sqrt(1/c)/c) - b*x*sqrt(-1/c)*sqrt(1/c)*log(x + sqrt(-1/c))/(c*x**5*sq
rt(-1/c) + c*x**5*sqrt(1/c) - x*sqrt(-1/c)/c - x*sqrt(1/c)/c) + b*x*sqr...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.33

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^2} dx$$

$$= \frac{1}{2} \left(c \left(\frac{2 \arctan(\sqrt{c}x)}{\sqrt{c}} - \frac{\log\left(\frac{cx-\sqrt{c}}{cx+\sqrt{c}}\right)}{\sqrt{c}} \right) - \frac{2 \operatorname{arctanh}(cx^2)}{x} \right) b - \frac{a}{x}$$

input

```
integrate((a+b*arctanh(c*x^2))/x^2,x, algorithm="maxima")
```

output

```
1/2*(c*(2*arctan(sqrt(c)*x)/sqrt(c) - log((c*x - sqrt(c))/(c*x + sqrt(c)))
/sqrt(c)) - 2*arctanh(c*x^2)/x)*b - a/x
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 79 vs. 2(38) = 76.

Time = 0.14 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.72

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^2} dx$$

$$= \frac{1}{2} bc \left(\frac{2 \arctan \left(x \sqrt{|c|} \right)}{\sqrt{|c|}} + \frac{\log \left(\left| x + \frac{1}{\sqrt{|c|}} \right| \right)}{\sqrt{|c|}} - \frac{\log \left(\left| x - \frac{1}{\sqrt{|c|}} \right| \right)}{\sqrt{|c|}} \right)$$

$$- \frac{b \log \left(-\frac{cx^2+1}{cx^2-1} \right)}{2x} - \frac{a}{x}$$

input `integrate((a+b*arctanh(c*x^2))/x^2,x, algorithm="giac")`

output `1/2*b*c*(2*arctan(x*sqrt(abs(c)))/sqrt(abs(c)) + log(abs(x + 1/sqrt(abs(c))))/sqrt(abs(c)) - log(abs(x - 1/sqrt(abs(c))))/sqrt(abs(c))) - 1/2*b*log(-(c*x^2 + 1)/(c*x^2 - 1))/x - a/x`

Mupad [B] (verification not implemented)

Time = 3.74 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.35

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^2} dx = b \sqrt{c} \operatorname{atan}(\sqrt{c} x) - \frac{a}{x} - \frac{b \ln(cx^2 + 1)}{2x}$$

$$+ \frac{b \ln(1 - cx^2)}{2x} - b \sqrt{c} \operatorname{atan}(\sqrt{c} x \operatorname{li}) \operatorname{li}$$

input `int((a + b*atanh(c*x^2))/x^2,x)`

output `b*c^(1/2)*atan(c^(1/2)*x) - a/x - b*c^(1/2)*atan(c^(1/2)*x*1i)*1i - (b*log(c*x^2 + 1))/(2*x) + (b*log(1 - c*x^2))/(2*x)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.52

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^2} dx$$

$$= \frac{2\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) bx - 2\sqrt{c} \operatorname{atanh}(cx^2) bx - 2 \operatorname{atanh}(cx^2) b - 2\sqrt{c} \log(\sqrt{c}x - 1) bx + \sqrt{c} \log(cx^2 + 1) bx - 2a}{2x}$$

input

```
int((a+b*atanh(c*x^2))/x^2,x)
```

output

```
(2*sqrt(c)*atan((c*x)/sqrt(c))*b*x - 2*sqrt(c)*atanh(c*x**2)*b*x - 2*atanh(c*x**2)*b - 2*sqrt(c)*log(sqrt(c)*x - 1)*b*x + sqrt(c)*log(c*x**2 + 1)*b*x - 2*a)/(2*x)
```

3.62 $\int \frac{a+b\operatorname{arctanh}(cx^2)}{x^4} dx$

Optimal result	537
Mathematica [A] (verified)	537
Rubi [A] (verified)	538
Maple [A] (verified)	540
Fricas [A] (verification not implemented)	540
Sympy [B] (verification not implemented)	541
Maxima [A] (verification not implemented)	542
Giac [B] (verification not implemented)	542
Mupad [B] (verification not implemented)	543
Reduce [B] (verification not implemented)	543

Optimal result

Integrand size = 14, antiderivative size = 63

$$\int \frac{a + b\operatorname{arctanh}(cx^2)}{x^4} dx = -\frac{2bc}{3x} - \frac{1}{3}bc^{3/2} \arctan(\sqrt{cx}) + \frac{1}{3}bc^{3/2}\operatorname{arctanh}(\sqrt{cx}) - \frac{a + b\operatorname{arctanh}(cx^2)}{3x^3}$$

output

```
-2/3*b*c/x-1/3*b*c^(3/2)*arctan(c^(1/2)*x)+1/3*b*c^(3/2)*arctanh(c^(1/2)*x)-1/3*(a+b*arctanh(c*x^2))/x^3
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.44

$$\int \frac{a + b\operatorname{arctanh}(cx^2)}{x^4} dx = -\frac{a}{3x^3} - \frac{2bc}{3x} - \frac{1}{3}bc^{3/2} \arctan(\sqrt{cx}) - \frac{b\operatorname{arctanh}(cx^2)}{3x^3} - \frac{1}{6}bc^{3/2} \log(1 - \sqrt{cx}) + \frac{1}{6}bc^{3/2} \log(1 + \sqrt{cx})$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])/x^4, x]
```

output

$$-1/3*a/x^3 - (2*b*c)/(3*x) - (b*c^{(3/2)}*ArcTan[Sqrt[c]*x])/3 - (b*ArcTanh[c*x^2])/(3*x^3) - (b*c^{(3/2)}*Log[1 - Sqrt[c]*x])/6 + (b*c^{(3/2)}*Log[1 + Sqrt[c]*x])/6$$
Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.10, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {6452, 847, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^4} dx$$

$$\downarrow 6452$$

$$\frac{2}{3}bc \int \frac{1}{x^2(1-c^2x^4)} dx - \frac{a + b \operatorname{arctanh}(cx^2)}{3x^3}$$

$$\downarrow 847$$

$$\frac{2}{3}bc \left(c^2 \int \frac{x^2}{1-c^2x^4} dx - \frac{1}{x} \right) - \frac{a + b \operatorname{arctanh}(cx^2)}{3x^3}$$

$$\downarrow 827$$

$$\frac{2}{3}bc \left(c^2 \left(\int \frac{1}{1-cx^2} dx - \int \frac{1}{cx^2+1} dx \right) - \frac{1}{x} \right) - \frac{a + b \operatorname{arctanh}(cx^2)}{3x^3}$$

$$\downarrow 216$$

$$\frac{2}{3}bc \left(c^2 \left(\int \frac{1}{1-cx^2} dx - \frac{\arctan(\sqrt{cx})}{2c^{3/2}} \right) - \frac{1}{x} \right) - \frac{a + b \operatorname{arctanh}(cx^2)}{3x^3}$$

$$\downarrow 219$$

$$\frac{2}{3}bc \left(c^2 \left(\frac{\operatorname{arctanh}(\sqrt{cx})}{2c^{3/2}} - \frac{\arctan(\sqrt{cx})}{2c^{3/2}} \right) - \frac{1}{x} \right) - \frac{a + b \operatorname{arctanh}(cx^2)}{3x^3}$$

input

$$\text{Int}[(a + b*ArcTanh[c*x^2])/x^4, x]$$

output $(2*b*c*(-x^{(-1)} + c^2*(-1/2*ArcTan[Sqrt[c]*x]/c^{(3/2)} + ArcTanh[Sqrt[c]*x]/(2*c^{(3/2)})))/3 - (a + b*ArcTanh[c*x^2])/(3*x^3)$

Defintions of rubi rules used

rule 216 $Int[((a_) + (b_)*(x_)^2)^{(-1)}, x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] \&\& PosQ[a/b] \&\& (GtQ[a, 0] || GtQ[b, 0])$

rule 219 $Int[((a_) + (b_)*(x_)^2)^{(-1)}, x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] \&\& NegQ[a/b] \&\& (GtQ[a, 0] || LtQ[b, 0])$

rule 827 $Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x], x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] \&\& !GtQ[a/b, 0]$

rule 847 $Int[((c_)*(x_))^{(m_)*((a_) + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] := Simp[(c*x)^{(m+1)*((a + b*x^n)^{(p+1))/(a*c*(m+1))}, x] - Simp[b*(m+n*(p+1)+1)/(a*c^n*(m+1)) Int[(c*x)^{(m+n)*(a + b*x^n)^p}, x], x] /; FreeQ[{a, b, c, p}, x] \&\& IGtQ[n, 0] \&\& LtQ[m, -1] \&\& IntBinomialQ[a, b, c, n, m, p, x]$

rule 6452 $Int[((a_) + ArcTanh[(c_)*(x_)^{(n_)}]*(b_))^{(p_)*(x_)^{(m_)}}, x_Symbol] := Simp[x^{(m+1)*((a + b*ArcTanh[c*x^n])^p/(m+1))}, x] - Simp[b*c*n*(p/(m+1)) Int[x^{(m+n)*((a + b*ArcTanh[c*x^n])^{(p-1)/(1 - c^2*x^{(2*n)})}), x], x] /; FreeQ[{a, b, c, m, n}, x] \&\& IGtQ[p, 0] \&\& (EqQ[p, 1] || (EqQ[n, 1] \&\& IntegerQ[m])) \&\& NeQ[m, -1]$

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.81

method	result	size
default	$-\frac{a}{3x^3} - \frac{b \operatorname{arctanh}(cx^2)}{3x^3} - \frac{bc^{\frac{3}{2}} \operatorname{arctan}(\sqrt{cx})}{3} - \frac{2bc}{3x} + \frac{bc^{\frac{3}{2}} \operatorname{arctanh}(\sqrt{cx})}{3}$	51
parts	$-\frac{a}{3x^3} - \frac{b \operatorname{arctanh}(cx^2)}{3x^3} - \frac{bc^{\frac{3}{2}} \operatorname{arctan}(\sqrt{cx})}{3} - \frac{2bc}{3x} + \frac{bc^{\frac{3}{2}} \operatorname{arctanh}(\sqrt{cx})}{3}$	51
risch	$-\frac{b \ln(cx^2+1)}{6x^3} - \frac{a}{3x^3} + \frac{b \ln(-cx^2+1)}{6x^3} - \frac{2bc}{3x} + \frac{bc^{\frac{3}{2}} \operatorname{arctanh}(\sqrt{cx})}{3} - \frac{bc^{\frac{3}{2}} \operatorname{arctan}(\sqrt{cx})}{3}$	68

input `int((a+b*arctanh(c*x^2))/x^4,x,method=_RETURNVERBOSE)`

output
$$-1/3*a/x^3-1/3*b/x^3*arctanh(c*x^2)-1/3*b*c^{(3/2)}*arctan(c^{(1/2)}*x)-2/3*b*c/x+1/3*b*c^{(3/2)}*arctanh(c^{(1/2)}*x)$$

Fricas [A] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs. 2(47) = 94.

Time = 0.09 (sec) , antiderivative size = 181, normalized size of antiderivative = 2.87

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^4} dx$$

$$= \left[\frac{2bc^{\frac{3}{2}}x^3 \operatorname{arctan}(\sqrt{cx}) - bc^{\frac{3}{2}}x^3 \log\left(\frac{cx^2+2\sqrt{cx}+1}{cx^2-1}\right) + 4bcx^2 + b \log\left(-\frac{cx^2+1}{cx^2-1}\right) + 2a}{6x^3}, \right.$$

$$\left. \frac{2b\sqrt{-cx}x^3 \operatorname{arctan}(\sqrt{-cx}) - b\sqrt{-cx}x^3 \log\left(\frac{cx^2-2\sqrt{-cx}-1}{cx^2+1}\right) + 4bcx^2 + b \log\left(-\frac{cx^2+1}{cx^2-1}\right) + 2a}{6x^3} \right]$$

input `integrate((a+b*arctanh(c*x^2))/x^4,x, algorithm="fricas")`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.03

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^4} dx = -\frac{1}{6} \left(\left(2\sqrt{c} \arctan(\sqrt{c}x) + \sqrt{c} \log\left(\frac{cx - \sqrt{c}}{cx + \sqrt{c}}\right) + \frac{4}{x} \right) c + \frac{2 \operatorname{arctanh}(cx^2)}{x^3} \right) b - \frac{a}{3x^3}$$

input `integrate((a+b*arctanh(c*x^2))/x^4,x, algorithm="maxima")`

output `-1/6*((2*sqrt(c)*arctan(sqrt(c)*x) + sqrt(c)*log((c*x - sqrt(c))/(c*x + sqrt(c)))) + 4/x)*c + 2*arctanh(c*x^2)/x^3)*b - 1/3*a/x^3`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs. $2(47) = 94$.

Time = 0.14 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.51

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^4} dx = -\frac{bc^3 \arctan\left(x\sqrt{|c|}\right)}{3|c|^{\frac{3}{2}}} + \frac{bc^3 \log\left(\left|x + \frac{1}{\sqrt{|c|}}\right|\right)}{6|c|^{\frac{3}{2}}} - \frac{bc^3 \log\left(\left|x - \frac{1}{\sqrt{|c|}}\right|\right)}{6|c|^{\frac{3}{2}}} - \frac{b \log\left(-\frac{cx^2+1}{cx^2-1}\right)}{6x^3} - \frac{2bcx^2 + a}{3x^3}$$

input `integrate((a+b*arctanh(c*x^2))/x^4,x, algorithm="giac")`

output `-1/3*b*c^3*arctan(x*sqrt(abs(c)))/abs(c)^(3/2) + 1/6*b*c^3*log(abs(x + 1/sqrt(abs(c))))/abs(c)^(3/2) - 1/6*b*c^3*log(abs(x - 1/sqrt(abs(c))))/abs(c)^(3/2) - 1/6*b*log(-(c*x^2 + 1)/(c*x^2 - 1))/x^3 - 1/3*(2*b*c*x^2 + a)/x^3`

Mupad [B] (verification not implemented)

Time = 3.85 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.13

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^4} dx = \frac{b \ln(1 - cx^2)}{6x^3} - \frac{bc^{3/2} \operatorname{atan}(\sqrt{c}x)}{3} - \frac{b \ln(cx^2 + 1)}{6x^3} - \frac{2bcx^2 + a}{3x^3} - \frac{bc^{3/2} \operatorname{atan}(\sqrt{c}x \operatorname{li}) \operatorname{li}}{3}$$

input `int((a + b*atanh(c*x^2))/x^4,x)`output `(b*log(1 - c*x^2))/(6*x^3) - (b*c^(3/2)*atan(c^(1/2)*x))/3 - (b*c^(3/2)*atan(c^(1/2)*x*1i)*1i)/3 - (b*log(c*x^2 + 1))/(6*x^3) - (a + 2*b*c*x^2)/(3*x^3)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.41

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^4} dx = \frac{-2\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) bcx^3 - 2\sqrt{c} \operatorname{atanh}(cx^2) bcx^3 - 2 \operatorname{atanh}(cx^2) b - 2\sqrt{c} \log(\sqrt{c}x - 1) bcx^3 + \sqrt{c} \log(cx^2 + 1) bcx^3 - 2a - 4bcx^2}{6x^3}$$

input `int((a+b*atanh(c*x^2))/x^4,x)`output `(- 2*sqrt(c)*atan((c*x)/sqrt(c))*b*c*x**3 - 2*sqrt(c)*atanh(c*x**2)*b*c*x**3 - 2*atanh(c*x**2)*b - 2*sqrt(c)*log(sqrt(c)*x - 1)*b*c*x**3 + sqrt(c)*log(c*x**2 + 1)*b*c*x**3 - 2*a - 4*b*c*x**2)/(6*x**3)`

3.63 $\int \frac{a+b\operatorname{arctanh}(cx^2)}{x^6} dx$

Optimal result	544
Mathematica [A] (verified)	544
Rubi [A] (verified)	545
Maple [A] (verified)	547
Fricas [A] (verification not implemented)	547
Sympy [B] (verification not implemented)	548
Maxima [A] (verification not implemented)	549
Giac [A] (verification not implemented)	549
Mupad [B] (verification not implemented)	550
Reduce [B] (verification not implemented)	550

Optimal result

Integrand size = 14, antiderivative size = 63

$$\int \frac{a + b\operatorname{arctanh}(cx^2)}{x^6} dx = -\frac{2bc}{15x^3} + \frac{1}{5}bc^{5/2} \arctan(\sqrt{cx}) + \frac{1}{5}bc^{5/2}\operatorname{arctanh}(\sqrt{cx}) - \frac{a + b\operatorname{arctanh}(cx^2)}{5x^5}$$

output

```
-2/15*b*c/x^3+1/5*b*c^(5/2)*arctan(c^(1/2)*x)+1/5*b*c^(5/2)*arctanh(c^(1/2)*x)-1/5*(a+b*arctanh(c*x^2))/x^5
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.44

$$\int \frac{a + b\operatorname{arctanh}(cx^2)}{x^6} dx = -\frac{a}{5x^5} - \frac{2bc}{15x^3} + \frac{1}{5}bc^{5/2} \arctan(\sqrt{cx}) - \frac{b\operatorname{arctanh}(cx^2)}{5x^5} - \frac{1}{10}bc^{5/2} \log(1 - \sqrt{cx}) + \frac{1}{10}bc^{5/2} \log(1 + \sqrt{cx})$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])/x^6, x]
```

output

```
-1/5*a/x^5 - (2*b*c)/(15*x^3) + (b*c^(5/2)*ArcTan[Sqrt[c]*x])/5 - (b*ArcTanh[c*x^2])/(5*x^5) - (b*c^(5/2)*Log[1 - Sqrt[c]*x])/10 + (b*c^(5/2)*Log[1 + Sqrt[c]*x])/10
```

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.13, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {6452, 847, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^6} dx$$

$$\downarrow 6452$$

$$\frac{2}{5}bc \int \frac{1}{x^4(1-c^2x^4)} dx - \frac{a + b \operatorname{arctanh}(cx^2)}{5x^5}$$

$$\downarrow 847$$

$$\frac{2}{5}bc \left(c^2 \int \frac{1}{1-c^2x^4} dx - \frac{1}{3x^3} \right) - \frac{a + b \operatorname{arctanh}(cx^2)}{5x^5}$$

$$\downarrow 756$$

$$\frac{2}{5}bc \left(c^2 \left(\frac{1}{2} \int \frac{1}{1-cx^2} dx + \frac{1}{2} \int \frac{1}{cx^2+1} dx \right) - \frac{1}{3x^3} \right) - \frac{a + b \operatorname{arctanh}(cx^2)}{5x^5}$$

$$\downarrow 216$$

$$\frac{2}{5}bc \left(c^2 \left(\frac{1}{2} \int \frac{1}{1-cx^2} dx + \frac{\arctan(\sqrt{cx})}{2\sqrt{c}} \right) - \frac{1}{3x^3} \right) - \frac{a + b \operatorname{arctanh}(cx^2)}{5x^5}$$

$$\downarrow 219$$

$$\frac{2}{5}bc \left(c^2 \left(\frac{\arctan(\sqrt{cx})}{2\sqrt{c}} + \frac{\operatorname{arctanh}(\sqrt{cx})}{2\sqrt{c}} \right) - \frac{1}{3x^3} \right) - \frac{a + b \operatorname{arctanh}(cx^2)}{5x^5}$$

input

```
Int[(a + b*ArcTanh[c*x^2])/x^6,x]
```

output $(2*b*c*(-1/3*1/x^3 + c^2*(ArcTan[Sqrt[c]*x]/(2*Sqrt[c]) + ArcTanh[Sqrt[c]*x]/(2*Sqrt[c])))/5 - (a + b*ArcTanh[c*x^2])/(5*x^5)$

Defintions of rubi rules used

rule 216 $Int[((a_) + (b_)*(x_)^2)^{-1}, x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] \&\& PosQ[a/b] \&\& (GtQ[a, 0] || GtQ[b, 0])$

rule 219 $Int[((a_) + (b_)*(x_)^2)^{-1}, x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] \&\& NegQ[a/b] \&\& (GtQ[a, 0] || LtQ[b, 0])$

rule 756 $Int[((a_) + (b_)*(x_)^4)^{-1}, x_Symbol] :> With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] \&\& !GtQ[a/b, 0]$

rule 847 $Int[((c_)*(x_))^{(m_)}*((a_) + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] :> Simp[(c*x)^{(m+1)}*((a + b*x^n)^{(p+1)}/(a*c*(m+1))), x] - Simp[b*(m+n*(p+1)+1)/(a*c^n*(m+1)) Int[(c*x)^{(m+n)}*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] \&\& IGtQ[n, 0] \&\& LtQ[m, -1] \&\& IntBinomialQ[a, b, c, n, m, p, x]$

rule 6452 $Int[((a_) + ArcTanh[(c_)*(x_)^{(n_)}])*(b_)^{(p_)}*(x_)^{(m_)}, x_Symbol] :> Simp[x^{(m+1)}*((a + b*ArcTanh[c*x^n])^p/(m+1)), x] - Simp[b*c*n*(p/(m+1)) Int[x^{(m+n)}*((a + b*ArcTanh[c*x^n])^{(p-1)})/(1 - c^2*x^{(2*n)}), x], x] /; FreeQ[{a, b, c, m, n}, x] \&\& IGtQ[p, 0] \&\& (EqQ[p, 1] || (EqQ[n, 1] \&\& IntegerQ[m])) \&\& NeQ[m, -1]$

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.81

method	result	size
default	$-\frac{a}{5x^5} - \frac{b \operatorname{arctanh}(cx^2)}{5x^5} + \frac{bc^{\frac{5}{2}} \arctan(\sqrt{c}x)}{5} + \frac{bc^{\frac{5}{2}} \operatorname{arctanh}(\sqrt{c}x)}{5} - \frac{2bc}{15x^3}$	51
parts	$-\frac{a}{5x^5} - \frac{b \operatorname{arctanh}(cx^2)}{5x^5} + \frac{bc^{\frac{5}{2}} \arctan(\sqrt{c}x)}{5} + \frac{bc^{\frac{5}{2}} \operatorname{arctanh}(\sqrt{c}x)}{5} - \frac{2bc}{15x^3}$	51
risch	$-\frac{b \ln(cx^2+1)}{10x^5} - \frac{a}{5x^5} + \frac{b \ln(-cx^2+1)}{10x^5} - \frac{2bc}{15x^3} + \frac{bc^{\frac{5}{2}} \operatorname{arctanh}(\sqrt{c}x)}{5} + \frac{bc^{\frac{5}{2}} \arctan(\sqrt{c}x)}{5}$	68

input `int((a+b*arctanh(c*x^2))/x^6,x,method=_RETURNVERBOSE)`

output
$$-1/5*a/x^5 - 1/5*b/x^5*\operatorname{arctanh}(c*x^2) + 1/5*b*c^{(5/2)}*\arctan(c^{(1/2)}*x) + 1/5*b*c^{(5/2)}*\operatorname{arctanh}(c^{(1/2)}*x) - 2/15*b*c/x^3$$

Fricas [A] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 100 vs. $2(47) = 94$.

Time = 0.09 (sec) , antiderivative size = 187, normalized size of antiderivative = 2.97

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^6} dx$$

$$= \left[\frac{6bc^{\frac{5}{2}}x^5 \arctan(\sqrt{c}x) + 3bc^{\frac{5}{2}}x^5 \log\left(\frac{cx^2+2\sqrt{c}x+1}{cx^2-1}\right) - 4bcx^2 - 3b \log\left(-\frac{cx^2+1}{cx^2-1}\right) - 6a}{30x^5}, \right.$$

$$\left. \frac{6b\sqrt{-c}x^5 \arctan(\sqrt{-c}x) - 3b\sqrt{-c}x^5 \log\left(\frac{cx^2+2\sqrt{-c}x-1}{cx^2+1}\right) + 4bcx^2 + 3b \log\left(-\frac{cx^2+1}{cx^2-1}\right) + 6a}{30x^5} \right]$$

input `integrate((a+b*arctanh(c*x^2))/x^6,x, algorithm="fricas")`

output

```
[1/30*(6*b*c^(5/2)*x^5*arctan(sqrt(c)*x) + 3*b*c^(5/2)*x^5*log((c*x^2 + 2*sqrt(c)*x + 1)/(c*x^2 - 1)) - 4*b*c*x^2 - 3*b*log(-(c*x^2 + 1)/(c*x^2 - 1)) - 6*a)/x^5, -1/30*(6*b*sqrt(-c)*c^2*x^5*arctan(sqrt(-c)*x) - 3*b*sqrt(-c)*c^2*x^5*log((c*x^2 + 2*sqrt(-c)*x - 1)/(c*x^2 + 1)) + 4*b*c*x^2 + 3*b*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 6*a)/x^5]
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1948 vs. $2(60) = 120$.

Time = 6.92 (sec) , antiderivative size = 1948, normalized size of antiderivative = 30.92

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^6} dx = \text{Too large to display}$$

input

```
integrate((a+b*atanh(c*x**2))/x**6,x)
```

output

```
Piecewise((-a/(5*x**5), Eq(c, 0)), (-a - oo*b)/(5*x**5), Eq(c, -1/x**2)), (-a + oo*b)/(5*x**5), Eq(c, x**(-2))), (-3*a*c*x**4*sqrt(-1/c)/(15*c*x**9*sqrt(-1/c) + 15*c*x**9*sqrt(1/c) - 15*x**5*sqrt(-1/c)/c - 15*x**5*sqrt(1/c)/c) - 3*a*c*x**4*sqrt(1/c)/(15*c*x**9*sqrt(-1/c) + 15*c*x**9*sqrt(1/c) - 15*x**5*sqrt(-1/c)/c - 15*x**5*sqrt(1/c)/c) + 3*a*sqrt(-1/c)/(15*c**2*x**9*sqrt(-1/c) + 15*c**2*x**9*sqrt(1/c) - 15*x**5*sqrt(-1/c) - 15*x**5*sqrt(1/c)) + 3*a*sqrt(1/c)/(15*c**2*x**9*sqrt(-1/c) + 15*c**2*x**9*sqrt(1/c) - 15*x**5*sqrt(-1/c) - 15*x**5*sqrt(1/c)) + 3*b*c**4*x**9*sqrt(-1/c)*sqrt(1/c)*log(x + sqrt(-1/c))/(15*c*x**9*sqrt(-1/c) + 15*c*x**9*sqrt(1/c) - 15*x**5*sqrt(-1/c)/c - 15*x**5*sqrt(1/c)/c) - 3*b*c**4*x**9*sqrt(-1/c)*sqrt(1/c)*log(x - sqrt(1/c))/(15*c*x**9*sqrt(-1/c) + 15*c*x**9*sqrt(1/c) - 15*x**5*sqrt(-1/c)/c - 15*x**5*sqrt(1/c)/c) - 3*b*c**4*x**9*sqrt(-1/c)*sqrt(1/c)*atanh(c*x**2)/(15*c*x**9*sqrt(-1/c) + 15*c*x**9*sqrt(1/c) - 15*x**5*sqrt(-1/c)/c - 15*x**5*sqrt(1/c)/c) + 3*b*c**3*x**9*log(x - sqrt(-1/c))/(15*c*x**9*sqrt(-1/c) + 15*c*x**9*sqrt(1/c) - 15*x**5*sqrt(-1/c)/c - 15*x**5*sqrt(1/c)/c) - 3*b*c**3*x**9*log(x + sqrt(1/c))/(15*c*x**9*sqrt(-1/c) + 15*c*x**9*sqrt(1/c) - 15*x**5*sqrt(-1/c)/c - 15*x**5*sqrt(1/c)/c) - 3*b*c**3*x**9*atanh(c*x**2)/(15*c*x**9*sqrt(-1/c) + 15*c*x**9*sqrt(1/c) - 15*x**5*sqrt(-1/c)/c - 15*x**5*sqrt(1/c)/c) - 2*b*c**2*x**6*sqrt(-1/c)/(15*c*x**9*sqrt(-1/c) + 15*c*x**9*sqrt(1/c) - 15*x**5*sqrt(-1/c)/c - 15*x**5*sqrt(1/c)/c)...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.05

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^6} dx$$

$$= \frac{1}{30} \left(\left(6c^{\frac{3}{2}} \arctan(\sqrt{cx}) - 3c^{\frac{3}{2}} \log\left(\frac{cx - \sqrt{c}}{cx + \sqrt{c}}\right) - \frac{4}{x^3} \right) c - \frac{6 \operatorname{arctanh}(cx^2)}{x^5} \right) b - \frac{a}{5x^5}$$

input `integrate((a+b*arctanh(c*x^2))/x^6,x, algorithm="maxima")`output `1/30*((6*c^(3/2)*arctan(sqrt(c)*x) - 3*c^(3/2)*log((c*x - sqrt(c))/(c*x + sqrt(c)))) - 4/x^3)*c - 6*arctanh(c*x^2)/x^5)*b - 1/5*a/x^5`**Giac [A] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.44

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^6} dx$$

$$= \frac{1}{10} bc^3 \left(\frac{2 \arctan\left(x\sqrt{|c|}\right)}{\sqrt{|c|}} + \frac{\log\left(\left|x + \frac{1}{\sqrt{|c|}}\right|\right)}{\sqrt{|c|}} - \frac{\log\left(\left|x - \frac{1}{\sqrt{|c|}}\right|\right)}{\sqrt{|c|}} \right)$$

$$- \frac{b \log\left(-\frac{cx^2+1}{cx^2-1}\right)}{10x^5} - \frac{2bcx^2 + 3a}{15x^5}$$

input `integrate((a+b*arctanh(c*x^2))/x^6,x, algorithm="giac")`output `1/10*b*c^3*(2*arctan(x*sqrt(abs(c)))/sqrt(abs(c)) + log(abs(x + 1/sqrt(abs(c))))/sqrt(abs(c)) - log(abs(x - 1/sqrt(abs(c))))/sqrt(abs(c))) - 1/10*b*log(-(c*x^2 + 1)/(c*x^2 - 1))/x^5 - 1/15*(2*b*c*x^2 + 3*a)/x^5`

Mupad [B] (verification not implemented)

Time = 3.88 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.13

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^6} dx = \frac{b c^{5/2} \operatorname{atan}(\sqrt{c} x)}{5} - \frac{\frac{2bcx^2}{3} + a}{5x^5} - \frac{b \ln(cx^2 + 1)}{10x^5} + \frac{b \ln(1 - cx^2)}{10x^5} - \frac{b c^{5/2} \operatorname{atan}(\sqrt{c} x \operatorname{li} \operatorname{li})}{5}$$

input `int((a + b*atanh(c*x^2))/x^6,x)`output `(b*c^(5/2)*atan(c^(1/2)*x))/5 - (a + (2*b*c*x^2)/3)/(5*x^5) - (b*c^(5/2)*atan(c^(1/2)*x*1i)*1i)/5 - (b*log(c*x^2 + 1))/(10*x^5) + (b*log(1 - c*x^2))/(10*x^5)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.56

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{x^6} dx = \frac{6\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) b c^2 x^5 - 6\sqrt{c} \operatorname{atanh}(cx^2) b c^2 x^5 - 6 \operatorname{atanh}(cx^2) b - 6\sqrt{c} \log(\sqrt{c} x - 1) b c^2 x^5 + 3\sqrt{c} \log(c)}{30x^5}$$

input `int((a+b*atanh(c*x^2))/x^6,x)`output `(6*sqrt(c)*atan((c*x)/sqrt(c))*b*c**2*x**5 - 6*sqrt(c)*atanh(c*x**2)*b*c**2*x**5 - 6*atanh(c*x**2)*b - 6*sqrt(c)*log(sqrt(c)*x - 1)*b*c**2*x**5 + 3*sqrt(c)*log(c*x**2 + 1)*b*c**2*x**5 - 6*a - 4*b*c*x**2)/(30*x**5)`

3.64 $\int x^7(a + b \operatorname{arctanh}(cx^2))^2 dx$

Optimal result	551
Mathematica [A] (verified)	551
Rubi [A] (verified)	552
Maple [A] (verified)	555
Fricas [A] (verification not implemented)	555
Sympy [A] (verification not implemented)	556
Maxima [A] (verification not implemented)	556
Giac [A] (verification not implemented)	557
Mupad [B] (verification not implemented)	558
Reduce [B] (verification not implemented)	559

Optimal result

Integrand size = 16, antiderivative size = 125

$$\int x^7(a + b \operatorname{arctanh}(cx^2))^2 dx = \frac{abx^2}{4c^3} + \frac{b^2x^4}{24c^2} + \frac{b^2x^2 \operatorname{arctanh}(cx^2)}{4c^3} + \frac{bx^6(a + b \operatorname{arctanh}(cx^2))}{12c} - \frac{(a + b \operatorname{arctanh}(cx^2))^2}{8c^4} + \frac{1}{8}x^8(a + b \operatorname{arctanh}(cx^2))^2 + \frac{b^2 \log(1 - c^2x^4)}{6c^4}$$

output

```
1/4*a*b*x^2/c^3+1/24*b^2*x^4/c^2+1/4*b^2*x^2*arctanh(c*x^2)/c^3+1/12*b*x^6*(a+b*arctanh(c*x^2))/c-1/8*(a+b*arctanh(c*x^2))^2/c^4+1/8*x^8*(a+b*arctanh(c*x^2))^2+1/6*b^2*ln(-c^2*x^4+1)/c^4
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.17

$$\int x^7(a + b \operatorname{arctanh}(cx^2))^2 dx = \frac{6abcx^2 + b^2c^2x^4 + 2abc^3x^6 + 3a^2c^4x^8 + 2bcx^2(3ac^3x^6 + b(3 + c^2x^4)) \operatorname{arctanh}(cx^2) + 3b^2(-1 + c^4x^8) \operatorname{arctanh}(cx^2)}{24c^4}$$

input `Integrate[x^7*(a + b*ArcTanh[c*x^2])^2,x]`

output $(6*a*b*c*x^2 + b^2*c^2*x^4 + 2*a*b*c^3*x^6 + 3*a^2*c^4*x^8 + 2*b*c*x^2*(3*a*c^3*x^6 + b*(3 + c^2*x^4))*ArcTanh[c*x^2] + 3*b^2*(-1 + c^4*x^8)*ArcTanh[c*x^2]^2 + b*(3*a + 4*b)*Log[1 - c*x^2] - 3*a*b*Log[1 + c*x^2] + 4*b^2*Log[1 + c*x^2])/(24*c^4)$

Rubi [A] (verified)

Time = 1.03 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.22, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {6454, 6452, 6542, 6452, 243, 49, 2009, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^7 (a + \operatorname{barctanh}(cx^2))^2 dx$$

$$\downarrow 6454$$

$$\frac{1}{2} \int x^6 (a + \operatorname{barctanh}(cx^2))^2 dx^2$$

$$\downarrow 6452$$

$$\frac{1}{2} \left(\frac{1}{4} x^8 (a + \operatorname{barctanh}(cx^2))^2 - \frac{1}{2} bc \int \frac{x^8 (a + \operatorname{barctanh}(cx^2))}{1 - c^2 x^4} dx^2 \right)$$

$$\downarrow 6542$$

$$\frac{1}{2} \left(\frac{1}{4} x^8 (a + \operatorname{barctanh}(cx^2))^2 - \frac{1}{2} bc \left(\frac{\int \frac{x^4 (a + \operatorname{barctanh}(cx^2))}{1 - c^2 x^4} dx^2}{c^2} - \frac{\int x^4 (a + \operatorname{barctanh}(cx^2)) dx^2}{c^2} \right) \right)$$

$$\downarrow 6452$$

$$\frac{1}{2} \left(\frac{1}{4} x^8 (a + \operatorname{barctanh}(cx^2))^2 - \frac{1}{2} bc \left(\frac{\int \frac{x^4 (a + \operatorname{barctanh}(cx^2))}{1 - c^2 x^4} dx^2}{c^2} - \frac{\frac{1}{3} x^6 (a + \operatorname{barctanh}(cx^2)) - \frac{1}{3} bc \int \frac{x^6}{1 - c^2 x^4} dx^2}{c^2} \right) \right)$$

↓ 243

$$\frac{1}{2} \left(\frac{1}{4} x^8 (a + \operatorname{barctanh}(cx^2))^2 - \frac{1}{2} bc \left(\frac{\int \frac{x^4 (a + \operatorname{barctanh}(cx^2))}{1 - c^2 x^4} dx^2}{c^2} - \frac{\frac{1}{3} x^6 (a + \operatorname{barctanh}(cx^2)) - \frac{1}{6} bc \int \frac{x^4}{1 - c^2 x^4} dx^4}{c^2} \right) \right)$$

↓ 49

$$\frac{1}{2} \left(\frac{1}{4} x^8 (a + \operatorname{barctanh}(cx^2))^2 - \frac{1}{2} bc \left(\frac{\int \frac{x^4 (a + \operatorname{barctanh}(cx^2))}{1 - c^2 x^4} dx^2}{c^2} - \frac{\frac{1}{3} x^6 (a + \operatorname{barctanh}(cx^2)) - \frac{1}{6} bc \int \left(-\frac{1}{c^2} - \frac{1}{c^2 c} \right)}{c^2} \right) \right)$$

↓ 2009

$$\frac{1}{2} \left(\frac{1}{4} x^8 (a + \operatorname{barctanh}(cx^2))^2 - \frac{1}{2} bc \left(\frac{\int \frac{x^4 (a + \operatorname{barctanh}(cx^2))}{1 - c^2 x^4} dx^2}{c^2} - \frac{\frac{1}{3} x^6 (a + \operatorname{barctanh}(cx^2)) - \frac{1}{6} bc \left(-\frac{x^4}{c^2} - \frac{\log(1 - c^2 x^4)}{c} \right)}{c^2} \right) \right)$$

↓ 6542

$$\frac{1}{2} \left(\frac{1}{4} x^8 (a + \operatorname{barctanh}(cx^2))^2 - \frac{1}{2} bc \left(\frac{\frac{\int \frac{a + \operatorname{barctanh}(cx^2)}{1 - c^2 x^4} dx^2}{c^2} - \frac{\int (a + \operatorname{barctanh}(cx^2)) dx^2}{c^2}}{c^2} - \frac{\frac{1}{3} x^6 (a + \operatorname{barctanh}(cx^2))}{c^2} \right) \right)$$

↓ 2009

$$\frac{1}{2} \left(\frac{1}{4} x^8 (a + \operatorname{barctanh}(cx^2))^2 - \frac{1}{2} bc \left(\frac{\frac{\int \frac{a + \operatorname{barctanh}(cx^2)}{1 - c^2 x^4} dx^2}{c^2} - \frac{ax^2 + bx^2 \operatorname{arctanh}(cx^2) + \frac{b \log(1 - c^2 x^4)}{2c}}{c^2}}{c^2} - \frac{\frac{1}{3} x^6 (a + \operatorname{barctanh}(cx^2))}{c^2} \right) \right)$$

↓ 6510

$$\frac{1}{2} \left(\frac{1}{4} x^8 (a + \operatorname{barctanh}(cx^2))^2 - \frac{1}{2} bc \left(\frac{\frac{(a + \operatorname{barctanh}(cx^2))^2}{2bc^3} - \frac{ax^2 + bx^2 \operatorname{arctanh}(cx^2) + \frac{b \log(1 - c^2 x^4)}{2c}}{c^2}}{c^2} - \frac{\frac{1}{3} x^6 (a + \operatorname{barctanh}(cx^2))}{c^2} \right) \right)$$

input `Int [x^7*(a + b*ArcTanh[c*x^2])^2,x]`

output
$$\frac{((x^8(a + b \operatorname{ArcTanh}[c x^2])^2)/4 - (b c (-((x^6(a + b \operatorname{ArcTanh}[c x^2]))) / 3 - (b c (-x^4/c^2) - \operatorname{Log}[1 - c^2 x^4]/c^4))/6)/c^2) + ((a + b \operatorname{ArcTanh}[c x^2])^2/(2 b c^3) - (a x^2 + b x^2 \operatorname{ArcTanh}[c x^2] + (b \operatorname{Log}[1 - c^2 x^4])/(2 c))/c^2)/c^2)/2}{2}$$

Defintions of rubi rules used

rule 49
$$\operatorname{Int}[(a + b(x))^{m_1} (c + d(x))^{n_1}, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{ExpandIntegrand}[(a + b x)^m (c + d x)^n, x], x] /; \operatorname{FreeQ}\{a, b, c, d, x\} \&\& \operatorname{IGtQ}[m, 0] \&\& \operatorname{IGtQ}[m + n + 2, 0]$$

rule 243
$$\operatorname{Int}(x)^{m_1} (a + b(x)^2)^{p_1}, x_Symbol] \rightarrow \operatorname{Simp}[1/2 \operatorname{Subst}[\operatorname{Int}[x^{(m-1)/2} (a + b x)^p, x], x, x^2], x] /; \operatorname{FreeQ}\{a, b, m, p, x\} \&\& \operatorname{IntegerQ}[(m-1)/2]$$

rule 2009
$$\operatorname{Int}[u, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$$

rule 6452
$$\operatorname{Int}[(a + \operatorname{ArcTanh}[c(x)^n] b)^{p_1} (x)^{m_1}, x_Symbol] \rightarrow \operatorname{Simp}[x^{m+1} (a + b \operatorname{ArcTanh}[c x^n])^p / (m+1), x] - \operatorname{Simp}[b c^n (p/(m+1)) \operatorname{Int}[x^{m+n} (a + b \operatorname{ArcTanh}[c x^n])^{p-1} / (1 - c^2 x^{2n})], x], x] /; \operatorname{FreeQ}\{a, b, c, m, n, x\} \&\& \operatorname{IGtQ}[p, 0] \&\& (\operatorname{EqQ}[p, 1] \|\| (\operatorname{EqQ}[n, 1] \&\& \operatorname{IntegerQ}[m])) \&\& \operatorname{NeQ}[m, -1]$$

rule 6454
$$\operatorname{Int}[(a + \operatorname{ArcTanh}[c(x)^n] b)^{p_1} (x)^{m_1}, x_Symbol] \rightarrow \operatorname{Simp}[1/n \operatorname{Subst}[\operatorname{Int}[x^{(\operatorname{Simplify}[(m+1)/n] - 1) (a + b \operatorname{ArcTanh}[c x])^p}, x], x, x^n], x] /; \operatorname{FreeQ}\{a, b, c, m, n, x\} \&\& \operatorname{IGtQ}[p, 1] \&\& \operatorname{IntegerQ}[\operatorname{Simplify}[(m+1)/n]]$$

rule 6510
$$\operatorname{Int}[(a + \operatorname{ArcTanh}[c(x)] b)^{p_1} / ((d) + (e)(x)^2), x_Symbol] \rightarrow \operatorname{Simp}[(a + b \operatorname{ArcTanh}[c x])^{p+1} / (b c d (p+1)), x] /; \operatorname{FreeQ}\{a, b, c, d, e, p, x\} \&\& \operatorname{EqQ}[c^2 d + e, 0] \&\& \operatorname{NeQ}[p, -1]$$

rule 6542

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^ (p_.)*((f_.)*(x_))^(m_)/((d_.) + (
e_.)*(x_)^2), x_Symbol] :> Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*
x])^p, x], x] - Simp[d*(f^2/e Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/
(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m,
1]
```

Maple [A] (verified)

Time = 0.73 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.30

method	result
parallelrisch	$\frac{3b^2 \operatorname{arctanh}(cx^2)^2 x^8 c^4 + 6ab \operatorname{arctanh}(cx^2) x^8 c^4 + 3a^2 c^4 x^8 + 2b^2 \operatorname{arctanh}(cx^2) x^6 c^3 + 2ab c^3 x^6 + b^2 c^2 x^4 + 6b^2 \operatorname{arctanh}(cx^2) x^2}{24c^4}$
risch	$\frac{b^2(c^4 x^8 - 1) \ln(cx^2 + 1)^2}{32c^4} + \frac{b(-3x^8 b \ln(-cx^2 + 1) c^4 + 6a c^4 x^8 + 2b c^3 x^6 + 6bc x^2 + 3b \ln(-cx^2 + 1)) \ln(cx^2 + 1)}{48c^4} + \frac{b^2 x^8 \ln(-cx^2 + 1)}{32c^4}$
default	Expression too large to display
parts	Expression too large to display

input

```
int(x^7*(a+b*arctanh(c*x^2))^2,x,method=_RETURNVERBOSE)
```

output

```
1/24*(3*b^2*arctanh(c*x^2)^2*x^8*c^4+6*a*b*arctanh(c*x^2)*x^8*c^4+3*a^2*c^
4*x^8+2*b^2*arctanh(c*x^2)*x^6*c^3+2*a*b*c^3*x^6+b^2*c^2*x^4+6*b^2*arctanh
(c*x^2)*x^2*c+6*a*b*c*x^2-3*b^2*arctanh(c*x^2)^2+8*ln(c*x^2-1)*b^2-6*arcta
nh(c*x^2)*a+b+8*arctanh(c*x^2)*b^2+b^2)/c^4
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.41

$$\int x^7 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$= \frac{12 a^2 c^4 x^8 + 8 a b c^3 x^6 + 4 b^2 c^2 x^4 + 24 a b c x^2 + 3 (b^2 c^4 x^8 - b^2) \log\left(-\frac{cx^2+1}{cx^2-1}\right)^2 - 4 (3 a b - 4 b^2) \log(cx^2 + 1)}{96 c^4}$$

input

```
integrate(x^7*(a+b*arctanh(c*x^2))^2,x, algorithm="fricas")
```


output

```
1/96*(12*a^2*c^4*x^8 + 8*a*b*c^3*x^6 + 4*b^2*c^2*x^4 + 24*a*b*c*x^2 + 3*(b^2*c^4*x^8 - b^2)*log(-(c*x^2 + 1)/(c*x^2 - 1))^2 - 4*(3*a*b - 4*b^2)*log(c*x^2 + 1) + 4*(3*a*b + 4*b^2)*log(c*x^2 - 1) + 4*(3*a*b*c^4*x^8 + b^2*c^3*x^6 + 3*b^2*c*x^2)*log(-(c*x^2 + 1)/(c*x^2 - 1)))/c^4
```

Sympy [A] (verification not implemented)

Time = 7.68 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.65

$$\int x^7 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$= \begin{cases} \frac{a^2 x^8}{8} + \frac{abx^8 \operatorname{atanh}(cx^2)}{4} + \frac{abx^6}{12c} + \frac{abx^2}{4c^3} - \frac{ab \operatorname{atanh}(cx^2)}{4c^4} + \frac{b^2 x^8 \operatorname{atanh}^2(cx^2)}{8} + \frac{b^2 x^6 \operatorname{atanh}(cx^2)}{12c} + \frac{b^2 x^4}{24c^2} + \frac{b^2 x^2 \operatorname{atanh}(cx^2)}{4c^3} \\ \frac{a^2 x^8}{8} \end{cases}$$

input

```
integrate(x**7*(a+b*atanh(c*x**2))**2,x)
```

output

```
Piecewise((a**2*x**8/8 + a*b*x**8*atanh(c*x**2)/4 + a*b*x**6/(12*c) + a*b*x**2/(4*c**3) - a*b*atanh(c*x**2)/(4*c**4) + b**2*x**8*atanh(c*x**2)**2/8 + b**2*x**6*atanh(c*x**2)/(12*c) + b**2*x**4/(24*c**2) + b**2*x**2*atanh(c*x**2)/(4*c**3) + b**2*log(x - sqrt(-1/c))/(3*c**4) + b**2*log(x + sqrt(-1/c))/(3*c**4) - b**2*atanh(c*x**2)**2/(8*c**4) - b**2*atanh(c*x**2)/(3*c**4), Ne(c, 0)), (a**2*x**8/8, True))
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.74

$$\int x^7 (a + b \operatorname{arctanh}(cx^2))^2 dx = \frac{1}{8} b^2 x^8 \operatorname{artanh}(cx^2)^2 + \frac{1}{8} a^2 x^8$$

$$+ \frac{1}{24} \left(6 x^8 \operatorname{artanh}(cx^2) + c \left(\frac{2(c^2 x^6 + 3 x^2)}{c^4} - \frac{3 \log(cx^2 + 1)}{c^5} + \frac{3 \log(cx^2 - 1)}{c^5} \right) \right) ab$$

$$+ \frac{1}{96} \left(4 c \left(\frac{2(c^2 x^6 + 3 x^2)}{c^4} - \frac{3 \log(cx^2 + 1)}{c^5} + \frac{3 \log(cx^2 - 1)}{c^5} \right) \operatorname{artanh}(cx^2) + \frac{4 c^2 x^4 - 2(3 \log(cx^2 - 1))}{c^5} \right)$$

input

```
integrate(x^7*(a+b*arctanh(c*x^2))^2,x, algorithm="maxima")
```

output

```
1/8*b^2*x^8*arctanh(c*x^2)^2 + 1/8*a^2*x^8 + 1/24*(6*x^8*arctanh(c*x^2) +
c*(2*(c^2*x^6 + 3*x^2)/c^4 - 3*log(c*x^2 + 1)/c^5 + 3*log(c*x^2 - 1)/c^5))
*a*b + 1/96*(4*c*(2*(c^2*x^6 + 3*x^2)/c^4 - 3*log(c*x^2 + 1)/c^5 + 3*log(c
*x^2 - 1)/c^5)*arctanh(c*x^2) + (4*c^2*x^4 - 2*(3*log(c*x^2 - 1) - 8)*log(
c*x^2 + 1) + 3*log(c*x^2 + 1)^2 + 3*log(c*x^2 - 1)^2 + 16*log(c*x^2 - 1))/
c^4)*b^2
```

Giac [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.40

$$\int x^7(a + b \operatorname{arctanh}(cx^2))^2 dx = \frac{1}{8} a^2 x^8 + \frac{abx^6}{12c} + \frac{b^2 x^4}{24c^2} + \frac{1}{32} \left(b^2 x^8 - \frac{b^2}{c^4} \right) \log \left(\frac{-cx^2 + 1}{cx^2 - 1} \right)^2 + \frac{1}{24} \left(3abx^8 + \frac{b^2 x^6}{c} + \frac{3b^2 x^2}{c^3} \right) \log \left(\frac{-cx^2 + 1}{cx^2 - 1} \right) + \frac{abx^2}{4c^3} - \frac{(3ab - 4b^2) \log(cx^2 + 1)}{24c^4} + \frac{(3ab + 4b^2) \log(cx^2 - 1)}{24c^4}$$

input

```
integrate(x^7*(a+b*arctanh(c*x^2))^2,x, algorithm="giac")
```

output

```
1/8*a^2*x^8 + 1/12*a*b*x^6/c + 1/24*b^2*x^4/c^2 + 1/32*(b^2*x^8 - b^2/c^4)
*log(-(c*x^2 + 1)/(c*x^2 - 1))^2 + 1/24*(3*a*b*x^8 + b^2*x^6/c + 3*b^2*x^2
/c^3)*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 1/4*a*b*x^2/c^3 - 1/24*(3*a*b - 4*b^
2)*log(c*x^2 + 1)/c^4 + 1/24*(3*a*b + 4*b^2)*log(c*x^2 - 1)/c^4
```

Mupad [B] (verification not implemented)

Time = 4.41 (sec) , antiderivative size = 335, normalized size of antiderivative = 2.68

$$\int x^7 (a + b \operatorname{arctanh}(cx^2))^2 dx = \frac{a^2 x^8}{8} + \frac{b^2 \ln(cx^2 - 1)}{6c^4} + \frac{b^2 \ln(cx^2 + 1)}{6c^4} - \frac{b^2 \ln(cx^2 + 1)^2}{32c^4} - \frac{b^2 \ln(1 - cx^2)^2}{32c^4} + \frac{b^2 x^4}{24c^2} + \frac{b^2 x^8 \ln(cx^2 + 1)^2}{32} + \frac{b^2 x^8 \ln(1 - cx^2)^2}{32} + \frac{b^2 x^2 \ln(cx^2 + 1)}{8c^3} - \frac{b^2 x^2 \ln(1 - cx^2)}{8c^3} + \frac{b^2 x^6 \ln(cx^2 + 1)}{24c} - \frac{b^2 x^6 \ln(1 - cx^2)}{24c} + \frac{ab \ln(cx^2 - 1)}{8c^4} - \frac{ab \ln(cx^2 + 1)}{8c^4} + \frac{abx^8 \ln(cx^2 + 1)}{8} - \frac{abx^8 \ln(1 - cx^2)}{8} + \frac{b^2 \ln(cx^2 + 1) \ln(1 - cx^2)}{16c^4} + \frac{abx^2}{4c^3} + \frac{abx^6}{12c} - \frac{b^2 x^8 \ln(cx^2 + 1) \ln(1 - cx^2)}{16}$$

input `int(x^7*(a + b*atanh(c*x^2))^2,x)`output `(a^2*x^8)/8 + (b^2*log(c*x^2 - 1))/(6*c^4) + (b^2*log(c*x^2 + 1))/(6*c^4) - (b^2*log(c*x^2 + 1)^2)/(32*c^4) - (b^2*log(1 - c*x^2)^2)/(32*c^4) + (b^2*x^4)/(24*c^2) + (b^2*x^8*log(c*x^2 + 1)^2)/32 + (b^2*x^8*log(1 - c*x^2)^2)/32 + (b^2*x^2*log(c*x^2 + 1))/(8*c^3) - (b^2*x^2*log(1 - c*x^2))/(8*c^3) + (b^2*x^6*log(c*x^2 + 1))/(24*c) - (b^2*x^6*log(1 - c*x^2))/(24*c) + (a*b*log(c*x^2 - 1))/(8*c^4) - (a*b*log(c*x^2 + 1))/(8*c^4) + (a*b*x^8*log(c*x^2 + 1))/8 - (a*b*x^8*log(1 - c*x^2))/8 + (b^2*log(c*x^2 + 1)*log(1 - c*x^2))/(16*c^4) + (a*b*x^2)/(4*c^3) + (a*b*x^6)/(12*c) - (b^2*x^8*log(c*x^2 + 1)*log(1 - c*x^2))/16`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.27

$$\int x^7 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$= \frac{3 \operatorname{atanh}(cx^2)^2 b^2 c^4 x^8 - 3 \operatorname{atanh}(cx^2)^2 b^2 + 6 \operatorname{atanh}(cx^2) ab c^4 x^8 - 6 \operatorname{atanh}(cx^2) ab + 2 \operatorname{atanh}(cx^2) b^2 c^3 x^6 + 2 a^2 c^3 x^6 + 6 a^2 c^2 x^4 + 2 a^2 c x^2 + 2 a^2}{24 c^4}$$

input

```
int(x^7*(a+b*atanh(c*x^2))^2,x)
```

output

```
(3*atanh(c*x**2)**2*b**2*c**4*x**8 - 3*atanh(c*x**2)**2*b**2 + 6*atanh(c*x**2)*a*b*c**4*x**8 - 6*atanh(c*x**2)*a*b + 2*atanh(c*x**2)*b**2*c**3*x**6 + 6*atanh(c*x**2)*b**2*c*x**2 - 8*atanh(c*x**2)*b**2 + 8*log(c*x**2 + 1)*b**2 + 3*a**2*c**4*x**8 + 2*a*b*c**3*x**6 + 6*a*b*c*x**2 + b**2*c**2*x**4)/(24*c**4)
```

3.65 $\int x^5(a + b\operatorname{arctanh}(cx^2))^2 dx$

Optimal result	560
Mathematica [A] (verified)	561
Rubi [A] (verified)	561
Maple [B] (verified)	565
Fricas [F]	565
Sympy [F]	566
Maxima [F]	566
Giac [F]	567
Mupad [F(-1)]	567
Reduce [F]	567

Optimal result

Integrand size = 16, antiderivative size = 146

$$\int x^5(a + b\operatorname{arctanh}(cx^2))^2 dx = \frac{b^2x^2}{6c^2} - \frac{b^2\operatorname{arctanh}(cx^2)}{6c^3} + \frac{bx^4(a + b\operatorname{arctanh}(cx^2))}{6c} + \frac{(a + b\operatorname{arctanh}(cx^2))^2}{6c^3} + \frac{1}{6}x^6(a + b\operatorname{arctanh}(cx^2))^2 - \frac{b(a + b\operatorname{arctanh}(cx^2))\log\left(\frac{2}{1-cx^2}\right)}{3c^3} - \frac{b^2\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^2}\right)}{6c^3}$$

output

```
1/6*b^2*x^2/c^2-1/6*b^2*arctanh(c*x^2)/c^3+1/6*b*x^4*(a+b*arctanh(c*x^2))/
c+1/6*(a+b*arctanh(c*x^2))^2/c^3+1/6*x^6*(a+b*arctanh(c*x^2))^2-1/3*b*(a+b
*arctanh(c*x^2))*ln(2/(-c*x^2+1))/c^3-1/6*b^2*polylog(2,1-2/(-c*x^2+1))/c^
3
```

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.90

$$\int x^5 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$= \frac{b^2 cx^2 + abc^2 x^4 + a^2 c^3 x^6 + b^2 (-1 + c^3 x^6) \operatorname{arctanh}(cx^2)^2 + b \operatorname{arctanh}(cx^2) \left(-b + bc^2 x^4 + 2ac^3 x^6 - 2b \log \right)}{6c^3}$$

input `Integrate[x^5*(a + b*ArcTanh[c*x^2])^2,x]`

output `(b^2*c*x^2 + a*b*c^2*x^4 + a^2*c^3*x^6 + b^2*(-1 + c^3*x^6)*ArcTanh[c*x^2]^2 + b*ArcTanh[c*x^2]*(-b + b*c^2*x^4 + 2*a*c^3*x^6 - 2*b*Log[1 + E^(-2*ArcTanh[c*x^2])])) + a*b*Log[-1 + c^2*x^4] + b^2*PolyLog[2, -E^(-2*ArcTanh[c*x^2])])/(6*c^3)`

Rubi [A] (verified)

Time = 0.96 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.11, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {6454, 6452, 6542, 6452, 262, 219, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^5 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$\downarrow 6454$$

$$\frac{1}{2} \int x^4 (a + b \operatorname{arctanh}(cx^2))^2 dx^2$$

$$\downarrow 6452$$

$$\frac{1}{2} \left(\frac{1}{3} x^6 (a + b \operatorname{arctanh}(cx^2))^2 - \frac{2}{3} bc \int \frac{x^6 (a + b \operatorname{arctanh}(cx^2))}{1 - c^2 x^4} dx^2 \right)$$

$$\downarrow 6542$$

$$\frac{1}{2} \left(\frac{1}{3} x^6 (a + \operatorname{barctanh}(cx^2))^2 - \frac{2}{3} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(cx^2))}{1 - c^2 x^4} dx^2}{c^2} - \frac{\int x^2 (a + \operatorname{barctanh}(cx^2)) dx^2}{c^2} \right) \right)$$

↓ 6452

$$\frac{1}{2} \left(\frac{1}{3} x^6 (a + \operatorname{barctanh}(cx^2))^2 - \frac{2}{3} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(cx^2))}{1 - c^2 x^4} dx^2}{c^2} - \frac{\frac{1}{2} x^4 (a + \operatorname{barctanh}(cx^2)) - \frac{1}{2} bc \int \frac{x^4}{1 - c^2 x^4} dx^2}{c^2} \right) \right)$$

↓ 262

$$\frac{1}{2} \left(\frac{1}{3} x^6 (a + \operatorname{barctanh}(cx^2))^2 - \frac{2}{3} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(cx^2))}{1 - c^2 x^4} dx^2}{c^2} - \frac{\frac{1}{2} x^4 (a + \operatorname{barctanh}(cx^2)) - \frac{1}{2} bc \left(\frac{\int \frac{1}{1 - c^2 x^4} dx^2}{c^2} \right)}{c^2} \right) \right)$$

↓ 219

$$\frac{1}{2} \left(\frac{1}{3} x^6 (a + \operatorname{barctanh}(cx^2))^2 - \frac{2}{3} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(cx^2))}{1 - c^2 x^4} dx^2}{c^2} - \frac{\frac{1}{2} x^4 (a + \operatorname{barctanh}(cx^2)) - \frac{1}{2} bc \left(\frac{\operatorname{arctanh}(cx^2)}{c^3} \right)}{c^2} \right) \right)$$

↓ 6546

$$\frac{1}{2} \left(\frac{1}{3} x^6 (a + \operatorname{barctanh}(cx^2))^2 - \frac{2}{3} bc \left(\frac{\int \frac{a + \operatorname{barctanh}(cx^2)}{1 - cx^2} dx^2}{c} - \frac{(a + \operatorname{barctanh}(cx^2))^2}{2bc^2} - \frac{\frac{1}{2} x^4 (a + \operatorname{barctanh}(cx^2)) - \frac{1}{2} bc \int \frac{1}{1 - c^2 x^4} dx^2}{c^2} \right) \right)$$

↓ 6470

$$\frac{1}{2} \left(\frac{1}{3} x^6 (a + \operatorname{barctanh}(cx^2))^2 - \frac{2}{3} bc \left(\frac{\frac{\log\left(\frac{2}{1 - cx^2}\right) (a + \operatorname{barctanh}(cx^2))}{c} - b \int \frac{\log\left(\frac{2}{1 - cx^2}\right)}{1 - c^2 x^4} dx^2}{c} - \frac{(a + \operatorname{barctanh}(cx^2))^2}{2bc^2} - \frac{\frac{1}{2} x^4 (a + \operatorname{barctanh}(cx^2)) - \frac{1}{2} bc \int \frac{1}{1 - c^2 x^4} dx^2}{c^2} \right) \right)$$

↓ 2849

$$\frac{1}{2} \left(\frac{1}{3} x^6 (a + \operatorname{arctanh}(cx^2))^2 - \frac{2}{3} bc \left(\frac{\int \frac{\log\left(\frac{2}{1-cx^2}\right) d\frac{1}{1-cx^2}}{\frac{1}{c}} + \frac{\log\left(\frac{2}{1-cx^2}\right) (a + b \operatorname{arctanh}(cx^2))}{c}}{c^2} - \frac{(a + b \operatorname{arctanh}(cx^2))^2}{2bc^2} \right) \right)$$

↓ 2752

$$\frac{1}{2} \left(\frac{1}{3} x^6 (a + \operatorname{arctanh}(cx^2))^2 - \frac{2}{3} bc \left(\frac{\frac{\log\left(\frac{2}{1-cx^2}\right) (a + b \operatorname{arctanh}(cx^2))}{c} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^2}\right)}{2c}}{c^2} - \frac{(a + b \operatorname{arctanh}(cx^2))^2}{2bc^2} \right) \right)$$

input `Int[x^5*(a + b*ArcTanh[c*x^2])^2,x]`

output `((x^6*(a + b*ArcTanh[c*x^2])^2)/3 - (2*b*c*(-((x^4*(a + b*ArcTanh[c*x^2]))/2 - (b*c*(-(x^2/c^2) + ArcTanh[c*x^2]/c^3))/2)/c^2) + (-1/2*(a + b*ArcTanh[c*x^2])^2/(b*c^2) + (((a + b*ArcTanh[c*x^2])*Log[2/(1 - c*x^2)])/c + (b*PolyLog[2, 1 - 2/(1 - c*x^2)]/(2*c))/c)/c^2))/3)/2`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 262 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*(m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e.)*(x_))]/((f_) + (g.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 6452 `Int[((a_.) + ArcTanh[(c.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_.) + ArcTanh[(c.)*(x_)^(n)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6470 `Int[((a_.) + ArcTanh[(c.)*(x_)]*(b_.))^(p_.)/((d_) + (e.)*(x_)), x_Symbol] := Simp[(-a + b*ArcTanh[c*x])^p*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6542 `Int[((a_.) + ArcTanh[(c.)*(x_)]*(b_.))^(p_.)*((f.)*(x_)^(m_))/((d_) + (e.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

rule 6546 `Int[((a_.) + ArcTanh[(c.)*(x_)]*(b_.))^(p_.)*(x_)/((d_) + (e.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 379 vs. $2(132) = 264$.

Time = 1.01 (sec) , antiderivative size = 380, normalized size of antiderivative = 2.60

method	result
risch	$\frac{x^6 a^2}{6} + \frac{abx^4}{6c} - \frac{b^2 \operatorname{dilog}\left(\frac{cx^2}{2} + \frac{1}{2}\right)}{6c^3} - \frac{2b^2 \ln(cx^2-1)}{9c^3} + \frac{bax^6 \ln(cx^2+1)}{6} + \frac{ba \ln(cx^2+1)}{6c^3} - \frac{b^2 \ln(-cx^2+1) \ln(cx^2+1)x}{12}$
default	Expression too large to display
parts	Expression too large to display

input `int(x^5*(a+b*arctanh(c*x^2))^2,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/6*x^6*a^2+1/6/c*a*b*x^4-1/6*b^2/c^3*\operatorname{dilog}(1/2*c*x^2+1/2)-2/9*b^2/c^3*\ln(c*x^2-1) \\ & +1/6*b*a*x^6*\ln(c*x^2+1)+1/6*b*a/c^3*\ln(c*x^2+1)-1/12*b^2*\ln(-c*x^2+1)*\ln(c*x^2+1) \\ & *x^6-1/12*b^2/c^3*\ln(-c*x^2+1)*\ln(c*x^2+1)+1/6*b^2/c^3*\ln(1/2-1/2*c*x^2)*\ln(c*x^2+1) \\ & -1/6*b^2/c^3*\ln(1/2-1/2*c*x^2)*\ln(1/2*c*x^2+1/2)+1/12*b^2/c*x^4*\ln(c*x^2+1) \\ & -17/108/c^3*b^2-1/12*b^2/c*x^4*\ln(-c*x^2+1)-1/6*a*b*x^6*\ln(-c*x^2+1) \\ & +1/6*a*b/c^3*\ln(c*x^2-1)+1/24*b^2*x^6*\ln(-c*x^2+1)^2+11/36/c^3*b^2*\ln(-c*x^2+1) \\ & -1/24/c^3*b^2*\ln(-c*x^2+1)^2+1/24*b^2*x^6*\ln(c*x^2+1)^2-1/12*b^2/c^3*\ln(c*x^2+1) \\ & +1/24/c^3*b^2*\ln(c*x^2+1)^2+1/6*b^2*x^2/c^2 \end{aligned}$$
Fricas [F]

$$\int x^5 (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{arctanh}(cx^2) + a)^2 x^5 dx$$

input `integrate(x^5*(a+b*arctanh(c*x^2))^2,x, algorithm="fricas")`

output `integral(b^2*x^5*arctanh(c*x^2)^2 + 2*a*b*x^5*arctanh(c*x^2) + a^2*x^5, x)`

Sympy [F]

$$\int x^5 (a + b \operatorname{arctanh}(cx^2))^2 dx = \int x^5 (a + b \operatorname{atanh}(cx^2))^2 dx$$

input `integrate(x**5*(a+b*atanh(c*x**2))**2,x)`

output `Integral(x**5*(a + b*atanh(c*x**2))**2, x)`

Maxima [F]

$$\int x^5 (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 x^5 dx$$

input `integrate(x^5*(a+b*arctanh(c*x^2))^2,x, algorithm="maxima")`

output `1/6*a^2*x^6 + 1/6*(2*x^6*arctanh(c*x^2) + (x^4/c^2 + log(c^2*x^4 - 1)/c^4)
*c)*a*b + 1/432*(18*x^6*log(-c*x^2 + 1)^2 - 2*c^4*(2*(c^2*x^6 + 3*x^2)/c^6
- 3*log(c*x^2 + 1)/c^7 + 3*log(c*x^2 - 1)/c^7) + 3*c^3*(x^4/c^4 + log(c^2
*x^4 - 1)/c^6) + 1296*c^3*integrate(1/9*x^7*log(c*x^2 + 1)/(c^4*x^4 - c^2)
, x) - 9*c^2*(2*x^2/c^4 - log(c*x^2 + 1)/c^5 + log(c*x^2 - 1)/c^5) - 6*c*(
(2*c^2*x^6 + 3*c*x^4 + 6*x^2)/c^3 + 6*log(c*x^2 - 1)/c^4)*log(-c*x^2 + 1)
+ 648*c*integrate(1/9*x^3*log(c*x^2 + 1)/(c^4*x^4 - c^2), x) + 6*(3*c^3*x^
6*log(c*x^2 + 1)^2 + (2*c^3*x^6 - 3*c^2*x^4 + 6*c*x^2 - 6*(c^3*x^6 + 1)*lo
g(c*x^2 + 1))*log(-c*x^2 + 1))/c^3 + (4*c^3*x^6 + 15*c^2*x^4 + 66*c*x^2 +
18*log(c*x^2 - 1)^2 + 66*log(c*x^2 - 1))/c^3 - 18*log(9*c^4*x^4 - 9*c^2)/c
^3 + 648*integrate(1/9*x*log(c*x^2 + 1)/(c^4*x^4 - c^2), x))*b^2`

Giac [F]

$$\int x^5 (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 x^5 dx$$

input `integrate(x^5*(a+b*arctanh(c*x^2))^2,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)^2*x^5, x)`

Mupad [F(-1)]

Timed out.

$$\int x^5 (a + b \operatorname{arctanh}(cx^2))^2 dx = \int x^5 (a + b \operatorname{atanh}(cx^2))^2 dx$$

input `int(x^5*(a + b*atanh(c*x^2))^2,x)`

output `int(x^5*(a + b*atanh(c*x^2))^2, x)`

Reduce [F]

$$\int x^5 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$= \frac{\operatorname{atanh}(cx^2)^2 b^2 c^3 x^6 - \operatorname{atanh}(cx^2)^2 b^2 c x^2 + 2 \operatorname{atanh}(cx^2) a b c^3 x^6 - 2 \operatorname{atanh}(cx^2) a b + \operatorname{atanh}(cx^2) b^2 c^2 x^4}{6c^3}$$

input `int(x^5*(a+b*atanh(c*x^2))^2,x)`

output `(atanh(c*x**2)**2*b**2*c**3*x**6 - atanh(c*x**2)**2*b**2*c*x**2 + 2*atanh(c*x**2)*a*b*c**3*x**6 - 2*atanh(c*x**2)*a*b + atanh(c*x**2)*b**2*c**2*x**4 - atanh(c*x**2)*b**2 + 2*int(atanh(c*x**2)**2*x,x)*b**2*c + 2*log(c*x**2 + 1)*a*b + a**2*c**3*x**6 + a*b*c**2*x**4 + b**2*c*x**2)/(6*c**3)`

3.66 $\int x^3(a + \operatorname{barctanh}(cx^2))^2 dx$

Optimal result	568
Mathematica [A] (verified)	568
Rubi [A] (verified)	569
Maple [A] (verified)	571
Fricas [A] (verification not implemented)	571
Sympy [B] (verification not implemented)	572
Maxima [B] (verification not implemented)	572
Giac [B] (verification not implemented)	573
Mupad [B] (verification not implemented)	574
Reduce [B] (verification not implemented)	575

Optimal result

Integrand size = 16, antiderivative size = 91

$$\int x^3(a + \operatorname{barctanh}(cx^2))^2 dx = \frac{abx^2}{2c} + \frac{b^2x^2\operatorname{arctanh}(cx^2)}{2c} - \frac{(a + \operatorname{barctanh}(cx^2))^2}{4c^2} + \frac{1}{4}x^4(a + \operatorname{barctanh}(cx^2))^2 + \frac{b^2 \log(1 - c^2x^4)}{4c^2}$$

output

$$\frac{1}{2}a*b*x^2/c + \frac{1}{2}*b^2*x^2*\operatorname{arctanh}(c*x^2)/c - \frac{1}{4}*(a+b*\operatorname{arctanh}(c*x^2))^2/c^2 + \frac{1}{4}*x^4*(a+b*\operatorname{arctanh}(c*x^2))^2 + \frac{1}{4}*b^2*\ln(-c^2*x^4+1)/c^2$$

Mathematica [A] (verified)

Time = 0.04 (sec), antiderivative size = 106, normalized size of antiderivative = 1.16

$$\int x^3(a + \operatorname{barctanh}(cx^2))^2 dx = \frac{2abcx^2 + a^2c^2x^4 + 2bcx^2(b + acx^2)\operatorname{arctanh}(cx^2) + b^2(-1 + c^2x^4)\operatorname{arctanh}(cx^2)^2 + b(a + b)\log(1 - cx^2)}{4c^2}$$

input

```
Integrate[x^3*(a + b*ArcTanh[c*x^2])^2, x]
```

output

$$(2*a*b*c*x^2 + a^2*c^2*x^4 + 2*b*c*x^2*(b + a*c*x^2)*\text{ArcTanh}[c*x^2] + b^2*(-1 + c^2*x^4)*\text{ArcTanh}[c*x^2]^2 + b*(a + b)*\text{Log}[1 - c*x^2] - a*b*\text{Log}[1 + c*x^2] + b^2*\text{Log}[1 + c*x^2])/(4*c^2)$$
Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.01, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6454, 6452, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$\downarrow 6454$$

$$\frac{1}{2} \int x^2 (a + b \operatorname{arctanh}(cx^2))^2 dx^2$$

$$\downarrow 6452$$

$$\frac{1}{2} \left(\frac{1}{2} x^4 (a + b \operatorname{arctanh}(cx^2))^2 - bc \int \frac{x^4 (a + b \operatorname{arctanh}(cx^2))}{1 - c^2 x^4} dx^2 \right)$$

$$\downarrow 6542$$

$$\frac{1}{2} \left(\frac{1}{2} x^4 (a + b \operatorname{arctanh}(cx^2))^2 - bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(cx^2)}{1 - c^2 x^4} dx^2}{c^2} - \frac{\int (a + b \operatorname{arctanh}(cx^2)) dx^2}{c^2} \right) \right)$$

$$\downarrow 2009$$

$$\frac{1}{2} \left(\frac{1}{2} x^4 (a + b \operatorname{arctanh}(cx^2))^2 - bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(cx^2)}{1 - c^2 x^4} dx^2}{c^2} - \frac{ax^2 + bx^2 \operatorname{arctanh}(cx^2) + \frac{b \log(1 - c^2 x^4)}{2c}}{c^2} \right) \right)$$

$$\downarrow 6510$$

$$\frac{1}{2} \left(\frac{1}{2} x^4 (a + b \operatorname{arctanh}(cx^2))^2 - bc \left(\frac{(a + b \operatorname{arctanh}(cx^2))^2}{2bc^3} - \frac{ax^2 + bx^2 \operatorname{arctanh}(cx^2) + \frac{b \log(1 - c^2 x^4)}{2c}}{c^2} \right) \right)$$

input `Int[x^3*(a + b*ArcTanh[c*x^2])^2,x]`

output `((x^4*(a + b*ArcTanh[c*x^2])^2)/2 - b*c*((a + b*ArcTanh[c*x^2])^2/(2*b*c^3) - (a*x^2 + b*x^2*ArcTanh[c*x^2] + (b*Log[1 - c^2*x^4])/(2*c))/c^2))/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6510 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6542 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((f_.)*(x_)^(m_.))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.33

method	result
parallelrisch	$\frac{b^2 \operatorname{arctanh}(cx^2)^2 x^4 c^2 + 2x^4 \operatorname{arctanh}(cx^2) abc^2 + a^2 c^2 x^4 + 2b^2 \operatorname{arctanh}(cx^2) x^2 c + 2abcx^2 - b^2 \operatorname{arctanh}(cx^2)^2 + 2 \ln(cx^2 - 1) b^2}{4c^2}$
risch	$\frac{b^2(c^2x^4-1)\ln(cx^2+1)^2}{16c^2} + \frac{b(-2bx^4\ln(-cx^2+1)ac^2+4a^2c^2x^4+4abcx^2+2b\ln(-cx^2+1)a+b^2)\ln(cx^2+1)}{16ac^2} + \frac{b^2x^4\ln(-1)}{1}$
default	Expression too large to display
parts	Expression too large to display

input `int(x^3*(a+b*arctanh(c*x^2))^2,x,method=_RETURNVERBOSE)`

output $\frac{1}{4}*(b^2*\operatorname{arctanh}(c*x^2)^2*x^4*c^2+2*x^4*\operatorname{arctanh}(c*x^2)*a*b*c^2+a^2*c^2*x^4+2*b^2*\operatorname{arctanh}(c*x^2)*x^2*c+2*a*b*c*x^2-b^2*\operatorname{arctanh}(c*x^2)^2+2*\ln(c*x^2-1)*b^2-2*\operatorname{arctanh}(c*x^2)*a*b+2*\operatorname{arctanh}(c*x^2)*b^2)/c^2$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.52

$$\int x^3 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$= \frac{4a^2c^2x^4 + 8abcx^2 + (b^2c^2x^4 - b^2) \log\left(-\frac{cx^2+1}{cx^2-1}\right)^2 - 4(ab - b^2) \log(cx^2 + 1) + 4(ab + b^2) \log(cx^2 - 1)}{16c^2}$$

input `integrate(x^3*(a+b*arctanh(c*x^2))^2,x, algorithm="fricas")`

output $\frac{1}{16}*(4*a^2*c^2*x^4 + 8*a*b*c*x^2 + (b^2*c^2*x^4 - b^2)*\log(-(c*x^2 + 1)/(c*x^2 - 1))^2 - 4*(a*b - b^2)*\log(c*x^2 + 1) + 4*(a*b + b^2)*\log(c*x^2 - 1) + 4*(a*b*c^2*x^4 + b^2*c*x^2)*\log(-(c*x^2 + 1)/(c*x^2 - 1)))/c^2$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 163 vs. $2(78) = 156$.

Time = 4.51 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.79

$$\int x^3 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$= \begin{cases} \frac{a^2 x^4}{4} + \frac{abx^4 \operatorname{atanh}(cx^2)}{2} + \frac{abx^2}{2c} - \frac{ab \operatorname{atanh}(cx^2)}{2c^2} + \frac{b^2 x^4 \operatorname{atanh}^2(cx^2)}{4} + \frac{b^2 x^2 \operatorname{atanh}(cx^2)}{2c} + \frac{b^2 \log\left(x - \sqrt{-\frac{1}{c}}\right)}{2c^2} + \frac{b^2 \log\left(x + \sqrt{-\frac{1}{c}}\right)}{2c^2} \\ \frac{a^2 x^4}{4} \end{cases}$$

input `integrate(x**3*(a+b*atanh(c*x**2))**2,x)`

output `Piecewise((a**2*x**4/4 + a*b*x**4*atanh(c*x**2)/2 + a*b*x**2/(2*c) - a*b*atanh(c*x**2)/(2*c**2) + b**2*x**4*atanh(c*x**2)**2/4 + b**2*x**2*atanh(c*x**2)/(2*c) + b**2*log(x - sqrt(-1/c))/(2*c**2) + b**2*log(x + sqrt(-1/c))/(2*c**2) - b**2*atanh(c*x**2)**2/(4*c**2) - b**2*atanh(c*x**2)/(2*c**2), Ne(c, 0)), (a**2*x**4/4, True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 186 vs. $2(81) = 162$.

Time = 0.03 (sec) , antiderivative size = 186, normalized size of antiderivative = 2.04

$$\int x^3 (a + b \operatorname{arctanh}(cx^2))^2 dx = \frac{1}{4} b^2 x^4 \operatorname{artanh}(cx^2)^2 + \frac{1}{4} a^2 x^4$$

$$+ \frac{1}{4} \left(2x^4 \operatorname{artanh}(cx^2) + c \left(\frac{2x^2}{c^2} - \frac{\log(cx^2 + 1)}{c^3} + \frac{\log(cx^2 - 1)}{c^3} \right) \right) ab$$

$$+ \frac{1}{16} \left(4c \left(\frac{2x^2}{c^2} - \frac{\log(cx^2 + 1)}{c^3} + \frac{\log(cx^2 - 1)}{c^3} \right) \operatorname{artanh}(cx^2) - \frac{2(\log(cx^2 - 1) - 2)\log(cx^2 + 1) - 1}{c^3} \right) b^2$$

input `integrate(x^3*(a+b*arctanh(c*x^2))^2,x, algorithm="maxima")`

output

$$\begin{aligned} & 1/4*b^2*x^4*\operatorname{arctanh}(c*x^2)^2 + 1/4*a^2*x^4 + 1/4*(2*x^4*\operatorname{arctanh}(c*x^2) + c \\ & *(2*x^2/c^2 - \log(c*x^2 + 1)/c^3 + \log(c*x^2 - 1)/c^3))*a*b + 1/16*(4*c*(2 \\ & *x^2/c^2 - \log(c*x^2 + 1)/c^3 + \log(c*x^2 - 1)/c^3)*\operatorname{arctanh}(c*x^2) - (2*(1 \\ & \log(c*x^2 - 1) - 2)*\log(c*x^2 + 1) - \log(c*x^2 + 1)^2 - \log(c*x^2 - 1)^2 - \\ & 4*\log(c*x^2 - 1))/c^2)*b^2 \end{aligned}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 361 vs. $2(81) = 162$.

Time = 0.15 (sec) , antiderivative size = 361, normalized size of antiderivative = 3.97

$$\begin{aligned} & \int x^3(a + b\operatorname{arctanh}(cx^2))^2 dx \\ & = \frac{1}{4} \left(\frac{(cx^2 + 1)b^2 \log\left(-\frac{cx^2+1}{cx^2-1}\right)^2}{\left(\frac{(cx^2+1)^2c^3}{(cx^2-1)^2} - \frac{2(cx^2+1)c^3}{cx^2-1} + c^3\right)(cx^2 - 1)} + \frac{2\left(\frac{2(cx^2+1)ab}{cx^2-1} + \frac{(cx^2+1)b^2}{cx^2-1} - b^2\right) \log\left(-\frac{cx^2+1}{cx^2-1}\right)}{\frac{(cx^2+1)^2c^3}{(cx^2-1)^2} - \frac{2(cx^2+1)c^3}{cx^2-1} + c^3} + \frac{4\left(\frac{(cx^2+1)a}{cx^2-1}\right)}{\frac{(cx^2+1)^2c^3}{(cx^2-1)^2}} \right) \end{aligned}$$

input

```
integrate(x^3*(a+b*arctanh(c*x^2))^2,x, algorithm="giac")
```

output

$$\begin{aligned} & 1/4*((c*x^2 + 1)*b^2*\log(-(c*x^2 + 1)/(c*x^2 - 1))^2/(((c*x^2 + 1)^2*c^3/(c \\ & *x^2 - 1)^2 - 2*(c*x^2 + 1)*c^3/(c*x^2 - 1) + c^3)*(c*x^2 - 1)) + 2*(2*(c \\ & *x^2 + 1)*a*b/(c*x^2 - 1) + (c*x^2 + 1)*b^2/(c*x^2 - 1) - b^2)*\log(-(c*x^2 \\ & + 1)/(c*x^2 - 1))/((c*x^2 + 1)^2*c^3/(c*x^2 - 1)^2 - 2*(c*x^2 + 1)*c^3/(c \\ & *x^2 - 1) + c^3) + 4*((c*x^2 + 1)*a^2/(c*x^2 - 1) + (c*x^2 + 1)*a*b/(c*x^2 \\ & - 1) - a*b)/((c*x^2 + 1)^2*c^3/(c*x^2 - 1)^2 - 2*(c*x^2 + 1)*c^3/(c*x^2 - \\ & 1) + c^3) - 2*b^2*\log(-(c*x^2 + 1)/(c*x^2 - 1) + 1)/c^3 + 2*b^2*\log(-(c*x \\ & ^2 + 1)/(c*x^2 - 1))/c^3)*c \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 4.15 (sec) , antiderivative size = 275, normalized size of antiderivative = 3.02

$$\begin{aligned}
\int x^3(a + b \operatorname{arctanh}(cx^2))^2 dx &= \frac{a^2 x^4}{4} + \frac{b^2 \ln(cx^2 - 1)}{4c^2} + \frac{b^2 \ln(cx^2 + 1)}{4c^2} \\
&\quad - \frac{b^2 \ln(cx^2 + 1)^2}{16c^2} - \frac{b^2 \ln(1 - cx^2)^2}{16c^2} \\
&\quad + \frac{b^2 x^4 \ln(cx^2 + 1)^2}{16} + \frac{b^2 x^4 \ln(1 - cx^2)^2}{16} \\
&\quad + \frac{b^2 x^2 \ln(cx^2 + 1)}{4c} - \frac{b^2 x^2 \ln(1 - cx^2)}{4c} \\
&\quad + \frac{ab \ln(cx^2 - 1)}{4c^2} - \frac{ab \ln(cx^2 + 1)}{4c^2} \\
&\quad + \frac{abx^4 \ln(cx^2 + 1)}{4} - \frac{abx^4 \ln(1 - cx^2)}{4} \\
&\quad + \frac{b^2 \ln(cx^2 + 1) \ln(1 - cx^2)}{8c^2} + \frac{abx^2}{2c} \\
&\quad - \frac{b^2 x^4 \ln(cx^2 + 1) \ln(1 - cx^2)}{8}
\end{aligned}$$

input

```
int(x^3*(a + b*atanh(c*x^2))^2,x)
```

output

```
(a^2*x^4)/4 + (b^2*log(c*x^2 - 1))/(4*c^2) + (b^2*log(c*x^2 + 1))/(4*c^2)
- (b^2*log(c*x^2 + 1)^2)/(16*c^2) - (b^2*log(1 - c*x^2)^2)/(16*c^2) + (b^2
*x^4*log(c*x^2 + 1)^2)/16 + (b^2*x^4*log(1 - c*x^2)^2)/16 + (b^2*x^2*log(c
*x^2 + 1))/(4*c) - (b^2*x^2*log(1 - c*x^2))/(4*c) + (a*b*log(c*x^2 - 1))/(
4*c^2) - (a*b*log(c*x^2 + 1))/(4*c^2) + (a*b*x^4*log(c*x^2 + 1))/4 - (a*b*
x^4*log(1 - c*x^2))/4 + (b^2*log(c*x^2 + 1)*log(1 - c*x^2))/(8*c^2) + (a*b
*x^2)/(2*c) - (b^2*x^4*log(c*x^2 + 1)*log(1 - c*x^2))/8
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.32

$$\int x^3 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$= \frac{\operatorname{atanh}(cx^2)^2 b^2 c^2 x^4 - \operatorname{atanh}(cx^2)^2 b^2 + 2 \operatorname{atanh}(cx^2) ab c^2 x^4 - 2 \operatorname{atanh}(cx^2) ab + 2 \operatorname{atanh}(cx^2) b^2 c x^2 - 2}{4c^2}$$

input

```
int(x^3*(a+b*atanh(c*x^2))^2,x)
```

output

```
(atanh(c*x**2)**2*b**2*c**2*x**4 - atanh(c*x**2)**2*b**2 + 2*atanh(c*x**2)
*a*b*c**2*x**4 - 2*atanh(c*x**2)*a*b + 2*atanh(c*x**2)*b**2*c*x**2 - 2*ata
nh(c*x**2)*b**2 + 2*log(c*x**2 + 1)*b**2 + a**2*c**2*x**4 + 2*a*b*c*x**2)/
(4*c**2)
```

3.67 $\int x(a + b \operatorname{arctanh}(cx^2))^2 dx$

Optimal result	576
Mathematica [A] (verified)	576
Rubi [A] (verified)	577
Maple [A] (verified)	579
Fricas [F]	580
Sympy [F]	580
Maxima [F]	580
Giac [F]	581
Mupad [F(-1)]	581
Reduce [F]	582

Optimal result

Integrand size = 14, antiderivative size = 94

$$\int x(a + b \operatorname{arctanh}(cx^2))^2 dx = \frac{(a + b \operatorname{arctanh}(cx^2))^2}{2c} + \frac{1}{2}x^2(a + b \operatorname{arctanh}(cx^2))^2 - \frac{b(a + b \operatorname{arctanh}(cx^2)) \log\left(\frac{2}{1-cx^2}\right)}{c} - \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^2}\right)}{2c}$$

```
output 1/2*(a+b*arctanh(c*x^2))^2/c+1/2*x^2*(a+b*arctanh(c*x^2))^2-b*(a+b*arctanh(c*x^2))*ln(2/(-c*x^2+1))/c-1/2*b^2*polylog(2,1-2/(-c*x^2+1))/c
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.05

$$\int x(a + b \operatorname{arctanh}(cx^2))^2 dx = \frac{b^2(-1 + cx^2) \operatorname{arctanh}(cx^2)^2 + 2b \operatorname{arctanh}(cx^2) \left(acx^2 - b \log\left(1 + e^{-2 \operatorname{arctanh}(cx^2)}\right)\right) + a(acx^2 + b \log(1 -$$

```
input Integrate[x*(a + b*ArcTanh[c*x^2])^2,x]
```

output

```
(b^2*(-1 + c*x^2)*ArcTanh[c*x^2]^2 + 2*b*ArcTanh[c*x^2]*(a*c*x^2 - b*Log[1
+ E^(-2*ArcTanh[c*x^2])]) + a*(a*c*x^2 + b*Log[1 - c^2*x^4]) + b^2*PolyLo
g[2, -E^(-2*ArcTanh[c*x^2])])/(2*c)
```

Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.11, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {6454, 6436, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(a + \operatorname{barctanh}(cx^2))^2 dx$$

$$\downarrow 6454$$

$$\frac{1}{2} \int (a + \operatorname{barctanh}(cx^2))^2 dx^2$$

$$\downarrow 6436$$

$$\frac{1}{2} \left(x^2(a + \operatorname{barctanh}(cx^2))^2 - 2bc \int \frac{x^2(a + \operatorname{barctanh}(cx^2))}{1 - c^2x^4} dx^2 \right)$$

$$\downarrow 6546$$

$$\frac{1}{2} \left(x^2(a + \operatorname{barctanh}(cx^2))^2 - 2bc \left(\frac{\int \frac{a + \operatorname{barctanh}(cx^2)}{1 - cx^2} dx^2}{c} - \frac{(a + \operatorname{barctanh}(cx^2))^2}{2bc^2} \right) \right)$$

$$\downarrow 6470$$

$$\frac{1}{2} \left(x^2(a + \operatorname{barctanh}(cx^2))^2 - 2bc \left(\frac{\frac{\log\left(\frac{2}{1 - cx^2}\right)(a + \operatorname{barctanh}(cx^2))}{c}}{c} - b \int \frac{\log\left(\frac{2}{1 - cx^2}\right)}{1 - c^2x^4} dx^2 - \frac{(a + \operatorname{barctanh}(cx^2))^2}{2bc^2} \right) \right)$$

$$\downarrow 2849$$

$$\frac{1}{2} \left(x^2 (a + \operatorname{arctanh}(cx^2))^2 - 2bc \left(\frac{b \int \frac{\log\left(\frac{2}{1-cx^2}\right) d\frac{1}{1-cx^2}}{c} + \frac{\log\left(\frac{2}{1-cx^2}\right) (a + \operatorname{arctanh}(cx^2))}{c}}{c} - \frac{(a + \operatorname{arctanh}(cx^2))^2}{2bc^2} \right) \right)$$

↓ 2752

$$\frac{1}{2} \left(x^2 (a + \operatorname{arctanh}(cx^2))^2 - 2bc \left(\frac{\log\left(\frac{2}{1-cx^2}\right) (a + \operatorname{arctanh}(cx^2))}{c} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^2}\right)}{2c} - \frac{(a + \operatorname{arctanh}(cx^2))^2}{2bc^2} \right) \right)$$

input `Int[x*(a + b*ArcTanh[c*x^2])^2,x]`

output `(x^2*(a + b*ArcTanh[c*x^2])^2 - 2*b*c*(-1/2*(a + b*ArcTanh[c*x^2])^2/(b*c^2) + ((a + b*ArcTanh[c*x^2])*Log[2/(1 - c*x^2)])/c + (b*PolyLog[2, 1 - 2/(1 - c*x^2)])/(2*c))/c)/2`

Defintions of rubi rules used

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 6436 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^p, x_Symbol] := Simp[x*(a + b*ArcTanh[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTanh[c*x^n])^(p-1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`

rule 6454 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x
], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simpl
ify[(m + 1)/n]]`

rule 6470 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_.) + (e_.)*(x_)), x_Symbol
] := Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c
*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^
2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2
, 0]`

rule 6546 `Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*(x_)/((d_.) + (e_.)*(x_)^2),
x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/
(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 1.06 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.43

method	result
derivativeldivides	$\frac{c x^2 a^2 + b^2 \left(\operatorname{arctanh}(c x^2)^2 (c x^2 - 1) + 2 \operatorname{arctanh}(c x^2)^2 - 2 \operatorname{arctanh}(c x^2) \ln \left(1 + \frac{(c x^2 + 1)^2}{-c^2 x^4 + 1} \right) - \operatorname{polylog} \left(2, -\frac{(c x^2 + 1)^2}{-c^2 x^4 + 1} \right) \right)}{2c}$
default	$\frac{c x^2 a^2 + b^2 \left(\operatorname{arctanh}(c x^2)^2 (c x^2 - 1) + 2 \operatorname{arctanh}(c x^2)^2 - 2 \operatorname{arctanh}(c x^2) \ln \left(1 + \frac{(c x^2 + 1)^2}{-c^2 x^4 + 1} \right) - \operatorname{polylog} \left(2, -\frac{(c x^2 + 1)^2}{-c^2 x^4 + 1} \right) \right)}{2c}$
parts	$\frac{a^2 x^2}{2} + \frac{b^2 \left(\operatorname{arctanh}(c x^2)^2 (c x^2 - 1) + 2 \operatorname{arctanh}(c x^2)^2 - 2 \operatorname{arctanh}(c x^2) \ln \left(1 + \frac{(c x^2 + 1)^2}{-c^2 x^4 + 1} \right) - \operatorname{polylog} \left(2, -\frac{(c x^2 + 1)^2}{-c^2 x^4 + 1} \right) \right)}{2c}$
risch	$\frac{b^2 \ln(c x^2 + 1)^2}{8c} + \frac{b a \ln(c x^2 + 1) x^2}{2} + \frac{b a \ln(c x^2 + 1)}{2c} - \frac{b^2 \ln(-c x^2 + 1) \ln(c x^2 + 1) x^2}{4} - \frac{b^2 \ln(-c x^2 + 1) \ln(c x^2 + 1)}{4c}$

input `int(x*(a+b*arctanh(c*x^2))^2,x,method=_RETURNVERBOSE)`

output

```
1/2/c*(c*x^2*a^2+b^2*(arctanh(c*x^2))^2*(c*x^2-1)+2*arctanh(c*x^2)^2-2*arctanh(c*x^2)*ln(1+(c*x^2+1)^2/(-c^2*x^4+1))-polylog(2,-(c*x^2+1)^2/(-c^2*x^4+1)))+2*a*b*c*x^2*arctanh(c*x^2)+a*b*ln(-c^2*x^4+1))
```

Fricas [F]

$$\int x(a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 x dx$$

input

```
integrate(x*(a+b*arctanh(c*x^2))^2,x, algorithm="fricas")
```

output

```
integral(b^2*x*arctanh(c*x^2)^2 + 2*a*b*x*arctanh(c*x^2) + a^2*x, x)
```

Sympy [F]

$$\int x(a + b \operatorname{arctanh}(cx^2))^2 dx = \int x(a + b \operatorname{atanh}(cx^2))^2 dx$$

input

```
integrate(x*(a+b*atanh(c*x**2))**2,x)
```

output

```
Integral(x*(a + b*atanh(c*x**2))**2, x)
```

Maxima [F]

$$\int x(a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 x dx$$

input

```
integrate(x*(a+b*arctanh(c*x^2))^2,x, algorithm="maxima")
```

output

```
1/2*a^2*x^2 + 1/8*(x^2*log(-c*x^2 + 1)^2 - c^2*(2*x^2/c^2 - log(c*x^2 + 1)
/c^3 + log(c*x^2 - 1)/c^3) - 2*c*(x^2/c + log(c*x^2 - 1)/c^2)*log(-c*x^2 +
1) + 12*c*integrate(x^3*log(c*x^2 + 1)/(c^2*x^4 - 1), x) + (c*x^2*log(c*x
^2 + 1)^2 + 2*(c*x^2 - (c*x^2 + 1)*log(c*x^2 + 1))*log(-c*x^2 + 1))/c + (2
*c*x^2 + log(c*x^2 - 1)^2 + 2*log(c*x^2 - 1))/c - log(c^2*x^4 - 1)/c + 4*i
ntegrate(x*log(c*x^2 + 1)/(c^2*x^4 - 1), x))*b^2 + 1/2*(2*c*x^2*arctanh(c*
x^2) + log(-c^2*x^4 + 1))*a*b/c
```

Giac [F]

$$\int x(a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 x dx$$

input

```
integrate(x*(a+b*arctanh(c*x^2))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^2) + a)^2*x, x)
```

Mupad [F(-1)]

Timed out.

$$\int x(a + b \operatorname{arctanh}(cx^2))^2 dx = \int x(a + b \operatorname{atanh}(cx^2))^2 dx$$

input

```
int(x*(a + b*atanh(c*x^2))^2,x)
```

output

```
int(x*(a + b*atanh(c*x^2))^2, x)
```

Reduce [F]

$$\int x(a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$= \frac{2 \operatorname{atanh}(cx^2) abc x^2 - 2 \operatorname{atanh}(cx^2) ab + 2 \left(\int \operatorname{atanh}(cx^2)^2 x dx \right) b^2 c + 2 \log(cx^2 + 1) ab + a^2 c x^2}{2c}$$

input `int(x*(a+b*atanh(c*x^2))^2,x)`

output `(2*atanh(c*x**2)*a*b*c*x**2 - 2*atanh(c*x**2)*a*b + 2*int(atanh(c*x**2)**2*x,x)*b**2*c + 2*log(c*x**2 + 1)*a*b + a**2*c*x**2)/(2*c)`

$$3.68 \quad \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x} dx$$

Optimal result	583
Mathematica [C] (verified)	584
Rubi [A] (verified)	585
Maple [F]	587
Fricas [F]	587
Sympy [F]	587
Maxima [F]	588
Giac [F]	588
Mupad [F(-1)]	588
Reduce [F]	589

Optimal result

Integrand size = 16, antiderivative size = 137

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x} dx &= (a + b \operatorname{arctanh}(cx^2))^2 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx^2}\right) \\ &\quad - \frac{1}{2}b(a + b \operatorname{arctanh}(cx^2)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx^2}\right) \\ &\quad + \frac{1}{2}b(a + b \operatorname{arctanh}(cx^2)) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - cx^2}\right) \\ &\quad + \frac{1}{4}b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - cx^2}\right) \\ &\quad - \frac{1}{4}b^2 \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 - cx^2}\right) \end{aligned}$$

output

```
-(a+b*arctanh(c*x^2))^2*arctanh(-1+2/(-c*x^2+1))-1/2*b*(a+b*arctanh(c*x^2)
)*polylog(2,1-2/(-c*x^2+1))+1/2*b*(a+b*arctanh(c*x^2))*polylog(2,-1+2/(-c*
x^2+1))+1/4*b^2*polylog(3,1-2/(-c*x^2+1))-1/4*b^2*polylog(3,-1+2/(-c*x^2+1
))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.32

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x} dx = a^2 \log(x) + \frac{1}{2} ab (-\operatorname{PolyLog}(2, -cx^2) + \operatorname{PolyLog}(2, cx^2))$$

$$+ \frac{1}{2} b^2 \left(\frac{i\pi^3}{24} - \frac{2}{3} \operatorname{arctanh}(cx^2)^3 \right.$$

$$\quad - \operatorname{arctanh}(cx^2)^2 \log(1 + e^{-2\operatorname{arctanh}(cx^2)})$$

$$\quad + \operatorname{arctanh}(cx^2)^2 \log(1 - e^{2\operatorname{arctanh}(cx^2)})$$

$$+ \operatorname{arctanh}(cx^2) \operatorname{PolyLog}(2, -e^{-2\operatorname{arctanh}(cx^2)})$$

$$+ \operatorname{arctanh}(cx^2) \operatorname{PolyLog}(2, e^{2\operatorname{arctanh}(cx^2)})$$

$$\quad + \frac{1}{2} \operatorname{PolyLog}(3, -e^{-2\operatorname{arctanh}(cx^2)})$$

$$\quad \left. - \frac{1}{2} \operatorname{PolyLog}(3, e^{2\operatorname{arctanh}(cx^2)}) \right)$$

input `Integrate[(a + b*ArcTanh[c*x^2])^2/x,x]`

output `a^2*Log[x] + (a*b*(-PolyLog[2, -(c*x^2)] + PolyLog[2, c*x^2]))/2 + (b^2*((I/24)*Pi^3 - (2*ArcTanh[c*x^2]^3)/3 - ArcTanh[c*x^2]^2*Log[1 + E^(-2*ArcTanh[c*x^2])] + ArcTanh[c*x^2]^2*Log[1 - E^(2*ArcTanh[c*x^2])] + ArcTanh[c*x^2]*PolyLog[2, -E^(-2*ArcTanh[c*x^2])] + ArcTanh[c*x^2]*PolyLog[2, E^(2*ArcTanh[c*x^2])] + PolyLog[3, -E^(-2*ArcTanh[c*x^2])]/2 - PolyLog[3, E^(2*ArcTanh[c*x^2])]/2))/2`

Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.19, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6450, 6448, 6614, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x} dx$$

$$\downarrow 6450$$

$$\frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^2} dx^2$$

$$\downarrow 6448$$

$$\frac{1}{2} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^2} \right) (a + b \operatorname{arctanh}(cx^2))^2 - 4bc \int \frac{(a + b \operatorname{arctanh}(cx^2)) \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^2} \right)}{1 - c^2 x^4} dx^2 \right)$$

$$\downarrow 6614$$

$$\frac{1}{2} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^2} \right) (a + b \operatorname{arctanh}(cx^2))^2 - 4bc \left(\frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(cx^2)) \log \left(2 - \frac{2}{1 - cx^2} \right)}{1 - c^2 x^4} dx^2 - \frac{1}{2} \int \right) \right)$$

$$\downarrow 6620$$

$$\frac{1}{2} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^2} \right) (a + b \operatorname{arctanh}(cx^2))^2 - 4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - cx^2} \right) (a + b \operatorname{arctanh}(cx^2))}{2c} - \frac{1}{2} \right) \right) \right)$$

$$\downarrow 7164$$

$$\frac{1}{2} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^2} \right) (a + b \operatorname{arctanh}(cx^2))^2 - 4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - cx^2} \right) (a + b \operatorname{arctanh}(cx^2))}{2c} - \frac{1}{2} \right) \right) \right)$$

input

```
Int[(a + b*ArcTanh[c*x^2])^2/x, x]
```

output

$$(2*(a + b*\text{ArcTanh}[c*x^2])^2*\text{ArcTanh}[1 - 2/(1 - c*x^2)] - 4*b*c*((a + b*\text{ArcTanh}[c*x^2])*PolyLog[2, 1 - 2/(1 - c*x^2)]/(2*c) - (b*PolyLog[3, 1 - 2/(1 - c*x^2)]/(4*c))/2 + (-1/2*(a + b*\text{ArcTanh}[c*x^2])*PolyLog[2, -1 + 2/(1 - c*x^2)]/c + (b*PolyLog[3, -1 + 2/(1 - c*x^2)]/(4*c))/2))/2$$

Defintions of rubi rules used

rule 6448

$$\text{Int}[(a + \text{ArcTanh}[c*x])^p/(x), x_Symbol] \rightarrow \text{Simp}[2*(a + b*\text{ArcTanh}[c*x])^p*\text{ArcTanh}[1 - 2/(1 - c*x)], x] - \text{Simp}[2*b*c*p \text{ Int}[(a + b*\text{ArcTanh}[c*x])^{p-1}*(\text{ArcTanh}[1 - 2/(1 - c*x)]/(1 - c^2*x^2)), x], x] /;$$

$$\text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{IGtQ}[p, 1]$$

rule 6450

$$\text{Int}[(a + \text{ArcTanh}[c*x]^n)^p/(x), x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[(a + b*\text{ArcTanh}[c*x])^p/x, x], x, x^n], x] /;$$

$$\text{FreeQ}[\{a, b, c, n\}, x] \ \&\& \ \text{IGtQ}[p, 0]$$

rule 6614

$$\text{Int}[(\text{ArcTanh}[u]*(a + \text{ArcTanh}[c*x])^p)/((d + e*x^2), x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Int}[\text{Log}[1 + u]*(a + b*\text{ArcTanh}[c*x])^p/(d + e*x^2), x], x] - \text{Simp}[1/2 \text{ Int}[\text{Log}[1 - u]*(a + b*\text{ArcTanh}[c*x])^p/(d + e*x^2), x], x] /;$$

$$\text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{EqQ}[u^2 - (1 - 2/(1 - c*x))^2, 0]$$

rule 6620

$$\text{Int}[(\text{Log}[u]*(a + \text{ArcTanh}[c*x])^p)/((d + e*x^2), x_Symbol] \rightarrow \text{Simp}[(-a + b*\text{ArcTanh}[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)), x] + \text{Simp}[b*(p/2 \text{ Int}[(a + b*\text{ArcTanh}[c*x])^{p-1}*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /;$$

$$\text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{EqQ}[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]$$

rule 7164

$$\text{Int}[u*PolyLog[n, v], x_Symbol] \rightarrow \text{With}[\{w = \text{DerivativeDivides}[v, u*v, x]\}, \text{Simp}[w*PolyLog[n + 1, v], x] /;$$

$$\text{!FalseQ}[w] /;$$

$$\text{FreeQ}[n, x]$$

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x} dx$$

input `int((a+b*arctanh(c*x^2))^2/x,x)`

output `int((a+b*arctanh(c*x^2))^2/x,x)`

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/x,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x^2)^2 + 2*a*b*arctanh(c*x^2) + a^2)/x, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{x} dx$$

input `integrate((a+b*atanh(c*x**2))**2/x,x)`

output `Integral((a + b*atanh(c*x**2))**2/x, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/x,x, algorithm="maxima")`

output `a^2*log(x) + integrate(1/4*b^2*(log(c*x^2 + 1) - log(-c*x^2 + 1))^2/x + a*b*(log(c*x^2 + 1) - log(-c*x^2 + 1))/x, x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/x,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)^2/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{x} dx$$

input `int((a + b*atanh(c*x^2))^2/x,x)`

output `int((a + b*atanh(c*x^2))^2/x, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x} dx = 2 \left(\int \frac{\operatorname{atanh}(cx^2)}{x} dx \right) ab + \left(\int \frac{\operatorname{atanh}(cx^2)^2}{x} dx \right) b^2 + \log(x) a^2$$

input `int((a+b*atanh(c*x^2))^2/x,x)`

output `2*int(atanh(c*x**2)/x,x)*a*b + int(atanh(c*x**2)**2/x,x)*b**2 + log(x)*a**2`

3.69
$$\int \frac{(a+b\operatorname{arctanh}(cx^2))^2}{x^3} dx$$

Optimal result	590
Mathematica [A] (verified)	591
Rubi [A] (verified)	591
Maple [C] (warning: unable to verify)	593
Fricas [F]	594
Sympy [F]	595
Maxima [F]	595
Giac [F]	595
Mupad [F(-1)]	596
Reduce [F]	596

Optimal result

Integrand size = 16, antiderivative size = 87

$$\int \frac{(a + b\operatorname{arctanh}(cx^2))^2}{x^3} dx = \frac{1}{2}c(a + b\operatorname{arctanh}(cx^2))^2 - \frac{(a + b\operatorname{arctanh}(cx^2))^2}{2x^2} + bc(a + b\operatorname{arctanh}(cx^2)) \log\left(2 - \frac{2}{1 + cx^2}\right) - \frac{1}{2}b^2c \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + cx^2}\right)$$

output `1/2*c*(a+b*arctanh(c*x^2))^2-1/2*(a+b*arctanh(c*x^2))^2/x^2+b*c*(a+b*arctanh(c*x^2))*ln(2-2/(c*x^2+1))-1/2*b^2*c*polylog(2,-1+2/(c*x^2+1))`

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.37

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^3} dx = -\frac{a^2}{2x^2} + abc \left(-\frac{\operatorname{arctanh}(cx^2)}{cx^2} + \log(cx^2) - \frac{1}{2} \log(1 - c^2x^4) \right) \\ + \frac{1}{2} b^2 c \left(\operatorname{arctanh}(cx^2) \left(\operatorname{arctanh}(cx^2) - \frac{\operatorname{arctanh}(cx^2)}{cx^2} \right) \right. \\ \left. + 2 \log(1 - e^{-2 \operatorname{arctanh}(cx^2)}) \right) \\ - \operatorname{PolyLog}\left(2, e^{-2 \operatorname{arctanh}(cx^2)}\right)$$

input `Integrate[(a + b*ArcTanh[c*x^2])^2/x^3, x]`

output `-1/2*a^2/x^2 + a*b*c*(-(ArcTanh[c*x^2]/(c*x^2)) + Log[c*x^2] - Log[1 - c^2*x^4]/2) + (b^2*c*(ArcTanh[c*x^2]*(ArcTanh[c*x^2] - ArcTanh[c*x^2]/(c*x^2)) + 2*Log[1 - E^(-2*ArcTanh[c*x^2])]) - PolyLog[2, E^(-2*ArcTanh[c*x^2])]) /2`

Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6454, 6452, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^3} dx \\ \downarrow 6454 \\ \frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^4} dx^2 \\ \downarrow 6452$$

$$\frac{1}{2} \left(2bc \int \frac{a + \operatorname{arctanh}(cx^2)}{x^2(1-c^2x^4)} dx^2 - \frac{(a + \operatorname{arctanh}(cx^2))^2}{x^2} \right)$$

↓ 6550

$$\frac{1}{2} \left(2bc \left(\int \frac{a + \operatorname{arctanh}(cx^2)}{x^2(cx^2+1)} dx^2 + \frac{(a + \operatorname{arctanh}(cx^2))^2}{2b} \right) - \frac{(a + \operatorname{arctanh}(cx^2))^2}{x^2} \right)$$

↓ 6494

$$\frac{1}{2} \left(2bc \left(-bc \int \frac{\log\left(2 - \frac{2}{cx^2+1}\right)}{1-c^2x^4} dx^2 + \frac{(a + \operatorname{arctanh}(cx^2))^2}{2b} + \log\left(2 - \frac{2}{cx^2+1}\right) (a + \operatorname{arctanh}(cx^2)) \right) - \frac{(a + \operatorname{arctanh}(cx^2))^2}{x^2} \right)$$

↓ 2897

$$\frac{1}{2} \left(2bc \left(\frac{(a + \operatorname{arctanh}(cx^2))^2}{2b} + \log\left(2 - \frac{2}{cx^2+1}\right) (a + \operatorname{arctanh}(cx^2)) - \frac{1}{2} b \operatorname{PolyLog}\left(2, \frac{2}{cx^2+1} - 1\right) \right) - \frac{(a + \operatorname{arctanh}(cx^2))^2}{x^2} \right)$$

input `Int[(a + b*ArcTanh[c*x^2])^2/x^3,x]`

output `(-((a + b*ArcTanh[c*x^2])^2/x^2) + 2*b*c*((a + b*ArcTanh[c*x^2])^2/(2*b) + (a + b*ArcTanh[c*x^2])*Log[2 - 2/(1 + c*x^2)] - (b*PolyLog[2, -1 + 2/(1 + c*x^2)])/2))/2`

Defintions of rubi rules used

rule 2897 `Int[Log[u_]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1-u)/D[u, x])]}, Simp[C*PolyLog[2, 1-u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m+1)*((a + b*ArcTanh[c*x^n])^p/(m+1)), x] - Simp[b*c*n*(p/(m+1)) Int[x^(m+n)*((a + b*ArcTanh[c*x^n])^(p-1)/(1-c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x
, x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simpl
ify[(m + 1)/n]`

rule 6494 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_))), x
_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] -
Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]
/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c
^2*d^2 - e^2, 0]`

rule 6550 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_)^2)),
x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/
d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.86 (sec) , antiderivative size = 820, normalized size of antiderivative = 9.43

method	result	size
default	Expression too large to display	820
parts	Expression too large to display	820

input `int((a+b*arctanh(c*x^2))^2/x^3,x,method=_RETURNVERBOSE)`

output

```
-1/2/x^2*a^2+b^2*(-1/2/x^2*arctanh(c*x^2)^2+2*c*(ln(x)*arctanh(c*x^2)-1/4*
arctanh(c*x^2)*ln(c*x^2+1)-1/4*arctanh(c*x^2)*ln(c*x^2-1)-1/2*c*(Sum(-1/4*
(ln(x-_alpha)*ln(c*x^2-1)-2*c*(1/2*ln(x-_alpha)*(ln((RootOf(_Z^2*c+2*_Z*_a
lpha*c-2,index=1)-x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=1))+ln((Ro
otOf(_Z^2*c+2*_Z*_alpha*c-2,index=2)-x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c
-2,index=2)))/c+1/2*(dilog((RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=1)-x+_alp
ha)/RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=1))+dilog((RootOf(_Z^2*c+2*_Z*_alp
ha*c-2,index=2)-x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=2)))/c))/c,_a
lpha=RootOf(_Z^2*c+1))+Sum(1/4*(ln(x-_alpha)*ln(c*x^2-1)-2*c*(1/4/_alpha/c
*ln(x-_alpha)^2-1/2*_alpha*ln(x-_alpha)*ln(1/2*(x+_alpha)/_alpha)-1/2*_alp
ha*dilog(1/2*(x+_alpha)/_alpha)))/c,_alpha=RootOf(_Z^2*c-1))+Sum(-1/4*(ln(
x-_alpha)*ln(c*x^2+1)-2*c*(1/4/_alpha/c*ln(x-_alpha)^2+1/2*_alpha*ln(x_al
pha)*ln(1/2*(x+_alpha)/_alpha)+1/2*_alpha*dilog(1/2*(x+_alpha)/_alpha)))/c
,_alpha=RootOf(_Z^2*c+1))+Sum(1/4*(ln(x-_alpha)*ln(c*x^2+1)-2*c*(1/2*ln(x-
_alpha)*(ln((RootOf(_Z^2*c+2*_Z*_alpha*c+2,index=1)-x+_alpha)/RootOf(_Z^2*
c+2*_Z*_alpha*c+2,index=1))+ln((RootOf(_Z^2*c+2*_Z*_alpha*c+2,index=2)-x+_
alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c+2,index=2)))/c+1/2*(dilog((RootOf(_Z^2*
c+2*_Z*_alpha*c+2,index=1)-x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c+2,index=1
))+dilog((RootOf(_Z^2*c+2*_Z*_alpha*c+2,index=2)-x+_alpha)/RootOf(_Z^2*c+2
*_Z*_alpha*c+2,index=2)))/c))/c,_alpha=RootOf(_Z^2*c-1))+ln(x)*(ln(1+(-...
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^3} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^2}{x^3} dx$$

input

```
integrate((a+b*arctanh(c*x^2))^2/x^3,x, algorithm="fricas")
```

output

```
integral((b^2*arctanh(c*x^2)^2 + 2*a*b*arctanh(c*x^2) + a^2)/x^3, x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^3} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{x^3} dx$$

input `integrate((a+b*atanh(c*x**2))**2/x**3,x)`

output `Integral((a + b*atanh(c*x**2))**2/x**3, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^3} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{x^3} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/x^3,x, algorithm="maxima")`

output `-1/2*(c*(log(c^2*x^4 - 1) - log(x^4)) + 2*arctanh(c*x^2)/x^2)*a*b - 1/8*b^2*(log(-c*x^2 + 1)^2/x^2 + 2*integrate(-((c*x^2 - 1)*log(c*x^2 + 1)^2 + 2*(c*x^2 - (c*x^2 - 1)*log(c*x^2 + 1))*log(-c*x^2 + 1))/(c*x^5 - x^3), x)) - 1/2*a^2/x^2`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^3} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{x^3} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/x^3,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)^2/x^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^3} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{x^3} dx$$

input `int((a + b*atanh(c*x^2))^2/x^3,x)`output `int((a + b*atanh(c*x^2))^2/x^3, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^3} dx$$

$$= \frac{-\operatorname{atanh}(cx^2)^2 b^2 + 2 \operatorname{atanh}(cx^2) abc x^2 - 2 \operatorname{atanh}(cx^2) ab - 4 \left(\int \frac{\operatorname{atanh}(cx^2)}{c^2 x^5 - x} dx \right) b^2 c x^2 - 2 \log(cx^2 + 1) ab}{2x^2}$$

input `int((a+b*atanh(c*x^2))^2/x^3,x)`output `(- atanh(c*x**2)**2*b**2 + 2*atanh(c*x**2)*a*b*c*x**2 - 2*atanh(c*x**2)*a*b - 4*int(atanh(c*x**2)/(c**2*x**5 - x),x)*b**2*c*x**2 - 2*log(c*x**2 + 1)*a*b*c*x**2 + 4*log(x)*a*b*c*x**2 - a**2)/(2*x**2)`

3.70
$$\int \frac{(a+b\operatorname{arctanh}(cx^2))^2}{x^5} dx$$

Optimal result	597
Mathematica [A] (verified)	597
Rubi [A] (verified)	598
Maple [A] (verified)	601
Fricas [A] (verification not implemented)	601
Sympy [B] (verification not implemented)	602
Maxima [B] (verification not implemented)	603
Giac [F]	603
Mupad [B] (verification not implemented)	604
Reduce [B] (verification not implemented)	604

Optimal result

Integrand size = 16, antiderivative size = 88

$$\int \frac{(a + \operatorname{arctanh}(cx^2))^2}{x^5} dx = -\frac{bc(a + \operatorname{arctanh}(cx^2))}{2x^2} + \frac{1}{4}c^2(a + \operatorname{arctanh}(cx^2))^2 - \frac{(a + \operatorname{arctanh}(cx^2))^2}{4x^4} + b^2c^2 \log(x) - \frac{1}{4}b^2c^2 \log(1 - c^2x^4)$$

output

$$-1/2*b*c*(a+b*\operatorname{arctanh}(c*x^2))/x^2+1/4*c^2*(a+b*\operatorname{arctanh}(c*x^2))^2-1/4*(a+b*\operatorname{arctanh}(c*x^2))^2/x^4+b^2*c^2*\ln(x)-1/4*b^2*c^2*\ln(-c^2*x^4+1)$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.26

$$\int \frac{(a + \operatorname{arctanh}(cx^2))^2}{x^5} dx = \frac{1}{4} \left(-\frac{a^2}{x^4} - \frac{2abc}{x^2} - \frac{2b(a + bcx^2) \operatorname{arctanh}(cx^2)}{x^4} + \frac{b^2(-1 + c^2x^4) \operatorname{arctanh}(cx^2)^2}{x^4} + 4b^2c^2 \log(x) - b(a + b)c^2 \log(1 - cx^2) + (a - b)bc^2 \log(1 + cx^2) \right)$$

input `Integrate[(a + b*ArcTanh[c*x^2])^2/x^5, x]`

output $(-a^2/x^4) - (2*a*b*c)/x^2 - (2*b*(a + b*c*x^2)*ArcTanh[c*x^2])/x^4 + (b^2*(-1 + c^2*x^4)*ArcTanh[c*x^2]^2)/x^4 + 4*b^2*c^2*Log[x] - b*(a + b)*c^2*Log[1 - c*x^2] + (a - b)*b*c^2*Log[1 + c*x^2])/4$

Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.99, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.562$, Rules used = {6454, 6452, 6544, 6452, 243, 47, 14, 16, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^5} dx \\
 & \quad \downarrow \text{6454} \\
 & \frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^6} dx^2 \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{2} \left(bc \int \frac{a + b \operatorname{arctanh}(cx^2)}{x^4(1 - c^2x^4)} dx^2 - \frac{(a + b \operatorname{arctanh}(cx^2))^2}{2x^4} \right) \\
 & \quad \downarrow \text{6544} \\
 & \frac{1}{2} \left(bc \left(c^2 \int \frac{a + b \operatorname{arctanh}(cx^2)}{1 - c^2x^4} dx^2 + \int \frac{a + b \operatorname{arctanh}(cx^2)}{x^4} dx^2 \right) - \frac{(a + b \operatorname{arctanh}(cx^2))^2}{2x^4} \right) \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{2} \left(bc \left(c^2 \int \frac{a + b \operatorname{arctanh}(cx^2)}{1 - c^2x^4} dx^2 + bc \int \frac{1}{x^2(1 - c^2x^4)} dx^2 - \frac{a + b \operatorname{arctanh}(cx^2)}{x^2} \right) - \frac{(a + b \operatorname{arctanh}(cx^2))^2}{2x^4} \right) \\
 & \quad \downarrow \text{243}
 \end{aligned}$$

$$\frac{1}{2} \left(bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^2)}{1 - c^2x^4} dx^2 + \frac{1}{2} bc \int \frac{1}{x^2(1 - c^2x^4)} dx^4 - \frac{a + \operatorname{barctanh}(cx^2)}{x^2} \right) - \frac{(a + \operatorname{barctanh}(cx^2))^2}{2x^4} \right)$$

↓ 47

$$\frac{1}{2} \left(bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^2)}{1 - c^2x^4} dx^2 + \frac{1}{2} bc \left(c^2 \int \frac{1}{1 - c^2x^4} dx^4 + \int \frac{1}{x^2} dx^4 \right) - \frac{a + \operatorname{barctanh}(cx^2)}{x^2} \right) - \frac{(a + \operatorname{barctanh}(cx^2))^2}{2x^4} \right)$$

↓ 14

$$\frac{1}{2} \left(bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^2)}{1 - c^2x^4} dx^2 + \frac{1}{2} bc \left(c^2 \int \frac{1}{1 - c^2x^4} dx^4 + \log(x^4) \right) - \frac{a + \operatorname{barctanh}(cx^2)}{x^2} \right) - \frac{(a + \operatorname{barctanh}(cx^2))^2}{2x^4} \right)$$

↓ 16

$$\frac{1}{2} \left(bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^2)}{1 - c^2x^4} dx^2 - \frac{a + \operatorname{barctanh}(cx^2)}{x^2} + \frac{1}{2} bc (\log(x^4) - \log(1 - c^2x^4)) \right) - \frac{(a + \operatorname{barctanh}(cx^2))^2}{2x^4} \right)$$

↓ 6510

$$\frac{1}{2} \left(bc \left(\frac{c(a + \operatorname{barctanh}(cx^2))^2}{2b} - \frac{a + \operatorname{barctanh}(cx^2)}{x^2} + \frac{1}{2} bc (\log(x^4) - \log(1 - c^2x^4)) \right) - \frac{(a + \operatorname{barctanh}(cx^2))^2}{2x^4} \right)$$

input

```
Int[(a + b*ArcTanh[c*x^2])^2/x^5,x]
```

output

```
(-1/2*(a + b*ArcTanh[c*x^2])^2/x^4 + b*c*(-((a + b*ArcTanh[c*x^2])/x^2) +
(c*(a + b*ArcTanh[c*x^2])^2)/(2*b) + (b*c*(Log[x^4] - Log[1 - c^2*x^4]))/2
)/2
```

Definitions of rubi rules used

- rule 14 $\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ ; FreeQ}[a, x]$
- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$
- rule 47 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x]$
- rule 243 $\text{Int}[(x_)^{(m_)*((a_)+(b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] \text{ ; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$
- rule 6452 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)]*(b_)]^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{2*n})}), x], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$
- rule 6454 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)]*(b_)]^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n] - 1)*(a + b*\text{ArcTanh}[c*x])^p}, x], x, x^n], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$
- rule 6510 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)]*(b_)]^{(p_)/((d_)+(e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^{(p+1)}/(b*c*d*(p+1)), x] \text{ ; FreeQ}[\{a, b, c, d, e, p\}, x] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{NeQ}[p, -1]$

rule 6544

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^ (p_.)*((f_.)*(x_))^(m_)]/((d_) + (
e_.)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x
], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x
^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

Maple [A] (verified)

Time = 0.53 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.73

method	result
parallelrisch	$\frac{b^2 \operatorname{arctanh}(cx^2)^2 x^4 c^2 + 4b^2 c^2 \ln(x) x^4 - 2 \ln(cx^2 - 1) b^2 c^2 x^4 + 2x^4 \operatorname{arctanh}(cx^2) ab c^2 - 2 \operatorname{arctanh}(cx^2) x^4 b^2 c^2 - a^2 c^2 x^4 - 2b^2 a c^2 x^4}{4x^4}$
risch	$\frac{b^2 (c^2 x^4 - 1) \ln(cx^2 + 1)^2}{16x^4} - \frac{b (b c^2 \ln(-cx^2 + 1) x^4 + 2bc x^2 - b \ln(-cx^2 + 1) + 2a) \ln(cx^2 + 1)}{8x^4} + \frac{b^2 c^2 x^4 \ln(-cx^2 + 1)^2 + 4b c^2 x^4 \ln(-cx^2 + 1) + 2a^2 c^2 x^4}{16x^4}$
default	Expression too large to display
parts	Expression too large to display

input

```
int((a+b*arctanh(c*x^2))^2/x^5,x,method=_RETURNVERBOSE)
```

output

```
1/4*(b^2*arctanh(c*x^2)^2*x^4*c^2+4*b^2*c^2*ln(x)*x^4-2*ln(c*x^2-1)*b^2*c^
2*x^4+2*x^4*arctanh(c*x^2)*a*b*c^2-2*arctanh(c*x^2)*x^4*b^2*c^2-a^2*c^2*x^
4-2*b^2*arctanh(c*x^2)*x^2*c-2*a*b*c*x^2-b^2*arctanh(c*x^2)^2-2*arctanh(c*
x^2)*a*b-a^2)/x^4
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.72

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^5} dx = \frac{16 b^2 c^2 x^4 \log(x) + 4 (ab - b^2) c^2 x^4 \log(cx^2 + 1) - 4 (ab + b^2) c^2 x^4 \log(cx^2 - 1) - 8 abcx^2 + (b^2 c^2 x^4 - b^2)}{16 x^4}$$

input

```
integrate((a+b*arctanh(c*x^2))^2/x^5,x, algorithm="fricas")
```

output

```
1/16*(16*b^2*c^2*x^4*log(x) + 4*(a*b - b^2)*c^2*x^4*log(c*x^2 + 1) - 4*(a*
b + b^2)*c^2*x^4*log(c*x^2 - 1) - 8*a*b*c*x^2 + (b^2*c^2*x^4 - b^2)*log(-(
c*x^2 + 1)/(c*x^2 - 1))^2 - 4*a^2 - 4*(b^2*c*x^2 + a*b)*log(-(c*x^2 + 1)/(
c*x^2 - 1)))/x^4
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 175 vs. $2(80) = 160$.

Time = 6.56 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.99

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^5} dx$$

$$= \begin{cases} -\frac{a^2}{4x^4} + \frac{abc^2 \operatorname{atanh}(cx^2)}{2} - \frac{abc}{2x^2} - \frac{ab \operatorname{atanh}(cx^2)}{2x^4} + b^2 c^2 \log(x) - \frac{b^2 c^2 \log\left(x - \sqrt{-\frac{1}{c}}\right)}{2} - \frac{b^2 c^2 \log\left(x + \sqrt{-\frac{1}{c}}\right)}{2} + \frac{b^2 c^2 \operatorname{atanh}^2}{4} \\ -\frac{a^2}{4x^4} \end{cases}$$

input

```
integrate((a+b*atanh(c*x**2))**2/x**5,x)
```

output

```
Piecewise((-a**2/(4*x**4) + a*b*c**2*atanh(c*x**2)/2 - a*b*c/(2*x**2) - a*
b*atanh(c*x**2)/(2*x**4) + b**2*c**2*log(x) - b**2*c**2*log(x - sqrt(-1/c)
)/2 - b**2*c**2*log(x + sqrt(-1/c))/2 + b**2*c**2*atanh(c*x**2)**2/4 + b**
2*c**2*atanh(c*x**2)/2 - b**2*c*atanh(c*x**2)/(2*x**2) - b**2*atanh(c*x**2
)**2/(4*x**4), Ne(c, 0)), (-a**2/(4*x**4), True))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 175 vs. $2(80) = 160$.

Time = 0.03 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.99

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^5} dx$$

$$= \frac{1}{4} \left(\left(c \log(cx^2 + 1) - c \log(cx^2 - 1) - \frac{2}{x^2} \right) c - \frac{2 \operatorname{artanh}(cx^2)}{x^4} \right) ab$$

$$+ \frac{1}{16} \left(\left(2(\log(cx^2 - 1) - 2) \log(cx^2 + 1) - \log(cx^2 + 1)^2 - \log(cx^2 - 1)^2 - 4 \log(cx^2 - 1) + 16 \log(x) \right) c^2 + 4(c \log(cx^2 + 1) - c \log(cx^2 - 1) - 2/x^2) * c * \operatorname{arctanh}(cx^2) \right) * b^2 - \frac{b^2 \operatorname{artanh}(cx^2)^2}{4x^4} - \frac{a^2}{4x^4}$$

input `integrate((a+b*arctanh(c*x^2))^2/x^5,x, algorithm="maxima")`

output `1/4*((c*log(c*x^2 + 1) - c*log(c*x^2 - 1) - 2/x^2)*c - 2*arctanh(c*x^2)/x^4)*a*b + 1/16*((2*(log(c*x^2 - 1) - 2)*log(c*x^2 + 1) - log(c*x^2 + 1)^2 - log(c*x^2 - 1)^2 - 4*log(c*x^2 - 1) + 16*log(x))*c^2 + 4*(c*log(c*x^2 + 1) - c*log(c*x^2 - 1) - 2/x^2)*c*arctanh(c*x^2))*b^2 - 1/4*b^2*arctanh(c*x^2)^2/x^4 - 1/4*a^2/x^4`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^5} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{x^5} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/x^5,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)^2/x^5, x)`

Mupad [B] (verification not implemented)

Time = 3.96 (sec) , antiderivative size = 278, normalized size of antiderivative = 3.16

$$\int \frac{(a + \operatorname{barctanh}(cx^2))^2}{x^5} dx = \frac{b^2 c^2 \ln(cx^2 + 1)^2}{16} - \frac{b^2 c^2 \ln(cx^2 - 1)}{4} - \frac{b^2 c^2 \ln(cx^2 + 1)}{4} - \frac{a^2}{4x^4} + \frac{b^2 c^2 \ln(1 - cx^2)^2}{16} - \frac{b^2 \ln(cx^2 + 1)^2}{16x^4} - \frac{b^2 \ln(1 - cx^2)^2}{16x^4} + b^2 c^2 \ln(x) - \frac{ab c^2 \ln(cx^2 - 1)}{4} + \frac{ab c^2 \ln(cx^2 + 1)}{4} - \frac{abc}{2x^2} - \frac{ab \ln(cx^2 + 1)}{4x^4} + \frac{ab \ln(1 - cx^2)}{4x^4} - \frac{b^2 c^2 \ln(cx^2 + 1) \ln(1 - cx^2)}{8} - \frac{b^2 c \ln(cx^2 + 1)}{4x^2} + \frac{b^2 c \ln(1 - cx^2)}{4x^2} + \frac{b^2 \ln(cx^2 + 1) \ln(1 - cx^2)}{8x^4}$$

input `int((a + b*atanh(c*x^2))^2/x^5,x)`output $(b^2 c^2 \log(cx^2 + 1)^2)/16 - (b^2 c^2 \log(cx^2 - 1))/4 - (b^2 c^2 \log(cx^2 + 1))/4 - a^2/(4x^4) + (b^2 c^2 \log(1 - cx^2)^2)/16 - (b^2 \log(cx^2 + 1)^2)/(16x^4) - (b^2 \log(1 - cx^2)^2)/(16x^4) + b^2 c^2 \log(x) - (abc^2 \log(cx^2 - 1))/4 + (abc^2 \log(cx^2 + 1))/4 - (abc)/(2x^2) - (ab \log(cx^2 + 1))/(4x^4) + (ab \log(1 - cx^2))/(4x^4) - (b^2 c^2 \log(cx^2 + 1) \log(1 - cx^2))/8 - (b^2 c \log(cx^2 + 1))/(4x^2) + (b^2 c \log(1 - cx^2))/(4x^2) + (b^2 \log(cx^2 + 1) \log(1 - cx^2))/(8x^4)$ **Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.59

$$\int \frac{(a + \operatorname{barctanh}(cx^2))^2}{x^5} dx = \frac{\operatorname{atanh}(cx^2)^2 b^2 c^2 x^4 - \operatorname{atanh}(cx^2)^2 b^2 + 2 \operatorname{atanh}(cx^2) ab c^2 x^4 - 2 \operatorname{atanh}(cx^2) ab + 2 \operatorname{atanh}(cx^2) b^2 c^2 x^4 - 2}{4x^4}$$

input `int((a+b*atanh(c*x^2))^2/x^5,x)`

output `(atanh(c*x**2)**2*b**2*c**2*x**4 - atanh(c*x**2)**2*b**2 + 2*atanh(c*x**2)*a*b*c**2*x**4 - 2*atanh(c*x**2)*a*b + 2*atanh(c*x**2)*b**2*c**2*x**4 - 2*atanh(c*x**2)*b**2*c*x**2 - 2*log(c*x**2 + 1)*b**2*c**2*x**4 + 4*log(x)*b**2*c**2*x**4 - a**2 - 2*a*b*c*x**2)/(4*x**4)`

3.71 $\int x^4(a + \operatorname{barctanh}(cx^2))^2 dx$

Optimal result	606
Mathematica [F]	607
Rubi [A] (verified)	608
Maple [F]	610
Fricas [F]	611
Sympy [F]	611
Maxima [F]	611
Giac [F]	612
Mupad [F(-1)]	612
Reduce [F]	613

Optimal result

Integrand size = 16, antiderivative size = 1173

$$\int x^4(a + \operatorname{barctanh}(cx^2))^2 dx = \text{Too large to display}$$

output

```

8/15*b^2*x/c^2-1/10*b^2*polylog(2,1+2*c^(1/2)*(1-(-c)^(1/2)*x)/((-c)^(1/2)
-c^(1/2))/(1+c^(1/2)*x))/c^(5/2)+1/20*b^2*x^5*ln(c*x^2+1)^2+1/25*b^2*x^5*1
n(-c*x^2+1)+1/25*b*x^5*(2*a-b*ln(-c*x^2+1))+1/5*I*b^2*polylog(2,1-2/(1+I*c
^(1/2)*x))/c^(5/2)-2/25*a*b*x^5+2/15*b^2*x^3*ln(c*x^2+1)/c+1/5*a*b*x^5*ln(
c*x^2+1)-1/10*b^2*x^5*ln(-c*x^2+1)*ln(c*x^2+1)-1/15*b^2*x^3*ln(-c*x^2+1)/c
+1/15*b*x^3*(2*a-b*ln(-c*x^2+1))/c+2/15*a*b*x^3/c+1/20*x^5*(2*a-b*ln(-c*x^
2+1))^2+1/5*b^2*polylog(2,1-2/(1+c^(1/2)*x))/c^(5/2)+1/5*b^2*polylog(2,1-2
/(1-c^(1/2)*x))/c^(5/2)-4/15*b^2*arctanh(c^(1/2)*x)/c^(5/2)-1/5*b^2*arctan
h(c^(1/2)*x)^2/c^(5/2)-4/15*b^2*arctan(c^(1/2)*x)/c^(5/2)-1/10*b^2*polylog
(2,1-2*c^(1/2)*(1+(-c)^(1/2)*x)/((-c)^(1/2)+c^(1/2))/(1+c^(1/2)*x))/c^(5/2
)+1/5*I*b^2*polylog(2,1-2/(1-I*c^(1/2)*x))/c^(5/2)+1/5*I*b^2*arctan(c^(1/2
)*x)^2/c^(5/2)-2/5*b^2*arctanh(c^(1/2)*x)*ln(2/(1+c^(1/2)*x))/c^(5/2)+2/5*
b^2*arctanh(c^(1/2)*x)*ln(2/(1-c^(1/2)*x))/c^(5/2)+1/5*b^2*arctanh(c^(1/2)
*x)*ln(2*c^(1/2)*(1+(-c)^(1/2)*x)/((-c)^(1/2)+c^(1/2))/(1+c^(1/2)*x))/c^(5
/2)+1/5*b^2*arctanh(c^(1/2)*x)*ln(-2*c^(1/2)*(1-(-c)^(1/2)*x)/((-c)^(1/2)-
c^(1/2))/(1+c^(1/2)*x))/c^(5/2)+1/5*b^2*arctan(c^(1/2)*x)*ln((1+I)*(1-c^(1
/2)*x)/(1-I*c^(1/2)*x))/c^(5/2)+1/5*b^2*arctan(c^(1/2)*x)*ln((1-I)*(1+c^(1
/2)*x)/(1-I*c^(1/2)*x))/c^(5/2)+2/5*b^2*arctan(c^(1/2)*x)*ln(2/(1+I*c^(1/2
)*x))/c^(5/2)-2/5*b^2*arctan(c^(1/2)*x)*ln(2/(1-I*c^(1/2)*x))/c^(5/2)+2/5*
a*b*arctan(c^(1/2)*x)/c^(5/2)-1/5*b^2*arctanh(c^(1/2)*x)*ln(c*x^2+1)/c^...

```

Mathematica [F]

$$\int x^4(a + b \operatorname{arctanh}(cx^2))^2 dx = \int x^4(a + b \operatorname{arctanh}(cx^2))^2 dx$$

input

```
Integrate[x^4*(a + b*ArcTanh[c*x^2])^2,x]
```

output

```
Integrate[x^4*(a + b*ArcTanh[c*x^2])^2, x]
```

Rubi [A] (verified)

Time = 2.62 (sec) , antiderivative size = 1173, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {6456, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

↓ 6456

$$\int \left(\frac{1}{4} x^4 (2a - b \log(1 - cx^2))^2 - \frac{1}{2} b x^4 \log(cx^2 + 1) (b \log(1 - cx^2) - 2a) + \frac{1}{4} b^2 x^4 \log^2(cx^2 + 1) \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{1}{20}(2a - b \log(1 - cx^2))^2 x^5 + \frac{1}{20}b^2 \log^2(cx^2 + 1) x^5 - \frac{2}{25}abx^5 + \frac{1}{25}b^2 \log(1 - cx^2) x^5 + \\
& \frac{1}{25}b(2a - b \log(1 - cx^2)) x^5 + \frac{1}{5}ab \log(cx^2 + 1) x^5 - \frac{1}{10}b^2 \log(1 - cx^2) \log(cx^2 + 1) x^5 - \\
& \frac{b^2 \log(1 - cx^2) x^3}{15c} + \frac{b(2a - b \log(1 - cx^2)) x^3}{15c} + \frac{2b^2 \log(cx^2 + 1) x^3}{15c^2} + \frac{2abx^3}{15c} + \frac{8b^2 x}{15c^2} + \\
& \frac{ib^2 \arctan(\sqrt{cx})^2}{5c^{5/2}} - \frac{b^2 \operatorname{arctanh}(\sqrt{cx})^2}{5c^{5/2}} - \frac{4b^2 \arctan(\sqrt{cx})}{15c^{5/2}} + \frac{2ab \arctan(\sqrt{cx})}{5c^{5/2}} - \\
& \frac{4b^2 \operatorname{arctanh}(\sqrt{cx})}{15c^{5/2}} + \frac{2b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(\frac{2}{1-\sqrt{cx}}\right)}{5c^{5/2}} - \frac{2b^2 \arctan(\sqrt{cx}) \log\left(\frac{2}{1-i\sqrt{cx}}\right)}{5c^{5/2}} + \\
& \frac{b^2 \arctan(\sqrt{cx}) \log\left(\frac{(1+i)(1-\sqrt{cx})}{1-i\sqrt{cx}}\right)}{5c^{5/2}} + \frac{2b^2 \arctan(\sqrt{cx}) \log\left(\frac{2}{i\sqrt{cx}+1}\right)}{5c^{5/2}} - \\
& \frac{2b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(\frac{2}{\sqrt{cx}+1}\right)}{5c^{5/2}} + \frac{b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(-\frac{2\sqrt{c}(1-\sqrt{-cx})}{(\sqrt{-c}-\sqrt{c})(\sqrt{cx}+1)}\right)}{5c^{5/2}} + \\
& \frac{b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(\frac{2\sqrt{c}(\sqrt{-cx}+1)}{(\sqrt{-c}+\sqrt{c})(\sqrt{cx}+1)}\right)}{5c^{5/2}} + \frac{b^2 \arctan(\sqrt{cx}) \log\left(\frac{(1-i)(\sqrt{cx}+1)}{1-i\sqrt{cx}}\right)}{5c^{5/2}} - \\
& \frac{b^2 \arctan(\sqrt{cx}) \log(1 - cx^2)}{5c^{5/2}} - \frac{b \operatorname{arctanh}(\sqrt{cx}) (2a - b \log(1 - cx^2))}{5c^{5/2}} + \\
& \frac{b^2 \arctan(\sqrt{cx}) \log(cx^2 + 1)}{5c^{5/2}} - \frac{b^2 \operatorname{arctanh}(\sqrt{cx}) \log(cx^2 + 1)}{5c^{5/2}} + \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-\sqrt{cx}}\right)}{5c^{5/2}} + \\
& \frac{ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i\sqrt{cx}}\right)}{5c^{5/2}} - \frac{ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{(1+i)(1-\sqrt{cx})}{1-i\sqrt{cx}}\right)}{10c^{5/2}} + \\
& \frac{ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{i\sqrt{cx}+1}\right)}{5c^{5/2}} + \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{\sqrt{cx}+1}\right)}{5c^{5/2}} - \\
& \frac{b^2 \operatorname{PolyLog}\left(2, \frac{2\sqrt{c}(1-\sqrt{-cx})}{(\sqrt{-c}-\sqrt{c})(\sqrt{cx}+1)} + 1\right)}{10c^{5/2}} - \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{c}(\sqrt{-cx}+1)}{(\sqrt{-c}+\sqrt{c})(\sqrt{cx}+1)}\right)}{10c^{5/2}} - \\
& \frac{ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{(1-i)(\sqrt{cx}+1)}{1-i\sqrt{cx}}\right)}{10c^{5/2}}
\end{aligned}$$

input `Int[x^4*(a + b*ArcTanh[c*x^2])^2,x]`

output

```
(8*b^2*x)/(15*c^2) + (2*a*b*x^3)/(15*c) - (2*a*b*x^5)/25 + (2*a*b*ArcTan[Sqrt[c]*x])/(5*c^(5/2)) - (4*b^2*ArcTan[Sqrt[c]*x])/(15*c^(5/2)) + ((I/5)*b^2*ArcTan[Sqrt[c]*x]^2)/c^(5/2) - (4*b^2*ArcTanh[Sqrt[c]*x])/(15*c^(5/2)) - (b^2*ArcTanh[Sqrt[c]*x]^2)/(5*c^(5/2)) + (2*b^2*ArcTanh[Sqrt[c]*x]*Log[2/(1 - Sqrt[c]*x)])/(5*c^(5/2)) - (2*b^2*ArcTan[Sqrt[c]*x]*Log[2/(1 - I*Sqrt[c]*x)])/(5*c^(5/2)) + (b^2*ArcTan[Sqrt[c]*x]*Log[((1 + I)*(1 - Sqrt[c]*x))/(1 - I*Sqrt[c]*x)])/(5*c^(5/2)) + (2*b^2*ArcTan[Sqrt[c]*x]*Log[2/(1 + I*Sqrt[c]*x)])/(5*c^(5/2)) - (2*b^2*ArcTanh[Sqrt[c]*x]*Log[2/(1 + Sqrt[c]*x)])/(5*c^(5/2)) + (b^2*ArcTanh[Sqrt[c]*x]*Log[(-2*Sqrt[c]*(1 - Sqrt[-c]*x))/((Sqrt[-c] - Sqrt[c])*(1 + Sqrt[c]*x))])/(5*c^(5/2)) + (b^2*ArcTanh[Sqrt[c]*x]*Log[(2*Sqrt[c]*(1 + Sqrt[-c]*x))/((Sqrt[-c] + Sqrt[c])*(1 + Sqrt[c]*x))])/(5*c^(5/2)) + (b^2*ArcTan[Sqrt[c]*x]*Log[((1 - I)*(1 + Sqrt[c]*x))/(1 - I*Sqrt[c]*x)])/(5*c^(5/2)) - (b^2*x^3*Log[1 - c*x^2])/(15*c) + (b^2*x^5*Log[1 - c*x^2])/25 - (b^2*ArcTan[Sqrt[c]*x]*Log[1 - c*x^2])/(5*c^(5/2)) + (b*x^3*(2*a - b*Log[1 - c*x^2]))/(15*c) + (b*x^5*(2*a - b*Log[1 - c*x^2]))/25 - (b*ArcTanh[Sqrt[c]*x]*(2*a - b*Log[1 - c*x^2]))/(5*c^(5/2)) + (x^5*(2*a - b*Log[1 - c*x^2])^2)/20 + (2*b^2*x^3*Log[1 + c*x^2])/(15*c) + (a*b*x^5*Log[1 + c*x^2])/5 + (b^2*ArcTan[Sqrt[c]*x]*Log[1 + c*x^2])/(5*c^(5/2)) - (b^2*ArcTanh[Sqrt[c]*x]*Log[1 + c*x^2])/(5*c^(5/2)) - (b^2*x^5*Log[1 - c*x^2]*Log[1 + c*x^2])/10 + (b^2*x^5*Log[1 + c*x^2]^2)/20 + (b^2*Poly...
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6456

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Int[ExpandIntegrand[x^m*(a + b*(Log[1 + c*x^n]/2) - b*(Log[1 - c*x^n]/2))^p, x], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1] && IGtQ[n, 0] && IntegerQ[m]
```

Maple [F]

$$\int x^4 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

input

```
int(x^4*(a+b*arctanh(c*x^2))^2,x)
```

output `int(x^4*(a+b*arctanh(c*x^2))^2,x)`

Fricas [F]

$$\int x^4(a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 x^4 dx$$

input `integrate(x^4*(a+b*arctanh(c*x^2))^2,x, algorithm="fricas")`

output `integral(b^2*x^4*arctanh(c*x^2)^2 + 2*a*b*x^4*arctanh(c*x^2) + a^2*x^4, x)`

Sympy [F]

$$\int x^4(a + b \operatorname{arctanh}(cx^2))^2 dx = \int x^4(a + b \operatorname{atanh}(cx^2))^2 dx$$

input `integrate(x**4*(a+b*atanh(c*x**2))**2,x)`

output `Integral(x**4*(a + b*atanh(c*x**2))**2, x)`

Maxima [F]

$$\int x^4(a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 x^4 dx$$

input `integrate(x^4*(a+b*arctanh(c*x^2))^2,x, algorithm="maxima")`

output

```
1/5*a^2*x^5 + 1/15*(6*x^5*arctanh(c*x^2) + c*(4*x^3/c^2 + 6*arctan(sqrt(c)
*x)/c^(7/2) + 3*log((c*x - sqrt(c))/(c*x + sqrt(c)))/c^(7/2)))*a*b + 1/20*
(x^5*log(-c*x^2 + 1)^2 - 5*integrate(-1/5*(5*(c*x^6 - x^4)*log(c*x^2 + 1)^
2 - 2*(2*c*x^6 + 5*(c*x^6 - x^4)*log(c*x^2 + 1))*log(-c*x^2 + 1)/(c*x^2 -
1), x))*b^2
```

Giac [F]

$$\int x^4 (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{arctanh}(cx^2) + a)^2 x^4 dx$$

input

```
integrate(x^4*(a+b*arctanh(c*x^2))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^2) + a)^2*x^4, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^4 (a + b \operatorname{arctanh}(cx^2))^2 dx = \int x^4 (a + b \operatorname{atanh}(cx^2))^2 dx$$

input

```
int(x^4*(a + b*atanh(c*x^2))^2,x)
```

output

```
int(x^4*(a + b*atanh(c*x^2))^2, x)
```

Reduce [F]

$$\int x^4 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$= \frac{6\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) ab - 4\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) b^2 + 3 \operatorname{atanh}(cx^2)^2 b^2 c^3 x^5 - 3 \operatorname{atanh}(cx^2)^2 b^2 cx + 6\sqrt{c} \operatorname{atanh}(cx^2) ab}{15c^3}$$

input `int(x^4*(a+b*atanh(c*x^2))^2,x)`

output `(6*sqrt(c)*atan((c*x)/sqrt(c))*a*b - 4*sqrt(c)*atan((c*x)/sqrt(c))*b**2 + 3*atanh(c*x**2)**2*b**2*c**3*x**5 - 3*atanh(c*x**2)**2*b**2*c*x + 6*sqrt(c)*atanh(c*x**2)*a*b + 4*sqrt(c)*atanh(c*x**2)*b**2 + 6*atanh(c*x**2)*a*b*c**3*x**5 + 4*atanh(c*x**2)*b**2*c**2*x**3 + 6*sqrt(c)*log(sqrt(c)*x - 1)*a*b + 4*sqrt(c)*log(sqrt(c)*x - 1)*b**2 - 3*sqrt(c)*log(c*x**2 + 1)*a*b - 2*sqrt(c)*log(c*x**2 + 1)*b**2 + 3*int(atanh(c*x**2)**2,x)*b**2*c + 3*a**2*c**3*x**5 + 4*a*b*c**2*x**3 + 8*b**2*c*x)/(15*c**3)`

3.72 $\int x^2(a + \operatorname{barctanh}(cx^2))^2 dx$

Optimal result	614
Mathematica [F]	615
Rubi [A] (verified)	616
Maple [F]	618
Fricas [F]	619
Sympy [F]	619
Maxima [F]	619
Giac [F]	620
Mupad [F(-1)]	620
Reduce [F]	621

Optimal result

Integrand size = 16, antiderivative size = 1129

$$\int x^2(a + \operatorname{barctanh}(cx^2))^2 dx = \text{Too large to display}$$

output

```

-2/9*a*b*x^3-1/6*b^2*polylog(2,1-2*c^(1/2)*(1+(-c)^(1/2)*x)/((-c)^(1/2)+c^(1/2))/(1+c^(1/2)*x))/c^(3/2)-1/6*b^2*polylog(2,1+2*c^(1/2)*(1-(-c)^(1/2)*x)/((-c)^(1/2)-c^(1/2))/(1+c^(1/2)*x))/c^(3/2)+4/3*b^2*arctan(c^(1/2)*x)/c^(3/2)-1/3*b^2*arctanh(c^(1/2)*x)^2/c^(3/2)-4/3*b^2*arctanh(c^(1/2)*x)/c^(3/2)+1/3*b^2*polylog(2,1-2/(1-c^(1/2)*x))/c^(3/2)+1/3*b^2*polylog(2,1-2/(1+c^(1/2)*x))/c^(3/2)+1/9*b*x^3*(2*a-b*ln(-c*x^2+1))+1/9*b^2*x^3*ln(-c*x^2+1)+1/12*b^2*x^3*ln(c*x^2+1)^2+2/3*b^2*x*ln(c*x^2+1)/c+1/3*a*b*x^3*ln(c*x^2+1)-1/6*b^2*x^3*ln(-c*x^2+1)*ln(c*x^2+1)-2/3*b^2*x*ln(-c*x^2+1)/c+1/12*x^3*(2*a-b*ln(-c*x^2+1))^2+4/3*a*b*x/c+1/6*I*b^2*polylog(2,1+(-1+I)*(1+c^(1/2)*x)/(1-I*c^(1/2)*x))/c^(3/2)+1/6*I*b^2*polylog(2,1-(1+I)*(1-c^(1/2)*x)/(1-I*c^(1/2)*x))/c^(3/2)+1/3*b^2*arctanh(c^(1/2)*x)*ln(2*c^(1/2)*(1+(-c)^(1/2)*x)/((-c)^(1/2)+c^(1/2))/(1+c^(1/2)*x))/c^(3/2)+1/3*b^2*arctanh(c^(1/2)*x)*ln(-2*c^(1/2)*(1-(-c)^(1/2)*x)/((-c)^(1/2)-c^(1/2))/(1+c^(1/2)*x))/c^(3/2)-2/3*b^2*arctanh(c^(1/2)*x)*ln(2/(1+c^(1/2)*x))/c^(3/2)+2/3*b^2*arctanh(c^(1/2)*x)*ln(2/(1-c^(1/2)*x))/c^(3/2)-1/3*b^2*arctan(c^(1/2)*x)*ln((1+I)*(1-c^(1/2)*x)/(1-I*c^(1/2)*x))/c^(3/2)-1/3*b^2*arctan(c^(1/2)*x)*ln((1-I)*(1+c^(1/2)*x)/(1-I*c^(1/2)*x))/c^(3/2)-2/3*b^2*arctan(c^(1/2)*x)*ln(2/(1+I*c^(1/2)*x))/c^(3/2)+2/3*b^2*arctan(c^(1/2)*x)*ln(2/(1-I*c^(1/2)*x))/c^(3/2)-2/3*a*b*arctan(c^(1/2)*x)/c^(3/2)-1/3*b^2*arctanh(c^(1/2)*x)*ln(c*x^2+1)/c^(3/2)-1/3*b^2*arctan(c^(1/2)*x)*ln(c*x^2+1)/c^(3/2)-1/3*b*arctanh(...

```

Mathematica [F]

$$\int x^2(a + b \operatorname{arctanh}(cx^2))^2 dx = \int x^2(a + b \operatorname{arctanh}(cx^2))^2 dx$$

input

```
Integrate[x^2*(a + b*ArcTanh[c*x^2])^2,x]
```

output

```
Integrate[x^2*(a + b*ArcTanh[c*x^2])^2, x]
```

Rubi [A] (verified)

Time = 2.50 (sec) , antiderivative size = 1129, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {6456, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

↓ 6456

$$\int \left(\frac{1}{4} x^2 (2a - b \log(1 - cx^2))^2 - \frac{1}{2} b x^2 \log(cx^2 + 1) (b \log(1 - cx^2) - 2a) + \frac{1}{4} b^2 x^2 \log^2(cx^2 + 1) \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{1}{12}(2a - b \log(1 - cx^2))^2 x^3 + \frac{1}{12}b^2 \log^2(cx^2 + 1) x^3 - \frac{2}{9}abx^3 + \frac{1}{9}b^2 \log(1 - cx^2) x^3 + \\
& \frac{1}{9}b(2a - b \log(1 - cx^2)) x^3 + \frac{1}{3}ab \log(cx^2 + 1) x^3 - \frac{1}{6}b^2 \log(1 - cx^2) \log(cx^2 + 1) x^3 - \\
& \frac{2b^2 \log(1 - cx^2) x}{3c} + \frac{2b^2 \log(cx^2 + 1) x}{3c} + \frac{4abx}{3c} - \frac{ib^2 \arctan(\sqrt{cx})^2}{3c^{3/2}} - \frac{b^2 \operatorname{arctanh}(\sqrt{cx})^2}{3c^{3/2}} + \\
& \frac{4b^2 \arctan(\sqrt{cx})}{3c^{3/2}} - \frac{2ab \arctan(\sqrt{cx})}{3c^{3/2}} - \frac{4b^2 \operatorname{arctanh}(\sqrt{cx})}{3c^{3/2}} + \frac{2b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(\frac{2}{1-\sqrt{cx}}\right)}{3c^{3/2}} + \\
& \frac{2b^2 \arctan(\sqrt{cx}) \log\left(\frac{2}{1-i\sqrt{cx}}\right)}{3c^{3/2}} - \frac{b^2 \arctan(\sqrt{cx}) \log\left(\frac{(1+i)(1-\sqrt{cx})}{1-i\sqrt{cx}}\right)}{3c^{3/2}} - \\
& \frac{2b^2 \arctan(\sqrt{cx}) \log\left(\frac{2}{i\sqrt{cx}+1}\right)}{3c^{3/2}} - \frac{2b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(\frac{2}{\sqrt{cx}+1}\right)}{3c^{3/2}} + \\
& \frac{b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(-\frac{2\sqrt{c}(1-\sqrt{-cx})}{(\sqrt{-c}-\sqrt{c})(\sqrt{cx}+1)}\right)}{3c^{3/2}} + \frac{b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(\frac{2\sqrt{c}(\sqrt{-cx}+1)}{(\sqrt{-c}+\sqrt{c})(\sqrt{cx}+1)}\right)}{3c^{3/2}} - \\
& \frac{b^2 \arctan(\sqrt{cx}) \log\left(\frac{(1-i)(\sqrt{cx}+1)}{1-i\sqrt{cx}}\right)}{3c^{3/2}} + \frac{b^2 \arctan(\sqrt{cx}) \log(1 - cx^2)}{3c^{3/2}} - \\
& \frac{b \operatorname{arctanh}(\sqrt{cx}) (2a - b \log(1 - cx^2))}{3c^{3/2}} - \frac{b^2 \arctan(\sqrt{cx}) \log(cx^2 + 1)}{3c^{3/2}} - \\
& \frac{b^2 \operatorname{arctanh}(\sqrt{cx}) \log(cx^2 + 1)}{3c^{3/2}} + \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-\sqrt{cx}}\right)}{3c^{3/2}} - \frac{ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i\sqrt{cx}}\right)}{3c^{3/2}} + \\
& \frac{ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{(1+i)(1-\sqrt{cx})}{1-i\sqrt{cx}}\right)}{6c^{3/2}} - \frac{ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{i\sqrt{cx}+1}\right)}{3c^{3/2}} + \\
& \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{\sqrt{cx}+1}\right)}{6c^{3/2}} - \frac{b^2 \operatorname{PolyLog}\left(2, \frac{2\sqrt{c}(1-\sqrt{-cx})}{(\sqrt{-c}-\sqrt{c})(\sqrt{cx}+1)} + 1\right)}{6c^{3/2}} - \\
& \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{c}(\sqrt{-cx}+1)}{(\sqrt{-c}+\sqrt{c})(\sqrt{cx}+1)}\right)}{6c^{3/2}} + \frac{ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{(1-i)(\sqrt{cx}+1)}{1-i\sqrt{cx}}\right)}{6c^{3/2}}
\end{aligned}$$

input `Int[x^2*(a + b*ArcTanh[c*x^2])^2,x]`

output

```
(4*a*b*x)/(3*c) - (2*a*b*x^3)/9 - (2*a*b*ArcTan[Sqrt[c]*x])/(3*c^(3/2)) +
(4*b^2*ArcTan[Sqrt[c]*x])/(3*c^(3/2)) - ((I/3)*b^2*ArcTan[Sqrt[c]*x]^2)/c^
(3/2) - (4*b^2*ArcTanh[Sqrt[c]*x])/(3*c^(3/2)) - (b^2*ArcTanh[Sqrt[c]*x]^2
)/(3*c^(3/2)) + (2*b^2*ArcTanh[Sqrt[c]*x]*Log[2/(1 - Sqrt[c]*x)])/(3*c^(3/
2)) + (2*b^2*ArcTan[Sqrt[c]*x]*Log[2/(1 - I*Sqrt[c]*x)])/(3*c^(3/2)) - (b^
2*ArcTan[Sqrt[c]*x]*Log[((1 + I)*(1 - Sqrt[c]*x))/(1 - I*Sqrt[c]*x)])/(3*c
^(3/2)) - (2*b^2*ArcTan[Sqrt[c]*x]*Log[2/(1 + I*Sqrt[c]*x)])/(3*c^(3/2)) -
(2*b^2*ArcTanh[Sqrt[c]*x]*Log[2/(1 + Sqrt[c]*x)])/(3*c^(3/2)) + (b^2*ArcT
anh[Sqrt[c]*x]*Log[(-2*Sqrt[c]*(1 - Sqrt[-c]*x))/((Sqrt[-c] - Sqrt[c])*(1
+ Sqrt[c]*x))])/(3*c^(3/2)) + (b^2*ArcTanh[Sqrt[c]*x]*Log[(2*Sqrt[c]*(1 +
Sqrt[-c]*x))/((Sqrt[-c] + Sqrt[c])*(1 + Sqrt[c]*x))])/(3*c^(3/2)) - (b^2*A
rcTan[Sqrt[c]*x]*Log[((1 - I)*(1 + Sqrt[c]*x))/(1 - I*Sqrt[c]*x)])/(3*c^(3
/2)) - (2*b^2*x*Log[1 - c*x^2])/(3*c) + (b^2*x^3*Log[1 - c*x^2])/9 + (b^2*
ArcTan[Sqrt[c]*x]*Log[1 - c*x^2])/(3*c^(3/2)) + (b*x^3*(2*a - b*Log[1 - c*
x^2]))/9 - (b*ArcTanh[Sqrt[c]*x]*(2*a - b*Log[1 - c*x^2]))/(3*c^(3/2)) + (
x^3*(2*a - b*Log[1 - c*x^2])^2)/12 + (2*b^2*x*Log[1 + c*x^2])/(3*c) + (a*b
*x^3*Log[1 + c*x^2])/3 - (b^2*ArcTan[Sqrt[c]*x]*Log[1 + c*x^2])/(3*c^(3/2)
) - (b^2*ArcTanh[Sqrt[c]*x]*Log[1 + c*x^2])/(3*c^(3/2)) - (b^2*x^3*Log[1 -
c*x^2]*Log[1 + c*x^2])/6 + (b^2*x^3*Log[1 + c*x^2]^2)/12 + (b^2*PolyLog[2
, 1 - 2/(1 - Sqrt[c]*x)])/(3*c^(3/2)) - ((I/3)*b^2*PolyLog[2, 1 - 2/(1 ...
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6456

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
Int[ExpandIntegrand[x^m*(a + b*(Log[1 + c*x^n]/2) - b*(Log[1 - c*x^n]/2))^p
, x], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1] && IGtQ[n, 0] && IntegerQ[m]
```

Maple [F]

$$\int x^2 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

input

```
int(x^2*(a+b*arctanh(c*x^2))^2,x)
```

output `int(x^2*(a+b*arctanh(c*x^2))^2,x)`

Fricas [F]

$$\int x^2(a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c*x^2))^2,x, algorithm="fricas")`

output `integral(b^2*x^2*arctanh(c*x^2)^2 + 2*a*b*x^2*arctanh(c*x^2) + a^2*x^2, x)`

Sympy [F]

$$\int x^2(a + b \operatorname{arctanh}(cx^2))^2 dx = \int x^2(a + b \operatorname{atanh}(cx^2))^2 dx$$

input `integrate(x**2*(a+b*atanh(c*x**2))**2,x)`

output `Integral(x**2*(a + b*atanh(c*x**2))**2, x)`

Maxima [F]

$$\int x^2(a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c*x^2))^2,x, algorithm="maxima")`

output

```
1/3*a^2*x^3 + 1/3*(2*x^3*arctanh(c*x^2) + c*(4*x/c^2 - 2*arctan(sqrt(c)*x)
/c^(5/2) + log((c*x - sqrt(c))/(c*x + sqrt(c)))/c^(5/2)))*a*b + 1/12*(x^3*
log(-c*x^2 + 1)^2 - 3*integrate(-1/3*(3*(c*x^4 - x^2)*log(c*x^2 + 1)^2 - 2
*(2*c*x^4 + 3*(c*x^4 - x^2)*log(c*x^2 + 1))*log(-c*x^2 + 1))/(c*x^2 - 1),
x))*b^2
```

Giac [F]

$$\int x^2 (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 x^2 dx$$

input

```
integrate(x^2*(a+b*arctanh(c*x^2))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^2) + a)^2*x^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 (a + b \operatorname{arctanh}(cx^2))^2 dx = \int x^2 (a + b \operatorname{atanh}(cx^2))^2 dx$$

input

```
int(x^2*(a + b*atanh(c*x^2))^2,x)
```

output

```
int(x^2*(a + b*atanh(c*x^2))^2, x)
```

Reduce [F]

$$\int x^2 (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$= \frac{-2\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) ab + 4\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) b^2 + \operatorname{atanh}(cx^2)^2 b^2 c^2 x^3 + 2\sqrt{c} \operatorname{atanh}(cx^2) ab + 4\sqrt{c} \operatorname{atanh}(cx^2) b^2}{}$$

input `int(x^2*(a+b*atanh(c*x^2))^2,x)`

output `(- 2*sqrt(c)*atan((c*x)/sqrt(c))*a*b + 4*sqrt(c)*atan((c*x)/sqrt(c))*b**2 + atanh(c*x**2)**2*b**2*c**2*x**3 + 2*sqrt(c)*atanh(c*x**2)*a*b + 4*sqrt(c)*atanh(c*x**2)*b**2 + 2*atanh(c*x**2)*a*b*c**2*x**3 + 4*atanh(c*x**2)*b**2*c*x + 2*sqrt(c)*log(sqrt(c)*x - 1)*a*b + 4*sqrt(c)*log(sqrt(c)*x - 1)*b**2 - sqrt(c)*log(c*x**2 + 1)*a*b - 2*sqrt(c)*log(c*x**2 + 1)*b**2 + 4*int(atanh(c*x**2)/(c**2*x**4 - 1),x)*b**2*c + a**2*c**2*x**3 + 4*a*b*c*x)/(3*c**2)`

3.73 $\int (a + b \operatorname{arctanh}(cx^2))^2 dx$

Optimal result	622
Mathematica [A] (verified)	623
Rubi [A] (verified)	624
Maple [F]	626
Fricas [F]	627
Sympy [F]	627
Maxima [F]	627
Giac [F]	628
Mupad [F(-1)]	628
Reduce [F]	628

Optimal result

Integrand size = 12, antiderivative size = 958

$$\int (a + b \operatorname{arctanh}(cx^2))^2 dx = \text{Too large to display}$$

output

```

-1/2*b^2*polylog(2,1-2*c^(1/2)*(1+(-c)^(1/2)*x)/((-c)^(1/2)+c^(1/2))/(1+c^(1/2)*x))/c^(1/2)-1/2*b^2*polylog(2,1+2*c^(1/2)*(1-(-c)^(1/2)*x)/((-c)^(1/2)-c^(1/2))/(1+c^(1/2)*x))/c^(1/2)-b^2*arctanh(c^(1/2)*x)^2/c^(1/2)+b^2*polylog(2,1-2/(1+c^(1/2)*x))/c^(1/2)+b^2*polylog(2,1-2/(1-c^(1/2)*x))/c^(1/2)+1/4*b^2*x*ln(c*x^2+1)^2+1/4*b^2*x*ln(-c*x^2+1)^2-1/2*b^2*x*ln(-c*x^2+1)*ln(c*x^2+1)+a*b*x*ln(c*x^2+1)-a*b*x*ln(-c*x^2+1)+a^2*x+b^2*arctan(c^(1/2)*x)*ln((1-I)*(1+c^(1/2)*x)/(1-I*c^(1/2)*x))/c^(1/2)+2*b^2*arctanh(c^(1/2)*x)*ln(2/(1-c^(1/2)*x))/c^(1/2)-2*b^2*arctanh(c^(1/2)*x)*ln(2/(1+c^(1/2)*x))/c^(1/2)-2*a*b*arctanh(c^(1/2)*x)/c^(1/2)+2*b^2*arctan(c^(1/2)*x)*ln(2/(1+I*c^(1/2)*x))/c^(1/2)-2*b^2*arctan(c^(1/2)*x)*ln(2/(1-I*c^(1/2)*x))/c^(1/2)+2*a*b*arctan(c^(1/2)*x)/c^(1/2)-1/2*I*b^2*polylog(2,1+(-1+I)*(1+c^(1/2)*x)/(1-I*c^(1/2)*x))/c^(1/2)-1/2*I*b^2*polylog(2,1-(1+I)*(1-c^(1/2)*x)/(1-I*c^(1/2)*x))/c^(1/2)+I*b^2*polylog(2,1-2/(1+I*c^(1/2)*x))/c^(1/2)+I*b^2*polylog(2,1-2/(1-I*c^(1/2)*x))/c^(1/2)-b^2*arctanh(c^(1/2)*x)*ln(c*x^2+1)/c^(1/2)+b^2*arctan(c^(1/2)*x)*ln(c*x^2+1)/c^(1/2)-b^2*arctan(c^(1/2)*x)*ln(-c*x^2+1)/c^(1/2)+b^2*arctanh(c^(1/2)*x)*ln(-c*x^2+1)/c^(1/2)+b^2*arctanh(c^(1/2)*x)*ln(2*c^(1/2)*(1+(-c)^(1/2)*x)/((-c)^(1/2)+c^(1/2))/(1+c^(1/2)*x))/c^(1/2)+b^2*arctanh(c^(1/2)*x)*ln(-2*c^(1/2)*(1-(-c)^(1/2)*x)/((-c)^(1/2)-c^(1/2))/(1+c^(1/2)*x))/c^(1/2)+I*b^2*arctan(c^(1/2)*x)^2/c^(1/2)+b^2*arctan(c^(1/2)*x)*ln((1+I)*(1-c^(1/2)*x)/(1-I*c^(1/2)*x))/c^(1/2)

```

Mathematica [A] (verified)

Time = 1.81 (sec) , antiderivative size = 566, normalized size of antiderivative = 0.59

$$\begin{aligned}
& \int (a + b \operatorname{arctanh}(cx^2))^2 dx \\
&= \frac{1}{2} x \left(2a^2 + 4ab \operatorname{arctanh}(cx^2) + \frac{4ab \left(\arctan(\sqrt{cx^2}) - \operatorname{arctanh}(\sqrt{cx^2}) \right)}{\sqrt{cx^2}} \right) \\
&+ \frac{b^2 \left(-2i \arctan(\sqrt{cx^2})^2 + 4 \arctan(\sqrt{cx^2}) \operatorname{arctanh}(cx^2) + 2\sqrt{cx^2} \operatorname{arctanh}(cx^2)^2 + 2 \arctan(\sqrt{cx^2}) \right)}{2}
\end{aligned}$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])^2,x]
```

output

```
(x*(2*a^2 + 4*a*b*ArcTanh[c*x^2] + (4*a*b*(ArcTan[Sqrt[c*x^2]] - ArcTanh[Sqrt[c*x^2]]))/Sqrt[c*x^2] + (b^2*((-2*I)*ArcTan[Sqrt[c*x^2]]^2 + 4*ArcTan[Sqrt[c*x^2]]*ArcTanh[c*x^2] + 2*Sqrt[c*x^2]*ArcTanh[c*x^2]^2 + 2*ArcTan[Sqrt[c*x^2]]*Log[1 + E^((4*I)*ArcTan[Sqrt[c*x^2]])] + 2*ArcTanh[c*x^2]*Log[1 - Sqrt[c*x^2]] - Log[2]*Log[1 - Sqrt[c*x^2]] + Log[1 - Sqrt[c*x^2]]^2/2 - Log[1 - Sqrt[c*x^2]]*Log[(1/2 + I/2)*(-I + Sqrt[c*x^2])]) - 2*ArcTanh[c*x^2]*Log[1 + Sqrt[c*x^2]] + Log[2]*Log[1 + Sqrt[c*x^2]] + Log[((1 + I) - (1 - I)*Sqrt[c*x^2])/2]*Log[1 + Sqrt[c*x^2]] + Log[(-1/2 - I/2)*(I + Sqrt[c*x^2])]*Log[1 + Sqrt[c*x^2]] - Log[1 + Sqrt[c*x^2]]^2/2 - Log[1 - Sqrt[c*x^2]]*Log[((1 + I) + (1 - I)*Sqrt[c*x^2])/2] - (I/2)*PolyLog[2, -E^((4*I)*ArcTan[Sqrt[c*x^2]])] + PolyLog[2, (1 - Sqrt[c*x^2])/2] - PolyLog[2, (-1/2 - I/2)*(-1 + Sqrt[c*x^2])] - PolyLog[2, (-1/2 + I/2)*(-1 + Sqrt[c*x^2])] - PolyLog[2, (1 + Sqrt[c*x^2])/2] + PolyLog[2, (1/2 - I/2)*(1 + Sqrt[c*x^2])] + PolyLog[2, (1/2 + I/2)*(1 + Sqrt[c*x^2])])/Sqrt[c*x^2])/2
```

Rubi [A] (verified)

Time = 2.66 (sec) , antiderivative size = 958, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6438, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$\downarrow 6438$$

$$\int \left(a^2 - ab \log(1 - cx^2) + ab \log(cx^2 + 1) + \frac{1}{4}b^2 \log^2(1 - cx^2) + \frac{1}{4}b^2 \log^2(cx^2 + 1) - \frac{1}{2}b^2 \log(1 - cx^2) \log(cx^2 + 1) \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& xa^2 + \frac{2b \arctan(\sqrt{cx}) a}{\sqrt{c}} - \frac{2b \operatorname{arctanh}(\sqrt{cx}) a}{\sqrt{c}} - bx \log(1 - cx^2) a + bx \log(cx^2 + 1) a + \\
& \frac{ib^2 \arctan(\sqrt{cx})^2}{\sqrt{c}} - \frac{b^2 \operatorname{arctanh}(\sqrt{cx})^2}{\sqrt{c}} + \frac{1}{4} b^2 x \log^2(1 - cx^2) + \frac{1}{4} b^2 x \log^2(cx^2 + 1) + \\
& \frac{2b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(\frac{2}{1-\sqrt{cx}}\right)}{\sqrt{c}} - \frac{2b^2 \arctan(\sqrt{cx}) \log\left(\frac{2}{1-i\sqrt{cx}}\right)}{\sqrt{c}} + \\
& \frac{b^2 \arctan(\sqrt{cx}) \log\left(\frac{(1+i)(1-\sqrt{cx})}{1-i\sqrt{cx}}\right)}{\sqrt{c}} + \frac{2b^2 \arctan(\sqrt{cx}) \log\left(\frac{2}{i\sqrt{cx}+1}\right)}{\sqrt{c}} - \\
& \frac{2b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(\frac{2}{\sqrt{cx}+1}\right)}{\sqrt{c}} + \frac{b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(-\frac{2\sqrt{c}(1-\sqrt{-cx})}{(\sqrt{-c}-\sqrt{c})(\sqrt{cx}+1)}\right)}{\sqrt{c}} + \\
& \frac{b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(\frac{2\sqrt{c}(\sqrt{-cx}+1)}{(\sqrt{-c}+\sqrt{c})(\sqrt{cx}+1)}\right)}{\sqrt{c}} + \frac{b^2 \arctan(\sqrt{cx}) \log\left(\frac{(1-i)(\sqrt{cx}+1)}{1-i\sqrt{cx}}\right)}{\sqrt{c}} - \\
& \frac{b^2 \arctan(\sqrt{cx}) \log(1 - cx^2)}{\sqrt{c}} + \frac{b^2 \operatorname{arctanh}(\sqrt{cx}) \log(1 - cx^2)}{\sqrt{c}} + \\
& \frac{b^2 \arctan(\sqrt{cx}) \log(cx^2 + 1)}{\sqrt{c}} - \frac{b^2 \operatorname{arctanh}(\sqrt{cx}) \log(cx^2 + 1)}{\sqrt{c}} - \\
& \frac{1}{2} b^2 x \log(1 - cx^2) \log(cx^2 + 1) + \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-\sqrt{cx}}\right)}{\sqrt{c}} + \frac{ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i\sqrt{cx}}\right)}{\sqrt{c}} - \\
& \frac{ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{(1+i)(1-\sqrt{cx})}{1-i\sqrt{cx}}\right)}{2\sqrt{c}} + \frac{ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{i\sqrt{cx}+1}\right)}{\sqrt{c}} + \\
& \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{\sqrt{cx}+1}\right)}{\sqrt{c}} - \frac{b^2 \operatorname{PolyLog}\left(2, \frac{2\sqrt{c}(1-\sqrt{-cx})}{(\sqrt{-c}-\sqrt{c})(\sqrt{cx}+1)} + 1\right)}{2\sqrt{c}} - \\
& \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{c}(\sqrt{-cx}+1)}{(\sqrt{-c}+\sqrt{c})(\sqrt{cx}+1)}\right)}{2\sqrt{c}} - \frac{ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{(1-i)(\sqrt{cx}+1)}{1-i\sqrt{cx}}\right)}{2\sqrt{c}}
\end{aligned}$$

input `Int[(a + b*ArcTanh[c*x^2])^2,x]`

output

```
a^2*x + (2*a*b*ArcTan[Sqrt[c]*x])/Sqrt[c] + (I*b^2*ArcTan[Sqrt[c]*x]^2)/Sqrt[c] - (2*a*b*ArcTanh[Sqrt[c]*x])/Sqrt[c] - (b^2*ArcTanh[Sqrt[c]*x]^2)/Sqrt[c] + (2*b^2*ArcTanh[Sqrt[c]*x]*Log[2/(1 - Sqrt[c]*x)])/Sqrt[c] - (2*b^2*ArcTan[Sqrt[c]*x]*Log[2/(1 - I*Sqrt[c]*x)])/Sqrt[c] + (b^2*ArcTan[Sqrt[c]*x]*Log[((1 + I)*(1 - Sqrt[c]*x))/(1 - I*Sqrt[c]*x)])/Sqrt[c] + (2*b^2*ArcTan[Sqrt[c]*x]*Log[2/(1 + I*Sqrt[c]*x)])/Sqrt[c] - (2*b^2*ArcTanh[Sqrt[c]*x]*Log[2/(1 + Sqrt[c]*x)])/Sqrt[c] + (b^2*ArcTanh[Sqrt[c]*x]*Log[(-2*Sqrt[c]*(1 - Sqrt[-c]*x))/((Sqrt[-c] - Sqrt[c])*(1 + Sqrt[c]*x))])/Sqrt[c] + (b^2*ArcTanh[Sqrt[c]*x]*Log[(2*Sqrt[c]*(1 + Sqrt[-c]*x))/((Sqrt[-c] + Sqrt[c])*(1 + Sqrt[c]*x))])/Sqrt[c] + (b^2*ArcTan[Sqrt[c]*x]*Log[((1 - I)*(1 + Sqrt[c]*x))/(1 - I*Sqrt[c]*x)])/Sqrt[c] - a*b*x*Log[1 - c*x^2] - (b^2*ArcTan[Sqrt[c]*x]*Log[1 - c*x^2])/Sqrt[c] + (b^2*ArcTanh[Sqrt[c]*x]*Log[1 - c*x^2])/Sqrt[c] + (b^2*x*Log[1 - c*x^2]^2)/4 + a*b*x*Log[1 + c*x^2] + (b^2*ArcTan[Sqrt[c]*x]*Log[1 + c*x^2])/Sqrt[c] - (b^2*ArcTanh[Sqrt[c]*x]*Log[1 + c*x^2])/Sqrt[c] - (b^2*x*Log[1 - c*x^2]*Log[1 + c*x^2])/2 + (b^2*x*Log[1 + c*x^2]^2)/4 + (b^2*PolyLog[2, 1 - 2/(1 - Sqrt[c]*x)])/Sqrt[c] + (I*b^2*PolyLog[2, 1 - 2/(1 - I*Sqrt[c]*x)])/Sqrt[c] - ((I/2)*b^2*PolyLog[2, 1 - ((1 + I)*(1 - Sqrt[c]*x))/(1 - I*Sqrt[c]*x)])/Sqrt[c] + (I*b^2*PolyLog[2, 1 - 2/(1 + I*Sqrt[c]*x)])/Sqrt[c] + (b^2*PolyLog[2, 1 - 2/(1 + Sqrt[c]*x)])/Sqrt[c] - (b^2*PolyLog[2, 1 + (2*Sqrt[c]*(1 - Sqrt[-c]*x))/((Sqrt[-c] - ...
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6438

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*(Log[1 + c*x^n]/2) - b*(Log[1 - c*x^n]/2))^p, x], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1] && IGtQ[n, 0]
```

Maple [F]

$$\int (a + b \operatorname{arctanh}(cx^2))^2 dx$$

input

```
int((a+b*arctanh(c*x^2))^2,x)
```

output `int((a+b*arctanh(c*x^2))^2,x)`

Fricas [F]

$$\int (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 dx$$

input `integrate((a+b*arctanh(c*x^2))^2,x, algorithm="fricas")`

output `integral(b^2*arctanh(c*x^2)^2 + 2*a*b*arctanh(c*x^2) + a^2, x)`

Sympy [F]

$$\int (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (a + b \operatorname{atanh}(cx^2))^2 dx$$

input `integrate((a+b*atanh(c*x**2))**2,x)`

output `Integral((a + b*atanh(c*x**2))**2, x)`

Maxima [F]

$$\int (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 dx$$

input `integrate((a+b*arctanh(c*x^2))^2,x, algorithm="maxima")`

output `(c*(2*arctan(sqrt(c)*x)/c^(3/2) + log((c*x - sqrt(c))/(c*x + sqrt(c))))/c^(3/2)) + 2*x*arctanh(c*x^2))*a*b + 1/4*(x*log(-c*x^2 + 1)^2 - integrate(-((c*x^2 - 1)*log(c*x^2 + 1)^2 - 2*(2*c*x^2 + (c*x^2 - 1)*log(c*x^2 + 1))*log(-c*x^2 + 1))/(c*x^2 - 1), x))*b^2 + a^2*x`

Giac [F]

$$\int (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 dx$$

input `integrate((a+b*arctanh(c*x^2))^2,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (a + b \operatorname{atanh}(cx^2))^2 dx$$

input `int((a + b*atanh(c*x^2))^2,x)`

output `int((a + b*atanh(c*x^2))^2, x)`

Reduce [F]

$$\int (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$= \frac{2\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) ab + 2\sqrt{c} \operatorname{atanh}(cx^2) ab + 2 \operatorname{atanh}(cx^2) abcx + 2\sqrt{c} \log(\sqrt{c}x - 1) ab - \sqrt{c} \log(cx^2 + 1) ab}{c}$$

input `int((a+b*atanh(c*x^2))^2,x)`

output `(2*sqrt(c)*atan((c*x)/sqrt(c))*a*b + 2*sqrt(c)*atanh(c*x**2)*a*b + 2*atanh(c*x**2)*a*b*c*x + 2*sqrt(c)*log(sqrt(c)*x - 1)*a*b - sqrt(c)*log(c*x**2 + 1)*a*b + int(atanh(c*x**2)**2,x)*b**2*c + a**2*c*x)/c`

$$3.74 \quad \int \frac{(a+b\operatorname{arctanh}(cx^2))^2}{x^2} dx$$

Optimal result	629
Mathematica [A] (verified)	630
Rubi [A] (verified)	631
Maple [F]	633
Fricas [F]	634
Sympy [F]	634
Maxima [F]	634
Giac [F]	635
Mupad [F(-1)]	635
Reduce [F]	636

Optimal result

Integrand size = 16, antiderivative size = 942

$$\int \frac{(a + b\operatorname{arctanh}(cx^2))^2}{x^2} dx = \text{Too large to display}$$

output

```

-1/4*b^2*ln(c*x^2+1)^2/x-b^2*c^(1/2)*polylog(2,1-2/(1+c^(1/2)*x))-b^2*c^(1/2)*polylog(2,1-2/(1-c^(1/2)*x))+b^2*c^(1/2)*arctanh(c^(1/2)*x)^2+1/2*b^2*c^(1/2)*polylog(2,1-2*c^(1/2)*(1+(-c)^(1/2)*x)/((-c)^(1/2)+c^(1/2)))/(1+c^(1/2)*x))+1/2*b^2*c^(1/2)*polylog(2,1+2*c^(1/2)*(1-(-c)^(1/2)*x)/((-c)^(1/2)-c^(1/2)))/(1+c^(1/2)*x))-a*b*ln(c*x^2+1)/x+1/2*b^2*ln(-c*x^2+1)*ln(c*x^2+1)/x-1/4*(2*a-b*ln(-c*x^2+1))^2/x+I*b^2*c^(1/2)*polylog(2,1-2/(1-I*c^(1/2)*x))-b^2*c^(1/2)*arctanh(c^(1/2)*x)*ln(2*c^(1/2)*(1+(-c)^(1/2)*x)/((-c)^(1/2)+c^(1/2)))/(1+c^(1/2)*x))-b^2*c^(1/2)*arctanh(c^(1/2)*x)*ln(-2*c^(1/2)*(1-(-c)^(1/2)*x)/((-c)^(1/2)-c^(1/2)))/(1+c^(1/2)*x))+b^2*c^(1/2)*arctan(c^(1/2)*x)*ln((1+I)*(1-c^(1/2)*x)/(1-I*c^(1/2)*x))+b^2*c^(1/2)*arctan(c^(1/2)*x)*ln((1-I)*(1+c^(1/2)*x)/(1-I*c^(1/2)*x))+I*b^2*c^(1/2)*arctan(c^(1/2)*x)^2+b^2*c^(1/2)*arctanh(c^(1/2)*x)*ln(c*x^2+1)+b^2*c^(1/2)*arctan(c^(1/2)*x)*ln(c*x^2+1)-b^2*c^(1/2)*arctan(c^(1/2)*x)*ln(-c*x^2+1)+b*c^(1/2)*arctanh(c^(1/2)*x)*(2*a-b*ln(-c*x^2+1))+I*b^2*c^(1/2)*polylog(2,1-2/(1+I*c^(1/2)*x))+2*b^2*c^(1/2)*arctanh(c^(1/2)*x)*ln(2/(1+c^(1/2)*x))-2*b^2*c^(1/2)*arctanh(c^(1/2)*x)*ln(2/(1-c^(1/2)*x))+2*b^2*c^(1/2)*arctan(c^(1/2)*x)*ln(2/(1+I*c^(1/2)*x))-2*b^2*c^(1/2)*arctan(c^(1/2)*x)*ln(2/(1-I*c^(1/2)*x))+2*a*b*c^(1/2)*arctan(c^(1/2)*x)-1/2*I*b^2*c^(1/2)*polylog(2,1+(-1+I)*(1+c^(1/2)*x)/(1-I*c^(1/2)*x))-1/2*I*b^2*c^(1/2)*polylog(2,1-(1+I)*(1-c^(1/2)*x)/(1-I*c^(1/2)*x))

```

Mathematica [A] (verified)

Time = 2.32 (sec) , antiderivative size = 566, normalized size of antiderivative = 0.60

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^2} dx$$

$$= \frac{-2a^2 - 4ab \operatorname{arctanh}(cx^2) + 4ab\sqrt{cx^2} \left(\arctan(\sqrt{cx^2}) + \operatorname{arctanh}(\sqrt{cx^2}) \right) + b^2\sqrt{cx^2} \left(-2i \arctan(\sqrt{cx^2}) \right)}{x^2}$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])^2/x^2, x]
```

output

```
(-2*a^2 - 4*a*b*ArcTanh[c*x^2] + 4*a*b*Sqrt[c*x^2]*(ArcTan[Sqrt[c*x^2]] +
ArcTanh[Sqrt[c*x^2]]) + b^2*Sqrt[c*x^2]*((-2*I)*ArcTan[Sqrt[c*x^2]]^2 + 4*
ArcTan[Sqrt[c*x^2]]*ArcTanh[c*x^2] - (2*ArcTanh[c*x^2]^2)/Sqrt[c*x^2] + 2*
ArcTan[Sqrt[c*x^2]]*Log[1 + E^((4*I)*ArcTan[Sqrt[c*x^2]])] - 2*ArcTanh[c*x
^2]*Log[1 - Sqrt[c*x^2]] + Log[2]*Log[1 - Sqrt[c*x^2]] - Log[1 - Sqrt[c*x
^2]]^2/2 + Log[1 - Sqrt[c*x^2]]*Log[(1/2 + I/2)*(-I + Sqrt[c*x^2])] + 2*Arc
Tanh[c*x^2]*Log[1 + Sqrt[c*x^2]] - Log[2]*Log[1 + Sqrt[c*x^2]] - Log[((1 +
I) - (1 - I)*Sqrt[c*x^2])/2]*Log[1 + Sqrt[c*x^2]] - Log[(-1/2 - I/2)*(I +
Sqrt[c*x^2])]*Log[1 + Sqrt[c*x^2]] + Log[1 + Sqrt[c*x^2]]^2/2 + Log[1 - S
qrt[c*x^2]]*Log[((1 + I) + (1 - I)*Sqrt[c*x^2])/2] - (I/2)*PolyLog[2, -E^(
(4*I)*ArcTan[Sqrt[c*x^2]])] - PolyLog[2, (1 - Sqrt[c*x^2])/2] + PolyLog[2,
(-1/2 - I/2)*(-1 + Sqrt[c*x^2])] + PolyLog[2, (-1/2 + I/2)*(-1 + Sqrt[c*x
^2])] + PolyLog[2, (1 + Sqrt[c*x^2])/2] - PolyLog[2, (1/2 - I/2)*(1 + Sqrt
[c*x^2])] - PolyLog[2, (1/2 + I/2)*(1 + Sqrt[c*x^2])])]/(2*x)
```

Rubi [A] (verified)

Time = 1.72 (sec) , antiderivative size = 942, normalized size of antiderivative = 1.00,
 number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules
 used = {6456, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the
 transformation is given above next to the arrow. The rules definitions used are listed
 below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^2} dx$$

↓ 6456

$$\int \left(\frac{(2a - b \log(1 - cx^2))^2}{4x^2} - \frac{b \log(cx^2 + 1) (b \log(1 - cx^2) - 2a)}{2x^2} + \frac{b^2 \log^2(cx^2 + 1)}{4x^2} \right) dx$$

↓ 2009

$$\begin{aligned}
& i\sqrt{c} \arctan(\sqrt{cx})^2 b^2 + \sqrt{c} \operatorname{arctanh}(\sqrt{cx})^2 b^2 - \frac{\log^2(cx^2 + 1) b^2}{4x} - \\
& 2\sqrt{c} \operatorname{arctanh}(\sqrt{cx}) \log\left(\frac{2}{1 - \sqrt{cx}}\right) b^2 - 2\sqrt{c} \arctan(\sqrt{cx}) \log\left(\frac{2}{1 - i\sqrt{cx}}\right) b^2 + \\
& \sqrt{c} \arctan(\sqrt{cx}) \log\left(\frac{(1+i)(1-\sqrt{cx})}{1 - i\sqrt{cx}}\right) b^2 + 2\sqrt{c} \arctan(\sqrt{cx}) \log\left(\frac{2}{i\sqrt{cx} + 1}\right) b^2 + \\
& 2\sqrt{c} \operatorname{arctanh}(\sqrt{cx}) \log\left(\frac{2}{\sqrt{cx} + 1}\right) b^2 - \sqrt{c} \operatorname{arctanh}(\sqrt{cx}) \log\left(-\frac{2\sqrt{c}(1 - \sqrt{-cx})}{(\sqrt{-c} - \sqrt{c})(\sqrt{cx} + 1)}\right) b^2 - \\
& \sqrt{c} \operatorname{arctanh}(\sqrt{cx}) \log\left(\frac{2\sqrt{c}(\sqrt{-cx} + 1)}{(\sqrt{-c} + \sqrt{c})(\sqrt{cx} + 1)}\right) b^2 + \\
& \sqrt{c} \arctan(\sqrt{cx}) \log\left(\frac{(1-i)(\sqrt{cx} + 1)}{1 - i\sqrt{cx}}\right) b^2 - \sqrt{c} \arctan(\sqrt{cx}) \log(1 - cx^2) b^2 + \\
& \sqrt{c} \arctan(\sqrt{cx}) \log(cx^2 + 1) b^2 + \sqrt{c} \operatorname{arctanh}(\sqrt{cx}) \log(cx^2 + 1) b^2 + \\
& \frac{\log(1 - cx^2) \log(cx^2 + 1) b^2}{2x} - \sqrt{c} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \sqrt{cx}}\right) b^2 + \\
& i\sqrt{c} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - i\sqrt{cx}}\right) b^2 - \frac{1}{2} i\sqrt{c} \operatorname{PolyLog}\left(2, 1 - \frac{(1+i)(1-\sqrt{cx})}{1 - i\sqrt{cx}}\right) b^2 + \\
& i\sqrt{c} \operatorname{PolyLog}\left(2, 1 - \frac{2}{i\sqrt{cx} + 1}\right) b^2 - \sqrt{c} \operatorname{PolyLog}\left(2, 1 - \frac{2}{\sqrt{cx} + 1}\right) b^2 + \\
& \frac{1}{2} \sqrt{c} \operatorname{PolyLog}\left(2, \frac{2\sqrt{c}(1 - \sqrt{-cx})}{(\sqrt{-c} - \sqrt{c})(\sqrt{cx} + 1)} + 1\right) b^2 + \\
& \frac{1}{2} \sqrt{c} \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{c}(\sqrt{-cx} + 1)}{(\sqrt{-c} + \sqrt{c})(\sqrt{cx} + 1)}\right) b^2 - \\
& \frac{1}{2} i\sqrt{c} \operatorname{PolyLog}\left(2, 1 - \frac{(1-i)(\sqrt{cx} + 1)}{1 - i\sqrt{cx}}\right) b^2 + 2a\sqrt{c} \arctan(\sqrt{cx}) b + \\
& \sqrt{c} \operatorname{arctanh}(\sqrt{cx}) (2a - b \log(1 - cx^2)) b - \frac{a \log(cx^2 + 1) b}{x} - \frac{(2a - b \log(1 - cx^2))^2}{4x}
\end{aligned}$$

input `Int[(a + b*ArcTanh[c*x^2])^2/x^2,x]`

output

```

2*a*b*Sqrt[c]*ArcTan[Sqrt[c]*x] + I*b^2*Sqrt[c]*ArcTan[Sqrt[c]*x]^2 + b^2*
Sqrt[c]*ArcTanh[Sqrt[c]*x]^2 - 2*b^2*Sqrt[c]*ArcTanh[Sqrt[c]*x]*Log[2/(1 -
Sqrt[c]*x)] - 2*b^2*Sqrt[c]*ArcTan[Sqrt[c]*x]*Log[2/(1 - I*Sqrt[c]*x)] +
b^2*Sqrt[c]*ArcTan[Sqrt[c]*x]*Log[((1 + I)*(1 - Sqrt[c]*x))/(1 - I*Sqrt[c]
*x)] + 2*b^2*Sqrt[c]*ArcTan[Sqrt[c]*x]*Log[2/(1 + I*Sqrt[c]*x)] + 2*b^2*Sq
rt[c]*ArcTanh[Sqrt[c]*x]*Log[2/(1 + Sqrt[c]*x)] - b^2*Sqrt[c]*ArcTanh[Sqrt
[c]*x]*Log[(-2*Sqrt[c]*(1 - Sqrt[-c]*x))/((Sqrt[-c] - Sqrt[c])*(1 + Sqrt[c]
*x))] - b^2*Sqrt[c]*ArcTanh[Sqrt[c]*x]*Log[(2*Sqrt[c]*(1 + Sqrt[-c]*x))/
((Sqrt[-c] + Sqrt[c])*(1 + Sqrt[c]*x))] + b^2*Sqrt[c]*ArcTan[Sqrt[c]*x]*Log
[((1 - I)*(1 + Sqrt[c]*x))/(1 - I*Sqrt[c]*x)] - b^2*Sqrt[c]*ArcTan[Sqrt[c]
*x]*Log[1 - c*x^2] + b*Sqrt[c]*ArcTanh[Sqrt[c]*x]*(2*a - b*Log[1 - c*x^2])
- (2*a - b*Log[1 - c*x^2])^2/(4*x) - (a*b*Log[1 + c*x^2])/x + b^2*Sqrt[c]
*ArcTan[Sqrt[c]*x]*Log[1 + c*x^2] + b^2*Sqrt[c]*ArcTanh[Sqrt[c]*x]*Log[1 +
c*x^2] + (b^2*Log[1 - c*x^2]*Log[1 + c*x^2])/(2*x) - (b^2*Log[1 + c*x^2]^
2)/(4*x) - b^2*Sqrt[c]*PolyLog[2, 1 - 2/(1 - Sqrt[c]*x)] + I*b^2*Sqrt[c]*P
olyLog[2, 1 - 2/(1 - I*Sqrt[c]*x)] - (I/2)*b^2*Sqrt[c]*PolyLog[2, 1 - ((
1 + I)*(1 - Sqrt[c]*x))/(1 - I*Sqrt[c]*x)] + I*b^2*Sqrt[c]*PolyLog[2, 1 - 2/
(1 + I*Sqrt[c]*x)] - b^2*Sqrt[c]*PolyLog[2, 1 - 2/(1 + Sqrt[c]*x)] + (b^2*
Sqrt[c]*PolyLog[2, 1 + (2*Sqrt[c]*(1 - Sqrt[-c]*x))/((Sqrt[-c] - Sqrt[c])*
(1 + Sqrt[c]*x)))]/2 + (b^2*Sqrt[c]*PolyLog[2, 1 - (2*Sqrt[c]*(1 + Sqrt...

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6456

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.)^(p_)*(x_)^(m_.), x_Symbol] :=
Int[ExpandIntegrand[x^m*(a + b*(Log[1 + c*x^n]/2) - b*(Log[1 - c*x^n]/2))^p
, x], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1] && IGtQ[n, 0] && IntegerQ[m]
```

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^2} dx$$

input

```
int((a+b*arctanh(c*x^2))^2/x^2,x)
```

output `int((a+b*arctanh(c*x^2))^2/x^2,x)`

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^2} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{x^2} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/x^2,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x^2)^2 + 2*a*b*arctanh(c*x^2) + a^2)/x^2, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^2} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{x^2} dx$$

input `integrate((a+b*atanh(c*x**2))**2/x**2,x)`

output `Integral((a + b*atanh(c*x**2))**2/x**2, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^2} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{x^2} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/x^2,x, algorithm="maxima")`

output

```
(c*(2*arctan(sqrt(c)*x)/sqrt(c) - log((c*x - sqrt(c))/(c*x + sqrt(c)))/sqrt(c)) - 2*arctanh(c*x^2)/x)*a*b - 1/4*b^2*(log(-c*x^2 + 1)^2/x + integrate(-((c*x^2 - 1)*log(c*x^2 + 1)^2 + 2*(2*c*x^2 - (c*x^2 - 1)*log(c*x^2 + 1))*log(-c*x^2 + 1))/(c*x^4 - x^2), x)) - a^2/x
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^2} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{x^2} dx$$

input

```
integrate((a+b*arctanh(c*x^2))^2/x^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^2) + a)^2/x^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^2} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{x^2} dx$$

input

```
int((a + b*atanh(c*x^2))^2/x^2,x)
```

output

```
int((a + b*atanh(c*x^2))^2/x^2, x)
```


Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^2} dx$$

$$= \frac{2\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) abx - 2\sqrt{c} \operatorname{atanh}(cx^2) abx - 2 \operatorname{atanh}(cx^2) ab - 2\sqrt{c} \log(\sqrt{c}x - 1) abx + \sqrt{c} \log(cx^2 + 1) abx}{x}$$

input `int((a+b*atanh(c*x^2))^2/x^2,x)`

output `(2*sqrt(c)*atan((c*x)/sqrt(c))*a*b*x - 2*sqrt(c)*atanh(c*x**2)*a*b*x - 2*atanh(c*x**2)*a*b - 2*sqrt(c)*log(sqrt(c)*x - 1)*a*b*x + sqrt(c)*log(c*x**2 + 1)*a*b*x + int(atanh(c*x**2)**2/x**2,x)*b**2*x - a**2)/x`

$$3.75 \quad \int \frac{(a+b\operatorname{arctanh}(cx^2))^2}{x^4} dx$$

Optimal result	637
Mathematica [F]	638
Rubi [A] (verified)	639
Maple [F]	641
Fricas [F]	642
Sympy [F]	642
Maxima [F]	642
Giac [F]	643
Mupad [F(-1)]	643
Reduce [F]	644

Optimal result

Integrand size = 16, antiderivative size = 1102

$$\int \frac{(a + b\operatorname{arctanh}(cx^2))^2}{x^4} dx = \text{Too large to display}$$

output

```

-1/12*b^2*ln(c*x^2+1)^2/x^3+4/3*b^2*c^(3/2)*arctan(c^(1/2)*x)-1/3*b^2*c^(3/2)*polylog(2,1-2/(1+c^(1/2)*x))-1/3*b^2*c^(3/2)*polylog(2,1-2/(1-c^(1/2)*x))+4/3*b^2*c^(3/2)*arctanh(c^(1/2)*x)+1/3*b^2*c^(3/2)*arctanh(c^(1/2)*x)^2+1/6*b^2*c^(3/2)*polylog(2,1-2*c^(1/2)*(1+(-c)^(1/2)*x)/((-c)^(1/2)+c^(1/2)))/(1+c^(1/2)*x))+1/6*b^2*c^(3/2)*polylog(2,1+2*c^(1/2)*(1-(-c)^(1/2)*x)/((-c)^(1/2)-c^(1/2)))/(1+c^(1/2)*x))-1/3*b^2*c^(3/2)*arctan(c^(1/2)*x)*ln(c*x^2+1)-1/3*b^2*c^(3/2)*arctanh(c^(1/2)*x)*ln(2*c^(1/2)*(1+(-c)^(1/2)*x)/((-c)^(1/2)+c^(1/2)))/(1+c^(1/2)*x))-1/3*I*b^2*c^(3/2)*arctan(c^(1/2)*x)^2-1/3*I*b^2*c^(3/2)*polylog(2,1-2/(1+I*c^(1/2)*x))-1/3*I*b^2*c^(3/2)*polylog(2,1-2/(1-I*c^(1/2)*x))-1/3*a*b*ln(c*x^2+1)/x^3-2/3*b^2*c*ln(c*x^2+1)/x+1/6*b^2*ln(-c*x^2+1)*ln(c*x^2+1)/x^3+1/3*b^2*c*ln(-c*x^2+1)/x-1/3*b*c*(2*a-b*ln(-c*x^2+1))/x-2/3*a*b*c/x-1/12*(2*a-b*ln(-c*x^2+1))^2/x^3+1/6*I*b^2*c^(3/2)*polylog(2,1+(-1+I)*(1+c^(1/2)*x)/(1-I*c^(1/2)*x))+1/6*I*b^2*c^(3/2)*polylog(2,1-(1+I)*(1-c^(1/2)*x)/(1-I*c^(1/2)*x))+1/3*b*c^(3/2)*arctanh(c^(1/2)*x)*(2*a-b*ln(-c*x^2+1))+1/3*b^2*c^(3/2)*arctan(c^(1/2)*x)*ln(-c*x^2+1)-1/3*b^2*c^(3/2)*arctanh(c^(1/2)*x)*ln(-2*c^(1/2)*(1-(-c)^(1/2)*x)/((-c)^(1/2)-c^(1/2)))/(1+c^(1/2)*x))+2/3*b^2*c^(3/2)*arctanh(c^(1/2)*x)*ln(2/(1+c^(1/2)*x))-2/3*b^2*c^(3/2)*arctanh(c^(1/2)*x)*ln(2/(1-c^(1/2)*x))-1/3*b^2*c^(3/2)*arctan(c^(1/2)*x)*ln((1+I)*(1-c^(1/2)*x)/(1-I*c^(1/2)*x))-1/3*b^2*c^(3/2)*arctan(c^(1/2)*x)*ln((1-I)*(1+c^(1/2)*x)/(1-I*c^(1/2)*x))-2/3*b^2...

```

Mathematica [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^4} dx = \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^4} dx$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])^2/x^4, x]
```

output

```
Integrate[(a + b*ArcTanh[c*x^2])^2/x^4, x]
```

Rubi [A] (verified)

Time = 2.10 (sec) , antiderivative size = 1102, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {6456, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^4} dx$$

↓ 6456

$$\int \left(\frac{(2a - b \log(1 - cx^2))^2}{4x^4} - \frac{b \log(cx^2 + 1) (b \log(1 - cx^2) - 2a)}{2x^4} + \frac{b^2 \log^2(cx^2 + 1)}{4x^4} \right) dx$$

↓ 2009

$$\begin{aligned}
& -\frac{1}{3}ic^{3/2}\arctan(\sqrt{cx})^2b^2 + \frac{1}{3}c^{3/2}\operatorname{arctanh}(\sqrt{cx})^2b^2 - \frac{\log^2(cx^2+1)b^2}{12x^3} + \\
& \frac{4}{3}c^{3/2}\arctan(\sqrt{cx})b^2 + \frac{4}{3}c^{3/2}\operatorname{arctanh}(\sqrt{cx})b^2 - \frac{2}{3}c^{3/2}\operatorname{arctanh}(\sqrt{cx})\log\left(\frac{2}{1-\sqrt{cx}}\right)b^2 + \\
& \frac{2}{3}c^{3/2}\arctan(\sqrt{cx})\log\left(\frac{2}{1-i\sqrt{cx}}\right)b^2 - \frac{1}{3}c^{3/2}\arctan(\sqrt{cx})\log\left(\frac{(1+i)(1-\sqrt{cx})}{1-i\sqrt{cx}}\right)b^2 - \\
& \frac{2}{3}c^{3/2}\arctan(\sqrt{cx})\log\left(\frac{2}{i\sqrt{cx}+1}\right)b^2 + \frac{2}{3}c^{3/2}\operatorname{arctanh}(\sqrt{cx})\log\left(\frac{2}{\sqrt{cx}+1}\right)b^2 - \\
& \frac{1}{3}c^{3/2}\operatorname{arctanh}(\sqrt{cx})\log\left(-\frac{2\sqrt{c}(1-\sqrt{-cx})}{(\sqrt{-c}-\sqrt{c})(\sqrt{cx}+1)}\right)b^2 - \\
& \frac{1}{3}c^{3/2}\operatorname{arctanh}(\sqrt{cx})\log\left(\frac{2\sqrt{c}(\sqrt{-cx}+1)}{(\sqrt{-c}+\sqrt{c})(\sqrt{cx}+1)}\right)b^2 - \\
& \frac{1}{3}c^{3/2}\arctan(\sqrt{cx})\log\left(\frac{(1-i)(\sqrt{cx}+1)}{1-i\sqrt{cx}}\right)b^2 + \frac{1}{3}c^{3/2}\arctan(\sqrt{cx})\log(1-cx^2)b^2 + \\
& \frac{c\log(1-cx^2)b^2}{3x} - \frac{1}{3}c^{3/2}\arctan(\sqrt{cx})\log(cx^2+1)b^2 + \\
& \frac{1}{3}c^{3/2}\operatorname{arctanh}(\sqrt{cx})\log(cx^2+1)b^2 + \frac{\log(1-cx^2)\log(cx^2+1)b^2}{6x^3} - \frac{2c\log(cx^2+1)b^2}{3x} - \\
& \frac{1}{3}c^{3/2}\operatorname{PolyLog}\left(2,1-\frac{2}{1-\sqrt{cx}}\right)b^2 - \frac{1}{3}ic^{3/2}\operatorname{PolyLog}\left(2,1-\frac{2}{1-i\sqrt{cx}}\right)b^2 + \\
& \frac{1}{6}ic^{3/2}\operatorname{PolyLog}\left(2,1-\frac{(1+i)(1-\sqrt{cx})}{1-i\sqrt{cx}}\right)b^2 - \frac{1}{3}ic^{3/2}\operatorname{PolyLog}\left(2,1-\frac{2}{i\sqrt{cx}+1}\right)b^2 - \\
& \frac{1}{3}c^{3/2}\operatorname{PolyLog}\left(2,1-\frac{2}{\sqrt{cx}+1}\right)b^2 + \frac{1}{6}c^{3/2}\operatorname{PolyLog}\left(2,\frac{2\sqrt{c}(1-\sqrt{-cx})}{(\sqrt{-c}-\sqrt{c})(\sqrt{cx}+1)}+1\right)b^2 + \\
& \frac{1}{6}c^{3/2}\operatorname{PolyLog}\left(2,1-\frac{2\sqrt{c}(\sqrt{-cx}+1)}{(\sqrt{-c}+\sqrt{c})(\sqrt{cx}+1)}\right)b^2 + \\
& \frac{1}{6}ic^{3/2}\operatorname{PolyLog}\left(2,1-\frac{(1-i)(\sqrt{cx}+1)}{1-i\sqrt{cx}}\right)b^2 - \frac{2}{3}ac^{3/2}\arctan(\sqrt{cx})b + \\
& \frac{1}{3}c^{3/2}\operatorname{arctanh}(\sqrt{cx})(2a-b\log(1-cx^2))b - \frac{c(2a-b\log(1-cx^2))b}{3x} - \frac{a\log(cx^2+1)b}{3x^3} - \\
& \frac{2acb}{3x} - \frac{(2a-b\log(1-cx^2))^2}{12x^3}
\end{aligned}$$

input

```
Int[(a + b*ArcTanh[c*x^2])^2/x^4,x]
```

output

```
(-2*a*b*c)/(3*x) - (2*a*b*c^(3/2)*ArcTan[Sqrt[c]*x])/3 + (4*b^2*c^(3/2)*ArcTan[Sqrt[c]*x])/3 - (I/3)*b^2*c^(3/2)*ArcTan[Sqrt[c]*x]^2 + (4*b^2*c^(3/2)*ArcTanh[Sqrt[c]*x])/3 + (b^2*c^(3/2)*ArcTanh[Sqrt[c]*x]^2)/3 - (2*b^2*c^(3/2)*ArcTanh[Sqrt[c]*x]*Log[2/(1 - Sqrt[c]*x)])/3 + (2*b^2*c^(3/2)*ArcTan[Sqrt[c]*x]*Log[2/(1 - I*Sqrt[c]*x)])/3 - (b^2*c^(3/2)*ArcTan[Sqrt[c]*x]*Log[((1 + I)*(1 - Sqrt[c]*x))/(1 - I*Sqrt[c]*x)])/3 - (2*b^2*c^(3/2)*ArcTan[Sqrt[c]*x]*Log[2/(1 + I*Sqrt[c]*x)])/3 + (2*b^2*c^(3/2)*ArcTanh[Sqrt[c]*x]*Log[2/(1 + Sqrt[c]*x)])/3 - (b^2*c^(3/2)*ArcTanh[Sqrt[c]*x]*Log[(-2*Sqrt[c]*(1 - Sqrt[-c]*x))/((Sqrt[-c] - Sqrt[c])*(1 + Sqrt[c]*x))])/3 - (b^2*c^(3/2)*ArcTanh[Sqrt[c]*x]*Log[(2*Sqrt[c]*(1 + Sqrt[-c]*x))/((Sqrt[-c] + Sqrt[c])*(1 + Sqrt[c]*x))])/3 - (b^2*c^(3/2)*ArcTan[Sqrt[c]*x]*Log[((1 - I)*(1 + Sqrt[c]*x))/(1 - I*Sqrt[c]*x)])/3 + (b^2*c*Log[1 - c*x^2])/(3*x) + (b^2*c^(3/2)*ArcTan[Sqrt[c]*x]*Log[1 - c*x^2])/3 - (b*c*(2*a - b*Log[1 - c*x^2]))/(3*x) + (b*c^(3/2)*ArcTanh[Sqrt[c]*x]*(2*a - b*Log[1 - c*x^2]))/3 - (2*a - b*Log[1 - c*x^2])^2/(12*x^3) - (a*b*Log[1 + c*x^2])/(3*x^3) - (2*b^2*c*Log[1 + c*x^2])/(3*x) - (b^2*c^(3/2)*ArcTan[Sqrt[c]*x]*Log[1 + c*x^2])/3 + (b^2*c^(3/2)*ArcTanh[Sqrt[c]*x]*Log[1 + c*x^2])/3 + (b^2*Log[1 - c*x^2]*Log[1 + c*x^2])/(6*x^3) - (b^2*Log[1 + c*x^2]^2)/(12*x^3) - (b^2*c^(3/2)*PolyLog[2, 1 - 2/(1 - Sqrt[c]*x)])/3 - (I/3)*b^2*c^(3/2)*PolyLog[2, 1 - 2/(1 - I*Sqrt[c]*x)] + (I/6)*b^2*c^(3/2)*PolyLog[2, 1 - ((1 + I)*(1 - Sqrt[c]*x))]
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6456

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.)^(p_)*(x_)^(m_.), x_Symbol] := Int[ExpandIntegrand[x^m*(a + b*(Log[1 + c*x^n]/2) - b*(Log[1 - c*x^n]/2))^p, x], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1] && IGtQ[n, 0] && IntegerQ[m]
```

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^4} dx$$

input

```
int((a+b*arctanh(c*x^2))^2/x^4,x)
```

output `int((a+b*arctanh(c*x^2))^2/x^4,x)`

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^4} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{x^4} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/x^4,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x^2)^2 + 2*a*b*arctanh(c*x^2) + a^2)/x^4, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^4} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{x^4} dx$$

input `integrate((a+b*atanh(c*x**2))**2/x**4,x)`

output `Integral((a + b*atanh(c*x**2))**2/x**4, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^4} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{x^4} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/x^4,x, algorithm="maxima")`

output

```
-1/3*((2*sqrt(c)*arctan(sqrt(c)*x) + sqrt(c)*log((c*x - sqrt(c))/(c*x + sqrt(c))) + 4/x)*c + 2*arctanh(c*x^2)/x^3)*a*b - 1/12*b^2*(log(-c*x^2 + 1)^2/x^3 + 3*integrate(-1/3*(3*(c*x^2 - 1)*log(c*x^2 + 1)^2 + 2*(2*c*x^2 - 3*(c*x^2 - 1)*log(c*x^2 + 1))*log(-c*x^2 + 1))/(c*x^6 - x^4), x)) - 1/3*a^2/x^3
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^4} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{x^4} dx$$

input

```
integrate((a+b*arctanh(c*x^2))^2/x^4,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^2) + a)^2/x^4, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^4} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{x^4} dx$$

input

```
int((a + b*atanh(c*x^2))^2/x^4,x)
```

output

```
int((a + b*atanh(c*x^2))^2/x^4, x)
```


Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^4} dx$$

$$= \frac{-2\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) abc x^3 - 2\sqrt{c} \operatorname{atanh}(cx^2) abc x^3 - 2 \operatorname{atanh}(cx^2) ab - 2\sqrt{c} \log(\sqrt{c}x - 1) abc x^3 + \sqrt{c} \log(\sqrt{c}x + 1) abc x^3}{3x^3}$$

input `int((a+b*atanh(c*x^2))^2/x^4,x)`

output `(- 2*sqrt(c)*atan((c*x)/sqrt(c))*a*b*c*x**3 - 2*sqrt(c)*atanh(c*x**2)*a*b*c*x**3 - 2*atanh(c*x**2)*a*b - 2*sqrt(c)*log(sqrt(c)*x - 1)*a*b*c*x**3 + sqrt(c)*log(c*x**2 + 1)*a*b*c*x**3 + 3*int(atanh(c*x**2)**2/x**4,x)*b**2*x**3 - a**2 - 4*a*b*c*x**2)/(3*x**3)`

$$3.76 \quad \int \frac{(a+b\operatorname{arctanh}(cx^2))^2}{x^6} dx$$

Optimal result	645
Mathematica [F]	646
Rubi [A] (verified)	647
Maple [F]	649
Fricas [F]	650
Sympy [F]	650
Maxima [F]	650
Giac [F]	651
Mupad [F(-1)]	651
Reduce [F]	652

Optimal result

Integrand size = 16, antiderivative size = 1176

$$\int \frac{(a + b\operatorname{arctanh}(cx^2))^2}{x^6} dx = \text{Too large to display}$$

output

```

-1/5*a*b*ln(c*x^2+1)/x^5-2/15*b^2*c*ln(c*x^2+1)/x^3+1/10*b^2*ln(-c*x^2+1)*
ln(c*x^2+1)/x^5+1/15*b^2*c*ln(-c*x^2+1)/x^3-1/5*b^2*c^2*ln(-c*x^2+1)/x-1/1
5*b*c*(2*a-b*ln(-c*x^2+1))/x^3-1/5*b*c^2*(2*a-b*ln(-c*x^2+1))/x-2/15*a*b*c
/x^3+2/5*a*b*c^2/x+4/15*b^2*c^(5/2)*arctanh(c^(1/2)*x)+1/5*b^2*c^(5/2)*arc
tanh(c^(1/2)*x)^2-4/15*b^2*c^(5/2)*arctan(c^(1/2)*x)+1/10*b^2*c^(5/2)*poly
log(2,1-2*c^(1/2)*(1+(-c)^(1/2)*x)/((-c)^(1/2)+c^(1/2))/(1+c^(1/2)*x))+1/1
0*b^2*c^(5/2)*polylog(2,1+2*c^(1/2)*(1-(-c)^(1/2)*x)/((-c)^(1/2)-c^(1/2))/
(1+c^(1/2)*x))-1/5*b^2*c^(5/2)*polylog(2,1-2/(1+c^(1/2)*x))+1/5*I*b^2*c^(5
/2)*polylog(2,1-2/(1+I*c^(1/2)*x))+1/5*I*b^2*c^(5/2)*polylog(2,1-2/(1-I*c^
(1/2)*x))+1/5*I*b^2*c^(5/2)*arctan(c^(1/2)*x)^2-2/5*b^2*c^(5/2)*arctanh(c^
(1/2)*x)*ln(2/(1-c^(1/2)*x))+1/5*b^2*c^(5/2)*arctan(c^(1/2)*x)*ln((1+I)*(1
-c^(1/2)*x)/(1-I*c^(1/2)*x))+1/5*b^2*c^(5/2)*arctan(c^(1/2)*x)*ln((1-I)*(1
+c^(1/2)*x)/(1-I*c^(1/2)*x))+2/5*b^2*c^(5/2)*arctan(c^(1/2)*x)*ln(2/(1+I*c
^(1/2)*x))-2/5*b^2*c^(5/2)*arctan(c^(1/2)*x)*ln(2/(1-I*c^(1/2)*x))+2/5*a*b
*c^(5/2)*arctan(c^(1/2)*x)+1/5*b^2*c^(5/2)*arctanh(c^(1/2)*x)*ln(c*x^2+1)+
1/5*b^2*c^(5/2)*arctan(c^(1/2)*x)*ln(c*x^2+1)+1/5*b*c^(5/2)*arctanh(c^(1/2
)*x)*(2*a-b*ln(-c*x^2+1))-1/5*b^2*c^(5/2)*arctan(c^(1/2)*x)*ln(-c*x^2+1)-1
/10*I*b^2*c^(5/2)*polylog(2,1+(-1+I)*(1+c^(1/2)*x)/(1-I*c^(1/2)*x))-1/10*I
*b^2*c^(5/2)*polylog(2,1-(1+I)*(1-c^(1/2)*x)/(1-I*c^(1/2)*x))-1/5*b^2*c^(5
/2)*arctanh(c^(1/2)*x)*ln(2*c^(1/2)*(1+(-c)^(1/2)*x)/((-c)^(1/2)+c^(1/2)...

```

Mathematica [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^6} dx = \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^6} dx$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])^2/x^6, x]
```

output

```
Integrate[(a + b*ArcTanh[c*x^2])^2/x^6, x]
```

Rubi [A] (verified)

Time = 2.36 (sec) , antiderivative size = 1176, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {6456, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^6} dx$$

↓ 6456

$$\int \left(\frac{(2a - b \log(1 - cx^2))^2}{4x^6} - \frac{b \log(cx^2 + 1) (b \log(1 - cx^2) - 2a)}{2x^6} + \frac{b^2 \log^2(cx^2 + 1)}{4x^6} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{1}{5}ib^2 \arctan(\sqrt{cx})^2 c^{5/2} + \frac{1}{5}b^2 \operatorname{arctanh}(\sqrt{cx})^2 c^{5/2} - \frac{4}{15}b^2 \arctan(\sqrt{cx}) c^{5/2} + \\
& \frac{2}{5}ab \arctan(\sqrt{cx}) c^{5/2} + \frac{4}{15}b^2 \operatorname{arctanh}(\sqrt{cx}) c^{5/2} - \frac{2}{5}b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(\frac{2}{1-\sqrt{cx}}\right) c^{5/2} - \\
& \frac{2}{5}b^2 \arctan(\sqrt{cx}) \log\left(\frac{2}{1-i\sqrt{cx}}\right) c^{5/2} + \frac{1}{5}b^2 \arctan(\sqrt{cx}) \log\left(\frac{(1+i)(1-\sqrt{cx})}{1-i\sqrt{cx}}\right) c^{5/2} + \\
& \frac{2}{5}b^2 \arctan(\sqrt{cx}) \log\left(\frac{2}{i\sqrt{cx}+1}\right) c^{5/2} + \frac{2}{5}b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(\frac{2}{\sqrt{cx}+1}\right) c^{5/2} - \\
& \frac{1}{5}b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(-\frac{2\sqrt{c}(1-\sqrt{-cx})}{(\sqrt{-c}-\sqrt{c})(\sqrt{cx}+1)}\right) c^{5/2} - \\
& \frac{1}{5}b^2 \operatorname{arctanh}(\sqrt{cx}) \log\left(\frac{2\sqrt{c}(\sqrt{-cx}+1)}{(\sqrt{-c}+\sqrt{c})(\sqrt{cx}+1)}\right) c^{5/2} + \\
& \frac{1}{5}b^2 \arctan(\sqrt{cx}) \log\left(\frac{(1-i)(\sqrt{cx}+1)}{1-i\sqrt{cx}}\right) c^{5/2} - \frac{1}{5}b^2 \arctan(\sqrt{cx}) \log(1-cx^2) c^{5/2} + \\
& \frac{1}{5}b \operatorname{arctanh}(\sqrt{cx}) (2a - b \log(1-cx^2)) c^{5/2} + \frac{1}{5}b^2 \arctan(\sqrt{cx}) \log(cx^2+1) c^{5/2} + \\
& \frac{1}{5}b^2 \operatorname{arctanh}(\sqrt{cx}) \log(cx^2+1) c^{5/2} - \frac{1}{5}b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-\sqrt{cx}}\right) c^{5/2} + \\
& \frac{1}{5}ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i\sqrt{cx}}\right) c^{5/2} - \frac{1}{10}ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{(1+i)(1-\sqrt{cx})}{1-i\sqrt{cx}}\right) c^{5/2} + \\
& \frac{1}{5}ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{i\sqrt{cx}+1}\right) c^{5/2} - \frac{1}{5}b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{\sqrt{cx}+1}\right) c^{5/2} + \\
& \frac{1}{10}b^2 \operatorname{PolyLog}\left(2, \frac{2\sqrt{c}(1-\sqrt{-cx})}{(\sqrt{-c}-\sqrt{c})(\sqrt{cx}+1)} + 1\right) c^{5/2} + \\
& \frac{1}{10}b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{c}(\sqrt{-cx}+1)}{(\sqrt{-c}+\sqrt{c})(\sqrt{cx}+1)}\right) c^{5/2} - \\
& \frac{1}{10}ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{(1-i)(\sqrt{cx}+1)}{1-i\sqrt{cx}}\right) c^{5/2} - \frac{b^2 \log(1-cx^2) c^2}{5x} - \\
& \frac{b(2a - b \log(1-cx^2)) c^2}{5x} - \frac{8b^2 c^2}{15x} + \frac{2abc^2}{5x} + \frac{b^2 \log(1-cx^2) c}{15x^3} - \frac{b(2a - b \log(1-cx^2)) c}{15x^3} - \\
& \frac{2b^2 \log(cx^2+1) c}{15x^3} - \frac{2abc}{15x^3} - \frac{(2a - b \log(1-cx^2))^2}{20x^5} - \frac{b^2 \log^2(cx^2+1)}{20x^5} + \\
& \frac{b^2 \log(1-cx^2) \log(cx^2+1)}{10x^5} - \frac{ab \log(cx^2+1)}{5x^5}
\end{aligned}$$

input

Int[(a + b*ArcTanh[c*x^2])^2/x^6,x]

output

```
(-2*a*b*c)/(15*x^3) + (2*a*b*c^2)/(5*x) - (8*b^2*c^2)/(15*x) + (2*a*b*c^(5/2)*ArcTan[Sqrt[c]*x])/5 - (4*b^2*c^(5/2)*ArcTan[Sqrt[c]*x])/15 + (I/5)*b^2*c^(5/2)*ArcTan[Sqrt[c]*x]^2 + (4*b^2*c^(5/2)*ArcTanh[Sqrt[c]*x])/15 + (b^2*c^(5/2)*ArcTanh[Sqrt[c]*x]^2)/5 - (2*b^2*c^(5/2)*ArcTanh[Sqrt[c]*x]*Log[2/(1 - Sqrt[c]*x)])/5 - (2*b^2*c^(5/2)*ArcTan[Sqrt[c]*x]*Log[2/(1 - I*Sqrt[c]*x)])/5 + (b^2*c^(5/2)*ArcTan[Sqrt[c]*x]*Log[((1 + I)*(1 - Sqrt[c]*x))/(1 - I*Sqrt[c]*x)])/5 + (2*b^2*c^(5/2)*ArcTan[Sqrt[c]*x]*Log[2/(1 + I*Sqrt[c]*x)])/5 + (2*b^2*c^(5/2)*ArcTanh[Sqrt[c]*x]*Log[2/(1 + Sqrt[c]*x)])/5 - (b^2*c^(5/2)*ArcTanh[Sqrt[c]*x]*Log[(-2*Sqrt[c]*(1 - Sqrt[-c]*x))/((Sqrt[-c] - Sqrt[c])*(1 + Sqrt[c]*x))])/5 - (b^2*c^(5/2)*ArcTanh[Sqrt[c]*x]*Log[(2*Sqrt[c]*(1 + Sqrt[-c]*x))/((Sqrt[-c] + Sqrt[c])*(1 + Sqrt[c]*x))])/5 + (b^2*c^(5/2)*ArcTan[Sqrt[c]*x]*Log[((1 - I)*(1 + Sqrt[c]*x))/(1 - I*Sqrt[c]*x)])/5 + (b^2*c*Log[1 - c*x^2])/(15*x^3) - (b^2*c^2*Log[1 - c*x^2])/(5*x) - (b^2*c^(5/2)*ArcTan[Sqrt[c]*x]*Log[1 - c*x^2])/5 - (b*c*(2*a - b*Log[1 - c*x^2]))/(15*x^3) - (b*c^2*(2*a - b*Log[1 - c*x^2]))/(5*x) + (b*c^(5/2)*ArcTanh[Sqrt[c]*x]*(2*a - b*Log[1 - c*x^2]))/5 - (2*a - b*Log[1 - c*x^2])^2/(20*x^5) - (a*b*Log[1 + c*x^2])/(5*x^5) - (2*b^2*c*Log[1 + c*x^2])/(15*x^3) + (b^2*c^(5/2)*ArcTan[Sqrt[c]*x]*Log[1 + c*x^2])/5 + (b^2*c^(5/2)*ArcTanh[Sqrt[c]*x]*Log[1 + c*x^2])/5 + (b^2*Log[1 - c*x^2]*Log[1 + c*x^2])/(10*x^5) - (b^2*Log[1 + c*x^2]^2)/(20*x^5) - (b^2*c^(5/2)*PolyLog[2, 1 - ...
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6456

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.)^(p_)*(x_)^(m_.), x_Symbol] := Int[ExpandIntegrand[x^m*(a + b*(Log[1 + c*x^n]/2) - b*(Log[1 - c*x^n]/2))^p, x], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1] && IGtQ[n, 0] && IntegerQ[m]
```

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^6} dx$$

input

```
int((a+b*arctanh(c*x^2))^2/x^6,x)
```

output `int((a+b*arctanh(c*x^2))^2/x^6,x)`

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^6} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{x^6} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/x^6,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x^2)^2 + 2*a*b*arctanh(c*x^2) + a^2)/x^6, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^6} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{x^6} dx$$

input `integrate((a+b*atanh(c*x**2))**2/x**6,x)`

output `Integral((a + b*atanh(c*x**2))**2/x**6, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^6} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{x^6} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/x^6,x, algorithm="maxima")`

output

```
1/15*((6*c^(3/2)*arctan(sqrt(c)*x) - 3*c^(3/2)*log((c*x +
sqrt(c))) - 4/x^3)*c - 6*arctanh(c*x^2)/x^5)*a*b - 1/20*b^2*(log(-c*x^2 +
1)^2/x^5 + 5*integrate(-1/5*(5*(c*x^2 - 1)*log(c*x^2 + 1)^2 + 2*(2*c*x^2 -
5*(c*x^2 - 1)*log(c*x^2 + 1))*log(-c*x^2 + 1))/(c*x^8 - x^6), x)) - 1/5*a
^2/x^5
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^6} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{x^6} dx$$

input

```
integrate((a+b*arctanh(c*x^2))^2/x^6,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^2) + a)^2/x^6, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^6} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{x^6} dx$$

input

```
int((a + b*atanh(c*x^2))^2/x^6,x)
```

output

```
int((a + b*atanh(c*x^2))^2/x^6, x)
```


Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^6} dx$$

$$= \frac{6\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) ab c^2 x^5 - 6\sqrt{c} \operatorname{atanh}(cx^2) ab c^2 x^5 - 6 \operatorname{atanh}(cx^2) ab - 6\sqrt{c} \log(\sqrt{c}x - 1) ab c^2 x^5 + 3\sqrt{c}}{15x^5}$$

input `int((a+b*atanh(c*x^2))^2/x^6,x)`

output `(6*sqrt(c)*atan((c*x)/sqrt(c))*a*b*c**2*x**5 - 6*sqrt(c)*atanh(c*x**2)*a*b*c**2*x**5 - 6*atanh(c*x**2)*a*b - 6*sqrt(c)*log(sqrt(c)*x - 1)*a*b*c**2*x**5 + 3*sqrt(c)*log(c*x**2 + 1)*a*b*c**2*x**5 + 15*int(atanh(c*x**2)**2/x**6,x)*b**2*x**5 - 3*a**2 - 4*a*b*c*x**2)/(15*x**5)`

3.77 $\int x^3(a + \operatorname{barctanh}(cx^2))^3 dx$

Optimal result	653
Mathematica [A] (verified)	654
Rubi [A] (verified)	654
Maple [C] (warning: unable to verify)	658
Fricas [F]	659
Sympy [F]	659
Maxima [F]	659
Giac [F]	660
Mupad [F(-1)]	660
Reduce [F]	661

Optimal result

Integrand size = 16, antiderivative size = 141

$$\int x^3(a + \operatorname{barctanh}(cx^2))^3 dx = \frac{3b(a + \operatorname{barctanh}(cx^2))^2}{4c^2} + \frac{3bx^2(a + \operatorname{barctanh}(cx^2))^2}{4c} - \frac{(a + \operatorname{barctanh}(cx^2))^3}{4c^2} + \frac{1}{4}x^4(a + \operatorname{barctanh}(cx^2))^3 - \frac{3b^2(a + \operatorname{barctanh}(cx^2)) \log\left(\frac{2}{1-cx^2}\right)}{2c^2} - \frac{3b^3 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^2}\right)}{4c^2}$$

output

```
3/4*b*(a+b*arctanh(c*x^2))^2/c^2+3/4*b*x^2*(a+b*arctanh(c*x^2))^2/c-1/4*(a+b*arctanh(c*x^2))^3/c^2+1/4*x^4*(a+b*arctanh(c*x^2))^3-3/2*b^2*(a+b*arctanh(c*x^2))*ln(2/(-c*x^2+1))/c^2-3/4*b^3*polylog(2,1-2/(-c*x^2+1))/c^2
```

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.31

$$\int x^3 (a + b \operatorname{arctanh}(cx^2))^3 dx$$

$$= \frac{6b^2(-1 + cx^2)(a + b + acx^2) \operatorname{arctanh}(cx^2)^2 + 2b^3(-1 + c^2x^4) \operatorname{arctanh}(cx^2)^3 + 6b \operatorname{arctanh}(cx^2) (acx^2(2b$$

input `Integrate[x^3*(a + b*ArcTanh[c*x^2])^3,x]`

output

```
(6*b^2*(-1 + c*x^2)*(a + b + a*c*x^2)*ArcTanh[c*x^2]^2 + 2*b^3*(-1 + c^2*x^4)*ArcTanh[c*x^2]^3 + 6*b*ArcTanh[c*x^2]*(a*c*x^2*(2*b + a*c*x^2) - 2*b^2*Log[1 + E^(-2*ArcTanh[c*x^2])]) + a*(6*a*b*c*x^2 + 2*a^2*c^2*x^4 + 3*a*b*Log[1 - c*x^2] - 3*a*b*Log[1 + c*x^2] + 6*b^2*Log[1 - c^2*x^4]) + 6*b^3*PolyLog[2, -E^(-2*ArcTanh[c*x^2])])/(8*c^2)
```

Rubi [A] (verified)

Time = 1.09 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.12, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.562$, Rules used = {6454, 6452, 6542, 6436, 6510, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 (a + b \operatorname{arctanh}(cx^2))^3 dx$$

$$\downarrow \text{6454}$$

$$\frac{1}{2} \int x^2 (a + b \operatorname{arctanh}(cx^2))^3 dx^2$$

$$\downarrow \text{6452}$$

$$\frac{1}{2} \left(\frac{1}{2} x^4 (a + b \operatorname{arctanh}(cx^2))^3 - \frac{3}{2} bc \int \frac{x^4 (a + b \operatorname{arctanh}(cx^2))^2}{1 - c^2 x^4} dx^2 \right)$$

$$\downarrow \text{6542}$$

$$\frac{1}{2} \left(\frac{1}{2} x^4 (a + \operatorname{barctanh}(cx^2))^3 - \frac{3}{2} bc \left(\frac{\int \frac{(a + \operatorname{barctanh}(cx^2))^2}{1 - c^2 x^4} dx^2}{c^2} - \frac{\int (a + \operatorname{barctanh}(cx^2))^2 dx^2}{c^2} \right) \right)$$

↓ 6436

$$\frac{1}{2} \left(\frac{1}{2} x^4 (a + \operatorname{barctanh}(cx^2))^3 - \frac{3}{2} bc \left(\frac{\int \frac{(a + \operatorname{barctanh}(cx^2))^2}{1 - c^2 x^4} dx^2}{c^2} - \frac{x^2 (a + \operatorname{barctanh}(cx^2))^2 - 2bc \int \frac{x^2 (a + \operatorname{barctanh}(cx^2))}{1 - c^2 x^4}}{c^2} \right) \right)$$

↓ 6510

$$\frac{1}{2} \left(\frac{1}{2} x^4 (a + \operatorname{barctanh}(cx^2))^3 - \frac{3}{2} bc \left(\frac{(a + \operatorname{barctanh}(cx^2))^3}{3bc^3} - \frac{x^2 (a + \operatorname{barctanh}(cx^2))^2 - 2bc \int \frac{x^2 (a + \operatorname{barctanh}(cx^2))}{1 - c^2 x^4}}{c^2} \right) \right)$$

↓ 6546

$$\frac{1}{2} \left(\frac{1}{2} x^4 (a + \operatorname{barctanh}(cx^2))^3 - \frac{3}{2} bc \left(\frac{(a + \operatorname{barctanh}(cx^2))^3}{3bc^3} - \frac{x^2 (a + \operatorname{barctanh}(cx^2))^2 - 2bc \left(\frac{\int \frac{a + \operatorname{barctanh}(cx^2)}{1 - cx^2}}{c} \right)}{c^2} \right) \right)$$

↓ 6470

$$\frac{1}{2} \left(\frac{1}{2} x^4 (a + \operatorname{barctanh}(cx^2))^3 - \frac{3}{2} bc \left(\frac{(a + \operatorname{barctanh}(cx^2))^3}{3bc^3} - \frac{x^2 (a + \operatorname{barctanh}(cx^2))^2 - 2bc \left(\frac{\log\left(\frac{2}{1 - cx^2}\right) (a + \operatorname{barctanh}(cx^2))}{c} \right)}{c^2} \right) \right)$$

↓ 2849

$$\frac{1}{2} \left(\frac{1}{2} x^4 (a + \operatorname{arctanh}(cx^2))^3 - \frac{3}{2} bc \left(\frac{(a + \operatorname{arctanh}(cx^2))^3}{3bc^3} - \frac{x^2 (a + \operatorname{arctanh}(cx^2))^2 - 2bc \left(\frac{b \int \frac{\log\left(\frac{2}{1-cx^2}\right) d \frac{1}{1-cx^2}}{\frac{1-cx^2}{c}}}{c} \right)}{c} \right) \right)$$

↓ 2752

$$\frac{1}{2} \left(\frac{1}{2} x^4 (a + \operatorname{arctanh}(cx^2))^3 - \frac{3}{2} bc \left(\frac{(a + \operatorname{arctanh}(cx^2))^3}{3bc^3} - \frac{x^2 (a + \operatorname{arctanh}(cx^2))^2 - 2bc \left(\frac{\log\left(\frac{2}{1-cx^2}\right) (a + \operatorname{arctanh}(cx^2))}{c} \right)}{c} \right) \right)$$

input `Int[x^3*(a + b*ArcTanh[c*x^2])^3,x]`

output `((x^4*(a + b*ArcTanh[c*x^2])^3)/2 - (3*b*c*((a + b*ArcTanh[c*x^2])^3/(3*b*c^3) - (x^2*(a + b*ArcTanh[c*x^2])^2 - 2*b*c*(-1/2*(a + b*ArcTanh[c*x^2])^2/(b*c^2) + (((a + b*ArcTanh[c*x^2])*Log[2/(1 - c*x^2)])/c + (b*PolyLog[2, 1 - 2/(1 - c*x^2)])/(2*c))/c))/c^2))/2/2`

Defintions of rubi rules used

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] :> Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] :> Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 6436 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot b)^p, x_Symbol] \rightarrow \text{Simp}[x \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^p, x] - \text{Simp}[b \cdot c \cdot n \cdot p \cdot \text{Int}[x^n \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2n})], x], x] /;$ $\text{FreeQ}\{a, b, c, n, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[n, 1] \ || \ \text{EqQ}[p, 1])$

rule 6452 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot b)^p \cdot x^m, x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^p / (m+1), x] - \text{Simp}[b \cdot c \cdot n \cdot (p/(m+1)) \cdot \text{Int}[x^{m+n} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2n})], x], x] /;$ $\text{FreeQ}\{a, b, c, m, n, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 6454 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot b)^p \cdot x^m, x_Symbol] \rightarrow \text{Simp}[1/n \cdot \text{Subst}[\text{Int}[x^{\text{Simplify}[(m+1)/n} - 1] \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x, x^n], x] /;$ $\text{FreeQ}\{a, b, c, m, n, x\} \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$

rule 6470 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((d) + (e) \cdot x), x_Symbol] \rightarrow \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{Log}[2/(1 + e \cdot (x/d))]/e), x] + \text{Simp}[b \cdot c \cdot (p/e) \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{Log}[2/(1 + e \cdot (x/d))]/(1 - c^2 \cdot x^2)), x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d^2 - e^2, 0]$

rule 6510 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((d) + (e) \cdot x^2), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot c \cdot d \cdot (p+1)), x] /;$ $\text{FreeQ}\{a, b, c, d, e, p, x\} \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{NeQ}[p, -1]$

rule 6542 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p \cdot (f \cdot x)^m / ((d) + (e) \cdot x^2), x_Symbol] \rightarrow \text{Simp}[f^2/e \cdot \text{Int}[(f \cdot x)^{m-2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x] - \text{Simp}[d \cdot (f^2/e) \cdot \text{Int}[(f \cdot x)^{m-2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2)], x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, x\} \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{GtQ}[m, 1]$

rule 6546

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)*(x_.)/((d_.) + (e_.)*(x_.)^2),
 x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/
 (c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.18 (sec) , antiderivative size = 798, normalized size of antiderivative = 5.66

method	result	size
risch	Expression too large to display	798

input

```
int(x^3*(a+b*arctanh(c*x^2))^3,x,method=_RETURNVERBOSE)
```

output

```
1/32*b^3*(c^2*x^4-1)/c^2*ln(c*x^2+1)^3+3/32*b^2*(-b*c^2*ln(-c*x^2+1)*x^4+2
*a*c^2*x^4+2*b*c*x^2+b*ln(-c*x^2+1)-2*a+2*b)/c^2*ln(c*x^2+1)^2+(3/32*b^3*(
c^2*x^4-1)/c^2*ln(-c*x^2+1)^2-3/32*b^2*(2*a*c*x^2+b)^2/c^2/a*ln(-c*x^2+1)-
3/32*b*(-4*a^3*c^2*x^4-8*a^2*b*c*x^2-4*ln(-c*x^2+1)*a^2*b-4*ln(-c*x^2+1)*a
*b^2-ln(-c*x^2+1)*b^3-4*a*b^2)/a/c^2)*ln(c*x^2+1)-3/8/c^2*b^3*ln(c*x^2-1)+
3/8/c^2*b^3*ln(-c*x^2+1)-1/32*b^3*x^4*ln(-c*x^2+1)^3-3/16*b^3/c^2*ln(-c*x^
2+1)^2+1/32*b^3/c^2*ln(-c*x^2+1)^3+3/4/c*b^2*Sum(-ln(x-_alpha)*ln(-c*x^2+
1)+2*c*(-1/2*ln(x-_alpha)*(ln((RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=1)-x+_a
lpha)/RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=1))+ln((RootOf(_Z^2*c+2*_Z*_alph
a*c-2,index=2)-x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=2)))/c-1/2*(d
ilog((RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=1)-x+_alpha)/RootOf(_Z^2*c+2*_Z*
_alpha*c-2,index=1))+dilog((RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=2)-x+_alph
a)/RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=2)))/c)*b/c,_alpha=RootOf(_Z^2*c+1
))-3/8/c^2*b^3*ln(c*x^2+1)+3/16*b^3/c*x^2*ln(-c*x^2+1)^2-3/8*a^2*b*x^4*ln(
-c*x^2+1)+3/4*a^2*b*x^2/c+3/8*a^2*b/c^2*ln(c*x^2-1)+3/16*a*b^2*x^4*ln(-c*x
^2+1)^2+9/16/c^2*a*b^2*ln(-c*x^2+1)-3/16/c^2*a*b^2*ln(-c*x^2+1)^2-3/8*b/c^
2*ln(c*x^2+1)*a^2+3/4*b^2/c^2*ln(c*x^2+1)*a+3/16/c^2*b^2*a*ln(c*x^2-1)-3/4
*a*b^2/c*x^2*ln(-c*x^2+1)-3/16*b^3/c^2+1/4*a^3*x^4
```

Fricas [F]

$$\int x^3 (a + b \operatorname{arctanh}(cx^2))^3 dx = \int (b \operatorname{artanh}(cx^2) + a)^3 x^3 dx$$

input `integrate(x^3*(a+b*arctanh(c*x^2))^3,x, algorithm="fricas")`

output `integral(b^3*x^3*arctanh(c*x^2)^3 + 3*a*b^2*x^3*arctanh(c*x^2)^2 + 3*a^2*b*x^3*arctanh(c*x^2) + a^3*x^3, x)`

Sympy [F]

$$\int x^3 (a + b \operatorname{arctanh}(cx^2))^3 dx = \int x^3 (a + b \operatorname{atanh}(cx^2))^3 dx$$

input `integrate(x**3*(a+b*atanh(c*x**2))**3,x)`

output `Integral(x**3*(a + b*atanh(c*x**2))**3, x)`

Maxima [F]

$$\int x^3 (a + b \operatorname{arctanh}(cx^2))^3 dx = \int (b \operatorname{artanh}(cx^2) + a)^3 x^3 dx$$

input `integrate(x^3*(a+b*arctanh(c*x^2))^3,x, algorithm="maxima")`

output

```

3/4*a*b^2*x^4*arctanh(c*x^2)^2 + 1/4*a^3*x^4 + 3/8*(2*x^4*arctanh(c*x^2) +
c*(2*x^2/c^2 - log(c*x^2 + 1)/c^3 + log(c*x^2 - 1)/c^3))*a^2*b + 3/16*(4*
c*(2*x^2/c^2 - log(c*x^2 + 1)/c^3 + log(c*x^2 - 1)/c^3)*arctanh(c*x^2) - (
2*(log(c*x^2 - 1) - 2)*log(c*x^2 + 1) - log(c*x^2 + 1)^2 - log(c*x^2 - 1)^
2 - 4*log(c*x^2 - 1))/c^2)*a*b^2 - 1/128*(4*x^4*log(-c*x^2 + 1)^3 + 3*c^3*
(x^4/c^3 + log(c^2*x^4 - 1)/c^5) - 6*c*((c*x^4 + 2*x^2)/c^2 + 2*log(c*x^2
- 1)/c^3)*log(-c*x^2 + 1)^2 + 21*c^2*(2*x^2/c^3 - log(c*x^2 + 1)/c^4 + log
(c*x^2 - 1)/c^4) + c*(6*(c^2*x^4 + 6*c*x^2 + 2*log(c*x^2 - 1)^2 + 6*log(c*
x^2 - 1))*log(-c*x^2 + 1)/c^3 - (3*c^2*x^4 + 42*c*x^2 + 4*log(c*x^2 - 1)^3
+ 18*log(c*x^2 - 1)^2 + 42*log(c*x^2 - 1))/c^3) - 1152*c*integrate(1/4*x^
3*log(c*x^2 + 1)/(c^3*x^4 - c), x) - 2*(12*c*x^2*log(c*x^2 + 1)^2 + 2*(c^2
*x^4 - 1)*log(c*x^2 + 1)^3 - 3*(c^2*x^4 - 2*c*x^2 - 2*(c^2*x^4 - 1)*log(c*
x^2 + 1) + 1)*log(-c*x^2 + 1)^2 + 3*(c^2*x^4 + 6*c*x^2 - 2*(c^2*x^4 - 1)*l
og(c*x^2 + 1)^2 - 8*(c*x^2 + 1)*log(c*x^2 + 1))*log(-c*x^2 + 1))/c^2 + 18*
log(4*c^3*x^4 - 4*c)/c^2 - 384*integrate(1/4*x*log(c*x^2 + 1)/(c^3*x^4 - c
), x))*b^3

```

Giac [F]

$$\int x^3 (a + b \operatorname{arctanh}(cx^2))^3 dx = \int (b \operatorname{artanh}(cx^2) + a)^3 x^3 dx$$

input

```
integrate(x^3*(a+b*arctanh(c*x^2))^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^2) + a)^3*x^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^3 (a + b \operatorname{arctanh}(cx^2))^3 dx = \int x^3 (a + b \operatorname{atanh}(cx^2))^3 dx$$

input

```
int(x^3*(a + b*atanh(c*x^2))^3,x)
```

output `int(x^3*(a + b*atanh(c*x^2))^3, x)`

Reduce [F]

$$\int x^3 (a + b \operatorname{arctanh}(cx^2))^3 dx$$

$$= \frac{\operatorname{atanh}(cx^2)^3 b^3 c^2 x^4 - \operatorname{atanh}(cx^2)^3 b^3 + 3 \operatorname{atanh}(cx^2)^2 a b^2 c^2 x^4 - 3 \operatorname{atanh}(cx^2)^2 a b^2 + 3 \operatorname{atanh}(cx^2)^2 b^3 c x^4 + 3 \operatorname{atanh}(cx^2)^2 a b^2 c x^4 - 3 \operatorname{atanh}(cx^2)^2 a b^2 + 3 \operatorname{atanh}(cx^2)^2 b^3 c x^4 + 3 \operatorname{atanh}(cx^2)^2 a b^2 c x^4 - 3 \operatorname{atanh}(cx^2)^2 a b^2 + 3 \operatorname{atanh}(cx^2)^2 b^3 c x^4 + \dots}{4c^2}$$

input `int(x^3*(a+b*atanh(c*x^2))^3,x)`

output `(atanh(c*x**2)**3*b**3*c**2*x**4 - atanh(c*x**2)**3*b**3 + 3*atanh(c*x**2)**2*a*b**2*c**2*x**4 - 3*atanh(c*x**2)**2*a*b**2 + 3*atanh(c*x**2)**2*b**3*c*x**2 + 3*atanh(c*x**2)*a**2*b*c**2*x**4 - 3*atanh(c*x**2)*a**2*b + 6*atanh(c*x**2)*a*b**2*c*x**2 - 6*atanh(c*x**2)*a*b**2 + 12*int((atanh(c*x**2)*x**3)/(c**2*x**4 - 1),x)*b**3*c**2 + 6*log(c*x**2 + 1)*a*b**2 + a**3*c**2*x**4 + 3*a**2*b*c*x**2)/(4*c**2)`

3.78 $\int x(a + \operatorname{barctanh}(cx^2))^3 dx$

Optimal result	662
Mathematica [A] (verified)	663
Rubi [A] (verified)	663
Maple [B] (verified)	666
Fricas [F]	666
Sympy [F]	667
Maxima [F]	667
Giac [F]	668
Mupad [F(-1)]	668
Reduce [F]	668

Optimal result

Integrand size = 14, antiderivative size = 134

$$\int x(a + \operatorname{barctanh}(cx^2))^3 dx = \frac{(a + \operatorname{barctanh}(cx^2))^3}{2c} + \frac{1}{2}x^2(a + \operatorname{barctanh}(cx^2))^3 - \frac{3b(a + \operatorname{barctanh}(cx^2))^2 \log\left(\frac{2}{1-cx^2}\right)}{2c} - \frac{3b^2(a + \operatorname{barctanh}(cx^2)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^2}\right)}{2c} + \frac{3b^3 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx^2}\right)}{4c}$$

output

```
1/2*(a+b*arctanh(c*x^2))^3/c+1/2*x^2*(a+b*arctanh(c*x^2))^3-3/2*b*(a+b*arctanh(c*x^2))^2*ln(2/(-c*x^2+1))/c-3/2*b^2*(a+b*arctanh(c*x^2))*polylog(2,1-2/(-c*x^2+1))/c+3/4*b^3*polylog(3,1-2/(-c*x^2+1))/c
```

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.59

$$\int x(a + b \operatorname{arctanh}(cx^2))^3 dx = \frac{a^3 x^2}{2} + \frac{3}{2} a^2 b x^2 \operatorname{arctanh}(cx^2) + \frac{3a^2 b \log(1 - c^2 x^4)}{4c} + \frac{3ab^2 \left(\operatorname{arctanh}(cx^2) \left(-\operatorname{arctanh}(cx^2) + cx^2 \operatorname{arctanh}(cx^2) - 2 \log(1 + e^{-2 \operatorname{arctanh}(cx^2)}) \right) \right) + \operatorname{PolyLog}(2, -e^{-2 \operatorname{arctanh}(cx^2)})}{2c} + \frac{b^3 \left(\operatorname{arctanh}(cx^2)^2 \left(-\operatorname{arctanh}(cx^2) + cx^2 \operatorname{arctanh}(cx^2) - 3 \log(1 + e^{-2 \operatorname{arctanh}(cx^2)}) \right) \right) + 3 \operatorname{arctanh}(cx^2) \operatorname{PolyLog}(3, -e^{-2 \operatorname{arctanh}(cx^2)})}{2c}$$

input `Integrate[x*(a + b*ArcTanh[c*x^2])^3,x]`

output
$$\frac{(a^3 x^2)/2 + (3 a^2 b x^2 \operatorname{ArcTanh}[c x^2])/2 + (3 a^2 b \operatorname{Log}[1 - c^2 x^4])/(4 c) + (3 a b^2 (\operatorname{ArcTanh}[c x^2] (-\operatorname{ArcTanh}[c x^2] + c x^2 \operatorname{ArcTanh}[c x^2] - 2 \operatorname{Log}[1 + E^{-2 \operatorname{ArcTanh}[c x^2]})]) + \operatorname{PolyLog}[2, -E^{-2 \operatorname{ArcTanh}[c x^2]}])/(2 c) + (b^3 (\operatorname{ArcTanh}[c x^2]^2 (-\operatorname{ArcTanh}[c x^2] + c x^2 \operatorname{ArcTanh}[c x^2] - 3 \operatorname{Log}[1 + E^{-2 \operatorname{ArcTanh}[c x^2]})]) + 3 \operatorname{ArcTanh}[c x^2] \operatorname{PolyLog}[2, -E^{-2 \operatorname{ArcTanh}[c x^2]}]) + (3 \operatorname{PolyLog}[3, -E^{-2 \operatorname{ArcTanh}[c x^2]}])/(2))}{2 c}$$

Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.07, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {6454, 6436, 6546, 6470, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(a + b \operatorname{arctanh}(cx^2))^3 dx$$

$$\downarrow 6454$$

$$\frac{1}{2} \int (a + b \operatorname{arctanh}(cx^2))^3 dx^2$$

$$\downarrow 6436$$

$$\begin{aligned}
 & \frac{1}{2} \left(x^2 (a + \operatorname{barctanh}(cx^2))^3 - 3bc \int \frac{x^2 (a + \operatorname{barctanh}(cx^2))^2}{1 - c^2x^4} dx^2 \right) \\
 & \quad \downarrow 6546 \\
 & \frac{1}{2} \left(x^2 (a + \operatorname{barctanh}(cx^2))^3 - 3bc \left(\frac{\int \frac{(a + \operatorname{barctanh}(cx^2))^2}{1 - cx^2} dx^2}{c} - \frac{(a + \operatorname{barctanh}(cx^2))^3}{3bc^2} \right) \right) \\
 & \quad \downarrow 6470 \\
 & \frac{1}{2} \left(x^2 (a + \operatorname{barctanh}(cx^2))^3 - 3bc \left(\frac{\log\left(\frac{2}{1 - cx^2}\right) (a + \operatorname{barctanh}(cx^2))^2}{c} - \frac{2b \int \frac{(a + \operatorname{barctanh}(cx^2)) \log\left(\frac{2}{1 - cx^2}\right)}{1 - c^2x^4} dx^2}{c} - \frac{(a + \operatorname{barctanh}(cx^2))^3}{3bc^2} \right) \right) \\
 & \quad \downarrow 6620 \\
 & \frac{1}{2} \left(x^2 (a + \operatorname{barctanh}(cx^2))^3 - 3bc \left(\frac{\log\left(\frac{2}{1 - cx^2}\right) (a + \operatorname{barctanh}(cx^2))^2}{c} - 2b \left(\frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx^2}\right)}{1 - c^2x^4} dx^2 - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx^2}\right)}{c} \right) - \frac{(a + \operatorname{barctanh}(cx^2))^3}{3bc^2} \right) \right) \\
 & \quad \downarrow 7164 \\
 & \frac{1}{2} \left(x^2 (a + \operatorname{barctanh}(cx^2))^3 - 3bc \left(\frac{\log\left(\frac{2}{1 - cx^2}\right) (a + \operatorname{barctanh}(cx^2))^2}{c} - 2b \left(\frac{b \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - cx^2}\right)}{4c} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx^2}\right)}{2c} \right) - \frac{(a + \operatorname{barctanh}(cx^2))^3}{3bc^2} \right) \right)
 \end{aligned}$$

input

```
Int [x*(a + b*ArcTanh[c*x^2])^3,x]
```

output

```
(x^2*(a + b*ArcTanh[c*x^2])^3 - 3*b*c*(-1/3*(a + b*ArcTanh[c*x^2])^3/(b*c^2) + (((a + b*ArcTanh[c*x^2])^2*Log[2/(1 - c*x^2)])/c - 2*b*(-1/2*((a + b*ArcTanh[c*x^2])*PolyLog[2, 1 - 2/(1 - c*x^2)])/c + (b*PolyLog[3, 1 - 2/(1 - c*x^2)])/(4*c)))/c)/2
```

Definitions of rubi rules used

rule 6436 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot b)^p, x_Symbol] \rightarrow \text{Simp}[x \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^p, x] - \text{Simp}[b \cdot c \cdot n \cdot p \cdot \text{Int}[x^n \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2n})], x], x] /;$ $\text{FreeQ}\{a, b, c, n, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[n, 1] \ || \ \text{EqQ}[p, 1])$

rule 6454 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot b)^p \cdot x^m, x_Symbol] \rightarrow \text{Simp}[1/n \cdot \text{Subst}[\text{Int}[x^{(\text{Simplify}[m+1]/n - 1) \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p}, x], x, x^n], x] /;$ $\text{FreeQ}\{a, b, c, m, n, x\} \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[m+1]/n]$

rule 6470 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((d + e \cdot x)), x_Symbol] \rightarrow \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{Log}[2/(1 + e \cdot (x/d))]/e), x] + \text{Simp}[b \cdot c \cdot (p/e) \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{Log}[2/(1 + e \cdot (x/d))]) / (1 - c^2 \cdot x^2)], x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d^2 - e^2, 0]$

rule 6546 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p \cdot x / ((d + e \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot e \cdot (p+1)), x] + \text{Simp}[1/(c \cdot d) \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p / (1 - c \cdot x)], x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{IGtQ}[p, 0]$

rule 6620 $\text{Int}[(\text{Log}[u] \cdot (a + \text{ArcTanh}[c \cdot x] \cdot b)^p) / ((d + e \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{PolyLog}[2, 1 - u] / (2 \cdot c \cdot d)), x] + \text{Simp}[b \cdot (p/2) \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{PolyLog}[2, 1 - u] / (d + e \cdot x^2))], x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{EqQ}[(1 - u)^2 - (1 - 2/(1 - c \cdot x))^2, 0]$

rule 7164 $\text{Int}[u \cdot \text{PolyLog}[n, v], x_Symbol] \rightarrow \text{With}\{w = \text{DerivativeDivides}[v, u \cdot v, x]\}, \text{Simp}[w \cdot \text{PolyLog}[n+1, v], x] /;$ $! \text{FalseQ}[w] /;$ $\text{FreeQ}[n, x]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 264 vs. $2(124) = 248$.

Time = 1.67 (sec) , antiderivative size = 265, normalized size of antiderivative = 1.98

method	result
derivativedivides	$a^3 c x^2 + b^3 \left(\operatorname{arctanh}(c x^2)^3 (c x^2 - 1) + 2 \operatorname{arctanh}(c x^2)^3 - 3 \operatorname{arctanh}(c x^2)^2 \ln \left(1 + \frac{(c x^2 + 1)^2}{-c^2 x^4 + 1} \right) - 3 \operatorname{arctanh}(c x^2) \operatorname{polylog} \right)$
default	$a^3 c x^2 + b^3 \left(\operatorname{arctanh}(c x^2)^3 (c x^2 - 1) + 2 \operatorname{arctanh}(c x^2)^3 - 3 \operatorname{arctanh}(c x^2)^2 \ln \left(1 + \frac{(c x^2 + 1)^2}{-c^2 x^4 + 1} \right) - 3 \operatorname{arctanh}(c x^2) \operatorname{polylog} \right)$
parts	$\frac{a^3 x^2}{2} + \frac{b^3 \left(\operatorname{arctanh}(c x^2)^3 (c x^2 - 1) + 2 \operatorname{arctanh}(c x^2)^3 - 3 \operatorname{arctanh}(c x^2)^2 \ln \left(1 + \frac{(c x^2 + 1)^2}{-c^2 x^4 + 1} \right) - 3 \operatorname{arctanh}(c x^2) \operatorname{polylog} \right)}{2c}$

input `int(x*(a+b*arctanh(c*x^2))^3,x,method=_RETURNVERBOSE)`

output $\frac{1}{2} / c * (a^3 * c * x^2 + b^3 * (\operatorname{arctanh}(c * x^2)^3 * (c * x^2 - 1) + 2 * \operatorname{arctanh}(c * x^2)^3 - 3 * \operatorname{arctanh}(c * x^2)^2 * \ln(1 + (c * x^2 + 1)^2 / (-c^2 * x^4 + 1)) - 3 * \operatorname{arctanh}(c * x^2) * \operatorname{polylog}(2, -(c * x^2 + 1)^2 / (-c^2 * x^4 + 1))) + 3 / 2 * \operatorname{polylog}(3, -(c * x^2 + 1)^2 / (-c^2 * x^4 + 1))) + 3 * a * b^2 * (\operatorname{arctanh}(c * x^2)^2 * (c * x^2 - 1) + 2 * \operatorname{arctanh}(c * x^2)^2 - 2 * \operatorname{arctanh}(c * x^2) * \ln(1 + (c * x^2 + 1)^2 / (-c^2 * x^4 + 1)) - \operatorname{polylog}(2, -(c * x^2 + 1)^2 / (-c^2 * x^4 + 1))) + 3 * a^2 * b * (c * x^2 * \operatorname{arctanh}(c * x^2) + 1 / 2 * \ln(-c^2 * x^4 + 1)))$

Fricas [F]

$$\int x(a + b \operatorname{arctanh}(c x^2))^3 dx = \int (b \operatorname{arctanh}(c x^2) + a)^3 x dx$$

input `integrate(x*(a+b*arctanh(c*x^2))^3,x, algorithm="fricas")`

output

```
integral(b^3*x*arctanh(c*x^2)^3 + 3*a*b^2*x*arctanh(c*x^2)^2 + 3*a^2*b*x*arctanh(c*x^2) + a^3*x, x)
```

Sympy [F]

$$\int x(a + b \operatorname{arctanh}(cx^2))^3 dx = \int x(a + b \operatorname{atanh}(cx^2))^3 dx$$

input

```
integrate(x*(a+b*atanh(c*x**2))**3,x)
```

output

```
Integral(x*(a + b*atanh(c*x**2))**3, x)
```

Maxima [F]

$$\int x(a + b \operatorname{arctanh}(cx^2))^3 dx = \int (b \operatorname{artanh}(cx^2) + a)^3 x dx$$

input

```
integrate(x*(a+b*arctanh(c*x^2))^3,x, algorithm="maxima")
```

output

```
1/2*a^3*x^2 + 3/4*(2*c*x^2*arctanh(c*x^2) + log(-c^2*x^4 + 1))*a^2*b/c - 1/16*((b^3*c*x^2 - b^3)*log(-c*x^2 + 1)^3 - 3*(2*a*b^2*c*x^2 + (b^3*c*x^2 + b^3)*log(c*x^2 + 1))*log(-c*x^2 + 1)^2)/c - integrate(-1/8*((b^3*c*x^3 - b^3*x)*log(c*x^2 + 1)^3 + 6*(a*b^2*c*x^3 - a*b^2*x)*log(c*x^2 + 1)^2 - 3*(4*a*b^2*c*x^3 + (b^3*c*x^3 - b^3*x)*log(c*x^2 + 1)^2 + 2*((2*a*b^2*c + b^3*c)*x^3 - (2*a*b^2 - b^3)*x)*log(c*x^2 + 1))*log(-c*x^2 + 1))/(c*x^2 - 1), x)
```


Giac [F]

$$\int x(a + b \operatorname{arctanh}(cx^2))^3 dx = \int (b \operatorname{artanh}(cx^2) + a)^3 x dx$$

input `integrate(x*(a+b*arctanh(c*x^2))^3,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)^3*x, x)`

Mupad [F(-1)]

Timed out.

$$\int x(a + b \operatorname{arctanh}(cx^2))^3 dx = \int x(a + b \operatorname{atanh}(cx^2))^3 dx$$

input `int(x*(a + b*atanh(c*x^2))^3,x)`

output `int(x*(a + b*atanh(c*x^2))^3, x)`

Reduce [F]

$$\int x(a + b \operatorname{arctanh}(cx^2))^3 dx$$

$$= \frac{3 \operatorname{atanh}(cx^2) a^2 b c x^2 - 3 \operatorname{atanh}(cx^2) a^2 b + 2 \left(\int \operatorname{atanh}(cx^2)^3 x dx \right) b^3 c + 6 \left(\int \operatorname{atanh}(cx^2)^2 x dx \right) a b^2 c + 3 \int \operatorname{atanh}(cx^2) x dx a^2 b + a^3 c x^2}{2c}$$

input `int(x*(a+b*atanh(c*x^2))^3,x)`

output `(3*atanh(c*x**2)*a**2*b*c*x**2 - 3*atanh(c*x**2)*a**2*b + 2*int(atanh(c*x**2)**3*x,x)*b**3*c + 6*int(atanh(c*x**2)**2*x,x)*a*b**2*c + 3*log(c*x**2 + 1)*a**2*b + a**3*c*x**2)/(2*c)`

3.79 $\int \frac{(a+b\operatorname{arctanh}(cx^2))^3}{x} dx$

Optimal result	669
Mathematica [C] (verified)	670
Rubi [A] (verified)	671
Maple [F]	673
Fricas [F]	674
Sympy [F]	674
Maxima [F]	674
Giac [F]	675
Mupad [F(-1)]	675
Reduce [F]	675

Optimal result

Integrand size = 16, antiderivative size = 207

$$\int \frac{(a + b\operatorname{arctanh}(cx^2))^3}{x} dx = (a + b\operatorname{arctanh}(cx^2))^3 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx^2}\right) - \frac{3}{4}b(a + b\operatorname{arctanh}(cx^2))^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx^2}\right) + \frac{3}{4}b(a + b\operatorname{arctanh}(cx^2))^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - cx^2}\right) + \frac{3}{4}b^2(a + b\operatorname{arctanh}(cx^2)) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - cx^2}\right) - \frac{3}{4}b^2(a + b\operatorname{arctanh}(cx^2)) \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 - cx^2}\right) - \frac{3}{8}b^3 \operatorname{PolyLog}\left(4, 1 - \frac{2}{1 - cx^2}\right) + \frac{3}{8}b^3 \operatorname{PolyLog}\left(4, -1 + \frac{2}{1 - cx^2}\right)$$

output

```
-(a+b*arctanh(c*x^2))^3*arctanh(-1+2/(-c*x^2+1))-3/4*b*(a+b*arctanh(c*x^2))^2*polylog(2,1-2/(-c*x^2+1))+3/4*b*(a+b*arctanh(c*x^2))^2*polylog(2,-1+2/(-c*x^2+1))+3/4*b^2*(a+b*arctanh(c*x^2))*polylog(3,1-2/(-c*x^2+1))-3/4*b^2*(a+b*arctanh(c*x^2))*polylog(3,-1+2/(-c*x^2+1))-3/8*b^3*polylog(4,1-2/(-c*x^2+1))+3/8*b^3*polylog(4,-1+2/(-c*x^2+1))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 371, normalized size of antiderivative = 1.79

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x} dx = a^3 \log(x) + \frac{3}{4} a^2 b (-\operatorname{PolyLog}(2, -cx^2) + \operatorname{PolyLog}(2, cx^2))$$

$$+ \frac{3}{2} ab^2 \left(\frac{i\pi^3}{24} - \frac{2}{3} \operatorname{arctanh}(cx^2)^3 \right.$$

$$\quad - \operatorname{arctanh}(cx^2)^2 \log(1 + e^{-2\operatorname{arctanh}(cx^2)})$$

$$\quad + \operatorname{arctanh}(cx^2)^2 \log(1 - e^{2\operatorname{arctanh}(cx^2)})$$

$$+ \operatorname{arctanh}(cx^2) \operatorname{PolyLog}(2, -e^{-2\operatorname{arctanh}(cx^2)})$$

$$+ \operatorname{arctanh}(cx^2) \operatorname{PolyLog}(2, e^{2\operatorname{arctanh}(cx^2)})$$

$$\quad + \frac{1}{2} \operatorname{PolyLog}(3, -e^{-2\operatorname{arctanh}(cx^2)})$$

$$\quad \left. - \frac{1}{2} \operatorname{PolyLog}(3, e^{2\operatorname{arctanh}(cx^2)}) \right)$$

$$+ \frac{1}{128} b^3 \left(\pi^4 - 32 \operatorname{arctanh}(cx^2)^4 \right.$$

$$\quad - 64 \operatorname{arctanh}(cx^2)^3 \log(1 + e^{-2\operatorname{arctanh}(cx^2)})$$

$$\quad + 64 \operatorname{arctanh}(cx^2)^3 \log(1 - e^{2\operatorname{arctanh}(cx^2)})$$

$$+ 96 \operatorname{arctanh}(cx^2)^2 \operatorname{PolyLog}(2, -e^{-2\operatorname{arctanh}(cx^2)})$$

$$+ 96 \operatorname{arctanh}(cx^2)^2 \operatorname{PolyLog}(2, e^{2\operatorname{arctanh}(cx^2)})$$

$$+ 96 \operatorname{arctanh}(cx^2) \operatorname{PolyLog}(3, -e^{-2\operatorname{arctanh}(cx^2)})$$

$$\quad - 96 \operatorname{arctanh}(cx^2) \operatorname{PolyLog}(3, e^{2\operatorname{arctanh}(cx^2)})$$

$$\quad + 48 \operatorname{PolyLog}(4, -e^{-2\operatorname{arctanh}(cx^2)})$$

$$\quad \left. + 48 \operatorname{PolyLog}(4, e^{2\operatorname{arctanh}(cx^2)}) \right)$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])^3/x,x]
```

output

```

a^3*Log[x] + (3*a^2*b*(-PolyLog[2, -(c*x^2)] + PolyLog[2, c*x^2]))/4 + (3*
a*b^2*((I/24)*Pi^3 - (2*ArcTanh[c*x^2]^3)/3 - ArcTanh[c*x^2]^2*Log[1 + E^(
-2*ArcTanh[c*x^2]]) + ArcTanh[c*x^2]^2*Log[1 - E^(2*ArcTanh[c*x^2]]) + Arc
Tanh[c*x^2]*PolyLog[2, -E^(-2*ArcTanh[c*x^2]]) + ArcTanh[c*x^2]*PolyLog[2,
E^(2*ArcTanh[c*x^2]]) + PolyLog[3, -E^(-2*ArcTanh[c*x^2]])/2 - PolyLog[3,
E^(2*ArcTanh[c*x^2]])/2)))/2 + (b^3*(Pi^4 - 32*ArcTanh[c*x^2]^4 - 64*ArcTa
nh[c*x^2]^3*Log[1 + E^(-2*ArcTanh[c*x^2]]) + 64*ArcTanh[c*x^2]^3*Log[1 - E
^(2*ArcTanh[c*x^2]]) + 96*ArcTanh[c*x^2]^2*PolyLog[2, -E^(-2*ArcTanh[c*x^2
]]) + 96*ArcTanh[c*x^2]^2*PolyLog[2, E^(2*ArcTanh[c*x^2]]) + 96*ArcTanh[c*
x^2]*PolyLog[3, -E^(-2*ArcTanh[c*x^2]]) - 96*ArcTanh[c*x^2]*PolyLog[3, E^(
2*ArcTanh[c*x^2]]) + 48*PolyLog[4, -E^(-2*ArcTanh[c*x^2]]) + 48*PolyLog[4,
E^(2*ArcTanh[c*x^2])])))/128

```

Rubi [A] (verified)

Time = 1.32 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.16, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6450, 6448, 6614, 6620, 6624, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x} dx \\
 & \quad \downarrow \text{6450} \\
 & \frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^2} dx^2 \\
 & \quad \downarrow \text{6448} \\
 & \frac{1}{2} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^2} \right) (a + b \operatorname{arctanh}(cx^2))^3 - 6bc \int \frac{(a + b \operatorname{arctanh}(cx^2))^2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^2} \right)}{1 - c^2 x^4} dx^2 \right) \\
 & \quad \downarrow \text{6614} \\
 & \frac{1}{2} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^2} \right) (a + b \operatorname{arctanh}(cx^2))^3 - 6bc \left(\frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(cx^2))^2 \log \left(2 - \frac{2}{1 - cx^2} \right)}{1 - c^2 x^4} dx^2 - \frac{1}{2} \int \right) \right)
 \end{aligned}$$

↓ 6620

$$\frac{1}{2} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^2} \right) (a + b \operatorname{arctanh}(cx^2))^3 - 6bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - cx^2} \right) (a + b \operatorname{arctanh}(cx^2))^2}{2c} - b \right) \right) \right)$$

↓ 6624

$$\frac{1}{2} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^2} \right) (a + b \operatorname{arctanh}(cx^2))^3 - 6bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - cx^2} \right) (a + b \operatorname{arctanh}(cx^2))^2}{2c} - b \right) \right) \right)$$

↓ 7164

$$\frac{1}{2} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^2} \right) (a + b \operatorname{arctanh}(cx^2))^3 - 6bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - cx^2} \right) (a + b \operatorname{arctanh}(cx^2))^2}{2c} - b \right) \right) \right)$$

input `Int[(a + b*ArcTanh[c*x^2])^3/x,x]`

output `(2*(a + b*ArcTanh[c*x^2])^3*ArcTanh[1 - 2/(1 - c*x^2)] - 6*b*c*(((a + b*ArcTanh[c*x^2])^2*PolyLog[2, 1 - 2/(1 - c*x^2)])/(2*c) - b*(((a + b*ArcTanh[c*x^2])*PolyLog[3, 1 - 2/(1 - c*x^2)])/(2*c) - (b*PolyLog[4, 1 - 2/(1 - c*x^2)])/(4*c))))/2 + (-1/2*((a + b*ArcTanh[c*x^2])^2*PolyLog[2, -1 + 2/(1 - c*x^2)])/c + b*(((a + b*ArcTanh[c*x^2])*PolyLog[3, -1 + 2/(1 - c*x^2)])/(2*c) - (b*PolyLog[4, -1 + 2/(1 - c*x^2)])/(4*c))))/2)`

Defintions of rubi rules used

rule 6448 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_)/(x_), x_Symbol] := Simp[2*(a + b*ArcTanh[c*x])^p*ArcTanh[1 - 2/(1 - c*x)], x] - Simp[2*b*c*p Int[(a + b*ArcTanh[c*x])^(p - 1)*(ArcTanh[1 - 2/(1 - c*x)]/(1 - c^2*x^2)), x], x] /;`
`FreeQ[{a, b, c}, x] && IGtQ[p, 1]`

rule 6450 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_)/(x_), x_Symbol] := Simp[1/n Subst[Int[(a + b*ArcTanh[c*x])^p/x, x], x, x^n], x] /;`
`FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]`

rule 6614

```
Int[(ArcTanh[u_]*((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/2 Int[Log[1 + u]*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] - Simp[1/2 Int[Log[1 - u]*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[u^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 6620

```
Int[(Log[u_]*((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(-a + b*ArcTanh[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)), x] + Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 6624

```
Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*PolyLog[k_, u_])/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(PolyLog[k + 1, u]/(2*c*d)), x] - Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[k + 1, u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, k}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[u^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v, x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x} dx$$

input

```
int((a+b*arctanh(c*x^2))^3/x,x)
```

output

```
int((a+b*arctanh(c*x^2))^3/x,x)
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^3}{x} dx$$

input `integrate((a+b*arctanh(c*x^2))^3/x,x, algorithm="fricas")`

output `integral((b^3*arctanh(c*x^2)^3 + 3*a*b^2*arctanh(c*x^2)^2 + 3*a^2*b*arctanh(c*x^2) + a^3)/x, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^3}{x} dx$$

input `integrate((a+b*atanh(c*x**2))**3/x,x)`

output `Integral((a + b*atanh(c*x**2))**3/x, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^3}{x} dx$$

input `integrate((a+b*arctanh(c*x^2))^3/x,x, algorithm="maxima")`

output `a^3*log(x) + integrate(1/8*b^3*(log(c*x^2 + 1) - log(-c*x^2 + 1))^3/x + 3/4*a*b^2*(log(c*x^2 + 1) - log(-c*x^2 + 1))^2/x + 3/2*a^2*b*(log(c*x^2 + 1) - log(-c*x^2 + 1))/x, x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x} dx = \int \frac{(b \operatorname{atanh}(cx^2) + a)^3}{x} dx$$

input `integrate((a+b*arctanh(c*x^2))^3/x,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)^3/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^3}{x} dx$$

input `int((a + b*atanh(c*x^2))^3/x,x)`

output `int((a + b*atanh(c*x^2))^3/x, x)`

Reduce [F]

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x} dx &= 3 \left(\int \frac{\operatorname{atanh}(cx^2)}{x} dx \right) a^2 b + \left(\int \frac{\operatorname{atanh}(cx^2)^3}{x} dx \right) b^3 \\ &\quad + 3 \left(\int \frac{\operatorname{atanh}(cx^2)^2}{x} dx \right) a b^2 + \log(x) a^3 \end{aligned}$$

input `int((a+b*atanh(c*x^2))^3/x,x)`

output `3*int(atanh(c*x**2)/x,x)*a**2*b + int(atanh(c*x**2)**3/x,x)*b**3 + 3*int(a
tanh(c*x**2)**2/x,x)*a*b**2 + log(x)*a**3`

$$3.80 \quad \int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^3} dx$$

Optimal result	676
Mathematica [C] (verified)	677
Rubi [A] (verified)	677
Maple [F]	680
Fricas [F]	680
Sympy [F]	680
Maxima [F]	681
Giac [F]	681
Mupad [F(-1)]	681
Reduce [F]	682

Optimal result

Integrand size = 16, antiderivative size = 125

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^3} dx &= \frac{1}{2}c(a + b \operatorname{arctanh}(cx^2))^3 - \frac{(a + b \operatorname{arctanh}(cx^2))^3}{2x^2} \\ &\quad + \frac{3}{2}bc(a + b \operatorname{arctanh}(cx^2))^2 \log\left(2 - \frac{2}{1 + cx^2}\right) \\ &\quad - \frac{3}{2}b^2c(a + b \operatorname{arctanh}(cx^2)) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + cx^2}\right) \\ &\quad - \frac{3}{4}b^3c \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 + cx^2}\right) \end{aligned}$$

output

```
1/2*c*(a+b*arctanh(c*x^2))^3-1/2*(a+b*arctanh(c*x^2))^3/x^2+3/2*b*c*(a+b*arctanh(c*x^2))^2*ln(2-2/(c*x^2+1))-3/2*b^2*c*(a+b*arctanh(c*x^2))*polylog(2,-1+2/(c*x^2+1))-3/4*b^3*c*polylog(3,-1+2/(c*x^2+1))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.78

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^3} dx$$

$$= \frac{1}{4} \left(-\frac{2a^3}{x^2} - \frac{6a^2 b \operatorname{arctanh}(cx^2)}{x^2} + 12a^2 b c \log(x) - 3a^2 b c \log(1 - c^2 x^4) \right. \\ \left. + 6ab^2 c \left(\operatorname{arctanh}(cx^2) \left(\left(1 - \frac{1}{cx^2}\right) \operatorname{arctanh}(cx^2) + 2 \log(1 - e^{-2 \operatorname{arctanh}(cx^2)}) \right) \right) \right. \\ \left. - \operatorname{PolyLog}\left(2, e^{-2 \operatorname{arctanh}(cx^2)}\right) \right) + 2b^3 c \left(\frac{i\pi^3}{8} - \operatorname{arctanh}(cx^2)^3 - \frac{\operatorname{arctanh}(cx^2)^3}{cx^2} \right. \\ \left. + 3 \operatorname{arctanh}(cx^2)^2 \log(1 - e^{2 \operatorname{arctanh}(cx^2)}) \right. \\ \left. + 3 \operatorname{arctanh}(cx^2) \operatorname{PolyLog}\left(2, e^{2 \operatorname{arctanh}(cx^2)}\right) - \frac{3}{2} \operatorname{PolyLog}\left(3, e^{2 \operatorname{arctanh}(cx^2)}\right) \right) \Bigg)$$

input `Integrate[(a + b*ArcTanh[c*x^2])^3/x^3,x]`

output `((-2*a^3)/x^2 - (6*a^2*b*ArcTanh[c*x^2])/x^2 + 12*a^2*b*c*Log[x] - 3*a^2*b*c*Log[1 - c^2*x^4] + 6*a*b^2*c*(ArcTanh[c*x^2]*((1 - 1/(c*x^2))*ArcTanh[c*x^2] + 2*Log[1 - E^(-2*ArcTanh[c*x^2])])) - PolyLog[2, E^(-2*ArcTanh[c*x^2])]) + 2*b^3*c*((I/8)*Pi^3 - ArcTanh[c*x^2]^3 - ArcTanh[c*x^2]^3/(c*x^2) + 3*ArcTanh[c*x^2]^2*Log[1 - E^(2*ArcTanh[c*x^2])] + 3*ArcTanh[c*x^2]*PolyLog[2, E^(2*ArcTanh[c*x^2])] - (3*PolyLog[3, E^(2*ArcTanh[c*x^2])])/2)/4`

Rubi [A] (verified)

Time = 1.41 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6454, 6452, 6550, 6494, 6618, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^3} dx \\
& \quad \downarrow \text{6454} \\
& \frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^4} dx^2 \\
& \quad \downarrow \text{6452} \\
& \frac{1}{2} \left(3bc \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^2(1 - c^2x^4)} dx^2 - \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^2} \right) \\
& \quad \downarrow \text{6550} \\
& \frac{1}{2} \left(3bc \left(\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^2(cx^2 + 1)} dx^2 + \frac{(a + b \operatorname{arctanh}(cx^2))^3}{3b} \right) - \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^2} \right) \\
& \quad \downarrow \text{6494} \\
& \frac{1}{2} \left(3bc \left(-2bc \int \frac{(a + b \operatorname{arctanh}(cx^2)) \log\left(2 - \frac{2}{cx^2 + 1}\right)}{1 - c^2x^4} dx^2 + \frac{(a + b \operatorname{arctanh}(cx^2))^3}{3b} + \log\left(2 - \frac{2}{cx^2 + 1}\right) (a + b \operatorname{arctanh}(cx^2)) \right) \right) \\
& \quad \downarrow \text{6618} \\
& \frac{1}{2} \left(3bc \left(-2bc \left(\frac{\operatorname{PolyLog}\left(2, \frac{2}{cx^2 + 1} - 1\right) (a + b \operatorname{arctanh}(cx^2))}{2c} - \frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, \frac{2}{cx^2 + 1} - 1\right)}{1 - c^2x^4} dx^2 \right) + \frac{(a + b \operatorname{arctanh}(cx^2))^3}{3b} + \log\left(2 - \frac{2}{cx^2 + 1}\right) (a + b \operatorname{arctanh}(cx^2)) \right) \right) \\
& \quad \downarrow \text{7164} \\
& \frac{1}{2} \left(3bc \left(-2bc \left(\frac{\operatorname{PolyLog}\left(2, \frac{2}{cx^2 + 1} - 1\right) (a + b \operatorname{arctanh}(cx^2))}{2c} + \frac{b \operatorname{PolyLog}\left(3, \frac{2}{cx^2 + 1} - 1\right)}{4c} \right) + \frac{(a + b \operatorname{arctanh}(cx^2))^3}{3b} + \log\left(2 - \frac{2}{cx^2 + 1}\right) (a + b \operatorname{arctanh}(cx^2)) \right) \right)
\end{aligned}$$

input `Int[(a + b*ArcTanh[c*x^2])^3/x^3,x]`

output `(-((a + b*ArcTanh[c*x^2])^3/x^2) + 3*b*c*((a + b*ArcTanh[c*x^2])^3/(3*b) + (a + b*ArcTanh[c*x^2])^2*Log[2 - 2/(1 + c*x^2)] - 2*b*c*((a + b*ArcTanh[c*x^2])*PolyLog[2, -1 + 2/(1 + c*x^2)]/(2*c) + (b*PolyLog[3, -1 + 2/(1 + c*x^2)]/(4*c))))/2`

Definitions of rubi rules used

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]`

rule 6494 `Int[((a_.) + ArcTanh[(c_.)*(x_)*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_))), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6550 `Int[((a_.) + ArcTanh[(c_.)*(x_)*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_)^2)), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]`

rule 6618 `Int[(Log[u_]*((a_.) + ArcTanh[(c_.)*(x_)*(b_.))^(p_.))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)), x] - Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 + c*x))^2, 0]`

rule 7164 `Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v, x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]`

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^3} dx$$

input `int((a+b*arctanh(c*x^2))^3/x^3,x)`

output `int((a+b*arctanh(c*x^2))^3/x^3,x)`

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^3} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^3}{x^3} dx$$

input `integrate((a+b*arctanh(c*x^2))^3/x^3,x, algorithm="fricas")`

output `integral((b^3*arctanh(c*x^2)^3 + 3*a*b^2*arctanh(c*x^2)^2 + 3*a^2*b*arctanh(c*x^2) + a^3)/x^3, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^3} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^3}{x^3} dx$$

input `integrate((a+b*atanh(c*x**2))**3/x**3,x)`

output `Integral((a + b*atanh(c*x**2))**3/x**3, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^3} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^3}{x^3} dx$$

input `integrate((a+b*arctanh(c*x^2))^3/x^3,x, algorithm="maxima")`

output `-3/4*(c*(log(c^2*x^4 - 1) - log(x^4)) + 2*arctanh(c*x^2)/x^2)*a^2*b - 1/2*a^3/x^2 - 1/16*((b^3*c*x^2 - b^3)*log(-c*x^2 + 1)^3 + 3*(2*a*b^2 + (b^3*c*x^2 + b^3)*log(c*x^2 + 1))*log(-c*x^2 + 1)^2)/x^2 - integrate(-1/8*((b^3*c*x^2 - b^3)*log(c*x^2 + 1)^3 + 6*(a*b^2*c*x^2 - a*b^2)*log(c*x^2 + 1)^2 + 3*(4*a*b^2*c*x^2 - (b^3*c*x^2 - b^3)*log(c*x^2 + 1)^2 + 2*(b^3*c^2*x^4 + 2*a*b^2 - (2*a*b^2*c - b^3*c)*x^2)*log(c*x^2 + 1))*log(-c*x^2 + 1))/(c*x^5 - x^3), x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^3} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^3}{x^3} dx$$

input `integrate((a+b*arctanh(c*x^2))^3/x^3,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)^3/x^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^3} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^3}{x^3} dx$$

input `int((a + b*atanh(c*x^2))^3/x^3,x)`

output `int((a + b*atanh(c*x^2))^3/x^3, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{atanh}(cx^2))^3}{x^3} dx$$

$$= \frac{-\operatorname{atanh}(cx^2)^3 b^3 - 3\operatorname{atanh}(cx^2)^2 a b^2 + 3\operatorname{atanh}(cx^2) a^2 b c x^2 - 3\operatorname{atanh}(cx^2) a^2 b - 12 \left(\int \frac{\operatorname{atanh}(cx^2)}{c^2 x^5 - x} dx \right) a b}{2x^2}$$

input `int((a+b*atanh(c*x^2))^3/x^3,x)`

output `(- atanh(c*x**2)**3*b**3 - 3*atanh(c*x**2)**2*a*b**2 + 3*atanh(c*x**2)*a*
*2*b*c*x**2 - 3*atanh(c*x**2)*a**2*b - 12*int(atanh(c*x**2)/(c**2*x**5 - x
,x)*a*b**2*c*x**2 - 6*int(atanh(c*x**2)**2/(c**2*x**5 - x),x)*b**3*c*x**2
- 3*log(c*x**2 + 1)*a**2*b*c*x**2 + 6*log(x)*a**2*b*c*x**2 - a**3)/(2*x**
2)`

$$3.81 \quad \int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^5} dx$$

Optimal result	683
Mathematica [A] (verified)	684
Rubi [A] (verified)	684
Maple [F]	687
Fricas [F]	687
Sympy [F]	687
Maxima [F]	688
Giac [F]	688
Mupad [F(-1)]	689
Reduce [F]	689

Optimal result

Integrand size = 16, antiderivative size = 139

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^5} dx = & \frac{3}{4}bc^2(a + b \operatorname{arctanh}(cx^2))^2 - \frac{3bc(a + b \operatorname{arctanh}(cx^2))^2}{4x^2} \\ & + \frac{1}{4}c^2(a + b \operatorname{arctanh}(cx^2))^3 - \frac{(a + b \operatorname{arctanh}(cx^2))^3}{4x^4} \\ & + \frac{3}{2}b^2c^2(a + b \operatorname{arctanh}(cx^2)) \log\left(2 - \frac{2}{1 + cx^2}\right) \\ & - \frac{3}{4}b^3c^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + cx^2}\right) \end{aligned}$$

output

```
3/4*b*c^2*(a+b*arctanh(c*x^2))^2-3/4*b*c*(a+b*arctanh(c*x^2))^2/x^2+1/4*c^
2*(a+b*arctanh(c*x^2))^3-1/4*(a+b*arctanh(c*x^2))^3/x^4+3/2*b^2*c^2*(a+b*a
rctanh(c*x^2))*ln(2-2/(c*x^2+1))-3/4*b^3*c^2*polylog(2,-1+2/(c*x^2+1))
```


Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.57

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^5} dx$$

$$= \frac{6b^2(-1 + cx^2)(a + acx^2 + bcx^2) \operatorname{arctanh}(cx^2)^2 + 2b^3(-1 + c^2x^4) \operatorname{arctanh}(cx^2)^3 - 6b \operatorname{arctanh}(cx^2) (a^2 +$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])^3/x^5,x]
```

output

```
(6*b^2*(-1 + c*x^2)*(a + a*c*x^2 + b*c*x^2)*ArcTanh[c*x^2]^2 + 2*b^3*(-1 + c^2*x^4)*ArcTanh[c*x^2]^3 - 6*b*ArcTanh[c*x^2]*(a^2 + 2*a*b*c*x^2 - 2*b^2*c^2*x^4*Log[1 - E^(-2*ArcTanh[c*x^2])]) + a*(-2*a^2 - 6*a*b*c*x^2 - 3*a*b*c^2*x^4*Log[1 - c*x^2] + 3*a*b*c^2*x^4*Log[1 + c*x^2] + 12*b^2*c^2*x^4*Log[(c*x^2)/Sqrt[1 - c^2*x^4]]) - 6*b^3*c^2*x^4*PolyLog[2, E^(-2*ArcTanh[c*x^2])])/(8*x^4)
```

Rubi [A] (verified)

Time = 1.69 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.99, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6454, 6452, 6544, 6452, 6510, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^5} dx$$

$$\downarrow 6454$$

$$\frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^6} dx^2$$

$$\downarrow 6452$$

$$\frac{1}{2} \left(\frac{3}{2} bc \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^4(1 - c^2x^4)} dx^2 - \frac{(a + b \operatorname{arctanh}(cx^2))^3}{2x^4} \right)$$

↓ 6544

$$\frac{1}{2} \left(\frac{3}{2} bc \left(c^2 \int \frac{(a + \operatorname{arctanh}(cx^2))^2}{1 - c^2x^4} dx^2 + \int \frac{(a + \operatorname{arctanh}(cx^2))^2}{x^4} dx^2 \right) - \frac{(a + \operatorname{arctanh}(cx^2))^3}{2x^4} \right)$$

↓ 6452

$$\frac{1}{2} \left(\frac{3}{2} bc \left(c^2 \int \frac{(a + \operatorname{arctanh}(cx^2))^2}{1 - c^2x^4} dx^2 + 2bc \int \frac{a + \operatorname{arctanh}(cx^2)}{x^2(1 - c^2x^4)} dx^2 - \frac{(a + \operatorname{arctanh}(cx^2))^2}{x^2} \right) - \frac{(a + \operatorname{arctanh}(cx^2))^3}{2x^4} \right)$$

↓ 6510

$$\frac{1}{2} \left(\frac{3}{2} bc \left(2bc \int \frac{a + \operatorname{arctanh}(cx^2)}{x^2(1 - c^2x^4)} dx^2 + \frac{c(a + \operatorname{arctanh}(cx^2))^3}{3b} - \frac{(a + \operatorname{arctanh}(cx^2))^2}{x^2} \right) - \frac{(a + \operatorname{arctanh}(cx^2))^3}{2x^4} \right)$$

↓ 6550

$$\frac{1}{2} \left(\frac{3}{2} bc \left(2bc \left(\int \frac{a + \operatorname{arctanh}(cx^2)}{x^2(cx^2 + 1)} dx^2 + \frac{(a + \operatorname{arctanh}(cx^2))^2}{2b} \right) + \frac{c(a + \operatorname{arctanh}(cx^2))^3}{3b} - \frac{(a + \operatorname{arctanh}(cx^2))^2}{x^2} \right) \right)$$

↓ 6494

$$\frac{1}{2} \left(\frac{3}{2} bc \left(2bc \left(-bc \int \frac{\log\left(2 - \frac{2}{cx^2 + 1}\right)}{1 - c^2x^4} dx^2 + \frac{(a + \operatorname{arctanh}(cx^2))^2}{2b} + \log\left(2 - \frac{2}{cx^2 + 1}\right) (a + \operatorname{arctanh}(cx^2)) \right) \right)$$

↓ 2897

$$\frac{1}{2} \left(\frac{3}{2} bc \left(2bc \left(\frac{(a + \operatorname{arctanh}(cx^2))^2}{2b} + \log\left(2 - \frac{2}{cx^2 + 1}\right) (a + \operatorname{arctanh}(cx^2)) - \frac{1}{2} b \operatorname{PolyLog}\left(2, \frac{2}{cx^2 + 1} - 1\right) \right) \right)$$

input `Int[(a + b*ArcTanh[c*x^2])^3/x^5,x]`

output `(-1/2*(a + b*ArcTanh[c*x^2])^3/x^4 + (3*b*c*(-((a + b*ArcTanh[c*x^2])^2/x^2) + (c*(a + b*ArcTanh[c*x^2])^3)/(3*b) + 2*b*c*((a + b*ArcTanh[c*x^2])^2/(2*b) + (a + b*ArcTanh[c*x^2])*Log[2 - 2/(1 + c*x^2)] - (b*PolyLog[2, -1 + 2/(1 + c*x^2)]/2)))/2)/2`

Definitions of rubi rules used

rule 2897 `Int[Log[u_]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6494 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((x_)*((d_) + (e_)*(x_))), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6510 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6544 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)*((f_)*(x_)^(m_))/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]`

rule 6550

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)/((x_.)*((d_.) + (e_.)*(x_)^2)),
  x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/
d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]
```

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^5} dx$$

input

```
int((a+b*arctanh(c*x^2))^3/x^5,x)
```

output

```
int((a+b*arctanh(c*x^2))^3/x^5,x)
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^5} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^3}{x^5} dx$$

input

```
integrate((a+b*arctanh(c*x^2))^3/x^5,x, algorithm="fricas")
```

output

```
integral((b^3*arctanh(c*x^2)^3 + 3*a*b^2*arctanh(c*x^2)^2 + 3*a^2*b*arctan
h(c*x^2) + a^3)/x^5, x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^5} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^3}{x^5} dx$$

input

```
integrate((a+b*atanh(c*x**2))**3/x**5,x)
```

output `Integral((a + b*atanh(c*x**2))**3/x**5, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^5} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^3}{x^5} dx$$

input `integrate((a+b*arctanh(c*x^2))^3/x^5,x, algorithm="maxima")`

output `3/8*((c*log(c*x^2 + 1) - c*log(c*x^2 - 1) - 2/x^2)*c - 2*arctanh(c*x^2)/x^4)*a^2*b + 3/16*((2*(log(c*x^2 - 1) - 2)*log(c*x^2 + 1) - log(c*x^2 + 1)^2 - log(c*x^2 - 1)^2 - 4*log(c*x^2 - 1) + 16*log(x))*c^2 + 4*(c*log(c*x^2 + 1) - c*log(c*x^2 - 1) - 2/x^2)*c*arctanh(c*x^2))*a*b^2 - 1/32*b^3*(((c^2*x^4 - 1)*log(-c*x^2 + 1)^3 + 3*(2*c*x^2 - (c^2*x^4 - 1)*log(c*x^2 + 1))*log(-c*x^2 + 1)^2)/x^4 + 4*integrate(-((c*x^2 - 1)*log(c*x^2 + 1)^3 + 3*(2*c^2*x^4 - (c*x^2 - 1)*log(c*x^2 + 1)^2 - (c^3*x^6 - c*x^2)*log(c*x^2 + 1))*log(-c*x^2 + 1))/(c*x^7 - x^5), x)) - 3/4*a*b^2*arctanh(c*x^2)^2/x^4 - 1/4*a^3/x^4`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^5} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^3}{x^5} dx$$

input `integrate((a+b*arctanh(c*x^2))^3/x^5,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)^3/x^5, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^5} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^3}{x^5} dx$$

input `int((a + b*atanh(c*x^2))^3/x^5,x)`output `int((a + b*atanh(c*x^2))^3/x^5, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^3}{x^5} dx$$

$$= \frac{\operatorname{atanh}(cx^2)^3 b^3 c^2 x^4 - \operatorname{atanh}(cx^2)^3 b^3 + 3 \operatorname{atanh}(cx^2)^2 a b^2 c^2 x^4 - 3 \operatorname{atanh}(cx^2)^2 a b^2 - 3 \operatorname{atanh}(cx^2)^2 b^3 c x^4}{4 x^4}$$

input `int((a+b*atanh(c*x^2))^3/x^5,x)`output `(atanh(c*x**2)**3*b**3*c**2*x**4 - atanh(c*x**2)**3*b**3 + 3*atanh(c*x**2)**2*a*b**2*c**2*x**4 - 3*atanh(c*x**2)**2*a*b**2 - 3*atanh(c*x**2)**2*b**3*c*x**4 + 3*atanh(c*x**2)*a**2*b*c**2*x**4 - 3*atanh(c*x**2)*a**2*b + 6*atanh(c*x**2)*a*b**2*c**2*x**4 - 6*atanh(c*x**2)*a*b**2*c*x**2 - 3*atanh(c*x**2)*b**3*c**2*x**4 + 3*atanh(c*x**2)*b**3 - 12*int(atanh(c*x**2)/(c**2*x**9 - x**5),x)*b**3*x**4 - 6*log(c*x**2 + 1)*a*b**2*c**2*x**4 + 12*log(x)*a*b**2*c**2*x**4 - a**3 - 3*a**2*b*c*x**2 + 3*b**3*c*x**2)/(4*x**4)`

3.82 $\int (dx)^{5/2} (a + b \operatorname{arctanh}(cx^2)) dx$

Optimal result	690
Mathematica [A] (verified)	691
Rubi [A] (verified)	691
Maple [A] (verified)	699
Fricas [C] (verification not implemented)	701
Sympy [F(-1)]	702
Maxima [A] (verification not implemented)	702
Giac [F]	703
Mupad [F(-1)]	703
Reduce [B] (verification not implemented)	704

Optimal result

Integrand size = 18, antiderivative size = 257

$$\begin{aligned} \int (dx)^{5/2} (a + b \operatorname{arctanh}(cx^2)) dx &= \frac{8bd(dx)^{3/2}}{21c} \\ &+ \frac{2bd^{5/2} \arctan\left(\frac{\sqrt[4]{c\sqrt{dx}}}{\sqrt{d}}\right)}{7c^{7/4}} + \frac{\sqrt{2}bd^{5/2} \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{c\sqrt{dx}}}{\sqrt{d}}\right)}{7c^{7/4}} \\ &- \frac{\sqrt{2}bd^{5/2} \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{c\sqrt{dx}}}{\sqrt{d}}\right)}{7c^{7/4}} + \frac{2(dx)^{7/2} (a + b \operatorname{arctanh}(cx^2))}{7d} \\ &- \frac{2bd^{5/2} \operatorname{arctanh}\left(\frac{\sqrt[4]{c\sqrt{dx}}}{\sqrt{d}}\right)}{7c^{7/4}} + \frac{\sqrt{2}bd^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{c\sqrt{dx}}}{\sqrt{d}(1+\sqrt{cx})}\right)}{7c^{7/4}} \end{aligned}$$

output

```
8/21*b*d*(d*x)^(3/2)/c+2/7*b*d^(5/2)*arctan(c^(1/4)*(d*x)^(1/2)/d^(1/2))/c
^(7/4)-1/7*2^(1/2)*b*d^(5/2)*arctan(-1+2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/2)
)/c^(7/4)-1/7*2^(1/2)*b*d^(5/2)*arctan(1+2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/
2))/c^(7/4)+2/7*(d*x)^(7/2)*(a+b*arctanh(c*x^2))/d-2/7*b*d^(5/2)*arctanh(c
^(1/4)*(d*x)^(1/2)/d^(1/2))/c^(7/4)+1/7*2^(1/2)*b*d^(5/2)*arctanh(2^(1/2)*
c^(1/4)*(d*x)^(1/2)/d^(1/2)/(1+c^(1/2)*x))/c^(7/4)
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 241, normalized size of antiderivative = 0.94

$$\int (dx)^{5/2} (a + b \operatorname{arctanh}(cx^2)) dx = \frac{(dx)^{5/2} (16bc^{3/4}x^{3/2} + 12ac^{7/4}x^{7/2} + 6\sqrt{2}b \arctan(1 - \sqrt{2}\sqrt[4]{c}\sqrt{x}) - 6\sqrt{2}b \arctan(1 + \sqrt{2}\sqrt[4]{c}\sqrt{x}))}{42c^{7/4}x^{5/2}}$$

input

```
Integrate[(d*x)^(5/2)*(a + b*ArcTanh[c*x^2]),x]
```

output

```
((d*x)^(5/2)*(16*b*c^(3/4)*x^(3/2) + 12*a*c^(7/4)*x^(7/2) + 6*Sqrt[2]*b*ArcTan[1 - Sqrt[2]*c^(1/4)*Sqrt[x]] - 6*Sqrt[2]*b*ArcTan[1 + Sqrt[2]*c^(1/4)*Sqrt[x]] + 12*b*ArcTan[c^(1/4)*Sqrt[x]] + 12*b*c^(7/4)*x^(7/2)*ArcTanh[c*x^2] + 6*b*Log[1 - c^(1/4)*Sqrt[x]] - 6*b*Log[1 + c^(1/4)*Sqrt[x]] - 3*Sqrt[2]*b*Log[1 - Sqrt[2]*c^(1/4)*Sqrt[x] + Sqrt[c]*x] + 3*Sqrt[2]*b*Log[1 + Sqrt[2]*c^(1/4)*Sqrt[x] + Sqrt[c]*x]))/(42*c^(7/4)*x^(5/2))
```

Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 356, normalized size of antiderivative = 1.39, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.889$, Rules used = {6464, 843, 851, 27, 829, 826, 827, 218, 221, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^{5/2} (a + b \operatorname{arctanh}(cx^2)) dx$$

$$\downarrow 6464$$

$$\frac{2(dx)^{7/2} (a + b \operatorname{arctanh}(cx^2))}{7d} - \frac{4bc \int \frac{(dx)^{9/2}}{1-c^2x^4} dx}{7d^2}$$

$$\downarrow 843$$

$$\frac{2(dx)^{7/2} (a + b \operatorname{arctanh}(cx^2))}{7d} - \frac{4bc \left(\frac{d^4 \int \frac{\sqrt{dx}}{1-c^2x^4} dx}{c^2} - \frac{2d^3(dx)^{3/2}}{3c^2} \right)}{7d^2}$$

$$\begin{aligned}
 & \downarrow 851 \\
 & \frac{2(dx)^{7/2} (a + \operatorname{barctanh}(cx^2))}{7d} - \frac{4bc \left(\frac{2d^3 \int \frac{d^5 x}{d^4 - c^2 d^4 x^4} d\sqrt{dx} - \frac{2d^3(dx)^{3/2}}{3c^2} \right)}{7d^2} \\
 & \downarrow 27 \\
 & \frac{2(dx)^{7/2} (a + \operatorname{barctanh}(cx^2))}{7d} - \frac{4bc \left(\frac{2d^7 \int \frac{dx}{d^4 - c^2 d^4 x^4} d\sqrt{dx} - \frac{2d^3(dx)^{3/2}}{3c^2} \right)}{7d^2} \\
 & \downarrow 829 \\
 & \frac{2(dx)^{7/2} (a + \operatorname{barctanh}(cx^2))}{7d} - \frac{4bc \left(\frac{2d^7 \left(\frac{\int \frac{dx}{d^2 - cd^2 x^2} d\sqrt{dx}}{2d^2} + \frac{\int \frac{dx}{cx^2 d^2 + d^2} d\sqrt{dx}}{2d^2} \right)}{c^2} - \frac{2d^3(dx)^{3/2}}{3c^2} \right)}{7d^2} \\
 & \downarrow 826 \\
 & \frac{2(dx)^{7/2} (a + \operatorname{barctanh}(cx^2))}{7d} - \frac{4bc \left(\frac{2d^7 \left(\frac{\int \frac{dx}{d^2 - cd^2 x^2} d\sqrt{dx}}{2d^2} + \frac{\frac{\int \frac{\sqrt{cx}d+d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{c}dx}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}}}{2d^2} \right)}{c^2} - \frac{2d^3(dx)^{3/2}}{3c^2} \right)}{7d^2} \\
 & \downarrow 827 \\
 & \frac{2(dx)^{7/2} (a + \operatorname{barctanh}(cx^2))}{7d} - \frac{4bc \left(\frac{2d^7 \left(\frac{\frac{\int \frac{\sqrt{cx}d+d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{c}dx}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}}}{2d^2} + \frac{\frac{\int \frac{1}{d-\sqrt{c}dx} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{1}{\sqrt{cx}d+d} d\sqrt{dx}}{2\sqrt{c}}}{2d^2} \right)}{c^2} - \frac{2d^3(dx)^{3/2}}{3c^2} \right)}{7d^2} \\
 & \downarrow 218
 \end{aligned}$$

$$\begin{array}{c}
 \frac{2(dx)^{7/2} (a + \operatorname{barctanh}(cx^2))}{7d} - \\
 \left(\frac{2d^7 \left(\frac{\int \frac{1}{d-\sqrt{cdx}} d\sqrt{dx}}{2\sqrt{c}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} + \frac{\int \frac{\sqrt{cdx+d}}{cx^2d^2+d^2} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2\sqrt{c}} \right)}{c^2} \right) - \frac{2d^3(dx)^{3/2}}{3c^2} \\
 \hline
 7d^2 \\
 \downarrow \text{221} \\
 \frac{2(dx)^{7/2} (a + \operatorname{barctanh}(cx^2))}{7d} - \\
 \left(\frac{2d^7 \left(\frac{\int \frac{\sqrt{cdx+d}}{cx^2d^2+d^2} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2\sqrt{c}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} \right)}{c^2} \right) - \frac{2d^3(dx)^{3/2}}{3c^2} \\
 \hline
 7d^2 \\
 \downarrow \text{1476} \\
 \frac{2(dx)^{7/2} (a + \operatorname{barctanh}(cx^2))}{7d} - \\
 \left(\frac{2d^7 \left(\frac{\int \frac{1}{xd+\frac{d}{\sqrt{c}}-\sqrt{2}\sqrt{dx}\sqrt{d}} d\sqrt{dx}}{2\sqrt{c}} + \frac{\int \frac{1}{xd+\frac{d}{\sqrt{c}}+\sqrt{2}\sqrt{dx}\sqrt{d}} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2\sqrt{c}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} \right)}{c^2} \right) - \frac{2d^3(dx)^{3/2}}{3c^2} \\
 \hline
 7d^2 \\
 \downarrow \text{1082}
 \end{array}$$

$$\frac{2(dx)^{7/2} (a + \operatorname{arctanh}(cx^2))}{c^2} - \frac{7d}{c^2} \left(\frac{\int \frac{1}{-dx-1} d \left(1 - \frac{\sqrt{2} \sqrt[4]{c} \sqrt{dx}}{\sqrt{d}} \right)}{\sqrt{2} \sqrt[4]{c} \sqrt{d}} - \frac{\int \frac{1}{-dx-1} d \left(\frac{\sqrt{2} \sqrt[4]{c} \sqrt{dx}}{\sqrt{d}} + 1 \right)}{\sqrt{2} \sqrt[4]{c} \sqrt{d}} - \frac{\int \frac{d - \sqrt{c} dx}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} + \frac{\operatorname{arctanh} \left(\frac{\sqrt[4]{c} \sqrt{dx}}{\sqrt{d}} \right)}{2c^{3/4} \sqrt{d}} - \frac{\operatorname{arctan} \left(\frac{\sqrt[4]{c} \sqrt{dx}}{\sqrt{d}} \right)}{2c^{3/4} \sqrt{d}} \right)$$

7d²

217

$$\frac{2(dx)^{7/2} (a + \operatorname{arctanh}(cx^2))}{c^2} - \frac{7d}{c^2} \left(\frac{\operatorname{arctan} \left(\frac{\sqrt{2} \sqrt[4]{c} \sqrt{dx}}{\sqrt{d}} + 1 \right)}{\sqrt{2} \sqrt[4]{c} \sqrt{d}} - \frac{\operatorname{arctan} \left(1 - \frac{\sqrt{2} \sqrt[4]{c} \sqrt{dx}}{\sqrt{d}} \right)}{\sqrt{2} \sqrt[4]{c} \sqrt{d}} - \frac{\int \frac{d - \sqrt{c} dx}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} + \frac{\operatorname{arctanh} \left(\frac{\sqrt[4]{c} \sqrt{dx}}{\sqrt{d}} \right)}{2c^{3/4} \sqrt{d}} - \frac{\operatorname{arctan} \left(\frac{\sqrt[4]{c} \sqrt{dx}}{\sqrt{d}} \right)}{2c^{3/4} \sqrt{d}} \right) - \frac{2d^3(dx)^{3/2}}{3c^2}$$

7d²

1479

$$\frac{2(dx)^{7/2} (a + \operatorname{barctanh}(cx^2))}{7d} -$$

$$\left(\frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx} + 1}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\operatorname{arctan}\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{\sqrt{2}\sqrt{d} - 2\sqrt[4]{c}\sqrt{dx}}{\sqrt[4]{c}\left(xd + \frac{d}{\sqrt{c}} - \frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{\sqrt{2}(\sqrt{d} + \sqrt{2}\sqrt[4]{c}\sqrt{dx})}{\sqrt[4]{c}\left(xd + \frac{d}{\sqrt{c}} + \frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} \right) \frac{2d^7}{2\sqrt{c}}$$

$$\frac{4bc}{c^2}$$

7d²

25

$$\frac{2(dx)^{7/2} (a + \operatorname{barctanh}(cx^2))}{7d} -$$

$$\left(\frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx} + 1}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\operatorname{arctan}\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{\sqrt{2}\sqrt{d} - 2\sqrt[4]{c}\sqrt{dx}}{\sqrt[4]{c}\left(xd + \frac{d}{\sqrt{c}} - \frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} + \frac{\int \frac{\sqrt{2}(\sqrt{d} + \sqrt{2}\sqrt[4]{c}\sqrt{dx})}{\sqrt[4]{c}\left(xd + \frac{d}{\sqrt{c}} + \frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} \right) \frac{2d^7}{2\sqrt{c}}$$

$$\frac{4bc}{c^2}$$

7d²

27

$$\frac{2(dx)^{7/2} (a + b \operatorname{arctanh}(cx^2))}{7d} -$$

$$\frac{2d^7 \left(\frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}} + 1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\operatorname{arctan}\left(1 - \frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\int \frac{\sqrt{2}\sqrt{d} - 2\sqrt[4]{C}\sqrt{dx}}{xd + \frac{d}{\sqrt{c}} - \frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt{c}}} d\sqrt{dx}}{2\sqrt{2}\sqrt{c}\sqrt{d}} + \frac{\int \frac{\sqrt{d} + \sqrt{2}\sqrt[4]{C}\sqrt{dx}}{xd + \frac{d}{\sqrt{c}} + \frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt{c}}} d\sqrt{dx}}{2\sqrt{c}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} \right)}{4bc} = \frac{7d^2}{c^2}$$

7d²

1103

$$\frac{2(dx)^{7/2} (a + b \operatorname{arctanh}(cx^2))}{7d} -$$

$$\frac{2d^7 \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} + \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}} + 1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\operatorname{arctan}\left(1 - \frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\log\left(\sqrt{cdx} + \sqrt{2}\sqrt[4]{C}\sqrt{d}\sqrt{dx} + d\right)}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\log\left(\sqrt{cdx} - \sqrt{2}\sqrt[4]{C}\sqrt{d}\sqrt{dx} + d\right)}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right)}{4bc} = \frac{7d^2}{c^2}$$

7d²

input `Int[(d*x)^(5/2)*(a + b*ArcTanh[c*x^2]),x]`

output

$$\begin{aligned} & (2*(d*x)^{(7/2)}*(a + b*\text{ArcTanh}[c*x^2]))/(7*d) - (4*b*c*((-2*d^3*(d*x)^{(3/2)}) \\ &)/(3*c^2) + (2*d^7*((-1/2*\text{ArcTan}[(c^{(1/4)}*\text{Sqrt}[d*x])/ \text{Sqrt}[d]]/(c^{(3/4)}*\text{Sqr} \\ & \text{t}[d]) + \text{ArcTanh}[(c^{(1/4)}*\text{Sqrt}[d*x])/ \text{Sqrt}[d]]/(2*c^{(3/4)}*\text{Sqrt}[d]))/(2*d^2) \\ & + ((-\text{ArcTan}[1 - (\text{Sqrt}[2]*c^{(1/4)}*\text{Sqrt}[d*x])/ \text{Sqrt}[d]]/(\text{Sqrt}[2]*c^{(1/4)}*\text{Sqr} \\ & \text{t}[d])) + \text{ArcTan}[1 + (\text{Sqrt}[2]*c^{(1/4)}*\text{Sqrt}[d*x])/ \text{Sqrt}[d]]/(\text{Sqrt}[2]*c^{(1/4)}* \\ & \text{Sqrt}[d]))/(2*\text{Sqrt}[c]) - (-1/2*\text{Log}[d + \text{Sqrt}[c]*d*x - \text{Sqrt}[2]*c^{(1/4)}*\text{Sqrt}[d] \\ &]*\text{Sqrt}[d*x])/(\text{Sqrt}[2]*c^{(1/4)}*\text{Sqrt}[d]) + \text{Log}[d + \text{Sqrt}[c]*d*x + \text{Sqrt}[2]*c^{(1/4)} \\ &]*\text{Sqrt}[d]*\text{Sqrt}[d*x])/ (2*\text{Sqrt}[2]*c^{(1/4)}*\text{Sqrt}[d]))/(2*\text{Sqrt}[c]))/(2*d^2) \\ &)/c^2)/(7*d^2) \end{aligned}$$

Defintions of rubi rules used

rule 25

$$\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$$

rule 27

$$\text{Int}[(a_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] /; \text{FreeQ}[a, \text{x}] \&\& \text{!Ma} \\ \text{tchQ}[\text{Fx}, (b_)*(\text{Gx}_) /; \text{FreeQ}[b, \text{x}]]$$

rule 217

$$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)} \\ (-1)*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& \text{PosQ}[a/b] \& \\ \& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])$$

rule 218

$$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{R} \\ \text{t}[a/b, 2]], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& \text{PosQ}[a/b]$$

rule 221

$$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x \\ / \text{Rt}[-a/b, 2]], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& \text{NegQ}[a/b]$$

rule 826

$$\text{Int}[(x_)^2/((a_ + (b_)*(x_)^4), \text{x_Symbol}] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, \\ 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*s) \quad \text{Int}[(r + s*x^2)/(a + b*x^ \\ 4), \text{x}], \text{x}] - \text{Simp}[1/(2*s) \quad \text{Int}[(r - s*x^2)/(a + b*x^4), \text{x}], \text{x}]] /; \text{FreeQ}[\{ \\ a, b\}, \text{x}] \&\& (\text{GtQ}[a/b, 0] \parallel (\text{PosQ}[a/b] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \\ \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$$

rule 827 $\text{Int}[(x_)^2/((a_)+(b_)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[s/(2*b) \text{ Int}[1/(r + s*x^2), x], x] - \text{Simp}[s/(2*b) \text{ Int}[1/(r - s*x^2), x], x]] /; \text{FreeQ}\{a, b, x\} \&\& \text{!GtQ}[a/b, 0]$

rule 829 $\text{Int}[(x_)^m/((a_)+(b_)*(x_)^n), x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[r/(2*a) \text{ Int}[x^m/(r + s*x^{n/2}), x], x] + \text{Simp}[r/(2*a) \text{ Int}[x^m/(r - s*x^{n/2}), x], x]] /; \text{FreeQ}\{a, b, x\} \&\& \text{IGtQ}[n/4, 0] \&\& \text{IGtQ}[m, 0] \&\& \text{LtQ}[m, n/2] \&\& \text{!GtQ}[a/b, 0]$

rule 843 $\text{Int}[((c_)*(x_))^m*((a_)+(b_)*(x_)^n)^p, x_Symbol] \rightarrow \text{Simp}[c^{(n-1)}*(c*x)^{m-n+1}*((a + b*x^n)^{p+1}/(b*(m+n*p+1))), x] - \text{Simp}[a*c^n*((m-n+1)/(b*(m+n*p+1))) \text{ Int}[(c*x)^{m-n}*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, p, x\} \&\& \text{IGtQ}[n, 0] \&\& \text{GtQ}[m, n-1] \&\& \text{NeQ}[m+n*p+1, 0] \&\& \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 851 $\text{Int}[((c_)*(x_))^m*((a_)+(b_)*(x_)^n)^p, x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \text{ Subst}[\text{Int}[x^{(k*(m+1)-1)}*(a + b*(x^{k*n})/c^n)^p, x], x, (c*x)^{1/k}], x]] /; \text{FreeQ}\{a, b, c, p, x\} \&\& \text{IGtQ}[n, 0] \&\& \text{FractionQ}[m] \&\& \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 1082 $\text{Int}[((a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S\text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \|\| \text{!RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}\{a, b, c, x\}$

rule 1103 $\text{Int}[((d_)+(e_)*(x_))/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}\{a, b, c, d, e, x\} \&\& \text{EqQ}[2*c*d - b*e, 0]$

rule 1476 $\text{Int}[((d_)+(e_)*(x_)^2)/((a_)+(c_)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2*(d/e), 2]\}, \text{Simp}[e/(2*c) \text{ Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Simp}[e/(2*c) \text{ Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /; \text{FreeQ}\{a, c, d, e, x\} \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[d*e]$

rule 1479

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

rule 6464

```
Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)*((d_)*(x_)^(m_)), x_Symbol] :
> Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n
/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a,
b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]
```

Maple [A] (verified)

Time = 4.39 (sec) , antiderivative size = 256, normalized size of antiderivative = 1.00

method	result
derivativedivides	$\frac{2a(dx)^{\frac{7}{2}}}{7} + 2b \left(\frac{(dx)^{\frac{7}{2}} \operatorname{arctanh}(cx^2)}{7} - \frac{4cd^2 \left(-\frac{(dx)^{\frac{3}{2}}}{3c^2} + \frac{d^2\sqrt{2} \left(\ln \left(\frac{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx}\sqrt{2} + \sqrt{\frac{d^2}{c}} \right)}{dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx}\sqrt{2} + \sqrt{\frac{d^2}{c}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2}\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 1} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2}\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} - 1} \right)}{16c^3 \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)}{7} \right)$
default	$\frac{2a(dx)^{\frac{7}{2}}}{7} + 2b \left(\frac{(dx)^{\frac{7}{2}} \operatorname{arctanh}(cx^2)}{7} - \frac{4cd^2 \left(-\frac{(dx)^{\frac{3}{2}}}{3c^2} + \frac{d^2\sqrt{2} \left(\ln \left(\frac{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx}\sqrt{2} + \sqrt{\frac{d^2}{c}} \right)}{dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx}\sqrt{2} + \sqrt{\frac{d^2}{c}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2}\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 1} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2}\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} - 1} \right)}{16c^3 \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)}{7} \right)$
parts	$\frac{2a(dx)^{\frac{7}{2}}}{7d} + 2b \left(\frac{(dx)^{\frac{7}{2}} \operatorname{arctanh}(cx^2)}{7} - \frac{4cd^2 \left(-\frac{(dx)^{\frac{3}{2}}}{3c^2} + \frac{d^2\sqrt{2} \left(\ln \left(\frac{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx}\sqrt{2} + \sqrt{\frac{d^2}{c}} \right)}{dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx}\sqrt{2} + \sqrt{\frac{d^2}{c}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2}\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 1} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2}\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} - 1} \right)}{16c^3 \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)}{7} \right)$

input `int((d*x)^(5/2)*(a+b*arctanh(c*x^2)),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 2/d*(1/7*a*(d*x)^(7/2)+b*(1/7*(d*x)^(7/2)*\operatorname{arctanh}(c*x^2)-4/7*c*d^2*(-1/3*(\\ & d*x)^(3/2)/c^2+1/16*d^2/c^3/(d^2/c)^(1/4)*2^(1/2)*(\ln((d*x-(d^2/c)^(1/4)*(\\ & d*x)^(1/2)*2^(1/2)+(d^2/c)^(1/2)))/(d*x+(d^2/c)^(1/4)*(d*x)^(1/2)*2^(1/2)+(\\ & d^2/c)^(1/2)))+2*\operatorname{arctan}(2^(1/2)/(d^2/c)^(1/4)*(d*x)^(1/2)+1)+2*\operatorname{arctan}(2^(1 \\ & /2)/(d^2/c)^(1/4)*(d*x)^(1/2)-1))-1/8*d^2/c^3/(d^2/c)^(1/4)*(2*\operatorname{arctan}((d*x \\ &)^(1/2)/(d^2/c)^(1/4))-\ln(((d*x)^(1/2)+(d^2/c)^(1/4))/((d*x)^(1/2)-(d^2/c) \\ & ^{(1/4)})))))) \end{aligned}$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 434, normalized size of antiderivative = 1.69

$$\int (dx)^{5/2} (a + b \operatorname{arctanh}(cx^2)) dx =$$

$$3 \left(\frac{b^4 d^{10}}{c^7} \right)^{\frac{1}{4}} c \log \left(\sqrt{dx} b^3 d^7 + \left(\frac{b^4 d^{10}}{c^7} \right)^{\frac{3}{4}} c^5 \right) - 3i \left(\frac{b^4 d^{10}}{c^7} \right)^{\frac{1}{4}} c \log \left(\sqrt{dx} b^3 d^7 + i \left(\frac{b^4 d^{10}}{c^7} \right)^{\frac{3}{4}} c^5 \right) + 3i \left(\frac{b^4 d^{10}}{c^7} \right)^{\frac{1}{4}} c$$

input

```
integrate((d*x)^(5/2)*(a+b*arctanh(c*x^2)),x, algorithm="fricas")
```

output

$$\begin{aligned} & -1/21*(3*(b^4*d^10/c^7)^(1/4)*c*\log(\operatorname{sqrt}(d*x)*b^3*d^7 + (b^4*d^10/c^7)^(3/ \\ & 4)*c^5) - 3*I*(b^4*d^10/c^7)^(1/4)*c*\log(\operatorname{sqrt}(d*x)*b^3*d^7 + I*(b^4*d^10/c \\ & ^7)^(3/4)*c^5) + 3*I*(b^4*d^10/c^7)^(1/4)*c*\log(\operatorname{sqrt}(d*x)*b^3*d^7 - I*(b^4 \\ & *d^10/c^7)^(3/4)*c^5) - 3*(b^4*d^10/c^7)^(1/4)*c*\log(\operatorname{sqrt}(d*x)*b^3*d^7 - (\\ & b^4*d^10/c^7)^(3/4)*c^5) + 3*(-b^4*d^10/c^7)^(1/4)*c*\log(\operatorname{sqrt}(d*x)*b^3*d^7 \\ & + (-b^4*d^10/c^7)^(3/4)*c^5) - 3*I*(-b^4*d^10/c^7)^(1/4)*c*\log(\operatorname{sqrt}(d*x)* \\ & b^3*d^7 + I*(-b^4*d^10/c^7)^(3/4)*c^5) + 3*I*(-b^4*d^10/c^7)^(1/4)*c*\log(\operatorname{s} \\ & \operatorname{qrt}(d*x)*b^3*d^7 - I*(-b^4*d^10/c^7)^(3/4)*c^5) - 3*(-b^4*d^10/c^7)^(1/4)* \\ & c*\log(\operatorname{sqrt}(d*x)*b^3*d^7 - (-b^4*d^10/c^7)^(3/4)*c^5) - (3*b*c*d^2*x^3*\log(\\ & -(c*x^2 + 1)/(c*x^2 - 1)) + 6*a*c*d^2*x^3 + 8*b*d^2*x)*\operatorname{sqrt}(d*x))/c \end{aligned}$$

Sympy [F(-1)]

Timed out.

$$\int (dx)^{5/2} (a + b \operatorname{arctanh}(cx^2)) dx = \text{Timed out}$$

```
input integrate((d*x)**(5/2)*(a+b*atanh(c*x**2)),x)
```

output Timed out

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 316, normalized size of antiderivative = 1.23

$$\int (dx)^{5/2} (a$$

$$+ b \operatorname{arctanh}(cx^2)) dx = \frac{12 (dx)^{\frac{7}{2}} a + 12 (dx)^{\frac{7}{2}} \operatorname{arctanh}(cx^2) - \frac{3 a^6 \left(2 \sqrt{2} \operatorname{arctan} \left(\frac{\sqrt{2} \left(\sqrt{2} c^{\frac{1}{4}} \sqrt{d} + 2 \sqrt{dx} \sqrt{c} \right)}{2 \sqrt{cd}} \right) \right) + 2 \sqrt{2} \operatorname{arctan} \left(\dots \right)}{\sqrt{cd} \sqrt{c}}}{\dots}$$

```
input integrate((d*x)^(5/2)*(a+b*arctanh(c*x^2)),x, algorithm="maxima")
```

output

```
1/42*(12*(d*x)^(7/2)*a + (12*(d*x)^(7/2)*arctanh(c*x^2) - (3*d^6*(2*sqrt(2)
)*arctan(1/2*sqrt(2)*(sqrt(2)*c^(1/4)*sqrt(d) + 2*sqrt(d*x)*sqrt(c))/sqrt(
sqrt(c)*d))/(sqrt(sqrt(c)*d)*sqrt(c)) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sq
rt(2)*c^(1/4)*sqrt(d) - 2*sqrt(d*x)*sqrt(c))/sqrt(sqrt(c)*d))/(sqrt(sqrt(c)
*d)*sqrt(c)) - sqrt(2)*log(sqrt(c)*d*x + sqrt(2)*sqrt(d*x)*c^(1/4)*sqrt(d)
+ d)/(c^(3/4)*sqrt(d)) + sqrt(2)*log(sqrt(c)*d*x - sqrt(2)*sqrt(d*x)*c^(1
/4)*sqrt(d) + d)/(c^(3/4)*sqrt(d)))/c^2 - 6*d^6*(2*arctan(sqrt(d*x)*sqrt(c)
)/sqrt(sqrt(c)*d))/(sqrt(sqrt(c)*d)*sqrt(c)) + log((sqrt(d*x)*sqrt(c) - sq
rt(sqrt(c)*d))/(sqrt(d*x)*sqrt(c) + sqrt(sqrt(c)*d)))/(sqrt(sqrt(c)*d)*sq
rt(c)))/c^2 - 16*(d*x)^(3/2)*d^4/c^2)*c/d^2)*b/d
```

Giac [F]

$$\int (dx)^{5/2} (a + b \operatorname{arctanh}(cx^2)) dx = \int (dx)^{5/2} (b \operatorname{arctanh}(cx^2) + a) dx$$

input

```
integrate((d*x)^(5/2)*(a+b*arctanh(c*x^2)),x, algorithm="giac")
```

output

```
integrate((d*x)^(5/2)*(b*arctanh(c*x^2) + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int (dx)^{5/2} (a + b \operatorname{arctanh}(cx^2)) dx = \int (dx)^{5/2} (a + b \operatorname{atanh}(cx^2)) dx$$

input

```
int((d*x)^(5/2)*(a + b*atanh(c*x^2)),x)
```

output

```
int((d*x)^(5/2)*(a + b*atanh(c*x^2)), x)
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 246, normalized size of antiderivative = 0.96

$$\int (dx)^{5/2} (a + b \operatorname{arctanh}(cx^2)) dx = \frac{\sqrt{d} d^2 \left(6c^{1/4} \sqrt{2} \operatorname{atan} \left(\frac{c^{1/4} \sqrt{2} - 2\sqrt{x} \sqrt{c}}{c^{1/4} \sqrt{2}} \right) b - 6c^{1/4} \sqrt{2} \operatorname{atan} \left(\frac{c^{1/4} \sqrt{2} + 2\sqrt{x} \sqrt{c}}{c^{1/4} \sqrt{2}} \right) b + 12c^{1/4} \operatorname{atan} \left(\frac{\sqrt{x} \sqrt{c}}{c^{1/4} \sqrt{2}} \right) b \right)}{42c^{3/2}}$$

input `int((d*x)^(5/2)*(a+b*atanh(c*x^2)),x)`

output `(sqrt(d)*d**2*(6*c**(1/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*b - 6*c**(1/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*b + 12*c**(1/4)*atan((sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*b + 6*c**(1/4)*sqrt(2)*atanh(c*x**2)*b + 12*sqrt(x)*atanh(c*x**2)*b*c**2*x**3 + 3*c**(1/4)*sqrt(2)*log(c**(1/4) + sqrt(x)*sqrt(c))*b + 3*c**(1/4)*sqrt(2)*log(-c**(1/4) + sqrt(x)*sqrt(c))*b - 6*c**(1/4)*sqrt(2)*log(-sqrt(x)*c**(1/4)*sqrt(2) + sqrt(c)*x + 1)*b + 3*c**(1/4)*sqrt(2)*log(sqrt(c)*x + 1)*b - 6*c**(1/4)*log(c**(1/4) + sqrt(x)*sqrt(c))*b + 6*c**(1/4)*log(-c**(1/4) + sqrt(x)*sqrt(c))*b + 12*sqrt(x)*a*c**2*x**3 + 16*sqrt(x)*b*c*x)/(42*c**2)`

3.83 $\int (dx)^{3/2} (a + \operatorname{barctanh}(cx^2)) dx$

Optimal result	705
Mathematica [A] (verified)	706
Rubi [A] (verified)	706
Maple [A] (verified)	713
Fricas [C] (verification not implemented)	715
Sympy [F]	716
Maxima [A] (verification not implemented)	716
Giac [F]	717
Mupad [F(-1)]	717
Reduce [B] (verification not implemented)	718

Optimal result

Integrand size = 18, antiderivative size = 257

$$\int (dx)^{3/2} (a + \operatorname{barctanh}(cx^2)) dx = \frac{8bd\sqrt{dx}}{5c} - \frac{2bd^{3/2} \arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{5c^{5/4}} + \frac{\sqrt{2}bd^{3/2} \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{5c^{5/4}} - \frac{\sqrt{2}bd^{3/2} \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{5c^{5/4}} + \frac{2(dx)^{5/2} (a + \operatorname{barctanh}(cx^2))}{5d} - \frac{2bd^{3/2} \operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{5c^{5/4}} - \frac{\sqrt{2}bd^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}(1+\sqrt{cx})}\right)}{5c^{5/4}}$$

output

```
8/5*b*d*(d*x)^(1/2)/c-2/5*b*d^(3/2)*arctan(c^(1/4)*(d*x)^(1/2)/d^(1/2))/c^(5/4)-1/5*2^(1/2)*b*d^(3/2)*arctan(-1+2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/2))/c^(5/4)-1/5*2^(1/2)*b*d^(3/2)*arctan(1+2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/2))/c^(5/4)+2/5*(d*x)^(5/2)*(a+b*arctanh(c*x^2))/d-2/5*b*d^(3/2)*arctanh(c^(1/4)*(d*x)^(1/2)/d^(1/2))/c^(5/4)-1/5*2^(1/2)*b*d^(3/2)*arctanh(2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/2)/(1+c^(1/2)*x))/c^(5/4)
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 240, normalized size of antiderivative = 0.93

$$\int (dx)^{3/2} (a + b \operatorname{arctanh}(cx^2)) dx = \frac{(dx)^{3/2} (16b\sqrt{c}\sqrt{x} + 4ac^{5/4}x^{5/2} + 2\sqrt{2}b \arctan(1 - \sqrt{2}\sqrt{c}\sqrt{x}) - 2\sqrt{2}b \arctan(1 + \sqrt{2}\sqrt{c}\sqrt{x}))}{10c^{5/4}x^{3/2}}$$

input

```
Integrate[(d*x)^(3/2)*(a + b*ArcTanh[c*x^2]),x]
```

output

```
((d*x)^(3/2)*(16*b*c^(1/4)*Sqrt[x] + 4*a*c^(5/4)*x^(5/2) + 2*Sqrt[2]*b*ArcTan[1 - Sqrt[2]*c^(1/4)*Sqrt[x]] - 2*Sqrt[2]*b*ArcTan[1 + Sqrt[2]*c^(1/4)*Sqrt[x]] - 4*b*ArcTan[c^(1/4)*Sqrt[x]] + 4*b*c^(5/4)*x^(5/2)*ArcTanh[c*x^2] + 2*b*Log[1 - c^(1/4)*Sqrt[x]] - 2*b*Log[1 + c^(1/4)*Sqrt[x]] + Sqrt[2]*b*Log[1 - Sqrt[2]*c^(1/4)*Sqrt[x] + Sqrt[c]*x] - Sqrt[2]*b*Log[1 + Sqrt[2]*c^(1/4)*Sqrt[x] + Sqrt[c]*x]))/(10*c^(5/4)*x^(3/2))
```

Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 350, normalized size of antiderivative = 1.36, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {6464, 843, 851, 758, 755, 756, 218, 221, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^{3/2} (a + b \operatorname{arctanh}(cx^2)) dx$$

$$\downarrow 6464$$

$$\frac{2(dx)^{5/2} (a + b \operatorname{arctanh}(cx^2))}{5d} - \frac{4bc \int \frac{(dx)^{7/2}}{1-c^2x^4} dx}{5d^2}$$

$$\downarrow 843$$

$$\frac{2(dx)^{5/2} (a + b \operatorname{arctanh}(cx^2))}{5d} - \frac{4bc \left(\frac{d^4 \int \frac{1}{\sqrt{dx}(1-c^2x^4)} dx}{c^2} - \frac{2d^3\sqrt{dx}}{c^2} \right)}{5d^2}$$

$$\begin{array}{c}
\downarrow 851 \\
\frac{2(dx)^{5/2} (a + \operatorname{barctanh}(cx^2))}{5d} - \frac{4bc \left(\frac{2d^3 \int \frac{1}{1-c^2x^4} d\sqrt{dx}}{c^2} - \frac{2d^3 \sqrt{dx}}{c^2} \right)}{5d^2} \\
\downarrow 758 \\
\frac{2(dx)^{5/2} (a + \operatorname{barctanh}(cx^2))}{5d} - \frac{4bc \left(\frac{2d^3 \left(\frac{1}{2} d^2 \int \frac{1}{d^2 - cd^2x^2} d\sqrt{dx} + \frac{1}{2} d^2 \int \frac{1}{cx^2d^2 + d^2} d\sqrt{dx} \right)}{c^2} - \frac{2d^3 \sqrt{dx}}{c^2} \right)}{5d^2} \\
\downarrow 755 \\
\frac{2(dx)^{5/2} (a + \operatorname{barctanh}(cx^2))}{5d} - \frac{4bc \left(\frac{2d^3 \left(\frac{1}{2} d^2 \int \frac{1}{d^2 - cd^2x^2} d\sqrt{dx} + \frac{1}{2} d^2 \left(\frac{\int \frac{d - \sqrt{cdx}}{cx^2d^2 + d^2} d\sqrt{dx}}{2d} + \frac{\int \frac{\sqrt{cdx} + d}{cx^2d^2 + d^2} d\sqrt{dx}}{2d} \right) \right)}{c^2} - \frac{2d^3 \sqrt{dx}}{c^2} \right)}{5d^2} \\
\downarrow 756 \\
\frac{2(dx)^{5/2} (a + \operatorname{barctanh}(cx^2))}{5d} - \frac{4bc \left(\frac{2d^3 \left(\frac{1}{2} d^2 \left(\frac{\int \frac{d - \sqrt{cdx}}{cx^2d^2 + d^2} d\sqrt{dx}}{2d} + \frac{\int \frac{\sqrt{cdx} + d}{cx^2d^2 + d^2} d\sqrt{dx}}{2d} \right) \right) + \frac{1}{2} d^2 \left(\frac{\int \frac{1}{d - \sqrt{cdx}} d\sqrt{dx}}{2d} + \frac{\int \frac{1}{\sqrt{cdx} + d} d\sqrt{dx}}{2d} \right) \right)}{c^2} - \frac{2d^3 \sqrt{dx}}{c^2} \right)}{5d^2} \\
\downarrow 218 \\
\frac{2(dx)^{5/2} (a + \operatorname{barctanh}(cx^2))}{5d} - \frac{4bc \left(\frac{2d^3 \left(\frac{1}{2} d^2 \left(\frac{\int \frac{1}{d - \sqrt{cdx}} d\sqrt{dx}}{2d} + \frac{\arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{cd^{3/2}}} \right) \right) + \frac{1}{2} d^2 \left(\frac{\int \frac{d - \sqrt{cdx}}{cx^2d^2 + d^2} d\sqrt{dx}}{2d} + \frac{\int \frac{\sqrt{cdx} + d}{cx^2d^2 + d^2} d\sqrt{dx}}{2d} \right) \right)}{c^2} - \frac{2d^3 \sqrt{dx}}{c^2} \right)}{5d^2} \\
\downarrow 221
\end{array}$$

$$4bc \left(\frac{2d^3 \left(\frac{1}{2}d^2 \left(\frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2d} + \frac{\int \frac{\sqrt{cdx+d}}{cx^2d^2+d^2} d\sqrt{dx}}{2d} \right) + \frac{1}{2}d^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}}\right) \right)}{c^2} - \frac{2d^3\sqrt{dx}}{c^2} \right)$$

5d²

↓ 1476

$$4bc \left(\frac{2d^3 \left(\frac{1}{2}d^2 \left(\frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2d} + \frac{\int \frac{1}{xd+\frac{d}{\sqrt{c}}-\sqrt{2}\sqrt{dx}\sqrt{d}} \frac{5d}{4\sqrt{c}} d\sqrt{dx}}{2d} + \frac{\int \frac{1}{xd+\frac{d}{\sqrt{c}}+\sqrt{2}\sqrt{dx}\sqrt{d}} \frac{5d}{4\sqrt{c}} d\sqrt{dx}}{2d} \right) + \frac{1}{2}d^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}}\right) \right)}{c^2} \right)$$

5d²

↓ 1082

$$4bc \left(\frac{2d^3 \left(\frac{1}{2}d^2 \left(\frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2d} + \frac{\int \frac{1}{-dx-1} d \left(1 - \frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}} \right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\int \frac{1}{-dx-1} d \left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}} + 1 \right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right) + \frac{1}{2}d^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}}\right) \right)}{c^2} \right)$$

5d²

↓ 217

$$\frac{2(dx)^{5/2} (a + \operatorname{barctanh}(cx^2))}{c^2} - \frac{5d}{c^2} \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}} + 1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} \right) + \frac{1}{2}d^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{c}d^{3/2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{c}d^{3/2}} \right)$$

$5d^2$

↓ 1479

$$\frac{2(dx)^{5/2} (a + \operatorname{barctanh}(cx^2))}{c^2} - \frac{5d}{c^2} \left(\frac{\int -\frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{c}\sqrt{dx}}{\sqrt[4]{c}\left(xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int -\frac{\sqrt{2}(\sqrt{d}+\sqrt{2}\sqrt[4]{c}\sqrt{dx})}{\sqrt[4]{c}\left(xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}} + 1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} \right)$$

$5d^2$

↓ 25

$$\begin{array}{c}
 \frac{2(dx)^{5/2} (a + \operatorname{barctanh}(cx^2))}{5d} \\
 \left(\frac{2d^3}{4bc} \left(\frac{1}{2}d^2 \left(\frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{C}\sqrt{dx}}{xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} + \frac{\int \frac{\sqrt{2}(\sqrt{d}+\sqrt{2}\sqrt[4]{C}\sqrt{dx})}{xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right) \right) \right) \\
 \hline
 c^2
 \end{array}$$

$5d^2$

↓ 27

$$\begin{array}{c}
 \frac{2(dx)^{5/2} (a + \operatorname{barctanh}(cx^2))}{5d} \\
 \left(\frac{2d^3}{4bc} \left(\frac{1}{2}d^2 \left(\frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{C}\sqrt{dx}}{xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}} d\sqrt{dx}}{2\sqrt{2}\sqrt{c}\sqrt{d}} + \frac{\int \frac{\sqrt{d}+\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}} d\sqrt{dx}}{2\sqrt{c}\sqrt{d}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right) \right) \right) + \frac{1}{2}d^2 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}+1\right)}{2\sqrt[4]{C}} \right) \\
 \hline
 c^2
 \end{array}$$

$5d^2$

↓ 1103

$$\frac{2(dx)^{5/2} (a + \operatorname{arctanh}(cx^2))}{5d} - \frac{2d^3 \left(\frac{1}{2}d^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}}\right) + \frac{1}{2}d^2 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}} + 1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right) + \frac{\log\left(\sqrt{cdx} + \sqrt{2}\sqrt[4]{C}\sqrt{d}\right)}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right)}{4bc} - \frac{\phantom{2d^3 \left(\frac{1}{2}d^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}}\right) + \frac{1}{2}d^2 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}} + 1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right) + \frac{\log\left(\sqrt{cdx} + \sqrt{2}\sqrt[4]{C}\sqrt{d}\right)}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right)}}{c^2} - \frac{\phantom{2d^3 \left(\frac{1}{2}d^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}}\right) + \frac{1}{2}d^2 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}} + 1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right) + \frac{\log\left(\sqrt{cdx} + \sqrt{2}\sqrt[4]{C}\sqrt{d}\right)}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right)}}{5d^2}$$

input `Int[(d*x)^(3/2)*(a + b*ArcTanh[c*x^2]), x]`

output `(2*(d*x)^(5/2)*(a + b*ArcTanh[c*x^2]))/(5*d) - (4*b*c*((-2*d^3*Sqrt[d*x])/c^2 + (2*d^3*((d^2*(ArcTan[(c^(1/4)*Sqrt[d*x])/Sqrt[d]]/(2*c^(1/4)*d^(3/2)) + ArcTanh[(c^(1/4)*Sqrt[d*x])/Sqrt[d]]/(2*c^(1/4)*d^(3/2)))))/2 + (d^2*((-ArcTan[1 - (Sqrt[2]*c^(1/4)*Sqrt[d*x])/Sqrt[d]]/(Sqrt[2]*c^(1/4)*Sqrt[d])) + ArcTan[1 + (Sqrt[2]*c^(1/4)*Sqrt[d*x])/Sqrt[d]]/(Sqrt[2]*c^(1/4)*Sqrt[d])))/(2*d) + (-1/2*Log[d + Sqrt[c]*d*x - Sqrt[2]*c^(1/4)*Sqrt[d]*Sqrt[d*x]]/(Sqrt[2]*c^(1/4)*Sqrt[d]) + Log[d + Sqrt[c]*d*x + Sqrt[2]*c^(1/4)*Sqrt[d]*Sqrt[d*x]]/(2*Sqrt[2]*c^(1/4)*Sqrt[d]))/(2*d))/c^2)/(5*d^2)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 218 $\text{Int}[\{(a_)+(b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b]$

rule 221 $\text{Int}[\{(a_)+(b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b]$

rule 755 $\text{Int}[\{(a_)+(b_)*(x_)^4\}^{-1}, x_Symbol] \rightarrow \text{With}\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*r) \ \text{Int}[(r - s*x^2)/(a + b*x^4), x], x] + \text{Simp}[1/(2*r) \ \text{Int}[(r + s*x^2)/(a + b*x^4), x], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

rule 756 $\text{Int}[\{(a_)+(b_)*(x_)^4\}^{-1}, x_Symbol] \rightarrow \text{With}\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[r/(2*a) \ \text{Int}[1/(r - s*x^2), x], x] + \text{Simp}[r/(2*a) \ \text{Int}[1/(r + s*x^2), x], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ !\text{GtQ}[a/b, 0]$

rule 758 $\text{Int}[\{(a_)+(b_)*(x_)^{n_}\}^{-1}, x_Symbol] \rightarrow \text{With}\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[r/(2*a) \ \text{Int}[1/(r - s*x^{(n/2)}), x], x] + \text{Simp}[r/(2*a) \ \text{Int}[1/(r + s*x^{(n/2)}), x], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{IGtQ}[n/4, 1] \ \&\& \ !\text{GtQ}[a/b, 0]$

rule 843 $\text{Int}[\{(c_)*(x_)\}^{m_}*\{(a_)+(b_)*(x_)^{n_}\}^{p_}, x_Symbol] \rightarrow \text{Simp}[c^{(n-1)}*(c*x)^{(m-n+1)}*((a + b*x^n)^{(p+1)}/(b*(m+n*p+1))), x] - \text{Simp}[a*c^n*((m-n+1)/(b*(m+n*p+1))) \ \text{Int}[(c*x)^{(m-n)}*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{GtQ}[m, n-1] \ \&\& \ \text{NeQ}[m+n*p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 851 $\text{Int}[\{(c_)*(x_)\}^{m_}*\{(a_)+(b_)*(x_)^{n_}\}^{p_}, x_Symbol] \rightarrow \text{With}\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \ \text{Subst}[\text{Int}[x^{(k*(m+1)-1)}*(a + b*(x^{(k*n)}/c^n)]^p, x], x, (c*x)^{(1/k)}], x] /; \text{FreeQ}\{a, b, c, p\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 6464 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)*((d_)*(x_)^(m_)), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 248, normalized size of antiderivative = 0.96

method	result
derivativedivides	$\frac{2(dx)^{\frac{5}{2}}a + 2b}{5} \left(\frac{(dx)^{\frac{5}{2}} \operatorname{arctanh}(cx^2)}{5} - \frac{4cd^2}{c^2} + \frac{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \left(\ln \left(\frac{\sqrt{dx} + \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}{\sqrt{dx} - \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) \right)}{8c^2} + \frac{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{2} \left(\ln \left(\frac{\sqrt{dx} + \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}{\sqrt{dx} - \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) \right)}{8c^2} \right)$
default	$\frac{2(dx)^{\frac{5}{2}}a + 2b}{5} \left(\frac{(dx)^{\frac{5}{2}} \operatorname{arctanh}(cx^2)}{5} - \frac{4cd^2}{c^2} + \frac{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \left(\ln \left(\frac{\sqrt{dx} + \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}{\sqrt{dx} - \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) \right)}{8c^2} + \frac{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{2} \left(\ln \left(\frac{\sqrt{dx} + \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}{\sqrt{dx} - \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) \right)}{8c^2} \right)$
parts	$\frac{2a(dx)^{\frac{5}{2}}}{5d} + \frac{2b}{5} \left(\frac{(dx)^{\frac{5}{2}} \operatorname{arctanh}(cx^2)}{5} - \frac{4cd^2}{c^2} + \frac{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \left(\ln \left(\frac{\sqrt{dx} + \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}{\sqrt{dx} - \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) \right)}{8c^2} + \frac{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{2} \left(\ln \left(\frac{\sqrt{dx} + \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}{\sqrt{dx} - \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) \right)}{8c^2} \right)$

```
input int((d*x)^(3/2)*(a+b*arctanh(c*x^2)),x,method=_RETURNVERBOSE)
```

output

```
2/d*(1/5*(d*x)^(5/2)*a+b*(1/5*(d*x)^(5/2)*arctanh(c*x^2)-4/5*c*d^2*(-1/c^2
*(d*x)^(1/2)+1/8/c^2*(d^2/c)^(1/4)*(ln(((d*x)^(1/2)+(d^2/c)^(1/4))/((d*x)^(
1/2)-(d^2/c)^(1/4)))+2*arctan((d*x)^(1/2)/(d^2/c)^(1/4)))+1/16/c^2*(d^2/c
)^(1/4)*2^(1/2)*(ln((d*x+(d^2/c)^(1/4)*(d*x)^(1/2)*2^(1/2)+(d^2/c)^(1/2)))/
(d*x-(d^2/c)^(1/4)*(d*x)^(1/2)*2^(1/2)+(d^2/c)^(1/2)))+2*arctan(2^(1/2)/(d
^2/c)^(1/4)*(d*x)^(1/2)+1)+2*arctan(2^(1/2)/(d^2/c)^(1/4)*(d*x)^(1/2)-1))
))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 376, normalized size of antiderivative = 1.46

$$\int (dx)^{3/2} (a + b \operatorname{arctanh}(cx^2)) dx =$$

$$\frac{\left(\frac{b^4 d^6}{c^5}\right)^{\frac{1}{4}} c \log\left(\sqrt{dx}bd + \left(\frac{b^4 d^6}{c^5}\right)^{\frac{1}{4}} c\right) + i \left(\frac{b^4 d^6}{c^5}\right)^{\frac{1}{4}} c \log\left(\sqrt{dx}bd + i \left(\frac{b^4 d^6}{c^5}\right)^{\frac{1}{4}} c\right) - i \left(\frac{b^4 d^6}{c^5}\right)^{\frac{1}{4}} c \log\left(\sqrt{dx}bd - i \left(\frac{b^4 d^6}{c^5}\right)^{\frac{1}{4}} c\right) - \left(\frac{b^4 d^6}{c^5}\right)^{\frac{1}{4}} c \log\left(\sqrt{dx}bd - \left(\frac{b^4 d^6}{c^5}\right)^{\frac{1}{4}} c\right)}{c}$$

input

```
integrate((d*x)^(3/2)*(a+b*arctanh(c*x^2)),x, algorithm="fricas")
```

output

```
-1/5*((b^4*d^6/c^5)^(1/4)*c*log(sqrt(d*x)*b*d + (b^4*d^6/c^5)^(1/4)*c) + I
*(b^4*d^6/c^5)^(1/4)*c*log(sqrt(d*x)*b*d + I*(b^4*d^6/c^5)^(1/4)*c) - I*(b
^4*d^6/c^5)^(1/4)*c*log(sqrt(d*x)*b*d - I*(b^4*d^6/c^5)^(1/4)*c) - (b^4*d^
6/c^5)^(1/4)*c*log(sqrt(d*x)*b*d - (b^4*d^6/c^5)^(1/4)*c) + (-b^4*d^6/c^5)
^(1/4)*c*log(sqrt(d*x)*b*d + (-b^4*d^6/c^5)^(1/4)*c) + I*(-b^4*d^6/c^5)^(1
/4)*c*log(sqrt(d*x)*b*d + I*(-b^4*d^6/c^5)^(1/4)*c) - I*(-b^4*d^6/c^5)^(1/
4)*c*log(sqrt(d*x)*b*d - I*(-b^4*d^6/c^5)^(1/4)*c) - (-b^4*d^6/c^5)^(1/4)*
c*log(sqrt(d*x)*b*d - (-b^4*d^6/c^5)^(1/4)*c) - (b*c*d*x^2*log(-(c*x^2 + 1
)/(c*x^2 - 1)) + 2*a*c*d*x^2 + 8*b*d)*sqrt(d*x))/c
```


Sympy [F]

$$\int (dx)^{3/2} (a + b \operatorname{arctanh}(cx^2)) dx = \int (dx)^{3/2} (a + b \operatorname{atanh}(cx^2)) dx$$

```
input integrate((d*x)**(3/2)*(a+b*atanh(c*x**2)), x)
```

```
output Integral((d*x)**(3/2)*(a + b*atanh(c*x**2)), x)
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 310, normalized size of antiderivative = 1.21

$$\int (dx)^{3/2} (a$$

$$+ b \operatorname{arctanh}(cx^2)) dx = 4(dx)^{5/2} a + 4(dx)^{5/2} \operatorname{arctanh}(cx^2) + \frac{16\sqrt{dx}d^4}{c^2} \frac{2\sqrt{2}d^5 \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}c\frac{1}{4}\sqrt{d}+2\sqrt{dx}\sqrt{c}\right)}{2\sqrt{cd}}\right) + 2\sqrt{2}d^5 a}{\sqrt{cd}}$$

```
input integrate((d*x)^(3/2)*(a+b*arctanh(c*x^2)), x, algorithm="maxima")
```

output

```
1/10*(4*(d*x)^(5/2)*a + (4*(d*x)^(5/2)*arctanh(c*x^2) + (16*sqrt(d*x)*d^4/
c^2 - (2*sqrt(2)*d^5*arctan(1/2*sqrt(2)*(sqrt(2)*c^(1/4)*sqrt(d) + 2*sqrt(
d*x)*sqrt(c))/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d) + 2*sqrt(2)*d^5*arctan(-1/2
*sqrt(2)*(sqrt(2)*c^(1/4)*sqrt(d) - 2*sqrt(d*x)*sqrt(c))/sqrt(sqrt(c)*d))/
sqrt(sqrt(c)*d) + sqrt(2)*d^(9/2)*log(sqrt(c)*d*x + sqrt(2)*sqrt(d*x)*c^(1
/4)*sqrt(d) + d)/c^(1/4) - sqrt(2)*d^(9/2)*log(sqrt(c)*d*x - sqrt(2)*sqrt(
d*x)*c^(1/4)*sqrt(d) + d)/c^(1/4))/c^2 - 2*(2*d^5*arctan(sqrt(d*x)*sqrt(c)
/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d) - d^5*log((sqrt(d*x)*sqrt(c) - sqrt(sqrt(
c)*d))/(sqrt(d*x)*sqrt(c) + sqrt(sqrt(c)*d)))/sqrt(sqrt(c)*d))/c^2*c/d^2
)*b)/d
```

Giac [F]

$$\int (dx)^{3/2} (a + b \operatorname{arctanh}(cx^2)) dx = \int (dx)^{3/2} (b \operatorname{arctanh}(cx^2) + a) dx$$

input

```
integrate((d*x)^(3/2)*(a+b*arctanh(c*x^2)),x, algorithm="giac")
```

output

```
integrate((d*x)^(3/2)*(b*arctanh(c*x^2) + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int (dx)^{3/2} (a + b \operatorname{arctanh}(cx^2)) dx = \int (dx)^{3/2} (a + b \operatorname{atanh}(cx^2)) dx$$

input

```
int((d*x)^(3/2)*(a + b*atanh(c*x^2)),x)
```

output

```
int((d*x)^(3/2)*(a + b*atanh(c*x^2)), x)
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 243, normalized size of antiderivative = 0.95

$$\int (dx)^{3/2} (a + b \operatorname{arctanh}(cx^2)) dx = \frac{\sqrt{d} d \left(2c^{3/4} \sqrt{2} \operatorname{atan} \left(\frac{c^{1/4} \sqrt{2} - 2\sqrt{x} \sqrt{c}}{c^{1/4} \sqrt{2}} \right) b - 2c^{3/4} \sqrt{2} \operatorname{atan} \left(\frac{c^{1/4} \sqrt{2} + 2\sqrt{x} \sqrt{c}}{c^{1/4} \sqrt{2}} \right) b - 4c^{3/4} \operatorname{atan} \left(\frac{\sqrt{x}}{c^{1/4}} \right) \right)}{10c^{3/2}}$$

input `int((d*x)^(3/2)*(a+b*atanh(c*x^2)),x)`output `(sqrt(d)*d*(2*c**(3/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*b - 2*c**(3/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*b - 4*c**(3/4)*atan((sqrt(x)*sqrt(c))/c**(1/4))*b - 2*c**(3/4)*sqrt(2)*atanh(c*x**2)*b + 4*sqrt(x)*atanh(c*x**2)*b*c**2*x**2 - c**(3/4)*sqrt(2)*log(c**(1/4) + sqrt(x)*sqrt(c))*b - c**(3/4)*sqrt(2)*log(- c**(1/4) + sqrt(x)*sqrt(c))*b + 2*c**(3/4)*sqrt(2)*log(- sqrt(x)*c**(1/4)*sqrt(2) + sqrt(c)*x + 1)*b - c**(3/4)*sqrt(2)*log(sqrt(c)*x + 1)*b - 2*c**(3/4)*log(c**(1/4) + sqrt(x)*sqrt(c))*b + 2*c**(3/4)*log(- c**(1/4) + sqrt(x)*sqrt(c))*b + 4*sqrt(x)*a*c**2*x**2 + 16*sqrt(x)*b*c)/(10*c**2)`

3.84 $\int \sqrt{dx}(a + b \operatorname{arctanh}(cx^2)) dx$

Optimal result	719
Mathematica [A] (verified)	720
Rubi [A] (verified)	720
Maple [A] (verified)	726
Fricas [C] (verification not implemented)	728
Sympy [F]	729
Maxima [A] (verification not implemented)	730
Giac [F]	731
Mupad [F(-1)]	731
Reduce [B] (verification not implemented)	731

Optimal result

Integrand size = 18, antiderivative size = 241

$$\begin{aligned}
 \int \sqrt{dx}(a + b \operatorname{arctanh}(cx^2)) dx = & \frac{2b\sqrt{d} \arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{3c^{3/4}} \\
 & - \frac{\sqrt{2}b\sqrt{d} \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{3c^{3/4}} \\
 & + \frac{\sqrt{2}b\sqrt{d} \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{3c^{3/4}} \\
 & + \frac{2(dx)^{3/2}(a + b \operatorname{arctanh}(cx^2))}{3d} \\
 & - \frac{2b\sqrt{d} \operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{3c^{3/4}} \\
 & - \frac{\sqrt{2}b\sqrt{d} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}(1+\sqrt{cx})}\right)}{3c^{3/4}}
 \end{aligned}$$

output

$$\begin{aligned} & \frac{2}{3} b d^{1/2} \arctan(c^{1/4} (d x)^{1/2} / d^{1/2}) / c^{3/4} + \frac{1}{3} 2^{1/2} b d^{1/2} \\ & \arctan(-1 + 2^{1/2} c^{1/4} (d x)^{1/2} / d^{1/2}) / c^{3/4} + \frac{1}{3} 2^{1/2} b \\ & d^{1/2} \arctan(1 + 2^{1/2} c^{1/4} (d x)^{1/2} / d^{1/2}) / c^{3/4} + \frac{2}{3} (d x)^{3/2} \\ & (a + b \operatorname{arctanh}(c x^2)) / d - \frac{2}{3} b d^{1/2} \operatorname{arctanh}(c^{1/4} (d x)^{1/2} / d^{1/2}) / c^{3/4} \\ & - \frac{1}{3} 2^{1/2} b d^{1/2} \operatorname{arctanh}(2^{1/2} c^{1/4} (d x)^{1/2} / d^{1/2}) / (1 + c^{1/2} x) / c^{3/4} \end{aligned}$$
Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 227, normalized size of antiderivative = 0.94

$$\int \sqrt{dx} (a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{\sqrt{dx} (4ac^{3/4} x^{3/2} - 2\sqrt{2}b \arctan(1 - \sqrt{2}\sqrt[4]{c}\sqrt{x}) + 2\sqrt{2}b \arctan(1 + \sqrt{2}\sqrt[4]{c}\sqrt{x}) + 4b \arctan(\sqrt[4]{c}\sqrt{x}) + 4a\sqrt{dx})}{6c^{3/4}\sqrt{dx}}$$

input

`Integrate[Sqrt[d*x]*(a + b*ArcTanh[c*x^2]),x]`

output

$$\begin{aligned} & (\operatorname{Sqrt}[d x] * (4 a c^{3/4} x^{3/2} - 2 \operatorname{Sqrt}[2] * b \operatorname{ArcTan}[1 - \operatorname{Sqrt}[2] * c^{1/4} * \\ & \operatorname{Sqrt}[x]] + 2 \operatorname{Sqrt}[2] * b \operatorname{ArcTan}[1 + \operatorname{Sqrt}[2] * c^{1/4} * \operatorname{Sqrt}[x]] + 4 * b \operatorname{ArcTan}[c^{1/4} * \\ & \operatorname{Sqrt}[x]] + 4 * b c^{3/4} x^{3/2} * \operatorname{ArcTanh}[c x^2] + 2 * b \operatorname{Log}[1 - c^{1/4} * \operatorname{Sqrt}[x]] \\ & - 2 * b \operatorname{Log}[1 + c^{1/4} * \operatorname{Sqrt}[x]] + \operatorname{Sqrt}[2] * b \operatorname{Log}[1 - \operatorname{Sqrt}[2] * c^{1/4} * \\ & \operatorname{Sqrt}[x] + \operatorname{Sqrt}[c] * x] - \operatorname{Sqrt}[2] * b \operatorname{Log}[1 + \operatorname{Sqrt}[2] * c^{1/4} * \operatorname{Sqrt}[x] + \operatorname{Sqrt}[c] \\ & * x])) / (6 * c^{3/4} * \operatorname{Sqrt}[x]) \end{aligned}$$
Rubi [A] (verified)Time = 0.61 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.36, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {6464, 851, 27, 830, 826, 827, 218, 221, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{dx} (a + b \operatorname{arctanh}(cx^2)) dx$$

$$\begin{aligned}
& \downarrow 6464 \\
& \frac{2(dx)^{3/2} (a + \operatorname{barctanh}(cx^2))}{3d} - \frac{4bc \int \frac{(dx)^{5/2}}{1-c^2x^4} dx}{3d^2} \\
& \downarrow 851 \\
& \frac{2(dx)^{3/2} (a + \operatorname{barctanh}(cx^2))}{3d} - \frac{8bc \int \frac{d^7 x^3}{d^4 - c^2 d^4 x^4} d\sqrt{dx}}{3d^3} \\
& \downarrow 27 \\
& \frac{2(dx)^{3/2} (a + \operatorname{barctanh}(cx^2))}{3d} - \frac{8}{3}bcd \int \frac{d^3 x^3}{d^4 - c^2 d^4 x^4} d\sqrt{dx} \\
& \downarrow 830 \\
& \frac{2(dx)^{3/2} (a + \operatorname{barctanh}(cx^2))}{3d} - \frac{8}{3}bcd \left(\frac{\int \frac{dx}{d^2 - cd^2 x^2} d\sqrt{dx}}{2c} - \frac{\int \frac{dx}{cx^2 d^2 + d^2} d\sqrt{dx}}{2c} \right) \\
& \downarrow 826 \\
& \frac{2(dx)^{3/2} (a + \operatorname{barctanh}(cx^2))}{3d} - \frac{8}{3}bcd \left(\frac{\int \frac{dx}{d^2 - cd^2 x^2} d\sqrt{dx}}{2c} - \frac{\int \frac{\sqrt{cx}d+d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} \right) \\
& \downarrow 827 \\
& \frac{2(dx)^{3/2} (a + \operatorname{barctanh}(cx^2))}{3d} - \frac{8}{3}bcd \left(\frac{\int \frac{1}{d-\sqrt{cdx}} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{1}{\sqrt{cx}d+d} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{\sqrt{cx}d+d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} \right) \\
& \downarrow 218 \\
& \frac{2(dx)^{3/2} (a + \operatorname{barctanh}(cx^2))}{3d} - \frac{8}{3}bcd \left(\frac{\int \frac{1}{d-\sqrt{cdx}} d\sqrt{dx}}{2\sqrt{c}} - \frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\int \frac{\sqrt{cx}d+d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} \right) \\
& \downarrow 221
\end{aligned}$$

$$\begin{aligned}
 & \frac{2(dx)^{3/2} (a + \operatorname{barctanh}(cx^2))}{3d} - \\
 & \frac{8}{3}bcd \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\int \frac{\sqrt{cx^2+d}}{cx^2d^2+d^2} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2\sqrt{c}} \right) \\
 & \quad \downarrow 1476 \\
 & \frac{2(dx)^{3/2} (a + \operatorname{barctanh}(cx^2))}{3d} - \\
 & \frac{8}{3}bcd \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\int \frac{1}{xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}} d\sqrt{dx}}{2\sqrt{c}} + \frac{\int \frac{1}{xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2\sqrt{c}} \right) \\
 & \quad \downarrow 1082 \\
 & \frac{2(dx)^{3/2} (a + \operatorname{barctanh}(cx^2))}{3d} - \\
 & \frac{8}{3}bcd \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\int \frac{1}{-dx-1} d\left(1-\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\int \frac{1}{-dx-1} d\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2\sqrt{c}} \right) \\
 & \quad \downarrow 217 \\
 & \frac{2(dx)^{3/2} (a + \operatorname{barctanh}(cx^2))}{3d} - \\
 & \frac{8}{3}bcd \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\operatorname{arctan}\left(1-\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2\sqrt{c}} \right) \\
 & \quad \downarrow 1479
 \end{aligned}$$

$$\frac{2(dx)^{3/2} (a + \operatorname{barctanh}(cx^2))}{3d} - \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}} + 1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\operatorname{arctan}\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{\sqrt{2}\sqrt{d} - 2\sqrt[4]{c}\sqrt{dx}}{\sqrt[4]{c}\left(xd + \frac{d}{c} - \frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}\right)}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} dx}{2c} \right)$$

25

$$\frac{2(dx)^{3/2} (a + \operatorname{barctanh}(cx^2))}{3d} - \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}} + 1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\operatorname{arctan}\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{\sqrt{2}\sqrt{d} - 2\sqrt[4]{c}\sqrt{dx}}{\sqrt[4]{c}\left(xd + \frac{d}{c} - \frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}\right)}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} dx}{2c} \right)$$

27

$$\frac{2(dx)^{3/2} (a + \operatorname{barctanh}(cx^2))}{3d} - \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}} + 1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\operatorname{arctan}\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{\sqrt{2}\sqrt{d} - 2\sqrt[4]{c}\sqrt{dx}}{xd + \frac{d}{c} - \frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}} d\sqrt{dx}}{2\sqrt{2}\sqrt{c}\sqrt{d}} + \frac{\int}{2\sqrt{c}} \right)$$

1103

$$\frac{2(dx)^{3/2} (a + b \operatorname{arctanh}(cx^2))}{3d} - \frac{8}{3}bcd \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}+1}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\operatorname{arctan}\left(1-\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\log\left(\sqrt{cdx}+\sqrt{2}\sqrt[4]{c}\sqrt{d}\sqrt{dx}+d\right)}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} \right)$$

```
input Int[Sqrt[d*x]*(a + b*ArcTanh[c*x^2]), x]
```

```
output (2*(d*x)^(3/2)*(a + b*ArcTanh[c*x^2]))/(3*d) - (8*b*c*d*((-1/2*ArcTan[(c^(1/4)*Sqrt[d*x])/Sqrt[d]]/Sqrt[d]]/(c^(3/4)*Sqrt[d]) + ArcTan[(c^(1/4)*Sqrt[d*x])/Sqrt[d]]/(2*c^(3/4)*Sqrt[d]))/(2*c) - ((-ArcTan[1 - (Sqrt[2]*c^(1/4)*Sqrt[d*x])/Sqrt[d]]/Sqrt[d]]/(Sqrt[2]*c^(1/4)*Sqrt[d])) + ArcTan[1 + (Sqrt[2]*c^(1/4)*Sqrt[d*x])/Sqrt[d]]/Sqrt[d]]/(Sqrt[2]*c^(1/4)*Sqrt[d]))/(2*Sqrt[c]) - (-1/2*Log[d + Sqrt[c]*d*x - Sqrt[2]*c^(1/4)*Sqrt[d]*Sqrt[d*x]]/(Sqrt[2]*c^(1/4)*Sqrt[d]) + Log[d + Sqrt[c]*d*x + Sqrt[2]*c^(1/4)*Sqrt[d]*Sqrt[d*x]]/(2*Sqrt[2]*c^(1/4)*Sqrt[d]))/(2*Sqrt[c]))/(2*c))/3
```

Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

rule 221 $\text{Int}[(a_ + (b_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) \cdot \text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$

rule 826 $\text{Int}[(x_)^2/((a_) + (b_ \cdot)(x_)^4), x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*s) \ \text{Int}[(r + s*x^2)/(a + b*x^4), x], x] - \text{Simp}[1/(2*s) \ \text{Int}[(r - s*x^2)/(a + b*x^4), x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

rule 827 $\text{Int}[(x_)^2/((a_) + (b_ \cdot)(x_)^4), x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[s/(2*b) \ \text{Int}[1/(r + s*x^2), x], x] - \text{Simp}[s/(2*b) \ \text{Int}[1/(r - s*x^2), x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ !\text{GtQ}[a/b, 0]$

rule 830 $\text{Int}[(x_)^m/((a_) + (b_ \cdot)(x_)^n), x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[s/(2*b) \ \text{Int}[x^{(m-n/2)}/(r + s*x^{(n/2)}), x], x] - \text{Simp}[s/(2*b) \ \text{Int}[x^{(m-n/2)}/(r - s*x^{(n/2)}), x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{IGtQ}[n/4, 0] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{LeQ}[n/2, m] \ \&\& \ \text{LtQ}[m, n] \ \&\& \ !\text{GtQ}[a/b, 0]$

rule 851 $\text{Int}[(c_ \cdot)(x_)^m/((a_) + (b_ \cdot)(x_)^n)^p, x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \ \text{Subst}[\text{Int}[x^{(k*(m+1)-1)}*(a + b*(x^{(k*n)}/c^n))^p, x], x, (c*x)^{(1/k)}], x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 1082 $\text{Int}[(a_) + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S \ \text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}[(d_) + (e_ \cdot)(x_)/((a_ \cdot) + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$

rule 1476

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

rule 1479

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

rule 6464

```
Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)*((d_)*(x_)^(m_)), x_Symbol] :
> Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n
/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a,
b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.00

method	result
derivativedivides	$\frac{2(dx)^{\frac{3}{2}}a}{3} + 2b \left(\frac{(dx)^{\frac{3}{2}} \operatorname{arctanh}(cx^2)}{3} - \frac{4cd^2 \left(\frac{\sqrt{2} \left(\ln \left(\frac{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}} \right)}{dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 1} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)}{16c^2 \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)}{3} \right)$
default	$\frac{2(dx)^{\frac{3}{2}}a}{3} + 2b \left(\frac{(dx)^{\frac{3}{2}} \operatorname{arctanh}(cx^2)}{3} - \frac{4cd^2 \left(\frac{\sqrt{2} \left(\ln \left(\frac{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}} \right)}{dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 1} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)}{16c^2 \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)}{3} \right)$
parts	$\frac{2a(dx)^{\frac{3}{2}}}{3d} + \left(\frac{(dx)^{\frac{3}{2}} \operatorname{arctanh}(cx^2)}{3} - \frac{4cd^2 \left(\frac{\sqrt{2} \left(\ln \left(\frac{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}} \right)}{dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 1} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)}{16c^2 \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)}{3} \right)$

input `int((d*x)^(1/2)*(a+b*arctanh(c*x^2)),x,method=_RETURNVERBOSE)`

output

```
2/d*(1/3*(d*x)^(3/2)*a+b*(1/3*(d*x)^(3/2)*arctanh(c*x^2)-4/3*c*d^2*(-1/16/
c^2/(d^2/c)^(1/4)*2^(1/2)*(ln((d*x-(d^2/c)^(1/4)*(d*x)^(1/2)*2^(1/2)+(d^2/
c)^(1/2))/(d*x+(d^2/c)^(1/4)*(d*x)^(1/2)*2^(1/2)+(d^2/c)^(1/2))))+2*arctan(
2^(1/2)/(d^2/c)^(1/4)*(d*x)^(1/2)+1)+2*arctan(2^(1/2)/(d^2/c)^(1/4)*(d*x)^(
1/2)-1))-1/8/c^2/(d^2/c)^(1/4)*(2*arctan((d*x)^(1/2)/(d^2/c)^(1/4))-ln(((
d*x)^(1/2)+(d^2/c)^(1/4))/((d*x)^(1/2)-(d^2/c)^(1/4))))))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 385, normalized size of antiderivative = 1.60

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx^2)) dx = \frac{1}{3} \left(bx \log \left(-\frac{cx^2 + 1}{cx^2 - 1} \right) + 2ax \right) \sqrt{dx} \\ - \frac{1}{3} \left(\frac{b^4 d^2}{c^3} \right)^{\frac{1}{4}} \log \left(\sqrt{dx} b^3 d + \left(\frac{b^4 d^2}{c^3} \right)^{\frac{3}{4}} c^2 \right) \\ + \frac{1}{3} i \left(\frac{b^4 d^2}{c^3} \right)^{\frac{1}{4}} \log \left(\sqrt{dx} b^3 d + i \left(\frac{b^4 d^2}{c^3} \right)^{\frac{3}{4}} c^2 \right) \\ - \frac{1}{3} i \left(\frac{b^4 d^2}{c^3} \right)^{\frac{1}{4}} \log \left(\sqrt{dx} b^3 d - i \left(\frac{b^4 d^2}{c^3} \right)^{\frac{3}{4}} c^2 \right) \\ + \frac{1}{3} \left(\frac{b^4 d^2}{c^3} \right)^{\frac{1}{4}} \log \left(\sqrt{dx} b^3 d - \left(\frac{b^4 d^2}{c^3} \right)^{\frac{3}{4}} c^2 \right) \\ + \frac{1}{3} \left(-\frac{b^4 d^2}{c^3} \right)^{\frac{1}{4}} \log \left(\sqrt{dx} b^3 d + \left(-\frac{b^4 d^2}{c^3} \right)^{\frac{3}{4}} c^2 \right) \\ - \frac{1}{3} i \left(-\frac{b^4 d^2}{c^3} \right)^{\frac{1}{4}} \log \left(\sqrt{dx} b^3 d + i \left(-\frac{b^4 d^2}{c^3} \right)^{\frac{3}{4}} c^2 \right) \\ + \frac{1}{3} i \left(-\frac{b^4 d^2}{c^3} \right)^{\frac{1}{4}} \log \left(\sqrt{dx} b^3 d - i \left(-\frac{b^4 d^2}{c^3} \right)^{\frac{3}{4}} c^2 \right) \\ - \frac{1}{3} \left(-\frac{b^4 d^2}{c^3} \right)^{\frac{1}{4}} \log \left(\sqrt{dx} b^3 d - \left(-\frac{b^4 d^2}{c^3} \right)^{\frac{3}{4}} c^2 \right)$$

input

```
integrate((d*x)^(1/2)*(a+b*arctanh(c*x^2)),x, algorithm="fricas")
```

output

```
1/3*(b*x*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 2*a*x)*sqrt(d*x) - 1/3*(b^4*d^2/c^3)^(1/4)*log(sqrt(d*x)*b^3*d + (b^4*d^2/c^3)^(3/4)*c^2) + 1/3*I*(b^4*d^2/c^3)^(1/4)*log(sqrt(d*x)*b^3*d + I*(b^4*d^2/c^3)^(3/4)*c^2) - 1/3*I*(b^4*d^2/c^3)^(1/4)*log(sqrt(d*x)*b^3*d - I*(b^4*d^2/c^3)^(3/4)*c^2) + 1/3*(b^4*d^2/c^3)^(1/4)*log(sqrt(d*x)*b^3*d - (b^4*d^2/c^3)^(3/4)*c^2) + 1/3*(-b^4*d^2/c^3)^(1/4)*log(sqrt(d*x)*b^3*d + (-b^4*d^2/c^3)^(3/4)*c^2) - 1/3*I*(-b^4*d^2/c^3)^(1/4)*log(sqrt(d*x)*b^3*d + I*(-b^4*d^2/c^3)^(3/4)*c^2) + 1/3*I*(-b^4*d^2/c^3)^(1/4)*log(sqrt(d*x)*b^3*d - I*(-b^4*d^2/c^3)^(3/4)*c^2) - 1/3*(-b^4*d^2/c^3)^(1/4)*log(sqrt(d*x)*b^3*d - (-b^4*d^2/c^3)^(3/4)*c^2)
```

Sympy [F]

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx^2)) dx = \int \sqrt{dx}(a + b \operatorname{atanh}(cx^2)) dx$$

input

```
integrate((d*x)**(1/2)*(a+b*atanh(c*x**2)), x)
```

output

```
Integral(sqrt(d*x)*(a + b*atanh(c*x**2)), x)
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 301, normalized size of antiderivative = 1.25

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx^2)) dx$$

$$4(dx)^{\frac{3}{2}} a + 4(dx)^{\frac{3}{2}} \operatorname{arctanh}(cx^2) + \frac{d^4 \left(\frac{2\sqrt{2} \operatorname{arctan}\left(\frac{\sqrt{2}\left(\sqrt{2}c^{\frac{1}{4}}\sqrt{d}+2\sqrt{dx}\sqrt{c}\right)}{2\sqrt{\sqrt{cd}}}\right)}{\sqrt{\sqrt{cd}\sqrt{c}}}\right) + \frac{2\sqrt{2} \operatorname{arctan}\left(-\frac{\sqrt{2}\left(\sqrt{2}c^{\frac{1}{4}}\sqrt{d}-2\sqrt{dx}\sqrt{c}\right)}{2\sqrt{\sqrt{cd}}}\right)}{\sqrt{\sqrt{cd}\sqrt{c}}}}{c} - \frac{\sqrt{2} \log\left(\frac{\sqrt{2}\left(\sqrt{2}c^{\frac{1}{4}}\sqrt{d}+2\sqrt{dx}\sqrt{c}\right)}{\sqrt{2}\left(\sqrt{2}c^{\frac{1}{4}}\sqrt{d}-2\sqrt{dx}\sqrt{c}\right)}\right)}{c} \right)}{6d}$$

```
input integrate((d*x)^(1/2)*(a+b*arctanh(c*x^2)),x, algorithm="maxima")
```

```
output 1/6*(4*(d*x)^(3/2)*a + (4*(d*x)^(3/2)*arctanh(c*x^2) + (d^4*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*c^(1/4)*sqrt(d) + 2*sqrt(d*x)*sqrt(c))/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*c^(1/4)*sqrt(d) - 2*sqrt(d*x)*sqrt(c))/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d)*sqrt(c) - sqrt(2)*log(sqrt(c)*d*x + sqrt(2)*sqrt(d*x)*c^(1/4)*sqrt(d) + d)/(c^(3/4)*sqrt(d)) + sqrt(2)*log(sqrt(c)*d*x - sqrt(2)*sqrt(d*x)*c^(1/4)*sqrt(d) + d)/(c^(3/4)*sqrt(d)))/c + 2*d^4*(2*arctan(sqrt(d*x)*sqrt(c)/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d)*sqrt(c) + log((sqrt(d*x)*sqrt(c) - sqrt(sqrt(c)*d))/sqrt(d*x)*sqrt(c) + sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d)*sqrt(c)))/c)*c/d^2)*b)/d
```

Giac [F]

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx^2)) dx = \int \sqrt{dx}(b \operatorname{artanh}(cx^2) + a) dx$$

input `integrate((d*x)^(1/2)*(a+b*arctanh(c*x^2)),x, algorithm="giac")`

output `integrate(sqrt(d*x)*(b*arctanh(c*x^2) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx^2)) dx = \int \sqrt{dx}(a + b \operatorname{atanh}(cx^2)) dx$$

input `int((d*x)^(1/2)*(a + b*atanh(c*x^2)),x)`

output `int((d*x)^(1/2)*(a + b*atanh(c*x^2)), x)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.95

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{\sqrt{d} \left(-2c^{\frac{1}{4}} \sqrt{2} \operatorname{atan} \left(\frac{c^{\frac{1}{4}} \sqrt{2} - 2\sqrt{x} \sqrt{c}}{c^{\frac{1}{4}} \sqrt{2}} \right) b + 2c^{\frac{1}{4}} \sqrt{2} \operatorname{atan} \left(\frac{c^{\frac{1}{4}} \sqrt{2} + 2\sqrt{x} \sqrt{c}}{c^{\frac{1}{4}} \sqrt{2}} \right) b + 4c^{\frac{1}{4}} \operatorname{atan} \left(\frac{\sqrt{x} \sqrt{c}}{c^{\frac{1}{4}}} \right) b - 2c^{\frac{1}{4}} \sqrt{2} \operatorname{atanh} \right)}{1}$$

input `int((d*x)^(1/2)*(a+b*atanh(c*x^2)),x)`

output

```
(sqrt(d)*(- 2*c**(1/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(c)
)/(c**(1/4)*sqrt(2)))*b + 2*c**(1/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) + 2*sq
rt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*b + 4*c**(1/4)*atan((sqrt(x)*sqrt(c))/c
**(1/4))*b - 2*c**(1/4)*sqrt(2)*atanh(c*x**2)*b + 4*sqrt(x)*atanh(c*x**2)*
b*c*x - c**(1/4)*sqrt(2)*log(c**(1/4) + sqrt(x)*sqrt(c))*b - c**(1/4)*sqrt
(2)*log(- c**(1/4) + sqrt(x)*sqrt(c))*b + 2*c**(1/4)*sqrt(2)*log(- sqrt(
x)*c**(1/4)*sqrt(2) + sqrt(c)*x + 1)*b - c**(1/4)*sqrt(2)*log(sqrt(c)*x +
1)*b - 2*c**(1/4)*log(c**(1/4) + sqrt(x)*sqrt(c))*b + 2*c**(1/4)*log(- c*
*(1/4) + sqrt(x)*sqrt(c))*b + 4*sqrt(x)*a*c*x)/(6*c)
```

3.85 $\int \frac{a+b\operatorname{arctanh}(cx^2)}{\sqrt{dx}} dx$

Optimal result	733
Mathematica [A] (verified)	734
Rubi [A] (verified)	734
Maple [A] (verified)	740
Fricas [C] (verification not implemented)	741
Sympy [F]	741
Maxima [A] (verification not implemented)	742
Giac [B] (verification not implemented)	743
Mupad [F(-1)]	744
Reduce [B] (verification not implemented)	744

Optimal result

Integrand size = 18, antiderivative size = 227

$$\int \frac{a + b\operatorname{arctanh}(cx^2)}{\sqrt{dx}} dx = -\frac{2b \arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt[4]{c}\sqrt{d}} - \frac{\sqrt{2}b \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt[4]{c}\sqrt{d}} + \frac{\sqrt{2}b \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt[4]{c}\sqrt{d}} + \frac{2\sqrt{dx}(a + b\operatorname{arctanh}(cx^2))}{d} - \frac{2b\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt[4]{c}\sqrt{d}} + \frac{\sqrt{2}b\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}(1+\sqrt{cx})}\right)}{\sqrt[4]{c}\sqrt{d}}$$

output

```
-2*b*arctan(c^(1/4)*(d*x)^(1/2)/d^(1/2))/c^(1/4)/d^(1/2)+2^(1/2)*b*arctan(-1+2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/2))/c^(1/4)/d^(1/2)+2^(1/2)*b*arctan(1+2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/2))/c^(1/4)/d^(1/2)+2*(d*x)^(1/2)*(a+b*arctanh(c*x^2))/d-2*b*arctanh(c^(1/4)*(d*x)^(1/2)/d^(1/2))/c^(1/4)/d^(1/2)+2^(1/2)*b*arctanh(2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/2)/(1+c^(1/2)*x))/c^(1/4)/d^(1/2)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{\sqrt{dx}} dx$$

$$= \frac{\sqrt{x}(4a\sqrt[4]{c}\sqrt{x} - 2\sqrt{2}b \arctan(1 - \sqrt{2}\sqrt[4]{c}\sqrt{x}) + 2\sqrt{2}b \arctan(1 + \sqrt{2}\sqrt[4]{c}\sqrt{x}) - 4b \arctan(\sqrt[4]{c}\sqrt{x}) + 4b \arctan(\sqrt[4]{c}\sqrt{x}))}{2\sqrt{d}}$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])/Sqrt[d*x], x]
```

output

```
(Sqrt[x]*(4*a*c^(1/4)*Sqrt[x] - 2*Sqrt[2]*b*ArcTan[1 - Sqrt[2]*c^(1/4)*Sqrt[x]] + 2*Sqrt[2]*b*ArcTan[1 + Sqrt[2]*c^(1/4)*Sqrt[x]] - 4*b*ArcTan[c^(1/4)*Sqrt[x]] + 4*b*c^(1/4)*Sqrt[x]*ArcTanh[c*x^2] + 2*b*Log[1 - c^(1/4)*Sqrt[x]] - 2*b*Log[1 + c^(1/4)*Sqrt[x]] - Sqrt[2]*b*Log[1 - Sqrt[2]*c^(1/4)*Sqrt[x] + Sqrt[c]*x] + Sqrt[2]*b*Log[1 + Sqrt[2]*c^(1/4)*Sqrt[x] + Sqrt[c]*x]))/(2*c^(1/4)*Sqrt[d*x])
```

Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.41, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {6464, 851, 27, 830, 755, 756, 218, 221, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{\sqrt{dx}} dx$$

$$\downarrow 6464$$

$$\frac{2\sqrt{dx}(a + b \operatorname{arctanh}(cx^2))}{d} - \frac{4bc \int \frac{(dx)^{3/2}}{1-c^2x^4} dx}{d^2}$$

$$\downarrow 851$$

$$\frac{2\sqrt{dx}(a + b \operatorname{arctanh}(cx^2))}{d} - \frac{8bc \int \frac{d^6 x^2}{d^4 - c^2 d^4 x^4} d\sqrt{dx}}{d^3}$$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{2\sqrt{dx}(a + \operatorname{barctanh}(cx^2))}{d} - 8bcd \int \frac{d^2 x^2}{d^4 - c^2 d^4 x^4} d\sqrt{dx} \\
& \downarrow 830 \\
& \frac{2\sqrt{dx}(a + \operatorname{barctanh}(cx^2))}{d} - 8bcd \left(\frac{\int \frac{1}{d^2 - cd^2 x^2} d\sqrt{dx}}{2c} - \frac{\int \frac{1}{cx^2 d^2 + d^2} d\sqrt{dx}}{2c} \right) \\
& \downarrow 755 \\
& \frac{2\sqrt{dx}(a + \operatorname{barctanh}(cx^2))}{d} - 8bcd \left(\frac{\int \frac{1}{d^2 - cd^2 x^2} d\sqrt{dx}}{2c} - \frac{\int \frac{d - \sqrt{cdx}}{cx^2 d^2 + d^2} d\sqrt{dx}}{2d} + \frac{\int \frac{\sqrt{cdx} + d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2c} \right) \\
& \downarrow 756 \\
& \frac{2\sqrt{dx}(a + \operatorname{barctanh}(cx^2))}{d} - 8bcd \left(\frac{\int \frac{1}{d - \sqrt{cdx}} d\sqrt{dx}}{2c} + \frac{\int \frac{1}{\sqrt{cdx} + d} d\sqrt{dx}}{2c} - \frac{\int \frac{d - \sqrt{cdx}}{cx^2 d^2 + d^2} d\sqrt{dx}}{2c} + \frac{\int \frac{\sqrt{cdx} + d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2c} \right) \\
& \downarrow 218 \\
& \frac{2\sqrt{dx}(a + \operatorname{barctanh}(cx^2))}{d} - 8bcd \left(\frac{\int \frac{1}{d - \sqrt{cdx}} d\sqrt{dx}}{2c} + \frac{\arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{cd^{3/2}}} - \frac{\int \frac{d - \sqrt{cdx}}{cx^2 d^2 + d^2} d\sqrt{dx}}{2c} + \frac{\int \frac{\sqrt{cdx} + d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2c} \right) \\
& \downarrow 221 \\
& \frac{2\sqrt{dx}(a + \operatorname{barctanh}(cx^2))}{d} - 8bcd \left(\frac{\arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{cd^{3/2}}} - \frac{\int \frac{d - \sqrt{cdx}}{cx^2 d^2 + d^2} d\sqrt{dx}}{2c} + \frac{\int \frac{\sqrt{cdx} + d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2c} \right) \\
& \downarrow 1476
\end{aligned}$$

$$8bcd \left(\frac{\frac{2\sqrt{dx}(a + b\operatorname{arctanh}(cx^2))}{d}}{2c} + \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2d} + \frac{\int \frac{1}{xd+\frac{d}{\sqrt{c}}-\sqrt{2}\sqrt{dx}\sqrt{d}} d\sqrt{dx}}{2\sqrt{c}} + \frac{\int \frac{1}{xd+\frac{d}{\sqrt{c}}+\sqrt{2}\sqrt{dx}\sqrt{d}} d\sqrt{dx}}{2\sqrt{c}}}{2c} \right)$$

1082

$$8bcd \left(\frac{\frac{2\sqrt{dx}(a + b\operatorname{arctanh}(cx^2))}{d}}{2c} + \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2d} + \frac{\int \frac{1}{-dx-1} d\left(1-\sqrt{2}\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\int \frac{1}{-dx-1} d\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}}}{2c} \right)$$

217

$$8bcd \left(\frac{\frac{2\sqrt{dx}(a + b\operatorname{arctanh}(cx^2))}{d}}{2c} + \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2d} + \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\operatorname{arctan}\left(1-\sqrt{2}\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}}}{2c} \right)$$

1479

$$8bcd \left(\frac{\frac{2\sqrt{dx}(a + b\operatorname{arctanh}(cx^2))}{d}}{2c} + \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} - \frac{\int -\frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{C}\sqrt{dx}}{\sqrt[4]{C}\left(xd+\frac{d}{\sqrt{c}}-\sqrt{2}\sqrt{dx}\sqrt{d}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\int -\frac{\sqrt{2}(\sqrt{d}+\sqrt{2}\sqrt[4]{C}\sqrt{dx})}{\sqrt[4]{C}\left(xd+\frac{d}{\sqrt{c}}+\sqrt{2}\sqrt{dx}\sqrt{d}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}}}{2c} + \dots \right)$$

25

$$8bcd \left(\frac{2\sqrt{dx}(a + b\operatorname{arctanh}(cx^2))}{d} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} - \frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{C}\sqrt{dx}}{\sqrt[4]{C}\left(xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}\right)}d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} + \frac{\int \frac{\sqrt{2}(\sqrt{d}+\sqrt{2}\sqrt[4]{C}\sqrt{dx})}{\sqrt[4]{C}\left(xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}\right)}d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} + \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right)$$

27

$$8bcd \left(\frac{2\sqrt{dx}(a + b\operatorname{arctanh}(cx^2))}{d} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} - \frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{C}\sqrt{dx}}{xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}}d\sqrt{dx}}{2\sqrt{2}\sqrt{c}\sqrt{d}} + \frac{\int \frac{\sqrt{d}+\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}}d\sqrt{dx}}{2\sqrt{c}\sqrt{d}} + \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right)$$

1103

$$8bcd \left(\frac{2\sqrt{dx}(a + b\operatorname{arctanh}(cx^2))}{d} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} - \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\operatorname{arctan}\left(1-\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} + \frac{\log\left(\sqrt{c}dx+\sqrt{2}\sqrt[4]{C}\sqrt{d}\sqrt{dx}+d\right)}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right)$$

input `Int[(a + b*ArcTanh[c*x^2])/Sqrt[d*x], x]`

output
$$\begin{aligned} & (2\sqrt{d*x}*(a + b*\text{ArcTanh}[c*x^2]))/d - 8*b*c*d*((\text{ArcTan}[(c^{1/4})*\sqrt{d*x}]/\sqrt{d}]/(2*c^{1/4}*d^{3/2})) + \text{ArcTanh}[(c^{1/4})*\sqrt{d*x}]/\sqrt{d}]/(2*c^{1/4}*d^{3/2}))/ (2*c) - ((-\text{ArcTan}[1 - (\sqrt{2}*c^{1/4})*\sqrt{d*x}]/\sqrt{d}]/(\sqrt{2}*c^{1/4})*\sqrt{d}))/ (2*d) + \text{ArcTan}[1 + (\sqrt{2}*c^{1/4})*\sqrt{d*x}]/\sqrt{d}]/(\sqrt{2}*c^{1/4})*\sqrt{d}))/ (2*d) + (-1/2*\text{Log}[d + \sqrt{c}*d*x - \sqrt{2}*c^{1/4}*\sqrt{d}*\sqrt{d*x}]/(\sqrt{2}*c^{1/4})*\sqrt{d}))/ (2*d) + \text{Log}[d + \sqrt{c}*d*x + \sqrt{2}*c^{1/4}*\sqrt{d}*\sqrt{d*x}]/(2*\sqrt{2}*c^{1/4}*\sqrt{d}))/ (2*d))/ (2*c) \end{aligned}$$

Defintions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 27 $\text{Int}[(a_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] /; \text{FreeQ}[a, \text{x}] \&\& \text{!MatchQ}[\text{Fx}, (b_)*(\text{Gx}_) /; \text{FreeQ}[b, \text{x}]$

rule 217 $\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& \text{PosQ}[a/b] \&\& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])$

rule 218 $\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& \text{PosQ}[a/b]$

rule 221 $\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& \text{NegQ}[a/b]$

rule 755 $\text{Int}[((a_) + (b_.)*(x_)^4)^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*r) \quad \text{Int}[(r - s*x^2)/(a + b*x^4), \text{x}], \text{x}] + \text{Simp}[1/(2*r) \quad \text{Int}[(r + s*x^2)/(a + b*x^4), \text{x}], \text{x}]] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& (\text{GtQ}[a/b, 0] \parallel (\text{PosQ}[a/b] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

rule 756 $\text{Int}[(a_ + (b_ \cdot x^4)^{-1}), x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[r/(2*a) \text{ Int}[1/(r - s*x^2), x], x] + \text{Simp}[r/(2*a) \text{ Int}[1/(r + s*x^2), x], x]] /; \text{FreeQ}[\{a, b\}, x] \&\& !\text{GtQ}[a/b, 0]$

rule 830 $\text{Int}[(x_)^{(m_)}/((a_ + (b_ \cdot x_)^{(n_)})], x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[s/(2*b) \text{ Int}[x^{(m - n/2)}/(r + s*x^{(n/2)})], x], x] - \text{Simp}[s/(2*b) \text{ Int}[x^{(m - n/2)}/(r - s*x^{(n/2)})], x], x]] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{IGtQ}[n/4, 0] \&\& \text{IGtQ}[m, 0] \&\& \text{LeQ}[n/2, m] \&\& \text{LtQ}[m, n] \&\& !\text{GtQ}[a/b, 0]$

rule 851 $\text{Int}[(c_ \cdot x_)^{(m_)} \cdot ((a_ + (b_ \cdot x_)^{(n_)})^{(p_)}], x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \text{ Subst}[\text{Int}[x^{(k*(m + 1) - 1)} \cdot (a + b \cdot x^{(k*n)/c^n})^p, x], x, (c*x)^{1/k}], x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{FractionQ}[m] \&\& \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 1082 $\text{Int}[(a_ + (b_ \cdot x_) + (c_ \cdot x_)^2)^{-1}], x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S \text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2)], x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \parallel !\text{RationalQ}[b^2 - 4*a*c]) /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}[(d_ + (e_ \cdot x_))/((a_ + (b_ \cdot x_) + (c_ \cdot x_)^2)], x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$

rule 1476 $\text{Int}[(d_ + (e_ \cdot x_)^2)/((a_ + (c_ \cdot x_)^4)], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2*(d/e), 2]\}, \text{Simp}[e/(2*c) \text{ Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Simp}[e/(2*c) \text{ Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[d*e]$

rule 1479 $\text{Int}[(d_ + (e_ \cdot x_)^2)/((a_ + (c_ \cdot x_)^4)], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2*(d/e), 2]\}, \text{Simp}[e/(2*c*q) \text{ Int}[(q - 2*x)/\text{Simp}[d/e + q*x - x^2, x], x], x] + \text{Simp}[e/(2*c*q) \text{ Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{NegQ}[d*e]$

rule 6464

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))*((d_)*(x_)^(m_), x_Symbol] :
> Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n
/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a,
b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 242, normalized size of antiderivative = 1.07

method	result
derivativedivides	$2\sqrt{dx} a+2b \left(\sqrt{dx} \operatorname{arctanh}(cx^2) - 4cd^2 \left(\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{2} \left(\ln \left(\frac{dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}}{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 1} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} - 1} \right) \right) \right) \frac{d}{16cd^2}$
default	$2\sqrt{dx} a+2b \left(\sqrt{dx} \operatorname{arctanh}(cx^2) - 4cd^2 \left(\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{2} \left(\ln \left(\frac{dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}}{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 1} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} - 1} \right) \right) \right) \frac{d}{16cd^2}$
parts	$\frac{2a\sqrt{dx}}{d} + 2b \left(\sqrt{dx} \operatorname{arctanh}(cx^2) - 4cd^2 \left(\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{2} \left(\ln \left(\frac{dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}}{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 1} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} - 1} \right) \right) \right) \frac{d}{16cd^2}$

input

```
int((a+b*arctanh(c*x^2))/(d*x)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
2/d*((d*x)^(1/2)*a+b*((d*x)^(1/2)*arctanh(c*x^2)-4*c*d^2*(-1/16/c*(d^2/c)^(1/4)/d^2*2^(1/2)*(ln((d*x+(d^2/c)^(1/4)*(d*x)^(1/2)*2^(1/2)+(d^2/c)^(1/2)))/(d*x-(d^2/c)^(1/4)*(d*x)^(1/2)*2^(1/2)+(d^2/c)^(1/2))))+2*arctan(2^(1/2)/(d^2/c)^(1/4)*(d*x)^(1/2)+1)+2*arctan(2^(1/2)/(d^2/c)^(1/4)*(d*x)^(1/2)-1))+1/8/c*(d^2/c)^(1/4)/d^2*(ln(((d*x)^(1/2)+(d^2/c)^(1/4))/((d*x)^(1/2)-(d^2/c)^(1/4))))+2*arctan((d*x)^(1/2)/(d^2/c)^(1/4))))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 354, normalized size of antiderivative = 1.56

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{\sqrt{dx}} dx = \frac{\left(\frac{b^4}{cd^2}\right)^{\frac{1}{4}} d \log\left(\sqrt{dx}b + \left(\frac{b^4}{cd^2}\right)^{\frac{1}{4}} d\right) + i \left(\frac{b^4}{cd^2}\right)^{\frac{1}{4}} d \log\left(\sqrt{dx}b + i \left(\frac{b^4}{cd^2}\right)^{\frac{1}{4}} d\right) - i \left(\frac{b^4}{cd^2}\right)^{\frac{1}{4}} d \log\left(\sqrt{dx}b - i \left(\frac{b^4}{cd^2}\right)^{\frac{1}{4}} d\right)}{d}$$

input `integrate((a+b*arctanh(c*x^2))/(d*x)^(1/2),x, algorithm="fricas")`

output `-((b^4/(c*d^2))^(1/4)*d*log(sqrt(d*x)*b + (b^4/(c*d^2))^(1/4)*d) + I*(b^4/(c*d^2))^(1/4)*d*log(sqrt(d*x)*b + I*(b^4/(c*d^2))^(1/4)*d) - I*(b^4/(c*d^2))^(1/4)*d*log(sqrt(d*x)*b - I*(b^4/(c*d^2))^(1/4)*d) - (b^4/(c*d^2))^(1/4)*d*log(sqrt(d*x)*b - (b^4/(c*d^2))^(1/4)*d) - (-b^4/(c*d^2))^(1/4)*d*log(sqrt(d*x)*b + (-b^4/(c*d^2))^(1/4)*d) - I*(-b^4/(c*d^2))^(1/4)*d*log(sqrt(d*x)*b + I*(-b^4/(c*d^2))^(1/4)*d) + I*(-b^4/(c*d^2))^(1/4)*d*log(sqrt(d*x)*b - I*(-b^4/(c*d^2))^(1/4)*d) + (-b^4/(c*d^2))^(1/4)*d*log(sqrt(d*x)*b - (-b^4/(c*d^2))^(1/4)*d) - sqrt(d*x)*(b*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 2*a))/d`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{\sqrt{dx}} dx = \int \frac{a + b \operatorname{atanh}(cx^2)}{\sqrt{dx}} dx$$

input `integrate((a+b*atanh(c*x**2))/(d*x)**(1/2),x)`

output `Integral((a + b*atanh(c*x**2))/sqrt(d*x), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 296, normalized size of antiderivative = 1.30

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{\sqrt{dx}} dx$$

$$= \frac{4\sqrt{dx} \operatorname{artanh}(cx^2) + c \left(\frac{2\sqrt{2}d^3 \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}c^{\frac{1}{4}}\sqrt{d} + 2\sqrt{dx}\sqrt{c}\right)}{2\sqrt{\sqrt{cd}}}\right)}{\sqrt{\sqrt{cd}}} + \frac{2\sqrt{2}d^3 \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}c^{\frac{1}{4}}\sqrt{d} - 2\sqrt{dx}\sqrt{c}\right)}{2\sqrt{\sqrt{cd}}}\right)}{\sqrt{\sqrt{cd}}} + \frac{\sqrt{2}d^{\frac{5}{2}} \log\left(\sqrt{cdx} + \sqrt{2}\sqrt{\frac{d}{c^{\frac{1}{4}}}}\right)}{c} \right)}{d^2}$$

input `integrate((a+b*arctanh(c*x^2))/(d*x)^(1/2),x, algorithm="maxima")`

output `1/2*((4*sqrt(d*x)*arctanh(c*x^2) + c*((2*sqrt(2)*d^3*arctan(1/2*sqrt(2)*(sqrt(2)*c^(1/4)*sqrt(d) + 2*sqrt(d*x)*sqrt(c))/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d) + 2*sqrt(2)*d^3*arctan(-1/2*sqrt(2)*(sqrt(2)*c^(1/4)*sqrt(d) - 2*sqrt(d*x)*sqrt(c))/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d) + sqrt(2)*d^(5/2)*log(sqrt(c)*d*x + sqrt(2)*sqrt(d*x)*c^(1/4)*sqrt(d) + d)/c^(1/4) - sqrt(2)*d^(5/2)*log(sqrt(c)*d*x - sqrt(2)*sqrt(d*x)*c^(1/4)*sqrt(d) + d)/c^(1/4))/c - 2*(2*d^3*arctan(sqrt(d*x)*sqrt(c)/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d) - d^3*log((sqrt(d*x)*sqrt(c) - sqrt(sqrt(c)*d))/(sqrt(d*x)*sqrt(c) + sqrt(sqrt(c)*d)))/sqrt(sqrt(c)*d))/c)/d^2)*b + 4*sqrt(d*x)*a)/d`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 493 vs. $2(159) = 318$.

Time = 0.13 (sec) , antiderivative size = 493, normalized size of antiderivative = 2.17

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{\sqrt{dx}} dx$$

$$= \left(cd^2 \left(\frac{2\sqrt{2}(c^3d^2)^{\frac{1}{4}} \operatorname{arctan}\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 2\sqrt{dx}\right)}{2\left(\frac{d^2}{c}\right)^{\frac{1}{4}}}\right)}{c^2d^2} + \frac{2\sqrt{2}(c^3d^2)^{\frac{1}{4}} \operatorname{arctan}\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{d^2}{c}\right)^{\frac{1}{4}} - 2\sqrt{dx}\right)}{2\left(\frac{d^2}{c}\right)^{\frac{1}{4}}}\right)}{c^2d^2} - \frac{2\sqrt{2}(-c^3d^2)^{\frac{1}{4}} \operatorname{arctan}\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 2\sqrt{dx}\right)}{2\left(\frac{d^2}{c}\right)^{\frac{1}{4}}}\right)}{c^2d^2} \right) \right)$$

input `integrate((a+b*arctanh(c*x^2))/(d*x)^(1/2),x, algorithm="giac")`

output

```
1/2*((c*d^2*(2*sqrt(2)*(c^3*d^2)^(1/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(d^2/c)^(1/4) + 2*sqrt(d*x))/(d^2/c)^(1/4)))/(c^2*d^2) + 2*sqrt(2)*(c^3*d^2)^(1/4)*arctan(-1/2*sqrt(2)*(sqrt(2)*(d^2/c)^(1/4) - 2*sqrt(d*x))/(d^2/c)^(1/4)))/(c^2*d^2) - 2*sqrt(2)*(-c^3*d^2)^(1/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(-d^2/c)^(1/4) + 2*sqrt(d*x))/(-d^2/c)^(1/4)))/(c^2*d^2) - 2*sqrt(2)*(-c^3*d^2)^(1/4)*arctan(-1/2*sqrt(2)*(sqrt(2)*(-d^2/c)^(1/4) - 2*sqrt(d*x))/(-d^2/c)^(1/4)))/(c^2*d^2) + sqrt(2)*(c^3*d^2)^(1/4)*log(d*x + sqrt(2)*sqrt(d*x)*(d^2/c)^(1/4) + sqrt(d^2/c))/(c^2*d^2) - sqrt(2)*(c^3*d^2)^(1/4)*log(d*x - sqrt(2)*sqrt(d*x)*(d^2/c)^(1/4) + sqrt(d^2/c))/(c^2*d^2) - sqrt(2)*(-c^3*d^2)^(1/4)*log(d*x + sqrt(2)*sqrt(d*x)*(-d^2/c)^(1/4) + sqrt(-d^2/c))/(c^2*d^2) + sqrt(2)*(-c^3*d^2)^(1/4)*log(d*x - sqrt(2)*sqrt(d*x)*(-d^2/c)^(1/4) + sqrt(-d^2/c))/(c^2*d^2) + 2*sqrt(d*x)*log(-(c*x^2 + 1)/(c*x^2 - 1))*b + 4*sqrt(d*x)*a)/d
```

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{\sqrt{dx}} dx = \int \frac{a + b \operatorname{atanh}(cx^2)}{\sqrt{dx}} dx$$

input `int((a + b*atanh(c*x^2))/(d*x)^(1/2), x)`output `int((a + b*atanh(c*x^2))/(d*x)^(1/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{\sqrt{dx}} dx$$

$$= \sqrt{d} \left(-2c^{\frac{3}{4}} \sqrt{2} \operatorname{atan} \left(\frac{c^{\frac{1}{4}} \sqrt{2} - 2\sqrt{x} \sqrt{c}}{c^{\frac{1}{4}} \sqrt{2}} \right) b + 2c^{\frac{3}{4}} \sqrt{2} \operatorname{atan} \left(\frac{c^{\frac{1}{4}} \sqrt{2} + 2\sqrt{x} \sqrt{c}}{c^{\frac{1}{4}} \sqrt{2}} \right) b - 4c^{\frac{3}{4}} \operatorname{atan} \left(\frac{\sqrt{x} \sqrt{c}}{c^{\frac{1}{4}}} \right) b + 2c^{\frac{3}{4}} \sqrt{2} \operatorname{atanh} \right.$$

input `int((a+b*atanh(c*x^2))/(d*x)^(1/2), x)`output `(sqrt(d)*(-2*c**(3/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*b + 2*c**(3/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*b - 4*c**(3/4)*atan((sqrt(x)*sqrt(c))/c**(1/4))*b + 2*c**(3/4)*sqrt(2)*atanh(c*x**2)*b + 4*sqrt(x)*atanh(c*x**2)*b*c + c**(3/4)*sqrt(2)*log(c**(1/4) + sqrt(x)*sqrt(c))*b + c**(3/4)*sqrt(2)*log(-c**(1/4) + sqrt(x)*sqrt(c))*b - 2*c**(3/4)*sqrt(2)*log(-sqrt(x)*c**(1/4)*sqrt(2) + sqrt(c)*x + 1)*b + c**(3/4)*sqrt(2)*log(sqrt(c)*x + 1)*b - 2*c**(3/4)*log(c**(1/4) + sqrt(x)*sqrt(c))*b + 2*c**(3/4)*log(-c**(1/4) + sqrt(x)*sqrt(c))*b + 4*sqrt(x)*a*c))/(2*c*d)`

3.86 $\int \frac{a+b\operatorname{arctanh}(cx^2)}{(dx)^{3/2}} dx$

Optimal result	745
Mathematica [A] (verified)	746
Rubi [A] (verified)	746
Maple [A] (verified)	752
Fricas [C] (verification not implemented)	753
Sympy [F]	754
Maxima [A] (verification not implemented)	754
Giac [B] (verification not implemented)	755
Mupad [F(-1)]	756
Reduce [B] (verification not implemented)	756

Optimal result

Integrand size = 18, antiderivative size = 228

$$\int \frac{a + b\operatorname{arctanh}(cx^2)}{(dx)^{3/2}} dx = -\frac{2b\sqrt[4]{c} \arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{d^{3/2}} - \frac{\sqrt{2}b\sqrt[4]{c} \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{d^{3/2}} + \frac{\sqrt{2}b\sqrt[4]{c} \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{d^{3/2}} - \frac{2(a + b\operatorname{arctanh}(cx^2))}{d\sqrt{dx}} + \frac{2b\sqrt[4]{c} \operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{d^{3/2}} - \frac{\sqrt{2}b\sqrt[4]{c} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}(1+\sqrt{cx})}\right)}{d^{3/2}}$$

output

```
-2*b*c^(1/4)*arctan(c^(1/4)*(d*x)^(1/2)/d^(1/2))/d^(3/2)+2^(1/2)*b*c^(1/4)
*arctan(-1+2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/2))/d^(3/2)+2^(1/2)*b*c^(1/4)*
arctan(1+2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/2))/d^(3/2)-2*(a+b*arctanh(c*x^2
))/d/(d*x)^(1/2)+2*b*c^(1/4)*arctanh(c^(1/4)*(d*x)^(1/2)/d^(1/2))/d^(3/2)-
2^(1/2)*b*c^(1/4)*arctanh(2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/2)/(1+c^(1/2)*x
))/d^(3/2)
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.18

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{3/2}} dx =$$

$$x(4a + 2\sqrt{2}b\sqrt[4]{c}\sqrt{x} \arctan(1 - \sqrt{2}\sqrt[4]{c}\sqrt{x}) - 2\sqrt{2}b\sqrt[4]{c}\sqrt{x} \arctan(1 + \sqrt{2}\sqrt[4]{c}\sqrt{x}) + 4b\sqrt[4]{c}\sqrt{x} \arctan(\sqrt[4]{c}\sqrt{x})) / (dx)^{3/2}$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])/(d*x)^(3/2), x]
```

output

```
-1/2*(x*(4*a + 2*Sqrt[2]*b*c^(1/4)*Sqrt[x]*ArcTan[1 - Sqrt[2]*c^(1/4)*Sqrt[x]] - 2*Sqrt[2]*b*c^(1/4)*Sqrt[x]*ArcTan[1 + Sqrt[2]*c^(1/4)*Sqrt[x]] + 4*b*c^(1/4)*Sqrt[x]*ArcTan[c^(1/4)*Sqrt[x]] + 4*b*ArcTanh[c*x^2] + 2*b*c^(1/4)*Sqrt[x]*Log[1 - c^(1/4)*Sqrt[x]] - 2*b*c^(1/4)*Sqrt[x]*Log[1 + c^(1/4)*Sqrt[x]] - Sqrt[2]*b*c^(1/4)*Sqrt[x]*Log[1 - Sqrt[2]*c^(1/4)*Sqrt[x]] + Sqrt[c]*x + Sqrt[2]*b*c^(1/4)*Sqrt[x]*Log[1 + Sqrt[2]*c^(1/4)*Sqrt[x]] + Sqrt[c]*x)))/(d*x)^(3/2)
```

Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 324, normalized size of antiderivative = 1.42, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {6464, 851, 27, 829, 826, 827, 218, 221, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{3/2}} dx$$

$$\downarrow 6464$$

$$\frac{4bc \int \frac{\sqrt{dx}}{1-c^2x^4} dx}{d^2} - \frac{2(a + b \operatorname{arctanh}(cx^2))}{d\sqrt{dx}}$$

$$\downarrow 851$$

$$\begin{aligned}
& \frac{8bc \int \frac{d^5 x}{d^4 - c^2 d^4 x^4} d\sqrt{dx}}{d^3} - \frac{2(a + \operatorname{barctanh}(cx^2))}{d\sqrt{dx}} \\
& \quad \downarrow 27 \\
& 8bcd \int \frac{dx}{d^4 - c^2 d^4 x^4} d\sqrt{dx} - \frac{2(a + \operatorname{barctanh}(cx^2))}{d\sqrt{dx}} \\
& \quad \downarrow 829 \\
& 8bcd \left(\frac{\int \frac{dx}{d^2 - cd^2 x^2} d\sqrt{dx}}{2d^2} + \frac{\int \frac{dx}{cx^2 d^2 + d^2} d\sqrt{dx}}{2d^2} \right) - \frac{2(a + \operatorname{barctanh}(cx^2))}{d\sqrt{dx}} \\
& \quad \downarrow 826 \\
& 8bcd \left(\frac{\int \frac{dx}{d^2 - cd^2 x^2} d\sqrt{dx}}{2d^2} + \frac{\int \frac{\sqrt{cx}d+d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} \right) - \frac{2(a + \operatorname{barctanh}(cx^2))}{d\sqrt{dx}} \\
& \quad \downarrow 827 \\
& 8bcd \left(\frac{\int \frac{\sqrt{cx}d+d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} + \frac{\int \frac{1}{d-\sqrt{cdx}} d\sqrt{dx}}{2d^2} - \frac{\int \frac{1}{\sqrt{cx}d+d} d\sqrt{dx}}{2d^2} \right) - \\
& \quad \frac{2(a + \operatorname{barctanh}(cx^2))}{d\sqrt{dx}} \\
& \quad \downarrow 218 \\
& 8bcd \left(\frac{\int \frac{1}{d-\sqrt{cdx}} d\sqrt{dx}}{2\sqrt{c}} - \frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} + \frac{\int \frac{\sqrt{cx}d+d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} \right) - \\
& \quad \frac{2(a + \operatorname{barctanh}(cx^2))}{d\sqrt{dx}} \\
& \quad \downarrow 221 \\
& 8bcd \left(\frac{\int \frac{\sqrt{cx}d+d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} \right) - \\
& \quad \frac{2(a + \operatorname{barctanh}(cx^2))}{d\sqrt{dx}}
\end{aligned}$$

$$\begin{aligned}
 & \downarrow 1476 \\
 8bcd & \left(\frac{\int \frac{1}{xd + \frac{d}{\sqrt{c}} - \frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}} d\sqrt{dx}}{2\sqrt{c}} + \frac{\int \frac{1}{xd + \frac{d}{\sqrt{c}} + \frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d - \sqrt{c}dx}{cx^2d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right) - \operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} \right) \\
 & \frac{2(a + \operatorname{barctanh}(cx^2))}{d\sqrt{dx}} \\
 & \downarrow 1082 \\
 8bcd & \left(\frac{\int \frac{1}{-dx-1} d\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{1}{-dx-1} d\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}} + 1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{d - \sqrt{c}dx}{cx^2d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right) - \operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} \right) \\
 & \frac{2(a + \operatorname{barctanh}(cx^2))}{d\sqrt{dx}} \\
 & \downarrow 217 \\
 8bcd & \left(\frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}} + 1\right) - \operatorname{arctan}\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{d - \sqrt{c}dx}{cx^2d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right) - \operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} \right) \\
 & \frac{2(a + \operatorname{barctanh}(cx^2))}{d\sqrt{dx}} \\
 & \downarrow 1479
 \end{aligned}$$

$$8bcd \left(\frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}}}{2\sqrt{c}} - \frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{C}\sqrt{dx}}{\sqrt[4]{C}\left(xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\int \frac{\sqrt{2}(\sqrt{d}+\sqrt{2}\sqrt[4]{C}\sqrt{dx})}{\sqrt[4]{C}\left(xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right) + \dots$$

$$\frac{2(a + b \operatorname{arctanh}(cx^2))}{d\sqrt{dx}}$$

25

$$8bcd \left(\frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}}}{2\sqrt{c}} - \frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{C}\sqrt{dx}}{\sqrt[4]{C}\left(xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} + \frac{\int \frac{\sqrt{2}(\sqrt{d}+\sqrt{2}\sqrt[4]{C}\sqrt{dx})}{\sqrt[4]{C}\left(xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right) + \dots$$

$$\frac{2(a + b \operatorname{arctanh}(cx^2))}{d\sqrt{dx}}$$

27

$$8bcd \left(\frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}}}{2\sqrt{c}} - \frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{C}\sqrt{dx}}{xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}} d\sqrt{dx}}{2\sqrt{2}\sqrt{c}\sqrt{d}} + \frac{\int \frac{\sqrt{d}+\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}} d\sqrt{dx}}{2\sqrt{c}\sqrt{d}} \right) + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}}$$

$$\frac{2(a + b \operatorname{arctanh}(cx^2))}{d\sqrt{dx}}$$

1103

$$8bcd \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} + \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\operatorname{arctan}\left(1-\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\log\left(\sqrt{cdx}+\sqrt{2}\sqrt[4]{c}\sqrt{d}\sqrt{dx}+d\right)}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} \right) - \frac{2(a + b\operatorname{arctanh}(cx^2))}{d\sqrt{dx}}$$

input `Int[(a + b*ArcTanh[c*x^2])/(d*x)^(3/2), x]`

output $(-2*(a + b*\operatorname{ArcTanh}[c*x^2]))/(d*\operatorname{Sqrt}[d*x]) + 8*b*c*d*((-1/2*\operatorname{ArcTan}[(c^{1/4})*\operatorname{Sqrt}[d*x])/ \operatorname{Sqrt}[d]]/(\operatorname{Sqrt}[d]) + \operatorname{ArcTanh}[(c^{1/4})*\operatorname{Sqrt}[d*x])/ \operatorname{Sqrt}[d]]/(2*c^{3/4}*\operatorname{Sqrt}[d]))/(2*d^2) + ((-\operatorname{ArcTan}[1 - (\operatorname{Sqrt}[2]*c^{1/4})*\operatorname{Sqrt}[d*x])/ \operatorname{Sqrt}[d]]/(\operatorname{Sqrt}[2]*c^{1/4}*\operatorname{Sqrt}[d])) + \operatorname{ArcTan}[1 + (\operatorname{Sqrt}[2]*c^{1/4})*\operatorname{Sqrt}[d*x])/ \operatorname{Sqrt}[d]]/(\operatorname{Sqrt}[2]*c^{1/4}*\operatorname{Sqrt}[d]))/(2*\operatorname{Sqrt}[c]) - (-1/2*\operatorname{Log}[d + \operatorname{Sqrt}[c]*d*x - \operatorname{Sqrt}[2]*c^{1/4}*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[d*x]]/(\operatorname{Sqrt}[2]*c^{1/4}*\operatorname{Sqrt}[d]) + \operatorname{Log}[d + \operatorname{Sqrt}[c]*d*x + \operatorname{Sqrt}[2]*c^{1/4}*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[d*x]]/(\operatorname{Sqrt}[2]*c^{1/4}*\operatorname{Sqrt}[d]))/(2*\operatorname{Sqrt}[c]))/(2*d^2)$

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 221 $\text{Int}[(a_ + (b_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) \cdot \text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b]$

rule 826 $\text{Int}[(x_)^2/((a_ + (b_ \cdot)(x_)^4), x_Symbol] \rightarrow \text{With}\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*s) \ \text{Int}[(r + s*x^2)/(a + b*x^4), x], x] - \text{Simp}[1/(2*s) \ \text{Int}[(r - s*x^2)/(a + b*x^4), x], x]] /; \text{FreeQ}\{a, b, x\} \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

rule 827 $\text{Int}[(x_)^2/((a_ + (b_ \cdot)(x_)^4), x_Symbol] \rightarrow \text{With}\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[s/(2*b) \ \text{Int}[1/(r + s*x^2), x], x] - \text{Simp}[s/(2*b) \ \text{Int}[1/(r - s*x^2), x], x]] /; \text{FreeQ}\{a, b, x\} \ \&\& \ !\text{GtQ}[a/b, 0]$

rule 829 $\text{Int}[(x_)^m/((a_ + (b_ \cdot)(x_)^n), x_Symbol] \rightarrow \text{With}\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[r/(2*a) \ \text{Int}[x^m/(r + s*x^{n/2}), x], x] + \text{Simp}[r/(2*a) \ \text{Int}[x^m/(r - s*x^{n/2}), x], x]] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{IGtQ}[n/4, 0] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{LtQ}[m, n/2] \ \&\& \ !\text{GtQ}[a/b, 0]$

rule 851 $\text{Int}[(c_ \cdot)(x_)^m \cdot ((a_ + (b_ \cdot)(x_)^n)^p), x_Symbol] \rightarrow \text{With}\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \ \text{Subst}[\text{Int}[x^{(k*(m+1)-1)} \cdot (a + b \cdot (x^{k*n})/c^n)^p, x], x, (c*x)^{1/k}], x]] /; \text{FreeQ}\{a, b, c, p, x\} \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 1082 $\text{Int}[(a_ + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}\{q = 1 - 4*S \ \text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c]) /; \text{FreeQ}\{a, b, c, x\}$

rule 1103 $\text{Int}[(d_ + (e_ \cdot)(x_))/((a_ + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$

```
rule 1476 Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

```
rule 1479 Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

```
rule 6464 Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)*((d_)*(x_)^(m_)), x_Symbol] :
> Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n
/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a,
b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.08

method	result
derivativedivides	$-\frac{2a}{\sqrt{dx}} + 2b \left(-\frac{\operatorname{arctanh}(cx^2)}{\sqrt{dx}} + 4cd^2 \frac{\sqrt{2} \left(\ln \left(\frac{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}}{dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}}\right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 1} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} - 1} \right)}{16d^2 c \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)$
default	$-\frac{2a}{\sqrt{dx}} + 2b \left(-\frac{\operatorname{arctanh}(cx^2)}{\sqrt{dx}} + 4cd^2 \frac{\sqrt{2} \left(\ln \left(\frac{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}}{dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}}\right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 1} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} - 1} \right)}{16d^2 c \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)$
parts	$-\frac{2a}{\sqrt{dx} d} + \frac{2b \left(-\frac{\operatorname{arctanh}(cx^2)}{\sqrt{dx}} + 4cd^2 \frac{\sqrt{2} \left(\ln \left(\frac{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}}{dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}}\right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 1} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} - 1} \right)}{16d^2 c \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)}{d}$

input `int((a+b*arctanh(c*x^2))/(d*x)^(3/2),x,method=_RETURNVERBOSE)`

output
$$\frac{2/d*(-a/(d*x)^{(1/2)}+b*(-1/(d*x)^{(1/2)}*arctanh(c*x^2)+4*c*d^2*(1/16/d^2/c/(d^2/c)^{(1/4)}*2^{(1/2)}*(ln((d*x-(d^2/c)^{(1/4)}*(d*x)^{(1/2)}*2^{(1/2)}+(d^2/c)^{(1/2)))/(d*x+(d^2/c)^{(1/4)}*(d*x)^{(1/2)}*2^{(1/2)}+(d^2/c)^{(1/2))))+2*arctan(2^{(1/2)}/(d^2/c)^{(1/4)}*(d*x)^{(1/2)}+1)+2*arctan(2^{(1/2)}/(d^2/c)^{(1/4)}*(d*x)^{(1/2)}-1))-1/8/d^2/c/(d^2/c)^{(1/4)}*(2*arctan((d*x)^{(1/2)}/(d^2/c)^{(1/4)})-ln(((d*x)^{(1/2)}+(d^2/c)^{(1/4)))/((d*x)^{(1/2)}-(d^2/c)^{(1/4)))))}{(d*x)^{(3/2)}}$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 388, normalized size of antiderivative = 1.70

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{3/2}} dx = \frac{d^2 x \left(\frac{b^4 c}{d^6}\right)^{\frac{1}{4}} \log\left(d^5 \left(\frac{b^4 c}{d^6}\right)^{\frac{3}{4}} + \sqrt{dxb^3c}\right) - i d^2 x \left(\frac{b^4 c}{d^6}\right)^{\frac{1}{4}} \log\left(i d^5 \left(\frac{b^4 c}{d^6}\right)^{\frac{3}{4}} + \sqrt{dxb^3c}\right)}{(dx)^{3/2}}$$

input `integrate((a+b*arctanh(c*x^2))/(d*x)^(3/2),x, algorithm="fricas")`

output
$$\frac{(d^2*x*(b^4*c/d^6)^{(1/4)}*\log(d^5*(b^4*c/d^6)^{(3/4)} + \operatorname{sqrt}(d*x)*b^3*c) - I*d^2*x*(b^4*c/d^6)^{(1/4)}*\log(I*d^5*(b^4*c/d^6)^{(3/4)} + \operatorname{sqrt}(d*x)*b^3*c) + I*d^2*x*(b^4*c/d^6)^{(1/4)}*\log(-I*d^5*(b^4*c/d^6)^{(3/4)} + \operatorname{sqrt}(d*x)*b^3*c) - d^2*x*(b^4*c/d^6)^{(1/4)}*\log(-d^5*(b^4*c/d^6)^{(3/4)} + \operatorname{sqrt}(d*x)*b^3*c) + d^2*x*(-b^4*c/d^6)^{(1/4)}*\log(d^5*(-b^4*c/d^6)^{(3/4)} + \operatorname{sqrt}(d*x)*b^3*c) - I*d^2*x*(-b^4*c/d^6)^{(1/4)}*\log(I*d^5*(-b^4*c/d^6)^{(3/4)} + \operatorname{sqrt}(d*x)*b^3*c) + I*d^2*x*(-b^4*c/d^6)^{(1/4)}*\log(-I*d^5*(-b^4*c/d^6)^{(3/4)} + \operatorname{sqrt}(d*x)*b^3*c) - d^2*x*(-b^4*c/d^6)^{(1/4)}*\log(-d^5*(-b^4*c/d^6)^{(3/4)} + \operatorname{sqrt}(d*x)*b^3*c) - \operatorname{sqrt}(d*x)*(b*\log(-(c*x^2 + 1)/(c*x^2 - 1)) + 2*a))/(d^2*x)}{(d^2*x)}$$

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{3/2}} dx = \int \frac{a + b \operatorname{atanh}(cx^2)}{(dx)^{\frac{3}{2}}} dx$$

input `integrate((a+b*atanh(c*x**2))/(d*x)**(3/2), x)`

output `Integral((a + b*atanh(c*x**2))/(d*x)**(3/2), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 296, normalized size of antiderivative = 1.30

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{3/2}} dx =$$

$$b \left(\frac{4 \operatorname{arctanh}(cx^2)}{\sqrt{dx}} - \frac{d^2 \left(\frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}c^{\frac{1}{4}}\sqrt{d}+2\sqrt{dx}\sqrt{c}\right)}{2\sqrt{\sqrt{cd}}}\right)}{\sqrt{\sqrt{cd}}\sqrt{c}} \right) + \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}c^{\frac{1}{4}}\sqrt{d}-2\sqrt{dx}\sqrt{c}\right)}{2\sqrt{\sqrt{cd}}}\right)}{\sqrt{\sqrt{cd}}\sqrt{c}} - \frac{\sqrt{2} \log\left(\sqrt{cd}x + \sqrt{2}\sqrt{dx}c^{\frac{1}{4}}\sqrt{d}+d\right)}{c^{\frac{3}{4}}\sqrt{d}}}{d^2} \right)$$

input `integrate((a+b*arctanh(c*x^2))/(d*x)^(3/2), x, algorithm="maxima")`

output

```
-1/2*(b*(4*arctanh(c*x^2)/sqrt(d*x) - (d^2*(2*sqrt(2)*arctan(1/2*sqrt(2)*
sqrt(2)*c^(1/4)*sqrt(d) + 2*sqrt(d*x)*sqrt(c))/sqrt(sqrt(c)*d))/sqrt(sqrt
(c)*d)*sqrt(c)) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*c^(1/4)*sqrt(d) -
2*sqrt(d*x)*sqrt(c))/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d)*sqrt(c) - sqrt(2)
*log(sqrt(c)*d*x + sqrt(2)*sqrt(d*x)*c^(1/4)*sqrt(d) + d)/(c^(3/4)*sqrt(d)
) + sqrt(2)*log(sqrt(c)*d*x - sqrt(2)*sqrt(d*x)*c^(1/4)*sqrt(d) + d)/(c^(3
/4)*sqrt(d))) - 2*d^2*(2*arctan(sqrt(d*x)*sqrt(c)/sqrt(sqrt(c)*d))/sqrt(s
qrt(c)*d)*sqrt(c) + log((sqrt(d*x)*sqrt(c) - sqrt(sqrt(c)*d))/sqrt(d*x)*
sqrt(c) + sqrt(sqrt(c)*d)))/sqrt(sqrt(c)*d)*sqrt(c))) * c/d^2 + 4*a/sqrt(
d*x))/d
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 505 vs. $2(160) = 320$.

Time = 0.29 (sec) , antiderivative size = 505, normalized size of antiderivative = 2.21

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{3/2}} dx =$$

$$\frac{2b \log\left(-\frac{cd^2x^2+d^2}{cd^2x^2-d^2}\right)}{\sqrt{dx}} + \frac{4a}{\sqrt{dx}} - \frac{2\sqrt{2}(c^3d^2)^{\frac{3}{4}}b \operatorname{arctan}\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{d^2}{c}\right)^{\frac{1}{4}}+2\sqrt{dx}\right)}{2\left(\frac{d^2}{c}\right)^{\frac{1}{4}}}\right)}{c^2d^2} - \frac{2\sqrt{2}(c^3d^2)^{\frac{3}{4}}b \operatorname{arctan}\left(-\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{d^2}{c}\right)^{\frac{1}{4}}-2\sqrt{dx}\right)}{2\left(\frac{d^2}{c}\right)^{\frac{1}{4}}}\right)}{c^2d^2}$$

input

```
integrate((a+b*arctanh(c*x^2))/(d*x)^(3/2),x, algorithm="giac")
```


output

```
-1/2*(2*b*log(-(c*d^2*x^2 + d^2)/(c*d^2*x^2 - d^2))/sqrt(d*x) + 4*a/sqrt(d
*x) - 2*sqrt(2)*(c^3*d^2)^(3/4)*b*arctan(1/2*sqrt(2)*(sqrt(2)*(d^2/c)^(1/4
) + 2*sqrt(d*x))/(d^2/c)^(1/4))/(c^2*d^2) - 2*sqrt(2)*(c^3*d^2)^(3/4)*b*ar
ctan(-1/2*sqrt(2)*(sqrt(2)*(d^2/c)^(1/4) - 2*sqrt(d*x))/(d^2/c)^(1/4))/(c^
2*d^2) - 2*sqrt(2)*(-c^3*d^2)^(3/4)*b*arctan(1/2*sqrt(2)*(sqrt(2)*(-d^2/c)
^(1/4) + 2*sqrt(d*x))/(-d^2/c)^(1/4))/(c^2*d^2) - 2*sqrt(2)*(-c^3*d^2)^(3/
4)*b*arctan(-1/2*sqrt(2)*(sqrt(2)*(-d^2/c)^(1/4) - 2*sqrt(d*x))/(-d^2/c)^(
1/4))/(c^2*d^2) + sqrt(2)*(c^3*d^2)^(3/4)*b*log(d*x + sqrt(2)*sqrt(d*x)*(d
^2/c)^(1/4) + sqrt(d^2/c))/(c^2*d^2) - sqrt(2)*(c^3*d^2)^(3/4)*b*log(d*x -
sqrt(2)*sqrt(d*x)*(d^2/c)^(1/4) + sqrt(d^2/c))/(c^2*d^2) + sqrt(2)*(-c^3*
d^2)^(3/4)*b*log(d*x + sqrt(2)*sqrt(d*x)*(-d^2/c)^(1/4) + sqrt(-d^2/c))/(c
^2*d^2) - sqrt(2)*(-c^3*d^2)^(3/4)*b*log(d*x - sqrt(2)*sqrt(d*x)*(-d^2/c)^(
1/4) + sqrt(-d^2/c))/(c^2*d^2))/d
```

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{3/2}} dx = \int \frac{a + b \operatorname{atanh}(cx^2)}{(dx)^{3/2}} dx$$

input

```
int((a + b*atanh(c*x^2))/(d*x)^(3/2), x)
```

output

```
int((a + b*atanh(c*x^2))/(d*x)^(3/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.07

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{3/2}} dx = \frac{\sqrt{d} \left(-2\sqrt{x} c^{\frac{1}{4}} \sqrt{2} \operatorname{atan} \left(\frac{c^{\frac{1}{4}} \sqrt{2} - 2\sqrt{x} \sqrt{c}}{c^{\frac{1}{4}} \sqrt{2}} \right) b + 2\sqrt{x} c^{\frac{1}{4}} \sqrt{2} \operatorname{atan} \left(\frac{c^{\frac{1}{4}} \sqrt{2} + 2\sqrt{x} \sqrt{c}}{c^{\frac{1}{4}} \sqrt{2}} \right) b - 4 \right)}{d}$$

input

```
int((a+b*atanh(c*x^2))/(d*x)^(3/2), x)
```

output

```
(sqrt(d)*(- 2*sqrt(x)*c**(1/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) - 2*sqrt(x)
*sqrt(c))/(c**(1/4)*sqrt(2)))*b + 2*sqrt(x)*c**(1/4)*sqrt(2)*atan((c**(1/4)
)*sqrt(2) + 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*b - 4*sqrt(x)*c**(1/4)*
atan((sqrt(x)*sqrt(c))/c**(1/4))*b - 2*sqrt(x)*c**(1/4)*sqrt(2)*atanh(c*x*
*2)*b - 4*atanh(c*x**2)*b - sqrt(x)*c**(1/4)*sqrt(2)*log(c**(1/4) + sqrt(x)
)*sqrt(c))*b - sqrt(x)*c**(1/4)*sqrt(2)*log(- c**(1/4) + sqrt(x)*sqrt(c))
*b + 2*sqrt(x)*c**(1/4)*sqrt(2)*log(- sqrt(x)*c**(1/4)*sqrt(2) + sqrt(c)*
x + 1)*b - sqrt(x)*c**(1/4)*sqrt(2)*log(sqrt(c)*x + 1)*b + 2*sqrt(x)*c**(1
/4)*log(c**(1/4) + sqrt(x)*sqrt(c))*b - 2*sqrt(x)*c**(1/4)*log(- c**(1/4)
+ sqrt(x)*sqrt(c))*b - 4*a))/(2*sqrt(x)*d**2)
```

3.87 $\int \frac{a+b\operatorname{arctanh}(cx^2)}{(dx)^{5/2}} dx$

Optimal result	758
Mathematica [A] (verified)	759
Rubi [A] (verified)	759
Maple [A] (verified)	765
Fricas [C] (verification not implemented)	767
Sympy [F]	768
Maxima [A] (verification not implemented)	768
Giac [B] (verification not implemented)	769
Mupad [F(-1)]	769
Reduce [B] (verification not implemented)	770

Optimal result

Integrand size = 18, antiderivative size = 241

$$\int \frac{a + b\operatorname{arctanh}(cx^2)}{(dx)^{5/2}} dx = \frac{2bc^{3/4} \arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{3d^{5/2}} - \frac{\sqrt{2}bc^{3/4} \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{3d^{5/2}} + \frac{\sqrt{2}bc^{3/4} \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{3d^{5/2}} - \frac{2(a + b\operatorname{arctanh}(cx^2))}{3d(dx)^{3/2}} + \frac{2bc^{3/4}\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{3d^{5/2}} + \frac{\sqrt{2}bc^{3/4}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}(1+\sqrt{cx})}\right)}{3d^{5/2}}$$

output

```
2/3*b*c^(3/4)*arctan(c^(1/4)*(d*x)^(1/2)/d^(1/2))/d^(5/2)+1/3*2^(1/2)*b*c^(3/4)*arctan(-1+2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/2))/d^(5/2)+1/3*2^(1/2)*b*c^(3/4)*arctan(1+2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/2))/d^(5/2)-2/3*(a+b*arctanh(c*x^2))/d/(d*x)^(3/2)+2/3*b*c^(3/4)*arctanh(c^(1/4)*(d*x)^(1/2)/d^(1/2))/d^(5/2)+1/3*2^(1/2)*b*c^(3/4)*arctanh(2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/2)/(1+c^(1/2)*x))/d^(5/2)
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.11

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{5/2}} dx =$$

$$x(4a + 2\sqrt{2}bc^{3/4}x^{3/2} \arctan(1 - \sqrt{2}\sqrt[4]{c}\sqrt{x}) - 2\sqrt{2}bc^{3/4}x^{3/2} \arctan(1 + \sqrt{2}\sqrt[4]{c}\sqrt{x}) - 4bc^{3/4}x^{3/2} \arctan$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])/(d*x)^(5/2), x]
```

output

```
-1/6*(x*(4*a + 2*Sqrt[2]*b*c^(3/4)*x^(3/2)*ArcTan[1 - Sqrt[2]*c^(1/4)*Sqrt[x]] - 2*Sqrt[2]*b*c^(3/4)*x^(3/2)*ArcTan[1 + Sqrt[2]*c^(1/4)*Sqrt[x]] - 4*b*c^(3/4)*x^(3/2)*ArcTan[c^(1/4)*Sqrt[x]] + 4*b*ArcTanh[c*x^2] + 2*b*c^(3/4)*x^(3/2)*Log[1 - c^(1/4)*Sqrt[x]] - 2*b*c^(3/4)*x^(3/2)*Log[1 + c^(1/4)*Sqrt[x]] + Sqrt[2]*b*c^(3/4)*x^(3/2)*Log[1 - Sqrt[2]*c^(1/4)*Sqrt[x]] + Sqrt[c]*x - Sqrt[2]*b*c^(3/4)*x^(3/2)*Log[1 + Sqrt[2]*c^(1/4)*Sqrt[x]] + Sqrt[c]*x)))/(d*x)^(5/2)
```

Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 326, normalized size of antiderivative = 1.35, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.778$, Rules used = {6464, 851, 758, 755, 756, 218, 221, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{5/2}} dx$$

$$\downarrow 6464$$

$$\frac{4bc \int \frac{1}{\sqrt{dx(1-c^2x^4)}} dx}{3d^2} - \frac{2(a + b \operatorname{arctanh}(cx^2))}{3d(dx)^{3/2}}$$

$$\downarrow 851$$

$$\frac{8bc \int \frac{1}{1-c^2x^4} d\sqrt{dx}}{3d^3} - \frac{2(a + \operatorname{barctanh}(cx^2))}{3d(dx)^{3/2}}$$

↓ 758

$$\frac{8bc \left(\frac{1}{2}d^2 \int \frac{1}{d^2 - cd^2x^2} d\sqrt{dx} + \frac{1}{2}d^2 \int \frac{1}{cx^2d^2 + d^2} d\sqrt{dx} \right)}{3d^3} - \frac{2(a + \operatorname{barctanh}(cx^2))}{3d(dx)^{3/2}}$$

↓ 755

$$\frac{8bc \left(\frac{1}{2}d^2 \int \frac{1}{d^2 - cd^2x^2} d\sqrt{dx} + \frac{1}{2}d^2 \left(\int \frac{\frac{d - \sqrt{cdx}}{cx^2d^2 + d^2} d\sqrt{dx}}{2d} + \int \frac{\frac{\sqrt{cdx} + d}{cx^2d^2 + d^2} d\sqrt{dx}}{2d} \right) \right)}{3d^3} - \frac{2(a + \operatorname{barctanh}(cx^2))}{3d(dx)^{3/2}}$$

↓ 756

$$\frac{8bc \left(\frac{1}{2}d^2 \left(\int \frac{\frac{d - \sqrt{cdx}}{cx^2d^2 + d^2} d\sqrt{dx}}{2d} + \int \frac{\frac{\sqrt{cdx} + d}{cx^2d^2 + d^2} d\sqrt{dx}}{2d} \right) + \frac{1}{2}d^2 \left(\int \frac{\frac{1}{d - \sqrt{cdx}} d\sqrt{dx}}{2d} + \int \frac{\frac{1}{\sqrt{cdx} + d} d\sqrt{dx}}{2d} \right) \right)}{3d^3} - \frac{2(a + \operatorname{barctanh}(cx^2))}{3d(dx)^{3/2}}$$

↓ 218

$$\frac{8bc \left(\frac{1}{2}d^2 \left(\int \frac{\frac{1}{d - \sqrt{cdx}} d\sqrt{dx}}{2d} + \frac{\arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{cd^3/2}} \right) + \frac{1}{2}d^2 \left(\int \frac{\frac{d - \sqrt{cdx}}{cx^2d^2 + d^2} d\sqrt{dx}}{2d} + \int \frac{\frac{\sqrt{cdx} + d}{cx^2d^2 + d^2} d\sqrt{dx}}{2d} \right) \right)}{3d^3} - \frac{2(a + \operatorname{barctanh}(cx^2))}{3d(dx)^{3/2}}$$

↓ 221

$$\frac{8bc \left(\frac{1}{2}d^2 \left(\int \frac{\frac{d - \sqrt{cdx}}{cx^2d^2 + d^2} d\sqrt{dx}}{2d} + \int \frac{\frac{\sqrt{cdx} + d}{cx^2d^2 + d^2} d\sqrt{dx}}{2d} \right) + \frac{1}{2}d^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{cd^3/2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{cd^3/2}} \right) \right)}{3d^3} - \frac{2(a + \operatorname{barctanh}(cx^2))}{3d(dx)^{3/2}}$$

↓ 1476

$$8bc \left(\frac{1}{2} d^2 \left(\frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2d} + \frac{\int \frac{1}{xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{4\sqrt{c}}} d\sqrt{dx}}{2\sqrt{c}} + \frac{\int \frac{1}{xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{4\sqrt{c}}} d\sqrt{dx}}{2\sqrt{c}} \right) + \frac{1}{2} d^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{cd^{3/2}}}\right) \right)$$

$$\frac{2(a + b\operatorname{arctanh}(cx^2))}{3d(dx)^{3/2}} \quad 3d^3$$

1082

$$8bc \left(\frac{1}{2} d^2 \left(\frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2d} + \frac{\int \frac{1}{-dx-1} d\left(1-\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{1}{-dx-1} d\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} \right) + \frac{1}{2} d^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{cd^{3/2}}}\right) \right)$$

$$\frac{2(a + b\operatorname{arctanh}(cx^2))}{3d(dx)^{3/2}} \quad 3d^3$$

217

$$8bc \left(\frac{1}{2} d^2 \left(\frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2d} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} \right) + \frac{1}{2} d^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{cd^{3/2}}}\right) \right)$$

$$\frac{2(a + b\operatorname{arctanh}(cx^2))}{3d(dx)^{3/2}} \quad 3d^3$$

1479

$$8bc \left(\frac{1}{2} d^2 \left(\frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{c}\sqrt{dx}}{\sqrt[4]{c}\left(xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{4\sqrt{c}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{\sqrt{2}\left(\sqrt{d}+\sqrt{2}\sqrt[4]{c}\sqrt{dx}\right)}{\sqrt[4]{c}\left(xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{4\sqrt{c}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} \right) \right)$$

$$\frac{2(a + b\operatorname{arctanh}(cx^2))}{3d(dx)^{3/2}} \quad 3d^3$$

$$\begin{aligned}
 & \downarrow 25 \\
 & 8bc \left(\frac{1}{2} d^2 \left(\frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{c}\sqrt{dx}}{\sqrt[4]{c}\left(xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} + \frac{\int \frac{\sqrt{2}(\sqrt{d}+\sqrt{2}\sqrt[4]{c}\sqrt{dx})}{\sqrt[4]{c}\left(xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} \right) + \right. \\
 & \left. \frac{2(a + b\operatorname{arctanh}(cx^2))}{3d(dx)^{3/2}} \right) \\
 & \downarrow 27
 \end{aligned}$$

$$\begin{aligned}
 & 8bc \left(\frac{1}{2} d^2 \left(\frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{c}\sqrt{dx}}{xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}} d\sqrt{dx}}{2\sqrt{2}\sqrt{c}\sqrt{d}} + \frac{\int \frac{\sqrt{d}+\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}} d\sqrt{dx}}{2\sqrt{c}\sqrt{d}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} \right) + \frac{1}{2} d^2 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} \right) \right) \\
 & \left. \frac{2(a + b\operatorname{arctanh}(cx^2))}{3d(dx)^{3/2}} \right) \\
 & \downarrow 1103
 \end{aligned}$$

$$\begin{aligned}
 & 8bc \left(\frac{1}{2} d^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{c}d^{3/2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{c}d^{3/2}} \right) + \frac{1}{2} d^2 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} + \frac{\log\left(\sqrt{cdx}+\sqrt{2}\sqrt[4]{c}\sqrt{dx}\right)}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} \right) \right) \\
 & \left. \frac{2(a + b\operatorname{arctanh}(cx^2))}{3d(dx)^{3/2}} \right)
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c*x^2])/(d*x)^(5/2), x]`

output

$$\begin{aligned} & (-2*(a + b*\text{ArcTanh}[c*x^2]))/(3*d*(d*x)^{(3/2)}) + (8*b*c*((d^2*(\text{ArcTan}[(c^{(1/4)}*\text{Sqrt}[d*x])/ \text{Sqrt}[d]])/(2*c^{(1/4)}*d^{(3/2)}) + \text{ArcTanh}[(c^{(1/4)}*\text{Sqrt}[d*x])/ \text{Sqrt}[d]])/(2*c^{(1/4)}*d^{(3/2)})))/2 + (d^2*((-\text{ArcTan}[1 - (\text{Sqrt}[2]*c^{(1/4)}*\text{Sqrt}[d*x])/ \text{Sqrt}[d]])/(\text{Sqrt}[2]*c^{(1/4)}*\text{Sqrt}[d])) + \text{ArcTan}[1 + (\text{Sqrt}[2]*c^{(1/4)}*\text{Sqrt}[d*x])/ \text{Sqrt}[d]])/(\text{Sqrt}[2]*c^{(1/4)}*\text{Sqrt}[d]))/(2*d) + (-1/2*\text{Log}[d + \text{Sqrt}[c]*d*x - \text{Sqrt}[2]*c^{(1/4)}*\text{Sqrt}[d]*\text{Sqrt}[d*x]]/(\text{Sqrt}[2]*c^{(1/4)}*\text{Sqrt}[d]) + \text{Log}[d + \text{Sqrt}[c]*d*x + \text{Sqrt}[2]*c^{(1/4)}*\text{Sqrt}[d]*\text{Sqrt}[d*x]]/(2*\text{Sqrt}[2]*c^{(1/4)}*\text{Sqrt}[d]))/(2*d)))/(3*d^3) \end{aligned}$$

Defintions of rubi rules used

rule 25

$$\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$$

rule 27

$$\text{Int}[(a_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] /; \text{FreeQ}[a, \text{x}] \&\& \text{!MatchQ}[\text{Fx}, (b_)*(\text{Gx}_) /; \text{FreeQ}[b, \text{x}]]$$

rule 217

$$\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1} * \text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& \text{PosQ}[a/b] \& \& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])$$

rule 218

$$\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& \text{PosQ}[a/b]$$

rule 221

$$\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& \text{NegQ}[a/b]$$

rule 755

$$\text{Int}[((a_) + (b_.)*(x_)^4)^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*r) \quad \text{Int}[(r - s*x^2)/(a + b*x^4), \text{x}], \text{x}] + \text{Simp}[1/(2*r) \quad \text{Int}[(r + s*x^2)/(a + b*x^4), \text{x}], \text{x}]] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& (\text{GtQ}[a/b, 0] \parallel (\text{PosQ}[a/b] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \& \& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$$

rule 756 $\text{Int}[(a_ + (b_ \cdot x_)^4)^{-1}, x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[r/(2*a) \text{ Int}[1/(r - s*x^2), x], x] + \text{Simp}[r/(2*a) \text{ Int}[1/(r + s*x^2), x], x]] /; \text{FreeQ}[\{a, b\}, x] \&\& !\text{GtQ}[a/b, 0]$

rule 758 $\text{Int}[(a_ + (b_ \cdot x_)^{n_})^{-1}, x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[r/(2*a) \text{ Int}[1/(r - s*x^{(n/2)}), x], x] + \text{Simp}[r/(2*a) \text{ Int}[1/(r + s*x^{(n/2)}), x], x]] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{IGtQ}[n/4, 1] \&\& !\text{GtQ}[a/b, 0]$

rule 851 $\text{Int}[(c_ \cdot x_)^m \cdot (a_ + (b_ \cdot x_)^{n_})^p, x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \text{ Subst}[\text{Int}[x^{(k*(m+1)-1)} \cdot (a + b \cdot x^{(k*n)/c})^p, x], x, (c*x)^{1/k}], x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{FractionQ}[m] \&\& \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 1082 $\text{Int}[(a_ + (b_ \cdot x_) + (c_ \cdot x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S \text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \|\ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}[(d_ + (e_ \cdot x_))/((a_ + (b_ \cdot x_) + (c_ \cdot x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$

rule 1476 $\text{Int}[(d_ + (e_ \cdot x_)^2)/((a_ + (c_ \cdot x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2*(d/e), 2]\}, \text{Simp}[e/(2*c) \text{ Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Simp}[e/(2*c) \text{ Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[d*e]$

rule 1479 $\text{Int}[(d_ + (e_ \cdot x_)^2)/((a_ + (c_ \cdot x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2*(d/e), 2]\}, \text{Simp}[e/(2*c*q) \text{ Int}[(q - 2*x)/\text{Simp}[d/e + q*x - x^2, x], x], x] + \text{Simp}[e/(2*c*q) \text{ Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{NegQ}[d*e]$

rule 6464

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))*((d_)*(x_)^(m_)), x_Symbol] :  
> Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n  
/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a,  
b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 238, normalized size of antiderivative = 0.99

method	result
derivativedivides	$-\frac{2a}{3(dx)^{\frac{3}{2}}} + 2b - \frac{\operatorname{arctanh}(cx^2)}{3(dx)^{\frac{3}{2}}} + \frac{4cd^2}{8d^4} \left(\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \ln \left(\frac{\sqrt{dx} + \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}{\sqrt{dx} - \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) \right) + \frac{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{2} \ln \left(\frac{dx + \left(\frac{d^2}{c}\right)}{dx - \left(\frac{d^2}{c}\right)} \right)}{3}$
default	$-\frac{2a}{3(dx)^{\frac{3}{2}}} + 2b - \frac{\operatorname{arctanh}(cx^2)}{3(dx)^{\frac{3}{2}}} + \frac{4cd^2}{8d^4} \left(\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \ln \left(\frac{\sqrt{dx} + \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}{\sqrt{dx} - \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) \right) + \frac{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{2} \ln \left(\frac{dx + \left(\frac{d^2}{c}\right)}{dx - \left(\frac{d^2}{c}\right)} \right)}{3}$
parts	$-\frac{2a}{3(dx)^{\frac{3}{2}}d} + 2b - \frac{\operatorname{arctanh}(cx^2)}{3(dx)^{\frac{3}{2}}} + \frac{4cd^2}{8d^4} \left(\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \ln \left(\frac{\sqrt{dx} + \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}{\sqrt{dx} - \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) \right) + \frac{\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{2} \ln \left(\frac{dx + \left(\frac{d^2}{c}\right)}{dx - \left(\frac{d^2}{c}\right)} \right)}{3}$

input

```
int((a+b*arctanh(c*x^2))/(d*x)^(5/2),x,method=_RETURNVERBOSE)
```

output

```
2/d*(-1/3*a/(d*x)^(3/2)+b*(-1/3/(d*x)^(3/2)*arctanh(c*x^2)+4/3*c*d^2*(1/8/
d^4*(d^2/c)^(1/4)*(ln(((d*x)^(1/2)+(d^2/c)^(1/4))/((d*x)^(1/2)-(d^2/c)^(1/
4))))+2*arctan((d*x)^(1/2)/(d^2/c)^(1/4)))+1/16/d^4*(d^2/c)^(1/4)*2^(1/2)*(
ln((d*x+(d^2/c)^(1/4)*(d*x)^(1/2)*2^(1/2)+(d^2/c)^(1/2)))/(d*x-(d^2/c)^(1/4
))*(d*x)^(1/2)*2^(1/2)+(d^2/c)^(1/2)))+2*arctan(2^(1/2)/(d^2/c)^(1/4)*(d*x)
^(1/2)+1)+2*arctan(2^(1/2)/(d^2/c)^(1/4)*(d*x)^(1/2)-1))))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 421, normalized size of antiderivative = 1.75

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{5/2}} dx = \frac{d^3 x^2 \left(\frac{b^4 c^3}{d^{10}}\right)^{\frac{1}{4}} \log\left(d^3 \left(\frac{b^4 c^3}{d^{10}}\right)^{\frac{1}{4}} + \sqrt{dxbc}\right) + i d^3 x^2 \left(\frac{b^4 c^3}{d^{10}}\right)^{\frac{1}{4}} \log\left(i d^3 \left(\frac{b^4 c^3}{d^{10}}\right)^{\frac{1}{4}} + \sqrt{dxbc}\right)}{d^3 x^2 \left(\frac{b^4 c^3}{d^{10}}\right)^{\frac{1}{4}} \log\left(d^3 \left(\frac{b^4 c^3}{d^{10}}\right)^{\frac{1}{4}} + \sqrt{dxbc}\right) + i d^3 x^2 \left(\frac{b^4 c^3}{d^{10}}\right)^{\frac{1}{4}} \log\left(i d^3 \left(\frac{b^4 c^3}{d^{10}}\right)^{\frac{1}{4}} + \sqrt{dxbc}\right)}$$

input

```
integrate((a+b*arctanh(c*x^2))/(d*x)^(5/2),x, algorithm="fricas")
```

output

```
1/3*(d^3*x^2*(b^4*c^3/d^10)^(1/4)*log(d^3*(b^4*c^3/d^10)^(1/4) + sqrt(d*x)
*b*c) + I*d^3*x^2*(b^4*c^3/d^10)^(1/4)*log(I*d^3*(b^4*c^3/d^10)^(1/4) + sq
rt(d*x)*b*c) - I*d^3*x^2*(b^4*c^3/d^10)^(1/4)*log(-I*d^3*(b^4*c^3/d^10)^(1
/4) + sqrt(d*x)*b*c) - d^3*x^2*(b^4*c^3/d^10)^(1/4)*log(-d^3*(b^4*c^3/d^10
)^(1/4) + sqrt(d*x)*b*c) + d^3*x^2*(-b^4*c^3/d^10)^(1/4)*log(d^3*(-b^4*c^3
/d^10)^(1/4) + sqrt(d*x)*b*c) + I*d^3*x^2*(-b^4*c^3/d^10)^(1/4)*log(I*d^3*
(-b^4*c^3/d^10)^(1/4) + sqrt(d*x)*b*c) - I*d^3*x^2*(-b^4*c^3/d^10)^(1/4)*l
og(-I*d^3*(-b^4*c^3/d^10)^(1/4) + sqrt(d*x)*b*c) - d^3*x^2*(-b^4*c^3/d^10)
^(1/4)*log(-d^3*(-b^4*c^3/d^10)^(1/4) + sqrt(d*x)*b*c) - sqrt(d*x)*(b*log(
-(c*x^2 + 1)/(c*x^2 - 1)) + 2*a))/(d^3*x^2)
```

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{5/2}} dx = \int \frac{a + b \operatorname{atanh}(cx^2)}{(dx)^{5/2}} dx$$

input `integrate((a+b*atanh(c*x**2))/(d*x)**(5/2), x)`

output `Integral((a + b*atanh(c*x**2))/(d*x)**(5/2), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.15

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{5/2}} dx = \frac{b}{d^2} \left(\frac{2\sqrt{2}d \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}c^{1/4}\sqrt{d} + 2\sqrt{dx}\sqrt{c}\right)}{2\sqrt{\sqrt{cd}}}\right)}{\sqrt{\sqrt{cd}}} + \frac{2\sqrt{2}d \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}c^{1/4}\sqrt{d} - 2\sqrt{dx}\sqrt{c}\right)}{2\sqrt{\sqrt{cd}}}\right)}{\sqrt{\sqrt{cd}}} + \frac{\sqrt{2}\sqrt{d} \log\left(\frac{\sqrt{cd}x + \sqrt{c}}{c^{1/4}}\right)}{d^2} \right)$$

input `integrate((a+b*arctanh(c*x^2))/(d*x)^(5/2), x, algorithm="maxima")`

output `1/6*(b*((2*sqrt(2)*d*arctan(1/2*sqrt(2)*(sqrt(2)*c^(1/4)*sqrt(d) + 2*sqrt(d*x)*sqrt(c))/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d) + 2*sqrt(2)*d*arctan(-1/2*sqrt(2)*(sqrt(2)*c^(1/4)*sqrt(d) - 2*sqrt(d*x)*sqrt(c))/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d) + sqrt(2)*sqrt(d)*log(sqrt(c)*d*x + sqrt(2)*sqrt(d*x)*c^(1/4)*sqrt(d) + d)/c^(1/4) - sqrt(2)*sqrt(d)*log(sqrt(c)*d*x - sqrt(2)*sqrt(d*x)*c^(1/4)*sqrt(d) + d)/c^(1/4) + 4*d*arctan(sqrt(d*x)*sqrt(c)/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d) - 2*d*log((sqrt(d*x)*sqrt(c) - sqrt(sqrt(c)*d))/(sqrt(d*x)*sqrt(c) + sqrt(sqrt(c)*d)))/sqrt(sqrt(c)*d)*c/d^2 - 4*arctanh(c*x^2)/(d*x)^(3/2)) - 4*a/(d*x)^(3/2))/d`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 516 vs. $2(162) = 324$.

Time = 0.47 (sec) , antiderivative size = 516, normalized size of antiderivative = 2.14

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{5/2}} dx = \frac{bcd^2 \left(\frac{2\sqrt{2}(c^3d^2)^{1/4} \operatorname{arctan}\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{d^2}{c}\right)^{1/4} + 2\sqrt{dx}\right)}{2\left(\frac{d^2}{c}\right)^{1/4}}\right)}{cd^4} \right) + \frac{2\sqrt{2}(c^3d^2)^{1/4} \operatorname{arctan}\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{d^2}{c}\right)^{1/4} - 2\sqrt{dx}\right)}{2\left(\frac{d^2}{c}\right)^{1/4}}\right)}{cd^4}}{cd^4}$$

input `integrate((a+b*arctanh(c*x^2))/(d*x)^(5/2),x, algorithm="giac")`

output `1/6*(b*c*d^2*(2*sqrt(2)*(c^3*d^2)^(1/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(d^2/c)^(1/4) + 2*sqrt(d*x))/(d^2/c)^(1/4)))/(c*d^4) + 2*sqrt(2)*(c^3*d^2)^(1/4)*arctan(-1/2*sqrt(2)*(sqrt(2)*(d^2/c)^(1/4) - 2*sqrt(d*x))/(d^2/c)^(1/4)))/(c*d^4) + 2*sqrt(2)*(-c^3*d^2)^(1/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(-d^2/c)^(1/4) + 2*sqrt(d*x))/(-d^2/c)^(1/4)))/(c*d^4) + 2*sqrt(2)*(-c^3*d^2)^(1/4)*arctan(-1/2*sqrt(2)*(sqrt(2)*(-d^2/c)^(1/4) - 2*sqrt(d*x))/(-d^2/c)^(1/4)))/(c*d^4) + sqrt(2)*(c^3*d^2)^(1/4)*log(d*x + sqrt(2)*sqrt(d*x)*(d^2/c)^(1/4) + sqrt(d^2/c)))/(c*d^4) - sqrt(2)*(c^3*d^2)^(1/4)*log(d*x - sqrt(2)*sqrt(d*x)*(d^2/c)^(1/4) + sqrt(d^2/c)))/(c*d^4) + sqrt(2)*(-c^3*d^2)^(1/4)*log(d*x + sqrt(2)*sqrt(d*x)*(-d^2/c)^(1/4) + sqrt(-d^2/c)))/(c*d^4) - sqrt(2)*(-c^3*d^2)^(1/4)*log(d*x - sqrt(2)*sqrt(d*x)*(-d^2/c)^(1/4) + sqrt(-d^2/c)))/(c*d^4) - 2*b*log(-(c*d^2*x^2 + d^2)/(c*d^2*x^2 - d^2))/(sqrt(d*x)*d*x) - 4*a/(sqrt(d*x)*d*x))/d`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{5/2}} dx = \int \frac{a + b \operatorname{atanh}(cx^2)}{(dx)^{5/2}} dx$$

input `int((a + b*atanh(c*x^2))/(d*x)^(5/2),x)`

output `int((a + b*atanh(c*x^2))/(d*x)^(5/2), x)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 254, normalized size of antiderivative = 1.05

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{5/2}} dx = \frac{\sqrt{d} \left(-2\sqrt{x} c^{\frac{3}{4}} \sqrt{2} \operatorname{atan} \left(\frac{c^{\frac{1}{4}} \sqrt{2} - 2\sqrt{x} \sqrt{c}}{c^{\frac{1}{4}} \sqrt{2}} \right) bx + 2\sqrt{x} c^{\frac{3}{4}} \sqrt{2} \operatorname{atan} \left(\frac{c^{\frac{1}{4}} \sqrt{2} + 2\sqrt{x} \sqrt{c}}{c^{\frac{1}{4}} \sqrt{2}} \right) bx - \right.}{}$$

input `int((a+b*atanh(c*x^2))/(d*x)^(5/2), x)`

output `(sqrt(d)*(- 2*sqrt(x)*c**(3/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*b*x + 2*sqrt(x)*c**(3/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*b*x + 4*sqrt(x)*c**(3/4)*atan((sqrt(x)*sqrt(c))/c**(1/4))*b*x + 2*sqrt(x)*c**(3/4)*sqrt(2)*atanh(c*x**2)*b*x - 4*atanh(c*x**2)*b + sqrt(x)*c**(3/4)*sqrt(2)*log(c**(1/4) + sqrt(x)*sqrt(c))*b*x + sqrt(x)*c**(3/4)*sqrt(2)*log(- c**(1/4) + sqrt(x)*sqrt(c))*b*x - 2*sqrt(x)*c**(3/4)*sqrt(2)*log(- sqrt(x)*c**(1/4)*sqrt(2) + sqrt(c)*x + 1)*b*x + sqrt(x)*c**(3/4)*sqrt(2)*log(sqrt(c)*x + 1)*b*x + 2*sqrt(x)*c**(3/4)*log(c**(1/4) + sqrt(x)*sqrt(c))*b*x - 2*sqrt(x)*c**(3/4)*log(- c**(1/4) + sqrt(x)*sqrt(c))*b*x - 4*a))/(6*sqrt(x)*d**3*x)`

3.88 $\int \frac{a+b\operatorname{arctanh}(cx^2)}{(dx)^{7/2}} dx$

Optimal result	771
Mathematica [A] (verified)	772
Rubi [A] (verified)	772
Maple [A] (verified)	780
Fricas [C] (verification not implemented)	782
Sympy [F(-1)]	783
Maxima [A] (verification not implemented)	783
Giac [B] (verification not implemented)	784
Mupad [F(-1)]	785
Reduce [B] (verification not implemented)	785

Optimal result

Integrand size = 18, antiderivative size = 257

$$\int \frac{a + b\operatorname{arctanh}(cx^2)}{(dx)^{7/2}} dx = -\frac{8bc}{5d^3\sqrt{dx}} - \frac{2bc^{5/4} \arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{5d^{7/2}}$$

$$+ \frac{\sqrt{2}bc^{5/4} \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{5d^{7/2}} - \frac{\sqrt{2}bc^{5/4} \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{5d^{7/2}}$$

$$- \frac{2(a + b\operatorname{arctanh}(cx^2))}{5d(dx)^{5/2}} + \frac{2bc^{5/4}\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{5d^{7/2}} + \frac{\sqrt{2}bc^{5/4}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}(1+\sqrt{cx})}\right)}{5d^{7/2}}$$

output

```
-8/5*b*c/d^3/(d*x)^(1/2)-2/5*b*c^(5/4)*arctan(c^(1/4)*(d*x)^(1/2)/d^(1/2))
/d^(7/2)-1/5*2^(1/2)*b*c^(5/4)*arctan(-1+2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/2))
/d^(7/2)-1/5*2^(1/2)*b*c^(5/4)*arctan(1+2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1/2))
/d^(7/2)-2/5*(a+b*arctanh(c*x^2))/d/(d*x)^(5/2)+2/5*b*c^(5/4)*arctanh
(c^(1/4)*(d*x)^(1/2)/d^(1/2))/d^(7/2)+1/5*2^(1/2)*b*c^(5/4)*arctanh(2^(1/2)
)*c^(1/4)*(d*x)^(1/2)/d^(1/2)/(1+c^(1/2)*x))/d^(7/2)
```


Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.07

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{7/2}} dx = \frac{x(-4a - 16bcx^2 + 2\sqrt{2}bc^{5/4}x^{5/2} \arctan(1 - \sqrt{2}\sqrt{c}\sqrt{x}) - 2\sqrt{2}bc^{5/4}x^{5/2} \arctan$$

input `Integrate[(a + b*ArcTanh[c*x^2])/(d*x)^(7/2), x]`

output $(x*(-4*a - 16*b*c*x^2 + 2*\sqrt{2}*b*c^{5/4}*x^{5/2}*ArcTan[1 - \sqrt{2}*c^{1/4}*\sqrt{x}] - 2*\sqrt{2}*b*c^{5/4}*x^{5/2}*ArcTan[1 + \sqrt{2}*c^{1/4}*\sqrt{x}] - 4*b*c^{5/4}*x^{5/2}*ArcTan[c^{1/4}*\sqrt{x}] - 4*b*ArcTanh[c*x^2] - 2*b*c^{5/4}*x^{5/2}*Log[1 - c^{1/4}*\sqrt{x}] + 2*b*c^{5/4}*x^{5/2}*Log[1 + c^{1/4}*\sqrt{x}] - \sqrt{2}*b*c^{5/4}*x^{5/2}*Log[1 - \sqrt{2}*c^{1/4}*\sqrt{x}] + \sqrt{2}*b*c^{5/4}*x^{5/2}*Log[1 + \sqrt{2}*c^{1/4}*\sqrt{x}] + \sqrt{c}*x) / (10*(d*x)^{7/2})$

Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 351, normalized size of antiderivative = 1.37, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.889$, Rules used = {6464, 847, 851, 27, 830, 826, 827, 218, 221, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{7/2}} dx \\ & \quad \downarrow \text{6464} \\ & \frac{4bc \int \frac{1}{(dx)^{3/2}(1-c^2x^4)} dx}{5d^2} - \frac{2(a + b \operatorname{arctanh}(cx^2))}{5d(dx)^{5/2}} \\ & \quad \downarrow \text{847} \\ & \frac{4bc \left(\frac{c^2 \int \frac{(dx)^{5/2}}{1-c^2x^4} dx}{d^4} - \frac{2}{d\sqrt{dx}} \right)}{5d^2} - \frac{2(a + b \operatorname{arctanh}(cx^2))}{5d(dx)^{5/2}} \end{aligned}$$

$$\begin{array}{c} \downarrow 851 \\ 4bc \left(\frac{2c^2 \int \frac{d^7 x^3}{d^4 - c^2 d^4 x^4} d\sqrt{dx} - \frac{2}{d\sqrt{dx}}}{5d^2} \right) - \frac{2(a + \operatorname{barctanh}(cx^2))}{5d(dx)^{5/2}} \end{array}$$

$$\begin{array}{c} \downarrow 27 \\ 4bc \left(\frac{2c^2 \int \frac{d^3 x^3}{d^4 - c^2 d^4 x^4} d\sqrt{dx} - \frac{2}{d\sqrt{dx}}}{5d^2} \right) - \frac{2(a + \operatorname{barctanh}(cx^2))}{5d(dx)^{5/2}} \end{array}$$

$$\begin{array}{c} \downarrow 830 \\ 4bc \left(\frac{2c^2 \left(\frac{\int \frac{dx}{d^2 - cd^2 x^2} d\sqrt{dx}}{2c} - \frac{\int \frac{dx}{cx^2 d^2 + d^2} d\sqrt{dx}}{2c} \right) - \frac{2}{d\sqrt{dx}}}{5d^2} \right) - \frac{2(a + \operatorname{barctanh}(cx^2))}{5d(dx)^{5/2}} \end{array}$$

$$\begin{array}{c} \downarrow 826 \\ 4bc \left(\frac{2c^2 \left(\frac{\int \frac{dx}{d^2 - cd^2 x^2} d\sqrt{dx}}{2c} - \frac{\int \frac{\sqrt{cx}d+d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{c}dx}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} \right) - \frac{2}{d\sqrt{dx}}}{5d^2} \right) - \frac{2(a + \operatorname{barctanh}(cx^2))}{5d(dx)^{5/2}} \end{array}$$

$$\begin{array}{c} \downarrow 827 \\ 4bc \left(\frac{2c^2 \left(\frac{\int \frac{1}{d-\sqrt{c}dx} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{1}{\sqrt{cx}d+d} d\sqrt{dx}}{2c} - \frac{\int \frac{\sqrt{cx}d+d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} - \frac{\int \frac{d-\sqrt{c}dx}{cx^2 d^2 + d^2} d\sqrt{dx}}{2\sqrt{c}} \right) - \frac{2}{d\sqrt{dx}}}{5d^2} \right) - \frac{2(a + \operatorname{barctanh}(cx^2))}{5d(dx)^{5/2}} \end{array}$$

$$\downarrow 218$$

$$\begin{aligned}
 & \left(\frac{4bc}{2c^2} \left(\frac{\int \frac{1}{d-\sqrt{cdx}} d\sqrt{dx} - \frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}}}{2c} - \frac{\int \frac{\sqrt{cdx}d+d}{cx^2d^2+d^2} d\sqrt{dx} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2c}}{d} \right) - \frac{2}{d\sqrt{dx}} \right) \\
 & \frac{5d^2}{2(a + b\operatorname{arctanh}(cx^2))} \\
 & \frac{5d(dx)^{5/2}}{5d(dx)^{5/2}} \\
 & \downarrow \text{221} \\
 & \left(\frac{4bc}{2c^2} \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}}}{2c} - \frac{\int \frac{\sqrt{cdx}d+d}{cx^2d^2+d^2} d\sqrt{dx} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2c}}{d} \right) - \frac{2}{d\sqrt{dx}} \right) \\
 & \frac{5d^2}{2(a + b\operatorname{arctanh}(cx^2))} \\
 & \frac{5d(dx)^{5/2}}{5d(dx)^{5/2}} \\
 & \downarrow \text{1476} \\
 & \left(\frac{4bc}{2c^2} \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}}}{2c} - \frac{\int \frac{1}{xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{4\sqrt{c}}} d\sqrt{dx} - \int \frac{1}{xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{4\sqrt{c}}} d\sqrt{dx}}{2\sqrt{c}} + \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2c}}{d} \right) - \frac{2}{d\sqrt{dx}} \right) \\
 & \frac{5d^2}{2(a + b\operatorname{arctanh}(cx^2))} \\
 & \frac{5d(dx)^{5/2}}{5d(dx)^{5/2}}
 \end{aligned}$$

↓ 1082

$$4bc \left(\frac{2c^2 \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c} - \frac{\int \frac{-\frac{1}{dx-1}d\left(1-\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{-\frac{1}{dx-1}d\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{d-\sqrt{c}dx}{cx^2d^2+d^2}d\sqrt{dx}}{2\sqrt{c}}}{d} \right)}{d} \right) - \frac{d\sqrt{dx}}{d\sqrt{dx}}$$

$$\frac{2(a + b\operatorname{arctanh}(cx^2))}{5d(dx)^{5/2}} \frac{5d^2}{5d(dx)^{5/2}}$$

↓ 217

$$4bc \left(\frac{2c^2 \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c} - \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\operatorname{arctan}\left(1-\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{d-\sqrt{c}dx}{cx^2d^2+d^2}d\sqrt{dx}}{2\sqrt{c}}}{d} \right) - \frac{2}{d\sqrt{dx}}$$

$$\frac{2(a + b\operatorname{arctanh}(cx^2))}{5d(dx)^{5/2}} \frac{5d^2}{5d(dx)^{5/2}}$$

↓ 1479

$$\left. \begin{array}{l} 2c^2 \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c} - \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\operatorname{arctan}\left(1-\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{c}\sqrt{dx}}{\sqrt[4]{c}\left(xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}\right)} dx}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} \right) \\ 4bc \end{array} \right\} d$$

$$\frac{2(a + b\operatorname{arctanh}(cx^2))}{5d(dx)^{5/2}} \quad 5d^2$$

↓ 25

$$\left. \begin{array}{l} 2c^2 \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c} - \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\operatorname{arctan}\left(1-\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{c}\sqrt{dx}}{\sqrt[4]{c}\left(xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{c}}\right)} dx}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} \right) \\ 4bc \end{array} \right\} d$$

$$\frac{2(a + b\operatorname{arctanh}(cx^2))}{5d(dx)^{5/2}} \quad 5d^2$$

↓ 27

$$\left(\begin{array}{l} 2c^2 \\ 4bc \end{array} \right) \left(\begin{array}{l} \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c} - \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\operatorname{arctan}\left(1-\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{c}\sqrt{dx}}{xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt{c}}}\sqrt{dx}}{2\sqrt{2}\sqrt{c}\sqrt{d}} - \frac{\int \frac{\sqrt{dx}}{xd+\frac{d}{\sqrt{c}}}}{2\sqrt{c}} \end{array} \right)$$

$$\frac{2(a + b\operatorname{arctanh}(cx^2))}{5d(dx)^{5/2}} \quad 5d^2$$

↓ 1103

$$\left(\begin{array}{l} 2c^2 \\ 4bc \end{array} \right) \left(\begin{array}{l} \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c^{3/4}\sqrt{d}} - \frac{\operatorname{arctan}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{2c} - \frac{\operatorname{arctan}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}+1\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\operatorname{arctan}\left(1-\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\log\left(\sqrt{cdx}+\sqrt{2}\sqrt[4]{c}\sqrt{d}\sqrt{dx}+d\right)}{2\sqrt{2}\sqrt[4]{c}\sqrt{d}} - \frac{\log\left(\sqrt{cdx}+\frac{d}{\sqrt{c}}\right)}{2\sqrt{c}} \end{array} \right)$$

$$\frac{2(a + b\operatorname{arctanh}(cx^2))}{5d(dx)^{5/2}} \quad 5d^2$$

input `Int[(a + b*ArcTanh[c*x^2])/(d*x)^(7/2), x]`

output

$$\begin{aligned} & (-2*(a + b*\text{ArcTanh}[c*x^2]))/(5*d*(d*x)^{(5/2)}) + (4*b*c*(-2/(d*\text{Sqrt}[d*x]) + \\ & (2*c^2*(-1/2*\text{ArcTan}[(c^{1/4}*\text{Sqrt}[d*x])/ \text{Sqrt}[d]]/(c^{3/4}*\text{Sqrt}[d]) + \text{Arc} \\ & \text{Tanh}[(c^{1/4}*\text{Sqrt}[d*x])/ \text{Sqrt}[d]]/(2*c^{3/4}*\text{Sqrt}[d]))/(2*c) - ((-\text{ArcTan}[\\ & 1 - (\text{Sqrt}[2]*c^{1/4}*\text{Sqrt}[d*x])/ \text{Sqrt}[d])/(\text{Sqrt}[2]*c^{1/4}*\text{Sqrt}[d])) + \text{ArcT} \\ & \text{an}[1 + (\text{Sqrt}[2]*c^{1/4}*\text{Sqrt}[d*x])/ \text{Sqrt}[d])/(\text{Sqrt}[2]*c^{1/4}*\text{Sqrt}[d]))/(2* \\ & \text{Sqrt}[c]) - (-1/2*\text{Log}[d + \text{Sqrt}[c]*d*x - \text{Sqrt}[2]*c^{1/4}*\text{Sqrt}[d]*\text{Sqrt}[d*x]]/ \\ & (\text{Sqrt}[2]*c^{1/4}*\text{Sqrt}[d]) + \text{Log}[d + \text{Sqrt}[c]*d*x + \text{Sqrt}[2]*c^{1/4}*\text{Sqrt}[d]* \\ & \text{Sqrt}[d*x]]/(2*\text{Sqrt}[2]*c^{1/4}*\text{Sqrt}[d]))/(2*\text{Sqrt}[c]))/(2*c))/d)/(5*d^2) \end{aligned}$$

Defintions of rubi rules used

rule 25

$$\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$$

rule 27

$$\text{Int}[(a_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] /; \text{FreeQ}[a, \text{x}] \&\& \text{!Ma} \\ \text{tchQ}[\text{Fx}, (b_)*(\text{Gx}_) /; \text{FreeQ}[b, \text{x}]$$

rule 217

$$\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)} \\ * \text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& \text{PosQ}[a/b] \& \\ \& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])$$

rule 218

$$\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{R} \\ \text{t}[a/b, 2]], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& \text{PosQ}[a/b]$$

rule 221

$$\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x \\ / \text{Rt}[-a/b, 2]], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& \text{NegQ}[a/b]$$

rule 826

$$\text{Int}[(x_)^2/((a_) + (b_.)*(x_)^4), \text{x_Symbol}] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, \\ 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*s) \quad \text{Int}[(r + s*x^2)/(a + b*x^ \\ 4), \text{x}], \text{x}] - \text{Simp}[1/(2*s) \quad \text{Int}[(r - s*x^2)/(a + b*x^4), \text{x}], \text{x}]] /; \text{FreeQ}[\{ \\ a, b\}, \text{x}] \&\& (\text{GtQ}[a/b, 0] \parallel (\text{PosQ}[a/b] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \\ \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$$

rule 827 $\text{Int}[(x_)^2/((a_)+(b_)*(x_)^4), x_Symbol] := \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[s/(2*b) \text{Int}[1/(r + s*x^2), x], x] - \text{Simp}[s/(2*b) \text{Int}[1/(r - s*x^2), x], x]] /; \text{FreeQ}[\{a, b\}, x] \&\& !\text{GtQ}[a/b, 0]$

rule 830 $\text{Int}[(x_)^{(m_)} / ((a_)+(b_)*(x_)^{(n_)}), x_Symbol] := \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[s/(2*b) \text{Int}[x^{(m-n/2)} / (r + s*x^{(n/2)}), x], x] - \text{Simp}[s/(2*b) \text{Int}[x^{(m-n/2)} / (r - s*x^{(n/2)}), x], x]] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{IGtQ}[n/4, 0] \&\& \text{IGtQ}[m, 0] \&\& \text{LeQ}[n/2, m] \&\& \text{LtQ}[m, n] \&\& !\text{GtQ}[a/b, 0]$

rule 847 $\text{Int}[(c_)*(x_)^{(m_)} * ((a_)+(b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] := \text{Simp}[(c*x)^{(m+1)} * ((a + b*x^n)^{(p+1)} / (a*c*(m+1))), x] - \text{Simp}[b*(m+n*(p+1) + 1) / (a*c^n*(m+1)) \text{Int}[(c*x)^{(m+n)} * (a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, p\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{LtQ}[m, -1] \&\& \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 851 $\text{Int}[(c_)*(x_)^{(m_)} * ((a_)+(b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] := \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \text{Subst}[\text{Int}[x^{(k*(m+1)-1)} * (a + b*(x^{(k*n)})/c^n)^p, x], x, (c*x)^{(1/k)}], x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{FractionQ}[m] \&\& \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 1082 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] := \text{With}[\{q = 1 - 4*S\text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] || !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] := \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$

rule 1476

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

rule 1479

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

rule 6464

```
Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)*((d_)*(x_)^(m_)), x_Symbol] :
> Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n
/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a,
b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 250, normalized size of antiderivative = 0.97

method	result
derivativedivides	$-\frac{2a}{5(dx)^{\frac{5}{2}}} + 2b - \frac{\operatorname{arctanh}(cx^2)}{5(dx)^{\frac{5}{2}}} + \frac{4cd^2}{d^4\sqrt{dx}} - \frac{\sqrt{2} \left(\ln \left(\frac{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}} {dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx} + 1}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx} + 1}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)}{16d^4 \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}$
default	$-\frac{2a}{5(dx)^{\frac{5}{2}}} + 2b - \frac{\operatorname{arctanh}(cx^2)}{5(dx)^{\frac{5}{2}}} + \frac{4cd^2}{d^4\sqrt{dx}} - \frac{\sqrt{2} \left(\ln \left(\frac{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}} {dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx} + 1}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx} + 1}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)}{16d^4 \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}$
parts	$-\frac{2a}{5(dx)^{\frac{5}{2}}} + 2b - \frac{\operatorname{arctanh}(cx^2)}{5(dx)^{\frac{5}{2}}} + \frac{4cd^2}{d^4\sqrt{dx}} - \frac{\sqrt{2} \left(\ln \left(\frac{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}} {dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{dx} \sqrt{2} + \sqrt{\frac{d^2}{c}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx} + 1}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{2} \sqrt{dx} + 1}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)}{16d^4 \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}$

input

```
int((a+b*arctanh(c*x^2))/(d*x)^(7/2),x,method=_RETURNVERBOSE)
```

output

```
2/d*(-1/5*a/(d*x)^(5/2)+b*(-1/5/(d*x)^(5/2)*arctanh(c*x^2)+4/5*c*d^2*(-1/d
^4/(d*x)^(1/2)-1/16/d^4/(d^2/c)^(1/4)*2^(1/2)*(ln((d*x-(d^2/c)^(1/4)*(d*x)
^(1/2)*2^(1/2)+(d^2/c)^(1/2)))/(d*x+(d^2/c)^(1/4)*(d*x)^(1/2)*2^(1/2)+(d^2/
c)^(1/2))))+2*arctan(2^(1/2)/(d^2/c)^(1/4)*(d*x)^(1/2)+1)+2*arctan(2^(1/2)/
(d^2/c)^(1/4)*(d*x)^(1/2)-1))-1/8/d^4/(d^2/c)^(1/4)*(2*arctan((d*x)^(1/2)/
(d^2/c)^(1/4))-ln(((d*x)^(1/2)+(d^2/c)^(1/4))/((d*x)^(1/2)-(d^2/c)^(1/4))
))))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 460, normalized size of antiderivative = 1.79

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{7/2}} dx = \frac{d^4 x^3 \left(\frac{b^4 c^5}{d^{14}}\right)^{\frac{1}{4}} \log\left(d^{11} \left(\frac{b^4 c^5}{d^{14}}\right)^{\frac{3}{4}} + \sqrt{d} x b^3 c^4\right) - i d^4 x^3 \left(\frac{b^4 c^5}{d^{14}}\right)^{\frac{1}{4}} \log\left(i d^{11} \left(\frac{b^4 c^5}{d^{14}}\right)^{\frac{3}{4}} + \sqrt{d} x b^3 c^4\right)}{d^4 x^3 \left(\frac{b^4 c^5}{d^{14}}\right)^{\frac{1}{4}} \log\left(d^{11} \left(\frac{b^4 c^5}{d^{14}}\right)^{\frac{3}{4}} + \sqrt{d} x b^3 c^4\right) - i d^4 x^3 \left(\frac{b^4 c^5}{d^{14}}\right)^{\frac{1}{4}} \log\left(i d^{11} \left(\frac{b^4 c^5}{d^{14}}\right)^{\frac{3}{4}} + \sqrt{d} x b^3 c^4\right)}$$

input

```
integrate((a+b*arctanh(c*x^2))/(d*x)^(7/2),x, algorithm="fricas")
```

output

```
1/5*(d^4*x^3*(b^4*c^5/d^14)^(1/4)*log(d^11*(b^4*c^5/d^14)^(3/4) + sqrt(d*x)
)*b^3*c^4) - I*d^4*x^3*(b^4*c^5/d^14)^(1/4)*log(I*d^11*(b^4*c^5/d^14)^(3/4)
) + sqrt(d*x)*b^3*c^4) + I*d^4*x^3*(b^4*c^5/d^14)^(1/4)*log(-I*d^11*(b^4*c
^5/d^14)^(3/4) + sqrt(d*x)*b^3*c^4) - d^4*x^3*(b^4*c^5/d^14)^(1/4)*log(-d
^11*(b^4*c^5/d^14)^(3/4) + sqrt(d*x)*b^3*c^4) - d^4*x^3*(-b^4*c^5/d^14)^(1/
4)*log(d^11*(-b^4*c^5/d^14)^(3/4) + sqrt(d*x)*b^3*c^4) + I*d^4*x^3*(-b^4*c
^5/d^14)^(1/4)*log(I*d^11*(-b^4*c^5/d^14)^(3/4) + sqrt(d*x)*b^3*c^4) - I*d
^4*x^3*(-b^4*c^5/d^14)^(1/4)*log(-I*d^11*(-b^4*c^5/d^14)^(3/4) + sqrt(d*x)
)*b^3*c^4) + d^4*x^3*(-b^4*c^5/d^14)^(1/4)*log(-d^11*(-b^4*c^5/d^14)^(3/4)
+ sqrt(d*x)*b^3*c^4) - (8*b*c*x^2 + b*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 2*a)
*sqrt(d*x))/(d^4*x^3)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{7/2}} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**2))/(d*x)**(7/2), x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 298, normalized size of antiderivative = 1.16

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{7/2}} dx =$$

$$\frac{c \left(\frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}c^{\frac{1}{4}}\sqrt{d}+2\sqrt{dx}\sqrt{c}\right)}{2\sqrt{cd}}\right)}{\sqrt{cd}\sqrt{c}} + \frac{2\sqrt{2} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}c^{\frac{1}{4}}\sqrt{d}-2\sqrt{dx}\sqrt{c}\right)}{2\sqrt{cd}}\right)}{\sqrt{cd}\sqrt{c}} \right) - \frac{\sqrt{2} \log\left(\sqrt{cdx}+\sqrt{2}\sqrt{dx}c^{\frac{1}{4}}\sqrt{d}+d\right)}{c^{\frac{3}{4}}\sqrt{d}} + \frac{\sqrt{2} \log\left(\sqrt{cdx}-\sqrt{2}\sqrt{dx}c^{\frac{1}{4}}\sqrt{d}+d\right)}{c^{\frac{3}{4}}\sqrt{d}}}{d^2}$$

$10d$

input `integrate((a+b*arctanh(c*x^2))/(d*x)^(7/2), x, algorithm="maxima")`

output

```

-1/10*(b*((c*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*c^(1/4)*sqrt(d) + 2*sqrt(d*x)*sqrt(c))/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d)*sqrt(c)) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*c^(1/4)*sqrt(d) - 2*sqrt(d*x)*sqrt(c))/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d)*sqrt(c) - sqrt(2)*log(sqrt(c)*d*x + sqrt(2)*sqrt(d*x)*c^(1/4)*sqrt(d) + d)/(c^(3/4)*sqrt(d)) + sqrt(2)*log(sqrt(c)*d*x - sqrt(2)*sqrt(d*x)*c^(1/4)*sqrt(d) + d)/(c^(3/4)*sqrt(d))) + 2*c*(2*arctan(sqrt(d*x)*sqrt(c)/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d)*sqrt(c) + log((sqrt(d*x)*sqrt(c) - sqrt(sqrt(c)*d))/sqrt(d*x)*sqrt(c) + sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d)*sqrt(c)) + 16/sqrt(d*x)*c/d^2 + 4*arctanh(c*x^2)/(d*x)^(5/2)) + 4*a/(d*x)^(5/2))/d

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 532 vs. $2(174) = 348$.

Time = 1.59 (sec) , antiderivative size = 532, normalized size of antiderivative = 2.07

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{7/2}} dx = \frac{2\sqrt{2}(c^3d^2)^{\frac{3}{4}} b \operatorname{arctan}\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 2\sqrt{dx}\right)}{2\left(\frac{d^2}{c}\right)^{\frac{1}{4}}}\right)}{cd^4} + \frac{2\sqrt{2}(c^3d^2)^{\frac{3}{4}} b \operatorname{arctan}\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{d^2}{c}\right)^{\frac{1}{4}} - 2\sqrt{dx}\right)}{2\left(\frac{d^2}{c}\right)^{\frac{1}{4}}}\right)}{cd^4} - \frac{2\sqrt{2}(-c^3d^2)^{\frac{3}{4}} b \operatorname{arctan}\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 2\sqrt{dx}\right)}{2\left(\frac{d^2}{c}\right)^{\frac{1}{4}}}\right)}{cd^4}$$

input

```
integrate((a+b*arctanh(c*x^2))/(d*x)^(7/2),x, algorithm="giac")
```

output

```
-1/10*(2*sqrt(2)*(c^3*d^2)^(3/4)*b*arctan(1/2*sqrt(2)*(sqrt(2)*(d^2/c)^(1/4) + 2*sqrt(d*x))/(d^2/c)^(1/4))/(c*d^4) + 2*sqrt(2)*(c^3*d^2)^(3/4)*b*arctan(-1/2*sqrt(2)*(sqrt(2)*(d^2/c)^(1/4) - 2*sqrt(d*x))/(d^2/c)^(1/4))/(c*d^4) - 2*sqrt(2)*(-c^3*d^2)^(3/4)*b*arctan(1/2*sqrt(2)*(sqrt(2)*(-d^2/c)^(1/4) + 2*sqrt(d*x))/(-d^2/c)^(1/4))/(c*d^4) - 2*sqrt(2)*(-c^3*d^2)^(3/4)*b*arctan(-1/2*sqrt(2)*(sqrt(2)*(-d^2/c)^(1/4) - 2*sqrt(d*x))/(-d^2/c)^(1/4))/(c*d^4) - sqrt(2)*(c^3*d^2)^(3/4)*b*log(d*x + sqrt(2)*sqrt(d*x)*(d^2/c)^(1/4) + sqrt(d^2/c))/(c*d^4) + sqrt(2)*(c^3*d^2)^(3/4)*b*log(d*x - sqrt(2)*sqrt(d*x)*(d^2/c)^(1/4) + sqrt(d^2/c))/(c*d^4) + sqrt(2)*(-c^3*d^2)^(3/4)*b*log(d*x + sqrt(2)*sqrt(d*x)*(-d^2/c)^(1/4) + sqrt(-d^2/c))/(c*d^4) - sqrt(2)*(-c^3*d^2)^(3/4)*b*log(d*x - sqrt(2)*sqrt(d*x)*(-d^2/c)^(1/4) + sqrt(-d^2/c))/(c*d^4) + 2*b*log(-(c*d^2*x^2 + d^2)/(c*d^2*x^2 - d^2))/(sqrt(d*x)*d^2*x^2) + 4*(4*b*c*d^2*x^2 + a*d^2)/(sqrt(d*x)*d^4*x^2))/d
```

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{7/2}} dx = \int \frac{a + b \operatorname{atanh}(cx^2)}{(dx)^{7/2}} dx$$

input

```
int((a + b*atanh(c*x^2))/(d*x)^(7/2), x)
```

output

```
int((a + b*atanh(c*x^2))/(d*x)^(7/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.09

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{7/2}} dx = \frac{\sqrt{d} \left(2\sqrt{x} c^{\frac{5}{4}} \sqrt{2} \operatorname{atan} \left(\frac{c^{\frac{1}{4}} \sqrt{2} - 2\sqrt{x} \sqrt{c}}{c^{\frac{1}{4}} \sqrt{2}} \right) b x^2 - 2\sqrt{x} c^{\frac{5}{4}} \sqrt{2} \operatorname{atan} \left(\frac{c^{\frac{1}{4}} \sqrt{2} + 2\sqrt{x} \sqrt{c}}{c^{\frac{1}{4}} \sqrt{2}} \right) b x^2 \right)}{(dx)^{7/2}}$$

input

```
int((a+b*atanh(c*x^2))/(d*x)^(7/2), x)
```

output

```
(sqrt(d)*(2*sqrt(x)*c**(1/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*b*c*x**2 - 2*sqrt(x)*c**(1/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*b*c*x**2 - 4*sqrt(x)*c**(1/4)*atan((sqrt(x)*sqrt(c))/c**(1/4))*b*c*x**2 + 2*sqrt(x)*c**(1/4)*sqrt(2)*atanh(c*x**2)*b*c*x**2 - 4*atanh(c*x**2)*b + sqrt(x)*c**(1/4)*sqrt(2)*log(c**(1/4) + sqrt(x)*sqrt(c))*b*c*x**2 + sqrt(x)*c**(1/4)*sqrt(2)*log(-c**(1/4) + sqrt(x)*sqrt(c))*b*c*x**2 - 2*sqrt(x)*c**(1/4)*sqrt(2)*log(-sqrt(x)*c**(1/4)*sqrt(2) + sqrt(c)*x + 1)*b*c*x**2 + sqrt(x)*c**(1/4)*sqrt(2)*log(sqrt(c)*x + 1)*b*c*x**2 + 2*sqrt(x)*c**(1/4)*log(c**(1/4) + sqrt(x)*sqrt(c))*b*c*x**2 - 2*sqrt(x)*c**(1/4)*log(-c**(1/4) + sqrt(x)*sqrt(c))*b*c*x**2 - 4*a - 16*b*c*x**2)/(10*sqrt(x)*d**4*x**2)
```

3.89 $\int \frac{a+b\operatorname{arctanh}(cx^2)}{(dx)^{9/2}} dx$

Optimal result	787
Mathematica [A] (verified)	788
Rubi [A] (verified)	788
Maple [A] (verified)	796
Fricas [C] (verification not implemented)	798
Sympy [F(-1)]	799
Maxima [A] (verification not implemented)	799
Giac [B] (verification not implemented)	800
Mupad [F(-1)]	801
Reduce [B] (verification not implemented)	801

Optimal result

Integrand size = 18, antiderivative size = 257

$$\int \frac{a + b\operatorname{arctanh}(cx^2)}{(dx)^{9/2}} dx = -\frac{8bc}{21d^3(dx)^{3/2}} + \frac{2bc^{7/4} \arctan\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{7d^{9/2}} + \frac{\sqrt{2}bc^{7/4} \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{7d^{9/2}} - \frac{\sqrt{2}bc^{7/4} \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{7d^{9/2}} - \frac{2(a + b\operatorname{arctanh}(cx^2))}{7d(dx)^{7/2}} + \frac{2bc^{7/4}\operatorname{arctanh}\left(\frac{\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}}\right)}{7d^{9/2}} - \frac{\sqrt{2}bc^{7/4}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{c}\sqrt{dx}}{\sqrt{d}(1+\sqrt{cx})}\right)}{7d^{9/2}}$$

output

```
-8/21*b*c/d^3/(d*x)^(3/2)+2/7*b*c^(7/4)*arctan(c^(1/4)*(d*x)^(1/2)/d^(1/2)
)/d^(9/2)-1/7*2^(1/2)*b*c^(7/4)*arctan(-1+2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^(1
/2))/d^(9/2)-1/7*2^(1/2)*b*c^(7/4)*arctan(1+2^(1/2)*c^(1/4)*(d*x)^(1/2)/d^
(1/2))/d^(9/2)-2/7*(a+b*arctanh(c*x^2))/d/(d*x)^(7/2)+2/7*b*c^(7/4)*arctan
h(c^(1/4)*(d*x)^(1/2)/d^(1/2))/d^(9/2)-1/7*2^(1/2)*b*c^(7/4)*arctanh(2^(1/
2)*c^(1/4)*(d*x)^(1/2)/d^(1/2)/(1+c^(1/2)*x))/d^(9/2)
```


Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.09

$$\int \frac{a + \operatorname{barctanh}(cx^2)}{(dx)^{9/2}} dx = \frac{\sqrt{dx}(-12a - 16bcx^2 + 6\sqrt{2}bc^{7/4}x^{7/2} \arctan(1 - \sqrt{2}\sqrt[4]{c}\sqrt{x}) - 6\sqrt{2}bc^{7/4}x^{7/2} \arctan(1 + \sqrt{2}\sqrt[4]{c}\sqrt{x}))}{(42d^5x^4)}$$

input `Integrate[(a + b*ArcTanh[c*x^2])/(d*x)^(9/2), x]`

output

```
(Sqrt[d*x]*(-12*a - 16*b*c*x^2 + 6*Sqrt[2]*b*c^(7/4)*x^(7/2)*ArcTan[1 - Sqrt[2]*c^(1/4)*Sqrt[x]] - 6*Sqrt[2]*b*c^(7/4)*x^(7/2)*ArcTan[1 + Sqrt[2]*c^(1/4)*Sqrt[x]] + 12*b*c^(7/4)*x^(7/2)*ArcTan[c^(1/4)*Sqrt[x]] - 12*b*ArcTanh[c*x^2] - 6*b*c^(7/4)*x^(7/2)*Log[1 - c^(1/4)*Sqrt[x]] + 6*b*c^(7/4)*x^(7/2)*Log[1 + c^(1/4)*Sqrt[x]] + 3*Sqrt[2]*b*c^(7/4)*x^(7/2)*Log[1 - Sqrt[2]*c^(1/4)*Sqrt[x] + Sqrt[c]*x] - 3*Sqrt[2]*b*c^(7/4)*x^(7/2)*Log[1 + Sqrt[2]*c^(1/4)*Sqrt[x] + Sqrt[c]*x]))/(42*d^5*x^4)
```

Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 349, normalized size of antiderivative = 1.36, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.889$, Rules used = {6464, 847, 851, 27, 830, 755, 756, 218, 221, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{barctanh}(cx^2)}{(dx)^{9/2}} dx$$

$$\downarrow 6464$$

$$\frac{4bc \int \frac{1}{(dx)^{5/2}(1-c^2x^4)} dx}{7d^2} - \frac{2(a + \operatorname{barctanh}(cx^2))}{7d(dx)^{7/2}}$$

$$\downarrow 847$$

$$\frac{4bc \left(\frac{c^2 \int \frac{(dx)^{3/2}}{1-c^2x^4} dx}{d^4} - \frac{2}{3d(dx)^{3/2}} \right)}{7d^2} - \frac{2(a + \operatorname{barctanh}(cx^2))}{7d(dx)^{7/2}}$$

$$\begin{aligned}
 & \downarrow 851 \\
 & \frac{4bc \left(\frac{2c^2 \int \frac{d^6 x^2}{d^4 - c^2 d^4 x^4} d\sqrt{dx} - \frac{2}{3d(dx)^{3/2}} \right)}{7d^2} - \frac{2(a + \operatorname{barctanh}(cx^2))}{7d(dx)^{7/2}} \\
 & \downarrow 27 \\
 & \frac{4bc \left(\frac{2c^2 \int \frac{d^2 x^2}{d^4 - c^2 d^4 x^4} d\sqrt{dx} - \frac{2}{3d(dx)^{3/2}} \right)}{7d^2} - \frac{2(a + \operatorname{barctanh}(cx^2))}{7d(dx)^{7/2}} \\
 & \downarrow 830 \\
 & \frac{4bc \left(\frac{2c^2 \left(\frac{\int \frac{1}{d^2 - cd^2 x^2} d\sqrt{dx}}{2c} - \frac{\int \frac{1}{cx^2 d^2 + d^2} d\sqrt{dx}}{2c} \right)}{d} - \frac{2}{3d(dx)^{3/2}} \right)}{7d^2} - \frac{2(a + \operatorname{barctanh}(cx^2))}{7d(dx)^{7/2}} \\
 & \downarrow 755 \\
 & \frac{4bc \left(\frac{2c^2 \left(\frac{\int \frac{1}{d^2 - cd^2 x^2} d\sqrt{dx}}{2c} - \frac{\int \frac{d - \sqrt{cdx}}{cx^2 d^2 + d^2} d\sqrt{dx}}{2d} + \frac{\int \frac{-\sqrt{cdx} + d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2c} \right)}{d} - \frac{2}{3d(dx)^{3/2}} \right)}{7d^2} - \frac{2(a + \operatorname{barctanh}(cx^2))}{7d(dx)^{7/2}} \\
 & \downarrow 756 \\
 & \frac{4bc \left(\frac{2c^2 \left(\frac{\int \frac{1}{d - \sqrt{cdx}} d\sqrt{dx}}{2d} + \frac{\int \frac{1}{\sqrt{cdx} + d} d\sqrt{dx}}{2c} - \frac{\int \frac{d - \sqrt{cdx}}{cx^2 d^2 + d^2} d\sqrt{dx}}{2d} + \frac{\int \frac{-\sqrt{cdx} + d}{cx^2 d^2 + d^2} d\sqrt{dx}}{2c} \right)}{d} - \frac{2}{3d(dx)^{3/2}} \right)}{7d^2} - \frac{2(a + \operatorname{barctanh}(cx^2))}{7d(dx)^{7/2}} \\
 & \downarrow 218
 \end{aligned}$$

$$4bc \left(\frac{2c^2 \left(\frac{\int \frac{1}{d-\sqrt{cdx}} d\sqrt{dx} + \frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx} + \frac{\int \frac{\sqrt{cx}d+d}{cx^2d^2+d^2} d\sqrt{dx}}{2c}}{d} \right) - \frac{2}{3d(dx)^{3/2}}}{d} \right)$$

$$\frac{7d^2}{2(a + \operatorname{barctanh}(cx^2))} \frac{1}{7d(dx)^{7/2}}$$

221

$$4bc \left(\frac{2c^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx} + \frac{\int \frac{\sqrt{cx}d+d}{cx^2d^2+d^2} d\sqrt{dx}}{2c}}{d} \right) - \frac{2}{3d(dx)^{3/2}}}{d} \right)$$

$$\frac{7d^2}{2(a + \operatorname{barctanh}(cx^2))} \frac{1}{7d(dx)^{7/2}}$$

1476

$$4bc \left(\frac{2c^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} - \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2d} + \frac{\int \frac{1}{xd+\frac{d}{\sqrt{c}}-\frac{1}{\sqrt{2}\sqrt{dx}\sqrt{d}}} d\sqrt{dx}}{2\sqrt{c}} + \frac{\int \frac{1}{xd+\frac{d}{\sqrt{c}}+\frac{1}{\sqrt{2}\sqrt{dx}\sqrt{d}}} d\sqrt{dx}}{2\sqrt{c}}}{d} \right) - \frac{2}{3d(dx)^{3/2}}}{d} \right)$$

$$\frac{7d^2}{2(a + \operatorname{barctanh}(cx^2))} \frac{1}{7d(dx)^{7/2}}$$

↓ 1082

$$4bc \left(\frac{2c^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right) + \operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2d} + \frac{\int \frac{1}{-dx-1} d \left(1 - \frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\int \frac{1}{-dx-1} d \left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}} + 1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}}}{2c} \right)}{d} - \frac{3d}{2c} \right)$$

$$\frac{2(a + b\operatorname{arctanh}(cx^2))}{7d(dx)^{7/2}} \cdot 7d^2$$

↓ 217

$$4bc \left(\frac{2c^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right) + \operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\int \frac{d-\sqrt{cdx}}{cx^2d^2+d^2} d\sqrt{dx}}{2d} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}} + 1\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}}}{2c} \right)}{d} - \frac{2}{3d(dx)^{3/2}} \right)$$

$$\frac{2(a + b\operatorname{arctanh}(cx^2))}{7d(dx)^{7/2}} \cdot 7d^2$$

↓ 1479

$$\left. \begin{array}{l} 2c^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} - \frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{C}\sqrt{dx}}{\sqrt[4]{C}\left(xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\int \frac{\sqrt{2}(\sqrt{d}+\sqrt{2}\sqrt[4]{C}\sqrt{dx})}{\sqrt[4]{C}\left(xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right) \\ 4bc \end{array} \right\} d$$

$$\frac{2(a + \operatorname{barctanh}(cx^2))}{7d(dx)^{7/2}} \quad 7d^2$$

↓ 25

$$\left. \begin{array}{l} 2c^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} - \frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{C}\sqrt{dx}}{\sqrt[4]{C}\left(xd+\frac{d}{\sqrt{c}}-\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} + \frac{\int \frac{\sqrt{2}(\sqrt{d}+\sqrt{2}\sqrt[4]{C}\sqrt{dx})}{\sqrt[4]{C}\left(xd+\frac{d}{\sqrt{c}}+\frac{\sqrt{2}\sqrt{dx}\sqrt{d}}{\sqrt[4]{C}}\right)} d\sqrt{dx}}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right) \\ 4bc \end{array} \right\} d$$

$$\frac{2(a + \operatorname{barctanh}(cx^2))}{7d(dx)^{7/2}} \quad 7d^2$$

↓ 27

$$\left. \begin{array}{l} 2c^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} - \frac{\int \frac{\sqrt{2}\sqrt{d}-2\sqrt[4]{C}\sqrt{dx}}{xd+\frac{d}{\sqrt{c}}-\sqrt{2}\sqrt{dx}\sqrt{d}} d\sqrt{dx}}{2\sqrt{2}\sqrt{c}\sqrt{d}} + \frac{\int \frac{\sqrt{d}+\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{xd+\frac{d}{\sqrt{c}}+\sqrt{2}\sqrt{dx}\sqrt{d}} d\sqrt{dx}}{2\sqrt{c}\sqrt{d}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}+1}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}-1}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right) \\ 4bc \end{array} \right\} d$$

$$\frac{2(a + b\operatorname{arctanh}(cx^2))}{7d(dx)^{7/2}} \quad 7d^2$$

1103

$$\left. \begin{array}{l} 2c^2 \left(\frac{\arctan\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{2\sqrt[4]{Cd^{3/2}}} - \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}+1}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{C}\sqrt{dx}}{\sqrt{d}}\right)}{\sqrt{2}\sqrt[4]{C}\sqrt{d}} + \frac{\log(\sqrt{cdx}+\sqrt{2}\sqrt[4]{C}\sqrt{d}\sqrt{dx}+d)}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} - \frac{\log(\sqrt{cdx}+\sqrt{2}\sqrt[4]{C}\sqrt{d}\sqrt{dx}-d)}{2\sqrt{2}\sqrt[4]{C}\sqrt{d}} \right) \\ 4bc \end{array} \right\} d$$

$$\frac{2(a + b\operatorname{arctanh}(cx^2))}{7d(dx)^{7/2}} \quad 7d^2$$

input

```
Int[(a + b*ArcTanh[c*x^2])/(d*x)^(9/2), x]
```

output

$$\begin{aligned} & (-2*(a + b*\text{ArcTanh}[c*x^2]))/(7*d*(d*x)^{(7/2)}) + (4*b*c*(-2/(3*d*(d*x)^{(3/2)})) \\ & + (2*c^2*((\text{ArcTan}[(c^{1/4})*\text{Sqrt}[d*x])/ \text{Sqrt}[d]]/(2*c^{1/4}*d^{(3/2)}) + \text{Arc} \\ & \text{Tanh}[(c^{1/4})*\text{Sqrt}[d*x])/ \text{Sqrt}[d]]/(2*c^{1/4}*d^{(3/2)})))/(2*c) - ((-\text{ArcTan} \\ & [1 - (\text{Sqrt}[2]*c^{1/4})*\text{Sqrt}[d*x])/ \text{Sqrt}[d]]/(\text{Sqrt}[2]*c^{1/4})*\text{Sqrt}[d])) + \text{Arc} \\ & \text{Tan}[1 + (\text{Sqrt}[2]*c^{1/4})*\text{Sqrt}[d*x])/ \text{Sqrt}[d]]/(\text{Sqrt}[2]*c^{1/4})*\text{Sqrt}[d]]/(2 \\ & *d) + (-1/2*\text{Log}[d + \text{Sqrt}[c]*d*x - \text{Sqrt}[2]*c^{1/4})*\text{Sqrt}[d]*\text{Sqrt}[d*x]]/(\text{Sqrt} \\ & [2]*c^{1/4})*\text{Sqrt}[d] + \text{Log}[d + \text{Sqrt}[c]*d*x + \text{Sqrt}[2]*c^{1/4})*\text{Sqrt}[d]*\text{Sqrt} \\ & [d*x]]/(2*\text{Sqrt}[2]*c^{1/4})*\text{Sqrt}[d]]/(2*d))/(2*c))/d)/(7*d^2) \end{aligned}$$

Defintions of rubi rules used

rule 25

$$\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$$

rule 27

$$\text{Int}[(a_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] /; \text{FreeQ}[a, \text{x}] \&\& \text{!MatchQ}[\text{Fx}, (b_)*(\text{Gx}_) /; \text{FreeQ}[b, \text{x}]$$

rule 217

$$\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& \text{PosQ}[a/b] \&\& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])$$

rule 218

$$\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& \text{PosQ}[a/b]$$

rule 221

$$\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& \text{NegQ}[a/b]$$

rule 755

$$\text{Int}[((a_) + (b_.)*(x_)^4)^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*r) \quad \text{Int}[(r - s*x^2)/(a + b*x^4), \text{x}], \text{x}] + \text{Simp}[1/(2*r) \quad \text{Int}[(r + s*x^2)/(a + b*x^4), \text{x}], \text{x}]] /; \text{FreeQ}[\{a, b\}, \text{x}] \&\& (\text{GtQ}[a/b, 0] \parallel (\text{PosQ}[a/b] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$$

rule 756 $\text{Int}[(a + (b \cdot x)^4)^{-1}, x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[r/(2 \cdot a) \text{Int}[1/(r - s \cdot x^2), x], x] + \text{Simp}[r/(2 \cdot a) \text{Int}[1/(r + s \cdot x^2), x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ !\text{GtQ}[a/b, 0]$

rule 830 $\text{Int}[(x^m)/((a + (b \cdot x)^n)), x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[s/(2 \cdot b) \text{Int}[x^{(m - n/2)} / (r + s \cdot x^{(n/2)})], x], x] - \text{Simp}[s/(2 \cdot b) \text{Int}[x^{(m - n/2)} / (r - s \cdot x^{(n/2)})], x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{IGtQ}[n/4, 0] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{LeQ}[n/2, m] \ \&\& \ \text{LtQ}[m, n] \ \&\& \ !\text{GtQ}[a/b, 0]$

rule 847 $\text{Int}[(c \cdot x)^m \cdot (a + (b \cdot x)^n)^p, x_Symbol] \rightarrow \text{Simp}[(c \cdot x)^{(m + 1)} \cdot ((a + b \cdot x^n)^{(p + 1)} / (a \cdot c \cdot (m + 1))), x] - \text{Simp}[b \cdot ((m + n \cdot (p + 1) + 1) / (a \cdot c^n \cdot (m + 1))) \text{Int}[(c \cdot x)^{(m + n)} \cdot (a + b \cdot x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 851 $\text{Int}[(c \cdot x)^m \cdot (a + (b \cdot x)^n)^p, x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \text{Subst}[\text{Int}[x^{(k \cdot (m + 1) - 1)} \cdot (a + b \cdot (x^{(k \cdot n)}) / c^n)^p], x, (c \cdot x)^{(1/k)}], x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 1082 $\text{Int}[(a + (b \cdot x) + (c \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4 \cdot S\text{implify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4 \cdot a \cdot c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}[(d + (e \cdot x)) / ((a + (b \cdot x) + (c \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]] / b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$

rule 1476

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

rule 1479

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

rule 6464

```
Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)*((d_)*(x_)^(m_)), x_Symbol] :
> Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n
/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a,
b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 250, normalized size of antiderivative = 0.97

method	result
derivativedivides	$-\frac{2a}{7(dx)^{\frac{7}{2}}} + 2b \left[-\frac{\operatorname{arctanh}(cx^2)}{7(dx)^{\frac{7}{2}}} + \frac{4cd^2}{8d^6} \left(c\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \left(\ln \left(\frac{\sqrt{dx} + \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}{\sqrt{dx} - \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) \right) \right] - \frac{c\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{2}}{d} \ln \left(\frac{dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)$
default	$-\frac{2a}{7(dx)^{\frac{7}{2}}} + 2b \left[-\frac{\operatorname{arctanh}(cx^2)}{7(dx)^{\frac{7}{2}}} + \frac{4cd^2}{8d^6} \left(c\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \left(\ln \left(\frac{\sqrt{dx} + \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}{\sqrt{dx} - \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) \right) \right] - \frac{c\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{2}}{d} \ln \left(\frac{dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)$
parts	$-\frac{2a}{7(dx)^{\frac{7}{2}}d} + 2b \left[-\frac{\operatorname{arctanh}(cx^2)}{7(dx)^{\frac{7}{2}}} + \frac{4cd^2}{8d^6} \left(c\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \left(\ln \left(\frac{\sqrt{dx} + \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}{\sqrt{dx} - \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) + 2 \operatorname{arctan} \left(\frac{\sqrt{dx}}{\left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right) \right) \right] - \frac{c\left(\frac{d^2}{c}\right)^{\frac{1}{4}} \sqrt{2}}{d} \ln \left(\frac{dx + \left(\frac{d^2}{c}\right)^{\frac{1}{4}}}{dx - \left(\frac{d^2}{c}\right)^{\frac{1}{4}}} \right)$

input

```
int((a+b*arctanh(c*x^2))/(d*x)^(9/2),x,method=_RETURNVERBOSE)
```

output

```
2/d*(-1/7*a/(d*x)^(7/2)+b*(-1/7/(d*x)^(7/2)*arctanh(c*x^2)+4/7*c*d^2*(1/8/
d^6*c*(d^2/c)^(1/4)*(ln(((d*x)^(1/2)+(d^2/c)^(1/4))/((d*x)^(1/2)-(d^2/c)^(
1/4))))+2*arctan((d*x)^(1/2)/(d^2/c)^(1/4))-1/16/d^6*c*(d^2/c)^(1/4)*2^(1/
2)*(ln((d*x+(d^2/c)^(1/4)*(d*x)^(1/2)*2^(1/2)+(d^2/c)^(1/2)))/(d*x-(d^2/c)^(
1/4)*(d*x)^(1/2)*2^(1/2)+(d^2/c)^(1/2))))+2*arctan(2^(1/2)/(d^2/c)^(1/4)*(
d*x)^(1/2)+1)+2*arctan(2^(1/2)/(d^2/c)^(1/4)*(d*x)^(1/2)-1))-1/3/d^4/(d*x)
^(3/2))))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 447, normalized size of antiderivative = 1.74

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{9/2}} dx = \frac{3 d^5 x^4 \left(\frac{b^4 c^7}{d^{18}}\right)^{\frac{1}{4}} \log \left(d^5 \left(\frac{b^4 c^7}{d^{18}}\right)^{\frac{1}{4}} + \sqrt{d x b c^2} \right) + 3 i d^5 x^4 \left(\frac{b^4 c^7}{d^{18}}\right)^{\frac{1}{4}} \log \left(i d^5 \left(\frac{b^4 c^7}{d^{18}}\right)^{\frac{1}{4}} + \sqrt{d x b c^2} \right)}{d^5 x^4 \left(\frac{b^4 c^7}{d^{18}}\right)^{\frac{1}{4}} \log \left(d^5 \left(\frac{b^4 c^7}{d^{18}}\right)^{\frac{1}{4}} + \sqrt{d x b c^2} \right) + 3 i d^5 x^4 \left(\frac{b^4 c^7}{d^{18}}\right)^{\frac{1}{4}} \log \left(i d^5 \left(\frac{b^4 c^7}{d^{18}}\right)^{\frac{1}{4}} + \sqrt{d x b c^2} \right)}$$

input

```
integrate((a+b*arctanh(c*x^2))/(d*x)^(9/2),x, algorithm="fricas")
```

output

```
1/21*(3*d^5*x^4*(b^4*c^7/d^18)^(1/4)*log(d^5*(b^4*c^7/d^18)^(1/4) + sqrt(d
*x)*b*c^2) + 3*I*d^5*x^4*(b^4*c^7/d^18)^(1/4)*log(I*d^5*(b^4*c^7/d^18)^(1/
4) + sqrt(d*x)*b*c^2) - 3*I*d^5*x^4*(b^4*c^7/d^18)^(1/4)*log(-I*d^5*(b^4*c
^7/d^18)^(1/4) + sqrt(d*x)*b*c^2) - 3*d^5*x^4*(b^4*c^7/d^18)^(1/4)*log(-d^
5*(b^4*c^7/d^18)^(1/4) + sqrt(d*x)*b*c^2) - 3*d^5*x^4*(-b^4*c^7/d^18)^(1/4
)*log(d^5*(-b^4*c^7/d^18)^(1/4) + sqrt(d*x)*b*c^2) - 3*I*d^5*x^4*(-b^4*c^7
/d^18)^(1/4)*log(I*d^5*(-b^4*c^7/d^18)^(1/4) + sqrt(d*x)*b*c^2) + 3*I*d^5*
x^4*(-b^4*c^7/d^18)^(1/4)*log(-I*d^5*(-b^4*c^7/d^18)^(1/4) + sqrt(d*x)*b*c
^2) + 3*d^5*x^4*(-b^4*c^7/d^18)^(1/4)*log(-d^5*(-b^4*c^7/d^18)^(1/4) + sqr
t(d*x)*b*c^2) - (8*b*c*x^2 + 3*b*log(-(c*x^2 + 1)/(c*x^2 - 1)) + 6*a)*sqrt
(d*x))/(d^5*x^4)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{9/2}} dx = \text{Timed out}$$

```
input integrate((a+b*atanh(c*x**2))/(d*x)**(9/2), x)
```

```
output Timed out
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.16

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{9/2}} dx =$$

$$\frac{c \left(\frac{6 \sqrt{2} c \arctan \left(\frac{\sqrt{2} \left(\sqrt{2} c^{\frac{1}{4}} \sqrt{d} + 2 \sqrt{dx} \sqrt{c} \right)}{2 \sqrt{\sqrt{cd}}} \right)}{\sqrt{\sqrt{cd} d}} \right) + \frac{6 \sqrt{2} c \arctan \left(-\frac{\sqrt{2} \left(\sqrt{2} c^{\frac{1}{4}} \sqrt{d} - 2 \sqrt{dx} \sqrt{c} \right)}{2 \sqrt{\sqrt{cd}}} \right)}{\sqrt{\sqrt{cd} d}} + \frac{3 \sqrt{2} c^{\frac{3}{4}} \log \left(\sqrt{cd} x + \sqrt{2} \sqrt{dx} c^{\frac{1}{4}} \sqrt{d} + d \right)}{d^{\frac{3}{2}}} - \frac{3 \sqrt{2} c^{\frac{3}{4}} \log \left(\sqrt{cd} x - \sqrt{2} \sqrt{dx} c^{\frac{1}{4}} \sqrt{d} + d \right)}{d^{\frac{3}{2}}} \right)}{b d^2}$$

$42 d$

```
input integrate((a+b*arctanh(c*x^2))/(d*x)^(9/2), x, algorithm="maxima")
```

output

```
-1/42*(b*(c*(6*sqrt(2)*c*arctan(1/2*sqrt(2)*(sqrt(2)*c^(1/4)*sqrt(d) + 2*sqrt(d*x)*sqrt(c))/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d)*d) + 6*sqrt(2)*c*arctan(-1/2*sqrt(2)*(sqrt(2)*c^(1/4)*sqrt(d) - 2*sqrt(d*x)*sqrt(c))/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d)*d) + 3*sqrt(2)*c^(3/4)*log(sqrt(c)*d*x + sqrt(2)*sqrt(d*x)*c^(1/4)*sqrt(d) + d)/d^(3/2) - 3*sqrt(2)*c^(3/4)*log(sqrt(c)*d*x - sqrt(2)*sqrt(d*x)*c^(1/4)*sqrt(d) + d)/d^(3/2) - 12*c*arctan(sqrt(d*x)*sqrt(c)/sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d)*d) + 6*c*log((sqrt(d*x)*sqrt(c) - sqrt(sqrt(c)*d))/sqrt(d*x)*sqrt(c) + sqrt(sqrt(c)*d))/sqrt(sqrt(c)*d)*d) + 16/(d*x)^(3/2))/d^2 + 12*arctanh(c*x^2)/(d*x)^(7/2) + 12*a/(d*x)^(7/2))/d
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 519 vs. $2(174) = 348$.

Time = 5.07 (sec) , antiderivative size = 519, normalized size of antiderivative = 2.02

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{9/2}} dx =$$

$$\frac{6\sqrt{2}(c^3d^2)^{\frac{1}{4}}bc \operatorname{arctan}\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 2\sqrt{dx}\right)}{2\left(\frac{d^2}{c}\right)^{\frac{1}{4}}}\right)}{d^4} + \frac{6\sqrt{2}(c^3d^2)^{\frac{1}{4}}bc \operatorname{arctan}\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{d^2}{c}\right)^{\frac{1}{4}} - 2\sqrt{dx}\right)}{2\left(\frac{d^2}{c}\right)^{\frac{1}{4}}}\right)}{d^4} - \frac{6\sqrt{2}(-c^3d^2)^{\frac{1}{4}}bc \operatorname{arctan}\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{d^2}{c}\right)^{\frac{1}{4}} + 2\sqrt{dx}\right)}{2\left(\frac{d^2}{c}\right)^{\frac{1}{4}}}\right)}{d^4}$$

input

```
integrate((a+b*arctanh(c*x^2))/(d*x)^(9/2),x, algorithm="giac")
```

output

```
-1/42*(6*sqrt(2)*(c^3*d^2)^(1/4)*b*c*arctan(1/2*sqrt(2)*(sqrt(2)*(d^2/c)^(1/4) + 2*sqrt(d*x))/(d^2/c)^(1/4))/d^4 + 6*sqrt(2)*(c^3*d^2)^(1/4)*b*c*arctan(-1/2*sqrt(2)*(sqrt(2)*(d^2/c)^(1/4) - 2*sqrt(d*x))/(d^2/c)^(1/4))/d^4 - 6*sqrt(2)*(-c^3*d^2)^(1/4)*b*c*arctan(1/2*sqrt(2)*(sqrt(2)*(-d^2/c)^(1/4) + 2*sqrt(d*x))/(-d^2/c)^(1/4))/d^4 - 6*sqrt(2)*(-c^3*d^2)^(1/4)*b*c*arctan(-1/2*sqrt(2)*(sqrt(2)*(-d^2/c)^(1/4) - 2*sqrt(d*x))/(-d^2/c)^(1/4))/d^4 + 3*sqrt(2)*(c^3*d^2)^(1/4)*b*c*log(d*x + sqrt(2)*sqrt(d*x)*(d^2/c)^(1/4) + sqrt(d^2/c))/d^4 - 3*sqrt(2)*(c^3*d^2)^(1/4)*b*c*log(d*x - sqrt(2)*sqrt(d*x)*(d^2/c)^(1/4) + sqrt(d^2/c))/d^4 - 3*sqrt(2)*(-c^3*d^2)^(1/4)*b*c*log(d*x + sqrt(2)*sqrt(d*x)*(-d^2/c)^(1/4) + sqrt(-d^2/c))/d^4 + 3*sqrt(2)*(-c^3*d^2)^(1/4)*b*c*log(d*x - sqrt(2)*sqrt(d*x)*(-d^2/c)^(1/4) + sqrt(-d^2/c))/d^4 + 6*b*log(-(c*d^2*x^2 + d^2)/(c*d^2*x^2 - d^2))/(sqrt(d*x)*d^3*x^3) + 4*(4*b*c*d^2*x^2 + 3*a*d^2)/(sqrt(d*x)*d^5*x^3))/d
```

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{9/2}} dx = \int \frac{a + b \operatorname{atanh}(cx^2)}{(dx)^{9/2}} dx$$

input

```
int((a + b*atanh(c*x^2))/(d*x)^(9/2), x)
```

output

```
int((a + b*atanh(c*x^2))/(d*x)^(9/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.11

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(dx)^{9/2}} dx = \frac{\sqrt{d} \left(6\sqrt{x} c^{\frac{7}{4}} \sqrt{2} \operatorname{atan} \left(\frac{c^{\frac{1}{4}} \sqrt{2} - 2\sqrt{x} \sqrt{c}}{c^{\frac{1}{4}} \sqrt{2}} \right) b x^3 - 6\sqrt{x} c^{\frac{7}{4}} \sqrt{2} \operatorname{atan} \left(\frac{c^{\frac{1}{4}} \sqrt{2} + 2\sqrt{x} \sqrt{c}}{c^{\frac{1}{4}} \sqrt{2}} \right) b x^3 \right)}{d^{\frac{7}{2}}}$$

input

```
int((a+b*atanh(c*x^2))/(d*x)^(9/2), x)
```

output

```
(sqrt(d)*(6*sqrt(x)*c**(3/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*b*c*x**3 - 6*sqrt(x)*c**(3/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*b*c*x**3 + 12*sqrt(x)*c**(3/4)*atan((sqrt(x)*sqrt(c))/c**(1/4))*b*c*x**3 - 6*sqrt(x)*c**(3/4)*sqrt(2)*atanh(c*x**2)*b*c*x**3 - 12*atanh(c*x**2)*b - 3*sqrt(x)*c**(3/4)*sqrt(2)*log(c**(1/4) + sqrt(x)*sqrt(c))*b*c*x**3 - 3*sqrt(x)*c**(3/4)*sqrt(2)*log(-c**(1/4) + sqrt(x)*sqrt(c))*b*c*x**3 + 6*sqrt(x)*c**(3/4)*sqrt(2)*log(-sqrt(x)*c**(1/4)*sqrt(2) + sqrt(c)*x + 1)*b*c*x**3 - 3*sqrt(x)*c**(3/4)*sqrt(2)*log(sqrt(c)*x + 1)*b*c*x**3 + 6*sqrt(x)*c**(3/4)*log(c**(1/4) + sqrt(x)*sqrt(c))*b*c*x**3 - 6*sqrt(x)*c**(3/4)*log(-c**(1/4) + sqrt(x)*sqrt(c))*b*c*x**3 - 12*a - 16*b*c*x**2))/(42*sqrt(x)*d**5*x**3)
```

3.90 $\int \sqrt{dx}(a + \operatorname{barctanh}(cx^2))^2 dx$

Optimal result	803
Mathematica [F]	804
Rubi [A] (warning: unable to verify)	805
Maple [F]	807
Fricas [F]	807
Sympy [F]	807
Maxima [F]	808
Giac [F]	808
Mupad [F(-1)]	809
Reduce [F]	809

Optimal result

Integrand size = 20, antiderivative size = 6274

$$\int \sqrt{dx}(a + \operatorname{barctanh}(cx^2))^2 dx = \text{Too large to display}$$

output

```

4/3*b^2*(d*x)^(1/2)*arctan(c^(1/4)*x^(1/2))*ln(2/(1-I*c^(1/4)*x^(1/2)))/c^(
(3/4)/x^(1/2)+2/3*b^2*(d*x)^(1/2)*arctan(c^(1/4)*x^(1/2))*ln(c*x^2+1)/c^(3
/4)/x^(1/2)+2/3*b^2*(d*x)^(1/2)*arctanh((-c)^(1/4)*x^(1/2))*ln(c*x^2+1)/(-
c)^(3/4)/x^(1/2)-2/3*b^2*(d*x)^(1/2)*arctan((-c)^(1/4)*x^(1/2))*ln(c*x^2+1
)/(-c)^(3/4)/x^(1/2)-2/3*b^2*(d*x)^(1/2)*arctanh(c^(1/4)*x^(1/2))*ln(c*x^2
+1)/c^(3/4)/x^(1/2)+2/3*b*(d*x)^(1/2)*arctan(c^(1/4)*x^(1/2))*(2*a-b*ln(-c
*x^2+1))/c^(3/4)/x^(1/2)-2/3*b^2*(d*x)^(1/2)*arctanh((-c)^(1/4)*x^(1/2))*l
n(-c*x^2+1)/(-c)^(3/4)/x^(1/2)+2/3*b^2*(d*x)^(1/2)*arctan((-c)^(1/4)*x^(1/
2))*ln(-c*x^2+1)/(-c)^(3/4)/x^(1/2)-2/3*b*(d*x)^(1/2)*arctanh(c^(1/4)*x^(1
/2))*(2*a-b*ln(-c*x^2+1))/c^(3/4)/x^(1/2)-2/3*b^2*(d*x)^(1/2)*arctanh((-c)
^(1/4)*x^(1/2))*ln(2*(-c)^(1/4)*(1+((-c)^(1/2))^(1/2)*x^(1/2)))/((-c)^(1
/2))^(1/2)+(-c)^(1/4))/(1+(-c)^(1/4)*x^(1/2)))/(-c)^(3/4)/x^(1/2)-2/3*b^2*
(d*x)^(1/2)*arctanh((-c)^(1/4)*x^(1/2))*ln(-2*(-c)^(1/4)*(1-((-c)^(1/2))^(
1/2)*x^(1/2)))/((-c)^(1/2))^(1/2)-(-c)^(1/4))/(1+(-c)^(1/4)*x^(1/2)))/(-
c)^(3/4)/x^(1/2)+2/3*b^2*(d*x)^(1/2)*arctanh((-c)^(1/4)*x^(1/2))*ln(2*(-c)
^(1/4)*(1+(-c)^(1/2))^(1/2)*x^(1/2)))/((-c)^(1/2))^(1/2)+(-c)^(1/4))/(1+(-c)
^(1/4)*x^(1/2)))/(-c)^(3/4)/x^(1/2)+2/3*b^2*(d*x)^(1/2)*arctanh((-c)^(1/4)*
x^(1/2))*ln(-2*(-c)^(1/4)*(1-(-c)^(1/2))^(1/2)*x^(1/2)))/((-c)^(1/2))-(-
c)^(1/4))/(1+(-c)^(1/4)*x^(1/2)))/(-c)^(3/4)/x^(1/2)+2/3*b^2*(d*x)^(1/2)*
arctanh((-c)^(1/4)*x^(1/2))*ln(2*(-c)^(1/4)*(1+c^(1/4)*x^(1/2)))/((-c)^(...

```

Mathematica [F]

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx^2))^2 dx = \int \sqrt{dx}(a + b \operatorname{arctanh}(cx^2))^2 dx$$

input

```
Integrate[Sqrt[d*x]*(a + b*ArcTanh[c*x^2])^2,x]
```

output

```
Integrate[Sqrt[d*x]*(a + b*ArcTanh[c*x^2])^2, x]
```

Rubi [A] (warning: unable to verify)

Time = 12.42 (sec) , antiderivative size = 5360, normalized size of antiderivative = 0.85, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {6466, 6458, 6456, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{dx} (a + b \operatorname{arctanh}(cx^2))^2 dx \\
 & \quad \downarrow \text{6466} \\
 & \frac{\sqrt{dx} \int \sqrt{x} (a + b \operatorname{arctanh}(cx^2))^2 dx}{\sqrt{x}} \\
 & \quad \downarrow \text{6458} \\
 & \frac{2\sqrt{dx} \int x (a + b \operatorname{arctanh}(cx^2))^2 d\sqrt{x}}{\sqrt{x}} \\
 & \quad \downarrow \text{6456} \\
 & \frac{2\sqrt{dx} \int \left(\frac{1}{4}x(2a - b \log(1 - cx^2))^2 + \frac{1}{4}b^2x \log^2(cx^2 + 1) - \frac{1}{2}bx(b \log(1 - cx^2) - 2a) \log(cx^2 + 1) \right) d\sqrt{x}}{\sqrt{x}} \\
 & \quad \downarrow \text{2009} \\
 & 2\sqrt{dx} \left(-\frac{i \arctan\left(\sqrt[4]{-c}\sqrt{x}\right)^2 b^2}{3(-c)^{3/4}} - \frac{i \arctan\left(\sqrt[4]{c}\sqrt{x}\right)^2 b^2}{3c^{3/4}} - \frac{\operatorname{arctanh}\left(\sqrt[4]{-c}\sqrt{x}\right)^2 b^2}{3(-c)^{3/4}} - \frac{\operatorname{arctanh}\left(\sqrt[4]{c}\sqrt{x}\right)^2 b^2}{3c^{3/4}} + \frac{1}{12}x^{3/2} \log^2(cx^2 + 1) \right)
 \end{aligned}$$

input `Int[Sqrt[d*x]*(a + b*ArcTanh[c*x^2])^2,x]`

output

```
(2*Sqrt[d*x]*((-4*a*b*x^(3/2))/9 - (Sqrt[2]*a*b*ArcTan[1 - Sqrt[2]*c^(1/4)
*Sqrt[x]])/(3*c^(3/4)) + (Sqrt[2]*a*b*ArcTan[1 + Sqrt[2]*c^(1/4)*Sqrt[x]])
/(3*c^(3/4)) - ((I/3)*b^2*ArcTan[(-c)^(1/4)*Sqrt[x]]^2)/(-c)^(3/4) - ((I/3)
)*b^2*ArcTan[c^(1/4)*Sqrt[x]]^2/c^(3/4) - (b^2*ArcTanh[(-c)^(1/4)*Sqrt[x]
]^2)/(3*(-c)^(3/4)) - (b^2*ArcTanh[c^(1/4)*Sqrt[x]]^2)/(3*c^(3/4)) + (2*b^
2*ArcTanh[(-c)^(1/4)*Sqrt[x]]*Log[2/(1 - (-c)^(1/4)*Sqrt[x])])/(3*(-c)^(3/
4)) + (2*b^2*ArcTan[(-c)^(1/4)*Sqrt[x]]*Log[2/(1 - I*(-c)^(1/4)*Sqrt[x])])
/(3*(-c)^(3/4)) - (b^2*ArcTan[(-c)^(1/4)*Sqrt[x]]*Log[(-2*(-c)^(1/4)*(1 -
Sqrt[-Sqrt[c]]*Sqrt[x])]/((I*Sqrt[-Sqrt[c]] - (-c)^(1/4))*(1 - I*(-c)^(1/4)
)*Sqrt[x])))/(3*(-c)^(3/4)) - (b^2*ArcTan[(-c)^(1/4)*Sqrt[x]]*Log[(2*(-c)
^(1/4)*(1 + Sqrt[-Sqrt[c]]*Sqrt[x])]/((I*Sqrt[-Sqrt[c]] + (-c)^(1/4))*(1 -
I*(-c)^(1/4)*Sqrt[x])))/(3*(-c)^(3/4)) + (b^2*ArcTan[(-c)^(1/4)*Sqrt[x]]
*Log[((1 + I)*(1 - (-c)^(1/4)*Sqrt[x])/(1 - I*(-c)^(1/4)*Sqrt[x])))/(3*(-
c)^(3/4)) - (2*b^2*ArcTan[(-c)^(1/4)*Sqrt[x]]*Log[2/(1 + I*(-c)^(1/4)*Sqrt
[x])])/(3*(-c)^(3/4)) - (2*b^2*ArcTanh[(-c)^(1/4)*Sqrt[x]]*Log[2/(1 + (-c)
^(1/4)*Sqrt[x])])/(3*(-c)^(3/4)) - (b^2*ArcTanh[(-c)^(1/4)*Sqrt[x]]*Log[(-
2*(-c)^(1/4)*(1 - Sqrt[-Sqrt[-c]]*Sqrt[x])]/((Sqrt[-Sqrt[-c]] - (-c)^(1/4)
)*(1 + (-c)^(1/4)*Sqrt[x])))/(3*(-c)^(3/4)) - (b^2*ArcTanh[(-c)^(1/4)*Sqr
t[x]]*Log[(2*(-c)^(1/4)*(1 + Sqrt[-Sqrt[-c]]*Sqrt[x])]/((Sqrt[-Sqrt[-c]] +
(-c)^(1/4))*(1 + (-c)^(1/4)*Sqrt[x])))/(3*(-c)^(3/4)) + (b^2*ArcTanh[...
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6456 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.)^(p_)*(x_)^(m_.), x_Symbol] :=
Int[ExpandIntegrand[x^m*(a + b*(Log[1 + c*x^n]/2) - b*(Log[1 - c*x^n]/2))^p
, x], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1] && IGtQ[n, 0] && IntegerQ[m]`

rule 6458 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.)^(p_)*(x_)^(m_.), x_Symbol] :=
With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a + b*ArcT
anh[c*x^(k*n)]]^p, x], x, x^(1/k)], x]] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1
] && IGtQ[n, 0] && FractionQ[m]`

rule 6466

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((d_.)*(x_)^(m_.), x_Sym
bol] :> Simp[d^IntPart[m]*((d*x)^FracPart[m]/x^FracPart[m]) Int[x^m*(a +
b*ArcTanh[c*x^n])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0] &
& (EqQ[p, 1] || RationalQ[m, n])
```

Maple [F]

$$\int \sqrt{dx} (a + b \operatorname{arctanh}(cx^2))^2 dx$$

input

```
int((d*x)^(1/2)*(a+b*arctanh(c*x^2))^2,x)
```

output

```
int((d*x)^(1/2)*(a+b*arctanh(c*x^2))^2,x)
```

Fricas [F]

$$\int \sqrt{dx} (a + b \operatorname{arctanh}(cx^2))^2 dx = \int \sqrt{dx} (b \operatorname{artanh}(cx^2) + a)^2 dx$$

input

```
integrate((d*x)^(1/2)*(a+b*arctanh(c*x^2))^2,x, algorithm="fricas")
```

output

```
integral((b^2*arctanh(c*x^2)^2 + 2*a*b*arctanh(c*x^2) + a^2)*sqrt(d*x), x)
```

Sympy [F]

$$\int \sqrt{dx} (a + b \operatorname{arctanh}(cx^2))^2 dx = \int \sqrt{dx} (a + b \operatorname{atanh}(cx^2))^2 dx$$

input

```
integrate((d*x)**(1/2)*(a+b*atanh(c*x**2))**2,x)
```

output

```
Integral(sqrt(d*x)*(a + b*atanh(c*x**2))**2, x)
```

Maxima [F]

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx^2))^2 dx = \int \sqrt{dx}(b \operatorname{artanh}(cx^2) + a)^2 dx$$

input `integrate((d*x)^(1/2)*(a+b*arctanh(c*x^2))^2,x, algorithm="maxima")`

output `1/6*b^2*sqrt(d)*x^(3/2)*log(-c*x^2 + 1)^2 + 1/6*a^2*c*sqrt(d)*(4*x^(3/2)/c - 3*(I*(log(I*c^(1/4)*sqrt(x) + 1) - log(-I*c^(1/4)*sqrt(x) + 1))/c^(3/4) - log((sqrt(c)*sqrt(x) - c^(1/4))/(sqrt(c)*sqrt(x) + c^(1/4)))/c^(3/4))/c) + 3*b^2*c*sqrt(d)*integrate(1/12*x^(5/2)*log(c*x^2 + 1)^2/(c*x^2 - 1), x) - 6*b^2*c*sqrt(d)*integrate(1/12*x^(5/2)*log(c*x^2 + 1)*log(-c*x^2 + 1)/(c*x^2 - 1), x) + 12*a*b*c*sqrt(d)*integrate(1/12*x^(5/2)*log(c*x^2 + 1)/(c*x^2 - 1), x) - 12*a*b*c*sqrt(d)*integrate(1/12*x^(5/2)*log(-c*x^2 + 1)/(c*x^2 - 1), x) - 8*b^2*c*sqrt(d)*integrate(1/12*x^(5/2)*log(-c*x^2 + 1)/(c*x^2 - 1), x) + 1/2*a^2*sqrt(d)*(I*(log(I*c^(1/4)*sqrt(x) + 1) - log(-I*c^(1/4)*sqrt(x) + 1))/c^(3/4) - log((sqrt(c)*sqrt(x) - c^(1/4))/(sqrt(c)*sqrt(x) + c^(1/4)))/c^(3/4)) - 3*b^2*sqrt(d)*integrate(1/12*sqrt(x)*log(c*x^2 + 1)^2/(c*x^2 - 1), x) + 6*b^2*sqrt(d)*integrate(1/12*sqrt(x)*log(c*x^2 + 1)*log(-c*x^2 + 1)/(c*x^2 - 1), x) - 12*a*b*sqrt(d)*integrate(1/12*sqrt(x)*log(c*x^2 + 1)/(c*x^2 - 1), x) + 12*a*b*sqrt(d)*integrate(1/12*sqrt(x)*log(-c*x^2 + 1)/(c*x^2 - 1), x)`

Giac [F]

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx^2))^2 dx = \int \sqrt{dx}(b \operatorname{artanh}(cx^2) + a)^2 dx$$

input `integrate((d*x)^(1/2)*(a+b*arctanh(c*x^2))^2,x, algorithm="giac")`

output `integrate(sqrt(d*x)*(b*arctanh(c*x^2) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx^2))^2 dx = \int \sqrt{dx}(a + b \operatorname{atanh}(cx^2))^2 dx$$

input `int((d*x)^(1/2)*(a + b*atanh(c*x^2))^2,x)`

output `int((d*x)^(1/2)*(a + b*atanh(c*x^2))^2, x)`

Reduce [F]

$$\int \sqrt{dx}(a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$= \frac{\sqrt{d} \left(-2c^{\frac{1}{4}}\sqrt{2} \operatorname{atan}\left(\frac{c^{\frac{1}{4}}\sqrt{2}-2\sqrt{x}\sqrt{c}}{c^{\frac{1}{4}}\sqrt{2}}\right) ab + 2c^{\frac{1}{4}}\sqrt{2} \operatorname{atan}\left(\frac{c^{\frac{1}{4}}\sqrt{2}+2\sqrt{x}\sqrt{c}}{c^{\frac{1}{4}}\sqrt{2}}\right) ab + 4c^{\frac{1}{4}} \operatorname{atan}\left(\frac{\sqrt{x}\sqrt{c}}{c^{\frac{1}{4}}}\right) ab - 2c^{\frac{1}{4}}\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{x}\sqrt{c}}{c^{\frac{1}{4}}}\right) ab \right)}{3c}$$

input `int((d*x)^(1/2)*(a+b*atanh(c*x^2))^2,x)`

output `(sqrt(d)*(-2*c**(1/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*a*b + 2*c**(1/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*a*b + 4*c**(1/4)*atan((sqrt(x)*sqrt(c))/(c**(1/4)))*a*b - 2*c**(1/4)*sqrt(2)*atanh(c*x**2)*a*b + 4*sqrt(x)*atanh(c*x**2)*a*b*c*x - c**(1/4)*sqrt(2)*log(c**(1/4) + sqrt(x)*sqrt(c))*a*b - c**(1/4)*sqrt(2)*log(-c**(1/4) + sqrt(x)*sqrt(c))*a*b + 2*c**(1/4)*sqrt(2)*log(-sqrt(x)*c**(1/4)*sqrt(2) + sqrt(c)*x + 1)*a*b - c**(1/4)*sqrt(2)*log(sqrt(c)*x + 1)*a*b - 2*c**(1/4)*log(c**(1/4) + sqrt(x)*sqrt(c))*a*b + 2*c**(1/4)*log(-c**(1/4) + sqrt(x)*sqrt(c))*a*b + 2*sqrt(x)*a**2*c*x + 3*int(sqrt(x)*atanh(c*x**2)**2,x)*b**2*c))/(3*c)`

$$3.91 \quad \int \frac{(a+b\operatorname{arctanh}(cx^2))^2}{\sqrt{dx}} dx$$

Optimal result	810
Mathematica [F]	811
Rubi [A] (verified)	812
Maple [F]	814
Fricas [F]	814
Sympy [F]	814
Maxima [F]	815
Giac [F]	815
Mupad [F(-1)]	816
Reduce [F]	816

Optimal result

Integrand size = 20, antiderivative size = 6127

$$\int \frac{(a + b\operatorname{arctanh}(cx^2))^2}{\sqrt{dx}} dx = \text{Too large to display}$$

output

```

-2*b^2*x^(1/2)*arctanh(c^(1/4)*x^(1/2))*ln(c*x^2+1)/c^(1/4)/(d*x)^(1/2)-2*
b^2*x^(1/2)*arctan(c^(1/4)*x^(1/2))*ln(c*x^2+1)/c^(1/4)/(d*x)^(1/2)+2*b^2*
x^(1/2)*arctanh((-c)^(1/4)*x^(1/2))*ln(c*x^2+1)/(-c)^(1/4)/(d*x)^(1/2)-2*b
^2*x^(1/2)*arctan((-c)^(1/4)*x^(1/2))*ln(-c*x^2+1)/(-c)^(1/4)/(d*x)^(1/2)+
2*b^2*x^(1/2)*arctanh(c^(1/4)*x^(1/2))*ln(-c*x^2+1)/c^(1/4)/(d*x)^(1/2)+2*
b^2*x^(1/2)*arctan(c^(1/4)*x^(1/2))*ln(-c*x^2+1)/c^(1/4)/(d*x)^(1/2)-2*b^2
*x^(1/2)*arctanh((-c)^(1/4)*x^(1/2))*ln(-c*x^2+1)/(-c)^(1/4)/(d*x)^(1/2)-I
*b^2*x^(1/2)*polylog(2,1-2*c^(1/4)*(1+(-(-c)^(1/2))^(1/2)*x^(1/2))/(I*(-(-
c)^(1/2))^(1/2)+c^(1/4))/(1-I*c^(1/4)*x^(1/2)))/c^(1/4)/(d*x)^(1/2)-I*b^2*
x^(1/2)*polylog(2,1+2*c^(1/4)*(1-(-(-c)^(1/2))^(1/2)*x^(1/2))/(I*(-(-c)^(1
/2))^(1/2)-c^(1/4))/(1-I*c^(1/4)*x^(1/2)))/c^(1/4)/(d*x)^(1/2)-I*b^2*x^(1/
2)*polylog(2,1-2*(-c)^(1/4)*(1+c^(1/4)*x^(1/2))/((-c)^(1/4)+I*c^(1/4))/(1-
I*(-c)^(1/4)*x^(1/2))/(-c)^(1/4)/(d*x)^(1/2)-I*b^2*x^(1/2)*polylog(2,1-2*
(-c)^(1/4)*(1-c^(1/4)*x^(1/2))/((-c)^(1/4)-I*c^(1/4))/(1-I*(-c)^(1/4)*x^(1
/2)))/(-c)^(1/4)/(d*x)^(1/2)-I*b^2*x^(1/2)*polylog(2,1-2*c^(1/4)*(1+(-c)^(
1/4)*x^(1/2))/(I*(-c)^(1/4)+c^(1/4))/(1-I*c^(1/4)*x^(1/2)))/c^(1/4)/(d*x)^(
1/2)-I*b^2*x^(1/2)*polylog(2,1+2*c^(1/4)*(1-(-c)^(1/4)*x^(1/2))/(I*(-c)^(
1/4)-c^(1/4))/(1-I*c^(1/4)*x^(1/2)))/c^(1/4)/(d*x)^(1/2)-I*b^2*x^(1/2)*pol
ylog(2,1-2*(-c)^(1/4)*(1+(-c^(1/2))^(1/2)*x^(1/2))/(I*(-c^(1/2))^(1/2)+(-c
)^(1/4))/(1-I*(-c)^(1/4)*x^(1/2)))/(-c)^(1/4)/(d*x)^(1/2)-I*b^2*x^(1/2)...

```

Mathematica [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{\sqrt{dx}} dx = \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{\sqrt{dx}} dx$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])^2/Sqrt[d*x], x]
```

output

```
Integrate[(a + b*ArcTanh[c*x^2])^2/Sqrt[d*x], x]
```


Rubi [A] (verified)

Time = 11.69 (sec) , antiderivative size = 5216, normalized size of antiderivative = 0.85, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {6466, 6458, 6438, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{\sqrt{dx}} dx \\
 & \quad \downarrow \text{6466} \\
 & \frac{\sqrt{x} \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{\sqrt{x}} dx}{\sqrt{dx}} \\
 & \quad \downarrow \text{6458} \\
 & \frac{2\sqrt{x} \int (a + b \operatorname{arctanh}(cx^2))^2 d\sqrt{x}}{\sqrt{dx}} \\
 & \quad \downarrow \text{6438} \\
 & \frac{2\sqrt{x} \int (a^2 - b \log(1 - cx^2) a + b \log(cx^2 + 1) a + \frac{1}{4}b^2 \log^2(1 - cx^2) + \frac{1}{4}b^2 \log^2(cx^2 + 1) - \frac{1}{2}b^2 \log(1 - cx^2) \log(cx^2 + 1)) d\sqrt{x}}{\sqrt{dx}} \\
 & \quad \downarrow \text{2009} \\
 & 2\sqrt{x} \left(\sqrt{x} a^2 - \frac{\sqrt{2} b \arctan(1 - \sqrt{2} \sqrt[4]{c} \sqrt{x}) a}{\sqrt[4]{c}} + \frac{\sqrt{2} b \arctan(\sqrt{2} \sqrt[4]{c} \sqrt{x} + 1) a}{\sqrt[4]{c}} - \frac{2 b \arctan(\sqrt[4]{c} \sqrt{x}) a}{\sqrt[4]{c}} - \frac{2 b \operatorname{arctanh}(\sqrt[4]{c} \sqrt{x}) a}{\sqrt[4]{c}} - \frac{b \log(1 - cx^2) \log(cx^2 + 1)}{\sqrt{dx}} \right)
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c*x^2])^2/Sqrt[d*x], x]`

output

```
(2*Sqrt[x]*(a^2*Sqrt[x] - (Sqrt[2]*a*b*ArcTan[1 - Sqrt[2]*c^(1/4)*Sqrt[x]]
)/c^(1/4) + (Sqrt[2]*a*b*ArcTan[1 + Sqrt[2]*c^(1/4)*Sqrt[x]])/c^(1/4) + (I
*b^2*ArcTan[(-c)^(1/4)*Sqrt[x]]^2)/(-c)^(1/4) - (2*a*b*ArcTan[c^(1/4)*Sqrt
[x]])/c^(1/4) + (I*b^2*ArcTan[c^(1/4)*Sqrt[x]]^2)/c^(1/4) - (b^2*ArcTanh[(
-c)^(1/4)*Sqrt[x]]^2)/(-c)^(1/4) - (2*a*b*ArcTanh[c^(1/4)*Sqrt[x]])/c^(1/4
) - (b^2*ArcTanh[c^(1/4)*Sqrt[x]]^2)/c^(1/4) + (2*b^2*ArcTanh[(-c)^(1/4)*S
qrt[x]]*Log[2/(1 - (-c)^(1/4)*Sqrt[x])]/(-c)^(1/4) - (2*b^2*ArcTan[(-c)^(
1/4)*Sqrt[x]]*Log[2/(1 - I*(-c)^(1/4)*Sqrt[x])]/(-c)^(1/4) + (b^2*ArcTan[
(-c)^(1/4)*Sqrt[x]]*Log[(-2*(-c)^(1/4)*(1 - Sqrt[-Sqrt[c]]*Sqrt[x])]/((I*S
qrt[-Sqrt[c]] - (-c)^(1/4))*(1 - I*(-c)^(1/4)*Sqrt[x]))]/(-c)^(1/4) + (b^
2*ArcTan[(-c)^(1/4)*Sqrt[x]]*Log[(2*(-c)^(1/4)*(1 + Sqrt[-Sqrt[c]]*Sqrt[x]
))]/((I*Sqrt[-Sqrt[c]] + (-c)^(1/4))*(1 - I*(-c)^(1/4)*Sqrt[x]))]/(-c)^(1/
4) - (b^2*ArcTan[(-c)^(1/4)*Sqrt[x]]*Log[((1 + I)*(1 - (-c)^(1/4)*Sqrt[x])
)/(1 - I*(-c)^(1/4)*Sqrt[x])]/(-c)^(1/4) + (2*b^2*ArcTan[(-c)^(1/4)*Sqrt[
x]]*Log[2/(1 + I*(-c)^(1/4)*Sqrt[x])]/(-c)^(1/4) - (2*b^2*ArcTanh[(-c)^(1
/4)*Sqrt[x]]*Log[2/(1 + (-c)^(1/4)*Sqrt[x])]/(-c)^(1/4) - (b^2*ArcTanh[(
-c)^(1/4)*Sqrt[x]]*Log[(-2*(-c)^(1/4)*(1 - Sqrt[-Sqrt[-c]]*Sqrt[x])]/((Sqrt
[-Sqrt[-c]] - (-c)^(1/4))*(1 + (-c)^(1/4)*Sqrt[x]))]/(-c)^(1/4) - (b^2*Ar
cTanh[(-c)^(1/4)*Sqrt[x]]*Log[(2*(-c)^(1/4)*(1 + Sqrt[-Sqrt[-c]]*Sqrt[x])
]/((Sqrt[-Sqrt[-c]] + (-c)^(1/4))*(1 + (-c)^(1/4)*Sqrt[x]))]/(-c)^(1/4)...
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6438 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_), x_Symbol] := Int[ExpandI
ntegrand[(a + b*(Log[1 + c*x^n]/2) - b*(Log[1 - c*x^n]/2))^p, x], x] /; Fre
eQ[{a, b, c}, x] && IGtQ[p, 1] && IGtQ[n, 0]`

rule 6458 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_)*(x_)^(m_.), x_Symbol] :=
With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a + b*ArcT
anh[c*x^(k*n)])^p, x], x, x^(1/k)], x]] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1
] && IGtQ[n, 0] && FractionQ[m]`

rule 6466

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((d_.)*(x_)^(m_), x_Sym
bol] :> Simp[d^IntPart[m]*((d*x)^FracPart[m]/x^FracPart[m]) Int[x^m*(a +
b*ArcTanh[c*x^n])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0] &
& (EqQ[p, 1] || RationalQ[m, n])
```

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{\sqrt{dx}} dx$$

input

```
int((a+b*arctanh(c*x^2))^2/(d*x)^(1/2), x)
```

output

```
int((a+b*arctanh(c*x^2))^2/(d*x)^(1/2), x)
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{\sqrt{dx}} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^2}{\sqrt{dx}} dx$$

input

```
integrate((a+b*arctanh(c*x^2))^2/(d*x)^(1/2), x, algorithm="fricas")
```

output

```
integral((b^2*arctanh(c*x^2)^2 + 2*a*b*arctanh(c*x^2) + a^2)*sqrt(d*x)/(d*
x), x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{\sqrt{dx}} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{\sqrt{dx}} dx$$

input

```
integrate((a+b*atanh(c*x**2))**2/(d*x)**(1/2), x)
```

output `Integral((a + b*atanh(c*x**2))**2/sqrt(d*x), x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{\sqrt{dx}} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^2}{\sqrt{dx}} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/(d*x)^(1/2),x, algorithm="maxima")`

output `-1/2*a^2*c*((-I*(log(I*c^(1/4)*sqrt(x) + 1) - log(-I*c^(1/4)*sqrt(x) + 1))/c^(1/4) - log((sqrt(c)*sqrt(x) - c^(1/4))/(sqrt(c)*sqrt(x) + c^(1/4)))/c^(1/4))/(c*sqrt(d)) - 4*sqrt(x)/(c*sqrt(d)) + b^2*c*integrate(1/4*x^(3/2)*log(c*x^2 + 1)^2/(c*sqrt(d)*x^2 - sqrt(d)), x) - 2*b^2*c*integrate(1/4*x^(3/2)*log(c*x^2 + 1)*log(-c*x^2 + 1)/(c*sqrt(d)*x^2 - sqrt(d)), x) + 4*a*b*c*integrate(1/4*x^(3/2)*log(c*x^2 + 1)/(c*sqrt(d)*x^2 - sqrt(d)), x) - 4*a*b*c*integrate(1/4*x^(3/2)*log(-c*x^2 + 1)/(c*sqrt(d)*x^2 - sqrt(d)), x) - 8*b^2*c*integrate(1/4*x^(3/2)*log(-c*x^2 + 1)/(c*sqrt(d)*x^2 - sqrt(d)), x) + 1/2*b^2*sqrt(x)*log(-c*x^2 + 1)^2/sqrt(d) - b^2*integrate(1/4*log(c*x^2 + 1)^2/((c*sqrt(d)*x^2 - sqrt(d))*sqrt(x)), x) + 2*b^2*integrate(1/4*log(c*x^2 + 1)*log(-c*x^2 + 1)/((c*sqrt(d)*x^2 - sqrt(d))*sqrt(x)), x) - 4*a*b*integrate(1/4*log(c*x^2 + 1)/((c*sqrt(d)*x^2 - sqrt(d))*sqrt(x)), x) + 4*a*b*integrate(1/4*log(-c*x^2 + 1)/((c*sqrt(d)*x^2 - sqrt(d))*sqrt(x)), x) + 1/2*a^2*(-I*(log(I*c^(1/4)*sqrt(x) + 1) - log(-I*c^(1/4)*sqrt(x) + 1))/c^(1/4) - log((sqrt(c)*sqrt(x) - c^(1/4))/(sqrt(c)*sqrt(x) + c^(1/4)))/c^(1/4))/sqrt(d)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{\sqrt{dx}} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^2}{\sqrt{dx}} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/(d*x)^(1/2),x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)^2/sqrt(d*x), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{\sqrt{dx}} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{\sqrt{dx}} dx$$

input `int((a + b*atanh(c*x^2))^2/(d*x)^(1/2), x)`

output `int((a + b*atanh(c*x^2))^2/(d*x)^(1/2), x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{\sqrt{dx}} dx$$

$$= \frac{-2c^{\frac{3}{4}}\sqrt{2} \operatorname{atan}\left(\frac{c^{\frac{1}{4}}\sqrt{2}-2\sqrt{x}\sqrt{c}}{c^{\frac{1}{4}}\sqrt{2}}\right) ab + 2c^{\frac{3}{4}}\sqrt{2} \operatorname{atan}\left(\frac{c^{\frac{1}{4}}\sqrt{2}+2\sqrt{x}\sqrt{c}}{c^{\frac{1}{4}}\sqrt{2}}\right) ab - 4c^{\frac{3}{4}} \operatorname{atan}\left(\frac{\sqrt{x}\sqrt{c}}{c^{\frac{1}{4}}}\right) ab + 2c^{\frac{3}{4}}\sqrt{2} \operatorname{atanh}(c)}$$

input `int((a+b*atanh(c*x^2))^2/(d*x)^(1/2), x)`

output `(- 2*c**(3/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*a*b + 2*c**(3/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*a*b - 4*c**(3/4)*atan((sqrt(x)*sqrt(c))/c**(1/4))*a*b + 2*c**(3/4)*sqrt(2)*atanh(c*x**2)*a*b + 4*sqrt(x)*atanh(c*x**2)*a*b*c + c**(3/4)*sqrt(2)*log(c**(1/4) + sqrt(x)*sqrt(c))*a*b + c**(3/4)*sqrt(2)*log(- c**(1/4) + sqrt(x)*sqrt(c))*a*b - 2*c**(3/4)*sqrt(2)*log(- sqrt(x)*c**(1/4)*sqrt(2) + sqrt(c)*x + 1)*a*b + c**(3/4)*sqrt(2)*log(sqrt(c)*x + 1)*a*b - 2*c**(3/4)*log(c**(1/4) + sqrt(x)*sqrt(c))*a*b + 2*c**(3/4)*log(- c**(1/4) + sqrt(x)*sqrt(c))*a*b + 2*sqrt(x)*a**2*c + int(atanh(c*x**2)**2/sqrt(x), x)*b**2*c)/(sqrt(d)*c)`

$$3.92 \quad \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{3/2}} dx$$

Optimal result	817
Mathematica [F]	818
Rubi [A] (verified)	819
Maple [F]	821
Fricas [F]	821
Sympy [F]	821
Maxima [F]	822
Giac [F]	822
Mupad [F(-1)]	823
Reduce [F]	823

Optimal result

Integrand size = 20, antiderivative size = 6281

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{3/2}} dx = \text{Too large to display}$$

output

```

-1/2*b^2*ln(c*x^2+1)^2/d/(d*x)^(1/2)+2*2^(1/2)*a*b*c^(1/4)*x^(1/2)*arctan(
-1+2^(1/2)*c^(1/4)*x^(1/2))/d/(d*x)^(1/2)-2*2^(1/2)*a*b*c^(1/4)*x^(1/2)*ar
ctanh(2^(1/2)*c^(1/4)*x^(1/2)/(1+c^(1/2)*x))/d/(d*x)^(1/2)+2*2^(1/2)*a*b*c
^(1/4)*x^(1/2)*arctan(1+2^(1/2)*c^(1/4)*x^(1/2))/d/(d*x)^(1/2)-2*b^2*(-c)^(
1/4)*x^(1/2)*polylog(2,1-2/(1+(-c)^(1/4)*x^(1/2)))/d/(d*x)^(1/2)-2*b^2*(-c)
^(1/4)*x^(1/2)*polylog(2,1-2/(1-(-c)^(1/4)*x^(1/2)))/d/(d*x)^(1/2)-2*b^2*
c^(1/4)*x^(1/2)*polylog(2,1-2/(1+c^(1/4)*x^(1/2)))/d/(d*x)^(1/2)-2*b^2*c^(
1/4)*x^(1/2)*polylog(2,1-2/(1-c^(1/4)*x^(1/2)))/d/(d*x)^(1/2)+2*b^2*(-c)^(
1/4)*x^(1/2)*arctanh((-c)^(1/4)*x^(1/2))^2/d/(d*x)^(1/2)+2*b^2*c^(1/4)*x^(
1/2)*arctanh(c^(1/4)*x^(1/2))^2/d/(d*x)^(1/2)-b^2*c^(1/4)*x^(1/2)*polylog
(2,1-2*c^(1/4)*(1+(-c^(1/2))^(1/2)*x^(1/2))/((-c^(1/2))^(1/2)+c^(1/4))/(1+
c^(1/4)*x^(1/2)))/d/(d*x)^(1/2)-b^2*c^(1/4)*x^(1/2)*polylog(2,1+2*c^(1/4)*
(1-(-c^(1/2))^(1/2)*x^(1/2))/((-c^(1/2))^(1/2)-c^(1/4))/(1+c^(1/4)*x^(1/2)
))/d/(d*x)^(1/2)+b^2*c^(1/4)*x^(1/2)*polylog(2,1-2*c^(1/4)*(1+(-c)^(1/2)
)^(1/2)*x^(1/2))/((-c)^(1/2))^(1/2)+c^(1/4))/(1+c^(1/4)*x^(1/2)))/d/(d*x)
^(1/2)+b^2*c^(1/4)*x^(1/2)*polylog(2,1+2*c^(1/4)*(1-(-c)^(1/2))^(1/2)*x
^(1/2))/((-c)^(1/2))^(1/2)-c^(1/4))/(1+c^(1/4)*x^(1/2)))/d/(d*x)^(1/2)+b
^2*(-c)^(1/4)*x^(1/2)*polylog(2,1-2*(-c)^(1/4)*(1+(-c^(1/2))^(1/2)*x^(1/2)
))/((-c^(1/2))^(1/2)+(-c)^(1/4))/(1+(-c)^(1/4)*x^(1/2)))/d/(d*x)^(1/2)+b^2*
(-c)^(1/4)*x^(1/2)*polylog(2,1+2*(-c)^(1/4)*(1-(-c^(1/2))^(1/2)*x^(1/2))...

```

Mathematica [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{3/2}} dx = \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{3/2}} dx$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])^2/(d*x)^(3/2), x]
```

output

```
Integrate[(a + b*ArcTanh[c*x^2])^2/(d*x)^(3/2), x]
```

Rubi [A] (verified)

Time = 11.44 (sec) , antiderivative size = 5171, normalized size of antiderivative = 0.82, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {6466, 6458, 6456, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{3/2}} dx$$

$$\downarrow 6466$$

$$\frac{\sqrt{x} \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^{3/2}} dx}{d\sqrt{dx}}$$

$$\downarrow 6458$$

$$\frac{2\sqrt{x} \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x} d\sqrt{x}}{d\sqrt{dx}}$$

$$\downarrow 6456$$

$$\frac{2\sqrt{x} \int \left(\frac{(2a - b \log(1 - cx^2))^2}{4x} + \frac{b^2 \log^2(cx^2 + 1)}{4x} - \frac{b(b \log(1 - cx^2) - 2a) \log(cx^2 + 1)}{2x} \right) d\sqrt{x}}{d\sqrt{dx}}$$

$$\downarrow 2009$$

$$2\sqrt{x} \left(i \sqrt[4]{-c} \arctan(\sqrt[4]{-c}\sqrt{x})^2 b^2 + i \sqrt[4]{c} \arctan(\sqrt[4]{c}\sqrt{x})^2 b^2 + \sqrt[4]{-c} \operatorname{arctanh}(\sqrt[4]{-c}\sqrt{x})^2 b^2 + \sqrt[4]{c} \operatorname{arctanh}(\sqrt[4]{c}\sqrt{x})^2 b^2 \right)$$

input

```
Int[(a + b*ArcTanh[c*x^2])^2/(d*x)^(3/2), x]
```


output

```
(2*Sqrt[x]*(-Sqrt[2]*a*b*c^(1/4)*ArcTan[1 - Sqrt[2]*c^(1/4)*Sqrt[x]]) + S
qrt[2]*a*b*c^(1/4)*ArcTan[1 + Sqrt[2]*c^(1/4)*Sqrt[x]] + I*b^2*(-c)^(1/4)*
ArcTan[(-c)^(1/4)*Sqrt[x]]^2 + I*b^2*c^(1/4)*ArcTan[c^(1/4)*Sqrt[x]]^2 + b
^2*(-c)^(1/4)*ArcTanh[(-c)^(1/4)*Sqrt[x]]^2 + b^2*c^(1/4)*ArcTanh[c^(1/4)*
Sqrt[x]]^2 - 2*b^2*(-c)^(1/4)*ArcTanh[(-c)^(1/4)*Sqrt[x]]*Log[2/(1 - (-c)^(
1/4)*Sqrt[x])] - 2*b^2*(-c)^(1/4)*ArcTan[(-c)^(1/4)*Sqrt[x]]*Log[2/(1 - I
*(-c)^(1/4)*Sqrt[x])] + b^2*(-c)^(1/4)*ArcTan[(-c)^(1/4)*Sqrt[x]]*Log[(-2*
(-c)^(1/4)*(1 - Sqrt[-Sqrt[c]]*Sqrt[x]))/((I*Sqrt[-Sqrt[c]] - (-c)^(1/4))*
(1 - I*(-c)^(1/4)*Sqrt[x]))] + b^2*(-c)^(1/4)*ArcTan[(-c)^(1/4)*Sqrt[x]]*L
og[(2*(-c)^(1/4)*(1 + Sqrt[-Sqrt[c]]*Sqrt[x]))/((I*Sqrt[-Sqrt[c]] + (-c)^(
1/4))*(1 - I*(-c)^(1/4)*Sqrt[x]))] - b^2*(-c)^(1/4)*ArcTan[(-c)^(1/4)*Sqrt
[x]]*Log[((1 + I)*(1 - (-c)^(1/4)*Sqrt[x]))/(1 - I*(-c)^(1/4)*Sqrt[x])] +
2*b^2*(-c)^(1/4)*ArcTan[(-c)^(1/4)*Sqrt[x]]*Log[2/(1 + I*(-c)^(1/4)*Sqrt[x
])] + 2*b^2*(-c)^(1/4)*ArcTanh[(-c)^(1/4)*Sqrt[x]]*Log[2/(1 + (-c)^(1/4)*S
qrt[x])] + b^2*(-c)^(1/4)*ArcTanh[(-c)^(1/4)*Sqrt[x]]*Log[(-2*(-c)^(1/4)*
(1 - Sqrt[-Sqrt[-c]]*Sqrt[x]))/((Sqrt[-Sqrt[-c]] - (-c)^(1/4))*(1 + (-c)^(1
/4)*Sqrt[x]))] + b^2*(-c)^(1/4)*ArcTanh[(-c)^(1/4)*Sqrt[x]]*Log[(2*(-c)^(1
/4)*(1 + Sqrt[-Sqrt[-c]]*Sqrt[x]))/((Sqrt[-Sqrt[-c]] + (-c)^(1/4))*(1 + (-
c)^(1/4)*Sqrt[x]))] - b^2*(-c)^(1/4)*ArcTanh[(-c)^(1/4)*Sqrt[x]]*Log[(-2*(
-c)^(1/4)*(1 - Sqrt[-Sqrt[c]]*Sqrt[x]))/((Sqrt[-Sqrt[c]] - (-c)^(1/4))*...
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6456

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.)^(p_)*(x_)^(m_.), x_Symbol] :=
Int[ExpandIntegrand[x^m*(a + b*(Log[1 + c*x^n]/2) - b*(Log[1 - c*x^n]/2))^p
, x], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1] && IGtQ[n, 0] && IntegerQ[m]
```

rule 6458

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.)^(p_)*(x_)^(m_.), x_Symbol] :=
With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a + b*ArcT
anh[c*x^(k*n)])^p, x], x, x^(1/k)], x]] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1
] && IGtQ[n, 0] && FractionQ[m]
```

rule 6466

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((d_.)*(x_)^(m_), x_Sym
bol] := Simp[d^IntPart[m]*((d*x)^FracPart[m]/x^FracPart[m]) Int[x^m*(a +
b*ArcTanh[c*x^n])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0] &
& (EqQ[p, 1] || RationalQ[m, n])
```

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{\frac{3}{2}}} dx$$

input

```
int((a+b*arctanh(c*x^2))^2/(d*x)^(3/2), x)
```

output

```
int((a+b*arctanh(c*x^2))^2/(d*x)^(3/2), x)
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{3/2}} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^2}{(dx)^{\frac{3}{2}}} dx$$

input

```
integrate((a+b*arctanh(c*x^2))^2/(d*x)^(3/2), x, algorithm="fricas")
```

output

```
integral((b^2*arctanh(c*x^2)^2 + 2*a*b*arctanh(c*x^2) + a^2)*sqrt(d*x)/(d^
2*x^2), x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{3/2}} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{(dx)^{\frac{3}{2}}} dx$$

input

```
integrate((a+b*atanh(c*x**2))**2/(d*x)**(3/2), x)
```

output `Integral((a + b*atanh(c*x**2))**2/(d*x)**(3/2), x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{3/2}} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^2}{(dx)^{3/2}} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/(d*x)^(3/2),x, algorithm="maxima")`

output `b^2*c*integrate(1/4*x^(3/2)*log(c*x^2 + 1)^2/(c*d^(3/2)*x^3 - d^(3/2)*x), x) - 2*b^2*c*integrate(1/4*x^(3/2)*log(c*x^2 + 1)*log(-c*x^2 + 1)/(c*d^(3/2)*x^3 - d^(3/2)*x), x) + 4*a*b*c*integrate(1/4*x^(3/2)*log(c*x^2 + 1)/(c*d^(3/2)*x^3 - d^(3/2)*x), x) - 4*a*b*c*integrate(1/4*x^(3/2)*log(-c*x^2 + 1)/(c*d^(3/2)*x^3 - d^(3/2)*x), x) + 8*b^2*c*integrate(1/4*x^(3/2)*log(-c*x^2 + 1)/(c*d^(3/2)*x^3 - d^(3/2)*x), x) + 1/2*a^2*(c*(I*(log(I*c^(1/4)*sqrt(x) + 1) - log(-I*c^(1/4)*sqrt(x) + 1))/c^(3/4) - log((sqrt(c)*sqrt(x) - c^(1/4))/(sqrt(c)*sqrt(x) + c^(1/4)))/c^(3/4))/d^(3/2) - 4/(d^(3/2)*sqrt(x))) - b^2*integrate(1/4*log(c*x^2 + 1)^2/((c*d^(3/2)*x^3 - d^(3/2)*x)*sqrt(x)), x) + 2*b^2*integrate(1/4*log(c*x^2 + 1)*log(-c*x^2 + 1)/((c*d^(3/2)*x^3 - d^(3/2)*x)*sqrt(x)), x) - 4*a*b*integrate(1/4*log(c*x^2 + 1)/((c*d^(3/2)*x^3 - d^(3/2)*x)*sqrt(x)), x) + 4*a*b*integrate(1/4*log(-c*x^2 + 1)/((c*d^(3/2)*x^3 - d^(3/2)*x)*sqrt(x)), x) - 1/2*a^2*c*(I*(log(I*c^(1/4)*sqrt(x) + 1) - log(-I*c^(1/4)*sqrt(x) + 1))/c^(3/4) - log((sqrt(c)*sqrt(x) - c^(1/4))/(sqrt(c)*sqrt(x) + c^(1/4)))/c^(3/4))/d^(3/2) - 1/2*b^2*log(-c*x^2 + 1)^2/(d^(3/2)*sqrt(x))`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{3/2}} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^2}{(dx)^{3/2}} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/(d*x)^(3/2),x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)^2/(d*x)^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{3/2}} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{(dx)^{3/2}} dx$$

input `int((a + b*atanh(c*x^2))^2/(d*x)^(3/2), x)`

output `int((a + b*atanh(c*x^2))^2/(d*x)^(3/2), x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{3/2}} dx = \frac{-2\sqrt{x} c^{1/4} \sqrt{2} \operatorname{atan}\left(\frac{c^{1/4} \sqrt{2} - 2\sqrt{x} \sqrt{c}}{c^{1/4} \sqrt{2}}\right) ab + 2\sqrt{x} c^{1/4} \sqrt{2} \operatorname{atan}\left(\frac{c^{1/4} \sqrt{2} + 2\sqrt{x} \sqrt{c}}{c^{1/4} \sqrt{2}}\right) ab - 4}{\dots}$$

input `int((a+b*atanh(c*x^2))^2/(d*x)^(3/2), x)`

output `(- 2*sqrt(x)*c**(1/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*a*b + 2*sqrt(x)*c**(1/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*a*b - 4*sqrt(x)*c**(1/4)*atan(sqrt(x)*sqrt(c)/c**(1/4))*a*b - 2*sqrt(x)*c**(1/4)*sqrt(2)*atanh(c*x**2)*a*b - 4*atanh(c*x**2)*a*b - sqrt(x)*c**(1/4)*sqrt(2)*log(c**(1/4) + sqrt(x)*sqrt(c))*a*b - sqrt(x)*c**(1/4)*sqrt(2)*log(-c**(1/4) + sqrt(x)*sqrt(c))*a*b + 2*sqrt(x)*c**(1/4)*sqrt(2)*log(-sqrt(x)*c**(1/4)*sqrt(2) + sqrt(c)*x + 1)*a*b - sqrt(x)*c**(1/4)*sqrt(2)*log(sqrt(c)*x + 1)*a*b + 2*sqrt(x)*c**(1/4)*log(c**(1/4) + sqrt(x)*sqrt(c))*a*b - 2*sqrt(x)*c**(1/4)*log(-c**(1/4) + sqrt(x)*sqrt(c))*a*b + sqrt(x)*int(atanh(c*x**2)**2/(sqrt(x)*x), x)*b**2 - 2*a**2)/(sqrt(x)*sqrt(d)*d)`

$$3.93 \quad \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{5/2}} dx$$

Optimal result	824
Mathematica [F]	825
Rubi [A] (warning: unable to verify)	826
Maple [F]	828
Fricas [F]	828
Sympy [F]	828
Maxima [F]	829
Giac [F]	829
Mupad [F(-1)]	830
Reduce [F]	830

Optimal result

Integrand size = 20, antiderivative size = 6464

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{5/2}} dx = \text{Too large to display}$$

output

```

-1/6*(2*a-b*ln(-c*x^2+1))^2/d^2/x/(d*x)^(1/2)+2/3*2^(1/2)*a*b*c^(3/4)*x^(1/2)*arctan(-1+2^(1/2)*c^(1/4)*x^(1/2))/d^2/(d*x)^(1/2)+2/3*2^(1/2)*a*b*c^(3/4)*x^(1/2)*arctanh(2^(1/2)*c^(1/4)*x^(1/2)/(1+c^(1/2)*x))/d^2/(d*x)^(1/2)+2/3*2^(1/2)*a*b*c^(3/4)*x^(1/2)*arctan(1+2^(1/2)*c^(1/4)*x^(1/2))/d^2/(d*x)^(1/2)+1/3*b^2*(-c)^(3/4)*x^(1/2)*polylog(2,1+2*(-c)^(1/4)*(1-(-c)^(1/2))^(1/2)*x^(1/2))/((-c)^(1/2))^(1/2)-(-c)^(1/4))/(1+(-c)^(1/4)*x^(1/2)))/d^2/(d*x)^(1/2)+1/3*b^2*c^(3/4)*x^(1/2)*polylog(2,1-2*c^(1/4)*(1+(-c)^(1/2))^(1/2)*x^(1/2))/((-(-c)^(1/2))^(1/2)+c^(1/4))/(1+c^(1/4)*x^(1/2)))/d^2/(d*x)^(1/2)+1/3*b^2*c^(3/4)*x^(1/2)*polylog(2,1+2*c^(1/4)*(1-(-c)^(1/2))^(1/2)*x^(1/2))/((-(-c)^(1/2))^(1/2)-c^(1/4))/(1+c^(1/4)*x^(1/2)))/d^2/(d*x)^(1/2)+2/3*b^2*(-c)^(3/4)*x^(1/2)*arctanh((-c)^(1/4)*x^(1/2))^2/d^2/(d*x)^(1/2)-2/3*b^2*c^(3/4)*x^(1/2)*polylog(2,1-2/(1+c^(1/4)*x^(1/2)))/d^2/(d*x)^(1/2)-2/3*b^2*c^(3/4)*x^(1/2)*polylog(2,1-2/(1-c^(1/4)*x^(1/2)))/d^2/(d*x)^(1/2)+1/3*b^2*(-c)^(3/4)*x^(1/2)*polylog(2,1-2*(-c)^(1/4)*(1+c^(1/4)*x^(1/2)))/((-c)^(1/4)+c^(1/4))/(1+(-c)^(1/4)*x^(1/2)))/d^2/(d*x)^(1/2)+1/3*b^2*(-c)^(3/4)*x^(1/2)*polylog(2,1-2*(-c)^(1/4)*(1-c^(1/4)*x^(1/2)))/((-c)^(1/4)-c^(1/4))/(1+(-c)^(1/4)*x^(1/2)))/d^2/(d*x)^(1/2)-2/3*b^2*(-c)^(3/4)*x^(1/2)*arctan((-c)^(1/4)*x^(1/2))*ln(2*(-c)^(1/4)*(1+c^(1/4)*x^(1/2)))/((-c)^(1/4)+I*c^(1/4))/(1-I*(-c)^(1/4)*x^(1/2)))/d^2/(d*x)^(1/2)-2/3*b^2*(-c)^(3/4)*x^(1/2)*arctan((-c)^(1/4)*x^(1/2))*ln(2*(-c)^(1/4)*(1-c^(1/4)*x^(1/2)))/((-c)^(1/4)-I*c^(1/4))/(1-I*(-c)^(1/4)*x^(1/2)))/d^2/(d*x)^(1/2)

```

Mathematica [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{5/2}} dx = \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{5/2}} dx$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])^2/(d*x)^(5/2), x]
```

output

```
Integrate[(a + b*ArcTanh[c*x^2])^2/(d*x)^(5/2), x]
```

Rubi [A] (warning: unable to verify)

Time = 10.21 (sec) , antiderivative size = 5305, normalized size of antiderivative = 0.82, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {6466, 6458, 6456, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{5/2}} dx \\
 & \quad \downarrow \text{6466} \\
 & \frac{\sqrt{x} \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^{5/2}} dx}{d^2 \sqrt{dx}} \\
 & \quad \downarrow \text{6458} \\
 & \frac{2\sqrt{x} \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{x^2} d\sqrt{x}}{d^2 \sqrt{dx}} \\
 & \quad \downarrow \text{6456} \\
 & \frac{2\sqrt{x} \int \left(\frac{(2a - b \log(1 - cx^2))^2}{4x^2} + \frac{b^2 \log^2(cx^2 + 1)}{4x^2} - \frac{b(b \log(1 - cx^2) - 2a) \log(cx^2 + 1)}{2x^2} \right) d\sqrt{x}}{d^2 \sqrt{dx}} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$2\sqrt{x} \left(-\frac{1}{3}i(-c)^{3/4} \arctan(\sqrt[4]{-c}\sqrt{x})^2 b^2 - \frac{1}{3}ic^{3/4} \arctan(\sqrt[4]{c}\sqrt{x})^2 b^2 + \frac{1}{3}(-c)^{3/4} \operatorname{arctanh}(\sqrt[4]{-c}\sqrt{x})^2 b^2 + \frac{1}{3}c^{3/4} \operatorname{arctanh}(\sqrt[4]{c}\sqrt{x})^2 b^2 \right)$$

input `Int[(a + b*ArcTanh[c*x^2])^2/(d*x)^(5/2), x]`

output

$$\begin{aligned}
& (2\sqrt{x}*(-1/3*(\sqrt{2}*a*b*c^{3/4}*\text{ArcTan}[1 - \sqrt{2}*c^{1/4}*\sqrt{x}]) \\
& + (\sqrt{2}*a*b*c^{3/4}*\text{ArcTan}[1 + \sqrt{2}*c^{1/4}*\sqrt{x}])/3 - (I/3)*b^2 \\
& *(-c)^{3/4}*\text{ArcTan}[(-c)^{1/4}*\sqrt{x}]^2 - (I/3)*b^2*c^{3/4}*\text{ArcTan}[c^{1/4} \\
&)*\sqrt{x}]^2 + (b^2*(-c)^{3/4}*\text{ArcTanh}[(-c)^{1/4}*\sqrt{x}]^2)/3 + (b^2*c^{3/4} \\
&)*\text{ArcTanh}[c^{1/4}*\sqrt{x}]^2)/3 - (2*b^2*(-c)^{3/4}*\text{ArcTanh}[(-c)^{1/4} \\
&)*\sqrt{x}]*\text{Log}[2/(1 - (-c)^{1/4}*\sqrt{x})]/3 + (2*b^2*(-c)^{3/4}*\text{ArcTan}[(-c) \\
&)^{1/4}*\sqrt{x}]*\text{Log}[2/(1 - I*(-c)^{1/4}*\sqrt{x})]/3 - (b^2*(-c)^{3/4}*\text{Ar} \\
&)*\text{cTan}[(-c)^{1/4}*\sqrt{x}]*\text{Log}[(-2*(-c)^{1/4}*(1 - \sqrt{-\text{Sqrt}[c]}*\sqrt{x})) / \\
& ((I*\sqrt{-\text{Sqrt}[c]} - (-c)^{1/4})*(1 - I*(-c)^{1/4}*\sqrt{x}))]/3 - (b^2*(-c) \\
&)^{3/4}*\text{ArcTan}[(-c)^{1/4}*\sqrt{x}]*\text{Log}[(2*(-c)^{1/4}*(1 + \sqrt{-\text{Sqrt}[c]}* \\
&)*\sqrt{x})]/((I*\sqrt{-\text{Sqrt}[c]} + (-c)^{1/4})*(1 - I*(-c)^{1/4}*\sqrt{x}))/3 \\
& + (b^2*(-c)^{3/4}*\text{ArcTan}[(-c)^{1/4}*\sqrt{x}]*\text{Log}[(1 + I)*(1 - (-c)^{1/4} \\
&)*\sqrt{x}])/(1 - I*(-c)^{1/4}*\sqrt{x}))/3 - (2*b^2*(-c)^{3/4}*\text{ArcTan}[(-c) \\
&)^{1/4}*\sqrt{x}]*\text{Log}[2/(1 + I*(-c)^{1/4}*\sqrt{x})]/3 + (2*b^2*(-c)^{3/4}*\text{Ar} \\
&)*\text{cTanh}[(-c)^{1/4}*\sqrt{x}]*\text{Log}[2/(1 + (-c)^{1/4}*\sqrt{x})]/3 + (b^2*(-c)^{3/4} \\
&)*\text{ArcTanh}[(-c)^{1/4}*\sqrt{x}]*\text{Log}[(-2*(-c)^{1/4}*(1 - \sqrt{-\text{Sqrt}[-c]}* \\
&)*\sqrt{x})]/((\sqrt{-\text{Sqrt}[-c]} - (-c)^{1/4})*(1 + (-c)^{1/4}*\sqrt{x}))/3 + (\\
&)^{3/4}*\text{ArcTanh}[(-c)^{1/4}*\sqrt{x}]*\text{Log}[(2*(-c)^{1/4}*(1 + \sqrt{-\text{Sqrt}[-c]}* \\
&)*\sqrt{x})]/((\sqrt{-\text{Sqrt}[-c]} + (-c)^{1/4})*(1 + (-c)^{1/4}*\sqrt{x}))) \\
&)]/3 - (b^2*(-c)^{3/4}*\text{ArcTanh}[(-c)^{1/4}*\sqrt{x}]*\text{Log}[(-2*(-c)^{1/4})*\dots
\end{aligned}$$

Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \text{ :> } \text{Simp}[\text{IntSum}[u, x], x] \text{ /; } \text{SumQ}[u]$$

rule 6456

$$\begin{aligned}
& \text{Int}[(a_.) + \text{ArcTanh}[(c_.)*(x_)^{(n_)}]*(b_.))^{(p_)}*(x_)^{(m_.)}, x_Symbol] \text{ :>} \\
& \text{Int}[\text{ExpandIntegrand}[x^m*(a + b*(\text{Log}[1 + c*x^n]/2) - b*(\text{Log}[1 - c*x^n]/2))]^p \\
& , x], x] \text{ /; } \text{FreeQ}\{a, b, c\}, x \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]
\end{aligned}$$

rule 6458

$$\begin{aligned}
& \text{Int}[(a_.) + \text{ArcTanh}[(c_.)*(x_)^{(n_)}]*(b_.))^{(p_)}*(x_)^{(m_.)}, x_Symbol] \text{ :>} \\
& \text{With}\{k = \text{Denominator}[m]\}, \text{Simp}[k \ \text{Subst}[\text{Int}[x^{(k*(m + 1) - 1)}*(a + b*\text{ArcT} \\
&)^{(k*n)}], x], x, x^{(1/k)}], x]] \text{ /; } \text{FreeQ}\{a, b, c\}, x \ \&\& \ \text{IGtQ}[p, 1] \\
& \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{FractionQ}[m]
\end{aligned}$$

rule 6466

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((d_.)*(x_)^(m_), x_Sym
bol] := Simp[d^IntPart[m]*((d*x)^FracPart[m]/x^FracPart[m]) Int[x^m*(a +
b*ArcTanh[c*x^n])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0] &
& (EqQ[p, 1] || RationalQ[m, n])
```

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{\frac{5}{2}}} dx$$

input

```
int((a+b*arctanh(c*x^2))^2/(d*x)^(5/2), x)
```

output

```
int((a+b*arctanh(c*x^2))^2/(d*x)^(5/2), x)
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{5/2}} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^2}{(dx)^{\frac{5}{2}}} dx$$

input

```
integrate((a+b*arctanh(c*x^2))^2/(d*x)^(5/2), x, algorithm="fricas")
```

output

```
integral((b^2*arctanh(c*x^2)^2 + 2*a*b*arctanh(c*x^2) + a^2)*sqrt(d*x)/(d^
3*x^3), x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{5/2}} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{(dx)^{\frac{5}{2}}} dx$$

input

```
integrate((a+b*atanh(c*x**2))**2/(d*x)**(5/2), x)
```

output `Integral((a + b*atanh(c*x**2))**2/(d*x)**(5/2), x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{5/2}} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^2}{(dx)^{5/2}} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/(d*x)^(5/2),x, algorithm="maxima")`

output `3*b^2*c*integrate(1/12*x^(3/2)*log(c*x^2 + 1)^2/(c*d^(5/2)*x^4 - d^(5/2)*x^2), x) - 6*b^2*c*integrate(1/12*x^(3/2)*log(c*x^2 + 1)*log(-c*x^2 + 1)/(c*d^(5/2)*x^4 - d^(5/2)*x^2), x) + 12*a*b*c*integrate(1/12*x^(3/2)*log(c*x^2 + 1)/(c*d^(5/2)*x^4 - d^(5/2)*x^2), x) - 12*a*b*c*integrate(1/12*x^(3/2)*log(-c*x^2 + 1)/(c*d^(5/2)*x^4 - d^(5/2)*x^2), x) + 8*b^2*c*integrate(1/12*x^(3/2)*log(-c*x^2 + 1)/(c*d^(5/2)*x^4 - d^(5/2)*x^2), x) + 1/6*a^2*(3*(-I*c^(3/4)*(log(I*c^(1/4)*sqrt(x) + 1) - log(-I*c^(1/4)*sqrt(x) + 1)) - c^(3/4)*log((sqrt(c)*sqrt(x) - c^(1/4))/(sqrt(c)*sqrt(x) + c^(1/4))))/d^(5/2) - 4/(d^(5/2)*x^(3/2))) - 3*b^2*integrate(1/12*log(c*x^2 + 1)^2/((c*d^(5/2)*x^4 - d^(5/2)*x^2)*sqrt(x)), x) + 6*b^2*integrate(1/12*log(c*x^2 + 1)*log(-c*x^2 + 1)/((c*d^(5/2)*x^4 - d^(5/2)*x^2)*sqrt(x)), x) - 12*a*b*integrate(1/12*log(c*x^2 + 1)/((c*d^(5/2)*x^4 - d^(5/2)*x^2)*sqrt(x)), x) + 12*a*b*integrate(1/12*log(-c*x^2 + 1)/((c*d^(5/2)*x^4 - d^(5/2)*x^2)*sqrt(x)), x) - 1/2*a^2*c*(-I*(log(I*c^(1/4)*sqrt(x) + 1) - log(-I*c^(1/4)*sqrt(x) + 1))/c^(1/4) - log((sqrt(c)*sqrt(x) - c^(1/4))/(sqrt(c)*sqrt(x) + c^(1/4)))/c^(1/4))/d^(5/2) - 1/6*b^2*log(-c*x^2 + 1)^2/(d^(5/2)*x^(3/2))`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{5/2}} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^2}{(dx)^{5/2}} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/(d*x)^(5/2),x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)^2/(d*x)^(5/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{5/2}} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{(dx)^{5/2}} dx$$

input `int((a + b*atanh(c*x^2))^2/(d*x)^(5/2), x)`

output `int((a + b*atanh(c*x^2))^2/(d*x)^(5/2), x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(dx)^{5/2}} dx = \frac{-2\sqrt{x} c^{3/4} \sqrt{2} \operatorname{atan}\left(\frac{c^{1/4} \sqrt{2} - 2\sqrt{x} \sqrt{c}}{c^{1/4} \sqrt{2}}\right) abx + 2\sqrt{x} c^{3/4} \sqrt{2} \operatorname{atan}\left(\frac{c^{1/4} \sqrt{2} + 2\sqrt{x} \sqrt{c}}{c^{1/4} \sqrt{2}}\right) abx - \dots}{\dots}$$

input `int((a+b*atanh(c*x^2))^2/(d*x)^(5/2), x)`

output `(- 2*sqrt(x)*c**(3/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*a*b*x + 2*sqrt(x)*c**(3/4)*sqrt(2)*atan((c**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(c))/(c**(1/4)*sqrt(2)))*a*b*x + 4*sqrt(x)*c**(3/4)*atan((sqrt(x)*sqrt(c))/c**(1/4))*a*b*x + 2*sqrt(x)*c**(3/4)*sqrt(2)*atanh(c*x**2)*a*b*x - 4*atanh(c*x**2)*a*b + sqrt(x)*c**(3/4)*sqrt(2)*log(c**(1/4) + sqrt(x)*sqrt(c))*a*b*x + sqrt(x)*c**(3/4)*sqrt(2)*log(- c**(1/4) + sqrt(x)*sqrt(c))*a*b*x - 2*sqrt(x)*c**(3/4)*sqrt(2)*log(- sqrt(x)*c**(1/4)*sqrt(2) + sqrt(c)*x + 1)*a*b*x + sqrt(x)*c**(3/4)*sqrt(2)*log(sqrt(c)*x + 1)*a*b*x + 2*sqrt(x)*c**(3/4)*log(c**(1/4) + sqrt(x)*sqrt(c))*a*b*x - 2*sqrt(x)*c**(3/4)*log(- c**(1/4) + sqrt(x)*sqrt(c))*a*b*x + 3*sqrt(x)*int(atanh(c*x**2)**2/(sqrt(x)*x**2), x)*b**2*x - 2*a**2)/(3*sqrt(x)*sqrt(d)*d**2*x)`

3.94 $\int (dx)^m (a + \operatorname{barctanh}(cx^2))^3 dx$

Optimal result	831
Mathematica [N/A]	831
Rubi [N/A]	832
Maple [N/A]	832
Fricas [N/A]	833
Sympy [N/A]	833
Maxima [N/A]	833
Giac [N/A]	834
Mupad [N/A]	834
Reduce [N/A]	835

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int (dx)^m (a + \operatorname{barctanh}(cx^2))^3 dx = \operatorname{Int}\left((dx)^m (a + \operatorname{barctanh}(cx^2))^3, x\right)$$

output `Defer(Int)((d*x)^m*(a+b*arctanh(c*x^2))^3,x)`

Mathematica [N/A]

Not integrable

Time = 1.48 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + \operatorname{barctanh}(cx^2))^3 dx = \int (dx)^m (a + \operatorname{barctanh}(cx^2))^3 dx$$

input `Integrate[(d*x)^m*(a + b*ArcTanh[c*x^2])^3,x]`

output `Integrate[(d*x)^m*(a + b*ArcTanh[c*x^2])^3, x]`

Rubi [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^3 dx$$

↓ 6468

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^3 dx$$

input `Int[(d*x)^m*(a + b*ArcTanh[c*x^2])^3,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^3 dx$$

input `int((d*x)^m*(a+b*arctanh(c*x^2))^3,x)`

output `int((d*x)^m*(a+b*arctanh(c*x^2))^3,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 50, normalized size of antiderivative = 2.78

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^3 dx = \int (b \operatorname{artanh}(cx^2) + a)^3 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^2))^3,x, algorithm="fricas")`

output `integral((b^3*arctanh(c*x^2)^3 + 3*a*b^2*arctanh(c*x^2)^2 + 3*a^2*b*arctanh(c*x^2) + a^3)*(d*x)^m, x)`

Sympy [N/A]

Not integrable

Time = 52.96 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^3 dx = \int (dx)^m (a + b \operatorname{atanh}(cx^2))^3 dx$$

input `integrate((d*x)**m*(a+b*atanh(c*x**2))**3,x)`

output `Integral((d*x)**m*(a + b*atanh(c*x**2))**3, x)`

Maxima [N/A]

Not integrable

Time = 2.48 (sec) , antiderivative size = 413, normalized size of antiderivative = 22.94

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^3 dx = \int (b \operatorname{artanh}(cx^2) + a)^3 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^2))^3,x, algorithm="maxima")`

output

```
-1/8*b^3*d^m*x*x^m*log(-c*x^2 + 1)^3/(m + 1) + (d*x)^(m + 1)*a^3/(d*(m + 1)) + integrate(1/8*((b^3*c*d^m*(m + 1)*x^2 - b^3*d^m*(m + 1))*x^m*log(c*x^2 + 1)^3 + 6*(a*b^2*c*d^m*(m + 1)*x^2 - a*b^2*d^m*(m + 1))*x^m*log(c*x^2 + 1)^2 + 12*(a^2*b*c*d^m*(m + 1)*x^2 - a^2*b*d^m*(m + 1))*x^m*log(c*x^2 + 1) + 3*((b^3*c*d^m*(m + 1)*x^2 - b^3*d^m*(m + 1))*x^m*log(c*x^2 + 1) - 2*(a*b^2*d^m*(m + 1) - (a*b^2*c*d^m*(m + 1) + b^3*c*d^m)*x^2)*x^m*log(-c*x^2 + 1)^2 - 3*((b^3*c*d^m*(m + 1)*x^2 - b^3*d^m*(m + 1))*x^m*log(c*x^2 + 1)^2 + 4*(a*b^2*c*d^m*(m + 1)*x^2 - a*b^2*d^m*(m + 1))*x^m*log(c*x^2 + 1) + 4*(a^2*b*c*d^m*(m + 1)*x^2 - a^2*b*d^m*(m + 1))*x^m*log(-c*x^2 + 1))/(c*(m + 1)*x^2 - m - 1), x)
```

Giac [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^3 dx = \int (b \operatorname{artanh}(cx^2) + a)^3 (dx)^m dx$$

input

```
integrate((d*x)^m*(a+b*arctanh(c*x^2))^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^2) + a)^3*(d*x)^m, x)
```

Mupad [N/A]

Not integrable

Time = 3.29 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^3 dx = \int (dx)^m (a + b \operatorname{atanh}(cx^2))^3 dx$$

input

```
int((d*x)^m*(a + b*atanh(c*x^2))^3,x)
```

output

```
int((d*x)^m*(a + b*atanh(c*x^2))^3, x)
```

Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 133, normalized size of antiderivative = 7.39

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^3 dx$$

$$= \frac{d^m \left(x^m a^3 x + 3 \left(\int x^m \operatorname{atanh}(cx^2) dx \right) a^2 b m + 3 \left(\int x^m \operatorname{atanh}(cx^2) dx \right) a^2 b + \left(\int x^m \operatorname{atanh}(cx^2) dx \right)^3 b^3 m + \dots \right)}{m + 1}$$

input

```
int((d*x)^m*(a+b*atanh(c*x^2))^3,x)
```

output

```
(d**m*(x**m*a**3*x + 3*int(x**m*atanh(c*x**2),x)*a**2*b*m + 3*int(x**m*atanh(c*x**2),x)*a**2*b + int(x**m*atanh(c*x**2)**3,x)*b**3*m + int(x**m*atanh(c*x**2)**3,x)*b**3 + 3*int(x**m*atanh(c*x**2)**2,x)*a*b**2*m + 3*int(x**m*atanh(c*x**2)**2,x)*a*b**2))/(m + 1)
```


3.95 $\int (dx)^m (a + \operatorname{barctanh}(cx^2))^2 dx$

Optimal result	836
Mathematica [N/A]	836
Rubi [N/A]	837
Maple [N/A]	837
Fricas [N/A]	838
Sympy [N/A]	838
Maxima [N/A]	838
Giac [N/A]	839
Mupad [N/A]	839
Reduce [N/A]	840

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int (dx)^m (a + \operatorname{barctanh}(cx^2))^2 dx = \operatorname{Int}\left((dx)^m (a + \operatorname{barctanh}(cx^2))^2, x\right)$$

output `Defer(Int)((d*x)^m*(a+b*arctanh(c*x^2))^2,x)`

Mathematica [N/A]

Not integrable

Time = 0.94 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + \operatorname{barctanh}(cx^2))^2 dx = \int (dx)^m (a + \operatorname{barctanh}(cx^2))^2 dx$$

input `Integrate[(d*x)^m*(a + b*ArcTanh[c*x^2])^2,x]`

output `Integrate[(d*x)^m*(a + b*ArcTanh[c*x^2])^2, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^2 dx$$

↓ 6468

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^2 dx$$

input `Int[(d*x)^m*(a + b*ArcTanh[c*x^2])^2,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^2 dx$$

input `int((d*x)^m*(a+b*arctanh(c*x^2))^2,x)`

output `int((d*x)^m*(a+b*arctanh(c*x^2))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.89

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^2))^2,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x^2)^2 + 2*a*b*arctanh(c*x^2) + a^2)*(d*x)^m, x)`

Sympy [N/A]

Not integrable

Time = 38.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (dx)^m (a + b \operatorname{atanh}(cx^2))^2 dx$$

input `integrate((d*x)**m*(a+b*atanh(c*x**2))**2,x)`

output `Integral((d*x)**m*(a + b*atanh(c*x**2))**2, x)`

Maxima [N/A]

Not integrable

Time = 1.30 (sec) , antiderivative size = 236, normalized size of antiderivative = 13.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^2))^2,x, algorithm="maxima")`

output

```
1/4*b^2*d^m*x*x^m*log(-c*x^2 + 1)^2/(m + 1) + (d*x)^(m + 1)*a^2/(d*(m + 1)
) - integrate(-1/4*((b^2*c*d^m*(m + 1)*x^2 - b^2*d^m*(m + 1))*x^m*log(c*x^
2 + 1)^2 + 4*(a*b*c*d^m*(m + 1)*x^2 - a*b*d^m*(m + 1))*x^m*log(c*x^2 + 1)
- 2*((b^2*c*d^m*(m + 1)*x^2 - b^2*d^m*(m + 1))*x^m*log(c*x^2 + 1) - 2*(a*b
*d^m*(m + 1) - (a*b*c*d^m*(m + 1) + b^2*c*d^m)*x^2)*x^m*log(-c*x^2 + 1))/
(c*(m + 1)*x^2 - m - 1), x)
```

Giac [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (b \operatorname{artanh}(cx^2) + a)^2 (dx)^m dx$$

input

```
integrate((d*x)^m*(a+b*arctanh(c*x^2))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^2) + a)^2*(d*x)^m, x)
```

Mupad [N/A]

Not integrable

Time = 3.34 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (dx)^m (a + b \operatorname{atanh}(cx^2))^2 dx$$

input

```
int((d*x)^m*(a + b*atanh(c*x^2))^2,x)
```

output

```
int((d*x)^m*(a + b*atanh(c*x^2))^2, x)
```

Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 88, normalized size of antiderivative = 4.89

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$= \frac{d^m \left(x^m a^2 x + 2 \left(\int x^m \operatorname{atanh}(cx^2) dx \right) abm + 2 \left(\int x^m \operatorname{atanh}(cx^2) dx \right) ab + \left(\int x^m \operatorname{atanh}(cx^2)^2 dx \right) b^2 m + \dots \right)}{m + 1}$$

input

```
int((d*x)^m*(a+b*atanh(c*x^2))^2,x)
```

output

```
(d**m*(x**m*a**2*x + 2*int(x**m*atanh(c*x**2),x)*a*b*m + 2*int(x**m*atanh(c*x**2),x)*a*b + int(x**m*atanh(c*x**2)**2,x)*b**2*m + int(x**m*atanh(c*x**2)**2,x)*b**2))/(m + 1)
```

3.96 $\int (dx)^m (a + \operatorname{barctanh}(cx^2)) dx$

Optimal result	841
Mathematica [A] (verified)	841
Rubi [A] (verified)	842
Maple [F]	843
Fricas [F]	843
Sympy [F]	844
Maxima [F]	844
Giac [F]	844
Mupad [F(-1)]	845
Reduce [F]	845

Optimal result

Integrand size = 16, antiderivative size = 74

$$\int (dx)^m (a + \operatorname{barctanh}(cx^2)) dx = \frac{(dx)^{1+m} (a + \operatorname{barctanh}(cx^2))}{d(1+m)} - \frac{2bc(dx)^{3+m} \operatorname{Hypergeometric2F1}\left(1, \frac{3+m}{4}, \frac{7+m}{4}, c^2x^4\right)}{d^3(1+m)(3+m)}$$

output

```
(d*x)^(1+m)*(a+b*arctanh(c*x^2))/d/(1+m)-2*b*c*(d*x)^(3+m)*hypergeom([1, 3/4+1/4*m], [7/4+1/4*m], c^2*x^4)/d^3/(1+m)/(3+m)
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.86

$$\int (dx)^m (a + \operatorname{barctanh}(cx^2)) dx = \frac{x(dx)^m (-(3+m)(a + \operatorname{barctanh}(cx^2))) + 2bcx^2 \operatorname{Hypergeometric2F1}\left(1, \frac{3+m}{4}, \frac{7+m}{4}, c^2x^4\right)}{(1+m)(3+m)}$$

input

```
Integrate[(d*x)^m*(a + b*ArcTanh[c*x^2]), x]
```

output

$$-\left(\frac{x(d*x)^m \left(-((3+m)(a+b*\text{ArcTanh}[c*x^2])\right) + 2*b*c*x^2*\text{Hypergeometric2F1}\left[1, (3+m)/4, (7+m)/4, c^2*x^4\right]\right)}{(1+m)(3+m)}\right)$$
Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {6464, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2)) dx$$

$$\downarrow 6464$$

$$\frac{(dx)^{m+1} (a + b \operatorname{arctanh}(cx^2))}{d(m+1)} - \frac{2bc \int \frac{(dx)^{m+2}}{1-c^2x^4} dx}{d^2(m+1)}$$

$$\downarrow 888$$

$$\frac{(dx)^{m+1} (a + b \operatorname{arctanh}(cx^2))}{d(m+1)} - \frac{2bc(dx)^{m+3} \operatorname{Hypergeometric2F1}\left(1, \frac{m+3}{4}, \frac{m+7}{4}, c^2x^4\right)}{d^3(m+1)(m+3)}$$

input

$$\text{Int}[(d*x)^m*(a + b*\text{ArcTanh}[c*x^2]), x]$$

output

$$\left(\frac{(d*x)^{(1+m)*(a + b*\text{ArcTanh}[c*x^2])}}{d*(1+m)} - \frac{(2*b*c*(d*x)^{(3+m)*\text{Hypergeometric2F1}\left[1, (3+m)/4, (7+m)/4, c^2*x^4\right]}}{d^3*(1+m)*(3+m)}\right)$$

Definitions of rubi rules used

rule 888 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p *((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 6464 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))*((d_)*(x_))^(m_), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]`

Maple [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2)) dx$$

input `int((d*x)^m*(a+b*arctanh(c*x^2)),x)`

output `int((d*x)^m*(a+b*arctanh(c*x^2)),x)`

Fricas [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2)) dx = \int (b \operatorname{arctanh}(cx^2) + a)(dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^2)),x, algorithm="fricas")`

output `integral((b*arctanh(c*x^2) + a)*(d*x)^m, x)`

Sympy [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2)) dx = \int (dx)^m (a + b \operatorname{atanh}(cx^2)) dx$$

input `integrate((d*x)**m*(a+b*atanh(c*x**2)),x)`

output `Integral((d*x)**m*(a + b*atanh(c*x**2)), x)`

Maxima [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2)) dx = \int (b \operatorname{artanh}(cx^2) + a)(dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^2)),x, algorithm="maxima")`

output `1/2*(4*c*d^m*integrate(x^2*x^m/(c^2*(m+1)*x^4 - m - 1), x) + (d^m*x*x^m*log(c*x^2 + 1) - d^m*x*x^m*log(-c*x^2 + 1))/(m + 1))*b + (d*x)^(m + 1)*a/(d*(m + 1))`

Giac [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2)) dx = \int (b \operatorname{artanh}(cx^2) + a)(dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^2)),x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)*(d*x)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^2)) dx = \int (dx)^m (a + b \operatorname{atanh}(cx^2)) dx$$

input `int((d*x)^m*(a + b*atanh(c*x^2)),x)`output `int((d*x)^m*(a + b*atanh(c*x^2)), x)`**Reduce [F]**

$$\begin{aligned} & \int (dx)^m (a + b \operatorname{arctanh}(cx^2)) dx \\ &= \frac{d^m (x^m a x + (\int x^m \operatorname{atanh}(c x^2) dx) b m + (\int x^m \operatorname{atanh}(c x^2) dx) b)}{m + 1} \end{aligned}$$

input `int((d*x)^m*(a+b*atanh(c*x^2)),x)`output `(d**m*(x**m*a*x + int(x**m*atanh(c*x**2),x)*b*m + int(x**m*atanh(c*x**2),x)*b))/(m + 1)`

$$3.97 \quad \int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^2)} dx$$

Optimal result	846
Mathematica [N/A]	846
Rubi [N/A]	847
Maple [N/A]	847
Fricas [N/A]	848
Sympy [N/A]	848
Maxima [N/A]	848
Giac [N/A]	849
Mupad [N/A]	849
Reduce [N/A]	850

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^2)} dx = \operatorname{Int}\left(\frac{(dx)^m}{a + b \operatorname{arctanh}(cx^2)}, x\right)$$

output `Defer(Int)((d*x)^m/(a+b*arctanh(c*x^2)),x)`

Mathematica [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^2)} dx = \int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^2)} dx$$

input `Integrate[(d*x)^m/(a + b*ArcTanh[c*x^2]),x]`

output `Integrate[(d*x)^m/(a + b*ArcTanh[c*x^2]), x]`

Rubi [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^2)} dx$$

↓ 6468

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^2)} dx$$

input `Int[(d*x)^m/(a + b*ArcTanh[c*x^2]),x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^2)} dx$$

input `int((d*x)^m/(a+b*arctanh(c*x^2)),x)`

output `int((d*x)^m/(a+b*arctanh(c*x^2)),x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^2)} dx = \int \frac{(dx)^m}{b \operatorname{artanh}(cx^2) + a} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x^2)),x, algorithm="fricas")`

output `integral((d*x)^m/(b*arctanh(c*x^2) + a), x)`

Sympy [N/A]

Not integrable

Time = 45.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^2)} dx = \int \frac{(dx)^m}{a + b \operatorname{atanh}(cx^2)} dx$$

input `integrate((d*x)**m/(a+b*atanh(c*x**2)),x)`

output `Integral((d*x)**m/(a + b*atanh(c*x**2)), x)`

Maxima [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^2)} dx = \int \frac{(dx)^m}{b \operatorname{artanh}(cx^2) + a} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x^2)),x, algorithm="maxima")`

output `integrate((d*x)^m/(b*arctanh(c*x^2) + a), x)`

Giac [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^2)} dx = \int \frac{(dx)^m}{b \operatorname{artanh}(cx^2) + a} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x^2)),x, algorithm="giac")`

output `integrate((d*x)^m/(b*arctanh(c*x^2) + a), x)`

Mupad [N/A]

Not integrable

Time = 3.47 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^2)} dx = \int \frac{(dx)^m}{a + b \operatorname{atanh}(cx^2)} dx$$

input `int((d*x)^m/(a + b*atanh(c*x^2)),x)`

output `int((d*x)^m/(a + b*atanh(c*x^2)), x)`

Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^2)} dx = d^m \left(\int \frac{x^m}{\operatorname{atanh}(cx^2) b + a} dx \right)$$

input

```
int((d*x)^m/(a+b*atanh(c*x^2)),x)
```

output

```
d**m*int(x**m/(atanh(c*x**2)*b + a),x)
```

3.98
$$\int \frac{(dx)^m}{(a+b\operatorname{arctanh}(cx^2))^2} dx$$

Optimal result	851
Mathematica [N/A]	851
Rubi [N/A]	852
Maple [N/A]	852
Fricas [N/A]	853
Sympy [F(-1)]	853
Maxima [N/A]	853
Giac [N/A]	854
Mupad [N/A]	854
Reduce [N/A]	855

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{(dx)^m}{(a + b\operatorname{arctanh}(cx^2))^2} dx = \operatorname{Int}\left(\frac{(dx)^m}{(a + b\operatorname{arctanh}(cx^2))^2}, x\right)$$

output

```
Defer(Int)((d*x)^m/(a+b*arctanh(c*x^2))^2,x)
```

Mathematica [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{(a + b\operatorname{arctanh}(cx^2))^2} dx = \int \frac{(dx)^m}{(a + b\operatorname{arctanh}(cx^2))^2} dx$$

input

```
Integrate[(d*x)^m/(a + b*ArcTanh[c*x^2])^2,x]
```

output

```
Integrate[(d*x)^m/(a + b*ArcTanh[c*x^2])^2, x]
```


Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^2))^2} dx$$

↓ 6468

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^2))^2} dx$$

input `Int[(d*x)^m/(a + b*ArcTanh[c*x^2])^2,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^2))^2} dx$$

input `int((d*x)^m/(a+b*arctanh(c*x^2))^2,x)`

output `int((d*x)^m/(a+b*arctanh(c*x^2))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.00

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^2))^2} dx = \int \frac{(dx)^m}{(b \operatorname{artanh}(cx^2) + a)^2} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x^2))^2,x, algorithm="fricas")`

output `integral((d*x)^m/(b^2*arctanh(c*x^2)^2 + 2*a*b*arctanh(c*x^2) + a^2), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^2))^2} dx = \text{Timed out}$$

input `integrate((d*x)**m/(a+b*atanh(c*x**2))**2,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 134, normalized size of antiderivative = 7.44

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^2))^2} dx = \int \frac{(dx)^m}{(b \operatorname{artanh}(cx^2) + a)^2} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x^2))^2,x, algorithm="maxima")`

output

```
(c^2*d^m*x^4 - d^m)*x^m/(b^2*c*x*log(c*x^2 + 1) - b^2*c*x*log(-c*x^2 + 1)
+ 2*a*b*c*x) + integrate(-(c^2*d^m*(m + 3)*x^4 - d^m*(m - 1))*x^m/(b^2*c*x
^2*log(c*x^2 + 1) - b^2*c*x^2*log(-c*x^2 + 1) + 2*a*b*c*x^2), x)
```

Giac [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^2))^2} dx = \int \frac{(dx)^m}{(b \operatorname{artanh}(cx^2) + a)^2} dx$$

input

```
integrate((d*x)^m/(a+b*arctanh(c*x^2))^2,x, algorithm="giac")
```

output

```
integrate((d*x)^m/(b*arctanh(c*x^2) + a)^2, x)
```

Mupad [N/A]

Not integrable

Time = 3.49 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^2))^2} dx = \int \frac{(dx)^m}{(a + b \operatorname{atanh}(cx^2))^2} dx$$

input

```
int((d*x)^m/(a + b*atanh(c*x^2))^2,x)
```

output

```
int((d*x)^m/(a + b*atanh(c*x^2))^2, x)
```

Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.11

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^2))^2} dx = d^m \left(\int \frac{x^m}{\operatorname{atanh}(cx^2)^2 b^2 + 2 \operatorname{atanh}(cx^2) ab + a^2} dx \right)$$

input

```
int((d*x)^m/(a+b*atanh(c*x^2))^2,x)
```

output

```
d**m*int(x**m/(atanh(c*x**2)**2*b**2 + 2*atanh(c*x**2)*a*b + a**2),x)
```

3.99 $\int x^{11}(a + b \operatorname{arctanh}(cx^3)) dx$

Optimal result	856
Mathematica [A] (verified)	856
Rubi [A] (verified)	857
Maple [A] (verified)	858
Fricas [A] (verification not implemented)	859
Sympy [F(-1)]	859
Maxima [A] (verification not implemented)	860
Giac [A] (verification not implemented)	860
Mupad [B] (verification not implemented)	861
Reduce [B] (verification not implemented)	861

Optimal result

Integrand size = 14, antiderivative size = 54

$$\int x^{11}(a + b \operatorname{arctanh}(cx^3)) dx = \frac{bx^3}{12c^3} + \frac{bx^9}{36c} - \frac{b \operatorname{arctanh}(cx^3)}{12c^4} + \frac{1}{12}x^{12}(a + b \operatorname{arctanh}(cx^3))$$

output

```
1/12*b*x^3/c^3+1/36*b*x^9/c-1/12*b*arctanh(c*x^3)/c^4+1/12*x^12*(a+b*arctanh(c*x^3))
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.44

$$\int x^{11}(a + b \operatorname{arctanh}(cx^3)) dx = \frac{bx^3}{12c^3} + \frac{bx^9}{36c} + \frac{ax^{12}}{12} + \frac{1}{12}bx^{12}\operatorname{arctanh}(cx^3) + \frac{b \log(1 - cx^3)}{24c^4} - \frac{b \log(1 + cx^3)}{24c^4}$$

input

```
Integrate[x^11*(a + b*ArcTanh[c*x^3]),x]
```

output

```
(b*x^3)/(12*c^3) + (b*x^9)/(36*c) + (a*x^12)/12 + (b*x^12*ArcTanh[c*x^3])/12 + (b*Log[1 - c*x^3])/(24*c^4) - (b*Log[1 + c*x^3])/(24*c^4)
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.98, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6452, 807, 254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^{11}(a + b \operatorname{arctanh}(cx^3)) dx \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{12}x^{12}(a + b \operatorname{arctanh}(cx^3)) - \frac{1}{4}bc \int \frac{x^{14}}{1 - c^2x^6} dx \\
 & \quad \downarrow \text{807} \\
 & \frac{1}{12}x^{12}(a + b \operatorname{arctanh}(cx^3)) - \frac{1}{12}bc \int \frac{x^{12}}{1 - c^2x^6} dx^3 \\
 & \quad \downarrow \text{254} \\
 & \frac{1}{12}x^{12}(a + b \operatorname{arctanh}(cx^3)) - \frac{1}{12}bc \int \left(-\frac{x^6}{c^2} + \frac{1}{c^4(1 - c^2x^6)} - \frac{1}{c^4} \right) dx^3 \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{12}x^{12}(a + b \operatorname{arctanh}(cx^3)) - \frac{1}{12}bc \left(\frac{\operatorname{arctanh}(cx^3)}{c^5} - \frac{x^3}{c^4} - \frac{x^9}{3c^2} \right)
 \end{aligned}$$

input `Int[x^11*(a + b*ArcTanh[c*x^3]),x]`

output `(x^12*(a + b*ArcTanh[c*x^3]))/12 - (b*c*(-(x^3/c^4) - x^9/(3*c^2) + ArcTanh[c*x^3]/c^5))/12`

Defintions of rubi rules used

rule 254 `Int[(x_)^(m_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^2, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 3]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.04

method	result	si
parallelrisch	$-\frac{-3b \operatorname{arctanh}(cx^3)x^{12}c^4 - 3ac^4x^{12} - bc^3x^9 - 3bcx^3 + 3b \operatorname{arctanh}(cx^3)}{36c^4}$	56
default	$\frac{ax^{12}}{12} + \frac{bx^{12} \operatorname{arctanh}(cx^3)}{12} + \frac{bx^9}{36c} + \frac{bx^3}{12c^3} - \frac{b \ln(cx^3+1)}{24c^4} + \frac{b \ln(cx^3-1)}{24c^4}$	60
parts	$\frac{ax^{12}}{12} + \frac{bx^{12} \operatorname{arctanh}(cx^3)}{12} + \frac{bx^9}{36c} + \frac{bx^3}{12c^3} - \frac{b \ln(cx^3+1)}{24c^4} + \frac{b \ln(cx^3-1)}{24c^4}$	60
risch	$\frac{x^{12}b \ln(cx^3+1)}{24} - \frac{x^{12}b \ln(-cx^3+1)}{24} + \frac{ax^{12}}{12} + \frac{bx^9}{36c} + \frac{bx^3}{12c^3} - \frac{b \ln(cx^3+1)}{24c^4} + \frac{b \ln(cx^3-1)}{24c^4}$	83
orering	$\frac{(10c^4x^{12} + 11c^2x^6 - 21)(a + b \operatorname{arctanh}(cx^3))}{54c^4} - \frac{(c^2x^6 + 3)(cx^3 - 1)(cx^3 + 1)(11x^{10}(a + b \operatorname{arctanh}(cx^3)) + \frac{3x^{13}bc}{-c^2x^6 + 1})}{108x^{10}c^4}$	100

input `int(x^11*(a+b*arctanh(c*x^3)),x,method=_RETURNVERBOSE)`

output

```
-1/36*(-3*b*arctanh(c*x^3)*x^12*c^4-3*a*c^4*x^12-b*c^3*x^9-3*b*c*x^3+3*b*a
rctanh(c*x^3))/c^4
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.19

$$\int x^{11}(a+b\operatorname{arctanh}(cx^3)) dx = \frac{6ac^4x^{12} + 2bc^3x^9 + 6bcx^3 + 3(bc^4x^{12} - b)\log\left(-\frac{cx^3+1}{cx^3-1}\right)}{72c^4}$$

input

```
integrate(x^11*(a+b*arctanh(c*x^3)),x, algorithm="fricas")
```

output

```
1/72*(6*a*c^4*x^12 + 2*b*c^3*x^9 + 6*b*c*x^3 + 3*(b*c^4*x^12 - b)*log(-(c*
x^3 + 1)/(c*x^3 - 1)))/c^4
```

Sympy [F(-1)]

Timed out.

$$\int x^{11}(a + b\operatorname{arctanh}(cx^3)) dx = \text{Timed out}$$

input

```
integrate(x**11*(a+b*atanh(c*x**3)),x)
```

output

```
Timed out
```


Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.28

$$\int x^{11}(a + b \operatorname{arctanh}(cx^3)) dx = \frac{1}{12} ax^{12} + \frac{1}{72} \left(6x^{12} \operatorname{artanh}(cx^3) + c \left(\frac{2(c^2x^9 + 3x^3)}{c^4} - \frac{3 \log(cx^3 + 1)}{c^5} + \frac{3 \log(cx^3 - 1)}{c^5} \right) \right) b$$

input `integrate(x^11*(a+b*arctanh(c*x^3)),x, algorithm="maxima")`output `1/12*a*x^12 + 1/72*(6*x^12*arctanh(c*x^3) + c*(2*(c^2*x^9 + 3*x^3)/c^4 - 3*log(c*x^3 + 1)/c^5 + 3*log(c*x^3 - 1)/c^5))*b`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.44

$$\int x^{11}(a + b \operatorname{arctanh}(cx^3)) dx = \frac{1}{24} bx^{12} \log\left(-\frac{cx^3 + 1}{cx^3 - 1}\right) + \frac{1}{12} ax^{12} + \frac{bx^9}{36c} + \frac{bx^3}{12c^3} - \frac{b \log(cx^3 + 1)}{24c^4} + \frac{b \log(cx^3 - 1)}{24c^4}$$

input `integrate(x^11*(a+b*arctanh(c*x^3)),x, algorithm="giac")`output `1/24*b*x^12*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 1/12*a*x^12 + 1/36*b*x^9/c + 1/12*b*x^3/c^3 - 1/24*b*log(c*x^3 + 1)/c^4 + 1/24*b*log(c*x^3 - 1)/c^4`

Mupad [B] (verification not implemented)

Time = 3.82 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.28

$$\int x^{11}(a + b \operatorname{arctanh}(cx^3)) dx = \frac{ax^{12}}{12} + \frac{bx^3}{12c^3} + \frac{bx^9}{36c} + \frac{bx^{12} \ln(cx^3 + 1)}{24} - \frac{bx^{12} \ln(1 - cx^3)}{24} + \frac{b \operatorname{atan}(cx^3 \operatorname{li}) \operatorname{li}}{12c^4}$$

input `int(x^11*(a + b*atanh(c*x^3)),x)`output `(a*x^12)/12 + (b*x^3)/(12*c^3) + (b*x^9)/(36*c) + (b*atan(c*x^3*1i)*1i)/(12*c^4) + (b*x^12*log(c*x^3 + 1))/24 - (b*x^12*log(1 - c*x^3))/24`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00

$$\int x^{11}(a + b \operatorname{arctanh}(cx^3)) dx = \frac{3 \operatorname{atanh}(cx^3) b c^4 x^{12} - 3 \operatorname{atanh}(cx^3) b + 3 a c^4 x^{12} + b c^3 x^9 + 3 b c x^3}{36 c^4}$$

input `int(x^11*(a+b*atanh(c*x^3)),x)`output `(3*atanh(c*x**3)*b*c**4*x**12 - 3*atanh(c*x**3)*b + 3*a*c**4*x**12 + b*c**3*x**9 + 3*b*c*x**3)/(36*c**4)`

3.100 $\int x^8(a + b \operatorname{arctanh}(cx^3)) dx$

Optimal result	862
Mathematica [A] (verified)	862
Rubi [A] (verified)	863
Maple [A] (verified)	864
Fricas [A] (verification not implemented)	865
Sympy [F(-1)]	865
Maxima [A] (verification not implemented)	865
Giac [A] (verification not implemented)	866
Mupad [B] (verification not implemented)	866
Reduce [B] (verification not implemented)	866

Optimal result

Integrand size = 14, antiderivative size = 48

$$\int x^8(a + b \operatorname{arctanh}(cx^3)) dx = \frac{bx^6}{18c} + \frac{1}{9}x^9(a + b \operatorname{arctanh}(cx^3)) + \frac{b \log(1 - c^2x^6)}{18c^3}$$

output $1/18*b*x^6/c+1/9*x^9*(a+b*\operatorname{arctanh}(c*x^3))+1/18*b*\ln(-c^2*x^6+1)/c^3$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.10

$$\int x^8(a + b \operatorname{arctanh}(cx^3)) dx = \frac{bx^6}{18c} + \frac{ax^9}{9} + \frac{1}{9}bx^9 \operatorname{arctanh}(cx^3) + \frac{b \log(1 - c^2x^6)}{18c^3}$$

input $\operatorname{Integrate}[x^8*(a + b*\operatorname{ArcTanh}[c*x^3]),x]$

output $(b*x^6)/(18*c) + (a*x^9)/9 + (b*x^9*\operatorname{ArcTanh}[c*x^3])/9 + (b*\operatorname{Log}[1 - c^2*x^6])/18*c^3$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6452, 798, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^8 (a + b \operatorname{arctanh}(cx^3)) dx \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{9} x^9 (a + b \operatorname{arctanh}(cx^3)) - \frac{1}{3} bc \int \frac{x^{11}}{1 - c^2 x^6} dx \\
 & \quad \downarrow \text{798} \\
 & \frac{1}{9} x^9 (a + b \operatorname{arctanh}(cx^3)) - \frac{1}{18} bc \int \frac{x^6}{1 - c^2 x^6} dx^6 \\
 & \quad \downarrow \text{49} \\
 & \frac{1}{9} x^9 (a + b \operatorname{arctanh}(cx^3)) - \frac{1}{18} bc \int \left(-\frac{1}{c^2} - \frac{1}{c^2 (c^2 x^6 - 1)} \right) dx^6 \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{9} x^9 (a + b \operatorname{arctanh}(cx^3)) - \frac{1}{18} bc \left(-\frac{x^6}{c^2} - \frac{\log(1 - c^2 x^6)}{c^4} \right)
 \end{aligned}$$

input `Int [x^8*(a + b*ArcTanh[c*x^3]), x]`

output `(x^9*(a + b*ArcTanh[c*x^3]))/9 - (b*c*(-(x^6/c^2) - Log[1 - c^2*x^6]/c^4))/18`

Definitions of rubi rules used

rule 49 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}*((c_.) + (d_.)(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{IGtQ}[m, 0] \&\& \text{IGtQ}[m + n + 2, 0]$

rule 798 $\text{Int}[(x_)^{(m_.)}*((a_) + (b_.)(x_)^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)*(a + b*x)^p, x}], x, x^n], x] /; \text{FreeQ}[\{a, b, m, n, p\}, x] \&\& \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 6452 $\text{Int}[(a_.) + \text{ArcTanh}[(c_.)(x_)^{(n_.)}]*(b_.)^{(p_.)}(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m + 1)), x] - \text{Simp}[b*c*n*(p/(m + 1)) \text{ Int}[x^{(m + n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p - 1)/(1 - c^2*x^{(2*n)})}), x], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \&\& \text{IGtQ}[p, 0] \&\& (\text{EqQ}[p, 1] \|\| (\text{EqQ}[n, 1] \&\& \text{IntegerQ}[m])) \&\& \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.94

method	result	size
default	$\frac{ax^9}{9} + \frac{bx^9 \arctanh(cx^3)}{9} + \frac{bx^6}{18c} + \frac{b \ln(c^2x^6-1)}{18c^3}$	45
parts	$\frac{ax^9}{9} + \frac{bx^9 \arctanh(cx^3)}{9} + \frac{bx^6}{18c} + \frac{b \ln(c^2x^6-1)}{18c^3}$	45
parallelrisch	$\frac{2b \arctanh(cx^3)x^9c^3+2ac^3x^9+c^2bx^6+2b \ln(cx^3-1)+2b \arctanh(cx^3)}{18c^3}$	59
risch	$\frac{x^9b \ln(cx^3+1)}{18} - \frac{x^9b \ln(-cx^3+1)}{18} + \frac{ax^9}{9} + \frac{bx^6}{18c} + \frac{b \ln(c^2x^6-1)}{18c^3}$	62

input $\text{int}(x^8*(a+b*\arctanh(c*x^3)), x, \text{method}=_RETURNVERBOSE)$

output $1/9*a*x^9+1/9*b*x^9*\arctanh(c*x^3)+1/18*b*x^6/c+1/18*b/c^3*\ln(c^2*x^6-1)$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.29

$$\int x^8(a + \operatorname{arctanh}(cx^3)) dx = \frac{bc^3x^9 \log\left(-\frac{cx^3+1}{cx^3-1}\right) + 2ac^3x^9 + bc^2x^6 + b \log(c^2x^6 - 1)}{18c^3}$$

input `integrate(x^8*(a+b*arctanh(c*x^3)),x, algorithm="fricas")`output `1/18*(b*c^3*x^9*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 2*a*c^3*x^9 + b*c^2*x^6 + b*log(c^2*x^6 - 1))/c^3`**Sympy [F(-1)]**

Timed out.

$$\int x^8(a + \operatorname{arctanh}(cx^3)) dx = \text{Timed out}$$

input `integrate(x**8*(a+b*atanh(c*x**3)),x)`output `Timed out`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.96

$$\begin{aligned} \int x^8(a + \operatorname{arctanh}(cx^3)) dx \\ = \frac{1}{9}ax^9 + \frac{1}{18} \left(2x^9 \operatorname{artanh}(cx^3) + \left(\frac{x^6}{c^2} + \frac{\log(c^2x^6 - 1)}{c^4} \right) c \right) b \end{aligned}$$

input `integrate(x^8*(a+b*arctanh(c*x^3)),x, algorithm="maxima")`output `1/9*a*x^9 + 1/18*(2*x^9*arctanh(c*x^3) + (x^6/c^2 + log(c^2*x^6 - 1)/c^4)*c)*b`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.19

$$\int x^8 (a + b \operatorname{arctanh}(cx^3)) dx = \frac{1}{18} bx^9 \log\left(-\frac{cx^3 + 1}{cx^3 - 1}\right) + \frac{1}{9} ax^9 + \frac{bx^6}{18c} + \frac{b \log(c^2 x^6 - 1)}{18c^3}$$

input `integrate(x^8*(a+b*arctanh(c*x^3)),x, algorithm="giac")`

output `1/18*b*x^9*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 1/9*a*x^9 + 1/18*b*x^6/c + 1/18*b*log(c^2*x^6 - 1)/c^3`

Mupad [B] (verification not implemented)

Time = 3.39 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.27

$$\int x^8 (a + b \operatorname{arctanh}(cx^3)) dx = \frac{ax^9}{9} + \frac{b \ln(c^2 x^6 - 1)}{18c^3} + \frac{bx^6}{18c} + \frac{bx^9 \ln(cx^3 + 1)}{18} - \frac{bx^9 \ln(1 - cx^3)}{18}$$

input `int(x^8*(a + b*atanh(c*x^3)),x)`

output `(a*x^9)/9 + (b*log(c^2*x^6 - 1))/(18*c^3) + (b*x^6)/(18*c) + (b*x^9*log(c*x^3 + 1))/18 - (b*x^9*log(1 - c*x^3))/18`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.65

$$\int x^8 (a + b \operatorname{arctanh}(cx^3)) dx = \frac{2a \operatorname{atanh}(cx^3) b c^3 x^9 - 2a \operatorname{atanh}(cx^3) b + 2 \log\left(c^{\frac{2}{3}} x^2 - c^{\frac{1}{3}} x + 1\right) b + 2 \log\left(c^{\frac{2}{3}} x + c^{\frac{1}{3}}\right) b + 2a c^3 x^9 + b c^2 x^6}{18c^3}$$

input `int(x^8*(a+b*atanh(c*x^3)),x)`

output
$$\frac{(2*\operatorname{atanh}(c*x**3)*b*c**3*x**9 - 2*\operatorname{atanh}(c*x**3)*b + 2*\log(c**(2/3)*x**2 - c**(1/3)*x + 1)*b + 2*\log(c**(2/3)*x + c**(1/3))*b + 2*a*c**3*x**9 + b*c**2*x**6)/(18*c**3)}$$

3.101 $\int x^5(a + \operatorname{arctanh}(cx^3)) dx$

Optimal result	868
Mathematica [A] (verified)	868
Rubi [A] (verified)	869
Maple [A] (verified)	870
Fricas [A] (verification not implemented)	871
Sympy [F(-1)]	871
Maxima [A] (verification not implemented)	872
Giac [B] (verification not implemented)	872
Mupad [B] (verification not implemented)	873
Reduce [B] (verification not implemented)	873

Optimal result

Integrand size = 14, antiderivative size = 43

$$\int x^5(a + \operatorname{arctanh}(cx^3)) dx = \frac{bx^3}{6c} - \frac{\operatorname{arctanh}(cx^3)}{6c^2} + \frac{1}{6}x^6(a + \operatorname{arctanh}(cx^3))$$

output $1/6*b*x^3/c-1/6*b*\operatorname{arctanh}(c*x^3)/c^2+1/6*x^6*(a+b*\operatorname{arctanh}(c*x^3))$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.56

$$\int x^5(a + \operatorname{arctanh}(cx^3)) dx = \frac{bx^3}{6c} + \frac{ax^6}{6} + \frac{1}{6}bx^6\operatorname{arctanh}(cx^3) + \frac{b \log(1 - cx^3)}{12c^2} - \frac{b \log(1 + cx^3)}{12c^2}$$

input $\operatorname{Integrate}[x^5*(a + b*\operatorname{ArcTanh}[c*x^3]), x]$

output $(b*x^3)/(6*c) + (a*x^6)/6 + (b*x^6*\operatorname{ArcTanh}[c*x^3])/6 + (b*\operatorname{Log}[1 - c*x^3])/(12*c^2) - (b*\operatorname{Log}[1 + c*x^3])/(12*c^2)$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6452, 807, 262, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^5(a + b \operatorname{arctanh}(cx^3)) dx \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{6}x^6(a + b \operatorname{arctanh}(cx^3)) - \frac{1}{2}bc \int \frac{x^8}{1 - c^2x^6} dx \\
 & \quad \downarrow \text{807} \\
 & \frac{1}{6}x^6(a + b \operatorname{arctanh}(cx^3)) - \frac{1}{6}bc \int \frac{x^6}{1 - c^2x^6} dx \\
 & \quad \downarrow \text{262} \\
 & \frac{1}{6}x^6(a + b \operatorname{arctanh}(cx^3)) - \frac{1}{6}bc \left(\frac{\int \frac{1}{1 - c^2x^6} dx^3}{c^2} - \frac{x^3}{c^2} \right) \\
 & \quad \downarrow \text{219} \\
 & \frac{1}{6}x^6(a + b \operatorname{arctanh}(cx^3)) - \frac{1}{6}bc \left(\frac{\operatorname{arctanh}(cx^3)}{c^3} - \frac{x^3}{c^2} \right)
 \end{aligned}$$

input `Int [x^5*(a + b*ArcTanh[c*x^3]), x]`

output `(x^6*(a + b*ArcTanh[c*x^3]))/6 - (b*c*(-(x^3/c^2) + ArcTanh[c*x^3]/c^3))/6`

Definitions of rubi rules used

rule 219 $\text{Int}[(a_ + (b_)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 262 $\text{Int}[(c_)(x_)^m((a_ + (b_)(x_)^2)^p), x_Symbol] \rightarrow \text{Simp}[c*(c*x)^{m-1}((a + b*x^2)^{p+1}/(b*(m+2*p+1))), x] - \text{Simp}[a*c^2*((m-1)/(b*(m+2*p+1))) \ \text{Int}[(c*x)^{m-2}*(a + b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{GtQ}[m, 2-1] \ \&\& \ \text{NeQ}[m+2*p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 807 $\text{Int}(x_)^{m_}((a_ + (b_)(x_)^n)^p), x_Symbol] \rightarrow \text{With}\{k = \text{GCD}[m+1, n]\}, \text{Simp}[1/k \ \text{Subst}[\text{Int}[x^{(m+1)/k-1}*(a + b*x^{n/k})^p, x], x, x^k], x] /; k \neq 1 /; \text{FreeQ}\{a, b, p\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$

rule 6452 $\text{Int}[(a_ + \text{ArcTanh}(c_)(x_)^n)*(b_)^p(x_)^m), x_Symbol] \rightarrow \text{Simp}[x^{m+1}((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \ \text{Int}[x^{m+n}((a + b*\text{ArcTanh}[c*x^n])^{p-1}/(1 - c^2*x^{2*n})), x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.07

method	result	size
parallelrisch	$-\frac{\arctanh(cx^3)bc^2x^6 - ac^2x^6 - bcx^3 + b \arctanh(cx^3)}{6c^2}$	46
default	$\frac{ax^6}{6} + \frac{bx^6 \arctanh(cx^3)}{6} + \frac{bx^3}{6c} - \frac{b \ln(cx^3+1)}{12c^2} + \frac{b \ln(cx^3-1)}{12c^2}$	57
parts	$\frac{ax^6}{6} + \frac{bx^6 \arctanh(cx^3)}{6} + \frac{bx^3}{6c} - \frac{b \ln(cx^3+1)}{12c^2} + \frac{b \ln(cx^3-1)}{12c^2}$	57
oring	$\frac{4(c^2x^6-1)(a+b \arctanh(cx^3))}{9c^2} - \frac{(cx^3-1)(cx^3+1)(5x^4(a+b \arctanh(cx^3))+\frac{3x^7bc}{-c^2x^6+1})}{18c^2x^4}$	83
risch	$\frac{bx^6 \ln(cx^3+1)}{12} - \frac{bx^6 \ln(-cx^3+1)}{12} + \frac{ax^6}{6} + \frac{bx^3}{6c} + \frac{b \ln(cx^3-1)}{12c^2} - \frac{b \ln(cx^3+1)}{12c^2} + \frac{b^2}{24ac^2}$	85

input `int(x^5*(a+b*arctanh(c*x^3)),x,method=_RETURNVERBOSE)`

output `-1/6*(-arctanh(c*x^3)*b*c^2*x^6-a*c^2*x^6-b*c*x^3+b*arctanh(c*x^3))/c^2`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.26

$$\int x^5 (a + b \operatorname{arctanh}(cx^3)) dx = \frac{2ac^2x^6 + 2bcx^3 + (bc^2x^6 - b) \log\left(-\frac{cx^3+1}{cx^3-1}\right)}{12c^2}$$

input `integrate(x^5*(a+b*arctanh(c*x^3)),x, algorithm="fricas")`

output `1/12*(2*a*c^2*x^6 + 2*b*c*x^3 + (b*c^2*x^6 - b)*log(-(c*x^3 + 1)/(c*x^3 - 1)))/c^2`

Sympy [F(-1)]

Timed out.

$$\int x^5 (a + b \operatorname{arctanh}(cx^3)) dx = \text{Timed out}$$

input `integrate(x**5*(a+b*atanh(c*x**3)),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.35

$$\int x^5 (a + b \operatorname{arctanh}(cx^3)) dx$$

$$= \frac{1}{6} ax^6 + \frac{1}{12} \left(2x^6 \operatorname{artanh}(cx^3) + c \left(\frac{2x^3}{c^2} - \frac{\log(cx^3 + 1)}{c^3} + \frac{\log(cx^3 - 1)}{c^3} \right) \right) b$$

input `integrate(x^5*(a+b*arctanh(c*x^3)),x, algorithm="maxima")`

output `1/6*a*x^6 + 1/12*(2*x^6*arctanh(c*x^3) + c*(2*x^3/c^2 - log(c*x^3 + 1)/c^3 + log(c*x^3 - 1)/c^3))*b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 181 vs. $2(37) = 74$.

Time = 0.13 (sec) , antiderivative size = 181, normalized size of antiderivative = 4.21

$$\int x^5 (a + b \operatorname{arctanh}(cx^3)) dx$$

$$= \frac{1}{3} c \left(\frac{(cx^3 + 1)b \log\left(-\frac{cx^3 + 1}{cx^3 - 1}\right)}{(cx^3 - 1) \left(\frac{(cx^3 + 1)^2 c^3}{(cx^3 - 1)^2} - \frac{2(cx^3 + 1)c^3}{cx^3 - 1} + c^3 \right)} + \frac{\frac{2(cx^3 + 1)a}{cx^3 - 1} + \frac{(cx^3 + 1)b}{cx^3 - 1} - b}{\frac{(cx^3 + 1)^2 c^3}{(cx^3 - 1)^2} - \frac{2(cx^3 + 1)c^3}{cx^3 - 1} + c^3} \right)$$

input `integrate(x^5*(a+b*arctanh(c*x^3)),x, algorithm="giac")`

output `1/3*c*((c*x^3 + 1)*b*log(-(c*x^3 + 1)/(c*x^3 - 1))/((c*x^3 - 1)*((c*x^3 + 1)^2*c^3/(c*x^3 - 1)^2 - 2*(c*x^3 + 1)*c^3/(c*x^3 - 1) + c^3)) + (2*(c*x^3 + 1)*a/(c*x^3 - 1) + (c*x^3 + 1)*b/(c*x^3 - 1) - b)/((c*x^3 + 1)^2*c^3/(c*x^3 - 1)^2 - 2*(c*x^3 + 1)*c^3/(c*x^3 - 1) + c^3))`

Mupad [B] (verification not implemented)

Time = 3.49 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.40

$$\int x^5 (a + b \operatorname{arctanh}(cx^3)) dx = \frac{ax^6}{6} + \frac{bx^3}{6c} + \frac{bx^6 \ln(cx^3 + 1)}{12} - \frac{bx^6 \ln(1 - cx^3)}{12} + \frac{b \operatorname{atan}(cx^3) \operatorname{li}}{6c^2}$$

input `int(x^5*(a + b*atanh(c*x^3)),x)`output `(a*x^6)/6 + (b*x^3)/(6*c) + (b*atan(c*x^3*1i)*1i)/(6*c^2) + (b*x^6*log(c*x^3 + 1))/12 - (b*x^6*log(1 - c*x^3))/12`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int x^5 (a + b \operatorname{arctanh}(cx^3)) dx = \frac{\operatorname{atanh}(cx^3) b c^2 x^6 - \operatorname{atanh}(cx^3) b + a c^2 x^6 + b c x^3}{6c^2}$$

input `int(x^5*(a+b*atanh(c*x^3)),x)`output `(atanh(c*x**3)*b*c**2*x**6 - atanh(c*x**3)*b + a*c**2*x**6 + b*c*x**3)/(6*c**2)`

3.102 $\int x^2(a + \operatorname{barctanh}(cx^3)) dx$

Optimal result	874
Mathematica [A] (verified)	874
Rubi [A] (verified)	875
Maple [A] (verified)	876
Fricas [A] (verification not implemented)	876
Sympy [F(-1)]	877
Maxima [A] (verification not implemented)	877
Giac [B] (verification not implemented)	877
Mupad [B] (verification not implemented)	878
Reduce [B] (verification not implemented)	878

Optimal result

Integrand size = 14, antiderivative size = 42

$$\int x^2(a + \operatorname{barctanh}(cx^3)) dx = \frac{ax^3}{3} + \frac{1}{3}bx^3\operatorname{arctanh}(cx^3) + \frac{b \log(1 - c^2x^6)}{6c}$$

output

```
1/3*a*x^3+1/3*b*x^3*arctanh(c*x^3)+1/6*b*ln(-c^2*x^6+1)/c
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int x^2(a + \operatorname{barctanh}(cx^3)) dx = \frac{ax^3}{3} + \frac{1}{3}bx^3\operatorname{arctanh}(cx^3) + \frac{b \log(1 - c^2x^6)}{6c}$$

input

```
Integrate[x^2*(a + b*ArcTanh[c*x^3]),x]
```

output

```
(a*x^3)/3 + (b*x^3*ArcTanh[c*x^3])/3 + (b*Log[1 - c^2*x^6])/(6*c)
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6452, 792}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2(a + \operatorname{arctanh}(cx^3)) dx$$

$$\downarrow 6452$$

$$\frac{1}{3}x^3(a + \operatorname{arctanh}(cx^3)) - bc \int \frac{x^5}{1 - c^2x^6} dx$$

$$\downarrow 792$$

$$\frac{1}{3}x^3(a + \operatorname{arctanh}(cx^3)) + \frac{b \log(1 - c^2x^6)}{6c}$$

input `Int[x^2*(a + b*ArcTanh[c*x^3]),x]`

output `(x^3*(a + b*ArcTanh[c*x^3]))/3 + (b*Log[1 - c^2*x^6])/(6*c)`

Defintions of rubi rules used

rule 792 `Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_.)), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

method	result	size
parts	$\frac{ax^3}{3} + \frac{bx^3 \operatorname{arctanh}(cx^3)}{3} + \frac{b \ln(-c^2x^6+1)}{6c}$	37
derivativdivides	$\frac{acx^3+b\left(cx^3 \operatorname{arctanh}(cx^3)+\frac{\ln(-c^2x^6+1)}{2}\right)}{3c}$	40
default	$\frac{acx^3+b\left(cx^3 \operatorname{arctanh}(cx^3)+\frac{\ln(-c^2x^6+1)}{2}\right)}{3c}$	40
parallelrisc	$\frac{b \operatorname{arctanh}(cx^3)x^3c+acx^3+b \ln(cx^3-1)+b \operatorname{arctanh}(cx^3)}{3c}$	43
risc	$\frac{bx^3 \ln(cx^3+1)}{6} - \frac{bx^3 \ln(-cx^3+1)}{6} + \frac{ax^3}{3} + \frac{b \ln(c^2x^6-1)}{6c}$	53

input `int(x^2*(a+b*arctanh(c*x^3)),x,method=_RETURNVERBOSE)`

output `1/3*a*x^3+1/3*b*x^3*arctanh(c*x^3)+1/6*b*ln(-c^2*x^6+1)/c`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.19

$$\int x^2(a + b \operatorname{arctanh}(cx^3)) dx = \frac{bcx^3 \log\left(-\frac{cx^3+1}{cx^3-1}\right) + 2acx^3 + b \log(c^2x^6 - 1)}{6c}$$

input `integrate(x^2*(a+b*arctanh(c*x^3)),x, algorithm="fricas")`

output `1/6*(b*c*x^3*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 2*a*c*x^3 + b*log(c^2*x^6 - 1))/c`

Sympy [F(-1)]

Timed out.

$$\int x^2(a + \operatorname{barctanh}(cx^3)) dx = \text{Timed out}$$

input `integrate(x**2*(a+b*atanh(c*x**3)),x)`output `Timed out`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

$$\int x^2(a + \operatorname{barctanh}(cx^3)) dx = \frac{1}{3}ax^3 + \frac{(2cx^3 \operatorname{artanh}(cx^3) + \log(-c^2x^6 + 1))b}{6c}$$

input `integrate(x^2*(a+b*arctanh(c*x^3)),x, algorithm="maxima")`output `1/3*a*x^3 + 1/6*(2*c*x^3*arctanh(c*x^3) + log(-c^2*x^6 + 1))*b/c`**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 188 vs. 2(36) = 72.

Time = 0.13 (sec) , antiderivative size = 188, normalized size of antiderivative = 4.48

$$\int x^2(a + \operatorname{barctanh}(cx^3)) dx = \frac{1}{3}ax^3 + \frac{1}{3}bc \left(\frac{\log\left(\frac{|-cx^3-1|}{|cx^3-1|}\right)}{c^2} - \frac{\log\left(\left|-\frac{cx^3+1}{cx^3-1} + 1\right|\right)}{c^2} + \frac{\log\left(\frac{\frac{c\left(\frac{cx^3+1}{cx^3-1}+1\right)}{(cx^3+1)c}-c}{\frac{c\left(\frac{cx^3+1}{cx^3-1}-1\right)}{(cx^3+1)c}-c}\right)}{c^2\left(\frac{cx^3+1}{cx^3-1}-1\right)} \right)$$

input `integrate(x^2*(a+b*arctanh(c*x^3)),x, algorithm="giac")`

output `1/3*a*x^3 + 1/3*b*c*(log(abs(-c*x^3 - 1)/abs(c*x^3 - 1))/c^2 - log(abs(-(c*x^3 + 1)/(c*x^3 - 1) + 1))/c^2 + log(-(c*((c*x^3 + 1)/(c*x^3 - 1) + 1)/((c*x^3 + 1)*c/(c*x^3 - 1) - c) + 1)/(c*((c*x^3 + 1)/(c*x^3 - 1) + 1)/((c*x^3 + 1)*c/(c*x^3 - 1) - c) - 1))/c^2*((c*x^3 + 1)/(c*x^3 - 1) - 1))`

Mupad [B] (verification not implemented)

Time = 3.32 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.24

$$\int x^2(a + b \operatorname{arctanh}(cx^3)) dx = \frac{ax^3}{3} + \frac{b \ln(c^2 x^6 - 1)}{6c} + \frac{bx^3 \ln(cx^3 + 1)}{6} - \frac{bx^3 \ln(1 - cx^3)}{6}$$

input `int(x^2*(a + b*atanh(c*x^3)),x)`

output `(a*x^3)/3 + (b*log(c^2*x^6 - 1))/(6*c) + (b*x^3*log(c*x^3 + 1))/6 - (b*x^3*log(1 - c*x^3))/6`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.50

$$\int x^2(a + b \operatorname{arctanh}(cx^3)) dx = \frac{\operatorname{atanh}(cx^3)bcx^3 - \operatorname{atanh}(cx^3)b + \log\left(c^{\frac{2}{3}}x^2 - c^{\frac{1}{3}}x + 1\right)b + \log\left(c^{\frac{2}{3}}x + c^{\frac{1}{3}}\right)b + acx^3}{3c}$$

input `int(x^2*(a+b*atanh(c*x^3)),x)`

output `(atanh(c*x**3)*b*c*x**3 - atanh(c*x**3)*b + log(c**(2/3)*x**2 - c**(1/3)*x + 1)*b + log(c**(2/3)*x + c**(1/3))*b + a*c*x**3)/(3*c)`

3.103 $\int \frac{a + b \operatorname{arctanh}(cx^3)}{x} dx$

Optimal result	879
Mathematica [A] (verified)	879
Rubi [A] (verified)	880
Maple [C] (verified)	881
Fricas [F]	881
Sympy [F]	882
Maxima [F]	882
Giac [F]	882
Mupad [F(-1)]	883
Reduce [F]	883

Optimal result

Integrand size = 14, antiderivative size = 30

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x} dx = a \log(x) - \frac{1}{6} b \operatorname{PolyLog}(2, -cx^3) + \frac{1}{6} b \operatorname{PolyLog}(2, cx^3)$$

output `a*ln(x)-1/6*b*polylog(2,-c*x^3)+1/6*b*polylog(2,c*x^3)`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x} dx = a \log(x) + \frac{1}{6} b (-\operatorname{PolyLog}(2, -cx^3) + \operatorname{PolyLog}(2, cx^3))$$

input `Integrate[(a + b*ArcTanh[c*x^3])/x,x]`

output `a*Log[x] + (b*(-PolyLog[2, -(c*x^3)] + PolyLog[2, c*x^3]))/6`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.20, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6450, 6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x} dx$$

↓ 6450

$$\frac{1}{3} \int \frac{a + b \operatorname{arctanh}(cx^3)}{x^3} dx^3$$

↓ 6446

$$\frac{1}{3} \left(a \log(x^3) - \frac{1}{2} b \operatorname{PolyLog}(2, -cx^3) + \frac{1}{2} b \operatorname{PolyLog}(2, cx^3) \right)$$

input `Int[(a + b*ArcTanh[c*x^3])/x,x]`

output `(a*Log[x^3] - (b*PolyLog[2, -(c*x^3)])/2 + (b*PolyLog[2, c*x^3])/2)/3`

Defintions of rubi rules used

rule 6446 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))/(x_), x_Symbol] := Simp[a*Log[x], x] + (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) / ; FreeQ[{a, b, c}, x]`

rule 6450 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_.)/(x_), x_Symbol] := Simp[1/n Subst[Int[(a + b*ArcTanh[c*x])^p/x, x], x, x^n], x] / ; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.20 (sec) , antiderivative size = 92, normalized size of antiderivative = 3.07

method	result
default	$a \ln(x) + b \ln(x) \operatorname{arctanh}(cx^3) + \frac{b \left(\sum_{-R1=\operatorname{RootOf}(cZ^3-1)} \left(\ln(x) \ln\left(\frac{R1-x}{-R1}\right) + \operatorname{dilog}\left(\frac{R1-x}{-R1}\right) \right) \right)}{2} - \frac{b \left(\sum_{-R1=\operatorname{RootOf}(cZ^3-1)} \left(\ln(x) \ln\left(\frac{R1-x}{-R1}\right) + \operatorname{dilog}\left(\frac{R1-x}{-R1}\right) \right) \right)}{2}$
parts	$a \ln(x) + b \ln(x) \operatorname{arctanh}(cx^3) + \frac{b \left(\sum_{-R1=\operatorname{RootOf}(cZ^3-1)} \left(\ln(x) \ln\left(\frac{R1-x}{-R1}\right) + \operatorname{dilog}\left(\frac{R1-x}{-R1}\right) \right) \right)}{2} - \frac{b \left(\sum_{-R1=\operatorname{RootOf}(cZ^3-1)} \left(\ln(x) \ln\left(\frac{R1-x}{-R1}\right) + \operatorname{dilog}\left(\frac{R1-x}{-R1}\right) \right) \right)}{2}$
risch	$a \ln(x) - \frac{\ln(x) \ln(-cx^3+1)b}{2} + \frac{b \left(\sum_{-R1=\operatorname{RootOf}(cZ^3-1)} \left(\ln(x) \ln\left(\frac{R1-x}{-R1}\right) + \operatorname{dilog}\left(\frac{R1-x}{-R1}\right) \right) \right)}{2} + \frac{\ln(x) \ln(cx^3+1)}{2}$

input `int((a+b*arctanh(c*x^3))/x,x,method=_RETURNVERBOSE)`

output `a*ln(x)+b*ln(x)*arctanh(c*x^3)+1/2*b*sum(ln(x)*ln((_R1-x)/_R1)+dilog((_R1-x)/_R1),_R1=RootOf(_Z^3*c-1))-1/2*b*sum(ln(x)*ln((_R1-x)/_R1)+dilog((_R1-x)/_R1),_R1=RootOf(_Z^3*c+1))`

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x} dx = \int \frac{b \operatorname{arctanh}(cx^3) + a}{x} dx$$

input `integrate((a+b*arctanh(c*x^3))/x,x, algorithm="fricas")`

output `integral((b*arctanh(c*x^3) + a)/x, x)`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x} dx = \int \frac{a + b \operatorname{atanh}(cx^3)}{x} dx$$

input `integrate((a+b*atanh(c*x**3))/x,x)`

output `Integral((a + b*atanh(c*x**3))/x, x)`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x} dx = \int \frac{b \operatorname{artanh}(cx^3) + a}{x} dx$$

input `integrate((a+b*arctanh(c*x^3))/x,x, algorithm="maxima")`

output `1/2*b*integrate((log(c*x^3 + 1) - log(-c*x^3 + 1))/x, x) + a*log(x)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x} dx = \int \frac{b \operatorname{artanh}(cx^3) + a}{x} dx$$

input `integrate((a+b*arctanh(c*x^3))/x,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^3) + a)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x} dx = \int \frac{a + b \operatorname{atanh}(cx^3)}{x} dx$$

input `int((a + b*atanh(c*x^3))/x,x)`output `int((a + b*atanh(c*x^3))/x, x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x} dx = \left(\int \frac{\operatorname{atanh}(cx^3)}{x} dx \right) b + \log(x) a$$

input `int((a+b*atanh(c*x^3))/x,x)`output `int(atanh(c*x**3)/x,x)*b + log(x)*a`

3.104 $\int \frac{a+b\operatorname{arctanh}(cx^3)}{x^4} dx$

Optimal result	884
Mathematica [A] (verified)	884
Rubi [A] (verified)	885
Maple [A] (verified)	886
Fricas [A] (verification not implemented)	887
Sympy [F(-1)]	887
Maxima [A] (verification not implemented)	888
Giac [A] (verification not implemented)	888
Mupad [B] (verification not implemented)	888
Reduce [B] (verification not implemented)	889

Optimal result

Integrand size = 14, antiderivative size = 40

$$\int \frac{a + b\operatorname{arctanh}(cx^3)}{x^4} dx = -\frac{a + b\operatorname{arctanh}(cx^3)}{3x^3} + bc \log(x) - \frac{1}{6}bc \log(1 - c^2x^6)$$

output `-1/3*(a+b*arctanh(c*x^3))/x^3+b*c*ln(x)-1/6*b*c*ln(-c^2*x^6+1)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.12

$$\int \frac{a + b\operatorname{arctanh}(cx^3)}{x^4} dx = -\frac{a}{3x^3} - \frac{b\operatorname{arctanh}(cx^3)}{3x^3} + bc \log(x) - \frac{1}{6}bc \log(1 - c^2x^6)$$

input `Integrate[(a + b*ArcTanh[c*x^3])/x^4,x]`

output `-1/3*a/x^3 - (b*ArcTanh[c*x^3])/(3*x^3) + b*c*Log[x] - (b*c*Log[1 - c^2*x^6])/6`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {6452, 798, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{arctanh}(cx^3)}{x^4} dx \\
 & \quad \downarrow 6452 \\
 & bc \int \frac{1}{x(1-c^2x^6)} dx - \frac{a + \operatorname{arctanh}(cx^3)}{3x^3} \\
 & \quad \downarrow 798 \\
 & \frac{1}{6}bc \int \frac{1}{x^6(1-c^2x^6)} dx^6 - \frac{a + \operatorname{arctanh}(cx^3)}{3x^3} \\
 & \quad \downarrow 47 \\
 & \frac{1}{6}bc \left(c^2 \int \frac{1}{1-c^2x^6} dx^6 + \int \frac{1}{x^6} dx^6 \right) - \frac{a + \operatorname{arctanh}(cx^3)}{3x^3} \\
 & \quad \downarrow 14 \\
 & \frac{1}{6}bc \left(c^2 \int \frac{1}{1-c^2x^6} dx^6 + \log(x^6) \right) - \frac{a + \operatorname{arctanh}(cx^3)}{3x^3} \\
 & \quad \downarrow 16 \\
 & \frac{1}{6}bc (\log(x^6) - \log(1-c^2x^6)) - \frac{a + \operatorname{arctanh}(cx^3)}{3x^3}
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c*x^3])/x^4,x]`

output `-1/3*(a + b*ArcTanh[c*x^3])/x^3 + (b*c*(Log[x^6] - Log[1 - c^2*x^6]))/6`

Definitions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.18

method	result	size
default	$-\frac{a}{3x^3} + b\left(-\frac{\operatorname{arctanh}(cx^3)}{3x^3} + c\left(\ln(x) - \frac{\ln(cx^3-1)}{6} - \frac{\ln(cx^3+1)}{6}\right)\right)$	47
parts	$-\frac{a}{3x^3} + b\left(-\frac{\operatorname{arctanh}(cx^3)}{3x^3} + c\left(\ln(x) - \frac{\ln(cx^3-1)}{6} - \frac{\ln(cx^3+1)}{6}\right)\right)$	47
paralelrisch	$\frac{3bc\ln(x)x^3 - \ln(cx^3-1)x^3bc - b\operatorname{arctanh}(cx^3)x^3c - b\operatorname{arctanh}(cx^3) - a}{3x^3}$	56
risch	$-\frac{b\ln(cx^3+1)}{6x^3} + \frac{6bc\ln(x)x^3 - bc\ln(c^2x^6-1)x^3 + b\ln(-cx^3+1) - 2a}{6x^3}$	62

input `int((a+b*arctanh(c*x^3))/x^4,x,method=_RETURNVERBOSE)`

output

```
-1/3*a/x^3+b*(-1/3/x^3*arctanh(c*x^3)+c*(ln(x)-1/6*ln(c*x^3-1)-1/6*ln(c*x^3+1)))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.38

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^4} dx = -\frac{bcx^3 \log(c^2x^6 - 1) - 6bcx^3 \log(x) + b \log\left(-\frac{cx^3+1}{cx^3-1}\right) + 2a}{6x^3}$$

input

```
integrate((a+b*arctanh(c*x^3))/x^4,x, algorithm="fricas")
```

output

```
-1/6*(b*c*x^3*log(c^2*x^6 - 1) - 6*b*c*x^3*log(x) + b*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 2*a)/x^3
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^4} dx = \text{Timed out}$$

input

```
integrate((a+b*atanh(c*x**3))/x**4,x)
```

output

```
Timed out
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.02

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^4} dx = -\frac{1}{6} \left(c(\log(c^2x^6 - 1) - \log(x^6)) + \frac{2 \operatorname{artanh}(cx^3)}{x^3} \right) b - \frac{a}{3x^3}$$

input `integrate((a+b*arctanh(c*x^3))/x^4,x, algorithm="maxima")`output `-1/6*(c*(log(c^2*x^6 - 1) - log(x^6)) + 2*arctanh(c*x^3)/x^3)*b - 1/3*a/x^3`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.28

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^4} dx = -\frac{1}{6} bc \log(c^2x^6 - 1) + bc \log(x) - \frac{b \log\left(-\frac{cx^3+1}{cx^3-1}\right)}{6x^3} - \frac{a}{3x^3}$$

input `integrate((a+b*arctanh(c*x^3))/x^4,x, algorithm="giac")`output `-1/6*b*c*log(c^2*x^6 - 1) + b*c*log(x) - 1/6*b*log(-(c*x^3 + 1)/(c*x^3 - 1))/x^3 - 1/3*a/x^3`**Mupad [B] (verification not implemented)**

Time = 3.40 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.38

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^4} dx = bc \ln(x) - \frac{a}{3x^3} - \frac{bc \ln(c^2x^6 - 1)}{6} - \frac{b \ln(cx^3 + 1)}{6x^3} + \frac{b \ln(1 - cx^3)}{6x^3}$$

input `int((a + b*atanh(c*x^3))/x^4,x)`

output

```
b*c*log(x) - a/(3*x^3) - (b*c*log(c^2*x^6 - 1))/6 - (b*log(c*x^3 + 1))/(6*x^3) + (b*log(1 - c*x^3))/(6*x^3)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.98

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^4} dx$$

$$= \frac{\operatorname{atanh}(cx^3)bcx^3 - \operatorname{atanh}(cx^3)b - \log\left(c^{\frac{2}{3}}x^2 - c^{\frac{1}{3}}x + 1\right)bcx^3 - \log\left(c^{\frac{2}{3}}x + c^{\frac{1}{3}}\right)bcx^3 + 3\log(x)bcx^3 - a}{3x^3}$$

input

```
int((a+b*atanh(c*x^3))/x^4,x)
```

output

```
(atanh(c*x**3)*b*c*x**3 - atanh(c*x**3)*b - log(c**(2/3)*x**2 - c**(1/3)*x + 1)*b*c*x**3 - log(c**(2/3)*x + c**(1/3))*b*c*x**3 + 3*log(x)*b*c*x**3 - a)/(3*x**3)
```

3.105 $\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^7} dx$

Optimal result	890
Mathematica [A] (verified)	890
Rubi [A] (verified)	891
Maple [A] (verified)	892
Fricas [A] (verification not implemented)	893
Sympy [F(-1)]	893
Maxima [A] (verification not implemented)	894
Giac [A] (verification not implemented)	894
Mupad [B] (verification not implemented)	895
Reduce [B] (verification not implemented)	895

Optimal result

Integrand size = 14, antiderivative size = 41

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^7} dx = -\frac{bc}{6x^3} + \frac{1}{6}bc^2 \operatorname{arctanh}(cx^3) - \frac{a + b \operatorname{arctanh}(cx^3)}{6x^6}$$

output

```
-1/6*b*c/x^3+1/6*b*c^2*arctanh(c*x^3)-1/6*(a+b*arctanh(c*x^3))/x^6
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.59

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^7} dx = -\frac{a}{6x^6} - \frac{bc}{6x^3} - \frac{b \operatorname{arctanh}(cx^3)}{6x^6} - \frac{1}{12}bc^2 \log(1 - cx^3) + \frac{1}{12}bc^2 \log(1 + cx^3)$$

input

```
Integrate[(a + b*ArcTanh[c*x^3])/x^7, x]
```

output

```
-1/6*a/x^6 - (b*c)/(6*x^3) - (b*ArcTanh[c*x^3])/(6*x^6) - (b*c^2*Log[1 - c*x^3])/12 + (b*c^2*Log[1 + c*x^3])/12
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.93, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6452, 807, 264, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^7} dx$$

$$\downarrow 6452$$

$$\frac{1}{2}bc \int \frac{1}{x^4(1-c^2x^6)} dx - \frac{a + b \operatorname{arctanh}(cx^3)}{6x^6}$$

$$\downarrow 807$$

$$\frac{1}{6}bc \int \frac{1}{x^6(1-c^2x^6)} dx^3 - \frac{a + b \operatorname{arctanh}(cx^3)}{6x^6}$$

$$\downarrow 264$$

$$\frac{1}{6}bc \left(c^2 \int \frac{1}{1-c^2x^6} dx^3 - \frac{1}{x^3} \right) - \frac{a + b \operatorname{arctanh}(cx^3)}{6x^6}$$

$$\downarrow 219$$

$$\frac{1}{6}bc \left(\operatorname{arctanh}(cx^3) - \frac{1}{x^3} \right) - \frac{a + b \operatorname{arctanh}(cx^3)}{6x^6}$$

input `Int[(a + b*ArcTanh[c*x^3])/x^7,x]`

output `-1/6*(a + b*ArcTanh[c*x^3])/x^6 + (b*c*(-x^(-3) + c*ArcTanh[c*x^3]))/6`

Defintions of rubi rules used

rule 219 $\text{Int}[\{(a_)+(b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 264 $\text{Int}[\{(c_)*(x_)\}^{(m_)}*\{(a_)+(b_)*(x_)^2\}^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(c*x)^{(m+1)}*\{(a+b*x^2)^{(p+1)}/(a*c*(m+1))\}, x] - \text{Simp}[b*((m+2*p+3)/(a*c^{2*(m+1)})) \ \text{Int}[(c*x)^{(m+2)}*(a+b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 807 $\text{Int}[(x_)^{(m_)}*\{(a_)+(b_)*(x_)^{(n_)}\}^{(p_)}, x_Symbol] \rightarrow \text{With}\{k = \text{GCD}[m+1, n]\}, \text{Simp}[1/k \ \text{Subst}[\text{Int}[x^{(m+1)/k-1}*(a+b*x^{(n/k)})^p, x], x, x^k], x] /; k \neq 1 /; \text{FreeQ}\{a, b, p\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$

rule 6452 $\text{Int}[\{(a_)+\text{ArcTanh}[(c_)*(x_)^{(n_)}]\}*(b_)\}^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*\{(a+b*\text{ArcTanh}[c*x^n])^p/(m+1)\}, x] - \text{Simp}[b*c*n*(p/(m+1)) \ \text{Int}[x^{(m+n)}*\{(a+b*\text{ArcTanh}[c*x^n])^{(p-1)}/(1-c^2*x^{(2*n)})\}, x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.10

method	result	size
parallelrisch	$-\frac{\arctanh(cx^3)bc^2x^6+ac^2x^6+bcx^3+b\arctanh(cx^3)+a}{6x^6}$	45
default	$-\frac{a}{6x^6} - \frac{b\arctanh(cx^3)}{6x^6} - \frac{bc^2\ln(cx^3-1)}{12} - \frac{bc}{6x^3} + \frac{bc^2\ln(cx^3+1)}{12}$	55
parts	$-\frac{a}{6x^6} - \frac{b\arctanh(cx^3)}{6x^6} - \frac{bc^2\ln(cx^3-1)}{12} - \frac{bc}{6x^3} + \frac{bc^2\ln(cx^3+1)}{12}$	55
risch	$-\frac{b\ln(cx^3+1)}{12x^6} - \frac{bc^2\ln(cx^3-1)x^6-bc^2\ln(cx^3+1)x^6+2bcx^3-b\ln(-cx^3+1)+2a}{12x^6}$	77
orering	$\frac{(\frac{5}{9}c^2x^7-\frac{5}{9}x)(a+b\arctanh(cx^3))}{x^7} + \frac{(cx^3-1)(cx^3+1)x^2\left(\frac{3bc}{x^5(-c^2x^6+1)}-\frac{7(a+b\arctanh(cx^3))}{x^8}\right)}{18}$	82

input `int((a+b*arctanh(c*x^3))/x^7,x,method=_RETURNVERBOSE)`

output `-1/6*(-arctanh(c*x^3)*b*c^2*x^6+a*c^2*x^6+b*c*x^3+b*arctanh(c*x^3)+a)/x^6`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.20

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^7} dx = -\frac{2bcx^3 - (bc^2x^6 - b) \log\left(-\frac{cx^3+1}{cx^3-1}\right) + 2a}{12x^6}$$

input `integrate((a+b*arctanh(c*x^3))/x^7,x, algorithm="fricas")`

output `-1/12*(2*b*c*x^3 - (b*c^2*x^6 - b)*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 2*a)/x^6`

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^7} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**3))/x**7,x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.24

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^7} dx$$

$$= \frac{1}{12} \left(\left(c \log(cx^3 + 1) - c \log(cx^3 - 1) - \frac{2}{x^3} \right) c - \frac{2 \operatorname{artanh}(cx^3)}{x^6} \right) b - \frac{a}{6x^6}$$

input `integrate((a+b*arctanh(c*x^3))/x^7,x, algorithm="maxima")`output `1/12*((c*log(c*x^3 + 1) - c*log(c*x^3 - 1) - 2/x^3)*c - 2*arctanh(c*x^3)/x^6)*b - 1/6*a/x^6`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.63

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^7} dx = \frac{1}{12} bc^2 \log(cx^3 + 1) - \frac{1}{12} bc^2 \log(cx^3 - 1)$$

$$- \frac{b \log\left(-\frac{cx^3+1}{cx^3-1}\right)}{12x^6} - \frac{bcx^3 + a}{6x^6}$$

input `integrate((a+b*arctanh(c*x^3))/x^7,x, algorithm="giac")`output `1/12*b*c^2*log(c*x^3 + 1) - 1/12*b*c^2*log(c*x^3 - 1) - 1/12*b*log(-(c*x^3 + 1)/(c*x^3 - 1))/x^6 - 1/6*(b*c*x^3 + a)/x^6`

Mupad [B] (verification not implemented)

Time = 3.57 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.27

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^7} dx = \frac{b c^2 \operatorname{atanh}(c x^3)}{6} - \frac{a}{6} + \frac{b \ln(cx^3+1)}{12} - \frac{b \ln(1-cx^3)}{12} + \frac{b c x^3}{6}$$

input `int((a + b*atanh(c*x^3))/x^7,x)`output `(b*c^2*atanh(c*x^3))/6 - (a/6 + (b*log(c*x^3 + 1))/12 - (b*log(1 - c*x^3))/12 + (b*c*x^3)/6)/x^6`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.95

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^7} dx = \frac{\operatorname{atanh}(c x^3) b c^2 x^6 - \operatorname{atanh}(c x^3) b - a - b c x^3}{6 x^6}$$

input `int((a+b*atanh(c*x^3))/x^7,x)`output `(atanh(c*x**3)*b*c**2*x**6 - atanh(c*x**3)*b - a - b*c*x**3)/(6*x**6)`

3.106 $\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^{10}} dx$

Optimal result	896
Mathematica [A] (verified)	896
Rubi [A] (verified)	897
Maple [A] (verified)	898
Fricas [A] (verification not implemented)	899
Sympy [F(-1)]	899
Maxima [A] (verification not implemented)	900
Giac [A] (verification not implemented)	900
Mupad [B] (verification not implemented)	901
Reduce [B] (verification not implemented)	901

Optimal result

Integrand size = 14, antiderivative size = 56

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^{10}} dx = -\frac{bc}{18x^6} - \frac{a + b \operatorname{arctanh}(cx^3)}{9x^9} + \frac{1}{3}bc^3 \log(x) - \frac{1}{18}bc^3 \log(1 - c^2x^6)$$

output

```
-1/18*b*c/x^6-1/9*(a+b*arctanh(c*x^3))/x^9+1/3*b*c^3*ln(x)-1/18*b*c^3*ln(-c^2*x^6+1)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.09

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^{10}} dx = -\frac{a}{9x^9} - \frac{bc}{18x^6} - \frac{b \operatorname{arctanh}(cx^3)}{9x^9} + \frac{1}{3}bc^3 \log(x) - \frac{1}{18}bc^3 \log(1 - c^2x^6)$$

input

```
Integrate[(a + b*ArcTanh[c*x^3])/x^10, x]
```

output

$$-1/9*a/x^9 - (b*c)/(18*x^6) - (b*ArcTanh[c*x^3])/(9*x^9) + (b*c^3*Log[x])/3 - (b*c^3*Log[1 - c^2*x^6])/18$$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6452, 798, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + \operatorname{barctanh}(cx^3)}{x^{10}} dx \\ & \quad \downarrow \text{6452} \\ & \frac{1}{3}bc \int \frac{1}{x^7(1-c^2x^6)} dx - \frac{a + \operatorname{barctanh}(cx^3)}{9x^9} \\ & \quad \downarrow \text{798} \\ & \frac{1}{18}bc \int \frac{1}{x^{12}(1-c^2x^6)} dx^6 - \frac{a + \operatorname{barctanh}(cx^3)}{9x^9} \\ & \quad \downarrow \text{54} \\ & \frac{1}{18}bc \int \left(-\frac{c^4}{c^2x^6-1} + \frac{c^2}{x^6} + \frac{1}{x^{12}} \right) dx^6 - \frac{a + \operatorname{barctanh}(cx^3)}{9x^9} \\ & \quad \downarrow \text{2009} \\ & \frac{1}{18}bc \left(c^2 \log(x^6) - c^2 \log(1-c^2x^6) - \frac{1}{x^6} \right) - \frac{a + \operatorname{barctanh}(cx^3)}{9x^9} \end{aligned}$$

input

$$\text{Int}[(a + b*ArcTanh[c*x^3])/x^10, x]$$

output

$$-1/9*(a + b*ArcTanh[c*x^3])/x^9 + (b*c*(-x^(-6) + c^2*Log[x^6] - c^2*Log[1 - c^2*x^6]))/18$$

Defintions of rubi rules used

rule 54 $\text{Int}[(a + (b \cdot x)^m) \cdot ((c + (d \cdot x)^n)^p), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b \cdot x)^m \cdot (c + d \cdot x)^n, x], x] /;$ FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])

rule 798 $\text{Int}[(x)^m \cdot ((a + (b \cdot x)^n)^p), x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n) - 1}] \cdot (a + b \cdot x)^p, x], x, x^n], x] /;$ FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

rule 2009 $\text{Int}[u, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /;$ SumQ[u]

rule 6452 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n]) \cdot (b \cdot x)^m, x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot ((a + b \cdot \text{ArcTanh}[c \cdot x^n])^p / (m + 1)), x] - \text{Simp}[b \cdot c \cdot n \cdot (p / (m + 1)) \cdot \text{Int}[x^{m+n} \cdot ((a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2n}))], x] /;$ FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]

Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.12

method	result	size
default	$-\frac{a}{9x^9} + b \left(-\frac{\text{arctanh}(cx^3)}{9x^9} + \frac{c \left(-\frac{1}{6x^6} + c^2 \ln(x) - \frac{c^2 \ln(cx^3-1)}{6} - \frac{c^2 \ln(cx^3+1)}{6} \right)}{3} \right)$	63
parts	$-\frac{a}{9x^9} + b \left(-\frac{\text{arctanh}(cx^3)}{9x^9} + \frac{c \left(-\frac{1}{6x^6} + c^2 \ln(x) - \frac{c^2 \ln(cx^3-1)}{6} - \frac{c^2 \ln(cx^3+1)}{6} \right)}{3} \right)$	63
risch	$-\frac{b \ln(cx^3+1)}{18x^9} + \frac{6bc^3 \ln(x)x^9 - bc^3 \ln(c^2x^6-1)x^9 - bcx^3 + b \ln(-cx^3+1) - 2a}{18x^9}$	73
parallelrisch	$\frac{6bc^3 \ln(x)x^9 - 2 \ln(cx^3-1)x^9 bc^3 - 2b \text{arctanh}(cx^3)x^9 c^3 - bc^3 x^9 - bcx^3 - 2b \text{arctanh}(cx^3) - 2a}{18x^9}$	78

input $\text{int}((a+b \cdot \text{arctanh}(c \cdot x^3))/x^{10}, x, \text{method}=_RETURNVERBOSE)$

output

```
-1/9*a/x^9+b*(-1/9/x^9*arctanh(c*x^3)+1/3*c*(-1/6/x^6+c^2*ln(x)-1/6*c^2*ln
(c*x^3-1)-1/6*c^2*ln(c*x^3+1)))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.16

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^{10}} dx$$

$$= -\frac{bc^3x^9 \log(c^2x^6 - 1) - 6bc^3x^9 \log(x) + bcx^3 + b \log\left(-\frac{cx^3+1}{cx^3-1}\right) + 2a}{18x^9}$$

input

```
integrate((a+b*arctanh(c*x^3))/x^10,x, algorithm="fricas")
```

output

```
-1/18*(b*c^3*x^9*log(c^2*x^6 - 1) - 6*b*c^3*x^9*log(x) + b*c*x^3 + b*log(-
(c*x^3 + 1)/(c*x^3 - 1)) + 2*a)/x^9
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^{10}} dx = \text{Timed out}$$

input

```
integrate((a+b*atanh(c*x**3))/x**10,x)
```

output

```
Timed out
```


Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.91

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^{10}} dx$$

$$= -\frac{1}{18} \left(\left(c^2 \log(c^2 x^6 - 1) - c^2 \log(x^6) + \frac{1}{x^6} \right) c + \frac{2 \operatorname{artanh}(cx^3)}{x^9} \right) b - \frac{a}{9 x^9}$$

input `integrate((a+b*arctanh(c*x^3))/x^10,x, algorithm="maxima")`output `-1/18*((c^2*log(c^2*x^6 - 1) - c^2*log(x^6) + 1/x^6)*c + 2*arctanh(c*x^3)/x^9)*b - 1/9*a/x^9`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.16

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^{10}} dx = -\frac{1}{18} bc^3 \log(c^2 x^6 - 1) + \frac{1}{3} bc^3 \log(x)$$

$$- \frac{b \log\left(-\frac{cx^3+1}{cx^3-1}\right)}{18 x^9} - \frac{bcx^3 + 2a}{18 x^9}$$

input `integrate((a+b*arctanh(c*x^3))/x^10,x, algorithm="giac")`output `-1/18*b*c^3*log(c^2*x^6 - 1) + 1/3*b*c^3*log(x) - 1/18*b*log(-(c*x^3 + 1)/(c*x^3 - 1))/x^9 - 1/18*(b*c*x^3 + 2*a)/x^9`

Mupad [B] (verification not implemented)

Time = 3.44 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.20

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^{10}} dx = \frac{bc^3 \ln(x)}{3} - \frac{bc^3 \ln(c^2 x^6 - 1)}{18} - \frac{a}{9x^9} - \frac{bc}{18x^6} - \frac{b \ln(cx^3 + 1)}{18x^9} + \frac{b \ln(1 - cx^3)}{18x^9}$$

input `int((a + b*atanh(c*x^3))/x^10,x)`output `(b*c^3*log(x))/3 - (b*c^3*log(c^2*x^6 - 1))/18 - a/(9*x^9) - (b*c)/(18*x^6) - (b*log(c*x^3 + 1))/(18*x^9) + (b*log(1 - c*x^3))/(18*x^9)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.70

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^{10}} dx = \frac{2 \operatorname{atanh}(cx^3) b c^3 x^9 - 2 \operatorname{atanh}(cx^3) b - 2 \log\left(c^{\frac{2}{3}} x^2 - c^{\frac{1}{3}} x + 1\right) b c^3 x^9 - 2 \log\left(c^{\frac{2}{3}} x + c^{\frac{1}{3}}\right) b c^3 x^9 + 6 \log(x)}{18x^9}$$

input `int((a+b*atanh(c*x^3))/x^10,x)`output `(2*atanh(c*x**3)*b*c**3*x**9 - 2*atanh(c*x**3)*b - 2*log(c**(2/3)*x**2 - c**(1/3)*x + 1)*b*c**3*x**9 - 2*log(c**(2/3)*x + c**(1/3))*b*c**3*x**9 + 6*log(x)*b*c**3*x**9 - 2*a - b*c*x**3)/(18*x**9)`

3.107 $\int x^3(a + \operatorname{barctanh}(cx^3)) dx$

Optimal result	902
Mathematica [A] (verified)	903
Rubi [A] (verified)	903
Maple [A] (verified)	908
Fricas [B] (verification not implemented)	909
Sympy [F(-1)]	910
Maxima [A] (verification not implemented)	910
Giac [A] (verification not implemented)	911
Mupad [B] (verification not implemented)	911
Reduce [B] (verification not implemented)	912

Optimal result

Integrand size = 14, antiderivative size = 140

$$\int x^3(a + \operatorname{barctanh}(cx^3)) dx = \frac{3bx}{4c} + \frac{\sqrt{3}b \arctan\left(\frac{1-2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{8c^{4/3}} - \frac{\sqrt{3}b \arctan\left(\frac{1+2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{8c^{4/3}} - \frac{\operatorname{barctanh}(\sqrt[3]{cx})}{4c^{4/3}} + \frac{1}{4}x^4(a + \operatorname{barctanh}(cx^3)) - \frac{\operatorname{barctanh}\left(\frac{\sqrt[3]{Cx}}{1+c^{2/3}x^2}\right)}{8c^{4/3}}$$

output

```
3/4*b*x/c+1/8*3^(1/2)*b*arctan(1/3*(1-2*c^(1/3)*x)*3^(1/2))/c^(4/3)-1/8*3^(1/2)*b*arctan(1/3*(1+2*c^(1/3)*x)*3^(1/2))/c^(4/3)-1/4*b*arctanh(c^(1/3)*x)/c^(4/3)+1/4*x^4*(a+b*arctanh(c*x^3))-1/8*b*arctanh(c^(1/3)*x/(1+c^(2/3)*x^2))/c^(4/3)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.40

$$\int x^3(a + b \operatorname{arctanh}(cx^3)) dx = \frac{3bx}{4c} + \frac{ax^4}{4} - \frac{\sqrt{3}b \arctan\left(\frac{-1+2\sqrt[3]{cx}}{\sqrt{3}}\right)}{8c^{4/3}}$$

$$- \frac{\sqrt{3}b \arctan\left(\frac{1+2\sqrt[3]{cx}}{\sqrt{3}}\right)}{8c^{4/3}} + \frac{1}{4}bx^4 \operatorname{arctanh}(cx^3)$$

$$+ \frac{b \log(1 - \sqrt[3]{cx})}{8c^{4/3}} - \frac{b \log(1 + \sqrt[3]{cx})}{8c^{4/3}}$$

$$+ \frac{b \log(1 - \sqrt[3]{cx} + c^{2/3}x^2)}{16c^{4/3}} - \frac{b \log(1 + \sqrt[3]{cx} + c^{2/3}x^2)}{16c^{4/3}}$$

input `Integrate[x^3*(a + b*ArcTanh[c*x^3]),x]`

output `(3*b*x)/(4*c) + (a*x^4)/4 - (Sqrt[3]*b*ArcTan[(-1 + 2*c^(1/3)*x)/Sqrt[3]])/(8*c^(4/3)) - (Sqrt[3]*b*ArcTan[(1 + 2*c^(1/3)*x)/Sqrt[3]])/(8*c^(4/3)) + (b*x^4*ArcTanh[c*x^3])/4 + (b*Log[1 - c^(1/3)*x])/(8*c^(4/3)) - (b*Log[1 + c^(1/3)*x])/(8*c^(4/3)) + (b*Log[1 - c^(1/3)*x + c^(2/3)*x^2])/(16*c^(4/3)) - (b*Log[1 + c^(1/3)*x + c^(2/3)*x^2])/(16*c^(4/3))`

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.26, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.786$, Rules used = {6452, 843, 754, 27, 219, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3(a + b \operatorname{arctanh}(cx^3)) dx$$

$$\downarrow 6452$$

$$\frac{1}{4}x^4(a + b \operatorname{arctanh}(cx^3)) - \frac{3}{4}bc \int \frac{x^6}{1 - c^2x^6} dx$$

↓ 843

$$\frac{1}{4}x^4(a + \operatorname{barctanh}(cx^3)) - \frac{3}{4}bc \left(\frac{\int \frac{1}{1-c^2x^6} dx}{c^2} - \frac{x}{c^2} \right)$$

↓ 754

$$\frac{1}{4}x^4(a + \operatorname{barctanh}(cx^3)) - \frac{3}{4}bc \left(\frac{\frac{1}{3} \int \frac{1}{1-c^{2/3}x^2} dx + \frac{1}{3} \int \frac{2-\sqrt[3]{cx}}{2(c^{2/3}x^2-\sqrt[3]{cx+1})} dx + \frac{1}{3} \int \frac{\sqrt[3]{cx+2}}{2(c^{2/3}x^2+\sqrt[3]{cx+1})} dx}{c^2} - \frac{x}{c^2} \right)$$

↓ 27

$$\frac{1}{4}x^4(a + \operatorname{barctanh}(cx^3)) - \frac{3}{4}bc \left(\frac{\frac{1}{3} \int \frac{1}{1-c^{2/3}x^2} dx + \frac{1}{6} \int \frac{2-\sqrt[3]{cx}}{c^{2/3}x^2-\sqrt[3]{cx+1}} dx + \frac{1}{6} \int \frac{\sqrt[3]{cx+2}}{c^{2/3}x^2+\sqrt[3]{cx+1}} dx}{c^2} - \frac{x}{c^2} \right)$$

↓ 219

$$\frac{1}{4}x^4(a + \operatorname{barctanh}(cx^3)) - \frac{3}{4}bc \left(\frac{\frac{1}{6} \int \frac{2-\sqrt[3]{cx}}{c^{2/3}x^2-\sqrt[3]{cx+1}} dx + \frac{1}{6} \int \frac{\sqrt[3]{cx+2}}{c^{2/3}x^2+\sqrt[3]{cx+1}} dx + \frac{\operatorname{arctanh}(\sqrt[3]{cx})}{3\sqrt[3]{c}}}{c^2} - \frac{x}{c^2} \right)$$

↓ 1142

$$\frac{1}{4}x^4(a + \operatorname{barctanh}(cx^3)) - \frac{3}{4}bc \left(\frac{\frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x^2-\sqrt[3]{cx+1}} dx - \frac{\int \frac{\sqrt[3]{c}(1-2\sqrt[3]{cx})}{c^{2/3}x^2-\sqrt[3]{cx+1}} dx}{2\sqrt[3]{c}} \right) + \frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x^2+\sqrt[3]{cx+1}} dx + \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{cx+1})}{c^{2/3}x^2+\sqrt[3]{cx+1}} dx}{2\sqrt[3]{c}} \right)}{c^2} + \frac{\operatorname{arctanh}(\sqrt[3]{cx})}{3\sqrt[3]{c}} \right)$$

↓ 25

$$\frac{3}{4}bc \left(\frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx^3)) - \frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx + \frac{\int \frac{\sqrt[3]{c}(1-2\sqrt[3]{Cx})}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx}{2\sqrt[3]{c}} \right) + \frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx + \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{Cx+1})}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx}{2\sqrt[3]{c}} \right) + \frac{\operatorname{arctanh}\left(\frac{2\sqrt[3]{Cx+1}}{\sqrt[3]{c}}\right)}{3\sqrt[3]{c}}}{c^2} \right)$$

↓ 27

$$\frac{3}{4}bc \left(\frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx^3)) - \frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx + \frac{1}{2} \int \frac{1-2\sqrt[3]{Cx}}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx \right) + \frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx + \frac{1}{2} \int \frac{2\sqrt[3]{Cx+1}}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx \right) + \frac{\operatorname{arctanh}\left(\frac{2\sqrt[3]{Cx+1}}{\sqrt[3]{c}}\right)}{3\sqrt[3]{c}}}{c^2} \right)$$

↓ 1082

$$\frac{3}{4}bc \left(\frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx^3)) - \frac{1}{6} \left(\frac{1}{2} \int \frac{1-2\sqrt[3]{Cx}}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx + \frac{3 \int \frac{1}{(1-2\sqrt[3]{Cx})^2} d(1-2\sqrt[3]{Cx})}{\sqrt[3]{c}} \right) + \frac{1}{6} \left(\frac{1}{2} \int \frac{2\sqrt[3]{Cx+1}}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx - \frac{3 \int \frac{1}{(2\sqrt[3]{Cx+1})^2} d(2\sqrt[3]{Cx+1})}{\sqrt[3]{c}} \right) + \frac{\operatorname{arctanh}\left(\frac{2\sqrt[3]{Cx+1}}{\sqrt[3]{c}}\right)}{3\sqrt[3]{c}}}{c^2} \right)$$

↓ 217

$$\frac{3}{4}bc \left(\frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx^3)) - \frac{1}{6} \left(\frac{1}{2} \int \frac{1-2\sqrt[3]{Cx}}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx - \frac{\sqrt{3} \arctan\left(\frac{1-2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{\sqrt[3]{c}} \right) + \frac{1}{6} \left(\frac{1}{2} \int \frac{2\sqrt[3]{Cx+1}}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx + \frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{Cx+1}}{\sqrt{3}}\right)}{\sqrt[3]{c}} \right) + \frac{\operatorname{arctanh}\left(\frac{2\sqrt[3]{Cx+1}}{\sqrt[3]{c}}\right)}{3\sqrt[3]{c}}}{c^2} \right)$$

↓ 1103

$$\frac{1}{4}x^4(a + \operatorname{arctanh}(cx^3)) - \frac{3}{4}bc \left(\frac{\frac{1}{6} \left(-\frac{\sqrt{3} \arctan\left(\frac{1-2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{\log(c^{2/3}x^2 - \sqrt[3]{Cx+1})}{2\sqrt[3]{c}} \right)}{c^2} + \frac{1}{6} \left(\frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{Cx+1}}{\sqrt{3}}\right)}{\sqrt[3]{c}} + \frac{\log(c^{2/3}x^2 + \sqrt[3]{Cx+1})}{2\sqrt[3]{c}} \right) + \frac{\operatorname{arctanh}\left(\frac{1-2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{3\sqrt[3]{c}} \right)$$

input `Int[x^3*(a + b*ArcTanh[c*x^3]),x]`

output `(x^4*(a + b*ArcTanh[c*x^3]))/4 - (3*b*c*(-(x/c^2) + (ArcTanh[c^(1/3)*x]/(3*c^(1/3))) + (-((Sqrt[3]*ArcTan[(1 - 2*c^(1/3)*x]/Sqrt[3]))/c^(1/3)) - Log[1 - c^(1/3)*x + c^(2/3)*x^2]/(2*c^(1/3)))/6 + ((Sqrt[3]*ArcTan[(1 + 2*c^(1/3)*x]/Sqrt[3]))/c^(1/3) + Log[1 + c^(1/3)*x + c^(2/3)*x^2]/(2*c^(1/3)))/6)/c^2)/4`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 754

```
Int[((a_) + (b_.)*(x_)^(n_))^(n_)*(-1), x_Symbol] := Module[{r = Numerator[Rt[-a/b, n]], s = Denominator[Rt[-a/b, n]], k, u}, Simp[u = Int[(r - s*Cos[(2*k*Pi)/n]*x)/(r^2 - 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x] + Int[(r + s*Cos[(2*k*Pi)/n]*x)/(r^2 + 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x]; 2*(r^2/(a*n)) Int[1/(r^2 - s^2*x^2), x] + 2*(r/(a*n)) Sum[u, {k, 1, (n - 2)/4}], x] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && NegQ[a/b]
```

rule 843

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Simp[a*c^n*(m - n + 1)/(b*(m + n*p + 1)) Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

rule 1082

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]
```

rule 1103

```
Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

rule 1142

```
Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]
```

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c^n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]
```


Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.31

method	result
default	$\frac{x^4 a}{4} + \frac{b x^4 \operatorname{arctanh}(c x^3)}{4} + \frac{3 b x}{4 c} + \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{8 c^2 \left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{b \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}} x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{16 c^2 \left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{b \sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2 x^{\frac{1}{3}} + 1\right)}{\left(\frac{1}{c}\right)^{\frac{1}{3}}}\right)}{8 c^2 \left(\frac{1}{c}\right)^{\frac{2}{3}}} - b \ln\left(\frac{1}{c}\right)$
parts	$\frac{x^4 a}{4} + \frac{b x^4 \operatorname{arctanh}(c x^3)}{4} + \frac{3 b x}{4 c} + \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{8 c^2 \left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{b \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}} x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{16 c^2 \left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{b \sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2 x^{\frac{1}{3}} + 1\right)}{\left(\frac{1}{c}\right)^{\frac{1}{3}}}\right)}{8 c^2 \left(\frac{1}{c}\right)^{\frac{2}{3}}} - b \ln\left(\frac{1}{c}\right)$
risch	$\frac{b x^4 \ln(c x^3 + 1)}{8} + \frac{x^4 a}{4} - \frac{b x^4 \ln(-c x^3 + 1)}{8} + \frac{3 b x}{4 c} + \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{8 c^2 \left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{b \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}} x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{16 c^2 \left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{b \sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2 x^{\frac{1}{3}} + 1\right)}{\left(\frac{1}{c}\right)^{\frac{1}{3}}}\right)}{8 c^2 \left(\frac{1}{c}\right)^{\frac{2}{3}}} - b \ln\left(\frac{1}{c}\right)$

```
input int(x^3*(a+b*arctanh(c*x^3)),x,method=_RETURNVERBOSE)
```

```
output 1/4*x^4*a+1/4*b*x^4*arctanh(c*x^3)+3/4*b*x/c+1/8*b/c^2/(1/c)^(2/3)*ln(x-(1/c)^(1/3))-1/16*b/c^2/(1/c)^(2/3)*ln(x^2+(1/c)^(1/3)*x+(1/c)^(2/3))-1/8*b/c^2/(1/c)^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x+1))-1/8*b/c^2/(1/c)^(2/3)*ln(x+(1/c)^(1/3))+1/16*b/c^2/(1/c)^(2/3)*ln(x^2-(1/c)^(1/3)*x+(1/c)^(2/3))-1/8*b/c^2/(1/c)^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x-1))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 211 vs. $2(104) = 208$.

Time = 0.11 (sec) , antiderivative size = 981, normalized size of antiderivative = 7.01

$$\int x^3(a + b \operatorname{arctanh}(cx^3)) dx = \text{Too large to display}$$

input `integrate(x^3*(a+b*arctanh(c*x^3)),x, algorithm="fricas")`

output

```
[1/16*(2*b*c^2*x^4*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 4*a*c^2*x^4 + sqrt(3)*b*c*sqrt((-c)^(1/3)/c)*log((2*c*x^3 - sqrt(3)*(2*c*x^2 + (-c)^(2/3)*x + (-c)^(1/3)))*sqrt((-c)^(1/3)/c) + 3*(-c)^(1/3)*x - 1)/(c*x^3 + 1)) + sqrt(3)*b*c*sqrt(-1/c^(2/3))*log((2*c*x^3 - sqrt(3)*(2*c*x^2 - c^(2/3)*x - c^(1/3)))*sqrt(-1/c^(2/3)) - 3*c^(1/3)*x + 1)/(c*x^3 - 1)) + 12*b*c*x + b*(-c)^(2/3)*log(c*x^2 - (-c)^(2/3)*x - (-c)^(1/3)) - b*c^(2/3)*log(c*x^2 + c^(2/3)*x + c^(1/3)) - 2*b*(-c)^(2/3)*log(c*x + (-c)^(2/3)) + 2*b*c^(2/3)*log(c*x - c^(2/3)))/c^2, 1/16*(2*b*c^2*x^4*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 4*a*c^2*x^4 - 2*sqrt(3)*b*c*sqrt(-(-c)^(1/3)/c)*arctan(1/3*sqrt(3)*(2*(-c)^(2/3)*x + (-c)^(1/3))*sqrt(-(-c)^(1/3)/c)) + sqrt(3)*b*c*sqrt(-1/c^(2/3))*log((2*c*x^3 - sqrt(3)*(2*c*x^2 - c^(2/3)*x - c^(1/3)))*sqrt(-1/c^(2/3)) - 3*c^(1/3)*x + 1)/(c*x^3 - 1)) + 12*b*c*x + b*(-c)^(2/3)*log(c*x^2 - (-c)^(2/3)*x - (-c)^(1/3)) - b*c^(2/3)*log(c*x^2 + c^(2/3)*x + c^(1/3)) - 2*b*(-c)^(2/3)*log(c*x + (-c)^(2/3)) + 2*b*c^(2/3)*log(c*x - c^(2/3)))/c^2, 1/16*(2*b*c^2*x^4*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 4*a*c^2*x^4 + sqrt(3)*b*c*sqrt((-c)^(1/3)/c)*log((2*c*x^3 - sqrt(3)*(2*c*x^2 + (-c)^(2/3)*x + (-c)^(1/3)))*sqrt((-c)^(1/3)/c) + 3*(-c)^(1/3)*x - 1)/(c*x^3 + 1)) - 2*sqrt(3)*b*c^(2/3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*x + c^(1/3))/c^(1/3)) + 12*b*c*x + b*(-c)^(2/3)*log(c*x^2 - (-c)^(2/3)*x - (-c)^(1/3)) - b*c^(2/3)*log(c*x^2 + c^(2/3)*x + c^(1/3)) - 2*b*(-c)^(2/3)*log(c*x + (-c)^(2/3)) + 2*b*c^(2/3)*log(c...
```

Sympy [F(-1)]

Timed out.

$$\int x^3(a + b \operatorname{arctanh}(cx^3)) dx = \text{Timed out}$$

input `integrate(x**3*(a+b*atanh(c*x**3)),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.16

$$\int x^3(a + b \operatorname{arctanh}(cx^3)) dx = \frac{1}{4} ax^4 + \frac{1}{16} \left(4x^4 \operatorname{arctanh}(cx^3) - c \left(\frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{\frac{2}{3}}x + c^{\frac{1}{3}})}{3c^{\frac{1}{3}}}\right)}{c^{\frac{7}{3}}} + \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{\frac{2}{3}}x - c^{\frac{1}{3}})}{3c^{\frac{1}{3}}}\right)}{c^{\frac{7}{3}}} - \frac{12x}{c^2} + \log\left(\frac{c^{\frac{2}{3}}x + c^{\frac{1}{3}}}{c^{\frac{1}{3}}}\right) - \log\left(\frac{c^{\frac{2}{3}}x - c^{\frac{1}{3}}}{c^{\frac{1}{3}}}\right) + 2\log\left(\frac{c^{\frac{1}{3}}x + 1}{c^{\frac{1}{3}}}\right) - 2\log\left(\frac{c^{\frac{1}{3}}x - 1}{c^{\frac{1}{3}}}\right) \right) \right)$$

input `integrate(x^3*(a+b*arctanh(c*x^3)),x, algorithm="maxima")`

output `1/4*a*x^4 + 1/16*(4*x^4*arctanh(c*x^3) - c*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*x + c^(1/3))/c^(1/3))/c^(7/3) + 2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*x - c^(1/3))/c^(1/3))/c^(7/3) - 12*x/c^2 + log(c^(2/3)*x^2 + c^(1/3)*x + 1)/c^(7/3) - log(c^(2/3)*x^2 - c^(1/3)*x + 1)/c^(7/3) + 2*log((c^(1/3)*x + 1)/c^(1/3))/c^(7/3) - 2*log((c^(1/3)*x - 1)/c^(1/3))/c^(7/3))*b`

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.48

$$\int x^3(a + b \operatorname{arctanh}(cx^3)) dx$$

$$= \frac{1}{16} bc^7 \left(\frac{2(-\frac{1}{c})^{\frac{1}{3}} \log\left(\left|x - (-\frac{1}{c})^{\frac{1}{3}}\right|\right)}{c^8} - \frac{2\sqrt{3}|c|^{\frac{2}{3}} \arctan\left(\frac{1}{3}\sqrt{3}c^{\frac{1}{3}}\left(2x + \frac{1}{c^{\frac{1}{3}}}\right)\right)}{c^9} - \frac{2\sqrt{3}(-c^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}}{2}\left(2x + \frac{1}{c^{\frac{1}{3}}}\right)\right)}{c^9} \right.$$

$$\left. + \frac{1}{8} bx^4 \log\left(-\frac{cx^3 + 1}{cx^3 - 1}\right) + \frac{1}{4} ax^4 + \frac{3bx}{4c} \right)$$

input `integrate(x^3*(a+b*arctanh(c*x^3)),x, algorithm="giac")`

output

```
1/16*b*c^7*(2*(-1/c)^(1/3)*log(abs(x - (-1/c)^(1/3)))/c^8 - 2*sqrt(3)*abs(c)^(2/3)*arctan(1/3*sqrt(3)*c^(1/3)*(2*x + 1/c^(1/3)))/c^9 - 2*sqrt(3)*(-c^2)^(1/3)*arctan(1/3*sqrt(3)*(2*x + (-1/c)^(1/3))/(-1/c)^(1/3))/c^9 - abs(c)^(2/3)*log(x^2 + x/c^(1/3) + 1/c^(2/3))/c^9 + 2*log(abs(x - 1/c^(1/3)))/c^(25/3) - (-c^2)^(1/3)*log(x^2 + x*(-1/c)^(1/3) + (-1/c)^(2/3))/c^9) + 1/8*b*x^4*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 1/4*a*x^4 + 3/4*b*x/c
```

Mupad [B] (verification not implemented)

Time = 4.00 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.89

$$\int x^3(a + b \operatorname{arctanh}(cx^3)) dx$$

$$= \frac{ax^4}{4} + \frac{b \left(-\frac{\operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}-i)}{2}\right)}{2} + \frac{\operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}+i)}{2}\right)}{2} + \operatorname{atan}(c^{1/3}x \operatorname{li}) \right) \operatorname{li}}{4c^{4/3}}$$

$$+ \frac{3bx}{4c} + \frac{bx^4 \ln(cx^3 + 1)}{8} - \frac{bx^4 \ln(1 - cx^3)}{8}$$

$$- \frac{\sqrt{3}b \left(\operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}-i)}{2}\right) + \operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}+i)}{2}\right) \right)}{8c^{4/3}}$$

input `int(x^3*(a + b*atanh(c*x^3)),x)`

output $(a*x^4)/4 + (b*(atan((c^{1/3}*x*(3^{1/2} + 1i))/2)/2 - atan((c^{1/3}*x*(3^{1/2} - 1i))/2)/2 + atan(c^{1/3}*x*1i))*1i)/(4*c^{4/3}) + (3*b*x)/(4*c) + (b*x^4*\log(c*x^3 + 1))/8 - (b*x^4*\log(1 - c*x^3))/8 - (3^{1/2}*b*(atan((c^{1/3}*x*(3^{1/2} - 1i))/2) + atan((c^{1/3}*x*(3^{1/2} + 1i))/2)))/(8*c^{4/3}))$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.80

$$\int x^3(a + \operatorname{arctanh}(cx^3)) dx$$

$$= \frac{-2\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x-1}{\sqrt{3}}\right) b - 2\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x+1}{\sqrt{3}}\right) b + 4c^{\frac{4}{3}} \operatorname{atanh}(cx^3) b x^4 + 2\operatorname{atanh}(cx^3) b + 4c^{\frac{4}{3}} a x^4 + 12c^{\frac{1}{3}} b}{16c^{\frac{4}{3}}}$$

input `int(x^3*(a+b*atanh(c*x^3)),x)`

output $(- 2*\sqrt{3}*atan((2*c**(1/3)*x - 1)/\sqrt{3})*b - 2*\sqrt{3}*atan((2*c**(1/3)*x + 1)/\sqrt{3})*b + 4*c**(1/3)*atanh(c*x**3)*b*c*x**4 + 2*atanh(c*x**3)*b + 4*c**(1/3)*a*c*x**4 + 12*c**(1/3)*b*x - 3*\log(c**(2/3)*x + c**(1/3))*b + 3*\log(c**(2/3)*x - c**(1/3))*b)/(16*c**(1/3)*c)$

3.108 $\int (a + b \operatorname{arctanh}(cx^3)) dx$

Optimal result	913
Mathematica [A] (verified)	913
Rubi [A] (verified)	914
Maple [A] (verified)	915
Fricas [A] (verification not implemented)	916
Sympy [F(-1)]	916
Maxima [A] (verification not implemented)	917
Giac [A] (verification not implemented)	917
Mupad [B] (verification not implemented)	918
Reduce [B] (verification not implemented)	918

Optimal result

Integrand size = 10, antiderivative size = 101

$$\int (a + b \operatorname{arctanh}(cx^3)) dx = ax + \frac{\sqrt{3}b \arctan\left(\frac{1+2c^{2/3}x^2}{\sqrt{3}}\right)}{2\sqrt[3]{c}} + b \operatorname{arctanh}(cx^3) + \frac{b \log(1 - c^{2/3}x^2)}{2\sqrt[3]{c}} - \frac{b \log(1 + c^{2/3}x^2 + c^{4/3}x^4)}{4\sqrt[3]{c}}$$

output

```
a*x+1/2*3^(1/2)*b*arctan(1/3*(1+2*c^(2/3)*x^2)*3^(1/2))/c^(1/3)+b*x*arctan
h(c*x^3)+1/2*b*ln(1-c^(2/3)*x^2)/c^(1/3)-1/4*b*ln(1+c^(2/3)*x^2+c^(4/3)*x^
4)/c^(1/3)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.35

$$\int (a + b \operatorname{arctanh}(cx^3)) dx = ax + b \operatorname{arctanh}(cx^3) + \frac{b \left(-2\sqrt{3} \arctan\left(\frac{-1+2\sqrt[3]{c}x}{\sqrt{3}}\right) + 2\sqrt{3} \arctan\left(\frac{1+2\sqrt[3]{c}x}{\sqrt{3}}\right) - 2 \log(1 - \sqrt[3]{c}x) - 2 \log(1 + \sqrt[3]{c}x) + \log(1 - 4\sqrt[3]{c}x^4) \right)}{4\sqrt[3]{c}}$$

input `Integrate[a + b*ArcTanh[c*x^3], x]`

output `a*x + b*x*ArcTanh[c*x^3] - (b*(-2*Sqrt[3]*ArcTan[(-1 + 2*c^(1/3)*x)/Sqrt[3]] + 2*Sqrt[3]*ArcTan[(1 + 2*c^(1/3)*x)/Sqrt[3]] - 2*Log[1 - c^(1/3)*x] - 2*Log[1 + c^(1/3)*x] + Log[1 - c^(1/3)*x + c^(2/3)*x^2] + Log[1 + c^(1/3)*x + c^(2/3)*x^2]))/(4*c^(1/3))`

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{arctanh}(cx^3)) dx$$

$$\downarrow \text{2009}$$

$$ax + \frac{\sqrt{3}b \operatorname{arctan}\left(\frac{2c^{2/3}x^2+1}{\sqrt{3}}\right)}{2\sqrt[3]{c}} + b \operatorname{arctanh}(cx^3) + \frac{b \log(1 - c^{2/3}x^2)}{2\sqrt[3]{c}} - \frac{b \log(c^{4/3}x^4 + c^{2/3}x^2 + 1)}{4\sqrt[3]{c}}$$

input `Int[a + b*ArcTanh[c*x^3], x]`

output `a*x + (Sqrt[3]*b*ArcTan[(1 + 2*c^(2/3)*x^2)/Sqrt[3]])/(2*c^(1/3)) + b*x*ArcTanh[c*x^3] + (b*Log[1 - c^(2/3)*x^2])/(2*c^(1/3)) - (b*Log[1 + c^(2/3)*x^2 + c^(4/3)*x^4])/(4*c^(1/3))`

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.98

method	result
default	$ax + bx \operatorname{arctanh}(cx^3) + \frac{b \ln\left(x^2 - \left(\frac{1}{c^2}\right)^{\frac{1}{3}}\right)}{2c\left(\frac{1}{c^2}\right)^{\frac{1}{3}}} - \frac{b \ln\left(x^4 + \left(\frac{1}{c^2}\right)^{\frac{1}{3}}x^2 + \left(\frac{1}{c^2}\right)^{\frac{2}{3}}\right)}{4c\left(\frac{1}{c^2}\right)^{\frac{1}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2x^2 - \left(\frac{1}{c^2}\right)^{\frac{1}{3}} + 1\right)}{3}\right)}{2c\left(\frac{1}{c^2}\right)^{\frac{1}{3}}}$
parts	$ax + bx \operatorname{arctanh}(cx^3) + \frac{b \ln\left(x^2 - \left(\frac{1}{c^2}\right)^{\frac{1}{3}}\right)}{2c\left(\frac{1}{c^2}\right)^{\frac{1}{3}}} - \frac{b \ln\left(x^4 + \left(\frac{1}{c^2}\right)^{\frac{1}{3}}x^2 + \left(\frac{1}{c^2}\right)^{\frac{2}{3}}\right)}{4c\left(\frac{1}{c^2}\right)^{\frac{1}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2x^2 - \left(\frac{1}{c^2}\right)^{\frac{1}{3}} + 1\right)}{3}\right)}{2c\left(\frac{1}{c^2}\right)^{\frac{1}{3}}}$
risch	$ax + \frac{bx \ln(cx^3+1)}{2} + \frac{b \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{2c\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{b \ln\left(x^2 - \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{4c\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(-2x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{3\left(\frac{1}{c}\right)^{\frac{1}{3}}}\right)}{2c\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{bx \ln(-cx^3+1)}{2}$

```
input int(a+b*arctanh(c*x^3),x,method=_RETURNVERBOSE)
```

```
output a*x+b*x*arctanh(c*x^3)+1/2*b/c/(1/c^2)^(1/3)*ln(x^2-(1/c^2)^(1/3))-1/4*b/c/(1/c^2)^(1/3)*ln(x^4+(1/c^2)^(1/3)*x^2+(1/c^2)^(2/3))+1/2*b*3^(1/2)/c/(1/c^2)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c^2)^(1/3)*x^2+1))
```


Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 260, normalized size of antiderivative = 2.57

$$\int (a + \operatorname{barctanh}(cx^3)) dx$$

$$= \frac{\sqrt{3}bc\sqrt{-\frac{1}{c^{\frac{2}{3}}}} \log\left(\frac{2c^2x^6 - 3c^{\frac{2}{3}}x^2 + \sqrt{3}(2c^{\frac{5}{3}}x^4 - cx^2 - c^{\frac{1}{3}})\sqrt{-\frac{1}{c^{\frac{2}{3}}}} + 1}{c^2x^6 - 1}\right) + 2bcx \log\left(-\frac{cx^3 + 1}{cx^3 - 1}\right) + 4acx - bc^{\frac{2}{3}} \log(c^2x^2 + c^{\frac{2}{3}}) + 2bc^{\frac{2}{3}} \log(cx^2 - c^{\frac{1}{3}})}{4c}$$

input `integrate(a+b*arctanh(c*x^3),x, algorithm="fricas")`output `[1/4*(sqrt(3)*b*c*sqrt(-1/c^(2/3))*log((2*c^2*x^6 - 3*c^(2/3)*x^2 + sqrt(3)*(2*c^(5/3)*x^4 - c*x^2 - c^(1/3))*sqrt(-1/c^(2/3)) + 1)/(c^2*x^6 - 1)) + 2*b*c*x*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 4*a*c*x - b*c^(2/3)*log(c^2*x^4 + c^(4/3)*x^2 + c^(2/3)) + 2*b*c^(2/3)*log(c*x^2 - c^(1/3)))/c, 1/4*(2*b*c*x*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 2*sqrt(3)*b*c^(2/3)*arctan(1/3*sqrt(3)*(2*c*x^2 + c^(1/3))/c^(1/3)) + 4*a*c*x - b*c^(2/3)*log(c^2*x^4 + c^(4/3)*x^2 + c^(2/3)) + 2*b*c^(2/3)*log(c*x^2 - c^(1/3)))/c]`**Sympy [F(-1)]**

Timed out.

$$\int (a + \operatorname{barctanh}(cx^3)) dx = \text{Timed out}$$

input `integrate(a+b*atanh(c*x**3),x)`output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.89

$$\int (a + b \operatorname{arctanh}(cx^3)) dx$$

$$= \frac{1}{4} \left(c \left(\frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{\frac{4}{3}}x^2 + c^{\frac{2}{3}})}{3c^{\frac{2}{3}}}\right)}{c^{\frac{4}{3}}} - \frac{\log\left(c^{\frac{4}{3}}x^4 + c^{\frac{2}{3}}x^2 + 1\right)}{c^{\frac{4}{3}}} + \frac{2 \log\left(\frac{c^{\frac{2}{3}}x^2 - 1}{c^{\frac{2}{3}}}\right)}{c^{\frac{4}{3}}}\right) + 4x \operatorname{artanh}(cx^3) \right) + ax$$

input `integrate(a+b*arctanh(c*x^3),x, algorithm="maxima")`output `1/4*(c*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(4/3)*x^2 + c^(2/3))/c^(2/3))/c^(4/3) - log(c^(4/3)*x^4 + c^(2/3)*x^2 + 1)/c^(4/3) + 2*log((c^(2/3)*x^2 - 1)/c^(2/3))/c^(4/3)) + 4*x*arctanh(c*x^3)*b + a*x`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.08

$$\int (a + b \operatorname{arctanh}(cx^3)) dx$$

$$= \frac{1}{4} \left(c \left(\frac{2\sqrt{3}|c|^{\frac{2}{3}} \arctan\left(\frac{1}{3}\sqrt{3}\left(2x^2 + \frac{1}{|c|^{\frac{2}{3}}}\right)|c|^{\frac{2}{3}}\right)}{c^2} - \frac{|c|^{\frac{2}{3}} \log\left(x^4 + \frac{x^2}{|c|^{\frac{2}{3}}} + \frac{1}{|c|^{\frac{4}{3}}}\right)}{c^2} + \frac{2 \log\left(\left|x^2 - \frac{1}{|c|^{\frac{2}{3}}}\right|\right)}{|c|^{\frac{4}{3}}}\right) \right) + ax$$

input `integrate(a+b*arctanh(c*x^3),x, algorithm="giac")`output `1/4*(c*(2*sqrt(3)*abs(c)^(2/3)*arctan(1/3*sqrt(3)*(2*x^2 + 1/abs(c)^(2/3))*abs(c)^(2/3))/c^2 - abs(c)^(2/3)*log(x^4 + x^2/abs(c)^(2/3) + 1/abs(c)^(4/3))/c^2 + 2*log(abs(x^2 - 1/abs(c)^(2/3)))/abs(c)^(4/3) + 2*x*log(-(c*x^3 + 1)/(c*x^3 - 1)))*b + a*x`

Mupad [B] (verification not implemented)

Time = 5.54 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.06

$$\int (a + b \operatorname{arctanh}(cx^3)) dx = ax + \frac{b \ln(c^{2/3}x^2 - 1)}{2c^{1/3}} - \frac{\ln(4c^{2/3}x^2 + 2 - \sqrt{3}2i)(b + \sqrt{3}bi)}{4c^{1/3}} - \frac{\ln(4c^{2/3}x^2 + 2 + \sqrt{3}2i)(b - \sqrt{3}bi)}{4c^{1/3}} + \frac{bx \ln(cx^3 + 1)}{2} - \frac{bx \ln(1 - cx^3)}{2}$$

input `int(a + b*atanh(c*x^3),x)`output `a*x + (b*log(c^(2/3)*x^2 - 1))/(2*c^(1/3)) - (log(4*c^(2/3)*x^2 - 3^(1/2)*2i + 2)*(b + 3^(1/2)*b*1i))/(4*c^(1/3)) - (log(3^(1/2)*2i + 4*c^(2/3)*x^2 + 2)*(b - 3^(1/2)*b*1i))/(4*c^(1/3)) + (b*x*log(c*x^3 + 1))/2 - (b*x*log(1 - c*x^3))/2`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.18

$$\int (a + b \operatorname{arctanh}(cx^3)) dx = \frac{2\sqrt{3} \operatorname{atan}\left(\frac{2c^{1/3}x-1}{\sqrt{3}}\right) b - 2\sqrt{3} \operatorname{atan}\left(\frac{2c^{1/3}x+1}{\sqrt{3}}\right) b + 4c^{1/3} \operatorname{atanh}(cx^3) bx + 2 \operatorname{atanh}(cx^3) b + 4c^{1/3} ax - 2 \log\left(c^{2/3}x^2\right)}{4c^{1/3}}$$

input `int(a+b*atanh(c*x^3),x)`output `(2*sqrt(3)*atan((2*c**(1/3)*x - 1)/sqrt(3))*b - 2*sqrt(3)*atan((2*c**(1/3)*x + 1)/sqrt(3))*b + 4*c**(1/3)*atanh(c*x**3)*b*x + 2*atanh(c*x**3)*b + 4*c**(1/3)*a*x - 2*log(c**(2/3)*x**2 - c**(1/3)*x + 1)*b + log(c**(2/3)*x + c**(1/3))*b + 3*log(c**(2/3)*x - c**(1/3))*b)/(4*c**(1/3))`

3.109 $\int \frac{a+b\operatorname{arctanh}(cx^3)}{x^3} dx$

Optimal result	919
Mathematica [A] (verified)	920
Rubi [A] (verified)	920
Maple [A] (verified)	924
Fricas [B] (verification not implemented)	925
Sympy [F(-1)]	925
Maxima [A] (verification not implemented)	926
Giac [A] (verification not implemented)	926
Mupad [B] (verification not implemented)	927
Reduce [B] (verification not implemented)	928

Optimal result

Integrand size = 14, antiderivative size = 131

$$\int \frac{a + b\operatorname{arctanh}(cx^3)}{x^3} dx$$

$$= -\frac{1}{4}\sqrt{3}bc^{2/3} \arctan\left(\frac{1-2\sqrt[3]{cx}}{\sqrt{3}}\right) + \frac{1}{4}\sqrt{3}bc^{2/3} \arctan\left(\frac{1+2\sqrt[3]{cx}}{\sqrt{3}}\right)$$

$$+ \frac{1}{2}bc^{2/3}\operatorname{arctanh}(\sqrt[3]{cx}) - \frac{a + b\operatorname{arctanh}(cx^3)}{2x^2} + \frac{1}{4}bc^{2/3}\operatorname{arctanh}\left(\frac{\sqrt[3]{cx}}{1+c^{2/3}x^2}\right)$$

output

```
-1/4*3^(1/2)*b*c^(2/3)*arctan(1/3*(1-2*c^(1/3)*x)*3^(1/2))+1/4*3^(1/2)*b*c^(2/3)*arctan(1/3*(1+2*c^(1/3)*x)*3^(1/2))+1/2*b*c^(2/3)*arctanh(c^(1/3)*x)-1/2*(a+b*arctanh(c*x^3))/x^2+1/4*b*c^(2/3)*arctanh(c^(1/3)*x/(1+c^(2/3)*x^2))
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.43

$$\int \frac{a + \operatorname{barctanh}(cx^3)}{x^3} dx = -\frac{a}{2x^2} + \frac{1}{4}\sqrt{3}bc^{2/3} \arctan\left(\frac{-1 + 2\sqrt[3]{cx}}{\sqrt{3}}\right) + \frac{1}{4}\sqrt{3}bc^{2/3} \arctan\left(\frac{1 + 2\sqrt[3]{cx}}{\sqrt{3}}\right) - \frac{\operatorname{barctanh}(cx^3)}{2x^2} - \frac{1}{4}bc^{2/3} \log(1 - \sqrt[3]{cx}) + \frac{1}{4}bc^{2/3} \log(1 + \sqrt[3]{cx}) - \frac{1}{8}bc^{2/3} \log(1 - \sqrt[3]{cx} + c^{2/3}x^2) + \frac{1}{8}bc^{2/3} \log(1 + \sqrt[3]{cx} + c^{2/3}x^2)$$

input

```
Integrate[(a + b*ArcTanh[c*x^3])/x^3, x]
```

output

```
-1/2*a/x^2 + (Sqrt[3]*b*c^(2/3)*ArcTan[(-1 + 2*c^(1/3)*x)/Sqrt[3]])/4 + (Sqrt[3]*b*c^(2/3)*ArcTan[(1 + 2*c^(1/3)*x)/Sqrt[3]])/4 - (b*ArcTanh[c*x^3])/(2*x^2) - (b*c^(2/3)*Log[1 - c^(1/3)*x])/4 + (b*c^(2/3)*Log[1 + c^(1/3)*x])/4 - (b*c^(2/3)*Log[1 - c^(1/3)*x + c^(2/3)*x^2])/8 + (b*c^(2/3)*Log[1 + c^(1/3)*x + c^(2/3)*x^2])/8
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.27, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {6452, 754, 27, 219, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{barctanh}(cx^3)}{x^3} dx$$

↓ 6452

$$\frac{3}{2}bc \int \frac{1}{1 - c^2x^6} dx - \frac{a + \operatorname{barctanh}(cx^3)}{2x^2}$$

↓ 754

$$\frac{3}{2}bc \left(\frac{1}{3} \int \frac{1}{1 - c^{2/3}x^2} dx + \frac{1}{3} \int \frac{2 - \sqrt[3]{cx}}{2(c^{2/3}x^2 - \sqrt[3]{cx} + 1)} dx + \frac{1}{3} \int \frac{\sqrt[3]{cx} + 2}{2(c^{2/3}x^2 + \sqrt[3]{cx} + 1)} dx \right) - \frac{a + \operatorname{barctanh}(cx^3)}{2x^2}$$

↓ 27

$$\frac{3}{2}bc \left(\frac{1}{3} \int \frac{1}{1 - c^{2/3}x^2} dx + \frac{1}{6} \int \frac{2 - \sqrt[3]{cx}}{c^{2/3}x^2 - \sqrt[3]{cx} + 1} dx + \frac{1}{6} \int \frac{\sqrt[3]{cx} + 2}{c^{2/3}x^2 + \sqrt[3]{cx} + 1} dx \right) - \frac{a + \operatorname{barctanh}(cx^3)}{2x^2}$$

↓ 219

$$\frac{3}{2}bc \left(\frac{1}{6} \int \frac{2 - \sqrt[3]{cx}}{c^{2/3}x^2 - \sqrt[3]{cx} + 1} dx + \frac{1}{6} \int \frac{\sqrt[3]{cx} + 2}{c^{2/3}x^2 + \sqrt[3]{cx} + 1} dx + \frac{\operatorname{arctanh}(\sqrt[3]{cx})}{3\sqrt[3]{c}} \right) - \frac{a + \operatorname{barctanh}(cx^3)}{2x^2}$$

↓ 1142

$$\frac{3}{2}bc \left(\frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x^2 - \sqrt[3]{cx} + 1} dx - \frac{\int \frac{\sqrt[3]{c}(1 - 2\sqrt[3]{cx})}{c^{2/3}x^2 - \sqrt[3]{cx} + 1} dx}{2\sqrt[3]{c}} \right) + \frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x^2 + \sqrt[3]{cx} + 1} dx + \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{cx} + 1)}{c^{2/3}x^2 + \sqrt[3]{cx} + 1} dx}{2\sqrt[3]{c}} \right) \right) - \frac{a + \operatorname{barctanh}(cx^3)}{2x^2}$$

↓ 25

$$\frac{3}{2}bc \left(\frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x^2 - \sqrt[3]{cx} + 1} dx + \frac{\int \frac{\sqrt[3]{c}(1 - 2\sqrt[3]{cx})}{c^{2/3}x^2 - \sqrt[3]{cx} + 1} dx}{2\sqrt[3]{c}} \right) + \frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x^2 + \sqrt[3]{cx} + 1} dx + \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{cx} + 1)}{c^{2/3}x^2 + \sqrt[3]{cx} + 1} dx}{2\sqrt[3]{c}} \right) \right) - \frac{a + \operatorname{barctanh}(cx^3)}{2x^2}$$

↓ 27

$$\frac{3}{2}bc \left(\frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x^2 - \sqrt[3]{cx} + 1} dx + \frac{1}{2} \int \frac{1 - 2\sqrt[3]{cx}}{c^{2/3}x^2 - \sqrt[3]{cx} + 1} dx \right) + \frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x^2 + \sqrt[3]{cx} + 1} dx + \frac{1}{2} \int \frac{2\sqrt[3]{cx} + 1}{c^{2/3}x^2 + \sqrt[3]{cx} + 1} dx \right) \right) - \frac{a + \operatorname{barctanh}(cx^3)}{2x^2}$$

↓ 1082

$$\frac{3}{2}bc \left(\frac{1}{6} \left(\frac{1}{2} \int \frac{1 - 2\sqrt[3]{cx}}{c^{2/3}x^2 - \sqrt[3]{cx} + 1} dx + \frac{3 \int \frac{1}{-(1-2\sqrt[3]{cx})^2-3} d(1-2\sqrt[3]{cx})}{\sqrt[3]{c}} \right) + \frac{1}{6} \left(\frac{1}{2} \int \frac{2\sqrt[3]{cx} + 1}{c^{2/3}x^2 + \sqrt[3]{cx} + 1} dx - \frac{3 \int \frac{1}{(1+2\sqrt[3]{cx})^2-3} d(1+2\sqrt[3]{cx})}{\sqrt[3]{c}} \right) \right) + \frac{a + b \operatorname{arctanh}(cx^3)}{2x^2}$$

↓ 217

$$\frac{3}{2}bc \left(\frac{1}{6} \left(\frac{1}{2} \int \frac{1 - 2\sqrt[3]{cx}}{c^{2/3}x^2 - \sqrt[3]{cx} + 1} dx - \frac{\sqrt{3} \operatorname{arctan} \left(\frac{1-2\sqrt[3]{cx}}{\sqrt{3}} \right)}{\sqrt[3]{c}} \right) + \frac{1}{6} \left(\frac{1}{2} \int \frac{2\sqrt[3]{cx} + 1}{c^{2/3}x^2 + \sqrt[3]{cx} + 1} dx + \frac{\sqrt{3} \operatorname{arctan} \left(\frac{2\sqrt[3]{cx} + 1}{\sqrt{3}} \right)}{\sqrt[3]{c}} \right) \right) + \frac{a + b \operatorname{arctanh}(cx^3)}{2x^2}$$

↓ 1103

$$\frac{3}{2}bc \left(\frac{1}{6} \left(-\frac{\sqrt{3} \operatorname{arctan} \left(\frac{1-2\sqrt[3]{cx}}{\sqrt{3}} \right)}{\sqrt[3]{c}} - \frac{\log(c^{2/3}x^2 - \sqrt[3]{cx} + 1)}{2\sqrt[3]{c}} \right) + \frac{1}{6} \left(\frac{\sqrt{3} \operatorname{arctan} \left(\frac{2\sqrt[3]{cx} + 1}{\sqrt{3}} \right)}{\sqrt[3]{c}} + \frac{\log(c^{2/3}x^2 + \sqrt[3]{cx} + 1)}{2\sqrt[3]{c}} \right) \right) + \frac{a + b \operatorname{arctanh}(cx^3)}{2x^2}$$

input `Int[(a + b*ArcTanh[c*x^3])/x^3,x]`

output `-1/2*(a + b*ArcTanh[c*x^3])/x^2 + (3*b*c*(ArcTanh[c^(1/3)*x]/(3*c^(1/3)) + ((Sqrt[3]*ArcTan[(1 - 2*c^(1/3)*x]/Sqrt[3])]/c^(1/3)) - Log[1 - c^(1/3)*x + c^(2/3)*x^2]/(2*c^(1/3)))/6 + ((Sqrt[3]*ArcTan[(1 + 2*c^(1/3)*x]/Sqrt[3])]/c^(1/3) + Log[1 + c^(1/3)*x + c^(2/3)*x^2]/(2*c^(1/3)))/6)/2`

Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 754 `Int[((a_) + (b_.)*(x_)^(n_))(-1), x_Symbol] := Module[{r = Numerator[Rt[-a/b, n]], s = Denominator[Rt[-a/b, n]], k, u}, Simp[u = Int[(r - s*Cos[(2*k*Pi)/n]*x)/(r^2 - 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x] + Int[(r + s*Cos[(2*k*Pi)/n]*x)/(r^2 + 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x]; 2*(r^2/(a*n)) Int[1/(r^2 - s^2*x^2), x] + 2*(r/(a*n)) Sum[u, {k, 1, (n - 2)/4}, x]] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && NegQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142

```
Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c)
Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]
```

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.21

method	result
default	$-\frac{a}{2x^2} - \frac{b \operatorname{arctanh}(cx^3)}{2x^2} - \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{b \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{8\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{1}{c}\right)^{\frac{1}{3}} + 1\right)}\right)}{4\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{b \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4\left(\frac{1}{c}\right)^{\frac{2}{3}}}$
parts	$-\frac{a}{2x^2} - \frac{b \operatorname{arctanh}(cx^3)}{2x^2} - \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{b \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{8\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{1}{c}\right)^{\frac{1}{3}} + 1\right)}\right)}{4\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{b \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4\left(\frac{1}{c}\right)^{\frac{2}{3}}}$
risch	$-\frac{b \ln(cx^3+1)}{4x^2} - \frac{a}{2x^2} + \frac{b \ln(-cx^3+1)}{4x^2} - \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{b \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{8\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{1}{c}\right)^{\frac{1}{3}} + 1\right)}\right)}{4\left(\frac{1}{c}\right)^{\frac{2}{3}}}$

input

```
int((a+b*arctanh(c*x^3))/x^3,x,method=_RETURNVERBOSE)
```

output

```
-1/2*a/x^2-1/2*b/x^2*arctanh(c*x^3)-1/4*b/(1/c)^(2/3)*ln(x-(1/c)^(1/3))+1/
8*b/(1/c)^(2/3)*ln(x^2+(1/c)^(1/3)*x+(1/c)^(2/3))+1/4*b/(1/c)^(2/3)*3^(1/2
)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x+1))+1/4*b/(1/c)^(2/3)*ln(x+(1/c)^(1/
3))-1/8*b/(1/c)^(2/3)*ln(x^2-(1/c)^(1/3)*x+(1/c)^(2/3))+1/4*b/(1/c)^(2/3)*
3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x-1))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 228 vs. 2(97) = 194.

Time = 0.11 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.74

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^3} dx =$$

$$2\sqrt{3}(-c^2)^{\frac{1}{3}}bx^2 \arctan\left(\frac{2\sqrt{3}(-c^2)^{\frac{2}{3}}x + \sqrt{3}c}{3c}\right) - 2\sqrt{3}b(c^2)^{\frac{1}{3}}x^2 \arctan\left(\frac{2\sqrt{3}(c^2)^{\frac{2}{3}}x - \sqrt{3}c}{3c}\right) + (-c^2)^{\frac{1}{3}}bx^2 \log\left(\frac{2\sqrt{3}(-c^2)^{\frac{2}{3}}x + \sqrt{3}c}{3c}\right) - \frac{2\sqrt{3}b(c^2)^{\frac{1}{3}}x^2 \arctan\left(\frac{2\sqrt{3}(c^2)^{\frac{2}{3}}x - \sqrt{3}c}{3c}\right)}{(-c^2)^{\frac{1}{3}}}$$

input

```
integrate((a+b*arctanh(c*x^3))/x^3,x, algorithm="fricas")
```

output

```
-1/8*(2*sqrt(3)*(-c^2)^(1/3)*b*x^2*arctan(1/3*(2*sqrt(3)*(-c^2)^(2/3)*x +
sqrt(3)*c)/c) - 2*sqrt(3)*b*(c^2)^(1/3)*x^2*arctan(1/3*(2*sqrt(3)*(c^2)^(2/
3)*x - sqrt(3)*c)/c) + (-c^2)^(1/3)*b*x^2*log(c^2*x^2 - (-c^2)^(1/3)*c*x
+ (-c^2)^(2/3)) + b*(c^2)^(1/3)*x^2*log(c^2*x^2 - (c^2)^(1/3)*c*x + (c^2)^(
2/3)) - 2*(-c^2)^(1/3)*b*x^2*log(c*x + (-c^2)^(1/3)) - 2*b*(c^2)^(1/3)*x^
2*log(c*x + (c^2)^(1/3)) + 2*b*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 4*a)/x^2
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^3} dx = \text{Timed out}$$

input

```
integrate((a+b*atanh(c*x**3))/x**3,x)
```

output

```
Timed out
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.18

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^3} dx$$

$$= \frac{1}{8} \left(\frac{2\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}(2c^{\frac{2}{3}}x + c^{\frac{1}{3}})}{3c^{\frac{1}{3}}}\right)}{c^{\frac{1}{3}}} + \frac{2\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}(2c^{\frac{2}{3}}x - c^{\frac{1}{3}})}{3c^{\frac{1}{3}}}\right)}{c^{\frac{1}{3}}} + \frac{\log\left(c^{\frac{2}{3}}x^2 + c^{\frac{1}{3}}x + 1\right)}{c^{\frac{1}{3}}} - \frac{\log\left(c^{\frac{2}{3}}x^2 - c^{\frac{1}{3}}x + 1\right)}{c^{\frac{1}{3}}} \right) - \frac{a}{2x^2}$$

input `integrate((a+b*arctanh(c*x^3))/x^3,x, algorithm="maxima")`output `1/8*((2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*x + c^(1/3))/c^(1/3))/c^(1/3) + 2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*x - c^(1/3))/c^(1/3))/c^(1/3) + log(c^(2/3)*x^2 + c^(1/3)*x + 1)/c^(1/3) - log(c^(2/3)*x^2 - c^(1/3)*x + 1)/c^(1/3) + 2*log((c^(1/3)*x + 1)/c^(1/3))/c^(1/3) - 2*log((c^(1/3)*x - 1)/c^(1/3))/c^(1/3))*c - 4*arctanh(c*x^3)/x^2)*b - 1/2*a/x^2`**Giac [A] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.26

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^3} dx$$

$$= \frac{1}{8} \left(\frac{2\sqrt{3} \operatorname{arctan}\left(\frac{\frac{1}{3}\sqrt{3}\left(2x + \frac{1}{|c|^{\frac{1}{3}}}\right)|c|^{\frac{1}{3}}}{|c|^{\frac{1}{3}}}\right)}{|c|^{\frac{1}{3}}} + \frac{2\sqrt{3} \operatorname{arctan}\left(\frac{\frac{1}{3}\sqrt{3}\left(2x - \frac{1}{|c|^{\frac{1}{3}}}\right)|c|^{\frac{1}{3}}}{|c|^{\frac{1}{3}}}\right)}{|c|^{\frac{1}{3}}} + \frac{\log\left(x^2 + \frac{x}{|c|^{\frac{1}{3}}} + \frac{1}{|c|^{\frac{2}{3}}}\right)}{|c|^{\frac{1}{3}}} - \frac{\log\left(x^2 + \frac{x}{|c|^{\frac{1}{3}}} - \frac{1}{|c|^{\frac{2}{3}}}\right)}{|c|^{\frac{1}{3}}} \right) - \frac{b \log\left(\frac{-cx^3+1}{cx^3-1}\right)}{4x^2} - \frac{a}{2x^2}$$

input `integrate((a+b*arctanh(c*x^3))/x^3,x, algorithm="giac")`

output

```
1/8*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1/abs(c)^(1/3))*abs(c)^(1/3))/abs
(c)^(1/3) + 2*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1/abs(c)^(1/3))*abs(c)^(1/
3))/abs(c)^(1/3) + log(x^2 + x/abs(c)^(1/3) + 1/abs(c)^(2/3))/abs(c)^(1/3)
- log(x^2 - x/abs(c)^(1/3) + 1/abs(c)^(2/3))/abs(c)^(1/3) + 2*log(abs(x +
1/abs(c)^(1/3)))/abs(c)^(1/3) - 2*log(abs(x - 1/abs(c)^(1/3)))/abs(c)^(1/
3))*b*c - 1/4*b*log(-(c*x^3 + 1)/(c*x^3 - 1))/x^2 - 1/2*a/x^2
```

Mupad [B] (verification not implemented)

Time = 4.08 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.90

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^3} dx$$

$$= \frac{b \ln(1 - cx^3)}{4x^2}$$

$$- \frac{bc^{2/3} \left(-\frac{\operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}-i)}{2}\right)}{2} + \frac{\operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}+i)}{2}\right)}{2} + \operatorname{atan}(c^{1/3}x \operatorname{li}) \right) \operatorname{li}}{2}$$

$$- \frac{b \ln(cx^3 + 1)}{4x^2} - \frac{a}{2x^2} + \frac{\sqrt{3}bc^{2/3} \left(\operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}-i)}{2}\right) + \operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}+i)}{2}\right) \right)}{4}$$

input

```
int((a + b*atanh(c*x^3))/x^3,x)
```

output

```
(b*log(1 - c*x^3))/(4*x^2) - (b*c^(2/3)*(atan((c^(1/3)*x*(3^(1/2) + 1i))/2
)/2 - atan((c^(1/3)*x*(3^(1/2) - 1i))/2)/2 + atan(c^(1/3)*x*1i))*1i)/2 - (
b*log(c*x^3 + 1))/(4*x^2) - a/(2*x^2) + (3^(1/2)*b*c^(2/3)*(atan((c^(1/3)*
x*(3^(1/2) - 1i))/2) + atan((c^(1/3)*x*(3^(1/2) + 1i))/2)))/4
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.93

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^3} dx$$

$$= \frac{2\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x-1}{\sqrt{3}}\right) bcx^2 + 2\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x+1}{\sqrt{3}}\right) bcx^2 - 4c^{\frac{1}{3}} \operatorname{atanh}(cx^3)b - 2\operatorname{atanh}(cx^3)bcx^2 - 4c^{\frac{1}{3}}a + 3b}{8c^{\frac{1}{3}}x^2}$$

input `int((a+b*atanh(c*x^3))/x^3,x)`

output

```
(2*sqrt(3)*atan((2*c**(1/3)*x - 1)/sqrt(3))*b*c*x**2 + 2*sqrt(3)*atan((2*c
**(1/3)*x + 1)/sqrt(3))*b*c*x**2 - 4*c**(1/3)*atanh(c*x**3)*b - 2*atanh(c*
x**3)*b*c*x**2 - 4*c**(1/3)*a + 3*log(c**(2/3)*x + c**(1/3))*b*c*x**2 - 3*
log(c**(2/3)*x - c**(1/3))*b*c*x**2)/(8*c**(1/3)*x**2)
```

3.110 $\int \frac{a+b\operatorname{arctanh}(cx^3)}{x^6} dx$

Optimal result	929
Mathematica [A] (verified)	929
Rubi [A] (verified)	930
Maple [A] (verified)	933
Fricas [A] (verification not implemented)	934
Sympy [F(-1)]	935
Maxima [A] (verification not implemented)	935
Giac [A] (verification not implemented)	936
Mupad [B] (verification not implemented)	936
Reduce [B] (verification not implemented)	937

Optimal result

Integrand size = 14, antiderivative size = 115

$$\int \frac{a + b\operatorname{arctanh}(cx^3)}{x^6} dx = -\frac{3bc}{10x^2} - \frac{1}{10}\sqrt{3}bc^{5/3}\arctan\left(\frac{1 + 2c^{2/3}x^2}{\sqrt{3}}\right) - \frac{a + b\operatorname{arctanh}(cx^3)}{5x^5} - \frac{1}{10}bc^{5/3}\log(1 - c^{2/3}x^2) + \frac{1}{20}bc^{5/3}\log(1 + c^{2/3}x^2 + c^{4/3}x^4)$$

output

$$-3/10*b*c/x^2-1/10*3^{(1/2)}*b*c^{(5/3)}*\arctan(1/3*(1+2*c^{(2/3)}*x^2)*3^{(1/2)})-1/5*(a+b*\operatorname{arctanh}(c*x^3))/x^5-1/10*b*c^{(5/3)}*\ln(1-c^{(2/3)}*x^2)+1/20*b*c^{(5/3)}*\ln(1+c^{(2/3)}*x^2+c^{(4/3)}*x^4)$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.70

$$\int \frac{a + b\operatorname{arctanh}(cx^3)}{x^6} dx = -\frac{a}{5x^5} - \frac{3bc}{10x^2} - \frac{1}{10}\sqrt{3}bc^{5/3}\arctan\left(\frac{-1 + 2\sqrt[3]{cx}}{\sqrt{3}}\right) + \frac{1}{10}\sqrt{3}bc^{5/3}\arctan\left(\frac{1 + 2\sqrt[3]{cx}}{\sqrt{3}}\right) - \frac{b\operatorname{arctanh}(cx^3)}{5x^5} - \frac{1}{10}bc^{5/3}\log(1 - \sqrt[3]{cx}) - \frac{1}{10}bc^{5/3}\log(1 + \sqrt[3]{cx}) + \frac{1}{20}bc^{5/3}\log(1 - \sqrt[3]{cx} + c^{2/3}x^2) + \frac{1}{20}bc^{5/3}\log(1 + \sqrt[3]{cx} + c^{2/3}x^2)$$

input `Integrate[(a + b*ArcTanh[c*x^3])/x^6,x]`

output
$$-1/5*a/x^5 - (3*b*c)/(10*x^2) - (\text{Sqrt}[3]*b*c^{(5/3)}*\text{ArcTan}[(-1 + 2*c^{(1/3)}*x)/\text{Sqrt}[3]])/10 + (\text{Sqrt}[3]*b*c^{(5/3)}*\text{ArcTan}[(1 + 2*c^{(1/3)}*x)/\text{Sqrt}[3]])/10 - (b*\text{ArcTanh}[c*x^3])/(5*x^5) - (b*c^{(5/3)}*\text{Log}[1 - c^{(1/3)}*x])/10 - (b*c^{(5/3)}*\text{Log}[1 + c^{(1/3)}*x])/10 + (b*c^{(5/3)}*\text{Log}[1 - c^{(1/3)}*x + c^{(2/3)}*x^2])/20 + (b*c^{(5/3)}*\text{Log}[1 + c^{(1/3)}*x + c^{(2/3)}*x^2])/20$$

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.10, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {6452, 807, 847, 821, 16, 1142, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + \text{barctanh}(cx^3)}{x^6} dx \\ & \quad \downarrow \text{6452} \\ & \frac{3}{5}bc \int \frac{1}{x^3(1-c^2x^6)} dx - \frac{a + \text{barctanh}(cx^3)}{5x^5} \\ & \quad \downarrow \text{807} \\ & \frac{3}{10}bc \int \frac{1}{x^4(1-c^2x^6)} dx^2 - \frac{a + \text{barctanh}(cx^3)}{5x^5} \\ & \quad \downarrow \text{847} \\ & \frac{3}{10}bc \left(c^2 \int \frac{x^2}{1-c^2x^6} dx^2 - \frac{1}{x^2} \right) - \frac{a + \text{barctanh}(cx^3)}{5x^5} \\ & \quad \downarrow \text{821} \\ & \frac{3}{10}bc \left(c^2 \left(\int \frac{1}{1-c^{2/3}x^2} dx^2 - \int \frac{1-c^{2/3}x^2}{c^{4/3}x^4+c^{2/3}x^2+1} dx^2 \right) - \frac{1}{x^2} \right) - \frac{a + \text{barctanh}(cx^3)}{5x^5} \\ & \quad \downarrow \text{16} \end{aligned}$$

$$\begin{aligned}
& \frac{3}{10}bc \left(c^2 \left(-\frac{\int \frac{1-c^{2/3}x^2}{c^{4/3}x^4+c^{2/3}x^2+1} dx^2}{3c^{2/3}} - \frac{\log(1-c^{2/3}x^2)}{3c^{4/3}} \right) - \frac{1}{x^2} \right) - \frac{a + \operatorname{barctanh}(cx^3)}{5x^5} \\
& \quad \downarrow \text{1142} \\
& \frac{3}{10}bc \left(c^2 \left(-\frac{\frac{3}{2} \int \frac{1}{c^{4/3}x^4+c^{2/3}x^2+1} dx^2 - \frac{\int \frac{c^{2/3}(2c^{2/3}x^2+1)}{c^{4/3}x^4+c^{2/3}x^2+1} dx^2}{2c^{2/3}} - \frac{\log(1-c^{2/3}x^2)}{3c^{4/3}} \right) - \frac{1}{x^2} \right) - \frac{a + \operatorname{barctanh}(cx^3)}{5x^5} \\
& \quad \downarrow \text{27} \\
& \frac{3}{10}bc \left(c^2 \left(-\frac{\frac{3}{2} \int \frac{1}{c^{4/3}x^4+c^{2/3}x^2+1} dx^2 - \frac{1}{2} \int \frac{2c^{2/3}x^2+1}{c^{4/3}x^4+c^{2/3}x^2+1} dx^2 - \frac{\log(1-c^{2/3}x^2)}{3c^{4/3}} \right) - \frac{1}{x^2} \right) - \frac{a + \operatorname{barctanh}(cx^3)}{5x^5} \\
& \quad \downarrow \text{1082} \\
& \frac{3}{10}bc \left(c^2 \left(-\frac{3 \int \frac{1}{-x^4-3} d(2c^{2/3}x^2+1)}{c^{2/3}} - \frac{1}{2} \int \frac{2c^{2/3}x^2+1}{c^{4/3}x^4+c^{2/3}x^2+1} dx^2 - \frac{\log(1-c^{2/3}x^2)}{3c^{4/3}} \right) - \frac{1}{x^2} \right) - \frac{a + \operatorname{barctanh}(cx^3)}{5x^5} \\
& \quad \downarrow \text{217} \\
& \frac{3}{10}bc \left(c^2 \left(-\frac{\frac{\sqrt{3} \arctan\left(\frac{2c^{2/3}x^2+1}{\sqrt{3}}\right)}{c^{2/3}} - \frac{1}{2} \int \frac{2c^{2/3}x^2+1}{c^{4/3}x^4+c^{2/3}x^2+1} dx^2 - \frac{\log(1-c^{2/3}x^2)}{3c^{4/3}} \right) - \frac{1}{x^2} \right) - \frac{a + \operatorname{barctanh}(cx^3)}{5x^5} \\
& \quad \downarrow \text{1103} \\
& \frac{3}{10}bc \left(c^2 \left(-\frac{\frac{\sqrt{3} \arctan\left(\frac{2c^{2/3}x^2+1}{\sqrt{3}}\right)}{c^{2/3}} - \frac{\log(c^{4/3}x^4+c^{2/3}x^2+1)}{2c^{2/3}} - \frac{\log(1-c^{2/3}x^2)}{3c^{4/3}} \right) - \frac{1}{x^2} \right) - \frac{a + \operatorname{barctanh}(cx^3)}{5x^5}
\end{aligned}$$

input `Int[(a + b*ArcTanh[c*x^3])/x^6,x]`

output `-1/5*(a + b*ArcTanh[c*x^3])/x^5 + (3*b*c*(-x^(-2) + c^2*(-1/3*Log[1 - c^(2/3)*x^2]/c^(4/3) - ((Sqrt[3]*ArcTan[(1 + 2*c^(2/3)*x^2]/Sqrt[3]))/c^(2/3) - Log[1 + c^(2/3)*x^2 + c^(4/3)*x^4]/(2*c^(2/3)))/(3*c^(2/3))))/10`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 821 `Int[(x_)/((a_) + (b_.)*(x_)^3), x_Symbol] := Simp[-(3*Rt[a, 3]*Rt[b, 3])^(-1) Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Simp[1/(3*Rt[a, 3]*Rt[b, 3]) Int[(Rt[a, 3] + Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]`

rule 847 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + n*(p + 1) + 1)/(a*c^n*(m + 1)) Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.50

method	result
default	$-\frac{a}{5x^5} - \frac{b \operatorname{arctanh}(cx^3)}{5x^5} - \frac{bc \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{10\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{bc \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{20\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{bc\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2x - \left(\frac{1}{c}\right)^{\frac{1}{3}} + 1\right)}{3}\right)}{10\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{3bc}{10x^2}$
parts	$-\frac{a}{5x^5} - \frac{b \operatorname{arctanh}(cx^3)}{5x^5} - \frac{bc \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{10\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{bc \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{20\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{bc\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2x - \left(\frac{1}{c}\right)^{\frac{1}{3}} + 1\right)}{3}\right)}{10\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{3bc}{10x^2}$
risch	$-\frac{b \ln(cx^3+1)}{10x^5} - \frac{a}{5x^5} + \frac{b \ln(-cx^3+1)}{10x^5} - \frac{bc \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{10\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{bc \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{20\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{bc\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2x - \left(\frac{1}{c}\right)^{\frac{1}{3}} + 1\right)}{3}\right)}{10\left(\frac{1}{c}\right)^{\frac{2}{3}}}$

input

```
int((a+b*arctanh(c*x^3))/x^6,x,method=_RETURNVERBOSE)
```

output

```
-1/5*a/x^5-1/5*b/x^5*arctanh(c*x^3)-1/10*b*c/(1/c)^(2/3)*ln(x-(1/c)^(1/3))
+1/20*b*c/(1/c)^(2/3)*ln(x^2+(1/c)^(1/3)*x+(1/c)^(2/3))+1/10*b*c/(1/c)^(2/3)
*3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x+1))-3/10*b*c/x^2-1/10*b*c/(1/c)^(2/3)
*ln(x+(1/c)^(1/3))+1/20*b*c/(1/c)^(2/3)*ln(x^2-(1/c)^(1/3)*x+(1/c)^(2/3))
-1/10*b*c/(1/c)^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x-1))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.31

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^6} dx = \frac{2\sqrt{3}(-c^2)^{\frac{1}{3}}bcx^5 \operatorname{arctan}\left(\frac{2}{3}\sqrt{3}(-c^2)^{\frac{1}{3}}x^2 - \frac{1}{3}\sqrt{3}\right) + (-c^2)^{\frac{1}{3}}bcx^5 \log\left(c^2x^4 + (-c^2)^{\frac{2}{3}}x^2 - (-c^2)^{\frac{1}{3}}\right) - 2}{10x^5}$$

input

```
integrate((a+b*arctanh(c*x^3))/x^6,x, algorithm="fricas")
```

output

```
-1/20*(2*sqrt(3)*(-c^2)^(1/3)*b*c*x^5*arctan(2/3*sqrt(3)*(-c^2)^(1/3)*x^2
- 1/3*sqrt(3)) + (-c^2)^(1/3)*b*c*x^5*log(c^2*x^4 + (-c^2)^(2/3)*x^2 - (-c
^2)^(1/3)) - 2*(-c^2)^(1/3)*b*c*x^5*log(c^2*x^2 - (-c^2)^(2/3)) + 6*b*c*x^
3 + 2*b*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 4*a)/x^5
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^6} dx = \text{Timed out}$$

input

```
integrate((a+b*atanh(c*x**3))/x**6,x)
```

output

Timed out

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.87

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^6} dx =$$

$$-\frac{1}{20} \left(\left(2 \sqrt{3} c^{\frac{2}{3}} \arctan \left(\frac{\sqrt{3} (2 c^{\frac{4}{3}} x^2 + c^{\frac{2}{3}})}{3 c^{\frac{2}{3}}} \right) - c^{\frac{2}{3}} \log (c^{\frac{4}{3}} x^4 + c^{\frac{2}{3}} x^2 + 1) + 2 c^{\frac{2}{3}} \log \left(\frac{c^{\frac{2}{3}} x^2 - 1}{c^{\frac{2}{3}}} \right) + \frac{6}{x^2} \right) - \frac{a}{5 x^5} \right)$$

input

```
integrate((a+b*arctanh(c*x^3))/x^6,x, algorithm="maxima")
```

output

```
-1/20*((2*sqrt(3)*c^(2/3)*arctan(1/3*sqrt(3)*(2*c^(4/3)*x^2 + c^(2/3))/c^(
2/3)) - c^(2/3)*log(c^(4/3)*x^4 + c^(2/3)*x^2 + 1) + 2*c^(2/3)*log((c^(2/3
)*x^2 - 1)/c^(2/3)) + 6/x^2)*c + 4*arctanh(c*x^3)/x^5)*b - 1/5*a/x^5
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.09

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^6} dx =$$

$$-\frac{1}{20} bc^3 \left(\frac{2\sqrt{3}|c|^{\frac{2}{3}} \arctan\left(\frac{1}{3}\sqrt{3}\left(2x^2 + \frac{1}{|c|^{\frac{2}{3}}}\right)|c|^{\frac{2}{3}}\right)}{c^2} - \frac{|c|^{\frac{2}{3}} \log\left(x^4 + \frac{x^2}{|c|^{\frac{2}{3}}} + \frac{1}{|c|^{\frac{4}{3}}}\right)}{c^2} + \frac{2 \log\left(\left|x^2 - \frac{1}{|c|^{\frac{2}{3}}}\right|\right)}{|c|^{\frac{4}{3}}}\right)$$

$$-\frac{b \log\left(-\frac{cx^3+1}{cx^3-1}\right)}{10x^5} - \frac{3bcx^3 + 2a}{10x^5}$$

input `integrate((a+b*arctanh(c*x^3))/x^6,x, algorithm="giac")`output `-1/20*b*c^3*(2*sqrt(3)*abs(c)^(2/3)*arctan(1/3*sqrt(3)*(2*x^2 + 1/abs(c)^(2/3))*abs(c)^(2/3))/c^2 - abs(c)^(2/3)*log(x^4 + x^2/abs(c)^(2/3) + 1/abs(c)^(4/3))/c^2 + 2*log(abs(x^2 - 1/abs(c)^(2/3)))/abs(c)^(4/3) - 1/10*b*log(-(c*x^3 + 1)/(c*x^3 - 1))/x^5 - 1/10*(3*b*c*x^3 + 2*a)/x^5`**Mupad [B] (verification not implemented)**

Time = 6.00 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.17

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^6} dx = \frac{b \ln(1 - cx^3)}{10x^5} - \frac{bc^{5/3} \ln(c^{2/3}x^2 - 1)}{10}$$

$$-\frac{b \ln(cx^3 + 1)}{10x^5} - \frac{\frac{3bcx^3}{2} + a}{5x^5}$$

$$+ \frac{bc^{5/3} \ln(\sqrt{3}c^{2/3}x^2 - c^{2/3}x^2 \operatorname{li} - 2i)(1 + \sqrt{3} \operatorname{li})}{20}$$

$$- \frac{bc^{5/3} \ln(-c^{2/3}x^2 \operatorname{li} - \sqrt{3}c^{2/3}x^2 - 2i)(-1 + \sqrt{3} \operatorname{li})}{20}$$

input `int((a + b*atanh(c*x^3))/x^6,x)`

output

```
(b*log(1 - c*x^3))/(10*x^5) - (b*c^(5/3)*log(c^(2/3)*x^2 - 1))/10 - (b*log
(c*x^3 + 1))/(10*x^5) - (a + (3*b*c*x^3)/2)/(5*x^5) + (b*c^(5/3)*log(3^(1/
2)*c^(2/3)*x^2 - c^(2/3)*x^2*i - 2i)*(3^(1/2)*i + 1))/20 - (b*c^(5/3)*lo
g(- c^(2/3)*x^2*i - 3^(1/2)*c^(2/3)*x^2 - 2i)*(3^(1/2)*i - 1))/20
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.44

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^6} dx$$

$$= \frac{-2\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x-1}{\sqrt{3}}\right) b c^2 x^5 + 2\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x+1}{\sqrt{3}}\right) b c^2 x^5 - 4c^{\frac{1}{3}} \operatorname{atanh}(cx^3) b - 2 \operatorname{atanh}(cx^3) b c^2 x^5 - 4c^{\frac{1}{3}} a}{20c^{\frac{1}{3}}x^5}$$

input

```
int((a+b*atanh(c*x^3))/x^6,x)
```

output

```
( - 2*sqrt(3)*atan((2*c**(1/3)*x - 1)/sqrt(3))*b*c**2*x**5 + 2*sqrt(3)*ata
n((2*c**(1/3)*x + 1)/sqrt(3))*b*c**2*x**5 - 4*c**(1/3)*atanh(c*x**3)*b - 2
*atanh(c*x**3)*b*c**2*x**5 - 4*c**(1/3)*a - 6*c**(1/3)*b*c*x**3 + 2*log(c*
*(2/3)*x**2 - c**(1/3)*x + 1)*b*c**2*x**5 - log(c**(2/3)*x + c**(1/3))*b*c
**2*x**5 - 3*log(c**(2/3)*x - c**(1/3))*b*c**2*x**5)/(20*c**(1/3)*x**5)
```

3.111 $\int x^7(a + b \operatorname{arctanh}(cx^3)) dx$

Optimal result	938
Mathematica [A] (verified)	939
Rubi [A] (verified)	939
Maple [A] (verified)	944
Fricas [B] (verification not implemented)	945
Sympy [F(-1)]	945
Maxima [A] (verification not implemented)	946
Giac [A] (verification not implemented)	947
Mupad [B] (verification not implemented)	948
Reduce [B] (verification not implemented)	948

Optimal result

Integrand size = 14, antiderivative size = 142

$$\int x^7(a + b \operatorname{arctanh}(cx^3)) dx = \frac{3bx^5}{40c} - \frac{\sqrt{3}b \arctan\left(\frac{1-2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{16c^{8/3}} + \frac{\sqrt{3}b \arctan\left(\frac{1+2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{16c^{8/3}} - \frac{b \operatorname{arctanh}(\sqrt[3]{cx})}{8c^{8/3}} + \frac{1}{8}x^8(a + b \operatorname{arctanh}(cx^3)) - \frac{b \operatorname{arctanh}\left(\frac{\sqrt[3]{Cx}}{1+c^{2/3}x^2}\right)}{16c^{8/3}}$$

output

```
3/40*b*x^5/c-1/16*3^(1/2)*b*arctan(1/3*(1-2*c^(1/3)*x)*3^(1/2))/c^(8/3)+1/16*3^(1/2)*b*arctan(1/3*(1+2*c^(1/3)*x)*3^(1/2))/c^(8/3)-1/8*b*arctanh(c^(1/3)*x)/c^(8/3)+1/8*x^8*(a+b*arctanh(c*x^3))-1/16*b*arctanh(c^(1/3)*x/(1+c^(2/3)*x^2))/c^(8/3)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.39

$$\int x^7(a + b \operatorname{arctanh}(cx^3)) dx = \frac{3bx^5}{40c} + \frac{ax^8}{8} + \frac{\sqrt{3}b \arctan\left(\frac{-1+2\sqrt[3]{cx}}{\sqrt{3}}\right)}{16c^{8/3}}$$

$$+ \frac{\sqrt{3}b \arctan\left(\frac{1+2\sqrt[3]{cx}}{\sqrt{3}}\right)}{16c^{8/3}} + \frac{1}{8}bx^8 \operatorname{arctanh}(cx^3)$$

$$+ \frac{b \log(1 - \sqrt[3]{cx})}{16c^{8/3}} - \frac{b \log(1 + \sqrt[3]{cx})}{16c^{8/3}}$$

$$+ \frac{b \log(1 - \sqrt[3]{cx} + c^{2/3}x^2)}{32c^{8/3}} - \frac{b \log(1 + \sqrt[3]{cx} + c^{2/3}x^2)}{32c^{8/3}}$$

input

```
Integrate[x^7*(a + b*ArcTanh[c*x^3]),x]
```

output

```
(3*b*x^5)/(40*c) + (a*x^8)/8 + (Sqrt[3]*b*ArcTan[(-1 + 2*c^(1/3)*x)/Sqrt[3]])/(16*c^(8/3)) + (Sqrt[3]*b*ArcTan[(1 + 2*c^(1/3)*x)/Sqrt[3]])/(16*c^(8/3)) + (b*x^8*ArcTanh[c*x^3])/8 + (b*Log[1 - c^(1/3)*x])/(16*c^(8/3)) - (b*Log[1 + c^(1/3)*x])/(16*c^(8/3)) + (b*Log[1 - c^(1/3)*x + c^(2/3)*x^2])/(32*c^(8/3)) - (b*Log[1 + c^(1/3)*x + c^(2/3)*x^2])/(32*c^(8/3))
```

Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.35, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.786$, Rules used = {6452, 843, 825, 27, 219, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^7(a + b \operatorname{arctanh}(cx^3)) dx$$

$$\downarrow \text{6452}$$

$$\frac{1}{8}x^8(a + b \operatorname{arctanh}(cx^3)) - \frac{3}{8}bc \int \frac{x^{10}}{1 - c^2x^6} dx$$

↓ 843

$$\frac{1}{8}x^8(a + \operatorname{barctanh}(cx^3)) - \frac{3}{8}bc \left(\frac{\int \frac{x^4}{1-c^2x^6} dx}{c^2} - \frac{x^5}{5c^2} \right)$$

↓ 825

$$\frac{1}{8}x^8(a + \operatorname{barctanh}(cx^3)) - \frac{3}{8}bc \left(\frac{\int \frac{1}{1-c^{2/3}x^2} dx}{3c^{4/3}} + \frac{\int -\frac{\sqrt[3]{Cx+1}}{2(c^{2/3}x^2 - \sqrt[3]{Cx+1})} dx}{3c^{4/3}} + \frac{\int -\frac{1-\sqrt[3]{Cx}}{2(c^{2/3}x^2 + \sqrt[3]{Cx+1})} dx}{3c^{4/3}} - \frac{x^5}{5c^2} \right)$$

↓ 27

$$\frac{1}{8}x^8(a + \operatorname{barctanh}(cx^3)) - \frac{3}{8}bc \left(\frac{\int \frac{1}{1-c^{2/3}x^2} dx}{3c^{4/3}} - \frac{\int \frac{\sqrt[3]{Cx+1}}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx}{6c^{4/3}} - \frac{\int \frac{1-\sqrt[3]{Cx}}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx}{6c^{4/3}} - \frac{x^5}{5c^2} \right)$$

↓ 219

$$\frac{1}{8}x^8(a + \operatorname{barctanh}(cx^3)) - \frac{3}{8}bc \left(\frac{\int \frac{\sqrt[3]{Cx+1}}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx}{6c^{4/3}} - \frac{\int \frac{1-\sqrt[3]{Cx}}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{Cx})}{3c^{5/3}} - \frac{x^5}{5c^2} \right)$$

↓ 1142

$$\frac{1}{8}x^8(a + \operatorname{barctanh}(cx^3)) - \frac{3}{8}bc \left(\frac{\int -\frac{\sqrt[3]{C}(1-2\sqrt[3]{Cx})}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx}{6c^{4/3}} - \frac{\int \frac{\sqrt[3]{C}(2\sqrt[3]{Cx+1})}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx}{2\sqrt[3]{C}} - \frac{\frac{3}{2} \int \frac{1}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{Cx})}{3c^{5/3}} - \frac{x^5}{5c^2} \right)$$

↓ 25

$$\frac{1}{8}x^8(a + \operatorname{barctanh}(cx^3)) - \frac{3}{8}bc \left(\frac{\int \frac{\sqrt[3]{c}(1-2\sqrt[3]{Cx})}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx}{6c^{4/3}} - \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{Cx+1})}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx}{2\sqrt[3]{c}} + \frac{\operatorname{arctanh}(\sqrt[3]{Cx})}{3c^{5/3}} - \frac{x^5}{5c^2} \right)$$

27

$$\frac{1}{8}x^8(a + \operatorname{barctanh}(cx^3)) - \frac{3}{8}bc \left(\frac{\frac{3}{2} \int \frac{1}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx - \frac{1}{2} \int \frac{1-2\sqrt[3]{Cx}}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx}{6c^{4/3}} - \frac{\frac{3}{2} \int \frac{1}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx - \frac{1}{2} \int \frac{2\sqrt[3]{Cx+1}}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{Cx})}{3c^{5/3}} - \frac{x^5}{5c^2} \right)$$

1082

$$\frac{1}{8}x^8(a + \operatorname{barctanh}(cx^3)) - \frac{3}{8}bc \left(\frac{\frac{3 \int \frac{1}{(1-2\sqrt[3]{Cx})^2} d(1-2\sqrt[3]{Cx})}{\sqrt[3]{c}} - \frac{1}{2} \int \frac{1-2\sqrt[3]{Cx}}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx}{6c^{4/3}} - \frac{\frac{3 \int \frac{1}{(2\sqrt[3]{Cx+1})^2} d(2\sqrt[3]{Cx+1})}{\sqrt[3]{c}} - \frac{1}{2} \int \frac{2\sqrt[3]{Cx+1}}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{Cx})}{3c^{5/3}} - \frac{x^5}{5c^2} \right)$$

217

$$\frac{1}{8}x^8(a + \operatorname{barctanh}(cx^3)) - \frac{3}{8}bc \left(\frac{-\frac{1}{2} \int \frac{1-2\sqrt[3]{Cx}}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx - \frac{\sqrt{3} \operatorname{arctan}\left(\frac{1-2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{\sqrt[3]{c}}}{6c^{4/3}} - \frac{\frac{\sqrt{3} \operatorname{arctan}\left(\frac{2\sqrt[3]{Cx+1}}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{1}{2} \int \frac{2\sqrt[3]{Cx+1}}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{Cx})}{3c^{5/3}} - \frac{x^5}{5c^2} \right)$$

1103

$$\frac{3}{8}bc \left(\frac{\frac{1}{8}x^8(a + \operatorname{arctanh}(cx^3)) - \frac{\log(c^{2/3}x^2 - \sqrt[3]{Cx+1})}{2\sqrt[3]{c}} - \frac{\sqrt{3}\arctan\left(\frac{1-2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{\sqrt{3}\arctan\left(\frac{2\sqrt[3]{Cx+1}}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{\log(c^{2/3}x^2 + \sqrt[3]{Cx+1})}{2\sqrt[3]{c}}}{6c^{4/3}} - \frac{1}{c^2} + \frac{\operatorname{arctanh}(\sqrt[3]{Cx})}{3c^{5/3}} - \frac{x^5}{5c^2} \right)$$

input `Int[x^7*(a + b*ArcTanh[c*x^3]),x]`

output `(x^8*(a + b*ArcTanh[c*x^3]))/8 - (3*b*c*(-1/5*x^5/c^2 + (ArcTanh[c^(1/3)*x]/(3*c^(5/3)) - ((Sqrt[3]*ArcTan[(1 - 2*c^(1/3)*x]/Sqrt[3])]/c^(1/3)) + Log[1 - c^(1/3)*x + c^(2/3)*x^2]/(2*c^(1/3)))/(6*c^(4/3)) - ((Sqrt[3]*ArcTan[(1 + 2*c^(1/3)*x]/Sqrt[3])]/c^(1/3) - Log[1 + c^(1/3)*x + c^(2/3)*x^2]/(2*c^(1/3)))/(6*c^(4/3)))/c^2)/8`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(- (Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 825 `Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Module[{r = Numerator[Rt[-a/b, n]], s = Denominator[Rt[-a/b, n]], k, u}, Simp[u = Int[(r*Cos[2*k*m*(Pi/n)] - s*Cos[2*k*(m + 1)*(Pi/n)]*x)/(r^2 - 2*r*s*Cos[2*k*(Pi/n)]*x + s^2*x^2), x] + Int[(r*Cos[2*k*m*(Pi/n)] + s*Cos[2*k*(m + 1)*(Pi/n)]*x)/(r^2 + 2*r*s*Cos[2*k*(Pi/n)]*x + s^2*x^2), x]; 2*(r^(m + 2)/(a*n*s^m)) Int[1/(r^2 - s^2*x^2), x] + 2*(r^(m + 1)/(a*n*s^m)) Sum[u, {k, 1, (n - 2)/4}], x] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && NegQ[a/b]`

rule 843 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Simp[a*c^n*((m - n + 1)/(b*(m + n*p + 1))) Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)]*(b_))^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.31

method	result
default	$\frac{x^8 a}{8} + \frac{b x^8 \operatorname{arctanh}(c x^3)}{8} + \frac{3 b x^5}{40 c} + \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{16 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{b \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}} x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{32 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b \sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2 x}{\left(\frac{1}{c}\right)^{\frac{1}{3}} + 1\right)}\right)}{16 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}}$
parts	$\frac{x^8 a}{8} + \frac{b x^8 \operatorname{arctanh}(c x^3)}{8} + \frac{3 b x^5}{40 c} + \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{16 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{b \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}} x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{32 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b \sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2 x}{\left(\frac{1}{c}\right)^{\frac{1}{3}} + 1\right)}\right)}{16 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}}$
risch	$\frac{x^8 b \ln(c x^3 + 1)}{16} + \frac{x^8 a}{8} - \frac{b x^8 \ln(-c x^3 + 1)}{16} + \frac{3 b x^5}{40 c} + \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{16 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{b \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}} x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{32 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b \sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2 x}{\left(\frac{1}{c}\right)^{\frac{1}{3}} + 1\right)}\right)}{16 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}}$

```
input int(x^7*(a+b*arctanh(c*x^3)),x,method=_RETURNVERBOSE)
```

```
output 1/8*x^8*a+1/8*b*x^8*arctanh(c*x^3)+3/40*b*x^5/c+1/16*b/c^3/(1/c)^(1/3)*ln(x-(1/c)^(1/3))-1/32*b/c^3/(1/c)^(1/3)*ln(x^2+(1/c)^(1/3)*x+(1/c)^(2/3))+1/16*b/c^3*3^(1/2)/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x+1))-1/16*b/c^3/(1/c)^(1/3)*ln(x+(1/c)^(1/3))+1/32*b/c^3/(1/c)^(1/3)*ln(x^2-(1/c)^(1/3)*x+(1/c)^(2/3))+1/16*b/c^3*3^(1/2)/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x-1))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 248 vs. $2(106) = 212$.

Time = 0.10 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.75

$$\int x^7(a + \operatorname{barctanh}(cx^3)) dx$$

$$= \frac{10bc^4x^8 \log\left(-\frac{cx^3+1}{cx^3-1}\right) + 20ac^4x^8 + 12bc^3x^5 + 10\sqrt{3}bc\sqrt{-(-c^2)^{\frac{1}{3}}}\arctan\left(\frac{\sqrt{3}\left(2cx+(-c^2)^{\frac{1}{3}}\right)\sqrt{-(-c^2)^{\frac{1}{3}}}}{3c}\right)}{c^4} + \dots$$

input `integrate(x^7*(a+b*arctanh(c*x^3)),x, algorithm="fricas")`

output `1/160*(10*b*c^4*x^8*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 20*a*c^4*x^8 + 12*b*c^3*x^5 + 10*sqrt(3)*b*c*sqrt(-(-c^2)^(1/3))*arctan(1/3*sqrt(3)*(2*c*x + (-c^2)^(1/3))*sqrt(-(-c^2)^(1/3))/c) + 10*sqrt(3)*b*(c^2)^(1/6)*c*arctan(1/3*sqrt(3)*(c^2)^(1/6)*(2*c*x + (c^2)^(1/3))/c) + 5*(-c^2)^(2/3)*b*log(c^2*x^2 + (-c^2)^(1/3)*c*x + (-c^2)^(2/3)) - 5*b*(c^2)^(2/3)*log(c^2*x^2 + (c^2)^(1/3)*c*x + (c^2)^(2/3)) - 10*(-c^2)^(2/3)*b*log(c*x - (-c^2)^(1/3)) + 10*b*(c^2)^(2/3)*log(c*x - (c^2)^(1/3)))/c^4`

Sympy [F(-1)]

Timed out.

$$\int x^7(a + \operatorname{barctanh}(cx^3)) dx = \text{Timed out}$$

input `integrate(x**7*(a+b*atanh(c*x**3)),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.15

$$\int x^7 (a + b \operatorname{arctanh}(cx^3)) dx = \frac{1}{8} ax^8 + \frac{1}{160} \left(20x^8 \operatorname{arctanh}(cx^3) + \left(\frac{12x^5}{c^2} + \frac{10\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}(2c^{2/3}x + c^{1/3})}{3c^{1/3}}\right)}{c^{11/3}} + \frac{10\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}(2c^{2/3}x - c^{1/3})}{3c^{1/3}}\right)}{c^{11/3}} \right) \right)$$

input `integrate(x^7*(a+b*arctanh(c*x^3)),x, algorithm="maxima")`

output `1/8*a*x^8 + 1/160*(20*x^8*arctanh(c*x^3) + (12*x^5/c^2 + 10*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*x + c^(1/3))/c^(1/3))/c^(11/3) + 10*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*x - c^(1/3))/c^(1/3))/c^(11/3) - 5*log(c^(2/3)*x^2 + c^(1/3)*x + 1)/c^(11/3) + 5*log(c^(2/3)*x^2 - c^(1/3)*x + 1)/c^(11/3) - 10*log((c^(1/3)*x + 1)/c^(1/3))/c^(11/3) + 10*log((c^(1/3)*x - 1)/c^(1/3))/c^(11/3))*c)*b`

Giac [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.46

$$\begin{aligned}
\int x^7(a + b \operatorname{arctanh}(cx^3)) dx &= \frac{1}{16} bx^8 \log\left(-\frac{cx^3 + 1}{cx^3 - 1}\right) + \frac{1}{8} ax^8 \\
&+ \frac{3bx^5}{40c} - \frac{b(-\frac{1}{c})^{\frac{2}{3}} \log\left(\left|x - (-\frac{1}{c})^{\frac{1}{3}}\right|\right)}{16c^2} \\
&+ \frac{\sqrt{3}b \arctan\left(\frac{\sqrt{3}\left(2x + (-\frac{1}{c})^{\frac{1}{3}}\right)}{3(-\frac{1}{c})^{\frac{1}{3}}}\right)}{16(-c^2)^{\frac{1}{3}}c^2} \\
&+ \frac{\sqrt{3}b \arctan\left(\frac{1}{3}\sqrt{3}c^{\frac{1}{3}}\left(2x + \frac{1}{c^{\frac{1}{3}}}\right)\right)}{16c^2|c|^{\frac{2}{3}}} \\
&- \frac{b \log\left(x^2 + x(-\frac{1}{c})^{\frac{1}{3}} + (-\frac{1}{c})^{\frac{2}{3}}\right)}{32(-c^2)^{\frac{1}{3}}c^2} \\
&- \frac{b \log\left(x^2 + \frac{x}{c^{\frac{1}{3}}} + \frac{1}{c^{\frac{2}{3}}}\right)}{32c^2|c|^{\frac{2}{3}}} + \frac{b \log\left(\left|x - \frac{1}{c^{\frac{1}{3}}}\right|\right)}{16c^{\frac{8}{3}}}
\end{aligned}$$

input `integrate(x^7*(a+b*arctanh(c*x^3)),x, algorithm="giac")`

output

```

1/16*b*x^8*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 1/8*a*x^8 + 3/40*b*x^5/c - 1/16
*b*(-1/c)^(2/3)*log(abs(x - (-1/c)^(1/3)))/c^2 + 1/16*sqrt(3)*b*arctan(1/3
*sqrt(3)*(2*x + (-1/c)^(1/3))/(-1/c)^(1/3))/((-c^2)^(1/3)*c^2) + 1/16*sqrt
(3)*b*arctan(1/3*sqrt(3)*c^(1/3)*(2*x + 1/c^(1/3)))/(c^2*abs(c)^(2/3)) - 1
/32*b*log(x^2 + x*(-1/c)^(1/3) + (-1/c)^(2/3))/((-c^2)^(1/3)*c^2) - 1/32*b
*log(x^2 + x/c^(1/3) + 1/c^(2/3))/(c^2*abs(c)^(2/3)) + 1/16*b*log(abs(x -
1/c^(1/3)))/c^(8/3)

```


Mupad [B] (verification not implemented)

Time = 4.01 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.89

$$\int x^7 (a + \operatorname{barctanh}(cx^3)) dx$$

$$= \frac{ax^8}{8} + \frac{b \left(-\frac{\operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}-i)}{2}\right)}{2} + \frac{\operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}+i)}{2}\right)}{2} + \operatorname{atan}(c^{1/3}xi) \right) 1i}{8c^{8/3}}$$

$$+ \frac{3bx^5}{40c} + \frac{bx^8 \ln(cx^3 + 1)}{16} - \frac{bx^8 \ln(1 - cx^3)}{16}$$

$$+ \frac{\sqrt{3}b \left(\operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}-i)}{2}\right) + \operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}+i)}{2}\right) \right)}{16c^{8/3}}$$

input `int(x^7*(a + b*atanh(c*x^3)),x)`output `(a*x^8)/8 + (b*(atan((c^(1/3)*x*(3^(1/2) + 1i))/2)/2 - atan((c^(1/3)*x*(3^(1/2) - 1i))/2)/2 + atan(c^(1/3)*x*1i))*1i)/(8*c^(8/3)) + (3*b*x^5)/(40*c) + (b*x^8*log(c*x^3 + 1))/16 - (b*x^8*log(1 - c*x^3))/16 + (3^(1/2)*b*(atan((c^(1/3)*x*(3^(1/2) - 1i))/2) + atan((c^(1/3)*x*(3^(1/2) + 1i))/2)))/(16*c^(8/3))`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.80

$$\int x^7 (a + \operatorname{barctanh}(cx^3)) dx$$

$$= \frac{10\sqrt{3} \operatorname{atan}\left(\frac{2c^{1/3}x-1}{\sqrt{3}}\right) b + 10\sqrt{3} \operatorname{atan}\left(\frac{2c^{1/3}x+1}{\sqrt{3}}\right) b + 20c^{8/3} \operatorname{atanh}(cx^3) bx^8 + 10 \operatorname{atanh}(cx^3) b + 20c^{8/3} ax^8 + 12}{160c^{8/3}}$$

input `int(x^7*(a+b*atanh(c*x^3)),x)`

output

```
(10*sqrt(3)*atan((2*c**(1/3)*x - 1)/sqrt(3))*b + 10*sqrt(3)*atan((2*c**(1/3)*x + 1)/sqrt(3))*b + 20*c**(2/3)*atanh(c*x**3)*b*c**2*x**8 + 10*atanh(c*x**3)*b + 20*c**(2/3)*a*c**2*x**8 + 12*c**(2/3)*b*c*x**5 - 15*log(c**(2/3)*x + c**(1/3))*b + 15*log(c**(2/3)*x - c**(1/3))*b)/(160*c**(2/3)*c**2)
```

3.112 $\int x^4(a + \operatorname{barctanh}(cx^3)) dx$

Optimal result	950
Mathematica [A] (verified)	951
Rubi [A] (verified)	951
Maple [A] (verified)	955
Fricas [A] (verification not implemented)	956
Sympy [F(-1)]	956
Maxima [A] (verification not implemented)	957
Giac [A] (verification not implemented)	957
Mupad [B] (verification not implemented)	958
Reduce [B] (verification not implemented)	958

Optimal result

Integrand size = 14, antiderivative size = 117

$$\int x^4(a + \operatorname{barctanh}(cx^3)) dx = \frac{3bx^2}{10c} - \frac{\sqrt{3}b \arctan\left(\frac{1+2c^{2/3}x^2}{\sqrt{3}}\right)}{10c^{5/3}} + \frac{1}{5}x^5(a + \operatorname{barctanh}(cx^3))$$

$$+ \frac{b \log(1 - c^{2/3}x^2)}{10c^{5/3}} - \frac{b \log(1 + c^{2/3}x^2 + c^{4/3}x^4)}{20c^{5/3}}$$

output

```
3/10*b*x^2/c-1/10*3^(1/2)*b*arctan(1/3*(1+2*c^(2/3)*x^2)*3^(1/2))/c^(5/3)+
1/5*x^5*(a+b*arctanh(c*x^3))+1/10*b*ln(1-c^(2/3)*x^2)/c^(5/3)-1/20*b*ln(1+
c^(2/3)*x^2+c^(4/3)*x^4)/c^(5/3)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.69

$$\int x^4(a + b \operatorname{arctanh}(cx^3)) dx = \frac{3bx^2}{10c} + \frac{ax^5}{5} - \frac{\sqrt{3}b \arctan\left(\frac{-1+2\sqrt[3]{cx}}{\sqrt{3}}\right)}{10c^{5/3}}$$

$$+ \frac{\sqrt{3}b \arctan\left(\frac{1+2\sqrt[3]{cx}}{\sqrt{3}}\right)}{10c^{5/3}} + \frac{1}{5}bx^5 \operatorname{arctanh}(cx^3)$$

$$+ \frac{b \log(1 - \sqrt[3]{cx})}{10c^{5/3}} + \frac{b \log(1 + \sqrt[3]{cx})}{10c^{5/3}}$$

$$- \frac{b \log(1 - \sqrt[3]{cx} + c^{2/3}x^2)}{20c^{5/3}} - \frac{b \log(1 + \sqrt[3]{cx} + c^{2/3}x^2)}{20c^{5/3}}$$

input

```
Integrate[x^4*(a + b*ArcTanh[c*x^3]),x]
```

output

```
(3*b*x^2)/(10*c) + (a*x^5)/5 - (Sqrt[3]*b*ArcTan[(-1 + 2*c^(1/3)*x)/Sqrt[3]])/(10*c^(5/3)) + (Sqrt[3]*b*ArcTan[(1 + 2*c^(1/3)*x)/Sqrt[3]])/(10*c^(5/3)) + (b*x^5*ArcTanh[c*x^3])/5 + (b*Log[1 - c^(1/3)*x])/(10*c^(5/3)) + (b*Log[1 + c^(1/3)*x])/(10*c^(5/3)) - (b*Log[1 - c^(1/3)*x + c^(2/3)*x^2])/(20*c^(5/3)) - (b*Log[1 + c^(1/3)*x + c^(2/3)*x^2])/(20*c^(5/3))
```

Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.07, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {6452, 807, 843, 750, 16, 1142, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4(a + b \operatorname{arctanh}(cx^3)) dx$$

$$\downarrow 6452$$

$$\frac{1}{5}x^5(a + b \operatorname{arctanh}(cx^3)) - \frac{3}{5}bc \int \frac{x^7}{1 - c^2x^6} dx$$

↓ 807

$$\frac{1}{5}x^5(a + \operatorname{barctanh}(cx^3)) - \frac{3}{10}bc \int \frac{x^6}{1 - c^2x^6} dx^2$$

↓ 843

$$\frac{1}{5}x^5(a + \operatorname{barctanh}(cx^3)) - \frac{3}{10}bc \left(\frac{\int \frac{1}{1 - c^2x^6} dx^2}{c^2} - \frac{x^2}{c^2} \right)$$

↓ 750

$$\frac{1}{5}x^5(a + \operatorname{barctanh}(cx^3)) - \frac{3}{10}bc \left(\frac{\frac{1}{3} \int \frac{1}{1 - c^{2/3}x^2} dx^2 + \frac{1}{3} \int \frac{c^{2/3}x^2 + 2}{c^{4/3}x^4 + c^{2/3}x^2 + 1} dx^2}{c^2} - \frac{x^2}{c^2} \right)$$

↓ 16

$$\frac{1}{5}x^5(a + \operatorname{barctanh}(cx^3)) - \frac{3}{10}bc \left(\frac{\frac{1}{3} \int \frac{c^{2/3}x^2 + 2}{c^{4/3}x^4 + c^{2/3}x^2 + 1} dx^2 - \frac{\log(1 - c^{2/3}x^2)}{3c^{2/3}}}{c^2} - \frac{x^2}{c^2} \right)$$

↓ 1142

$$\frac{1}{5}x^5(a + \operatorname{barctanh}(cx^3)) - \frac{3}{10}bc \left(\frac{\frac{1}{3} \left(\frac{3}{2} \int \frac{1}{c^{4/3}x^4 + c^{2/3}x^2 + 1} dx^2 + \frac{\int \frac{c^{2/3}(2c^{2/3}x^2 + 1)}{c^{4/3}x^4 + c^{2/3}x^2 + 1} dx^2}{2c^{2/3}} \right) - \frac{\log(1 - c^{2/3}x^2)}{3c^{2/3}}}{c^2} - \frac{x^2}{c^2} \right)$$

↓ 27

$$\frac{1}{5}x^5(a + \operatorname{barctanh}(cx^3)) - \frac{3}{10}bc \left(\frac{\frac{1}{3} \left(\frac{3}{2} \int \frac{1}{c^{4/3}x^4 + c^{2/3}x^2 + 1} dx^2 + \frac{1}{2} \int \frac{2c^{2/3}x^2 + 1}{c^{4/3}x^4 + c^{2/3}x^2 + 1} dx^2 \right) - \frac{\log(1 - c^{2/3}x^2)}{3c^{2/3}}}{c^2} - \frac{x^2}{c^2} \right)$$

↓ 1082

$$\frac{1}{5}x^5(a + \operatorname{barctanh}(cx^3)) - \frac{3}{10}bc \left(\frac{\frac{1}{3} \left(\frac{1}{2} \int \frac{2c^{2/3}x^2 + 1}{c^{4/3}x^4 + c^{2/3}x^2 + 1} dx^2 - \frac{3 \int \frac{-1}{-x^4 - 3} d(2c^{2/3}x^2 + 1)}{c^{2/3}} \right) - \frac{\log(1 - c^{2/3}x^2)}{3c^{2/3}}}{c^2} - \frac{x^2}{c^2} \right)$$

$$\begin{array}{c} \downarrow 217 \\ \frac{1}{5}x^5(a + b\operatorname{arctanh}(cx^3)) - \\ \frac{3}{10}bc \left(\frac{\frac{1}{3} \left(\frac{1}{2} \int \frac{2c^{2/3}x^2+1}{c^{4/3}x^4+c^{2/3}x^2+1} dx^2 + \frac{\sqrt{3} \arctan\left(\frac{2c^{2/3}x^2+1}{\sqrt{3}}\right)}{c^{2/3}} \right) - \frac{\log(1-c^{2/3}x^2)}{3c^{2/3}}}{c^2} - \frac{x^2}{c^2} \right) \end{array}$$

$$\begin{array}{c} \downarrow 1103 \\ \frac{1}{5}x^5(a + b\operatorname{arctanh}(cx^3)) - \\ \frac{3}{10}bc \left(\frac{\frac{1}{3} \left(\frac{\sqrt{3} \arctan\left(\frac{2c^{2/3}x^2+1}{\sqrt{3}}\right)}{c^{2/3}} + \frac{\log(c^{4/3}x^4+c^{2/3}x^2+1)}{2c^{2/3}} \right) - \frac{\log(1-c^{2/3}x^2)}{3c^{2/3}}}{c^2} - \frac{x^2}{c^2} \right) \end{array}$$

input `Int[x^4*(a + b*ArcTanh[c*x^3]),x]`

output `(x^5*(a + b*ArcTanh[c*x^3]))/5 - (3*b*c*(-(x^2/c^2) + (-1/3*Log[1 - c^(2/3)*x^2]/c^(2/3) + ((Sqrt[3]*ArcTan[(1 + 2*c^(2/3)*x^2]/Sqrt[3])]/c^(2/3) + Log[1 + c^(2/3)*x^2 + c^(4/3)*x^4]/(2*c^(2/3)))/3)/c^2)/10`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 $\text{Int}[\{(a_)+(b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /;$ $\text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 750 $\text{Int}[\{(a_)+(b_)*(x_)^3\}^{-1}, x_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \ \text{Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \ \text{Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /;$ $\text{FreeQ}\{a, b\}, x]$

rule 807 $\text{Int}[(x_)^{(m_)*((a_)+(b_)*(x_)^{n_})}^{(p_)}, x_Symbol] \rightarrow \text{With}\{k = \text{GCD}[m + 1, n]\}, \text{Simp}[1/k \ \text{Subst}[\text{Int}[x^{(m+1)/k - 1}*(a + b*x^{(n/k)})^p, x], x, x^k], x] /;$ $k \neq 1] /;$ $\text{FreeQ}\{a, b, p\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$

rule 843 $\text{Int}[\{(c_)*(x_)\}^{(m_)*((a_)+(b_)*(x_)^{n_})}^{(p_)}, x_Symbol] \rightarrow \text{Simp}[c^{(n-1)}*(c*x)^{(m-n+1)}*((a + b*x^n)^{(p+1})/(b*(m+n*p+1))), x] - \text{Simp}[a*c^n*((m-n+1)/(b*(m+n*p+1))) \ \text{Int}[(c*x)^{(m-n)}*(a + b*x^n)^p, x], x] /;$ $\text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{GtQ}[m, n-1] \ \&\& \ \text{NeQ}[m+n*p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 1082 $\text{Int}[\{(a_)+(b_)*(x_)+(c_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{With}\{q = 1 - 4*S\text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /;$ $\text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c]) /;$ $\text{FreeQ}\{a, b, c\}, x]$

rule 1103 $\text{Int}[\{(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2)\}, x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /;$ $\text{FreeQ}\{a, b, c, d, e\}, x \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$

rule 1142 $\text{Int}[\{(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2)\}, x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \ \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \ \text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /;$ $\text{FreeQ}\{a, b, c, d, e\}, x]$

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.97

method	result
default	$\frac{ax^5}{5} + \frac{bx^5 \operatorname{arctanh}(cx^3)}{5} + \frac{3bx^2}{10c} + \frac{b \ln\left(x^2 - \left(\frac{1}{c^2}\right)^{\frac{1}{3}}\right)}{10c^3 \left(\frac{1}{c^2}\right)^{\frac{2}{3}}} - \frac{b \ln\left(x^4 + \left(\frac{1}{c^2}\right)^{\frac{1}{3}}x^2 + \left(\frac{1}{c^2}\right)^{\frac{2}{3}}\right)}{20c^3 \left(\frac{1}{c^2}\right)^{\frac{2}{3}}} - \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2x^2}{\left(\frac{1}{c^2}\right)^{\frac{1}{3}} + 1\right)}\right)}{10c^3 \left(\frac{1}{c^2}\right)^{\frac{2}{3}}}$
parts	$\frac{ax^5}{5} + \frac{bx^5 \operatorname{arctanh}(cx^3)}{5} + \frac{3bx^2}{10c} + \frac{b \ln\left(x^2 - \left(\frac{1}{c^2}\right)^{\frac{1}{3}}\right)}{10c^3 \left(\frac{1}{c^2}\right)^{\frac{2}{3}}} - \frac{b \ln\left(x^4 + \left(\frac{1}{c^2}\right)^{\frac{1}{3}}x^2 + \left(\frac{1}{c^2}\right)^{\frac{2}{3}}\right)}{20c^3 \left(\frac{1}{c^2}\right)^{\frac{2}{3}}} - \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2x^2}{\left(\frac{1}{c^2}\right)^{\frac{1}{3}} + 1\right)}\right)}{10c^3 \left(\frac{1}{c^2}\right)^{\frac{2}{3}}}$
risch	$\frac{bx^5 \ln(cx^3 + 1)}{10} + \frac{ax^5}{5} - \frac{bx^5 \ln(-cx^3 + 1)}{10} + \frac{3bx^2}{10c} + \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{10c^2 \left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{b \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{20c^2 \left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2x^2}{\left(\frac{1}{c^2}\right)^{\frac{1}{3}} + 1\right)}\right)}{10c^2 \left(\frac{1}{c}\right)^{\frac{1}{3}}}$

input

```
int(x^4*(a+b*arctanh(c*x^3)),x,method=_RETURNVERBOSE)
```

output

```
1/5*a*x^5+1/5*b*x^5*arctanh(c*x^3)+3/10*b*x^2/c+1/10*b/c^3/(1/c^2)^(2/3)*
ln(x^2-(1/c^2)^(1/3))-1/20*b/c^3/(1/c^2)^(2/3)*ln(x^4+(1/c^2)^(1/3)*x^2+(1/
c^2)^(2/3))-1/10*b/c^3/(1/c^2)^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c^2)
^(1/3)*x^2+1))
```


Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.27

$$\int x^4(a + \operatorname{arctanh}(cx^3)) dx$$

$$= \frac{2bc^3x^5 \log\left(-\frac{cx^3+1}{cx^3-1}\right) + 4ac^3x^5 + 6bc^2x^2 - 2\sqrt{3}b(c^2)^{\frac{1}{6}}c \arctan\left(\frac{\sqrt{3}\left(2(c^2)^{\frac{2}{3}}x^2+(c^2)^{\frac{1}{3}}\right)(c^2)^{\frac{1}{6}}}{3c}\right) - b(c^2)^{\frac{2}{3}} \log\left(\dots\right)}{20c^3}$$

input `integrate(x^4*(a+b*arctanh(c*x^3)),x, algorithm="fricas")`output `1/20*(2*b*c^3*x^5*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 4*a*c^3*x^5 + 6*b*c^2*x^2 - 2*sqrt(3)*b*(c^2)^(1/6)*c*arctan(1/3*sqrt(3)*(2*(c^2)^(2/3)*x^2 + (c^2)^(1/3))*(c^2)^(1/6)/c) - b*(c^2)^(2/3)*log(c^2*x^4 + (c^2)^(2/3)*x^2 + (c^2)^(1/3)) + 2*b*(c^2)^(2/3)*log(c^2*x^2 - (c^2)^(2/3)))/c^3`**Sympy [F(-1)]**

Timed out.

$$\int x^4(a + \operatorname{arctanh}(cx^3)) dx = \text{Timed out}$$

input `integrate(x**4*(a+b*atanh(c*x**3)),x)`output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.88

$$\int x^4(a + b \operatorname{arctanh}(cx^3)) dx = \frac{1}{5} ax^5 + \frac{1}{20} \left(4x^5 \operatorname{arctanh}(cx^3) + c \left(\frac{6x^2}{c^2} - \frac{2\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}(2c^{\frac{4}{3}}x^2 + c^{\frac{2}{3}})}{3c^{\frac{2}{3}}}\right)}{c^{\frac{8}{3}}} - \frac{\log\left(c^{\frac{4}{3}}x^4 + c^{\frac{2}{3}}x^2 + 1\right)}{c^{\frac{8}{3}}} + \frac{2 \log\left(\frac{x^2 - \frac{1}{|c|^{\frac{2}{3}}}}{\left|x^2 + \frac{1}{|c|^{\frac{4}{3}}}\right|}\right)}{c^{\frac{10}{3}}}\right) \right)$$

input `integrate(x^4*(a+b*arctanh(c*x^3)),x, algorithm="maxima")`output `1/5*a*x^5 + 1/20*(4*x^5*arctanh(c*x^3) + c*(6*x^2/c^2 - 2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(4/3)*x^2 + c^(2/3))/c^(2/3))/c^(8/3) - log(c^(4/3)*x^4 + c^(2/3)*x^2 + 1)/c^(8/3) + 2*log((c^(2/3)*x^2 - 1)/c^(2/3))/c^(8/3))*b`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.08

$$\int x^4(a + b \operatorname{arctanh}(cx^3)) dx = -\frac{1}{20} bc^9 \left(\frac{2\sqrt{3} \operatorname{arctan}\left(\frac{\frac{1}{3}\sqrt{3}\left(2x^2 + \frac{1}{|c|^{\frac{2}{3}}}\right)|c|^{\frac{2}{3}}}{c^{10}|c|^{\frac{2}{3}}}\right)}{c^{10}|c|^{\frac{2}{3}}} + \frac{\log\left(x^4 + \frac{x^2}{|c|^{\frac{2}{3}}} + \frac{1}{|c|^{\frac{4}{3}}}\right)}{c^{10}|c|^{\frac{2}{3}}} - \frac{2 \log\left(\left|x^2 - \frac{1}{|c|^{\frac{2}{3}}}\right|\right)}{c^{10}|c|^{\frac{2}{3}}}\right) + \frac{1}{10} bx^5 \log\left(-\frac{cx^3 + 1}{cx^3 - 1}\right) + \frac{1}{5} ax^5 + \frac{3bx^2}{10c}$$

input `integrate(x^4*(a+b*arctanh(c*x^3)),x, algorithm="giac")`output `-1/20*b*c^9*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*x^2 + 1/abs(c)^(2/3))*abs(c)^(2/3))/(c^10*abs(c)^(2/3)) + log(x^4 + x^2/abs(c)^(2/3) + 1/abs(c)^(4/3))/(c^10*abs(c)^(2/3)) - 2*log(abs(x^2 - 1/abs(c)^(2/3)))/(c^10*abs(c)^(2/3)) + 1/10*b*x^5*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 1/5*a*x^5 + 3/10*b*x^2/c`

Mupad [B] (verification not implemented)

Time = 5.22 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.06

$$\int x^4(a + \operatorname{barctanh}(cx^3)) dx = \frac{ax^5}{5} + \frac{b \ln(1 - c^{2/3}x^2)}{10c^{5/3}} + \frac{3bx^2}{10c} - \frac{\ln(2c^{2/3}x^2 + 1 - \sqrt{3}i)(b - \sqrt{3}bi)}{20c^{5/3}} - \frac{\ln(2c^{2/3}x^2 + 1 + \sqrt{3}i)(b + \sqrt{3}bi)}{20c^{5/3}} + \frac{bx^5 \ln(cx^3 + 1)}{10} - \frac{bx^5 \ln(1 - cx^3)}{10}$$

input `int(x^4*(a + b*atanh(c*x^3)),x)`

output

```
(a*x^5)/5 + (b*log(1 - c^(2/3)*x^2))/(10*c^(5/3)) + (3*b*x^2)/(10*c) - (log(2*c^(2/3)*x^2 - 3^(1/2)*1i + 1)*(b - 3^(1/2)*b*1i))/(20*c^(5/3)) - (log(3^(1/2)*1i + 2*c^(2/3)*x^2 + 1)*(b + 3^(1/2)*b*1i))/(20*c^(5/3)) + (b*x^5*log(c*x^3 + 1))/10 - (b*x^5*log(1 - c*x^3))/10
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.13

$$\int x^4(a + \operatorname{barctanh}(cx^3)) dx = \frac{-2\sqrt{3} \operatorname{atan}\left(\frac{2c^{1/3}x-1}{\sqrt{3}}\right)b + 2\sqrt{3} \operatorname{atan}\left(\frac{2c^{1/3}x+1}{\sqrt{3}}\right)b + 4c^{5/3} \operatorname{atanh}(cx^3)bx^5 + 2 \operatorname{atanh}(cx^3)b + 4c^{5/3}ax^5 + 6c^{2/3}bx}{20c^{5/3}}$$

input `int(x^4*(a+b*atanh(c*x^3)),x)`

output

```
( - 2*sqrt(3)*atan((2*c**(1/3)*x - 1)/sqrt(3))*b + 2*sqrt(3)*atan((2*c**(1/3)*x + 1)/sqrt(3))*b + 4*c**(2/3)*atanh(c*x**3)*b*c*x**5 + 2*atanh(c*x**3)*b + 4*c**(2/3)*a*c*x**5 + 6*c**(2/3)*b*x**2 - 2*log(c**(2/3)*x**2 - c**(1/3)*x + 1)*b + log(c**(2/3)*x + c**(1/3))*b + 3*log(c**(2/3)*x - c**(1/3))*b)/(20*c**(2/3)*c)
```

3.113 $\int x(a + b \operatorname{arctanh}(cx^3)) dx$

Optimal result	959
Mathematica [A] (verified)	960
Rubi [A] (verified)	960
Maple [A] (verified)	964
Fricas [B] (verification not implemented)	965
Sympy [F(-1)]	966
Maxima [A] (verification not implemented)	966
Giac [A] (verification not implemented)	967
Mupad [B] (verification not implemented)	968
Reduce [B] (verification not implemented)	968

Optimal result

Integrand size = 12, antiderivative size = 131

$$\int x(a + b \operatorname{arctanh}(cx^3)) dx = -\frac{\sqrt{3}b \arctan\left(\frac{1-2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{4c^{2/3}} + \frac{\sqrt{3}b \arctan\left(\frac{1+2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{4c^{2/3}} - \frac{b \operatorname{arctanh}(\sqrt[3]{Cx})}{2c^{2/3}} + \frac{1}{2}x^2(a + b \operatorname{arctanh}(cx^3)) - \frac{b \operatorname{arctanh}\left(\frac{\sqrt[3]{Cx}}{1+c^{2/3}x^2}\right)}{4c^{2/3}}$$

output

```
-1/4*3^(1/2)*b*arctan(1/3*(1-2*c^(1/3)*x)*3^(1/2))/c^(2/3)+1/4*3^(1/2)*b*
rctan(1/3*(1+2*c^(1/3)*x)*3^(1/2))/c^(2/3)-1/2*b*arctanh(c^(1/3)*x)/c^(2/3
)+1/2*x^2*(a+b*arctanh(c*x^3))-1/4*b*arctanh(c^(1/3)*x/(1+c^(2/3)*x^2))/c^
(2/3)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.43

$$\int x(a + b \operatorname{arctanh}(cx^3)) dx = \frac{ax^2}{2} + \frac{\sqrt{3}b \operatorname{arctan}\left(\frac{-1+2\sqrt[3]{cx}}{\sqrt{3}}\right)}{4c^{2/3}} + \frac{\sqrt{3}b \operatorname{arctan}\left(\frac{1+2\sqrt[3]{cx}}{\sqrt{3}}\right)}{4c^{2/3}} \\ + \frac{1}{2}bx^2 \operatorname{arctanh}(cx^3) + \frac{b \log(1 - \sqrt[3]{cx})}{4c^{2/3}} - \frac{b \log(1 + \sqrt[3]{cx})}{4c^{2/3}} \\ + \frac{b \log(1 - \sqrt[3]{cx} + c^{2/3}x^2)}{8c^{2/3}} - \frac{b \log(1 + \sqrt[3]{cx} + c^{2/3}x^2)}{8c^{2/3}}$$

input `Integrate[x*(a + b*ArcTanh[c*x^3]),x]`

output $(a*x^2)/2 + (\operatorname{Sqrt}[3]*b*\operatorname{ArcTan}[(-1 + 2*c^{(1/3)}*x)/\operatorname{Sqrt}[3]])/(4*c^{(2/3)}) + (\operatorname{Sqrt}[3]*b*\operatorname{ArcTan}[(1 + 2*c^{(1/3)}*x)/\operatorname{Sqrt}[3]])/(4*c^{(2/3)}) + (b*x^2*\operatorname{ArcTanh}[c*x^3])/2 + (b*\operatorname{Log}[1 - c^{(1/3)}*x])/(4*c^{(2/3)}) - (b*\operatorname{Log}[1 + c^{(1/3)}*x])/(4*c^{(2/3)}) + (b*\operatorname{Log}[1 - c^{(1/3)}*x + c^{(2/3)}*x^2])/(8*c^{(2/3)}) - (b*\operatorname{Log}[1 + c^{(1/3)}*x + c^{(2/3)}*x^2])/(8*c^{(2/3)})$

Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.34, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {6452, 825, 27, 219, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(a + b \operatorname{arctanh}(cx^3)) dx \\ \downarrow 6452 \\ \frac{1}{2}x^2(a + b \operatorname{arctanh}(cx^3)) - \frac{3}{2}bc \int \frac{x^4}{1 - c^2x^6} dx \\ \downarrow 825$$

$$\frac{1}{2}x^2(a + \operatorname{barctanh}(cx^3)) - \frac{3}{2}bc \left(\frac{\int \frac{1}{1-c^{2/3}x^2} dx}{3c^{4/3}} + \frac{\int -\frac{\sqrt[3]{cx+1}}{2(c^{2/3}x^2 - \sqrt[3]{cx+1})} dx}{3c^{4/3}} + \frac{\int -\frac{1-\sqrt[3]{cx}}{2(c^{2/3}x^2 + \sqrt[3]{cx+1})} dx}{3c^{4/3}} \right)$$

↓ 27

$$\frac{1}{2}x^2(a + \operatorname{barctanh}(cx^3)) - \frac{3}{2}bc \left(\frac{\int \frac{1}{1-c^{2/3}x^2} dx}{3c^{4/3}} - \frac{\int \frac{\sqrt[3]{cx+1}}{c^{2/3}x^2 - \sqrt[3]{cx+1}} dx}{6c^{4/3}} - \frac{\int \frac{1-\sqrt[3]{cx}}{c^{2/3}x^2 + \sqrt[3]{cx+1}} dx}{6c^{4/3}} \right)$$

↓ 219

$$\frac{1}{2}x^2(a + \operatorname{barctanh}(cx^3)) - \frac{3}{2}bc \left(-\frac{\int \frac{\sqrt[3]{cx+1}}{c^{2/3}x^2 - \sqrt[3]{cx+1}} dx}{6c^{4/3}} - \frac{\int \frac{1-\sqrt[3]{cx}}{c^{2/3}x^2 + \sqrt[3]{cx+1}} dx}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{cx})}{3c^{5/3}} \right)$$

↓ 1142

$$\frac{1}{2}x^2(a + \operatorname{barctanh}(cx^3)) - \frac{3}{2}bc \left(-\frac{\frac{3}{2} \int \frac{1}{c^{2/3}x^2 - \sqrt[3]{cx+1}} dx + \frac{\int -\frac{\sqrt[3]{c}(1-2\sqrt[3]{cx})}{c^{2/3}x^2 - \sqrt[3]{cx+1}} dx}{2\sqrt[3]{c}}}{6c^{4/3}} - \frac{\frac{3}{2} \int \frac{1}{c^{2/3}x^2 + \sqrt[3]{cx+1}} dx - \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{cx+1})}{c^{2/3}x^2 + \sqrt[3]{cx+1}} dx}{2\sqrt[3]{c}}}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{cx})}{3c^{5/3}} \right)$$

↓ 25

$$\frac{1}{2}x^2(a + \operatorname{barctanh}(cx^3)) - \frac{3}{2}bc \left(-\frac{\frac{3}{2} \int \frac{1}{c^{2/3}x^2 - \sqrt[3]{cx+1}} dx - \frac{\int \frac{\sqrt[3]{c}(1-2\sqrt[3]{cx})}{c^{2/3}x^2 - \sqrt[3]{cx+1}} dx}{2\sqrt[3]{c}}}{6c^{4/3}} - \frac{\frac{3}{2} \int \frac{1}{c^{2/3}x^2 + \sqrt[3]{cx+1}} dx - \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{cx+1})}{c^{2/3}x^2 + \sqrt[3]{cx+1}} dx}{2\sqrt[3]{c}}}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{cx})}{3c^{5/3}} \right)$$

↓ 27

$$\frac{3}{2}bc \left(\frac{\frac{1}{2}x^2(a + \operatorname{arctanh}(cx^3)) - \frac{3}{2} \int \frac{1}{c^{2/3}x^2 - \sqrt[3]{cx+1}} dx - \frac{1}{2} \int \frac{1-2\sqrt[3]{cx}}{c^{2/3}x^2 - \sqrt[3]{cx+1}} dx}{6c^{4/3}} - \frac{\frac{3}{2} \int \frac{1}{c^{2/3}x^2 + \sqrt[3]{cx+1}} dx - \frac{1}{2} \int \frac{2\sqrt[3]{cx+1}}{c^{2/3}x^2 + \sqrt[3]{cx+1}} dx}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{cx})}{3c^{5/3}} \right)$$

↓ 1082

$$\frac{3}{2}bc \left(\frac{\frac{1}{2}x^2(a + \operatorname{arctanh}(cx^3)) - \frac{3 \int \frac{1}{(1-2\sqrt[3]{cx})^2} d(1-2\sqrt[3]{cx})}{\sqrt[3]{c}}}{6c^{4/3}} - \frac{\frac{1}{2} \int \frac{1-2\sqrt[3]{cx}}{c^{2/3}x^2 - \sqrt[3]{cx+1}} dx}{6c^{4/3}} - \frac{\frac{1}{2} \int \frac{2\sqrt[3]{cx+1}}{c^{2/3}x^2 + \sqrt[3]{cx+1}} dx - \frac{3 \int \frac{1}{(2\sqrt[3]{cx+1})^2} d(2\sqrt[3]{cx+1})}{\sqrt[3]{c}}}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{cx})}{3c^{5/3}} \right)$$

↓ 217

$$\frac{3}{2}bc \left(\frac{\frac{1}{2}x^2(a + \operatorname{arctanh}(cx^3)) - \frac{-\frac{1}{2} \int \frac{1-2\sqrt[3]{cx}}{c^{2/3}x^2 - \sqrt[3]{cx+1}} dx - \frac{\sqrt{3} \arctan\left(\frac{1-2\sqrt[3]{cx}}{\sqrt{3}}\right)}{\sqrt[3]{c}}}{6c^{4/3}} - \frac{\frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{cx+1}}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{1}{2} \int \frac{2\sqrt[3]{cx+1}}{c^{2/3}x^2 + \sqrt[3]{cx+1}} dx}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{cx})}{3c^{5/3}} \right)$$

↓ 1103

$$\frac{3}{2}bc \left(\frac{\frac{1}{2}x^2(a + \operatorname{arctanh}(cx^3)) - \frac{\frac{\log(c^{2/3}x^2 - \sqrt[3]{cx+1})}{2\sqrt[3]{c}} - \frac{\sqrt{3} \arctan\left(\frac{1-2\sqrt[3]{cx}}{\sqrt{3}}\right)}{\sqrt[3]{c}}}{6c^{4/3}} - \frac{\frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{cx+1}}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{\log(c^{2/3}x^2 + \sqrt[3]{cx+1})}{2\sqrt[3]{c}}}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{cx})}{3c^{5/3}} \right)$$

input

```
Int[x*(a + b*ArcTanh[c*x^3]),x]
```

output

$$\begin{aligned} & (x^2(a + b \operatorname{ArcTanh}[c x^3]))/2 - (3 b c (\operatorname{ArcTanh}[c^{1/3} x]/(3 c^{5/3})) - \\ & - ((\sqrt{3} \operatorname{ArcTan}[(1 - 2 c^{1/3} x)/\sqrt{3}])/c^{1/3}) + \operatorname{Log}[1 - c^{1/3} x \\ & + c^{2/3} x^2]/(2 c^{1/3}))/ (6 c^{4/3}) - ((\sqrt{3} \operatorname{ArcTan}[(1 + 2 c^{1/3} \\ &) x]/\sqrt{3}))/c^{1/3} - \operatorname{Log}[1 + c^{1/3} x + c^{2/3} x^2]/(2 c^{1/3}))/ (6 c^{4/3}))/2 \end{aligned}$$

Defintions of rubi rules used

rule 25

$$\operatorname{Int}[-(F x), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F x, x], x]$$

rule 27

$$\operatorname{Int}[(a)(F x), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[a \operatorname{Int}[F x, x], x] /; \operatorname{FreeQ}[a, x] \&\& !\operatorname{MatchQ}[F x, (b)(G x) /; \operatorname{FreeQ}[b, x]]$$

rule 217

$$\operatorname{Int}[(a) + (b)(x)^2)^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(-\operatorname{Rt}[-a, 2] \operatorname{Rt}[-b, 2])^{-1} \operatorname{ArcTan}[\operatorname{Rt}[-b, 2] (x/\operatorname{Rt}[-a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&\& \operatorname{PosQ}[a/b] \&\& (\operatorname{LtQ}[a, 0] \parallel \operatorname{LtQ}[b, 0])$$

rule 219

$$\operatorname{Int}[(a) + (b)(x)^2)^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2] \operatorname{Rt}[-b, 2])) \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2] (x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&\& \operatorname{NegQ}[a/b] \&\& (\operatorname{GtQ}[a, 0] \parallel \operatorname{LtQ}[b, 0])$$

rule 825

$$\operatorname{Int}[(x)^{(m)} / ((a) + (b)(x)^{(n)}), x_{\text{Symbol}}] \rightarrow \operatorname{Module}[\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, n]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, n]], k, u\}, \operatorname{Simp}[u = \operatorname{Int}[(r \operatorname{Cos}[2 k m (\operatorname{Pi}/n)] - s \operatorname{Cos}[2 k (m + 1) (\operatorname{Pi}/n)] x) / (r^2 - 2 r s \operatorname{Cos}[2 k (\operatorname{Pi}/n)] x + s^2 x^2), x] + \operatorname{Int}[(r \operatorname{Cos}[2 k m (\operatorname{Pi}/n)] + s \operatorname{Cos}[2 k (m + 1) (\operatorname{Pi}/n)] x) / (r^2 + 2 r s \operatorname{Cos}[2 k (\operatorname{Pi}/n)] x + s^2 x^2), x]; 2 (r^{(m + 2)} / (a n s^m)) \operatorname{Int}[1 / (r^2 - s^2 x^2), x] + 2 (r^{(m + 1)} / (a n s^m)) \operatorname{Sum}[u, \{k, 1, (n - 2)/4\}], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&\& \operatorname{IGtQ}[(n - 2)/4, 0] \&\& \operatorname{IGtQ}[m, 0] \&\& \operatorname{LtQ}[m, n - 1] \&\& \operatorname{NegQ}[a/b]$$

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.35

method	result
default	$\frac{ax^2}{2} + \frac{bx^2 \operatorname{arctanh}(cx^3)}{2} + \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4c\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{b \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{8c\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{-2x}{\left(\frac{1}{c}\right)^{\frac{1}{3}} + 1\right)}\right)}{4c\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{b \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4c\left(\frac{1}{c}\right)^{\frac{1}{3}}}$
parts	$\frac{ax^2}{2} + \frac{bx^2 \operatorname{arctanh}(cx^3)}{2} + \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4c\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{b \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{8c\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{-2x}{\left(\frac{1}{c}\right)^{\frac{1}{3}} + 1\right)}\right)}{4c\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{b \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4c\left(\frac{1}{c}\right)^{\frac{1}{3}}}$
risch	$\frac{x^2 b \ln(cx^3 + 1)}{4} + \frac{ax^2}{2} - \frac{bx^2 \ln(-cx^3 + 1)}{4} + \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4c\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{b \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{8c\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{-2x}{\left(\frac{1}{c}\right)^{\frac{1}{3}} + 1\right)}\right)}{4c\left(\frac{1}{c}\right)^{\frac{1}{3}}}$

input

```
int(x*(a+b*arctanh(c*x^3)),x,method=_RETURNVERBOSE)
```

output

```
1/2*a*x^2+1/2*b*x^2*arctanh(c*x^3)+1/4*b/c/(1/c)^(1/3)*ln(x-(1/c)^(1/3))-1/8*b/c/(1/c)^(1/3)*ln(x^2+(1/c)^(1/3)*x+(1/c)^(2/3))+1/4*b*3^(1/2)/c/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x+1))-1/4*b/c/(1/c)^(1/3)*ln(x+(1/c)^(1/3))+1/8*b/c/(1/c)^(1/3)*ln(x^2-(1/c)^(1/3)*x+(1/c)^(2/3))+1/4*b*3^(1/2)/c/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x-1))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 238 vs. 2(97) = 194.

Time = 0.08 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.82

$$\int x(a + b \operatorname{arctanh}(cx^3)) dx$$

$$2bc^2x^2 \log\left(\frac{-cx^3+1}{cx^3-1}\right) + 4ac^2x^2 + 2\sqrt{3}bc\sqrt{-(-c^2)^{\frac{1}{3}}} \operatorname{arctan}\left(\frac{\sqrt{3}\left(2cx+(-c^2)^{\frac{1}{3}}\right)\sqrt{-(-c^2)^{\frac{1}{3}}}}{3c}\right) + 2\sqrt{3}b(c^2)^{\frac{1}{6}}ca$$

input `integrate(x*(a+b*arctanh(c*x^3)),x, algorithm="fricas")`

output
$$\frac{1}{8}*(2*b*c^2*x^2*\log(-(c*x^3 + 1)/(c*x^3 - 1)) + 4*a*c^2*x^2 + 2*\sqrt{3}*b*c*\sqrt{-(-c^2)^{(1/3)}}*\arctan(1/3*\sqrt{3}*(2*c*x + (-c^2)^{(1/3)})*\sqrt{-(-c^2)^{(1/3)}}/c) + 2*\sqrt{3}*b*(c^2)^{(1/6)}*c*\arctan(1/3*\sqrt{3}*(c^2)^{(1/6)}*(2*c*x + (c^2)^{(1/3)})/c) + (-c^2)^{(2/3)}*b*\log(c^2*x^2 + (-c^2)^{(1/3)}*c*x + (-c^2)^{(2/3)}) - b*(c^2)^{(2/3)}*\log(c^2*x^2 + (c^2)^{(1/3)}*c*x + (c^2)^{(2/3)}) - 2*(-c^2)^{(2/3)}*b*\log(c*x - (-c^2)^{(1/3)}) + 2*b*(c^2)^{(2/3)}*\log(c*x - (c^2)^{(1/3)}))/c^2$$

Sympy [F(-1)]

Timed out.

$$\int x(a + b\operatorname{arctanh}(cx^3)) dx = \text{Timed out}$$

input `integrate(x*(a+b*atanh(c*x**3)),x)`

output Timed out

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.18

$$\int x(a + b\operatorname{arctanh}(cx^3)) dx = \frac{1}{2}ax^2 + \frac{1}{8} \left(4x^2 \operatorname{arctanh}(cx^3) + c \left(\frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{\frac{2}{3}}x + c^{\frac{1}{3}})}{3c^{\frac{1}{3}}}\right)}{c^{\frac{5}{3}}} + \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{\frac{2}{3}}x - c^{\frac{1}{3}})}{3c^{\frac{1}{3}}}\right)}{c^{\frac{5}{3}}} - \frac{\log\left(c^{\frac{2}{3}}x^2 + \dots\right)}{c^{\frac{5}{3}}}\right) \right)$$

input `integrate(x*(a+b*arctanh(c*x^3)),x, algorithm="maxima")`

output

```
1/2*a*x^2 + 1/8*(4*x^2*arctanh(c*x^3) + c*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2
*c^(2/3)*x + c^(1/3))/c^(1/3))/c^(5/3) + 2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c
^(2/3)*x - c^(1/3))/c^(1/3))/c^(5/3) - log(c^(2/3)*x^2 + c^(1/3)*x + 1)/c^
(5/3) + log(c^(2/3)*x^2 - c^(1/3)*x + 1)/c^(5/3) - 2*log((c^(1/3)*x + 1)/c
^(1/3))/c^(5/3) + 2*log((c^(1/3)*x - 1)/c^(1/3))/c^(5/3))*b
```

Giac [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.32

$$\int x(a + b \operatorname{arctanh}(cx^3)) dx = \frac{1}{4} bx^2 \log\left(-\frac{cx^3 + 1}{cx^3 - 1}\right) + \frac{1}{2} ax^2$$

$$+ \frac{\sqrt{3}bc \operatorname{arctan}\left(\frac{1}{3}\sqrt{3}\left(2x + \frac{1}{|c|^{1/3}}\right)|c|^{1/3}\right)}{4|c|^{5/3}}$$

$$+ \frac{\sqrt{3}bc \operatorname{arctan}\left(\frac{1}{3}\sqrt{3}\left(2x - \frac{1}{|c|^{1/3}}\right)|c|^{1/3}\right)}{4|c|^{5/3}}$$

$$- \frac{bc \log\left(x^2 + \frac{x}{|c|^{1/3}} + \frac{1}{|c|^{2/3}}\right)}{8|c|^{5/3}} + \frac{bc \log\left(x^2 - \frac{x}{|c|^{1/3}} + \frac{1}{|c|^{2/3}}\right)}{8|c|^{5/3}}$$

$$- \frac{bc \log\left(\left|x + \frac{1}{|c|^{1/3}}\right|\right)}{4|c|^{5/3}} + \frac{bc \log\left(\left|x - \frac{1}{|c|^{1/3}}\right|\right)}{4|c|^{5/3}}$$

input

```
integrate(x*(a+b*arctanh(c*x^3)),x, algorithm="giac")
```

output

```
1/4*b*x^2*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 1/2*a*x^2 + 1/4*sqrt(3)*b*c*arct
an(1/3*sqrt(3)*(2*x + 1/abs(c)^(1/3))*abs(c)^(1/3))/abs(c)^(5/3) + 1/4*sqrt
(3)*b*c*arctan(1/3*sqrt(3)*(2*x - 1/abs(c)^(1/3))*abs(c)^(1/3))/abs(c)^(5
/3) - 1/8*b*c*log(x^2 + x/abs(c)^(1/3) + 1/abs(c)^(2/3))/abs(c)^(5/3) + 1/
8*b*c*log(x^2 - x/abs(c)^(1/3) + 1/abs(c)^(2/3))/abs(c)^(5/3) - 1/4*b*c*lo
g(abs(x + 1/abs(c)^(1/3)))/abs(c)^(5/3) + 1/4*b*c*log(abs(x - 1/abs(c)^(1/
3)))/abs(c)^(5/3)
```

Mupad [B] (verification not implemented)

Time = 4.01 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.90

$$\int x(a + b \operatorname{arctanh}(cx^3)) dx$$

$$= \frac{ax^2}{2} + \frac{b \left(-\frac{\operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}-i)}{2}\right)}{2} + \frac{\operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}+i)}{2}\right)}{2} + \operatorname{atan}(c^{1/3}xi) \right) i}{2c^{2/3}}$$

$$+ \frac{bx^2 \ln(cx^3 + 1)}{4} - \frac{bx^2 \ln(1 - cx^3)}{4}$$

$$+ \frac{\sqrt{3}b \left(\operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}-i)}{2}\right) + \operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}+i)}{2}\right) \right)}{4c^{2/3}}$$

input `int(x*(a + b*atanh(c*x^3)),x)`output `(a*x^2)/2 + (b*(atan((c^(1/3)*x*(3^(1/2) + 1i))/2)/2 - atan((c^(1/3)*x*(3^(1/2) - 1i))/2)/2 + atan(c^(1/3)*x*1i))*i)/(2*c^(2/3)) + (b*x^2*log(c*x^3 + 1))/4 - (b*x^2*log(1 - c*x^3))/4 + (3^(1/2)*b*(atan((c^(1/3)*x*(3^(1/2) - 1i))/2) + atan((c^(1/3)*x*(3^(1/2) + 1i))/2)))/(4*c^(2/3))`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.80

$$\int x(a + b \operatorname{arctanh}(cx^3)) dx$$

$$= \frac{2\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x-1}{\sqrt{3}}\right) b + 2\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x+1}{\sqrt{3}}\right) b + 4c^{\frac{2}{3}} \operatorname{atanh}(cx^3) b x^2 + 2 \operatorname{atanh}(cx^3) b + 4c^{\frac{2}{3}} a x^2 - 3 \log\left(c^{\frac{2}{3}}\right)}{8c^{\frac{2}{3}}}$$

input `int(x*(a+b*atanh(c*x^3)),x)`

output

```
(2*sqrt(3)*atan((2*c**(1/3)*x - 1)/sqrt(3))*b + 2*sqrt(3)*atan((2*c**(1/3)
*x + 1)/sqrt(3))*b + 4*c**(2/3)*atanh(c*x**3)*b*x**2 + 2*atanh(c*x**3)*b +
4*c**(2/3)*a*x**2 - 3*log(c**(2/3)*x + c**(1/3))*b + 3*log(c**(2/3)*x - c
**(1/3))*b)/(8*c**(2/3))
```

3.114 $\int \frac{a+b\operatorname{arctanh}(cx^3)}{x^2} dx$

Optimal result	970
Mathematica [A] (verified)	970
Rubi [A] (verified)	971
Maple [A] (verified)	974
Fricas [A] (verification not implemented)	975
Sympy [F(-1)]	975
Maxima [A] (verification not implemented)	976
Giac [A] (verification not implemented)	976
Mupad [B] (verification not implemented)	977
Reduce [B] (verification not implemented)	977

Optimal result

Integrand size = 14, antiderivative size = 104

$$\int \frac{a + b\operatorname{arctanh}(cx^3)}{x^2} dx = \frac{1}{2}\sqrt{3}b\sqrt[3]{c} \arctan\left(\frac{1 + 2c^{2/3}x^2}{\sqrt{3}}\right) - \frac{a + b\operatorname{arctanh}(cx^3)}{x} - \frac{1}{2}b\sqrt[3]{c} \log(1 - c^{2/3}x^2) + \frac{1}{4}b\sqrt[3]{c} \log(1 + c^{2/3}x^2 + c^{4/3}x^4)$$

output

```
1/2*3^(1/2)*b*c^(1/3)*arctan(1/3*(1+2*c^(2/3)*x^2)*3^(1/2))-(a+b*arctanh(c*x^3))/x-1/2*b*c^(1/3)*ln(1-c^(2/3)*x^2)+1/4*b*c^(1/3)*ln(1+c^(2/3)*x^2+c^(4/3)*x^4)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.76

$$\int \frac{a + b\operatorname{arctanh}(cx^3)}{x^2} dx = -\frac{a}{x} + \frac{1}{2}\sqrt{3}b\sqrt[3]{c} \arctan\left(\frac{-1 + 2\sqrt[3]{cx}}{\sqrt{3}}\right) - \frac{1}{2}\sqrt{3}b\sqrt[3]{c} \arctan\left(\frac{1 + 2\sqrt[3]{cx}}{\sqrt{3}}\right) - \frac{b\operatorname{arctanh}(cx^3)}{x} - \frac{1}{2}b\sqrt[3]{c} \log(1 - \sqrt[3]{cx}) - \frac{1}{2}b\sqrt[3]{c} \log(1 + \sqrt[3]{cx}) + \frac{1}{4}b\sqrt[3]{c} \log(1 - \sqrt[3]{cx} + c^{2/3}x^2) + \frac{1}{4}b\sqrt[3]{c} \log(1 + \sqrt[3]{cx} + c^{2/3}x^2)$$

input `Integrate[(a + b*ArcTanh[c*x^3])/x^2,x]`

output $-(a/x) + (\text{Sqrt}[3]*b*c^{(1/3)}*\text{ArcTan}[(-1 + 2*c^{(1/3)}*x)/\text{Sqrt}[3]])/2 - (\text{Sqrt}[3]*b*c^{(1/3)}*\text{ArcTan}[(1 + 2*c^{(1/3)}*x)/\text{Sqrt}[3]])/2 - (b*\text{ArcTanh}[c*x^3])/x - (b*c^{(1/3)}*\text{Log}[1 - c^{(1/3)}*x])/2 - (b*c^{(1/3)}*\text{Log}[1 + c^{(1/3)}*x])/2 + (b*c^{(1/3)}*\text{Log}[1 - c^{(1/3)}*x + c^{(2/3)}*x^2])/4 + (b*c^{(1/3)}*\text{Log}[1 + c^{(1/3)}*x + c^{(2/3)}*x^2])/4$

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.06, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {6452, 807, 750, 16, 1142, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}(cx^3)}{x^2} dx \\ & \quad \downarrow 6452 \\ & 3bc \int \frac{x}{1 - c^2 x^6} dx - \frac{a + b \operatorname{arctanh}(cx^3)}{x} \\ & \quad \downarrow 807 \\ & \frac{3}{2}bc \int \frac{1}{1 - c^2 x^6} dx^2 - \frac{a + b \operatorname{arctanh}(cx^3)}{x} \\ & \quad \downarrow 750 \\ & \frac{3}{2}bc \left(\frac{1}{3} \int \frac{1}{1 - c^{2/3} x^2} dx^2 + \frac{1}{3} \int \frac{c^{2/3} x^2 + 2}{c^{4/3} x^4 + c^{2/3} x^2 + 1} dx^2 \right) - \frac{a + b \operatorname{arctanh}(cx^3)}{x} \\ & \quad \downarrow 16 \\ & \frac{3}{2}bc \left(\frac{1}{3} \int \frac{c^{2/3} x^2 + 2}{c^{4/3} x^4 + c^{2/3} x^2 + 1} dx^2 - \frac{\log(1 - c^{2/3} x^2)}{3c^{2/3}} \right) - \frac{a + b \operatorname{arctanh}(cx^3)}{x} \\ & \quad \downarrow 1142 \end{aligned}$$

$$\begin{aligned}
& \frac{3}{2}bc \left(\frac{1}{3} \left(\frac{3}{2} \int \frac{1}{c^{4/3}x^4 + c^{2/3}x^2 + 1} dx^2 + \frac{\int \frac{c^{2/3}(2c^{2/3}x^2+1)}{c^{4/3}x^4+c^{2/3}x^2+1} dx^2}{2c^{2/3}} \right) - \frac{\log(1 - c^{2/3}x^2)}{3c^{2/3}} \right) - \\
& \qquad \qquad \qquad \frac{x}{a + \operatorname{barctanh}(cx^3)} \\
& \qquad \qquad \qquad \downarrow 27 \\
& \frac{3}{2}bc \left(\frac{1}{3} \left(\frac{3}{2} \int \frac{1}{c^{4/3}x^4 + c^{2/3}x^2 + 1} dx^2 + \frac{1}{2} \int \frac{2c^{2/3}x^2 + 1}{c^{4/3}x^4 + c^{2/3}x^2 + 1} dx^2 \right) - \frac{\log(1 - c^{2/3}x^2)}{3c^{2/3}} \right) - \\
& \qquad \qquad \qquad \frac{x}{a + \operatorname{barctanh}(cx^3)} \\
& \qquad \qquad \qquad \downarrow 1082 \\
& \frac{3}{2}bc \left(\frac{1}{3} \left(\frac{1}{2} \int \frac{2c^{2/3}x^2 + 1}{c^{4/3}x^4 + c^{2/3}x^2 + 1} dx^2 - \frac{3 \int \frac{1}{-x^4-3} d(2c^{2/3}x^2 + 1)}{c^{2/3}} \right) - \frac{\log(1 - c^{2/3}x^2)}{3c^{2/3}} \right) - \\
& \qquad \qquad \qquad \frac{x}{a + \operatorname{barctanh}(cx^3)} \\
& \qquad \qquad \qquad \downarrow 217 \\
& \frac{3}{2}bc \left(\frac{1}{3} \left(\frac{1}{2} \int \frac{2c^{2/3}x^2 + 1}{c^{4/3}x^4 + c^{2/3}x^2 + 1} dx^2 + \frac{\sqrt{3} \arctan\left(\frac{2c^{2/3}x^2+1}{\sqrt{3}}\right)}{c^{2/3}} \right) - \frac{\log(1 - c^{2/3}x^2)}{3c^{2/3}} \right) - \\
& \qquad \qquad \qquad \frac{x}{a + \operatorname{barctanh}(cx^3)} \\
& \qquad \qquad \qquad \downarrow 1103 \\
& \frac{3}{2}bc \left(\frac{1}{3} \left(\frac{\sqrt{3} \arctan\left(\frac{2c^{2/3}x^2+1}{\sqrt{3}}\right)}{c^{2/3}} + \frac{\log(c^{4/3}x^4 + c^{2/3}x^2 + 1)}{2c^{2/3}} \right) - \frac{\log(1 - c^{2/3}x^2)}{3c^{2/3}} \right) - \\
& \qquad \qquad \qquad \frac{x}{a + \operatorname{barctanh}(cx^3)}
\end{aligned}$$

input

```
Int[(a + b*ArcTanh[c*x^3])/x^2,x]
```

output

```
-((a + b*ArcTanh[c*x^3])/x) + (3*b*c*(-1/3*Log[1 - c^(2/3)*x^2]/c^(2/3) +
((Sqrt[3]*ArcTan[(1 + 2*c^(2/3)*x^2)/Sqrt[3]])/c^(2/3) + Log[1 + c^(2/3)*x
^2 + c^(4/3)*x^4]/(2*c^(2/3)))/3)/2
```

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] /; \text{FreeQ}[b, x]$
- rule 217 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 750 $\text{Int}[(a_)+(b_)*(x_)^3)^{-1}, x_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$
- rule 807 $\text{Int}[(x_)^{(m_)}*((a_)+(b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{With}[\{k = \text{GCD}[m + 1, n]\}, \text{Simp}[1/k \text{ Subst}[\text{Int}[x^{((m + 1)/k - 1)*(a + b*x^{(n/k)})^p}, x], x, x^k], x] /; k \neq 1] /; \text{FreeQ}[\{a, b, p\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$
- rule 1082 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$
- rule 1142 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{ Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{ Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.01

method	result
default	$-\frac{a}{x} - \frac{b \operatorname{arctanh}(cx^3)}{x} - \frac{b \ln\left(x^2 - \left(\frac{1}{c^2}\right)^{\frac{1}{3}}\right)}{2c\left(\frac{1}{c^2}\right)^{\frac{2}{3}}} + \frac{b \ln\left(x^4 + \left(\frac{1}{c^2}\right)^{\frac{1}{3}}x^2 + \left(\frac{1}{c^2}\right)^{\frac{2}{3}}\right)}{4c\left(\frac{1}{c^2}\right)^{\frac{2}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{-2x^2}{\left(\frac{1}{c^2}\right)^{\frac{1}{3}} + 1\right)}\right)}{2c\left(\frac{1}{c^2}\right)^{\frac{2}{3}}}$
parts	$-\frac{a}{x} - \frac{b \operatorname{arctanh}(cx^3)}{x} - \frac{b \ln\left(x^2 - \left(\frac{1}{c^2}\right)^{\frac{1}{3}}\right)}{2c\left(\frac{1}{c^2}\right)^{\frac{2}{3}}} + \frac{b \ln\left(x^4 + \left(\frac{1}{c^2}\right)^{\frac{1}{3}}x^2 + \left(\frac{1}{c^2}\right)^{\frac{2}{3}}\right)}{4c\left(\frac{1}{c^2}\right)^{\frac{2}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{-2x^2}{\left(\frac{1}{c^2}\right)^{\frac{1}{3}} + 1\right)}\right)}{2c\left(\frac{1}{c^2}\right)^{\frac{2}{3}}}$
risch	$-\frac{b \ln(cx^3+1)}{2x} - \frac{a}{x} + \frac{b \ln(-cx^3+1)}{2x} - \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{2\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{4\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{-2x}{\left(\frac{1}{c}\right)^{\frac{1}{3}} + 1\right)}\right)}{2\left(\frac{1}{c}\right)^{\frac{1}{3}}}$

input

```
int((a+b*arctanh(c*x^3))/x^2,x,method=_RETURNVERBOSE)
```

output

```
-a/x-b/x*arctanh(c*x^3)-1/2*b/c/(1/c^2)^(2/3)*ln(x^2-(1/c^2)^(1/3))+1/4*b/c/(1/c^2)^(2/3)*ln(x^4+(1/c^2)^(1/3)*x^2+(1/c^2)^(2/3))+1/2*b/c/(1/c^2)^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c^2)^(1/3)*x^2+1))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.12

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^2} dx = \frac{2\sqrt{3}b(-c)^{\frac{1}{3}} x \arctan\left(\frac{2}{3}\sqrt{3}(-c)^{\frac{2}{3}}x^2 + \frac{1}{3}\sqrt{3}\right) + b(-c)^{\frac{1}{3}} x \log\left(c^2x^4 - (-c)^{\frac{1}{3}}cx^2 + (-c)^{\frac{2}{3}}\right) - 2b(-c)^{\frac{1}{3}}}{4x}$$

input `integrate((a+b*arctanh(c*x^3))/x^2,x, algorithm="fricas")`

output

```
-1/4*(2*sqrt(3)*b*(-c)^(1/3)*x*arctan(2/3*sqrt(3)*(-c)^(2/3)*x^2 + 1/3*sqrt(3)) + b*(-c)^(1/3)*x*log(c^2*x^4 - (-c)^(1/3)*c*x^2 + (-c)^(2/3)) - 2*b*(-c)^(1/3)*x*log(c*x^2 + (-c)^(1/3)) + 2*b*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 4*a)/x
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^2} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**3))/x**2,x)`

output

Timed out

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.90

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^2} dx$$

$$= \frac{1}{4} \left(c \left(\frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{\frac{4}{3}}x^2 + c^{\frac{2}{3}})}{3c^{\frac{2}{3}}}\right)}{c^{\frac{2}{3}}} + \frac{\log\left(c^{\frac{4}{3}}x^4 + c^{\frac{2}{3}}x^2 + 1\right)}{c^{\frac{2}{3}}} - \frac{2 \log\left(\frac{c^{\frac{2}{3}}x^2 - 1}{c^{\frac{2}{3}}}\right)}{c^{\frac{2}{3}}} \right) - \frac{4 \operatorname{arctanh}(cx^3)}{x} \right) b - \frac{a}{x}$$

input `integrate((a+b*arctanh(c*x^3))/x^2,x, algorithm="maxima")`output `1/4*(c*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(4/3)*x^2 + c^(2/3))/c^(2/3))/c^(2/3) + log(c^(4/3)*x^4 + c^(2/3)*x^2 + 1)/c^(2/3) - 2*log((c^(2/3)*x^2 - 1)/c^(2/3))/c^(2/3)) - 4*arctanh(c*x^3)/x)*b - a/x`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.02

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^2} dx$$

$$= \frac{1}{4} bc \left(\frac{2\sqrt{3} \arctan\left(\frac{\frac{1}{3}\sqrt{3}\left(2x^2 + \frac{1}{|c|^{\frac{2}{3}}}\right)|c|^{\frac{2}{3}}}{|c|^{\frac{2}{3}}}\right)}{|c|^{\frac{2}{3}}} + \frac{\log\left(x^4 + \frac{x^2}{|c|^{\frac{2}{3}}} + \frac{1}{|c|^{\frac{4}{3}}}\right)}{|c|^{\frac{2}{3}}} - \frac{2 \log\left(\left|x^2 - \frac{1}{|c|^{\frac{2}{3}}}\right|\right)}{|c|^{\frac{2}{3}}}\right) - \frac{b \log\left(-\frac{cx^3+1}{cx^3-1}\right)}{2x} - \frac{a}{x}$$

input `integrate((a+b*arctanh(c*x^3))/x^2,x, algorithm="giac")`

output

```
1/4*b*c*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*x^2 + 1/abs(c)^(2/3))*abs(c)^(2/3
))/abs(c)^(2/3) + log(x^4 + x^2/abs(c)^(2/3) + 1/abs(c)^(4/3))/abs(c)^(2/3
) - 2*log(abs(x^2 - 1/abs(c)^(2/3)))/abs(c)^(2/3) - 1/2*b*log(-(c*x^3 + 1
)/(c*x^3 - 1))/x - a/x
```

Mupad [B] (verification not implemented)

Time = 5.31 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.12

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^2} dx = \frac{b \ln(1 - cx^3)}{2x} - \frac{bc^{1/3} \ln(1 - c^{2/3}x^2)}{2} - \frac{b \ln(cx^3 + 1)}{2x} - \frac{a}{x} - \frac{bc^{1/3} \ln(-\sqrt{3} - c^{2/3}x^2 2i - i)(-1 + \sqrt{3} 1i)}{4} + \frac{bc^{1/3} \ln(-\sqrt{3} + c^{2/3}x^2 2i + 1i)(1 + \sqrt{3} 1i)}{4}$$

input

```
int((a + b*atanh(c*x^3))/x^2,x)
```

output

```
(b*log(1 - c*x^3))/(2*x) - (b*c^(1/3)*log(1 - c^(2/3)*x^2))/2 - (b*log(c*x
^3 + 1))/(2*x) - a/x - (b*c^(1/3)*log(- 3^(1/2) - c^(2/3)*x^2*2i - 1i)*(3^(
1/2)*1i - 1))/4 + (b*c^(1/3)*log(c^(2/3)*x^2*2i - 3^(1/2) + 1i)*(3^(1/2)*
1i + 1))/4
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.28

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^2} dx = \frac{2\sqrt{3} \operatorname{atan}\left(\frac{2c^{1/3}x-1}{\sqrt{3}}\right)bcx - 2\sqrt{3} \operatorname{atan}\left(\frac{2c^{1/3}x+1}{\sqrt{3}}\right)bcx - 4c^{2/3} \operatorname{atanh}(cx^3)b - 2 \operatorname{atanh}(cx^3)bcx - 4c^{2/3}a + 2 \log\left(\frac{2c^{1/3}x-1}{\sqrt{3}}\right)}{4c^{2/3}x}$$

input

```
int((a+b*atanh(c*x^3))/x^2,x)
```

output

```
(2*sqrt(3)*atan((2*c**(1/3)*x - 1)/sqrt(3))*b*c*x - 2*sqrt(3)*atan((2*c**(1/3)*x + 1)/sqrt(3))*b*c*x - 4*c**(2/3)*atanh(c*x**3)*b - 2*atanh(c*x**3)*b*c*x - 4*c**(2/3)*a + 2*log(c**(2/3)*x**2 - c**(1/3)*x + 1)*b*c*x - log(c**(2/3)*x + c**(1/3))*b*c*x - 3*log(c**(2/3)*x - c**(1/3))*b*c*x)/(4*c**(2/3)*x)
```

3.115 $\int \frac{a+b\operatorname{arctanh}(cx^3)}{x^5} dx$

Optimal result	979
Mathematica [A] (verified)	980
Rubi [A] (verified)	980
Maple [A] (verified)	985
Fricas [A] (verification not implemented)	985
Sympy [F(-1)]	986
Maxima [A] (verification not implemented)	986
Giac [A] (verification not implemented)	987
Mupad [B] (verification not implemented)	988
Reduce [B] (verification not implemented)	988

Optimal result

Integrand size = 14, antiderivative size = 140

$$\int \frac{a + b\operatorname{arctanh}(cx^3)}{x^5} dx$$

$$= -\frac{3bc}{4x} + \frac{1}{8}\sqrt{3}bc^{4/3} \arctan\left(\frac{1 - 2\sqrt[3]{cx}}{\sqrt{3}}\right) - \frac{1}{8}\sqrt{3}bc^{4/3} \arctan\left(\frac{1 + 2\sqrt[3]{cx}}{\sqrt{3}}\right)$$

$$+ \frac{1}{4}bc^{4/3}\operatorname{arctanh}(\sqrt[3]{cx}) - \frac{a + b\operatorname{arctanh}(cx^3)}{4x^4} + \frac{1}{8}bc^{4/3}\operatorname{arctanh}\left(\frac{\sqrt[3]{cx}}{1 + c^{2/3}x^2}\right)$$

output

```
-3/4*b*c/x+1/8*3^(1/2)*b*c^(4/3)*arctan(1/3*(1-2*c^(1/3)*x)*3^(1/2))-1/8*3^(1/2)*b*c^(4/3)*arctan(1/3*(1+2*c^(1/3)*x)*3^(1/2))+1/4*b*c^(4/3)*arctanh(c^(1/3)*x)-1/4*(a+b*arctanh(c*x^3))/x^4+1/8*b*c^(4/3)*arctanh(c^(1/3)*x/(1+c^(2/3)*x^2))
```


Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.40

$$\int \frac{a + \operatorname{barctanh}(cx^3)}{x^5} dx = -\frac{a}{4x^4} - \frac{3bc}{4x} - \frac{1}{8}\sqrt{3}bc^{4/3} \arctan\left(\frac{-1 + 2\sqrt[3]{cx}}{\sqrt{3}}\right) - \frac{1}{8}\sqrt{3}bc^{4/3} \arctan\left(\frac{1 + 2\sqrt[3]{cx}}{\sqrt{3}}\right) - \frac{\operatorname{barctanh}(cx^3)}{4x^4} - \frac{1}{8}bc^{4/3} \log(1 - \sqrt[3]{cx}) + \frac{1}{8}bc^{4/3} \log(1 + \sqrt[3]{cx}) - \frac{1}{16}bc^{4/3} \log(1 - \sqrt[3]{cx} + c^{2/3}x^2) + \frac{1}{16}bc^{4/3} \log(1 + \sqrt[3]{cx} + c^{2/3}x^2)$$

input

```
Integrate[(a + b*ArcTanh[c*x^3])/x^5, x]
```

output

```
-1/4*a/x^4 - (3*b*c)/(4*x) - (Sqrt[3]*b*c^(4/3)*ArcTan[(-1 + 2*c^(1/3)*x)/Sqrt[3]])/8 - (Sqrt[3]*b*c^(4/3)*ArcTan[(1 + 2*c^(1/3)*x)/Sqrt[3]])/8 - (b*ArcTanh[c*x^3])/(4*x^4) - (b*c^(4/3)*Log[1 - c^(1/3)*x])/8 + (b*c^(4/3)*Log[1 + c^(1/3)*x])/8 - (b*c^(4/3)*Log[1 - c^(1/3)*x + c^(2/3)*x^2])/16 + (b*c^(4/3)*Log[1 + c^(1/3)*x + c^(2/3)*x^2])/16
```

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.33, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.786$, Rules used = {6452, 847, 825, 27, 219, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + \operatorname{barctanh}(cx^3)}{x^5} dx \\ & \quad \downarrow \text{6452} \\ & \frac{3}{4}bc \int \frac{1}{x^2(1 - c^2x^6)} dx - \frac{a + \operatorname{barctanh}(cx^3)}{4x^4} \\ & \quad \downarrow \text{847} \\ & \frac{3}{4}bc \left(c^2 \int \frac{x^4}{1 - c^2x^6} dx - \frac{1}{x} \right) - \frac{a + \operatorname{barctanh}(cx^3)}{4x^4} \end{aligned}$$

$$\begin{aligned}
& \downarrow 825 \\
& \frac{3}{4}bc \left(c^2 \left(\frac{\int \frac{1}{1-c^{2/3}x^2} dx}{3c^{4/3}} + \frac{\int -\frac{\sqrt[3]{Cx+1}}{2(c^{2/3}x^2-\sqrt[3]{Cx+1})} dx}{3c^{4/3}} + \frac{\int -\frac{1-\sqrt[3]{Cx}}{2(c^{2/3}x^2+\sqrt[3]{Cx+1})} dx}{3c^{4/3}} \right) - \frac{1}{x} \right) - \\
& \quad \frac{a + \operatorname{barctanh}(cx^3)}{4x^4} \\
& \downarrow 27 \\
& \frac{3}{4}bc \left(c^2 \left(\frac{\int \frac{1}{1-c^{2/3}x^2} dx}{3c^{4/3}} - \frac{\int \frac{\sqrt[3]{Cx+1}}{c^{2/3}x^2-\sqrt[3]{Cx+1}} dx}{6c^{4/3}} - \frac{\int \frac{1-\sqrt[3]{Cx}}{c^{2/3}x^2+\sqrt[3]{Cx+1}} dx}{6c^{4/3}} \right) - \frac{1}{x} \right) - \\
& \quad \frac{a + \operatorname{barctanh}(cx^3)}{4x^4} \\
& \downarrow 219 \\
& \frac{3}{4}bc \left(c^2 \left(-\frac{\int \frac{\sqrt[3]{Cx+1}}{c^{2/3}x^2-\sqrt[3]{Cx+1}} dx}{6c^{4/3}} - \frac{\int \frac{1-\sqrt[3]{Cx}}{c^{2/3}x^2+\sqrt[3]{Cx+1}} dx}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{Cx})}{3c^{5/3}} \right) - \frac{1}{x} \right) - \\
& \quad \frac{a + \operatorname{barctanh}(cx^3)}{4x^4} \\
& \downarrow 1142 \\
& \frac{3}{4}bc \left(c^2 \left(-\frac{\frac{3}{2} \int \frac{1}{c^{2/3}x^2-\sqrt[3]{Cx+1}} dx + \frac{\int -\frac{\sqrt[3]{C}(1-2\sqrt[3]{Cx})}{c^{2/3}x^2-\sqrt[3]{Cx+1}} dx}{2\sqrt[3]{C}}}{6c^{4/3}} - \frac{\frac{3}{2} \int \frac{1}{c^{2/3}x^2+\sqrt[3]{Cx+1}} dx - \frac{\int \frac{\sqrt[3]{C}(2\sqrt[3]{Cx+1})}{c^{2/3}x^2+\sqrt[3]{Cx+1}} dx}{2\sqrt[3]{C}}}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{Cx})}{3c^{5/3}} \right) \right) - \\
& \quad \frac{a + \operatorname{barctanh}(cx^3)}{4x^4} \\
& \downarrow 25
\end{aligned}$$

$$\frac{3}{4}bc \left(c^2 \left(-\frac{\frac{3}{2} \int \frac{1}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx - \frac{\int \frac{\sqrt[3]{c}(1-2\sqrt[3]{Cx})}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx}{2\sqrt[3]{c}}}{6c^{4/3}} - \frac{\frac{3}{2} \int \frac{1}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx - \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{Cx+1})}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx}{2\sqrt[3]{c}}}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{Cx})}{3c^{5/3}} \right) \right)$$

$$\frac{a + \operatorname{barctanh}(cx^3)}{4x^4}$$

↓ 27

$$\frac{3}{4}bc \left(c^2 \left(-\frac{\frac{3}{2} \int \frac{1}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx - \frac{1}{2} \int \frac{1-2\sqrt[3]{Cx}}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx}{6c^{4/3}} - \frac{\frac{3}{2} \int \frac{1}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx - \frac{1}{2} \int \frac{2\sqrt[3]{Cx+1}}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{Cx})}{3c^{5/3}} \right) \right)$$

$$\frac{a + \operatorname{barctanh}(cx^3)}{4x^4}$$

↓ 1082

$$\frac{3}{4}bc \left(c^2 \left(-\frac{\frac{3 \int \frac{1}{-(1-2\sqrt[3]{Cx})^2 - 3}}{\sqrt[3]{c}} d(1-2\sqrt[3]{Cx})}{6c^{4/3}} - \frac{1}{2} \int \frac{1-2\sqrt[3]{Cx}}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx - \frac{1}{2} \int \frac{2\sqrt[3]{Cx+1}}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx - \frac{3 \int \frac{1}{-(2\sqrt[3]{Cx+1})^2 - 3}}{\sqrt[3]{c}} d(2\sqrt[3]{Cx+1})}{6c^{4/3}} \right) \right)$$

$$\frac{a + \operatorname{barctanh}(cx^3)}{4x^4}$$

↓ 217

$$\frac{3}{4}bc \left(c^2 \left(-\frac{-\frac{1}{2} \int \frac{1-2\sqrt[3]{Cx}}{c^{2/3}x^2 - \sqrt[3]{Cx+1}} dx - \frac{\sqrt{3} \operatorname{arctan}\left(\frac{1-2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{\sqrt[3]{c}}}{6c^{4/3}} - \frac{\sqrt{3} \operatorname{arctan}\left(\frac{2\sqrt[3]{Cx+1}}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{1}{2} \int \frac{2\sqrt[3]{Cx+1}}{c^{2/3}x^2 + \sqrt[3]{Cx+1}} dx}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{Cx})}{3c^{5/3}} \right) \right)$$

$$\frac{a + \operatorname{barctanh}(cx^3)}{4x^4}$$

↓ 1103

$$\frac{3}{4}bc \left(c^2 \left(-\frac{\log(c^{2/3}x^2 - \sqrt[3]{c}x+1)}{2\sqrt[3]{c}} - \frac{\sqrt{3} \arctan\left(\frac{1-2\sqrt[3]{c}x}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{c}x+1}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{\log(c^{2/3}x^2 + \sqrt[3]{c}x+1)}{2\sqrt[3]{c}} \right) + \frac{\operatorname{arctanh}(\sqrt[3]{c}x)}{3c^{5/3}} \right) + \frac{a + b \operatorname{arctanh}(cx^3)}{4x^4}$$

input `Int[(a + b*ArcTanh[c*x^3])/x^5, x]`

output `-1/4*(a + b*ArcTanh[c*x^3])/x^4 + (3*b*c*(-x^(-1) + c^2*(ArcTanh[c^(1/3)*x]/(3*c^(5/3)) - ((Sqrt[3]*ArcTan[(1 - 2*c^(1/3)*x]/Sqrt[3])/c^(1/3)) + Log[1 - c^(1/3)*x + c^(2/3)*x^2]/(2*c^(1/3)))/(6*c^(4/3)) - ((Sqrt[3]*ArcTan[(1 + 2*c^(1/3)*x]/Sqrt[3])/c^(1/3) - Log[1 + c^(1/3)*x + c^(2/3)*x^2]/(2*c^(1/3)))/(6*c^(4/3))))/4`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 825 `Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Module[{r = Numerator[Rt[-a/b, n]], s = Denominator[Rt[-a/b, n]], k, u}, Simp[u = Int[(r*Cos[2*k*m*(Pi/n)] - s*Cos[2*k*(m + 1)*(Pi/n)]*x)/(r^2 - 2*r*s*Cos[2*k*(Pi/n)]*x + s^2*x^2), x] + Int[(r*Cos[2*k*m*(Pi/n)] + s*Cos[2*k*(m + 1)*(Pi/n)]*x)/(r^2 + 2*r*s*Cos[2*k*(Pi/n)]*x + s^2*x^2), x]; 2*(r^(m + 2)/(a*n*s^m)) Int[1/(r^2 - s^2*x^2), x] + 2*(r^(m + 1)/(a*n*s^m)) Sum[u, {k, 1, (n - 2)/4}], x] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && NegQ[a/b]`

rule 847 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + n*(p + 1) + 1)/(a*c^n*(m + 1)) Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.23

method	result
default	$-\frac{a}{4x^4} - \frac{b \operatorname{arctanh}(cx^3)}{4x^4} - \frac{bc \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{8\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{bc \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{16\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{bc\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{1}{c}\right)^{\frac{1}{3}}+1\right)}\right)}{8\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{3bc}{4x} + \dots$
parts	$-\frac{a}{4x^4} - \frac{b \operatorname{arctanh}(cx^3)}{4x^4} - \frac{bc \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{8\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{bc \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{16\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{bc\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{1}{c}\right)^{\frac{1}{3}}+1\right)}\right)}{8\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{3bc}{4x} + \dots$
risch	$-\frac{b \ln(cx^3+1)}{8x^4} - \frac{a}{4x^4} + \frac{b \ln(-cx^3+1)}{8x^4} - \frac{bc \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{8\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{bc \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{16\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{bc\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{1}{c}\right)^{\frac{1}{3}}+1\right)}\right)}{8\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \dots$

```
input int((a+b*arctanh(c*x^3))/x^5,x,method=_RETURNVERBOSE)
```

```
output -1/4*a/x^4-1/4*b/x^4*arctanh(c*x^3)-1/8*b*c/(1/c)^(1/3)*ln(x-(1/c)^(1/3))+
1/16*b*c/(1/c)^(1/3)*ln(x^2+(1/c)^(1/3)*x+(1/c)^(2/3))-1/8*b*c*3^(1/2)/(1/
c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x+1))-3/4*b*c/x+1/8*b*c/(1/c)^(
1/3)*ln(x+(1/c)^(1/3))-1/16*b*c/(1/c)^(1/3)*ln(x^2-(1/c)^(1/3)*x+(1/c)^(2/
3))-1/8*b*c*3^(1/2)/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x-1))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.40

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^5} dx = \frac{2\sqrt{3}b(-c)^{\frac{1}{3}}cx^4 \operatorname{arctan}\left(\frac{2}{3}\sqrt{3}(-c)^{\frac{1}{3}}x - \frac{1}{3}\sqrt{3}\right) + 2\sqrt{3}bc^{\frac{4}{3}}x^4 \operatorname{arctan}\left(\frac{2}{3}\sqrt{3}c^{\frac{1}{3}}x - \frac{1}{3}\sqrt{3}\right) + b(-c)^{\frac{1}{3}}cx^4 \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right) + \dots}{x^4}$$

input `integrate((a+b*arctanh(c*x^3))/x^5,x, algorithm="fricas")`

output `-1/16*(2*sqrt(3)*b*(-c)^(1/3)*c*x^4*arctan(2/3*sqrt(3)*(-c)^(1/3)*x - 1/3*sqrt(3)) + 2*sqrt(3)*b*c^(4/3)*x^4*arctan(2/3*sqrt(3)*c^(1/3)*x - 1/3*sqrt(3)) + b*(-c)^(1/3)*c*x^4*log(c*x^2 + (-c)^(2/3)*x - (-c)^(1/3)) + b*c^(4/3)*x^4*log(c*x^2 - c^(2/3)*x + c^(1/3)) - 2*b*(-c)^(1/3)*c*x^4*log(c*x - (-c)^(2/3)) - 2*b*c^(4/3)*x^4*log(c*x + c^(2/3)) + 12*b*c*x^3 + 2*b*log(-(c*x^3 + 1)/(c*x^3 - 1)) + 4*a)/x^4`

Sympy [F(-1)]

Timed out.

$$\int \frac{a + \operatorname{arctanh}(cx^3)}{x^5} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**3))/x**5,x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.14

$$\int \frac{a + \operatorname{arctanh}(cx^3)}{x^5} dx =$$

$$-\frac{1}{16} \left(\left(2\sqrt{3}c^{\frac{1}{3}} \arctan \left(\frac{\sqrt{3}(2c^{\frac{2}{3}}x + c^{\frac{1}{3}})}{3c^{\frac{1}{3}}} \right) + 2\sqrt{3}c^{\frac{1}{3}} \arctan \left(\frac{\sqrt{3}(2c^{\frac{2}{3}}x - c^{\frac{1}{3}})}{3c^{\frac{1}{3}}} \right) - c^{\frac{1}{3}} \log \left(c^{\frac{2}{3}}x^2 + c \right) \right) - \frac{a}{4x^4} \right)$$

input `integrate((a+b*arctanh(c*x^3))/x^5,x, algorithm="maxima")`

output

```
-1/16*((2*sqrt(3)*c^(1/3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*x + c^(1/3))/c^(1/3)) + 2*sqrt(3)*c^(1/3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*x - c^(1/3))/c^(1/3)) - c^(1/3)*log(c^(2/3)*x^2 + c^(1/3)*x + 1) + c^(1/3)*log(c^(2/3)*x^2 - c^(1/3)*x + 1) - 2*c^(1/3)*log((c^(1/3)*x + 1)/c^(1/3)) + 2*c^(1/3)*log((c^(1/3)*x - 1)/c^(1/3)) + 12/x)*c + 4*arctanh(c*x^3)/x^4)*b - 1/4*a/x^4
```

Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.38

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^5} dx = -\frac{\sqrt{3}bc^3 \operatorname{arctan}\left(\frac{1}{3}\sqrt{3}\left(2x + \frac{1}{|c|^{1/3}}\right)|c|^{1/3}\right)}{8|c|^{5/3}} - \frac{\sqrt{3}bc^3 \operatorname{arctan}\left(\frac{1}{3}\sqrt{3}\left(2x - \frac{1}{|c|^{1/3}}\right)|c|^{1/3}\right)}{8|c|^{5/3}} + \frac{bc^3 \log\left(x^2 + \frac{x}{|c|^{1/3}} + \frac{1}{|c|^{2/3}}\right)}{16|c|^{5/3}} - \frac{bc^3 \log\left(x^2 - \frac{x}{|c|^{1/3}} + \frac{1}{|c|^{2/3}}\right)}{16|c|^{5/3}} + \frac{bc^3 \log\left(\left|x + \frac{1}{|c|^{1/3}}\right|\right)}{8|c|^{5/3}} - \frac{bc^3 \log\left(\left|x - \frac{1}{|c|^{1/3}}\right|\right)}{8|c|^{5/3}} - \frac{b \log\left(\frac{-cx^3+1}{cx^3-1}\right)}{8x^4} - \frac{3bcx^3 + a}{4x^4}$$

input

```
integrate((a+b*arctanh(c*x^3))/x^5,x, algorithm="giac")
```

output

```
-1/8*sqrt(3)*b*c^3*arctan(1/3*sqrt(3)*(2*x + 1/abs(c)^(1/3))*abs(c)^(1/3))/abs(c)^(5/3) - 1/8*sqrt(3)*b*c^3*arctan(1/3*sqrt(3)*(2*x - 1/abs(c)^(1/3))*abs(c)^(1/3))/abs(c)^(5/3) + 1/16*b*c^3*log(x^2 + x/abs(c)^(1/3) + 1/abs(c)^(2/3))/abs(c)^(5/3) - 1/16*b*c^3*log(x^2 - x/abs(c)^(1/3) + 1/abs(c)^(2/3))/abs(c)^(5/3) + 1/8*b*c^3*log(abs(x + 1/abs(c)^(1/3)))/abs(c)^(5/3) - 1/8*b*c^3*log(abs(x - 1/abs(c)^(1/3)))/abs(c)^(5/3) - 1/8*b*log(-(c*x^3 + 1)/(c*x^3 - 1))/x^4 - 1/4*(3*b*c*x^3 + a)/x^4
```


Mupad [B] (verification not implemented)

Time = 4.15 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.89

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^5} dx$$

$$= \frac{b \ln(1 - cx^3)}{8x^4}$$

$$- \frac{bc^{4/3} \left(-\frac{\operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}-i)}{2}\right)}{2} + \frac{\operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}+i)}{2}\right)}{2} + \operatorname{atan}(c^{1/3}x \operatorname{li}) \right) \operatorname{li}}{4} - \frac{3bc}{4x}$$

$$- \frac{b \ln(cx^3 + 1)}{8x^4} - \frac{a}{4x^4} - \frac{\sqrt{3}bc^{4/3} \left(\operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}-i)}{2}\right) + \operatorname{atan}\left(\frac{c^{1/3}x(\sqrt{3}+i)}{2}\right) \right)}{8}$$

input `int((a + b*atanh(c*x^3))/x^5,x)`output `(b*log(1 - c*x^3))/(8*x^4) - (b*c^(4/3)*(atan((c^(1/3)*x*(3^(1/2) + 1i))/2)/2 - atan((c^(1/3)*x*(3^(1/2) - 1i))/2)/2 + atan(c^(1/3)*x*1i))*1i)/4 - (3*b*c)/(4*x) - (b*log(c*x^3 + 1))/(8*x^4) - a/(4*x^4) - (3^(1/2)*b*c^(4/3)*(atan((c^(1/3)*x*(3^(1/2) - 1i))/2) + atan((c^(1/3)*x*(3^(1/2) + 1i))/2)))/8`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.01

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^5} dx$$

$$= \frac{-2\sqrt{3} \operatorname{atan}\left(\frac{2c^{1/3}x-1}{\sqrt{3}}\right) bc^2x^4 - 2\sqrt{3} \operatorname{atan}\left(\frac{2c^{1/3}x+1}{\sqrt{3}}\right) bc^2x^4 - 4c^{2/3} \operatorname{atanh}(cx^3) b - 2 \operatorname{atanh}(cx^3) bc^2x^4 - 4c^{2/3} a}{16c^{2/3}x^4}$$

input `int((a+b*atanh(c*x^3))/x^5,x)`

output

```
( - 2*sqrt(3)*atan((2*c**(1/3)*x - 1)/sqrt(3))*b*c**2*x**4 - 2*sqrt(3)*atan((2*c**(1/3)*x + 1)/sqrt(3))*b*c**2*x**4 - 4*c**(2/3)*atanh(c*x**3)*b - 2*atanh(c*x**3)*b*c**2*x**4 - 4*c**(2/3)*a - 12*c**(2/3)*b*c*x**3 + 3*log(c**(2/3)*x + c**(1/3))*b*c**2*x**4 - 3*log(c**(2/3)*x - c**(1/3))*b*c**2*x**4)/(16*c**(2/3)*x**4)
```

3.116 $\int x^{11}(a + b \operatorname{arctanh}(cx^3))^2 dx$

Optimal result	990
Mathematica [A] (verified)	990
Rubi [A] (verified)	991
Maple [A] (verified)	994
Fricas [A] (verification not implemented)	994
Sympy [F(-1)]	995
Maxima [A] (verification not implemented)	995
Giac [A] (verification not implemented)	996
Mupad [B] (verification not implemented)	997
Reduce [B] (verification not implemented)	998

Optimal result

Integrand size = 16, antiderivative size = 125

$$\int x^{11}(a + b \operatorname{arctanh}(cx^3))^2 dx = \frac{abx^3}{6c^3} + \frac{b^2x^6}{36c^2} + \frac{b^2x^3 \operatorname{arctanh}(cx^3)}{6c^3} + \frac{bx^9(a + b \operatorname{arctanh}(cx^3))}{18c} - \frac{(a + b \operatorname{arctanh}(cx^3))^2}{12c^4} + \frac{1}{12}x^{12}(a + b \operatorname{arctanh}(cx^3))^2 + \frac{b^2 \log(1 - c^2x^6)}{9c^4}$$

output

```
1/6*a*b*x^3/c^3+1/36*b^2*x^6/c^2+1/6*b^2*x^3*arctanh(c*x^3)/c^3+1/18*b*x^9
*(a+b*arctanh(c*x^3))/c-1/12*(a+b*arctanh(c*x^3))^2/c^4+1/12*x^12*(a+b*arc
tanh(c*x^3))^2+1/9*b^2*ln(-c^2*x^6+1)/c^4
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.17

$$\int x^{11}(a + b \operatorname{arctanh}(cx^3))^2 dx = \frac{6abcx^3 + b^2c^2x^6 + 2abc^3x^9 + 3a^2c^4x^{12} + 2bcx^3(3ac^3x^9 + b(3 + c^2x^6)) \operatorname{arctanh}(cx^3) + 3b^2(-1 + c^4x^{12}) \operatorname{arctanh}^2(cx^3)}{36c^4}$$

input `Integrate[x^11*(a + b*ArcTanh[c*x^3])^2,x]`

output $(6*a*b*c*x^3 + b^2*c^2*x^6 + 2*a*b*c^3*x^9 + 3*a^2*c^4*x^{12} + 2*b*c*x^3*(3*a*c^3*x^9 + b*(3 + c^2*x^6))*ArcTanh[c*x^3] + 3*b^2*(-1 + c^4*x^{12})*ArcTanh[c*x^3]^2 + b*(3*a + 4*b)*Log[1 - c*x^3] - 3*a*b*Log[1 + c*x^3] + 4*b^2*Log[1 + c*x^3])/(36*c^4)$

Rubi [A] (verified)

Time = 0.91 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.22, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {6454, 6452, 6542, 6452, 243, 49, 2009, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{11}(a + b \operatorname{arctanh}(cx^3))^2 dx$$

$$\downarrow 6454$$

$$\frac{1}{3} \int x^9 (a + b \operatorname{arctanh}(cx^3))^2 dx^3$$

$$\downarrow 6452$$

$$\frac{1}{3} \left(\frac{1}{4} x^{12} (a + b \operatorname{arctanh}(cx^3))^2 - \frac{1}{2} bc \int \frac{x^{12} (a + b \operatorname{arctanh}(cx^3))}{1 - c^2 x^6} dx^3 \right)$$

$$\downarrow 6542$$

$$\frac{1}{3} \left(\frac{1}{4} x^{12} (a + b \operatorname{arctanh}(cx^3))^2 - \frac{1}{2} bc \left(\frac{\int \frac{x^6 (a + b \operatorname{arctanh}(cx^3))}{1 - c^2 x^6} dx^3}{c^2} - \frac{\int x^6 (a + b \operatorname{arctanh}(cx^3)) dx^3}{c^2} \right) \right)$$

$$\downarrow 6452$$

$$\frac{1}{3} \left(\frac{1}{4} x^{12} (a + b \operatorname{arctanh}(cx^3))^2 - \frac{1}{2} bc \left(\frac{\int \frac{x^6 (a + b \operatorname{arctanh}(cx^3))}{1 - c^2 x^6} dx^3}{c^2} - \frac{\frac{1}{3} x^9 (a + b \operatorname{arctanh}(cx^3)) - \frac{1}{3} bc \int \frac{x^9}{1 - c^2 x^6} dx^3}{c^2} \right) \right)$$

↓ 243

$$\frac{1}{3} \left(\frac{1}{4} x^{12} (a + \operatorname{barctanh}(cx^3))^2 - \frac{1}{2} bc \left(\frac{\int \frac{x^6 (a + \operatorname{barctanh}(cx^3))}{1 - c^2 x^6} dx^3}{c^2} - \frac{\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3)) - \frac{1}{6} bc \int \frac{x^6}{1 - c^2 x^6} dx^6}{c^2} \right) \right)$$

↓ 49

$$\frac{1}{3} \left(\frac{1}{4} x^{12} (a + \operatorname{barctanh}(cx^3))^2 - \frac{1}{2} bc \left(\frac{\int \frac{x^6 (a + \operatorname{barctanh}(cx^3))}{1 - c^2 x^6} dx^3}{c^2} - \frac{\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3)) - \frac{1}{6} bc \int \left(-\frac{1}{c^2} - \frac{1}{c^2} \right)}{c^2} \right) \right)$$

↓ 2009

$$\frac{1}{3} \left(\frac{1}{4} x^{12} (a + \operatorname{barctanh}(cx^3))^2 - \frac{1}{2} bc \left(\frac{\int \frac{x^6 (a + \operatorname{barctanh}(cx^3))}{1 - c^2 x^6} dx^3}{c^2} - \frac{\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3)) - \frac{1}{6} bc \left(-\frac{x^6}{c^2} - \frac{\log(1 - c^2 x^6)}{c^2} \right)}{c^2} \right) \right)$$

↓ 6542

$$\frac{1}{3} \left(\frac{1}{4} x^{12} (a + \operatorname{barctanh}(cx^3))^2 - \frac{1}{2} bc \left(\frac{\int \frac{a + \operatorname{barctanh}(cx^3)}{1 - c^2 x^6} dx^3}{c^2} - \frac{\int (a + \operatorname{barctanh}(cx^3)) dx^3}{c^2} - \frac{\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))}{c^2} \right) \right)$$

↓ 2009

$$\frac{1}{3} \left(\frac{1}{4} x^{12} (a + \operatorname{barctanh}(cx^3))^2 - \frac{1}{2} bc \left(\frac{\int \frac{a + \operatorname{barctanh}(cx^3)}{1 - c^2 x^6} dx^3}{c^2} - \frac{ax^3 + bx^3 \operatorname{arctanh}(cx^3) + \frac{b \log(1 - c^2 x^6)}{2c}}{c^2} - \frac{\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))}{c^2} \right) \right)$$

↓ 6510

$$\frac{1}{3} \left(\frac{1}{4} x^{12} (a + \operatorname{barctanh}(cx^3))^2 - \frac{1}{2} bc \left(\frac{(a + \operatorname{barctanh}(cx^3))^2}{2bc^3} - \frac{ax^3 + bx^3 \operatorname{arctanh}(cx^3) + \frac{b \log(1 - c^2 x^6)}{2c}}{c^2} - \frac{\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))}{c^2} \right) \right)$$

input `Int[x^11*(a + b*ArcTanh[c*x^3])^2,x]`

output
$$\frac{((x^{12}(a + b \operatorname{ArcTanh}[c x^3])^2)/4 - (b c (-((x^9(a + b \operatorname{ArcTanh}[c x^3]))/3 - (b c (-x^6/c^2) - \operatorname{Log}[1 - c^2 x^6]/c^4))/6)/c^2) + ((a + b \operatorname{ArcTanh}[c x^3])^2/(2 b c^3) - (a x^3 + b x^3 \operatorname{ArcTanh}[c x^3] + (b \operatorname{Log}[1 - c^2 x^6])/(2 c))/c^2)/c^2)/2)/3}$$

Defintions of rubi rules used

rule 49
$$\operatorname{Int}[(a + b x)^m (c + d x)^n, x] \rightarrow \operatorname{Int}[\operatorname{ExpandIntegrand}[(a + b x)^m (c + d x)^n, x], x] /; \operatorname{FreeQ}\{a, b, c, d, x\} \&\& \operatorname{IGtQ}[m, 0] \&\& \operatorname{IGtQ}[m + n + 2, 0]$$

rule 243
$$\operatorname{Int}[x^m (a + b x)^p, x] \rightarrow \operatorname{Simp}[1/2 \operatorname{Subst}[\operatorname{Int}[x^{(m-1)/2} (a + b x)^p, x], x, x^2], x] /; \operatorname{FreeQ}\{a, b, m, p, x\} \&\& \operatorname{IntegerQ}[(m-1)/2]$$

rule 2009
$$\operatorname{Int}[u, x] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$$

rule 6452
$$\operatorname{Int}[(a + \operatorname{ArcTanh}[c x^n])^p (b x)^m, x] \rightarrow \operatorname{Simp}[x^{m+1} ((a + b \operatorname{ArcTanh}[c x^n])^p / (m+1)), x] - \operatorname{Simp}[b c^n (p/(m+1)) \operatorname{Int}[x^{m+n} ((a + b \operatorname{ArcTanh}[c x^n])^{p-1} / (1 - c^2 x^{2n}))], x], x] /; \operatorname{FreeQ}\{a, b, c, m, n, x\} \&\& \operatorname{IGtQ}[p, 0] \&\& (\operatorname{EqQ}[p, 1] \|\| (\operatorname{EqQ}[n, 1] \&\& \operatorname{IntegerQ}[m])) \&\& \operatorname{NeQ}[m, -1]$$

rule 6454
$$\operatorname{Int}[(a + \operatorname{ArcTanh}[c x^n])^p (b x)^m, x] \rightarrow \operatorname{Simp}[1/n \operatorname{Subst}[\operatorname{Int}[x^{(\operatorname{Simplify}[(m+1)/n] - 1) (a + b \operatorname{ArcTanh}[c x])^p}, x], x, x^n], x] /; \operatorname{FreeQ}\{a, b, c, m, n, x\} \&\& \operatorname{IGtQ}[p, 1] \&\& \operatorname{IntegerQ}[\operatorname{Simplify}[(m+1)/n]]$$

rule 6510
$$\operatorname{Int}[(a + \operatorname{ArcTanh}[c x])^p / ((d + e x)^2), x] \rightarrow \operatorname{Simp}[(a + b \operatorname{ArcTanh}[c x])^{p+1} / (b c d (p+1)), x] /; \operatorname{FreeQ}\{a, b, c, d, e, p, x\} \&\& \operatorname{EqQ}[c^2 d + e, 0] \&\& \operatorname{NeQ}[p, -1]$$

rule 6542

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*((f_.)*(x_))^(m_)/((d_.) + (
e_.)*(x_)^2), x_Symbol] :> Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*
x])^p, x], x] - Simp[d*(f^2/e Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/
(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m,
1]
```

Maple [A] (verified)

Time = 1.53 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.30

method	result
parallelrisch	$\frac{3b^2 \operatorname{arctanh}(cx^3)^2 x^{12} c^4 + 6ab \operatorname{arctanh}(cx^3) x^{12} c^4 + 3a^2 c^4 x^{12} + 2b^2 \operatorname{arctanh}(cx^3) x^9 c^3 + 2ab c^3 x^9 + b^2 c^2 x^6 + 6b^2 \operatorname{arctanh}(cx^3)}{36c^4}$
risch	$\frac{b^2(x^{12}c^4 - 1) \ln(cx^3 + 1)^2}{48c^4} + \frac{b(-3x^{12}b \ln(-cx^3 + 1)c^4 + 6ac^4x^{12} + 2bc^3x^9 + 6bcx^3 + 3b \ln(-cx^3 + 1)) \ln(cx^3 + 1)}{72c^4} + \frac{b^2x^{12}}{36c^4}$
default	Expression too large to display
parts	Expression too large to display

input

```
int(x^11*(a+b*arctanh(c*x^3))^2,x,method=_RETURNVERBOSE)
```

output

```
1/36*(3*b^2*arctanh(c*x^3)^2*x^12*c^4+6*a*b*arctanh(c*x^3)*x^12*c^4+3*a^2*
c^4*x^12+2*b^2*arctanh(c*x^3)*x^9*c^3+2*a*b*c^3*x^9+b^2*c^2*x^6+6*b^2*arct
anh(c*x^3)*x^3*c+6*a*b*c*x^3-3*b^2*arctanh(c*x^3)^2+8*ln(c*x^3-1)*b^2-6*ar
ctanh(c*x^3)*a*b+8*arctanh(c*x^3)*b^2+b^2)/c^4
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.41

$$\int x^{11} (a + b \operatorname{arctanh}(cx^3))^2 dx$$

$$= \frac{12 a^2 c^4 x^{12} + 8 a b c^3 x^9 + 4 b^2 c^2 x^6 + 24 a b c x^3 + 3 (b^2 c^4 x^{12} - b^2) \log\left(-\frac{cx^3+1}{cx^3-1}\right)^2 - 4 (3 a b - 4 b^2) \log(cx^3 + 1)}{144 c^4}$$

input

```
integrate(x^11*(a+b*arctanh(c*x^3))^2,x, algorithm="fricas")
```

output

```
1/144*(12*a^2*c^4*x^12 + 8*a*b*c^3*x^9 + 4*b^2*c^2*x^6 + 24*a*b*c*x^3 + 3*
(b^2*c^4*x^12 - b^2)*log(-(c*x^3 + 1)/(c*x^3 - 1))^2 - 4*(3*a*b - 4*b^2)*l
og(c*x^3 + 1) + 4*(3*a*b + 4*b^2)*log(c*x^3 - 1) + 4*(3*a*b*c^4*x^12 + b^2
*c^3*x^9 + 3*b^2*c*x^3)*log(-(c*x^3 + 1)/(c*x^3 - 1)))/c^4
```

Sympy [F(-1)]

Timed out.

$$\int x^{11} (a + b \operatorname{arctanh}(cx^3))^2 dx = \text{Timed out}$$

input

```
integrate(x**11*(a+b*atanh(c*x**3))**2,x)
```

output

Timed out

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.74

$$\int x^{11} (a + b \operatorname{arctanh}(cx^3))^2 dx = \frac{1}{12} b^2 x^{12} \operatorname{arctanh}(cx^3)^2 + \frac{1}{12} a^2 x^{12} + \frac{1}{36} \left(6 x^{12} \operatorname{arctanh}(cx^3) + c \left(\frac{2(c^2 x^9 + 3x^3)}{c^4} - \frac{3 \log(cx^3 + 1)}{c^5} + \frac{3 \log(cx^3 - 1)}{c^5} \right) \right) ab + \frac{1}{144} \left(4c \left(\frac{2(c^2 x^9 + 3x^3)}{c^4} - \frac{3 \log(cx^3 + 1)}{c^5} + \frac{3 \log(cx^3 - 1)}{c^5} \right) \operatorname{arctanh}(cx^3) + \frac{4c^2 x^6 - 2(3 \log(cx^3 - 1))}{c^4} \right) b^2$$

input

```
integrate(x^11*(a+b*arctanh(c*x^3))^2,x, algorithm="maxima")
```

output

```
1/12*b^2*x^12*arctanh(c*x^3)^2 + 1/12*a^2*x^12 + 1/36*(6*x^12*arctanh(c*x^
3) + c*(2*(c^2*x^9 + 3*x^3)/c^4 - 3*log(c*x^3 + 1)/c^5 + 3*log(c*x^3 - 1)/
c^5))*a*b + 1/144*(4*c*(2*(c^2*x^9 + 3*x^3)/c^4 - 3*log(c*x^3 + 1)/c^5 + 3
*log(c*x^3 - 1)/c^5)*arctanh(c*x^3) + (4*c^2*x^6 - 2*(3*log(c*x^3 - 1) - 8
)*log(c*x^3 + 1) + 3*log(c*x^3 + 1)^2 + 3*log(c*x^3 - 1)^2 + 16*log(c*x^3
- 1))/c^4)*b^2
```


Giac [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.40

$$\begin{aligned}
\int x^{11}(a + b \operatorname{arctanh}(cx^3))^2 dx &= \frac{1}{12} a^2 x^{12} + \frac{abx^9}{18c} + \frac{b^2 x^6}{36c^2} \\
&+ \frac{1}{48} \left(b^2 x^{12} - \frac{b^2}{c^4} \right) \log \left(-\frac{cx^3 + 1}{cx^3 - 1} \right)^2 + \frac{abx^3}{6c^3} \\
&+ \frac{1}{36} \left(3abx^{12} + \frac{b^2 x^9}{c} + \frac{3b^2 x^3}{c^3} \right) \log \left(-\frac{cx^3 + 1}{cx^3 - 1} \right) \\
&- \frac{(3ab - 4b^2) \log(cx^3 + 1)}{36c^4} \\
&+ \frac{(3ab + 4b^2) \log(cx^3 - 1)}{36c^4}
\end{aligned}$$

input `integrate(x^11*(a+b*arctanh(c*x^3))^2,x, algorithm="giac")`

output `1/12*a^2*x^12 + 1/18*a*b*x^9/c + 1/36*b^2*x^6/c^2 + 1/48*(b^2*x^12 - b^2/c^4)*log(-(c*x^3 + 1)/(c*x^3 - 1))^2 + 1/6*a*b*x^3/c^3 + 1/36*(3*a*b*x^12 + b^2*x^9/c + 3*b^2*x^3/c^3)*log(-(c*x^3 + 1)/(c*x^3 - 1)) - 1/36*(3*a*b - 4*b^2)*log(c*x^3 + 1)/c^4 + 1/36*(3*a*b + 4*b^2)*log(c*x^3 - 1)/c^4`

Mupad [B] (verification not implemented)

Time = 4.34 (sec) , antiderivative size = 335, normalized size of antiderivative = 2.68

$$\begin{aligned}
\int x^{11}(a + b \operatorname{arctanh}(cx^3))^2 dx = & \frac{a^2 x^{12}}{12} + \frac{b^2 \ln(cx^3 - 1)}{9c^4} + \frac{b^2 \ln(cx^3 + 1)}{9c^4} \\
& - \frac{b^2 \ln(cx^3 + 1)^2}{48c^4} - \frac{b^2 \ln(1 - cx^3)^2}{48c^4} + \frac{b^2 x^6}{36c^2} \\
& + \frac{b^2 x^{12} \ln(cx^3 + 1)^2}{48} + \frac{b^2 x^{12} \ln(1 - cx^3)^2}{48} \\
& + \frac{b^2 x^3 \ln(cx^3 + 1)}{12c^3} - \frac{b^2 x^3 \ln(1 - cx^3)}{12c^3} \\
& + \frac{b^2 x^9 \ln(cx^3 + 1)}{36c} - \frac{b^2 x^9 \ln(1 - cx^3)}{36c} \\
& + \frac{ab \ln(cx^3 - 1)}{12c^4} - \frac{ab \ln(cx^3 + 1)}{12c^4} \\
& + \frac{abx^{12} \ln(cx^3 + 1)}{12} - \frac{abx^{12} \ln(1 - cx^3)}{12} \\
& + \frac{b^2 \ln(cx^3 + 1) \ln(1 - cx^3)}{24c^4} + \frac{abx^3}{6c^3} \\
& + \frac{abx^9}{18c} - \frac{b^2 x^{12} \ln(cx^3 + 1) \ln(1 - cx^3)}{24}
\end{aligned}$$

input `int(x^11*(a + b*atanh(c*x^3))^2,x)`output `(a^2*x^12)/12 + (b^2*log(c*x^3 - 1))/(9*c^4) + (b^2*log(c*x^3 + 1))/(9*c^4) - (b^2*log(c*x^3 + 1)^2)/(48*c^4) - (b^2*log(1 - c*x^3)^2)/(48*c^4) + (b^2*x^6)/(36*c^2) + (b^2*x^12*log(c*x^3 + 1)^2)/48 + (b^2*x^12*log(1 - c*x^3)^2)/48 + (b^2*x^3*log(c*x^3 + 1))/(12*c^3) - (b^2*x^3*log(1 - c*x^3))/(12*c^3) + (b^2*x^9*log(c*x^3 + 1))/(36*c) - (b^2*x^9*log(1 - c*x^3))/(36*c) + (a*b*log(c*x^3 - 1))/(12*c^4) - (a*b*log(c*x^3 + 1))/(12*c^4) + (a*b*x^12*log(c*x^3 + 1))/12 - (a*b*x^12*log(1 - c*x^3))/12 + (b^2*log(c*x^3 + 1)*log(1 - c*x^3))/(24*c^4) + (a*b*x^3)/(6*c^3) + (a*b*x^9)/(18*c) - (b^2*x^12*log(c*x^3 + 1)*log(1 - c*x^3))/24`

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.46

$$\int x^{11} (a + b \operatorname{arctanh}(cx^3))^2 dx$$

$$= \frac{3 \operatorname{atanh}(cx^3)^2 b^2 c^4 x^{12} - 3 \operatorname{atanh}(cx^3)^2 b^2 + 6 \operatorname{atanh}(cx^3) ab c^4 x^{12} - 6 \operatorname{atanh}(cx^3) ab + 2 \operatorname{atanh}(cx^3) b^2 c^3 x^9}{36 c^4}$$

input

```
int(x^11*(a+b*atanh(c*x^3))^2,x)
```

output

```
(3*atanh(c*x**3)**2*b**2*c**4*x**12 - 3*atanh(c*x**3)**2*b**2 + 6*atanh(c*x**3)*a*b*c**4*x**12 - 6*atanh(c*x**3)*a*b + 2*atanh(c*x**3)*b**2*c**3*x**9 + 6*atanh(c*x**3)*b**2*c*x**3 - 8*atanh(c*x**3)*b**2 + 8*log(c**(2/3)*x**2 - c**(1/3)*x + 1)*b**2 + 8*log(c**(2/3)*x + c**(1/3))*b**2 + 3*a**2*c**4*x**12 + 2*a*b*c**3*x**9 + 6*a*b*c*x**3 + b**2*c**2*x**6)/(36*c**4)
```

3.117 $\int x^8(a + b \operatorname{arctanh}(cx^3))^2 dx$

Optimal result	999
Mathematica [A] (verified)	1000
Rubi [A] (verified)	1000
Maple [C] (warning: unable to verify)	1004
Fricas [F]	1005
Sympy [F(-1)]	1005
Maxima [F]	1005
Giac [F]	1006
Mupad [F(-1)]	1006
Reduce [F]	1007

Optimal result

Integrand size = 16, antiderivative size = 146

$$\int x^8(a + b \operatorname{arctanh}(cx^3))^2 dx = \frac{b^2 x^3}{9c^2} - \frac{b^2 \operatorname{arctanh}(cx^3)}{9c^3} + \frac{bx^6(a + b \operatorname{arctanh}(cx^3))}{9c} + \frac{(a + b \operatorname{arctanh}(cx^3))^2}{9c^3} + \frac{1}{9} x^9 (a + b \operatorname{arctanh}(cx^3))^2 - \frac{2b(a + b \operatorname{arctanh}(cx^3)) \log\left(\frac{2}{1-cx^3}\right)}{9c^3} - \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^3}\right)}{9c^3}$$

output

```
1/9*b^2*x^3/c^2-1/9*b^2*arctanh(c*x^3)/c^3+1/9*b*x^6*(a+b*arctanh(c*x^3))/
c+1/9*(a+b*arctanh(c*x^3))^2/c^3+1/9*x^9*(a+b*arctanh(c*x^3))^2-2/9*b*(a+b
*arctanh(c*x^3))*ln(2/(-c*x^3+1))/c^3-1/9*b^2*polylog(2,1-2/(-c*x^3+1))/c^
3
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.90

$$\int x^8 (a + b \operatorname{arctanh}(cx^3))^2 dx$$

$$= \frac{b^2 cx^3 + abc^2 x^6 + a^2 c^3 x^9 + b^2 (-1 + c^3 x^9) \operatorname{arctanh}(cx^3)^2 + b \operatorname{arctanh}(cx^3) \left(-b + bc^2 x^6 + 2ac^3 x^9 - 2b \log \right)}{9c^3}$$

input

```
Integrate[x^8*(a + b*ArcTanh[c*x^3])^2,x]
```

output

```
(b^2*c*x^3 + a*b*c^2*x^6 + a^2*c^3*x^9 + b^2*(-1 + c^3*x^9)*ArcTanh[c*x^3]^2 + b*ArcTanh[c*x^3]*(-b + b*c^2*x^6 + 2*a*c^3*x^9 - 2*b*Log[1 + E^(-2*ArcTanh[c*x^3])])) + a*b*Log[-1 + c^2*x^6] + b^2*PolyLog[2, -E^(-2*ArcTanh[c*x^3])])/(9*c^3)
```

Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.11, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {6454, 6452, 6542, 6452, 262, 219, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^8 (a + b \operatorname{arctanh}(cx^3))^2 dx$$

$$\downarrow 6454$$

$$\frac{1}{3} \int x^6 (a + b \operatorname{arctanh}(cx^3))^2 dx^3$$

$$\downarrow 6452$$

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + b \operatorname{arctanh}(cx^3))^2 - \frac{2}{3} bc \int \frac{x^9 (a + b \operatorname{arctanh}(cx^3))}{1 - c^2 x^6} dx^3 \right)$$

$$\downarrow 6542$$

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))^2 - \frac{2}{3} bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(cx^3))}{1 - c^2 x^6} dx^3}{c^2} - \frac{\int x^3 (a + \operatorname{barctanh}(cx^3)) dx^3}{c^2} \right) \right)$$

↓ 6452

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))^2 - \frac{2}{3} bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(cx^3))}{1 - c^2 x^6} dx^3}{c^2} - \frac{\frac{1}{2} x^6 (a + \operatorname{barctanh}(cx^3)) - \frac{1}{2} bc \int \frac{x^6}{1 - c^2 x^6} dx^3}{c^2} \right) \right)$$

↓ 262

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))^2 - \frac{2}{3} bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(cx^3))}{1 - c^2 x^6} dx^3}{c^2} - \frac{\frac{1}{2} x^6 (a + \operatorname{barctanh}(cx^3)) - \frac{1}{2} bc \left(\frac{\int \frac{1}{1 - c^2 x^6} dx^3}{c^2} \right)}{c^2} \right) \right)$$

↓ 219

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))^2 - \frac{2}{3} bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(cx^3))}{1 - c^2 x^6} dx^3}{c^2} - \frac{\frac{1}{2} x^6 (a + \operatorname{barctanh}(cx^3)) - \frac{1}{2} bc \left(\frac{\operatorname{arctanh}(cx^3)}{c^3} \right)}{c^2} \right) \right)$$

↓ 6546

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))^2 - \frac{2}{3} bc \left(\frac{\int \frac{a + \operatorname{barctanh}(cx^3)}{1 - cx^3} dx^3}{c} - \frac{(a + \operatorname{barctanh}(cx^3))^2}{2bc^2} - \frac{\frac{1}{2} x^6 (a + \operatorname{barctanh}(cx^3)) - \frac{1}{2} bc \left(\frac{\operatorname{arctanh}(cx^3)}{c^3} \right)}{c^2} \right) \right)$$

↓ 6470

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))^2 - \frac{2}{3} bc \left(\frac{\frac{\log\left(\frac{2}{1 - cx^3}\right) (a + \operatorname{barctanh}(cx^3))}{c} - b \int \frac{\log\left(\frac{2}{1 - cx^3}\right)}{1 - c^2 x^6} dx^3}{c} - \frac{(a + \operatorname{barctanh}(cx^3))^2}{2bc^2} - \frac{\frac{1}{2} x^6 (a + \operatorname{barctanh}(cx^3)) - \frac{1}{2} bc \left(\frac{\operatorname{arctanh}(cx^3)}{c^3} \right)}{c^2} \right) \right)$$

↓ 2849

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))^2 - \frac{2}{3} bc \left(\frac{\int \frac{\log\left(\frac{2}{1-cx^3}\right) d\frac{1}{1-cx^3}}{\frac{1-2}{1-cx^3} c} + \frac{\log\left(\frac{2}{1-cx^3}\right) (a + \operatorname{barctanh}(cx^3))}{c}}{c^2} - \frac{(a + \operatorname{barctanh}(cx^3))^2}{2bc^2} \right) \right)$$

↓ 2752

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))^2 - \frac{2}{3} bc \left(\frac{\frac{\log\left(\frac{2}{1-cx^3}\right) (a + \operatorname{barctanh}(cx^3))}{c} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^3}\right)}{2c}}{c^2} - \frac{(a + \operatorname{barctanh}(cx^3))^2}{2bc^2} \right) \right)$$

input `Int[x^8*(a + b*ArcTanh[c*x^3])^2,x]`

output `((x^9*(a + b*ArcTanh[c*x^3])^2)/3 - (2*b*c*(-((x^6*(a + b*ArcTanh[c*x^3]))/2 - (b*c*(-(x^3/c^2) + ArcTanh[c*x^3]/c^3))/2)/c^2) + (-1/2*(a + b*ArcTanh[c*x^3])^2/(b*c^2) + (((a + b*ArcTanh[c*x^3])*Log[2/(1 - c*x^3)])/c + (b*PolyLog[2, 1 - 2/(1 - c*x^3)]/(2*c))/c)/c^2))/3/3`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 262 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*(m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e.)*(x_))]/((f_) + (g.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 6452 `Int[((a_.) + ArcTanh[(c.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_.) + ArcTanh[(c.)*(x_)^(n)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6470 `Int[((a_.) + ArcTanh[(c.)*(x_)]*(b_.))^(p_.)/((d_) + (e.)*(x_)), x_Symbol] := Simp[(-a + b*ArcTanh[c*x])^p*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6542 `Int[((a_.) + ArcTanh[(c.)*(x_)]*(b_.))^(p_.)*((f.)*(x_)^(m_))/((d_) + (e.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

rule 6546 `Int[((a_.) + ArcTanh[(c.)*(x_)]*(b_.))^(p_.)*(x_)/((d_) + (e.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.36 (sec) , antiderivative size = 2906, normalized size of antiderivative = 19.90

method	result	size
default	Expression too large to display	2906
parts	Expression too large to display	2906

input `int(x^8*(a+b*arctanh(c*x^3))^2,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/9*a^2*x^9+b^2*(1/9*x^9*arctanh(c*x^3)^2-2/3*c*(-1/6*arctanh(c*x^3)/c^2*x \\ & ^6-1/6*arctanh(c*x^3)/c^4*\ln(c^2*x^6-1)-1/2*c*(1/3/c^4*x^3-1/6/c^5*\ln(c*x^ \\ & 3+1)+1/6/c^5*\ln(c*x^3-1)+1/c^4*(\text{Sum}(1/6*(\ln(x_alpha)*\ln(c^2*x^6-1)-6*c^2* \\ & (1/2/c*(1/3*\ln(x_alpha))*(\ln(1/2*(x_alpha)/_alpha)+\ln((\text{RootOf}(_Z^2+_Z*_al \\ & pha+_alpha^2,\text{index}=1)-x_alpha)/\text{RootOf}(_Z^2+_Z*_alpha+_alpha^2,\text{index}=1))+\ln \\ & ((\text{RootOf}(_Z^2+_Z*_alpha+_alpha^2,\text{index}=2)-x_alpha)/\text{RootOf}(_Z^2+_Z*_alpha \\ & +_alpha^2,\text{index}=2))))/c+1/3*(\text{dilog}(1/2*(x_alpha)/_alpha)+\text{dilog}((\text{RootOf}(_Z^ \\ & 2+_Z*_alpha+_alpha^2,\text{index}=1)-x_alpha)/\text{RootOf}(_Z^2+_Z*_alpha+_alpha^2,\text{ind} \\ & ex=1))+\text{dilog}((\text{RootOf}(_Z^2+_Z*_alpha+_alpha^2,\text{index}=2)-x_alpha)/\text{RootOf}(_Z^ \\ & 2+_Z*_alpha+_alpha^2,\text{index}=2))))/c)+1/2*c*(1/6/_alpha^2/c*\ln(x_alpha)^2-1/ \\ & 3*_alpha*\ln(x_alpha)*(2*\ln((\text{RootOf}(_Z^2+3*_Z*_alpha+3*_alpha^2,\text{index}=1)-x \\ & +_alpha)/\text{RootOf}(_Z^2+3*_Z*_alpha+3*_alpha^2,\text{index}=1))*\text{RootOf}(_Z^2+3*_Z*_al \\ & pha+3*_alpha^2,\text{index}=1)*\text{RootOf}(_Z^2+3*_Z*_alpha+3*_alpha^2,\text{index}=2)+3*\ln((\\ & \text{RootOf}(_Z^2+3*_Z*_alpha+3*_alpha^2,\text{index}=1)-x_alpha)/\text{RootOf}(_Z^2+3*_Z*_al \\ & pha+3*_alpha^2,\text{index}=1))*\text{RootOf}(_Z^2+3*_Z*_alpha+3*_alpha^2,\text{index}=1)*_alph \\ & a+6*\ln((\text{RootOf}(_Z^2+3*_Z*_alpha+3*_alpha^2,\text{index}=1)-x_alpha)/\text{RootOf}(_Z^2+ \\ & 3*_Z*_alpha+3*_alpha^2,\text{index}=1))*\text{RootOf}(_Z^2+3*_Z*_alpha+3*_alpha^2,\text{index}= \\ & 2)*_alpha+9*\ln((\text{RootOf}(_Z^2+3*_Z*_alpha+3*_alpha^2,\text{index}=1)-x_alpha)/\text{Root} \\ & \text{Of}(_Z^2+3*_Z*_alpha+3*_alpha^2,\text{index}=1))*_alpha^2+2*\ln((\text{RootOf}(_Z^2+3*_Z* \\ & alpha+3*_alpha^2,\text{index}=2)-x_alpha)/\text{RootOf}(_Z^2+3*_Z*_alpha+3*_alpha^2,\dots \end{aligned}$$

Fricas [F]

$$\int x^8 (a + b \operatorname{arctanh}(cx^3))^2 dx = \int (b \operatorname{arctanh}(cx^3) + a)^2 x^8 dx$$

input `integrate(x^8*(a+b*arctanh(c*x^3))^2,x, algorithm="fricas")`

output `integral(b^2*x^8*arctanh(c*x^3)^2 + 2*a*b*x^8*arctanh(c*x^3) + a^2*x^8, x)`

Sympy [F(-1)]

Timed out.

$$\int x^8 (a + b \operatorname{arctanh}(cx^3))^2 dx = \text{Timed out}$$

input `integrate(x**8*(a+b*atanh(c*x**3))**2,x)`

output `Timed out`

Maxima [F]

$$\int x^8 (a + b \operatorname{arctanh}(cx^3))^2 dx = \int (b \operatorname{arctanh}(cx^3) + a)^2 x^8 dx$$

input `integrate(x^8*(a+b*arctanh(c*x^3))^2,x, algorithm="maxima")`

output

```

1/9*a^2*x^9 + 1/9*(2*x^9*arctanh(c*x^3) + (x^6/c^2 + log(c^2*x^6 - 1)/c^4)
*c)*a*b + 1/648*(18*x^9*log(-c*x^3 + 1)^2 - 2*c^4*(2*(c^2*x^9 + 3*x^3)/c^6
- 3*log(c*x^3 + 1)/c^7 + 3*log(c*x^3 - 1)/c^7) + 3*(x^6/c^4 + log(c^2*x^6
- 1)/c^6)*c^3 + 1944*c^3*integrate(1/9*x^11*log(c*x^3 + 1)/(c^4*x^6 - c^2
), x) - 9*c^2*(2*x^3/c^4 - log(c*x^3 + 1)/c^5 + log(c*x^3 - 1)/c^5) - 6*c*
((2*c^2*x^9 + 3*c*x^6 + 6*x^3)/c^3 + 6*log(c*x^3 - 1)/c^4)*log(-c*x^3 + 1)
+ 972*c*integrate(1/9*x^5*log(c*x^3 + 1)/(c^4*x^6 - c^2), x) + 6*(3*c^3*x
^9*log(c*x^3 + 1)^2 + (2*c^3*x^9 - 3*c^2*x^6 + 6*c*x^3 - 6*(c^3*x^9 + 1)*l
og(c*x^3 + 1))*log(-c*x^3 + 1))/c^3 + (4*c^3*x^9 + 15*c^2*x^6 + 66*c*x^3 +
18*log(c*x^3 - 1)^2 + 66*log(c*x^3 - 1))/c^3 - 18*log(9*c^4*x^6 - 9*c^2)/
c^3 + 972*integrate(1/9*x^2*log(c*x^3 + 1)/(c^4*x^6 - c^2), x))*b^2

```

Giac [F]

$$\int x^8 (a + b \operatorname{arctanh}(cx^3))^2 dx = \int (b \operatorname{artanh}(cx^3) + a)^2 x^8 dx$$

input

```
integrate(x^8*(a+b*arctanh(c*x^3))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^3) + a)^2*x^8, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^8 (a + b \operatorname{arctanh}(cx^3))^2 dx = \int x^8 (a + b \operatorname{atanh}(cx^3))^2 dx$$

input

```
int(x^8*(a + b*atanh(c*x^3))^2,x)
```

output

```
int(x^8*(a + b*atanh(c*x^3))^2, x)
```

Reduce [F]

$$\int x^8 (a + b \operatorname{arctanh}(cx^3))^2 dx$$

$$= \frac{\operatorname{atanh}(cx^3)^2 b^2 c^3 x^9 - \operatorname{atanh}(cx^3)^2 b^2 c x^3 + 2 \operatorname{atanh}(cx^3) a b c^3 x^9 - 2 \operatorname{atanh}(cx^3) a b + \operatorname{atanh}(cx^3) b^2 c^2 x^6}{9 c^3}$$

input `int(x^8*(a+b*atanh(c*x^3))^2,x)`

output `(atanh(c*x**3)**2*b**2*c**3*x**9 - atanh(c*x**3)**2*b**2*c*x**3 + 2*atanh(c*x**3)*a*b*c**3*x**9 - 2*atanh(c*x**3)*a*b + atanh(c*x**3)*b**2*c**2*x**6 - atanh(c*x**3)*b**2 + 3*int(atanh(c*x**3)**2*x**2,x)*b**2*c + 2*log(c**(2/3)*x**2 - c**(1/3)*x + 1)*a*b + 2*log(c**(2/3)*x + c**(1/3))*a*b + a**2*c**3*x**9 + a*b*c**2*x**6 + b**2*c*x**3)/(9*c**3)`

3.118 $\int x^5(a + b \operatorname{arctanh}(cx^3))^2 dx$

Optimal result	1008
Mathematica [A] (verified)	1008
Rubi [A] (verified)	1009
Maple [A] (verified)	1011
Fricas [A] (verification not implemented)	1011
Sympy [F(-1)]	1012
Maxima [B] (verification not implemented)	1012
Giac [B] (verification not implemented)	1013
Mupad [B] (verification not implemented)	1014
Reduce [B] (verification not implemented)	1015

Optimal result

Integrand size = 16, antiderivative size = 91

$$\int x^5(a + b \operatorname{arctanh}(cx^3))^2 dx = \frac{abx^3}{3c} + \frac{b^2x^3 \operatorname{arctanh}(cx^3)}{3c} - \frac{(a + b \operatorname{arctanh}(cx^3))^2}{6c^2} + \frac{1}{6}x^6(a + b \operatorname{arctanh}(cx^3))^2 + \frac{b^2 \log(1 - c^2x^6)}{6c^2}$$

output

$1/3*a*b*x^3/c+1/3*b^2*x^3*\operatorname{arctanh}(c*x^3)/c-1/6*(a+b*\operatorname{arctanh}(c*x^3))^2/c^2+1/6*x^6*(a+b*\operatorname{arctanh}(c*x^3))^2+1/6*b^2*\ln(-c^2*x^6+1)/c^2$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.16

$$\int x^5(a + b \operatorname{arctanh}(cx^3))^2 dx = \frac{2abcx^3 + a^2c^2x^6 + 2bcx^3(b + acx^3) \operatorname{arctanh}(cx^3) + b^2(-1 + c^2x^6) \operatorname{arctanh}(cx^3)^2 + b(a + b) \log(1 - cx^3)}{6c^2}$$

input

`Integrate[x^5*(a + b*ArcTanh[c*x^3])^2,x]`

output

$$(2*a*b*c*x^3 + a^2*c^2*x^6 + 2*b*c*x^3*(b + a*c*x^3)*\text{ArcTanh}[c*x^3] + b^2*(-1 + c^2*x^6)*\text{ArcTanh}[c*x^3]^2 + b*(a + b)*\text{Log}[1 - c*x^3] - a*b*\text{Log}[1 + c*x^3] + b^2*\text{Log}[1 + c*x^3])/(6*c^2)$$
Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.01, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6454, 6452, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^5 (a + b \operatorname{arctanh}(cx^3))^2 dx$$

$$\downarrow 6454$$

$$\frac{1}{3} \int x^3 (a + b \operatorname{arctanh}(cx^3))^2 dx^3$$

$$\downarrow 6452$$

$$\frac{1}{3} \left(\frac{1}{2} x^6 (a + b \operatorname{arctanh}(cx^3))^2 - bc \int \frac{x^6 (a + b \operatorname{arctanh}(cx^3))}{1 - c^2 x^6} dx^3 \right)$$

$$\downarrow 6542$$

$$\frac{1}{3} \left(\frac{1}{2} x^6 (a + b \operatorname{arctanh}(cx^3))^2 - bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(cx^3)}{1 - c^2 x^6} dx^3}{c^2} - \frac{\int (a + b \operatorname{arctanh}(cx^3)) dx^3}{c^2} \right) \right)$$

$$\downarrow 2009$$

$$\frac{1}{3} \left(\frac{1}{2} x^6 (a + b \operatorname{arctanh}(cx^3))^2 - bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(cx^3)}{1 - c^2 x^6} dx^3}{c^2} - \frac{ax^3 + bx^3 \operatorname{arctanh}(cx^3) + \frac{b \log(1 - c^2 x^6)}{2c}}{c^2} \right) \right)$$

$$\downarrow 6510$$

$$\frac{1}{3} \left(\frac{1}{2} x^6 (a + b \operatorname{arctanh}(cx^3))^2 - bc \left(\frac{(a + b \operatorname{arctanh}(cx^3))^2}{2bc^3} - \frac{ax^3 + bx^3 \operatorname{arctanh}(cx^3) + \frac{b \log(1 - c^2 x^6)}{2c}}{c^2} \right) \right)$$

input `Int[x^5*(a + b*ArcTanh[c*x^3])^2,x]`

output `((x^6*(a + b*ArcTanh[c*x^3])^2)/2 - b*c*((a + b*ArcTanh[c*x^3])^2/(2*b*c^3) - (a*x^3 + b*x^3*ArcTanh[c*x^3] + (b*Log[1 - c^2*x^6])/(2*c))/c^2))/3`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6510 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6542 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((f_.)*(x_)^(m_.))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

Maple [A] (verified)

Time = 1.03 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.33

method	result
parallelrisch	$\frac{b^2 \operatorname{arctanh}(cx^3)^2 x^6 c^2 + 2x^6 \operatorname{arctanh}(cx^3) abc^2 + a^2 c^2 x^6 + 2b^2 \operatorname{arctanh}(cx^3) x^3 c + 2abcx^3 - b^2 \operatorname{arctanh}(cx^3)^2 + 2 \ln(cx^3 - 1) b^2}{6c^2}$
risch	$\frac{b^2 (c^2 x^6 - 1) \ln(cx^3 + 1)^2}{24c^2} + \frac{b(-2bx^6 \ln(-cx^3 + 1) a c^2 + 4a^2 c^2 x^6 + 4abcx^3 + 2b \ln(-cx^3 + 1) a + b^2) \ln(cx^3 + 1)}{24ac^2} + \frac{b^2 x^6 \ln(-1)}{2}$
default	Expression too large to display
parts	Expression too large to display

input `int(x^5*(a+b*arctanh(c*x^3))^2,x,method=_RETURNVERBOSE)`

output $\frac{1}{6} * (b^2 * \operatorname{arctanh}(c * x^3)^2 * x^6 * c^2 + 2 * x^6 * \operatorname{arctanh}(c * x^3) * a * b * c^2 + a^2 * c^2 * x^6 + 2 * b^2 * \operatorname{arctanh}(c * x^3) * x^3 * c + 2 * a * b * c * x^3 - b^2 * \operatorname{arctanh}(c * x^3)^2 + 2 * \ln(c * x^3 - 1) * b^2 - 2 * \operatorname{arctanh}(c * x^3) * a * b + 2 * \operatorname{arctanh}(c * x^3) * b^2) / c^2$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.52

$$\int x^5 (a + b \operatorname{arctanh}(cx^3))^2 dx$$

$$= \frac{4a^2 c^2 x^6 + 8abcx^3 + (b^2 c^2 x^6 - b^2) \log\left(-\frac{cx^3+1}{cx^3-1}\right)^2 - 4(ab - b^2) \log(cx^3 + 1) + 4(ab + b^2) \log(cx^3 - 1)}{24c^2}$$

input `integrate(x^5*(a+b*arctanh(c*x^3))^2,x, algorithm="fricas")`

output $\frac{1}{24} * (4 * a^2 * c^2 * x^6 + 8 * a * b * c * x^3 + (b^2 * c^2 * x^6 - b^2) * \log(-(c * x^3 + 1) / (c * x^3 - 1))^2 - 4 * (a * b - b^2) * \log(c * x^3 + 1) + 4 * (a * b + b^2) * \log(c * x^3 - 1) + 4 * (a * b * c^2 * x^6 + b^2 * c * x^3) * \log(-(c * x^3 + 1) / (c * x^3 - 1))) / c^2$

Sympy [F(-1)]

Timed out.

$$\int x^5(a + \operatorname{arctanh}(cx^3))^2 dx = \text{Timed out}$$

input `integrate(x**5*(a+b*atanh(c*x**3))**2,x)`

output `Timed out`

Maxima [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 186 vs. $2(81) = 162$.

Time = 0.04 (sec) , antiderivative size = 186, normalized size of antiderivative = 2.04

$$\begin{aligned} \int x^5(a + \operatorname{arctanh}(cx^3))^2 dx &= \frac{1}{6} b^2 x^6 \operatorname{arctanh}(cx^3)^2 + \frac{1}{6} a^2 x^6 \\ &+ \frac{1}{6} \left(2x^6 \operatorname{arctanh}(cx^3) + c \left(\frac{2x^3}{c^2} - \frac{\log(cx^3+1)}{c^3} + \frac{\log(cx^3-1)}{c^3} \right) \right) ab \\ &+ \frac{1}{24} \left(4c \left(\frac{2x^3}{c^2} - \frac{\log(cx^3+1)}{c^3} + \frac{\log(cx^3-1)}{c^3} \right) \operatorname{arctanh}(cx^3) - \frac{2(\log(cx^3-1)-2)\log(cx^3+1)-\log^2(cx^3-1)-\log^2(cx^3+1)}{c^2} \right) b^2 \end{aligned}$$

input `integrate(x^5*(a+b*arctanh(c*x^3))^2,x, algorithm="maxima")`

output `1/6*b^2*x^6*arctanh(c*x^3)^2 + 1/6*a^2*x^6 + 1/6*(2*x^6*arctanh(c*x^3) + c*(2*x^3/c^2 - log(c*x^3 + 1)/c^3 + log(c*x^3 - 1)/c^3))*a*b + 1/24*(4*c*(2*x^3/c^2 - log(c*x^3 + 1)/c^3 + log(c*x^3 - 1)/c^3)*arctanh(c*x^3) - (2*(log(c*x^3 - 1) - 2)*log(c*x^3 + 1) - log(c*x^3 + 1)^2 - log(c*x^3 - 1)^2 - 4*log(c*x^3 - 1))/c^2)*b^2`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 361 vs. $2(81) = 162$.

Time = 0.16 (sec) , antiderivative size = 361, normalized size of antiderivative = 3.97

$$\int x^5 (a + b \operatorname{arctanh}(cx^3))^2 dx$$

$$= \frac{1}{6} \left(\frac{(cx^3 + 1)b^2 \log\left(-\frac{cx^3+1}{cx^3-1}\right)^2}{(cx^3 - 1) \left(\frac{(cx^3+1)^2 c^3}{(cx^3-1)^2} - \frac{2(cx^3+1)c^3}{cx^3-1} + c^3 \right)} + \frac{2 \left(\frac{2(cx^3+1)ab}{cx^3-1} + \frac{(cx^3+1)b^2}{cx^3-1} - b^2 \right) \log\left(-\frac{cx^3+1}{cx^3-1}\right)}{\frac{(cx^3+1)^2 c^3}{(cx^3-1)^2} - \frac{2(cx^3+1)c^3}{cx^3-1} + c^3} + \frac{4 \left(\frac{(cx^3+1)a}{cx^3-1} \right)}{\frac{(cx^3+1)^2 c^3}{(cx^3-1)^2}} \right)$$

input `integrate(x^5*(a+b*arctanh(c*x^3))^2,x, algorithm="giac")`

output

```
1/6*((c*x^3 + 1)*b^2*log(-(c*x^3 + 1)/(c*x^3 - 1))^2/((c*x^3 - 1)*((c*x^3
+ 1)^2*c^3/(c*x^3 - 1)^2 - 2*(c*x^3 + 1)*c^3/(c*x^3 - 1) + c^3)) + 2*(2*(c
*x^3 + 1)*a*b/(c*x^3 - 1) + (c*x^3 + 1)*b^2/(c*x^3 - 1) - b^2)*log(-(c*x^3
+ 1)/(c*x^3 - 1))/((c*x^3 + 1)^2*c^3/(c*x^3 - 1)^2 - 2*(c*x^3 + 1)*c^3/(c
*x^3 - 1) + c^3) + 4*((c*x^3 + 1)*a^2/(c*x^3 - 1) + (c*x^3 + 1)*a*b/(c*x^3
- 1) - a*b)/((c*x^3 + 1)^2*c^3/(c*x^3 - 1)^2 - 2*(c*x^3 + 1)*c^3/(c*x^3 -
1) + c^3) - 2*b^2*log(-(c*x^3 + 1)/(c*x^3 - 1))/c^3 + 2*b^2*log(-(c*x
^3 + 1)/(c*x^3 - 1))/c^3)*c
```

Mupad [B] (verification not implemented)

Time = 4.17 (sec) , antiderivative size = 275, normalized size of antiderivative = 3.02

$$\begin{aligned}
\int x^5 (a + b \operatorname{arctanh}(cx^3))^2 dx &= \frac{a^2 x^6}{6} + \frac{b^2 \ln(cx^3 - 1)}{6c^2} + \frac{b^2 \ln(cx^3 + 1)}{6c^2} \\
&\quad - \frac{b^2 \ln(cx^3 + 1)^2}{24c^2} - \frac{b^2 \ln(1 - cx^3)^2}{24c^2} \\
&\quad + \frac{b^2 x^6 \ln(cx^3 + 1)^2}{24} + \frac{b^2 x^6 \ln(1 - cx^3)^2}{24} \\
&\quad + \frac{b^2 x^3 \ln(cx^3 + 1)}{6c} - \frac{b^2 x^3 \ln(1 - cx^3)}{6c} \\
&\quad + \frac{ab \ln(cx^3 - 1)}{6c^2} - \frac{ab \ln(cx^3 + 1)}{6c^2} \\
&\quad + \frac{abx^6 \ln(cx^3 + 1)}{6} - \frac{abx^6 \ln(1 - cx^3)}{6} \\
&\quad + \frac{b^2 \ln(cx^3 + 1) \ln(1 - cx^3)}{12c^2} + \frac{abx^3}{3c} \\
&\quad - \frac{b^2 x^6 \ln(cx^3 + 1) \ln(1 - cx^3)}{12}
\end{aligned}$$

input `int(x^5*(a + b*atanh(c*x^3))^2,x)`

output

```

(a^2*x^6)/6 + (b^2*log(c*x^3 - 1))/(6*c^2) + (b^2*log(c*x^3 + 1))/(6*c^2)
- (b^2*log(c*x^3 + 1)^2)/(24*c^2) - (b^2*log(1 - c*x^3)^2)/(24*c^2) + (b^2
*x^6*log(c*x^3 + 1)^2)/24 + (b^2*x^6*log(1 - c*x^3)^2)/24 + (b^2*x^3*log(c
*x^3 + 1))/(6*c) - (b^2*x^3*log(1 - c*x^3))/(6*c) + (a*b*log(c*x^3 - 1))/(
6*c^2) - (a*b*log(c*x^3 + 1))/(6*c^2) + (a*b*x^6*log(c*x^3 + 1))/6 - (a*b*
x^6*log(1 - c*x^3))/6 + (b^2*log(c*x^3 + 1)*log(1 - c*x^3))/(12*c^2) + (a*
b*x^3)/(3*c) - (b^2*x^6*log(c*x^3 + 1)*log(1 - c*x^3))/12

```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.57

$$\int x^5 (a + b \operatorname{arctanh}(cx^3))^2 dx$$

$$= \frac{a \operatorname{tanh}(cx^3)^2 b^2 c^2 x^6 - a \operatorname{tanh}(cx^3)^2 b^2 + 2 a \operatorname{tanh}(cx^3) a b c^2 x^6 - 2 a \operatorname{tanh}(cx^3) a b + 2 a \operatorname{tanh}(cx^3) b^2 c x^3 - 2 a^2 c^2 x^6 + 2 a^2 b c x^3}{6c^2}$$

input

```
int(x^5*(a+b*atanh(c*x^3))^2,x)
```

output

```
(atanh(c*x**3)**2*b**2*c**2*x**6 - atanh(c*x**3)**2*b**2 + 2*atanh(c*x**3)
*a*b*c**2*x**6 - 2*atanh(c*x**3)*a*b + 2*atanh(c*x**3)*b**2*c*x**3 - 2*ata
nh(c*x**3)*b**2 + 2*log(c**(2/3)*x**2 - c**(1/3)*x + 1)*b**2 + 2*log(c**(2
/3)*x + c**(1/3))*b**2 + a**2*c**2*x**6 + 2*a*b*c*x**3)/(6*c**2)
```

3.119 $\int x^2(a + \operatorname{barctanh}(cx^3))^2 dx$

Optimal result	1016
Mathematica [A] (verified)	1016
Rubi [A] (verified)	1017
Maple [A] (verified)	1019
Fricas [F]	1020
Sympy [F(-1)]	1020
Maxima [F]	1020
Giac [F]	1021
Mupad [F(-1)]	1021
Reduce [F]	1022

Optimal result

Integrand size = 16, antiderivative size = 96

$$\int x^2(a + \operatorname{barctanh}(cx^3))^2 dx = \frac{(a + \operatorname{barctanh}(cx^3))^2}{3c} + \frac{1}{3}x^3(a + \operatorname{barctanh}(cx^3))^2 - \frac{2b(a + \operatorname{barctanh}(cx^3)) \log\left(\frac{2}{1-cx^3}\right)}{3c} - \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^3}\right)}{3c}$$

output `1/3*(a+b*arctanh(c*x^3))^2/c+1/3*x^3*(a+b*arctanh(c*x^3))^2-2/3*b*(a+b*arctanh(c*x^3))*ln(2/(-c*x^3+1))/c-1/3*b^2*polylog(2,1-2/(-c*x^3+1))/c`

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.03

$$\int x^2(a + \operatorname{barctanh}(cx^3))^2 dx = \frac{b^2(-1 + cx^3) \operatorname{arctanh}(cx^3)^2 + 2\operatorname{barctanh}(cx^3) \left(acx^3 - b \log\left(1 + e^{-2\operatorname{arctanh}(cx^3)}\right)\right) + a(acx^3 + b \log(1 -$$

3c

input `Integrate[x^2*(a + b*ArcTanh[c*x^3])^2,x]`

output

```
(b^2*(-1 + c*x^3)*ArcTanh[c*x^3]^2 + 2*b*ArcTanh[c*x^3]*(a*c*x^3 - b*Log[1
+ E^(-2*ArcTanh[c*x^3])]) + a*(a*c*x^3 + b*Log[1 - c^2*x^6]) + b^2*PolyLo
g[2, -E^(-2*ArcTanh[c*x^3])])/(3*c)
```

Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.08, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6454, 6436, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 (a + \operatorname{barctanh}(cx^3))^2 dx$$

$$\downarrow 6454$$

$$\frac{1}{3} \int (a + \operatorname{barctanh}(cx^3))^2 dx^3$$

$$\downarrow 6436$$

$$\frac{1}{3} \left(x^3 (a + \operatorname{barctanh}(cx^3))^2 - 2bc \int \frac{x^3 (a + \operatorname{barctanh}(cx^3))}{1 - c^2 x^6} dx^3 \right)$$

$$\downarrow 6546$$

$$\frac{1}{3} \left(x^3 (a + \operatorname{barctanh}(cx^3))^2 - 2bc \left(\frac{\int \frac{a + \operatorname{barctanh}(cx^3)}{1 - cx^3} dx^3}{c} - \frac{(a + \operatorname{barctanh}(cx^3))^2}{2bc^2} \right) \right)$$

$$\downarrow 6470$$

$$\frac{1}{3} \left(x^3 (a + \operatorname{barctanh}(cx^3))^2 - 2bc \left(\frac{\frac{\log\left(\frac{2}{1 - cx^3}\right) (a + \operatorname{barctanh}(cx^3))}{c} - b \int \frac{\log\left(\frac{2}{1 - cx^3}\right)}{1 - c^2 x^6} dx^3}{c} - \frac{(a + \operatorname{barctanh}(cx^3))^2}{2bc^2} \right) \right)$$

$$\downarrow 2849$$

$$\frac{1}{3} \left(x^3 (a + \operatorname{arctanh}(cx^3))^2 - 2bc \left(\frac{b \int \frac{\log\left(\frac{2}{1-cx^3}\right) d\frac{1}{1-cx^3}}{c} + \frac{\log\left(\frac{2}{1-cx^3}\right) (a + \operatorname{arctanh}(cx^3))}{c}}{c} - \frac{(a + \operatorname{arctanh}(cx^3))^2}{2bc^2} \right) \right)$$

↓ 2752

$$\frac{1}{3} \left(x^3 (a + \operatorname{arctanh}(cx^3))^2 - 2bc \left(\frac{\log\left(\frac{2}{1-cx^3}\right) (a + \operatorname{arctanh}(cx^3))}{c} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^3}\right)}{2c} - \frac{(a + \operatorname{arctanh}(cx^3))^2}{2bc^2} \right) \right)$$

input `Int[x^2*(a + b*ArcTanh[c*x^3])^2,x]`

output `(x^3*(a + b*ArcTanh[c*x^3])^2 - 2*b*c*(-1/2*(a + b*ArcTanh[c*x^3])^2/(b*c^2) + ((a + b*ArcTanh[c*x^3])*Log[2/(1 - c*x^3)])/c + (b*PolyLog[2, 1 - 2/(1 - c*x^3)])/(2*c))/c)/3`

Defintions of rubi rules used

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 6436 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^p, x_Symbol] := Simp[x*(a + b*ArcTanh[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTanh[c*x^n])^(p-1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`

```
rule 6454 Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
  Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x
], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simpl
ify[(m + 1)/n]]
```

```
rule 6470 Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_.) + (e_.)*(x_)), x_Symbol
] := Simp[(-a + b*ArcTanh[c*x])^p*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c
*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^
2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2
, 0]
```

```
rule 6546 Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*(x_)/((d_.) + (e_.)*(x_)^2),
x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/
(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

Maple [A] (verified)

Time = 1.10 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.40

method	result
derivativedivides	$\frac{cx^3a^2+b^2 \left(\operatorname{arctanh}(cx^3)^2(cx^3-1)+2\operatorname{arctanh}(cx^3)^2-2\operatorname{arctanh}(cx^3)\ln\left(1+\frac{(cx^3+1)^2}{-c^2x^6+1}\right)-\operatorname{polylog}\left(2,-\frac{(cx^3+1)^2}{-c^2x^6+1}\right)}{3c}$
default	$\frac{cx^3a^2+b^2 \left(\operatorname{arctanh}(cx^3)^2(cx^3-1)+2\operatorname{arctanh}(cx^3)^2-2\operatorname{arctanh}(cx^3)\ln\left(1+\frac{(cx^3+1)^2}{-c^2x^6+1}\right)-\operatorname{polylog}\left(2,-\frac{(cx^3+1)^2}{-c^2x^6+1}\right)}{3c}$
parts	$\frac{a^2x^3}{3} + \frac{b^2 \left(\operatorname{arctanh}(cx^3)^2(cx^3-1)+2\operatorname{arctanh}(cx^3)^2-2\operatorname{arctanh}(cx^3)\ln\left(1+\frac{(cx^3+1)^2}{-c^2x^6+1}\right)-\operatorname{polylog}\left(2,-\frac{(cx^3+1)^2}{-c^2x^6+1}\right)}{3c}$
risch	$-\frac{b^2 \operatorname{dilog}\left(\frac{cx^3}{2}+\frac{1}{2}\right)}{3c} - \frac{b^2 \ln(cx^3-1)}{3c} + \frac{a^2x^3}{3} - \frac{\ln(-cx^3+1)abx^3}{3} + \frac{\ln(-cx^3+1)ab}{3c} + \frac{b^2 \ln(cx^3+1)^2x^3}{12} +$

```
input int(x^2*(a+b*arctanh(c*x^3))^2,x,method=_RETURNVERBOSE)
```


output

```
1/3/c*(c*x^3*a^2+b^2*(arctanh(c*x^3)^2*(c*x^3-1)+2*arctanh(c*x^3)^2-2*arctanh(c*x^3)*ln(1+(c*x^3+1)^2/(-c^2*x^6+1))-polylog(2,-(c*x^3+1)^2/(-c^2*x^6+1)))+2*a*b*c*x^3*arctanh(c*x^3)+a*b*ln(-c^2*x^6+1))
```

Fricas [F]

$$\int x^2(a + b \operatorname{arctanh}(cx^3))^2 dx = \int (b \operatorname{arctanh}(cx^3) + a)^2 x^2 dx$$

input

```
integrate(x^2*(a+b*arctanh(c*x^3))^2,x, algorithm="fricas")
```

output

```
integral(b^2*x^2*arctanh(c*x^3)^2 + 2*a*b*x^2*arctanh(c*x^3) + a^2*x^2, x)
```

Sympy [F(-1)]

Timed out.

$$\int x^2(a + b \operatorname{arctanh}(cx^3))^2 dx = \text{Timed out}$$

input

```
integrate(x**2*(a+b*atanh(c*x**3))**2,x)
```

output

Timed out

Maxima [F]

$$\int x^2(a + b \operatorname{arctanh}(cx^3))^2 dx = \int (b \operatorname{arctanh}(cx^3) + a)^2 x^2 dx$$

input

```
integrate(x^2*(a+b*arctanh(c*x^3))^2,x, algorithm="maxima")
```

output

```
1/3*a^2*x^3 + 1/12*(x^3*log(-c*x^3 + 1)^2 - c^2*(2*x^3/c^2 - log(c*x^3 + 1)
)/c^3 + log(c*x^3 - 1)/c^3) - 2*(x^3/c + log(c*x^3 - 1)/c^2)*c*log(-c*x^3
+ 1) + 18*c*integrate(x^5*log(c*x^3 + 1)/(c^2*x^6 - 1), x) + (c*x^3*log(c*
x^3 + 1)^2 + 2*(c*x^3 - (c*x^3 + 1)*log(c*x^3 + 1))*log(-c*x^3 + 1))/c + (
2*c*x^3 + log(c*x^3 - 1)^2 + 2*log(c*x^3 - 1))/c - log(c^2*x^6 - 1)/c + 6*
integrate(x^2*log(c*x^3 + 1)/(c^2*x^6 - 1), x))*b^2 + 1/3*(2*c*x^3*arctanh
(c*x^3) + log(-c^2*x^6 + 1))*a*b/c
```

Giac [F]

$$\int x^2 (a + b \operatorname{arctanh}(cx^3))^2 dx = \int (b \operatorname{artanh}(cx^3) + a)^2 x^2 dx$$

input

```
integrate(x^2*(a+b*arctanh(c*x^3))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^3) + a)^2*x^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 (a + b \operatorname{arctanh}(cx^3))^2 dx = \int x^2 (a + b \operatorname{atanh}(cx^3))^2 dx$$

input

```
int(x^2*(a + b*atanh(c*x^3))^2,x)
```

output

```
int(x^2*(a + b*atanh(c*x^3))^2, x)
```

Reduce [F]

$$\int x^2 (a + b \operatorname{arctanh}(cx^3))^2 dx$$

$$= \frac{2 \operatorname{atanh}(cx^3) abc x^3 - 2 \operatorname{atanh}(cx^3) ab + 3 \left(\int \operatorname{atanh}(cx^3)^2 x^2 dx \right) b^2 c + 2 \log\left(c^{\frac{2}{3}} x^2 - c^{\frac{1}{3}} x + 1\right) ab + 2 \log\left(c^{\frac{2}{3}} x^2 - c^{\frac{1}{3}} x + 1\right) ab}{3c}$$

input `int(x^2*(a+b*atanh(c*x^3))^2,x)`

output `(2*atanh(c*x**3)*a*b*c*x**3 - 2*atanh(c*x**3)*a*b + 3*int(atanh(c*x**3)**2*x**2,x)*b**2*c + 2*log(c**(2/3)*x**2 - c**(1/3)*x + 1)*a*b + 2*log(c**(2/3)*x + c**(1/3))*a*b + a**2*c*x**3)/(3*c)`

$$3.120 \quad \int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x} dx$$

Optimal result	1023
Mathematica [C] (verified)	1024
Rubi [A] (verified)	1025
Maple [F]	1027
Fricas [F]	1027
Sympy [F]	1027
Maxima [F]	1028
Giac [F]	1028
Mupad [F(-1)]	1028
Reduce [F]	1029

Optimal result

Integrand size = 16, antiderivative size = 140

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x} dx = & \frac{2}{3} (a + b \operatorname{arctanh}(cx^3))^2 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx^3}\right) \\ & - \frac{1}{3} b (a + b \operatorname{arctanh}(cx^3)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx^3}\right) \\ & + \frac{1}{3} b (a + b \operatorname{arctanh}(cx^3)) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - cx^3}\right) \\ & + \frac{1}{6} b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - cx^3}\right) \\ & - \frac{1}{6} b^2 \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 - cx^3}\right) \end{aligned}$$

output

```
-2/3*(a+b*arctanh(c*x^3))^2*arctanh(-1+2/(-c*x^3+1))-1/3*b*(a+b*arctanh(c*
x^3))*polylog(2,1-2/(-c*x^3+1))+1/3*b*(a+b*arctanh(c*x^3))*polylog(2,-1+2/
(-c*x^3+1))+1/6*b^2*polylog(3,1-2/(-c*x^3+1))-1/6*b^2*polylog(3,-1+2/(-c*x
^3+1))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.29

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x} dx = a^2 \log(x) + \frac{1}{3} ab (-\operatorname{PolyLog}(2, -cx^3) + \operatorname{PolyLog}(2, cx^3))$$

$$+ \frac{1}{3} b^2 \left(\frac{i\pi^3}{24} - \frac{2}{3} \operatorname{arctanh}(cx^3)^3 \right.$$

$$\quad - \operatorname{arctanh}(cx^3)^2 \log(1 + e^{-2\operatorname{arctanh}(cx^3)})$$

$$\quad + \operatorname{arctanh}(cx^3)^2 \log(1 - e^{2\operatorname{arctanh}(cx^3)})$$

$$+ \operatorname{arctanh}(cx^3) \operatorname{PolyLog}(2, -e^{-2\operatorname{arctanh}(cx^3)})$$

$$+ \operatorname{arctanh}(cx^3) \operatorname{PolyLog}(2, e^{2\operatorname{arctanh}(cx^3)})$$

$$\quad + \frac{1}{2} \operatorname{PolyLog}(3, -e^{-2\operatorname{arctanh}(cx^3)})$$

$$\quad \left. - \frac{1}{2} \operatorname{PolyLog}(3, e^{2\operatorname{arctanh}(cx^3)}) \right)$$

input `Integrate[(a + b*ArcTanh[c*x^3])^2/x,x]`

output `a^2*Log[x] + (a*b*(-PolyLog[2, -(c*x^3)] + PolyLog[2, c*x^3]))/3 + (b^2*((I/24)*Pi^3 - (2*ArcTanh[c*x^3]^3)/3 - ArcTanh[c*x^3]^2*Log[1 + E^(-2*ArcTanh[c*x^3])] + ArcTanh[c*x^3]^2*Log[1 - E^(2*ArcTanh[c*x^3])] + ArcTanh[c*x^3]*PolyLog[2, -E^(-2*ArcTanh[c*x^3])] + ArcTanh[c*x^3]*PolyLog[2, E^(2*ArcTanh[c*x^3])] + PolyLog[3, -E^(-2*ArcTanh[c*x^3])]/2 - PolyLog[3, E^(2*ArcTanh[c*x^3])]/2))/3`

Rubi [A] (verified)

Time = 0.84 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.16, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6450, 6448, 6614, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x} dx$$

$$\downarrow 6450$$

$$\frac{1}{3} \int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^3} dx^3$$

$$\downarrow 6448$$

$$\frac{1}{3} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^3} \right) (a + b \operatorname{arctanh}(cx^3))^2 - 4bc \int \frac{(a + b \operatorname{arctanh}(cx^3)) \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^3} \right)}{1 - c^2 x^6} dx^3 \right)$$

$$\downarrow 6614$$

$$\frac{1}{3} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^3} \right) (a + b \operatorname{arctanh}(cx^3))^2 - 4bc \left(\frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(cx^3)) \log \left(2 - \frac{2}{1 - cx^3} \right)}{1 - c^2 x^6} dx^3 - \frac{1}{2} \int \right) \right)$$

$$\downarrow 6620$$

$$\frac{1}{3} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^3} \right) (a + b \operatorname{arctanh}(cx^3))^2 - 4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - cx^3} \right) (a + b \operatorname{arctanh}(cx^3))}{2c} - \frac{1}{2} \right) \right) \right)$$

$$\downarrow 7164$$

$$\frac{1}{3} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^3} \right) (a + b \operatorname{arctanh}(cx^3))^2 - 4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - cx^3} \right) (a + b \operatorname{arctanh}(cx^3))}{2c} - \frac{1}{2} \right) \right) \right)$$

input

```
Int[(a + b*ArcTanh[c*x^3])^2/x, x]
```

output

$$(2*(a + b*\text{ArcTanh}[c*x^3])^2*\text{ArcTanh}[1 - 2/(1 - c*x^3)] - 4*b*c*((a + b*\text{ArcTanh}[c*x^3])*PolyLog[2, 1 - 2/(1 - c*x^3)]/(2*c) - (b*PolyLog[3, 1 - 2/(1 - c*x^3)]/(4*c))/2 + (-1/2*(a + b*\text{ArcTanh}[c*x^3])*PolyLog[2, -1 + 2/(1 - c*x^3)]/c + (b*PolyLog[3, -1 + 2/(1 - c*x^3)]/(4*c))/2))/3$$

Defintions of rubi rules used

rule 6448

$$\text{Int}[(a + \text{ArcTanh}[c*x])^p/(x), x_Symbol] \rightarrow \text{Simp}[2*(a + b*\text{ArcTanh}[c*x])^p*\text{ArcTanh}[1 - 2/(1 - c*x)], x] - \text{Simp}[2*b*c*p \text{ Int}[(a + b*\text{ArcTanh}[c*x])^{p-1}*(\text{ArcTanh}[1 - 2/(1 - c*x)]/(1 - c^2*x^2)), x], x] /;$$

$$\text{FreeQ}\{a, b, c, x\} \ \&\& \ \text{IGtQ}[p, 1]$$

rule 6450

$$\text{Int}[(a + \text{ArcTanh}[c*x]^n)/(x), x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[(a + b*\text{ArcTanh}[c*x])^p/x, x], x, x^n], x] /;$$

$$\text{FreeQ}\{a, b, c, n, x\} \ \&\& \ \text{IGtQ}[p, 0]$$

rule 6614

$$\text{Int}[(\text{ArcTanh}[u]*(a + \text{ArcTanh}[c*x])^p)/(d + e*x^2), x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Int}[\text{Log}[1 + u]*(a + b*\text{ArcTanh}[c*x])^p/(d + e*x^2), x], x] - \text{Simp}[1/2 \text{ Int}[\text{Log}[1 - u]*(a + b*\text{ArcTanh}[c*x])^p/(d + e*x^2), x], x] /;$$

$$\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{EqQ}[u^2 - (1 - 2/(1 - c*x))^2, 0]$$

rule 6620

$$\text{Int}[(\text{Log}[u]*(a + \text{ArcTanh}[c*x])^p)/(d + e*x^2), x_Symbol] \rightarrow \text{Simp}[(-a + b*\text{ArcTanh}[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)), x] + \text{Simp}[b*(p/2 \text{ Int}[(a + b*\text{ArcTanh}[c*x])^{p-1}*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /;$$

$$\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{EqQ}[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]$$

rule 7164

$$\text{Int}[u*PolyLog[n, v], x_Symbol] \rightarrow \text{With}\{w = \text{DerivativeDivides}[v, u*v, x]\}, \text{Simp}[w*PolyLog[n + 1, v], x] /;$$

$$\text{!FalseQ}[w] /;$$

$$\text{FreeQ}[n, x]$$

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x} dx$$

input `int((a+b*arctanh(c*x^3))^2/x,x)`

output `int((a+b*arctanh(c*x^3))^2/x,x)`

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x} dx = \int \frac{(b \operatorname{arctanh}(cx^3) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x^3))^2/x,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x^3)^2 + 2*a*b*arctanh(c*x^3) + a^2)/x, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x} dx = \int \frac{(a + b \operatorname{atanh}(cx^3))^2}{x} dx$$

input `integrate((a+b*atanh(c*x**3))**2/x,x)`

output `Integral((a + b*atanh(c*x**3))**2/x, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x} dx = \int \frac{(b \operatorname{artanh}(cx^3) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x^3))^2/x,x, algorithm="maxima")`

output `a^2*log(x) + integrate(1/4*b^2*(log(c*x^3 + 1) - log(-c*x^3 + 1))^2/x + a*b*(log(c*x^3 + 1) - log(-c*x^3 + 1))/x, x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x} dx = \int \frac{(b \operatorname{artanh}(cx^3) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x^3))^2/x,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^3) + a)^2/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x} dx = \int \frac{(a + b \operatorname{atanh}(cx^3))^2}{x} dx$$

input `int((a + b*atanh(c*x^3))^2/x,x)`

output `int((a + b*atanh(c*x^3))^2/x, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x} dx = 2 \left(\int \frac{\operatorname{atanh}(cx^3)}{x} dx \right) ab + \left(\int \frac{\operatorname{atanh}(cx^3)^2}{x} dx \right) b^2 + \log(x) a^2$$

input `int((a+b*atanh(c*x^3))^2/x,x)`

output `2*int(atanh(c*x**3)/x,x)*a*b + int(atanh(c*x**3)**2/x,x)*b**2 + log(x)*a**2`

3.121 $\int \frac{(a+b\operatorname{arctanh}(cx^3))^2}{x^4} dx$

Optimal result	1030
Mathematica [A] (verified)	1030
Rubi [A] (verified)	1031
Maple [C] (warning: unable to verify)	1033
Fricas [F]	1034
Sympy [F(-1)]	1034
Maxima [F]	1034
Giac [F]	1035
Mupad [F(-1)]	1035
Reduce [F]	1035

Optimal result

Integrand size = 16, antiderivative size = 90

$$\int \frac{(a + b\operatorname{arctanh}(cx^3))^2}{x^4} dx = \frac{1}{3}c(a + b\operatorname{arctanh}(cx^3))^2 - \frac{(a + b\operatorname{arctanh}(cx^3))^2}{3x^3} + \frac{2}{3}bc(a + b\operatorname{arctanh}(cx^3)) \log\left(2 - \frac{2}{1 + cx^3}\right) - \frac{1}{3}b^2c \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + cx^3}\right)$$

output `1/3*c*(a+b*arctanh(c*x^3))^2-1/3*(a+b*arctanh(c*x^3))^2/x^3+2/3*b*c*(a+b*arctanh(c*x^3))*ln(2-2/(c*x^3+1))-1/3*b^2*c*polylog(2,-1+2/(c*x^3+1))`

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.30

$$\int \frac{(a + b\operatorname{arctanh}(cx^3))^2}{x^4} dx = \frac{b^2(-1 + cx^3) \operatorname{arctanh}(cx^3)^2 + 2b\operatorname{arctanh}(cx^3) \left(-a + bcx^3 \log\left(1 - e^{-2\operatorname{arctanh}(cx^3)}\right)\right) - a(a - 2bcx^3 \log(c))}{3x^3}$$

input `Integrate[(a + b*ArcTanh[c*x^3])^2/x^4, x]`

output `(b^2*(-1 + c*x^3)*ArcTanh[c*x^3]^2 + 2*b*ArcTanh[c*x^3]*(-a + b*c*x^3*Log[1 - E^(-2*ArcTanh[c*x^3])]) - a*(a - 2*b*c*x^3*Log[c*x^3] + b*c*x^3*Log[1 - c^2*x^6]) - b^2*c*x^3*PolyLog[2, E^(-2*ArcTanh[c*x^3])])/(3*x^3)`

Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.01, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6454, 6452, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^4} dx \\
 & \quad \downarrow 6454 \\
 & \frac{1}{3} \int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^6} dx^3 \\
 & \quad \downarrow 6452 \\
 & \frac{1}{3} \left(2bc \int \frac{a + b \operatorname{arctanh}(cx^3)}{x^3(1 - c^2x^6)} dx^3 - \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^3} \right) \\
 & \quad \downarrow 6550 \\
 & \frac{1}{3} \left(2bc \left(\int \frac{a + b \operatorname{arctanh}(cx^3)}{x^3(cx^3 + 1)} dx^3 + \frac{(a + b \operatorname{arctanh}(cx^3))^2}{2b} \right) - \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^3} \right) \\
 & \quad \downarrow 6494 \\
 & \frac{1}{3} \left(2bc \left(-bc \int \frac{\log\left(2 - \frac{2}{cx^3+1}\right)}{1 - c^2x^6} dx^3 + \frac{(a + b \operatorname{arctanh}(cx^3))^2}{2b} + \log\left(2 - \frac{2}{cx^3+1}\right) (a + b \operatorname{arctanh}(cx^3)) \right) - \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^3} \right) \\
 & \quad \downarrow 2897
 \end{aligned}$$

$$\frac{1}{3} \left(2bc \left(\frac{(a + b \operatorname{arctanh}(cx^3))^2}{2b} + \log \left(2 - \frac{2}{cx^3 + 1} \right) (a + b \operatorname{arctanh}(cx^3)) - \frac{1}{2} b \operatorname{PolyLog} \left(2, \frac{2}{cx^3 + 1} - 1 \right) \right) - \right)$$

input `Int[(a + b*ArcTanh[c*x^3])^2/x^4,x]`

output `(-((a + b*ArcTanh[c*x^3])^2/x^3) + 2*b*c*((a + b*ArcTanh[c*x^3])^2/(2*b) + (a + b*ArcTanh[c*x^3])*Log[2 - 2/(1 + c*x^3)] - (b*PolyLog[2, -1 + 2/(1 + c*x^3)]))/2)/3`

Defintions of rubi rules used

rule 2897 `Int[Log[u]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6494 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((x)*((d_) + (e_)*(x))), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6550

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)/((x_.)*((d_.) + (e_.)*(x_)^2)),
 x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/
 d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.87 (sec) , antiderivative size = 2993, normalized size of antiderivative = 33.26

method	result	size
default	Expression too large to display	2993
parts	Expression too large to display	2993

input

```
int((a+b*arctanh(c*x^3))^2/x^4,x,method=_RETURNVERBOSE)
```

output

```
-1/3*a^2/x^3+b^2*(-1/3/x^3*arctanh(c*x^3)^2+2*c*(arctanh(c*x^3)*ln(x)-1/6*
arctanh(c*x^3)*ln(c*x^3-1)-1/6*arctanh(c*x^3)*ln(c*x^3+1)-1/2*c*(Sum(1/6*(
ln(x-_alpha)*ln(c*x^3-1)-3*c*(1/6/_alpha^2/c*ln(x-_alpha)^2-1/3*_alpha*ln(
x-_alpha)*(2*ln((RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1)-x+_alpha)/Ro
otOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1))*RootOf(_Z^2+3*_Z*_alpha+3*_alpha
^2,index=1)*RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=2)+3*ln((RootOf(_Z^2+
3*_Z*_alpha+3*_alpha^2,index=1)-x+_alpha)/RootOf(_Z^2+3*_Z*_alpha+3*_alpha
^2,index=1))*RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1)*_alpha+6*ln((Root
Of(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1)-x+_alpha)/RootOf(_Z^2+3*_Z*_alpha+
3*_alpha^2,index=1))*RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=2)*_alpha+9*
ln((RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1)-x+_alpha)/RootOf(_Z^2+3*_Z
*_alpha+3*_alpha^2,index=1))*_alpha^2+2*ln((RootOf(_Z^2+3*_Z*_alpha+3*_alp
ha^2,index=2)-x+_alpha)/RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=2))*Root0
f(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1)*RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,
index=2)+6*ln((RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=2)-x+_alpha)/Root0
f(_Z^2+3*_Z*_alpha+3*_alpha^2,index=2))*RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2
,index=1)*_alpha+3*ln((RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=2)-x+_alph
a)/RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=2))*RootOf(_Z^2+3*_Z*_alpha+3*
_alpha^2,index=2)*_alpha+9*ln((RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=2)
-x+_alpha)/RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=2))*_alpha^2)/(3*_a...
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^4} dx = \int \frac{(b \operatorname{artanh}(cx^3) + a)^2}{x^4} dx$$

input `integrate((a+b*arctanh(c*x^3))^2/x^4,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x^3)^2 + 2*a*b*arctanh(c*x^3) + a^2)/x^4, x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^4} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**3))**2/x**4,x)`

output `Timed out`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^4} dx = \int \frac{(b \operatorname{artanh}(cx^3) + a)^2}{x^4} dx$$

input `integrate((a+b*arctanh(c*x^3))^2/x^4,x, algorithm="maxima")`

output `-1/3*(c*(log(c^2*x^6 - 1) - log(x^6)) + 2*arctanh(c*x^3)/x^3)*a*b - 1/12*b^2*(log(-c*x^3 + 1)^2/x^3 + 3*integrate(-((c*x^3 - 1)*log(c*x^3 + 1)^2 + 2*(c*x^3 - (c*x^3 - 1)*log(c*x^3 + 1))*log(-c*x^3 + 1))/(c*x^7 - x^4), x)) - 1/3*a^2/x^3`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^4} dx = \int \frac{(b \operatorname{artanh}(cx^3) + a)^2}{x^4} dx$$

input `integrate((a+b*arctanh(c*x^3))^2/x^4,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^3) + a)^2/x^4, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^4} dx = \int \frac{(a + b \operatorname{atanh}(cx^3))^2}{x^4} dx$$

input `int((a + b*atanh(c*x^3))^2/x^4,x)`

output `int((a + b*atanh(c*x^3))^2/x^4, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^4} dx = \frac{-\operatorname{atanh}(cx^3)^2 b^2 + 2 \operatorname{atanh}(cx^3) abc x^3 - 2 \operatorname{atanh}(cx^3) ab - 6 \left(\int \frac{\operatorname{atanh}(cx^3)}{c^2 x^7 - x} dx \right) b^2 c x^3 - 2 \log\left(c^{\frac{2}{3}} x^2 - c^{\frac{1}{3}} x\right)}{3x^3}$$

input `int((a+b*atanh(c*x^3))^2/x^4,x)`

output `(- atanh(c*x**3)**2*b**2 + 2*atanh(c*x**3)*a*b*c*x**3 - 2*atanh(c*x**3)*a*b - 6*int(atanh(c*x**3)/(c**2*x**7 - x),x)*b**2*c*x**3 - 2*log(c**(2/3)*x**2 - c**(1/3)*x + 1)*a*b*c*x**3 - 2*log(c**(2/3)*x + c**(1/3))*a*b*c*x**3 + 6*log(x)*a*b*c*x**3 - a**2)/(3*x**3)`

$$3.122 \quad \int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^7} dx$$

Optimal result	1036
Mathematica [A] (verified)	1036
Rubi [A] (verified)	1037
Maple [A] (verified)	1040
Fricas [A] (verification not implemented)	1040
Sympy [F(-1)]	1041
Maxima [B] (verification not implemented)	1041
Giac [F]	1042
Mupad [B] (verification not implemented)	1042
Reduce [B] (verification not implemented)	1043

Optimal result

Integrand size = 16, antiderivative size = 88

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^7} dx = -\frac{bc(a + b \operatorname{arctanh}(cx^3))}{3x^3} + \frac{1}{6}c^2(a + b \operatorname{arctanh}(cx^3))^2 - \frac{(a + b \operatorname{arctanh}(cx^3))^2}{6x^6} + b^2c^2 \log(x) - \frac{1}{6}b^2c^2 \log(1 - c^2x^6)$$

output

```
-1/3*b*c*(a+b*arctanh(c*x^3))/x^3+1/6*c^2*(a+b*arctanh(c*x^3))^2-1/6*(a+b*arctanh(c*x^3))^2/x^6+b^2*c^2*ln(x)-1/6*b^2*c^2*ln(-c^2*x^6+1)
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.26

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^7} dx = \frac{1}{6} \left(-\frac{a^2}{x^6} - \frac{2abc}{x^3} - \frac{2b(a + bcx^3) \operatorname{arctanh}(cx^3)}{x^6} + \frac{b^2(-1 + c^2x^6) \operatorname{arctanh}(cx^3)^2}{x^6} + 6b^2c^2 \log(x) - b(a + b)c^2 \log(1 - cx^3) + (a - b)bc^2 \log(1 + cx^3) \right)$$

input `Integrate[(a + b*ArcTanh[c*x^3])^2/x^7,x]`

output $(-a^2/x^6) - (2*a*b*c)/x^3 - (2*b*(a + b*c*x^3)*ArcTanh[c*x^3])/x^6 + (b^2*(-1 + c^2*x^6)*ArcTanh[c*x^3]^2)/x^6 + 6*b^2*c^2*Log[x] - b*(a + b)*c^2*Log[1 - c*x^3] + (a - b)*b*c^2*Log[1 + c*x^3])/6$

Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.99, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.562$, Rules used = {6454, 6452, 6544, 6452, 243, 47, 14, 16, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^7} dx$$

$$\downarrow 6454$$

$$\frac{1}{3} \int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^9} dx^3$$

$$\downarrow 6452$$

$$\frac{1}{3} \left(bc \int \frac{a + b \operatorname{arctanh}(cx^3)}{x^6 (1 - c^2 x^6)} dx^3 - \frac{(a + b \operatorname{arctanh}(cx^3))^2}{2x^6} \right)$$

$$\downarrow 6544$$

$$\frac{1}{3} \left(bc \left(c^2 \int \frac{a + b \operatorname{arctanh}(cx^3)}{1 - c^2 x^6} dx^3 + \int \frac{a + b \operatorname{arctanh}(cx^3)}{x^6} dx^3 \right) - \frac{(a + b \operatorname{arctanh}(cx^3))^2}{2x^6} \right)$$

$$\downarrow 6452$$

$$\frac{1}{3} \left(bc \left(c^2 \int \frac{a + b \operatorname{arctanh}(cx^3)}{1 - c^2 x^6} dx^3 + bc \int \frac{1}{x^3 (1 - c^2 x^6)} dx^3 - \frac{a + b \operatorname{arctanh}(cx^3)}{x^3} \right) - \frac{(a + b \operatorname{arctanh}(cx^3))^2}{2x^6} \right)$$

$$\downarrow 243$$

$$\frac{1}{3} \left(bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^3)}{1 - c^2x^6} dx^3 + \frac{1}{2} bc \int \frac{1}{x^3(1 - c^2x^6)} dx^6 - \frac{a + \operatorname{barctanh}(cx^3)}{x^3} \right) - \frac{(a + \operatorname{barctanh}(cx^3))^2}{2x^6} \right)$$

↓ 47

$$\frac{1}{3} \left(bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^3)}{1 - c^2x^6} dx^3 + \frac{1}{2} bc \left(c^2 \int \frac{1}{1 - c^2x^6} dx^6 + \int \frac{1}{x^3} dx^6 \right) - \frac{a + \operatorname{barctanh}(cx^3)}{x^3} \right) - \frac{(a + \operatorname{barctanh}(cx^3))^2}{2x^6} \right)$$

↓ 14

$$\frac{1}{3} \left(bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^3)}{1 - c^2x^6} dx^3 + \frac{1}{2} bc \left(c^2 \int \frac{1}{1 - c^2x^6} dx^6 + \log(x^6) \right) - \frac{a + \operatorname{barctanh}(cx^3)}{x^3} \right) - \frac{(a + \operatorname{barctanh}(cx^3))^2}{2x^6} \right)$$

↓ 16

$$\frac{1}{3} \left(bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^3)}{1 - c^2x^6} dx^3 - \frac{a + \operatorname{barctanh}(cx^3)}{x^3} + \frac{1}{2} bc (\log(x^6) - \log(1 - c^2x^6)) \right) - \frac{(a + \operatorname{barctanh}(cx^3))^2}{2x^6} \right)$$

↓ 6510

$$\frac{1}{3} \left(bc \left(\frac{c(a + \operatorname{barctanh}(cx^3))^2}{2b} - \frac{a + \operatorname{barctanh}(cx^3)}{x^3} + \frac{1}{2} bc (\log(x^6) - \log(1 - c^2x^6)) \right) - \frac{(a + \operatorname{barctanh}(cx^3))^2}{2x^6} \right)$$

input `Int[(a + b*ArcTanh[c*x^3])^2/x^7,x]`

output `(-1/2*(a + b*ArcTanh[c*x^3])^2/x^6 + b*c*(-((a + b*ArcTanh[c*x^3])/x^3) + (c*(a + b*ArcTanh[c*x^3])^2)/(2*b) + (b*c*(Log[x^6] - Log[1 - c^2*x^6]))/2))/3`

Definitions of rubi rules used

- rule 14 $\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ ; FreeQ}[a, x]$
- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$
- rule 47 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x]$
- rule 243 $\text{Int}[(x_)^{(m_)}*((a_)+(b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] \text{ ; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$
- rule 6452 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{2n})}), x], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$
- rule 6454 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n] - 1)}*(a + b*\text{ArcTanh}[c*x])^p, x], x, x^n], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$
- rule 6510 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)]*(b_))^{(p_)}((d_)+(e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^{(p+1)}/(b*c*d*(p+1)), x] \text{ ; FreeQ}[\{a, b, c, d, e, p\}, x] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{NeQ}[p, -1]$

rule 6544

```
Int[(((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^ (p_.)*((f_.)*(x_.))^ (m_.))/((d_.) + (
e_.)*(x_)^2), x_Symbol] :> Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x
], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x
^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.73

method	result
parallelrisch	$\frac{b^2 \operatorname{arctanh}(cx^3)^2 x^6 c^2 + 6b^2 c^2 \ln(x) x^6 - 2 \ln(cx^3 - 1) b^2 c^2 x^6 + 2x^6 \operatorname{arctanh}(cx^3) a b c^2 - 2 \operatorname{arctanh}(cx^3) b^2 c^2 x^6 - a^2 c^2 x^6 - 2b^2 a c^2 x^6}{6x^6}$
risch	$\frac{b^2 (c^2 x^6 - 1) \ln(cx^3 + 1)^2}{24x^6} - \frac{b (b c^2 \ln(-cx^3 + 1) x^6 + 2bcx^3 - b \ln(-cx^3 + 1) + 2a) \ln(cx^3 + 1)}{12x^6} + \frac{b^2 c^2 x^6 \ln(-cx^3 + 1)^2 + 24b^2 a c^2 x^6}{12x^6}$
default	Expression too large to display
parts	Expression too large to display

```
input int((a+b*arctanh(c*x^3))^2/x^7,x,method=_RETURNVERBOSE)
```

```
output 1/6*(b^2*arctanh(c*x^3)^2*x^6*c^2+6*b^2*c^2*ln(x)*x^6-2*ln(c*x^3-1)*b^2*c^
2*x^6+2*x^6*arctanh(c*x^3)*a*b*c^2-2*arctanh(c*x^3)*b^2*c^2*x^6-a^2*c^2*x^
6-2*b^2*arctanh(c*x^3)*x^3*c-2*a*b*c*x^3-b^2*arctanh(c*x^3)^2-2*arctanh(c*
x^3)*a*b-a^2)/x^6
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.72

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^7} dx$$

$$= \frac{24 b^2 c^2 x^6 \log(x) + 4 (ab - b^2) c^2 x^6 \log(cx^3 + 1) - 4 (ab + b^2) c^2 x^6 \log(cx^3 - 1) - 8 abcx^3 + (b^2 c^2 x^6 - b^2)}{24 x^6}$$

```
input integrate((a+b*arctanh(c*x^3))^2/x^7,x, algorithm="fricas")
```

output

```
1/24*(24*b^2*c^2*x^6*log(x) + 4*(a*b - b^2)*c^2*x^6*log(c*x^3 + 1) - 4*(a*
b + b^2)*c^2*x^6*log(c*x^3 - 1) - 8*a*b*c*x^3 + (b^2*c^2*x^6 - b^2)*log(-(
c*x^3 + 1)/(c*x^3 - 1))^2 - 4*a^2 - 4*(b^2*c*x^3 + a*b)*log(-(c*x^3 + 1)/(
c*x^3 - 1)))/x^6
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^7} dx = \text{Timed out}$$

input

```
integrate((a+b*atanh(c*x**3))**2/x**7,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 175 vs. 2(80) = 160.

Time = 0.03 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.99

$$\begin{aligned} & \int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^7} dx \\ &= \frac{1}{6} \left(\left(c \log(cx^3 + 1) - c \log(cx^3 - 1) - \frac{2}{x^3} \right) c - \frac{2 \operatorname{arctanh}(cx^3)}{x^6} \right) ab \\ & \quad + \frac{1}{24} \left(\left(2(\log(cx^3 - 1) - 2) \log(cx^3 + 1) - \log(cx^3 + 1)^2 - \log(cx^3 - 1)^2 - 4 \log(cx^3 - 1) + 24 \log \right. \right. \\ & \quad \left. \left. - \frac{b^2 \operatorname{arctanh}(cx^3)^2}{6x^6} - \frac{a^2}{6x^6} \right) \right) \end{aligned}$$

input

```
integrate((a+b*arctanh(c*x^3))^2/x^7,x, algorithm="maxima")
```

output

```
1/6*((c*log(c*x^3 + 1) - c*log(c*x^3 - 1) - 2/x^3)*c - 2*arctanh(c*x^3)/x^
6)*a*b + 1/24*((2*(log(c*x^3 - 1) - 2)*log(c*x^3 + 1) - log(c*x^3 + 1)^2 -
log(c*x^3 - 1)^2 - 4*log(c*x^3 - 1) + 24*log(x))*c^2 + 4*(c*log(c*x^3 + 1
) - c*log(c*x^3 - 1) - 2/x^3)*c*arctanh(c*x^3))*b^2 - 1/6*b^2*arctanh(c*x^
3)^2/x^6 - 1/6*a^2/x^6
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^7} dx = \int \frac{(b \operatorname{arctanh}(cx^3) + a)^2}{x^7} dx$$

input

```
integrate((a+b*arctanh(c*x^3))^2/x^7,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^3) + a)^2/x^7, x)
```

Mupad [B] (verification not implemented)

Time = 4.27 (sec) , antiderivative size = 278, normalized size of antiderivative = 3.16

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^7} dx = & \frac{b^2 c^2 \ln(cx^3 + 1)^2}{24} - \frac{b^2 c^2 \ln(cx^3 - 1)}{6} \\ & - \frac{b^2 c^2 \ln(cx^3 + 1)}{6} - \frac{a^2}{6x^6} + \frac{b^2 c^2 \ln(1 - cx^3)^2}{24} \\ & - \frac{b^2 \ln(cx^3 + 1)^2}{24x^6} - \frac{b^2 \ln(1 - cx^3)^2}{24x^6} + b^2 c^2 \ln(x) \\ & - \frac{ab c^2 \ln(cx^3 - 1)}{6} + \frac{ab c^2 \ln(cx^3 + 1)}{6} \\ & - \frac{abc}{3x^3} - \frac{ab \ln(cx^3 + 1)}{6x^6} + \frac{ab \ln(1 - cx^3)}{6x^6} \\ & - \frac{b^2 c^2 \ln(cx^3 + 1) \ln(1 - cx^3)}{12} - \frac{b^2 c \ln(cx^3 + 1)}{6x^3} \\ & + \frac{b^2 c \ln(1 - cx^3)}{6x^3} + \frac{b^2 \ln(cx^3 + 1) \ln(1 - cx^3)}{12x^6} \end{aligned}$$

input

```
int((a + b*atanh(c*x^3))^2/x^7,x)
```


$$3.123 \quad \int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^{10}} dx$$

Optimal result	1044
Mathematica [A] (verified)	1045
Rubi [A] (verified)	1045
Maple [C] (warning: unable to verify)	1048
Fricas [F]	1049
Sympy [F(-1)]	1050
Maxima [F]	1050
Giac [F]	1050
Mupad [F(-1)]	1051
Reduce [F]	1051

Optimal result

Integrand size = 16, antiderivative size = 144

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^{10}} dx = & -\frac{b^2 c^2}{9x^3} + \frac{1}{9} b^2 c^3 \operatorname{arctanh}(cx^3) - \frac{bc(a + b \operatorname{arctanh}(cx^3))}{9x^6} \\ & + \frac{1}{9} c^3 (a + b \operatorname{arctanh}(cx^3))^2 - \frac{(a + b \operatorname{arctanh}(cx^3))^2}{9x^9} \\ & + \frac{2}{9} b c^3 (a + b \operatorname{arctanh}(cx^3)) \log\left(2 - \frac{2}{1 + cx^3}\right) \\ & - \frac{1}{9} b^2 c^3 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + cx^3}\right) \end{aligned}$$

output

```
-1/9*b^2*c^2/x^3+1/9*b^2*c^3*arctanh(c*x^3)-1/9*b*c*(a+b*arctanh(c*x^3))/x^6+1/9*c^3*(a+b*arctanh(c*x^3))^2-1/9*(a+b*arctanh(c*x^3))^2/x^9+2/9*b*c^3*(a+b*arctanh(c*x^3))*ln(2-2/(c*x^3+1))-1/9*b^2*c^3*polylog(2,-1+2/(c*x^3+1))
```

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^{10}} dx = \frac{a^2 + abcx^3 + b^2c^2x^6 + b^2(1 - c^3x^9) \operatorname{arctanh}(cx^3)^2 + b \operatorname{arctanh}(cx^3) (2a + bcbx^3 - bc^3x^9 - 2bc^3x^9 \log(1 - c^3x^9))}{9x^9}$$

input

```
Integrate[(a + b*ArcTanh[c*x^3])^2/x^10,x]
```

output

```
-1/9*(a^2 + a*b*c*x^3 + b^2*c^2*x^6 + b^2*(1 - c^3*x^9)*ArcTanh[c*x^3]^2 +
b*ArcTanh[c*x^3]*(2*a + b*c*x^3 - b*c^3*x^9 - 2*b*c^3*x^9*Log[1 - E^(-2*ArcTanh[c*x^3])]) -
2*a*b*c^3*x^9*Log[c*x^3] + a*b*c^3*x^9*Log[1 - c^2*x^6] + b^2*c^3*x^9*PolyLog[2, E^(-2*ArcTanh[c*x^3])])/x^9
```

Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.95, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.562$, Rules used = {6454, 6452, 6544, 6452, 264, 219, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^{10}} dx \\ & \quad \downarrow \text{6454} \\ & \frac{1}{3} \int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^{12}} dx^3 \\ & \quad \downarrow \text{6452} \\ & \frac{1}{3} \left(\frac{2}{3} bc \int \frac{a + b \operatorname{arctanh}(cx^3)}{x^9(1 - c^2x^6)} dx^3 - \frac{(a + b \operatorname{arctanh}(cx^3))^2}{3x^9} \right) \\ & \quad \downarrow \text{6544} \end{aligned}$$

$$\frac{1}{3} \left(\frac{2}{3} bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^3)}{x^3(1-c^2x^6)} dx^3 + \int \frac{a + \operatorname{barctanh}(cx^3)}{x^9} dx^3 \right) - \frac{(a + \operatorname{barctanh}(cx^3))^2}{3x^9} \right)$$

↓ 6452

$$\frac{1}{3} \left(\frac{2}{3} bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^3)}{x^3(1-c^2x^6)} dx^3 + \frac{1}{2} bc \int \frac{1}{x^6(1-c^2x^6)} dx^3 - \frac{a + \operatorname{barctanh}(cx^3)}{2x^6} \right) - \frac{(a + \operatorname{barctanh}(cx^3))^2}{3x^9} \right)$$

↓ 264

$$\frac{1}{3} \left(\frac{2}{3} bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^3)}{x^3(1-c^2x^6)} dx^3 + \frac{1}{2} bc \left(c^2 \int \frac{1}{1-c^2x^6} dx^3 - \frac{1}{x^3} \right) - \frac{a + \operatorname{barctanh}(cx^3)}{2x^6} \right) - \frac{(a + \operatorname{barctanh}(cx^3))^2}{3x^9} \right)$$

↓ 219

$$\frac{1}{3} \left(\frac{2}{3} bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^3)}{x^3(1-c^2x^6)} dx^3 - \frac{a + \operatorname{barctanh}(cx^3)}{2x^6} + \frac{1}{2} bc \left(\operatorname{carctanh}(cx^3) - \frac{1}{x^3} \right) \right) - \frac{(a + \operatorname{barctanh}(cx^3))^2}{3x^9} \right)$$

↓ 6550

$$\frac{1}{3} \left(\frac{2}{3} bc \left(c^2 \left(\int \frac{a + \operatorname{barctanh}(cx^3)}{x^3(cx^3+1)} dx^3 + \frac{(a + \operatorname{barctanh}(cx^3))^2}{2b} \right) - \frac{a + \operatorname{barctanh}(cx^3)}{2x^6} + \frac{1}{2} bc \left(\operatorname{carctanh}(cx^3) - \frac{1}{x^3} \right) \right) \right)$$

↓ 6494

$$\frac{1}{3} \left(\frac{2}{3} bc \left(c^2 \left(-bc \int \frac{\log\left(2 - \frac{2}{cx^3+1}\right)}{1-c^2x^6} dx^3 + \frac{(a + \operatorname{barctanh}(cx^3))^2}{2b} + \log\left(2 - \frac{2}{cx^3+1}\right) (a + \operatorname{barctanh}(cx^3)) \right) \right) \right)$$

↓ 2897

$$\frac{1}{3} \left(\frac{2}{3} bc \left(c^2 \left(\frac{(a + \operatorname{barctanh}(cx^3))^2}{2b} + \log\left(2 - \frac{2}{cx^3+1}\right) (a + \operatorname{barctanh}(cx^3)) - \frac{1}{2} b \operatorname{PolyLog}\left(2, \frac{2}{cx^3+1} - 1\right) \right) \right) \right)$$

input `Int[(a + b*ArcTanh[c*x^3])^2/x^10,x]`

output

$$\frac{(-1/3*(a + b*\text{ArcTanh}[c*x^3])^2/x^9 + (2*b*c*(-1/2*(a + b*\text{ArcTanh}[c*x^3]))/x^6 + (b*c*(-x^{(-3)} + c*\text{ArcTanh}[c*x^3]))/2 + c^2*((a + b*\text{ArcTanh}[c*x^3])^2/(2*b) + (a + b*\text{ArcTanh}[c*x^3])*Log[2 - 2/(1 + c*x^3)] - (b*\text{PolyLog}[2, -1 + 2/(1 + c*x^3)]/2))/3)/3}$$

Defintions of rubi rules used

rule 219

$$\text{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[-b, 2])) \cdot \text{ArcTanh}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[a, 2])], x] \text{ ; FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}\{a/b\} \ \&\& \ (\text{GtQ}\{a, 0\} \ || \ \text{LtQ}\{b, 0\})$$

rule 264

$$\text{Int}[(c \cdot x)^m \cdot (a + (b \cdot x)^2)^p, x_Symbol] \rightarrow \text{Simp}[(c \cdot x)^{m+1} \cdot ((a + b \cdot x^2)^{p+1} / (a \cdot c \cdot (m+1))), x] - \text{Simp}[b \cdot ((m+2 \cdot p+3) / (a \cdot c^{2 \cdot (m+1)})) \cdot \text{Int}[(c \cdot x)^{m+2} \cdot (a + b \cdot x^2)^p, x], x] \text{ ; FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{LtQ}\{m, -1\} \ \&\& \ \text{IntBinomialQ}\{a, b, c, 2, m, p, x\}$$

rule 2897

$$\text{Int}[\text{Log}[u] \cdot (Pq)^m, x_Symbol] \rightarrow \text{With}\{C = \text{FullSimplify}[Pq^m \cdot ((1-u)/D[u, x])]\}, \text{Simp}[C \cdot \text{PolyLog}[2, 1-u], x] \text{ ; FreeQ}\{C, x\} \text{ ; IntegerQ}\{m\} \ \&\& \ \text{PolyQ}\{Pq, x\} \ \&\& \ \text{RationalFunctionQ}\{u, x\} \ \&\& \ \text{LeQ}\{\text{RationalFunctionExponents}[u, x][[2]], \text{Expon}[Pq, x]\}$$

rule 6452

$$\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot (b \cdot x)^p) \cdot (x)^m, x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot ((a + b \cdot \text{ArcTanh}[c \cdot x^n])^p / (m+1)), x] - \text{Simp}[b \cdot c \cdot n \cdot (p / (m+1)) \cdot \text{Int}[x^{m+n} \cdot ((a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2 \cdot n}))], x], x] \text{ ; FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}\{p, 0\} \ \&\& \ (\text{EqQ}\{p, 1\} \ || \ (\text{EqQ}\{n, 1\} \ \&\& \ \text{IntegerQ}\{m\})) \ \&\& \ \text{NeQ}\{m, -1\}$$

rule 6454

$$\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot (b \cdot x)^p) \cdot (x)^m, x_Symbol] \rightarrow \text{Simp}[1/n \cdot \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n] - 1) \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p}, x], x, x^n], x] \text{ ; FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}\{p, 1\} \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$$

rule 6494

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_))), x
_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] -
Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]]
/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c
^2*d^2 - e^2, 0]
```

rule 6544

```
Int((((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*((f_.)*(x_))^(m_))/((d_) + (
e_.)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x
], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x
^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

rule 6550

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_)^2)),
x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/
d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.24 (sec) , antiderivative size = 3062, normalized size of antiderivative = 21.26

method	result	size
default	Expression too large to display	3062
parts	Expression too large to display	3062

input

```
int((a+b*arctanh(c*x^3))^2/x^10,x,method=_RETURNVERBOSE)
```

output

```

-1/9*a^2/x^9+b^2*(-1/9/x^9*arctanh(c*x^3)^2+2/3*c*(-1/6/x^6*arctanh(c*x^3)
+arctanh(c*x^3)*c^2*ln(x)-1/6*arctanh(c*x^3)*c^2*ln(c*x^3-1)-1/6*arctanh(c
*x^3)*c^2*ln(c*x^3+1)-1/2*c*(1/6*c*ln(c*x^3-1)+1/3/x^3-1/6*c*ln(c*x^3+1)+c
^2*(Sum(1/6*(ln(x-_alpha)*ln(c*x^3-1)-3*c*(1/6/_alpha^2/c*ln(x-_alpha)^2-1
/3*_alpha*ln(x-_alpha))*(2*ln((RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1)-
x+_alpha)/RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1))*RootOf(_Z^2+3*_Z*_a
lpha+3*_alpha^2,index=1)*RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=2))+3*ln(
(RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1)-x+_alpha)/RootOf(_Z^2+3*_Z*_a
lpha+3*_alpha^2,index=1))*RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1)*_alp
ha+6*ln((RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1)-x+_alpha)/RootOf(_Z^2
+3*_Z*_alpha+3*_alpha^2,index=1))*RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index
=2)*_alpha+9*ln((RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1)-x+_alpha)/Ro
otOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1))*_alpha^2+2*ln((RootOf(_Z^2+3*_Z*
_alpha+3*_alpha^2,index=2)-x+_alpha)/RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,in
dex=2))*RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1)*RootOf(_Z^2+3*_Z*_alp
ha+3*_alpha^2,index=2)+6*ln((RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=2)-x+
_alpha)/RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=2))*RootOf(_Z^2+3*_Z*_alp
ha+3*_alpha^2,index=1)*_alpha+3*ln((RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,ind
ex=2)-x+_alpha)/RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=2))*RootOf(_Z^2+3
*_Z*_alpha+3*_alpha^2,index=2)*_alpha+9*ln((RootOf(_Z^2+3*_Z*_alpha+3*_...

```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^{10}} dx = \int \frac{(b \operatorname{arctanh}(cx^3) + a)^2}{x^{10}} dx$$

input

```
integrate((a+b*arctanh(c*x^3))^2/x^10,x, algorithm="fricas")
```

output

```
integral((b^2*arctanh(c*x^3)^2 + 2*a*b*arctanh(c*x^3) + a^2)/x^10, x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^{10}} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**3))**2/x**10,x)`output `Timed out`**Maxima [F]**

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^{10}} dx = \int \frac{(b \operatorname{arctanh}(cx^3) + a)^2}{x^{10}} dx$$

input `integrate((a+b*arctanh(c*x^3))^2/x^10,x, algorithm="maxima")`output `-1/9*((c^2*log(c^2*x^6 - 1) - c^2*log(x^6) + 1/x^6)*c + 2*arctanh(c*x^3)/x^9)*a*b - 1/36*b^2*(log(-c*x^3 + 1)^2/x^9 + 9*integrate(-1/3*(3*(c*x^3 - 1)*log(c*x^3 + 1)^2 + 2*(c*x^3 - 3*(c*x^3 - 1)*log(c*x^3 + 1))*log(-c*x^3 + 1))/(c*x^13 - x^10), x)) - 1/9*a^2/x^9`**Giac [F]**

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^{10}} dx = \int \frac{(b \operatorname{arctanh}(cx^3) + a)^2}{x^{10}} dx$$

input `integrate((a+b*arctanh(c*x^3))^2/x^10,x, algorithm="giac")`output `integrate((b*arctanh(c*x^3) + a)^2/x^10, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^{10}} dx = \int \frac{(a + b \operatorname{atanh}(cx^3))^2}{x^{10}} dx$$

input `int((a + b*atanh(c*x^3))^2/x^10,x)`output `int((a + b*atanh(c*x^3))^2/x^10, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^{10}} dx$$

$$= \frac{-\operatorname{atanh}(cx^3)^2 b^2 + 2\operatorname{atanh}(cx^3) ab c^3 x^9 - 2\operatorname{atanh}(cx^3) ab + \operatorname{atanh}(cx^3) b^2 c^3 x^9 - \operatorname{atanh}(cx^3) b^2 c x^3 - 6 \int \frac{\operatorname{atanh}(cx^3)}{c^2 x^7 - x} dx}{9x^9}$$

input `int((a+b*atanh(c*x^3))^2/x^10,x)`output `(- atanh(c*x**3)**2*b**2 + 2*atanh(c*x**3)*a*b*c**3*x**9 - 2*atanh(c*x**3)*a*b + atanh(c*x**3)*b**2*c**3*x**9 - atanh(c*x**3)*b**2*c*x**3 - 6*int(atanh(c*x**3)/(c**2*x**7 - x),x)*b**2*c**3*x**9 - 2*log(c**(2/3)*x**2 - c**(1/3)*x + 1)*a*b*c**3*x**9 - 2*log(c**(2/3)*x + c**(1/3))*a*b*c**3*x**9 + 6*log(x)*a*b*c**3*x**9 - a**2 - a*b*c*x**3 - b**2*c**2*x**6)/(9*x**9)`

3.124 $\int x^8(a + b \operatorname{arctanh}(cx^3))^3 dx$

Optimal result	1052
Mathematica [A] (verified)	1053
Rubi [A] (verified)	1053
Maple [F]	1057
Fricas [F]	1057
Sympy [F(-1)]	1058
Maxima [F]	1058
Giac [F]	1058
Mupad [F(-1)]	1059
Reduce [F]	1059

Optimal result

Integrand size = 16, antiderivative size = 231

$$\int x^8(a + b \operatorname{arctanh}(cx^3))^3 dx = \frac{ab^2x^3}{3c^2} + \frac{b^3x^3\operatorname{arctanh}(cx^3)}{3c^2} - \frac{b(a + b \operatorname{arctanh}(cx^3))^2}{6c^3} + \frac{bx^6(a + b \operatorname{arctanh}(cx^3))^2}{6c} + \frac{(a + b \operatorname{arctanh}(cx^3))^3}{9c^3} + \frac{1}{9}x^9(a + b \operatorname{arctanh}(cx^3))^3 - \frac{b(a + b \operatorname{arctanh}(cx^3))^2 \log\left(\frac{2}{1-cx^3}\right)}{3c^3} + \frac{b^3 \log(1 - c^2x^6)}{6c^3} - \frac{b^2(a + b \operatorname{arctanh}(cx^3)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^3}\right)}{3c^3} + \frac{b^3 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx^3}\right)}{6c^3}$$

output `1/3*a*b^2*x^3/c^2+1/3*b^3*x^3*arctanh(c*x^3)/c^2-1/6*b*(a+b*arctanh(c*x^3))^2/c^3+1/6*b*x^6*(a+b*arctanh(c*x^3))^2/c+1/9*(a+b*arctanh(c*x^3))^3/c^3+1/9*x^9*(a+b*arctanh(c*x^3))^3-1/3*b*(a+b*arctanh(c*x^3))^2*ln(2/(-c*x^3+1))/c^3+1/6*b^3*ln(-c^2*x^6+1)/c^3-1/3*b^2*(a+b*arctanh(c*x^3))*polylog(2,1-2/(-c*x^3+1))/c^3+1/6*b^3*polylog(3,1-2/(-c*x^3+1))/c^3`

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 334, normalized size of antiderivative = 1.45

$$\int x^8 (a + b \operatorname{arctanh}(cx^3))^3 dx$$

$$= \frac{6ab^2cx^3 + 3a^2bc^2x^6 + 2a^3c^3x^9 - 6ab^2\operatorname{arctanh}(cx^3) + 6b^3cx^3\operatorname{arctanh}(cx^3) + 6ab^2c^2x^6\operatorname{arctanh}(cx^3) + 6a^2bc^3x^9\operatorname{arctanh}(cx^3) - 6ab^2\operatorname{arctanh}(cx^3)^2 - 3b^3\operatorname{arctanh}(cx^3)^2 + 3b^3c^2x^6\operatorname{arctanh}(cx^3)^2 + 6ab^2c^3x^9\operatorname{arctanh}(cx^3)^2 - 2b^3\operatorname{arctanh}(cx^3)^3 + 2b^3c^3x^9\operatorname{arctanh}(cx^3)^3 - 12ab^2\operatorname{arctanh}(cx^3)\operatorname{Log}[1 + E^{-2\operatorname{arctanh}(cx^3)}] - 6b^3\operatorname{arctanh}(cx^3)^2\operatorname{Log}[1 + E^{-2\operatorname{arctanh}(cx^3)}] + 3a^2b\operatorname{Log}[1 - c^2x^6] + 3b^3\operatorname{Log}[1 - c^2x^6] + 6b^2(a + b\operatorname{arctanh}(cx^3))\operatorname{PolyLog}[2, -E^{-2\operatorname{arctanh}(cx^3)}] + 3b^3\operatorname{PolyLog}[3, -E^{-2\operatorname{arctanh}(cx^3)}]}{18c^3}$$

input

```
Integrate[x^8*(a + b*ArcTanh[c*x^3])^3,x]
```

output

```
(6*a*b^2*c*x^3 + 3*a^2*b*c^2*x^6 + 2*a^3*c^3*x^9 - 6*a*b^2*ArcTanh[c*x^3]
+ 6*b^3*c*x^3*ArcTanh[c*x^3] + 6*a*b^2*c^2*x^6*ArcTanh[c*x^3] + 6*a^2*b*c^
3*x^9*ArcTanh[c*x^3] - 6*a*b^2*ArcTanh[c*x^3]^2 - 3*b^3*ArcTanh[c*x^3]^2 +
3*b^3*c^2*x^6*ArcTanh[c*x^3]^2 + 6*a*b^2*c^3*x^9*ArcTanh[c*x^3]^2 - 2*b^3
*ArcTanh[c*x^3]^3 + 2*b^3*c^3*x^9*ArcTanh[c*x^3]^3 - 12*a*b^2*ArcTanh[c*x^
3]*Log[1 + E^(-2*ArcTanh[c*x^3])] - 6*b^3*ArcTanh[c*x^3]^2*Log[1 + E^(-2*A
rcTanh[c*x^3])] + 3*a^2*b*Log[1 - c^2*x^6] + 3*b^3*Log[1 - c^2*x^6] + 6*b^
2*(a + b*ArcTanh[c*x^3])*PolyLog[2, -E^(-2*ArcTanh[c*x^3])] + 3*b^3*PolyLo
g[3, -E^(-2*ArcTanh[c*x^3])])/(18*c^3)
```

Rubi [A] (verified)

Time = 2.24 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.688$, Rules used = {6454, 6452, 6542, 6452, 6542, 2009, 6510, 6546, 6470, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^8 (a + b \operatorname{arctanh}(cx^3))^3 dx$$

$$\downarrow 6454$$

$$\frac{1}{3} \int x^6 (a + b \operatorname{arctanh}(cx^3))^3 dx^3$$

$$\downarrow 6452$$

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))^3 - bc \int \frac{x^9 (a + \operatorname{barctanh}(cx^3))^2}{1 - c^2 x^6} dx^3 \right)$$

↓ 6542

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))^3 - bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(cx^3))^2}{1 - c^2 x^6} dx^3}{c^2} - \frac{\int x^3 (a + \operatorname{barctanh}(cx^3))^2 dx^3}{c^2} \right) \right)$$

↓ 6452

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))^3 - bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(cx^3))^2}{1 - c^2 x^6} dx^3}{c^2} - \frac{\frac{1}{2} x^6 (a + \operatorname{barctanh}(cx^3))^2 - bc \int \frac{x^6 (a + \operatorname{barctanh}(cx^3))}{1 - c^2 x^6}}{c^2} \right) \right)$$

↓ 6542

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))^3 - bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(cx^3))^2}{1 - c^2 x^6} dx^3}{c^2} - \frac{\frac{1}{2} x^6 (a + \operatorname{barctanh}(cx^3))^2 - bc \left(\frac{\int \frac{a + \operatorname{barctanh}(cx^3)}{1 - c^2 x^6}}{c^2} \right)}{c^2} \right) \right)$$

↓ 2009

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))^3 - bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(cx^3))^2}{1 - c^2 x^6} dx^3}{c^2} - \frac{\frac{1}{2} x^6 (a + \operatorname{barctanh}(cx^3))^2 - bc \left(\frac{\int \frac{a + \operatorname{barctanh}(cx^3)}{1 - c^2 x^6}}{c^2} \right)}{c^2} \right) \right)$$

↓ 6510

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{barctanh}(cx^3))^3 - bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(cx^3))^2}{1 - c^2 x^6} dx^3}{c^2} - \frac{\frac{1}{2} x^6 (a + \operatorname{barctanh}(cx^3))^2 - bc \left(\frac{(a + \operatorname{barctanh}(cx^3))}{2bc^3} \right)}{c^2} \right) \right)$$

↓ 6546

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{arctanh}(cx^3))^3 - bc \left(\frac{\int \frac{(a + \operatorname{arctanh}(cx^3))^2}{1 - cx^3} dx^3}{c} - \frac{(a + \operatorname{arctanh}(cx^3))^3}{3bc^2} - \frac{\frac{1}{2} x^6 (a + \operatorname{arctanh}(cx^3))^2}{c^2} \right) \right)$$

↓ 6470

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{arctanh}(cx^3))^3 - bc \left(\frac{\frac{\log\left(\frac{2}{1 - cx^3}\right) (a + \operatorname{arctanh}(cx^3))^2}{c} - 2b \int \frac{(a + \operatorname{arctanh}(cx^3)) \log\left(\frac{2}{1 - cx^3}\right)}{1 - c^2 x^6} dx^3}{c} - \frac{(a + \operatorname{arctanh}(cx^3))^3}{3bc^2} \right) \right)$$

↓ 6620

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{arctanh}(cx^3))^3 - bc \left(\frac{\frac{\log\left(\frac{2}{1 - cx^3}\right) (a + \operatorname{arctanh}(cx^3))^2}{c} - 2b \left(\frac{\frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx^3}\right)}{1 - c^2 x^6} dx^3}{c} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx^3}\right) (a + \operatorname{arctanh}(cx^3))^2}{2c} \right)}{c} \right) \right)$$

↓ 7164

$$\frac{1}{3} \left(\frac{1}{3} x^9 (a + \operatorname{arctanh}(cx^3))^3 - bc \left(\frac{\frac{\log\left(\frac{2}{1 - cx^3}\right) (a + \operatorname{arctanh}(cx^3))^2}{c} - 2b \left(\frac{\frac{b \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - cx^3}\right)}{4c} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx^3}\right) (a + \operatorname{arctanh}(cx^3))^2}{2c} \right)}{c} \right) \right)$$

input `Int [x^8*(a + b*ArcTanh[c*x^3])^3,x]`

output `((x^9*(a + b*ArcTanh[c*x^3])^3)/3 - b*c*(-(((x^6*(a + b*ArcTanh[c*x^3])^2)/2 - b*c*((a + b*ArcTanh[c*x^3])^2/(2*b*c^3) - (a*x^3 + b*x^3*ArcTanh[c*x^3] + (b*Log[1 - c^2*x^6])/(2*c))/c^2))/c^2) + (-1/3*(a + b*ArcTanh[c*x^3])^3/(b*c^2) + ((a + b*ArcTanh[c*x^3])^2*Log[2/(1 - c*x^3)]/c - 2*b*(-1/2*(a + b*ArcTanh[c*x^3])*PolyLog[2, 1 - 2/(1 - c*x^3)]/c + (b*PolyLog[3, 1 - 2/(1 - c*x^3)]/(4*c))/c)/c^2))/3`

Defintions of rubi rules used

rule 2009 $\text{Int}[u_, x_Symbol] \text{ :> Simp}[\text{IntSum}[u, x], x] \text{ /; SumQ}[u]$

rule 6452 $\text{Int}[\{(a_.) + \text{ArcTanh}[(c_.)(x_)^{(n_.)}] * (b_.)\}^{(p_.)} * (x_)^{(m_.)}, x_Symbol] \text{ :> Simp}[x^{(m+1)} * \{(a + b * \text{ArcTanh}[c * x^n]\}^{p/(m+1)}, x] - \text{Simp}[b * c * n * \{(p/(m+1)) \text{ Int}[x^{(m+n)} * \{(a + b * \text{ArcTanh}[c * x^n]\}^{(p-1)/(1-c^2 * x^{2*n})}\}, x], x] \text{ /; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 6454 $\text{Int}[\{(a_.) + \text{ArcTanh}[(c_.)(x_)^{(n_.)}] * (b_.)\}^{(p_.)} * (x_)^{(m_.)}, x_Symbol] \text{ :> Simp}[1/n \ \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n] - 1) * (a + b * \text{ArcTanh}[c * x])^p}, x], x, x^n], x] \text{ /; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$

rule 6470 $\text{Int}[\{(a_.) + \text{ArcTanh}[(c_.)(x_) * (b_.)]\}^{(p_.)} / \{(d_.) + (e_.)(x_)\}, x_Symbol] \text{ :> Simp}[-(a + b * \text{ArcTanh}[c * x])^p * (\text{Log}[2/(1 + e * (x/d))]/e), x] + \text{Simp}[b * c * \{(p/e) \ \text{Int}[(a + b * \text{ArcTanh}[c * x])^{(p-1)} * (\text{Log}[2/(1 + e * (x/d))]/(1 - c^2 * x^2))\}, x], x] \text{ /; FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 * d^2 - e^2, 0]$

rule 6510 $\text{Int}[\{(a_.) + \text{ArcTanh}[(c_.)(x_) * (b_.)]\}^{(p_.)} / \{(d_.) + (e_.)(x_)^2\}, x_Symbol] \text{ :> Simp}[(a + b * \text{ArcTanh}[c * x])^{(p+1)} / (b * c * d * (p+1)), x] \text{ /; FreeQ}[\{a, b, c, d, e, p\}, x] \ \&\& \ \text{EqQ}[c^2 * d + e, 0] \ \&\& \ \text{NeQ}[p, -1]$

rule 6542 $\text{Int}[\{((a_.) + \text{ArcTanh}[(c_.)(x_) * (b_.)]\}^{(p_.)} * \{(f_.)(x_)^{(m_.)}\} / \{(d_.) + (e_.)(x_)^2\}, x_Symbol] \text{ :> Simp}[f^2/e \ \text{Int}[(f * x)^{(m-2)} * (a + b * \text{ArcTanh}[c * x])^p, x], x] - \text{Simp}[d * (f^2/e) \ \text{Int}[(f * x)^{(m-2)} * \{(a + b * \text{ArcTanh}[c * x])^p / (d + e * x^2)\}, x], x] \text{ /; FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{GtQ}[m, 1]$

rule 6546

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*(x_)/((d_) + (e_.)*(x_)^2),
 x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/
 (c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

rule 6620

```
Int[(Log[u_]*)((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)^
 2), x_Symbol] := Simp[(-(a + b*ArcTanh[c*x])^p)*(PolyLog[2, 1 - u]/(2*c*d))
 , x] + Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(
 d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d
 + e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
 x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [F]

$$\int x^8 (a + b \operatorname{arctanh}(cx^3))^3 dx$$

input

```
int(x^8*(a+b*arctanh(c*x^3))^3,x)
```

output

```
int(x^8*(a+b*arctanh(c*x^3))^3,x)
```

Fricas [F]

$$\int x^8 (a + b \operatorname{arctanh}(cx^3))^3 dx = \int (b \operatorname{arctanh}(cx^3) + a)^3 x^8 dx$$

input

```
integrate(x^8*(a+b*arctanh(c*x^3))^3,x, algorithm="fricas")
```

output

```
integral(b^3*x^8*arctanh(c*x^3)^3 + 3*a*b^2*x^8*arctanh(c*x^3)^2 + 3*a^2*b
*x^8*arctanh(c*x^3) + a^3*x^8, x)
```

Sympy [F(-1)]

Timed out.

$$\int x^8(a + \operatorname{arctanh}(cx^3))^3 dx = \text{Timed out}$$

input `integrate(x**8*(a+b*atanh(c*x**3))**3,x)`

output `Timed out`

Maxima [F]

$$\int x^8(a + \operatorname{arctanh}(cx^3))^3 dx = \int (b \operatorname{arctanh}(cx^3) + a)^3 x^8 dx$$

input `integrate(x^8*(a+b*arctanh(c*x^3))^3,x, algorithm="maxima")`

output `1/9*a^3*x^9 + 1/6*(2*x^9*arctanh(c*x^3) + (x^6/c^2 + log(c^2*x^6 - 1)/c^4)*c)*a^2*b - 1/72*((b^3*c^3*x^9 - b^3)*log(-c*x^3 + 1)^3 - 3*(2*a*b^2*c^3*x^9 + b^3*c^2*x^6 + (b^3*c^3*x^9 + b^3)*log(c*x^3 + 1))*log(-c*x^3 + 1)^2)/c^3 - integrate(-1/8*((b^3*c^3*x^11 - b^3*c^2*x^8)*log(c*x^3 + 1)^3 + 6*(a*b^2*c^3*x^11 - a*b^2*c^2*x^8)*log(c*x^3 + 1)^2 - (4*a*b^2*c^3*x^11 + 2*b^3*c^2*x^8 + 3*(b^3*c^3*x^11 - b^3*c^2*x^8)*log(c*x^3 + 1)^2 - 2*(6*a*b^2*c^2*x^8 - (6*a*b^2*c^3 + b^3*c^3)*x^11 - b^3*x^2)*log(c*x^3 + 1))*log(-c*x^3 + 1))/(c^3*x^3 - c^2), x)`

Giac [F]

$$\int x^8(a + \operatorname{arctanh}(cx^3))^3 dx = \int (b \operatorname{arctanh}(cx^3) + a)^3 x^8 dx$$

input `integrate(x^8*(a+b*arctanh(c*x^3))^3,x, algorithm="giac")`

3.125 $\int x^5(a + \operatorname{barctanh}(cx^3))^3 dx$

Optimal result	1060
Mathematica [A] (verified)	1061
Rubi [A] (verified)	1061
Maple [C] (warning: unable to verify)	1065
Fricas [F]	1066
Sympy [F(-1)]	1066
Maxima [F]	1066
Giac [F]	1067
Mupad [F(-1)]	1067
Reduce [F]	1068

Optimal result

Integrand size = 16, antiderivative size = 139

$$\int x^5(a + \operatorname{barctanh}(cx^3))^3 dx = \frac{b(a + \operatorname{barctanh}(cx^3))^2}{2c^2} + \frac{bx^3(a + \operatorname{barctanh}(cx^3))^2}{2c} - \frac{(a + \operatorname{barctanh}(cx^3))^3}{6c^2} + \frac{1}{6}x^6(a + \operatorname{barctanh}(cx^3))^3 - \frac{b^2(a + \operatorname{barctanh}(cx^3)) \log\left(\frac{2}{1-cx^3}\right)}{c^2} - \frac{b^3 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^3}\right)}{2c^2}$$

output

```
1/2*b*(a+b*arctanh(c*x^3))^2/c^2+1/2*b*x^3*(a+b*arctanh(c*x^3))^2/c-1/6*(a+b*arctanh(c*x^3))^3/c^2+1/6*x^6*(a+b*arctanh(c*x^3))^3-b^2*(a+b*arctanh(c*x^3))*ln(2/(-c*x^3+1))/c^2-1/2*b^3*polylog(2,1-2/(-c*x^3+1))/c^2
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.33

$$\int x^5 (a + \operatorname{barctanh}(cx^3))^3 dx$$

$$= \frac{6b^2(-1 + cx^3)(a + b + acx^3) \operatorname{arctanh}(cx^3)^2 + 2b^3(-1 + c^2x^6) \operatorname{arctanh}(cx^3)^3 + 6\operatorname{barctanh}(cx^3) \left(acx^3(2b \right.$$

input

```
Integrate[x^5*(a + b*ArcTanh[c*x^3])^3,x]
```

output

```
(6*b^2*(-1 + c*x^3)*(a + b + a*c*x^3)*ArcTanh[c*x^3]^2 + 2*b^3*(-1 + c^2*x^6)*ArcTanh[c*x^3]^3 + 6*b*ArcTanh[c*x^3]*(a*c*x^3*(2*b + a*c*x^3) - 2*b^2*Log[1 + E^(-2*ArcTanh[c*x^3])]) + a*(6*a*b*c*x^3 + 2*a^2*c^2*x^6 + 3*a*b*Log[1 - c*x^3] - 3*a*b*Log[1 + c*x^3] + 6*b^2*Log[1 - c^2*x^6]) + 6*b^3*PolyLog[2, -E^(-2*ArcTanh[c*x^3])])/(12*c^2)
```

Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.14, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.562$, Rules used = {6454, 6452, 6542, 6436, 6510, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^5 (a + \operatorname{barctanh}(cx^3))^3 dx$$

$$\downarrow 6454$$

$$\frac{1}{3} \int x^3 (a + \operatorname{barctanh}(cx^3))^3 dx^3$$

$$\downarrow 6452$$

$$\frac{1}{3} \left(\frac{1}{2} x^6 (a + \operatorname{barctanh}(cx^3))^3 - \frac{3}{2} bc \int \frac{x^6 (a + \operatorname{barctanh}(cx^3))^2}{1 - c^2 x^6} dx^3 \right)$$

$$\downarrow 6542$$

$$\frac{1}{3} \left(\frac{1}{2} x^6 (a + \operatorname{barctanh}(cx^3))^3 - \frac{3}{2} bc \left(\frac{\int \frac{(a + \operatorname{barctanh}(cx^3))^2}{1 - c^2 x^6} dx^3}{c^2} - \frac{\int (a + \operatorname{barctanh}(cx^3))^2 dx^3}{c^2} \right) \right)$$

↓ 6436

$$\frac{1}{3} \left(\frac{1}{2} x^6 (a + \operatorname{barctanh}(cx^3))^3 - \frac{3}{2} bc \left(\frac{\int \frac{(a + \operatorname{barctanh}(cx^3))^2}{1 - c^2 x^6} dx^3}{c^2} - \frac{x^3 (a + \operatorname{barctanh}(cx^3))^2 - 2bc \int \frac{x^3 (a + \operatorname{barctanh}(cx^3))}{1 - c^2 x^6}}{c^2} \right) \right)$$

↓ 6510

$$\frac{1}{3} \left(\frac{1}{2} x^6 (a + \operatorname{barctanh}(cx^3))^3 - \frac{3}{2} bc \left(\frac{(a + \operatorname{barctanh}(cx^3))^3}{3bc^3} - \frac{x^3 (a + \operatorname{barctanh}(cx^3))^2 - 2bc \int \frac{x^3 (a + \operatorname{barctanh}(cx^3))}{1 - c^2 x^6}}{c^2} \right) \right)$$

↓ 6546

$$\frac{1}{3} \left(\frac{1}{2} x^6 (a + \operatorname{barctanh}(cx^3))^3 - \frac{3}{2} bc \left(\frac{(a + \operatorname{barctanh}(cx^3))^3}{3bc^3} - \frac{x^3 (a + \operatorname{barctanh}(cx^3))^2 - 2bc \left(\frac{\int \frac{a + \operatorname{barctanh}(cx^3)}{1 - cx^3}}{c} \right)}{c^2} \right) \right)$$

↓ 6470

$$\frac{1}{3} \left(\frac{1}{2} x^6 (a + \operatorname{barctanh}(cx^3))^3 - \frac{3}{2} bc \left(\frac{(a + \operatorname{barctanh}(cx^3))^3}{3bc^3} - \frac{x^3 (a + \operatorname{barctanh}(cx^3))^2 - 2bc \left(\frac{\log\left(\frac{2}{1 - cx^3}\right)(a + \operatorname{barctanh}(cx^3))}{c} \right)}{c^2} \right) \right)$$

↓ 2849

$$\frac{1}{3} \left(\frac{1}{2} x^6 (a + \operatorname{arctanh}(cx^3))^3 - \frac{3}{2} bc \left(\frac{(a + \operatorname{arctanh}(cx^3))^3}{3bc^3} - \frac{x^3 (a + \operatorname{arctanh}(cx^3))^2 - 2bc \left(\frac{b \int \frac{\log\left(\frac{2}{1-cx^3}\right) d \frac{1}{1-cx^3}}{\frac{1-2}{1-cx^3}}}{c}}{c} \right)}{3bc^3} \right) \right)$$

↓ 2752

$$\frac{1}{3} \left(\frac{1}{2} x^6 (a + \operatorname{arctanh}(cx^3))^3 - \frac{3}{2} bc \left(\frac{(a + \operatorname{arctanh}(cx^3))^3}{3bc^3} - \frac{x^3 (a + \operatorname{arctanh}(cx^3))^2 - 2bc \left(\frac{\log\left(\frac{2}{1-cx^3}\right) (a + \operatorname{arctanh}(cx^3))}{c} \right)}{3bc^3} \right) \right)$$

input `Int[x^5*(a + b*ArcTanh[c*x^3])^3,x]`

output `((x^6*(a + b*ArcTanh[c*x^3])^3)/2 - (3*b*c*((a + b*ArcTanh[c*x^3])^3/(3*b*c^3) - (x^3*(a + b*ArcTanh[c*x^3])^2 - 2*b*c*(-1/2*(a + b*ArcTanh[c*x^3])^2/(b*c^2) + (((a + b*ArcTanh[c*x^3])*Log[2/(1 - c*x^3)])/c + (b*PolyLog[2, 1 - 2/(1 - c*x^3)])/(2*c))/c))/c^2))/2)/3`

Defintions of rubi rules used

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] :> Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] :> Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 6436 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot b)^p, x_Symbol] \rightarrow \text{Simp}[x \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^p, x] - \text{Simp}[b \cdot c \cdot n \cdot p \cdot \text{Int}[x^n \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2n})], x], x] /;$ $\text{FreeQ}\{a, b, c, n, x\} \ \&\& \ \text{IGtQ}\{p, 0\} \ \&\& \ (\text{EqQ}\{n, 1\} \ || \ \text{EqQ}\{p, 1\})$

rule 6452 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot b)^p \cdot x^m, x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^p / (m+1), x] - \text{Simp}[b \cdot c \cdot n \cdot (p/(m+1)) \cdot \text{Int}[x^{m+n} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2n})], x], x] /;$ $\text{FreeQ}\{a, b, c, m, n, x\} \ \&\& \ \text{IGtQ}\{p, 0\} \ \&\& \ (\text{EqQ}\{p, 1\} \ || \ (\text{EqQ}\{n, 1\} \ \&\& \ \text{IntegerQ}\{m\})) \ \&\& \ \text{NeQ}\{m, -1\}$

rule 6454 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot b)^p \cdot x^m, x_Symbol] \rightarrow \text{Simp}[1/n \cdot \text{Subst}[\text{Int}[x^{\text{Simplify}[(m+1)/n} - 1] \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x, x^n], x] /;$ $\text{FreeQ}\{a, b, c, m, n, x\} \ \&\& \ \text{IGtQ}\{p, 1\} \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$

rule 6470 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((d) + (e) \cdot x), x_Symbol] \rightarrow \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{Log}[2/(1 + e \cdot (x/d))]/e), x] + \text{Simp}[b \cdot c \cdot (p/e) \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{Log}[2/(1 + e \cdot (x/d))]/(1 - c^2 \cdot x^2)), x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}\{p, 0\} \ \&\& \ \text{EqQ}[c^2 \cdot d^2 - e^2, 0]$

rule 6510 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((d) + (e) \cdot x^2), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot c \cdot d \cdot (p+1)), x] /;$ $\text{FreeQ}\{a, b, c, d, e, p, x\} \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{NeQ}\{p, -1\}$

rule 6542 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p \cdot (f \cdot x)^m / ((d) + (e) \cdot x^2), x_Symbol] \rightarrow \text{Simp}[f^2/e \cdot \text{Int}[(f \cdot x)^{m-2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x] - \text{Simp}[d \cdot (f^2/e) \cdot \text{Int}[(f \cdot x)^{m-2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2)], x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, x\} \ \&\& \ \text{GtQ}\{p, 0\} \ \&\& \ \text{GtQ}\{m, 1\}$

rule 6546

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)*(x_.)/((d_.) + (e_.)*(x_)^2),
 x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/
 (c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.61 (sec) , antiderivative size = 800, normalized size of antiderivative = 5.76

method	result	size
risch	Expression too large to display	800

input

```
int(x^5*(a+b*arctanh(c*x^3))^3,x,method=_RETURNVERBOSE)
```

output

```
1/48*b^3*(c^2*x^6-1)/c^2*ln(c*x^3+1)^3+1/16*b^2*(-b*c^2*ln(-c*x^3+1)*x^6+2
*a*c^2*x^6+2*b*c*x^3+b*ln(-c*x^3+1)-2*a+2*b)/c^2*ln(c*x^3+1)^2+(1/16*b^3*(
c^2*x^6-1)/c^2*ln(-c*x^3+1)^2-1/16*b^2*(2*a*c*x^3+b)^2/c^2/a*ln(-c*x^3+1)-
1/16*b*(-4*a^3*c^2*x^6-8*a^2*b*c*x^3-4*ln(-c*x^3+1)*a^2*b-4*ln(-c*x^3+1)*a
*b^2-ln(-c*x^3+1)*b^3-4*a*b^2)/a/c^2)*ln(c*x^3+1)-1/48*b^3*x^6*ln(-c*x^3+1
)^3-1/8*b^3/c^2*ln(-c*x^3+1)^2+1/48*b^3/c^2*ln(-c*x^3+1)^3+1/4/c^2*b^3*ln(
-c*x^3+1)+1/6*a^3*x^6-1/4/c^2*b^3*ln(c*x^3+1)-1/4/c^2*b^3*ln(c*x^3-1)+1/8*
b^3/c*x^3*ln(-c*x^3+1)^2-1/4*a^2*b*x^6*ln(-c*x^3+1)+1/4*a^2*b/c^2*ln(c*x^3
-1)+1/8*a*b^2*x^6*ln(-c*x^3+1)^2+3/8/c^2*a*b^2*ln(-c*x^3+1)-1/8/c^2*a*b^2*
ln(-c*x^3+1)^2-1/2*a*b^2/c*x^3*ln(-c*x^3+1)-1/4*b/c^2*ln(c*x^3+1)*a^2+1/2*
b^2/c^2*ln(c*x^3+1)*a+1/8/c^2*b^2*a*ln(c*x^3-1)+3/4/c*b^2*Sum(-2/3*(ln(x-
_alpha)*ln(-c*x^3+1)+3*c*(-1/3*ln(x-_alpha)*(ln(1/2*(x+_alpha)/_alpha)+ln((
RootOf(_Z^2+_Z*_alpha+_alpha^2,index=1)-x+_alpha)/RootOf(_Z^2+_Z*_alpha+_a
lpha^2,index=1))+ln((RootOf(_Z^2+_Z*_alpha+_alpha^2,index=2)-x+_alpha)/Roo
tOf(_Z^2+_Z*_alpha+_alpha^2,index=2)))/c-1/3*(dilog(1/2*(x+_alpha)/_alpha)
+dilog((RootOf(_Z^2+_Z*_alpha+_alpha^2,index=1)-x+_alpha)/RootOf(_Z^2+_Z*_
alpha+_alpha^2,index=1))+dilog((RootOf(_Z^2+_Z*_alpha+_alpha^2,index=2)-x+
_alpha)/RootOf(_Z^2+_Z*_alpha+_alpha^2,index=2)))/c))*b/c,_alpha=RootOf(_Z
^3*c+1))-1/8*b^3/c^2+1/2/c*a^2*b*x^3
```

Fricas [F]

$$\int x^5 (a + b \operatorname{arctanh}(cx^3))^3 dx = \int (b \operatorname{artanh}(cx^3) + a)^3 x^5 dx$$

input `integrate(x^5*(a+b*arctanh(c*x^3))^3,x, algorithm="fricas")`

output `integral(b^3*x^5*arctanh(c*x^3)^3 + 3*a*b^2*x^5*arctanh(c*x^3)^2 + 3*a^2*b*x^5*arctanh(c*x^3) + a^3*x^5, x)`

Sympy [F(-1)]

Timed out.

$$\int x^5 (a + b \operatorname{arctanh}(cx^3))^3 dx = \text{Timed out}$$

input `integrate(x**5*(a+b*atanh(c*x**3))**3,x)`

output `Timed out`

Maxima [F]

$$\int x^5 (a + b \operatorname{arctanh}(cx^3))^3 dx = \int (b \operatorname{artanh}(cx^3) + a)^3 x^5 dx$$

input `integrate(x^5*(a+b*arctanh(c*x^3))^3,x, algorithm="maxima")`

output

```

1/2*a*b^2*x^6*arctanh(c*x^3)^2 + 1/6*a^3*x^6 + 1/4*(2*x^6*arctanh(c*x^3) +
c*(2*x^3/c^2 - log(c*x^3 + 1)/c^3 + log(c*x^3 - 1)/c^3))*a^2*b + 1/8*(4*c
*(2*x^3/c^2 - log(c*x^3 + 1)/c^3 + log(c*x^3 - 1)/c^3)*arctanh(c*x^3) - (2
*(log(c*x^3 - 1) - 2)*log(c*x^3 + 1) - log(c*x^3 + 1)^2 - log(c*x^3 - 1)^2
- 4*log(c*x^3 - 1))/c^2)*a*b^2 - 1/192*(4*x^6*log(-c*x^3 + 1)^3 + 3*(x^6/
c^3 + log(c^2*x^6 - 1)/c^5)*c^3 - 6*c*((c*x^6 + 2*x^3)/c^2 + 2*log(c*x^3 -
1)/c^3)*log(-c*x^3 + 1)^2 + 21*c^2*(2*x^3/c^3 - log(c*x^3 + 1)/c^4 + log(
c*x^3 - 1)/c^4) + c*(6*(c^2*x^6 + 6*c*x^3 + 2*log(c*x^3 - 1)^2 + 6*log(c*x
^3 - 1))*log(-c*x^3 + 1)/c^3 - (3*c^2*x^6 + 42*c*x^3 + 4*log(c*x^3 - 1)^3
+ 18*log(c*x^3 - 1)^2 + 42*log(c*x^3 - 1))/c^3) - 1728*c*integrate(1/4*x^5
*log(c*x^3 + 1)/(c^3*x^6 - c), x) - 2*(12*c*x^3*log(c*x^3 + 1)^2 + 2*(c^2*
x^6 - 1)*log(c*x^3 + 1)^3 - 3*(c^2*x^6 - 2*c*x^3 - 2*(c^2*x^6 - 1)*log(c*x
^3 + 1) + 1)*log(-c*x^3 + 1)^2 + 3*(c^2*x^6 + 6*c*x^3 - 2*(c^2*x^6 - 1)*lo
g(c*x^3 + 1)^2 - 8*(c*x^3 + 1)*log(c*x^3 + 1))*log(-c*x^3 + 1))/c^2 + 18*1
og(4*c^3*x^6 - 4*c)/c^2 - 576*integrate(1/4*x^2*log(c*x^3 + 1)/(c^3*x^6 -
c), x))*b^3

```

Giac [F]

$$\int x^5 (a + b \operatorname{arctanh}(cx^3))^3 dx = \int (b \operatorname{artanh}(cx^3) + a)^3 x^5 dx$$

input

```
integrate(x^5*(a+b*arctanh(c*x^3))^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^3) + a)^3*x^5, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^5 (a + b \operatorname{arctanh}(cx^3))^3 dx = \int x^5 (a + b \operatorname{atanh}(cx^3))^3 dx$$

input

```
int(x^5*(a + b*atanh(c*x^3))^3,x)
```


output `int(x^5*(a + b*atanh(c*x^3))^3, x)`

Reduce [F]

$$\int x^5 (a + b \operatorname{arctanh}(cx^3))^3 dx$$

$$= \frac{\operatorname{atanh}(cx^3)^3 b^3 c^2 x^6 - \operatorname{atanh}(cx^3)^3 b^3 + 3 \operatorname{atanh}(cx^3)^2 a b^2 c^2 x^6 - 3 \operatorname{atanh}(cx^3)^2 a b^2 + 3 \operatorname{atanh}(cx^3)^2 b^3 c x^3}{6 c^2}$$

input `int(x^5*(a+b*atanh(c*x^3))^3,x)`

output `(atanh(c*x**3)**3*b**3*c**2*x**6 - atanh(c*x**3)**3*b**3 + 3*atanh(c*x**3)**2*a*b**2*c**2*x**6 - 3*atanh(c*x**3)**2*a*b**2 + 3*atanh(c*x**3)**2*b**3*c*x**3 + 3*atanh(c*x**3)*a**2*b*c**2*x**6 - 3*atanh(c*x**3)*a**2*b + 6*atanh(c*x**3)*a*b**2*c*x**3 - 6*atanh(c*x**3)*a*b**2 + 18*int((atanh(c*x**3)*x**5)/(c**2*x**6 - 1),x)*b**3*c**2 + 6*log(c**(2/3)*x**2 - c**(1/3)*x + 1)*a*b**2 + 6*log(c**(2/3)*x + c**(1/3))*a*b**2 + a**3*c**2*x**6 + 3*a**2*b*c*x**3)/(6*c**2)`

3.126 $\int x^2(a + \operatorname{barctanh}(cx^3))^3 dx$

Optimal result	1069
Mathematica [A] (verified)	1070
Rubi [A] (verified)	1070
Maple [B] (verified)	1073
Fricas [F]	1073
Sympy [F(-1)]	1074
Maxima [F]	1074
Giac [F]	1075
Mupad [F(-1)]	1075
Reduce [F]	1075

Optimal result

Integrand size = 16, antiderivative size = 130

$$\int x^2(a + \operatorname{barctanh}(cx^3))^3 dx = \frac{(a + \operatorname{barctanh}(cx^3))^3}{3c} + \frac{1}{3}x^3(a + \operatorname{barctanh}(cx^3))^3 - \frac{b(a + \operatorname{barctanh}(cx^3))^2 \log\left(\frac{2}{1-cx^3}\right)}{c} - \frac{b^2(a + \operatorname{barctanh}(cx^3)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^3}\right)}{c} + \frac{b^3 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx^3}\right)}{2c}$$

output

```
1/3*(a+b*arctanh(c*x^3))^3/c+1/3*x^3*(a+b*arctanh(c*x^3))^3-b*(a+b*arctanh(c*x^3))^2*ln(2/(-c*x^3+1))/c-b^2*(a+b*arctanh(c*x^3))*polylog(2,1-2/(-c*x^3+1))/c+1/2*b^3*polylog(3,1-2/(-c*x^3+1))/c
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.47

$$\int x^2 (a + b \operatorname{arctanh}(cx^3))^3 dx$$

$$= \frac{2a^3 cx^3 + 6a^2 bcx^3 \operatorname{arctanh}(cx^3) + 3a^2 b \log(1 - c^2 x^6) + 6ab^2 \left(\operatorname{arctanh}(cx^3) \left((-1 + cx^3) \operatorname{arctanh}(cx^3) - 2 \log(1 + E^{-2 \operatorname{arctanh}(cx^3)}) \right) \right) + 6b^3 \operatorname{PolyLog}[2, -E^{-2 \operatorname{arctanh}(cx^3)}]}{6c}$$

input

```
Integrate[x^2*(a + b*ArcTanh[c*x^3])^3,x]
```

output

```
(2*a^3*c*x^3 + 6*a^2*b*c*x^3*ArcTanh[c*x^3] + 3*a^2*b*Log[1 - c^2*x^6] + 6*a*b^2*(ArcTanh[c*x^3]*((-1 + c*x^3)*ArcTanh[c*x^3] - 2*Log[1 + E^(-2*ArcTanh[c*x^3])])) + PolyLog[2, -E^(-2*ArcTanh[c*x^3])]) + b^3*(2*ArcTanh[c*x^3]^2*((-1 + c*x^3)*ArcTanh[c*x^3] - 3*Log[1 + E^(-2*ArcTanh[c*x^3])]) + 6*ArcTanh[c*x^3]*PolyLog[2, -E^(-2*ArcTanh[c*x^3])]) + 3*PolyLog[3, -E^(-2*ArcTanh[c*x^3])])/(6*c)
```

Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.10, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6454, 6436, 6546, 6470, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 (a + b \operatorname{arctanh}(cx^3))^3 dx$$

$$\downarrow 6454$$

$$\frac{1}{3} \int (a + b \operatorname{arctanh}(cx^3))^3 dx^3$$

$$\downarrow 6436$$

$$\frac{1}{3} \left(x^3 (a + b \operatorname{arctanh}(cx^3))^3 - 3bc \int \frac{x^3 (a + b \operatorname{arctanh}(cx^3))^2}{1 - c^2 x^6} dx^3 \right)$$

$$\frac{1}{3} \left(x^3 (a + \operatorname{barctanh}(cx^3))^3 - 3bc \left(\int \frac{(a + \operatorname{barctanh}(cx^3))^2}{1 - cx^3} dx^3 - \frac{(a + \operatorname{barctanh}(cx^3))^3}{3bc^2} \right) \right)$$

6546

$$\frac{1}{3} \left(x^3 (a + \operatorname{barctanh}(cx^3))^3 - 3bc \left(\frac{\log\left(\frac{2}{1 - cx^3}\right) (a + \operatorname{barctanh}(cx^3))^2}{c} - \frac{2b \int \frac{(a + \operatorname{barctanh}(cx^3)) \log\left(\frac{2}{1 - cx^3}\right)}{1 - c^2 x^6} dx^3}{c} - (a + \operatorname{barctanh}(cx^3))^3 \right) \right)$$

6470

$$\frac{1}{3} \left(x^3 (a + \operatorname{barctanh}(cx^3))^3 - 3bc \left(\frac{\log\left(\frac{2}{1 - cx^3}\right) (a + \operatorname{barctanh}(cx^3))^2}{c} - 2b \left(\frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx^3}\right)}{1 - c^2 x^6} dx^3 - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx^3}\right)}{c} \right) \right) \right)$$

6620

$$\frac{1}{3} \left(x^3 (a + \operatorname{barctanh}(cx^3))^3 - 3bc \left(\frac{\log\left(\frac{2}{1 - cx^3}\right) (a + \operatorname{barctanh}(cx^3))^2}{c} - 2b \left(\frac{b \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - cx^3}\right)}{4c} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx^3}\right)}{2c} \right) \right) \right)$$

7164

input `Int [x^2*(a + b*ArcTanh[c*x^3])^3,x]`

output `(x^3*(a + b*ArcTanh[c*x^3])^3 - 3*b*c*(-1/3*(a + b*ArcTanh[c*x^3])^3/(b*c^2) + (((a + b*ArcTanh[c*x^3])^2*Log[2/(1 - c*x^3)])/c - 2*b*(-1/2*((a + b*ArcTanh[c*x^3])*PolyLog[2, 1 - 2/(1 - c*x^3)])/c + (b*PolyLog[3, 1 - 2/(1 - c*x^3)])/(4*c)))/c))/3`

Defintions of rubi rules used

rule 6436 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot b)^p, x_Symbol] \rightarrow \text{Simp}[x \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^p, x] - \text{Simp}[b \cdot c \cdot n \cdot p \cdot \text{Int}[x^n \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2n})], x], x] /;$ $\text{FreeQ}\{a, b, c, n, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[n, 1] \ || \ \text{EqQ}[p, 1])$

rule 6454 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot b)^p \cdot x^m, x_Symbol] \rightarrow \text{Simp}[1/n \cdot \text{Subst}[\text{Int}[x^{(\text{Simplify}[m+1]/n) - 1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x, x^n], x] /;$ $\text{FreeQ}\{a, b, c, m, n, x\} \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[m+1]/n]$

rule 6470 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((d + e \cdot x)), x_Symbol] \rightarrow \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{Log}[2/(1 + e \cdot (x/d))]/e), x] + \text{Simp}[b \cdot c \cdot (p/e) \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{Log}[2/(1 + e \cdot (x/d))]) / (1 - c^2 \cdot x^2)], x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d^2 - e^2, 0]$

rule 6546 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p \cdot x / ((d + e \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot e \cdot (p+1)), x] + \text{Simp}[1/(c \cdot d) \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p / (1 - c \cdot x), x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{IGtQ}[p, 0]$

rule 6620 $\text{Int}[(\text{Log}[u] \cdot (a + \text{ArcTanh}[c \cdot x] \cdot b)^p) / ((d + e \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{PolyLog}[2, 1 - u] / (2 \cdot c \cdot d)), x] + \text{Simp}[b \cdot (p/2) \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{PolyLog}[2, 1 - u] / (d + e \cdot x^2))], x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{EqQ}[(1 - u)^2 - (1 - 2/(1 - c \cdot x))^2, 0]$

rule 7164 $\text{Int}[u \cdot \text{PolyLog}[n, v], x_Symbol] \rightarrow \text{With}\{w = \text{DerivativeDivides}[v, u \cdot v, x]\}, \text{Simp}[w \cdot \text{PolyLog}[n+1, v], x] /;$ $! \text{FalseQ}[w] /;$ $\text{FreeQ}[n, x]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 264 vs. $2(124) = 248$.

Time = 1.94 (sec) , antiderivative size = 265, normalized size of antiderivative = 2.04

method	result
derivativedivides	$a^3 c x^3 + b^3 \left(\operatorname{arctanh}(c x^3)^3 (c x^3 - 1) + 2 \operatorname{arctanh}(c x^3)^3 - 3 \operatorname{arctanh}(c x^3)^2 \ln \left(1 + \frac{(c x^3 + 1)^2}{-c^2 x^6 + 1} \right) - 3 \operatorname{arctanh}(c x^3) \operatorname{polylog} \right)$
default	$a^3 c x^3 + b^3 \left(\operatorname{arctanh}(c x^3)^3 (c x^3 - 1) + 2 \operatorname{arctanh}(c x^3)^3 - 3 \operatorname{arctanh}(c x^3)^2 \ln \left(1 + \frac{(c x^3 + 1)^2}{-c^2 x^6 + 1} \right) - 3 \operatorname{arctanh}(c x^3) \operatorname{polylog} \right)$
parts	$\frac{a^3 x^3}{3} + \frac{b^3 \left(\operatorname{arctanh}(c x^3)^3 (c x^3 - 1) + 2 \operatorname{arctanh}(c x^3)^3 - 3 \operatorname{arctanh}(c x^3)^2 \ln \left(1 + \frac{(c x^3 + 1)^2}{-c^2 x^6 + 1} \right) - 3 \operatorname{arctanh}(c x^3) \operatorname{polylog} \right)}{3c}$

input `int(x^2*(a+b*arctanh(c*x^3))^3,x,method=_RETURNVERBOSE)`

output `1/3/c*(a^3*c*x^3+b^3*(arctanh(c*x^3)^3*(c*x^3-1)+2*arctanh(c*x^3)^3-3*arctanh(c*x^3)^2*ln(1+(c*x^3+1)^2/(-c^2*x^6+1))-3*arctanh(c*x^3)*polylog(2,-(c*x^3+1)^2/(-c^2*x^6+1))+3/2*polylog(3,-(c*x^3+1)^2/(-c^2*x^6+1)))+3*a*b^2*(arctanh(c*x^3)^2*(c*x^3-1)+2*arctanh(c*x^3)^2-2*arctanh(c*x^3)*ln(1+(c*x^3+1)^2/(-c^2*x^6+1))-polylog(2,-(c*x^3+1)^2/(-c^2*x^6+1)))+3*a^2*b*(c*x^3*arctanh(c*x^3)+1/2*ln(-c^2*x^6+1))`

Fricas [F]

$$\int x^2 (a + b \operatorname{arctanh}(c x^3))^3 dx = \int (b \operatorname{arctanh}(c x^3) + a)^3 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c*x^3))^3,x, algorithm="fricas")`

output

```
integral(b^3*x^2*arctanh(c*x^3)^3 + 3*a*b^2*x^2*arctanh(c*x^3)^2 + 3*a^2*b*x^2*arctanh(c*x^3) + a^3*x^2, x)
```

Sympy [F(-1)]

Timed out.

$$\int x^2(a + b \operatorname{arctanh}(cx^3))^3 dx = \text{Timed out}$$

input

```
integrate(x**2*(a+b*atanh(c*x**3))**3,x)
```

output

Timed out

Maxima [F]

$$\int x^2(a + b \operatorname{arctanh}(cx^3))^3 dx = \int (b \operatorname{arctanh}(cx^3) + a)^3 x^2 dx$$

input

```
integrate(x^2*(a+b*arctanh(c*x^3))^3,x, algorithm="maxima")
```

output

```
1/3*a^3*x^3 + 1/2*(2*c*x^3*arctanh(c*x^3) + log(-c^2*x^6 + 1))*a^2*b/c - 1/24*((b^3*c*x^3 - b^3)*log(-c*x^3 + 1)^3 - 3*(2*a*b^2*c*x^3 + (b^3*c*x^3 + b^3)*log(c*x^3 + 1))*log(-c*x^3 + 1)^2)/c - integrate(-1/8*((b^3*c*x^5 - b^3*x^2)*log(c*x^3 + 1)^3 + 6*(a*b^2*c*x^5 - a*b^2*x^2)*log(c*x^3 + 1)^2 - 3*(4*a*b^2*c*x^5 + (b^3*c*x^5 - b^3*x^2)*log(c*x^3 + 1)^2 + 2*((2*a*b^2*c + b^3*c)*x^5 - (2*a*b^2 - b^3)*x^2)*log(c*x^3 + 1))*log(-c*x^3 + 1))/(c*x^3 - 1), x)
```

Giac [F]

$$\int x^2 (a + b \operatorname{arctanh}(cx^3))^3 dx = \int (b \operatorname{artanh}(cx^3) + a)^3 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c*x^3))^3,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^3) + a)^3*x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 (a + b \operatorname{arctanh}(cx^3))^3 dx = \int x^2 (a + b \operatorname{atanh}(cx^3))^3 dx$$

input `int(x^2*(a + b*atanh(c*x^3))^3,x)`

output `int(x^2*(a + b*atanh(c*x^3))^3, x)`

Reduce [F]

$$\int x^2 (a + b \operatorname{arctanh}(cx^3))^3 dx$$

$$= \frac{3 \operatorname{atanh}(cx^3) a^2 b c x^3 - 3 \operatorname{atanh}(cx^3) a^2 b + 3 \left(\int \operatorname{atanh}(cx^3)^3 x^2 dx \right) b^3 c + 9 \left(\int \operatorname{atanh}(cx^3)^2 x^2 dx \right) a b^2 c + \dots}{3c}$$

input `int(x^2*(a+b*atanh(c*x^3))^3,x)`

output `(3*atanh(c*x**3)*a**2*b*c*x**3 - 3*atanh(c*x**3)*a**2*b + 3*int(atanh(c*x**3)**3*x**2,x)*b**3*c + 9*int(atanh(c*x**3)**2*x**2,x)*a*b**2*c + 3*log(c**(2/3)*x**2 - c**(1/3)*x + 1)*a**2*b + 3*log(c**(2/3)*x + c**(1/3))*a**2*b + a**3*c*x**3)/(3*c)`

$$3.127 \quad \int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x} dx$$

Optimal result	1076
Mathematica [C] (verified)	1077
Rubi [A] (verified)	1078
Maple [F]	1080
Fricas [F]	1081
Sympy [F(-1)]	1081
Maxima [F]	1081
Giac [F]	1082
Mupad [F(-1)]	1082
Reduce [F]	1082

Optimal result

Integrand size = 16, antiderivative size = 210

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x} dx = & \frac{2}{3}(a + b \operatorname{arctanh}(cx^3))^3 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx^3}\right) \\ & - \frac{1}{2}b(a + b \operatorname{arctanh}(cx^3))^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx^3}\right) \\ & + \frac{1}{2}b(a + b \operatorname{arctanh}(cx^3))^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - cx^3}\right) \\ & + \frac{1}{2}b^2(a + b \operatorname{arctanh}(cx^3)) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - cx^3}\right) \\ & - \frac{1}{2}b^2(a + b \operatorname{arctanh}(cx^3)) \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 - cx^3}\right) \\ & - \frac{1}{4}b^3 \operatorname{PolyLog}\left(4, 1 - \frac{2}{1 - cx^3}\right) \\ & + \frac{1}{4}b^3 \operatorname{PolyLog}\left(4, -1 + \frac{2}{1 - cx^3}\right) \end{aligned}$$

output

```
-2/3*(a+b*arctanh(c*x^3))^3*arctanh(-1+2/(-c*x^3+1))-1/2*b*(a+b*arctanh(c*
x^3))^2*polylog(2,1-2/(-c*x^3+1))+1/2*b*(a+b*arctanh(c*x^3))^2*polylog(2,-
1+2/(-c*x^3+1))+1/2*b^2*(a+b*arctanh(c*x^3))*polylog(3,1-2/(-c*x^3+1))-1/2
*b^2*(a+b*arctanh(c*x^3))*polylog(3,-1+2/(-c*x^3+1))-1/4*b^3*polylog(4,1-2
/(-c*x^3+1))+1/4*b^3*polylog(4,-1+2/(-c*x^3+1))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.32 (sec) , antiderivative size = 368, normalized size of antiderivative = 1.75

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x} dx = a^3 \log(x) + \frac{1}{2} a^2 b (-\operatorname{PolyLog}(2, -cx^3) + \operatorname{PolyLog}(2, cx^3))$$

$$+ ab^2 \left(\frac{i\pi^3}{24} - \frac{2}{3} \operatorname{arctanh}(cx^3)^3 \right.$$

$$\quad - \operatorname{arctanh}(cx^3)^2 \log(1 + e^{-2\operatorname{arctanh}(cx^3)})$$

$$\quad + \operatorname{arctanh}(cx^3)^2 \log(1 - e^{2\operatorname{arctanh}(cx^3)})$$

$$+ \operatorname{arctanh}(cx^3) \operatorname{PolyLog}(2, -e^{-2\operatorname{arctanh}(cx^3)})$$

$$+ \operatorname{arctanh}(cx^3) \operatorname{PolyLog}(2, e^{2\operatorname{arctanh}(cx^3)})$$

$$\quad + \frac{1}{2} \operatorname{PolyLog}(3, -e^{-2\operatorname{arctanh}(cx^3)})$$

$$\quad \left. - \frac{1}{2} \operatorname{PolyLog}(3, e^{2\operatorname{arctanh}(cx^3)}) \right)$$

$$+ \frac{1}{192} b^3 \left(\pi^4 - 32 \operatorname{arctanh}(cx^3)^4 \right.$$

$$\quad - 64 \operatorname{arctanh}(cx^3)^3 \log(1 + e^{-2\operatorname{arctanh}(cx^3)})$$

$$\quad + 64 \operatorname{arctanh}(cx^3)^3 \log(1 - e^{2\operatorname{arctanh}(cx^3)})$$

$$+ 96 \operatorname{arctanh}(cx^3)^2 \operatorname{PolyLog}(2, -e^{-2\operatorname{arctanh}(cx^3)})$$

$$+ 96 \operatorname{arctanh}(cx^3)^2 \operatorname{PolyLog}(2, e^{2\operatorname{arctanh}(cx^3)})$$

$$+ 96 \operatorname{arctanh}(cx^3) \operatorname{PolyLog}(3, -e^{-2\operatorname{arctanh}(cx^3)})$$

$$\quad - 96 \operatorname{arctanh}(cx^3) \operatorname{PolyLog}(3, e^{2\operatorname{arctanh}(cx^3)})$$

$$\quad + 48 \operatorname{PolyLog}(4, -e^{-2\operatorname{arctanh}(cx^3)})$$

$$\quad \left. + 48 \operatorname{PolyLog}(4, e^{2\operatorname{arctanh}(cx^3)}) \right)$$

input

```
Integrate[(a + b*ArcTanh[c*x^3])^3/x,x]
```

output

```

a^3*Log[x] + (a^2*b*(-PolyLog[2, -(c*x^3)] + PolyLog[2, c*x^3])/2 + a*b^2
*((I/24)*Pi^3 - (2*ArcTanh[c*x^3]^3)/3 - ArcTanh[c*x^3]^2*Log[1 + E^(-2*Ar
cTanh[c*x^3])] + ArcTanh[c*x^3]^2*Log[1 - E^(2*ArcTanh[c*x^3])] + ArcTanh[
c*x^3]*PolyLog[2, -E^(-2*ArcTanh[c*x^3])] + ArcTanh[c*x^3]*PolyLog[2, E^(2
*ArcTanh[c*x^3])] + PolyLog[3, -E^(-2*ArcTanh[c*x^3])/2 - PolyLog[3, E^(2
*ArcTanh[c*x^3])/2] + (b^3*(Pi^4 - 32*ArcTanh[c*x^3]^4 - 64*ArcTanh[c*x^3
]^3*Log[1 + E^(-2*ArcTanh[c*x^3])] + 64*ArcTanh[c*x^3]^3*Log[1 - E^(2*ArcT
anh[c*x^3])] + 96*ArcTanh[c*x^3]^2*PolyLog[2, -E^(-2*ArcTanh[c*x^3])] + 96
*ArcTanh[c*x^3]^2*PolyLog[2, E^(2*ArcTanh[c*x^3])] + 96*ArcTanh[c*x^3]*Pol
yLog[3, -E^(-2*ArcTanh[c*x^3])] - 96*ArcTanh[c*x^3]*PolyLog[3, E^(2*ArcTan
h[c*x^3])] + 48*PolyLog[4, -E^(-2*ArcTanh[c*x^3])] + 48*PolyLog[4, E^(2*Ar
cTanh[c*x^3])])))/192

```

Rubi [A] (verified)

Time = 1.18 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.14, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6450, 6448, 6614, 6620, 6624, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x} dx \\
 & \quad \downarrow \text{6450} \\
 & \frac{1}{3} \int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^3} dx^3 \\
 & \quad \downarrow \text{6448} \\
 & \frac{1}{3} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^3} \right) (a + b \operatorname{arctanh}(cx^3))^3 - 6bc \int \frac{(a + b \operatorname{arctanh}(cx^3))^2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^3} \right)}{1 - c^2 x^6} dx^3 \right) \\
 & \quad \downarrow \text{6614} \\
 & \frac{1}{3} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^3} \right) (a + b \operatorname{arctanh}(cx^3))^3 - 6bc \left(\frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(cx^3))^2 \log \left(2 - \frac{2}{1 - cx^3} \right)}{1 - c^2 x^6} dx^3 - \frac{1}{2} \int \right) \right)
 \end{aligned}$$

↓ 6620

$$\frac{1}{3} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^3} \right) (a + b \operatorname{arctanh}(cx^3))^3 - 6bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - cx^3} \right) (a + b \operatorname{arctanh}(cx^3))^2}{2c} - b \right) \right) \right)$$

↓ 6624

$$\frac{1}{3} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^3} \right) (a + b \operatorname{arctanh}(cx^3))^3 - 6bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - cx^3} \right) (a + b \operatorname{arctanh}(cx^3))^2}{2c} - b \right) \right) \right)$$

↓ 7164

$$\frac{1}{3} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^3} \right) (a + b \operatorname{arctanh}(cx^3))^3 - 6bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - cx^3} \right) (a + b \operatorname{arctanh}(cx^3))^2}{2c} - b \right) \right) \right)$$

input `Int[(a + b*ArcTanh[c*x^3])^3/x,x]`

output `(2*(a + b*ArcTanh[c*x^3])^3*ArcTanh[1 - 2/(1 - c*x^3)] - 6*b*c*(((a + b*ArcTanh[c*x^3])^2*PolyLog[2, 1 - 2/(1 - c*x^3)])/(2*c) - b*(((a + b*ArcTanh[c*x^3])*PolyLog[3, 1 - 2/(1 - c*x^3)])/(2*c) - (b*PolyLog[4, 1 - 2/(1 - c*x^3)])/(4*c))))/2 + (-1/2*((a + b*ArcTanh[c*x^3])^2*PolyLog[2, -1 + 2/(1 - c*x^3)])/c + b*(((a + b*ArcTanh[c*x^3])*PolyLog[3, -1 + 2/(1 - c*x^3)])/(2*c) - (b*PolyLog[4, -1 + 2/(1 - c*x^3)])/(4*c))))/2)/3`

Defintions of rubi rules used

rule 6448 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_)/(x_), x_Symbol] := Simp[2*(a + b*ArcTanh[c*x])^p*ArcTanh[1 - 2/(1 - c*x)], x] - Simp[2*b*c*p Int[(a + b*ArcTanh[c*x])^(p - 1)*(ArcTanh[1 - 2/(1 - c*x)]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1]`

rule 6450 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_)/(x_), x_Symbol] := Simp[1/n Subst[Int[(a + b*ArcTanh[c*x])^p/x, x], x, x^n], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]`

rule 6614 `Int[(ArcTanh[u_]*((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/2 Int[Log[1 + u]*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] - Simp[1/2 Int[Log[1 - u]*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[u^2 - (1 - 2/(1 - c*x))^2, 0]`

rule 6620 `Int[(Log[u_]*((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(-a + b*ArcTanh[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)), x] + Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]`

rule 6624 `Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*PolyLog[k_, u_])/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(PolyLog[k + 1, u]/(2*c*d)), x] - Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[k + 1, u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, k}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[u^2 - (1 - 2/(1 - c*x))^2, 0]`

rule 7164 `Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v, x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]`

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x} dx$$

input `int((a+b*arctanh(c*x^3))^3/x,x)`

output `int((a+b*arctanh(c*x^3))^3/x,x)`

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x} dx = \int \frac{(b \operatorname{arctanh}(cx^3) + a)^3}{x} dx$$

input `integrate((a+b*arctanh(c*x^3))^3/x,x, algorithm="fricas")`

output `integral((b^3*arctanh(c*x^3)^3 + 3*a*b^2*arctanh(c*x^3)^2 + 3*a^2*b*arctanh(c*x^3) + a^3)/x, x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**3))**3/x,x)`

output `Timed out`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x} dx = \int \frac{(b \operatorname{arctanh}(cx^3) + a)^3}{x} dx$$

input `integrate((a+b*arctanh(c*x^3))^3/x,x, algorithm="maxima")`

output `a^3*log(x) + integrate(1/8*b^3*(log(c*x^3 + 1) - log(-c*x^3 + 1))^3/x + 3/4*a*b^2*(log(c*x^3 + 1) - log(-c*x^3 + 1))^2/x + 3/2*a^2*b*(log(c*x^3 + 1) - log(-c*x^3 + 1))/x, x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x} dx = \int \frac{(b \operatorname{artanh}(cx^3) + a)^3}{x} dx$$

input `integrate((a+b*arctanh(c*x^3))^3/x,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^3) + a)^3/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x} dx = \int \frac{(a + b \operatorname{atanh}(cx^3))^3}{x} dx$$

input `int((a + b*atanh(c*x^3))^3/x,x)`

output `int((a + b*atanh(c*x^3))^3/x, x)`

Reduce [F]

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x} dx &= 3 \left(\int \frac{\operatorname{atanh}(cx^3)}{x} dx \right) a^2 b + \left(\int \frac{\operatorname{atanh}(cx^3)^3}{x} dx \right) b^3 \\ &\quad + 3 \left(\int \frac{\operatorname{atanh}(cx^3)^2}{x} dx \right) a b^2 + \log(x) a^3 \end{aligned}$$

input `int((a+b*atanh(c*x^3))^3/x,x)`

output `3*int(atanh(c*x**3)/x,x)*a**2*b + int(atanh(c*x**3)**3/x,x)*b**3 + 3*int(a
tanh(c*x**3)**2/x,x)*a*b**2 + log(x)*a**3`

$$3.128 \quad \int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^4} dx$$

Optimal result	1083
Mathematica [C] (verified)	1084
Rubi [A] (verified)	1085
Maple [F]	1087
Fricas [F]	1087
Sympy [F(-1)]	1088
Maxima [F]	1088
Giac [F]	1088
Mupad [F(-1)]	1089
Reduce [F]	1089

Optimal result

Integrand size = 16, antiderivative size = 120

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^4} dx = & \frac{1}{3}c(a + b \operatorname{arctanh}(cx^3))^3 - \frac{(a + b \operatorname{arctanh}(cx^3))^3}{3x^3} \\ & + bc(a + b \operatorname{arctanh}(cx^3))^2 \log\left(2 - \frac{2}{1 + cx^3}\right) \\ & - b^2c(a + b \operatorname{arctanh}(cx^3)) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + cx^3}\right) \\ & - \frac{1}{2}b^3c \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 + cx^3}\right) \end{aligned}$$

output

```
1/3*c*(a+b*arctanh(c*x^3))^3-1/3*(a+b*arctanh(c*x^3))^3/x^3+b*c*(a+b*arctanh(c*x^3))^2*ln(2-2/(c*x^3+1))-b^2*c*(a+b*arctanh(c*x^3))*polylog(2,-1+2/(c*x^3+1))-1/2*b^3*c*polylog(3,-1+2/(c*x^3+1))
```


Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.86

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^4} dx = -\frac{a^3}{3x^3} - \frac{a^2 b \operatorname{arctanh}(cx^3)}{x^3} + 3a^2 b c \log(x) - \frac{1}{2} a^2 b c \log(1 - c^2 x^6) + ab^2 c \left(\operatorname{arctanh}(cx^3) \left(\left(1 - \frac{1}{cx^3}\right) \operatorname{arctanh}(cx^3) + 2 \log\left(1 - e^{-2 \operatorname{arctanh}(cx^3)}\right)\right) - \operatorname{PolyLog}\left(2, e^{-2 \operatorname{arctanh}(cx^3)}\right)\right) + \frac{1}{3} b^3 c \left(\frac{i\pi^3}{8} - \operatorname{arctanh}(cx^3)^3 - \frac{\operatorname{arctanh}(cx^3)^3}{cx^3} + 3 \operatorname{arctanh}(cx^3)^2 \log\left(1 - e^{2 \operatorname{arctanh}(cx^3)}\right) + 3 \operatorname{arctanh}(cx^3) \operatorname{PolyLog}\left(2, e^{2 \operatorname{arctanh}(cx^3)}\right) - \frac{3}{2} \operatorname{PolyLog}\left(3, e^{2 \operatorname{arctanh}(cx^3)}\right)\right)$$

input `Integrate[(a + b*ArcTanh[c*x^3])^3/x^4,x]`

output `-1/3*a^3/x^3 - (a^2*b*ArcTanh[c*x^3])/x^3 + 3*a^2*b*c*Log[x] - (a^2*b*c*Log[1 - c^2*x^6])/2 + a*b^2*c*(ArcTanh[c*x^3]*((1 - 1/(c*x^3))*ArcTanh[c*x^3] + 2*Log[1 - E^(-2*ArcTanh[c*x^3])]) - PolyLog[2, E^(-2*ArcTanh[c*x^3])]) + (b^3*c*((I/8)*Pi^3 - ArcTanh[c*x^3]^3 - ArcTanh[c*x^3]^3/(c*x^3) + 3*ArcTanh[c*x^3]^2*Log[1 - E^(2*ArcTanh[c*x^3])] + 3*ArcTanh[c*x^3]*PolyLog[2, E^(2*ArcTanh[c*x^3])] - (3*PolyLog[3, E^(2*ArcTanh[c*x^3])])/2))/3`

Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.11, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6454, 6452, 6550, 6494, 6618, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^4} dx$$

$$\downarrow 6454$$

$$\frac{1}{3} \int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^6} dx^3$$

$$\downarrow 6452$$

$$\frac{1}{3} \left(3bc \int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^3(1 - c^2x^6)} dx^3 - \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^3} \right)$$

$$\downarrow 6550$$

$$\frac{1}{3} \left(3bc \left(\int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^3(cx^3 + 1)} dx^3 + \frac{(a + b \operatorname{arctanh}(cx^3))^3}{3b} \right) - \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^3} \right)$$

$$\downarrow 6494$$

$$\frac{1}{3} \left(3bc \left(-2bc \int \frac{(a + b \operatorname{arctanh}(cx^3)) \log \left(2 - \frac{2}{cx^3 + 1} \right)}{1 - c^2x^6} dx^3 + \frac{(a + b \operatorname{arctanh}(cx^3))^3}{3b} + \log \left(2 - \frac{2}{cx^3 + 1} \right) (a + b \operatorname{arctanh}(cx^3)) \right) - \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^3} \right)$$

$$\downarrow 6618$$

$$\frac{1}{3} \left(3bc \left(-2bc \left(\frac{\operatorname{PolyLog} \left(2, \frac{2}{cx^3 + 1} - 1 \right) (a + b \operatorname{arctanh}(cx^3))}{2c} - \frac{1}{2} b \int \frac{\operatorname{PolyLog} \left(2, \frac{2}{cx^3 + 1} - 1 \right)}{1 - c^2x^6} dx^3 \right) + \frac{(a + b \operatorname{arctanh}(cx^3))^3}{3b} + \log \left(2 - \frac{2}{cx^3 + 1} \right) (a + b \operatorname{arctanh}(cx^3)) \right) - \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^3} \right)$$

$$\downarrow 7164$$

$$\frac{1}{3} \left(3bc \left(-2bc \left(\frac{\operatorname{PolyLog} \left(2, \frac{2}{cx^3 + 1} - 1 \right) (a + b \operatorname{arctanh}(cx^3))}{2c} + \frac{b \operatorname{PolyLog} \left(3, \frac{2}{cx^3 + 1} - 1 \right)}{4c} \right) + \frac{(a + b \operatorname{arctanh}(cx^3))^3}{3b} + \log \left(2 - \frac{2}{cx^3 + 1} \right) (a + b \operatorname{arctanh}(cx^3)) \right) - \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^3} \right)$$

input `Int[(a + b*ArcTanh[c*x^3])^3/x^4,x]`

output `(-((a + b*ArcTanh[c*x^3])^3/x^3) + 3*b*c*((a + b*ArcTanh[c*x^3])^3/(3*b) + (a + b*ArcTanh[c*x^3])^2*Log[2 - 2/(1 + c*x^3)] - 2*b*c*((a + b*ArcTanh[c*x^3])*PolyLog[2, -1 + 2/(1 + c*x^3)])/(2*c) + (b*PolyLog[3, -1 + 2/(1 + c*x^3)])/(4*c)))/3`

Defintions of rubi rules used

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]`

rule 6494 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_))), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6550 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_)^2)), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]`

rule 6618

```
Int[(Log[u_]*((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.))/((d_) + (e_.)*(x_)^
2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)), x
] - Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d +
e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d +
e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 + c*x))^2, 0]
```

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^4} dx$$

input

```
int((a+b*arctanh(c*x^3))^3/x^4,x)
```

output

```
int((a+b*arctanh(c*x^3))^3/x^4,x)
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^4} dx = \int \frac{(b \operatorname{arctanh}(cx^3) + a)^3}{x^4} dx$$

input

```
integrate((a+b*arctanh(c*x^3))^3/x^4,x, algorithm="fricas")
```

output

```
integral((b^3*arctanh(c*x^3)^3 + 3*a*b^2*arctanh(c*x^3)^2 + 3*a^2*b*arctan
h(c*x^3) + a^3)/x^4, x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^4} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**3))**3/x**4,x)`output `Timed out`**Maxima [F]**

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^4} dx = \int \frac{(b \operatorname{arctanh}(cx^3) + a)^3}{x^4} dx$$

input `integrate((a+b*arctanh(c*x^3))^3/x^4,x, algorithm="maxima")`output `-1/2*(c*(log(c^2*x^6 - 1) - log(x^6)) + 2*arctanh(c*x^3)/x^3)*a^2*b - 1/3*a^3/x^3 - 1/24*((b^3*c*x^3 - b^3)*log(-c*x^3 + 1)^3 + 3*(2*a*b^2 + (b^3*c*x^3 + b^3)*log(c*x^3 + 1))*log(-c*x^3 + 1)^2)/x^3 - integrate(-1/8*((b^3*c*x^3 - b^3)*log(c*x^3 + 1)^3 + 6*(a*b^2*c*x^3 - a*b^2)*log(c*x^3 + 1)^2 + 3*(4*a*b^2*c*x^3 - (b^3*c*x^3 - b^3)*log(c*x^3 + 1)^2 + 2*(b^3*c^2*x^6 - (2*a*b^2*c - b^3*c)*x^3 + 2*a*b^2)*log(c*x^3 + 1))*log(-c*x^3 + 1))/(c*x^7 - x^4), x)`**Giac [F]**

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^4} dx = \int \frac{(b \operatorname{arctanh}(cx^3) + a)^3}{x^4} dx$$

input `integrate((a+b*arctanh(c*x^3))^3/x^4,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^3) + a)^3/x^4, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^4} dx = \int \frac{(a + b \operatorname{atanh}(cx^3))^3}{x^4} dx$$

input `int((a + b*atanh(c*x^3))^3/x^4,x)`

output `int((a + b*atanh(c*x^3))^3/x^4, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^4} dx$$

$$= \frac{-\operatorname{atanh}(cx^3)^3 b^3 - 3 \operatorname{atanh}(cx^3)^2 a b^2 + 3 \operatorname{atanh}(cx^3) a^2 b c x^3 - 3 \operatorname{atanh}(cx^3) a^2 b - 18 \left(\int \frac{\operatorname{atanh}(cx^3)}{c^2 x^7 - x} dx \right) a b}{1}$$

input `int((a+b*atanh(c*x^3))^3/x^4,x)`

output `(- atanh(c*x**3)**3*b**3 - 3*atanh(c*x**3)**2*a*b**2 + 3*atanh(c*x**3)*a*
*2*b*c*x**3 - 3*atanh(c*x**3)*a**2*b - 18*int(atanh(c*x**3)/(c**2*x**7 - x
,x)*a*b**2*c*x**3 - 9*int(atanh(c*x**3)**2/(c**2*x**7 - x),x)*b**3*c*x**3
- 3*log(c**(2/3)*x**2 - c**(1/3)*x + 1)*a**2*b*c*x**3 - 3*log(c**(2/3)*x
+ c**(1/3))*a**2*b*c*x**3 + 9*log(x)*a**2*b*c*x**3 - a**3)/(3*x**3)`

$$3.129 \quad \int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^7} dx$$

Optimal result	1090
Mathematica [A] (verified)	1091
Rubi [A] (verified)	1091
Maple [F]	1094
Fricas [F]	1094
Sympy [F(-1)]	1094
Maxima [F]	1095
Giac [F]	1095
Mupad [F(-1)]	1096
Reduce [F]	1096

Optimal result

Integrand size = 16, antiderivative size = 136

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^7} dx &= \frac{1}{2}bc^2(a + b \operatorname{arctanh}(cx^3))^2 - \frac{bc(a + b \operatorname{arctanh}(cx^3))^2}{2x^3} \\ &+ \frac{1}{6}c^2(a + b \operatorname{arctanh}(cx^3))^3 - \frac{(a + b \operatorname{arctanh}(cx^3))^3}{6x^6} \\ &+ b^2c^2(a + b \operatorname{arctanh}(cx^3)) \log\left(2 - \frac{2}{1 + cx^3}\right) \\ &- \frac{1}{2}b^3c^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + cx^3}\right) \end{aligned}$$

output

```
1/2*b*c^2*(a+b*arctanh(c*x^3))^2-1/2*b*c*(a+b*arctanh(c*x^3))^2/x^3+1/6*c^
2*(a+b*arctanh(c*x^3))^3-1/6*(a+b*arctanh(c*x^3))^3/x^6+b^2*c^2*(a+b*arcta
nh(c*x^3))*ln(2-2/(c*x^3+1))-1/2*b^3*c^2*polylog(2,-1+2/(c*x^3+1))
```

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.60

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^7} dx$$

$$= \frac{6b^2(-1 + cx^3)(a + acx^3 + bcx^3) \operatorname{arctanh}(cx^3)^2 + 2b^3(-1 + c^2x^6) \operatorname{arctanh}(cx^3)^3 - 6b \operatorname{arctanh}(cx^3) (a^2 +$$

input `Integrate[(a + b*ArcTanh[c*x^3])^3/x^7,x]`

output

```
(6*b^2*(-1 + c*x^3)*(a + a*c*x^3 + b*c*x^3)*ArcTanh[c*x^3]^2 + 2*b^3*(-1 +
c^2*x^6)*ArcTanh[c*x^3]^3 - 6*b*ArcTanh[c*x^3]*(a^2 + 2*a*b*c*x^3 - 2*b^2
*c^2*x^6*Log[1 - E^(-2*ArcTanh[c*x^3])]) + a*(-2*a^2 - 6*a*b*c*x^3 - 3*a*b
*c^2*x^6*Log[1 - c*x^3] + 3*a*b*c^2*x^6*Log[1 + c*x^3] + 12*b^2*c^2*x^6*Lo
g[(c*x^3)/Sqrt[1 - c^2*x^6]]) - 6*b^3*c^2*x^6*PolyLog[2, E^(-2*ArcTanh[c*x
^3])])/(12*x^6)
```

Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.01, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6454, 6452, 6544, 6452, 6510, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^7} dx$$

$$\downarrow 6454$$

$$\frac{1}{3} \int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^9} dx^3$$

$$\downarrow 6452$$

$$\frac{1}{3} \left(\frac{3}{2} bc \int \frac{(a + b \operatorname{arctanh}(cx^3))^2}{x^6(1 - c^2x^6)} dx^3 - \frac{(a + b \operatorname{arctanh}(cx^3))^3}{2x^6} \right)$$

↓ 6544

$$\frac{1}{3} \left(\frac{3}{2} bc \left(c^2 \int \frac{(a + \operatorname{arctanh}(cx^3))^2}{1 - c^2 x^6} dx^3 + \int \frac{(a + \operatorname{arctanh}(cx^3))^2}{x^6} dx^3 \right) - \frac{(a + \operatorname{arctanh}(cx^3))^3}{2x^6} \right)$$

↓ 6452

$$\frac{1}{3} \left(\frac{3}{2} bc \left(c^2 \int \frac{(a + \operatorname{arctanh}(cx^3))^2}{1 - c^2 x^6} dx^3 + 2bc \int \frac{a + \operatorname{arctanh}(cx^3)}{x^3 (1 - c^2 x^6)} dx^3 - \frac{(a + \operatorname{arctanh}(cx^3))^2}{x^3} \right) - \frac{(a + \operatorname{arctanh}(cx^3))^3}{2x^6} \right)$$

↓ 6510

$$\frac{1}{3} \left(\frac{3}{2} bc \left(2bc \int \frac{a + \operatorname{arctanh}(cx^3)}{x^3 (1 - c^2 x^6)} dx^3 + \frac{c(a + \operatorname{arctanh}(cx^3))^3}{3b} - \frac{(a + \operatorname{arctanh}(cx^3))^2}{x^3} \right) - \frac{(a + \operatorname{arctanh}(cx^3))^3}{2x^6} \right)$$

↓ 6550

$$\frac{1}{3} \left(\frac{3}{2} bc \left(2bc \left(\int \frac{a + \operatorname{arctanh}(cx^3)}{x^3 (cx^3 + 1)} dx^3 + \frac{(a + \operatorname{arctanh}(cx^3))^2}{2b} \right) + \frac{c(a + \operatorname{arctanh}(cx^3))^3}{3b} - \frac{(a + \operatorname{arctanh}(cx^3))^3}{x^3} \right) \right)$$

↓ 6494

$$\frac{1}{3} \left(\frac{3}{2} bc \left(2bc \left(-bc \int \frac{\log \left(2 - \frac{2}{cx^3 + 1} \right)}{1 - c^2 x^6} dx^3 + \frac{(a + \operatorname{arctanh}(cx^3))^2}{2b} + \log \left(2 - \frac{2}{cx^3 + 1} \right) (a + \operatorname{arctanh}(cx^3)) \right) \right)$$

↓ 2897

$$\frac{1}{3} \left(\frac{3}{2} bc \left(2bc \left(\frac{(a + \operatorname{arctanh}(cx^3))^2}{2b} + \log \left(2 - \frac{2}{cx^3 + 1} \right) (a + \operatorname{arctanh}(cx^3)) - \frac{1}{2} b \operatorname{PolyLog} \left(2, \frac{2}{cx^3 + 1} - 1 \right) \right) \right)$$

input `Int[(a + b*ArcTanh[c*x^3])^3/x^7,x]`

output `(-1/2*(a + b*ArcTanh[c*x^3])^3/x^6 + (3*b*c*(-((a + b*ArcTanh[c*x^3])^2/x^3) + (c*(a + b*ArcTanh[c*x^3])^3)/(3*b) + 2*b*c*((a + b*ArcTanh[c*x^3])^2/(2*b) + (a + b*ArcTanh[c*x^3])*Log[2 - 2/(1 + c*x^3)] - (b*PolyLog[2, -1 + 2/(1 + c*x^3)]/2)))/2)/3`

Defintions of rubi rules used

rule 2897 `Int[Log[u_]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6494 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((x_)*((d_) + (e_)*(x_))), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6510 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6544 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)*((f_)*(x_)^(m_))/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]`

rule 6550

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)/((x_.)*((d_.) + (e_.)*(x_)^2)),
 x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/
 d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]
```

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^7} dx$$

input

```
int((a+b*arctanh(c*x^3))^3/x^7,x)
```

output

```
int((a+b*arctanh(c*x^3))^3/x^7,x)
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^7} dx = \int \frac{(b \operatorname{arctanh}(cx^3) + a)^3}{x^7} dx$$

input

```
integrate((a+b*arctanh(c*x^3))^3/x^7,x, algorithm="fricas")
```

output

```
integral((b^3*arctanh(c*x^3)^3 + 3*a*b^2*arctanh(c*x^3)^2 + 3*a^2*b*arctan
h(c*x^3) + a^3)/x^7, x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^7} dx = \text{Timed out}$$

input

```
integrate((a+b*atanh(c*x**3))**3/x**7,x)
```

output Timed out

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^7} dx = \int \frac{(b \operatorname{arctanh}(cx^3) + a)^3}{x^7} dx$$

input `integrate((a+b*arctanh(c*x^3))^3/x^7,x, algorithm="maxima")`

output

```
1/4*((c*log(c*x^3 + 1) - c*log(c*x^3 - 1) - 2/x^3)*c - 2*arctanh(c*x^3)/x^6)*a^2*b + 1/8*((2*(log(c*x^3 - 1) - 2)*log(c*x^3 + 1) - log(c*x^3 + 1)^2 - log(c*x^3 - 1)^2 - 4*log(c*x^3 - 1) + 24*log(x))*c^2 + 4*(c*log(c*x^3 + 1) - c*log(c*x^3 - 1) - 2/x^3)*c*arctanh(c*x^3))*a*b^2 - 1/48*b^3*(((c^2*x^6 - 1)*log(-c*x^3 + 1)^3 + 3*(2*c*x^3 - (c^2*x^6 - 1)*log(c*x^3 + 1))*log(-c*x^3 + 1)^2)/x^6 + 6*integrate(-((c*x^3 - 1)*log(c*x^3 + 1)^3 + 3*(2*c^2*x^6 - (c*x^3 - 1)*log(c*x^3 + 1)^2 - (c^3*x^9 - c*x^3)*log(c*x^3 + 1))*log(-c*x^3 + 1))/(c*x^10 - x^7), x)) - 1/2*a*b^2*arctanh(c*x^3)^2/x^6 - 1/6*a^3/x^6
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^3))^3}{x^7} dx = \int \frac{(b \operatorname{arctanh}(cx^3) + a)^3}{x^7} dx$$

input `integrate((a+b*arctanh(c*x^3))^3/x^7,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^3) + a)^3/x^7, x)`

3.130 $\int (dx)^m (a + \operatorname{arctanh}(cx^3))^3 dx$

Optimal result	1097
Mathematica [N/A]	1097
Rubi [N/A]	1098
Maple [N/A]	1098
Fricas [N/A]	1099
Sympy [F(-1)]	1099
Maxima [N/A]	1099
Giac [N/A]	1100
Mupad [N/A]	1100
Reduce [N/A]	1101

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int (dx)^m (a + \operatorname{arctanh}(cx^3))^3 dx = \operatorname{Int}\left((dx)^m (a + \operatorname{arctanh}(cx^3))^3, x\right)$$

output `Defer(Int)((d*x)^m*(a+b*arctanh(c*x^3))^3,x)`

Mathematica [N/A]

Not integrable

Time = 1.59 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + \operatorname{arctanh}(cx^3))^3 dx = \int (dx)^m (a + \operatorname{arctanh}(cx^3))^3 dx$$

input `Integrate[(d*x)^m*(a + b*ArcTanh[c*x^3])^3,x]`

output `Integrate[(d*x)^m*(a + b*ArcTanh[c*x^3])^3, x]`

Rubi [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^3 dx$$

↓ 6468

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^3 dx$$

input `Int[(d*x)^m*(a + b*ArcTanh[c*x^3])^3,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.10 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^3 dx$$

input `int((d*x)^m*(a+b*arctanh(c*x^3))^3,x)`

output `int((d*x)^m*(a+b*arctanh(c*x^3))^3,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 50, normalized size of antiderivative = 2.78

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^3 dx = \int (b \operatorname{artanh}(cx^3) + a)^3 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^3))^3,x, algorithm="fricas")`

output `integral((b^3*arctanh(c*x^3)^3 + 3*a*b^2*arctanh(c*x^3)^2 + 3*a^2*b*arctanh(c*x^3) + a^3)*(d*x)^m, x)`

Sympy [F(-1)]

Timed out.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^3 dx = \text{Timed out}$$

input `integrate((d*x)**m*(a+b*atanh(c*x**3))**3,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 2.49 (sec) , antiderivative size = 416, normalized size of antiderivative = 23.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^3 dx = \int (b \operatorname{artanh}(cx^3) + a)^3 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^3))^3,x, algorithm="maxima")`

output

```
-1/8*b^3*d^m*x*x^m*log(-c*x^3 + 1)^3/(m + 1) + (d*x)^(m + 1)*a^3/(d*(m + 1)) + integrate(1/8*((b^3*c*d^m*(m + 1)*x^3 - b^3*d^m*(m + 1))*x^m*log(c*x^3 + 1)^3 + 6*(a*b^2*c*d^m*(m + 1)*x^3 - a*b^2*d^m*(m + 1))*x^m*log(c*x^3 + 1)^2 + 12*(a^2*b*c*d^m*(m + 1)*x^3 - a^2*b*d^m*(m + 1))*x^m*log(c*x^3 + 1) + 3*((b^3*c*d^m*(m + 1)*x^3 - b^3*d^m*(m + 1))*x^m*log(c*x^3 + 1) - (2*a*b^2*d^m*(m + 1) - (2*a*b^2*c*d^m*(m + 1) + 3*b^3*c*d^m)*x^3)*x^m*log(-c*x^3 + 1)^2 - 3*((b^3*c*d^m*(m + 1)*x^3 - b^3*d^m*(m + 1))*x^m*log(c*x^3 + 1)^2 + 4*(a*b^2*c*d^m*(m + 1)*x^3 - a*b^2*d^m*(m + 1))*x^m*log(c*x^3 + 1) + 4*(a^2*b*c*d^m*(m + 1)*x^3 - a^2*b*d^m*(m + 1))*x^m*log(-c*x^3 + 1))/(c*(m + 1)*x^3 - m - 1), x)
```

Giac [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^3 dx = \int (b \operatorname{artanh}(cx^3) + a)^3 (dx)^m dx$$

input

```
integrate((d*x)^m*(a+b*arctanh(c*x^3))^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^3) + a)^3*(d*x)^m, x)
```

Mupad [N/A]

Not integrable

Time = 3.56 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^3 dx = \int (dx)^m (a + b \operatorname{atanh}(cx^3))^3 dx$$

input

```
int((d*x)^m*(a + b*atanh(c*x^3))^3,x)
```

output

```
int((d*x)^m*(a + b*atanh(c*x^3))^3, x)
```

Reduce [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 133, normalized size of antiderivative = 7.39

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^3 dx$$

$$= \frac{d^m \left(x^m a^3 x + 3 \left(\int x^m \operatorname{atanh}(cx^3) dx \right) a^2 b m + 3 \left(\int x^m \operatorname{atanh}(cx^3) dx \right) a^2 b + \left(\int x^m \operatorname{atanh}(cx^3)^3 dx \right) b^3 m + \dots \right)}{m + 1}$$

input

```
int((d*x)^m*(a+b*atanh(c*x^3))^3,x)
```

output

```
(d**m*(x**m*a**3*x + 3*int(x**m*atanh(c*x**3),x)*a**2*b*m + 3*int(x**m*atanh(c*x**3),x)*a**2*b + int(x**m*atanh(c*x**3)**3,x)*b**3*m + int(x**m*atanh(c*x**3)**3,x)*b**3 + 3*int(x**m*atanh(c*x**3)**2,x)*a*b**2*m + 3*int(x**m*atanh(c*x**3)**2,x)*a*b**2))/(m + 1)
```

3.131 $\int (dx)^m (a + \operatorname{arctanh}(cx^3))^2 dx$

Optimal result	1102
Mathematica [N/A]	1102
Rubi [N/A]	1103
Maple [N/A]	1103
Fricas [N/A]	1104
Sympy [F(-1)]	1104
Maxima [N/A]	1104
Giac [N/A]	1105
Mupad [N/A]	1105
Reduce [N/A]	1106

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int (dx)^m (a + \operatorname{arctanh}(cx^3))^2 dx = \operatorname{Int}\left((dx)^m (a + \operatorname{arctanh}(cx^3))^2, x\right)$$

output `Defer(Int)((d*x)^m*(a+b*arctanh(c*x^3))^2,x)`

Mathematica [N/A]

Not integrable

Time = 0.99 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + \operatorname{arctanh}(cx^3))^2 dx = \int (dx)^m (a + \operatorname{arctanh}(cx^3))^2 dx$$

input `Integrate[(d*x)^m*(a + b*ArcTanh[c*x^3])^2,x]`

output `Integrate[(d*x)^m*(a + b*ArcTanh[c*x^3])^2, x]`

Rubi [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^2 dx$$

↓ 6468

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^2 dx$$

input `Int[(d*x)^m*(a + b*ArcTanh[c*x^3])^2,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^2 dx$$

input `int((d*x)^m*(a+b*arctanh(c*x^3))^2,x)`

output `int((d*x)^m*(a+b*arctanh(c*x^3))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.89

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^2 dx = \int (b \operatorname{artanh}(cx^3) + a)^2 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^3))^2,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x^3)^2 + 2*a*b*arctanh(c*x^3) + a^2)*(d*x)^m, x)`

Sympy [F(-1)]

Timed out.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^2 dx = \text{Timed out}$$

input `integrate((d*x)**m*(a+b*atanh(c*x**3))**2,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 1.29 (sec) , antiderivative size = 239, normalized size of antiderivative = 13.28

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^2 dx = \int (b \operatorname{artanh}(cx^3) + a)^2 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^3))^2,x, algorithm="maxima")`

output

```
1/4*b^2*d^m*x^m*log(-c*x^3 + 1)^2/(m + 1) + (d*x)^(m + 1)*a^2/(d*(m + 1)
) - integrate(-1/4*((b^2*c*d^m*(m + 1)*x^3 - b^2*d^m*(m + 1))*x^m*log(c*x^
3 + 1)^2 + 4*(a*b*c*d^m*(m + 1)*x^3 - a*b*d^m*(m + 1))*x^m*log(c*x^3 + 1)
- 2*((b^2*c*d^m*(m + 1)*x^3 - b^2*d^m*(m + 1))*x^m*log(c*x^3 + 1) - (2*a*b
*d^m*(m + 1) - (2*a*b*c*d^m*(m + 1) + 3*b^2*c*d^m)*x^3)*x^m*log(-c*x^3 +
1))/(c*(m + 1)*x^3 - m - 1), x)
```

Giac [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^2 dx = \int (b \operatorname{arctanh}(cx^3) + a)^2 (dx)^m dx$$

input

```
integrate((d*x)^m*(a+b*arctanh(c*x^3))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^3) + a)^2*(d*x)^m, x)
```

Mupad [N/A]

Not integrable

Time = 3.55 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3))^2 dx = \int (dx)^m (a + b \operatorname{atanh}(cx^3))^2 dx$$

input

```
int((d*x)^m*(a + b*atanh(c*x^3))^2,x)
```

output

```
int((d*x)^m*(a + b*atanh(c*x^3))^2, x)
```

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 88, normalized size of antiderivative = 4.89

$$\int (dx)^m (a + b \operatorname{atanh}(cx^3))^2 dx$$

$$= \frac{d^m \left(x^m a^2 x + 2 \left(\int x^m \operatorname{atanh}(cx^3) dx \right) abm + 2 \left(\int x^m \operatorname{atanh}(cx^3) dx \right) ab + \left(\int x^m \operatorname{atanh}(cx^3)^2 dx \right) b^2 m + \dots \right)}{m + 1}$$

input

```
int((d*x)^m*(a+b*atanh(c*x^3))^2,x)
```

output

```
(d**m*(x**m*a**2*x + 2*int(x**m*atanh(c*x**3),x)*a*b*m + 2*int(x**m*atanh(c*x**3),x)*a*b + int(x**m*atanh(c*x**3)**2,x)*b**2*m + int(x**m*atanh(c*x**3)**2,x)*b**2))/(m + 1)
```

3.132 $\int (dx)^m (a + \operatorname{barctanh}(cx^3)) dx$

Optimal result	1107
Mathematica [A] (verified)	1107
Rubi [A] (verified)	1108
Maple [F]	1109
Fricas [F]	1109
Sympy [F(-1)]	1110
Maxima [F]	1110
Giac [F]	1110
Mupad [F(-1)]	1111
Reduce [F]	1111

Optimal result

Integrand size = 16, antiderivative size = 74

$$\int (dx)^m (a + \operatorname{barctanh}(cx^3)) dx = \frac{(dx)^{1+m} (a + \operatorname{barctanh}(cx^3))}{d(1+m)} - \frac{3bc(dx)^{4+m} \operatorname{Hypergeometric2F1}\left(1, \frac{4+m}{6}, \frac{10+m}{6}, c^2x^6\right)}{d^4(1+m)(4+m)}$$

output

```
(d*x)^(1+m)*(a+b*arctanh(c*x^3))/d/(1+m)-3*b*c*(d*x)^(4+m)*hypergeom([1, 2
/3+1/6*m], [5/3+1/6*m], c^2*x^6)/d^4/(1+m)/(4+m)
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.86

$$\int (dx)^m (a + \operatorname{barctanh}(cx^3)) dx = -\frac{x(dx)^m (-(4+m)(a + \operatorname{barctanh}(cx^3))) + 3bcx^3 \operatorname{Hypergeometric2F1}\left(1, \frac{4+m}{6}, \frac{10+m}{6}, c^2x^6\right)}{(1+m)(4+m)}$$

input

```
Integrate[(d*x)^m*(a + b*ArcTanh[c*x^3]), x]
```


output

```

-((x*(d*x)^m*(-((4 + m)*(a + b*ArcTanh[c*x^3])) + 3*b*c*x^3*Hypergeometric
2F1[1, (4 + m)/6, (10 + m)/6, c^2*x^6]))/((1 + m)*(4 + m))

```

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {6464, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (dx)^m (a + b \operatorname{arctanh}(cx^3)) dx \\
 & \quad \downarrow \text{6464} \\
 & \frac{(dx)^{m+1} (a + b \operatorname{arctanh}(cx^3))}{d(m+1)} - \frac{3bc \int \frac{(dx)^{m+3}}{1-c^2x^6} dx}{d^3(m+1)} \\
 & \quad \downarrow \text{888} \\
 & \frac{(dx)^{m+1} (a + b \operatorname{arctanh}(cx^3))}{d(m+1)} - \frac{3bc(dx)^{m+4} \operatorname{Hypergeometric2F1}\left(1, \frac{m+4}{6}, \frac{m+10}{6}, c^2x^6\right)}{d^4(m+1)(m+4)}
 \end{aligned}$$

input

```

Int[(d*x)^m*(a + b*ArcTanh[c*x^3]),x]

```

output

```

((d*x)^(1 + m)*(a + b*ArcTanh[c*x^3]))/(d*(1 + m)) - (3*b*c*(d*x)^(4 + m)*
Hypergeometric2F1[1, (4 + m)/6, (10 + m)/6, c^2*x^6])/(d^4*(1 + m)*(4 + m)
)

```

Definitions of rubi rules used

rule 888 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p *((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 6464 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))*((d_)*(x_))^(m_), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]`

Maple [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3)) dx$$

input `int((d*x)^m*(a+b*arctanh(c*x^3)),x)`

output `int((d*x)^m*(a+b*arctanh(c*x^3)),x)`

Fricas [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3)) dx = \int (b \operatorname{arctanh}(cx^3) + a)(dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^3)),x, algorithm="fricas")`

output `integral((b*arctanh(c*x^3) + a)*(d*x)^m, x)`

Sympy [F(-1)]

Timed out.

$$\int (dx)^m (a + \operatorname{barctanh}(cx^3)) dx = \text{Timed out}$$

input `integrate((d*x)**m*(a+b*atanh(c*x**3)),x)`

output `Timed out`

Maxima [F]

$$\int (dx)^m (a + \operatorname{barctanh}(cx^3)) dx = \int (\operatorname{bartanh}(cx^3) + a)(dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^3)),x, algorithm="maxima")`

output `1/2*(6*c*d^m*integrate(x^3*x^m/(c^2*(m+1)*x^6 - m - 1), x) + (d^m*x*x^m*log(c*x^3 + 1) - d^m*x*x^m*log(-c*x^3 + 1))/(m + 1))*b + (d*x)^(m + 1)*a/(d*(m + 1))`

Giac [F]

$$\int (dx)^m (a + \operatorname{barctanh}(cx^3)) dx = \int (\operatorname{bartanh}(cx^3) + a)(dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^3)),x, algorithm="giac")`

output `integrate((b*arctanh(c*x^3) + a)*(d*x)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^3)) dx = \int (dx)^m (a + b \operatorname{atanh}(cx^3)) dx$$

input `int((d*x)^m*(a + b*atanh(c*x^3)),x)`output `int((d*x)^m*(a + b*atanh(c*x^3)), x)`**Reduce [F]**

$$\begin{aligned} & \int (dx)^m (a + b \operatorname{arctanh}(cx^3)) dx \\ &= \frac{d^m (x^m a x + (\int x^m \operatorname{atanh}(c x^3) dx) b m + (\int x^m \operatorname{atanh}(c x^3) dx) b)}{m + 1} \end{aligned}$$

input `int((d*x)^m*(a+b*atanh(c*x^3)),x)`output `(d**m*(x**m*a*x + int(x**m*atanh(c*x**3),x)*b*m + int(x**m*atanh(c*x**3),x)*b))/(m + 1)`

$$3.133 \quad \int \frac{(dx)^m}{a+b\operatorname{arctanh}(cx^3)} dx$$

Optimal result	1112
Mathematica [N/A]	1112
Rubi [N/A]	1113
Maple [N/A]	1113
Fricas [N/A]	1114
Sympy [F(-1)]	1114
Maxima [N/A]	1114
Giac [N/A]	1115
Mupad [N/A]	1115
Reduce [N/A]	1115

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{(dx)^m}{a + b\operatorname{arctanh}(cx^3)} dx = \operatorname{Int}\left(\frac{(dx)^m}{a + b\operatorname{arctanh}(cx^3)}, x\right)$$

output `Defer(Int)((d*x)^m/(a+b*arctanh(c*x^3)), x)`

Mathematica [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b\operatorname{arctanh}(cx^3)} dx = \int \frac{(dx)^m}{a + b\operatorname{arctanh}(cx^3)} dx$$

input `Integrate[(d*x)^m/(a + b*ArcTanh[c*x^3]), x]`

output `Integrate[(d*x)^m/(a + b*ArcTanh[c*x^3]), x]`

Rubi [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^3)} dx$$

↓ 6468

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^3)} dx$$

input `Int[(d*x)^m/(a + b*ArcTanh[c*x^3]),x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.10 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^3)} dx$$

input `int((d*x)^m/(a+b*arctanh(c*x^3)),x)`

output `int((d*x)^m/(a+b*arctanh(c*x^3)),x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^3)} dx = \int \frac{(dx)^m}{b \operatorname{artanh}(cx^3) + a} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x^3)),x, algorithm="fricas")`

output `integral((d*x)^m/(b*arctanh(c*x^3) + a), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^3)} dx = \text{Timed out}$$

input `integrate((d*x)**m/(a+b*atanh(c*x**3)),x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^3)} dx = \int \frac{(dx)^m}{b \operatorname{artanh}(cx^3) + a} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x^3)),x, algorithm="maxima")`

output `integrate((d*x)^m/(b*arctanh(c*x^3) + a), x)`

Giac [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^3)} dx = \int \frac{(dx)^m}{b \operatorname{artanh}(cx^3) + a} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x^3)),x, algorithm="giac")`

output `integrate((d*x)^m/(b*arctanh(c*x^3) + a), x)`

Mupad [N/A]

Not integrable

Time = 3.33 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^3)} dx = \int \frac{(dx)^m}{a + b \operatorname{atanh}(cx^3)} dx$$

input `int((d*x)^m/(a + b*atanh(c*x^3)),x)`

output `int((d*x)^m/(a + b*atanh(c*x^3)), x)`

Reduce [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^3)} dx = d^m \left(\int \frac{x^m}{\operatorname{atanh}(cx^3) b + a} dx \right)$$

input `int((d*x)^m/(a+b*atanh(c*x^3)),x)`

output `d**m*int(x**m/(atanh(c*x**3)*b + a),x)`

$$3.134 \quad \int \frac{(dx)^m}{(a+b\operatorname{arctanh}(cx^3))^2} dx$$

Optimal result	1117
Mathematica [N/A]	1117
Rubi [N/A]	1118
Maple [N/A]	1118
Fricas [N/A]	1119
Sympy [F(-1)]	1119
Maxima [N/A]	1119
Giac [N/A]	1120
Mupad [N/A]	1120
Reduce [N/A]	1121

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{(dx)^m}{(a + b\operatorname{arctanh}(cx^3))^2} dx = \operatorname{Int}\left(\frac{(dx)^m}{(a + b\operatorname{arctanh}(cx^3))^2}, x\right)$$

output `Defer(Int)((d*x)^m/(a+b*arctanh(c*x^3))^2,x)`

Mathematica [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{(a + b\operatorname{arctanh}(cx^3))^2} dx = \int \frac{(dx)^m}{(a + b\operatorname{arctanh}(cx^3))^2} dx$$

input `Integrate[(d*x)^m/(a + b*ArcTanh[c*x^3])^2,x]`

output `Integrate[(d*x)^m/(a + b*ArcTanh[c*x^3])^2, x]`

Rubi [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^3))^2} dx$$

↓ 6468

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^3))^2} dx$$

input `Int[(d*x)^m/(a + b*ArcTanh[c*x^3])^2,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^3))^2} dx$$

input `int((d*x)^m/(a+b*arctanh(c*x^3))^2,x)`

output `int((d*x)^m/(a+b*arctanh(c*x^3))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.00

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^3))^2} dx = \int \frac{(dx)^m}{(b \operatorname{artanh}(cx^3) + a)^2} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x^3))^2,x, algorithm="fricas")`

output `integral((d*x)^m/(b^2*arctanh(c*x^3)^2 + 2*a*b*arctanh(c*x^3) + a^2), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^3))^2} dx = \text{Timed out}$$

input `integrate((d*x)**m/(a+b*atanh(c*x**3))**2,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 141, normalized size of antiderivative = 7.83

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^3))^2} dx = \int \frac{(dx)^m}{(b \operatorname{artanh}(cx^3) + a)^2} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x^3))^2,x, algorithm="maxima")`

output

```
2/3*(c^2*d^m*x^6 - d^m)*x^m/(b^2*c*x^2*log(c*x^3 + 1) - b^2*c*x^2*log(-c*x^3 + 1) + 2*a*b*c*x^2) + integrate(-2/3*(c^2*d^m*(m + 4)*x^6 - d^m*(m - 2)*x^m/(b^2*c*x^3*log(c*x^3 + 1) - b^2*c*x^3*log(-c*x^3 + 1) + 2*a*b*c*x^3), x)
```

Giac [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^3))^2} dx = \int \frac{(dx)^m}{(b \operatorname{artanh}(cx^3) + a)^2} dx$$

input

```
integrate((d*x)^m/(a+b*arctanh(c*x^3))^2,x, algorithm="giac")
```

output

```
integrate((d*x)^m/(b*arctanh(c*x^3) + a)^2, x)
```

Mupad [N/A]

Not integrable

Time = 3.60 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^3))^2} dx = \int \frac{(dx)^m}{(a + b \operatorname{atanh}(cx^3))^2} dx$$

input

```
int((d*x)^m/(a + b*atanh(c*x^3))^2,x)
```

output

```
int((d*x)^m/(a + b*atanh(c*x^3))^2, x)
```

Reduce [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.11

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^3))^2} dx = d^m \left(\int \frac{x^m}{\operatorname{atanh}(cx^3)^2 b^2 + 2 \operatorname{atanh}(cx^3) ab + a^2} dx \right)$$

input `int((d*x)^m/(a+b*atanh(c*x^3))^2,x)`output `d**m*int(x**m/(atanh(c*x**3)**2*b**2 + 2*atanh(c*x**3)*a*b + a**2),x)`

3.135 $\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx$

Optimal result	1122
Mathematica [A] (verified)	1122
Rubi [A] (verified)	1123
Maple [A] (verified)	1124
Fricas [A] (verification not implemented)	1125
Sympy [A] (verification not implemented)	1125
Maxima [A] (verification not implemented)	1126
Giac [B] (verification not implemented)	1126
Mupad [B] (verification not implemented)	1127
Reduce [B] (verification not implemented)	1127

Optimal result

Integrand size = 14, antiderivative size = 50

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{1}{4} b c^3 x + \frac{1}{12} b c x^3 + \frac{1}{4} x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) - \frac{1}{4} b c^4 \operatorname{arctanh} \left(\frac{x}{c} \right)$$

output `1/4*b*c^3*x+1/12*b*c*x^3+1/4*x^4*(a+b*arctanh(c/x))-1/4*b*c^4*arctanh(x/c)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.34

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{1}{4} b c^3 x + \frac{1}{12} b c x^3 + \frac{a x^4}{4} + \frac{1}{4} b x^4 \operatorname{arctanh} \left(\frac{c}{x} \right) + \frac{1}{8} b c^4 \log(-c + x) - \frac{1}{8} b c^4 \log(c + x)$$

input `Integrate[x^3*(a + b*ArcTanh[c/x]),x]`

output `(b*c^3*x)/4 + (b*c*x^3)/12 + (a*x^4)/4 + (b*x^4*ArcTanh[c/x])/4 + (b*c^4*Log[-c + x])/8 - (b*c^4*Log[c + x])/8`

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.96, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6452, 795, 254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right) \right) dx \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{4}bc \int \frac{x^2}{1 - \frac{c^2}{x^2}} dx + \frac{1}{4}x^4 \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right) \right) \\
 & \quad \downarrow \text{795} \\
 & \frac{1}{4}bc \int \frac{x^4}{x^2 - c^2} dx + \frac{1}{4}x^4 \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right) \right) \\
 & \quad \downarrow \text{254} \\
 & \frac{1}{4}bc \int \left(\frac{c^4}{x^2 - c^2} + c^2 + x^2 \right) dx + \frac{1}{4}x^4 \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right) \right) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{4}x^4 \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right) \right) + \frac{1}{4}bc \left(c^3 \left(-\operatorname{arctanh}\left(\frac{x}{c}\right) \right) + c^2x + \frac{x^3}{3} \right)
 \end{aligned}$$

input `Int [x^3*(a + b*ArcTanh[c/x]),x]`

output `(x^4*(a + b*ArcTanh[c/x]))/4 + (b*c*(c^2*x + x^3/3 - c^3*ArcTanh[x/c]))/4`

Definitions of rubi rules used

rule 254 $\text{Int}[(x_)^{(m_)} / ((a_) + (b_)*(x_)^2), x_Symbol] \rightarrow \text{Int}[\text{PolynomialDivide}[x^m, a + b*x^2, x], x] /;$ FreeQ[{a, b}, x] && IGtQ[m, 3]

rule 795 $\text{Int}[(x_)^{(m_)} * ((a_) + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Int}[x^{(m + n*p)} * (b + a/x^n)^p, x] /;$ FreeQ[{a, b, m, n}, x] && IntegerQ[p] && NegQ[n]

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /;$ SumQ[u]

rule 6452 $\text{Int}[(a_) + \text{ArcTanh}[(c_)*(x_)^{(n_)}] * (b_)^{(p_)} * (x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)} * ((a + b*\text{ArcTanh}[c*x^n])^p / (m + 1)), x] - \text{Simp}[b*c*n*(p/(m + 1)) \text{Int}[x^{(m + n)} * ((a + b*\text{ArcTanh}[c*x^n])^{(p - 1)} / (1 - c^2*x^{(2*n)})), x], x] /;$ FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]

Maple [A] (verified)

Time = 0.56 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.92

method	result
paralelrisch	$\frac{x^4 \operatorname{arctanh}(\frac{c}{x})b}{4} - \frac{\operatorname{arctanh}(\frac{c}{x})bc^4}{4} + \frac{x^4a}{4} + \frac{bcx^3}{12} + \frac{bc^3x}{4}$
parts	$\frac{x^4a}{4} - bc^4 \left(-\frac{x^4 \operatorname{arctanh}(\frac{c}{x})}{4c^4} + \frac{\ln(1+\frac{c}{x})}{8} - \frac{\ln(\frac{c}{x}-1)}{8} - \frac{x^3}{12c^3} - \frac{x}{4c} \right)$
derivativedivides	$-c^4 \left(-\frac{ax^4}{4c^4} + b \left(-\frac{x^4 \operatorname{arctanh}(\frac{c}{x})}{4c^4} + \frac{\ln(1+\frac{c}{x})}{8} - \frac{\ln(\frac{c}{x}-1)}{8} - \frac{x^3}{12c^3} - \frac{x}{4c} \right) \right)$
default	$-c^4 \left(-\frac{ax^4}{4c^4} + b \left(-\frac{x^4 \operatorname{arctanh}(\frac{c}{x})}{4c^4} + \frac{\ln(1+\frac{c}{x})}{8} - \frac{\ln(\frac{c}{x}-1)}{8} - \frac{x^3}{12c^3} - \frac{x}{4c} \right) \right)$
orering	$-\frac{(2c^4 - c^2x^2 - x^4)(a + b \operatorname{arctanh}(\frac{c}{x}))}{2} + \frac{(3c^2 + x^2)(c-x)(x+c) \left(3x^2(a + b \operatorname{arctanh}(\frac{c}{x})) - \frac{xbc}{1 - \frac{c^2}{x^2}} \right)}{12x^2}$
risch	$\frac{bx^4 \ln(x+c)}{8} - \frac{bx^4 \ln(c-x)}{8} - \frac{i\pi b x^4 \operatorname{csgn}(\frac{i}{x}) \operatorname{csgn}(i(x+c)) \operatorname{csgn}(\frac{i(x+c)}{x})}{16} - \frac{i\pi b x^4 \operatorname{csgn}(i(c-x)) \operatorname{csgn}(\frac{i(c-x)}{x})^2}{16}$

input $\text{int}(x^3*(a+b*\operatorname{arctanh}(c/x)), x, \text{method}=_RETURNVERBOSE)$

output $1/4*x^4*\operatorname{arctanh}(c/x)*b-1/4*\operatorname{arctanh}(c/x)*b*c^4+1/4*x^4*a+1/12*b*c*x^3+1/4*b*c^3*x$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.96

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{1}{4} b c^3 x + \frac{1}{12} b c x^3 + \frac{1}{4} a x^4 - \frac{1}{8} (b c^4 - b x^4) \log \left(-\frac{c+x}{c-x} \right)$$

input `integrate(x^3*(a+b*arctanh(c/x)),x, algorithm="fricas")`

output $1/4*b*c^3*x + 1/12*b*c*x^3 + 1/4*a*x^4 - 1/8*(b*c^4 - b*x^4)*\log(-(c+x)/(c-x))$

Sympy [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.92

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{a x^4}{4} - \frac{b c^4 \operatorname{atanh} \left(\frac{c}{x} \right)}{4} + \frac{b c^3 x}{4} + \frac{b c x^3}{12} + \frac{b x^4 \operatorname{atanh} \left(\frac{c}{x} \right)}{4}$$

input `integrate(x**3*(a+b*atanh(c/x)),x)`

output $a*x**4/4 - b*c**4*atanh(c/x)/4 + b*c**3*x/4 + b*c*x**3/12 + b*x**4*atanh(c/x)/4$

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.14

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx$$

$$= \frac{1}{4} a x^4$$

$$+ \frac{1}{24} \left(6 x^4 \operatorname{artanh} \left(\frac{c}{x} \right) - (3 c^3 \log (c + x) - 3 c^3 \log (-c + x) - 6 c^2 x - 2 x^3) c \right) b$$

input `integrate(x^3*(a+b*arctanh(c/x)),x, algorithm="maxima")`

output `1/4*a*x^4 + 1/24*(6*x^4*arctanh(c/x) - (3*c^3*log(c + x) - 3*c^3*log(-c + x) - 6*c^2*x - 2*x^3)*c)*b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 262 vs. 2(42) = 84.

Time = 0.12 (sec) , antiderivative size = 262, normalized size of antiderivative = 5.24

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx$$

$$= \frac{3 \left(\frac{b(c+x)^3 c^5}{(c-x)^3} + \frac{b(c+x)c^5}{c-x} \right) \log \left(-\frac{c+x}{c-x} \right)}{\frac{(c+x)^4}{(c-x)^4} + \frac{4(c+x)^3}{(c-x)^3} + \frac{6(c+x)^2}{(c-x)^2} + \frac{4(c+x)}{c-x} + 1} + \frac{2bc^5 + \frac{6a(c+x)^3 c^5}{(c-x)^3} + \frac{3b(c+x)^3 c^5}{(c-x)^3} + \frac{6b(c+x)^2 c^5}{(c-x)^2} + \frac{6a(c+x)c^5}{c-x} + \frac{5b(c+x)c^5}{c-x}}{\frac{(c+x)^4}{(c-x)^4} + \frac{4(c+x)^3}{(c-x)^3} + \frac{6(c+x)^2}{(c-x)^2} + \frac{4(c+x)}{c-x} + 1}$$

$$3c$$

input `integrate(x^3*(a+b*arctanh(c/x)),x, algorithm="giac")`

output `-1/3*(3*(b*(c + x)^3*c^5/(c - x)^3 + b*(c + x)*c^5/(c - x))*log(-(c + x)/(c - x)))/((c + x)^4/(c - x)^4 + 4*(c + x)^3/(c - x)^3 + 6*(c + x)^2/(c - x)^2 + 4*(c + x)/(c - x) + 1) + (2*b*c^5 + 6*a*(c + x)^3*c^5/(c - x)^3 + 3*b*(c + x)^3*c^5/(c - x)^3 + 6*b*(c + x)^2*c^5/(c - x)^2 + 6*a*(c + x)*c^5/(c - x) + 5*b*(c + x)*c^5/(c - x))/((c + x)^4/(c - x)^4 + 4*(c + x)^3/(c - x)^3 + 6*(c + x)^2/(c - x)^2 + 4*(c + x)/(c - x) + 1))/c`

Mupad [B] (verification not implemented)

Time = 3.62 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.90

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{a x^4}{4} - \frac{b c^4 \operatorname{atanh} \left(\frac{c}{x} \right)}{4} + \frac{b x^4 \operatorname{atanh} \left(\frac{c}{x} \right)}{4} + \frac{b c x^3}{12} + \frac{b c^3 x}{4}$$

input `int(x^3*(a + b*atanh(c/x)),x)`output `(a*x^4)/4 - (b*c^4*atanh(c/x))/4 + (b*x^4*atanh(c/x))/4 + (b*c*x^3)/12 + (b*c^3*x)/4`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.90

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = -\frac{\operatorname{atanh} \left(\frac{c}{x} \right) b c^4}{4} + \frac{\operatorname{atanh} \left(\frac{c}{x} \right) b x^4}{4} + \frac{a x^4}{4} + \frac{b c^3 x}{4} + \frac{b c x^3}{12}$$

input `int(x^3*(a+b*atanh(c/x)),x)`output `(- 3*atanh(c/x)*b*c**4 + 3*atanh(c/x)*b*x**4 + 3*a*x**4 + 3*b*c**3*x + b*c*x**3)/12`

3.136 $\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx$

Optimal result	1128
Mathematica [A] (verified)	1128
Rubi [A] (verified)	1129
Maple [A] (verified)	1131
Fricas [A] (verification not implemented)	1131
Sympy [A] (verification not implemented)	1132
Maxima [A] (verification not implemented)	1132
Giac [B] (verification not implemented)	1132
Mupad [B] (verification not implemented)	1133
Reduce [B] (verification not implemented)	1133

Optimal result

Integrand size = 14, antiderivative size = 45

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{1}{6} b c x^2 + \frac{1}{3} x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) + \frac{1}{6} b c^3 \log (c^2 - x^2)$$

output $1/6*b*c*x^2+1/3*x^3*(a+b*\operatorname{arctanh}(c/x))+1/6*b*c^3*\ln(c^2-x^2)$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.11

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{1}{6} b c x^2 + \frac{a x^3}{3} + \frac{1}{3} b x^3 \operatorname{arctanh} \left(\frac{c}{x} \right) + \frac{1}{6} b c^3 \log (-c^2 + x^2)$$

input $\operatorname{Integrate}[x^2*(a + b*\operatorname{ArcTanh}[c/x]),x]$

output $(b*c*x^2)/6 + (a*x^3)/3 + (b*x^3*\operatorname{ArcTanh}[c/x])/3 + (b*c^3*\operatorname{Log}[-c^2 + x^2])/6$

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.93, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {6452, 795, 243, 25, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right) \right) dx \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{3}bc \int \frac{x}{1 - \frac{c^2}{x^2}} dx + \frac{1}{3}x^3 \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right) \right) \\
 & \quad \downarrow \text{795} \\
 & \frac{1}{3}bc \int \frac{x^3}{x^2 - c^2} dx + \frac{1}{3}x^3 \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right) \right) \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{6}bc \int -\frac{x^2}{c^2 - x^2} dx^2 + \frac{1}{3}x^3 \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right) \right) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{3}x^3 \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right) \right) - \frac{1}{6}bc \int \frac{x^2}{c^2 - x^2} dx^2 \\
 & \quad \downarrow \text{49} \\
 & \frac{1}{3}x^3 \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right) \right) - \frac{1}{6}bc \int \left(\frac{c^2}{c^2 - x^2} - 1 \right) dx^2 \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{3}x^3 \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right) \right) + \frac{1}{6}bc(c^2 \log(c^2 - x^2) + x^2)
 \end{aligned}$$

input `Int[x^2*(a + b*ArcTanh[c/x]),x]`

output `(x^3*(a + b*ArcTanh[c/x]))/3 + (b*c*(x^2 + c^2*Log[c^2 - x^2]))/6`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 49 $\text{Int}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.)^{(\text{m}_.)} * ((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{(\text{n}_.)}), \text{x_Symbol}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(\text{a} + \text{b} * \text{x})^{\text{m}} * (\text{c} + \text{d} * \text{x})^{\text{n}}, \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{IGtQ}[\text{m}, 0] \&\& \text{IGtQ}[\text{m} + \text{n} + 2, 0]$
- rule 243 $\text{Int}[(\text{x}_.)^{(\text{m}_.)} * ((\text{a}_.) + (\text{b}_.) * (\text{x}_.)^2)^{(\text{p}_.)}, \text{x_Symbol}] \rightarrow \text{Simp}[1/2 \quad \text{Subst}[\text{Int}[\text{x}^{(\text{m} - 1)/2} * (\text{a} + \text{b} * \text{x})^{\text{p}}, \text{x}], \text{x}, \text{x}^2], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{m}, \text{p}\}, \text{x}] \&\& \text{IntegerQ}[(\text{m} - 1)/2]$
- rule 795 $\text{Int}[(\text{x}_.)^{(\text{m}_.)} * ((\text{a}_.) + (\text{b}_.) * (\text{x}_.)^{(\text{n}_.)})^{(\text{p}_.)}, \text{x_Symbol}] \rightarrow \text{Int}[\text{x}^{(\text{m} + \text{n} * \text{p})} * (\text{b} + \text{a} / \text{x}^{\text{n}})^{\text{p}}, \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{m}, \text{n}\}, \text{x}] \&\& \text{IntegerQ}[\text{p}] \&\& \text{NegQ}[\text{n}]$
- rule 2009 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Simp}[\text{IntSum}[\text{u}, \text{x}], \text{x}] \text{ ; SumQ}[\text{u}]$
- rule 6452 $\text{Int}[(\text{a}_.) + \text{ArcTanh}[(\text{c}_.) * (\text{x}_.)^{(\text{n}_.)}] * (\text{b}_.)^{(\text{p}_.)} * (\text{x}_.)^{(\text{m}_.)}, \text{x_Symbol}] \rightarrow \text{Simp}[\text{x}^{(\text{m} + 1)} * ((\text{a} + \text{b} * \text{ArcTanh}[\text{c} * \text{x}^{\text{n}}])^{\text{p}} / (\text{m} + 1)), \text{x}] - \text{Simp}[\text{b} * \text{c} * \text{n} * (\text{p} / (\text{m} + 1)) \quad \text{Int}[\text{x}^{(\text{m} + \text{n})} * ((\text{a} + \text{b} * \text{ArcTanh}[\text{c} * \text{x}^{\text{n}}])^{(\text{p} - 1)} / (1 - \text{c}^2 * \text{x}^{(2 * \text{n})})), \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{m}, \text{n}\}, \text{x}] \&\& \text{IGtQ}[\text{p}, 0] \&\& (\text{EqQ}[\text{p}, 1] \text{ || } (\text{EqQ}[\text{n}, 1] \&\& \text{IntegerQ}[\text{m}])) \&\& \text{NeQ}[\text{m}, -1]$

Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.27

method	result
parallelrisc	$\frac{\ln(x-c)bc^3}{3} + \frac{bx^3 \operatorname{arctanh}\left(\frac{c}{x}\right)}{3} + \frac{\operatorname{arctanh}\left(\frac{c}{x}\right)bc^3}{3} + \frac{ax^3}{3} + \frac{bcx^2}{6} + \frac{bc^3}{6}$
parts	$\frac{ax^3}{3} - bc^3 \left(-\frac{x^3 \operatorname{arctanh}\left(\frac{c}{x}\right)}{3c^3} - \frac{\ln\left(\frac{c}{x}-1\right)}{6} - \frac{\ln\left(1+\frac{c}{x}\right)}{6} - \frac{x^2}{6c^2} + \frac{\ln\left(\frac{c}{x}\right)}{3} \right)$
derivativedivides	$-c^3 \left(-\frac{ax^3}{3c^3} + b \left(-\frac{x^3 \operatorname{arctanh}\left(\frac{c}{x}\right)}{3c^3} - \frac{\ln\left(\frac{c}{x}-1\right)}{6} - \frac{\ln\left(1+\frac{c}{x}\right)}{6} - \frac{x^2}{6c^2} + \frac{\ln\left(\frac{c}{x}\right)}{3} \right) \right)$
default	$-c^3 \left(-\frac{ax^3}{3c^3} + b \left(-\frac{x^3 \operatorname{arctanh}\left(\frac{c}{x}\right)}{3c^3} - \frac{\ln\left(\frac{c}{x}-1\right)}{6} - \frac{\ln\left(1+\frac{c}{x}\right)}{6} - \frac{x^2}{6c^2} + \frac{\ln\left(\frac{c}{x}\right)}{3} \right) \right)$
risc	$\frac{bx^3 \ln(x+c)}{6} - \frac{bx^3 \ln(c-x)}{6} - \frac{i\pi b x^3 \operatorname{csgn}\left(\frac{i(x+c)}{x}\right)^3}{12} + \frac{i\pi b x^3 \operatorname{csgn}\left(\frac{i}{x}\right) \operatorname{csgn}(i(c-x)) \operatorname{csgn}\left(\frac{i(c-x)}{x}\right)}{12} - \frac{i\pi b x^3}{6}$

input `int(x^2*(a+b*arctanh(c/x)),x,method=_RETURNVERBOSE)`output $\frac{1}{3}*\ln(x-c)*b*c^3 + \frac{1}{3}*b*x^3*\operatorname{arctanh}(c/x) + \frac{1}{3}*\operatorname{arctanh}(c/x)*b*c^3 + \frac{1}{3}*a*x^3 + \frac{1}{6}*b*c*x^2 + \frac{1}{6}*b*c^3$ **Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.09

$$\int x^2 \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right) dx = \frac{1}{6} bc^3 \log(-c^2 + x^2) + \frac{1}{6} bx^3 \log\left(-\frac{c+x}{c-x}\right) + \frac{1}{6} bcx^2 + \frac{1}{3} ax^3$$

input `integrate(x^2*(a+b*arctanh(c/x)),x, algorithm="fricas")`output $\frac{1}{6}*b*c^3*\log(-c^2 + x^2) + \frac{1}{6}*b*x^3*\log(-(c+x)/(c-x)) + \frac{1}{6}*b*c*x^2 + \frac{1}{3}*a*x^3$

Sympy [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.09

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{ax^3}{3} + \frac{bc^3 \log(-c+x)}{3} + \frac{bc^3 \operatorname{atanh} \left(\frac{c}{x} \right)}{3} + \frac{bcx^2}{6} + \frac{bx^3 \operatorname{atanh} \left(\frac{c}{x} \right)}{3}$$

input `integrate(x**2*(a+b*atanh(c/x)),x)`

output `a*x**3/3 + b*c**3*log(-c + x)/3 + b*c**3*atanh(c/x)/3 + b*c*x**2/6 + b*x**3*atanh(c/x)/3`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.93

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{1}{3} ax^3 + \frac{1}{6} \left(2x^3 \operatorname{artanh} \left(\frac{c}{x} \right) + (c^2 \log(-c^2 + x^2) + x^2)c \right) b$$

input `integrate(x^2*(a+b*arctanh(c/x)),x, algorithm="maxima")`

output `1/3*a*x^3 + 1/6*(2*x^3*arctanh(c/x) + (c^2*log(-c^2 + x^2) + x^2)*c)*b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 227 vs. 2(39) = 78.

Time = 0.12 (sec) , antiderivative size = 227, normalized size of antiderivative = 5.04

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{bc^4 \log \left(-\frac{c+x}{c-x} - 1 \right) - bc^4 \log \left(-\frac{c+x}{c-x} \right) + \frac{\left(bc^4 + \frac{3b(c+x)^2 c^4}{(c-x)^2} \right) \log \left(-\frac{c+x}{c-x} \right)}{\frac{(c+x)^3}{(c-x)^3} + \frac{3(c+x)^2}{(c-x)^2} + \frac{3(c+x)}{c-x} + 1} + \frac{2 \left(ac^4 + \frac{3a(c+x)^2 c^4}{(c-x)^2} + \frac{b(c+x)^2 c^4}{(c-x)^2} + \frac{b(c+x)c^4}{c-x} \right)}{\frac{(c+x)^3}{(c-x)^3} + \frac{3(c+x)^2}{(c-x)^2} + \frac{3(c+x)}{c-x} + 1}}{3c}$$

input `integrate(x^2*(a+b*arctanh(c/x)),x, algorithm="giac")`

output `-1/3*(b*c^4*log(-(c + x)/(c - x) - 1) - b*c^4*log(-(c + x)/(c - x)) + (b*c^4 + 3*b*(c + x)^2*c^4/(c - x)^2)*log(-(c + x)/(c - x))/((c + x)^3/(c - x)^3 + 3*(c + x)^2/(c - x)^2 + 3*(c + x)/(c - x) + 1) + 2*(a*c^4 + 3*a*(c + x)^2*c^4/(c - x)^2 + b*(c + x)^2*c^4/(c - x)^2 + b*(c + x)*c^4/(c - x))/((c + x)^3/(c - x)^3 + 3*(c + x)^2/(c - x)^2 + 3*(c + x)/(c - x) + 1))/c`

Mupad [B] (verification not implemented)

Time = 3.44 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.93

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{a x^3}{3} + \frac{b c^3 \ln(x^2 - c^2)}{6} + \frac{b x^3 \operatorname{atanh} \left(\frac{c}{x} \right)}{3} + \frac{b c x^2}{6}$$

input `int(x^2*(a + b*atanh(c/x)),x)`

output `(a*x^3)/3 + (b*c^3*log(x^2 - c^2))/6 + (b*x^3*atanh(c/x))/3 + (b*c*x^2)/6`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.16

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = -\frac{\operatorname{atanh} \left(\frac{c}{x} \right) b c^3}{3} + \frac{\operatorname{atanh} \left(\frac{c}{x} \right) b x^3}{3} + \frac{\log(-c - x) b c^3}{3} + \frac{a x^3}{3} + \frac{b c x^2}{6}$$

input `int(x^2*(a+b*atanh(c/x)),x)`

output `(- 2*atanh(c/x)*b*c**3 + 2*atanh(c/x)*b*x**3 + 2*log(- c - x)*b*c**3 + 2*a*x**3 + b*c*x**2)/6`

3.137 $\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx$

Optimal result	1134
Mathematica [A] (verified)	1134
Rubi [A] (verified)	1135
Maple [A] (verified)	1136
Fricas [A] (verification not implemented)	1137
Sympy [A] (verification not implemented)	1138
Maxima [A] (verification not implemented)	1138
Giac [B] (verification not implemented)	1138
Mupad [B] (verification not implemented)	1139
Reduce [B] (verification not implemented)	1139

Optimal result

Integrand size = 12, antiderivative size = 39

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{bcx}{2} + \frac{1}{2}x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) - \frac{1}{2}bc^2 \operatorname{arctanh} \left(\frac{x}{c} \right)$$

output `1/2*b*c*x+1/2*x^2*(a+b*arctanh(c/x))-1/2*b*c^2*arctanh(x/c)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.44

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{bcx}{2} + \frac{ax^2}{2} + \frac{1}{2}bx^2 \operatorname{arctanh} \left(\frac{c}{x} \right) + \frac{1}{4}bc^2 \log(-c+x) - \frac{1}{4}bc^2 \log(c+x)$$

input `Integrate[x*(a + b*ArcTanh[c/x]),x]`

output `(b*c*x)/2 + (a*x^2)/2 + (b*x^2*ArcTanh[c/x])/2 + (b*c^2*Log[-c + x])/4 - (b*c^2*Log[c + x])/4`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.90, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6452, 772, 262, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{2} bc \int \frac{1}{1 - \frac{c^2}{x^2}} dx + \frac{1}{2} x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) \\
 & \quad \downarrow \text{772} \\
 & \frac{1}{2} bc \int \frac{x^2}{x^2 - c^2} dx + \frac{1}{2} x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) \\
 & \quad \downarrow \text{262} \\
 & \frac{1}{2} bc \left(c^2 \int \frac{1}{x^2 - c^2} dx + x \right) + \frac{1}{2} x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) \\
 & \quad \downarrow \text{220} \\
 & \frac{1}{2} x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) + \frac{1}{2} bc \left(x - c \operatorname{arctanh} \left(\frac{x}{c} \right) \right)
 \end{aligned}$$

input `Int[x*(a + b*ArcTanh[c/x]),x]`

output `(x^2*(a + b*ArcTanh[c/x]))/2 + (b*c*(x - c*ArcTanh[x/c]))/2`

Definitions of rubi rules used

rule 220 $\text{Int}[(a_ + (b_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[b, 2])^{-1} \cdot \text{ArcTanh}[\text{Rt}[b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 262 $\text{Int}[(c_ \cdot)(x_)^{m_} \cdot (a_ + (b_ \cdot)(x_)^2)^{p_}], x_Symbol] \rightarrow \text{Simp}[c \cdot (c \cdot x)^{m-1} \cdot ((a + b \cdot x^2)^{p+1} / (b \cdot (m + 2 \cdot p + 1))), x] - \text{Simp}[a \cdot c^2 \cdot ((m-1) / (b \cdot (m + 2 \cdot p + 1))) \ \text{Int}[(c \cdot x)^{m-2} \cdot (a + b \cdot x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{GtQ}[m, 2 - 1] \ \&\& \ \text{NeQ}[m + 2 \cdot p + 1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 772 $\text{Int}[(a_ + (b_ \cdot)(x_)^{n_})^{p_}], x_Symbol] \rightarrow \text{Int}[x^{n \cdot p} \cdot (b + a/x^n)^p, x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{ILtQ}[n, 0] \ \&\& \ \text{IntegerQ}[p]$

rule 6452 $\text{Int}[(a_ + \text{ArcTanh}[c_ \cdot)(x_)^{n_}] \cdot (b_)^{p_} \cdot (x_)^{m_}], x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot ((a + b \cdot \text{ArcTanh}[c \cdot x^n])^p / (m+1)), x] - \text{Simp}[b \cdot c \cdot n \cdot (p / (m+1)) \ \text{Int}[x^{m+n} \cdot ((a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2 \cdot n}))], x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.10

method	result
parallelrisch	$\frac{\operatorname{arctanh}\left(\frac{c}{x}\right) b x^2}{2} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) b c^2}{2} + \frac{a x^2}{2} + \frac{b c x}{2} + \frac{a c^2}{2}$
parts	$\frac{a x^2}{2} - b c^2 \left(-\frac{x^2 \operatorname{arctanh}\left(\frac{c}{x}\right)}{2 c^2} - \frac{\ln\left(\frac{c}{x}-1\right)}{4} - \frac{x}{2 c} + \frac{\ln\left(1+\frac{c}{x}\right)}{4} \right)$
derivativedivides	$-c^2 \left(-\frac{a x^2}{2 c^2} + b \left(-\frac{x^2 \operatorname{arctanh}\left(\frac{c}{x}\right)}{2 c^2} - \frac{\ln\left(\frac{c}{x}-1\right)}{4} - \frac{x}{2 c} + \frac{\ln\left(1+\frac{c}{x}\right)}{4} \right) \right)$
default	$-c^2 \left(-\frac{a x^2}{2 c^2} + b \left(-\frac{x^2 \operatorname{arctanh}\left(\frac{c}{x}\right)}{2 c^2} - \frac{\ln\left(\frac{c}{x}-1\right)}{4} - \frac{x}{2 c} + \frac{\ln\left(1+\frac{c}{x}\right)}{4} \right) \right)$
oring	$-(c^2 - x^2) \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right) + \frac{(c-x)(x+c) \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) - \frac{b c}{x \left(1 - \frac{c^2}{x^2} \right)} \right)}{2}$
risch	$\frac{x^2 b \ln(x+c)}{4} - \frac{x^2 b \ln(c-x)}{4} + \frac{i \pi b x^2 \operatorname{csgn}\left(\frac{i}{x}\right) \operatorname{csgn}(i(c-x)) \operatorname{csgn}\left(\frac{i(c-x)}{x}\right)}{8} - \frac{i \pi b x^2 \operatorname{csgn}\left(\frac{i(x+c)}{x}\right)^3}{8} - \frac{i \pi b x^2 \operatorname{csgn}\left(\frac{i(x+c)}{x}\right)}{8}$

input

```
int(x*(a+b*arctanh(c/x)),x,method=_RETURNVERBOSE)
```

output

```
1/2*arctanh(c/x)*b*x^2-1/2*arctanh(c/x)*b*c^2+1/2*a*x^2+1/2*b*c*x+1/2*a*c^2
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

$$\int x \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right) dx = \frac{1}{2} b c x + \frac{1}{2} a x^2 - \frac{1}{4} (b c^2 - b x^2) \log\left(-\frac{c+x}{c-x}\right)$$

input

```
integrate(x*(a+b*arctanh(c/x)),x, algorithm="fricas")
```

output

```
1/2*b*c*x + 1/2*a*x^2 - 1/4*(b*c^2 - b*x^2)*log(-(c + x)/(c - x))
```

Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.92

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{ax^2}{2} - \frac{bc^2 \operatorname{atanh} \left(\frac{c}{x} \right)}{2} + \frac{bcx}{2} + \frac{bx^2 \operatorname{atanh} \left(\frac{c}{x} \right)}{2}$$

input `integrate(x*(a+b*atanh(c/x)),x)`

output `a*x**2/2 - b*c**2*atanh(c/x)/2 + b*c*x/2 + b*x**2*atanh(c/x)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.13

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx \\ = \frac{1}{2} ax^2 + \frac{1}{4} \left(2x^2 \operatorname{artanh} \left(\frac{c}{x} \right) - (c \log(c+x) - c \log(-c+x) - 2x)c \right) b$$

input `integrate(x*(a+b*arctanh(c/x)),x, algorithm="maxima")`

output `1/2*a*x^2 + 1/4*(2*x^2*arctanh(c/x) - (c*log(c + x) - c*log(-c + x) - 2*x)*c)*b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 130 vs. 2(33) = 66.

Time = 0.12 (sec) , antiderivative size = 130, normalized size of antiderivative = 3.33

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = - \frac{b(c+x)c^3 \log \left(-\frac{c+x}{c-x} \right)}{(c-x) \left(\frac{(c+x)^2}{(c-x)^2} + \frac{2(c+x)}{c-x} + 1 \right)} + \frac{bc^3 + \frac{2a(c+x)c^3}{c-x} + \frac{b(c+x)c^3}{c-x}}{\frac{(c+x)^2}{(c-x)^2} + \frac{2(c+x)}{c-x} + 1}$$

input `integrate(x*(a+b*arctanh(c/x)),x, algorithm="giac")`

output

$$\frac{-(b(c+x)c^3 \log(-(c+x)/(c-x)))/((c-x)((c+x)^2/(c-x)^2 + 2(c+x)/(c-x) + 1)) + (b^2c^3 + 2a(c+x)c^3/(c-x) + b^2(c+x)c^3/(c-x))/((c+x)^2/(c-x)^2 + 2(c+x)/(c-x) + 1))/c$$

Mupad [B] (verification not implemented)

Time = 3.58 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.92

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{ax^2}{2} - \frac{bc^2 \operatorname{atanh} \left(\frac{c}{x} \right)}{2} + \frac{bx^2 \operatorname{atanh} \left(\frac{c}{x} \right)}{2} + \frac{bcx}{2}$$

input

int(x*(a + b*atanh(c/x)),x)

output

$$(a*x^2)/2 - (b*c^2*atanh(c/x))/2 + (b*x^2*atanh(c/x))/2 + (b*c*x)/2$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.92

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = -\frac{\operatorname{atanh} \left(\frac{c}{x} \right) bc^2}{2} + \frac{\operatorname{atanh} \left(\frac{c}{x} \right) bx^2}{2} + \frac{ax^2}{2} + \frac{bcx}{2}$$

input

int(x*(a+b*atanh(c/x)),x)

output

$$(-\operatorname{atanh}(c/x)*b*c**2 + \operatorname{atanh}(c/x)*b*x**2 + a*x**2 + b*c*x)/2$$

3.138 $\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx$

Optimal result	1140
Mathematica [A] (verified)	1140
Rubi [A] (verified)	1141
Maple [A] (verified)	1142
Fricas [A] (verification not implemented)	1142
Sympy [A] (verification not implemented)	1143
Maxima [A] (verification not implemented)	1143
Giac [B] (verification not implemented)	1143
Mupad [B] (verification not implemented)	1144
Reduce [B] (verification not implemented)	1144

Optimal result

Integrand size = 10, antiderivative size = 29

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = ax + b \operatorname{arctanh} \left(\frac{c}{x} \right) + \frac{1}{2} bc \log (c^2 - x^2)$$

output `a*x+b*x*arctanh(c/x)+1/2*b*c*ln(c^2-x^2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = ax + b \operatorname{arctanh} \left(\frac{c}{x} \right) + \frac{1}{2} bc \log (c^2 - x^2)$$

input `Integrate[a + b*ArcTanh[c/x],x]`

output `a*x + b*x*ArcTanh[c/x] + (b*c*Log[c^2 - x^2])/2`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right) dx$$

$$\downarrow \text{2009}$$

$$ax + b \operatorname{arctanh}\left(\frac{c}{x}\right) + \frac{1}{2}bc \log(c^2 - x^2)$$

input

```
Int[a + b*ArcTanh[c/x],x]
```

output

```
a*x + b*x*ArcTanh[c/x] + (b*c*Log[c^2 - x^2])/2
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.10

method	result
parallelrisch	$b(\ln(x-c)c + \operatorname{arctanh}\left(\frac{c}{x}\right)x + \operatorname{arctanh}\left(\frac{c}{x}\right)c) + ax$
default	$ax - bc\left(-\frac{x \operatorname{arctanh}\left(\frac{c}{x}\right)}{c} - \frac{\ln\left(\frac{c}{x}-1\right)}{2} - \frac{\ln\left(1+\frac{c}{x}\right)}{2} + \ln\left(\frac{c}{x}\right)\right)$
parts	$ax - bc\left(-\frac{x \operatorname{arctanh}\left(\frac{c}{x}\right)}{c} - \frac{\ln\left(\frac{c}{x}-1\right)}{2} - \frac{\ln\left(1+\frac{c}{x}\right)}{2} + \ln\left(\frac{c}{x}\right)\right)$
derivativedivides	$-c\left(-\frac{ax}{c} + b\left(-\frac{x \operatorname{arctanh}\left(\frac{c}{x}\right)}{c} - \frac{\ln\left(\frac{c}{x}-1\right)}{2} - \frac{\ln\left(1+\frac{c}{x}\right)}{2} + \ln\left(\frac{c}{x}\right)\right)\right)$
risch	$ax + \frac{bx \ln(x+c)}{2} - \frac{b \ln(c-x)x}{2} - \frac{ib\pi x}{2} + \frac{ib\pi \operatorname{csgn}(i(x+c)) \operatorname{csgn}\left(\frac{i(x+c)}{x}\right)^2 x}{4} + \frac{ib\pi \operatorname{csgn}\left(\frac{i(c-x)}{x}\right)^2 x}{2} - \frac{ib\pi \operatorname{csgn}\left(\frac{i(c-x)}{x}\right)^2 x}{2}$

input `int(a+b*arctanh(c/x),x,method=_RETURNVERBOSE)`output `b*(ln(x-c)*c+arctanh(c/x)*x+arctanh(c/x)*c)+a*x`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.21

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right)\right) dx = \frac{1}{2} bc \log(-c^2 + x^2) + \frac{1}{2} bx \log\left(-\frac{c+x}{c-x}\right) + ax$$

input `integrate(a+b*arctanh(c/x),x, algorithm="fricas")`output `1/2*b*c*log(-c^2 + x^2) + 1/2*b*x*log(-(c + x)/(c - x)) + a*x`

Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = ax + b \left(c \log(-c + x) + c \operatorname{atanh} \left(\frac{c}{x} \right) + x \operatorname{atanh} \left(\frac{c}{x} \right) \right)$$

input `integrate(a+b*atanh(c/x),x)`

output `a*x + b*(c*log(-c + x) + c*atanh(c/x) + x*atanh(c/x))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx = \frac{1}{2} \left(2x \operatorname{arctanh} \left(\frac{c}{x} \right) + c \log(-c^2 + x^2) \right) b + ax$$

input `integrate(a+b*arctanh(c/x),x, algorithm="maxima")`

output `1/2*(2*x*arctanh(c/x) + c*log(-c^2 + x^2))*b + a*x`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 150 vs. 2(27) = 54.

Time = 0.12 (sec) , antiderivative size = 150, normalized size of antiderivative = 5.17

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right) dx$$

$$= ax + \frac{c^2 \left(\log \left(\frac{|-c-x|}{|c-x|} \right) - \log \left(\left| -\frac{c+x}{c-x} - 1 \right| \right) \right) - \frac{c^2 \log \left(-\frac{\frac{c}{c-x} + \frac{1}{c}}{\frac{c}{c-x} - 1} + 1 \right)}{\frac{\frac{c+x}{c-x} - 1}{\frac{c+x}{c-x} + 1}}}{c} b$$

input `integrate(a+b*arctanh(c/x),x, algorithm="giac")`

output `a*x + (c^2*(log(abs(-c - x)/abs(c - x)) - log(abs(-(c + x)/(c - x) - 1))) - c^2*log(-(c*((c + x)/((c - x)*c) + 1/c)/((c + x)/(c - x) - 1) + 1)/(c*((c + x)/((c - x)*c) + 1/c)/((c + x)/(c - x) - 1) - 1))/((c + x)/(c - x) + 1))*b/c`

Mupad [B] (verification not implemented)

Time = 3.43 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right) dx = a x + b x \operatorname{atanh}\left(\frac{c}{x}\right) + \frac{b c \ln(x^2 - c^2)}{2}$$

input `int(a + b*atanh(c/x),x)`

output `a*x + b*x*atanh(c/x) + (b*c*log(x^2 - c^2))/2`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.17

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right) dx = -\operatorname{atanh}\left(\frac{c}{x}\right) b c + \operatorname{atanh}\left(\frac{c}{x}\right) b x + \log(-c - x) b c + a x$$

input `int(a+b*atanh(c/x),x)`

output `- atanh(c/x)*b*c + atanh(c/x)*b*x + log(- c - x)*b*c + a*x`

3.139 $\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} dx$

Optimal result	1145
Mathematica [A] (verified)	1145
Rubi [A] (verified)	1146
Maple [B] (verified)	1147
Fricas [F]	1147
Sympy [F]	1148
Maxima [F]	1148
Giac [F]	1148
Mupad [F(-1)]	1149
Reduce [F]	1149

Optimal result

Integrand size = 14, antiderivative size = 30

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} dx = a \log(x) + \frac{1}{2} b \operatorname{PolyLog}\left(2, -\frac{c}{x}\right) - \frac{1}{2} b \operatorname{PolyLog}\left(2, \frac{c}{x}\right)$$

output `a*ln(x)+1/2*b*polylog(2,-c/x)-1/2*b*polylog(2,c/x)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} dx = a \log(x) + \frac{1}{2} b \left(\operatorname{PolyLog}\left(2, -\frac{c}{x}\right) - \operatorname{PolyLog}\left(2, \frac{c}{x}\right) \right)$$

input `Integrate[(a + b*ArcTanh[c/x])/x,x]`

output `a*Log[x] + (b*(PolyLog[2, -(c/x)] - PolyLog[2, c/x]))/2`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.10, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6450, 6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} dx$$

↓ 6450

$$- \int x \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right) d\frac{1}{x}$$

↓ 6446

$$-a \log\left(\frac{1}{x}\right) + \frac{1}{2}b \operatorname{PolyLog}\left(2, -\frac{c}{x}\right) - \frac{1}{2}b \operatorname{PolyLog}\left(2, \frac{c}{x}\right)$$

input `Int[(a + b*ArcTanh[c/x])/x,x]`

output `-(a*Log[x^(-1)]) + (b*PolyLog[2, -(c/x)])/2 - (b*PolyLog[2, c/x])/2`

Defintions of rubi rules used

rule 6446 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))/(x_), x_Symbol] := Simp[a*Log[x], x] + (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) / ; FreeQ[{a, b, c}, x]`

rule 6450 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)]*(b_.))^(p_.)/(x_), x_Symbol] := Simp[1/n Subst[Int[(a + b*ArcTanh[c*x])^p/x, x], x, x^n], x] / ; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. $2(26) = 52$.

Time = 0.40 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.90

method	result
parts	$a \ln(x) + b \left(-\ln\left(\frac{c}{x}\right) \operatorname{arctanh}\left(\frac{c}{x}\right) + \frac{\operatorname{dilog}\left(\frac{c}{x}\right)}{2} + \frac{\operatorname{dilog}\left(1+\frac{c}{x}\right)}{2} + \frac{\ln\left(\frac{c}{x}\right) \ln\left(1+\frac{c}{x}\right)}{2} \right)$
derivativedivides	$-a \ln\left(\frac{c}{x}\right) - b \left(\ln\left(\frac{c}{x}\right) \operatorname{arctanh}\left(\frac{c}{x}\right) - \frac{\operatorname{dilog}\left(\frac{c}{x}\right)}{2} - \frac{\operatorname{dilog}\left(1+\frac{c}{x}\right)}{2} - \frac{\ln\left(\frac{c}{x}\right) \ln\left(1+\frac{c}{x}\right)}{2} \right)$
default	$-a \ln\left(\frac{c}{x}\right) - b \left(\ln\left(\frac{c}{x}\right) \operatorname{arctanh}\left(\frac{c}{x}\right) - \frac{\operatorname{dilog}\left(\frac{c}{x}\right)}{2} - \frac{\operatorname{dilog}\left(1+\frac{c}{x}\right)}{2} - \frac{\ln\left(\frac{c}{x}\right) \ln\left(1+\frac{c}{x}\right)}{2} \right)$
risch	$\frac{b \ln(x) \ln(x+c)}{2} - \frac{\left(-2ib\pi \operatorname{csgn}\left(\frac{i(c-x)}{x}\right)^2 - ib\pi \operatorname{csgn}\left(\frac{i}{x}\right) \operatorname{csgn}(i(c-x)) \operatorname{csgn}\left(\frac{i(c-x)}{x}\right) + ib\pi \operatorname{csgn}\left(\frac{i}{x}\right) \operatorname{csgn}(i(x+c)) \operatorname{csgn}\left(\frac{i(c-x)}{x}\right) \right)}{2}$

input `int((a+b*arctanh(c/x))/x,x,method=_RETURNVERBOSE)`

output `a*ln(x)+b*(-ln(c/x)*arctanh(c/x)+1/2*dilog(c/x)+1/2*dilog(1+c/x)+1/2*ln(c/x)*ln(1+c/x))`

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} dx = \int \frac{b \operatorname{arctanh}\left(\frac{c}{x}\right) + a}{x} dx$$

input `integrate((a+b*arctanh(c/x))/x,x, algorithm="fricas")`

output `integral((b*arctanh(c/x) + a)/x, x)`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} dx = \int \frac{a + b \operatorname{atanh}\left(\frac{c}{x}\right)}{x} dx$$

input `integrate((a+b*atanh(c/x))/x,x)`

output `Integral((a + b*atanh(c/x))/x, x)`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} dx = \int \frac{b \operatorname{artanh}\left(\frac{c}{x}\right) + a}{x} dx$$

input `integrate((a+b*arctanh(c/x))/x,x, algorithm="maxima")`

output `1/2*b*integrate((log(c/x + 1) - log(-c/x + 1))/x, x) + a*log(x)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} dx = \int \frac{b \operatorname{artanh}\left(\frac{c}{x}\right) + a}{x} dx$$

input `integrate((a+b*arctanh(c/x))/x,x, algorithm="giac")`

output `integrate((b*arctanh(c/x) + a)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} dx = \int \frac{a + b \operatorname{atanh}\left(\frac{c}{x}\right)}{x} dx$$

input `int((a + b*atanh(c/x))/x,x)`output `int((a + b*atanh(c/x))/x, x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} dx = \left(\int \frac{\operatorname{atanh}\left(\frac{c}{x}\right)}{x} dx \right) b + \log(x) a$$

input `int((a+b*atanh(c/x))/x,x)`output `int(atanh(c/x)/x,x)*b + log(x)*a`

3.140 $\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^2} dx$

Optimal result	1150
Mathematica [A] (verified)	1150
Rubi [A] (verified)	1151
Maple [A] (verified)	1152
Fricas [A] (verification not implemented)	1152
Sympy [A] (verification not implemented)	1153
Maxima [A] (verification not implemented)	1153
Giac [B] (verification not implemented)	1153
Mupad [B] (verification not implemented)	1154
Reduce [B] (verification not implemented)	1154

Optimal result

Integrand size = 14, antiderivative size = 35

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^2} dx = -\frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} - \frac{b \log\left(1 - \frac{c^2}{x^2}\right)}{2c}$$

output $-(a+b*\operatorname{arctanh}(c/x))/x-1/2*b*\ln(1-c^2/x^2)/c$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.09

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^2} dx = -\frac{a}{x} - \frac{b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} - \frac{b \log\left(1 - \frac{c^2}{x^2}\right)}{2c}$$

input `Integrate[(a + b*ArcTanh[c/x])/x^2,x]`

output $-(a/x) - (b*\operatorname{ArcTanh}[c/x])/x - (b*\operatorname{Log}[1 - c^2/x^2])/(2*c)$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6452, 792}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^2} dx$$

$$\downarrow 6452$$

$$-bc \int \frac{1}{\left(1 - \frac{c^2}{x^2}\right) x^3} dx - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x}$$

$$\downarrow 792$$

$$-\frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} - \frac{b \log\left(1 - \frac{c^2}{x^2}\right)}{2c}$$

input `Int[(a + b*ArcTanh[c/x])/x^2,x]`

output `-((a + b*ArcTanh[c/x])/x) - (b*Log[1 - c^2/x^2])/(2*c)`

Defintions of rubi rules used

rule 792 `Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_.)), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

method	result
parts	$-\frac{a}{x} - \frac{b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} - \frac{b \ln\left(1 - \frac{c^2}{x^2}\right)}{2c}$
derivativedivides	$-\frac{\frac{ca}{x} + b \left(\frac{c \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} + \frac{\ln\left(1 - \frac{c^2}{x^2}\right)}{2} \right)}{c}$
default	$-\frac{\frac{ca}{x} + b \left(\frac{c \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} + \frac{\ln\left(1 - \frac{c^2}{x^2}\right)}{2} \right)}{c}$
parallelrisc	$\frac{\ln(x)xb - \ln(x-c)xb - bx \operatorname{arctanh}\left(\frac{c}{x}\right) - \operatorname{arctanh}\left(\frac{c}{x}\right)bc - ac}{xc}$
risc	$-\frac{b \ln(x+c)}{2x} + \frac{-i\pi bc \operatorname{csgn}(i(x+c)) \operatorname{csgn}\left(\frac{i(x+c)}{x}\right)^2 - i\pi bc \operatorname{csgn}\left(\frac{i}{x}\right) \operatorname{csgn}(i(c-x)) \operatorname{csgn}\left(\frac{i(c-x)}{x}\right) + i\pi bc \operatorname{csgn}\left(\frac{i(x+c)}{x}\right)^3}{2x}$

input `int((a+b*arctanh(c/x))/x^2,x,method=_RETURNVERBOSE)`

output `-a/x-b/x*arctanh(c/x)-1/2*b*ln(1-c^2/x^2)/c`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.37

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^2} dx = -\frac{bx \log(-c^2 + x^2) - 2bx \log(x) + bc \log\left(-\frac{c+x}{c-x}\right) + 2ac}{2cx}$$

input `integrate((a+b*arctanh(c/x))/x^2,x, algorithm="fricas")`

output `-1/2*(b*x*log(-c^2 + x^2) - 2*b*x*log(x) + b*c*log(-(c + x)/(c - x)) + 2*a*c)/(c*x)`

Sympy [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.11

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^2} dx = \begin{cases} -\frac{a}{x} - \frac{b \operatorname{atanh}\left(\frac{c}{x}\right)}{x} + \frac{b \log(x)}{c} - \frac{b \log(-c+x)}{c} - \frac{b \operatorname{atanh}\left(\frac{c}{x}\right)}{c} & \text{for } c \neq 0 \\ -\frac{a}{x} & \text{otherwise} \end{cases}$$

input `integrate((a+b*atanh(c/x))/x**2,x)`

output `Piecewise((-a/x - b*atanh(c/x)/x + b*log(x)/c - b*log(-c + x)/c - b*atanh(c/x)/c, Ne(c, 0)), (-a/x, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^2} dx = -\frac{b \left(\frac{2c \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} + \log\left(-\frac{c^2}{x^2} + 1\right) \right)}{2c} - \frac{a}{x}$$

input `integrate((a+b*arctanh(c/x))/x^2,x, algorithm="maxima")`

output `-1/2*b*(2*c*arctanh(c/x)/x + log(-c^2/x^2 + 1))/c - a/x`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 87 vs. 2(33) = 66.

Time = 0.12 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.49

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^2} dx = \frac{b \log\left(-\frac{c+x}{c-x} + 1\right) - b \log\left(-\frac{c+x}{c-x}\right) - \frac{b \log\left(-\frac{c+x}{c-x}\right)}{\frac{c+x}{c-x} - 1} - \frac{2a}{\frac{c+x}{c-x} - 1}}{c}$$

input `integrate((a+b*arctanh(c/x))/x^2,x, algorithm="giac")`

output $(b \cdot \log(-(c + x)/(c - x) + 1) - b \cdot \log(-(c + x)/(c - x)) - b \cdot \log(-(c + x)/(c - x)))/((c + x)/(c - x) - 1) - 2 \cdot a/((c + x)/(c - x) - 1))/c$

Mupad [B] (verification not implemented)

Time = 3.50 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.23

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^2} dx = \frac{b x \ln(x) - \frac{b x \ln(x^2 - c^2)}{2}}{c x} - \frac{a + b \operatorname{atanh}\left(\frac{c}{x}\right)}{x}$$

input `int((a + b*atanh(c/x))/x^2,x)`

output $(b \cdot x \cdot \log(x) - (b \cdot x \cdot \log(x^2 - c^2))/2)/(c \cdot x) - (a + b \cdot \operatorname{atanh}(c/x))/x$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.37

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^2} dx = \frac{-\operatorname{atanh}\left(\frac{c}{x}\right) b c + \operatorname{atanh}\left(\frac{c}{x}\right) b x - \log(-c - x) b x + \log(x) b x - a c}{c x}$$

input `int((a+b*atanh(c/x))/x^2,x)`

output $(- \operatorname{atanh}(c/x) \cdot b \cdot c + \operatorname{atanh}(c/x) \cdot b \cdot x - \log(-c - x) \cdot b \cdot x + \log(x) \cdot b \cdot x - a \cdot c) / (c \cdot x)$

3.141 $\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^3} dx$

Optimal result	1155
Mathematica [A] (verified)	1155
Rubi [A] (verified)	1156
Maple [A] (verified)	1157
Fricas [A] (verification not implemented)	1158
Sympy [A] (verification not implemented)	1159
Maxima [A] (verification not implemented)	1159
Giac [B] (verification not implemented)	1159
Mupad [B] (verification not implemented)	1160
Reduce [B] (verification not implemented)	1160

Optimal result

Integrand size = 14, antiderivative size = 43

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^3} dx = -\frac{b}{2cx} - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{2x^2} + \frac{b \operatorname{arctanh}\left(\frac{x}{c}\right)}{2c^2}$$

output

$$-1/2*b/c/x - 1/2*(a + b*\operatorname{arctanh}(c/x))/x^2 + 1/2*b*\operatorname{arctanh}(x/c)/c^2$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.40

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^3} dx = -\frac{a}{2x^2} - \frac{b}{2cx} - \frac{b \operatorname{arctanh}\left(\frac{c}{x}\right)}{2x^2} - \frac{b \log(-c + x)}{4c^2} + \frac{b \log(c + x)}{4c^2}$$

input

```
Integrate[(a + b*ArcTanh[c/x])/x^3, x]
```

output

$$-1/2*a/x^2 - b/(2*c*x) - (b*ArcTanh[c/x])/(2*x^2) - (b*Log[-c + x])/(4*c^2) + (b*Log[c + x])/(4*c^2)$$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6452, 795, 264, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^3} dx \\
 & \quad \downarrow \text{6452} \\
 & -\frac{1}{2}bc \int \frac{1}{\left(1 - \frac{c^2}{x^2}\right)x^4} dx - \frac{a + \operatorname{arctanh}\left(\frac{c}{x}\right)}{2x^2} \\
 & \quad \downarrow \text{795} \\
 & -\frac{1}{2}bc \int \frac{1}{x^2(x^2 - c^2)} dx - \frac{a + \operatorname{arctanh}\left(\frac{c}{x}\right)}{2x^2} \\
 & \quad \downarrow \text{264} \\
 & -\frac{1}{2}bc \left(\frac{\int \frac{1}{x^2 - c^2} dx}{c^2} + \frac{1}{c^2 x} \right) - \frac{a + \operatorname{arctanh}\left(\frac{c}{x}\right)}{2x^2} \\
 & \quad \downarrow \text{220} \\
 & -\frac{a + \operatorname{arctanh}\left(\frac{c}{x}\right)}{2x^2} - \frac{1}{2}bc \left(\frac{1}{c^2 x} - \frac{\operatorname{arctanh}\left(\frac{x}{c}\right)}{c^3} \right)
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c/x])/x^3,x]`

output `-1/2*(a + b*ArcTanh[c/x])/x^2 - (b*c*(1/(c^2*x) - ArcTanh[x/c]/c^3))/2`

Definitions of rubi rules used

rule 220 $\text{Int}[(a_.) + (b_.) \cdot (x_.)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[b, 2])^{-1}) \cdot \text{ArcTanh}[\text{Rt}[b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 264 $\text{Int}[(c_.) \cdot (x_.)^m] \cdot ((a_.) + (b_.) \cdot (x_.)^2)^{p_}, x_Symbol] \rightarrow \text{Simp}[(c \cdot x)^{m+1} \cdot ((a + b \cdot x^2)^{p+1} / (a \cdot c \cdot (m+1))), x] - \text{Simp}[b \cdot ((m+2 \cdot p+3) / (a \cdot c^{2 \cdot (m+1)})) \ \text{Int}[(c \cdot x)^{m+2} \cdot (a + b \cdot x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 795 $\text{Int}[(x_.)^{m_}] \cdot ((a_.) + (b_.) \cdot (x_.)^{n_})^{p_}, x_Symbol] \rightarrow \text{Int}[x^{m+n \cdot p} \cdot (b + a/x^n)^p, x] /; \text{FreeQ}[\{a, b, m, n\}, x] \ \&\& \ \text{IntegerQ}[p] \ \&\& \ \text{NegQ}[n]$

rule 6452 $\text{Int}[(a_.) + \text{ArcTanh}[(c_.) \cdot (x_.)^{n_}]] \cdot (b_.)^{p_} \cdot (x_.)^{m_}, x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot ((a + b \cdot \text{ArcTanh}[c \cdot x^n])^p / (m+1)), x] - \text{Simp}[b \cdot c \cdot n \cdot (p / (m+1)) \ \text{Int}[x^{m+n} \cdot ((a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2 \cdot n}))], x], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.98

method	result
parallelrisch	$-\frac{-\operatorname{arctanh}\left(\frac{c}{x}\right)bx^2+\operatorname{arctanh}\left(\frac{c}{x}\right)bc^2+bcx+ac^2}{2x^2c^2}$
parts	$-\frac{a}{2x^2}-\frac{b\left(\frac{c^2\operatorname{arctanh}\left(\frac{c}{x}\right)}{2x^2}+\frac{c}{2x}+\frac{\ln\left(\frac{c}{x}-1\right)}{4}-\frac{\ln\left(1+\frac{c}{x}\right)}{4}\right)}{c^2}$
derivativedivides	$-\frac{\frac{ac^2}{2x^2}+b\left(\frac{c^2\operatorname{arctanh}\left(\frac{c}{x}\right)}{2x^2}+\frac{c}{2x}+\frac{\ln\left(\frac{c}{x}-1\right)}{4}-\frac{\ln\left(1+\frac{c}{x}\right)}{4}\right)}{c^2}$
default	$-\frac{\frac{ac^2}{2x^2}+b\left(\frac{c^2\operatorname{arctanh}\left(\frac{c}{x}\right)}{2x^2}+\frac{c}{2x}+\frac{\ln\left(\frac{c}{x}-1\right)}{4}-\frac{\ln\left(1+\frac{c}{x}\right)}{4}\right)}{c^2}$
orering	$-\frac{2(c^2-x^2)(a+b\operatorname{arctanh}\left(\frac{c}{x}\right))}{x^2c^2}-\frac{(c-x)(x+c)x^2\left(-\frac{bc}{x^5\left(1-\frac{c^2}{x^2}\right)}-\frac{3(a+b\operatorname{arctanh}\left(\frac{c}{x}\right))}{x^4}\right)}{2c^2}$
risch	$-\frac{b\ln(x+c)}{4x^2}-\frac{-i\pi bc^2\operatorname{csgn}\left(\frac{i}{x}\right)\operatorname{csgn}(i(x+c))\operatorname{csgn}\left(\frac{i(x+c)}{x}\right)+i\pi bc^2\operatorname{csgn}(i(x+c))\operatorname{csgn}\left(\frac{i(x+c)}{x}\right)^2+i\pi bc^2\operatorname{csgn}\left(\frac{i}{x}\right)\operatorname{csgn}(i(x+c))}{c^2}$

input `int((a+b*arctanh(c/x))/x^3,x,method=_RETURNVERBOSE)`

output `-1/2*(-arctanh(c/x)*b*x^2+arctanh(c/x)*b*c^2+b*c*x+a*c^2)/x^2/c^2`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.07

$$\int \frac{a + b\operatorname{arctanh}\left(\frac{c}{x}\right)}{x^3} dx = -\frac{2ac^2 + 2bcx + (bc^2 - bx^2)\log\left(-\frac{c+x}{c-x}\right)}{4c^2x^2}$$

input `integrate((a+b*arctanh(c/x))/x^3,x, algorithm="fricas")`

output `-1/4*(2*a*c^2 + 2*b*c*x + (b*c^2 - b*x^2)*log(-(c + x)/(c - x)))/(c^2*x^2)`

Sympy [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.02

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^3} dx = \begin{cases} -\frac{a}{2x^2} - \frac{b \operatorname{atanh}\left(\frac{c}{x}\right)}{2x^2} - \frac{b}{2cx} + \frac{b \operatorname{atanh}\left(\frac{c}{x}\right)}{2c^2} & \text{for } c \neq 0 \\ -\frac{a}{2x^2} & \text{otherwise} \end{cases}$$

input `integrate((a+b*atanh(c/x))/x**3,x)`

output `Piecewise((-a/(2*x**2) - b*atanh(c/x)/(2*x**2) - b/(2*c*x) + b*atanh(c/x)/(2*c**2), Ne(c, 0)), (-a/(2*x**2), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.21

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^3} dx = \frac{1}{4} \left(c \left(\frac{\log(c+x)}{c^3} - \frac{\log(-c+x)}{c^3} - \frac{2}{c^2 x} \right) - \frac{2 \operatorname{artanh}\left(\frac{c}{x}\right)}{x^2} \right) b - \frac{a}{2x^2}$$

input `integrate((a+b*arctanh(c/x))/x^3,x, algorithm="maxima")`

output `1/4*(c*(log(c + x)/c^3 - log(-c + x)/c^3 - 2/(c^2*x)) - 2*arctanh(c/x)/x^2)*b - 1/2*a/x^2`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 123 vs. $2(37) = 74$.

Time = 0.12 (sec) , antiderivative size = 123, normalized size of antiderivative = 2.86

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^3} dx = -\frac{\frac{b(c+x) \log\left(-\frac{c+x}{c-x}\right)}{\left(\frac{(c+x)^2 c}{(c-x)^2} - \frac{2(c+x)c}{c-x} + c\right)(c-x)} - \frac{b - \frac{2a(c+x)}{c-x} - \frac{b(c+x)}{c-x}}{\frac{(c+x)^2 c}{(c-x)^2} - \frac{2(c+x)c}{c-x} + c}}{c}$$

input `integrate((a+b*arctanh(c/x))/x^3,x, algorithm="giac")`

output `-(b*(c + x)*log(-(c + x)/(c - x)))/((c + x)^2*c/(c - x)^2 - 2*(c + x)*c/(c - x) + c)*(c - x) - (b - 2*a*(c + x)/(c - x) - b*(c + x)/(c - x))/((c + x)^2*c/(c - x)^2 - 2*(c + x)*c/(c - x) + c)/c`

Mupad [B] (verification not implemented)

Time = 3.52 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.14

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^3} dx = \frac{b c \operatorname{atan}\left(\frac{x}{\sqrt{-c^2}}\right)}{2(-c^2)^{3/2}} - \frac{b}{2cx} - \frac{b \operatorname{atanh}\left(\frac{c}{x}\right)}{2x^2} - \frac{a}{2x^2}$$

input `int((a + b*atanh(c/x))/x^3,x)`

output `(b*c*atan(x/(-c^2)^(1/2)))/(2*(-c^2)^(3/2)) - b/(2*c*x) - (b*atanh(c/x))/(2*x^2) - a/(2*x^2)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^3} dx = \frac{-\operatorname{atanh}\left(\frac{c}{x}\right) b c^2 + \operatorname{atanh}\left(\frac{c}{x}\right) b x^2 - a c^2 - b c x}{2c^2 x^2}$$

input `int((a+b*atanh(c/x))/x^3,x)`

output `(- atanh(c/x)*b*c**2 + atanh(c/x)*b*x**2 - a*c**2 - b*c*x)/(2*c**2*x**2)`

3.142 $\int \frac{a+b\operatorname{arctanh}\left(\frac{c}{x}\right)}{x^4} dx$

Optimal result	1161
Mathematica [A] (verified)	1161
Rubi [A] (verified)	1162
Maple [A] (verified)	1164
Fricas [A] (verification not implemented)	1164
Sympy [A] (verification not implemented)	1165
Maxima [A] (verification not implemented)	1165
Giac [B] (verification not implemented)	1166
Mupad [B] (verification not implemented)	1166
Reduce [B] (verification not implemented)	1167

Optimal result

Integrand size = 14, antiderivative size = 48

$$\int \frac{a + b\operatorname{arctanh}\left(\frac{c}{x}\right)}{x^4} dx = -\frac{b}{6cx^2} - \frac{a + b\operatorname{arctanh}\left(\frac{c}{x}\right)}{3x^3} - \frac{b \log\left(1 - \frac{c^2}{x^2}\right)}{6c^3}$$

output `-1/6*b/c/x^2-1/3*(a+b*arctanh(c/x))/x^3-1/6*b*ln(1-c^2/x^2)/c^3`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.29

$$\int \frac{a + b\operatorname{arctanh}\left(\frac{c}{x}\right)}{x^4} dx = -\frac{a}{3x^3} - \frac{b}{6cx^2} - \frac{b\operatorname{arctanh}\left(\frac{c}{x}\right)}{3x^3} + \frac{b \log(x)}{3c^3} - \frac{b \log(-c^2 + x^2)}{6c^3}$$

input `Integrate[(a + b*ArcTanh[c/x])/x^4,x]`

output `-1/3*a/x^3 - b/(6*c*x^2) - (b*ArcTanh[c/x])/(3*x^3) + (b*Log[x])/(3*c^3) - (b*Log[-c^2 + x^2])/(6*c^3)`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.15, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {6452, 795, 243, 25, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barctanh}\left(\frac{c}{x}\right)}{x^4} dx \\
 & \quad \downarrow \text{6452} \\
 & -\frac{1}{3}bc \int \frac{1}{\left(1 - \frac{c^2}{x^2}\right)x^5} dx - \frac{a + \operatorname{barctanh}\left(\frac{c}{x}\right)}{3x^3} \\
 & \quad \downarrow \text{795} \\
 & -\frac{1}{3}bc \int \frac{1}{x^3(x^2 - c^2)} dx - \frac{a + \operatorname{barctanh}\left(\frac{c}{x}\right)}{3x^3} \\
 & \quad \downarrow \text{243} \\
 & -\frac{1}{6}bc \int -\frac{1}{x^4(c^2 - x^2)} dx^2 - \frac{a + \operatorname{barctanh}\left(\frac{c}{x}\right)}{3x^3} \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{6}bc \int \frac{1}{x^4(c^2 - x^2)} dx^2 - \frac{a + \operatorname{barctanh}\left(\frac{c}{x}\right)}{3x^3} \\
 & \quad \downarrow \text{54} \\
 & \frac{1}{6}bc \int \left(\frac{1}{c^4 x^2} + \frac{1}{c^2 x^4} + \frac{1}{c^4 (c^2 - x^2)} \right) dx^2 - \frac{a + \operatorname{barctanh}\left(\frac{c}{x}\right)}{3x^3} \\
 & \quad \downarrow \text{2009} \\
 & -\frac{a + \operatorname{barctanh}\left(\frac{c}{x}\right)}{3x^3} - \frac{1}{6}bc \left(-\frac{\log(x^2)}{c^4} + \frac{1}{c^2 x^2} + \frac{\log(c^2 - x^2)}{c^4} \right)
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c/x])/x^4,x]`

output
$$-1/3*(a + b*\text{ArcTanh}[c/x])/x^3 - (b*c*(1/(c^2*x^2) - \text{Log}[x^2]/c^4 + \text{Log}[c^2 - x^2]/c^4))/6$$

Defintions of rubi rules used

rule 25
$$\text{Int}[-(\text{Fx}_), x_Symbol] \text{ :> } \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, x], x]$$

rule 54
$$\text{Int}[(a_) + (b_)*(x_)^{(m_)}*((c_) + (d_)*(x_)^{(n_)}), x_Symbol] \text{ :> } \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] \text{ ; FreeQ}\{a, b, c, d\}, x] \ \&\& \ \&\& \ \text{IntegerQ}[m, 0] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ !(\text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[m + n + 2, 0])$$

rule 243
$$\text{Int}[(x_)^{(m_)}*((a_) + (b_)*(x_)^2)^{(p_)}], x_Symbol] \text{ :> } \text{Simp}[1/2 \quad \text{Subst}[\text{Int}[x^{((m-1)/2)*(a+b*x)^p}, x], x, x^2], x] \text{ ; FreeQ}\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$$

rule 795
$$\text{Int}[(x_)^{(m_)}*((a_) + (b_)*(x_)^{(n_)})^{(p_)}], x_Symbol] \text{ :> } \text{Int}[x^{(m+n*p)}*(b + a/x^n)^p, x] \text{ ; FreeQ}\{a, b, m, n\}, x] \ \&\& \ \text{IntegerQ}[p] \ \&\& \ \text{NegQ}[n]$$

rule 2009
$$\text{Int}[u_, x_Symbol] \text{ :> } \text{Simp}[\text{IntSum}[u, x], x] \text{ ; SumQ}[u]$$

rule 6452
$$\text{Int}[(a_) + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_)]^{(p_)}*(x_)^{(m_)}], x_Symbol] \text{ :> } \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \quad \text{Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{2*n})}), x], x] \text{ ; FreeQ}\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$$

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.19

method	result
parts	$-\frac{a}{3x^3} - \frac{b \left(\frac{c^3 \operatorname{arctanh}\left(\frac{c}{x}\right)}{3x^3} + \frac{c^2}{6x^2} + \frac{\ln\left(\frac{c}{x}-1\right)}{6} + \frac{\ln\left(1+\frac{c}{x}\right)}{6} \right)}{c^3}$
derivativedivides	$-\frac{\frac{a}{3x^3} + b \left(\frac{c^3 \operatorname{arctanh}\left(\frac{c}{x}\right)}{3x^3} + \frac{c^2}{6x^2} + \frac{\ln\left(\frac{c}{x}-1\right)}{6} + \frac{\ln\left(1+\frac{c}{x}\right)}{6} \right)}{c^3}$
default	$-\frac{\frac{a}{3x^3} + b \left(\frac{c^3 \operatorname{arctanh}\left(\frac{c}{x}\right)}{3x^3} + \frac{c^2}{6x^2} + \frac{\ln\left(\frac{c}{x}-1\right)}{6} + \frac{\ln\left(1+\frac{c}{x}\right)}{6} \right)}{c^3}$
parallelrisc	$\frac{2b \ln(x)x^3 - 2 \ln(x-c)x^3 b - 2b x^3 \operatorname{arctanh}\left(\frac{c}{x}\right) - 2 \operatorname{arctanh}\left(\frac{c}{x}\right) b c^3 - b c^2 x - 2a c^3}{6x^3 c^3}$
risc	$-\frac{b \ln(x+c)}{6x^3} - \frac{2i\pi b c^3 \operatorname{csgn}\left(\frac{i(c-x)}{x}\right)^2 + i\pi b c^3 \operatorname{csgn}\left(\frac{i}{x}\right) \operatorname{csgn}(i(c-x)) \operatorname{csgn}\left(\frac{i(c-x)}{x}\right) + i\pi b c^3 \operatorname{csgn}(i(x+c)) \operatorname{csgn}\left(\frac{i(x+c)}{x}\right)}{6x^3}$

input `int((a+b*arctanh(c/x))/x^4,x,method=_RETURNVERBOSE)`

output `-1/3*a/x^3-b/c^3*(1/3*c^3/x^3*arctanh(c/x)+1/6*c^2/x^2+1/6*ln(c/x-1)+1/6*ln(1+c/x))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.29

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^4} dx$$

$$= -\frac{bx^3 \log(-c^2 + x^2) - 2bx^3 \log(x) + bc^3 \log\left(-\frac{c+x}{c-x}\right) + 2ac^3 + bc^2x}{6c^3x^3}$$

input `integrate((a+b*arctanh(c/x))/x^4,x, algorithm="fricas")`

output `-1/6*(b*x^3*log(-c^2 + x^2) - 2*b*x^3*log(x) + b*c^3*log(-(c + x)/(c - x)) + 2*a*c^3 + b*c^2*x)/(c^3*x^3)`

Sympy [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.42

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^4} dx$$

$$= \begin{cases} -\frac{a}{3x^3} - \frac{b \operatorname{arctanh}\left(\frac{c}{x}\right)}{3x^3} - \frac{b}{6cx^2} + \frac{b \log(x)}{3c^3} - \frac{b \log(-c+x)}{3c^3} - \frac{b \operatorname{arctanh}\left(\frac{c}{x}\right)}{3c^3} & \text{for } c \neq 0 \\ -\frac{a}{3x^3} & \text{otherwise} \end{cases}$$

input `integrate((a+b*atanh(c/x))/x**4,x)`output `Piecewise((-a/(3*x**3) - b*atanh(c/x)/(3*x**3) - b/(6*c*x**2) + b*log(x)/(3*c**3) - b*log(-c + x)/(3*c**3) - b*atanh(c/x)/(3*c**3), Ne(c, 0)), (-a/(3*x**3), True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.15

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^4} dx =$$

$$-\frac{1}{6} \left(c \left(\frac{\log(-c^2 + x^2)}{c^4} - \frac{\log(x^2)}{c^4} + \frac{1}{c^2 x^2} \right) + \frac{2 \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^3} \right) b$$

$$- \frac{a}{3x^3}$$

input `integrate((a+b*arctanh(c/x))/x^4,x, algorithm="maxima")`output `-1/6*(c*(log(-c^2 + x^2)/c^4 - log(x^2)/c^4 + 1/(c^2*x^2)) + 2*arctanh(c/x)/x^3)*b - 1/3*a/x^3`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 234 vs. $2(42) = 84$.

Time = 0.13 (sec) , antiderivative size = 234, normalized size of antiderivative = 4.88

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^4} dx = \frac{\left(b + \frac{3b(c+x)^2}{(c-x)^2}\right) \log\left(-\frac{c+x}{c-x}\right)}{\frac{(c+x)^3 c^2}{(c-x)^3} - \frac{3(c+x)^2 c^2}{(c-x)^2} + \frac{3(c+x)c^2}{c-x} - c^2} + \frac{2\left(a + \frac{3a(c+x)^2}{(c-x)^2} + \frac{b(c+x)^2}{(c-x)^2} - \frac{b(c+x)}{c-x}\right)}{\frac{(c+x)^3 c^2}{(c-x)^3} - \frac{3(c+x)^2 c^2}{(c-x)^2} + \frac{3(c+x)c^2}{c-x} - c^2} - \frac{b \log\left(-\frac{c+x}{c-x} + 1\right)}{c^2} + \frac{b \log\left(-\frac{c+x}{c-x}\right)}{c^2}$$

$3c$

input `integrate((a+b*arctanh(c/x))/x^4,x, algorithm="giac")`

output `-1/3*((b + 3*b*(c + x)^2/(c - x)^2)*log(-(c + x)/(c - x)))/((c + x)^3*c^2/(c - x)^3 - 3*(c + x)^2*c^2/(c - x)^2 + 3*(c + x)*c^2/(c - x) - c^2) + 2*(a + 3*a*(c + x)^2/(c - x)^2 + b*(c + x)^2/(c - x)^2 - b*(c + x)/(c - x))/((c + x)^3*c^2/(c - x)^3 - 3*(c + x)^2*c^2/(c - x)^2 + 3*(c + x)*c^2/(c - x) - c^2) - b*log(-(c + x)/(c - x) + 1)/c^2 + b*log(-(c + x)/(c - x))/c^2/c`

Mupad [B] (verification not implemented)

Time = 3.49 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.23

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^4} dx = -\frac{a}{3} + \frac{b \operatorname{atanh}\left(\frac{c}{x}\right)}{3} - \frac{b x^3 \ln(x^2 - c^2)}{6} - \frac{b x^3 \ln(x)}{3} + \frac{b c^2 x}{6}$$

input `int((a + b*atanh(c/x))/x^4,x)`

output `-(a/3 + (b*atanh(c/x))/3)/x^3 - ((b*x^3*log(x^2 - c^2))/6 - (b*x^3*log(x))/3 + (b*c^2*x)/6)/(c^3*x^3)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.42

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x}\right)}{x^4} dx$$

$$= \frac{-2 \operatorname{atanh}\left(\frac{c}{x}\right) b c^3 + 2 \operatorname{atanh}\left(\frac{c}{x}\right) b x^3 - 2 \log(-c - x) b x^3 + 2 \log(x) b x^3 - 2 a c^3 - b c^2 x}{6 c^3 x^3}$$

input `int((a+b*atanh(c/x))/x^4,x)`

output `(- 2*atanh(c/x)*b*c**3 + 2*atanh(c/x)*b*x**3 - 2*log(- c - x)*b*x**3 + 2*log(x)*b*x**3 - 2*a*c**3 - b*c**2*x)/(6*c**3*x**3)`

3.143 $\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx$

Optimal result	1168
Mathematica [A] (verified)	1169
Rubi [A] (warning: unable to verify)	1169
Maple [A] (verified)	1173
Fricas [A] (verification not implemented)	1174
Sympy [A] (verification not implemented)	1174
Maxima [A] (verification not implemented)	1175
Giac [B] (verification not implemented)	1175
Mupad [B] (verification not implemented)	1176
Reduce [B] (verification not implemented)	1177

Optimal result

Integrand size = 16, antiderivative size = 123

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \frac{1}{12} b^2 c^2 x^2 + \frac{1}{2} b c^3 x \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right) + \frac{1}{6} b c x^3 \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right) - \frac{1}{4} c^4 \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right)^2 + \frac{1}{4} x^4 \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right)^2 + \frac{1}{3} b^2 c^4 \log \left(1 - \frac{c^2}{x^2} \right) + \frac{2}{3} b^2 c^4 \log(x)$$

output

```
1/12*b^2*c^2*x^2+1/2*b*c^3*x*(a+b*arccoth(x/c))+1/6*b*c*x^3*(a+b*arccoth(x/c))-1/4*c^4*(a+b*arccoth(x/c))^2+1/4*x^4*(a+b*arccoth(x/c))^2+1/3*b^2*c^4*ln(1-c^2/x^2)+2/3*b^2*c^4*ln(x)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.07

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \frac{1}{12} \left(6abc^3x + b^2c^2x^2 + 2abcx^3 + 3a^2x^4 \right. \\ \left. + 2bx(3ax^3 + bc(3c^2 + x^2)) \operatorname{arctanh} \left(\frac{c}{x} \right) \right. \\ \left. + 3b^2(-c^4 + x^4) \operatorname{arctanh} \left(\frac{c}{x} \right)^2 + b(3a + 4b)c^4 \log(-c + x) \right. \\ \left. - 3abc^4 \log(c + x) + 4b^2c^4 \log(c + x) \right)$$

input

```
Integrate[x^3*(a + b*ArcTanh[c/x])^2,x]
```

output

```
(6*a*b*c^3*x + b^2*c^2*x^2 + 2*a*b*c*x^3 + 3*a^2*x^4 + 2*b*x*(3*a*x^3 + b*c*(3*c^2 + x^2))*ArcTanh[c/x] + 3*b^2*(-c^4 + x^4)*ArcTanh[c/x]^2 + b*(3*a + 4*b)*c^4*Log[-c + x] - 3*a*b*c^4*Log[c + x] + 4*b^2*c^4*Log[c + x])/12
```

Rubi [A] (warning: unable to verify)

Time = 1.01 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.14, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {6454, 6452, 6544, 6452, 243, 54, 2009, 6544, 6452, 243, 47, 14, 16, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx \\ \downarrow 6454 \\ - \int x^5 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 d \frac{1}{x} \\ \downarrow 6452 \\ \frac{1}{4} x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 - \frac{1}{2} bc \int \frac{x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)}{1 - \frac{c^2}{x^2}} d \frac{1}{x}$$

↓ 6544

$$\frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^2 - \frac{1}{2}bc\left(c^2 \int \frac{x^2(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \int x^4(a + \operatorname{barctanh}(\frac{c}{x})) d\frac{1}{x}\right)$$

↓ 6452

$$\frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^2 - \frac{1}{2}bc\left(c^2 \int \frac{x^2(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{1}{3}bc \int \frac{x^3}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - \frac{1}{3}x^3\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)\right)$$

↓ 243

$$\frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^2 - \frac{1}{2}bc\left(c^2 \int \frac{x^2(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{1}{6}bc \int \frac{x^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x^2} - \frac{1}{3}x^3\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)\right)$$

↓ 54

$$\frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^2 - \frac{1}{2}bc\left(c^2 \int \frac{x^2(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{1}{6}bc \int \left(-\frac{c^4}{\frac{c^2}{x^2} - 1} + xc^2 + x^2\right) d\frac{1}{x^2} - \frac{1}{3}x^3\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)\right)$$

↓ 2009

$$\frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^2 - \frac{1}{2}bc\left(c^2 \int \frac{x^2(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - \frac{1}{3}x^3\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right) + \frac{1}{6}bc\left(c^2\left(-\log\left(1 - \frac{c^2}{x^2}\right)\right) + c^2 \log\left(\frac{1}{x^2}\right) - x\right)\right)$$

↓ 6544

$$\frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^2 - \frac{1}{2}bc\left(c^2\left(c^2 \int \frac{a + \operatorname{barctanh}(\frac{c}{x})}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \int x^2\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right) d\frac{1}{x}\right) - \frac{1}{3}x^3\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right) + \frac{1}{6}bc\left(c^2\left(-\log\left(1 - \frac{c^2}{x^2}\right)\right) + c^2 \log\left(\frac{1}{x^2}\right) - x\right)\right)$$

↓ 6452

$$\frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^2 - \frac{1}{2}bc\left(c^2\left(c^2 \int \frac{a + \operatorname{barctanh}(\frac{c}{x})}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + bc \int \frac{x}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - x\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)\right) - \frac{1}{3}x^3\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right) + \frac{1}{6}bc\left(c^2\left(-\log\left(1 - \frac{c^2}{x^2}\right)\right) + c^2 \log\left(\frac{1}{x^2}\right) - x\right)\right)$$

$$\begin{aligned}
 & \downarrow 243 \\
 & \frac{1}{4}x^4\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2 - \\
 & \frac{1}{2}bc\left(c^2\left(c^2\int\frac{a + \operatorname{arctanh}\left(\frac{c}{x}\right)}{1 - \frac{c^2}{x^2}}d\frac{1}{x} + \frac{1}{2}bc\int\frac{x}{1 - \frac{c^2}{x^2}}d\frac{1}{x^2} - x\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)\right) - \frac{1}{3}x^3\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right) + \right. \\
 & \downarrow 47 \\
 & \frac{1}{4}x^4\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2 - \\
 & \frac{1}{2}bc\left(c^2\left(c^2\int\frac{a + \operatorname{arctanh}\left(\frac{c}{x}\right)}{1 - \frac{c^2}{x^2}}d\frac{1}{x} + \frac{1}{2}bc\left(c^2\int\frac{1}{1 - \frac{c^2}{x^2}}d\frac{1}{x^2} + \int xd\frac{1}{x^2}\right) - x\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)\right) - \frac{1}{3}x^3\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right) + \right. \\
 & \downarrow 14 \\
 & \frac{1}{4}x^4\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2 - \\
 & \frac{1}{2}bc\left(c^2\left(c^2\int\frac{a + \operatorname{arctanh}\left(\frac{c}{x}\right)}{1 - \frac{c^2}{x^2}}d\frac{1}{x} + \frac{1}{2}bc\left(c^2\int\frac{1}{1 - \frac{c^2}{x^2}}d\frac{1}{x^2} + \log\left(\frac{1}{x^2}\right)\right) - x\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)\right) - \frac{1}{3}x^3\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right) + \right. \\
 & \downarrow 16 \\
 & \frac{1}{4}x^4\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2 - \\
 & \frac{1}{2}bc\left(c^2\left(c^2\int\frac{a + \operatorname{arctanh}\left(\frac{c}{x}\right)}{1 - \frac{c^2}{x^2}}d\frac{1}{x} - x\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right) + \frac{1}{2}bc\left(\log\left(\frac{1}{x^2}\right) - \log\left(1 - \frac{c^2}{x^2}\right)\right)\right) - \frac{1}{3}x^3\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right) + \right. \\
 & \downarrow 6510 \\
 & \frac{1}{4}x^4\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2 - \\
 & \frac{1}{2}bc\left(c^2\left(\frac{c\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2}{2b} - x\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right) + \frac{1}{2}bc\left(\log\left(\frac{1}{x^2}\right) - \log\left(1 - \frac{c^2}{x^2}\right)\right)\right) - \frac{1}{3}x^3\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right) + \right.
 \end{aligned}$$

input

`Int [x^3*(a + b*ArcTanh [c/x])^2,x]`

output

$(x^4*(a + b*ArcTanh [c/x])^2)/4 - (b*c*(-1/3*(x^3*(a + b*ArcTanh [c/x])) + (b*c*(-x - c^2*Log [1 - c^2/x^2] + c^2*Log [x^(-2)])))/6 + c^2*(-(x*(a + b*ArcTanh [c/x])) + (c*(a + b*ArcTanh [c/x])^2)/(2*b) + (b*c*(-Log [1 - c^2/x^2] + Log [x^(-2)])))/2)))/2$

Definitions of rubi rules used

- rule 14 $\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ ; FreeQ}[a, x]$
- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$
- rule 47 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x]$
- rule 54 $\text{Int}(((a_)+(b_)*(x_))^{(m_)}*((c_)+(d_)*(x_))^{(n_)}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{ILtQ}[m, 0] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ !(\text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[m + n + 2, 0])$
- rule 243 $\text{Int}[(x_)^{(m_)}*((a_)+(b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] \text{ ; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$
- rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ ; SumQ}[u]$
- rule 6452 $\text{Int}(((a_)+\text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}), x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{(2*n)})}), x], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$
- rule 6454 $\text{Int}(((a_)+\text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}), x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n] - 1)}*(a + b*\text{ArcTanh}[c*x])^p, x], x, x^n], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$

```
rule 6510 Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol]
  := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x]
  && EqQ[c^2*d + e, 0] && NeQ[p, -1]
```

```
rule 6544 Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*((f_.)*(x_)^(m_))/((d_) + (e_.)*(x_)^2), x_Symbol]
  := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /;
  FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

Maple [A] (verified)

Time = 1.83 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.31

method	result
parallelrisch	$\frac{x^4 \operatorname{arctanh}\left(\frac{c}{x}\right)^2 b^2}{4} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right)^2 b^2 c^4}{4} + \frac{2b^2 c^4 \ln(x-c)}{3} + \frac{x^4 \operatorname{arctanh}\left(\frac{c}{x}\right) ab}{2} + \frac{x^3 \operatorname{arctanh}\left(\frac{c}{x}\right) b^2 c}{6} + \frac{x \operatorname{arctanh}\left(\frac{c}{x}\right) b^2 c^3}{2}$
parts	$\frac{a^2 x^4}{4} - b^2 c^4 \left(-\frac{x^4 \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{4c^4} - \frac{x^3 \operatorname{arctanh}\left(\frac{c}{x}\right)}{6c^3} - \frac{x \operatorname{arctanh}\left(\frac{c}{x}\right)}{2c} + \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(1+\frac{c}{x}\right)}{4} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right)}{4} \right)$
derivativedivides	$-c^4 \left(-\frac{a^2 x^4}{4c^4} + b^2 \left(-\frac{x^4 \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{4c^4} - \frac{x^3 \operatorname{arctanh}\left(\frac{c}{x}\right)}{6c^3} - \frac{x \operatorname{arctanh}\left(\frac{c}{x}\right)}{2c} + \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(1+\frac{c}{x}\right)}{4} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right)}{4} \right) \right)$
default	$-c^4 \left(-\frac{a^2 x^4}{4c^4} + b^2 \left(-\frac{x^4 \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{4c^4} - \frac{x^3 \operatorname{arctanh}\left(\frac{c}{x}\right)}{6c^3} - \frac{x \operatorname{arctanh}\left(\frac{c}{x}\right)}{2c} + \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(1+\frac{c}{x}\right)}{4} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right)}{4} \right) \right)$
risch	Expression too large to display

```
input int(x^3*(a+b*arctanh(c/x))^2,x,method=_RETURNVERBOSE)
```

```
output 1/4*x^4*arctanh(c/x)^2*b^2-1/4*arctanh(c/x)^2*b^2*c^4+2/3*b^2*c^4*ln(x-c)+
1/2*x^4*arctanh(c/x)*a*b+1/6*x^3*arctanh(c/x)*b^2*c+1/2*x*arctanh(c/x)*b^2
*c^3-1/2*arctanh(c/x)*a*b*c^4+2/3*arctanh(c/x)*b^2*c^4+1/4*a^2*x^4+1/6*a*b
*c*x^3+1/12*b^2*c^2*x^2+1/2*c^3*a*b*x+1/12*b^2*c^4
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.21

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \frac{1}{2} abc^3 x + \frac{1}{12} b^2 c^2 x^2 + \frac{1}{6} abc x^3 + \frac{1}{4} a^2 x^4$$

$$- \frac{1}{12} (3ab - 4b^2) c^4 \log(c + x)$$

$$+ \frac{1}{12} (3ab + 4b^2) c^4 \log(-c + x)$$

$$- \frac{1}{16} (b^2 c^4 - b^2 x^4) \log \left(-\frac{c+x}{c-x} \right)^2$$

$$+ \frac{1}{12} (3b^2 c^3 x + b^2 c x^3 + 3abx^4) \log \left(-\frac{c+x}{c-x} \right)$$

input `integrate(x^3*(a+b*arctanh(c/x))^2,x, algorithm="fricas")`

output `1/2*a*b*c^3*x + 1/12*b^2*c^2*x^2 + 1/6*a*b*c*x^3 + 1/4*a^2*x^4 - 1/12*(3*a*b - 4*b^2)*c^4*log(c + x) + 1/12*(3*a*b + 4*b^2)*c^4*log(-c + x) - 1/16*(b^2*c^4 - b^2*x^4)*log(-(c + x)/(c - x))^2 + 1/12*(3*b^2*c^3*x + b^2*c*x^3 + 3*a*b*x^4)*log(-(c + x)/(c - x))`

Sympy [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.28

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \frac{a^2 x^4}{4} - \frac{abc^4 \operatorname{atanh} \left(\frac{c}{x} \right)}{2} + \frac{abc^3 x}{2} + \frac{abc x^3}{6}$$

$$+ \frac{abx^4 \operatorname{atanh} \left(\frac{c}{x} \right)}{2} + \frac{2b^2 c^4 \log(-c + x)}{3}$$

$$- \frac{b^2 c^4 \operatorname{atanh}^2 \left(\frac{c}{x} \right)}{4} + \frac{2b^2 c^4 \operatorname{atanh} \left(\frac{c}{x} \right)}{3} + \frac{b^2 c^3 x \operatorname{atanh} \left(\frac{c}{x} \right)}{2}$$

$$+ \frac{b^2 c^2 x^2}{12} + \frac{b^2 c x^3 \operatorname{atanh} \left(\frac{c}{x} \right)}{6} + \frac{b^2 x^4 \operatorname{atanh}^2 \left(\frac{c}{x} \right)}{4}$$

input `integrate(x**3*(a+b*atanh(c/x))**2,x)`

output

```
a**2*x**4/4 - a*b*c**4*atanh(c/x)/2 + a*b*c**3*x/2 + a*b*c*x**3/6 + a*b*x*
*4*atanh(c/x)/2 + 2*b**2*c**4*log(-c + x)/3 - b**2*c**4*atanh(c/x)**2/4 +
2*b**2*c**4*atanh(c/x)/3 + b**2*c**3*x*atanh(c/x)/2 + b**2*c**2*x**2/12 +
b**2*c*x**3*atanh(c/x)/6 + b**2*x**4*atanh(c/x)**2/4
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.54

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \frac{1}{4} b^2 x^4 \operatorname{artanh} \left(\frac{c}{x} \right)^2 + \frac{1}{4} a^2 x^4$$

$$+ \frac{1}{12} \left(6 x^4 \operatorname{artanh} \left(\frac{c}{x} \right) - (3 c^3 \log(c+x) - 3 c^3 \log(-c+x) - 6 c^2 x - 2 x^3) c \right) a b$$

$$+ \frac{1}{48} \left((3 c^2 \log(c+x))^2 + 3 c^2 \log(-c+x)^2 + 16 c^2 \log(c+x) + 4 x^2 - 2 (3 c^2 \log(c+x) - 8 c^2) \log(-c+x) \right)$$

input

```
integrate(x^3*(a+b*arctanh(c/x))^2,x, algorithm="maxima")
```

output

```
1/4*b^2*x^4*arctanh(c/x)^2 + 1/4*a^2*x^4 + 1/12*(6*x^4*arctanh(c/x) - (3*c
^3*log(c + x) - 3*c^3*log(-c + x) - 6*c^2*x - 2*x^3)*c)*a*b + 1/48*((3*c^2
*log(c + x)^2 + 3*c^2*log(-c + x)^2 + 16*c^2*log(c + x) + 4*x^2 - 2*(3*c^2
*log(c + x) - 8*c^2)*log(-c + x))*c^2 - 4*(3*c^3*log(c + x) - 3*c^3*log(-c
+ x) - 6*c^2*x - 2*x^3)*c*arctanh(c/x))*b^2
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 552 vs. 2(109) = 218.

Time = 0.13 (sec) , antiderivative size = 552, normalized size of antiderivative = 4.49

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \text{Too large to display}$$

input

```
integrate(x^3*(a+b*arctanh(c/x))^2,x, algorithm="giac")
```

output

```

-1/6*(4*b^2*c^5*log(-(c+x)/(c-x) - 1) - 4*b^2*c^5*log(-(c+x)/(c-x)
) + 3*(b^2*(c+x)^3*c^5/(c-x)^3 + b^2*(c+x)*c^5/(c-x))*log(-(c+x)
/(c-x))^2/((c+x)^4/(c-x)^4 + 4*(c+x)^3/(c-x)^3 + 6*(c+x)^2/(c
-x)^2 + 4*(c+x)/(c-x) + 1) + 2*(2*b^2*c^5 + 6*a*b*(c+x)^3*c^5/(c-
x)^3 + 3*b^2*(c+x)^3*c^5/(c-x)^3 + 6*b^2*(c+x)^2*c^5/(c-x)^2 + 6*a
*b*(c+x)*c^5/(c-x) + 5*b^2*(c+x)*c^5/(c-x))*log(-(c+x)/(c-x))/
((c+x)^4/(c-x)^4 + 4*(c+x)^3/(c-x)^3 + 6*(c+x)^2/(c-x)^2 + 4*(
c+x)/(c-x) + 1) + 2*(4*a*b*c^5 + 6*a^2*(c+x)^3*c^5/(c-x)^3 + 6*a*b
*(c+x)^3*c^5/(c-x)^3 + b^2*(c+x)^3*c^5/(c-x)^3 + 12*a*b*(c+x)^2*
c^5/(c-x)^2 + 2*b^2*(c+x)^2*c^5/(c-x)^2 + 6*a^2*(c+x)*c^5/(c-x)
+ 10*a*b*(c+x)*c^5/(c-x) + b^2*(c+x)*c^5/(c-x))/((c+x)^4/(c-x)
^4 + 4*(c+x)^3/(c-x)^3 + 6*(c+x)^2/(c-x)^2 + 4*(c+x)/(c-x) + 1
))/c

```

Mupad [B] (verification not implemented)

Time = 3.61 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.15

$$\begin{aligned}
\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx &= \frac{a^2 x^4}{4} - \frac{b^2 c^4 \operatorname{atanh} \left(\frac{c}{x} \right)^2}{4} + \frac{b^2 x^4 \operatorname{atanh} \left(\frac{c}{x} \right)^2}{4} \\
&+ \frac{b^2 c^4 \ln(x^2 - c^2)}{3} + \frac{b^2 c^2 x^2}{12} + \frac{b^2 c x^3 \operatorname{atanh} \left(\frac{c}{x} \right)}{6} \\
&+ \frac{b^2 c^3 x \operatorname{atanh} \left(\frac{c}{x} \right)}{2} + \frac{a b c x^3}{6} + \frac{a b c^3 x}{2} \\
&- \frac{a b c^4 \operatorname{atanh} \left(\frac{c}{x} \right)}{2} + \frac{a b x^4 \operatorname{atanh} \left(\frac{c}{x} \right)}{2}
\end{aligned}$$

input

```
int(x^3*(a + b*atanh(c/x))^2,x)
```

output

```

(a^2*x^4)/4 - (b^2*c^4*atanh(c/x)^2)/4 + (b^2*x^4*atanh(c/x)^2)/4 + (b^2*c
^4*log(x^2 - c^2))/3 + (b^2*c^2*x^2)/12 + (b^2*c*x^3*atanh(c/x))/6 + (b^2*
c^3*x*atanh(c/x))/2 + (a*b*c*x^3)/6 + (a*b*c^3*x)/2 - (a*b*c^4*atanh(c/x))
/2 + (a*b*x^4*atanh(c/x))/2

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.25

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = -\frac{\operatorname{atanh} \left(\frac{c}{x} \right)^2 b^2 c^4}{4} + \frac{\operatorname{atanh} \left(\frac{c}{x} \right)^2 b^2 x^4}{4} - \frac{\operatorname{atanh} \left(\frac{c}{x} \right) a b c^4}{2}$$

$$+ \frac{\operatorname{atanh} \left(\frac{c}{x} \right) a b x^4}{2} - \frac{2 \operatorname{atanh} \left(\frac{c}{x} \right) b^2 c^4}{3} + \frac{\operatorname{atanh} \left(\frac{c}{x} \right) b^2 c^3 x}{2}$$

$$+ \frac{\operatorname{atanh} \left(\frac{c}{x} \right) b^2 c x^3}{6} + \frac{2 \log(-c-x) b^2 c^4}{3}$$

$$+ \frac{a^2 x^4}{4} + \frac{a b c^3 x}{2} + \frac{a b c x^3}{6} + \frac{b^2 c^2 x^2}{12}$$

input `int(x^3*(a+b*atanh(c/x))^2,x)`output `(- 3*atanh(c/x)**2*b**2*c**4 + 3*atanh(c/x)**2*b**2*x**4 - 6*atanh(c/x)*a*b*c**4 + 6*atanh(c/x)*a*b*x**4 - 8*atanh(c/x)*b**2*c**4 + 6*atanh(c/x)*b**2*c**3*x + 2*atanh(c/x)*b**2*c*x**3 + 8*log(- c - x)*b**2*c**4 + 3*a**2*x**4 + 6*a*b*c**3*x + 2*a*b*c*x**3 + b**2*c**2*x**2)/12`

3.144 $\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx$

Optimal result	1178
Mathematica [A] (verified)	1179
Rubi [A] (verified)	1179
Maple [B] (verified)	1182
Fricas [F]	1183
Sympy [F]	1183
Maxima [F]	1184
Giac [F]	1184
Mupad [F(-1)]	1184
Reduce [F]	1185

Optimal result

Integrand size = 16, antiderivative size = 142

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \frac{1}{3} b^2 c^2 x - \frac{1}{3} b^2 c^3 \coth^{-1} \left(\frac{x}{c} \right) + \frac{1}{3} b c x^2 \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right) - \frac{1}{3} c^3 \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2 + \frac{1}{3} x^3 \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2 - \frac{2}{3} b c^3 \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right) \log \left(2 - \frac{2}{1 + \frac{c}{x}} \right) + \frac{1}{3} b^2 c^3 \operatorname{PolyLog} \left(2, -1 + \frac{2}{1 + \frac{c}{x}} \right)$$

output `1/3*b^2*c^2*x-1/3*b^2*c^3*arccoth(x/c)+1/3*b*c*x^2*(a+b*arccoth(x/c))-1/3*c^3*(a+b*arccoth(x/c))^2+1/3*x^3*(a+b*arccoth(x/c))^2-2/3*b*c^3*(a+b*arccoth(x/c))*ln(2-2/(1+c/x))+1/3*b^2*c^3*polylog(2,-1+2/(1+c/x))`

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.02

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \frac{1}{3} \left(b^2 c^2 x + abc x^2 + a^2 x^3 + b^2 (-c^3 + x^3) \operatorname{arctanh} \left(\frac{c}{x} \right)^2 \right. \\ \left. + b \operatorname{arctanh} \left(\frac{c}{x} \right) \left(-bc^3 + bcx^2 + 2ax^3 \right. \right. \\ \left. \left. - 2bc^3 \log \left(1 - e^{-2 \operatorname{arctanh} \left(\frac{c}{x} \right)} \right) \right) + abc^3 \log \left(1 - \frac{c^2}{x^2} \right) \right. \\ \left. - 2abc^3 \log \left(\frac{c}{x} \right) + b^2 c^3 \operatorname{PolyLog} \left(2, e^{-2 \operatorname{arctanh} \left(\frac{c}{x} \right)} \right) \right)$$

input `Integrate[x^2*(a + b*ArcTanh[c/x])^2,x]`

output `(b^2*c^2*x + a*b*c*x^2 + a^2*x^3 + b^2*(-c^3 + x^3)*ArcTanh[c/x]^2 + b*ArcTanh[c/x]*(-b*c^3) + b*c*x^2 + 2*a*x^3 - 2*b*c^3*Log[1 - E^(-2*ArcTanh[c/x])]) + a*b*c^3*Log[1 - c^2/x^2] - 2*a*b*c^3*Log[c/x] + b^2*c^3*PolyLog[2, E^(-2*ArcTanh[c/x])])/3`

Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.92, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.562$, Rules used = {6454, 6452, 6544, 6452, 264, 219, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx \\ \downarrow 6454 \\ - \int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 d \frac{1}{x} \\ \downarrow 6452$$

$$\begin{aligned}
& \frac{1}{3}x^3 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2 - \frac{2}{3}bc \int \frac{x^3 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} \\
& \quad \downarrow \text{6544} \\
& \frac{1}{3}x^3 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2 - \\
& \frac{2}{3}bc \left(c^2 \int \frac{x \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \int x^3 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right) d\frac{1}{x} \right) \\
& \quad \downarrow \text{6452} \\
& \frac{1}{3}x^3 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2 - \\
& \frac{2}{3}bc \left(c^2 \int \frac{x \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{1}{2}bc \int \frac{x^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - \frac{1}{2}x^2 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right) \right) \\
& \quad \downarrow \text{264} \\
& \frac{1}{3}x^3 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2 - \\
& \frac{2}{3}bc \left(c^2 \int \frac{x \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{1}{2}bc \left(c^2 \int \frac{1}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - x \right) - \frac{1}{2}x^2 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right) \right) \\
& \quad \downarrow \text{219} \\
& \frac{1}{3}x^3 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2 - \\
& \frac{2}{3}bc \left(c^2 \int \frac{x \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - \frac{1}{2}x^2 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right) + \frac{1}{2}bc \left(\operatorname{carctanh}\left(\frac{c}{x}\right) - x \right) \right) \\
& \quad \downarrow \text{6550} \\
& \frac{1}{3}x^3 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2 - \\
& \frac{2}{3}bc \left(c^2 \left(\int \frac{x \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)}{\frac{c}{x} + 1} d\frac{1}{x} + \frac{\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2}{2b} \right) - \frac{1}{2}x^2 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right) + \frac{1}{2}bc \left(\operatorname{carctanh}\left(\frac{c}{x}\right) - x \right) \right) \\
& \quad \downarrow \text{6494} \\
& \frac{1}{3}x^3 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2 - \\
& \frac{2}{3}bc \left(c^2 \left(-bc \int \frac{\log\left(2 - \frac{2}{\frac{c}{x} + 1}\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2}{2b} + \log\left(2 - \frac{2}{\frac{c}{x} + 1}\right) \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right) \right) - \frac{1}{2}x^2 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right) \right) \\
& \quad \downarrow \text{2897}
\end{aligned}$$

$$\frac{1}{3}x^3\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2 - \frac{2}{3}bc\left(c^2\left(\frac{(a + \operatorname{arctanh}\left(\frac{c}{x}\right))^2}{2b} + \log\left(2 - \frac{2}{\frac{c}{x} + 1}\right)\left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right) - \frac{1}{2}b\operatorname{PolyLog}\left(2, \frac{2}{\frac{c}{x} + 1} - 1\right)\right) - \frac{1}{2}x^2\left(\frac{c}{x} + 1\right)\right)$$

input `Int[x^2*(a + b*ArcTanh[c/x])^2,x]`

output `(x^3*(a + b*ArcTanh[c/x])^2)/3 - (2*b*c*(-1/2*(x^2*(a + b*ArcTanh[c/x])) + (b*c*(-x + c*ArcTanh[c/x]))/2 + c^2*((a + b*ArcTanh[c/x])^2/(2*b) + (a + b*ArcTanh[c/x])*Log[2 - 2/(1 + c/x)] - (b*PolyLog[2, -1 + 2/(1 + c/x)]))/2))/3`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 264 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + 2*p + 3)/(a*c^(2*(m + 1))) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2897 `Int[Log[u]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 $\text{Int}[(a + \text{ArcTanh}[c \cdot x]^n] \cdot (b \cdot x)^p \cdot x^m, x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n) - 1}] \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x, x^n], x] /; \text{FreeQ}\{a, b, c, m, n\}, x\} \&\& \text{IGtQ}[p, 1] \&\& \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$

rule 6494 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x))^p / ((x) \cdot ((d) + (e) \cdot (x))), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{Log}[2 - 2/(1 + e \cdot (x/d))]/d), x] - \text{Simp}[b \cdot c \cdot (p/d) \text{ Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{Log}[2 - 2/(1 + e \cdot (x/d))]) / (1 - c^2 \cdot x^2)], x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x\} \&\& \text{IGtQ}[p, 0] \&\& \text{EqQ}[c^2 \cdot d^2 - e^2, 0]$

rule 6544 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x))^p \cdot ((f) \cdot (x))^m / ((d) + (e) \cdot (x)^2), x_Symbol] \rightarrow \text{Simp}[1/d \text{ Int}[(f \cdot x)^m \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x] - \text{Simp}[e/(d \cdot f^2) \text{ Int}[(f \cdot x)^{m+2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2)], x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x\} \&\& \text{GtQ}[p, 0] \&\& \text{LtQ}[m, -1]$

rule 6550 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x))^p / ((x) \cdot ((d) + (e) \cdot (x)^2)), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot d \cdot (p+1)), x] + \text{Simp}[1/d \text{ Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p / (x \cdot (1 + c \cdot x)), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x\} \&\& \text{EqQ}[c^2 \cdot d + e, 0] \&\& \text{GtQ}[p, 0]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 297 vs. $2(128) = 256$.

Time = 1.48 (sec) , antiderivative size = 298, normalized size of antiderivative = 2.10

method	result
parts	$\frac{a^2 x^3}{3} - b^2 c^3 \left(-\frac{x^3 \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{3c^3} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(\frac{c}{x}-1\right)}{3} - \frac{x^2 \operatorname{arctanh}\left(\frac{c}{x}\right)}{3c^2} + \frac{2 \ln\left(\frac{c}{x}\right) \operatorname{arctanh}\left(\frac{c}{x}\right)}{3} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right)}{3} \right)$
derivativedivides	$-c^3 \left(-\frac{a^2 x^3}{3c^3} + b^2 \left(-\frac{x^3 \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{3c^3} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(\frac{c}{x}-1\right)}{3} - \frac{x^2 \operatorname{arctanh}\left(\frac{c}{x}\right)}{3c^2} + \frac{2 \ln\left(\frac{c}{x}\right) \operatorname{arctanh}\left(\frac{c}{x}\right)}{3} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right)}{3} \right) \right)$
default	$-c^3 \left(-\frac{a^2 x^3}{3c^3} + b^2 \left(-\frac{x^3 \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{3c^3} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(\frac{c}{x}-1\right)}{3} - \frac{x^2 \operatorname{arctanh}\left(\frac{c}{x}\right)}{3c^2} + \frac{2 \ln\left(\frac{c}{x}\right) \operatorname{arctanh}\left(\frac{c}{x}\right)}{3} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right)}{3} \right) \right)$
risch	Expression too large to display

input `int(x^2*(a+b*arctanh(c/x))^2,x,method=_RETURNVERBOSE)`

output `1/3*a^2*x^3-b^2*c^3*(-1/3/c^3*x^3*arctanh(c/x)^2-1/3*arctanh(c/x)*ln(c/x-1)-1/3/c^2*x^2*arctanh(c/x)+2/3*ln(c/x)*arctanh(c/x)-1/3*arctanh(c/x)*ln(1+c/x)-1/6*ln(c/x-1)-1/3*x/c+1/6*ln(1+c/x)+1/3*dilog(1/2*c/x+1/2)+1/6*ln(c/x-1)*ln(1/2*c/x+1/2)-1/12*ln(c/x-1)^2+1/12*ln(1+c/x)^2-1/6*(ln(1+c/x)-ln(1/2*c/x+1/2))*ln(-1/2*c/x+1/2)-1/3*dilog(c/x)-1/3*dilog(1+c/x)-1/3*ln(c/x)*ln(1+c/x))-2*a*b*c^3*(-1/3/c^3*x^3*arctanh(c/x)-1/6*ln(c/x-1)-1/6*ln(1+c/x)-1/6/c^2*x^2+1/3*ln(c/x))`

Fricas [F]

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x} \right) + a \right)^2 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c/x))^2,x, algorithm="fricas")`

output `integral(b^2*x^2*arctanh(c/x)^2 + 2*a*b*x^2*arctanh(c/x) + a^2*x^2, x)`

Sympy [F]

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \int x^2 \left(a + b \operatorname{atanh} \left(\frac{c}{x} \right) \right)^2 dx$$

input `integrate(x**2*(a+b*atanh(c/x))**2,x)`

output `Integral(x**2*(a + b*atanh(c/x))**2, x)`

Maxima [F]

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x} \right) + a \right)^2 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c/x))^2,x, algorithm="maxima")`

output `1/3*a^2*x^3 + 1/3*(2*x^3*arctanh(c/x) + (c^2*log(-c^2 + x^2) + x^2)*c)*a*b + 1/12*(6*c^4*integrate(-1/3*log(c + x)/(c^2 - x^2), x) + x^3*log(c + x)^2 + 6*c^3*integrate(-1/3*x*log(c + x)/(c^2 - x^2), x) - (c*log(c + x) - c*log(-c + x) - 2*x)*c^2 - (c^3 - x^3)*log(-c + x)^2 + (c^2*log(-c^2 + x^2) + x^2)*c + 12*c*integrate(-1/3*x^3*log(c + x)/(c^2 - x^2), x) - 2*(c*x^2 + (c^3 + x^3)*log(c + x))*log(-c + x))*b^2`

Giac [F]

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x} \right) + a \right)^2 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c/x))^2,x, algorithm="giac")`

output `integrate((b*arctanh(c/x) + a)^2*x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \int x^2 \left(a + b \operatorname{atanh} \left(\frac{c}{x} \right) \right)^2 dx$$

input `int(x^2*(a + b*atanh(c/x))^2,x)`

output `int(x^2*(a + b*atanh(c/x))^2, x)`

Reduce [F]

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = -\frac{\operatorname{atanh} \left(\frac{c}{x} \right)^2 b^2 c^2 x}{3} + \frac{\operatorname{atanh} \left(\frac{c}{x} \right)^2 b^2 x^3}{3}$$

$$- \frac{2 \operatorname{atanh} \left(\frac{c}{x} \right) a b c^3}{3} + \frac{2 \operatorname{atanh} \left(\frac{c}{x} \right) a b x^3}{3} - \frac{\operatorname{atanh} \left(\frac{c}{x} \right) b^2 c^3}{3}$$

$$+ \frac{\operatorname{atanh} \left(\frac{c}{x} \right) b^2 c x^2}{3} + \frac{\left(\int \operatorname{atanh} \left(\frac{c}{x} \right)^2 dx \right) b^2 c^2}{3}$$

$$+ \frac{2 \log(-c-x) a b c^3}{3} + \frac{a^2 x^3}{3} + \frac{a b c x^2}{3} + \frac{b^2 c^2 x}{3}$$

input

```
int(x^2*(a+b*atanh(c/x))^2,x)
```

output

```
( - atanh(c/x)**2*b**2*c**2*x + atanh(c/x)**2*b**2*x**3 - 2*atanh(c/x)*a*b
*c**3 + 2*atanh(c/x)*a*b*x**3 - atanh(c/x)*b**2*c**3 + atanh(c/x)*b**2*c*x
**2 + int(atanh(c/x)**2,x)*b**2*c**2 + 2*log(-c-x)*a*b*c**3 + a**2*x**
3 + a*b*c*x**2 + b**2*c**2*x)/3
```

3.145 $\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx$

Optimal result	1186
Mathematica [A] (verified)	1186
Rubi [A] (verified)	1187
Maple [A] (verified)	1190
Fricas [A] (verification not implemented)	1191
Sympy [A] (verification not implemented)	1191
Maxima [A] (verification not implemented)	1192
Giac [B] (verification not implemented)	1192
Mupad [B] (verification not implemented)	1193
Reduce [B] (verification not implemented)	1193

Optimal result

Integrand size = 14, antiderivative size = 83

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = bcx \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right) - \frac{1}{2} c^2 \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right)^2 + \frac{1}{2} x^2 \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right)^2 + \frac{1}{2} b^2 c^2 \log \left(1 - \frac{c^2}{x^2} \right) + b^2 c^2 \log(x)$$

output

```
b*c*x*(a+b*arccoth(x/c))-1/2*c^2*(a+b*arccoth(x/c))^2+1/2*x^2*(a+b*arccoth(x/c))^2+1/2*b^2*c^2*ln(1-c^2/x^2)+b^2*c^2*ln(x)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.11

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \frac{1}{2} \left(2abcx + a^2 x^2 + 2bx(bc + ax) \operatorname{arctanh} \left(\frac{c}{x} \right) + b^2 (-c^2 + x^2) \operatorname{arctanh} \left(\frac{c}{x} \right)^2 + b(a + b)c^2 \log(-c + x) - abc^2 \log(c + x) + b^2 c^2 \log(c + x) \right)$$

input `Integrate[x*(a + b*ArcTanh[c/x])^2,x]`

output $(2*a*b*c*x + a^2*x^2 + 2*b*x*(b*c + a*x)*\text{ArcTanh}[c/x] + b^2*(-c^2 + x^2)*\text{ArcTanh}[c/x]^2 + b*(a + b)*c^2*\text{Log}[-c + x] - a*b*c^2*\text{Log}[c + x] + b^2*c^2*\text{Log}[c + x])/2$

Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.99, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {6454, 6452, 6544, 6452, 243, 47, 14, 16, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \left(a + \text{arctanh}\left(\frac{c}{x}\right) \right)^2 dx \\
 & \quad \downarrow 6454 \\
 & - \int x^3 \left(a + \text{arctanh}\left(\frac{c}{x}\right) \right)^2 d\frac{1}{x} \\
 & \quad \downarrow 6452 \\
 & \frac{1}{2}x^2 \left(a + \text{arctanh}\left(\frac{c}{x}\right) \right)^2 - bc \int \frac{x^2 \left(a + \text{arctanh}\left(\frac{c}{x}\right) \right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} \\
 & \quad \downarrow 6544 \\
 & \frac{1}{2}x^2 \left(a + \text{arctanh}\left(\frac{c}{x}\right) \right)^2 - bc \left(c^2 \int \frac{a + \text{arctanh}\left(\frac{c}{x}\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \int x^2 \left(a + \text{arctanh}\left(\frac{c}{x}\right) \right) d\frac{1}{x} \right) \\
 & \quad \downarrow 6452 \\
 & \frac{1}{2}x^2 \left(a + \text{arctanh}\left(\frac{c}{x}\right) \right)^2 - \\
 & bc \left(c^2 \int \frac{a + \text{arctanh}\left(\frac{c}{x}\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + bc \int \frac{x}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - x \left(a + \text{arctanh}\left(\frac{c}{x}\right) \right) \right) \\
 & \quad \downarrow 243
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{2}x^2\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^2 - \\
& bc\left(c^2 \int \frac{a + \operatorname{barctanh}\left(\frac{c}{x}\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{1}{2}bc \int \frac{x}{1 - \frac{c^2}{x^2}} d\frac{1}{x^2} - x\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)\right) \\
& \quad \downarrow 47 \\
& \frac{1}{2}x^2\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^2 - \\
& bc\left(c^2 \int \frac{a + \operatorname{barctanh}\left(\frac{c}{x}\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{1}{2}bc\left(c^2 \int \frac{1}{1 - \frac{c^2}{x^2}} d\frac{1}{x^2} + \int x d\frac{1}{x^2}\right) - x\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)\right) \\
& \quad \downarrow 14 \\
& \frac{1}{2}x^2\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^2 - \\
& bc\left(c^2 \int \frac{a + \operatorname{barctanh}\left(\frac{c}{x}\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{1}{2}bc\left(c^2 \int \frac{1}{1 - \frac{c^2}{x^2}} d\frac{1}{x^2} + \log\left(\frac{1}{x^2}\right)\right) - x\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)\right) \\
& \quad \downarrow 16 \\
& \frac{1}{2}x^2\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^2 - \\
& bc\left(c^2 \int \frac{a + \operatorname{barctanh}\left(\frac{c}{x}\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - x\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right) + \frac{1}{2}bc\left(\log\left(\frac{1}{x^2}\right) - \log\left(1 - \frac{c^2}{x^2}\right)\right)\right) \\
& \quad \downarrow 6510 \\
& \frac{1}{2}x^2\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^2 - \\
& bc\left(\frac{c\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^2}{2b} - x\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right) + \frac{1}{2}bc\left(\log\left(\frac{1}{x^2}\right) - \log\left(1 - \frac{c^2}{x^2}\right)\right)\right)
\end{aligned}$$

input

```
Int [x*(a + b*ArcTanh[c/x])^2,x]
```

output

```
(x^2*(a + b*ArcTanh[c/x])^2)/2 - b*c*(-(x*(a + b*ArcTanh[c/x])) + (c*(a + b*ArcTanh[c/x])^2)/(2*b) + (b*c*(-Log[1 - c^2/x^2] + Log[x^(-2)]))/2)
```

Definitions of rubi rules used

- rule 14 $\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ ; FreeQ}[a, x]$
- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$
- rule 47 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x]$
- rule 243 $\text{Int}[(x_)^{(m_)}*((a_)+(b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] \text{ ; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$
- rule 6452 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{2n})}), x], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$
- rule 6454 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n] - 1)}*(a + b*\text{ArcTanh}[c*x])^p, x], x, x^n], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$
- rule 6510 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)]*(b_))^{(p_)}((d_)+(e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^{(p+1)}/(b*c*d*(p+1)), x] \text{ ; FreeQ}[\{a, b, c, d, e, p\}, x] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{NeQ}[p, -1]$

rule 6544

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^ (p_.)*((f_.)*(x_))^(m_)/((d_) + (
e_.)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x
], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x
^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

Maple [A] (verified)

Time = 1.51 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.42

method	result
parallelrisch	$\frac{b^2 x^2 \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{2} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right)^2 b^2 c^2}{2} + b^2 c^2 \ln(x - c) + x^2 a b \operatorname{arctanh}\left(\frac{c}{x}\right) + x \operatorname{arctanh}\left(\frac{c}{x}\right) b$
parts	$\frac{a^2 x^2}{2} - b^2 c^2 \left(-\frac{x^2 \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{2c^2} - \frac{x \operatorname{arctanh}\left(\frac{c}{x}\right)}{c} + \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(1 + \frac{c}{x}\right)}{2} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(\frac{c}{x} - 1\right)}{2} + \frac{\ln\left(\frac{c}{x} - 1\right)}{2} \right)$
derivativedivides	$-c^2 \left(-\frac{a^2 x^2}{2c^2} + b^2 \left(-\frac{x^2 \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{2c^2} - \frac{x \operatorname{arctanh}\left(\frac{c}{x}\right)}{c} + \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(1 + \frac{c}{x}\right)}{2} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(\frac{c}{x} - 1\right)}{2} + \frac{\ln\left(\frac{c}{x} - 1\right)}{2} \right) \right)$
default	$-c^2 \left(-\frac{a^2 x^2}{2c^2} + b^2 \left(-\frac{x^2 \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{2c^2} - \frac{x \operatorname{arctanh}\left(\frac{c}{x}\right)}{c} + \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(1 + \frac{c}{x}\right)}{2} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(\frac{c}{x} - 1\right)}{2} + \frac{\ln\left(\frac{c}{x} - 1\right)}{2} \right) \right)$
risch	Expression too large to display

input

```
int(x*(a+b*arctanh(c/x))^2,x,method=_RETURNVERBOSE)
```

output

```
1/2*b^2*x^2*arctanh(c/x)^2-1/2*arctanh(c/x)^2*b^2*c^2+b^2*c^2*ln(x-c)+x^2*
a*b*arctanh(c/x)+x*arctanh(c/x)*b^2*c-arctanh(c/x)*a*b*c^2+arctanh(c/x)*b^
2*c^2+1/2*a^2*x^2+a*b*c*x+1/2*a^2*c^2
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.34

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = abcx + \frac{1}{2} a^2 x^2 - \frac{1}{2} (ab - b^2) c^2 \log(c + x) \\ + \frac{1}{2} (ab + b^2) c^2 \log(-c + x) \\ - \frac{1}{8} (b^2 c^2 - b^2 x^2) \log \left(-\frac{c + x}{c - x} \right)^2 \\ + \frac{1}{2} (b^2 cx + abx^2) \log \left(-\frac{c + x}{c - x} \right)$$

input `integrate(x*(a+b*arctanh(c/x))^2,x, algorithm="fricas")`output `a*b*c*x + 1/2*a^2*x^2 - 1/2*(a*b - b^2)*c^2*log(c + x) + 1/2*(a*b + b^2)*c^2*log(-c + x) - 1/8*(b^2*c^2 - b^2*x^2)*log(-(c + x)/(c - x))^2 + 1/2*(b^2*c*x + a*b*x^2)*log(-(c + x)/(c - x))`**Sympy [A] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.25

$$\int x \left(a + b \operatorname{atanh} \left(\frac{c}{x} \right) \right)^2 dx = \frac{a^2 x^2}{2} - abc^2 \operatorname{atanh} \left(\frac{c}{x} \right) + abcx + abx^2 \operatorname{atanh} \left(\frac{c}{x} \right) \\ + b^2 c^2 \log(-c + x) - \frac{b^2 c^2 \operatorname{atanh}^2 \left(\frac{c}{x} \right)}{2} \\ + b^2 c^2 \operatorname{atanh} \left(\frac{c}{x} \right) + b^2 cx \operatorname{atanh} \left(\frac{c}{x} \right) + \frac{b^2 x^2 \operatorname{atanh}^2 \left(\frac{c}{x} \right)}{2}$$

input `integrate(x*(a+b*atanh(c/x))**2,x)`output `a**2*x**2/2 - a*b*c**2*atanh(c/x) + a*b*c*x + a*b*x**2*atanh(c/x) + b**2*c**2*log(-c + x) - b**2*c**2*atanh(c/x)**2/2 + b**2*c**2*atanh(c/x) + b**2*c*x*atanh(c/x) + b**2*x**2*atanh(c/x)**2/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.64

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \frac{1}{2} b^2 x^2 \operatorname{arctanh} \left(\frac{c}{x} \right)^2 + \frac{1}{2} a^2 x^2 + \frac{1}{2} \left(2 x^2 \operatorname{arctanh} \left(\frac{c}{x} \right) - (c \log(c+x) - c \log(-c+x) - 2x)c \right) ab + \frac{1}{8} \left((\log(c+x))^2 - 2(\log(c+x) - 2) \log(-c+x) + \log(-c+x)^2 + 4 \log(c+x) \right) c^2 - 4(c \log(c+x) - c \log(-c+x)) c$$

input `integrate(x*(a+b*arctanh(c/x))^2,x, algorithm="maxima")`

output `1/2*b^2*x^2*arctanh(c/x)^2 + 1/2*a^2*x^2 + 1/2*(2*x^2*arctanh(c/x) - (c*log(c+x) - c*log(-c+x) - 2*x)*c)*a*b + 1/8*((log(c+x)^2 - 2*(log(c+x) - 2)*log(-c+x) + log(-c+x)^2 + 4*log(c+x))*c^2 - 4*(c*log(c+x) - c*log(-c+x) - 2*x)*c*arctanh(c/x))*b^2`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 268 vs. 2(77) = 154.

Time = 0.12 (sec) , antiderivative size = 268, normalized size of antiderivative = 3.23

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \frac{2 b^2 c^3 \log \left(-\frac{c+x}{c-x} - 1 \right) - 2 b^2 c^3 \log \left(-\frac{c+x}{c-x} \right) + \frac{b^2 (c+x) c^3 \log \left(\frac{-c+x}{c-x} \right)^2}{(c-x) \left(\frac{(c+x)^2}{(c-x)^2} + \frac{2(c+x)}{c-x} + 1 \right)} + \frac{2 \left(b^2 c^3 + \frac{2 a b (c+x) c^3}{c-x} + \frac{b^2 (c+x) c^3}{c-x} \right) \log \left(-\frac{c+x}{c-x} \right)}{\frac{(c+x)^2}{(c-x)^2} + \frac{2(c+x)}{c-x} + 1}}{2 c}$$

input `integrate(x*(a+b*arctanh(c/x))^2,x, algorithm="giac")`

output `-1/2*(2*b^2*c^3*log(-(c+x)/(c-x) - 1) - 2*b^2*c^3*log(-(c+x)/(c-x)) + b^2*(c+x)*c^3*log(-(c+x)/(c-x))^2/((c-x)*((c+x)^2/(c-x)^2 + 2*(c+x)/(c-x) + 1)) + 2*(b^2*c^3 + 2*a*b*(c+x)*c^3/(c-x) + b^2*(c+x)*c^3/(c-x))*log(-(c+x)/(c-x))/((c+x)^2/(c-x)^2 + 2*(c+x)/(c-x) + 1) + 4*(a*b*c^3 + a^2*(c+x)*c^3/(c-x) + a*b*(c+x)*c^3/(c-x))/((c+x)^2/(c-x)^2 + 2*(c+x)/(c-x) + 1))/c`

Mupad [B] (verification not implemented)

Time = 3.47 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.22

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = \frac{a^2 x^2}{2} - \frac{b^2 c^2 \operatorname{atanh} \left(\frac{c}{x} \right)^2}{2} + \frac{b^2 x^2 \operatorname{atanh} \left(\frac{c}{x} \right)^2}{2} + \frac{b^2 c^2 \ln(x^2 - c^2)}{2} - a b c^2 \operatorname{atanh} \left(\frac{c}{x} \right) + a b x^2 \operatorname{atanh} \left(\frac{c}{x} \right) + b^2 c x \operatorname{atanh} \left(\frac{c}{x} \right) + a b c x$$

input `int(x*(a + b*atanh(c/x))^2,x)`output `(a^2*x^2)/2 - (b^2*c^2*atanh(c/x)^2)/2 + (b^2*x^2*atanh(c/x)^2)/2 + (b^2*c^2*log(x^2 - c^2))/2 - a*b*c^2*atanh(c/x) + a*b*x^2*atanh(c/x) + b^2*c*x*atanh(c/x) + a*b*c*x`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.35

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx = -\frac{\operatorname{atanh} \left(\frac{c}{x} \right)^2 b^2 c^2}{2} + \frac{\operatorname{atanh} \left(\frac{c}{x} \right)^2 b^2 x^2}{2} - \operatorname{atanh} \left(\frac{c}{x} \right) a b c^2 + \operatorname{atanh} \left(\frac{c}{x} \right) a b x^2 - \operatorname{atanh} \left(\frac{c}{x} \right) b^2 c^2 + \operatorname{atanh} \left(\frac{c}{x} \right) b^2 c x + \log(-c - x) b^2 c^2 + \frac{a^2 x^2}{2} + a b c x$$

input `int(x*(a+b*atanh(c/x))^2,x)`output `(- atanh(c/x)**2*b**2*c**2 + atanh(c/x)**2*b**2*x**2 - 2*atanh(c/x)*a*b*c**2 + 2*atanh(c/x)*a*b*x**2 - 2*atanh(c/x)*b**2*c**2 + 2*atanh(c/x)*b**2*c*x + 2*log(-c - x)*b**2*c**2 + a**2*x**2 + 2*a*b*c*x)/2`

3.146 $\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2 dx$

Optimal result	1194
Mathematica [A] (verified)	1194
Rubi [A] (verified)	1195
Maple [B] (verified)	1198
Fricas [F]	1198
Sympy [F]	1199
Maxima [F]	1199
Giac [F]	1199
Mupad [F(-1)]	1200
Reduce [F]	1200

Optimal result

Integrand size = 12, antiderivative size = 74

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2 dx = c \left(a + b \operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^2 + x \left(a + b \operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^2 - 2bc \left(a + b \operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right) \log\left(\frac{2c}{c-x}\right) - b^2 c \operatorname{PolyLog}\left(2, 1 - \frac{2c}{c-x}\right)$$

output `c*(a+b*arccoth(x/c))^2+x*(a+b*arccoth(x/c))^2-2*b*c*(a+b*arccoth(x/c))*ln(2*c/(c-x))-b^2*c*polylog(2,1-2*c/(c-x))`

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.31

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2 dx = b^2(-c + x) \operatorname{arctanh}\left(\frac{c}{x}\right)^2 + 2b \operatorname{arctanh}\left(\frac{c}{x}\right) \left(ax - bc \log\left(1 - e^{-2 \operatorname{arctanh}\left(\frac{c}{x}\right)}\right)\right) + a \left(ax + bc \log\left(1 - \frac{c^2}{x^2}\right) - 2bc \log\left(\frac{c}{x}\right)\right) + b^2 c \operatorname{PolyLog}\left(2, e^{-2 \operatorname{arctanh}\left(\frac{c}{x}\right)}\right)$$

input `Integrate[(a + b*ArcTanh[c/x])^2,x]`

output `b^2*(-c + x)*ArcTanh[c/x]^2 + 2*b*ArcTanh[c/x]*(a*x - b*c*Log[1 - E^(-2*ArcTanh[c/x])]) + a*(a*x + b*c*Log[1 - c^2/x^2] - 2*b*c*Log[c/x]) + b^2*c*PolyLog[2, E^(-2*ArcTanh[c/x])]`

Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.08, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6440, 6437, 27, 6547, 27, 6471, 27, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 dx \\
 & \quad \downarrow 6440 \\
 & \int \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right)^2 dx \\
 & \quad \downarrow 6437 \\
 & x \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right)^2 - \frac{2b \int \frac{c^2 x (a + b \operatorname{coth}^{-1}(\frac{x}{c}))}{c^2 - x^2} dx}{c} \\
 & \quad \downarrow 27 \\
 & x \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right)^2 - 2bc \int \frac{x (a + b \operatorname{coth}^{-1}(\frac{x}{c}))}{c^2 - x^2} dx \\
 & \quad \downarrow 6547 \\
 & x \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right)^2 - 2bc \left(\frac{\int \frac{c(a + b \operatorname{coth}^{-1}(\frac{x}{c}))}{c - x} dx}{c} - \frac{(a + b \operatorname{coth}^{-1}(\frac{x}{c}))^2}{2b} \right) \\
 & \quad \downarrow 27 \\
 & x \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right)^2 - 2bc \left(\int \frac{a + b \operatorname{coth}^{-1}(\frac{x}{c})}{c - x} dx - \frac{(a + b \operatorname{coth}^{-1}(\frac{x}{c}))^2}{2b} \right)
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 6471 \\
& x \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2 - \\
2bc & \left(-\frac{b \int \frac{c^2 \log \left(\frac{2c}{c-x} \right) dx}{c^2 - x^2} - \frac{\left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2}{2b} + \log \left(\frac{2c}{c-x} \right) \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)}{c} \right) \\
& \downarrow 27 \\
& x \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2 - \\
2bc & \left(-bc \int \frac{\log \left(\frac{2c}{c-x} \right)}{c^2 - x^2} dx - \frac{\left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2}{2b} + \log \left(\frac{2c}{c-x} \right) \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right) \right) \\
& \downarrow 2849 \\
& x \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2 - \\
2bc & \left(bc \int \frac{\log \left(\frac{2c}{c-x} \right)}{1 - \frac{2c}{c-x}} d \frac{1}{c-x} - \frac{\left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2}{2b} + \log \left(\frac{2c}{c-x} \right) \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right) \right) \\
& \downarrow 2752 \\
& x \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2 - \\
2bc & \left(-\frac{\left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2}{2b} + \log \left(\frac{2c}{c-x} \right) \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right) + \frac{1}{2} b \operatorname{PolyLog} \left(2, 1 - \frac{2c}{c-x} \right) \right)
\end{aligned}$$

input `Int[(a + b*ArcTanh[c/x])^2,x]`

output `x*(a + b*ArcCoth[x/c])^2 - 2*b*c*(-1/2*(a + b*ArcCoth[x/c])^2/b + (a + b*ArcCoth[x/c])*Log[(2*c)/(c - x)] + (b*PolyLog[2, 1 - (2*c)/(c - x)])/2)`

Defintions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 2752 $\text{Int}[\text{Log}[(c_*)(x_)]/((d_) + (e_*)(x_)), x_Symbol] \rightarrow \text{Simp}[(-e^{(-1)})*\text{PolyLog}[2, 1 - c*x], x] /; \text{FreeQ}[\{c, d, e\}, x] \ \&\& \ \text{EqQ}[e + c*d, 0]$
- rule 2849 $\text{Int}[\text{Log}[(c_)]/((d_) + (e_*)(x_))]/((f_) + (g_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[-e/g \text{ Subst}[\text{Int}[\text{Log}[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; \text{FreeQ}[\{c, d, e, f, g\}, x] \ \&\& \ \text{EqQ}[c, 2*d] \ \&\& \ \text{EqQ}[e^2*f + d^2*g, 0]$
- rule 6437 $\text{Int}[(a_.) + \text{ArcCoth}[(c_*)(x_)^(n_)]*(b_.)]^(p_), x_Symbol] \rightarrow \text{Simp}[x*(a + b*\text{ArcCoth}[c*x^n])^p, x] - \text{Simp}[b*c*n*p \text{ Int}[x^n*((a + b*\text{ArcCoth}[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; \text{FreeQ}[\{a, b, c, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[n, 1] \ || \ \text{EqQ}[p, 1])$
- rule 6440 $\text{Int}[(a_.) + \text{ArcTanh}[(c_*)(x_)^(n_)]*(b_.)]^(p_), x_Symbol] \rightarrow \text{Int}[(a + b*\text{ArcCoth}[1/(x^n*c)])^p, x] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{ILtQ}[n, 0]$
- rule 6471 $\text{Int}[(a_.) + \text{ArcCoth}[(c_*)(x_)]*(b_.)]^(p_)/((d_) + (e_*)(x_)), x_Symbol] \rightarrow \text{Simp}[(-a + b*\text{ArcCoth}[c*x])^p*(\text{Log}[2/(1 + e*(x/d))]/e), x] + \text{Simp}[b*c*(p/e) \text{ Int}[(a + b*\text{ArcCoth}[c*x])^(p - 1)*(\text{Log}[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d^2 - e^2, 0]$
- rule 6547 $\text{Int}[(((a_.) + \text{ArcCoth}[(c_*)(x_)]*(b_.)]^(p_))*(x_)/((d_) + (e_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcCoth}[c*x])^(p + 1)/(b*e*(p + 1)), x] + \text{Simp}[1/(c*d) \text{ Int}[(a + b*\text{ArcCoth}[c*x])^p/(1 - c*x), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{IGtQ}[p, 0]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 234 vs. $2(74) = 148$.

Time = 0.81 (sec) , antiderivative size = 235, normalized size of antiderivative = 3.18

method	result
parts	$xa^2 - b^2c \left(-\frac{x \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{c} - \operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(\frac{c}{x} - 1\right) + 2 \ln\left(\frac{c}{x}\right) \operatorname{arctanh}\left(\frac{c}{x}\right) - \operatorname{arctanh}\left(\frac{c}{x}\right) \right)$
derivativedivides	$-c \left(-\frac{a^2x}{c} + b^2 \left(-\frac{x \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{c} - \operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(\frac{c}{x} - 1\right) + 2 \ln\left(\frac{c}{x}\right) \operatorname{arctanh}\left(\frac{c}{x}\right) - \operatorname{arctanh}\left(\frac{c}{x}\right) \right) \right)$
default	$-c \left(-\frac{a^2x}{c} + b^2 \left(-\frac{x \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{c} - \operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(\frac{c}{x} - 1\right) + 2 \ln\left(\frac{c}{x}\right) \operatorname{arctanh}\left(\frac{c}{x}\right) - \operatorname{arctanh}\left(\frac{c}{x}\right) \right) \right)$
risch	Expression too large to display

input `int((a+b*arctanh(c/x))^2,x,method=_RETURNVERBOSE)`

output `x*a^2-b^2*c*(-1/c*x*arctanh(c/x)^2-arctanh(c/x)*ln(c/x-1)+2*ln(c/x)*arctanh(c/x)-arctanh(c/x)*ln(1+c/x)+dilog(1/2*c/x+1/2)+1/2*ln(c/x-1)*ln(1/2*c/x+1/2)-1/4*ln(c/x-1)^2+1/4*ln(1+c/x)^2-1/2*(ln(1+c/x)-ln(1/2*c/x+1/2))*ln(-1/2*c/x+1/2)-dilog(c/x)-dilog(1+c/x)-ln(c/x)*ln(1+c/x))-2*a*b*c*(-1/c*x*arctanh(c/x)-1/2*ln(c/x-1)-1/2*ln(1+c/x)+ln(c/x))`

Fricas [F]

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right)^2 dx = \int \left(b \operatorname{arctanh}\left(\frac{c}{x}\right) + a \right)^2 dx$$

input `integrate((a+b*arctanh(c/x))^2,x, algorithm="fricas")`

output `integral(b^2*arctanh(c/x)^2 + 2*a*b*arctanh(c/x) + a^2, x)`

Sympy [F]

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right)^2 dx = \int \left(a + b \operatorname{atanh}\left(\frac{c}{x}\right) \right)^2 dx$$

input `integrate((a+b*atanh(c/x))**2,x)`

output `Integral((a + b*atanh(c/x))**2, x)`

Maxima [F]

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right)^2 dx = \int \left(b \operatorname{artanh}\left(\frac{c}{x}\right) + a \right)^2 dx$$

input `integrate((a+b*arctanh(c/x))^2,x, algorithm="maxima")`

output `(2*x*arctanh(c/x) + c*log(-c^2 + x^2))*a*b + 1/4*(x*log(c + x)^2 - 2*(c + x)*log(c + x)*log(-c + x) - (c - x)*log(-c + x)^2 + integrate(-2*(c^2 + 3*c*x)*log(c + x)/(c^2 - x^2), x))*b^2 + a^2*x`

Giac [F]

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right)^2 dx = \int \left(b \operatorname{artanh}\left(\frac{c}{x}\right) + a \right)^2 dx$$

input `integrate((a+b*arctanh(c/x))^2,x, algorithm="giac")`

output `integrate((b*arctanh(c/x) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right)^2 dx = \int \left(a + b \operatorname{atanh}\left(\frac{c}{x}\right) \right)^2 dx$$

input `int((a + b*atanh(c/x))^2,x)`output `int((a + b*atanh(c/x))^2, x)`**Reduce [F]**

$$\begin{aligned} \int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right)^2 dx &= -2 \operatorname{atanh}\left(\frac{c}{x}\right) abc + 2 \operatorname{atanh}\left(\frac{c}{x}\right) abx \\ &+ \left(\int \operatorname{atanh}\left(\frac{c}{x}\right)^2 dx \right) b^2 + 2 \log(-c - x) abc + a^2 x \end{aligned}$$

input `int((a+b*atanh(c/x))^2,x)`output `- 2*atanh(c/x)*a*b*c + 2*atanh(c/x)*a*b*x + int(atanh(c/x)**2,x)*b**2 + 2*log(-c-x)*a*b*c + a**2*x`

$$3.147 \quad \int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2}{x} dx$$

Optimal result	1201
Mathematica [C] (verified)	1202
Rubi [A] (verified)	1202
Maple [C] (warning: unable to verify)	1205
Fricas [F]	1206
Sympy [F]	1206
Maxima [F]	1207
Giac [F]	1207
Mupad [F(-1)]	1207
Reduce [F]	1208

Optimal result

Integrand size = 16, antiderivative size = 133

$$\begin{aligned} \int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2}{x} dx = & -2 \left(a + b \operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^2 \operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{c}{x}}\right) \\ & + b \left(a + b \operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{c}{x}}\right) \\ & - b \left(a + b \operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - \frac{c}{x}}\right) \\ & - \frac{1}{2} b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - \frac{c}{x}}\right) \\ & + \frac{1}{2} b^2 \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 - \frac{c}{x}}\right) \end{aligned}$$

output

```
2*(a+b*arccoth(x/c))^2*arctanh(-1+2/(1-c/x))+b*(a+b*arccoth(x/c))*polylog(
2,1-2/(1-c/x))-b*(a+b*arccoth(x/c))*polylog(2,-1+2/(1-c/x))-1/2*b^2*polylo
g(3,1-2/(1-c/x))+1/2*b^2*polylog(3,-1+2/(1-c/x))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.33

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x} dx = a^2 \log(x) + ab \left(\operatorname{PolyLog}\left(2, -\frac{c}{x}\right) - \operatorname{PolyLog}\left(2, \frac{c}{x}\right) \right) + b^2 \left(-\frac{i\pi^3}{24} + \frac{2}{3} \operatorname{arctanh}\left(\frac{c}{x}\right)^3 + \operatorname{arctanh}\left(\frac{c}{x}\right)^2 \log\left(1 + e^{-2\operatorname{arctanh}(\frac{c}{x})}\right) - \operatorname{arctanh}\left(\frac{c}{x}\right)^2 \log\left(1 - e^{2\operatorname{arctanh}(\frac{c}{x})}\right) - \operatorname{arctanh}\left(\frac{c}{x}\right) \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arctanh}(\frac{c}{x})}\right) - \operatorname{arctanh}\left(\frac{c}{x}\right) \operatorname{PolyLog}\left(2, e^{2\operatorname{arctanh}(\frac{c}{x})}\right) - \frac{1}{2} \operatorname{PolyLog}\left(3, -e^{-2\operatorname{arctanh}(\frac{c}{x})}\right) + \frac{1}{2} \operatorname{PolyLog}\left(3, e^{2\operatorname{arctanh}(\frac{c}{x})}\right) \right)$$

input `Integrate[(a + b*ArcTanh[c/x])^2/x,x]`

output `a^2*Log[x] + a*b*(PolyLog[2, -(c/x)] - PolyLog[2, c/x]) + b^2*((-1/24*I)*Pi^3 + (2*ArcTanh[c/x]^3)/3 + ArcTanh[c/x]^2*Log[1 + E^(-2*ArcTanh[c/x])] - ArcTanh[c/x]^2*Log[1 - E^(2*ArcTanh[c/x])] - ArcTanh[c/x]*PolyLog[2, -E^(-2*ArcTanh[c/x])] - ArcTanh[c/x]*PolyLog[2, E^(2*ArcTanh[c/x])] - PolyLog[3, -E^(-2*ArcTanh[c/x])]/2 + PolyLog[3, E^(2*ArcTanh[c/x])]/2)`

Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.20, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6450, 6448, 6614, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{(a + \operatorname{barctanh}(\frac{c}{x}))^2}{x} dx \\
& \quad \downarrow \text{6450} \\
& - \int x \left(a + \operatorname{barctanh}(\frac{c}{x}) \right)^2 d\frac{1}{x} \\
& \quad \downarrow \text{6448} \\
& 4bc \int \frac{\operatorname{arctanh}\left(1 - \frac{2}{1-\frac{c}{x}}\right) \left(a + \operatorname{barctanh}(\frac{c}{x}) \right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - 2\operatorname{arctanh}\left(1 - \frac{2}{1-\frac{c}{x}}\right) \left(a + \operatorname{barctanh}(\frac{c}{x}) \right)^2 \\
& \quad \downarrow \text{6614} \\
& 4bc \left(\frac{1}{2} \int \frac{\left(a + \operatorname{barctanh}(\frac{c}{x}) \right) \log\left(2 - \frac{2}{1-\frac{c}{x}}\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - \frac{1}{2} \int \frac{\left(a + \operatorname{barctanh}(\frac{c}{x}) \right) \log\left(\frac{2}{1-\frac{c}{x}}\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} \right) - \\
& \quad 2\operatorname{arctanh}\left(1 - \frac{2}{1-\frac{c}{x}}\right) \left(a + \operatorname{barctanh}(\frac{c}{x}) \right)^2 \\
& \quad \downarrow \text{6620} \\
& 4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-\frac{c}{x}}\right) \left(a + \operatorname{barctanh}(\frac{c}{x}) \right)}{2c} - \frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-\frac{c}{x}}\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} \right) + \frac{1}{2} \left(\frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, \frac{2}{1-\frac{c}{x}}\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} \right) \right) - \\
& \quad 2\operatorname{arctanh}\left(1 - \frac{2}{1-\frac{c}{x}}\right) \left(a + \operatorname{barctanh}(\frac{c}{x}) \right)^2 \\
& \quad \downarrow \text{7164} \\
& 4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-\frac{c}{x}}\right) \left(a + \operatorname{barctanh}(\frac{c}{x}) \right)}{2c} - \frac{b \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-\frac{c}{x}}\right)}{4c} \right) + \frac{1}{2} \left(\frac{b \operatorname{PolyLog}\left(3, \frac{2}{1-\frac{c}{x}} - 1\right)}{4c} \right) \right) - \\
& \quad 2\operatorname{arctanh}\left(1 - \frac{2}{1-\frac{c}{x}}\right) \left(a + \operatorname{barctanh}(\frac{c}{x}) \right)^2
\end{aligned}$$

input `Int[(a + b*ArcTanh[c/x])^2/x,x]`

output

```
-2*ArcTanh[1 - 2/(1 - c/x)]*(a + b*ArcTanh[c/x])^2 + 4*b*c*(((a + b*ArcTanh[c/x])*PolyLog[2, 1 - 2/(1 - c/x)]/(2*c) - (b*PolyLog[3, 1 - 2/(1 - c/x)])/(4*c))/2 + (-1/2*((a + b*ArcTanh[c/x])*PolyLog[2, -1 + 2/(1 - c/x)]/c + (b*PolyLog[3, -1 + 2/(1 - c/x)]/(4*c))/2)
```

Defintions of rubi rules used

rule 6448

```
Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_)/(x_), x_Symbol] := Simp[2*(a + b*ArcTanh[c*x])^p*ArcTanh[1 - 2/(1 - c*x)], x] - Simp[2*b*c*p Int[(a + b*ArcTanh[c*x])^(p - 1)*(ArcTanh[1 - 2/(1 - c*x)]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1]
```

rule 6450

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_)/(x_), x_Symbol] := Simp[1/n Subst[Int[(a + b*ArcTanh[c*x])^p/x, x], x, x^n], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]
```

rule 6614

```
Int[(ArcTanh[u_] * ((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/2 Int[Log[1 + u]*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] - Simp[1/2 Int[Log[1 - u]*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[u^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 6620

```
Int[(Log[u_] * ((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(-(a + b*ArcTanh[c*x])^p)*(PolyLog[2, 1 - u]/(2*c*d)), x] + Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v, x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 13.60 (sec) , antiderivative size = 704, normalized size of antiderivative = 5.29

method	result
parts	$a^2 \ln(x) + b^2 \left(-\ln\left(\frac{c}{x}\right) \operatorname{arctanh}\left(\frac{c}{x}\right)^2 + \operatorname{arctanh}\left(\frac{c}{x}\right) \operatorname{polylog}\left(2, -\frac{(1+\frac{c}{x})^2}{1-\frac{c^2}{x^2}}\right) - \frac{\operatorname{polylog}\left(3, -\frac{(1+\frac{c}{x})^2}{1-\frac{c^2}{x^2}}\right)}{2} \right)$
derivativedivides	$-a^2 \ln\left(\frac{c}{x}\right) - b^2 \left(\ln\left(\frac{c}{x}\right) \operatorname{arctanh}\left(\frac{c}{x}\right)^2 - \operatorname{arctanh}\left(\frac{c}{x}\right) \operatorname{polylog}\left(2, -\frac{(1+\frac{c}{x})^2}{1-\frac{c^2}{x^2}}\right) + \frac{\operatorname{polylog}\left(3, -\frac{(1+\frac{c}{x})^2}{1-\frac{c^2}{x^2}}\right)}{2} \right)$
default	$-a^2 \ln\left(\frac{c}{x}\right) - b^2 \left(\ln\left(\frac{c}{x}\right) \operatorname{arctanh}\left(\frac{c}{x}\right)^2 - \operatorname{arctanh}\left(\frac{c}{x}\right) \operatorname{polylog}\left(2, -\frac{(1+\frac{c}{x})^2}{1-\frac{c^2}{x^2}}\right) + \frac{\operatorname{polylog}\left(3, -\frac{(1+\frac{c}{x})^2}{1-\frac{c^2}{x^2}}\right)}{2} \right)$

input

```
int((a+b*arctanh(c/x))^2/x,x,method=_RETURNVERBOSE)
```

output

```
a^2*ln(x)+b^2*(-ln(c/x)*arctanh(c/x)^2+arctanh(c/x)*polylog(2,-(1+c/x)^2/(1-c^2/x^2))-1/2*polylog(3,-(1+c/x)^2/(1-c^2/x^2))+arctanh(c/x)^2*ln((1+c/x)^2/(1-c^2/x^2)-1)-arctanh(c/x)^2*ln(1-(1+c/x)/(1-c^2/x^2)^(1/2))-2*arctanh(c/x)*polylog(2,(1+c/x)/(1-c^2/x^2)^(1/2))+2*polylog(3,(1+c/x)/(1-c^2/x^2)^(1/2))-arctanh(c/x)^2*ln(1+(1+c/x)/(1-c^2/x^2)^(1/2))-2*arctanh(c/x)*polylog(2,-(1+c/x)/(1-c^2/x^2)^(1/2))+2*polylog(3,-(1+c/x)/(1-c^2/x^2)^(1/2))-1/2*I*Pi*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))*(csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1))*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))-csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1))*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))-csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))+csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^2)*arctanh(c/x)^2+2*a*b*(-ln(c/x)*arctanh(c/x)+1/2*dilog(c/x)+1/2*dilog(1+c/x)+1/2*ln(c/x)*ln(1+c/x))
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x} dx = \int \frac{(b \operatorname{arctanh}(\frac{c}{x}) + a)^2}{x} dx$$

input

```
integrate((a+b*arctanh(c/x))^2/x,x, algorithm="fricas")
```

output

```
integral((b^2*arctanh(c/x)^2 + 2*a*b*arctanh(c/x) + a^2)/x, x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x}))^2}{x} dx$$

input

```
integrate((a+b*atanh(c/x))**2/x,x)
```

output

```
Integral((a + b*atanh(c/x))**2/x, x)
```

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x}) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c/x))^2/x,x, algorithm="maxima")`

output `a^2*log(x) + integrate(1/4*b^2*(log(c/x + 1) - log(-c/x + 1))^2/x + a*b*(log(c/x + 1) - log(-c/x + 1))/x, x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x}) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c/x))^2/x,x, algorithm="giac")`

output `integrate((b*arctanh(c/x) + a)^2/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x}))^2}{x} dx$$

input `int((a + b*atanh(c/x))^2/x,x)`

output `int((a + b*atanh(c/x))^2/x, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x} dx = 2 \left(\int \frac{\operatorname{atanh}(\frac{c}{x})}{x} dx \right) ab + \left(\int \frac{\operatorname{atanh}(\frac{c}{x})^2}{x} dx \right) b^2 + \log(x) a^2$$

input `int((a+b*atanh(c/x))^2/x,x)`

output `2*int(atanh(c/x)/x,x)*a*b + int(atanh(c/x)**2/x,x)*b**2 + log(x)*a**2`

3.148 $\int \frac{\left(a+b\operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2}{x^2} dx$

Optimal result	1209
Mathematica [A] (verified)	1209
Rubi [A] (verified)	1210
Maple [A] (verified)	1213
Fricas [F]	1213
Sympy [F]	1214
Maxima [F]	1214
Giac [F]	1215
Mupad [F(-1)]	1215
Reduce [F]	1215

Optimal result

Integrand size = 16, antiderivative size = 87

$$\int \frac{\left(a+b\operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2}{x^2} dx = -\frac{\left(a+b\operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^2}{c} - \frac{\left(a+b\operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^2}{x} + \frac{2b\left(a+b\operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)\log\left(\frac{2}{1-\frac{c}{x}}\right)}{c} + \frac{b^2\operatorname{PolyLog}\left(2,1-\frac{2}{1-\frac{c}{x}}\right)}{c}$$

output `-(a+b*arccoth(x/c))^2/c-(a+b*arccoth(x/c))^2/x+2*b*(a+b*arccoth(x/c))*ln(2/(1-c/x))/c+b^2*polylog(2,1-2/(1-c/x))/c`

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.16

$$\int \frac{\left(a+b\operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2}{x^2} dx = \frac{b^2(-c+x)\operatorname{arctanh}\left(\frac{c}{x}\right)^2 + 2b\operatorname{arctanh}\left(\frac{c}{x}\right)\left(-ac+bx\log\left(1+e^{-2\operatorname{arctanh}\left(\frac{c}{x}\right)}\right)\right) + a\left(-ac+2bx\log\left(\frac{1}{\sqrt{1-\frac{c^2}{x^2}}}\right)\right)}{cx}$$

input `Integrate[(a + b*ArcTanh[c/x])^2/x^2,x]`

output `(b^2*(-c + x)*ArcTanh[c/x]^2 + 2*b*ArcTanh[c/x]*(-(a*c) + b*x*Log[1 + E^(-2*ArcTanh[c/x])]) + a*(-(a*c) + 2*b*x*Log[1/Sqrt[1 - c^2/x^2]]) - b^2*x*PolyLog[2, -E^(-2*ArcTanh[c/x])])/(c*x)`

Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.16, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6454, 6436, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^2} dx \\
 & \quad \downarrow \text{6454} \\
 & - \int (a + b \operatorname{arctanh}(\frac{c}{x}))^2 d\frac{1}{x} \\
 & \quad \downarrow \text{6436} \\
 & 2bc \int \frac{a + b \operatorname{arctanh}(\frac{c}{x})}{(1 - \frac{c^2}{x^2})x} d\frac{1}{x} - \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x} \\
 & \quad \downarrow \text{6546} \\
 & 2bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(\frac{c}{x})}{1 - \frac{c}{x}} d\frac{1}{x}}{c} - \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{2bc^2} \right) - \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x} \\
 & \quad \downarrow \text{6470}
 \end{aligned}$$

$$\begin{aligned}
 & 2bc \left(\frac{\frac{\log\left(\frac{2}{1-\frac{c}{x}}\right)(a+b\operatorname{arctanh}\left(\frac{c}{x}\right))}{c} - b \int \frac{\log\left(\frac{2}{1-\frac{c}{x}}\right) d\frac{1}{x}}{1-\frac{c^2}{x^2}}}{c} - \frac{(a+b\operatorname{arctanh}\left(\frac{c}{x}\right))^2}{2bc^2} \right) - \\
 & \qquad \qquad \qquad \frac{(a+b\operatorname{arctanh}\left(\frac{c}{x}\right))^2}{x} \\
 & \qquad \qquad \qquad \downarrow \text{2849} \\
 & 2bc \left(\frac{b \int \frac{\log\left(\frac{2}{1-\frac{c}{x}}\right) d\frac{1}{x}}{1-\frac{c^2}{x^2}} + \frac{\log\left(\frac{2}{1-\frac{c}{x}}\right)(a+b\operatorname{arctanh}\left(\frac{c}{x}\right))}{c}}{c} - \frac{(a+b\operatorname{arctanh}\left(\frac{c}{x}\right))^2}{2bc^2} \right) - \\
 & \qquad \qquad \qquad \frac{(a+b\operatorname{arctanh}\left(\frac{c}{x}\right))^2}{x} \\
 & \qquad \qquad \qquad \downarrow \text{2752} \\
 & 2bc \left(\frac{\frac{\log\left(\frac{2}{1-\frac{c}{x}}\right)(a+b\operatorname{arctanh}\left(\frac{c}{x}\right))}{c} + \frac{b \operatorname{PolyLog}\left(2, 1-\frac{2}{1-\frac{c}{x}}\right)}{2c}}{c} - \frac{(a+b\operatorname{arctanh}\left(\frac{c}{x}\right))^2}{2bc^2} \right) - \\
 & \qquad \qquad \qquad \frac{(a+b\operatorname{arctanh}\left(\frac{c}{x}\right))^2}{x}
 \end{aligned}$$

input

```
Int[(a + b*ArcTanh[c/x])^2/x^2,x]
```

output

```
-((a + b*ArcTanh[c/x])^2/x) + 2*b*c*(-1/2*(a + b*ArcTanh[c/x])^2/(b*c^2) +
((a + b*ArcTanh[c/x])*Log[2/(1 - c/x)]/c + (b*PolyLog[2, 1 - 2/(1 - c/x)
]))/(2*c))/c)
```

Defintions of rubi rules used

rule 2752

```
Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] :> Simp[(-e^(-1))*PolyLo
g[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]
```


rule 2849 $\text{Int}[\text{Log}[(c_)/((d_)+(e_)(x_))]/((f_)+(g_)(x_)^2), x_Symbol] \rightarrow \text{Simp}[-e/g \text{ Subst}[\text{Int}[\text{Log}[2*d*x]/(1-2*d*x), x], x, 1/(d+e*x)], x] /;$ $\text{FreeQ}[\{c, d, e, f, g\}, x] \ \&\& \ \text{EqQ}[c, 2*d] \ \&\& \ \text{EqQ}[e^2*f + d^2*g, 0]$

rule 6436 $\text{Int}[(a_)+\text{ArcTanh}[(c_)(x_)]*(b_)]^{(p_)}, x_Symbol] \rightarrow \text{Simp}[x*(a + b*\text{ArcTanh}[c*x^n])^p, x] - \text{Simp}[b*c*n*p \text{ Int}[x^n*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{2*n})})], x], x] /;$ $\text{FreeQ}[\{a, b, c, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[n, 1] \ || \ \text{EqQ}[p, 1])$

rule 6454 $\text{Int}[(a_)+\text{ArcTanh}[(c_)(x_)]*(b_)]^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n]-1)*(a + b*\text{ArcTanh}[c*x])^p}, x], x, x^n], x] /;$ $\text{FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$

rule 6470 $\text{Int}[(a_)+\text{ArcTanh}[(c_)(x_)]*(b_)]^{(p_)} / ((d_)+(e_)(x_)), x_Symbol] \rightarrow \text{Simp}[(-a + b*\text{ArcTanh}[c*x])^p*(\text{Log}[2/(1+e*(x/d))]/e), x] + \text{Simp}[b*c*(p/e) \text{ Int}[(a + b*\text{ArcTanh}[c*x])^{(p-1)}*(\text{Log}[2/(1+e*(x/d))]/(1-c^2*x^2)), x], x] /;$ $\text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d^2 - e^2, 0]$

rule 6546 $\text{Int}[(a_)+\text{ArcTanh}[(c_)(x_)]*(b_)]^{(p_)}*(x_)/((d_)+(e_)(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^{(p+1)}/(b*e*(p+1)), x] + \text{Simp}[1/(c*d) \text{ Int}[(a + b*\text{ArcTanh}[c*x])^p/(1-c*x), x], x] /;$ $\text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{IGtQ}[p, 0]$

Maple [A] (verified)

Time = 1.36 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.54

method	result
derivativedivides	$-\frac{\frac{c a^2}{x} + b^2 \left(\operatorname{arctanh}\left(\frac{c}{x}\right)^2 \left(\frac{c}{x} - 1\right) + 2 \operatorname{arctanh}\left(\frac{c}{x}\right)^2 - 2 \operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(1 + \frac{\left(1 + \frac{c}{x}\right)^2}{1 - \frac{c^2}{x^2}}\right) - \operatorname{polylog}\left(2, -\frac{\left(1 + \frac{c}{x}\right)^2}{1 - \frac{c^2}{x^2}}\right) \right)}{c} + \frac{2abc}{x}$
default	$-\frac{\frac{c a^2}{x} + b^2 \left(\operatorname{arctanh}\left(\frac{c}{x}\right)^2 \left(\frac{c}{x} - 1\right) + 2 \operatorname{arctanh}\left(\frac{c}{x}\right)^2 - 2 \operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(1 + \frac{\left(1 + \frac{c}{x}\right)^2}{1 - \frac{c^2}{x^2}}\right) - \operatorname{polylog}\left(2, -\frac{\left(1 + \frac{c}{x}\right)^2}{1 - \frac{c^2}{x^2}}\right) \right)}{c} + \frac{2abc}{x}$
parts	$-\frac{a^2}{x} - \frac{b^2 \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{x} - \frac{b^2 \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{c} + \frac{2b^2 \operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(1 + \frac{\left(1 + \frac{c}{x}\right)^2}{1 - \frac{c^2}{x^2}}\right)}{c} + \frac{b^2 \operatorname{polylog}\left(2, -\frac{\left(1 + \frac{c}{x}\right)^2}{1 - \frac{c^2}{x^2}}\right)}{c}$

input `int((a+b*arctanh(c/x))^2/x^2,x,method=_RETURNVERBOSE)`

output `-1/c*(c/x*a^2+b^2*(arctanh(c/x)^2*(c/x-1)+2*arctanh(c/x)^2-2*arctanh(c/x)*ln(1+(1+c/x)^2/(1-c^2/x^2))-polylog(2,-(1+c/x)^2/(1-c^2/x^2)))+2*a*b*c/x*arctanh(c/x)+a*b*ln(1-c^2/x^2))`

Fricas [F]

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2}{x^2} dx = \int \frac{\left(b \operatorname{arctanh}\left(\frac{c}{x}\right) + a\right)^2}{x^2} dx$$

input `integrate((a+b*arctanh(c/x))^2/x^2,x, algorithm="fricas")`

output `integral((b^2*arctanh(c/x)^2 + 2*a*b*arctanh(c/x) + a^2)/x^2, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^2} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x}))^2}{x^2} dx$$

input `integrate((a+b*atanh(c/x))**2/x**2,x)`

output `Integral((a + b*atanh(c/x))**2/x**2, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^2} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x}) + a)^2}{x^2} dx$$

input `integrate((a+b*arctanh(c/x))^2/x^2,x, algorithm="maxima")`

output `1/4*(c^3*integrate(-log(x)^2/(c^3*x^2 - c*x^4), x) + c^2*(log(-c^2 + x^2)/c^3 - log(x^2)/c^3) - 4*c^2*integrate(-x*log(c + x)/(c^3*x^2 - c*x^4), x) + 2*c^2*integrate(-x*log(x)/(c^3*x^2 - c*x^4), x) + 2*c*(log(-c + x)/c^2 - log(x)/c^2 + 1/(c*x))*log(-c/x + 1) - c*(log(c + x)/c^2 - log(-c + x)/c^2) - c*integrate(-x^2*log(x)^2/(c^3*x^2 - c*x^4), x) - 2*c*integrate(-x^2*log(c + x)/(c^3*x^2 - c*x^4), x) + 4*c*integrate(-x^2*log(x)/(c^3*x^2 - c*x^4), x) - log(-c/x + 1)^2/x - (c*log(c + x)^2 - 2*((c + x)*log(c + x) - (c + x)*log(x) - c)*log(-c + x))/(c*x) - (x*log(-c + x)^2 + x*log(x)^2 - 2*(x*log(x) - x)*log(-c + x) - 2*x*log(x) + 2*c)/(c*x) - 2*integrate(-x^3*log(c + x)/(c^3*x^2 - c*x^4), x) + 2*integrate(-x^3*log(x)/(c^3*x^2 - c*x^4), x))*b^2 - a*b*(2*c*arctanh(c/x)/x + log(-c^2/x^2 + 1))/c - a^2/x`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^2} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x}) + a)^2}{x^2} dx$$

input `integrate((a+b*arctanh(c/x))^2/x^2,x, algorithm="giac")`

output `integrate((b*arctanh(c/x) + a)^2/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^2} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x}))^2}{x^2} dx$$

input `int((a + b*atanh(c/x))^2/x^2,x)`

output `int((a + b*atanh(c/x))^2/x^2, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^2} dx$$

$$= \frac{-\operatorname{atanh}(\frac{c}{x})^2 b^2 c - 2 \operatorname{atanh}(\frac{c}{x}) abc + 2 \operatorname{atanh}(\frac{c}{x}) abx + 2 \left(\int \frac{\operatorname{atanh}(\frac{c}{x})}{c^2 x - x^3} dx \right) b^2 c^2 x - 2 \log(-c - x) abx + 2 \log(cx)}{cx}$$

input `int((a+b*atanh(c/x))^2/x^2,x)`

output `(- atanh(c/x)**2*b**2*c - 2*atanh(c/x)*a*b*c + 2*atanh(c/x)*a*b*x + 2*int(atanh(c/x)/(c**2*x - x**3),x)*b**2*c**2*x - 2*log(- c - x)*a*b*x + 2*log(x)*a*b*x - a**2*c)/(c*x)`

3.149 $\int \frac{\left(a+b\operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2}{x^3} dx$

Optimal result	1216
Mathematica [A] (verified)	1216
Rubi [A] (verified)	1217
Maple [A] (verified)	1219
Fricas [A] (verification not implemented)	1219
Sympy [A] (verification not implemented)	1220
Maxima [B] (verification not implemented)	1220
Giac [B] (verification not implemented)	1221
Mupad [B] (verification not implemented)	1222
Reduce [B] (verification not implemented)	1222

Optimal result

Integrand size = 16, antiderivative size = 87

$$\int \frac{\left(a+b\operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2}{x^3} dx = -\frac{ab}{cx} - \frac{b^2 \operatorname{coth}^{-1}\left(\frac{x}{c}\right)}{cx} + \frac{\left(a+b\operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^2}{2c^2} - \frac{\left(a+b\operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^2}{2x^2} - \frac{b^2 \log\left(1-\frac{c^2}{x^2}\right)}{2c^2}$$

output

```
-a*b/c/x-b^2*arccoth(x/c)/c/x+1/2*(a+b*arccoth(x/c))^2/c^2-1/2*(a+b*arccot
h(x/c))^2/x^2-1/2*b^2*ln(1-c^2/x^2)/c^2
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.37

$$\int \frac{\left(a+b\operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2}{x^3} dx = \frac{a^2c^2 + 2abcx + 2bc(ac + bx)\operatorname{arctanh}\left(\frac{c}{x}\right) + b^2(c^2 - x^2)\operatorname{arctanh}\left(\frac{c}{x}\right)^2 - 2b^2x^2 \log(x) + abx^2 \log(-c + x)}{2c^2x^2}$$

input

```
Integrate[(a + b*ArcTanh[c/x])^2/x^3,x]
```

output

$$-1/2*(a^2*c^2 + 2*a*b*c*x + 2*b*c*(a*c + b*x)*\text{ArcTanh}[c/x] + b^2*(c^2 - x^2)*\text{ArcTanh}[c/x]^2 - 2*b^2*x^2*\text{Log}[x] + a*b*x^2*\text{Log}[-c + x] + b^2*x^2*\text{Log}[-c + x] - a*b*x^2*\text{Log}[c + x] + b^2*x^2*\text{Log}[c + x])/(c^2*x^2)$$
Rubi [A] (verified)

Time = 0.86 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6454, 6452, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^3} dx \\ & \quad \downarrow \text{6454} \\ & - \int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x} d\frac{1}{x} \\ & \quad \downarrow \text{6452} \\ & bc \int \frac{a + b \operatorname{arctanh}(\frac{c}{x})}{(1 - \frac{c^2}{x^2}) x^2} d\frac{1}{x} - \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{2x^2} \\ & \quad \downarrow \text{6542} \\ & bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(\frac{c}{x})}{1 - \frac{c^2}{x^2}} d\frac{1}{x}}{c^2} - \frac{\int (a + b \operatorname{arctanh}(\frac{c}{x})) d\frac{1}{x}}{c^2} \right) - \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{2x^2} \\ & \quad \downarrow \text{2009} \\ & bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(\frac{c}{x})}{1 - \frac{c^2}{x^2}} d\frac{1}{x}}{c^2} - \frac{\frac{a}{x} + \frac{b \operatorname{arctanh}(\frac{c}{x})}{x} + \frac{b \log(1 - \frac{c^2}{x^2})}{2c}}{c^2} \right) - \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{2x^2} \\ & \quad \downarrow \text{6510} \end{aligned}$$

$$bc \left(\frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{2bc^3} - \frac{\frac{a}{x} + \frac{b \operatorname{arctanh}(\frac{c}{x})}{x} + \frac{b \log(1 - \frac{c^2}{x^2})}{2c}}{c^2} \right) - \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{2x^2}$$

input `Int[(a + b*ArcTanh[c/x])^2/x^3,x]`

output `-1/2*(a + b*ArcTanh[c/x])^2/x^2 + b*c*((a + b*ArcTanh[c/x])^2/(2*b*c^3) - (a/x + (b*ArcTanh[c/x])/x + (b*Log[1 - c^2/x^2])/(2*c))/c^2)`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6510 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/((d_.) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6542 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((f_.)*(x_)^(m_.))/((d_.) + (e_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

Maple [A] (verified)

Time = 2.04 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.52

method	result
parallelrisc	$\frac{b^2 x^2 \operatorname{arctanh}\left(\frac{c}{x}\right)^2 - \operatorname{arctanh}\left(\frac{c}{x}\right)^2 b^2 c^2 + 2b^2 x^2 \ln(x) - 2 \ln(x-c)x^2 b^2 + 2x^2 ab \operatorname{arctanh}\left(\frac{c}{x}\right) - 2x^2 \operatorname{arctanh}\left(\frac{c}{x}\right) b^2 - 2x \operatorname{arctanh}\left(\frac{c}{x}\right) b^2 c^2}{2x^2 c^2}$
parts	$-\frac{a^2}{2x^2} - \frac{b^2 \left(\frac{c^2 \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{2x^2} + \frac{c \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} + \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(\frac{c}{x}-1\right)}{2} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(1+\frac{c}{x}\right)}{2} - \frac{\ln\left(\frac{c}{x}-1\right) \ln\left(\frac{c}{2x}+\frac{1}{2}\right)}{4} + \frac{\ln\left(\frac{c}{x}\right) \ln\left(\frac{c}{2x}+\frac{1}{2}\right)}{4} \right)}{c^2}$
derivativedivides	$-\frac{\frac{a^2 c^2}{2x^2} + b^2 \left(\frac{c^2 \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{2x^2} + \frac{c \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} + \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(\frac{c}{x}-1\right)}{2} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(1+\frac{c}{x}\right)}{2} - \frac{\ln\left(\frac{c}{x}-1\right) \ln\left(\frac{c}{2x}+\frac{1}{2}\right)}{4} + \frac{\ln\left(\frac{c}{x}\right) \ln\left(\frac{c}{2x}+\frac{1}{2}\right)}{4} \right)}{c^2}$
default	$-\frac{\frac{a^2 c^2}{2x^2} + b^2 \left(\frac{c^2 \operatorname{arctanh}\left(\frac{c}{x}\right)^2}{2x^2} + \frac{c \operatorname{arctanh}\left(\frac{c}{x}\right)}{x} + \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(\frac{c}{x}-1\right)}{2} - \frac{\operatorname{arctanh}\left(\frac{c}{x}\right) \ln\left(1+\frac{c}{x}\right)}{2} - \frac{\ln\left(\frac{c}{x}-1\right) \ln\left(\frac{c}{2x}+\frac{1}{2}\right)}{4} + \frac{\ln\left(\frac{c}{x}\right) \ln\left(\frac{c}{2x}+\frac{1}{2}\right)}{4} \right)}{c^2}$
risc	Expression too large to display

input `int((a+b*arctanh(c/x))^2/x^3,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{2} * (b^2 * x^2 * \operatorname{arctanh}(c/x)^2 - \operatorname{arctanh}(c/x)^2 * b^2 * c^2 + 2 * b^2 * x^2 * \ln(x) - 2 * \ln(x-c) * x^2 * b^2 + 2 * x^2 * a * b * \operatorname{arctanh}(c/x) - 2 * x^2 * \operatorname{arctanh}(c/x) * b^2 - 2 * x * \operatorname{arctanh}(c/x) * b^2 * c^2 - 2 * \operatorname{arctanh}(c/x) * a * b * c^2 - 2 * a * b * c * x - a^2 * c^2) / x^2 / c^2$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.49

$$\int \frac{(a + b \operatorname{arctanh}\left(\frac{c}{x}\right))^2}{x^3} dx$$

$$= \frac{8 b^2 x^2 \log(x) - 4 a^2 c^2 - 8 a b c x + 4 (a b - b^2) x^2 \log(c + x) - 4 (a b + b^2) x^2 \log(-c + x) - (b^2 c^2 - b^2 x^2) \log\left(\frac{c-x}{c+x}\right)}{8 c^2 x^2}$$

input `integrate((a+b*arctanh(c/x))^2/x^3,x, algorithm="fricas")`

output
$$\frac{1}{8} * (8 * b^2 * x^2 * \log(x) - 4 * a^2 * c^2 - 8 * a * b * c * x + 4 * (a * b - b^2) * x^2 * \log(c + x) - 4 * (a * b + b^2) * x^2 * \log(-c + x) - (b^2 * c^2 - b^2 * x^2) * \log\left(\frac{c-x}{c+x}\right) - 4 * (a * b * c^2 + b^2 * c * x) * \log\left(\frac{c-x}{c+x}\right)) / (c^2 * x^2)$$

Sympy [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.43

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^3} dx$$

$$= \begin{cases} -\frac{a^2}{2x^2} - \frac{ab \operatorname{atanh}(\frac{c}{x})}{x^2} - \frac{ab}{cx} + \frac{ab \operatorname{atanh}(\frac{c}{x})}{c^2} - \frac{b^2 \operatorname{atanh}^2(\frac{c}{x})}{2x^2} - \frac{b^2 \operatorname{atanh}(\frac{c}{x})}{cx} + \frac{b^2 \log(x)}{c^2} - \frac{b^2 \log(-c+x)}{c^2} + \frac{b^2 \operatorname{atanh}^2(\frac{c}{x})}{2c^2} \\ -\frac{a^2}{2x^2} \end{cases}$$

input `integrate((a+b*atanh(c/x))**2/x**3,x)`

output `Piecewise((-a**2/(2*x**2) - a*b*atanh(c/x)/x**2 - a*b/(c*x) + a*b*atanh(c/x)/c**2 - b**2*atanh(c/x)**2/(2*x**2) - b**2*atanh(c/x)/(c*x) + b**2*log(x)/c**2 - b**2*log(-c + x)/c**2 + b**2*atanh(c/x)**2/(2*c**2) - b**2*atanh(c/x)/c**2, Ne(c, 0)), (-a**2/(2*x**2), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 165 vs. 2(81) = 162.

Time = 0.04 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.90

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^3} dx$$

$$= \frac{1}{2} \left(c \left(\frac{\log(c+x)}{c^3} - \frac{\log(-c+x)}{c^3} - \frac{2}{c^2 x} \right) - \frac{2 \operatorname{artanh}(\frac{c}{x})}{x^2} \right) ab$$

$$- \frac{1}{8} \left(c^2 \left(\frac{\log(c+x)^2 - 2(\log(c+x) - 2)\log(-c+x) + \log(-c+x)^2 + 4\log(c+x)}{c^4} - \frac{8\log(x)}{c^4} \right) - \frac{b^2 \operatorname{artanh}(\frac{c}{x})^2}{2x^2} - \frac{a^2}{2x^2} \right)$$

input `integrate((a+b*arctanh(c/x))^2/x^3,x, algorithm="maxima")`

output

```
1/2*(c*(log(c + x)/c^3 - log(-c + x)/c^3 - 2/(c^2*x)) - 2*arctanh(c/x)/x^2
)*a*b - 1/8*(c^2*((log(c + x)^2 - 2*(log(c + x) - 2)*log(-c + x) + log(-c
+ x)^2 + 4*log(c + x))/c^4 - 8*log(x)/c^4) - 4*c*(log(c + x)/c^3 - log(-c
+ x)/c^3 - 2/(c^2*x))*arctanh(c/x))*b^2 - 1/2*b^2*arctanh(c/x)^2/x^2 - 1/2
*a^2/x^2
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 255 vs. $2(81) = 162$.

Time = 0.13 (sec) , antiderivative size = 255, normalized size of antiderivative = 2.93

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^3} dx =$$

$$\frac{b^2(c+x) \log\left(-\frac{c+x}{c-x}\right)^2}{\left(\frac{(c+x)^2 c}{(c-x)^2} - \frac{2(c+x)c}{c-x} + c\right)(c-x)} - \frac{2b^2 \log\left(-\frac{c+x}{c-x} + 1\right)}{c} + \frac{2b^2 \log\left(-\frac{c+x}{c-x}\right)}{c} - \frac{2\left(b^2 - \frac{2ab(c+x)}{c-x} - \frac{b^2(c+x)}{c-x}\right) \log\left(-\frac{c+x}{c-x}\right)}{\frac{(c+x)^2 c}{(c-x)^2} - \frac{2(c+x)c}{c-x} + c} - \frac{4\left(ab - \frac{a^2(c+x)}{c-x}\right)}{\frac{(c+x)^2 c}{(c-x)^2} - \frac{2(c+x)c}{c-x} + c}$$

$2c$

input

```
integrate((a+b*arctanh(c/x))^2/x^3,x, algorithm="giac")
```

output

```
-1/2*(b^2*(c + x)*log(-(c + x)/(c - x))^2/(((c + x)^2*c/(c - x)^2 - 2*(c +
x)*c/(c - x) + c)*(c - x)) - 2*b^2*log(-(c + x)/(c - x) + 1)/c + 2*b^2*lo
g(-(c + x)/(c - x))/c - 2*(b^2 - 2*a*b*(c + x)/(c - x) - b^2*(c + x)/(c -
x))*log(-(c + x)/(c - x))/((c + x)^2*c/(c - x)^2 - 2*(c + x)*c/(c - x) + c
) - 4*(a*b - a^2*(c + x)/(c - x) - a*b*(c + x)/(c - x))/((c + x)^2*c/(c -
x)^2 - 2*(c + x)*c/(c - x) + c))/c
```

Mupad [B] (verification not implemented)

Time = 4.30 (sec) , antiderivative size = 235, normalized size of antiderivative = 2.70

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^3} dx = \ln\left(1 - \frac{c}{x}\right) \left(\frac{ab}{2x^2} - \ln\left(\frac{c}{x} + 1\right) \left(\frac{b^2}{4c^2} - \frac{b^2}{4x^2}\right) + \frac{b^2(2cx - c^2)}{8c^2x^2} + \frac{b^2(2c^2 + 4xc)}{16c^2x^2}\right) - \frac{\frac{a^2}{2} + \frac{abx}{c}}{x^2} + \ln\left(\frac{c}{x} + 1\right)^2 \left(\frac{b^2}{8c^2} - \frac{b^2}{8x^2}\right) + \ln\left(1 - \frac{c}{x}\right)^2 \left(\frac{b^2}{8c^2} - \frac{b^2}{8x^2}\right) - \frac{\ln(x - c)(b^2 + ab)}{2c^2} + \frac{\ln(c + x)(ab - b^2)}{2c^2} - \frac{\ln(\frac{c}{x} + 1) \left(\frac{ab}{2} + \frac{b^2x}{2c}\right)}{x^2} + \frac{b^2 \ln(x)}{c^2}$$

input `int((a + b*atanh(c/x))^2/x^3,x)`output `log(1 - c/x)*((a*b)/(2*x^2) - log(c/x + 1)*(b^2/(4*c^2) - b^2/(4*x^2)) + (b^2*(2*c*x - c^2))/(8*c^2*x^2) + (b^2*(4*c*x + 2*c^2))/(16*c^2*x^2)) - (a^2/2 + (a*b*x)/c)/x^2 + log(c/x + 1)^2*(b^2/(8*c^2) - b^2/(8*x^2)) + log(1 - c/x)^2*(b^2/(8*c^2) - b^2/(8*x^2)) - (log(x - c)*(a*b + b^2))/(2*c^2) + (log(c + x)*(a*b - b^2))/(2*c^2) - (log(c/x + 1)*((a*b)/2 + (b^2*x)/(2*c)))/x^2 + (b^2*log(x))/c^2`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.53

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{x^3} dx = \frac{-\operatorname{atanh}(\frac{c}{x})^2 b^2 c^2 + \operatorname{atanh}(\frac{c}{x})^2 b^2 x^2 - 2 \operatorname{atanh}(\frac{c}{x}) ab c^2 + 2 \operatorname{atanh}(\frac{c}{x}) ab x^2 - 2 \operatorname{atanh}(\frac{c}{x}) b^2 cx + 2 \operatorname{atanh}(\frac{c}{x}) b^2 \ln(x)}{2c^2 x^2}$$

input `int((a+b*atanh(c/x))^2/x^3,x)`

output

```
( - atanh(c/x)**2*b**2*c**2 + atanh(c/x)**2*b**2*x**2 - 2*atanh(c/x)*a*b*c
**2 + 2*atanh(c/x)*a*b*x**2 - 2*atanh(c/x)*b**2*c*x + 2*atanh(c/x)*b**2*x*
*2 - 2*log( - c - x)*b**2*x**2 + 2*log(x)*b**2*x**2 - a**2*c**2 - 2*a*b*c*
x)/(2*c**2*x**2)
```

3.150 $\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx$

Optimal result	1224
Mathematica [A] (verified)	1225
Rubi [A] (verified)	1226
Maple [C] (warning: unable to verify)	1230
Fricas [F]	1231
Sympy [F]	1231
Maxima [F]	1231
Giac [F]	1232
Mupad [F(-1)]	1232
Reduce [F]	1233

Optimal result

Integrand size = 16, antiderivative size = 203

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \frac{1}{4} b^3 c^3 x - \frac{1}{4} b^3 c^4 \coth^{-1} \left(\frac{x}{c} \right) + \frac{1}{4} b^2 c^2 x^2 \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right) - b c^4 \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2 + \frac{3}{4} b c^3 x \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2 + \frac{1}{4} b c x^3 \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2 - \frac{1}{4} c^4 \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^3 + \frac{1}{4} x^4 \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^3 - 2 b^2 c^4 \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right) \log \left(2 - \frac{2}{1 + \frac{c}{x}} \right) + b^3 c^4 \operatorname{PolyLog} \left(2, -1 + \frac{2}{1 + \frac{c}{x}} \right)$$

output

```
1/4*b^3*c^3*x-1/4*b^3*c^4*arccoth(x/c)+1/4*b^2*c^2*x^2*(a+b*arccoth(x/c))-
b*c^4*(a+b*arccoth(x/c))^2+3/4*b*c^3*x*(a+b*arccoth(x/c))^2+1/4*b*c*x^3*(a
+b*arccoth(x/c))^2-1/4*c^4*(a+b*arccoth(x/c))^3+1/4*x^4*(a+b*arccoth(x/c))
^3-2*b^2*c^4*(a+b*arccoth(x/c))*ln(2-2/(1+c/x))+b^3*c^4*polylog(2,-1+2/(1+
c/x))
```

Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.41

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \frac{1}{8} \left(-2ab^2c^4 + 6a^2bc^3x + 2b^3c^3x + 2ab^2c^2x^2 + 2a^2bcx^3 \right. \\ \left. + 2a^3x^4 + 2b^2(bc(-4c^3 + 3c^2x + x^3) \right. \\ \left. + 3a(-c^4 + x^4)) \operatorname{arctanh} \left(\frac{c}{x} \right)^2 \right. \\ \left. + 2b^3(-c^4 + x^4) \operatorname{arctanh} \left(\frac{c}{x} \right)^3 \right. \\ \left. + 2b \operatorname{arctanh} \left(\frac{c}{x} \right) (3a^2x^4 + 2abcx(3c^2 + x^2) \right. \\ \left. + b^2(-c^4 + c^2x^2) - 8b^2c^4 \log \left(1 - e^{-2 \operatorname{arctanh} \left(\frac{c}{x} \right)} \right) \right) \\ \left. + 3a^2bc^4 \log \left(1 - \frac{c}{x} \right) - 16ab^2c^4 \log \left(\frac{c}{\sqrt{1 - \frac{c^2}{x^2}x}} \right) \right. \\ \left. - 3a^2bc^4 \log \left(\frac{c+x}{x} \right) \right. \\ \left. + 8b^3c^4 \operatorname{PolyLog} \left(2, e^{-2 \operatorname{arctanh} \left(\frac{c}{x} \right)} \right) \right)$$

input `Integrate[x^3*(a + b*ArcTanh[c/x])^3,x]`output `(-2*a*b^2*c^4 + 6*a^2*b*c^3*x + 2*b^3*c^3*x + 2*a*b^2*c^2*x^2 + 2*a^2*b*c*x^3 + 2*a^3*x^4 + 2*b^2*(b*c*(-4*c^3 + 3*c^2*x + x^3) + 3*a*(-c^4 + x^4))*ArcTanh[c/x]^2 + 2*b^3*(-c^4 + x^4)*ArcTanh[c/x]^3 + 2*b*ArcTanh[c/x]*(3*a^2*x^4 + 2*a*b*c*x*(3*c^2 + x^2) + b^2*(-c^4 + c^2*x^2) - 8*b^2*c^4*Log[1 - E^(-2*ArcTanh[c/x])]) + 3*a^2*b*c^4*Log[1 - c/x] - 16*a*b^2*c^4*Log[c/(Sqrt[1 - c^2/x^2]*x)] - 3*a^2*b*c^4*Log[(c + x)/x] + 8*b^3*c^4*PolyLog[2, E^(-2*ArcTanh[c/x])])/8`

Rubi [A] (verified)

Time = 2.55 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.31, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6454, 6452, 6544, 6452, 6544, 6452, 264, 219, 6510, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^3 dx \\
 & \quad \downarrow \text{6454} \\
 & - \int x^5 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^3 d\frac{1}{x} \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{4}x^4 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^3 - \frac{3}{4}bc \int \frac{x^4 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} \\
 & \quad \downarrow \text{6544} \\
 & \frac{1}{4}x^4 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^3 - \\
 & \frac{3}{4}bc \left(c^2 \int \frac{x^2 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \int x^4 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2 d\frac{1}{x} \right) \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{4}x^4 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^3 - \\
 & \frac{3}{4}bc \left(c^2 \int \frac{x^2 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{2}{3}bc \int \frac{x^3 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - \frac{1}{3}x^3 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2 \right) \\
 & \quad \downarrow \text{6544} \\
 & \frac{1}{4}x^4 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^3 - \\
 & \frac{3}{4}bc \left(c^2 \left(c^2 \int \frac{\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \int x^2 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2 d\frac{1}{x} \right) + \frac{2}{3}bc \left(c^2 \int \frac{x \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \right. \right. \\
 & \quad \left. \left. \int x^3 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right) d\frac{1}{x} - \frac{1}{3}x^3 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2 \right) \right) \\
 & \quad \downarrow \text{6452}
 \end{aligned}$$

$$\frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^3 - \frac{3}{4}bc\left(\frac{2}{3}bc\left(c^2 \int \frac{x(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{1}{2}bc \int \frac{x^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - \frac{1}{2}x^2(a + \operatorname{barctanh}(\frac{c}{x}))\right) + c^2\left(c^2 \int \frac{(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x}\right)\right)$$

↓ 264

$$\frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^3 - \frac{3}{4}bc\left(\frac{2}{3}bc\left(c^2 \int \frac{x(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{1}{2}bc\left(c^2 \int \frac{1}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - x\right) - \frac{1}{2}x^2(a + \operatorname{barctanh}(\frac{c}{x}))\right) + c^2\left(c^2 \int \frac{(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x}\right)\right)$$

↓ 219

$$\frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^3 - \frac{3}{4}bc\left(\frac{2}{3}bc\left(c^2 \int \frac{x(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - \frac{1}{2}x^2(a + \operatorname{barctanh}(\frac{c}{x})) + \frac{1}{2}bc(\operatorname{carctanh}(\frac{c}{x}) - x)\right) + c^2\left(c^2 \int \frac{(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x}\right)\right)$$

↓ 6510

$$\frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^3 - \frac{3}{4}bc\left(c^2\left(2bc \int \frac{x(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{c(a + \operatorname{barctanh}(\frac{c}{x}))^3}{3b} - x(a + \operatorname{barctanh}(\frac{c}{x}))^2\right) + \frac{2}{3}bc\left(c^2 \int \frac{x(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x}\right)\right)$$

↓ 6550

$$\frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^3 - \frac{3}{4}bc\left(\frac{2}{3}bc\left(c^2\left(\int \frac{x(a + \operatorname{barctanh}(\frac{c}{x}))}{\frac{c}{x} + 1} d\frac{1}{x} + \frac{(a + \operatorname{barctanh}(\frac{c}{x}))^2}{2b}\right) - \frac{1}{2}x^2(a + \operatorname{barctanh}(\frac{c}{x})) + \frac{1}{2}bc(\operatorname{carctanh}(\frac{c}{x}) - x)\right) + c^2\left(c^2 \int \frac{(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x}\right)\right)$$

↓ 6494

$$\frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^3 - \frac{3}{4}bc\left(c^2\left(2bc\left(-bc \int \frac{\log\left(2 - \frac{2}{\frac{c}{x} + 1}\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{(a + \operatorname{barctanh}(\frac{c}{x}))^2}{2b} + \log\left(2 - \frac{2}{\frac{c}{x} + 1}\right)(a + \operatorname{barctanh}(\frac{c}{x}))\right) + c^2\left(c^2 \int \frac{(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x}\right)\right)\right)$$

↓ 2897

$$\frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right)\right)^3 - \frac{3}{4}bc\left(\frac{2}{3}bc\left(c^2\left(\frac{(a + \operatorname{barctanh}(\frac{c}{x}))^2}{2b} + \log\left(2 - \frac{2}{\frac{c}{x} + 1}\right)(a + \operatorname{barctanh}(\frac{c}{x})) - \frac{1}{2}b \operatorname{PolyLog}\left(2, \frac{2}{\frac{c}{x} + 1} - 1\right)\right) + c^2\left(c^2 \int \frac{(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x}\right)\right)\right)$$

input `Int[x^3*(a + b*ArcTanh[c/x])^3,x]`

output `(x^4*(a + b*ArcTanh[c/x])^3)/4 - (3*b*c*(-1/3*(x^3*(a + b*ArcTanh[c/x])^2) + c^2*(-(x*(a + b*ArcTanh[c/x])^2) + (c*(a + b*ArcTanh[c/x])^3)/(3*b) + 2*b*c*((a + b*ArcTanh[c/x])^2/(2*b) + (a + b*ArcTanh[c/x])*Log[2 - 2/(1 + c/x)] - (b*PolyLog[2, -1 + 2/(1 + c/x)]/2)) + (2*b*c*(-1/2*(x^2*(a + b*ArcTanh[c/x])) + (b*c*(-x + c*ArcTanh[c/x]))/2 + c^2*((a + b*ArcTanh[c/x])^2/(2*b) + (a + b*ArcTanh[c/x])*Log[2 - 2/(1 + c/x)] - (b*PolyLog[2, -1 + 2/(1 + c/x)]/2)))/3)/4`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 264 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2897 `Int[Log[u_]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x
, x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simpl
ify[(m + 1)/n]`

rule 6494 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_))), x
_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] -
Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]
/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c
^2*d^2 - e^2, 0]`

rule 6510 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symb
ol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b
, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6544 `Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_))/((d_) + (e
.)*(x)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x
, x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x
^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]`

rule 6550 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_)^2)),
x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/
d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 20.36 (sec) , antiderivative size = 1267, normalized size of antiderivative = 6.24

method	result	size
derivativedivides	Expression too large to display	1267
default	Expression too large to display	1267
parts	Expression too large to display	1322
risch	Expression too large to display	44754

input `int(x^3*(a+b*arctanh(c/x))^3,x,method=_RETURNVERBOSE)`

output

```
-c^4*(-3/16*I*b^3*Pi*csgn(I*(1+c/x)^2/(c^2/x^2-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^3*arctanh(c/x)^2+3/8*I*b^3*Pi*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))^3*arctanh(c/x)^2-3/16*I*b^3*Pi*csgn(I*(1+c/x)/(1-c^2/x^2)^(1/2))^2*csgn(I*(1+c/x)^2/(c^2/x^2-1))*arctanh(c/x)^2-3/8*I*b^3*Pi*csgn(I*(1+c/x)/(1-c^2/x^2)^(1/2))*csgn(I*(1+c/x)^2/(c^2/x^2-1))^2*arctanh(c/x)^2-3/16*I*b^3*Pi*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))^2*arctanh(c/x)^2+3/16*I*b^3*Pi*csgn(I*(1+c/x)^2/(c^2/x^2-1))*csgn(I*(1+c/x)^2/(c^2/x^2-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^2*arctanh(c/x)^2+3/16*I*b^3*Pi*csgn(I*(1+c/x)^2/(c^2/x^2-1))*csgn(I*(1+c/x)^2/(c^2/x^2-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^2*arctanh(c/x)^2+3/8*I*b^3*Pi*arctanh(c/x)^2-1/4*b^3/c^4*x^4*arctanh(c/x)^3-1/4*b^3*arctanh(c/x)^2/c^3*x^3-3/4*b^3*arctanh(c/x)^2/c*x-1/4*b^3*arctanh(c/x)/c^2*x^2-3/16*I*b^3*Pi*csgn(I*(1+c/x)^2/(c^2/x^2-1))^3*arctanh(c/x)^2-3/8*I*b^3*Pi*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))^2*arctanh(c/x)^2-1/4*a^3/c^4*x^4+1/4*b^3*arctanh(c/x)^3-b^3*arctanh(c/x)^2+1/4*b^3*arctanh(c/x)+3*a*b^2*(-1/4/c^4*x^4*arctanh(c/x)^2-1/6/c^3*x^3*arctanh(c/x)-1/2/c*x*arctanh(c/x)+1/4*arctanh(c/x)*ln(1+c/x)-1/4*arctanh(c/x)*ln(c/x-1)-1/16*ln(c/x-1)^2+1/8*ln(c/x-1)*ln(1/2*c/x+1/2)+1/8*(ln(1+c/x)-ln(1/2*c/x+1/2))*ln(-1/2*c/x+1/2)-1/16*ln(1+c/x)^2-1/12/c^2*x^2+2/3*ln(c/x)-1/3*ln(1+c/x)-1/3*ln(c/x-1))+3*a^2*b*(-1/4/c^4*x^4*arctanh(c/x)+1/8*ln(1+c/x)-1/8*ln(c/x-1)-1/12/c^3*x^3-1/4*x/c)+1/4*...
```

Fricas [F]

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x} \right) + a \right)^3 x^3 dx$$

input `integrate(x^3*(a+b*arctanh(c/x))^3,x, algorithm="fricas")`

output `integral(b^3*x^3*arctanh(c/x)^3 + 3*a*b^2*x^3*arctanh(c/x)^2 + 3*a^2*b*x^3*arctanh(c/x) + a^3*x^3, x)`

Sympy [F]

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int x^3 \left(a + b \operatorname{atanh} \left(\frac{c}{x} \right) \right)^3 dx$$

input `integrate(x**3*(a+b*atanh(c/x))**3,x)`

output `Integral(x**3*(a + b*atanh(c/x))**3, x)`

Maxima [F]

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x} \right) + a \right)^3 x^3 dx$$

input `integrate(x^3*(a+b*arctanh(c/x))^3,x, algorithm="maxima")`

output

```

3/4*a*b^2*x^4*arctanh(c/x)^2 + 1/4*a^3*x^4 + 1/8*(6*x^4*arctanh(c/x) - (3*
c^3*log(c + x) - 3*c^3*log(-c + x) - 6*c^2*x - 2*x^3)*c)*a^2*b + 1/16*((3*
c^2*log(c + x)^2 + 3*c^2*log(-c + x)^2 + 16*c^2*log(c + x) + 4*x^2 - 2*(3*
c^2*log(c + x) - 8*c^2)*log(-c + x))*c^2 - 4*(3*c^3*log(c + x) - 3*c^3*log
(-c + x) - 6*c^2*x - 2*x^3)*c*arctanh(c/x))*a*b^2 + 1/32*(16*c^5*integrate
(-log(c + x)/(c^2 - x^2), x) + 40*c^4*integrate(-x*log(c + x)/(c^2 - x^2),
x) - 2*(c*log(c + x) - c*log(-c + x) - 2*x)*c^3 - (c^4 - x^4)*log(c + x)^
3 + (c^4 - x^4)*log(-c + x)^3 + 2*(c^2*log(-c^2 + x^2) + x^2)*c^2 + 8*c^2*
integrate(-x^3*log(c + x)/(c^2 - x^2), x) + 2*(3*c^3*x + c*x^3)*log(c + x)
^2 - (8*c^4 - 6*c^3*x - 2*c*x^3 + 3*(c^4 - x^4)*log(c + x))*log(-c + x)^2
- (4*c^2*x^2 - 3*(c^4 - x^4)*log(c + x)^2 + 4*(4*c^4 + 3*c^3*x + c*x^3)*lo
g(c + x))*log(-c + x))*b^3

```

Giac [F]

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x} \right) + a \right)^3 x^3 dx$$

input

```
integrate(x^3*(a+b*arctanh(c/x))^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c/x) + a)^3*x^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int x^3 \left(a + b \operatorname{atanh} \left(\frac{c}{x} \right) \right)^3 dx$$

input

```
int(x^3*(a + b*atanh(c/x))^3,x)
```

output

```
int(x^3*(a + b*atanh(c/x))^3, x)
```

Reduce [F]

$$\begin{aligned}
\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = & -\frac{\operatorname{atanh} \left(\frac{c}{x} \right)^3 b^3 c^4}{4} + \frac{\operatorname{atanh} \left(\frac{c}{x} \right)^3 b^3 x^4}{4} - \frac{3 \operatorname{atanh} \left(\frac{c}{x} \right)^2 a b^2 c^4}{4} \\
& + \frac{3 \operatorname{atanh} \left(\frac{c}{x} \right)^2 a b^2 x^4}{4} + \frac{3 \operatorname{atanh} \left(\frac{c}{x} \right)^2 b^3 c^3 x}{4} \\
& + \frac{\operatorname{atanh} \left(\frac{c}{x} \right)^2 b^3 c x^3}{4} - \frac{3 \operatorname{atanh} \left(\frac{c}{x} \right) a^2 b c^4}{4} \\
& + \frac{3 \operatorname{atanh} \left(\frac{c}{x} \right) a^2 b x^4}{4} - 2 \operatorname{atanh} \left(\frac{c}{x} \right) a b^2 c^4 \\
& + \frac{3 \operatorname{atanh} \left(\frac{c}{x} \right) a b^2 c^3 x}{2} + \frac{\operatorname{atanh} \left(\frac{c}{x} \right) a b^2 c x^3}{2} \\
& - \frac{\operatorname{atanh} \left(\frac{c}{x} \right) b^3 c^4}{4} + \frac{\operatorname{atanh} \left(\frac{c}{x} \right) b^3 c^2 x^2}{4} \\
& - 2 \left(\int \frac{\operatorname{atanh} \left(\frac{c}{x} \right) x}{c^2 - x^2} dx \right) b^3 c^4 + 2 \log(-c - x) a b^2 c^4 \\
& + \frac{a^3 x^4}{4} + \frac{3 a^2 b c^3 x}{4} + \frac{a^2 b c x^3}{4} + \frac{a b^2 c^2 x^2}{4} + \frac{b^3 c^3 x}{4}
\end{aligned}$$

input `int(x^3*(a+b*atanh(c/x))^3,x)`

output `(- atanh(c/x)**3*b**3*c**4 + atanh(c/x)**3*b**3*x**4 - 3*atanh(c/x)**2*a*b**2*c**4 + 3*atanh(c/x)**2*a*b**2*x**4 + 3*atanh(c/x)**2*b**3*c**3*x + atanh(c/x)**2*b**3*c*x**3 - 3*atanh(c/x)*a**2*b*c**4 + 3*atanh(c/x)*a**2*b*x**4 - 8*atanh(c/x)*a*b**2*c**4 + 6*atanh(c/x)*a*b**2*c**3*x + 2*atanh(c/x)*a*b**2*c*x**3 - atanh(c/x)*b**3*c**4 + atanh(c/x)*b**3*c**2*x**2 - 8*int((atanh(c/x)*x)/(c**2 - x**2),x)*b**3*c**4 + 8*log(-c - x)*a*b**2*c**4 + a**3*x**4 + 3*a**2*b*c**3*x + a**2*b*c*x**3 + a*b**2*c**2*x**2 + b**3*c**3*x)/4`

3.151 $\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx$

Optimal result	1234
Mathematica [C] (verified)	1235
Rubi [A] (verified)	1236
Maple [C] (warning: unable to verify)	1240
Fricas [F]	1241
Sympy [F]	1242
Maxima [F]	1242
Giac [F]	1242
Mupad [F(-1)]	1243
Reduce [F]	1243

Optimal result

Integrand size = 16, antiderivative size = 217

$$\begin{aligned}
 \int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx &= b^2 c^2 x \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right) - \frac{1}{2} b c^3 \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right)^2 \\
 &\quad + \frac{1}{2} b c x^2 \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right)^2 \\
 &\quad - \frac{1}{3} c^3 \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right)^3 + \frac{1}{3} x^3 \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right)^3 \\
 &\quad - b c^3 \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right)^2 \log \left(2 - \frac{2}{1 + \frac{c}{x}} \right) \\
 &\quad + \frac{1}{2} b^3 c^3 \log \left(1 - \frac{c^2}{x^2} \right) + b^3 c^3 \log(x) \\
 &\quad + b^2 c^3 \left(a + b \operatorname{coth}^{-1} \left(\frac{x}{c} \right) \right) \operatorname{PolyLog} \left(2, -1 + \frac{2}{1 + \frac{c}{x}} \right) \\
 &\quad + \frac{1}{2} b^3 c^3 \operatorname{PolyLog} \left(3, -1 + \frac{2}{1 + \frac{c}{x}} \right)
 \end{aligned}$$

output

```

b^2*c^2*x*(a+b*arccoth(x/c))-1/2*b*c^3*(a+b*arccoth(x/c))^2+1/2*b*c*x^2*(a
+b*arccoth(x/c))^2-1/3*c^3*(a+b*arccoth(x/c))^3+1/3*x^3*(a+b*arccoth(x/c))
^3-b*c^3*(a+b*arccoth(x/c))^2*ln(2-2/(1+c/x))+1/2*b^3*c^3*ln(1-c^2/x^2)+b^
3*c^3*ln(x)+b^2*c^3*(a+b*arccoth(x/c))*polylog(2,-1+2/(1+c/x))+1/2*b^3*c^3
*polylog(3,-1+2/(1+c/x))

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.56 (sec) , antiderivative size = 316, normalized size of antiderivative = 1.46

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \frac{1}{6} \left(3a^2bcx^2 + 2a^3x^3 + 6a^2bx^3 \operatorname{arctanh} \left(\frac{c}{x} \right) \right. \\ \left. + 3a^2bc^3 \log(-c^2 + x^2) \right. \\ \left. + 6ab^2 \left(c^2x + (-c^3 + x^3) \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 \right. \\ \left. + c \operatorname{arctanh} \left(\frac{c}{x} \right) \left(-c^2 + x^2 - 2c^2 \log \left(1 - e^{-2\operatorname{arctanh} \left(\frac{c}{x} \right)} \right) \right) \right. \\ \left. + c^3 \operatorname{PolyLog} \left(2, e^{-2\operatorname{arctanh} \left(\frac{c}{x} \right)} \right) \right. \\ \left. + \frac{1}{4}b^3 \left(-ic^3\pi^3 + 24c^2x \operatorname{arctanh} \left(\frac{c}{x} \right) \right. \right. \\ \left. \left. - 12c^3 \operatorname{arctanh} \left(\frac{c}{x} \right)^2 + 12cx^2 \operatorname{arctanh} \left(\frac{c}{x} \right)^2 \right. \right. \\ \left. \left. + 8c^3 \operatorname{arctanh} \left(\frac{c}{x} \right)^3 + 8x^3 \operatorname{arctanh} \left(\frac{c}{x} \right)^3 \right. \right. \\ \left. \left. - 24c^3 \operatorname{arctanh} \left(\frac{c}{x} \right)^2 \log \left(1 - e^{2\operatorname{arctanh} \left(\frac{c}{x} \right)} \right) \right. \right. \\ \left. \left. - 24c^3 \log \left(\frac{c}{\sqrt{1 - \frac{c^2}{x^2}}x} \right) \right. \right. \\ \left. \left. - 24c^3 \operatorname{arctanh} \left(\frac{c}{x} \right) \operatorname{PolyLog} \left(2, e^{2\operatorname{arctanh} \left(\frac{c}{x} \right)} \right) \right. \right. \\ \left. \left. + 12c^3 \operatorname{PolyLog} \left(3, e^{2\operatorname{arctanh} \left(\frac{c}{x} \right)} \right) \right) \right)$$

input

```
Integrate[x^2*(a + b*ArcTanh[c/x])^3,x]
```


output

```
(3*a^2*b*c*x^2 + 2*a^3*x^3 + 6*a^2*b*x^3*ArcTanh[c/x] + 3*a^2*b*c^3*Log[-c^2 + x^2] + 6*a*b^2*(c^2*x + (-c^3 + x^3)*ArcTanh[c/x]^2 + c*ArcTanh[c/x]*(-c^2 + x^2 - 2*c^2*Log[1 - E^(-2*ArcTanh[c/x])])) + c^3*PolyLog[2, E^(-2*ArcTanh[c/x])]) + (b^3*((-I)*c^3*Pi^3 + 24*c^2*x*ArcTanh[c/x] - 12*c^3*ArcTanh[c/x]^2 + 12*c*x^2*ArcTanh[c/x]^2 + 8*c^3*ArcTanh[c/x]^3 + 8*x^3*ArcTanh[c/x]^3 - 24*c^3*ArcTanh[c/x]^2*Log[1 - E^(2*ArcTanh[c/x])]) - 24*c^3*Log[c/(Sqrt[1 - c^2/x^2]*x)] - 24*c^3*ArcTanh[c/x]*PolyLog[2, E^(2*ArcTanh[c/x])] + 12*c^3*PolyLog[3, E^(2*ArcTanh[c/x])]))/4/6
```

Rubi [A] (verified)

Time = 2.03 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.938$, Rules used = {6454, 6452, 6544, 6452, 6544, 6452, 243, 47, 14, 16, 6510, 6550, 6494, 6618, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx \\
 & \quad \downarrow \text{6454} \\
 & - \int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 d \frac{1}{x} \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{3} x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 - bc \int \frac{x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2}{1 - \frac{c^2}{x^2}} d \frac{1}{x} \\
 & \quad \downarrow \text{6544} \\
 & \frac{1}{3} x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 - \\
 & bc \left(c^2 \int \frac{x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2}{1 - \frac{c^2}{x^2}} d \frac{1}{x} + \int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^2 d \frac{1}{x} \right) \\
 & \quad \downarrow \text{6452}
 \end{aligned}$$

$$bc \left(c^2 \int \frac{x(a + \operatorname{barctanh}(\frac{c}{x}))^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + bc \int \frac{x^2(a + \operatorname{barctanh}(\frac{c}{x}))}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - \frac{1}{2}x^2(a + \operatorname{barctanh}(\frac{c}{x}))^2 \right)$$

↓ 6544

$$bc \left(c^2 \int \frac{x(a + \operatorname{barctanh}(\frac{c}{x}))^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + bc \left(c^2 \int \frac{a + \operatorname{barctanh}(\frac{c}{x})}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \int x^2(a + \operatorname{barctanh}(\frac{c}{x})) d\frac{1}{x} \right) - \frac{1}{2}x^2(a + \operatorname{barctanh}(\frac{c}{x}))^2 \right)$$

↓ 6452

$$bc \left(c^2 \int \frac{x(a + \operatorname{barctanh}(\frac{c}{x}))^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + bc \left(c^2 \int \frac{a + \operatorname{barctanh}(\frac{c}{x})}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + bc \int \frac{x}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - x(a + \operatorname{barctanh}(\frac{c}{x})) \right) - \frac{1}{2}x^2(a + \operatorname{barctanh}(\frac{c}{x}))^2 \right)$$

↓ 243

$$bc \left(c^2 \int \frac{x(a + \operatorname{barctanh}(\frac{c}{x}))^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + bc \left(c^2 \int \frac{a + \operatorname{barctanh}(\frac{c}{x})}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{1}{2}bc \int \frac{x}{1 - \frac{c^2}{x^2}} d\frac{1}{x^2} - x(a + \operatorname{barctanh}(\frac{c}{x})) \right) - \frac{1}{2}x^2(a + \operatorname{barctanh}(\frac{c}{x}))^2 \right)$$

↓ 47

$$bc \left(c^2 \int \frac{x(a + \operatorname{barctanh}(\frac{c}{x}))^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + bc \left(c^2 \int \frac{a + \operatorname{barctanh}(\frac{c}{x})}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{1}{2}bc \left(c^2 \int \frac{1}{1 - \frac{c^2}{x^2}} d\frac{1}{x^2} + \int x d\frac{1}{x^2} \right) - x(a + \operatorname{barctanh}(\frac{c}{x})) \right) - \frac{1}{2}x^2(a + \operatorname{barctanh}(\frac{c}{x}))^2 \right)$$

↓ 14

$$bc \left(c^2 \int \frac{x(a + \operatorname{barctanh}(\frac{c}{x}))^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + bc \left(c^2 \int \frac{a + \operatorname{barctanh}(\frac{c}{x})}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{1}{2}bc \left(c^2 \int \frac{1}{1 - \frac{c^2}{x^2}} d\frac{1}{x^2} + \log\left(\frac{1}{x^2}\right) \right) - x(a + \operatorname{barctanh}(\frac{c}{x})) \right) - \frac{1}{2}x^2(a + \operatorname{barctanh}(\frac{c}{x}))^2 \right)$$

↓ 16

$$bc \left(c^2 \int \frac{x(a + \operatorname{barctanh}(\frac{c}{x}))^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + bc \left(c^2 \int \frac{a + \operatorname{barctanh}(\frac{c}{x})}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - x(a + \operatorname{barctanh}(\frac{c}{x})) + \frac{1}{2}bc \left(\log\left(\frac{1}{x^2}\right) - \log\left(\frac{1}{x^2}\right) \right) \right) - \frac{1}{2}x^2(a + \operatorname{barctanh}(\frac{c}{x}))^2 \right)$$

↓ 6510

$$bc \left(c^2 \int \frac{x(a + \operatorname{arctanh}(\frac{c}{x}))^2}{1 - \frac{c^2}{x^2}} dx + bc \left(\frac{c(a + \operatorname{arctanh}(\frac{c}{x}))^2}{2b} - x(a + \operatorname{arctanh}(\frac{c}{x})) + \frac{1}{2}bc \left(\log\left(\frac{1}{x^2}\right) - \log\left(\frac{c}{x}\right) \right) \right) \right)$$

↓ 6550

$$bc \left(c^2 \left(\int \frac{x(a + \operatorname{arctanh}(\frac{c}{x}))^2}{\frac{c}{x} + 1} dx + \frac{(a + \operatorname{arctanh}(\frac{c}{x}))^3}{3b} \right) + bc \left(\frac{c(a + \operatorname{arctanh}(\frac{c}{x}))^2}{2b} - x(a + \operatorname{arctanh}(\frac{c}{x})) \right) \right)$$

↓ 6494

$$bc \left(c^2 \left(-2bc \int \frac{(a + \operatorname{arctanh}(\frac{c}{x})) \log\left(2 - \frac{2}{\frac{c}{x} + 1}\right)}{1 - \frac{c^2}{x^2}} dx + \frac{(a + \operatorname{arctanh}(\frac{c}{x}))^3}{3b} + \log\left(2 - \frac{2}{\frac{c}{x} + 1}\right) (a + \operatorname{arctanh}(\frac{c}{x})) \right) \right)$$

↓ 6618

$$bc \left(c^2 \left(-2bc \left(\frac{\operatorname{PolyLog}\left(2, \frac{2}{\frac{c}{x} + 1} - 1\right) (a + \operatorname{arctanh}(\frac{c}{x}))}{2c} - \frac{1}{2}b \int \frac{\operatorname{PolyLog}\left(2, \frac{2}{\frac{c}{x} + 1} - 1\right)}{1 - \frac{c^2}{x^2}} dx \right) + \frac{(a + \operatorname{arctanh}(\frac{c}{x}))^3}{3b} \right) \right)$$

↓ 7164

$$bc \left(c^2 \left(-2bc \left(\frac{\operatorname{PolyLog}\left(2, \frac{2}{\frac{c}{x} + 1} - 1\right) (a + \operatorname{arctanh}(\frac{c}{x}))}{2c} + \frac{b \operatorname{PolyLog}\left(3, \frac{2}{\frac{c}{x} + 1} - 1\right)}{4c} \right) + \frac{(a + \operatorname{arctanh}(\frac{c}{x}))^3}{3b} \right) \right)$$

input

```
Int [x^2*(a + b*ArcTanh[c/x])^3,x]
```

output

```
(x^3*(a + b*ArcTanh[c/x])^3)/3 - b*c*(-1/2*(x^2*(a + b*ArcTanh[c/x])^2) +
b*c*(-(x*(a + b*ArcTanh[c/x])) + (c*(a + b*ArcTanh[c/x])^2)/(2*b) + (b*c*(
-Log[1 - c^2/x^2] + Log[x^(-2)]))/2) + c^2*((a + b*ArcTanh[c/x])^3/(3*b) +
(a + b*ArcTanh[c/x])^2*Log[2 - 2/(1 + c/x)] - 2*b*c*((a + b*ArcTanh[c/x]
)*PolyLog[2, -1 + 2/(1 + c/x)]/(2*c) + (b*PolyLog[3, -1 + 2/(1 + c/x)]/(
4*c))))
```

Definitions of rubi rules used

- rule 14 $\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ ; FreeQ}[a, x]$
- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$
- rule 47 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x]$
- rule 243 $\text{Int}[(x_)^{(m_)}*((a_)+(b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] \text{ ; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$
- rule 6452 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{2n})}), x], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$
- rule 6454 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n] - 1)*(a + b*\text{ArcTanh}[c*x])^p}, x], x, x^n], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$
- rule 6494 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)]*(b_))^{(p_)}*((d_)+(e_)*(x_)), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^p*(\text{Log}[2 - 2/(1 + e*(x/d))]/d), x] - \text{Simp}[b*c*(p/d) \text{ Int}[(a + b*\text{ArcTanh}[c*x])^{(p-1)}*(\text{Log}[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] \text{ ; FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d^2 - e^2, 0]$

rule 6510 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / (d + e \cdot x^2), x_{\text{Symbol}}] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot c \cdot d \cdot (p+1)), x] /;$ $\text{FreeQ}[\{a, b, c, d, e, p\}, x]$ && $\text{EqQ}[c^2 \cdot d + e, 0]$ && $\text{NeQ}[p, -1]$

rule 6544 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p \cdot (f \cdot x)^m / (d + e \cdot x^2), x_{\text{Symbol}}] \rightarrow \text{Simp}[1/d \int (f \cdot x)^m (a + b \cdot \text{ArcTanh}[c \cdot x])^p dx, x] - \text{Simp}[e/(d \cdot f^2) \int (f \cdot x)^{m+2} (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2) dx, x] /;$ $\text{FreeQ}[\{a, b, c, d, e, f\}, x]$ && $\text{GtQ}[p, 0]$ && $\text{LtQ}[m, -1]$

rule 6550 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / (x \cdot (d + e \cdot x^2)), x_{\text{Symbol}}] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot d \cdot (p+1)), x] + \text{Simp}[1/d \int (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (x \cdot (1 + c \cdot x)) dx, x] /;$ $\text{FreeQ}[\{a, b, c, d, e\}, x]$ && $\text{EqQ}[c^2 \cdot d + e, 0]$ && $\text{GtQ}[p, 0]$

rule 6618 $\text{Int}[(\text{Log}[u] \cdot (a + \text{ArcTanh}[c \cdot x] \cdot b)^p) / (d + e \cdot x^2), x_{\text{Symbol}}] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{PolyLog}[2, 1 - u] / (2 \cdot c \cdot d)), x] - \text{Simp}[b \cdot (p/2) \int (a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{PolyLog}[2, 1 - u] / (d + e \cdot x^2)) dx, x] /;$ $\text{FreeQ}[\{a, b, c, d, e\}, x]$ && $\text{IGtQ}[p, 0]$ && $\text{EqQ}[c^2 \cdot d + e, 0]$ && $\text{EqQ}[(1 - u)^2 - (1 - 2/(1 + c \cdot x))^2, 0]$

rule 7164 $\text{Int}[u \cdot \text{PolyLog}[n, v], x_{\text{Symbol}}] \rightarrow \text{With}[\{w = \text{DerivativeDivides}[v, u \cdot v, x]\}, \text{Simp}[w \cdot \text{PolyLog}[n + 1, v], x] /;$ $\text{!FalseQ}[w]] /;$ $\text{FreeQ}[n, x]$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 36.31 (sec) , antiderivative size = 1780, normalized size of antiderivative = 8.20

method	result	size
derivativedivides	Expression too large to display	1780
default	Expression too large to display	1780
parts	Expression too large to display	1782

input `int(x^2*(a+b*arctanh(c/x))^3,x,method=_RETURNVERBOSE)`

output

```
-c^3*(-1/3*a^3/c^3*x^3+b^3*(1/4*I*Pi*csgn(I*(1+c/x)^2/(c^2/x^2-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^3*arctanh(c/x)^2+1/4*I*Pi*csgn(I*(1+c/x)^2/(c^2/x^2-1))^3*arctanh(c/x)^2-1/2*(-(1-c^2/x^2)^(1/2)+c/x+1)/c*x*arctanh(c/x)+1/2*I*Pi*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^3*arctanh(c/x)^2+1/2*I*Pi*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))^3*arctanh(c/x)^2-1/2*I*Pi*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))^2*arctanh(c/x)^2+ln(c/x)*arctanh(c/x)^2-arctanh(c/x)^2*ln((1+c/x)^2/(1-c^2/x^2)-1)+arctanh(c/x)^2*ln(1+(1+c/x)/(1-c^2/x^2)^(1/2))+2*arctanh(c/x)*polylog(2,-(1+c/x)/(1-c^2/x^2)^(1/2))+arctanh(c/x)^2*ln(1-(1+c/x)/(1-c^2/x^2)^(1/2))+2*arctanh(c/x)*polylog(2,(1+c/x)/(1-c^2/x^2)^(1/2))+1/2*I*Pi*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1))*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^2*arctanh(c/x)^2+ln((1+c/x)/(1-c^2/x^2)^(1/2)-1)+1/2*arctanh(c/x)^2-1/2/c^2*x^2*arctanh(c/x)^2+ln(1+(1+c/x)/(1-c^2/x^2)^(1/2))-2*polylog(3,-(1+c/x)/(1-c^2/x^2)^(1/2))-2*polylog(3,(1+c/x)/(1-c^2/x^2)^(1/2))+1/4*I*Pi*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))*csgn(I*(1+c/x)^2/(c^2/x^2-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^2*arctanh(c/x)^2-1/2*I*Pi*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1))*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^2*arctanh(c/x)^2+1/2*I*Pi*csgn(I*(1+c/x)/(1-c^2/x^2)^(1/2))*csgn(I*(1+c/x)^2/(c^2/x^2-1))^2*arctanh(c/x)^2-1/2*I*Pi*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^2*arctanh(c/x)^2...
```

Fricas [F]

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int \left(b \operatorname{arctanh} \left(\frac{c}{x} \right) + a \right)^3 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c/x))^3,x, algorithm="fricas")`

output `integral(b^3*x^2*arctanh(c/x)^3 + 3*a*b^2*x^2*arctanh(c/x)^2 + 3*a^2*b*x^2*arctanh(c/x) + a^3*x^2, x)`

Sympy [F]

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int x^2 \left(a + b \operatorname{atanh} \left(\frac{c}{x} \right) \right)^3 dx$$

input `integrate(x**2*(a+b*atanh(c/x))**3,x)`

output `Integral(x**2*(a + b*atanh(c/x))**3, x)`

Maxima [F]

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x} \right) + a \right)^3 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c/x))^3,x, algorithm="maxima")`

output `1/3*a^3*x^3 + 1/2*(2*x^3*arctanh(c/x) + (c^2*log(-c^2 + x^2) + x^2)*c)*a^2 *b + 1/24*(b^3*c^3 - b^3*x^3)*log(-c + x)^3 + 1/8*(b^3*c*x^2 + 2*a*b^2*x^3 + (b^3*c^3 + b^3*x^3)*log(c + x))*log(-c + x)^2 - integrate(-1/8*((b^3*c*x^2 - b^3*x^3)*log(c + x)^3 + 6*(a*b^2*c*x^2 - a*b^2*x^3)*log(c + x)^2 + (2*b^3*c*x^2 + 4*a*b^2*x^3 - 3*(b^3*c*x^2 - b^3*x^3)*log(c + x)^2 + 2*(b^3*c^3 - 6*a*b^2*c*x^2 + (6*a*b^2 + b^3)*x^3)*log(c + x))*log(-c + x))/(c - x), x)`

Giac [F]

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x} \right) + a \right)^3 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c/x))^3,x, algorithm="giac")`

output `integrate((b*arctanh(c/x) + a)^3*x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int x^2 \left(a + b \operatorname{atanh} \left(\frac{c}{x} \right) \right)^3 dx$$

input `int(x^2*(a + b*atanh(c/x))^3,x)`output `int(x^2*(a + b*atanh(c/x))^3, x)`**Reduce [F]**

$$\begin{aligned} \int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx &= -\frac{\operatorname{atanh} \left(\frac{c}{x} \right)^3 b^3 c^2 x}{3} + \frac{\operatorname{atanh} \left(\frac{c}{x} \right)^3 b^3 x^3}{3} \\ &\quad - \operatorname{atanh} \left(\frac{c}{x} \right)^2 a b^2 c^2 x + \operatorname{atanh} \left(\frac{c}{x} \right)^2 a b^2 x^3 \\ &\quad - \frac{\operatorname{atanh} \left(\frac{c}{x} \right)^2 b^3 c^3}{2} + \frac{\operatorname{atanh} \left(\frac{c}{x} \right)^2 b^3 c x^2}{2} - \operatorname{atanh} \left(\frac{c}{x} \right) a^2 b c^3 \\ &\quad + \operatorname{atanh} \left(\frac{c}{x} \right) a^2 b x^3 - \operatorname{atanh} \left(\frac{c}{x} \right) a b^2 c^3 \\ &\quad + \operatorname{atanh} \left(\frac{c}{x} \right) a b^2 c x^2 - \operatorname{atanh} \left(\frac{c}{x} \right) b^3 c^3 \\ &\quad + \operatorname{atanh} \left(\frac{c}{x} \right) b^3 c^2 x + \frac{\left(\int \operatorname{atanh} \left(\frac{c}{x} \right)^3 dx \right) b^3 c^2}{3} \\ &\quad + \left(\int \operatorname{atanh} \left(\frac{c}{x} \right)^2 dx \right) a b^2 c^2 + \log(-c - x) a^2 b c^3 \\ &\quad + \log(-c - x) b^3 c^3 + \frac{a^3 x^3}{3} + \frac{a^2 b c x^2}{2} + a b^2 c^2 x \end{aligned}$$

input `int(x^2*(a+b*atanh(c/x))^3,x)`

output

```
( - 2*atanh(c/x)**3*b**3*c**2*x + 2*atanh(c/x)**3*b**3*x**3 - 6*atanh(c/x)
**2*a*b**2*c**2*x + 6*atanh(c/x)**2*a*b**2*x**3 - 3*atanh(c/x)**2*b**3*c**
3 + 3*atanh(c/x)**2*b**3*c*x**2 - 6*atanh(c/x)*a**2*b*c**3 + 6*atanh(c/x)*
a**2*b*x**3 - 6*atanh(c/x)*a*b**2*c**3 + 6*atanh(c/x)*a*b**2*c*x**2 - 6*at
anh(c/x)*b**3*c**3 + 6*atanh(c/x)*b**3*c**2*x + 2*int(atanh(c/x)**3,x)*b**
3*c**2 + 6*int(atanh(c/x)**2,x)*a*b**2*c**2 + 6*log( - c - x)*a**2*b*c**3
+ 6*log( - c - x)*b**3*c**3 + 2*a**3*x**3 + 3*a**2*b*c*x**2 + 6*a*b**2*c**
2*x)/6
```

3.152 $\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx$

Optimal result	1245
Mathematica [A] (verified)	1246
Rubi [A] (verified)	1247
Maple [C] (warning: unable to verify)	1250
Fricas [F]	1250
Sympy [F]	1251
Maxima [F]	1251
Giac [F]	1251
Mupad [F(-1)]	1252
Reduce [F]	1252

Optimal result

Integrand size = 14, antiderivative size = 135

$$\begin{aligned} \int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = & -\frac{3}{2} b c^2 \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2 + \frac{3}{2} b c x \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2 \\ & - \frac{1}{2} c^2 \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^3 + \frac{1}{2} x^2 \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^3 \\ & - 3 b^2 c^2 \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right) \log \left(2 - \frac{2}{1 + \frac{c}{x}} \right) \\ & + \frac{3}{2} b^3 c^2 \operatorname{PolyLog} \left(2, -1 + \frac{2}{1 + \frac{c}{x}} \right) \end{aligned}$$

output

```
-3/2*b*c^2*(a+b*arccoth(x/c))^2+3/2*b*c*x*(a+b*arccoth(x/c))^2-1/2*c^2*(a+
b*arccoth(x/c))^3+1/2*x^2*(a+b*arccoth(x/c))^3-3*b^2*c^2*(a+b*arccoth(x/c)
)*ln(2-2/(1+c/x))+3/2*b^3*c^2*polylog(2,-1+2/(1+c/x))
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.43

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \frac{1}{4} \left(6b^2(-c+x)(bc+a(c+x)) \operatorname{arctanh} \left(\frac{c}{x} \right)^2 \right. \\ \left. + 2b^3(-c^2+x^2) \operatorname{arctanh} \left(\frac{c}{x} \right)^3 \right. \\ \left. + 6b \operatorname{arctanh} \left(\frac{c}{x} \right) \left(ax(2bc+ax) - 2b^2c^2 \log \left(1 - e^{-2 \operatorname{arctanh} \left(\frac{c}{x} \right)} \right) \right) \right. \\ \left. + a \left(3abc^2 \log \left(1 - \frac{c}{x} \right) - 12b^2c^2 \log \left(\frac{c}{\sqrt{1 - \frac{c^2}{x^2}}x} \right) \right) \right. \\ \left. + a \left(6bcx + 2ax^2 - 3bc^2 \log \left(\frac{c+x}{x} \right) \right) \right) \\ \left. + 6b^3c^2 \operatorname{PolyLog} \left(2, e^{-2 \operatorname{arctanh} \left(\frac{c}{x} \right)} \right) \right)$$

input `Integrate[x*(a + b*ArcTanh[c/x])^3,x]`output `(6*b^2*(-c + x)*(b*c + a*(c + x))*ArcTanh[c/x]^2 + 2*b^3*(-c^2 + x^2)*ArcTanh[c/x]^3 + 6*b*ArcTanh[c/x]*(a*x*(2*b*c + a*x) - 2*b^2*c^2*Log[1 - E^(-2*ArcTanh[c/x])]) + a*(3*a*b*c^2*Log[1 - c/x] - 12*b^2*c^2*Log[c/(Sqrt[1 - c^2/x^2]*x)] + a*(6*b*c*x + 2*a*x^2 - 3*b*c^2*Log[(c + x)/x])) + 6*b^3*c^2*PolyLog[2, E^(-2*ArcTanh[c/x])])/4`

Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.97, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {6454, 6452, 6544, 6452, 6510, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^3 dx \\
 & \quad \downarrow \text{6454} \\
 & - \int x^3 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^3 d\frac{1}{x} \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{2}x^2 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^3 - \frac{3}{2}bc \int \frac{x^2 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} \\
 & \quad \downarrow \text{6544} \\
 & \frac{1}{2}x^2 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^3 - \\
 & \frac{3}{2}bc \left(c^2 \int \frac{\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \int x^2 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2 d\frac{1}{x} \right) \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{2}x^2 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^3 - \\
 & \frac{3}{2}bc \left(c^2 \int \frac{\left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + 2bc \int \frac{x \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - x \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2 \right) \\
 & \quad \downarrow \text{6510} \\
 & \frac{1}{2}x^2 \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^3 - \\
 & \frac{3}{2}bc \left(2bc \int \frac{x \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{c \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^3}{3b} - x \left(a + \operatorname{barctanh}\left(\frac{c}{x}\right) \right)^2 \right) \\
 & \quad \downarrow \text{6550}
 \end{aligned}$$

$$\frac{3}{2}bc \left(2bc \left(\int \frac{x(a + \operatorname{arctanh}(\frac{c}{x}))}{\frac{c}{x} + 1} d\frac{1}{x} + \frac{(a + \operatorname{arctanh}(\frac{c}{x}))^2}{2b} \right) + \frac{c(a + \operatorname{arctanh}(\frac{c}{x}))^3}{3b} - x(a + \operatorname{arctanh}(\frac{c}{x}))^2 \right)$$

↓ 6494

$$\frac{3}{2}bc \left(2bc \left(-bc \int \frac{\log\left(2 - \frac{2}{\frac{c}{x} + 1}\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} + \frac{(a + \operatorname{arctanh}(\frac{c}{x}))^2}{2b} + \log\left(2 - \frac{2}{\frac{c}{x} + 1}\right) (a + \operatorname{arctanh}(\frac{c}{x})) \right) + \frac{c(a + \operatorname{arctanh}(\frac{c}{x}))^3}{3b} - x(a + \operatorname{arctanh}(\frac{c}{x}))^2 \right)$$

↓ 2897

$$\frac{3}{2}bc \left(2bc \left(\frac{(a + \operatorname{arctanh}(\frac{c}{x}))^2}{2b} + \log\left(2 - \frac{2}{\frac{c}{x} + 1}\right) (a + \operatorname{arctanh}(\frac{c}{x})) - \frac{1}{2}b \operatorname{PolyLog}\left(2, \frac{2}{\frac{c}{x} + 1} - 1\right) \right) + \frac{c(a + \operatorname{arctanh}(\frac{c}{x}))^3}{3b} - x(a + \operatorname{arctanh}(\frac{c}{x}))^2 \right)$$

input `Int [x*(a + b*ArcTanh [c/x])^3,x]`

output `(x^2*(a + b*ArcTanh [c/x])^3)/2 - (3*b*c*(-(x*(a + b*ArcTanh [c/x])^2) + (c*(a + b*ArcTanh [c/x])^3)/(3*b) + 2*b*c*((a + b*ArcTanh [c/x])^2/(2*b) + (a + b*ArcTanh [c/x])*Log [2 - 2/(1 + c/x)] - (b*PolyLog [2, -1 + 2/(1 + c/x)])/2)))/2`

Defintions of rubi rules used

rule 2897 `Int [Log [u_]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh [c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh [c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 $\text{Int}[(a + \text{ArcTanh}[c \cdot x]^n] \cdot b)^p \cdot x^m, x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{\text{Simplify}[(m+1)/n] - 1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x, x^n], x] /; \text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$

rule 6494 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((x) \cdot ((d) + (e) \cdot x)), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{Log}[2 - 2/(1 + e \cdot (x/d))]/d), x] - \text{Simp}[b \cdot c \cdot (p/d) \text{ Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{Log}[2 - 2/(1 + e \cdot (x/d))]) / (1 - c^2 \cdot x^2)], x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d^2 - e^2, 0]$

rule 6510 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((d) + (e) \cdot x^2), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot c \cdot d \cdot (p+1)), x] /; \text{FreeQ}\{a, b, c, d, e, p\}, x \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{NeQ}[p, -1]$

rule 6544 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p \cdot (f \cdot x)^m / ((d) + (e) \cdot x^2), x_Symbol] \rightarrow \text{Simp}[1/d \text{ Int}[(f \cdot x)^m \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x] - \text{Simp}[e/(d \cdot f^2) \text{ Int}[(f \cdot x)^{m+2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2)], x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{LtQ}[m, -1]$

rule 6550 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((x) \cdot ((d) + (e) \cdot x^2)), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot d \cdot (p+1)), x] + \text{Simp}[1/d \text{ Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p / (x \cdot (1 + c \cdot x)), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{GtQ}[p, 0]$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 22.59 (sec) , antiderivative size = 5036, normalized size of antiderivative = 37.30

method	result	size
derivativedivides	Expression too large to display	5036
default	Expression too large to display	5036
parts	Expression too large to display	5038
risch	Expression too large to display	48058

input `int(x*(a+b*arctanh(c/x))^3,x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F]

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int \left(b \operatorname{arctanh} \left(\frac{c}{x} \right) + a \right)^3 x dx$$

input `integrate(x*(a+b*arctanh(c/x))^3,x, algorithm="fricas")`

output `integral(b^3*x*arctanh(c/x)^3 + 3*a*b^2*x*arctanh(c/x)^2 + 3*a^2*b*x*arctanh(c/x) + a^3*x, x)`

Sympy [F]

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int x \left(a + b \operatorname{atanh} \left(\frac{c}{x} \right) \right)^3 dx$$

input `integrate(x*(a+b*atanh(c/x))**3,x)`

output `Integral(x*(a + b*atanh(c/x))**3, x)`

Maxima [F]

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x} \right) + a \right)^3 x dx$$

input `integrate(x*(a+b*arctanh(c/x))^3,x, algorithm="maxima")`

output `3/2*a*b^2*x^2*arctanh(c/x)^2 + 1/2*a^3*x^2 + 3/4*(2*x^2*arctanh(c/x) - (c*log(c + x) - c*log(-c + x) - 2*x)*c)*a^2*b + 3/8*((log(c + x)^2 - 2*(log(c + x) - 2)*log(-c + x) + log(-c + x)^2 + 4*log(c + x))*c^2 - 4*(c*log(c + x) - c*log(-c + x) - 2*x)*c*arctanh(c/x))*a*b^2 + 1/16*(6*c*x*log(c + x)^2 - (c^2 - x^2)*log(c + x)^3 + (c^2 - x^2)*log(-c + x)^3 - 3*(2*c^2 - 2*c*x + (c^2 - x^2)*log(c + x))*log(-c + x)^2 + 3*((c^2 - x^2)*log(c + x)^2 - 4*(c^2 + c*x)*log(c + x))*log(-c + x) + 2*integrate(-6*(c^3 + 3*c^2*x)*log(c + x)/(c^2 - x^2), x))*b^3`

Giac [F]

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x} \right) + a \right)^3 x dx$$

input `integrate(x*(a+b*arctanh(c/x))^3,x, algorithm="giac")`

output `integrate((b*arctanh(c/x) + a)^3*x, x)`

Mupad [F(-1)]

Timed out.

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int x \left(a + b \operatorname{atanh} \left(\frac{c}{x} \right) \right)^3 dx$$

input `int(x*(a + b*atanh(c/x))^3,x)`

output `int(x*(a + b*atanh(c/x))^3, x)`

Reduce [F]

$$\begin{aligned} \int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = & -\frac{\operatorname{atanh} \left(\frac{c}{x} \right)^3 b^3 c^2}{2} + \frac{\operatorname{atanh} \left(\frac{c}{x} \right)^3 b^3 x^2}{2} \\ & - \frac{3 \operatorname{atanh} \left(\frac{c}{x} \right)^2 a b^2 c^2}{2} + \frac{3 \operatorname{atanh} \left(\frac{c}{x} \right)^2 a b^2 x^2}{2} \\ & + \frac{3 \operatorname{atanh} \left(\frac{c}{x} \right)^2 b^3 c x}{2} - \frac{3 \operatorname{atanh} \left(\frac{c}{x} \right) a^2 b c^2}{2} \\ & + \frac{3 \operatorname{atanh} \left(\frac{c}{x} \right) a^2 b x^2}{2} - 3 \operatorname{atanh} \left(\frac{c}{x} \right) a b^2 c^2 \\ & + 3 \operatorname{atanh} \left(\frac{c}{x} \right) a b^2 c x - 3 \left(\int \frac{\operatorname{atanh} \left(\frac{c}{x} \right) x}{c^2 - x^2} dx \right) b^3 c^2 \\ & + 3 \log(-c - x) a b^2 c^2 + \frac{a^3 x^2}{2} + \frac{3 a^2 b c x}{2} \end{aligned}$$

input `int(x*(a+b*atanh(c/x))^3,x)`

output

```
( - atanh(c/x)**3*b**3*c**2 + atanh(c/x)**3*b**3*x**2 - 3*atanh(c/x)**2*a*
b**2*c**2 + 3*atanh(c/x)**2*a*b**2*x**2 + 3*atanh(c/x)**2*b**3*c*x - 3*ata
nh(c/x)*a**2*b*c**2 + 3*atanh(c/x)*a**2*b*x**2 - 6*atanh(c/x)*a*b**2*c**2
+ 6*atanh(c/x)*a*b**2*c*x - 6*int((atanh(c/x)*x)/(c**2 - x**2),x)*b**3*c**
2 + 6*log( - c - x)*a*b**2*c**2 + a**3*x**2 + 3*a**2*b*c*x)/2
```

3.153 $\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^3 dx$

Optimal result	1254
Mathematica [C] (verified)	1255
Rubi [A] (verified)	1256
Maple [C] (warning: unable to verify)	1258
Fricas [F]	1259
Sympy [F]	1260
Maxima [F]	1260
Giac [F]	1260
Mupad [F(-1)]	1261
Reduce [F]	1261

Optimal result

Integrand size = 12, antiderivative size = 108

$$\begin{aligned} \int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^3 dx &= c \left(a + b \operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^3 + x \left(a + b \operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^3 \\ &\quad - 3bc \left(a + b \operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^2 \log\left(\frac{2c}{c-x}\right) \\ &\quad - 3b^2c \left(a + b \operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right) \operatorname{PolyLog}\left(2, 1 - \frac{2c}{c-x}\right) \\ &\quad + \frac{3}{2}b^3c \operatorname{PolyLog}\left(3, 1 - \frac{2c}{c-x}\right) \end{aligned}$$

output

```
c*(a+b*arccoth(x/c))^3+x*(a+b*arccoth(x/c))^3-3*b*c*(a+b*arccoth(x/c))^2*ln(2*c/(c-x))-3*b^2*c*(a+b*arccoth(x/c))*polylog(2,1-2*c/(c-x))+3/2*b^3*c*polylog(3,1-2*c/(c-x))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.20 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.83

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right)^3 dx = a^3 x + 3a^2 b x \operatorname{arctanh}\left(\frac{c}{x}\right) + \frac{3}{2} a^2 b c \log(-c^2 + x^2) - 3ab^2 \left(\operatorname{arctanh}\left(\frac{c}{x}\right) \left((c-x) \operatorname{arctanh}\left(\frac{c}{x}\right) + 2c \log\left(1 - e^{-2\operatorname{arctanh}\left(\frac{c}{x}\right)}\right) - c \operatorname{PolyLog}\left(2, e^{-2\operatorname{arctanh}\left(\frac{c}{x}\right)}\right) \right) + \frac{1}{8} b^3 \left(-ic\pi^3 + 8c \operatorname{arctanh}\left(\frac{c}{x}\right)^3 + 8x \operatorname{arctanh}\left(\frac{c}{x}\right)^3 - 24c \operatorname{arctanh}\left(\frac{c}{x}\right)^2 \log\left(1 - e^{2\operatorname{arctanh}\left(\frac{c}{x}\right)}\right) - 24c \operatorname{arctanh}\left(\frac{c}{x}\right) \operatorname{PolyLog}\left(2, e^{2\operatorname{arctanh}\left(\frac{c}{x}\right)}\right) + 12c \operatorname{PolyLog}\left(3, e^{2\operatorname{arctanh}\left(\frac{c}{x}\right)}\right) \right)$$

input `Integrate[(a + b*ArcTanh[c/x])^3,x]`

output `a^3*x + 3*a^2*b*x*ArcTanh[c/x] + (3*a^2*b*c*Log[-c^2 + x^2])/2 - 3*a*b^2*(ArcTanh[c/x]*((c - x)*ArcTanh[c/x] + 2*c*Log[1 - E^(-2*ArcTanh[c/x])]) - c*PolyLog[2, E^(-2*ArcTanh[c/x])]) + (b^3*((-I)*c*Pi^3 + 8*c*ArcTanh[c/x]^3 + 8*x*ArcTanh[c/x]^3 - 24*c*ArcTanh[c/x]^2*Log[1 - E^(2*ArcTanh[c/x])] - 24*c*ArcTanh[c/x]*PolyLog[2, E^(2*ArcTanh[c/x])] + 12*c*PolyLog[3, E^(2*ArcTanh[c/x])]))/8`

Rubi [A] (verified)

Time = 1.07 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.12, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6440, 6437, 27, 6547, 27, 6471, 27, 6621, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx \\
 & \quad \downarrow 6440 \\
 & \int \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^3 dx \\
 & \quad \downarrow 6437 \\
 & x \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^3 - \frac{3b \int \frac{c^2 x (a + b \coth^{-1}(\frac{x}{c}))^2}{c^2 - x^2} dx}{c} \\
 & \quad \downarrow 27 \\
 & x \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^3 - 3bc \int \frac{x (a + b \coth^{-1}(\frac{x}{c}))^2}{c^2 - x^2} dx \\
 & \quad \downarrow 6547 \\
 & x \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^3 - 3bc \left(\frac{\int \frac{c (a + b \coth^{-1}(\frac{x}{c}))^2}{c - x} dx}{c} - \frac{(a + b \coth^{-1}(\frac{x}{c}))^3}{3b} \right) \\
 & \quad \downarrow 27 \\
 & x \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^3 - 3bc \left(\int \frac{(a + b \coth^{-1}(\frac{x}{c}))^2}{c - x} dx - \frac{(a + b \coth^{-1}(\frac{x}{c}))^3}{3b} \right) \\
 & \quad \downarrow 6471 \\
 & x \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^3 - \\
 & 3bc \left(-\frac{2b \int \frac{c^2 (a + b \coth^{-1}(\frac{x}{c})) \log(\frac{2c}{c-x})}{c^2 - x^2} dx}{c} - \frac{(a + b \coth^{-1}(\frac{x}{c}))^3}{3b} + \log \left(\frac{2c}{c-x} \right) \left(a + b \coth^{-1} \left(\frac{x}{c} \right) \right)^2 \right) \\
 & \quad \downarrow 27
 \end{aligned}$$

$$\begin{aligned}
& 3bc \left(-2bc \int \frac{(a + b \coth^{-1}(\frac{x}{c})) \log\left(\frac{2c}{c-x}\right)}{c^2 - x^2} dx - \frac{(a + b \coth^{-1}(\frac{x}{c}))^3}{3b} + \log\left(\frac{2c}{c-x}\right) (a + b \coth^{-1}(\frac{x}{c}))^2 \right) \\
& \quad \downarrow \text{6621} \\
& 3bc \left(-2bc \left(\frac{1}{2} b \int \frac{\text{PolyLog}\left(2, 1 - \frac{2c}{c-x}\right)}{c^2 - x^2} dx - \frac{\text{PolyLog}\left(2, 1 - \frac{2c}{c-x}\right) (a + b \coth^{-1}(\frac{x}{c}))}{2c} \right) - \frac{(a + b \coth^{-1}(\frac{x}{c}))^3}{3b} \right) \\
& \quad \downarrow \text{7164} \\
& 3bc \left(-2bc \left(\frac{b \text{PolyLog}\left(3, 1 - \frac{2c}{c-x}\right)}{4c} - \frac{\text{PolyLog}\left(2, 1 - \frac{2c}{c-x}\right) (a + b \coth^{-1}(\frac{x}{c}))}{2c} \right) - \frac{(a + b \coth^{-1}(\frac{x}{c}))^3}{3b} + \log\left(\frac{2c}{c-x}\right) (a + b \coth^{-1}(\frac{x}{c}))^2 \right)
\end{aligned}$$

input `Int[(a + b*ArcTanh[c/x])^3,x]`

output `x*(a + b*ArcCoth[x/c])^3 - 3*b*c*(-1/3*(a + b*ArcCoth[x/c])^3/b + (a + b*ArcCoth[x/c])^2*Log[(2*c)/(c - x)] - 2*b*c*(-1/2*((a + b*ArcCoth[x/c])*PolyLog[2, 1 - (2*c)/(c - x)])/c + (b*PolyLog[3, 1 - (2*c)/(c - x)]/(4*c)))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 6437 `Int[((a_) + ArcCoth[(c_)*(x_)^(n_)])*(b_)^(p_), x_Symbol] := Simp[x*(a + b*ArcCoth[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcCoth[c*x^n])^(p-1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`

rule 6440 $\text{Int}[(a + b \text{ArcTanh}[c \cdot x]^n) \cdot (b \cdot x)^p, x_Symbol] \rightarrow \text{Int}[(a + b \text{ArcCoth}[1/(x^n \cdot c)])^p, x] /;$ FreeQ[{a, b, c}, x] && IGtQ[p, 1] && ILtQ[n, 0]

rule 6471 $\text{Int}[(a + b \text{ArcCoth}[c \cdot x] \cdot (b \cdot x))^p / ((d + e \cdot x) \cdot x), x_Symbol] \rightarrow \text{Simp}[(-a + b \text{ArcCoth}[c \cdot x])^p \cdot (\text{Log}[2/(1 + e \cdot (x/d))]) / e, x] + \text{Simp}[b \cdot c \cdot (p/e) \text{Int}[(a + b \text{ArcCoth}[c \cdot x])^{p-1} \cdot (\text{Log}[2/(1 + e \cdot (x/d))]) / (1 - c^2 \cdot x^2), x], x] /;$ FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2 \cdot d^2 - e^2, 0]

rule 6547 $\text{Int}[(a + b \text{ArcCoth}[c \cdot x] \cdot (b \cdot x))^p \cdot (x) / ((d + e \cdot x) \cdot x^2), x_Symbol] \rightarrow \text{Simp}[(a + b \text{ArcCoth}[c \cdot x])^{p+1} / (b \cdot e \cdot (p+1)), x] + \text{Simp}[1 / (c \cdot d) \text{Int}[(a + b \text{ArcCoth}[c \cdot x])^p / (1 - c \cdot x), x], x] /;$ FreeQ[{a, b, c, d, e}, x] && EqQ[c^2 \cdot d + e, 0] && IGtQ[p, 0]

rule 6621 $\text{Int}[(\text{Log}[u] \cdot (a + b \text{ArcCoth}[c \cdot x] \cdot (b \cdot x))^p) / ((d + e \cdot x) \cdot x^2), x_Symbol] \rightarrow \text{Simp}[(-a + b \text{ArcCoth}[c \cdot x])^p \cdot (\text{PolyLog}[2, 1 - u] / (2 \cdot c \cdot d)), x] + \text{Simp}[b \cdot (p/2) \text{Int}[(a + b \text{ArcCoth}[c \cdot x])^{p-1} \cdot (\text{PolyLog}[2, 1 - u] / (d + e \cdot x^2)), x], x] /;$ FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2 \cdot d + e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c \cdot x))^2, 0]

rule 7164 $\text{Int}(u \cdot \text{PolyLog}[n, v], x_Symbol) \rightarrow \text{With}[\{w = \text{DerivativeDivides}[v, u \cdot v, x]\}, \text{Simp}[w \cdot \text{PolyLog}[n + 1, v], x] /;$!FalseQ[w]] /;

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 27.39 (sec) , antiderivative size = 1475, normalized size of antiderivative = 13.66

method	result	size
parts	Expression too large to display	1475
derivativedivides	Expression too large to display	1478
default	Expression too large to display	1478

input `int((a+b*arctanh(c/x))^3,x,method=_RETURNVERBOSE)`

output `x*a^3-b^3*c*(-1/c*x*arctanh(c/x)^3+3*ln(c/x)*arctanh(c/x)^2-3/2*arctanh(c/x)^2*ln(c/x-1)-3/2*arctanh(c/x)^2*ln(1+c/x)+3*arctanh(c/x)^2*ln((1+c/x)/(1-c^2/x^2)^(1/2))-arctanh(c/x)^3+3/4*(I*Pi*csgn(I*(1+c/x)^2/(c^2/x^2-1))^3+2*I*Pi-2*I*Pi*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^2+2*I*Pi*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^3+2*I*Pi*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))^3+I*Pi*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))*csgn(I*(1+c/x)^2/(c^2/x^2-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^2-2*I*Pi*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1))*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^2-2*I*Pi*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))^2-I*Pi*csgn(I*(1+c/x)^2/(c^2/x^2-1))*csgn(I*(1+c/x)^2/(c^2/x^2-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^2+I*Pi*csgn(I*(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1))^3+2*I*Pi*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1))*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^2+2*I*Pi*csgn(I*(1+c/x)/(1-c^2/x^2)^(1/2))*csgn(I*(1+c/x)^2/(c^2/x^2-1))^2+I*Pi*csgn(I*(1+c/x)/(1-c^2/x^2)^(1/2))^2*csgn(I*(1+c/x)^2/(c^2/x^2-1))-I*Pi*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))*csgn(I*(1+c/x)^2/(c^2/x^2-1))*csgn(I*(1+c/x)^2/(c^2/x^2-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^2+4*ln(2)*arctanh(c/x)^2-3*arctanh(c/x)^2*ln((1+c/x)^2/(1-c^2/x^2)-1)+3*arctanh(c/x)^2*ln(1+(1+c/x)/(1-c^2/x^2)^(1/2))+6*arctanh(c/x)*polylog(2,-(1+c/x)/(1-c^2/x^2)^(1/2))-6*polylog(3,-(1+c/x)/(1-c^2/x^2)^(1/2))+3*arctanh(c/x)...`

Fricas [F]

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int \left(b \operatorname{arctanh} \left(\frac{c}{x} \right) + a \right)^3 dx$$

input `integrate((a+b*arctanh(c/x))^3,x, algorithm="fricas")`

output `integral(b^3*arctanh(c/x)^3 + 3*a*b^2*arctanh(c/x)^2 + 3*a^2*b*arctanh(c/x) + a^3, x)`

Sympy [F]

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int \left(a + b \operatorname{atanh} \left(\frac{c}{x} \right) \right)^3 dx$$

input `integrate((a+b*atanh(c/x))**3,x)`

output `Integral((a + b*atanh(c/x))**3, x)`

Maxima [F]

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x} \right) + a \right)^3 dx$$

input `integrate((a+b*arctanh(c/x))^3,x, algorithm="maxima")`

output `3/2*(2*x*arctanh(c/x) + c*log(-c^2 + x^2))*a^2*b + a^3*x + 1/8*(b^3*c - b^3*x)*log(-c + x)^3 + 3/8*(2*a*b^2*x + (b^3*c + b^3*x)*log(c + x))*log(-c + x)^2 - integrate(-1/8*((b^3*c - b^3*x)*log(c + x)^3 + 6*(a*b^2*c - a*b^2*x)*log(c + x)^2 + 3*(4*a*b^2*x - (b^3*c - b^3*x)*log(c + x)^2 - 2*(2*a*b^2*c - b^3*c - (2*a*b^2 + b^3)*x)*log(c + x))*log(-c + x))/(c - x), x)`

Giac [F]

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x} \right) \right)^3 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x} \right) + a \right)^3 dx$$

input `integrate((a+b*arctanh(c/x))^3,x, algorithm="giac")`

output `integrate((b*arctanh(c/x) + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right)^3 dx = \int \left(a + b \operatorname{atanh}\left(\frac{c}{x}\right) \right)^3 dx$$

input `int((a + b*atanh(c/x))^3,x)`output `int((a + b*atanh(c/x))^3, x)`**Reduce [F]**

$$\begin{aligned} \int \left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right) \right)^3 dx &= -3 \operatorname{atanh}\left(\frac{c}{x}\right) a^2 b c + 3 \operatorname{atanh}\left(\frac{c}{x}\right) a^2 b x \\ &+ \left(\int \operatorname{atanh}\left(\frac{c}{x}\right)^3 dx \right) b^3 + 3 \left(\int \operatorname{atanh}\left(\frac{c}{x}\right)^2 dx \right) a b^2 \\ &+ 3 \log(-c - x) a^2 b c + a^3 x \end{aligned}$$

input `int((a+b*atanh(c/x))^3,x)`output `- 3*atanh(c/x)*a**2*b*c + 3*atanh(c/x)*a**2*b*x + int(atanh(c/x)**3,x)*b*
*3 + 3*int(atanh(c/x)**2,x)*a*b**2 + 3*log(- c - x)*a**2*b*c + a**3*x`

$$3.154 \quad \int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^3}{x} dx$$

Optimal result	1262
Mathematica [C] (verified)	1263
Rubi [A] (verified)	1265
Maple [C] (warning: unable to verify)	1268
Fricas [F]	1269
Sympy [F]	1269
Maxima [F]	1269
Giac [F]	1270
Mupad [F(-1)]	1270
Reduce [F]	1270

Optimal result

Integrand size = 16, antiderivative size = 208

$$\begin{aligned} \int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^3}{x} dx = & -2 \left(a + b \coth^{-1}\left(\frac{x}{c}\right)\right)^3 \operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{c}{x}}\right) \\ & + \frac{3}{2} b \left(a + b \coth^{-1}\left(\frac{x}{c}\right)\right)^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{c}{x}}\right) \\ & - \frac{3}{2} b \left(a + b \coth^{-1}\left(\frac{x}{c}\right)\right)^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - \frac{c}{x}}\right) \\ & - \frac{3}{2} b^2 \left(a + b \coth^{-1}\left(\frac{x}{c}\right)\right) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - \frac{c}{x}}\right) \\ & + \frac{3}{2} b^2 \left(a + b \coth^{-1}\left(\frac{x}{c}\right)\right) \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 - \frac{c}{x}}\right) \\ & + \frac{3}{4} b^3 \operatorname{PolyLog}\left(4, 1 - \frac{2}{1 - \frac{c}{x}}\right) \\ & - \frac{3}{4} b^3 \operatorname{PolyLog}\left(4, -1 + \frac{2}{1 - \frac{c}{x}}\right) \end{aligned}$$

output

```
2*(a+b*arccoth(x/c))^3*arctanh(-1+2/(1-c/x))+3/2*b*(a+b*arccoth(x/c))^2*po
lylog(2,1-2/(1-c/x))-3/2*b*(a+b*arccoth(x/c))^2*polylog(2,-1+2/(1-c/x))-3/
2*b^2*(a+b*arccoth(x/c))*polylog(3,1-2/(1-c/x))+3/2*b^2*(a+b*arccoth(x/c))
*polylog(3,-1+2/(1-c/x))+3/4*b^3*polylog(4,1-2/(1-c/x))-3/4*b^3*polylog(4,
-1+2/(1-c/x))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.32 (sec) , antiderivative size = 373, normalized size of antiderivative = 1.79

$$\begin{aligned}
\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x} dx = & a^3 \log(x) + \frac{3}{2} a^2 b \left(\operatorname{PolyLog}\left(2, -\frac{c}{x}\right) - \operatorname{PolyLog}\left(2, \frac{c}{x}\right) \right) \\
& + 3ab^2 \left(-\frac{i\pi^3}{24} + \frac{2}{3} \operatorname{arctanh}\left(\frac{c}{x}\right)^3 \right. \\
& \quad + \operatorname{arctanh}\left(\frac{c}{x}\right)^2 \log\left(1 + e^{-2\operatorname{arctanh}(\frac{c}{x})}\right) \\
& \quad - \operatorname{arctanh}\left(\frac{c}{x}\right)^2 \log\left(1 - e^{2\operatorname{arctanh}(\frac{c}{x})}\right) \\
& \quad - \operatorname{arctanh}\left(\frac{c}{x}\right) \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arctanh}(\frac{c}{x})}\right) \\
& \quad - \operatorname{arctanh}\left(\frac{c}{x}\right) \operatorname{PolyLog}\left(2, e^{2\operatorname{arctanh}(\frac{c}{x})}\right) \\
& \quad - \frac{1}{2} \operatorname{PolyLog}\left(3, -e^{-2\operatorname{arctanh}(\frac{c}{x})}\right) \\
& \quad \left. + \frac{1}{2} \operatorname{PolyLog}\left(3, e^{2\operatorname{arctanh}(\frac{c}{x})}\right) \right) + \frac{1}{64} b^3 \left(-\pi^4 \right. \\
& + 32 \operatorname{arctanh}\left(\frac{c}{x}\right)^4 + 64 \operatorname{arctanh}\left(\frac{c}{x}\right)^3 \log\left(1 + e^{-2\operatorname{arctanh}(\frac{c}{x})}\right) \\
& \quad - 64 \operatorname{arctanh}\left(\frac{c}{x}\right)^3 \log\left(1 - e^{2\operatorname{arctanh}(\frac{c}{x})}\right) \\
& \quad - 96 \operatorname{arctanh}\left(\frac{c}{x}\right)^2 \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arctanh}(\frac{c}{x})}\right) \\
& \quad - 96 \operatorname{arctanh}\left(\frac{c}{x}\right)^2 \operatorname{PolyLog}\left(2, e^{2\operatorname{arctanh}(\frac{c}{x})}\right) \\
& \quad - 96 \operatorname{arctanh}\left(\frac{c}{x}\right) \operatorname{PolyLog}\left(3, -e^{-2\operatorname{arctanh}(\frac{c}{x})}\right) \\
& \quad + 96 \operatorname{arctanh}\left(\frac{c}{x}\right) \operatorname{PolyLog}\left(3, e^{2\operatorname{arctanh}(\frac{c}{x})}\right) \\
& \quad - 48 \operatorname{PolyLog}\left(4, -e^{-2\operatorname{arctanh}(\frac{c}{x})}\right) \\
& \quad \left. - 48 \operatorname{PolyLog}\left(4, e^{2\operatorname{arctanh}(\frac{c}{x})}\right) \right)
\end{aligned}$$

input

```
Integrate[(a + b*ArcTanh[c/x])^3/x, x]
```

output

```

a^3*Log[x] + (3*a^2*b*(PolyLog[2, -(c/x)] - PolyLog[2, c/x]))/2 + 3*a*b^2*
((-1/24*I)*Pi^3 + (2*ArcTanh[c/x]^3)/3 + ArcTanh[c/x]^2*Log[1 + E^(-2*ArcT
anh[c/x])] - ArcTanh[c/x]^2*Log[1 - E^(2*ArcTanh[c/x])] - ArcTanh[c/x]*Pol
yLog[2, -E^(-2*ArcTanh[c/x])] - ArcTanh[c/x]*PolyLog[2, E^(2*ArcTanh[c/x])
] - PolyLog[3, -E^(-2*ArcTanh[c/x])]/2 + PolyLog[3, E^(2*ArcTanh[c/x])]/2)
+ (b^3*(-Pi^4 + 32*ArcTanh[c/x]^4 + 64*ArcTanh[c/x]^3*Log[1 + E^(-2*ArcTa
nh[c/x])] - 64*ArcTanh[c/x]^3*Log[1 - E^(2*ArcTanh[c/x])] - 96*ArcTanh[c/x
]^2*PolyLog[2, -E^(-2*ArcTanh[c/x])] - 96*ArcTanh[c/x]^2*PolyLog[2, E^(2*A
rcTanh[c/x])] - 96*ArcTanh[c/x]*PolyLog[3, -E^(-2*ArcTanh[c/x])] + 96*ArcT
anh[c/x]*PolyLog[3, E^(2*ArcTanh[c/x])] - 48*PolyLog[4, -E^(-2*ArcTanh[c/x
])] - 48*PolyLog[4, E^(2*ArcTanh[c/x])]))/64

```

Rubi [A] (verified)

Time = 1.72 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.13, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6450, 6448, 6614, 6620, 6624, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x} dx \\
 & \quad \downarrow 6450 \\
 & - \int x (a + b \operatorname{arctanh}(\frac{c}{x}))^3 d\frac{1}{x} \\
 & \quad \downarrow 6448 \\
 & 6bc \int \frac{\operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{c}{x}}\right) (a + b \operatorname{arctanh}(\frac{c}{x}))^2}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - \\
 & \quad 2 \operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{c}{x}}\right) (a + b \operatorname{arctanh}(\frac{c}{x}))^3 \\
 & \quad \downarrow 6614
 \end{aligned}$$

$$6bc \left(\frac{1}{2} \int \frac{(a + \operatorname{arctanh}(\frac{c}{x}))^2 \log\left(2 - \frac{2}{1 - \frac{c}{x}}\right) d\frac{1}{x}}{1 - \frac{c^2}{x^2}} - \frac{1}{2} \int \frac{(a + \operatorname{arctanh}(\frac{c}{x}))^2 \log\left(\frac{2}{1 - \frac{c}{x}}\right) d\frac{1}{x}}{1 - \frac{c^2}{x^2}} \right) - \frac{2 \operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^3}{2c}$$

↓ 6620

$$6bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2}{2c} - b \int \frac{(a + \operatorname{arctanh}(\frac{c}{x})) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{c}{x}}\right) d\frac{1}{x}}{1 - \frac{c^2}{x^2}} \right) + \frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)}{2c} - \frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(4, 1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} \right) \right) - \frac{2 \operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^3}{2c}$$

↓ 6624

$$6bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2}{2c} - b \left(\frac{\operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)}{2c} - \frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(4, 1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} \right) \right) + \frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)}{2c} - \frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(4, 1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} \right) \right) - \frac{2 \operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^3}{2c}$$

↓ 7164

$$6bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^2}{2c} - b \left(\frac{\operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)}{2c} - \frac{b \operatorname{PolyLog}\left(4, 1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)}{1 - \frac{c^2}{x^2}} \right) \right) + \frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)}{2c} - \frac{b \operatorname{PolyLog}\left(4, 1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)}{1 - \frac{c^2}{x^2}} \right) \right) - \frac{2 \operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{c}{x}}\right) \left(a + \operatorname{arctanh}\left(\frac{c}{x}\right)\right)^3}{2c}$$

input `Int[(a + b*ArcTanh[c/x])^3/x,x]`

output `-2*ArcTanh[1 - 2/(1 - c/x)]*(a + b*ArcTanh[c/x])^3 + 6*b*c*(((a + b*ArcTanh[c/x])^2*PolyLog[2, 1 - 2/(1 - c/x)]/(2*c) - b*(((a + b*ArcTanh[c/x])*PolyLog[3, 1 - 2/(1 - c/x)]/(2*c) - (b*PolyLog[4, 1 - 2/(1 - c/x)]/(4*c)))/2 + (-1/2*((a + b*ArcTanh[c/x])^2*PolyLog[2, -1 + 2/(1 - c/x)]/c + b*((a + b*ArcTanh[c/x])*PolyLog[3, -1 + 2/(1 - c/x)]/(2*c) - (b*PolyLog[4, -1 + 2/(1 - c/x)]/(4*c))))/2)`

Defintions of rubi rules used

rule 6448 $\text{Int}[(a + \text{ArcTanh}[c \cdot x])^p \cdot \text{ArcTanh}[1 - 2/(1 - c \cdot x)], x] - \text{Simp}[2 \cdot b \cdot c \cdot p \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot \text{ArcTanh}[1 - 2/(1 - c \cdot x)]/(1 - c^2 \cdot x^2)], x] /;$
 $\text{FreeQ}\{a, b, c, x\} \ \&\& \ \text{IGtQ}[p, 1]$

rule 6450 $\text{Int}[(a + \text{ArcTanh}[c \cdot x]^n) \cdot (b + \text{ArcTanh}[c \cdot x])^p / x, x] /;$
 $\text{FreeQ}\{a, b, c, n, x\} \ \&\& \ \text{IGtQ}[p, 0]$

rule 6614 $\text{Int}[\text{ArcTanh}[u] \cdot (a + \text{ArcTanh}[c \cdot x])^p / ((d + e \cdot x)^2), x] /;$
 $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{EqQ}[u^2 - (1 - 2/(1 - c \cdot x))^2, 0]$

rule 6620 $\text{Int}[(\text{Log}[u] \cdot (a + \text{ArcTanh}[c \cdot x])^p) / ((d + e \cdot x)^2), x] /;$
 $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{EqQ}[(1 - u)^2 - (1 - 2/(1 - c \cdot x))^2, 0]$

rule 6624 $\text{Int}[(a + \text{ArcTanh}[c \cdot x])^p \cdot \text{PolyLog}[k, u] / ((d + e \cdot x)^2), x] /;$
 $\text{FreeQ}\{a, b, c, d, e, k, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{EqQ}[u^2 - (1 - 2/(1 - c \cdot x))^2, 0]$

rule 7164 $\text{Int}[u \cdot \text{PolyLog}[n, v], x] /;$
 $\text{FreeQ}[n, x] \ \&\& \ \text{EqQ}[v, u \cdot v]$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 15.62 (sec) , antiderivative size = 1452, normalized size of antiderivative = 6.98

method	result	size
parts	Expression too large to display	1452
derivativedivides	Expression too large to display	1454
default	Expression too large to display	1454

input `int((a+b*arctanh(c/x))^3/x,x,method=_RETURNVERBOSE)`

output

```

a^3*ln(x)+b^3*(-ln(c/x)*arctanh(c/x)^3+arctanh(c/x)^3*ln((1+c/x)^2/(1-c^2/x^2)-1)-arctanh(c/x)^3*ln(1-(1+c/x)/(1-c^2/x^2)^(1/2))-3*arctanh(c/x)^2*polylog(2,(1+c/x)/(1-c^2/x^2)^(1/2))+6*arctanh(c/x)*polylog(3,(1+c/x)/(1-c^2/x^2)^(1/2))-6*polylog(4,(1+c/x)/(1-c^2/x^2)^(1/2))-arctanh(c/x)^3*ln(1+(1+c/x)/(1-c^2/x^2)^(1/2))-3*arctanh(c/x)^2*polylog(2,-(1+c/x)/(1-c^2/x^2)^(1/2))+6*arctanh(c/x)*polylog(3,-(1+c/x)/(1-c^2/x^2)^(1/2))-6*polylog(4,-(1+c/x)/(1-c^2/x^2)^(1/2))-1/2*I*Pi*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))*(csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1))*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1))))-csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1))*csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))-csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))*csgn(I/(1-(1+c/x)^2/(c^2/x^2-1)))+csgn(I*(-(1+c/x)^2/(c^2/x^2-1)-1)/(1-(1+c/x)^2/(c^2/x^2-1)))^2)*arctanh(c/x)^3+3/2*arctanh(c/x)^2*polylog(2,-(1+c/x)^2/(1-c^2/x^2))-3/2*arctanh(c/x)*polylog(3,-(1+c/x)^2/(1-c^2/x^2))+3/4*polylog(4,-(1+c/x)^2/(1-c^2/x^2)))+3*a*b^2*(-ln(c/x)*arctanh(c/x)^2+arctanh(c/x)*polylog(2,-(1+c/x)^2/(1-c^2/x^2))-1/2*polylog(3,-(1+c/x)^2/(1-c^2/x^2))+arctanh(c/x)^2*ln((1+c/x)^2/(1-c^2/x^2)-1)-arctanh(c/x)^2*ln(1-(1+c/x)/(1-c^2/x^2)^(1/2))-2*arctanh(c/x)*polylog(2,(1+c/x)/(1-c^2/x^2)^(1/2))+2*polylog(3,(1+c/x)/(1-c^2/x^2)^(1/2))-arctanh(c/x)^2*ln(1+(1+c/x)/(1-c^2/x^2)^(1/2))-2*arctanh(c/x)*polylog(2,-(1+c/x)/(1-c^2/x^2)^(1/2))+2*polylog(3,-(1+c/x)/(1-c^2/x^2)^(1/2))-1/2*I*Pi*csgn(I*...

```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x}) + a)^3}{x} dx$$

input `integrate((a+b*arctanh(c/x))^3/x,x, algorithm="fricas")`

output `integral((b^3*arctanh(c/x)^3 + 3*a*b^2*arctanh(c/x)^2 + 3*a^2*b*arctanh(c/x) + a^3)/x, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x}))^3}{x} dx$$

input `integrate((a+b*atanh(c/x))**3/x,x)`

output `Integral((a + b*atanh(c/x))**3/x, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x}) + a)^3}{x} dx$$

input `integrate((a+b*arctanh(c/x))^3/x,x, algorithm="maxima")`

output `a^3*log(x) + integrate(1/8*b^3*(log(c/x + 1) - log(-c/x + 1))^3/x + 3/4*a*b^2*(log(c/x + 1) - log(-c/x + 1))^2/x + 3/2*a^2*b*(log(c/x + 1) - log(-c/x + 1))/x, x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x}) + a)^3}{x} dx$$

input `integrate((a+b*arctanh(c/x))^3/x,x, algorithm="giac")`

output `integrate((b*arctanh(c/x) + a)^3/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x}))^3}{x} dx$$

input `int((a + b*atanh(c/x))^3/x,x)`

output `int((a + b*atanh(c/x))^3/x, x)`

Reduce [F]

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x} dx &= 3 \left(\int \frac{\operatorname{atanh}(\frac{c}{x})}{x} dx \right) a^2 b + \left(\int \frac{\operatorname{atanh}(\frac{c}{x})^3}{x} dx \right) b^3 \\ &\quad + 3 \left(\int \frac{\operatorname{atanh}(\frac{c}{x})^2}{x} dx \right) a b^2 + \log(x) a^3 \end{aligned}$$

input `int((a+b*atanh(c/x))^3/x,x)`

output `3*int(atanh(c/x)/x,x)*a**2*b + int(atanh(c/x)**3/x,x)*b**3 + 3*int(atanh(c/x)**2/x,x)*a*b**2 + log(x)*a**3`

3.155 $\int \frac{\left(a+b\operatorname{arctanh}\left(\frac{c}{x}\right)\right)^3}{x^2} dx$

Optimal result	1271
Mathematica [A] (verified)	1272
Rubi [A] (verified)	1272
Maple [B] (verified)	1275
Fricas [F]	1276
Sympy [F]	1276
Maxima [F]	1276
Giac [F]	1277
Mupad [F(-1)]	1277
Reduce [F]	1277

Optimal result

Integrand size = 16, antiderivative size = 126

$$\int \frac{\left(a+b\operatorname{arctanh}\left(\frac{c}{x}\right)\right)^3}{x^2} dx = -\frac{\left(a+b\operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^3}{c} - \frac{\left(a+b\operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^3}{x} + \frac{3b\left(a+b\operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^2 \log\left(\frac{2}{1-\frac{c}{x}}\right)}{c} + \frac{3b^2\left(a+b\operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right) \operatorname{PolyLog}\left(2, 1-\frac{2}{1-\frac{c}{x}}\right)}{c} - \frac{3b^3 \operatorname{PolyLog}\left(3, 1-\frac{2}{1-\frac{c}{x}}\right)}{2c}$$

output

```
-(a+b*arccoth(x/c))^3/c-(a+b*arccoth(x/c))^3/x+3*b*(a+b*arccoth(x/c))^2*ln
(2/(1-c/x))/c+3*b^2*(a+b*arccoth(x/c))*polylog(2,1-2/(1-c/x))/c-3/2*b^3*po
lylog(3,1-2/(1-c/x))/c
```

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.63

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^2} dx = -\frac{a^3}{x} - \frac{3a^2 b \operatorname{arctanh}(\frac{c}{x})}{x} - \frac{3a^2 b \log\left(1 - \frac{c^2}{x^2}\right)}{2c}$$

$$- \frac{3ab^2 \left(\operatorname{arctanh}(\frac{c}{x}) \left(-\operatorname{arctanh}(\frac{c}{x}) + \frac{c \operatorname{arctanh}(\frac{c}{x})}{x} - 2 \log\left(1 + e^{-2 \operatorname{arctanh}(\frac{c}{x})}\right) \right) + \operatorname{PolyLog}\left(2, -e^{-2 \operatorname{arctanh}(\frac{c}{x})}\right) \right)}{c}$$

$$- \frac{b^3 \left(\operatorname{arctanh}(\frac{c}{x})^2 \left(-\operatorname{arctanh}(\frac{c}{x}) + \frac{c \operatorname{arctanh}(\frac{c}{x})}{x} - 3 \log\left(1 + e^{-2 \operatorname{arctanh}(\frac{c}{x})}\right) \right) + 3 \operatorname{arctanh}(\frac{c}{x}) \operatorname{PolyLog}\left(2, -e^{-2 \operatorname{arctanh}(\frac{c}{x})}\right) \right)}{c}$$

input `Integrate[(a + b*ArcTanh[c/x])^3/x^2, x]`

output `-(a^3/x) - (3*a^2*b*ArcTanh[c/x])/x - (3*a^2*b*Log[1 - c^2/x^2])/(2*c) - (3*a*b^2*(ArcTanh[c/x]*(-ArcTanh[c/x] + (c*ArcTanh[c/x])/x - 2*Log[1 + E^(-2*ArcTanh[c/x])]) + PolyLog[2, -E^(-2*ArcTanh[c/x])]))/c - (b^3*(ArcTanh[c/x]^2*(-ArcTanh[c/x] + (c*ArcTanh[c/x])/x - 3*Log[1 + E^(-2*ArcTanh[c/x])]) + 3*ArcTanh[c/x]*PolyLog[2, -E^(-2*ArcTanh[c/x])]) + (3*PolyLog[3, -E^(-2*ArcTanh[c/x])]))/2)/c`

Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.11, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6454, 6436, 6546, 6470, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^2} dx$$

↓ 6454

$$- \int \left(a + b \operatorname{arctanh}(\frac{c}{x}) \right)^3 d\frac{1}{x}$$

$$\begin{aligned}
 & \downarrow \text{6436} \\
 & 3bc \int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{(1 - \frac{c^2}{x^2})x} d\frac{1}{x} - \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x} \\
 & \downarrow \text{6546} \\
 & 3bc \left(\frac{\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{1 - \frac{c}{x}} d\frac{1}{x}}{c} - \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{3bc^2} \right) - \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x} \\
 & \downarrow \text{6470} \\
 & 3bc \left(\frac{\left(\frac{\log\left(\frac{2}{1 - \frac{c}{x}}\right)(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{c} - 2b \int \frac{(a + b \operatorname{arctanh}(\frac{c}{x})) \log\left(\frac{2}{1 - \frac{c}{x}}\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} \right)}{c} - \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{3bc^2} \right) - \\
 & \quad \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x} \\
 & \downarrow \text{6620} \\
 & 3bc \left(\frac{\left(\frac{\log\left(\frac{2}{1 - \frac{c}{x}}\right)(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{c} - 2b \left(\frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{c}{x}}\right)}{1 - \frac{c^2}{x^2}} d\frac{1}{x} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{c}{x}}\right)(a + b \operatorname{arctanh}(\frac{c}{x}))}{2c} \right) \right)}{c} - \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{3bc^2} \right) - \\
 & \quad \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x} \\
 & \downarrow \text{7164} \\
 & 3bc \left(\frac{\left(\frac{\log\left(\frac{2}{1 - \frac{c}{x}}\right)(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{c} - 2b \left(\frac{b \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - \frac{c}{x}}\right)}{4c} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{c}{x}}\right)(a + b \operatorname{arctanh}(\frac{c}{x}))}{2c} \right) \right)}{c} - \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{3bc^2} \right) - \\
 & \quad \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x}
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c/x])^3/x^2,x]`

output `-((a + b*ArcTanh[c/x])^3/x) + 3*b*c*(-1/3*(a + b*ArcTanh[c/x])^3/(b*c^2) + ((a + b*ArcTanh[c/x])^2*Log[2/(1 - c/x)]/c - 2*b*(-1/2*((a + b*ArcTanh[c/x])*PolyLog[2, 1 - 2/(1 - c/x)]/c + (b*PolyLog[3, 1 - 2/(1 - c/x)]/(4*c))))/c)`

Defintions of rubi rules used

rule 6436 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.), x_Symbol] := Simp[x*(a + b*ArcTanh[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`

rule 6454 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6470 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6546 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6620

```
Int[(Log[u_]*((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.))/((d_) + (e_.)*(x_)^
2), x_Symbol] := Simp[(- (a + b*ArcTanh[c*x])^p)*(PolyLog[2, 1 - u]/(2*c*d))
, x] + Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(
d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d
+ e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 264 vs. $2(124) = 248$.

Time = 2.18 (sec) , antiderivative size = 265, normalized size of antiderivative = 2.10

method	result
derivativedivides	$\frac{\frac{c}{x}a^3 + b^3}{c} \left(\operatorname{arctanh}\left(\frac{c}{x}\right)^3 \left(\frac{c}{x} - 1\right) + 2 \operatorname{arctanh}\left(\frac{c}{x}\right)^3 - 3 \operatorname{arctanh}\left(\frac{c}{x}\right)^2 \ln\left(1 + \frac{\left(\frac{1+c}{x}\right)^2}{1 - \frac{c^2}{x^2}}\right) - 3 \operatorname{arctanh}\left(\frac{c}{x}\right) \operatorname{polylog}\left(2, -\frac{1+c}{1-c}\right) \right)$
default	$\frac{\frac{c}{x}a^3 + b^3}{c} \left(\operatorname{arctanh}\left(\frac{c}{x}\right)^3 \left(\frac{c}{x} - 1\right) + 2 \operatorname{arctanh}\left(\frac{c}{x}\right)^3 - 3 \operatorname{arctanh}\left(\frac{c}{x}\right)^2 \ln\left(1 + \frac{\left(\frac{1+c}{x}\right)^2}{1 - \frac{c^2}{x^2}}\right) - 3 \operatorname{arctanh}\left(\frac{c}{x}\right) \operatorname{polylog}\left(2, -\frac{1+c}{1-c}\right) \right)$
parts	$-\frac{a^3}{x} - \frac{b^3}{c} \left(\operatorname{arctanh}\left(\frac{c}{x}\right)^3 \left(\frac{c}{x} - 1\right) + 2 \operatorname{arctanh}\left(\frac{c}{x}\right)^3 - 3 \operatorname{arctanh}\left(\frac{c}{x}\right)^2 \ln\left(1 + \frac{\left(\frac{1+c}{x}\right)^2}{1 - \frac{c^2}{x^2}}\right) - 3 \operatorname{arctanh}\left(\frac{c}{x}\right) \operatorname{polylog}\left(2, -\frac{1+c}{1-c}\right) \right)$

input

```
int((a+b*arctanh(c/x))^3/x^2,x,method=_RETURNVERBOSE)
```

output

```
-1/c*(c/x*a^3+b^3*(arctanh(c/x)^3*(c/x-1)+2*arctanh(c/x)^3-3*arctanh(c/x)^
2*ln(1+(1+c/x)^2/(1-c^2/x^2))-3*arctanh(c/x)*polylog(2,-(1+c/x)^2/(1-c^2/x
^2))+3/2*polylog(3,-(1+c/x)^2/(1-c^2/x^2)))+3*a*b^2*(arctanh(c/x)^2*(c/x-1
)+2*arctanh(c/x)^2-2*arctanh(c/x)*ln(1+(1+c/x)^2/(1-c^2/x^2))-polylog(2,-(
1+c/x)^2/(1-c^2/x^2)))+3*a^2*b*(c/x*arctanh(c/x)+1/2*ln(1-c^2/x^2)))
```


Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^2} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x}) + a)^3}{x^2} dx$$

input `integrate((a+b*arctanh(c/x))^3/x^2,x, algorithm="fricas")`

output `integral((b^3*arctanh(c/x)^3 + 3*a*b^2*arctanh(c/x)^2 + 3*a^2*b*arctanh(c/x) + a^3)/x^2, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^2} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x}))^3}{x^2} dx$$

input `integrate((a+b*atanh(c/x))**3/x**2,x)`

output `Integral((a + b*atanh(c/x))**3/x**2, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^2} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x}) + a)^3}{x^2} dx$$

input `integrate((a+b*arctanh(c/x))^3/x^2,x, algorithm="maxima")`

output `-3/2*a^2*b*(2*c*arctanh(c/x)/x + log(-c^2/x^2 + 1))/c - a^3/x + 1/8*((b^3*c - b^3*x)*log(-c + x)^3 - 3*(2*a*b^2*c + (b^3*c + b^3*x)*log(c + x))*log(-c + x)^2)/(c*x) - integrate(-1/8*((b^3*c^2 - b^3*c*x)*log(c + x)^3 + 6*(a*b^2*c^2 - a*b^2*c*x)*log(c + x)^2 - 3*(4*a*b^2*c*x + (b^3*c^2 - b^3*c*x)*log(c + x)^2 + 2*(2*a*b^2*c^2 + b^3*x^2 - (2*a*b^2*c - b^3*c)*x)*log(c + x))*log(-c + x))/(c^2*x^2 - c*x^3), x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^2} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x}) + a)^3}{x^2} dx$$

input `integrate((a+b*arctanh(c/x))^3/x^2,x, algorithm="giac")`

output `integrate((b*arctanh(c/x) + a)^3/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^2} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x}))^3}{x^2} dx$$

input `int((a + b*atanh(c/x))^3/x^2,x)`

output `int((a + b*atanh(c/x))^3/x^2, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^2} dx$$

$$= \frac{-\operatorname{atanh}(\frac{c}{x})^3 b^3 c - 3 \operatorname{atanh}(\frac{c}{x})^2 a b^2 c - 3 \operatorname{atanh}(\frac{c}{x}) a^2 b c + 3 \operatorname{atanh}(\frac{c}{x}) a^2 b x + 6 \left(\int \frac{\operatorname{atanh}(\frac{c}{x})}{c^2 x - x^3} dx \right) a b^2 c^2 x + 3}{c x}$$

input `int((a+b*atanh(c/x))^3/x^2,x)`

output

```
( - atanh(c/x)**3*b**3*c - 3*atanh(c/x)**2*a*b**2*c - 3*atanh(c/x)*a**2*b*  
c + 3*atanh(c/x)*a**2*b*x + 6*int(atanh(c/x)/(c**2*x - x**3),x)*a*b**2*c**  
2*x + 3*int(atanh(c/x)**2/(c**2*x - x**3),x)*b**3*c**2*x - 3*log( - c - x)  
*a**2*b*x + 3*log(x)*a**2*b*x - a**3*c)/(c*x)
```

3.156 $\int \frac{\left(a+b\operatorname{arctanh}\left(\frac{c}{x}\right)\right)^3}{x^3} dx$

Optimal result	1279
Mathematica [A] (verified)	1280
Rubi [A] (verified)	1280
Maple [C] (warning: unable to verify)	1284
Fricas [F]	1285
Sympy [F]	1285
Maxima [F]	1285
Giac [F]	1286
Mupad [F(-1)]	1287
Reduce [F]	1287

Optimal result

Integrand size = 16, antiderivative size = 139

$$\int \frac{\left(a+b\operatorname{arctanh}\left(\frac{c}{x}\right)\right)^3}{x^3} dx = -\frac{3b\left(a+b\operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^2}{2c^2} - \frac{3b\left(a+b\operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^2}{2cx} + \frac{\left(a+b\operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^3}{2c^2} - \frac{\left(a+b\operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)^3}{2x^2} + \frac{3b^2\left(a+b\operatorname{coth}^{-1}\left(\frac{x}{c}\right)\right)\log\left(\frac{2}{1-\frac{c}{x}}\right)}{c^2} + \frac{3b^3\operatorname{PolyLog}\left(2,1-\frac{2}{1-\frac{c}{x}}\right)}{2c^2}$$

output

```
-3/2*b*(a+b*arccoth(x/c))^2/c^2-3/2*b*(a+b*arccoth(x/c))^2/c/x+1/2*(a+b*arccoth(x/c))^3/c^2-1/2*(a+b*arccoth(x/c))^3/x^2+3*b^2*(a+b*arccoth(x/c))*ln(2/(1-c/x))/c^2+3/2*b^3*polylog(2,1-2/(1-c/x))/c^2
```

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.40

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^3} dx$$

$$= \frac{6b^2(-c+x)(bx+a(c+x))\operatorname{arctanh}(\frac{c}{x})^2 + 2b^3(-c^2+x^2)\operatorname{arctanh}(\frac{c}{x})^3 + 6b\operatorname{arctanh}(\frac{c}{x})(-ac(ac+2bx))}{x^3}$$

input

```
Integrate[(a + b*ArcTanh[c/x])^3/x^3,x]
```

output

```
(6*b^2*(-c+x)*(b*x+a*(c+x))*ArcTanh[c/x]^2 + 2*b^3*(-c^2+x^2)*ArcTanh[c/x]^3 + 6*b*ArcTanh[c/x]*(-(a*c*(a*c+2*b*x)) + 2*b^2*x^2*Log[1+E^(-2*ArcTanh[c/x])]) + a*(12*b^2*x^2*Log[1/Sqrt[1-c^2/x^2]] - a*(2*a*c^2 + 6*b*c*x + 3*b*x^2*Log[1-c/x] - 3*b*x^2*Log[(c+x)/x])) - 6*b^3*x^2*PolyLog[2, -E^(-2*ArcTanh[c/x])])/(4*c^2*x^2)
```

Rubi [A] (verified)

Time = 1.13 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.11, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.562$, Rules used = {6454, 6452, 6542, 6436, 6510, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^3} dx$$

$$\downarrow 6454$$

$$- \int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x} d\frac{1}{x}$$

$$\downarrow 6452$$

$$\frac{3}{2}bc \int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^2}{(1 - \frac{c^2}{x^2})x^2} d\frac{1}{x} - \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{2x^2}$$

$$\begin{array}{c}
 \downarrow 6542 \\
 \frac{3}{2}bc \left(\frac{\int \frac{(a+b\operatorname{arctanh}(\frac{c}{x}))^2}{1-\frac{c^2}{x^2}} d\frac{1}{x}}{c^2} - \frac{\int (a+b\operatorname{arctanh}(\frac{c}{x}))^2 d\frac{1}{x}}{c^2} \right) - \frac{(a+b\operatorname{arctanh}(\frac{c}{x}))^3}{2x^2} \\
 \downarrow 6436 \\
 \frac{3}{2}bc \left(\frac{\int \frac{(a+b\operatorname{arctanh}(\frac{c}{x}))^2}{1-\frac{c^2}{x^2}} d\frac{1}{x}}{c^2} - \frac{\frac{(a+b\operatorname{arctanh}(\frac{c}{x}))^2}{x} - 2bc \int \frac{a+b\operatorname{arctanh}(\frac{c}{x})}{(1-\frac{c^2}{x^2})x} d\frac{1}{x}}{c^2} \right) - \\
 \frac{(a+b\operatorname{arctanh}(\frac{c}{x}))^3}{2x^2} \\
 \downarrow 6510 \\
 \frac{3}{2}bc \left(\frac{(a+b\operatorname{arctanh}(\frac{c}{x}))^3}{3bc^3} - \frac{\frac{(a+b\operatorname{arctanh}(\frac{c}{x}))^2}{x} - 2bc \int \frac{a+b\operatorname{arctanh}(\frac{c}{x})}{(1-\frac{c^2}{x^2})x} d\frac{1}{x}}{c^2} \right) - \\
 \frac{(a+b\operatorname{arctanh}(\frac{c}{x}))^3}{2x^2} \\
 \downarrow 6546 \\
 \frac{3}{2}bc \left(\frac{(a+b\operatorname{arctanh}(\frac{c}{x}))^3}{3bc^3} - \frac{\frac{(a+b\operatorname{arctanh}(\frac{c}{x}))^2}{x} - 2bc \left(\frac{\int \frac{a+b\operatorname{arctanh}(\frac{c}{x})}{1-\frac{c}{x}} d\frac{1}{x}}{c} - \frac{(a+b\operatorname{arctanh}(\frac{c}{x}))^2}{2bc^2} \right)}{c^2} \right) - \\
 \frac{(a+b\operatorname{arctanh}(\frac{c}{x}))^3}{2x^2} \\
 \downarrow 6470
 \end{array}$$

$$\frac{3}{2}bc \left(\frac{(a + \operatorname{arctanh}(\frac{c}{x}))^3}{3bc^3} - \frac{\frac{(a + \operatorname{arctanh}(\frac{c}{x}))^2}{x} - 2bc \left(\frac{\log(\frac{2}{1-\frac{c}{x}})(a + \operatorname{arctanh}(\frac{c}{x}))}{c} - b \int \frac{\log(\frac{2}{1-\frac{c}{x}})}{1-\frac{c^2}{x^2}} d\frac{1}{x} \right)}{c^2} - \frac{(a + \operatorname{arctanh}(\frac{c}{x}))}{2bc^2} \right)$$

$$\frac{(a + \operatorname{arctanh}(\frac{c}{x}))^3}{2x^2}$$

↓ 2849

$$\frac{3}{2}bc \left(\frac{(a + \operatorname{arctanh}(\frac{c}{x}))^3}{3bc^3} - \frac{\frac{(a + \operatorname{arctanh}(\frac{c}{x}))^2}{x} - 2bc \left(\frac{b \int \frac{\log(\frac{2}{1-\frac{c}{x}})}{1-\frac{2}{1-\frac{c}{x}}} d\frac{1}{x}}{c} + \frac{\log(\frac{2}{1-\frac{c}{x}})(a + \operatorname{arctanh}(\frac{c}{x}))}{c} \right)}{c^2} - \frac{(a + \operatorname{arctanh}(\frac{c}{x}))}{2bc^2} \right)$$

$$\frac{(a + \operatorname{arctanh}(\frac{c}{x}))^3}{2x^2}$$

↓ 2752

$$\frac{3}{2}bc \left(\frac{(a + \operatorname{arctanh}(\frac{c}{x}))^3}{3bc^3} - \frac{\frac{(a + \operatorname{arctanh}(\frac{c}{x}))^2}{x} - 2bc \left(\frac{\log(\frac{2}{1-\frac{c}{x}})(a + \operatorname{arctanh}(\frac{c}{x}))}{c} + \frac{b \operatorname{PolyLog}(2, 1 - \frac{2}{1-\frac{c}{x}})}{2c} \right)}{c^2} - \frac{(a + \operatorname{arctanh}(\frac{c}{x}))}{2bc^2} \right)$$

$$\frac{(a + \operatorname{arctanh}(\frac{c}{x}))^3}{2x^2}$$

input `Int[(a + b*ArcTanh[c/x])^3/x^3, x]`

output

```
-1/2*(a + b*ArcTanh[c/x])^3/x^2 + (3*b*c*((a + b*ArcTanh[c/x])^3/(3*b*c^3)
- ((a + b*ArcTanh[c/x])^2/x - 2*b*c*(-1/2*(a + b*ArcTanh[c/x])^2/(b*c^2)
+ (((a + b*ArcTanh[c/x])*Log[2/(1 - c/x)]/c + (b*PolyLog[2, 1 - 2/(1 - c/
x]))/(2*c))/c))/c^2))/2
```

Defintions of rubi rules used

rule 2752

```
Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLo
g[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]
```

rule 2849

```
Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp
[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[
{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]
```

rule 6436

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^p, x_Symbol] := Simp[x*(a
+ b*ArcTanh[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTanh[c*x^n])
^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]
&& (EqQ[n, 1] || EqQ[p, 1])
```

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^p*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

rule 6454

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^p*(x_)^(m_.), x_Symbol] :=
Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x
], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simpl
ify[(m + 1)/n]
```


rule 6470

```
Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol]
  := Simp[(-a + b*ArcTanh[c*x])^p*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c
  *(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^
  2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]
```

rule 6510

```
Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol]
  := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x]
  && EqQ[c^2*d + e, 0] && NeQ[p, -1]
```

rule 6542

```
Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_))/((d_) + (e_.)*(x_)^2), x_Symbol]
  := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]
```

rule 6546

```
Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*(x_)/((d_) + (e_.)*(x_)^2), x_Symbol]
  := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 22.87 (sec) , antiderivative size = 6081, normalized size of antiderivative = 43.75

method	result	size
derivativeldivides	Expression too large to display	6081
default	Expression too large to display	6081
parts	Expression too large to display	6083

input

```
int((a+b*arctanh(c/x))^3/x^3,x,method=_RETURNVERBOSE)
```

output result too large to display

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^3} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x}) + a)^3}{x^3} dx$$

input `integrate((a+b*arctanh(c/x))^3/x^3,x, algorithm="fricas")`

output `integral((b^3*arctanh(c/x)^3 + 3*a*b^2*arctanh(c/x)^2 + 3*a^2*b*arctanh(c/x) + a^3)/x^3, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^3} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x}))^3}{x^3} dx$$

input `integrate((a+b*atanh(c/x))**3/x**3,x)`

output `Integral((a + b*atanh(c/x))**3/x**3, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^3} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x}) + a)^3}{x^3} dx$$

input `integrate((a+b*arctanh(c/x))^3/x^3,x, algorithm="maxima")`

output

```

3/4*(c*(log(c + x)/c^3 - log(-c + x)/c^3 - 2/(c^2*x)) - 2*arctanh(c/x)/x^2
)*a^2*b - 3/8*(c^2*((log(c + x)^2 - 2*(log(c + x) - 2)*log(-c + x) + log(-
c + x)^2 + 4*log(c + x))/c^4 - 8*log(x)/c^4) - 4*c*(log(c + x)/c^3 - log(-
c + x)/c^3 - 2/(c^2*x))*arctanh(c/x))*a*b^2 + 1/64*(32*c^4*integrate(-1/4*
log(x)^3/(c^4*x^3 - c^2*x^5), x) - 3*c^3*(log(c + x)/c^5 - log(-c + x)/c^5
- 2/(c^4*x)) + 48*c^3*integrate(-1/4*x*log(x)^2/(c^4*x^3 - c^2*x^5), x) +
48*c^3*integrate(-1/4*x*log(x)/(c^4*x^3 - c^2*x^5), x) - 6*c*(2*log(-c +
x)/c^3 - 2*log(x)/c^3 + (c + 2*x)/(c^2*x^2))*log(-c/x + 1)^2 + 21*c^2*(log
(c + x)/c^4 + log(-c + x)/c^4 - 2*log(x)/c^4) - 32*c^2*integrate(-1/4*x^2*
log(x)^3/(c^4*x^3 - c^2*x^5), x) + 48*c^2*integrate(-1/4*x^2*log(x)^2/(c^4
*x^3 - c^2*x^5), x) - 384*c^2*integrate(-1/4*x^2*log(c + x)/(c^4*x^3 - c^2
*x^5), x) + 144*c^2*integrate(-1/4*x^2*log(x)/(c^4*x^3 - c^2*x^5), x) - 18
*c*(log(c + x)/c^3 - log(-c + x)/c^3) + c*(6*(2*x^2*log(-c + x)^2 + 2*x^2*
log(x)^2 - 6*x^2*log(x) + c^2 + 6*c*x - 2*(2*x^2*log(x) - 3*x^2)*log(-c +
x))*log(-c/x + 1)/(c^3*x^2) - (4*x^2*log(-c + x)^3 - 4*x^2*log(x)^3 + 18*x
^2*log(x)^2 - 6*(2*x^2*log(x) - 3*x^2)*log(-c + x)^2 - 42*x^2*log(x) + 3*c
^2 + 42*c*x + 6*(2*x^2*log(x)^2 - 6*x^2*log(x) + 7*x^2)*log(-c + x))/(c^3*
x^2)) - 48*c*integrate(-1/4*x^3*log(x)^2/(c^4*x^3 - c^2*x^5), x) - 192*c*i
ntegrate(-1/4*x^3*log(c + x)/(c^4*x^3 - c^2*x^5), x) + 336*c*integrate(-1/
4*x^3*log(x)/(c^4*x^3 - c^2*x^5), x) + 4*log(-c/x + 1)^3/x^2 - 2*(12*c*...

```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^3} dx = \int \frac{(b \operatorname{arctanh}(\frac{c}{x}) + a)^3}{x^3} dx$$

input

```
integrate((a+b*arctanh(c/x))^3/x^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c/x) + a)^3/x^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^3} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x}))^3}{x^3} dx$$

input `int((a + b*atanh(c/x))^3/x^3,x)`output `int((a + b*atanh(c/x))^3/x^3, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x}))^3}{x^3} dx$$

$$= \frac{-\operatorname{atanh}(\frac{c}{x})^3 b^3 c^2 + \operatorname{atanh}(\frac{c}{x})^3 b^3 x^2 - 3 \operatorname{atanh}(\frac{c}{x})^2 a b^2 c^2 + 3 \operatorname{atanh}(\frac{c}{x})^2 a b^2 x^2 - 3 \operatorname{atanh}(\frac{c}{x})^2 b^3 c x - 3 \operatorname{atanh}(\frac{c}{x})^2 a b^2 c x + 3 \operatorname{atanh}(\frac{c}{x})^2 a b^2 x^2 - 3 \operatorname{atanh}(\frac{c}{x})^2 b^3 c x - 3 \operatorname{atanh}(\frac{c}{x})^2 a b^2 c x + 3 \operatorname{atanh}(\frac{c}{x})^2 a b^2 x^2 - 6 \operatorname{atanh}(\frac{c}{x}) a b^2 c x + 6 \operatorname{atanh}(\frac{c}{x}) a b^2 x^2 + 3 \operatorname{atanh}(\frac{c}{x}) b^3 c^2 - 3 \operatorname{atanh}(\frac{c}{x}) b^3 x^2 + 2 + 6 \int (\operatorname{atanh}(\frac{c}{x}) / (c^2 x^3 - x^5), x) b^3 c^4 x^2 - 6 \log(-c - x) a b^2 x^2 + 6 \log(x) a b^2 x^2 - a^3 c^2 - 3 a^2 b c x + 3 b^3 c x}{(2 c^2 x^2)}$$

input `int((a+b*atanh(c/x))^3/x^3,x)`output `(- atanh(c/x)**3*b**3*c**2 + atanh(c/x)**3*b**3*x**2 - 3*atanh(c/x)**2*a*b**2*c**2 + 3*atanh(c/x)**2*a*b**2*x**2 - 3*atanh(c/x)**2*b**3*c*x - 3*atanh(c/x)**2*a*b**2*c*x + 3*atanh(c/x)**2*a*b**2*x**2 - 6*atanh(c/x)*a*b**2*c*x + 6*atanh(c/x)*a*b**2*x**2 + 3*atanh(c/x)*b**3*c**2 - 3*atanh(c/x)*b**3*x**2 + 2 + 6*int(atanh(c/x)/(c**2*x**3 - x**5),x)*b**3*c**4*x**2 - 6*log(-c - x)*a*b**2*x**2 + 6*log(x)*a*b**2*x**2 - a**3*c**2 - 3*a**2*b*c*x + 3*b**3*c*x)/(2*c**2*x**2)`

3.157 $\int x^7 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx$

Optimal result	1288
Mathematica [A] (verified)	1288
Rubi [A] (verified)	1289
Maple [A] (verified)	1291
Fricas [A] (verification not implemented)	1292
Sympy [A] (verification not implemented)	1292
Maxima [A] (verification not implemented)	1292
Giac [A] (verification not implemented)	1293
Mupad [B] (verification not implemented)	1293
Reduce [B] (verification not implemented)	1294

Optimal result

Integrand size = 14, antiderivative size = 54

$$\int x^7 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{1}{8} b c^3 x^2 + \frac{1}{24} b c x^6 + \frac{1}{8} x^8 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) - \frac{1}{8} b c^4 \operatorname{arctanh} \left(\frac{x^2}{c} \right)$$

output

```
1/8*b*c^3*x^2+1/24*b*c*x^6+1/8*x^8*(a+b*arctanh(c/x^2))-1/8*b*c^4*arctanh(x^2/c)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.35

$$\int x^7 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{1}{8} b c^3 x^2 + \frac{1}{24} b c x^6 + \frac{a x^8}{8} + \frac{1}{8} b x^8 \operatorname{arctanh} \left(\frac{c}{x^2} \right) + \frac{1}{16} b c^4 \log(-c + x^2) - \frac{1}{16} b c^4 \log(c + x^2)$$

input

```
Integrate[x^7*(a + b*ArcTanh[c/x^2]),x]
```

output

$$(b*c^3*x^2)/8 + (b*c*x^6)/24 + (a*x^8)/8 + (b*x^8*ArcTanh[c/x^2])/8 + (b*c^4*Log[-c + x^2])/16 - (b*c^4*Log[c + x^2])/16$$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.96, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {6452, 795, 807, 25, 254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^7 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx \\ & \quad \downarrow \text{6452} \\ & \frac{1}{4} bc \int \frac{x^5}{1 - \frac{c^2}{x^4}} dx + \frac{1}{8} x^8 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) \\ & \quad \downarrow \text{795} \\ & \frac{1}{4} bc \int \frac{x^9}{x^4 - c^2} dx + \frac{1}{8} x^8 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) \\ & \quad \downarrow \text{807} \\ & \frac{1}{8} bc \int -\frac{x^8}{c^2 - x^4} dx^2 + \frac{1}{8} x^8 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) \\ & \quad \downarrow \text{25} \\ & \frac{1}{8} x^8 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) - \frac{1}{8} bc \int \frac{x^8}{c^2 - x^4} dx^2 \\ & \quad \downarrow \text{254} \\ & \frac{1}{8} x^8 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) - \frac{1}{8} bc \int \left(\frac{c^4}{c^2 - x^4} - c^2 - x^4 \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{8} x^8 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) + \frac{1}{8} bc \left(-c^3 \operatorname{arctanh} \left(\frac{x^2}{c} \right) + c^2 x^2 + \frac{x^6}{3} \right) \end{aligned}$$

input `Int[x^7*(a + b*ArcTanh[c/x^2]),x]`

output `(x^8*(a + b*ArcTanh[c/x^2]))/8 + (b*c*(c^2*x^2 + x^6/3 - c^3*ArcTanh[x^2/c]))/8`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 254 `Int[(x_)^(m_)/((a_) + (b_)*(x_)^2), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^2, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 3]`

rule 795 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Int[x^(m + n*p)*(b + a/x^n)^p, x] /; FreeQ[{a, b, m, n}, x] && IntegerQ[p] && NegQ[n]`

rule 807 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.89

method	result
parallelrisc	$\frac{x^8 \operatorname{arctanh}\left(\frac{c}{x^2}\right)b}{8} + \frac{x^8 a}{8} + \frac{bcx^6}{24} + \frac{bc^3x^2}{8} - \frac{\operatorname{arctanh}\left(\frac{c}{x^2}\right)bc^4}{8}$
parts	$\frac{x^8 a}{8} + b \left(\frac{x^8 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{8} - \frac{c \left(-\frac{x^6}{6} - \frac{c^2 x^2}{2} + \frac{c^3 \ln\left(1 + \frac{c}{x^2}\right)}{4} - \frac{c^3 \ln\left(\frac{c}{x^2} - 1\right)}{4} \right)}{4} \right)$
derivativedivides	$\frac{x^8 a}{8} - b \left(-\frac{x^8 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{8} + \frac{c \left(-\frac{x^6}{6} - \frac{c^2 x^2}{2} + \frac{c^3 \ln\left(1 + \frac{c}{x^2}\right)}{4} - \frac{c^3 \ln\left(\frac{c}{x^2} - 1\right)}{4} \right)}{4} \right)$
default	$\frac{x^8 a}{8} - b \left(-\frac{x^8 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{8} + \frac{c \left(-\frac{x^6}{6} - \frac{c^2 x^2}{2} + \frac{c^3 \ln\left(1 + \frac{c}{x^2}\right)}{4} - \frac{c^3 \ln\left(\frac{c}{x^2} - 1\right)}{4} \right)}{4} \right)$
orering	$-\frac{(-13x^8 - 14c^2x^4 + 27c^4)(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right))}{48} + \frac{(x^4 + 3c^2)(x^2 + c)(-x^2 + c) \left(7x^6(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)) - \frac{2x^4bc}{1 - \frac{c^2}{x^4}} \right)}{48x^6}$
risc	$\frac{x^8 b \ln(x^2 + c)}{16} - \frac{x^8 b \ln(-x^2 + c)}{16} + \frac{i\pi b x^8 \operatorname{csgn}\left(\frac{i(-x^2 + c)}{x^2}\right)^2}{16} - \frac{i\pi b x^8 \operatorname{csgn}(i(-x^2 + c)) \operatorname{csgn}\left(\frac{i(-x^2 + c)}{x^2}\right)^2}{32} + \dots$

input `int(x^7*(a+b*arctanh(c/x^2)),x,method=_RETURNVERBOSE)`

output `1/8*x^8*arctanh(c/x^2)*b+1/8*x^8*a+1/24*b*c*x^6+1/8*b*c^3*x^2-1/8*arctanh(c/x^2)*b*c^4`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.98

$$\int x^7 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{1}{8} ax^8 + \frac{1}{24} bcx^6 + \frac{1}{8} bc^3x^2 + \frac{1}{16} (bx^8 - bc^4) \log \left(\frac{x^2 + c}{x^2 - c} \right)$$

input `integrate(x^7*(a+b*arctanh(c/x^2)),x, algorithm="fricas")`

output `1/8*a*x^8 + 1/24*b*c*x^6 + 1/8*b*c^3*x^2 + 1/16*(b*x^8 - b*c^4)*log((x^2 + c)/(x^2 - c))`

Sympy [A] (verification not implemented)

Time = 3.16 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.94

$$\int x^7 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{ax^8}{8} - \frac{bc^4 \operatorname{atanh} \left(\frac{c}{x^2} \right)}{8} + \frac{bc^3x^2}{8} + \frac{bcx^6}{24} + \frac{bx^8 \operatorname{atanh} \left(\frac{c}{x^2} \right)}{8}$$

input `integrate(x**7*(a+b*atanh(c/x**2)),x)`

output `a*x**8/8 - b*c**4*atanh(c/x**2)/8 + b*c**3*x**2/8 + b*c*x**6/24 + b*x**8*atanh(c/x**2)/8`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.15

$$\begin{aligned} \int x^7 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx \\ = \frac{1}{8} ax^8 \\ + \frac{1}{48} \left(6x^8 \operatorname{artanh} \left(\frac{c}{x^2} \right) + (2x^6 + 6c^2x^2 - 3c^3 \log(x^2 + c) + 3c^3 \log(x^2 - c))c \right) b \end{aligned}$$

input `integrate(x^7*(a+b*arctanh(c/x^2)),x, algorithm="maxima")`

output $1/8*a*x^8 + 1/48*(6*x^8*\operatorname{arctanh}(c/x^2) + (2*x^6 + 6*c^2*x^2 - 3*c^3*\log(x^2 + c) + 3*c^3*\log(x^2 - c))*c)*b$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.31

$$\int x^7 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{1}{16} b x^8 \log \left(\frac{x^2 + c}{x^2 - c} \right) + \frac{1}{8} a x^8 + \frac{1}{24} b c x^6 + \frac{1}{8} b c^3 x^2 - \frac{1}{16} b c^4 \log (x^2 + c) + \frac{1}{16} b c^4 \log (-x^2 + c)$$

input `integrate(x^7*(a+b*arctanh(c/x^2)),x, algorithm="giac")`

output $1/16*b*x^8*\log((x^2 + c)/(x^2 - c)) + 1/8*a*x^8 + 1/24*b*c*x^6 + 1/8*b*c^3*x^2 - 1/16*b*c^4*\log(x^2 + c) + 1/16*b*c^4*\log(-x^2 + c)$

Mupad [B] (verification not implemented)

Time = 4.35 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.22

$$\int x^7 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{a x^8}{8} + \frac{b c^3 x^2}{8} + \frac{b x^8 \ln (x^2 + c)}{16} + \frac{b c x^6}{24} - \frac{b x^8 \ln (x^2 - c)}{16} + \frac{b c^4 \operatorname{atan} \left(\frac{x^2 i}{c} \right) i}{8}$$

input `int(x^7*(a + b*atanh(c/x^2)),x)`

output $(a*x^8)/8 + (b*c^3*x^2)/8 + (b*x^8*\log(c + x^2))/16 + (b*c^4*\operatorname{atan}((x^2*i)/c)*i)/8 + (b*c*x^6)/24 - (b*x^8*\log(x^2 - c))/16$

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.87

$$\int x^7 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = -\frac{\operatorname{atanh} \left(\frac{c}{x^2} \right) b c^4}{8} + \frac{\operatorname{atanh} \left(\frac{c}{x^2} \right) b x^8}{8} + \frac{a x^8}{8} + \frac{b c^3 x^2}{8} + \frac{b c x^6}{24}$$

input `int(x^7*(a+b*atanh(c/x^2)),x)`

output `(- 3*atanh(c/x**2)*b*c**4 + 3*atanh(c/x**2)*b*x**8 + 3*a*x**8 + 3*b*c**3*x**2 + b*c*x**6)/24`

3.158 $\int x^5 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx$

Optimal result	1295
Mathematica [A] (verified)	1295
Rubi [A] (verified)	1296
Maple [A] (verified)	1298
Fricas [A] (verification not implemented)	1298
Sympy [B] (verification not implemented)	1299
Maxima [A] (verification not implemented)	1299
Giac [A] (verification not implemented)	1300
Mupad [B] (verification not implemented)	1300
Reduce [B] (verification not implemented)	1301

Optimal result

Integrand size = 14, antiderivative size = 45

$$\int x^5 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{1}{12} b c x^4 + \frac{1}{6} x^6 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) + \frac{1}{12} b c^3 \log (c^2 - x^4)$$

output $1/12*b*c*x^4+1/6*x^6*(a+b*\operatorname{arctanh}(c/x^2))+1/12*b*c^3*\ln(-x^4+c^2)$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.11

$$\int x^5 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{1}{12} b c x^4 + \frac{a x^6}{6} + \frac{1}{6} b x^6 \operatorname{arctanh} \left(\frac{c}{x^2} \right) + \frac{1}{12} b c^3 \log (-c^2 + x^4)$$

input $\operatorname{Integrate}[x^5*(a + b*\operatorname{ArcTanh}[c/x^2]),x]$

output $(b*c*x^4)/12 + (a*x^6)/6 + (b*x^6*\operatorname{ArcTanh}[c/x^2])/6 + (b*c^3*\operatorname{Log}[-c^2 + x^4])/12$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.93, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {6452, 795, 798, 25, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^5 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) dx \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{3} bc \int \frac{x^3}{1 - \frac{c^2}{x^4}} dx + \frac{1}{6} x^6 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) \\
 & \quad \downarrow \text{795} \\
 & \frac{1}{3} bc \int \frac{x^7}{x^4 - c^2} dx + \frac{1}{6} x^6 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) \\
 & \quad \downarrow \text{798} \\
 & \frac{1}{12} bc \int -\frac{x^4}{c^2 - x^4} dx^4 + \frac{1}{6} x^6 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{6} x^6 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) - \frac{1}{12} bc \int \frac{x^4}{c^2 - x^4} dx^4 \\
 & \quad \downarrow \text{49} \\
 & \frac{1}{6} x^6 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) - \frac{1}{12} bc \int \left(\frac{c^2}{c^2 - x^4} - 1 \right) dx^4 \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{6} x^6 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) + \frac{1}{12} bc (c^2 \log (c^2 - x^4) + x^4)
 \end{aligned}$$

input `Int [x^5*(a + b*ArcTanh[c/x^2]), x]`

output `(x^6*(a + b*ArcTanh[c/x^2]))/6 + (b*c*(x^4 + c^2*Log[c^2 - x^4]))/12`

Defintions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 49 $\text{Int}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.)^{(\text{m}_.)} * ((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{(\text{n}_.)}), \text{x_Symbol}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(\text{a} + \text{b} * \text{x})^{\text{m}} * (\text{c} + \text{d} * \text{x})^{\text{n}}, \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{IGtQ}[\text{m}, 0] \&\& \text{IGtQ}[\text{m} + \text{n} + 2, 0]$
- rule 795 $\text{Int}[(\text{x}_.)^{(\text{m}_.)} * ((\text{a}_.) + (\text{b}_.) * (\text{x}_.)^{(\text{n}_.)})^{(\text{p}_.)}, \text{x_Symbol}] \rightarrow \text{Int}[\text{x}^{(\text{m} + \text{n} * \text{p})} * (\text{b} + \text{a} / \text{x}^{\text{n}})^{\text{p}}, \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{m}, \text{n}\}, \text{x}] \&\& \text{IntegerQ}[\text{p}] \&\& \text{NegQ}[\text{n}]$
- rule 798 $\text{Int}[(\text{x}_.)^{(\text{m}_.)} * ((\text{a}_.) + (\text{b}_.) * (\text{x}_.)^{(\text{n}_.)})^{(\text{p}_.)}, \text{x_Symbol}] \rightarrow \text{Simp}[1 / \text{n} \quad \text{Subst}[\text{Int}[\text{x}^{(\text{Simplify}[(\text{m} + 1) / \text{n}] - 1) * (\text{a} + \text{b} * \text{x})^{\text{p}}, \text{x}], \text{x}, \text{x}^{\text{n}}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{m}, \text{n}, \text{p}\}, \text{x}] \&\& \text{IntegerQ}[\text{Simplify}[(\text{m} + 1) / \text{n}]]$
- rule 2009 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Simp}[\text{IntSum}[\text{u}, \text{x}], \text{x}] \text{ ; SumQ}[\text{u}]$
- rule 6452 $\text{Int}[(\text{a}_.) + \text{ArcTanh}[(\text{c}_.) * (\text{x}_.)^{(\text{n}_.)}] * (\text{b}_.)^{(\text{p}_.)} * (\text{x}_.)^{(\text{m}_.)}, \text{x_Symbol}] \rightarrow \text{Simp}[\text{x}^{(\text{m} + 1)} * ((\text{a} + \text{b} * \text{ArcTanh}[\text{c} * \text{x}^{\text{n}}])^{\text{p}} / (\text{m} + 1)), \text{x}] - \text{Simp}[\text{b} * \text{c} * \text{n} * (\text{p} / (\text{m} + 1)) \quad \text{Int}[\text{x}^{(\text{m} + \text{n})} * ((\text{a} + \text{b} * \text{ArcTanh}[\text{c} * \text{x}^{\text{n}}])^{(\text{p} - 1)} / (1 - \text{c}^2 * \text{x}^{(2 * \text{n})})), \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{m}, \text{n}\}, \text{x}] \&\& \text{IGtQ}[\text{p}, 0] \&\& (\text{EqQ}[\text{p}, 1] \text{ || } (\text{EqQ}[\text{n}, 1] \&\& \text{IntegerQ}[\text{m}])) \&\& \text{NeQ}[\text{m}, -1]$

Maple [A] (verified)

Time = 0.73 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.31

method	result
parallelrisc	$\frac{x^6 \operatorname{arctanh}\left(\frac{c}{x^2}\right)b}{6} + \frac{ax^6}{6} + \frac{bcx^4}{12} + \frac{\ln(x^2-c)bc^3}{6} + \frac{\operatorname{arctanh}\left(\frac{c}{x^2}\right)bc^3}{6} + \frac{bc^3}{12}$
parts	$\frac{ax^6}{6} + b \left(\frac{x^6 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{6} - \frac{c \left(-\frac{c^2 \ln\left(\frac{c}{x^2}-1\right)}{4} - \frac{c^2 \ln\left(1+\frac{c}{x^2}\right)}{4} - \frac{x^4}{4} + c^2 \ln\left(\frac{1}{x}\right) \right)}{3} \right)$
derivativedivides	$\frac{ax^6}{6} - b \left(-\frac{x^6 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{6} + \frac{c \left(-\frac{c^2 \ln\left(\frac{c}{x^2}-1\right)}{4} - \frac{c^2 \ln\left(1+\frac{c}{x^2}\right)}{4} - \frac{x^4}{4} + c^2 \ln\left(\frac{1}{x}\right) \right)}{3} \right)$
default	$\frac{ax^6}{6} - b \left(-\frac{x^6 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{6} + \frac{c \left(-\frac{c^2 \ln\left(\frac{c}{x^2}-1\right)}{4} - \frac{c^2 \ln\left(1+\frac{c}{x^2}\right)}{4} - \frac{x^4}{4} + c^2 \ln\left(\frac{1}{x}\right) \right)}{3} \right)$
risch	$\frac{bx^6 \ln(x^2+c)}{12} - \frac{bx^6 \ln(-x^2+c)}{12} - \frac{i\pi b x^6 \operatorname{csgn}\left(\frac{i(-x^2+c)}{x^2}\right)^3}{24} - \frac{i\pi b x^6}{12} + \frac{i\pi b x^6 \operatorname{csgn}\left(\frac{i}{x^2}\right) \operatorname{csgn}\left(\frac{i(x^2+c)}{x^2}\right)^2}{24}$

input `int(x^5*(a+b*arctanh(c/x^2)),x,method=_RETURNVERBOSE)`output `1/6*x^6*arctanh(c/x^2)*b+1/6*a*x^6+1/12*b*c*x^4+1/6*ln(x^2-c)*b*c^3+1/6*arctanh(c/x^2)*b*c^3+1/12*b*c^3`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.16

$$\int x^5 \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right) \right) dx = \frac{1}{12} bx^6 \log\left(\frac{x^2+c}{x^2-c}\right) + \frac{1}{6} ax^6 + \frac{1}{12} bcx^4 + \frac{1}{12} bc^3 \log(x^4 - c^2)$$

input `integrate(x^5*(a+b*arctanh(c/x^2)),x, algorithm="fricas")`

output

```
1/12*b*x^6*log((x^2 + c)/(x^2 - c)) + 1/6*a*x^6 + 1/12*b*c*x^4 + 1/12*b*c^3*log(x^4 - c^2)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 75 vs. $2(37) = 74$.

Time = 2.34 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.67

$$\int x^5 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{ax^6}{6} + \frac{bc^3 \log(x - \sqrt{-c})}{6} + \frac{bc^3 \log(x + \sqrt{-c})}{6} - \frac{bc^3 \operatorname{atanh} \left(\frac{c}{x^2} \right)}{6} + \frac{bcx^4}{12} + \frac{bx^6 \operatorname{atanh} \left(\frac{c}{x^2} \right)}{6}$$

input

```
integrate(x**5*(a+b*atanh(c/x**2)),x)
```

output

```
a*x**6/6 + b*c**3*log(x - sqrt(-c))/6 + b*c**3*log(x + sqrt(-c))/6 - b*c**3*atanh(c/x**2)/6 + b*c*x**4/12 + b*x**6*atanh(c/x**2)/6
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.93

$$\int x^5 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{1}{6} ax^6 + \frac{1}{12} \left(2x^6 \operatorname{artanh} \left(\frac{c}{x^2} \right) + (x^4 + c^2 \log(x^4 - c^2))c \right) b$$

input

```
integrate(x^5*(a+b*arctanh(c/x^2)),x, algorithm="maxima")
```

output

```
1/6*a*x^6 + 1/12*(2*x^6*arctanh(c/x^2) + (x^4 + c^2*log(x^4 - c^2))*c)*b
```


Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.16

$$\int x^5 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{1}{12} b x^6 \log \left(\frac{x^2 + c}{x^2 - c} \right) + \frac{1}{6} a x^6 \\ + \frac{1}{12} b c x^4 + \frac{1}{12} b c^3 \log (x^4 - c^2)$$

input `integrate(x^5*(a+b*arctanh(c/x^2)),x, algorithm="giac")`

output `1/12*b*x^6*log((x^2 + c)/(x^2 - c)) + 1/6*a*x^6 + 1/12*b*c*x^4 + 1/12*b*c^3*log(x^4 - c^2)`

Mupad [B] (verification not implemented)

Time = 4.00 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.24

$$\int x^5 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{a x^6}{6} + \frac{b c^3 \ln (x^4 - c^2)}{12} + \frac{b x^6 \ln (x^2 + c)}{12} \\ + \frac{b c x^4}{12} - \frac{b x^6 \ln (x^2 - c)}{12}$$

input `int(x^5*(a + b*atanh(c/x^2)),x)`

output `(a*x^6)/6 + (b*c^3*log(x^4 - c^2))/12 + (b*x^6*log(c + x^2))/12 + (b*c*x^4)/12 - (b*x^6*log(x^2 - c))/12`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.11

$$\int x^5 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = -\frac{\operatorname{atanh} \left(\frac{c}{x^2} \right) b c^3}{6} + \frac{\operatorname{atanh} \left(\frac{c}{x^2} \right) b x^6}{6} + \frac{\log(x^2 + c) b c^3}{6} + \frac{a x^6}{6} + \frac{b c x^4}{12}$$

input `int(x^5*(a+b*atanh(c/x^2)),x)`output `(- 2*atanh(c/x**2)*b*c**3 + 2*atanh(c/x**2)*b*x**6 + 2*log(c + x**2)*b*c*
*3 + 2*a*x**6 + b*c*x**4)/12`

3.159 $\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx$

Optimal result	1302
Mathematica [A] (verified)	1302
Rubi [A] (verified)	1303
Maple [A] (verified)	1305
Fricas [A] (verification not implemented)	1305
Sympy [A] (verification not implemented)	1306
Maxima [A] (verification not implemented)	1306
Giac [B] (verification not implemented)	1306
Mupad [B] (verification not implemented)	1307
Reduce [B] (verification not implemented)	1307

Optimal result

Integrand size = 14, antiderivative size = 43

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{1}{4} b c x^2 + \frac{1}{4} x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) - \frac{1}{4} b c^2 \operatorname{arctanh} \left(\frac{x^2}{c} \right)$$

output

```
1/4*b*c*x^2+1/4*x^4*(a+b*arctanh(c/x^2))-1/4*b*c^2*arctanh(x^2/c)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.44

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{1}{4} b c x^2 + \frac{a x^4}{4} + \frac{1}{4} b x^4 \operatorname{arctanh} \left(\frac{c}{x^2} \right) + \frac{1}{8} b c^2 \log(-c + x^2) - \frac{1}{8} b c^2 \log(c + x^2)$$

input

```
Integrate[x^3*(a + b*ArcTanh[c/x^2]),x]
```

output

```
(b*c*x^2)/4 + (a*x^4)/4 + (b*x^4*ArcTanh[c/x^2])/4 + (b*c^2*Log[-c + x^2])/8 - (b*c^2*Log[c + x^2])/8
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.91, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {6452, 795, 807, 25, 262, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) dx \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{2}bc \int \frac{x}{1 - \frac{c^2}{x^4}} dx + \frac{1}{4}x^4 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) \\
 & \quad \downarrow \text{795} \\
 & \frac{1}{2}bc \int \frac{x^5}{x^4 - c^2} dx + \frac{1}{4}x^4 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) \\
 & \quad \downarrow \text{807} \\
 & \frac{1}{4}bc \int -\frac{x^4}{c^2 - x^4} dx^2 + \frac{1}{4}x^4 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{4}x^4 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) - \frac{1}{4}bc \int \frac{x^4}{c^2 - x^4} dx^2 \\
 & \quad \downarrow \text{262} \\
 & \frac{1}{4}bc \left(x^2 - c^2 \int \frac{1}{c^2 - x^4} dx^2 \right) + \frac{1}{4}x^4 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) \\
 & \quad \downarrow \text{219} \\
 & \frac{1}{4}x^4 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) + \frac{1}{4}bc \left(x^2 - \operatorname{carctanh} \left(\frac{x^2}{c} \right) \right)
 \end{aligned}$$

input `Int[x^3*(a + b*ArcTanh[c/x^2]),x]`

output `(x^4*(a + b*ArcTanh[c/x^2]))/4 + (b*c*(x^2 - c*ArcTanh[x^2/c]))/4`

Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 262 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 795 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Int[x^(m + n*p)*(b + a/x^n)^p, x] /; FreeQ[{a, b, m, n}, x] && IntegerQ[p] && NegQ[n]`
- rule 807 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`
- rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)]*(b_))^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.05

method	result	size
parallelrisc	$\frac{\operatorname{arctanh}\left(\frac{c}{x^2}\right)bx^4}{4} + \frac{x^4a}{4} + \frac{bcx^2}{4} - \frac{\operatorname{arctanh}\left(\frac{c}{x^2}\right)bc^2}{4} + \frac{ac^2}{4}$	45
derivativedivides	$\frac{x^4a}{4} + \frac{\operatorname{arctanh}\left(\frac{c}{x^2}\right)bx^4}{4} + \frac{bcx^2}{4} + \frac{bc^2 \ln\left(\frac{c}{x^2}-1\right)}{8} - \frac{bc^2 \ln\left(1+\frac{c}{x^2}\right)}{8}$	55
default	$\frac{x^4a}{4} + \frac{\operatorname{arctanh}\left(\frac{c}{x^2}\right)bx^4}{4} + \frac{bcx^2}{4} + \frac{bc^2 \ln\left(\frac{c}{x^2}-1\right)}{8} - \frac{bc^2 \ln\left(1+\frac{c}{x^2}\right)}{8}$	55
parts	$\frac{x^4a}{4} + \frac{\operatorname{arctanh}\left(\frac{c}{x^2}\right)bx^4}{4} + \frac{bcx^2}{4} + \frac{bc^2 \ln\left(\frac{c}{x^2}-1\right)}{8} - \frac{bc^2 \ln\left(1+\frac{c}{x^2}\right)}{8}$	55
oring	$-\frac{5\left(a+b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)\left(-x^4+c^2\right)}{8} + \frac{\left(x^2+c\right)\left(-x^2+c\right)\left(3x^2\left(a+b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)-\frac{2bc}{1-\frac{c^2}{x^4}}\right)}{8x^2}$	72
risc	Expression too large to display	4322

input `int(x^3*(a+b*arctanh(c/x^2)),x,method=_RETURNVERBOSE)`output `1/4*arctanh(c/x^2)*b*x^4+1/4*x^4*a+1/4*b*c*x^2-1/4*arctanh(c/x^2)*b*c^2+1/4*a*c^2`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.02

$$\int x^3 \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right) \right) dx = \frac{1}{4} ax^4 + \frac{1}{4} bcx^2 + \frac{1}{8} (bx^4 - bc^2) \log\left(\frac{x^2 + c}{x^2 - c}\right)$$

input `integrate(x^3*(a+b*arctanh(c/x^2)),x, algorithm="fricas")`output `1/4*a*x^4 + 1/4*b*c*x^2 + 1/8*(b*x^4 - b*c^2)*log((x^2 + c)/(x^2 - c))`

Sympy [A] (verification not implemented)

Time = 1.69 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.95

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{ax^4}{4} - \frac{bc^2 \operatorname{atanh} \left(\frac{c}{x^2} \right)}{4} + \frac{bcx^2}{4} + \frac{bx^4 \operatorname{atanh} \left(\frac{c}{x^2} \right)}{4}$$

input `integrate(x**3*(a+b*atanh(c/x**2)),x)`

output `a*x**4/4 - b*c**2*atanh(c/x**2)/4 + b*c*x**2/4 + b*x**4*atanh(c/x**2)/4`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.14

$$\begin{aligned} & \int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx \\ &= \frac{1}{4} ax^4 + \frac{1}{8} \left(2x^4 \operatorname{artanh} \left(\frac{c}{x^2} \right) + (2x^2 - c \log(x^2 + c) + c \log(x^2 - c))c \right) b \end{aligned}$$

input `integrate(x^3*(a+b*arctanh(c/x^2)),x, algorithm="maxima")`

output `1/4*a*x^4 + 1/8*(2*x^4*arctanh(c/x^2) + (2*x^2 - c*log(x^2 + c) + c*log(x^2 - c))*c)*b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 162 vs. $2(37) = 74$.

Time = 0.12 (sec) , antiderivative size = 162, normalized size of antiderivative = 3.77

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{\frac{(x^2+c)bc^3 \log\left(\frac{x^2+c}{x^2-c}\right)}{(x^2-c)\left(\frac{(x^2+c)^2}{(x^2-c)^2} - \frac{2(x^2+c)}{x^2-c} + 1\right)} + \frac{\frac{2(x^2+c)ac^3}{x^2-c} + \frac{(x^2+c)bc^3}{x^2-c} - bc^3}{\frac{(x^2+c)^2}{(x^2-c)^2} - \frac{2(x^2+c)}{x^2-c} + 1}}{2c}$$

input `integrate(x^3*(a+b*arctanh(c/x^2)),x, algorithm="giac")`

output
$$\frac{1}{2}((x^2 + c)*b*c^3*\log((x^2 + c)/(x^2 - c)))/((x^2 - c)*((x^2 + c)^2/(x^2 - c)^2 - 2*(x^2 + c)/(x^2 - c) + 1)) + (2*(x^2 + c)*a*c^3/(x^2 - c) + (x^2 + c)*b*c^3/(x^2 - c) - b*c^3)/((x^2 + c)^2/(x^2 - c)^2 - 2*(x^2 + c)/(x^2 - c) + 1))/c$$

Mupad [B] (verification not implemented)

Time = 3.89 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.33

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{a x^4}{4} + \frac{b x^4 \ln(x^2 + c)}{8} + \frac{b c x^2}{4} - \frac{b x^4 \ln(x^2 - c)}{8} + \frac{b c^2 \operatorname{atan} \left(\frac{x^2 \operatorname{li}}{c} \right) \operatorname{li}}{4}$$

input `int(x^3*(a + b*atanh(c/x^2)),x)`

output
$$\frac{(a*x^4)}{4} + \frac{(b*x^4*\log(c + x^2))}{8} + \frac{(b*c^2*\operatorname{atan}((x^2*\operatorname{li})/c)*\operatorname{li})}{4} + \frac{(b*c*x^2)}{4} - \frac{(b*x^4*\log(x^2 - c))}{8}$$

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.88

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = -\frac{\operatorname{atanh} \left(\frac{c}{x^2} \right) b c^2}{4} + \frac{\operatorname{atanh} \left(\frac{c}{x^2} \right) b x^4}{4} + \frac{a x^4}{4} + \frac{b c x^2}{4}$$

input `int(x^3*(a+b*atanh(c/x^2)),x)`

output
$$(-\operatorname{atanh}(c/x^{**2})*b*c^{**2} + \operatorname{atanh}(c/x^{**2})*b*x^{**4} + a*x^{**4} + b*c*x^{**2})/4$$

3.160 $\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx$

Optimal result	1308
Mathematica [A] (verified)	1308
Rubi [A] (verified)	1309
Maple [A] (verified)	1310
Fricas [A] (verification not implemented)	1310
Sympy [A] (verification not implemented)	1311
Maxima [A] (verification not implemented)	1311
Giac [B] (verification not implemented)	1312
Mupad [B] (verification not implemented)	1312
Reduce [B] (verification not implemented)	1313

Optimal result

Integrand size = 12, antiderivative size = 39

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{ax^2}{2} + \frac{1}{2}bx^2 \operatorname{arctanh} \left(\frac{c}{x^2} \right) + \frac{1}{4}bc \log(c^2 - x^4)$$

output `1/2*a*x^2+1/2*b*x^2*arctanh(c/x^2)+1/4*b*c*ln(-x^4+c^2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{ax^2}{2} + \frac{1}{2}bx^2 \operatorname{arctanh} \left(\frac{c}{x^2} \right) + \frac{1}{4}bc \log(-c^2 + x^4)$$

input `Integrate[x*(a + b*ArcTanh[c/x^2]),x]`

output `(a*x^2)/2 + (b*x^2*ArcTanh[c/x^2])/2 + (b*c*Log[-c^2 + x^4])/4`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.87, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6452, 795, 792}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \left(a + \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx$$

$$\downarrow 6452$$

$$bc \int \frac{1}{\left(1 - \frac{c^2}{x^4}\right) x} dx + \frac{1}{2} x^2 \left(a + \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)$$

$$\downarrow 795$$

$$bc \int \frac{x^3}{x^4 - c^2} dx + \frac{1}{2} x^2 \left(a + \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)$$

$$\downarrow 792$$

$$\frac{1}{2} x^2 \left(a + \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) + \frac{1}{4} bc \log (c^2 - x^4)$$

input `Int[x*(a + b*ArcTanh[c/x^2]),x]`

output `(x^2*(a + b*ArcTanh[c/x^2]))/2 + (b*c*Log[c^2 - x^4])/4`

Defintions of rubi rules used

rule 792 `Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_.)), x_Symbol] := Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]`

rule 795 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^p, x_Symbol] := Int[x^(m + n*p)*(b + a/x^n)^p, x] /; FreeQ[{a, b, m, n}, x] && IntegerQ[p] && NegQ[n]`

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.08

method	result
parallelrisch	$\frac{bx^2 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{2} + \frac{\ln(x^2-c)bc}{2} + \frac{ax^2}{2} + \frac{\operatorname{arctanh}\left(\frac{c}{x^2}\right)bc}{2}$
derivativdivides	$\frac{ax^2}{2} - b\left(-\frac{x^2 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{2} + c\left(\ln\left(\frac{1}{x}\right) - \frac{\ln\left(1+\frac{c}{x^2}\right)}{4} - \frac{\ln\left(\frac{c}{x^2}-1\right)}{4}\right)\right)$
default	$\frac{ax^2}{2} - b\left(-\frac{x^2 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{2} + c\left(\ln\left(\frac{1}{x}\right) - \frac{\ln\left(1+\frac{c}{x^2}\right)}{4} - \frac{\ln\left(\frac{c}{x^2}-1\right)}{4}\right)\right)$
parts	$\frac{ax^2}{2} + b\left(\frac{x^2 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{2} - c\left(\ln\left(\frac{1}{x}\right) - \frac{\ln\left(1+\frac{c}{x^2}\right)}{4} - \frac{\ln\left(\frac{c}{x^2}-1\right)}{4}\right)\right)$
risch	$\frac{x^2 b \ln(x^2+c)}{4} - \frac{bx^2 \ln(-x^2+c)}{4} - \frac{i\pi b x^2 \operatorname{csgn}\left(\frac{i}{x^2}\right) \operatorname{csgn}(i(x^2+c)) \operatorname{csgn}\left(\frac{i(x^2+c)}{x^2}\right)}{8} - \frac{i\pi b x^2}{4} + \frac{i\pi b x^2 \operatorname{csgn}\left(\frac{1}{x}\right)}{4}$

input

```
int(x*(a+b*arctanh(c/x^2)),x,method=_RETURNVERBOSE)
```

output

```
1/2*b*x^2*arctanh(c/x^2)+1/2*ln(x^2-c)*b*c+1/2*a*x^2+1/2*arctanh(c/x^2)*b*
c
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.10

$$\int x\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right) dx = \frac{1}{4} b x^2 \log\left(\frac{x^2 + c}{x^2 - c}\right) + \frac{1}{2} a x^2 + \frac{1}{4} b c \log(x^4 - c^2)$$

input

```
integrate(x*(a+b*arctanh(c/x^2)),x, algorithm="fricas")
```

output $1/4*b*x^2*\log((x^2 + c)/(x^2 - c)) + 1/2*a*x^2 + 1/4*b*c*\log(x^4 - c^2)$

Sympy [A] (verification not implemented)

Time = 1.23 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.56

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{ax^2}{2} + \frac{bc \log(x - \sqrt{-c})}{2} + \frac{bc \log(x + \sqrt{-c})}{2} - \frac{bc \operatorname{atanh} \left(\frac{c}{x^2} \right)}{2} + \frac{bx^2 \operatorname{atanh} \left(\frac{c}{x^2} \right)}{2}$$

input `integrate(x*(a+b*atanh(c/x**2)),x)`

output $a*x**2/2 + b*c*\log(x - \sqrt{-c})/2 + b*c*\log(x + \sqrt{-c})/2 - b*c*atanh(c/x**2)/2 + b*x**2*atanh(c/x**2)/2$

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.87

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{1}{2} ax^2 + \frac{1}{4} \left(2x^2 \operatorname{artanh} \left(\frac{c}{x^2} \right) + c \log(x^4 - c^2) \right) b$$

input `integrate(x*(a+b*arctanh(c/x^2)),x, algorithm="maxima")`

output $1/2*a*x^2 + 1/4*(2*x^2*arctanh(c/x^2) + c*\log(x^4 - c^2))*b$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 184 vs. $2(33) = 66$.

Time = 0.15 (sec) , antiderivative size = 184, normalized size of antiderivative = 4.72

$$\int x \left(a + \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx$$

$$= \frac{1}{2} ax^2 + \frac{c^2 \left(\log \left(\frac{|-x^2-c|}{|-x^2+c|} \right) - \log \left(\left| \frac{x^2+c}{x^2-c} - 1 \right| \right) \right) + \frac{c^2 \log \left(\frac{c \left(\frac{x^2+c}{(x^2-c)c} - \frac{1}{c} \right)}{\frac{x^2+c+1}{x^2-c} + 1} \right) - \frac{c \left(\frac{x^2+c}{(x^2-c)c} - \frac{1}{c} \right)}{\frac{x^2+c+1}{x^2-c} - 1}}{2c} b$$

input `integrate(x*(a+b*arctanh(c/x^2)),x, algorithm="giac")`

output `1/2*a*x^2 + 1/2*(c^2*(log(abs(-x^2 - c)/abs(-x^2 + c)) - log(abs((x^2 + c)/(x^2 - c) - 1))) + c^2*log(-(c*((x^2 + c)/((x^2 - c)*c) - 1/c)/((x^2 + c)/(x^2 - c) + 1) + 1)/(c*((x^2 + c)/((x^2 - c)*c) - 1/c)/((x^2 + c)/(x^2 - c) + 1) - 1))/((x^2 + c)/(x^2 - c) - 1))*b/c`

Mupad [B] (verification not implemented)

Time = 3.70 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.21

$$\int x \left(a + \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{ax^2}{2} + \frac{bx^2 \ln(x^2 + c)}{4} + \frac{bc \ln(x^4 - c^2)}{4} - \frac{bx^2 \ln(x^2 - c)}{4}$$

input `int(x*(a + b*atanh(c/x^2)),x)`

output `(a*x^2)/2 + (b*x^2*log(c + x^2))/4 + (b*c*log(x^4 - c^2))/4 - (b*x^2*log(x^2 - c))/4`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = -\frac{\operatorname{atanh} \left(\frac{c}{x^2} \right) bc}{2} + \frac{\operatorname{atanh} \left(\frac{c}{x^2} \right) b x^2}{2} + \frac{\log(x^2 + c) bc}{2} + \frac{a x^2}{2}$$

input

```
int(x*(a+b*atanh(c/x^2)),x)
```

output

```
( - atanh(c/x**2)*b*c + atanh(c/x**2)*b*x**2 + log(c + x**2)*b*c + a*x**2 )  
/2
```

$$3.161 \quad \int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} dx$$

Optimal result	1314
Mathematica [A] (verified)	1314
Rubi [A] (verified)	1315
Maple [B] (verified)	1316
Fricas [F]	1316
Sympy [F]	1317
Maxima [F]	1317
Giac [F]	1317
Mupad [F(-1)]	1318
Reduce [F]	1318

Optimal result

Integrand size = 14, antiderivative size = 30

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} dx = a \log(x) + \frac{1}{4}b \operatorname{PolyLog}\left(2, -\frac{c}{x^2}\right) - \frac{1}{4}b \operatorname{PolyLog}\left(2, \frac{c}{x^2}\right)$$

output `a*ln(x)+1/4*b*polylog(2,-c/x^2)-1/4*b*polylog(2,c/x^2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} dx = a \log(x) + \frac{1}{4}b \left(\operatorname{PolyLog}\left(2, -\frac{c}{x^2}\right) - \operatorname{PolyLog}\left(2, \frac{c}{x^2}\right) \right)$$

input `Integrate[(a + b*ArcTanh[c/x^2])/x,x]`

output `a*Log[x] + (b*(PolyLog[2, -(c/x^2)] - PolyLog[2, c/x^2]))/4`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.23, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6450, 6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} dx$$

$$\downarrow 6450$$

$$-\frac{1}{2} \int x^2 \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right) \right) d\frac{1}{x^2}$$

$$\downarrow 6446$$

$$\frac{1}{2} \left(-a \log\left(\frac{1}{x^2}\right) + \frac{1}{2} b \operatorname{PolyLog}\left(2, -\frac{c}{x^2}\right) - \frac{1}{2} b \operatorname{PolyLog}\left(2, \frac{c}{x^2}\right) \right)$$

input `Int[(a + b*ArcTanh[c/x^2])/x,x]`

output `(-(a*Log[x^(-2)]) + (b*PolyLog[2, -(c/x^2)])/2 - (b*PolyLog[2, c/x^2])/2)/2`

Defintions of rubi rules used

rule 6446 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))/(x_), x_Symbol] :> Simp[a*Log[x], x] + (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) / ; FreeQ[{a, b, c}, x]`

rule 6450 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)]*(b_.))^(p_.)/(x_), x_Symbol] :> Simp[1/n Subst[Int[(a + b*ArcTanh[c*x])^p/x, x], x, x^n], x] / ; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 148 vs. $2(26) = 52$.

Time = 0.37 (sec) , antiderivative size = 149, normalized size of antiderivative = 4.97

method	result
parts	$a \ln(x) + b \left(-\ln\left(\frac{1}{x}\right) \operatorname{arctanh}\left(\frac{c}{x^2}\right) + 2c \left(-\frac{\ln\left(\frac{1}{x}\right) \left(\ln\left(1 - \frac{\sqrt{c}}{x}\right) + \ln\left(1 + \frac{\sqrt{c}}{x}\right) \right)}{4c} - \frac{\operatorname{dilog}\left(1 - \frac{\sqrt{c}}{x}\right) + \operatorname{dilog}\left(1 + \frac{\sqrt{c}}{x}\right)}{4c} \right) \right)$
derivativedivides	$-a \ln\left(\frac{1}{x}\right) - b \left(\ln\left(\frac{1}{x}\right) \operatorname{arctanh}\left(\frac{c}{x^2}\right) - 2c \left(-\frac{\ln\left(\frac{1}{x}\right) \left(\ln\left(1 - \frac{\sqrt{c}}{x}\right) + \ln\left(1 + \frac{\sqrt{c}}{x}\right) \right)}{4c} - \frac{\operatorname{dilog}\left(1 - \frac{\sqrt{c}}{x}\right) + \operatorname{dilog}\left(1 + \frac{\sqrt{c}}{x}\right)}{4c} \right) \right)$
default	$-a \ln\left(\frac{1}{x}\right) - b \left(\ln\left(\frac{1}{x}\right) \operatorname{arctanh}\left(\frac{c}{x^2}\right) - 2c \left(-\frac{\ln\left(\frac{1}{x}\right) \left(\ln\left(1 - \frac{\sqrt{c}}{x}\right) + \ln\left(1 + \frac{\sqrt{c}}{x}\right) \right)}{4c} - \frac{\operatorname{dilog}\left(1 - \frac{\sqrt{c}}{x}\right) + \operatorname{dilog}\left(1 + \frac{\sqrt{c}}{x}\right)}{4c} \right) \right)$
risch	$\frac{b \ln(x) \ln(x^2+c)}{2} + \frac{\left(4a - 2ib\pi - ib\pi \operatorname{csgn}\left(\frac{i(x^2+c)}{x^2}\right) \right)^3 + 2ib\pi \operatorname{csgn}\left(\frac{i(-x^2+c)}{x^2}\right)^2 + ib\pi \operatorname{csgn}\left(\frac{i}{x^2}\right) \operatorname{csgn}(i(-x^2+c)) \operatorname{csgn}(i)}{\dots}$

input `int((a+b*arctanh(c/x^2))/x,x,method=_RETURNVERBOSE)`

output `a*ln(x)+b*(-ln(1/x)*arctanh(c/x^2)+2*c*(-1/4*ln(1/x)*(ln(1-1/x*c^(1/2))+ln(1+1/x*c^(1/2)))/c-1/4*(dilog(1-1/x*c^(1/2))+dilog(1+1/x*c^(1/2)))/c+1/4*ln(1/x)*(ln(1+1/x*(-c)^(1/2))+ln(1-1/x*(-c)^(1/2)))/c+1/4*(dilog(1+1/x*(-c)^(1/2))+dilog(1-1/x*(-c)^(1/2)))/c)`

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} dx = \int \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right) + a}{x} dx$$

input `integrate((a+b*arctanh(c/x^2))/x,x, algorithm="fricas")`

output `integral((b*arctanh(c/x^2) + a)/x, x)`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} dx = \int \frac{a + b \operatorname{atanh}\left(\frac{c}{x^2}\right)}{x} dx$$

input `integrate((a+b*atanh(c/x**2))/x,x)`

output `Integral((a + b*atanh(c/x**2))/x, x)`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} dx = \int \frac{b \operatorname{artanh}\left(\frac{c}{x^2}\right) + a}{x} dx$$

input `integrate((a+b*arctanh(c/x^2))/x,x, algorithm="maxima")`

output `1/2*b*integrate((log(c/x^2 + 1) - log(-c/x^2 + 1))/x, x) + a*log(x)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} dx = \int \frac{b \operatorname{artanh}\left(\frac{c}{x^2}\right) + a}{x} dx$$

input `integrate((a+b*arctanh(c/x^2))/x,x, algorithm="giac")`

output `integrate((b*arctanh(c/x^2) + a)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} dx = \int \frac{a + b \operatorname{atanh}\left(\frac{c}{x^2}\right)}{x} dx$$

input `int((a + b*atanh(c/x^2))/x,x)`output `int((a + b*atanh(c/x^2))/x, x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} dx = \left(\int \frac{\operatorname{atanh}\left(\frac{c}{x^2}\right)}{x} dx \right) b + \log(x) a$$

input `int((a+b*atanh(c/x^2))/x,x)`output `int(atanh(c/x**2)/x,x)*b + log(x)*a`

$$3.162 \quad \int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^3} dx$$

Optimal result	1319
Mathematica [A] (verified)	1319
Rubi [A] (verified)	1320
Maple [A] (verified)	1321
Fricas [A] (verification not implemented)	1321
Sympy [B] (verification not implemented)	1322
Maxima [A] (verification not implemented)	1322
Giac [A] (verification not implemented)	1323
Mupad [B] (verification not implemented)	1323
Reduce [B] (verification not implemented)	1323

Optimal result

Integrand size = 14, antiderivative size = 37

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^3} dx = -\frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{2x^2} - \frac{b \log\left(1 - \frac{c^2}{x^4}\right)}{4c}$$

output `-1/2*(a+b*arctanh(c/x^2))/x^2-1/4*b*ln(1-c^2/x^4)/c`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.14

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^3} dx = -\frac{a}{2x^2} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{2x^2} - \frac{b \log\left(1 - \frac{c^2}{x^4}\right)}{4c}$$

input `Integrate[(a + b*ArcTanh[c/x^2])/x^3,x]`

output `-1/2*a/x^2 - (b*ArcTanh[c/x^2])/(2*x^2) - (b*Log[1 - c^2/x^4])/(4*c)`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6452, 792}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^3} dx$$

↓ 6452

$$-bc \int \frac{1}{\left(1 - \frac{c^2}{x^4}\right) x^5} dx - \frac{a + \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{2x^2}$$

↓ 792

$$-\frac{a + \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{2x^2} - \frac{b \log\left(1 - \frac{c^2}{x^4}\right)}{4c}$$

input

```
Int[(a + b*ArcTanh[c/x^2])/x^3,x]
```

output

```
-1/2*(a + b*ArcTanh[c/x^2])/x^2 - (b*Log[1 - c^2/x^4])/(4*c)
```

Defintions of rubi rules used

rule 792

```
Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_.)), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]
```

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

method	result
parts	$-\frac{a}{2x^2} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{2x^2} - \frac{b \ln\left(1 - \frac{c^2}{x^4}\right)}{4c}$
derivativedivides	$-\frac{\frac{ca}{x^2} + b \left(\frac{c \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^2} + \frac{\ln\left(1 - \frac{c^2}{x^4}\right)}{2} \right)}{2c}$
default	$-\frac{\frac{ca}{x^2} + b \left(\frac{c \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^2} + \frac{\ln\left(1 - \frac{c^2}{x^4}\right)}{2} \right)}{2c}$
parallelrisc	$\frac{2b \ln(x)x^2 - \ln(x^2 - c)x^2 b - b x^2 \operatorname{arctanh}\left(\frac{c}{x^2}\right) - \operatorname{arctanh}\left(\frac{c}{x^2}\right) bc - ac}{2cx^2}$
risc	$-\frac{b \ln(x^2 + c)}{4x^2} - \frac{-2i\pi bc + i\pi bc \operatorname{csgn}\left(\frac{i}{x^2}\right) \operatorname{csgn}(i(-x^2 + c)) \operatorname{csgn}\left(\frac{i(-x^2 + c)}{x^2}\right) + i\pi bc \operatorname{csgn}\left(\frac{i}{x^2}\right) \operatorname{csgn}\left(\frac{i(x^2 + c)}{x^2}\right)^2 - i\pi bc}{4x^2}$

input `int((a+b*arctanh(c/x^2))/x^3,x,method=_RETURNVERBOSE)`output `-1/2*a/x^2-1/2*b*arctanh(c/x^2)/x^2-1/4*b*ln(1-c^2/x^4)/c`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.49

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^3} dx = -\frac{bx^2 \log(x^4 - c^2) - 4bx^2 \log(x) + bc \log\left(\frac{x^2 + c}{x^2 - c}\right) + 2ac}{4cx^2}$$

input `integrate((a+b*arctanh(c/x^2))/x^3,x, algorithm="fricas")`output `-1/4*(b*x^2*log(x^4 - c^2) - 4*b*x^2*log(x) + b*c*log((x^2 + c)/(x^2 - c)) + 2*a*c)/(c*x^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 76 vs. $2(32) = 64$.

Time = 4.18 (sec) , antiderivative size = 76, normalized size of antiderivative = 2.05

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^3} dx$$

$$= \begin{cases} -\frac{a}{2x^2} - \frac{b \operatorname{atanh}\left(\frac{c}{x^2}\right)}{2x^2} + \frac{b \log(x)}{c} - \frac{b \log(x - \sqrt{-c})}{2c} - \frac{b \log(x + \sqrt{-c})}{2c} + \frac{b \operatorname{atanh}\left(\frac{c}{x^2}\right)}{2c} & \text{for } c \neq 0 \\ -\frac{a}{2x^2} & \text{otherwise} \end{cases}$$

input `integrate((a+b*atanh(c/x**2))/x**3,x)`

output `Piecewise((-a/(2*x**2) - b*atanh(c/x**2)/(2*x**2) + b*log(x)/c - b*log(x - sqrt(-c))/(2*c) - b*log(x + sqrt(-c))/(2*c) + b*atanh(c/x**2)/(2*c), Ne(c, 0)), (-a/(2*x**2), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^3} dx = -\frac{b \left(\frac{2c \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^2} + \log\left(-\frac{c^2}{x^4} + 1\right) \right)}{4c} - \frac{a}{2x^2}$$

input `integrate((a+b*arctanh(c/x^2))/x^3,x, algorithm="maxima")`

output `-1/4*b*(2*c*arctanh(c/x^2)/x^2 + log(-c^2/x^4 + 1))/c - 1/2*a/x^2`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.41

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^3} dx = -\frac{b \log(x^4 - c^2)}{4c} + \frac{b \log(x)}{c} - \frac{b \log\left(\frac{x^2+c}{x^2-c}\right)}{4x^2} - \frac{a}{2x^2}$$

input `integrate((a+b*arctanh(c/x^2))/x^3,x, algorithm="giac")`output `-1/4*b*log(x^4 - c^2)/c + b*log(x)/c - 1/4*b*log((x^2 + c)/(x^2 - c))/x^2 - 1/2*a/x^2`**Mupad [B] (verification not implemented)**

Time = 3.82 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.51

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^3} dx = \frac{b \ln(x)}{c} - \frac{b \ln(x^4 - c^2)}{4c} - \frac{a}{2x^2} - \frac{b \ln(x^2 + c)}{4x^2} + \frac{b \ln(x^2 - c)}{4x^2}$$

input `int((a + b*atanh(c/x^2))/x^3,x)`output `(b*log(x))/c - (b*log(x^4 - c^2))/(4*c) - a/(2*x^2) - (b*log(c + x^2))/(4*x^2) + (b*log(x^2 - c))/(4*x^2)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.46

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^3} dx \\ &= \frac{-\operatorname{atanh}\left(\frac{c}{x^2}\right) bc + \operatorname{atanh}\left(\frac{c}{x^2}\right) b x^2 - \log(x^2 + c) b x^2 + 2 \log(x) b x^2 - ac}{2c x^2} \end{aligned}$$

input `int((a+b*atanh(c/x^2))/x^3,x)`

output $(- \operatorname{atanh}(c/x^{**2}) * b * c + \operatorname{atanh}(c/x^{**2}) * b * x^{**2} - \log(c + x^{**2}) * b * x^{**2} + 2 * \log(x) * b * x^{**2} - a * c) / (2 * c * x^{**2})$

$$3.163 \quad \int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^5} dx$$

Optimal result	1325
Mathematica [A] (verified)	1325
Rubi [A] (verified)	1326
Maple [A] (verified)	1328
Fricas [A] (verification not implemented)	1328
Sympy [A] (verification not implemented)	1329
Maxima [A] (verification not implemented)	1329
Giac [A] (verification not implemented)	1330
Mupad [B] (verification not implemented)	1330
Reduce [B] (verification not implemented)	1330

Optimal result

Integrand size = 14, antiderivative size = 45

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^5} dx = -\frac{b}{4cx^2} - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{4x^4} + \frac{b \operatorname{arctanh}\left(\frac{x^2}{c}\right)}{4c^2}$$

output $-1/4*b/c/x^2 - 1/4*(a + b*\operatorname{arctanh}(c/x^2))/x^4 + 1/4*b*\operatorname{arctanh}(x^2/c)/c^2$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.42

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^5} dx = -\frac{a}{4x^4} - \frac{b}{4cx^2} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{4x^4} - \frac{b \log(-c + x^2)}{8c^2} + \frac{b \log(c + x^2)}{8c^2}$$

input `Integrate[(a + b*ArcTanh[c/x^2])/x^5, x]`

output $-1/4*a/x^4 - b/(4*c*x^2) - (b*\operatorname{ArcTanh}[c/x^2])/(4*x^4) - (b*\operatorname{Log}[-c + x^2])/(8*c^2) + (b*\operatorname{Log}[c + x^2])/(8*c^2)$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {6452, 795, 807, 25, 264, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^5} dx \\
 & \quad \downarrow \text{6452} \\
 & -\frac{1}{2}bc \int \frac{1}{\left(1 - \frac{c^2}{x^4}\right) x^7} dx - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{4x^4} \\
 & \quad \downarrow \text{795} \\
 & -\frac{1}{2}bc \int \frac{1}{x^3 (x^4 - c^2)} dx - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{4x^4} \\
 & \quad \downarrow \text{807} \\
 & -\frac{1}{4}bc \int -\frac{1}{x^4 (c^2 - x^4)} dx^2 - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{4x^4} \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{4}bc \int \frac{1}{x^4 (c^2 - x^4)} dx^2 - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{4x^4} \\
 & \quad \downarrow \text{264} \\
 & -\frac{1}{4}bc \left(\frac{1}{c^2 x^2} - \frac{\int \frac{1}{c^2 - x^4} dx^2}{c^2} \right) - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{4x^4} \\
 & \quad \downarrow \text{219} \\
 & -\frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{4x^4} - \frac{1}{4}bc \left(\frac{1}{c^2 x^2} - \frac{\operatorname{arctanh}\left(\frac{x^2}{c}\right)}{c^3} \right)
 \end{aligned}$$

input

```
Int[(a + b*ArcTanh[c/x^2])/x^5, x]
```

output
$$-1/4*(a + b*\text{ArcTanh}[c/x^2])/x^4 - (b*c*(1/(c^2*x^2) - \text{ArcTanh}[x^2/c]/c^3))/4$$

Defintions of rubi rules used

rule 25
$$\text{Int}[-(F x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F x, x], x]$$

rule 219
$$\text{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[-b, 2])) * \text{ArcTanh}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 264
$$\text{Int}[(c \cdot x)^m \cdot (a + (b \cdot x)^2)^p, x_Symbol] \rightarrow \text{Simp}[(c \cdot x)^{m+1} \cdot (a + b \cdot x^2)^{p+1} / (a \cdot c \cdot (m+1)), x] - \text{Simp}[b \cdot (m+2 \cdot p+3) / (a \cdot c^2 \cdot (m+1)) \quad \text{Int}[(c \cdot x)^{m+2} \cdot (a + b \cdot x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 795
$$\text{Int}[(x)^m \cdot (a + (b \cdot x)^n)^p, x_Symbol] \rightarrow \text{Int}[x^{m+n \cdot p} \cdot (b + a/x^n)^p, x] /; \text{FreeQ}\{a, b, m, n\}, x \ \&\& \ \text{IntegerQ}[p] \ \&\& \ \text{NegQ}[n]$$

rule 807
$$\text{Int}[(x)^m \cdot (a + (b \cdot x)^n)^p, x_Symbol] \rightarrow \text{With}\{k = \text{GCD}[m+1, n]\}, \text{Simp}[1/k \quad \text{Subst}[\text{Int}[x^{(m+1)/k-1} \cdot (a + b \cdot x^{n/k})^p, x], x, x^k], x] /; k \neq 1] /; \text{FreeQ}\{a, b, p\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$$

rule 6452
$$\text{Int}[(a + \text{ArcTanh}[c \cdot x^n]) \cdot (b \cdot x)^m, x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^p / (m+1), x] - \text{Simp}[b \cdot c \cdot n \cdot (p/(m+1)) \quad \text{Int}[x^{m+n} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2 \cdot n})], x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$$

Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.98

method	result
parallelrisc	$-\frac{-\operatorname{arctanh}\left(\frac{c}{x^2}\right) b x^4 + b c x^2 + \operatorname{arctanh}\left(\frac{c}{x^2}\right) b c^2 + a c^2}{4 x^4 c^2}$
derivativedivides	$-\frac{a}{4 x^4} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{4 x^4} - \frac{b}{4 c x^2} + \frac{b \ln\left(1 + \frac{c}{x^2}\right)}{8 c^2} - \frac{b \ln\left(\frac{c}{x^2} - 1\right)}{8 c^2}$
default	$-\frac{a}{4 x^4} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{4 x^4} - \frac{b}{4 c x^2} + \frac{b \ln\left(1 + \frac{c}{x^2}\right)}{8 c^2} - \frac{b \ln\left(\frac{c}{x^2} - 1\right)}{8 c^2}$
parts	$-\frac{a}{4 x^4} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{4 x^4} - \frac{b}{4 c x^2} + \frac{b \ln\left(1 + \frac{c}{x^2}\right)}{8 c^2} - \frac{b \ln\left(\frac{c}{x^2} - 1\right)}{8 c^2}$
oring	$-\frac{7(-x^4+c^2)\left(a+b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)}{8 x^4 c^2} - \frac{x^2(-x^2+c)(x^2+c)\left(-\frac{2 b c}{x^8\left(1-\frac{c^2}{x^4}\right)} - \frac{5\left(a+b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)}{x^6}\right)}{8 c^2}$
risc	$-\frac{b \ln\left(x^2+c\right)}{8 x^4} - \frac{2 i \pi b c^2 \operatorname{csgn}\left(\frac{i\left(-x^2+c\right)}{x^2}\right)^2}{8 x^4} + i \pi b c^2 \operatorname{csgn}\left(i\left(x^2+c\right)\right) \operatorname{csgn}\left(\frac{i\left(x^2+c\right)}{x^2}\right)^2 - i \pi b c^2 \operatorname{csgn}\left(\frac{i\left(-x^2+c\right)}{x^2}\right)^3 + i \pi b c^2 \operatorname{csgn}\left(i\left(x^2+c\right)\right) \operatorname{csgn}\left(\frac{i\left(x^2+c\right)}{x^2}\right)^3$

input `int((a+b*arctanh(c/x^2))/x^5,x,method=_RETURNVERBOSE)`

output `-1/4*(-arctanh(c/x^2)*b*x^4+b*c*x^2+arctanh(c/x^2)*b*c^2+a*c^2)/x^4/c^2`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.16

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^5} dx = -\frac{2 b c x^2 + 2 a c^2 - (b x^4 - b c^2) \log\left(\frac{x^2+c}{x^2-c}\right)}{8 c^2 x^4}$$

input `integrate((a+b*arctanh(c/x^2))/x^5,x, algorithm="fricas")`

output `-1/8*(2*b*c*x^2 + 2*a*c^2 - (b*x^4 - b*c^2)*log((x^2 + c)/(x^2 - c)))/(c^2*x^4)`

Sympy [A] (verification not implemented)

Time = 5.21 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.09

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^5} dx = \begin{cases} -\frac{a}{4x^4} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{4x^4} - \frac{b}{4cx^2} + \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{4c^2} & \text{for } c \neq 0 \\ -\frac{a}{4x^4} & \text{otherwise} \end{cases}$$

input `integrate((a+b*atanh(c/x**2))/x**5,x)`output `Piecewise((-a/(4*x**4) - b*atanh(c/x**2)/(4*x**4) - b/(4*c*x**2) + b*atanh(c/x**2)/(4*c**2), Ne(c, 0)), (-a/(4*x**4), True))`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.24

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^5} dx = \frac{1}{8} \left(c \left(\frac{\log(x^2 + c)}{c^3} - \frac{\log(x^2 - c)}{c^3} - \frac{2}{c^2 x^2} \right) - \frac{2 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^4} \right) b - \frac{a}{4x^4}$$

input `integrate((a+b*arctanh(c/x^2))/x^5,x, algorithm="maxima")`output `1/8*(c*(log(x^2 + c)/c^3 - log(x^2 - c)/c^3 - 2/(c^2*x^2)) - 2*arctanh(c/x^2)/x^4)*b - 1/4*a/x^4`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.47

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^5} dx = \frac{b \log(x^2 + c)}{8c^2} - \frac{b \log(-x^2 + c)}{8c^2} - \frac{b \log\left(\frac{x^2+c}{x^2-c}\right)}{8x^4} - \frac{bx^2 + ac}{4cx^4}$$

input `integrate((a+b*arctanh(c/x^2))/x^5,x, algorithm="giac")`output `1/8*b*log(x^2 + c)/c^2 - 1/8*b*log(-x^2 + c)/c^2 - 1/8*b*log((x^2 + c)/(x^2 - c))/x^4 - 1/4*(b*x^2 + a*c)/(c*x^4)`**Mupad [B] (verification not implemented)**

Time = 3.81 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.31

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^5} dx = \frac{bx^4 \operatorname{atanh}\left(\frac{x^2}{c}\right)}{4} - \frac{bcx^2}{4} - \frac{a}{4} - \frac{b \ln(x^2-c)}{8} + \frac{b \ln(x^2+c)}{8}$$

input `int((a + b*atanh(c/x^2))/x^5,x)`output `((b*x^4*atanh(x^2/c))/4 - (b*c*x^2)/4)/(c^2*x^4) - (a/4 - (b*log(x^2 - c))/8 + (b*log(c + x^2))/8)/x^4`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^5} dx = \frac{-\operatorname{atanh}\left(\frac{c}{x^2}\right) bc^2 + \operatorname{atanh}\left(\frac{c}{x^2}\right) bx^4 - ac^2 - bcx^2}{4c^2x^4}$$

input `int((a+b*atanh(c/x^2))/x^5,x)`

output $(- \operatorname{atanh}(c/x^{**2}) * b * c^{**2} + \operatorname{atanh}(c/x^{**2}) * b * x^{**4} - a * c^{**2} - b * c * x^{**2}) / (4 * c * x^{**4})$

$$3.164 \quad \int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^7} dx$$

Optimal result	1332
Mathematica [A] (verified)	1332
Rubi [A] (verified)	1333
Maple [A] (verified)	1335
Fricas [A] (verification not implemented)	1335
Sympy [B] (verification not implemented)	1336
Maxima [A] (verification not implemented)	1336
Giac [A] (verification not implemented)	1337
Mupad [B] (verification not implemented)	1337
Reduce [B] (verification not implemented)	1337

Optimal result

Integrand size = 14, antiderivative size = 48

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^7} dx = -\frac{b}{12cx^4} - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{6x^6} - \frac{b \log\left(1 - \frac{c^2}{x^4}\right)}{12c^3}$$

output `-1/12*b/c/x^4-1/6*(a+b*arctanh(c/x^2))/x^6-1/12*b*ln(1-c^2/x^4)/c^3`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.29

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^7} dx = -\frac{a}{6x^6} - \frac{b}{12cx^4} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{6x^6} + \frac{b \log(x)}{3c^3} - \frac{b \log(-c^2 + x^4)}{12c^3}$$

input `Integrate[(a + b*ArcTanh[c/x^2])/x^7,x]`

output `-1/6*a/x^6 - b/(12*c*x^4) - (b*ArcTanh[c/x^2])/(6*x^6) + (b*Log[x])/(3*c^3) - (b*Log[-c^2 + x^4])/(12*c^3)`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.15, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {6452, 795, 798, 25, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barctanh}\left(\frac{c}{x^2}\right)}{x^7} dx \\
 & \quad \downarrow \text{6452} \\
 & -\frac{1}{3}bc \int \frac{1}{\left(1 - \frac{c^2}{x^4}\right)x^9} dx - \frac{a + \operatorname{barctanh}\left(\frac{c}{x^2}\right)}{6x^6} \\
 & \quad \downarrow \text{795} \\
 & -\frac{1}{3}bc \int \frac{1}{x^5(x^4 - c^2)} dx - \frac{a + \operatorname{barctanh}\left(\frac{c}{x^2}\right)}{6x^6} \\
 & \quad \downarrow \text{798} \\
 & -\frac{1}{12}bc \int -\frac{1}{x^8(c^2 - x^4)} dx^4 - \frac{a + \operatorname{barctanh}\left(\frac{c}{x^2}\right)}{6x^6} \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{12}bc \int \frac{1}{x^8(c^2 - x^4)} dx^4 - \frac{a + \operatorname{barctanh}\left(\frac{c}{x^2}\right)}{6x^6} \\
 & \quad \downarrow \text{54} \\
 & \frac{1}{12}bc \int \left(\frac{1}{c^4 x^4} + \frac{1}{c^2 x^8} + \frac{1}{c^4(c^2 - x^4)} \right) dx^4 - \frac{a + \operatorname{barctanh}\left(\frac{c}{x^2}\right)}{6x^6} \\
 & \quad \downarrow \text{2009} \\
 & -\frac{a + \operatorname{barctanh}\left(\frac{c}{x^2}\right)}{6x^6} - \frac{1}{12}bc \left(-\frac{\log(x^4)}{c^4} + \frac{1}{c^2 x^4} + \frac{\log(c^2 - x^4)}{c^4} \right)
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c/x^2])/x^7,x]`

output

$$-1/6*(a + b*\text{ArcTanh}[c/x^2])/x^6 - (b*c*(1/(c^2*x^4) - \text{Log}[x^4]/c^4 + \text{Log}[c^2 - x^4]/c^4))/12$$
Defintions of rubi rules used

rule 25

$$\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$$

rule 54

$$\text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m * (c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, x\} \&\& \text{ILtQ}[m, 0] \&\& \text{IntegerQ}[n] \&\& !(\text{IGtQ}[n, 0] \&\& \text{LtQ}[m + n + 2, 0])$$

rule 795

$$\text{Int}[(x)^m * (a + b*x)^n]^p, x_Symbol] \rightarrow \text{Int}[x^{m + n*p} * (b + a/x^n)^p, x] /; \text{FreeQ}\{a, b, m, n, x\} \&\& \text{IntegerQ}[p] \&\& \text{NegQ}[n]$$

rule 798

$$\text{Int}[(x)^m * (a + b*x)^n]^p, x_Symbol] \rightarrow \text{Simp}[1/n \quad \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1) * (a + b*x)^p}, x], x, x^n], x] /; \text{FreeQ}\{a, b, m, n, p, x\} \&\& \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]$$

rule 2009

$$\text{Int}[u, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

rule 6452

$$\text{Int}[(a + \text{ArcTanh}[c*x^n]) * (b)^p * (x)^m, x_Symbol] \rightarrow \text{Simp}[x^{m + 1} * (a + b*\text{ArcTanh}[c*x^n])^p / (m + 1), x] - \text{Simp}[b*c*n * (p / (m + 1)) \quad \text{Int}[x^{m + n} * (a + b*\text{ArcTanh}[c*x^n])^{p - 1} / (1 - c^2*x^{2*n})], x], x] /; \text{FreeQ}\{a, b, c, m, n, x\} \&\& \text{IGtQ}[p, 0] \&\& (\text{EqQ}[p, 1] \|\| (\text{EqQ}[n, 1] \&\& \text{IntegerQ}[m])) \&\& \text{NeQ}[m, -1]$$

Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.94

method	result
derivativedivides	$-\frac{a}{6x^6} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{6x^6} - \frac{b}{12cx^4} - \frac{b \ln\left(\frac{c^2}{x^4}-1\right)}{12c^3}$
default	$-\frac{a}{6x^6} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{6x^6} - \frac{b}{12cx^4} - \frac{b \ln\left(\frac{c^2}{x^4}-1\right)}{12c^3}$
parts	$-\frac{a}{6x^6} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{6x^6} - \frac{b}{12cx^4} - \frac{b \ln\left(\frac{c^2}{x^4}-1\right)}{12c^3}$
parallelrisch	$\frac{4b \ln(x)x^6 - 2 \ln(x^2 - c)x^6 - 2x^6 \operatorname{arctanh}\left(\frac{c}{x^2}\right) - b - b c^2 x^2 - 2 \operatorname{arctanh}\left(\frac{c}{x^2}\right) b c^3 - 2 a c^3}{12x^6 c^3}$
risch	$-\frac{b \ln(x^2 + c)}{12x^6} - \frac{-2i\pi b c^3 - i\pi b c^3 \operatorname{csgn}\left(\frac{i}{x^2}\right) \operatorname{csgn}\left(\frac{i(-x^2 + c)}{x^2}\right)^2 + i\pi b c^3 \operatorname{csgn}\left(\frac{i}{x^2}\right) \operatorname{csgn}(i(-x^2 + c)) \operatorname{csgn}\left(\frac{i(-x^2 + c)}{x^2}\right)}{12x^6 c^3}$

input `int((a+b*arctanh(c/x^2))/x^7,x,method=_RETURNVERBOSE)`

output `-1/6*a/x^6-1/6*b/x^6*arctanh(c/x^2)-1/12*b/c/x^4-1/12*b/c^3*ln(c^2/x^4-1)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.40

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^7} dx = -\frac{bx^6 \log(x^4 - c^2) - 4bx^6 \log(x) + bc^2x^2 + bc^3 \log\left(\frac{x^2+c}{x^2-c}\right) + 2ac^3}{12c^3x^6}$$

input `integrate((a+b*arctanh(c/x^2))/x^7,x, algorithm="fricas")`

output `-1/12*(b*x^6*log(x^4 - c^2) - 4*b*x^6*log(x) + b*c^2*x^2 + b*c^3*log((x^2 + c)/(x^2 - c)) + 2*a*c^3)/(c^3*x^6)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 94 vs. $2(42) = 84$.

Time = 7.97 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.96

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^7} dx$$

$$= \begin{cases} -\frac{a}{6x^6} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{6x^6} - \frac{b}{12cx^4} + \frac{b \log(x)}{3c^3} - \frac{b \log(x - \sqrt{-c})}{6c^3} - \frac{b \log(x + \sqrt{-c})}{6c^3} + \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{6c^3} & \text{for } c \neq 0 \\ -\frac{a}{6x^6} & \text{otherwise} \end{cases}$$

input `integrate((a+b*atanh(c/x**2))/x**7,x)`

output `Piecewise((-a/(6*x**6) - b*atanh(c/x**2)/(6*x**6) - b/(12*c*x**4) + b*log(x)/(3*c**3) - b*log(x - sqrt(-c))/(6*c**3) - b*log(x + sqrt(-c))/(6*c**3) + b*atanh(c/x**2)/(6*c**3), Ne(c, 0)), (-a/(6*x**6), True))`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.15

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^7} dx$$

$$= -\frac{1}{12} \left(c \left(\frac{\log(x^4 - c^2)}{c^4} - \frac{\log(x^4)}{c^4} + \frac{1}{c^2 x^4} \right) + \frac{2 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^6} \right) b - \frac{a}{6x^6}$$

input `integrate((a+b*arctanh(c/x^2))/x^7,x, algorithm="maxima")`

output `-1/12*(c*(log(x^4 - c^2)/c^4 - log(x^4)/c^4 + 1/(c^2*x^4)) + 2*arctanh(c/x^2)/x^6)*b - 1/6*a/x^6`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.35

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^7} dx = -\frac{b \log(x^4 - c^2)}{12 c^3} + \frac{b \log(x)}{3 c^3} - \frac{b \log\left(\frac{x^2+c}{x^2-c}\right)}{12 x^6} - \frac{b x^2 + 2 a c}{12 c x^6}$$

input `integrate((a+b*arctanh(c/x^2))/x^7,x, algorithm="giac")`

output `-1/12*b*log(x^4 - c^2)/c^3 + 1/3*b*log(x)/c^3 - 1/12*b*log((x^2 + c)/(x^2 - c))/x^6 - 1/12*(b*x^2 + 2*a*c)/(c*x^6)`

Mupad [B] (verification not implemented)

Time = 3.79 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.38

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^7} dx = \frac{b \ln(x)}{3 c^3} - \frac{b \ln(x^4 - c^2)}{12 c^3} - \frac{b}{12 c x^4} - \frac{a}{6 x^6} - \frac{b \ln(x^2 + c)}{12 x^6} + \frac{b \ln(x^2 - c)}{12 x^6}$$

input `int((a + b*atanh(c/x^2))/x^7,x)`

output `(b*log(x))/(3*c^3) - (b*log(x^4 - c^2))/(12*c^3) - b/(12*c*x^4) - a/(6*x^6) - (b*log(c + x^2))/(12*x^6) + (b*log(x^2 - c))/(12*x^6)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.42

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^7} dx = \frac{-2 \operatorname{atanh}\left(\frac{c}{x^2}\right) b c^3 + 2 \operatorname{atanh}\left(\frac{c}{x^2}\right) b x^6 - 2 \log(x^2 + c) b x^6 + 4 \log(x) b x^6 - 2 a c^3 - b c^2 x^2}{12 c^3 x^6}$$

input `int((a+b*atanh(c/x^2))/x^7,x)`

output `(- 2*atanh(c/x**2)*b*c**3 + 2*atanh(c/x**2)*b*x**6 - 2*log(c + x**2)*b*x*
*6 + 4*log(x)*b*x**6 - 2*a*c**3 - b*c**2*x**2)/(12*c**3*x**6)`

3.165 $\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx$

Optimal result	1339
Mathematica [A] (verified)	1339
Rubi [A] (verified)	1340
Maple [A] (verified)	1342
Fricas [A] (verification not implemented)	1343
Sympy [B] (verification not implemented)	1343
Maxima [A] (verification not implemented)	1344
Giac [A] (verification not implemented)	1345
Mupad [B] (verification not implemented)	1345
Reduce [B] (verification not implemented)	1346

Optimal result

Integrand size = 14, antiderivative size = 63

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{2}{15} b c x^3 + \frac{1}{5} b c^{5/2} \arctan \left(\frac{x}{\sqrt{c}} \right) + \frac{1}{5} x^5 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) - \frac{1}{5} b c^{5/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right)$$

output

```
2/15*b*c*x^3+1/5*b*c^(5/2)*arctan(x/c^(1/2))+1/5*x^5*(a+b*arctanh(c/x^2))-1/5*b*c^(5/2)*arctanh(x/c^(1/2))
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.40

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{2}{15} b c x^3 + \frac{a x^5}{5} + \frac{1}{5} b c^{5/2} \arctan \left(\frac{x}{\sqrt{c}} \right) + \frac{1}{5} b x^5 \operatorname{arctanh} \left(\frac{c}{x^2} \right) + \frac{1}{10} b c^{5/2} \log(\sqrt{c} - x) - \frac{1}{10} b c^{5/2} \log(\sqrt{c} + x)$$

input

```
Integrate[x^4*(a + b*ArcTanh[c/x^2]),x]
```


output

$$(2*b*c*x^3)/15 + (a*x^5)/5 + (b*c^{(5/2)*ArcTan[x/Sqrt[c]]})/5 + (b*x^5*ArcTanh[c/x^2])/5 + (b*c^{(5/2)*Log[Sqrt[c] - x]})/10 - (b*c^{(5/2)*Log[Sqrt[c] + x]})/10$$
Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.14, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6452, 795, 843, 25, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4 \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right) \right) dx$$

$$\downarrow 6452$$

$$\frac{2}{5}bc \int \frac{x^2}{1 - \frac{c^2}{x^4}} dx + \frac{1}{5}x^5 \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right) \right)$$

$$\downarrow 795$$

$$\frac{2}{5}bc \int \frac{x^6}{x^4 - c^2} dx + \frac{1}{5}x^5 \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right) \right)$$

$$\downarrow 843$$

$$\frac{2}{5}bc \left(c^2 \int -\frac{x^2}{c^2 - x^4} dx + \frac{x^3}{3} \right) + \frac{1}{5}x^5 \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right) \right)$$

$$\downarrow 25$$

$$\frac{2}{5}bc \left(\frac{x^3}{3} - c^2 \int \frac{x^2}{c^2 - x^4} dx \right) + \frac{1}{5}x^5 \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right) \right)$$

$$\downarrow 827$$

$$\frac{2}{5}bc \left(\frac{x^3}{3} - c^2 \left(\frac{1}{2} \int \frac{1}{c - x^2} dx - \frac{1}{2} \int \frac{1}{x^2 + c} dx \right) \right) + \frac{1}{5}x^5 \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right) \right)$$

$$\downarrow 216$$

$$\frac{2}{5}bc \left(\frac{x^3}{3} - c^2 \left(\frac{1}{2} \int \frac{1}{c - x^2} dx - \frac{\arctan\left(\frac{x}{\sqrt{c}}\right)}{2\sqrt{c}} \right) \right) + \frac{1}{5}x^5 \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right) \right)$$

$$\frac{1}{5}x^5\left(a + b\operatorname{arctanh}\left(\frac{c}{x^2}\right)\right) + \frac{2}{5}bc\left(\frac{x^3}{3} - c^2\left(\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right)}{2\sqrt{c}} - \frac{\operatorname{arctan}\left(\frac{x}{\sqrt{c}}\right)}{2\sqrt{c}}\right)\right)$$

input `Int[x^4*(a + b*ArcTanh[c/x^2]),x]`

output `(x^5*(a + b*ArcTanh[c/x^2]))/5 + (2*b*c*(x^3/3 - c^2*(-1/2*ArcTan[x/Sqrt[c]]/Sqrt[c] + ArcTanh[x/Sqrt[c]]/(2*Sqrt[c]))))/5`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 795 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Int[x^(m + n*p)*(b + a/x^n)^p, x] /; FreeQ[{a, b, m, n}, x] && IntegerQ[p] && NegQ[n]`

rule 827 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x], x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 843

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Simp[
a*c^n*(m - n + 1)/(b*(m + n*p + 1)) Int[(c*x)^(m - n)*(a + b*x^n)^p, x]
, x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*
p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.70 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.84

method	result
parts	$\frac{ax^5}{5} + \frac{bx^5 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{5} + \frac{bc^{\frac{5}{2}} \arctan\left(\frac{x}{\sqrt{c}}\right)}{5} - \frac{bc^{\frac{5}{2}} \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right)}{5} + \frac{2bcx^3}{15}$
derivativedivides	$\frac{ax^5}{5} + \frac{bx^5 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{5} - \frac{bc^{\frac{5}{2}} \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right)}{5} - \frac{bc^{\frac{5}{2}} \arctan\left(\frac{\sqrt{c}}{x}\right)}{5} + \frac{2bcx^3}{15}$
default	$\frac{ax^5}{5} + \frac{bx^5 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{5} - \frac{bc^{\frac{5}{2}} \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right)}{5} - \frac{bc^{\frac{5}{2}} \arctan\left(\frac{\sqrt{c}}{x}\right)}{5} + \frac{2bcx^3}{15}$
risch	$\frac{bx^5 \ln(x^2+c)}{10} - \frac{bx^5 \ln(-x^2+c)}{10} - \frac{i\pi bx^5}{10} - \frac{i\pi bx^5 \operatorname{csgn}(i(-x^2+c)) \operatorname{csgn}\left(\frac{i(-x^2+c)}{x^2}\right)^2}{20} + \frac{i\pi bx^5 \operatorname{csgn}\left(\frac{i}{x^2}\right) \operatorname{csgn}(c)}{20}$

input

```
int(x^4*(a+b*arctanh(c/x^2)),x,method=_RETURNVERBOSE)
```

output

```
1/5*a*x^5+1/5*b*x^5*arctanh(c/x^2)+1/5*b*c^(5/2)*arctan(x/c^(1/2))-1/5*b*c
^(5/2)*arctanh(1/x*c^(1/2))+2/15*b*c*x^3
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 170, normalized size of antiderivative = 2.70

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \left[\frac{1}{10} b x^5 \log \left(\frac{x^2 + c}{x^2 - c} \right) + \frac{1}{5} a x^5 + \frac{2}{15} b c x^3 \right. \\ \left. + \frac{1}{5} b c^{\frac{5}{2}} \arctan \left(\frac{x}{\sqrt{c}} \right) \right. \\ \left. + \frac{1}{10} b c^{\frac{5}{2}} \log \left(\frac{x^2 - 2\sqrt{c}x + c}{x^2 - c} \right), \frac{1}{10} b x^5 \log \left(\frac{x^2 + c}{x^2 - c} \right) \right. \\ \left. + \frac{1}{5} a x^5 + \frac{2}{15} b c x^3 + \frac{1}{5} b \sqrt{-c} c^2 \arctan \left(\frac{\sqrt{-c}x}{c} \right) \right. \\ \left. + \frac{1}{10} b \sqrt{-c} c^2 \log \left(\frac{x^2 + 2\sqrt{-c}x - c}{x^2 + c} \right) \right]$$

input `integrate(x^4*(a+b*arctanh(c/x^2)),x, algorithm="fricas")`

output `[1/10*b*x^5*log((x^2 + c)/(x^2 - c)) + 1/5*a*x^5 + 2/15*b*c*x^3 + 1/5*b*c^(5/2)*arctan(x/sqrt(c)) + 1/10*b*c^(5/2)*log((x^2 - 2*sqrt(c)*x + c)/(x^2 - c)), 1/10*b*x^5*log((x^2 + c)/(x^2 - c)) + 1/5*a*x^5 + 2/15*b*c*x^3 + 1/5*b*sqrt(-c)*c^2*arctan(sqrt(-c)*x/c) + 1/10*b*sqrt(-c)*c^2*log((x^2 + 2*sqrt(-c)*x - c)/(x^2 + c))]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 845 vs. 2(58) = 116.

Time = 3.85 (sec) , antiderivative size = 845, normalized size of antiderivative = 13.41

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \text{Too large to display}$$

input `integrate(x**4*(a+b*atanh(c/x**2)),x)`

output

```
Piecewise((a*x**5/5, Eq(c, 0)), (x**5*(a - oo*b)/5, Eq(c, -x**2)), (x**5*(
a + oo*b)/5, Eq(c, x**2)), (-6*a*c**2*x**5*sqrt(-c)/(-30*c**2*sqrt(-c) + 3
0*x**4*sqrt(-c)) + 6*a*x**9*sqrt(-c)/(-30*c**2*sqrt(-c) + 30*x**4*sqrt(-c)
) - 6*b*c**(9/2)*sqrt(-c)*log(-sqrt(c) + x)/(-30*c**2*sqrt(-c) + 30*x**4*s
qrt(-c)) + 3*b*c**(9/2)*sqrt(-c)*log(x - sqrt(-c))/(-30*c**2*sqrt(-c) + 30
*x**4*sqrt(-c)) + 3*b*c**(9/2)*sqrt(-c)*log(x + sqrt(-c))/(-30*c**2*sqrt(-
c) + 30*x**4*sqrt(-c)) - 6*b*c**(9/2)*sqrt(-c)*atanh(c/x**2)/(-30*c**2*sq
rt(-c) + 30*x**4*sqrt(-c)) + 6*b*c**(5/2)*x**4*sqrt(-c)*log(-sqrt(c) + x)/(
-30*c**2*sqrt(-c) + 30*x**4*sqrt(-c)) - 3*b*c**(5/2)*x**4*sqrt(-c)*log(x -
sqrt(-c))/(-30*c**2*sqrt(-c) + 30*x**4*sqrt(-c)) - 3*b*c**(5/2)*x**4*sqrt
(-c)*log(x + sqrt(-c))/(-30*c**2*sqrt(-c) + 30*x**4*sqrt(-c)) + 6*b*c**(5/
2)*x**4*sqrt(-c)*atanh(c/x**2)/(-30*c**2*sqrt(-c) + 30*x**4*sqrt(-c)) - 3*
b*c**5*log(x - sqrt(-c))/(-30*c**2*sqrt(-c) + 30*x**4*sqrt(-c)) + 3*b*c**5
*log(x + sqrt(-c))/(-30*c**2*sqrt(-c) + 30*x**4*sqrt(-c)) + 3*b*c**3*x**4*
log(x - sqrt(-c))/(-30*c**2*sqrt(-c) + 30*x**4*sqrt(-c)) - 3*b*c**3*x**4*
log(x + sqrt(-c))/(-30*c**2*sqrt(-c) + 30*x**4*sqrt(-c)) - 4*b*c**3*x**3*sq
rt(-c)/(-30*c**2*sqrt(-c) + 30*x**4*sqrt(-c)) - 6*b*c**2*x**5*sqrt(-c)*ata
nh(c/x**2)/(-30*c**2*sqrt(-c) + 30*x**4*sqrt(-c)) + 4*b*c*x**7*sqrt(-c)/(-
30*c**2*sqrt(-c) + 30*x**4*sqrt(-c)) + 6*b*x**9*sqrt(-c)*atanh(c/x**2)/(-3
0*c**2*sqrt(-c) + 30*x**4*sqrt(-c)), True))
```

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.98

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx$$

$$= \frac{1}{5} a x^5$$

$$+ \frac{1}{30} \left(6 x^5 \operatorname{arctanh} \left(\frac{c}{x^2} \right) + \left(4 x^3 + 6 c^{\frac{3}{2}} \operatorname{arctan} \left(\frac{x}{\sqrt{c}} \right) + 3 c^{\frac{3}{2}} \log \left(\frac{x - \sqrt{c}}{x + \sqrt{c}} \right) \right) c \right) b$$

input

```
integrate(x^4*(a+b*arctanh(c/x^2)),x, algorithm="maxima")
```

output

```
1/5*a*x^5 + 1/30*(6*x^5*arctanh(c/x^2) + (4*x^3 + 6*c^(3/2)*arctan(x/sqrt(
c)) + 3*c^(3/2)*log((x - sqrt(c))/(x + sqrt(c))))*c)*b
```

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.06

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{1}{10} b x^5 \log \left(\frac{x^2 + c}{x^2 - c} \right) + \frac{1}{5} a x^5 + \frac{2}{15} b c x^3$$

$$+ \frac{b c^3 \arctan \left(\frac{x}{\sqrt{-c}} \right)}{5 \sqrt{-c}} + \frac{1}{5} b c^{5/2} \arctan \left(\frac{x}{\sqrt{c}} \right)$$

input `integrate(x^4*(a+b*arctanh(c/x^2)),x, algorithm="giac")`

output `1/10*b*x^5*log((x^2 + c)/(x^2 - c)) + 1/5*a*x^5 + 2/15*b*c*x^3 + 1/5*b*c^3*arctan(x/sqrt(-c))/sqrt(-c) + 1/5*b*c^(5/2)*arctan(x/sqrt(c))`

Mupad [B] (verification not implemented)

Time = 3.87 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.06

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{a x^5}{5} + \frac{b c^{5/2} \operatorname{atan} \left(\frac{x}{\sqrt{c}} \right)}{5} + \frac{b x^5 \ln(x^2 + c)}{10}$$

$$+ \frac{2 b c x^3}{15} - \frac{b x^5 \ln(x^2 - c)}{10} + \frac{b c^{5/2} \operatorname{atan} \left(\frac{x \operatorname{li}}{\sqrt{c}} \right) \operatorname{li}}{5}$$

input `int(x^4*(a + b*atanh(c/x^2)),x)`

output `(a*x^5)/5 + (b*c^(5/2)*atan(x/c^(1/2)))/5 + (b*c^(5/2)*atan((x*li)/c^(1/2))*li)/5 + (b*x^5*log(c + x^2))/10 + (2*b*c*x^3)/15 - (b*x^5*log(x^2 - c))/10`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.33

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{\sqrt{c} \operatorname{atan} \left(\frac{x}{\sqrt{c}} \right) b c^2}{5} + \frac{\sqrt{c} \operatorname{atanh} \left(\frac{c}{x^2} \right) b c^2}{5} \\ + \frac{\operatorname{atanh} \left(\frac{c}{x^2} \right) b x^5}{5} + \frac{\sqrt{c} \log(\sqrt{c} - x) b c^2}{5} \\ - \frac{\sqrt{c} \log(x^2 + c) b c^2}{10} + \frac{a x^5}{5} + \frac{2 b c x^3}{15}$$

input `int(x^4*(a+b*atanh(c/x^2)),x)`output `(6*sqrt(c)*atan(x/sqrt(c))*b*c**2 + 6*sqrt(c)*atanh(c/x**2)*b*c**2 + 6*atanh(c/x**2)*b*x**5 + 6*sqrt(c)*log(sqrt(c) - x)*b*c**2 - 3*sqrt(c)*log(c + x**2)*b*c**2 + 6*a*x**5 + 4*b*c*x**3)/30`

3.166 $\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx$

Optimal result	1347
Mathematica [A] (verified)	1347
Rubi [A] (verified)	1348
Maple [A] (verified)	1350
Fricas [A] (verification not implemented)	1351
Sympy [B] (verification not implemented)	1351
Maxima [A] (verification not implemented)	1352
Giac [A] (verification not implemented)	1353
Mupad [B] (verification not implemented)	1353
Reduce [B] (verification not implemented)	1354

Optimal result

Integrand size = 14, antiderivative size = 61

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{2bcx}{3} - \frac{1}{3} bc^{3/2} \arctan \left(\frac{x}{\sqrt{c}} \right) + \frac{1}{3} x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) - \frac{1}{3} bc^{3/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right)$$

output

```
2/3*b*c*x-1/3*b*c^(3/2)*arctan(x/c^(1/2))+1/3*x^3*(a+b*arctanh(c/x^2))-1/3
*b*c^(3/2)*arctanh(x/c^(1/2))
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.41

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{2bcx}{3} + \frac{ax^3}{3} - \frac{1}{3} bc^{3/2} \arctan \left(\frac{x}{\sqrt{c}} \right) + \frac{1}{3} bx^3 \operatorname{arctanh} \left(\frac{c}{x^2} \right) + \frac{1}{6} bc^{3/2} \log(\sqrt{c} - x) - \frac{1}{6} bc^{3/2} \log(\sqrt{c} + x)$$

input

```
Integrate[x^2*(a + b*ArcTanh[c/x^2]), x]
```


output

$$\frac{(2bcx)/3 + (ax^3)/3 - (bc^{3/2})\text{ArcTan}[x/\text{Sqrt}[c]]/3 + (bx^3\text{ArcTanh}[c/x^2])/3 + (bc^{3/2})\text{Log}[\text{Sqrt}[c] - x]/6 - (bc^{3/2})\text{Log}[\text{Sqrt}[c] + x])/6$$
Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.07, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {6452, 772, 843, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \left(a + \text{barctanh} \left(\frac{c}{x^2} \right) \right) dx$$

$$\downarrow 6452$$

$$\frac{2}{3}bc \int \frac{1}{1 - \frac{c^2}{x^4}} dx + \frac{1}{3}x^3 \left(a + \text{barctanh} \left(\frac{c}{x^2} \right) \right)$$

$$\downarrow 772$$

$$\frac{2}{3}bc \int \frac{x^4}{x^4 - c^2} dx + \frac{1}{3}x^3 \left(a + \text{barctanh} \left(\frac{c}{x^2} \right) \right)$$

$$\downarrow 843$$

$$\frac{2}{3}bc \left(c^2 \int \frac{1}{x^4 - c^2} dx + x \right) + \frac{1}{3}x^3 \left(a + \text{barctanh} \left(\frac{c}{x^2} \right) \right)$$

$$\downarrow 756$$

$$\frac{2}{3}bc \left(c^2 \left(-\frac{\int \frac{1}{c-x^2} dx}{2c} - \frac{\int \frac{1}{x^2+c} dx}{2c} \right) + x \right) + \frac{1}{3}x^3 \left(a + \text{barctanh} \left(\frac{c}{x^2} \right) \right)$$

$$\downarrow 216$$

$$\frac{2}{3}bc \left(c^2 \left(-\frac{\int \frac{1}{c-x^2} dx}{2c} - \frac{\arctan \left(\frac{x}{\sqrt{c}} \right)}{2c^{3/2}} \right) + x \right) + \frac{1}{3}x^3 \left(a + \text{barctanh} \left(\frac{c}{x^2} \right) \right)$$

$$\downarrow 219$$

$$\frac{1}{3}x^3\left(a + b\operatorname{arctanh}\left(\frac{c}{x^2}\right)\right) + \frac{2}{3}bc\left(c^2\left(-\frac{\arctan\left(\frac{x}{\sqrt{c}}\right)}{2c^{3/2}} - \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right)}{2c^{3/2}}\right) + x\right)$$

input `Int[x^2*(a + b*ArcTanh[c/x^2]),x]`

output `(x^3*(a + b*ArcTanh[c/x^2]))/3 + (2*b*c*(x + c^2*(-1/2*ArcTan[x/Sqrt[c]]/c^(3/2) - ArcTanh[x/Sqrt[c]]/(2*c^(3/2))))/3`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 756 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 772 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Int[x^(n*p)*(b + a/x^n)^p, x] /; FreeQ[{a, b}, x] && ILtQ[n, 0] && IntegerQ[p]`

rule 843 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n-1)*(c*x)^(m-n+1)*((a + b*x^n)^(p+1)/(b*(m+n*p+1))), x] - Simp[a*c^n*(m-n+1)/(b*(m+n*p+1)) Int[(c*x)^(m-n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n-1] && NeQ[m+n*p+1, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.84

method	result
parts	$\frac{ax^3}{3} + \frac{bx^3 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{3} - \frac{bc^{\frac{3}{2}} \arctan\left(\frac{x}{\sqrt{c}}\right)}{3} + \frac{2bcx}{3} - \frac{bc^{\frac{3}{2}} \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right)}{3}$
derivativedivides	$\frac{ax^3}{3} + \frac{bx^3 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{3} + \frac{bc^{\frac{3}{2}} \arctan\left(\frac{\sqrt{c}}{x}\right)}{3} + \frac{2bcx}{3} - \frac{bc^{\frac{3}{2}} \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right)}{3}$
default	$\frac{ax^3}{3} + \frac{bx^3 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{3} + \frac{bc^{\frac{3}{2}} \arctan\left(\frac{\sqrt{c}}{x}\right)}{3} + \frac{2bcx}{3} - \frac{bc^{\frac{3}{2}} \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right)}{3}$
risch	$\frac{bx^3 \ln(x^2+c)}{6} - \frac{bx^3 \ln(-x^2+c)}{6} + \frac{i\pi b x^3 \operatorname{csgn}\left(\frac{i}{x^2}\right) \operatorname{csgn}\left(\frac{i(x^2+c)}{x^2}\right)^2}{12} + \frac{i\pi b x^3 \operatorname{csgn}(i(x^2+c)) \operatorname{csgn}\left(\frac{i(x^2+c)}{x^2}\right)^2}{12}$

input

```
int(x^2*(a+b*arctanh(c/x^2)),x,method=_RETURNVERBOSE)
```

output

```
1/3*a*x^3+1/3*b*x^3*arctanh(c/x^2)-1/3*b*c^(3/2)*arctan(x/c^(1/2))+2/3*b*c
*x-1/3*b*c^(3/2)*arctanh(1/x*c^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 162, normalized size of antiderivative = 2.66

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \left[\frac{1}{6} b x^3 \log \left(\frac{x^2 + c}{x^2 - c} \right) + \frac{1}{3} a x^3 - \frac{1}{3} b c^{\frac{3}{2}} \arctan \left(\frac{x}{\sqrt{c}} \right) \right. \\ \left. + \frac{1}{6} b c^{\frac{3}{2}} \log \left(\frac{x^2 - 2\sqrt{cx} + c}{x^2 - c} \right) \right. \\ \left. + \frac{2}{3} b c x, \frac{1}{6} b x^3 \log \left(\frac{x^2 + c}{x^2 - c} \right) + \frac{1}{3} a x^3 \right. \\ \left. + \frac{1}{3} b \sqrt{-c} \arctan \left(\frac{\sqrt{-c} x}{c} \right) \right. \\ \left. + \frac{1}{6} b \sqrt{-c} \log \left(\frac{x^2 - 2\sqrt{-c} x - c}{x^2 + c} \right) + \frac{2}{3} b c x \right]$$

input `integrate(x^2*(a+b*arctanh(c/x^2)),x, algorithm="fricas")`

output `[1/6*b*x^3*log((x^2 + c)/(x^2 - c)) + 1/3*a*x^3 - 1/3*b*c^(3/2)*arctan(x/sqrt(c)) + 1/6*b*c^(3/2)*log((x^2 - 2*sqrt(c)*x + c)/(x^2 - c)) + 2/3*b*c*x, 1/6*b*x^3*log((x^2 + c)/(x^2 - c)) + 1/3*a*x^3 + 1/3*b*sqrt(-c)*c*arctan(sqrt(-c)*x/c) + 1/6*b*sqrt(-c)*c*log((x^2 - 2*sqrt(-c)*x - c)/(x^2 + c)) + 2/3*b*c*x]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 830 vs. 2(56) = 112.

Time = 2.80 (sec) , antiderivative size = 830, normalized size of antiderivative = 13.61

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \text{Too large to display}$$

input `integrate(x**2*(a+b*atanh(c/x**2)),x)`

output

```
Piecewise((a*x**3/3, Eq(c, 0)), (x**3*(a - oo*b)/3, Eq(c, -x**2)), (x**3*(a + oo*b)/3, Eq(c, x**2)), (-2*a*c**2*x**3*sqrt(-c)/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) + 2*a*x**7*sqrt(-c)/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) - 2*b*c**(7/2)*sqrt(-c)*log(-sqrt(c) + x)/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) + b*c**(7/2)*sqrt(-c)*log(x - sqrt(-c))/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) + b*c**(7/2)*sqrt(-c)*log(x + sqrt(-c))/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) - 2*b*c**(7/2)*sqrt(-c)*atanh(c/x**2)/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) + 2*b*c**(3/2)*x**4*sqrt(-c)*log(-sqrt(c) + x)/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) - b*c**(3/2)*x**4*sqrt(-c)*log(x - sqrt(-c))/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) - b*c**(3/2)*x**4*sqrt(-c)*log(x + sqrt(-c))/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) + 2*b*c**(3/2)*x**4*sqrt(-c)*atanh(c/x**2)/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) + b*c**4*log(x - sqrt(-c))/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) - b*c**4*log(x + sqrt(-c))/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) - 4*b*c**3*x*sqrt(-c)/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) - b*c**2*x**4*log(x - sqrt(-c))/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) + b*c**2*x**4*log(x + sqrt(-c))/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) - 2*b*c**2*x**3*sqrt(-c)*atanh(c/x**2)/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) + 4*b*c*x**5*sqrt(-c)/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)) + 2*b*x**7*sqrt(-c)*atanh(c/x**2)/(-6*c**2*sqrt(-c) + 6*x**4*sqrt(-c)), True))
```

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx$$

$$= \frac{1}{3} a x^3 + \frac{1}{6} \left(2 x^3 \operatorname{arctanh} \left(\frac{c}{x^2} \right) - \left(2 \sqrt{c} \arctan \left(\frac{x}{\sqrt{c}} \right) - \sqrt{c} \log \left(\frac{x - \sqrt{c}}{x + \sqrt{c}} \right) - 4 x \right) c \right) b$$

input

```
integrate(x^2*(a+b*arctanh(c/x^2)),x, algorithm="maxima")
```

output

```
1/3*a*x^3 + 1/6*(2*x^3*arctanh(c/x^2) - (2*sqrt(c)*arctan(x/sqrt(c)) - sqrt(c)*log((x - sqrt(c))/(x + sqrt(c))) - 4*x)*c)*b
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.13

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{1}{3} bc^3 \left(\frac{\arctan \left(\frac{x}{\sqrt{-c}} \right)}{\sqrt{-c}} - \frac{\arctan \left(\frac{x}{\sqrt{c}} \right)}{c^{3/2}} \right) + \frac{1}{6} bx^3 \log \left(\frac{x^2 + c}{x^2 - c} \right) + \frac{1}{3} ax^3 + \frac{2}{3} bcx$$

input `integrate(x^2*(a+b*arctanh(c/x^2)),x, algorithm="giac")`

output `1/3*b*c^3*(arctan(x/sqrt(-c))/(sqrt(-c)*c) - arctan(x/sqrt(c))/c^(3/2)) + 1/6*b*x^3*log((x^2 + c)/(x^2 - c)) + 1/3*a*x^3 + 2/3*b*c*x`

Mupad [B] (verification not implemented)

Time = 3.73 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.07

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \frac{ax^3}{3} - \frac{bc^{3/2} \operatorname{atan} \left(\frac{x}{\sqrt{c}} \right)}{3} + \frac{2bcx}{3} + \frac{bx^3 \ln(x^2 + c)}{6} - \frac{bx^3 \ln(x^2 - c)}{6} + \frac{bc^{3/2} \operatorname{atan} \left(\frac{x \operatorname{li}}{\sqrt{c}} \right) \operatorname{li}}{3}$$

input `int(x^2*(a + b*atanh(c/x^2)),x)`

output `(a*x^3)/3 - (b*c^(3/2)*atan(x/c^(1/2)))/3 + (b*c^(3/2)*atan((x*li)/c^(1/2))*li)/3 + (2*b*c*x)/3 + (b*x^3*log(c + x^2))/6 - (b*x^3*log(x^2 - c))/6`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.21

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = -\frac{\sqrt{c} \operatorname{atan} \left(\frac{x}{\sqrt{c}} \right) bc}{3} + \frac{\sqrt{c} \operatorname{atanh} \left(\frac{c}{x^2} \right) bc}{3} + \frac{\operatorname{atanh} \left(\frac{c}{x^2} \right) b x^3}{3} \\ + \frac{\sqrt{c} \log(\sqrt{c} - x) bc}{3} - \frac{\sqrt{c} \log(x^2 + c) bc}{6} + \frac{a x^3}{3} + \frac{2bcx}{3}$$

input

```
int(x^2*(a+b*atanh(c/x^2)),x)
```

output

```
( - 2*sqrt(c)*atan(x/sqrt(c))*b*c + 2*sqrt(c)*atanh(c/x**2)*b*c + 2*atanh(
c/x**2)*b*x**3 + 2*sqrt(c)*log(sqrt(c) - x)*b*c - sqrt(c)*log(c + x**2)*b*
c + 2*a*x**3 + 4*b*c*x)/6
```

3.167 $\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx$

Optimal result	1355
Mathematica [A] (verified)	1355
Rubi [A] (verified)	1356
Maple [A] (verified)	1357
Fricas [A] (verification not implemented)	1357
Sympy [A] (verification not implemented)	1358
Maxima [A] (verification not implemented)	1359
Giac [A] (verification not implemented)	1359
Mupad [B] (verification not implemented)	1360
Reduce [B] (verification not implemented)	1360

Optimal result

Integrand size = 10, antiderivative size = 44

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = ax + b\sqrt{c} \arctan \left(\frac{x}{\sqrt{c}} \right) + bx \operatorname{arctanh} \left(\frac{c}{x^2} \right) - b\sqrt{c} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right)$$

output

```
a*x+b*c^(1/2)*arctan(x/c^(1/2))+b*x*arctanh(c/x^2)-b*c^(1/2)*arctanh(x/c^(1/2))
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.23

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = ax + bx \operatorname{arctanh} \left(\frac{c}{x^2} \right) + \frac{1}{2} b \sqrt{c} \left(2 \arctan \left(\frac{x}{\sqrt{c}} \right) + \log(\sqrt{c} - x) - \log(\sqrt{c} + x) \right)$$

input

```
Integrate[a + b*ArcTanh[c/x^2], x]
```


output

```
a*x + b*x*ArcTanh[c/x^2] + (b*Sqrt[c]*(2*ArcTan[x/Sqrt[c]] + Log[Sqrt[c] -
x] - Log[Sqrt[c] + x]))/2
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx$$

↓ 2009

$$ax + b\sqrt{c} \arctan \left(\frac{x}{\sqrt{c}} \right) + bx \operatorname{arctanh} \left(\frac{c}{x^2} \right) - b\sqrt{c} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right)$$

input

```
Int[a + b*ArcTanh[c/x^2],x]
```

output

```
a*x + b*Sqrt[c]*ArcTan[x/Sqrt[c]] + b*x*ArcTanh[c/x^2] - b*Sqrt[c]*ArcTanh
[x/Sqrt[c]]
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.89

method	result
default	$ax + bx \operatorname{arctanh}\left(\frac{c}{x^2}\right) - b\sqrt{c} \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right) + b\sqrt{c} \operatorname{arctan}\left(\frac{x}{\sqrt{c}}\right)$
parts	$ax + bx \operatorname{arctanh}\left(\frac{c}{x^2}\right) - b\sqrt{c} \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right) + b\sqrt{c} \operatorname{arctan}\left(\frac{x}{\sqrt{c}}\right)$
derivativedivides	$ax + bx \operatorname{arctanh}\left(\frac{c}{x^2}\right) - b\sqrt{c} \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right) - b\sqrt{c} \operatorname{arctan}\left(\frac{\sqrt{c}}{x}\right)$
risch	$ax + \frac{bx \ln(x^2+c)}{2} - \frac{bx \ln(-x^2+c)}{2} + \frac{ib\pi x \operatorname{csgn}\left(\frac{i}{x^2}\right) \operatorname{csgn}(i(-x^2+c)) \operatorname{csgn}\left(\frac{i(-x^2+c)}{x^2}\right)}{4} - \frac{ib\pi x \operatorname{csgn}(i(-x^2+c))}{4}$

input

```
int(a+b*arctanh(c/x^2),x,method=_RETURNVERBOSE)
```

output

```
a*x+b*x*arctanh(c/x^2)-b*c^(1/2)*arctanh(1/x*c^(1/2))+b*c^(1/2)*arctan(x/c^(1/2))
```

Fricas [A] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 74 vs. 2(36) = 72.

Time = 0.09 (sec) , antiderivative size = 138, normalized size of antiderivative = 3.14

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right) \right) dx = \left[\frac{1}{2} bx \log\left(\frac{x^2+c}{x^2-c}\right) + b\sqrt{c} \operatorname{arctan}\left(\frac{x}{\sqrt{c}}\right) + \frac{1}{2} b\sqrt{c} \log\left(\frac{x^2-2\sqrt{cx}+c}{x^2-c}\right) + ax, \frac{1}{2} bx \log\left(\frac{x^2+c}{x^2-c}\right) + b\sqrt{-c} \operatorname{arctan}\left(\frac{\sqrt{-cx}}{c}\right) + \frac{1}{2} b\sqrt{-c} \log\left(\frac{x^2+2\sqrt{-cx}-c}{x^2+c}\right) + ax \right]$$

input `integrate(a+b*arctanh(c/x^2),x, algorithm="fricas")`

output `[1/2*b*x*log((x^2 + c)/(x^2 - c)) + b*sqrt(c)*arctan(x/sqrt(c)) + 1/2*b*sqrt(c)*log((x^2 - 2*sqrt(c)*x + c)/(x^2 - c)) + a*x, 1/2*b*x*log((x^2 + c)/(x^2 - c)) + b*sqrt(-c)*arctan(sqrt(-c)*x/c) + 1/2*b*sqrt(-c)*log((x^2 + 2*sqrt(-c)*x - c)/(x^2 + c)) + a*x]`

Sympy [A] (verification not implemented)

Time = 2.55 (sec) , antiderivative size = 632, normalized size of antiderivative = 14.36

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = ax$$

$$+ b \begin{cases} 0 \\ -\infty x \\ \infty x \\ -\frac{2c^{\frac{5}{2}}\sqrt{-c}\log(-\sqrt{-c}+x)}{-2c^2\sqrt{-c}+2x^4\sqrt{-c}} + \frac{c^{\frac{5}{2}}\sqrt{-c}\log(x-\sqrt{-c})}{-2c^2\sqrt{-c}+2x^4\sqrt{-c}} + \frac{c^{\frac{5}{2}}\sqrt{-c}\log(x+\sqrt{-c})}{-2c^2\sqrt{-c}+2x^4\sqrt{-c}} - \frac{2c^{\frac{5}{2}}\sqrt{-c}\operatorname{atanh}\left(\frac{c}{x^2}\right)}{-2c^2\sqrt{-c}+2x^4\sqrt{-c}} + \frac{2\sqrt{c}x^4\sqrt{-c}\log(-\sqrt{-c}+x)}{-2c^2\sqrt{-c}+2x^4\sqrt{-c}} \end{cases}$$

input `integrate(a+b*atanh(c/x**2),x)`

output `a*x + b*Piecewise((0, Eq(c, 0)), (-oo*x, Eq(c, -x**2)), (oo*x, Eq(c, x**2)), (-2*c**(5/2)*sqrt(-c)*log(-sqrt(c) + x)/(-2*c**2*sqrt(-c) + 2*x**4*sqrt(-c)) + c**(5/2)*sqrt(-c)*log(x - sqrt(-c))/(-2*c**2*sqrt(-c) + 2*x**4*sqrt(-c)) + c**(5/2)*sqrt(-c)*log(x + sqrt(-c))/(-2*c**2*sqrt(-c) + 2*x**4*sqrt(-c)) - 2*c**(5/2)*sqrt(-c)*atanh(c/x**2)/(-2*c**2*sqrt(-c) + 2*x**4*sqrt(-c)) + 2*sqrt(c)*x**4*sqrt(-c)*log(-sqrt(c) + x)/(-2*c**2*sqrt(-c) + 2*x**4*sqrt(-c)) - sqrt(c)*x**4*sqrt(-c)*log(x - sqrt(-c))/(-2*c**2*sqrt(-c) + 2*x**4*sqrt(-c)) - sqrt(c)*x**4*sqrt(-c)*log(x + sqrt(-c))/(-2*c**2*sqrt(-c) + 2*x**4*sqrt(-c)) + 2*sqrt(c)*x**4*sqrt(-c)*atanh(c/x**2)/(-2*c**2*sqrt(-c) + 2*x**4*sqrt(-c)) - c**3*log(x - sqrt(-c))/(-2*c**2*sqrt(-c) + 2*x**4*sqrt(-c)) + c**3*log(x + sqrt(-c))/(-2*c**2*sqrt(-c) + 2*x**4*sqrt(-c)) - 2*c**2*x*sqrt(-c)*atanh(c/x**2)/(-2*c**2*sqrt(-c) + 2*x**4*sqrt(-c)) + c*x**4*log(x - sqrt(-c))/(-2*c**2*sqrt(-c) + 2*x**4*sqrt(-c)) - c*x**4*log(x + sqrt(-c))/(-2*c**2*sqrt(-c) + 2*x**4*sqrt(-c)) + 2*x**5*sqrt(-c)*atanh(c/x**2)/(-2*c**2*sqrt(-c) + 2*x**4*sqrt(-c)), True))`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.16

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx$$

$$= \frac{1}{2} \left(c \left(\frac{2 \arctan \left(\frac{x}{\sqrt{c}} \right)}{\sqrt{c}} + \frac{\log \left(\frac{x-\sqrt{c}}{x+\sqrt{c}} \right)}{\sqrt{c}} \right) + 2x \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) b + ax$$

input `integrate(a+b*arctanh(c/x^2),x, algorithm="maxima")`output `1/2*(c*(2*arctan(x/sqrt(c))/sqrt(c) + log((x - sqrt(c))/(x + sqrt(c)))/sqrt(c)) + 2*x*arctanh(c/x^2))*b + a*x`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.30

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx$$

$$= \frac{1}{2} \left(2c \left(\frac{\arctan \left(\frac{x}{\sqrt{-c}} \right)}{\sqrt{-c}} + \frac{\arctan \left(\frac{x}{\sqrt{c}} \right)}{\sqrt{c}} \right) + x \log \left(-\frac{\frac{c}{x^2} + 1}{\frac{c}{x^2} - 1} \right) \right) b + ax$$

input `integrate(a+b*arctanh(c/x^2),x, algorithm="giac")`output `1/2*(2*c*(arctan(x/sqrt(-c))/sqrt(-c) + arctan(x/sqrt(c))/sqrt(c)) + x*log(-(c/x^2 + 1)/(c/x^2 - 1)))*b + a*x`

Mupad [B] (verification not implemented)

Time = 3.61 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.18

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = ax + \frac{bx \ln(x^2 + c)}{2} + b\sqrt{c} \operatorname{atan} \left(\frac{x}{\sqrt{c}} \right) - \frac{bx \ln(x^2 - c)}{2} + b\sqrt{c} \operatorname{atan} \left(\frac{x}{\sqrt{c}} \right) \operatorname{li}$$

input `int(a + b*atanh(c/x^2),x)`output `a*x + (b*x*log(c + x^2))/2 + b*c^(1/2)*atan(x/c^(1/2)) + b*c^(1/2)*atan((x*1i)/c^(1/2))*1i - (b*x*log(x^2 - c))/2`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.27

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \sqrt{c} \operatorname{atan} \left(\frac{x}{\sqrt{c}} \right) b + \sqrt{c} \operatorname{atanh} \left(\frac{c}{x^2} \right) b + \operatorname{atanh} \left(\frac{c}{x^2} \right) bx + \sqrt{c} \log(\sqrt{c} - x) b - \frac{\sqrt{c} \log(x^2 + c) b}{2} + ax$$

input `int(a+b*atanh(c/x^2),x)`output `(2*sqrt(c)*atan(x/sqrt(c))*b + 2*sqrt(c)*atanh(c/x**2)*b + 2*atanh(c/x**2)*b*x + 2*sqrt(c)*log(sqrt(c) - x)*b - sqrt(c)*log(c + x**2)*b + 2*a*x)/2`

3.168 $\int \frac{a+b\operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^2} dx$

Optimal result	1361
Mathematica [A] (verified)	1361
Rubi [A] (verified)	1362
Maple [A] (verified)	1364
Fricas [A] (verification not implemented)	1364
Sympy [B] (verification not implemented)	1365
Maxima [A] (verification not implemented)	1366
Giac [A] (verification not implemented)	1366
Mupad [B] (verification not implemented)	1367
Reduce [B] (verification not implemented)	1367

Optimal result

Integrand size = 14, antiderivative size = 46

$$\int \frac{a + b\operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^2} dx = \frac{b \arctan\left(\frac{x}{\sqrt{c}}\right)}{\sqrt{c}} - \frac{a + b\operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} + \frac{b\operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right)}{\sqrt{c}}$$

output

`b*arctan(x/c^(1/2))/c^(1/2)-(a+b*arctanh(c/x^2))/x+b*arctanh(x/c^(1/2))/c^(1/2)`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.57

$$\int \frac{a + b\operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^2} dx = -\frac{a}{x} + \frac{b \arctan\left(\frac{x}{\sqrt{c}}\right)}{\sqrt{c}} - \frac{b\operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} - \frac{b \log(\sqrt{c} - x)}{2\sqrt{c}} + \frac{b \log(\sqrt{c} + x)}{2\sqrt{c}}$$

input

`Integrate[(a + b*ArcTanh[c/x^2])/x^2,x]`

output

$$-(a/x) + (b*\text{ArcTan}[x/\text{Sqrt}[c]])/\text{Sqrt}[c] - (b*\text{ArcTanh}[c/x^2])/x - (b*\text{Log}[\text{Sqrt}[c] - x])/(2*\text{Sqrt}[c]) + (b*\text{Log}[\text{Sqrt}[c] + x])/(2*\text{Sqrt}[c])$$
Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.20, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {6452, 795, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^2} dx \\ & \quad \downarrow \text{6452} \\ & -2bc \int \frac{1}{\left(1 - \frac{c^2}{x^4}\right) x^4} dx - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} \\ & \quad \downarrow \text{795} \\ & -2bc \int \frac{1}{x^4 - c^2} dx - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} \\ & \quad \downarrow \text{756} \\ & -2bc \left(-\frac{\int \frac{1}{c-x^2} dx}{2c} - \frac{\int \frac{1}{x^2+c} dx}{2c} \right) - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} \\ & \quad \downarrow \text{216} \\ & -2bc \left(-\frac{\int \frac{1}{c-x^2} dx}{2c} - \frac{\arctan\left(\frac{x}{\sqrt{c}}\right)}{2c^{3/2}} \right) - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} \\ & \quad \downarrow \text{219} \\ & \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} - 2bc \left(-\frac{\arctan\left(\frac{x}{\sqrt{c}}\right)}{2c^{3/2}} - \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right)}{2c^{3/2}} \right) \end{aligned}$$

input `Int[(a + b*ArcTanh[c/x^2])/x^2,x]`

output `-((a + b*ArcTanh[c/x^2])/x) - 2*b*c*(-1/2*ArcTan[x/Sqrt[c]]/c^(3/2) - ArcTanh[x/Sqrt[c]]/(2*c^(3/2)))`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 756 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 795 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Int[x^(m + n*p)*(b + a/x^n)^p, x] /; FreeQ[{a, b, m, n}, x] && IntegerQ[p] && NegQ[n]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.96

method	result
parts	$-\frac{a}{x} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} + \frac{b \operatorname{arctan}\left(\frac{x}{\sqrt{c}}\right)}{\sqrt{c}} + \frac{b \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right)}{\sqrt{c}}$
derivativedivides	$-\frac{a}{x} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} + \frac{b \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right)}{\sqrt{c}} - \frac{b \operatorname{arctan}\left(\frac{\sqrt{c}}{x}\right)}{\sqrt{c}}$
default	$-\frac{a}{x} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} + \frac{b \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right)}{\sqrt{c}} - \frac{b \operatorname{arctan}\left(\frac{\sqrt{c}}{x}\right)}{\sqrt{c}}$
risch	$-\frac{b \ln(x^2+c)}{2x} - \frac{-i\pi bc \operatorname{csgn}\left(\frac{i}{x^2}\right) \operatorname{csgn}\left(\frac{i(-x^2+c)}{x^2}\right)^2}{2} - 2i\pi bc - i\pi bc \operatorname{csgn}\left(\frac{i(-x^2+c)}{x^2}\right)^3 + 2i\pi bc \operatorname{csgn}\left(\frac{i(-x^2+c)}{x^2}\right)^2 - i$

input

```
int((a+b*arctanh(c/x^2))/x^2,x,method=_RETURNVERBOSE)
```

output

```
-a/x-b/x*arctanh(c/x^2)+b*arctan(x/c^(1/2))/c^(1/2)+b/c^(1/2)*arctanh(1/x*
c^(1/2))
```

Fricas [A] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 84 vs. 2(38) = 76.

Time = 0.09 (sec) , antiderivative size = 159, normalized size of antiderivative = 3.46

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^2} dx$$

$$= \left[\frac{2b\sqrt{cx} \operatorname{arctan}\left(\frac{x}{\sqrt{c}}\right) + b\sqrt{cx} \log\left(\frac{x^2+2\sqrt{cx}+c}{x^2-c}\right) - bc \log\left(\frac{x^2+c}{x^2-c}\right) - 2ac}{2cx}, \right.$$

$$\left. - \frac{2b\sqrt{-cx} \operatorname{arctan}\left(\frac{\sqrt{-cx}}{c}\right) + b\sqrt{-cx} \log\left(\frac{x^2-2\sqrt{-cx}-c}{x^2+c}\right) + bc \log\left(\frac{x^2+c}{x^2-c}\right) + 2ac}{2cx} \right]$$

input

```
integrate((a+b*arctanh(c/x^2))/x^2,x, algorithm="fricas")
```

output

```
[1/2*(2*b*sqrt(c)*x*arctan(x/sqrt(c)) + b*sqrt(c)*x*log((x^2 + 2*sqrt(c)*x + c)/(x^2 - c)) - b*c*log((x^2 + c)/(x^2 - c)) - 2*a*c)/(c*x), -1/2*(2*b*sqrt(-c)*x*arctan(sqrt(-c)*x/c) + b*sqrt(-c)*x*log((x^2 - 2*sqrt(-c)*x - c)/(x^2 + c)) + b*c*log((x^2 + c)/(x^2 - c)) + 2*a*c)/(c*x)]
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 886 vs. $2(41) = 82$.

Time = 3.53 (sec) , antiderivative size = 886, normalized size of antiderivative = 19.26

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^2} dx = \text{Too large to display}$$

input

```
integrate((a+b*atanh(c/x**2))/x**2,x)
```

output

```
Piecewise((-a/x, Eq(c, 0)), (-(a - oo*b)/x, Eq(c, -x**2)), (-(a + oo*b)/x, Eq(c, x**2)), (2*a*c**(7/2)*sqrt(-c)/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c)) - 2*a*c**(3/2)*x**4*sqrt(-c)/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c)) - b*c**(7/2)*x*log(x - sqrt(-c))/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c)) + b*c**(7/2)*x*log(x + sqrt(-c))/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c)) + 2*b*c**(7/2)*sqrt(-c)*atanh(c/x**2)/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c)) + b*c**(3/2)*x**5*log(x - sqrt(-c))/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c)) - b*c**(3/2)*x**5*log(x + sqrt(-c))/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c)) - 2*b*c**(3/2)*x**4*sqrt(-c)*atanh(c/x**2)/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c)) + 2*b*c**3*x*sqrt(-c)*log(-sqrt(c) + x)/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c)) - b*c**3*x*sqrt(-c)*log(x - sqrt(-c))/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c)) - b*c**3*x*sqrt(-c)*log(x + sqrt(-c))/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c)) + 2*b*c**3*x*sqrt(-c)*atanh(c/x**2)/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c)) - 2*b*c*x**5*sqrt(-c)*log(-sqrt(c) + x)/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c)) + b*c*x**5*sqrt(-c)*log(x - sqrt(-c))/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c)) + b*c*x**5*sqrt(-c)*log(x + sqrt(-c))/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c)) - 2*b*c*x**5*sqrt(-c)*atanh(c/x**2)/(-2*c**(7/2)*x*sqrt(-c) + 2*c**(3/2)*x**5*sqrt(-c))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.24

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^2} dx = \frac{1}{2} \left(c \left(\frac{2 \arctan\left(\frac{x}{\sqrt{c}}\right)}{c^{\frac{3}{2}}} - \frac{\log\left(\frac{x-\sqrt{c}}{x+\sqrt{c}}\right)}{c^{\frac{3}{2}}}\right) - \frac{2 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x} \right) b - \frac{a}{x}$$

input `integrate((a+b*arctanh(c/x^2))/x^2,x, algorithm="maxima")`output `1/2*(c*(2*arctan(x/sqrt(c))/c^(3/2) - log((x - sqrt(c))/(x + sqrt(c)))/c^(3/2)) - 2*arctanh(c/x^2)/x)*b - a/x`**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.35

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^2} dx = -bc \left(\frac{\arctan\left(\frac{x}{\sqrt{-c}}\right)}{\sqrt{-c}} - \frac{\arctan\left(\frac{x}{\sqrt{c}}\right)}{c^{\frac{3}{2}}}\right) - \frac{b \log\left(\frac{x^2+c}{x^2-c}\right)}{2x} - \frac{a}{x}$$

input `integrate((a+b*arctanh(c/x^2))/x^2,x, algorithm="giac")`output `-b*c*(arctan(x/sqrt(-c))/(sqrt(-c)*c) - arctan(x/sqrt(c))/c^(3/2)) - 1/2*b*log((x^2 + c)/(x^2 - c))/x - a/x`

Mupad [B] (verification not implemented)

Time = 3.82 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.28

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^2} dx = \frac{b \operatorname{atan}\left(\frac{x}{\sqrt{c}}\right)}{\sqrt{c}} - \frac{a}{x} - \frac{b \ln(x^2 + c)}{2x} + \frac{b \ln(x^2 - c)}{2x} - \frac{b \operatorname{atan}\left(\frac{x}{\sqrt{c}}\right)}{\sqrt{c}}$$

input `int((a + b*atanh(c/x^2))/x^2,x)`output `(b*atan(x/c^(1/2)))/c^(1/2) - a/x - (b*atan((x*1i)/c^(1/2))*1i)/c^(1/2) - (b*log(c + x^2))/(2*x) + (b*log(x^2 - c))/(2*x)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.57

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^2} dx = \frac{2\sqrt{c} \operatorname{atan}\left(\frac{x}{\sqrt{c}}\right) bx - 2\sqrt{c} \operatorname{atanh}\left(\frac{c}{x^2}\right) bx - 2 \operatorname{atanh}\left(\frac{c}{x^2}\right) bc - 2\sqrt{c} \log(\sqrt{c} - x) bx + \sqrt{c} \log(x^2 + c) bx - 2a}{2cx}$$

input `int((a+b*atanh(c/x^2))/x^2,x)`output `(2*sqrt(c)*atan(x/sqrt(c))*b*x - 2*sqrt(c)*atanh(c/x**2)*b*x - 2*atanh(c/x**2)*b*c - 2*sqrt(c)*log(sqrt(c) - x)*b*x + sqrt(c)*log(c + x**2)*b*x - 2*a*c)/(2*c*x)`

3.169 $\int \frac{a+b\operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^4} dx$

Optimal result	1368
Mathematica [A] (verified)	1368
Rubi [A] (verified)	1369
Maple [A] (verified)	1371
Fricas [A] (verification not implemented)	1372
Sympy [B] (verification not implemented)	1372
Maxima [A] (verification not implemented)	1373
Giac [A] (verification not implemented)	1374
Mupad [B] (verification not implemented)	1374
Reduce [B] (verification not implemented)	1375

Optimal result

Integrand size = 14, antiderivative size = 65

$$\int \frac{a + b\operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^4} dx = -\frac{2b}{3cx} - \frac{b \arctan\left(\frac{x}{\sqrt{c}}\right)}{3c^{3/2}} - \frac{a + b\operatorname{arctanh}\left(\frac{c}{x^2}\right)}{3x^3} + \frac{b\operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right)}{3c^{3/2}}$$

output `-2/3*b/c/x-1/3*b*arctan(x/c^(1/2))/c^(3/2)-1/3*(a+b*arctanh(c/x^2))/x^3+1/3*b*arctanh(x/c^(1/2))/c^(3/2)`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.38

$$\int \frac{a + b\operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^4} dx = -\frac{a}{3x^3} - \frac{2b}{3cx} - \frac{b \arctan\left(\frac{x}{\sqrt{c}}\right)}{3c^{3/2}} - \frac{b\operatorname{arctanh}\left(\frac{c}{x^2}\right)}{3x^3} - \frac{b \log(\sqrt{c} - x)}{6c^{3/2}} + \frac{b \log(\sqrt{c} + x)}{6c^{3/2}}$$

input `Integrate[(a + b*ArcTanh[c/x^2])/x^4,x]`

output

```
-1/3*a/x^3 - (2*b)/(3*c*x) - (b*ArcTan[x/Sqrt[c]])/(3*c^(3/2)) - (b*ArcTan
h[c/x^2])/(3*x^3) - (b*Log[Sqrt[c] - x])/(6*c^(3/2)) + (b*Log[Sqrt[c] + x
)/(6*c^(3/2))
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.11, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6452, 795, 847, 25, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^4} dx \\
 & \quad \downarrow \text{6452} \\
 & -\frac{2}{3}bc \int \frac{1}{\left(1 - \frac{c^2}{x^4}\right)x^6} dx - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{3x^3} \\
 & \quad \downarrow \text{795} \\
 & -\frac{2}{3}bc \int \frac{1}{x^2(x^4 - c^2)} dx - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{3x^3} \\
 & \quad \downarrow \text{847} \\
 & -\frac{2}{3}bc \left(\frac{\int -\frac{x^2}{c^2 - x^4} dx}{c^2} + \frac{1}{c^2 x} \right) - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{3x^3} \\
 & \quad \downarrow \text{25} \\
 & -\frac{2}{3}bc \left(\frac{1}{c^2 x} - \frac{\int \frac{x^2}{c^2 - x^4} dx}{c^2} \right) - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{3x^3} \\
 & \quad \downarrow \text{827} \\
 & -\frac{2}{3}bc \left(\frac{1}{c^2 x} - \frac{\frac{1}{2} \int \frac{1}{c - x^2} dx - \frac{1}{2} \int \frac{1}{x^2 + c} dx}{c^2} \right) - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{3x^3} \\
 & \quad \downarrow \text{216}
 \end{aligned}$$

$$-\frac{2}{3}bc \left(\frac{1}{c^2x} - \frac{\frac{1}{2} \int \frac{1}{c-x^2} dx - \frac{\arctan\left(\frac{x}{\sqrt{c}}\right)}{2\sqrt{c}}}{c^2} \right) - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{3x^3}$$

↓ 219

$$-\frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{3x^3} - \frac{2}{3}bc \left(\frac{1}{c^2x} - \frac{\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right)}{2\sqrt{c}} - \frac{\arctan\left(\frac{x}{\sqrt{c}}\right)}{2\sqrt{c}}}{c^2} \right)$$

input `Int[(a + b*ArcTanh[c/x^2])/x^4,x]`

output `-1/3*(a + b*ArcTanh[c/x^2])/x^3 - (2*b*c*(1/(c^2*x) - (-1/2*ArcTan[x/Sqrt[c]]/Sqrt[c] + ArcTanh[x/Sqrt[c]]/(2*Sqrt[c]))/c^2))/3`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 795 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Int[x^(m + n*p)*(b + a/x^n)^p, x] /; FreeQ[{a, b, m, n}, x] && IntegerQ[p] && NegQ[n]`

rule 827 `Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x], x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 847 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + n*(p + 1) + 1)/(a*c^n*(m + 1)) Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)]*(b_))^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.85

method	result
parts	$-\frac{a}{3x^3} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{3x^3} - \frac{2b}{3cx} - \frac{b \operatorname{arctan}\left(\frac{x}{\sqrt{c}}\right)}{3c^{\frac{3}{2}}} + \frac{b \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right)}{3c^{\frac{3}{2}}}$
derivativedivides	$-\frac{a}{3x^3} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{3x^3} - \frac{2b}{3cx} + \frac{b \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right)}{3c^{\frac{3}{2}}} + \frac{b \operatorname{arctan}\left(\frac{\sqrt{c}}{x}\right)}{3c^{\frac{3}{2}}}$
default	$-\frac{a}{3x^3} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{3x^3} - \frac{2b}{3cx} + \frac{b \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right)}{3c^{\frac{3}{2}}} + \frac{b \operatorname{arctan}\left(\frac{\sqrt{c}}{x}\right)}{3c^{\frac{3}{2}}}$
risch	$-\frac{b \ln(x^2+c)}{6x^3} - \frac{4a-2ib\pi-ib\pi\operatorname{csgn}\left(\frac{i(x^2+c)}{x^2}\right)^3}{6x^3} + 2ib\pi\operatorname{csgn}\left(\frac{i(-x^2+c)}{x^2}\right)^2 + ib\pi \operatorname{csgn}\left(\frac{i}{x^2}\right) \operatorname{csgn}(i(-x^2+c)) \operatorname{csgn}\left(\frac{i}{x^2}\right)$

input `int((a+b*arctanh(c/x^2))/x^4,x,method=_RETURNVERBOSE)`

output

```
-1/3*a/x^3-1/3*b/x^3*arctanh(c/x^2)-2/3*b/c/x-1/3*b*arctan(x/c^(1/2))/c^(3/2)+1/3*b/c^(3/2)*arctanh(1/x*c^(1/2))
```

Fricas [A] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 99 vs. $2(49) = 98$.

Time = 0.11 (sec) , antiderivative size = 189, normalized size of antiderivative = 2.91

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^4} dx$$

$$= \left[\frac{2b\sqrt{cx^3} \arctan\left(\frac{x}{\sqrt{c}}\right) - b\sqrt{cx^3} \log\left(\frac{x^2+2\sqrt{cx}+c}{x^2-c}\right) + 4bcx^2 + bc^2 \log\left(\frac{x^2+c}{x^2-c}\right) + 2ac^2}{6c^2x^3}, \right.$$

$$\left. \frac{2b\sqrt{-cx^3} \arctan\left(\frac{\sqrt{-cx}}{c}\right) + b\sqrt{-cx^3} \log\left(\frac{x^2+2\sqrt{-cx}-c}{x^2+c}\right) + 4bcx^2 + bc^2 \log\left(\frac{x^2+c}{x^2-c}\right) + 2ac^2}{6c^2x^3} \right]$$

input

```
integrate((a+b*arctanh(c/x^2))/x^4,x, algorithm="fricas")
```

output

```
[-1/6*(2*b*sqrt(c)*x^3*arctan(x/sqrt(c)) - b*sqrt(c)*x^3*log((x^2 + 2*sqrt(c)*x + c)/(x^2 - c)) + 4*b*c*x^2 + b*c^2*log((x^2 + c)/(x^2 - c)) + 2*a*c^2)/(c^2*x^3), -1/6*(2*b*sqrt(-c)*x^3*arctan(sqrt(-c)*x/c) + b*sqrt(-c)*x^3*log((x^2 + 2*sqrt(-c)*x - c)/(x^2 + c)) + 4*b*c*x^2 + b*c^2*log((x^2 + c)/(x^2 - c)) + 2*a*c^2)/(c^2*x^3)]
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1046 vs. $2(58) = 116$.

Time = 4.83 (sec) , antiderivative size = 1046, normalized size of antiderivative = 16.09

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^4} dx = \text{Too large to display}$$

input

```
integrate((a+b*atanh(c/x**2))/x**4,x)
```

output

```
Piecewise((-a/(3*x**3), Eq(c, 0)), (-a - oo*b)/(3*x**3), Eq(c, -x**2)), (-a + oo*b)/(3*x**3), Eq(c, x**2)), (2*a*c**(17/2)*sqrt(-c)/(-6*c**(17/2)*x**3*sqrt(-c) + 6*c**(13/2)*x**7*sqrt(-c)) - 2*a*c**(13/2)*x**4*sqrt(-c)/(-6*c**(17/2)*x**3*sqrt(-c) + 6*c**(13/2)*x**7*sqrt(-c)) + 2*b*c**(17/2)*sqrt(-c)*atanh(c/x**2)/(-6*c**(17/2)*x**3*sqrt(-c) + 6*c**(13/2)*x**7*sqrt(-c)) + b*c**(15/2)*x**3*log(x - sqrt(-c))/(-6*c**(17/2)*x**3*sqrt(-c) + 6*c**(13/2)*x**7*sqrt(-c)) - b*c**(15/2)*x**3*log(x + sqrt(-c))/(-6*c**(17/2)*x**3*sqrt(-c) + 6*c**(13/2)*x**7*sqrt(-c)) + 4*b*c**(15/2)*x**2*sqrt(-c)/(-6*c**(17/2)*x**3*sqrt(-c) + 6*c**(13/2)*x**7*sqrt(-c)) - 2*b*c**(13/2)*x**4*sqrt(-c)*atanh(c/x**2)/(-6*c**(17/2)*x**3*sqrt(-c) + 6*c**(13/2)*x**7*sqrt(-c)) - b*c**(11/2)*x**7*log(x - sqrt(-c))/(-6*c**(17/2)*x**3*sqrt(-c) + 6*c**(13/2)*x**7*sqrt(-c)) + b*c**(11/2)*x**7*log(x + sqrt(-c))/(-6*c**(17/2)*x**3*sqrt(-c) + 6*c**(13/2)*x**7*sqrt(-c)) - 4*b*c**(11/2)*x**6*sqrt(-c)/(-6*c**(17/2)*x**3*sqrt(-c) + 6*c**(13/2)*x**7*sqrt(-c)) + 2*b*c**7*x**3*sqrt(-c)*log(-sqrt(c) + x)/(-6*c**(17/2)*x**3*sqrt(-c) + 6*c**(13/2)*x**7*sqrt(-c)) - b*c**7*x**3*sqrt(-c)*log(x - sqrt(-c))/(-6*c**(17/2)*x**3*sqrt(-c) + 6*c**(13/2)*x**7*sqrt(-c)) - b*c**7*x**3*sqrt(-c)*log(x + sqrt(-c))/(-6*c**(17/2)*x**3*sqrt(-c) + 6*c**(13/2)*x**7*sqrt(-c)) + 2*b*c**7*x**3*sqrt(-c)*atanh(c/x**2)/(-6*c**(17/2)*x**3*sqrt(-c) + 6*c**(13/2)*x**7*sqrt(-c)) - 2*b*c**5*x**7*sqrt(-c)*log(-sqrt(c) + x)/(-6*c**(17/2)*x**...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.98

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^4} dx$$

$$= -\frac{1}{6} \left(c \left(\frac{2 \arctan\left(\frac{x}{\sqrt{c}}\right)}{c^{\frac{5}{2}}} + \frac{\log\left(\frac{x-\sqrt{c}}{x+\sqrt{c}}\right)}{c^{\frac{5}{2}}} + \frac{4}{c^2 x} \right) + \frac{2 \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^3} \right) b - \frac{a}{3x^3}$$

input

```
integrate((a+b*arctanh(c/x^2))/x^4,x, algorithm="maxima")
```

output

```
-1/6*(c*(2*arctan(x/sqrt(c))/c^(5/2) + log((x - sqrt(c))/(x + sqrt(c))))/c^(5/2) + 4/(c^2*x)) + 2*arctanh(c/x^2)/x^3)*b - 1/3*a/x^3
```

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.11

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^4} dx = -\frac{b \arctan\left(\frac{x}{\sqrt{-c}}\right)}{3\sqrt{-c}} - \frac{b \arctan\left(\frac{x}{\sqrt{c}}\right)}{3c^{3/2}} - \frac{b \log\left(\frac{x^2+c}{x^2-c}\right)}{6x^3} - \frac{2bx^2 + ac}{3cx^3}$$

input `integrate((a+b*arctanh(c/x^2))/x^4,x, algorithm="giac")`

output `-1/3*b*arctan(x/sqrt(-c))/(sqrt(-c)*c) - 1/3*b*arctan(x/sqrt(c))/c^(3/2) - 1/6*b*log((x^2 + c)/(x^2 - c))/x^3 - 1/3*(2*b*x^2 + a*c)/(c*x^3)`

Mupad [B] (verification not implemented)

Time = 3.98 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.06

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^4} dx = \frac{b \ln(x^2 - c)}{6x^3} - \frac{2b}{3cx} - \frac{b \operatorname{atan}\left(\frac{x}{\sqrt{c}}\right)}{3c^{3/2}} - \frac{b \ln(x^2 + c)}{6x^3} - \frac{a}{3x^3} - \frac{b \operatorname{atan}\left(\frac{x \operatorname{li}}{\sqrt{c}}\right) \operatorname{li}}{3c^{3/2}}$$

input `int((a + b*atanh(c/x^2))/x^4,x)`

output `(b*log(x^2 - c))/(6*x^3) - (2*b)/(3*c*x) - (b*atan(x/c^(1/2)))/(3*c^(3/2)) - (b*atan((x*1i)/c^(1/2))*1i)/(3*c^(3/2)) - (b*log(c + x^2))/(6*x^3) - a/(3*x^3)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.40

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^4} dx$$

$$= \frac{-2\sqrt{c} \operatorname{atan}\left(\frac{x}{\sqrt{c}}\right) b x^3 - 2\sqrt{c} \operatorname{atanh}\left(\frac{c}{x^2}\right) b x^3 - 2 \operatorname{atanh}\left(\frac{c}{x^2}\right) b c^2 - 2\sqrt{c} \log(\sqrt{c} - x) b x^3 + \sqrt{c} \log(x^2 + c) b x^3}{6c^2 x^3}$$

input `int((a+b*atanh(c/x^2))/x^4,x)`output `(- 2*sqrt(c)*atan(x/sqrt(c))*b*x**3 - 2*sqrt(c)*atanh(c/x**2)*b*x**3 - 2*atanh(c/x**2)*b*c**2 - 2*sqrt(c)*log(sqrt(c) - x)*b*x**3 + sqrt(c)*log(c + x**2)*b*x**3 - 2*a*c**2 - 4*b*c*x**2)/(6*c**2*x**3)`

$$3.170 \quad \int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^6} dx$$

Optimal result	1376
Mathematica [A] (verified)	1376
Rubi [A] (verified)	1377
Maple [A] (verified)	1379
Fricas [A] (verification not implemented)	1380
Sympy [B] (verification not implemented)	1380
Maxima [A] (verification not implemented)	1381
Giac [A] (verification not implemented)	1382
Mupad [B] (verification not implemented)	1382
Reduce [B] (verification not implemented)	1383

Optimal result

Integrand size = 14, antiderivative size = 65

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^6} dx = -\frac{2b}{15cx^3} + \frac{b \arctan\left(\frac{x}{\sqrt{c}}\right)}{5c^{5/2}} - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{5x^5} + \frac{b \operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right)}{5c^{5/2}}$$

output

```
-2/15*b/c/x^3+1/5*b*arctan(x/c^(1/2))/c^(5/2)-1/5*(a+b*arctanh(c/x^2))/x^5
+1/5*b*arctanh(x/c^(1/2))/c^(5/2)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.38

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^6} dx = -\frac{a}{5x^5} - \frac{2b}{15cx^3} + \frac{b \arctan\left(\frac{x}{\sqrt{c}}\right)}{5c^{5/2}} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{5x^5} \\ - \frac{b \log(\sqrt{c} - x)}{10c^{5/2}} + \frac{b \log(\sqrt{c} + x)}{10c^{5/2}}$$

input

```
Integrate[(a + b*ArcTanh[c/x^2])/x^6, x]
```

output

$$-1/5*a/x^5 - (2*b)/(15*c*x^3) + (b*ArcTan[x/Sqrt[c]])/(5*c^(5/2)) - (b*ArcTanh[c/x^2])/(5*x^5) - (b*Log[Sqrt[c] - x])/(10*c^(5/2)) + (b*Log[Sqrt[c] + x])/(10*c^(5/2))$$
Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.14, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {6452, 795, 847, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^6} dx \\ & \quad \downarrow \text{6452} \\ & -\frac{2}{5}bc \int \frac{1}{\left(1 - \frac{c^2}{x^4}\right)x^8} dx - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{5x^5} \\ & \quad \downarrow \text{795} \\ & -\frac{2}{5}bc \int \frac{1}{x^4(x^4 - c^2)} dx - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{5x^5} \\ & \quad \downarrow \text{847} \\ & -\frac{2}{5}bc \left(\frac{\int \frac{1}{x^4 - c^2} dx}{c^2} + \frac{1}{3c^2x^3} \right) - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{5x^5} \\ & \quad \downarrow \text{756} \\ & -\frac{2}{5}bc \left(\frac{-\int \frac{1}{c-x^2} dx}{2c} - \frac{\int \frac{1}{x^2+c} dx}{2c} + \frac{1}{3c^2x^3} \right) - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{5x^5} \\ & \quad \downarrow \text{216} \\ & -\frac{2}{5}bc \left(\frac{-\int \frac{1}{c-x^2} dx}{2c} - \frac{\arctan\left(\frac{x}{\sqrt{c}}\right)}{2c^{3/2}} + \frac{1}{3c^2x^3} \right) - \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{5x^5} \\ & \quad \downarrow \text{219} \end{aligned}$$

$$-\frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{5x^5} - \frac{2}{5}bc \left(\frac{-\frac{\arctan\left(\frac{x}{\sqrt{c}}\right)}{2c^{3/2}} - \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right)}{2c^{3/2}}}{c^2} + \frac{1}{3c^2x^3} \right)$$

input `Int[(a + b*ArcTanh[c/x^2])/x^6,x]`

output `-1/5*(a + b*ArcTanh[c/x^2])/x^5 - (2*b*c*(1/(3*c^2*x^3) + (-1/2*ArcTan[x/Sqrt[c]]/c^(3/2) - ArcTanh[x/Sqrt[c]]/(2*c^(3/2)))/c^2))/5`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 756 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 795 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Int[x^(m + n*p)*(b + a/x^n)^p, x] /; FreeQ[{a, b, m, n}, x] && IntegerQ[p] && NegQ[n]`

rule 847 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + n*(p + 1) + 1)/(a*c^n*(m + 1)) Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.85

method	result
parts	$-\frac{a}{5x^5} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{5x^5} - \frac{2b}{15cx^3} + \frac{b \operatorname{arctan}\left(\frac{x}{\sqrt{c}}\right)}{5c^{\frac{5}{2}}} + \frac{b \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right)}{5c^{\frac{5}{2}}}$
derivativedivides	$-\frac{a}{5x^5} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{5x^5} - \frac{2b}{15cx^3} + \frac{b \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right)}{5c^{\frac{5}{2}}} - \frac{b \operatorname{arctan}\left(\frac{\sqrt{c}}{x}\right)}{5c^{\frac{5}{2}}}$
default	$-\frac{a}{5x^5} - \frac{b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{5x^5} - \frac{2b}{15cx^3} + \frac{b \operatorname{arctanh}\left(\frac{\sqrt{c}}{x}\right)}{5c^{\frac{5}{2}}} - \frac{b \operatorname{arctan}\left(\frac{\sqrt{c}}{x}\right)}{5c^{\frac{5}{2}}}$
risch	$-\frac{b \ln(x^2+c)}{10x^5} - \frac{4a-2ib\pi-ib\pi\operatorname{csgn}\left(\frac{i(x^2+c)}{x^2}\right)^3}{10x^5} + 2ib\pi\operatorname{csgn}\left(\frac{i(-x^2+c)}{x^2}\right)^2 + ib\pi \operatorname{csgn}\left(\frac{i}{x^2}\right) \operatorname{csgn}(i(-x^2+c)) \operatorname{csgn}\left(\frac{i}{x^2}\right)$

input

```
int((a+b*arctanh(c/x^2))/x^6,x,method=_RETURNVERBOSE)
```

output

```
-1/5*a/x^5-1/5*b/x^5*arctanh(c/x^2)-2/15*b/c/x^3+1/5*b*arctan(x/c^(1/2))/c
^(5/2)+1/5*b/c^(5/2)*arctanh(1/x*c^(1/2))
```


Fricas [A] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 103 vs. $2(49) = 98$.

Time = 0.09 (sec) , antiderivative size = 196, normalized size of antiderivative = 3.02

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^6} dx$$

$$= \left[\frac{6b\sqrt{cx^5} \arctan\left(\frac{x}{\sqrt{c}}\right) + 3b\sqrt{cx^5} \log\left(\frac{x^2+2\sqrt{cx}+c}{x^2-c}\right) - 4bc^2x^2 - 3bc^3 \log\left(\frac{x^2+c}{x^2-c}\right) - 6ac^3}{30c^3x^5}, \right.$$

$$\left. - \frac{6b\sqrt{-cx^5} \arctan\left(\frac{\sqrt{-cx}}{c}\right) + 3b\sqrt{-cx^5} \log\left(\frac{x^2-2\sqrt{-cx}-c}{x^2+c}\right) + 4bc^2x^2 + 3bc^3 \log\left(\frac{x^2+c}{x^2-c}\right) + 6ac^3}{30c^3x^5} \right]$$

input `integrate((a+b*arctanh(c/x^2))/x^6,x, algorithm="fricas")`

output `[1/30*(6*b*sqrt(c)*x^5*arctan(x/sqrt(c)) + 3*b*sqrt(c)*x^5*log((x^2 + 2*sqrt(c)*x + c)/(x^2 - c)) - 4*b*c^2*x^2 - 3*b*c^3*log((x^2 + c)/(x^2 - c)) - 6*a*c^3)/(c^3*x^5), -1/30*(6*b*sqrt(-c)*x^5*arctan(sqrt(-c)*x/c) + 3*b*sqrt(-c)*x^5*log((x^2 - 2*sqrt(-c)*x - c)/(x^2 + c)) + 4*b*c^2*x^2 + 3*b*c^3*log((x^2 + c)/(x^2 - c)) + 6*a*c^3)/(c^3*x^5)]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 994 vs. $2(60) = 120$.

Time = 6.72 (sec) , antiderivative size = 994, normalized size of antiderivative = 15.29

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^6} dx = \text{Too large to display}$$

input `integrate((a+b*atanh(c/x**2))/x**6,x)`

output

```
Piecewise((-a/(5*x**5), Eq(c, 0)), (-a - oo*b)/(5*x**5), Eq(c, -x**2)), (-a + oo*b)/(5*x**5), Eq(c, x**2)), (6*a*c**13*sqrt(-c)/(-30*c**13*x**5*sqrt(-c) + 30*c**11*x**9*sqrt(-c)) - 6*a*c**11*x**4*sqrt(-c)/(-30*c**13*x**5*sqrt(-c) + 30*c**11*x**9*sqrt(-c)) + 6*b*c**(21/2)*x**5*sqrt(-c)*log(-sqrt(c) + x)/(-30*c**13*x**5*sqrt(-c) + 30*c**11*x**9*sqrt(-c)) - 3*b*c**(21/2)*x**5*sqrt(-c)*log(x - sqrt(-c))/(-30*c**13*x**5*sqrt(-c) + 30*c**11*x**9*sqrt(-c)) - 3*b*c**(21/2)*x**5*sqrt(-c)*log(x + sqrt(-c))/(-30*c**13*x**5*sqrt(-c) + 30*c**11*x**9*sqrt(-c)) + 6*b*c**(21/2)*x**5*sqrt(-c)*atanh(c/x**2)/(-30*c**13*x**5*sqrt(-c) + 30*c**11*x**9*sqrt(-c)) - 6*b*c**(17/2)*x**9*sqrt(-c)*log(-sqrt(c) + x)/(-30*c**13*x**5*sqrt(-c) + 30*c**11*x**9*sqrt(-c)) + 3*b*c**(17/2)*x**9*sqrt(-c)*log(x - sqrt(-c))/(-30*c**13*x**5*sqrt(-c) + 30*c**11*x**9*sqrt(-c)) + 3*b*c**(17/2)*x**9*sqrt(-c)*log(x + sqrt(-c))/(-30*c**13*x**5*sqrt(-c) + 30*c**11*x**9*sqrt(-c)) - 6*b*c**(17/2)*x**9*sqrt(-c)*atanh(c/x**2)/(-30*c**13*x**5*sqrt(-c) + 30*c**11*x**9*sqrt(-c)) + 6*b*c**13*sqrt(-c)*atanh(c/x**2)/(-30*c**13*x**5*sqrt(-c) + 30*c**11*x**9*sqrt(-c)) + 4*b*c**12*x**2*sqrt(-c)/(-30*c**13*x**5*sqrt(-c) + 30*c**11*x**9*sqrt(-c)) - 3*b*c**11*x**5*log(x - sqrt(-c))/(-30*c**13*x**5*sqrt(-c) + 30*c**11*x**9*sqrt(-c)) + 3*b*c**11*x**5*log(x + sqrt(-c))/(-30*c**13*x**5*sqrt(-c) + 30*c**11*x**9*sqrt(-c)) - 6*b*c**11*x**4*sqrt(-c)*atanh(c/x**2)/(-30*c**13*x**5*sqrt(-c) + 30*c**11*x**9*sqrt(-c)) - 4*b*c**...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^6} dx$$

$$= \frac{1}{30} \left(c \left(\frac{6 \arctan\left(\frac{x}{\sqrt{c}}\right)}{c^{\frac{7}{2}}} - \frac{3 \log\left(\frac{x-\sqrt{c}}{x+\sqrt{c}}\right)}{c^{\frac{7}{2}}} - \frac{4}{c^2 x^3} \right) - \frac{6 \operatorname{artanh}\left(\frac{c}{x^2}\right)}{x^5} \right) b - \frac{a}{5 x^5}$$

input

```
integrate((a+b*arctanh(c/x^2))/x^6,x, algorithm="maxima")
```

output

```
1/30*(c*(6*arctan(x/sqrt(c))/c^(7/2) - 3*log((x - sqrt(c))/(x + sqrt(c))))/c^(7/2) - 4/(c^2*x^3)) - 6*arctanh(c/x^2)/x^5)*b - 1/5*a/x^5
```

Giac [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.14

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^6} dx = -\frac{1}{5} b \left(\frac{\arctan\left(\frac{x}{\sqrt{-c}}\right)}{\sqrt{-c}c^2} - \frac{\arctan\left(\frac{x}{\sqrt{c}}\right)}{c^{\frac{5}{2}}}\right) - \frac{b \log\left(\frac{x^2+c}{x^2-c}\right)}{10x^5} - \frac{2bx^2 + 3ac}{15cx^5}$$

input `integrate((a+b*arctanh(c/x^2))/x^6,x, algorithm="giac")`output `-1/5*b*(arctan(x/sqrt(-c))/(sqrt(-c)*c^2) - arctan(x/sqrt(c))/c^(5/2)) - 1/10*b*log((x^2 + c)/(x^2 - c))/x^5 - 1/15*(2*b*x^2 + 3*a*c)/(c*x^5)`**Mupad [B] (verification not implemented)**

Time = 3.92 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.06

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^6} dx = \frac{b \operatorname{atan}\left(\frac{x}{\sqrt{c}}\right)}{5c^{5/2}} - \frac{2b}{15cx^3} - \frac{a}{5x^5} - \frac{b \ln(x^2 + c)}{10x^5} + \frac{b \ln(x^2 - c)}{10x^5} - \frac{b \operatorname{atan}\left(\frac{x1i}{\sqrt{c}}\right) 1i}{5c^{5/2}}$$

input `int((a + b*atanh(c/x^2))/x^6,x)`output `(b*atan(x/c^(1/2)))/(5*c^(5/2)) - (2*b)/(15*c*x^3) - a/(5*x^5) - (b*atan((x*1i)/c^(1/2))*1i)/(5*c^(5/2)) - (b*log(c + x^2))/(10*x^5) + (b*log(x^2 - c))/(10*x^5)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.45

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{x^6} dx$$

$$= \frac{6\sqrt{c} \operatorname{atan}\left(\frac{x}{\sqrt{c}}\right) b x^5 - 6\sqrt{c} \operatorname{atanh}\left(\frac{c}{x^2}\right) b x^5 - 6 \operatorname{atanh}\left(\frac{c}{x^2}\right) b c^3 - 6\sqrt{c} \log(\sqrt{c} - x) b x^5 + 3\sqrt{c} \log(x^2 + c) b}{30c^3x^5}$$

input `int((a+b*atanh(c/x^2))/x^6,x)`output `(6*sqrt(c)*atan(x/sqrt(c))*b*x**5 - 6*sqrt(c)*atanh(c/x**2)*b*x**5 - 6*atanh(c/x**2)*b*c**3 - 6*sqrt(c)*log(sqrt(c) - x)*b*x**5 + 3*sqrt(c)*log(c + x**2)*b*x**5 - 6*a*c**3 - 4*b*c**2*x**2)/(30*c**3*x**5)`

3.171 $\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx$

Optimal result	1384
Mathematica [A] (verified)	1385
Rubi [A] (verified)	1385
Maple [C] (warning: unable to verify)	1388
Fricas [A] (verification not implemented)	1389
Sympy [A] (verification not implemented)	1390
Maxima [A] (verification not implemented)	1390
Giac [B] (verification not implemented)	1391
Mupad [B] (verification not implemented)	1392
Reduce [B] (verification not implemented)	1392

Optimal result

Integrand size = 16, antiderivative size = 94

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \frac{1}{2} b c x^2 \left(a + b \operatorname{coth}^{-1} \left(\frac{x^2}{c} \right) \right) - \frac{1}{4} c^2 \left(a + b \operatorname{coth}^{-1} \left(\frac{x^2}{c} \right) \right)^2 + \frac{1}{4} x^4 \left(a + b \operatorname{coth}^{-1} \left(\frac{x^2}{c} \right) \right)^2 + \frac{1}{4} b^2 c^2 \log \left(1 - \frac{c^2}{x^4} \right) + b^2 c^2 \log(x)$$

output

```
1/2*b*c*x^2*(a+b*arccoth(x^2/c))-1/4*c^2*(a+b*arccoth(x^2/c))^2+1/4*x^4*(a+b*arccoth(x^2/c))^2+1/4*b^2*c^2*ln(1-c^2/x^4)+b^2*c^2*ln(x)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.11

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \frac{1}{4} \left(2abcx^2 + a^2x^4 + 2bx^2(bc + ax^2) \operatorname{arctanh} \left(\frac{c}{x^2} \right) + b^2(-c^2 + x^4) \operatorname{arctanh} \left(\frac{c}{x^2} \right)^2 + b(a+b)c^2 \log(-c+x^2) - abc^2 \log(c+x^2) + b^2c^2 \log(c+x^2) \right)$$

input

```
Integrate[x^3*(a + b*ArcTanh[c/x^2])^2,x]
```

output

```
(2*a*b*c*x^2 + a^2*x^4 + 2*b*x^2*(b*c + a*x^2)*ArcTanh[c/x^2] + b^2*(-c^2 + x^4)*ArcTanh[c/x^2]^2 + b*(a + b)*c^2*Log[-c + x^2] - a*b*c^2*Log[c + x^2] + b^2*c^2*Log[c + x^2])/4
```

Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.94, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.562$, Rules used = {6454, 6452, 6544, 6452, 243, 47, 14, 16, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx \\ & \quad \downarrow \text{6454} \\ & -\frac{1}{2} \int x^6 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 d \frac{1}{x^2} \\ & \quad \downarrow \text{6452} \\ & \frac{1}{2} \left(\frac{1}{2} x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 - bc \int \frac{x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)}{1 - \frac{c^2}{x^4}} d \frac{1}{x^2} \right) \\ & \quad \downarrow \text{6544} \end{aligned}$$

$$\frac{1}{2} \left(\frac{1}{2} x^4 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right)^2 - bc \left(c^2 \int \frac{a + \operatorname{barctanh} \left(\frac{c}{x^2} \right)}{1 - \frac{c^2}{x^4}} d \frac{1}{x^2} + \int x^4 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) d \frac{1}{x^2} \right) \right)$$

↓ 6452

$$\frac{1}{2} \left(\frac{1}{2} x^4 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right)^2 - bc \left(c^2 \int \frac{a + \operatorname{barctanh} \left(\frac{c}{x^2} \right)}{1 - \frac{c^2}{x^4}} d \frac{1}{x^2} + bc \int \frac{x^2}{1 - \frac{c^2}{x^4}} d \frac{1}{x^2} - x^2 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) \right) \right)$$

↓ 243

$$\frac{1}{2} \left(\frac{1}{2} x^4 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right)^2 - bc \left(c^2 \int \frac{a + \operatorname{barctanh} \left(\frac{c}{x^2} \right)}{1 - \frac{c^2}{x^4}} d \frac{1}{x^2} + \frac{1}{2} bc \int \frac{x^2}{1 - \frac{c^2}{x^4}} d \frac{1}{x^4} - x^2 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) \right) \right)$$

↓ 47

$$\frac{1}{2} \left(\frac{1}{2} x^4 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right)^2 - bc \left(c^2 \int \frac{a + \operatorname{barctanh} \left(\frac{c}{x^2} \right)}{1 - \frac{c^2}{x^4}} d \frac{1}{x^2} + \frac{1}{2} bc \left(c^2 \int \frac{1}{1 - \frac{c^2}{x^4}} d \frac{1}{x^4} + \int x^2 d \frac{1}{x^4} \right) - x^2 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) \right) \right)$$

↓ 14

$$\frac{1}{2} \left(\frac{1}{2} x^4 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right)^2 - bc \left(c^2 \int \frac{a + \operatorname{barctanh} \left(\frac{c}{x^2} \right)}{1 - \frac{c^2}{x^4}} d \frac{1}{x^2} + \frac{1}{2} bc \left(c^2 \int \frac{1}{1 - \frac{c^2}{x^4}} d \frac{1}{x^4} + \log \left(\frac{1}{x^4} \right) \right) - x^2 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) \right) \right)$$

↓ 16

$$\frac{1}{2} \left(\frac{1}{2} x^4 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right)^2 - bc \left(c^2 \int \frac{a + \operatorname{barctanh} \left(\frac{c}{x^2} \right)}{1 - \frac{c^2}{x^4}} d \frac{1}{x^2} - x^2 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) + \frac{1}{2} bc \left(\log \left(\frac{1}{x^4} \right) - \log \left(\frac{1}{x^4} \right) \right) \right) \right)$$

↓ 6510

$$\frac{1}{2} \left(\frac{1}{2} x^4 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right)^2 - bc \left(- \left(x^2 \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right) \right) + \frac{c \left(a + \operatorname{barctanh} \left(\frac{c}{x^2} \right) \right)^2}{2b} + \frac{1}{2} bc \left(\log \left(\frac{1}{x^4} \right) - \log \left(\frac{1}{x^4} \right) \right) \right) \right)$$

input `Int[x^3*(a + b*ArcTanh[c/x^2])^2,x]`

output
$$\frac{((x^4*(a + b*\text{ArcTanh}[c/x^2])^2)/2 - b*c*(-(x^2*(a + b*\text{ArcTanh}[c/x^2]))) + (c*(a + b*\text{ArcTanh}[c/x^2])^2)/(2*b) + (b*c*(-\text{Log}[1 - c^2/x^4] + \text{Log}[x^{(-4)}]))/2)/2}$$

Defintions of rubi rules used

rule 14
$$\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ /; FreeQ}[a, x]$$

rule 16
$$\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ /; FreeQ}[\{a, b, c\}, x]$$

rule 47
$$\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ /; FreeQ}[\{a, b, c, d\}, x]$$

rule 243
$$\text{Int}[(x_)^{(m_)}*((a_)+(b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{((m-1)/2)*(a + b*x)^p}, x], x, x^2], x] \text{ /; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$$

rule 6452
$$\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{(2*n)})}), x], x] \text{ /; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$$

rule 6454
$$\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n] - 1)*(a + b*\text{ArcTanh}[c*x])^p}, x], x, x^n], x] \text{ /; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$$

rule 6510

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol]
:= Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x]
&& EqQ[c^2*d + e, 0] && NeQ[p, -1]
```

rule 6544

```
Int((((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*((f_.)*(x_)^(m_)))/((d_) + (e_.)*(x_)^2), x_Symbol]
:= Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.27 (sec) , antiderivative size = 805, normalized size of antiderivative = 8.56

Expression too large to display

input

```
int(x^3*(a+b*arctanh(c/x^2))^2,x)
```

output

```

1/4*a^2*x^4-b^2*(-1/4*x^4*arctanh(c/x^2)^2+c*(-1/4*arctanh(c/x^2)*c*ln(c/x
^2-1)-1/2*x^2*arctanh(c/x^2)+1/4*arctanh(c/x^2)*c*ln(1+c/x^2)-1/2*c*(c*(Su
m(1/4*(ln(1/x-_alpha)*ln(c/x^2-1)-2*c*(1/4/_alpha/c*ln(1/x-_alpha)^2-1/2*_
alpha*ln(1/x-_alpha)*ln(1/2*(1/x+_alpha)/_alpha)-1/2*_alpha*dilog(1/2*(1/x
+_alpha)/_alpha)))/c,_alpha=RootOf(_Z^2*c-1))+Sum(-1/4*(ln(1/x-_alpha)*ln(
c/x^2-1)-2*c*(1/2*ln(1/x-_alpha)*(ln((RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=
1)-1/x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=1))+ln((RootOf(_Z^2*c+2
*_Z*_alpha*c-2,index=2)-1/x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=2)
))/c+1/2*(dilog((RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=1)-1/x+_alpha)/RootOf
(_Z^2*c+2*_Z*_alpha*c-2,index=1))+dilog((RootOf(_Z^2*c+2*_Z*_alpha*c-2,ind
ex=2)-1/x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=2)))/c))/c,_alpha=Ro
otOf(_Z^2*c+1))-2*ln(1/x)+1/2*ln(c/x^2-1)+1/2*ln(1+c/x^2)-c*(Sum(1/4*(ln(
1/x-_alpha)*ln(1+c/x^2)-2*c*(1/2*ln(1/x-_alpha)*(ln((RootOf(_Z^2*c+2*_Z*_a
lpha*c+2,index=1)-1/x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c+2,index=1))+ln((
RootOf(_Z^2*c+2*_Z*_alpha*c+2,index=2)-1/x+_alpha)/RootOf(_Z^2*c+2*_Z*_alp
ha*c+2,index=2)))/c+1/2*(dilog((RootOf(_Z^2*c+2*_Z*_alpha*c+2,index=1)-1/x
+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c+2,index=1))+dilog((RootOf(_Z^2*c+2*_Z
*_alpha*c+2,index=2)-1/x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c+2,index=2)))/
c))/c,_alpha=RootOf(_Z^2*c-1))+Sum(-1/4*(ln(1/x-_alpha)*ln(1+c/x^2)-2*c*(1
/4/_alpha/c*ln(1/x-_alpha)^2+1/2*_alpha*ln(1/x-_alpha)*ln(1/2*(1/x+_alp...

```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.34

$$\begin{aligned}
\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx &= \frac{1}{4} a^2 x^4 + \frac{1}{2} abc x^2 - \frac{1}{4} (ab - b^2) c^2 \log(x^2 + c) \\
&+ \frac{1}{4} (ab + b^2) c^2 \log(x^2 - c) \\
&+ \frac{1}{16} (b^2 x^4 - b^2 c^2) \log \left(\frac{x^2 + c}{x^2 - c} \right)^2 \\
&+ \frac{1}{4} (abx^4 + b^2 cx^2) \log \left(\frac{x^2 + c}{x^2 - c} \right)
\end{aligned}$$

input

```
integrate(x^3*(a+b*arctanh(c/x^2))^2,x, algorithm="fricas")
```

output

```
1/4*a^2*x^4 + 1/2*a*b*c*x^2 - 1/4*(a*b - b^2)*c^2*log(x^2 + c) + 1/4*(a*b
+ b^2)*c^2*log(x^2 - c) + 1/16*(b^2*x^4 - b^2*c^2)*log((x^2 + c)/(x^2 - c)
)^2 + 1/4*(a*b*x^4 + b^2*c*x^2)*log((x^2 + c)/(x^2 - c))
```

Sympy [A] (verification not implemented)

Time = 2.10 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.61

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \frac{a^2 x^4}{4} - \frac{abc^2 \operatorname{atanh} \left(\frac{c}{x^2} \right)}{2} + \frac{abcx^2}{2} + \frac{abx^4 \operatorname{atanh} \left(\frac{c}{x^2} \right)}{2} + \frac{b^2 c^2 \log(x - \sqrt{-c})}{2} + \frac{b^2 c^2 \log(x + \sqrt{-c})}{2} - \frac{b^2 c^2 \operatorname{atanh}^2 \left(\frac{c}{x^2} \right)}{4} - \frac{b^2 c^2 \operatorname{atanh} \left(\frac{c}{x^2} \right)}{2} + \frac{b^2 cx^2 \operatorname{atanh} \left(\frac{c}{x^2} \right)}{2} + \frac{b^2 x^4 \operatorname{atanh}^2 \left(\frac{c}{x^2} \right)}{4}$$

input

```
integrate(x**3*(a+b*atanh(c/x**2))**2,x)
```

output

```
a**2*x**4/4 - a*b*c**2*atanh(c/x**2)/2 + a*b*c*x**2/2 + a*b*x**4*atanh(c/x
**2)/2 + b**2*c**2*log(x - sqrt(-c))/2 + b**2*c**2*log(x + sqrt(-c))/2 - b
**2*c**2*atanh(c/x**2)**2/4 - b**2*c**2*atanh(c/x**2)/2 + b**2*c*x**2*atan
h(c/x**2)/2 + b**2*x**4*atanh(c/x**2)**2/4
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.67

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \frac{1}{4} b^2 x^4 \operatorname{artanh} \left(\frac{c}{x^2} \right)^2 + \frac{1}{4} a^2 x^4 + \frac{1}{4} \left(2x^4 \operatorname{artanh} \left(\frac{c}{x^2} \right) + (2x^2 - c \log(x^2 + c) + c \log(x^2 - c))c \right) ab + \frac{1}{16} \left((\log(x^2 + c))^2 - 2(\log(x^2 + c) - 2) \log(x^2 - c) + \log(x^2 - c)^2 + 4 \log(x^2 + c) \right) c^2 + 4(2x^2 - c) \log(x^2 + c) c$$

input

```
integrate(x^3*(a+b*arctanh(c/x^2))^2,x, algorithm="maxima")
```

output

```
1/4*b^2*x^4*arctanh(c/x^2)^2 + 1/4*a^2*x^4 + 1/4*(2*x^4*arctanh(c/x^2) + (
2*x^2 - c*log(x^2 + c) + c*log(x^2 - c))*c)*a*b + 1/16*((log(x^2 + c)^2 -
2*(log(x^2 + c) - 2)*log(x^2 - c) + log(x^2 - c)^2 + 4*log(x^2 + c))*c^2 +
4*(2*x^2 - c*log(x^2 + c) + c*log(x^2 - c))*c*arctanh(c/x^2))*b^2
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 327 vs. $2(86) = 172$.

Time = 0.14 (sec) , antiderivative size = 327, normalized size of antiderivative = 3.48

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx =$$

$$\frac{2b^2c^3 \log\left(\frac{x^2+c}{x^2-c} - 1\right) - 2b^2c^3 \log\left(\frac{x^2+c}{x^2-c}\right) - \frac{(x^2+c)b^2c^3 \log\left(\frac{x^2+c}{x^2-c}\right)^2}{(x^2-c)\left(\frac{(x^2+c)^2}{(x^2-c)^2} - \frac{2(x^2+c)}{x^2-c} + 1\right)} - \frac{2\left(\frac{2(x^2+c)abc^3}{x^2-c} + \frac{(x^2+c)b^2c^3}{x^2-c} - b^2c^3\right) \log\left(\frac{x^2+c}{x^2-c}\right)}{\frac{(x^2+c)^2}{(x^2-c)^2} - \frac{2(x^2+c)}{x^2-c} + 1}}{4c}$$

input

```
integrate(x^3*(a+b*arctanh(c/x^2))^2,x, algorithm="giac")
```

output

```
-1/4*(2*b^2*c^3*log((x^2 + c)/(x^2 - c) - 1) - 2*b^2*c^3*log((x^2 + c)/(x^2
- c)) - (x^2 + c)*b^2*c^3*log((x^2 + c)/(x^2 - c))^2/((x^2 - c)*((x^2 +
c)^2/(x^2 - c)^2 - 2*(x^2 + c)/(x^2 - c) + 1)) - 2*(2*(x^2 + c)*a*b*c^3/(x
^2 - c) + (x^2 + c)*b^2*c^3/(x^2 - c) - b^2*c^3)*log((x^2 + c)/(x^2 - c))/
((x^2 + c)^2/(x^2 - c)^2 - 2*(x^2 + c)/(x^2 - c) + 1) - 4*((x^2 + c)*a^2*c
^3/(x^2 - c) + (x^2 + c)*a*b*c^3/(x^2 - c) - a*b*c^3)/((x^2 + c)^2/(x^2 -
c)^2 - 2*(x^2 + c)/(x^2 - c) + 1))/c
```

Mupad [B] (verification not implemented)

Time = 4.07 (sec) , antiderivative size = 247, normalized size of antiderivative = 2.63

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \frac{a^2 x^4}{4} - \frac{a b c^2 \ln(x^2 + c)}{4} + \frac{a b c^2 \ln(x^2 - c)}{4} + \frac{a b c x^2}{2} + \frac{a b x^4 \ln(x^2 + c)}{4} - \frac{a b x^4 \ln(x^2 - c)}{4} - \frac{b^2 c^2 \ln(x^2 + c)^2}{16} + \frac{b^2 c^2 \ln(x^2 + c) \ln(x^2 - c)}{8} + \frac{b^2 c^2 \ln(x^2 + c)}{4} - \frac{b^2 c^2 \ln(x^2 - c)^2}{16} + \frac{b^2 c^2 \ln(x^2 - c)}{4} + \frac{b^2 c x^2 \ln(x^2 + c)}{4} - \frac{b^2 c x^2 \ln(x^2 - c)}{4} + \frac{b^2 x^4 \ln(x^2 + c)^2}{16} - \frac{b^2 x^4 \ln(x^2 + c) \ln(x^2 - c)}{8} + \frac{b^2 x^4 \ln(x^2 - c)^2}{16}$$

input `int(x^3*(a + b*atanh(c/x^2))^2,x)`output $(a^2 x^4)/4 + (b^2 c^2 \log(x^2 - c))/4 - (b^2 c^2 \log(c + x^2)^2)/16 + (b^2 x^4 \log(c + x^2)^2)/16 - (b^2 c^2 \log(x^2 - c)^2)/16 + (b^2 x^4 \log(x^2 - c)^2)/16 + (b^2 c^2 \log(c + x^2))/4 + (a b x^4 \log(c + x^2))/4 + (a b c^2 \log(x^2 - c))/4 + (b^2 c^2 \log(c + x^2) \log(x^2 - c))/8 + (a b c x^2)/2 - (a b x^4 \log(x^2 - c))/4 + (b^2 c x^2 \log(c + x^2))/4 - (b^2 x^4 \log(c + x^2) \log(x^2 - c))/8 - (b^2 c x^2 \log(x^2 - c))/4 - (a b c^2 \log(c + x^2))/4$ **Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.26

$$\int x^3 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = -\frac{\operatorname{atanh} \left(\frac{c}{x^2} \right)^2 b^2 c^2}{4} + \frac{\operatorname{atanh} \left(\frac{c}{x^2} \right)^2 b^2 x^4}{4} - \frac{\operatorname{atanh} \left(\frac{c}{x^2} \right) a b c^2}{2} + \frac{\operatorname{atanh} \left(\frac{c}{x^2} \right) a b x^4}{2} - \frac{\operatorname{atanh} \left(\frac{c}{x^2} \right) b^2 c^2}{2} + \frac{\operatorname{atanh} \left(\frac{c}{x^2} \right) b^2 c x^2}{2} + \frac{\log(x^2 + c) b^2 c^2}{2} + \frac{a^2 x^4}{4} + \frac{a b c x^2}{2}$$

input `int(x^3*(a+b*atanh(c/x^2))^2,x)`

output `(- atanh(c/x**2)**2*b**2*c**2 + atanh(c/x**2)**2*b**2*x**4 - 2*atanh(c/x**2)*a*b*c**2 + 2*atanh(c/x**2)*a*b*x**4 - 2*atanh(c/x**2)*b**2*c**2 + 2*atanh(c/x**2)*b**2*c*x**2 + 2*log(c + x**2)*b**2*c**2 + a**2*x**4 + 2*a*b*c*x**2)/4`

3.172 $\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx$

Optimal result	1394
Mathematica [A] (verified)	1395
Rubi [A] (verified)	1395
Maple [C] (warning: unable to verify)	1397
Fricas [F]	1398
Sympy [F]	1399
Maxima [F]	1399
Giac [F]	1399
Mupad [F(-1)]	1400
Reduce [F]	1400

Optimal result

Integrand size = 14, antiderivative size = 94

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \frac{1}{2}c \left(a + b \operatorname{coth}^{-1} \left(\frac{x^2}{c} \right) \right)^2 + \frac{1}{2}x^2 \left(a + b \operatorname{coth}^{-1} \left(\frac{x^2}{c} \right) \right)^2 - bc \left(a + b \operatorname{coth}^{-1} \left(\frac{x^2}{c} \right) \right) \log \left(\frac{2c}{c - x^2} \right) - \frac{1}{2}b^2c \operatorname{PolyLog} \left(2, 1 - \frac{2c}{c - x^2} \right)$$

output

```
1/2*c*(a+b*arccoth(x^2/c))^2+1/2*x^2*(a+b*arccoth(x^2/c))^2-b*c*(a+b*arcco
th(x^2/c))*ln(2*c/(-x^2+c))-1/2*b^2*c*polylog(2,1-2*c/(-x^2+c))
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.14

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \frac{1}{2} \left(b^2 (-c + x^2) \operatorname{arctanh} \left(\frac{c}{x^2} \right)^2 \right. \\ \left. + 2b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \left(ax^2 - bc \log \left(1 - e^{-2 \operatorname{arctanh} \left(\frac{c}{x^2} \right)} \right) \right) \right. \\ \left. + a \left(ax^2 + bc \log \left(1 - \frac{c^2}{x^4} \right) - 2bc \log \left(\frac{c}{x^2} \right) \right) \right. \\ \left. + b^2 c \operatorname{PolyLog} \left(2, e^{-2 \operatorname{arctanh} \left(\frac{c}{x^2} \right)} \right) \right)$$

input `Integrate[x*(a + b*ArcTanh[c/x^2])^2,x]`

output `(b^2*(-c + x^2)*ArcTanh[c/x^2]^2 + 2*b*ArcTanh[c/x^2]*(a*x^2 - b*c*Log[1 - E^(-2*ArcTanh[c/x^2])]) + a*(a*x^2 + b*c*Log[1 - c^2/x^4] - 2*b*c*Log[c/x^2]) + b^2*c*PolyLog[2, E^(-2*ArcTanh[c/x^2])])/2`

Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.96, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {6454, 6452, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx \\ \downarrow 6454 \\ -\frac{1}{2} \int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 d \frac{1}{x^2} \\ \downarrow 6452 \\ \frac{1}{2} \left(x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 - 2bc \int \frac{x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)}{1 - \frac{c^2}{x^4}} d \frac{1}{x^2} \right)$$

$$\frac{1}{2} \left(x^2 \left(a + \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 - 2bc \left(\int \frac{x^2 \left(a + \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)}{\frac{c}{x^2} + 1} d \frac{1}{x^2} + \frac{\left(a + \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2}{2b} \right) \right)$$

↓ 6494

$$\frac{1}{2} \left(x^2 \left(a + \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 - 2bc \left(-bc \int \frac{\log \left(2 - \frac{2}{\frac{c}{x^2} + 1} \right)}{1 - \frac{c^2}{x^4}} d \frac{1}{x^2} + \frac{\left(a + \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2}{2b} + \log \left(2 - \frac{2}{\frac{c}{x^2} + 1} \right) \left(a + \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) \right) \right)$$

↓ 2897

$$\frac{1}{2} \left(x^2 \left(a + \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 - 2bc \left(\frac{\left(a + \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2}{2b} + \log \left(2 - \frac{2}{\frac{c}{x^2} + 1} \right) \left(a + \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) - \frac{1}{2} b \operatorname{PolyLog} \left(2, -1 + \frac{2}{1 + \frac{c}{x^2}} \right) \right) \right)$$

input `Int[x*(a + b*ArcTanh[c/x^2])^2,x]`

output `(x^2*(a + b*ArcTanh[c/x^2])^2 - 2*b*c*((a + b*ArcTanh[c/x^2])^2/(2*b) + (a + b*ArcTanh[c/x^2])*Log[2 - 2/(1 + c/x^2)] - (b*PolyLog[2, -1 + 2/(1 + c/x^2)])/2))/2`

Defintions of rubi rules used

rule 2897 `Int[Log[u]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x
], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simpl
ify[(m + 1)/n]]`

rule 6494 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_))), x
_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] -
Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]
/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c
^2*d^2 - e^2, 0]`

rule 6550 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_)^2)),
x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/
d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.80 (sec) , antiderivative size = 889, normalized size of antiderivative = 9.46

method	result	size
derivativedivides	Expression too large to display	889
default	Expression too large to display	889
parts	Expression too large to display	889
risch	Expression too large to display	4795

input `int(x*(a+b*arctanh(c/x^2))^2,x,method=_RETURNVERBOSE)`

output

```

1/2*a^2*x^2-b^2*(-1/2*x^2*arctanh(c/x^2)^2+2*c*(-1/4*arctanh(c/x^2)*ln(1+c
/x^2)-1/4*arctanh(c/x^2)*ln(c/x^2-1)+ln(1/x)*arctanh(c/x^2)-1/2*c*(Sum(-1/
4*(ln(1/x-_alpha)*ln(c/x^2-1)-2*c*(1/2*ln(1/x-_alpha)*(ln((RootOf(_Z^2*c+2
*_Z*_alpha*c-2,index=1)-1/x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=1)
)+ln((RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=2)-1/x+_alpha)/RootOf(_Z^2*c+2*_
Z*_alpha*c-2,index=2))))/c+1/2*(dilog((RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=
1)-1/x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c-2,index=1))+dilog((RootOf(_Z^2*
c+2*_Z*_alpha*c-2,index=2)-1/x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c-2,index
=2)))/c)/c,_alpha=RootOf(_Z^2*c+1))+Sum(1/4*(ln(1/x-_alpha)*ln(c/x^2-1)-2
*c*(1/4/_alpha/c*ln(1/x-_alpha)^2-1/2*_alpha*ln(1/x-_alpha)*ln(1/2*(1/x+_a
lpha)/_alpha)-1/2*_alpha*dilog(1/2*(1/x+_alpha)/_alpha)))/c,_alpha=RootOf(
_Z^2*c-1))+Sum(-1/4*(ln(1/x-_alpha)*ln(1+c/x^2)-2*c*(1/4/_alpha/c*ln(1/x-_
alpha)^2+1/2*_alpha*ln(1/x-_alpha)*ln(1/2*(1/x+_alpha)/_alpha)+1/2*_alpha*
dilog(1/2*(1/x+_alpha)/_alpha)))/c,_alpha=RootOf(_Z^2*c+1))+Sum(1/4*(ln(1/
x-_alpha)*ln(1+c/x^2)-2*c*(1/2*ln(1/x-_alpha)*(ln((RootOf(_Z^2*c+2*_Z*_alp
ha*c+2,index=1)-1/x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c+2,index=1))+ln((Ro
otOf(_Z^2*c+2*_Z*_alpha*c+2,index=2)-1/x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha
*c+2,index=2))))/c+1/2*(dilog((RootOf(_Z^2*c+2*_Z*_alpha*c+2,index=1)-1/x+_
alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c+2,index=1))+dilog((RootOf(_Z^2*c+2*_Z*_
alpha*c+2,index=2)-1/x+_alpha)/RootOf(_Z^2*c+2*_Z*_alpha*c+2,index=2)))...

```

Fricas [F]

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(b \operatorname{arctanh} \left(\frac{c}{x^2} \right) + a \right)^2 x dx$$

input

```
integrate(x*(a+b*arctanh(c/x^2))^2,x, algorithm="fricas")
```

output

```
integral(b^2*x*arctanh(c/x^2)^2 + 2*a*b*x*arctanh(c/x^2) + a^2*x, x)
```

Sympy [F]

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int x \left(a + b \operatorname{atanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input `integrate(x*(a+b*atanh(c/x**2))**2,x)`

output `Integral(x*(a + b*atanh(c/x**2))**2, x)`

Maxima [F]

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x^2} \right) + a \right)^2 x dx$$

input `integrate(x*(a+b*arctanh(c/x^2))^2,x, algorithm="maxima")`

output `1/2*a^2*x^2 + 1/2*(2*x^2*arctanh(c/x^2) + c*log(x^4 - c^2))*a*b + 1/8*(x^2 *log(x^2 + c)^2 - 2*(x^2 + c)*log(x^2 + c)*log(x^2 - c) + (x^2 - c)*log(x^2 - c)^2 + 2*integrate(2*(3*c*x^3 + c^2*x)*log(x^2 + c)/(x^4 - c^2), x))*b^2`

Giac [F]

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x^2} \right) + a \right)^2 x dx$$

input `integrate(x*(a+b*arctanh(c/x^2))^2,x, algorithm="giac")`

output `integrate((b*arctanh(c/x^2) + a)^2*x, x)`

Mupad [F(-1)]

Timed out.

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int x \left(a + b \operatorname{atanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input `int(x*(a + b*atanh(c/x^2))^2,x)`output `int(x*(a + b*atanh(c/x^2))^2, x)`**Reduce [F]**

$$\int x \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = -\operatorname{atanh} \left(\frac{c}{x^2} \right) abc + \operatorname{atanh} \left(\frac{c}{x^2} \right) ab x^2 + \left(\int \operatorname{atanh} \left(\frac{c}{x^2} \right)^2 x dx \right) b^2 + \log(x^2 + c) abc + \frac{a^2 x^2}{2}$$

input `int(x*(a+b*atanh(c/x^2))^2,x)`output `(- 2*atanh(c/x**2)*a*b*c + 2*atanh(c/x**2)*a*b*x**2 + 2*int(atanh(c/x**2)**2*x,x)*b**2 + 2*log(c + x**2)*a*b*c + a**2*x**2)/2`

$$3.173 \quad \int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x} dx$$

Optimal result	1401
Mathematica [C] (verified)	1402
Rubi [A] (verified)	1403
Maple [F]	1405
Fricas [F]	1405
Sympy [F]	1405
Maxima [F]	1406
Giac [F]	1406
Mupad [F(-1)]	1406
Reduce [F]	1407

Optimal result

Integrand size = 16, antiderivative size = 144

$$\begin{aligned} \int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x} dx = & -\left(a + b \operatorname{coth}^{-1}\left(\frac{x^2}{c}\right)\right)^2 \operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{c}{x^2}}\right) \\ & + \frac{1}{2}b \left(a + b \operatorname{coth}^{-1}\left(\frac{x^2}{c}\right)\right) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{c}{x^2}}\right) \\ & - \frac{1}{2}b \left(a + b \operatorname{coth}^{-1}\left(\frac{x^2}{c}\right)\right) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - \frac{c}{x^2}}\right) \\ & - \frac{1}{4}b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - \frac{c}{x^2}}\right) \\ & + \frac{1}{4}b^2 \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 - \frac{c}{x^2}}\right) \end{aligned}$$

output

```
(a+b*arccoth(x^2/c))^2*arctanh(-1+2/(1-c/x^2))+1/2*b*(a+b*arccoth(x^2/c))*
polylog(2,1-2/(1-c/x^2))-1/2*b*(a+b*arccoth(x^2/c))*polylog(2,-1+2/(1-c/x^
2))-1/4*b^2*polylog(3,1-2/(1-c/x^2))+1/4*b^2*polylog(3,-1+2/(1-c/x^2))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.27

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x} dx = a^2 \log(x) + \frac{1}{2} ab \left(\operatorname{PolyLog}\left(2, -\frac{c}{x^2}\right) - \operatorname{PolyLog}\left(2, \frac{c}{x^2}\right) \right) + \frac{1}{2} b^2 \left(-\frac{i\pi^3}{24} + \frac{2}{3} \operatorname{arctanh}\left(\frac{c}{x^2}\right)^3 + \operatorname{arctanh}\left(\frac{c}{x^2}\right)^2 \log\left(1 + e^{-2\operatorname{arctanh}\left(\frac{c}{x^2}\right)}\right) - \operatorname{arctanh}\left(\frac{c}{x^2}\right)^2 \log\left(1 - e^{2\operatorname{arctanh}\left(\frac{c}{x^2}\right)}\right) - \operatorname{arctanh}\left(\frac{c}{x^2}\right) \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arctanh}\left(\frac{c}{x^2}\right)}\right) - \operatorname{arctanh}\left(\frac{c}{x^2}\right) \operatorname{PolyLog}\left(2, e^{2\operatorname{arctanh}\left(\frac{c}{x^2}\right)}\right) - \frac{1}{2} \operatorname{PolyLog}\left(3, -e^{-2\operatorname{arctanh}\left(\frac{c}{x^2}\right)}\right) + \frac{1}{2} \operatorname{PolyLog}\left(3, e^{2\operatorname{arctanh}\left(\frac{c}{x^2}\right)}\right) \right)$$

input `Integrate[(a + b*ArcTanh[c/x^2])^2/x,x]`

output `a^2*Log[x] + (a*b*(PolyLog[2, -(c/x^2)] - PolyLog[2, c/x^2]))/2 + (b^2*((-1/24*I)*Pi^3 + (2*ArcTanh[c/x^2]^3)/3 + ArcTanh[c/x^2]^2*Log[1 + E^(-2*ArcTanh[c/x^2])] - ArcTanh[c/x^2]^2*Log[1 - E^(2*ArcTanh[c/x^2])] - ArcTanh[c/x^2]*PolyLog[2, -E^(-2*ArcTanh[c/x^2])] - ArcTanh[c/x^2]*PolyLog[2, E^(2*ArcTanh[c/x^2])] - PolyLog[3, -E^(-2*ArcTanh[c/x^2])]/2 + PolyLog[3, E^(2*ArcTanh[c/x^2])]/2))/2`

Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.13, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6450, 6448, 6614, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{barctanh}(\frac{c}{x^2}))^2}{x} dx$$

↓ 6450

$$-\frac{1}{2} \int x^2 (a + \operatorname{barctanh}(\frac{c}{x^2}))^2 d\frac{1}{x^2}$$

↓ 6448

$$\frac{1}{2} \left(4bc \int \frac{\operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{c}{x^2}}\right) (a + \operatorname{barctanh}(\frac{c}{x^2}))}{1 - \frac{c^2}{x^4}} d\frac{1}{x^2} - 2\operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{c}{x^2}}\right) (a + \operatorname{barctanh}(\frac{c}{x^2}))^2 \right)$$

↓ 6614

$$\frac{1}{2} \left(4bc \left(\frac{1}{2} \int \frac{(a + \operatorname{barctanh}(\frac{c}{x^2})) \log\left(2 - \frac{2}{1 - \frac{c}{x^2}}\right)}{1 - \frac{c^2}{x^4}} d\frac{1}{x^2} - \frac{1}{2} \int \frac{(a + \operatorname{barctanh}(\frac{c}{x^2})) \log\left(\frac{2}{1 - \frac{c}{x^2}}\right)}{1 - \frac{c^2}{x^4}} d\frac{1}{x^2} \right) - 2\operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{c}{x^2}}\right) (a + \operatorname{barctanh}(\frac{c}{x^2}))^2 \right)$$

↓ 6620

$$\frac{1}{2} \left(4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{c}{x^2}}\right) (a + \operatorname{barctanh}(\frac{c}{x^2}))}{2c} - \frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{c}{x^2}}\right)}{1 - \frac{c^2}{x^4}} d\frac{1}{x^2} \right) + \frac{1}{2} \left(\frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, \frac{2}{1 - \frac{c}{x^2}}\right)}{1 - \frac{c^2}{x^4}} d\frac{1}{x^2} \right) \right) - 2\operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{c}{x^2}}\right) (a + \operatorname{barctanh}(\frac{c}{x^2}))^2 \right)$$

↓ 7164

$$\frac{1}{2} \left(4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{c}{x^2}}\right) (a + \operatorname{barctanh}(\frac{c}{x^2}))}{2c} - \frac{b \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - \frac{c}{x^2}}\right)}{4c} \right) + \frac{1}{2} \left(\frac{b \operatorname{PolyLog}\left(3, \frac{2}{1 - \frac{c}{x^2}}\right)}{4c} \right) \right) - 2\operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{c}{x^2}}\right) (a + \operatorname{barctanh}(\frac{c}{x^2}))^2 \right)$$

input $\text{Int}[(a + b \cdot \text{ArcTanh}[c/x^2])^2/x, x]$

output $(-2 \cdot \text{ArcTanh}[1 - 2/(1 - c/x^2)] \cdot (a + b \cdot \text{ArcTanh}[c/x^2])^2 + 4 \cdot b \cdot c \cdot (((a + b \cdot \text{ArcTanh}[c/x^2]) \cdot \text{PolyLog}[2, 1 - 2/(1 - c/x^2)])/(2 \cdot c) - (b \cdot \text{PolyLog}[3, 1 - 2/(1 - c/x^2)])/(4 \cdot c))/2 + (-1/2 \cdot ((a + b \cdot \text{ArcTanh}[c/x^2]) \cdot \text{PolyLog}[2, -1 + 2/(1 - c/x^2)])/c + (b \cdot \text{PolyLog}[3, -1 + 2/(1 - c/x^2)])/(4 \cdot c))/2)/2$

Defintions of rubi rules used

rule 6448 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x)^p)/x, x_Symbol] \rightarrow \text{Simp}[2 \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot \text{ArcTanh}[1 - 2/(1 - c \cdot x)], x] - \text{Simp}[2 \cdot b \cdot c \cdot p \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{ArcTanh}[1 - 2/(1 - c \cdot x)]/(1 - c^2 \cdot x^2)), x], x] /;$
 $\text{FreeQ}\{a, b, c, x\} \ \&\& \ \text{IGtQ}[p, 1]$

rule 6450 $\text{Int}[(a + \text{ArcTanh}[c \cdot x]^n] \cdot (b \cdot x)^p/x, x_Symbol] \rightarrow \text{Simp}[1/n \cdot \text{Subst}[\text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p/x, x], x, x^n], x] /;$
 $\text{FreeQ}\{a, b, c, n\}, x\} \ \&\& \ \text{IGtQ}[p, 0]$

rule 6614 $\text{Int}[(\text{ArcTanh}[u] \cdot (a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x)^p))/((d + e \cdot x^2)), x_Symbol] \rightarrow \text{Simp}[1/2 \cdot \text{Int}[\text{Log}[1 + u] \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p/(d + e \cdot x^2)), x], x] - \text{Simp}[1/2 \cdot \text{Int}[\text{Log}[1 - u] \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p/(d + e \cdot x^2)), x], x] /;$
 $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{EqQ}[u^2 - (1 - 2/(1 - c \cdot x))^2, 0]$

rule 6620 $\text{Int}[(\text{Log}[u] \cdot (a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x)^p))/((d + e \cdot x^2)), x_Symbol] \rightarrow \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{PolyLog}[2, 1 - u]/(2 \cdot c \cdot d)), x] + \text{Simp}[b \cdot (p/2) \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{PolyLog}[2, 1 - u]/(d + e \cdot x^2)), x], x] /;$
 $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{EqQ}[(1 - u)^2 - (1 - 2/(1 - c \cdot x))^2, 0]$

rule 7164 $\text{Int}[u \cdot \text{PolyLog}[n, v], x_Symbol] \rightarrow \text{With}\{w = \text{DerivativeDivides}[v, u \cdot v, x]\}, \text{Simp}[w \cdot \text{PolyLog}[n + 1, v], x] /;$
 $! \text{FalseQ}[w] /;$
 $\text{FreeQ}[n, x]$

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x} dx$$

input `int((a+b*arctanh(c/x^2))^2/x,x)`

output `int((a+b*arctanh(c/x^2))^2/x,x)`

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x} dx = \int \frac{(b \operatorname{arctanh}(\frac{c}{x^2}) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c/x^2))^2/x,x, algorithm="fricas")`

output `integral((b^2*arctanh(c/x^2)^2 + 2*a*b*arctanh(c/x^2) + a^2)/x, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x^2}))^2}{x} dx$$

input `integrate((a+b*atanh(c/x**2))**2/x,x)`

output `Integral((a + b*atanh(c/x**2))**2/x, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x^2}) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c/x^2))^2/x,x, algorithm="maxima")`

output `a^2*log(x) + integrate(1/4*b^2*(log(c/x^2 + 1) - log(-c/x^2 + 1))^2/x + a*b*(log(c/x^2 + 1) - log(-c/x^2 + 1))/x, x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x^2}) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c/x^2))^2/x,x, algorithm="giac")`

output `integrate((b*arctanh(c/x^2) + a)^2/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x^2}))^2}{x} dx$$

input `int((a + b*atanh(c/x^2))^2/x,x)`

output `int((a + b*atanh(c/x^2))^2/x, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x} dx$$

$$= 2 \left(\int \frac{\operatorname{atanh}(\frac{c}{x^2})}{x} dx \right) ab + \left(\int \frac{\operatorname{atanh}(\frac{c}{x^2})^2}{x} dx \right) b^2 + \log(x) a^2$$

input

```
int((a+b*atanh(c/x^2))^2/x,x)
```

output

```
2*int(atanh(c/x**2)/x,x)*a*b + int(atanh(c/x**2)**2/x,x)*b**2 + log(x)*a**2
```

3.174 $\int \frac{\left(a+b\operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^3} dx$

Optimal result	1408
Mathematica [A] (verified)	1409
Rubi [A] (verified)	1409
Maple [A] (verified)	1412
Fricas [F]	1412
Sympy [F]	1413
Maxima [F]	1413
Giac [F]	1414
Mupad [F(-1)]	1414
Reduce [F]	1414

Optimal result

Integrand size = 16, antiderivative size = 99

$$\int \frac{\left(a+b\operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^3} dx = -\frac{\left(a+b\operatorname{coth}^{-1}\left(\frac{x^2}{c}\right)\right)^2}{2c} - \frac{\left(a+b\operatorname{coth}^{-1}\left(\frac{x^2}{c}\right)\right)^2}{2x^2} + \frac{b\left(a+b\operatorname{coth}^{-1}\left(\frac{x^2}{c}\right)\right)\log\left(\frac{2}{1-\frac{c}{x^2}}\right)}{c} + \frac{b^2\operatorname{PolyLog}\left(2,1-\frac{2}{1-\frac{c}{x^2}}\right)}{2c}$$

output

$-1/2*(a+b*\operatorname{arccoth}(x^2/c))^2/c-1/2*(a+b*\operatorname{arccoth}(x^2/c))^2/x^2+b*(a+b*\operatorname{arccoth}(x^2/c))*\ln(2/(1-c/x^2))/c+1/2*b^2*\operatorname{polylog}(2,1-2/(1-c/x^2))/c$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.15

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^3} dx = -\frac{a^2}{2x^2} - \frac{ab \left(\frac{\operatorname{arctanh}(\frac{c}{x^2})}{x^2} - \log \left(\frac{1}{\sqrt{1 - \frac{c^2}{x^4}}} \right) \right)}{c} - \frac{b^2 \left(\operatorname{arctanh}(\frac{c}{x^2}) \left(-\operatorname{arctanh}(\frac{c}{x^2}) + \frac{\operatorname{arctanh}(\frac{c}{x^2})}{x^2} - 2 \log \left(1 + e^{-2 \operatorname{arctanh}(\frac{c}{x^2})} \right) \right) \right)}{2c} + \operatorname{PolyLog} \left(2, -e^{-2 \operatorname{arctanh}(\frac{c}{x^2})} \right)$$

input `Integrate[(a + b*ArcTanh[c/x^2])^2/x^3, x]`

output

```
-1/2*a^2/x^2 - (a*b*((c*ArcTanh[c/x^2])/x^2 - Log[1/Sqrt[1 - c^2/x^4]]))/c
- (b^2*(ArcTanh[c/x^2]*(-ArcTanh[c/x^2] + (c*ArcTanh[c/x^2])/x^2 - 2*Log[
1 + E^(-2*ArcTanh[c/x^2])])) + PolyLog[2, -E^(-2*ArcTanh[c/x^2])])/(2*c)
```

Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6454, 6436, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^3} dx \\ & \quad \downarrow \text{6454} \\ & -\frac{1}{2} \int (a + b \operatorname{arctanh}(\frac{c}{x^2}))^2 d \frac{1}{x^2} \\ & \quad \downarrow \text{6436} \\ & \frac{1}{2} \left(2bc \int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{(1 - \frac{c^2}{x^4}) x^2} d \frac{1}{x^2} - \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^2} \right) \end{aligned}$$

$$\frac{1}{2} \left(2bc \left(\frac{\int \frac{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)}{1 - \frac{c}{x^2}} d\frac{1}{x^2}}{c} - \frac{(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right))^2}{2bc^2} \right) - \frac{(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right))^2}{x^2} \right)$$

↓ 6546

$$\frac{1}{2} \left(2bc \left(\frac{\frac{\log\left(\frac{2}{1 - \frac{c}{x^2}}\right)(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right))}{c} - b \int \frac{\log\left(\frac{2}{1 - \frac{c}{x^2}}\right)}{1 - \frac{c^2}{x^4}} d\frac{1}{x^2}}{c} - \frac{(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right))^2}{2bc^2} \right) - \frac{(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right))^2}{x^2} \right)$$

↓ 6470

$$\frac{1}{2} \left(2bc \left(\frac{b \int \frac{\log\left(\frac{2}{1 - \frac{c}{x^2}}\right)}{1 - \frac{c}{x^2}} d\frac{1}{x^2} + \frac{\log\left(\frac{2}{1 - \frac{c}{x^2}}\right)(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right))}{c}}{c} - \frac{(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right))^2}{2bc^2} \right) - \frac{(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right))^2}{x^2} \right)$$

↓ 2849

$$\frac{1}{2} \left(2bc \left(\frac{\frac{\log\left(\frac{2}{1 - \frac{c}{x^2}}\right)(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right))}{c} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{c}{x^2}}\right)}{2c}}{c} - \frac{(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right))^2}{2bc^2} \right) - \frac{(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right))^2}{x^2} \right)$$

↓ 2752

input `Int[(a + b*ArcTanh[c/x^2])^2/x^3,x]`

output `(-((a + b*ArcTanh[c/x^2])^2/x^2) + 2*b*c*(-1/2*(a + b*ArcTanh[c/x^2])^2/(b*c^2) + (((a + b*ArcTanh[c/x^2])*Log[2/(1 - c/x^2)])/c + (b*PolyLog[2, 1 - 2/(1 - c/x^2)])/(2*c))/c)/2`

Definitions of rubi rules used

rule 2752 $\text{Int}[\text{Log}[(c_)*(x_)]/((d_)+(e_)*(x_)), x_Symbol] \rightarrow \text{Simp}[(-e^{(-1)})*\text{PolyLog}[2, 1 - c*x], x] /; \text{FreeQ}[\{c, d, e\}, x] \ \&\& \ \text{EqQ}[e + c*d, 0]$

rule 2849 $\text{Int}[\text{Log}[(c_)/((d_)+(e_)*(x_))]/((f_)+(g_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[-e/g \ \text{Subst}[\text{Int}[\text{Log}[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; \text{FreeQ}[\{c, d, e, f, g\}, x] \ \&\& \ \text{EqQ}[c, 2*d] \ \&\& \ \text{EqQ}[e^2*f + d^2*g, 0]$

rule 6436 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^(n_)]*(b_))^(p_), x_Symbol] \rightarrow \text{Simp}[x*(a + b*\text{ArcTanh}[c*x^n])^p, x] - \text{Simp}[b*c*n*p \ \text{Int}[x^n*((a + b*\text{ArcTanh}[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; \text{FreeQ}[\{a, b, c, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[n, 1] \ || \ \text{EqQ}[p, 1])$

rule 6454 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] \rightarrow \text{Simp}[1/n \ \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)*(a + b*\text{ArcTanh}[c*x])^p}, x], x, x^n], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]$

rule 6470 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)])*(b_)^(p_)/((d_)+(e_)*(x_)), x_Symbol] \rightarrow \text{Simp}[(-a + b*\text{ArcTanh}[c*x])^p*(\text{Log}[2/(1 + e*(x/d))]/e), x] + \text{Simp}[b*c*(p/e) \ \text{Int}[(a + b*\text{ArcTanh}[c*x])^(p - 1)*(\text{Log}[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d^2 - e^2, 0]$

rule 6546 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)])*(b_)^(p_)*(x_)/((d_)+(e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^(p + 1)/(b*e*(p + 1)), x] + \text{Simp}[1/(c*d) \ \text{Int}[(a + b*\text{ArcTanh}[c*x])^p/(1 - c*x), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{IGtQ}[p, 0]$

Maple [A] (verified)

Time = 1.60 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.35

method	result
derivativedivides	$\frac{\frac{c a^2}{x^2} + b^2 \left(\operatorname{arctanh}\left(\frac{c}{x^2}\right)^2 \left(\frac{c}{x^2} - 1\right) + 2 \operatorname{arctanh}\left(\frac{c}{x^2}\right)^2 - 2 \operatorname{arctanh}\left(\frac{c}{x^2}\right) \ln\left(1 + \frac{\left(1 + \frac{c}{x^2}\right)^2}{1 - \frac{c^2}{x^4}}\right) - \operatorname{polylog}\left(2, -\frac{\left(1 + \frac{c}{x^2}\right)^2}{1 - \frac{c^2}{x^4}}\right)}{2c}}$
default	$\frac{\frac{c a^2}{x^2} + b^2 \left(\operatorname{arctanh}\left(\frac{c}{x^2}\right)^2 \left(\frac{c}{x^2} - 1\right) + 2 \operatorname{arctanh}\left(\frac{c}{x^2}\right)^2 - 2 \operatorname{arctanh}\left(\frac{c}{x^2}\right) \ln\left(1 + \frac{\left(1 + \frac{c}{x^2}\right)^2}{1 - \frac{c^2}{x^4}}\right) - \operatorname{polylog}\left(2, -\frac{\left(1 + \frac{c}{x^2}\right)^2}{1 - \frac{c^2}{x^4}}\right)}{2c}}$
parts	$-\frac{a^2}{2x^2} - \frac{b^2 \left(\operatorname{arctanh}\left(\frac{c}{x^2}\right)^2 \left(\frac{c}{x^2} - 1\right) + 2 \operatorname{arctanh}\left(\frac{c}{x^2}\right)^2 - 2 \operatorname{arctanh}\left(\frac{c}{x^2}\right) \ln\left(1 + \frac{\left(1 + \frac{c}{x^2}\right)^2}{1 - \frac{c^2}{x^4}}\right) - \operatorname{polylog}\left(2, -\frac{\left(1 + \frac{c}{x^2}\right)^2}{1 - \frac{c^2}{x^4}}\right)}{2c}$

input `int((a+b*arctanh(c/x^2))^2/x^3,x,method=_RETURNVERBOSE)`

output
$$-1/2/c*(c/x^2*a^2+b^2*(\operatorname{arctanh}(c/x^2)^2*(c/x^2-1)+2*\operatorname{arctanh}(c/x^2)^2-2*\operatorname{arctanh}(c/x^2)*\ln(1+(1+c/x^2)^2/(1-c^2/x^4))-\operatorname{polylog}(2,-(1+c/x^2)^2/(1-c^2/x^4))))+2*a*b*c/x^2*\operatorname{arctanh}(c/x^2)+a*b*\ln(1-c^2/x^4))$$

Fricas [F]

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^3} dx = \int \frac{\left(b \operatorname{arctanh}\left(\frac{c}{x^2}\right) + a\right)^2}{x^3} dx$$

input `integrate((a+b*arctanh(c/x^2))^2/x^3,x, algorithm="fricas")`

output `integral((b^2*arctanh(c/x^2)^2 + 2*a*b*arctanh(c/x^2) + a^2)/x^3, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^3} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x^2}))^2}{x^3} dx$$

input `integrate((a+b*atanh(c/x**2))**2/x**3,x)`

output `Integral((a + b*atanh(c/x**2))**2/x**3, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^3} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x^2}) + a)^2}{x^3} dx$$

input `integrate((a+b*arctanh(c/x^2))^2/x^3,x, algorithm="maxima")`

output `1/8*(8*c^3*integrate(log(x)^2/(c*x^7 - c^3*x^3), x) + c^2*(log(x^2 + c)/c^3 + log(x^2 - c)/c^3 - 4*log(x)/c^3) - 8*c^2*integrate(x^2*log(x^2 + c)/(c*x^7 - c^3*x^3), x) + 8*c^2*integrate(x^2*log(x)/(c*x^7 - c^3*x^3), x) + 2*c*(log(x^2 - c)/c^2 - log(x^2)/c^2 + 1/(c*x^2))*log(-c/x^2 + 1) - c*(log(x^2 + c)/c^2 - log(x^2 - c)/c^2) - 8*c*integrate(x^4*log(x)^2/(c*x^7 - c^3*x^3), x) - 4*c*integrate(x^4*log(x^2 + c)/(c*x^7 - c^3*x^3), x) + 16*c*integrate(x^4*log(x)/(c*x^7 - c^3*x^3), x) - log(-c/x^2 + 1)^2/x^2 - (x^2*log(x^2 - c)^2 + 4*x^2*log(x)^2 - 4*x^2*log(x) - 2*(2*x^2*log(x) - x^2)*log(x^2 - c) + 2*c)/(c*x^2) - (c*log(x^2 + c)^2 - 2*((x^2 + c)*log(x^2 + c) - 2*(x^2 + c)*log(x) - c)*log(x^2 - c))/(c*x^2) - 4*integrate(x^6*log(x^2 + c)/(c*x^7 - c^3*x^3), x) + 8*integrate(x^6*log(x)/(c*x^7 - c^3*x^3), x))*b^2 - 1/2*a*b*(2*c*arctanh(c/x^2)/x^2 + log(-c^2/x^4 + 1))/c - 1/2*a^2/x^2`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^3} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x^2}) + a)^2}{x^3} dx$$

input `integrate((a+b*arctanh(c/x^2))^2/x^3,x, algorithm="giac")`

output `integrate((b*arctanh(c/x^2) + a)^2/x^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^3} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x^2}))^2}{x^3} dx$$

input `int((a + b*atanh(c/x^2))^2/x^3,x)`

output `int((a + b*atanh(c/x^2))^2/x^3, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^3} dx$$

$$= \frac{-\operatorname{atanh}(\frac{c}{x^2})^2 b^2 c - 2 \operatorname{atanh}(\frac{c}{x^2}) abc + 2 \operatorname{atanh}(\frac{c}{x^2}) ab x^2 + 4 \left(\int \frac{\operatorname{atanh}(\frac{c}{x^2})}{-x^5 + c^2 x} dx \right) b^2 c^2 x^2 - 2 \log(x^2 + c) ab x^2}{2c x^2}$$

input `int((a+b*atanh(c/x^2))^2/x^3,x)`

output `(- atanh(c/x**2)**2*b**2*c - 2*atanh(c/x**2)*a*b*c + 2*atanh(c/x**2)*a*b*x**2 + 4*int(atanh(c/x**2)/(c**2*x - x**5),x)*b**2*c**2*x**2 - 2*log(c + x**2)*a*b*x**2 + 4*log(x)*a*b*x**2 - a**2*c)/(2*c*x**2)`

3.175 $\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^5} dx$

Optimal result	1415
Mathematica [A] (verified)	1415
Rubi [A] (verified)	1416
Maple [A] (verified)	1418
Fricas [A] (verification not implemented)	1418
Sympy [B] (verification not implemented)	1419
Maxima [B] (verification not implemented)	1419
Giac [F]	1420
Mupad [B] (verification not implemented)	1420
Reduce [B] (verification not implemented)	1421

Optimal result

Integrand size = 16, antiderivative size = 97

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^5} dx = -\frac{ab}{2cx^2} - \frac{b^2 \operatorname{coth}^{-1}\left(\frac{x^2}{c}\right)}{2cx^2} + \frac{\left(a + b \operatorname{coth}^{-1}\left(\frac{x^2}{c}\right)\right)^2}{4c^2} - \frac{\left(a + b \operatorname{coth}^{-1}\left(\frac{x^2}{c}\right)\right)^2}{4x^4} - \frac{b^2 \log\left(1 - \frac{c^2}{x^4}\right)}{4c^2}$$

```
output -1/2*a*b/c/x^2-1/2*b^2*arccoth(x^2/c)/c/x^2+1/4*(a+b*arccoth(x^2/c))^2/c^2
-1/4*(a+b*arccoth(x^2/c))^2/x^4-1/4*b^2*ln(1-c^2/x^4)/c^2
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.35

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^5} dx = \frac{a^2c^2 + 2abcx^2 + 2bc(ac + bx^2) \operatorname{arctanh}\left(\frac{c}{x^2}\right) + b^2(c^2 - x^4) \operatorname{arctanh}\left(\frac{c}{x^2}\right)^2 - 4b^2x^4 \log(x) + abx^4 \log(-c/x^2)}{4c^2x^4}$$

```
input Integrate[(a + b*ArcTanh[c/x^2])^2/x^5,x]
```

output

```
-1/4*(a^2*c^2 + 2*a*b*c*x^2 + 2*b*c*(a*c + b*x^2)*ArcTanh[c/x^2] + b^2*(c^2 - x^4)*ArcTanh[c/x^2]^2 - 4*b^2*x^4*Log[x] + a*b*x^4*Log[-c + x^2] + b^2*x^4*Log[-c + x^2] - a*b*x^4*Log[c + x^2] + b^2*x^4*Log[c + x^2))/(c^2*x^4)
```

Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.94, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6454, 6452, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^5} dx$$

$$\downarrow 6454$$

$$-\frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^2} d\frac{1}{x^2}$$

$$\downarrow 6452$$

$$\frac{1}{2} \left(bc \int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{(1 - \frac{c^2}{x^4}) x^4} d\frac{1}{x^2} - \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{2x^4} \right)$$

$$\downarrow 6542$$

$$\frac{1}{2} \left(bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{1 - \frac{c^2}{x^4}} d\frac{1}{x^2}}{c^2} - \frac{\int (a + b \operatorname{arctanh}(\frac{c}{x^2})) d\frac{1}{x^2}}{c^2} \right) - \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{2x^4} \right)$$

$$\downarrow 2009$$

$$\frac{1}{2} \left(bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(\frac{c}{x^2})}{1 - \frac{c^2}{x^4}} d\frac{1}{x^2}}{c^2} - \frac{\frac{a}{x^2} + \frac{b \operatorname{arctanh}(\frac{c}{x^2})}{x^2} + \frac{b \log(1 - \frac{c^2}{x^4})}{2c}}{c^2} \right) - \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{2x^4} \right)$$

$$\downarrow 6510$$

$$\frac{1}{2} \left(bc \left(\frac{(a + \operatorname{barctanh}(\frac{c}{x^2}))^2}{2bc^3} - \frac{a}{x^2} + \frac{\operatorname{barctanh}(\frac{c}{x^2})}{x^2} + \frac{b \log(1 - \frac{c^2}{x^4})}{2c} \right) - \frac{(a + \operatorname{barctanh}(\frac{c}{x^2}))^2}{2x^4} \right)$$

input `Int[(a + b*ArcTanh[c/x^2])^2/x^5,x]`

output `(-1/2*(a + b*ArcTanh[c/x^2])^2/x^4 + b*c*((a + b*ArcTanh[c/x^2])^2/(2*b*c^3) - (a/x^2 + (b*ArcTanh[c/x^2])/x^2 + (b*Log[1 - c^2/x^4])/(2*c))/c^2))/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6510 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/((d_.) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6542 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((f_.)*(x_)^(m_.))/((d_.) + (e_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

Maple [A] (verified)

Time = 244.92 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.42

method	result
paralelrisch	$\frac{b^2 \operatorname{arctanh}\left(\frac{c}{x^2}\right)^2 x^4 + 4b^2 \ln(x)x^4 - 2 \ln(x^2 - c)x^4 b^2 + 2x^4 \operatorname{arctanh}\left(\frac{c}{x^2}\right) ab - 2x^4 \operatorname{arctanh}\left(\frac{c}{x^2}\right) b^2 - 2b^2 c \operatorname{arctanh}\left(\frac{c}{x^2}\right) x^2}{4x^4 c^2}$
parts	Expression too large to display
derivativedivides	Expression too large to display
default	Expression too large to display
risch	Expression too large to display

input `int((a+b*arctanh(c/x^2))^2/x^5,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{4} * (b^2 * \operatorname{arctanh}(c/x^2)^2 * x^4 + 4 * b^2 * \ln(x) * x^4 - 2 * \ln(x^2 - c) * x^4 * b^2 + 2 * x^4 * \operatorname{arctanh}(c/x^2) * a * b - 2 * x^4 * \operatorname{arctanh}(c/x^2) * b^2 - 2 * b^2 * c * \operatorname{arctanh}(c/x^2) * x^2 - b^2 * c^2 * \operatorname{arctanh}(c/x^2)^2 - 2 * a * b * c * x^2 - 2 * a * b * c^2 * \operatorname{arctanh}(c/x^2) - a^2 * c^2) / x^4 / c^2$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.47

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^5} dx$$

$$= \frac{16 b^2 x^4 \log(x) + 4 (ab - b^2) x^4 \log(x^2 + c) - 4 (ab + b^2) x^4 \log(x^2 - c) - 8 abc x^2 - 4 a^2 c^2 + (b^2 x^4 - b^2 c^2)}{16 c^2 x^4}$$

input `integrate((a+b*arctanh(c/x^2))^2/x^5,x, algorithm="fricas")`

output
$$\frac{1}{16} * (16 * b^2 * x^4 * \log(x) + 4 * (a * b - b^2) * x^4 * \log(x^2 + c) - 4 * (a * b + b^2) * x^4 * \log(x^2 - c) - 8 * a * b * c * x^2 - 4 * a^2 * c^2 + (b^2 * x^4 - b^2 * c^2) * \log((x^2 + c) / (x^2 - c)) - 4 * (b^2 * c * x^2 + a * b * c^2) * \log((x^2 + c) / (x^2 - c))) / (c^2 * x^4)$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 172 vs. $2(78) = 156$.

Time = 5.88 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.77

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^5} dx$$

$$= \begin{cases} -\frac{a^2}{4x^4} - \frac{ab \operatorname{atanh}(\frac{c}{x^2})}{2x^4} - \frac{ab}{2cx^2} + \frac{ab \operatorname{atanh}(\frac{c}{x^2})}{2c^2} - \frac{b^2 \operatorname{atanh}^2(\frac{c}{x^2})}{4x^4} - \frac{b^2 \operatorname{atanh}(\frac{c}{x^2})}{2cx^2} + \frac{b^2 \log(x)}{c^2} - \frac{b^2 \log(x - \sqrt{-c})}{2c^2} - \frac{b^2 \log(x + \sqrt{-c})}{2c^2} \\ -\frac{a^2}{4x^4} \end{cases}$$

input `integrate((a+b*atanh(c/x**2))**2/x**5,x)`

output `Piecewise((-a**2/(4*x**4) - a*b*atanh(c/x**2)/(2*x**4) - a*b/(2*c*x**2) + a*b*atanh(c/x**2)/(2*c**2) - b**2*atanh(c/x**2)**2/(4*x**4) - b**2*atanh(c/x**2)/(2*c*x**2) + b**2*log(x)/c**2 - b**2*log(x - sqrt(-c))/(2*c**2) - b**2*log(x + sqrt(-c))/(2*c**2) + b**2*atanh(c/x**2)**2/(4*c**2) + b**2*atanh(c/x**2)/(2*c**2), Ne(c, 0)), (-a**2/(4*x**4), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 183 vs. $2(87) = 174$.

Time = 0.03 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.89

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^5} dx$$

$$= \frac{1}{4} \left(c \left(\frac{\log(x^2 + c)}{c^3} - \frac{\log(x^2 - c)}{c^3} - \frac{2}{c^2 x^2} \right) - \frac{2 \operatorname{artanh}(\frac{c}{x^2})}{x^4} \right) ab$$

$$- \frac{1}{16} \left(c^2 \left(\frac{\log(x^2 + c)^2 - 2(\log(x^2 + c) - 2) \log(x^2 - c) + \log(x^2 - c)^2 + 4 \log(x^2 + c)}{c^4} - \frac{16 \log(x)}{c^4} \right) \right)$$

$$- \frac{b^2 \operatorname{artanh}(\frac{c}{x^2})^2}{4x^4} - \frac{a^2}{4x^4}$$

input `integrate((a+b*arctanh(c/x^2))^2/x^5,x, algorithm="maxima")`

output

```
1/4*(c*(log(x^2 + c)/c^3 - log(x^2 - c)/c^3 - 2/(c^2*x^2)) - 2*arctanh(c/x^2)/x^4)*a*b - 1/16*(c^2*((log(x^2 + c)^2 - 2*(log(x^2 + c) - 2)*log(x^2 - c) + log(x^2 - c)^2 + 4*log(x^2 + c))/c^4 - 16*log(x)/c^4) - 4*c*(log(x^2 + c)/c^3 - log(x^2 - c)/c^3 - 2/(c^2*x^2))*arctanh(c/x^2))*b^2 - 1/4*b^2*arctanh(c/x^2)^2/x^4 - 1/4*a^2/x^4
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^5} dx = \int \frac{(b \operatorname{arctanh}(\frac{c}{x^2}) + a)^2}{x^5} dx$$

input

```
integrate((a+b*arctanh(c/x^2))^2/x^5,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c/x^2) + a)^2/x^5, x)
```

Mupad [B] (verification not implemented)

Time = 4.45 (sec) , antiderivative size = 262, normalized size of antiderivative = 2.70

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^5} dx &= \frac{b^2 \ln(x^2 + c)^2}{16 c^2} - \frac{b^2 \ln(x^2 - c)}{4 c^2} - \frac{a^2}{4 x^4} - \frac{b^2 \ln(x^2 + c)^2}{16 x^4} \\ &+ \frac{b^2 \ln(x^2 - c)^2}{16 c^2} - \frac{b^2 \ln(x^2 - c)^2}{16 x^4} + \frac{b^2 \ln(x)}{c^2} \\ &- \frac{b^2 \ln(x^2 + c)}{4 c^2} - \frac{a b \ln(x^2 + c)}{4 x^4} + \frac{b^2 \ln(x^2 - c)}{4 c x^2} \\ &- \frac{a b \ln(x^2 - c)}{4 c^2} - \frac{b^2 \ln(x^2 + c) \ln(x^2 - c)}{8 c^2} \\ &+ \frac{a b \ln(x^2 - c)}{4 x^4} + \frac{b^2 \ln(x^2 + c) \ln(x^2 - c)}{8 x^4} \\ &- \frac{a b}{2 c x^2} - \frac{b^2 \ln(x^2 + c)}{4 c x^2} + \frac{a b \ln(x^2 + c)}{4 c^2} \end{aligned}$$

input

```
int((a + b*atanh(c/x^2))^2/x^5,x)
```

output

```
(b^2*log(c + x^2)^2)/(16*c^2) - (b^2*log(x^2 - c))/(4*c^2) - a^2/(4*x^4) -
(b^2*log(c + x^2)^2)/(16*x^4) + (b^2*log(x^2 - c)^2)/(16*c^2) - (b^2*log(
x^2 - c)^2)/(16*x^4) + (b^2*log(x))/c^2 - (b^2*log(c + x^2))/(4*c^2) - (a*
b*log(c + x^2))/(4*x^4) + (b^2*log(x^2 - c))/(4*c*x^2) - (a*b*log(x^2 - c)
)/(4*c^2) - (b^2*log(c + x^2)*log(x^2 - c))/(8*c^2) + (a*b*log(x^2 - c))/(
4*x^4) + (b^2*log(c + x^2)*log(x^2 - c))/(8*x^4) - (a*b)/(2*c*x^2) - (b^2*
log(c + x^2))/(4*c*x^2) + (a*b*log(c + x^2))/(4*c^2)
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.39

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^5} dx$$

$$= \frac{-\operatorname{atanh}(\frac{c}{x^2})^2 b^2 c^2 + \operatorname{atanh}(\frac{c}{x^2})^2 b^2 x^4 - 2 \operatorname{atanh}(\frac{c}{x^2}) a b c^2 + 2 \operatorname{atanh}(\frac{c}{x^2}) a b x^4 - 2 \operatorname{atanh}(\frac{c}{x^2}) b^2 c x^2 + 2 a^2}{4 c^2 x^4}$$

input

```
int((a+b*atanh(c/x^2))^2/x^5,x)
```

output

```
( - atanh(c/x**2)**2*b**2*c**2 + atanh(c/x**2)**2*b**2*x**4 - 2*atanh(c/x*
*2)*a*b*c**2 + 2*atanh(c/x**2)*a*b*x**4 - 2*atanh(c/x**2)*b**2*c*x**2 + 2*
atanh(c/x**2)*b**2*x**4 - 2*log(c + x**2)*b**2*x**4 + 4*log(x)*b**2*x**4 -
a**2*c**2 - 2*a*b*c*x**2)/(4*c**2*x**4)
```

3.176 $\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx$

Optimal result	1422
Mathematica [F]	1423
Rubi [A] (verified)	1424
Maple [F]	1427
Fricas [F]	1427
Sympy [F]	1427
Maxima [F]	1428
Giac [F]	1428
Mupad [F(-1)]	1428
Reduce [F]	1429

Optimal result

Integrand size = 16, antiderivative size = 1214

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \text{Too large to display}$$

output

```

1/20*b^2*x^5*ln(1+c/x^2)^2+1/5*I*b^2*c^(5/2)*polylog(2,1-2*c^(1/2)/(c^(1/2)-I*x))+1/5*I*b^2*c^(5/2)*polylog(2,I*x/c^(1/2))+1/5*b^2*c^(5/2)*arctanh(x/c^(1/2))*ln(2*c^(1/2)*((-c)^(1/2)+x)/((-c)^(1/2)+c^(1/2))/(c^(1/2)+x))+1/5*b^2*c^(5/2)*arctanh(x/c^(1/2))*ln(2*c^(1/2)*((-c)^(1/2)-x)/((-c)^(1/2)-c^(1/2))/(c^(1/2)+x))+2/5*b^2*c^(5/2)*arctanh(x/c^(1/2))*ln(2-2*c^(1/2)/(c^(1/2)+x))-2/5*b^2*c^(5/2)*arctanh(x/c^(1/2))*ln(2*c^(1/2)/(c^(1/2)+x))+2/5*a*b*c^(5/2)*arctan(x/c^(1/2))+1/5*b^2*c^(5/2)*arctan(x/c^(1/2))*ln((1+I)*(c^(1/2)-x)/(c^(1/2)-I*x))+1/5*b^2*c^(5/2)*arctan(x/c^(1/2))*ln((1-I)*(c^(1/2)+x)/(c^(1/2)-I*x))+2/5*b^2*c^(5/2)*arctan(x/c^(1/2))*ln(2-2*c^(1/2)/(c^(1/2)-I*x))-2/5*b^2*c^(5/2)*arctan(x/c^(1/2))*ln(2*c^(1/2)/(c^(1/2)-I*x))-1/5*b^2*c^(5/2)*arctanh(x/c^(1/2))*ln(1+c/x^2)+1/5*b^2*c^(5/2)*arctan(x/c^(1/2))*ln(1+c/x^2)-1/5*b*c^(5/2)*arctanh(x/c^(1/2))*(2*a-b*ln(1-c/x^2))-1/5*b^2*c^(5/2)*arctan(x/c^(1/2))*ln(1-c/x^2)-1/10*I*b^2*c^(5/2)*polylog(2,1+(-1+I)*(c^(1/2)+x)/(c^(1/2)-I*x))-1/10*I*b^2*c^(5/2)*polylog(2,1-(1+I)*(c^(1/2)-x)/(c^(1/2)-I*x))-1/5*I*b^2*c^(5/2)*polylog(2,-1+2*c^(1/2)/(c^(1/2)-I*x))-1/5*I*b^2*c^(5/2)*polylog(2,-I*x/c^(1/2))-1/5*I*b^2*c^(5/2)*arctan(x/c^(1/2))^2+8/15*b^2*c^2*x+1/5*b^2*c^(5/2)*polylog(2,-x/c^(1/2))-1/5*b^2*c^(5/2)*polylog(2,x/c^(1/2))-4/15*b^2*c^(5/2)*arctanh(x/c^(1/2))+1/5*b^2*c^(5/2)*arctanh(x/c^(1/2))^2-4/15*b^2*c^(5/2)*arctan(x/c^(1/2))-1/10*b^2*c^(5/2)*polylog(2,1-2*c^(1/2)*((-c)^(1/2)+x)/((-c)^(1/2)+c^(1/2))/(c^(1/2)+x))

```

Mathematica [F]

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input

```
Integrate[x^4*(a + b*ArcTanh[c/x^2])^2,x]
```

output

```
Integrate[x^4*(a + b*ArcTanh[c/x^2])^2, x]
```

Rubi [A] (verified)

Time = 3.16 (sec) , antiderivative size = 1214, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6460, 6457, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

↓ 6460

$$\int x^4 \left(a + b \operatorname{coth}^{-1} \left(\frac{x^2}{c} \right) \right)^2 dx$$

↓ 6457

$$\int \left(\frac{1}{4} x^4 \left(2a - b \log \left(1 - \frac{c}{x^2} \right) \right)^2 - \frac{1}{2} b x^4 \log \left(\frac{c}{x^2} + 1 \right) \left(b \log \left(1 - \frac{c}{x^2} \right) - 2a \right) + \frac{1}{4} b^2 x^4 \log^2 \left(\frac{c}{x^2} + 1 \right) \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{1}{20} \left(2a - b \log \left(1 - \frac{c}{x^2} \right) \right)^2 x^5 + \frac{1}{20} b^2 \log^2 \left(\frac{c}{x^2} + 1 \right) x^5 + \frac{1}{5} ab \log \left(\frac{c}{x^2} + 1 \right) x^5 - \\
& \frac{1}{10} b^2 \log \left(1 - \frac{c}{x^2} \right) \log \left(\frac{c}{x^2} + 1 \right) x^5 + \frac{2}{15} abc x^3 - \frac{1}{15} b^2 c \log \left(1 - \frac{c}{x^2} \right) x^3 + \\
& \frac{1}{15} bc \left(2a - b \log \left(1 - \frac{c}{x^2} \right) \right) x^3 + \frac{2}{15} b^2 c \log \left(\frac{c}{x^2} + 1 \right) x^3 + \frac{8}{15} b^2 c^2 x - \frac{1}{5} i b^2 c^{5/2} \arctan \left(\frac{x}{\sqrt{c}} \right)^2 + \\
& \frac{1}{5} b^2 c^{5/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right)^2 - \frac{4}{15} b^2 c^{5/2} \arctan \left(\frac{x}{\sqrt{c}} \right) + \frac{2}{5} abc^{5/2} \arctan \left(\frac{x}{\sqrt{c}} \right) - \\
& \frac{4}{15} b^2 c^{5/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right) + \frac{2}{5} b^2 c^{5/2} \arctan \left(\frac{x}{\sqrt{c}} \right) \log \left(2 - \frac{2\sqrt{c}}{\sqrt{c} - ix} \right) - \\
& \frac{1}{5} b^2 c^{5/2} \arctan \left(\frac{x}{\sqrt{c}} \right) \log \left(1 - \frac{c}{x^2} \right) - \frac{1}{5} bc^{5/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right) \left(2a - b \log \left(1 - \frac{c}{x^2} \right) \right) + \\
& \frac{1}{5} b^2 c^{5/2} \arctan \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{c}{x^2} + 1 \right) - \frac{1}{5} b^2 c^{5/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{c}{x^2} + 1 \right) - \\
& \frac{2}{5} b^2 c^{5/2} \arctan \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{2\sqrt{c}}{\sqrt{c} - ix} \right) + \frac{1}{5} b^2 c^{5/2} \arctan \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{(1+i)(\sqrt{c}-x)}{\sqrt{c}-ix} \right) - \\
& \frac{2}{5} b^2 c^{5/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{2\sqrt{c}}{x+\sqrt{c}} \right) + \frac{1}{5} b^2 c^{5/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{2\sqrt{c}(\sqrt{-c}-x)}{(\sqrt{-c}-\sqrt{c})(x+\sqrt{c})} \right) + \\
& \frac{1}{5} b^2 c^{5/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{2\sqrt{c}(x+\sqrt{-c})}{(\sqrt{-c}+\sqrt{c})(x+\sqrt{c})} \right) + \\
& \frac{1}{5} b^2 c^{5/2} \arctan \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{(1-i)(x+\sqrt{c})}{\sqrt{c}-ix} \right) + \frac{2}{5} b^2 c^{5/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right) \log \left(2 - \frac{2\sqrt{c}}{x+\sqrt{c}} \right) + \\
& \frac{1}{5} i b^2 c^{5/2} \operatorname{PolyLog} \left(2, 1 - \frac{2\sqrt{c}}{\sqrt{c}-ix} \right) - \frac{1}{5} i b^2 c^{5/2} \operatorname{PolyLog} \left(2, \frac{2\sqrt{c}}{\sqrt{c}-ix} - 1 \right) - \\
& \frac{1}{10} i b^2 c^{5/2} \operatorname{PolyLog} \left(2, 1 - \frac{(1+i)(\sqrt{c}-x)}{\sqrt{c}-ix} \right) + \frac{1}{5} b^2 c^{5/2} \operatorname{PolyLog} \left(2, -\frac{x}{\sqrt{c}} \right) - \\
& \frac{1}{5} i b^2 c^{5/2} \operatorname{PolyLog} \left(2, -\frac{ix}{\sqrt{c}} \right) + \frac{1}{5} i b^2 c^{5/2} \operatorname{PolyLog} \left(2, \frac{ix}{\sqrt{c}} \right) - \frac{1}{5} b^2 c^{5/2} \operatorname{PolyLog} \left(2, \frac{x}{\sqrt{c}} \right) + \\
& \frac{1}{5} b^2 c^{5/2} \operatorname{PolyLog} \left(2, 1 - \frac{2\sqrt{c}}{x+\sqrt{c}} \right) - \frac{1}{5} b^2 c^{5/2} \operatorname{PolyLog} \left(2, \frac{2\sqrt{c}}{x+\sqrt{c}} - 1 \right) - \\
& \frac{1}{10} b^2 c^{5/2} \operatorname{PolyLog} \left(2, 1 - \frac{2\sqrt{c}(\sqrt{-c}-x)}{(\sqrt{-c}-\sqrt{c})(x+\sqrt{c})} \right) - \\
& \frac{1}{10} b^2 c^{5/2} \operatorname{PolyLog} \left(2, 1 - \frac{2\sqrt{c}(x+\sqrt{-c})}{(\sqrt{-c}+\sqrt{c})(x+\sqrt{c})} \right) - \\
& \frac{1}{10} i b^2 c^{5/2} \operatorname{PolyLog} \left(2, 1 - \frac{(1-i)(x+\sqrt{c})}{\sqrt{c}-ix} \right)
\end{aligned}$$

input

Int[x^4*(a + b*ArcTanh[c/x^2])^2,x]

output

$$\begin{aligned}
& (8b^2c^2x)/15 + (2ab^2cx^3)/15 + (2ab^2c^{5/2}\text{ArcTan}[x/\text{Sqrt}[c]])/5 \\
& - (4b^2c^{5/2}\text{ArcTan}[x/\text{Sqrt}[c]])/15 - (I/5)b^2c^{5/2}\text{ArcTan}[x/\text{Sqrt}[c]]^2 \\
& - (4b^2c^{5/2}\text{ArcTanh}[x/\text{Sqrt}[c]])/15 + (b^2c^{5/2}\text{ArcTanh}[x/\text{Sqrt}[c]]^2)/5 \\
& + (2b^2c^{5/2}\text{ArcTan}[x/\text{Sqrt}[c]]\text{Log}[2 - (2\text{Sqrt}[c])/(c - Ix)])/5 \\
& - (b^2c^2x^3\text{Log}[1 - c/x^2])/15 - (b^2c^{5/2}\text{ArcTan}[x/\text{Sqrt}[c]]\text{Log}[1 - c/x^2])/5 \\
& + (b^2c^2x^3(2a - b\text{Log}[1 - c/x^2]))/15 - (b^2c^{5/2}\text{ArcTanh}[x/\text{Sqrt}[c]](2a - b\text{Log}[1 - c/x^2]))/5 \\
& + (x^5(2a - b\text{Log}[1 - c/x^2])^2)/20 + (2b^2c^2x^3\text{Log}[1 + c/x^2])/15 + (ab^2x^5\text{Log}[1 + c/x^2])/5 \\
& + (b^2c^{5/2}\text{ArcTan}[x/\text{Sqrt}[c]]\text{Log}[1 + c/x^2])/5 - (b^2c^{5/2}\text{ArcTanh}[x/\text{Sqrt}[c]]\text{Log}[1 + c/x^2])/5 \\
& - (b^2x^5\text{Log}[1 - c/x^2]\text{Log}[1 + c/x^2])/10 + (b^2x^5\text{Log}[1 + c/x^2]^2)/20 \\
& - (2b^2c^{5/2}\text{ArcTan}[x/\text{Sqrt}[c]]\text{Log}[(2\text{Sqrt}[c])/(c - Ix)])/5 + (b^2c^{5/2}\text{ArcTan}[x/\text{Sqrt}[c]]\text{Log}[(1 + I)(\text{Sqrt}[c] - x)/(c - Ix)])/5 \\
& - (2b^2c^{5/2}\text{ArcTanh}[x/\text{Sqrt}[c]]\text{Log}[(2\text{Sqrt}[c])/(c + x)])/5 + (b^2c^{5/2}\text{ArcTanh}[x/\text{Sqrt}[c]]\text{Log}[(2\text{Sqrt}[c](\text{Sqrt}[-c] - x)/((\text{Sqrt}[-c] - \text{Sqrt}[c])(\text{Sqrt}[c] + x))])/5 \\
& + (b^2c^{5/2}\text{ArcTanh}[x/\text{Sqrt}[c]]\text{Log}[(2\text{Sqrt}[c](\text{Sqrt}[-c] + x)/((\text{Sqrt}[-c] + \text{Sqrt}[c])(\text{Sqrt}[c] + x))])/5 \\
& + (b^2c^{5/2}\text{ArcTan}[x/\text{Sqrt}[c]]\text{Log}[(1 - I)(\text{Sqrt}[c] + x)/(c - Ix)])/5 + (2b^2c^{5/2}\text{ArcTanh}[x/\text{Sqrt}[c]]\text{Log}[2 - (2\text{Sqrt}[c])/(c + x)])/5 \\
& + (I/5)b^2c^{5/2}\text{PolyLog}[2, 1 - (2\text{Sqrt}[c])/(c - Ix)] - (I/5)b^2c^{5/2}\text{PolyLog}[2, -1 + (2\text{Sqrt}[c])/(c - Ix)]
\end{aligned}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 6457 $\text{Int}[(a_.) + \text{ArcCoth}[(c_.)*(x_)^(n_)]*(b_.)^(p_)*(x_)^(m_.), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[x^m*(a + b*(\text{Log}[1 + 1/(x^n*c)]/2) - b*(\text{Log}[1 - 1/(x^n*c)]/2))^p, x], x] /; \text{FreeQ}\{a, b, c\}, x \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$

rule 6460 $\text{Int}[(a_.) + \text{ArcTanh}[(c_.)*(x_)^(n_)]*(b_.)^(p_)*(x_)^(m_.), x_Symbol] \rightarrow \text{Int}[x^m*(a + b*\text{ArcCoth}[1/(x^n*c)])^p, x] /; \text{FreeQ}\{a, b, c, m\}, x \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{ILtQ}[n, 0]$

Maple [F]

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input `int(x^4*(a+b*arctanh(c/x^2))^2,x)`

output `int(x^4*(a+b*arctanh(c/x^2))^2,x)`

Fricas [F]

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(b \operatorname{arctanh} \left(\frac{c}{x^2} \right) + a \right)^2 x^4 dx$$

input `integrate(x^4*(a+b*arctanh(c/x^2))^2,x, algorithm="fricas")`

output `integral(b^2*x^4*arctanh(c/x^2)^2 + 2*a*b*x^4*arctanh(c/x^2) + a^2*x^4, x)`

Sympy [F]

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int x^4 \left(a + b \operatorname{atanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input `integrate(x**4*(a+b*atanh(c/x**2))**2,x)`

output `Integral(x**4*(a + b*atanh(c/x**2))**2, x)`

Maxima [F]

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(b \operatorname{arctanh} \left(\frac{c}{x^2} \right) + a \right)^2 x^4 dx$$

input `integrate(x^4*(a+b*arctanh(c/x^2))^2,x, algorithm="maxima")`

output `1/5*a^2*x^5 + 1/15*(6*x^5*arctanh(c/x^2) + (4*x^3 + 6*c^(3/2)*arctan(x/sqrt(c)) + 3*c^(3/2)*log((x - sqrt(c))/(x + sqrt(c))))*c)*a*b + 1/20*(x^5*log(x^2 - c)^2 - 5*integrate(-1/5*(5*(x^6 - c*x^4)*log(x^2 + c)^2 - 2*(2*x^6 + 5*(x^6 - c*x^4)*log(x^2 + c))*log(x^2 - c))/(x^2 - c), x))*b^2`

Giac [F]

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(b \operatorname{arctanh} \left(\frac{c}{x^2} \right) + a \right)^2 x^4 dx$$

input `integrate(x^4*(a+b*arctanh(c/x^2))^2,x, algorithm="giac")`

output `integrate((b*arctanh(c/x^2) + a)^2*x^4, x)`

Mupad [F(-1)]

Timed out.

$$\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int x^4 \left(a + b \operatorname{atanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input `int(x^4*(a + b*atanh(c/x^2))^2,x)`

output `int(x^4*(a + b*atanh(c/x^2))^2, x)`

Reduce [F]

$$\begin{aligned}
\int x^4 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = & \frac{2\sqrt{c} \operatorname{atan} \left(\frac{x}{\sqrt{c}} \right) ab c^2}{5} - \frac{4\sqrt{c} \operatorname{atan} \left(\frac{x}{\sqrt{c}} \right) b^2 c^2}{15} \\
& - \frac{\operatorname{atanh} \left(\frac{c}{x^2} \right)^2 b^2 c^2 x}{5} + \frac{\operatorname{atanh} \left(\frac{c}{x^2} \right)^2 b^2 x^5}{5} \\
& + \frac{2\sqrt{c} \operatorname{atanh} \left(\frac{c}{x^2} \right) ab c^2}{5} + \frac{4\sqrt{c} \operatorname{atanh} \left(\frac{c}{x^2} \right) b^2 c^2}{15} \\
& + \frac{2 \operatorname{atanh} \left(\frac{c}{x^2} \right) ab x^5}{5} + \frac{4 \operatorname{atanh} \left(\frac{c}{x^2} \right) b^2 c x^3}{15} \\
& + \frac{2\sqrt{c} \log(\sqrt{c} - x) ab c^2}{5} + \frac{4\sqrt{c} \log(\sqrt{c} - x) b^2 c^2}{15} \\
& - \frac{\sqrt{c} \log(x^2 + c) ab c^2}{5} - \frac{2\sqrt{c} \log(x^2 + c) b^2 c^2}{15} \\
& + \frac{\left(\int \operatorname{atanh} \left(\frac{c}{x^2} \right)^2 dx \right) b^2 c^2}{5} + \frac{a^2 x^5}{5} + \frac{4abc x^3}{15} + \frac{8b^2 c^2 x}{15}
\end{aligned}$$

input `int(x^4*(a+b*atanh(c/x^2))^2,x)`

output `(6*sqrt(c)*atan(x/sqrt(c))*a*b*c**2 - 4*sqrt(c)*atan(x/sqrt(c))*b**2*c**2 - 3*atanh(c/x**2)**2*b**2*c**2*x + 3*atanh(c/x**2)**2*b**2*x**5 + 6*sqrt(c)*atanh(c/x**2)*a*b*c**2 + 4*sqrt(c)*atanh(c/x**2)*b**2*c**2 + 6*atanh(c/x**2)*a*b*x**5 + 4*atanh(c/x**2)*b**2*c*x**3 + 6*sqrt(c)*log(sqrt(c) - x)*a*b*c**2 + 4*sqrt(c)*log(sqrt(c) - x)*b**2*c**2 - 3*sqrt(c)*log(c + x**2)*a*b*c**2 - 2*sqrt(c)*log(c + x**2)*b**2*c**2 + 3*int(atanh(c/x**2)**2,x)*b**2*c**2 + 3*a**2*x**5 + 4*a*b*c*x**3 + 8*b**2*c**2*x)/15`

3.177 $\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx$

Optimal result	1430
Mathematica [F]	1431
Rubi [A] (verified)	1432
Maple [F]	1435
Fricas [F]	1435
Sympy [F]	1435
Maxima [F]	1436
Giac [F]	1436
Mupad [F(-1)]	1436
Reduce [F]	1437

Optimal result

Integrand size = 16, antiderivative size = 1172

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \text{Too large to display}$$

output

```

1/12*x^3*(2*a-b*ln(1-c/x^2))^2-1/3*I*b^2*c^(3/2)*polylog(2,I*x/c^(1/2))+1/
6*I*b^2*c^(3/2)*polylog(2,1+(-1+I)*(c^(1/2)+x)/(c^(1/2)-I*x))+1/6*I*b^2*c^
(3/2)*polylog(2,1-(-1+I)*(c^(1/2)-x)/(c^(1/2)-I*x))+1/3*I*b^2*c^(3/2)*polyl
og(2,-1+2*c^(1/2)/(c^(1/2)-I*x))+1/3*I*b^2*c^(3/2)*polylog(2,-I*x/c^(1/2))
+1/3*I*b^2*c^(3/2)*arctan(x/c^(1/2))^2+1/3*b^2*c^(3/2)*arctanh(x/c^(1/2))*
ln(2*c^(1/2)*((-c)^(1/2)+x)/((-c)^(1/2)+c^(1/2))/(c^(1/2)+x))+1/3*b^2*c^(3
/2)*arctanh(x/c^(1/2))*ln(2*c^(1/2)*((-c)^(1/2)-x)/((-c)^(1/2)-c^(1/2))/(c
^(1/2)+x))+2/3*b^2*c^(3/2)*arctanh(x/c^(1/2))*ln(2-2*c^(1/2)/(c^(1/2)+x))-
2/3*b^2*c^(3/2)*arctanh(x/c^(1/2))*ln(2*c^(1/2)/(c^(1/2)+x))-1/3*b^2*c^(3/
2)*arctan(x/c^(1/2))*ln((1+I)*(c^(1/2)-x)/(c^(1/2)-I*x))-1/3*b^2*c^(3/2)*a
rctan(x/c^(1/2))*ln((1-I)*(c^(1/2)+x)/(c^(1/2)-I*x))-2/3*b^2*c^(3/2)*arcta
n(x/c^(1/2))*ln(2-2*c^(1/2)/(c^(1/2)-I*x))+2/3*b^2*c^(3/2)*arctan(x/c^(1/2
))*ln(2*c^(1/2)/(c^(1/2)-I*x))-2/3*a*b*c^(3/2)*arctan(x/c^(1/2))-1/3*b^2*c
^(3/2)*arctanh(x/c^(1/2))*ln(1+c/x^2)-1/3*b^2*c^(3/2)*arctan(x/c^(1/2))*ln
(1+c/x^2)-1/3*b*c^(3/2)*arctanh(x/c^(1/2))*(2*a-b*ln(1-c/x^2))+1/3*b^2*c^(
3/2)*arctan(x/c^(1/2))*ln(1-c/x^2)-1/3*I*b^2*c^(3/2)*polylog(2,1-2*c^(1/2)
/(c^(1/2)-I*x))-1/6*b^2*c^(3/2)*polylog(2,1-2*c^(1/2)*((-c)^(1/2)+x)/((-c)
^(1/2)+c^(1/2))/(c^(1/2)+x))-1/6*b^2*c^(3/2)*polylog(2,1-2*c^(1/2)*((-c)^(
1/2)-x)/((-c)^(1/2)-c^(1/2))/(c^(1/2)+x))+1/3*b^2*c^(3/2)*polylog(2,1-2*c^
(1/2)/(c^(1/2)+x))-1/3*b^2*c^(3/2)*polylog(2,-1+2*c^(1/2)/(c^(1/2)+x))+...

```

Mathematica [F]

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input

```
Integrate[x^2*(a + b*ArcTanh[c/x^2])^2,x]
```

output

```
Integrate[x^2*(a + b*ArcTanh[c/x^2])^2, x]
```

Rubi [A] (verified)

Time = 2.63 (sec) , antiderivative size = 1172, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6460, 6457, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

↓ 6460

$$\int x^2 \left(a + b \operatorname{coth}^{-1} \left(\frac{x^2}{c} \right) \right)^2 dx$$

↓ 6457

$$\int \left(\frac{1}{4} x^2 \left(2a - b \log \left(1 - \frac{c}{x^2} \right) \right)^2 - \frac{1}{2} b x^2 \log \left(\frac{c}{x^2} + 1 \right) \left(b \log \left(1 - \frac{c}{x^2} \right) - 2a \right) + \frac{1}{4} b^2 x^2 \log^2 \left(\frac{c}{x^2} + 1 \right) \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{1}{12} \left(2a - b \log \left(1 - \frac{c}{x^2} \right) \right)^2 x^3 + \frac{1}{12} b^2 \log^2 \left(\frac{c}{x^2} + 1 \right) x^3 + \frac{1}{3} ab \log \left(\frac{c}{x^2} + 1 \right) x^3 - \\
& \frac{1}{6} b^2 \log \left(1 - \frac{c}{x^2} \right) \log \left(\frac{c}{x^2} + 1 \right) x^3 + \frac{4}{3} abc x - \frac{2}{3} b^2 c \log \left(1 - \frac{c}{x^2} \right) x + \frac{2}{3} b^2 c \log \left(\frac{c}{x^2} + 1 \right) x + \\
& \frac{1}{3} i b^2 c^{3/2} \arctan \left(\frac{x}{\sqrt{c}} \right)^2 + \frac{1}{3} b^2 c^{3/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right)^2 + \frac{4}{3} b^2 c^{3/2} \arctan \left(\frac{x}{\sqrt{c}} \right) - \\
& \frac{2}{3} abc^{3/2} \arctan \left(\frac{x}{\sqrt{c}} \right) - \frac{4}{3} b^2 c^{3/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right) - \frac{2}{3} b^2 c^{3/2} \arctan \left(\frac{x}{\sqrt{c}} \right) \log \left(2 - \frac{2\sqrt{c}}{\sqrt{c} - ix} \right) + \\
& \frac{1}{3} b^2 c^{3/2} \arctan \left(\frac{x}{\sqrt{c}} \right) \log \left(1 - \frac{c}{x^2} \right) - \frac{1}{3} bc^{3/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right) \left(2a - b \log \left(1 - \frac{c}{x^2} \right) \right) - \\
& \frac{1}{3} b^2 c^{3/2} \arctan \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{c}{x^2} + 1 \right) - \frac{1}{3} b^2 c^{3/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{c}{x^2} + 1 \right) + \\
& \frac{2}{3} b^2 c^{3/2} \arctan \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{2\sqrt{c}}{\sqrt{c} - ix} \right) - \frac{1}{3} b^2 c^{3/2} \arctan \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{(1+i)(\sqrt{c}-x)}{\sqrt{c}-ix} \right) - \\
& \frac{2}{3} b^2 c^{3/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{2\sqrt{c}}{x+\sqrt{c}} \right) + \frac{1}{3} b^2 c^{3/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{2\sqrt{c}(\sqrt{-c}-x)}{(\sqrt{-c}-\sqrt{c})(x+\sqrt{c})} \right) + \\
& \frac{1}{3} b^2 c^{3/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{2\sqrt{c}(x+\sqrt{-c})}{(\sqrt{-c}+\sqrt{c})(x+\sqrt{c})} \right) - \\
& \frac{1}{3} b^2 c^{3/2} \arctan \left(\frac{x}{\sqrt{c}} \right) \log \left(\frac{(1-i)(x+\sqrt{c})}{\sqrt{c}-ix} \right) + \frac{2}{3} b^2 c^{3/2} \operatorname{arctanh} \left(\frac{x}{\sqrt{c}} \right) \log \left(2 - \frac{2\sqrt{c}}{x+\sqrt{c}} \right) - \\
& \frac{1}{3} i b^2 c^{3/2} \operatorname{PolyLog} \left(2, 1 - \frac{2\sqrt{c}}{\sqrt{c}-ix} \right) + \frac{1}{3} i b^2 c^{3/2} \operatorname{PolyLog} \left(2, \frac{2\sqrt{c}}{\sqrt{c}-ix} - 1 \right) + \\
& \frac{1}{6} i b^2 c^{3/2} \operatorname{PolyLog} \left(2, 1 - \frac{(1+i)(\sqrt{c}-x)}{\sqrt{c}-ix} \right) + \frac{1}{3} b^2 c^{3/2} \operatorname{PolyLog} \left(2, -\frac{x}{\sqrt{c}} \right) + \\
& \frac{1}{3} i b^2 c^{3/2} \operatorname{PolyLog} \left(2, -\frac{ix}{\sqrt{c}} \right) - \frac{1}{3} i b^2 c^{3/2} \operatorname{PolyLog} \left(2, \frac{ix}{\sqrt{c}} \right) - \frac{1}{3} b^2 c^{3/2} \operatorname{PolyLog} \left(2, \frac{x}{\sqrt{c}} \right) + \\
& \frac{1}{3} b^2 c^{3/2} \operatorname{PolyLog} \left(2, 1 - \frac{2\sqrt{c}}{x+\sqrt{c}} \right) - \frac{1}{3} b^2 c^{3/2} \operatorname{PolyLog} \left(2, \frac{2\sqrt{c}}{x+\sqrt{c}} - 1 \right) - \\
& \frac{1}{6} b^2 c^{3/2} \operatorname{PolyLog} \left(2, 1 - \frac{2\sqrt{c}(\sqrt{-c}-x)}{(\sqrt{-c}-\sqrt{c})(x+\sqrt{c})} \right) - \\
& \frac{1}{6} b^2 c^{3/2} \operatorname{PolyLog} \left(2, 1 - \frac{2\sqrt{c}(x+\sqrt{-c})}{(\sqrt{-c}+\sqrt{c})(x+\sqrt{c})} \right) + \\
& \frac{1}{6} i b^2 c^{3/2} \operatorname{PolyLog} \left(2, 1 - \frac{(1-i)(x+\sqrt{c})}{\sqrt{c}-ix} \right)
\end{aligned}$$

input `Int[x^2*(a + b*ArcTanh[c/x^2])^2,x]`

output

$$\begin{aligned}
& (4*a*b*c*x)/3 - (2*a*b*c^{(3/2)}*ArcTan[x/Sqrt[c]])/3 + (4*b^2*c^{(3/2)}*ArcTan[x/Sqrt[c]])/3 + (I/3)*b^2*c^{(3/2)}*ArcTan[x/Sqrt[c]]^2 - (4*b^2*c^{(3/2)}*ArcTanh[x/Sqrt[c]])/3 + (b^2*c^{(3/2)}*ArcTanh[x/Sqrt[c]]^2)/3 - (2*b^2*c^{(3/2)}*ArcTan[x/Sqrt[c]]*Log[2 - (2*Sqrt[c])/(Sqrt[c] - I*x)])/3 - (2*b^2*c*x*Log[1 - c/x^2])/3 + (b^2*c^{(3/2)}*ArcTan[x/Sqrt[c]]*Log[1 - c/x^2])/3 - (b*c^{(3/2)}*ArcTanh[x/Sqrt[c]]*(2*a - b*Log[1 - c/x^2]))/3 + (x^3*(2*a - b*Log[1 - c/x^2])^2)/12 + (2*b^2*c*x*Log[1 + c/x^2])/3 + (a*b*x^3*Log[1 + c/x^2])/3 - (b^2*c^{(3/2)}*ArcTan[x/Sqrt[c]]*Log[1 + c/x^2])/3 - (b^2*c^{(3/2)}*ArcTanh[x/Sqrt[c]]*Log[1 + c/x^2])/3 - (b^2*x^3*Log[1 - c/x^2]*Log[1 + c/x^2])/6 + (b^2*x^3*Log[1 + c/x^2]^2)/12 + (2*b^2*c^{(3/2)}*ArcTan[x/Sqrt[c]]*Log[(2*Sqrt[c])/(Sqrt[c] - I*x)])/3 - (b^2*c^{(3/2)}*ArcTan[x/Sqrt[c]]*Log[((1 + I)*(Sqrt[c] - x))/(Sqrt[c] - I*x)])/3 - (2*b^2*c^{(3/2)}*ArcTanh[x/Sqrt[c]]*Log[(2*Sqrt[c])/(Sqrt[c] + x)])/3 + (b^2*c^{(3/2)}*ArcTanh[x/Sqrt[c]]*Log[(2*Sqrt[c]*(Sqrt[-c] - x))/((Sqrt[-c] - Sqrt[c])*(Sqrt[c] + x))])/3 + (b^2*c^{(3/2)}*ArcTanh[x/Sqrt[c]]*Log[(2*Sqrt[c]*(Sqrt[-c] + x))/((Sqrt[-c] + Sqrt[c])*(Sqrt[c] + x))])/3 - (b^2*c^{(3/2)}*ArcTan[x/Sqrt[c]]*Log[((1 - I)*(Sqrt[c] + x))/(Sqrt[c] - I*x)])/3 + (2*b^2*c^{(3/2)}*ArcTanh[x/Sqrt[c]]*Log[2 - (2*Sqrt[c])/(Sqrt[c] + x)])/3 - (I/3)*b^2*c^{(3/2)}*PolyLog[2, 1 - (2*Sqrt[c])/(Sqrt[c] - I*x)] + (I/3)*b^2*c^{(3/2)}*PolyLog[2, -1 + (2*Sqrt[c])/(Sqrt[c] - I*x)] + (I/6)*b^2*c^{(3/2)}*PolyLog[2, 1 - ((1 + I)*(Sqrt[c] - x)...
\end{aligned}$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6457 `Int[((a_.) + ArcCoth[(c_.)*(x_)^(n_)])*(b_.))^(p_)*(x_)^(m_.), x_Symbol] := Int[ExpandIntegrand[x^m*(a + b*(Log[1 + 1/(x^n*c)]/2) - b*(Log[1 - 1/(x^n*c)]/2))^p, x], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1] && IGtQ[n, 0] && IntegerQ[m]`

rule 6460 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_)*(x_)^(m_.), x_Symbol] := Int[x^m*(a + b*ArcCoth[1/(x^n*c)])^p, x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 1] && ILtQ[n, 0]`

Maple [F]

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input `int(x^2*(a+b*arctanh(c/x^2))^2,x)`

output `int(x^2*(a+b*arctanh(c/x^2))^2,x)`

Fricas [F]

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(b \operatorname{arctanh} \left(\frac{c}{x^2} \right) + a \right)^2 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c/x^2))^2,x, algorithm="fricas")`

output `integral(b^2*x^2*arctanh(c/x^2)^2 + 2*a*b*x^2*arctanh(c/x^2) + a^2*x^2, x)`

Sympy [F]

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int x^2 \left(a + b \operatorname{atanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input `integrate(x**2*(a+b*atanh(c/x**2))**2,x)`

output `Integral(x**2*(a + b*atanh(c/x**2))**2, x)`

Maxima [F]

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(b \operatorname{arctanh} \left(\frac{c}{x^2} \right) + a \right)^2 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c/x^2))^2,x, algorithm="maxima")`

output `1/3*a^2*x^3 + 1/3*(2*x^3*arctanh(c/x^2) - (2*sqrt(c)*arctan(x/sqrt(c)) - sqrt(c)*log((x - sqrt(c))/(x + sqrt(c)))) - 4*x)*c)*a*b + 1/12*(x^3*log(x^2 - c)^2 - 3*integrate(-1/3*(3*(x^4 - c*x^2)*log(x^2 + c)^2 - 2*(2*x^4 + 3*(x^4 - c*x^2)*log(x^2 + c))*log(x^2 - c))/(x^2 - c), x))*b^2`

Giac [F]

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(b \operatorname{arctanh} \left(\frac{c}{x^2} \right) + a \right)^2 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c/x^2))^2,x, algorithm="giac")`

output `integrate((b*arctanh(c/x^2) + a)^2*x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int x^2 \left(a + b \operatorname{atanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input `int(x^2*(a + b*atanh(c/x^2))^2,x)`

output `int(x^2*(a + b*atanh(c/x^2))^2, x)`

Reduce [F]

$$\begin{aligned}
\int x^2 \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = & -\frac{2\sqrt{c} \operatorname{atan} \left(\frac{x}{\sqrt{c}} \right) abc}{3} + \frac{4\sqrt{c} \operatorname{atan} \left(\frac{x}{\sqrt{c}} \right) b^2 c}{3} \\
& + \frac{\operatorname{atanh} \left(\frac{c}{x^2} \right)^2 b^2 x^3}{3} + \frac{2\sqrt{c} \operatorname{atanh} \left(\frac{c}{x^2} \right) abc}{3} \\
& + \frac{4\sqrt{c} \operatorname{atanh} \left(\frac{c}{x^2} \right) b^2 c}{3} + \frac{2 \operatorname{atanh} \left(\frac{c}{x^2} \right) ab x^3}{3} \\
& + \frac{4 \operatorname{atanh} \left(\frac{c}{x^2} \right) b^2 cx}{3} + \frac{2\sqrt{c} \log(\sqrt{c} - x) abc}{3} \\
& + \frac{4\sqrt{c} \log(\sqrt{c} - x) b^2 c}{3} \\
& - \frac{\sqrt{c} \log(x^2 + c) abc}{3} - \frac{2\sqrt{c} \log(x^2 + c) b^2 c}{3} \\
& - \frac{4 \left(\int \frac{\operatorname{atanh} \left(\frac{c}{x^2} \right)}{-x^4 + c^2} dx \right) b^2 c^3}{3} + \frac{a^2 x^3}{3} + \frac{4abcx}{3}
\end{aligned}$$

input `int(x^2*(a+b*atanh(c/x^2))^2,x)`

output `(- 2*sqrt(c)*atan(x/sqrt(c))*a*b*c + 4*sqrt(c)*atan(x/sqrt(c))*b**2*c + a
tanh(c/x**2)**2*b**2*x**3 + 2*sqrt(c)*atanh(c/x**2)*a*b*c + 4*sqrt(c)*atan
h(c/x**2)*b**2*c + 2*atanh(c/x**2)*a*b*x**3 + 4*atanh(c/x**2)*b**2*c*x + 2
*sqrt(c)*log(sqrt(c) - x)*a*b*c + 4*sqrt(c)*log(sqrt(c) - x)*b**2*c - sqrt
(c)*log(c + x**2)*a*b*c - 2*sqrt(c)*log(c + x**2)*b**2*c - 4*int(atanh(c/x
2)/(c2 - x**4),x)*b**2*c**3 + a**2*x**3 + 4*a*b*c*x)/3`

3.178 $\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2 dx$

Optimal result	1438
Mathematica [A] (verified)	1439
Rubi [A] (verified)	1440
Maple [F]	1443
Fricas [F]	1443
Sympy [F]	1443
Maxima [F]	1444
Giac [F]	1444
Mupad [F(-1)]	1444
Reduce [F]	1445

Optimal result

Integrand size = 12, antiderivative size = 1050

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2 dx = \text{Too large to display}$$

output

```

-1/2*I*b^2*c^(1/2)*polylog(2,1-(1+I)*(c^(1/2)-x)/(c^(1/2)-I*x))-I*b^2*c^(1/2)*polylog(2,-1+2*c^(1/2)/(c^(1/2)-I*x))-I*b^2*c^(1/2)*polylog(2,-I*x/c^(1/2))+b^2*c^(1/2)*arctan(x/c^(1/2))*ln((1+I)*(c^(1/2)-x)/(c^(1/2)-I*x))+b^2*c^(1/2)*arctan(x/c^(1/2))*ln((1-I)*(c^(1/2)+x)/(c^(1/2)-I*x))-b^2*c^(1/2)*arctanh(x/c^(1/2))*ln(1+c/x^2)+b^2*c^(1/2)*arctan(x/c^(1/2))*ln(1+c/x^2)-b^2*c^(1/2)*arctan(x/c^(1/2))*ln(1-c/x^2)+b^2*c^(1/2)*arctanh(x/c^(1/2))*ln(1-c/x^2)+I*b^2*c^(1/2)*polylog(2,1-2*c^(1/2)/(c^(1/2)-I*x))+I*b^2*c^(1/2)*polylog(2,I*x/c^(1/2))+b^2*c^(1/2)*arctanh(x/c^(1/2))*ln(2*c^(1/2)*((-c)^(1/2)+x)/((-c)^(1/2)+c^(1/2))/(c^(1/2)+x))+b^2*c^(1/2)*arctanh(x/c^(1/2))*ln(2*c^(1/2)*((-c)^(1/2)-x)/((-c)^(1/2)-c^(1/2))/(c^(1/2)+x))+2*a*b*c^(1/2)*arctan(x/c^(1/2))+2*b^2*c^(1/2)*arctan(x/c^(1/2))*ln(2-2*c^(1/2)/(c^(1/2)-I*x))-2*b^2*c^(1/2)*arctan(x/c^(1/2))*ln(2*c^(1/2)/(c^(1/2)-I*x))-2*a*b*c^(1/2)*arctanh(x/c^(1/2))+2*b^2*c^(1/2)*arctanh(x/c^(1/2))*ln(2-2*c^(1/2)/(c^(1/2)+x))-2*b^2*c^(1/2)*arctanh(x/c^(1/2))*ln(2*c^(1/2)/(c^(1/2)+x))-I*b^2*c^(1/2)*arctan(x/c^(1/2))^2-1/2*I*b^2*c^(1/2)*polylog(2,1+(-1+I)*(c^(1/2)+x)/(c^(1/2)-I*x))-1/2*b^2*c^(1/2)*polylog(2,1-2*c^(1/2)*((-c)^(1/2)+x)/((-c)^(1/2)+c^(1/2))/(c^(1/2)+x))-1/2*b^2*c^(1/2)*polylog(2,1-2*c^(1/2)*((-c)^(1/2)-x)/((-c)^(1/2)-c^(1/2))/(c^(1/2)+x))+1/4*b^2*x*ln(1+c/x^2)^2+1/4*b^2*x*ln(1-c/x^2)^2+a*b*x*ln(1+c/x^2)-a*b*x*ln(1-c/x^2)+b^2*c^(1/2)*polylog(2,-x/c^(1/2))-b^2*c^(1/2)*polylog(2,x/c^(1/2))+b^2*c^(1/2)*arcta...

```

Mathematica [A] (verified)

Time = 2.25 (sec) , antiderivative size = 565, normalized size of antiderivative = 0.54

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \text{Too large to display}$$

input

```
Integrate[(a + b*ArcTanh[c/x^2])^2,x]
```

output

```

a^2*x - 2*a*b*Sqrt[c/x^2]*x*(ArcTan[Sqrt[c/x^2]] + ArcTanh[Sqrt[c/x^2]]) +
2*a*b*x*ArcTanh[c/x^2] - (b^2*Sqrt[c/x^2]*x*((-2*I)*ArcTan[Sqrt[c/x^2]]^2
+ 4*ArcTan[Sqrt[c/x^2]]*ArcTanh[c/x^2] - (2*ArcTanh[c/x^2]^2)/Sqrt[c/x^2]
+ 2*ArcTan[Sqrt[c/x^2]]*Log[1 + E^((4*I)*ArcTan[Sqrt[c/x^2]])] - 2*ArcTan
h[c/x^2]*Log[1 - Sqrt[c/x^2]] + Log[2]*Log[1 - Sqrt[c/x^2]] - Log[1 - Sqrt
[c/x^2]]^2/2 + Log[1 - Sqrt[c/x^2]]*Log[(1/2 + I/2)*(-I + Sqrt[c/x^2])] +
2*ArcTanh[c/x^2]*Log[1 + Sqrt[c/x^2]] - Log[2]*Log[1 + Sqrt[c/x^2]] - Log[
((1 + I) - (1 - I)*Sqrt[c/x^2])/2]*Log[1 + Sqrt[c/x^2]] - Log[(-1/2 - I/2)
*(I + Sqrt[c/x^2])]*Log[1 + Sqrt[c/x^2]] + Log[1 + Sqrt[c/x^2]]^2/2 + Log[
1 - Sqrt[c/x^2]]*Log[((1 + I) + (1 - I)*Sqrt[c/x^2])/2] - (I/2)*PolyLog[2,
-E^((4*I)*ArcTan[Sqrt[c/x^2]])] - PolyLog[2, (1 - Sqrt[c/x^2])/2] + PolyL
og[2, (-1/2 - I/2)*(-1 + Sqrt[c/x^2])] + PolyLog[2, (-1/2 + I/2)*(-1 + Sqr
t[c/x^2])] + PolyLog[2, (1 + Sqrt[c/x^2])/2] - PolyLog[2, (1/2 - I/2)*(1 + Sqr
t[c/x^2])] - PolyLog[2, (1/2 + I/2)*(1 + Sqrt[c/x^2])]))/2

```

Rubi [A] (verified)

Time = 1.75 (sec) , antiderivative size = 1050, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6440, 6439, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx \\
 & \quad \downarrow \text{6440} \\
 & \int \left(a + b \operatorname{coth}^{-1} \left(\frac{x^2}{c} \right) \right)^2 dx \\
 & \quad \downarrow \text{6439} \\
 & \int \left(a^2 - ab \log \left(1 - \frac{c}{x^2} \right) + ab \log \left(\frac{c}{x^2} + 1 \right) + \frac{1}{4} b^2 \log^2 \left(1 - \frac{c}{x^2} \right) + \frac{1}{4} b^2 \log^2 \left(\frac{c}{x^2} + 1 \right) - \frac{1}{2} b^2 \log \left(1 - \frac{c}{x^2} \right) \log \left(\frac{c}{x^2} + 1 \right) \right) dx \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\begin{aligned}
& xa^2 + 2b\sqrt{c} \arctan\left(\frac{x}{\sqrt{c}}\right) a - 2b\sqrt{c} \operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) a - bx \log\left(1 - \frac{c}{x^2}\right) a + bx \log\left(\frac{c}{x^2} + 1\right) a - \\
& ib^2\sqrt{c} \arctan\left(\frac{x}{\sqrt{c}}\right)^2 + b^2\sqrt{c} \operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right)^2 + \frac{1}{4}b^2x \log^2\left(1 - \frac{c}{x^2}\right) + \frac{1}{4}b^2x \log^2\left(\frac{c}{x^2} + 1\right) + \\
& 2b^2\sqrt{c} \arctan\left(\frac{x}{\sqrt{c}}\right) \log\left(2 - \frac{2\sqrt{c}}{\sqrt{c} - ix}\right) - b^2\sqrt{c} \arctan\left(\frac{x}{\sqrt{c}}\right) \log\left(1 - \frac{c}{x^2}\right) + \\
& b^2\sqrt{c} \operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) \log\left(1 - \frac{c}{x^2}\right) + b^2\sqrt{c} \arctan\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{c}{x^2} + 1\right) - \\
& b^2\sqrt{c} \operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{c}{x^2} + 1\right) - \frac{1}{2}b^2x \log\left(1 - \frac{c}{x^2}\right) \log\left(\frac{c}{x^2} + 1\right) - \\
& 2b^2\sqrt{c} \arctan\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{2\sqrt{c}}{\sqrt{c} - ix}\right) + b^2\sqrt{c} \arctan\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{(1+i)(\sqrt{c}-x)}{\sqrt{c}-ix}\right) - \\
& 2b^2\sqrt{c} \operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{2\sqrt{c}}{x+\sqrt{c}}\right) + b^2\sqrt{c} \operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{2\sqrt{c}(\sqrt{-c}-x)}{(\sqrt{-c}-\sqrt{c})(x+\sqrt{c})}\right) + \\
& b^2\sqrt{c} \operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{2\sqrt{c}(x+\sqrt{-c})}{(\sqrt{-c}+\sqrt{c})(x+\sqrt{c})}\right) + \\
& b^2\sqrt{c} \arctan\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{(1-i)(x+\sqrt{c})}{\sqrt{c}-ix}\right) + 2b^2\sqrt{c} \operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) \log\left(2 - \frac{2\sqrt{c}}{x+\sqrt{c}}\right) + \\
& ib^2\sqrt{c} \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{c}}{\sqrt{c}-ix}\right) - ib^2\sqrt{c} \operatorname{PolyLog}\left(2, \frac{2\sqrt{c}}{\sqrt{c}-ix} - 1\right) - \\
& \frac{1}{2}ib^2\sqrt{c} \operatorname{PolyLog}\left(2, 1 - \frac{(1+i)(\sqrt{c}-x)}{\sqrt{c}-ix}\right) + b^2\sqrt{c} \operatorname{PolyLog}\left(2, -\frac{x}{\sqrt{c}}\right) - \\
& ib^2\sqrt{c} \operatorname{PolyLog}\left(2, -\frac{ix}{\sqrt{c}}\right) + ib^2\sqrt{c} \operatorname{PolyLog}\left(2, \frac{ix}{\sqrt{c}}\right) - b^2\sqrt{c} \operatorname{PolyLog}\left(2, \frac{x}{\sqrt{c}}\right) + \\
& b^2\sqrt{c} \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{c}}{x+\sqrt{c}}\right) - b^2\sqrt{c} \operatorname{PolyLog}\left(2, \frac{2\sqrt{c}}{x+\sqrt{c}} - 1\right) - \\
& \frac{1}{2}b^2\sqrt{c} \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{c}(\sqrt{-c}-x)}{(\sqrt{-c}-\sqrt{c})(x+\sqrt{c})}\right) - \\
& \frac{1}{2}b^2\sqrt{c} \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{c}(x+\sqrt{-c})}{(\sqrt{-c}+\sqrt{c})(x+\sqrt{c})}\right) - \frac{1}{2}ib^2\sqrt{c} \operatorname{PolyLog}\left(2, 1 - \frac{(1-i)(x+\sqrt{c})}{\sqrt{c}-ix}\right)
\end{aligned}$$

input

Int[(a + b*ArcTanh[c/x^2])^2,x]

output

```

a^2*x + 2*a*b*Sqrt[c]*ArcTan[x/Sqrt[c]] - I*b^2*Sqrt[c]*ArcTan[x/Sqrt[c]]^
2 - 2*a*b*Sqrt[c]*ArcTanh[x/Sqrt[c]] + b^2*Sqrt[c]*ArcTanh[x/Sqrt[c]]^2 +
2*b^2*Sqrt[c]*ArcTan[x/Sqrt[c]]*Log[2 - (2*Sqrt[c])/(Sqrt[c] - I*x)] - a*b
*x*Log[1 - c/x^2] - b^2*Sqrt[c]*ArcTan[x/Sqrt[c]]*Log[1 - c/x^2] + b^2*Sqr
t[c]*ArcTanh[x/Sqrt[c]]*Log[1 - c/x^2] + (b^2*x*Log[1 - c/x^2]^2)/4 + a*b*
x*Log[1 + c/x^2] + b^2*Sqrt[c]*ArcTan[x/Sqrt[c]]*Log[1 + c/x^2] - b^2*Sqrt
[c]*ArcTanh[x/Sqrt[c]]*Log[1 + c/x^2] - (b^2*x*Log[1 - c/x^2]*Log[1 + c/x^
2])/2 + (b^2*x*Log[1 + c/x^2]^2)/4 - 2*b^2*Sqrt[c]*ArcTan[x/Sqrt[c]]*Log[(
2*Sqrt[c])/(Sqrt[c] - I*x)] + b^2*Sqrt[c]*ArcTan[x/Sqrt[c]]*Log[((1 + I)*(
Sqrt[c] - x))/(Sqrt[c] - I*x)] - 2*b^2*Sqrt[c]*ArcTanh[x/Sqrt[c]]*Log[(2*S
qrt[c])/(Sqrt[c] + x)] + b^2*Sqrt[c]*ArcTanh[x/Sqrt[c]]*Log[(2*Sqrt[c]*(Sq
rt[-c] - x))/((Sqrt[-c] - Sqrt[c])*(Sqrt[c] + x))] + b^2*Sqrt[c]*ArcTanh[x
/Sqrt[c]]*Log[(2*Sqrt[c]*(Sqrt[-c] + x))/((Sqrt[-c] + Sqrt[c])*(Sqrt[c] +
x))] + b^2*Sqrt[c]*ArcTan[x/Sqrt[c]]*Log[((1 - I)*(Sqrt[c] + x))/(Sqrt[c]
- I*x)] + 2*b^2*Sqrt[c]*ArcTanh[x/Sqrt[c]]*Log[2 - (2*Sqrt[c])/(Sqrt[c] +
x)] + I*b^2*Sqrt[c]*PolyLog[2, 1 - (2*Sqrt[c])/(Sqrt[c] - I*x)] - I*b^2*Sqr
t[c]*PolyLog[2, -1 + (2*Sqrt[c])/(Sqrt[c] - I*x)] - (I/2)*b^2*Sqrt[c]*Pol
yLog[2, 1 - ((1 + I)*(Sqrt[c] - x))/(Sqrt[c] - I*x)] + b^2*Sqrt[c]*PolyLog
[2, -(x/Sqrt[c])] - I*b^2*Sqrt[c]*PolyLog[2, ((-I)*x)/Sqrt[c]] + I*b^2*Sqr
t[c]*PolyLog[2, (I*x)/Sqrt[c]] - b^2*Sqrt[c]*PolyLog[2, x/Sqrt[c]] + b^...

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6439

```
Int[((a_.) + ArcCoth[(c_.)*(x_)^(n_)])*(b_.))^p_, x_Symbol] := Int[ExpandI
ntegrand[(a + b*(Log[1 + 1/(x^n*c)])/2) - b*(Log[1 - 1/(x^n*c)])/2]^p, x]
/; FreeQ[{a, b, c}, x] && IGtQ[p, 1] && IGtQ[n, 0]
```

rule 6440

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^p_, x_Symbol] := Int[(a + b*
ArcCoth[1/(x^n*c)])^p, x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1] && ILtQ[n, 0]
]
```

Maple [F]

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input `int((a+b*arctanh(c/x^2))^2,x)`

output `int((a+b*arctanh(c/x^2))^2,x)`

Fricas [F]

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(b \operatorname{arctanh} \left(\frac{c}{x^2} \right) + a \right)^2 dx$$

input `integrate((a+b*arctanh(c/x^2))^2,x, algorithm="fricas")`

output `integral(b^2*arctanh(c/x^2)^2 + 2*a*b*arctanh(c/x^2) + a^2, x)`

Sympy [F]

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(a + b \operatorname{atanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input `integrate((a+b*atanh(c/x**2))**2,x)`

output `Integral((a + b*atanh(c/x**2))**2, x)`

Maxima [F]

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x^2} \right) + a \right)^2 dx$$

input `integrate((a+b*arctanh(c/x^2))^2,x, algorithm="maxima")`

output `(c*(2*arctan(x/sqrt(c))/sqrt(c) + log((x - sqrt(c))/(x + sqrt(c)))/sqrt(c)) + 2*x*arctanh(c/x^2))*a*b + 1/4*(x*log(x^2 - c)^2 - integrate(-((x^2 - c)*log(x^2 + c)^2 - 2*(2*x^2 + (x^2 - c)*log(x^2 + c))*log(x^2 - c))/(x^2 - c), x))*b^2 + a^2*x`

Giac [F]

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x^2} \right) + a \right)^2 dx$$

input `integrate((a+b*arctanh(c/x^2))^2,x, algorithm="giac")`

output `integrate((b*arctanh(c/x^2) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(a + b \operatorname{atanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input `int((a + b*atanh(c/x^2))^2,x)`

output `int((a + b*atanh(c/x^2))^2, x)`

Reduce [F]

$$\int \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right) \right)^2 dx = 2\sqrt{c} \operatorname{atan}\left(\frac{x}{\sqrt{c}}\right) ab + 2\sqrt{c} \operatorname{atanh}\left(\frac{c}{x^2}\right) ab \\ + 2 \operatorname{atanh}\left(\frac{c}{x^2}\right) abx + 2\sqrt{c} \log(\sqrt{c} - x) ab \\ - \sqrt{c} \log(x^2 + c) ab + \left(\int \operatorname{atanh}\left(\frac{c}{x^2}\right)^2 dx \right) b^2 + a^2 x$$

input `int((a+b*atanh(c/x^2))^2,x)`

output `2*sqrt(c)*atan(x/sqrt(c))*a*b + 2*sqrt(c)*atanh(c/x**2)*a*b + 2*atanh(c/x**2)*a*b*x + 2*sqrt(c)*log(sqrt(c) - x)*a*b - sqrt(c)*log(c + x**2)*a*b + int(atanh(c/x**2)**2,x)*b**2 + a**2*x`

$$3.179 \quad \int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^2} dx$$

Optimal result	1446
Mathematica [A] (verified)	1447
Rubi [A] (verified)	1448
Maple [F]	1451
Fricas [F]	1451
Sympy [F]	1451
Maxima [F]	1452
Giac [F]	1452
Mupad [F(-1)]	1452
Reduce [F]	1453

Optimal result

Integrand size = 16, antiderivative size = 1080

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^2} dx = \text{Too large to display}$$

output

```

-1/4*(2*a-b*ln(1-c/x^2))^2/x-b^2*arctan(x/c^(1/2))*ln(1-c/x^2)/c^(1/2)+I*b
^2*polylog(2,1-2*c^(1/2)/(c^(1/2)-I*x))/c^(1/2)+I*b^2*polylog(2,I*x/c^(1/2
))/c^(1/2)-b^2*arctanh(x/c^(1/2))*ln(2*c^(1/2)*((-c)^(1/2)+x)/((-c)^(1/2)+
c^(1/2))/(c^(1/2)+x))/c^(1/2)-b^2*arctanh(x/c^(1/2))*ln(2*c^(1/2)*((-c)^(1
/2)-x)/((-c)^(1/2)-c^(1/2))/(c^(1/2)+x))/c^(1/2)+b^2*arctanh(x/c^(1/2))*ln
(1+c/x^2)/c^(1/2)+b^2*arctan(x/c^(1/2))*ln((1+I)*(c^(1/2)-x)/(c^(1/2)-I*x)
)/c^(1/2)+b^2*arctan(x/c^(1/2))*ln((1-I)*(c^(1/2)+x)/(c^(1/2)-I*x))/c^(1/2
)+2*b^2*arctanh(x/c^(1/2))*ln(2*c^(1/2)/(c^(1/2)+x))/c^(1/2)+2*a*b*arctan(
x/c^(1/2))/c^(1/2)+2*b^2*arctan(x/c^(1/2))*ln(2-2*c^(1/2)/(c^(1/2)-I*x))/c
^(1/2)-2*b^2*arctan(x/c^(1/2))*ln(2*c^(1/2)/(c^(1/2)-I*x))/c^(1/2)-2*b^2*a
rctanh(x/c^(1/2))*ln(2-2*c^(1/2)/(c^(1/2)+x))/c^(1/2)-I*b^2*arctan(x/c^(1/
2))^2/c^(1/2)-1/2*I*b^2*polylog(2,1+(-1+I)*(c^(1/2)+x)/(c^(1/2)-I*x))/c^(1
/2)-1/2*I*b^2*polylog(2,1-(1+I)*(c^(1/2)-x)/(c^(1/2)-I*x))/c^(1/2)-I*b^2*p
olylog(2,-1+2*c^(1/2)/(c^(1/2)-I*x))/c^(1/2)-I*b^2*polylog(2,-I*x/c^(1/2)
)/c^(1/2)+b^2*arctan(x/c^(1/2))*ln(1+c/x^2)/c^(1/2)+b*arctanh(x/c^(1/2))*(2
*a-b*ln(1-c/x^2))/c^(1/2)+1/2*b^2*polylog(2,1-2*c^(1/2)*((-c)^(1/2)+x)/((-
c)^(1/2)+c^(1/2))/(c^(1/2)+x))/c^(1/2)+1/2*b^2*polylog(2,1-2*c^(1/2)*((-c)
^(1/2)-x)/((-c)^(1/2)-c^(1/2))/(c^(1/2)+x))/c^(1/2)-b^2*polylog(2,1-2*c^(1
/2)/(c^(1/2)+x))/c^(1/2)+b^2*polylog(2,-1+2*c^(1/2)/(c^(1/2)+x))/c^(1/2)-b
^2*polylog(2,-x/c^(1/2))/c^(1/2)+b^2*polylog(2,x/c^(1/2))/c^(1/2)-b^2*a...

```

Mathematica [A] (verified)

Time = 1.97 (sec) , antiderivative size = 568, normalized size of antiderivative = 0.53

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^2} dx = \text{Too large to display}$$

input

```
Integrate[(a + b*ArcTanh[c/x^2])^2/x^2,x]
```

output

```

(-2*a^2 - (4*a*b*(ArcTan[Sqrt[c/x^2]] - ArcTanh[Sqrt[c/x^2]]))/Sqrt[c/x^2]
- 4*a*b*ArcTanh[c/x^2] + (b^2*((2*I)*ArcTan[Sqrt[c/x^2]]^2 - 4*ArcTan[Sqr
t[c/x^2]]*ArcTanh[c/x^2] - 2*Sqrt[c/x^2]*ArcTanh[c/x^2]^2 - 2*ArcTan[Sqrt[
c/x^2]]*Log[1 + E^((4*I)*ArcTan[Sqrt[c/x^2]])] - 2*ArcTanh[c/x^2]*Log[1 -
Sqrt[c/x^2]] + Log[2]*Log[1 - Sqrt[c/x^2]] - Log[1 - Sqrt[c/x^2]]^2/2 + Lo
g[1 - Sqrt[c/x^2]]*Log[(1/2 + I/2)*(-I + Sqrt[c/x^2])]) + 2*ArcTanh[c/x^2]*
Log[1 + Sqrt[c/x^2]] - Log[2]*Log[1 + Sqrt[c/x^2]] - Log[((1 + I) - (1 - I
)*Sqrt[c/x^2])/2]*Log[1 + Sqrt[c/x^2]] - Log[(-1/2 - I/2)*(I + Sqrt[c/x^2]
)]*Log[1 + Sqrt[c/x^2]] + Log[1 + Sqrt[c/x^2]]^2/2 + Log[1 - Sqrt[c/x^2]]*
Log[((1 + I) + (1 - I)*Sqrt[c/x^2])/2] + (I/2)*PolyLog[2, -E^((4*I)*ArcTan
[Sqrt[c/x^2]])] - PolyLog[2, (1 - Sqrt[c/x^2])/2] + PolyLog[2, (-1/2 - I/2
)*(-1 + Sqrt[c/x^2])] + PolyLog[2, (-1/2 + I/2)*(-1 + Sqrt[c/x^2])] + Poly
Log[2, (1 + Sqrt[c/x^2])/2] - PolyLog[2, (1/2 - I/2)*(1 + Sqrt[c/x^2])] -
PolyLog[2, (1/2 + I/2)*(1 + Sqrt[c/x^2])])/Sqrt[c/x^2])/(2*x)

```

Rubi [A] (verified)

Time = 2.43 (sec) , antiderivative size = 1117, normalized size of antiderivative = 1.03, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6460, 6457, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^2} dx \\
 & \quad \downarrow \text{6460} \\
 & \int \frac{\left(a + b \operatorname{coth}^{-1}\left(\frac{x^2}{c}\right)\right)^2}{x^2} dx \\
 & \quad \downarrow \text{6457} \\
 & \int \left(\frac{\left(2a - b \log\left(1 - \frac{c}{x^2}\right)\right)^2}{4x^2} - \frac{b \log\left(\frac{c}{x^2} + 1\right) \left(b \log\left(1 - \frac{c}{x^2}\right) - 2a\right)}{2x^2} + \frac{b^2 \log^2\left(\frac{c}{x^2} + 1\right)}{4x^2} \right) dx \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\begin{aligned}
& \frac{i \arctan\left(\frac{x}{\sqrt{c}}\right)^2 b^2}{\sqrt{c}} - \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right)^2 b^2}{\sqrt{c}} - \frac{\log^2\left(\frac{c}{x^2} + 1\right) b^2}{4x} - \frac{2 \cot^{-1}\left(\frac{x}{\sqrt{c}}\right) b^2}{\sqrt{c}} - \\
& \frac{2 \coth^{-1}\left(\frac{x}{\sqrt{c}}\right) b^2}{\sqrt{c}} - \frac{2 \arctan\left(\frac{x}{\sqrt{c}}\right) b^2}{\sqrt{c}} + \frac{2 \operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) b^2}{\sqrt{c}} + \\
& \frac{2 \arctan\left(\frac{x}{\sqrt{c}}\right) \log\left(2 - \frac{2\sqrt{c}}{\sqrt{c-ix}}\right) b^2}{\sqrt{c}} + \frac{\cot^{-1}\left(\frac{x}{\sqrt{c}}\right) \log\left(1 - \frac{c}{x^2}\right) b^2}{\sqrt{c}} - \frac{\log\left(1 - \frac{c}{x^2}\right) b^2}{x} + \\
& \frac{\coth^{-1}\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{c}{x^2} + 1\right) b^2}{\sqrt{c}} + \frac{\arctan\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{c}{x^2} + 1\right) b^2}{\sqrt{c}} + \frac{\log\left(1 - \frac{c}{x^2}\right) \log\left(\frac{c}{x^2} + 1\right) b^2}{2x} + \\
& \frac{2 \cot^{-1}\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{2}{1 - \frac{i\sqrt{c}}{x}}\right) b^2}{\sqrt{c}} - \frac{\cot^{-1}\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{(1+i)\left(1 - \frac{\sqrt{c}}{x}\right)}{1 - \frac{i\sqrt{c}}{x}}\right) b^2}{\sqrt{c}} + \\
& \frac{2 \coth^{-1}\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{2}{\frac{\sqrt{c}}{x} + 1}\right) b^2}{\sqrt{c}} - \frac{\coth^{-1}\left(\frac{x}{\sqrt{c}}\right) \log\left(-\frac{2\sqrt{c}\left(1 - \frac{\sqrt{c}}{x}\right)}{(\sqrt{c} - \sqrt{c})\left(\frac{\sqrt{c}}{x} + 1\right)}\right) b^2}{\sqrt{c}} - \\
& \frac{\coth^{-1}\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{2\sqrt{c}\left(\frac{\sqrt{c}}{x} + 1\right)}{(\sqrt{c} + \sqrt{c})\left(\frac{\sqrt{c}}{x} + 1\right)}\right) b^2}{\sqrt{c}} - \frac{\cot^{-1}\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{(1-i)\left(\frac{\sqrt{c}}{x} + 1\right)}{1 - \frac{i\sqrt{c}}{x}}\right) b^2}{\sqrt{c}} - \\
& \frac{2 \operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) \log\left(2 - \frac{2\sqrt{c}}{x + \sqrt{c}}\right) b^2}{\sqrt{c}} - \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{i\sqrt{c}}{x}}\right) b^2}{\sqrt{c}} + \\
& \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{(1+i)\left(1 - \frac{\sqrt{c}}{x}\right)}{1 - \frac{i\sqrt{c}}{x}}\right) b^2}{2\sqrt{c}} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{\frac{\sqrt{c}}{x} + 1}\right) b^2}{\sqrt{c}} + \\
& \frac{\operatorname{PolyLog}\left(2, \frac{2\sqrt{c}\left(1 - \frac{\sqrt{c}}{x}\right)}{(\sqrt{c} - \sqrt{c})\left(\frac{\sqrt{c}}{x} + 1\right)} + 1\right) b^2}{2\sqrt{c}} + \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{c}\left(\frac{\sqrt{c}}{x} + 1\right)}{(\sqrt{c} + \sqrt{c})\left(\frac{\sqrt{c}}{x} + 1\right)}\right) b^2}{2\sqrt{c}} + \\
& \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{(1-i)\left(\frac{\sqrt{c}}{x} + 1\right)}{1 - \frac{i\sqrt{c}}{x}}\right) b^2}{2\sqrt{c}} - \frac{i \operatorname{PolyLog}\left(2, \frac{2\sqrt{c}}{\sqrt{c-ix}} - 1\right) b^2}{\sqrt{c}} + \\
& \frac{\operatorname{PolyLog}\left(2, \frac{2\sqrt{c}}{x + \sqrt{c}} - 1\right) b^2}{\sqrt{c}} - \frac{2a \cot^{-1}\left(\frac{x}{\sqrt{c}}\right) b}{\sqrt{c}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) (2a - b \log\left(1 - \frac{c}{x^2}\right)) b}{\sqrt{c}} - \\
& \frac{(2a - b \log\left(1 - \frac{c}{x^2}\right)) b}{x} - \frac{a \log\left(\frac{c}{x^2} + 1\right) b}{x} + \frac{2ab}{x} - \frac{(2a - b \log\left(1 - \frac{c}{x^2}\right))^2}{4x}
\end{aligned}$$

input

Int[(a + b*ArcTanh[c/x^2])^2/x^2, x]

output

$$\begin{aligned}
& (2*a*b)/x - (2*a*b*\text{ArcCot}[x/\text{Sqrt}[c]])/\text{Sqrt}[c] - (2*b^2*\text{ArcCot}[x/\text{Sqrt}[c]])/\text{Sqrt}[c] \\
& - (2*b^2*\text{ArcCoth}[x/\text{Sqrt}[c]])/\text{Sqrt}[c] - (2*b^2*\text{ArcTan}[x/\text{Sqrt}[c]])/\text{Sqrt}[c] \\
& - (I*b^2*\text{ArcTan}[x/\text{Sqrt}[c]]^2)/\text{Sqrt}[c] + (2*b^2*\text{ArcTanh}[x/\text{Sqrt}[c]])/\text{Sqrt}[c] \\
& - (b^2*\text{ArcTanh}[x/\text{Sqrt}[c]]^2)/\text{Sqrt}[c] + (2*b^2*\text{ArcTan}[x/\text{Sqrt}[c]]*\text{Log}[2 - (2*\text{Sqrt}[c])/(\text{Sqrt}[c] - I*x)])/ \text{Sqrt}[c] \\
& - (b^2*\text{Log}[1 - c/x^2])/x + (b^2*\text{ArcCot}[x/\text{Sqrt}[c]]*\text{Log}[1 - c/x^2])/ \text{Sqrt}[c] - (b*(2*a - b*\text{Log}[1 - c/x^2]))/x \\
& + (b*\text{ArcTanh}[x/\text{Sqrt}[c]]*(2*a - b*\text{Log}[1 - c/x^2]))/\text{Sqrt}[c] - (2*a - b*\text{Log}[1 - c/x^2])^2/(4*x) \\
& - (a*b*\text{Log}[1 + c/x^2])/x + (b^2*\text{ArcCoth}[x/\text{Sqrt}[c]]*\text{Log}[1 + c/x^2])/ \text{Sqrt}[c] + (b^2*\text{ArcTan}[x/\text{Sqrt}[c]]*\text{Log}[1 + c/x^2])/ \text{Sqrt}[c] \\
& + (b^2*\text{Log}[1 - c/x^2]*\text{Log}[1 + c/x^2])/(2*x) - (b^2*\text{Log}[1 + c/x^2]^2)/(4*x) + (2*b^2*\text{ArcCot}[x/\text{Sqrt}[c]]*\text{Log}[2/(1 - (I*\text{Sqrt}[c])/x)])/ \text{Sqrt}[c] \\
& - (b^2*\text{ArcCoth}[x/\text{Sqrt}[c]]*\text{Log}[(1 + I)*(1 - \text{Sqrt}[c]/x)/(1 - (I*\text{Sqrt}[c])/x)])/ \text{Sqrt}[c] + (2*b^2*\text{ArcCoth}[x/\text{Sqrt}[c]]*\text{Log}[2/(1 + \text{Sqrt}[c]/x)])/ \text{Sqrt}[c] \\
& - (b^2*\text{ArcCoth}[x/\text{Sqrt}[c]]*\text{Log}[(-2*\text{Sqrt}[c]*(1 - \text{Sqrt}[-c]/x))/((\text{Sqrt}[-c] - \text{Sqrt}[c])*(1 + \text{Sqrt}[c]/x))])/ \text{Sqrt}[c] \\
& - (b^2*\text{ArcCoth}[x/\text{Sqrt}[c]]*\text{Log}[(2*\text{Sqrt}[c]*(1 + \text{Sqrt}[-c]/x))/((\text{Sqrt}[-c] + \text{Sqrt}[c])*(1 + \text{Sqrt}[c]/x))])/ \text{Sqrt}[c] \\
& - (b^2*\text{ArcCot}[x/\text{Sqrt}[c]]*\text{Log}[(1 - I)*(1 + \text{Sqrt}[c]/x)/(1 - (I*\text{Sqrt}[c])/x)])/ \text{Sqrt}[c] - (2*b^2*\text{ArcTanh}[x/\text{Sqrt}[c]]*\text{Log}[2 - (2*\text{Sqrt}[c])/(\text{Sqrt}[c] + x)])/ \text{Sqrt}[c] \\
& - (I*b^2*\text{PolyLog}[2, 1 - 2/(1 - (I*\text{Sqrt}[c])/x)])/ \text{Sqrt}[c] + ((I/2)*b^2*\text{PolyLog}[2, 1 - (1 + I)*(1 - \text{Sqrt}[c]/x)/(1 - (I*\text{Sqrt}[c])/x)])/ \text{Sqrt}[c] - (b^2*\text{PolyLog}[2, \dots
\end{aligned}$$

Defintions of rubi rules used

rule 2009 $\text{Int}[u_, x_Symbol] \text{ :> } \text{Simp}[\text{IntSum}[u, x], x] \text{ /; } \text{SumQ}[u]$

rule 6457 $\text{Int}[(a_.) + \text{ArcCoth}[(c_.)*(x_)^(n_)]*(b_.)^(p_)*(x_)^(m_.), x_Symbol] \text{ :> } \text{Int}[\text{ExpandIntegrand}[x^m*(a + b*(\text{Log}[1 + 1/(x^n*c)]/2) - b*(\text{Log}[1 - 1/(x^n*c)]/2))^p, x], x] \text{ /; } \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$

rule 6460 $\text{Int}[(a_.) + \text{ArcTanh}[(c_.)*(x_)^(n_)]*(b_.)^(p_)*(x_)^(m_.), x_Symbol] \text{ :> } \text{Int}[x^m*(a + b*\text{ArcCoth}[1/(x^n*c)])^p, x] \text{ /; } \text{FreeQ}[\{a, b, c, m\}, x] \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{ILtQ}[n, 0]$

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^2} dx$$

input `int((a+b*arctanh(c/x^2))^2/x^2,x)`

output `int((a+b*arctanh(c/x^2))^2/x^2,x)`

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^2} dx = \int \frac{(b \operatorname{arctanh}(\frac{c}{x^2}) + a)^2}{x^2} dx$$

input `integrate((a+b*arctanh(c/x^2))^2/x^2,x, algorithm="fricas")`

output `integral((b^2*arctanh(c/x^2)^2 + 2*a*b*arctanh(c/x^2) + a^2)/x^2, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^2} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x^2}))^2}{x^2} dx$$

input `integrate((a+b*atanh(c/x**2))**2/x**2,x)`

output `Integral((a + b*atanh(c/x**2))**2/x**2, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^2} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x^2}) + a)^2}{x^2} dx$$

input `integrate((a+b*arctanh(c/x^2))^2/x^2,x, algorithm="maxima")`

output `(c*(2*arctan(x/sqrt(c))/c^(3/2) - log((x - sqrt(c))/(x + sqrt(c)))/c^(3/2)) - 2*arctanh(c/x^2)/x)*a*b - 1/4*b^2*(log(x^2 - c)^2/x + integrate(-((x^2 - c)*log(x^2 + c)^2 + 2*(2*x^2 - (x^2 - c)*log(x^2 + c))*log(x^2 - c))/(x^4 - c*x^2), x)) - a^2/x`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^2} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x^2}) + a)^2}{x^2} dx$$

input `integrate((a+b*arctanh(c/x^2))^2/x^2,x, algorithm="giac")`

output `integrate((b*arctanh(c/x^2) + a)^2/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^2} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x^2}))^2}{x^2} dx$$

input `int((a + b*atanh(c/x^2))^2/x^2,x)`

output `int((a + b*atanh(c/x^2))^2/x^2, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^2} dx$$

$$= \frac{2\sqrt{c} \operatorname{atan}\left(\frac{x}{\sqrt{c}}\right) abx - 2\sqrt{c} \operatorname{atanh}\left(\frac{c}{x^2}\right) abx - 2 \operatorname{atanh}\left(\frac{c}{x^2}\right) abc - 2\sqrt{c} \log(\sqrt{c} - x) abx + \sqrt{c} \log(x^2 + c) abx}{cx}$$

input `int((a+b*atanh(c/x^2))^2/x^2,x)`

output `(2*sqrt(c)*atan(x/sqrt(c))*a*b*x - 2*sqrt(c)*atanh(c/x**2)*a*b*x - 2*atanh(c/x**2)*a*b*c - 2*sqrt(c)*log(sqrt(c) - x)*a*b*x + sqrt(c)*log(c + x**2)*a*b*x + int(atanh(c/x**2)**2/x**2,x)*b**2*c*x - a**2*c)/(c*x)`

$$3.180 \quad \int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^4} dx$$

Optimal result	1454
Mathematica [F]	1455
Rubi [A] (verified)	1456
Maple [F]	1459
Fricas [F]	1459
Sympy [F]	1459
Maxima [F]	1460
Giac [F]	1460
Mupad [F(-1)]	1460
Reduce [F]	1461

Optimal result

Integrand size = 16, antiderivative size = 1263

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^4} dx = \text{Too large to display}$$

output

```

1/3*b^2*arctanh(x/c^(1/2))*ln(1+c/x^2)/c^(3/2)-1/3*b^2*arctan(x/c^(1/2))*l
n(1+c/x^2)/c^(3/2)+1/3*b*arctanh(x/c^(1/2))*(2*a-b*ln(1-c/x^2))/c^(3/2)+1/
3*b^2*arctan(x/c^(1/2))*ln(1-c/x^2)/c^(3/2)-1/3*I*b^2*polylog(2,1-2*c^(1/2
))/(c^(1/2)-I*x))/c^(3/2)-1/3*I*b^2*polylog(2,I*x/c^(1/2))/c^(3/2)-1/3*b^2*
arctanh(x/c^(1/2))*ln(2*c^(1/2)*((-c)^(1/2)+x)/((-c)^(1/2)+c^(1/2))/(c^(1/
2)+x))/c^(3/2)-1/3*b^2*arctanh(x/c^(1/2))*ln(2*c^(1/2)*((-c)^(1/2)-x)/((-c
)^(1/2)-c^(1/2))/(c^(1/2)+x))/c^(3/2)-2/3*b^2*arctanh(x/c^(1/2))*ln(2-2*c^
(1/2)/(c^(1/2)+x))/c^(3/2)+2/3*b^2*arctanh(x/c^(1/2))*ln(2*c^(1/2)/(c^(1/2
)+x))/c^(3/2)+1/6*I*b^2*polylog(2,1+(-1+I)*(c^(1/2)+x)/(c^(1/2)-I*x))/c^(3
/2)+1/6*I*b^2*polylog(2,1-(1+I)*(c^(1/2)-x)/(c^(1/2)-I*x))/c^(3/2)+1/3*I*b
^2*polylog(2,-1+2*c^(1/2)/(c^(1/2)-I*x))/c^(3/2)+1/3*I*b^2*polylog(2,-I*x/
c^(1/2))/c^(3/2)+1/3*I*b^2*arctan(x/c^(1/2))^2/c^(3/2)-1/3*b^2*arctan(x/c^
(1/2))*ln((1+I)*(c^(1/2)-x)/(c^(1/2)-I*x))/c^(3/2)-1/3*b^2*arctan(x/c^(1/2
))*ln((1-I)*(c^(1/2)+x)/(c^(1/2)-I*x))/c^(3/2)-2/3*b^2*arctan(x/c^(1/2))*l
n(2-2*c^(1/2)/(c^(1/2)-I*x))/c^(3/2)+2/3*b^2*arctan(x/c^(1/2))*ln(2*c^(1/2
))/(c^(1/2)-I*x))/c^(3/2)-2/3*a*b*arctan(x/c^(1/2))/c^(3/2)-1/12*(2*a-b*ln(
1-c/x^2))^2/x^3+4/3*b^2*arctanh(x/c^(1/2))/c^(3/2)-1/3*b^2*arctanh(x/c^(1/
2))^2/c^(3/2)+4/3*b^2*arctan(x/c^(1/2))/c^(3/2)+1/3*b^2*polylog(2,x/c^(1/2
))/c^(3/2)-1/3*b^2*polylog(2,-x/c^(1/2))/c^(3/2)-1/3*b^2*polylog(2,1-2*c^(
1/2)/(c^(1/2)+x))/c^(3/2)+1/3*b^2*polylog(2,-1+2*c^(1/2)/(c^(1/2)+x))/c...

```

Mathematica [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^4} dx = \int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^4} dx$$

input

```
Integrate[(a + b*ArcTanh[c/x^2])^2/x^4, x]
```

output

```
Integrate[(a + b*ArcTanh[c/x^2])^2/x^4, x]
```

Rubi [A] (verified)

Time = 2.53 (sec) , antiderivative size = 1263, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6460, 6457, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}{x^4} dx$$

↓ 6460

$$\int \frac{\left(a + b \operatorname{coth}^{-1}\left(\frac{x^2}{c}\right)\right)^2}{x^4} dx$$

↓ 6457

$$\int \left(\frac{\left(2a - b \log\left(1 - \frac{c}{x^2}\right)\right)^2}{4x^4} - \frac{b \log\left(\frac{c}{x^2} + 1\right) \left(b \log\left(1 - \frac{c}{x^2}\right) - 2a\right)}{2x^4} + \frac{b^2 \log^2\left(\frac{c}{x^2} + 1\right)}{4x^4} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{i \arctan\left(\frac{x}{\sqrt{c}}\right)^2 b^2}{3c^{3/2}} - \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right)^2 b^2}{3c^{3/2}} - \frac{\log^2\left(\frac{c}{x^2} + 1\right) b^2}{12x^3} + \frac{4 \arctan\left(\frac{x}{\sqrt{c}}\right) b^2}{3c^{3/2}} + \\
& \frac{4 \operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) b^2}{3c^{3/2}} - \frac{2 \arctan\left(\frac{x}{\sqrt{c}}\right) \log\left(2 - \frac{2\sqrt{c}}{\sqrt{c-ix}}\right) b^2}{3c^{3/2}} + \frac{\arctan\left(\frac{x}{\sqrt{c}}\right) \log\left(1 - \frac{c}{x^2}\right) b^2}{3c^{3/2}} + \\
& \frac{\log\left(1 - \frac{c}{x^2}\right) b^2}{3cx} - \frac{\log\left(1 - \frac{c}{x^2}\right) b^2}{9x^3} - \frac{\arctan\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{c}{x^2} + 1\right) b^2}{3c^{3/2}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{c}{x^2} + 1\right) b^2}{3c^{3/2}} + \\
& \frac{\log\left(1 - \frac{c}{x^2}\right) \log\left(\frac{c}{x^2} + 1\right) b^2}{6x^3} - \frac{2 \log\left(\frac{c}{x^2} + 1\right) b^2}{3cx} + \frac{2 \arctan\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{2\sqrt{c}}{\sqrt{c-ix}}\right) b^2}{3c^{3/2}} - \\
& \frac{\arctan\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{(1+i)(\sqrt{c-x})}{\sqrt{c-ix}}\right) b^2}{3c^{3/2}} + \frac{2 \operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{2\sqrt{c}}{x+\sqrt{c}}\right) b^2}{3c^{3/2}} - \\
& \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{2\sqrt{c}(\sqrt{c-x})}{(\sqrt{-c-\sqrt{c}})(x+\sqrt{c})}\right) b^2}{3c^{3/2}} - \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{2\sqrt{c}(x+\sqrt{-c})}{(\sqrt{-c+\sqrt{c}})(x+\sqrt{c})}\right) b^2}{3c^{3/2}} - \\
& \frac{\arctan\left(\frac{x}{\sqrt{c}}\right) \log\left(\frac{(1-i)(x+\sqrt{c})}{\sqrt{c-ix}}\right) b^2}{3c^{3/2}} - \frac{2 \operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) \log\left(2 - \frac{2\sqrt{c}}{x+\sqrt{c}}\right) b^2}{3c^{3/2}} - \\
& \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{c}}{\sqrt{c-ix}}\right) b^2}{3c^{3/2}} + \frac{i \operatorname{PolyLog}\left(2, \frac{2\sqrt{c}}{\sqrt{c-ix}} - 1\right) b^2}{3c^{3/2}} + \\
& \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{(1+i)(\sqrt{c-x})}{\sqrt{c-ix}}\right) b^2}{6c^{3/2}} - \frac{\operatorname{PolyLog}\left(2, -\frac{x}{\sqrt{c}}\right) b^2}{3c^{3/2}} + \frac{i \operatorname{PolyLog}\left(2, -\frac{ix}{\sqrt{c}}\right) b^2}{3c^{3/2}} - \\
& \frac{i \operatorname{PolyLog}\left(2, \frac{ix}{\sqrt{c}}\right) b^2}{3c^{3/2}} + \frac{\operatorname{PolyLog}\left(2, \frac{x}{\sqrt{c}}\right) b^2}{3c^{3/2}} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{c}}{x+\sqrt{c}}\right) b^2}{3c^{3/2}} + \\
& \frac{\operatorname{PolyLog}\left(2, \frac{2\sqrt{c}}{x+\sqrt{c}} - 1\right) b^2}{3c^{3/2}} + \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{c}(\sqrt{c-x})}{(\sqrt{-c-\sqrt{c}})(x+\sqrt{c})}\right) b^2}{6c^{3/2}} + \\
& \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{c}(x+\sqrt{-c})}{(\sqrt{-c+\sqrt{c}})(x+\sqrt{c})}\right) b^2}{6c^{3/2}} + \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{(1-i)(x+\sqrt{c})}{\sqrt{c-ix}}\right) b^2}{6c^{3/2}} - \\
& \frac{2a \arctan\left(\frac{x}{\sqrt{c}}\right) b}{3c^{3/2}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{c}}\right) (2a - b \log\left(1 - \frac{c}{x^2}\right)) b}{3c^{3/2}} - \frac{(2a - b \log\left(1 - \frac{c}{x^2}\right)) b}{3cx} \\
& \frac{(2a - b \log\left(1 - \frac{c}{x^2}\right)) b}{9x^3} - \frac{a \log\left(\frac{c}{x^2} + 1\right) b}{3x^3} - \frac{2ab}{3cx} + \frac{2ab}{9x^3} - \frac{(2a - b \log\left(1 - \frac{c}{x^2}\right))^2}{12x^3}
\end{aligned}$$

input

Int[(a + b*ArcTanh[c/x^2])^2/x^4, x]

output

$$\begin{aligned}
& (2ab)/(9x^3) - (2ab)/(3cx) - (2ab \operatorname{ArcTan}[x/\sqrt{c}])/(3c^{3/2}) \\
& + (4b^2 \operatorname{ArcTan}[x/\sqrt{c}])/(3c^{3/2}) + ((I/3)b^2 \operatorname{ArcTan}[x/\sqrt{c}]^2)/ \\
& c^{3/2} + (4b^2 \operatorname{ArcTanh}[x/\sqrt{c}])/(3c^{3/2}) - (b^2 \operatorname{ArcTanh}[x/\sqrt{c}] \\
& ^2)/(3c^{3/2}) - (2b^2 \operatorname{ArcTan}[x/\sqrt{c}] \operatorname{Log}[2 - (2\sqrt{c})/(\sqrt{c} - \\
& Ix)])/(3c^{3/2}) - (b^2 \operatorname{Log}[1 - c/x^2])/(9x^3) + (b^2 \operatorname{Log}[1 - c/x^2])/(\\
& 3cx) + (b^2 \operatorname{ArcTan}[x/\sqrt{c}] \operatorname{Log}[1 - c/x^2])/(3c^{3/2}) - (b(2a - b \\
& \operatorname{Log}[1 - c/x^2]))/(9x^3) - (b(2a - b \operatorname{Log}[1 - c/x^2]))/(3cx) + (b \operatorname{ArcTa} \\
& \operatorname{nh}[x/\sqrt{c}] * (2a - b \operatorname{Log}[1 - c/x^2]))/(3c^{3/2}) - (2a - b \operatorname{Log}[1 - c/x \\
& ^2])^2/(12x^3) - (ab \operatorname{Log}[1 + c/x^2])/(3x^3) - (2b^2 \operatorname{Log}[1 + c/x^2])/(3 \\
& cx) - (b^2 \operatorname{ArcTan}[x/\sqrt{c}] \operatorname{Log}[1 + c/x^2])/(3c^{3/2}) + (b^2 \operatorname{ArcTanh}[\\
& x/\sqrt{c}] \operatorname{Log}[1 + c/x^2])/(3c^{3/2}) + (b^2 \operatorname{Log}[1 - c/x^2] \operatorname{Log}[1 + c/x^2 \\
&])/(6x^3) - (b^2 \operatorname{Log}[1 + c/x^2]^2)/(12x^3) + (2b^2 \operatorname{ArcTan}[x/\sqrt{c}] \operatorname{Lo} \\
& \operatorname{g}[(2\sqrt{c})/(\sqrt{c} - Ix)])/(3c^{3/2}) - (b^2 \operatorname{ArcTan}[x/\sqrt{c}] \operatorname{Log}[(\\
& (1 + I)(\sqrt{c} - x))/(\sqrt{c} - Ix)])/(3c^{3/2}) + (2b^2 \operatorname{ArcTanh}[x/\sqrt{c} \\
&] \operatorname{Log}[(2\sqrt{c})/(\sqrt{c} + x)])/(3c^{3/2}) - (b^2 \operatorname{ArcTanh}[x/\sqrt{c} \\
&] \operatorname{Log}[(2\sqrt{c} * (\sqrt{-c} - x))/((\sqrt{-c} - \sqrt{c}) * (\sqrt{c} + x))])/(\\
& 3c^{3/2}) - (b^2 \operatorname{ArcTanh}[x/\sqrt{c}] \operatorname{Log}[(2\sqrt{c} * (\sqrt{-c} + x))/((\sqrt{c} \\
& - \sqrt{-c}) * (\sqrt{c} + x))])/(3c^{3/2}) - (b^2 \operatorname{ArcTan}[x/\sqrt{c}] \operatorname{Log}[\\
& ((1 - I)(\sqrt{c} + x))/(\sqrt{c} - Ix)])/(3c^{3/2}) - (2b^2 \operatorname{ArcTanh}[x/\sqrt{c}] \\
& \operatorname{Log}[2 - (2\sqrt{c})/(\sqrt{c} + x)])/(3c^{3/2}) - ((I/3)b^2 \operatorname{Po...}
\end{aligned}$$

Defintions of rubi rules used

rule 2009 $\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$

rule 6457 $\operatorname{Int}[(a_.) + \operatorname{ArcCoth}[(c_.)(x_)^{(n)}] * (b_.)^{(p)} * (x_)^{(m_.)}, x_Symbol] \rightarrow$
 $\operatorname{Int}[\operatorname{ExpandIntegrand}[x^m * (a + b * (\operatorname{Log}[1 + 1/(x^n * c)]/2) - b * (\operatorname{Log}[1 - 1/(x^n * c)]/2))^{(p)}, x], x] /; \operatorname{FreeQ}[\{a, b, c\}, x] \&\& \operatorname{IGtQ}[p, 1] \&\& \operatorname{IGtQ}[n, 0] \&\& \operatorname{IntegerQ}[m]$

rule 6460 $\operatorname{Int}[(a_.) + \operatorname{ArcTanh}[(c_.)(x_)^{(n)}] * (b_.)^{(p)} * (x_)^{(m_.)}, x_Symbol] \rightarrow$
 $\operatorname{Int}[x^m * (a + b * \operatorname{ArcCoth}[1/(x^n * c)])^{(p)}, x] /; \operatorname{FreeQ}[\{a, b, c, m\}, x] \&\& \operatorname{IGtQ}[p, 1] \&\& \operatorname{ILtQ}[n, 0]$

Maple [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^4} dx$$

input `int((a+b*arctanh(c/x^2))^2/x^4,x)`

output `int((a+b*arctanh(c/x^2))^2/x^4,x)`

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^4} dx = \int \frac{(b \operatorname{arctanh}(\frac{c}{x^2}) + a)^2}{x^4} dx$$

input `integrate((a+b*arctanh(c/x^2))^2/x^4,x, algorithm="fricas")`

output `integral((b^2*arctanh(c/x^2)^2 + 2*a*b*arctanh(c/x^2) + a^2)/x^4, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^4} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x^2}))^2}{x^4} dx$$

input `integrate((a+b*atanh(c/x**2))**2/x**4,x)`

output `Integral((a + b*atanh(c/x**2))**2/x**4, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^4} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x^2}) + a)^2}{x^4} dx$$

input `integrate((a+b*arctanh(c/x^2))^2/x^4,x, algorithm="maxima")`

output `-1/3*(c*(2*arctan(x/sqrt(c))/c^(5/2) + log((x - sqrt(c))/(x + sqrt(c))))/c^(5/2) + 4/(c^2*x)) + 2*arctanh(c/x^2)/x^3)*a*b - 1/12*b^2*(log(x^2 - c)^2/x^3 + 3*integrate(-1/3*(3*(x^2 - c)*log(x^2 + c)^2 + 2*(2*x^2 - 3*(x^2 - c))*log(x^2 + c))*log(x^2 - c))/(x^6 - c*x^4), x)) - 1/3*a^2/x^3`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^4} dx = \int \frac{(b \operatorname{artanh}(\frac{c}{x^2}) + a)^2}{x^4} dx$$

input `integrate((a+b*arctanh(c/x^2))^2/x^4,x, algorithm="giac")`

output `integrate((b*arctanh(c/x^2) + a)^2/x^4, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^4} dx = \int \frac{(a + b \operatorname{atanh}(\frac{c}{x^2}))^2}{x^4} dx$$

input `int((a + b*atanh(c/x^2))^2/x^4,x)`

output `int((a + b*atanh(c/x^2))^2/x^4, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(\frac{c}{x^2}))^2}{x^4} dx$$

$$= \frac{-2\sqrt{c} \operatorname{atan}\left(\frac{x}{\sqrt{c}}\right) ab x^3 - 2\sqrt{c} \operatorname{atanh}\left(\frac{c}{x^2}\right) ab x^3 - 2 \operatorname{atanh}\left(\frac{c}{x^2}\right) ab c^2 - 2\sqrt{c} \log(\sqrt{c} - x) ab x^3 + \sqrt{c} \log(x^2)}{3c^2 x^3}$$

input `int((a+b*atanh(c/x^2))^2/x^4,x)`

output `(- 2*sqrt(c)*atan(x/sqrt(c))*a*b*x**3 - 2*sqrt(c)*atanh(c/x**2)*a*b*x**3 - 2*atanh(c/x**2)*a*b*c**2 - 2*sqrt(c)*log(sqrt(c)- x)*a*b*x**3 + sqrt(c)*log(c + x**2)*a*b*x**3 + 3*int(atanh(c/x**2)**2/x**4,x)*b**2*c**2*x**3 - a**2*c**2 - 4*a*b*c*x**2)/(3*c**2*x**3)`

3.181 $\int (dx)^m \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right)\right)^3 dx$

Optimal result	1462
Mathematica [N/A]	1462
Rubi [N/A]	1463
Maple [N/A]	1463
Fricas [N/A]	1464
Sympy [N/A]	1464
Maxima [N/A]	1464
Giac [N/A]	1465
Mupad [N/A]	1465
Reduce [N/A]	1466

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int (dx)^m \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right)\right)^3 dx = \operatorname{Int}\left(\left(dx\right)^m \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right)\right)^3, x\right)$$

output `Defer(Int)((d*x)^m*(a+b*arctanh(c/x^2))^3,x)`

Mathematica [N/A]

Not integrable

Time = 1.98 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right)\right)^3 dx = \int (dx)^m \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right)\right)^3 dx$$

input `Integrate[(d*x)^m*(a + b*ArcTanh[c/x^2])^3,x]`

output `Integrate[(d*x)^m*(a + b*ArcTanh[c/x^2])^3, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^3 dx$$

↓ 6468

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^3 dx$$

input `Int[(d*x)^m*(a + b*ArcTanh[c/x^2])^3,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^3 dx$$

input `int((d*x)^m*(a+b*arctanh(c/x^2))^3,x)`

output `int((d*x)^m*(a+b*arctanh(c/x^2))^3,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 50, normalized size of antiderivative = 2.78

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^3 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x^2} \right) + a \right)^3 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c/x^2))^3,x, algorithm="fricas")`

output `integral((b^3*arctanh(c/x^2)^3 + 3*a*b^2*arctanh(c/x^2)^2 + 3*a^2*b*arctanh(c/x^2) + a^3)*(d*x)^m, x)`

Sympy [N/A]

Not integrable

Time = 54.85 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^3 dx = \int (dx)^m \left(a + b \operatorname{atanh} \left(\frac{c}{x^2} \right) \right)^3 dx$$

input `integrate((d*x)**m*(a+b*atanh(c/x**2))**3,x)`

output `Integral((d*x)**m*(a + b*atanh(c/x**2))**3, x)`

Maxima [N/A]

Not integrable

Time = 2.54 (sec) , antiderivative size = 398, normalized size of antiderivative = 22.11

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^3 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x^2} \right) + a \right)^3 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c/x^2))^3,x, algorithm="maxima")`

output

```
-1/8*b^3*d^m*x*x^m*log(x^2 - c)^3/(m + 1) + (d*x)^(m + 1)*a^3/(d*(m + 1))
+ integrate(1/8*((b^3*d^m*(m + 1)*x^2 - b^3*c*d^m*(m + 1))*x^m*log(x^2 + c)
)^3 + 6*(a*b^2*d^m*(m + 1)*x^2 - a*b^2*c*d^m*(m + 1))*x^m*log(x^2 + c)^2 +
12*(a^2*b*d^m*(m + 1)*x^2 - a^2*b*c*d^m*(m + 1))*x^m*log(x^2 + c) + 3*((b
^3*d^m*(m + 1)*x^2 - b^3*c*d^m*(m + 1))*x^m*log(x^2 + c) - 2*(a*b^2*c*d^m*
(m + 1) - (a*b^2*d^m*(m + 1) + b^3*d^m)*x^2)*x^m*log(x^2 - c)^2 - 3*((b^3
*d^m*(m + 1)*x^2 - b^3*c*d^m*(m + 1))*x^m*log(x^2 + c)^2 + 4*(a*b^2*d^m*(m
+ 1)*x^2 - a*b^2*c*d^m*(m + 1))*x^m*log(x^2 + c) + 4*(a^2*b*d^m*(m + 1)*x
^2 - a^2*b*c*d^m*(m + 1))*x^m*log(x^2 - c))/((m + 1)*x^2 - c*(m + 1)), x)
```

Giac [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^3 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x^2} \right) + a \right)^3 (dx)^m dx$$

input

```
integrate((d*x)^m*(a+b*arctanh(c/x^2))^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c/x^2) + a)^3*(d*x)^m, x)
```

Mupad [N/A]

Not integrable

Time = 3.90 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^3 dx = \int (dx)^m \left(a + b \operatorname{atanh} \left(\frac{c}{x^2} \right) \right)^3 dx$$

input

```
int((d*x)^m*(a + b*atanh(c/x^2))^3,x)
```

output

```
int((d*x)^m*(a + b*atanh(c/x^2))^3, x)
```

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 133, normalized size of antiderivative = 7.39

$$\int (dx)^m \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right) \right)^3 dx$$

$$= \frac{d^m \left(x^m a^3 x + 3 \left(\int x^m \operatorname{atanh}\left(\frac{c}{x^2}\right) dx \right) a^2 b m + 3 \left(\int x^m \operatorname{atanh}\left(\frac{c}{x^2}\right) dx \right) a^2 b + \left(\int x^m \operatorname{atanh}\left(\frac{c}{x^2}\right)^3 dx \right) b^3 m + \dots \right)}{m + 1}$$

input

```
int((d*x)^m*(a+b*atanh(c/x^2))^3,x)
```

output

```
(d**m*(x**m*a**3*x + 3*int(x**m*atanh(c/x**2),x)*a**2*b*m + 3*int(x**m*atanh(c/x**2),x)*a**2*b + int(x**m*atanh(c/x**2)**3,x)*b**3*m + int(x**m*atanh(c/x**2)**3,x)*b**3 + 3*int(x**m*atanh(c/x**2)**2,x)*a*b**2*m + 3*int(x**m*atanh(c/x**2)**2,x)*a*b**2))/(m + 1)
```

3.182 $\int (dx)^m \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2 dx$

Optimal result	1467
Mathematica [N/A]	1467
Rubi [N/A]	1468
Maple [N/A]	1468
Fricas [N/A]	1469
Sympy [N/A]	1469
Maxima [N/A]	1469
Giac [N/A]	1470
Mupad [N/A]	1470
Reduce [N/A]	1471

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int (dx)^m \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2 dx = \operatorname{Int}\left(\left(dx\right)^m \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2, x\right)$$

output `Defer(Int)((d*x)^m*(a+b*arctanh(c/x^2))^2,x)`

Mathematica [N/A]

Not integrable

Time = 1.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2 dx = \int (dx)^m \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2 dx$$

input `Integrate[(d*x)^m*(a + b*ArcTanh[c/x^2])^2,x]`

output `Integrate[(d*x)^m*(a + b*ArcTanh[c/x^2])^2, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

↓ 6468

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input `Int[(d*x)^m*(a + b*ArcTanh[c/x^2])^2,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input `int((d*x)^m*(a+b*arctanh(c/x^2))^2,x)`

output `int((d*x)^m*(a+b*arctanh(c/x^2))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.89

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(b \operatorname{arctanh} \left(\frac{c}{x^2} \right) + a \right)^2 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c/x^2))^2,x, algorithm="fricas")`

output `integral((b^2*arctanh(c/x^2)^2 + 2*a*b*arctanh(c/x^2) + a^2)*(d*x)^m, x)`

Sympy [N/A]

Not integrable

Time = 35.63 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int (dx)^m \left(a + b \operatorname{atanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input `integrate((d*x)**m*(a+b*atanh(c/x**2))**2,x)`

output `Integral((d*x)**m*(a + b*atanh(c/x**2))**2, x)`

Maxima [N/A]

Not integrable

Time = 1.40 (sec) , antiderivative size = 228, normalized size of antiderivative = 12.67

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(b \operatorname{arctanh} \left(\frac{c}{x^2} \right) + a \right)^2 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c/x^2))^2,x, algorithm="maxima")`

output

```
1/4*b^2*d^m*x*x^m*log(x^2 - c)^2/(m + 1) + (d*x)^(m + 1)*a^2/(d*(m + 1)) -
integrate(-1/4*((b^2*d^m*(m + 1)*x^2 - b^2*c*d^m*(m + 1))*x^m*log(x^2 + c)
)^2 + 4*(a*b*d^m*(m + 1)*x^2 - a*b*c*d^m*(m + 1))*x^m*log(x^2 + c) - 2*((b
^2*d^m*(m + 1)*x^2 - b^2*c*d^m*(m + 1))*x^m*log(x^2 + c) - 2*(a*b*c*d^m*(m
+ 1) - (a*b*d^m*(m + 1) + b^2*d^m)*x^2)*x^m*log(x^2 - c))/((m + 1)*x^2 -
c*(m + 1)), x)
```

Giac [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x^2} \right) + a \right)^2 (dx)^m dx$$

input

```
integrate((d*x)^m*(a+b*arctanh(c/x^2))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c/x^2) + a)^2*(d*x)^m, x)
```

Mupad [N/A]

Not integrable

Time = 3.97 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right)^2 dx = \int (dx)^m \left(a + b \operatorname{atanh} \left(\frac{c}{x^2} \right) \right)^2 dx$$

input

```
int((d*x)^m*(a + b*atanh(c/x^2))^2,x)
```

output

```
int((d*x)^m*(a + b*atanh(c/x^2))^2, x)
```

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 88, normalized size of antiderivative = 4.89

$$\int (dx)^m \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right) \right)^2 dx$$

$$= \frac{d^m \left(x^m a^2 x + 2 \left(\int x^m \operatorname{atanh}\left(\frac{c}{x^2}\right) dx \right) abm + 2 \left(\int x^m \operatorname{atanh}\left(\frac{c}{x^2}\right) dx \right) ab + \left(\int x^m \operatorname{atanh}\left(\frac{c}{x^2}\right)^2 dx \right) b^2 m + \left(\int x^m \operatorname{atanh}\left(\frac{c}{x^2}\right) dx \right) b^2 + \left(\int x^m \operatorname{atanh}\left(\frac{c}{x^2}\right) dx \right)^2 \right)}{m + 1}$$

input

```
int((d*x)^m*(a+b*atanh(c/x^2))^2,x)
```

output

```
(d**m*(x**m*a**2*x + 2*int(x**m*atanh(c/x**2),x)*a*b*m + 2*int(x**m*atanh(c/x**2),x)*a*b + int(x**m*atanh(c/x**2)**2,x)*b**2*m + int(x**m*atanh(c/x**2)**2,x)*b**2))/(m + 1)
```

3.183 $\int (dx)^m \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right) \right) dx$

Optimal result	1472
Mathematica [A] (verified)	1472
Rubi [A] (verified)	1473
Maple [F]	1474
Fricas [F]	1474
Sympy [F]	1475
Maxima [F]	1475
Giac [F]	1475
Mupad [F(-1)]	1476
Reduce [F]	1476

Optimal result

Integrand size = 16, antiderivative size = 75

$$\int (dx)^m \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right) \right) dx$$

$$= \frac{(dx)^{1+m} \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right) \right)}{d(1+m)} - \frac{2bcd(dx)^{-1+m} \operatorname{Hypergeometric2F1}\left(1, \frac{1-m}{4}, \frac{5-m}{4}, \frac{c^2}{x^4}\right)}{1-m^2}$$

output $(d*x)^{(1+m)}*(a+b*\operatorname{arctanh}(c/x^2))/d/(1+m)-2*b*c*d*(d*x)^{(-1+m)}*\operatorname{hypergeom}\left(1, 1/4-1/4*m\right), [5/4-1/4*m], c^2/x^4)/(-m^2+1)$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.91

$$\int (dx)^m \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right) \right) dx$$

$$= \frac{(dx)^m \left((-1+m)x^2 \left(a + \operatorname{barctanh}\left(\frac{c}{x^2}\right) \right) + 2bc \operatorname{Hypergeometric2F1}\left(1, \frac{1}{4} - \frac{m}{4}, \frac{5}{4} - \frac{m}{4}, \frac{c^2}{x^4}\right) \right)}{(-1+m)(1+m)x}$$

input $\operatorname{Integrate}[(d*x)^m*(a + b*\operatorname{ArcTanh}[c/x^2]), x]$

output

$$\frac{((dx)^m((-1+m)x^2(a+b\text{ArcTanh}[c/x^2]))+2bc\text{Hypergeometric2F1}[1, 1/4-m/4, 5/4-m/4, c^2/x^4])}{((-1+m)(1+m)x)}$$
Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.04, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6464, 862, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right) \right) dx$$

$$\downarrow 6464$$

$$\frac{2bcd^2 \int \frac{(dx)^{m-2}}{1-\frac{c^2}{x^4}} dx}{m+1} + \frac{(dx)^{m+1} \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right) \right)}{d(m+1)}$$

$$\downarrow 862$$

$$\frac{(dx)^{m+1} \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right) \right)}{d(m+1)} - \frac{2bcd \left(\frac{1}{x}\right)^{m-1} (dx)^{m-1} \int \frac{\left(\frac{1}{x}\right)^{-m}}{1-\frac{c^2}{x^4}} d\frac{1}{x}}{m+1}$$

$$\downarrow 888$$

$$\frac{(dx)^{m+1} \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right) \right)}{d(m+1)} - \frac{2bcd(dx)^{m-1} \text{Hypergeometric2F1}\left(1, \frac{1-m}{4}, \frac{5-m}{4}, \frac{c^2}{x^4}\right)}{(1-m)(m+1)}$$

input

$$\text{Int}[(dx)^m(a + b\text{ArcTanh}[c/x^2]), x]$$

output

$$\frac{(dx)^{(1+m)}(a + b\text{ArcTanh}[c/x^2])}{d(1+m)} - \frac{(2bc\text{Hypergeometric2F1}[1, (1-m)/4, (5-m)/4, c^2/x^4])}{((1-m)(1+m))}$$

Definitions of rubi rules used

rule 862 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-c^(-1))*(c*x)^(m + 1)*(1/x)^(m + 1) Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x], x] /; FreeQ[{a, b, c, m, p}, x] && ILtQ[n, 0] && !RationalQ[m]`

rule 888 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 6464 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_))*((d_)*(x_)^(m_), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(d*(m + 1))), x] - Simp[b*c*(n/(d^n*(m + 1))) Int[(d*x)^(m + n)/(1 - c^2*x^(2*n)), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegerQ[n] && NeQ[m, -1]`

Maple [F]

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx$$

input `int((d*x)^m*(a+b*arctanh(c/x^2)),x)`

output `int((d*x)^m*(a+b*arctanh(c/x^2)),x)`

Fricas [F]

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \int \left(b \operatorname{arctanh} \left(\frac{c}{x^2} \right) + a \right) (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c/x^2)),x, algorithm="fricas")`

output `integral((b*arctanh(c/x^2) + a)*(d*x)^m, x)`

Sympy [F]

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \int (dx)^m \left(a + b \operatorname{atanh} \left(\frac{c}{x^2} \right) \right) dx$$

input `integrate((d*x)**m*(a+b*atanh(c/x**2)),x)`

output `Integral((d*x)**m*(a + b*atanh(c/x**2)), x)`

Maxima [F]

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x^2} \right) + a \right) (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c/x^2)),x, algorithm="maxima")`

output `1/2*(4*c*d^m*integrate(x^2*x^m/((m + 1)*x^4 - c^2*(m + 1)), x) + (d^m*x*x^m*log(x^2 + c) - d^m*x*x^m*log(x^2 - c))/(m + 1)*b + (d*x)^(m + 1)*a/(d*(m + 1))`

Giac [F]

$$\int (dx)^m \left(a + b \operatorname{arctanh} \left(\frac{c}{x^2} \right) \right) dx = \int \left(b \operatorname{artanh} \left(\frac{c}{x^2} \right) + a \right) (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c/x^2)),x, algorithm="giac")`

output `integrate((b*arctanh(c/x^2) + a)*(d*x)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (dx)^m \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right) \right) dx = \int (dx)^m \left(a + b \operatorname{atanh}\left(\frac{c}{x^2}\right) \right) dx$$

input `int((d*x)^m*(a + b*atanh(c/x^2)),x)`output `int((d*x)^m*(a + b*atanh(c/x^2)), x)`**Reduce [F]**

$$\begin{aligned} & \int (dx)^m \left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right) \right) dx \\ &= \frac{d^m \left(x^m a x + \left(\int x^m \operatorname{atanh}\left(\frac{c}{x^2}\right) dx \right) b m + \left(\int x^m \operatorname{atanh}\left(\frac{c}{x^2}\right) dx \right) b \right)}{m + 1} \end{aligned}$$

input `int((d*x)^m*(a+b*atanh(c/x^2)),x)`output `(d**m*(x**m*a*x + int(x**m*atanh(c/x**2),x)*b*m + int(x**m*atanh(c/x**2),x)*b))/(m + 1)`

$$3.184 \quad \int \frac{(dx)^m}{a+b\operatorname{arctanh}\left(\frac{c}{x^2}\right)} dx$$

Optimal result	1477
Mathematica [N/A]	1477
Rubi [N/A]	1478
Maple [N/A]	1478
Fricas [N/A]	1479
Sympy [N/A]	1479
Maxima [N/A]	1479
Giac [N/A]	1480
Mupad [N/A]	1480
Reduce [N/A]	1481

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{(dx)^m}{a + \operatorname{arctanh}\left(\frac{c}{x^2}\right)} dx = \operatorname{Int}\left(\frac{(dx)^m}{a + \operatorname{arctanh}\left(\frac{c}{x^2}\right)}, x\right)$$

output `Defer(Int)((d*x)^m/(a+b*arctanh(c/x^2)), x)`

Mathematica [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + \operatorname{arctanh}\left(\frac{c}{x^2}\right)} dx = \int \frac{(dx)^m}{a + \operatorname{arctanh}\left(\frac{c}{x^2}\right)} dx$$

input `Integrate[(d*x)^m/(a + b*ArcTanh[c/x^2]), x]`

output `Integrate[(d*x)^m/(a + b*ArcTanh[c/x^2]), x]`

Rubi [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)} dx$$

↓ 6468

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)} dx$$

input `Int[(d*x)^m/(a + b*ArcTanh[c/x^2]),x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)} dx$$

input `int((d*x)^m/(a+b*arctanh(c/x^2)),x)`

output `int((d*x)^m/(a+b*arctanh(c/x^2)),x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)} dx = \int \frac{(dx)^m}{b \operatorname{artanh}\left(\frac{c}{x^2}\right) + a} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c/x^2)),x, algorithm="fricas")`

output `integral((d*x)^m/(b*arctanh(c/x^2) + a), x)`

Sympy [N/A]

Not integrable

Time = 45.61 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)} dx = \int \frac{(dx)^m}{a + b \operatorname{atanh}\left(\frac{c}{x^2}\right)} dx$$

input `integrate((d*x)**m/(a+b*atanh(c/x**2)),x)`

output `Integral((d*x)**m/(a + b*atanh(c/x**2)), x)`

Maxima [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)} dx = \int \frac{(dx)^m}{b \operatorname{artanh}\left(\frac{c}{x^2}\right) + a} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c/x^2)),x, algorithm="maxima")`

output `integrate((d*x)^m/(b*arctanh(c/x^2) + a), x)`

Giac [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)} dx = \int \frac{(dx)^m}{b \operatorname{artanh}\left(\frac{c}{x^2}\right) + a} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c/x^2)),x, algorithm="giac")`

output `integrate((d*x)^m/(b*arctanh(c/x^2) + a), x)`

Mupad [N/A]

Not integrable

Time = 3.42 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)} dx = \int \frac{(dx)^m}{a + b \operatorname{atanh}\left(\frac{c}{x^2}\right)} dx$$

input `int((d*x)^m/(a + b*atanh(c/x^2)),x)`

output `int((d*x)^m/(a + b*atanh(c/x^2)), x)`

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)} dx = d^m \left(\int \frac{x^m}{\operatorname{atanh}\left(\frac{c}{x^2}\right) b + a} dx \right)$$

input `int((d*x)^m/(a+b*atanh(c/x^2)),x)`output `d**m*int(x**m/(atanh(c/x**2)*b + a),x)`

$$3.185 \quad \int \frac{(dx)^m}{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2} dx$$

Optimal result	1482
Mathematica [N/A]	1482
Rubi [N/A]	1483
Maple [N/A]	1483
Fricas [N/A]	1484
Sympy [F(-1)]	1484
Maxima [N/A]	1484
Giac [N/A]	1485
Mupad [N/A]	1485
Reduce [N/A]	1486

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{(dx)^m}{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2} dx = \operatorname{Int}\left(\frac{(dx)^m}{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2}, x\right)$$

output `Defer(Int)((d*x)^m/(a+b*arctanh(c/x^2))^2,x)`

Mathematica [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2} dx = \int \frac{(dx)^m}{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2} dx$$

input `Integrate[(d*x)^m/(a + b*ArcTanh[c/x^2])^2,x]`

output `Integrate[(d*x)^m/(a + b*ArcTanh[c/x^2])^2, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(dx)^m}{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2} dx$$

↓ 6468

$$\int \frac{(dx)^m}{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2} dx$$

input `Int[(d*x)^m/(a + b*ArcTanh[c/x^2])^2,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{(dx)^m}{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2} dx$$

input `int((d*x)^m/(a+b*arctanh(c/x^2))^2,x)`

output `int((d*x)^m/(a+b*arctanh(c/x^2))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.00

$$\int \frac{(dx)^m}{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2} dx = \int \frac{(dx)^m}{\left(b \operatorname{arctanh}\left(\frac{c}{x^2}\right) + a\right)^2} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c/x^2))^2,x, algorithm="fricas")`

output `integral((d*x)^m/(b^2*arctanh(c/x^2)^2 + 2*a*b*arctanh(c/x^2) + a^2), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(dx)^m}{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2} dx = \text{Timed out}$$

input `integrate((d*x)**m/(a+b*atanh(c/x**2))**2,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 128, normalized size of antiderivative = 7.11

$$\int \frac{(dx)^m}{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2} dx = \int \frac{(dx)^m}{\left(b \operatorname{arctanh}\left(\frac{c}{x^2}\right) + a\right)^2} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c/x^2))^2,x, algorithm="maxima")`

output

```
(d^m*x^4 - c^2*d^m)*x^m/(b^2*c*x*log(x^2 + c) - b^2*c*x*log(x^2 - c) + 2*a
*b*c*x) + integrate(-(d^m*(m + 3)*x^4 - c^2*d^m*(m - 1))*x^m/(b^2*c*x^2*lo
g(x^2 + c) - b^2*c*x^2*log(x^2 - c) + 2*a*b*c*x^2), x)
```

Giac [N/A]

Not integrable

Time = 0.13 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2} dx = \int \frac{(dx)^m}{\left(b \operatorname{artanh}\left(\frac{c}{x^2}\right) + a\right)^2} dx$$

input

```
integrate((d*x)^m/(a+b*arctanh(c/x^2))^2,x, algorithm="giac")
```

output

```
integrate((d*x)^m/(b*arctanh(c/x^2) + a)^2, x)
```

Mupad [N/A]

Not integrable

Time = 3.77 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2} dx = \int \frac{(dx)^m}{\left(a + b \operatorname{atanh}\left(\frac{c}{x^2}\right)\right)^2} dx$$

input

```
int((d*x)^m/(a + b*atanh(c/x^2))^2,x)
```

output

```
int((d*x)^m/(a + b*atanh(c/x^2))^2, x)
```

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.11

$$\int \frac{(dx)^m}{\left(a + b \operatorname{arctanh}\left(\frac{c}{x^2}\right)\right)^2} dx = d^m \left(\int \frac{x^m}{\operatorname{atanh}\left(\frac{c}{x^2}\right)^2 b^2 + 2 \operatorname{atanh}\left(\frac{c}{x^2}\right) ab + a^2} dx \right)$$

input `int((d*x)^m/(a+b*atanh(c/x^2))^2,x)`output `d**m*int(x**m/(atanh(c/x**2)**2*b**2 + 2*atanh(c/x**2)*a*b + a**2),x)`

3.186 $\int x^3(a + b \operatorname{arctanh}(c\sqrt{x})) dx$

Optimal result	1487
Mathematica [A] (verified)	1487
Rubi [A] (verified)	1488
Maple [A] (verified)	1490
Fricas [A] (verification not implemented)	1491
Sympy [F]	1491
Maxima [A] (verification not implemented)	1491
Giac [B] (verification not implemented)	1492
Mupad [B] (verification not implemented)	1493
Reduce [B] (verification not implemented)	1493

Optimal result

Integrand size = 16, antiderivative size = 88

$$\int x^3(a + b \operatorname{arctanh}(c\sqrt{x})) dx = \frac{b\sqrt{x}}{4c^7} + \frac{bx^{3/2}}{12c^5} + \frac{bx^{5/2}}{20c^3} + \frac{bx^{7/2}}{28c} - \frac{b \operatorname{arctanh}(c\sqrt{x})}{4c^8} + \frac{1}{4}x^4(a + b \operatorname{arctanh}(c\sqrt{x}))$$

output

```
1/4*b*x^(1/2)/c^7+1/12*b*x^(3/2)/c^5+1/20*b*x^(5/2)/c^3+1/28*b*x^(7/2)/c-1/4*b*arctanh(c*x^(1/2))/c^8+1/4*x^4*(a+b*arctanh(c*x^(1/2)))
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.30

$$\begin{aligned} & \int x^3(a + b \operatorname{arctanh}(c\sqrt{x})) dx \\ &= \frac{b\sqrt{x}}{4c^7} + \frac{bx^{3/2}}{12c^5} + \frac{bx^{5/2}}{20c^3} + \frac{bx^{7/2}}{28c} + \frac{ax^4}{4} \\ &+ \frac{1}{4}bx^4 \operatorname{arctanh}(c\sqrt{x}) + \frac{b \log(1 - c\sqrt{x})}{8c^8} - \frac{b \log(1 + c\sqrt{x})}{8c^8} \end{aligned}$$

input

```
Integrate[x^3*(a + b*ArcTanh[c*Sqrt[x]]),x]
```

output

```
(b*Sqrt[x])/(4*c^7) + (b*x^(3/2))/(12*c^5) + (b*x^(5/2))/(20*c^3) + (b*x^(7/2))/(28*c) + (a*x^4)/4 + (b*x^4*ArcTanh[c*Sqrt[x]])/4 + (b*Log[1 - c*Sqrt[x]])/(8*c^8) - (b*Log[1 + c*Sqrt[x]])/(8*c^8)
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.15, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {6452, 60, 60, 60, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 (a + b \operatorname{arctanh}(c\sqrt{x})) \, dx \\
 & \quad \downarrow 6452 \\
 & \frac{1}{4} x^4 (a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{8} bc \int \frac{x^{7/2}}{1 - c^2 x} \, dx \\
 & \quad \downarrow 60 \\
 & \frac{1}{4} x^4 (a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{8} bc \left(\frac{\int \frac{x^{5/2}}{1 - c^2 x} \, dx}{c^2} - \frac{2x^{7/2}}{7c^2} \right) \\
 & \quad \downarrow 60 \\
 & \frac{1}{4} x^4 (a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{8} bc \left(\frac{\int \frac{x^{3/2}}{1 - c^2 x} \, dx}{c^2} - \frac{2x^{5/2}}{5c^2} - \frac{2x^{7/2}}{7c^2} \right) \\
 & \quad \downarrow 60 \\
 & \frac{1}{4} x^4 (a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{8} bc \left(\frac{\int \frac{\sqrt{x}}{1 - c^2 x} \, dx}{c^2} - \frac{2x^{3/2}}{3c^2} - \frac{2x^{5/2}}{5c^2} - \frac{2x^{7/2}}{7c^2} \right) \\
 & \quad \downarrow 60
 \end{aligned}$$

$$\frac{1}{4}x^4(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{8}bc \left(\frac{\int \frac{1}{\sqrt{x}(1-c^2x)} dx}{c^2} - \frac{2\sqrt{x}}{c^2} - \frac{2x^{3/2}}{3c^2} - \frac{2x^{5/2}}{5c^2} - \frac{2x^{7/2}}{7c^2} \right)$$

↓ 73

$$\frac{1}{4}x^4(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{8}bc \left(\frac{2 \int \frac{1}{1-c^2x} d\sqrt{x}}{c^2} - \frac{2\sqrt{x}}{c^2} - \frac{2x^{3/2}}{3c^2} - \frac{2x^{5/2}}{5c^2} - \frac{2x^{7/2}}{7c^2} \right)$$

↓ 219

$$\frac{1}{4}x^4(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{8}bc \left(\frac{\frac{2\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{2\sqrt{x}}{c^2} - \frac{2x^{3/2}}{3c^2} - \frac{2x^{5/2}}{5c^2} - \frac{2x^{7/2}}{7c^2}}{c^2} \right)$$

input `Int[x^3*(a + b*ArcTanh[c*Sqrt[x]]),x]`

output `(x^4*(a + b*ArcTanh[c*Sqrt[x]]))/4 - (b*c*((-2*x^(7/2))/(7*c^2) + ((-2*x^(5/2))/(5*c^2) + ((-2*x^(3/2))/(3*c^2) + ((-2*Sqrt[x])/c^2 + (2*ArcTanh[c*Sqrt[x]])/c^3)/c^2)/c^2)/8`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
 ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
 Q[a, 0] || LtQ[b, 0])`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
 > Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
 + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.90

method	result	size
parts	$\frac{x^4 a}{4} + \frac{2b \left(\frac{c^8 x^4 \operatorname{arctanh}(c\sqrt{x})}{8} + \frac{c^7 x^{\frac{7}{2}}}{56} + \frac{c^5 x^{\frac{5}{2}}}{40} + \frac{c^3 x^{\frac{3}{2}}}{24} + \frac{c\sqrt{x}}{8} + \frac{\ln(c\sqrt{x}-1)}{16} - \frac{\ln(1+c\sqrt{x})}{16} \right)}{c^8}$	79
derivativedivides	$\frac{a c^8 x^4}{4} + 2b \left(\frac{c^8 x^4 \operatorname{arctanh}(c\sqrt{x})}{8} + \frac{c^7 x^{\frac{7}{2}}}{56} + \frac{c^5 x^{\frac{5}{2}}}{40} + \frac{c^3 x^{\frac{3}{2}}}{24} + \frac{c\sqrt{x}}{8} + \frac{\ln(c\sqrt{x}-1)}{16} - \frac{\ln(1+c\sqrt{x})}{16} \right)$	83
default	$\frac{a c^8 x^4}{4} + 2b \left(\frac{c^8 x^4 \operatorname{arctanh}(c\sqrt{x})}{8} + \frac{c^7 x^{\frac{7}{2}}}{56} + \frac{c^5 x^{\frac{5}{2}}}{40} + \frac{c^3 x^{\frac{3}{2}}}{24} + \frac{c\sqrt{x}}{8} + \frac{\ln(c\sqrt{x}-1)}{16} - \frac{\ln(1+c\sqrt{x})}{16} \right)$	83

input `int(x^3*(a+b*arctanh(c*x^(1/2))),x,method=_RETURNVERBOSE)`

output `1/4*x^4*a+2*b/c^8*(1/8*c^8*x^4*arctanh(c*x^(1/2))+1/56*c^7*x^(7/2)+1/40*c^5*x^(5/2)+1/24*c^3*x^(3/2)+1/8*c*x^(1/2)+1/16*ln(c*x^(1/2)-1)-1/16*ln(1+c*x^(1/2)))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.01

$$\int x^3(a + \operatorname{barctanh}(c\sqrt{x})) dx$$

$$= \frac{210 ac^8 x^4 + 105 (bc^8 x^4 - b) \log\left(-\frac{c^2 x + 2c\sqrt{x} + 1}{c^2 x - 1}\right) + 2(15 bc^7 x^3 + 21 bc^5 x^2 + 35 bc^3 x + 105 bc)\sqrt{x}}{840 c^8}$$

input `integrate(x^3*(a+b*arctanh(c*x^(1/2))),x, algorithm="fricas")`output `1/840*(210*a*c^8*x^4 + 105*(b*c^8*x^4 - b)*log(-(c^2*x + 2*c*sqrt(x) + 1)/(c^2*x - 1)) + 2*(15*b*c^7*x^3 + 21*b*c^5*x^2 + 35*b*c^3*x + 105*b*c)*sqrt(x))/c^8`**Sympy [F]**

$$\int x^3(a + \operatorname{barctanh}(c\sqrt{x})) dx = \int x^3(a + b \operatorname{atanh}(c\sqrt{x})) dx$$

input `integrate(x**3*(a+b*atanh(c*x**(1/2))),x)`output `Integral(x**3*(a + b*atanh(c*sqrt(x))), x)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.98

$$\int x^3(a + \operatorname{barctanh}(c\sqrt{x})) dx = \frac{1}{4} ax^4$$

$$+ \frac{1}{840} \left(210 x^4 \operatorname{artanh}(c\sqrt{x}) + c \left(\frac{2 \left(15 c^6 x^{\frac{7}{2}} + 21 c^4 x^{\frac{5}{2}} + 35 c^2 x^{\frac{3}{2}} + 105 \sqrt{x} \right)}{c^8} - \frac{105 \log(c\sqrt{x} + 1)}{c^9} \right) \right)$$

input `integrate(x^3*(a+b*arctanh(c*x^(1/2))),x, algorithm="maxima")`

output `1/4*a*x^4 + 1/840*(210*x^4*arctanh(c*sqrt(x)) + c*(2*(15*c^6*x^(7/2) + 21*c^4*x^(5/2) + 35*c^2*x^(3/2) + 105*sqrt(x))/c^8 - 105*log(c*sqrt(x) + 1)/c^9 + 105*log(c*sqrt(x) - 1)/c^9))*b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 359 vs. $2(64) = 128$.

Time = 0.14 (sec) , antiderivative size = 359, normalized size of antiderivative = 4.08

$$\int x^3(a + b \operatorname{arctanh}(c\sqrt{x})) dx = \frac{1}{4} ax^4 + \frac{2}{105} bc \left(\frac{\frac{105(c\sqrt{x}+1)^6}{(c\sqrt{x}-1)^6} - \frac{315(c\sqrt{x}+1)^5}{(c\sqrt{x}-1)^5} + \frac{770(c\sqrt{x}+1)^4}{(c\sqrt{x}-1)^4} - \frac{770(c\sqrt{x}+1)^3}{(c\sqrt{x}-1)^3} + \frac{609(c\sqrt{x}+1)^2}{(c\sqrt{x}-1)^2} - \frac{203(c\sqrt{x}+1)}{c\sqrt{x}-1} + 44}{c^9 \left(\frac{c\sqrt{x}+1}{c\sqrt{x}-1} - 1 \right)^7} + \frac{105}{c^9} \right)$$

input `integrate(x^3*(a+b*arctanh(c*x^(1/2))),x, algorithm="giac")`

output `1/4*a*x^4 + 2/105*b*c*((105*(c*sqrt(x) + 1)^6/(c*sqrt(x) - 1)^6 - 315*(c*sqrt(x) + 1)^5/(c*sqrt(x) - 1)^5 + 770*(c*sqrt(x) + 1)^4/(c*sqrt(x) - 1)^4 - 770*(c*sqrt(x) + 1)^3/(c*sqrt(x) - 1)^3 + 609*(c*sqrt(x) + 1)^2/(c*sqrt(x) - 1)^2 - 203*(c*sqrt(x) + 1)/(c*sqrt(x) - 1) + 44)/(c^9*((c*sqrt(x) + 1)/(c*sqrt(x) - 1) - 1)^7) + 105*((c*sqrt(x) + 1)^7/(c*sqrt(x) - 1)^7 + 7*(c*sqrt(x) + 1)^5/(c*sqrt(x) - 1)^5 + 7*(c*sqrt(x) + 1)^3/(c*sqrt(x) - 1)^3 + (c*sqrt(x) + 1)/(c*sqrt(x) - 1))*log(-(c*((c*sqrt(x) + 1)/(c*sqrt(x) - 1) + 1))/((c*sqrt(x) + 1)*c/(c*sqrt(x) - 1) - c) + 1)/(c*((c*sqrt(x) + 1)/(c*sqrt(x) - 1) + 1))/((c*sqrt(x) + 1)*c/(c*sqrt(x) - 1) - c) - 1))/c^9*((c*sqrt(x) + 1)/(c*sqrt(x) - 1) - 1)^8))`

Mupad [B] (verification not implemented)

Time = 4.62 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.98

$$\int x^3 (a + b \operatorname{arctanh}(c\sqrt{x})) dx = \frac{\frac{bc^3 x^{3/2}}{12} - \frac{b \operatorname{atanh}(c\sqrt{x})}{4} + \frac{bc^5 x^{5/2}}{20} + \frac{bc^7 x^{7/2}}{28} + \frac{bc\sqrt{x}}{4}}{c^8} + \frac{b(105x^4 \ln(c\sqrt{x} + 1) - 105x^4 \ln(1 - c\sqrt{x}))}{840} + \frac{ax^4}{4}$$

input `int(x^3*(a + b*atanh(c*x^(1/2))),x)`output `((b*c^3*x^(3/2))/12 - (b*atanh(c*x^(1/2)))/4 + (b*c^5*x^(5/2))/20 + (b*c^7*x^(7/2))/28 + (b*c*x^(1/2))/4)/c^8 + (b*(105*x^4*log(c*x^(1/2) + 1) - 105*x^4*log(1 - c*x^(1/2)))/840 + (a*x^4)/4`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.84

$$\int x^3 (a + b \operatorname{arctanh}(c\sqrt{x})) dx = \frac{105 \operatorname{atanh}(\sqrt{x}c) b c^8 x^4 - 105 \operatorname{atanh}(\sqrt{x}c) b + 15\sqrt{x} b c^7 x^3 + 21\sqrt{x} b c^5 x^2 + 35\sqrt{x} b c^3 x + 105\sqrt{x} b c + 105 a c^8 x^4}{420 c^8}$$

input `int(x^3*(a+b*atanh(c*x^(1/2))),x)`output `(105*atanh(sqrt(x)*c)*b*c**8*x**4 - 105*atanh(sqrt(x)*c)*b + 15*sqrt(x)*b*c**7*x**3 + 21*sqrt(x)*b*c**5*x**2 + 35*sqrt(x)*b*c**3*x + 105*sqrt(x)*b*c + 105*a*c**8*x**4)/(420*c**8)`

3.187 $\int x^2(a + b \operatorname{arctanh}(c\sqrt{x})) dx$

Optimal result	1494
Mathematica [A] (verified)	1494
Rubi [A] (verified)	1495
Maple [A] (verified)	1497
Fricas [A] (verification not implemented)	1497
Sympy [F]	1498
Maxima [A] (verification not implemented)	1498
Giac [B] (verification not implemented)	1499
Mupad [B] (verification not implemented)	1499
Reduce [B] (verification not implemented)	1500

Optimal result

Integrand size = 16, antiderivative size = 75

$$\int x^2(a + b \operatorname{arctanh}(c\sqrt{x})) dx = \frac{b\sqrt{x}}{3c^5} + \frac{bx^{3/2}}{9c^3} + \frac{bx^{5/2}}{15c} - \frac{b \operatorname{arctanh}(c\sqrt{x})}{3c^6} + \frac{1}{3}x^3(a + b \operatorname{arctanh}(c\sqrt{x}))$$

output `1/3*b*x^(1/2)/c^5+1/9*b*x^(3/2)/c^3+1/15*b*x^(5/2)/c-1/3*b*arctanh(c*x^(1/2))/c^6+1/3*x^3*(a+b*arctanh(c*x^(1/2)))`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.35

$$\int x^2(a + b \operatorname{arctanh}(c\sqrt{x})) dx = \frac{b\sqrt{x}}{3c^5} + \frac{bx^{3/2}}{9c^3} + \frac{bx^{5/2}}{15c} + \frac{ax^3}{3} + \frac{1}{3}bx^3 \operatorname{arctanh}(c\sqrt{x}) + \frac{b \log(1 - c\sqrt{x})}{6c^6} - \frac{b \log(1 + c\sqrt{x})}{6c^6}$$

input `Integrate[x^2*(a + b*ArcTanh[c*Sqrt[x]]),x]`

output

```
(b*Sqrt[x])/(3*c^5) + (b*x^(3/2))/(9*c^3) + (b*x^(5/2))/(15*c) + (a*x^3)/3
+ (b*x^3*ArcTanh[c*Sqrt[x]])/3 + (b*Log[1 - c*Sqrt[x]])/(6*c^6) - (b*Log[
1 + c*Sqrt[x]])/(6*c^6)
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6452, 60, 60, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 (a + b \operatorname{arctanh}(c\sqrt{x})) \, dx \\
 & \quad \downarrow 6452 \\
 & \frac{1}{3} x^3 (a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6} bc \int \frac{x^{5/2}}{1 - c^2 x} \, dx \\
 & \quad \downarrow 60 \\
 & \frac{1}{3} x^3 (a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6} bc \left(\frac{\int \frac{x^{3/2}}{1 - c^2 x} \, dx}{c^2} - \frac{2x^{5/2}}{5c^2} \right) \\
 & \quad \downarrow 60 \\
 & \frac{1}{3} x^3 (a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6} bc \left(\frac{\frac{\int \frac{\sqrt{x}}{1 - c^2 x} \, dx}{c^2} - \frac{2x^{3/2}}{3c^2}}{c^2} - \frac{2x^{5/2}}{5c^2} \right) \\
 & \quad \downarrow 60 \\
 & \frac{1}{3} x^3 (a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6} bc \left(\frac{\frac{\frac{\int \frac{1}{\sqrt{x}(1 - c^2 x)} \, dx}{c^2} - \frac{2\sqrt{x}}{c^2}}{c^2} - \frac{2x^{3/2}}{3c^2}}{c^2} - \frac{2x^{5/2}}{5c^2} \right) \\
 & \quad \downarrow 73
 \end{aligned}$$

$$\frac{1}{3}x^3(a + \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6}bc \left(\frac{\frac{2 \int \frac{1}{1-c^2x} d\sqrt{x} - \frac{2\sqrt{x}}{c^2} - \frac{2x^{3/2}}{3c^2}}{c^2}}{c^2} - \frac{2x^{5/2}}{5c^2} \right)$$

↓ 219

$$\frac{1}{3}x^3(a + \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6}bc \left(\frac{\frac{2\operatorname{arctanh}(c\sqrt{x}) - \frac{2\sqrt{x}}{c^2} - \frac{2x^{3/2}}{3c^2}}{c^2}}{c^2} - \frac{2x^{5/2}}{5c^2} \right)$$

input `Int[x^2*(a + b*ArcTanh[c*Sqrt[x]]),x]`

output `(x^3*(a + b*ArcTanh[c*Sqrt[x]]))/3 - (b*c*((-2*x^(5/2))/(5*c^2) + ((-2*x^(3/2))/(3*c^2) + ((-2*Sqrt[x])/c^2 + (2*ArcTanh[c*Sqrt[x]]/c^3)/c^2)/c^2))/6`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.95

method	result	size
parts	$\frac{ax^3}{3} + \frac{2b \left(\frac{c^6 x^3 \operatorname{arctanh}(c\sqrt{x})}{6} + \frac{c^5 x^{\frac{5}{2}}}{30} + \frac{c^3 x^{\frac{3}{2}}}{18} + \frac{c\sqrt{x}}{6} + \frac{\ln(c\sqrt{x}-1)}{12} - \frac{\ln(1+c\sqrt{x})}{12} \right)}{c^6}$	71
derivativedivides	$\frac{\frac{ac^6x^3}{3} + 2b \left(\frac{c^6 x^3 \operatorname{arctanh}(c\sqrt{x})}{6} + \frac{c^5 x^{\frac{5}{2}}}{30} + \frac{c^3 x^{\frac{3}{2}}}{18} + \frac{c\sqrt{x}}{6} + \frac{\ln(c\sqrt{x}-1)}{12} - \frac{\ln(1+c\sqrt{x})}{12} \right)}{c^6}$	75
default	$\frac{\frac{ac^6x^3}{3} + 2b \left(\frac{c^6 x^3 \operatorname{arctanh}(c\sqrt{x})}{6} + \frac{c^5 x^{\frac{5}{2}}}{30} + \frac{c^3 x^{\frac{3}{2}}}{18} + \frac{c\sqrt{x}}{6} + \frac{\ln(c\sqrt{x}-1)}{12} - \frac{\ln(1+c\sqrt{x})}{12} \right)}{c^6}$	75

input

```
int(x^2*(a+b*arctanh(c*x^(1/2))),x,method=_RETURNVERBOSE)
```

output

```
1/3*a*x^3+2*b/c^6*(1/6*c^6*x^3*arctanh(c*x^(1/2))+1/30*c^5*x^(5/2)+1/18*c^
3*x^(3/2)+1/6*c*x^(1/2)+1/12*ln(c*x^(1/2)-1)-1/12*ln(1+c*x^(1/2)))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.07

$$\int x^2 (a + b \operatorname{arctanh}(c\sqrt{x})) dx$$

$$= \frac{30ac^6x^3 + 15(bc^6x^3 - b) \log\left(-\frac{c^2x+2c\sqrt{x}+1}{c^2x-1}\right) + 2(3bc^5x^2 + 5bc^3x + 15bc)\sqrt{x}}{90c^6}$$

input

```
integrate(x^2*(a+b*arctanh(c*x^(1/2))),x, algorithm="fricas")
```

output $\frac{1}{90}(30ac^6x^3 + 15(b^2c^6x^3 - b)\log(-c^2x + 2c\sqrt{x} + 1)/(c^2x - 1)) + 2(3b^2c^5x^2 + 5b^2c^3x + 15b^2c)\sqrt{x}/c^6$

Sympy [F]

$$\int x^2(a + b \operatorname{arctanh}(c\sqrt{x})) dx = \int x^2(a + b \operatorname{atanh}(c\sqrt{x})) dx$$

input `integrate(x**2*(a+b*atanh(c*x**(1/2))),x)`

output `Integral(x**2*(a + b*atanh(c*sqrt(x))), x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.04

$$\int x^2(a + b \operatorname{arctanh}(c\sqrt{x})) dx = \frac{1}{3}ax^3 + \frac{1}{90} \left(30x^3 \operatorname{arctanh}(c\sqrt{x}) + c \left(\frac{2(3c^4x^{\frac{5}{2}} + 5c^2x^{\frac{3}{2}} + 15\sqrt{x})}{c^6} - \frac{15 \log(c\sqrt{x} + 1)}{c^7} + \frac{15 \log(c\sqrt{x} - 1)}{c^7} \right) \right)$$

input `integrate(x^2*(a+b*arctanh(c*x^(1/2))),x, algorithm="maxima")`

output $\frac{1}{3}ax^3 + \frac{1}{90}(30x^3\operatorname{arctanh}(c\sqrt{x}) + c(2(3c^4x^{5/2} + 5c^2x^{3/2} + 15\sqrt{x})/c^6 - 15\log(c\sqrt{x} + 1)/c^7 + 15\log(c\sqrt{x} - 1)/c^7))*b$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 301 vs. $2(55) = 110$.

Time = 0.14 (sec) , antiderivative size = 301, normalized size of antiderivative = 4.01

$$\int x^2(a + \operatorname{barctanh}(c\sqrt{x})) dx = \frac{1}{3} ax^3 + \frac{2}{45} bc \left(\frac{\frac{45(c\sqrt{x}+1)^4}{(c\sqrt{x}-1)^4} - \frac{90(c\sqrt{x}+1)^3}{(c\sqrt{x}-1)^3} + \frac{140(c\sqrt{x}+1)^2}{(c\sqrt{x}-1)^2} - \frac{70(c\sqrt{x}+1)}{c\sqrt{x}-1} + 23}{c^7 \left(\frac{c\sqrt{x}+1}{c\sqrt{x}-1} - 1 \right)^5} + \frac{15 \left(\frac{3(c\sqrt{x}+1)^5}{(c\sqrt{x}-1)^5} + \frac{10(c\sqrt{x}+1)^3}{(c\sqrt{x}-1)^3} + \frac{3(c\sqrt{x}+1)}{c\sqrt{x}-1} \right)}{c^7 \left(\frac{c\sqrt{x}+1}{c\sqrt{x}-1} - 1 \right)^6} \right)$$

input `integrate(x^2*(a+b*arctanh(c*x^(1/2))),x, algorithm="giac")`

output

```
1/3*a*x^3 + 2/45*b*c*((45*(c*sqrt(x) + 1)^4/(c*sqrt(x) - 1)^4 - 90*(c*sqrt(x) + 1)^3/(c*sqrt(x) - 1)^3 + 140*(c*sqrt(x) + 1)^2/(c*sqrt(x) - 1)^2 - 70*(c*sqrt(x) + 1)/(c*sqrt(x) - 1) + 23)/(c^7*((c*sqrt(x) + 1)/(c*sqrt(x) - 1) - 1)^5) + 15*(3*(c*sqrt(x) + 1)^5/(c*sqrt(x) - 1)^5 + 10*(c*sqrt(x) + 1)^3/(c*sqrt(x) - 1)^3 + 3*(c*sqrt(x) + 1)/(c*sqrt(x) - 1))*log(-(c*((c*sqrt(x) + 1)/(c*sqrt(x) - 1) + 1)/((c*sqrt(x) + 1)*c/(c*sqrt(x) - 1) - c) + 1)/(c*((c*sqrt(x) + 1)/(c*sqrt(x) - 1) + 1)/((c*sqrt(x) + 1)*c/(c*sqrt(x) - 1) - c) - 1))/(c^7*((c*sqrt(x) + 1)/(c*sqrt(x) - 1) - 1)^6))
```

Mupad [B] (verification not implemented)

Time = 4.25 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.77

$$\int x^2(a + \operatorname{barctanh}(c\sqrt{x})) dx = \frac{ax^3}{3} + \frac{bc^3 x^{3/2}}{9} - \frac{b \operatorname{atanh}(c\sqrt{x})}{3} + \frac{bc^5 x^{5/2}}{15} + \frac{bc\sqrt{x}}{3} + \frac{bx^3 \operatorname{atanh}(c\sqrt{x})}{3}$$

input `int(x^2*(a + b*atanh(c*x^(1/2))),x)`

output

$$\frac{(a*x^3)/3 + ((b*c^3*x^(3/2))/9 - (b*atanh(c*x^(1/2)))/3 + (b*c^5*x^(5/2))/15 + (b*c*x^(1/2))/3)/c^6 + (b*x^3*atanh(c*x^(1/2)))/3}$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.84

$$\int x^2(a + b \operatorname{arctanh}(c\sqrt{x})) dx$$

$$= \frac{15 \operatorname{atanh}(\sqrt{x}c) b c^6 x^3 - 15 \operatorname{atanh}(\sqrt{x}c) b + 3\sqrt{x} b c^5 x^2 + 5\sqrt{x} b c^3 x + 15\sqrt{x} b c + 15a c^6 x^3}{45c^6}$$

input

```
int(x^2*(a+b*atanh(c*x^(1/2))),x)
```

output

$$(15*\operatorname{atanh}(\operatorname{sqrt}(x)*c)*b*c**6*x**3 - 15*\operatorname{atanh}(\operatorname{sqrt}(x)*c)*b + 3*\operatorname{sqrt}(x)*b*c**5*x**2 + 5*\operatorname{sqrt}(x)*b*c**3*x + 15*\operatorname{sqrt}(x)*b*c + 15*a*c**6*x**3)/(45*c**6)$$

3.188 $\int x(a + b \operatorname{arctanh}(c\sqrt{x})) dx$

Optimal result	1501
Mathematica [A] (verified)	1501
Rubi [A] (verified)	1502
Maple [A] (verified)	1504
Fricas [A] (verification not implemented)	1504
Sympy [F]	1505
Maxima [A] (verification not implemented)	1505
Giac [B] (verification not implemented)	1505
Mupad [B] (verification not implemented)	1506
Reduce [B] (verification not implemented)	1507

Optimal result

Integrand size = 14, antiderivative size = 62

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x})) dx = \frac{b\sqrt{x}}{2c^3} + \frac{bx^{3/2}}{6c} - \frac{b \operatorname{arctanh}(c\sqrt{x})}{2c^4} + \frac{1}{2}x^2(a + b \operatorname{arctanh}(c\sqrt{x}))$$

output

```
1/2*b*x^(1/2)/c^3+1/6*b*x^(3/2)/c-1/2*b*arctanh(c*x^(1/2))/c^4+1/2*x^2*(a+b*arctanh(c*x^(1/2)))
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.42

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x})) dx = \frac{b\sqrt{x}}{2c^3} + \frac{bx^{3/2}}{6c} + \frac{ax^2}{2} + \frac{1}{2}bx^2 \operatorname{arctanh}(c\sqrt{x}) + \frac{b \log(1 - c\sqrt{x})}{4c^4} - \frac{b \log(1 + c\sqrt{x})}{4c^4}$$

input

```
Integrate[x*(a + b*ArcTanh[c*Sqrt[x]]),x]
```

output

```
(b*Sqrt[x])/(2*c^3) + (b*x^(3/2))/(6*c) + (a*x^2)/2 + (b*x^2*ArcTanh[c*Sqrt[x]])/2 + (b*Log[1 - c*Sqrt[x]])/(4*c^4) - (b*Log[1 + c*Sqrt[x]])/(4*c^4)
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.08, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {6452, 60, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x(a + b \operatorname{arctanh}(c\sqrt{x})) dx \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{2}x^2(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \int \frac{x^{3/2}}{1 - c^2x} dx \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2}x^2(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\int \frac{\sqrt{x}}{1 - c^2x} dx}{c^2} - \frac{2x^{3/2}}{3c^2} \right) \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2}x^2(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\int \frac{1}{\sqrt{x}(1 - c^2x)} dx}{c^2} - \frac{2\sqrt{x}}{c^2} - \frac{2x^{3/2}}{3c^2} \right) \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{2}x^2(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{2 \int \frac{1}{1 - c^2x} d\sqrt{x}}{c^2} - \frac{2\sqrt{x}}{c^2} - \frac{2x^{3/2}}{3c^2} \right) \\
 & \quad \downarrow \text{219} \\
 & \frac{1}{2}x^2(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{2 \operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{2\sqrt{x}}{c^2} - \frac{2x^{3/2}}{3c^2} \right)
 \end{aligned}$$

input `Int[x*(a + b*ArcTanh[c*sqrt[x]]),x]`

output

$$\frac{(x^2(a + b \operatorname{ArcTanh}[c \sqrt{x}]))}{2} - \frac{(b*c*((-2*x^{(3/2)))/(3*c^2) + ((-2*\operatorname{Sqrt}[x])/c^2 + (2*\operatorname{ArcTanh}[c*\operatorname{Sqrt}[x]])/c^3)/c^2))/4}$$

Defintions of rubi rules used

rule 60

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(
b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]
```

rule 73

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 219

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.02

method	result	size
parts	$\frac{ax^2}{2} + \frac{2b \left(\frac{c^4 x^2 \operatorname{arctanh}(c\sqrt{x})}{4} + \frac{c^3 x^{\frac{3}{2}}}{12} + \frac{c\sqrt{x}}{4} + \frac{\ln(c\sqrt{x}-1)}{8} - \frac{\ln(1+c\sqrt{x})}{8} \right)}{c^4}$	63
derivativedivides	$\frac{ac^4 x^2 + 2b \left(\frac{c^4 x^2 \operatorname{arctanh}(c\sqrt{x})}{4} + \frac{c^3 x^{\frac{3}{2}}}{12} + \frac{c\sqrt{x}}{4} + \frac{\ln(c\sqrt{x}-1)}{8} - \frac{\ln(1+c\sqrt{x})}{8} \right)}{c^4}$	67
default	$\frac{ac^4 x^2 + 2b \left(\frac{c^4 x^2 \operatorname{arctanh}(c\sqrt{x})}{4} + \frac{c^3 x^{\frac{3}{2}}}{12} + \frac{c\sqrt{x}}{4} + \frac{\ln(c\sqrt{x}-1)}{8} - \frac{\ln(1+c\sqrt{x})}{8} \right)}{c^4}$	67

input `int(x*(a+b*arctanh(c*x^(1/2))),x,method=_RETURNVERBOSE)`

output `1/2*a*x^2+2*b/c^4*(1/4*c^4*x^2*arctanh(c*x^(1/2))+1/12*c^3*x^(3/2)+1/4*c*x^(1/2)+1/8*ln(c*x^(1/2)-1)-1/8*ln(1+c*x^(1/2)))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.13

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x})) dx$$

$$= \frac{6ac^4x^2 + 3(bc^4x^2 - b) \log\left(-\frac{c^2x+2c\sqrt{x}+1}{c^2x-1}\right) + 2(bc^3x + 3bc)\sqrt{x}}{12c^4}$$

input `integrate(x*(a+b*arctanh(c*x^(1/2))),x, algorithm="fricas")`

output `1/12*(6*a*c^4*x^2 + 3*(b*c^4*x^2 - b)*log(-(c^2*x + 2*c*sqrt(x) + 1)/(c^2*x - 1)) + 2*(b*c^3*x + 3*b*c)*sqrt(x))/c^4`

Sympy [F]

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x})) dx = \int x(a + b \operatorname{atanh}(c\sqrt{x})) dx$$

input `integrate(x*(a+b*atanh(c*x**(1/2))),x)`

output `Integral(x*(a + b*atanh(c*sqrt(x))), x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.11

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x})) dx = \frac{1}{2} ax^2 + \frac{1}{12} \left(6x^2 \operatorname{arctanh}(c\sqrt{x}) + c \left(\frac{2(c^2 x^{\frac{3}{2}} + 3\sqrt{x})}{c^4} - \frac{3 \log(c\sqrt{x} + 1)}{c^5} + \frac{3 \log(c\sqrt{x} - 1)}{c^5} \right) \right) b$$

input `integrate(x*(a+b*arctanh(c*x^(1/2))),x, algorithm="maxima")`

output `1/2*a*x^2 + 1/12*(6*x^2*arctanh(c*sqrt(x)) + c*(2*(c^2*x^(3/2) + 3*sqrt(x))/c^4 - 3*log(c*sqrt(x) + 1)/c^5 + 3*log(c*sqrt(x) - 1)/c^5))*b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 239 vs. 2(46) = 92.

Time = 0.13 (sec) , antiderivative size = 239, normalized size of antiderivative = 3.85

$$\int x(a + \operatorname{arctanh}(c\sqrt{x})) dx$$

$$= \frac{1}{2} ax^2$$

$$+ \frac{2}{3} bc \left(\frac{\frac{3(c\sqrt{x}+1)^2}{(c\sqrt{x}-1)^2} - \frac{3(c\sqrt{x}+1)}{c\sqrt{x}-1} + 2}{c^5 \left(\frac{c\sqrt{x}+1}{c\sqrt{x}-1} - 1 \right)^3} + \frac{3 \left(\frac{(c\sqrt{x}+1)^3}{(c\sqrt{x}-1)^3} + \frac{c\sqrt{x}+1}{c\sqrt{x}-1} \right) \log \left(-\frac{\frac{c \left(\frac{c\sqrt{x}+1}{c\sqrt{x}-1} + 1 \right)}{(c\sqrt{x}+1)c - c} + 1}{\frac{c \left(\frac{c\sqrt{x}+1}{c\sqrt{x}-1} + 1 \right)}{(c\sqrt{x}+1)c - c} - 1} \right)}{c^5 \left(\frac{c\sqrt{x}+1}{c\sqrt{x}-1} - 1 \right)^4} \right)$$

input `integrate(x*(a+b*arctanh(c*x^(1/2))),x, algorithm="giac")`

output `1/2*a*x^2 + 2/3*b*c*((3*(c*sqrt(x) + 1)^2/(c*sqrt(x) - 1)^2 - 3*(c*sqrt(x) + 1)/(c*sqrt(x) - 1) + 2)/(c^5*((c*sqrt(x) + 1)/(c*sqrt(x) - 1) - 1)^3) + 3*((c*sqrt(x) + 1)^3/(c*sqrt(x) - 1)^3 + (c*sqrt(x) + 1)/(c*sqrt(x) - 1)) *log(-(c*((c*sqrt(x) + 1)/(c*sqrt(x) - 1) + 1)/((c*sqrt(x) + 1)*c/(c*sqrt(x) - 1) - c) + 1)/(c*((c*sqrt(x) + 1)/(c*sqrt(x) - 1) + 1)/((c*sqrt(x) + 1)*c/(c*sqrt(x) - 1) - c) - 1)))/(c^5*((c*sqrt(x) + 1)/(c*sqrt(x) - 1) - 1)^4))`

Mupad [B] (verification not implemented)

Time = 4.23 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.79

$$\int x(a + \operatorname{arctanh}(c\sqrt{x})) dx = \frac{bc^3 x^{3/2}}{6} - \frac{b \operatorname{atanh}(c\sqrt{x})}{2} + \frac{bc\sqrt{x}}{2} + \frac{ax^2}{2} + \frac{bx^2 \operatorname{atanh}(c\sqrt{x})}{2}$$

input `int(x*(a + b*atanh(c*x^(1/2))),x)`

output `((b*c^3*x^(3/2))/6 - (b*atanh(c*x^(1/2)))/2 + (b*c*x^(1/2))/2)/c^4 + (a*x^2)/2 + (b*x^2*atanh(c*x^(1/2)))/2`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.82

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x})) dx$$

$$= \frac{3 \operatorname{atanh}(\sqrt{x}c) b c^4 x^2 - 3 \operatorname{atanh}(\sqrt{x}c) b + \sqrt{x} b c^3 x + 3\sqrt{x} b c + 3a c^4 x^2}{6c^4}$$

input `int(x*(a+b*atanh(c*x^(1/2))),x)`output `(3*atanh(sqrt(x)*c)*b*c**4*x**2 - 3*atanh(sqrt(x)*c)*b + sqrt(x)*b*c**3*x + 3*sqrt(x)*b*c + 3*a*c**4*x**2)/(6*c**4)`

3.189 $\int (a + b \operatorname{arctanh}(c\sqrt{x})) dx$

Optimal result	1508
Mathematica [A] (verified)	1508
Rubi [A] (verified)	1509
Maple [A] (verified)	1509
Fricas [A] (verification not implemented)	1510
Sympy [F]	1510
Maxima [A] (verification not implemented)	1511
Giac [B] (verification not implemented)	1511
Mupad [B] (verification not implemented)	1512
Reduce [B] (verification not implemented)	1512

Optimal result

Integrand size = 12, antiderivative size = 39

$$\int (a + b \operatorname{arctanh}(c\sqrt{x})) dx = \frac{b\sqrt{x}}{c} + ax - \frac{b \operatorname{arctanh}(c\sqrt{x})}{c^2} + b x \operatorname{arctanh}(c\sqrt{x})$$

output

```
b*x^(1/2)/c+a*x-b*arctanh(c*x^(1/2))/c^2+b*x*arctanh(c*x^(1/2))
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.08

$$\int (a + b \operatorname{arctanh}(c\sqrt{x})) dx = ax + b x \operatorname{arctanh}(c\sqrt{x}) - bc \left(-\frac{\sqrt{x}}{c^2} + \frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} \right)$$

input

```
Integrate[a + b*ArcTanh[c*Sqrt[x]], x]
```

output

```
a*x + b*x*ArcTanh[c*Sqrt[x]] - b*c*(-(Sqrt[x]/c^2) + ArcTanh[c*Sqrt[x]]/c^3)
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{arctanh}(c\sqrt{x})) dx$$

$$\downarrow \text{2009}$$

$$ax - \frac{b \operatorname{arctanh}(c\sqrt{x})}{c^2} + b x \operatorname{arctanh}(c\sqrt{x}) + \frac{b\sqrt{x}}{c}$$

input `Int[a + b*ArcTanh[c*Sqrt[x]],x]`

output `(b*Sqrt[x])/c + a*x - (b*ArcTanh[c*Sqrt[x]])/c^2 + b*x*ArcTanh[c*Sqrt[x]]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.28

method	result	size
default	$ax + \frac{2b \left(\frac{c^2 x \operatorname{arctanh}(c\sqrt{x})}{2} + \frac{c\sqrt{x}}{2} + \frac{\ln(c\sqrt{x}-1)}{4} - \frac{\ln(1+c\sqrt{x})}{4} \right)}{c^2}$	50
parts	$ax + \frac{2b \left(\frac{c^2 x \operatorname{arctanh}(c\sqrt{x})}{2} + \frac{c\sqrt{x}}{2} + \frac{\ln(c\sqrt{x}-1)}{4} - \frac{\ln(1+c\sqrt{x})}{4} \right)}{c^2}$	50
derivativedivides	$\frac{ax c^2 + 2b \left(\frac{c^2 x \operatorname{arctanh}(c\sqrt{x})}{2} + \frac{c\sqrt{x}}{2} + \frac{\ln(c\sqrt{x}-1)}{4} - \frac{\ln(1+c\sqrt{x})}{4} \right)}{c^2}$	55

input `int(a+b*arctanh(c*x^(1/2)),x,method=_RETURNVERBOSE)`

output `a*x+2*b/c^2*(1/2*c^2*x*arctanh(c*x^(1/2))+1/2*c*x^(1/2)+1/4*ln(c*x^(1/2)-1)-1/4*ln(1+c*x^(1/2)))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.44

$$\int (a + b \operatorname{arctanh}(c\sqrt{x})) dx = \frac{2ac^2x + 2bc\sqrt{x} + (bc^2x - b) \log\left(-\frac{c^2x + 2c\sqrt{x} + 1}{c^2x - 1}\right)}{2c^2}$$

input `integrate(a+b*arctanh(c*x^(1/2)),x, algorithm="fricas")`

output `1/2*(2*a*c^2*x + 2*b*c*sqrt(x) + (b*c^2*x - b)*log(-(c^2*x + 2*c*sqrt(x) + 1)/(c^2*x - 1)))/c^2`

Sympy [F]

$$\int (a + b \operatorname{arctanh}(c\sqrt{x})) dx = \int (a + b \operatorname{atanh}(c\sqrt{x})) dx$$

input `integrate(a+b*atanh(c*x**(1/2)),x)`

output `Integral(a + b*atanh(c*sqrt(x)), x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.36

$$\int (a + b \operatorname{arctanh}(c\sqrt{x})) dx$$

$$= \frac{1}{2} \left(c \left(\frac{2\sqrt{x}}{c^2} - \frac{\log(c\sqrt{x}+1)}{c^3} + \frac{\log(c\sqrt{x}-1)}{c^3} \right) + 2x \operatorname{artanh}(c\sqrt{x}) \right) b + ax$$

input `integrate(a+b*arctanh(c*x^(1/2)),x, algorithm="maxima")`

output `1/2*(c*(2*sqrt(x)/c^2 - log(c*sqrt(x) + 1)/c^3 + log(c*sqrt(x) - 1)/c^3) + 2*x*arctanh(c*sqrt(x)))*b + a*x`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 174 vs. 2(33) = 66.

Time = 0.13 (sec) , antiderivative size = 174, normalized size of antiderivative = 4.46

$$\int (a + b \operatorname{arctanh}(c\sqrt{x})) dx$$

$$= 2bc \left(\frac{1}{c^3 \left(\frac{c\sqrt{x}+1}{c\sqrt{x}-1} - 1 \right)} + \frac{(c\sqrt{x}+1) \log \left(\frac{\frac{c \left(\frac{c\sqrt{x}+1}{c\sqrt{x}-1} + 1 \right)}{\frac{(c\sqrt{x}+1)c}{c\sqrt{x}-1} - c} + 1}{\frac{c \left(\frac{c\sqrt{x}+1}{c\sqrt{x}-1} + 1 \right)}{\frac{(c\sqrt{x}+1)c}{c\sqrt{x}-1} - c} - 1} \right)}{(c\sqrt{x}-1)c^3 \left(\frac{c\sqrt{x}+1}{c\sqrt{x}-1} - 1 \right)^2} \right) + ax$$

input `integrate(a+b*arctanh(c*x^(1/2)),x, algorithm="giac")`

output

```
2*b*c*(1/(c^3*((c*sqrt(x) + 1)/(c*sqrt(x) - 1) - 1)) + (c*sqrt(x) + 1)*log
(-(c*((c*sqrt(x) + 1)/(c*sqrt(x) - 1) + 1)/((c*sqrt(x) + 1)*c/(c*sqrt(x) -
1) - c) + 1)/(c*((c*sqrt(x) + 1)/(c*sqrt(x) - 1) + 1)/((c*sqrt(x) + 1)*c/
(c*sqrt(x) - 1) - c) - 1))/((c*sqrt(x) - 1)*c^3*((c*sqrt(x) + 1)/(c*sqrt(x)
) - 1) - 1)^2)) + a*x
```

Mupad [B] (verification not implemented)

Time = 4.05 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.82

$$\int (a + b \operatorname{arctanh}(c\sqrt{x})) dx = ax + bx \operatorname{atanh}(c\sqrt{x}) - \frac{b(\operatorname{atanh}(c\sqrt{x}) - c\sqrt{x})}{c^2}$$

input

```
int(a + b*atanh(c*x^(1/2)),x)
```

output

```
a*x + b*x*atanh(c*x^(1/2)) - (b*(atanh(c*x^(1/2)) - c*x^(1/2)))/c^2
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.90

$$\int (a + b \operatorname{arctanh}(c\sqrt{x})) dx = \frac{\operatorname{atanh}(\sqrt{x}c) b c^2 x - \operatorname{atanh}(\sqrt{x}c) b + \sqrt{x} b c + a c^2 x}{c^2}$$

input

```
int(a+b*atanh(c*x^(1/2)),x)
```

output

```
(atanh(sqrt(x)*c)*b*c**2*x - atanh(sqrt(x)*c)*b + sqrt(x)*b*c + a*c**2*x)/
c**2
```

3.190 $\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x} dx$

Optimal result	1513
Mathematica [A] (verified)	1513
Rubi [A] (verified)	1514
Maple [B] (verified)	1515
Fricas [F]	1515
Sympy [F]	1516
Maxima [B] (verification not implemented)	1516
Giac [F]	1516
Mupad [F(-1)]	1517
Reduce [F]	1517

Optimal result

Integrand size = 16, antiderivative size = 29

$$\int \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{x} dx = a \log(x) - b \operatorname{PolyLog}(2, -c\sqrt{x}) + b \operatorname{PolyLog}(2, c\sqrt{x})$$

output `a*ln(x)-b*polylog(2,-c*x^(1/2))+b*polylog(2,c*x^(1/2))`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{x} dx = a \log(x) - b \operatorname{PolyLog}(2, -c\sqrt{x}) + b \operatorname{PolyLog}(2, c\sqrt{x})$$

input `Integrate[(a + b*ArcTanh[c*Sqrt[x]])/x,x]`

output `a*Log[x] - b*PolyLog[2, -(c*Sqrt[x])] + b*PolyLog[2, c*Sqrt[x]]`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.38, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {6450, 6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x} dx$$

↓ 6450

$$2 \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}} d\sqrt{x}$$

↓ 6446

$$2 \left(a \log(\sqrt{x}) - \frac{1}{2} b \operatorname{PolyLog}(2, -c\sqrt{x}) + \frac{1}{2} b \operatorname{PolyLog}(2, c\sqrt{x}) \right)$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])/x,x]`

output `2*(a*Log[Sqrt[x]] - (b*PolyLog[2, -(c*Sqrt[x])])/2 + (b*PolyLog[2, c*Sqrt[x]])/2)`

Defintions of rubi rules used

rule 6446 `Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))/(x_), x_Symbol] :> Simp[a*Log[x], x] + (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) / ; FreeQ[{a, b, c}, x]`

rule 6450 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)]*(b_.))^(p_.)/(x_), x_Symbol] :> Simp[1/n Subst[Int[(a + b*ArcTanh[c*x])^p/x, x], x, x^n], x] / ; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. $2(25) = 50$.

Time = 0.36 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.97

method	result
parts	$a \ln(x) + b(2 \ln(c\sqrt{x}) \operatorname{arctanh}(c\sqrt{x}) - \operatorname{dilog}(c\sqrt{x}) - \operatorname{dilog}(1 + c\sqrt{x}) - \ln(c\sqrt{x}))$
derivativedivides	$2a \ln(c\sqrt{x}) + 2b \left(\ln(c\sqrt{x}) \operatorname{arctanh}(c\sqrt{x}) - \frac{\operatorname{dilog}(c\sqrt{x})}{2} - \frac{\operatorname{dilog}(1+c\sqrt{x})}{2} - \frac{\ln(c\sqrt{x}) \ln(1+c\sqrt{x})}{2} \right)$
default	$2a \ln(c\sqrt{x}) + 2b \left(\ln(c\sqrt{x}) \operatorname{arctanh}(c\sqrt{x}) - \frac{\operatorname{dilog}(c\sqrt{x})}{2} - \frac{\operatorname{dilog}(1+c\sqrt{x})}{2} - \frac{\ln(c\sqrt{x}) \ln(1+c\sqrt{x})}{2} \right)$

input `int((a+b*arctanh(c*x^(1/2)))/x,x,method=_RETURNVERBOSE)`

output `a*ln(x)+b*(2*ln(c*x^(1/2))*arctanh(c*x^(1/2))-dilog(c*x^(1/2))-dilog(1+c*x^(1/2))-ln(c*x^(1/2))*ln(1+c*x^(1/2)))`

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x} dx = \int \frac{b \operatorname{arctanh}(c\sqrt{x}) + a}{x} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x,x, algorithm="fricas")`

output `integral((b*arctanh(c*sqrt(x)) + a)/x, x)`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x} dx = \int \frac{a + b \operatorname{atanh}(c\sqrt{x})}{x} dx$$

input `integrate((a+b*atanh(c*x**(1/2)))/x,x)`

output `Integral((a + b*atanh(c*sqrt(x)))/x, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs. $2(23) = 46$.

Time = 0.12 (sec) , antiderivative size = 61, normalized size of antiderivative = 2.10

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x} dx = -(\log(c\sqrt{x}) \log(-c\sqrt{x} + 1) + \operatorname{Li}_2(-c\sqrt{x} + 1))b \\ + (\log(c\sqrt{x} + 1) \log(-c\sqrt{x}) + \operatorname{Li}_2(c\sqrt{x} + 1))b + a \log(x)$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x,x, algorithm="maxima")`

output `-(log(c*sqrt(x))*log(-c*sqrt(x) + 1) + dilog(-c*sqrt(x) + 1))*b + (log(c*sqrt(x) + 1)*log(-c*sqrt(x)) + dilog(c*sqrt(x) + 1))*b + a*log(x)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x} dx = \int \frac{b \operatorname{artanh}(c\sqrt{x}) + a}{x} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x,x, algorithm="giac")`

output `integrate((b*arctanh(c*sqrt(x)) + a)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x} dx = \int \frac{a + b \operatorname{atanh}(c\sqrt{x})}{x} dx$$

input `int((a + b*atanh(c*x^(1/2)))/x,x)`output `int((a + b*atanh(c*x^(1/2)))/x, x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x} dx = \left(\int \frac{\operatorname{atanh}(\sqrt{x}c)}{x} dx \right) b + \log(x) a$$

input `int((a+b*atanh(c*x^(1/2)))/x,x)`output `int(atanh(sqrt(x)*c)/x,x)*b + log(x)*a`

3.191 $\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2} dx$

Optimal result	1518
Mathematica [A] (verified)	1518
Rubi [A] (verified)	1519
Maple [A] (verified)	1520
Fricas [A] (verification not implemented)	1521
Sympy [B] (verification not implemented)	1521
Maxima [A] (verification not implemented)	1522
Giac [B] (verification not implemented)	1522
Mupad [B] (verification not implemented)	1523
Reduce [B] (verification not implemented)	1523

Optimal result

Integrand size = 16, antiderivative size = 40

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2} dx = -\frac{bc}{\sqrt{x}} + bc^2 \operatorname{arctanh}(c\sqrt{x}) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x}$$

output

```
-b*c/x^(1/2)+b*c^2*arctanh(c*x^(1/2))-(a+b*arctanh(c*x^(1/2)))/x
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.68

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2} dx = -\frac{a}{x} - \frac{bc}{\sqrt{x}} - \frac{b \operatorname{arctanh}(c\sqrt{x})}{x} - \frac{1}{2}bc^2 \log(1 - c\sqrt{x}) + \frac{1}{2}bc^2 \log(1 + c\sqrt{x})$$

input

```
Integrate[(a + b*ArcTanh[c*Sqrt[x]])/x^2,x]
```

output

```
-(a/x) - (b*c)/Sqrt[x] - (b*ArcTanh[c*Sqrt[x]])/x - (b*c^2*Log[1 - c*Sqrt[x]])/2 + (b*c^2*Log[1 + c*Sqrt[x]])/2
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.08, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6452, 61, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2} dx$$

$$\downarrow 6452$$

$$\frac{1}{2}bc \int \frac{1}{x^{3/2}(1-c^2x)} dx - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x}$$

$$\downarrow 61$$

$$\frac{1}{2}bc \left(c^2 \int \frac{1}{\sqrt{x}(1-c^2x)} dx - \frac{2}{\sqrt{x}} \right) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x}$$

$$\downarrow 73$$

$$\frac{1}{2}bc \left(2c^2 \int \frac{1}{1-c^2x} d\sqrt{x} - \frac{2}{\sqrt{x}} \right) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x}$$

$$\downarrow 219$$

$$\frac{1}{2}bc \left(2c \operatorname{arctanh}(c\sqrt{x}) - \frac{2}{\sqrt{x}} \right) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x}$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])/x^2,x]`

output `-((a + b*ArcTanh[c*Sqrt[x]])/x) + (b*c*(-2/Sqrt[x] + 2*c*ArcTanh[c*Sqrt[x]]))/2`

Definitions of rubi rules used

rule 61

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((
m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x],
x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0]
) || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n])) && IntLinearQ[a, b, c, d
, m, n, x]
```

rule 73

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 219

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1]
) && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.42

method	result	size
parts	$-\frac{a}{x} + 2b c^2 \left(-\frac{\operatorname{arctanh}(c\sqrt{x})}{2c^2 x} - \frac{\ln(c\sqrt{x}-1)}{4} - \frac{1}{2c\sqrt{x}} + \frac{\ln(1+c\sqrt{x})}{4} \right)$	57
derivativedivides	$2c^2 \left(-\frac{a}{2c^2 x} + b \left(-\frac{\operatorname{arctanh}(c\sqrt{x})}{2c^2 x} - \frac{\ln(c\sqrt{x}-1)}{4} - \frac{1}{2c\sqrt{x}} + \frac{\ln(1+c\sqrt{x})}{4} \right) \right)$	61
default	$2c^2 \left(-\frac{a}{2c^2 x} + b \left(-\frac{\operatorname{arctanh}(c\sqrt{x})}{2c^2 x} - \frac{\ln(c\sqrt{x}-1)}{4} - \frac{1}{2c\sqrt{x}} + \frac{\ln(1+c\sqrt{x})}{4} \right) \right)$	61

input `int((a+b*arctanh(c*x^(1/2)))/x^2,x,method=_RETURNVERBOSE)`

output `-a/x+2*b*c^2*(-1/2/c^2/x*arctanh(c*x^(1/2))-1/4*ln(c*x^(1/2)-1)-1/2/c/x^(1/2)+1/4*ln(1+c*x^(1/2)))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.32

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2} dx = -\frac{2bc\sqrt{x} - (bc^2x - b) \log\left(-\frac{c^2x+2c\sqrt{x}+1}{c^2x-1}\right) + 2a}{2x}$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^2,x, algorithm="fricas")`

output `-1/2*(2*b*c*sqrt(x) - (b*c^2*x - b)*log(-(c^2*x + 2*c*sqrt(x) + 1)/(c^2*x - 1)) + 2*a)/x`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 231 vs. 2(34) = 68.

Time = 2.56 (sec) , antiderivative size = 231, normalized size of antiderivative = 5.78

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2} dx = \begin{cases} -\frac{a}{x} + \frac{b \operatorname{atanh}\left(\sqrt{x}\sqrt{\frac{1}{x}}\right)}{x} & \text{for } c \\ -\frac{a}{x} - \frac{b \operatorname{atanh}\left(\sqrt{x}\sqrt{\frac{1}{x}}\right)}{x} & \text{for } c \\ -\frac{ac^2x^{\frac{3}{2}}}{c^2x^{\frac{5}{2}}-x^{\frac{3}{2}}} + \frac{a\sqrt{x}}{c^2x^{\frac{5}{2}}-x^{\frac{3}{2}}} + \frac{bc^4x^{\frac{5}{2}} \operatorname{atanh}(c\sqrt{x})}{c^2x^{\frac{5}{2}}-x^{\frac{3}{2}}} - \frac{bc^3x^2}{c^2x^{\frac{5}{2}}-x^{\frac{3}{2}}} - \frac{2bc^2x^{\frac{3}{2}} \operatorname{atanh}(c\sqrt{x})}{c^2x^{\frac{5}{2}}-x^{\frac{3}{2}}} + \frac{bcx}{c^2x^{\frac{5}{2}}-x^{\frac{3}{2}}} + \frac{b\sqrt{x} \operatorname{atanh}(c\sqrt{x})}{c^2x^{\frac{5}{2}}-x^{\frac{3}{2}}} & \text{other} \end{cases}$$

input `integrate((a+b*atanh(c*x**(1/2)))/x**2,x)`

output

```
Piecewise((-a/x + b*atanh(sqrt(x)*sqrt(1/x))/x, Eq(c, -sqrt(1/x))), (-a/x - b*atanh(sqrt(x)*sqrt(1/x))/x, Eq(c, sqrt(1/x))), (-a*c**2*x**(3/2)/(c**2*x**(5/2) - x**(3/2)) + a*sqrt(x)/(c**2*x**(5/2) - x**(3/2)) + b*c**4*x**(5/2)*atanh(c*sqrt(x))/(c**2*x**(5/2) - x**(3/2)) - b*c**3*x**2/(c**2*x**(5/2) - x**(3/2)) - 2*b*c**2*x**(3/2)*atanh(c*sqrt(x))/(c**2*x**(5/2) - x**(3/2)) + b*c*x/(c**2*x**(5/2) - x**(3/2)) + b*sqrt(x)*atanh(c*sqrt(x))/(c**2*x**(5/2) - x**(3/2)), True))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.28

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2} dx$$

$$= \frac{1}{2} \left(\left(c \log(c\sqrt{x} + 1) - c \log(c\sqrt{x} - 1) - \frac{2}{\sqrt{x}} \right) c - \frac{2 \operatorname{arctanh}(c\sqrt{x})}{x} \right) b - \frac{a}{x}$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))/x^2,x, algorithm="maxima")
```

output

```
1/2*((c*log(c*sqrt(x) + 1) - c*log(c*sqrt(x) - 1) - 2/sqrt(x))*c - 2*arctanh(c*sqrt(x))/x)*b - a/x
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 168 vs. 2(34) = 68.

Time = 0.13 (sec) , antiderivative size = 168, normalized size of antiderivative = 4.20

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2} dx$$

$$= 2 \left(\frac{(c\sqrt{x} + 1)bc \log\left(-\frac{c\sqrt{x}+1}{c\sqrt{x}-1}\right)}{(c\sqrt{x} - 1) \left(\frac{(c\sqrt{x}+1)^2}{(c\sqrt{x}-1)^2} + \frac{2(c\sqrt{x}+1)}{c\sqrt{x}-1} + 1 \right)} + \frac{\frac{2(c\sqrt{x}+1)ac}{c\sqrt{x}-1} + \frac{(c\sqrt{x}+1)bc}{c\sqrt{x}-1} + bc}{\frac{(c\sqrt{x}+1)^2}{(c\sqrt{x}-1)^2} + \frac{2(c\sqrt{x}+1)}{c\sqrt{x}-1} + 1} \right) c$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))/x^2,x, algorithm="giac")
```

output

$$2*((c*\sqrt{x} + 1)*b*c*\log(-(c*\sqrt{x} + 1)/(c*\sqrt{x} - 1)))/((c*\sqrt{x} - 1)*((c*\sqrt{x} + 1)^2/(c*\sqrt{x} - 1)^2 + 2*(c*\sqrt{x} + 1)/(c*\sqrt{x} - 1) + 1)) + (2*(c*\sqrt{x} + 1)*a*c/(c*\sqrt{x} - 1) + (c*\sqrt{x} + 1)*b*c/(c*\sqrt{x} - 1) + b*c)/((c*\sqrt{x} + 1)^2/(c*\sqrt{x} - 1)^2 + 2*(c*\sqrt{x} + 1)/(c*\sqrt{x} - 1) + 1)*c$$
Mupad [B] (verification not implemented)

Time = 4.35 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.30

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2} dx = b c \operatorname{atan}\left(\frac{c^2 \sqrt{x}}{\sqrt{-c^2}}\right) \sqrt{-c^2} - \frac{a}{x} - \frac{b \operatorname{atanh}(c\sqrt{x}) + b c \sqrt{x}}{x}$$

input

$$\operatorname{int}((a + b*\operatorname{atanh}(c*x^{(1/2)}))/x^2, x)$$

output

$$b*c*\operatorname{atan}((c^2*x^{(1/2)})/(-c^2)^{(1/2)})*(-c^2)^{(1/2)} - a/x - (b*\operatorname{atanh}(c*x^{(1/2)}) + b*c*x^{(1/2)})/x$$
Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.82

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2} dx = \frac{\operatorname{atanh}(\sqrt{x}c) b c^2 x - \operatorname{atanh}(\sqrt{x}c) b - \sqrt{x} b c - a}{x}$$

input

$$\operatorname{int}((a+b*\operatorname{atanh}(c*x^{(1/2)}))/x^2, x)$$

output

$$(\operatorname{atanh}(\sqrt{x}*c)*b*c**2*x - \operatorname{atanh}(\sqrt{x}*c)*b - \sqrt{x}*b*c - a)/x$$

3.192 $\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x^3} dx$

Optimal result	1524
Mathematica [A] (verified)	1524
Rubi [A] (verified)	1525
Maple [A] (verified)	1526
Fricas [A] (verification not implemented)	1527
Sympy [B] (verification not implemented)	1527
Maxima [A] (verification not implemented)	1528
Giac [B] (verification not implemented)	1528
Mupad [B] (verification not implemented)	1529
Reduce [B] (verification not implemented)	1530

Optimal result

Integrand size = 16, antiderivative size = 60

$$\int \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{x^3} dx = -\frac{bc}{6x^{3/2}} - \frac{bc^3}{2\sqrt{x}} + \frac{1}{2}bc^4\operatorname{arctanh}(c\sqrt{x}) - \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{2x^2}$$

output

$$-1/6*b*c/x^(3/2)-1/2*b*c^3/x^(1/2)+1/2*b*c^4*\operatorname{arctanh}(c*x^(1/2))-1/2*(a+b*\operatorname{arctanh}(c*x^(1/2)))/x^2$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.43

$$\int \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{x^3} dx = -\frac{a}{2x^2} - \frac{bc}{6x^{3/2}} - \frac{bc^3}{2\sqrt{x}} - \frac{b\operatorname{arctanh}(c\sqrt{x})}{2x^2} - \frac{1}{4}bc^4 \log(1 - c\sqrt{x}) + \frac{1}{4}bc^4 \log(1 + c\sqrt{x})$$

input

$$\operatorname{Integrate}[(a + b*\operatorname{ArcTanh}[c*\operatorname{Sqrt}[x]])/x^3, x]$$

output

$$-1/2*a/x^2 - (b*c)/(6*x^(3/2)) - (b*c^3)/(2*\operatorname{Sqrt}[x]) - (b*\operatorname{ArcTanh}[c*\operatorname{Sqrt}[x]])/(2*x^2) - (b*c^4*\operatorname{Log}[1 - c*\operatorname{Sqrt}[x]])/4 + (b*c^4*\operatorname{Log}[1 + c*\operatorname{Sqrt}[x]])/4$$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.98, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6452, 61, 61, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3} dx \\
 & \quad \downarrow 6452 \\
 & \frac{1}{4}bc \int \frac{1}{x^{5/2}(1-c^2x)} dx - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{2x^2} \\
 & \quad \downarrow 61 \\
 & \frac{1}{4}bc \left(c^2 \int \frac{1}{x^{3/2}(1-c^2x)} dx - \frac{2}{3x^{3/2}} \right) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{2x^2} \\
 & \quad \downarrow 61 \\
 & \frac{1}{4}bc \left(c^2 \left(c^2 \int \frac{1}{\sqrt{x}(1-c^2x)} dx - \frac{2}{\sqrt{x}} \right) - \frac{2}{3x^{3/2}} \right) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{2x^2} \\
 & \quad \downarrow 73 \\
 & \frac{1}{4}bc \left(c^2 \left(2c^2 \int \frac{1}{1-c^2x} d\sqrt{x} - \frac{2}{\sqrt{x}} \right) - \frac{2}{3x^{3/2}} \right) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{2x^2} \\
 & \quad \downarrow 219 \\
 & \frac{1}{4}bc \left(c^2 \left(2 \operatorname{arctanh}(c\sqrt{x}) - \frac{2}{\sqrt{x}} \right) - \frac{2}{3x^{3/2}} \right) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{2x^2}
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])/x^3,x]`

output `-1/2*(a + b*ArcTanh[c*Sqrt[x]])/x^2 + (b*c*(-2/(3*x^(3/2)) + c^2*(-2/Sqrt[x] + 2*c*ArcTanh[c*Sqrt[x]])))/4`

Defintions of rubi rules used

```
rule 61 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((
m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x],
x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0
] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d
, m, n, x]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 6452 Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.08

method	result	size
parts	$-\frac{a}{2x^2} + 2bc^4 \left(-\frac{\operatorname{arctanh}(c\sqrt{x})}{4c^4x^2} - \frac{\ln(c\sqrt{x}-1)}{8} + \frac{\ln(1+c\sqrt{x})}{8} - \frac{1}{12c^3x^{\frac{3}{2}}} - \frac{1}{4c\sqrt{x}} \right)$	65
derivativedivides	$2c^4 \left(-\frac{a}{4c^4x^2} + b \left(-\frac{\operatorname{arctanh}(c\sqrt{x})}{4c^4x^2} - \frac{\ln(c\sqrt{x}-1)}{8} + \frac{\ln(1+c\sqrt{x})}{8} - \frac{1}{12c^3x^{\frac{3}{2}}} - \frac{1}{4c\sqrt{x}} \right) \right)$	69
default	$2c^4 \left(-\frac{a}{4c^4x^2} + b \left(-\frac{\operatorname{arctanh}(c\sqrt{x})}{4c^4x^2} - \frac{\ln(c\sqrt{x}-1)}{8} + \frac{\ln(1+c\sqrt{x})}{8} - \frac{1}{12c^3x^{\frac{3}{2}}} - \frac{1}{4c\sqrt{x}} \right) \right)$	69

input `int((a+b*arctanh(c*x^(1/2)))/x^3,x,method=_RETURNVERBOSE)`

output
$$-1/2*a/x^2+2*b*c^4*(-1/4/c^4/x^2*arctanh(c*x^(1/2))-1/8*\ln(c*x^(1/2)-1)+1/8*\ln(1+c*x^(1/2))-1/12/c^3/x^(3/2)-1/4/c/x^(1/2))$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.07

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3} dx = \frac{3(bc^4x^2 - b) \log\left(-\frac{c^2x+2c\sqrt{x}+1}{c^2x-1}\right) - 2(3bc^3x + bc)\sqrt{x} - 6a}{12x^2}$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^3,x, algorithm="fricas")`

output
$$1/12*(3*(b*c^4*x^2 - b)*\log(-(c^2*x + 2*c*\sqrt{x} + 1)/(c^2*x - 1)) - 2*(3*b*c^3*x + b*c)*\sqrt{x} - 6*a)/x^2$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 342 vs. 2(54) = 108.

Time = 7.34 (sec) , antiderivative size = 342, normalized size of antiderivative = 5.70

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3} dx = \begin{cases} -\frac{a}{2x^2} + \frac{b \operatorname{atanh}\left(\sqrt{x}\sqrt{\frac{1}{x}}\right)}{2x^2} \\ -\frac{a}{2x^2} - \frac{b \operatorname{atanh}\left(\sqrt{x}\sqrt{\frac{1}{x}}\right)}{2x^2} \\ -\frac{3ac^2x^{\frac{3}{2}}}{6c^2x^{\frac{7}{2}}-6x^{\frac{5}{2}}} + \frac{3a\sqrt{x}}{6c^2x^{\frac{7}{2}}-6x^{\frac{5}{2}}} + \frac{3bc^6x^{\frac{7}{2}} \operatorname{atanh}(c\sqrt{x})}{6c^2x^{\frac{7}{2}}-6x^{\frac{5}{2}}} - \frac{3bc^5x^3}{6c^2x^{\frac{7}{2}}-6x^{\frac{5}{2}}} - \frac{3bc^4x^{\frac{5}{2}} \operatorname{atanh}(c\sqrt{x})}{6c^2x^{\frac{7}{2}}-6x^{\frac{5}{2}}} + \frac{2bc^3x^2}{6c^2x^{\frac{7}{2}}-6x^{\frac{5}{2}}} - \frac{3bc^2x^{\frac{3}{2}} \operatorname{atanh}(c\sqrt{x})}{6c^2x^{\frac{7}{2}}-6x^{\frac{5}{2}}} \end{cases}$$

input `integrate((a+b*atanh(c*x**(1/2)))/x**3,x)`

output

```
Piecewise((-a/(2*x**2) + b*atanh(sqrt(x)*sqrt(1/x))/(2*x**2), Eq(c, -sqrt(1/x))), (-a/(2*x**2) - b*atanh(sqrt(x)*sqrt(1/x))/(2*x**2), Eq(c, sqrt(1/x))), (-3*a*c**2*x**(3/2)/(6*c**2*x**(7/2) - 6*x**(5/2)) + 3*a*sqrt(x)/(6*c**2*x**(7/2) - 6*x**(5/2)) + 3*b*c**6*x**(7/2)*atanh(c*sqrt(x))/(6*c**2*x**(7/2) - 6*x**(5/2)) - 3*b*c**5*x**3/(6*c**2*x**(7/2) - 6*x**(5/2)) - 3*b*c**4*x**(5/2)*atanh(c*sqrt(x))/(6*c**2*x**(7/2) - 6*x**(5/2)) + 2*b*c**3*x**2/(6*c**2*x**(7/2) - 6*x**(5/2)) - 3*b*c**2*x**(3/2)*atanh(c*sqrt(x))/(6*c**2*x**(7/2) - 6*x**(5/2)) + b*c*x/(6*c**2*x**(7/2) - 6*x**(5/2)) + 3*b*sqrt(x)*atanh(c*sqrt(x))/(6*c**2*x**(7/2) - 6*x**(5/2)), True))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.07

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3} dx$$

$$= \frac{1}{12} \left(\left(3c^3 \log(c\sqrt{x} + 1) - 3c^3 \log(c\sqrt{x} - 1) - \frac{2(3c^2x + 1)}{x^{\frac{3}{2}}} \right) c - \frac{6 \operatorname{arctanh}(c\sqrt{x})}{x^2} \right) b - \frac{a}{2x^2}$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))/x^3,x, algorithm="maxima")
```

output

```
1/12*((3*c^3*log(c*sqrt(x) + 1) - 3*c^3*log(c*sqrt(x) - 1) - 2*(3*c^2*x + 1)/x^(3/2))*c - 6*arctanh(c*sqrt(x))/x^2)*b - 1/2*a/x^2
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 356 vs. 2(44) = 88.

Time = 0.14 (sec) , antiderivative size = 356, normalized size of antiderivative = 5.93

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3} dx$$

$$= \frac{2}{3} c \left(\frac{3 \left(\frac{(c\sqrt{x}+1)^3 bc^3}{(c\sqrt{x}-1)^3} + \frac{(c\sqrt{x}+1)bc^3}{c\sqrt{x}-1} \right) \log \left(-\frac{c\sqrt{x}+1}{c\sqrt{x}-1} \right)}{\frac{(c\sqrt{x}+1)^4}{(c\sqrt{x}-1)^4} + \frac{4(c\sqrt{x}+1)^3}{(c\sqrt{x}-1)^3} + \frac{6(c\sqrt{x}+1)^2}{(c\sqrt{x}-1)^2} + \frac{4(c\sqrt{x}+1)}{c\sqrt{x}-1} + 1} + \frac{\frac{6(c\sqrt{x}+1)^3 ac^3}{(c\sqrt{x}-1)^3} + \frac{6(c\sqrt{x}+1)ac^3}{c\sqrt{x}-1} + \frac{3(c\sqrt{x}+1)^3 bc^3}{(c\sqrt{x}-1)^3} + \frac{6(c\sqrt{x}+1)^2 bc^3}{(c\sqrt{x}-1)^2}}{\frac{(c\sqrt{x}+1)^4}{(c\sqrt{x}-1)^4} + \frac{4(c\sqrt{x}+1)^3}{(c\sqrt{x}-1)^3} + \frac{6(c\sqrt{x}+1)^2}{(c\sqrt{x}-1)^2}} \right)$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^3,x, algorithm="giac")`

output `2/3*c*(3*((c*sqrt(x) + 1)^3*b*c^3/(c*sqrt(x) - 1)^3 + (c*sqrt(x) + 1)*b*c^3/(c*sqrt(x) - 1))*log(-(c*sqrt(x) + 1)/(c*sqrt(x) - 1)))/((c*sqrt(x) + 1)^4/(c*sqrt(x) - 1)^4 + 4*(c*sqrt(x) + 1)^3/(c*sqrt(x) - 1)^3 + 6*(c*sqrt(x) + 1)^2/(c*sqrt(x) - 1)^2 + 4*(c*sqrt(x) + 1)/(c*sqrt(x) - 1) + 1) + (6*(c*sqrt(x) + 1)^3*a*c^3/(c*sqrt(x) - 1)^3 + 6*(c*sqrt(x) + 1)*a*c^3/(c*sqrt(x) - 1) + 3*(c*sqrt(x) + 1)^3*b*c^3/(c*sqrt(x) - 1)^3 + 6*(c*sqrt(x) + 1)^2*b*c^3/(c*sqrt(x) - 1)^2 + 5*(c*sqrt(x) + 1)*b*c^3/(c*sqrt(x) - 1) + 2*b*c^3)/((c*sqrt(x) + 1)^4/(c*sqrt(x) - 1)^4 + 4*(c*sqrt(x) + 1)^3/(c*sqrt(x) - 1)^3 + 6*(c*sqrt(x) + 1)^2/(c*sqrt(x) - 1)^2 + 4*(c*sqrt(x) + 1)/(c*sqrt(x) - 1) + 1))`

Mupad [B] (verification not implemented)

Time = 4.30 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.02

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3} dx = \frac{b c^4 \operatorname{atanh}(c\sqrt{x})}{2} - \frac{b (3 \ln(c\sqrt{x} + 1) - 3 \ln(1 - c\sqrt{x}) + 2c\sqrt{x} + 6c^3 x^{3/2})}{12x^2} - \frac{a}{2x^2}$$

input `int((a + b*atanh(c*x^(1/2)))/x^3,x)`

output `(b*c^4*atanh(c*x^(1/2)))/2 - (b*(3*log(c*x^(1/2) + 1) - 3*log(1 - c*x^(1/2))) + 2*c*x^(1/2) + 6*c^3*x^(3/2))/(12*x^2) - a/(2*x^2)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.77

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3} dx$$

$$= \frac{3 \operatorname{atanh}(\sqrt{x} c) b c^4 x^2 - 3 \operatorname{atanh}(\sqrt{x} c) b - 3\sqrt{x} b c^3 x - \sqrt{x} b c - 3a}{6x^2}$$

input `int((a+b*atanh(c*x^(1/2)))/x^3,x)`output `(3*atanh(sqrt(x)*c)*b*c**4*x**2 - 3*atanh(sqrt(x)*c)*b - 3*sqrt(x)*b*c**3*x - sqrt(x)*b*c - 3*a)/(6*x**2)`

3.193 $\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x^4} dx$

Optimal result	1531
Mathematica [A] (verified)	1531
Rubi [A] (verified)	1532
Maple [A] (verified)	1534
Fricas [A] (verification not implemented)	1534
Sympy [B] (verification not implemented)	1535
Maxima [A] (verification not implemented)	1535
Giac [B] (verification not implemented)	1536
Mupad [B] (verification not implemented)	1537
Reduce [B] (verification not implemented)	1537

Optimal result

Integrand size = 16, antiderivative size = 73

$$\int \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{x^4} dx = -\frac{bc}{15x^{5/2}} - \frac{bc^3}{9x^{3/2}} - \frac{bc^5}{3\sqrt{x}} + \frac{1}{3}bc^6\operatorname{arctanh}(c\sqrt{x}) - \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{3x^3}$$

output

```
-1/15*b*c/x^(5/2)-1/9*b*c^3/x^(3/2)-1/3*b*c^5/x^(1/2)+1/3*b*c^6*arctanh(c*
x^(1/2))-1/3*(a+b*arctanh(c*x^(1/2)))/x^3
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.36

$$\int \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{x^4} dx = -\frac{a}{3x^3} - \frac{bc}{15x^{5/2}} - \frac{bc^3}{9x^{3/2}} - \frac{bc^5}{3\sqrt{x}} - \frac{b\operatorname{arctanh}(c\sqrt{x})}{3x^3} - \frac{1}{6}bc^6 \log(1 - c\sqrt{x}) + \frac{1}{6}bc^6 \log(1 + c\sqrt{x})$$

input

```
Integrate[(a + b*ArcTanh[c*Sqrt[x]])/x^4,x]
```


output

$$-1/3*a/x^3 - (b*c)/(15*x^(5/2)) - (b*c^3)/(9*x^(3/2)) - (b*c^5)/(3*\text{Sqrt}[x]) - (b*\text{ArcTanh}[c*\text{Sqrt}[x]])/(3*x^3) - (b*c^6*\text{Log}[1 - c*\text{Sqrt}[x]])/6 + (b*c^6*\text{Log}[1 + c*\text{Sqrt}[x]])/6$$
Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6452, 61, 61, 61, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4} dx$$

$$\downarrow 6452$$

$$\frac{1}{6}bc \int \frac{1}{x^{7/2}(1-c^2x)} dx - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{3x^3}$$

$$\downarrow 61$$

$$\frac{1}{6}bc \left(c^2 \int \frac{1}{x^{5/2}(1-c^2x)} dx - \frac{2}{5x^{5/2}} \right) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{3x^3}$$

$$\downarrow 61$$

$$\frac{1}{6}bc \left(c^2 \left(c^2 \int \frac{1}{x^{3/2}(1-c^2x)} dx - \frac{2}{3x^{3/2}} \right) - \frac{2}{5x^{5/2}} \right) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{3x^3}$$

$$\downarrow 61$$

$$\frac{1}{6}bc \left(c^2 \left(c^2 \left(c^2 \int \frac{1}{\sqrt{x}(1-c^2x)} dx - \frac{2}{\sqrt{x}} \right) - \frac{2}{3x^{3/2}} \right) - \frac{2}{5x^{5/2}} \right) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{3x^3}$$

$$\downarrow 73$$

$$\frac{1}{6}bc \left(c^2 \left(c^2 \left(2c^2 \int \frac{1}{1-c^2x} d\sqrt{x} - \frac{2}{\sqrt{x}} \right) - \frac{2}{3x^{3/2}} \right) - \frac{2}{5x^{5/2}} \right) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{3x^3}$$

$$\downarrow 219$$

$$\frac{1}{6}bc \left(c^2 \left(c^2 \left(2 \operatorname{arctanh}(c\sqrt{x}) - \frac{2}{\sqrt{x}} \right) - \frac{2}{3x^{3/2}} \right) - \frac{2}{5x^{5/2}} \right) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{3x^3}$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])/x^4,x]`

output `-1/3*(a + b*ArcTanh[c*Sqrt[x]])/x^3 + (b*c*(-2/(5*x^(5/2)) + c^2*(-2/(3*x^(3/2)) + c^2*(-2/Sqrt[x] + 2*c*ArcTanh[c*Sqrt[x]]))))/6`

Defintions of rubi rules used

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00

method	result	si
parts	$-\frac{a}{3x^3} + 2bc^6 \left(-\frac{\operatorname{arctanh}(c\sqrt{x})}{6c^6x^3} - \frac{1}{30c^5x^{\frac{5}{2}}} - \frac{1}{18c^3x^{\frac{3}{2}}} - \frac{1}{6c\sqrt{x}} - \frac{\ln(c\sqrt{x}-1)}{12} + \frac{\ln(1+c\sqrt{x})}{12} \right)$	73
derivativedivides	$2c^6 \left(-\frac{a}{6c^6x^3} + b \left(-\frac{\operatorname{arctanh}(c\sqrt{x})}{6c^6x^3} - \frac{1}{30c^5x^{\frac{5}{2}}} - \frac{1}{18c^3x^{\frac{3}{2}}} - \frac{1}{6c\sqrt{x}} - \frac{\ln(c\sqrt{x}-1)}{12} + \frac{\ln(1+c\sqrt{x})}{12} \right) \right)$	73
default	$2c^6 \left(-\frac{a}{6c^6x^3} + b \left(-\frac{\operatorname{arctanh}(c\sqrt{x})}{6c^6x^3} - \frac{1}{30c^5x^{\frac{5}{2}}} - \frac{1}{18c^3x^{\frac{3}{2}}} - \frac{1}{6c\sqrt{x}} - \frac{\ln(c\sqrt{x}-1)}{12} + \frac{\ln(1+c\sqrt{x})}{12} \right) \right)$	73

input `int((a+b*arctanh(c*x^(1/2)))/x^4,x,method=_RETURNVERBOSE)`

output
$$-1/3*a/x^3+2*b*c^6*(-1/6/c^6/x^3*arctanh(c*x^(1/2))-1/30/c^5/x^(5/2)-1/18/c^3/x^(3/2)-1/6/c/x^(1/2)-1/12*\ln(c*x^(1/2)-1)+1/12*\ln(1+c*x^(1/2)))$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.01

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4} dx$$

$$= \frac{15(bc^6x^3 - b) \log\left(-\frac{c^2x+2c\sqrt{x}+1}{c^2x-1}\right) - 2(15bc^5x^2 + 5bc^3x + 3bc)\sqrt{x} - 30a}{90x^3}$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^4,x, algorithm="fricas")`

output
$$1/90*(15*(b*c^6*x^3 - b)*\log(-(c^2*x + 2*c*\sqrt{x} + 1)/(c^2*x - 1)) - 2*(15*b*c^5*x^2 + 5*b*c^3*x + 3*b*c)*\sqrt{x} - 30*a)/x^3$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 371 vs. $2(66) = 132$.

Time = 21.14 (sec) , antiderivative size = 371, normalized size of antiderivative = 5.08

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4} dx$$

$$= \begin{cases} -\frac{a}{3x^3} + \frac{b \operatorname{arctanh}\left(\sqrt{x}\sqrt{\frac{1}{x}}\right)}{3x^3} \\ -\frac{a}{3x^3} - \frac{b \operatorname{arctanh}\left(\sqrt{x}\sqrt{\frac{1}{x}}\right)}{3x^3} \\ -\frac{15ac^2x^{\frac{3}{2}}}{45c^2x^{\frac{9}{2}}-45x^{\frac{7}{2}}} + \frac{15a\sqrt{x}}{45c^2x^{\frac{9}{2}}-45x^{\frac{7}{2}}} + \frac{15bc^8x^{\frac{9}{2}} \operatorname{arctanh}(c\sqrt{x})}{45c^2x^{\frac{9}{2}}-45x^{\frac{7}{2}}} - \frac{15bc^7x^4}{45c^2x^{\frac{9}{2}}-45x^{\frac{7}{2}}} - \frac{15bc^6x^{\frac{7}{2}} \operatorname{arctanh}(c\sqrt{x})}{45c^2x^{\frac{9}{2}}-45x^{\frac{7}{2}}} + \frac{10bc^5x^3}{45c^2x^{\frac{9}{2}}-45x^{\frac{7}{2}}} + \frac{2}{45c^2} \end{cases}$$

input `integrate((a+b*atanh(c*x**(1/2)))/x**4,x)`

output `Piecewise((-a/(3*x**3) + b*atanh(sqrt(x)*sqrt(1/x))/(3*x**3), Eq(c, -sqrt(1/x))), (-a/(3*x**3) - b*atanh(sqrt(x)*sqrt(1/x))/(3*x**3), Eq(c, sqrt(1/x))), (-15*a*c**2*x**(3/2)/(45*c**2*x**(9/2) - 45*x**(7/2)) + 15*a*sqrt(x)/(45*c**2*x**(9/2) - 45*x**(7/2)) + 15*b*c**8*x**(9/2)*atanh(c*sqrt(x))/(45*c**2*x**(9/2) - 45*x**(7/2)) - 15*b*c**7*x**4/(45*c**2*x**(9/2) - 45*x**(7/2)) - 15*b*c**6*x**(7/2)*atanh(c*sqrt(x))/(45*c**2*x**(9/2) - 45*x**(7/2)) + 10*b*c**5*x**3/(45*c**2*x**(9/2) - 45*x**(7/2)) + 2*b*c**3*x**2/(45*c**2*x**(9/2) - 45*x**(7/2)) - 15*b*c**2*x**(3/2)*atanh(c*sqrt(x))/(45*c**2*x**(9/2) - 45*x**(7/2)) + 3*b*c*x/(45*c**2*x**(9/2) - 45*x**(7/2)) + 15*b*sqrt(x)*atanh(c*sqrt(x))/(45*c**2*x**(9/2) - 45*x**(7/2)), True))`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.99

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4} dx$$

$$= \frac{1}{90} \left(\left(15c^5 \log(c\sqrt{x} + 1) - 15c^5 \log(c\sqrt{x} - 1) - \frac{2(15c^4x^2 + 5c^2x + 3)}{x^{\frac{5}{2}}} \right) c - \frac{30 \operatorname{arctanh}(c\sqrt{x})}{x^3} \right) b - \frac{a}{3x^3}$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^4,x, algorithm="maxima")`

output `1/90*((15*c^5*log(c*sqrt(x) + 1) - 15*c^5*log(c*sqrt(x) - 1) - 2*(15*c^4*x^2 + 5*c^2*x + 3)/x^(5/2))*c - 30*arctanh(c*sqrt(x))/x^3)*b - 1/3*a/x^3`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 534 vs. 2(53) = 106.

Time = 0.14 (sec) , antiderivative size = 534, normalized size of antiderivative = 7.32

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4} dx = \text{Too large to display}$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^4,x, algorithm="giac")`

output `2/45*c*(15*(3*(c*sqrt(x) + 1)^5*b*c^5/(c*sqrt(x) - 1)^5 + 10*(c*sqrt(x) + 1)^3*b*c^5/(c*sqrt(x) - 1)^3 + 3*(c*sqrt(x) + 1)*b*c^5/(c*sqrt(x) - 1))*log(-(c*sqrt(x) + 1)/(c*sqrt(x) - 1))/((c*sqrt(x) + 1)^6/(c*sqrt(x) - 1)^6 + 6*(c*sqrt(x) + 1)^5/(c*sqrt(x) - 1)^5 + 15*(c*sqrt(x) + 1)^4/(c*sqrt(x) - 1)^4 + 20*(c*sqrt(x) + 1)^3/(c*sqrt(x) - 1)^3 + 15*(c*sqrt(x) + 1)^2/(c*sqrt(x) - 1)^2 + 6*(c*sqrt(x) + 1)/(c*sqrt(x) - 1) + 1) + (90*(c*sqrt(x) + 1)^5*a*c^5/(c*sqrt(x) - 1)^5 + 300*(c*sqrt(x) + 1)^3*a*c^5/(c*sqrt(x) - 1)^3 + 90*(c*sqrt(x) + 1)*a*c^5/(c*sqrt(x) - 1) + 45*(c*sqrt(x) + 1)^5*b*c^5/(c*sqrt(x) - 1)^5 + 135*(c*sqrt(x) + 1)^4*b*c^5/(c*sqrt(x) - 1)^4 + 230*(c*sqrt(x) + 1)^3*b*c^5/(c*sqrt(x) - 1)^3 + 210*(c*sqrt(x) + 1)^2*b*c^5/(c*sqrt(x) - 1)^2 + 93*(c*sqrt(x) + 1)*b*c^5/(c*sqrt(x) - 1) + 23*b*c^5)/((c*sqrt(x) + 1)^6/(c*sqrt(x) - 1)^6 + 6*(c*sqrt(x) + 1)^5/(c*sqrt(x) - 1)^5 + 15*(c*sqrt(x) + 1)^4/(c*sqrt(x) - 1)^4 + 20*(c*sqrt(x) + 1)^3/(c*sqrt(x) - 1)^3 + 15*(c*sqrt(x) + 1)^2/(c*sqrt(x) - 1)^2 + 6*(c*sqrt(x) + 1)/(c*sqrt(x) - 1) + 1))`

Mupad [B] (verification not implemented)

Time = 4.07 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.95

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4} dx$$

$$= \frac{b c^6 \operatorname{atanh}(c\sqrt{x})}{3} - \frac{b (15 \ln(c\sqrt{x} + 1) - 15 \ln(1 - c\sqrt{x}) + 6c\sqrt{x} + 10c^3 x^{3/2} + 30c^5 x^{5/2})}{90x^3} - \frac{a}{3x^3}$$

input `int((a + b*atanh(c*x^(1/2)))/x^4,x)`output `(b*c^6*atanh(c*x^(1/2)))/3 - (b*(15*log(c*x^(1/2) + 1) - 15*log(1 - c*x^(1/2)) + 6*c*x^(1/2) + 10*c^3*x^(3/2) + 30*c^5*x^(5/2)))/(90*x^3) - a/(3*x^3)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.78

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4} dx$$

$$= \frac{15 \operatorname{atanh}(\sqrt{x}c) b c^6 x^3 - 15 \operatorname{atanh}(\sqrt{x}c) b - 15\sqrt{x} b c^5 x^2 - 5\sqrt{x} b c^3 x - 3\sqrt{x} b c - 15a}{45x^3}$$

input `int((a+b*atanh(c*x^(1/2)))/x^4,x)`output `(15*atanh(sqrt(x)*c)*b*c**6*x**3 - 15*atanh(sqrt(x)*c)*b - 15*sqrt(x)*b*c**5*x**2 - 5*sqrt(x)*b*c**3*x - 3*sqrt(x)*b*c - 15*a)/(45*x**3)`

3.194 $\int x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx$

Optimal result	1538
Mathematica [A] (verified)	1539
Rubi [A] (warning: unable to verify)	1539
Maple [A] (verified)	1544
Fricas [A] (verification not implemented)	1545
Sympy [F]	1546
Maxima [A] (verification not implemented)	1546
Giac [F]	1547
Mupad [B] (verification not implemented)	1548
Reduce [B] (verification not implemented)	1549

Optimal result

Integrand size = 18, antiderivative size = 211

$$\int x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \frac{ab\sqrt{x}}{2c^7} + \frac{71b^2x}{420c^6} + \frac{3b^2x^2}{70c^4} + \frac{b^2x^3}{84c^2} + \frac{b^2\sqrt{x}\operatorname{arctanh}(c\sqrt{x})}{2c^7} + \frac{bx^{3/2}(a + b \operatorname{arctanh}(c\sqrt{x}))}{6c^5} + \frac{bx^{5/2}(a + b \operatorname{arctanh}(c\sqrt{x}))}{10c^3} + \frac{bx^{7/2}(a + b \operatorname{arctanh}(c\sqrt{x}))}{14c} - \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{4c^8} + \frac{1}{4}x^4(a + b \operatorname{arctanh}(c\sqrt{x}))^2 + \frac{44b^2 \log(1 - c^2x)}{105c^8}$$

output

```
1/2*a*b*x^(1/2)/c^7+71/420*b^2*x/c^6+3/70*b^2*x^2/c^4+1/84*b^2*x^3/c^2+1/2
*b^2*x^(1/2)*arctanh(c*x^(1/2))/c^7+1/6*b*x^(3/2)*(a+b*arctanh(c*x^(1/2)))
/c^5+1/10*b*x^(5/2)*(a+b*arctanh(c*x^(1/2)))/c^3+1/14*b*x^(7/2)*(a+b*arcta
nh(c*x^(1/2)))/c-1/4*(a+b*arctanh(c*x^(1/2)))^2/c^8+1/4*x^4*(a+b*arctanh(c
*x^(1/2)))^2+44/105*b^2*ln(-c^2*x+1)/c^8
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.06

$$\int x^3 (a + \operatorname{arctanh}(c\sqrt{x}))^2 dx$$

$$= \frac{210abc\sqrt{x} + 71b^2c^2x + 70abc^3x^{3/2} + 18b^2c^4x^2 + 42abc^5x^{5/2} + 5b^2c^6x^3 + 30abc^7x^{7/2} + 105a^2c^8x^4 + 2bc\sqrt{x}}$$

input

```
Integrate[x^3*(a + b*ArcTanh[c*Sqrt[x]])^2,x]
```

output

```
(210*a*b*c*Sqrt[x] + 71*b^2*c^2*x + 70*a*b*c^3*x^(3/2) + 18*b^2*c^4*x^2 +
42*a*b*c^5*x^(5/2) + 5*b^2*c^6*x^3 + 30*a*b*c^7*x^(7/2) + 105*a^2*c^8*x^4
+ 2*b*c*Sqrt[x]*(105*a*c^7*x^(7/2) + b*(105 + 35*c^2*x + 21*c^4*x^2 + 15*c
^6*x^3))*ArcTanh[c*Sqrt[x]] + 105*b^2*(-1 + c^8*x^4)*ArcTanh[c*Sqrt[x]]^2
+ b*(105*a + 176*b)*Log[1 - c*Sqrt[x]] - 105*a*b*Log[1 + c*Sqrt[x]] + 176*
b^2*Log[1 + c*Sqrt[x]])/(420*c^8)
```

Rubi [A] (warning: unable to verify)

Time = 2.04 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.45, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 1.111$, Rules used = {6454, 6452, 6542, 6452, 243, 49, 2009, 6542, 6452, 243, 49, 2009, 6542, 6452, 243, 49, 2009, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 (a + \operatorname{arctanh}(c\sqrt{x}))^2 dx$$

$$\downarrow \text{6454}$$

$$2 \int x^{7/2} (a + \operatorname{arctanh}(c\sqrt{x}))^2 d\sqrt{x}$$

$$\downarrow \text{6452}$$

$$2 \left(\frac{1}{8} x^4 (a + \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \int \frac{x^4 (a + \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x} \right)$$

↓ 6542

$$2 \left(\frac{1}{8} x^4 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\int x^3 (a + \operatorname{barctanh}(c\sqrt{x})) d\sqrt{x}}{c^2} \right) \right)$$

↓ 6452

$$2 \left(\frac{1}{8} x^4 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{7} x^{7/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{7} bc \int \frac{x^{7/2}}{1 - c^2 x} d\sqrt{x}}{c^2} \right) \right)$$

↓ 243

$$2 \left(\frac{1}{8} x^4 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{7} x^{7/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{14} bc \int \frac{x^{3/2}}{1 - c^2 x} d\sqrt{x}}{c^2} \right) \right)$$

↓ 49

$$2 \left(\frac{1}{8} x^4 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{7} x^{7/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{14} bc \int \left(-\frac{x}{c^2} - \frac{1}{c^2} \right) d\sqrt{x}}{c^2} \right) \right)$$

↓ 2009

$$2 \left(\frac{1}{8} x^4 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{7} x^{7/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{14} bc \left(-\frac{x}{c^6} - \frac{1}{c^2} \right)}{c^2} \right) \right)$$

↓ 6542

$$2 \left(\frac{1}{8} x^4 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\int x^2 (a + \operatorname{barctanh}(c\sqrt{x})) d\sqrt{x}}{c^2} - \frac{\frac{1}{7} x^{7/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{14} bc \left(-\frac{x}{c^6} - \frac{1}{c^2} \right)}{c^2} \right) \right)$$

↓ 6452

$$2 \left(\frac{1}{8} x^4 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{5} bc \int \frac{x^{5/2}}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{1}{7} x^7 \right) \right)$$

↓ 243

$$2 \left(\frac{1}{8} x^4 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{10} bc \int \frac{x}{1 - c^2 x} dx}{c^2} - \frac{1}{7} x^7 \right) \right)$$

↓ 49

$$2 \left(\frac{1}{8} x^4 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{10} bc \int \left(-\frac{x}{c^2} - \frac{1}{c^4 (c^2 x - 1)} \right)}{c^2} \right) \right)$$

↓ 2009

$$2 \left(\frac{1}{8} x^4 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{10} bc \left(-\frac{x}{c^4} - \frac{x}{2c^2} - \frac{\log(1 - c^2 x)}{c^6} \right)}{c^2} \right) \right)$$

↓ 6542

$$2 \left(\frac{1}{8} x^4 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\frac{\int \frac{x (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\int x (a + \operatorname{barctanh}(c\sqrt{x})) d\sqrt{x}}{c^2}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + \operatorname{barctanh}(c\sqrt{x}))}{c^2} \right) \right)$$

↓ 6452

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{x(a+b \operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x} - \frac{\frac{1}{3} x^{3/2} (a+b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6} bc \int \frac{x^{3/2}}{1-c^2x} d\sqrt{x}}{c^2}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a+b \operatorname{arctanh}(c\sqrt{x}))}{c^2}}{c^2} \right) \right)$$

↓ 243

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{x(a+b \operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x} - \frac{\frac{1}{3} x^{3/2} (a+b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6} bc \int \frac{x}{1-c^2x} dx}{c^2}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a+b \operatorname{arctanh}(c\sqrt{x}))}{c^2}}{c^2} \right) \right)$$

↓ 49

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{x(a+b \operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x} - \frac{\frac{1}{3} x^{3/2} (a+b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6} bc \int \left(-\frac{1}{c^2} - \frac{1}{c^2(c^2x-1)} \right) dx}{c^2}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a+b \operatorname{arctanh}(c\sqrt{x}))}{c^2}}{c^2} \right) \right)$$

↓ 2009

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{x(a+b \operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x} - \frac{\frac{1}{3} x^{3/2} (a+b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6} bc \left(-\frac{x}{c^2} - \frac{\log(1-c^2x)}{c^4} \right)}{c^2}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a+b \operatorname{arctanh}(c\sqrt{x}))}{c^2}}{c^2} \right) \right)$$

↓ 6542

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{a+b \operatorname{arctanh}(c\sqrt{x})}{1-c^2x} d\sqrt{x} - \frac{\int (a+b \operatorname{arctanh}(c\sqrt{x})) d\sqrt{x}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a+b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6} bc \left(-\frac{x}{c^2} - \frac{\log(1-c^2x)}{c^4} \right)}{c^2}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a+b \operatorname{arctanh}(c\sqrt{x}))}{c^2}}{c^2} \right) \right)$$

↓ 2009

$$2 \left(\frac{1}{8} x^4 (a + \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{1 - c^2 x} d\sqrt{x} - \frac{a\sqrt{x} + b\sqrt{x} \operatorname{arctanh}(c\sqrt{x}) + \frac{b \log(1 - c^2 x)}{2c}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x}))}{c^2}}{c^2} \right) \right)$$

↓ 6510

$$2 \left(\frac{1}{8} x^4 (a + \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{4} bc \left(\frac{\frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{2bc^3} - \frac{a\sqrt{x} + b\sqrt{x} \operatorname{arctanh}(c\sqrt{x}) + \frac{b \log(1 - c^2 x)}{2c}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x}))}{c^2}}{c^2} \right) \right)$$

input `Int[x^3*(a + b*ArcTanh[c*Sqrt[x]])^2,x]`

output `2*((x^4*(a + b*ArcTanh[c*Sqrt[x]])^2)/8 - (b*c*(-((x^(7/2)*(a + b*ArcTanh[c*Sqrt[x]]))/7 - (b*c*(-(x/c^6) - x/(2*c^4) - x^(3/2)/(3*c^2) - Log[1 - c^2*x]/c^8))/14)/c^2) + (-((x^(5/2)*(a + b*ArcTanh[c*Sqrt[x]]))/5 - (b*c*(-(x/c^4) - x/(2*c^2) - Log[1 - c^2*x]/c^6))/10)/c^2) + (-((x^(3/2)*(a + b*ArcTanh[c*Sqrt[x]]))/3 - (b*c*(-(x/c^2) - Log[1 - c^2*x]/c^4))/6)/c^2) + ((a + b*ArcTanh[c*Sqrt[x]])^2/(2*b*c^3) - (a*Sqrt[x] + b*Sqrt[x]*ArcTanh[c*Sqrt[x] + (b*Log[1 - c^2*x])/(2*c))/c^2)/c^2)/4)`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6510 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6542 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((f_.)*(x_)^(m_.))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

Maple [A] (verified)

Time = 1.03 (sec) , antiderivative size = 309, normalized size of antiderivative = 1.46

method	result
parts	$\frac{a^2 x^4}{4} + \frac{2b^2 \left(\frac{c^8 x^4 \operatorname{arctanh}(c\sqrt{x})^2}{8} + \frac{\operatorname{arctanh}(c\sqrt{x})c^7 x^{\frac{7}{2}}}{28} + \frac{\operatorname{arctanh}(c\sqrt{x})c^5 x^{\frac{5}{2}}}{20} + \frac{\operatorname{arctanh}(c\sqrt{x})c^3 x^{\frac{3}{2}}}{12} + \frac{\operatorname{arctanh}(c\sqrt{x})c\sqrt{x}}{4} + \operatorname{arctanh}(c\sqrt{x}) \right)}{4}$
derivativedivides	$\frac{a^2 c^8 x^4}{4} + 2b^2 \left(\frac{c^8 x^4 \operatorname{arctanh}(c\sqrt{x})^2}{8} + \frac{\operatorname{arctanh}(c\sqrt{x})c^7 x^{\frac{7}{2}}}{28} + \frac{\operatorname{arctanh}(c\sqrt{x})c^5 x^{\frac{5}{2}}}{20} + \frac{\operatorname{arctanh}(c\sqrt{x})c^3 x^{\frac{3}{2}}}{12} + \frac{\operatorname{arctanh}(c\sqrt{x})c\sqrt{x}}{4} + \operatorname{arctanh}(c\sqrt{x}) \right)$
default	$\frac{a^2 c^8 x^4}{4} + 2b^2 \left(\frac{c^8 x^4 \operatorname{arctanh}(c\sqrt{x})^2}{8} + \frac{\operatorname{arctanh}(c\sqrt{x})c^7 x^{\frac{7}{2}}}{28} + \frac{\operatorname{arctanh}(c\sqrt{x})c^5 x^{\frac{5}{2}}}{20} + \frac{\operatorname{arctanh}(c\sqrt{x})c^3 x^{\frac{3}{2}}}{12} + \frac{\operatorname{arctanh}(c\sqrt{x})c\sqrt{x}}{4} + \operatorname{arctanh}(c\sqrt{x}) \right)$

input `int(x^3*(a+b*arctanh(c*x^(1/2)))^2,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 1/4*a^2*x^4+2*b^2/c^8*(1/8*c^8*x^4*arctanh(c*x^(1/2))^2+1/28*arctanh(c*x^(1/2))*c^7*x^(7/2)+1/20*arctanh(c*x^(1/2))*c^5*x^(5/2)+1/12*arctanh(c*x^(1/2))*c^3*x^(3/2)+1/4*arctanh(c*x^(1/2))*c*x^(1/2)+1/8*arctanh(c*x^(1/2))*\ln(c*x^(1/2)-1)-1/8*arctanh(c*x^(1/2))*\ln(1+c*x^(1/2))+1/32*\ln(c*x^(1/2)-1)^2-1/16*\ln(c*x^(1/2)-1)*\ln(1/2*c*x^(1/2)+1/2)-1/16*(\ln(1+c*x^(1/2))-\ln(1/2*c*x^(1/2)+1/2))*\ln(-1/2*c*x^(1/2)+1/2)+1/32*\ln(1+c*x^(1/2))^2+1/168*c^6*x^3+3/140*c^4*x^2+71/840*c^2*x+22/105*\ln(c*x^(1/2)-1)+22/105*\ln(1+c*x^(1/2))+4*a*b/c^8*(1/8*c^8*x^4*arctanh(c*x^(1/2))+1/56*c^7*x^(7/2)+1/40*c^5*x^(5/2)+1/24*c^3*x^(3/2)+1/8*c*x^(1/2)+1/16*\ln(c*x^(1/2)-1)-1/16*\ln(1+c*x^(1/2))) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.29

$$\int x^3(a + b\operatorname{arctanh}(c\sqrt{x}))^2 dx = \frac{420 a^2 c^8 x^4 + 20 b^2 c^6 x^3 + 72 b^2 c^4 x^2 + 284 b^2 c^2 x + 105 (b^2 c^8 x^4 - b^2) \log\left(-\frac{c^2 x + 2c\sqrt{x} + 1}{c^2 x - 1}\right)^2 + 4(105 abc^8 - 105 abc^6 x + 105 abc^4 x^2 - 105 abc^2 x^3 + 105 abc x^4)}{4}$$

input `integrate(x^3*(a+b*arctanh(c*x^(1/2)))^2,x, algorithm="fricas")`

output

```
1/1680*(420*a^2*c^8*x^4 + 20*b^2*c^6*x^3 + 72*b^2*c^4*x^2 + 284*b^2*c^2*x
+ 105*(b^2*c^8*x^4 - b^2)*log(-(c^2*x + 2*c*sqrt(x) + 1)/(c^2*x - 1))^2 +
4*(105*a*b*c^8 - 105*a*b + 176*b^2)*log(c*sqrt(x) + 1) - 4*(105*a*b*c^8 -
105*a*b - 176*b^2)*log(c*sqrt(x) - 1) + 4*(105*a*b*c^8*x^4 - 105*a*b*c^8 +
(15*b^2*c^7*x^3 + 21*b^2*c^5*x^2 + 35*b^2*c^3*x + 105*b^2*c)*sqrt(x))*log
(-(c^2*x + 2*c*sqrt(x) + 1)/(c^2*x - 1)) + 8*(15*a*b*c^7*x^3 + 21*a*b*c^5*
x^2 + 35*a*b*c^3*x + 105*a*b*c)*sqrt(x))/c^8
```

Sympy [F]

$$\int x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \int x^3 (a + b \operatorname{atanh}(c\sqrt{x}))^2 dx$$

input

```
integrate(x**3*(a+b*atanh(c*x**(1/2)))**2,x)
```

output

```
Integral(x**3*(a + b*atanh(c*sqrt(x)))**2, x)
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 265, normalized size of antiderivative = 1.26

$$\int x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \frac{1}{4} b^2 x^4 \operatorname{artanh}(c\sqrt{x})^2 + \frac{1}{4} a^2 x^4$$

$$+ \frac{1}{420} \left(210 x^4 \operatorname{artanh}(c\sqrt{x}) + c \left(\frac{2 \left(15 c^6 x^{\frac{7}{2}} + 21 c^4 x^{\frac{5}{2}} + 35 c^2 x^{\frac{3}{2}} + 105 \sqrt{x} \right)}{c^8} - \frac{105 \log(c\sqrt{x} + 1)}{c^9} + \frac{105 \log(c\sqrt{x} - 1)}{c^9} \right) \right)$$

$$+ \frac{1}{1680} \left(4 c \left(\frac{2 \left(15 c^6 x^{\frac{7}{2}} + 21 c^4 x^{\frac{5}{2}} + 35 c^2 x^{\frac{3}{2}} + 105 \sqrt{x} \right)}{c^8} - \frac{105 \log(c\sqrt{x} + 1)}{c^9} + \frac{105 \log(c\sqrt{x} - 1)}{c^9} \right) \right)$$

input

```
integrate(x^3*(a+b*arctanh(c*x^(1/2)))^2,x, algorithm="maxima")
```

output

```
1/4*b^2*x^4*arctanh(c*sqrt(x))^2 + 1/4*a^2*x^4 + 1/420*(210*x^4*arctanh(c*
sqrt(x)) + c*(2*(15*c^6*x^(7/2) + 21*c^4*x^(5/2) + 35*c^2*x^(3/2) + 105*sq
rt(x))/c^8 - 105*log(c*sqrt(x) + 1)/c^9 + 105*log(c*sqrt(x) - 1)/c^9))*a*b
+ 1/1680*(4*c*(2*(15*c^6*x^(7/2) + 21*c^4*x^(5/2) + 35*c^2*x^(3/2) + 105*
sqrt(x))/c^8 - 105*log(c*sqrt(x) + 1)/c^9 + 105*log(c*sqrt(x) - 1)/c^9)*ar
ctanh(c*sqrt(x)) + (20*c^6*x^3 + 72*c^4*x^2 + 284*c^2*x - 2*(105*log(c*sq
rt(x) - 1) - 352)*log(c*sqrt(x) + 1) + 105*log(c*sqrt(x) + 1)^2 + 105*log(c
*sqrt(x) - 1)^2 + 704*log(c*sqrt(x) - 1))/c^8)*b^2
```

Giac [F]

$$\int x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \int (b \operatorname{arctanh}(c\sqrt{x}) + a)^2 x^3 dx$$

input

```
integrate(x^3*(a+b*arctanh(c*x^(1/2)))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*sqrt(x)) + a)^2*x^3, x)
```


Mupad [B] (verification not implemented)

Time = 6.49 (sec) , antiderivative size = 453, normalized size of antiderivative = 2.15

$$\begin{aligned}
\int x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = & \frac{a^2 x^4}{4} + \frac{44 b^2 \ln(c\sqrt{x} - 1)}{105 c^8} + \frac{44 b^2 \ln(c\sqrt{x} + 1)}{105 c^8} \\
& + \frac{71 b^2 x}{420 c^6} - \frac{b^2 \ln(c\sqrt{x} + 1)^2}{16 c^8} \\
& - \frac{b^2 \ln(1 - c\sqrt{x})^2}{16 c^8} + \frac{b^2 x^3}{84 c^2} + \frac{3 b^2 x^2}{70 c^4} \\
& + \frac{b^2 x^4 \ln(c\sqrt{x} + 1)^2}{16} + \frac{b^2 x^4 \ln(1 - c\sqrt{x})^2}{16} \\
& + \frac{b^2 x^{7/2} \ln(c\sqrt{x} + 1)}{28 c} + \frac{b^2 x^{5/2} \ln(c\sqrt{x} + 1)}{20 c^3} \\
& + \frac{b^2 x^{3/2} \ln(c\sqrt{x} + 1)}{12 c^5} + \frac{b^2 \sqrt{x} \ln(c\sqrt{x} + 1)}{4 c^7} \\
& - \frac{b^2 x^{7/2} \ln(1 - c\sqrt{x})}{28 c} - \frac{b^2 x^{5/2} \ln(1 - c\sqrt{x})}{20 c^3} \\
& - \frac{b^2 x^{3/2} \ln(1 - c\sqrt{x})}{12 c^5} - \frac{b^2 \sqrt{x} \ln(1 - c\sqrt{x})}{4 c^7} \\
& + \frac{a b \ln(c\sqrt{x} - 1)}{4 c^8} - \frac{a b \ln(c\sqrt{x} + 1)}{4 c^8} \\
& + \frac{a b x^4 \ln(c\sqrt{x} + 1)}{4} - \frac{a b x^4 \ln(1 - c\sqrt{x})}{4} \\
& + \frac{b^2 \ln(c\sqrt{x} + 1) \ln(1 - c\sqrt{x})}{8 c^8} \\
& + \frac{a b x^{7/2}}{14 c} + \frac{a b x^{5/2}}{10 c^3} + \frac{a b x^{3/2}}{6 c^5} + \frac{a b \sqrt{x}}{2 c^7} \\
& - \frac{b^2 x^4 \ln(c\sqrt{x} + 1) \ln(1 - c\sqrt{x})}{8}
\end{aligned}$$

input `int(x^3*(a + b*atanh(c*x^(1/2)))^2,x)`

output

```
(a^2*x^4)/4 + (44*b^2*log(c*x^(1/2) - 1))/(105*c^8) + (44*b^2*log(c*x^(1/2)
) + 1))/(105*c^8) + (71*b^2*x)/(420*c^6) - (b^2*log(c*x^(1/2) + 1)^2)/(16*
c^8) - (b^2*log(1 - c*x^(1/2))^2)/(16*c^8) + (b^2*x^3)/(84*c^2) + (3*b^2*x
^2)/(70*c^4) + (b^2*x^4*log(c*x^(1/2) + 1)^2)/16 + (b^2*x^4*log(1 - c*x^(1
/2))^2)/16 + (b^2*x^(7/2)*log(c*x^(1/2) + 1))/(28*c) + (b^2*x^(5/2)*log(c*
x^(1/2) + 1))/(20*c^3) + (b^2*x^(3/2)*log(c*x^(1/2) + 1))/(12*c^5) + (b^2*
x^(1/2)*log(c*x^(1/2) + 1))/(4*c^7) - (b^2*x^(7/2)*log(1 - c*x^(1/2)))/(28
*c) - (b^2*x^(5/2)*log(1 - c*x^(1/2)))/(20*c^3) - (b^2*x^(3/2)*log(1 - c*x
^(1/2)))/(12*c^5) - (b^2*x^(1/2)*log(1 - c*x^(1/2)))/(4*c^7) + (a*b*log(c*
x^(1/2) - 1))/(4*c^8) - (a*b*log(c*x^(1/2) + 1))/(4*c^8) + (a*b*x^4*log(c*
x^(1/2) + 1))/4 - (a*b*x^4*log(1 - c*x^(1/2)))/4 + (b^2*log(c*x^(1/2) + 1)
*log(1 - c*x^(1/2)))/(8*c^8) + (a*b*x^(7/2))/(14*c) + (a*b*x^(5/2))/(10*c
^3) + (a*b*x^(3/2))/(6*c^5) + (a*b*x^(1/2))/(2*c^7) - (b^2*x^4*log(c*x^(1/2)
) + 1)*log(1 - c*x^(1/2))/8
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.09

$$\int x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx$$

$$= \frac{105 a \operatorname{atanh}(\sqrt{x} c)^2 b^2 c^8 x^4 - 105 a \operatorname{atanh}(\sqrt{x} c)^2 b^2 + 30 \sqrt{x} a \operatorname{atanh}(\sqrt{x} c) b^2 c^7 x^3 + 42 \sqrt{x} a \operatorname{atanh}(\sqrt{x} c) b^2 c^5 x^2}{1}$$

input

```
int(x^3*(a+b*atanh(c*x^(1/2)))^2,x)
```

output

```
(105*atanh(sqrt(x)*c)**2*b**2*c**8*x**4 - 105*atanh(sqrt(x)*c)**2*b**2 + 3
0*sqrt(x)*atanh(sqrt(x)*c)*b**2*c**7*x**3 + 42*sqrt(x)*atanh(sqrt(x)*c)*b*
**2*c**5*x**2 + 70*sqrt(x)*atanh(sqrt(x)*c)*b**2*c**3*x + 210*sqrt(x)*atanh
(sqrt(x)*c)*b**2*c + 210*atanh(sqrt(x)*c)*a*b*c**8*x**4 - 210*atanh(sqrt(x)
)*c)*a*b + 352*atanh(sqrt(x)*c)*b**2 + 30*sqrt(x)*a*b*c**7*x**3 + 42*sqrt(
x)*a*b*c**5*x**2 + 70*sqrt(x)*a*b*c**3*x + 210*sqrt(x)*a*b*c + 352*log(sqrt
(x)*c - 1)*b**2 + 105*a**2*c**8*x**4 + 5*b**2*c**6*x**3 + 18*b**2*c**4*x*
**2 + 71*b**2*c**2*x)/(420*c**8)
```

3.195 $\int x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx$

Optimal result	1550
Mathematica [A] (verified)	1551
Rubi [A] (warning: unable to verify)	1551
Maple [B] (verified)	1555
Fricas [A] (verification not implemented)	1556
Sympy [F]	1556
Maxima [A] (verification not implemented)	1557
Giac [F]	1557
Mupad [B] (verification not implemented)	1558
Reduce [B] (verification not implemented)	1558

Optimal result

Integrand size = 18, antiderivative size = 173

$$\int x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \frac{2ab\sqrt{x}}{3c^5} + \frac{8b^2x}{45c^4} + \frac{b^2x^2}{30c^2} + \frac{2b^2\sqrt{x}\operatorname{arctanh}(c\sqrt{x})}{3c^5}$$

$$+ \frac{2bx^{3/2}(a + b \operatorname{arctanh}(c\sqrt{x}))}{9c^3}$$

$$+ \frac{2bx^{5/2}(a + b \operatorname{arctanh}(c\sqrt{x}))}{15c}$$

$$- \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{3c^6}$$

$$+ \frac{1}{3}x^3(a + b \operatorname{arctanh}(c\sqrt{x}))^2 + \frac{23b^2 \log(1 - c^2x)}{45c^6}$$

output

```
2/3*a*b*x^(1/2)/c^5+8/45*b^2*x/c^4+1/30*b^2*x^2/c^2+2/3*b^2*x^(1/2)*arctan
h(c*x^(1/2))/c^5+2/9*b*x^(3/2)*(a+b*arctanh(c*x^(1/2)))/c^3+2/15*b*x^(5/2)
*(a+b*arctanh(c*x^(1/2)))/c-1/3*(a+b*arctanh(c*x^(1/2)))^2/c^6+1/3*x^3*(a+
b*arctanh(c*x^(1/2)))^2+23/45*b^2*ln(-c^2*x+1)/c^6
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.12

$$\int x^2(a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx$$

$$= \frac{60abc\sqrt{x} + 16b^2c^2x + 20abc^3x^{3/2} + 3b^2c^4x^2 + 12abc^5x^{5/2} + 30a^2c^6x^3 + 4bc\sqrt{x}(15ac^5x^{5/2} + b(15 + 5c^2x$$

input

```
Integrate[x^2*(a + b*ArcTanh[c*Sqrt[x]])^2,x]
```

output

```
(60*a*b*c*Sqrt[x] + 16*b^2*c^2*x + 20*a*b*c^3*x^(3/2) + 3*b^2*c^4*x^2 + 12
*a*b*c^5*x^(5/2) + 30*a^2*c^6*x^3 + 4*b*c*Sqrt[x]*(15*a*c^5*x^(5/2) + b*(1
5 + 5*c^2*x + 3*c^4*x^2))*ArcTanh[c*Sqrt[x]] + 30*b^2*(-1 + c^6*x^3)*ArcTa
nh[c*Sqrt[x]]^2 + 2*b*(15*a + 23*b)*Log[1 - c*Sqrt[x]] - 30*a*b*Log[1 + c*
Sqrt[x]] + 46*b^2*Log[1 + c*Sqrt[x]])/(90*c^6)
```

Rubi [A] (warning: unable to verify)

Time = 2.35 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.31, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {6454, 6452, 6542, 6452, 243, 49, 2009, 6542, 6452, 243, 49, 2009, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2(a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx$$

$$\downarrow 6454$$

$$2 \int x^{5/2}(a + b \operatorname{arctanh}(c\sqrt{x}))^2 d\sqrt{x}$$

$$\downarrow 6452$$

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{3} bc \int \frac{x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x} \right)$$

↓ 6542

$$2 \left(\frac{1}{6} x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{3} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\int x^2 (a + \operatorname{barctanh}(c\sqrt{x})) d\sqrt{x}}{c^2} \right) \right)$$

↓ 6452

$$2 \left(\frac{1}{6} x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{3} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{5} bc \int \frac{x^{5/2}}{1 - c^2 x} d\sqrt{x}}{c^2} \right) \right)$$

↓ 243

$$2 \left(\frac{1}{6} x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{3} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{10} bc \int \frac{x}{1 - c^2 x} d\sqrt{x}}{c^2} \right) \right)$$

↓ 49

$$2 \left(\frac{1}{6} x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{3} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{10} bc \int \left(-\frac{x}{c^2} \right)}{c^2} \right) \right)$$

↓ 2009

$$2 \left(\frac{1}{6} x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{3} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{10} bc \left(-\frac{x}{c^4} - \right)}{c^2} \right) \right)$$

↓ 6542

$$2 \left(\frac{1}{6} x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{3} bc \left(\frac{\int \frac{x (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\int x (a + \operatorname{barctanh}(c\sqrt{x})) d\sqrt{x}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + \operatorname{barctanh}(c\sqrt{x}))}{c^2} \right) \right)$$

↓ 6452

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{3} bc \left(\frac{\int \frac{x^{a+b \operatorname{arctanh}(c\sqrt{x})}}{1-c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a+b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{3} bc \int \frac{x^{3/2}}{1-c^2 x} d\sqrt{x}}{c^2} - \frac{1}{5} x^{5/2} \right) \right)$$

↓ 243

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{3} bc \left(\frac{\int \frac{x^{a+b \operatorname{arctanh}(c\sqrt{x})}}{1-c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a+b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6} bc \int \frac{x}{1-c^2 x} dx}{c^2} - \frac{1}{5} x^{5/2} \right) \right)$$

↓ 49

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{3} bc \left(\frac{\int \frac{x^{a+b \operatorname{arctanh}(c\sqrt{x})}}{1-c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a+b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6} bc \int \left(-\frac{1}{c^2} - \frac{1}{c^2(c^2 x - 1)} \right) dx}{c^2} \right) \right)$$

↓ 2009

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{3} bc \left(\frac{\int \frac{x^{a+b \operatorname{arctanh}(c\sqrt{x})}}{1-c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a+b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6} bc \left(-\frac{x}{c^2} - \frac{\log(1-c^2 x)}{c^4} \right)}{c^2} \right) \right)$$

↓ 6542

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{3} bc \left(\frac{\frac{\int \frac{a+b \operatorname{arctanh}(c\sqrt{x})}{1-c^2 x} d\sqrt{x}}{c^2} - \frac{\int (a+b \operatorname{arctanh}(c\sqrt{x})) d\sqrt{x}}{c^2}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a+b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6} bc}{c^2} \right) \right)$$

↓ 2009

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{3} bc \left(\frac{\frac{\frac{\int \frac{a+b \operatorname{arctanh}(c\sqrt{x})}{1-c^2 x} d\sqrt{x}}{c^2} - \frac{a\sqrt{x} + b\sqrt{x} \operatorname{arctanh}(c\sqrt{x}) + \frac{b \log(1-c^2 x)}{2c}}{c^2}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a+b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{6} bc}{c^2} \right) \right)$$

↓ 6510

$$2 \left(\frac{1}{6} x^3 (a + \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{3} bc \left(\frac{\frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{2bc^3} - \frac{a\sqrt{x} + b\sqrt{x} \operatorname{arctanh}(c\sqrt{x}) + \frac{b \log(1-c^2x)}{2c}}{c^2}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} \right) \right)$$

input `Int[x^2*(a + b*ArcTanh[c*Sqrt[x]])^2,x]`

output `2*((x^3*(a + b*ArcTanh[c*Sqrt[x]])^2)/6 - (b*c*(-((x^(5/2)*(a + b*ArcTanh[c*Sqrt[x]]))/5 - (b*c*(-(x/c^4) - x/(2*c^2) - Log[1 - c^2*x]/c^6))/10)/c^2) + (-((x^(3/2)*(a + b*ArcTanh[c*Sqrt[x]]))/3 - (b*c*(-(x/c^2) - Log[1 - c^2*x]/c^4))/6)/c^2) + ((a + b*ArcTanh[c*Sqrt[x]])^2/(2*b*c^3) - (a*Sqrt[x] + b*Sqrt[x]*ArcTanh[c*Sqrt[x]] + (b*Log[1 - c^2*x])/(2*c))/c^2)/c^2)/3)`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

```
rule 6454 Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
  Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x
], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simpl
ify[(m + 1)/n]]
```

```
rule 6510 Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_.) + (e_.)*(x_)^2), x_Symb
ol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b
, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]
```

```
rule 6542 Int((((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_)))/((d_.) + (
e_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*
x])^p, x], x] - Simp[d*(f^2/e Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/
(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m,
1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 278 vs. 2(137) = 274.

Time = 1.01 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.61

method	result
parts	$\frac{a^2 x^3}{3} + \frac{2b^2 \left(\frac{c^6 x^3 \operatorname{arctanh}(c\sqrt{x})^2}{6} + \frac{\operatorname{arctanh}(c\sqrt{x})c^5 x^{\frac{5}{2}}}{15} + \frac{\operatorname{arctanh}(c\sqrt{x})c^3 x^{\frac{3}{2}}}{9} + \frac{\operatorname{arctanh}(c\sqrt{x})c\sqrt{x}}{3} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{6} \right)}{1}$
derivativedivides	$\frac{a^2 c^6 x^3}{3} + 2b^2 \left(\frac{c^6 x^3 \operatorname{arctanh}(c\sqrt{x})^2}{6} + \frac{\operatorname{arctanh}(c\sqrt{x})c^5 x^{\frac{5}{2}}}{15} + \frac{\operatorname{arctanh}(c\sqrt{x})c^3 x^{\frac{3}{2}}}{9} + \frac{\operatorname{arctanh}(c\sqrt{x})c\sqrt{x}}{3} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{6} \right)$
default	$\frac{a^2 c^6 x^3}{3} + 2b^2 \left(\frac{c^6 x^3 \operatorname{arctanh}(c\sqrt{x})^2}{6} + \frac{\operatorname{arctanh}(c\sqrt{x})c^5 x^{\frac{5}{2}}}{15} + \frac{\operatorname{arctanh}(c\sqrt{x})c^3 x^{\frac{3}{2}}}{9} + \frac{\operatorname{arctanh}(c\sqrt{x})c\sqrt{x}}{3} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{6} \right)$

```
input int(x^2*(a+b*arctanh(c*x^(1/2)))^2,x,method=_RETURNVERBOSE)
```


output

```
1/3*a^2*x^3+2*b^2/c^6*(1/6*c^6*x^3*arctanh(c*x^(1/2))^2+1/15*arctanh(c*x^(1/2))*c^5*x^(5/2)+1/9*arctanh(c*x^(1/2))*c^3*x^(3/2)+1/3*arctanh(c*x^(1/2))*c*x^(1/2)+1/6*arctanh(c*x^(1/2))*ln(c*x^(1/2)-1)-1/6*arctanh(c*x^(1/2))*ln(1+c*x^(1/2))+1/24*ln(c*x^(1/2)-1)^2-1/12*ln(c*x^(1/2)-1)*ln(1/2*c*x^(1/2)+1/2)+1/24*ln(1+c*x^(1/2))^2-1/12*(ln(1+c*x^(1/2))-ln(1/2*c*x^(1/2)+1/2))*ln(-1/2*c*x^(1/2)+1/2)+1/60*c^4*x^2+4/45*c^2*x+23/90*ln(c*x^(1/2)-1)+23/90*ln(1+c*x^(1/2)))+4*a*b/c^6*(1/6*c^6*x^3*arctanh(c*x^(1/2))+1/30*c^5*x^(5/2)+1/18*c^3*x^(3/2)+1/6*c*x^(1/2)+1/12*ln(c*x^(1/2)-1)-1/12*ln(1+c*x^(1/2)))
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.39

$$\int x^2(a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx$$

$$= \frac{60 a^2 c^6 x^3 + 6 b^2 c^4 x^2 + 32 b^2 c^2 x + 15 (b^2 c^6 x^3 - b^2) \log\left(-\frac{c^2 x + 2 c \sqrt{x} + 1}{c^2 x - 1}\right)^2 + 4 (15 a b c^6 - 15 a b + 23 b^2) \log\left(-\frac{c^2 x + 2 c \sqrt{x} + 1}{c^2 x - 1}\right) + 8 (3 a^2 b c^5 x^2 + 5 a b^2 c^3 x + 15 a b^2 c) \sqrt{x}}{c^6}$$

input

```
integrate(x^2*(a+b*arctanh(c*x^(1/2)))^2,x, algorithm="fricas")
```

output

```
1/180*(60*a^2*c^6*x^3 + 6*b^2*c^4*x^2 + 32*b^2*c^2*x + 15*(b^2*c^6*x^3 - b^2)*log(-(c^2*x + 2*c*sqrt(x) + 1)/(c^2*x - 1))^2 + 4*(15*a*b*c^6 - 15*a*b + 23*b^2)*log(c*sqrt(x) + 1) - 4*(15*a*b*c^6 - 15*a*b - 23*b^2)*log(c*sqrt(x) - 1) + 4*(15*a*b*c^6*x^3 - 15*a*b*c^6 + (3*b^2*c^5*x^2 + 5*b^2*c^3*x + 15*b^2*c)*sqrt(x))*log(-(c^2*x + 2*c*sqrt(x) + 1)/(c^2*x - 1)) + 8*(3*a*b*c^5*x^2 + 5*a*b*c^3*x + 15*a*b*c)*sqrt(x))/c^6
```

Sympy [F]

$$\int x^2(a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \int x^2(a + b \operatorname{atanh}(c\sqrt{x}))^2 dx$$

input

```
integrate(x**2*(a+b*atanh(c*x**(1/2)))**2,x)
```

output `Integral(x**2*(a + b*atanh(c*sqrt(x)))**2, x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.39

$$\int x^2(a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \frac{1}{3} b^2 x^3 \operatorname{arctanh}(c\sqrt{x})^2 + \frac{1}{3} a^2 x^3 + \frac{1}{45} \left(30 x^3 \operatorname{arctanh}(c\sqrt{x}) + c \left(\frac{2(3c^4 x^{\frac{5}{2}} + 5c^2 x^{\frac{3}{2}} + 15\sqrt{x})}{c^6} - \frac{15 \log(c\sqrt{x} + 1)}{c^7} + \frac{15 \log(c\sqrt{x} - 1)}{c^7} \right) \right) \operatorname{arctanh}(c\sqrt{x}) + \frac{1}{180} \left(4c \left(\frac{2(3c^4 x^{\frac{5}{2}} + 5c^2 x^{\frac{3}{2}} + 15\sqrt{x})}{c^6} - \frac{15 \log(c\sqrt{x} + 1)}{c^7} + \frac{15 \log(c\sqrt{x} - 1)}{c^7} \right) \right) \operatorname{arctanh}(c\sqrt{x}) + \dots$$

input `integrate(x^2*(a+b*arctanh(c*x^(1/2)))^2,x, algorithm="maxima")`

output `1/3*b^2*x^3*arctanh(c*sqrt(x))^2 + 1/3*a^2*x^3 + 1/45*(30*x^3*arctanh(c*sqrt(x)) + c*(2*(3*c^4*x^(5/2) + 5*c^2*x^(3/2) + 15*sqrt(x))/c^6 - 15*log(c*sqrt(x) + 1)/c^7 + 15*log(c*sqrt(x) - 1)/c^7))*a*b + 1/180*(4*c*(2*(3*c^4*x^(5/2) + 5*c^2*x^(3/2) + 15*sqrt(x))/c^6 - 15*log(c*sqrt(x) + 1)/c^7 + 15*log(c*sqrt(x) - 1)/c^7)*arctanh(c*sqrt(x)) + (6*c^4*x^2 + 32*c^2*x - 2*(15*log(c*sqrt(x) - 1) - 46)*log(c*sqrt(x) + 1) + 15*log(c*sqrt(x) + 1)^2 + 15*log(c*sqrt(x) - 1)^2 + 92*log(c*sqrt(x) - 1))/c^6)*b^2`

Giac [F]

$$\int x^2(a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \int (b \operatorname{arctanh}(c\sqrt{x}) + a)^2 x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c*x^(1/2)))^2,x, algorithm="giac")`

output `integrate((b*arctanh(c*sqrt(x)) + a)^2*x^2, x)`

Mupad [B] (verification not implemented)

Time = 4.26 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.07

$$\int x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx$$

$$= \frac{46 b^2 \ln(c^2 x - 1) - 30 b^2 \operatorname{atanh}(c\sqrt{x})^2 - 60 a b \operatorname{atanh}(c\sqrt{x}) + 16 b^2 c^2 x + 30 a^2 c^6 x^3 + 3 b^2 c^4 x^2 + 30 b^2 c^2 x}{90 c^6}$$

input `int(x^2*(a + b*atanh(c*x^(1/2)))^2,x)`output
$$\frac{(46*b^2*\log(c^2*x - 1) - 30*b^2*\operatorname{atanh}(c*x^{(1/2)})^2 - 60*a*b*\operatorname{atanh}(c*x^{(1/2)}) + 16*b^2*c^2*x + 30*a^2*c^6*x^3 + 3*b^2*c^4*x^2 + 30*b^2*c^6*x^3*\operatorname{atanh}(c*x^{(1/2)})^2 + 60*b^2*c*x^{(1/2)}*\operatorname{atanh}(c*x^{(1/2)}) + 60*a*b*c*x^{(1/2)} + 20*b^2*c^3*x^{(3/2)}*\operatorname{atanh}(c*x^{(1/2)}) + 12*b^2*c^5*x^{(5/2)}*\operatorname{atanh}(c*x^{(1/2)}) + 20*a*b*c^3*x^{(3/2)} + 12*a*b*c^5*x^{(5/2)} + 60*a*b*c^6*x^3*\operatorname{atanh}(c*x^{(1/2)}))/90*c^6}$$
Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.09

$$\int x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx$$

$$= \frac{30 \operatorname{atanh}(\sqrt{x} c)^2 b^2 c^6 x^3 - 30 \operatorname{atanh}(\sqrt{x} c)^2 b^2 + 12 \sqrt{x} \operatorname{atanh}(\sqrt{x} c) b^2 c^5 x^2 + 20 \sqrt{x} \operatorname{atanh}(\sqrt{x} c) b^2 c^3 x + 30 a^2 c^6 x^3 + 3 b^2 c^4 x^2 + 16 b^2 c^2 x}{90 c^6}$$

input `int(x^2*(a+b*atanh(c*x^(1/2)))^2,x)`output
$$(30*\operatorname{atanh}(\operatorname{sqrt}(x)*c)**2*b**2*c**6*x**3 - 30*\operatorname{atanh}(\operatorname{sqrt}(x)*c)**2*b**2 + 12*\operatorname{sqrt}(x)*\operatorname{atanh}(\operatorname{sqrt}(x)*c)*b**2*c**5*x**2 + 20*\operatorname{sqrt}(x)*\operatorname{atanh}(\operatorname{sqrt}(x)*c)*b**2*c**3*x + 60*\operatorname{sqrt}(x)*\operatorname{atanh}(\operatorname{sqrt}(x)*c)*b**2*c + 60*\operatorname{atanh}(\operatorname{sqrt}(x)*c)*a*b*c**6*x**3 - 60*\operatorname{atanh}(\operatorname{sqrt}(x)*c)*a*b + 92*\operatorname{atanh}(\operatorname{sqrt}(x)*c)*b**2 + 12*\operatorname{sqrt}(x)*a*b*c**5*x**2 + 20*\operatorname{sqrt}(x)*a*b*c**3*x + 60*\operatorname{sqrt}(x)*a*b*c + 92*\log(\operatorname{sqrt}(x)*c - 1)*b**2 + 30*a**2*c**6*x**3 + 3*b**2*c**4*x**2 + 16*b**2*c**2*x)/(90*c**6)$$

3.196 $\int x(a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx$

Optimal result	1559
Mathematica [A] (verified)	1559
Rubi [A] (verified)	1560
Maple [B] (verified)	1563
Fricas [A] (verification not implemented)	1564
Sympy [F]	1564
Maxima [B] (verification not implemented)	1564
Giac [F]	1565
Mupad [B] (verification not implemented)	1566
Reduce [B] (verification not implemented)	1566

Optimal result

Integrand size = 16, antiderivative size = 129

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \frac{ab\sqrt{x}}{c^3} + \frac{b^2x}{6c^2} + \frac{b^2\sqrt{x}\operatorname{arctanh}(c\sqrt{x})}{c^3} + \frac{bx^{3/2}(a + b \operatorname{arctanh}(c\sqrt{x}))}{3c} - \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{2c^4} + \frac{1}{2}x^2(a + b \operatorname{arctanh}(c\sqrt{x}))^2 + \frac{2b^2 \log(1 - c^2x)}{3c^4}$$

output

```
a*b*x^(1/2)/c^3+1/6*b^2*x/c^2+b^2*x^(1/2)*arctanh(c*x^(1/2))/c^3+1/3*b*x^(3/2)*(a+b*arctanh(c*x^(1/2)))/c-1/2*(a+b*arctanh(c*x^(1/2)))^2/c^4+1/2*x^2*(a+b*arctanh(c*x^(1/2)))^2+2/3*b^2*ln(-c^2*x+1)/c^4
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.24

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \frac{6abc\sqrt{x} + b^2c^2x + 2abc^3x^{3/2} + 3a^2c^4x^2 + 2bc\sqrt{x}(3ac^3x^{3/2} + b(3 + c^2x)) \operatorname{arctanh}(c\sqrt{x}) + 3b^2(-1 + c^4x)}{6c^4}$$

input `Integrate[x*(a + b*ArcTanh[c*Sqrt[x]])^2,x]`

output $(6*a*b*c*\text{Sqrt}[x] + b^2*c^2*x + 2*a*b*c^3*x^{(3/2)} + 3*a^2*c^4*x^2 + 2*b*c*\text{Sqrt}[x]*(3*a*c^3*x^{(3/2)} + b*(3 + c^2*x))*\text{ArcTanh}[c*\text{Sqrt}[x]] + 3*b^2*(-1 + c^4*x^2)*\text{ArcTanh}[c*\text{Sqrt}[x]]^2 + b*(3*a + 4*b)*\text{Log}[1 - c*\text{Sqrt}[x]] - 3*a*b*\text{Log}[1 + c*\text{Sqrt}[x]] + 4*b^2*\text{Log}[1 + c*\text{Sqrt}[x]])/(6*c^4)$

Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.23, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {6454, 6452, 6542, 6452, 243, 49, 2009, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx$$

$$\downarrow 6454$$

$$2 \int x^{3/2}(a + b \operatorname{arctanh}(c\sqrt{x}))^2 d\sqrt{x}$$

$$\downarrow 6452$$

$$2 \left(\frac{1}{4} x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{2} bc \int \frac{x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x} \right)$$

$$\downarrow 6542$$

$$2 \left(\frac{1}{4} x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{2} bc \left(\frac{\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\int x(a + b \operatorname{arctanh}(c\sqrt{x})) d\sqrt{x}}{c^2} \right) \right)$$

$$\downarrow 6452$$

$$2 \left(\frac{1}{4} x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{1}{2} bc \left(\frac{\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{3} bc \int \frac{x^{3/2}}{1 - c^2 x} d\sqrt{x}}{c^2} \right) \right)$$

↓ 243

$$2 \left(\frac{1}{4} x^2 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{2} bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{6} bc \int \frac{x}{1 - c^2 x} dx}{c^2} \right) \right)$$

↓ 49

$$2 \left(\frac{1}{4} x^2 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{2} bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{6} bc \int \left(-\frac{1}{c^2} - \right)}{c^2} \right) \right)$$

↓ 2009

$$2 \left(\frac{1}{4} x^2 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{2} bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{6} bc \left(-\frac{x}{c^2} - \log \right)}{c^2} \right) \right)$$

↓ 6542

$$2 \left(\frac{1}{4} x^2 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{2} bc \left(\frac{\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\int (a + \operatorname{barctanh}(c\sqrt{x})) d\sqrt{x}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a + \operatorname{barctanh}(c\sqrt{x}))}{c^2} \right) \right)$$

↓ 2009

$$2 \left(\frac{1}{4} x^2 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{2} bc \left(\frac{\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{a\sqrt{x} + b\sqrt{x} \operatorname{arctanh}(c\sqrt{x}) + \frac{b \log(1 - c^2 x)}{2c}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a + \operatorname{barctanh}(c\sqrt{x}))}{c^2} \right) \right)$$

↓ 6510

$$2 \left(\frac{1}{4} x^2 (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{1}{2} bc \left(\frac{\left(\frac{a + \operatorname{barctanh}(c\sqrt{x})}{2bc^3} \right)^2 - \frac{a\sqrt{x} + b\sqrt{x} \operatorname{arctanh}(c\sqrt{x}) + \frac{b \log(1 - c^2 x)}{2c}}{c^2}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a + \operatorname{barctanh}(c\sqrt{x}))}{c^2} \right) \right)$$

input `Int[x*(a + b*ArcTanh[c*Sqrt[x]])^2,x]`

output

$$2*((x^2*(a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])^2)/4 - (b*c*(-((x^{3/2})*(a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])))/3 - (b*c*(-(x/c^2) - \text{Log}[1 - c^2*x]/c^4))/6)/c^2) + ((a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])^2/(2*b*c^3) - (a*\text{Sqrt}[x] + b*\text{Sqrt}[x]*\text{ArcTanh}[c*\text{Sqrt}[x]] + (b*\text{Log}[1 - c^2*x])/(2*c))/c^2)/c^2)/2$$

Defintions of rubi rules used

rule 49

$$\text{Int}[(a + b*x)^m*(c + d*x)^n, x] \text{Symbol} \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] \text{ ; FreeQ}\{a, b, c, d\}, x \text{ \&\& IGtQ}\{m, 0\} \text{ \&\& IGtQ}\{m + n + 2, 0\}$$

rule 243

$$\text{Int}(x^m*(a + b*x^2)^p, x) \text{Symbol} \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] \text{ ; FreeQ}\{a, b, m, p\}, x \text{ \&\& IntegerQ}\{m-1\}/2$$

rule 2009

$$\text{Int}[u, x] \text{Symbol} \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ ; SumQ}[u]$$

rule 6452

$$\text{Int}[(a + \text{ArcTanh}[c*x^n])^p*(b*x^m), x] \text{Symbol} \rightarrow \text{Simp}[x^{m+1}*(a + b*\text{ArcTanh}[c*x^n])^p/(m+1), x] - \text{Simp}[b*c^n*(p/(m+1)) \text{ Int}[x^{m+n}*(a + b*\text{ArcTanh}[c*x^n])^{p-1}/(1 - c^2*x^{2n}), x], x] \text{ ; FreeQ}\{a, b, c, m, n\}, x \text{ \&\& IGtQ}\{p, 0\} \text{ \&\& (EqQ}\{p, 1\} \text{ || (EqQ}\{n, 1\} \text{ \&\& IntegerQ}\{m\})) \text{ \&\& NeQ}\{m, -1\}$$

rule 6454

$$\text{Int}[(a + \text{ArcTanh}[c*x^n])^p*(b*x^m), x] \text{Symbol} \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}\{m+1\}/n - 1)*(a + b*\text{ArcTanh}[c*x])^p}, x], x, x^n], x] \text{ ; FreeQ}\{a, b, c, m, n\}, x \text{ \&\& IGtQ}\{p, 1\} \text{ \&\& IntegerQ}\{\text{Simplify}\{m+1\}/n\}$$

rule 6510

$$\text{Int}[(a + \text{ArcTanh}[c*x])^p/((d + e*x^2)), x] \text{Symbol} \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^p/(b*c*d*(p+1)), x] \text{ ; FreeQ}\{a, b, c, d, e, p\}, x \text{ \&\& EqQ}\{c^2*d + e, 0\} \text{ \&\& NeQ}\{p, -1\}$$

rule 6542

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*((f_.)*(x_))^(m_)/((d_) + (
e_.)*(x_)^2), x_Symbol] :> Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*
x])^p, x], x] - Simp[d*(f^2/e Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/
(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m,
1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 248 vs. $2(105) = 210$.

Time = 1.04 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.93

method	result
parts	$\frac{a^2 x^2}{2} + \frac{2b^2 \left(\frac{c^4 x^2 \operatorname{arctanh}(c\sqrt{x})^2}{4} + \frac{\operatorname{arctanh}(c\sqrt{x}) c^3 x^{\frac{3}{2}}}{6} + \frac{\operatorname{arctanh}(c\sqrt{x}) c\sqrt{x}}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{4} - \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1+c\sqrt{x})}{4} \right)}{1}$
derivativedivides	$\frac{a^2 c^4 x^2}{2} + 2b^2 \left(\frac{c^4 x^2 \operatorname{arctanh}(c\sqrt{x})^2}{4} + \frac{\operatorname{arctanh}(c\sqrt{x}) c^3 x^{\frac{3}{2}}}{6} + \frac{\operatorname{arctanh}(c\sqrt{x}) c\sqrt{x}}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{4} - \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1+c\sqrt{x})}{4} \right)$
default	$\frac{a^2 c^4 x^2}{2} + 2b^2 \left(\frac{c^4 x^2 \operatorname{arctanh}(c\sqrt{x})^2}{4} + \frac{\operatorname{arctanh}(c\sqrt{x}) c^3 x^{\frac{3}{2}}}{6} + \frac{\operatorname{arctanh}(c\sqrt{x}) c\sqrt{x}}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{4} - \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1+c\sqrt{x})}{4} \right)$

input

```
int(x*(a+b*arctanh(c*x^(1/2)))^2,x,method=_RETURNVERBOSE)
```

output

```
1/2*a^2*x^2+2*b^2/c^4*(1/4*c^4*x^2*arctanh(c*x^(1/2))^2+1/6*arctanh(c*x^(1/2))*c^3*x^(3/2)+1/2*arctanh(c*x^(1/2))*c*x^(1/2)+1/4*arctanh(c*x^(1/2))*ln(c*x^(1/2)-1)-1/4*arctanh(c*x^(1/2))*ln(1+c*x^(1/2))-1/8*ln(c*x^(1/2)-1)*ln(1/2*c*x^(1/2)+1/2)+1/16*ln(c*x^(1/2)-1)^2+1/16*ln(1+c*x^(1/2))^2-1/8*(ln(1+c*x^(1/2))-ln(1/2*c*x^(1/2)+1/2))*ln(-1/2*c*x^(1/2)+1/2)+1/12*c^2*x+1/3*ln(c*x^(1/2)-1)+1/3*ln(1+c*x^(1/2)))+4*a*b/c^4*(1/4*c^4*x^2*arctanh(c*x^(1/2))+1/12*c^3*x^(3/2)+1/4*c*x^(1/2)+1/8*ln(c*x^(1/2)-1)-1/8*ln(1+c*x^(1/2)))
```


Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.60

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx$$

$$= \frac{12a^2c^4x^2 + 4b^2c^2x + 3(b^2c^4x^2 - b^2) \log\left(-\frac{c^2x+2c\sqrt{x}+1}{c^2x-1}\right)^2 + 4(3abc^4 - 3ab + 4b^2) \log(c\sqrt{x} + 1) - 4(3$$

input `integrate(x*(a+b*arctanh(c*x^(1/2)))^2,x, algorithm="fricas")`

output

```
1/24*(12*a^2*c^4*x^2 + 4*b^2*c^2*x + 3*(b^2*c^4*x^2 - b^2)*log(-(c^2*x + 2
*c*sqrt(x) + 1)/(c^2*x - 1))^2 + 4*(3*a*b*c^4 - 3*a*b + 4*b^2)*log(c*sqrt(
x) + 1) - 4*(3*a*b*c^4 - 3*a*b - 4*b^2)*log(c*sqrt(x) - 1) + 4*(3*a*b*c^4*
x^2 - 3*a*b*c^4 + (b^2*c^3*x + 3*b^2*c)*sqrt(x))*log(-(c^2*x + 2*c*sqrt(x)
+ 1)/(c^2*x - 1)) + 8*(a*b*c^3*x + 3*a*b*c)*sqrt(x))/c^4
```

Sympy [F]

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \int x(a + b \operatorname{atanh}(c\sqrt{x}))^2 dx$$

input `integrate(x*(a+b*atanh(c*x**(1/2)))**2,x)`

output

```
Integral(x*(a + b*atanh(c*sqrt(x)))**2, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 215 vs. 2(105) = 210.

Time = 0.03 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.67

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \frac{1}{2} b^2 x^2 \operatorname{arctanh}(c\sqrt{x})^2 + \frac{1}{2} a^2 x^2 + \frac{1}{6} \left(6x^2 \operatorname{arctanh}(c\sqrt{x}) + c \left(\frac{2(c^2 x^{3/2} + 3\sqrt{x})}{c^4} - \frac{3 \log(c\sqrt{x} + 1)}{c^5} + \frac{3 \log(c\sqrt{x} - 1)}{c^5} \right) \right) ab + \frac{1}{24} \left(4c \left(\frac{2(c^2 x^{3/2} + 3\sqrt{x})}{c^4} - \frac{3 \log(c\sqrt{x} + 1)}{c^5} + \frac{3 \log(c\sqrt{x} - 1)}{c^5} \right) \operatorname{arctanh}(c\sqrt{x}) + \frac{4c^2 x - 2(3 \log(c\sqrt{x} + 1) - 3 \log(c\sqrt{x} - 1))}{c^4} \right) b^2$$

input `integrate(x*(a+b*arctanh(c*x^(1/2)))^2,x, algorithm="maxima")`

output `1/2*b^2*x^2*arctanh(c*sqrt(x))^2 + 1/2*a^2*x^2 + 1/6*(6*x^2*arctanh(c*sqrt(x)) + c*(2*(c^2*x^(3/2) + 3*sqrt(x))/c^4 - 3*log(c*sqrt(x) + 1)/c^5 + 3*log(c*sqrt(x) - 1)/c^5))*a*b + 1/24*(4*c*(2*(c^2*x^(3/2) + 3*sqrt(x))/c^4 - 3*log(c*sqrt(x) + 1)/c^5 + 3*log(c*sqrt(x) - 1)/c^5)*arctanh(c*sqrt(x)) + (4*c^2*x - 2*(3*log(c*sqrt(x) - 1) - 3*log(c*sqrt(x) + 1) - 8)*log(c*sqrt(x) + 1) + 3*log(c*sqrt(x) + 1)^2 + 3*log(c*sqrt(x) - 1)^2 + 16*log(c*sqrt(x) - 1))/c^4)*b^2`

Giac [F]

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \int (b \operatorname{arctanh}(c\sqrt{x}) + a)^2 x dx$$

input `integrate(x*(a+b*arctanh(c*x^(1/2)))^2,x, algorithm="giac")`

output `integrate((b*arctanh(c*sqrt(x)) + a)^2*x, x)`

3.197 $\int (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx$

Optimal result	1567
Mathematica [A] (verified)	1567
Rubi [A] (verified)	1568
Maple [B] (verified)	1570
Fricas [B] (verification not implemented)	1570
Sympy [F]	1571
Maxima [B] (verification not implemented)	1571
Giac [F]	1572
Mupad [B] (verification not implemented)	1572
Reduce [B] (verification not implemented)	1573

Optimal result

Integrand size = 14, antiderivative size = 85

$$\int (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \frac{2ab\sqrt{x}}{c} + \frac{2b^2\sqrt{x}\operatorname{arctanh}(c\sqrt{x})}{c} - \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} + x(a + b \operatorname{arctanh}(c\sqrt{x}))^2 + \frac{b^2 \log(1 - c^2x)}{c^2}$$

output `2*a*b*x^(1/2)/c+2*b^2*x^(1/2)*arctanh(c*x^(1/2))/c-(a+b*arctanh(c*x^(1/2)))^2/c^2+x*(a+b*arctanh(c*x^(1/2)))^2+b^2*ln(-c^2*x+1)/c^2`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.35

$$\int (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \frac{2abc\sqrt{x} + a^2c^2x + 2bc(b + ac\sqrt{x})\sqrt{x}\operatorname{arctanh}(c\sqrt{x}) + b^2(-1 + c^2x)\operatorname{arctanh}(c\sqrt{x})^2 + b(a + b)\log(1 - c^2x)}{c^2}$$

input `Integrate[(a + b*ArcTanh[c*Sqrt[x]])^2, x]`

output

```
(2*a*b*c*Sqrt[x] + a^2*c^2*x + 2*b*c*(b + a*c*Sqrt[x])*Sqrt[x]*ArcTanh[c*Sqrt[x]] + b^2*(-1 + c^2*x)*ArcTanh[c*Sqrt[x]]^2 + b*(a + b)*Log[1 - c*Sqrt[x]] - a*b*Log[1 + c*Sqrt[x]] + b^2*Log[1 + c*Sqrt[x]])/c^2
```

Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.13, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {6442, 6452, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx$$

$$\downarrow 6442$$

$$2 \int \sqrt{x} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 d\sqrt{x}$$

$$\downarrow 6452$$

$$2 \left(\frac{1}{2} x (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - bc \int \frac{x (a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x} \right)$$

$$\downarrow 6542$$

$$2 \left(\frac{1}{2} x (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\int (a + b \operatorname{arctanh}(c\sqrt{x})) d\sqrt{x}}{c^2} \right) \right)$$

$$\downarrow 2009$$

$$2 \left(\frac{1}{2} x (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{a\sqrt{x} + b\sqrt{x} \operatorname{arctanh}(c\sqrt{x}) + \frac{b \log(1 - c^2 x)}{2c}}{c^2} \right) \right)$$

$$\downarrow 6510$$

$$2 \left(\frac{1}{2} x (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - bc \left(\frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{2bc^3} - \frac{a\sqrt{x} + b\sqrt{x} \operatorname{arctanh}(c\sqrt{x}) + \frac{b \log(1 - c^2 x)}{2c}}{c^2} \right) \right)$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])^2,x]`

output `2*((x*(a + b*ArcTanh[c*Sqrt[x]])^2)/2 - b*c*((a + b*ArcTanh[c*Sqrt[x]])^2/(2*b*c^3) - (a*Sqrt[x] + b*Sqrt[x]*ArcTanh[c*Sqrt[x]] + (b*Log[1 - c^2*x])/(2*c))/c^2))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6442 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^(p_), x_Symbol] := With[{k = Denominator[n]}, Simp[k Subst[Int[x^(k - 1)*(a + b*ArcTanh[c*x^(k*n)])^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1] && FractionQ[n]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6510 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6542 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_.))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 212 vs. 2(75) = 150.

Time = 1.00 (sec) , antiderivative size = 213, normalized size of antiderivative = 2.51

method	result
parts	$x a^2 + \frac{2b^2 \left(\frac{c^2 x \operatorname{arctanh}(c\sqrt{x})^2}{2} + \operatorname{arctanh}(c\sqrt{x})c\sqrt{x} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{2} - \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1+c\sqrt{x})}{2} - \frac{\ln(c\sqrt{x}-1) \ln(c\sqrt{x}+1)}{4} \right)}{a^2 c^2 x + 2b^2}$
derivativedivides	$\frac{a^2 c^2 x + 2b^2 \left(\frac{c^2 x \operatorname{arctanh}(c\sqrt{x})^2}{2} + \operatorname{arctanh}(c\sqrt{x})c\sqrt{x} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{2} - \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1+c\sqrt{x})}{2} - \frac{\ln(c\sqrt{x}-1) \ln(c\sqrt{x}+1)}{4} \right)}{a^2 c^2 x + 2b^2}$
default	$\frac{a^2 c^2 x + 2b^2 \left(\frac{c^2 x \operatorname{arctanh}(c\sqrt{x})^2}{2} + \operatorname{arctanh}(c\sqrt{x})c\sqrt{x} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{2} - \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1+c\sqrt{x})}{2} - \frac{\ln(c\sqrt{x}-1) \ln(c\sqrt{x}+1)}{4} \right)}{a^2 c^2 x + 2b^2}$

input `int((a+b*arctanh(c*x^(1/2)))^2,x,method=_RETURNVERBOSE)`

output `x*a^2+2*b^2/c^2*(1/2*c^2*x*arctanh(c*x^(1/2))^2+arctanh(c*x^(1/2))*c*x^(1/2)+1/2*arctanh(c*x^(1/2))*ln(c*x^(1/2)-1)-1/2*arctanh(c*x^(1/2))*ln(1+c*x^(1/2))-1/4*ln(c*x^(1/2)-1)*ln(1/2*c*x^(1/2)+1/2)+1/8*ln(c*x^(1/2)-1)^2+1/2*ln(c*x^(1/2)-1)+1/2*ln(1+c*x^(1/2))+1/8*ln(1+c*x^(1/2))^2-1/4*(ln(1+c*x^(1/2))-ln(1/2*c*x^(1/2)+1/2))*ln(-1/2*c*x^(1/2)+1/2))+4*a*b/c^2*(1/2*c^2*x*arctanh(c*x^(1/2))+1/2*c*x^(1/2)+1/4*ln(c*x^(1/2)-1)-1/4*ln(1+c*x^(1/2)))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 165 vs. 2(75) = 150.

Time = 0.10 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.94

$$\int (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx$$

$$= \frac{4 a^2 c^2 x + 8 a b c \sqrt{x} + (b^2 c^2 x - b^2) \log\left(-\frac{c^2 x + 2 c \sqrt{x} + 1}{c^2 x - 1}\right)^2 + 4 (a b c^2 - a b + b^2) \log(c\sqrt{x} + 1) - 4 (a b c^2 - a b)}{4 c^2}$$

input `integrate((a+b*arctanh(c*x^(1/2)))^2,x, algorithm="fricas")`

output

```
1/4*(4*a^2*c^2*x + 8*a*b*c*sqrt(x) + (b^2*c^2*x - b^2)*log(-(c^2*x + 2*c*sqrt(x) + 1)/(c^2*x - 1))^2 + 4*(a*b*c^2 - a*b + b^2)*log(c*sqrt(x) + 1) - 4*(a*b*c^2 - a*b - b^2)*log(c*sqrt(x) - 1) + 4*(a*b*c^2*x - a*b*c^2 + b^2*c*sqrt(x))*log(-(c^2*x + 2*c*sqrt(x) + 1)/(c^2*x - 1)))/c^2
```

Sympy [F]

$$\int (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \int (a + b \operatorname{atanh}(c\sqrt{x}))^2 dx$$

input

```
integrate((a+b*atanh(c*x**(1/2)))**2,x)
```

output

```
Integral((a + b*atanh(c*sqrt(x)))**2, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 175 vs. $2(75) = 150$.

Time = 0.04 (sec) , antiderivative size = 175, normalized size of antiderivative = 2.06

$$\begin{aligned} & \int (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx \\ &= \left(c \left(\frac{2\sqrt{x}}{c^2} - \frac{\log(c\sqrt{x} + 1)}{c^3} + \frac{\log(c\sqrt{x} - 1)}{c^3} \right) + 2x \operatorname{artanh}(c\sqrt{x}) \right) ab \\ &+ \frac{1}{4} \left(4c \left(\frac{2\sqrt{x}}{c^2} - \frac{\log(c\sqrt{x} + 1)}{c^3} + \frac{\log(c\sqrt{x} - 1)}{c^3} \right) \operatorname{artanh}(c\sqrt{x}) + 4x \operatorname{artanh}(c\sqrt{x})^2 - \frac{2(\log(c\sqrt{x} + 1) - \log(c\sqrt{x} - 1))^2}{4} \right) \\ &+ a^2x \end{aligned}$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))^2,x, algorithm="maxima")
```


output

```
(c*(2*sqrt(x)/c^2 - log(c*sqrt(x) + 1)/c^3 + log(c*sqrt(x) - 1)/c^3) + 2*x
*arctanh(c*sqrt(x)))*a*b + 1/4*(4*c*(2*sqrt(x)/c^2 - log(c*sqrt(x) + 1)/c^
3 + log(c*sqrt(x) - 1)/c^3)*arctanh(c*sqrt(x)) + 4*x*arctanh(c*sqrt(x))^2
- (2*(log(c*sqrt(x) - 1) - 2)*log(c*sqrt(x) + 1) - log(c*sqrt(x) + 1)^2 -
log(c*sqrt(x) - 1)^2 - 4*log(c*sqrt(x) - 1))/c^2)*b^2 + a^2*x
```

Giac [F]

$$\int (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = \int (b \operatorname{artanh}(c\sqrt{x}) + a)^2 dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*sqrt(x)) + a)^2, x)
```

Mupad [B] (verification not implemented)

Time = 3.77 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.11

$$\int (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx = a^2 x + \frac{c(2b^2 \sqrt{x} \operatorname{atanh}(c\sqrt{x}) + 2ab\sqrt{x}) - b^2 \operatorname{atanh}(c\sqrt{x})^2 + b^2 \ln(c^2 x - 1) - 2ab \operatorname{atanh}(c\sqrt{x})}{c^2} + b^2 x \operatorname{atanh}(c\sqrt{x})^2 + 2abx \operatorname{atanh}(c\sqrt{x})$$

input

```
int((a + b*atanh(c*x^(1/2)))^2,x)
```

output

```
a^2*x + (c*(2*b^2*x^(1/2)*atanh(c*x^(1/2)) + 2*a*b*x^(1/2)) - b^2*atanh(c*
x^(1/2))^2 + b^2*log(c^2*x - 1) - 2*a*b*atanh(c*x^(1/2)))/c^2 + b^2*x*atan
h(c*x^(1/2))^2 + 2*a*b*x*atanh(c*x^(1/2))
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.22

$$\int (a + b \operatorname{arctanh}(c\sqrt{x}))^2 dx$$

$$= \frac{\operatorname{atanh}(\sqrt{x}c)^2 b^2 c^2 x - \operatorname{atanh}(\sqrt{x}c)^2 b^2 + 2\sqrt{x} \operatorname{atanh}(\sqrt{x}c) b^2 c + 2 \operatorname{atanh}(\sqrt{x}c) a b c^2 x - 2 \operatorname{atanh}(\sqrt{x}c) a b c^2 x}{c^2}$$

input `int((a+b*atanh(c*x^(1/2)))^2,x)`output `(atanh(sqrt(x)*c)**2*b**2*c**2*x - atanh(sqrt(x)*c)**2*b**2 + 2*sqrt(x)*atanh(sqrt(x)*c)*b**2*c + 2*atanh(sqrt(x)*c)*a*b*c**2*x - 2*atanh(sqrt(x)*c)*a*b + 2*atanh(sqrt(x)*c)*b**2 + 2*sqrt(x)*a*b*c + 2*log(sqrt(x)*c - 1)*b**2 + a**2*c**2*x)/c**2`

$$3.198 \quad \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x} dx$$

Optimal result	1574
Mathematica [C] (verified)	1575
Rubi [A] (verified)	1576
Maple [C] (warning: unable to verify)	1578
Fricas [F]	1579
Sympy [F]	1579
Maxima [F]	1580
Giac [F]	1580
Mupad [F(-1)]	1580
Reduce [F]	1581

Optimal result

Integrand size = 18, antiderivative size = 145

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x} dx = & 4 \operatorname{arctanh}\left(1 - \frac{2}{1 - c\sqrt{x}}\right) (a + b \operatorname{arctanh}(c\sqrt{x}))^2 \\ & - 2b(a + b \operatorname{arctanh}(c\sqrt{x})) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - c\sqrt{x}}\right) \\ & + 2b(a + b \operatorname{arctanh}(c\sqrt{x})) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - c\sqrt{x}}\right) \\ & + b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - c\sqrt{x}}\right) \\ & - b^2 \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 - c\sqrt{x}}\right) \end{aligned}$$

output

```
-4*arctanh(-1+2/(1-c*x^(1/2)))*(a+b*arctanh(c*x^(1/2)))^2-2*b*(a+b*arctanh
(c*x^(1/2))*polylog(2,1-2/(1-c*x^(1/2)))+2*b*(a+b*arctanh(c*x^(1/2))*pol
ylog(2,-1+2/(1-c*x^(1/2)))+b^2*polylog(3,1-2/(1-c*x^(1/2)))-b^2*polylog(3,
-1+2/(1-c*x^(1/2)))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.40

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x} dx = a^2 \log(x) + 2ab(-\operatorname{PolyLog}(2, -c\sqrt{x}) + \operatorname{PolyLog}(2, c\sqrt{x})) + 2b^2 \left(\frac{i\pi^3}{24} - \frac{2}{3} \operatorname{arctanh}(c\sqrt{x})^3 - \operatorname{arctanh}(c\sqrt{x})^2 \log(1 + e^{-2\operatorname{arctanh}(c\sqrt{x})}) + \operatorname{arctanh}(c\sqrt{x})^2 \log(1 - e^{2\operatorname{arctanh}(c\sqrt{x})}) + \operatorname{arctanh}(c\sqrt{x}) \operatorname{PolyLog}(2, -e^{-2\operatorname{arctanh}(c\sqrt{x})}) + \operatorname{arctanh}(c\sqrt{x}) \operatorname{PolyLog}(2, e^{2\operatorname{arctanh}(c\sqrt{x})}) + \frac{1}{2} \operatorname{PolyLog}(3, -e^{-2\operatorname{arctanh}(c\sqrt{x})}) - \frac{1}{2} \operatorname{PolyLog}(3, e^{2\operatorname{arctanh}(c\sqrt{x})}) \right)$$

input `Integrate[(a + b*ArcTanh[c*Sqrt[x]])^2/x,x]`

output `a^2*Log[x] + 2*a*b*(-PolyLog[2, -(c*Sqrt[x])] + PolyLog[2, c*Sqrt[x]]) + 2*b^2*((I/24)*Pi^3 - (2*ArcTanh[c*Sqrt[x]]^3)/3 - ArcTanh[c*Sqrt[x]]^2*Log[1 + E^(-2*ArcTanh[c*Sqrt[x]])] + ArcTanh[c*Sqrt[x]]^2*Log[1 - E^(2*ArcTanh[c*Sqrt[x]])] + ArcTanh[c*Sqrt[x]]*PolyLog[2, -E^(-2*ArcTanh[c*Sqrt[x]])] + ArcTanh[c*Sqrt[x]]*PolyLog[2, E^(2*ArcTanh[c*Sqrt[x]])] + PolyLog[3, -E^(-2*ArcTanh[c*Sqrt[x]])]/2 - PolyLog[3, E^(2*ArcTanh[c*Sqrt[x]])]/2)`

Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.22, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {6450, 6448, 6614, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x} dx$$

$$\downarrow 6450$$

$$2 \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{\sqrt{x}} d\sqrt{x}$$

$$\downarrow 6448$$

$$2 \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - c\sqrt{x}} \right) (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - 4bc \int \frac{\operatorname{arctanh} \left(1 - \frac{2}{1 - c\sqrt{x}} \right) (a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x} \right)$$

$$\downarrow 6614$$

$$2 \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - c\sqrt{x}} \right) (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - 4bc \left(\frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x})) \log \left(2 - \frac{2}{1 - c\sqrt{x}} \right)}{1 - c^2 x} d\sqrt{x} - \frac{1}{2} \right) \right)$$

$$\downarrow 6620$$

$$2 \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - c\sqrt{x}} \right) (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - 4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - c\sqrt{x}} \right) (a + b \operatorname{arctanh}(c\sqrt{x}))}{2c} \right) \right) \right)$$

$$\downarrow 7164$$

$$2 \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - c\sqrt{x}} \right) (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - 4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - c\sqrt{x}} \right) (a + b \operatorname{arctanh}(c\sqrt{x}))}{2c} \right) \right) \right)$$

input

```
Int[(a + b*ArcTanh[c*Sqrt[x]])^2/x,x]
```

output

$$2*(2*\text{ArcTanh}[1 - 2/(1 - c*\text{Sqrt}[x])]*(a + b*\text{ArcTanh}[c*\text{Sqrt}[x]))^2 - 4*b*c*((a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])*\text{PolyLog}[2, 1 - 2/(1 - c*\text{Sqrt}[x])])/(2*c) - (b*\text{PolyLog}[3, 1 - 2/(1 - c*\text{Sqrt}[x])])/(4*c))/2 + (-1/2*((a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])*\text{PolyLog}[2, -1 + 2/(1 - c*\text{Sqrt}[x])])/c + (b*\text{PolyLog}[3, -1 + 2/(1 - c*\text{Sqrt}[x])])/(4*c))/2)$$

Defintions of rubi rules used

rule 6448

$$\text{Int}[(a + \text{ArcTanh}[c*x])^p * \text{ArcTanh}[1 - 2/(1 - c*x)], x] - \text{Simp}[2*b*c*p \text{Int}[(a + b*\text{ArcTanh}[c*x])^{p-1} * \text{ArcTanh}[1 - 2/(1 - c*x)]/(1 - c^2*x^2)], x] /;$$

$$\text{FreeQ}\{a, b, c, x\} \ \&\& \ \text{IGtQ}[p, 1]$$

rule 6450

$$\text{Int}[(a + \text{ArcTanh}[c*x]^n) * \text{ArcTanh}[1 - 2/(1 - c*x)]^p / x, x] - \text{Simp}[1/n \text{Subst}[\text{Int}[(a + b*\text{ArcTanh}[c*x])^p / x, x], x, x^n], x] /;$$

$$\text{FreeQ}\{a, b, c, n, x\} \ \&\& \ \text{IGtQ}[p, 0]$$

rule 6614

$$\text{Int}[\text{ArcTanh}[u] * (a + \text{ArcTanh}[c*x])^p / ((d + e*x^2)), x] - \text{Simp}[1/2 \text{Int}[\text{Log}[1 + u] * (a + b*\text{ArcTanh}[c*x])^p / (d + e*x^2)], x] - \text{Simp}[1/2 \text{Int}[\text{Log}[1 - u] * (a + b*\text{ArcTanh}[c*x])^p / (d + e*x^2)], x] /;$$

$$\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{EqQ}[u^2 - (1 - 2/(1 - c*x))^2, 0]$$

rule 6620

$$\text{Int}[(\text{Log}[u] * (a + \text{ArcTanh}[c*x])^p) / ((d + e*x^2)), x] - \text{Simp}[(-a + b*\text{ArcTanh}[c*x])^p * (\text{PolyLog}[2, 1 - u] / (2*c*d)), x] + \text{Simp}[b*(p/2) \text{Int}[(a + b*\text{ArcTanh}[c*x])^{p-1} * (\text{PolyLog}[2, 1 - u] / (d + e*x^2)), x], x] /;$$

$$\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{EqQ}[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]$$

rule 7164

$$\text{Int}[u * \text{PolyLog}[n, v], x] - \text{Simp}[w * \text{PolyLog}[n + 1, v], x] /;$$

$$\text{With}\{w = \text{DerivativeDivides}[v, u*v, x]\}, \text{Simp}[w * \text{PolyLog}[n + 1, v], x] /;$$

$$\text{!FalseQ}[w] /;$$

$$\text{FreeQ}[n, x]$$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 12.20 (sec) , antiderivative size = 662, normalized size of antiderivative = 4.57

method	result
parts	$a^2 \ln(x) + b^2 \left(2 \ln(c\sqrt{x}) \operatorname{arctanh}(c\sqrt{x})^2 - 2 \operatorname{arctanh}(c\sqrt{x}) \operatorname{polylog}\left(2, -\frac{(1+c\sqrt{x})^2}{-c^2x+1}\right) \right)$
derivativedivides	$2a^2 \ln(c\sqrt{x}) + 2b^2 \left(\ln(c\sqrt{x}) \operatorname{arctanh}(c\sqrt{x})^2 - \operatorname{arctanh}(c\sqrt{x}) \operatorname{polylog}\left(2, -\frac{(1+c\sqrt{x})^2}{-c^2x+1}\right) \right)$
default	$2a^2 \ln(c\sqrt{x}) + 2b^2 \left(\ln(c\sqrt{x}) \operatorname{arctanh}(c\sqrt{x})^2 - \operatorname{arctanh}(c\sqrt{x}) \operatorname{polylog}\left(2, -\frac{(1+c\sqrt{x})^2}{-c^2x+1}\right) \right)$

input

```
int((a+b*arctanh(c*x^(1/2)))^2/x,x,method=_RETURNVERBOSE)
```

output

```

a^2*ln(x)+b^2*(2*ln(c*x^(1/2))*arctanh(c*x^(1/2))^2-2*arctanh(c*x^(1/2))*p
olylog(2,-(1+c*x^(1/2))^2/(-c^2*x+1))+polylog(3,-(1+c*x^(1/2))^2/(-c^2*x+1
))-2*arctanh(c*x^(1/2))^2*ln((1+c*x^(1/2))^2/(-c^2*x+1)-1)+2*arctanh(c*x^(
1/2))^2*ln(1-(1+c*x^(1/2))/(-c^2*x+1)^(1/2))+4*arctanh(c*x^(1/2))*polylog(
2,(1+c*x^(1/2))/(-c^2*x+1)^(1/2))-4*polylog(3,(1+c*x^(1/2))/(-c^2*x+1)^(1/
2))+2*arctanh(c*x^(1/2))^2*ln(1+(1+c*x^(1/2))/(-c^2*x+1)^(1/2))+4*arctanh(
c*x^(1/2))*polylog(2,-(1+c*x^(1/2))/(-c^2*x+1)^(1/2))-4*polylog(3,-(1+c*x^
(1/2))/(-c^2*x+1)^(1/2))+I*Pi*csgn(I*(-(1+c*x^(1/2))^2/(c^2*x-1)-1)/(1-(1+
c*x^(1/2))^2/(c^2*x-1)))*csgn(I*(-(1+c*x^(1/2))^2/(c^2*x-1)-1))*csgn(I/(1
-(1+c*x^(1/2))^2/(c^2*x-1)))-csgn(I*(-(1+c*x^(1/2))^2/(c^2*x-1)-1))*csgn(I
*(-(1+c*x^(1/2))^2/(c^2*x-1)-1)/(1-(1+c*x^(1/2))^2/(c^2*x-1)))-csgn(I*(-(1
+c*x^(1/2))^2/(c^2*x-1)-1)/(1-(1+c*x^(1/2))^2/(c^2*x-1)))*csgn(I/(1-(1+c*x
^(1/2))^2/(c^2*x-1)))+csgn(I*(-(1+c*x^(1/2))^2/(c^2*x-1)-1)/(1-(1+c*x^(1/2
))^2/(c^2*x-1)))^2)*arctanh(c*x^(1/2))^2+2*a*b*(2*ln(c*x^(1/2))*arctanh(c
*x^(1/2))-dilog(c*x^(1/2))-dilog(1+c*x^(1/2))-ln(c*x^(1/2))*ln(1+c*x^(1/2
)))

```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x} dx = \int \frac{(b \operatorname{artanh}(c\sqrt{x}) + a)^2}{x} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))^2/x,x, algorithm="fricas")
```

output

```
integral((b^2*arctanh(c*sqrt(x))^2 + 2*a*b*arctanh(c*sqrt(x)) + a^2)/x, x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x} dx = \int \frac{(a + b \operatorname{atanh}(c\sqrt{x}))^2}{x} dx$$

input

```
integrate((a+b*atanh(c*x**(1/2)))**2/x,x)
```


output `Integral((a + b*atanh(c*sqrt(x)))**2/x, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x} dx = \int \frac{(b \operatorname{arctanh}(c\sqrt{x}) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))^2/x,x, algorithm="maxima")`

output `1/4*b^2*integrate(log(c*sqrt(x) + 1)^2/x, x) - 1/2*b^2*integrate(log(c*sqrt(x) + 1)*log(-c*sqrt(x) + 1)/x, x) + 1/4*b^2*integrate(log(-c*sqrt(x) + 1)^2/x, x) + a*b*integrate(log(c*sqrt(x) + 1)/x, x) - a*b*integrate(log(-c*sqrt(x) + 1)/x, x) + a^2*log(x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x} dx = \int \frac{(b \operatorname{arctanh}(c\sqrt{x}) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))^2/x,x, algorithm="giac")`

output `integrate((b*arctanh(c*sqrt(x)) + a)^2/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x} dx = \int \frac{(a + b \operatorname{atanh}(c\sqrt{x}))^2}{x} dx$$

input `int((a + b*atanh(c*x^(1/2)))^2/x,x)`

output `int((a + b*atanh(c*x^(1/2)))^2/x, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x} dx = 2 \left(\int \frac{\operatorname{atanh}(\sqrt{x}c)}{x} dx \right) ab + \left(\int \frac{\operatorname{atanh}(\sqrt{x}c)^2}{x} dx \right) b^2 + \log(x) a^2$$

input `int((a+b*atanh(c*x^(1/2)))^2/x,x)`

output `2*int(atanh(sqrt(x)*c)/x,x)*a*b + int(atanh(sqrt(x)*c)**2/x,x)*b**2 + log(x)*a**2`

3.199 $\int \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{x^2} dx$

Optimal result	1582
Mathematica [A] (verified)	1582
Rubi [A] (verified)	1583
Maple [B] (verified)	1586
Fricas [B] (verification not implemented)	1587
Sympy [B] (verification not implemented)	1587
Maxima [B] (verification not implemented)	1588
Giac [F]	1589
Mupad [B] (verification not implemented)	1589
Reduce [B] (verification not implemented)	1590

Optimal result

Integrand size = 18, antiderivative size = 85

$$\int \frac{(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{x^2} dx = -\frac{2bc(a + b\operatorname{arctanh}(c\sqrt{x}))}{\sqrt{x}} + c^2(a + b\operatorname{arctanh}(c\sqrt{x}))^2 - \frac{(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{x} + b^2c^2 \log(x) - b^2c^2 \log(1 - c^2x)$$

output

```
-2*b*c*(a+b*arctanh(c*x^(1/2)))/x^(1/2)+c^2*(a+b*arctanh(c*x^(1/2)))^2-(a+b*arctanh(c*x^(1/2)))^2/x+b^2*c^2*ln(x)-b^2*c^2*ln(-c^2*x+1)
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.52

$$\int \frac{(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{x^2} dx = \frac{a^2 + 2abc\sqrt{x} + 2b(a + bc\sqrt{x}) \operatorname{arctanh}(c\sqrt{x}) - b^2(-1 + c^2x) \operatorname{arctanh}(c\sqrt{x})^2 + b(a + b)c^2x \log(1 - c^2x)}{x}$$

input `Integrate[(a + b*ArcTanh[c*Sqrt[x]])^2/x^2,x]`

output
$$-\left(\frac{a^2 + 2ab\sqrt{x} + 2b(a + b\sqrt{x})\operatorname{ArcTanh}[c\sqrt{x}] - b^2(-1 + c^2x)\operatorname{ArcTanh}[c\sqrt{x}]^2 + b(a + b)c^2x\log[1 - c\sqrt{x}] - abc^2x\log[1 + c\sqrt{x}] + b^2c^2x\log[1 + c\sqrt{x}] - b^2c^2x\log[x]}{x}\right)$$

Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.05, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6454, 6452, 6544, 6452, 243, 47, 14, 16, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{x^2} dx \\ & \quad \downarrow \text{6454} \\ & 2 \int \frac{(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{x^{3/2}} d\sqrt{x} \\ & \quad \downarrow \text{6452} \\ & 2 \left(bc \int \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{x(1 - c^2x)} d\sqrt{x} - \frac{(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{2x} \right) \\ & \quad \downarrow \text{6544} \\ & 2 \left(bc \left(c^2 \int \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{1 - c^2x} d\sqrt{x} + \int \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{x} d\sqrt{x} \right) - \frac{(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{2x} \right) \\ & \quad \downarrow \text{6452} \\ & 2 \left(bc \left(c^2 \int \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{1 - c^2x} d\sqrt{x} + bc \int \frac{1}{\sqrt{x}(1 - c^2x)} d\sqrt{x} - \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}} \right) - \frac{(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{2x} \right) \\ & \quad \downarrow \text{243} \end{aligned}$$

$$2 \left(bc \left(c^2 \int \frac{a + \operatorname{arctanh}(c\sqrt{x})}{1 - c^2x} d\sqrt{x} + \frac{1}{2} bc \int \frac{1}{\sqrt{x}(1 - c^2x)} dx - \frac{a + \operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}} \right) - \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^2}{2x} \right)$$

↓ 47

$$2 \left(bc \left(c^2 \int \frac{a + \operatorname{arctanh}(c\sqrt{x})}{1 - c^2x} d\sqrt{x} + \frac{1}{2} bc \left(c^2 \int \frac{1}{1 - c^2x} dx + \int \frac{1}{\sqrt{x}} dx \right) - \frac{a + \operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}} \right) - \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^2}{2x} \right)$$

↓ 14

$$2 \left(bc \left(c^2 \int \frac{a + \operatorname{arctanh}(c\sqrt{x})}{1 - c^2x} d\sqrt{x} + \frac{1}{2} bc \left(c^2 \int \frac{1}{1 - c^2x} dx + \log(x) \right) - \frac{a + \operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}} \right) - \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^2}{2x} \right)$$

↓ 16

$$2 \left(bc \left(c^2 \int \frac{a + \operatorname{arctanh}(c\sqrt{x})}{1 - c^2x} d\sqrt{x} - \frac{a + \operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}} + \frac{1}{2} bc (\log(x) - \log(1 - c^2x)) \right) - \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^2}{2x} \right)$$

↓ 6510

$$2 \left(bc \left(\frac{c(a + \operatorname{arctanh}(c\sqrt{x}))^2}{2b} - \frac{a + \operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}} + \frac{1}{2} bc (\log(x) - \log(1 - c^2x)) \right) - \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^2}{2x} \right)$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])^2/x^2,x]`

output `2*(-1/2*(a + b*ArcTanh[c*Sqrt[x]])^2/x + b*c*(-((a + b*ArcTanh[c*Sqrt[x]])/Sqrt[x]) + (c*(a + b*ArcTanh[c*Sqrt[x]])^2)/(2*b) + (b*c*(Log[x] - Log[1 - c^2*x]))/2)`

Definitions of rubi rules used

- rule 14 $\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ ; FreeQ}[a, x]$
- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$
- rule 47 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x]$
- rule 243 $\text{Int}[(x_)^{(m_)}*((a_)+(b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] \text{ ; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$
- rule 6452 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{2n})}), x], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$
- rule 6454 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n] - 1)}*(a + b*\text{ArcTanh}[c*x])^p, x], x, x^n], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$
- rule 6510 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)]*(b_))^{(p_)}((d_)+(e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^{(p+1)}/(b*c*d*(p+1)), x] \text{ ; FreeQ}[\{a, b, c, d, e, p\}, x] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{NeQ}[p, -1]$

rule 6544

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^ (p_.)*((f_.)*(x_))^(m_)/((d_) + (
e_.)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x
], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x
^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 230 vs. $2(77) = 154$.

Time = 0.46 (sec) , antiderivative size = 231, normalized size of antiderivative = 2.72

method	result
parts	$-\frac{a^2}{x} + 2b^2c^2 \left(-\frac{\operatorname{arctanh}(c\sqrt{x})^2}{2c^2x} - \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1+c\sqrt{x})}{2} - \frac{\operatorname{arctanh}(c\sqrt{x})}{c\sqrt{x}} \right)$
derivativedivides	$2c^2 \left(-\frac{a^2}{2c^2x} + b^2 \left(-\frac{\operatorname{arctanh}(c\sqrt{x})^2}{2c^2x} - \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1+c\sqrt{x})}{2} - \frac{\operatorname{arctanh}(c\sqrt{x})}{c\sqrt{x}} \right) \right)$
default	$2c^2 \left(-\frac{a^2}{2c^2x} + b^2 \left(-\frac{\operatorname{arctanh}(c\sqrt{x})^2}{2c^2x} - \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1+c\sqrt{x})}{2} - \frac{\operatorname{arctanh}(c\sqrt{x})}{c\sqrt{x}} \right) \right)$

input

```
int((a+b*arctanh(c*x^(1/2)))^2/x^2,x,method=_RETURNVERBOSE)
```

output

```
-1/x*a^2+2*b^2*c^2*(-1/2/c^2/x*arctanh(c*x^(1/2))^2-1/2*arctanh(c*x^(1/2))
*ln(c*x^(1/2)-1)+1/2*arctanh(c*x^(1/2))*ln(1+c*x^(1/2))-arctanh(c*x^(1/2))
/c/x^(1/2)+1/4*ln(c*x^(1/2)-1)*ln(1/2*c*x^(1/2)+1/2)-1/8*ln(c*x^(1/2)-1)^2
-1/2*ln(c*x^(1/2)-1)-1/2*ln(1+c*x^(1/2))+ln(c*x^(1/2))-1/8*ln(1+c*x^(1/2))
^2+1/4*(ln(1+c*x^(1/2))-ln(1/2*c*x^(1/2)+1/2))*ln(-1/2*c*x^(1/2)+1/2))+4*a
*b*c^2*(-1/2/c^2/x*arctanh(c*x^(1/2))-1/4*ln(c*x^(1/2)-1)-1/2/c/x^(1/2)+1/
4*ln(1+c*x^(1/2)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 157 vs. 2(77) = 154.

Time = 0.10 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.85

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x^2} dx$$

$$= \frac{8b^2c^2x \log(\sqrt{x}) + 4(ab - b^2)c^2x \log(c\sqrt{x} + 1) - 4(ab + b^2)c^2x \log(c\sqrt{x} - 1) - 8abc\sqrt{x} + (b^2c^2x - b^2)}{4x}$$

input `integrate((a+b*arctanh(c*x^(1/2)))^2/x^2,x, algorithm="fricas")`

output `1/4*(8*b^2*c^2*x*log(sqrt(x)) + 4*(a*b - b^2)*c^2*x*log(c*sqrt(x) + 1) - 4*(a*b + b^2)*c^2*x*log(c*sqrt(x) - 1) - 8*a*b*c*sqrt(x) + (b^2*c^2*x - b^2)*log(-(c^2*x + 2*c*sqrt(x) + 1)/(c^2*x - 1))^2 - 4*a^2 - 4*(b^2*c*sqrt(x) + a*b)*log(-(c^2*x + 2*c*sqrt(x) + 1)/(c^2*x - 1)))/x`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 680 vs. 2(78) = 156.

Time = 2.62 (sec) , antiderivative size = 680, normalized size of antiderivative = 8.00

$$\int \frac{(a + b \operatorname{atanh}(c\sqrt{x}))^2}{x^2} dx = \text{Too large to display}$$

input `integrate((a+b*atanh(c*x**(1/2)))**2/x**2,x)`

output

```
Piecewise((-a**2/x, Eq(c, 0)), (-a**2/x + 2*a*b*atanh(sqrt(x)*sqrt(1/x))/x
- b**2*atanh(sqrt(x)*sqrt(1/x))**2/x, Eq(c, -sqrt(1/x))), (-a**2/x - 2*a*
b*atanh(sqrt(x)*sqrt(1/x))/x - b**2*atanh(sqrt(x)*sqrt(1/x))**2/x, Eq(c, s
qrt(1/x))), (-a**2*c**2*x**(3/2)/(c**2*x**(5/2) - x**(3/2)) + a**2*sqrt(x)
/(c**2*x**(5/2) - x**(3/2)) + 2*a*b*c**4*x**(5/2)*atanh(c*sqrt(x))/(c**2*x
**(5/2) - x**(3/2)) - 2*a*b*c**3*x**2/(c**2*x**(5/2) - x**(3/2)) - 4*a*b*c
**2*x**(3/2)*atanh(c*sqrt(x))/(c**2*x**(5/2) - x**(3/2)) + 2*a*b*c*x/(c**2
*x**(5/2) - x**(3/2)) + 2*a*b*sqrt(x)*atanh(c*sqrt(x))/(c**2*x**(5/2) - x*
*(3/2)) + b**2*c**4*x**(5/2)*log(x)/(c**2*x**(5/2) - x**(3/2)) - 2*b**2*c*
**4*x**(5/2)*log(sqrt(x) - 1/c)/(c**2*x**(5/2) - x**(3/2)) + b**2*c**4*x**(
5/2)*atanh(c*sqrt(x))**2/(c**2*x**(5/2) - x**(3/2)) - 2*b**2*c**4*x**(5/2)
*atanh(c*sqrt(x))/(c**2*x**(5/2) - x**(3/2)) - 2*b**2*c**3*x**2*atanh(c*sq
rt(x))/(c**2*x**(5/2) - x**(3/2)) - b**2*c**2*x**(3/2)*log(x)/(c**2*x**(5/
2) - x**(3/2)) + 2*b**2*c**2*x**(3/2)*log(sqrt(x) - 1/c)/(c**2*x**(5/2) -
x**(3/2)) - 2*b**2*c**2*x**(3/2)*atanh(c*sqrt(x))**2/(c**2*x**(5/2) - x**(
3/2)) + 2*b**2*c**2*x**(3/2)*atanh(c*sqrt(x))/(c**2*x**(5/2) - x**(3/2)) +
2*b**2*c*x*atanh(c*sqrt(x))/(c**2*x**(5/2) - x**(3/2)) + b**2*sqrt(x)*ata
nh(c*sqrt(x))**2/(c**2*x**(5/2) - x**(3/2)), True))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 174 vs. $2(77) = 154$.

Time = 0.04 (sec) , antiderivative size = 174, normalized size of antiderivative = 2.05

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x^2} dx$$

$$= \left(\left(c \log(c\sqrt{x} + 1) - c \log(c\sqrt{x} - 1) - \frac{2}{\sqrt{x}} \right) c - \frac{2 \operatorname{arctanh}(c\sqrt{x})}{x} \right) ab$$

$$+ \frac{1}{4} \left(\left(2 (\log(c\sqrt{x} - 1) - 2) \log(c\sqrt{x} + 1) - \log(c\sqrt{x} + 1)^2 - \log(c\sqrt{x} - 1)^2 - 4 \log(c\sqrt{x} - 1) + 4 \right) \right.$$

$$\left. - \frac{b^2 \operatorname{arctanh}(c\sqrt{x})^2}{x} - \frac{a^2}{x} \right)$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))^2/x^2,x, algorithm="maxima")
```

output

```
((c*log(c*sqrt(x) + 1) - c*log(c*sqrt(x) - 1) - 2/sqrt(x))*c - 2*arctanh(c*sqrt(x))/x)*a*b + 1/4*((2*(log(c*sqrt(x) - 1) - 2)*log(c*sqrt(x) + 1) - log(c*sqrt(x) + 1)^2 - log(c*sqrt(x) - 1)^2 - 4*log(c*sqrt(x) - 1) + 4*log(x))*c^2 + 4*(c*log(c*sqrt(x) + 1) - c*log(c*sqrt(x) - 1) - 2/sqrt(x))*c*arctanh(c*sqrt(x)))*b^2 - b^2*arctanh(c*sqrt(x))^2/x - a^2/x
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x^2} dx = \int \frac{(b \operatorname{arctanh}(c\sqrt{x}) + a)^2}{x^2} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))^2/x^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*sqrt(x)) + a)^2/x^2, x)
```

Mupad [B] (verification not implemented)

Time = 4.43 (sec) , antiderivative size = 278, normalized size of antiderivative = 3.27

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x^2} dx &= 2b^2c^2 \ln(\sqrt{x}) - \frac{a^2}{x} - b^2c^2 \ln(c\sqrt{x} - 1) \\ &\quad - b^2c^2 \ln(c\sqrt{x} + 1) + \frac{b^2c^2 \ln(c\sqrt{x} + 1)^2}{4} \\ &\quad + \frac{b^2c^2 \ln(1 - c\sqrt{x})^2}{4} - \frac{b^2 \ln(c\sqrt{x} + 1)^2}{4x} \\ &\quad - \frac{b^2 \ln(1 - c\sqrt{x})^2}{4x} - ab^2c^2 \ln(c\sqrt{x} - 1) \\ &\quad + ab^2c^2 \ln(c\sqrt{x} + 1) - \frac{2abc}{\sqrt{x}} - \frac{ab \ln(c\sqrt{x} + 1)}{x} \\ &\quad + \frac{ab \ln(1 - c\sqrt{x})}{x} - \frac{b^2c^2 \ln(c\sqrt{x} + 1) \ln(1 - c\sqrt{x})}{2} \\ &\quad - \frac{b^2c \ln(c\sqrt{x} + 1)}{\sqrt{x}} + \frac{b^2c \ln(1 - c\sqrt{x})}{\sqrt{x}} \\ &\quad + \frac{b^2 \ln(c\sqrt{x} + 1) \ln(1 - c\sqrt{x})}{2x} \end{aligned}$$

input `int((a + b*atanh(c*x^(1/2)))^2/x^2,x)`

output $2*b^2*c^2*\log(x^{1/2}) - a^2/x - b^2*c^2*\log(c*x^{1/2} - 1) - b^2*c^2*\log(c*x^{1/2} + 1) + (b^2*c^2*\log(c*x^{1/2} + 1)^2)/4 + (b^2*c^2*\log(1 - c*x^{1/2}))^2/4 - (b^2*\log(c*x^{1/2} + 1)^2)/(4*x) - (b^2*\log(1 - c*x^{1/2}))^2/(4*x) - a*b*c^2*\log(c*x^{1/2} - 1) + a*b*c^2*\log(c*x^{1/2} + 1) - (2*a*b*c)/x^{1/2} - (a*b*\log(c*x^{1/2} + 1))/x + (a*b*\log(1 - c*x^{1/2}))/x - (b^2*c^2*\log(c*x^{1/2} + 1)*\log(1 - c*x^{1/2}))/2 - (b^2*c*\log(c*x^{1/2} + 1))/x^{1/2} + (b^2*c*\log(1 - c*x^{1/2}))/x^{1/2} + (b^2*\log(c*x^{1/2} + 1)*\log(1 - c*x^{1/2}))/2$

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.42

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x^2} dx$$

$$= \frac{\operatorname{atanh}(\sqrt{x}c)^2 b^2 c^2 x - \operatorname{atanh}(\sqrt{x}c)^2 b^2 - 2\sqrt{x} \operatorname{atanh}(\sqrt{x}c) b^2 c + 2\operatorname{atanh}(\sqrt{x}c) a b c^2 x - 2\operatorname{atanh}(\sqrt{x}c) a^2}{x}$$

input `int((a+b*atanh(c*x^(1/2)))^2/x^2,x)`

output $(\operatorname{atanh}(\sqrt{x}c)**2*b**2*c**2*x - \operatorname{atanh}(\sqrt{x}c)**2*b**2 - 2*\sqrt{x}*\operatorname{atanh}(\sqrt{x}c)*b**2*c + 2*\operatorname{atanh}(\sqrt{x}c)*a*b*c**2*x - 2*\operatorname{atanh}(\sqrt{x}c)*a*b - 2*\operatorname{atanh}(\sqrt{x}c)*b**2*c**2*x - 2*\sqrt{x}*a*b*c - 2*\log(\sqrt{x}c - 1)*b**2*c**2*x + 2*\log(\sqrt{x}c)*b**2*c**2*x - a**2)/x$

3.200 $\int \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{x^3} dx$

Optimal result	1591
Mathematica [A] (verified)	1592
Rubi [A] (warning: unable to verify)	1592
Maple [B] (verified)	1596
Fricas [A] (verification not implemented)	1597
Sympy [B] (verification not implemented)	1597
Maxima [B] (verification not implemented)	1598
Giac [F]	1599
Mupad [B] (verification not implemented)	1600
Reduce [B] (verification not implemented)	1601

Optimal result

Integrand size = 18, antiderivative size = 133

$$\int \frac{(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{x^3} dx = -\frac{b^2c^2}{6x} - \frac{bc(a + b\operatorname{arctanh}(c\sqrt{x}))}{3x^{3/2}} - \frac{bc^3(a + b\operatorname{arctanh}(c\sqrt{x}))}{\sqrt{x}} + \frac{1}{2}c^4(a + b\operatorname{arctanh}(c\sqrt{x}))^2 - \frac{(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{2x^2} + \frac{2}{3}b^2c^4 \log(x) - \frac{2}{3}b^2c^4 \log(1 - c^2x)$$

output

```
-1/6*b^2*c^2/x-1/3*b*c*(a+b*arctanh(c*x^(1/2)))/x^(3/2)-b*c^3*(a+b*arctanh(c*x^(1/2)))/x^(1/2)+1/2*c^4*(a+b*arctanh(c*x^(1/2)))^2-1/2*(a+b*arctanh(c*x^(1/2)))^2/x^2+2/3*b^2*c^4*ln(x)-2/3*b^2*c^4*ln(-c^2*x+1)
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.34

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x^3} dx = \frac{3a^2 + 2abc\sqrt{x} + b^2c^2x + 6abc^3x^{3/2} + 2b(3a + bc\sqrt{x}(1 + 3c^2x)) \operatorname{arctanh}(c\sqrt{x}) - 3b^2(-1 + c^4x^2) \operatorname{arctanh}(c\sqrt{x})}{x^2}$$

input

```
Integrate[(a + b*ArcTanh[c*Sqrt[x]])^2/x^3,x]
```

output

```
-1/6*(3*a^2 + 2*a*b*c*Sqrt[x] + b^2*c^2*x + 6*a*b*c^3*x^(3/2) + 2*b*(3*a +
b*c*Sqrt[x]*(1 + 3*c^2*x))*ArcTanh[c*Sqrt[x]] - 3*b^2*(-1 + c^4*x^2)*ArcT
anh[c*Sqrt[x]]^2 + b*(3*a + 4*b)*c^4*x^2*Log[1 - c*Sqrt[x]] - 3*a*b*c^4*x^
2*Log[1 + c*Sqrt[x]] + 4*b^2*c^4*x^2*Log[1 + c*Sqrt[x]] - 4*b^2*c^4*x^2*Lo
g[x])/x^2
```

Rubi [A] (warning: unable to verify)

Time = 1.04 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.14, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.778$, Rules used = {6454, 6452, 6544, 6452, 243, 54, 2009, 6544, 6452, 243, 47, 14, 16, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x^3} dx \\ & \quad \downarrow \text{6454} \\ & 2 \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x^{5/2}} d\sqrt{x} \\ & \quad \downarrow \text{6452} \\ & 2 \left(\frac{1}{2} bc \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(1 - c^2x)} d\sqrt{x} - \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{4x^2} \right) \end{aligned}$$

↓ 6544

$$2 \left(\frac{1}{2} bc \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x(1-c^2x)} d\sqrt{x} + \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^2} d\sqrt{x} \right) - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{4x^2} \right)$$

↓ 6452

$$2 \left(\frac{1}{2} bc \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x(1-c^2x)} d\sqrt{x} + \frac{1}{3} bc \int \frac{1}{x^{3/2}(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{3x^{3/2}} \right) - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{4x^2} \right)$$

↓ 243

$$2 \left(\frac{1}{2} bc \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x(1-c^2x)} d\sqrt{x} + \frac{1}{6} bc \int \frac{1}{x(1-c^2x)} dx - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{3x^{3/2}} \right) - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{4x^2} \right)$$

↓ 54

$$2 \left(\frac{1}{2} bc \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x(1-c^2x)} d\sqrt{x} + \frac{1}{6} bc \int \left(-\frac{c^4}{c^2x-1} + \frac{c^2}{\sqrt{x}} + \frac{1}{x} \right) dx - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{3x^{3/2}} \right) - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{4x^2} \right)$$

↓ 2009

$$2 \left(\frac{1}{2} bc \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{3x^{3/2}} + \frac{1}{6} bc \left(c^2 \log(x) - c^2 \log(1-c^2x) - \frac{1}{\sqrt{x}} \right) \right) - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{4x^2} \right)$$

↓ 6544

$$2 \left(\frac{1}{2} bc \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{1-c^2x} d\sqrt{x} + \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x} d\sqrt{x} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{3x^{3/2}} + \frac{1}{6} bc \left(c^2 \log(x) - c^2 \log(1-c^2x) - \frac{1}{\sqrt{x}} \right) \right) - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{4x^2} \right)$$

↓ 6452

$$2 \left(\frac{1}{2} bc \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{1-c^2x} d\sqrt{x} + bc \int \frac{1}{\sqrt{x}(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{3x^{3/2}} \right) - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{4x^2} \right)$$

↓ 243

$$2 \left(\frac{1}{2} bc \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{1-c^2x} d\sqrt{x} + \frac{1}{2} bc \int \frac{1}{\sqrt{x}(1-c^2x)} dx - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{3x^{3/2}} \right) - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{4x^2} \right)$$

$$\begin{aligned}
& \downarrow 47 \\
& 2 \left(\frac{1}{2} bc \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{1 - c^2 x} d\sqrt{x} + \frac{1}{2} bc \left(c^2 \int \frac{1}{1 - c^2 x} dx + \int \frac{1}{\sqrt{x}} dx \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}} \right) \\
& \downarrow 14 \\
& 2 \left(\frac{1}{2} bc \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{1 - c^2 x} d\sqrt{x} + \frac{1}{2} bc \left(c^2 \int \frac{1}{1 - c^2 x} dx + \log(x) \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}} \right) \\
& \downarrow 16 \\
& 2 \left(\frac{1}{2} bc \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{1 - c^2 x} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}} + \frac{1}{2} bc (\log(x) - \log(1 - c^2 x)) \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}} \right) \\
& \downarrow 6510 \\
& 2 \left(\frac{1}{2} bc \left(c^2 \left(\frac{c(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}} + \frac{1}{2} bc (\log(x) - \log(1 - c^2 x)) \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}} \right) \right)
\end{aligned}$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])^2/x^3,x]`

output `2*(-1/4*(a + b*ArcTanh[c*Sqrt[x]])^2/x^2 + (b*c*(-1/3*(a + b*ArcTanh[c*Sqrt[x]])/x^(3/2) + c^2*(-((a + b*ArcTanh[c*Sqrt[x]])/Sqrt[x]) + (c*(a + b*ArcTanh[c*Sqrt[x]])^2)/(2*b) + (b*c*(Log[x] - Log[1 - c^2*x]))/2) + (b*c*(-(1/Sqrt[x]) + c^2*Log[x] - c^2*Log[1 - c^2*x]))/6))/2`

Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

- rule 47 $\text{Int}[1/((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.)), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x]$
- rule 54 $\text{Int}(((a_.) + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol) \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x]$ && $\text{ILtQ}[m, 0]$ && $\text{IntegerQ}[n]$ && $!(\text{IGtQ}[n, 0] \text{ \&\& } \text{LtQ}[m + n + 2, 0])$
- rule 243 $\text{Int}[(x_.)^{(m_.)}*((a_.) + (b_.)*(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] /;$ $\text{FreeQ}\{a, b, m, p\}, x]$ && $\text{IntegerQ}[(m-1)/2]$
- rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /;$ $\text{SumQ}[u]$
- rule 6452 $\text{Int}(((a_.) + \text{ArcTanh}[(c_.)*(x_.)^{(n_.)}]*(b_.))^{(p_.)}*(x_.)^{(m_.)}, x_Symbol) \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{(2*n)})}), x], x] /;$ $\text{FreeQ}\{a, b, c, m, n\}, x]$ && $\text{IGtQ}[p, 0]$ && $(\text{EqQ}[p, 1] \text{ || } (\text{EqQ}[n, 1] \text{ \&\& } \text{IntegerQ}[m]))$ && $\text{NeQ}[m, -1]$
- rule 6454 $\text{Int}(((a_.) + \text{ArcTanh}[(c_.)*(x_.)^{(n_.)}]*(b_.))^{(p_.)}*(x_.)^{(m_.)}, x_Symbol) \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n] - 1)*(a + b*\text{ArcTanh}[c*x])^p}, x], x, x^n], x] /;$ $\text{FreeQ}\{a, b, c, m, n\}, x]$ && $\text{IGtQ}[p, 1]$ && $\text{IntegerQ}[\text{Simplify}[(m+1)/n]]$
- rule 6510 $\text{Int}(((a_.) + \text{ArcTanh}[(c_.)*(x_.)]*(b_.))^{(p_.)}/((d_.) + (e_.)*(x_.)^2), x_Symbol) \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^{(p+1)}/(b*c*d*(p+1)), x] /;$ $\text{FreeQ}\{a, b, c, d, e, p\}, x]$ && $\text{EqQ}[c^2*d + e, 0]$ && $\text{NeQ}[p, -1]$

rule 6544

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*((f_.)*(x_))^(m_)/((d_) + (
e_.)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x
], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x
^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 262 vs. 2(109) = 218.

Time = 0.45 (sec) , antiderivative size = 263, normalized size of antiderivative = 1.98

method	result
parts	$-\frac{a^2}{2x^2} + 2b^2c^4 \left(-\frac{\operatorname{arctanh}(c\sqrt{x})^2}{4c^4x^2} - \frac{\operatorname{arctanh}(c\sqrt{x})}{6c^3x^{\frac{3}{2}}} - \frac{\operatorname{arctanh}(c\sqrt{x})}{2c\sqrt{x}} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1+c\sqrt{x})}{4} - \frac{\operatorname{arctanh}(c\sqrt{x})}{4} \right)$
derivativedivides	$2c^4 \left(-\frac{a^2}{4c^4x^2} + b^2 \left(-\frac{\operatorname{arctanh}(c\sqrt{x})^2}{4c^4x^2} - \frac{\operatorname{arctanh}(c\sqrt{x})}{6c^3x^{\frac{3}{2}}} - \frac{\operatorname{arctanh}(c\sqrt{x})}{2c\sqrt{x}} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1+c\sqrt{x})}{4} - \frac{\operatorname{arctanh}(c\sqrt{x})}{4} \right) \right)$
default	$2c^4 \left(-\frac{a^2}{4c^4x^2} + b^2 \left(-\frac{\operatorname{arctanh}(c\sqrt{x})^2}{4c^4x^2} - \frac{\operatorname{arctanh}(c\sqrt{x})}{6c^3x^{\frac{3}{2}}} - \frac{\operatorname{arctanh}(c\sqrt{x})}{2c\sqrt{x}} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1+c\sqrt{x})}{4} - \frac{\operatorname{arctanh}(c\sqrt{x})}{4} \right) \right)$

input

```
int((a+b*arctanh(c*x^(1/2)))^2/x^3,x,method=_RETURNVERBOSE)
```

output

```
-1/2/x^2*a^2+2*b^2*c^4*(-1/4/c^4/x^2*arctanh(c*x^(1/2))^2-1/6*arctanh(c*x^(1/2))/c^3/x^(3/2)-1/2*arctanh(c*x^(1/2))/c/x^(1/2)+1/4*arctanh(c*x^(1/2))*ln(1+c*x^(1/2))-1/4*arctanh(c*x^(1/2))*ln(c*x^(1/2)-1)-1/16*ln(c*x^(1/2)-1)^2+1/8*ln(c*x^(1/2)-1)*ln(1/2*c*x^(1/2)+1/2)+1/8*(ln(1+c*x^(1/2))-ln(1/2*c*x^(1/2)+1/2))*ln(-1/2*c*x^(1/2)+1/2)-1/16*ln(1+c*x^(1/2))^2-1/12/c^2/x+2/3*ln(c*x^(1/2))-1/3*ln(1+c*x^(1/2))-1/3*ln(c*x^(1/2)-1))+4*a*b*c^4*(-1/4/c^4/x^2*arctanh(c*x^(1/2))-1/12/c^3/x^(3/2)-1/4/c/x^(1/2)+1/8*ln(1+c*x^(1/2))-1/8*ln(c*x^(1/2)-1))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.51

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x^3} dx$$

$$= \frac{32 b^2 c^4 x^2 \log(\sqrt{x}) + 4(3ab - 4b^2)c^4 x^2 \log(c\sqrt{x} + 1) - 4(3ab + 4b^2)c^4 x^2 \log(c\sqrt{x} - 1) - 4b^2 c^2 x + 3}{x^3}$$

input `integrate((a+b*arctanh(c*x^(1/2)))^2/x^3,x, algorithm="fricas")`

output `1/24*(32*b^2*c^4*x^2*log(sqrt(x)) + 4*(3*a*b - 4*b^2)*c^4*x^2*log(c*sqrt(x) + 1) - 4*(3*a*b + 4*b^2)*c^4*x^2*log(c*sqrt(x) - 1) - 4*b^2*c^2*x + 3*(b^2*c^4*x^2 - b^2)*log(-(c^2*x + 2*c*sqrt(x) + 1)/(c^2*x - 1))^2 - 12*a^2 - 4*(3*a*b + (3*b^2*c^3*x + b^2*c)*sqrt(x))*log(-(c^2*x + 2*c*sqrt(x) + 1)/(c^2*x - 1)) - 8*(3*a*b*c^3*x + a*b*c)*sqrt(x))/x^2`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 972 vs. 2(122) = 244.

Time = 7.44 (sec) , antiderivative size = 972, normalized size of antiderivative = 7.31

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x^3} dx = \text{Too large to display}$$

input `integrate((a+b*atanh(c*x**(1/2)))**2/x**3,x)`

output

```
Piecewise((-a**2/(2*x**2), Eq(c, 0)), (-a**2/(2*x**2) + a*b*atanh(sqrt(x)*
sqrt(1/x))/x**2 - b**2*atanh(sqrt(x)*sqrt(1/x))**2/(2*x**2), Eq(c, -sqrt(1
/x))), (-a**2/(2*x**2) - a*b*atanh(sqrt(x)*sqrt(1/x))/x**2 - b**2*atanh(sq
rt(x)*sqrt(1/x))**2/(2*x**2), Eq(c, sqrt(1/x))), (-3*a**2*c**2*x**(3/2)/(6
*c**2*x**(7/2) - 6*x**(5/2)) + 3*a**2*sqrt(x)/(6*c**2*x**(7/2) - 6*x**(5/2
)) + 6*a*b*c**6*x**(7/2)*atanh(c*sqrt(x))/(6*c**2*x**(7/2) - 6*x**(5/2)) -
6*a*b*c**5*x**3/(6*c**2*x**(7/2) - 6*x**(5/2)) - 6*a*b*c**4*x**(5/2)*atan
h(c*sqrt(x))/(6*c**2*x**(7/2) - 6*x**(5/2)) + 4*a*b*c**3*x**2/(6*c**2*x**(
7/2) - 6*x**(5/2)) - 6*a*b*c**2*x**(3/2)*atanh(c*sqrt(x))/(6*c**2*x**(7/2)
- 6*x**(5/2)) + 2*a*b*c*x/(6*c**2*x**(7/2) - 6*x**(5/2)) + 6*a*b*sqrt(x)*
atanh(c*sqrt(x))/(6*c**2*x**(7/2) - 6*x**(5/2)) + 4*b**2*c**6*x**(7/2)*log
(x)/(6*c**2*x**(7/2) - 6*x**(5/2)) - 8*b**2*c**6*x**(7/2)*log(sqrt(x) - 1/
c)/(6*c**2*x**(7/2) - 6*x**(5/2)) + 3*b**2*c**6*x**(7/2)*atanh(c*sqrt(x))*
*2/(6*c**2*x**(7/2) - 6*x**(5/2)) - 8*b**2*c**6*x**(7/2)*atanh(c*sqrt(x))/
(6*c**2*x**(7/2) - 6*x**(5/2)) - 6*b**2*c**5*x**3*atanh(c*sqrt(x))/(6*c**2
*x**(7/2) - 6*x**(5/2)) - 4*b**2*c**4*x**(5/2)*log(x)/(6*c**2*x**(7/2) - 6
*x**(5/2)) + 8*b**2*c**4*x**(5/2)*log(sqrt(x) - 1/c)/(6*c**2*x**(7/2) - 6*
x**(5/2)) - 3*b**2*c**4*x**(5/2)*atanh(c*sqrt(x))**2/(6*c**2*x**(7/2) - 6*
x**(5/2)) + 8*b**2*c**4*x**(5/2)*atanh(c*sqrt(x))/(6*c**2*x**(7/2) - 6*x**
(5/2)) - b**2*c**4*x**(5/2)/(6*c**2*x**(7/2) - 6*x**(5/2)) + 4*b**2*c**...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 234 vs. $2(109) = 218$.

Time = 0.04 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.76

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x^3} dx$$

$$= \frac{1}{6} \left(\left(3c^3 \log(c\sqrt{x} + 1) - 3c^3 \log(c\sqrt{x} - 1) - \frac{2(3c^2x + 1)}{x^{\frac{3}{2}}} \right) c - \frac{6 \operatorname{artanh}(c\sqrt{x})}{x^2} \right) ab$$

$$+ \frac{1}{24} \left(\frac{16c^2 \log(x) - \frac{3c^2x \log(c\sqrt{x} + 1)^2 + 3c^2x \log(c\sqrt{x} - 1)^2 + 16c^2x \log(c\sqrt{x} - 1) - 2(3c^2x \log(c\sqrt{x} + 1) - 3c^2x \log(c\sqrt{x} - 1))}{x} - \frac{b^2 \operatorname{artanh}(c\sqrt{x})^2}{2x^2} - \frac{a^2}{2x^2} \right)$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))^2/x^3,x, algorithm="maxima")
```

output

```
1/6*((3*c^3*log(c*sqrt(x) + 1) - 3*c^3*log(c*sqrt(x) - 1) - 2*(3*c^2*x + 1)
)/x^(3/2))*c - 6*arctanh(c*sqrt(x))/x^2)*a*b + 1/24*((16*c^2*log(x) - (3*c
^2*x*log(c*sqrt(x) + 1)^2 + 3*c^2*x*log(c*sqrt(x) - 1)^2 + 16*c^2*x*log(c*
sqrt(x) - 1) - 2*(3*c^2*x*log(c*sqrt(x) - 1) - 8*c^2*x)*log(c*sqrt(x) + 1)
+ 4)/x)*c^2 + 4*(3*c^3*log(c*sqrt(x) + 1) - 3*c^3*log(c*sqrt(x) - 1) - 2*
(3*c^2*x + 1)/x^(3/2))*c*arctanh(c*sqrt(x)))*b^2 - 1/2*b^2*arctanh(c*sqrt(
x))^2/x^2 - 1/2*a^2/x^2
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x^3} dx = \int \frac{(b \operatorname{arctanh}(c\sqrt{x}) + a)^2}{x^3} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))^2/x^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*sqrt(x)) + a)^2/x^3, x)
```

Mupad [B] (verification not implemented)

Time = 5.05 (sec) , antiderivative size = 341, normalized size of antiderivative = 2.56

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x^3} dx = \frac{4b^2 c^4 \ln(\sqrt{x})}{3} - \frac{a^2}{2x^2} - \frac{2b^2 c^4 \ln(c\sqrt{x} - 1)}{3}$$

$$- \frac{2b^2 c^4 \ln(c\sqrt{x} + 1)}{3} + \frac{b^2 c^4 \ln(c\sqrt{x} + 1)^2}{8}$$

$$+ \frac{b^2 c^4 \ln(1 - c\sqrt{x})^2}{8} - \frac{b^2 c^2}{6x} - \frac{b^2 \ln(c\sqrt{x} + 1)^2}{8x^2}$$

$$- \frac{b^2 \ln(1 - c\sqrt{x})^2}{8x^2} - \frac{b^2 c^3 \ln(c\sqrt{x} + 1)}{2\sqrt{x}}$$

$$+ \frac{b^2 c^3 \ln(1 - c\sqrt{x})}{2\sqrt{x}} - \frac{ab c^4 \ln(c\sqrt{x} - 1)}{2}$$

$$+ \frac{ab c^4 \ln(c\sqrt{x} + 1)}{2} - \frac{abc}{3x^{3/2}} - \frac{ab \ln(c\sqrt{x} + 1)}{2x^2}$$

$$+ \frac{ab \ln(1 - c\sqrt{x})}{2x^2} - \frac{b^2 c^4 \ln(c\sqrt{x} + 1) \ln(1 - c\sqrt{x})}{4}$$

$$- \frac{ab c^3}{\sqrt{x}} - \frac{b^2 c \ln(c\sqrt{x} + 1)}{6x^{3/2}} + \frac{b^2 c \ln(1 - c\sqrt{x})}{6x^{3/2}}$$

$$+ \frac{b^2 \ln(c\sqrt{x} + 1) \ln(1 - c\sqrt{x})}{4x^2}$$

input `int((a + b*atanh(c*x^(1/2)))^2/x^3,x)`output

```
(4*b^2*c^4*log(x^(1/2)))/3 - a^2/(2*x^2) - (2*b^2*c^4*log(c*x^(1/2) - 1))/3 - (2*b^2*c^4*log(c*x^(1/2) + 1))/3 + (b^2*c^4*log(c*x^(1/2) + 1)^2)/8 + (b^2*c^4*log(1 - c*x^(1/2))^2)/8 - (b^2*c^2)/(6*x) - (b^2*log(c*x^(1/2) + 1)^2)/(8*x^2) - (b^2*log(1 - c*x^(1/2))^2)/(8*x^2) - (b^2*c^3*log(c*x^(1/2) + 1))/(2*x^(1/2)) + (b^2*c^3*log(1 - c*x^(1/2)))/(2*x^(1/2)) - (a*b*c^4*log(c*x^(1/2) - 1))/2 + (a*b*c^4*log(c*x^(1/2) + 1))/2 - (a*b*c)/(3*x^(3/2)) - (a*b*log(c*x^(1/2) + 1))/(2*x^2) + (a*b*log(1 - c*x^(1/2)))/(2*x^2) - (b^2*c^4*log(c*x^(1/2) + 1)*log(1 - c*x^(1/2)))/4 - (a*b*c^3)/x^(1/2) - (b^2*c*log(c*x^(1/2) + 1))/(6*x^(3/2)) + (b^2*c*log(1 - c*x^(1/2)))/(6*x^(3/2)) + (b^2*log(c*x^(1/2) + 1)*log(1 - c*x^(1/2)))/(4*x^2)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.26

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x^3} dx$$

$$= \frac{3 \operatorname{atanh}(\sqrt{x}c)^2 b^2 c^4 x^2 - 3 \operatorname{atanh}(\sqrt{x}c)^2 b^2 - 6\sqrt{x} \operatorname{atanh}(\sqrt{x}c) b^2 c^3 x - 2\sqrt{x} \operatorname{atanh}(\sqrt{x}c) b^2 c + 6 \operatorname{atanh}(\sqrt{x}c) a b c^3 x - 2 \operatorname{atanh}(\sqrt{x}c) a b c + 6 \operatorname{atanh}(\sqrt{x}c) a^2 c^3 x - 2 \operatorname{atanh}(\sqrt{x}c) a^2 c + 6 \operatorname{atanh}(\sqrt{x}c) a^2 c^3 x - 2 \operatorname{atanh}(\sqrt{x}c) a^2 c}{6x^3}$$

input

```
int((a+b*atanh(c*x^(1/2)))^2/x^3,x)
```

output

```
(3*atanh(sqrt(x)*c)**2*b**2*c**4*x**2 - 3*atanh(sqrt(x)*c)**2*b**2 - 6*sqrt(x)*atanh(sqrt(x)*c)*b**2*c**3*x - 2*sqrt(x)*atanh(sqrt(x)*c)*b**2*c + 6*atanh(sqrt(x)*c)*a*b*c**4*x**2 - 6*atanh(sqrt(x)*c)*a*b - 8*atanh(sqrt(x)*c)*b**2*c**4*x**2 - 6*sqrt(x)*a*b*c**3*x - 2*sqrt(x)*a*b*c - 8*log(sqrt(x)*c - 1)*b**2*c**4*x**2 + 8*log(sqrt(x))*b**2*c**4*x**2 - 3*a**2 - b**2*c**2*x)/(6*x**2)
```

3.201 $\int x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx$

Optimal result	1602
Mathematica [A] (verified)	1603
Rubi [B] (verified)	1604
Maple [C] (warning: unable to verify)	1613
Fricas [F]	1614
Sympy [F]	1614
Maxima [B] (verification not implemented)	1614
Giac [F]	1615
Mupad [F(-1)]	1616
Reduce [F]	1616

Optimal result

Integrand size = 18, antiderivative size = 374

$$\begin{aligned}
 & \int x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx \\
 &= \frac{47b^3\sqrt{x}}{70c^7} + \frac{23b^3x^{3/2}}{420c^5} + \frac{b^3x^{5/2}}{140c^3} - \frac{47b^3\operatorname{arctanh}(c\sqrt{x})}{70c^8} + \frac{71b^2x(a + b\operatorname{arctanh}(c\sqrt{x}))}{140c^6} \\
 &+ \frac{9b^2x^2(a + b\operatorname{arctanh}(c\sqrt{x}))}{70c^4} + \frac{b^2x^3(a + b\operatorname{arctanh}(c\sqrt{x}))}{28c^2} \\
 &+ \frac{44b(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{35c^8} + \frac{3b\sqrt{x}(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{4c^7} \\
 &+ \frac{bx^{3/2}(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{4c^5} + \frac{3bx^{5/2}(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{20c^3} \\
 &+ \frac{3bx^{7/2}(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{28c} - \frac{(a + b\operatorname{arctanh}(c\sqrt{x}))^3}{4c^8} \\
 &+ \frac{1}{4}x^4(a + b\operatorname{arctanh}(c\sqrt{x}))^3 - \frac{88b^2(a + b\operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1-c\sqrt{x}}\right)}{35c^8} - \frac{44b^3 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-c\sqrt{x}}\right)}{35c^8}
 \end{aligned}$$

output

$$\frac{47/70*b^3*x^{(1/2)}/c^7+23/420*b^3*x^{(3/2)}/c^5+1/140*b^3*x^{(5/2)}/c^3-47/70*b^3*\operatorname{arctanh}(c*x^{(1/2)})/c^8+71/140*b^2*x*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))/c^6+9/70*b^2*x^2*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))/c^4+1/28*b^2*x^3*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))/c^2+44/35*b*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))^2/c^8+3/4*b*x^{(1/2)}*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))^2/c^7+1/4*b*x^{(3/2)}*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))^2/c^5+3/20*b*x^{(5/2)}*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))^2/c^3+3/28*b*x^{(7/2)}*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))^2/c-1/4*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))^3/c^8+1/4*x^4*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))^3-88/35*b^2*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))*\ln(2/(1-c*x^{(1/2)}))/c^8-44/35*b^3*\operatorname{polylog}(2,1-2/(1-c*x^{(1/2)}))/c^8}{c^8}$$
Mathematica [A] (verified)

Time = 0.75 (sec) , antiderivative size = 418, normalized size of antiderivative = 1.12

$$\int x^3(a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx$$

$$= \frac{-564ab^2 + 630a^2bc\sqrt{x} + 564b^3c\sqrt{x} + 426ab^2c^2x + 210a^2bc^3x^{3/2} + 46b^3c^3x^{3/2} + 108ab^2c^4x^2 + 126a^2bc^5x^2}{c^8}$$

input

`Integrate[x^3*(a + b*ArcTanh[c*Sqrt[x]])^3,x]`

output

$$\frac{(-564*a*b^2 + 630*a^2*b*c*\operatorname{Sqrt}[x] + 564*b^3*c*\operatorname{Sqrt}[x] + 426*a*b^2*c^2*x + 210*a^2*b*c^3*x^{(3/2)} + 46*b^3*c^3*x^{(3/2)} + 108*a*b^2*c^4*x^2 + 126*a^2*b*c^5*x^{(5/2)} + 6*b^3*c^5*x^{(5/2)} + 30*a*b^2*c^6*x^3 + 90*a^2*b*c^7*x^{(7/2)} + 210*a^3*c^8*x^4 + 6*b^2*(b*(-176 + 105*c*\operatorname{Sqrt}[x] + 35*c^3*x^{(3/2)} + 21*c^5*x^{(5/2)} + 15*c^7*x^{(7/2)})) + 105*a*(-1 + c^8*x^4))*\operatorname{ArcTanh}[c*\operatorname{Sqrt}[x]]^2 + 210*b^3*(-1 + c^8*x^4)*\operatorname{ArcTanh}[c*\operatorname{Sqrt}[x]]^3 + 6*b*\operatorname{ArcTanh}[c*\operatorname{Sqrt}[x]]*(105*a^2*c^8*x^4 + b^2*(-94 + 71*c^2*x + 18*c^4*x^2 + 5*c^6*x^3) + 2*a*b*c*\operatorname{Sqrt}[x]*(105 + 35*c^2*x + 21*c^4*x^2 + 15*c^6*x^3) - 352*b^2*\operatorname{Log}[1 + E^{(-2*\operatorname{ArcTanh}[c*\operatorname{Sqrt}[x]])}]) + 315*a^2*b*\operatorname{Log}[1 - c*\operatorname{Sqrt}[x]] - 315*a^2*b*\operatorname{Log}[1 + c*\operatorname{Sqrt}[x]] + 1056*a*b^2*\operatorname{Log}[1 - c^2*x] + 1056*b^3*\operatorname{PolyLog}[2, -E^{(-2*\operatorname{ArcTanh}[c*\operatorname{Sqrt}[x]])}])]/(840*c^8)$$

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 949 vs. $2(374) = 748$.

Time = 5.93 (sec) , antiderivative size = 949, normalized size of antiderivative = 2.54, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 1.222$, Rules used = {6454, 6452, 6542, 6452, 6542, 6452, 254, 2009, 6542, 6452, 254, 2009, 6542, 6436, 6452, 262, 219, 6510, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^3 dx \\
 & \quad \downarrow \text{6454} \\
 & 2 \int x^{7/2} (a + \operatorname{barctanh}(c\sqrt{x}))^3 d\sqrt{x} \\
 & \quad \downarrow \text{6452} \\
 & 2 \left(\frac{1}{8} x^4 (a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \int \frac{x^4 (a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x} \right) \\
 & \quad \downarrow \text{6542} \\
 & 2 \left(\frac{1}{8} x^4 (a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\int x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^2 d\sqrt{x}}{c^2} \right) \right) \\
 & \quad \downarrow \text{6452} \\
 & 2 \left(\frac{1}{8} x^4 (a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \left(\frac{\int \frac{x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{7} x^{7/2} (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{2}{7} bc \int \frac{x^{7/2}}{c^2}}{c^2} \right) \right) \\
 & \quad \downarrow \text{6542}
 \end{aligned}$$

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \left(\frac{\int \frac{x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\int x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 d\sqrt{x}}{c^2} - \frac{\frac{1}{7} x^{7/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} \right) \right)$$

↓ 6452

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \left(\frac{\int \frac{x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{2}{5} bc \int \frac{x^{5/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} \right) \right)$$

↓ 254

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \left(\frac{\int \frac{x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{2}{5} bc \int \frac{x^{5/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} \right) \right)$$

↓ 2009

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \left(\frac{\int \frac{x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{2}{5} bc \int \frac{x^{5/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} \right) \right)$$

↓ 6542

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \int \frac{x (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x} - \frac{\int x (a + b \operatorname{arctanh}(c\sqrt{x}))^2 d\sqrt{x}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} \right)$$

↓ 6452

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \int \frac{x (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x} - \frac{\frac{1}{3} x^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{2}{3} bc \int \frac{x^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x}}{c^2} \right)$$

↓ 254

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \int \frac{x (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x} - \frac{\frac{1}{3} x^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{2}{3} bc \int \frac{x^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x}}{c^2} \right)$$

↓ 2009

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \int \frac{x (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x} - \frac{\frac{1}{3} x^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{2}{3} bc \int \frac{x^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x}}{c^2} \right)$$

↓ 6542

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \right) \left(\frac{\int \frac{(a+b \operatorname{arctanh}(c\sqrt{x}))^2}{1-c^2 x} d\sqrt{x}}{c^2} - \frac{\int (a+b \operatorname{arctanh}(c\sqrt{x}))^2 d\sqrt{x}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a+b \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} \right)$$

↓ 6436

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \right) \left(\frac{\int \frac{(a+b \operatorname{arctanh}(c\sqrt{x}))^2}{1-c^2 x} d\sqrt{x}}{c^2} - \frac{\sqrt{x} (a+b \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} - 2bc \int \frac{\sqrt{x} (a+b \operatorname{arctanh}(c\sqrt{x}))}{1-c^2 x} d\sqrt{x} \right)$$

↓ 6452

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \right) \left(\frac{\int \frac{(a+b \operatorname{arctanh}(c\sqrt{x}))^2}{1-c^2 x} d\sqrt{x}}{c^2} - \frac{\sqrt{x} (a+b \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} - 2bc \int \frac{\sqrt{x} (a+b \operatorname{arctanh}(c\sqrt{x}))}{1-c^2 x} d\sqrt{x} \right)$$

↓ 262

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \right) \left(\frac{\int \frac{(a+b \operatorname{arctanh}(c\sqrt{x}))^2}{1-c^2 x} d\sqrt{x}}{c^2} - \frac{\sqrt{x} (a+b \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} - 2bc \int \frac{\sqrt{x} (a+b \operatorname{arctanh}(c\sqrt{x}))}{1-c^2 x} d\sqrt{x}}{c^2} \right)$$

↓ 219

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \right) \left(\frac{\int \frac{(a+b \operatorname{arctanh}(c\sqrt{x}))^2}{1-c^2 x} d\sqrt{x}}{c^2} - \frac{\sqrt{x} (a+b \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} - 2bc \int \frac{\sqrt{x} (a+b \operatorname{arctanh}(c\sqrt{x}))}{1-c^2 x} d\sqrt{x}}{c^2} \right)$$

↓ 6510

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \right) \left(\frac{(a+b \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x} (a+b \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} - 2bc \int \frac{\sqrt{x} (a+b \operatorname{arctanh}(c\sqrt{x}))}{1-c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{3} x^{\frac{5}{2}}}{c^2} \right)$$

↓ 6546

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \right) \left(\frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{1 - c\sqrt{x}} d\sqrt{x} - (a + b \operatorname{arctanh}(c\sqrt{x})) \right)}{c^2}}{c^2} \right)$$

↓ 6470

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \right) \left(\frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \left(\frac{(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1 - c\sqrt{x}}\right)}{c} \right)}{c^2}}{c^2} \right)$$

↓ 2849

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \right) \left(\frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \left(\frac{(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1 - c\sqrt{x}}\right)}{c} \right)}{c^2}}{c^2} \right)$$

↓ 2752

$$2 \left(\frac{1}{8} x^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{8} bc \right) \left(\frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc}{c^2} \left(\frac{(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1 - c\sqrt{x}}\right)}{c} \right) \right)$$

input `Int [x^3*(a + b*ArcTanh[c*Sqrt[x]])^3,x]`

output

```

2*((x^4*(a + b*ArcTanh[c*Sqrt[x]])^3)/8 - (3*b*c*(-((x^(7/2)*(a + b*ArcTanh[c*Sqrt[x]])^2)/7 - (2*b*c*(-((x^3*(a + b*ArcTanh[c*Sqrt[x]]))/6 - (b*c*(-(Sqrt[x]/c^6) - x^(3/2)/(3*c^4) - x^(5/2)/(5*c^2) + ArcTanh[c*Sqrt[x]]/c^7))/6)/c^2) + (-((x^2*(a + b*ArcTanh[c*Sqrt[x]]))/4 - (b*c*(-(Sqrt[x]/c^4) - x^(3/2)/(3*c^2) + ArcTanh[c*Sqrt[x]]/c^5))/4)/c^2) + (-((x*(a + b*ArcTanh[c*Sqrt[x]]))/2 - (b*c*(-(Sqrt[x]/c^2) + ArcTanh[c*Sqrt[x]]/c^3))/2)/c^2) + (-1/2*(a + b*ArcTanh[c*Sqrt[x]])^2/(b*c^2) + ((a + b*ArcTanh[c*Sqrt[x]])*Log[2/(1 - c*Sqrt[x]])]/c + (b*PolyLog[2, 1 - 2/(1 - c*Sqrt[x]])]/(2*c))/c)/c^2)/7)/c^2) + (-((x^(5/2)*(a + b*ArcTanh[c*Sqrt[x]])^2)/5 - (2*b*c*(-((x^2*(a + b*ArcTanh[c*Sqrt[x]]))/4 - (b*c*(-(Sqrt[x]/c^4) - x^(3/2)/(3*c^2) + ArcTanh[c*Sqrt[x]]/c^5))/4)/c^2) + (-((x*(a + b*ArcTanh[c*Sqrt[x]]))/2 - (b*c*(-(Sqrt[x]/c^2) + ArcTanh[c*Sqrt[x]]/c^3))/2)/c^2) + (-1/2*(a + b*ArcTanh[c*Sqrt[x]])^2/(b*c^2) + ((a + b*ArcTanh[c*Sqrt[x]])*Log[2/(1 - c*Sqrt[x]])]/c + (b*PolyLog[2, 1 - 2/(1 - c*Sqrt[x]])]/(2*c))/c)/c^2)/5)/c^2) + (-((x^(3/2)*(a + b*ArcTanh[c*Sqrt[x]])^2)/3 - (2*b*c*(-((x*(a + b*ArcTanh[c*Sqrt[x]]))/2 - (b*c*(-(Sqrt[x]/c^2) + ArcTanh[c*Sqrt[x]]/c^3))/2)/c^2) + (-1/2*(a + b*ArcTanh[c*Sqrt[x]])^2/(b*c^2) + ((a + b*ArcTanh[c*Sqrt[x]])*Log[2/(1 - c*Sqrt[x]])]/c + (b*PolyLog[2, 1 - 2/(1 - c*Sqrt[x]])]/(2*c))/c)/c^2)/3)/c^2) + ((a + b*ArcTanh[c*Sqrt[x]])^3/(3*b*c^3) - (Sqrt[x]*(a + b*ArcTanh[c*Sqrt[x]])^2 - 2*b*c*(-1/2...
    
```

Definitions of rubi rules used

rule 219 $\text{Int}[(a_+) + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))* \text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 254 $\text{Int}[(x_+)^{m_+}/((a_+) + (b_+)(x_+)^2), x_Symbol] \rightarrow \text{Int}[\text{PolynomialDivide}[x^m, a + b*x^2, x], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{IGtQ}[m, 3]$

rule 262 $\text{Int}[(c_+)(x_+)^{m_+}((a_+) + (b_+)(x_+)^2)^{p_+}, x_Symbol] \rightarrow \text{Simp}[c*(c*x)^{m-1}*((a + b*x^2)^{p+1}/(b*(m+2*p+1))), x] - \text{Simp}[a*c^2*((m-1)/(b*(m+2*p+1))) \ \text{Int}[(c*x)^{m-2}*(a + b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{GtQ}[m, 2-1] \ \&\& \ \text{NeQ}[m+2*p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 2009 $\text{Int}[u_+, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 2752 $\text{Int}[\text{Log}[(c_+)(x_+)]/((d_+) + (e_+)(x_+)), x_Symbol] \rightarrow \text{Simp}[(-e^{-1})*\text{PolyLog}[2, 1 - c*x], x] /; \text{FreeQ}\{c, d, e\}, x \ \&\& \ \text{EqQ}[e + c*d, 0]$

rule 2849 $\text{Int}[\text{Log}[(c_+)/((d_+) + (e_+)(x_+))]/((f_+) + (g_+)(x_+)^2), x_Symbol] \rightarrow \text{Simp}[-e/g \ \text{Subst}[\text{Int}[\text{Log}[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; \text{FreeQ}\{c, d, e, f, g\}, x \ \&\& \ \text{EqQ}[c, 2*d] \ \&\& \ \text{EqQ}[e^2*f + d^2*g, 0]$

rule 6436 $\text{Int}[(a_+ + \text{ArcTanh}[(c_+)(x_+)^{n_+}]*(b_+))^{p_+}, x_Symbol] \rightarrow \text{Simp}[x*(a + b*\text{ArcTanh}[c*x^n])^p, x] - \text{Simp}[b*c*n*p \ \text{Int}[x^n*((a + b*\text{ArcTanh}[c*x^n])^{p-1}/(1 - c^2*x^{2*n}))], x], x] /; \text{FreeQ}\{a, b, c, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[n, 1] \ || \ \text{EqQ}[p, 1])$

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6470 `Int[((a_.) + ArcTanh[(c_.)*(x_)*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol] :> Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6510 `Int[((a_.) + ArcTanh[(c_.)*(x_)*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6542 `Int((((a_.) + ArcTanh[(c_.)*(x_)*(b_.))^(p_.)*((f_.)*(x_)^(m_)))/((d_) + (e_.)*(x_)^2), x_Symbol] :> Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

rule 6546 `Int((((a_.) + ArcTanh[(c_.)*(x_)*(b_.))^(p_.)*(x_))/((d_) + (e_.)*(x_)^2), x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 19.24 (sec) , antiderivative size = 1341, normalized size of antiderivative = 3.59

method	result	size
derivativedivides	Expression too large to display	1341
default	Expression too large to display	1341
parts	Expression too large to display	1403

input `int(x^3*(a+b*arctanh(c*x^(1/2)))^3,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 2/c^8*(-11/30*b^3+3/16*I*b^3*Pi*csgn(I*(1+c*x^(1/2))/(-c^2*x+1)^(1/2))*csgn \\ & n(I*(1+c*x^(1/2))^2/(c^2*x-1))^2*arctanh(c*x^(1/2))^2+1/8*a^3*c^8*x^4+3/32 \\ & *I*b^3*Pi*csgn(I/(1-(1+c*x^(1/2))^2/(c^2*x-1)))*csgn(I*(1+c*x^(1/2))^2/(c^ \\ & 2*x-1)/(1-(1+c*x^(1/2))^2/(c^2*x-1)))^2*arctanh(c*x^(1/2))^2-3/32*I*b^3*Pi \\ & *csgn(I*(1+c*x^(1/2))^2/(c^2*x-1))*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1)/(1-(1+ \\ & c*x^(1/2))^2/(c^2*x-1)))^2*arctanh(c*x^(1/2))^2+3/32*I*b^3*Pi*csgn(I*(1+c* \\ & x^(1/2))/(-c^2*x+1)^(1/2))^2*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1))*arctanh(c*x \\ & ^{(1/2)})^2-3/32*I*b^3*Pi*csgn(I/(1-(1+c*x^(1/2))^2/(c^2*x-1)))*csgn(I*(1+c* \\ & x^(1/2))^2/(c^2*x-1))*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1)/(1-(1+c*x^(1/2))^2/ \\ & (c^2*x-1)))*arctanh(c*x^(1/2))^2-44/35*b^3*arctanh(c*x^(1/2))*ln(1+I*(1+c* \\ & x^(1/2))/(-c^2*x+1)^(1/2))-44/35*b^3*arctanh(c*x^(1/2))*ln(1-I*(1+c*x^(1/2) \\ &))/(-c^2*x+1)^(1/2))+3/8*b^3*arctanh(c*x^(1/2))^2*ln((1+c*x^(1/2))/(-c^2*x \\ & +1)^(1/2))+3/16*b^3*arctanh(c*x^(1/2))^2*ln(c*x^(1/2)-1)-3/16*b^3*arctanh(\\ & c*x^(1/2))^2*ln(1+c*x^(1/2))+3*a*b^2*(1/8*c^8*x^4*arctanh(c*x^(1/2))^2+1/2 \\ & 8*arctanh(c*x^(1/2))*c^7*x^(7/2)+1/20*arctanh(c*x^(1/2))*c^5*x^(5/2)+1/12* \\ & arctanh(c*x^(1/2))*c^3*x^(3/2)+1/4*arctanh(c*x^(1/2))*c*x^(1/2)+1/8*arctan \\ & h(c*x^(1/2))*ln(c*x^(1/2)-1)-1/8*arctanh(c*x^(1/2))*ln(1+c*x^(1/2))+1/32*ln \\ & n(c*x^(1/2)-1)^2-1/16*ln(c*x^(1/2)-1)*ln(1/2*c*x^(1/2)+1/2)-1/16*(ln(1+c*x \\ & ^{(1/2)})-ln(1/2*c*x^(1/2)+1/2))*ln(-1/2*c*x^(1/2)+1/2)+1/32*ln(1+c*x^(1/2)) \\ & ^2+1/168*c^6*x^3+3/140*c^4*x^2+71/840*c^2*x+22/105*ln(c*x^(1/2)-1)+22/1\dots \end{aligned}$$

Fricas [F]

$$\int x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \int (b \operatorname{artanh}(c\sqrt{x}) + a)^3 x^3 dx$$

input `integrate(x^3*(a+b*arctanh(c*x^(1/2)))^3,x, algorithm="fricas")`

output `integral(b^3*x^3*arctanh(c*sqrt(x))^3 + 3*a*b^2*x^3*arctanh(c*sqrt(x))^2 + 3*a^2*b*x^3*arctanh(c*sqrt(x)) + a^3*x^3, x)`

Sympy [F]

$$\int x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \int x^3 (a + b \operatorname{atanh}(c\sqrt{x}))^3 dx$$

input `integrate(x**3*(a+b*atanh(c*x**(1/2)))**3,x)`

output `Integral(x**3*(a + b*atanh(c*sqrt(x)))**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1972 vs. $2(297) = 594$.

Time = 0.67 (sec) , antiderivative size = 1972, normalized size of antiderivative = 5.27

$$\int x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \text{Too large to display}$$

input `integrate(x^3*(a+b*arctanh(c*x^(1/2)))^3,x, algorithm="maxima")`

output

```

1/4*a^3*x^4 - 1/26880*a*b^2*c*((315*c^7*x^4 + 500*c^5*x^3 + 1002*c^3*x^2 +
3684*c*x - 12*(105*c^7*x^4 + 120*c^6*x^(7/2) + 140*c^5*x^3 + 168*c^4*x^(5
/2) + 210*c^3*x^2 + 280*c^2*x^(3/2) + 420*c*x + 840*sqrt(x))*log(c*sqrt(x)
+ 1))/c^8 - 6396*log(c*sqrt(x) + 1)/c^9 - 6396*log(c*sqrt(x) - 1)/c^9 -
1/2240*(840*x^4*log(c*sqrt(x) + 1) - c*((105*c^7*x^4 - 120*c^6*x^(7/2) + 1
40*c^5*x^3 - 168*c^4*x^(5/2) + 210*c^3*x^2 - 280*c^2*x^(3/2) + 420*c*x - 8
40*sqrt(x))/c^8 + 840*log(c*sqrt(x) + 1)/c^9))*a*b^2*log(-c*sqrt(x) + 1) +
1/2240*(840*x^4*log(c*sqrt(x) + 1) - c*((105*c^7*x^4 - 120*c^6*x^(7/2) +
140*c^5*x^3 - 168*c^4*x^(5/2) + 210*c^3*x^2 - 280*c^2*x^(3/2) + 420*c*x -
840*sqrt(x))/c^8 + 840*log(c*sqrt(x) + 1)/c^9))*a^2*b - 1/2240*(840*x^4*lo
g(-c*sqrt(x) + 1) - c*((105*c^7*x^4 + 120*c^6*x^(7/2) + 140*c^5*x^3 + 168*
c^4*x^(5/2) + 210*c^3*x^2 + 280*c^2*x^(3/2) + 420*c*x + 840*sqrt(x))/c^8 +
840*log(c*sqrt(x) - 1)/c^9))*a^2*b + 1/1881600*(11025*(32*log(-c*sqrt(x)
+ 1)^2 - 8*log(-c*sqrt(x) + 1) + 1)*(c*sqrt(x) - 1)^8 + 57600*(49*log(-c*s
qrt(x) + 1)^2 - 14*log(-c*sqrt(x) + 1) + 2)*(c*sqrt(x) - 1)^7 + 548800*(18
*log(-c*sqrt(x) + 1)^2 - 6*log(-c*sqrt(x) + 1) + 1)*(c*sqrt(x) - 1)^6 + 79
0272*(25*log(-c*sqrt(x) + 1)^2 - 10*log(-c*sqrt(x) + 1) + 2)*(c*sqrt(x) -
1)^5 + 3087000*(8*log(-c*sqrt(x) + 1)^2 - 4*log(-c*sqrt(x) + 1) + 1)*(c*sqr
t(x) - 1)^4 + 2195200*(9*log(-c*sqrt(x) + 1)^2 - 6*log(-c*sqrt(x) + 1) +
2)*(c*sqrt(x) - 1)^3 + 4939200*(2*log(-c*sqrt(x) + 1)^2 - 2*log(-c*sqrt...

```

Giac [F]

$$\int x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \int (b \operatorname{artanh}(c\sqrt{x}) + a)^3 x^3 dx$$

input

```
integrate(x^3*(a+b*arctanh(c*x^(1/2)))^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*sqrt(x)) + a)^3*x^3, x)
```


3.202 $\int x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx$

Optimal result	1617
Mathematica [A] (verified)	1618
Rubi [B] (verified)	1618
Maple [C] (warning: unable to verify)	1626
Fricas [F]	1627
Sympy [F]	1628
Maxima [B] (verification not implemented)	1628
Giac [F]	1629
Mupad [F(-1)]	1630
Reduce [F]	1630

Optimal result

Integrand size = 18, antiderivative size = 304

$$\int x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \frac{19b^3 \sqrt{x}}{30c^5} + \frac{b^3 x^{3/2}}{30c^3} - \frac{19b^3 \operatorname{arctanh}(c\sqrt{x})}{30c^6} + \frac{8b^2 x (a + b \operatorname{arctanh}(c\sqrt{x}))}{15c^4} + \frac{b^2 x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))}{10c^2} + \frac{23b (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{15c^6} + \frac{b\sqrt{x} (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{c^5} + \frac{bx^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{3c^3} + \frac{bx^{5/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{5c} - \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{3c^6} + \frac{1}{3} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{46b^2 (a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1-c\sqrt{x}}\right)}{15c^6} - \frac{23b^3 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-c\sqrt{x}}\right)}{15c^6}$$

output

```
19/30*b^3*x^(1/2)/c^5+1/30*b^3*x^(3/2)/c^3-19/30*b^3*arctanh(c*x^(1/2))/c^6+8/15*b^2*x*(a+b*arctanh(c*x^(1/2)))/c^4+1/10*b^2*x^2*(a+b*arctanh(c*x^(1/2)))/c^2+23/15*b*(a+b*arctanh(c*x^(1/2)))^2/c^6+b*x^(1/2)*(a+b*arctanh(c*x^(1/2)))^2/c^5+1/3*b*x^(3/2)*(a+b*arctanh(c*x^(1/2)))^2/c^3+1/5*b*x^(5/2)*(a+b*arctanh(c*x^(1/2)))^2/c-1/3*(a+b*arctanh(c*x^(1/2)))^3/c^6+1/3*x^3*(a+b*arctanh(c*x^(1/2)))^3-46/15*b^2*(a+b*arctanh(c*x^(1/2)))*ln(2/(1-c*x^(1/2)))/c^6-23/15*b^3*polylog(2,1-2/(1-c*x^(1/2)))/c^6
```

Mathematica [A] (verified)

Time = 0.53 (sec) , antiderivative size = 351, normalized size of antiderivative = 1.15

$$\int x^2 (a + \operatorname{arctanh}(c\sqrt{x}))^3 dx$$

$$= \frac{-19ab^2 + 30a^2bc\sqrt{x} + 19b^3c\sqrt{x} + 16ab^2c^2x + 10a^2bc^3x^{3/2} + b^3c^3x^{3/2} + 3ab^2c^4x^2 + 6a^2bc^5x^{5/2} + 10a^3c^6x^3}{30c^6}$$

input

```
Integrate[x^2*(a + b*ArcTanh[c*Sqrt[x]])^3,x]
```

output

```
(-19*a*b^2 + 30*a^2*b*c*Sqrt[x] + 19*b^3*c*Sqrt[x] + 16*a*b^2*c^2*x + 10*a^2*b*c^3*x^(3/2) + b^3*c^3*x^(3/2) + 3*a*b^2*c^4*x^2 + 6*a^2*b*c^5*x^(5/2) + 10*a^3*c^6*x^3 + 2*b^2*(b*(-23 + 15*c*Sqrt[x] + 5*c^3*x^(3/2) + 3*c^5*x^(5/2)) + 15*a*(-1 + c^6*x^3))*ArcTanh[c*Sqrt[x]]^2 + 10*b^3*(-1 + c^6*x^3)*ArcTanh[c*Sqrt[x]]^3 + b*ArcTanh[c*Sqrt[x]]*(30*a^2*c^6*x^3 + 4*a*b*c*Sqrt[x]*(15 + 5*c^2*x + 3*c^4*x^2) + b^2*(-19 + 16*c^2*x + 3*c^4*x^2) - 92*b^2*Log[1 + E^(-2*ArcTanh[c*Sqrt[x]])]) + 15*a^2*b*Log[1 - c*Sqrt[x]] - 15*a^2*b*Log[1 + c*Sqrt[x]] + 46*a*b^2*Log[1 - c^2*x] + 46*b^3*PolyLog[2, -E^(-2*ArcTanh[c*Sqrt[x]])])/(30*c^6)
```

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 611 vs. $2(304) = 608$.

Time = 4.13 (sec) , antiderivative size = 611, normalized size of antiderivative = 2.01, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {6454, 6452, 6542, 6452, 6542, 6452, 254, 2009, 6542, 6436, 6452, 262, 219, 6510, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 (a + \operatorname{arctanh}(c\sqrt{x}))^3 dx$$

↓ 6454

$$2 \int x^{5/2} (a + \operatorname{barctanh}(c\sqrt{x}))^3 d\sqrt{x}$$

↓ 6452

$$2 \left(\frac{1}{6} x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \int \frac{x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x} \right)$$

↓ 6542

$$2 \left(\frac{1}{6} x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\int x^2 (a + \operatorname{barctanh}(c\sqrt{x}))^2 d\sqrt{x}}{c^2} \right) \right)$$

↓ 6452

$$2 \left(\frac{1}{6} x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \left(\frac{\int \frac{x^2 (a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{2}{5} bc \int \frac{x^{5/2}}{1 - c^2 x} d\sqrt{x}}{c^2} \right) \right)$$

↓ 6542

$$2 \left(\frac{1}{6} x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \left(\frac{\int \frac{x (a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\int x (a + \operatorname{barctanh}(c\sqrt{x}))^2 d\sqrt{x}}{c^2} - \frac{\frac{1}{5} x^{5/2} (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{2}{5} bc \int \frac{x^{5/2}}{1 - c^2 x} d\sqrt{x}}{c^2} \right) \right)$$

↓ 6452

$$2 \left(\frac{1}{6} x^3 (a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \left(\frac{\int \frac{x (a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{2}{3} bc \int \frac{x^{3/2} (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} \right) \right)$$

↓ 254

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \left(\frac{\int \frac{x (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{2}{3} bc \int \frac{x^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x}}{c^2} \right) \right)$$

↓ 2009

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \left(\frac{\int \frac{x (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{2}{3} bc \int \frac{x^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x}}{c^2} \right) \right)$$

↓ 6542

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \left(\frac{\frac{\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\int (a + b \operatorname{arctanh}(c\sqrt{x}))^2 d\sqrt{x}}{c^2}}{c^2} - \frac{\frac{1}{3} x^{3/2} (a + b \operatorname{arctanh}(c\sqrt{x}))}{c^2} \right) \right)$$

↓ 6436

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \left(\frac{\frac{\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\sqrt{x} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \int \frac{\sqrt{x} (a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2}}{c^2} \right) \right)$$

↓ 6452

$$2 \left(\frac{1}{6} x^3 (a + \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \left(\frac{\int \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\sqrt{x} (a + \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} - 2bc \int \frac{\sqrt{x} (a + \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x} \right) \right)$$

↓ 262

$$2 \left(\frac{1}{6} x^3 (a + \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \left(\frac{\int \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\sqrt{x} (a + \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} - 2bc \int \frac{\sqrt{x} (a + \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x} \right) \right)$$

↓ 219

$$2 \left(\frac{1}{6} x^3 (a + \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \left(\frac{\int \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\sqrt{x} (a + \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} - 2bc \int \frac{\sqrt{x} (a + \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x} \right) \right)$$

↓ 6510

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \left(\frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \int \frac{\sqrt{x} (a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} \right) \right)$$

↓ 6546

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \left(\frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \left(\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{1 - c\sqrt{x}} d\sqrt{x} - (a + b \operatorname{arctanh}(c\sqrt{x})) \right)}{c^2} \right) \right)$$

↓ 6470

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \left(\frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \left(\frac{\log\left(\frac{2}{1 - c\sqrt{x}}\right) (a + b \operatorname{arctanh}(c\sqrt{x}))}{c} \right)}{c^2} \right) \right)$$

↓ 2849

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \right) \left(\frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc}{c^2} \left(\frac{b \int \frac{\log\left(\frac{2}{1-c\sqrt{x}}\right) d\frac{1}{1-c\sqrt{x}}}{e} + \frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)}{c} \right) \right)$$

↓ 2752

$$2 \left(\frac{1}{6} x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{1}{2} bc \right) \left(\frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x} (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc}{c^2} \left(\frac{\log\left(\frac{2}{1-c\sqrt{x}}\right) (a + b \operatorname{arctanh}(c\sqrt{x}))}{e} + \frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)}{c} \right) \right)$$

input Int [x^2*(a + b*ArcTanh [c*Sqrt [x]])^3, x]

output

$$2*((x^3*(a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])^3)/6 - (b*c*(-((x^{5/2})*(a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])^2)/5 - (2*b*c*(-((x^2*(a + b*\text{ArcTanh}[c*\text{Sqrt}[x]]))/4 - (b*c*(-(Sqrt[x]/c^4) - x^{3/2}/(3*c^2) + \text{ArcTanh}[c*\text{Sqrt}[x]]/c^5))/4)/c^2) + (-((x*(a + b*\text{ArcTanh}[c*\text{Sqrt}[x]]))/2 - (b*c*(-(Sqrt[x]/c^2) + \text{ArcTanh}[c*\text{Sqrt}[x]]/c^3))/2)/c^2) + (-1/2*(a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])^2/(b*c^2) + ((a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])*\text{Log}[2/(1 - c*\text{Sqrt}[x])])/c + (b*\text{PolyLog}[2, 1 - 2/(1 - c*\text{Sqrt}[x])])/(2*c))/c/c^2/c^2)/5)/c^2) + (-((x^{3/2})*(a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])^2)/3 - (2*b*c*(-((x*(a + b*\text{ArcTanh}[c*\text{Sqrt}[x]]))/2 - (b*c*(-(Sqrt[x]/c^2) + \text{ArcTanh}[c*\text{Sqrt}[x]]/c^3))/2)/c^2) + (-1/2*(a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])^2/(b*c^2) + ((a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])*\text{Log}[2/(1 - c*\text{Sqrt}[x])])/c + (b*\text{PolyLog}[2, 1 - 2/(1 - c*\text{Sqrt}[x])])/(2*c))/c/c^2)/3)/c^2) + ((a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])^3/(3*b*c^3) - (Sqrt[x]*(a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])^2 - 2*b*c*(-1/2*(a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])^2/(b*c^2) + ((a + b*\text{ArcTanh}[c*\text{Sqrt}[x]])*\text{Log}[2/(1 - c*\text{Sqrt}[x])])/c + (b*\text{PolyLog}[2, 1 - 2/(1 - c*\text{Sqrt}[x])])/(2*c))/c))/c^2)/c^2)/2)$$

Defintions of rubi rules used

rule 219

$$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 254

$$\text{Int}[(x_)^{(m)}/((a_ + (b_)*(x_)^2), x_Symbol] \rightarrow \text{Int}[\text{PolynomialDivide}[x^m, a + b*x^2, x], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{IGtQ}[m, 3]$$

rule 262

$$\text{Int}[(c_)*(x_)^{(m)}*((a_ + (b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[c*(c*x)^{(m-1)}*((a + b*x^2)^{(p+1)}/(b*(m+2*p+1))), x] - \text{Simp}[a*c^2*((m-1)/(b*(m+2*p+1))) \ \text{Int}[(c*x)^{(m-2)}*(a + b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{GtQ}[m, 2-1] \ \&\& \ \text{NeQ}[m+2*p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 2009

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

rule 2752 $\text{Int}[\text{Log}[(c_)*(x_)]/((d_)+(e_)*(x_)), x_Symbol] \rightarrow \text{Simp}[(-e^{(-1)})*\text{PolyLog}[2, 1 - c*x], x] /; \text{FreeQ}[\{c, d, e\}, x] \ \&\& \ \text{EqQ}[e + c*d, 0]$

rule 2849 $\text{Int}[\text{Log}[(c_)/((d_)+(e_)*(x_))]/((f_)+(g_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[-e/g \ \text{Subst}[\text{Int}[\text{Log}[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; \text{FreeQ}[\{c, d, e, f, g\}, x] \ \&\& \ \text{EqQ}[c, 2*d] \ \&\& \ \text{EqQ}[e^2*f + d^2*g, 0]$

rule 6436 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^(n_)]*(b_))^(p_), x_Symbol] \rightarrow \text{Simp}[x*(a + b*\text{ArcTanh}[c*x^n])^p, x] - \text{Simp}[b*c*n*p \ \text{Int}[x^n*((a + b*\text{ArcTanh}[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; \text{FreeQ}[\{a, b, c, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[n, 1] \ || \ \text{EqQ}[p, 1])$

rule 6452 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^(n_)]*(b_))^(p_)*(x_)^(m_), x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m + 1)), x] - \text{Simp}[b*c*n*(p/(m + 1)) \ \text{Int}[x^{(m + n)}*((a + b*\text{ArcTanh}[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 6454 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^(n_)]*(b_))^(p_)*(x_)^(m_), x_Symbol] \rightarrow \text{Simp}[1/n \ \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)*(a + b*\text{ArcTanh}[c*x])^p}, x], x, x^n], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]$

rule 6470 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)]*(b_))^(p_)/((d_)+(e_)*(x_)), x_Symbol] \rightarrow \text{Simp}[(-a + b*\text{ArcTanh}[c*x])^p*(\text{Log}[2/(1 + e*(x/d))]/e), x] + \text{Simp}[b*c*(p/e) \ \text{Int}[(a + b*\text{ArcTanh}[c*x])^(p - 1)*(\text{Log}[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d^2 - e^2, 0]$

rule 6510 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)]*(b_))^(p_)/((d_)+(e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; \text{FreeQ}[\{a, b, c, d, e, p\}, x] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{NeQ}[p, -1]$

rule 6542

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*((f_.)*(x_))^(m_)/((d_) + (
e_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*
x])^p, x], x] - Simp[d*(f^2/e Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/
(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m,
1]
```

rule 6546

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*(x_)/((d_) + (e_.)*(x_)^2),
x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/
(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 16.20 (sec) , antiderivative size = 1264, normalized size of antiderivative = 4.16

method	result	size
derivativeldivides	Expression too large to display	1264
default	Expression too large to display	1264
parts	Expression too large to display	1324

input

```
int(x^2*(a+b*arctanh(c*x^(1/2)))^3,x,method=_RETURNVERBOSE)
```

output

```

2/c^6*(-1/3*b^3-1/8*I*b^3*Pi*csgn(I/(1-(1+c*x^(1/2))^2/(c^2*x-1)))*csgn(I*
(1+c*x^(1/2))^2/(c^2*x-1))*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1)/(1-(1+c*x^(1/2)
))^2/(c^2*x-1)))*arctanh(c*x^(1/2))^2-1/4*I*b^3*Pi*csgn(I/(1-(1+c*x^(1/2)
))^2/(c^2*x-1)))^3*arctanh(c*x^(1/2))^2+1/8*I*b^3*Pi*csgn(I*(1+c*x^(1/2))^2/
(c^2*x-1)/(1-(1+c*x^(1/2))^2/(c^2*x-1)))^3*arctanh(c*x^(1/2))^2+1/4*I*b^3*
Pi*csgn(I/(1-(1+c*x^(1/2))^2/(c^2*x-1)))^2*arctanh(c*x^(1/2))^2+1/8*I*b^3*
Pi*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1))^3*arctanh(c*x^(1/2))^2+3*a*b^2*(1/6*c
^6*x^3*arctanh(c*x^(1/2))^2+1/15*arctanh(c*x^(1/2))*c^5*x^(5/2)+1/9*arctan
h(c*x^(1/2))*c^3*x^(3/2)+1/3*arctanh(c*x^(1/2))*c*x^(1/2)+1/6*arctanh(c*x^
(1/2))*ln(c*x^(1/2)-1)-1/6*arctanh(c*x^(1/2))*ln(1+c*x^(1/2))+1/24*ln(c*x^
(1/2)-1)^2-1/12*ln(c*x^(1/2)-1)*ln(1/2*c*x^(1/2)+1/2)+1/24*ln(1+c*x^(1/2)
)^2-1/12*(ln(1+c*x^(1/2))-ln(1/2*c*x^(1/2)+1/2))*ln(-1/2*c*x^(1/2)+1/2)+1/6
0*c^4*x^2+4/45*c^2*x+23/90*ln(c*x^(1/2)-1)+23/90*ln(1+c*x^(1/2)))+3*a^2*b*
(1/6*c^6*x^3*arctanh(c*x^(1/2))+1/30*c^5*x^(5/2)+1/18*c^3*x^(3/2)+1/6*c*x^
(1/2)+1/12*ln(c*x^(1/2)-1)-1/12*ln(1+c*x^(1/2)))+1/6*a^3*c^6*x^3-23/15*b^3
*arctanh(c*x^(1/2))*ln(1+I*(1+c*x^(1/2))/(-c^2*x+1)^(1/2))-23/15*b^3*arcta
nh(c*x^(1/2))*ln(1-I*(1+c*x^(1/2))/(-c^2*x+1)^(1/2))+1/2*b^3*arctanh(c*x^
(1/2))^2*ln((1+c*x^(1/2))/(-c^2*x+1)^(1/2))+1/4*b^3*arctanh(c*x^(1/2))^2*ln
(c*x^(1/2)-1)-1/4*b^3*arctanh(c*x^(1/2))^2*ln(1+c*x^(1/2))+19/60*b^3*c*x^
(1/2)+1/60*b^3*c^3*x^(3/2)+1/8*I*b^3*Pi*csgn(I*(1+c*x^(1/2))/(-c^2*x+1)^...

```

Fricas [F]

$$\int x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \int (b \operatorname{arctanh}(c\sqrt{x}) + a)^3 x^2 dx$$

input

```
integrate(x^2*(a+b*arctanh(c*x^(1/2)))^3,x, algorithm="fricas")
```

output

```
integral(b^3*x^2*arctanh(c*sqrt(x))^3 + 3*a*b^2*x^2*arctanh(c*sqrt(x))^2 +
3*a^2*b*x^2*arctanh(c*sqrt(x)) + a^3*x^2, x)
```


Sympy [F]

$$\int x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \int x^2 (a + b \operatorname{atanh}(c\sqrt{x}))^3 dx$$

input `integrate(x**2*(a+b*atanh(c*x**(1/2)))**3,x)`

output `Integral(x**2*(a + b*atanh(c*sqrt(x)))**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1579 vs. 2(243) = 486.

Time = 0.59 (sec) , antiderivative size = 1579, normalized size of antiderivative = 5.19

$$\int x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \text{Too large to display}$$

input `integrate(x^2*(a+b*arctanh(c*x^(1/2)))^3,x, algorithm="maxima")`

output

```

1/3*a^3*x^3 - 1/720*a*b^2*c*((20*c^5*x^3 + 39*c^3*x^2 + 138*c*x - 6*(10*c^
5*x^3 + 12*c^4*x^(5/2) + 15*c^3*x^2 + 20*c^2*x^(3/2) + 30*c*x + 60*sqrt(x)
)*log(c*sqrt(x) + 1))/c^6 - 222*log(c*sqrt(x) + 1)/c^7 - 222*log(c*sqrt(x)
- 1)/c^7) - 1/120*(60*x^3*log(c*sqrt(x) + 1) - c*((10*c^5*x^3 - 12*c^4*x^
(5/2) + 15*c^3*x^2 - 20*c^2*x^(3/2) + 30*c*x - 60*sqrt(x))/c^6 + 60*log(c*
sqrt(x) + 1)/c^7))*a*b^2*log(-c*sqrt(x) + 1) + 1/120*(60*x^3*log(c*sqrt(x)
+ 1) - c*((10*c^5*x^3 - 12*c^4*x^(5/2) + 15*c^3*x^2 - 20*c^2*x^(3/2) + 30
*c*x - 60*sqrt(x))/c^6 + 60*log(c*sqrt(x) + 1)/c^7))*a^2*b - 1/120*(60*x^3
*log(-c*sqrt(x) + 1) - c*((10*c^5*x^3 + 12*c^4*x^(5/2) + 15*c^3*x^2 + 20*c
^2*x^(3/2) + 30*c*x + 60*sqrt(x))/c^6 + 60*log(c*sqrt(x) - 1)/c^7))*a^2*b
+ 1/7200*(100*(18*log(-c*sqrt(x) + 1)^2 - 6*log(-c*sqrt(x) + 1) + 1)*(c*sq
rt(x) - 1)^6 + 432*(25*log(-c*sqrt(x) + 1)^2 - 10*log(-c*sqrt(x) + 1) + 2)
*(c*sqrt(x) - 1)^5 + 3375*(8*log(-c*sqrt(x) + 1)^2 - 4*log(-c*sqrt(x) + 1)
+ 1)*(c*sqrt(x) - 1)^4 + 4000*(9*log(-c*sqrt(x) + 1)^2 - 6*log(-c*sqrt(x)
+ 1) + 2)*(c*sqrt(x) - 1)^3 + 13500*(2*log(-c*sqrt(x) + 1)^2 - 2*log(-c*s
qrt(x) + 1) + 1)*(c*sqrt(x) - 1)^2 + 10800*(log(-c*sqrt(x) + 1)^2 - 2*log(
-c*sqrt(x) + 1) + 2)*(c*sqrt(x) - 1))*a*b^2/c^6 - 1/864000*(1000*(36*log(-
c*sqrt(x) + 1)^3 - 18*log(-c*sqrt(x) + 1)^2 + 6*log(-c*sqrt(x) + 1) - 1)*(
c*sqrt(x) - 1)^6 + 1728*(125*log(-c*sqrt(x) + 1)^3 - 75*log(-c*sqrt(x) + 1
)^2 + 30*log(-c*sqrt(x) + 1) - 6)*(c*sqrt(x) - 1)^5 + 16875*(32*log(-c*...

```

Giac [F]

$$\int x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \int (b \operatorname{arctanh}(c\sqrt{x}) + a)^3 x^2 dx$$

input

```
integrate(x^2*(a+b*arctanh(c*x^(1/2)))^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*sqrt(x)) + a)^3*x^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \int x^2 (a + b \operatorname{atanh}(c\sqrt{x}))^3 dx$$

input `int(x^2*(a + b*atanh(c*x^(1/2)))^3,x)`output `int(x^2*(a + b*atanh(c*x^(1/2)))^3, x)`**Reduce [F]**

$$\int x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx$$

$$= \frac{46 \left(\int \frac{\operatorname{atanh}(\sqrt{x}c)}{c^2x-1} dx \right) b^3 c^2 + 12\sqrt{x} \operatorname{atanh}(\sqrt{x}c) a b^2 c^5 x^2 + 20\sqrt{x} \operatorname{atanh}(\sqrt{x}c) a b^2 c^3 x + 10 \operatorname{atanh}(\sqrt{x}c)^3 b^3 c^2}{1}$$

input `int(x^2*(a+b*atanh(c*x^(1/2)))^3,x)`output `(10*atanh(sqrt(x)*c)**3*b**3*c**6*x**3 - 10*atanh(sqrt(x)*c)**3*b**3 + 6*sqrt(x)*atanh(sqrt(x)*c)**2*b**3*c**5*x**2 + 10*sqrt(x)*atanh(sqrt(x)*c)**2*b**3*c**3*x + 30*sqrt(x)*atanh(sqrt(x)*c)**2*b**3*c + 30*atanh(sqrt(x)*c)**2*a*b**2*c**6*x**3 - 30*atanh(sqrt(x)*c)**2*a*b**2 + 12*sqrt(x)*atanh(sqrt(x)*c)*a*b**2*c**5*x**2 + 20*sqrt(x)*atanh(sqrt(x)*c)*a*b**2*c**3*x + 60*sqrt(x)*atanh(sqrt(x)*c)*a*b**2*c + 30*atanh(sqrt(x)*c)*a**2*b*c**6*x**3 - 30*atanh(sqrt(x)*c)*a**2*b + 92*atanh(sqrt(x)*c)*a*b**2 + 3*atanh(sqrt(x)*c)*b**3*c**4*x**2 + 16*atanh(sqrt(x)*c)*b**3*c**2*x - 19*atanh(sqrt(x)*c)*b**3 + 6*sqrt(x)*a**2*b*c**5*x**2 + 10*sqrt(x)*a**2*b*c**3*x + 30*sqrt(x)*a**2*b*c + sqrt(x)*b**3*c**3*x + 19*sqrt(x)*b**3*c + 46*int(atanh(sqrt(x)*c)/(c**2*x - 1),x)*b**3*c**2 + 92*log(sqrt(x)*c - 1)*a*b**2 + 10*a**3*c**6*x**3 + 3*a*b**2*c**4*x**2 + 16*a*b**2*c**2*x)/(30*c**6)`

3.203 $\int x(a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx$

Optimal result	1631
Mathematica [A] (verified)	1632
Rubi [A] (verified)	1632
Maple [C] (warning: unable to verify)	1638
Fricas [F]	1639
Sympy [F]	1639
Maxima [B] (verification not implemented)	1639
Giac [F]	1640
Mupad [F(-1)]	1641
Reduce [F]	1641

Optimal result

Integrand size = 16, antiderivative size = 234

$$\begin{aligned}
 \int x(a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = & \frac{b^3\sqrt{x}}{2c^3} - \frac{b^3 \operatorname{arctanh}(c\sqrt{x})}{2c^4} + \frac{b^2 x(a + b \operatorname{arctanh}(c\sqrt{x}))}{2c^2} \\
 & + \frac{2b(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{c^4} \\
 & + \frac{3b\sqrt{x}(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{2c^3} \\
 & + \frac{bx^{3/2}(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{2c} \\
 & - \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{2c^4} + \frac{1}{2}x^2(a + b \operatorname{arctanh}(c\sqrt{x}))^3 \\
 & - \frac{4b^2(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1-c\sqrt{x}}\right)}{c^4} \\
 & - \frac{2b^3 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-c\sqrt{x}}\right)}{c^4}
 \end{aligned}$$

output

$$\frac{1/2*b^3*x^{(1/2)}/c^3-1/2*b^3*\operatorname{arctanh}(c*x^{(1/2)})/c^4+1/2*b^2*x*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))/c^2+2*b*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))^2/c^4+3/2*b*x^{(1/2)}*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))^2/c^3+1/2*b*x^{(3/2)}*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))^2/c-1/2*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))^3/c^4+1/2*x^2*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))^3-4*b^2*(a+b*\operatorname{arctanh}(c*x^{(1/2)}))*\ln(2/(1-c*x^{(1/2)}))/c^4-2*b^3*\operatorname{polylog}(2,1-2/(1-c*x^{(1/2)}))/c^4}$$
Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.22

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx$$

$$= \frac{-2ab^2 + 6a^2bc\sqrt{x} + 2b^3c\sqrt{x} + 2ab^2c^2x + 2a^2bc^3x^{3/2} + 2a^3c^4x^2 + 2b^2(b(-4 + 3c\sqrt{x} + c^3x^{3/2}) + 3a(-1 - c^2x)) \operatorname{ArcTanh}[c\sqrt{x}]^2 + 2b^3(-1 + c^4x^2) \operatorname{ArcTanh}[c\sqrt{x}]^3 + 2b \operatorname{ArcTanh}[c\sqrt{x}] (3a^2c^4x^2 + b^2(-1 + c^2x) + 2a*b*c\sqrt{x}*(3 + c^2x) - 8b^2*\operatorname{Log}[1 + E^{(-2*\operatorname{ArcTanh}[c\sqrt{x}])}] + 3a^2*b*\operatorname{Log}[1 - c\sqrt{x}] - 3a^2*b*\operatorname{Log}[1 + c\sqrt{x}] + 8a*b^2*\operatorname{Log}[1 - c^2x] + 8b^3*\operatorname{PolyLog}[2, -E^{(-2*\operatorname{ArcTanh}[c\sqrt{x}])}])]}{4c^4}$$

input

$$\operatorname{Integrate}[x*(a + b*\operatorname{ArcTanh}[c*\operatorname{Sqrt}[x]])^3, x]$$

output

$$\frac{(-2*a*b^2 + 6*a^2*b*c*\operatorname{Sqrt}[x] + 2*b^3*c*\operatorname{Sqrt}[x] + 2*a*b^2*c^2*x + 2*a^2*b*c^3*x^{(3/2)} + 2*a^3*c^4*x^2 + 2*b^2*(b*(-4 + 3*c*\operatorname{Sqrt}[x] + c^3*x^{(3/2)}) + 3*a*(-1 + c^4*x^2))*\operatorname{ArcTanh}[c*\operatorname{Sqrt}[x]]^2 + 2*b^3*(-1 + c^4*x^2)*\operatorname{ArcTanh}[c*\operatorname{Sqrt}[x]]^3 + 2*b*\operatorname{ArcTanh}[c*\operatorname{Sqrt}[x]]*(3*a^2*c^4*x^2 + b^2*(-1 + c^2*x) + 2*a*b*c*\operatorname{Sqrt}[x]*(3 + c^2*x) - 8*b^2*\operatorname{Log}[1 + E^{(-2*\operatorname{ArcTanh}[c*\operatorname{Sqrt}[x]])}] + 3*a^2*b*\operatorname{Log}[1 - c*\operatorname{Sqrt}[x]] - 3*a^2*b*\operatorname{Log}[1 + c*\operatorname{Sqrt}[x]] + 8*a*b^2*\operatorname{Log}[1 - c^2*x] + 8*b^3*\operatorname{PolyLog}[2, -E^{(-2*\operatorname{ArcTanh}[c*\operatorname{Sqrt}[x]])}])]}{4*c^4}$$
Rubi [A] (verified)

Time = 2.18 (sec) , antiderivative size = 356, normalized size of antiderivative = 1.52, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {6454, 6452, 6542, 6452, 6542, 6436, 6452, 262, 219, 6510, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x(a + \operatorname{barctanh}(c\sqrt{x}))^3 dx \\
 & \quad \downarrow 6454 \\
 & 2 \int x^{3/2}(a + \operatorname{barctanh}(c\sqrt{x}))^3 d\sqrt{x} \\
 & \quad \downarrow 6452 \\
 & 2 \left(\frac{1}{4}x^2(a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{3}{4}bc \int \frac{x^2(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2x} d\sqrt{x} \right) \\
 & \quad \downarrow 6542 \\
 & 2 \left(\frac{1}{4}x^2(a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{3}{4}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2x} d\sqrt{x}}{c^2} - \frac{\int x(a + \operatorname{barctanh}(c\sqrt{x}))^2 d\sqrt{x}}{c^2} \right) \right) \\
 & \quad \downarrow 6452 \\
 & 2 \left(\frac{1}{4}x^2(a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{3}{4}bc \left(\frac{\int \frac{x(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{3}x^{3/2}(a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{2}{3}bc \int \frac{x^{3/2}(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2x} d\sqrt{x}}{c^2} \right) \right) \\
 & \quad \downarrow 6542 \\
 & 2 \left(\frac{1}{4}x^2(a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{3}{4}bc \left(\frac{\int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2x} d\sqrt{x}}{c^2} - \frac{\int (a + \operatorname{barctanh}(c\sqrt{x}))^2 d\sqrt{x}}{c^2} - \frac{\frac{1}{3}x^{3/2}(a + \operatorname{barctanh}(c\sqrt{x}))^2 - \frac{2}{3}bc \int \frac{x^{3/2}(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2x} d\sqrt{x}}{c^2} \right) \right) \\
 & \quad \downarrow 6436 \\
 & 2 \left(\frac{1}{4}x^2(a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{3}{4}bc \left(\frac{\int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2x} d\sqrt{x}}{c^2} - \frac{\sqrt{x}(a + \operatorname{barctanh}(c\sqrt{x}))^2 - 2bc \int \frac{\sqrt{x}(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2x} d\sqrt{x}}{c^2} \right) \right)
 \end{aligned}$$

↓ 6452

$$2 \left(\frac{1}{4} x^2 (a + \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{4} bc \left(\frac{\int \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\sqrt{x} (a + \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \int \frac{\sqrt{x} (a + \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x}}{c^2} \right) \right)$$

↓ 262

$$2 \left(\frac{1}{4} x^2 (a + \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{4} bc \left(\frac{\int \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\sqrt{x} (a + \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \int \frac{\sqrt{x} (a + \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x}}{c^2} \right) \right)$$

↓ 219

$$2 \left(\frac{1}{4} x^2 (a + \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{4} bc \left(\frac{\int \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\sqrt{x} (a + \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \int \frac{\sqrt{x} (a + \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x}}{c^2} \right) \right)$$

↓ 6510

$$2 \left(\frac{1}{4} x^2 (a + \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{4} bc \left(\frac{(a + \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x} (a + \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \int \frac{\sqrt{x} (a + \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} \right) \right)$$

↓ 6546

$$2 \left(\frac{1}{4} x^2 (a + \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{4} bc \left(\frac{(a + \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x}(a + \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \left(\frac{\int \frac{a + \operatorname{arctanh}(c\sqrt{x})}{1 - c\sqrt{x}} d\sqrt{x}}{c} - \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} \right)}{c^2} \right) \right)$$

↓ 6470

$$2 \left(\frac{1}{4} x^2 (a + \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{4} bc \left(\frac{(a + \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x}(a + \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \left(\frac{\log\left(\frac{2}{1 - c\sqrt{x}}\right)(a + \operatorname{arctanh}(c\sqrt{x}))}{c} - \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} \right)}{c^2} \right) \right)$$

↓ 2849

$$2 \left(\frac{1}{4} x^2 (a + \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{4} bc \left(\frac{(a + \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x}(a + \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \left(\frac{b \int \frac{\log\left(\frac{2}{1 - c\sqrt{x}}\right)}{1 - \frac{2}{1 - c\sqrt{x}}} d\frac{1}{1 - c\sqrt{x}}}{c} + \frac{\log\left(\frac{2}{1 - c\sqrt{x}}\right)(a + \operatorname{arctanh}(c\sqrt{x}))}{c} - \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} \right)}{c^2} \right) \right)$$

↓ 2752

$$2 \left(\frac{1}{4} x^2 (a + \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{4} bc \left(\frac{(a + \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x}(a + \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \left(\frac{\log\left(\frac{2}{1 - c\sqrt{x}}\right)(a + \operatorname{arctanh}(c\sqrt{x}))}{c} - \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} \right)}{c^2} \right) \right)$$

input `Int[x*(a + b*ArcTanh[c*Sqrt[x]])^3,x]`

output `2*((x^2*(a + b*ArcTanh[c*Sqrt[x]])^3)/4 - (3*b*c*(-((x^(3/2)*(a + b*ArcTanh[c*Sqrt[x]])^2)/3 - (2*b*c*(-((x*(a + b*ArcTanh[c*Sqrt[x]]))/2 - (b*c*(-(Sqrt[x]/c^2) + ArcTanh[c*Sqrt[x]]/c^3))/2)/c^2) + (-1/2*(a + b*ArcTanh[c*Sqrt[x]])^2/(b*c^2) + ((a + b*ArcTanh[c*Sqrt[x]])*Log[2/(1 - c*Sqrt[x]))]/c + (b*PolyLog[2, 1 - 2/(1 - c*Sqrt[x]))]/(2*c))/c)/c^2))/3)/c^2) + ((a + b*ArcTanh[c*Sqrt[x]])^3/(3*b*c^3) - (Sqrt[x]*(a + b*ArcTanh[c*Sqrt[x]])^2 - 2*b*c*(-1/2*(a + b*ArcTanh[c*Sqrt[x]])^2/(b*c^2) + ((a + b*ArcTanh[c*Sqrt[x]])*Log[2/(1 - c*Sqrt[x]))]/c + (b*PolyLog[2, 1 - 2/(1 - c*Sqrt[x]))]/(2*c))/c))/c^2)/c^2))/4)`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 262 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*(m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2752 `Int[Log[(c_)*(x_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_)]/((d_) + (e_)*(x_)]/((f_) + (g_)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 6436 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot b)^p, x_Symbol] \rightarrow \text{Simp}[x \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^p, x] - \text{Simp}[b \cdot c \cdot n \cdot p \cdot \text{Int}[x^n \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2n})], x], x] /;$ $\text{FreeQ}\{a, b, c, n, x\} \ \&\& \ \text{IGtQ}\{p, 0\} \ \&\& \ (\text{EqQ}\{n, 1\} \ || \ \text{EqQ}\{p, 1\})$

rule 6452 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot b)^p \cdot x^m, x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^p / (m+1), x] - \text{Simp}[b \cdot c \cdot n \cdot (p/(m+1)) \cdot \text{Int}[x^{m+n} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2n})], x], x] /;$ $\text{FreeQ}\{a, b, c, m, n, x\} \ \&\& \ \text{IGtQ}\{p, 0\} \ \&\& \ (\text{EqQ}\{p, 1\} \ || \ (\text{EqQ}\{n, 1\} \ \&\& \ \text{IntegerQ}\{m\})) \ \&\& \ \text{NeQ}\{m, -1\}$

rule 6454 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot b)^p \cdot x^m, x_Symbol] \rightarrow \text{Simp}[1/n \cdot \text{Subst}[\text{Int}[x^{\text{Simplify}[(m+1)/n} - 1] \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x, x^n], x] /;$ $\text{FreeQ}\{a, b, c, m, n, x\} \ \&\& \ \text{IGtQ}\{p, 1\} \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$

rule 6470 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((d) + (e) \cdot x), x_Symbol] \rightarrow \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{Log}[2/(1 + e \cdot (x/d))]/e), x] + \text{Simp}[b \cdot c \cdot (p/e) \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{Log}[2/(1 + e \cdot (x/d))]/(1 - c^2 \cdot x^2)), x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}\{p, 0\} \ \&\& \ \text{EqQ}[c^2 \cdot d^2 - e^2, 0]$

rule 6510 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((d) + (e) \cdot x^2), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot c \cdot d \cdot (p+1)), x] /;$ $\text{FreeQ}\{a, b, c, d, e, p, x\} \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{NeQ}\{p, -1\}$

rule 6542 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p \cdot (f \cdot x)^m / ((d) + (e) \cdot x^2), x_Symbol] \rightarrow \text{Simp}[f^2/e \cdot \text{Int}[(f \cdot x)^{m-2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x] - \text{Simp}[d \cdot (f^2/e) \cdot \text{Int}[(f \cdot x)^{m-2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2)], x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, x\} \ \&\& \ \text{GtQ}\{p, 0\} \ \&\& \ \text{GtQ}\{m, 1\}$

rule 6546

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*(x_)/((d_) + (e_.)*(x_)^2),
 x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/
 (c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 15.42 (sec) , antiderivative size = 1145, normalized size of antiderivative = 4.89

method	result	size
derivativeldivides	Expression too large to display	1145
default	Expression too large to display	1145
parts	Expression too large to display	1147

input

```
int(x*(a+b*arctanh(c*x^(1/2)))^3,x,method=_RETURNVERBOSE)
```

output

```
2/c^4*(1/4*a^3*c^4*x^2+b^3*(3/8*I*Pi*csgn(I/(1-(1+c*x^(1/2))^2/(c^2*x-1)))
^2*arctanh(c*x^(1/2))^2+3/16*I*Pi*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1))^3*arct
anh(c*x^(1/2))^2+3/16*I*Pi*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1)/(1-(1+c*x^(1/2)
))^2/(c^2*x-1))^3*arctanh(c*x^(1/2))^2-3/8*I*Pi*csgn(I/(1-(1+c*x^(1/2))^2
/(c^2*x-1)))^3*arctanh(c*x^(1/2))^2-1/4-3/16*I*Pi*csgn(I/(1-(1+c*x^(1/2))^
2/(c^2*x-1))) *csgn(I*(1+c*x^(1/2))^2/(c^2*x-1))*csgn(I*(1+c*x^(1/2))^2/(c
^2*x-1)/(1-(1+c*x^(1/2))^2/(c^2*x-1))) *arctanh(c*x^(1/2))^2+arctanh(c*x^(1/
2))^2+1/4*c*x^(1/2)-2*dilog(1-I*(1+c*x^(1/2))/(-c^2*x+1)^(1/2))-2*dilog(1+
I*(1+c*x^(1/2))/(-c^2*x+1)^(1/2))-1/4*arctanh(c*x^(1/2))^3+1/4*c^4*x^2*arc
tanh(c*x^(1/2))^3+1/4*arctanh(c*x^(1/2))^2*c^3*x^(3/2)+3/4*arctanh(c*x^(1/
2))^2*c*x^(1/2)+1/2*(c*x^(1/2)-1)*(1+c*x^(1/2))*arctanh(c*x^(1/2))-3/8*I*P
i*arctanh(c*x^(1/2))^2+3/8*I*Pi*csgn(I*(1+c*x^(1/2))/(-c^2*x+1)^(1/2))*csg
n(I*(1+c*x^(1/2))^2/(c^2*x-1))^2*arctanh(c*x^(1/2))^2+3/16*I*Pi*csgn(I/(1-
(1+c*x^(1/2))^2/(c^2*x-1))) *csgn(I*(1+c*x^(1/2))^2/(c^2*x-1)/(1-(1+c*x^(1/
2))^2/(c^2*x-1)))^2*arctanh(c*x^(1/2))^2+3/16*I*Pi*csgn(I*(1+c*x^(1/2))/(-
c^2*x+1)^(1/2))^2*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1))*arctanh(c*x^(1/2))^2-3
/16*I*Pi*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1))*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1
))/(1-(1+c*x^(1/2))^2/(c^2*x-1)))^2*arctanh(c*x^(1/2))^2+3/8*arctanh(c*x^(1
/2))^2*ln(c*x^(1/2)-1)-3/8*arctanh(c*x^(1/2))^2*ln(1+c*x^(1/2))-1/4*arctan
h(c*x^(1/2))*(1+c*x^(1/2))^2+1/2*(1+c*x^(1/2))*arctanh(c*x^(1/2))-2*arc...
```

Fricas [F]

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \int (b \operatorname{artanh}(c\sqrt{x}) + a)^3 x dx$$

input `integrate(x*(a+b*arctanh(c*x^(1/2)))^3,x, algorithm="fricas")`

output `integral(b^3*x*arctanh(c*sqrt(x))^3 + 3*a*b^2*x*arctanh(c*sqrt(x))^2 + 3*a^2*b*x*arctanh(c*sqrt(x)) + a^3*x, x)`

Sympy [F]

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \int x(a + b \operatorname{atanh}(c\sqrt{x}))^3 dx$$

input `integrate(x*(a+b*atanh(c*x**(1/2)))**3,x)`

output `Integral(x*(a + b*atanh(c*sqrt(x)))**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1184 vs. 2(191) = 382.

Time = 0.52 (sec) , antiderivative size = 1184, normalized size of antiderivative = 5.06

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \text{Too large to display}$$

input `integrate(x*(a+b*arctanh(c*x^(1/2)))^3,x, algorithm="maxima")`

output

```

1/2*a^3*x^2 - 1/32*a*b^2*c*((3*c^3*x^2 + 10*c*x - 2*(3*c^3*x^2 + 4*c^2*x^(
3/2) + 6*c*x + 12*sqrt(x))*log(c*sqrt(x) + 1))/c^4 - 14*log(c*sqrt(x) + 1)
/c^5 - 14*log(c*sqrt(x) - 1)/c^5) - 1/16*(12*x^2*log(c*sqrt(x) + 1) - c*((
3*c^3*x^2 - 4*c^2*x^(3/2) + 6*c*x - 12*sqrt(x))/c^4 + 12*log(c*sqrt(x) + 1
)/c^5))*a*b^2*log(-c*sqrt(x) + 1) + 1/16*(12*x^2*log(c*sqrt(x) + 1) - c*((
3*c^3*x^2 - 4*c^2*x^(3/2) + 6*c*x - 12*sqrt(x))/c^4 + 12*log(c*sqrt(x) + 1
)/c^5))*a^2*b - 1/16*(12*x^2*log(-c*sqrt(x) + 1) - c*((3*c^3*x^2 + 4*c^2*x
^(3/2) + 6*c*x + 12*sqrt(x))/c^4 + 12*log(c*sqrt(x) - 1)/c^5))*a^2*b + 1/1
92*(9*(8*log(-c*sqrt(x) + 1)^2 - 4*log(-c*sqrt(x) + 1) + 1)*(c*sqrt(x) - 1
)^4 + 32*(9*log(-c*sqrt(x) + 1)^2 - 6*log(-c*sqrt(x) + 1) + 2)*(c*sqrt(x)
- 1)^3 + 216*(2*log(-c*sqrt(x) + 1)^2 - 2*log(-c*sqrt(x) + 1) + 1)*(c*sqrt
(x) - 1)^2 + 288*(log(-c*sqrt(x) + 1)^2 - 2*log(-c*sqrt(x) + 1) + 2)*(c*sq
rt(x) - 1))*a*b^2/c^4 - 1/4608*(9*(32*log(-c*sqrt(x) + 1)^3 - 24*log(-c*sq
rt(x) + 1)^2 + 12*log(-c*sqrt(x) + 1) - 3)*(c*sqrt(x) - 1)^4 + 128*(9*log(
-c*sqrt(x) + 1)^3 - 9*log(-c*sqrt(x) + 1)^2 + 6*log(-c*sqrt(x) + 1) - 2)*(
c*sqrt(x) - 1)^3 + 432*(4*log(-c*sqrt(x) + 1)^3 - 6*log(-c*sqrt(x) + 1)^2
+ 6*log(-c*sqrt(x) + 1) - 3)*(c*sqrt(x) - 1)^2 + 1152*(log(-c*sqrt(x) + 1)
^3 - 3*log(-c*sqrt(x) + 1)^2 + 6*log(-c*sqrt(x) + 1) - 6)*(c*sqrt(x) - 1))
*b^3/c^4 + 2*(log(c*sqrt(x) + 1)*log(-1/2*c*sqrt(x) + 1/2) + dilog(1/2*c*s
qrt(x) + 1/2))*b^3/c^4 - 319/384*b^3*log(c*sqrt(x) - 1)/c^4 + 1/16*(25*...

```

Giac [F]

$$\int x(a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \int (b \operatorname{artanh}(c\sqrt{x}) + a)^3 x dx$$

input

```
integrate(x*(a+b*arctanh(c*x^(1/2)))^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*sqrt(x)) + a)^3*x, x)
```


3.204 $\int (a + \operatorname{arctanh}(c\sqrt{x}))^3 dx$

Optimal result	1642
Mathematica [A] (verified)	1643
Rubi [A] (verified)	1643
Maple [C] (warning: unable to verify)	1647
Fricas [F]	1647
Sympy [F]	1648
Maxima [F]	1648
Giac [F]	1649
Mupad [F(-1)]	1649
Reduce [F]	1649

Optimal result

Integrand size = 14, antiderivative size = 142

$$\int (a + \operatorname{arctanh}(c\sqrt{x}))^3 dx = \frac{3b(a + \operatorname{arctanh}(c\sqrt{x}))^2}{c^2} + \frac{3b\sqrt{x}(a + \operatorname{arctanh}(c\sqrt{x}))^2}{c} - \frac{(a + \operatorname{arctanh}(c\sqrt{x}))^3}{c^2} + x(a + \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{6b^2(a + \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1-c\sqrt{x}}\right)}{c^2} - \frac{3b^3 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-c\sqrt{x}}\right)}{c^2}$$

output

```
3*b*(a+b*arctanh(c*x^(1/2)))^2/c^2+3*b*x^(1/2)*(a+b*arctanh(c*x^(1/2)))^2/c-(a+b*arctanh(c*x^(1/2)))^3/c^2+x*(a+b*arctanh(c*x^(1/2)))^3-6*b^2*(a+b*arctanh(c*x^(1/2)))*ln(2/(1-c*x^(1/2)))/c^2-3*b^3*polylog(2,1-2/(1-c*x^(1/2)))/c^2
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.42

$$\int (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx$$

$$= \frac{6b^2(-1 + c\sqrt{x})(a + b + ac\sqrt{x}) \operatorname{arctanh}(c\sqrt{x})^2 + 2b^3(-1 + c^2x) \operatorname{arctanh}(c\sqrt{x})^3 + 6b \operatorname{arctanh}(c\sqrt{x}) (2$$

input

```
Integrate[(a + b*ArcTanh[c*Sqrt[x]])^3, x]
```

output

```
(6*b^2*(-1 + c*Sqrt[x])*(a + b + a*c*Sqrt[x])*ArcTanh[c*Sqrt[x]]^2 + 2*b^3
*(-1 + c^2*x)*ArcTanh[c*Sqrt[x]]^3 + 6*b*ArcTanh[c*Sqrt[x]]*(2*a*b*c*Sqrt[
x] + a^2*c^2*x - 2*b^2*Log[1 + E^(-2*ArcTanh[c*Sqrt[x]])]) + a*(6*a*b*c*Sq
rt[x] + 2*a^2*c^2*x + 3*a*b*Log[1 - c*Sqrt[x]] - 3*a*b*Log[1 + c*Sqrt[x]]
+ 6*b^2*Log[1 - c^2*x]) + 6*b^3*PolyLog[2, -E^(-2*ArcTanh[c*Sqrt[x]])])/(2
*c^2)
```

Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.20, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {6442, 6452, 6542, 6436, 6510, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx$$

$$\downarrow 6442$$

$$2 \int \sqrt{x} (a + b \operatorname{arctanh}(c\sqrt{x}))^3 d\sqrt{x}$$

$$\downarrow 6452$$

$$2 \left(\frac{1}{2} x (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{2} bc \int \frac{x (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x} \right)$$

↓ 6542

$$2 \left(\frac{1}{2}x(a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{3}{2}bc \left(\frac{\int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2x} d\sqrt{x}}{c^2} - \frac{\int (a + \operatorname{barctanh}(c\sqrt{x}))^2 d\sqrt{x}}{c^2} \right) \right)$$

↓ 6436

$$2 \left(\frac{1}{2}x(a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{3}{2}bc \left(\frac{\int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2x} d\sqrt{x}}{c^2} - \frac{\sqrt{x}(a + \operatorname{barctanh}(c\sqrt{x}))^2 - 2bc \int \frac{\sqrt{x}(a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2x} d\sqrt{x}}{c^2} \right) \right)$$

↓ 6510

$$2 \left(\frac{1}{2}x(a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{3}{2}bc \left(\frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x}(a + \operatorname{barctanh}(c\sqrt{x}))^2 - 2bc \int \frac{\sqrt{x}(a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2x} d\sqrt{x}}{c^2} \right) \right)$$

↓ 6546

$$2 \left(\frac{1}{2}x(a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{3}{2}bc \left(\frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x}(a + \operatorname{barctanh}(c\sqrt{x}))^2 - 2bc \left(\frac{\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{1 - c\sqrt{x}} d\sqrt{x}}{c} \right)}{c^2} \right) \right)$$

↓ 6470

$$2 \left(\frac{1}{2}x(a + \operatorname{barctanh}(c\sqrt{x}))^3 - \frac{3}{2}bc \left(\frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x}(a + \operatorname{barctanh}(c\sqrt{x}))^2 - 2bc \left(\frac{\log\left(\frac{2}{1 - c\sqrt{x}}\right)(a + b)}{c} \right)}{c^2} \right) \right)$$

↓ 2849

$$2 \left(\frac{1}{2}x(a + \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{2}bc \left(\frac{(a + \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x}(a + \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \left(\frac{b \int \frac{\log\left(\frac{2}{1-c\sqrt{x}}\right) d}{1-\frac{2}{1-c\sqrt{x}}} d}{c}} \right)}{\right)} \right)$$

↓ 2752

$$2 \left(\frac{1}{2}x(a + \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{3}{2}bc \left(\frac{(a + \operatorname{arctanh}(c\sqrt{x}))^3}{3bc^3} - \frac{\sqrt{x}(a + \operatorname{arctanh}(c\sqrt{x}))^2 - 2bc \left(\frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)(a+b)}{\frac{1-c\sqrt{x}}{c}} \right)}{\right)} \right)$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])^3,x]`

output `2*((x*(a + b*ArcTanh[c*Sqrt[x]])^3)/2 - (3*b*c*((a + b*ArcTanh[c*Sqrt[x]])^3/(3*b*c^3) - (Sqrt[x]*(a + b*ArcTanh[c*Sqrt[x]])^2 - 2*b*c*(-1/2*(a + b*ArcTanh[c*Sqrt[x]])^2/(b*c^2) + ((a + b*ArcTanh[c*Sqrt[x]])*Log[2/(1 - c*Sqrt[x]]))/c + (b*PolyLog[2, 1 - 2/(1 - c*Sqrt[x]]))/(2*c))/c)/c^2))/2)`

Defintions of rubi rules used

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 6436 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot (b \cdot x^n)^p, x_Symbol] \rightarrow \text{Simp}[x \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^p, x] - \text{Simp}[b \cdot c \cdot n \cdot p \cdot \text{Int}[x^n \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2n})], x], x] /;$ $\text{FreeQ}\{a, b, c, n, x\} \ \&\& \ \text{IGtQ}\{p, 0\} \ \&\& \ (\text{EqQ}\{n, 1\} \ || \ \text{EqQ}\{p, 1\})$

rule 6442 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot (b \cdot x^n)^p, x_Symbol] \rightarrow \text{With}\{k = \text{Denominator}[n]\}, \text{Simp}[k \cdot \text{Subst}[\text{Int}[x^{k-1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^{k \cdot n})]^p, x], x, x^{1/k}], x] /;$ $\text{FreeQ}\{a, b, c, x\} \ \&\& \ \text{IGtQ}\{p, 1\} \ \&\& \ \text{FractionQ}[n]$

rule 6452 $\text{Int}[(a + \text{ArcTanh}[c \cdot x^n] \cdot (b \cdot x^n)^p \cdot (x^m), x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^p / (m+1), x] - \text{Simp}[b \cdot c \cdot n \cdot (p/(m+1)) \cdot \text{Int}[x^{m+n} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2n})], x], x] /;$ $\text{FreeQ}\{a, b, c, m, n, x\} \ \&\& \ \text{IGtQ}\{p, 0\} \ \&\& \ (\text{EqQ}\{p, 1\} \ || \ (\text{EqQ}\{n, 1\} \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}\{m, -1\}$

rule 6470 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x)^p / ((d + (e \cdot x)), x_Symbol] \rightarrow \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{Log}[2/(1 + e \cdot (x/d))]/e), x] + \text{Simp}[b \cdot c \cdot (p/e) \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{Log}[2/(1 + e \cdot (x/d))]/(1 - c^2 \cdot x^2)), x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}\{p, 0\} \ \&\& \ \text{EqQ}[c^2 \cdot d^2 - e^2, 0]$

rule 6510 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x)^p / ((d + (e \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot c \cdot d \cdot (p+1)), x] /;$ $\text{FreeQ}\{a, b, c, d, e, p, x\} \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{NeQ}\{p, -1\}$

rule 6542 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x)^p \cdot ((f \cdot x)^m) / ((d + (e \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[f^2/e \cdot \text{Int}[(f \cdot x)^{m-2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x] - \text{Simp}[d \cdot (f^2/e) \cdot \text{Int}[(f \cdot x)^{m-2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2), x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, x\} \ \&\& \ \text{GtQ}\{p, 0\} \ \&\& \ \text{GtQ}\{m, 1\}$

rule 6546

```
Int[(((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^ (p_.)*(x_))/((d_) + (e_.)*(x_)^2),
 x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/
 (c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 14.08 (sec) , antiderivative size = 5673, normalized size of antiderivative = 39.95

method	result	size
derivativeldivides	Expression too large to display	5673
default	Expression too large to display	5673
parts	Expression too large to display	5674

input

```
int((a+b*arctanh(c*x^(1/2)))^3,x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [F]

$$\int (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \int (b \operatorname{arctanh}(c\sqrt{x}) + a)^3 dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))^3,x, algorithm="fricas")
```

output

```
integral(b^3*arctanh(c*sqrt(x))^3 + 3*a*b^2*arctanh(c*sqrt(x))^2 + 3*a^2*b
*arctanh(c*sqrt(x)) + a^3, x)
```

Sympy [F]

$$\int (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \int (a + b \operatorname{atanh}(c\sqrt{x}))^3 dx$$

input `integrate((a+b*atanh(c*x**(1/2)))**3,x)`

output `Integral((a + b*atanh(c*sqrt(x)))**3, x)`

Maxima [F]

$$\int (a + b \operatorname{arctanh}(c\sqrt{x}))^3 dx = \int (b \operatorname{artanh}(c\sqrt{x}) + a)^3 dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))^3,x, algorithm="maxima")`

output `3/2*(c*(2*sqrt(x)/c^2 - log(c*sqrt(x) + 1)/c^3 + log(c*sqrt(x) - 1)/c^3) + 2*x*arctanh(c*sqrt(x))*a^2*b + 3/4*(4*c*(2*sqrt(x)/c^2 - log(c*sqrt(x) + 1)/c^3 + log(c*sqrt(x) - 1)/c^3)*arctanh(c*sqrt(x)) + 4*x*arctanh(c*sqrt(x))^2 - (2*(log(c*sqrt(x) - 1) - 2)*log(c*sqrt(x) + 1) - log(c*sqrt(x) + 1)^2 - log(c*sqrt(x) - 1)^2 - 4*log(c*sqrt(x) - 1))/c^2)*a*b^2 + a^3*x - 1/32*b^3*(((4*log(-c*sqrt(x) + 1)^3 - 6*log(-c*sqrt(x) + 1)^2 + 6*log(-c*sqrt(x) + 1) - 3)*(c*sqrt(x) - 1)^2 + 8*(log(-c*sqrt(x) + 1)^3 - 3*log(-c*sqrt(x) + 1)^2 + 6*log(-c*sqrt(x) + 1) - 6)*(c*sqrt(x) - 1))/c^2 - 4*integrate(log(c*sqrt(x) + 1)^3 - 3*log(c*sqrt(x) + 1)^2*log(-c*sqrt(x) + 1) + 3*log(c*sqrt(x) + 1)*log(-c*sqrt(x) + 1)^2, x))`

output

```
(atanh(sqrt(x)*c)**3*b**3*c**2*x - atanh(sqrt(x)*c)**3*b**3 + 3*sqrt(x)*at  
anh(sqrt(x)*c)**2*b**3*c + 3*atanh(sqrt(x)*c)**2*a*b**2*c**2*x - 3*atanh(s  
qrt(x)*c)**2*a*b**2 + 6*sqrt(x)*atanh(sqrt(x)*c)*a*b**2*c + 3*atanh(sqrt(x  
)  
*c)*a**2*b*c**2*x - 3*atanh(sqrt(x)*c)*a**2*b + 6*atanh(sqrt(x)*c)*a*b**2  
+ 3*sqrt(x)*a**2*b*c + 3*int(atanh(sqrt(x)*c)/(c**2*x - 1),x)*b**3*c**2 +  
6*log(sqrt(x)*c - 1)*a*b**2 + a**3*c**2*x)/c**2
```

$$3.205 \quad \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x} dx$$

Optimal result	1651
Mathematica [C] (verified)	1652
Rubi [A] (verified)	1654
Maple [C] (warning: unable to verify)	1657
Fricas [F]	1658
Sympy [F]	1658
Maxima [F]	1658
Giac [F]	1659
Mupad [F(-1)]	1659
Reduce [F]	1660

Optimal result

Integrand size = 18, antiderivative size = 224

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x} dx = & 4 \operatorname{arctanh}\left(1 - \frac{2}{1 - c\sqrt{x}}\right) (a + b \operatorname{arctanh}(c\sqrt{x}))^3 \\ & - 3b(a + b \operatorname{arctanh}(c\sqrt{x}))^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - c\sqrt{x}}\right) \\ & + 3b(a + b \operatorname{arctanh}(c\sqrt{x}))^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - c\sqrt{x}}\right) \\ & + 3b^2(a + b \operatorname{arctanh}(c\sqrt{x})) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - c\sqrt{x}}\right) \\ & - 3b^2(a + b \operatorname{arctanh}(c\sqrt{x})) \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 - c\sqrt{x}}\right) \\ & - \frac{3}{2}b^3 \operatorname{PolyLog}\left(4, 1 - \frac{2}{1 - c\sqrt{x}}\right) \\ & + \frac{3}{2}b^3 \operatorname{PolyLog}\left(4, -1 + \frac{2}{1 - c\sqrt{x}}\right) \end{aligned}$$

output

```
-4*arctanh(-1+2/(1-c*x^(1/2)))*(a+b*arctanh(c*x^(1/2)))^3-3*b*(a+b*arctanh(c*x^(1/2)))^2*polylog(2,1-2/(1-c*x^(1/2)))+3*b*(a+b*arctanh(c*x^(1/2)))^2*polylog(2,-1+2/(1-c*x^(1/2)))+3*b^2*(a+b*arctanh(c*x^(1/2)))*polylog(3,1-2/(1-c*x^(1/2)))-3*b^2*(a+b*arctanh(c*x^(1/2)))*polylog(3,-1+2/(1-c*x^(1/2)))-3/2*b^3*polylog(4,1-2/(1-c*x^(1/2)))+3/2*b^3*polylog(4,-1+2/(1-c*x^(1/2)))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.32 (sec) , antiderivative size = 423, normalized size of antiderivative = 1.89

$$\begin{aligned}
 \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x} dx = & a^3 \log(x) \\
 & + 3a^2b(-\operatorname{PolyLog}(2, -c\sqrt{x}) + \operatorname{PolyLog}(2, c\sqrt{x})) \\
 & + 6ab^2 \left(\frac{i\pi^3}{24} - \frac{2}{3} \operatorname{arctanh}(c\sqrt{x})^3 \right. \\
 & \quad - \operatorname{arctanh}(c\sqrt{x})^2 \log(1 + e^{-2\operatorname{arctanh}(c\sqrt{x})}) \\
 & \quad + \operatorname{arctanh}(c\sqrt{x})^2 \log(1 - e^{2\operatorname{arctanh}(c\sqrt{x})}) \\
 & \quad + \operatorname{arctanh}(c\sqrt{x}) \operatorname{PolyLog}(2, -e^{-2\operatorname{arctanh}(c\sqrt{x})}) \\
 & \quad + \operatorname{arctanh}(c\sqrt{x}) \operatorname{PolyLog}(2, e^{2\operatorname{arctanh}(c\sqrt{x})}) \\
 & \quad + \frac{1}{2} \operatorname{PolyLog}(3, -e^{-2\operatorname{arctanh}(c\sqrt{x})}) \\
 & \quad \left. - \frac{1}{2} \operatorname{PolyLog}(3, e^{2\operatorname{arctanh}(c\sqrt{x})}) \right) \\
 & + \frac{1}{32} b^3 \left(\pi^4 - 32 \operatorname{arctanh}(c\sqrt{x})^4 \right. \\
 & \quad - 64 \operatorname{arctanh}(c\sqrt{x})^3 \log(1 + e^{-2\operatorname{arctanh}(c\sqrt{x})}) \\
 & \quad + 64 \operatorname{arctanh}(c\sqrt{x})^3 \log(1 - e^{2\operatorname{arctanh}(c\sqrt{x})}) \\
 & \quad + 96 \operatorname{arctanh}(c\sqrt{x})^2 \operatorname{PolyLog}(2, -e^{-2\operatorname{arctanh}(c\sqrt{x})}) \\
 & \quad + 96 \operatorname{arctanh}(c\sqrt{x})^2 \operatorname{PolyLog}(2, e^{2\operatorname{arctanh}(c\sqrt{x})}) \\
 & \quad + 96 \operatorname{arctanh}(c\sqrt{x}) \operatorname{PolyLog}(3, -e^{-2\operatorname{arctanh}(c\sqrt{x})}) \\
 & \quad - 96 \operatorname{arctanh}(c\sqrt{x}) \operatorname{PolyLog}(3, e^{2\operatorname{arctanh}(c\sqrt{x})}) \\
 & \quad + 48 \operatorname{PolyLog}(4, -e^{-2\operatorname{arctanh}(c\sqrt{x})}) \\
 & \quad \left. + 48 \operatorname{PolyLog}(4, e^{2\operatorname{arctanh}(c\sqrt{x})}) \right)
 \end{aligned}$$

input `Integrate[(a + b*ArcTanh[c*sqrt[x]])^3/x, x]`

output

```

a^3*Log[x] + 3*a^2*b*(-PolyLog[2, -(c*Sqrt[x])] + PolyLog[2, c*Sqrt[x]]) +
6*a*b^2*((I/24)*Pi^3 - (2*ArcTanh[c*Sqrt[x]]^3)/3 - ArcTanh[c*Sqrt[x]]^2*
Log[1 + E^(-2*ArcTanh[c*Sqrt[x]])] + ArcTanh[c*Sqrt[x]]^2*Log[1 - E^(2*Arc
Tanh[c*Sqrt[x]])] + ArcTanh[c*Sqrt[x]]*PolyLog[2, -E^(-2*ArcTanh[c*Sqrt[x]
]]) + ArcTanh[c*Sqrt[x]]*PolyLog[2, E^(2*ArcTanh[c*Sqrt[x]])] + PolyLog[3,
-E^(-2*ArcTanh[c*Sqrt[x]])]/2 - PolyLog[3, E^(2*ArcTanh[c*Sqrt[x]])]/2) +
(b^3*(Pi^4 - 32*ArcTanh[c*Sqrt[x]]^4 - 64*ArcTanh[c*Sqrt[x]]^3*Log[1 + E^
(-2*ArcTanh[c*Sqrt[x]])] + 64*ArcTanh[c*Sqrt[x]]^3*Log[1 - E^(2*ArcTanh[c*
Sqrt[x]])] + 96*ArcTanh[c*Sqrt[x]]^2*PolyLog[2, -E^(-2*ArcTanh[c*Sqrt[x]])]
] + 96*ArcTanh[c*Sqrt[x]]^2*PolyLog[2, E^(2*ArcTanh[c*Sqrt[x]])] + 96*ArcT
anh[c*Sqrt[x]]*PolyLog[3, -E^(-2*ArcTanh[c*Sqrt[x]])] - 96*ArcTanh[c*Sqrt[
x]]*PolyLog[3, E^(2*ArcTanh[c*Sqrt[x]])] + 48*PolyLog[4, -E^(-2*ArcTanh[c*
Sqrt[x]])] + 48*PolyLog[4, E^(2*ArcTanh[c*Sqrt[x]])])/32

```

Rubi [A] (verified)

Time = 1.20 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.17, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6450, 6448, 6614, 6620, 6624, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x} dx \\
& \quad \downarrow \text{6450} \\
& 2 \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{\sqrt{x}} d\sqrt{x} \\
& \quad \downarrow \text{6448} \\
& 2 \left(2 \operatorname{arctanh}\left(1 - \frac{2}{1 - c\sqrt{x}}\right) (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - 6bc \int \frac{\operatorname{arctanh}\left(1 - \frac{2}{1 - c\sqrt{x}}\right) (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{1 - c^2x} d\sqrt{x} \right) \\
& \quad \downarrow \text{6614}
\end{aligned}$$

$$2 \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - c\sqrt{x}} \right) (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - 6bc \left(\frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2 \log \left(2 - \frac{2}{1 - c\sqrt{x}} \right)}{1 - c^2 x} d\sqrt{x} - \right. \right.$$

↓ 6620

$$2 \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - c\sqrt{x}} \right) (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - 6bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - c\sqrt{x}} \right) (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{2c} \right. \right. \right.$$

↓ 6624

$$2 \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - c\sqrt{x}} \right) (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - 6bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - c\sqrt{x}} \right) (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{2c} \right. \right. \right.$$

↓ 7164

$$2 \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - c\sqrt{x}} \right) (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - 6bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - c\sqrt{x}} \right) (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{2c} \right. \right. \right.$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])^3/x,x]`

output `2*(2*ArcTanh[1 - 2/(1 - c*Sqrt[x])]*(a + b*ArcTanh[c*Sqrt[x]))^3 - 6*b*c*((((a + b*ArcTanh[c*Sqrt[x]])^2*PolyLog[2, 1 - 2/(1 - c*Sqrt[x])])/(2*c) - b*((a + b*ArcTanh[c*Sqrt[x]))*PolyLog[3, 1 - 2/(1 - c*Sqrt[x])])/(2*c) - (b*PolyLog[4, 1 - 2/(1 - c*Sqrt[x])])/(4*c)))/2 + (-1/2*((a + b*ArcTanh[c*Sqrt[x]])^2*PolyLog[2, -1 + 2/(1 - c*Sqrt[x])])/c + b*((a + b*ArcTanh[c*Sqrt[x]))*PolyLog[3, -1 + 2/(1 - c*Sqrt[x])])/(2*c) - (b*PolyLog[4, -1 + 2/(1 - c*Sqrt[x])])/(4*c)))/2)`

Definitions of rubi rules used

rule 6448 $\text{Int}[(a + \text{ArcTanh}[c \cdot x])^p / (x), x_Symbol] \rightarrow \text{Simp}[2 \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot \text{ArcTanh}[1 - 2/(1 - c \cdot x)], x] - \text{Simp}[2 \cdot b \cdot c \cdot p \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot \text{ArcTanh}[1 - 2/(1 - c \cdot x)] / (1 - c^2 \cdot x^2)], x, x] /;$
 $\text{FreeQ}\{a, b, c, x\} \ \&\& \ \text{IGtQ}[p, 1]$

rule 6450 $\text{Int}[(a + \text{ArcTanh}[c \cdot x]^n] \cdot (b + \text{ArcTanh}[c \cdot x])^p / (x), x_Symbol] \rightarrow \text{Simp}[1/n \cdot \text{Subst}[\text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p / x, x], x, x^n], x] /;$
 $\text{FreeQ}\{a, b, c, n, x\} \ \&\& \ \text{IGtQ}[p, 0]$

rule 6614 $\text{Int}[(\text{ArcTanh}[u] \cdot (a + \text{ArcTanh}[c \cdot x]) \cdot (b + \text{ArcTanh}[c \cdot x])^p) / ((d + e \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[1/2 \cdot \text{Int}[\text{Log}[1 + u] \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2)], x, x] - \text{Simp}[1/2 \cdot \text{Int}[\text{Log}[1 - u] \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2)], x, x] /;$
 $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{EqQ}[u^2 - (1 - 2/(1 - c \cdot x))^2, 0]$

rule 6620 $\text{Int}[(\text{Log}[u] \cdot (a + \text{ArcTanh}[c \cdot x]) \cdot (b + \text{ArcTanh}[c \cdot x])^p) / ((d + e \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{PolyLog}[2, 1 - u] / (2 \cdot c \cdot d)), x] + \text{Simp}[b \cdot (p/2) \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{PolyLog}[2, 1 - u] / (d + e \cdot x^2))], x, x] /;$
 $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{EqQ}[(1 - u)^2 - (1 - 2/(1 - c \cdot x))^2, 0]$

rule 6624 $\text{Int}[(a + \text{ArcTanh}[c \cdot x])^p \cdot \text{PolyLog}[k, u] / ((d + e \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{PolyLog}[k + 1, u] / (2 \cdot c \cdot d)), x] - \text{Simp}[b \cdot (p/2) \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{PolyLog}[k + 1, u] / (d + e \cdot x^2))], x, x] /;$
 $\text{FreeQ}\{a, b, c, d, e, k, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{EqQ}[u^2 - (1 - 2/(1 - c \cdot x))^2, 0]$

rule 7164 $\text{Int}[u \cdot \text{PolyLog}[n, v], x_Symbol] \rightarrow \text{With}\{w = \text{DerivativeDivides}[v, u \cdot v, x]\}, \text{Simp}[w \cdot \text{PolyLog}[n + 1, v], x] /;$
 $! \text{FalseQ}[w] /;$
 $\text{FreeQ}[n, x]$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 14.58 (sec) , antiderivative size = 1363, normalized size of antiderivative = 6.08

method	result	size
parts	Expression too large to display	1363
derivativedivides	Expression too large to display	1364
default	Expression too large to display	1364

input `int((a+b*arctanh(c*x^(1/2)))^3/x,x,method=_RETURNVERBOSE)`

output

```

a^3*ln(x)+b^3*(2*ln(c*x^(1/2))*arctanh(c*x^(1/2))^3-2*arctanh(c*x^(1/2))^3
*ln((1+c*x^(1/2))^2/(-c^2*x+1)-1)+2*arctanh(c*x^(1/2))^3*ln(1-(1+c*x^(1/2)
)/(-c^2*x+1)^(1/2))+6*arctanh(c*x^(1/2))^2*polylog(2,(1+c*x^(1/2))/(-c^2*x
+1)^(1/2))-12*arctanh(c*x^(1/2))*polylog(3,(1+c*x^(1/2))/(-c^2*x+1)^(1/2))
+12*polylog(4,(1+c*x^(1/2))/(-c^2*x+1)^(1/2))+2*arctanh(c*x^(1/2))^3*ln(1+
(1+c*x^(1/2))/(-c^2*x+1)^(1/2))+6*arctanh(c*x^(1/2))^2*polylog(2,-(1+c*x^(
1/2))/(-c^2*x+1)^(1/2))-12*arctanh(c*x^(1/2))*polylog(3,-(1+c*x^(1/2))/(-c
^2*x+1)^(1/2))+12*polylog(4,-(1+c*x^(1/2))/(-c^2*x+1)^(1/2))+I*Pi*csgn(I*(
-(1+c*x^(1/2))^2/(c^2*x-1)-1)/(1-(1+c*x^(1/2))^2/(c^2*x-1)))*(csgn(I*(-(1+
c*x^(1/2))^2/(c^2*x-1)-1))*csgn(I/(1-(1+c*x^(1/2))^2/(c^2*x-1)))-csgn(I*(-
(1+c*x^(1/2))^2/(c^2*x-1)-1))*csgn(I*(-(1+c*x^(1/2))^2/(c^2*x-1)-1)/(1-(1+
c*x^(1/2))^2/(c^2*x-1)))-csgn(I*(-(1+c*x^(1/2))^2/(c^2*x-1)-1)/(1-(1+c*x^(
1/2))^2/(c^2*x-1)))*csgn(I/(1-(1+c*x^(1/2))^2/(c^2*x-1)))+csgn(I*(-(1+c*x^(
1/2))^2/(c^2*x-1)-1)/(1-(1+c*x^(1/2))^2/(c^2*x-1)))^2)*arctanh(c*x^(1/2))
^3-3*arctanh(c*x^(1/2))^2*polylog(2,-(1+c*x^(1/2))^2/(-c^2*x+1))+3*arctanh
(c*x^(1/2))*polylog(3,-(1+c*x^(1/2))^2/(-c^2*x+1))-3/2*polylog(4,-(1+c*x^(
1/2))^2/(-c^2*x+1))+3*a*b^2*(2*ln(c*x^(1/2))*arctanh(c*x^(1/2))^2-2*arcta
nh(c*x^(1/2))*polylog(2,-(1+c*x^(1/2))^2/(-c^2*x+1))+polylog(3,-(1+c*x^(1/
2))^2/(-c^2*x+1))-2*arctanh(c*x^(1/2))^2*ln((1+c*x^(1/2))^2/(-c^2*x+1)-1)+
2*arctanh(c*x^(1/2))^2*ln(1-(1+c*x^(1/2))/(-c^2*x+1)^(1/2))+4*arctanh(c...

```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x} dx = \int \frac{(b \operatorname{artanh}(c\sqrt{x}) + a)^3}{x} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))^3/x,x, algorithm="fricas")`

output `integral((b^3*arctanh(c*sqrt(x))^3 + 3*a*b^2*arctanh(c*sqrt(x))^2 + 3*a^2*b*arctanh(c*sqrt(x)) + a^3)/x, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x} dx = \int \frac{(a + b \operatorname{atanh}(c\sqrt{x}))^3}{x} dx$$

input `integrate((a+b*atanh(c*x**(1/2)))**3/x,x)`

output `Integral((a + b*atanh(c*sqrt(x)))**3/x, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x} dx = \int \frac{(b \operatorname{artanh}(c\sqrt{x}) + a)^3}{x} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))^3/x,x, algorithm="maxima")`

output

```
1/8*b^3*integrate(log(c*sqrt(x) + 1)^3/x, x) - 3/8*b^3*integrate(log(c*sqrt(x) + 1)^2*log(-c*sqrt(x) + 1)/x, x) + 3/8*b^3*integrate(log(c*sqrt(x) + 1)*log(-c*sqrt(x) + 1)^2/x, x) - 1/8*b^3*integrate(log(-c*sqrt(x) + 1)^3/x, x) + 3/4*a*b^2*integrate(log(c*sqrt(x) + 1)^2/x, x) - 3/2*a*b^2*integrate(log(c*sqrt(x) + 1)*log(-c*sqrt(x) + 1)/x, x) + 3/4*a*b^2*integrate(log(-c*sqrt(x) + 1)^2/x, x) + 3/2*a^2*b*integrate(log(c*sqrt(x) + 1)/x, x) - 3/2*a^2*b*integrate(log(-c*sqrt(x) + 1)/x, x) + a^3*log(x)
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x} dx = \int \frac{(b \operatorname{artanh}(c\sqrt{x}) + a)^3}{x} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))^3/x,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*sqrt(x)) + a)^3/x, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x} dx = \int \frac{(a + b \operatorname{atanh}(c\sqrt{x}))^3}{x} dx$$

input

```
int((a + b*atanh(c*x^(1/2)))^3/x,x)
```

output

```
int((a + b*atanh(c*x^(1/2)))^3/x, x)
```


Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x} dx = 3 \left(\int \frac{\operatorname{atanh}(\sqrt{x}c)}{x} dx \right) a^2 b + \left(\int \frac{\operatorname{atanh}(\sqrt{x}c)^3}{x} dx \right) b^3$$

$$+ 3 \left(\int \frac{\operatorname{atanh}(\sqrt{x}c)^2}{x} dx \right) a b^2 + \log(x) a^3$$

input `int((a+b*atanh(c*x^(1/2)))^3/x,x)`

output `3*int(atanh(sqrt(x)*c)/x,x)*a**2*b + int(atanh(sqrt(x)*c)**3/x,x)*b**3 + 3*int(atanh(sqrt(x)*c)**2/x,x)*a*b**2 + log(x)*a**3`

$$3.206 \quad \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x^2} dx$$

Optimal result	1661
Mathematica [A] (verified)	1662
Rubi [A] (verified)	1662
Maple [C] (warning: unable to verify)	1665
Fricas [F]	1666
Sympy [F]	1667
Maxima [B] (verification not implemented)	1667
Giac [F]	1668
Mupad [F(-1)]	1668
Reduce [F]	1669

Optimal result

Integrand size = 18, antiderivative size = 142

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x^2} dx &= 3bc^2(a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{3bc(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{\sqrt{x}} \\ &\quad + c^2(a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x} \\ &\quad + 6b^2c^2(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(2 - \frac{2}{1 + c\sqrt{x}}\right) \\ &\quad - 3b^3c^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + c\sqrt{x}}\right) \end{aligned}$$

output

```
3*b*c^2*(a+b*arctanh(c*x^(1/2)))^2-3*b*c*(a+b*arctanh(c*x^(1/2)))^2/x^(1/2)
)+c^2*(a+b*arctanh(c*x^(1/2)))^3-(a+b*arctanh(c*x^(1/2)))^3/x+6*b^2*c^2*(a
+b*arctanh(c*x^(1/2)))*ln(2-2/(1+c*x^(1/2)))-3*b^3*c^2*polylog(2,-1+2/(1+c
*x^(1/2)))
```

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.62

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x^2} dx$$

$$= \frac{6b^2(-1 + c\sqrt{x})(a + ac\sqrt{x} + bc\sqrt{x}) \operatorname{arctanh}(c\sqrt{x})^2 + 2b^3(-1 + c^2x) \operatorname{arctanh}(c\sqrt{x})^3 - 6b \operatorname{arctanh}(c\sqrt{x})}{x^2}$$

input `Integrate[(a + b*ArcTanh[c*Sqrt[x]])^3/x^2,x]`

output `(6*b^2*(-1 + c*Sqrt[x])*(a + a*c*Sqrt[x] + b*c*Sqrt[x])*ArcTanh[c*Sqrt[x]]^2 + 2*b^3*(-1 + c^2*x)*ArcTanh[c*Sqrt[x]]^3 - 6*b*ArcTanh[c*Sqrt[x]]*(a^2 + 2*a*b*c*Sqrt[x] - 2*b^2*c^2*x*Log[1 - E^(-2*ArcTanh[c*Sqrt[x]])])) + a*(-2*a^2 - 6*a*b*c*Sqrt[x] - 3*a*b*c^2*x*Log[1 - c*Sqrt[x]] + 3*a*b*c^2*x*Log[1 + c*Sqrt[x]] + 12*b^2*c^2*x*Log[(c*Sqrt[x])/Sqrt[1 - c^2*x]]) - 6*b^3*c^2*x*PolyLog[2, E^(-2*ArcTanh[c*Sqrt[x]])])/(2*x)`

Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.06, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {6454, 6452, 6544, 6452, 6510, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x^2} dx$$

$$\downarrow 6454$$

$$2 \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x^{3/2}} d\sqrt{x}$$

$$\downarrow 6452$$

$$2 \left(\frac{3}{2} bc \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{x(1 - c^2x)} d\sqrt{x} - \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{2x} \right)$$

↓ 6544

$$2 \left(\frac{3}{2} bc \left(c^2 \int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x} + \int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{x} d\sqrt{x} \right) - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{2x} \right)$$

↓ 6452

$$2 \left(\frac{3}{2} bc \left(c^2 \int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1 - c^2 x} d\sqrt{x} + 2bc \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1 - c^2 x)} d\sqrt{x} - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{\sqrt{x}} \right) - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{2x} \right)$$

↓ 6510

$$2 \left(\frac{3}{2} bc \left(2bc \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1 - c^2 x)} d\sqrt{x} + \frac{c(a + \operatorname{barctanh}(c\sqrt{x}))^3}{3b} - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{\sqrt{x}} \right) - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{2x} \right)$$

↓ 6550

$$2 \left(\frac{3}{2} bc \left(2bc \left(\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{(\sqrt{xc} + 1)\sqrt{x}} d\sqrt{x} + \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} \right) + \frac{c(a + \operatorname{barctanh}(c\sqrt{x}))^3}{3b} - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{\sqrt{x}} \right) \right)$$

↓ 6494

$$2 \left(\frac{3}{2} bc \left(2bc \left(-bc \int \frac{\log\left(2 - \frac{2}{\sqrt{xc} + 1}\right)}{1 - c^2 x} d\sqrt{x} + \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} + \log\left(2 - \frac{2}{c\sqrt{x} + 1}\right) (a + \operatorname{barctanh}(c\sqrt{x})) \right) \right)$$

↓ 2897

$$2 \left(\frac{3}{2} bc \left(2bc \left(\frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} + \log\left(2 - \frac{2}{c\sqrt{x} + 1}\right) (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{2} b \operatorname{PolyLog}\left(2, \frac{2}{\sqrt{xc} + 1} - 1\right) \right) \right)$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])^3/x^2,x]`

output `2*(-1/2*(a + b*ArcTanh[c*Sqrt[x]])^3/x + (3*b*c*(-((a + b*ArcTanh[c*Sqrt[x]])^2/Sqrt[x]) + (c*(a + b*ArcTanh[c*Sqrt[x]])^3)/(3*b) + 2*b*c*((a + b*ArcTanh[c*Sqrt[x]])^2/(2*b) + (a + b*ArcTanh[c*Sqrt[x]])*Log[2 - 2/(1 + c*Sqrt[x]]) - (b*PolyLog[2, -1 + 2/(1 + c*Sqrt[x]])/2)))/2)`

Defintions of rubi rules used

rule 2897 `Int[Log[u_]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6494 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((x_)*((d_) + (e_)*(x_))), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6510 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6544 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)*((f_)*(x_)^(m_))/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]`

rule 6550

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)/((x_.)*((d_.) + (e_.)*(x_)^2)),
 x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/
d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 15.12 (sec) , antiderivative size = 4700, normalized size of antiderivative = 33.10

method	result	size
derivativedivides	Expression too large to display	4700
default	Expression too large to display	4700
parts	Expression too large to display	4702

input

```
int((a+b*arctanh(c*x^(1/2)))^3/x^2,x,method=_RETURNVERBOSE)
```

output

```

2*c^2*(-1/2*a^3/c^2/x+b^3*(3/8*I*Pi*csgn(I/(1-(1+c*x^(1/2))^2/(c^2*x-1)))*
csgn(I*(1+c*x^(1/2))^2/(c^2*x-1))*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1)/(1-(1+c
*x^(1/2))^2/(c^2*x-1)))*dilog(1+(1+c*x^(1/2))/(-c^2*x+1)^(1/2))-3/2*arctan
h(c*x^(1/2))^2+3/8*I*Pi*csgn(I/(1-(1+c*x^(1/2))^2/(c^2*x-1)))*csgn(I*(1+c*
*x^(1/2))^2/(c^2*x-1)/(1-(1+c*x^(1/2))^2/(c^2*x-1)))*2*arctanh(c*x^(1/2))*l
n(1-(1+c*x^(1/2))/(-c^2*x+1)^(1/2))-3/8*I*Pi*csgn(I*(1+c*x^(1/2))^2/(c^2*x
-1))*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1)/(1-(1+c*x^(1/2))^2/(c^2*x-1)))*2*arc
tanh(c*x^(1/2))*ln(1-(1+c*x^(1/2))/(-c^2*x+1)^(1/2))+3/8*I*Pi*csgn(I/(1-(1
+c*x^(1/2))^2/(c^2*x-1)))*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1))*csgn(I*(1+c*x^
(1/2))^2/(c^2*x-1)/(1-(1+c*x^(1/2))^2/(c^2*x-1)))*arctanh(c*x^(1/2))^2-3/8
*I*Pi*csgn(I/(1-(1+c*x^(1/2))^2/(c^2*x-1)))*csgn(I*(1+c*x^(1/2))^2/(c^2*x-
1))*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1)/(1-(1+c*x^(1/2))^2/(c^2*x-1)))*polylo
g(2,(1+c*x^(1/2))/(-c^2*x+1)^(1/2))-3/8*I*Pi*csgn(I/(1-(1+c*x^(1/2))^2/(c^
2*x-1)))*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1))*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1
)/(1-(1+c*x^(1/2))^2/(c^2*x-1)))*polylog(2,-(1+c*x^(1/2))/(-c^2*x+1)^(1/2)
)-3/8*I*Pi*csgn(I/(1-(1+c*x^(1/2))^2/(c^2*x-1)))*csgn(I*(1+c*x^(1/2))^2/(c
^2*x-1))*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1)/(1-(1+c*x^(1/2))^2/(c^2*x-1)))*d
ilog((1+c*x^(1/2))/(-c^2*x+1)^(1/2))+3/4*I*Pi*csgn(I*(1+c*x^(1/2))/(-c^2*x
+1)^(1/2))*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1))^2*arctanh(c*x^(1/2))*ln(1-(1+
c*x^(1/2))/(-c^2*x+1)^(1/2))+3/8*I*Pi*csgn(I*(1+c*x^(1/2))/(-c^2*x+1)^(...

```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x^2} dx = \int \frac{(b \operatorname{arctanh}(c\sqrt{x}) + a)^3}{x^2} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))^3/x^2,x, algorithm="fricas")
```

output

```
integral((b^3*arctanh(c*sqrt(x))^3 + 3*a*b^2*arctanh(c*sqrt(x))^2 + 3*a^2*
b*arctanh(c*sqrt(x)) + a^3)/x^2, x)
```

Sympy [F]

$$\int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{x^2} dx = \int \frac{(a + b \operatorname{atanh}(c\sqrt{x}))^3}{x^2} dx$$

input `integrate((a+b*atanh(c*x**(1/2)))**3/x**2,x)`

output `Integral((a + b*atanh(c*sqrt(x)))**3/x**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 528 vs. $2(125) = 250$.

Time = 1.02 (sec) , antiderivative size = 528, normalized size of antiderivative = 3.72

$$\begin{aligned} & \int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{x^2} dx \\ &= -3 \left(\log(c\sqrt{x} + 1) \log\left(-\frac{1}{2}c\sqrt{x} + \frac{1}{2}\right) + \operatorname{Li}_2\left(\frac{1}{2}c\sqrt{x} + \frac{1}{2}\right) \right) b^3 c^2 \\ & \quad - 3 \left(\log(c\sqrt{x}) \log(-c\sqrt{x} + 1) + \operatorname{Li}_2(-c\sqrt{x} + 1) \right) b^3 c^2 \\ & \quad + 3 \left(\log(c\sqrt{x} + 1) \log(-c\sqrt{x}) + \operatorname{Li}_2(c\sqrt{x} + 1) \right) b^3 c^2 - 3ab^2c^2 \log(c\sqrt{x} - 1) \\ & \quad - \frac{3}{4} \left(\left(2c \log(c\sqrt{x} - 1) - c \log(x) + \frac{2}{\sqrt{x}} \right) c - \frac{2 \log(-c\sqrt{x} + 1)}{x} \right) a^2 b \\ & \quad - \frac{a^3}{x} + \frac{3}{2} (a^2bc^2 - 2ab^2c^2) \log(c\sqrt{x} + 1) - \frac{3}{4} (a^2bc^2 - 4ab^2c^2) \log(x) \\ & \quad - \frac{12a^2bc\sqrt{x} - (b^3c^2x - b^3) \log(c\sqrt{x} + 1)^3 + (b^3c^2x - b^3) \log(-c\sqrt{x} + 1)^3 + 6(b^3c\sqrt{x} + ab^2 - (ab^2c^2 - \dots)}{x^2} \end{aligned}$$

input `integrate((a+b*arctanh(c*x^(1/2)))^3/x^2,x, algorithm="maxima")`

output

```
-3*(log(c*sqrt(x) + 1)*log(-1/2*c*sqrt(x) + 1/2) + dilog(1/2*c*sqrt(x) + 1/2))*b^3*c^2 - 3*(log(c*sqrt(x))*log(-c*sqrt(x) + 1) + dilog(-c*sqrt(x) + 1))*b^3*c^2 + 3*(log(c*sqrt(x) + 1)*log(-c*sqrt(x)) + dilog(c*sqrt(x) + 1))*b^3*c^2 - 3*a*b^2*c^2*log(c*sqrt(x) - 1) - 3/4*((2*c*log(c*sqrt(x) - 1) - c*log(x) + 2/sqrt(x))*c - 2*log(-c*sqrt(x) + 1)/x)*a^2*b - a^3/x + 3/2*(a^2*b*c^2 - 2*a*b^2*c^2)*log(c*sqrt(x) + 1) - 3/4*(a^2*b*c^2 - 4*a*b^2*c^2)*log(x) - 1/8*(12*a^2*b*c*sqrt(x) - (b^3*c^2*x - b^3)*log(c*sqrt(x) + 1)^3 + (b^3*c^2*x - b^3)*log(-c*sqrt(x) + 1)^3 + 6*(b^3*c*sqrt(x) + a*b^2 - (a*b^2*c^2 - b^3*c^2)*x)*log(c*sqrt(x) + 1)^2 + 3*(2*b^3*c*sqrt(x) + 2*a*b^2 - 2*(a*b^2*c^2 + b^3*c^2)*x - (b^3*c^2*x - b^3)*log(c*sqrt(x) + 1))*log(-c*sqrt(x) + 1)^2 + 12*(2*a*b^2*c*sqrt(x) + a^2*b)*log(c*sqrt(x) + 1) - 3*(8*a*b^2*c*sqrt(x) - (b^3*c^2*x - b^3)*log(c*sqrt(x) + 1)^2 + 4*(b^3*c*sqrt(x) + a*b^2 - (a*b^2*c^2 - b^3*c^2)*x)*log(c*sqrt(x) + 1))*log(-c*sqrt(x) + 1))/x
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x^2} dx = \int \frac{(b \operatorname{artanh}(c\sqrt{x}) + a)^3}{x^2} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))^3/x^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*sqrt(x)) + a)^3/x^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x^2} dx = \int \frac{(a + b \operatorname{atanh}(c\sqrt{x}))^3}{x^2} dx$$

input

```
int((a + b*atanh(c*x^(1/2)))^3/x^2,x)
```

output

```
int((a + b*atanh(c*x^(1/2)))^3/x^2, x)
```


$$3.207 \quad \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x^3} dx$$

Optimal result	1670
Mathematica [A] (verified)	1671
Rubi [A] (verified)	1671
Maple [C] (warning: unable to verify)	1675
Fricas [F]	1676
Sympy [F]	1677
Maxima [B] (verification not implemented)	1677
Giac [F]	1678
Mupad [F(-1)]	1679
Reduce [F]	1679

Optimal result

Integrand size = 18, antiderivative size = 234

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x^3} dx = & -\frac{b^3 c^3}{2\sqrt{x}} + \frac{1}{2} b^3 c^4 \operatorname{arctanh}(c\sqrt{x}) \\ & - \frac{b^2 c^2 (a + b \operatorname{arctanh}(c\sqrt{x}))}{2x} \\ & + 2bc^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^2 - \frac{bc(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{2x^{3/2}} \\ & - \frac{3bc^3 (a + b \operatorname{arctanh}(c\sqrt{x}))^2}{2\sqrt{x}} \\ & + \frac{1}{2} c^4 (a + b \operatorname{arctanh}(c\sqrt{x}))^3 - \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{2x^2} \\ & + 4b^2 c^4 (a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(2 - \frac{2}{1 + c\sqrt{x}}\right) \\ & - 2b^3 c^4 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + c\sqrt{x}}\right) \end{aligned}$$

output

```
-1/2*b^3*c^3/x^(1/2)+1/2*b^3*c^4*arctanh(c*x^(1/2))-1/2*b^2*c^2*(a+b*arctanh(c*x^(1/2)))/x+2*b*c^4*(a+b*arctanh(c*x^(1/2)))^2-1/2*b*c*(a+b*arctanh(c*x^(1/2)))^2/x^(3/2)-3/2*b*c^3*(a+b*arctanh(c*x^(1/2)))^2/x^(1/2)+1/2*c^4*(a+b*arctanh(c*x^(1/2)))^3-1/2*(a+b*arctanh(c*x^(1/2)))^3/x^2+4*b^2*c^4*(a+b*arctanh(c*x^(1/2)))*ln(2-2/(1+c*x^(1/2)))-2*b^3*c^4*polylog(2,-1+2/(1+c*x^(1/2)))
```

Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 333, normalized size of antiderivative = 1.42

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x^3} dx =$$

$$\frac{2a^3 + 2a^2bc\sqrt{x} + 2ab^2c^2x + 6a^2bc^3x^{3/2} + 2b^3c^3x^{3/2} - 2ab^2c^4x^2 - 2b^2(bc\sqrt{x}(-1 - 3c^2x + 4c^3x^{3/2}) + \dots)}{x^3}$$

input

```
Integrate[(a + b*ArcTanh[c*Sqrt[x]])^3/x^3,x]
```

output

```
-1/4*(2*a^3 + 2*a^2*b*c*Sqrt[x] + 2*a*b^2*c^2*x + 6*a^2*b*c^3*x^(3/2) + 2*b^3*c^3*x^(3/2) - 2*a*b^2*c^4*x^2 - 2*b^2*(b*c*Sqrt[x]*(-1 - 3*c^2*x + 4*c^3*x^(3/2)) + 3*a*(-1 + c^4*x^2))*ArcTanh[c*Sqrt[x]]^2 - 2*b^3*(-1 + c^4*x^2)*ArcTanh[c*Sqrt[x]]^3 + 2*b*ArcTanh[c*Sqrt[x]]*(3*a^2 + b^2*c^2*x*(1 - c^2*x) + 2*a*b*c*Sqrt[x]*(1 + 3*c^2*x) - 8*b^2*c^4*x^2*Log[1 - E^(-2*ArcTanh[c*Sqrt[x]])]) + 3*a^2*b*c^4*x^2*Log[1 - c*Sqrt[x]] - 3*a^2*b*c^4*x^2*Log[1 + c*Sqrt[x]] - 16*a*b^2*c^4*x^2*Log[(c*Sqrt[x])/Sqrt[1 - c^2*x]] + 8*b^3*c^4*x^2*PolyLog[2, E^(-2*ArcTanh[c*Sqrt[x]])])/x^2
```

Rubi [A] (verified)Time = 2.01 (sec) , antiderivative size = 306, normalized size of antiderivative = 1.31, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {6454, 6452, 6544, 6452, 6544, 6452, 264, 219, 6510, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{x^3} dx \\
& \quad \downarrow \text{6454} \\
& 2 \int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{x^{5/2}} d\sqrt{x} \\
& \quad \downarrow \text{6452} \\
& 2 \left(\frac{3}{4} bc \int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{x^2(1-c^2x)} d\sqrt{x} - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{4x^2} \right) \\
& \quad \downarrow \text{6544} \\
& 2 \left(\frac{3}{4} bc \left(c^2 \int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{x(1-c^2x)} d\sqrt{x} + \int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{x^2} d\sqrt{x} \right) - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{4x^2} \right) \\
& \quad \downarrow \text{6452} \\
& 2 \left(\frac{3}{4} bc \left(\frac{2}{3} bc \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(1-c^2x)} d\sqrt{x} + c^2 \int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{x(1-c^2x)} d\sqrt{x} - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{3x^{3/2}} \right) - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^3}{4x^2} \right) \\
& \quad \downarrow \text{6544} \\
& 2 \left(\frac{3}{4} bc \left(\frac{2}{3} bc \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} + \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}} d\sqrt{x} \right) + c^2 \left(c^2 \int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1-c^2x} d\sqrt{x} \right) \right) \right. \\
& \quad \downarrow \text{6452} \\
& 2 \left(\frac{3}{4} bc \left(c^2 \left(c^2 \int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1-c^2x} d\sqrt{x} + 2bc \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{\sqrt{x}} \right) + \frac{2}{3} bc \right) \right. \\
& \quad \downarrow \text{264} \\
& 2 \left(\frac{3}{4} bc \left(c^2 \left(c^2 \int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1-c^2x} d\sqrt{x} + 2bc \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{\sqrt{x}} \right) + \frac{2}{3} bc \right) \right. \\
& \quad \downarrow \text{219} \\
& 2 \left(\frac{3}{4} bc \left(c^2 \left(c^2 \int \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{1-c^2x} d\sqrt{x} + 2bc \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} - \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{\sqrt{x}} \right) + \frac{2}{3} bc \right) \right.
\end{aligned}$$

↓ 6510

$$2\left(\frac{3}{4}bc\left(c^2\left(2bc\int\frac{a+\operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)}d\sqrt{x}+\frac{c(a+\operatorname{barctanh}(c\sqrt{x}))^3}{3b}-\frac{(a+\operatorname{barctanh}(c\sqrt{x}))^2}{\sqrt{x}}\right)+\frac{2}{3}bc\left(c^2\int\right.\right.$$

↓ 6550

$$2\left(\frac{3}{4}bc\left(c^2\left(2bc\left(\int\frac{a+\operatorname{barctanh}(c\sqrt{x})}{(\sqrt{xc}+1)\sqrt{x}}d\sqrt{x}+\frac{(a+\operatorname{barctanh}(c\sqrt{x}))^2}{2b}\right)+\frac{c(a+\operatorname{barctanh}(c\sqrt{x}))^3}{3b}-\frac{(a+\operatorname{barctanh}(c\sqrt{x}))^2}{\sqrt{x}}\right)\right.\right.$$

↓ 6494

$$2\left(\frac{3}{4}bc\left(c^2\left(2bc\left(-bc\int\frac{\log\left(2-\frac{2}{\sqrt{xc}+1}\right)}{1-c^2x}d\sqrt{x}+\frac{(a+\operatorname{barctanh}(c\sqrt{x}))^2}{2b}+\log\left(2-\frac{2}{c\sqrt{x}+1}\right)(a+\operatorname{barctanh}(c\sqrt{x}))\right)\right.\right.\right.$$

↓ 2897

$$2\left(\frac{3}{4}bc\left(c^2\left(2bc\left(\frac{(a+\operatorname{barctanh}(c\sqrt{x}))^2}{2b}+\log\left(2-\frac{2}{c\sqrt{x}+1}\right)(a+\operatorname{barctanh}(c\sqrt{x}))-\frac{1}{2}b\operatorname{PolyLog}\left(2,\frac{2}{\sqrt{xc}+1}\right)\right)\right)\right.\right.$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])^3/x^3,x]`

output

```
2*(-1/4*(a + b*ArcTanh[c*Sqrt[x]])^3/x^2 + (3*b*c*(-1/3*(a + b*ArcTanh[c*Sqrt[x]])^2/x^(3/2) + c^2*(-((a + b*ArcTanh[c*Sqrt[x]])^2/Sqrt[x]) + (c*(a + b*ArcTanh[c*Sqrt[x]])^3)/(3*b) + 2*b*c*((a + b*ArcTanh[c*Sqrt[x]])^2/(2*b) + (a + b*ArcTanh[c*Sqrt[x]])*Log[2 - 2/(1 + c*Sqrt[x])]) - (b*PolyLog[2, -1 + 2/(1 + c*Sqrt[x])])/2)) + (2*b*c*(-1/2*(a + b*ArcTanh[c*Sqrt[x]])/x + (b*c*(-(1/Sqrt[x]) + c*ArcTanh[c*Sqrt[x])))/2 + c^2*((a + b*ArcTanh[c*Sqrt[x]])^2/(2*b) + (a + b*ArcTanh[c*Sqrt[x]])*Log[2 - 2/(1 + c*Sqrt[x])]) - (b*PolyLog[2, -1 + 2/(1 + c*Sqrt[x])])/2))/3)/4
```

Definitions of rubi rules used

rule 219 $\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 264 $\text{Int}[(c_)*(x_))^{(m_)}*((a_ + (b_)*(x_)^2)^{(p_)}), x_Symbol] \rightarrow \text{Simp}[(c*x)^{(m+1)}*((a + b*x^2)^{(p+1)}/(a*c*(m+1))), x] - \text{Simp}[b*((m+2*p+3)/(a*c^{2*(m+1)})) \ \text{Int}[(c*x)^{(m+2)}*(a + b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 2897 $\text{Int}[\text{Log}[u_]*(Pq_)^{(m_)}), x_Symbol] \rightarrow \text{With}\{C = \text{FullSimplify}[Pq^m*((1-u)/D[u, x])]\}, \text{Simp}[C*\text{PolyLog}[2, 1-u], x] /; \text{FreeQ}[C, x] /; \text{IntegerQ}[m] \ \&\& \ \text{PolyQ}[Pq, x] \ \&\& \ \text{RationalFunctionQ}[u, x] \ \&\& \ \text{LeQ}[\text{RationalFunctionExponents}[u, x][[2]], \text{Expon}[Pq, x]]$

rule 6452 $\text{Int}[(a_ + \text{ArcTanh}[c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}), x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \ \text{Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)}/(1-c^2*x^{(2*n)})), x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 6454 $\text{Int}[(a_ + \text{ArcTanh}[c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}), x_Symbol] \rightarrow \text{Simp}[1/n \ \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n] - 1)*(a + b*\text{ArcTanh}[c*x])^p}, x], x, x^n], x] /; \text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$

rule 6494 $\text{Int}[(a_ + \text{ArcTanh}[c_)*(x_)]*(b_))^{(p_)}((x_)*((d_ + (e_)*(x_))), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^p*(\text{Log}[2 - 2/(1 + e*(x/d))]/d), x] - \text{Simp}[b*c*(p/d) \ \text{Int}[(a + b*\text{ArcTanh}[c*x])^{(p-1)}*(\text{Log}[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d^2 - e^2, 0]$

rule 6510 $\text{Int}[\left((a_{.}) + \text{ArcTanh}[c_{.}(x_{.})](b_{.})\right)^{p_{.}} / \left((d_{.}) + (e_{.})(x_{.})^2\right), x_{\text{Symbol}}] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot c \cdot d \cdot (p+1)), x] /;$ $\text{FreeQ}[\{a, b, c, d, e, p\}, x]$ && $\text{EqQ}[c^2 \cdot d + e, 0]$ && $\text{NeQ}[p, -1]$

rule 6544 $\text{Int}[\left((a_{.}) + \text{ArcTanh}[c_{.}(x_{.})](b_{.})\right)^{p_{.}} \cdot \left((f_{.})(x_{.})\right)^{m_{.}} / \left((d_{.}) + (e_{.})(x_{.})^2\right), x_{\text{Symbol}}] \rightarrow \text{Simp}[1/d \int [(f \cdot x)^m \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x] - \text{Simp}[e/(d \cdot f^2) \int [(f \cdot x)^{m+2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2), x], x] /;$ $\text{FreeQ}[\{a, b, c, d, e, f\}, x]$ && $\text{GtQ}[p, 0]$ && $\text{LtQ}[m, -1]$

rule 6550 $\text{Int}[\left((a_{.}) + \text{ArcTanh}[c_{.}(x_{.})](b_{.})\right)^{p_{.}} / \left((x_{.}) \cdot \left((d_{.}) + (e_{.})(x_{.})^2\right)\right), x_{\text{Symbol}}] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot d \cdot (p+1)), x] + \text{Simp}[1/d \int [(a + b \cdot \text{ArcTanh}[c \cdot x])^p / (x \cdot (1 + c \cdot x)), x], x] /;$ $\text{FreeQ}[\{a, b, c, d, e\}, x]$ && $\text{EqQ}[c^2 \cdot d + e, 0]$ && $\text{GtQ}[p, 0]$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 16.69 (sec) , antiderivative size = 1219, normalized size of antiderivative = 5.21

method	result	size
derivativedivides	Expression too large to display	1219
default	Expression too large to display	1219
parts	Expression too large to display	1275

input $\text{int}((a+b \cdot \arctanh(c \cdot x^{1/2}))^3 / x^3, x, \text{method}=_RETURNVERBOSE)$

output

```

2*c^4*(-1/4*b^3*arctanh(c*x^(1/2))^2/c^3/x^(3/2)-1/4*b^3/c^4/x^2*arctanh(c
*x^(1/2))^3-1/4*b^3*arctanh(c*x^(1/2))/c^2/x+3/8*I*b^3*Pi*arctanh(c*x^(1/2
))^2-3/8*I*b^3*Pi*csgn(I*(1+c*x^(1/2))/(-c^2*x+1)^(1/2))*csgn(I*(1+c*x^(1/
2))^2/(c^2*x-1))^2*arctanh(c*x^(1/2))^2+3/16*I*b^3*Pi*csgn(I*(1+c*x^(1/2))
^2/(c^2*x-1))*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1)/(1-(1+c*x^(1/2))^2/(c^2*x-
1)))^2*arctanh(c*x^(1/2))^2-3/16*I*b^3*Pi*csgn(I/(1-(1+c*x^(1/2))^2/(c^2*x-
1)))^2*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1)/(1-(1+c*x^(1/2))^2/(c^2*x-1)))^2*arc
tanh(c*x^(1/2))^2-3/16*I*b^3*Pi*csgn(I*(1+c*x^(1/2))/(-c^2*x+1)^(1/2))^2*c
sgn(I*(1+c*x^(1/2))^2/(c^2*x-1))*arctanh(c*x^(1/2))^2+3/16*I*b^3*Pi*csgn(I
/(1-(1+c*x^(1/2))^2/(c^2*x-1)))^2*csgn(I*(1+c*x^(1/2))^2/(c^2*x-1))*csgn(I*(
1+c*x^(1/2))^2/(c^2*x-1)/(1-(1+c*x^(1/2))^2/(c^2*x-1)))^2*arctanh(c*x^(1/2))
^2+3*a*b^2*(-1/4/c^4/x^2*arctanh(c*x^(1/2))^2-1/6*arctanh(c*x^(1/2))/c^3/x
^(3/2)-1/2*arctanh(c*x^(1/2))/c/x^(1/2)+1/4*arctanh(c*x^(1/2))*ln(1+c*x^(1
/2))-1/4*arctanh(c*x^(1/2))*ln(c*x^(1/2)-1)-1/16*ln(c*x^(1/2)-1)^2+1/8*ln(
c*x^(1/2)-1)*ln(1/2*c*x^(1/2)+1/2)+1/8*(ln(1+c*x^(1/2))-ln(1/2*c*x^(1/2)+1
/2))*ln(-1/2*c*x^(1/2)+1/2)-1/16*ln(1+c*x^(1/2))^2-1/12/c^2/x+2/3*ln(c*x^(
1/2))-1/3*ln(1+c*x^(1/2))-1/3*ln(c*x^(1/2)-1))-1/4*b^3/(c*x^(1/2)-(-c^2*x+
1)^(1/2)+1)*(-c^2*x+1)^(1/2)+1/4*b^3/(c*x^(1/2)+(-c^2*x+1)^(1/2)+1)*(-c^2*
x+1)^(1/2)+3*a^2*b*(-1/4/c^4/x^2*arctanh(c*x^(1/2))-1/12/c^3/x^(3/2)-1/4/c
/x^(1/2)+1/8*ln(1+c*x^(1/2))-1/8*ln(c*x^(1/2)-1))-1/4*a^3/c^4/x^2-3/4*b...

```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x^3} dx = \int \frac{(b \operatorname{arctanh}(c\sqrt{x}) + a)^3}{x^3} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))^3/x^3,x, algorithm="fricas")
```

output

```
integral((b^3*arctanh(c*sqrt(x))^3 + 3*a*b^2*arctanh(c*sqrt(x))^2 + 3*a^2*
b*arctanh(c*sqrt(x)) + a^3)/x^3, x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x^3} dx = \int \frac{(a + b \operatorname{atanh}(c\sqrt{x}))^3}{x^3} dx$$

input `integrate((a+b*atanh(c*x**(1/2)))**3/x**3,x)`

output `Integral((a + b*atanh(c*sqrt(x)))**3/x**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 703 vs. 2(193) = 386.

Time = 1.17 (sec) , antiderivative size = 703, normalized size of antiderivative = 3.00

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x^3} dx = \text{Too large to display}$$

input `integrate((a+b*arctanh(c*x^(1/2)))^3/x^3,x, algorithm="maxima")`

output

```

-2*(log(c*sqrt(x) + 1)*log(-1/2*c*sqrt(x) + 1/2) + dilog(1/2*c*sqrt(x) + 1/2))*b^3*c^4 - 2*(log(c*sqrt(x))*log(-c*sqrt(x) + 1) + dilog(-c*sqrt(x) + 1))*b^3*c^4 + 2*(log(c*sqrt(x) + 1)*log(-c*sqrt(x)) + dilog(c*sqrt(x) + 1))*b^3*c^4 - 1/8*((6*c^3*log(c*sqrt(x) - 1) - 3*c^3*log(x) + (6*c^2*x + 3*c*sqrt(x) + 2)/x^(3/2))*c - 6*log(-c*sqrt(x) + 1)/x^2)*a^2*b + 1/4*(3*a^2*b*c^4 - 8*a*b^2*c^4 + b^3*c^4)*log(c*sqrt(x) + 1) - 1/4*(8*a*b^2*c^4 + b^3*c^4)*log(c*sqrt(x) - 1) - 1/8*(3*a^2*b*c^4 - 16*a*b^2*c^4)*log(x) - 1/2*a^3/x^2 - 1/16*(4*a^2*b*c*sqrt(x) - (b^3*c^4*x^2 - b^3)*log(c*sqrt(x) + 1)^3 + (b^3*c^4*x^2 - b^3)*log(-c*sqrt(x) + 1)^3 + 2*(3*b^3*c^3*x^(3/2) + b^3*c*sqrt(x) + 3*a*b^2 - (3*a*b^2*c^4 - 4*b^3*c^4)*x^2)*log(c*sqrt(x) + 1)^2 + (6*b^3*c^3*x^(3/2) + 2*b^3*c*sqrt(x) + 6*a*b^2 - 2*(3*a*b^2*c^4 + 4*b^3*c^4)*x^2 - 3*(b^3*c^4*x^2 - b^3)*log(c*sqrt(x) + 1))*log(-c*sqrt(x) + 1)^2 + 4*(3*a^2*b*c^3 + 2*b^3*c^3)*x^(3/2) - 2*(3*a^2*b*c^2 - 4*a*b^2*c^2)*x + 4*(6*a*b^2*c^3*x^(3/2) + b^3*c^2*x + 2*a*b^2*c*sqrt(x) + 3*a^2*b)*log(c*sqrt(x) + 1) - (24*a*b^2*c^3*x^(3/2) + 4*b^3*c^2*x + 8*a*b^2*c*sqrt(x) - 3*(b^3*c^4*x^2 - b^3)*log(c*sqrt(x) + 1)^2 + 4*(3*b^3*c^3*x^(3/2) + b^3*c*sqrt(x) + 3*a*b^2 - (3*a*b^2*c^4 - 4*b^3*c^4)*x^2)*log(c*sqrt(x) + 1))*log(-c*sqrt(x) + 1))/x^2

```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^3}{x^3} dx = \int \frac{(b \operatorname{arctanh}(c\sqrt{x}) + a)^3}{x^3} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))^3/x^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*sqrt(x)) + a)^3/x^3, x)
```


3.208 $\int x^{3/2} \operatorname{arctanh}(\sqrt{x}) dx$

Optimal result	1680
Mathematica [A] (verified)	1680
Rubi [A] (verified)	1681
Maple [A] (verified)	1682
Fricas [A] (verification not implemented)	1682
Sympy [B] (verification not implemented)	1683
Maxima [A] (verification not implemented)	1683
Giac [B] (verification not implemented)	1684
Mupad [B] (verification not implemented)	1684
Reduce [B] (verification not implemented)	1685

Optimal result

Integrand size = 12, antiderivative size = 38

$$\int x^{3/2} \operatorname{arctanh}(\sqrt{x}) dx = \frac{x}{5} + \frac{x^2}{10} + \frac{2}{5} x^{5/2} \operatorname{arctanh}(\sqrt{x}) + \frac{1}{5} \log(1-x)$$

output

```
1/5*x+1/10*x^2+2/5*x^(5/2)*arctanh(x^(1/2))+1/5*ln(1-x)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int x^{3/2} \operatorname{arctanh}(\sqrt{x}) dx = \frac{1}{10} (x(2+x) + 4x^{5/2} \operatorname{arctanh}(\sqrt{x}) + 2 \log(1-x))$$

input

```
Integrate[x^(3/2)*ArcTanh[Sqrt[x]],x]
```

output

```
(x*(2 + x) + 4*x^(5/2)*ArcTanh[Sqrt[x]] + 2*Log[1 - x])/10
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.92, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6452, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{3/2} \operatorname{arctanh}(\sqrt{x}) dx$$

$$\downarrow 6452$$

$$\frac{2}{5} x^{5/2} \operatorname{arctanh}(\sqrt{x}) - \frac{1}{5} \int \frac{x^2}{1-x} dx$$

$$\downarrow 49$$

$$\frac{2}{5} x^{5/2} \operatorname{arctanh}(\sqrt{x}) - \frac{1}{5} \int \left(-x + \frac{1}{1-x} - 1 \right) dx$$

$$\downarrow 2009$$

$$\frac{2}{5} x^{5/2} \operatorname{arctanh}(\sqrt{x}) + \frac{1}{5} \left(\frac{x^2}{2} + x + \log(1-x) \right)$$

input `Int [x^(3/2)*ArcTanh[Sqrt [x]] ,x]`

output `(2*x^(5/2)*ArcTanh[Sqrt [x]])/5 + (x + x^2/2 + Log[1 - x])/5`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int [ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.92

method	result	size
derivativedivides	$\frac{2x^{\frac{5}{2}} \operatorname{arctanh}(\sqrt{x})}{5} + \frac{x^2}{10} + \frac{x}{5} + \frac{\ln(-1+\sqrt{x})}{5} + \frac{\ln(1+\sqrt{x})}{5}$	35
default	$\frac{2x^{\frac{5}{2}} \operatorname{arctanh}(\sqrt{x})}{5} + \frac{x^2}{10} + \frac{x}{5} + \frac{\ln(-1+\sqrt{x})}{5} + \frac{\ln(1+\sqrt{x})}{5}$	35
meijerg	$\frac{x(3x+6)}{30} - \frac{x^{\frac{5}{2}}(\ln(1-\sqrt{x})-\ln(1+\sqrt{x}))}{5} + \frac{\ln(1-x)}{5}$	40

input `int(x^(3/2)*arctanh(x^(1/2)),x,method=_RETURNVERBOSE)`

output `2/5*x^(5/2)*arctanh(x^(1/2))+1/10*x^2+1/5*x+1/5*ln(-1+x^(1/2))+1/5*ln(1+x^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.95

$$\int x^{3/2} \operatorname{arctanh}(\sqrt{x}) dx = \frac{1}{5} x^{\frac{5}{2}} \log\left(-\frac{x + 2\sqrt{x} + 1}{x - 1}\right) + \frac{1}{10} x^2 + \frac{1}{5} x + \frac{1}{5} \log(x - 1)$$

input `integrate(x^(3/2)*arctanh(x^(1/2)),x, algorithm="fricas")`

output `1/5*x^(5/2)*log(-(x + 2*sqrt(x) + 1)/(x - 1)) + 1/10*x^2 + 1/5*x + 1/5*log(x - 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. $2(29) = 58$.

Time = 0.79 (sec) , antiderivative size = 121, normalized size of antiderivative = 3.18

$$\int x^{3/2} \operatorname{arctanh}(\sqrt{x}) dx = \frac{4x^{7/2} \operatorname{atanh}(\sqrt{x})}{10x - 10} - \frac{4x^{5/2} \operatorname{atanh}(\sqrt{x})}{10x - 10} + \frac{x^3}{10x - 10} + \frac{x^2}{10x - 10} + \frac{4x \log(\sqrt{x} + 1)}{10x - 10} - \frac{4x \operatorname{atanh}(\sqrt{x})}{10x - 10} - \frac{4 \log(\sqrt{x} + 1)}{10x - 10} + \frac{4 \operatorname{atanh}(\sqrt{x})}{10x - 10} - \frac{2}{10x - 10}$$

input `integrate(x**(3/2)*atanh(x**(1/2)),x)`

output `4*x**(7/2)*atanh(sqrt(x))/(10*x - 10) - 4*x**(5/2)*atanh(sqrt(x))/(10*x - 10) + x**3/(10*x - 10) + x**2/(10*x - 10) + 4*x*log(sqrt(x) + 1)/(10*x - 10) - 4*x*atanh(sqrt(x))/(10*x - 10) - 4*log(sqrt(x) + 1)/(10*x - 10) + 4*atanh(sqrt(x))/(10*x - 10) - 2/(10*x - 10)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.63

$$\int x^{3/2} \operatorname{arctanh}(\sqrt{x}) dx = \frac{2}{5} x^{5/2} \operatorname{artanh}(\sqrt{x}) + \frac{1}{10} x^2 + \frac{1}{5} x + \frac{1}{5} \log(x - 1)$$

input `integrate(x^(3/2)*arctanh(x^(1/2)),x, algorithm="maxima")`

output `2/5*x^(5/2)*arctanh(sqrt(x)) + 1/10*x^2 + 1/5*x + 1/5*log(x - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 170 vs. $2(26) = 52$.

Time = 0.12 (sec) , antiderivative size = 170, normalized size of antiderivative = 4.47

$$\int x^{3/2} \operatorname{arctanh}(\sqrt{x}) dx = \frac{8 \left(\frac{(\sqrt{x+1})^3}{(\sqrt{x-1})^3} - \frac{(\sqrt{x+1})^2}{(\sqrt{x-1})^2} + \frac{\sqrt{x+1}}{\sqrt{x-1}} \right)}{5 \left(\frac{\sqrt{x+1}}{\sqrt{x-1}} - 1 \right)^4} + \frac{2 \left(\frac{5(\sqrt{x+1})^4}{(\sqrt{x-1})^4} + \frac{10(\sqrt{x+1})^2}{(\sqrt{x-1})^2} + 1 \right) \log \left(-\frac{\sqrt{x+1}}{\sqrt{x-1}} \right)}{5 \left(\frac{\sqrt{x+1}}{\sqrt{x-1}} - 1 \right)^5} + \frac{2}{5} \log \left(\frac{\sqrt{x+1}}{|\sqrt{x-1}|} \right) - \frac{2}{5} \log \left(\left| -\frac{\sqrt{x+1}}{\sqrt{x-1}} + 1 \right| \right)$$

input `integrate(x^(3/2)*arctanh(x^(1/2)),x, algorithm="giac")`

output `8/5*((sqrt(x) + 1)^3/(sqrt(x) - 1)^3 - (sqrt(x) + 1)^2/(sqrt(x) - 1)^2 + (sqrt(x) + 1)/(sqrt(x) - 1))/(sqrt(x) + 1)/(sqrt(x) - 1) - 1)^4 + 2/5*(5*(sqrt(x) + 1)^4/(sqrt(x) - 1)^4 + 10*(sqrt(x) + 1)^2/(sqrt(x) - 1)^2 + 1)*log(-(sqrt(x) + 1)/(sqrt(x) - 1))/(sqrt(x) + 1)/(sqrt(x) - 1) - 1)^5 + 2/5*log((sqrt(x) + 1)/abs(sqrt(x) - 1)) - 2/5*log(abs(-(sqrt(x) + 1)/(sqrt(x) - 1) + 1))`

Mupad [B] (verification not implemented)

Time = 3.24 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.63

$$\int x^{3/2} \operatorname{arctanh}(\sqrt{x}) dx = \frac{x}{5} + \frac{\ln(x-1)}{5} + \frac{2x^{5/2} \operatorname{atanh}(\sqrt{x})}{5} + \frac{x^2}{10}$$

input `int(x^(3/2)*atanh(x^(1/2)),x)`

output `x/5 + log(x - 1)/5 + (2*x^(5/2)*atanh(x^(1/2)))/5 + x^2/10`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int x^{3/2} \operatorname{arctanh}(\sqrt{x}) dx = \frac{2\sqrt{x} \operatorname{atanh}(\sqrt{x}) x^2}{5} + \frac{2 \operatorname{atanh}(\sqrt{x})}{5} + \frac{2 \log(\sqrt{x} - 1)}{5} + \frac{x^2}{10} + \frac{x}{5}$$

input `int(x^(3/2)*atanh(x^(1/2)),x)`

output `(4*sqrt(x)*atanh(sqrt(x))*x**2 + 4*atanh(sqrt(x)) + 4*log(sqrt(x) - 1) + x**2 + 2*x)/10`

3.209 $\int \sqrt{x} \operatorname{arctanh}(\sqrt{x}) dx$

Optimal result	1686
Mathematica [A] (verified)	1686
Rubi [A] (verified)	1687
Maple [A] (verified)	1688
Fricas [A] (verification not implemented)	1688
Sympy [F]	1689
Maxima [A] (verification not implemented)	1689
Giac [B] (verification not implemented)	1689
Mupad [F(-1)]	1690
Reduce [B] (verification not implemented)	1690

Optimal result

Integrand size = 12, antiderivative size = 31

$$\int \sqrt{x} \operatorname{arctanh}(\sqrt{x}) dx = \frac{x}{3} + \frac{2}{3} x^{3/2} \operatorname{arctanh}(\sqrt{x}) + \frac{1}{3} \log(1-x)$$

output $1/3*x+2/3*x^{(3/2)}*\operatorname{arctanh}(x^{(1/2)})+1/3*\ln(1-x)$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.81

$$\int \sqrt{x} \operatorname{arctanh}(\sqrt{x}) dx = \frac{1}{3} (x + 2x^{3/2} \operatorname{arctanh}(\sqrt{x}) + \log(1-x))$$

input $\operatorname{Integrate}[\operatorname{Sqrt}[x]*\operatorname{ArcTanh}[\operatorname{Sqrt}[x]], x]$

output $(x + 2*x^{(3/2)}*\operatorname{ArcTanh}[\operatorname{Sqrt}[x]] + \operatorname{Log}[1 - x])/3$

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6452, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x} \operatorname{arctanh}(\sqrt{x}) dx$$

$$\downarrow 6452$$

$$\frac{2}{3} x^{3/2} \operatorname{arctanh}(\sqrt{x}) - \frac{1}{3} \int \frac{x}{1-x} dx$$

$$\downarrow 49$$

$$\frac{2}{3} x^{3/2} \operatorname{arctanh}(\sqrt{x}) - \frac{1}{3} \int \left(\frac{1}{1-x} - 1 \right) dx$$

$$\downarrow 2009$$

$$\frac{2}{3} x^{3/2} \operatorname{arctanh}(\sqrt{x}) + \frac{1}{3} (x + \log(1-x))$$

input `Int[Sqrt[x]*ArcTanh[Sqrt[x]],x]`

output `(2*x^(3/2)*ArcTanh[Sqrt[x]])/3 + (x + Log[1 - x])/3`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.97

method	result	size
derivativdivides	$\frac{2x^{\frac{3}{2}} \operatorname{arctanh}(\sqrt{x})}{3} + \frac{x}{3} + \frac{\ln(-1+\sqrt{x})}{3} + \frac{\ln(1+\sqrt{x})}{3}$	30
default	$\frac{2x^{\frac{3}{2}} \operatorname{arctanh}(\sqrt{x})}{3} + \frac{x}{3} + \frac{\ln(-1+\sqrt{x})}{3} + \frac{\ln(1+\sqrt{x})}{3}$	30
meijerg	$\frac{x}{3} - \frac{x^{\frac{3}{2}} (\ln(1-\sqrt{x}) - \ln(1+\sqrt{x}))}{3} + \frac{\ln(1-x)}{3}$	35

input

```
int(x^(1/2)*arctanh(x^(1/2)),x,method=_RETURNVERBOSE)
```

output

```
2/3*x^(3/2)*arctanh(x^(1/2))+1/3*x+1/3*ln(-1+x^(1/2))+1/3*ln(1+x^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \sqrt{x} \operatorname{arctanh}(\sqrt{x}) dx = \frac{1}{3} x^{\frac{3}{2}} \log\left(-\frac{x + 2\sqrt{x} + 1}{x - 1}\right) + \frac{1}{3} x + \frac{1}{3} \log(x - 1)$$

input

```
integrate(x^(1/2)*arctanh(x^(1/2)),x, algorithm="fricas")
```

output

```
1/3*x^(3/2)*log(-(x + 2*sqrt(x) + 1)/(x - 1)) + 1/3*x + 1/3*log(x - 1)
```

Sympy [F]

$$\int \sqrt{x} \operatorname{arctanh}(\sqrt{x}) dx = \int \sqrt{x} \operatorname{atanh}(\sqrt{x}) dx$$

input `integrate(x**(1/2)*atanh(x**(1/2)),x)`

output `Integral(sqrt(x)*atanh(sqrt(x)), x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.61

$$\int \sqrt{x} \operatorname{arctanh}(\sqrt{x}) dx = \frac{2}{3} x^{\frac{3}{2}} \operatorname{artanh}(\sqrt{x}) + \frac{1}{3} x + \frac{1}{3} \log(x-1)$$

input `integrate(x^(1/2)*arctanh(x^(1/2)),x, algorithm="maxima")`

output `2/3*x^(3/2)*arctanh(sqrt(x)) + 1/3*x + 1/3*log(x - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. $2(21) = 42$.

Time = 0.12 (sec) , antiderivative size = 121, normalized size of antiderivative = 3.90

$$\int \sqrt{x} \operatorname{arctanh}(\sqrt{x}) dx = \frac{2 \left(\frac{3(\sqrt{x}+1)^2}{(\sqrt{x}-1)^2} + 1 \right) \log \left(-\frac{\sqrt{x}+1}{\sqrt{x}-1} \right)}{3 \left(\frac{\sqrt{x}+1}{\sqrt{x}-1} - 1 \right)^3} + \frac{4(\sqrt{x}+1)}{3(\sqrt{x}-1) \left(\frac{\sqrt{x}+1}{\sqrt{x}-1} - 1 \right)^2} \\ + \frac{2}{3} \log \left(\frac{\sqrt{x}+1}{|\sqrt{x}-1|} \right) - \frac{2}{3} \log \left(\left| -\frac{\sqrt{x}+1}{\sqrt{x}-1} + 1 \right| \right)$$

input `integrate(x^(1/2)*arctanh(x^(1/2)),x, algorithm="giac")`

output

```
2/3*(3*(sqrt(x) + 1)^2/(sqrt(x) - 1)^2 + 1)*log(-(sqrt(x) + 1)/(sqrt(x) - 1))/((sqrt(x) + 1)/(sqrt(x) - 1) - 1)^3 + 4/3*(sqrt(x) + 1)/((sqrt(x) - 1) * ((sqrt(x) + 1)/(sqrt(x) - 1) - 1)^2) + 2/3*log((sqrt(x) + 1)/abs(sqrt(x) - 1)) - 2/3*log(abs(-(sqrt(x) + 1)/(sqrt(x) - 1) + 1))
```

Mupad [F(-1)]

Timed out.

$$\int \sqrt{x} \operatorname{arctanh}(\sqrt{x}) \, dx = \int \sqrt{x} \operatorname{atanh}(\sqrt{x}) \, dx$$

input

```
int(x^(1/2)*atanh(x^(1/2)),x)
```

output

```
int(x^(1/2)*atanh(x^(1/2)), x)
```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.77

$$\int \sqrt{x} \operatorname{arctanh}(\sqrt{x}) \, dx = \frac{2\sqrt{x} \operatorname{atanh}(\sqrt{x}) x}{3} + \frac{2 \operatorname{atanh}(\sqrt{x})}{3} + \frac{2 \log(\sqrt{x} - 1)}{3} + \frac{x}{3}$$

input

```
int(x^(1/2)*atanh(x^(1/2)),x)
```

output

```
(2*sqrt(x)*atanh(sqrt(x))*x + 2*atanh(sqrt(x)) + 2*log(sqrt(x) - 1) + x)/3
```

3.210 $\int \frac{\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} dx$

Optimal result	1691
Mathematica [A] (verified)	1691
Rubi [A] (verified)	1692
Maple [A] (verified)	1693
Fricas [A] (verification not implemented)	1693
Sympy [B] (verification not implemented)	1693
Maxima [A] (verification not implemented)	1694
Giac [B] (verification not implemented)	1694
Mupad [B] (verification not implemented)	1695
Reduce [B] (verification not implemented)	1695

Optimal result

Integrand size = 12, antiderivative size = 20

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} dx = 2\sqrt{x}\operatorname{arctanh}(\sqrt{x}) + \log(1-x)$$

output `2*x^(1/2)*arctanh(x^(1/2))+ln(1-x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} dx = 2\sqrt{x}\operatorname{arctanh}(\sqrt{x}) + \log(1-x)$$

input `Integrate[ArcTanh[Sqrt[x]]/Sqrt[x],x]`

output `2*Sqrt[x]*ArcTanh[Sqrt[x]] + Log[1 - x]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6452, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} dx$$

$$\downarrow \text{6452}$$

$$2\sqrt{x}\operatorname{arctanh}(\sqrt{x}) - \int \frac{1}{1-x} dx$$

$$\downarrow \text{16}$$

$$2\sqrt{x}\operatorname{arctanh}(\sqrt{x}) + \log(1-x)$$

input `Int[ArcTanh[Sqrt[x]]/Sqrt[x],x]`

output `2*Sqrt[x]*ArcTanh[Sqrt[x]] + Log[1 - x]`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

method	result	size
derivativdivides	$2\sqrt{x} \operatorname{arctanh}(\sqrt{x}) + \ln(1-x)$	17
default	$2\sqrt{x} \operatorname{arctanh}(\sqrt{x}) + \ln(1-x)$	17
meijerg	$-\sqrt{x} (\ln(1-\sqrt{x}) - \ln(1+\sqrt{x})) + \ln(1-x)$	30

input `int(arctanh(x^(1/2))/x^(1/2),x,method=_RETURNVERBOSE)`

output `2*x^(1/2)*arctanh(x^(1/2))+ln(1-x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.25

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} dx = \sqrt{x} \log\left(-\frac{x+2\sqrt{x}+1}{x-1}\right) + \log(x-1)$$

input `integrate(arctanh(x^(1/2))/x^(1/2),x, algorithm="fricas")`

output `sqrt(x)*log(-(x+2*sqrt(x)+1)/(x-1))+log(x-1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 87 vs. $2(17) = 34$.

Time = 0.21 (sec) , antiderivative size = 87, normalized size of antiderivative = 4.35

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} dx = \frac{2x^{\frac{3}{2}} \operatorname{atanh}(\sqrt{x})}{x-1} - \frac{2\sqrt{x} \operatorname{atanh}(\sqrt{x})}{x-1} + \frac{2x \log(\sqrt{x}+1)}{x-1} - \frac{2x \operatorname{atanh}(\sqrt{x})}{x-1} - \frac{2 \log(\sqrt{x}+1)}{x-1} + \frac{2 \operatorname{atanh}(\sqrt{x})}{x-1}$$

input `integrate(atanh(x**(1/2))/x**(1/2),x)`

output `2*x**(3/2)*atanh(sqrt(x))/(x - 1) - 2*sqrt(x)*atanh(sqrt(x))/(x - 1) + 2*x*log(sqrt(x) + 1)/(x - 1) - 2*x*atanh(sqrt(x))/(x - 1) - 2*log(sqrt(x) + 1)/(x - 1) + 2*atanh(sqrt(x))/(x - 1)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} dx = 2\sqrt{x} \operatorname{arctanh}(\sqrt{x}) + \log(-x + 1)$$

input `integrate(arctanh(x^(1/2))/x^(1/2),x, algorithm="maxima")`

output `2*sqrt(x)*arctanh(sqrt(x)) + log(-x + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 72 vs. 2(16) = 32.

Time = 0.11 (sec) , antiderivative size = 72, normalized size of antiderivative = 3.60

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} dx = \frac{2 \log\left(-\frac{\sqrt{x}+1}{\sqrt{x}-1}\right)}{\frac{\sqrt{x}+1}{\sqrt{x}-1} - 1} + 2 \log\left(\frac{\sqrt{x}+1}{|\sqrt{x}-1|}\right) - 2 \log\left(\left|-\frac{\sqrt{x}+1}{\sqrt{x}-1} + 1\right|\right)$$

input `integrate(arctanh(x^(1/2))/x^(1/2),x, algorithm="giac")`

output `2*log(-(sqrt(x) + 1)/(sqrt(x) - 1))/((sqrt(x) + 1)/(sqrt(x) - 1) - 1) + 2*log((sqrt(x) + 1)/abs(sqrt(x) - 1)) - 2*log(abs(-(sqrt(x) + 1)/(sqrt(x) - 1) + 1))`

Mupad [B] (verification not implemented)

Time = 3.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} dx = \ln(x-1) + 2\sqrt{x} \operatorname{atanh}(\sqrt{x})$$

input `int(atanh(x^(1/2))/x^(1/2),x)`

output `log(x - 1) + 2*x^(1/2)*atanh(x^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} dx = 2\sqrt{x} \operatorname{atanh}(\sqrt{x}) + 2\operatorname{atanh}(\sqrt{x}) + 2\log(\sqrt{x}-1)$$

input `int(atanh(x^(1/2))/x^(1/2),x)`

output `2*(sqrt(x)*atanh(sqrt(x)) + atanh(sqrt(x)) + log(sqrt(x) - 1))`

3.211 $\int \frac{\operatorname{arctanh}(\sqrt{x})}{x^{3/2}} dx$

Optimal result	1696
Mathematica [A] (verified)	1696
Rubi [A] (verified)	1697
Maple [A] (verified)	1698
Fricas [A] (verification not implemented)	1699
Sympy [B] (verification not implemented)	1699
Maxima [A] (verification not implemented)	1700
Giac [B] (verification not implemented)	1700
Mupad [B] (verification not implemented)	1700
Reduce [B] (verification not implemented)	1701

Optimal result

Integrand size = 12, antiderivative size = 24

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{x^{3/2}} dx = -\frac{2\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} - \log(1-x) + \log(x)$$

output `-2*arctanh(x^(1/2))/x^(1/2)-ln(1-x)+ln(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{x^{3/2}} dx = -\frac{2\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} - \log(1-x) + \log(x)$$

input `Integrate[ArcTanh[Sqrt[x]]/x^(3/2),x]`

output `(-2*ArcTanh[Sqrt[x]])/Sqrt[x] - Log[1 - x] + Log[x]`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6452, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(\sqrt{x})}{x^{3/2}} dx \\ & \quad \downarrow \text{6452} \\ & \int \frac{1}{(1-x)x} dx - \frac{2\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} \\ & \quad \downarrow \text{47} \\ & \int \frac{1}{1-x} dx + \int \frac{1}{x} dx - \frac{2\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} \\ & \quad \downarrow \text{14} \\ & \int \frac{1}{1-x} dx - \frac{2\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} + \log(x) \\ & \quad \downarrow \text{16} \\ & -\frac{2\operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} - \log(1-x) + \log(x) \end{aligned}$$

input `Int[ArcTanh[Sqrt[x]]/x^(3/2),x]`

output `(-2*ArcTanh[Sqrt[x]])/Sqrt[x] - Log[1 - x] + Log[x]`

Definitions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} + \ln(x) - \ln(1 + \sqrt{x}) - \ln(-1 + \sqrt{x})$	29
default	$-\frac{2 \operatorname{arctanh}(\sqrt{x})}{\sqrt{x}} + \ln(x) - \ln(1 + \sqrt{x}) - \ln(-1 + \sqrt{x})$	29
meijerg	$\ln(x) + i\pi + \frac{\ln(1 - \sqrt{x}) - \ln(1 + \sqrt{x})}{\sqrt{x}} - \ln(1 - x)$	37

input `int(arctanh(x^(1/2))/x^(3/2), x, method=_RETURNVERBOSE)`

output `-2*arctanh(x^(1/2))/x^(1/2)+ln(x)-ln(1+x^(1/2))-ln(-1+x^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.54

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{x^{3/2}} dx = -\frac{x \log(x-1) - x \log(x) + \sqrt{x} \log\left(-\frac{x+2\sqrt{x}+1}{x-1}\right)}{x}$$

input `integrate(arctanh(x^(1/2))/x^(3/2),x, algorithm="fricas")`

output `-(x*log(x - 1) - x*log(x) + sqrt(x)*log(-(x + 2*sqrt(x) + 1)/(x - 1)))/x`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 126 vs. 2(20) = 40.

Time = 0.37 (sec) , antiderivative size = 126, normalized size of antiderivative = 5.25

$$\begin{aligned} \int \frac{\operatorname{arctanh}(\sqrt{x})}{x^{3/2}} dx &= -\frac{2x^{3/2} \operatorname{atanh}(\sqrt{x})}{x^2 - x} + \frac{2\sqrt{x} \operatorname{atanh}(\sqrt{x})}{x^2 - x} \\ &+ \frac{x^2 \log(x)}{x^2 - x} - \frac{2x^2 \log(\sqrt{x} + 1)}{x^2 - x} + \frac{2x^2 \operatorname{atanh}(\sqrt{x})}{x^2 - x} \\ &- \frac{x \log(x)}{x^2 - x} + \frac{2x \log(\sqrt{x} + 1)}{x^2 - x} - \frac{2x \operatorname{atanh}(\sqrt{x})}{x^2 - x} \end{aligned}$$

input `integrate(atanh(x**(1/2))/x**(3/2),x)`

output `-2*x**(3/2)*atanh(sqrt(x))/(x**2 - x) + 2*sqrt(x)*atanh(sqrt(x))/(x**2 - x) + x**2*log(x)/(x**2 - x) - 2*x**2*log(sqrt(x) + 1)/(x**2 - x) + 2*x**2*atanh(sqrt(x))/(x**2 - x) - x*log(x)/(x**2 - x) + 2*x*log(sqrt(x) + 1)/(x**2 - x) - 2*x*atanh(sqrt(x))/(x**2 - x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.75

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{x^{3/2}} dx = -\frac{2 \operatorname{artanh}(\sqrt{x})}{\sqrt{x}} - \log(x-1) + \log(x)$$

input `integrate(arctanh(x^(1/2))/x^(3/2),x, algorithm="maxima")`

output `-2*arctanh(sqrt(x))/sqrt(x) - log(x - 1) + log(x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 72 vs. 2(20) = 40.

Time = 0.12 (sec) , antiderivative size = 72, normalized size of antiderivative = 3.00

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{x^{3/2}} dx = \frac{2 \log\left(-\frac{\sqrt{x}+1}{\sqrt{x}-1}\right)}{\frac{\sqrt{x}+1}{\sqrt{x}-1} + 1} - 2 \log\left(\left|\frac{\sqrt{x}+1}{\sqrt{x}-1}\right|\right) + 2 \log\left(\left|-\frac{\sqrt{x}+1}{\sqrt{x}-1} - 1\right|\right)$$

input `integrate(arctanh(x^(1/2))/x^(3/2),x, algorithm="giac")`

output `2*log(-(sqrt(x) + 1)/(sqrt(x) - 1))/((sqrt(x) + 1)/(sqrt(x) - 1) + 1) - 2*log((sqrt(x) + 1)/abs(sqrt(x) - 1)) + 2*log(abs(-(sqrt(x) + 1)/(sqrt(x) - 1) - 1))`

Mupad [B] (verification not implemented)

Time = 3.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{x^{3/2}} dx = 2 \ln(\sqrt{x}) - \ln(x-1) - \frac{2 \operatorname{atanh}(\sqrt{x})}{\sqrt{x}}$$

input `int(atanh(x^(1/2))/x^(3/2),x)`

output $2*\log(x^{(1/2)}) - \log(x - 1) - (2*atanh(x^{(1/2)}))/x^{(1/2)}$

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.42

$$\int \frac{\operatorname{arctanh}(\sqrt{x})}{x^{3/2}} dx = \frac{-2\sqrt{x} \operatorname{atanh}(\sqrt{x}) - 2\operatorname{atanh}(\sqrt{x}) - 2\sqrt{x} \log(\sqrt{x} - 1) + 2\sqrt{x} \log(\sqrt{x})}{\sqrt{x}}$$

input $\operatorname{int}(\operatorname{atanh}(x^{(1/2)})/x^{(3/2)}, x)$

output $(2*(-\sqrt{x}*\operatorname{atanh}(\sqrt{x}) - \operatorname{atanh}(\sqrt{x}) - \sqrt{x}*\log(\sqrt{x} - 1) + \sqrt{x}*\log(\sqrt{x}))) / \sqrt{x}$

3.212 $\int x^3(a + b \operatorname{arctanh}(cx^{3/2})) dx$

Optimal result	1702
Mathematica [A] (verified)	1703
Rubi [A] (verified)	1703
Maple [A] (verified)	1709
Fricas [C] (verification not implemented)	1710
Sympy [F(-1)]	1711
Maxima [A] (verification not implemented)	1711
Giac [C] (verification not implemented)	1712
Mupad [B] (verification not implemented)	1712
Reduce [B] (verification not implemented)	1713

Optimal result

Integrand size = 16, antiderivative size = 160

$$\int x^3(a + b \operatorname{arctanh}(cx^{3/2})) dx = \frac{3bx^{5/2}}{20c} - \frac{\sqrt{3}b \arctan\left(\frac{1-2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{8c^{8/3}} + \frac{\sqrt{3}b \arctan\left(\frac{1+2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{8c^{8/3}} - \frac{b \operatorname{arctanh}(\sqrt[3]{c}\sqrt{x})}{4c^{8/3}} + \frac{1}{4}x^4(a + b \operatorname{arctanh}(cx^{3/2})) - \frac{b \operatorname{arctanh}\left(\frac{\sqrt[3]{c}\sqrt{x}}{1+c^{2/3}x}\right)}{8c^{8/3}}$$

output

```
3/20*b*x^(5/2)/c-1/8*3^(1/2)*b*arctan(1/3*(1-2*c^(1/3)*x^(1/2))*3^(1/2))/c
^(8/3)+1/8*3^(1/2)*b*arctan(1/3*(1+2*c^(1/3)*x^(1/2))*3^(1/2))/c^(8/3)-1/4
*b*arctanh(c^(1/3)*x^(1/2))/c^(8/3)+1/4*x^4*(a+b*arctanh(c*x^(3/2)))-1/8*b
*arctanh(c^(1/3)*x^(1/2)/(1+c^(2/3)*x))/c^(8/3)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.39

$$\int x^3(a + b \operatorname{arctanh}(cx^{3/2})) dx = \frac{3bx^{5/2}}{20c} + \frac{ax^4}{4} + \frac{\sqrt{3}b \arctan\left(\frac{-1+2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{8c^{8/3}} + \frac{\sqrt{3}b \arctan\left(\frac{1+2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{8c^{8/3}} + \frac{1}{4}bx^4 \operatorname{arctanh}(cx^{3/2}) + \frac{b \log(1 - \sqrt[3]{c}\sqrt{x})}{8c^{8/3}} - \frac{b \log(1 + \sqrt[3]{c}\sqrt{x})}{8c^{8/3}} + \frac{b \log(1 - \sqrt[3]{c}\sqrt{x} + c^{2/3}x)}{16c^{8/3}} - \frac{b \log(1 + \sqrt[3]{c}\sqrt{x} + c^{2/3}x)}{16c^{8/3}}$$

input `Integrate[x^3*(a + b*ArcTanh[c*x^(3/2)]),x]`

output $(3*b*x^{(5/2)})/(20*c) + (a*x^4)/4 + (\operatorname{Sqrt}[3]*b*\operatorname{ArcTan}[(-1 + 2*c^{(1/3)}*\operatorname{Sqrt}[x])/ \operatorname{Sqrt}[3]])/(8*c^{(8/3)}) + (\operatorname{Sqrt}[3]*b*\operatorname{ArcTan}[(1 + 2*c^{(1/3)}*\operatorname{Sqrt}[x])/ \operatorname{Sqrt}[3]])/(8*c^{(8/3)}) + (b*x^4*\operatorname{ArcTanh}[c*x^{(3/2)}])/4 + (b*\operatorname{Log}[1 - c^{(1/3)}*\operatorname{Sqrt}[x]])/(8*c^{(8/3)}) - (b*\operatorname{Log}[1 + c^{(1/3)}*\operatorname{Sqrt}[x]])/(8*c^{(8/3)}) + (b*\operatorname{Log}[1 - c^{(1/3)}*\operatorname{Sqrt}[x] + c^{(2/3)}*x])/(16*c^{(8/3)}) - (b*\operatorname{Log}[1 + c^{(1/3)}*\operatorname{Sqrt}[x] + c^{(2/3)}*x])/(16*c^{(8/3)})$

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.32, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6452, 843, 851, 825, 27, 219, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3(a + b \operatorname{arctanh}(cx^{3/2})) dx$$

$$\downarrow 6452$$

$$\frac{1}{4}x^4(a + b \operatorname{arctanh}(cx^{3/2})) - \frac{3}{8}bc \int \frac{x^{9/2}}{1 - c^2x^3} dx$$

$$\downarrow 843$$

$$\begin{aligned}
& \frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(cx^{3/2}\right)\right) - \frac{3}{8}bc\left(\frac{\int \frac{x^{3/2}}{1-c^2x^3}dx}{c^2} - \frac{2x^{5/2}}{5c^2}\right) \\
& \quad \downarrow \text{851} \\
& \frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(cx^{3/2}\right)\right) - \frac{3}{8}bc\left(\frac{2\int \frac{x^2}{1-c^2x^3}d\sqrt{x}}{c^2} - \frac{2x^{5/2}}{5c^2}\right) \\
& \quad \downarrow \text{825} \\
& \frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(cx^{3/2}\right)\right) - \\
& \left. \frac{3}{8}bc\left(\frac{2\left(\frac{\int \frac{1}{1-c^{2/3}x}d\sqrt{x}}{3c^{4/3}} + \frac{\int -\frac{\sqrt[3]{c\sqrt{x}+1}}{2(c^{2/3}x-\sqrt[3]{c\sqrt{x}+1})}d\sqrt{x}}{3c^{4/3}} + \frac{\int -\frac{1-\sqrt[3]{c\sqrt{x}}}{2(c^{2/3}x+\sqrt[3]{c\sqrt{x}+1})}d\sqrt{x}}{3c^{4/3}}\right)}{c^2} - \frac{2x^{5/2}}{5c^2}\right)\right) \\
& \quad \downarrow \text{27} \\
& \frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(cx^{3/2}\right)\right) - \\
& \left. \frac{3}{8}bc\left(\frac{2\left(\frac{\int \frac{1}{1-c^{2/3}x}d\sqrt{x}}{3c^{4/3}} - \frac{\int \frac{\sqrt[3]{c\sqrt{x}+1}}{c^{2/3}x-\sqrt[3]{c\sqrt{x}+1}}d\sqrt{x}}{6c^{4/3}} - \frac{\int \frac{1-\sqrt[3]{c\sqrt{x}}}{c^{2/3}x+\sqrt[3]{c\sqrt{x}+1}}d\sqrt{x}}{6c^{4/3}}\right)}{c^2} - \frac{2x^{5/2}}{5c^2}\right)\right) \\
& \quad \downarrow \text{219} \\
& \frac{1}{4}x^4\left(a + \operatorname{barctanh}\left(cx^{3/2}\right)\right) - \\
& \left. \frac{3}{8}bc\left(\frac{2\left(-\frac{\int \frac{\sqrt[3]{c\sqrt{x}+1}}{c^{2/3}x-\sqrt[3]{c\sqrt{x}+1}}d\sqrt{x}}{6c^{4/3}} - \frac{\int \frac{1-\sqrt[3]{c\sqrt{x}}}{c^{2/3}x+\sqrt[3]{c\sqrt{x}+1}}d\sqrt{x}}{6c^{4/3}} + \frac{\operatorname{arctanh}\left(\sqrt[3]{c\sqrt{x}}\right)}{3c^{5/3}}\right)}{c^2} - \frac{2x^{5/2}}{5c^2}\right)\right) \\
& \quad \downarrow \text{1142}
\end{aligned}$$

$$\frac{3}{8}bc \left(\frac{\frac{1}{4}x^4 \left(a + \operatorname{barctanh}(cx^{3/2}) \right) - \int \frac{\sqrt[3]{c}(1-2\sqrt[3]{c}\sqrt{x})}{c^{2/3}x - \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x}}{6c^{4/3}} - \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{c}\sqrt{x+1})}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x}}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{c}\sqrt{x+1})}{3c^{5/3}}}{c^2} \right)$$

25

$$\frac{3}{8}bc \left(\frac{\frac{1}{4}x^4 \left(a + \operatorname{barctanh}(cx^{3/2}) \right) - \int \frac{\sqrt[3]{c}(1-2\sqrt[3]{c}\sqrt{x})}{c^{2/3}x - \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x}}{6c^{4/3}} - \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{c}\sqrt{x+1})}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x}}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{c}\sqrt{x+1})}{3c^{5/3}}}{c^2} \right)$$

27

$$\frac{3}{8}bc \left(\frac{\frac{1}{4}x^4 \left(a + \operatorname{barctanh}(cx^{3/2}) \right) - \int \frac{\sqrt[3]{c}(1-2\sqrt[3]{c}\sqrt{x})}{c^{2/3}x - \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} - \frac{1}{2} \int \frac{2\sqrt[3]{c}\sqrt{x+1}}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x}}{6c^{4/3}} - \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{c}\sqrt{x+1})}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x}}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{c}\sqrt{x+1})}{3c^{5/3}}}{c^2} \right)$$

1082

$$\frac{3}{8}bc \left(\frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx^{3/2})) - 2 \left(\frac{\int \frac{1-2\sqrt[3]{c}\sqrt{x}}{c^{2/3}x-\sqrt[3]{c}\sqrt{x+1}} dx - \int \frac{2\sqrt[3]{c}\sqrt{x+1}}{c^{2/3}x+\sqrt[3]{c}\sqrt{x+1}} dx - \int \frac{1-2\sqrt[3]{c}\sqrt{x}}{c^{2/3}x-\sqrt[3]{c}\sqrt{x+1}} dx - \int \frac{2\sqrt[3]{c}\sqrt{x+1}}{c^{2/3}x+\sqrt[3]{c}\sqrt{x+1}} dx \right)}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{c}\sqrt{x})}{3c^{5/3}}}{c^2} \right)$$

217

$$\frac{3}{8}bc \left(\frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx^{3/2})) - 2 \left(\frac{\int \frac{1-2\sqrt[3]{c}\sqrt{x}}{c^{2/3}x-\sqrt[3]{c}\sqrt{x+1}} dx - \frac{\sqrt{3} \arctan\left(\frac{1-2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{c}\sqrt{x+1}}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \int \frac{2\sqrt[3]{c}\sqrt{x+1}}{c^{2/3}x+\sqrt[3]{c}\sqrt{x+1}} dx \right)}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{c}\sqrt{x})}{3c^{5/3}}}{c^2} \right)$$

1103

$$\frac{3}{8}bc \left(\frac{\frac{1}{4}x^4(a + \operatorname{barctanh}(cx^{3/2})) - 2 \left(\frac{\frac{\log(c^{2/3}x-\sqrt[3]{c}\sqrt{x+1})}{2\sqrt[3]{c}} - \frac{\sqrt{3} \arctan\left(\frac{1-2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{c}\sqrt{x+1}}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{\log(c^{2/3}x+\sqrt[3]{c}\sqrt{x+1})}{2\sqrt[3]{c}} \right)}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{c}\sqrt{x})}{3c^{5/3}}}{c^2} \right)$$

input `Int[x^3*(a + b*ArcTanh[c*x^(3/2)]), x]`

output

$$\frac{(x^4(a + b \operatorname{ArcTanh}[c x^{3/2}]))/4 - (3 b c ((-2 x^{5/2})/(5 c^2) + (2 (\operatorname{ArcTanh}[c^{1/3} \sqrt{x}]/(3 c^{5/3}) - ((\sqrt{3} \operatorname{ArcTan}[(1 - 2 c^{1/3}) \sqrt{x}]/\sqrt{3}])/c^{1/3})) + \operatorname{Log}[1 - c^{1/3} \sqrt{x} + c^{2/3} x]/(2 c^{1/3})))/(6 c^{4/3}) - ((\sqrt{3} \operatorname{ArcTan}[(1 + 2 c^{1/3}) \sqrt{x}]/\sqrt{3}])/c^{1/3}) - \operatorname{Log}[1 + c^{1/3} \sqrt{x} + c^{2/3} x]/(2 c^{1/3})))/(6 c^{4/3}))/c^2)/8$$

Defintions of rubi rules used

rule 25

$$\operatorname{Int}[-(F x), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F x, x], x]$$

rule 27

$$\operatorname{Int}[(a)(F x), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[a \operatorname{Int}[F x, x], x] /; \operatorname{FreeQ}[a, x] \&\& \operatorname{!MatchQ}[F x, (b)(G x) /; \operatorname{FreeQ}[b, x]]$$

rule 217

$$\operatorname{Int}[(a) + (b)(x)^2)^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(-(\operatorname{Rt}[-a, 2] \operatorname{Rt}[-b, 2])^{-1}) \operatorname{ArcTan}[\operatorname{Rt}[-b, 2] (x/\operatorname{Rt}[-a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&\& \operatorname{PosQ}[a/b] \&\& (\operatorname{LtQ}[a, 0] \operatorname{||} \operatorname{LtQ}[b, 0])$$

rule 219

$$\operatorname{Int}[(a) + (b)(x)^2)^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2] \operatorname{Rt}[-b, 2])) \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2] (x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&\& \operatorname{NegQ}[a/b] \&\& (\operatorname{GtQ}[a, 0] \operatorname{||} \operatorname{LtQ}[b, 0])$$

rule 825

$$\operatorname{Int}[(x)^{(m)} / ((a) + (b)(x)^{(n)}), x_{\text{Symbol}}] \rightarrow \operatorname{Module}[\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, n]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, n]], k, u\}, \operatorname{Simp}[u = \operatorname{Int}[(r \operatorname{Cos}[2 k m (\operatorname{Pi}/n)] - s \operatorname{Cos}[2 k (m + 1) (\operatorname{Pi}/n)] x) / (r^2 - 2 r s \operatorname{Cos}[2 k (\operatorname{Pi}/n)] x + s^2 x^2), x] + \operatorname{Int}[(r \operatorname{Cos}[2 k m (\operatorname{Pi}/n)] + s \operatorname{Cos}[2 k (m + 1) (\operatorname{Pi}/n)] x) / (r^2 + 2 r s \operatorname{Cos}[2 k (\operatorname{Pi}/n)] x + s^2 x^2), x]; 2 (r^{(m + 2)} / (a n s^m)) \operatorname{Int}[1 / (r^2 - s^2 x^2), x] + 2 (r^{(m + 1)} / (a n s^m)) \operatorname{Sum}[u, \{k, 1, (n - 2)/4\}], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&\& \operatorname{IGtQ}[(n - 2)/4, 0] \&\& \operatorname{IGtQ}[m, 0] \&\& \operatorname{LtQ}[m, n - 1] \&\& \operatorname{NegQ}[a/b]$$

rule 843 $\text{Int}[\text{((c_.)*(x_))}^{\text{(m_)}} * \text{((a_) + (b_.)*(x_)^{\text{(n_)}})^{\text{(p_)}}], \text{x_Symbol}] \text{:> Simp}[c^{\text{(n - 1)}} * (c*x)^{\text{(m - n + 1)}} * ((a + b*x^n)^{\text{(p + 1)}} / (b*(m + n*p + 1))), x] - \text{Simp}[a*c^{\text{n}} * ((m - n + 1) / (b*(m + n*p + 1))) \text{Int}[(c*x)^{\text{(m - n)}} * (a + b*x^n)^{\text{p}}, x], x] \text{/; FreeQ}\{a, b, c, p\}, x] \&\& \text{IGtQ}\{n, 0\} \&\& \text{GtQ}\{m, n - 1\} \&\& \text{NeQ}\{m + n*p + 1, 0\} \&\& \text{IntBinomialQ}\{a, b, c, n, m, p, x\}$

rule 851 $\text{Int}[\text{((c_.)*(x_))}^{\text{(m_)}} * \text{((a_) + (b_.)*(x_)^{\text{(n_)}})^{\text{(p_)}}], \text{x_Symbol}] \text{:> With}\{k = \text{Denominator}\{m\}\}, \text{Simp}[k/c \text{Subst}[\text{Int}[x^{\text{(k*(m + 1) - 1)}} * (a + b*(x^{\text{(k*n)}}/c^n))^{\text{p}}, x], x, (c*x)^{\text{(1/k)}}], x]] \text{/; FreeQ}\{a, b, c, p\}, x] \&\& \text{IGtQ}\{n, 0\} \&\& \text{FractionQ}\{m\} \&\& \text{IntBinomialQ}\{a, b, c, n, m, p, x\}$

rule 1082 $\text{Int}[\text{((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^{\text{-1}}}, \text{x_Symbol}] \text{:> With}\{q = 1 - 4*S\text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{Subst}[\text{Int}[1/(q - x^2)], x], x, 1 + 2*c*(x/b)], x] \text{/; RationalQ}\{q\} \&\& (\text{EqQ}\{q^2, 1\} \|\| \text{!RationalQ}\{b^2 - 4*a*c\}) \text{/; FreeQ}\{a, b, c\}, x]$

rule 1103 $\text{Int}[\text{((d_) + (e_.)*(x_))} / \text{((a_) + (b_.)*(x_) + (c_.)*(x_)^2)}, \text{x_Symbol}] \text{:> Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] \text{/; FreeQ}\{a, b, c, d, e\}, x] \&\& \text{EqQ}\{2*c*d - b*e, 0\}$

rule 1142 $\text{Int}[\text{((d_) + (e_.)*(x_))} / \text{((a_) + (b_.)*(x_) + (c_.)*(x_)^2)}, \text{x_Symbol}] \text{:> Simp}[(2*c*d - b*e)/(2*c) \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] \text{/; FreeQ}\{a, b, c, d, e\}, x]$

rule 6452 $\text{Int}[\text{((a_) + ArcTanh}\{c_.\} * (x_)^{\text{(n_.)}})^{\text{(p_.)}} * (b_.)^{\text{(m_.)}} * (x_)^{\text{(m_.)}}, \text{x_Symbol}] \text{:> Simp}[x^{\text{(m + 1)}} * ((a + b*\text{ArcTanh}[c*x^n])^{\text{p}} / (m + 1)), x] - \text{Simp}[b*c*n*(p/(m + 1)) \text{Int}[x^{\text{(m + n)}} * ((a + b*\text{ArcTanh}[c*x^n])^{\text{p - 1}} / (1 - c^2*x^{\text{(2*n)}})), x], x] \text{/; FreeQ}\{a, b, c, m, n\}, x] \&\& \text{IGtQ}\{p, 0\} \&\& (\text{EqQ}\{p, 1\} \|\| (\text{EqQ}\{n, 1\} \&\& \text{IntegerQ}\{m\})) \&\& \text{NeQ}\{m, -1\}$

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.21

method	result
derivativedivides	$\frac{x^4 a}{4} + \frac{b x^4 \operatorname{arctanh}(c x^{\frac{3}{2}})}{4} + \frac{3 b x^{\frac{5}{2}}}{20 c} - \frac{b \ln\left(\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{8 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}} \sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{16 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b \sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2}{\left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}\right)}{8 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}}$
default	$\frac{x^4 a}{4} + \frac{b x^4 \operatorname{arctanh}(c x^{\frac{3}{2}})}{4} + \frac{3 b x^{\frac{5}{2}}}{20 c} - \frac{b \ln\left(\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{8 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}} \sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{16 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b \sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2}{\left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}\right)}{8 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}}$
parts	$\frac{x^4 a}{4} + \frac{b x^4 \operatorname{arctanh}(c x^{\frac{3}{2}})}{4} + \frac{3 b x^{\frac{5}{2}}}{20 c} - \frac{b \ln\left(\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{8 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}} \sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{16 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b \sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2}{\left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}\right)}{8 c^3 \left(\frac{1}{c}\right)^{\frac{1}{3}}}$

input `int(x^3*(a+b*arctanh(c*x^(3/2))),x,method=_RETURNVERBOSE)`

output `1/4*x^4*a+1/4*b*x^4*arctanh(c*x^(3/2))+3/20*b*x^(5/2)/c-1/8*b/c^3/(1/c)^(1/3)*ln(x^(1/2)+(1/c)^(1/3))+1/16*b/c^3/(1/c)^(1/3)*ln(x-(1/c)^(1/3)*x^(1/2)+(1/c)^(2/3))+1/8*b/c^3*3^(1/2)/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x^(1/2)-1))+1/8*b/c^3/(1/c)^(1/3)*ln(x^(1/2)-(1/c)^(1/3))-1/16*b/c^3/(1/c)^(1/3)*ln(x+(1/c)^(1/3)*x^(1/2)+(1/c)^(2/3))+1/8*b/c^3*3^(1/2)/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x^(1/2)+1))`

Sympy [F(-1)]

Timed out.

$$\int x^3(a + \operatorname{barctanh}(cx^{3/2})) dx = \text{Timed out}$$

input `integrate(x**3*(a+b*atanh(c*x**(3/2))),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.08

$$\int x^3(a + \operatorname{barctanh}(cx^{3/2})) dx = \frac{1}{4}ax^4 + \frac{1}{80} \left(20x^4 \operatorname{artanh}\left(cx^{\frac{3}{2}}\right) + c \left(\frac{12x^{\frac{5}{2}}}{c^2} + \frac{10\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{\frac{2}{3}}\sqrt{x}+c^{\frac{1}{3}})}{3c^{\frac{1}{3}}}\right)}{c^{\frac{11}{3}}} + \frac{10\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{\frac{2}{3}}\sqrt{x}-c^{\frac{1}{3}})}{3c^{\frac{1}{3}}}\right)}{c^{\frac{11}{3}}} \right) \right)$$

input `integrate(x^3*(a+b*arctanh(c*x^(3/2))),x, algorithm="maxima")`

output `1/4*a*x^4 + 1/80*(20*x^4*arctanh(c*x^(3/2)) + c*(12*x^(5/2)/c^2 + 10*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*sqrt(x) + c^(1/3))/c^(1/3))/c^(11/3) + 10*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*sqrt(x) - c^(1/3))/c^(1/3))/c^(11/3) - 5*log(c^(2/3)*x + c^(1/3)*sqrt(x) + 1)/c^(11/3) + 5*log(c^(2/3)*x - c^(1/3)*sqrt(x) + 1)/c^(11/3) - 10*log((c^(1/3)*sqrt(x) + 1)/c^(1/3))/c^(11/3) + 10*log((c^(1/3)*sqrt(x) - 1)/c^(1/3))/c^(11/3)))*b`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.42

$$\int x^3(a + \operatorname{barctanh}(cx^{3/2})) dx = \frac{1}{4}ax^4 + \frac{1}{80} \left(10x^4 \log\left(-\frac{cx^{\frac{3}{2}} + 1}{cx^{\frac{3}{2}} - 1}\right) + c \left(\frac{12x^{\frac{5}{2}}}{c^2} - \frac{10\sqrt{3}\left(\frac{1}{2}i\sqrt{3} + \frac{1}{2}\right)^2 |c|^{\frac{4}{3}} \arctan\left(\frac{\sqrt{3}\left(2\sqrt{x} + (-\frac{1}{c})^{\frac{1}{3}}\right)}{3(-\frac{1}{c})^{\frac{1}{3}}}\right)}{c^5} + \frac{5\left(\frac{1}{2}i\sqrt{3}\right)}{c^5} \right) \right)$$

input `integrate(x^3*(a+b*arctanh(c*x^(3/2))),x, algorithm="giac")`

output `1/4*a*x^4 + 1/80*(10*x^4*log(-(c*x^(3/2) + 1)/(c*x^(3/2) - 1)) + c*(12*x^(5/2)/c^2 - 10*sqrt(3)*(1/2*I*sqrt(3) + 1/2)^2*abs(c)^(4/3)*arctan(1/3*sqrt(3)*(2*sqrt(x) + (-1/c)^(1/3))/(-1/c)^(1/3))/c^5 + 5*(1/2*I*sqrt(3) + 1/2)^2*abs(c)^(4/3)*log(x + sqrt(x)*(-1/c)^(1/3) + (-1/c)^(2/3))/c^5 - 10*(-1/c)^(2/3)*log(abs(sqrt(x) - (-1/c)^(1/3)))/c^3 + 10*sqrt(3)*abs(c)^(4/3)*arctan(1/3*sqrt(3)*c^(1/3)*(2*sqrt(x) + 1/c^(1/3)))/c^5 - 5*abs(c)^(4/3)*log(x + sqrt(x)/c^(1/3) + 1/c^(2/3))/c^5 + 10*log(abs(sqrt(x) - 1/c^(1/3)))/c^(11/3))*b`

Mupad [B] (verification not implemented)

Time = 16.65 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.44

$$\int x^3(a + \operatorname{barctanh}(cx^{3/2})) dx = \frac{ax^4}{4} + \frac{3bx^{5/2}}{20c} + \frac{b \ln\left(\frac{c^{1/3}\sqrt{x}-1}{c^{1/3}\sqrt{x}+1}\right)}{8c^{8/3}} + \frac{\ln(1 - cx^{3/2})\left(\frac{bx^4}{4} - \frac{bc^2x^7}{4}\right)}{2c^2x^3 - 2} + \frac{bx^4 \ln(cx^{3/2} + 1)}{8} + \frac{b \ln\left(\frac{\sqrt{3}+c^{2/3}x \operatorname{li}-c^{1/3}\sqrt{x}4i-\sqrt{3}c^{2/3}x+1i}{2c^{2/3}x+1+\sqrt{3}1i}\right)}{8c^{8/3}} \sqrt{-\frac{1}{2} + \frac{\sqrt{3}1i}{2}} + \frac{\sqrt{2}b \ln\left(\frac{\sqrt{3}c^{2/3}x+c^{2/3}x \operatorname{li}+c^{1/3}\sqrt{x}4i-\sqrt{3}+1i}{2c^{2/3}x+1-\sqrt{3}1i}\right)}{16c^{8/3}} \sqrt{1 + \sqrt{3}1i \operatorname{li}}$$

input `int(x^3*(a + b*atanh(c*x^(3/2))),x)`

output
$$\begin{aligned} & (a*x^4)/4 + (3*b*x^{(5/2)})/(20*c) + (b*\log((c^{(1/3)}*x^{(1/2)} - 1)/(c^{(1/3)}*x^{(1/2)} + 1)))/(8*c^{(8/3)}) + (\log(1 - c*x^{(3/2)})*((b*x^4)/4 - (b*c^2*x^7)/4))/((2*c^2*x^3 - 2) + (b*x^4*\log(c*x^{(3/2)} + 1))/8 + (b*\log((3^{(1/2)} + c^{(2/3)}*x*1i - c^{(1/3)}*x^{(1/2)}*4i - 3^{(1/2)}*c^{(2/3)}*x + 1i)/(3^{(1/2)}*1i + 2*c^{(2/3)}*x + 1))*((3^{(1/2)}*1i)/2 - 1/2)^{(1/2)})/(8*c^{(8/3)}) + (2^{(1/2)}*b*\log((c^{(2/3)}*x*1i - 3^{(1/2)} + c^{(1/3)}*x^{(1/2)}*4i + 3^{(1/2)}*c^{(2/3)}*x + 1i)/(2*c^{(2/3)}*x - 3^{(1/2)}*1i + 1))*(3^{(1/2)}*1i + 1)^{(1/2)*1i})/(16*c^{(8/3)}) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.75

$$\int x^3 (a + b \operatorname{arctanh}(cx^{3/2})) dx = \frac{10\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{x}c^{1/3}-1}{\sqrt{3}}\right) b + 10\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{x}c^{1/3}+1}{\sqrt{3}}\right) b + 20c^{2/3} \operatorname{atanh}(\sqrt{x}cx) b x^4 + 10a x^4}{10\sqrt{3}}$$

input `int(x^3*(a+b*atanh(c*x^(3/2))),x)`

output
$$\begin{aligned} & (10*\sqrt{3})*\operatorname{atan}((2*\sqrt{x})*c^{(1/3)} - 1)/\sqrt{3})*b + 10*\sqrt{3})*\operatorname{atan}((2*\sqrt{x})*c^{(1/3)} + 1)/\sqrt{3})*b + 20*c^{(2/3)}*\operatorname{atanh}(\sqrt{x}*c*x)*b*c^{(2/3)}*x^4 + 10*\operatorname{atanh}(\sqrt{x}*c*x)*b + 12*\sqrt{x}*c^{(2/3)}*b*c*x^2 + 20*c^{(2/3)}*a*c^{(2/3)}*x^4 - 15*\log(\sqrt{x}*c^{(2/3)} + c^{(1/3)})*b + 15*\log(\sqrt{x}*c^{(2/3)} - c^{(1/3)})*b)/(80*c^{(2/3)}*c^2) \end{aligned}$$

3.213 $\int x^2(a + b \operatorname{arctanh}(cx^{3/2})) dx$

Optimal result	1714
Mathematica [A] (verified)	1714
Rubi [A] (verified)	1715
Maple [A] (verified)	1717
Fricas [A] (verification not implemented)	1717
Sympy [F(-1)]	1718
Maxima [A] (verification not implemented)	1718
Giac [B] (verification not implemented)	1718
Mupad [B] (verification not implemented)	1719
Reduce [B] (verification not implemented)	1719

Optimal result

Integrand size = 16, antiderivative size = 49

$$\int x^2(a + b \operatorname{arctanh}(cx^{3/2})) dx = \frac{bx^{3/2}}{3c} - \frac{b \operatorname{arctanh}(cx^{3/2})}{3c^2} + \frac{1}{3}x^3(a + b \operatorname{arctanh}(cx^{3/2}))$$

output

```
1/3*b*x^(3/2)/c-1/3*b*arctanh(c*x^(3/2))/c^2+1/3*x^3*(a+b*arctanh(c*x^(3/2)))
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.53

$$\int x^2(a + b \operatorname{arctanh}(cx^{3/2})) dx = \frac{bx^{3/2}}{3c} + \frac{ax^3}{3} + \frac{1}{3}bx^3 \operatorname{arctanh}(cx^{3/2}) + \frac{b \log(1 - cx^{3/2})}{6c^2} - \frac{b \log(1 + cx^{3/2})}{6c^2}$$

input

```
Integrate[x^2*(a + b*ArcTanh[c*x^(3/2)]),x]
```

output

```
(b*x^(3/2))/(3*c) + (a*x^3)/3 + (b*x^3*ArcTanh[c*x^(3/2)])/3 + (b*Log[1 - c*x^(3/2)])/(6*c^2) - (b*Log[1 + c*x^(3/2)])/(6*c^2)
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.10, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6452, 843, 851, 807, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \left(a + \operatorname{barctanh}(cx^{3/2}) \right) dx \\
 & \quad \downarrow \text{6452} \\
 & \frac{1}{3}x^3 \left(a + \operatorname{barctanh}(cx^{3/2}) \right) - \frac{1}{2}bc \int \frac{x^{7/2}}{1 - c^2x^3} dx \\
 & \quad \downarrow \text{843} \\
 & \frac{1}{3}x^3 \left(a + \operatorname{barctanh}(cx^{3/2}) \right) - \frac{1}{2}bc \left(\frac{\int \frac{\sqrt{x}}{1 - c^2x^3} dx}{c^2} - \frac{2x^{3/2}}{3c^2} \right) \\
 & \quad \downarrow \text{851} \\
 & \frac{1}{3}x^3 \left(a + \operatorname{barctanh}(cx^{3/2}) \right) - \frac{1}{2}bc \left(\frac{2 \int \frac{x}{1 - c^2x^3} d\sqrt{x}}{c^2} - \frac{2x^{3/2}}{3c^2} \right) \\
 & \quad \downarrow \text{807} \\
 & \frac{1}{3}x^3 \left(a + \operatorname{barctanh}(cx^{3/2}) \right) - \frac{1}{2}bc \left(\frac{2 \int \frac{1}{1 - c^2x} dx^{3/2}}{3c^2} - \frac{2x^{3/2}}{3c^2} \right) \\
 & \quad \downarrow \text{219} \\
 & \frac{1}{3}x^3 \left(a + \operatorname{barctanh}(cx^{3/2}) \right) - \frac{1}{2}bc \left(\frac{2\operatorname{arctanh}(cx^{3/2})}{3c^3} - \frac{2x^{3/2}}{3c^2} \right)
 \end{aligned}$$

input `Int[x^2*(a + b*ArcTanh[c*x^(3/2)]),x]`

output `(x^3*(a + b*ArcTanh[c*x^(3/2)]))/3 - (b*c*((-2*x^(3/2))/(3*c^2) + (2*ArcTanh[c*x^(3/2)]/(3*c^3)))/2`

Definitions of rubi rules used

rule 219 $\text{Int}[(a_+) + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))* \text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 807 $\text{Int}(x_+)^{m_+}((a_+) + (b_+)(x_+)^{n_+})^{p_+}, x_Symbol] \rightarrow \text{With}\{k = \text{GCD}[m + 1, n]\}, \text{Simp}[1/k \ \text{Subst}[\text{Int}[x^{(m+1)/k - 1}(a + b*x^{(n/k)})^p], x, x^k], x] /; k \neq 1] /; \text{FreeQ}\{a, b, p\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$

rule 843 $\text{Int}(((c_+)(x_+))^{m_+}((a_+) + (b_+)(x_+)^{n_+})^{p_+}, x_Symbol] \rightarrow \text{Simp}[c^{(n-1)}*(c*x)^{m-n+1}*((a + b*x^n)^{(p+1})/(b*(m+n*p+1))), x] - \text{Simp}[a*c^n*((m-n+1)/(b*(m+n*p+1))) \ \text{Int}[(c*x)^{m-n}*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{GtQ}[m, n-1] \ \&\& \ \text{NeQ}[m+n*p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 851 $\text{Int}(((c_+)(x_+))^{m_+}((a_+) + (b_+)(x_+)^{n_+})^{p_+}, x_Symbol] \rightarrow \text{With}\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \ \text{Subst}[\text{Int}[x^{(k*(m+1)-1)}*(a + b*(x^{(k*n)})/c^n)^p], x, (c*x)^{1/k}], x] /; \text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 6452 $\text{Int}(((a_+) + \text{ArcTanh}[(c_+)(x_+)^{n_+}]*(b_+))^{p_+}(x_+)^{m_+}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c^n*(p/(m+1)) \ \text{Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)})/(1 - c^2*x^{(2*n)}), x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.12

method	result	size
parts	$\frac{ax^3}{3} + \frac{2b \left(\frac{c^2 x^3 \operatorname{arctanh}(cx^{\frac{3}{2}})}{2} + \frac{cx^{\frac{3}{2}}}{2} + \frac{\ln(cx^{\frac{3}{2}}-1)}{4} - \frac{\ln(cx^{\frac{3}{2}}+1)}{4} \right)}{3c^2}$	55
derivativedivides	$\frac{\frac{ac^2x^3}{3} + \left(\frac{c^2x^3 \operatorname{arctanh}(cx^{\frac{3}{2}})}{2} + \frac{cx^{\frac{3}{2}}}{2} + \frac{\ln(cx^{\frac{3}{2}}-1)}{4} - \frac{\ln(cx^{\frac{3}{2}}+1)}{4} \right)}{c^2}$	59
default	$\frac{ac^2x^3}{3} + \frac{2b \left(\frac{c^2x^3 \operatorname{arctanh}(cx^{\frac{3}{2}})}{2} + \frac{cx^{\frac{3}{2}}}{2} + \frac{\ln(cx^{\frac{3}{2}}-1)}{4} - \frac{\ln(cx^{\frac{3}{2}}+1)}{4} \right)}{c^2}$	59

input `int(x^2*(a+b*arctanh(c*x^(3/2))),x,method=_RETURNVERBOSE)`

output `1/3*a*x^3+2/3*b/c^2*(1/2*c^2*x^3*arctanh(c*x^(3/2))+1/2*c*x^(3/2)+1/4*ln(c*x^(3/2)-1)-1/4*ln(c*x^(3/2)+1))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.31

$$\int x^2(a + b \operatorname{arctanh}(cx^{3/2})) dx = \frac{2ac^2x^3 + 2bcx^{\frac{3}{2}} + (bc^2x^3 - b) \log\left(-\frac{c^2x^3 + 2cx^{\frac{3}{2}} + 1}{c^2x^3 - 1}\right)}{6c^2}$$

input `integrate(x^2*(a+b*arctanh(c*x^(3/2))),x, algorithm="fricas")`

output `1/6*(2*a*c^2*x^3 + 2*b*c*x^(3/2) + (b*c^2*x^3 - b)*log(-(c^2*x^3 + 2*c*x^(3/2) + 1)/(c^2*x^3 - 1)))/c^2`

Sympy [F(-1)]

Timed out.

$$\int x^2(a + \operatorname{barctanh}(cx^{3/2})) dx = \text{Timed out}$$

input `integrate(x**2*(a+b*atanh(c*x**(3/2))),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.18

$$\int x^2(a + \operatorname{barctanh}(cx^{3/2})) dx = \frac{1}{3}ax^3 + \frac{1}{6} \left(2x^3 \operatorname{artanh}(cx^{\frac{3}{2}}) + c \left(\frac{2x^{\frac{3}{2}}}{c^2} - \frac{\log(cx^{\frac{3}{2}} + 1)}{c^3} + \frac{\log(cx^{\frac{3}{2}} - 1)}{c^3} \right) \right) b$$

input `integrate(x^2*(a+b*arctanh(c*x^(3/2))),x, algorithm="maxima")`

output `1/3*a*x^3 + 1/6*(2*x^3*arctanh(c*x^(3/2)) + c*(2*x^(3/2)/c^2 - log(c*x^(3/2) + 1)/c^3 + log(c*x^(3/2) - 1)/c^3))*b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 97 vs. 2(37) = 74.

Time = 0.12 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.98

$$\int x^2(a + \operatorname{barctanh}(cx^{3/2})) dx = \frac{1}{3}ax^3 + \frac{2}{3}bc \left(\frac{1}{c^3 \left(\frac{cx^{\frac{3}{2}} + 1}{cx^{\frac{3}{2}} - 1} - 1 \right)} + \frac{\left(cx^{\frac{3}{2}} + 1 \right) \log \left(-\frac{cx^{\frac{3}{2}} + 1}{cx^{\frac{3}{2}} - 1} \right)}{\left(cx^{\frac{3}{2}} - 1 \right) c^3 \left(\frac{cx^{\frac{3}{2}} + 1}{cx^{\frac{3}{2}} - 1} - 1 \right)^2} \right)$$

input `integrate(x^2*(a+b*arctanh(c*x^(3/2))),x, algorithm="giac")`

output $\frac{1}{3}ax^3 + \frac{2}{3}bc \left(\frac{1}{c^3} \left(\frac{cx^{3/2} + 1}{cx^{3/2} - 1} - 1 \right) + \frac{cx^{3/2} + 1}{c^3} \log \left(\frac{-cx^{3/2} + 1}{cx^{3/2} - 1} \right) \right) / \left(\frac{cx^{3/2} - 1}{c^3} \left(\frac{cx^{3/2} + 1}{cx^{3/2} - 1} - 1 \right)^2 \right)$

Mupad [B] (verification not implemented)

Time = 4.42 (sec) , antiderivative size = 110, normalized size of antiderivative = 2.24

$$\int x^2 (a + b \operatorname{arctanh}(cx^{3/2})) dx = \frac{ax^3}{3} + \frac{bx^{3/2}}{3c} + \frac{b \ln \left(\frac{cx^{3/2}-1}{cx^{3/2}+1} \right)}{6c^2} + \frac{bx^3 \ln(cx^{3/2}+1)}{6} + \frac{bx^3 \ln(1-cx^{3/2})}{3(2c^2x^3-2)} - \frac{bc^2x^6 \ln(1-cx^{3/2})}{3(2c^2x^3-2)}$$

input `int(x^2*(a + b*atanh(c*x^(3/2))),x)`

output $\frac{(ax^3)/3 + (bx^{3/2})/(3c) + (b \log((cx^{3/2} - 1)/(cx^{3/2} + 1))) / (6c^2) + (bx^3 \log(cx^{3/2} + 1)) / 6 + (bx^3 \log(1 - cx^{3/2})) / (3(2c^2x^3 - 2)) - (bc^2x^6 \log(1 - cx^{3/2})) / (3(2c^2x^3 - 2))}{1}$

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.88

$$\int x^2 (a + b \operatorname{arctanh}(cx^{3/2})) dx = \frac{\operatorname{atanh}(\sqrt{x} cx) b c^2 x^3 - \operatorname{atanh}(\sqrt{x} cx) b + \sqrt{x} bcx + a c^2 x^3}{3c^2}$$

input `int(x^2*(a+b*atanh(c*x^(3/2))),x)`

output $\frac{(\operatorname{atanh}(\sqrt{x} cx) * b * c ** 2 * x ** 3 - \operatorname{atanh}(\sqrt{x} cx) * b + \sqrt{x} * b * c * x + a * c ** 2 * x ** 3) / (3 * c ** 2)}$

3.214 $\int x(a + b \operatorname{arctanh}(cx^{3/2})) dx$

Optimal result	1720
Mathematica [A] (verified)	1721
Rubi [A] (verified)	1721
Maple [A] (verified)	1726
Fricas [C] (verification not implemented)	1727
Sympy [F(-1)]	1728
Maxima [A] (verification not implemented)	1728
Giac [F]	1729
Mupad [B] (verification not implemented)	1729
Reduce [B] (verification not implemented)	1730

Optimal result

Integrand size = 14, antiderivative size = 160

$$\int x(a + b \operatorname{arctanh}(cx^{3/2})) dx = \frac{3b\sqrt{x}}{2c} + \frac{\sqrt{3}b \arctan\left(\frac{1-2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{4c^{4/3}} - \frac{\sqrt{3}b \arctan\left(\frac{1+2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{4c^{4/3}} - \frac{b \operatorname{arctanh}(\sqrt[3]{c}\sqrt{x})}{2c^{4/3}} + \frac{1}{2}x^2(a + b \operatorname{arctanh}(cx^{3/2})) - \frac{b \operatorname{arctanh}\left(\frac{\sqrt[3]{c}\sqrt{x}}{1+c^{2/3}x}\right)}{4c^{4/3}}$$

output

```
3/2*b*x^(1/2)/c+1/4*3^(1/2)*b*arctan(1/3*(1-2*c^(1/3)*x^(1/2))*3^(1/2))/c^(4/3)-1/4*3^(1/2)*b*arctan(1/3*(1+2*c^(1/3)*x^(1/2))*3^(1/2))/c^(4/3)-1/2*b*arctanh(c^(1/3)*x^(1/2))/c^(4/3)+1/2*x^2*(a+b*arctanh(c*x^(3/2)))-1/4*b*arctanh(c^(1/3)*x^(1/2)/(1+c^(2/3)*x))/c^(4/3)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.39

$$\int x(a + b \operatorname{arctanh}(cx^{3/2})) dx = \frac{3b\sqrt{x}}{2c} + \frac{ax^2}{2} - \frac{\sqrt{3}b \arctan\left(\frac{-1+2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{4c^{4/3}} - \frac{\sqrt{3}b \arctan\left(\frac{1+2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{4c^{4/3}} + \frac{1}{2}bx^2 \operatorname{arctanh}(cx^{3/2}) + \frac{b \log(1 - \sqrt[3]{c}\sqrt{x})}{4c^{4/3}} - \frac{b \log(1 + \sqrt[3]{c}\sqrt{x})}{4c^{4/3}} + \frac{b \log(1 - \sqrt[3]{c}\sqrt{x} + c^{2/3}x)}{8c^{4/3}} - \frac{b \log(1 + \sqrt[3]{c}\sqrt{x} + c^{2/3}x)}{8c^{4/3}}$$

input `Integrate[x*(a + b*ArcTanh[c*x^(3/2)]),x]`

output $(3*b*\operatorname{Sqrt}[x])/(2*c) + (a*x^2)/2 - (\operatorname{Sqrt}[3]*b*\operatorname{ArcTan}[(-1 + 2*c^{(1/3)}*\operatorname{Sqrt}[x])/ \operatorname{Sqrt}[3]])/(4*c^{(4/3)}) - (\operatorname{Sqrt}[3]*b*\operatorname{ArcTan}[(1 + 2*c^{(1/3)}*\operatorname{Sqrt}[x])/ \operatorname{Sqrt}[3]])/(4*c^{(4/3)}) + (b*x^2*\operatorname{ArcTanh}[c*x^{(3/2)}])/2 + (b*\operatorname{Log}[1 - c^{(1/3)}*\operatorname{Sqrt}[x]])/(4*c^{(4/3)}) - (b*\operatorname{Log}[1 + c^{(1/3)}*\operatorname{Sqrt}[x]])/(4*c^{(4/3)}) + (b*\operatorname{Log}[1 - c^{(1/3)}*\operatorname{Sqrt}[x] + c^{(2/3)}*x])/(8*c^{(4/3)}) - (b*\operatorname{Log}[1 + c^{(1/3)}*\operatorname{Sqrt}[x] + c^{(2/3)}*x])/(8*c^{(4/3)})$

Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.25, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$, Rules used = {6452, 843, 851, 754, 27, 219, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(a + b \operatorname{arctanh}(cx^{3/2})) dx$$

$$\downarrow 6452$$

$$\frac{1}{2}x^2(a + b \operatorname{arctanh}(cx^{3/2})) - \frac{3}{4}bc \int \frac{x^{5/2}}{1 - c^2x^3} dx$$

$$\downarrow 843$$

$$\begin{aligned}
& \frac{1}{2}x^2\left(a + \operatorname{barctanh}\left(cx^{3/2}\right)\right) - \frac{3}{4}bc\left(\frac{\int \frac{1}{\sqrt{x}(1-c^2x^3)}dx}{c^2} - \frac{2\sqrt{x}}{c^2}\right) \\
& \quad \downarrow 851 \\
& \frac{1}{2}x^2\left(a + \operatorname{barctanh}\left(cx^{3/2}\right)\right) - \frac{3}{4}bc\left(\frac{2\int \frac{1}{1-c^2x^3}d\sqrt{x}}{c^2} - \frac{2\sqrt{x}}{c^2}\right) \\
& \quad \downarrow 754 \\
& \frac{1}{2}x^2\left(a + \operatorname{barctanh}\left(cx^{3/2}\right)\right) - \\
& \frac{3}{4}bc\left(\frac{2\left(\frac{1}{3}\int \frac{1}{1-c^{2/3}x}d\sqrt{x} + \frac{1}{3}\int \frac{2-\sqrt[3]{c}\sqrt{x}}{2(c^{2/3}x-\sqrt[3]{c}\sqrt{x+1})}d\sqrt{x} + \frac{1}{3}\int \frac{\sqrt[3]{c}\sqrt{x+2}}{2(c^{2/3}x+\sqrt[3]{c}\sqrt{x+1})}d\sqrt{x}\right)}{c^2} - \frac{2\sqrt{x}}{c^2}\right) \\
& \quad \downarrow 27 \\
& \frac{1}{2}x^2\left(a + \operatorname{barctanh}\left(cx^{3/2}\right)\right) - \\
& \frac{3}{4}bc\left(\frac{2\left(\frac{1}{3}\int \frac{1}{1-c^{2/3}x}d\sqrt{x} + \frac{1}{6}\int \frac{2-\sqrt[3]{c}\sqrt{x}}{c^{2/3}x-\sqrt[3]{c}\sqrt{x+1}}d\sqrt{x} + \frac{1}{6}\int \frac{\sqrt[3]{c}\sqrt{x+2}}{c^{2/3}x+\sqrt[3]{c}\sqrt{x+1}}d\sqrt{x}\right)}{c^2} - \frac{2\sqrt{x}}{c^2}\right) \\
& \quad \downarrow 219 \\
& \frac{1}{2}x^2\left(a + \operatorname{barctanh}\left(cx^{3/2}\right)\right) - \\
& \frac{3}{4}bc\left(\frac{2\left(\frac{1}{6}\int \frac{2-\sqrt[3]{c}\sqrt{x}}{c^{2/3}x-\sqrt[3]{c}\sqrt{x+1}}d\sqrt{x} + \frac{1}{6}\int \frac{\sqrt[3]{c}\sqrt{x+2}}{c^{2/3}x+\sqrt[3]{c}\sqrt{x+1}}d\sqrt{x} + \frac{\operatorname{arctanh}\left(\sqrt[3]{c}\sqrt{x}\right)}{3\sqrt[3]{c}}\right)}{c^2} - \frac{2\sqrt{x}}{c^2}\right) \\
& \quad \downarrow 1142 \\
& \frac{1}{2}x^2\left(a + \operatorname{barctanh}\left(cx^{3/2}\right)\right) - \\
& \frac{3}{4}bc\left(\frac{2\left(\frac{1}{6}\left(\frac{3}{2}\int \frac{1}{c^{2/3}x-\sqrt[3]{c}\sqrt{x+1}}d\sqrt{x} - \frac{\int -\frac{\sqrt[3]{c}(1-2\sqrt[3]{c}\sqrt{x})}{c^{2/3}x-\sqrt[3]{c}\sqrt{x+1}}d\sqrt{x}}{2\sqrt[3]{c}}\right) + \frac{1}{6}\left(\frac{3}{2}\int \frac{1}{c^{2/3}x+\sqrt[3]{c}\sqrt{x+1}}d\sqrt{x} + \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{c}\sqrt{x+1})}{c^{2/3}x+\sqrt[3]{c}\sqrt{x+1}}d\sqrt{x}}{2\sqrt[3]{c}}\right)\right)}{c^2}\right)
\end{aligned}$$

$$\begin{array}{c} \downarrow 25 \\ \frac{1}{2}x^2(a + \operatorname{barctanh}(cx^{3/2})) - \\ \frac{3}{4}bc \left(\frac{2 \left(\frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x - \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} + \frac{\int \frac{\sqrt[3]{c}(1-2\sqrt[3]{c}\sqrt{x})}{c^{2/3}x - \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x}}{2\sqrt[3]{c}} \right) \right) + \frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} + \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{c}\sqrt{x+1})}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x}}{2\sqrt[3]{c}} \right)}{c^2} \right) \end{array}$$

$$\begin{array}{c} \downarrow 27 \\ \frac{1}{2}x^2(a + \operatorname{barctanh}(cx^{3/2})) - \\ \frac{3}{4}bc \left(\frac{2 \left(\frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x - \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} + \frac{1}{2} \int \frac{1-2\sqrt[3]{c}\sqrt{x}}{c^{2/3}x - \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} \right) \right) + \frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} + \frac{1}{2} \int \frac{2\sqrt[3]{c}\sqrt{x+1}}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} \right)}{c^2} \right) \end{array}$$

$$\begin{array}{c} \downarrow 1082 \\ \frac{1}{2}x^2(a + \operatorname{barctanh}(cx^{3/2})) - \\ \frac{3}{4}bc \left(\frac{2 \left(\frac{1}{6} \left(\frac{1}{2} \int \frac{1-2\sqrt[3]{c}\sqrt{x}}{c^{2/3}x - \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} + \frac{3 \int \frac{1}{-x-3} d(1-2\sqrt[3]{c}\sqrt{x})}{\sqrt[3]{c}} \right) \right) + \frac{1}{6} \left(\frac{1}{2} \int \frac{2\sqrt[3]{c}\sqrt{x+1}}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} - \frac{3 \int \frac{1}{-x-3} d(2\sqrt[3]{c}\sqrt{x+1})}{\sqrt[3]{c}} \right)}{c^2} \right) \end{array}$$

$$\begin{array}{c} \downarrow 217 \\ \frac{1}{2}x^2(a + \operatorname{barctanh}(cx^{3/2})) - \\ \frac{3}{4}bc \left(\frac{2 \left(\frac{1}{6} \left(\frac{1}{2} \int \frac{1-2\sqrt[3]{c}\sqrt{x}}{c^{2/3}x - \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} - \frac{\sqrt{3} \arctan\left(\frac{1-2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{\sqrt[3]{c}} \right) \right) + \frac{1}{6} \left(\frac{1}{2} \int \frac{2\sqrt[3]{c}\sqrt{x+1}}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} + \frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{c}\sqrt{x+1}}{\sqrt{3}}\right)}{\sqrt[3]{c}} \right)}{c^2} \right) \end{array}$$

$$\downarrow 1103$$

$$\frac{3}{4}bc \left(\frac{\frac{1}{2}x^2(a + \operatorname{arctanh}(cx^{3/2})) - 2 \left(\frac{1}{6} \left(-\frac{\sqrt{3} \arctan\left(\frac{1-2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right) - \log(c^{2/3}x - \sqrt[3]{c}\sqrt{x+1})}{\sqrt[3]{c}} \right) + \frac{1}{6} \left(\frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{c}\sqrt{x+1}}{\sqrt{3}}\right) + \log(c^{2/3}x + \sqrt[3]{c}\sqrt{x+1})}{\sqrt[3]{c}} \right)}{c^2} \right) + \dots \right)$$

input `Int[x*(a + b*ArcTanh[c*x^(3/2)]),x]`

output $(x^2*(a + b*ArcTanh[c*x^(3/2)]))/2 - (3*b*c*((-2*sqrt[x])/c^2 + (2*(ArcTanh[c^(1/3)*sqrt[x]]/(3*c^(1/3)) + (-((sqrt[3]*ArcTan[(1 - 2*c^(1/3)*sqrt[x])/sqrt[3]])/c^(1/3)) - Log[1 - c^(1/3)*sqrt[x] + c^(2/3)*x]/(2*c^(1/3)))/6 + ((sqrt[3]*ArcTan[(1 + 2*c^(1/3)*sqrt[x])/sqrt[3]])/c^(1/3) + Log[1 + c^(1/3)*sqrt[x] + c^(2/3)*x]/(2*c^(1/3)))/6))/c^2)/4$

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 754 `Int[((a_) + (b_.)*(x_)^(n_))^(n_)^(-1), x_Symbol] := Module[{r = Numerator[Rt[-a/b, n]], s = Denominator[Rt[-a/b, n]], k, u}, Simp[u = Int[(r - s*Cos[(2*k*Pi)/n]*x)/(r^2 - 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x] + Int[(r + s*Cos[(2*k*Pi)/n]*x)/(r^2 + 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x]; 2*(r^2/(a*n)) Int[1/(r^2 - s^2*x^2), x] + 2*(r/(a*n)) Sum[u, {k, 1, (n - 2)/4}], x] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && NegQ[a/b]`

rule 843 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Simp[a*c^n*(m - n + 1)/(b*(m + n*p + 1)) Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 851 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n)]^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.21

method	result
derivativedivides	$\frac{ax^2}{2} + \frac{x^2b \operatorname{arctanh}(cx^{\frac{3}{2}})}{2} + \frac{3b\sqrt{x}}{2c} + \frac{b \ln\left(\sqrt{x} - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4c^2\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{b \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{8c^2\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{1}{c}\right)^{\frac{1}{3}}\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}}{\left(\frac{1}{c}\right)^{\frac{1}{3}}}\right)}{4c^2\left(\frac{1}{c}\right)^{\frac{2}{3}}}$
default	$\frac{ax^2}{2} + \frac{x^2b \operatorname{arctanh}(cx^{\frac{3}{2}})}{2} + \frac{3b\sqrt{x}}{2c} + \frac{b \ln\left(\sqrt{x} - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4c^2\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{b \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{8c^2\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{1}{c}\right)^{\frac{1}{3}}\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}}{\left(\frac{1}{c}\right)^{\frac{1}{3}}}\right)}{4c^2\left(\frac{1}{c}\right)^{\frac{2}{3}}}$
parts	$\frac{ax^2}{2} + \frac{x^2b \operatorname{arctanh}(cx^{\frac{3}{2}})}{2} + \frac{3b\sqrt{x}}{2c} + \frac{b \ln\left(\sqrt{x} - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4c^2\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{b \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{8c^2\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{1}{c}\right)^{\frac{1}{3}}\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}}{\left(\frac{1}{c}\right)^{\frac{1}{3}}}\right)}{4c^2\left(\frac{1}{c}\right)^{\frac{2}{3}}}$

input

```
int(x*(a+b*arctanh(c*x^(3/2))),x,method=_RETURNVERBOSE)
```

output

```
1/2*a*x^2+1/2*x^2*b*arctanh(c*x^(3/2))+3/2*b*x^(1/2)/c+1/4*b/c^2/(1/c)^(2/
3)*ln(x^(1/2)-(1/c)^(1/3))-1/8*b/c^2/(1/c)^(2/3)*ln(x+(1/c)^(1/3)*x^(1/2)+
(1/c)^(2/3))-1/4*b/c^2/(1/c)^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/
3)*x^(1/2)+1))-1/4*b/c^2/(1/c)^(2/3)*ln(x^(1/2)+(1/c)^(1/3))+1/8*b/c^2/(1/
c)^(2/3)*ln(x-(1/c)^(1/3)*x^(1/2)+(1/c)^(2/3))-1/4*b/c^2/(1/c)^(2/3)*3^(1/
2)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x^(1/2)-1))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.78 (sec) , antiderivative size = 1488, normalized size of antiderivative = 9.30

$$\int x(a + \operatorname{arctanh}(cx^{3/2})) dx = \text{Too large to display}$$

input `integrate(x*(a+b*arctanh(c*x^(3/2))),x, algorithm="fricas")`

output

```

1/16*(8*a*c*x^2 - 2*((1/2)^(1/3)*(b^3 - (c^4 - 1)*b^3/c^4 + b^3/c^4)^(1/3)
*(I*sqrt(3) + 1) + 2*b)*c*log(1/2*((1/2)^(1/3)*(b^3 - (c^4 - 1)*b^3/c^4 +
b^3/c^4)^(1/3)*(I*sqrt(3) + 1) + 2*b)*c - b*c + b*sqrt(x)) - 4*(2*(-1/128*
b^3 + 1/128*(c^4 - 1)*b^3/c^4 + 1/128*b^3/c^4)^(1/3)*(I*sqrt(3) + 1) - b)*
c*log((2*(-1/128*b^3 + 1/128*(c^4 - 1)*b^3/c^4 + 1/128*b^3/c^4)^(1/3)*(I*sq
rt(3) + 1) - b)*c + b*c + b*sqrt(x)) + (((1/2)^(1/3)*(b^3 - (c^4 - 1)*b^3
/c^4 + b^3/c^4)^(1/3)*(I*sqrt(3) + 1) + 2*b)*c - 6*b*c - 2*sqrt(-3/4*((1/2)
)^(1/3)*(b^3 - (c^4 - 1)*b^3/c^4 + b^3/c^4)^(1/3)*(I*sqrt(3) + 1) + 2*b)^2
+ 3*((1/2)^(1/3)*(b^3 - (c^4 - 1)*b^3/c^4 + b^3/c^4)^(1/3)*(I*sqrt(3) + 1)
) + 2*b)*b - 3*b^2)*c*log(-1/2*((1/2)^(1/3)*(b^3 - (c^4 - 1)*b^3/c^4 + b^
3/c^4)^(1/3)*(I*sqrt(3) + 1) + 2*b)*c + b*c + sqrt(-3/4*((1/2)^(1/3)*(b^3
- (c^4 - 1)*b^3/c^4 + b^3/c^4)^(1/3)*(I*sqrt(3) + 1) + 2*b)^2 + 3*((1/2)^(
1/3)*(b^3 - (c^4 - 1)*b^3/c^4 + b^3/c^4)^(1/3)*(I*sqrt(3) + 1) + 2*b)*b -
3*b^2)*c + 2*b*sqrt(x)) + (((1/2)^(1/3)*(b^3 - (c^4 - 1)*b^3/c^4 + b^3/c^4)
)^(1/3)*(I*sqrt(3) + 1) + 2*b)*c - 6*b*c + 2*sqrt(-3/4*((1/2)^(1/3)*(b^3 -
(c^4 - 1)*b^3/c^4 + b^3/c^4)^(1/3)*(I*sqrt(3) + 1) + 2*b)^2 + 3*((1/2)^(1
/3)*(b^3 - (c^4 - 1)*b^3/c^4 + b^3/c^4)^(1/3)*(I*sqrt(3) + 1) + 2*b)*b - 3
*b^2)*c*log(-1/2*((1/2)^(1/3)*(b^3 - (c^4 - 1)*b^3/c^4 + b^3/c^4)^(1/3)*(
I*sqrt(3) + 1) + 2*b)*c + b*c - sqrt(-3/4*((1/2)^(1/3)*(b^3 - (c^4 - 1)*b^
3/c^4 + b^3/c^4)^(1/3)*(I*sqrt(3) + 1) + 2*b)^2 + 3*((1/2)^(1/3)*(b^3 - ...

```

Sympy [F(-1)]

Timed out.

$$\int x(a + \operatorname{barctanh}(cx^{3/2})) dx = \text{Timed out}$$

input `integrate(x*(a+b*atanh(c*x**(3/2))),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.08

$$\int x(a + \operatorname{barctanh}(cx^{3/2})) dx = \frac{1}{2} ax^2 + \frac{1}{8} \left(4x^2 \operatorname{artanh}\left(cx^{\frac{3}{2}}\right) - c \left(\frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{\frac{2}{3}}\sqrt{x}+c^{\frac{1}{3}})}{3c^{\frac{1}{3}}}\right)}{c^{\frac{7}{3}}} + \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{\frac{2}{3}}\sqrt{x}-c^{\frac{1}{3}})}{3c^{\frac{1}{3}}}\right)}{c^{\frac{7}{3}}} + \frac{\log\left(c^{\frac{2}{3}}x + \dots\right)}{c^{\frac{7}{3}}} \right) \right)$$

input `integrate(x*(a+b*arctanh(c*x^(3/2))),x, algorithm="maxima")`

output `1/2*a*x^2 + 1/8*(4*x^2*arctanh(c*x^(3/2)) - c*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*sqrt(x) + c^(1/3))/c^(1/3))/c^(7/3) + 2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*sqrt(x) - c^(1/3))/c^(1/3))/c^(7/3) + log(c^(2/3)*x + c^(1/3)*sqrt(x) + 1)/c^(7/3) - log(c^(2/3)*x - c^(1/3)*sqrt(x) + 1)/c^(7/3) + 2*log((c^(1/3)*sqrt(x) + 1)/c^(1/3))/c^(7/3) - 2*log((c^(1/3)*sqrt(x) - 1)/c^(1/3))/c^(7/3) - 12*sqrt(x)/c^2)*b`

Giac [F]

$$\int x(a + \operatorname{barctanh}(cx^{3/2})) dx = \int \left(b \operatorname{artanh}\left(cx^{\frac{3}{2}}\right) + a \right) x dx$$

input `integrate(x*(a+b*arctanh(c*x^(3/2))),x, algorithm="giac")`

output `integrate((b*arctanh(c*x^(3/2)) + a)*x, x)`

Mupad [B] (verification not implemented)

Time = 15.60 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.54

$$\begin{aligned} \int x(a + \operatorname{barctanh}(cx^{3/2})) dx &= \frac{ax^2}{2} + \frac{3b\sqrt{x}}{2c} + \frac{b \ln\left(\frac{c^{1/3}\sqrt{x}-1}{c^{1/3}\sqrt{x}+1}\right)}{4c^{4/3}} \\ &+ \frac{\ln(1 - cx^{3/2}) \left(\frac{bx^2}{2} - \frac{bc^2x^5}{2}\right)}{2c^2x^3 - 2} + \frac{bx^2 \ln(cx^{3/2} + 1)}{4} \\ &+ \frac{b \ln\left(\frac{\sqrt{3}c^{2/3}x + c^{2/3}x \operatorname{li} - c^{1/3}\sqrt{x}4i - \sqrt{3} + \operatorname{li}}{2c^{2/3}x + 1 - \sqrt{3} \operatorname{li}}\right)}{4c^{4/3}} \sqrt{-\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}} \\ &+ \frac{\sqrt{2}b \ln\left(\frac{2\sqrt{2}-c^{1/3}\sqrt{x}(1+\sqrt{3} \operatorname{li})^{5/2} \operatorname{li} - \sqrt{2}c^{2/3}x + \sqrt{2}\sqrt{3}c^{2/3}x \operatorname{li}}{2c^{2/3}x + 1 + \sqrt{3} \operatorname{li}}\right)}{8c^{4/3}} \sqrt{1 + \sqrt{3} \operatorname{li} \operatorname{li}} \end{aligned}$$

input `int(x*(a + b*atanh(c*x^(3/2))),x)`

output `(a*x^2)/2 + (3*b*x^(1/2))/(2*c) + (b*log((c^(1/3)*x^(1/2) - 1)/(c^(1/3)*x^(1/2) + 1)))/(4*c^(4/3)) + (log(1 - c*x^(3/2))*((b*x^2)/2 - (b*c^2*x^5)/2))/(2*c^2*x^3 - 2) + (b*x^2*log(c*x^(3/2) + 1))/4 + (b*log((c^(2/3)*x*1i - 3^(1/2) - c^(1/3)*x^(1/2)*4i + 3^(1/2)*c^(2/3)*x + 1i)/(2*c^(2/3)*x - 3^(1/2)*1i + 1))*((3^(1/2)*1i)/2 - 1/2)^(1/2))/(4*c^(4/3)) + (2^(1/2)*b*log((2*2^(1/2) - c^(1/3)*x^(1/2)*(3^(1/2)*1i + 1)^(5/2)*1i - 2^(1/2)*c^(2/3)*x + 2^(1/2)*3^(1/2)*c^(2/3)*x*1i)/(3^(1/2)*1i + 2*c^(2/3)*x + 1))*(3^(1/2)*1i + 1)^(1/2)*1i)/(8*c^(4/3))`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.73

$$\int x(a + b \operatorname{arctanh}(cx^{3/2})) dx = \frac{-2\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{x}c^{1/3}-1}{\sqrt{3}}\right)b - 2\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{x}c^{1/3}+1}{\sqrt{3}}\right)b + 4c^{4/3} \operatorname{atanh}(\sqrt{x}cx)bx^2 + 2a \operatorname{arctanh}(cx^{3/2})}{8c^{1/3}c}$$

input `int(x*(a+b*atanh(c*x^(3/2))),x)`output `(- 2*sqrt(3)*atan((2*sqrt(x)*c**(1/3) - 1)/sqrt(3))*b - 2*sqrt(3)*atan((2*sqrt(x)*c**(1/3) + 1)/sqrt(3))*b + 4*c**(1/3)*atanh(sqrt(x)*c*x)*b*c*x**2 + 2*atanh(sqrt(x)*c*x)*b + 12*sqrt(x)*c**(1/3)*b + 4*c**(1/3)*a*c*x**2 - 3*log(sqrt(x)*c**(2/3) + c**(1/3))*b + 3*log(sqrt(x)*c**(2/3) - c**(1/3))*b)/(8*c**(1/3)*c)`

3.215 $\int (a + b \operatorname{arctanh}(cx^{3/2})) dx$

Optimal result	1731
Mathematica [A] (verified)	1732
Rubi [A] (verified)	1732
Maple [A] (verified)	1733
Fricas [C] (verification not implemented)	1734
Sympy [F(-1)]	1735
Maxima [A] (verification not implemented)	1735
Giac [A] (verification not implemented)	1736
Mupad [B] (verification not implemented)	1736
Reduce [B] (verification not implemented)	1737

Optimal result

Integrand size = 12, antiderivative size = 140

$$\int (a + b \operatorname{arctanh}(cx^{3/2})) dx = ax - \frac{\sqrt{3}b \arctan\left(\frac{1-2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{2c^{2/3}} + \frac{\sqrt{3}b \arctan\left(\frac{1+2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{2c^{2/3}} - \frac{b \operatorname{arctanh}(\sqrt[3]{c}\sqrt{x})}{c^{2/3}} + b x \operatorname{arctanh}(cx^{3/2}) - \frac{b \operatorname{arctanh}\left(\frac{\sqrt[3]{c}\sqrt{x}}{1+c^{2/3}x}\right)}{2c^{2/3}}$$

output

```
a*x-1/2*3^(1/2)*b*arctan(1/3*(1-2*c^(1/3)*x^(1/2))*3^(1/2))/c^(2/3)+1/2*3^(1/2)*b*arctan(1/3*(1+2*c^(1/3)*x^(1/2))*3^(1/2))/c^(2/3)-b*arctanh(c^(1/3)*x^(1/2))/c^(2/3)+b*x*arctanh(c*x^(3/2))-1/2*b*arctanh(c^(1/3)*x^(1/2)/(1+c^(2/3)*x))/c^(2/3)
```


Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.81

$$\int (a + b \operatorname{arctanh}(cx^{3/2})) dx = ax + b x \operatorname{arctanh}(cx^{3/2}) - \frac{b \left(\sqrt{3} \left(\arctan \left(\frac{1-2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}} \right) - \arctan \left(\frac{1+2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}} \right) \right) + 2 \operatorname{arctanh}(\sqrt[3]{c}\sqrt{x}) + \operatorname{arctanh} \left(\frac{\sqrt[3]{c}\sqrt{x}}{1+c^{2/3}x} \right) \right)}{2c^{2/3}}$$

input `Integrate[a + b*ArcTanh[c*x^(3/2)],x]`

output `a*x + b*x*ArcTanh[c*x^(3/2)] - (b*(Sqrt[3]*(ArcTan[(1 - 2*c^(1/3)*Sqrt[x])/Sqrt[3]] - ArcTan[(1 + 2*c^(1/3)*Sqrt[x])/Sqrt[3]]) + 2*ArcTanh[c^(1/3)*Sqrt[x]] + ArcTanh[(c^(1/3)*Sqrt[x]/(1 + c^(2/3)*x)]))/(2*c^(2/3))`

Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.21, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{arctanh}(cx^{3/2})) dx$$

↓ 2009

$$ax - \frac{\sqrt{3}b \arctan \left(\frac{1-2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}} \right)}{2c^{2/3}} + \frac{\sqrt{3}b \arctan \left(\frac{2\sqrt[3]{c}\sqrt{x}+1}{\sqrt{3}} \right)}{2c^{2/3}} - \frac{b \operatorname{arctanh}(\sqrt[3]{c}\sqrt{x})}{c^{2/3}} + b x \operatorname{arctanh}(cx^{3/2}) + \frac{b \log(c^{2/3}x - \sqrt[3]{c}\sqrt{x} + 1)}{4c^{2/3}} - \frac{b \log(c^{2/3}x + \sqrt[3]{c}\sqrt{x} + 1)}{4c^{2/3}}$$

input `Int[a + b*ArcTanh[c*x^(3/2)],x]`

output

```
a*x - (Sqrt[3]*b*ArcTan[(1 - 2*c^(1/3)*Sqrt[x])/Sqrt[3]])/(2*c^(2/3)) + (Sqrt[3]*b*ArcTan[(1 + 2*c^(1/3)*Sqrt[x])/Sqrt[3]])/(2*c^(2/3)) - (b*ArcTanh[c^(1/3)*Sqrt[x]])/c^(2/3) + b*x*ArcTanh[c*x^(3/2)] + (b*Log[1 - c^(1/3)*Sqrt[x] + c^(2/3)*x])/(4*c^(2/3)) - (b*Log[1 + c^(1/3)*Sqrt[x] + c^(2/3)*x])/(4*c^(2/3))
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.28

method	result
derivativedivides	$ax + bx \operatorname{arctanh}\left(cx^{\frac{3}{2}}\right) + \frac{b \ln\left(\sqrt{x} - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{2c\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{b \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{4c\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2\sqrt{x}}{\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{2\sqrt{x}}{\left(\frac{1}{c}\right)^{\frac{1}{3}}}\right)}{3}\right)}{2c\left(\frac{1}{c}\right)^{\frac{1}{3}}}$
default	$ax + bx \operatorname{arctanh}\left(cx^{\frac{3}{2}}\right) + \frac{b \ln\left(\sqrt{x} - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{2c\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{b \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{4c\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2\sqrt{x}}{\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{2\sqrt{x}}{\left(\frac{1}{c}\right)^{\frac{1}{3}}}\right)}{3}\right)}{2c\left(\frac{1}{c}\right)^{\frac{1}{3}}}$
parts	$ax + bx \operatorname{arctanh}\left(cx^{\frac{3}{2}}\right) + \frac{b \ln\left(\sqrt{x} - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{2c\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{b \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{4c\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2\sqrt{x}}{\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{2\sqrt{x}}{\left(\frac{1}{c}\right)^{\frac{1}{3}}}\right)}{3}\right)}{2c\left(\frac{1}{c}\right)^{\frac{1}{3}}}$

input

```
int(a+b*arctanh(c*x^(3/2)),x,method=_RETURNVERBOSE)
```

output

```
a*x+b*x*arctanh(c*x^(3/2))+1/2*b/c/(1/c)^(1/3)*ln(x^(1/2)-(1/c)^(1/3))-1/4
*b/c/(1/c)^(1/3)*ln(x+(1/c)^(1/3)*x^(1/2)+(1/c)^(2/3))+1/2*b*3^(1/2)/c/(1/
c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x^(1/2)+1))-1/2*b/c/(1/c)^(1/3)
*ln(x^(1/2)+(1/c)^(1/3))+1/4*b/c/(1/c)^(1/3)*ln(x-(1/c)^(1/3)*x^(1/2)+(1/c
)^(2/3))+1/2*b*3^(1/2)/c/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x^(
1/2)-1))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.85 (sec) , antiderivative size = 1914, normalized size of antiderivative = 13.67

$$\int (a + b \operatorname{arctanh}(cx^{3/2})) dx = \text{Too large to display}$$

input

```
integrate(a+b*arctanh(c*x^(3/2)),x, algorithm="fricas")
```

output

```
a*x + 1/8*((1/2)^(1/3)*(b^3 - (c^2 - 1)*b^3/c^2 + b^3/c^2)^(1/3)*(I*sqrt(3
) + 1) - 4*b - 2*sqrt(-3/4*((1/2)^(1/3)*(b^3 - (c^2 - 1)*b^3/c^2 + b^3/c^2
)^(1/3)*(I*sqrt(3) + 1) + 2*b)^2 + 3*((1/2)^(1/3)*(b^3 - (c^2 - 1)*b^3/c^2
+ b^3/c^2)^(1/3)*(I*sqrt(3) + 1) + 2*b)*b - 3*b^2))*log(1/4*((1/2)^(1/3)*
(b^3 - (c^2 - 1)*b^3/c^2 + b^3/c^2)^(1/3)*(I*sqrt(3) + 1) + 2*b)^2*c - ((1
/2)^(1/3)*(b^3 - (c^2 - 1)*b^3/c^2 + b^3/c^2)^(1/3)*(I*sqrt(3) + 1) + 2*b)
*b*c + b^2*c + 2*b^2*sqrt(x) + 1/2*sqrt(-3/4*((1/2)^(1/3)*(b^3 - (c^2 - 1)
*b^3/c^2 + b^3/c^2)^(1/3)*(I*sqrt(3) + 1) + 2*b)^2 + 3*((1/2)^(1/3)*(b^3 -
(c^2 - 1)*b^3/c^2 + b^3/c^2)^(1/3)*(I*sqrt(3) + 1) + 2*b)*b - 3*b^2))*(((1
/2)^(1/3)*(b^3 - (c^2 - 1)*b^3/c^2 + b^3/c^2)^(1/3)*(I*sqrt(3) + 1) + 2*b)
*c - 2*b*c)) + 1/8*((1/2)^(1/3)*(b^3 - (c^2 - 1)*b^3/c^2 + b^3/c^2)^(1/3)*
(I*sqrt(3) + 1) - 4*b + 2*sqrt(-3/4*((1/2)^(1/3)*(b^3 - (c^2 - 1)*b^3/c^2
+ b^3/c^2)^(1/3)*(I*sqrt(3) + 1) + 2*b)^2 + 3*((1/2)^(1/3)*(b^3 - (c^2 - 1)
)*b^3/c^2 + b^3/c^2)^(1/3)*(I*sqrt(3) + 1) + 2*b)*b - 3*b^2))*log(1/4*((1/
2)^(1/3)*(b^3 - (c^2 - 1)*b^3/c^2 + b^3/c^2)^(1/3)*(I*sqrt(3) + 1) + 2*b)^
2*c - ((1/2)^(1/3)*(b^3 - (c^2 - 1)*b^3/c^2 + b^3/c^2)^(1/3)*(I*sqrt(3) +
1) + 2*b)*b*c + b^2*c + 2*b^2*sqrt(x) - 1/2*sqrt(-3/4*((1/2)^(1/3)*(b^3 -
(c^2 - 1)*b^3/c^2 + b^3/c^2)^(1/3)*(I*sqrt(3) + 1) + 2*b)^2 + 3*((1/2)^(1/
3)*(b^3 - (c^2 - 1)*b^3/c^2 + b^3/c^2)^(1/3)*(I*sqrt(3) + 1) + 2*b)*b - 3*
b^2))*(((1/2)^(1/3)*(b^3 - (c^2 - 1)*b^3/c^2 + b^3/c^2)^(1/3)*(I*sqrt(3)...
```

Sympy [F(-1)]

Timed out.

$$\int (a + b \operatorname{arctanh}(cx^{3/2})) dx = \text{Timed out}$$

input `integrate(a+b*atanh(c*x**(3/2)),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.13

$$\int (a + b \operatorname{arctanh}(cx^{3/2})) dx = \frac{1}{4} \left(c \left(\frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{2/3}\sqrt{x}+c^{1/3})}{3c^{1/3}}\right)}{c^{5/3}} + \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{2/3}\sqrt{x}-c^{1/3})}{3c^{1/3}}\right)}{c^{5/3}} - \frac{\log\left(c^{2/3}x + ax\right)}{c^{5/3}} \right) \right)$$

input `integrate(a+b*arctanh(c*x^(3/2)),x, algorithm="maxima")`

output `1/4*(c*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*sqrt(x) + c^(1/3))/c^(1/3)))/c^(5/3) + 2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*sqrt(x) - c^(1/3))/c^(1/3))/c^(5/3) - log(c^(2/3)*x + c^(1/3)*sqrt(x) + 1)/c^(5/3) + log(c^(2/3)*x - c^(1/3)*sqrt(x) + 1)/c^(5/3) - 2*log((c^(1/3)*sqrt(x) + 1)/c^(1/3))/c^(5/3) + 2*log((c^(1/3)*sqrt(x) - 1)/c^(1/3))/c^(5/3)) + 4*x*arctanh(c*x^(3/2))*b + a*x`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.33

$$\int (a + b \operatorname{arctanh}(cx^{3/2})) dx = \frac{1}{4} \left(c \left(\frac{2\sqrt{3}|c|^{1/3} \arctan\left(\frac{1}{3}\sqrt{3}\left(2\sqrt{x} + \frac{1}{|c|^{1/3}}\right)|c|^{1/3}\right)}{c^2} + \frac{2\sqrt{3}|c|^{1/3} \arctan\left(\frac{1}{3}\sqrt{3}\left(2\sqrt{x} - \frac{1}{|c|^{1/3}}\right)|c|^{1/3}\right)}{c^2} \right) + ax \right.$$

input `integrate(a+b*arctanh(c*x^(3/2)),x, algorithm="giac")`

output

```
1/4*(c*(2*sqrt(3)*abs(c)^(1/3)*arctan(1/3*sqrt(3)*(2*sqrt(x) + 1/abs(c)^(1/3)))*abs(c)^(1/3))/c^2 + 2*sqrt(3)*abs(c)^(1/3)*arctan(1/3*sqrt(3)*(2*sqrt(x) - 1/abs(c)^(1/3)))*abs(c)^(1/3))/c^2 - abs(c)^(1/3)*log(x + sqrt(x)/abs(c)^(1/3) + 1/abs(c)^(2/3))/c^2 + abs(c)^(1/3)*log(x - sqrt(x)/abs(c)^(1/3) + 1/abs(c)^(2/3))/c^2 - 2*abs(c)^(1/3)*log(sqrt(x) + 1/abs(c)^(1/3))/c^2 + 2*abs(c)^(1/3)*log(abs(sqrt(x) - 1/abs(c)^(1/3)))/c^2 + 2*x*log(-(c*x^(3/2) + 1)/(c*x^(3/2) - 1)))*b + a*x
```

Mupad [B] (verification not implemented)

Time = 8.19 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.76

$$\int (a + b \operatorname{atanh}(cx^{3/2})) dx = ax + bx \operatorname{atanh}(cx^{3/2}) - \frac{b \operatorname{atanh}(c^{1/3} \sqrt{x})}{c^{2/3}} + \frac{b \operatorname{atanh}\left(\frac{486 c^8 \sqrt{x}}{-243 c^{23/3} + \sqrt{3} c^{23/3} 243i}\right) (1 + \sqrt{3} i)}{2 c^{2/3}} + \frac{b \operatorname{atanh}\left(\frac{486 c^8 \sqrt{x}}{243 c^{23/3} + \sqrt{3} c^{23/3} 243i}\right) (-1 + \sqrt{3} i)}{2 c^{2/3}}$$

input `int(a + b*atanh(c*x^(3/2)),x)`

output

```
a*x + b*x*atanh(c*x^(3/2)) - (b*atanh(c^(1/3)*x^(1/2)))/c^(2/3) + (b*atanh
((486*c^8*x^(1/2))/(3^(1/2)*c^(23/3)*243i - 243*c^(23/3)))*(3^(1/2)*1i + 1
))/ (2*c^(2/3)) + (b*atanh((486*c^8*x^(1/2))/(3^(1/2)*c^(23/3)*243i + 243*c
^(23/3)))*(3^(1/2)*1i - 1))/(2*c^(2/3))
```

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.75

$$\int (a + b \operatorname{arctanh}(cx^{3/2})) dx = \frac{2\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{x}c^{1/3}-1}{\sqrt{3}}\right) b + 2\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{x}c^{1/3}+1}{\sqrt{3}}\right) b + 4c^{2/3} \operatorname{atanh}(\sqrt{x}cx) bx + 2 \operatorname{atanh}(\sqrt{x}cx) bx}{4c^{2/3}}$$

input

```
int(a+b*atanh(c*x^(3/2)),x)
```

output

```
(2*sqrt(3)*atan((2*sqrt(x)*c**(1/3) - 1)/sqrt(3))*b + 2*sqrt(3)*atan((2*sq
rt(x)*c**(1/3) + 1)/sqrt(3))*b + 4*c**(2/3)*atanh(sqrt(x)*c*x)*b*x + 2*ata
nh(sqrt(x)*c*x)*b + 4*c**(2/3)*a*x - 3*log(sqrt(x)*c**(2/3) + c**(1/3))*b
+ 3*log(sqrt(x)*c**(2/3) - c**(1/3))*b)/(4*c**(2/3))
```

$$3.216 \quad \int \frac{a+b \operatorname{arctanh}(cx^{3/2})}{x} dx$$

Optimal result	1738
Mathematica [A] (verified)	1738
Rubi [A] (verified)	1739
Maple [B] (verified)	1740
Fricas [F]	1740
Sympy [F(-1)]	1741
Maxima [B] (verification not implemented)	1741
Giac [F]	1741
Mupad [F(-1)]	1742
Reduce [F]	1742

Optimal result

Integrand size = 16, antiderivative size = 34

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x} dx = a \log(x) - \frac{1}{3}b \operatorname{PolyLog}(2, -cx^{3/2}) + \frac{1}{3}b \operatorname{PolyLog}(2, cx^{3/2})$$

output `a*ln(x)-1/3*b*polylog(2,-c*x^(3/2))+1/3*b*polylog(2,c*x^(3/2))`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.94

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x} dx = a \log(x) + \frac{1}{3}b(-\operatorname{PolyLog}(2, -cx^{3/2}) + \operatorname{PolyLog}(2, cx^{3/2}))$$

input `Integrate[(a + b*ArcTanh[c*x^(3/2)])/x,x]`

output `a*Log[x] + (b*(-PolyLog[2, -(c*x^(3/2))] + PolyLog[2, c*x^(3/2)]))/3`

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.24, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {6450, 6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x} dx$$

↓ 6450

$$\frac{2}{3} \int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^{3/2}} dx^{3/2}$$

↓ 6446

$$\frac{2}{3} \left(a \log(x^{3/2}) - \frac{1}{2} b \operatorname{PolyLog}\left(2, -cx^{3/2}\right) + \frac{1}{2} b \operatorname{PolyLog}\left(2, cx^{3/2}\right) \right)$$

input `Int[(a + b*ArcTanh[c*x^(3/2)])/x,x]`

output `(2*(a*Log[x^(3/2)] - (b*PolyLog[2, -(c*x^(3/2))])/2 + (b*PolyLog[2, c*x^(3/2)])/2)/3`

Defintions of rubi rules used

rule 6446 `Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))/(x_), x_Symbol] := Simp[a*Log[x], x] + (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) / ; FreeQ[{a, b, c}, x]`

rule 6450 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)]*(b_.))^(p_.)/(x_), x_Symbol] := Simp[1/n Subst[Int[(a + b*ArcTanh[c*x])^p/x, x], x, x^n], x] / ; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. 2(26) = 52.

Time = 0.44 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.68

method	result	si
parts	$a \ln(x) + b \left(\frac{2 \ln(cx^{\frac{3}{2}}) \operatorname{arctanh}(cx^{\frac{3}{2}})}{3} - \frac{\operatorname{dilog}(cx^{\frac{3}{2}})}{3} - \frac{\operatorname{dilog}(cx^{\frac{3}{2}}+1)}{3} - \frac{\ln(cx^{\frac{3}{2}}) \ln(cx^{\frac{3}{2}}+1)}{3} \right)$	57
derivativedivides	$\frac{2a \ln(cx^{\frac{3}{2}})}{3} + \frac{2b \left(\ln(cx^{\frac{3}{2}}) \operatorname{arctanh}(cx^{\frac{3}{2}}) - \frac{\operatorname{dilog}(cx^{\frac{3}{2}}+1)}{2} - \frac{\ln(cx^{\frac{3}{2}}) \ln(cx^{\frac{3}{2}}+1)}{2} - \frac{\operatorname{dilog}(cx^{\frac{3}{2}})}{2} \right)}{3}$	62
default	$\frac{2a \ln(cx^{\frac{3}{2}})}{3} + \frac{2b \left(\ln(cx^{\frac{3}{2}}) \operatorname{arctanh}(cx^{\frac{3}{2}}) - \frac{\operatorname{dilog}(cx^{\frac{3}{2}}+1)}{2} - \frac{\ln(cx^{\frac{3}{2}}) \ln(cx^{\frac{3}{2}}+1)}{2} - \frac{\operatorname{dilog}(cx^{\frac{3}{2}})}{2} \right)}{3}$	62

```
input int((a+b*arctanh(c*x^(3/2)))/x,x,method=_RETURNVERBOSE)
```

```
output a*ln(x)+b*(2/3*ln(c*x^(3/2))*arctanh(c*x^(3/2))-1/3*dilog(c*x^(3/2))-1/3*dilog(c*x^(3/2)+1)-1/3*ln(c*x^(3/2))*ln(c*x^(3/2)+1))
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x} dx = \int \frac{b \operatorname{artanh}(cx^{3/2}) + a}{x} dx$$

```
input integrate((a+b*arctanh(c*x^(3/2)))/x,x, algorithm="fricas")
```

```
output integral((b*arctanh(c*x^(3/2)) + a)/x, x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**(3/2)))/x,x)`output `Timed out`**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 62 vs. 2(24) = 48.

Time = 0.17 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.82

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x} dx = -\frac{1}{3} \left(\log \left(cx^{\frac{3}{2}} \right) \log \left(-cx^{\frac{3}{2}} + 1 \right) + \operatorname{Li}_2 \left(-cx^{\frac{3}{2}} + 1 \right) \right) b$$

$$+ \frac{1}{3} \left(\log \left(cx^{\frac{3}{2}} + 1 \right) \log \left(-cx^{\frac{3}{2}} \right) + \operatorname{Li}_2 \left(cx^{\frac{3}{2}} + 1 \right) \right) b + a \log(x)$$

input `integrate((a+b*arctanh(c*x^(3/2)))/x,x, algorithm="maxima")`output `-1/3*(log(c*x^(3/2))*log(-c*x^(3/2) + 1) + dilog(-c*x^(3/2) + 1))*b + 1/3*(log(c*x^(3/2) + 1)*log(-c*x^(3/2)) + dilog(c*x^(3/2) + 1))*b + a*log(x)`**Giac [F]**

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x} dx = \int \frac{b \operatorname{arctanh} \left(cx^{\frac{3}{2}} \right) + a}{x} dx$$

input `integrate((a+b*arctanh(c*x^(3/2)))/x,x, algorithm="giac")`output `integrate((b*arctanh(c*x^(3/2)) + a)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x} dx = \int \frac{a + b \operatorname{atanh}(cx^{3/2})}{x} dx$$

input `int((a + b*atanh(c*x^(3/2)))/x,x)`output `int((a + b*atanh(c*x^(3/2)))/x, x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x} dx = \left(\int \frac{\operatorname{atanh}(\sqrt{x} cx)}{x} dx \right) b + \log(x) a$$

input `int((a+b*atanh(c*x^(3/2)))/x,x)`output `int(atanh(sqrt(x)*c*x)/x,x)*b + log(x)*a`

3.217 $\int \frac{a+b\operatorname{arctanh}(cx^{3/2})}{x^2} dx$

Optimal result	1743
Mathematica [A] (verified)	1743
Rubi [A] (verified)	1744
Maple [A] (verified)	1748
Fricas [B] (verification not implemented)	1749
Sympy [F(-1)]	1749
Maxima [A] (verification not implemented)	1750
Giac [A] (verification not implemented)	1750
Mupad [B] (verification not implemented)	1751
Reduce [B] (verification not implemented)	1752

Optimal result

Integrand size = 16, antiderivative size = 142

$$\int \frac{a + b\operatorname{arctanh}(cx^{3/2})}{x^2} dx = -\frac{1}{2}\sqrt{3}bc^{2/3} \arctan\left(\frac{1 - 2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right) + \frac{1}{2}\sqrt{3}bc^{2/3} \arctan\left(\frac{1 + 2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right) + bc^{2/3} \operatorname{arctanh}(\sqrt[3]{c}\sqrt{x}) - \frac{a + b\operatorname{arctanh}(cx^{3/2})}{x} + \frac{1}{2}bc^{2/3} \operatorname{arctanh}\left(\frac{\sqrt[3]{c}\sqrt{x}}{1 + c^{2/3}}\right)$$

output

```
-1/2*3^(1/2)*b*c^(2/3)*arctan(1/3*(1-2*c^(1/3)*x^(1/2))*3^(1/2))+1/2*3^(1/2)*b*c^(2/3)*arctan(1/3*(1+2*c^(1/3)*x^(1/2))*3^(1/2))+b*c^(2/3)*arctanh(c^(1/3)*x^(1/2))-(a+b*arctanh(c*x^(3/2)))/x+1/2*b*c^(2/3)*arctanh(c^(1/3)*x^(1/2)/(1+c^(2/3)*x))
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.44

$$\int \frac{a + b\operatorname{arctanh}(cx^{3/2})}{x^2} dx = -\frac{a}{x} + \frac{1}{2}\sqrt{3}bc^{2/3} \arctan\left(\frac{-1 + 2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right) + \frac{1}{2}\sqrt{3}bc^{2/3} \arctan\left(\frac{1 + 2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right) - \frac{b\operatorname{arctanh}(cx^{3/2})}{x} - \frac{1}{2}bc^{2/3} \log(1 - \sqrt[3]{c}\sqrt{x}) + \frac{1}{2}bc^{2/3} \log(1 + \sqrt[3]{c}\sqrt{x}) - \frac{1}{4}b$$

input `Integrate[(a + b*ArcTanh[c*x^(3/2)])/x^2,x]`

output
$$-(a/x) + (\text{Sqrt}[3]*b*c^{(2/3)}*\text{ArcTan}[(-1 + 2*c^{(1/3)}*\text{Sqrt}[x])/ \text{Sqrt}[3]])/2 + (\text{Sqrt}[3]*b*c^{(2/3)}*\text{ArcTan}[(1 + 2*c^{(1/3)}*\text{Sqrt}[x])/ \text{Sqrt}[3]])/2 - (b*\text{ArcTanh}[c*x^{(3/2)}])/x - (b*c^{(2/3)}*\text{Log}[1 - c^{(1/3)}*\text{Sqrt}[x]])/2 + (b*c^{(2/3)}*\text{Log}[1 + c^{(1/3)}*\text{Sqrt}[x]])/2 - (b*c^{(2/3)}*\text{Log}[1 - c^{(1/3)}*\text{Sqrt}[x] + c^{(2/3)}*x])/4 + (b*c^{(2/3)}*\text{Log}[1 + c^{(1/3)}*\text{Sqrt}[x] + c^{(2/3)}*x])/4$$

Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.27, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.688$, Rules used = {6452, 851, 754, 27, 219, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^2} dx$$

$$\downarrow 6452$$

$$\frac{3}{2}bc \int \frac{1}{\sqrt{x}(1 - c^2x^3)} dx - \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x}$$

$$\downarrow 851$$

$$3bc \int \frac{1}{1 - c^2x^3} d\sqrt{x} - \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x}$$

$$\downarrow 754$$

$$3bc \left(\frac{1}{3} \int \frac{1}{1 - c^{2/3}x} d\sqrt{x} + \frac{1}{3} \int \frac{2 - \sqrt[3]{c}\sqrt{x}}{2(c^{2/3}x - \sqrt[3]{c}\sqrt{x} + 1)} d\sqrt{x} + \frac{1}{3} \int \frac{\sqrt[3]{c}\sqrt{x} + 2}{2(c^{2/3}x + \sqrt[3]{c}\sqrt{x} + 1)} d\sqrt{x} \right) - \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x}$$

$$\downarrow 27$$

$$3bc \left(\frac{1}{3} \int \frac{1}{1 - c^{2/3}x} d\sqrt{x} + \frac{1}{6} \int \frac{2 - \sqrt[3]{c}\sqrt{x}}{c^{2/3}x - \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} + \frac{1}{6} \int \frac{\sqrt[3]{c}\sqrt{x} + 2}{c^{2/3}x + \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} \right) -$$

$$\frac{x}{a + \operatorname{barctanh}(cx^{3/2})}$$

↓ 219

$$3bc \left(\frac{1}{6} \int \frac{2 - \sqrt[3]{c}\sqrt{x}}{c^{2/3}x - \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} + \frac{1}{6} \int \frac{\sqrt[3]{c}\sqrt{x} + 2}{c^{2/3}x + \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} + \frac{\operatorname{arctanh}(\sqrt[3]{c}\sqrt{x})}{3\sqrt[3]{c}} \right) -$$

$$\frac{x}{a + \operatorname{barctanh}(cx^{3/2})}$$

↓ 1142

$$3bc \left(\frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x - \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} - \frac{\int -\frac{\sqrt[3]{c}(1-2\sqrt[3]{c}\sqrt{x})}{c^{2/3}x - \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x}}{2\sqrt[3]{c}} \right) + \frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x + \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} + \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{c}\sqrt{x} - 1)}{c^{2/3}x + \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x}}{2\sqrt[3]{c}} \right) \right)$$

$$\frac{x}{a + \operatorname{barctanh}(cx^{3/2})}$$

↓ 25

$$3bc \left(\frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x - \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} + \frac{\int \frac{\sqrt[3]{c}(1-2\sqrt[3]{c}\sqrt{x})}{c^{2/3}x - \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x}}{2\sqrt[3]{c}} \right) + \frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x + \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} + \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{c}\sqrt{x} - 1)}{c^{2/3}x + \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x}}{2\sqrt[3]{c}} \right) \right)$$

$$\frac{x}{a + \operatorname{barctanh}(cx^{3/2})}$$

↓ 27

$$3bc \left(\frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x - \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} + \frac{1}{2} \int \frac{1 - 2\sqrt[3]{c}\sqrt{x}}{c^{2/3}x - \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} \right) + \frac{1}{6} \left(\frac{3}{2} \int \frac{1}{c^{2/3}x + \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} + \frac{1}{2} \int \frac{2\sqrt[3]{c}\sqrt{x} - 1}{c^{2/3}x + \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} \right) \right)$$

$$\frac{x}{a + \operatorname{barctanh}(cx^{3/2})}$$

↓ 1082

$$3bc \left(\frac{1}{6} \left(\frac{1}{2} \int \frac{1 - 2\sqrt[3]{c}\sqrt{x}}{c^{2/3}x - \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} + \frac{3 \int \frac{1}{-x-3} d(1 - 2\sqrt[3]{c}\sqrt{x})}{\sqrt[3]{c}} \right) + \frac{1}{6} \left(\frac{1}{2} \int \frac{2\sqrt[3]{c}\sqrt{x} + 1}{c^{2/3}x + \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} - \frac{3 \int \frac{1}{-x-3}}{1} \right) \right)$$

$$\frac{x}{a + \operatorname{barctanh}(cx^{3/2})}$$

x

↓ 217

$$3bc \left(\frac{1}{6} \left(\frac{1}{2} \int \frac{1 - 2\sqrt[3]{c}\sqrt{x}}{c^{2/3}x - \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} - \frac{\sqrt{3} \arctan\left(\frac{1-2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{\sqrt[3]{c}} \right) + \frac{1}{6} \left(\frac{1}{2} \int \frac{2\sqrt[3]{c}\sqrt{x} + 1}{c^{2/3}x + \sqrt[3]{c}\sqrt{x} + 1} d\sqrt{x} + \frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{c}\sqrt{x} + 1}{\sqrt{3}}\right)}{\sqrt[3]{c}} \right) \right) \frac{a + \operatorname{barctanh}(cx^{3/2})}{x}$$

↓ 1103

$$3bc \left(\frac{1}{6} \left(-\frac{\sqrt{3} \arctan\left(\frac{1-2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{\log(c^{2/3}x - \sqrt[3]{c}\sqrt{x} + 1)}{2\sqrt[3]{c}} \right) + \frac{1}{6} \left(\frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{c}\sqrt{x} + 1}{\sqrt{3}}\right)}{\sqrt[3]{c}} + \frac{\log(c^{2/3}x + \sqrt[3]{c}\sqrt{x} + 1)}{2\sqrt[3]{c}} \right) \right) \frac{a + \operatorname{barctanh}(cx^{3/2})}{x}$$

input `Int[(a + b*ArcTanh[c*x^(3/2)])/x^2,x]`

output `-((a + b*ArcTanh[c*x^(3/2)])/x) + 3*b*c*(ArcTanh[c^(1/3)*Sqrt[x]]/(3*c^(1/3))) + (-((Sqrt[3]*ArcTan[(1 - 2*c^(1/3)*Sqrt[x])/Sqrt[3]])/c^(1/3)) - Log[1 - c^(1/3)*Sqrt[x] + c^(2/3)*x]/(2*c^(1/3)))/6 + ((Sqrt[3]*ArcTan[(1 + 2*c^(1/3)*Sqrt[x])/Sqrt[3]])/c^(1/3) + Log[1 + c^(1/3)*Sqrt[x] + c^(2/3)*x]/(2*c^(1/3)))/6`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 219 $\text{Int}[(a_ + (b_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[-b, 2])) \cdot \text{ArcTanh}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 754 $\text{Int}[(a_ + (b_ \cdot)(x_)^n)^{-1}, x_Symbol] \rightarrow \text{Module}\{r = \text{Numerator}[\text{Rt}[-a/b, n]], s = \text{Denominator}[\text{Rt}[-a/b, n]], k, u\}, \text{Simp}[u = \text{Int}[(r - s \cdot \text{Cos}[(2 \cdot k \cdot \text{Pi})/n] \cdot x)/(r^2 - 2 \cdot r \cdot s \cdot \text{Cos}[(2 \cdot k \cdot \text{Pi})/n] \cdot x + s^2 \cdot x^2), x] + \text{Int}[(r + s \cdot \text{Cos}[(2 \cdot k \cdot \text{Pi})/n] \cdot x)/(r^2 + 2 \cdot r \cdot s \cdot \text{Cos}[(2 \cdot k \cdot \text{Pi})/n] \cdot x + s^2 \cdot x^2), x]; 2 \cdot (r^2/(a \cdot n)) \ \text{Int}[1/(r^2 - s^2 \cdot x^2), x] + 2 \cdot (r/(a \cdot n)) \ \text{Sum}[u, \{k, 1, (n - 2)/4\}], x] /; \text{FreeQ}\{a, b\}, x\} \ \&\& \ \text{IGtQ}[(n - 2)/4, 0] \ \&\& \ \text{NegQ}[a/b]$

rule 851 $\text{Int}[(c_ \cdot)(x_)^m \cdot (a_ + (b_ \cdot)(x_)^n)^p, x_Symbol] \rightarrow \text{With}\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \ \text{Subst}[\text{Int}[x^{(k \cdot (m + 1) - 1)} \cdot (a + b \cdot x^{k \cdot n})/c^n]^p, x], x, (c \cdot x)^{1/k}], x] /; \text{FreeQ}\{a, b, c, p\}, x\} \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 1082 $\text{Int}[(a_ + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}\{q = 1 - 4 \cdot S\text{implify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4 \cdot a \cdot c]) /; \text{FreeQ}\{a, b, c\}, x]$

rule 1103 $\text{Int}[(d_ + (e_ \cdot)(x_))/((a_ + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2), x_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /; \text{FreeQ}\{a, b, c, d, e\}, x\} \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$

rule 1142 $\text{Int}[(d_ + (e_ \cdot)(x_))/((a_ + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2 \cdot c \cdot d - b \cdot e)/(2 \cdot c) \ \text{Int}[1/(a + b \cdot x + c \cdot x^2), x], x] + \text{Simp}[e/(2 \cdot c) \ \text{Int}[(b + 2 \cdot c \cdot x)/(a + b \cdot x + c \cdot x^2), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x]$

rule 6452 $\text{Int}[(a_ + \text{ArcTanh}[c_ \cdot (x_)^n] \cdot (b_ \cdot)^{p_} \cdot (x_)^{m_}), x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^p/(m+1), x] - \text{Simp}[b \cdot c \cdot n \cdot (p/(m+1)) \ \text{Int}[x^{m+n} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1}/(1 - c^2 \cdot x^{2 \cdot n})], x] /; \text{FreeQ}\{a, b, c, m, n\}, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.18

method	result
derivativedivides	$-\frac{a}{x} - \frac{b \operatorname{arctanh}\left(cx^{\frac{3}{2}}\right)}{x} - \frac{b \ln\left(\sqrt{x} - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{2\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{b \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{4\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2\sqrt{x}}{\left(\frac{1}{c}\right)^{\frac{1}{3}}} + 1\right)}{3}\right)}{2\left(\frac{1}{c}\right)^{\frac{2}{3}}}$
default	$-\frac{a}{x} - \frac{b \operatorname{arctanh}\left(cx^{\frac{3}{2}}\right)}{x} - \frac{b \ln\left(\sqrt{x} - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{2\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{b \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{4\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2\sqrt{x}}{\left(\frac{1}{c}\right)^{\frac{1}{3}}} + 1\right)}{3}\right)}{2\left(\frac{1}{c}\right)^{\frac{2}{3}}}$
parts	$-\frac{a}{x} - \frac{b \operatorname{arctanh}\left(cx^{\frac{3}{2}}\right)}{x} - \frac{b \ln\left(\sqrt{x} - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{2\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{b \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{4\left(\frac{1}{c}\right)^{\frac{2}{3}}} + \frac{b\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2\sqrt{x}}{\left(\frac{1}{c}\right)^{\frac{1}{3}}} + 1\right)}{3}\right)}{2\left(\frac{1}{c}\right)^{\frac{2}{3}}}$

input `int((a+b*arctanh(c*x^(3/2)))/x^2,x,method=_RETURNVERBOSE)`

output `-a/x-b/x*arctanh(c*x^(3/2))-1/2*b/(1/c)^(2/3)*ln(x^(1/2)-(1/c)^(1/3))+1/4*b/(1/c)^(2/3)*ln(x+(1/c)^(1/3)*x^(1/2)+(1/c)^(2/3))+1/2*b/(1/c)^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x^(1/2)+1))+1/2*b/(1/c)^(2/3)*ln(x^(1/2)+(1/c)^(1/3))-1/4*b/(1/c)^(2/3)*ln(x-(1/c)^(1/3)*x^(1/2)+(1/c)^(2/3))+1/2*b/(1/c)^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x^(1/2)-1))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 234 vs. $2(102) = 204$.

Time = 0.10 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.65

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^2} dx =$$

$$\frac{2\sqrt{3}(-c^2)^{\frac{1}{3}} bx \arctan\left(\frac{2\sqrt{3}(-c^2)^{\frac{2}{3}}\sqrt{x} + \sqrt{3}c}{3c}\right) - 2\sqrt{3}b(c^2)^{\frac{1}{3}}x \arctan\left(\frac{2\sqrt{3}(c^2)^{\frac{2}{3}}\sqrt{x} - \sqrt{3}c}{3c}\right) + (-c^2)^{\frac{1}{3}} bx \log\left(c^2x - (-c^2)^{\frac{1}{3}}c\sqrt{x} + (-c^2)^{\frac{2}{3}}\right) + b(c^2)^{\frac{1}{3}}x \log\left(c^2x - (-c^2)^{\frac{1}{3}}c\sqrt{x} + (-c^2)^{\frac{2}{3}}\right) - 2(-c^2)^{\frac{1}{3}}b \log\left(c\sqrt{x} + (-c^2)^{\frac{1}{3}}\right) - 2b \log\left(-c^2x^3 + 2cx^{\frac{3}{2}} + 1\right)}{c^2x^3 - 1} + \frac{4a}{x}$$

input `integrate((a+b*arctanh(c*x^(3/2)))/x^2,x, algorithm="fricas")`

output `-1/4*(2*sqrt(3)*(-c^2)^(1/3)*b*x*arctan(1/3*(2*sqrt(3)*(-c^2)^(2/3)*sqrt(x) + sqrt(3)*c)/c) - 2*sqrt(3)*b*(c^2)^(1/3)*x*arctan(1/3*(2*sqrt(3)*(c^2)^(2/3)*sqrt(x) - sqrt(3)*c)/c) + (-c^2)^(1/3)*b*x*log(c^2*x - (-c^2)^(1/3)*c*sqrt(x) + (-c^2)^(2/3)) + b*(c^2)^(1/3)*x*log(c^2*x - (c^2)^(1/3)*c*sqrt(x) + (c^2)^(2/3)) - 2*(-c^2)^(1/3)*b*x*log(c*sqrt(x) + (-c^2)^(1/3)) - 2*b*(c^2)^(1/3)*x*log(c*sqrt(x) + (c^2)^(1/3)) + 2*b*log(-c^2*x^3 + 2*c*x^(3/2) + 1)/(c^2*x^3 - 1) + 4*a)/x`

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^2} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**(3/2)))/x**2,x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.15

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^2} dx = \frac{1}{4} \left(\left(\frac{2\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}(2c^{2/3}\sqrt{x} + c^{1/3})}{3c^{1/3}}\right)}{c^{1/3}} + \frac{2\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}(2c^{2/3}\sqrt{x} - c^{1/3})}{3c^{1/3}}\right)}{c^{1/3}} + \frac{\log\left(\frac{c^{1/3} + 2\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}(2c^{2/3}\sqrt{x} + c^{1/3})}{3c^{1/3}}\right)}{c^{1/3}}\right)}{c^{1/3}} \right) \right. \\ \left. - \frac{a}{x} \right)$$

input `integrate((a+b*arctanh(c*x^(3/2)))/x^2,x, algorithm="maxima")`

output `1/4*((2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*sqrt(x) + c^(1/3))/c^(1/3))/c^(1/3) + 2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*sqrt(x) - c^(1/3))/c^(1/3))/c^(1/3) + log(c^(2/3)*x + c^(1/3)*sqrt(x) + 1)/c^(1/3) - log(c^(2/3)*x - c^(1/3)*sqrt(x) + 1)/c^(1/3) + 2*log((c^(1/3)*sqrt(x) + 1)/c^(1/3))/c^(1/3) - 2*log((c^(1/3)*sqrt(x) - 1)/c^(1/3))/c^(1/3))*c - 4*arctanh(c*x^(3/2))/x)*b - a/x`

Giac [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.21

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^2} dx = \frac{1}{4} \left(\frac{2\sqrt{3} \operatorname{arctan}\left(\frac{\frac{1}{3}\sqrt{3}\left(2\sqrt{x} + \frac{1}{|c|^{1/3}}\right)|c|^{1/3}}{|c|^{1/3}}\right)}{|c|^{1/3}} + \frac{2\sqrt{3} \operatorname{arctan}\left(\frac{\frac{1}{3}\sqrt{3}\left(2\sqrt{x} - \frac{1}{|c|^{1/3}}\right)|c|^{1/3}}{|c|^{1/3}}\right)}{|c|^{1/3}} + \frac{b \log\left(\frac{-cx^{3/2} + 1}{cx^{3/2} - 1}\right)}{2x} - \frac{a}{x} \right)$$

input `integrate((a+b*arctanh(c*x^(3/2)))/x^2,x, algorithm="giac")`

output

```
1/4*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*sqrt(x) + 1/abs(c)^(1/3))*abs(c)^(1/3
))/abs(c)^(1/3) + 2*sqrt(3)*arctan(1/3*sqrt(3)*(2*sqrt(x) - 1/abs(c)^(1/3)
))*abs(c)^(1/3)/abs(c)^(1/3) + log(x + sqrt(x)/abs(c)^(1/3) + 1/abs(c)^(2/
3))/abs(c)^(1/3) - log(x - sqrt(x)/abs(c)^(1/3) + 1/abs(c)^(2/3))/abs(c)^(
1/3) + 2*log(sqrt(x) + 1/abs(c)^(1/3))/abs(c)^(1/3) - 2*log(abs(sqrt(x) -
1/abs(c)^(1/3)))/abs(c)^(1/3))*b*c - 1/2*b*log(-(c*x^(3/2) + 1)/(c*x^(3/2)
- 1))/x - a/x
```

Mupad [B] (verification not implemented)

Time = 10.40 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.55

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^2} dx = \frac{b c^{2/3} \ln\left(\frac{c^{1/3} \sqrt{x} + 1}{c^{1/3} \sqrt{x} - 1}\right)}{2} - \frac{a}{x}$$

$$+ \frac{\ln(1 - cx^{3/2})(bx - bc^2 x^4)}{2x^2 - 2c^2 x^5} - \frac{b \ln(cx^{3/2} + 1)}{2x}$$

$$+ \frac{b c^{2/3} \ln\left(\frac{\sqrt{3} + c^{2/3} x \operatorname{li} - c^{1/3} \sqrt{x} 4i - \sqrt{3} c^{2/3} x + \operatorname{li}}{2 c^{2/3} x + 1 + \sqrt{3} \operatorname{li}}\right)}{2} \sqrt{\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}} \operatorname{li}$$

$$+ \frac{b c^{2/3} \ln\left(\frac{\sqrt{3} c^{2/3} x + c^{2/3} x \operatorname{li} + c^{1/3} \sqrt{x} 4i - \sqrt{3} + \operatorname{li}}{2 c^{2/3} x + 1 - \sqrt{3} \operatorname{li}}\right)}{2} \sqrt{-\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}}$$

input

```
int((a + b*atanh(c*x^(3/2)))/x^2,x)
```

output

```
(b*c^(2/3)*log((c^(1/3)*x^(1/2) + 1)/(c^(1/3)*x^(1/2) - 1)))/2 - a/x + (lo
g(1 - c*x^(3/2))*(b*x - b*c^2*x^4))/(2*x^2 - 2*c^2*x^5) - (b*log(c*x^(3/2)
+ 1))/(2*x) + (b*c^(2/3)*log((3^(1/2) + c^(2/3)*x*1i - c^(1/3)*x^(1/2)*4i
- 3^(1/2)*c^(2/3)*x + 1i)/(3^(1/2)*1i + 2*c^(2/3)*x + 1))*((3^(1/2)*1i)/2
+ 1/2)^(1/2)*1i)/2 + (b*c^(2/3)*log((c^(2/3)*x*1i - 3^(1/2) + c^(1/3)*x^(
1/2)*4i + 3^(1/2)*c^(2/3)*x + 1i)/(2*c^(2/3)*x - 3^(1/2)*1i + 1))*((3^(1/2)
)*1i)/2 - 1/2)^(1/2))/2
```

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.82

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^2} dx = \frac{2\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{x}c^{1/3}-1}{\sqrt{3}}\right) b cx + 2\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{x}c^{1/3}+1}{\sqrt{3}}\right) b cx - 4c^{1/3} \operatorname{atanh}(\sqrt{x} cx) b}{4}$$

input `int((a+b*atanh(c*x^(3/2)))/x^2,x)`output `(2*sqrt(3)*atan((2*sqrt(x)*c**(1/3) - 1)/sqrt(3))*b*c*x + 2*sqrt(3)*atan((2*sqrt(x)*c**(1/3) + 1)/sqrt(3))*b*c*x - 4*c**(1/3)*atanh(sqrt(x)*c*x)*b - 2*atanh(sqrt(x)*c*x)*b*c*x - 4*c**(1/3)*a + 3*log(sqrt(x)*c**(2/3) + c**(1/3))*b*c*x - 3*log(sqrt(x)*c**(2/3) - c**(1/3))*b*c*x)/(4*c**(1/3)*x)`

3.218 $\int \frac{a+b\operatorname{arctanh}(cx^{3/2})}{x^3} dx$

Optimal result	1753
Mathematica [A] (verified)	1753
Rubi [A] (verified)	1754
Maple [A] (verified)	1759
Fricas [A] (verification not implemented)	1760
Sympy [F(-1)]	1760
Maxima [A] (verification not implemented)	1761
Giac [A] (verification not implemented)	1761
Mupad [B] (verification not implemented)	1762
Reduce [B] (verification not implemented)	1763

Optimal result

Integrand size = 16, antiderivative size = 158

$$\int \frac{a + b\operatorname{arctanh}(cx^{3/2})}{x^3} dx = -\frac{3bc}{2\sqrt{x}} + \frac{1}{4}\sqrt{3}bc^{4/3} \arctan\left(\frac{1 - 2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right) - \frac{1}{4}\sqrt{3}bc^{4/3} \arctan\left(\frac{1 + 2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right) + \frac{1}{2}bc^{4/3}\operatorname{arctanh}(\sqrt[3]{c}\sqrt{x}) - \frac{a + b\operatorname{arctanh}(cx^{3/2})}{2x^2} + \frac{1}{4}bc^{4/3}\operatorname{arctanh}\left(\frac{\sqrt[3]{c}}{1 + c}\right)$$

output

```
-3/2*b*c/x^(1/2)+1/4*3^(1/2)*b*c^(4/3)*arctan(1/3*(1-2*c^(1/3)*x^(1/2))*3^(1/2))-1/4*3^(1/2)*b*c^(4/3)*arctan(1/3*(1+2*c^(1/3)*x^(1/2))*3^(1/2))+1/2*b*c^(4/3)*arctanh(c^(1/3)*x^(1/2))-1/2*(a+b*arctanh(c*x^(3/2)))/x^2+1/4*b*c^(4/3)*arctanh(c^(1/3)*x^(1/2)/(1+c^(2/3)*x))
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.39

$$\int \frac{a + b\operatorname{arctanh}(cx^{3/2})}{x^3} dx = -\frac{a}{2x^2} - \frac{3bc}{2\sqrt{x}} - \frac{1}{4}\sqrt{3}bc^{4/3} \arctan\left(\frac{-1 + 2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right) - \frac{1}{4}\sqrt{3}bc^{4/3} \arctan\left(\frac{1 + 2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right) - \frac{b\operatorname{arctanh}(cx^{3/2})}{2x^2} - \frac{1}{4}bc^{4/3} \log(1 - \sqrt[3]{c}\sqrt{x}) + \frac{1}{4}bc^{4/3} \log(1 + \sqrt[3]{c}\sqrt{x}) - \frac{1}{8}$$

input `Integrate[(a + b*ArcTanh[c*x^(3/2)])/x^3,x]`

output
$$-1/2*a/x^2 - (3*b*c)/(2*\text{Sqrt}[x]) - (\text{Sqrt}[3]*b*c^{(4/3)}*\text{ArcTan}[(-1 + 2*c^{(1/3)}*\text{Sqrt}[x])/ \text{Sqrt}[3]])/4 - (\text{Sqrt}[3]*b*c^{(4/3)}*\text{ArcTan}[(1 + 2*c^{(1/3)}*\text{Sqrt}[x])/ \text{Sqrt}[3]])/4 - (b*\text{ArcTanh}[c*x^{(3/2)}])/(2*x^2) - (b*c^{(4/3)}*\text{Log}[1 - c^{(1/3)}*\text{Sqrt}[x]])/4 + (b*c^{(4/3)}*\text{Log}[1 + c^{(1/3)}*\text{Sqrt}[x]])/4 - (b*c^{(4/3)}*\text{Log}[1 - c^{(1/3)}*\text{Sqrt}[x] + c^{(2/3)}*x])/8 + (b*c^{(4/3)}*\text{Log}[1 + c^{(1/3)}*\text{Sqrt}[x] + c^{(2/3)}*x])/8$$

Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.31, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6452, 847, 851, 825, 27, 219, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^3} dx \\ & \quad \downarrow \text{6452} \\ & \frac{3}{4}bc \int \frac{1}{x^{3/2}(1-c^2x^3)} dx - \frac{a + b \operatorname{arctanh}(cx^{3/2})}{2x^2} \\ & \quad \downarrow \text{847} \\ & \frac{3}{4}bc \left(c^2 \int \frac{x^{3/2}}{1-c^2x^3} dx - \frac{2}{\sqrt{x}} \right) - \frac{a + b \operatorname{arctanh}(cx^{3/2})}{2x^2} \\ & \quad \downarrow \text{851} \\ & \frac{3}{4}bc \left(2c^2 \int \frac{x^2}{1-c^2x^3} d\sqrt{x} - \frac{2}{\sqrt{x}} \right) - \frac{a + b \operatorname{arctanh}(cx^{3/2})}{2x^2} \\ & \quad \downarrow \text{825} \end{aligned}$$

$$\frac{3}{4}bc \left(2c^2 \left(\frac{\int \frac{1}{1-c^{2/3}x} d\sqrt{x}}{3c^{4/3}} + \frac{\int -\frac{\sqrt[3]{c\sqrt{x}+1}}{2(c^{2/3}x-\sqrt[3]{c\sqrt{x}+1})} d\sqrt{x}}{3c^{4/3}} + \frac{\int -\frac{1-\sqrt[3]{c\sqrt{x}}}{2(c^{2/3}x+\sqrt[3]{c\sqrt{x}+1})} d\sqrt{x}}{3c^{4/3}} \right) - \frac{2}{\sqrt{x}} \right) -$$

$$\frac{a + \operatorname{barctanh}(cx^{3/2})}{2x^2}$$

↓ 27

$$\frac{3}{4}bc \left(2c^2 \left(\frac{\int \frac{1}{1-c^{2/3}x} d\sqrt{x}}{3c^{4/3}} - \frac{\int \frac{\sqrt[3]{c\sqrt{x}+1}}{c^{2/3}x-\sqrt[3]{c\sqrt{x}+1}} d\sqrt{x}}{6c^{4/3}} - \frac{\int \frac{1-\sqrt[3]{c\sqrt{x}}}{c^{2/3}x+\sqrt[3]{c\sqrt{x}+1}} d\sqrt{x}}{6c^{4/3}} \right) - \frac{2}{\sqrt{x}} \right) -$$

$$\frac{a + \operatorname{barctanh}(cx^{3/2})}{2x^2}$$

↓ 219

$$\frac{3}{4}bc \left(2c^2 \left(-\frac{\int \frac{\sqrt[3]{c\sqrt{x}+1}}{c^{2/3}x-\sqrt[3]{c\sqrt{x}+1}} d\sqrt{x}}{6c^{4/3}} - \frac{\int \frac{1-\sqrt[3]{c\sqrt{x}}}{c^{2/3}x+\sqrt[3]{c\sqrt{x}+1}} d\sqrt{x}}{6c^{4/3}} + \frac{\operatorname{arctanh}(\sqrt[3]{c\sqrt{x}})}{3c^{5/3}} \right) - \frac{2}{\sqrt{x}} \right) -$$

$$\frac{a + \operatorname{barctanh}(cx^{3/2})}{2x^2}$$

↓ 1142

$$\frac{3}{4}bc \left(2c^2 \left(-\frac{\frac{3}{2} \int \frac{1}{c^{2/3}x-\sqrt[3]{c\sqrt{x}+1}} d\sqrt{x} + \frac{\int -\frac{\sqrt[3]{c}(1-2\sqrt[3]{c\sqrt{x}})}{c^{2/3}x-\sqrt[3]{c\sqrt{x}+1}} d\sqrt{x}}{2\sqrt[3]{c}}}{6c^{4/3}} - \frac{\frac{3}{2} \int \frac{1}{c^{2/3}x+\sqrt[3]{c\sqrt{x}+1}} d\sqrt{x} - \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{c\sqrt{x}+1})}{c^{2/3}x+\sqrt[3]{c\sqrt{x}+1}} d\sqrt{x}}{2\sqrt[3]{c}}}{6c^{4/3}} + \right)$$

$$\frac{a + \operatorname{barctanh}(cx^{3/2})}{2x^2}$$

↓ 25

$$\frac{3}{4}bc \left(2c^2 \left(-\frac{\frac{3}{2} \int \frac{1}{c^{2/3}x - \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} - \frac{\int \frac{\sqrt[3]{c}(1-2\sqrt[3]{c}\sqrt{x})}{c^{2/3}x - \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x}}{2\sqrt[3]{c}}}{6c^{4/3}} - \frac{\frac{3}{2} \int \frac{1}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} - \frac{\int \frac{\sqrt[3]{c}(2\sqrt[3]{c}\sqrt{x+1})}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x}}{2\sqrt[3]{c}}}{6c^{4/3}} + a \right) \right)$$

$$\frac{a + \operatorname{barctanh}(cx^{3/2})}{2x^2}$$

↓ 27

$$\frac{3}{4}bc \left(2c^2 \left(-\frac{\frac{3}{2} \int \frac{1}{c^{2/3}x - \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} - \frac{1}{2} \int \frac{1-2\sqrt[3]{c}\sqrt{x}}{c^{2/3}x - \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x}}{6c^{4/3}} - \frac{\frac{3}{2} \int \frac{1}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} - \frac{1}{2} \int \frac{2\sqrt[3]{c}\sqrt{x+1}}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x}}{6c^{4/3}} + a \right) \right)$$

$$\frac{a + \operatorname{barctanh}(cx^{3/2})}{2x^2}$$

↓ 1082

$$\frac{3}{4}bc \left(2c^2 \left(-\frac{\frac{3 \int \frac{1}{-x-3} d(1-2\sqrt[3]{c}\sqrt{x})}{\sqrt[3]{c}} - \frac{1}{2} \int \frac{1-2\sqrt[3]{c}\sqrt{x}}{c^{2/3}x - \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x}}{6c^{4/3}} - \frac{-\frac{1}{2} \int \frac{2\sqrt[3]{c}\sqrt{x+1}}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} - \frac{3 \int \frac{1}{-x-3} d(2\sqrt[3]{c}\sqrt{x+1})}{\sqrt[3]{c}}}{6c^{4/3}} + a \right) \right)$$

$$\frac{a + \operatorname{barctanh}(cx^{3/2})}{2x^2}$$

↓ 217

$$\frac{3}{4}bc \left(2c^2 \left(-\frac{-\frac{1}{2} \int \frac{1-2\sqrt[3]{c}\sqrt{x}}{c^{2/3}x - \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x} - \frac{\sqrt{3} \arctan\left(\frac{1-2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{\sqrt[3]{c}}}{6c^{4/3}} - \frac{\frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{c}\sqrt{x+1}}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{1}{2} \int \frac{2\sqrt[3]{c}\sqrt{x+1}}{c^{2/3}x + \sqrt[3]{c}\sqrt{x+1}} d\sqrt{x}}{6c^{4/3}} + a \right) \right)$$

$$\frac{a + \operatorname{barctanh}(cx^{3/2})}{2x^2}$$

↓ 1103

$$\frac{3}{4}bc \left(2c^2 \left(-\frac{\log(c^{2/3}x - \sqrt[3]{c}\sqrt{x+1})}{2\sqrt[3]{c}} - \frac{\sqrt{3}\arctan\left(\frac{1-2\sqrt[3]{c}\sqrt{x}}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{\sqrt{3}\arctan\left(\frac{2\sqrt[3]{c}\sqrt{x+1}}{\sqrt{3}}\right)}{\sqrt[3]{c}} - \frac{\log(c^{2/3}x + \sqrt[3]{c}\sqrt{x+1})}{2\sqrt[3]{c}} \right) + \frac{\operatorname{arctanh}\left(\frac{a + b\operatorname{arctanh}(cx^{3/2})}{2x^2}\right)}{3c^5} \right)$$

input `Int[(a + b*ArcTanh[c*x^(3/2)])/x^3,x]`

output `-1/2*(a + b*ArcTanh[c*x^(3/2)])/x^2 + (3*b*c*(-2/Sqrt[x] + 2*c^2*(ArcTanh[c^(1/3)*Sqrt[x]]/(3*c^(5/3)) - ((Sqrt[3]*ArcTan[(1 - 2*c^(1/3)*Sqrt[x]]/Sqrt[3])/c^(1/3)) + Log[1 - c^(1/3)*Sqrt[x] + c^(2/3)*x]/(2*c^(1/3)))/(6*c^(4/3)) - ((Sqrt[3]*ArcTan[(1 + 2*c^(1/3)*Sqrt[x]]/Sqrt[3])/c^(1/3) - Log[1 + c^(1/3)*Sqrt[x] + c^(2/3)*x]/(2*c^(1/3)))/(6*c^(4/3))))/4`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 825 `Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Module[{r = Numerator[Rt[-a/b, n]], s = Denominator[Rt[-a/b, n]], k, u}, Simp[u = Int[(r*Cos[2*k*m*(Pi/n)] - s*Cos[2*k*(m + 1)*(Pi/n)]*x)/(r^2 - 2*r*s*Cos[2*k*(Pi/n)]*x + s^2*x^2), x] + Int[(r*Cos[2*k*m*(Pi/n)] + s*Cos[2*k*(m + 1)*(Pi/n)]*x)/(r^2 + 2*r*s*Cos[2*k*(Pi/n)]*x + s^2*x^2), x]; 2*(r^(m + 2)/(a*n*s^m)) Int[1/(r^2 - s^2*x^2), x] + 2*(r^(m + 1)/(a*n*s^m)) Sum[u, {k, 1, (n - 2)/4}], x] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && NegQ[a/b]`

rule 847 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + n*(p + 1) + 1)/(a*c^n*(m + 1)) Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 851 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n)]^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.14

method	result
derivativedivides	$-\frac{a}{2x^2} - \frac{b \operatorname{arctanh}\left(cx^{\frac{3}{2}}\right)}{2x^2} - \frac{bc \ln\left(\sqrt{x} - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{bc \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{8\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{bc\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2\sqrt{x}}{\left(\frac{1}{c}\right)^{\frac{1}{3}}} + 1\right)}{3}\right)}{4\left(\frac{1}{c}\right)^{\frac{1}{3}}}$
default	$-\frac{a}{2x^2} - \frac{b \operatorname{arctanh}\left(cx^{\frac{3}{2}}\right)}{2x^2} - \frac{bc \ln\left(\sqrt{x} - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{bc \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{8\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{bc\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2\sqrt{x}}{\left(\frac{1}{c}\right)^{\frac{1}{3}}} + 1\right)}{3}\right)}{4\left(\frac{1}{c}\right)^{\frac{1}{3}}}$
parts	$-\frac{a}{2x^2} - \frac{b \operatorname{arctanh}\left(cx^{\frac{3}{2}}\right)}{2x^2} - \frac{bc \ln\left(\sqrt{x} - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{4\left(\frac{1}{c}\right)^{\frac{1}{3}}} + \frac{bc \ln\left(x + \left(\frac{1}{c}\right)^{\frac{1}{3}}\sqrt{x} + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{8\left(\frac{1}{c}\right)^{\frac{1}{3}}} - \frac{bc\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{2\sqrt{x}}{\left(\frac{1}{c}\right)^{\frac{1}{3}}} + 1\right)}{3}\right)}{4\left(\frac{1}{c}\right)^{\frac{1}{3}}}$

input

```
int((a+b*arctanh(c*x^(3/2)))/x^3,x,method=_RETURNVERBOSE)
```

output

```
-1/2*a/x^2-1/2*b/x^2*arctanh(c*x^(3/2))-1/4*b*c/(1/c)^(1/3)*ln(x^(1/2)-(1/c)^(1/3))+1/8*b*c/(1/c)^(1/3)*ln(x+(1/c)^(1/3)*x^(1/2)+(1/c)^(2/3))-1/4*b*c*3^(1/2)/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x^(1/2)+1))+1/4*b*c/(1/c)^(1/3)*ln(x^(1/2)+(1/c)^(1/3))-1/8*b*c/(1/c)^(1/3)*ln(x-(1/c)^(1/3)*x^(1/2)+(1/c)^(2/3))-1/4*b*c*3^(1/2)/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x^(1/2)-1))-3/2*b*c/x^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.35

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^3} dx =$$

$$\frac{2\sqrt{3}b(-c)^{\frac{1}{3}}cx^2 \arctan\left(\frac{2}{3}\sqrt{3}(-c)^{\frac{1}{3}}\sqrt{x} - \frac{1}{3}\sqrt{3}\right) + 2\sqrt{3}bc^{\frac{4}{3}}x^2 \arctan\left(\frac{2}{3}\sqrt{3}c^{\frac{1}{3}}\sqrt{x} - \frac{1}{3}\sqrt{3}\right) + b(-c)^{\frac{1}{3}}cx^2}{x^3}$$

input `integrate((a+b*arctanh(c*x^(3/2)))/x^3,x, algorithm="fricas")`

output

$$\begin{aligned} & -1/8*(2*\sqrt{3})*b*(-c)^{(1/3)}*c*x^2*\arctan(2/3*\sqrt{3})*(-c)^{(1/3)}*\sqrt{x} - \\ & 1/3*\sqrt{3}) + 2*\sqrt{3})*b*c^{(4/3)}*x^2*\arctan(2/3*\sqrt{3})*c^{(1/3)}*\sqrt{x} - \\ & 1/3*\sqrt{3}) + b*(-c)^{(1/3)}*c*x^2*\log(c*x + (-c)^{(2/3)}*\sqrt{x} - (-c)^{(1/3)}) + \\ & b*c^{(4/3)}*x^2*\log(c*x - c^{(2/3)}*\sqrt{x} + c^{(1/3)}) - 2*b*(-c)^{(1/3)}* \\ & c*x^2*\log(c*\sqrt{x} - (-c)^{(2/3)}) - 2*b*c^{(4/3)}*x^2*\log(c*\sqrt{x} + c^{(2/3)}) + \\ & 12*b*c*x^{(3/2)} + 2*b*\log(-c^2*x^3 + 2*c*x^{(3/2)} + 1)/(c^2*x^3 - 1) \\ &) + 4*a)/x^2 \end{aligned}$$
Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^3} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**(3/2)))/x**3,x)`

output Timed out

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.06

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^3} dx =$$

$$-\frac{1}{8} \left(\left(2\sqrt{3}c^{1/3} \arctan \left(\frac{\sqrt{3}(2c^{2/3}\sqrt{x} + c^{1/3})}{3c^{1/3}} \right) + 2\sqrt{3}c^{1/3} \arctan \left(\frac{\sqrt{3}(2c^{2/3}\sqrt{x} - c^{1/3})}{3c^{1/3}} \right) - c^{1/3} \log(c^{2/3}x + c^{1/3}) \right) - \frac{a}{2x^2} \right)$$

input `integrate((a+b*arctanh(c*x^(3/2)))/x^3,x, algorithm="maxima")`

output

```
-1/8*((2*sqrt(3)*c^(1/3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*sqrt(x) + c^(1/3))/
c^(1/3)) + 2*sqrt(3)*c^(1/3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*sqrt(x) - c^(1/3)
)/c^(1/3)) - c^(1/3)*log(c^(2/3)*x + c^(1/3)*sqrt(x) + 1) + c^(1/3)*log(
c^(2/3)*x - c^(1/3)*sqrt(x) + 1) - 2*c^(1/3)*log((c^(1/3)*sqrt(x) + 1)/c^(
1/3)) + 2*c^(1/3)*log((c^(1/3)*sqrt(x) - 1)/c^(1/3)) + 12/sqrt(x))*c + 4*a
rctanh(c*x^(3/2))/x^2)*b - 1/2*a/x^2
```

Giac [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.27

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^3} dx = -\frac{\sqrt{3}bc^3 \arctan\left(\frac{1}{3}\sqrt{3}\left(2\sqrt{x} + \frac{1}{|c|^{1/3}}\right)|c|^{1/3}\right)}{4|c|^{5/3}}$$

$$-\frac{\sqrt{3}bc^3 \arctan\left(\frac{1}{3}\sqrt{3}\left(2\sqrt{x} - \frac{1}{|c|^{1/3}}\right)|c|^{1/3}\right)}{4|c|^{5/3}} + \frac{bc^3 \log\left(x + \frac{\sqrt{x}}{|c|^{1/3}} + \frac{1}{|c|^{2/3}}\right)}{8|c|^{5/3}}$$

$$-\frac{bc^3 \log\left(x - \frac{\sqrt{x}}{|c|^{1/3}} + \frac{1}{|c|^{2/3}}\right)}{8|c|^{5/3}} + \frac{bc^3 \log\left(\sqrt{x} + \frac{1}{|c|^{1/3}}\right)}{4|c|^{5/3}}$$

$$-\frac{bc^3 \log\left(\left|\sqrt{x} - \frac{1}{|c|^{1/3}}\right|\right)}{4|c|^{5/3}} - \frac{b \log\left(-\frac{cx^{3/2}+1}{cx^{3/2}-1}\right)}{4x^2} - \frac{3bcx^{3/2} + a}{2x^2}$$

input `integrate((a+b*arctanh(c*x^(3/2)))/x^3,x, algorithm="giac")`

output
$$\begin{aligned} & -1/4*\sqrt{3}*b*c^3*\arctan(1/3*\sqrt{3}*(2*\sqrt{x} + 1/\text{abs}(c)^{(1/3)})*\text{abs}(c)^{(1/3)})/\text{abs}(c)^{(5/3)} - 1/4*\sqrt{3}*b*c^3*\arctan(1/3*\sqrt{3}*(2*\sqrt{x} - 1/\text{abs}(c)^{(1/3)})*\text{abs}(c)^{(1/3)})/\text{abs}(c)^{(5/3)} \\ & + 1/8*b*c^3*\log(x + \sqrt{x}/\text{abs}(c)^{(1/3)} + 1/\text{abs}(c)^{(2/3)})/\text{abs}(c)^{(5/3)} - 1/8*b*c^3*\log(x - \sqrt{x}/\text{abs}(c)^{(1/3)} + 1/\text{abs}(c)^{(2/3)})/\text{abs}(c)^{(5/3)} \\ & + 1/4*b*c^3*\log(\sqrt{x} + 1/\text{abs}(c)^{(1/3)})/\text{abs}(c)^{(5/3)} - 1/4*b*c^3*\log(\text{abs}(\sqrt{x} - 1/\text{abs}(c)^{(1/3)}))/\text{abs}(c)^{(5/3)} \\ & - 1/4*b*\log(-(c*x^(3/2) + 1)/(c*x^(3/2) - 1))/x^2 - 1/2*(3*b*c*x^(3/2) + a)/x^2 \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 10.47 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.44

$$\begin{aligned} \int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^3} dx &= \frac{bc^{4/3} \ln\left(\frac{c^{1/3}\sqrt{x+1}}{c^{1/3}\sqrt{x-1}}\right)}{4} - \frac{a}{2x^2} \\ &+ \frac{\ln\left(1 - cx^{3/2}\right) \left(\frac{bx}{2} - \frac{bc^2x^4}{2}\right)}{2x^3 - 2c^2x^6} - \frac{3bc}{2\sqrt{x}} - \frac{b \ln(cx^{3/2} + 1)}{4x^2} \\ &+ \frac{bc^{4/3} \ln\left(\frac{\sqrt{3} + c^{2/3}x \operatorname{li} + c^{1/3}\sqrt{x}4i - \sqrt{3}c^{2/3}x + \operatorname{li}}{2c^{2/3}x + 1 + \sqrt{3} \operatorname{li}}\right)}{4} \sqrt{-\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}} \\ &+ \frac{bc^{4/3} \ln\left(\frac{\sqrt{3}c^{2/3}x + c^{2/3}x \operatorname{li} - c^{1/3}\sqrt{x}4i - \sqrt{3} + \operatorname{li}}{2c^{2/3}x + 1 - \sqrt{3} \operatorname{li}}\right)}{4} \sqrt{\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}} \operatorname{li} \end{aligned}$$

input `int((a + b*atanh(c*x^(3/2)))/x^3,x)`

output
$$\begin{aligned} & (b*c^{(4/3)}*\log((c^{(1/3)}*x^{(1/2)} + 1)/(c^{(1/3)}*x^{(1/2)} - 1)))/4 - a/(2*x^2) \\ & + (\log(1 - c*x^{(3/2)})*((b*x)/2 - (b*c^2*x^4)/2))/(2*x^3 - 2*c^2*x^6) - (3*b*c)/(2*x^{(1/2)}) - (b*\log(c*x^{(3/2)} + 1))/(4*x^2) + (b*c^{(4/3)}*\log((3^{(1/2)} + c^{(2/3)}*x*\operatorname{li} + c^{(1/3)}*x^{(1/2)}*4i - 3^{(1/2)}*c^{(2/3)}*x + \operatorname{li}))/((3^{(1/2)}* \operatorname{li} + 2*c^{(2/3)}*x + 1))*((3^{(1/2)}*\operatorname{li})/2 - 1/2)^{(1/2)})/4 + (b*c^{(4/3)}*\log((c^{(2/3)}*x*\operatorname{li} - 3^{(1/2)} - c^{(1/3)}*x^{(1/2)}*4i + 3^{(1/2)}*c^{(2/3)}*x + \operatorname{li}))/((2*c^{(2/3)}*x - 3^{(1/2)}*\operatorname{li} + 1))*((3^{(1/2)}*\operatorname{li})/2 + 1/2)^{(1/2)}*\operatorname{li})/4 \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.92

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^3} dx = \frac{-2\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{x}c^{1/3}-1}{\sqrt{3}}\right) b c^2 x^2 - 2\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{x}c^{1/3}+1}{\sqrt{3}}\right) b c^2 x^2 - 4c^{2/3} \operatorname{atanh}(\sqrt{x}c^{1/3}) b c^2 x^2 - 4c^{2/3} a}{8c^{2/3} x^2}$$

input `int((a+b*atanh(c*x^(3/2)))/x^3,x)`

output

```
( - 2*sqrt(3)*atan((2*sqrt(x)*c**(1/3) - 1)/sqrt(3))*b*c**2*x**2 - 2*sqrt(
3)*atan((2*sqrt(x)*c**(1/3) + 1)/sqrt(3))*b*c**2*x**2 - 4*c**(2/3)*atanh(s
qrt(x)*c*x)*b - 2*atanh(sqrt(x)*c*x)*b*c**2*x**2 - 12*sqrt(x)*c**(2/3)*b*c
*x - 4*c**(2/3)*a + 3*log(sqrt(x)*c**(2/3) + c**(1/3))*b*c**2*x**2 - 3*log
(sqrt(x)*c**(2/3) - c**(1/3))*b*c**2*x**2)/(8*c**(2/3)*x**2)
```


3.219 $\int \frac{a+b\operatorname{arctanh}(cx^{3/2})}{x^4} dx$

Optimal result	1764
Mathematica [B] (verified)	1764
Rubi [A] (verified)	1765
Maple [A] (verified)	1767
Fricas [A] (verification not implemented)	1767
Sympy [F(-1)]	1768
Maxima [A] (verification not implemented)	1768
Giac [A] (verification not implemented)	1768
Mupad [B] (verification not implemented)	1769
Reduce [B] (verification not implemented)	1769

Optimal result

Integrand size = 16, antiderivative size = 47

$$\int \frac{a + b\operatorname{arctanh}(cx^{3/2})}{x^4} dx = -\frac{bc}{3x^{3/2}} + \frac{1}{3}bc^2\operatorname{arctanh}(cx^{3/2}) - \frac{a + b\operatorname{arctanh}(cx^{3/2})}{3x^3}$$

output `-1/3*b*c/x^(3/2)+1/3*b*c^2*arctanh(c*x^(3/2))-1/3*(a+b*arctanh(c*x^(3/2)))/x^3`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 140 vs. 2(47) = 94.

Time = 0.03 (sec) , antiderivative size = 140, normalized size of antiderivative = 2.98

$$\int \frac{a + b\operatorname{arctanh}(cx^{3/2})}{x^4} dx = -\frac{a}{3x^3} - \frac{bc}{3x^{3/2}} - \frac{b\operatorname{arctanh}(cx^{3/2})}{3x^3} - \frac{1}{6}bc^2 \log(1 - \sqrt[3]{c}\sqrt{x}) + \frac{1}{6}bc^2 \log(1 + \sqrt[3]{c}\sqrt{x}) + \frac{1}{6}bc^2 \log(1 - \sqrt[3]{c}\sqrt{x} + c^{2/3}x) - \frac{1}{6}bc^2 \log(1 + \sqrt[3]{c}\sqrt{x} + c^{2/3}x)$$

input `Integrate[(a + b*ArcTanh[c*x^(3/2)])/x^4, x]`

output

$$-1/3*a/x^3 - (b*c)/(3*x^{(3/2)}) - (b*ArcTanh[c*x^{(3/2)}])/(3*x^3) - (b*c^2*Log[1 - c^{(1/3)*Sqrt[x]}])/6 + (b*c^2*Log[1 + c^{(1/3)*Sqrt[x]}])/6 + (b*c^2*Log[1 - c^{(1/3)*Sqrt[x]} + c^{(2/3)*x}])/6 - (b*c^2*Log[1 + c^{(1/3)*Sqrt[x]} + c^{(2/3)*x}])/6$$
Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.04, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6452, 847, 851, 807, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + \text{barctanh}(cx^{3/2})}{x^4} dx \\ & \quad \downarrow \text{6452} \\ & \frac{1}{2}bc \int \frac{1}{x^{5/2}(1 - c^2x^3)} dx - \frac{a + \text{barctanh}(cx^{3/2})}{3x^3} \\ & \quad \downarrow \text{847} \\ & \frac{1}{2}bc \left(c^2 \int \frac{\sqrt{x}}{1 - c^2x^3} dx - \frac{2}{3x^{3/2}} \right) - \frac{a + \text{barctanh}(cx^{3/2})}{3x^3} \\ & \quad \downarrow \text{851} \\ & \frac{1}{2}bc \left(2c^2 \int \frac{x}{1 - c^2x^3} d\sqrt{x} - \frac{2}{3x^{3/2}} \right) - \frac{a + \text{barctanh}(cx^{3/2})}{3x^3} \\ & \quad \downarrow \text{807} \\ & \frac{1}{2}bc \left(\frac{2}{3}c^2 \int \frac{1}{1 - c^2x} dx^{3/2} - \frac{2}{3x^{3/2}} \right) - \frac{a + \text{barctanh}(cx^{3/2})}{3x^3} \\ & \quad \downarrow \text{219} \\ & \frac{1}{2}bc \left(\frac{2}{3} \text{carctanh}(cx^{3/2}) - \frac{2}{3x^{3/2}} \right) - \frac{a + \text{barctanh}(cx^{3/2})}{3x^3} \end{aligned}$$

input

$$\text{Int}[(a + b*ArcTanh[c*x^{(3/2)}])/x^4, x]$$

output
$$-1/3*(a + b*\text{ArcTanh}[c*x^{(3/2)}])/x^3 + (b*c*(-2/(3*x^{(3/2)}) + (2*c*\text{ArcTanh}[c*x^{(3/2)}])/3))/2$$

Defintions of rubi rules used

rule 219
$$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \text{ :> } \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] \text{ ; FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 807
$$\text{Int}[(x_)^{(m_)}*((a_ + (b_)*(x_)^{(n_)})^{(p_)}), x_Symbol] \text{ :> } \text{With}\{k = \text{GCD}[m + 1, n]\}, \text{Simp}[1/k \ \text{Subst}[\text{Int}[x^{(m + 1)/k - 1}*(a + b*x^{(n/k)})^p, x], x, x^k], x] \text{ ; } k \neq 1 \text{ ; FreeQ}\{a, b, p\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$$

rule 847
$$\text{Int}[(c_)*(x_)^{(m_)}*((a_ + (b_)*(x_)^{(n_)})^{(p_)}), x_Symbol] \text{ :> } \text{Simp}[(c*x)^{(m + 1)}*((a + b*x^n)^{(p + 1)}/(a*c*(m + 1))), x] - \text{Simp}[b*(m + n*(p + 1) + 1)/(a*c^n*(m + 1)) \ \text{Int}[(c*x)^{(m + n)}*(a + b*x^n)^p, x], x] \text{ ; FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$$

rule 851
$$\text{Int}[(c_)*(x_)^{(m_)}*((a_ + (b_)*(x_)^{(n_)})^{(p_)}), x_Symbol] \text{ :> } \text{With}\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \ \text{Subst}[\text{Int}[x^{(k*(m + 1) - 1)}*(a + b*(x^{(k*n)}/c^n))^p, x], x, (c*x)^{(1/k)}], x] \text{ ; FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$$

rule 6452
$$\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}], x_Symbol] \text{ :> } \text{Simp}[x^{(m + 1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m + 1)), x] - \text{Simp}[b*c*n*(p/(m + 1)) \ \text{Int}[x^{(m + n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p - 1)})/(1 - c^2*x^{(2*n)}), x], x] \text{ ; FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$$

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.17

method	result	size
derivativedivides	$-\frac{a}{3x^3} - \frac{b \operatorname{arctanh}(cx^{\frac{3}{2}})}{3x^3} - \frac{bc}{3x^{\frac{3}{2}}} + \frac{bc^2 \ln(cx^{\frac{3}{2}}+1)}{6} - \frac{bc^2 \ln(cx^{\frac{3}{2}}-1)}{6}$	55
default	$-\frac{a}{3x^3} - \frac{b \operatorname{arctanh}(cx^{\frac{3}{2}})}{3x^3} - \frac{bc}{3x^{\frac{3}{2}}} + \frac{bc^2 \ln(cx^{\frac{3}{2}}+1)}{6} - \frac{bc^2 \ln(cx^{\frac{3}{2}}-1)}{6}$	55
parts	$-\frac{a}{3x^3} - \frac{b \operatorname{arctanh}(cx^{\frac{3}{2}})}{3x^3} - \frac{bc}{3x^{\frac{3}{2}}} + \frac{bc^2 \ln(cx^{\frac{3}{2}}+1)}{6} - \frac{bc^2 \ln(cx^{\frac{3}{2}}-1)}{6}$	55

input `int((a+b*arctanh(c*x^(3/2)))/x^4,x,method=_RETURNVERBOSE)`

output
$$-1/3*a/x^3-1/3*b/x^3*arctanh(c*x^(3/2))-1/3*b*c/x^(3/2)+1/6*b*c^2*\ln(c*x^(3/2)+1)-1/6*b*c^2*\ln(c*x^(3/2)-1)$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.26

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^4} dx = -\frac{2bcx^{\frac{3}{2}} - (bc^2x^3 - b) \log\left(-\frac{c^2x^3+2cx^{\frac{3}{2}}+1}{c^2x^3-1}\right) + 2a}{6x^3}$$

input `integrate((a+b*arctanh(c*x^(3/2)))/x^4,x, algorithm="fricas")`

output
$$-1/6*(2*b*c*x^(3/2) - (b*c^2*x^3 - b)*\log(-(c^2*x^3 + 2*c*x^(3/2) + 1)/(c^2*x^3 - 1)) + 2*a)/x^3$$

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^4} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**(3/2)))/x**4,x)`output `Timed out`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.09

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^4} dx = \frac{1}{6} \left(\left(c \log(cx^{\frac{3}{2}} + 1) - c \log(cx^{\frac{3}{2}} - 1) - \frac{2}{x^{\frac{3}{2}}} \right) c - \frac{2 \operatorname{artanh}(cx^{\frac{3}{2}})}{x^3} \right) b - \frac{a}{3x^3}$$

input `integrate((a+b*arctanh(c*x^(3/2)))/x^4,x, algorithm="maxima")`output `1/6*((c*log(c*x^(3/2) + 1) - c*log(c*x^(3/2) - 1) - 2/x^(3/2))*c - 2*arctanh(c*x^(3/2))/x^3)*b - 1/3*a/x^3`**Giac [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.43

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^4} dx = \frac{1}{6} bc^2 \log(cx^{\frac{3}{2}} + 1) - \frac{1}{6} bc^2 \log(cx^{\frac{3}{2}} - 1) - \frac{b \log\left(\frac{-cx^{\frac{3}{2}} + 1}{cx^{\frac{3}{2}} - 1}\right)}{6x^3} - \frac{bcx^{\frac{3}{2}} + a}{3x^3}$$

input `integrate((a+b*arctanh(c*x^(3/2)))/x^4,x, algorithm="giac")`

output $1/6*b*c^2*\log(c*x^{(3/2)} + 1) - 1/6*b*c^2*\log(c*x^{(3/2)} - 1) - 1/6*b*\log(-(c*x^{(3/2)} + 1)/(c*x^{(3/2)} - 1))/x^3 - 1/3*(b*c*x^{(3/2)} + a)/x^3$

Mupad [B] (verification not implemented)

Time = 3.92 (sec) , antiderivative size = 114, normalized size of antiderivative = 2.43

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^4} dx = \frac{b c^2 \ln\left(\frac{cx^{3/2}+1}{cx^{3/2}-1}\right)}{6} - \frac{a}{3x^3} - \frac{bc}{3x^{3/2}} - \frac{b \ln(cx^{3/2}+1)}{6x^3} + \frac{bx \ln(1-cx^{3/2})}{3(2x^4-2c^2x^7)} - \frac{bc^2x^4 \ln(1-cx^{3/2})}{3(2x^4-2c^2x^7)}$$

input `int((a + b*atanh(c*x^(3/2)))/x^4,x)`

output $(b*c^2*\log((c*x^{(3/2)} + 1)/(c*x^{(3/2)} - 1)))/6 - a/(3*x^3) - (b*c)/(3*x^{(3/2)}) - (b*\log(c*x^{(3/2)} + 1))/(6*x^3) + (b*x*\log(1 - c*x^{(3/2)}))/(3*(2*x^4 - 2*c^2*x^7)) - (b*c^2*x^4*\log(1 - c*x^{(3/2)}))/(3*(2*x^4 - 2*c^2*x^7))$

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.83

$$\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^4} dx = \frac{\operatorname{atanh}(\sqrt{x} cx) b c^2 x^3 - \operatorname{atanh}(\sqrt{x} cx) b - \sqrt{x} bcx - a}{3x^3}$$

input `int((a+b*atanh(c*x^(3/2)))/x^4,x)`

output $(\operatorname{atanh}(\sqrt{x}*c*x)*b*c**2*x**3 - \operatorname{atanh}(\sqrt{x}*c*x)*b - \sqrt{x}*b*c*x - a)/(3*x**3)$

3.220 $\int x^2 (a + b \operatorname{arctanh}(cx^{3/2}))^2 dx$

Optimal result	1770
Mathematica [A] (verified)	1770
Rubi [A] (verified)	1771
Maple [B] (verified)	1773
Fricas [B] (verification not implemented)	1773
Sympy [F(-1)]	1774
Maxima [B] (verification not implemented)	1774
Giac [F]	1775
Mupad [B] (verification not implemented)	1775
Reduce [B] (verification not implemented)	1776

Optimal result

Integrand size = 18, antiderivative size = 101

$$\int x^2 (a + b \operatorname{arctanh}(cx^{3/2}))^2 dx = \frac{2abx^{3/2}}{3c} + \frac{2b^2x^{3/2}\operatorname{arctanh}(cx^{3/2})}{3c} - \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{3c^2} + \frac{1}{3}x^3(a + b \operatorname{arctanh}(cx^{3/2}))^2 + \frac{b^2 \log(1 - c^2x^3)}{3c^2}$$

output $2/3*a*b*x^{(3/2)}/c+2/3*b^2*x^{(3/2)}*arctanh(c*x^{(3/2)})/c-1/3*(a+b*arctanh(c*x^{(3/2)}))^2/c^2+1/3*x^3*(a+b*arctanh(c*x^{(3/2)}))^2+1/3*b^2*\ln(-c^2*x^3+1)/c^2$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.21

$$\int x^2 (a + b \operatorname{arctanh}(cx^{3/2}))^2 dx = \frac{2abcx^{3/2} + a^2c^2x^3 + 2bcx^{3/2}(b + acx^{3/2}) \operatorname{arctanh}(cx^{3/2}) + b^2(-1 + c^2x^3) \operatorname{arctanh}(cx^{3/2})}{3c^2}$$

input `Integrate[x^2*(a + b*ArcTanh[c*x^(3/2)])^2,x]`

output

$$\frac{(2abcx^{3/2} + a^2c^2x^3 + 2bcx^{3/2}(b + acx^{3/2}))\operatorname{ArcTanh}[cx^{3/2}] + b^2(-1 + c^2x^3)\operatorname{ArcTanh}[cx^{3/2}]^2 + b(a + b)\operatorname{Log}[1 - cx^{3/2}] - ab\operatorname{Log}[1 + cx^{3/2}] + b^2\operatorname{Log}[1 + cx^{3/2}]}{3c^2}$$
Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.01, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {6454, 6452, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 (a + b \operatorname{arctanh}(cx^{3/2}))^2 dx$$

$$\downarrow 6454$$

$$\frac{2}{3} \int x^{3/2} (a + b \operatorname{arctanh}(cx^{3/2}))^2 dx^{3/2}$$

$$\downarrow 6452$$

$$\frac{2}{3} \left(\frac{1}{2} x^3 (a + b \operatorname{arctanh}(cx^{3/2}))^2 - bc \int \frac{x^3 (a + b \operatorname{arctanh}(cx^{3/2}))}{1 - c^2 x^3} dx^{3/2} \right)$$

$$\downarrow 6542$$

$$\frac{2}{3} \left(\frac{1}{2} x^3 (a + b \operatorname{arctanh}(cx^{3/2}))^2 - bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{1 - c^2 x^3} dx^{3/2}}{c^2} - \frac{\int (a + b \operatorname{arctanh}(cx^{3/2})) dx^{3/2}}{c^2} \right) \right)$$

$$\downarrow 2009$$

$$\frac{2}{3} \left(\frac{1}{2} x^3 (a + b \operatorname{arctanh}(cx^{3/2}))^2 - bc \left(\frac{\int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{1 - c^2 x^3} dx^{3/2}}{c^2} - \frac{ax^{3/2} + bx^{3/2} \operatorname{arctanh}(cx^{3/2}) + \frac{b \log(1 - c^2 x^3)}{2c}}{c^2} \right) \right)$$

$$\downarrow 6510$$

$$\frac{2}{3} \left(\frac{1}{2} x^3 (a + b \operatorname{arctanh}(cx^{3/2}))^2 - bc \left(\frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{2bc^3} - \frac{ax^{3/2} + bx^{3/2} \operatorname{arctanh}(cx^{3/2}) + \frac{b \log(1 - c^2 x^3)}{2c}}{c^2} \right) \right)$$

input `Int[x^2*(a + b*ArcTanh[c*x^(3/2)])^2,x]`

output `(2*((x^3*(a + b*ArcTanh[c*x^(3/2)])^2)/2 - b*c*((a + b*ArcTanh[c*x^(3/2)])^2/(2*b*c^3 - (a*x^(3/2) + b*x^(3/2)*ArcTanh[c*x^(3/2)] + (b*Log[1 - c^2*x^3])/(2*c))/c^2)))/3`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6454 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTanh[c*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify[(m + 1)/n]]`

rule 6510 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6542 `Int[(((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((f_.)*(x_)^(m_.)))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 219 vs. 2(81) = 162.

Time = 1.16 (sec) , antiderivative size = 220, normalized size of antiderivative = 2.18

method	result
parts	$\frac{a^2 x^3}{3} + \frac{2b^2 \left(\frac{c^2 x^3 \operatorname{arctanh}\left(c x^{\frac{3}{2}}\right)^2}{2} + \operatorname{arctanh}\left(c x^{\frac{3}{2}}\right) c x^{\frac{3}{2}} + \frac{\operatorname{arctanh}\left(c x^{\frac{3}{2}}\right) \ln\left(c x^{\frac{3}{2}} - 1\right)}{2} - \frac{\operatorname{arctanh}\left(c x^{\frac{3}{2}}\right) \ln\left(c x^{\frac{3}{2}} + 1\right)}{2} \right)}{3}$
derivativedivides	$\frac{a^2 c^2 x^3}{3} + \frac{2b^2 \left(\frac{c^2 x^3 \operatorname{arctanh}\left(c x^{\frac{3}{2}}\right)^2}{2} + \operatorname{arctanh}\left(c x^{\frac{3}{2}}\right) c x^{\frac{3}{2}} + \frac{\operatorname{arctanh}\left(c x^{\frac{3}{2}}\right) \ln\left(c x^{\frac{3}{2}} - 1\right)}{2} - \frac{\operatorname{arctanh}\left(c x^{\frac{3}{2}}\right) \ln\left(c x^{\frac{3}{2}} + 1\right)}{2} \right)}{3}$
default	$\frac{a^2 c^2 x^3}{3} + \frac{2b^2 \left(\frac{c^2 x^3 \operatorname{arctanh}\left(c x^{\frac{3}{2}}\right)^2}{2} + \operatorname{arctanh}\left(c x^{\frac{3}{2}}\right) c x^{\frac{3}{2}} + \frac{\operatorname{arctanh}\left(c x^{\frac{3}{2}}\right) \ln\left(c x^{\frac{3}{2}} - 1\right)}{2} - \frac{\operatorname{arctanh}\left(c x^{\frac{3}{2}}\right) \ln\left(c x^{\frac{3}{2}} + 1\right)}{2} \right)}{3}$

```
input int(x^2*(a+b*arctanh(c*x^(3/2)))^2,x,method=_RETURNVERBOSE)
```

```
output 1/3*a^2*x^3+2/3*b^2/c^2*(1/2*c^2*x^3*arctanh(c*x^(3/2))^2+arctanh(c*x^(3/2))
)*c*x^(3/2)+1/2*arctanh(c*x^(3/2))*ln(c*x^(3/2)-1)-1/2*arctanh(c*x^(3/2))
*ln(c*x^(3/2)+1)+1/8*ln(c*x^(3/2)-1)^2-1/4*ln(c*x^(3/2)-1)*ln(1/2*c*x^(3/2)
)+1/2)+1/2*ln(c*x^(3/2)-1)+1/2*ln(c*x^(3/2)+1)-1/4*(ln(c*x^(3/2)+1)-ln(1/2
*c*x^(3/2)+1/2))*ln(-1/2*c*x^(3/2)+1/2)+1/8*ln(c*x^(3/2)+1)^2+4/3*a*b/c^2
*(1/2*c^2*x^3*arctanh(c*x^(3/2))+1/2*c*x^(3/2)+1/4*ln(c*x^(3/2)-1)-1/4*ln(
c*x^(3/2)+1))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 179 vs. 2(81) = 162.

Time = 0.11 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.77

$$\int x^2(a + b \operatorname{arctanh}(cx^{3/2}))^2 dx = \frac{4a^2c^2x^3 + 8abcx^{\frac{3}{2}} + (b^2c^2x^3 - b^2) \log\left(-\frac{c^2x^3 + 2cx^{\frac{3}{2}} + 1}{c^2x^3 - 1}\right)^2 + 4(abc^2 - ab + b^2) \log\left(-\frac{c^2x^3 + 2cx^{\frac{3}{2}} + 1}{c^2x^3 - 1}\right) \log\left(\frac{c^2x^3 + 2cx^{\frac{3}{2}} + 1}{c^2x^3 - 1}\right)}{3}$$

input `integrate(x^2*(a+b*arctanh(c*x^(3/2)))^2,x, algorithm="fricas")`

output
$$\frac{1}{12}(4a^2c^2x^3 + 8abcx^{3/2} + (b^2c^2x^3 - b^2)\log(-(c^2x^3 + 2cx^{3/2} + 1)/(c^2x^3 - 1))^2 + 4(a^2c^2 - ab + b^2)\log(cx^{3/2} + 1) - 4(a^2c^2 - ab - b^2)\log(cx^{3/2} - 1) + 4(a^2c^2x^3 + b^2cx^{3/2} - abc^2)\log(-(c^2x^3 + 2cx^{3/2} + 1)/(c^2x^3 - 1)))/c^2$$

Sympy [F(-1)]

Timed out.

$$\int x^2(a + b\operatorname{arctanh}(cx^{3/2}))^2 dx = \text{Timed out}$$

input `integrate(x**2*(a+b*atanh(c*x**(3/2)))**2,x)`

output Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 186 vs. $2(81) = 162$.

Time = 0.04 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.84

$$\begin{aligned} \int x^2(a + b\operatorname{arctanh}(cx^{3/2}))^2 dx &= \frac{1}{3}b^2x^3 \operatorname{artanh}\left(cx^{\frac{3}{2}}\right)^2 + \frac{1}{3}a^2x^3 \\ &+ \frac{1}{3}\left(2x^3 \operatorname{artanh}\left(cx^{\frac{3}{2}}\right) + c\left(\frac{2x^{\frac{3}{2}}}{c^2} - \frac{\log\left(cx^{\frac{3}{2}} + 1\right)}{c^3} + \frac{\log\left(cx^{\frac{3}{2}} - 1\right)}{c^3}\right)\right)ab \\ &+ \frac{1}{12}\left(4c\left(\frac{2x^{\frac{3}{2}}}{c^2} - \frac{\log\left(cx^{\frac{3}{2}} + 1\right)}{c^3} + \frac{\log\left(cx^{\frac{3}{2}} - 1\right)}{c^3}\right) \operatorname{artanh}\left(cx^{\frac{3}{2}}\right) - \frac{2\left(\log\left(cx^{\frac{3}{2}} - 1\right) - 2\right)\log\left(cx^{\frac{3}{2}} + 1\right)}{c^3}\right) \end{aligned}$$

input `integrate(x^2*(a+b*arctanh(c*x^(3/2)))^2,x, algorithm="maxima")`

output

```
1/3*b^2*x^3*arctanh(c*x^(3/2))^2 + 1/3*a^2*x^3 + 1/3*(2*x^3*arctanh(c*x^(3/2)) + c*(2*x^(3/2)/c^2 - log(c*x^(3/2) + 1)/c^3 + log(c*x^(3/2) - 1)/c^3)*a*b + 1/12*(4*c*(2*x^(3/2)/c^2 - log(c*x^(3/2) + 1)/c^3 + log(c*x^(3/2) - 1)/c^3)*arctanh(c*x^(3/2)) - (2*(log(c*x^(3/2) - 1) - 2)*log(c*x^(3/2) + 1) - log(c*x^(3/2) + 1)^2 - log(c*x^(3/2) - 1)^2 - 4*log(c*x^(3/2) - 1))/c^2)*b^2
```

Giac [F]

$$\int x^2 (a + b \operatorname{arctanh}(cx^{3/2}))^2 dx = \int (b \operatorname{arctanh}(cx^{3/2}) + a)^2 x^2 dx$$

input

```
integrate(x^2*(a+b*arctanh(c*x^(3/2)))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^(3/2)) + a)^2*x^2, x)
```

Mupad [B] (verification not implemented)

Time = 3.89 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.04

$$\int x^2 (a + b \operatorname{arctanh}(cx^{3/2}))^2 dx = \frac{c \left(\frac{2b^2 x^{3/2} \operatorname{atanh}(cx^{3/2})}{3} + \frac{2abx^{3/2}}{3} \right) - \frac{b^2 \operatorname{atanh}(cx^{3/2})^2}{3} + \frac{b^2 \ln(c^2 x^3 - 1)}{3} - \frac{2ab \operatorname{atanh}(cx^{3/2})}{3}}{c^2} + \frac{a^2 x^3}{3} + \frac{b^2 x^3 \operatorname{atanh}(cx^{3/2})^2}{3} + \frac{2abx^3 \operatorname{atanh}(cx^{3/2})}{3}$$

input

```
int(x^2*(a + b*atanh(c*x^(3/2)))^2,x)
```

output

```
(c*((2*b^2*x^(3/2)*atanh(c*x^(3/2)))/3 + (2*a*b*x^(3/2))/3) - (b^2*atanh(c*x^(3/2))^2)/3 + (b^2*log(c^2*x^3 - 1))/3 - (2*a*b*atanh(c*x^(3/2)))/3)/c^2 + (a^2*x^3)/3 + (b^2*x^3*atanh(c*x^(3/2))^2)/3 + (2*a*b*x^3*atanh(c*x^(3/2)))/3
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.42

$$\int x^2 (a + b \operatorname{arctanh}(cx^{3/2}))^2 dx = \frac{\operatorname{atanh}(\sqrt{x} cx)^2 b^2 c^2 x^3 - \operatorname{atanh}(\sqrt{x} cx)^2 b^2 + 2\sqrt{x} \operatorname{atanh}(\sqrt{x} cx) b^2 cx + 2 \operatorname{atanh}(\sqrt{x} cx) a b c x^{3/2} + a^2 c x^{3/2}}{3c^2}$$

input

```
int(x^2*(a+b*atanh(c*x^(3/2)))^2,x)
```

output

```
(atanh(sqrt(x)*c*x)**2*b**2*c**2*x**3 - atanh(sqrt(x)*c*x)**2*b**2 + 2*sqrt(x)*atanh(sqrt(x)*c*x)*b**2*c*x + 2*atanh(sqrt(x)*c*x)*a*b*c**2*x**3 - 2*atanh(sqrt(x)*c*x)*a*b - 2*atanh(sqrt(x)*c*x)*b**2 + 2*sqrt(x)*a*b*c*x + 2*log(c**(2/3)*x - sqrt(x)*c**(1/3) + 1)*b**2 + 2*log(sqrt(x)*c**(2/3) + c**(1/3))*b**2 + a**2*c**2*x**3)/(3*c**2)
```

3.221
$$\int \frac{\left(a+b\operatorname{arctanh}\left(cx^{3/2}\right)\right)^2}{x} dx$$

Optimal result	1777
Mathematica [C] (verified)	1777
Rubi [A] (verified)	1778
Maple [C] (warning: unable to verify)	1780
Fricas [F]	1782
Sympy [F(-1)]	1782
Maxima [F]	1783
Giac [F]	1783
Mupad [F(-1)]	1784
Reduce [F]	1784

Optimal result

Integrand size = 18, antiderivative size = 156

$$\int \frac{(a + b\operatorname{arctanh}(cx^{3/2}))^2}{x} dx = \frac{4}{3}(a + b\operatorname{arctanh}(cx^{3/2}))^2 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx^{3/2}}\right) - \frac{2}{3}b(a + b\operatorname{arctanh}(cx^{3/2})) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx^{3/2}}\right) + \frac{2}{3}b(a + b\operatorname{arctanh}(cx^{3/2}))^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - cx^{3/2}}\right)$$

output

```
-4/3*(a+b*arctanh(c*x^(3/2)))^2*arctanh(-1+2/(1-c*x^(3/2)))-2/3*b*(a+b*arctanh(c*x^(3/2)))*polylog(2,1-2/(1-c*x^(3/2)))+2/3*b*(a+b*arctanh(c*x^(3/2)))*polylog(2,-1+2/(1-c*x^(3/2)))+1/3*b^2*polylog(3,1-2/(1-c*x^(3/2)))-1/3*b^2*polylog(3,-1+2/(1-c*x^(3/2)))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.33

$$\int \frac{(a + b\operatorname{arctanh}(cx^{3/2}))^2}{x} dx = a^2 \log(x) + \frac{2}{3}ab(-\operatorname{PolyLog}(2, -cx^{3/2}) + \operatorname{PolyLog}(2, cx^{3/2})) + \frac{2}{3}b^2\left(\frac{i\pi^3}{24} - \frac{2}{3}\operatorname{arctanh}(cx^{3/2})^3 - \operatorname{arctanh}(cx^{3/2})^2 \log(1 - cx^{3/2})\right)$$

input `Integrate[(a + b*ArcTanh[c*x^(3/2)])^2/x,x]`

output $a^2 \text{Log}[x] + (2ab(-\text{PolyLog}[2, -(cx^{3/2})] + \text{PolyLog}[2, cx^{3/2}]))/3 + (2b^2((I/24)\pi^3 - (2\text{ArcTanh}[cx^{3/2}]^3)/3 - \text{ArcTanh}[cx^{3/2}]^2 \text{Log}[1 + E^{-(2\text{ArcTanh}[cx^{3/2}])}] + \text{ArcTanh}[cx^{3/2}]^2 \text{Log}[1 - E^{(2\text{ArcTanh}[cx^{3/2}])}] + \text{ArcTanh}[cx^{3/2}]\text{PolyLog}[2, -E^{-(2\text{ArcTanh}[cx^{3/2}])}] + \text{ArcTanh}[cx^{3/2}]\text{PolyLog}[2, E^{(2\text{ArcTanh}[cx^{3/2}])}] + \text{PolyLog}[3, -E^{-(2\text{ArcTanh}[cx^{3/2}])}]/2 - \text{PolyLog}[3, E^{(2\text{ArcTanh}[cx^{3/2}])}]/2))/3$

Rubi [A] (verified)

Time = 0.86 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.15, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {6450, 6448, 6614, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{x} dx$$

↓ 6450

$$\frac{2}{3} \int \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{x^{3/2}} dx^{3/2}$$

↓ 6448

$$\frac{2}{3} \left(2 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx^{3/2}}\right) (a + b \operatorname{arctanh}(cx^{3/2}))^2 - 4bc \int \frac{(a + b \operatorname{arctanh}(cx^{3/2})) \operatorname{arctanh}\left(1 - \frac{2}{1 - cx^{3/2}}\right)}{1 - c^2 x^3} dx \right)$$

↓ 6614

$$\frac{2}{3} \left(2 \operatorname{arctanh}\left(1 - \frac{2}{1 - cx^{3/2}}\right) (a + b \operatorname{arctanh}(cx^{3/2}))^2 - 4bc \left(\frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(cx^{3/2})) \log\left(2 - \frac{2}{1 - cx^{3/2}}\right)}{1 - c^2 x^3} dx \right) \right)$$

↓ 6620

$$\frac{2}{3} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^{3/2}} \right) \left(a + \operatorname{arctanh} \left(cx^{3/2} \right) \right)^2 - 4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - cx^{3/2}} \right) \left(a + \operatorname{arctanh} \left(cx^{3/2} \right) \right)}{2c} \right) \right) \right)$$

↓ 7164

$$\frac{2}{3} \left(2 \operatorname{arctanh} \left(1 - \frac{2}{1 - cx^{3/2}} \right) \left(a + \operatorname{arctanh} \left(cx^{3/2} \right) \right)^2 - 4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog} \left(2, 1 - \frac{2}{1 - cx^{3/2}} \right) \left(a + \operatorname{arctanh} \left(cx^{3/2} \right) \right)}{2c} \right) \right) \right)$$

input `Int[(a + b*ArcTanh[c*x^(3/2)])^2/x,x]`

output `(2*(2*(a + b*ArcTanh[c*x^(3/2)])^2*ArcTanh[1 - 2/(1 - c*x^(3/2))] - 4*b*c*((((a + b*ArcTanh[c*x^(3/2)])*PolyLog[2, 1 - 2/(1 - c*x^(3/2))])/(2*c) - (b*PolyLog[3, 1 - 2/(1 - c*x^(3/2))])/(4*c))/2 + (-1/2*(a + b*ArcTanh[c*x^(3/2)])*PolyLog[2, -1 + 2/(1 - c*x^(3/2))])/c + (b*PolyLog[3, -1 + 2/(1 - c*x^(3/2))])/(4*c))/2))/3`

Defintions of rubi rules used

rule 6448 `Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^p_/(x_), x_Symbol] := Simp[2*(a + b*ArcTanh[c*x])^p*ArcTanh[1 - 2/(1 - c*x)], x] - Simp[2*b*c*p Int[(a + b*ArcTanh[c*x])^(p - 1)*(ArcTanh[1 - 2/(1 - c*x)]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1]`

rule 6450 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)]*(b_.))^p_/(x_), x_Symbol] := Simp[1/n Subst[Int[(a + b*ArcTanh[c*x])^p/x, x], x, x^n], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]`

rule 6614 `Int[(ArcTanh[u]*((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^p_)/((d_.) + (e_.)*(x_)^2), x_Symbol] := Simp[1/2 Int[Log[1 + u]*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] - Simp[1/2 Int[Log[1 - u]*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[u^2 - (1 - 2/(1 - c*x))^2, 0]`

rule 6620

```
Int[(Log[u_]*)((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.))/((d_) + (e_.)*(x_)^
2), x_Symbol] := Simp[(-(a + b*ArcTanh[c*x])^p)*(PolyLog[2, 1 - u]/(2*c*d))
, x] + Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(
d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d
+ e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 18.05 (sec) , antiderivative size = 706, normalized size of antiderivative = 4.53

method	result
parts	$a^2 \ln(x) + b^2 \left(\frac{2 \ln(cx^{\frac{3}{2}}) \operatorname{arctanh}(cx^{\frac{3}{2}})^2}{3} - \frac{2 \operatorname{arctanh}(cx^{\frac{3}{2}}) \operatorname{polylog}\left(2, -\frac{(cx^{\frac{3}{2}}+1)^2}{-c^2x^3+1}\right)}{3} + \frac{\operatorname{polylog}\left(3, -\frac{(cx^{\frac{3}{2}}+1)^2}{-c^2x^3+1}\right)}{3} \right)$
derivativedivides	$\frac{2a^2 \ln(cx^{\frac{3}{2}})}{3} + \frac{2b^2 \left(\ln(cx^{\frac{3}{2}}) \operatorname{arctanh}(cx^{\frac{3}{2}})^2 - \operatorname{arctanh}(cx^{\frac{3}{2}}) \operatorname{polylog}\left(2, -\frac{(cx^{\frac{3}{2}}+1)^2}{-c^2x^3+1}\right) + \frac{\operatorname{polylog}\left(3, -\frac{(cx^{\frac{3}{2}}+1)^2}{-c^2x^3+1}\right)}{2} \right)}{2}$
default	$\frac{2a^2 \ln(cx^{\frac{3}{2}})}{3} + \frac{2b^2 \left(\ln(cx^{\frac{3}{2}}) \operatorname{arctanh}(cx^{\frac{3}{2}})^2 - \operatorname{arctanh}(cx^{\frac{3}{2}}) \operatorname{polylog}\left(2, -\frac{(cx^{\frac{3}{2}}+1)^2}{-c^2x^3+1}\right) + \frac{\operatorname{polylog}\left(3, -\frac{(cx^{\frac{3}{2}}+1)^2}{-c^2x^3+1}\right)}{2} \right)}{2}$

input

```
int((a+b*arctanh(c*x^(3/2)))^2/x,x,method=_RETURNVERBOSE)
```

output

```
a^2*ln(x)+b^2*(2/3*ln(c*x^(3/2))*arctanh(c*x^(3/2))^2-2/3*arctanh(c*x^(3/2))
)*polylog(2,-(c*x^(3/2)+1)^2/(-c^2*x^3+1))+1/3*polylog(3,-(c*x^(3/2)+1)^2
/(-c^2*x^3+1))-2/3*arctanh(c*x^(3/2))^2*ln((c*x^(3/2)+1)^2/(-c^2*x^3+1)-1)
+2/3*arctanh(c*x^(3/2))^2*ln(1-(c*x^(3/2)+1)/(-c^2*x^3+1)^(1/2))+4/3*arcta
nh(c*x^(3/2))*polylog(2,(c*x^(3/2)+1)/(-c^2*x^3+1)^(1/2))-4/3*polylog(3,(c
*x^(3/2)+1)/(-c^2*x^3+1)^(1/2))+2/3*arctanh(c*x^(3/2))^2*ln(1+(c*x^(3/2)+1
)/(-c^2*x^3+1)^(1/2))+4/3*arctanh(c*x^(3/2))*polylog(2,-(c*x^(3/2)+1)/(-c^
2*x^3+1)^(1/2))-4/3*polylog(3,-(c*x^(3/2)+1)/(-c^2*x^3+1)^(1/2))+1/3*I*Pi*
csgn(I*(-(c*x^(3/2)+1)^2/(c^2*x^3-1)-1)/(1-(c*x^(3/2)+1)^2/(c^2*x^3-1)))*
csgn(I*(-(c*x^(3/2)+1)^2/(c^2*x^3-1)-1))*csgn(I/(1-(c*x^(3/2)+1)^2/(c^2*x^
3-1)))-csgn(I*(-(c*x^(3/2)+1)^2/(c^2*x^3-1)-1))*csgn(I*(-(c*x^(3/2)+1)^2/(
c^2*x^3-1)-1)/(1-(c*x^(3/2)+1)^2/(c^2*x^3-1)))-csgn(I*(-(c*x^(3/2)+1)^2/(c
^2*x^3-1)-1)/(1-(c*x^(3/2)+1)^2/(c^2*x^3-1)))*csgn(I/(1-(c*x^(3/2)+1)^2/(c
^2*x^3-1)))+csgn(I*(-(c*x^(3/2)+1)^2/(c^2*x^3-1)-1)/(1-(c*x^(3/2)+1)^2/(c^
2*x^3-1)))^2)*arctanh(c*x^(3/2))^2+2*a*b*(2/3*ln(c*x^(3/2))*arctanh(c*x^(
3/2))-1/3*dilog(c*x^(3/2))-1/3*dilog(c*x^(3/2)+1)-1/3*ln(c*x^(3/2))*ln(c*x
^(3/2)+1))
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{x} dx = \int \frac{(b \operatorname{arctanh}(cx^{3/2}) + a)^2}{x} dx$$

input

```
integrate((a+b*arctanh(c*x^(3/2)))^2/x,x, algorithm="fricas")
```

output

```
integral((b^2*arctanh(c*x^(3/2))^2 + 2*a*b*arctanh(c*x^(3/2)) + a^2)/x, x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{x} dx = \text{Timed out}$$

input

```
integrate((a+b*atanh(c*x**(3/2)))**2/x,x)
```

output Timed out

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{x} dx = \int \frac{(b \operatorname{arctanh}(cx^{\frac{3}{2}}) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x^(3/2)))^2/x,x, algorithm="maxima")`

output `1/4*b^2*integrate(log(c*x^(3/2) + 1)^2/x, x) - 1/2*b^2*integrate(log(c*x^(3/2) + 1)*log(-c*x^(3/2) + 1)/x, x) + 1/4*b^2*integrate(log(-c*x^(3/2) + 1)^2/x, x) + a*b*integrate(log(c*x^(3/2) + 1)/x, x) - a*b*integrate(log(-c*x^(3/2) + 1)/x, x) + a^2*log(x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{x} dx = \int \frac{(b \operatorname{arctanh}(cx^{\frac{3}{2}}) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x^(3/2)))^2/x,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^(3/2)) + a)^2/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{x} dx = \int \frac{(a + b \operatorname{atanh}(cx^{3/2}))^2}{x} dx$$

input `int((a + b*atanh(c*x^(3/2)))^2/x,x)`output `int((a + b*atanh(c*x^(3/2)))^2/x, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{x} dx = 2 \left(\int \frac{\operatorname{atanh}(\sqrt{x} cx)}{x} dx \right) ab$$

$$+ \left(\int \frac{\operatorname{atanh}(\sqrt{x} cx)^2}{x} dx \right) b^2 + \log(x) a^2$$

input `int((a+b*atanh(c*x^(3/2)))^2/x,x)`output `2*int(atanh(sqrt(x)*c*x)/x,x)*a*b + int(atanh(sqrt(x)*c*x)**2/x,x)*b**2 + log(x)*a**2`

3.222 $\int \frac{(a+b\operatorname{arctanh}(cx^{3/2}))^2}{x^4} dx$

Optimal result	1785
Mathematica [B] (verified)	1785
Rubi [A] (verified)	1786
Maple [C] (warning: unable to verify)	1789
Fricas [B] (verification not implemented)	1790
Sympy [F(-1)]	1790
Maxima [B] (verification not implemented)	1790
Giac [F]	1791
Mupad [B] (verification not implemented)	1792
Reduce [B] (verification not implemented)	1792

Optimal result

Integrand size = 18, antiderivative size = 96

$$\int \frac{(a + b\operatorname{arctanh}(cx^{3/2}))^2}{x^4} dx = -\frac{2bc(a + b\operatorname{arctanh}(cx^{3/2}))}{3x^{3/2}} + \frac{1}{3}c^2(a + b\operatorname{arctanh}(cx^{3/2}))^2 - \frac{(a + b\operatorname{arctanh}(cx^{3/2}))^2}{3x^3} + b^2c^2 \log(x) - \frac{1}{3}b^2c^2 \log(1 - c^2x^3)$$

```
output -2/3*b*c*(a+b*arctanh(c*x^(3/2)))/x^(3/2)+1/3*c^2*(a+b*arctanh(c*x^(3/2)))^2-1/3*(a+b*arctanh(c*x^(3/2)))^2/x^3+b^2*c^2*ln(x)-1/3*b^2*c^2*ln(-c^2*x^3+1)
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 210 vs. 2(96) = 192.

Time = 0.09 (sec) , antiderivative size = 210, normalized size of antiderivative = 2.19

$$\int \frac{(a + b\operatorname{arctanh}(cx^{3/2}))^2}{x^4} dx = \frac{a^2 + 2abcx^{3/2} + 2b(a + bcx^{3/2}) \operatorname{arctanh}(cx^{3/2}) - b^2(-1 + c^2x^3) \operatorname{arctanh}(cx^{3/2})^2 + abc^2x^3 \log(1 - \sqrt[3]{c}\sqrt{x})}{x^3}$$

input `Integrate[(a + b*ArcTanh[c*x^(3/2)])^2/x^4,x]`

output
$$-1/3*(a^2 + 2*a*b*c*x^(3/2) + 2*b*(a + b*c*x^(3/2))*ArcTanh[c*x^(3/2)] - b^2*(-1 + c^2*x^3)*ArcTanh[c*x^(3/2)]^2 + a*b*c^2*x^3*Log[1 - c^(1/3)*Sqrt[x]] - a*b*c^2*x^3*Log[1 + c^(1/3)*Sqrt[x]] - 3*b^2*c^2*x^3*Log[x] - a*b*c^2*x^3*Log[1 - c^(1/3)*Sqrt[x] + c^(2/3)*x] + a*b*c^2*x^3*Log[1 + c^(1/3)*Sqrt[x] + c^(2/3)*x] + b^2*c^2*x^3*Log[1 - c^2*x^3])/x^3$$

Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.99, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6454, 6452, 6544, 6452, 243, 47, 14, 16, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{x^4} dx$$

$$\downarrow 6454$$

$$\frac{2}{3} \int \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{x^{9/2}} dx^{3/2}$$

$$\downarrow 6452$$

$$\frac{2}{3} \left(bc \int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^3(1 - c^2x^3)} dx^{3/2} - \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{2x^3} \right)$$

$$\downarrow 6544$$

$$\frac{2}{3} \left(bc \left(c^2 \int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{1 - c^2x^3} dx^{3/2} + \int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^3} dx^{3/2} \right) - \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{2x^3} \right)$$

$$\downarrow 6452$$

$$\frac{2}{3} \left(bc \left(c^2 \int \frac{a + b \operatorname{arctanh}(cx^{3/2})}{1 - c^2x^3} dx^{3/2} + bc \int \frac{1}{x^{3/2}(1 - c^2x^3)} dx^{3/2} - \frac{a + b \operatorname{arctanh}(cx^{3/2})}{x^{3/2}} \right) - \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{2x^3} \right)$$

↓ 243

$$\frac{2}{3} \left(bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^{3/2})}{1 - c^2x^3} dx^{3/2} + \frac{1}{2} bc \int \frac{1}{x^{3/2}(1 - c^2x^3)} dx^3 - \frac{a + \operatorname{barctanh}(cx^{3/2})}{x^{3/2}} \right) - \frac{(a + \operatorname{barctanh}(cx^{3/2}))^2}{2x^3} \right)$$

↓ 47

$$\frac{2}{3} \left(bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^{3/2})}{1 - c^2x^3} dx^{3/2} + \frac{1}{2} bc \left(c^2 \int \frac{1}{1 - c^2x^3} dx^3 + \int \frac{1}{x^{3/2}} dx^3 \right) - \frac{a + \operatorname{barctanh}(cx^{3/2})}{x^{3/2}} \right) - \frac{(a + \operatorname{barctanh}(cx^{3/2}))^2}{2x^3} \right)$$

↓ 14

$$\frac{2}{3} \left(bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^{3/2})}{1 - c^2x^3} dx^{3/2} + \frac{1}{2} bc \left(c^2 \int \frac{1}{1 - c^2x^3} dx^3 + \log(x^3) \right) - \frac{a + \operatorname{barctanh}(cx^{3/2})}{x^{3/2}} \right) - \frac{(a + \operatorname{barctanh}(cx^{3/2}))^2}{2x^3} \right)$$

↓ 16

$$\frac{2}{3} \left(bc \left(c^2 \int \frac{a + \operatorname{barctanh}(cx^{3/2})}{1 - c^2x^3} dx^{3/2} - \frac{a + \operatorname{barctanh}(cx^{3/2})}{x^{3/2}} + \frac{1}{2} bc (\log(x^3) - \log(1 - c^2x^3)) \right) - \frac{(a + \operatorname{barctanh}(cx^{3/2}))^2}{2x^3} \right)$$

↓ 6510

$$\frac{2}{3} \left(bc \left(\frac{c(a + \operatorname{barctanh}(cx^{3/2}))^2}{2b} - \frac{a + \operatorname{barctanh}(cx^{3/2})}{x^{3/2}} + \frac{1}{2} bc (\log(x^3) - \log(1 - c^2x^3)) \right) - \frac{(a + \operatorname{barctanh}(cx^{3/2}))^2}{2x^3} \right)$$

input `Int[(a + b*ArcTanh[c*x^(3/2)])^2/x^4, x]`

output `(2*(-1/2*(a + b*ArcTanh[c*x^(3/2)])^2/x^3 + b*c*(-((a + b*ArcTanh[c*x^(3/2)])/x^(3/2)) + (c*(a + b*ArcTanh[c*x^(3/2)])^2)/(2*b) + (b*c*(Log[x^3] - Log[1 - c^2*x^3]))/2))/3`

Definitions of rubi rules used

- rule 14 $\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ ; FreeQ}[a, x]$
- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$
- rule 47 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x]$
- rule 243 $\text{Int}[(x_)^{(m_)}*((a_)+(b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] \text{ ; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$
- rule 6452 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{2n})}), x], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$
- rule 6454 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n] - 1)}*(a + b*\text{ArcTanh}[c*x])^p, x], x, x^n], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 1] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$
- rule 6510 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)]*(b_))^{(p_)}((d_)+(e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^{(p+1)}/(b*c*d*(p+1)), x] \text{ ; FreeQ}[\{a, b, c, d, e, p\}, x] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{NeQ}[p, -1]$

rule 6544

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*((f_.)*(x_))^(m_)/((d_) + (
e_.)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x
], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x
^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 2.24 (sec) , antiderivative size = 3062, normalized size of antiderivative = 31.90

method	result	size
parts	Expression too large to display	3062
derivativedivides	Expression too large to display	3063
default	Expression too large to display	3063

input

```
int((a+b*arctanh(c*x^(3/2)))^2/x^4,x,method=_RETURNVERBOSE)
```

output

```
-1/3*a^2/x^3+b^2*(-1/3/x^3*arctanh(c*x^(3/2))^2+2*c*(1/6*arctanh(c*x^(3/2)
)*c*ln(c*x^(3/2)+1)-1/3*arctanh(c*x^(3/2))/x^(3/2)-1/6*arctanh(c*x^(3/2))*
c*ln(c*x^(3/2)-1)-1/2*c*(c*(Sum(-1/6*(ln(x^(1/2))-_alpha)*ln(c*x^(3/2)-1)-3
*c*(1/3*ln(x^(1/2))-_alpha)*(ln(1/2*(x^(1/2))+_alpha)/_alpha)+ln((RootOf(_Z^
2+_Z*_alpha+_alpha^2,index=1)-x^(1/2)+_alpha)/RootOf(_Z^2+_Z*_alpha+_alpha
^2,index=1))+ln((RootOf(_Z^2+_Z*_alpha+_alpha^2,index=2)-x^(1/2)+_alpha)/R
ootOf(_Z^2+_Z*_alpha+_alpha^2,index=2)))/c+1/3*(dilog(1/2*(x^(1/2)+_alpha)
/_alpha)+dilog((RootOf(_Z^2+_Z*_alpha+_alpha^2,index=1)-x^(1/2)+_alpha)/Ro
otOf(_Z^2+_Z*_alpha+_alpha^2,index=1))+dilog((RootOf(_Z^2+_Z*_alpha+_alpha
^2,index=2)-x^(1/2)+_alpha)/RootOf(_Z^2+_Z*_alpha+_alpha^2,index=2)))/c)/
c,_alpha=RootOf(_Z^3*c+1))+Sum(1/6*(ln(x^(1/2))-_alpha)*ln(c*x^(3/2)-1)-3*c
*(1/6/_alpha^2/c*ln(x^(1/2))-_alpha)^2-1/3*_alpha*ln(x^(1/2))-_alpha)*(2*Ro
otOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1)*RootOf(_Z^2+3*_Z*_alpha+3*_alpha^
2,index=2)*ln((RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1)-x^(1/2)+_alpha)
/RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1))+2*RootOf(_Z^2+3*_Z*_alpha+3*
_alpha^2,index=1)*RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=2)*ln((RootOf(_
Z^2+3*_Z*_alpha+3*_alpha^2,index=2)-x^(1/2)+_alpha)/RootOf(_Z^2+3*_Z*_alph
a+3*_alpha^2,index=2))+3*RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1)*_alph
a*ln((RootOf(_Z^2+3*_Z*_alpha+3*_alpha^2,index=1)-x^(1/2)+_alpha)/RootOf(_
Z^2+3*_Z*_alpha+3*_alpha^2,index=1))+6*RootOf(_Z^2+3*_Z*_alpha+3*_alpha...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 173 vs. $2(80) = 160$.

Time = 0.11 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.80

$$\int \frac{(a + \operatorname{arctanh}(cx^{3/2}))^2}{x^4} dx = \frac{24b^2c^2x^3 \log(\sqrt{x}) + 4(ab - b^2)c^2x^3 \log(cx^{\frac{3}{2}} + 1) - 4(ab + b^2)c^2x^3 \log(\dots)}{x^4}$$

input `integrate((a+b*arctanh(c*x^(3/2)))^2/x^4,x, algorithm="fricas")`

output `1/12*(24*b^2*c^2*x^3*log(sqrt(x)) + 4*(a*b - b^2)*c^2*x^3*log(c*x^(3/2) + 1) - 4*(a*b + b^2)*c^2*x^3*log(c*x^(3/2) - 1) - 8*a*b*c*x^(3/2) + (b^2*c^2*x^3 - b^2)*log(-(c^2*x^3 + 2*c*x^(3/2) + 1)/(c^2*x^3 - 1))^2 - 4*a^2 - 4*(b^2*c*x^(3/2) + a*b)*log(-(c^2*x^3 + 2*c*x^(3/2) + 1)/(c^2*x^3 - 1)))/x^3`

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + \operatorname{arctanh}(cx^{3/2}))^2}{x^4} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**(3/2)))**2/x**4,x)`

output `Timed out`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 175 vs. $2(80) = 160$.

Time = 0.04 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.82

$$\int \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{x^4} dx = \frac{1}{3} \left(\left(c \log(cx^{3/2} + 1) - c \log(cx^{3/2} - 1) - \frac{2}{x^{3/2}} \right) c - \frac{2 \operatorname{artanh}(cx^{3/2})}{x^3} \right) ab$$

$$+ \frac{1}{12} \left(\left(2 \left(\log(cx^{3/2} - 1) - 2 \right) \log(cx^{3/2} + 1) - \log(cx^{3/2} + 1)^2 - \log(cx^{3/2} - 1)^2 - 4 \log(cx^{3/2} - 1) + 12 \log(x) \right) c^2 \right.$$

$$\left. - \frac{b^2 \operatorname{artanh}(cx^{3/2})^2}{3x^3} - \frac{a^2}{3x^3} \right)$$

input `integrate((a+b*arctanh(c*x^(3/2)))^2/x^4,x, algorithm="maxima")`

output

```
1/3*((c*log(c*x^(3/2) + 1) - c*log(c*x^(3/2) - 1) - 2/x^(3/2))*c - 2*arctanh(c*x^(3/2))/x^3)*a*b + 1/12*((2*(log(c*x^(3/2) - 1) - 2)*log(c*x^(3/2) + 1) - log(c*x^(3/2) + 1)^2 - log(c*x^(3/2) - 1)^2 - 4*log(c*x^(3/2) - 1) + 12*log(x))*c^2 + 4*(c*log(c*x^(3/2) + 1) - c*log(c*x^(3/2) - 1) - 2/x^(3/2))*c*arctanh(c*x^(3/2))*b^2 - 1/3*b^2*arctanh(c*x^(3/2))^2/x^3 - 1/3*a^2/x^3
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{x^4} dx = \int \frac{(b \operatorname{artanh}(cx^{3/2}) + a)^2}{x^4} dx$$

input `integrate((a+b*arctanh(c*x^(3/2)))^2/x^4,x, algorithm="giac")`

output

```
integrate((b*arctanh(c*x^(3/2)) + a)^2/x^4, x)
```

Mupad [B] (verification not implemented)

Time = 4.69 (sec) , antiderivative size = 281, normalized size of antiderivative = 2.93

$$\int \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{x^4} dx = \frac{2b^2 c^2 \ln(x^{3/2})}{3} - \frac{a^2}{3x^3} - \frac{b^2 c^2 \ln(cx^{3/2} - 1)}{3}$$

$$- \frac{b^2 c^2 \ln(cx^{3/2} + 1)}{3} + \frac{b^2 c^2 \ln(cx^{3/2} + 1)^2}{12} + \frac{b^2 c^2 \ln(1 - cx^{3/2})^2}{12}$$

$$- \frac{b^2 \ln(cx^{3/2} + 1)^2}{12x^3} - \frac{b^2 \ln(1 - cx^{3/2})^2}{12x^3} - \frac{abc^2 \ln(cx^{3/2} - 1)}{3}$$

$$+ \frac{abc^2 \ln(cx^{3/2} + 1)}{3} - \frac{2abc}{3x^{3/2}} - \frac{ab \ln(cx^{3/2} + 1)}{3x^3} + \frac{ab \ln(1 - cx^{3/2})}{3x^3}$$

$$- \frac{b^2 c^2 \ln(cx^{3/2} + 1) \ln(1 - cx^{3/2})}{6} - \frac{b^2 c \ln(cx^{3/2} + 1)}{3x^{3/2}}$$

$$+ \frac{b^2 c \ln(1 - cx^{3/2})}{3x^{3/2}} + \frac{b^2 \ln(cx^{3/2} + 1) \ln(1 - cx^{3/2})}{6x^3}$$

input `int((a + b*atanh(c*x^(3/2)))^2/x^4,x)`output

```
(2*b^2*c^2*log(x^(3/2)))/3 - a^2/(3*x^3) - (b^2*c^2*log(c*x^(3/2) - 1))/3
- (b^2*c^2*log(c*x^(3/2) + 1))/3 + (b^2*c^2*log(c*x^(3/2) + 1)^2)/12 + (b^
2*c^2*log(1 - c*x^(3/2))^2)/12 - (b^2*log(c*x^(3/2) + 1)^2)/(12*x^3) - (b^
2*log(1 - c*x^(3/2))^2)/(12*x^3) - (a*b*c^2*log(c*x^(3/2) - 1))/3 + (a*b*c
^2*log(c*x^(3/2) + 1))/3 - (2*a*b*c)/(3*x^(3/2)) - (a*b*log(c*x^(3/2) + 1)
)/(3*x^3) + (a*b*log(1 - c*x^(3/2)))/(3*x^3) - (b^2*c^2*log(c*x^(3/2) + 1)
*log(1 - c*x^(3/2)))/6 - (b^2*c*log(c*x^(3/2) + 1))/(3*x^(3/2)) + (b^2*c*log(1 - c*x^(3/2)))/(3*x^(3/2)) + (b^2*log(c*x^(3/2) + 1)*log(1 - c*x^(3/2)))/(6*x^3)
```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.77

$$\int \frac{(a + b \operatorname{arctanh}(cx^{3/2}))^2}{x^4} dx = \frac{\operatorname{atanh}(\sqrt{x} cx)^2 b^2 c^2 x^3 - \operatorname{atanh}(\sqrt{x} cx)^2 b^2 - 2\sqrt{x} \operatorname{atanh}(\sqrt{x} cx) b^2 cx + 2a \operatorname{atanh}(\sqrt{x} cx) b^2 cx - 2a^2}{x^4}$$

input `int((a+b*atanh(c*x^(3/2)))^2/x^4,x)`

output

```
(atanh(sqrt(x)*c*x)**2*b**2*c**2*x**3 - atanh(sqrt(x)*c*x)**2*b**2 - 2*sqrt(x)*atanh(sqrt(x)*c*x)*b**2*c*x + 2*atanh(sqrt(x)*c*x)*a*b*c**2*x**3 - 2*atanh(sqrt(x)*c*x)*a*b + 2*atanh(sqrt(x)*c*x)*b**2*c**2*x**3 - 2*sqrt(x)*a*b*c*x - 2*log(c**(2/3)*x - sqrt(x)*c**(1/3) + 1)*b**2*c**2*x**3 - 2*log(sqrt(x)*c**(2/3) + c**(1/3))*b**2*c**2*x**3 + 6*log(sqrt(x))*b**2*c**2*x**3 - a**2)/(3*x**3)
```

3.223 $\int x^2(a + \operatorname{barctanh}(cx^n)) dx$

Optimal result	1794
Mathematica [A] (verified)	1794
Rubi [A] (verified)	1795
Maple [F]	1796
Fricas [F]	1796
Sympy [F]	1797
Maxima [F]	1797
Giac [F]	1797
Mupad [F(-1)]	1798
Reduce [F]	1798

Optimal result

Integrand size = 14, antiderivative size = 66

$$\int x^2(a + \operatorname{barctanh}(cx^n)) dx = \frac{1}{3}x^3(a + \operatorname{barctanh}(cx^n)) - \frac{bcnx^{3+n} \operatorname{Hypergeometric2F1}\left(1, \frac{3+n}{2n}, \frac{3(1+n)}{2n}, c^2x^{2n}\right)}{3(3+n)}$$

output

```
1/3*x^3*(a+b*arctanh(c*x^n))-b*c*n*x^(3+n)*hypergeom([1, 1/2*(3+n)/n],[3/2*(1+n)/n],c^2*x^(2*n))/(9+3*n)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.11

$$\int x^2(a + \operatorname{barctanh}(cx^n)) dx = \frac{ax^3}{3} + \frac{1}{3}bx^3\operatorname{arctanh}(cx^n) - \frac{bcnx^{3+n} \operatorname{Hypergeometric2F1}\left(1, \frac{3+n}{2n}, 1 + \frac{3+n}{2n}, c^2x^{2n}\right)}{3(3+n)}$$

input

```
Integrate[x^2*(a + b*ArcTanh[c*x^n]),x]
```

output

$$\frac{(a*x^3)/3 + (b*x^3*ArcTanh[c*x^n])/3 - (b*c*n*x^(3 + n)*Hypergeometric2F1[1, (3 + n)/(2*n), 1 + (3 + n)/(2*n), c^2*x^(2*n)])/(3*(3 + n))}{1}$$
Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6452, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2(a + b \operatorname{arctanh}(cx^n)) dx$$

$$\downarrow 6452$$

$$\frac{1}{3}x^3(a + b \operatorname{arctanh}(cx^n)) - \frac{1}{3}bcn \int \frac{x^{n+2}}{1 - c^2x^{2n}} dx$$

$$\downarrow 888$$

$$\frac{1}{3}x^3(a + b \operatorname{arctanh}(cx^n)) - \frac{bcn x^{n+3} \operatorname{Hypergeometric2F1}\left(1, \frac{n+3}{2n}, \frac{3(n+1)}{2n}, c^2x^{2n}\right)}{3(n+3)}$$

input

$$\text{Int}[x^2*(a + b*ArcTanh[c*x^n]), x]$$

output

$$\frac{(x^3*(a + b*ArcTanh[c*x^n]))/3 - (b*c*n*x^(3 + n)*Hypergeometric2F1[1, (3 + n)/(2*n), (3*(1 + n))/(2*n), c^2*x^(2*n)])/(3*(3 + n))}{1}$$

Definitions of rubi rules used

rule 888 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [F]

$$\int x^2(a + b \operatorname{arctanh}(cx^n)) dx$$

input `int(x^2*(a+b*arctanh(c*x^n)),x)`

output `int(x^2*(a+b*arctanh(c*x^n)),x)`

Fricas [F]

$$\int x^2(a + b \operatorname{arctanh}(cx^n)) dx = \int (b \operatorname{arctanh}(cx^n) + a)x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c*x^n)),x, algorithm="fricas")`

output `integral(b*x^2*arctanh(c*x^n) + a*x^2, x)`

Sympy [F]

$$\int x^2(a + b \operatorname{arctanh}(cx^n)) dx = \int x^2(a + b \operatorname{atanh}(cx^n)) dx$$

input `integrate(x**2*(a+b*atanh(c*x**n)),x)`

output `Integral(x**2*(a + b*atanh(c*x**n)), x)`

Maxima [F]

$$\int x^2(a + b \operatorname{arctanh}(cx^n)) dx = \int (b \operatorname{atanh}(cx^n) + a)x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c*x^n)),x, algorithm="maxima")`

output `1/3*a*x^3 + 1/6*(x^3*log(c*x^n + 1) - x^3*log(-c*x^n + 1) + 3*n*integrate(1/3*x^2/(c*x^n + 1), x) + 3*n*integrate(1/3*x^2/(c*x^n - 1), x))*b`

Giac [F]

$$\int x^2(a + b \operatorname{arctanh}(cx^n)) dx = \int (b \operatorname{atanh}(cx^n) + a)x^2 dx$$

input `integrate(x^2*(a+b*arctanh(c*x^n)),x, algorithm="giac")`

output `integrate((b*arctanh(c*x^n) + a)*x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^2(a + b \operatorname{arctanh}(cx^n)) dx = \int x^2(a + b \operatorname{atanh}(cx^n)) dx$$

input `int(x^2*(a + b*atanh(c*x^n)),x)`output `int(x^2*(a + b*atanh(c*x^n)), x)`**Reduce [F]**

$$\int x^2(a + b \operatorname{arctanh}(cx^n)) dx = \left(\int \operatorname{atanh}(x^n c) x^2 dx \right) b + \frac{a x^3}{3}$$

input `int(x^2*(a+b*atanh(c*x^n)),x)`output `(3*int(atanh(x**n*c)*x**2,x)*b + a*x**3)/3`

3.224 $\int x(a + \operatorname{barctanh}(cx^n)) dx$

Optimal result	1799
Mathematica [A] (verified)	1799
Rubi [A] (verified)	1800
Maple [F]	1801
Fricas [F]	1801
Sympy [F]	1801
Maxima [F]	1802
Giac [F]	1802
Mupad [F(-1)]	1802
Reduce [F]	1803

Optimal result

Integrand size = 12, antiderivative size = 67

$$\int x(a + \operatorname{barctanh}(cx^n)) dx = \frac{1}{2}x^2(a + \operatorname{barctanh}(cx^n)) - \frac{bcnx^{2+n} \operatorname{Hypergeometric2F1}\left(1, \frac{2+n}{2n}, \frac{1}{2}\left(3 + \frac{2}{n}\right), c^2x^{2n}\right)}{2(2+n)}$$

output

```
1/2*x^2*(a+b*arctanh(c*x^n))-b*c*n*x^(2+n)*hypergeom([1, 1/2*(2+n)/n], [3/2+1/n], c^2*x^(2*n))/(4+2*n)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.09

$$\int x(a + \operatorname{barctanh}(cx^n)) dx = \frac{ax^2}{2} + \frac{1}{2}bx^2\operatorname{arctanh}(cx^n) - \frac{bcnx^{2+n} \operatorname{Hypergeometric2F1}\left(1, \frac{2+n}{2n}, 1 + \frac{2+n}{2n}, c^2x^{2n}\right)}{2(2+n)}$$

input

```
Integrate[x*(a + b*ArcTanh[c*x^n]), x]
```

output

```
(a*x^2)/2 + (b*x^2*ArcTanh[c*x^n])/2 - (b*c*n*x^(2 + n)*Hypergeometric2F1[
1, (2 + n)/(2*n), 1 + (2 + n)/(2*n), c^2*x^(2*n)])/(2*(2 + n))
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6452, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(a + b \operatorname{arctanh}(cx^n)) dx$$

$$\downarrow 6452$$

$$\frac{1}{2}x^2(a + b \operatorname{arctanh}(cx^n)) - \frac{1}{2}bcn \int \frac{x^{n+1}}{1 - c^2x^{2n}} dx$$

$$\downarrow 888$$

$$\frac{1}{2}x^2(a + b \operatorname{arctanh}(cx^n)) - \frac{bcn x^{n+2} \operatorname{Hypergeometric2F1}\left(1, \frac{n+2}{2n}, \frac{1}{2}\left(3 + \frac{2}{n}\right), c^2x^{2n}\right)}{2(n+2)}$$

input

```
Int[x*(a + b*ArcTanh[c*x^n]),x]
```

output

```
(x^2*(a + b*ArcTanh[c*x^n]))/2 - (b*c*n*x^(2 + n)*Hypergeometric2F1[1, (2
+ n)/(2*n), (3 + 2/n)/2, c^2*x^(2*n)])/(2*(2 + n))
```

Defintions of rubi rules used

rule 888

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p
*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1
, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILt
Q[p, 0] || GtQ[a, 0])
```

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [F]

$$\int x(a + b \operatorname{arctanh}(cx^n)) dx$$

input `int(x*(a+b*arctanh(c*x^n)),x)`

output `int(x*(a+b*arctanh(c*x^n)),x)`

Fricas [F]

$$\int x(a + b \operatorname{arctanh}(cx^n)) dx = \int (b \operatorname{artanh}(cx^n) + a)x dx$$

input `integrate(x*(a+b*arctanh(c*x^n)),x, algorithm="fricas")`

output `integral(b*x*arctanh(c*x^n) + a*x, x)`

Sympy [F]

$$\int x(a + b \operatorname{arctanh}(cx^n)) dx = \int x(a + b \operatorname{atanh}(cx^n)) dx$$

input `integrate(x*(a+b*atanh(c*x**n)),x)`

output `Integral(x*(a + b*atanh(c*x**n)), x)`

Maxima [F]

$$\int x(a + b \operatorname{arctanh}(cx^n)) dx = \int (b \operatorname{artanh}(cx^n) + a)x dx$$

input `integrate(x*(a+b*arctanh(c*x^n)),x, algorithm="maxima")`

output `1/2*a*x^2 + 1/4*(x^2*log(c*x^n + 1) - x^2*log(-c*x^n + 1) + 2*n*integrate(1/2*x/(c*x^n + 1), x) + 2*n*integrate(1/2*x/(c*x^n - 1), x))*b`

Giac [F]

$$\int x(a + b \operatorname{arctanh}(cx^n)) dx = \int (b \operatorname{artanh}(cx^n) + a)x dx$$

input `integrate(x*(a+b*arctanh(c*x^n)),x, algorithm="giac")`

output `integrate((b*arctanh(c*x^n) + a)*x, x)`

Mupad [F(-1)]

Timed out.

$$\int x(a + b \operatorname{arctanh}(cx^n)) dx = \int x(a + b \operatorname{atanh}(cx^n)) dx$$

input `int(x*(a + b*atanh(c*x^n)),x)`

output `int(x*(a + b*atanh(c*x^n)), x)`

Reduce [F]

$$\int x(a + b \operatorname{arctanh}(cx^n)) dx = \left(\int \operatorname{atanh}(x^n c) x dx \right) b + \frac{ax^2}{2}$$

input `int(x*(a+b*atanh(c*x^n)),x)`

output `(2*int(atanh(x**n*c)*x,x)*b + a*x**2)/2`

3.225 $\int (a + b \operatorname{arctanh}(cx^n)) dx$

Optimal result	1804
Mathematica [A] (verified)	1804
Rubi [A] (verified)	1805
Maple [F]	1806
Fricas [F]	1806
Sympy [F]	1806
Maxima [F]	1807
Giac [F]	1807
Mupad [F(-1)]	1807
Reduce [F]	1808

Optimal result

Integrand size = 10, antiderivative size = 58

$$\int (a + b \operatorname{arctanh}(cx^n)) dx = ax + b \operatorname{arctanh}(cx^n) - \frac{bcn x^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}\left(3 + \frac{1}{n}\right), c^2 x^{2n}\right)}{1+n}$$

output

```
a*x+b*x*arctanh(c*x^n)-b*c*n*x^(1+n)*hypergeom([1, 1/2*(1+n)/n], [3/2+1/2/n], c^2*x^(2*n))/(1+n)
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\int (a + b \operatorname{arctanh}(cx^n)) dx = ax + b \operatorname{arctanh}(cx^n) - \frac{bcn x^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}\left(3 + \frac{1}{n}\right), c^2 x^{2n}\right)}{1+n}$$

input

```
Integrate[a + b*ArcTanh[c*x^n], x]
```

output

$$a*x + b*x*ArcTanh[c*x^n] - (b*c*n*x^{(1+n)}*Hypergeometric2F1[1, (1+n)/(2*n), (3+n^{-1})/2, c^2*x^{(2*n)}])/(1+n)$$
Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{arctanh}(cx^n)) dx$$

↓ 2009

$$ax + b \operatorname{arctanh}(cx^n) - \frac{bcn x^{n+1} \operatorname{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}\left(3 + \frac{1}{n}\right), c^2 x^{2n}\right)}{n+1}$$

input

$$\text{Int}[a + b*ArcTanh[c*x^n], x]$$

output

$$a*x + b*x*ArcTanh[c*x^n] - (b*c*n*x^{(1+n)}*Hypergeometric2F1[1, (1+n)/(2*n), (3+n^{-1})/2, c^2*x^{(2*n)}])/(1+n)$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [F]

$$\int (a + b \operatorname{arctanh}(cx^n)) dx$$

input `int(a+b*arctanh(c*x^n),x)`

output `int(a+b*arctanh(c*x^n),x)`

Fricas [F]

$$\int (a + b \operatorname{arctanh}(cx^n)) dx = \int b \operatorname{artanh}(cx^n) + a dx$$

input `integrate(a+b*arctanh(c*x^n),x, algorithm="fricas")`

output `integral(b*arctanh(c*x^n) + a, x)`

Sympy [F]

$$\int (a + b \operatorname{arctanh}(cx^n)) dx = \int (a + b \operatorname{atanh}(cx^n)) dx$$

input `integrate(a+b*atanh(c*x**n),x)`

output `Integral(a + b*atanh(c*x**n), x)`

Maxima [F]

$$\int (a + b \operatorname{arctanh}(cx^n)) dx = \int b \operatorname{artanh}(cx^n) + a dx$$

input `integrate(a+b*arctanh(c*x^n),x, algorithm="maxima")`

output `1/2*(n*integrate(1/(c*x^n + 1), x) + n*integrate(1/(c*x^n - 1), x) + x*log(c*x^n + 1) - x*log(-c*x^n + 1))*b + a*x`

Giac [F]

$$\int (a + b \operatorname{arctanh}(cx^n)) dx = \int b \operatorname{artanh}(cx^n) + a dx$$

input `integrate(a+b*arctanh(c*x^n),x, algorithm="giac")`

output `integrate(b*arctanh(c*x^n) + a, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + b \operatorname{arctanh}(cx^n)) dx = \int a + b \operatorname{atanh}(cx^n) dx$$

input `int(a + b*atanh(c*x^n),x)`

output `int(a + b*atanh(c*x^n), x)`

Reduce [F]

$$\int (a + b \operatorname{arctanh}(cx^n)) dx = \left(\int \operatorname{atanh}(x^n c) dx \right) b + ax$$

input `int(a+b*atanh(c*x^n),x)`

output `int(atanh(x**n*c),x)*b + a*x`

3.226 $\int \frac{a+b\operatorname{arctanh}(cx^n)}{x} dx$

Optimal result	1809
Mathematica [C] (verified)	1809
Rubi [A] (verified)	1810
Maple [A] (verified)	1811
Fricas [B] (verification not implemented)	1811
Sympy [F]	1812
Maxima [F]	1812
Giac [F]	1812
Mupad [F(-1)]	1813
Reduce [F]	1813

Optimal result

Integrand size = 14, antiderivative size = 36

$$\int \frac{a + b\operatorname{arctanh}(cx^n)}{x} dx = a \log(x) - \frac{b \operatorname{PolyLog}(2, -cx^n)}{2n} + \frac{b \operatorname{PolyLog}(2, cx^n)}{2n}$$

output

$a*\ln(x)-1/2*b*polylog(2,-c*x^n)/n+1/2*b*polylog(2,c*x^n)/n$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.07 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.08

$$\int \frac{a + b\operatorname{arctanh}(cx^n)}{x} dx = \frac{bcx^n {}_3F_2\left(\frac{1}{2}, \frac{1}{2}, 1; \frac{3}{2}, \frac{3}{2}; c^2x^{2n}\right)}{n} + a \log(x)$$

input

`Integrate[(a + b*ArcTanh[c*x^n])/x,x]`

output

$(b*c*x^n*HypergeometricPFQ[\{1/2, 1/2, 1\}, \{3/2, 3/2\}, c^2*x^(2*n)])/n + a*Log[x]$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6450, 6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x} dx$$

↓ 6450

$$\int \frac{x^{-n}(a + b \operatorname{arctanh}(cx^n)) dx^n}{n}$$

↓ 6446

$$\frac{a \log(x^n) - \frac{1}{2}b \operatorname{PolyLog}(2, -cx^n) + \frac{1}{2}b \operatorname{PolyLog}(2, cx^n)}{n}$$

input `Int[(a + b*ArcTanh[c*x^n])/x,x]`

output `(a*Log[x^n] - (b*PolyLog[2, -(c*x^n)])/2 + (b*PolyLog[2, c*x^n])/2)/n`

Defintions of rubi rules used

rule 6446 `Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))/(x_), x_Symbol] := Simp[a*Log[x], x] + (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) / ; FreeQ[{a, b, c}, x]`

rule 6450 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)]*(b_.))^(p_.)/(x_), x_Symbol] := Simp[1/n Subst[Int[(a + b*ArcTanh[c*x])^p/x, x], x, x^n], x] / ; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 0.68 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.97

method	result	size
risch	$a \ln(x) + \frac{b \operatorname{dilog}(1-cx^n)}{2n} - \frac{b \operatorname{dilog}(cx^n+1)}{2n}$	35
parts	$a \ln(x) + \frac{b \left(\ln(cx^n) \operatorname{arctanh}(cx^n) - \frac{\operatorname{dilog}(cx^n+1)}{2} - \frac{\ln(cx^n) \ln(cx^n+1)}{2} - \frac{\operatorname{dilog}(cx^n)}{2} \right)}{n}$	59
derivativedivides	$\frac{a \ln(cx^n) + b \left(\ln(cx^n) \operatorname{arctanh}(cx^n) - \frac{\operatorname{dilog}(cx^n+1)}{2} - \frac{\ln(cx^n) \ln(cx^n+1)}{2} - \frac{\operatorname{dilog}(cx^n)}{2} \right)}{n}$	64
default	$\frac{a \ln(cx^n) + b \left(\ln(cx^n) \operatorname{arctanh}(cx^n) - \frac{\operatorname{dilog}(cx^n+1)}{2} - \frac{\ln(cx^n) \ln(cx^n+1)}{2} - \frac{\operatorname{dilog}(cx^n)}{2} \right)}{n}$	64

input `int((a+b*arctanh(c*x^n))/x,x,method=_RETURNVERBOSE)`

output `a*ln(x)+1/2*b/n*dilog(1-c*x^n)-1/2/n*b*dilog(c*x^n+1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(30) = 60.

Time = 0.09 (sec) , antiderivative size = 141, normalized size of antiderivative = 3.92

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x} dx = \frac{bn \log(c \cosh(n \log(x)) + c \sinh(n \log(x)) + 1) \log(x) - bn \log(-c \cosh(n \log(x)) - c \sinh(n \log(x)) - 1) \log(x) - 2a n \log(x) - b \operatorname{dilog}(c \cosh(n \log(x)) + c \sinh(n \log(x))) + b \operatorname{dilog}(-c \cosh(n \log(x)) - c \sinh(n \log(x)))}{n}$$

input `integrate((a+b*arctanh(c*x^n))/x,x, algorithm="fricas")`

output `-1/2*(b*n*log(c*cosh(n*log(x)) + c*sinh(n*log(x)) + 1)*log(x) - b*n*log(-c*cosh(n*log(x)) - c*sinh(n*log(x)) + 1)*log(x) - b*n*log(x)*log(-(c*cosh(n*log(x)) + c*sinh(n*log(x)) + 1)/(c*cosh(n*log(x)) + c*sinh(n*log(x)) - 1)) - 2*a*n*log(x) - b*dilog(c*cosh(n*log(x)) + c*sinh(n*log(x)))) + b*dilog(-c*cosh(n*log(x)) - c*sinh(n*log(x))))/n`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x} dx = \int \frac{a + b \operatorname{atanh}(cx^n)}{x} dx$$

input `integrate((a+b*atanh(c*x**n))/x,x)`

output `Integral((a + b*atanh(c*x**n))/x, x)`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x} dx = \int \frac{b \operatorname{artanh}(cx^n) + a}{x} dx$$

input `integrate((a+b*arctanh(c*x^n))/x,x, algorithm="maxima")`

output `1/2*(n*integrate(log(x)/(c*x*x^n + x), x) + n*integrate(log(x)/(c*x*x^n - x), x) + log(c*x^n + 1)*log(x) - log(-c*x^n + 1)*log(x))*b + a*log(x)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x} dx = \int \frac{b \operatorname{artanh}(cx^n) + a}{x} dx$$

input `integrate((a+b*arctanh(c*x^n))/x,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^n) + a)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x} dx = \int \frac{a + b \operatorname{atanh}(cx^n)}{x} dx$$

input `int((a + b*atanh(c*x^n))/x,x)`output `int((a + b*atanh(c*x^n))/x, x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x} dx = \left(\int \frac{\operatorname{atanh}(x^n c)}{x} dx \right) b + \log(x) a$$

input `int((a+b*atanh(c*x^n))/x,x)`output `int(atanh(x**n*c)/x,x)*b + log(x)*a`

3.227 $\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^2} dx$

Optimal result	1814
Mathematica [A] (verified)	1814
Rubi [A] (verified)	1815
Maple [F]	1816
Fricas [F]	1816
Sympy [F]	1816
Maxima [F]	1817
Giac [F]	1817
Mupad [F(-1)]	1817
Reduce [F]	1818

Optimal result

Integrand size = 14, antiderivative size = 67

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^2} dx = -\frac{a + b \operatorname{arctanh}(cx^n)}{x} - \frac{bcn x^{-1+n} \operatorname{Hypergeometric2F1}\left(1, -\frac{1-n}{2n}, \frac{1}{2}\left(3 - \frac{1}{n}\right), c^2 x^{2n}\right)}{1-n}$$

output

$-(a+b*\operatorname{arctanh}(c*x^n))/x-b*c*n*x^{(-1+n)}*\operatorname{hypergeom}([1, -1/2*(1-n)/n], [3/2-1/2/n], c^2*x^{(2*n)})/(1-n)$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.99

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^2} dx = -\frac{a}{x} - \frac{b \operatorname{arctanh}(cx^n)}{x} + \frac{bcn x^{-1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{-1+n}{2n}, 1 + \frac{-1+n}{2n}, c^2 x^{2n}\right)}{-1+n}$$

input

$\operatorname{Integrate}[(a + b*\operatorname{ArcTanh}[c*x^n])/x^2, x]$

output

$$-(a/x) - (b \operatorname{ArcTanh}[c x^n])/x + (b c n x^{-(1+n)} \operatorname{Hypergeometric2F1}[1, (-1+n)/(2n), 1 + (-1+n)/(2n), c^2 x^{2n}])/(-1+n)$$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6452, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}(cx^n)}{x^2} dx \\ & \quad \downarrow \text{6452} \\ & bcn \int \frac{x^{n-2}}{1 - c^2 x^{2n}} dx - \frac{a + b \operatorname{arctanh}(cx^n)}{x} \\ & \quad \downarrow \text{888} \\ & -\frac{a + b \operatorname{arctanh}(cx^n)}{x} - \frac{bcn x^{n-1} \operatorname{Hypergeometric2F1}\left(1, -\frac{1-n}{2n}, \frac{1}{2}\left(3 - \frac{1}{n}\right), c^2 x^{2n}\right)}{1-n} \end{aligned}$$

input

$$\operatorname{Int}[(a + b \operatorname{ArcTanh}[c x^n])/x^2, x]$$

output

$$-((a + b \operatorname{ArcTanh}[c x^n])/x) - (b c n x^{-(1+n)} \operatorname{Hypergeometric2F1}[1, -1/2*(1-n)/n, (3-n)/2, c^2 x^{2n}])/(1-n)$$

Defintions of rubi rules used

rule 888

$$\operatorname{Int}[(c x)^m (a + b (x^n)^p), x_Symbol] \rightarrow \operatorname{Simp}[a^p * ((c x)^{m+1} / (c^{m+1})) * \operatorname{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n + 1, (-b)(x^n/a)], x] /; \operatorname{FreeQ}[\{a, b, c, m, n, p\}, x] \&\& !\operatorname{IGtQ}[p, 0] \&\& (\operatorname{ILtQ}[p, 0] \mid \mid \operatorname{GtQ}[a, 0])$$

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^2} dx$$

input

```
int((a+b*arctanh(c*x^n))/x^2,x)
```

output

```
int((a+b*arctanh(c*x^n))/x^2,x)
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^2} dx = \int \frac{b \operatorname{arctanh}(cx^n) + a}{x^2} dx$$

input

```
integrate((a+b*arctanh(c*x^n))/x^2,x, algorithm="fricas")
```

output

```
integral((b*arctanh(c*x^n) + a)/x^2, x)
```

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^2} dx = \int \frac{a + b \operatorname{atanh}(cx^n)}{x^2} dx$$

input

```
integrate((a+b*atanh(c*x**n))/x**2,x)
```

output

```
Integral((a + b*atanh(c*x**n))/x**2, x)
```

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^2} dx = \int \frac{b \operatorname{artanh}(cx^n) + a}{x^2} dx$$

input `integrate((a+b*arctanh(c*x^n))/x^2,x, algorithm="maxima")`

output `-1/2*(n*integrate(1/(c*x^2*x^n + x^2), x) + n*integrate(1/(c*x^2*x^n - x^2), x) + (log(c*x^n + 1) - log(-c*x^n + 1))/x)*b - a/x`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^2} dx = \int \frac{b \operatorname{artanh}(cx^n) + a}{x^2} dx$$

input `integrate((a+b*arctanh(c*x^n))/x^2,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^n) + a)/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^2} dx = \int \frac{a + b \operatorname{atanh}(cx^n)}{x^2} dx$$

input `int((a + b*atanh(c*x^n))/x^2,x)`

output `int((a + b*atanh(c*x^n))/x^2, x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^2} dx = \frac{\left(\int \frac{\operatorname{atanh}(x^n c)}{x^2} dx \right) bx - a}{x}$$

input `int((a+b*atanh(c*x^n))/x^2,x)`

output `(int(atanh(x**n*c)/x**2,x)*b*x - a)/x`

3.228 $\int \frac{a+b\operatorname{arctanh}(cx^n)}{x^3} dx$

Optimal result	1819
Mathematica [A] (verified)	1819
Rubi [A] (verified)	1820
Maple [F]	1821
Fricas [F]	1821
Sympy [F(-1)]	1821
Maxima [F]	1822
Giac [F]	1822
Mupad [F(-1)]	1822
Reduce [F]	1823

Optimal result

Integrand size = 14, antiderivative size = 70

$$\int \frac{a + b\operatorname{arctanh}(cx^n)}{x^3} dx = -\frac{a + b\operatorname{arctanh}(cx^n)}{2x^2} - \frac{bcnx^{-2+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}\left(1 - \frac{2}{n}\right), \frac{1}{2}\left(3 - \frac{2}{n}\right), c^2x^{2n}\right)}{2(2 - n)}$$

output

$$-1/2*(a+b*\operatorname{arctanh}(c*x^n))/x^2-b*c*n*x^{(-2+n)}*\operatorname{hypergeom}([1, 1/2-1/n], [3/2-1/n], c^2*x^{(2*n)})/(4-2*n)$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.04

$$\int \frac{a + b\operatorname{arctanh}(cx^n)}{x^3} dx = -\frac{a}{2x^2} - \frac{b\operatorname{arctanh}(cx^n)}{2x^2} + \frac{bcnx^{-2+n} \operatorname{Hypergeometric2F1}\left(1, \frac{-2+n}{2n}, 1 + \frac{-2+n}{2n}, c^2x^{2n}\right)}{2(-2 + n)}$$

input

$$\operatorname{Integrate}[(a + b*\operatorname{ArcTanh}[c*x^n])/x^3, x]$$

output

$$-1/2*a/x^2 - (b*ArcTanh[c*x^n])/(2*x^2) + (b*c*n*x^{(-2 + n)}*Hypergeometric2F1[1, (-2 + n)/(2*n), 1 + (-2 + n)/(2*n), c^2*x^{(2*n)}])/(2*(-2 + n))$$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.01, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6452, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^3} dx$$

↓ 6452

$$\frac{1}{2}bcn \int \frac{x^{n-3}}{1 - c^2x^{2n}} dx - \frac{a + b \operatorname{arctanh}(cx^n)}{2x^2}$$

↓ 888

$$-\frac{a + b \operatorname{arctanh}(cx^n)}{2x^2} - \frac{bcn x^{n-2} \operatorname{Hypergeometric2F1}\left(1, -\frac{2-n}{2n}, \frac{1}{2}\left(3 - \frac{2}{n}\right), c^2x^{2n}\right)}{2(2-n)}$$

input

```
Int[(a + b*ArcTanh[c*x^n])/x^3,x]
```

output

$$-1/2*(a + b*ArcTanh[c*x^n])/x^2 - (b*c*n*x^{(-2 + n)}*Hypergeometric2F1[1, -1/2*(2 - n)/n, (3 - 2/n)/2, c^2*x^{(2*n)}])/(2*(2 - n))$$

Defintions of rubi rules used

rule 888

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])
```

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]
```

Maple [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^3} dx$$

input

```
int((a+b*arctanh(c*x^n))/x^3,x)
```

output

```
int((a+b*arctanh(c*x^n))/x^3,x)
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^3} dx = \int \frac{b \operatorname{arctanh}(cx^n) + a}{x^3} dx$$

input

```
integrate((a+b*arctanh(c*x^n))/x^3,x, algorithm="fricas")
```

output

```
integral((b*arctanh(c*x^n) + a)/x^3, x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^3} dx = \text{Timed out}$$

input

```
integrate((a+b*atanh(c*x**n))/x**3,x)
```

output Timed out

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^3} dx = \int \frac{b \operatorname{artanh}(cx^n) + a}{x^3} dx$$

input `integrate((a+b*arctanh(c*x^n))/x^3,x, algorithm="maxima")`

output `-1/4*(2*n*integrate(1/2/(c*x^3*x^n + x^3), x) + 2*n*integrate(1/2/(c*x^3*x^n - x^3), x) + (log(c*x^n + 1) - log(-c*x^n + 1))/x^2)*b - 1/2*a/x^2`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^3} dx = \int \frac{b \operatorname{artanh}(cx^n) + a}{x^3} dx$$

input `integrate((a+b*arctanh(c*x^n))/x^3,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^n) + a)/x^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^3} dx = \int \frac{a + b \operatorname{atanh}(cx^n)}{x^3} dx$$

input `int((a + b*atanh(c*x^n))/x^3,x)`

output `int((a + b*atanh(c*x^n))/x^3, x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^3} dx = \frac{2 \left(\int \frac{\operatorname{atanh}(x^n c)}{x^3} dx \right) b x^2 - a}{2x^2}$$

input `int((a+b*atanh(c*x^n))/x^3,x)`

output `(2*int(atanh(x**n*c)/x**3,x)*b*x**2 - a)/(2*x**2)`

3.229 $\int \frac{a+b\operatorname{arctanh}(cx^n)}{x^4} dx$

Optimal result	1824
Mathematica [A] (verified)	1824
Rubi [A] (verified)	1825
Maple [F]	1826
Fricas [F]	1826
Sympy [F]	1827
Maxima [F]	1827
Giac [F]	1827
Mupad [F(-1)]	1828
Reduce [F]	1828

Optimal result

Integrand size = 14, antiderivative size = 72

$$\int \frac{a + b\operatorname{arctanh}(cx^n)}{x^4} dx = -\frac{a + b\operatorname{arctanh}(cx^n)}{3x^3} - \frac{bcnx^{-3+n} \operatorname{Hypergeometric2F1}\left(1, -\frac{3-n}{2n}, -\frac{3(1-n)}{2n}, c^2x^{2n}\right)}{3(3-n)}$$

output `-1/3*(a+b*arctanh(c*x^n))/x^3-b*c*n*x^(-3+n)*hypergeom([1, -1/2*(3-n)/n], [1/2*(-3+3*n)/n], c^2*x^(2*n))/(9-3*n)`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.01

$$\int \frac{a + b\operatorname{arctanh}(cx^n)}{x^4} dx = -\frac{a}{3x^3} - \frac{b\operatorname{arctanh}(cx^n)}{3x^3} + \frac{bcnx^{-3+n} \operatorname{Hypergeometric2F1}\left(1, \frac{-3+n}{2n}, 1 + \frac{-3+n}{2n}, c^2x^{2n}\right)}{3(-3+n)}$$

input `Integrate[(a + b*ArcTanh[c*x^n])/x^4, x]`

output

$$-1/3*a/x^3 - (b*ArcTanh[c*x^n])/(3*x^3) + (b*c*n*x^{(-3 + n)}*Hypergeometric2F1[1, (-3 + n)/(2*n), 1 + (-3 + n)/(2*n), c^2*x^{(2*n)}])/(3*(-3 + n))$$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6452, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^4} dx$$

$$\downarrow 6452$$

$$\frac{1}{3}bcn \int \frac{x^{n-4}}{1 - c^2x^{2n}} dx - \frac{a + b \operatorname{arctanh}(cx^n)}{3x^3}$$

$$\downarrow 888$$

$$\frac{a + b \operatorname{arctanh}(cx^n)}{3x^3} - \frac{bcn x^{n-3} \operatorname{Hypergeometric2F1}\left(1, -\frac{3-n}{2n}, -\frac{3(1-n)}{2n}, c^2x^{2n}\right)}{3(3-n)}$$

input

$$\operatorname{Int}[(a + b*ArcTanh[c*x^n])/x^4, x]$$

output

$$-1/3*(a + b*ArcTanh[c*x^n])/x^3 - (b*c*n*x^{(-3 + n)}*Hypergeometric2F1[1, -1/2*(3 - n)/n, (-3*(1 - n))/(2*n), c^2*x^{(2*n)}])/(3*(3 - n))$$

Definitions of rubi rules used

rule 888 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

Maple [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^4} dx$$

input `int((a+b*arctanh(c*x^n))/x^4,x)`

output `int((a+b*arctanh(c*x^n))/x^4,x)`

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^4} dx = \int \frac{b \operatorname{arctanh}(cx^n) + a}{x^4} dx$$

input `integrate((a+b*arctanh(c*x^n))/x^4,x, algorithm="fricas")`

output `integral((b*arctanh(c*x^n) + a)/x^4, x)`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^4} dx = \int \frac{a + b \operatorname{atanh}(cx^n)}{x^4} dx$$

input `integrate((a+b*atanh(c*x**n))/x**4,x)`

output `Integral((a + b*atanh(c*x**n))/x**4, x)`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^4} dx = \int \frac{b \operatorname{artanh}(cx^n) + a}{x^4} dx$$

input `integrate((a+b*arctanh(c*x^n))/x^4,x, algorithm="maxima")`

output `-1/6*(3*n*integrate(1/3/(c*x^4*x^n + x^4), x) + 3*n*integrate(1/3/(c*x^4*x^n - x^4), x) + (log(c*x^n + 1) - log(-c*x^n + 1))/x^3)*b - 1/3*a/x^3`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^4} dx = \int \frac{b \operatorname{artanh}(cx^n) + a}{x^4} dx$$

input `integrate((a+b*arctanh(c*x^n))/x^4,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^n) + a)/x^4, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^4} dx = \int \frac{a + b \operatorname{atanh}(cx^n)}{x^4} dx$$

input `int((a + b*atanh(c*x^n))/x^4,x)`output `int((a + b*atanh(c*x^n))/x^4, x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{arctanh}(cx^n)}{x^4} dx = \frac{3 \left(\int \frac{\operatorname{atanh}(x^n c)}{x^4} dx \right) b x^3 - a}{3x^3}$$

input `int((a+b*atanh(c*x^n))/x^4,x)`output `(3*int(atanh(x**n*c)/x**4,x)*b*x**3 - a)/(3*x**3)`

3.230 $\int x(a + b \operatorname{arctanh}(cx^n))^2 dx$

Optimal result	1829
Mathematica [N/A]	1829
Rubi [N/A]	1830
Maple [N/A]	1830
Fricas [N/A]	1831
Sympy [N/A]	1831
Maxima [N/A]	1831
Giac [N/A]	1832
Mupad [N/A]	1832
Reduce [N/A]	1833

Optimal result

Integrand size = 14, antiderivative size = 14

$$\int x(a + b \operatorname{arctanh}(cx^n))^2 dx = \operatorname{Int}(x(a + b \operatorname{arctanh}(cx^n))^2, x)$$

output `Defer(Int)(x*(a+b*arctanh(c*x^n))^2,x)`

Mathematica [N/A]

Not integrable

Time = 12.17 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int x(a + b \operatorname{arctanh}(cx^n))^2 dx = \int x(a + b \operatorname{arctanh}(cx^n))^2 dx$$

input `Integrate[x*(a + b*ArcTanh[c*x^n])^2,x]`

output `Integrate[x*(a + b*ArcTanh[c*x^n])^2, x]`

Rubi [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(a + \text{barctanh}(cx^n))^2 dx$$

↓ 6468

$$\int x(a + \text{barctanh}(cx^n))^2 dx$$

input `Int[x*(a + b*ArcTanh[c*x^n])^2,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int x(a + b \operatorname{arctanh}(cx^n))^2 dx$$

input `int(x*(a+b*arctanh(c*x^n))^2,x)`

output `int(x*(a+b*arctanh(c*x^n))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 32, normalized size of antiderivative = 2.29

$$\int x(a + b \operatorname{arctanh}(cx^n))^2 dx = \int (b \operatorname{artanh}(cx^n) + a)^2 x dx$$

input `integrate(x*(a+b*arctanh(c*x^n))^2,x, algorithm="fricas")`

output `integral(b^2*x*arctanh(c*x^n)^2 + 2*a*b*x*arctanh(c*x^n) + a^2*x, x)`

Sympy [N/A]

Not integrable

Time = 30.21 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int x(a + b \operatorname{arctanh}(cx^n))^2 dx = \int x(a + b \operatorname{atanh}(cx^n))^2 dx$$

input `integrate(x*(a+b*atanh(c*x**n))**2,x)`

output `Integral(x*(a + b*atanh(c*x**n))**2, x)`

Maxima [N/A]

Not integrable

Time = 1.29 (sec) , antiderivative size = 155, normalized size of antiderivative = 11.07

$$\int x(a + b \operatorname{arctanh}(cx^n))^2 dx = \int (b \operatorname{artanh}(cx^n) + a)^2 x dx$$

input `integrate(x*(a+b*arctanh(c*x^n))^2,x, algorithm="maxima")`

output

```
1/8*b^2*x^2*log(-c*x^n + 1)^2 + 1/2*a^2*x^2 - integrate(-1/4*((b^2*c*x*x^n
- b^2*x)*log(c*x^n + 1)^2 + 4*(a*b*c*x*x^n - a*b*x)*log(c*x^n + 1) + (4*a
*b*x - (b^2*c*n + 4*a*b*c)*x*x^n - 2*(b^2*c*x*x^n - b^2*x)*log(c*x^n + 1))
*log(-c*x^n + 1))/(c*x^n - 1), x)
```

Giac [N/A]

Not integrable

Time = 0.41 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int x(a + b \operatorname{arctanh}(cx^n))^2 dx = \int (b \operatorname{artanh}(cx^n) + a)^2 x dx$$

input

```
integrate(x*(a+b*arctanh(c*x^n))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^n) + a)^2*x, x)
```

Mupad [N/A]

Not integrable

Time = 3.46 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int x(a + b \operatorname{arctanh}(cx^n))^2 dx = \int x(a + b \operatorname{atanh}(cx^n))^2 dx$$

input

```
int(x*(a + b*atanh(c*x^n))^2,x)
```

output

```
int(x*(a + b*atanh(c*x^n))^2, x)
```

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.79

$$\int x(a + b \operatorname{arctanh}(cx^n))^2 dx = 2 \left(\int \operatorname{atanh}(x^n c) x dx \right) ab + \left(\int \operatorname{atanh}(x^n c)^2 x dx \right) b^2 + \frac{a^2 x^2}{2}$$

input `int(x*(a+b*atanh(c*x^n))^2,x)`output `(4*int(atanh(x**n*c)*x,x)*a*b + 2*int(atanh(x**n*c)**2*x,x)*b**2 + a**2*x**2)/2`

3.231 $\int (a + b \operatorname{arctanh}(cx^n))^2 dx$

Optimal result	1834
Mathematica [N/A]	1834
Rubi [N/A]	1835
Maple [N/A]	1835
Fricas [N/A]	1836
Sympy [N/A]	1836
Maxima [N/A]	1836
Giac [N/A]	1837
Mupad [N/A]	1837
Reduce [N/A]	1838

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int (a + b \operatorname{arctanh}(cx^n))^2 dx = \operatorname{Int}((a + b \operatorname{arctanh}(cx^n))^2, x)$$

output `Defer(Int)((a+b*arctanh(c*x^n))^2,x)`

Mathematica [N/A]

Not integrable

Time = 1.55 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int (a + b \operatorname{arctanh}(cx^n))^2 dx = \int (a + b \operatorname{arctanh}(cx^n))^2 dx$$

input `Integrate[(a + b*ArcTanh[c*x^n])^2,x]`

output `Integrate[(a + b*ArcTanh[c*x^n])^2, x]`

Rubi [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{arctanh}(cx^n))^2 dx$$

↓ 6444

$$\int (a + b \operatorname{arctanh}(cx^n))^2 dx$$

input `Int[(a + b*ArcTanh[c*x^n])^2,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int (a + b \operatorname{arctanh}(cx^n))^2 dx$$

input `int((a+b*arctanh(c*x^n))^2,x)`

output `int((a+b*arctanh(c*x^n))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 28, normalized size of antiderivative = 2.33

$$\int (a + b \operatorname{arctanh}(cx^n))^2 dx = \int (b \operatorname{artanh}(cx^n) + a)^2 dx$$

input `integrate((a+b*arctanh(c*x^n))^2,x, algorithm="fricas")`

output `integral(b^2*arctanh(c*x^n)^2 + 2*a*b*arctanh(c*x^n) + a^2, x)`

Sympy [N/A]

Not integrable

Time = 18.78 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int (a + b \operatorname{arctanh}(cx^n))^2 dx = \int (a + b \operatorname{atanh}(cx^n))^2 dx$$

input `integrate((a+b*atanh(c*x**n))**2,x)`

output `Integral((a + b*atanh(c*x**n))**2, x)`

Maxima [N/A]

Not integrable

Time = 0.40 (sec) , antiderivative size = 143, normalized size of antiderivative = 11.92

$$\int (a + b \operatorname{arctanh}(cx^n))^2 dx = \int (b \operatorname{artanh}(cx^n) + a)^2 dx$$

input `integrate((a+b*arctanh(c*x^n))^2,x, algorithm="maxima")`

output

```
1/4*b^2*x*log(-c*x^n + 1)^2 + a^2*x - integrate(-1/4*((b^2*c*x^n - b^2)*log(c*x^n + 1)^2 + 4*(a*b*c*x^n - a*b)*log(c*x^n + 1) + 2*(2*a*b - (b^2*c*n + 2*a*b*c))*x^n - (b^2*c*x^n - b^2)*log(c*x^n + 1))*log(-c*x^n + 1)/(c*x^n - 1), x)
```

Giac [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int (a + b \operatorname{arctanh}(cx^n))^2 dx = \int (b \operatorname{artanh}(cx^n) + a)^2 dx$$

input

```
integrate((a+b*arctanh(c*x^n))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^n) + a)^2, x)
```

Mupad [N/A]

Not integrable

Time = 3.31 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int (a + b \operatorname{arctanh}(cx^n))^2 dx = \int (a + b \operatorname{atanh}(cx^n))^2 dx$$

input

```
int((a + b*atanh(c*x^n))^2,x)
```

output

```
int((a + b*atanh(c*x^n))^2, x)
```

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 32, normalized size of antiderivative = 2.67

$$\int (a + b \operatorname{arctanh}(cx^n))^2 dx = 2 \left(\int \operatorname{atanh}(x^n c) dx \right) ab + \left(\int \operatorname{atanh}(x^n c)^2 dx \right) b^2 + a^2 x$$

input

```
int((a+b*atanh(c*x^n))^2,x)
```

output

```
2*int(atanh(x**n*c),x)*a*b + int(atanh(x**n*c)**2,x)*b**2 + a**2*x
```

3.232
$$\int \frac{(a+b\operatorname{arctanh}(cx^n))^2}{x} dx$$

Optimal result	1839
Mathematica [C] (verified)	1840
Rubi [A] (verified)	1840
Maple [C] (warning: unable to verify)	1842
Fricas [F]	1844
Sympy [F]	1844
Maxima [F]	1844
Giac [F]	1845
Mupad [F(-1)]	1845
Reduce [F]	1845

Optimal result

Integrand size = 16, antiderivative size = 148

$$\int \frac{(a + b\operatorname{arctanh}(cx^n))^2}{x} dx = \frac{2(a + b\operatorname{arctanh}(cx^n))^2 \operatorname{arctanh}(1 - \frac{2}{1-cx^n})}{n} - \frac{b(a + b\operatorname{arctanh}(cx^n)) \operatorname{PolyLog}(2, 1 - \frac{2}{1-cx^n})}{n} + \frac{b(a + b\operatorname{arctanh}(cx^n)) \operatorname{PolyLog}(2, -1 + \frac{2}{1-cx^n})}{n} + \frac{b^2 \operatorname{PolyLog}(3, 1 - \frac{2}{1-cx^n})}{2n} - \frac{b^2 \operatorname{PolyLog}(3, -1 + \frac{2}{1-cx^n})}{2n}$$

output

```
-2*(a+b*arctanh(c*x^n))^2*arctanh(-1+2/(1-c*x^n))/n-b*(a+b*arctanh(c*x^n))
*polylog(2,1-2/(1-c*x^n))/n+b*(a+b*arctanh(c*x^n))*polylog(2,-1+2/(1-c*x^n
))/n+1/2*b^2*polylog(3,1-2/(1-c*x^n))/n-1/2*b^2*polylog(3,-1+2/(1-c*x^n))/
n
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.32 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.22

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x} dx = a^2 \log(x) + \frac{ab(-\operatorname{PolyLog}(2, -cx^n) + \operatorname{PolyLog}(2, cx^n))}{n} + \frac{b^2 \left(\frac{i\pi^3}{24} - \frac{2}{3} \operatorname{arctanh}(cx^n)^3 - \operatorname{arctanh}(cx^n)^2 \log(1 + e^{-2\operatorname{arctanh}(cx^n)}) + \operatorname{arctanh}(cx^n)^2 \log(1 - e^{2\operatorname{arctanh}(cx^n)}) \right)}{n}$$

input

```
Integrate[(a + b*ArcTanh[c*x^n])^2/x, x]
```

output

```
a^2*Log[x] + (a*b*(-PolyLog[2, -(c*x^n)] + PolyLog[2, c*x^n]))/n + (b^2*((I/24)*Pi^3 - (2*ArcTanh[c*x^n]^3)/3 - ArcTanh[c*x^n]^2*Log[1 + E^(-2*ArcTanh[c*x^n])] + ArcTanh[c*x^n]^2*Log[1 - E^(2*ArcTanh[c*x^n])] + ArcTanh[c*x^n]*PolyLog[2, -E^(-2*ArcTanh[c*x^n])] + ArcTanh[c*x^n]*PolyLog[2, E^(2*ArcTanh[c*x^n])] + PolyLog[3, -E^(-2*ArcTanh[c*x^n])]/2 - PolyLog[3, E^(2*ArcTanh[c*x^n])]/2))/n
```

Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.10, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6450, 6448, 6614, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x} dx$$

↓ 6450

$$\int \frac{x^{-n} (a + b \operatorname{arctanh}(cx^n))^2}{n} dx^n$$

↓ 6448

$$\frac{2\operatorname{arctanh}\left(1 - \frac{2}{1-cx^n}\right) (a + b\operatorname{arctanh}(cx^n))^2 - 4bc \int \frac{(a+b\operatorname{arctanh}(cx^n))\operatorname{arctanh}\left(1 - \frac{2}{1-cx^n}\right)}{1-c^2x^{2n}} dx^n}{n}$$

↓ 6614

$$2\operatorname{arctanh}\left(1 - \frac{2}{1-cx^n}\right) (a + b\operatorname{arctanh}(cx^n))^2 - 4bc \left(\frac{1}{2} \int \frac{(a+b\operatorname{arctanh}(cx^n)) \log\left(2 - \frac{2}{1-cx^n}\right)}{1-c^2x^{2n}} dx^n - \frac{1}{2} \int \frac{(a+b\operatorname{arctanh}(cx^n))}{1-c^2x^{2n}} dx^n \right)$$

↓ 6620

$$2\operatorname{arctanh}\left(1 - \frac{2}{1-cx^n}\right) (a + b\operatorname{arctanh}(cx^n))^2 - 4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^n}\right) (a+b\operatorname{arctanh}(cx^n))}{2c} - \frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^n}\right)}{1-c^2x^{2n}} dx^n \right) \right)$$

↓ 7164

$$2\operatorname{arctanh}\left(1 - \frac{2}{1-cx^n}\right) (a + b\operatorname{arctanh}(cx^n))^2 - 4bc \left(\frac{1}{2} \left(\frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^n}\right) (a+b\operatorname{arctanh}(cx^n))}{2c} - \frac{b \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx^n}\right)}{4c} \right) \right)$$

input `Int[(a + b*ArcTanh[c*x^n])^2/x, x]`

output `(2*(a + b*ArcTanh[c*x^n])^2*ArcTanh[1 - 2/(1 - c*x^n)] - 4*b*c*(((a + b*ArcTanh[c*x^n])*PolyLog[2, 1 - 2/(1 - c*x^n)])/(2*c) - (b*PolyLog[3, 1 - 2/(1 - c*x^n)]/(4*c))/2 + (-1/2*((a + b*ArcTanh[c*x^n])*PolyLog[2, -1 + 2/(1 - c*x^n)])/c + (b*PolyLog[3, -1 + 2/(1 - c*x^n)]/(4*c))/2))/n`

Defintions of rubi rules used

rule 6448

```
Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^p/(x_), x_Symbol] := Simp[2*(a + b*ArcTanh[c*x])^p*ArcTanh[1 - 2/(1 - c*x)], x] - Simp[2*b*c*p Int[(a + b*ArcTanh[c*x])^(p - 1)*(ArcTanh[1 - 2/(1 - c*x)]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1]
```

rule 6450 $\text{Int}[(a_.) + \text{ArcTanh}[(c_.)(x_)^{(n)}](b_.)]^{(p_.)}/(x_), x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[(a + b*\text{ArcTanh}[c*x])^p/x, x], x, x^n], x] /; \text{FreeQ}[\{a, b, c, n\}, x] \ \&\& \ \text{IGtQ}[p, 0]$

rule 6614 $\text{Int}[(\text{ArcTanh}[u_]*((a_.) + \text{ArcTanh}[(c_.)(x_)](b_.))^{(p_.)})/((d_) + (e_.)(x_)^2), x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Int}[\text{Log}[1 + u]*((a + b*\text{ArcTanh}[c*x])^p/(d + e*x^2)), x], x] - \text{Simp}[1/2 \text{ Int}[\text{Log}[1 - u]*((a + b*\text{ArcTanh}[c*x])^p/(d + e*x^2)), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{EqQ}[u^2 - (1 - 2/(1 - c*x))^2, 0]$

rule 6620 $\text{Int}[(\text{Log}[u_]*((a_.) + \text{ArcTanh}[(c_.)(x_)](b_.))^{(p_.)})/((d_) + (e_.)(x_)^2), x_Symbol] \rightarrow \text{Simp}[(-a + b*\text{ArcTanh}[c*x])^p*(\text{PolyLog}[2, 1 - u]/(2*c*d)), x] + \text{Simp}[b*(p/2) \text{ Int}[(a + b*\text{ArcTanh}[c*x])^{(p-1)}*(\text{PolyLog}[2, 1 - u]/(d + e*x^2)), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{EqQ}[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]$

rule 7164 $\text{Int}[(u_)*\text{PolyLog}[n_, v_], x_Symbol] \rightarrow \text{With}[\{w = \text{DerivativeDivides}[v, u*v, x]\}, \text{Simp}[w*\text{PolyLog}[n + 1, v], x] /; \text{!FalseQ}[w]] /; \text{FreeQ}[n, x]$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 13.48 (sec) , antiderivative size = 750, normalized size of antiderivative = 5.07

method	result
parts	$a^2 \ln(x) + \frac{b^2 \left(\ln(cx^n) \operatorname{arctanh}(cx^n)^2 - \operatorname{arctanh}(cx^n) \operatorname{polylog}\left(2, -\frac{(cx^n+1)^2}{1-c^2x^{2n}}\right) + \frac{\operatorname{polylog}\left(3, -\frac{(cx^n+1)^2}{1-c^2x^{2n}}\right)}{2} - \operatorname{arctanh}(cx^n) \right)}{1}$
derivativedivides	$a^2 \ln(cx^n) + b^2 \frac{\left(\ln(cx^n) \operatorname{arctanh}(cx^n)^2 - \operatorname{arctanh}(cx^n) \operatorname{polylog}\left(2, -\frac{(cx^n+1)^2}{1-c^2x^{2n}}\right) + \frac{\operatorname{polylog}\left(3, -\frac{(cx^n+1)^2}{1-c^2x^{2n}}\right)}{2} - \operatorname{arctanh}(cx^n) \right)}{1}$
default	$a^2 \ln(cx^n) + b^2 \frac{\left(\ln(cx^n) \operatorname{arctanh}(cx^n)^2 - \operatorname{arctanh}(cx^n) \operatorname{polylog}\left(2, -\frac{(cx^n+1)^2}{1-c^2x^{2n}}\right) + \frac{\operatorname{polylog}\left(3, -\frac{(cx^n+1)^2}{1-c^2x^{2n}}\right)}{2} - \operatorname{arctanh}(cx^n) \right)}{1}$

```
input int((a+b*arctanh(c*x^n))^2/x,x,method=_RETURNVERBOSE)
```

```
output a^2*ln(x)+b^2/n*(ln(c*x^n)*arctanh(c*x^n)^2-arctanh(c*x^n)*polylog(2,-(c*x^n+1)^2/(1-c^2*(x^n)^2))+1/2*polylog(3,-(c*x^n+1)^2/(1-c^2*(x^n)^2))-arctanh(c*x^n)^2*ln((c*x^n+1)^2/(1-c^2*(x^n)^2)-1)+arctanh(c*x^n)^2*ln(1-(c*x^n+1)/(1-c^2*(x^n)^2)^(1/2))+2*arctanh(c*x^n)*polylog(2,(c*x^n+1)/(1-c^2*(x^n)^2)^(1/2))-2*polylog(3,(c*x^n+1)/(1-c^2*(x^n)^2)^(1/2))+arctanh(c*x^n)^2*ln(1+(c*x^n+1)/(1-c^2*(x^n)^2)^(1/2))+2*arctanh(c*x^n)*polylog(2,-(c*x^n+1)/(1-c^2*(x^n)^2)^(1/2))-2*polylog(3,-(c*x^n+1)/(1-c^2*(x^n)^2)^(1/2))+1/2*I*Pi*csgn(I*(-(c*x^n+1)^2/(c^2*(x^n)^2-1)-1)/(1-(c*x^n+1)^2/(c^2*(x^n)^2-1)))*(csgn(I*(-(c*x^n+1)^2/(c^2*(x^n)^2-1)-1))*csgn(I/(1-(c*x^n+1)^2/(c^2*(x^n)^2-1)-1)/(1-(c*x^n+1)^2/(c^2*(x^n)^2-1))))-csgn(I*(-(c*x^n+1)^2/(c^2*(x^n)^2-1)-1))*csgn(I*(-(c*x^n+1)^2/(c^2*(x^n)^2-1)-1)/(1-(c*x^n+1)^2/(c^2*(x^n)^2-1))))-csgn(I*(-(c*x^n+1)^2/(c^2*(x^n)^2-1)-1)/(1-(c*x^n+1)^2/(c^2*(x^n)^2-1))))*csgn(I/(1-(c*x^n+1)^2/(c^2*(x^n)^2-1)-1)/(1-(c*x^n+1)^2/(c^2*(x^n)^2-1))))+csgn(I*(-(c*x^n+1)^2/(c^2*(x^n)^2-1)-1)/(1-(c*x^n+1)^2/(c^2*(x^n)^2-1))))^2)*arctanh(c*x^n)^2)+2*a*b/n*(ln(c*x^n)*arctanh(c*x^n)-1/2*dilog(c*x^n+1)-1/2*ln(c*x^n)*ln(c*x^n+1)-1/2*dilog(c*x^n))
```


Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x} dx = \int \frac{(b \operatorname{artanh}(cx^n) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x^n))^2/x,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x^n)^2 + 2*a*b*arctanh(c*x^n) + a^2)/x, x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x} dx = \int \frac{(a + b \operatorname{atanh}(cx^n))^2}{x} dx$$

input `integrate((a+b*atanh(c*x**n))**2/x,x)`

output `Integral((a + b*atanh(c*x**n))**2/x, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x} dx = \int \frac{(b \operatorname{artanh}(cx^n) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x^n))^2/x,x, algorithm="maxima")`

output `1/4*b^2*log(-c*x^n + 1)^2*log(x) + a^2*log(x) - integrate(-1/4*((b^2*c*x^n - b^2)*log(c*x^n + 1)^2 + 4*(a*b*c*x^n - a*b)*log(c*x^n + 1) + 2*(2*a*b - (b^2*c*n*log(x) + 2*a*b*c))*x^n - (b^2*c*x^n - b^2)*log(c*x^n + 1))*log(-c*x^n + 1)/(c*x*x^n - x), x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x} dx = \int \frac{(b \operatorname{artanh}(cx^n) + a)^2}{x} dx$$

input `integrate((a+b*arctanh(c*x^n))^2/x,x, algorithm="giac")`

output `integrate((b*arctanh(c*x^n) + a)^2/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x} dx = \int \frac{(a + b \operatorname{atanh}(cx^n))^2}{x} dx$$

input `int((a + b*atanh(c*x^n))^2/x,x)`

output `int((a + b*atanh(c*x^n))^2/x, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x} dx = 2 \left(\int \frac{\operatorname{atanh}(x^n c)}{x} dx \right) ab + \left(\int \frac{\operatorname{atanh}(x^n c)^2}{x} dx \right) b^2 + \log(x) a^2$$

input `int((a+b*atanh(c*x^n))^2/x,x)`

output `2*int(atanh(x**n*c)/x,x)*a*b + int(atanh(x**n*c)**2/x,x)*b**2 + log(x)*a**2`

$$3.233 \quad \int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^2} dx$$

Optimal result	1846
Mathematica [N/A]	1846
Rubi [N/A]	1847
Maple [N/A]	1847
Fricas [N/A]	1848
Sympy [N/A]	1848
Maxima [N/A]	1848
Giac [N/A]	1849
Mupad [N/A]	1849
Reduce [N/A]	1850

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^2} dx = \operatorname{Int}\left(\frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^2}, x\right)$$

output `Defer(Int)((a+b*arctanh(c*x^n))^2/x^2,x)`

Mathematica [N/A]

Not integrable

Time = 12.21 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^2} dx = \int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^2} dx$$

input `Integrate[(a + b*ArcTanh[c*x^n])^2/x^2,x]`

output `Integrate[(a + b*ArcTanh[c*x^n])^2/x^2, x]`

Rubi [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^2} dx$$

↓ 6468

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^2} dx$$

input `Int[(a + b*ArcTanh[c*x^n])^2/x^2,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^2} dx$$

input `int((a+b*arctanh(c*x^n))^2/x^2,x)`

output `int((a+b*arctanh(c*x^n))^2/x^2,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 32, normalized size of antiderivative = 2.00

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^2} dx = \int \frac{(b \operatorname{artanh}(cx^n) + a)^2}{x^2} dx$$

input `integrate((a+b*arctanh(c*x^n))^2/x^2,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x^n)^2 + 2*a*b*arctanh(c*x^n) + a^2)/x^2, x)`

Sympy [N/A]

Not integrable

Time = 19.73 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^2} dx = \int \frac{(a + b \operatorname{atanh}(cx^n))^2}{x^2} dx$$

input `integrate((a+b*atanh(c*x**n))**2/x**2,x)`

output `Integral((a + b*atanh(c*x**n))**2/x**2, x)`

Maxima [N/A]

Not integrable

Time = 0.51 (sec) , antiderivative size = 154, normalized size of antiderivative = 9.62

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^2} dx = \int \frac{(b \operatorname{artanh}(cx^n) + a)^2}{x^2} dx$$

input `integrate((a+b*arctanh(c*x^n))^2/x^2,x, algorithm="maxima")`

output

```
-1/4*b^2*log(-c*x^n + 1)^2/x - a^2/x - integrate(-1/4*((b^2*c*x^n - b^2)*log(c*x^n + 1)^2 + 4*(a*b*c*x^n - a*b)*log(c*x^n + 1) + 2*(2*a*b + (b^2*c*n - 2*a*b*c)*x^n - (b^2*c*x^n - b^2)*log(c*x^n + 1))*log(-c*x^n + 1)/(c*x^2*x^n - x^2), x)
```

Giac [N/A]

Not integrable

Time = 0.41 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^2} dx = \int \frac{(b \operatorname{artanh}(cx^n) + a)^2}{x^2} dx$$

input

```
integrate((a+b*arctanh(c*x^n))^2/x^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^n) + a)^2/x^2, x)
```

Mupad [N/A]

Not integrable

Time = 3.39 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^2} dx = \int \frac{(a + b \operatorname{atanh}(cx^n))^2}{x^2} dx$$

input

```
int((a + b*atanh(c*x^n))^2/x^2,x)
```

output

```
int((a + b*atanh(c*x^n))^2/x^2, x)
```

Reduce [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 46, normalized size of antiderivative = 2.88

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^2} dx = \frac{2 \left(\int \frac{\operatorname{atanh}(x^n c)}{x^2} dx \right) abx + \left(\int \frac{\operatorname{atanh}(x^n c)^2}{x^2} dx \right) b^2 x - a^2}{x}$$

input `int((a+b*atanh(c*x^n))^2/x^2,x)`output `(2*int(atanh(x**n*c)/x**2,x)*a*b*x + int(atanh(x**n*c)**2/x**2,x)*b**2*x - a**2)/x`

$$3.234 \quad \int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^3} dx$$

Optimal result	1851
Mathematica [N/A]	1851
Rubi [N/A]	1852
Maple [N/A]	1852
Fricas [N/A]	1853
Sympy [N/A]	1853
Maxima [N/A]	1853
Giac [N/A]	1854
Mupad [N/A]	1854
Reduce [N/A]	1855

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^3} dx = \operatorname{Int}\left(\frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^3}, x\right)$$

output `Defer(Int)((a+b*arctanh(c*x^n))^2/x^3,x)`

Mathematica [N/A]

Not integrable

Time = 12.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^3} dx = \int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^3} dx$$

input `Integrate[(a + b*ArcTanh[c*x^n])^2/x^3,x]`

output `Integrate[(a + b*ArcTanh[c*x^n])^2/x^3, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^3} dx$$

↓ 6468

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^3} dx$$

input `Int[(a + b*ArcTanh[c*x^n])^2/x^3,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^3} dx$$

input `int((a+b*arctanh(c*x^n))^2/x^3,x)`

output `int((a+b*arctanh(c*x^n))^2/x^3,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 32, normalized size of antiderivative = 2.00

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^3} dx = \int \frac{(b \operatorname{artanh}(cx^n) + a)^2}{x^3} dx$$

input `integrate((a+b*arctanh(c*x^n))^2/x^3,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x^n)^2 + 2*a*b*arctanh(c*x^n) + a^2)/x^3, x)`

Sympy [N/A]

Not integrable

Time = 43.17 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^3} dx = \int \frac{(a + b \operatorname{atanh}(cx^n))^2}{x^3} dx$$

input `integrate((a+b*atanh(c*x**n))**2/x**3,x)`

output `Integral((a + b*atanh(c*x**n))**2/x**3, x)`

Maxima [N/A]

Not integrable

Time = 0.50 (sec) , antiderivative size = 153, normalized size of antiderivative = 9.56

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^3} dx = \int \frac{(b \operatorname{artanh}(cx^n) + a)^2}{x^3} dx$$

input `integrate((a+b*arctanh(c*x^n))^2/x^3,x, algorithm="maxima")`

output

```
-1/8*b^2*log(-c*x^n + 1)^2/x^2 - 1/2*a^2/x^2 - integrate(-1/4*((b^2*c*x^n
- b^2)*log(c*x^n + 1)^2 + 4*(a*b*c*x^n - a*b)*log(c*x^n + 1) + (4*a*b + (b
^2*c*n - 4*a*b*c)*x^n - 2*(b^2*c*x^n - b^2)*log(c*x^n + 1))*log(-c*x^n + 1
))/(c*x^3*x^n - x^3), x)
```

Giac [N/A]

Not integrable

Time = 0.39 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^3} dx = \int \frac{(b \operatorname{artanh}(cx^n) + a)^2}{x^3} dx$$

input

```
integrate((a+b*arctanh(c*x^n))^2/x^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^n) + a)^2/x^3, x)
```

Mupad [N/A]

Not integrable

Time = 3.38 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^3} dx = \int \frac{(a + b \operatorname{atanh}(cx^n))^2}{x^3} dx$$

input

```
int((a + b*atanh(c*x^n))^2/x^3,x)
```

output

```
int((a + b*atanh(c*x^n))^2/x^3, x)
```

Reduce [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 52, normalized size of antiderivative = 3.25

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))^2}{x^3} dx = \frac{4 \left(\int \frac{\operatorname{atanh}(x^n c)}{x^3} dx \right) ab x^2 + 2 \left(\int \frac{\operatorname{atanh}(x^n c)^2}{x^3} dx \right) b^2 x^2 - a^2}{2x^2}$$

input `int((a+b*atanh(c*x^n))^2/x^3,x)`output `(4*int(atanh(x**n*c)/x**3,x)*a*b*x**2 + 2*int(atanh(x**n*c)**2/x**3,x)*b**2*x**2 - a**2)/(2*x**2)`

3.235 $\int \frac{\operatorname{arctanh}(ax^n)}{x} dx$

Optimal result	1856
Mathematica [C] (verified)	1856
Rubi [A] (verified)	1857
Maple [A] (verified)	1858
Fricas [B] (verification not implemented)	1858
Sympy [F]	1859
Maxima [B] (verification not implemented)	1859
Giac [F]	1860
Mupad [F(-1)]	1860
Reduce [F]	1860

Optimal result

Integrand size = 10, antiderivative size = 30

$$\int \frac{\operatorname{arctanh}(ax^n)}{x} dx = -\frac{\operatorname{PolyLog}(2, -ax^n)}{2n} + \frac{\operatorname{PolyLog}(2, ax^n)}{2n}$$

output `-1/2*polylog(2,-a*x^n)/n+1/2*polylog(2,a*x^n)/n`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.10

$$\int \frac{\operatorname{arctanh}(ax^n)}{x} dx = \frac{ax^n {}_3F_2\left(\frac{1}{2}, \frac{1}{2}, 1; \frac{3}{2}, \frac{3}{2}; a^2x^{2n}\right)}{n}$$

input `Integrate[ArcTanh[a*x^n]/x,x]`

output `(a*x^n*HypergeometricPFQ[{1/2, 1/2, 1}, {3/2, 3/2}, a^2*x^(2*n)])/n`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {6450, 6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(ax^n)}{x} dx$$

↓ 6450

$$\int x^{-n} \operatorname{arctanh}(ax^n) dx^n$$

↓ 6446

$$\frac{\operatorname{PolyLog}(2, ax^n)}{2} - \frac{1}{2} \operatorname{PolyLog}(2, -ax^n)$$

n

input `Int[ArcTanh[a*x^n]/x, x]`

output `(-1/2*PolyLog[2, -(a*x^n)] + PolyLog[2, a*x^n]/2)/n`

Defintions of rubi rules used

rule 6446 `Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))/(x_), x_Symbol] := Simp[a*Log[x], x] + (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) / ; FreeQ[{a, b, c}, x]`

rule 6450 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)]*(b_.))^(p_.)/(x_), x_Symbol] := Simp[1/n Subst[Int[(a + b*ArcTanh[c*x])^p/x, x], x, x^n], x] / ; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.97

method	result	size
risch	$\frac{\operatorname{dilog}(1-ax^n)}{2n} - \frac{\operatorname{dilog}(ax^n+1)}{2n}$	29
derivativdivides	$\frac{\ln(ax^n) \operatorname{arctanh}(ax^n) - \frac{\operatorname{dilog}(ax^n)}{2} - \frac{\operatorname{dilog}(ax^n+1)}{2} - \frac{\ln(ax^n) \ln(ax^n+1)}{2}}{n}$	53
default	$\frac{\ln(ax^n) \operatorname{arctanh}(ax^n) - \frac{\operatorname{dilog}(ax^n)}{2} - \frac{\operatorname{dilog}(ax^n+1)}{2} - \frac{\ln(ax^n) \ln(ax^n+1)}{2}}{n}$	53
meijerg	$-\frac{i \left(\frac{2iax^n \operatorname{polylog}(2, \sqrt{x^{2n}a^2})}{\sqrt{x^{2n}a^2}} - \frac{2iax^n \operatorname{polylog}(2, -\sqrt{x^{2n}a^2})}{\sqrt{x^{2n}a^2}} \right)}{4n}$	72

input `int(arctanh(a*x^n)/x,x,method=_RETURNVERBOSE)`

output `1/2/n*dilog(1-a*x^n)-1/2/n*dilog(a*x^n+1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 129 vs. 2(24) = 48.

Time = 0.09 (sec) , antiderivative size = 129, normalized size of antiderivative = 4.30

$$\int \frac{\operatorname{arctanh}(ax^n)}{x} dx =$$

$$\frac{n \log(a \cosh(n \log(x)) + a \sinh(n \log(x)) + 1) \log(x) - n \log(-a \cosh(n \log(x)) - a \sinh(n \log(x)))}{n}$$

input `integrate(arctanh(a*x^n)/x,x, algorithm="fricas")`

output `-1/2*(n*log(a*cosh(n*log(x)) + a*sinh(n*log(x)) + 1)*log(x) - n*log(-a*cosh(n*log(x)) - a*sinh(n*log(x)) + 1)*log(x) - n*log(x)*log(-a*cosh(n*log(x)) + a*sinh(n*log(x)) + 1)/(a*cosh(n*log(x)) + a*sinh(n*log(x)) - 1)) - dilog(a*cosh(n*log(x)) + a*sinh(n*log(x))) + dilog(-a*cosh(n*log(x)) - a*sinh(n*log(x))))/n`

Sympy [F]

$$\int \frac{\operatorname{arctanh}(ax^n)}{x} dx = \int \frac{\operatorname{atanh}(ax^n)}{x} dx$$

input `integrate(atanh(a*x**n)/x,x)`

output `Integral(atanh(a*x**n)/x, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 147 vs. $2(24) = 48$.

Time = 0.10 (sec) , antiderivative size = 147, normalized size of antiderivative = 4.90

$$\begin{aligned} \int \frac{\operatorname{arctanh}(ax^n)}{x} dx &= -\frac{1}{2} an \left(\frac{\log\left(\frac{ax^n+1}{a}\right)}{an} - \frac{\log\left(\frac{ax^n-1}{a}\right)}{an} \right) \log(x) \\ &+ \frac{1}{2} an \left(\frac{\log(ax^n+1)\log(x) - \log(ax^n-1)\log(x)}{an} - \frac{n \log(ax^n+1)\log(x) + \operatorname{Li}_2(-ax^n)}{an^2} \right) + \frac{n \log(-a}{an^2} \\ &+ \operatorname{artanh}(ax^n) \log(x) \end{aligned}$$

input `integrate(arctanh(a*x^n)/x,x, algorithm="maxima")`

output `-1/2*a*n*(log((a*x^n + 1)/a)/(a*n) - log((a*x^n - 1)/a)/(a*n))*log(x) + 1/2*a*n*((log(a*x^n + 1)*log(x) - log(a*x^n - 1)*log(x))/(a*n) - (n*log(a*x^n + 1)*log(x) + dilog(-a*x^n))/(a*n^2) + (n*log(-a*x^n + 1)*log(x) + dilog(a*x^n))/(a*n^2)) + arctanh(a*x^n)*log(x)`

Giac [F]

$$\int \frac{\operatorname{arctanh}(ax^n)}{x} dx = \int \frac{\operatorname{artanh}(ax^n)}{x} dx$$

input `integrate(arctanh(a*x^n)/x,x, algorithm="giac")`

output `integrate(arctanh(a*x^n)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(ax^n)}{x} dx = \int \frac{\operatorname{atanh}(ax^n)}{x} dx$$

input `int(atanh(a*x^n)/x,x)`

output `int(atanh(a*x^n)/x, x)`

Reduce [F]

$$\int \frac{\operatorname{arctanh}(ax^n)}{x} dx = \int \frac{\operatorname{atanh}(x^n a)}{x} dx$$

input `int(atanh(a*x^n)/x,x)`

output `int(atanh(x**n*a)/x,x)`

3.236 $\int \frac{\operatorname{arctanh}(ax^5)}{x} dx$

Optimal result	1861
Mathematica [A] (verified)	1861
Rubi [A] (verified)	1862
Maple [C] (verified)	1863
Fricas [F]	1863
Sympy [F(-1)]	1864
Maxima [B] (verification not implemented)	1864
Giac [F]	1865
Mupad [F(-1)]	1865
Reduce [F]	1865

Optimal result

Integrand size = 10, antiderivative size = 24

$$\int \frac{\operatorname{arctanh}(ax^5)}{x} dx = -\frac{1}{10} \operatorname{PolyLog}(2, -ax^5) + \frac{\operatorname{PolyLog}(2, ax^5)}{10}$$

output `-1/10*polylog(2,-a*x^5)+1/10*polylog(2,a*x^5)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{arctanh}(ax^5)}{x} dx = \frac{1}{10} (-\operatorname{PolyLog}(2, -ax^5) + \operatorname{PolyLog}(2, ax^5))$$

input `Integrate[ArcTanh[a*x^5]/x,x]`

output `(-PolyLog[2, -(a*x^5)] + PolyLog[2, a*x^5])/10`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {6450, 6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(ax^5)}{x} dx$$

↓ 6450

$$\frac{1}{5} \int \frac{\operatorname{arctanh}(ax^5)}{x^5} dx^5$$

↓ 6446

$$\frac{1}{5} \left(\frac{\operatorname{PolyLog}(2, ax^5)}{2} - \frac{1}{2} \operatorname{PolyLog}(2, -ax^5) \right)$$

input

```
Int[ArcTanh[a*x^5]/x, x]
```

output

```
(-1/2*PolyLog[2, -(a*x^5)] + PolyLog[2, a*x^5]/2)/5
```

Defintions of rubi rules used

rule 6446

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))/(x_), x_Symbol] := Simp[a*Log[x], x] + (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) / ; FreeQ[{a, b, c}, x]
```

rule 6450

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)]*(b_.))^(p_.)/(x_), x_Symbol] := Simp[1/n Subst[Int[(a + b*ArcTanh[c*x])^p/x, x], x, x^n], x] / ; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.34 (sec) , antiderivative size = 61, normalized size of antiderivative = 2.54

method	result
meijerg	$-\frac{i \left(\frac{2ia x^5 \operatorname{polylog}(2, \sqrt{a^2 x^{10}})}{\sqrt{a^2 x^{10}}} - \frac{2ia x^5 \operatorname{polylog}(2, -\sqrt{a^2 x^{10}})}{\sqrt{a^2 x^{10}}} \right)}{20}$
default	$\ln(x) \operatorname{arctanh}(ax^5) - 5a \left(-\frac{\sum_{-R1=\operatorname{RootOf}(a-Z^5-1)} \left(\ln(x) \ln\left(\frac{R1-x}{-R1}\right) + \operatorname{dilog}\left(\frac{R1-x}{-R1}\right) \right)}{10a} + \frac{\sum_{-R1=\operatorname{RootOf}(a-Z^5-1)} \left(\ln(x) \ln\left(\frac{R1-x}{-R1}\right) + \operatorname{dilog}\left(\frac{R1-x}{-R1}\right) \right)}{10a} \right)$
parts	$\ln(x) \operatorname{arctanh}(ax^5) - 5a \left(-\frac{\sum_{-R1=\operatorname{RootOf}(a-Z^5-1)} \left(\ln(x) \ln\left(\frac{R1-x}{-R1}\right) + \operatorname{dilog}\left(\frac{R1-x}{-R1}\right) \right)}{10a} + \frac{\sum_{-R1=\operatorname{RootOf}(a-Z^5-1)} \left(\ln(x) \ln\left(\frac{R1-x}{-R1}\right) + \operatorname{dilog}\left(\frac{R1-x}{-R1}\right) \right)}{10a} \right)$
risch	$\frac{\ln(x) \ln(ax^5+1)}{2} - \frac{\left(\sum_{-R1=\operatorname{RootOf}(a-Z^5+1)} \left(\ln(x) \ln\left(\frac{R1-x}{-R1}\right) + \operatorname{dilog}\left(\frac{R1-x}{-R1}\right) \right) \right)}{2} - \frac{\ln(x) \ln(-ax^5+1)}{2} + \frac{\left(\sum_{-R1=\operatorname{RootOf}(a-Z^5+1)} \left(\ln(x) \ln\left(\frac{R1-x}{-R1}\right) + \operatorname{dilog}\left(\frac{R1-x}{-R1}\right) \right) \right)}{2}$

input `int(arctanh(a*x^5)/x,x,method=_RETURNVERBOSE)`

output `-1/20*I*(2*I*a*x^5/(a^2*x^10)^(1/2)*polylog(2,(a^2*x^10)^(1/2))-2*I*a*x^5/(a^2*x^10)^(1/2)*polylog(2,-(a^2*x^10)^(1/2)))`

Fricas [F]

$$\int \frac{\operatorname{arctanh}(ax^5)}{x} dx = \int \frac{\operatorname{artanh}(ax^5)}{x} dx$$

input `integrate(arctanh(a*x^5)/x,x, algorithm="fricas")`

output `integral(arctanh(a*x^5)/x, x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(ax^5)}{x} dx = \text{Timed out}$$

input `integrate(atanh(a*x**5)/x,x)`output `Timed out`**Maxima [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 104 vs. $2(18) = 36$.

Time = 0.04 (sec) , antiderivative size = 104, normalized size of antiderivative = 4.33

$$\int \frac{\operatorname{arctanh}(ax^5)}{x} dx = -\frac{1}{2} a \left(\frac{\log(ax^5 + 1)}{a} - \frac{\log(ax^5 - 1)}{a} \right) \log(x) - \frac{1}{10} a \left(\frac{\log(ax^5 - 1) \log(ax^5) + \operatorname{Li}_2(-ax^5 + 1)}{a} - \frac{\log(ax^5 + 1) \log(-ax^5) + \operatorname{Li}_2(ax^5 + 1)}{a} \right) + \operatorname{artanh}(ax^5) \log(x)$$

input `integrate(arctanh(a*x^5)/x,x, algorithm="maxima")`output `-1/2*a*(log(a*x^5 + 1)/a - log(a*x^5 - 1)/a)*log(x) - 1/10*a*((log(a*x^5 - 1)*log(a*x^5) + dilog(-a*x^5 + 1))/a - (log(a*x^5 + 1)*log(-a*x^5) + dilog(a*x^5 + 1))/a) + arctanh(a*x^5)*log(x)`

Giac [F]

$$\int \frac{\operatorname{arctanh}(ax^5)}{x} dx = \int \frac{\operatorname{artanh}(ax^5)}{x} dx$$

input `integrate(arctanh(a*x^5)/x,x, algorithm="giac")`

output `integrate(arctanh(a*x^5)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(ax^5)}{x} dx = \int \frac{\operatorname{atanh}(ax^5)}{x} dx$$

input `int(atanh(a*x^5)/x,x)`

output `int(atanh(a*x^5)/x, x)`

Reduce [F]

$$\int \frac{\operatorname{arctanh}(ax^5)}{x} dx = \int \frac{\operatorname{atanh}(ax^5)}{x} dx$$

input `int(atanh(a*x^5)/x,x)`

output `int(atanh(a*x**5)/x,x)`

3.237 $\int \operatorname{arctanh}\left(\frac{1}{x}\right) dx$

Optimal result	1866
Mathematica [A] (verified)	1866
Rubi [A] (verified)	1867
Maple [A] (verified)	1868
Fricas [A] (verification not implemented)	1868
Sympy [A] (verification not implemented)	1869
Maxima [A] (verification not implemented)	1869
Giac [B] (verification not implemented)	1869
Mupad [B] (verification not implemented)	1870
Reduce [B] (verification not implemented)	1870

Optimal result

Integrand size = 4, antiderivative size = 19

$$\int \operatorname{arctanh}\left(\frac{1}{x}\right) dx = x \operatorname{arctanh}\left(\frac{1}{x}\right) + \frac{1}{2} \log(1 - x^2)$$

output

```
x*arctanh(1/x)+1/2*ln(-x^2+1)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \operatorname{arctanh}\left(\frac{1}{x}\right) dx = x \operatorname{arctanh}\left(\frac{1}{x}\right) + \frac{1}{2} \log(-1 + x^2)$$

input

```
Integrate[ArcTanh[x^(-1)],x]
```

output

```
x*ArcTanh[x^(-1)] + Log[-1 + x^2]/2
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6436, 795, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \operatorname{arctanh}\left(\frac{1}{x}\right) dx \\ & \quad \downarrow \text{6436} \\ & \int \frac{1}{\left(1 - \frac{1}{x^2}\right)x} dx + x \operatorname{arctanh}\left(\frac{1}{x}\right) \\ & \quad \downarrow \text{795} \\ & \int \frac{x}{x^2 - 1} dx + x \operatorname{arctanh}\left(\frac{1}{x}\right) \\ & \quad \downarrow \text{240} \\ & x \operatorname{arctanh}\left(\frac{1}{x}\right) + \frac{1}{2} \log(1 - x^2) \end{aligned}$$

input

```
Int[ArcTanh[x^(-1)], x]
```

output

```
x*ArcTanh[x^(-1)] + Log[1 - x^2]/2
```

Defintions of rubi rules used

rule 240

```
Int[(x_)/((a_) + (b_)*(x_)^2), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]
```

rule 795

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :> Int[x^(m + n*p)*(b + a/x^n)^p, x] /; FreeQ[{a, b, m, n}, x] && IntegerQ[p] && NegQ[n]
```


rule 6436

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.), x_Symbol] := Simp[x*(a
+ b*ArcTanh[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTanh[c*x^n])
^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]
&& (EqQ[n, 1] || EqQ[p, 1])
```

Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

method	result
parallelsch	$x \operatorname{arctanh}\left(\frac{1}{x}\right) + \ln(-1+x) + \operatorname{arctanh}\left(\frac{1}{x}\right)$
parts	$x \operatorname{arctanh}\left(\frac{1}{x}\right) + \frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$
derivativdivides	$x \operatorname{arctanh}\left(\frac{1}{x}\right) + \frac{\ln\left(\frac{1}{x}+1\right)}{2} - \ln\left(\frac{1}{x}\right) + \frac{\ln\left(\frac{1}{x}-1\right)}{2}$
default	$x \operatorname{arctanh}\left(\frac{1}{x}\right) + \frac{\ln\left(\frac{1}{x}+1\right)}{2} - \ln\left(\frac{1}{x}\right) + \frac{\ln\left(\frac{1}{x}-1\right)}{2}$
meijerg	$\ln(x) - \frac{i\pi}{2} - \frac{\ln\left(1-\sqrt{\frac{1}{x^2}}\right) - \ln\left(1+\sqrt{\frac{1}{x^2}}\right)}{2\sqrt{\frac{1}{x^2}}} + \frac{\ln\left(-\frac{1}{x^2}+1\right)}{2}$
risch	$\frac{x \ln(1+x)}{2} - \frac{\ln(-1+x)x}{2} + \frac{i\pi \operatorname{csgn}\left(\frac{i}{x}\right) \operatorname{csgn}(i(-1+x)) \operatorname{csgn}\left(\frac{i(-1+x)}{x}\right)x}{4} - \frac{i\pi \operatorname{csgn}\left(\frac{i}{x}\right) \operatorname{csgn}(i(1+x)) \operatorname{csgn}\left(\frac{i(1+x)}{x}\right)}{4}$

input `int(arctanh(1/x), x, method=_RETURNVERBOSE)`

output `x*arctanh(1/x)+ln(-1+x)+arctanh(1/x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.16

$$\int \operatorname{arctanh}\left(\frac{1}{x}\right) dx = \frac{1}{2} x \log\left(\frac{x+1}{x-1}\right) + \frac{1}{2} \log(x^2 - 1)$$

input `integrate(arctanh(1/x), x, algorithm="fricas")`

output `1/2*x*log((x + 1)/(x - 1)) + 1/2*log(x^2 - 1)`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \operatorname{arctanh}\left(\frac{1}{x}\right) dx = x \operatorname{atanh}\left(\frac{1}{x}\right) + \log(x+1) - \operatorname{atanh}\left(\frac{1}{x}\right)$$

input `integrate(atanh(1/x),x)`

output `x*atanh(1/x) + log(x + 1) - atanh(1/x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \operatorname{arctanh}\left(\frac{1}{x}\right) dx = x \operatorname{artanh}\left(\frac{1}{x}\right) + \frac{1}{2} \log(x^2 - 1)$$

input `integrate(arctanh(1/x),x, algorithm="maxima")`

output `x*arctanh(1/x) + 1/2*log(x^2 - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 101 vs. $2(17) = 34$.

Time = 0.11 (sec) , antiderivative size = 101, normalized size of antiderivative = 5.32

$$\int \operatorname{arctanh}\left(\frac{1}{x}\right) dx = \frac{\log\left(-\frac{\frac{\frac{x+1}{x-1}-1}{x-1}+1}{\frac{x+1}{x-1}-1}\right)}{\frac{x+1}{x-1}-1} + \log\left(\frac{|x+1|}{|x-1|}\right) - \log\left(\left|\frac{x+1}{x-1}-1\right|\right)$$

input `integrate(arctanh(1/x),x, algorithm="giac")`

output

```
log(-(((x + 1)/(x - 1) - 1)/((x + 1)/(x - 1) + 1) + 1)/(((x + 1)/(x - 1) - 1)/((x + 1)/(x - 1) + 1) - 1))/((x + 1)/(x - 1) - 1) + log(abs(x + 1)/abs(x - 1)) - log(abs((x + 1)/(x - 1) - 1))
```

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \operatorname{arctanh}\left(\frac{1}{x}\right) dx = \frac{\ln(x^2 - 1)}{2} + x \operatorname{atanh}\left(\frac{1}{x}\right)$$

input

```
int(atanh(1/x), x)
```

output

```
log(x^2 - 1)/2 + x*atanh(1/x)
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \operatorname{arctanh}\left(\frac{1}{x}\right) dx = \operatorname{atanh}\left(\frac{1}{x}\right) x + \operatorname{atanh}\left(\frac{1}{x}\right) + \log(x - 1)$$

input

```
int(atanh(1/x), x)
```

output

```
atanh(1/x)*x + atanh(1/x) + log(x - 1)
```

3.238 $\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^3 dx$

Optimal result	1871
Mathematica [N/A]	1871
Rubi [N/A]	1872
Maple [N/A]	1872
Fricas [N/A]	1873
Sympy [F(-1)]	1873
Maxima [N/A]	1873
Giac [N/A]	1874
Mupad [N/A]	1874
Reduce [N/A]	1875

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^3 dx = \operatorname{Int}((dx)^m (a + b \operatorname{arctanh}(cx^n))^3, x)$$

output `Defer(Int)((d*x)^m*(a+b*arctanh(c*x^n))^3,x)`

Mathematica [N/A]

Not integrable

Time = 6.76 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^3 dx = \int (dx)^m (a + b \operatorname{arctanh}(cx^n))^3 dx$$

input `Integrate[(d*x)^m*(a + b*ArcTanh[c*x^n])^3,x]`

output `Integrate[(d*x)^m*(a + b*ArcTanh[c*x^n])^3, x]`

Rubi [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^3 dx$$

↓ 6468

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^3 dx$$

input `Int[(d*x)^m*(a + b*ArcTanh[c*x^n])^3,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.05 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^3 dx$$

input `int((d*x)^m*(a+b*arctanh(c*x^n))^3,x)`

output `int((d*x)^m*(a+b*arctanh(c*x^n))^3,x)`

Fricas [N/A]

Not integrable

Time = 0.10 (sec) , antiderivative size = 65, normalized size of antiderivative = 3.61

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^3 dx = \int (b \operatorname{artanh}(cx^n) + a)^3 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^n))^3,x, algorithm="fricas")`

output `integral((d*x)^m*b^3*arctanh(c*x^n)^3 + 3*(d*x)^m*a*b^2*arctanh(c*x^n)^2 + 3*(d*x)^m*a^2*b*arctanh(c*x^n) + (d*x)^m*a^3, x)`

Sympy [F(-1)]

Timed out.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^3 dx = \text{Timed out}$$

input `integrate((d*x)**m*(a+b*atanh(c*x**n))**3,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 1.49 (sec) , antiderivative size = 468, normalized size of antiderivative = 26.00

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^3 dx = \int (b \operatorname{artanh}(cx^n) + a)^3 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^n))^3,x, algorithm="maxima")`

output

```
-1/8*b^3*d^m*x*x^m*log(-c*x^n + 1)^3/(m + 1) + (d*x)^(m + 1)*a^3/(d*(m + 1)) + integrate(1/8*((b^3*c*d^m*(m + 1)*e^(m*log(x) + n*log(x)) - b^3*d^m*(m + 1)*x^m)*log(c*x^n + 1)^3 + 6*(a*b^2*c*d^m*(m + 1)*e^(m*log(x) + n*log(x)) - a*b^2*d^m*(m + 1)*x^m)*log(c*x^n + 1)^2 - 3*(2*a*b^2*d^m*(m + 1)*x^m - (2*a*b^2*c*d^m*(m + 1) + b^3*c*d^m*n)*e^(m*log(x) + n*log(x)) - (b^3*c*d^m*(m + 1)*e^(m*log(x) + n*log(x)) - b^3*d^m*(m + 1)*x^m)*log(c*x^n + 1))*log(-c*x^n + 1)^2 + 12*(a^2*b*c*d^m*(m + 1)*e^(m*log(x) + n*log(x)) - a^2*b*d^m*(m + 1)*x^m)*log(c*x^n + 1) - 3*(4*a^2*b*c*d^m*(m + 1)*e^(m*log(x) + n*log(x)) - 4*a^2*b*d^m*(m + 1)*x^m + (b^3*c*d^m*(m + 1)*e^(m*log(x) + n*log(x)) - b^3*d^m*(m + 1)*x^m)*log(c*x^n + 1)^2 + 4*(a*b^2*c*d^m*(m + 1)*e^(m*log(x) + n*log(x)) - a*b^2*d^m*(m + 1)*x^m)*log(c*x^n + 1))*log(-c*x^n + 1))/(c*(m + 1)*x^n - m - 1), x)
```

Giac [N/A]

Not integrable

Time = 0.77 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^3 dx = \int (b \operatorname{artanh}(cx^n) + a)^3 (dx)^m dx$$

input

```
integrate((d*x)^m*(a+b*arctanh(c*x^n))^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^n) + a)^3*(d*x)^m, x)
```

Mupad [N/A]

Not integrable

Time = 3.62 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^3 dx = \int (dx)^m (a + b \operatorname{atanh}(cx^n))^3 dx$$

input

```
int((d*x)^m*(a + b*atanh(c*x^n))^3,x)
```

output `int((d*x)^m*(a + b*atanh(c*x^n))^3, x)`

Reduce [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 133, normalized size of antiderivative = 7.39

$$\int (dx)^m (a + b \operatorname{atanh}(cx^n))^3 dx$$

$$= \frac{d^m (x^m a^3 x + 3 \int x^m \operatorname{atanh}(x^n c) dx) a^2 b m + 3 \left(\int x^m \operatorname{atanh}(x^n c) dx \right) a^2 b + \left(\int x^m \operatorname{atanh}(x^n c)^3 dx \right) b^3 m + \dots}{m + 1}$$

input `int((d*x)^m*(a+b*atanh(c*x^n))^3,x)`

output `(d**m*(x**m*a**3*x + 3*int(x**m*atanh(x**n*c),x)*a**2*b*m + 3*int(x**m*atanh(x**n*c),x)*a**2*b + int(x**m*atanh(x**n*c)**3,x)*b**3*m + int(x**m*atanh(x**n*c)**3,x)*b**3 + 3*int(x**m*atanh(x**n*c)**2,x)*a*b**2*m + 3*int(x**m*atanh(x**n*c)**2,x)*a*b**2))/(m + 1)`

3.239 $\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^2 dx$

Optimal result	1876
Mathematica [N/A]	1876
Rubi [N/A]	1877
Maple [N/A]	1877
Fricas [N/A]	1878
Sympy [F(-1)]	1878
Maxima [N/A]	1878
Giac [N/A]	1879
Mupad [N/A]	1879
Reduce [N/A]	1880

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^2 dx = \operatorname{Int}((dx)^m (a + b \operatorname{arctanh}(cx^n))^2, x)$$

output `Defer(Int)((d*x)^m*(a+b*arctanh(c*x^n))^2,x)`

Mathematica [N/A]

Not integrable

Time = 14.30 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^2 dx = \int (dx)^m (a + b \operatorname{arctanh}(cx^n))^2 dx$$

input `Integrate[(d*x)^m*(a + b*ArcTanh[c*x^n])^2,x]`

output `Integrate[(d*x)^m*(a + b*ArcTanh[c*x^n])^2, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^2 dx$$

↓ 6468

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^2 dx$$

input `Int[(d*x)^m*(a + b*ArcTanh[c*x^n])^2,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^2 dx$$

input `int((d*x)^m*(a+b*arctanh(c*x^n))^2,x)`

output `int((d*x)^m*(a+b*arctanh(c*x^n))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.44

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^2 dx = \int (b \operatorname{artanh}(cx^n) + a)^2 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^n))^2,x, algorithm="fricas")`

output `integral((d*x)^m*b^2*arctanh(c*x^n)^2 + 2*(d*x)^m*a*b*arctanh(c*x^n) + (d*x)^m*a^2, x)`

Sympy [F(-1)]

Timed out.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^2 dx = \text{Timed out}$$

input `integrate((d*x)**m*(a+b*atanh(c*x**n))**2,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.88 (sec) , antiderivative size = 265, normalized size of antiderivative = 14.72

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^2 dx = \int (b \operatorname{artanh}(cx^n) + a)^2 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^n))^2,x, algorithm="maxima")`

output
$$\frac{1}{4}b^2d^mxx^m\log(-cx^n+1)^2/(m+1) + (dx)^{m+1}a^2/(d(m+1)) - \int (-1/4((b^2cd^m(m+1)e^{(m\log(x)+n\log(x))} - b^2d^m(m+1)x^m)\log(cx^n+1)^2 + 4(abcd^m(m+1)e^{(m\log(x)+n\log(x))} - abcd^m(m+1)x^m)\log(cx^n+1) + 2(2abd^m(m+1)x^m - (2ab^2cd^m(m+1) + b^2cd^m n)e^{(m\log(x)+n\log(x))} - (b^2cd^m(m+1)e^{(m\log(x)+n\log(x))} - b^2d^m(m+1)x^m)\log(cx^n+1))\log(-cx^n+1))/(c(m+1)x^n - m - 1), x)$$

Giac [N/A]

Not integrable

Time = 0.45 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^2 dx = \int (b \operatorname{artanh}(cx^n) + a)^2 (dx)^m dx$$

input `integrate((dx)^m*(a+b*arctanh(cx^n))^2,x, algorithm="giac")`

output `integrate((b*arctanh(cx^n) + a)^2*(dx)^m, x)`

Mupad [N/A]

Not integrable

Time = 3.61 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^2 dx = \int (dx)^m (a + b \operatorname{atanh}(cx^n))^2 dx$$

input `int((dx)^m*(a + b*atanh(cx^n))^2,x)`

output `int((dx)^m*(a + b*atanh(cx^n))^2, x)`

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 88, normalized size of antiderivative = 4.89

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n))^2 dx$$

$$= \frac{d^m (x^m a^2 x + 2(\int x^m \operatorname{atanh}(x^n c) dx) abm + 2(\int x^m \operatorname{atanh}(x^n c) dx) ab + (\int x^m \operatorname{atanh}(x^n c)^2 dx) b^2 m + (\int x^m \operatorname{atanh}(x^n c)^2 dx) b^2 m + (\int x^m \operatorname{atanh}(x^n c)^2 dx) b^2 m}{m + 1}$$

input `int((d*x)^m*(a+b*atanh(c*x^n))^2,x)`

output `(d**m*(x**m*a**2*x + 2*int(x**m*atanh(x**n*c),x)*a*b*m + 2*int(x**m*atanh(x**n*c),x)*a*b + int(x**m*atanh(x**n*c)**2,x)*b**2*m + int(x**m*atanh(x**n*c)**2,x)*b**2))/m + 1)`

3.240 $\int (dx)^m (a + \operatorname{barctanh}(cx^n)) dx$

Optimal result	1881
Mathematica [A] (verified)	1881
Rubi [A] (verified)	1882
Maple [F]	1883
Fricas [F]	1883
Sympy [F]	1884
Maxima [F]	1884
Giac [F]	1884
Mupad [F(-1)]	1885
Reduce [F]	1885

Optimal result

Integrand size = 16, antiderivative size = 84

$$\int (dx)^m (a + \operatorname{barctanh}(cx^n)) dx = \frac{x(dx)^m (a + \operatorname{barctanh}(cx^n))}{1 + m} - \frac{bcnx^{1+n}(dx)^m \operatorname{Hypergeometric2F1}\left(1, \frac{1+m+n}{2n}, \frac{1+m+3n}{2n}, c^2x^{2n}\right)}{(1+m)(1+m+n)}$$

output `x*(d*x)^m*(a+b*arctanh(c*x^n))/(1+m)-b*c*n*x^(1+n)*(d*x)^m*hypergeom([1, 1/2*(1+m+n)/n], [1/2*(1+m+3*n)/n], c^2*x^(2*n))/(1+m)/(1+m+n)`

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.92

$$\int (dx)^m (a + \operatorname{barctanh}(cx^n)) dx = \frac{x(dx)^m ((1+m+n)(a + \operatorname{barctanh}(cx^n)) - bcnx^n \operatorname{Hypergeometric2F1}\left(1, \frac{1+m+n}{2n}, \frac{1+m+3n}{2n}, c^2x^{2n}\right))}{(1+m)(1+m+n)}$$

input `Integrate[(d*x)^m*(a + b*ArcTanh[c*x^n]),x]`

output

```
(x*(d*x)^m*((1 + m + n)*(a + b*ArcTanh[c*x^n]) - b*c*n*x^n*Hypergeometric2
F1[1, (1 + m + n)/(2*n), (1 + m + 3*n)/(2*n), c^2*x^(2*n)]))/((1 + m)*(1 +
m + n))
```

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.07, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6466, 6452, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m (a + \text{barctanh}(cx^n)) dx$$

$$\downarrow 6466$$

$$x^{-m} (dx)^m \int x^m (a + \text{barctanh}(cx^n)) dx$$

$$\downarrow 6452$$

$$x^{-m} (dx)^m \left(\frac{x^{m+1} (a + \text{barctanh}(cx^n))}{m+1} - \frac{bcn \int \frac{x^{m+n}}{1-c^2 x^{2n}} dx}{m+1} \right)$$

$$\downarrow 888$$

$$x^{-m} (dx)^m \left(\frac{x^{m+1} (a + \text{barctanh}(cx^n))}{m+1} - \frac{bcn x^{m+n+1} \text{Hypergeometric2F1} \left(1, \frac{m+n+1}{2n}, \frac{m+3n+1}{2n}, c^2 x^{2n} \right)}{(m+1)(m+n+1)} \right)$$

input

```
Int[(d*x)^m*(a + b*ArcTanh[c*x^n]),x]
```

output

```
((d*x)^m*((x^(1 + m)*(a + b*ArcTanh[c*x^n]))/(1 + m) - (b*c*n*x^(1 + m + n)
)*Hypergeometric2F1[1, (1 + m + n)/(2*n), (1 + m + 3*n)/(2*n), c^2*x^(2*n)
])/((1 + m)*(1 + m + n)))/x^m
```

Definitions of rubi rules used

rule 888 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p *((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6466 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((d_.)*(x_)^(m_.), x_Symbol] :> Simp[d^IntPart[m]*((d*x)^FracPart[m]/x^FracPart[m]) Int[x^m*(a + b*ArcTanh[c*x^n])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || RationalQ[m, n])`

Maple [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n)) dx$$

input `int((d*x)^m*(a+b*arctanh(c*x^n)),x)`

output `int((d*x)^m*(a+b*arctanh(c*x^n)),x)`

Fricas [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n)) dx = \int (b \operatorname{arctanh}(cx^n) + a)(dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^n)),x, algorithm="fricas")`

output `integral((d*x)^m*b*arctanh(c*x^n) + (d*x)^m*a, x)`

Sympy [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n)) dx = \int (dx)^m (a + b \operatorname{atanh}(cx^n)) dx$$

input `integrate((d*x)**m*(a+b*atanh(c*x**n)), x)`

output `Integral((d*x)**m*(a + b*atanh(c*x**n)), x)`

Maxima [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n)) dx = \int (b \operatorname{artanh}(cx^n) + a)(dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^n)), x, algorithm="maxima")`

output `1/2*(d^m*n*integrate(x^m/(c*(m + 1)*x^n + m + 1), x) + d^m*n*integrate(x^m/(c*(m + 1)*x^n - m - 1), x) + (d^m*x*x^m*log(c*x^n + 1) - d^m*x*x^m*log(-c*x^n + 1))/(m + 1)*b + (d*x)^(m + 1)*a/(d*(m + 1))`

Giac [F]

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n)) dx = \int (b \operatorname{artanh}(cx^n) + a)(dx)^m dx$$

input `integrate((d*x)^m*(a+b*arctanh(c*x^n)), x, algorithm="giac")`

output `integrate((b*arctanh(c*x^n) + a)*(d*x)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (dx)^m (a + b \operatorname{arctanh}(cx^n)) dx = \int (dx)^m (a + b \operatorname{atanh}(cx^n)) dx$$

input `int((d*x)^m*(a + b*atanh(c*x^n)),x)`

output `int((d*x)^m*(a + b*atanh(c*x^n)), x)`

Reduce [F]

$$\begin{aligned} & \int (dx)^m (a + b \operatorname{arctanh}(cx^n)) dx \\ &= \frac{d^m (x^m a x + (\int x^m \operatorname{atanh}(x^n c) dx) b m + (\int x^m \operatorname{atanh}(x^n c) dx) b)}{m + 1} \end{aligned}$$

input `int((d*x)^m*(a+b*atanh(c*x^n)),x)`

output `(d**m*(x**m*a*x + int(x**m*atanh(x**n*c),x)*b*m + int(x**m*atanh(x**n*c),x)*b))/(m + 1)`

$$3.241 \quad \int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^n)} dx$$

Optimal result	1886
Mathematica [N/A]	1886
Rubi [N/A]	1887
Maple [N/A]	1887
Fricas [N/A]	1888
Sympy [N/A]	1888
Maxima [N/A]	1888
Giac [N/A]	1889
Mupad [N/A]	1889
Reduce [N/A]	1890

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^n)} dx = \operatorname{Int}\left(\frac{(dx)^m}{a + b \operatorname{arctanh}(cx^n)}, x\right)$$

output `Defer(Int)((d*x)^m/(a+b*arctanh(c*x^n)),x)`

Mathematica [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^n)} dx = \int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^n)} dx$$

input `Integrate[(d*x)^m/(a + b*ArcTanh[c*x^n]),x]`

output `Integrate[(d*x)^m/(a + b*ArcTanh[c*x^n]), x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^n)} dx$$

↓ 6468

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^n)} dx$$

input `Int[(d*x)^m/(a + b*ArcTanh[c*x^n]),x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^n)} dx$$

input `int((d*x)^m/(a+b*arctanh(c*x^n)),x)`

output `int((d*x)^m/(a+b*arctanh(c*x^n)),x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^n)} dx = \int \frac{(dx)^m}{b \operatorname{artanh}(cx^n) + a} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x^n)),x, algorithm="fricas")`

output `integral((d*x)^m/(b*arctanh(c*x^n) + a), x)`

Sympy [N/A]

Not integrable

Time = 36.48 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^n)} dx = \int \frac{(dx)^m}{a + b \operatorname{atanh}(cx^n)} dx$$

input `integrate((d*x)**m/(a+b*atanh(c*x**n)),x)`

output `Integral((d*x)**m/(a + b*atanh(c*x**n)), x)`

Maxima [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^n)} dx = \int \frac{(dx)^m}{b \operatorname{artanh}(cx^n) + a} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x^n)),x, algorithm="maxima")`

output `integrate((d*x)^m/(b*arctanh(c*x^n) + a), x)`

Giac [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^n)} dx = \int \frac{(dx)^m}{b \operatorname{artanh}(cx^n) + a} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x^n)),x, algorithm="giac")`

output `integrate((d*x)^m/(b*arctanh(c*x^n) + a), x)`

Mupad [N/A]

Not integrable

Time = 3.36 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^n)} dx = \int \frac{(dx)^m}{a + b \operatorname{atanh}(cx^n)} dx$$

input `int((d*x)^m/(a + b*atanh(c*x^n)),x)`

output `int((d*x)^m/(a + b*atanh(c*x^n)), x)`

Reduce [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

$$\int \frac{(dx)^m}{a + b \operatorname{arctanh}(cx^n)} dx = d^m \left(\int \frac{x^m}{\operatorname{atanh}(x^n c) b + a} dx \right)$$

input `int((d*x)^m/(a+b*atanh(c*x^n)),x)`output `d**m*int(x**m/(atanh(x**n*c)*b + a),x)`

3.242
$$\int \frac{(dx)^m}{(a+b\mathbf{arctanh}(cx^n))^2} dx$$

Optimal result	1891
Mathematica [N/A]	1891
Rubi [N/A]	1892
Maple [F(-1)]	1892
Fricas [N/A]	1893
Sympy [F(-1)]	1893
Maxima [N/A]	1893
Giac [N/A]	1894
Mupad [N/A]	1894
Reduce [N/A]	1895

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{(dx)^m}{(a + \mathbf{barctanh}(cx^n))^2} dx = \mathbf{Int}\left(\frac{(dx)^m}{(a + \mathbf{barctanh}(cx^n))^2}, x\right)$$

output

```
Defer(Int)((d*x)^m/(a+b*arctanh(c*x^n))^2,x)
```

Mathematica [N/A]

Not integrable

Time = 1.76 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{(a + \mathbf{barctanh}(cx^n))^2} dx = \int \frac{(dx)^m}{(a + \mathbf{barctanh}(cx^n))^2} dx$$

input

```
Integrate[(d*x)^m/(a + b*ArcTanh[c*x^n])^2,x]
```

output

```
Integrate[(d*x)^m/(a + b*ArcTanh[c*x^n])^2, x]
```


Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^n))^2} dx$$

↓ 6468

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^n))^2} dx$$

input `Int[(d*x)^m/(a + b*ArcTanh[c*x^n])^2,x]`

output `$Aborted`

Maple [F(-1)]

Timed out.

hanged

input `int((d*x)^m/(a+b*arctanh(c*x^n))^2,x)`

output `int((d*x)^m/(a+b*arctanh(c*x^n))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.00

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^n))^2} dx = \int \frac{(dx)^m}{(b \operatorname{artanh}(cx^n) + a)^2} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x^n))^2,x, algorithm="fricas")`

output `integral((d*x)^m/(b^2*arctanh(c*x^n)^2 + 2*a*b*arctanh(c*x^n) + a^2), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^n))^2} dx = \text{Timed out}$$

input `integrate((d*x)**m/(a+b*atanh(c*x**n))**2,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 169, normalized size of antiderivative = 9.39

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^n))^2} dx = \int \frac{(dx)^m}{(b \operatorname{artanh}(cx^n) + a)^2} dx$$

input `integrate((d*x)^m/(a+b*arctanh(c*x^n))^2,x, algorithm="maxima")`

output

```
2*(c^2*d^m*x*e^(m*log(x) + 2*n*log(x)) - d^m*x*x^m)/(b^2*c*n*x^n*log(c*x^n
+ 1) - b^2*c*n*x^n*log(-c*x^n + 1) + 2*a*b*c*n*x^n) + integrate(-2*(c^2*d
^m*(m + n + 1)*e^(m*log(x) + 2*n*log(x)) - d^m*(m - n + 1)*x^m)/(b^2*c*n*x
^n*log(c*x^n + 1) - b^2*c*n*x^n*log(-c*x^n + 1) + 2*a*b*c*n*x^n), x)
```

Giac [N/A]

Not integrable

Time = 0.39 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^n))^2} dx = \int \frac{(dx)^m}{(b \operatorname{artanh}(cx^n) + a)^2} dx$$

input

```
integrate((d*x)^m/(a+b*arctanh(c*x^n))^2,x, algorithm="giac")
```

output

```
integrate((d*x)^m/(b*arctanh(c*x^n) + a)^2, x)
```

Mupad [N/A]

Not integrable

Time = 3.47 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^n))^2} dx = \int \frac{(dx)^m}{(a + b \operatorname{atanh}(cx^n))^2} dx$$

input

```
int((d*x)^m/(a + b*atanh(c*x^n))^2,x)
```

output

```
int((d*x)^m/(a + b*atanh(c*x^n))^2, x)
```

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.11

$$\int \frac{(dx)^m}{(a + b \operatorname{arctanh}(cx^n))^2} dx = d^m \left(\int \frac{x^m}{\operatorname{atanh}(x^n c)^2 b^2 + 2 \operatorname{atanh}(x^n c) ab + a^2} dx \right)$$

input `int((d*x)^m/(a+b*atanh(c*x^n))^2,x)`output `d**m*int(x**m/(atanh(x**n*c)**2*b**2 + 2*atanh(x**n*c)*a*b + a**2),x)`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 1896
4.2 Links to plain text integration problems used in this report for each CAS . 1914

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
          finalresult={"C","Result contains complex when optimal does not."}
        ]
      ,(*ELSE*)(*result does not contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
          ]
        ]
      ,(*ELSE*)(*expnResult>expnOptimal*)
        If[FreeQ[result,Integrate] && FreeQ[result,Int],
          finalresult={"C","Result contains higher order function than in optimal. Order "
          ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]==Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]==Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]==Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]==Plus || Head[expn]==Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```
    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 5]],
  If [AppellFunctionQ [Head [expn]],
    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 6]],
  If [Head [expn] === RootSum,
    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 7]],
  If [Head [expn] === Integrate || Head [expn] === Int,
    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 8]],
  9]]]]]]]]]]]
```

```
ElementaryFunctionQ [func_] :=
  MemberQ [ {
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]
```

```
SpecialFunctionQ [func_] :=
  MemberQ [ {
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]
```

```
HypergeometricFunctionQ [func_] :=
  MemberQ [ {Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]
```

```
AppellFunctionQ [func_] :=
  MemberQ [ {AppellF1}, func]
```


Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

        end if
    else #result contains complex but optimal is not
        if debug then
            print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
    fi;
else # result do not contain complex
    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well
    fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co

        fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```



```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file