

Computer Algebra Independent Integration Tests

Summer 2024

7-Inverse-hyperbolic-functions/7.3-Inverse-hyperbolic-
tangent/338-7.3.3

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 8:25am

Contents

1	Introduction	4
1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23
2	detailed summary tables of results	24
2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	42
3	Listing of integrals	44
3.1	$\int (d + ex)^4(a + \operatorname{barctanh}(cx)) dx$	46
3.2	$\int (d + ex)^3(a + \operatorname{barctanh}(cx)) dx$	54
3.3	$\int (d + ex)^2(a + \operatorname{barctanh}(cx)) dx$	62
3.4	$\int (d + ex)(a + \operatorname{barctanh}(cx)) dx$	69
3.5	$\int \frac{a + \operatorname{barctanh}(cx)}{d + ex} dx$	75
3.6	$\int \frac{a + \operatorname{barctanh}(cx)}{(d + ex)^2} dx$	82

3.7	$\int \frac{a+\operatorname{barctanh}(cx)}{(d+ex)^3} dx$	89
3.8	$\int \frac{a+\operatorname{barctanh}(cx)}{(d+ex)^4} dx$	97
3.9	$\int (d+ex)^3(a+\operatorname{barctanh}(cx))^2 dx$	107
3.10	$\int (d+ex)^2(a+\operatorname{barctanh}(cx))^2 dx$	117
3.11	$\int (d+ex)(a+\operatorname{barctanh}(cx))^2 dx$	125
3.12	$\int \frac{(a+\operatorname{barctanh}(cx))^2}{d+ex} dx$	132
3.13	$\int \frac{(a+\operatorname{barctanh}(cx))^2}{(d+ex)^2} dx$	139
3.14	$\int \frac{(a+\operatorname{barctanh}(cx))^2}{(d+ex)^3} dx$	146
3.15	$\int (d+ex)^3(a+\operatorname{barctanh}(cx))^3 dx$	154
3.16	$\int (d+ex)^2(a+\operatorname{barctanh}(cx))^3 dx$	163
3.17	$\int (d+ex)(a+\operatorname{barctanh}(cx))^3 dx$	171
3.18	$\int \frac{(a+\operatorname{barctanh}(cx))^3}{d+ex} dx$	178
3.19	$\int \frac{(a+\operatorname{barctanh}(cx))^3}{(d+ex)^2} dx$	185
3.20	$\int \frac{(a+\operatorname{barctanh}(cx))^3}{(d+ex)^3} dx$	194
3.21	$\int \frac{a+\operatorname{barctanh}(cx)}{1+2cx} dx$	202
3.22	$\int \frac{\operatorname{arctanh}(x)}{1-\sqrt{2}x} dx$	208
3.23	$\int \frac{\operatorname{arctanh}(x)}{a+bx} dx$	215
3.24	$\int (d+ex)^3(a+\operatorname{barctanh}(cx^2)) dx$	221
3.25	$\int (d+ex)^2(a+\operatorname{barctanh}(cx^2)) dx$	231
3.26	$\int (d+ex)(a+\operatorname{barctanh}(cx^2)) dx$	240
3.27	$\int \frac{a+\operatorname{barctanh}(cx^2)}{d+ex} dx$	248
3.28	$\int \frac{a+\operatorname{barctanh}(cx^2)}{(d+ex)^2} dx$	255
3.29	$\int \frac{a+\operatorname{barctanh}(cx^2)}{(d+ex)^3} dx$	265
3.30	$\int (d+ex)(a+\operatorname{barctanh}(cx^2))^2 dx$	274
3.31	$\int \frac{(a+\operatorname{barctanh}(cx^2))^2}{d+ex} dx$	282
3.32	$\int \frac{(a+\operatorname{barctanh}(cx^2))^2}{(d+ex)^2} dx$	287
3.33	$\int (d+ex)^2(a+\operatorname{barctanh}(cx^3)) dx$	292
3.34	$\int (d+ex)(a+\operatorname{barctanh}(cx^3)) dx$	302
3.35	$\int \frac{a+\operatorname{barctanh}(cx^3)}{d+ex} dx$	311
3.36	$\int \frac{a+\operatorname{barctanh}(cx^3)}{(d+ex)^2} dx$	319
3.37	$\int \frac{x^3(a+\operatorname{barctanh}(c\sqrt{x}))}{1-c^2x} dx$	329
3.38	$\int \frac{x^2(a+\operatorname{barctanh}(c\sqrt{x}))}{1-c^2x} dx$	339

3.39	$\int \frac{x(a+b\operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} dx$	348
3.40	$\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{1-c^2x} dx$	356
3.41	$\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x(1-c^2x)} dx$	362
3.42	$\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x^2(1-c^2x)} dx$	368
3.43	$\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x^3(1-c^2x)} dx$	376
3.44	$\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x^4(1-c^2x)} dx$	385
3.45	$\int \frac{x^2(a+b\operatorname{arctanh}(c\sqrt{x}))}{d+ex} dx$	395
3.46	$\int \frac{x(a+b\operatorname{arctanh}(c\sqrt{x}))}{d+ex} dx$	405
3.47	$\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{d+ex} dx$	413
3.48	$\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x(d+ex)} dx$	420
3.49	$\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x^2(d+ex)} dx$	427
3.50	$\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x^3(d+ex)} dx$	436
4	Appendix	447
4.1	Listing of Grading functions	447
4.2	Links to plain text integration problems used in this report for each CAS465	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [50]. This is test number [338].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (50)	0.00 (0)
Mathematica	98.00 (49)	2.00 (1)
Maple	98.00 (49)	2.00 (1)
Maxima	60.00 (30)	40.00 (20)
Fricas	34.00 (17)	66.00 (33)
Mupad	34.00 (17)	66.00 (33)
Giac	34.00 (17)	66.00 (33)
Reduce	32.00 (16)	68.00 (34)
Sympy	20.00 (10)	80.00 (40)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	96.000	0.000	0.000	4.000
Maple	66.000	12.000	16.000	6.000
Mathematica	60.000	0.000	34.000	6.000
Maxima	38.000	18.000	0.000	44.000
Giac	16.000	14.000	0.000	70.000
Fricas	10.000	14.000	6.000	70.000
Sympy	4.000	16.000	0.000	80.000
Mupad	0.000	30.000	0.000	70.000
Reduce	0.000	28.000	0.000	72.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	1	100.00	0.00	0.00
Maple	1	100.00	0.00	0.00
Maxima	20	100.00	0.00	0.00
Fricas	33	100.00	0.00	0.00
Mupad	33	0.00	100.00	0.00
Giac	33	100.00	0.00	0.00
Reduce	34	100.00	0.00	0.00
Sympy	40	67.50	32.50	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.16
Reduce	0.19
Rubi	0.88
Maple	1.54
Sympy	3.00
Giac	3.69
Mupad	4.48
Mathematica	4.91
Fricas	9.77

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Maxima	243.03	2.06	224.50	1.44
Rubi	272.14	1.06	220.00	1.01
Mathematica	362.78	1.56	272.00	1.16
Reduce	468.31	4.91	283.00	2.19
Giac	502.35	3.45	269.00	1.60
Mupad	595.53	2.93	309.00	2.07
Maple	1721.57	3.60	273.00	1.40
Fricas	1884.76	6.98	401.00	2.54
Sympy	2275.90	14.90	708.00	6.78

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{31, 32}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {5, 12, 13, 14, 18, 19, 21, 23, 30, 45, 47, 48}

Maple {12, 15, 16, 17, 18, 19, 20}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	42

2.1 List of integrals sorted by grade for each CAS

Rubi	25
Mma	25
Maple	26
Fricas	26
Maxima	26
Giac	27
Mupad	27
Sympy	27
Reduce	28

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 15, 16, 17, 24, 25, 26, 28, 29, 30, 33, 34, 36, 37, 38, 39, 40, 41, 42, 43, 44 }

B grade { }

C grade { 5, 12, 13, 14, 18, 19, 21, 22, 23, 27, 35, 45, 46, 47, 48, 49, 50 }

F normal fail { 20 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 2, 3, 4, 5, 6, 7, 8, 11, 13, 14, 21, 22, 23, 24, 25, 26, 27, 28, 29, 33, 34, 36, 37, 38, 39, 40, 44, 45, 46, 47, 48, 49, 50 }

B grade { 1, 9, 10, 41, 42, 43 }

C grade { 12, 15, 16, 17, 18, 19, 20, 35 }

F normal fail { 30 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 3, 4, 24, 25, 26 }

B grade { 1, 2, 6, 7, 8, 28, 29 }

C grade { 33, 34, 36 }

F normal fail { 5, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 27, 30, 35, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maxima

A grade { 1, 2, 3, 4, 6, 7, 23, 24, 25, 26, 28, 29, 33, 34, 36, 37, 38, 39, 40 }

B grade { 8, 9, 10, 11, 22, 41, 42, 43, 44 }

C grade { }

F normal fail { 5, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 27, 30, 35, 45, 46, 47, 48, 49, 50 }

F(-1) timedout fail { }

F(-2) exception fail { }

Giac

A grade { 24, 25, 26, 28, 29, 33, 34, 36 }

B grade { 1, 2, 3, 4, 6, 7, 8 }

C grade { }

F normal fail { 5, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 27, 30, 35, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 6, 7, 8, 24, 25, 26, 28, 29, 33, 34, 36 }

C grade { }

F normal fail { }

F(-1) timedout fail { 5, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 27, 30, 35, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50 }

F(-2) exception fail { }

Sympy

A grade { 4, 26 }

B grade { 1, 2, 3, 6, 7, 8, 24, 25 }

C grade { }

F normal fail { 5, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 30, 37, 38, 39, 40, 41, 42, 43, 44, 46, 47 }

F(-1) timedout fail { 27, 28, 29, 31, 32, 33, 34, 35, 36, 45, 48, 49, 50 }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 6, 7, 8, 24, 25, 26, 28, 29, 33, 34 }

C grade { }

F normal fail { 5, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 27, 30, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	149	150	274	295	273	322	381	1576	348	272
N.S.	1	1.01	1.84	1.98	1.83	2.16	2.56	10.58	2.34	1.83
time (sec)	N/A	0.374	0.083	0.351	0.035	0.086	0.499	0.152	0.181	4.210

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	125	122	205	227	209	244	279	970	247	197
N.S.	1	0.98	1.64	1.82	1.67	1.95	2.23	7.76	1.98	1.58
time (sec)	N/A	0.343	0.064	0.191	0.028	0.095	0.501	0.136	0.172	3.798

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	96	99	129	165	137	163	178	532	172	127
N.S.	1	1.03	1.34	1.72	1.43	1.70	1.85	5.54	1.79	1.32
time (sec)	N/A	0.308	0.058	0.158	0.025	0.091	0.346	0.129	0.196	3.581

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	85	96	78	83	98	92	245	85	67
N.S.	1	1.01	1.14	0.93	0.99	1.17	1.10	2.92	1.01	0.80
time (sec)	N/A	0.280	0.018	0.103	0.034	0.089	0.344	0.117	0.175	3.521

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	114	114	257	145	0	0	0	0	30	0
N.S.	1	1.00	2.25	1.27	0.00	0.00	0.00	0.00	0.26	0.00
time (sec)	N/A	0.469	0.174	0.345	0.000	0.000	0.000	0.000	0.204	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	89	102	114	99	182	663	243	223	115
N.S.	1	0.96	1.10	1.23	1.06	1.96	7.13	2.61	2.40	1.24
time (sec)	N/A	0.306	0.090	0.265	0.026	0.102	1.421	0.124	0.217	6.171

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	130	124	133	142	190	454	3216	809	693	427
N.S.	1	0.95	1.02	1.09	1.46	3.49	24.74	6.22	5.33	3.28
time (sec)	N/A	0.370	0.102	0.318	0.033	0.141	3.312	0.129	0.188	6.712

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	174	171	173	184	339	859	10946	1792	1565	418
N.S.	1	0.98	0.99	1.06	1.95	4.94	62.91	10.30	8.99	2.40
time (sec)	N/A	0.428	0.182	0.438	0.045	0.290	5.155	0.165	0.184	5.043

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	359	362	506	804	782	0	0	0	619	0
N.S.	1	1.01	1.41	2.24	2.18	0.00	0.00	0.00	1.72	0.00
time (sec)	N/A	1.210	0.566	0.293	0.204	0.000	0.000	0.000	0.203	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	257	270	319	590	524	0	0	0	400	0
N.S.	1	1.05	1.24	2.30	2.04	0.00	0.00	0.00	1.56	0.00
time (sec)	N/A	1.024	0.398	0.200	0.194	0.000	0.000	0.000	0.183	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	160	177	174	298	313	0	0	0	208	0
N.S.	1	1.11	1.09	1.86	1.96	0.00	0.00	0.00	1.30	0.00
time (sec)	N/A	0.847	0.360	0.147	0.207	0.000	0.000	0.000	0.203	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	188	188	1055	1087	0	0	0	0	55	0
N.S.	1	1.00	5.61	5.78	0.00	0.00	0.00	0.00	0.29	0.00
time (sec)	N/A	0.363	10.044	2.880	0.000	0.000	0.000	0.000	0.183	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	321	310	317	451	0	0	0	0	0	0
N.S.	1	0.97	0.99	1.40	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.694	2.973	0.413	0.000	0.000	0.000	0.000	0.194	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	480	450	470	592	0	0	0	0	0	0
N.S.	1	0.94	0.98	1.23	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.868	4.507	0.746	0.000	0.000	0.000	0.000	0.210	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	614	608	830	5093	0	0	0	0	1081	0
N.S.	1	0.99	1.35	8.29	0.00	0.00	0.00	0.00	1.76	0.00
time (sec)	N/A	1.490	1.240	28.289	0.000	0.000	0.000	0.000	0.187	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	387	403	591	3848	0	0	0	0	696	0
N.S.	1	1.04	1.53	9.94	0.00	0.00	0.00	0.00	1.80	0.00
time (sec)	N/A	1.102	0.767	14.828	0.000	0.000	0.000	0.000	0.185	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	244	259	331	6347	0	0	0	0	345	0
N.S.	1	1.06	1.36	26.01	0.00	0.00	0.00	0.00	1.41	0.00
time (sec)	N/A	0.847	0.671	2.789	0.000	0.000	0.000	0.000	0.180	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	272	272	2160	2160	0	0	0	0	80	0
N.S.	1	1.00	7.94	7.94	0.00	0.00	0.00	0.00	0.29	0.00
time (sec)	N/A	0.376	26.718	2.857	0.000	0.000	0.000	0.000	0.189	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	517	487	1107	3154	0	0	0	0	0	0
N.S.	1	0.94	2.14	6.10	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.947	11.532	2.632	0.000	0.000	0.000	0.000	0.226	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	No	TBD	TBD	TBD	TBD	TBD	TBD
size	953	896	0	50477	0	0	0	0	0	0
N.S.	1	0.94	0.00	52.97	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.175	0.000	10.626	0.000	0.000	0.000	0.000	0.357	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	109	240	92	0	0	0	0	34	0
N.S.	1	1.63	3.58	1.37	0.00	0.00	0.00	0.00	0.51	0.00
time (sec)	N/A	0.657	0.207	0.222	0.000	0.000	0.000	0.000	0.178	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	108	272	127	144	0	0	0	47	0
N.S.	1	1.23	3.09	1.44	1.64	0.00	0.00	0.00	0.53	0.00
time (sec)	N/A	0.474	0.082	0.298	0.111	0.000	0.000	0.000	0.197	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	86	260	106	119	0	0	0	12	0
N.S.	1	1.00	3.02	1.23	1.38	0.00	0.00	0.00	0.14	0.00
time (sec)	N/A	0.406	0.062	0.291	0.026	0.000	0.000	0.000	0.177	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	154	160	254	242	240	519	2786	227	314	823
N.S.	1	1.04	1.65	1.57	1.56	3.37	18.09	1.47	2.04	5.34
time (sec)	N/A	0.527	0.136	0.497	0.114	0.147	7.477	7.467	0.188	4.696

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	158	140	170	199	171	401	3465	171	250	309
N.S.	1	0.89	1.08	1.26	1.08	2.54	21.93	1.08	1.58	1.96
time (sec)	N/A	0.423	0.098	0.369	0.113	0.101	6.063	1.100	0.188	3.940

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	103	104	92	95	249	753	133	118	242
N.S.	1	0.88	0.89	0.79	0.81	2.13	6.44	1.14	1.01	2.07
time (sec)	N/A	0.370	0.047	0.211	0.106	0.102	4.895	0.298	0.185	3.836

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	325	325	285	347	0	0	0	0	32	0
N.S.	1	1.00	0.88	1.07	0.00	0.00	0.00	0.00	0.10	0.00
time (sec)	N/A	0.804	17.327	0.276	0.000	0.000	0.000	0.000	0.189	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	166	168	261	155	172	634	0	269	506	727
N.S.	1	1.01	1.57	0.93	1.04	3.82	0.00	1.62	3.05	4.38
time (sec)	N/A	0.561	0.232	0.504	0.110	2.550	0.000	0.142	0.173	5.028

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	226	223	379	229	311	1639	0	362	1597	2016
N.S.	1	0.99	1.68	1.01	1.38	7.25	0.00	1.60	7.07	8.92
time (sec)	N/A	0.637	0.444	0.744	0.114	22.241	0.000	24.159	0.208	7.637

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	1085	1085	684	0	0	0	0	0	168	0
N.S.	1	1.00	0.63	0.00	0.00	0.00	0.00	0.00	0.15	0.00
time (sec)	N/A	2.088	2.080	0.000	0.000	0.000	0.000	0.000	0.169	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	81	36	0	22	59	22
N.S.	1	1.00	1.10	1.00	4.05	1.80	0.00	1.10	2.95	1.10
time (sec)	N/A	0.214	48.764	0.056	0.197	0.094	0.000	0.167	0.183	3.359

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	323	47	0	22	695	22
N.S.	1	1.00	1.10	1.00	16.15	2.35	0.00	1.10	34.75	1.10
time (sec)	N/A	0.207	85.623	0.056	1.277	0.075	0.000	0.363	0.205	3.487

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	300	326	299	467	295	9282	0	346	369	1081
N.S.	1	1.09	1.00	1.56	0.98	30.94	0.00	1.15	1.23	3.60
time (sec)	N/A	0.753	0.145	0.374	0.119	0.995	0.000	11.459	0.177	3.633

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	250	297	333	363	248	3928	0	267	252	621
N.S.	1	1.19	1.33	1.45	0.99	15.71	0.00	1.07	1.01	2.48
time (sec)	N/A	0.665	0.075	0.233	0.116	0.898	0.000	0.843	0.184	3.527

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	523	519	515	182	0	0	0	0	32	0
N.S.	1	0.99	0.98	0.35	0.00	0.00	0.00	0.00	0.06	0.00
time (sec)	N/A	1.244	10.781	0.326	0.000	0.000	0.000	0.000	0.225	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F(-1)	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	414	410	534	461	394	12984	0	554	20	2638
N.S.	1	0.99	1.29	1.11	0.95	31.36	0.00	1.34	0.05	6.37
time (sec)	N/A	1.059	0.343	0.814	0.113	137.992	0.000	15.723	200.017	3.925

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	195	300	160	247	246	0	0	0	68	0
N.S.	1	1.54	0.82	1.27	1.26	0.00	0.00	0.00	0.35	0.00
time (sec)	N/A	3.168	0.476	0.243	0.195	0.000	0.000	0.000	0.176	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	160	217	130	216	208	0	0	0	59	0
N.S.	1	1.36	0.81	1.35	1.30	0.00	0.00	0.00	0.37	0.00
time (sec)	N/A	1.389	0.353	0.074	0.207	0.000	0.000	0.000	0.182	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	146	96	185	166	0	0	0	47	0
N.S.	1	1.22	0.80	1.54	1.38	0.00	0.00	0.00	0.39	0.00
time (sec)	N/A	0.853	0.229	0.070	0.187	0.000	0.000	0.000	0.189	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	89	75	140	101	0	0	0	39	0
N.S.	1	1.14	0.96	1.79	1.29	0.00	0.00	0.00	0.50	0.00
time (sec)	N/A	0.488	0.137	0.061	0.234	0.000	0.000	0.000	0.186	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	75	74	180	159	0	0	0	40	0
N.S.	1	1.09	1.07	2.61	2.30	0.00	0.00	0.00	0.58	0.00
time (sec)	N/A	0.572	0.180	0.091	0.170	0.000	0.000	0.000	0.175	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	123	118	243	248	0	0	0	58	0
N.S.	1	1.05	1.01	2.08	2.12	0.00	0.00	0.00	0.50	0.00
time (sec)	N/A	0.901	0.301	0.098	0.192	0.000	0.000	0.000	0.161	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	185	158	273	291	0	0	0	123	0
N.S.	1	1.18	1.01	1.74	1.85	0.00	0.00	0.00	0.78	0.00
time (sec)	N/A	1.274	0.454	0.103	0.205	0.000	0.000	0.000	0.164	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	192	261	187	303	330	0	0	0	157	0
N.S.	1	1.36	0.97	1.58	1.72	0.00	0.00	0.00	0.82	0.00
time (sec)	N/A	1.733	0.600	0.104	0.205	0.000	0.000	0.000	0.165	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	460	459	558	564	0	0	0	0	56	0
N.S.	1	1.00	1.21	1.23	0.00	0.00	0.00	0.00	0.12	0.00
time (sec)	N/A	1.459	1.862	0.184	0.000	0.000	0.000	0.000	0.171	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	374	387	463	494	0	0	0	0	40	0
N.S.	1	1.03	1.24	1.32	0.00	0.00	0.00	0.00	0.11	0.00
time (sec)	N/A	0.925	1.648	0.066	0.000	0.000	0.000	0.000	0.183	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	318	329	432	433	0	0	0	0	31	0
N.S.	1	1.03	1.36	1.36	0.00	0.00	0.00	0.00	0.10	0.00
time (sec)	N/A	0.843	1.165	0.062	0.000	0.000	0.000	0.000	0.168	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	358	374	445	499	0	0	0	0	40	0
N.S.	1	1.04	1.24	1.39	0.00	0.00	0.00	0.00	0.11	0.00
time (sec)	N/A	1.357	1.499	0.092	0.000	0.000	0.000	0.000	0.183	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	413	429	479	577	0	0	0	0	56	0
N.S.	1	1.04	1.16	1.40	0.00	0.00	0.00	0.00	0.14	0.00
time (sec)	N/A	1.670	1.936	0.100	0.000	0.000	0.000	0.000	0.188	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	506	498	538	650	0	0	0	0	990	0
N.S.	1	0.98	1.06	1.28	0.00	0.00	0.00	0.00	1.96	0.00
time (sec)	N/A	1.558	2.624	0.117	0.000	0.000	0.000	0.000	0.190	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [44] had the largest ratio of [.769230999999999998]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	3	3	1.01	16	0.188
2	A	3	3	0.98	16	0.188
3	A	3	3	1.03	16	0.188
4	A	3	3	1.01	14	0.214
5	A	5	4	1.00	16	0.250
6	A	3	3	0.96	16	0.188
7	A	3	3	0.95	16	0.188
8	A	3	3	0.98	16	0.188
9	A	2	2	1.01	18	0.111
10	A	2	2	1.05	18	0.111
11	A	2	2	1.11	16	0.125
12	A	1	1	1.00	18	0.056
13	A	2	2	0.97	18	0.111
14	A	2	2	0.94	18	0.111
15	A	2	2	0.99	18	0.111
16	A	2	2	1.04	18	0.111
17	A	2	2	1.06	16	0.125
18	A	1	1	1.00	18	0.056
19	A	2	2	0.94	18	0.111
20	A	2	2	0.94	18	0.111
21	A	5	4	1.63	17	0.235

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	5	4	1.23	15	0.267
23	A	5	4	1.00	10	0.400
24	A	3	3	1.04	18	0.167
25	A	3	3	0.89	18	0.167
26	A	3	3	0.88	16	0.188
27	A	3	3	1.00	18	0.167
28	A	3	3	1.01	18	0.167
29	A	3	3	0.99	18	0.167
30	A	2	2	1.00	18	0.111
31	N/A	1	0	1.00	20	0.000
32	N/A	1	0	1.00	20	0.000
33	A	3	3	1.09	18	0.167
34	A	3	3	1.19	16	0.188
35	A	3	3	0.99	18	0.167
36	A	3	3	0.99	18	0.167
37	A	18	17	1.54	26	0.654
38	A	14	13	1.36	26	0.500
39	A	10	9	1.22	24	0.375
40	A	6	5	1.14	23	0.217
41	A	6	5	1.09	26	0.192
42	A	10	9	1.05	26	0.346
43	A	15	14	1.18	26	0.538
44	A	21	20	1.36	26	0.769
45	A	12	11	1.00	23	0.478
46	A	8	7	1.03	21	0.333
47	A	4	3	1.03	20	0.150
48	A	5	4	1.04	23	0.174
49	A	9	8	1.04	23	0.348
50	A	14	13	0.98	23	0.565

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int (d + ex)^4 (a + \operatorname{barctanh}(cx)) dx$	46
3.2	$\int (d + ex)^3 (a + \operatorname{barctanh}(cx)) dx$	54
3.3	$\int (d + ex)^2 (a + \operatorname{barctanh}(cx)) dx$	62
3.4	$\int (d + ex) (a + \operatorname{barctanh}(cx)) dx$	69
3.5	$\int \frac{a + \operatorname{barctanh}(cx)}{d + ex} dx$	75
3.6	$\int \frac{a + \operatorname{barctanh}(cx)}{(d + ex)^2} dx$	82
3.7	$\int \frac{a + \operatorname{barctanh}(cx)}{(d + ex)^3} dx$	89
3.8	$\int \frac{a + \operatorname{barctanh}(cx)}{(d + ex)^4} dx$	97
3.9	$\int (d + ex)^3 (a + \operatorname{barctanh}(cx))^2 dx$	107
3.10	$\int (d + ex)^2 (a + \operatorname{barctanh}(cx))^2 dx$	117
3.11	$\int (d + ex) (a + \operatorname{barctanh}(cx))^2 dx$	125
3.12	$\int \frac{(a + \operatorname{barctanh}(cx))^2}{d + ex} dx$	132
3.13	$\int \frac{(a + \operatorname{barctanh}(cx))^2}{(d + ex)^2} dx$	139
3.14	$\int \frac{(a + \operatorname{barctanh}(cx))^2}{(d + ex)^3} dx$	146
3.15	$\int (d + ex)^3 (a + \operatorname{barctanh}(cx))^3 dx$	154
3.16	$\int (d + ex)^2 (a + \operatorname{barctanh}(cx))^3 dx$	163
3.17	$\int (d + ex) (a + \operatorname{barctanh}(cx))^3 dx$	171
3.18	$\int \frac{(a + \operatorname{barctanh}(cx))^3}{d + ex} dx$	178
3.19	$\int \frac{(a + \operatorname{barctanh}(cx))^3}{(d + ex)^2} dx$	185
3.20	$\int \frac{(a + \operatorname{barctanh}(cx))^3}{(d + ex)^3} dx$	194
3.21	$\int \frac{a + \operatorname{barctanh}(cx)}{1 + 2cx} dx$	202
3.22	$\int \frac{\operatorname{arctanh}(x)}{1 - \sqrt{2}x} dx$	208
3.23	$\int \frac{\operatorname{arctanh}(x)}{a + bx} dx$	215

3.24	$\int (d + ex)^3 (a + \operatorname{barctanh}(cx^2)) dx$	221
3.25	$\int (d + ex)^2 (a + \operatorname{barctanh}(cx^2)) dx$	231
3.26	$\int (d + ex) (a + \operatorname{barctanh}(cx^2)) dx$	240
3.27	$\int \frac{a + \operatorname{barctanh}(cx^2)}{d + ex} dx$	248
3.28	$\int \frac{a + \operatorname{barctanh}(cx^2)}{(d + ex)^2} dx$	255
3.29	$\int \frac{a + \operatorname{barctanh}(cx^2)}{(d + ex)^3} dx$	265
3.30	$\int (d + ex) (a + \operatorname{barctanh}(cx^2))^2 dx$	274
3.31	$\int \frac{(a + \operatorname{barctanh}(cx^2))^2}{d + ex} dx$	282
3.32	$\int \frac{(a + \operatorname{barctanh}(cx^2))^2}{(d + ex)^2} dx$	287
3.33	$\int (d + ex)^2 (a + \operatorname{barctanh}(cx^3)) dx$	292
3.34	$\int (d + ex) (a + \operatorname{barctanh}(cx^3)) dx$	302
3.35	$\int \frac{a + \operatorname{barctanh}(cx^3)}{d + ex} dx$	311
3.36	$\int \frac{a + \operatorname{barctanh}(cx^3)}{(d + ex)^2} dx$	319
3.37	$\int \frac{x^3 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} dx$	329
3.38	$\int \frac{x^2 (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} dx$	339
3.39	$\int \frac{x (a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2 x} dx$	348
3.40	$\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{1 - c^2 x} dx$	356
3.41	$\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x(1 - c^2 x)} dx$	362
3.42	$\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^2(1 - c^2 x)} dx$	368
3.43	$\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^3(1 - c^2 x)} dx$	376
3.44	$\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^4(1 - c^2 x)} dx$	385
3.45	$\int \frac{x^2 (a + \operatorname{barctanh}(c\sqrt{x}))}{d + ex} dx$	395
3.46	$\int \frac{x (a + \operatorname{barctanh}(c\sqrt{x}))}{d + ex} dx$	405
3.47	$\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{d + ex} dx$	413
3.48	$\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x(d + ex)} dx$	420
3.49	$\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^2(d + ex)} dx$	427
3.50	$\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^3(d + ex)} dx$	436

3.1 $\int (d + ex)^4 (a + \operatorname{barctanh}(cx)) dx$

Optimal result	46
Mathematica [A] (verified)	46
Rubi [A] (verified)	47
Maple [B] (verified)	48
Fricas [B] (verification not implemented)	49
Sympy [B] (verification not implemented)	50
Maxima [A] (verification not implemented)	51
Giac [B] (verification not implemented)	51
Mupad [B] (verification not implemented)	52
Reduce [B] (verification not implemented)	53

Optimal result

Integrand size = 16, antiderivative size = 149

$$\int (d + ex)^4 (a + \operatorname{barctanh}(cx)) dx = \frac{bde(2c^2d^2 + e^2)x}{c^3} + \frac{be^2(10c^2d^2 + e^2)x^2}{10c^3} + \frac{bde^3x^3}{3c} + \frac{be^4x^4}{20c} + \frac{(d + ex)^5(a + \operatorname{barctanh}(cx))}{5e} + \frac{b(cd + e)^5 \log(1 - cx)}{10c^5e} - \frac{b(cd - e)^5 \log(1 + cx)}{10c^5e}$$

```
output b*d*e*(2*c^2*d^2+e^2)*x/c^3+1/10*b*e^2*(10*c^2*d^2+e^2)*x^2/c^3+1/3*b*d*e^3*x^3/c+1/20*b*e^4*x^4/c+1/5*(e*x+d)^5*(a+b*arctanh(c*x))/e+1/10*b*(c*d+e)^5*ln(-c*x+1)/c^5/e-1/10*b*(c*d-e)^5*ln(c*x+1)/c^5/e
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 274, normalized size of antiderivative = 1.84

$$\int (d + ex)^4 (a + \operatorname{barctanh}(cx)) dx = \frac{60c^2d(ac^3d^3 + be(2c^2d^2 + e^2))x + 6c^2e(20ac^3d^3 + be(10c^2d^2 + e^2))x^2 + 20c^4de^2(6acd + be)x^3 + 3c^4e^3(2$$

```
input Integrate[(d + e*x)^4*(a + b*ArcTanh[c*x]),x]
```

output

```
(60*c^2*d*(a*c^3*d^3 + b*e*(2*c^2*d^2 + e^2))*x + 6*c^2*e*(20*a*c^3*d^3 +
b*e*(10*c^2*d^2 + e^2))*x^2 + 20*c^4*d*e^2*(6*a*c*d + b*e)*x^3 + 3*c^4*e^3
*(20*a*c*d + b*e)*x^4 + 12*a*c^5*e^4*x^5 + 12*b*c^5*x*(5*d^4 + 10*d^3*e*x
+ 10*d^2*e^2*x^2 + 5*d*e^3*x^3 + e^4*x^4)*ArcTanh[c*x] + 6*b*(5*c^4*d^4 +
10*c^3*d^3*e + 10*c^2*d^2*e^2 + 5*c*d*e^3 + e^4)*Log[1 - c*x] + 6*b*(5*c^4
*d^4 - 10*c^3*d^3*e + 10*c^2*d^2*e^2 - 5*c*d*e^3 + e^4)*Log[1 + c*x])/(60*
c^5)
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.01, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6478, 477, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (d + ex)^4 (a + \operatorname{barctanh}(cx)) dx \\
 & \quad \downarrow 6478 \\
 & \frac{(d + ex)^5 (a + \operatorname{barctanh}(cx))}{5e} - \frac{bc \int \frac{(d+ex)^5}{1-c^2x^2} dx}{5e} \\
 & \quad \downarrow 477 \\
 & \frac{(d + ex)^5 (a + \operatorname{barctanh}(cx))}{5e} - \\
 & \frac{bc \int \left(\frac{(cd-e)^5}{2c^5(cx+1)} - \frac{e^5x^3}{c^2} - \frac{5de^4x^2}{c^2} - \frac{5de^2(2c^2d^2+e^2)}{c^4} - \frac{e^3(10c^2d^2+e^2)x}{c^4} + \frac{(cd+e)^5}{2c^5(1-cx)} \right) dx}{5e} \\
 & \quad \downarrow 2009 \\
 & \frac{(d + ex)^5 (a + \operatorname{barctanh}(cx))}{5e} - \\
 & \frac{bc \left(\frac{(cd-e)^5 \log(cx+1)}{2c^6} - \frac{(cd+e)^5 \log(1-cx)}{2c^6} - \frac{5de^4x^3}{3c^2} - \frac{e^5x^4}{4c^2} - \frac{5de^2x(2c^2d^2+e^2)}{c^4} - \frac{e^3x^2(10c^2d^2+e^2)}{2c^4} \right)}{5e}
 \end{aligned}$$

input

```
Int[(d + e*x)^4*(a + b*ArcTanh[c*x]), x]
```


output

$$\frac{((d + e*x)^5*(a + b*\text{ArcTanh}[c*x]))/(5*e) - (b*c*((-5*d*e^2*(2*c^2*d^2 + e^2)*x)/c^4 - (e^3*(10*c^2*d^2 + e^2)*x^2)/(2*c^4) - (5*d*e^4*x^3)/(3*c^2) - (e^5*x^4)/(4*c^2) - ((c*d + e)^5*\text{Log}[1 - c*x])/(2*c^6) + ((c*d - e)^5*\text{Log}[1 + c*x])/(2*c^6)))/(5*e)}$$
Defintions of rubi rules used

rule 477

```
Int[((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[
a^p Int[ExpandIntegrand[(c + d*x)^n*(1 - Rt[-b/a, 2]*x)^p*(1 + Rt[-b/a, 2]
*x)^p, x], x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IntegerQ[n] &
& NiceSqrtQ[-b/a] && !FractionalPowerFactorQ[Rt[-b/a, 2]]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6478

```
Int[((a_) + ArcTanh[(c_)*(x_)])*(b_))*((d_) + (e_)*(x_))^(q_), x_Symbol
] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])/(e*(q + 1))), x] - Simp[b
*(c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 - c^2*x^2), x], x] /; FreeQ[{a,
b, c, d, e, q}, x] && NeQ[q, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 294 vs. $2(137) = 274$.

Time = 0.35 (sec) , antiderivative size = 295, normalized size of antiderivative = 1.98

method	result
parts	$\frac{a(ex+d)^5}{5e} + \frac{b \left(\frac{ce^4 \operatorname{arctanh}(cx)x^5}{5} + ce^3 \operatorname{arctanh}(cx)x^4d + 2ce^2 \operatorname{arctanh}(cx)x^3d^2 + 2ce \operatorname{arctanh}(cx)x^2d^3 + \operatorname{arctanh}(cx) \right)}{5e}$
derivativdivides	$\frac{a(cex+cd)^5}{5c^4e} + \frac{b \left(\frac{\operatorname{arctanh}(cx)e^5d^5}{5e} + \operatorname{arctanh}(cx)e^5d^4x + 2e \operatorname{arctanh}(cx)e^5d^3x^2 + 2e^2 \operatorname{arctanh}(cx)e^5d^2x^3 + e^3 \operatorname{arctanh}(cx)e^5dx^4 + \frac{e^4}{5} \right)}{5c^4e}$
default	$\frac{a(cex+cd)^5}{5c^4e} + \frac{b \left(\frac{\operatorname{arctanh}(cx)e^5d^5}{5e} + \operatorname{arctanh}(cx)e^5d^4x + 2e \operatorname{arctanh}(cx)e^5d^3x^2 + 2e^2 \operatorname{arctanh}(cx)e^5d^2x^3 + e^3 \operatorname{arctanh}(cx)e^5dx^4 + \frac{e^4}{5} \right)}{5c^4e}$
parallelrisc	$6be^4 + 60b^2c^2d^2e^2 + 120ac^3d^3e + 12 \operatorname{arctanh}(cx)be^4 + 12 \ln(cx-1)be^4 + 60 \ln(cx-1)bc^4d^4 + 60xac^5d^4 + 3x^4bc^4e^4 + 12x^5$
risc	$\frac{be^4x^4}{20c} + \frac{bde^3x^3}{3c} + e^3ad^4x^4 + 2e^2ad^2x^3 + 2ead^3x^2 + ad^4x + \frac{e^2bd^2x^2}{c} + \frac{2ebd^3x}{c} + \frac{e^3bdx}{c^3} - \frac{e^4}{c^3}$

```
input int((e*x+d)^4*(a+b*arctanh(c*x)),x,method=_RETURNVERBOSE)
```

```
output 1/5*a*(e*x+d)^5/e+b/c*(1/5*c*e^4*arctanh(c*x)*x^5+c*e^3*arctanh(c*x)*x^4*d
+2*c*e^2*arctanh(c*x)*x^3*d^2+2*c*e*arctanh(c*x)*x^2*d^3+arctanh(c*x)*c*x*
d^4+1/5*c/e*arctanh(c*x)*d^5-1/5/c^4/e*(-1/4*e^5*c^4*x^4-5*c^4*d^2*e^3*x^2
-5/3*c^4*d*e^4*x^3-10*c^4*d^3*e^2*x-5*c^2*d*e^4*x+1/2*(c^5*d^5-5*c^4*d^4*e
+10*c^3*d^3*e^2-10*c^2*d^2*e^3+5*c*d*e^4-e^5)*ln(c*x+1)-1/2*(c^5*d^5+5*c^4
*d^4*e+10*c^3*d^3*e^2+10*c^2*d^2*e^3+5*c*d*e^4+e^5)*ln(c*x-1)-1/2*e^5*c^2*
x^2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 322 vs. 2(137) = 274.

Time = 0.09 (sec) , antiderivative size = 322, normalized size of antiderivative = 2.16

$$\int (d + ex)^4(a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{12ac^5e^4x^5 + 3(20ac^5de^3 + bc^4e^4)x^4 + 20(6ac^5d^2e^2 + bc^4de^3)x^3 + 6(20ac^5d^3e + 10bc^4d^2e^2 + bc^2e^4)x^2 + \dots}{1}$$

```
input integrate((e*x+d)^4*(a+b*arctanh(c*x)),x, algorithm="fricas")
```

output

```
1/60*(12*a*c^5*e^4*x^5 + 3*(20*a*c^5*d*e^3 + b*c^4*e^4)*x^4 + 20*(6*a*c^5*
d^2*e^2 + b*c^4*d*e^3)*x^3 + 6*(20*a*c^5*d^3*e + 10*b*c^4*d^2*e^2 + b*c^2*
e^4)*x^2 + 60*(a*c^5*d^4 + 2*b*c^4*d^3*e + b*c^2*d*e^3)*x + 6*(5*b*c^4*d^4
- 10*b*c^3*d^3*e + 10*b*c^2*d^2*e^2 - 5*b*c*d*e^3 + b*e^4)*log(c*x + 1) +
6*(5*b*c^4*d^4 + 10*b*c^3*d^3*e + 10*b*c^2*d^2*e^2 + 5*b*c*d*e^3 + b*e^4)
*log(c*x - 1) + 6*(b*c^5*e^4*x^5 + 5*b*c^5*d*e^3*x^4 + 10*b*c^5*d^2*e^2*x^
3 + 10*b*c^5*d^3*e*x^2 + 5*b*c^5*d^4*x)*log(-(c*x + 1)/(c*x - 1))/c^5
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 381 vs. $2(134) = 268$.

Time = 0.50 (sec) , antiderivative size = 381, normalized size of antiderivative = 2.56

$$\int (d + ex)^4 (a + b \operatorname{arctanh}(cx)) dx$$

$$= \begin{cases} ad^4x + 2ad^3ex^2 + 2ad^2e^2x^3 + ade^3x^4 + \frac{ae^4x^5}{5} + bd^4x \operatorname{atanh}(cx) + 2bd^3ex^2 \operatorname{atanh}(cx) + 2bd^2e^2x^3 \operatorname{atanh}(cx) \\ a\left(d^4x + 2d^3ex^2 + 2d^2e^2x^3 + de^3x^4 + \frac{e^4x^5}{5}\right) \end{cases}$$

input

```
integrate((e*x+d)**4*(a+b*atanh(c*x)),x)
```

output

```
Piecewise((a*d**4*x + 2*a*d**3*e*x**2 + 2*a*d**2*e**2*x**3 + a*d*e**3*x**4
+ a*e**4*x**5/5 + b*d**4*x*atanh(c*x) + 2*b*d**3*e*x**2*atanh(c*x) + 2*b*
d**2*e**2*x**3*atanh(c*x) + b*d*e**3*x**4*atanh(c*x) + b*e**4*x**5*atanh(c
*x)/5 + b*d**4*log(x - 1/c)/c + b*d**4*atanh(c*x)/c + 2*b*d**3*e*x/c + b*d
**2*e**2*x**2/c + b*d*e**3*x**3/(3*c) + b*e**4*x**4/(20*c) - 2*b*d**3*e*at
anh(c*x)/c**2 + 2*b*d**2*e**2*log(x - 1/c)/c**3 + 2*b*d**2*e**2*atanh(c*x)
/c**3 + b*d*e**3*x/c**3 + b*e**4*x**2/(10*c**3) - b*d*e**3*atanh(c*x)/c**4
+ b*e**4*log(x - 1/c)/(5*c**5) + b*e**4*atanh(c*x)/(5*c**5), Ne(c, 0)), (
a*(d**4*x + 2*d**3*e*x**2 + 2*d**2*e**2*x**3 + d*e**3*x**4 + e**4*x**5/5),
True))
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.83

$$\int (d + ex)^4 (a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{1}{5} a e^4 x^5 + a d e^3 x^4 + 2 a d^2 e^2 x^3 + 2 a d^3 e x^2$$

$$+ \left(2 x^2 \operatorname{artanh}(cx) + c \left(\frac{2x}{c^2} - \frac{\log(cx+1)}{c^3} + \frac{\log(cx-1)}{c^3} \right) \right) b d^3 e$$

$$+ \left(2 x^3 \operatorname{artanh}(cx) + c \left(\frac{x^2}{c^2} + \frac{\log(c^2 x^2 - 1)}{c^4} \right) \right) b d^2 e^2$$

$$+ \frac{1}{6} \left(6 x^4 \operatorname{artanh}(cx) + c \left(\frac{2(c^2 x^3 + 3x)}{c^4} - \frac{3 \log(cx+1)}{c^5} + \frac{3 \log(cx-1)}{c^5} \right) \right) b d e^3$$

$$+ \frac{1}{20} \left(4 x^5 \operatorname{artanh}(cx) + c \left(\frac{c^2 x^4 + 2x^2}{c^4} + \frac{2 \log(c^2 x^2 - 1)}{c^6} \right) \right) b e^4$$

$$+ a d^4 x + \frac{(2 c x \operatorname{artanh}(cx) + \log(-c^2 x^2 + 1)) b d^4}{2 c}$$

input `integrate((e*x+d)^4*(a+b*arctanh(c*x)),x, algorithm="maxima")`

output `1/5*a*e^4*x^5 + a*d*e^3*x^4 + 2*a*d^2*e^2*x^3 + 2*a*d^3*e*x^2 + (2*x^2*arc
tanh(c*x) + c*(2*x/c^2 - log(c*x + 1)/c^3 + log(c*x - 1)/c^3))*b*d^3*e + (
2*x^3*arctanh(c*x) + c*(x^2/c^2 + log(c^2*x^2 - 1)/c^4))*b*d^2*e^2 + 1/6*(
6*x^4*arctanh(c*x) + c*(2*(c^2*x^3 + 3*x)/c^4 - 3*log(c*x + 1)/c^5 + 3*log
(c*x - 1)/c^5))*b*d*e^3 + 1/20*(4*x^5*arctanh(c*x) + c*((c^2*x^4 + 2*x^2)/
c^4 + 2*log(c^2*x^2 - 1)/c^6))*b*e^4 + a*d^4*x + 1/2*(2*c*x*arctanh(c*x) +
log(-c^2*x^2 + 1))*b*d^4/c`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1576 vs. 2(137) = 274.

Time = 0.15 (sec) , antiderivative size = 1576, normalized size of antiderivative = 10.58

$$\int (d + ex)^4 (a + b \operatorname{arctanh}(cx)) dx = \text{Too large to display}$$

input `integrate((e*x+d)^4*(a+b*arctanh(c*x)),x, algorithm="giac")`

output

```

1/15*c*(3*(5*(c*x + 1)^4*b*c^4*d^4/(c*x - 1)^4 - 20*(c*x + 1)^3*b*c^4*d^4/
(c*x - 1)^3 + 30*(c*x + 1)^2*b*c^4*d^4/(c*x - 1)^2 - 20*(c*x + 1)*b*c^4*d^
4/(c*x - 1) + 5*b*c^4*d^4 + 20*(c*x + 1)^4*b*c^3*d^3*e/(c*x - 1)^4 - 60*(c
*x + 1)^3*b*c^3*d^3*e/(c*x - 1)^3 + 60*(c*x + 1)^2*b*c^3*d^3*e/(c*x - 1)^2
- 20*(c*x + 1)*b*c^3*d^3*e/(c*x - 1) + 30*(c*x + 1)^4*b*c^2*d^2*e^2/(c*x
- 1)^4 - 60*(c*x + 1)^3*b*c^2*d^2*e^2/(c*x - 1)^3 + 40*(c*x + 1)^2*b*c^2*d
^2*e^2/(c*x - 1)^2 - 20*(c*x + 1)*b*c^2*d^2*e^2/(c*x - 1) + 10*b*c^2*d^2*e
^2 + 20*(c*x + 1)^4*b*c*d*e^3/(c*x - 1)^4 - 20*(c*x + 1)^3*b*c*d*e^3/(c*x
- 1)^3 + 20*(c*x + 1)^2*b*c*d*e^3/(c*x - 1)^2 - 20*(c*x + 1)*b*c*d*e^3/(c
*x - 1) + 5*(c*x + 1)^4*b*e^4/(c*x - 1)^4 + 10*(c*x + 1)^2*b*e^4/(c*x - 1)^
2 + b*e^4)*log(-(c*x + 1)/(c*x - 1))/((c*x + 1)^5*c^6/(c*x - 1)^5 - 5*(c*x
+ 1)^4*c^6/(c*x - 1)^4 + 10*(c*x + 1)^3*c^6/(c*x - 1)^3 - 10*(c*x + 1)^2*
c^6/(c*x - 1)^2 + 5*(c*x + 1)*c^6/(c*x - 1) - c^6) + 2*(15*(c*x + 1)^4*a*c
^4*d^4/(c*x - 1)^4 - 60*(c*x + 1)^3*a*c^4*d^4/(c*x - 1)^3 + 90*(c*x + 1)^2
*a*c^4*d^4/(c*x - 1)^2 - 60*(c*x + 1)*a*c^4*d^4/(c*x - 1) + 15*a*c^4*d^4 +
60*(c*x + 1)^4*a*c^3*d^3*e/(c*x - 1)^4 - 180*(c*x + 1)^3*a*c^3*d^3*e/(c*x
- 1)^3 + 180*(c*x + 1)^2*a*c^3*d^3*e/(c*x - 1)^2 - 60*(c*x + 1)*a*c^3*d^3
*e/(c*x - 1) + 30*(c*x + 1)^4*b*c^3*d^3*e/(c*x - 1)^4 - 120*(c*x + 1)^3*b*
c^3*d^3*e/(c*x - 1)^3 + 180*(c*x + 1)^2*b*c^3*d^3*e/(c*x - 1)^2 - 120*(c*x
+ 1)*b*c^3*d^3*e/(c*x - 1) + 30*b*c^3*d^3*e + 90*(c*x + 1)^4*a*c^2*d^2...

```

Mupad [B] (verification not implemented)

Time = 4.21 (sec) , antiderivative size = 272, normalized size of antiderivative = 1.83

$$\begin{aligned}
\int (d + ex)^4(a + b \operatorname{arctanh}(cx)) dx &= \frac{a e^4 x^5}{5} + a d^4 x + \frac{b d^4 \ln(c^2 x^2 - 1)}{2c} \\
&+ \frac{b e^4 \ln(c^2 x^2 - 1)}{10 c^5} + 2 a d^2 e^2 x^3 + \frac{b e^4 x^4}{20 c} \\
&+ \frac{b e^4 x^2}{10 c^3} + b d^4 x \operatorname{atanh}(cx) + 2 a d^3 e x^2 \\
&+ a d e^3 x^4 + \frac{b e^4 x^5 \operatorname{atanh}(cx)}{5} + \frac{2 b d^3 e x}{c} \\
&+ \frac{b d e^3 x}{c^3} - \frac{2 b d^3 e \operatorname{atanh}(cx)}{c^2} - \frac{b d e^3 \operatorname{atanh}(cx)}{c^4} \\
&+ 2 b d^3 e x^2 \operatorname{atanh}(cx) + b d e^3 x^4 \operatorname{atanh}(cx) \\
&+ \frac{b d e^3 x^3}{3 c} + 2 b d^2 e^2 x^3 \operatorname{atanh}(cx) \\
&+ \frac{b d^2 e^2 \ln(c^2 x^2 - 1)}{c^3} + \frac{b d^2 e^2 x^2}{c}
\end{aligned}$$

3.2 $\int (d + ex)^3 (a + b \operatorname{arctanh}(cx)) dx$

Optimal result	54
Mathematica [A] (verified)	54
Rubi [A] (verified)	55
Maple [A] (verified)	56
Fricas [B] (verification not implemented)	57
Sympy [B] (verification not implemented)	57
Maxima [A] (verification not implemented)	58
Giac [B] (verification not implemented)	59
Mupad [B] (verification not implemented)	60
Reduce [B] (verification not implemented)	60

Optimal result

Integrand size = 16, antiderivative size = 125

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx)) dx = \frac{be(6c^2d^2 + e^2)x}{4c^3} + \frac{bde^2x^2}{2c} + \frac{be^3x^3}{12c} + \frac{(d + ex)^4 (a + b \operatorname{arctanh}(cx))}{4e} + \frac{b(cd + e)^4 \log(1 - cx)}{8c^4e} - \frac{b(cd - e)^4 \log(1 + cx)}{8c^4e}$$

output

```
1/4*b*e*(6*c^2*d^2+e^2)*x/c^3+1/2*b*d*e^2*x^2/c+1/12*b*e^3*x^3/c+1/4*(e*x+d)^4*(a+b*arctanh(c*x))/e+1/8*b*(c*d+e)^4*ln(-c*x+1)/c^4/e-1/8*b*(c*d-e)^4*ln(c*x+1)/c^4/e
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.64

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx)) dx = \frac{6c(4ac^3d^3 + be(6c^2d^2 + e^2))x + 12c^3de(3acd + be)x^2 + 2c^3e^2(12acd + be)x^3 + 6ac^4e^3x^4 + 6bc^4x(4d^3 + 6d^2e + 2de^2 + e^3)}{12c^4e}$$

input

```
Integrate[(d + e*x)^3*(a + b*ArcTanh[c*x]),x]
```

output

$$(6*c*(4*a*c^3*d^3 + b*e*(6*c^2*d^2 + e^2))*x + 12*c^3*d*e*(3*a*c*d + b*e)*x^2 + 2*c^3*e^2*(12*a*c*d + b*e)*x^3 + 6*a*c^4*e^3*x^4 + 6*b*c^4*x*(4*d^3 + 6*d^2*e*x + 4*d*e^2*x^2 + e^3*x^3)*ArcTanh[c*x] + 3*b*(4*c^3*d^3 + 6*c^2*d^2*e + 4*c*d*e^2 + e^3)*Log[1 - c*x] + 3*b*(4*c^3*d^3 - 6*c^2*d^2*e + 4*c*d*e^2 - e^3)*Log[1 + c*x])/(24*c^4)$$

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.98, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6478, 477, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx)) dx$$

$$\downarrow 6478$$

$$\frac{(d + ex)^4 (a + b \operatorname{arctanh}(cx))}{4e} - \frac{bc \int \frac{(d+ex)^4}{1-c^2x^2} dx}{4e}$$

$$\downarrow 477$$

$$\frac{(d + ex)^4 (a + b \operatorname{arctanh}(cx))}{4e} - \frac{bc \int \left(\frac{(cd-e)^4}{2c^4(cx+1)} - \frac{e^4x^2}{c^2} - \frac{e^2(6c^2d^2+e^2)}{c^4} - \frac{4de^3x}{c^2} + \frac{(cd+e)^4}{2c^4(1-cx)} \right) dx}{4e}$$

$$\downarrow 2009$$

$$\frac{(d + ex)^4 (a + b \operatorname{arctanh}(cx))}{4e} - \frac{bc \left(\frac{(cd-e)^4 \log(cx+1)}{2c^5} - \frac{(cd+e)^4 \log(1-cx)}{2c^5} - \frac{2de^3x^2}{c^2} - \frac{e^4x^3}{3c^2} - \frac{e^2x(6c^2d^2+e^2)}{c^4} \right)}{4e}$$

input

```
Int[(d + e*x)^3*(a + b*ArcTanh[c*x]), x]
```

output

$$\frac{((d + e*x)^4*(a + b*ArcTanh[c*x]))}{(4*e)} - \frac{(b*c*(-((e^2*(6*c^2*d^2 + e^2))*x)/c^4) - (2*d*e^3*x^2)/c^2 - (e^4*x^3)/(3*c^2) - ((c*d + e)^4*Log[1 - c*x])/(2*c^5) + ((c*d - e)^4*Log[1 + c*x])/(2*c^5))}{(4*e)}$$

output

```
1/4*a*(e*x+d)^4/e+b/c*(1/4*c*e^3*arctanh(c*x)*x^4+c*e^2*arctanh(c*x)*x^3*d
+3/2*c*e*arctanh(c*x)*x^2*d^2+arctanh(c*x)*c*x*d^3+1/4*c/e*arctanh(c*x)*d^
4-1/4/c^3/e*(-6*c^3*d^2*e^2*x-2*c^3*d*e^3*x^2-1/3*e^4*c^3*x^3-e^4*c*x-1/2*
(c^4*d^4+4*c^3*d^3*e+6*c^2*d^2*e^2+4*c*d*e^3+e^4)*ln(c*x-1)+1/2*(c^4*d^4-4
*c^3*d^3*e+6*c^2*d^2*e^2-4*c*d*e^3+e^4)*ln(c*x+1))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 244 vs. $2(113) = 226$.

Time = 0.10 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.95

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{6ac^4e^3x^4 + 2(12ac^4de^2 + bc^3e^3)x^3 + 12(3ac^4d^2e + bc^3de^2)x^2 + 6(4ac^4d^3 + 6bc^3d^2e + bce^3)x + 3(4bc^3d^4 + 4ac^4d^3 + 6bc^3d^2e + bce^3)}{c^4}$$

input

```
integrate((e*x+d)^3*(a+b*arctanh(c*x)),x, algorithm="fricas")
```

output

```
1/24*(6*a*c^4*e^3*x^4 + 2*(12*a*c^4*d*e^2 + b*c^3*e^3)*x^3 + 12*(3*a*c^4*d
^2*e + b*c^3*d*e^2)*x^2 + 6*(4*a*c^4*d^3 + 6*b*c^3*d^2*e + b*c*e^3)*x + 3*
(4*b*c^3*d^3 - 6*b*c^2*d^2*e + 4*b*c*d*e^2 - b*e^3)*log(c*x + 1) + 3*(4*b*
c^3*d^3 + 6*b*c^2*d^2*e + 4*b*c*d*e^2 + b*e^3)*log(c*x - 1) + 3*(b*c^4*e^3
*x^4 + 4*b*c^4*d*e^2*x^3 + 6*b*c^4*d^2*e*x^2 + 4*b*c^4*d^3*x)*log(-(c*x +
1)/(c*x - 1))/c^4
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 279 vs. $2(109) = 218$.

Time = 0.50 (sec) , antiderivative size = 279, normalized size of antiderivative = 2.23

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx)) dx$$

$$= \begin{cases} ad^3x + \frac{3ad^2ex^2}{2} + ade^2x^3 + \frac{ae^3x^4}{4} + bd^3x \operatorname{atanh}(cx) + \frac{3bd^2ex^2 \operatorname{atanh}(cx)}{2} + bde^2x^3 \operatorname{atanh}(cx) + \frac{be^3x^4 \operatorname{atanh}(cx)}{4} \\ a \left(d^3x + \frac{3d^2ex^2}{2} + de^2x^3 + \frac{e^3x^4}{4} \right) \end{cases}$$

input `integrate((e*x+d)**3*(a+b*atanh(c*x)),x)`

output `Piecewise((a*d**3*x + 3*a*d**2*e*x**2/2 + a*d*e**2*x**3 + a*e**3*x**4/4 + b*d**3*x*atanh(c*x) + 3*b*d**2*e*x**2*atanh(c*x)/2 + b*d*e**2*x**3*atanh(c*x) + b*e**3*x**4*atanh(c*x)/4 + b*d**3*log(x - 1/c)/c + b*d**3*atanh(c*x)/c + 3*b*d**2*e*x/(2*c) + b*d*e**2*x**2/(2*c) + b*e**3*x**3/(12*c) - 3*b*d**2*e*atanh(c*x)/(2*c**2) + b*d*e**2*log(x - 1/c)/c**3 + b*d*e**2*atanh(c*x)/c**3 + b*e**3*x/(4*c**3) - b*e**3*atanh(c*x)/(4*c**4), Ne(c, 0)), (a*(d**3*x + 3*d**2*e*x**2/2 + d*e**2*x**3 + e**3*x**4/4), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.67

$$\begin{aligned} & \int (d + ex)^3 (a + b \operatorname{arctanh}(cx)) dx \\ &= \frac{1}{4} ae^3 x^4 + ade^2 x^3 + \frac{3}{2} ad^2 ex^2 \\ &+ \frac{3}{4} \left(2x^2 \operatorname{arctanh}(cx) + c \left(\frac{2x}{c^2} - \frac{\log(cx+1)}{c^3} + \frac{\log(cx-1)}{c^3} \right) \right) bd^2 e \\ &+ \frac{1}{2} \left(2x^3 \operatorname{arctanh}(cx) + c \left(\frac{x^2}{c^2} + \frac{\log(c^2 x^2 - 1)}{c^4} \right) \right) bde^2 \\ &+ \frac{1}{24} \left(6x^4 \operatorname{arctanh}(cx) + c \left(\frac{2(c^2 x^3 + 3x)}{c^4} - \frac{3 \log(cx+1)}{c^5} + \frac{3 \log(cx-1)}{c^5} \right) \right) be^3 \\ &+ ad^3 x + \frac{(2cx \operatorname{arctanh}(cx) + \log(-c^2 x^2 + 1))bd^3}{2c} \end{aligned}$$

input `integrate((e*x+d)^3*(a+b*arctanh(c*x)),x, algorithm="maxima")`

output `1/4*a*e^3*x^4 + a*d*e^2*x^3 + 3/2*a*d^2*e*x^2 + 3/4*(2*x^2*arctanh(c*x) + c*(2*x/c^2 - log(c*x + 1)/c^3 + log(c*x - 1)/c^3))*b*d^2*e + 1/2*(2*x^3*arctanh(c*x) + c*(x^2/c^2 + log(c^2*x^2 - 1)/c^4))*b*d*e^2 + 1/24*(6*x^4*arctanh(c*x) + c*(2*(c^2*x^3 + 3*x)/c^4 - 3*log(c*x + 1)/c^5 + 3*log(c*x - 1)/c^5))*b*e^3 + a*d^3*x + 1/2*(2*c*x*arctanh(c*x) + log(-c^2*x^2 + 1))*b*d^3/c`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 970 vs. $2(113) = 226$.

Time = 0.14 (sec) , antiderivative size = 970, normalized size of antiderivative = 7.76

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx)) dx = \text{Too large to display}$$

input `integrate((e*x+d)^3*(a+b*arctanh(c*x)),x, algorithm="giac")`

output

```
1/3*c*(3*((c*x + 1)^3*b*c^3*d^3/(c*x - 1)^3 - 3*(c*x + 1)^2*b*c^3*d^3/(c*x
- 1)^2 + 3*(c*x + 1)*b*c^3*d^3/(c*x - 1) - b*c^3*d^3 + 3*(c*x + 1)^3*b*c^
2*d^2*e/(c*x - 1)^3 - 6*(c*x + 1)^2*b*c^2*d^2*e/(c*x - 1)^2 + 3*(c*x + 1)*
b*c^2*d^2*e/(c*x - 1) + 3*(c*x + 1)^3*b*c*d*e^2/(c*x - 1)^3 - 3*(c*x + 1)^
2*b*c*d*e^2/(c*x - 1)^2 + (c*x + 1)*b*c*d*e^2/(c*x - 1) - b*c*d*e^2 + (c*x
+ 1)^3*b*e^3/(c*x - 1)^3 + (c*x + 1)*b*e^3/(c*x - 1))*log(-(c*x + 1)/(c*x
- 1))/((c*x + 1)^4*c^5/(c*x - 1)^4 - 4*(c*x + 1)^3*c^5/(c*x - 1)^3 + 6*(c
*x + 1)^2*c^5/(c*x - 1)^2 - 4*(c*x + 1)*c^5/(c*x - 1) + c^5) + (6*(c*x + 1
)^3*a*c^3*d^3/(c*x - 1)^3 - 18*(c*x + 1)^2*a*c^3*d^3/(c*x - 1)^2 + 18*(c*x
+ 1)*a*c^3*d^3/(c*x - 1) - 6*a*c^3*d^3 + 18*(c*x + 1)^3*a*c^2*d^2*e/(c*x
- 1)^3 - 36*(c*x + 1)^2*a*c^2*d^2*e/(c*x - 1)^2 + 18*(c*x + 1)*a*c^2*d^2*e
/(c*x - 1) + 9*(c*x + 1)^3*b*c^2*d^2*e/(c*x - 1)^3 - 27*(c*x + 1)^2*b*c^2*d
^2*e/(c*x - 1)^2 + 27*(c*x + 1)*b*c^2*d^2*e/(c*x - 1) - 9*b*c^2*d^2*e + 1
8*(c*x + 1)^3*a*c*d*e^2/(c*x - 1)^3 - 18*(c*x + 1)^2*a*c*d*e^2/(c*x - 1)^2
+ 6*(c*x + 1)*a*c*d*e^2/(c*x - 1) - 6*a*c*d*e^2 + 6*(c*x + 1)^3*b*c*d*e^2
/(c*x - 1)^3 - 12*(c*x + 1)^2*b*c*d*e^2/(c*x - 1)^2 + 6*(c*x + 1)*b*c*d*e^
2/(c*x - 1) + 6*(c*x + 1)^3*a*e^3/(c*x - 1)^3 + 6*(c*x + 1)*a*e^3/(c*x - 1
) + 3*(c*x + 1)^3*b*e^3/(c*x - 1)^3 - 6*(c*x + 1)^2*b*e^3/(c*x - 1)^2 + 5*
(c*x + 1)*b*e^3/(c*x - 1) - 2*b*e^3)/((c*x + 1)^4*c^5/(c*x - 1)^4 - 4*(c*x
+ 1)^3*c^5/(c*x - 1)^3 + 6*(c*x + 1)^2*c^5/(c*x - 1)^2 - 4*(c*x + 1)*c...
```


output

```
(12*atanh(c*x)*b*c**4*d**3*x + 18*atanh(c*x)*b*c**4*d**2*e*x**2 + 12*atanh(c*x)*b*c**4*d*e**2*x**3 + 3*atanh(c*x)*b*c**4*e**3*x**4 + 12*atanh(c*x)*b*c**3*d**3 - 18*atanh(c*x)*b*c**2*d**2*e + 12*atanh(c*x)*b*c*d*e**2 - 3*atanh(c*x)*b*e**3 + 12*log(c**2*x - c)*b*c**3*d**3 + 12*log(c**2*x - c)*b*c*d*e**2 + 12*a*c**4*d**3*x + 18*a*c**4*d**2*e*x**2 + 12*a*c**4*d*e**2*x**3 + 3*a*c**4*e**3*x**4 + 18*b*c**3*d**2*e*x + 6*b*c**3*d*e**2*x**2 + b*c**3*e**3*x**3 + 3*b*c*e**3*x)/(12*c**4)
```

3.3 $\int (d + ex)^2 (a + b \operatorname{arctanh}(cx)) dx$

Optimal result	62
Mathematica [A] (verified)	62
Rubi [A] (verified)	63
Maple [A] (verified)	64
Fricas [A] (verification not implemented)	65
Sympy [B] (verification not implemented)	65
Maxima [A] (verification not implemented)	66
Giac [B] (verification not implemented)	66
Mupad [B] (verification not implemented)	67
Reduce [B] (verification not implemented)	68

Optimal result

Integrand size = 16, antiderivative size = 96

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx)) dx = \frac{bdex}{c} + \frac{be^2x^2}{6c} + \frac{(d + ex)^3 (a + b \operatorname{arctanh}(cx))}{3e} + \frac{b(cd + e)^3 \log(1 - cx)}{6c^3e} - \frac{b(cd - e)^3 \log(1 + cx)}{6c^3e}$$

output

```
b*d*e*x/c+1/6*b*e^2*x^2/c+1/3*(e*x+d)^3*(a+b*arctanh(c*x))/e+1/6*b*(c*d+e)^3*ln(-c*x+1)/c^3/e-1/6*b*(c*d-e)^3*ln(c*x+1)/c^3/e
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.34

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx)) dx = \frac{1}{6} \left(\frac{6d(acd + be)x}{c} + \frac{e(6acd + be)x^2}{c} + 2ae^2x^3 + 2bx(3d^2 + 3dex + e^2x^2) \operatorname{arctanh}(cx) + \frac{b(3c^2d^2 + 3cde + e^2) \log(1 - cx)}{c^3} + \frac{b(3c^2d^2 - 3cde + e^2) \log(1 + cx)}{c^3} \right)$$

input `Integrate[(d + e*x)^2*(a + b*ArcTanh[c*x]), x]`

output `((6*d*(a*c*d + b*e)*x)/c + (e*(6*a*c*d + b*e)*x^2)/c + 2*a*e^2*x^3 + 2*b*x*(3*d^2 + 3*d*e*x + e^2*x^2)*ArcTanh[c*x] + (b*(3*c^2*d^2 + 3*c*d*e + e^2)*Log[1 - c*x])/c^3 + (b*(3*c^2*d^2 - 3*c*d*e + e^2)*Log[1 + c*x])/c^3)/6`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.03, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6478, 477, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex)^2 (a + \operatorname{barctanh}(cx)) dx$$

$$\downarrow 6478$$

$$\frac{(d + ex)^3 (a + \operatorname{barctanh}(cx))}{3e} - \frac{bc \int \frac{(d+ex)^3}{1-c^2x^2} dx}{3e}$$

$$\downarrow 477$$

$$\frac{(d + ex)^3 (a + \operatorname{barctanh}(cx))}{3e} - \frac{bc \int \left(\frac{(cd-e)^3}{2c^3(cx+1)} - \frac{3de^2}{c^2} - \frac{e^3x}{c^2} + \frac{(cd+e)^3}{2c^3(1-cx)} \right) dx}{3e}$$

$$\downarrow 2009$$

$$\frac{(d + ex)^3 (a + \operatorname{barctanh}(cx))}{3e} - \frac{bc \left(\frac{(cd-e)^3 \log(cx+1)}{2c^4} - \frac{(cd+e)^3 \log(1-cx)}{2c^4} - \frac{3de^2x}{c^2} - \frac{e^3x^2}{2c^2} \right)}{3e}$$

input `Int[(d + e*x)^2*(a + b*ArcTanh[c*x]), x]`

output `((d + e*x)^3*(a + b*ArcTanh[c*x]))/(3*e) - (b*c*((-3*d*e^2*x)/c^2 - (e^3*x^2)/(2*c^2) - ((c*d + e)^3*Log[1 - c*x])/(2*c^4) + ((c*d - e)^3*Log[1 + c*x])/(2*c^4)))/(3*e)`

Defintions of rubi rules used

```
rule 477 Int[((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[
a^p Int[ExpandIntegrand[(c + d*x)^n*(1 - Rt[-b/a, 2]*x)^p*(1 + Rt[-b/a, 2
]*x)^p, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IntegerQ[n] &
& NiceSqrtQ[-b/a] && !FractionalPowerFactorQ[Rt[-b/a, 2]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6478 Int[((a_) + ArcTanh[(c_)*(x_)]*(b_))*((d_) + (e_)*(x_))^(q_), x_Symbol
] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])/(e*(q + 1))), x] - Simp[b
*(c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 - c^2*x^2), x], x] /; FreeQ[{a,
b, c, d, e, q}, x] && NeQ[q, -1]
```

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.72

method	result
parallelrisch	$\frac{2x^3 \operatorname{arctanh}(cx) b c^3 e^2 + 2x^3 a c^3 e^2 + 6x^2 \operatorname{arctanh}(cx) b c^3 d e + 6x^2 a c^3 d e + 6b d^2 \operatorname{arctanh}(cx) x c^3 + x^2 b c^2 e^2 + 6x a c^3 d^2 + 6b d^3 \operatorname{arctanh}(cx)}{6c^3}$
parts	$\frac{a(e x+d)^3}{3e} + \frac{b \left(\frac{c e^2 \operatorname{arctanh}(c x) x^3}{3} + c e \operatorname{arctanh}(c x) x^2 d + \operatorname{arctanh}(c x) c x d^2 + \frac{c \operatorname{arctanh}(c x) d^3}{3e} - 3c^2 d e^2 x - \frac{e^3 c^2 x^2}{2} - \frac{(c^3 d^3 + a c^3 d^2 + 6c^2 d^2 e x + 6c d^2 e^2 x^2 + 6a c^2 d^2 e x^2 + 6a c^2 d^2 e x^2 + 6a c^2 d^2 e x^2)}{6c^3} \right)}{c}$
derivativedivides	$\frac{a(c e x+c d)^3}{3c^2 e} + \frac{b \left(\frac{\operatorname{arctanh}(c x) c^3 d^3}{3e} + \operatorname{arctanh}(c x) c^3 d^2 x + e \operatorname{arctanh}(c x) c^3 d x^2 + \frac{e^2 \operatorname{arctanh}(c x) c^3 x^3}{3} - 3c^2 d e^2 x - \frac{e^3 c^2 x^2}{2} - \frac{(c^3 d^3 + a c^3 d^2 + 6c^2 d^2 e x + 6c d^2 e^2 x^2 + 6a c^2 d^2 e x^2 + 6a c^2 d^2 e x^2 + 6a c^2 d^2 e x^2)}{6c^3} \right)}{c^2}$
default	$\frac{a(c e x+c d)^3}{3c^2 e} + \frac{b \left(\frac{\operatorname{arctanh}(c x) c^3 d^3}{3e} + \operatorname{arctanh}(c x) c^3 d^2 x + e \operatorname{arctanh}(c x) c^3 d x^2 + \frac{e^2 \operatorname{arctanh}(c x) c^3 x^3}{3} - 3c^2 d e^2 x - \frac{e^3 c^2 x^2}{2} - \frac{(c^3 d^3 + a c^3 d^2 + 6c^2 d^2 e x + 6c d^2 e^2 x^2 + 6a c^2 d^2 e x^2 + 6a c^2 d^2 e x^2 + 6a c^2 d^2 e x^2)}{6c^3} \right)}{c^2}$
risch	$\frac{(e x+d)^3 b \ln(c x+1)}{6e} - \frac{e^2 b x^3 \ln(-c x+1)}{6} - \frac{e b d x^2 \ln(-c x+1)}{2} + \frac{e^2 a x^3}{3} - \frac{b d^2 x \ln(-c x+1)}{2} + e a d x^2 - \frac{\ln(c x+d)}{c}$

```
input int((e*x+d)^2*(a+b*arctanh(c*x)),x,method=_RETURNVERBOSE)
```

output

```
1/6*(2*x^3*arctanh(c*x)*b*c^3*e^2+2*x^3*a*c^3*e^2+6*x^2*arctanh(c*x)*b*c^3*d*e+6*x^2*a*c^3*d*e+6*b*d^2*arctanh(c*x)*x*c^3+x^2*b*c^2*e^2+6*x*a*c^3*d^2+6*ln(c*x-1)*b*c^2*d^2+6*x*b*c^2*d*e+6*arctanh(c*x)*b*c^2*d^2-6*arctanh(c*x)*b*c*d*e+2*ln(c*x-1)*b*e^2+2*arctanh(c*x)*b*e^2)/c^3
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.70

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{2ac^3e^2x^3 + (6ac^3de + bc^2e^2)x^2 + 6(ac^3d^2 + bc^2de)x + (3bc^2d^2 - 3bcde + be^2) \log(cx + 1) + (3bc^2d^2 + 3bcde - be^2) \log(cx - 1)}{6c^3}$$

input

```
integrate((e*x+d)^2*(a+b*arctanh(c*x)),x, algorithm="fricas")
```

output

```
1/6*(2*a*c^3*e^2*x^3 + (6*a*c^3*d*e + b*c^2*e^2)*x^2 + 6*(a*c^3*d^2 + b*c^2*d*e)*x + (3*b*c^2*d^2 - 3*b*c*d*e + b*e^2)*log(c*x + 1) + (3*b*c^2*d^2 + 3*b*c*d*e + b*e^2)*log(c*x - 1) + (b*c^3*e^2*x^3 + 3*b*c^3*d*e*x^2 + 3*b*c^3*d^2*x)*log(-(c*x + 1)/(c*x - 1)))/c^3
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 178 vs. 2(82) = 164.

Time = 0.35 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.85

$$\int (d + ex)^2 (a + b \operatorname{atanh}(cx)) dx$$

$$= \begin{cases} ad^2x + adex^2 + \frac{ae^2x^3}{3} + bd^2x \operatorname{atanh}(cx) + bdex^2 \operatorname{atanh}(cx) + \frac{be^2x^3 \operatorname{atanh}(cx)}{3} + \frac{bd^2 \log(x - \frac{1}{c})}{c} + \frac{bd^2 \operatorname{atanh}(cx)}{c} \\ a \left(d^2x + dex^2 + \frac{e^2x^3}{3} \right) \end{cases}$$

input

```
integrate((e*x+d)**2*(a+b*atanh(c*x)),x)
```

output

```
Piecewise((a*d**2*x + a*d*e*x**2 + a*e**2*x**3/3 + b*d**2*x*atanh(c*x) + b
*d*e*x**2*atanh(c*x) + b*e**2*x**3*atanh(c*x)/3 + b*d**2*log(x - 1/c)/c +
b*d**2*atanh(c*x)/c + b*d*e*x/c + b*e**2*x**2/(6*c) - b*d*e*atanh(c*x)/c**
2 + b*e**2*log(x - 1/c)/(3*c**3) + b*e**2*atanh(c*x)/(3*c**3), Ne(c, 0)),
(a*(d**2*x + d*e*x**2 + e**2*x**3/3), True))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.43

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{1}{3} a e^2 x^3 + a d e x^2 + \frac{1}{2} \left(2 x^2 \operatorname{arctanh}(cx) + c \left(\frac{2x}{c^2} - \frac{\log(cx+1)}{c^3} + \frac{\log(cx-1)}{c^3} \right) \right) b d e$$

$$+ \frac{1}{6} \left(2 x^3 \operatorname{arctanh}(cx) + c \left(\frac{x^2}{c^2} + \frac{\log(c^2 x^2 - 1)}{c^4} \right) \right) b e^2$$

$$+ a d^2 x + \frac{(2 c x \operatorname{arctanh}(cx) + \log(-c^2 x^2 + 1)) b d^2}{2 c}$$

input

```
integrate((e*x+d)^2*(a+b*arctanh(c*x)),x, algorithm="maxima")
```

output

```
1/3*a*e^2*x^3 + a*d*e*x^2 + 1/2*(2*x^2*arctanh(c*x) + c*(2*x/c^2 - log(c*x
+ 1)/c^3 + log(c*x - 1)/c^3))*b*d*e + 1/6*(2*x^3*arctanh(c*x) + c*(x^2/c^
2 + log(c^2*x^2 - 1)/c^4))*b*e^2 + a*d^2*x + 1/2*(2*c*x*arctanh(c*x) + log
(-c^2*x^2 + 1))*b*d^2/c
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 532 vs. 2(88) = 176.

Time = 0.13 (sec) , antiderivative size = 532, normalized size of antiderivative = 5.54

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{1}{3} c \left(\frac{\left(\frac{3(cx+1)^2 bc^2 d^2}{(cx-1)^2} - \frac{6(cx+1) bc^2 d^2}{cx-1} + 3 bc^2 d^2 + \frac{6(cx+1)^2 bcde}{(cx-1)^2} - \frac{6(cx+1) bcde}{cx-1} + \frac{3(cx+1)^2 be^2}{(cx-1)^2} + be^2 \right) \log\left(-\frac{cx+1}{cx-1}\right) + \frac{(cx+1)^3 c^4}{(cx-1)^3} - \frac{3(cx+1)^2 c^4}{(cx-1)^2} + \frac{3(cx+1) c^4}{cx-1} - c^4}{1} \right)$$

input `integrate((e*x+d)^2*(a+b*arctanh(c*x)),x, algorithm="giac")`

output

$$\frac{1}{3}c \left(\frac{3(c^2x^2 + 1)^2 b c^2 d^2}{(cx - 1)^2} - \frac{6(c^2x^2 + 1) b c^2 d^2}{(cx - 1)} + 3b c^2 d^2 + \frac{6(c^2x^2 + 1)^2 b c d e}{(cx - 1)^2} - \frac{6(c^2x^2 + 1) b c d e}{(cx - 1)} + \frac{3(c^2x^2 + 1)^2 b e^2}{(cx - 1)^2} + \frac{b e^2}{(cx - 1)} \right) \log\left(\frac{-(cx + 1)}{(cx - 1)}\right) + \frac{3(c^2x^2 + 1)^3 c^4}{(cx - 1)^3} - \frac{3(c^2x^2 + 1)^2 c^4}{(cx - 1)^2} + \frac{3(c^2x^2 + 1) c^4}{(cx - 1)} - c^4 + \frac{2(3(c^2x^2 + 1)^2 a c^2 d^2}{(cx - 1)^2} - \frac{6(c^2x^2 + 1) a c^2 d^2}{(cx - 1)} + \frac{3 a c^2 d^2}{(cx - 1)} + \frac{6(c^2x^2 + 1)^2 a c d e}{(cx - 1)^2} - \frac{6(c^2x^2 + 1) a c d e}{(cx - 1)} + \frac{3(c^2x^2 + 1)^2 b c d e}{(cx - 1)^2} - \frac{6(c^2x^2 + 1) b c d e}{(cx - 1)} + \frac{3 b c d e}{(cx - 1)} + \frac{3(c^2x^2 + 1)^2 a e^2}{(cx - 1)^2} + \frac{a e^2}{(cx - 1)} + \frac{(c^2x^2 + 1)^2 b e^2}{(cx - 1)^2} - \frac{(c^2x^2 + 1) b e^2}{(cx - 1)} \Big) / \left(\frac{(c^2x^2 + 1)^3 c^4}{(cx - 1)^3} - \frac{3(c^2x^2 + 1)^2 c^4}{(cx - 1)^2} + \frac{3(c^2x^2 + 1) c^4}{(cx - 1)} - c^4 \right) - \frac{(3b c^2 d^2 + b e^2) \log\left(\frac{-(cx + 1)}{(cx - 1)}\right) + 1}{c^4} + \frac{(3b c^2 d^2 + b e^2) \log\left(\frac{-(cx + 1)}{(cx - 1)}\right)}{c^4}$$

Mupad [B] (verification not implemented)

Time = 3.58 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.32

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx)) dx = \frac{a e^2 x^3}{3} + a d^2 x + \frac{b d^2 \ln(c^2 x^2 - 1)}{2c} + \frac{b e^2 \ln(c^2 x^2 - 1)}{6c^3} + \frac{b e^2 x^2}{6c} + a d e x^2 + b d^2 x \operatorname{atanh}(cx) + \frac{b e^2 x^3 \operatorname{atanh}(cx)}{3} + \frac{b d e x}{c} - \frac{b d e \operatorname{atanh}(cx)}{c^2} + b d e x^2 \operatorname{atanh}(cx)$$

input `int((a + b*atanh(c*x))*(d + e*x)^2,x)`

output

$$\frac{(a e^2 x^3)}{3} + a d^2 x + \frac{(b d^2 \log(c^2 x^2 - 1))}{(2c)} + \frac{(b e^2 \log(c^2 x^2 - 1))}{(6c^3)} + \frac{(b e^2 x^2)}{(6c)} + a d e x^2 + b d^2 x \operatorname{atanh}(c x) + \frac{(b e^2 x^3 \operatorname{atanh}(c x))}{3} + \frac{(b d e x)}{c} - \frac{(b d e \operatorname{atanh}(c x))}{c^2} + b d e x^2 \operatorname{atanh}(c x)$$

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.79

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{6 \operatorname{atanh}(cx) b c^3 d^2 x + 6 \operatorname{atanh}(cx) b c^3 d e x^2 + 2 \operatorname{atanh}(cx) b c^3 e^2 x^3 + 6 \operatorname{atanh}(cx) b c^2 d^2 - 6 \operatorname{atanh}(cx) b c d e}{6 c^3}$$

input `int((e*x+d)^2*(a+b*atanh(c*x)),x)`output `(6*atanh(c*x)*b*c**3*d**2*x + 6*atanh(c*x)*b*c**3*d*e*x**2 + 2*atanh(c*x)*b*c**3*e**2*x**3 + 6*atanh(c*x)*b*c**2*d**2 - 6*atanh(c*x)*b*c*d*e + 2*atanh(c*x)*b*e**2 + 6*log(c**2*x - c)*b*c**2*d**2 + 2*log(c**2*x - c)*b*e**2 + 6*a*c**3*d**2*x + 6*a*c**3*d*e*x**2 + 2*a*c**3*e**2*x**3 + 6*b*c**2*d*e*x + b*c**2*e**2*x**2)/(6*c**3)`

3.4 $\int (d + ex)(a + b \operatorname{arctanh}(cx)) dx$

Optimal result	69
Mathematica [A] (verified)	69
Rubi [A] (verified)	70
Maple [A] (verified)	71
Fricas [A] (verification not implemented)	72
Sympy [A] (verification not implemented)	72
Maxima [A] (verification not implemented)	73
Giac [B] (verification not implemented)	73
Mupad [B] (verification not implemented)	74
Reduce [B] (verification not implemented)	74

Optimal result

Integrand size = 14, antiderivative size = 84

$$\int (d + ex)(a + b \operatorname{arctanh}(cx)) dx = \frac{bex}{2c} + \frac{(d + ex)^2(a + b \operatorname{arctanh}(cx))}{2e} + \frac{b(cd + e)^2 \log(1 - cx)}{4c^2e} - \frac{b(cd - e)^2 \log(1 + cx)}{4c^2e}$$

output

```
1/2*b*e*x/c+1/2*(e*x+d)^2*(a+b*arctanh(c*x))/e+1/4*b*(c*d+e)^2*ln(-c*x+1)/c^2/e-1/4*b*(c*d-e)^2*ln(c*x+1)/c^2/e
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.14

$$\int (d + ex)(a + b \operatorname{arctanh}(cx)) dx = adx + \frac{bex}{2c} + \frac{1}{2}aex^2 + bdx \operatorname{arctanh}(cx) + \frac{1}{2}bex^2 \operatorname{arctanh}(cx) + \frac{be \log(1 - cx)}{4c^2} - \frac{be \log(1 + cx)}{4c^2} + \frac{bd \log(1 - c^2x^2)}{2c}$$

input

```
Integrate[(d + e*x)*(a + b*ArcTanh[c*x]),x]
```

output

```
a*d*x + (b*e*x)/(2*c) + (a*e*x^2)/2 + b*d*x*ArcTanh[c*x] + (b*e*x^2*ArcTan
h[c*x])/2 + (b*e*Log[1 - c*x])/(4*c^2) - (b*e*Log[1 + c*x])/(4*c^2) + (b*d
*Log[1 - c^2*x^2])/(2*c)
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.01, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {6478, 477, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex)(a + b \operatorname{arctanh}(cx)) dx$$

$$\downarrow 6478$$

$$\frac{(d + ex)^2(a + b \operatorname{arctanh}(cx))}{2e} - \frac{bc \int \frac{(d+ex)^2}{1-c^2x^2} dx}{2e}$$

$$\downarrow 477$$

$$\frac{(d + ex)^2(a + b \operatorname{arctanh}(cx))}{2e} - \frac{bc \int \left(\frac{(cd-e)^2}{2c^2(cx+1)} - \frac{e^2}{c^2} + \frac{(cd+e)^2}{2c^2(1-cx)} \right) dx}{2e}$$

$$\downarrow 2009$$

$$\frac{(d + ex)^2(a + b \operatorname{arctanh}(cx))}{2e} - \frac{bc \left(\frac{(cd-e)^2 \log(cx+1)}{2c^3} - \frac{(cd+e)^2 \log(1-cx)}{2c^3} - \frac{e^2 x}{c^2} \right)}{2e}$$

input

```
Int[(d + e*x)*(a + b*ArcTanh[c*x]),x]
```

output

```
((d + e*x)^2*(a + b*ArcTanh[c*x]))/(2*e) - (b*c*(-((e^2*x)/c^2) - ((c*d +
e)^2*Log[1 - c*x])/(2*c^3) + ((c*d - e)^2*Log[1 + c*x])/(2*c^3)))/(2*e)
```

Defintions of rubi rules used

```
rule 477 Int[((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[
a^p Int[ExpandIntegrand[(c + d*x)^n*(1 - Rt[-b/a, 2]*x)^p*(1 + Rt[-b/a, 2]
]*x)^p, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IntegerQ[n] &
& NiceSqrtQ[-b/a] && !FractionalPowerFactorQ[Rt[-b/a, 2]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6478 Int[((a_) + ArcTanh[(c_)*(x_)]*(b_))*((d_) + (e_)*(x_))^(q_), x_Symbol
] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])/(e*(q + 1))), x] - Simp[b
*(c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 - c^2*x^2), x], x] /; FreeQ[{a,
b, c, d, e, q}, x] && NeQ[q, -1]
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.93

method	result
parts	$a\left(\frac{1}{2}e x^2 + dx\right) + \frac{b\left(\frac{c \operatorname{arctanh}(cx)x^2 e}{2} + \operatorname{arctanh}(cx)dcx - \frac{-cex - \frac{(2cd+e)\ln(cx-1)}{2} + \frac{(-2cd+e)\ln(cx+1)}{2}}{2c}\right)}{c}$
parallelrisch	$\frac{\operatorname{arctanh}(cx)bc^2e x^2 + ac^2e x^2 + 2bd \operatorname{arctanh}(cx)x c^2 + 2ac^2 dx + 2\ln(cx-1)bcd + bce x + 2 \operatorname{arctanh}(cx)bcd - \operatorname{arctanh}(cx)bc^2}{2c^2}$
derivativdivides	$\frac{a\left(c^2 dx + \frac{1}{2}c^2 e x^2\right)}{c} + \frac{b\left(\operatorname{arctanh}(cx)dc^2 x + \frac{\operatorname{arctanh}(cx)e c^2 x^2}{2} + \frac{cex}{2} + \frac{(2cd+e)\ln(cx-1)}{4} - \frac{(-2cd+e)\ln(cx+1)}{4}\right)}{c}$
default	$\frac{a\left(c^2 dx + \frac{1}{2}c^2 e x^2\right)}{c} + \frac{b\left(\operatorname{arctanh}(cx)dc^2 x + \frac{\operatorname{arctanh}(cx)e c^2 x^2}{2} + \frac{cex}{2} + \frac{(2cd+e)\ln(cx-1)}{4} - \frac{(-2cd+e)\ln(cx+1)}{4}\right)}{c}$
risch	$\frac{bx(ex+2d)\ln(cx+1)}{4} - \frac{be x^2 \ln(-cx+1)}{4} - \frac{bdx \ln(-cx+1)}{2} + \frac{ae x^2}{2} + adx + \frac{\ln(cx+1)bd}{2c} + \frac{\ln(-cx+1)bd}{2c} +$

```
input int((e*x+d)*(a+b*arctanh(c*x)),x,method=_RETURNVERBOSE)
```

```
output a*(1/2*e*x^2+d*x)+b/c*(1/2*c*arctanh(c*x)*x^2*e+arctanh(c*x)*d*c*x-1/2/c*(-
-c*e*x-1/2*(2*c*d+e)*ln(c*x-1)+1/2*(-2*c*d+e)*ln(c*x+1))
```


Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.17

$$\int (d + ex)(a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{2ac^2ex^2 + 2(2ac^2d + bce)x + (2bcd - be) \log(cx + 1) + (2bcd + be) \log(cx - 1) + (bc^2ex^2 + 2bc^2dx) \log(-\frac{cx+1}{cx-1})}{4c^2}$$

input `integrate((e*x+d)*(a+b*arctanh(c*x)),x, algorithm="fricas")`

output `1/4*(2*a*c^2*e*x^2 + 2*(2*a*c^2*d + b*c*e)*x + (2*b*c*d - b*e)*log(c*x + 1) + (2*b*c*d + b*e)*log(c*x - 1) + (b*c^2*e*x^2 + 2*b*c^2*d*x)*log(-(c*x + 1)/(c*x - 1)))/c^2`

Sympy [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.10

$$\int (d + ex)(a + b \operatorname{atanh}(cx)) dx$$

$$= \begin{cases} adx + \frac{aex^2}{2} + bdx \operatorname{atanh}(cx) + \frac{bex^2 \operatorname{atanh}(cx)}{2} + \frac{bd \log(x - \frac{1}{c})}{c} + \frac{bd \operatorname{atanh}(cx)}{c} + \frac{bex}{2c} - \frac{be \operatorname{atanh}(cx)}{2c^2} & \text{for } c \neq 0 \\ a(dx + \frac{ex^2}{2}) & \text{otherwise} \end{cases}$$

input `integrate((e*x+d)*(a+b*atanh(c*x)),x)`

output `Piecewise((a*d*x + a*e*x**2/2 + b*d*x*atanh(c*x) + b*e*x**2*atanh(c*x)/2 + b*d*log(x - 1/c)/c + b*d*atanh(c*x)/c + b*e*x/(2*c) - b*e*atanh(c*x)/(2*c**2), Ne(c, 0)), (a*(d*x + e*x**2/2), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.99

$$\int (d + ex)(a + b \operatorname{arctanh}(cx)) dx$$

$$= \frac{1}{2} aex^2 + \frac{1}{4} \left(2x^2 \operatorname{arctanh}(cx) + c \left(\frac{2x}{c^2} - \frac{\log(cx+1)}{c^3} + \frac{\log(cx-1)}{c^3} \right) \right) be$$

$$+ adx + \frac{(2cx \operatorname{arctanh}(cx) + \log(-c^2x^2 + 1))bd}{2c}$$

input `integrate((e*x+d)*(a+b*arctanh(c*x)),x, algorithm="maxima")`

output `1/2*a*e*x^2 + 1/4*(2*x^2*arctanh(c*x) + c*(2*x/c^2 - log(c*x + 1)/c^3 + log(c*x - 1)/c^3))*b*e + a*d*x + 1/2*(2*c*x*arctanh(c*x) + log(-c^2*x^2 + 1))*b*d/c`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 245 vs. 2(76) = 152.

Time = 0.12 (sec) , antiderivative size = 245, normalized size of antiderivative = 2.92

$$\int (d + ex)(a + b \operatorname{arctanh}(cx)) dx =$$

$$-c \left(\frac{bd \log\left(-\frac{cx+1}{cx-1} + 1\right)}{c^2} - \frac{\left(\frac{(cx+1)bcd}{cx-1} - bcd + \frac{(cx+1)be}{cx-1}\right) \log\left(-\frac{cx+1}{cx-1}\right)}{\frac{(cx+1)^2c^3}{(cx-1)^2} - \frac{2(cx+1)c^3}{cx-1} + c^3} - \frac{bd \log\left(-\frac{cx+1}{cx-1}\right)}{c^2} - \frac{2(cx+1)acd}{cx-1} - 2a \frac{(cx+1)}{(cx-1)} \right)$$

input `integrate((e*x+d)*(a+b*arctanh(c*x)),x, algorithm="giac")`

output `-c*(b*d*log(-(c*x + 1)/(c*x - 1) + 1)/c^2 - ((c*x + 1)*b*c*d/(c*x - 1) - b*c*d + (c*x + 1)*b*e/(c*x - 1))*log(-(c*x + 1)/(c*x - 1))/((c*x + 1)^2*c^3/(c*x - 1)^2 - 2*(c*x + 1)*c^3/(c*x - 1) + c^3) - b*d*log(-(c*x + 1)/(c*x - 1))/c^2 - (2*(c*x + 1)*a*c*d/(c*x - 1) - 2*a*c*d + 2*(c*x + 1)*a*e/(c*x - 1) + (c*x + 1)*b*e/(c*x - 1) - b*e)/((c*x + 1)^2*c^3/(c*x - 1)^2 - 2*(c*x + 1)*c^3/(c*x - 1) + c^3))`

Mupad [B] (verification not implemented)

Time = 3.52 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.80

$$\int (d + ex)(a + b \operatorname{arctanh}(cx)) dx = a dx + \frac{a e x^2}{2} + b dx \operatorname{atanh}(cx) + \frac{b e x}{2c} - \frac{b e \operatorname{atanh}(cx)}{2c^2} + \frac{b e x^2 \operatorname{atanh}(cx)}{2} + \frac{b d \ln(c^2 x^2 - 1)}{2c}$$

input `int((a + b*atanh(c*x))*(d + e*x),x)`output `a*d*x + (a*e*x^2)/2 + b*d*x*atanh(c*x) + (b*e*x)/(2*c) - (b*e*atanh(c*x))/(2*c^2) + (b*e*x^2*atanh(c*x))/2 + (b*d*log(c^2*x^2 - 1))/(2*c)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.01

$$\int (d + ex)(a + b \operatorname{arctanh}(cx)) dx = \frac{2 \operatorname{atanh}(cx) b c^2 dx + \operatorname{atanh}(cx) b c^2 e x^2 + 2 \operatorname{atanh}(cx) b c d - \operatorname{atanh}(cx) b e + 2 \log(c^2 x - c) b c d + 2 a c^2 dx + \dots}{2c^2}$$

input `int((e*x+d)*(a+b*atanh(c*x)),x)`output `(2*atanh(c*x)*b*c**2*d*x + atanh(c*x)*b*c**2*e*x**2 + 2*atanh(c*x)*b*c*d - atanh(c*x)*b*e + 2*log(c**2*x - c)*b*c*d + 2*a*c**2*d*x + a*c**2*e*x**2 + b*c*e*x)/(2*c**2)`

3.5 $\int \frac{a+b\operatorname{arctanh}(cx)}{d+ex} dx$

Optimal result	75
Mathematica [C] (warning: unable to verify)	76
Rubi [A] (verified)	76
Maple [A] (verified)	78
Fricas [F]	79
Sympy [F]	79
Maxima [F]	80
Giac [F]	80
Mupad [F(-1)]	80
Reduce [F]	81

Optimal result

Integrand size = 16, antiderivative size = 114

$$\int \frac{a + b\operatorname{arctanh}(cx)}{d + ex} dx = -\frac{(a + b\operatorname{arctanh}(cx)) \log\left(\frac{2}{1+cx}\right)}{e} + \frac{(a + b\operatorname{arctanh}(cx)) \log\left(\frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{e} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+cx}\right)}{2e} - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{2e}$$

output `-(a+b*arctanh(c*x))*ln(2/(c*x+1))/e+(a+b*arctanh(c*x))*ln(2*c*(e*x+d)/(c*d+e)/(c*x+1))/e+1/2*b*polylog(2,1-2/(c*x+1))/e-1/2*b*polylog(2,1-2*c*(e*x+d)/(c*d+e)/(c*x+1))/e`

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.17 (sec) , antiderivative size = 257, normalized size of antiderivative = 2.25

$$\int \frac{a + b \operatorname{arctanh}(cx)}{d + ex} dx$$

$$= \frac{a \log(d + ex) + b \operatorname{arctanh}(cx) \left(\frac{1}{2} \log(1 - c^2 x^2) + \log \left(i \sinh \left(\operatorname{arctanh} \left(\frac{cd}{e} \right) + \operatorname{arctanh}(cx) \right) \right) \right) - \frac{1}{2} i b \left(-\frac{1}{4} i \right)}{1}$$

input `Integrate[(a + b*ArcTanh[c*x])/(d + e*x),x]`

output `(a*Log[d + e*x] + b*ArcTanh[c*x]*(Log[1 - c^2*x^2]/2 + Log[I*Sinh[ArcTanh[(c*d)/e] + ArcTanh[c*x]]]) - (I/2)*b*((-1/4*I)*(Pi - (2*I)*ArcTanh[c*x])^2 + I*(ArcTanh[(c*d)/e] + ArcTanh[c*x])^2 + (Pi - (2*I)*ArcTanh[c*x])*Log[1 + E^(2*ArcTanh[c*x])] + (2*I)*(ArcTanh[(c*d)/e] + ArcTanh[c*x])*Log[1 - E^(-2*(ArcTanh[(c*d)/e] + ArcTanh[c*x])]) - (Pi - (2*I)*ArcTanh[c*x])*Log[2/Sqrt[1 - c^2*x^2]] - (2*I)*(ArcTanh[(c*d)/e] + ArcTanh[c*x])*Log[(2*I)*Sinh[ArcTanh[(c*d)/e] + ArcTanh[c*x]]) - I*PolyLog[2, -E^(2*ArcTanh[c*x])] - I*PolyLog[2, E^(-2*(ArcTanh[(c*d)/e] + ArcTanh[c*x])])])/e`

Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6472, 2849, 2752, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx)}{d + ex} dx$$

↓ 6472

$$\begin{aligned}
& -\frac{bc \int \frac{\log\left(\frac{2c(d+ex)}{(cd+e)(cx+1)}\right) dx}{e} + bc \int \frac{\log\left(\frac{2}{cx+1}\right) dx}{1-c^2x^2} + \frac{(a + \operatorname{barctanh}(cx)) \log\left(\frac{2c(d+ex)}{(cx+1)(cd+e)}\right)}{e}}{\log\left(\frac{2}{cx+1}\right) (a + \operatorname{barctanh}(cx))} \\
& \quad \downarrow \text{2849} \\
& -\frac{bc \int \frac{\log\left(\frac{2c(d+ex)}{(cd+e)(cx+1)}\right) dx}{e} + b \int \frac{\log\left(\frac{2}{cx+1}\right) d\frac{1}{cx+1}}{1-\frac{2}{cx+1}} + \frac{(a + \operatorname{barctanh}(cx)) \log\left(\frac{2c(d+ex)}{(cx+1)(cd+e)}\right)}{e}}{\log\left(\frac{2}{cx+1}\right) (a + \operatorname{barctanh}(cx))} \\
& \quad \downarrow \text{2752} \\
& -\frac{bc \int \frac{\log\left(\frac{2c(d+ex)}{(cd+e)(cx+1)}\right) dx}{e} + \frac{(a + \operatorname{barctanh}(cx)) \log\left(\frac{2c(d+ex)}{(cx+1)(cd+e)}\right)}{e}}{\log\left(\frac{2}{cx+1}\right) (a + \operatorname{barctanh}(cx))} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{cx+1}\right)}{2e} \\
& \quad \downarrow \text{2897} \\
& \frac{(a + \operatorname{barctanh}(cx)) \log\left(\frac{2c(d+ex)}{(cx+1)(cd+e)}\right)}{e} - \frac{\log\left(\frac{2}{cx+1}\right) (a + \operatorname{barctanh}(cx))}{e} \\
& \quad - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2c(d+ex)}{(cd+e)(cx+1)}\right)}{2e} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{cx+1}\right)}{2e}
\end{aligned}$$

input `Int[(a + b*ArcTanh[c*x])/(d + e*x), x]`

output `-(((a + b*ArcTanh[c*x])*Log[2/(1 + c*x)])/e) + ((a + b*ArcTanh[c*x])*Log[(2*c*(d + e*x))/((c*d + e)*(1 + c*x))])/e + (b*PolyLog[2, 1 - 2/(1 + c*x)])/ (2*e) - (b*PolyLog[2, 1 - (2*c*(d + e*x))/((c*d + e)*(1 + c*x))])/ (2*e)`

Defintions of rubi rules used

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 2897 `Int[Log[u_]*(Pq_)^(m_.), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x]] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6472 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-a + b*ArcTanh[c*x])*(Log[2/(1 + c*x)]/e), x] + (Simp[(a + b*ArcTanh[c*x])*(Log[2*c*((d + e*x)/((c*d + e)*(1 + c*x))])/e), x] + Simp[b*(c/e) Int[Log[2/(1 + c*x)]/(1 - c^2*x^2), x], x] - Simp[b*(c/e) Int[Log[2*c*((d + e*x)/((c*d + e)*(1 + c*x)))/(1 - c^2*x^2), x], x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 - e^2, 0]`

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.27

method	result
parts	$\frac{a \ln(cx+d)}{e} + \frac{b \ln(cx+d) \operatorname{arctanh}(cx)}{e} + \frac{b \ln(cx+d) \ln\left(\frac{cex-e}{-cd-e}\right)}{2e} + \frac{b \operatorname{dilog}\left(\frac{cex-e}{-cd-e}\right)}{2e} - \frac{b \ln(cx+d) \ln\left(\frac{cex+e}{-cd+e}\right)}{2e}$
derivativedivides	$\frac{ac \ln(cx+d)}{e} + bc \left(\frac{\ln(cx+d) \operatorname{arctanh}(cx)}{e} - \frac{e \left(\operatorname{dilog}\left(\frac{cex-e}{-cd-e}\right) + \ln(cx+d) \ln\left(\frac{cex-e}{-cd-e}\right) \right)}{2} + \frac{e \left(\operatorname{dilog}\left(\frac{cex+e}{-cd+e}\right) + \ln(cx+d) \ln\left(\frac{cex+e}{-cd+e}\right) \right)}{2} \right)$
default	$\frac{ac \ln(cx+d)}{e} + bc \left(\frac{\ln(cx+d) \operatorname{arctanh}(cx)}{e} - \frac{e \left(\operatorname{dilog}\left(\frac{cex-e}{-cd-e}\right) + \ln(cx+d) \ln\left(\frac{cex-e}{-cd-e}\right) \right)}{2} + \frac{e \left(\operatorname{dilog}\left(\frac{cex+e}{-cd+e}\right) + \ln(cx+d) \ln\left(\frac{cex+e}{-cd+e}\right) \right)}{2} \right)$
risch	$-\frac{b \operatorname{dilog}\left(\frac{(-cx+1)e-cd-e}{-cd-e}\right)}{2e} - \frac{b \ln(-cx+1) \ln\left(\frac{(-cx+1)e-cd-e}{-cd-e}\right)}{2e} + \frac{a \ln((-cx+1)e-cd-e)}{e} + \frac{b \operatorname{dilog}\left(\frac{(cx+1)e+cd-e}{cd-e}\right)}{2e}$

input `int((a+b*arctanh(c*x))/(e*x+d),x,method=_RETURNVERBOSE)`

output `a*ln(e*x+d)/e+b*ln(c*e*x+c*d)/e*arctanh(c*x)+1/2*b/e*ln(c*e*x+c*d)*ln((c*e*x-e)/(-c*d-e))+1/2*b/e*dilog((c*e*x-e)/(-c*d-e))-1/2*b/e*ln(c*e*x+c*d)*ln((c*e*x+e)/(-c*d+e))-1/2*b/e*dilog((c*e*x+e)/(-c*d+e))`

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{d + ex} dx = \int \frac{b \operatorname{artanh}(cx) + a}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x))/(e*x+d),x, algorithm="fricas")`

output `integral((b*arctanh(c*x) + a)/(e*x + d), x)`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{d + ex} dx = \int \frac{a + b \operatorname{atanh}(cx)}{d + ex} dx$$

input `integrate((a+b*atanh(c*x))/(e*x+d),x)`

output `Integral((a + b*atanh(c*x))/(d + e*x), x)`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{d + ex} dx = \int \frac{b \operatorname{artanh}(cx) + a}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x))/(e*x+d),x, algorithm="maxima")`

output `1/2*b*integrate((log(c*x + 1) - log(-c*x + 1))/(e*x + d), x) + a*log(e*x + d)/e`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{d + ex} dx = \int \frac{b \operatorname{artanh}(cx) + a}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x))/(e*x+d),x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)/(e*x + d), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx)}{d + ex} dx = \int \frac{a + b \operatorname{atanh}(cx)}{d + ex} dx$$

input `int((a + b*atanh(c*x))/(d + e*x),x)`

output `int((a + b*atanh(c*x))/(d + e*x), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{d + ex} dx = \frac{\left(\int \frac{\operatorname{atanh}(cx)}{ex+d} dx \right) be + \log(ex + d) a}{e}$$

input `int((a+b*atanh(c*x))/(e*x+d),x)`

output `(int(atanh(c*x)/(d + e*x),x)*b*e + log(d + e*x)*a)/e`

3.6 $\int \frac{a+b\operatorname{arctanh}(cx)}{(d+ex)^2} dx$

Optimal result	82
Mathematica [A] (verified)	82
Rubi [A] (verified)	83
Maple [A] (verified)	84
Fricas [B] (verification not implemented)	85
Sympy [B] (verification not implemented)	85
Maxima [A] (verification not implemented)	86
Giac [B] (verification not implemented)	87
Mupad [B] (verification not implemented)	87
Reduce [B] (verification not implemented)	88

Optimal result

Integrand size = 16, antiderivative size = 93

$$\int \frac{a + b\operatorname{arctanh}(cx)}{(d + ex)^2} dx = -\frac{a + b\operatorname{arctanh}(cx)}{e(d + ex)} - \frac{bc \log(1 - cx)}{2e(cd + e)} + \frac{bc \log(1 + cx)}{2(cd - e)e} - \frac{bc \log(d + ex)}{c^2d^2 - e^2}$$

output

```
-(a+b*arctanh(c*x))/e/(e*x+d)-1/2*b*c*ln(-c*x+1)/e/(c*d+e)+1/2*b*c*ln(c*x+1)/(c*d-e)/e-b*c*ln(e*x+d)/(c^2*d^2-e^2)
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.10

$$\int \frac{a + b\operatorname{arctanh}(cx)}{(d + ex)^2} dx = -\frac{a}{e(d + ex)} - \frac{b\operatorname{arctanh}(cx)}{e(d + ex)} - \frac{bc \log(1 - cx)}{2e(cd + e)} - \frac{bc \log(1 + cx)}{2e(-cd + e)} - \frac{bc \log(d + ex)}{c^2d^2 - e^2}$$

input

```
Integrate[(a + b*ArcTanh[c*x])/(d + e*x)^2,x]
```

output

$$-(a/(e*(d + e*x))) - (b*ArcTanh[c*x])/(e*(d + e*x)) - (b*c*Log[1 - c*x])/(2*e*(c*d + e)) - (b*c*Log[1 + c*x])/(2*e*(-(c*d) + e)) - (b*c*Log[d + e*x])/(c^2*d^2 - e^2)$$

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.96, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6478, 477, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^2} dx$$

$$\downarrow 6478$$

$$\frac{bc \int \frac{1}{(d+ex)(1-c^2x^2)} dx}{e} - \frac{a + b \operatorname{arctanh}(cx)}{e(d + ex)}$$

$$\downarrow 477$$

$$\frac{bc \int \left(-\frac{e^2}{(c^2d^2 - e^2)(d+ex)} + \frac{c}{2(cd+e)(1-cx)} + \frac{c}{2(cd-e)(cx+1)} \right) dx}{e} - \frac{a + b \operatorname{arctanh}(cx)}{e(d + ex)}$$

$$\downarrow 2009$$

$$\frac{bc \left(-\frac{e \log(d+ex)}{c^2d^2 - e^2} - \frac{\log(1-cx)}{2(cd+e)} + \frac{\log(cx+1)}{2(cd-e)} \right)}{e} - \frac{a + b \operatorname{arctanh}(cx)}{e(d + ex)}$$

input

$$\text{Int}[(a + b*ArcTanh[c*x])/(d + e*x)^2, x]$$

output

$$-((a + b*ArcTanh[c*x])/(e*(d + e*x))) + (b*c*(-1/2*Log[1 - c*x]/(c*d + e) + Log[1 + c*x]/(2*(c*d - e)) - (e*Log[d + e*x])/(c^2*d^2 - e^2)))/e$$

Defintions of rubi rules used

```
rule 477 Int[((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[
a^p Int[ExpandIntegrand[(c + d*x)^n*(1 - Rt[-b/a, 2]*x)^p*(1 + Rt[-b/a, 2
]*x)^p, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IntegerQ[n] &
& NiceSqrtQ[-b/a] && !FractionalPowerFactorQ[Rt[-b/a, 2]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6478 Int[((a_) + ArcTanh[(c_)*(x_)]*(b_))*((d_) + (e_)*(x_))^(q_), x_Symbol
] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])/(e*(q + 1))), x] - Simp[b
*(c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 - c^2*x^2), x], x] /; FreeQ[{a,
b, c, d, e, q}, x] && NeQ[q, -1]
```

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.23

method	result
parts	$-\frac{a}{(ex+d)e} + \frac{b \left(-\frac{c^2 \operatorname{arctanh}(cx)}{(cex+cd)e} + \frac{c^2 \left(-\frac{e \ln(cex+cd)}{(cd+e)(cd-e)} - \frac{\ln(cx-1)}{2cd+2e} + \frac{\ln(cx+1)}{2cd-2e} \right)}{e} \right)}{c}$
derivativedivides	$-\frac{a c^2}{(cex+cd)e} + b c^2 \left(-\frac{\operatorname{arctanh}(cx)}{(cex+cd)e} + \frac{-\frac{e \ln(cex+cd)}{(cd+e)(cd-e)} - \frac{\ln(cx-1)}{2cd+2e} + \frac{\ln(cx+1)}{2cd-2e}}{e} \right)$
default	$-\frac{a c^2}{(cex+cd)e} + b c^2 \left(-\frac{\operatorname{arctanh}(cx)}{(cex+cd)e} + \frac{-\frac{e \ln(cex+cd)}{(cd+e)(cd-e)} - \frac{\ln(cx-1)}{2cd+2e} + \frac{\ln(cx+1)}{2cd-2e}}{e} \right)$
parallelrisc	$\frac{\ln(cx-1)bc d^2 - \ln(ex+d)bc d^2 + xa c^2 d^2 + \operatorname{arctanh}(cx)bc d^2 + \operatorname{arctanh}(cx)bde - xa e^2 + \ln(cx-1)xbcd e - \ln(ex+d)xbcd e + \ln(cx+1)bc d^2 - \ln(ex+d)bc d^2 + xa c^2 d^2 + \operatorname{arctanh}(cx)bc d^2 + \operatorname{arctanh}(cx)bde - xa e^2 + \ln(cx-1)xbcd e - \ln(ex+d)xbcd e}{(c^2 d^2 - e^2)(ex+d)d}$
risc	$-\frac{b \ln(cx+1)}{2e(ex+d)} + \frac{\ln(-cx-1)bc^2 dex - \ln(cx-1)bc^2 dex + \ln(-cx-1)bc^2 d^2 + \ln(-cx-1)bc e^2 x - 2 \ln(-ex-d)bc e^2 x - \ln(-ex-d)bc e^2 x}{2e(ex+d)}$

```
input int((a+b*arctanh(c*x))/(e*x+d)^2,x,method=_RETURNVERBOSE)
```

```
output -a/(e*x+d)/e+b/c*(-c^2/(c*e*x+c*d)/e*arctanh(c*x)+c^2/e*(-e/(c*d+e)/(c*d-e
)*ln(c*e*x+c*d)-1/(2*c*d+2*e)*ln(c*x-1)+1/(2*c*d-2*e)*ln(c*x+1)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 182 vs. $2(89) = 178$.

Time = 0.10 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.96

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^2} dx = \frac{2ac^2d^2 - 2ae^2 - (bc^2d^2 + bcde + (bc^2de + bce^2)x) \log(cx + 1) + (bc^2d^2 - bcde + (bc^2de - bce^2)x) \log(cx - 1)}{2(c^2d^3e - de^3 + (c^2d^2e^2 - e^4)x)}$$

input `integrate((a+b*arctanh(c*x))/(e*x+d)^2,x, algorithm="fricas")`

output `-1/2*(2*a*c^2*d^2 - 2*a*e^2 - (b*c^2*d^2 + b*c*d*e + (b*c^2*d*e + b*c*e^2)*x)*log(c*x + 1) + (b*c^2*d^2 - b*c*d*e + (b*c^2*d*e - b*c*e^2)*x)*log(c*x - 1) + 2*(b*c*e^2*x + b*c*d*e)*log(e*x + d) + (b*c^2*d^2 - b*e^2)*log(-(c*x + 1)/(c*x - 1)))/(c^2*d^3*e - d*e^3 + (c^2*d^2*e^2 - e^4)*x)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 663 vs. $2(73) = 146$.

Time = 1.42 (sec) , antiderivative size = 663, normalized size of antiderivative = 7.13

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^2} dx = \left\{ \begin{array}{l} \frac{ax}{d^2} \\ -\frac{a}{de+e^2x} \\ \frac{ax + b \operatorname{atanh}(cx) + \frac{b \log\left(x - \frac{1}{c}\right)}{c} + \frac{b \operatorname{atanh}(cx)}{c}}{d^2} \\ -\frac{2ad}{2d^2e+2de^2x} + \frac{bd \operatorname{atanh}\left(\frac{ex}{d}\right)}{2d^2e+2de^2x} + \frac{bd}{2d^2e+2de^2x} - \frac{bex \operatorname{atanh}\left(\frac{ex}{d}\right)}{2d^2e+2de^2x} \\ -\frac{2ad}{2d^2e+2de^2x} - \frac{bd \operatorname{atanh}\left(\frac{ex}{d}\right)}{2d^2e+2de^2x} - \frac{bd}{2d^2e+2de^2x} + \frac{bex \operatorname{atanh}\left(\frac{ex}{d}\right)}{2d^2e+2de^2x} \\ -\frac{ac^2d^2}{c^2d^3e+c^2d^2e^2x-de^3-e^4x} + \frac{ae^2}{c^2d^3e+c^2d^2e^2x-de^3-e^4x} + \frac{bc^2dex \operatorname{atanh}(cx)}{c^2d^3e+c^2d^2e^2x-de^3-e^4x} + \frac{bcde \log\left(x - \frac{1}{c}\right)}{c^2d^3e+c^2d^2e^2x-de^3-e^4x} - \frac{bcde \log}{c^2d^3e+c^2d^2e^2x-de^3-e^4x} \end{array} \right.$$

input `integrate((a+b*atanh(c*x))/(e*x+d)**2,x)`

output

```
Piecewise((a*x/d**2, Eq(c, 0) & Eq(e, 0)), (-a/(d*e + e**2*x), Eq(c, 0)),
((a*x + b*x*atanh(c*x) + b*log(x - 1/c)/c + b*atanh(c*x)/c)/d**2, Eq(e, 0)
), (-2*a*d/(2*d**2*e + 2*d*e**2*x) + b*d*atanh(e*x/d)/(2*d**2*e + 2*d*e**2*x)
+ b*d/(2*d**2*e + 2*d*e**2*x) - b*e*x*atanh(e*x/d)/(2*d**2*e + 2*d*e**2*x),
Eq(c, -e/d)), (-2*a*d/(2*d**2*e + 2*d*e**2*x) - b*d*atanh(e*x/d)/(2*d**2*e + 2*d*e**2*x)
- b*d/(2*d**2*e + 2*d*e**2*x) + b*e*x*atanh(e*x/d)/(2*d**2*e + 2*d*e**2*x),
Eq(c, e/d)), (-a*c**2*d**2/(c**2*d**3*e + c**2*d**2*e**2*x - d*e**3 - e**4*x)
+ a*e**2/(c**2*d**3*e + c**2*d**2*e**2*x - d*e**3 - e**4*x) + b*c**2*d*e*x*atanh(c*x)/(c**2*d**3*e + c**2*d**2*e**2*x - d*e**3 - e**4*x)
+ b*c*d*e*log(x - 1/c)/(c**2*d**3*e + c**2*d**2*e**2*x - d*e**3 - e**4*x) - b*c*d*e*log(d/e + x)/(c**2*d**3*e + c**2*d**2*e**2*x - d*e**3 - e**4*x)
+ b*c*d*e*atanh(c*x)/(c**2*d**3*e + c**2*d**2*e**2*x - d*e**3 - e**4*x) + b*c*e**2*x*log(x - 1/c)/(c**2*d**3*e + c**2*d**2*e**2*x - d*e**3 - e**4*x)
- b*c*e**2*x*log(d/e + x)/(c**2*d**3*e + c**2*d**2*e**2*x - d*e**3 - e**4*x) + b*c*e**2*x*atanh(c*x)/(c**2*d**3*e + c**2*d**2*e**2*x - d*e**3 - e**4*x)
+ b*e**2*atanh(c*x)/(c**2*d**3*e + c**2*d**2*e**2*x - d*e**3 - e**4*x), True))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.06

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^2} dx$$

$$= \frac{1}{2} \left(c \left(\frac{\log(cx + 1)}{cde - e^2} - \frac{\log(cx - 1)}{cde + e^2} - \frac{2 \log(ex + d)}{c^2 d^2 - e^2} \right) - \frac{2 \operatorname{artanh}(cx)}{e^2 x + de} \right) b - \frac{a}{e^2 x + de}$$

input

```
integrate((a+b*arctanh(c*x))/(e*x+d)^2,x, algorithm="maxima")
```

output

```
1/2*(c*(log(c*x + 1)/(c*d*e - e^2) - log(c*x - 1)/(c*d*e + e^2) - 2*log(e*x + d)/(c^2*d^2 - e^2)) - 2*arctanh(c*x)/(e^2*x + d*e))*b - a/(e^2*x + d*e)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 243 vs. $2(89) = 178$.

Time = 0.12 (sec) , antiderivative size = 243, normalized size of antiderivative = 2.61

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^2} dx =$$

$$-c \left(\frac{b \log \left(\frac{(cx+1)cd}{cx-1} - cd + \frac{(cx+1)e}{cx-1} + e \right)}{c^2 d^2 - e^2} - \frac{b \log \left(-\frac{cx+1}{cx-1} \right)}{\frac{(cx+1)c^2 d^2}{cx-1} - c^2 d^2 + \frac{2(cx+1)cde}{cx-1} + \frac{(cx+1)e^2}{cx-1} + e^2} - \frac{b \log \left(-\frac{cx+1}{cx-1} \right)}{c^2 d^2 - e^2} \right)$$

input `integrate((a+b*arctanh(c*x))/(e*x+d)^2,x, algorithm="giac")`

output

```
-c*(b*log((c*x + 1)*c*d/(c*x - 1) - c*d + (c*x + 1)*e/(c*x - 1) + e)/(c^2*d^2 - e^2) - b*log(-(c*x + 1)/(c*x - 1))/((c*x + 1)*c^2*d^2/(c*x - 1) - c^2*d^2 + 2*(c*x + 1)*c*d*e/(c*x - 1) + (c*x + 1)*e^2/(c*x - 1) + e^2) - b*log(-(c*x + 1)/(c*x - 1))/(c^2*d^2 - e^2) - 2*a/((c*x + 1)*c^2*d^2/(c*x - 1) - c^2*d^2 + 2*(c*x + 1)*c*d*e/(c*x - 1) + (c*x + 1)*e^2/(c*x - 1) + e^2))
```

Mupad [B] (verification not implemented)

Time = 6.17 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.24

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^2} dx =$$

$$\frac{d^2 \left(\frac{bc \ln(c^2 x^2 - 1)}{2} - bc \ln(d + ex) + ac^2 x + bc^2 x \operatorname{atanh}(cx) \right) + de \left(b \operatorname{atanh}(cx) - bcx \ln(d + ex) \right)}{d(e^2 - c^2 d^2)(d + ex)}$$

input `int((a + b*atanh(c*x))/(d + e*x)^2,x)`

output

```
-(d^2*((b*c*log(c^2*x^2 - 1))/2 - b*c*log(d + e*x) + a*c^2*x + b*c^2*x*atanh(c*x)) + d*e*(b*atanh(c*x) - b*c*x*log(d + e*x) + (b*c*x*log(c^2*x^2 - 1))/2) - a*e^2*x)/(d*(e^2 - c^2*d^2)*(d + e*x))
```


Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 223, normalized size of antiderivative = 2.40

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^2} dx$$

$$= \frac{2 \operatorname{atanh}(cx) b c^2 d^2 x - 2 \operatorname{atanh}(cx) b e^2 x + \log(c^2 x - c) b c d^2 + \log(c^2 x - c) b c d e x - \log(c^2 x - c) b d e - \log(c^2 x + c) b c^2 d^2 x + 2 \operatorname{atanh}(cx) b e^2 x + \log(c^2 x + c) b c d^2 + \log(c^2 x + c) b c d e x + \log(c^2 x + c) b d e + \log(c^2 x + c) b e^2 x - 2 \log(d + e x) b c d^2 - 2 \log(d + e x) b c d e x + 2 a c^2 d^2 x - 2 a e^2 x}{(2 d (c^2 d^3 + c^2 d^2 e x - d e^2 - e^3 x))}$$

input `int((a+b*atanh(c*x))/(e*x+d)^2,x)`output `(2*atanh(c*x)*b*c**2*d**2*x - 2*atanh(c*x)*b*e**2*x + log(c**2*x - c)*b*c*d**2 + log(c**2*x - c)*b*c*d*e*x - log(c**2*x - c)*b*d*e - log(c**2*x - c)*b*e**2*x + log(c**2*x + c)*b*c*d**2 + log(c**2*x + c)*b*c*d*e*x + log(c**2*x + c)*b*d*e + log(c**2*x + c)*b*e**2*x - 2*log(d + e*x)*b*c*d**2 - 2*log(d + e*x)*b*c*d*e*x + 2*a*c**2*d**2*x - 2*a*e**2*x)/(2*d*(c**2*d**3 + c**2*d**2*e*x - d*e**2 - e**3*x))`

3.7 $\int \frac{a+b\operatorname{arctanh}(cx)}{(d+ex)^3} dx$

Optimal result	89
Mathematica [A] (verified)	89
Rubi [A] (verified)	90
Maple [A] (verified)	91
Fricas [B] (verification not implemented)	92
Sympy [B] (verification not implemented)	92
Maxima [A] (verification not implemented)	93
Giac [B] (verification not implemented)	94
Mupad [B] (verification not implemented)	95
Reduce [B] (verification not implemented)	95

Optimal result

Integrand size = 16, antiderivative size = 130

$$\int \frac{a + b\operatorname{arctanh}(cx)}{(d + ex)^3} dx = \frac{bc}{2(c^2d^2 - e^2)(d + ex)} - \frac{a + b\operatorname{arctanh}(cx)}{2e(d + ex)^2} - \frac{bc^2 \log(1 - cx)}{4e(cd + e)^2} + \frac{bc^2 \log(1 + cx)}{4(cd - e)^2e} - \frac{bc^3d \log(d + ex)}{(c^2d^2 - e^2)^2}$$

output

```
1/2*b*c/(c^2*d^2-e^2)/(e*x+d)-1/2*(a+b*arctanh(c*x))/e/(e*x+d)^2-1/4*b*c^2
*ln(-c*x+1)/e/(c*d+e)^2+1/4*b*c^2*ln(c*x+1)/(c*d-e)^2/e-b*c^3*d*ln(e*x+d)/
(c^2*d^2-e^2)^2
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.02

$$\int \frac{a + b\operatorname{arctanh}(cx)}{(d + ex)^3} dx = \frac{1}{4} \left(-\frac{2a}{e(d + ex)^2} + \frac{2bc}{(c^2d^2 - e^2)(d + ex)} - \frac{2b\operatorname{arctanh}(cx)}{e(d + ex)^2} - \frac{bc^2 \log(1 - cx)}{e(cd + e)^2} + \frac{bc^2 \log(1 + cx)}{e(-cd + e)^2} - \frac{4bc^3d \log(d + ex)}{(-c^2d^2 + e^2)^2} \right)$$

input

```
Integrate[(a + b*ArcTanh[c*x])/(d + e*x)^3,x]
```

output

$$\frac{((-2*a)/(e*(d + e*x)^2) + (2*b*c)/((c^2*d^2 - e^2)*(d + e*x)) - (2*b*ArcTanh[c*x])/(e*(d + e*x)^2) - (b*c^2*Log[1 - c*x])/(e*(c*d + e)^2) + (b*c^2*Log[1 + c*x])/(e*(-(c*d) + e)^2) - (4*b*c^3*d*Log[d + e*x])/(-(c^2*d^2) + e^2)^2)/4$$

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.95, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6478, 477, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^3} dx$$

$$\downarrow 6478$$

$$\frac{bc \int \frac{1}{(d+ex)^2(1-c^2x^2)} dx}{2e} - \frac{a + b \operatorname{arctanh}(cx)}{2e(d + ex)^2}$$

$$\downarrow 477$$

$$\frac{bc \int \left(\frac{e^2}{2(cd+e)^2(1-cx)} + \frac{e^2}{2(cd-e)^2(cx+1)} - \frac{2de^2c^2}{(c^2d^2-e^2)^2(d+ex)} - \frac{e^2}{(c^2d^2-e^2)(d+ex)^2} \right) dx}{2e} - \frac{a + b \operatorname{arctanh}(cx)}{2e(d + ex)^2}$$

$$\downarrow 2009$$

$$\frac{bc \left(\frac{e}{(c^2d^2-e^2)(d+ex)} - \frac{2c^2de \log(d+ex)}{(c^2d^2-e^2)^2} - \frac{c \log(1-cx)}{2(cd+e)^2} + \frac{c \log(cx+1)}{2(cd-e)^2} \right)}{2e} - \frac{a + b \operatorname{arctanh}(cx)}{2e(d + ex)^2}$$

input

```
Int[(a + b*ArcTanh[c*x])/(d + e*x)^3,x]
```

output

$$-1/2*(a + b*ArcTanh[c*x])/(e*(d + e*x)^2) + (b*c*(e/((c^2*d^2 - e^2)*(d + e*x)) - (c*Log[1 - c*x])/(2*(c*d + e)^2) + (c*Log[1 + c*x])/(2*(c*d - e)^2) - (2*c^2*d*e*Log[d + e*x])/(c^2*d^2 - e^2)^2)/(2*e)$$

Defintions of rubi rules used

```
rule 477 Int[((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[
a^p Int[ExpandIntegrand[(c + d*x)^n*(1 - Rt[-b/a, 2]*x)^p*(1 + Rt[-b/a, 2
]*x)^p, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IntegerQ[n] &
& NiceSqrtQ[-b/a] && !FractionalPowerFactorQ[Rt[-b/a, 2]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6478 Int[((a_) + ArcTanh[(c_)*(x_)]*(b_))*((d_) + (e_)*(x_))^(q_), x_Symbol
] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])/(e*(q + 1))), x] - Simp[b
*(c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 - c^2*x^2), x], x] /; FreeQ[{a,
b, c, d, e, q}, x] && NeQ[q, -1]
```

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.09

method	result
parts	$-\frac{a}{2(ex+d)^2e} + \frac{b \left(-\frac{c^3 \operatorname{arctanh}(cx)}{2(cex+cd)^2e} + \frac{c^3 \left(\frac{e}{(cd+e)(cd-e)(cex+cd)} - \frac{2edc \ln(cx+cd)}{(cd+e)^2(cd-e)^2} - \frac{\ln(cx-1)}{2(cd+e)^2} + \frac{\ln(cx+1)}{2(cd-e)^2} \right)}{2e} \right)}{c}$
derivativedivides	$-\frac{ac^3}{2(cex+cd)^2e} + bc^3 \left(-\frac{\operatorname{arctanh}(cx)}{2(cex+cd)^2e} + \frac{e}{(cd+e)(cd-e)(cex+cd)} - \frac{2edc \ln(cx+cd)}{(cd+e)^2(cd-e)^2} - \frac{\ln(cx-1)}{2(cd+e)^2} + \frac{\ln(cx+1)}{2(cd-e)^2} \right)$
default	$-\frac{ac^3}{2(cex+cd)^2e} + bc^3 \left(-\frac{\operatorname{arctanh}(cx)}{2(cex+cd)^2e} + \frac{e}{(cd+e)(cd-e)(cex+cd)} - \frac{2edc \ln(cx+cd)}{(cd+e)^2(cd-e)^2} - \frac{\ln(cx-1)}{2(cd+e)^2} + \frac{\ln(cx+1)}{2(cd-e)^2} \right)$
parallelrisc	$-\frac{e^3 a - x^2 a c^2 e^3 - 2 \ln(ex+d) b c^3 d^3 + x^2 a c^4 d^2 e + 2 b d^3 \operatorname{arctanh}(cx) x c^4 - x^2 b c^3 d e^2 - x b c^3 d^2 e + 3 \operatorname{arctanh}(cx) b c^2 d^2 e - a c^3 d^2 e^2}{c}$
risc	$-\frac{b \ln(cx+1)}{4e(ex+d)^2} + \frac{\ln(-cx+1) b e^4 - b c^4 d^2 e^2 x^2 \ln(-cx+1) - 2 b c^4 d^3 e x \ln(-cx+1) + \ln(-cx-1) b c^2 d^2 e^2 + \ln(-cx-1) b c^2 d^2 e - a c^3 d^2 e^2}{4e(ex+d)^2}$

```
input int((a+b*arctanh(c*x))/(e*x+d)^3,x,method=_RETURNVERBOSE)
```

output

```
-1/2*a/(e*x+d)^2/e+b/c*(-1/2*c^3/(c*e*x+c*d)^2/e*arctanh(c*x)+1/2*c^3/e*(e
/(c*d+e)/(c*d-e)/(c*e*x+c*d)-2*e*d*c/(c*d+e)^2/(c*d-e)^2*ln(c*e*x+c*d)-1/2
/(c*d+e)^2*ln(c*x-1)+1/2/(c*d-e)^2*ln(c*x+1)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 454 vs. $2(122) = 244$.

Time = 0.14 (sec) , antiderivative size = 454, normalized size of antiderivative = 3.49

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^3} dx = \frac{2ac^4d^4 - 2bc^3d^3e - 4ac^2d^2e^2 + 2bcde^3 + 2ae^4 - 2(bc^3d^2e^2 - bce^4)x - (bc^4d^4 + 2bc^3d^3e + bc^2d^2e^2 +$$

input

```
integrate((a+b*arctanh(c*x))/(e*x+d)^3,x, algorithm="fricas")
```

output

```
-1/4*(2*a*c^4*d^4 - 2*b*c^3*d^3*e - 4*a*c^2*d^2*e^2 + 2*b*c*d*e^3 + 2*a*e^4
- 2*(b*c^3*d^2*e^2 - b*c*e^4)*x - (b*c^4*d^4 + 2*b*c^3*d^3*e + b*c^2*d^2
*e^2 + (b*c^4*d^2*e^2 + 2*b*c^3*d*e^3 + b*c^2*e^4)*x^2 + 2*(b*c^4*d^3*e +
2*b*c^3*d^2*e^2 + b*c^2*d*e^3)*x)*log(c*x + 1) + (b*c^4*d^4 - 2*b*c^3*d^3*
e + b*c^2*d^2*e^2 + (b*c^4*d^2*e^2 - 2*b*c^3*d*e^3 + b*c^2*e^4)*x^2 + 2*(b
*c^4*d^3*e - 2*b*c^3*d^2*e^2 + b*c^2*d*e^3)*x)*log(c*x - 1) + 4*(b*c^3*d*e
^3*x^2 + 2*b*c^3*d^2*e^2*x + b*c^3*d^3*e)*log(e*x + d) + (b*c^4*d^4 - 2*b*
c^2*d^2*e^2 + b*e^4)*log(-(c*x + 1)/(c*x - 1)))/(c^4*d^6*e - 2*c^2*d^4*e^3
+ d^2*e^5 + (c^4*d^4*e^3 - 2*c^2*d^2*e^5 + e^7)*x^2 + 2*(c^4*d^5*e^2 - 2*
c^2*d^3*e^4 + d*e^6)*x)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3216 vs. $2(109) = 218$.

Time = 3.31 (sec) , antiderivative size = 3216, normalized size of antiderivative = 24.74

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^3} dx = \text{Too large to display}$$

input `integrate((a+b*atanh(c*x))/(e*x+d)**3,x)`

output `Piecewise((a*x/d**3, Eq(c, 0) & Eq(e, 0)), (-a/(2*d**2*e + 4*d*e**2*x + 2*e**3*x**2), Eq(c, 0)), ((a*x + b*x*atanh(c*x) + b*log(x - 1/c)/c + b*atanh(c*x)/c)/d**3, Eq(e, 0)), (-4*a*d**2/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) + 3*b*d**2*atanh(e*x/d)/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) + 2*b*d**2/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) - 2*b*d*e*x*atanh(e*x/d)/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) + b*d*e*x/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) - b*e**2*x**2*atanh(e*x/d)/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2), Eq(c, -e/d)), (-4*a*d**2/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) - 3*b*d**2*atanh(e*x/d)/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) - 2*b*d**2/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) + 2*b*d*e*x*atanh(e*x/d)/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) - b*d*e*x/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) + b*e**2*x**2*atanh(e*x/d)/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2), Eq(c, e/d)), (-a*c**4*d**4/(2*c**4*d**6*e + 4*c**4*d**5*e**2*x + 2*c**4*d**4*e**3*x**2 - 4*c**2*d**4*e**3 - 8*c**2*d**3*e**4*x - 4*c**2*d**2*e**5*x**2 + 2*d**2*e**5 + 4*d*e**6*x + 2*e**7*x**2) + 2*a*c**2*d**2*e**2/(2*c**4*d**6*e + 4*c**4*d**5*e**2*x + 2*c**4*d**4*e**3*x**2 - 4*c**2*d**4*e**3 - 8*c**2*d**3*e**4*x - 4*c**2*d**2*e**5*x**2 + 2*d**2*e**5 + 4*d*e**6*x + 2*e**7*x**2) - a*e**4/(2*c**4*d**6*e + 4*c**4*d**5*e**2*x + 2*c**4*d**4*e**3*x**2 - 4*c**2*d**4*e**3 - 8*c**2*d**3*e**4*x - 4*c**2*d**2*e**5...`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.46

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^3} dx =$$

$$-\frac{1}{4} \left(\left(\frac{4c^2d \log(ex + d)}{c^4d^4 - 2c^2d^2e^2 + e^4} - \frac{c \log(cx + 1)}{c^2d^2e - 2cde^2 + e^3} + \frac{c \log(cx - 1)}{c^2d^2e + 2cde^2 + e^3} - \frac{2}{c^2d^3 - de^2 + (c^2d^2e - e^3)x} \right) \right.$$

$$\left. - \frac{a}{2(e^3x^2 + 2de^2x + d^2e)} \right)$$

input `integrate((a+b*arctanh(c*x))/(e*x+d)^3,x, algorithm="maxima")`

output

```
-1/4*((4*c^2*d*log(e*x + d)/(c^4*d^4 - 2*c^2*d^2*e^2 + e^4) - c*log(c*x +
1)/(c^2*d^2*e - 2*c*d*e^2 + e^3) + c*log(c*x - 1)/(c^2*d^2*e + 2*c*d*e^2 +
e^3) - 2/(c^2*d^3 - d*e^2 + (c^2*d^2*e - e^3)*x))*c + 2*arctanh(c*x)/(e^3
*x^2 + 2*d*e^2*x + d^2*e))*b - 1/2*a/(e^3*x^2 + 2*d*e^2*x + d^2*e)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 809 vs. $2(122) = 244$.

Time = 0.13 (sec) , antiderivative size = 809, normalized size of antiderivative = 6.22

$$\int \frac{a + \operatorname{arctanh}(cx)}{(d + ex)^3} dx = \text{Too large to display}$$

input

```
integrate((a+b*arctanh(c*x))/(e*x+d)^3,x, algorithm="giac")
```

output

```
-(b*c^2*d*log(-(c*x + 1)*c*d/(c*x - 1) + c*d - (c*x + 1)*e/(c*x - 1) - e)/
(c^4*d^4 - 2*c^2*d^2*e^2 + e^4) - b*c^2*d*log(-(c*x + 1)/(c*x - 1))/(c^4*d
^4 - 2*c^2*d^2*e^2 + e^4) - ((c*x + 1)*b*c^2*d/(c*x - 1) - b*c^2*d + (c*x
+ 1)*b*c*e/(c*x - 1))*log(-(c*x + 1)/(c*x - 1))/((c*x + 1)^2*c^4*d^4/(c*x
- 1)^2 - 2*(c*x + 1)*c^4*d^4/(c*x - 1) + c^4*d^4 + 4*(c*x + 1)^2*c^3*d^3*e
/(c*x - 1)^2 - 4*(c*x + 1)*c^3*d^3*e/(c*x - 1) + 6*(c*x + 1)^2*c^2*d^2*e^2
/(c*x - 1)^2 - 2*c^2*d^2*e^2 + 4*(c*x + 1)^2*c*d*e^3/(c*x - 1)^2 + 4*(c*x
+ 1)*c*d*e^3/(c*x - 1) + (c*x + 1)^2*e^4/(c*x - 1)^2 + 2*(c*x + 1)*e^4/(c*
x - 1) + e^4) - (2*(c*x + 1)*a*c^3*d^2/(c*x - 1) - 2*a*c^3*d^2 + 2*a*c^2*d
*e - (c*x + 1)*b*c^2*d*e/(c*x - 1) + b*c^2*d*e - 2*(c*x + 1)*a*c*e^2/(c*x
- 1) - (c*x + 1)*b*c*e^2/(c*x - 1) - b*c*e^2)/((c*x + 1)^2*c^5*d^5/(c*x -
1)^2 - 2*(c*x + 1)*c^5*d^5/(c*x - 1) + c^5*d^5 + 3*(c*x + 1)^2*c^4*d^4*e/(
c*x - 1)^2 - 2*(c*x + 1)*c^4*d^4*e/(c*x - 1) - c^4*d^4*e + 2*(c*x + 1)^2*c
^3*d^3*e^2/(c*x - 1)^2 + 4*(c*x + 1)*c^3*d^3*e^2/(c*x - 1) - 2*c^3*d^3*e^2
- 2*(c*x + 1)^2*c^2*d^2*e^3/(c*x - 1)^2 + 4*(c*x + 1)*c^2*d^2*e^3/(c*x -
1) + 2*c^2*d^2*e^3 - 3*(c*x + 1)^2*c*d*e^4/(c*x - 1)^2 - 2*(c*x + 1)*c*d*e
^4/(c*x - 1) + c*d*e^4 - (c*x + 1)^2*e^5/(c*x - 1)^2 - 2*(c*x + 1)*e^5/(c*
x - 1) - e^5))*c
```

Mupad [B] (verification not implemented)

Time = 6.71 (sec) , antiderivative size = 427, normalized size of antiderivative = 3.28

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^3} dx = \frac{bc^3 d \ln(c^2 x^2 - 1)}{2(c^4 d^4 - 2c^2 d^2 e^2 + e^4)}$$

$$- \frac{\operatorname{atan}\left(\frac{c^2 x}{\sqrt{-c^2}}\right) (bc^5 d^2 + bc^3 e^2)}{2e^5 \sqrt{-c^2} - c^2 (2d^4 e (-c^2)^{3/2} + 4d^2 e^3 \sqrt{-c^2})} - \frac{bc^3 d \ln(d + ex)}{c^4 d^4 - 2c^2 d^2 e^2 + e^4}$$

$$- \frac{\frac{b \operatorname{atanh}(cx)}{2e} - \frac{x(-ac^2 d^2 + \frac{bcde}{2} + ae^2)}{d(e^2 - c^2 d^2)} - \frac{x^2(-\frac{ac^2 d^2 e}{2} + \frac{bcde^2}{2} + \frac{ae^3}{2})}{d^2(e^2 - c^2 d^2)} + \frac{x^4(-\frac{ac^4 d^2 e}{2} + \frac{bc^3 de^2}{2} + \frac{ac^2 e^3}{2})}{d^2(e^2 - c^2 d^2)} + \frac{x^3(-ac^4 d^2 + \frac{bc^3 e^2}{2})}{d(e^2 - c^2 d^2)}}{-c^2 d^2 x^2 - 2c^2 dex^3 - c^2 e^2 x^4 + d^2 + 2dex + e^2 x^2}$$

input `int((a + b*atanh(c*x))/(d + e*x)^3,x)`output
$$\frac{(bc^3 d \log(c^2 x^2 - 1)) / (2(e^4 + c^4 d^4 - 2c^2 d^2 e^2)) - (\operatorname{atan}((c^2 x) / (-c^2)^{1/2}) * (bc^5 d^2 + bc^3 e^2)) / (2e^5 (-c^2)^{1/2} - c^2 (2d^4 e (-c^2)^{3/2} + 4d^2 e^3 (-c^2)^{1/2})) - (bc^3 d \log(d + ex)) / (e^4 + c^4 d^4 - 2c^2 d^2 e^2) - ((b \operatorname{atanh}(cx)) / (2e) - (x(ae^2 - ac^2 d^2 + (bcde)/2)) / (d(e^2 - c^2 d^2)) - (x^2((ae^3)/2 + (bcde^2)/2 - (ac^2 d^2 e)/2)) / (d^2(e^2 - c^2 d^2)) + (x^4((ac^2 e^3)/2 - (ac^4 d^2 e)/2 + (bc^3 de^2)/2)) / (d^2(e^2 - c^2 d^2)) + (x^3(ac^2 e^2 - ac^4 d^2 + (bc^3 de)/2)) / (d(e^2 - c^2 d^2)) - (bc^2 x^2 \operatorname{atanh}(cx)) / (2e)) / (d^2 + e^2 x^2 + 2dex - c^2 d^2 x^2 - c^2 e^2 x^4 - 2c^2 dex^3)}$$
Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 693, normalized size of antiderivative = 5.33

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^3} dx$$

$$= \frac{-2ac^4 d^6 - 2ad^2 e^4 - bc^3 d^3 e^3 x^2 + bcd e^5 x^2 + 4 \operatorname{atanh}(cx) bc^4 d^5 ex + 2 \operatorname{atanh}(cx) bc^4 d^4 e^2 x^2 + \log(c^2 x - c^2)}{-c^2 d^2 x^2 - 2c^2 dex^3 - c^2 e^2 x^4 + d^2 + 2dex + e^2 x^2}$$

input `int((a+b*atanh(c*x))/(e*x+d)^3,x)`

output

```
(4*atanh(c*x)*b*c**4*d**5*e*x + 2*atanh(c*x)*b*c**4*d**4*e**2*x**2 - 8*atanh(c*x)*b*c**2*d**3*e**3*x - 4*atanh(c*x)*b*c**2*d**2*e**4*x**2 + 4*atanh(c*x)*b*d*e**5*x + 2*atanh(c*x)*b*e**6*x**2 + 2*log(c**2*x - c)*b*c**3*d**5*e + 4*log(c**2*x - c)*b*c**3*d**4*e**2*x + 2*log(c**2*x - c)*b*c**3*d**3*e**3*x**2 - 3*log(c**2*x - c)*b*c**2*d**4*e**2 - 6*log(c**2*x - c)*b*c**2*d**3*e**3*x - 3*log(c**2*x - c)*b*c**2*d**2*e**4*x**2 + log(c**2*x - c)*b*d**2*e**4 + 2*log(c**2*x - c)*b*d*e**5*x + log(c**2*x - c)*b*e**6*x**2 + 2*log(c**2*x + c)*b*c**3*d**5*e + 4*log(c**2*x + c)*b*c**3*d**4*e**2*x + 2*log(c**2*x + c)*b*c**3*d**3*e**3*x**2 + 3*log(c**2*x + c)*b*c**2*d**4*e**2 + 6*log(c**2*x + c)*b*c**2*d**3*e**3*x + 3*log(c**2*x + c)*b*c**2*d**2*e**4*x**2 - log(c**2*x + c)*b*d**2*e**4 - 2*log(c**2*x + c)*b*d*e**5*x - log(c**2*x + c)*b*e**6*x**2 - 4*log(d + e*x)*b*c**3*d**5*e - 8*log(d + e*x)*b*c**3*d**4*e**2*x - 4*log(d + e*x)*b*c**3*d**3*e**3*x**2 - 2*a*c**4*d**6 + 4*a*c**2*d**4*e**2 - 2*a*d**2*e**4 + b*c**3*d**5*e - b*c**3*d**3*e**3*x**2 - b*c*d**3*e**3 + b*c*d*e**5*x**2)/(4*d**2*e*(c**4*d**6 + 2*c**4*d**5*e*x + c**4*d**4*e**2*x**2 - 2*c**2*d**4*e**2 - 4*c**2*d**3*e**3*x - 2*c**2*d**2*e**4*x**2 + d**2*e**4 + 2*d*e**5*x + e**6*x**2))
```

3.8 $\int \frac{a+b\operatorname{arctanh}(cx)}{(d+ex)^4} dx$

Optimal result	97
Mathematica [A] (verified)	98
Rubi [A] (verified)	98
Maple [A] (verified)	100
Fricas [B] (verification not implemented)	100
Sympy [B] (verification not implemented)	101
Maxima [B] (verification not implemented)	102
Giac [B] (verification not implemented)	103
Mupad [B] (verification not implemented)	104
Reduce [B] (verification not implemented)	105

Optimal result

Integrand size = 16, antiderivative size = 174

$$\int \frac{a + b\operatorname{arctanh}(cx)}{(d + ex)^4} dx = \frac{bc}{6(c^2d^2 - e^2)(d + ex)^2} + \frac{2bc^3d}{3(c^2d^2 - e^2)^2(d + ex)}$$

$$- \frac{a + b\operatorname{arctanh}(cx)}{3e(d + ex)^3} - \frac{bc^3 \log(1 - cx)}{6e(cd + e)^3}$$

$$+ \frac{bc^3 \log(1 + cx)}{6(cd - e)^3e} - \frac{bc^3(3c^2d^2 + e^2) \log(d + ex)}{3(c^2d^2 - e^2)^3}$$

output

```
1/6*b*c/(c^2*d^2-e^2)/(e*x+d)^2+2/3*b*c^3*d/(c^2*d^2-e^2)^2/(e*x+d)-1/3*(a
+b*arctanh(c*x))/e/(e*x+d)^3-1/6*b*c^3*ln(-c*x+1)/e/(c*d+e)^3+1/6*b*c^3*ln
(c*x+1)/(c*d-e)^3/e-1/3*b*c^3*(3*c^2*d^2+e^2)*ln(e*x+d)/(c^2*d^2-e^2)^3
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.99

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^4} dx = \frac{1}{6} \left(-\frac{2a}{e(d + ex)^3} + \frac{bc}{(c^2d^2 - e^2)(d + ex)^2} \right. \\ \left. + \frac{4bc^3d}{(-c^2d^2 + e^2)^2(d + ex)} - \frac{2b \operatorname{arctanh}(cx)}{e(d + ex)^3} - \frac{bc^3 \log(1 - cx)}{e(cd + e)^3} \right. \\ \left. + \frac{bc^3 \log(1 + cx)}{(cd - e)^3 e} - \frac{2bc^3(3c^2d^2 + e^2) \log(d + ex)}{(c^2d^2 - e^2)^3} \right)$$

input

```
Integrate[(a + b*ArcTanh[c*x])/(d + e*x)^4,x]
```

output

```
((-2*a)/(e*(d + e*x)^3) + (b*c)/((c^2*d^2 - e^2)*(d + e*x)^2) + (4*b*c^3*d) / ((- (c^2*d^2) + e^2)^2*(d + e*x)) - (2*b*ArcTanh[c*x])/(e*(d + e*x)^3) - (b*c^3*Log[1 - c*x])/(e*(c*d + e)^3) + (b*c^3*Log[1 + c*x])/((c*d - e)^3*e) - (2*b*c^3*(3*c^2*d^2 + e^2)*Log[d + e*x])/(c^2*d^2 - e^2)^3)/6
```

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.98, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6478, 477, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^4} dx \\ \downarrow 6478 \\ \frac{bc \int \frac{1}{(d+ex)^3(1-c^2x^2)} dx}{3e} - \frac{a + b \operatorname{arctanh}(cx)}{3e(d + ex)^3} \\ \downarrow 477$$

$$bc \int \left(\frac{c^3}{2(cd+e)^3(1-cx)} + \frac{c^3}{2(cd-e)^3(cx+1)} - \frac{e^2(3c^2d^2+e^2)c^2}{(c^2d^2-e^2)^3(d+ex)} - \frac{2de^2c^2}{(c^2d^2-e^2)^2(d+ex)^2} - \frac{e^2}{(c^2d^2-e^2)(d+ex)^3} \right) dx$$

$$\frac{3e}{a + \operatorname{arctanh}(cx)} \frac{1}{3e(d+ex)^3}$$

↓ 2009

$$bc \left(\frac{2c^2de}{(c^2d^2-e^2)^2(d+ex)} + \frac{e}{2(c^2d^2-e^2)(d+ex)^2} - \frac{c^2e(3c^2d^2+e^2) \log(d+ex)}{(c^2d^2-e^2)^3} - \frac{c^2 \log(1-cx)}{2(cd+e)^3} + \frac{c^2 \log(cx+1)}{2(cd-e)^3} \right)$$

$$\frac{3e}{a + \operatorname{arctanh}(cx)} \frac{1}{3e(d+ex)^3}$$

input `Int[(a + b*ArcTanh[c*x])/(d + e*x)^4,x]`

output `-1/3*(a + b*ArcTanh[c*x])/(e*(d + e*x)^3) + (b*c*(e/(2*(c^2*d^2 - e^2)*(d + e*x)^2) + (2*c^2*d*e)/((c^2*d^2 - e^2)^2*(d + e*x)) - (c^2*Log[1 - c*x])/(2*(c*d + e)^3) + (c^2*Log[1 + c*x])/(2*(c*d - e)^3) - (c^2*e*(3*c^2*d^2 + e^2)*Log[d + e*x])/(c^2*d^2 - e^2)^3)/(3*e)`

Defintions of rubi rules used

rule 477 `Int[((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p Int[ExpandIntegrand[(c + d*x)^n*(1 - Rt[-b/a, 2]*x)^p*(1 + Rt[-b/a, 2]*x)^p, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IntegerQ[n] && NiceSqrtQ[-b/a] && !FractionalPowerFactorQ[Rt[-b/a, 2]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6478 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)*((d_) + (e_)*(x_))^(q_), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])/(e*(q + 1))), x] - Simp[b*(c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 - c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[q, -1]`

Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.06

method	result
parts	$-\frac{a}{3(ex+d)^3e} + \frac{b}{c} \left(-\frac{c^4 \operatorname{arctanh}(cx)}{3(cex+cd)^3e} + \frac{c^4}{2(cd+e)(cd-e)(cex+cd)^2} - \frac{e(3c^2d^2+e^2) \ln(cex+cd)}{(cd+e)^3(cd-e)^3} + \frac{2edc}{(cd+e)^2(cd-e)^2(cex+cd)} - \frac{\ln(cx-1)}{2(cd-e)} + \frac{1}{2(cd+e)} \right)$
derivativedivides	$-\frac{ac^4}{3(cex+cd)^3e} + bc^4 \left(-\frac{\operatorname{arctanh}(cx)}{3(cex+cd)^3e} + \frac{e}{2(cd+e)(cd-e)(cex+cd)^2} - \frac{e(3c^2d^2+e^2) \ln(cex+cd)}{(cd+e)^3(cd-e)^3} + \frac{2edc}{(cd+e)^2(cd-e)^2(cex+cd)} - \frac{\ln(cx-1)}{2(cd-e)} + \frac{1}{2(cd+e)} \right)$
default	$-\frac{ac^4}{3(cex+cd)^3e} + bc^4 \left(-\frac{\operatorname{arctanh}(cx)}{3(cex+cd)^3e} + \frac{e}{2(cd+e)(cd-e)(cex+cd)^2} - \frac{e(3c^2d^2+e^2) \ln(cex+cd)}{(cd+e)^3(cd-e)^3} + \frac{2edc}{(cd+e)^2(cd-e)^2(cex+cd)} - \frac{\ln(cx-1)}{2(cd-e)} + \frac{1}{2(cd+e)} \right)$
parallelrisch	$\frac{6xa^6d^8 + 6 \operatorname{arctanh}(cx)bc^5d^8 - 6x^2ade^7 - 6xa^2d^2e^6 + 2 \operatorname{arctanh}(cx)bd^3e^5 + 6 \ln(cx-1)bc^5d^8 - 6 \ln(ex+d)bc^5d^8 + 2x^3a^6}{c}$
risch	Expression too large to display

```
input int((a+b*arctanh(c*x))/(e*x+d)^4,x,method=_RETURNVERBOSE)
```

```
output -1/3*a/(e*x+d)^3/e+b/c*(-1/3*c^4/(c*e*x+c*d)^3/e*arctanh(c*x)+1/3*c^4/e*(1/2*e/(c*d+e)/(c*d-e)/(c*e*x+c*d)^2-e*(3*c^2*d^2+e^2)/(c*d+e)^3/(c*d-e)^3*ln(c*e*x+c*d)+2*e*d*c/(c*d+e)^2/(c*d-e)^2/(c*e*x+c*d)-1/2/(c*d+e)^3*ln(c*x-1)+1/2/(c*d-e)^3*ln(c*x+1)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 859 vs. 2(162) = 324.

Time = 0.29 (sec) , antiderivative size = 859, normalized size of antiderivative = 4.94

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^4} dx = \text{Too large to display}$$

```
input integrate((a+b*arctanh(c*x))/(e*x+d)^4,x, algorithm="fricas")
```

output

```

-1/6*(2*a*c^6*d^6 - 5*b*c^5*d^5*e - 6*a*c^4*d^4*e^2 + 6*b*c^3*d^3*e^3 + 6*
a*c^2*d^2*e^4 - b*c*d*e^5 - 2*a*e^6 - 4*(b*c^5*d^3*e^3 - b*c^3*d*e^5)*x^2
- (9*b*c^5*d^4*e^2 - 10*b*c^3*d^2*e^4 + b*c*e^6)*x - (b*c^6*d^6 + 3*b*c^5*
d^5*e + 3*b*c^4*d^4*e^2 + b*c^3*d^3*e^3 + (b*c^6*d^3*e^3 + 3*b*c^5*d^2*e^4
+ 3*b*c^4*d*e^5 + b*c^3*e^6)*x^3 + 3*(b*c^6*d^4*e^2 + 3*b*c^5*d^3*e^3 + 3
*b*c^4*d^2*e^4 + b*c^3*d*e^5)*x^2 + 3*(b*c^6*d^5*e + 3*b*c^5*d^4*e^2 + 3*b
*c^4*d^3*e^3 + b*c^3*d^2*e^4)*x)*log(c*x + 1) + (b*c^6*d^6 - 3*b*c^5*d^5*e
+ 3*b*c^4*d^4*e^2 - b*c^3*d^3*e^3 + (b*c^6*d^3*e^3 - 3*b*c^5*d^2*e^4 + 3*
b*c^4*d*e^5 - b*c^3*e^6)*x^3 + 3*(b*c^6*d^4*e^2 - 3*b*c^5*d^3*e^3 + 3*b*c^
4*d^2*e^4 - b*c^3*d*e^5)*x^2 + 3*(b*c^6*d^5*e - 3*b*c^5*d^4*e^2 + 3*b*c^4*
d^3*e^3 - b*c^3*d^2*e^4)*x)*log(c*x - 1) + 2*(3*b*c^5*d^5*e + b*c^3*d^3*e^
3 + (3*b*c^5*d^2*e^4 + b*c^3*e^6)*x^3 + 3*(3*b*c^5*d^3*e^3 + b*c^3*d*e^5)*
x^2 + 3*(3*b*c^5*d^4*e^2 + b*c^3*d^2*e^4)*x)*log(e*x + d) + (b*c^6*d^6 - 3
*b*c^4*d^4*e^2 + 3*b*c^2*d^2*e^4 - b*e^6)*log(-(c*x + 1)/(c*x - 1))/(c^6*
d^9*e - 3*c^4*d^7*e^3 + 3*c^2*d^5*e^5 - d^3*e^7 + (c^6*d^6*e^4 - 3*c^4*d^4
*e^6 + 3*c^2*d^2*e^8 - e^10)*x^3 + 3*(c^6*d^7*e^3 - 3*c^4*d^5*e^5 + 3*c^2*
d^3*e^7 - d*e^9)*x^2 + 3*(c^6*d^8*e^2 - 3*c^4*d^6*e^4 + 3*c^2*d^4*e^6 - d^
2*e^8)*x)

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10946 vs. $2(150) = 300$.

Time = 5.15 (sec) , antiderivative size = 10946, normalized size of antiderivative = 62.91

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^4} dx = \text{Too large to display}$$

input

```
integrate((a+b*atanh(c*x))/(e*x+d)**4,x)
```

output

```
Piecewise((a*x/d**4, Eq(c, 0) & Eq(e, 0)), (-a/(3*d**3*e + 9*d**2*e**2*x +
9*d*e**3*x**2 + 3*e**4*x**3), Eq(c, 0)), ((a*x + b*x*atanh(c*x) + b*log(x
- 1/c)/c + b*atanh(c*x)/c)/d**4, Eq(e, 0)), (-24*a*d**3/(72*d**6*e + 216*
d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3) + 21*b*d**3*atanh(e*
x/d)/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3
) + 10*b*d**3/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*
e**4*x**3) - 9*b*d**2*e*x*atanh(e*x/d)/(72*d**6*e + 216*d**5*e**2*x + 216*
d**4*e**3*x**2 + 72*d**3*e**4*x**3) + 9*b*d**2*e*x/(72*d**6*e + 216*d**5*
e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3) - 9*b*d**2*x**2*atanh(e*
x/d)/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3
) + 3*b*d**2*x**2/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72
*d**3*e**4*x**3) - 3*b*e**3*x**3*atanh(e*x/d)/(72*d**6*e + 216*d**5*e**2*x
+ 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3), Eq(c, -e/d)), (-24*a*d**3/(72*
d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3) - 21*b*
d**3*atanh(e*x/d)/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d
**3*e**4*x**3) - 10*b*d**3/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x*
*2 + 72*d**3*e**4*x**3) + 9*b*d**2*e*x*atanh(e*x/d)/(72*d**6*e + 216*d**5*
e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3) - 9*b*d**2*e*x/(72*d**6*
e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3) + 9*b*d**2*
x**2*atanh(e*x/d)/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 7...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 339 vs. $2(162) = 324$.

Time = 0.04 (sec) , antiderivative size = 339, normalized size of antiderivative = 1.95

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^4} dx$$

$$= \frac{1}{6} \left(\left(\frac{c^2 \log(cx + 1)}{c^3 d^3 e - 3c^2 d^2 e^2 + 3c d e^3 - e^4} - \frac{c^2 \log(cx - 1)}{c^3 d^3 e + 3c^2 d^2 e^2 + 3c d e^3 + e^4} - \frac{2(3c^4 d^2 + c^2 e^2) \log(ex + d)}{c^6 d^6 - 3c^4 d^4 e^2 + 3c^2 d^2 e^4 - e^6} \right) - \frac{a}{3(e^4 x^3 + 3d e^3 x^2 + 3d^2 e^2 x + d^3 e)} \right)$$

input

```
integrate((a+b*arctanh(c*x))/(e*x+d)^4,x, algorithm="maxima")
```

output

```
1/6*((c^2*log(c*x + 1)/(c^3*d^3*e - 3*c^2*d^2*e^2 + 3*c*d*e^3 - e^4) - c^2
*log(c*x - 1)/(c^3*d^3*e + 3*c^2*d^2*e^2 + 3*c*d*e^3 + e^4) - 2*(3*c^4*d^2
+ c^2*e^2)*log(e*x + d)/(c^6*d^6 - 3*c^4*d^4*e^2 + 3*c^2*d^2*e^4 - e^6) +
(4*c^2*d*e*x + 5*c^2*d^2 - e^2)/(c^4*d^6 - 2*c^2*d^4*e^2 + d^2*e^4 + (c^4
*d^4*e^2 - 2*c^2*d^2*e^4 + e^6)*x^2 + 2*(c^4*d^5*e - 2*c^2*d^3*e^3 + d*e^5
)*x))*c - 2*arctanh(c*x)/(e^4*x^3 + 3*d*e^3*x^2 + 3*d^2*e^2*x + d^3*e))*b
- 1/3*a/(e^4*x^3 + 3*d*e^3*x^2 + 3*d^2*e^2*x + d^3*e)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1792 vs. $2(162) = 324$.

Time = 0.17 (sec) , antiderivative size = 1792, normalized size of antiderivative = 10.30

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^4} dx = \text{Too large to display}$$

input

```
integrate((a+b*arctanh(c*x))/(e*x+d)^4,x, algorithm="giac")
```


output

```

-1/3*c*((3*b*c^4*d^2 + b*c^2*e^2)*log(-(c*x + 1)*c*d/(c*x - 1) + c*d - (c*
x + 1)*e/(c*x - 1) - e)/(c^6*d^6 - 3*c^4*d^4*e^2 + 3*c^2*d^2*e^4 - e^6) -
(3*(c*x + 1)^2*b*c^4*d^2/(c*x - 1)^2 - 6*(c*x + 1)*b*c^4*d^2/(c*x - 1) + 3
*b*c^4*d^2 + 6*(c*x + 1)^2*b*c^3*d*e/(c*x - 1)^2 - 6*(c*x + 1)*b*c^3*d*e/(
c*x - 1) + 3*(c*x + 1)^2*b*c^2*e^2/(c*x - 1)^2 + b*c^2*e^2)*log(-(c*x + 1)
/(c*x - 1))/((c*x + 1)^3*c^6*d^6/(c*x - 1)^3 - 3*(c*x + 1)^2*c^6*d^6/(c*x
- 1)^2 + 3*(c*x + 1)*c^6*d^6/(c*x - 1) - c^6*d^6 + 6*(c*x + 1)^3*c^5*d^5*e
/(c*x - 1)^3 - 12*(c*x + 1)^2*c^5*d^5*e/(c*x - 1)^2 + 6*(c*x + 1)*c^5*d^5*
e/(c*x - 1) + 15*(c*x + 1)^3*c^4*d^4*e^2/(c*x - 1)^3 - 15*(c*x + 1)^2*c^4*
d^4*e^2/(c*x - 1)^2 - 3*(c*x + 1)*c^4*d^4*e^2/(c*x - 1) + 3*c^4*d^4*e^2 +
20*(c*x + 1)^3*c^3*d^3*e^3/(c*x - 1)^3 - 12*(c*x + 1)*c^3*d^3*e^3/(c*x - 1
) + 15*(c*x + 1)^3*c^2*d^2*e^4/(c*x - 1)^3 + 15*(c*x + 1)^2*c^2*d^2*e^4/(c
*x - 1)^2 - 3*(c*x + 1)*c^2*d^2*e^4/(c*x - 1) - 3*c^2*d^2*e^4 + 6*(c*x + 1
)^3*c*d*e^5/(c*x - 1)^3 + 12*(c*x + 1)^2*c*d*e^5/(c*x - 1)^2 + 6*(c*x + 1
)*c*d*e^5/(c*x - 1) + (c*x + 1)^3*e^6/(c*x - 1)^3 + 3*(c*x + 1)^2*e^6/(c*x
- 1)^2 + 3*(c*x + 1)*e^6/(c*x - 1) + e^6) - (3*b*c^4*d^2 + b*c^2*e^2)*log(
-(c*x + 1)/(c*x - 1))/(c^6*d^6 - 3*c^4*d^4*e^2 + 3*c^2*d^2*e^4 - e^6) - 2*
(3*(c*x + 1)^2*a*c^6*d^4/(c*x - 1)^2 - 6*(c*x + 1)*a*c^6*d^4/(c*x - 1) + 3
*a*c^6*d^4 + 6*(c*x + 1)*a*c^5*d^3*e/(c*x - 1) - 6*a*c^5*d^3*e - 3*(c*x +
1)^2*b*c^5*d^3*e/(c*x - 1)^2 + 6*(c*x + 1)*b*c^5*d^3*e/(c*x - 1) - 3*b*...

```

Mupad [B] (verification not implemented)

Time = 5.04 (sec) , antiderivative size = 418, normalized size of antiderivative = 2.40

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^4} dx \\
 &= \ln(d + ex) \left(\frac{bc^3}{6e(e + cd)^3} + \frac{bc^3}{6e(e - cd)^3} \right) \\
 & - \frac{2ac^4d^4 - 5bc^3d^3e - 4a^2c^2d^2e^2 + bcde^3 + 2ae^4}{2(c^4d^4 - 2c^2d^2e^2 + e^4)} + \frac{x(bc^4e^4 - 9bc^3d^2e^2)}{2(c^4d^4 - 2c^2d^2e^2 + e^4)} - \frac{2bc^3de^3x^2}{c^4d^4 - 2c^2d^2e^2 + e^4} \\
 & - \frac{3d^3e + 9d^2e^2x + 9de^3x^2 + 3e^4x^3}{bc^3 \ln(cx - 1)} - \frac{bc^3 \ln(cx + 1)}{6c^3d^3e + 18c^2d^2e^2 + 18cde^3 + 6e^4} - \frac{-6c^3d^3e + 18c^2d^2e^2 - 18cde^3 + 6e^4}{b \ln(cx + 1)} \\
 & - \frac{6e(d^3 + 3d^2ex + 3de^2x^2 + e^3x^3)}{6e(d^3 + 3d^2ex + 3de^2x^2 + e^3x^3)} + \frac{3e(2d^3 + 6d^2ex + 6de^2x^2 + 2e^3x^3)}{3e(2d^3 + 6d^2ex + 6de^2x^2 + 2e^3x^3)}
 \end{aligned}$$

input

```
int((a + b*atanh(c*x))/(d + e*x)^4,x)
```

output

```
log(d + e*x)*((b*c^3)/(6*e*(e + c*d)^3) + (b*c^3)/(6*e*(e - c*d)^3)) - ((2
*a*e^4 + 2*a*c^4*d^4 - 4*a*c^2*d^2*e^2 + b*c*d*e^3 - 5*b*c^3*d^3*e)/(2*(e^
4 + c^4*d^4 - 2*c^2*d^2*e^2)) + (x*(b*c*e^4 - 9*b*c^3*d^2*e^2))/(2*(e^4 +
c^4*d^4 - 2*c^2*d^2*e^2)) - (2*b*c^3*d*e^3*x^2)/(e^4 + c^4*d^4 - 2*c^2*d^2
*e^2))/(3*d^3*e + 3*e^4*x^3 + 9*d^2*e^2*x + 9*d*e^3*x^2) - (b*c^3*log(c*x
- 1))/(6*e^4 + 6*c^3*d^3*e + 18*c^2*d^2*e^2 + 18*c*d*e^3) - (b*c^3*log(c*x
+ 1))/(6*e^4 - 6*c^3*d^3*e + 18*c^2*d^2*e^2 - 18*c*d*e^3) - (b*log(c*x +
1))/(6*e*(d^3 + e^3*x^3 + 3*d*e^2*x^2 + 3*d^2*e*x)) + (b*log(1 - c*x))/(3*
e*(2*d^3 + 2*e^3*x^3 + 6*d*e^2*x^2 + 6*d^2*e*x))
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 1565, normalized size of antiderivative = 8.99

$$\int \frac{a + b \operatorname{arctanh}(cx)}{(d + ex)^4} dx = \text{Too large to display}$$

input

```
int((a+b*atanh(c*x))/(e*x+d)^4,x)
```

output

```
(18*atanh(c*x)*b*c**6*d**8*e*x + 18*atanh(c*x)*b*c**6*d**7*e**2*x**2 + 6*
tanh(c*x)*b*c**6*d**6*e**3*x**3 - 54*atanh(c*x)*b*c**4*d**6*e**3*x - 54*at
anh(c*x)*b*c**4*d**5*e**4*x**2 - 18*atanh(c*x)*b*c**4*d**4*e**5*x**3 + 54*
atanh(c*x)*b*c**2*d**4*e**5*x + 54*atanh(c*x)*b*c**2*d**3*e**6*x**2 + 18*a
tanh(c*x)*b*c**2*d**2*e**7*x**3 - 18*atanh(c*x)*b*d**2*e**7*x - 18*atanh(c
*x)*b*d*e**8*x**2 - 6*atanh(c*x)*b*e**9*x**3 + 9*log(c**2*x - c)*b*c**5*d*
**8*e + 27*log(c**2*x - c)*b*c**5*d**7*e**2*x + 27*log(c**2*x - c)*b*c**5*d
**6*e**3*x**2 + 9*log(c**2*x - c)*b*c**5*d**5*e**4*x**3 - 18*log(c**2*x -
c)*b*c**4*d**7*e**2 - 54*log(c**2*x - c)*b*c**4*d**6*e**3*x - 54*log(c**2*
x - c)*b*c**4*d**5*e**4*x**2 - 18*log(c**2*x - c)*b*c**4*d**4*e**5*x**3 +
3*log(c**2*x - c)*b*c**3*d**6*e**3 + 9*log(c**2*x - c)*b*c**3*d**5*e**4*x
+ 9*log(c**2*x - c)*b*c**3*d**4*e**5*x**2 + 3*log(c**2*x - c)*b*c**3*d**3*
e**6*x**3 + 9*log(c**2*x - c)*b*c**2*d**5*e**4 + 27*log(c**2*x - c)*b*c**2
*d**4*e**5*x + 27*log(c**2*x - c)*b*c**2*d**3*e**6*x**2 + 9*log(c**2*x - c
)*b*c**2*d**2*e**7*x**3 - 3*log(c**2*x - c)*b*d**3*e**6 - 9*log(c**2*x - c
)*b*d**2*e**7*x - 9*log(c**2*x - c)*b*d*e**8*x**2 - 3*log(c**2*x - c)*b*e
**9*x**3 + 9*log(c**2*x + c)*b*c**5*d**8*e + 27*log(c**2*x + c)*b*c**5*d**7
*e**2*x + 27*log(c**2*x + c)*b*c**5*d**6*e**3*x**2 + 9*log(c**2*x + c)*b*c
**5*d**5*e**4*x**3 + 18*log(c**2*x + c)*b*c**4*d**7*e**2 + 54*log(c**2*x +
c)*b*c**4*d**6*e**3*x + 54*log(c**2*x + c)*b*c**4*d**5*e**4*x**2 + 18*...
```

3.9 $\int (d + ex)^3 (a + b \operatorname{arctanh}(cx))^2 dx$

Optimal result	108
Mathematica [A] (verified)	109
Rubi [A] (verified)	109
Maple [B] (verified)	111
Fricas [F]	112
Sympy [F]	113
Maxima [B] (verification not implemented)	113
Giac [F]	114
Mupad [F(-1)]	115
Reduce [F]	115

Optimal result

Integrand size = 18, antiderivative size = 359

$$\begin{aligned}
 \int (d + ex)^3 (a + \operatorname{arctanh}(cx))^2 dx = & \frac{b^2 d e^2 x}{c^2} + \frac{a b e (6c^2 d^2 + e^2) x}{2c^3} \\
 & + \frac{b^2 e^3 x^2}{12c^2} - \frac{b^2 d e^2 \operatorname{arctanh}(cx)}{c^3} \\
 & + \frac{b^2 e (6c^2 d^2 + e^2) x \operatorname{arctanh}(cx)}{2c^3} \\
 & + \frac{b d e^2 x^2 (a + \operatorname{arctanh}(cx))}{c} \\
 & + \frac{b e^3 x^3 (a + \operatorname{arctanh}(cx))}{6c} \\
 & + \frac{d (c^2 d^2 + e^2) (a + \operatorname{arctanh}(cx))^2}{c^3} \\
 & - \frac{(c^4 d^4 + 6c^2 d^2 e^2 + e^4) (a + \operatorname{arctanh}(cx))^2}{4c^4 e} \\
 & + \frac{(d + ex)^4 (a + \operatorname{arctanh}(cx))^2}{4e} \\
 & - \frac{2bd(c^2 d^2 + e^2) (a + \operatorname{arctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{c^3} \\
 & + \frac{b^2 e^3 \log(1 - c^2 x^2)}{12c^4} \\
 & + \frac{b^2 e (6c^2 d^2 + e^2) \log(1 - c^2 x^2)}{4c^4} \\
 & - \frac{b^2 d (c^2 d^2 + e^2) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{c^3}
 \end{aligned}$$

output

```

b^2*d*e^2*x/c^2+1/2*a*b*e*(6*c^2*d^2+e^2)*x/c^3+1/12*b^2*e^3*x^2/c^2-b^2*d
*e^2*arctanh(c*x)/c^3+1/2*b^2*e*(6*c^2*d^2+e^2)*x*arctanh(c*x)/c^3+b*d*e^2
*x^2*(a+b*arctanh(c*x))/c+1/6*b*e^3*x^3*(a+b*arctanh(c*x))/c+d*(c^2*d^2+e^
2)*(a+b*arctanh(c*x))^2/c^3-1/4*(c^4*d^4+6*c^2*d^2*e^2+e^4)*(a+b*arctanh(c
*x))^2/c^4/e+1/4*(e*x+d)^4*(a+b*arctanh(c*x))^2/e-2*b*d*(c^2*d^2+e^2)*(a+b
*arctanh(c*x))*ln(2/(-c*x+1))/c^3+1/12*b^2*e^3*ln(-c^2*x^2+1)/c^4+1/4*b^2*
e*(6*c^2*d^2+e^2)*ln(-c^2*x^2+1)/c^4-b^2*d*(c^2*d^2+e^2)*polylog(2,1-2/(-c
*x+1))/c^3

```

Mathematica [A] (verified)

Time = 0.57 (sec) , antiderivative size = 506, normalized size of antiderivative = 1.41

$$\int (d + ex)^3 (a + \text{barctanh}(cx))^2 dx$$

$$= \frac{-b^2 e^3 + 12a^2 c^4 d^3 x + 36abc^3 d^2 ex + 12b^2 c^2 de^2 x + 6abce^3 x + 18a^2 c^4 d^2 ex^2 + 12abc^3 de^2 x^2 + b^2 c^2 e^3 x^2 + 12abc^3 d^2 ex^2 + 12abc^3 de^2 x^2 + b^2 c^2 e^3 x^2 + 12abc^3 d^2 ex^2 + 12abc^3 de^2 x^2 + b^2 c^2 e^3 x^2}{12c^4}$$

input `Integrate[(d + e*x)^3*(a + b*ArcTanh[c*x])^2,x]`

output
$$\frac{(-b^2 e^3 + 12a^2 c^4 d^3 x + 36a^2 b c^3 d^2 e x + 12b^2 c^2 d e^2 x + 6a^2 b c^3 d^2 e x + 12a^2 b c^3 d^2 e x + 12b^2 c^2 d e^2 x + b^2 c^2 e^3 x^2 + 12a^2 c^4 d^3 x + 2a^2 b c^3 d^2 e x + 2a^2 b c^3 d^2 e x + 3a^2 c^4 d^2 e x^2 + 3b^2 c^2 d e^2 x + 4d^3 e^2 x^2 + e^3 x^3) \text{ArcTanh}[c x]^2 + 2b c^3 \text{ArcTanh}[c x] (3a^2 c^3 d^2 e x^2 + 6d^2 e^2 x + 4d e^2 x^2 + e^3 x^3) + b e (18c^2 d^2 x + 6d e (-1 + c^2 x^2) + e^2 x (3 + c^2 x^2)) - 12b d (c^2 d^2 + e^2) \text{Log}[1 + E^{-2 \text{ArcTanh}[c x]}] + 18a^2 b c^2 d^2 e \text{Log}[1 - c x] + 3a^2 b e^3 \text{Log}[1 - c x] - 18a^2 b c^2 d^2 e \text{Log}[1 + c x] - 3a^2 b e^3 \text{Log}[1 + c x] + 12a^2 b c^3 d^3 \text{Log}[1 - c^2 x^2] + 18b^2 c^2 d^2 e \text{Log}[1 - c^2 x^2] + 4b^2 e^3 \text{Log}[1 - c^2 x^2] + 12a^2 b c d e^2 \text{Log}[-1 + c^2 x^2] + 12b^2 c d (c^2 d^2 + e^2) \text{PolyLog}[2, -E^{-2 \text{ArcTanh}[c x]}]]}{12c^4}$$

Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 362, normalized size of antiderivative = 1.01, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex)^3 (a + \text{barctanh}(cx))^2 dx$$

↓ 6480

$$bc \int \frac{(d+ex)^4(a+\operatorname{arctanh}(cx))^2}{4e} - \frac{x^2(a+\operatorname{arctanh}(cx))e^4}{c^2} - \frac{4dx(a+\operatorname{arctanh}(cx))e^3}{c^2} - \frac{(6c^2d^2+e^2)(a+\operatorname{arctanh}(cx))e^2}{c^4} + \frac{(c^4d^4+6c^2e^2d^2+4c^2e(c^2d^2+e^2)xd)}{c^4(1-c^2x^2)}$$

$$2e$$

↓ 2009

$$bc \left(-\frac{2de^3x^2(a+\operatorname{arctanh}(cx))}{c^2} - \frac{e^4x^3(a+\operatorname{arctanh}(cx))}{3c^2} - \frac{2de(c^2d^2+e^2)(a+\operatorname{arctanh}(cx))^2}{bc^4} + \frac{4de(c^2d^2+e^2)\log\left(\frac{2}{1-cx}\right)(a+\operatorname{arctanh}(cx))}{c^4} \right)$$

input `Int[(d + e*x)^3*(a + b*ArcTanh[c*x])^2,x]`

output `((d + e*x)^4*(a + b*ArcTanh[c*x])^2)/(4*e) - (b*c*((-2*b*d*e^3*x)/c^3 - (a*e^2*(6*c^2*d^2 + e^2)*x)/c^4 - (b*e^4*x^2)/(6*c^3) + (2*b*d*e^3*ArcTanh[c*x])/c^4 - (b*e^2*(6*c^2*d^2 + e^2)*x*ArcTanh[c*x])/c^4 - (2*d*e^3*x^2*(a + b*ArcTanh[c*x]))/c^2 - (e^4*x^3*(a + b*ArcTanh[c*x]))/(3*c^2) - (2*d*e*(c^2*d^2 + e^2)*(a + b*ArcTanh[c*x])^2)/(b*c^4) + ((c^4*d^4 + 6*c^2*d^2*e^2 + e^4)*(a + b*ArcTanh[c*x])^2)/(2*b*c^5) + (4*d*e*(c^2*d^2 + e^2)*(a + b*ArcTanh[c*x])*Log[2/(1 - c*x)])/c^4 - (b*e^4*Log[1 - c^2*x^2])/(6*c^5) - (b*e^2*(6*c^2*d^2 + e^2)*Log[1 - c^2*x^2])/(2*c^5) + (2*b*d*e*(c^2*d^2 + e^2)*PolyLog[2, 1 - 2/(1 - c*x)])/c^4)/(2*e)`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6480 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^p_*((d_.) + (e_.)*(x_))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 803 vs. 2(343) = 686.

Time = 0.29 (sec) , antiderivative size = 804, normalized size of antiderivative = 2.24

method	result
parts	$\frac{a^2(e x+d)^4}{4 e} + \frac{b^2 \left(\frac{c e^3 \operatorname{arctanh}(c x)^2 x^4}{4} + c e^2 \operatorname{arctanh}(c x)^2 x^3 d + \frac{3 c e \operatorname{arctanh}(c x)^2 x^2 d^2}{2} + \operatorname{arctanh}(c x)^2 c x d^3 + \frac{c \operatorname{arctanh}(c x)}{4 e} \right)}{4 e}$
derivativdivides	$\frac{a^2(c e x+c d)^4}{4 c^3 e} + \frac{b^2 \left(\frac{\operatorname{arctanh}(c x)^2 c^4 d^4}{4 e} + \operatorname{arctanh}(c x)^2 c^4 d^3 x + \frac{3 e \operatorname{arctanh}(c x)^2 c^4 d^2 x^2}{2} + e^2 \operatorname{arctanh}(c x)^2 c^4 d x^3 + \frac{e^3 \operatorname{arctanh}(c x)^2 c^4 x}{4} \right)}{4 c^3 e}$
default	$\frac{a^2(c e x+c d)^4}{4 c^3 e} + \frac{b^2 \left(\frac{\operatorname{arctanh}(c x)^2 c^4 d^4}{4 e} + \operatorname{arctanh}(c x)^2 c^4 d^3 x + \frac{3 e \operatorname{arctanh}(c x)^2 c^4 d^2 x^2}{2} + e^2 \operatorname{arctanh}(c x)^2 c^4 d x^3 + \frac{e^3 \operatorname{arctanh}(c x)^2 c^4 x}{4} \right)}{4 c^3 e}$
risch	Expression too large to display

input

```
int((e*x+d)^3*(a+b*arctanh(c*x))^2,x,method=_RETURNVERBOSE)
```


output

```

1/4*a^2*(e*x+d)^4/e+b^2/c*(1/4*c*e^3*arctanh(c*x)^2*x^4+c*e^2*arctanh(c*x)
^2*x^3*d+3/2*c*e*arctanh(c*x)^2*x^2*d^2+arctanh(c*x)^2*c*x*d^3+1/4*c/e*arc
tanh(c*x)^2*d^4-1/2/c^3/e*(-6*arctanh(c*x)*c^3*d^2*e^2*x-2*arctanh(c*x)*c^
3*d*e^3*x^2-1/3*arctanh(c*x)*e^4*c^3*x^3-arctanh(c*x)*e^4*c*x-1/2*arctanh(
c*x)*ln(c*x-1)*c^4*d^4-2*arctanh(c*x)*ln(c*x-1)*c^3*d^3*e-3*arctanh(c*x)*l
n(c*x-1)*c^2*d^2*e^2-2*arctanh(c*x)*ln(c*x-1)*c*d*e^3-1/2*arctanh(c*x)*ln(
c*x-1)*e^4+1/2*arctanh(c*x)*ln(c*x+1)*c^4*d^4-2*arctanh(c*x)*ln(c*x+1)*c^3
*d^3*e+3*arctanh(c*x)*ln(c*x+1)*c^2*d^2*e^2-2*arctanh(c*x)*ln(c*x+1)*c*d*e
^3+1/2*arctanh(c*x)*ln(c*x+1)*e^4-1/6*(-3*c^4*d^4+12*c^3*d^3*e-18*c^2*d^2*
e^2+12*c*d*e^3-3*e^4)*(-1/4*ln(c*x+1)^2+1/2*(ln(c*x+1)-ln(1/2*c*x+1/2))*ln
(-1/2*c*x+1/2)-1/2*dilog(1/2*c*x+1/2))-1/6*(3*c^4*d^4+12*c^3*d^3*e+18*c^2*
d^2*e^2+12*c*d*e^3+3*e^4)*(1/4*ln(c*x-1)^2-1/2*dilog(1/2*c*x+1/2)-1/2*ln(c
*x-1)*ln(1/2*c*x+1/2))-1/3*e^2*(6*c^2*d*e*x+1/2*e^2*c^2*x^2+1/2*(18*c^2*d^
2+6*c*d*e+4*e^2)*ln(c*x-1)-1/2*(-18*c^2*d^2+6*c*d*e-4*e^2)*ln(c*x+1))))+2*
a*b/c*(1/4*c*e^3*arctanh(c*x)*x^4+c*e^2*arctanh(c*x)*x^3*d+3/2*c*e*arctanh
(c*x)*x^2*d^2+arctanh(c*x)*c*x*d^3+1/4*c/e*arctanh(c*x)*d^4-1/4/c^3/e*(-6*
c^3*d^2*e^2*x-2*c^3*d*e^3*x^2-1/3*e^4*c^3*x^3-e^4*c*x-1/2*(c^4*d^4+4*c^3*d
^3*e+6*c^2*d^2*e^2+4*c*d*e^3+e^4)*ln(c*x-1)+1/2*(c^4*d^4-4*c^3*d^3*e+6*c^2
*d^2*e^2-4*c*d*e^3+e^4)*ln(c*x+1)))

```

Fricas [F]

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx))^2 dx = \int (ex + d)^3 (b \operatorname{arctanh}(cx) + a)^2 dx$$

input

```
integrate((e*x+d)^3*(a+b*arctanh(c*x))^2,x, algorithm="fricas")
```

output

```

integral(a^2*e^3*x^3 + 3*a^2*d*e^2*x^2 + 3*a^2*d^2*e*x + a^2*d^3 + (b^2*e^
3*x^3 + 3*b^2*d*e^2*x^2 + 3*b^2*d^2*e*x + b^2*d^3)*arctanh(c*x)^2 + 2*(a*b
*e^3*x^3 + 3*a*b*d*e^2*x^2 + 3*a*b*d^2*e*x + a*b*d^3)*arctanh(c*x), x)

```

Sympy [F]

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx))^2 dx = \int (a + b \operatorname{atanh}(cx))^2 (d + ex)^3 dx$$

input `integrate((e*x+d)**3*(a+b*atanh(c*x))**2,x)`

output `Integral((a + b*atanh(c*x))**2*(d + e*x)**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 782 vs. $2(340) = 680$.

Time = 0.20 (sec) , antiderivative size = 782, normalized size of antiderivative = 2.18

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx))^2 dx = \text{Too large to display}$$

input `integrate((e*x+d)^3*(a+b*arctanh(c*x))^2,x, algorithm="maxima")`

output

```

1/4*a^2*e^3*x^4 + a^2*d*e^2*x^3 + 3/2*a^2*d^2*e*x^2 + 3/2*(2*x^2*arctanh(c
*x) + c*(2*x/c^2 - log(c*x + 1)/c^3 + log(c*x - 1)/c^3))*a*b*d^2*e + (2*x^
3*arctanh(c*x) + c*(x^2/c^2 + log(c^2*x^2 - 1)/c^4))*a*b*d*e^2 + 1/12*(6*x
^4*arctanh(c*x) + c*(2*(c^2*x^3 + 3*x)/c^4 - 3*log(c*x + 1)/c^5 + 3*log(c*
x - 1)/c^5))*a*b*e^3 + a^2*d^3*x + (2*c*x*arctanh(c*x) + log(-c^2*x^2 + 1)
)*a*b*d^3/c + (c^2*d^3 + d*e^2)*(log(c*x + 1)*log(-1/2*c*x + 1/2) + dilog(
1/2*c*x + 1/2))*b^2/c^3 + 1/6*(9*c^2*d^2*e - 3*c*d*e^2 + 2*e^3)*b^2*log(c*
x + 1)/c^4 + 1/6*(9*c^2*d^2*e + 3*c*d*e^2 + 2*e^3)*b^2*log(c*x - 1)/c^4 +
1/48*(4*b^2*c^2*e^3*x^2 + 48*b^2*c^2*d*e^2*x + 3*(b^2*c^4*e^3*x^4 + 4*b^2*
c^4*d*e^2*x^3 + 6*b^2*c^4*d^2*e*x^2 + 4*b^2*c^4*d^3*x + (4*c^3*d^3 - 6*c^2
*d^2*e + 4*c*d*e^2 - e^3)*b^2)*log(c*x + 1)^2 + 3*(b^2*c^4*e^3*x^4 + 4*b^2
*c^4*d*e^2*x^3 + 6*b^2*c^4*d^2*e*x^2 + 4*b^2*c^4*d^3*x - (4*c^3*d^3 + 6*c^
2*d^2*e + 4*c*d*e^2 + e^3)*b^2)*log(-c*x + 1)^2 + 4*(b^2*c^3*e^3*x^3 + 6*b
^2*c^3*d*e^2*x^2 + 3*(6*c^3*d^2*e + c*e^3)*b^2*x)*log(c*x + 1) - 2*(2*b^2*
c^3*e^3*x^3 + 12*b^2*c^3*d*e^2*x^2 + 6*(6*c^3*d^2*e + c*e^3)*b^2*x + 3*(b^
2*c^4*e^3*x^4 + 4*b^2*c^4*d*e^2*x^3 + 6*b^2*c^4*d^2*e*x^2 + 4*b^2*c^4*d^3*
x + (4*c^3*d^3 - 6*c^2*d^2*e + 4*c*d*e^2 - e^3)*b^2)*log(c*x + 1))*log(-c*
x + 1))/c^4

```

Giac [F]

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx))^2 dx = \int (ex + d)^3 (b \operatorname{arctanh}(cx) + a)^2 dx$$

input

```
integrate((e*x+d)^3*(a+b*arctanh(c*x))^2,x, algorithm="giac")
```

output

```
integrate((e*x + d)^3*(b*arctanh(c*x) + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx))^2 dx = \int (a + b \operatorname{atanh}(cx))^2 (d + ex)^3 dx$$

input `int((a + b*atanh(c*x))^2*(d + e*x)^3,x)`output `int((a + b*atanh(c*x))^2*(d + e*x)^3, x)`**Reduce [F]**

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{8 \log(c^2 x - c) b^2 e^3 + 8 \operatorname{atanh}(cx) b^2 e^3 - 3 \operatorname{atanh}(cx)^2 b^2 e^3 + 24 \left(\int \frac{\operatorname{atanh}(cx) x}{c^2 x^2 - 1} dx \right) b^2 c^3 d e^2 + 12 \operatorname{atanh}(cx)^2 b^2 c^3 d e^2}{1}$$

input `int((e*x+d)^3*(a+b*atanh(c*x))^2,x)`

output

```
(12*atanh(c*x)**2*b**2*c**4*d**3*x + 18*atanh(c*x)**2*b**2*c**4*d**2*e*x**
2 + 12*atanh(c*x)**2*b**2*c**4*d*e**2*x**3 + 3*atanh(c*x)**2*b**2*c**4*e**
3*x**4 - 18*atanh(c*x)**2*b**2*c**2*d**2*e - 3*atanh(c*x)**2*b**2*e**3 + 2
4*atanh(c*x)*a*b*c**4*d**3*x + 36*atanh(c*x)*a*b*c**4*d**2*e*x**2 + 24*ata
nh(c*x)*a*b*c**4*d*e**2*x**3 + 6*atanh(c*x)*a*b*c**4*e**3*x**4 + 24*atanh(
c*x)*a*b*c**3*d**3 - 36*atanh(c*x)*a*b*c**2*d**2*e + 24*atanh(c*x)*a*b*c*d
*e**2 - 6*atanh(c*x)*a*b*e**3 + 36*atanh(c*x)*b**2*c**3*d**2*e*x + 12*atan
h(c*x)*b**2*c**3*d*e**2*x**2 + 2*atanh(c*x)*b**2*c**3*e**3*x**3 + 36*atanh
(c*x)*b**2*c**2*d**2*e - 12*atanh(c*x)*b**2*c*d*e**2 + 6*atanh(c*x)*b**2*c
*e**3*x + 8*atanh(c*x)*b**2*e**3 + 24*int((atanh(c*x)*x)/(c**2*x**2 - 1),x
)*b**2*c**5*d**3 + 24*int((atanh(c*x)*x)/(c**2*x**2 - 1),x)*b**2*c**3*d*e
**2 + 24*log(c**2*x - c)*a*b*c**3*d**3 + 24*log(c**2*x - c)*a*b*c*d*e**2 +
36*log(c**2*x - c)*b**2*c**2*d**2*e + 8*log(c**2*x - c)*b**2*e**3 + 12*a**
2*c**4*d**3*x + 18*a**2*c**4*d**2*e*x**2 + 12*a**2*c**4*d*e**2*x**3 + 3*a
**2*c**4*e**3*x**4 + 36*a*b*c**3*d**2*e*x + 12*a*b*c**3*d*e**2*x**2 + 2*a*b
*c**3*e**3*x**3 + 6*a*b*c*e**3*x + 12*b**2*c**2*d*e**2*x + b**2*c**2*e**3*
x**2)/(12*c**4)
```

3.10 $\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^2 dx$

Optimal result	117
Mathematica [A] (verified)	118
Rubi [A] (verified)	118
Maple [B] (verified)	120
Fricas [F]	121
Sympy [F]	121
Maxima [B] (verification not implemented)	122
Giac [F]	123
Mupad [F(-1)]	123
Reduce [F]	123

Optimal result

Integrand size = 18, antiderivative size = 257

$$\begin{aligned}
 \int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^2 dx = & \frac{2abdex}{c} + \frac{b^2 e^2 x}{3c^2} - \frac{b^2 e^2 \operatorname{arctanh}(cx)}{3c^3} \\
 & + \frac{2b^2 dex \operatorname{arctanh}(cx)}{c} + \frac{be^2 x^2 (a + b \operatorname{arctanh}(cx))}{3c} \\
 & + \frac{(3c^2 d^2 + e^2) (a + b \operatorname{arctanh}(cx))^2}{3c^3} \\
 & - \frac{d \left(d^2 + \frac{3e^2}{c^2} \right) (a + b \operatorname{arctanh}(cx))^2}{3e} \\
 & + \frac{(d + ex)^3 (a + b \operatorname{arctanh}(cx))^2}{3e} \\
 & - \frac{2b(3c^2 d^2 + e^2) (a + b \operatorname{arctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{3c^3} \\
 & + \frac{b^2 de \log(1 - c^2 x^2)}{c^2} \\
 & - \frac{b^2 (3c^2 d^2 + e^2) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{3c^3}
 \end{aligned}$$

output

```
2*a*b*d*e*x/c+1/3*b^2*e^2*x/c^2-1/3*b^2*e^2*arctanh(c*x)/c^3+2*b^2*d*e*x*a
rctanh(c*x)/c+1/3*b*e^2*x^2*(a+b*arctanh(c*x))/c+1/3*(3*c^2*d^2+e^2)*(a+b*
arctanh(c*x))^2/c^3-1/3*d*(d^2+3*e^2/c^2)*(a+b*arctanh(c*x))^2/e+1/3*(e*x+
d)^3*(a+b*arctanh(c*x))^2/e-2/3*b*(3*c^2*d^2+e^2)*(a+b*arctanh(c*x))*ln(2/
(-c*x+1))/c^3+b^2*d*e*ln(-c^2*x^2+1)/c^2-1/3*b^2*(3*c^2*d^2+e^2)*polylog(2
,1-2/(-c*x+1))/c^3
```

Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 319, normalized size of antiderivative = 1.24

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{3a^2c^3d^2x + 6abc^2dex + b^2ce^2x + 3a^2c^3dex^2 + abc^2e^2x^2 + a^2c^3e^2x^3 + b^2(-1 + cx)(e^2 + ce(3d + ex) + c^2)}{c^3}$$

input

```
Integrate[(d + e*x)^2*(a + b*ArcTanh[c*x])^2,x]
```

output

```
(3*a^2*c^3*d^2*x + 6*a*b*c^2*d*e*x + b^2*c*e^2*x + 3*a^2*c^3*d*e*x^2 + a*b
*c^2*e^2*x^2 + a^2*c^3*e^2*x^3 + b^2*(-1 + c*x)*(e^2 + c*e*(3*d + e*x) + c
^2*(3*d^2 + 3*d*e*x + e^2*x^2))*ArcTanh[c*x]^2 + b*ArcTanh[c*x]*(b*e*(-e +
6*c^2*d*x + c^2*e*x^2) + 2*a*c^3*x*(3*d^2 + 3*d*e*x + e^2*x^2) - 2*b*(3*c
^2*d^2 + e^2)*Log[1 + E^(-2*ArcTanh[c*x])]) + 3*a*b*c*d*e*Log[1 - c*x] - 3
*a*b*c*d*e*Log[1 + c*x] + 3*a*b*c^2*d^2*Log[1 - c^2*x^2] + 3*b^2*c*d*e*Log
[1 - c^2*x^2] + a*b*e^2*Log[-1 + c^2*x^2] + b^2*(3*c^2*d^2 + e^2)*PolyLog[
2, -E^(-2*ArcTanh[c*x])])/(3*c^3)
```

Rubi [A] (verified)Time = 1.02 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.05, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int (d + ex)^2 (a + \operatorname{arctanh}(cx))^2 dx \\
& \quad \downarrow \text{6480} \\
& \frac{(d + ex)^3 (a + \operatorname{arctanh}(cx))^2}{3e} - \\
& \frac{2bc \int \left(-\frac{x(a + \operatorname{arctanh}(cx))e^3}{c^2} - \frac{3d(a + \operatorname{arctanh}(cx))e^2}{c^2} + \frac{(d(c^2d^2 + 3e^2) + e(3c^2d^2 + e^2)x)(a + \operatorname{arctanh}(cx))}{c^2(1 - c^2x^2)} \right) dx}{3e} \\
& \quad \downarrow \text{2009} \\
& \frac{(d + ex)^3 (a + \operatorname{arctanh}(cx))^2}{3e} - \\
& \frac{2bc \left(-\frac{e^3x^2(a + \operatorname{arctanh}(cx))}{2c^2} - \frac{e(3c^2d^2 + e^2)(a + \operatorname{arctanh}(cx))^2}{2bc^4} + \frac{e(3c^2d^2 + e^2) \log\left(\frac{2}{1 - cx}\right)(a + \operatorname{arctanh}(cx))}{c^4} + \frac{d(c^2d^2 + 3e^2)(a + \operatorname{arctanh}(cx))}{2bc^4} \right)}{3e}
\end{aligned}$$

input `Int[(d + e*x)^2*(a + b*ArcTanh[c*x])^2,x]`

output `((d + e*x)^3*(a + b*ArcTanh[c*x])^2)/(3*e) - (2*b*c*((-3*a*d*e^2*x)/c^2 - (b*e^3*x)/(2*c^3) + (b*e^3*ArcTanh[c*x])/(2*c^4) - (3*b*d*e^2*x*ArcTanh[c*x])/c^2 - (e^3*x^2*(a + b*ArcTanh[c*x]))/(2*c^2) - (e*(3*c^2*d^2 + e^2)*(a + b*ArcTanh[c*x])^2)/(2*b*c^4) + (d*(c^2*d^2 + 3*e^2)*(a + b*ArcTanh[c*x])^2)/(2*b*c^3) + (e*(3*c^2*d^2 + e^2)*(a + b*ArcTanh[c*x])*Log[2/(1 - c*x)])/c^4 - (3*b*d*e^2*Log[1 - c^2*x^2])/(2*c^3) + (b*e*(3*c^2*d^2 + e^2)*PolyLog[2, 1 - 2/(1 - c*x)]/(2*c^4))/(3*e)`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6480 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^p*((d_) + (e_.)*(x_))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 - c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 589 vs. 2(241) = 482.

Time = 0.20 (sec) , antiderivative size = 590, normalized size of antiderivative = 2.30

method	result
parts	$\frac{a^2 (ex+d)^3}{3e} + b^2 \left(\frac{c e^2 \operatorname{arctanh}(cx)^2 x^3}{3} + ce \operatorname{arctanh}(cx)^2 x^2 d + \operatorname{arctanh}(cx)^2 cx d^2 + \frac{c \operatorname{arctanh}(cx)^2 d^3}{3e} - \frac{2 \left(-3 \operatorname{arctanh}(cx) c^2 d^2 \right)}{3e} \right)$
derivativedivides	$\frac{a^2 (cex+cd)^3}{3e^2} + b^2 \left(\frac{\operatorname{arctanh}(cx)^2 c^3 d^3}{3e} + \operatorname{arctanh}(cx)^2 c^3 d^2 x + e \operatorname{arctanh}(cx)^2 c^3 d x^2 + \frac{e^2 \operatorname{arctanh}(cx)^2 c^3 x^3}{3} - \frac{2 \left(-3 \operatorname{arctanh}(cx) c^2 d^2 \right)}{3e} \right)$
default	$\frac{a^2 (cex+cd)^3}{3e^2} + b^2 \left(\frac{\operatorname{arctanh}(cx)^2 c^3 d^3}{3e} + \operatorname{arctanh}(cx)^2 c^3 d^2 x + e \operatorname{arctanh}(cx)^2 c^3 d x^2 + \frac{e^2 \operatorname{arctanh}(cx)^2 c^3 x^3}{3} - \frac{2 \left(-3 \operatorname{arctanh}(cx) c^2 d^2 \right)}{3e} \right)$
risch	$\frac{2abdex}{c} + \frac{b^2 e^2 x}{3c^2} + \frac{b^2 \ln\left(-\frac{cx}{2} + \frac{1}{2}\right) \ln\left(\frac{cx}{2} + \frac{1}{2}\right) e^2}{3c^3} - \frac{b^2 \ln\left(\frac{cx}{2} + \frac{1}{2}\right) \ln(-cx+1) e^2}{3c^3} + \frac{b^2 \ln\left(-\frac{cx}{2} + \frac{1}{2}\right) \ln\left(\frac{cx}{2} + \frac{1}{2}\right) d^2}{c}$

input

```
int((e*x+d)^2*(a+b*arctanh(c*x))^2,x,method=_RETURNVERBOSE)
```

output

```
1/3*a^2*(e*x+d)^3/e+b^2/c*(1/3*c*e^2*arctanh(c*x)^2*x^3+c*e*arctanh(c*x)^2
*x^2*d+arctanh(c*x)^2*c*x*d^2+1/3*c/e*arctanh(c*x)^2*d^3-2/3/c^2/e*(-3*arc
tanh(c*x)*c^2*d*e^2*x-1/2*arctanh(c*x)*e^3*c^2*x^2-1/2*arctanh(c*x)*ln(c*x
-1)*c^3*d^3-3/2*arctanh(c*x)*ln(c*x-1)*c^2*d^2*e-3/2*arctanh(c*x)*ln(c*x-1
)*c*d*e^2-1/2*arctanh(c*x)*ln(c*x-1)*e^3+1/2*arctanh(c*x)*ln(c*x+1)*c^3*d^
3-3/2*arctanh(c*x)*ln(c*x+1)*c^2*d^2*e+3/2*arctanh(c*x)*ln(c*x+1)*c*d*e^2-
1/2*arctanh(c*x)*ln(c*x+1)*e^3-1/2*(c^3*d^3+3*c^2*d^2*e+3*c*d*e^2+e^3)*(1/
4*ln(c*x-1)^2-1/2*dilog(1/2*c*x+1/2)-1/2*ln(c*x-1)*ln(1/2*c*x+1/2))-1/2*e^
2*(c*e*x+1/2*(6*c*d+e)*ln(c*x-1)-1/2*(-6*c*d+e)*ln(c*x+1))-1/2*(-c^3*d^3+3
*c^2*d^2*e-3*c*d*e^2+e^3)*(-1/4*ln(c*x+1)^2+1/2*(ln(c*x+1)-ln(1/2*c*x+1/2
))*ln(-1/2*c*x+1/2)-1/2*dilog(1/2*c*x+1/2))))+2*a*b/c*(1/3*c*e^2*arctanh(c*
x)*x^3+c*e*arctanh(c*x)*x^2*d+arctanh(c*x)*c*x*d^2+1/3*c/e*arctanh(c*x)*d^
3-1/3/c^2/e*(-3*c^2*d*e^2*x-1/2*e^3*c^2*x^2-1/2*(c^3*d^3+3*c^2*d^2*e+3*c*d
*e^2+e^3)*ln(c*x-1)+1/2*(c^3*d^3-3*c^2*d^2*e+3*c*d*e^2-e^3)*ln(c*x+1)))
```

Fricas [F]

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^2 dx = \int (ex + d)^2 (b \operatorname{artanh}(cx) + a)^2 dx$$

input

```
integrate((e*x+d)^2*(a+b*arctanh(c*x))^2,x, algorithm="fricas")
```

output

```
integral(a^2*e^2*x^2 + 2*a^2*d*e*x + a^2*d^2 + (b^2*e^2*x^2 + 2*b^2*d*e*x
+ b^2*d^2)*arctanh(c*x)^2 + 2*(a*b*e^2*x^2 + 2*a*b*d*e*x + a*b*d^2)*arctan
h(c*x), x)
```

Sympy [F]

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^2 dx = \int (a + b \operatorname{atanh}(cx))^2 (d + ex)^2 dx$$

input

```
integrate((e*x+d)**2*(a+b*atanh(c*x))**2,x)
```

output

```
Integral((a + b*atanh(c*x))**2*(d + e*x)**2, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 524 vs. $2(238) = 476$.

Time = 0.19 (sec) , antiderivative size = 524, normalized size of antiderivative = 2.04

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{1}{3} a^2 e^2 x^3 + a^2 d e x^2 + \left(2 x^2 \operatorname{arctanh}(cx) + c \left(\frac{2x}{c^2} - \frac{\log(cx+1)}{c^3} + \frac{\log(cx-1)}{c^3} \right) \right) abde$$

$$+ \frac{1}{3} \left(2 x^3 \operatorname{arctanh}(cx) + c \left(\frac{x^2}{c^2} + \frac{\log(c^2 x^2 - 1)}{c^4} \right) \right) abe^2$$

$$+ a^2 d^2 x + \frac{(2cx \operatorname{arctanh}(cx) + \log(-c^2 x^2 + 1)) abd^2}{c}$$

$$+ \frac{(3c^2 d^2 + e^2) (\log(cx+1) \log(-\frac{1}{2}cx + \frac{1}{2}) + \operatorname{Li}_2(\frac{1}{2}cx + \frac{1}{2})) b^2}{3c^3}$$

$$+ \frac{(6cde - e^2) b^2 \log(cx+1)}{6c^3} + \frac{(6cde + e^2) b^2 \log(cx-1)}{6c^3}$$

$$+ \frac{4b^2 ce^2 x + (b^2 c^3 e^2 x^3 + 3b^2 c^3 d e x^2 + 3b^2 c^3 d^2 x + (3c^2 d^2 - 3cde + e^2) b^2) \log(cx+1)^2 + (b^2 c^3 e^2 x^3 + 3b^2 c^3 d e x^2 + 3b^2 c^3 d^2 x + (3c^2 d^2 - 3cde + e^2) b^2) \log(-cx+1)}{6c^3}$$

input `integrate((e*x+d)^2*(a+b*arctanh(c*x))^2,x, algorithm="maxima")`

output

```
1/3*a^2*e^2*x^3 + a^2*d*e*x^2 + (2*x^2*arctanh(c*x) + c*(2*x/c^2 - log(c*x
+ 1)/c^3 + log(c*x - 1)/c^3))*a*b*d*e + 1/3*(2*x^3*arctanh(c*x) + c*(x^2/
c^2 + log(c^2*x^2 - 1)/c^4))*a*b*e^2 + a^2*d^2*x + (2*c*x*arctanh(c*x) + l
og(-c^2*x^2 + 1))*a*b*d^2/c + 1/3*(3*c^2*d^2 + e^2)*(log(c*x + 1)*log(-1/2
*c*x + 1/2) + dilog(1/2*c*x + 1/2))*b^2/c^3 + 1/6*(6*c*d*e - e^2)*b^2*log(
c*x + 1)/c^3 + 1/6*(6*c*d*e + e^2)*b^2*log(c*x - 1)/c^3 + 1/12*(4*b^2*c*e^
2*x + (b^2*c^3*e^2*x^3 + 3*b^2*c^3*d*e*x^2 + 3*b^2*c^3*d^2*x + (3*c^2*d^2
- 3*c*d*e + e^2)*b^2)*log(c*x + 1)^2 + (b^2*c^3*e^2*x^3 + 3*b^2*c^3*d*e*x^
2 + 3*b^2*c^3*d^2*x - (3*c^2*d^2 + 3*c*d*e + e^2)*b^2)*log(-c*x + 1)^2 + 2
*(b^2*c^2*e^2*x^2 + 6*b^2*c^2*d*e*x)*log(c*x + 1) - 2*(b^2*c^2*e^2*x^2 + 6
*b^2*c^2*d*e*x + (b^2*c^3*e^2*x^3 + 3*b^2*c^3*d*e*x^2 + 3*b^2*c^3*d^2*x +
(3*c^2*d^2 - 3*c*d*e + e^2)*b^2)*log(c*x + 1))*log(-c*x + 1)/c^3
```

Giac [F]

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^2 dx = \int (ex + d)^2 (b \operatorname{artanh}(cx) + a)^2 dx$$

input `integrate((e*x+d)^2*(a+b*arctanh(c*x))^2,x, algorithm="giac")`

output `integrate((e*x + d)^2*(b*arctanh(c*x) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^2 dx = \int (a + b \operatorname{atanh}(cx))^2 (d + ex)^2 dx$$

input `int((a + b*atanh(c*x))^2*(d + e*x)^2,x)`

output `int((a + b*atanh(c*x))^2*(d + e*x)^2, x)`

Reduce [F]

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{\operatorname{atanh}(cx)^2 b^2 c^3 e^2 x^3 + \operatorname{atanh}(cx) b^2 c^2 e^2 x^2 + ab c^2 e^2 x^2 + 3 \operatorname{atanh}(cx)^2 b^2 c^3 d e x^2 + 6 \operatorname{atanh}(cx) ab c^3 d^2 x + \dots}{\dots}$$

input `int((e*x+d)^2*(a+b*atanh(c*x))^2,x)`

output

```
(3*atanh(c*x)**2*b**2*c**3*d**2*x + 3*atanh(c*x)**2*b**2*c**3*d*e*x**2 + a
tanh(c*x)**2*b**2*c**3*e**2*x**3 - 3*atanh(c*x)**2*b**2*c*d*e + 6*atanh(c*
x)*a*b*c**3*d**2*x + 6*atanh(c*x)*a*b*c**3*d*e*x**2 + 2*atanh(c*x)*a*b*c**
3*e**2*x**3 + 6*atanh(c*x)*a*b*c**2*d**2 - 6*atanh(c*x)*a*b*c*d*e + 2*atan
h(c*x)*a*b*e**2 + 6*atanh(c*x)*b**2*c**2*d*e*x + atanh(c*x)*b**2*c**2*e**2
*x**2 + 6*atanh(c*x)*b**2*c*d*e - atanh(c*x)*b**2*e**2 + 6*int((atanh(c*x)
*x)/(c**2*x**2 - 1),x)*b**2*c**4*d**2 + 2*int((atanh(c*x)*x)/(c**2*x**2 -
1),x)*b**2*c**2*e**2 + 6*log(c**2*x - c)*a*b*c**2*d**2 + 2*log(c**2*x - c)
*a*b*e**2 + 6*log(c**2*x - c)*b**2*c*d*e + 3*a**2*c**3*d**2*x + 3*a**2*c**
3*d*e*x**2 + a**2*c**3*e**2*x**3 + 6*a*b*c**2*d*e*x + a*b*c**2*e**2*x**2 +
b**2*c*e**2*x)/(3*c**3)
```

3.11 $\int (d + ex)(a + b \operatorname{arctanh}(cx))^2 dx$

Optimal result	125
Mathematica [A] (verified)	126
Rubi [A] (verified)	126
Maple [A] (verified)	128
Fricas [F]	129
Sympy [F]	129
Maxima [B] (verification not implemented)	130
Giac [F]	130
Mupad [F(-1)]	131
Reduce [F]	131

Optimal result

Integrand size = 16, antiderivative size = 160

$$\int (d + ex)(a + b \operatorname{arctanh}(cx))^2 dx = \frac{abex}{c} + \frac{b^2 ex \operatorname{arctanh}(cx)}{c} + \frac{d(a + b \operatorname{arctanh}(cx))^2}{c} - \frac{\left(d^2 + \frac{e^2}{c^2}\right)(a + b \operatorname{arctanh}(cx))^2}{2e} + \frac{(d + ex)^2(a + b \operatorname{arctanh}(cx))^2}{2e} - \frac{2bd(a + b \operatorname{arctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{c} + \frac{b^2 e \log(1 - c^2 x^2)}{2c^2} - \frac{b^2 d \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{c}$$

output

```
a*b*e*x/c+b^2*e*x*arctanh(c*x)/c+d*(a+b*arctanh(c*x))^2/c-1/2*(d^2+e^2/c^2)*(a+b*arctanh(c*x))^2/e+1/2*(e*x+d)^2*(a+b*arctanh(c*x))^2/e-2*b*d*(a+b*arctanh(c*x))*ln(2/(-c*x+1))/c+1/2*b^2*e*ln(-c^2*x^2+1)/c^2-b^2*d*polylog(2,1-2/(-c*x+1))/c
```

Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.09

$$\int (d + ex)(a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{2a^2c^2dx + 2abcex + a^2c^2ex^2 + b^2(-1 + cx)(2cd + e + cex)\operatorname{arctanh}(cx)^2 + 2bc\operatorname{arctanh}(cx)(bex + acx(2$$

input

```
Integrate[(d + e*x)*(a + b*ArcTanh[c*x])^2,x]
```

output

```
(2*a^2*c^2*d*x + 2*a*b*c*e*x + a^2*c^2*e*x^2 + b^2*(-1 + c*x)*(2*c*d + e +
c*e*x)*ArcTanh[c*x]^2 + 2*b*c*ArcTanh[c*x]*(b*e*x + a*c*x*(2*d + e*x) - 2
*b*d*Log[1 + E^(-2*ArcTanh[c*x])]) + a*b*e*Log[1 - c*x] - a*b*e*Log[1 + c*
x] + 2*a*b*c*d*Log[1 - c^2*x^2] + b^2*e*Log[1 - c^2*x^2] + 2*b^2*c*d*PolyL
og[2, -E^(-2*ArcTanh[c*x])])/(2*c^2)
```

Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.11, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex)(a + b \operatorname{arctanh}(cx))^2 dx$$

$$\downarrow \text{6480}$$

$$\frac{(d + ex)^2(a + b \operatorname{arctanh}(cx))^2}{2e}$$

$$\frac{bc \int \left(\frac{(d^2c^2 + 2dexc^2 + e^2)(a + b \operatorname{arctanh}(cx))}{c^2(1 - c^2x^2)} - \frac{e^2(a + b \operatorname{arctanh}(cx))}{c^2} \right) dx}{e}$$

$$\downarrow \text{2009}$$

$$\frac{(d + ex)^2(a + \operatorname{arctanh}(cx))^2}{2e} - bc \left(-\frac{de(a + \operatorname{arctanh}(cx))^2}{bc^2} + \frac{2de \log\left(\frac{2}{1-cx}\right)(a + \operatorname{arctanh}(cx))}{c^2} + \frac{(c^2d^2 + e^2)(a + \operatorname{arctanh}(cx))^2}{2bc^3} - \frac{ae^2x}{c^2} - \frac{be^2x \operatorname{arctanh}(cx)}{c^2} + \dots \right)$$

input `Int[(d + e*x)*(a + b*ArcTanh[c*x])^2, x]`

output `((d + e*x)^2*(a + b*ArcTanh[c*x])^2)/(2*e) - (b*c*(-((a*e^2*x)/c^2) - (b*e^2*x*ArcTanh[c*x])/c^2 - (d*e*(a + b*ArcTanh[c*x])^2)/(b*c^2) + ((c^2*d^2 + e^2)*(a + b*ArcTanh[c*x])^2)/(2*b*c^3) + (2*d*e*(a + b*ArcTanh[c*x])*Log[2/(1 - c*x)])/c^2 - (b*e^2*Log[1 - c^2*x^2])/(2*c^3) + (b*d*e*PolyLog[2, 1 - 2/(1 - c*x)])/c^2)/e`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6480 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^p*((d_) + (e_.)*(x_))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 298, normalized size of antiderivative = 1.86

method	result
parts	$a^2 \left(\frac{1}{2} e x^2 + dx \right) + \frac{b^2 \left(\frac{c \operatorname{arctanh}(cx)^2 x^2 e}{2} + \operatorname{arctanh}(cx)^2 dx - \frac{\operatorname{arctanh}(cx) e c x - \operatorname{arctanh}(cx) \ln(cx-1) c d - \frac{\operatorname{arctanh}(cx)}{2}}{2} \right)}{c}$
derivativedivides	$\frac{a^2 \left(c^2 dx + \frac{1}{2} c^2 e x^2 \right)}{c} + \frac{b^2 \left(\operatorname{arctanh}(cx)^2 d c^2 x + \frac{\operatorname{arctanh}(cx)^2 e c^2 x^2}{2} + \operatorname{arctanh}(cx) e c x + \operatorname{arctanh}(cx) \ln(cx-1) c d + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{2} \right)}{c}$
default	$\frac{a^2 \left(c^2 dx + \frac{1}{2} c^2 e x^2 \right)}{c} + \frac{b^2 \left(\operatorname{arctanh}(cx)^2 d c^2 x + \frac{\operatorname{arctanh}(cx)^2 e c^2 x^2}{2} + \operatorname{arctanh}(cx) e c x + \operatorname{arctanh}(cx) \ln(cx-1) c d + \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{2} \right)}{c}$
risch	$\frac{a b e x}{c} + a^2 dx + \frac{b^2 \ln(-cx-1) e}{2c^2} + \frac{b^2 \operatorname{dilog}\left(-\frac{cx}{2} + \frac{1}{2}\right) d}{c} + \frac{e a^2 x^2}{2} - \frac{b^2 e \ln(-cx+1)^2}{8c^2} + \frac{b^2 e \ln(-cx+1)}{2c^2} - \frac{\ln(-cx+1)}{2c}$

input `int((e*x+d)*(a+b*arctanh(c*x))^2,x,method=_RETURNVERBOSE)`

output
$$a^2 \left(\frac{1}{2} e x^2 + dx \right) + \frac{b^2}{c} \left(\frac{1}{2} c \operatorname{arctanh}(cx)^2 x^2 e + \operatorname{arctanh}(cx)^2 d c x - \frac{1}{c} \left(-\operatorname{arctanh}(cx) e c x - \operatorname{arctanh}(cx) \ln(cx-1) c d - \frac{1}{2} \operatorname{arctanh}(cx) \ln(cx-1) e - \operatorname{arctanh}(cx) \ln(cx+1) c d + \frac{1}{2} \operatorname{arctanh}(cx) \ln(cx+1) e - \frac{1}{2} (2 c d - e) \left(-\frac{1}{4} \ln^2(cx+1) + \frac{1}{2} (\ln(cx+1) - \ln(\frac{1}{2} c x + \frac{1}{2})) \ln(-\frac{1}{2} c x + \frac{1}{2}) - \frac{1}{2} \operatorname{dilog}(\frac{1}{2} c x + \frac{1}{2})) - \frac{1}{2} (2 c d + e) \left(\frac{1}{4} \ln^2(cx-1) - \frac{1}{2} \operatorname{dilog}(\frac{1}{2} c x + \frac{1}{2}) - \frac{1}{2} \ln(cx-1) \ln(\frac{1}{2} c x + \frac{1}{2}) \right) - \frac{1}{2} \ln(cx-1) e - \frac{1}{2} \ln(cx+1) e \right) \right) + 2 a b / c \left(\frac{1}{2} c \operatorname{arctanh}(cx) x^2 e + \operatorname{arctanh}(cx) d c x - \frac{1}{2} c \left(-c e x - \frac{1}{2} (2 c d + e) \ln(cx-1) + \frac{1}{2} (-2 c d + e) \ln(cx+1) \right) \right)$$

Fricas [F]

$$\int (d + ex)(a + b \operatorname{arctanh}(cx))^2 dx = \int (ex + d)(b \operatorname{artanh}(cx) + a)^2 dx$$

input `integrate((e*x+d)*(a+b*arctanh(c*x))^2,x, algorithm="fricas")`

output `integral(a^2*e*x + a^2*d + (b^2*e*x + b^2*d)*arctanh(c*x)^2 + 2*(a*b*e*x + a*b*d)*arctanh(c*x), x)`

Sympy [F]

$$\int (d + ex)(a + b \operatorname{arctanh}(cx))^2 dx = \int (a + b \operatorname{atanh}(cx))^2 (d + ex) dx$$

input `integrate((e*x+d)*(a+b*atanh(c*x))**2,x)`

output `Integral((a + b*atanh(c*x))**2*(d + e*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 313 vs. $2(151) = 302$.

Time = 0.21 (sec) , antiderivative size = 313, normalized size of antiderivative = 1.96

$$\begin{aligned} & \int (d + ex)(a + b \operatorname{arctanh}(cx))^2 dx \\ &= \frac{1}{2} a^2 ex^2 + \frac{1}{2} \left(2x^2 \operatorname{arctanh}(cx) + c \left(\frac{2x}{c^2} - \frac{\log(cx+1)}{c^3} + \frac{\log(cx-1)}{c^3} \right) \right) abe \\ &+ a^2 dx + \frac{(2cx \operatorname{arctanh}(cx) + \log(-c^2x^2 + 1))abd}{c} \\ &+ \frac{(\log(cx+1) \log(-\frac{1}{2}cx + \frac{1}{2}) + \operatorname{Li}_2(\frac{1}{2}cx + \frac{1}{2}))b^2d}{c} \\ &+ \frac{b^2e \log(cx+1)}{2c^2} + \frac{b^2e \log(cx-1)}{2c^2} \\ &+ \frac{4b^2cex \log(cx+1) + (b^2c^2ex^2 + 2b^2c^2dx + (2cd - e)b^2) \log(cx+1)^2 + (b^2c^2ex^2 + 2b^2c^2dx - (2cd + e)b^2) \log(cx-1)^2}{8c^2} \end{aligned}$$

input `integrate((e*x+d)*(a+b*arctanh(c*x))^2,x, algorithm="maxima")`

output `1/2*a^2*e*x^2 + 1/2*(2*x^2*arctanh(c*x) + c*(2*x/c^2 - log(c*x + 1)/c^3 + log(c*x - 1)/c^3))*a*b*e + a^2*d*x + (2*c*x*arctanh(c*x) + log(-c^2*x^2 + 1))*a*b*d/c + (log(c*x + 1)*log(-1/2*c*x + 1/2) + dilog(1/2*c*x + 1/2))*b^2*d/c + 1/2*b^2*e*log(c*x + 1)/c^2 + 1/2*b^2*e*log(c*x - 1)/c^2 + 1/8*(4*b^2*c*e*x*log(c*x + 1) + (b^2*c^2*e*x^2 + 2*b^2*c^2*d*x + (2*c*d - e)*b^2)*log(c*x + 1)^2 + (b^2*c^2*e*x^2 + 2*b^2*c^2*d*x - (2*c*d + e)*b^2)*log(-c*x + 1)^2 - 2*(2*b^2*c*e*x + (b^2*c^2*e*x^2 + 2*b^2*c^2*d*x + (2*c*d - e)*b^2)*log(c*x + 1))*log(-c*x + 1))/c^2`

Giac [F]

$$\int (d + ex)(a + b \operatorname{arctanh}(cx))^2 dx = \int (ex + d)(b \operatorname{arctanh}(cx) + a)^2 dx$$

input `integrate((e*x+d)*(a+b*arctanh(c*x))^2,x, algorithm="giac")`

output `integrate((e*x + d)*(b*arctanh(c*x) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int (d + ex)(a + b \operatorname{arctanh}(cx))^2 dx = \int (a + b \operatorname{atanh}(cx))^2 (d + ex) dx$$

input `int((a + b*atanh(c*x))^2*(d + e*x), x)`

output `int((a + b*atanh(c*x))^2*(d + e*x), x)`

Reduce [F]

$$\int (d + ex)(a + b \operatorname{arctanh}(cx))^2 dx$$

$$= \frac{2 \operatorname{atanh}(cx)^2 b^2 c^2 dx + \operatorname{atanh}(cx)^2 b^2 c^2 e x^2 - \operatorname{atanh}(cx)^2 b^2 e + 4 \operatorname{atanh}(cx) ab c^2 dx + 2 \operatorname{atanh}(cx) ab c^2 e x^2}{}$$

input `int((e*x+d)*(a+b*atanh(c*x))^2,x)`

output `(2*atanh(c*x)**2*b**2*c**2*d*x + atanh(c*x)**2*b**2*c**2*e*x**2 - atanh(c*x)**2*b**2*e + 4*atanh(c*x)*a*b*c**2*d*x + 2*atanh(c*x)*a*b*c**2*e*x**2 + 4*atanh(c*x)*a*b*c*d - 2*atanh(c*x)*a*b*e + 2*atanh(c*x)*b**2*c*e*x + 2*atanh(c*x)*b**2*e + 4*int((atanh(c*x)*x)/(c**2*x**2 - 1),x)*b**2*c**3*d + 4*log(c**2*x - c)*a*b*c*d + 2*log(c**2*x - c)*b**2*e + 2*a**2*c**2*d*x + a**2*c**2*e*x**2 + 2*a*b*c*e*x)/(2*c**2)`

3.12 $\int \frac{(a+b\operatorname{arctanh}(cx))^2}{d+ex} dx$

Optimal result	132
Mathematica [C] (warning: unable to verify)	133
Rubi [A] (verified)	134
Maple [C] (warning: unable to verify)	135
Fricas [F]	136
Sympy [F]	137
Maxima [F]	137
Giac [F]	137
Mupad [F(-1)]	138
Reduce [F]	138

Optimal result

Integrand size = 18, antiderivative size = 188

$$\int \frac{(a + b\operatorname{arctanh}(cx))^2}{d + ex} dx = -\frac{(a + b\operatorname{arctanh}(cx))^2 \log\left(\frac{2}{1+cx}\right)}{e} + \frac{(a + b\operatorname{arctanh}(cx))^2 \log\left(\frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{e} + \frac{b(a + b\operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+cx}\right)}{e} - \frac{b(a + b\operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{e} + \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1+cx}\right)}{2e} - \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{2e}$$

output

```
-(a+b*arctanh(c*x))^2*ln(2/(c*x+1))/e+(a+b*arctanh(c*x))^2*ln(2*c*(e*x+d)/(c*d+e)/(c*x+1))/e+b*(a+b*arctanh(c*x))*polylog(2,1-2/(c*x+1))/e-b*(a+b*arctanh(c*x))*polylog(2,1-2*c*(e*x+d)/(c*d+e)/(c*x+1))/e+1/2*b^2*polylog(3,1-2/(c*x+1))/e-1/2*b^2*polylog(3,1-2*c*(e*x+d)/(c*d+e)/(c*x+1))/e
```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 10.04 (sec) , antiderivative size = 1055, normalized size of antiderivative = 5.61

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{d + ex} dx = \text{Too large to display}$$

input `Integrate[(a + b*ArcTanh[c*x])^2/(d + e*x),x]`

output

```
(6*a^2*Log[d + e*x] + 6*a*b*ArcTanh[c*x]*(Log[1 - c^2*x^2] + 2*Log[I*Sinh[
ArcTanh[(c*d)/e] + ArcTanh[c*x]]]) - (6*I)*a*b*((-1/4*I)*(Pi - (2*I)*ArcTan
h[c*x])^2 + I*(ArcTanh[(c*d)/e] + ArcTanh[c*x])^2 + (Pi - (2*I)*ArcTanh[c
*x])*Log[1 + E^(2*ArcTanh[c*x])] + (2*I)*(ArcTanh[(c*d)/e] + ArcTanh[c*x])
*Log[1 - E^(-2*(ArcTanh[(c*d)/e] + ArcTanh[c*x]))] - (Pi - (2*I)*ArcTanh[c
*x])*Log[2/Sqrt[1 - c^2*x^2]] - (2*I)*(ArcTanh[(c*d)/e] + ArcTanh[c*x])*Lo
g[(2*I)*Sinh[ArcTanh[(c*d)/e] + ArcTanh[c*x]]] - I*PolyLog[2, -E^(2*ArcTan
h[c*x])] - I*PolyLog[2, E^(-2*(ArcTanh[(c*d)/e] + ArcTanh[c*x]))] + (b^2*
(-8*c*d*ArcTanh[c*x]^3 + 4*e*ArcTanh[c*x]^3 - (4*Sqrt[1 - (c^2*d^2)/e^2]*e
*ArcTanh[c*x]^3)/E^ArcTanh[(c*d)/e] - 6*c*d*ArcTanh[c*x]^2*Log[1 + E^(-2*A
rcTanh[c*x])] - (6*I)*c*d*Pi*ArcTanh[c*x]*Log[(E^(-ArcTanh[c*x]) + E^ArcTan
h[c*x])/2] - 6*c*d*ArcTanh[c*x]^2*Log[1 - (Sqrt[c*d + e]*E^ArcTanh[c*x])/
Sqrt[-(c*d) + e]] - 6*c*d*ArcTanh[c*x]^2*Log[1 + (Sqrt[c*d + e]*E^ArcTanh[
c*x])/Sqrt[-(c*d) + e]] + 6*c*d*ArcTanh[c*x]^2*Log[1 - E^(ArcTanh[(c*d)/e]
+ ArcTanh[c*x])] + 6*c*d*ArcTanh[c*x]^2*Log[1 + E^(ArcTanh[(c*d)/e] + Arc
Tanh[c*x])] + 6*c*d*ArcTanh[c*x]^2*Log[1 - E^(2*(ArcTanh[(c*d)/e] + ArcTan
h[c*x]))] + 12*c*d*ArcTanh[(c*d)/e]*ArcTanh[c*x]*Log[(I/2)*E^(-ArcTanh[(c*
d)/e] - ArcTanh[c*x])*(-1 + E^(2*(ArcTanh[(c*d)/e] + ArcTanh[c*x]))] + 6*
c*d*ArcTanh[c*x]^2*Log[(e*(-1 + E^(2*ArcTanh[c*x])) + c*d*(1 + E^(2*ArcTan
h[c*x]))]/(2*E^ArcTanh[c*x]))] - 6*c*d*ArcTanh[c*x]^2*Log[(c*(d + e*x))/...
```

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {6474}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{arctanh}(cx))^2}{d + ex} dx$$

↓ 6474

$$\begin{aligned} & - \frac{b(a + \operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2c(d+ex)}{(cd+e)(cx+1)}\right)}{e} + \\ & \frac{(a + \operatorname{arctanh}(cx))^2 \log\left(\frac{2c(d+ex)}{(cx+1)(cd+e)}\right)}{e} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{cx+1}\right) (a + \operatorname{arctanh}(cx))}{e} - \\ & \frac{\log\left(\frac{2}{cx+1}\right) (a + \operatorname{arctanh}(cx))^2}{e} - \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2c(d+ex)}{(cd+e)(cx+1)}\right)}{2e} + \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{cx+1}\right)}{2e} \end{aligned}$$

input `Int[(a + b*ArcTanh[c*x])^2/(d + e*x),x]`

output `-(((a + b*ArcTanh[c*x])^2*Log[2/(1 + c*x)])/e) + ((a + b*ArcTanh[c*x])^2*Log[(2*c*(d + e*x))/((c*d + e)*(1 + c*x))])/e + (b*(a + b*ArcTanh[c*x])*PolyLog[2, 1 - 2/(1 + c*x)])/e - (b*(a + b*ArcTanh[c*x])*PolyLog[2, 1 - (2*c*(d + e*x))/((c*d + e)*(1 + c*x))])/e + (b^2*PolyLog[3, 1 - 2/(1 + c*x)])/(2*e) - (b^2*PolyLog[3, 1 - (2*c*(d + e*x))/((c*d + e)*(1 + c*x))])/(2*e)`

Definitions of rubi rules used

rule 6474

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^2/((d_) + (e_.)*(x_.)), x_Symbol] :>
Simp[(-(a + b*ArcTanh[c*x])^2)*(Log[2/(1 + c*x)]/e), x] + (Simp[(a + b*ArcTanh[c*x])^2*(Log[2*c*((d + e*x)/((c*d + e)*(1 + c*x))])/e), x] + Simp[b*(a + b*ArcTanh[c*x])*(PolyLog[2, 1 - 2/(1 + c*x)]/e), x] - Simp[b*(a + b*ArcTanh[c*x])*(PolyLog[2, 1 - 2*c*((d + e*x)/((c*d + e)*(1 + c*x))])/e), x] + Simp[b^2*(PolyLog[3, 1 - 2/(1 + c*x)]/(2*e)), x] - Simp[b^2*(PolyLog[3, 1 - 2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/(2*e)), x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 - e^2, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.88 (sec) , antiderivative size = 1087, normalized size of antiderivative = 5.78

method	result	size
derivatividivides	Expression too large to display	1087
default	Expression too large to display	1087
parts	Expression too large to display	1090

input

```
int((a+b*arctanh(c*x))^2/(e*x+d),x,method=_RETURNVERBOSE)
```


output

```

1/c*(a^2*c*ln(c*e*x+c*d)/e+b^2*c*(ln(c*e*x+c*d)/e*arctanh(c*x)^2-2/e*(1/2*
arctanh(c*x)^2*ln(d*c*(1+(c*x+1)^2/(-c^2*x^2+1))+e*((c*x+1)^2/(-c^2*x^2+1)
-1))-1/4*I*Pi*csgn(I*(d*c*(1-(c*x+1)^2/(c^2*x^2-1))+e*(-(c*x+1)^2/(c^2*x^2
-1)-1))/(1-(c*x+1)^2/(c^2*x^2-1)))*(csgn(I*(d*c*(1-(c*x+1)^2/(c^2*x^2-1))+
e*(-(c*x+1)^2/(c^2*x^2-1)-1)))*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))-csgn(I*(d
*c*(1-(c*x+1)^2/(c^2*x^2-1))+e*(-(c*x+1)^2/(c^2*x^2-1)-1))/(1-(c*x+1)^2/(c
^2*x^2-1)))*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))-csgn(I*(d*c*(1-(c*x+1)^2/(c^
2*x^2-1))+e*(-(c*x+1)^2/(c^2*x^2-1)-1)))*csgn(I*(d*c*(1-(c*x+1)^2/(c^2*x^2
-1))+e*(-(c*x+1)^2/(c^2*x^2-1)-1))/(1-(c*x+1)^2/(c^2*x^2-1)))+csgn(I*(d*c*
(1-(c*x+1)^2/(c^2*x^2-1))+e*(-(c*x+1)^2/(c^2*x^2-1)-1))/(1-(c*x+1)^2/(c^2*
x^2-1)))^2)*arctanh(c*x)^2+1/2*arctanh(c*x)*polylog(2,-(c*x+1)^2/(-c^2*x^2
+1))-1/4*polylog(3,-(c*x+1)^2/(-c^2*x^2+1))-1/2/(c*d+e)*e*arctanh(c*x)^2*ln
n(1-(c*d+e)*(c*x+1)^2/(-c^2*x^2+1)/(-c*d+e))-1/2/(c*d+e)*e*arctanh(c*x)*po
lylog(2,(c*d+e)*(c*x+1)^2/(-c^2*x^2+1)/(-c*d+e))+1/4/(c*d+e)*e*polylog(3,(
c*d+e)*(c*x+1)^2/(-c^2*x^2+1)/(-c*d+e))-1/2/(c*d+e)*d*c*arctanh(c*x)^2*ln(
1-(c*d+e)*(c*x+1)^2/(-c^2*x^2+1)/(-c*d+e))-1/2/(c*d+e)*d*c*arctanh(c*x)*po
lylog(2,(c*d+e)*(c*x+1)^2/(-c^2*x^2+1)/(-c*d+e))+1/4/(c*d+e)*d*c*polylog(3
,(c*d+e)*(c*x+1)^2/(-c^2*x^2+1)/(-c*d+e))))+2*a*b*c*(ln(c*e*x+c*d)/e*arcta
nh(c*x)-1/e^2*(-1/2*e*(dilog((c*e*x-e)/(-c*d-e))+ln(c*e*x+c*d)*ln((c*e*x-e
)/(-c*d-e)))+1/2*e*(dilog((c*e*x+e)/(-c*d-e))+ln(c*e*x+c*d)*ln((c*e*x+e...

```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{d + ex} dx = \int \frac{(b \operatorname{arctanh}(cx) + a)^2}{ex + d} dx$$

input

```
integrate((a+b*arctanh(c*x))^2/(e*x+d),x, algorithm="fricas")
```

output

```
integral((b^2*arctanh(c*x)^2 + 2*a*b*arctanh(c*x) + a^2)/(e*x + d), x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{d + ex} dx = \int \frac{(a + b \operatorname{atanh}(cx))^2}{d + ex} dx$$

input `integrate((a+b*atanh(c*x))**2/(e*x+d), x)`

output `Integral((a + b*atanh(c*x))**2/(d + e*x), x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{d + ex} dx = \int \frac{(b \operatorname{atanh}(cx) + a)^2}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x))^2/(e*x+d), x, algorithm="maxima")`

output `a^2*log(e*x + d)/e + integrate(1/4*b^2*(log(c*x + 1) - log(-c*x + 1))^2/(e*x + d) + a*b*(log(c*x + 1) - log(-c*x + 1))/(e*x + d), x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{d + ex} dx = \int \frac{(b \operatorname{atanh}(cx) + a)^2}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x))^2/(e*x+d), x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^2/(e*x + d), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{d + ex} dx = \int \frac{(a + b \operatorname{atanh}(cx))^2}{d + ex} dx$$

input `int((a + b*atanh(c*x))^2/(d + e*x),x)`output `int((a + b*atanh(c*x))^2/(d + e*x), x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{d + ex} dx = \frac{2 \left(\int \frac{\operatorname{atanh}(cx)}{ex+d} dx \right) a b e + \left(\int \frac{\operatorname{atanh}(cx)^2}{ex+d} dx \right) b^2 e + \log(ex + d) a^2}{e}$$

input `int((a+b*atanh(c*x))^2/(e*x+d),x)`output `(2*int(atanh(c*x)/(d + e*x),x)*a*b*e + int(atanh(c*x)**2/(d + e*x),x)*b**2 *e + log(d + e*x)*a**2)/e`

3.13 $\int \frac{(a+b\operatorname{arctanh}(cx))^2}{(d+ex)^2} dx$

Optimal result	139
Mathematica [C] (warning: unable to verify)	140
Rubi [A] (verified)	141
Maple [A] (verified)	142
Fricas [F]	143
Sympy [F]	143
Maxima [F]	144
Giac [F]	144
Mupad [F(-1)]	144
Reduce [F]	145

Optimal result

Integrand size = 18, antiderivative size = 321

$$\begin{aligned}
 \int \frac{(a + b\operatorname{arctanh}(cx))^2}{(d + ex)^2} dx = & -\frac{(a + b\operatorname{arctanh}(cx))^2}{e(d + ex)} + \frac{bc(a + b\operatorname{arctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{e(cd + e)} \\
 & - \frac{bc(a + b\operatorname{arctanh}(cx)) \log\left(\frac{2}{1+cx}\right)}{(cd - e)e} \\
 & + \frac{2bc(a + b\operatorname{arctanh}(cx)) \log\left(\frac{2}{1+cx}\right)}{c^2d^2 - e^2} \\
 & - \frac{2bc(a + b\operatorname{arctanh}(cx)) \log\left(\frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{c^2d^2 - e^2} \\
 & + \frac{b^2c \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{2e(cd + e)} \\
 & + \frac{b^2c \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+cx}\right)}{2(cd - e)e} - \frac{b^2c \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+cx}\right)}{c^2d^2 - e^2} \\
 & + \frac{b^2c \operatorname{PolyLog}\left(2, 1 - \frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{c^2d^2 - e^2}
 \end{aligned}$$

output

```

-(a+b*arctanh(c*x))^2/e/(e*x+d)+b*c*(a+b*arctanh(c*x))*ln(2/(-c*x+1))/e/(c
*d+e)-b*c*(a+b*arctanh(c*x))*ln(2/(c*x+1))/(c*d-e)/e+2*b*c*(a+b*arctanh(c*
x))*ln(2/(c*x+1))/(c^2*d^2-e^2)-2*b*c*(a+b*arctanh(c*x))*ln(2*c*(e*x+d)/(c
*d+e)/(c*x+1))/(c^2*d^2-e^2)+1/2*b^2*c*polylog(2,1-2/(-c*x+1))/e/(c*d+e)+1
/2*b^2*c*polylog(2,1-2/(c*x+1))/(c*d-e)/e-b^2*c*polylog(2,1-2/(c*x+1))/(c^
2*d^2-e^2)+b^2*c*polylog(2,1-2*c*(e*x+d)/(c*d+e)/(c*x+1))/(c^2*d^2-e^2)

```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 2.97 (sec) , antiderivative size = 317, normalized size of antiderivative = 0.99

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^2} dx$$

$$= -\frac{a^2}{e(d + ex)} + \frac{abc \left(-\frac{2 \operatorname{arctanh}(cx)}{cd + cex} + \frac{(-cd + e) \log(1 - cx) + (cd + e) \log(1 + cx) - 2e \log(c(d + ex))}{(cd - e)(cd + e)} \right)}{e}$$

$$+ \frac{b^2 \left(-\frac{e^{-\operatorname{arctanh}\left(\frac{cd}{e}\right)} \operatorname{arctanh}(cx)^2}{\sqrt{1 - \frac{c^2 d^2}{e^2} e}} + \frac{x \operatorname{arctanh}(cx)^2}{d + ex} + \frac{cd \left(i\pi \log\left(1 + e^{2 \operatorname{arctanh}(cx)}\right) - 2 \operatorname{arctanh}(cx) \log\left(1 - e^{-2 \operatorname{arctanh}\left(\frac{cd}{e}\right)}\right) \right)}{d + ex} \right)}{e}$$

input

```
Integrate[(a + b*ArcTanh[c*x])^2/(d + e*x)^2,x]
```

output

```

-(a^2/(e*(d + e*x))) + (a*b*c*((-2*ArcTanh[c*x])/(c*d + c*e*x) + ((-(c*d)
+ e)*Log[1 - c*x] + (c*d + e)*Log[1 + c*x] - 2*e*Log[c*(d + e*x)])/((c*d -
e)*(c*d + e))))/e + (b^2*(-(ArcTanh[c*x]^2/(Sqrt[1 - (c^2*d^2)/e^2]*e*E^A
rcTanh[(c*d)/e])) + (x*ArcTanh[c*x]^2)/(d + e*x) + (c*d*(I*Pi*Log[1 + E^(2
*ArcTanh[c*x])] - 2*ArcTanh[c*x]*Log[1 - E^(-2*(ArcTanh[(c*d)/e] + ArcTanh
[c*x])]) - I*Pi*(ArcTanh[c*x] - Log[1 - c^2*x^2]/2) - 2*ArcTanh[(c*d)/e]*(
ArcTanh[c*x] + Log[1 - E^(-2*(ArcTanh[(c*d)/e] + ArcTanh[c*x])]) - Log[I*S
inh[ArcTanh[(c*d)/e] + ArcTanh[c*x]]]) + PolyLog[2, E^(-2*(ArcTanh[(c*d)/e
] + ArcTanh[c*x])])))))/(c^2*d^2 - e^2))/d

```

Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 310, normalized size of antiderivative = 0.97, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + \operatorname{arctanh}(cx))^2}{(d + ex)^2} dx \\
 & \quad \downarrow \text{6480} \\
 & \frac{2bc \int \left(-\frac{(a + \operatorname{arctanh}(cx))e^2}{(cd - e)(cd + e)(d + ex)} + \frac{c(a + \operatorname{arctanh}(cx))}{2(cd + e)(1 - cx)} + \frac{c(a + \operatorname{arctanh}(cx))}{2(cd - e)(cx + 1)} \right) dx}{e} - \frac{(a + \operatorname{arctanh}(cx))^2}{e(d + ex)} \\
 & \quad \downarrow \text{2009} \\
 & \frac{2bc \left(\frac{e \log\left(\frac{2}{cx + 1}\right)(a + \operatorname{arctanh}(cx))}{c^2 d^2 - e^2} - \frac{e(a + \operatorname{arctanh}(cx)) \log\left(\frac{2c(d + ex)}{(cx + 1)(cd + e)}\right)}{c^2 d^2 - e^2} + \frac{\log\left(\frac{2}{1 - cx}\right)(a + \operatorname{arctanh}(cx))}{2(cd + e)} - \frac{\log\left(\frac{2}{cx + 1}\right)(a + \operatorname{arctanh}(cx))}{2(cd - e)} \right)}{e} \\
 & \quad \frac{(a + \operatorname{arctanh}(cx))^2}{e(d + ex)}
 \end{aligned}$$

input

```
Int[(a + b*ArcTanh[c*x])^2/(d + e*x)^2, x]
```

output

```

-((a + b*ArcTanh[c*x])^2/(e*(d + e*x))) + (2*b*c*((a + b*ArcTanh[c*x])*Log[2/(1 - c*x)])/(2*(c*d + e)) - ((a + b*ArcTanh[c*x])*Log[2/(1 + c*x)])/(2*(c*d - e)) + (e*(a + b*ArcTanh[c*x])*Log[2/(1 + c*x)])/(c^2*d^2 - e^2) - (e*(a + b*ArcTanh[c*x])*Log[(2*c*(d + e*x))/((c*d + e)*(1 + c*x))])/(c^2*d^2 - e^2) + (b*PolyLog[2, 1 - 2/(1 - c*x)]/(4*(c*d + e)) + (b*PolyLog[2, 1 - 2/(1 + c*x)]/(4*(c*d - e)) - (b*e*PolyLog[2, 1 - 2/(1 + c*x)]/(2*(c^2*d^2 - e^2)) + (b*e*PolyLog[2, 1 - (2*c*(d + e*x))/((c*d + e)*(1 + c*x))])/(2*(c^2*d^2 - e^2))))/e

```

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6480 Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)*((d_.) + (e_.)*(x_.))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 451, normalized size of antiderivative = 1.40

method	result
derivativedivides	$-\frac{a^2 c^2}{(cex+cd)e} + b^2 c^2 \left(-\frac{\operatorname{arctanh}(cx)^2}{(cex+cd)e} + \frac{-2 \operatorname{arctanh}(cx) e \ln(cex+cd) - 2 \operatorname{arctanh}(cx) \ln(cx-1) + 2 \operatorname{arctanh}(cx) \ln(cx+1)}{(cd+e)(cd-e)} - \frac{2 \left(\frac{e(dil)}{2} \right)}{2cd-2e} \right)$
default	$-\frac{a^2 c^2}{(cex+cd)e} + b^2 c^2 \left(-\frac{\operatorname{arctanh}(cx)^2}{(cex+cd)e} + \frac{-2 \operatorname{arctanh}(cx) e \ln(cex+cd) - 2 \operatorname{arctanh}(cx) \ln(cx-1) + 2 \operatorname{arctanh}(cx) \ln(cx+1)}{(cd+e)(cd-e)} - \frac{2 \left(\frac{e(dil)}{2} \right)}{2cd-2e} \right)$
parts	$-\frac{a^2}{(ex+d)e} + b^2 \left(-\frac{c^2 \operatorname{arctanh}(cx)^2}{(cex+cd)e} + \frac{2c^2 \left(-\frac{\operatorname{arctanh}(cx) e \ln(cex+cd) - \operatorname{arctanh}(cx) \ln(cx-1) + \operatorname{arctanh}(cx) \ln(cx+1)}{(cd+e)(cd-e)} - \frac{e(dil)}{2cd-2e} \right)}{2cd-2e} \right)$

```
input int((a+b*arctanh(c*x))^2/(e*x+d)^2,x,method=_RETURNVERBOSE)
```

output

```
1/c*(-a^2*c^2/(c*e*x+c*d)/e+b^2*c^2*(-1/(c*e*x+c*d)/e*arctanh(c*x)^2+2/e*(-arctanh(c*x)*e/(c*d+e)/(c*d-e)*ln(c*e*x+c*d)-arctanh(c*x)/(2*c*d+2*e)*ln(c*x-1)+arctanh(c*x)/(2*c*d-2*e)*ln(c*x+1)-1/(c*d+e)/(c*d-e)*(1/2*e*(dilog((c*e*x-e)/(-c*d-e))+ln(c*e*x+c*d)*ln((c*e*x-e)/(-c*d-e)))-1/2*e*(dilog((c*e*x+e)/(-c*d+e))+ln(c*e*x+c*d)*ln((c*e*x+e)/(-c*d+e))))+1/2/(c*d-e)*(-1/4*ln(c*x+1)^2+1/2*(ln(c*x+1)-ln(1/2*c*x+1/2))*ln(-1/2*c*x+1/2)-1/2*dilog(1/2*c*x+1/2))-1/2/(c*d+e)*(1/4*ln(c*x-1)^2-1/2*dilog(1/2*c*x+1/2)-1/2*ln(c*x-1)*ln(1/2*c*x+1/2))))+2*a*b*c^2*(-1/(c*e*x+c*d)/e*arctanh(c*x)+1/e*(-e/(c*d+e)/(c*d-e)*ln(c*e*x+c*d)-1/(2*c*d+2*e)*ln(c*x-1)+1/(2*c*d-2*e)*ln(c*x+1)))
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^2} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^2}{(ex + d)^2} dx$$

input

```
integrate((a+b*arctanh(c*x))^2/(e*x+d)^2,x, algorithm="fricas")
```

output

```
integral((b^2*arctanh(c*x)^2 + 2*a*b*arctanh(c*x) + a^2)/(e^2*x^2 + 2*d*e*x + d^2), x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^2} dx = \int \frac{(a + b \operatorname{atanh}(cx))^2}{(d + ex)^2} dx$$

input

```
integrate((a+b*atanh(c*x))**2/(e*x+d)**2,x)
```

output

```
Integral((a + b*atanh(c*x))**2/(d + e*x)**2, x)
```


Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^2} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^2}{(ex + d)^2} dx$$

input `integrate((a+b*arctanh(c*x))^2/(e*x+d)^2,x, algorithm="maxima")`

output `(c*(log(c*x + 1)/(c*d*e - e^2) - log(c*x - 1)/(c*d*e + e^2) - 2*log(e*x + d)/(c^2*d^2 - e^2)) - 2*arctanh(c*x)/(e^2*x + d*e))*a*b - 1/4*b^2*(log(-c*x + 1)^2/(e^2*x + d*e) + integrate(-((c*e*x - e)*log(c*x + 1)^2 + 2*(c*e*x + c*d - (c*e*x - e)*log(c*x + 1))*log(-c*x + 1))/(c*e^3*x^3 - d^2*e + (2*c*d*e^2 - e^3)*x^2 + (c*d^2*e - 2*d*e^2)*x), x)) - a^2/(e^2*x + d*e)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^2} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^2}{(ex + d)^2} dx$$

input `integrate((a+b*arctanh(c*x))^2/(e*x+d)^2,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^2/(e*x + d)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^2} dx = \int \frac{(a + b \operatorname{atanh}(cx))^2}{(d + ex)^2} dx$$

input `int((a + b*atanh(c*x))^2/(d + e*x)^2,x)`

output `int((a + b*atanh(c*x))^2/(d + e*x)^2, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^2} dx = \text{too large to display}$$

input `int((a+b*atanh(c*x))^2/(e*x+d)^2,x)`

output

```
(atanh(c*x)**2*b**2*c**4*d**4*x - atanh(c*x)**2*b**2*c**2*d**3*e - atanh(c
*x)**2*b**2*c**2*d**2*e**2*x + atanh(c*x)**2*b**2*d*e**3 + 2*atanh(c*x)*a*
b*c**4*d**4*x - 2*atanh(c*x)*a*b*e**4*x + 2*atanh(c*x)*b**2*c**3*d**3*e*x
- 2*atanh(c*x)*b**2*c*d*e**3*x + 2*int((atanh(c*x)*x)/(c**4*d**4*x**2 + 2*
c**4*d**3*e*x**3 + c**4*d**2*e**2*x**4 - c**2*d**4 - 2*c**2*d**3*e*x + 2*c
**2*d*e**3*x**3 + c**2*e**4*x**4 - d**2*e**2 - 2*d*e**3*x - e**4*x**2),x)*
b**2*c**7*d**8 + 2*int((atanh(c*x)*x)/(c**4*d**4*x**2 + 2*c**4*d**3*e*x**3
+ c**4*d**2*e**2*x**4 - c**2*d**4 - 2*c**2*d**3*e*x + 2*c**2*d*e**3*x**3
+ c**2*e**4*x**4 - d**2*e**2 - 2*d*e**3*x - e**4*x**2),x)*b**2*c**7*d**7*e
*x - 2*int((atanh(c*x)*x)/(c**4*d**4*x**2 + 2*c**4*d**3*e*x**3 + c**4*d**2
*e**2*x**4 - c**2*d**4 - 2*c**2*d**3*e*x + 2*c**2*d*e**3*x**3 + c**2*e**4*
x**4 - d**2*e**2 - 2*d*e**3*x - e**4*x**2),x)*b**2*c**5*d**6*e**2 - 2*int(
(atanh(c*x)*x)/(c**4*d**4*x**2 + 2*c**4*d**3*e*x**3 + c**4*d**2*e**2*x**4
- c**2*d**4 - 2*c**2*d**3*e*x + 2*c**2*d*e**3*x**3 + c**2*e**4*x**4 - d**2
*e**2 - 2*d*e**3*x - e**4*x**2),x)*b**2*c**5*d**5*e**3*x - 2*int((atanh(c
*x)*x)/(c**4*d**4*x**2 + 2*c**4*d**3*e*x**3 + c**4*d**2*e**2*x**4 - c**2*d*
**4 - 2*c**2*d**3*e*x + 2*c**2*d*e**3*x**3 + c**2*e**4*x**4 - d**2*e**2 - 2
*d*e**3*x - e**4*x**2),x)*b**2*c**3*d**4*e**4 - 2*int((atanh(c*x)*x)/(c**4
*d**4*x**2 + 2*c**4*d**3*e*x**3 + c**4*d**2*e**2*x**4 - c**2*d**4 - 2*c**2
*d**3*e*x + 2*c**2*d*e**3*x**3 + c**2*e**4*x**4 - d**2*e**2 - 2*d*e**3*...
```

3.14 $\int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^3} dx$

Optimal result	146
Mathematica [C] (warning: unable to verify)	147
Rubi [A] (verified)	148
Maple [A] (verified)	150
Fricas [F]	151
Sympy [F]	151
Maxima [F]	151
Giac [F]	152
Mupad [F(-1)]	152
Reduce [F]	153

Optimal result

Integrand size = 18, antiderivative size = 480

$$\begin{aligned}
 \int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^3} dx = & \frac{bc(a + b \operatorname{arctanh}(cx))}{(c^2 d^2 - e^2)(d + ex)} - \frac{(a + b \operatorname{arctanh}(cx))^2}{2e(d + ex)^2} \\
 & + \frac{bc^2(a + b \operatorname{arctanh}(cx)) \log\left(\frac{2}{1 - cx}\right)}{2e(cd + e)^2} \\
 & + \frac{b^2 c^2 \log(1 - cx)}{2(cd - e)(cd + e)^2} - \frac{bc^2(a + b \operatorname{arctanh}(cx)) \log\left(\frac{2}{1 + cx}\right)}{2(cd - e)^2 e} \\
 & + \frac{2bc^3 d(a + b \operatorname{arctanh}(cx)) \log\left(\frac{2}{1 + cx}\right)}{(cd - e)^2 (cd + e)^2} \\
 & - \frac{b^2 c^2 \log(1 + cx)}{2(cd - e)^2 (cd + e)} + \frac{b^2 c^2 e \log(d + ex)}{(cd - e)^2 (cd + e)^2} \\
 & - \frac{2bc^3 d(a + b \operatorname{arctanh}(cx)) \log\left(\frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{(cd - e)^2 (cd + e)^2} \\
 & + \frac{b^2 c^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx}\right)}{4e(cd + e)^2} + \frac{b^2 c^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 + cx}\right)}{4(cd - e)^2 e} \\
 & - \frac{b^2 c^3 d \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 + cx}\right)}{(cd - e)^2 (cd + e)^2} \\
 & + \frac{b^2 c^3 d \operatorname{PolyLog}\left(2, 1 - \frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{(cd - e)^2 (cd + e)^2}
 \end{aligned}$$

output

```

b*c*(a+b*arctanh(c*x))/(c^2*d^2-e^2)/(e*x+d)-1/2*(a+b*arctanh(c*x))^2/e/(e
*x+d)^2+1/2*b*c^2*(a+b*arctanh(c*x))*ln(2/(-c*x+1))/e/(c*d+e)^2+1/2*b^2*c^
2*ln(-c*x+1)/(c*d-e)/(c*d+e)^2-1/2*b*c^2*(a+b*arctanh(c*x))*ln(2/(c*x+1))/
(c*d-e)^2/e+2*b*c^3*d*(a+b*arctanh(c*x))*ln(2/(c*x+1))/(c*d-e)^2/(c*d+e)^2
-1/2*b^2*c^2*ln(c*x+1)/(c*d-e)^2/(c*d+e)+b^2*c^2*e*ln(e*x+d)/(c*d-e)^2/(c*
d+e)^2-2*b*c^3*d*(a+b*arctanh(c*x))*ln(2*c*(e*x+d)/(c*d+e)/(c*x+1))/(c*d-e
)^2/(c*d+e)^2+1/4*b^2*c^2*polylog(2,1-2/(-c*x+1))/e/(c*d+e)^2+1/4*b^2*c^2*
polylog(2,1-2/(c*x+1))/(c*d-e)^2/e-b^2*c^3*d*polylog(2,1-2/(c*x+1))/(c*d-e
)^2/(c*d+e)^2+b^2*c^3*d*polylog(2,1-2*c*(e*x+d)/(c*d+e)/(c*x+1))/(c*d-e)^2
/(c*d+e)^2

```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 4.51 (sec) , antiderivative size = 470, normalized size of antiderivative = 0.98

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^3} dx = -\frac{a^2}{2e(d + ex)^2}$$

$$-\frac{abc^2 \left(\frac{2 \operatorname{arctanh}(cx)}{(cd+ce)^2} + \frac{\log(1-cx)}{(cd+e)^2} + \frac{-\log(1+cx) + \frac{2e(-c^2d^2+e^2+2c^2d(d+ex)\log(c(d+ex)))}{c(cd+e)^2(d+ex)}}{(-cd+e)^2} \right)}{2e}$$

$$+\frac{b^2c^2 \left(-\frac{2e^{-\operatorname{arctanh}(\frac{cd}{e})} \operatorname{arctanh}(cx)^2}{\sqrt{1-\frac{c^2d^2}{e^2}}e} - \frac{e(-1+c^2x^2) \operatorname{arctanh}(cx)^2}{c^2(d+ex)^2} + \frac{2x \operatorname{arctanh}(cx)(-e+cd \operatorname{arctanh}(cx))}{cd(d+ex)} + \frac{2e(-\operatorname{arctanh}(\frac{cd}{e}))}{cd(d+ex)} \right)}{cd(d+ex)}$$

input

```
Integrate[(a + b*ArcTanh[c*x])^2/(d + e*x)^3,x]
```

output

```
-1/2*a^2/(e*(d + e*x)^2) - (a*b*c^2*((2*ArcTanh[c*x])/(c*d + c*e*x)^2 + Log[1 - c*x]/(c*d + e)^2 + (-Log[1 + c*x] + (2*e*(-(c^2*d^2) + e^2 + 2*c^2*d*(d + e*x)*Log[c*(d + e*x)])))/(c*(c*d + e)^2*(d + e*x)))/((-c*d) + e)^2)/(2*e) + (b^2*c^2*(-2*ArcTanh[c*x]^2)/(Sqrt[1 - (c^2*d^2)/e^2]*e*E^ArcTanh[(c*d)/e]) - (e*(-1 + c^2*x^2)*ArcTanh[c*x]^2)/(c^2*(d + e*x)^2) + (2*x*ArcTanh[c*x]*(-e + c*d*ArcTanh[c*x]))/(c*d*(d + e*x)) + (2*e*(-e*ArcTanh[c*x]) + c*d*Log[(c*(d + e*x))/Sqrt[1 - c^2*x^2]]))/(c^3*d^3 - c*d*e^2) + (2*c*d*(I*Pi*Log[1 + E^(2*ArcTanh[c*x])] - 2*ArcTanh[c*x]*Log[1 - E^(-2*(ArcTanh[(c*d)/e] + ArcTanh[c*x]))] - I*Pi*(ArcTanh[c*x] - Log[1 - c^2*x^2]/2) - 2*ArcTanh[(c*d)/e]*(ArcTanh[c*x] + Log[1 - E^(-2*(ArcTanh[(c*d)/e] + ArcTanh[c*x]))] - Log[I*Sinh[ArcTanh[(c*d)/e] + ArcTanh[c*x]]]) + PolyLog[2, E^(-2*(ArcTanh[(c*d)/e] + ArcTanh[c*x]))]))/(c^2*d^2 - e^2))/(2*(c*d - e)*(c*d + e))
```

Rubi [A] (verified)

Time = 0.87 (sec) , antiderivative size = 450, normalized size of antiderivative = 0.94, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{barctanh}(cx))^2}{(d + ex)^3} dx$$

↓ 6480

$$\frac{bc \int \left(\frac{(a + \operatorname{barctanh}(cx))^2 c^2}{2(cd+e)^2(1-cx)} + \frac{(a + \operatorname{barctanh}(cx))^2 c^2}{2(cd-e)^2(cx+1)} - \frac{2de^2(a + \operatorname{barctanh}(cx))^2 c^2}{(cd-e)^2(cd+e)^2(d+ex)} - \frac{e^2(a + \operatorname{barctanh}(cx))^2}{(cd-e)(cd+e)(d+ex)^2} \right) dx}{(a + \operatorname{barctanh}(cx))^2}{2e(d + ex)^2}$$

↓ 2009

$$\frac{bc \left(\frac{e(a + \operatorname{barctanh}(cx))}{(c^2 d^2 - e^2)(d + ex)} + \frac{2c^2 de \log\left(\frac{2}{cx+1}\right)(a + \operatorname{barctanh}(cx))}{(cd-e)^2(cd+e)^2} - \frac{2c^2 de(a + \operatorname{barctanh}(cx)) \log\left(\frac{2c(d+ex)}{(cx+1)(cd+e)}\right)}{(cd-e)^2(cd+e)^2} + \frac{c \log\left(\frac{2}{1-cx}\right)(a + \operatorname{barctanh}(cx))}{2(cd+e)^2} \right)}{(a + \operatorname{barctanh}(cx))^2}{2e(d + ex)^2}$$

input `Int[(a + b*ArcTanh[c*x])^2/(d + e*x)^3,x]`

output `-1/2*(a + b*ArcTanh[c*x])^2/(e*(d + e*x)^2) + (b*c*((e*(a + b*ArcTanh[c*x]))/((c^2*d^2 - e^2)*(d + e*x)) + (c*(a + b*ArcTanh[c*x])*Log[2/(1 - c*x)])/(2*(c*d + e)^2) + (b*c*e*Log[1 - c*x])/(2*(c*d - e)*(c*d + e)^2) - (c*(a + b*ArcTanh[c*x])*Log[2/(1 + c*x)])/(2*(c*d - e)^2) + (2*c^2*d*e*(a + b*ArcTanh[c*x])*Log[2/(1 + c*x)])/((c*d - e)^2*(c*d + e)^2) - (b*c*e*Log[1 + c*x])/(2*(c*d - e)^2*(c*d + e)) + (b*c*e^2*Log[d + e*x])/((c*d - e)^2*(c*d + e)^2) - (2*c^2*d*e*(a + b*ArcTanh[c*x])*Log[(2*c*(d + e*x))/((c*d + e)*(1 + c*x))])/((c*d - e)^2*(c*d + e)^2) + (b*c*PolyLog[2, 1 - 2/(1 - c*x)])/(4*(c*d + e)^2) + (b*c*PolyLog[2, 1 - 2/(1 + c*x)])/(4*(c*d - e)^2) - (b*c^2*d*e*PolyLog[2, 1 - 2/(1 + c*x)])/((c*d - e)^2*(c*d + e)^2) + (b*c^2*d*e*PolyLog[2, 1 - (2*c*(d + e*x))/(c*d + e)])/((c*d - e)^2*(c*d + e)^2))/e`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6480 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^p_*((d_) + (e_.)*(x_))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]`

Maple [A] (verified)

Time = 0.75 (sec) , antiderivative size = 592, normalized size of antiderivative = 1.23

method	result
derivativedivides	$-\frac{a^2 e^3}{2(cex+cd)^2 e} + b^2 c^3 \left(-\frac{\operatorname{arctanh}(cx)^2}{2(cex+cd)^2 e} + \frac{\operatorname{arctanh}(cx)e}{(cd+e)(cd-e)(cex+cd)} - \frac{2 \operatorname{arctanh}(cx)edc \ln(cex+cd)}{(cd+e)^2 (cd-e)^2} - \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{2(cd+e)^2} + \frac{\operatorname{arctanh}(cx)}{2(cd+e)} \right)$
default	$-\frac{a^2 e^3}{2(cex+cd)^2 e} + b^2 c^3 \left(-\frac{\operatorname{arctanh}(cx)^2}{2(cex+cd)^2 e} + \frac{\operatorname{arctanh}(cx)e}{(cd+e)(cd-e)(cex+cd)} - \frac{2 \operatorname{arctanh}(cx)edc \ln(cex+cd)}{(cd+e)^2 (cd-e)^2} - \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{2(cd+e)^2} + \frac{\operatorname{arctanh}(cx)}{2(cd+e)} \right)$
parts	$-\frac{a^2}{2(ex+d)^2 e} + b^2 \left(-\frac{c^3 \operatorname{arctanh}(cx)^2}{2(cex+cd)^2 e} + c^3 \left(\frac{\operatorname{arctanh}(cx)e}{(cd+e)(cd-e)(cex+cd)} - \frac{2 \operatorname{arctanh}(cx)edc \ln(cex+cd)}{(cd+e)^2 (cd-e)^2} - \frac{\operatorname{arctanh}(cx) \ln(cx-1)}{2(cd+e)^2} + \frac{\operatorname{arctanh}(cx)}{2(cd+e)} \right) \right)$

input `int((a+b*arctanh(c*x))^2/(e*x+d)^3,x,method=_RETURNVERBOSE)`

output `1/c*(-1/2*a^2*c^3/(c*e*x+c*d)^2/e+b^2*c^3*(-1/2/(c*e*x+c*d)^2/e*arctanh(c*x)^2+1/e*(arctanh(c*x)*e/(c*d+e)/(c*d-e)/(c*e*x+c*d)-2*arctanh(c*x)*e*d*c/(c*d+e)^2/(c*d-e)^2*ln(c*e*x+c*d)-1/2*arctanh(c*x)/(c*d+e)^2*ln(c*x-1)+1/2*arctanh(c*x)/(c*d-e)^2*ln(c*x+1)-1/2/(c*d+e)^2*(1/4*ln(c*x-1)^2-1/2*dilog(1/2*c*x+1/2)-1/2*ln(c*x-1)*ln(1/2*c*x+1/2))+1/2/(c*d-e)^2*(-1/4*ln(c*x+1)^2+1/2*(ln(c*x+1)-ln(1/2*c*x+1/2))*ln(-1/2*c*x+1/2)-1/2*dilog(1/2*c*x+1/2))+e/(c*d+e)/(c*d-e)*(e/(c*d+e)/(c*d-e)*ln(c*e*x+c*d)+1/(2*c*d+2*e)*ln(c*x-1)-1/(2*c*d-2*e)*ln(c*x+1))-2*d*c/(c*d+e)^2/(c*d-e)^2*(1/2*e*(dilog((c*e*x-e)/(-c*d-e))+ln(c*e*x+c*d)*ln((c*e*x-e)/(-c*d-e)))-1/2*e*(dilog((c*e*x+e)/(-c*d+e))+ln(c*e*x+c*d)*ln((c*e*x+e)/(-c*d+e)))))+2*a*b*c^3*(-1/2/(c*e*x+c*d)^2/e*arctanh(c*x)+1/2/e*(e/(c*d+e)/(c*d-e)/(c*e*x+c*d)-2*e*d*c/(c*d+e)^2/(c*d-e)^2*ln(c*e*x+c*d)-1/2/(c*d+e)^2*ln(c*x-1)+1/2/(c*d-e)^2*ln(c*x+1))))`

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^3} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^2}{(ex + d)^3} dx$$

input `integrate((a+b*arctanh(c*x))^2/(e*x+d)^3,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x)^2 + 2*a*b*arctanh(c*x) + a^2)/(e^3*x^3 + 3*d*e^2*x^2 + 3*d^2*e*x + d^3), x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^3} dx = \int \frac{(a + b \operatorname{atanh}(cx))^2}{(d + ex)^3} dx$$

input `integrate((a+b*atanh(c*x))**2/(e*x+d)**3,x)`

output `Integral((a + b*atanh(c*x))**2/(d + e*x)**3, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^3} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^2}{(ex + d)^3} dx$$

input `integrate((a+b*arctanh(c*x))^2/(e*x+d)^3,x, algorithm="maxima")`

output

```
-1/2*((4*c^2*d*log(e*x + d)/(c^4*d^4 - 2*c^2*d^2*e^2 + e^4) - c*log(c*x +
1)/(c^2*d^2*e - 2*c*d*e^2 + e^3) + c*log(c*x - 1)/(c^2*d^2*e + 2*c*d*e^2 +
e^3) - 2/(c^2*d^3 - d*e^2 + (c^2*d^2*e - e^3)*x))*c + 2*arctanh(c*x)/(e^3
*x^2 + 2*d*e^2*x + d^2*e))*a*b - 1/8*b^2*(log(-c*x + 1)^2/(e^3*x^2 + 2*d*e
^2*x + d^2*e) + 2*integrate(-((c*e*x - e)*log(c*x + 1)^2 + (c*e*x + c*d -
2*(c*e*x - e)*log(c*x + 1))*log(-c*x + 1))/(c*e^4*x^4 - d^3*e + (3*c*d*e^3
- e^4)*x^3 + 3*(c*d^2*e^2 - d*e^3)*x^2 + (c*d^3*e - 3*d^2*e^2)*x), x)) -
1/2*a^2/(e^3*x^2 + 2*d*e^2*x + d^2*e)
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^3} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^2}{(ex + d)^3} dx$$

input

```
integrate((a+b*arctanh(c*x))^2/(e*x+d)^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x) + a)^2/(e*x + d)^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^3} dx = \int \frac{(a + b \operatorname{atanh}(cx))^2}{(d + ex)^3} dx$$

input

```
int((a + b*atanh(c*x))^2/(d + e*x)^3,x)
```

output

```
int((a + b*atanh(c*x))^2/(d + e*x)^3, x)
```

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^2}{(d + ex)^3} dx = \text{too large to display}$$

input `int((a+b*atanh(c*x))^2/(e*x+d)^3,x)`

output

```
(12*atanh(c*x)**2*b**2*c**8*d**8*e*x + 6*atanh(c*x)**2*b**2*c**8*d**7*e**2
*x**2 - 18*atanh(c*x)**2*b**2*c**6*d**7*e**2 - 20*atanh(c*x)**2*b**2*c**6*
d**6*e**3*x - 10*atanh(c*x)**2*b**2*c**6*d**5*e**4*x**2 + 30*atanh(c*x)**2
*b**2*c**4*d**5*e**4 + 4*atanh(c*x)**2*b**2*c**4*d**4*e**5*x + 2*atanh(c*x)
)**2*b**2*c**4*d**3*e**6*x**2 - 6*atanh(c*x)**2*b**2*c**2*d**3*e**6 + 4*at
anh(c*x)**2*b**2*c**2*d**2*e**7*x + 2*atanh(c*x)**2*b**2*c**2*d*e**8*x**2
- 6*atanh(c*x)**2*b**2*d*e**8 - 12*atanh(c*x)*a*b*c**8*d**9 - 16*atanh(c*x)
)*a*b*c**6*d**7*e**2 + 56*atanh(c*x)*a*b*c**4*d**5*e**4 - 16*atanh(c*x)*a*
b*c**2*d**3*e**6 - 12*atanh(c*x)*a*b*d*e**8 - 22*atanh(c*x)*b**2*c**7*d**8
*e - 8*atanh(c*x)*b**2*c**7*d**7*e**2*x + 2*atanh(c*x)*b**2*c**7*d**6*e**3
*x**2 + 42*atanh(c*x)*b**2*c**5*d**6*e**3 + 24*atanh(c*x)*b**2*c**5*d**5*e
**4*x + 2*atanh(c*x)*b**2*c**5*d**4*e**5*x**2 - 18*atanh(c*x)*b**2*c**3*d
**4*e**5 - 24*atanh(c*x)*b**2*c**3*d**3*e**6*x - 10*atanh(c*x)*b**2*c**3*d
**2*e**7*x**2 - 2*atanh(c*x)*b**2*c*d**2*e**7 + 8*atanh(c*x)*b**2*c*d*e**8*
x + 6*atanh(c*x)*b**2*c*e**9*x**2 + 24*int((atanh(c*x)*x)/(c**4*d**5*x**2
+ 3*c**4*d**4*e*x**3 + 3*c**4*d**3*e**2*x**4 + c**4*d**2*e**3*x**5 - c**2*
d**5 - 3*c**2*d**4*e*x + 8*c**2*d**2*e**3*x**3 + 9*c**2*d*e**4*x**4 + 3*c*
**2*e**5*x**5 - 3*d**3*e**2 - 9*d**2*e**3*x - 9*d*e**4*x**2 - 3*e**5*x**3),
x)*b**2*c**11*d**13*e + 48*int((atanh(c*x)*x)/(c**4*d**5*x**2 + 3*c**4*d**
4*e*x**3 + 3*c**4*d**3*e**2*x**4 + c**4*d**2*e**3*x**5 - c**2*d**5 - 3*...
```

3.15 $\int (d + ex)^3 (a + \operatorname{barctanh}(cx))^3 dx$

Optimal result	155
Mathematica [A] (verified)	156
Rubi [A] (verified)	157
Maple [C] (warning: unable to verify)	159
Fricas [F]	159
Sympy [F]	160
Maxima [F]	160
Giac [F]	161
Mupad [F(-1)]	161
Reduce [F]	161

Optimal result

Integrand size = 18, antiderivative size = 614

$$\begin{aligned}
& \int (d + ex)^3 (a + \operatorname{barctanh}(cx))^3 dx \\
&= \frac{3ab^2de^2x}{c^2} + \frac{b^3e^3x}{4c^3} - \frac{b^3e^3\operatorname{arctanh}(cx)}{4c^4} + \frac{3b^3de^2x\operatorname{arctanh}(cx)}{c^2} \\
&+ \frac{b^2e^3x^2(a + \operatorname{barctanh}(cx))}{4c^2} - \frac{3bde^2(a + \operatorname{barctanh}(cx))^2}{2c^3} + \frac{be^3(a + \operatorname{barctanh}(cx))^2}{4c^4} \\
&+ \frac{3be(6c^2d^2 + e^2)(a + \operatorname{barctanh}(cx))^2}{4c^4} + \frac{3be(6c^2d^2 + e^2)x(a + \operatorname{barctanh}(cx))^2}{4c^3} \\
&+ \frac{3bde^2x^2(a + \operatorname{barctanh}(cx))^2}{2c} + \frac{be^3x^3(a + \operatorname{barctanh}(cx))^2}{4c} \\
&+ \frac{d(c^2d^2 + e^2)(a + \operatorname{barctanh}(cx))^3}{c^3} - \frac{(c^4d^4 + 6c^2d^2e^2 + e^4)(a + \operatorname{barctanh}(cx))^3}{4c^4e} \\
&+ \frac{(d + ex)^4(a + \operatorname{barctanh}(cx))^3}{4e} - \frac{b^2e^3(a + \operatorname{barctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{2c^4} \\
&- \frac{3b^2e(6c^2d^2 + e^2)(a + \operatorname{barctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{2c^4} \\
&- \frac{3bd(c^2d^2 + e^2)(a + \operatorname{barctanh}(cx))^2 \log\left(\frac{2}{1-cx}\right)}{c^3} + \frac{3b^3de^2 \log(1 - c^2x^2)}{2c^3} \\
&- \frac{b^3e^3 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{4c^4} - \frac{3b^3e(6c^2d^2 + e^2) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{4c^4} \\
&- \frac{3b^2d(c^2d^2 + e^2)(a + \operatorname{barctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{c^3} \\
&+ \frac{3b^3d(c^2d^2 + e^2) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx}\right)}{2c^3}
\end{aligned}$$

output

```

3*a*b^2*d*e^2*x/c^2+1/4*b^3*e^3*x/c^3-1/4*b^3*e^3*arctanh(c*x)/c^4+3*b^3*d
*e^2*x*arctanh(c*x)/c^2+1/4*b^2*e^3*x^2*(a+b*arctanh(c*x))/c^2-3/2*b*d*e^2
*(a+b*arctanh(c*x))^2/c^3+1/4*b*e^3*(a+b*arctanh(c*x))^2/c^4+3/4*b*e*(6*c^
2*d^2+e^2)*(a+b*arctanh(c*x))^2/c^4+3/4*b*e*(6*c^2*d^2+e^2)*x*(a+b*arctanh
(c*x))^2/c^3+3/2*b*d*e^2*x^2*(a+b*arctanh(c*x))^2/c+1/4*b*e^3*x^3*(a+b*arc
tanh(c*x))^2/c+d*(c^2*d^2+e^2)*(a+b*arctanh(c*x))^3/c^3-1/4*(c^4*d^4+6*c^2
*d^2*e^2+e^4)*(a+b*arctanh(c*x))^3/c^4/e+1/4*(e*x+d)^4*(a+b*arctanh(c*x))^
3/e-1/2*b^2*e^3*(a+b*arctanh(c*x))*ln(2/(-c*x+1))/c^4-3/2*b^2*e*(6*c^2*d^2
+e^2)*(a+b*arctanh(c*x))*ln(2/(-c*x+1))/c^4-3*b*d*(c^2*d^2+e^2)*(a+b*arcta
nh(c*x))^2*ln(2/(-c*x+1))/c^3+3/2*b^3*d*e^2*ln(-c^2*x^2+1)/c^3-1/4*b^3*e^3
*polylog(2,1-2/(-c*x+1))/c^4-3/4*b^3*e*(6*c^2*d^2+e^2)*polylog(2,1-2/(-c*x
+1))/c^4-3*b^2*d*(c^2*d^2+e^2)*(a+b*arctanh(c*x))*polylog(2,1-2/(-c*x+1))/
c^3+3/2*b^3*d*(c^2*d^2+e^2)*polylog(3,1-2/(-c*x+1))/c^3

```

Mathematica [A] (verified)

Time = 1.24 (sec) , antiderivative size = 830, normalized size of antiderivative = 1.35

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx))^3 dx = \text{Too large to display}$$

input

```
Integrate[(d + e*x)^3*(a + b*ArcTanh[c*x])^3,x]
```

output

```
(2*a^2*c*(4*a*c^3*d^3 + 3*b*e*(6*c^2*d^2 + e^2))*x + 12*a^2*c^3*d*e*(a*c*d
+ b*e)*x^2 + 2*a^2*c^3*e^2*(4*a*c*d + b*e)*x^3 + 2*a^3*c^4*e^3*x^4 + 6*a^
2*b*c^4*x*(4*d^3 + 6*d^2*e*x + 4*d*e^2*x^2 + e^3*x^3)*ArcTanh[c*x] + 3*a^2
*b*(4*c^3*d^3 + 6*c^2*d^2*e + 4*c*d*e^2 + e^3)*Log[1 - c*x] + 3*a^2*b*(4*c
^3*d^3 - 6*c^2*d^2*e + 4*c*d*e^2 - e^3)*Log[1 + c*x] + 36*a*b^2*c^2*d^2*e*
(2*c*x*ArcTanh[c*x] + (-1 + c^2*x^2)*ArcTanh[c*x]^2 + Log[1 - c^2*x^2]) +
2*a*b^2*e^3*(-1 + c^2*x^2 + 2*c*x*(3 + c^2*x^2)*ArcTanh[c*x] + 3*(-1 + c^4
*x^4)*ArcTanh[c*x]^2 + 4*Log[1 - c^2*x^2]) - 12*b^3*c^2*d^2*e*(ArcTanh[c*x
]*((3 - 3*c*x)*ArcTanh[c*x] + (1 - c^2*x^2)*ArcTanh[c*x]^2 + 6*Log[1 + E^(-
2*ArcTanh[c*x]))] - 3*PolyLog[2, -E^(-2*ArcTanh[c*x]))] + 24*a*b^2*c*d*e^
2*(c*x + (-1 + c^3*x^3)*ArcTanh[c*x]^2 + ArcTanh[c*x]*(-1 + c^2*x^2 - 2*Lo
g[1 + E^(-2*ArcTanh[c*x]))] + PolyLog[2, -E^(-2*ArcTanh[c*x]))] + 24*a*b^2
*c^3*d^3*(ArcTanh[c*x]*((-1 + c*x)*ArcTanh[c*x] - 2*Log[1 + E^(-2*ArcTanh[
c*x]))] + PolyLog[2, -E^(-2*ArcTanh[c*x]))] + 2*b^3*e^3*(c*x + (-4 + 3*c*x
+ c^3*x^3)*ArcTanh[c*x]^2 + (-1 + c^4*x^4)*ArcTanh[c*x]^3 + ArcTanh[c*x]*
(-1 + c^2*x^2 - 8*Log[1 + E^(-2*ArcTanh[c*x]))] + 4*PolyLog[2, -E^(-2*ArcT
anh[c*x]))] + 8*b^3*c^3*d^3*(ArcTanh[c*x]^2*((-1 + c*x)*ArcTanh[c*x] - 3*L
og[1 + E^(-2*ArcTanh[c*x]))] + 3*ArcTanh[c*x]*PolyLog[2, -E^(-2*ArcTanh[c*
x]))] + (3*PolyLog[3, -E^(-2*ArcTanh[c*x]))]/2) + 4*b^3*c*d*e^2*(6*c*x*ArcT
anh[c*x] - 3*ArcTanh[c*x]^2 + 3*c^2*x^2*ArcTanh[c*x]^2 - 2*ArcTanh[c*x]...
```

Rubi [A] (verified)

Time = 1.49 (sec) , antiderivative size = 608, normalized size of antiderivative = 0.99, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex)^3 (a + \operatorname{barctanh}(cx))^3 dx$$

↓ 6480

$$\frac{(d + ex)^4 (a + \operatorname{barctanh}(cx))^3}{4e} - \frac{3bc \int \left(-\frac{x^2 (a + \operatorname{barctanh}(cx))^2 e^4}{c^2} - \frac{4dx (a + \operatorname{barctanh}(cx))^2 e^3}{c^2} - \frac{(6c^2 d^2 + e^2) (a + \operatorname{barctanh}(cx))^2 e^2}{c^4} + \frac{(c^4 d^4 + 6c^2 e^2 d^2 + 4c^2 e (c^2 d^2 + e^2))}{c^4 (1 - c^2 x^2)} \right) dx}{4e}$$

$$\begin{array}{c} \downarrow 2009 \\ \frac{(d+ex)^4(a+\operatorname{arctanh}(cx))^3}{4e} \\ \hline 3bc \left(-\frac{e^4(a+\operatorname{arctanh}(cx))^2}{3c^5} + \frac{2be^4 \log\left(\frac{2}{1-cx}\right)(a+\operatorname{arctanh}(cx))}{3c^5} + \frac{2de^3(a+\operatorname{arctanh}(cx))^2}{c^4} - \frac{be^4x^2(a+\operatorname{arctanh}(cx))}{3c^3} - \frac{2de^3x}{c^3} \right) \end{array}$$

input `Int[(d + e*x)^3*(a + b*ArcTanh[c*x])^3, x]`

output

```
((d + e*x)^4*(a + b*ArcTanh[c*x])^3)/(4*e) - (3*b*c*((-4*a*b*d*e^3*x)/c^3 - (b^2*e^4*x)/(3*c^4) + (b^2*e^4*ArcTanh[c*x])/(3*c^5) - (4*b^2*d*e^3*x*ArcTanh[c*x])/c^3 - (b*e^4*x^2*(a + b*ArcTanh[c*x]))/(3*c^3) + (2*d*e^3*(a + b*ArcTanh[c*x])^2)/c^4 - (e^4*(a + b*ArcTanh[c*x])^2)/(3*c^5) - (e^2*(6*c^2*d^2 + e^2)*(a + b*ArcTanh[c*x])^2)/c^5 - (e^2*(6*c^2*d^2 + e^2)*x*(a + b*ArcTanh[c*x])^2)/c^4 - (2*d*e^3*x^2*(a + b*ArcTanh[c*x])^2)/c^2 - (e^4*x^3*(a + b*ArcTanh[c*x])^2)/(3*c^2) - (4*d*e*(c^2*d^2 + e^2)*(a + b*ArcTanh[c*x])^3)/(3*b*c^4) + ((c^4*d^4 + 6*c^2*d^2*e^2 + e^4)*(a + b*ArcTanh[c*x])^3)/(3*b*c^5) + (2*b*e^4*(a + b*ArcTanh[c*x])*Log[2/(1 - c*x)]/(3*c^5) + (2*b*e^2*(6*c^2*d^2 + e^2)*(a + b*ArcTanh[c*x])*Log[2/(1 - c*x)]/c^5 + (4*d*e*(c^2*d^2 + e^2)*(a + b*ArcTanh[c*x])^2*Log[2/(1 - c*x)]/c^4 - (2*b^2*d*e^3*Log[1 - c^2*x^2])/c^4 + (b^2*e^4*PolyLog[2, 1 - 2/(1 - c*x)]/(3*c^5) + (b^2*e^2*(6*c^2*d^2 + e^2)*PolyLog[2, 1 - 2/(1 - c*x)]/c^5 + (4*b*d*e*(c^2*d^2 + e^2)*(a + b*ArcTanh[c*x])*PolyLog[2, 1 - 2/(1 - c*x)]/c^4 - (2*b^2*d*e*(c^2*d^2 + e^2)*PolyLog[3, 1 - 2/(1 - c*x)]/c^4))/(4*e)
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6480

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^p*((d_) + (e_.)*(x_.))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 28.29 (sec) , antiderivative size = 5093, normalized size of antiderivative = 8.29

method	result	size
parts	Expression too large to display	5093
derivativedivides	Expression too large to display	5124
default	Expression too large to display	5124

input `int((e*x+d)^3*(a+b*arctanh(c*x))^3,x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F]

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx))^3 dx = \int (ex + d)^3 (b \operatorname{arctanh}(cx) + a)^3 dx$$

input `integrate((e*x+d)^3*(a+b*arctanh(c*x))^3,x, algorithm="fricas")`

output `integral(a^3*e^3*x^3 + 3*a^3*d*e^2*x^2 + 3*a^3*d^2*e*x + a^3*d^3 + (b^3*e^3*x^3 + 3*b^3*d*e^2*x^2 + 3*b^3*d^2*e*x + b^3*d^3)*arctanh(c*x)^3 + 3*(a*b^2*e^3*x^3 + 3*a*b^2*d*e^2*x^2 + 3*a*b^2*d^2*e*x + a*b^2*d^3)*arctanh(c*x)^2 + 3*(a^2*b*e^3*x^3 + 3*a^2*b*d*e^2*x^2 + 3*a^2*b*d^2*e*x + a^2*b*d^3)*arctanh(c*x), x)`

Sympy [F]

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx))^3 dx = \int (a + b \operatorname{atanh}(cx))^3 (d + ex)^3 dx$$

input `integrate((e*x+d)**3*(a+b*atanh(c*x))**3,x)`

output `Integral((a + b*atanh(c*x))**3*(d + e*x)**3, x)`

Maxima [F]

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx))^3 dx = \int (ex + d)^3 (b \operatorname{artanh}(cx) + a)^3 dx$$

input `integrate((e*x+d)^3*(a+b*arctanh(c*x))^3,x, algorithm="maxima")`

output

```

1/4*a^3*e^3*x^4 + a^3*d*e^2*x^3 + 3/2*a^3*d^2*e*x^2 + 9/4*(2*x^2*arctanh(c
*x) + c*(2*x/c^2 - log(c*x + 1)/c^3 + log(c*x - 1)/c^3))*a^2*b*d^2*e + 3/2
*(2*x^3*arctanh(c*x) + c*(x^2/c^2 + log(c^2*x^2 - 1)/c^4))*a^2*b*d*e^2 + 1
/8*(6*x^4*arctanh(c*x) + c*(2*(c^2*x^3 + 3*x)/c^4 - 3*log(c*x + 1)/c^5 + 3
*log(c*x - 1)/c^5))*a^2*b*e^3 + a^3*d^3*x + 3/2*(2*c*x*arctanh(c*x) + log(
-c^2*x^2 + 1))*a^2*b*d^3/c - 1/32*((b^3*c^4*e^3*x^4 + 4*b^3*c^4*d*e^2*x^3
+ 6*b^3*c^4*d^2*e*x^2 + 4*b^3*c^4*d^3*x - (4*c^3*d^3 + 6*c^2*d^2*e + 4*c*d
*e^2 + e^3)*b^3)*log(-c*x + 1)^3 - (6*a*b^2*c^4*e^3*x^4 + 2*(12*a*b^2*c^4*
d*e^2 + b^3*c^3*e^3)*x^3 + 12*(3*a*b^2*c^4*d^2*e + b^3*c^3*d*e^2)*x^2 + 6*
(4*a*b^2*c^4*d^3 + (6*c^3*d^2*e + c*e^3)*b^3)*x + 3*(b^3*c^4*e^3*x^4 + 4*b
^3*c^4*d*e^2*x^3 + 6*b^3*c^4*d^2*e*x^2 + 4*b^3*c^4*d^3*x + (4*c^3*d^3 - 6*
c^2*d^2*e + 4*c*d*e^2 - e^3)*b^3)*log(c*x + 1))*log(-c*x + 1)^2/c^4 - int
egrate(-1/16*(2*(b^3*c^4*e^3*x^4 - b^3*c^3*d^3 + (3*c^4*d*e^2 - c^3*e^3)*b
^3*x^3 + 3*(c^4*d^2*e - c^3*d*e^2)*b^3*x^2 + (c^4*d^3 - 3*c^3*d^2*e)*b^3*x
)*log(c*x + 1)^3 + 12*(a*b^2*c^4*e^3*x^4 - a*b^2*c^3*d^3 + (3*c^4*d*e^2 -
c^3*e^3)*a*b^2*x^3 + 3*(c^4*d^2*e - c^3*d*e^2)*a*b^2*x^2 + (c^4*d^3 - 3*c^
3*d^2*e)*a*b^2*x)*log(c*x + 1)^2 - (6*a*b^2*c^4*e^3*x^4 + 2*(12*a*b^2*c^4*
d*e^2 + b^3*c^3*e^3)*x^3 + 12*(3*a*b^2*c^4*d^2*e + b^3*c^3*d*e^2)*x^2 + 6*
(b^3*c^4*e^3*x^4 - b^3*c^3*d^3 + (3*c^4*d*e^2 - c^3*e^3)*b^3*x^3 + 3*(c^4*
d^2*e - c^3*d*e^2)*b^3*x^2 + (c^4*d^3 - 3*c^3*d^2*e)*b^3*x)*log(c*x + 1...

```

Giac [F]

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx))^3 dx = \int (ex + d)^3 (b \operatorname{artanh}(cx) + a)^3 dx$$

input `integrate((e*x+d)^3*(a+b*arctanh(c*x))^3,x, algorithm="giac")`

output `integrate((e*x + d)^3*(b*arctanh(c*x) + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx))^3 dx = \int (a + b \operatorname{atanh}(cx))^3 (d + ex)^3 dx$$

input `int((a + b*atanh(c*x))^3*(d + e*x)^3,x)`

output `int((a + b*atanh(c*x))^3*(d + e*x)^3, x)`

Reduce [F]

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx))^3 dx = \text{Too large to display}$$

input `int((e*x+d)^3*(a+b*atanh(c*x))^3,x)`

output

```
(4*atanh(c*x)**3*b**3*c**4*d**3*x + 6*atanh(c*x)**3*b**3*c**4*d**2*e*x**2
+ 4*atanh(c*x)**3*b**3*c**4*d*e**2*x**3 + atanh(c*x)**3*b**3*c**4*e**3*x**
4 - 6*atanh(c*x)**3*b**3*c**2*d**2*e - atanh(c*x)**3*b**3*e**3 + 12*atanh(
c*x)**2*a*b**2*c**4*d**3*x + 18*atanh(c*x)**2*a*b**2*c**4*d**2*e*x**2 + 12
*atanh(c*x)**2*a*b**2*c**4*d*e**2*x**3 + 3*atanh(c*x)**2*a*b**2*c**4*e**3*
x**4 - 18*atanh(c*x)**2*a*b**2*c**2*d**2*e - 3*atanh(c*x)**2*a*b**2*e**3 +
18*atanh(c*x)**2*b**3*c**3*d**2*e*x + 6*atanh(c*x)**2*b**3*c**3*d*e**2*x*
*2 + atanh(c*x)**2*b**3*c**3*e**3*x**3 - 6*atanh(c*x)**2*b**3*c*d*e**2 + 3
*atanh(c*x)**2*b**3*c*e**3*x + 12*atanh(c*x)*a**2*b*c**4*d**3*x + 18*atanh
(c*x)*a**2*b*c**4*d**2*e*x**2 + 12*atanh(c*x)*a**2*b*c**4*d*e**2*x**3 + 3*
atanh(c*x)*a**2*b*c**4*e**3*x**4 + 12*atanh(c*x)*a**2*b*c**3*d**3 - 18*ata
nh(c*x)*a**2*b*c**2*d**2*e + 12*atanh(c*x)*a**2*b*c*d*e**2 - 3*atanh(c*x)*
a**2*b*e**3 + 36*atanh(c*x)*a*b**2*c**3*d**2*e*x + 12*atanh(c*x)*a*b**2*c*
*3*d*e**2*x**2 + 2*atanh(c*x)*a*b**2*c**3*e**3*x**3 + 36*atanh(c*x)*a*b**2
*c**2*d**2*e - 12*atanh(c*x)*a*b**2*c*d*e**2 + 6*atanh(c*x)*a*b**2*c*e**3*
x + 8*atanh(c*x)*a*b**2*e**3 + 12*atanh(c*x)*b**3*c**2*d*e**2*x + atanh(c*
x)*b**3*c**2*e**3*x**2 + 12*atanh(c*x)*b**3*c*d*e**2 - atanh(c*x)*b**3*e**
3 + 24*int((atanh(c*x)*x)/(c**2*x**2 - 1),x)*a*b**2*c**5*d**3 + 24*int((at
anh(c*x)*x)/(c**2*x**2 - 1),x)*a*b**2*c**3*d*e**2 + 36*int((atanh(c*x)*x)/
(c**2*x**2 - 1),x)*b**3*c**4*d**2*e + 8*int((atanh(c*x)*x)/(c**2*x**2 - ...
```

3.16 $\int (d + ex)^2 (a + \operatorname{barctanh}(cx))^3 dx$

Optimal result	163
Mathematica [A] (verified)	164
Rubi [A] (verified)	165
Maple [C] (warning: unable to verify)	166
Fricas [F]	167
Sympy [F]	168
Maxima [F]	168
Giac [F]	169
Mupad [F(-1)]	169
Reduce [F]	169

Optimal result

Integrand size = 18, antiderivative size = 387

$$\begin{aligned}
 & \int (d + ex)^2 (a + \operatorname{barctanh}(cx))^3 dx \\
 &= \frac{ab^2 e^2 x}{c^2} + \frac{b^3 e^2 x \operatorname{arctanh}(cx)}{c^2} + \frac{3bde(a + \operatorname{barctanh}(cx))^2}{c^2} - \frac{be^2(a + \operatorname{barctanh}(cx))^2}{2c^3} \\
 &+ \frac{3bdex(a + \operatorname{barctanh}(cx))^2}{c} + \frac{be^2 x^2 (a + \operatorname{barctanh}(cx))^2}{2c} \\
 &+ \frac{(3c^2 d^2 + e^2)(a + \operatorname{barctanh}(cx))^3}{3c^3} - \frac{d\left(d^2 + \frac{3e^2}{c^2}\right)(a + \operatorname{barctanh}(cx))^3}{3e} \\
 &+ \frac{(d + ex)^3 (a + \operatorname{barctanh}(cx))^3}{3e} - \frac{6b^2 de(a + \operatorname{barctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{c^2} \\
 &- \frac{b(3c^2 d^2 + e^2)(a + \operatorname{barctanh}(cx))^2 \log\left(\frac{2}{1-cx}\right)}{c^3} \\
 &+ \frac{b^3 e^2 \log(1 - c^2 x^2)}{2c^3} - \frac{3b^3 de \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{c^2} \\
 &- \frac{b^2(3c^2 d^2 + e^2)(a + \operatorname{barctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{c^3} \\
 &+ \frac{b^3(3c^2 d^2 + e^2) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx}\right)}{2c^3}
 \end{aligned}$$

```
output
a*b^2*e^2*x/c^2+b^3*e^2*x*arctanh(c*x)/c^2+3*b*d*e*(a+b*arctanh(c*x))^2/c^
2-1/2*b*e^2*(a+b*arctanh(c*x))^2/c^3+3*b*d*e*x*(a+b*arctanh(c*x))^2/c+1/2*
b*e^2*x^2*(a+b*arctanh(c*x))^2/c+1/3*(3*c^2*d^2+e^2)*(a+b*arctanh(c*x))^3/
c^3-1/3*d*(d^2+3*e^2/c^2)*(a+b*arctanh(c*x))^3/e+1/3*(e*x+d)^3*(a+b*arctan
h(c*x))^3/e-6*b^2*d*e*(a+b*arctanh(c*x))*ln(2/(-c*x+1))/c^2-b*(3*c^2*d^2+e
^2)*(a+b*arctanh(c*x))^2*ln(2/(-c*x+1))/c^3+1/2*b^3*e^2*ln(-c^2*x^2+1)/c^3
-3*b^3*d*e*polylog(2,1-2/(-c*x+1))/c^2-b^2*(3*c^2*d^2+e^2)*(a+b*arctanh(c*
x))*polylog(2,1-2/(-c*x+1))/c^3+1/2*b^3*(3*c^2*d^2+e^2)*polylog(3,1-2/(-c*
x+1))/c^3
```

Mathematica [A] (verified)

Time = 0.77 (sec) , antiderivative size = 591, normalized size of antiderivative = 1.53

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^3 dx$$

$$= \frac{6a^2 c^2 d(acd + 3be)x + 3a^2 c^2 e(2acd + be)x^2 + 2a^3 c^3 e^2 x^3 + 6a^2 bc^3 x(3d^2 + 3dex + e^2 x^2) \operatorname{arctanh}(cx) + 3a^3 d^2 c^2 \operatorname{arctanh}(cx)^2 + 3a^2 b^2 c^2 d e \operatorname{arctanh}(cx)^2 + 3a^2 b^2 c^2 e^2 \operatorname{arctanh}(cx)^2 \ln(2/(-c x + 1)) - b(3c^2 d^2 + e^2) \operatorname{arctanh}(cx)^2 \ln(2/(-c x + 1)) + 1/2 b^3 e^2 \ln(-c^2 x^2 + 1) - 3 b^3 d e \operatorname{polylog}(2, 1 - 2/(-c x + 1)) - b^2 (3c^2 d^2 + e^2) \operatorname{arctanh}(c x) \operatorname{polylog}(2, 1 - 2/(-c x + 1)) + 1/2 b^3 (3c^2 d^2 + e^2) \operatorname{polylog}(3, 1 - 2/(-c x + 1))}{6c^3}$$

```
input
Integrate[(d + e*x)^2*(a + b*ArcTanh[c*x])^3,x]
```

```
output
(6*a^2*c^2*d*(a*c*d + 3*b*e)*x + 3*a^2*c^2*e*(2*a*c*d + b*e)*x^2 + 2*a^3*c^
3*e^2*x^3 + 6*a^2*b*c^3*x*(3*d^2 + 3*d*e*x + e^2*x^2)*ArcTanh[c*x] + 3*a^
2*b*(3*c^2*d^2 + 3*c*d*e + e^2)*Log[1 - c*x] + 3*a^2*b*(3*c^2*d^2 - 3*c*d*
e + e^2)*Log[1 + c*x] + 18*a*b^2*c*d*e*(2*c*x*ArcTanh[c*x] + (-1 + c^2*x^2
)*ArcTanh[c*x]^2 + Log[1 - c^2*x^2]) - 6*b^3*c*d*e*(ArcTanh[c*x]*((3 - 3*c
*x)*ArcTanh[c*x] + (1 - c^2*x^2)*ArcTanh[c*x]^2 + 6*Log[1 + E^(-2*ArcTan
h[c*x])) - 3*PolyLog[2, -E^(-2*ArcTanh[c*x])) + 6*a*b^2*e^2*(c*x + (-1 + c
^3*x^3)*ArcTanh[c*x]^2 + ArcTanh[c*x]*(-1 + c^2*x^2 - 2*Log[1 + E^(-2*ArcT
anh[c*x])) + PolyLog[2, -E^(-2*ArcTanh[c*x])) + 18*a*b^2*c^2*d^2*(ArcTan
h[c*x]*((-1 + c*x)*ArcTanh[c*x] - 2*Log[1 + E^(-2*ArcTanh[c*x])) + PolyLo
g[2, -E^(-2*ArcTanh[c*x])) + 6*b^3*c^2*d^2*(ArcTanh[c*x]^2*((-1 + c*x)*Ar
cTanh[c*x] - 3*Log[1 + E^(-2*ArcTanh[c*x])) + 3*ArcTanh[c*x]*PolyLog[2, -
E^(-2*ArcTanh[c*x])) + (3*PolyLog[3, -E^(-2*ArcTanh[c*x]))/2) + b^3*e^2*(
6*c*x*ArcTanh[c*x] - 3*ArcTanh[c*x]^2 + 3*c^2*x^2*ArcTanh[c*x]^2 - 2*ArcTa
nh[c*x]^3 + 2*c^3*x^3*ArcTanh[c*x]^3 - 6*ArcTanh[c*x]^2*Log[1 + E^(-2*ArcT
anh[c*x])) + 3*Log[1 - c^2*x^2] + 6*ArcTanh[c*x]*PolyLog[2, -E^(-2*ArcTan
h[c*x])) + 3*PolyLog[3, -E^(-2*ArcTanh[c*x])))/(6*c^3)
```

Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 403, normalized size of antiderivative = 1.04, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^3 dx$$

↓ 6480

$$\frac{(d + ex)^3 (a + b \operatorname{arctanh}(cx))^3}{3e} - \frac{bc \int \left(-\frac{x(a + b \operatorname{arctanh}(cx))^2 e^3}{c^2} - \frac{3d(a + b \operatorname{arctanh}(cx))^2 e^2}{c^2} + \frac{(d(c^2 d^2 + 3e^2) + e(3c^2 d^2 + e^2)x)(a + b \operatorname{arctanh}(cx))^2}{c^2(1 - c^2 x^2)} \right) dx}{e}$$

↓ 2009

$$\frac{(d + ex)^3 (a + b \operatorname{arctanh}(cx))^3}{3e} - \frac{bc \left(\frac{e^3 (a + b \operatorname{arctanh}(cx))^2}{2c^4} - \frac{3de^2 (a + b \operatorname{arctanh}(cx))^2}{c^3} + \frac{6bde^2 \log\left(\frac{2}{1 - cx}\right) (a + b \operatorname{arctanh}(cx))}{c^3} - \frac{3de^2 x (a + b \operatorname{arctanh}(cx))^2}{c^2} - \frac{e^3 x^2 (a + b \operatorname{arctanh}(cx))^2}{c^2} \right)}{e}$$

input

```
Int[(d + e*x)^2*(a + b*ArcTanh[c*x])^3,x]
```

output

```
((d + e*x)^3*(a + b*ArcTanh[c*x])^3)/(3*e) - (b*c*(-((a*b*e^3*x)/c^3) - (b^2*e^3*x*ArcTanh[c*x])/c^3 - (3*d*e^2*(a + b*ArcTanh[c*x])^2)/c^3 + (e^3*(a + b*ArcTanh[c*x])^2)/(2*c^4) - (3*d*e^2*x*(a + b*ArcTanh[c*x])^2)/c^2 - (e^3*x^2*(a + b*ArcTanh[c*x])^2)/(2*c^2) - (e*(3*c^2*d^2 + e^2)*(a + b*ArcTanh[c*x])^3)/(3*b*c^4) + (d*(c^2*d^2 + 3*e^2)*(a + b*ArcTanh[c*x])^3)/(3*b*c^3) + (6*b*d*e^2*(a + b*ArcTanh[c*x])*Log[2/(1 - c*x)])/c^3 + (e*(3*c^2*d^2 + e^2)*(a + b*ArcTanh[c*x])^2*Log[2/(1 - c*x)])/c^4 - (b^2*e^3*Log[1 - c^2*x^2])/(2*c^4) + (3*b^2*d*e^2*PolyLog[2, 1 - 2/(1 - c*x)])/c^3 + (b*e*(3*c^2*d^2 + e^2)*(a + b*ArcTanh[c*x])*PolyLog[2, 1 - 2/(1 - c*x)])/c^4 - (b^2*e*(3*c^2*d^2 + e^2)*PolyLog[3, 1 - 2/(1 - c*x)]/(2*c^4))/e
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6480 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_)*((d_) + (e_.)*(x_))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 14.83 (sec) , antiderivative size = 3848, normalized size of antiderivative = 9.94

method	result	size
parts	Expression too large to display	3848
derivativedivides	Expression too large to display	3873
default	Expression too large to display	3873

input `int((e*x+d)^2*(a+b*arctanh(c*x))^3,x,method=_RETURNVERBOSE)`

output

```

1/3*a^3*(e*x+d)^3/e+b^3/c*(1/3*c*e^2*arctanh(c*x)^3*x^3+c*e*arctanh(c*x)^3
*x^2*d+arctanh(c*x)^3*c*x*d^2+1/3*c/e*arctanh(c*x)^3*d^3-1/c^2/e*(-1/2*arc
tanh(c*x)^2*ln(c*x-1)*e^-3-1/2*arctanh(c*x)^2*ln(c*x+1)*e^-3-(c*x+1)*arctanh
(c*x)*e^3+ln(2)*e^3*arctanh(c*x)^2+1/3*c^3*d^3*arctanh(c*x)^3+ln((c*x+1)/(-
c^2*x^2+1)^(1/2))*e^3*arctanh(c*x)^2+polylog(2,-(c*x+1)^2/(-c^2*x^2+1))*e
^3*arctanh(c*x)+3/2*arctanh(c*x)^2*ln(c*x+1)*c*d*e^2-3/2*arctanh(c*x)^2*ln
(c*x-1)*c^2*d^2*e+1/4*I*Pi*c^3*d^3*csgn(I*(c*x+1)^2/(c^2*x^2-1))*csgn(I*(c
*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I/(1-(c*x+1)^2/(c^2*x^
2-1)))*arctanh(c*x)^2+3/4*I*Pi*c*d*e^2*csgn(I*(c*x+1)^2/(c^2*x^2-1))*csgn(
I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^2*arctanh(c*x)^2-3/4*I*
Pi*c^2*d^2*e*csgn(I*(c*x+1)^2/(c^2*x^2-1))*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1
-(c*x+1)^2/(c^2*x^2-1)))^2*arctanh(c*x)^2-3/2*I*Pi*c*d*e^2*csgn(I*(c*x+1)/
(-c^2*x^2+1)^(1/2))*csgn(I*(c*x+1)^2/(c^2*x^2-1))^2*arctanh(c*x)^2+3/4*I*P
i*c^2*d^2*e*csgn(I*(c*x+1)/(-c^2*x^2+1)^(1/2))^2*csgn(I*(c*x+1)^2/(c^2*x^2
-1))*arctanh(c*x)^2-3/4*I*Pi*c*d*e^2*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+
1)^2/(c^2*x^2-1)))^2*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))*arctanh(c*x)^2-3/4*
I*Pi*c*d*e^2*csgn(I*(c*x+1)/(-c^2*x^2+1)^(1/2))^2*csgn(I*(c*x+1)^2/(c^2*x^
2-1))*arctanh(c*x)^2+3/2*I*Pi*c^2*d^2*e*csgn(I*(c*x+1)/(-c^2*x^2+1)^(1/2))
*csgn(I*(c*x+1)^2/(c^2*x^2-1))^2*arctanh(c*x)^2+3/4*I*Pi*c^2*d^2*e*csgn(I*
(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^2*csgn(I/(1-(c*x+1)^2/...

```

Fricas [F]

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^3 dx = \int (ex + d)^2 (b \operatorname{arctanh}(cx) + a)^3 dx$$

input

```
integrate((e*x+d)^2*(a+b*arctanh(c*x))^3,x, algorithm="fricas")
```

output

```

integral(a^3*e^2*x^2 + 2*a^3*d*e*x + a^3*d^2 + (b^3*e^2*x^2 + 2*b^3*d*e*x
+ b^3*d^2)*arctanh(c*x)^3 + 3*(a*b^2*e^2*x^2 + 2*a*b^2*d*e*x + a*b^2*d^2)*
arctanh(c*x)^2 + 3*(a^2*b*e^2*x^2 + 2*a^2*b*d*e*x + a^2*b*d^2)*arctanh(c*x
), x)

```


Sympy [F]

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^3 dx = \int (a + b \operatorname{atanh}(cx))^3 (d + ex)^2 dx$$

input `integrate((e*x+d)**2*(a+b*atanh(c*x))**3,x)`

output `Integral((a + b*atanh(c*x))**3*(d + e*x)**2, x)`

Maxima [F]

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^3 dx = \int (ex + d)^2 (b \operatorname{artanh}(cx) + a)^3 dx$$

input `integrate((e*x+d)^2*(a+b*arctanh(c*x))^3,x, algorithm="maxima")`

output `1/3*a^3*e^2*x^3 + a^3*d*e*x^2 + 3/2*(2*x^2*arctanh(c*x) + c*(2*x/c^2 - log(c*x + 1)/c^3 + log(c*x - 1)/c^3))*a^2*b*d*e + 1/2*(2*x^3*arctanh(c*x) + c*(x^2/c^2 + log(c^2*x^2 - 1)/c^4))*a^2*b*e^2 + a^3*d^2*x + 3/2*(2*c*x*arctanh(c*x) + log(-c^2*x^2 + 1))*a^2*b*d^2/c - 1/24*((b^3*c^3*e^2*x^3 + 3*b^3*c^3*d*e*x^2 + 3*b^3*c^3*d^2*x - (3*c^2*d^2 + 3*c*d*e + e^2)*b^3)*log(-c*x + 1)^3 - 3*(2*a*b^2*c^3*e^2*x^3 + (6*a*b^2*c^3*d*e + b^3*c^2*e^2)*x^2 + 6*(a*b^2*c^3*d^2 + b^3*c^2*d*e)*x + (b^3*c^3*e^2*x^3 + 3*b^3*c^3*d*e*x^2 + 3*b^3*c^3*d^2*x + (3*c^2*d^2 - 3*c*d*e + e^2)*b^3)*log(c*x + 1))*log(-c*x + 1)^2)/c^3 - integrate(-1/8*((b^3*c^3*e^2*x^3 - b^3*c^2*d^2 + (2*c^3*d*e - c^2*e^2)*b^3*x^2 + (c^3*d^2 - 2*c^2*d*e)*b^3*x)*log(c*x + 1)^3 + 6*(a*b^2*c^3*e^2*x^3 - a*b^2*c^2*d^2 + (2*c^3*d*e - c^2*e^2)*a*b^2*x^2 + (c^3*d^2 - 2*c^2*d*e)*a*b^2*x)*log(c*x + 1)^2 - (4*a*b^2*c^3*e^2*x^3 + 2*(6*a*b^2*c^3*d*e + b^3*c^2*e^2)*x^2 + 3*(b^3*c^3*e^2*x^3 - b^3*c^2*d^2 + (2*c^3*d*e - c^2*e^2)*b^3*x^2 + (c^3*d^2 - 2*c^2*d*e)*b^3*x)*log(c*x + 1)^2 + 12*(a*b^2*c^3*d^2 + b^3*c^2*d*e)*x - 2*(6*a*b^2*c^2*d^2 - (3*c^2*d^2 - 3*c*d*e + e^2)*b^3 - (6*a*b^2*c^3*e^2 + b^3*c^3*e^2)*x^3 - 3*(b^3*c^3*d*e + 2*(2*c^3*d*e - c^2*e^2)*a*b^2)*x^2 - 3*(b^3*c^3*d^2 + 2*(c^3*d^2 - 2*c^2*d*e)*a*b^2)*x)*log(c*x + 1))*log(-c*x + 1))/(c^3*x - c^2), x)`

Giac [F]

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^3 dx = \int (ex + d)^2 (b \operatorname{artanh}(cx) + a)^3 dx$$

input `integrate((e*x+d)^2*(a+b*arctanh(c*x))^3,x, algorithm="giac")`

output `integrate((e*x + d)^2*(b*arctanh(c*x) + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^3 dx = \int (a + b \operatorname{atanh}(cx))^3 (d + ex)^2 dx$$

input `int((a + b*atanh(c*x))^3*(d + e*x)^2,x)`

output `int((a + b*atanh(c*x))^3*(d + e*x)^2, x)`

Reduce [F]

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx))^3 dx = \text{Too large to display}$$

input `int((e*x+d)^2*(a+b*atanh(c*x))^3,x)`

output

```
(6*atanh(c*x)**3*b**3*c**3*d**2*x + 6*atanh(c*x)**3*b**3*c**3*d*e*x**2 + 2
*atanh(c*x)**3*b**3*c**3*e**2*x**3 - 6*atanh(c*x)**3*b**3*c*d*e + 18*atanh
(c*x)**2*a*b**2*c**3*d**2*x + 18*atanh(c*x)**2*a*b**2*c**3*d*e*x**2 + 6*at
anh(c*x)**2*a*b**2*c**3*e**2*x**3 - 18*atanh(c*x)**2*a*b**2*c*d*e + 18*ata
nh(c*x)**2*b**3*c**2*d*e*x + 3*atanh(c*x)**2*b**3*c**2*e**2*x**2 - 3*atanh
(c*x)**2*b**3*e**2 + 18*atanh(c*x)*a**2*b*c**3*d**2*x + 18*atanh(c*x)*a**2
*b*c**3*d*e*x**2 + 6*atanh(c*x)*a**2*b*c**3*e**2*x**3 + 18*atanh(c*x)*a**2
*b*c**2*d**2 - 18*atanh(c*x)*a**2*b*c*d*e + 6*atanh(c*x)*a**2*b*e**2 + 36*
atanh(c*x)*a*b**2*c**2*d*e*x + 6*atanh(c*x)*a*b**2*c**2*e**2*x**2 + 36*ata
nh(c*x)*a*b**2*c*d*e - 6*atanh(c*x)*a*b**2*e**2 + 6*atanh(c*x)*b**3*c*e**2
*x + 6*atanh(c*x)*b**3*e**2 + 36*int((atanh(c*x)*x)/(c**2*x**2 - 1),x)*a*b
**2*c**4*d**2 + 12*int((atanh(c*x)*x)/(c**2*x**2 - 1),x)*a*b**2*c**2*e**2
+ 36*int((atanh(c*x)*x)/(c**2*x**2 - 1),x)*b**3*c**3*d*e + 18*int((atanh(c
*x)**2*x)/(c**2*x**2 - 1),x)*b**3*c**4*d**2 + 6*int((atanh(c*x)**2*x)/(c**
2*x**2 - 1),x)*b**3*c**2*e**2 + 18*log(c**2*x - c)*a**2*b*c**2*d**2 + 6*lo
g(c**2*x - c)*a**2*b*e**2 + 36*log(c**2*x - c)*a*b**2*c*d*e + 6*log(c**2*x
- c)*b**3*e**2 + 6*a**3*c**3*d**2*x + 6*a**3*c**3*d*e*x**2 + 2*a**3*c**3*
e**2*x**3 + 18*a**2*b*c**2*d*e*x + 3*a**2*b*c**2*e**2*x**2 + 6*a*b**2*c*e
**2*x)/(6*c**3)
```

3.17 $\int (d + ex)(a + \operatorname{barctanh}(cx))^3 dx$

Optimal result	171
Mathematica [A] (verified)	172
Rubi [A] (verified)	173
Maple [C] (warning: unable to verify)	174
Fricas [F]	174
Sympy [F]	175
Maxima [F]	175
Giac [F]	176
Mupad [F(-1)]	176
Reduce [F]	176

Optimal result

Integrand size = 16, antiderivative size = 244

$$\begin{aligned}
 \int (d + ex)(a + \operatorname{barctanh}(cx))^3 dx = & \frac{3be(a + \operatorname{barctanh}(cx))^2}{2c^2} + \frac{3bex(a + \operatorname{barctanh}(cx))^2}{2c} \\
 & + \frac{d(a + \operatorname{barctanh}(cx))^3}{c} \\
 & - \frac{\left(d^2 + \frac{e^2}{c^2}\right)(a + \operatorname{barctanh}(cx))^3}{2e} \\
 & + \frac{(d + ex)^2(a + \operatorname{barctanh}(cx))^3}{2e} \\
 & - \frac{3b^2e(a + \operatorname{barctanh}(cx)) \log\left(\frac{2}{1-cx}\right)}{c^2} \\
 & - \frac{3bd(a + \operatorname{barctanh}(cx))^2 \log\left(\frac{2}{1-cx}\right)}{c} \\
 & - \frac{3b^3e \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{2c^2} \\
 & - \frac{3b^2d(a + \operatorname{barctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{c} \\
 & + \frac{3b^3d \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx}\right)}{2c}
 \end{aligned}$$

output

```
3/2*b*e*(a+b*arctanh(c*x))^2/c^2+3/2*b*e*x*(a+b*arctanh(c*x))^2/c+d*(a+b*arctanh(c*x))^3/c-1/2*(d^2+e^2/c^2)*(a+b*arctanh(c*x))^3/e+1/2*(e*x+d)^2*(a+b*arctanh(c*x))^3/e-3*b^2*e*(a+b*arctanh(c*x))*ln(2/(-c*x+1))/c^2-3*b*d*(a+b*arctanh(c*x))^2*ln(2/(-c*x+1))/c-3/2*b^3*e*polylog(2,1-2/(-c*x+1))/c^2-3*b^2*d*(a+b*arctanh(c*x))*polylog(2,1-2/(-c*x+1))/c+3/2*b^3*d*polylog(3,1-2/(-c*x+1))/c
```

Mathematica [A] (verified)

Time = 0.67 (sec) , antiderivative size = 331, normalized size of antiderivative = 1.36

$$\int (d + ex)(a + b \operatorname{arctanh}(cx))^3 dx$$

$$= \frac{2a^2c(2acd + 3be)x + 2a^3c^2ex^2 + 6a^2bc^2x(2d + ex)\operatorname{arctanh}(cx) + 3a^2b(2cd + e)\log(1 - cx) + 3a^2b(2cd - e)\log(1 + cx)}{c^3}$$

input

```
Integrate[(d + e*x)*(a + b*ArcTanh[c*x])^3,x]
```

output

```
(2*a^2*c*(2*a*c*d + 3*b*e)*x + 2*a^3*c^2*e*x^2 + 6*a^2*b*c^2*x*(2*d + e*x)*ArcTanh[c*x] + 3*a^2*b*(2*c*d + e)*Log[1 - c*x] + 3*a^2*b*(2*c*d - e)*Log[1 + c*x] + 6*a*b^2*e*(2*c*x*ArcTanh[c*x] + (-1 + c^2*x^2)*ArcTanh[c*x]^2 + Log[1 - c^2*x^2]) - 2*b^3*e*(ArcTanh[c*x]*((3 - 3*c*x)*ArcTanh[c*x] + (1 - c^2*x^2)*ArcTanh[c*x]^2 + 6*Log[1 + E^(-2*ArcTanh[c*x])]) - 3*PolyLog[2, -E^(-2*ArcTanh[c*x])]) + 12*a*b^2*c*d*(ArcTanh[c*x]*((-1 + c*x)*ArcTanh[c*x] - 2*Log[1 + E^(-2*ArcTanh[c*x])]) + PolyLog[2, -E^(-2*ArcTanh[c*x])]) + 4*b^3*c*d*(ArcTanh[c*x]^2*((-1 + c*x)*ArcTanh[c*x] - 3*Log[1 + E^(-2*ArcTanh[c*x])]) + 3*ArcTanh[c*x]*PolyLog[2, -E^(-2*ArcTanh[c*x])]) + (3*PolyLog[3, -E^(-2*ArcTanh[c*x])]))/2)/(4*c^2)
```

Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 259, normalized size of antiderivative = 1.06, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (d + ex)(a + b \operatorname{arctanh}(cx))^3 dx \\
 & \quad \downarrow \text{6480} \\
 & \frac{(d + ex)^2(a + b \operatorname{arctanh}(cx))^3}{2e} - \\
 & \frac{3bc \int \left(\frac{(d^2 c^2 + 2dexc^2 + e^2)(a + b \operatorname{arctanh}(cx))^2}{c^2(1 - c^2 x^2)} - \frac{e^2(a + b \operatorname{arctanh}(cx))^2}{c^2} \right) dx}{2e} \\
 & \quad \downarrow \text{2009} \\
 & \frac{(d + ex)^2(a + b \operatorname{arctanh}(cx))^3}{2e} - \\
 & \frac{3bc \left(-\frac{e^2(a + b \operatorname{arctanh}(cx))^2}{c^3} + \frac{2be^2 \log\left(\frac{2}{1 - cx}\right)(a + b \operatorname{arctanh}(cx))}{c^3} + \frac{2bde \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - cx}\right)(a + b \operatorname{arctanh}(cx))}{c^2} - \frac{2de(a + b \operatorname{arctanh}(cx))}{3bc^2} \right)}{2e}
 \end{aligned}$$

input

```
Int[(d + e*x)*(a + b*ArcTanh[c*x])^3, x]
```

output

```
((d + e*x)^2*(a + b*ArcTanh[c*x])^3)/(2*e) - (3*b*c*(-(e^2*(a + b*ArcTanh[c*x])^2)/c^3) - (e^2*x*(a + b*ArcTanh[c*x])^2)/c^2 - (2*d*e*(a + b*ArcTanh[c*x])^3)/(3*b*c^2) + ((c^2*d^2 + e^2)*(a + b*ArcTanh[c*x])^3)/(3*b*c^3) + (2*b*e^2*(a + b*ArcTanh[c*x])*Log[2/(1 - c*x)])/c^3 + (2*d*e*(a + b*ArcTanh[c*x])^2*Log[2/(1 - c*x)])/c^2 + (b^2*e^2*PolyLog[2, 1 - 2/(1 - c*x)])/c^3 + (2*b*d*e*(a + b*ArcTanh[c*x])*PolyLog[2, 1 - 2/(1 - c*x)])/c^2 - (b^2*d*e*PolyLog[3, 1 - 2/(1 - c*x)])/c^2)/(2*e)
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6480 `Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_)*((d_) + (e_.)*(x_.))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.79 (sec) , antiderivative size = 6347, normalized size of antiderivative = 26.01

method	result	size
parts	Expression too large to display	6347
derivativedivides	Expression too large to display	6355
default	Expression too large to display	6355

input `int((e*x+d)*(a+b*arctanh(c*x))^3,x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F]

$$\int (d + ex)(a + b \operatorname{arctanh}(cx))^3 dx = \int (ex + d)(b \operatorname{arctanh}(cx) + a)^3 dx$$

input `integrate((e*x+d)*(a+b*arctanh(c*x))^3,x, algorithm="fricas")`

output `integral(a^3*e*x + a^3*d + (b^3*e*x + b^3*d)*arctanh(c*x)^3 + 3*(a*b^2*e*x + a*b^2*d)*arctanh(c*x)^2 + 3*(a^2*b*e*x + a^2*b*d)*arctanh(c*x), x)`

Sympy [F]

$$\int (d + ex)(a + b \operatorname{arctanh}(cx))^3 dx = \int (a + b \operatorname{atanh}(cx))^3 (d + ex) dx$$

input `integrate((e*x+d)*(a+b*atanh(c*x))**3,x)`

output `Integral((a + b*atanh(c*x))**3*(d + e*x), x)`

Maxima [F]

$$\int (d + ex)(a + b \operatorname{arctanh}(cx))^3 dx = \int (ex + d)(b \operatorname{artanh}(cx) + a)^3 dx$$

input `integrate((e*x+d)*(a+b*arctanh(c*x))^3,x, algorithm="maxima")`

output `1/2*a^3*e*x^2 + 3/4*(2*x^2*arctanh(c*x) + c*(2*x/c^2 - log(c*x + 1)/c^3 + log(c*x - 1)/c^3))*a^2*b*e + a^3*d*x + 3/2*(2*c*x*arctanh(c*x) + log(-c^2*x^2 + 1))*a^2*b*d/c - 1/16*((b^3*c^2*e*x^2 + 2*b^3*c^2*d*x - (2*c*d + e)*b^3)*log(-c*x + 1)^3 - 3*(2*a*b^2*c^2*e*x^2 + 2*(2*a*b^2*c^2*d + b^3*c*e)*x + (b^3*c^2*e*x^2 + 2*b^3*c^2*d*x + (2*c*d - e)*b^3)*log(c*x + 1))*log(-c*x + 1)^2)/c^2 - integrate(-1/8*((b^3*c^2*e*x^2 - b^3*c*d + (c^2*d - c*e)*b^3*x)*log(c*x + 1)^3 + 6*(a*b^2*c^2*e*x^2 - a*b^2*c*d + (c^2*d - c*e)*a*b^2*x)*log(c*x + 1)^2 - 3*(2*a*b^2*c^2*e*x^2 + (b^3*c^2*e*x^2 - b^3*c*d + (c^2*d - c*e)*b^3*x)*log(c*x + 1)^2 + 2*(2*a*b^2*c^2*d + b^3*c*e)*x - (4*a*b^2*c*d - (2*c*d - e)*b^3 - (4*a*b^2*c^2*e + b^3*c^2*e)*x^2 - 2*(b^3*c^2*d + 2*(c^2*d - c*e)*a*b^2)*x)*log(c*x + 1))/c^2, x)`

Giac [F]

$$\int (d + ex)(a + b \operatorname{arctanh}(cx))^3 dx = \int (ex + d)(b \operatorname{artanh}(cx) + a)^3 dx$$

input `integrate((e*x+d)*(a+b*arctanh(c*x))^3,x, algorithm="giac")`

output `integrate((e*x + d)*(b*arctanh(c*x) + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int (d + ex)(a + b \operatorname{arctanh}(cx))^3 dx = \int (a + b \operatorname{atanh}(cx))^3 (d + ex) dx$$

input `int((a + b*atanh(c*x))^3*(d + e*x),x)`

output `int((a + b*atanh(c*x))^3*(d + e*x), x)`

Reduce [F]

$$\int (d + ex)(a + b \operatorname{arctanh}(cx))^3 dx$$

$$= \frac{2 \operatorname{atanh}(cx)^3 b^3 c^2 dx + \operatorname{atanh}(cx)^3 b^3 c^2 e x^2 - \operatorname{atanh}(cx)^3 b^3 e + 6 \operatorname{atanh}(cx)^2 a b^2 c^2 dx + 3 \operatorname{atanh}(cx)^2 a b^2 c^2 e x}{1}$$

input `int((e*x+d)*(a+b*atanh(c*x))^3,x)`

output

```
(2*atanh(c*x)**3*b**3*c**2*d*x + atanh(c*x)**3*b**3*c**2*e*x**2 - atanh(c*
x)**3*b**3*e + 6*atanh(c*x)**2*a*b**2*c**2*d*x + 3*atanh(c*x)**2*a*b**2*c*
**2*e*x**2 - 3*atanh(c*x)**2*a*b**2*e + 3*atanh(c*x)**2*b**3*c*e*x + 6*atan
h(c*x)*a**2*b*c**2*d*x + 3*atanh(c*x)*a**2*b*c**2*e*x**2 + 6*atanh(c*x)*a*
**2*b*c*d - 3*atanh(c*x)*a**2*b*e + 6*atanh(c*x)*a*b**2*c*e*x + 6*atanh(c*x
)*a*b**2*e + 12*int((atanh(c*x)*x)/(c**2*x**2 - 1),x)*a*b**2*c**3*d + 6*in
t((atanh(c*x)*x)/(c**2*x**2 - 1),x)*b**3*c**2*e + 6*int((atanh(c*x)**2*x)/
(c**2*x**2 - 1),x)*b**3*c**3*d + 6*log(c**2*x - c)*a**2*b*c*d + 6*log(c**2
*x - c)*a*b**2*e + 2*a**3*c**2*d*x + a**3*c**2*e*x**2 + 3*a**2*b*c*e*x)/(2
*c**2)
```

3.18 $\int \frac{(a+b\operatorname{arctanh}(cx))^3}{d+ex} dx$

Optimal result	178
Mathematica [C] (warning: unable to verify)	179
Rubi [A] (verified)	180
Maple [C] (warning: unable to verify)	181
Fricas [F]	182
Sympy [F]	183
Maxima [F]	183
Giac [F]	183
Mupad [F(-1)]	184
Reduce [F]	184

Optimal result

Integrand size = 18, antiderivative size = 272

$$\begin{aligned}
 \int \frac{(a + b\operatorname{arctanh}(cx))^3}{d + ex} dx = & -\frac{(a + b\operatorname{arctanh}(cx))^3 \log\left(\frac{2}{1+cx}\right)}{e} \\
 & + \frac{(a + b\operatorname{arctanh}(cx))^3 \log\left(\frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{e} \\
 & + \frac{3b(a + b\operatorname{arctanh}(cx))^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+cx}\right)}{2e} \\
 & - \frac{3b(a + b\operatorname{arctanh}(cx))^2 \operatorname{PolyLog}\left(2, 1 - \frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{2e} \\
 & + \frac{3b^2(a + b\operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1+cx}\right)}{2e} \\
 & - \frac{3b^2(a + b\operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(3, 1 - \frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{2e} \\
 & + \frac{3b^3 \operatorname{PolyLog}\left(4, 1 - \frac{2}{1+cx}\right)}{4e} \\
 & - \frac{3b^3 \operatorname{PolyLog}\left(4, 1 - \frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{4e}
 \end{aligned}$$

output

```

-(a+b*arctanh(c*x))^3*ln(2/(c*x+1))/e+(a+b*arctanh(c*x))^3*ln(2*c*(e*x+d)/
(c*d+e)/(c*x+1))/e+3/2*b*(a+b*arctanh(c*x))^2*polylog(2,1-2/(c*x+1))/e-3/2
*b*(a+b*arctanh(c*x))^2*polylog(2,1-2*c*(e*x+d)/(c*d+e)/(c*x+1))/e+3/2*b^2
*(a+b*arctanh(c*x))*polylog(3,1-2/(c*x+1))/e-3/2*b^2*(a+b*arctanh(c*x))*po
lylog(3,1-2*c*(e*x+d)/(c*d+e)/(c*x+1))/e+3/4*b^3*polylog(4,1-2/(c*x+1))/e-
3/4*b^3*polylog(4,1-2*c*(e*x+d)/(c*d+e)/(c*x+1))/e

```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 26.72 (sec) , antiderivative size = 2160, normalized size of antiderivative = 7.94

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{d + ex} dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*ArcTanh[c*x])^3/(d + e*x), x]
```

output

```

(a^3*Log[d + e*x])/e - ((3*I)*a^2*b*(I*ArcTanh[c*x]*(-Log[1/Sqrt[1 - c^2*x
^2]] + Log[I*Sinh[ArcTanh[(c*d)/e] + ArcTanh[c*x]]]) + ((-I)*(I*ArcTanh[(c
*d)/e] + I*ArcTanh[c*x])^2 - (I/4)*(Pi - (2*I)*ArcTanh[c*x])^2 + 2*(I*ArcT
anh[(c*d)/e] + I*ArcTanh[c*x])*Log[1 - E^((2*I)*(I*ArcTanh[(c*d)/e] + I*Ar
cTanh[c*x]))] + (Pi - (2*I)*ArcTanh[c*x])*Log[1 - E^(I*(Pi - (2*I)*ArcTan
h[c*x]))] - (Pi - (2*I)*ArcTanh[c*x])*Log[2*Sin[(Pi - (2*I)*ArcTanh[c*x])/2
]] - 2*(I*ArcTanh[(c*d)/e] + I*ArcTanh[c*x])*Log[(2*I)*Sinh[ArcTanh[(c*d)/
e] + ArcTanh[c*x]]) - I*PolyLog[2, E^((2*I)*(I*ArcTanh[(c*d)/e] + I*ArcTan
h[c*x]))] - I*PolyLog[2, E^(I*(Pi - (2*I)*ArcTanh[c*x]))]/2))/e + (a*b^2*
(-8*c*d*ArcTanh[c*x]^3 + 4*e*ArcTanh[c*x]^3 - (4*Sqrt[1 - (c^2*d^2)/e^2]*e
*ArcTanh[c*x]^3)/E^ArcTanh[(c*d)/e] - 6*c*d*ArcTanh[c*x]^2*Log[1 + E^(-2*A
rcTanh[c*x])] - 6*c*d*ArcTanh[c*x]^2*Log[1 - (Sqrt[c*d + e]*E^ArcTanh[c*x]
)/Sqrt[-(c*d) + e]] - 6*c*d*ArcTanh[c*x]^2*Log[1 + (Sqrt[c*d + e]*E^ArcTan
h[c*x])/Sqrt[-(c*d) + e]] - (6*I)*c*d*Pi*ArcTanh[c*x]*Log[(1 + E^(2*ArcTan
h[c*x]))/(2*E^ArcTanh[c*x])] + 6*c*d*ArcTanh[c*x]^2*Log[1 - E^(ArcTanh[(c*
d)/e] + ArcTanh[c*x])] + 6*c*d*ArcTanh[c*x]^2*Log[1 + E^(ArcTanh[(c*d)/e]
+ ArcTanh[c*x])] + 6*c*d*ArcTanh[c*x]^2*Log[1 - E^(2*(ArcTanh[(c*d)/e] + A
rcTanh[c*x]))] + 12*c*d*ArcTanh[(c*d)/e]*ArcTanh[c*x]*Log[(I/2)*E^(-ArcTan
h[(c*d)/e] - ArcTanh[c*x])*(-1 + E^(2*(ArcTanh[(c*d)/e] + ArcTanh[c*x])))]
+ 6*c*d*ArcTanh[c*x]^2*Log[(e*(-1 + E^(2*ArcTanh[c*x])) + c*d*(1 + E(...

```

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 272, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {6476}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{barctanh}(cx))^3}{d + ex} dx$$

↓ 6476

$$\begin{aligned} & -\frac{3b^2(a + \operatorname{barctanh}(cx)) \operatorname{PolyLog}\left(3, 1 - \frac{2c(d+ex)}{(cd+e)(cx+1)}\right)}{2e} + \\ & \frac{3b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{cx+1}\right) (a + \operatorname{barctanh}(cx))}{2e} - \\ & \frac{3b(a + \operatorname{barctanh}(cx))^2 \operatorname{PolyLog}\left(2, 1 - \frac{2c(d+ex)}{(cd+e)(cx+1)}\right)}{2e} + \frac{(a + \operatorname{barctanh}(cx))^3 \log\left(\frac{2c(d+ex)}{(cx+1)(cd+e)}\right)}{e} + \\ & \frac{3b \operatorname{PolyLog}\left(2, 1 - \frac{2}{cx+1}\right) (a + \operatorname{barctanh}(cx))^2}{2e} - \frac{\log\left(\frac{2}{cx+1}\right) (a + \operatorname{barctanh}(cx))^3}{e} - \\ & \frac{3b^3 \operatorname{PolyLog}\left(4, 1 - \frac{2c(d+ex)}{(cd+e)(cx+1)}\right)}{4e} + \frac{3b^3 \operatorname{PolyLog}\left(4, 1 - \frac{2}{cx+1}\right)}{4e} \end{aligned}$$

input

```
Int[(a + b*ArcTanh[c*x])^3/(d + e*x), x]
```

output

```
-(((a + b*ArcTanh[c*x])^3*Log[2/(1 + c*x)])/e) + ((a + b*ArcTanh[c*x])^3*Log[(2*c*(d + e*x))/((c*d + e)*(1 + c*x))])/e + (3*b*(a + b*ArcTanh[c*x])^2*PolyLog[2, 1 - 2/(1 + c*x)]/(2*e) - (3*b*(a + b*ArcTanh[c*x])^2*PolyLog[2, 1 - (2*c*(d + e*x))/((c*d + e)*(1 + c*x))]/(2*e) + (3*b^2*(a + b*ArcTanh[c*x])*PolyLog[3, 1 - 2/(1 + c*x)]/(2*e) - (3*b^2*(a + b*ArcTanh[c*x])*PolyLog[3, 1 - (2*c*(d + e*x))/((c*d + e)*(1 + c*x))]/(2*e) + (3*b^3*PolyLog[4, 1 - 2/(1 + c*x)]/(4*e) - (3*b^3*PolyLog[4, 1 - (2*c*(d + e*x))/((c*d + e)*(1 + c*x))]/(4*e)
```

Definitions of rubi rules used

rule 6476

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^3/((d_) + (e_.)*(x_.)), x_Symbol] :=
  Simp[(-(a + b*ArcTanh[c*x])^3)*(Log[2/(1 + c*x)]/e), x] + (Simp[(a + b*Arc
  Tanh[c*x])^3*(Log[2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/e), x] + Simp[3*b*
  (a + b*ArcTanh[c*x])^2*(PolyLog[2, 1 - 2/(1 + c*x)]/(2*e)), x] - Simp[3*b*(
  a + b*ArcTanh[c*x])^2*(PolyLog[2, 1 - 2*c*((d + e*x)/((c*d + e)*(1 + c*x))
  ]/(2*e)), x] + Simp[3*b^2*(a + b*ArcTanh[c*x])*(PolyLog[3, 1 - 2/(1 + c*x)
  ]/(2*e)), x] - Simp[3*b^2*(a + b*ArcTanh[c*x])*(PolyLog[3, 1 - 2*c*((d + e*x
  )/((c*d + e)*(1 + c*x)))]/(2*e)), x] + Simp[3*b^3*(PolyLog[4, 1 - 2/(1 + c*
  x)]/(4*e)), x] - Simp[3*b^3*(PolyLog[4, 1 - 2*c*((d + e*x)/((c*d + e)*(1 +
  c*x)))]/(4*e)), x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 - e^2, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.86 (sec) , antiderivative size = 2160, normalized size of antiderivative = 7.94

method	result	size
derivativdivides	Expression too large to display	2160
default	Expression too large to display	2160
parts	Expression too large to display	2164

input

```
int((a+b*arctanh(c*x))^3/(e*x+d),x,method=_RETURNVERBOSE)
```

output

```

1/c*(a^3*c*ln(c*e*x+c*d)/e+b^3*c*(ln(c*e*x+c*d)/e*arctanh(c*x)^3-3/e*(1/3*
arctanh(c*x)^3*ln(d*c*(1+(c*x+1)^2/(-c^2*x^2+1))+e*((c*x+1)^2/(-c^2*x^2+1)
-1))-1/6*I*Pi*csgn(I*(d*c*(1-(c*x+1)^2/(c^2*x^2-1))+e*(-(c*x+1)^2/(c^2*x^2
-1)-1))/(1-(c*x+1)^2/(c^2*x^2-1)))*(csgn(I*(d*c*(1-(c*x+1)^2/(c^2*x^2-1))+
e*(-(c*x+1)^2/(c^2*x^2-1)-1)))*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))-csgn(I*(d
*c*(1-(c*x+1)^2/(c^2*x^2-1))+e*(-(c*x+1)^2/(c^2*x^2-1)-1))/(1-(c*x+1)^2/(c
^2*x^2-1)))*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))-csgn(I*(d*c*(1-(c*x+1)^2/(c^
2*x^2-1))+e*(-(c*x+1)^2/(c^2*x^2-1)-1)))*csgn(I*(d*c*(1-(c*x+1)^2/(c^2*x^2
-1))+e*(-(c*x+1)^2/(c^2*x^2-1)-1))/(1-(c*x+1)^2/(c^2*x^2-1)))+csgn(I*(d*c*
(1-(c*x+1)^2/(c^2*x^2-1))+e*(-(c*x+1)^2/(c^2*x^2-1)-1))/(1-(c*x+1)^2/(c^2*
x^2-1)))^2)*arctanh(c*x)^3+1/2*arctanh(c*x)^2*polylog(2,-(c*x+1)^2/(-c^2*x
^2+1))-1/2*arctanh(c*x)*polylog(3,-(c*x+1)^2/(-c^2*x^2+1))+1/4*polylog(4,-
(c*x+1)^2/(-c^2*x^2+1))-1/3*e/(c*d+e)*arctanh(c*x)^3*ln(1-(c*d+e)*(c*x+1)^
2/(-c^2*x^2+1)/(-c*d+e))-1/2*e/(c*d+e)*arctanh(c*x)^2*polylog(2,(c*d+e)*(c
*x+1)^2/(-c^2*x^2+1)/(-c*d+e))+1/2*e/(c*d+e)*arctanh(c*x)*polylog(3,(c*d+e
)*(c*x+1)^2/(-c^2*x^2+1)/(-c*d+e))-1/4*e/(c*d+e)*polylog(4,(c*d+e)*(c*x+1)
^2/(-c^2*x^2+1)/(-c*d+e))-1/3*d*c/(c*d+e)*arctanh(c*x)^3*ln(1-(c*d+e)*(c*x
+1)^2/(-c^2*x^2+1)/(-c*d+e))-1/2*d*c/(c*d+e)*arctanh(c*x)^2*polylog(2,(c*d
+e)*(c*x+1)^2/(-c^2*x^2+1)/(-c*d+e))+1/2*d*c/(c*d+e)*arctanh(c*x)*polylog(
3,(c*d+e)*(c*x+1)^2/(-c^2*x^2+1)/(-c*d+e))-1/4*d*c/(c*d+e)*polylog(4,(c...

```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{d + ex} dx = \int \frac{(b \operatorname{arctanh}(cx) + a)^3}{ex + d} dx$$

input

```
integrate((a+b*arctanh(c*x))^3/(e*x+d),x, algorithm="fricas")
```

output

```
integral((b^3*arctanh(c*x)^3 + 3*a*b^2*arctanh(c*x)^2 + 3*a^2*b*arctanh(c*
x) + a^3)/(e*x + d), x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{d + ex} dx = \int \frac{(a + b \operatorname{atanh}(cx))^3}{d + ex} dx$$

input `integrate((a+b*atanh(c*x))**3/(e*x+d), x)`

output `Integral((a + b*atanh(c*x))**3/(d + e*x), x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{d + ex} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x))^3/(e*x+d), x, algorithm="maxima")`

output `a^3*log(e*x + d)/e + integrate(1/8*b^3*(log(c*x + 1) - log(-c*x + 1))^3/(e*x + d) + 3/4*a*b^2*(log(c*x + 1) - log(-c*x + 1))^2/(e*x + d) + 3/2*a^2*b*(log(c*x + 1) - log(-c*x + 1))/(e*x + d), x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{d + ex} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x))^3/(e*x+d), x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^3/(e*x + d), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{d + ex} dx = \int \frac{(a + b \operatorname{atanh}(cx))^3}{d + ex} dx$$

input `int((a + b*atanh(c*x))^3/(d + e*x),x)`output `int((a + b*atanh(c*x))^3/(d + e*x), x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{d + ex} dx$$

$$= \frac{3 \left(\int \frac{\operatorname{atanh}(cx)}{ex+d} dx \right) a^2 b e + \left(\int \frac{\operatorname{atanh}(cx)^3}{ex+d} dx \right) b^3 e + 3 \left(\int \frac{\operatorname{atanh}(cx)^2}{ex+d} dx \right) a b^2 e + \log(ex + d) a^3}{e}$$

input `int((a+b*atanh(c*x))^3/(e*x+d),x)`output `(3*int(atanh(c*x)/(d + e*x),x)*a**2*b*e + int(atanh(c*x)**3/(d + e*x),x)*b**3*e + 3*int(atanh(c*x)**2/(d + e*x),x)*a*b**2*e + log(d + e*x)*a**3)/e`

$$3.19 \quad \int \frac{(a+b \operatorname{arctanh}(cx))^3}{(d+ex)^2} dx$$

Optimal result	186
Mathematica [C] (warning: unable to verify)	187
Rubi [A] (verified)	188
Maple [C] (warning: unable to verify)	189
Fricas [F]	190
Sympy [F]	191
Maxima [F]	191
Giac [F]	192
Mupad [F(-1)]	192
Reduce [F]	192

Optimal result

Integrand size = 18, antiderivative size = 517

$$\begin{aligned}
 \int \frac{(a + \operatorname{barctanh}(cx))^3}{(d + ex)^2} dx = & -\frac{(a + \operatorname{barctanh}(cx))^3}{e(d + ex)} + \frac{3bc(a + \operatorname{barctanh}(cx))^2 \log\left(\frac{2}{1-cx}\right)}{2e(cd + e)} \\
 & - \frac{3bc(a + \operatorname{barctanh}(cx))^2 \log\left(\frac{2}{1+cx}\right)}{2(cd - e)e} \\
 & + \frac{3bc(a + \operatorname{barctanh}(cx))^2 \log\left(\frac{2}{1+cx}\right)}{c^2d^2 - e^2} \\
 & - \frac{3bc(a + \operatorname{barctanh}(cx))^2 \log\left(\frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{c^2d^2 - e^2} \\
 & + \frac{3b^2c(a + \operatorname{barctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right)}{2e(cd + e)} \\
 & + \frac{3b^2c(a + \operatorname{barctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+cx}\right)}{2(cd - e)e} \\
 & - \frac{3b^2c(a + \operatorname{barctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+cx}\right)}{c^2d^2 - e^2} \\
 & + \frac{3b^2c(a + \operatorname{barctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{c^2d^2 - e^2} \\
 & - \frac{3b^3c \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-cx}\right)}{4e(cd + e)} + \frac{3b^3c \operatorname{PolyLog}\left(3, 1 - \frac{2}{1+cx}\right)}{4(cd - e)e} \\
 & - \frac{3b^3c \operatorname{PolyLog}\left(3, 1 - \frac{2}{1+cx}\right)}{2(c^2d^2 - e^2)} \\
 & + \frac{3b^3c \operatorname{PolyLog}\left(3, 1 - \frac{2c(d+ex)}{(cd+e)(1+cx)}\right)}{2(c^2d^2 - e^2)}
 \end{aligned}$$

output

```

-(a+b*arctanh(c*x))^3/e/(e*x+d)+3/2*b*c*(a+b*arctanh(c*x))^2*ln(2/(-c*x+1))
)/e/(c*d+e)-3/2*b*c*(a+b*arctanh(c*x))^2*ln(2/(c*x+1))/(c*d-e)/e+3*b*c*(a
+b*arctanh(c*x))^2*ln(2/(c*x+1))/(c^2*d^2-e^2)-3*b*c*(a+b*arctanh(c*x))^2*l
n(2*c*(e*x+d)/(c*d+e)/(c*x+1))/(c^2*d^2-e^2)+3/2*b^2*c*(a+b*arctanh(c*x))*
polylog(2,1-2/(-c*x+1))/e/(c*d+e)+3/2*b^2*c*(a+b*arctanh(c*x))*polylog(2,1
-2/(c*x+1))/(c*d-e)/e-3*b^2*c*(a+b*arctanh(c*x))*polylog(2,1-2/(c*x+1))/(c
^2*d^2-e^2)+3*b^2*c*(a+b*arctanh(c*x))*polylog(2,1-2*c*(e*x+d)/(c*d+e)/(c*
x+1))/(c^2*d^2-e^2)-3/4*b^3*c*polylog(3,1-2/(-c*x+1))/e/(c*d+e)+3/4*b^3*c*
polylog(3,1-2/(c*x+1))/(c*d-e)/e-3*b^3*c*polylog(3,1-2/(c*x+1))/(2*c^2*d^2
-2*e^2)+3*b^3*c*polylog(3,1-2*c*(e*x+d)/(c*d+e)/(c*x+1))/(2*c^2*d^2-2*e^2)

```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 11.53 (sec) , antiderivative size = 1107, normalized size of antiderivative = 2.14

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^2} dx = \text{Too large to display}$$

input `Integrate[(a + b*ArcTanh[c*x])^3/(d + e*x)^2,x]`

output

```

-(a^3/(e*(d + e*x))) - (3*a^2*b*ArcTanh[c*x])/(e*(d + e*x)) - (3*a^2*b*c*Log[1 - c*x])/(2*e*(c*d + e)) + (3*a^2*b*c*Log[1 + c*x])/(2*c*d*e - 2*e^2) - (3*a^2*b*c*Log[d + e*x])/(c^2*d^2 - e^2) + (3*a*b^2*(-(ArcTanh[c*x]^2/(Sqrt[1 - (c^2*d^2)/e^2]*e*E^ArcTanh[(c*d)/e])) + (x*ArcTanh[c*x]^2)/(d + e*x) + (c*d*(I*Pi*Log[1 + E^(2*ArcTanh[c*x])]) - 2*ArcTanh[c*x]*Log[1 - E^(-2*(ArcTanh[(c*d)/e] + ArcTanh[c*x]))] - I*Pi*(ArcTanh[c*x] - Log[1 - c^2*x^2]/2) - 2*ArcTanh[(c*d)/e]*(ArcTanh[c*x] + Log[1 - E^(-2*(ArcTanh[(c*d)/e] + ArcTanh[c*x]))] - Log[I*Sinh[ArcTanh[(c*d)/e] + ArcTanh[c*x]]]) + PolyLog[2, E^(-2*(ArcTanh[(c*d)/e] + ArcTanh[c*x]))]))/(c^2*d^2 - e^2))/d + (b^3*((x*ArcTanh[c*x]^3)/(d + e*x) + (3*(6*c*d*ArcTanh[c*x]^3 - 2*e*ArcTanh[c*x]^3 + (4*Sqrt[1 - (c^2*d^2)/e^2]*e*ArcTanh[c*x]^3)/E^ArcTanh[(c*d)/e] + (6*I)*c*d*Pi*ArcTanh[c*x]*Log[(E^(-ArcTanh[c*x]) + E^ArcTanh[c*x])/2] + 6*c*d*ArcTanh[c*x]^2*Log[1 - (Sqrt[c*d + e]*E^ArcTanh[c*x])/Sqrt[-(c*d) + e]] + 6*c*d*ArcTanh[c*x]^2*Log[1 + (Sqrt[c*d + e]*E^ArcTanh[c*x])/Sqrt[-(c*d) + e]] - 6*c*d*ArcTanh[c*x]^2*Log[1 - E^(ArcTanh[(c*d)/e] + ArcTanh[c*x])] - 6*c*d*ArcTanh[c*x]^2*Log[1 + E^(ArcTanh[(c*d)/e] + ArcTanh[c*x])] - 6*c*d*ArcTanh[c*x]^2*Log[1 - E^(2*(ArcTanh[(c*d)/e] + ArcTanh[c*x]))] - 12*c*d*ArcTanh[(c*d)/e]*ArcTanh[c*x]*Log[(I/2)*E^(-ArcTanh[(c*d)/e] - ArcTanh[c*x])*(-1 + E^(2*(ArcTanh[(c*d)/e] + ArcTanh[c*x]))]) - 6*c*d*ArcTanh[c*x]^2*Log[(e*(-1 + E^(2*ArcTanh[c*x])) + c*d*(1 + E^(2*ArcTanh[c*x])))]/(2...

```

Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 487, normalized size of antiderivative = 0.94, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^2} dx$$

↓ 6480

$$\frac{3bc \int \left(\frac{c(a+b \operatorname{arctanh}(cx))^2}{2(cd+e)(1-cx)} + \frac{c(a+b \operatorname{arctanh}(cx))^2}{2(cd-e)(cx+1)} - \frac{e^2(a+b \operatorname{arctanh}(cx))^2}{(cd-e)(cd+e)(d+ex)} \right) dx}{e} - \frac{(a + b \operatorname{arctanh}(cx))^3}{e(d + ex)}$$

↓ 2009

$$\frac{3bc \left(-\frac{be \operatorname{PolyLog}\left(2, 1 - \frac{2}{cx+1}\right)(a+b \operatorname{arctanh}(cx))}{c^2 d^2 - e^2} + \frac{be(a+b \operatorname{arctanh}(cx)) \operatorname{PolyLog}\left(2, 1 - \frac{2c(d+ex)}{(cd+e)(cx+1)}\right)}{c^2 d^2 - e^2} + \frac{e \log\left(\frac{2}{cx+1}\right)(a+b \operatorname{arctanh}(cx))}{c^2 d^2 - e^2} \right)}{e} - \frac{(a + b \operatorname{arctanh}(cx))^3}{e(d + ex)}$$

input

```
Int[(a + b*ArcTanh[c*x])^3/(d + e*x)^2,x]
```

output

```

-((a + b*ArcTanh[c*x])^3/(e*(d + e*x))) + (3*b*c*((a + b*ArcTanh[c*x])^2*
Log[2/(1 - c*x)]/(2*(c*d + e)) - ((a + b*ArcTanh[c*x])^2*Log[2/(1 + c*x)]
)/(2*(c*d - e)) + (e*(a + b*ArcTanh[c*x])^2*Log[2/(1 + c*x)]/(c^2*d^2 - e
^2) - (e*(a + b*ArcTanh[c*x])^2*Log[(2*c*(d + e*x))/((c*d + e)*(1 + c*x))]
)/(c^2*d^2 - e^2) + (b*(a + b*ArcTanh[c*x])*PolyLog[2, 1 - 2/(1 - c*x)]/(
2*(c*d + e)) + (b*(a + b*ArcTanh[c*x])*PolyLog[2, 1 - 2/(1 + c*x)]/(2*(c*
d - e)) - (b*e*(a + b*ArcTanh[c*x])*PolyLog[2, 1 - 2/(1 + c*x)]/(c^2*d^2
- e^2) + (b*e*(a + b*ArcTanh[c*x])*PolyLog[2, 1 - (2*c*(d + e*x))/((c*d +
e)*(1 + c*x))]/(c^2*d^2 - e^2) - (b^2*PolyLog[3, 1 - 2/(1 - c*x)]/(4*(c*
d + e)) + (b^2*PolyLog[3, 1 - 2/(1 + c*x)]/(4*(c*d - e)) - (b^2*e*PolyLog
[3, 1 - 2/(1 + c*x)]/(2*(c^2*d^2 - e^2)) + (b^2*e*PolyLog[3, 1 - (2*c*(d
+ e*x))/((c*d + e)*(1 + c*x))]/(2*(c^2*d^2 - e^2)))))/e

```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6480 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_)*((d_) + (e_.)*(x_))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.63 (sec) , antiderivative size = 3154, normalized size of antiderivative = 6.10

method	result	size
derivativedivides	Expression too large to display	3154
default	Expression too large to display	3154
parts	Expression too large to display	3162

input `int((a+b*arctanh(c*x))^3/(e*x+d)^2,x,method=_RETURNVERBOSE)`

output

```

1/c*(-a^3*c^2/(c*e*x+c*d)/e+b^3*c^2*(-1/(c*e*x+c*d)/e*arctanh(c*x)^3+3/e*
-1/2*I/(c*d+e)/(c*d-e)*Pi*e*arctanh(c*x)^2+1/3/(c*d-e)*arctanh(c*x)^3-e/(c
*d+e)^2/(c*d-e)*d*c*arctanh(c*x)^2*ln(1-(c*d+e)*(c*x+1)^2/(-c^2*x^2+1)/(-c
*d+e))-1/2*I/(c*d+e)/(c*d-e)*Pi*csgn(I*(d*c*(1-(c*x+1)^2/(c^2*x^2-1))+e*(-
(c*x+1)^2/(c^2*x^2-1)-1))/(1-(c*x+1)^2/(c^2*x^2-1)))^3*e*arctanh(c*x)^2+1/
2*I/(c*d+e)/(c*d-e)*Pi*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))^2*e*arctanh(c*x)^
2+1/2*I/(c*d+e)/(c*d-e)*Pi*c*d*arctanh(c*x)^2-1/2*I/(c*d+e)/(c*d-e)*Pi*csg
n(I/(1-(c*x+1)^2/(c^2*x^2-1)))^3*e*arctanh(c*x)^2-e/(c*d+e)^2/(c*d-e)*d*c*
arctanh(c*x)*polylog(2,(c*d+e)*(c*x+1)^2/(-c^2*x^2+1)/(-c*d+e))-1/4*I/(c*d
+e)/(c*d-e)*Pi*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))^3*e
*arctanh(c*x)^2-1/4*I/(c*d+e)/(c*d-e)*Pi*csgn(I*(c*x+1)^2/(c^2*x^2-1))^3*e
*arctanh(c*x)^2+1/4*I/(c*d+e)/(c*d-e)*Pi*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(
c*x+1)^2/(c^2*x^2-1)))*csgn(I/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(c*x+1)^2/
(c^2*x^2-1))*c*d*arctanh(c*x)^2-arctanh(c*x)^2*e/(c*d+e)/(c*d-e)*ln(c*e*x+
c*d)+1/2*e^2/(c*d+e)^2/(c*d-e)*polylog(3,(c*d+e)*(c*x+1)^2/(-c^2*x^2+1)/(-
c*d+e))+arctanh(c*x)^2/(2*c*d-2*e)*ln(c*x+1)-arctanh(c*x)^2/(2*c*d+2*e)*ln
(c*x-1)-arctanh(c*x)^2/(c*d-e)*ln((c*x+1)/(-c^2*x^2+1)^(1/2))+1/4*I/(c*d+e
)/(c*d-e)*Pi*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(
I/(1-(c*x+1)^2/(c^2*x^2-1)))*csgn(I*(c*x+1)^2/(c^2*x^2-1))*e*arctanh(c*x)^
2+1/4*I/(c*d+e)/(c*d-e)*Pi*csgn(I*(c*x+1)^2/(c^2*x^2-1)/(1-(c*x+1)^2/(c...

```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^2} dx = \int \frac{(b \operatorname{arctanh}(cx) + a)^3}{(ex + d)^2} dx$$

input

```
integrate((a+b*arctanh(c*x))^3/(e*x+d)^2,x, algorithm="fricas")
```

output

```
integral((b^3*arctanh(c*x)^3 + 3*a*b^2*arctanh(c*x)^2 + 3*a^2*b*arctanh(c*
x) + a^3)/(e^2*x^2 + 2*d*e*x + d^2), x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^2} dx = \int \frac{(a + b \operatorname{atanh}(cx))^3}{(d + ex)^2} dx$$

input `integrate((a+b*atanh(c*x))**3/(e*x+d)**2,x)`

output `Integral((a + b*atanh(c*x))**3/(d + e*x)**2, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^2} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{(ex + d)^2} dx$$

input `integrate((a+b*arctanh(c*x))^3/(e*x+d)^2,x, algorithm="maxima")`

output `3/2*(c*(log(c*x + 1)/(c*d*e - e^2) - log(c*x - 1)/(c*d*e + e^2) - 2*log(e*x + d)/(c^2*d^2 - e^2)) - 2*arctanh(c*x)/(e^2*x + d*e))*a^2*b - a^3/(e^2*x + d*e) - 1/8*(((c^2*d*e - c*e^2)*b^3*x - (c*d*e - e^2)*b^3)*log(-c*x + 1)^3 + 3*(2*(c^2*d^2 - e^2)*a*b^2 - ((c^2*d*e + c*e^2)*b^3*x + (c*d*e + e^2)*b^3)*log(c*x + 1))*log(-c*x + 1)^2)/(c^2*d^3*e - d*e^3 + (c^2*d^2*e^2 - e^4)*x) - integrate(1/8*(((c^2*d*e - c*e^2)*b^3*x - (c*d*e - e^2)*b^3)*log(c*x + 1)^3 + 6*((c^2*d*e - c*e^2)*a*b^2*x - (c*d*e - e^2)*a*b^2)*log(c*x + 1)^2 + 3*(4*(c^2*d*e - c*e^2)*a*b^2*x + 4*(c^2*d^2 - c*d*e)*a*b^2 - ((c^2*d*e - c*e^2)*b^3*x - (c*d*e - e^2)*b^3)*log(c*x + 1)^2 - 2*(b^3*c^2*e^2*x^2 + b^3*c*d*e - 2*(c*d*e - e^2)*a*b^2 + (2*(c^2*d*e - c*e^2)*a*b^2 + (c^2*d*e + c*e^2)*b^3)*x)*log(c*x + 1))*log(-c*x + 1))/(c*d^3*e - d^2*e^2 - (c^2*d*e^3 - c*e^4)*x^3 - (2*c^2*d^2*e^2 - 3*c*d*e^3 + e^4)*x^2 - (c^2*d^3*e - 3*c*d^2*e^2 + 2*d*e^3)*x), x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^2} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{(ex + d)^2} dx$$

input `integrate((a+b*arctanh(c*x))^3/(e*x+d)^2,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^3/(e*x + d)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^2} dx = \int \frac{(a + b \operatorname{atanh}(cx))^3}{(d + ex)^2} dx$$

input `int((a + b*atanh(c*x))^3/(d + e*x)^2,x)`

output `int((a + b*atanh(c*x))^3/(d + e*x)^2, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^2} dx = \text{too large to display}$$

input `int((a+b*atanh(c*x))^3/(e*x+d)^2,x)`

output

```

(4*atanh(c*x)**3*b**3*c**6*d**6*x - 4*atanh(c*x)**3*b**3*c**4*d**5*e - 4*a
tanh(c*x)**3*b**3*c**4*d**4*e**2*x + 4*atanh(c*x)**3*b**3*c**2*d**3*e**3 +
  12*atanh(c*x)**2*a*b**2*c**6*d**6*x - 12*atanh(c*x)**2*a*b**2*c**4*d**5*e
- 12*atanh(c*x)**2*a*b**2*c**4*d**4*e**2*x + 12*atanh(c*x)**2*a*b**2*c**2
*d**3*e**3 - 3*atanh(c*x)**2*b**3*c**5*d**6 + 9*atanh(c*x)**2*b**3*c**5*d*
*5*e*x - 12*atanh(c*x)**2*b**3*c**3*d**3*e**3*x + 3*atanh(c*x)**2*b**3*c*d
**2*e**4 + 3*atanh(c*x)**2*b**3*c*d*e**5*x + 12*atanh(c*x)*a**2*b*c**6*d**
6*x - 12*atanh(c*x)*a**2*b*c**2*d**2*e**4*x + 24*atanh(c*x)*a*b**2*c**5*d*
*5*e*x - 24*atanh(c*x)*a*b**2*c**3*d**3*e**3*x + 18*atanh(c*x)*b**3*c**4*d
**4*e**2*x - 24*atanh(c*x)*b**3*c**2*d**2*e**4*x + 6*atanh(c*x)*b**3*e**6*
x - 6*int(atanh(c*x)/(c**4*d**4*x**2 + 2*c**4*d**3*e*x**3 + c**4*d**2*e**2
*x**4 - c**2*d**4 - 2*c**2*d**3*e*x + 2*c**2*d*e**3*x**3 + c**2*e**4*x**4
- d**2*e**2 - 2*d*e**3*x - e**4*x**2),x)*b**3*c**8*d**10 - 6*int(atanh(c*x
)/(c**4*d**4*x**2 + 2*c**4*d**3*e*x**3 + c**4*d**2*e**2*x**4 - c**2*d**4 -
2*c**2*d**3*e*x + 2*c**2*d*e**3*x**3 + c**2*e**4*x**4 - d**2*e**2 - 2*d*e
**3*x - e**4*x**2),x)*b**3*c**8*d**9*e*x + 12*int(atanh(c*x)/(c**4*d**4*x*
*2 + 2*c**4*d**3*e*x**3 + c**4*d**2*e**2*x**4 - c**2*d**4 - 2*c**2*d**3*e*
x + 2*c**2*d*e**3*x**3 + c**2*e**4*x**4 - d**2*e**2 - 2*d*e**3*x - e**4*x*
*2),x)*b**3*c**6*d**8*e**2 + 12*int(atanh(c*x)/(c**4*d**4*x**2 + 2*c**4*d*
*3*e*x**3 + c**4*d**2*e**2*x**4 - c**2*d**4 - 2*c**2*d**3*e*x + 2*c**2*...

```

$$3.20 \quad \int \frac{(a+b\operatorname{arctanh}(cx))^3}{(d+ex)^3} dx$$

Optimal result	194
Mathematica [F]	195
Rubi [A] (verified)	196
Maple [C] (warning: unable to verify)	198
Fricas [F]	198
Sympy [F]	198
Maxima [F]	199
Giac [F]	200
Mupad [F(-1)]	200
Reduce [F]	200

Optimal result

Integrand size = 18, antiderivative size = 953

$$\int \frac{(a + b\operatorname{arctanh}(cx))^3}{(d + ex)^3} dx = \text{Too large to display}$$

output

```

3/2*b*c*(a+b*arctanh(c*x))^2/(c^2*d^2-e^2)/(e*x+d)-1/2*(a+b*arctanh(c*x))^
3/e/(e*x+d)^2-3/2*b^2*c^2*(a+b*arctanh(c*x))*ln(2/(-c*x+1))/(c*d-e)/(c*d+e
)^2+3/4*b*c^2*(a+b*arctanh(c*x))^2*ln(2/(-c*x+1))/e/(c*d+e)^2-3*b^2*c^2*e*
(a+b*arctanh(c*x))*ln(2/(c*x+1))/(c*d-e)^2/(c*d+e)^2+3/2*b^2*c^2*(a+b*arct
anh(c*x))*ln(2/(c*x+1))/(c*d-e)^2/(c*d+e)-3/4*b*c^2*(a+b*arctanh(c*x))^2*ln
(2/(c*x+1))/(c*d-e)^2/e+3*b*c^3*d*(a+b*arctanh(c*x))^2*ln(2/(c*x+1))/(c*d
-e)^2/(c*d+e)^2+3*b^2*c^2*e*(a+b*arctanh(c*x))*ln(2*c*(e*x+d)/(c*d+e)/(c*x
+1))/(c*d-e)^2/(c*d+e)^2-3*b*c^3*d*(a+b*arctanh(c*x))^2*ln(2*c*(e*x+d)/(c*
d+e)/(c*x+1))/(c*d-e)^2/(c*d+e)^2-3/4*b^3*c^2*polylog(2,1-2/(-c*x+1))/(c*d
-e)/(c*d+e)^2+3/4*b^2*c^2*(a+b*arctanh(c*x))*polylog(2,1-2/(-c*x+1))/e/(c*
d+e)^2+3/2*b^3*c^2*e*polylog(2,1-2/(c*x+1))/(c*d-e)^2/(c*d+e)^2-3/4*b^3*c^
2*polylog(2,1-2/(c*x+1))/(c*d-e)^2/(c*d+e)+3/4*b^2*c^2*(a+b*arctanh(c*x))*
polylog(2,1-2/(c*x+1))/(c*d-e)^2/e-3*b^2*c^3*d*(a+b*arctanh(c*x))*polylog(
2,1-2/(c*x+1))/(c*d-e)^2/(c*d+e)^2-3/2*b^3*c^2*e*polylog(2,1-2*c*(e*x+d)/(
c*d+e)/(c*x+1))/(c*d-e)^2/(c*d+e)^2+3*b^2*c^3*d*(a+b*arctanh(c*x))*polylog
(2,1-2*c*(e*x+d)/(c*d+e)/(c*x+1))/(c*d-e)^2/(c*d+e)^2-3/8*b^3*c^2*polylog(
3,1-2/(-c*x+1))/e/(c*d+e)^2+3/8*b^3*c^2*polylog(3,1-2/(c*x+1))/(c*d-e)^2/e
-3/2*b^3*c^3*d*polylog(3,1-2/(c*x+1))/(c*d-e)^2/(c*d+e)^2+3/2*b^3*c^3*d*po
lylog(3,1-2*c*(e*x+d)/(c*d+e)/(c*x+1))/(c*d-e)^2/(c*d+e)^2

```

Mathematica [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^3} dx = \int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^3} dx$$

input

```
Integrate[(a + b*ArcTanh[c*x])^3/(d + e*x)^3,x]
```

output

```
Integrate[(a + b*ArcTanh[c*x])^3/(d + e*x)^3, x]
```

Rubi [A] (verified)

Time = 2.17 (sec) , antiderivative size = 896, normalized size of antiderivative = 0.94, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^3} dx$$

↓ 6480

$$\frac{3bc \int \left(\frac{c^2(a+b \operatorname{arctanh}(cx))^2}{2(cd+e)^2(1-cx)} + \frac{c^2(a+b \operatorname{arctanh}(cx))^2}{2(cd-e)^2(cx+1)} - \frac{2c^2de^2(a+b \operatorname{arctanh}(cx))^2}{(cd-e)^2(cd+e)^2(d+ex)} - \frac{e^2(a+b \operatorname{arctanh}(cx))^2}{(cd-e)(cd+e)(d+ex)^2} \right) dx}{\frac{2e(a+b \operatorname{arctanh}(cx))^3}{2e(d+ex)^2}}$$

↓ 2009

$$\frac{3bc \left(-\frac{ce \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx}\right) b^2}{2(cd-e)(cd+e)^2} - \frac{ce \operatorname{PolyLog}\left(2, 1 - \frac{2}{cx+1}\right) b^2}{2(cd-e)^2(cd+e)} + \frac{ce^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{cx+1}\right) b^2}{(cd-e)^2(cd+e)^2} - \frac{ce^2 \operatorname{PolyLog}\left(2, 1 - \frac{2c(d+ex)}{(cd+e)(cx+1)}\right) b^2}{(cd-e)^2(cd+e)^2} - \frac{c^2 d^2 \operatorname{PolyLog}\left(2, 1 - \frac{2c(d+ex)}{(cd+e)(cx+1)}\right) b^2}{(cd-e)^2(cd+e)^2} \right)}{\frac{2e(a+b \operatorname{arctanh}(cx))^3}{2e(d+ex)^2}}$$

input `Int[(a + b*ArcTanh[c*x])^3/(d + e*x)^3, x]`

output

```

-1/2*(a + b*ArcTanh[c*x])^3/(e*(d + e*x)^2) + (3*b*c*((e*(a + b*ArcTanh[c*
x])^2)/((c^2*d^2 - e^2)*(d + e*x)) - (b*c*e*(a + b*ArcTanh[c*x])*Log[2/(1
- c*x)])/((c*d - e)*(c*d + e)^2) + (c*(a + b*ArcTanh[c*x])^2*Log[2/(1 - c*
x)])/(2*(c*d + e)^2) - (2*b*c*e^2*(a + b*ArcTanh[c*x])*Log[2/(1 + c*x)])/(
(c*d - e)^2*(c*d + e)^2) + (b*c*e*(a + b*ArcTanh[c*x])*Log[2/(1 + c*x)])/(
(c*d - e)^2*(c*d + e)) - (c*(a + b*ArcTanh[c*x])^2*Log[2/(1 + c*x)])/(2*(c
*d - e)^2) + (2*c^2*d*e*(a + b*ArcTanh[c*x])^2*Log[2/(1 + c*x)])/((c*d - e
)^2*(c*d + e)^2) + (2*b*c*e^2*(a + b*ArcTanh[c*x])*Log[(2*c*(d + e*x))/((c
*d + e)*(1 + c*x))])/((c*d - e)^2*(c*d + e)^2) - (2*c^2*d*e*(a + b*ArcTanh
[c*x])^2*Log[(2*c*(d + e*x))/((c*d + e)*(1 + c*x))])/((c*d - e)^2*(c*d + e
)^2) - (b^2*c*e*PolyLog[2, 1 - 2/(1 - c*x)])/(2*(c*d - e)*(c*d + e)^2) + (
b*c*(a + b*ArcTanh[c*x])*PolyLog[2, 1 - 2/(1 - c*x)])/(2*(c*d + e)^2) + (b
^2*c*e^2*PolyLog[2, 1 - 2/(1 + c*x)])/((c*d - e)^2*(c*d + e)^2) - (b^2*c*e
*PolyLog[2, 1 - 2/(1 + c*x)])/(2*(c*d - e)^2*(c*d + e)) + (b*c*(a + b*ArcT
anh[c*x])*PolyLog[2, 1 - 2/(1 + c*x)])/(2*(c*d - e)^2) - (2*b*c^2*d*e*(a +
b*ArcTanh[c*x])*PolyLog[2, 1 - 2/(1 + c*x)])/((c*d - e)^2*(c*d + e)^2) -
(b^2*c*e^2*PolyLog[2, 1 - (2*c*(d + e*x))/((c*d + e)*(1 + c*x))])/((c*d -
e)^2*(c*d + e)^2) + (2*b*c^2*d*e*(a + b*ArcTanh[c*x])*PolyLog[2, 1 - (2*c*
(d + e*x))/((c*d + e)*(1 + c*x))])/((c*d - e)^2*(c*d + e)^2) - (b^2*c*Poly
Log[3, 1 - 2/(1 - c*x)])/(4*(c*d + e)^2) + (b^2*c*PolyLog[3, 1 - 2/(1 +...

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6480

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^p_*((d_) + (e_.)*(x_))^(q_.), x_S
ymbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] -
Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1
), (d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x]
&& IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 10.63 (sec) , antiderivative size = 50477, normalized size of antiderivative = 52.97

method	result	size
derivativedivides	Expression too large to display	50477
default	Expression too large to display	50477
parts	Expression too large to display	50485

input `int((a+b*arctanh(c*x))^3/(e*x+d)^3,x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^3} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{(ex + d)^3} dx$$

input `integrate((a+b*arctanh(c*x))^3/(e*x+d)^3,x, algorithm="fricas")`

output `integral((b^3*arctanh(c*x)^3 + 3*a*b^2*arctanh(c*x)^2 + 3*a^2*b*arctanh(c*x) + a^3)/(e^3*x^3 + 3*d*e^2*x^2 + 3*d^2*e*x + d^3), x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^3} dx = \int \frac{(a + b \operatorname{atanh}(cx))^3}{(d + ex)^3} dx$$

input `integrate((a+b*atanh(c*x))**3/(e*x+d)**3,x)`

output `Integral((a + b*atanh(c*x))**3/(d + e*x)**3, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^3} dx = \int \frac{(b \operatorname{arctanh}(cx) + a)^3}{(ex + d)^3} dx$$

input `integrate((a+b*arctanh(c*x))^3/(e*x+d)^3,x, algorithm="maxima")`

output

```
-3/4*((4*c^2*d*log(e*x + d)/(c^4*d^4 - 2*c^2*d^2*e^2 + e^4) - c*log(c*x +
1)/(c^2*d^2*e - 2*c*d*e^2 + e^3) + c*log(c*x - 1)/(c^2*d^2*e + 2*c*d*e^2 +
e^3) - 2/(c^2*d^3 - d*e^2 + (c^2*d^2*e - e^3)*x))*c + 2*arctanh(c*x)/(e^3
*x^2 + 2*d*e^2*x + d^2*e))*a^2*b - 1/2*a^3/(e^3*x^2 + 2*d*e^2*x + d^2*e) -
1/16*(((c^4*d^2*e^2 - 2*c^3*d*e^3 + c^2*e^4)*b^3*x^2 + 2*(c^4*d^3*e - 2*c
^3*d^2*e^2 + c^2*d*e^3)*b^3*x - (2*c^3*d^3*e - 3*c^2*d^2*e^2 + e^4)*b^3)*1
og(-c*x + 1)^3 - 3*(2*(c^3*d^2*e^2 - c*e^4)*b^3*x - 2*(c^4*d^4 - 2*c^2*d^
2*e^2 + e^4)*a*b^2 + 2*(c^3*d^3*e - c*d*e^3)*b^3 + ((c^4*d^2*e^2 + 2*c^3*d*
e^3 + c^2*e^4)*b^3*x^2 + 2*(c^4*d^3*e + 2*c^3*d^2*e^2 + c^2*d*e^3)*b^3*x +
(2*c^3*d^3*e + 3*c^2*d^2*e^2 - e^4)*b^3)*log(c*x + 1))*log(-c*x + 1)^2)/(
c^4*d^6*e - 2*c^2*d^4*e^3 + d^2*e^5 + (c^4*d^4*e^3 - 2*c^2*d^2*e^5 + e^7)*
x^2 + 2*(c^4*d^5*e^2 - 2*c^2*d^3*e^4 + d*e^6)*x) - integrate(1/8*(((c^4*d^
3*e - c^3*d^2*e^2 - c^2*d*e^3 + c*e^4)*b^3*x - (c^3*d^3*e - c^2*d^2*e^2 -
c*d*e^3 + e^4)*b^3)*log(c*x + 1)^3 + 6*(((c^4*d^3*e - c^3*d^2*e^2 - c^2*d*e
^3 + c*e^4)*a*b^2*x - (c^3*d^3*e - c^2*d^2*e^2 - c*d*e^3 + e^4)*a*b^2)*log
(c*x + 1)^2 - 3*(2*(c^3*d*e^3 - c^2*e^4)*b^3*x^2 - 2*(c^4*d^4 - c^3*d^3*e
- c^2*d^2*e^2 + c*d*e^3)*a*b^2 + 2*(c^3*d^3*e - c^2*d^2*e^2)*b^3 + ((c^4*d
^3*e - c^3*d^2*e^2 - c^2*d*e^3 + c*e^4)*b^3*x - (c^3*d^3*e - c^2*d^2*e^2 -
c*d*e^3 + e^4)*b^3)*log(c*x + 1)^2 - 2*(((c^4*d^3*e - c^3*d^2*e^2 - c^2*d*
e^3 + c*e^4)*a*b^2 - 2*(c^3*d^2*e^2 - c^2*d*e^3)*b^3)*x + ((c^4*d*e^3 + ...
```


Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^3} dx = \int \frac{(b \operatorname{artanh}(cx) + a)^3}{(ex + d)^3} dx$$

input `integrate((a+b*arctanh(c*x))^3/(e*x+d)^3,x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)^3/(e*x + d)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^3} dx = \int \frac{(a + b \operatorname{atanh}(cx))^3}{(d + ex)^3} dx$$

input `int((a + b*atanh(c*x))^3/(d + e*x)^3,x)`

output `int((a + b*atanh(c*x))^3/(d + e*x)^3, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(cx))^3}{(d + ex)^3} dx = \text{too large to display}$$

input `int((a+b*atanh(c*x))^3/(e*x+d)^3,x)`

output

```
(36*atanh(c*x)**3*b**3*c**10*d**10*e*x + 18*atanh(c*x)**3*b**3*c**10*d**9*
e**2*x**2 - 54*atanh(c*x)**3*b**3*c**8*d**9*e**2 - 48*atanh(c*x)**3*b**3*c
**8*d**8*e**3*x - 24*atanh(c*x)**3*b**3*c**8*d**7*e**4*x**2 + 72*atanh(c*x
)**3*b**3*c**6*d**7*e**4 - 8*atanh(c*x)**3*b**3*c**6*d**6*e**5*x - 4*atanh
(c*x)**3*b**3*c**6*d**5*e**6*x**2 + 12*atanh(c*x)**3*b**3*c**4*d**5*e**6 +
16*atanh(c*x)**3*b**3*c**4*d**4*e**7*x + 8*atanh(c*x)**3*b**3*c**4*d**3*e
**8*x**2 - 24*atanh(c*x)**3*b**3*c**2*d**3*e**8 + 4*atanh(c*x)**3*b**3*c**
2*d**2*e**9*x + 2*atanh(c*x)**3*b**3*c**2*d*e**10*x**2 - 6*atanh(c*x)**3*b
**3*d*e**10 + 108*atanh(c*x)**2*a*b**2*c**10*d**10*e*x + 54*atanh(c*x)**2*
a*b**2*c**10*d**9*e**2*x**2 - 162*atanh(c*x)**2*a*b**2*c**8*d**9*e**2 - 14
4*atanh(c*x)**2*a*b**2*c**8*d**8*e**3*x - 72*atanh(c*x)**2*a*b**2*c**8*d**
7*e**4*x**2 + 216*atanh(c*x)**2*a*b**2*c**6*d**7*e**4 - 24*atanh(c*x)**2*a
*b**2*c**6*d**6*e**5*x - 12*atanh(c*x)**2*a*b**2*c**6*d**5*e**6*x**2 + 36*
atanh(c*x)**2*a*b**2*c**4*d**5*e**6 + 48*atanh(c*x)**2*a*b**2*c**4*d**4*e
**7*x + 24*atanh(c*x)**2*a*b**2*c**4*d**3*e**8*x**2 - 72*atanh(c*x)**2*a*b
**2*c**2*d**3*e**8 + 12*atanh(c*x)**2*a*b**2*c**2*d**2*e**9*x + 6*atanh(c*x
)**2*a*b**2*c**2*d*e**10*x**2 - 18*atanh(c*x)**2*a*b**2*d*e**10 - 36*atanh
(c*x)**2*b**3*c**9*d**10*e + 90*atanh(c*x)**2*b**3*c**9*d**9*e**2*x + 72*a
tanh(c*x)**2*b**3*c**9*d**8*e**3*x**2 - 6*atanh(c*x)**2*b**3*c**7*d**8*e**
3 - 228*atanh(c*x)**2*b**3*c**7*d**7*e**4*x - 150*atanh(c*x)**2*b**3*c...
```

3.21 $\int \frac{a+b\operatorname{arctanh}(cx)}{1+2cx} dx$

Optimal result	202
Mathematica [C] (warning: unable to verify)	202
Rubi [A] (verified)	203
Maple [A] (verified)	205
Fricas [F]	205
Sympy [F]	206
Maxima [F]	206
Giac [F]	206
Mupad [F(-1)]	207
Reduce [F]	207

Optimal result

Integrand size = 17, antiderivative size = 67

$$\int \frac{a + \operatorname{arctanh}(cx)}{1 + 2cx} dx = \frac{(a - \operatorname{arctanh}(\frac{1}{2})) \log(-\frac{1+2cx}{2d})}{2c} - \frac{b \operatorname{PolyLog}(2, -1 - 2cx)}{4c} + \frac{b \operatorname{PolyLog}(2, \frac{1}{3}(1 + 2cx))}{4c}$$

output `1/2*(a-b*arctanh(1/2))*ln(-1/2*(2*c*x+1)/d)/c-1/4*b*polylog(2,-2*c*x-1)/c+1/4*b*polylog(2,2/3*c*x+1/3)/c`

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.21 (sec) , antiderivative size = 240, normalized size of antiderivative = 3.58

$$\int \frac{a + \operatorname{arctanh}(cx)}{1 + 2cx} dx = \frac{a \log(1 + 2cx) + \operatorname{arctanh}(cx) (\frac{1}{2} \log(1 - c^2 x^2) + \log(i \sinh(\operatorname{arctanh}(\frac{1}{2}) + \operatorname{arctanh}(cx)))) - \frac{1}{2} i b (-\frac{1}{4} i \dots)}{\dots}$$

input `Integrate[(a + b*ArcTanh[c*x])/(1 + 2*c*x), x]`

output

```
(a*Log[1 + 2*c*x] + b*ArcTanh[c*x]*(Log[1 - c^2*x^2]/2 + Log[I*Sinh[ArcTanh[1/2] + ArcTanh[c*x]]]) - (I/2)*b*((-1/4*I)*(Pi - (2*I)*ArcTanh[c*x])^2 + I*(ArcTanh[1/2] + ArcTanh[c*x])^2 + (Pi - (2*I)*ArcTanh[c*x])*Log[1 + E^(2*ArcTanh[c*x])] + (2*I)*(ArcTanh[1/2] + ArcTanh[c*x])*Log[1 - E^(-2*(ArcTanh[1/2] + ArcTanh[c*x])])]) - (Pi - (2*I)*ArcTanh[c*x])*Log[2/Sqrt[1 - c^2*x^2]] - (2*I)*(ArcTanh[1/2] + ArcTanh[c*x])*Log[(2*I)*Sinh[ArcTanh[1/2] + ArcTanh[c*x]]] - I*PolyLog[2, -E^(2*ArcTanh[c*x])] - I*PolyLog[2, E^(-2*(ArcTanh[1/2] + ArcTanh[c*x])])])/(2*c)
```

Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.63, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {6472, 2849, 2752, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}(cx)}{2cx + 1} dx \\
 & \quad \downarrow \text{6472} \\
 & \frac{1}{2}b \int \frac{\log\left(\frac{2}{cx+1}\right)}{1 - c^2x^2} dx - \frac{1}{2}b \int \frac{\log\left(\frac{2(2cx+1)}{3(cx+1)}\right)}{1 - c^2x^2} dx - \frac{\log\left(\frac{2}{cx+1}\right)(a + b \operatorname{arctanh}(cx))}{2c} + \\
 & \quad \frac{\log\left(\frac{2(2cx+1)}{3(cx+1)}\right)(a + b \operatorname{arctanh}(cx))}{2c} \\
 & \quad \downarrow \text{2849} \\
 & -\frac{1}{2}b \int \frac{\log\left(\frac{2(2cx+1)}{3(cx+1)}\right)}{1 - c^2x^2} dx + \frac{b \int \frac{\log\left(\frac{2}{cx+1}\right) d\frac{1}{cx+1}}{1 - \frac{2}{cx+1}}}{2c} - \frac{\log\left(\frac{2}{cx+1}\right)(a + b \operatorname{arctanh}(cx))}{2c} + \\
 & \quad \frac{\log\left(\frac{2(2cx+1)}{3(cx+1)}\right)(a + b \operatorname{arctanh}(cx))}{2c} \\
 & \quad \downarrow \text{2752}
 \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{2}b \int \frac{\log\left(\frac{2(2cx+1)}{3(cx+1)}\right)}{1-c^2x^2} dx - \frac{\log\left(\frac{2}{cx+1}\right)(a + \operatorname{barctanh}(cx))}{2c} + \\
& \frac{\log\left(\frac{2(2cx+1)}{3(cx+1)}\right)(a + \operatorname{barctanh}(cx))}{2c} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{cx+1}\right)}{4c} \\
& \quad \downarrow \text{2897} \\
& -\frac{\log\left(\frac{2}{cx+1}\right)(a + \operatorname{barctanh}(cx))}{2c} + \frac{\log\left(\frac{2(2cx+1)}{3(cx+1)}\right)(a + \operatorname{barctanh}(cx))}{2c} + \\
& \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{cx+1}\right)}{4c} - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2(2cx+1)}{3(cx+1)}\right)}{4c}
\end{aligned}$$

input `Int[(a + b*ArcTanh[c*x])/(1 + 2*c*x), x]`

output `-1/2*((a + b*ArcTanh[c*x])*Log[2/(1 + c*x)])/c + ((a + b*ArcTanh[c*x])*Log[(2*(1 + 2*c*x))/(3*(1 + c*x))])/(2*c) + (b*PolyLog[2, 1 - 2/(1 + c*x)])/(4*c) - (b*PolyLog[2, 1 - (2*(1 + 2*c*x))/(3*(1 + c*x))])/(4*c)`

Defintions of rubi rules used

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 2897 `Int[Log[u_]*(Pq_)^(m_.), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]]`

rule 6472

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))/((d_.) + (e_.)*(x_.)), x_Symbol] := S
imp[(-(a + b*ArcTanh[c*x]))*(Log[2/(1 + c*x)]/e), x] + (Simp[(a + b*ArcTanh
[c*x])*(Log[2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/e), x] + Simp[b*(c/e)
Int[Log[2/(1 + c*x)]/(1 - c^2*x^2), x], x] - Simp[b*(c/e) Int[Log[2*c*((d
+ e*x)/((c*d + e)*(1 + c*x)))]/(1 - c^2*x^2), x], x]) /; FreeQ[{a, b, c, d
, e}, x] && NeQ[c^2*d^2 - e^2, 0]
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.37

method	result
derivativedivides	$\frac{\frac{a \ln(2cx+1)}{2} + b \left(\frac{\ln(2cx+1) \operatorname{arctanh}(cx)}{2} + \frac{(\ln(2cx+1) - \ln(\frac{2cx}{3} + \frac{1}{3})) \ln(\frac{2}{3} - \frac{2cx}{3})}{4} - \frac{\operatorname{dilog}(\frac{2cx}{3} + \frac{1}{3})}{4} - \frac{\operatorname{dilog}(2cx+2)}{4} - \frac{\ln(2cx+1) \ln(2cx+2)}{4} \right)}{c}$
default	$\frac{\frac{a \ln(2cx+1)}{2} + b \left(\frac{\ln(2cx+1) \operatorname{arctanh}(cx)}{2} + \frac{(\ln(2cx+1) - \ln(\frac{2cx}{3} + \frac{1}{3})) \ln(\frac{2}{3} - \frac{2cx}{3})}{4} - \frac{\operatorname{dilog}(\frac{2cx}{3} + \frac{1}{3})}{4} - \frac{\operatorname{dilog}(2cx+2)}{4} - \frac{\ln(2cx+1) \ln(2cx+2)}{4} \right)}{c}$
parts	$\frac{a \ln(2cx+1)}{2c} + \frac{b \left(\frac{\ln(2cx+1) \operatorname{arctanh}(cx)}{2} + \frac{(\ln(2cx+1) - \ln(\frac{2cx}{3} + \frac{1}{3})) \ln(\frac{2}{3} - \frac{2cx}{3})}{4} - \frac{\operatorname{dilog}(\frac{2cx}{3} + \frac{1}{3})}{4} - \frac{\operatorname{dilog}(2cx+2)}{4} - \frac{\ln(2cx+1) \ln(2cx+2)}{4} \right)}{c}$
risch	$\frac{b \ln(\frac{2cx}{3} + \frac{1}{3}) \ln(\frac{2}{3} - \frac{2cx}{3})}{4c} - \frac{b \ln(\frac{2cx}{3} + \frac{1}{3}) \ln(-cx+1)}{4c} + \frac{b \operatorname{dilog}(\frac{2}{3} - \frac{2cx}{3})}{4c} + \frac{a \ln(-2cx-1)}{2c} - \frac{b \ln(-2cx-1) \ln(2cx+2)}{4c}$

input

```
int((a+b*arctanh(c*x))/(2*c*x+1),x,method=_RETURNVERBOSE)
```

output

```
1/c*(1/2*a*ln(2*c*x+1)+b*(1/2*ln(2*c*x+1)*arctanh(c*x)+1/4*(ln(2*c*x+1)-ln
(2/3*c*x+1/3))*ln(2/3-2/3*c*x)-1/4*dilog(2/3*c*x+1/3)-1/4*dilog(2*c*x+2)-1
/4*ln(2*c*x+1)*ln(2*c*x+2)))
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{1 + 2cx} dx = \int \frac{b \operatorname{arctanh}(cx) + a}{2cx + 1} dx$$

input

```
integrate((a+b*arctanh(c*x))/(2*c*x+1),x, algorithm="fricas")
```

output `integral((b*arctanh(c*x) + a)/(2*c*x + 1), x)`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{1 + 2cx} dx = \int \frac{a + b \operatorname{atanh}(cx)}{2cx + 1} dx$$

input `integrate((a+b*atanh(c*x))/(2*c*x+1), x)`

output `Integral((a + b*atanh(c*x))/(2*c*x + 1), x)`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{1 + 2cx} dx = \int \frac{b \operatorname{artanh}(cx) + a}{2cx + 1} dx$$

input `integrate((a+b*arctanh(c*x))/(2*c*x+1), x, algorithm="maxima")`

output `1/2*b*integrate((log(c*x + 1) - log(-c*x + 1))/(2*c*x + 1), x) + 1/2*a*log(2*c*x + 1)/c`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(cx)}{1 + 2cx} dx = \int \frac{b \operatorname{artanh}(cx) + a}{2cx + 1} dx$$

input `integrate((a+b*arctanh(c*x))/(2*c*x+1), x, algorithm="giac")`

output `integrate((b*arctanh(c*x) + a)/(2*c*x + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx)}{1 + 2cx} dx = \int \frac{a + b \operatorname{atanh}(cx)}{2cx + 1} dx$$

input `int((a + b*atanh(c*x))/(2*c*x + 1),x)`output `int((a + b*atanh(c*x))/(2*c*x + 1), x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{arctanh}(cx)}{1 + 2cx} dx = \frac{2 \left(\int \frac{\operatorname{atanh}(cx)}{2cx+1} dx \right) bc + \log(2cx + 1) a}{2c}$$

input `int((a+b*atanh(c*x))/(2*c*x+1),x)`output `(2*int(atanh(c*x)/(2*c*x + 1),x)*b*c + log(2*c*x + 1)*a)/(2*c)`

3.22 $\int \frac{\operatorname{arctanh}(x)}{1-\sqrt{2}x} dx$

Optimal result	208
Mathematica [C] (verified)	208
Rubi [A] (verified)	209
Maple [A] (verified)	211
Fricas [F]	212
Sympy [F]	212
Maxima [B] (verification not implemented)	212
Giac [F]	213
Mupad [F(-1)]	213
Reduce [F]	214

Optimal result

Integrand size = 15, antiderivative size = 88

$$\int \frac{\operatorname{arctanh}(x)}{1-\sqrt{2}x} dx = -\frac{\operatorname{arctanh}\left(\frac{1}{\sqrt{2}}\right) \log(1-\sqrt{2}x)}{\sqrt{2}} - \frac{\operatorname{PolyLog}\left(2, -\frac{\sqrt{2}-2x}{2-\sqrt{2}}\right)}{2\sqrt{2}} + \frac{\operatorname{PolyLog}\left(2, \frac{\sqrt{2}-2x}{2+\sqrt{2}}\right)}{2\sqrt{2}}$$

output

```
-1/2*arctanh(1/2*2^(1/2))*ln(1-x*2^(1/2))*2^(1/2)-1/4*polylog(2,-(2^(1/2)-2*x)/(2-2^(1/2)))*2^(1/2)+1/4*polylog(2,(2^(1/2)-2*x)/(2+2^(1/2)))*2^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 272, normalized size of antiderivative = 3.09

$$\int \frac{\operatorname{arctanh}(x)}{1-\sqrt{2}x} dx$$

$$= \frac{\pi^2 - 4\operatorname{arctanh}\left(\frac{1}{\sqrt{2}}\right)^2 - 4i\pi\operatorname{arctanh}(x) + 8\operatorname{arctanh}\left(\frac{1}{\sqrt{2}}\right)\operatorname{arctanh}(x) - 8\operatorname{arctanh}(x)^2 + 8\operatorname{arctanh}\left(\frac{1}{\sqrt{2}}\right)\log(1-\sqrt{2}x)}{2\sqrt{2}}$$

input `Integrate[ArcTanh[x]/(1 - Sqrt[2]*x), x]`

output `(Pi^2 - 4*ArcTanh[1/Sqrt[2]]^2 - (4*I)*Pi*ArcTanh[x] + 8*ArcTanh[1/Sqrt[2]]*ArcTanh[x] - 8*ArcTanh[x]^2 + 8*ArcTanh[1/Sqrt[2]]*Log[1 - E^(2*ArcTanh[1/Sqrt[2]] - 2*ArcTanh[x])] - 8*ArcTanh[x]*Log[1 - E^(2*ArcTanh[1/Sqrt[2]] - 2*ArcTanh[x])] + (4*I)*Pi*Log[1 + E^(2*ArcTanh[x])] + 8*ArcTanh[x]*Log[1 + E^(2*ArcTanh[x])] - (4*I)*Pi*Log[2/Sqrt[1 - x^2]] - 8*ArcTanh[x]*Log[2/Sqrt[1 - x^2]] - 4*ArcTanh[x]*Log[1 - x^2] - 8*ArcTanh[x]*Log[(-I)*Sinh[ArcTanh[1/Sqrt[2]] - ArcTanh[x]]] - 8*ArcTanh[1/Sqrt[2]]*Log[(-2*I)*Sinh[ArcTanh[1/Sqrt[2]] - ArcTanh[x]]] + 8*ArcTanh[x]*Log[(-2*I)*Sinh[ArcTanh[1/Sqrt[2]] - ArcTanh[x]]] + 4*PolyLog[2, E^(2*ArcTanh[1/Sqrt[2]] - 2*ArcTanh[x])] + 4*PolyLog[2, -E^(2*ArcTanh[x])])/(8*Sqrt[2])`

Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.23, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {6472, 2849, 2752, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arctanh}(x)}{1 - \sqrt{2}x} dx \\
 & \quad \downarrow \text{6472} \\
 & -\frac{\int \frac{\log\left(\frac{2}{x+1}\right) dx}{1-x^2}}{\sqrt{2}} + \frac{\int \frac{\log\left(-\frac{2(1+\sqrt{2})(1-\sqrt{2}x)}{x+1}\right) dx}{1-x^2}}{\sqrt{2}} + \frac{\operatorname{arctanh}(x) \log\left(\frac{2}{x+1}\right)}{\sqrt{2}} - \\
 & \quad \frac{\operatorname{arctanh}(x) \log\left(-\frac{2(1+\sqrt{2})(1-\sqrt{2}x)}{x+1}\right)}{\sqrt{2}} \\
 & \quad \downarrow \text{2849}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{\int \frac{\log\left(-\frac{2(1+\sqrt{2})(1-\sqrt{2}x)}{x+1}\right)}{1-x^2} dx}{\sqrt{2}} - \frac{\int \frac{\log\left(\frac{2}{x+1}\right)}{1-\frac{2}{x+1}} d\frac{1}{x+1}}{\sqrt{2}} + \frac{\operatorname{arctanh}(x) \log\left(\frac{2}{x+1}\right)}{\sqrt{2}} - \\
 & \frac{\operatorname{arctanh}(x) \log\left(-\frac{2(1+\sqrt{2})(1-\sqrt{2}x)}{x+1}\right)}{\sqrt{2}} \\
 & \quad \downarrow \text{2752} \\
 & \frac{\int \frac{\log\left(-\frac{2(1+\sqrt{2})(1-\sqrt{2}x)}{x+1}\right)}{1-x^2} dx}{\sqrt{2}} + \frac{\operatorname{arctanh}(x) \log\left(\frac{2}{x+1}\right)}{\sqrt{2}} - \frac{\operatorname{arctanh}(x) \log\left(-\frac{2(1+\sqrt{2})(1-\sqrt{2}x)}{x+1}\right)}{\sqrt{2}} - \\
 & \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{x+1}\right)}{2\sqrt{2}} \\
 & \quad \downarrow \text{2897} \\
 & \frac{\operatorname{arctanh}(x) \log\left(\frac{2}{x+1}\right)}{\sqrt{2}} - \frac{\operatorname{arctanh}(x) \log\left(-\frac{2(1+\sqrt{2})(1-\sqrt{2}x)}{x+1}\right)}{\sqrt{2}} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{x+1}\right)}{2\sqrt{2}} + \\
 & \frac{\operatorname{PolyLog}\left(2, \frac{2(1+\sqrt{2})(1-\sqrt{2}x)}{x+1} + 1\right)}{2\sqrt{2}}
 \end{aligned}$$

input `Int[ArcTanh[x]/(1 - Sqrt[2]*x), x]`

output `(ArcTanh[x]*Log[2/(1 + x)]/Sqrt[2] - (ArcTanh[x]*Log[(-2*(1 + Sqrt[2])*(1 - Sqrt[2]*x))/(1 + x)]/Sqrt[2] - PolyLog[2, 1 - 2/(1 + x)]/(2*Sqrt[2]) + PolyLog[2, 1 + (2*(1 + Sqrt[2])*(1 - Sqrt[2]*x))/(1 + x)]/(2*Sqrt[2]))`

Defintions of rubi rules used

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] :> Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 $\text{Int}[\text{Log}[(c_)/((d_)+(e_)*(x_))]/((f_)+(g_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[-e/g \text{ Subst}[\text{Int}[\text{Log}[2*d*x]/(1-2*d*x), x], x, 1/(d+e*x)], x] /; \text{FreeQ}[\{c, d, e, f, g\}, x] \&\& \text{EqQ}[c, 2*d] \&\& \text{EqQ}[e^2*f + d^2*g, 0]$

rule 2897 $\text{Int}[\text{Log}[u_]*(Pq_)^{(m_.)}, x_Symbol] \rightarrow \text{With}[\{C = \text{FullSimplify}[Pq^m*((1-u)/D[u, x])]\}, \text{Simp}[C*\text{PolyLog}[2, 1-u], x] /; \text{FreeQ}[C, x] /; \text{IntegerQ}[m] \&\& \text{PolyQ}[Pq, x] \&\& \text{RationalFunctionQ}[u, x] \&\& \text{LeQ}[\text{RationalFunctionExponents}[u, x][[2]], \text{Expon}[Pq, x]]]$

rule 6472 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)]*(b_))/((d_)+(e_)*(x_)), x_Symbol] \rightarrow \text{Simp}[(-a + b*\text{ArcTanh}[c*x])*(\text{Log}[2/(1+c*x)]/e), x] + (\text{Simp}[(a + b*\text{ArcTanh}[c*x])*(\text{Log}[2*c*((d+e*x)/((c*d+e)*(1+c*x))])]/e), x] + \text{Simp}[b*(c/e) \text{ Int}[\text{Log}[2/(1+c*x)]/(1-c^2*x^2), x], x] - \text{Simp}[b*(c/e) \text{ Int}[\text{Log}[2*c*((d+e*x)/((c*d+e)*(1+c*x)))]/(1-c^2*x^2), x], x)] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{NeQ}[c^2*d^2 - e^2, 0]$

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.44

method	result
default	$-\frac{\ln(\sqrt{2}x-1)\sqrt{2} \operatorname{arctanh}(x)}{2} - \frac{\sqrt{2} \ln(\sqrt{2}x-1) \ln\left(\frac{\sqrt{2}-\sqrt{2}x}{\sqrt{2}-1}\right)}{4} + \frac{\sqrt{2} \ln(\sqrt{2}x-1) \ln\left(\frac{\sqrt{2}+\sqrt{2}x}{1+\sqrt{2}}\right)}{4} - \frac{\sqrt{2} \operatorname{dilog}\left(\frac{\sqrt{2}-\sqrt{2}x}{\sqrt{2}-1}\right)}{4} +$
parts	$-\frac{\ln(\sqrt{2}x-1)\sqrt{2} \operatorname{arctanh}(x)}{2} - \frac{\sqrt{2} \ln(\sqrt{2}x-1) \ln\left(\frac{\sqrt{2}-\sqrt{2}x}{\sqrt{2}-1}\right)}{4} + \frac{\sqrt{2} \ln(\sqrt{2}x-1) \ln\left(\frac{\sqrt{2}+\sqrt{2}x}{1+\sqrt{2}}\right)}{4} - \frac{\sqrt{2} \operatorname{dilog}\left(\frac{\sqrt{2}-\sqrt{2}x}{\sqrt{2}-1}\right)}{4} +$
risch	$\frac{\sqrt{2} \ln\left(\frac{\sqrt{2}-2x}{2+\sqrt{2}}\right) \ln\left(\frac{2+2x}{2+\sqrt{2}}\right)}{4} - \frac{\sqrt{2} \ln\left(\frac{\sqrt{2}-2x}{2+\sqrt{2}}\right) \ln(1+x)}{4} + \frac{\sqrt{2} \operatorname{dilog}\left(\frac{2+2x}{2+\sqrt{2}}\right)}{4} - \frac{\sqrt{2} \ln\left(\frac{2x-\sqrt{2}}{2-\sqrt{2}}\right) \ln\left(\frac{2-2x}{2-\sqrt{2}}\right)}{4} + \frac{\sqrt{2} \ln\left(\frac{2x-\sqrt{2}}{2-\sqrt{2}}\right)}{4}$

input $\text{int}(\operatorname{arctanh}(x)/(1-2^{(1/2)*x}), x, \text{method}=_RETURNVERBOSE)$

output
$$-1/2*\ln(2^{(1/2)*x}-1)*2^{(1/2)*\operatorname{arctanh}(x)}-1/4*2^{(1/2)*\ln(2^{(1/2)*x}-1)*\ln((2^{(1/2)}-2^{(1/2)*x})/(2^{(1/2)}-1))+1/4*2^{(1/2)*\ln(2^{(1/2)*x}-1)*\ln((2^{(1/2)}+2^{(1/2)*x})/(1+2^{(1/2)}))}-1/4*2^{(1/2)*\operatorname{dilog}((2^{(1/2)}-2^{(1/2)*x})/(2^{(1/2)}-1))+1/4*2^{(1/2)*\operatorname{dilog}((2^{(1/2)}+2^{(1/2)*x})/(1+2^{(1/2)}))}$$

Fricas [F]

$$\int \frac{\operatorname{arctanh}(x)}{1 - \sqrt{2}x} dx = \int -\frac{\operatorname{artanh}(x)}{\sqrt{2}x - 1} dx$$

input `integrate(arctanh(x)/(1-2^(1/2)*x),x, algorithm="fricas")`

output `integral(-(sqrt(2)*x + 1)*arctanh(x)/(2*x^2 - 1), x)`

Sympy [F]

$$\int \frac{\operatorname{arctanh}(x)}{1 - \sqrt{2}x} dx = - \int \frac{\operatorname{atanh}(x)}{\sqrt{2}x - 1} dx$$

input `integrate(atanh(x)/(1-2**(1/2)*x),x)`

output `-Integral(atanh(x)/(sqrt(2)*x - 1), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 144 vs. 2(69) = 138.

Time = 0.11 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.64

$$\begin{aligned} \int \frac{\operatorname{arctanh}(x)}{1 - \sqrt{2}x} dx &= \frac{1}{4} \sqrt{2} (\log(x+1) - \log(x-1)) \log(\sqrt{2}x - 1) \\ &\quad - \frac{1}{2} \sqrt{2} \operatorname{artanh}(x) \log(\sqrt{2}x - 1) \\ &\quad - \frac{1}{4} \sqrt{2} \left(\log(x+1) \log\left(-\frac{\sqrt{2}x + \sqrt{2}}{\sqrt{2} + 1} + 1\right) + \operatorname{Li}_2\left(\frac{\sqrt{2}x + \sqrt{2}}{\sqrt{2} + 1}\right) \right) \\ &\quad + \frac{1}{4} \sqrt{2} \left(\log(x-1) \log\left(\frac{\sqrt{2}x - \sqrt{2}}{\sqrt{2} - 1} + 1\right) + \operatorname{Li}_2\left(-\frac{\sqrt{2}x - \sqrt{2}}{\sqrt{2} - 1}\right) \right) \end{aligned}$$

input `integrate(arctanh(x)/(1-2^(1/2)*x),x, algorithm="maxima")`

output `1/4*sqrt(2)*(log(x + 1) - log(x - 1))*log(sqrt(2)*x - 1) - 1/2*sqrt(2)*arctanh(x)*log(sqrt(2)*x - 1) - 1/4*sqrt(2)*(log(x + 1)*log(-(sqrt(2)*x + sqrt(2))/(sqrt(2) + 1) + 1) + dilog((sqrt(2)*x + sqrt(2))/(sqrt(2) + 1))) + 1/4*sqrt(2)*(log(x - 1)*log((sqrt(2)*x - sqrt(2))/(sqrt(2) - 1) + 1) + dilog(-(sqrt(2)*x - sqrt(2))/(sqrt(2) - 1)))`

Giac [F]

$$\int \frac{\operatorname{arctanh}(x)}{1 - \sqrt{2}x} dx = \int -\frac{\operatorname{artanh}(x)}{\sqrt{2}x - 1} dx$$

input `integrate(arctanh(x)/(1-2^(1/2)*x),x, algorithm="giac")`

output `integrate(-arctanh(x)/(sqrt(2)*x - 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(x)}{1 - \sqrt{2}x} dx = -\int \frac{\operatorname{atanh}(x)}{\sqrt{2}x - 1} dx$$

input `int(-atanh(x)/(2^(1/2)*x - 1),x)`

output `-int(atanh(x)/(2^(1/2)*x - 1), x)`

Reduce [F]

$$\int \frac{\operatorname{arctanh}(x)}{1 - \sqrt{2}x} dx = \frac{\operatorname{atanh}(x)^2}{4} - \sqrt{2} \left(\int \frac{\operatorname{atanh}(x) x}{2x^2 - 1} dx \right) + \frac{\left(\int \frac{\operatorname{atanh}(x)}{2x^4 - 3x^2 + 1} dx \right)}{2}$$

input `int(atanh(x)/(1-2^(1/2)*x),x)`

output `(atanh(x)**2 - 4*sqrt(2)*int((atanh(x)*x)/(2*x**2 - 1),x) + 2*int(atanh(x)/(2*x**4 - 3*x**2 + 1),x))/4`

3.23 $\int \frac{\operatorname{arctanh}(x)}{a+bx} dx$

Optimal result	215
Mathematica [C] (warning: unable to verify)	215
Rubi [A] (verified)	216
Maple [A] (verified)	218
Fricas [F]	218
Sympy [F]	218
Maxima [A] (verification not implemented)	219
Giac [F]	219
Mupad [F(-1)]	220
Reduce [F]	220

Optimal result

Integrand size = 10, antiderivative size = 86

$$\int \frac{\operatorname{arctanh}(x)}{a+bx} dx = -\frac{\operatorname{arctanh}(x) \log\left(\frac{2}{1+x}\right)}{b} + \frac{\operatorname{arctanh}(x) \log\left(\frac{2(a+bx)}{(a+b)(1+x)}\right)}{b} + \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1+x}\right)}{2b} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2(a+bx)}{(a+b)(1+x)}\right)}{2b}$$

output

```
-arctanh(x)*ln(2/(1+x))/b+arctanh(x)*ln(2*(b*x+a)/(a+b)/(1+x))/b+1/2*polylog(2,1-2/(1+x))/b-1/2*polylog(2,1-2*(b*x+a)/(a+b)/(1+x))/b
```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.06 (sec) , antiderivative size = 260, normalized size of antiderivative = 3.02

$$\int \frac{\operatorname{arctanh}(x)}{a+bx} dx = \frac{-\pi^2 + 4\operatorname{arctanh}\left(\frac{a}{b}\right)^2 + 4i\pi\operatorname{arctanh}(x) + 8\operatorname{arctanh}\left(\frac{a}{b}\right)\operatorname{arctanh}(x) + 8\operatorname{arctanh}(x)^2 - 4i\pi \log\left(1 + e^{2\operatorname{arctanh}(x)}\right)}{2b}$$

input `Integrate[ArcTanh[x]/(a + b*x),x]`

output
$$\begin{aligned} & (-\text{Pi}^2 + 4*\text{ArcTanh}[a/b]^2 + (4*I)*\text{Pi}*\text{ArcTanh}[x] + 8*\text{ArcTanh}[a/b]*\text{ArcTanh}[x] \\ & + 8*\text{ArcTanh}[x]^2 - (4*I)*\text{Pi}*\text{Log}[1 + \text{E}^{(2*\text{ArcTanh}[x])}] - 8*\text{ArcTanh}[x]*\text{Log} \\ & [1 + \text{E}^{(2*\text{ArcTanh}[x])}] + 8*\text{ArcTanh}[a/b]*\text{Log}[1 - \text{E}^{(-2*(\text{ArcTanh}[a/b] + \text{ArcT} \\ & \text{anh}[x]))}] + 8*\text{ArcTanh}[x]*\text{Log}[1 - \text{E}^{(-2*(\text{ArcTanh}[a/b] + \text{ArcTanh}[x]))}] + (4* \\ & I)*\text{Pi}*\text{Log}[2/\text{Sqrt}[1 - x^2]] + 8*\text{ArcTanh}[x]*\text{Log}[2/\text{Sqrt}[1 - x^2]] + 4*\text{ArcTanh} \\ & [x]*\text{Log}[1 - x^2] + 8*\text{ArcTanh}[x]*\text{Log}[I*\text{Sinh}[\text{ArcTanh}[a/b] + \text{ArcTanh}[x]]] - 8 \\ & * \text{ArcTanh}[a/b]*\text{Log}[(2*I)*\text{Sinh}[\text{ArcTanh}[a/b] + \text{ArcTanh}[x]]] - 8*\text{ArcTanh}[x]*\text{Lo} \\ & \text{g}[(2*I)*\text{Sinh}[\text{ArcTanh}[a/b] + \text{ArcTanh}[x]]] - 4*\text{PolyLog}[2, -\text{E}^{(2*\text{ArcTanh}[x])}] \\ & - 4*\text{PolyLog}[2, \text{E}^{(-2*(\text{ArcTanh}[a/b] + \text{ArcTanh}[x]))}])]/(8*b) \end{aligned}$$

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {6472, 2849, 2752, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\text{arctanh}(x)}{a + bx} dx \\ & \quad \downarrow 6472 \\ & -\frac{\int \frac{\log\left(\frac{2(a+bx)}{(a+b)(x+1)}\right) dx}{b}}{b} + \frac{\int \frac{\log\left(\frac{2}{x+1}\right) dx}{b}}{b} + \frac{\text{arctanh}(x) \log\left(\frac{2(a+bx)}{(x+1)(a+b)}\right)}{b} - \frac{\text{arctanh}(x) \log\left(\frac{2}{x+1}\right)}{b} \\ & \quad \downarrow 2849 \\ & -\frac{\int \frac{\log\left(\frac{2(a+bx)}{(a+b)(x+1)}\right) dx}{b}}{b} + \frac{\int \frac{\log\left(\frac{2}{x+1}\right) d\frac{1}{x+1}}{b}}{b} + \frac{\text{arctanh}(x) \log\left(\frac{2(a+bx)}{(x+1)(a+b)}\right)}{b} - \frac{\text{arctanh}(x) \log\left(\frac{2}{x+1}\right)}{b} \\ & \quad \downarrow 2752 \\ & -\frac{\int \frac{\log\left(\frac{2(a+bx)}{(a+b)(x+1)}\right) dx}{b}}{b} + \frac{\text{arctanh}(x) \log\left(\frac{2(a+bx)}{(x+1)(a+b)}\right)}{b} - \frac{\text{arctanh}(x) \log\left(\frac{2}{x+1}\right)}{b} + \\ & \quad \frac{\text{PolyLog}\left(2, 1 - \frac{2}{x+1}\right)}{2b} \end{aligned}$$

$$\frac{\operatorname{arctanh}(x) \log\left(\frac{2(a+bx)}{(x+1)(a+b)}\right)}{b} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2(a+bx)}{(a+b)(x+1)}\right)}{2b} - \frac{\operatorname{arctanh}(x) \log\left(\frac{2}{x+1}\right)}{b} + \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{x+1}\right)}{2b}$$

input `Int[ArcTanh[x]/(a + b*x), x]`

output `-((ArcTanh[x]*Log[2/(1 + x)])/b) + (ArcTanh[x]*Log[(2*(a + b*x))/((a + b)*(1 + x))])/b + PolyLog[2, 1 - 2/(1 + x)]/(2*b) - PolyLog[2, 1 - (2*(a + b*x))/((a + b)*(1 + x))]/(2*b)`

Defintions of rubi rules used

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 2897 `Int[Log[u_]*(Pq_)^(m_.), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]]`

rule 6472 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-a + b*ArcTanh[c*x])*(Log[2/(1 + c*x)]/e), x] + (Simp[(a + b*ArcTanh[c*x])*(Log[2*c*((d + e*x)/((c*d + e)*(1 + c*x))])/e), x] + Simp[b*(c/e) Int[Log[2/(1 + c*x)]/(1 - c^2*x^2), x], x] - Simp[b*(c/e) Int[Log[2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/(1 - c^2*x^2), x], x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 - e^2, 0]`

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.23

method	result	size
default	$\frac{\ln(bx+a) \operatorname{arctanh}(x)}{b} - \frac{b \left(\operatorname{dilog}\left(\frac{bx+b}{-a+b}\right) + \ln(bx+a) \ln\left(\frac{bx+b}{-a+b}\right) \right)}{2} - \frac{b \left(\operatorname{dilog}\left(\frac{bx-b}{-a-b}\right) + \ln(bx+a) \ln\left(\frac{bx-b}{-a-b}\right) \right)}{2}$	106
parts	$\frac{\ln(bx+a) \operatorname{arctanh}(x)}{b} - \frac{b \left(\operatorname{dilog}\left(\frac{bx+b}{-a+b}\right) + \ln(bx+a) \ln\left(\frac{bx+b}{-a+b}\right) \right)}{2} - \frac{b \left(\operatorname{dilog}\left(\frac{bx-b}{-a-b}\right) + \ln(bx+a) \ln\left(\frac{bx-b}{-a-b}\right) \right)}{2}$	106
risch	$-\frac{\operatorname{dilog}\left(\frac{(1-x)b-a-b}{-a-b}\right)}{2b} - \frac{\ln(1-x) \ln\left(\frac{(1-x)b-a-b}{-a-b}\right)}{2b} + \frac{\operatorname{dilog}\left(\frac{(1+x)b+a-b}{a-b}\right)}{2b} + \frac{\ln(1+x) \ln\left(\frac{(1+x)b+a-b}{a-b}\right)}{2b}$	120

input `int(arctanh(x)/(b*x+a), x, method=_RETURNVERBOSE)`

output `ln(b*x+a)/b*arctanh(x)-1/b^2*(1/2*b*(dilog((b*x+b)/(-a+b))+ln(b*x+a)*ln((b*x+b)/(-a+b)))-1/2*b*(dilog((b*x-b)/(-a-b))+ln(b*x+a)*ln((b*x-b)/(-a-b))))`

Fricas [F]

$$\int \frac{\operatorname{arctanh}(x)}{a+bx} dx = \int \frac{\operatorname{artanh}(x)}{bx+a} dx$$

input `integrate(arctanh(x)/(b*x+a), x, algorithm="fricas")`

output `integral(arctanh(x)/(b*x + a), x)`

Sympy [F]

$$\int \frac{\operatorname{arctanh}(x)}{a+bx} dx = \int \frac{\operatorname{atanh}(x)}{a+bx} dx$$

input `integrate(atanh(x)/(b*x+a), x)`

output `Integral(atanh(x)/(a + b*x), x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.38

$$\int \frac{\operatorname{arctanh}(x)}{a + bx} dx = -\frac{(\log(x + 1) - \log(x - 1)) \log(bx + a)}{2b} + \frac{\operatorname{artanh}(x) \log(bx + a)}{b} - \frac{\log(x - 1) \log\left(\frac{bx-b}{a+b} + 1\right) + \operatorname{Li}_2\left(-\frac{bx-b}{a+b}\right)}{2b} + \frac{\log(x + 1) \log\left(\frac{bx+b}{a-b} + 1\right) + \operatorname{Li}_2\left(-\frac{bx+b}{a-b}\right)}{2b}$$

input `integrate(arctanh(x)/(b*x+a),x, algorithm="maxima")`

output `-1/2*(log(x + 1) - log(x - 1))*log(b*x + a)/b + arctanh(x)*log(b*x + a)/b - 1/2*(log(x - 1)*log((b*x - b)/(a + b) + 1) + dilog(-(b*x - b)/(a + b)))/b + 1/2*(log(x + 1)*log((b*x + b)/(a - b) + 1) + dilog(-(b*x + b)/(a - b)))/b`

Giac [F]

$$\int \frac{\operatorname{arctanh}(x)}{a + bx} dx = \int \frac{\operatorname{artanh}(x)}{bx + a} dx$$

input `integrate(arctanh(x)/(b*x+a),x, algorithm="giac")`

output `integrate(arctanh(x)/(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(x)}{a + bx} dx = \int \frac{\operatorname{atanh}(x)}{a + bx} dx$$

input `int(atanh(x)/(a + b*x), x)`output `int(atanh(x)/(a + b*x), x)`**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(x)}{a + bx} dx = \int \frac{\operatorname{atanh}(x)}{bx + a} dx$$

input `int(atanh(x)/(b*x+a), x)`output `int(atanh(x)/(a + b*x), x)`

3.24 $\int (d + ex)^3 (a + b \operatorname{arctanh}(cx^2)) dx$

Optimal result	221
Mathematica [A] (verified)	222
Rubi [A] (verified)	222
Maple [A] (verified)	224
Fricas [A] (verification not implemented)	225
Sympy [B] (verification not implemented)	225
Maxima [A] (verification not implemented)	226
Giac [A] (verification not implemented)	227
Mupad [B] (verification not implemented)	228
Reduce [B] (verification not implemented)	229

Optimal result

Integrand size = 18, antiderivative size = 154

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx^2)) dx = \frac{2bde^2x}{c} + \frac{be^3x^2}{4c} + \frac{bd(cd^2 - e^2) \arctan(\sqrt{cx})}{c^{3/2}} - \frac{bd(cd^2 + e^2) \operatorname{arctanh}(\sqrt{cx})}{c^{3/2}} - \frac{b(c^2d^4 + e^4) \operatorname{arctanh}(cx^2)}{4c^2e} + \frac{(d + ex)^4 (a + b \operatorname{arctanh}(cx^2))}{4e} + \frac{3bd^2e \log(1 - c^2x^4)}{4c}$$

output

```
2*b*d*e^2*x/c+1/4*b*e^3*x^2/c+b*d*(c*d^2-e^2)*arctan(c^(1/2)*x)/c^(3/2)-b*d*(c*d^2+e^2)*arctanh(c^(1/2)*x)/c^(3/2)-1/4*b*(c^2*d^4+e^4)*arctanh(c*x^2)/c^2/e+1/4*(e*x+d)^4*(a+b*arctanh(c*x^2))/e+3/4*b*d^2*e*ln(-c^2*x^4+1)/c
```

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 254, normalized size of antiderivative = 1.65

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx^2)) dx = \frac{1}{8} \left(\frac{8d(acd^2 + 2be^2)x}{c} + \frac{2e(6acd^2 + be^2)x^2}{c} + \frac{8ade^2x^3 + 2ae^3x^4}{c^{3/2}} + \frac{8bd(cd^2 - e^2) \arctan(\sqrt{cx})}{c^{3/2}} + 2bx(4d^3 + 6d^2ex + 4de^2x^2 + e^3x^3) \operatorname{arctanh}(cx^2) + \frac{b(4c^{3/2}d^3 + 4\sqrt{c}de^2 + e^3) \log(1 - \sqrt{cx})}{c^2} + \frac{b(-4c^2d^3 - 4cde^2 + \sqrt{c}e^3) \log(1 + \sqrt{cx})}{c^{5/2}} - \frac{be^3 \log(1 + cx^2)}{c^2} + \frac{6bd^2e \log(1 - c^2x^4)}{c} \right)$$

input `Integrate[(d + e*x)^3*(a + b*ArcTanh[c*x^2]),x]`

output `((8*d*(a*c*d^2 + 2*b*e^2)*x)/c + (2*e*(6*a*c*d^2 + b*e^2)*x^2)/c + 8*a*d*e^2*x^3 + 2*a*e^3*x^4 + (8*b*d*(c*d^2 - e^2)*ArcTan[Sqrt[c]*x])/c^(3/2) + 2*b*x*(4*d^3 + 6*d^2*e*x + 4*d*e^2*x^2 + e^3*x^3)*ArcTanh[c*x^2] + (b*(4*c^(3/2)*d^3 + 4*Sqrt[c]*d*e^2 + e^3)*Log[1 - Sqrt[c]*x])/c^2 + (b*(-4*c^2*d^3 - 4*c*d*e^2 + Sqrt[c]*e^3)*Log[1 + Sqrt[c]*x])/c^(5/2) - (b*e^3*Log[1 + c*x^2])/c^2 + (6*b*d^2*e*Log[1 - c^2*x^4])/c)/8`

Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.04, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6486, 2372, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int (d + ex)^3 (a + \operatorname{arctanh}(cx^2)) dx \\
& \quad \downarrow \text{6486} \\
& \frac{(d + ex)^4 (a + \operatorname{arctanh}(cx^2))}{4e} - \frac{bc \int \frac{x(d+ex)^4}{1-c^2x^4} dx}{2e} \\
& \quad \downarrow \text{2372} \\
& \frac{(d + ex)^4 (a + \operatorname{arctanh}(cx^2))}{4e} - \frac{bc \int \left(\frac{(4ed^3 + 4e^3x^2d)x^2}{1-c^2x^4} + \frac{(d^4 + 6e^2x^2d^2 + e^4x^4)x}{1-c^2x^4} \right) dx}{2e} \\
& \quad \downarrow \text{2009} \\
& \frac{(d + ex)^4 (a + \operatorname{arctanh}(cx^2))}{4e} - \\
& \frac{bc \left(-\frac{2de \arctan(\sqrt{cx})(cd^2 - e^2)}{c^{5/2}} + \frac{2de \operatorname{arctanh}(\sqrt{cx})(cd^2 + e^2)}{c^{5/2}} + \frac{\operatorname{arctanh}(cx^2)(c^2d^4 + e^4)}{2c^3} - \frac{3d^2e^2 \log(1 - c^2x^4)}{2c^2} - \frac{4de^3x}{c^2} - \frac{e^4x^2}{2c^2} \right)}{2e}
\end{aligned}$$

input `Int[(d + e*x)^3*(a + b*ArcTanh[c*x^2]),x]`

output `((d + e*x)^4*(a + b*ArcTanh[c*x^2]))/(4*e) - (b*c*((-4*d*e^3*x)/c^2 - (e^4*x^2)/(2*c^2) - (2*d*e*(c*d^2 - e^2)*ArcTan[Sqrt[c]*x])/c^(5/2) + (2*d*e*(c*d^2 + e^2)*ArcTanh[Sqrt[c]*x])/c^(5/2) + ((c^2*d^4 + e^4)*ArcTanh[c*x^2])/(2*c^3) - (3*d^2*e^2*Log[1 - c^2*x^4])/(2*c^2)))/(2*e)`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2372 `Int[(Pq_)*((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Module[{q = Expon[Pq, x], j, k}, Int[Sum[((c*x)^(m + j)/c^j)*Sum[Coeff[Pq, x, j + k*(n/2)]*x^(k*(n/2)), {k, 0, 2*((q - j)/n) + 1}]*(a + b*x^n)^p, {j, 0, n/2 - 1}], x] /; FreeQ[{a, b, c, m, p}, x] && PolyQ[Pq, x] && IGtQ[n/2, 0] && !PolyQ[Pq, x^(n/2)]`

rule 6486

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.)*((d_.) + (e_.)*(x_)^(m_.), x_
Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(e*(m + 1))), x]
- Simp[b*c*(n/(e*(m + 1))) Int[x^(n - 1)*((d + e*x)^(m + 1)/(1 - c^2*x^(2
*n))), x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 242, normalized size of antiderivative = 1.57

method	result
default	$\frac{a(ex+d)^4}{4e} + b \left(\frac{e^3 \operatorname{arctanh}(cx^2)x^4}{4} + e^2 \operatorname{arctanh}(cx^2)x^3d + \frac{3e \operatorname{arctanh}(cx^2)x^2d^2}{2} + \operatorname{arctanh}(cx^2)xd^3 + \dots \right)$
parts	$\frac{a(ex+d)^4}{4e} + b \left(\frac{e^3 \operatorname{arctanh}(cx^2)x^4}{4} + e^2 \operatorname{arctanh}(cx^2)x^3d + \frac{3e \operatorname{arctanh}(cx^2)x^2d^2}{2} + \operatorname{arctanh}(cx^2)xd^3 + \dots \right)$
risch	Expression too large to display

input

```
int((e*x+d)^3*(a+b*arctanh(c*x^2)),x,method=_RETURNVERBOSE)
```

output

```
1/4*a*(e*x+d)^4/e+b*(1/4*e^3*arctanh(c*x^2)*x^4+e^2*arctanh(c*x^2)*x^3*d+3
/2*e*arctanh(c*x^2)*x^2*d^2+arctanh(c*x^2)*x*d^3+1/4/e*arctanh(c*x^2)*d^4-
1/2/e*c*(-e^3/c^2*(1/2*e*x^2+4*d*x)+1/2/c^2*(1/2*(-c^2*d^4-6*c*d^2*e^2-e^4
)/c*ln(c*x^2-1)-(-4*c*d^3*e-4*d*e^3)/c^(1/2)*arctanh(c^(1/2)*x))+1/2/c^2*(
1/2*(c^2*d^4-6*c*d^2*e^2+e^4)/c*ln(c*x^2+1)+(-4*c*d^3*e+4*d*e^3)/c^(1/2)*a
rctan(c^(1/2)*x)))
```

Fricas [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 519, normalized size of antiderivative = 3.37

$$\int (d + ex)^3 (a + \operatorname{arctanh}(cx^2)) dx$$

$$= \left[\frac{2ac^2e^3x^4 + 8ac^2de^2x^3 + 2(6ac^2d^2e + bce^3)x^2 + 8(bcd^3 - bde^2)\sqrt{c} \arctan(\sqrt{cx}) + 4(bcd^3 + bde^2)\sqrt{c}}{\dots} \right]$$

input `integrate((e*x+d)^3*(a+b*arctanh(c*x^2)),x, algorithm="fricas")`

output `[1/8*(2*a*c^2*e^3*x^4 + 8*a*c^2*d*e^2*x^3 + 2*(6*a*c^2*d^2*e + b*c*e^3)*x^2 + 8*(b*c*d^3 - b*d*e^2)*sqrt(c)*arctan(sqrt(c)*x) + 4*(b*c*d^3 + b*d*e^2)*sqrt(c)*log((c*x^2 - 2*sqrt(c)*x + 1)/(c*x^2 - 1)) + 8*(a*c^2*d^3 + 2*b*c*d*e^2)*x + (6*b*c*d^2*e - b*e^3)*log(c*x^2 + 1) + (6*b*c*d^2*e + b*e^3)*log(c*x^2 - 1) + (b*c^2*e^3*x^4 + 4*b*c^2*d*e^2*x^3 + 6*b*c^2*d^2*e*x^2 + 4*b*c^2*d^3*x)*log(-(c*x^2 + 1)/(c*x^2 - 1)))/c^2, 1/8*(2*a*c^2*e^3*x^4 + 8*a*c^2*d*e^2*x^3 + 2*(6*a*c^2*d^2*e + b*c*e^3)*x^2 + 8*(b*c*d^3 + b*d*e^2)*sqrt(-c)*arctan(sqrt(-c)*x) + 4*(b*c*d^3 - b*d*e^2)*sqrt(-c)*log((c*x^2 + 2*sqrt(-c)*x - 1)/(c*x^2 + 1)) + 8*(a*c^2*d^3 + 2*b*c*d*e^2)*x + (6*b*c*d^2*e - b*e^3)*log(c*x^2 + 1) + (6*b*c*d^2*e + b*e^3)*log(c*x^2 - 1) + (b*c^2*e^3*x^4 + 4*b*c^2*d*e^2*x^3 + 6*b*c^2*d^2*e*x^2 + 4*b*c^2*d^3*x)*log(-(c*x^2 + 1)/(c*x^2 - 1)))/c^2]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2786 vs. 2(141) = 282.

Time = 7.48 (sec) , antiderivative size = 2786, normalized size of antiderivative = 18.09

$$\int (d + ex)^3 (a + \operatorname{arctanh}(cx^2)) dx = \text{Too large to display}$$

input `integrate((e*x+d)**3*(a+b*atanh(c*x**2)),x)`

output

```
Piecewise((8*a*c**2*d**3*x*sqrt(-1/c)*sqrt(1/c)/(2*c**3*(-1/c)**(3/2)*sqrt(1/c) - 2*c**3*sqrt(-1/c)*(1/c)**(3/2) + 12*c**2*sqrt(-1/c)*sqrt(1/c)) + 12*a*c**2*d**2*e*x**2*sqrt(-1/c)*sqrt(1/c)/(2*c**3*(-1/c)**(3/2)*sqrt(1/c) - 2*c**3*sqrt(-1/c)*(1/c)**(3/2) + 12*c**2*sqrt(-1/c)*sqrt(1/c)) + 8*a*c**2*d*e**2*x**3*sqrt(-1/c)*sqrt(1/c)/(2*c**3*(-1/c)**(3/2)*sqrt(1/c) - 2*c**3*sqrt(-1/c)*(1/c)**(3/2) + 12*c**2*sqrt(-1/c)*sqrt(1/c)) + 2*a*c**2*e**3*x**4*sqrt(-1/c)*sqrt(1/c)/(2*c**3*(-1/c)**(3/2)*sqrt(1/c) - 2*c**3*sqrt(-1/c)*(1/c)**(3/2) + 12*c**2*sqrt(-1/c)*sqrt(1/c)) + 8*b*c**2*d**3*x*sqrt(-1/c)*sqrt(1/c)*atanh(c*x**2)/(2*c**3*(-1/c)**(3/2)*sqrt(1/c) - 2*c**3*sqrt(-1/c)*(1/c)**(3/2) + 12*c**2*sqrt(-1/c)*sqrt(1/c)) - 2*b*c**2*d**3*(-1/c)**(3/2)*log(x + sqrt(-1/c))/(2*c**3*(-1/c)**(3/2)*sqrt(1/c) - 2*c**3*sqrt(-1/c)*(1/c)**(3/2) + 12*c**2*sqrt(-1/c)*sqrt(1/c)) + 2*b*c**2*d**3*(1/c)**(3/2)*log(x + sqrt(-1/c))/(2*c**3*(-1/c)**(3/2)*sqrt(1/c) - 2*c**3*sqrt(-1/c)*(1/c)**(3/2) + 12*c**2*sqrt(-1/c)*sqrt(1/c)) + 12*b*c**2*d**2*e*x**2*sqrt(-1/c)*sqrt(1/c)*atanh(c*x**2)/(2*c**3*(-1/c)**(3/2)*sqrt(1/c) - 2*c**3*sqrt(-1/c)*(1/c)**(3/2) + 12*c**2*sqrt(-1/c)*sqrt(1/c)) + 3*b*c**2*d**2*e*(-1/c)**(3/2)*sqrt(1/c)*log(x + sqrt(-1/c))/(2*c**3*(-1/c)**(3/2)*sqrt(1/c) - 2*c**3*sqrt(-1/c)*(1/c)**(3/2) + 12*c**2*sqrt(-1/c)*sqrt(1/c)) - 3*b*c**2*d**2*e*sqrt(-1/c)*(1/c)**(3/2)*log(x + sqrt(-1/c))/(2*c**3*(-1/c)**(3/2)*sqrt(1/c) - 2*c**3*sqrt(-1/c)*(1/c)**(3/2) + 12*c**2*sqrt(-1/c)*sqrt(1/c)) - 2*c**3*sqrt(-1/c)*(1/c)**(3/2) + 12*c**2*sqrt(-1/c)*sqrt(1/c)...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.56

$$\begin{aligned}
 & \int (d + ex)^3 (a + b \operatorname{arctanh}(cx^2)) dx \\
 &= \frac{1}{4} ae^3 x^4 + ade^2 x^3 + \frac{3}{2} ad^2 ex^2 \\
 &+ \frac{1}{2} \left(c \left(\frac{2 \arctan(\sqrt{cx})}{c^{\frac{3}{2}}} + \frac{\log\left(\frac{cx-\sqrt{c}}{cx+\sqrt{c}}\right)}{c^{\frac{3}{2}}} \right) + 2x \operatorname{artanh}(cx^2) \right) bd^3 \\
 &+ \frac{1}{2} \left(2x^3 \operatorname{artanh}(cx^2) + c \left(\frac{4x}{c^2} - \frac{2 \arctan(\sqrt{cx})}{c^{\frac{5}{2}}} + \frac{\log\left(\frac{cx-\sqrt{c}}{cx+\sqrt{c}}\right)}{c^{\frac{5}{2}}} \right) \right) bde^2 \\
 &+ \frac{1}{8} \left(2x^4 \operatorname{artanh}(cx^2) + c \left(\frac{2x^2}{c^2} - \frac{\log(cx^2+1)}{c^3} + \frac{\log(cx^2-1)}{c^3} \right) \right) be^3 \\
 &+ ad^3 x + \frac{3(2cx^2 \operatorname{artanh}(cx^2) + \log(-c^2x^4+1))bd^2e}{4c}
 \end{aligned}$$

input `integrate((e*x+d)^3*(a+b*arctanh(c*x^2)),x, algorithm="maxima")`

output
$$\begin{aligned} & 1/4*a*e^3*x^4 + a*d*e^2*x^3 + 3/2*a*d^2*e*x^2 + 1/2*(c*(2*\arctan(\sqrt{c}*x) \\ &)/c^{(3/2)} + \log((c*x - \sqrt{c})/(c*x + \sqrt{c}))/c^{(3/2)}) + 2*x*\arctanh(c* \\ & x^2))*b*d^3 + 1/2*(2*x^3*\arctanh(c*x^2) + c*(4*x/c^2 - 2*\arctan(\sqrt{c}*x) \\ & /c^{(5/2)} + \log((c*x - \sqrt{c})/(c*x + \sqrt{c}))/c^{(5/2)}))*b*d*e^2 + 1/8*(2 \\ & *x^4*\arctanh(c*x^2) + c*(2*x^2/c^2 - \log(c*x^2 + 1)/c^3 + \log(c*x^2 - 1)/c \\ & ^3))*b*e^3 + a*d^3*x + 3/4*(2*c*x^2*\arctanh(c*x^2) + \log(-c^2*x^4 + 1))*b* \\ & d^2*e/c \end{aligned}$$

Giac [A] (verification not implemented)

Time = 7.47 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.47

$$\begin{aligned} & \int (d + ex)^3 (a + b \operatorname{arctanh}(cx^2)) dx \\ & = \frac{1}{4} a e^3 x^4 + a d e^2 x^3 + \frac{(6 a c d^2 e + b e^3) x^2}{4 c} \\ & + \frac{1}{8} (b e^3 x^4 + 4 b d e^2 x^3 + 6 b d^2 e x^2 + 4 b d^3 x) \log \left(-\frac{c x^2 + 1}{c x^2 - 1} \right) + \frac{(a c d^3 + 2 b d e^2) x}{c} \\ & + \frac{(b c d^3 - b d e^2) \arctan(\sqrt{c x})}{c^{\frac{3}{2}}} + \frac{(b c d^3 + b d e^2) \arctan\left(\frac{c x}{\sqrt{-c}}\right)}{\sqrt{-c c}} \\ & + \frac{(6 b c d^2 e - b e^3) \log(c x^2 + 1)}{8 c^2} + \frac{(6 b c d^2 e + b e^3) \log(c x^2 - 1)}{8 c^2} \end{aligned}$$

input `integrate((e*x+d)^3*(a+b*arctanh(c*x^2)),x, algorithm="giac")`

output
$$\begin{aligned} & 1/4*a*e^3*x^4 + a*d*e^2*x^3 + 1/4*(6*a*c*d^2*e + b*e^3)*x^2/c + 1/8*(b*e^3 \\ & *x^4 + 4*b*d*e^2*x^3 + 6*b*d^2*e*x^2 + 4*b*d^3*x)*\log(-(c*x^2 + 1)/(c*x^2 \\ & - 1)) + (a*c*d^3 + 2*b*d*e^2)*x/c + (b*c*d^3 - b*d*e^2)*\arctan(\sqrt{c}*x)/ \\ & c^{(3/2)} + (b*c*d^3 + b*d*e^2)*\arctan(c*x/\sqrt{-c})/(\sqrt{-c}*c) + 1/8*(6*b \\ & *c*d^2*e - b*e^3)*\log(c*x^2 + 1)/c^2 + 1/8*(6*b*c*d^2*e + b*e^3)*\log(c*x^2 \\ & - 1)/c^2 \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 4.70 (sec) , antiderivative size = 823, normalized size of antiderivative = 5.34

$$\begin{aligned}
& \int (d + ex)^3 (a + \operatorname{arctanh}(cx^2)) dx \\
&= \ln(cx^2 + 1) \left(\frac{bd^3x}{2} + \frac{3bd^2ex^2}{4} + \frac{bde^2x^3}{2} + \frac{be^3x^4}{8} \right) \\
&\quad - \ln(1 - cx^2) \left(\frac{bd^3x}{2} + \frac{3bd^2ex^2}{4} + \frac{bde^2x^3}{2} + \frac{be^3x^4}{8} \right) + \frac{ae^3x^4}{4} \\
&\quad - \frac{\ln(8c^5d^6 - c^2e^6 - 4de^5\sqrt{-c^5} + e^6x\sqrt{-c^5} + 8c^3d^2e^4 + 4c^4d^3e^3x - 4c^3de^5x + 4cd^3e^3\sqrt{-c^5} - 8c^4}{8c^4}}{\ln(8c^5d^6 - c^2e^6 + 4de^5\sqrt{-c^5} - e^6x\sqrt{-c^5} + 8c^3d^2e^4 + 4c^4d^3e^3x - 4c^3de^5x - 4cd^3e^3\sqrt{-c^5} + 8c^4)} \\
&\quad + \frac{x(2ac^2d^3 + 4bcde^2)}{2c^2} \\
&\quad + \frac{\ln(64c^2d^{12}(c^5)^{7/2} + 128d^8e^4(c^5)^{7/2} - 64c^{20}d^{12}x - c^{14}e^{12}x + ce^{12}(c^5)^{5/2} - 32c^{16}d^4e^8x - 128c^{18}}{8c^4}}{\ln(8c^{10}d^6 + c^7e^6 + 8c^8d^2e^4 - 4de^5(c^5)^{3/2} + e^6x(c^5)^{3/2} - 4c^9d^3e^3x - 4cd^3e^3(c^5)^{3/2} + 8c^3d^6x)} \\
&\quad + \frac{x^2(6ac^2d^2e + bce^3)}{4c^2} + ade^2x^3
\end{aligned}$$

input `int((a + b*atanh(c*x^2))*(d + e*x)^3,x)`

output

```

log(c*x^2 + 1)*((b*e^3*x^4)/8 + (b*d^3*x)/2 + (3*b*d^2*e*x^2)/4 + (b*d*e^2
*x^3)/2) - log(1 - c*x^2)*((b*e^3*x^4)/8 + (b*d^3*x)/2 + (3*b*d^2*e*x^2)/4
+ (b*d*e^2*x^3)/2) + (a*e^3*x^4)/4 - (log(8*c^5*d^6 - c^2*e^6 - 4*d*e^5*(
-c^5)^(1/2) + e^6*x*(-c^5)^(1/2) + 8*c^3*d^2*e^4 + 4*c^4*d^3*e^3*x - 4*c^3
*d*e^5*x + 4*c*d^3*e^3*(-c^5)^(1/2) - 8*c^3*d^6*x*(-c^5)^(1/2) - 8*c*d^2*e
^4*x*(-c^5)^(1/2))*(b*c^2*e^3 - 4*b*c*d^3*(-c^5)^(1/2) + 4*b*d*e^2*(-c^5)^(
1/2) - 6*b*c^3*d^2*e))/(8*c^4) - (log(8*c^5*d^6 - c^2*e^6 + 4*d*e^5*(-c^5)
^(1/2) - e^6*x*(-c^5)^(1/2) + 8*c^3*d^2*e^4 + 4*c^4*d^3*e^3*x - 4*c^3*d*e
^5*x - 4*c*d^3*e^3*(-c^5)^(1/2) + 8*c^3*d^6*x*(-c^5)^(1/2) + 8*c*d^2*e^4*x
*(-c^5)^(1/2))*(b*c^2*e^3 + 4*b*c*d^3*(-c^5)^(1/2) - 4*b*d*e^2*(-c^5)^(1/2)
- 6*b*c^3*d^2*e))/(8*c^4) + (x*(2*a*c^2*d^3 + 4*b*c*d*e^2))/(2*c^2) + (1
og(64*c^2*d^12*(c^5)^(7/2) + 128*d^8*e^4*(c^5)^(7/2) - 64*c^20*d^12*x - c^
14*e^12*x + c*e^12*(c^5)^(5/2) - 32*c^16*d^4*e^8*x - 128*c^18*d^8*e^4*x +
32*c^3*d^4*e^8*(c^5)^(5/2))*(b*c^2*e^3 + 4*b*c*d^3*(c^5)^(1/2) + 4*b*d*e^2
*(c^5)^(1/2) + 6*b*c^3*d^2*e))/(8*c^4) + (log(8*c^10*d^6 + c^7*e^6 + 8*c^8
*d^2*e^4 - 4*d*e^5*(c^5)^(3/2) + e^6*x*(c^5)^(3/2) - 4*c^9*d^3*e^3*x - 4*c
*d^3*e^3*(c^5)^(3/2) + 8*c^3*d^6*x*(c^5)^(3/2) - 4*c^8*d*e^5*x + 8*c*d^2*e
^4*x*(c^5)^(3/2))*(b*c^2*e^3 - 4*b*c*d^3*(c^5)^(1/2) - 4*b*d*e^2*(c^5)^(1/
2) + 6*b*c^3*d^2*e))/(8*c^4) + (x^2*(b*c*e^3 + 6*a*c^2*d^2*e))/(4*c^2) + a
*d*e^2*x^3

```

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 314, normalized size of antiderivative = 2.04

$$\int (d + ex)^3 (a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{4\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) bcd^3 - 4\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) bde^2 + 4\sqrt{c} \operatorname{atanh}(cx^2) bcd^3 + 4\sqrt{c} \operatorname{atanh}(cx^2) bde^2 + 4 \operatorname{atanh}(cx^2) a d^3}{1}$$

input

```
int((e*x+d)^3*(a+b*atanh(c*x^2)),x)
```

output

```
(4*sqrt(c)*atan((c*x)/sqrt(c))*b*c*d**3 - 4*sqrt(c)*atan((c*x)/sqrt(c))*b*
d*e**2 + 4*sqrt(c)*atanh(c*x**2)*b*c*d**3 + 4*sqrt(c)*atanh(c*x**2)*b*d*e*
*2 + 4*atanh(c*x**2)*b*c**2*d**3*x + 6*atanh(c*x**2)*b*c**2*d**2*e*x**2 +
4*atanh(c*x**2)*b*c**2*d*e**2*x**3 + atanh(c*x**2)*b*c**2*e**3*x**4 - 6*at
anh(c*x**2)*b*c*d**2*e - atanh(c*x**2)*b*e**3 + 4*sqrt(c)*log(sqrt(c)*x -
1)*b*c*d**3 + 4*sqrt(c)*log(sqrt(c)*x - 1)*b*d*e**2 - 2*sqrt(c)*log(c*x**2
+ 1)*b*c*d**3 - 2*sqrt(c)*log(c*x**2 + 1)*b*d*e**2 + 6*log(c*x**2 + 1)*b*
c*d**2*e + 4*a*c**2*d**3*x + 6*a*c**2*d**2*e*x**2 + 4*a*c**2*d*e**2*x**3 +
a*c**2*e**3*x**4 + 8*b*c*d*e**2*x + b*c*e**3*x**2)/(4*c**2)
```

3.25 $\int (d + ex)^2 (a + b \operatorname{arctanh}(cx^2)) dx$

Optimal result	231
Mathematica [A] (verified)	232
Rubi [A] (verified)	232
Maple [A] (verified)	234
Fricas [A] (verification not implemented)	234
Sympy [B] (verification not implemented)	235
Maxima [A] (verification not implemented)	236
Giac [A] (verification not implemented)	237
Mupad [B] (verification not implemented)	238
Reduce [B] (verification not implemented)	239

Optimal result

Integrand size = 18, antiderivative size = 158

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx^2)) dx = \frac{2be^2x}{3c} + \frac{b(3cd^2 - e^2) \arctan(\sqrt{cx})}{3c^{3/2}} - \frac{b(3cd^2 + e^2) \operatorname{arctanh}(\sqrt{cx})}{3c^{3/2}} + \frac{(d + ex)^3 (a + b \operatorname{arctanh}(cx^2))}{3e} + \frac{bd(cd^2 + 3e^2) \log(1 - cx^2)}{6ce} - \frac{bd(cd^2 - 3e^2) \log(1 + cx^2)}{6ce}$$

output

```
2/3*b*e^2*x/c+1/3*b*(3*c*d^2-e^2)*arctan(c^(1/2)*x)/c^(3/2)-1/3*b*(3*c*d^2
+e^2)*arctanh(c^(1/2)*x)/c^(3/2)+1/3*(e*x+d)^3*(a+b*arctanh(c*x^2))/e+1/6*
b*d*(c*d^2+3*e^2)*ln(-c*x^2+1)/c/e-1/6*b*d*(c*d^2-3*e^2)*ln(c*x^2+1)/c/e
```


Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.08

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx^2)) dx = \frac{1}{6} \left(6ad^2x + \frac{4be^2x}{c} + 6adex^2 + 2ae^2x^3 + \frac{2b(3cd^2 - e^2) \arctan(\sqrt{cx})}{c^{3/2}} + 2bx(3d^2 + 3dex + e^2x^2) \operatorname{arctanh}(cx^2) + \frac{b(3cd^2 + e^2) \log(1 - \sqrt{cx})}{c^{3/2}} - \frac{b(3cd^2 + e^2) \log(1 + \sqrt{cx})}{c^{3/2}} + \frac{3bde \log(1 - c^2x^4)}{c} \right)$$

input

```
Integrate[(d + e*x)^2*(a + b*ArcTanh[c*x^2]),x]
```

output

```
(6*a*d^2*x + (4*b*e^2*x)/c + 6*a*d*e*x^2 + 2*a*e^2*x^3 + (2*b*(3*c*d^2 - e^2)*ArcTan[Sqrt[c]*x])/c^(3/2) + 2*b*x*(3*d^2 + 3*d*e*x + e^2*x^2)*ArcTanh[c*x^2] + (b*(3*c*d^2 + e^2)*Log[1 - Sqrt[c]*x])/c^(3/2) - (b*(3*c*d^2 + e^2)*Log[1 + Sqrt[c]*x])/c^(3/2) + (3*b*d*e*Log[1 - c^2*x^4])/c/6
```

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.89, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6486, 2370, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx^2)) dx$$

↓ 6486

$$\frac{(d+ex)^3(a+\operatorname{arctanh}(cx^2))}{3e} - \frac{2bc \int \frac{x(d+ex)^3}{1-c^2x^4} dx}{3e}$$

↓ 2370

$$\frac{(d+ex)^3(a+\operatorname{arctanh}(cx^2))}{3e} - \frac{2bc \int \left(\frac{(x^2e^3+3d^2e)x^2}{1-c^2x^4} + \frac{(d^3+3e^2x^2d)x}{1-c^2x^4} \right) dx}{3e}$$

↓ 2009

$$\frac{(d+ex)^3(a+\operatorname{arctanh}(cx^2))}{3e} - \frac{2bc \left(-\frac{e \operatorname{arctan}(\sqrt{cx})(3cd^2-e^2)}{2c^{5/2}} + \frac{e \operatorname{arctanh}(\sqrt{cx})(3cd^2+e^2)}{2c^{5/2}} + \frac{d^3 \operatorname{arctanh}(cx^2)}{2c} - \frac{3de^2 \log(1-c^2x^4)}{4c^2} - \frac{e^3x}{c^2} \right)}{3e}$$

input `Int[(d + e*x)^2*(a + b*ArcTanh[c*x^2]), x]`

output `((d + e*x)^3*(a + b*ArcTanh[c*x^2]))/(3*e) - (2*b*c*(-((e^3*x)/c^2) - (e*(3*c*d^2 - e^2)*ArcTan[Sqrt[c]*x])/(2*c^(5/2))) + (e*(3*c*d^2 + e^2)*ArcTanh[Sqrt[c]*x])/(2*c^(5/2)) + (d^3*ArcTanh[c*x^2])/(2*c) - (3*d*e^2*Log[1 - c^2*x^4])/(4*c^2))/(3*e)`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2370 `Int[((Pq_)*((c_)*(x_)^(m_)))/((a_) + (b_)*(x_)^(n_)), x_Symbol] := With[{v = Sum[(c*x)^(m + ii)*((Coeff[Pq, x, ii] + Coeff[Pq, x, n/2 + ii])*x^(n/2))]/(c^ii*(a + b*x^n))}, {ii, 0, n/2 - 1}], Int[v, x] /; SumQ[v] /; FreeQ[{a, b, c, m}, x] && PolyQ[Pq, x] && IGtQ[n/2, 0] && Expon[Pq, x] < n`

rule 6486 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)*((d_) + (e_)*(x_)^(m_)), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(e*(m + 1))), x] - Simp[b*c*(n/(e*(m + 1))) Int[x^(n - 1)*((d + e*x)^(m + 1)/(1 - c^2*x^(2*n)))] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.26

method	result
default	$\frac{a(ex+d)^3}{3e} + b \left(\frac{e^2 \operatorname{arctanh}(cx^2)x^3}{3} + e \operatorname{arctanh}(cx^2)x^2d + \operatorname{arctanh}(cx^2)xd^2 + \frac{\operatorname{arctanh}(cx^2)d^3}{3e} - \frac{2c}{3} \left(\frac{d^2}{e} \operatorname{arctanh}(cx^2) + \frac{d}{e} \operatorname{arctanh}(cx^2) \right) \right)$
parts	$\frac{a(ex+d)^3}{3e} + b \left(\frac{e^2 \operatorname{arctanh}(cx^2)x^3}{3} + e \operatorname{arctanh}(cx^2)x^2d + \operatorname{arctanh}(cx^2)xd^2 + \frac{\operatorname{arctanh}(cx^2)d^3}{3e} - \frac{2c}{3} \left(\frac{d^2}{e} \operatorname{arctanh}(cx^2) + \frac{d}{e} \operatorname{arctanh}(cx^2) \right) \right)$
risch	Expression too large to display

input `int((e*x+d)^2*(a+b*arctanh(c*x^2)),x,method=_RETURNVERBOSE)`

output `1/3*a*(e*x+d)^3/e+b*(1/3*e^2*arctanh(c*x^2)*x^3+e*arctanh(c*x^2)*x^2*d+arctanh(c*x^2)*x*d^2+1/3/e*arctanh(c*x^2)*d^3-2/3/e*c*(-e^3/c^2*x+1/2/c^2*(1/2*(-c^2*d^3-3*c*d*e^2)/c*ln(c*x^2-1)-(-3*c*d^2*e-e^3)/c^(1/2)*arctanh(c^(1/2)*x))+1/2/c^2*(1/2*(c^2*d^3-3*c*d*e^2)/c*ln(c*x^2+1)+(-3*c*d^2*e+e^3)/c^(1/2)*arctan(c^(1/2)*x)))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 401, normalized size of antiderivative = 2.54

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \left[\frac{2ac^2e^2x^3 + 6ac^2dex^2 + 3bcde \log(cx^2 + 1) + 3bcde \log(cx^2 - 1) + 2(3bcd^2 - be^2)\sqrt{c} \operatorname{arctan}(\sqrt{cx})}{3e} \right]$$

input `integrate((e*x+d)^2*(a+b*arctanh(c*x^2)),x, algorithm="fricas")`

output

```
[1/6*(2*a*c^2*e^2*x^3 + 6*a*c^2*d*e*x^2 + 3*b*c*d*e*log(c*x^2 + 1) + 3*b*c*d*e*log(c*x^2 - 1) + 2*(3*b*c*d^2 - b*e^2)*sqrt(c)*arctan(sqrt(c)*x) + (3*b*c*d^2 + b*e^2)*sqrt(c)*log((c*x^2 - 2*sqrt(c)*x + 1)/(c*x^2 - 1)) + 2*(3*a*c^2*d^2 + 2*b*c*e^2)*x + (b*c^2*e^2*x^3 + 3*b*c^2*d*e*x^2 + 3*b*c^2*d^2*x)*log(-(c*x^2 + 1)/(c*x^2 - 1)))/c^2, 1/6*(2*a*c^2*e^2*x^3 + 6*a*c^2*d*e*x^2 + 3*b*c*d*e*log(c*x^2 + 1) + 3*b*c*d*e*log(c*x^2 - 1) + 2*(3*b*c*d^2 + b*e^2)*sqrt(-c)*arctan(sqrt(-c)*x) + (3*b*c*d^2 - b*e^2)*sqrt(-c)*log((c*x^2 + 2*sqrt(-c)*x - 1)/(c*x^2 + 1)) + 2*(3*a*c^2*d^2 + 2*b*c*e^2)*x + (b*c^2*e^2*x^3 + 3*b*c^2*d*e*x^2 + 3*b*c^2*d^2*x)*log(-(c*x^2 + 1)/(c*x^2 - 1)))/c^2]
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3465 vs. $2(139) = 278$.

Time = 6.06 (sec) , antiderivative size = 3465, normalized size of antiderivative = 21.93

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx^2)) dx = \text{Too large to display}$$

input

```
integrate((e*x+d)**2*(a+b*atanh(c*x**2)),x)
```

output

```

Piecewise((12*a*c**2*d**2*x*sqrt(-1/c)/(12*c**3*(-1/c)**(3/2) - 12*c**3*(1/c)**(3/2) + 24*c**2*sqrt(-1/c) + 24*c**2*sqrt(1/c)) + 12*a*c**2*d**2*x*sqrt(1/c)/(12*c**3*(-1/c)**(3/2) - 12*c**3*(1/c)**(3/2) + 24*c**2*sqrt(-1/c) + 24*c**2*sqrt(1/c)) + 12*a*c**2*d*e*x**2*sqrt(-1/c)/(12*c**3*(-1/c)**(3/2) - 12*c**3*(1/c)**(3/2) + 24*c**2*sqrt(-1/c) + 24*c**2*sqrt(1/c)) + 12*a*c**2*d*e*x**2*sqrt(1/c)/(12*c**3*(-1/c)**(3/2) - 12*c**3*(1/c)**(3/2) + 24*c**2*sqrt(-1/c) + 24*c**2*sqrt(1/c)) + 4*a*c**2*e**2*x**3*sqrt(-1/c)/(12*c**3*(-1/c)**(3/2) - 12*c**3*(1/c)**(3/2) + 24*c**2*sqrt(-1/c) + 24*c**2*sqrt(1/c)) + 4*a*c**2*e**2*x**3*sqrt(1/c)/(12*c**3*(-1/c)**(3/2) - 12*c**3*(1/c)**(3/2) + 24*c**2*sqrt(-1/c) + 24*c**2*sqrt(1/c)) - 3*b*c**3*d**2*(-1/c)**(3/2)*sqrt(1/c)*log(x - sqrt(-1/c))/(12*c**3*(-1/c)**(3/2) - 12*c**3*(1/c)**(3/2) + 24*c**2*sqrt(-1/c) + 24*c**2*sqrt(1/c)) - 12*b*c**3*d**2*(-1/c)**(3/2)*sqrt(1/c)*log(x + sqrt(-1/c))/(12*c**3*(-1/c)**(3/2) - 12*c**3*(1/c)**(3/2) + 24*c**2*sqrt(-1/c) + 24*c**2*sqrt(1/c)) + 3*b*c**3*d**2*(-1/c)**(3/2)*sqrt(1/c)*log(x - sqrt(1/c))/(12*c**3*(-1/c)**(3/2) - 12*c**3*(1/c)**(3/2) + 24*c**2*sqrt(-1/c) + 24*c**2*sqrt(1/c)) + 3*b*c**3*d**2*sqrt(-1/c)*(1/c)**(3/2)*log(x - sqrt(-1/c))/(12*c**3*(-1/c)**(3/2) - 12*c**3*(1/c)**(3/2) + 24*c**2*sqrt(-1/c) + 24*c**2*sqrt(1/c)) - 12*b*c**3*d**2*sqrt(-1/c)*(1/c)**(3/2)*log(x + sqrt(-1/c))/(12*c**3*(-1/c)**(3/2) - 12*c**3*(1/c)**(3/2) + 24*c**2*sqrt(-1/c) + 24*c**2*sqrt(1/c)) - 3*b*c**3*d**...

```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.08

$$\begin{aligned}
& \int (d + ex)^2 (a + b \operatorname{arctanh}(cx^2)) dx \\
&= \frac{1}{3} ae^2 x^3 + adex^2 + \frac{1}{2} \left(c \left(\frac{2 \arctan(\sqrt{cx})}{c^{\frac{3}{2}}} + \frac{\log\left(\frac{cx - \sqrt{c}}{cx + \sqrt{c}}\right)}{c^{\frac{3}{2}}} \right) + 2x \operatorname{artanh}(cx^2) \right) bd^2 \\
&+ \frac{1}{6} \left(2x^3 \operatorname{artanh}(cx^2) + c \left(\frac{4x}{c^2} - \frac{2 \arctan(\sqrt{cx})}{c^{\frac{5}{2}}} + \frac{\log\left(\frac{cx - \sqrt{c}}{cx + \sqrt{c}}\right)}{c^{\frac{5}{2}}} \right) \right) be^2 \\
&+ ad^2 x + \frac{(2cx^2 \operatorname{artanh}(cx^2) + \log(-c^2 x^4 + 1))bde}{2c}
\end{aligned}$$

input

```
integrate((e*x+d)^2*(a+b*arctanh(c*x^2)),x, algorithm="maxima")
```

output

```
1/3*a*e^2*x^3 + a*d*e*x^2 + 1/2*(c*(2*arctan(sqrt(c)*x)/c^(3/2) + log((c*x
- sqrt(c))/(c*x + sqrt(c)))/c^(3/2)) + 2*x*arctanh(c*x^2))*b*d^2 + 1/6*(2
*x^3*arctanh(c*x^2) + c*(4*x/c^2 - 2*arctan(sqrt(c)*x)/c^(5/2) + log((c*x
- sqrt(c))/(c*x + sqrt(c)))/c^(5/2)))*b*e^2 + a*d^2*x + 1/2*(2*c*x^2*arcta
nh(c*x^2) + log(-c^2*x^4 + 1))*b*d*e/c
```

Giac [A] (verification not implemented)

Time = 1.10 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.08

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx^2)) dx = \frac{1}{3} ae^2 x^3 + adex^2 + \frac{bde \log(cx^2 + 1)}{2c}$$

$$+ \frac{bde \log(cx^2 - 1)}{2c}$$

$$+ \frac{1}{6} (be^2 x^3 + 3bde x^2 + 3bd^2 x) \log\left(-\frac{cx^2 + 1}{cx^2 - 1}\right)$$

$$+ \frac{(3acd^2 + 2be^2)x}{3c}$$

$$+ \frac{(3bcd^2 - be^2) \arctan(\sqrt{cx})}{3c^{\frac{3}{2}}}$$

$$+ \frac{(3bcd^2 + be^2) \arctan\left(\frac{cx}{\sqrt{-c}}\right)}{3\sqrt{-cc}}$$

input

```
integrate((e*x+d)^2*(a+b*arctanh(c*x^2)),x, algorithm="giac")
```

output

```
1/3*a*e^2*x^3 + a*d*e*x^2 + 1/2*b*d*e*log(c*x^2 + 1)/c + 1/2*b*d*e*log(c*x
^2 - 1)/c + 1/6*(b*e^2*x^3 + 3*b*d*e*x^2 + 3*b*d^2*x)*log(-(c*x^2 + 1)/(c*
x^2 - 1)) + 1/3*(3*a*c*d^2 + 2*b*e^2)*x/c + 1/3*(3*b*c*d^2 - b*e^2)*arctan
(sqrt(c)*x)/c^(3/2) + 1/3*(3*b*c*d^2 + b*e^2)*arctan(c*x/sqrt(-c))/(sqrt(-
c)*c)
```

Mupad [B] (verification not implemented)

Time = 3.94 (sec) , antiderivative size = 309, normalized size of antiderivative = 1.96

$$\begin{aligned}
& \int (d + ex)^2 (a + b \operatorname{arctanh}(cx^2)) dx \\
&= \ln(cx^2 + 1) \left(\frac{bd^2x}{2} + \frac{bde x^2}{2} + \frac{be^2 x^3}{6} \right) \\
&\quad - \ln(1 - cx^2) \left(\frac{bd^2x}{2} + \frac{bde x^2}{2} + \frac{be^2 x^3}{6} \right) + \frac{x(3ac^2d^2 + 2bce^2)}{3c^2} \\
&\quad + \frac{ae^2 x^3}{3} - \frac{\ln(c + x\sqrt{c^3}) (be^2\sqrt{c^3} + 3bcd^2\sqrt{c^3} - 3bc^2de)}{6c^3} \\
&\quad + \frac{\ln(c - x\sqrt{c^3}) (be^2\sqrt{c^3} + 3bcd^2\sqrt{c^3} + 3bc^2de)}{6c^3} \\
&\quad + \frac{\ln(c + x\sqrt{-c^3}) (be^2\sqrt{-c^3} + 3bc^2de - 3bcd^2\sqrt{-c^3})}{6c^3} \\
&\quad + \frac{\ln(c - x\sqrt{-c^3}) (3bc^2de - be^2\sqrt{-c^3} + 3bcd^2\sqrt{-c^3})}{6c^3} + a d e x^2
\end{aligned}$$

input `int((a + b*atanh(c*x^2))*(d + e*x)^2,x)`output `log(c*x^2 + 1)*((b*e^2*x^3)/6 + (b*d^2*x)/2 + (b*d*e*x^2)/2) - log(1 - c*x^2)*((b*e^2*x^3)/6 + (b*d^2*x)/2 + (b*d*e*x^2)/2) + (x*(3*a*c^2*d^2 + 2*b*c*e^2))/(3*c^2) + (a*e^2*x^3)/3 - (log(c + x*(c^3)^(1/2))*(b*e^2*(c^3)^(1/2) + 3*b*c*d^2*(c^3)^(1/2) - 3*b*c^2*d*e))/(6*c^3) + (log(c - x*(c^3)^(1/2))*(b*e^2*(c^3)^(1/2) + 3*b*c*d^2*(c^3)^(1/2) + 3*b*c^2*d*e))/(6*c^3) + (log(c + x*(-c^3)^(1/2))*(b*e^2*(-c^3)^(1/2) + 3*b*c^2*d*e - 3*b*c*d^2*(-c^3)^(1/2)))/(6*c^3) + (log(c - x*(-c^3)^(1/2))*(3*b*c^2*d*e - b*e^2*(-c^3)^(1/2) + 3*b*c*d^2*(-c^3)^(1/2)))/(6*c^3) + a*d*e*x^2`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.58

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{6\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) b c d^2 - 2\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) b e^2 + 6\sqrt{c} \operatorname{atanh}(cx^2) b c d^2 + 2\sqrt{c} \operatorname{atanh}(cx^2) b e^2 + 6 \operatorname{atanh}(cx^2)}$$

input

```
int((e*x+d)^2*(a+b*atanh(c*x^2)),x)
```

output

```
(6*sqrt(c)*atan((c*x)/sqrt(c))*b*c*d**2 - 2*sqrt(c)*atan((c*x)/sqrt(c))*b*
e**2 + 6*sqrt(c)*atanh(c*x**2)*b*c*d**2 + 2*sqrt(c)*atanh(c*x**2)*b*e**2 +
6*atanh(c*x**2)*b*c**2*d**2*x + 6*atanh(c*x**2)*b*c**2*d*e*x**2 + 2*atanh
(c*x**2)*b*c**2*e**2*x**3 - 6*atanh(c*x**2)*b*c*d*e + 6*sqrt(c)*log(sqrt(c)
)*x - 1)*b*c*d**2 + 2*sqrt(c)*log(sqrt(c)*x - 1)*b*e**2 - 3*sqrt(c)*log(c*
x**2 + 1)*b*c*d**2 - sqrt(c)*log(c*x**2 + 1)*b*e**2 + 6*log(c*x**2 + 1)*b*
c*d*e + 6*a*c**2*d**2*x + 6*a*c**2*d*e*x**2 + 2*a*c**2*e**2*x**3 + 4*b*c*e
**2*x)/(6*c**2)
```


3.26 $\int (d + ex) (a + b \operatorname{arctanh}(cx^2)) dx$

Optimal result	240
Mathematica [A] (verified)	241
Rubi [A] (verified)	241
Maple [A] (verified)	242
Fricas [A] (verification not implemented)	243
Sympy [A] (verification not implemented)	244
Maxima [A] (verification not implemented)	244
Giac [A] (verification not implemented)	245
Mupad [B] (verification not implemented)	246
Reduce [B] (verification not implemented)	247

Optimal result

Integrand size = 16, antiderivative size = 117

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^2)) dx = \frac{bd \arctan(\sqrt{cx})}{\sqrt{c}} - \frac{bd \operatorname{arctanh}(\sqrt{cx})}{\sqrt{c}} + \frac{(d + ex)^2 (a + b \operatorname{arctanh}(cx^2))}{2e} + \frac{b(cd^2 + e^2) \log(1 - cx^2)}{4ce} - \frac{b(cd^2 - e^2) \log(1 + cx^2)}{4ce}$$

```
output b*d*arctan(c^(1/2)*x)/c^(1/2)-b*d*arctanh(c^(1/2)*x)/c^(1/2)+1/2*(e*x+d)^2
*(a+b*arctanh(c*x^2))/e+1/4*b*(c*d^2+e^2)*ln(-c*x^2+1)/c/e-1/4*b*(c*d^2-e^
2)*ln(c*x^2+1)/c/e
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.89

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^2)) dx$$

$$= adx + \frac{1}{2} aex^2 + bdx \operatorname{arctanh}(cx^2) + \frac{1}{2} bex^2 \operatorname{arctanh}(cx^2)$$

$$+ \frac{bd(2 \arctan(\sqrt{cx}) + \log(1 - \sqrt{cx}) - \log(1 + \sqrt{cx}))}{2\sqrt{c}} + \frac{be \log(1 - c^2x^4)}{4c}$$

input `Integrate[(d + e*x)*(a + b*ArcTanh[c*x^2]), x]`

output `a*d*x + (a*e*x^2)/2 + b*d*x*ArcTanh[c*x^2] + (b*e*x^2*ArcTanh[c*x^2])/2 + (b*d*(2*ArcTan[Sqrt[c]*x] + Log[1 - Sqrt[c]*x] - Log[1 + Sqrt[c]*x]))/(2*Sqrt[c]) + (b*e*Log[1 - c^2*x^4])/(4*c)`

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.88, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6486, 2370, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^2)) dx$$

$$\downarrow \text{6486}$$

$$\frac{(d + ex)^2 (a + b \operatorname{arctanh}(cx^2))}{2e} - \frac{bc \int \frac{x(d+ex)^2}{1-c^2x^4} dx}{e}$$

$$\downarrow \text{2370}$$

$$\frac{(d + ex)^2 (a + b \operatorname{arctanh}(cx^2))}{2e} - \frac{bc \int \left(\frac{2dex^2}{1-c^2x^4} + \frac{(d^2+e^2x^2)x}{1-c^2x^4} \right) dx}{e}$$

$$\downarrow \text{2009}$$

$$\frac{(d + ex)^2 (a + b \operatorname{arctanh}(cx^2))}{2e} - \frac{bc \left(-\frac{de \arctan(\sqrt{cx})}{c^{3/2}} + \frac{de \operatorname{arctanh}(\sqrt{cx})}{c^{3/2}} + \frac{d^2 \operatorname{arctanh}(cx^2)}{2c} - \frac{e^2 \log(1 - c^2 x^4)}{4c^2} \right)}{e}$$

input `Int[(d + e*x)*(a + b*ArcTanh[c*x^2]),x]`

output `((d + e*x)^2*(a + b*ArcTanh[c*x^2]))/(2*e) - (b*c*(-((d*e*ArcTan[Sqrt[c]*x])/c^(3/2)) + (d*e*ArcTanh[Sqrt[c]*x])/c^(3/2) + (d^2*ArcTanh[c*x^2])/(2*c) - (e^2*Log[1 - c^2*x^4])/(4*c^2)))/e`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2370 `Int[((Pq_)*((c_)*(x_)^(m_)))/((a_) + (b_)*(x_)^(n_)), x_Symbol] := With[{v = Sum[(c*x)^(m + ii)*((Coeff[Pq, x, ii] + Coeff[Pq, x, n/2 + ii]*x^(n/2)))/(c^ii*(a + b*x^n))], {ii, 0, n/2 - 1}}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b, c, m}, x] && PolyQ[Pq, x] && IGtQ[n/2, 0] && Expon[Pq, x] < n`

rule 6486 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)*((d_) + (e_)*(x_)^(m_)), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(e*(m + 1))), x] - Simp[b*c*(n/(e*(m + 1))) Int[x^(n - 1)*((d + e*x)^(m + 1)/(1 - c^2*x^(2*n)))]], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.79

method	result
default	$a\left(\frac{1}{2}ex^2 + dx\right) + \frac{b \operatorname{arctanh}(cx^2)x^2e}{2} + b \operatorname{arctanh}(cx^2) dx + \frac{be \ln(cx^2-1)}{4c} - \frac{bd \operatorname{arctanh}(\sqrt{cx})}{\sqrt{c}} + \frac{be \ln(cx^2+1)}{4c}$
parts	$a\left(\frac{1}{2}ex^2 + dx\right) + \frac{b \operatorname{arctanh}(cx^2)x^2e}{2} + b \operatorname{arctanh}(cx^2) dx + \frac{be \ln(cx^2-1)}{4c} - \frac{bd \operatorname{arctanh}(\sqrt{cx})}{\sqrt{c}} + \frac{be \ln(cx^2+1)}{4c}$
risch	$\frac{bx(ex+2d) \ln(cx^2+1)}{4} - \frac{be x^2 \ln(-cx^2+1)}{4} - \frac{bdx \ln(-cx^2+1)}{2} + \frac{ae x^2}{2} - \frac{b \ln(1+\sqrt{cx})d}{2\sqrt{c}} + \frac{b \ln(1-\sqrt{-cx})\sqrt{-c}d}{2c} - \frac{b \ln(cx^2-1)}{4c}$

input `int((e*x+d)*(a+b*arctanh(c*x^2)),x,method=_RETURNVERBOSE)`

output `a*(1/2*e*x^2+d*x)+1/2*b*arctanh(c*x^2)*x^2*e+b*arctanh(c*x^2)*d*x+1/4*b*e/c*ln(c*x^2-1)-b*d*arctanh(c^(1/2)*x)/c^(1/2)+1/4*b*e/c*ln(c*x^2+1)+b*d*arctanh(c^(1/2)*x)/c^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 249, normalized size of antiderivative = 2.13

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{2acex^2 + 4acdx + 4b\sqrt{cd} \arctan(\sqrt{cx}) + 2b\sqrt{cd} \log\left(\frac{cx^2 - 2\sqrt{cx} + 1}{cx^2 - 1}\right) + be \log(cx^2 + 1) + be \log(cx^2 - 1)}{4c}$$

input `integrate((e*x+d)*(a+b*arctanh(c*x^2)),x, algorithm="fricas")`

output `[1/4*(2*a*c*e*x^2 + 4*a*c*d*x + 4*b*sqrt(c)*d*arctan(sqrt(c)*x) + 2*b*sqrt(c)*d*log((c*x^2 - 2*sqrt(c)*x + 1)/(c*x^2 - 1)) + b*e*log(c*x^2 + 1) + b*e*log(c*x^2 - 1) + (b*c*e*x^2 + 2*b*c*d*x)*log(-(c*x^2 + 1)/(c*x^2 - 1)))/c, 1/4*(2*a*c*e*x^2 + 4*a*c*d*x + 4*b*sqrt(-c)*d*arctan(sqrt(-c)*x) - 2*b*sqrt(-c)*d*log((c*x^2 - 2*sqrt(-c)*x - 1)/(c*x^2 + 1)) + b*e*log(c*x^2 + 1) + b*e*log(c*x^2 - 1) + (b*c*e*x^2 + 2*b*c*d*x)*log(-(c*x^2 + 1)/(c*x^2 - 1)))/c]`

Sympy [A] (verification not implemented)

Time = 4.89 (sec) , antiderivative size = 753, normalized size of antiderivative = 6.44

$$\int (d + ex) (a + \operatorname{barctanh}(cx^2)) dx = \text{Too large to display}$$

input `integrate((e*x+d)*(a+b*atanh(c*x**2)),x)`

output

```
a*d*x + a*e*x**2/2 + b*d*Piecewise((4*c*x*sqrt(-1/c)*sqrt(1/c)*atanh(c*x**2)/(c**2*(-1/c)**(3/2)*sqrt(1/c) - c**2*sqrt(-1/c)*(1/c)**(3/2) + 6*c*sqrt(-1/c)*sqrt(1/c)) - c*(-1/c)**(3/2)*log(x + sqrt(-1/c))/(c**2*(-1/c)**(3/2)*sqrt(1/c) - c**2*sqrt(-1/c)*(1/c)**(3/2) + 6*c*sqrt(-1/c)*sqrt(1/c)) + c*(1/c)**(3/2)*log(x + sqrt(-1/c))/(c**2*(-1/c)**(3/2)*sqrt(1/c) - c**2*sqrt(-1/c)*(1/c)**(3/2) + 6*c*sqrt(-1/c)*sqrt(1/c)) - 2*sqrt(-1/c)*log(x - sqrt(-1/c))/(c**2*(-1/c)**(3/2)*sqrt(1/c) - c**2*sqrt(-1/c)*(1/c)**(3/2) + 6*c*sqrt(-1/c)*sqrt(1/c)) - 3*sqrt(-1/c)*log(x + sqrt(-1/c))/(c**2*(-1/c)**(3/2)*sqrt(1/c) - c**2*sqrt(-1/c)*(1/c)**(3/2) + 6*c*sqrt(-1/c)*sqrt(1/c)) + 4*sqrt(-1/c)*log(x - sqrt(1/c))/(c**2*(-1/c)**(3/2)*sqrt(1/c) - c**2*sqrt(-1/c)*(1/c)**(3/2) + 6*c*sqrt(-1/c)*sqrt(1/c)) + 4*sqrt(-1/c)*atanh(c*x**2)/(c**2*(-1/c)**(3/2)*sqrt(1/c) - c**2*sqrt(-1/c)*(1/c)**(3/2) + 6*c*sqrt(-1/c)*sqrt(1/c)) + 2*sqrt(1/c)*log(x - sqrt(-1/c))/(c**2*(-1/c)**(3/2)*sqrt(1/c) - c**2*sqrt(-1/c)*(1/c)**(3/2) + 6*c*sqrt(-1/c)*sqrt(1/c)) - 3*sqrt(1/c)*log(x + sqrt(-1/c))/(c**2*(-1/c)**(3/2)*sqrt(1/c) - c**2*sqrt(-1/c)*(1/c)**(3/2) + 6*c*sqrt(-1/c)*sqrt(1/c)), Ne(c, 0)), (0, True)) + b*e*Piecewise((x**2*atanh(c*x**2)/2 + log(c*x**2 + 1)/(2*c) - atanh(c*x**2)/(2*c), Ne(c, 0)), (0, True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.81

$$\begin{aligned} & \int (d + ex) (a + \operatorname{barctanh}(cx^2)) dx \\ &= \frac{1}{2} aex^2 + \frac{1}{2} \left(c \left(\frac{2 \arctan(\sqrt{c}x)}{c^{\frac{3}{2}}} + \frac{\log\left(\frac{cx-\sqrt{c}}{cx+\sqrt{c}}\right)}{c^{\frac{3}{2}}} \right) + 2x \operatorname{artanh}(cx^2) \right) bd \\ &+ adx + \frac{(2cx^2 \operatorname{artanh}(cx^2) + \log(-c^2x^4 + 1))be}{4c} \end{aligned}$$

input `integrate((e*x+d)*(a+b*arctanh(c*x^2)),x, algorithm="maxima")`

output `1/2*a*e*x^2 + 1/2*(c*(2*arctan(sqrt(c)*x)/c^(3/2) + log((c*x - sqrt(c))/(c*x + sqrt(c))))/c^(3/2)) + 2*x*arctanh(c*x^2)*b*d + a*d*x + 1/4*(2*c*x^2*a*rctanh(c*x^2) + log(-c^2*x^4 + 1))*b*e/c`

Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.14

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^2)) dx = \frac{1}{2} aex^2 + adx + \frac{bcd \operatorname{arctan}(x\sqrt{|c|})}{|c|^{\frac{3}{2}}} + \frac{be \log\left(x^2 + \frac{1}{|c|}\right)}{4c} + \frac{1}{4} (bex^2 + 2bdx) \log\left(-\frac{cx^2 + 1}{cx^2 - 1}\right) - \frac{(2bcd\sqrt{|c|} - bce) \log\left(\left|x + \frac{1}{\sqrt{|c|}}\right|\right)}{4c^2} + \frac{(2bcd\sqrt{|c|} + bce) \log\left(\left|x - \frac{1}{\sqrt{|c|}}\right|\right)}{4c^2}$$

input `integrate((e*x+d)*(a+b*arctanh(c*x^2)),x, algorithm="giac")`

output `1/2*a*e*x^2 + a*d*x + b*c*d*arctan(x*sqrt(abs(c)))/abs(c)^(3/2) + 1/4*b*e*log(x^2 + 1/abs(c))/c + 1/4*(b*e*x^2 + 2*b*d*x)*log(-(c*x^2 + 1)/(c*x^2 - 1)) - 1/4*(2*b*c*d*sqrt(abs(c)) - b*c*e)*log(abs(x + 1/sqrt(abs(c))))/c^2 + 1/4*(2*b*c*d*sqrt(abs(c)) + b*c*e)*log(abs(x - 1/sqrt(abs(c))))/c^2`

Mupad [B] (verification not implemented)

Time = 3.84 (sec) , antiderivative size = 242, normalized size of antiderivative = 2.07

$$\begin{aligned}
\int (d + ex) (a + \operatorname{barctanh}(cx^2)) dx &= a dx + \frac{ae x^2}{2} + \frac{bdx \ln(cx^2 + 1)}{2} \\
&\quad - \frac{bdx \ln(1 - cx^2)}{2} + \frac{be \ln(c + x\sqrt{c^3})}{4c} \\
&\quad + \frac{be \ln(c - x\sqrt{c^3})}{4c} + \frac{be x^2 \ln(cx^2 + 1)}{4} \\
&\quad - \frac{be x^2 \ln(1 - cx^2)}{4} + \frac{be \ln(c + x\sqrt{-c^3})}{4c} \\
&\quad + \frac{be \ln(c - x\sqrt{-c^3})}{4c} - \frac{bd \ln(c + x\sqrt{c^3}) \sqrt{c^3}}{2c^2} \\
&\quad + \frac{bd \ln(c - x\sqrt{c^3}) \sqrt{c^3}}{2c^2} \\
&\quad - \frac{bd \ln(c + x\sqrt{-c^3}) \sqrt{-c^3}}{2c^2} \\
&\quad + \frac{bd \ln(c - x\sqrt{-c^3}) \sqrt{-c^3}}{2c^2}
\end{aligned}$$

input `int((a + b*atanh(c*x^2))*(d + e*x),x)`output

```

a*d*x + (a*e*x^2)/2 + (b*d*x*log(c*x^2 + 1))/2 - (b*d*x*log(1 - c*x^2))/2
+ (b*e*log(c + x*(c^3)^(1/2)))/(4*c) + (b*e*log(c - x*(c^3)^(1/2)))/(4*c)
+ (b*e*x^2*log(c*x^2 + 1))/4 - (b*e*x^2*log(1 - c*x^2))/4 + (b*e*log(c + x
*(-c^3)^(1/2)))/(4*c) + (b*e*log(c - x*(-c^3)^(1/2)))/(4*c) - (b*d*log(c +
x*(c^3)^(1/2))*(c^3)^(1/2))/(2*c^2) + (b*d*log(c - x*(c^3)^(1/2))*(c^3)^(
1/2))/(2*c^2) - (b*d*log(c + x*(-c^3)^(1/2))*(-c^3)^(1/2))/(2*c^2) + (b*d*
log(c - x*(-c^3)^(1/2))*(-c^3)^(1/2))/(2*c^2)

```

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.01

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^2)) dx$$

$$= \frac{2\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) bd + 2\sqrt{c} \operatorname{atanh}(cx^2) bd + 2 \operatorname{atanh}(cx^2) bcdx + \operatorname{atanh}(cx^2) bce x^2 - \operatorname{atanh}(cx^2) be + 2\sqrt{c}}{2c}$$

input

```
int((e*x+d)*(a+b*atanh(c*x^2)),x)
```

output

```
(2*sqrt(c)*atan((c*x)/sqrt(c))*b*d + 2*sqrt(c)*atanh(c*x**2)*b*d + 2*atanh
(c*x**2)*b*c*d*x + atanh(c*x**2)*b*c*e*x**2 - atanh(c*x**2)*b*e + 2*sqrt(c
)*log(sqrt(c)*x - 1)*b*d - sqrt(c)*log(c*x**2 + 1)*b*d + log(c*x**2 + 1)*b
*e + 2*a*c*d*x + a*c*e*x**2)/(2*c)
```


3.27 $\int \frac{a+b\operatorname{arctanh}(cx^2)}{d+ex} dx$

Optimal result	248
Mathematica [C] (verified)	249
Rubi [A] (verified)	250
Maple [A] (verified)	251
Fricas [F]	252
Sympy [F(-1)]	252
Maxima [F]	253
Giac [F]	253
Mupad [F(-1)]	253
Reduce [F]	254

Optimal result

Integrand size = 18, antiderivative size = 325

$$\int \frac{a + b\operatorname{arctanh}(cx^2)}{d + ex} dx = \frac{(a + b\operatorname{arctanh}(cx^2)) \log(d + ex)}{e} - \frac{b \log\left(\frac{e(1-\sqrt{-cx})}{\sqrt{-cd+e}}\right) \log(d + ex)}{2e} - \frac{b \log\left(-\frac{e(1+\sqrt{-cx})}{\sqrt{-cd-e}}\right) \log(d + ex)}{2e} + \frac{b \log\left(\frac{e(1-\sqrt{cx})}{\sqrt{cd+e}}\right) \log(d + ex)}{2e} + \frac{b \log\left(-\frac{e(1+\sqrt{cx})}{\sqrt{cd-e}}\right) \log(d + ex)}{2e} - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{-c}(d+ex)}{\sqrt{-cd-e}}\right)}{2e} + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{c}(d+ex)}{\sqrt{cd-e}}\right)}{2e} - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{-c}(d+ex)}{\sqrt{-cd+e}}\right)}{2e} + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{c}(d+ex)}{\sqrt{cd+e}}\right)}{2e}$$

output

```
(a+b*arctanh(c*x^2))*ln(e*x+d)/e-1/2*b*ln(e*(1-(-c)^(1/2)*x)/((-c)^(1/2)*d+e))*ln(e*x+d)/e-1/2*b*ln(-e*(1+(-c)^(1/2)*x)/((-c)^(1/2)*d-e))*ln(e*x+d)/e+1/2*b*ln(e*(1-c^(1/2)*x)/(c^(1/2)*d+e))*ln(e*x+d)/e+1/2*b*ln(-e*(1+c^(1/2)*x)/(c^(1/2)*d-e))*ln(e*x+d)/e-1/2*b*polylog(2,(-c)^(1/2)*(e*x+d)/((-c)^(1/2)*d-e))/e+1/2*b*polylog(2,c^(1/2)*(e*x+d)/(c^(1/2)*d-e))/e-1/2*b*polylog(2,(-c)^(1/2)*(e*x+d)/((-c)^(1/2)*d+e))/e+1/2*b*polylog(2,c^(1/2)*(e*x+d)/(c^(1/2)*d+e))/e
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 17.33 (sec) , antiderivative size = 285, normalized size of antiderivative = 0.88

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{d + ex} dx = \frac{a \log(d + ex)}{e} + \frac{b \left(2 \operatorname{arctanh}(cx^2) \log(d + ex) - \log\left(\frac{e(i - \sqrt{cx})}{\sqrt{cd + ie}}\right) \log(d + ex) - \log\left(-\frac{e(i + \sqrt{cx})}{\sqrt{cd - ie}}\right) \log(d + ex) + \log\left(-\frac{e(1 - \sqrt{cx})}{\sqrt{cd + ie}}\right) \log(d + ex) + \log\left(-\frac{e(1 + \sqrt{cx})}{\sqrt{cd - ie}}\right) \log(d + ex) \right)}{e}$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])/(d + e*x), x]
```

output

```
(a*Log[d + e*x])/e + (b*(2*ArcTanh[c*x^2]*Log[d + e*x] - Log[(e*(I - Sqrt[c]*x))/(Sqrt[c]*d + I*e)]*Log[d + e*x] - Log[-((e*(I + Sqrt[c]*x))/(Sqrt[c]*d - I*e))]*Log[d + e*x] + Log[-((e*(1 + Sqrt[c]*x))/(Sqrt[c]*d - e))]*Log[d + e*x] + Log[d + e*x]*Log[(e - Sqrt[c]*e*x)/(Sqrt[c]*d + e)] + PolyLog[2, (Sqrt[c]*(d + e*x))/(Sqrt[c]*d - e)] - PolyLog[2, (Sqrt[c]*(d + e*x))/(Sqrt[c]*d + I*e)] + PolyLog[2, (Sqrt[c]*(d + e*x))/(Sqrt[c]*d + e)]))/(2*e)
```

Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 325, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6482, 2863, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barctanh}(cx^2)}{d + ex} dx \\
 & \quad \downarrow \text{6482} \\
 & \frac{\log(d + ex) (a + \operatorname{barctanh}(cx^2))}{e} - \frac{2bc \int \frac{x \log(d+ex)}{1-c^2x^4} dx}{e} \\
 & \quad \downarrow \text{2863} \\
 & \frac{\log(d + ex) (a + \operatorname{barctanh}(cx^2))}{e} - \frac{2bc \int \left(\frac{cx \log(d+ex)}{2(c-c^2x^2)} + \frac{cx \log(d+ex)}{2(c^2x^2+c)} \right) dx}{e} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\log(d + ex) (a + \operatorname{barctanh}(cx^2))}{e} - \\
 & \frac{2bc \left(\frac{\operatorname{PolyLog}\left(2, \frac{\sqrt{-c}(d+ex)}{\sqrt{-cd-e}}\right)}{4c} - \frac{\operatorname{PolyLog}\left(2, \frac{\sqrt{c}(d+ex)}{\sqrt{cd-e}}\right)}{4c} + \frac{\operatorname{PolyLog}\left(2, \frac{\sqrt{-c}(d+ex)}{\sqrt{-cd+e}}\right)}{4c} - \frac{\operatorname{PolyLog}\left(2, \frac{\sqrt{c}(d+ex)}{\sqrt{cd+e}}\right)}{4c} + \frac{\log(d+ex) \log\left(\frac{e(1-\sqrt{-c}x)}{\sqrt{-cd+e}}\right)}{4c} \right)}{e}
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c*x^2])/(d + e*x),x]`

output `((a + b*ArcTanh[c*x^2])*Log[d + e*x])/e - (2*b*c*((Log[(e*(1 - Sqrt[-c]*x))/(Sqrt[-c]*d + e)]*Log[d + e*x])/(4*c) + (Log[-((e*(1 + Sqrt[-c]*x))/(Sqrt[-c]*d - e))]*Log[d + e*x])/(4*c) - (Log[(e*(1 - Sqrt[c]*x))/(Sqrt[c]*d + e)]*Log[d + e*x])/(4*c) - (Log[-((e*(1 + Sqrt[c]*x))/(Sqrt[c]*d - e))]*Log[d + e*x])/(4*c) + PolyLog[2, (Sqrt[-c]*(d + e*x))/(Sqrt[-c]*d - e])/(4*c) - PolyLog[2, (Sqrt[c]*(d + e*x))/(Sqrt[c]*d - e])/(4*c) + PolyLog[2, (Sqrt[-c]*(d + e*x))/(Sqrt[-c]*d + e])/(4*c) - PolyLog[2, (Sqrt[c]*(d + e*x))/(Sqrt[c]*d + e])/(4*c)))/e`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2863 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]*(b_.))^(p_.)*((h_.)*(x_)^(m_.))*((f_) + (g_.)*(x_)^(r_.))^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log[c*(d + e*x)^n])^p, (h*x)^m*(f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n, p, q, r}, x] && IntegerQ[m] && IntegerQ[q]`

rule 6482 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.)/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[Log[d + e*x]*((a + b*ArcTanh[c*x^n])/e), x] - Simp[b*c*(n/e) Int[x^(n - 1)*(Log[d + e*x]/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, d, e, n}, x] && IntegerQ[n]`

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 347, normalized size of antiderivative = 1.07

method	result
default	$\frac{a \ln(ex+d)}{e} + b \left(\frac{\ln(ex+d) \operatorname{arctanh}(cx^2)}{e} - \frac{e^2 \left(\frac{\ln(ex+d) \left(\ln \left(\frac{e\sqrt{c} - (ex+d)c+cd}{e\sqrt{c+cd}} \right) + \ln \left(\frac{e\sqrt{c} + (ex+d)c-cd}{e\sqrt{c-cd}} \right) \right)}{2c} \right) + \operatorname{dilog} \left(\frac{e\sqrt{c} - (ex+d)c+cd}{e\sqrt{c+cd}} \right)}{2} \right)$
parts	$\frac{a \ln(ex+d)}{e} + b \left(\frac{\ln(ex+d) \operatorname{arctanh}(cx^2)}{e} - \frac{e^2 \left(\frac{\ln(ex+d) \left(\ln \left(\frac{e\sqrt{c} - (ex+d)c+cd}{e\sqrt{c+cd}} \right) + \ln \left(\frac{e\sqrt{c} + (ex+d)c-cd}{e\sqrt{c-cd}} \right) \right)}{2c} \right) + \operatorname{dilog} \left(\frac{e\sqrt{c} - (ex+d)c+cd}{e\sqrt{c+cd}} \right)}{2} \right)$
risch	$\frac{a \ln(ex+d)}{e} - \frac{b \ln(ex+d) \ln(-cx^2+1)}{2e} + \frac{b \ln(ex+d) \ln \left(\frac{e\sqrt{c} - (ex+d)c+cd}{e\sqrt{c+cd}} \right)}{2e} + \frac{b \ln(ex+d) \ln \left(\frac{e\sqrt{c} + (ex+d)c-cd}{e\sqrt{c-cd}} \right)}{2e} + \frac{b \operatorname{dilog} \left(\frac{e\sqrt{c} - (ex+d)c+cd}{e\sqrt{c+cd}} \right)}{2e}$

input `int((a+b*arctanh(c*x^2))/(e*x+d),x,method=_RETURNVERBOSE)`

output `a*ln(e*x+d)/e+b*(ln(e*x+d)/e*arctanh(c*x^2)-2/e^3*c*(-1/2*e^2*(1/2*ln(e*x+d)*ln((e*c^(1/2)-(e*x+d)*c+c*d)/(e*c^(1/2)+c*d))+ln((e*c^(1/2)+(e*x+d)*c-c*d)/(e*c^(1/2)-c*d)))/c+1/2*(dilog((e*c^(1/2)-(e*x+d)*c+c*d)/(e*c^(1/2)+c*d))+dilog((e*c^(1/2)+(e*x+d)*c-c*d)/(e*c^(1/2)-c*d)))/c)+1/2*e^2*(1/2*ln(e*x+d)*(ln((e*(-c)^(1/2)-(e*x+d)*c+c*d)/(e*(-c)^(1/2)+c*d))+ln((e*(-c)^(1/2)+(e*x+d)*c-c*d)/(e*(-c)^(1/2)-c*d)))/c+1/2*(dilog((e*(-c)^(1/2)-(e*x+d)*c+c*d)/(e*(-c)^(1/2)+c*d))+dilog((e*(-c)^(1/2)+(e*x+d)*c-c*d)/(e*(-c)^(1/2)-c*d)))/c))`

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{d + ex} dx = \int \frac{b \operatorname{artanh}(cx^2) + a}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x^2))/(e*x+d),x, algorithm="fricas")`

output `integral((b*arctanh(c*x^2) + a)/(e*x + d), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{d + ex} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**2))/(e*x+d),x)`

output `Timed out`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{d + ex} dx = \int \frac{b \operatorname{artanh}(cx^2) + a}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x^2))/(e*x+d),x, algorithm="maxima")`

output `1/2*b*integrate((log(c*x^2 + 1) - log(-c*x^2 + 1))/(e*x + d), x) + a*log(e*x + d)/e`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{d + ex} dx = \int \frac{b \operatorname{artanh}(cx^2) + a}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x^2))/(e*x+d),x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)/(e*x + d), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{d + ex} dx = \int \frac{a + b \operatorname{atanh}(cx^2)}{d + ex} dx$$

input `int((a + b*atanh(c*x^2))/(d + e*x),x)`

output `int((a + b*atanh(c*x^2))/(d + e*x), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{d + ex} dx = \frac{\left(\int \frac{\operatorname{atanh}(cx^2)}{ex+d} dx \right) be + \log(ex + d) a}{e}$$

input `int((a+b*atanh(c*x^2))/(e*x+d),x)`

output `(int(atanh(c*x**2)/(d + e*x),x)*b*e + log(d + e*x)*a)/e`

3.28 $\int \frac{a+b\operatorname{arctanh}(cx^2)}{(d+ex)^2} dx$

Optimal result	255
Mathematica [A] (verified)	256
Rubi [A] (verified)	256
Maple [A] (verified)	258
Fricas [B] (verification not implemented)	258
Sympy [F(-1)]	259
Maxima [A] (verification not implemented)	260
Giac [A] (verification not implemented)	260
Mupad [B] (verification not implemented)	262
Reduce [B] (verification not implemented)	263

Optimal result

Integrand size = 18, antiderivative size = 166

$$\int \frac{a + b\operatorname{arctanh}(cx^2)}{(d + ex)^2} dx = \frac{b\sqrt{c} \arctan(\sqrt{cx})}{cd^2 + e^2} - \frac{b\sqrt{c}\operatorname{arctanh}(\sqrt{cx})}{cd^2 - e^2} - \frac{a + b\operatorname{arctanh}(cx^2)}{e(d + ex)} + \frac{2bcde \log(d + ex)}{c^2d^4 - e^4} - \frac{bcd \log(1 - cx^2)}{2e(cd^2 - e^2)} + \frac{bcd \log(1 + cx^2)}{2e(cd^2 + e^2)}$$

output

```
b*c^(1/2)*arctan(c^(1/2)*x)/(c*d^2+e^2)-b*c^(1/2)*arctanh(c^(1/2)*x)/(c*d^2-e^2)-(a+b*arctanh(c*x^2))/e/(e*x+d)+2*b*c*d*e*ln(e*x+d)/(c^2*d^4-e^4)-1/2*b*c*d*ln(-c*x^2+1)/e/(c*d^2-e^2)+1/2*b*c*d*ln(c*x^2+1)/e/(c*d^2+e^2)
```


Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.57

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^2} dx = \frac{1}{2} \left(-\frac{2a}{e(d + ex)} + \frac{2b\sqrt{c} \arctan(\sqrt{cx})}{cd^2 + e^2} - \frac{2b \operatorname{arctanh}(cx^2)}{e(d + ex)} \right. \\ \left. + \frac{b\sqrt{c}(c^{3/2}d^3 - cd^2e - e^3) \log(1 - \sqrt{cx})}{-c^2d^4e + e^5} \right. \\ \left. + \frac{b\sqrt{c}(c^{3/2}d^3 + cd^2e + e^3) \log(1 + \sqrt{cx})}{-c^2d^4e + e^5} \right. \\ \left. + \frac{4bcde \log(d + ex)}{c^2d^4 - e^4} + \frac{bc^2d^3 \log(1 + cx^2)}{c^2d^4e - e^5} \right. \\ \left. - \frac{bcde \log(1 - c^2x^4)}{c^2d^4 - e^4} \right)$$

input

```
Integrate[(a + b*ArcTanh[c*x^2])/(d + e*x)^2,x]
```

output

```
((-2*a)/(e*(d + e*x)) + (2*b*Sqrt[c]*ArcTan[Sqrt[c]*x])/(c*d^2 + e^2) - (2
*b*ArcTanh[c*x^2])/(e*(d + e*x)) + (b*Sqrt[c]*(c^(3/2)*d^3 - c*d^2*e - e^3
)*Log[1 - Sqrt[c]*x])/(-c^2*d^4*e) + e^5) + (b*Sqrt[c]*(c^(3/2)*d^3 + c*d
^2*e + e^3)*Log[1 + Sqrt[c]*x])/(-c^2*d^4*e) + e^5) + (4*b*c*d*e*Log[d +
e*x])/(c^2*d^4 - e^4) + (b*c^2*d^3*Log[1 + c*x^2])/(c^2*d^4*e - e^5) - (b*
c*d*e*Log[1 - c^2*x^4])/(c^2*d^4 - e^4))/2
```

Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.01, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6486, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^2} dx$$

↓ 6486

$$\frac{2bc \int \frac{x}{(d+ex)(1-c^2x^4)} dx}{e} - \frac{a + \operatorname{arctanh}(cx^2)}{e(d+ex)}$$

↓ 7276

$$\frac{2bc \int \left(-\frac{de^3}{(e^4-c^2d^4)(d+ex)} + \frac{e-cdx}{2(cd^2-e^2)(cx^2-1)} + \frac{e+cdx}{2(cd^2+e^2)(cx^2+1)} \right) dx}{e} - \frac{a + \operatorname{arctanh}(cx^2)}{e(d+ex)}$$

↓ 2009

$$\frac{2bc \left(\frac{e \arctan(\sqrt{cx})}{2\sqrt{c}(cd^2+e^2)} - \frac{e \operatorname{arctanh}(\sqrt{cx})}{2\sqrt{c}(cd^2-e^2)} + \frac{de^2 \log(d+ex)}{c^2d^4-e^4} - \frac{d \log(1-cx^2)}{4(cd^2-e^2)} + \frac{d \log(cx^2+1)}{4(cd^2+e^2)} \right)}{e} - \frac{a + \operatorname{arctanh}(cx^2)}{e(d+ex)}$$

input `Int[(a + b*ArcTanh[c*x^2])/(d + e*x)^2,x]`

output `-((a + b*ArcTanh[c*x^2])/(e*(d + e*x))) + (2*b*c*((e*ArcTan[Sqrt[c]*x])/(2*sqrt[c]*(c*d^2 + e^2)) - (e*ArcTanh[Sqrt[c]*x])/(2*sqrt[c]*(c*d^2 - e^2))) + (d*e^2*Log[d + e*x])/(c^2*d^4 - e^4) - (d*Log[1 - c*x^2])/(4*(c*d^2 - e^2)) + (d*Log[1 + c*x^2])/(4*(c*d^2 + e^2)))/e`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6486 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.)*((d_.) + (e_.)*(x_)^(m_.)), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(e*(m + 1))), x] - Simp[b*c*(n/(e*(m + 1))) Int[x^(n - 1)*((d + e*x)^(m + 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.93

method	result
default	$-\frac{a}{(ex+d)e} + b \left(-\frac{\operatorname{arctanh}(cx^2)}{(ex+d)e} + \frac{2c \left(\frac{-\frac{d \ln(cx^2-1)}{2} - \frac{e \operatorname{arctanh}(\sqrt{c}x)}{\sqrt{c}}}{2cd^2-2e^2} + \frac{\frac{d \ln(cx^2+1)}{2} + \frac{e \operatorname{arctan}(\sqrt{c}x)}{\sqrt{c}}}{2cd^2+2e^2} + \frac{de^2 \ln(ex+d)}{(cd^2-e^2)(cd^2+e^2)} \right)}{e} \right)$
parts	$-\frac{a}{(ex+d)e} + b \left(-\frac{\operatorname{arctanh}(cx^2)}{(ex+d)e} + \frac{2c \left(\frac{-\frac{d \ln(cx^2-1)}{2} - \frac{e \operatorname{arctanh}(\sqrt{c}x)}{\sqrt{c}}}{2cd^2-2e^2} + \frac{\frac{d \ln(cx^2+1)}{2} + \frac{e \operatorname{arctan}(\sqrt{c}x)}{\sqrt{c}}}{2cd^2+2e^2} + \frac{de^2 \ln(ex+d)}{(cd^2-e^2)(cd^2+e^2)} \right)}{e} \right)$

input `int((a+b*arctanh(c*x^2))/(e*x+d)^2,x,method=_RETURNVERBOSE)`

output `-a/(e*x+d)/e+b*(-1/(e*x+d)/e*arctanh(c*x^2)+2/e*c*(1/(2*c*d^2-2*e^2)*(-1/2*d*ln(c*x^2-1)-e/c^(1/2)*arctanh(c^(1/2)*x))+1/(2*c*d^2+2*e^2)*(1/2*d*ln(c*x^2+1)+e/c^(1/2)*arctan(c^(1/2)*x))+d*e^2/(c*d^2-e^2)/(c*d^2+e^2)*ln(e*x+d))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 321 vs. 2(154) = 308.

Time = 2.55 (sec) , antiderivative size = 634, normalized size of antiderivative = 3.82

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^2} dx$$

$$= \left[\frac{2ac^2d^4 - 2ae^4 - 2(bcd^3e - bde^3 + (bcd^2e^2 - be^4)x)\sqrt{c} \operatorname{arctan}(\sqrt{c}x) + (bcd^3e + bde^3 + (bcd^2e^2 + be^4)x)\sqrt{-c} \operatorname{arctan}(\sqrt{-c}x) - (bcd^3e - bde^3 + (bcd^2e^2 - be^4)x)\sqrt{-c} \operatorname{arctan}(\sqrt{-c}x)}{\dots} \right]$$

input `integrate((a+b*arctanh(c*x^2))/(e*x+d)^2,x, algorithm="fricas")`

output `[-1/2*(2*a*c^2*d^4 - 2*a*e^4 - 2*(b*c*d^3*e - b*d*e^3 + (b*c*d^2*e^2 - b*e^4)*x)*sqrt(c)*arctan(sqrt(c)*x) + (b*c*d^3*e + b*d*e^3 + (b*c*d^2*e^2 + b*e^4)*x)*sqrt(c)*log((c*x^2 + 2*sqrt(c)*x + 1)/(c*x^2 - 1)) - (b*c^2*d^4 - b*c*d^2*e^2 + (b*c^2*d^3*e - b*c*d*e^3)*x)*log(c*x^2 + 1) + (b*c^2*d^4 + b*c*d^2*e^2 + (b*c^2*d^3*e + b*c*d*e^3)*x)*log(c*x^2 - 1) - 4*(b*c*d*e^3*x + b*c*d^2*e^2)*log(e*x + d) + (b*c^2*d^4 - b*e^4)*log(-(c*x^2 + 1)/(c*x^2 - 1)))/(c^2*d^5*e - d*e^5 + (c^2*d^4*e^2 - e^6)*x), -1/2*(2*a*c^2*d^4 - 2*a*e^4 - 2*(b*c*d^3*e + b*d*e^3 + (b*c*d^2*e^2 + b*e^4)*x)*sqrt(-c)*arctan(sqrt(-c)*x) - (b*c*d^3*e - b*d*e^3 + (b*c*d^2*e^2 - b*e^4)*x)*sqrt(-c)*log((c*x^2 + 2*sqrt(-c)*x - 1)/(c*x^2 + 1)) - (b*c^2*d^4 - b*c*d^2*e^2 + (b*c^2*d^3*e - b*c*d*e^3)*x)*log(c*x^2 + 1) + (b*c^2*d^4 + b*c*d^2*e^2 + (b*c^2*d^3*e + b*c*d*e^3)*x)*log(c*x^2 - 1) - 4*(b*c*d*e^3*x + b*c*d^2*e^2)*log(e*x + d) + (b*c^2*d^4 - b*e^4)*log(-(c*x^2 + 1)/(c*x^2 - 1)))/(c^2*d^5*e - d*e^5 + (c^2*d^4*e^2 - e^6)*x)]`

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^2} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**2))/(e*x+d)**2,x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.04

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^2} dx$$

$$= \frac{1}{2} \left(\left(\frac{4de \log(ex + d)}{c^2 d^4 - e^4} + \frac{d \log(cx^2 + 1)}{cd^2 e + e^3} - \frac{d \log(cx^2 - 1)}{cd^2 e - e^3} + \frac{2 \arctan(\sqrt{c}x)}{(cd^2 + e^2)\sqrt{c}} + \frac{\log\left(\frac{cx - \sqrt{c}}{cx + \sqrt{c}}\right)}{(cd^2 - e^2)\sqrt{c}} \right) c - \frac{2a}{e} \right) - \frac{a}{e^2 x + de}$$

input `integrate((a+b*arctanh(c*x^2))/(e*x+d)^2,x, algorithm="maxima")`

output `1/2*((4*d*e*log(e*x + d)/(c^2*d^4 - e^4) + d*log(c*x^2 + 1)/(c*d^2*e + e^3) - d*log(c*x^2 - 1)/(c*d^2*e - e^3) + 2*arctan(sqrt(c)*x)/((c*d^2 + e^2)*sqrt(c)) + log((c*x - sqrt(c))/(c*x + sqrt(c)))/((c*d^2 - e^2)*sqrt(c)))*c - 2*arctanh(c*x^2)/(e^2*x + d*e))*b - a/(e^2*x + d*e)`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.62

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^2} dx$$

$$= \frac{1}{2} \left(\frac{cd \log\left(c - \frac{2cd}{ex+d} + \frac{cd^2}{(ex+d)^2} + \frac{e^2}{(ex+d)^2}\right)}{cd^2 e + e^3} - \frac{cd \log\left(c - \frac{2cd}{ex+d} + \frac{cd^2}{(ex+d)^2} - \frac{e^2}{(ex+d)^2}\right)}{cd^2 e - e^3} + \frac{2c \arctan\left(\frac{cd - \frac{cd^2}{ex+d} + \frac{e^2}{(ex+d)^2}}{\sqrt{-c}}\right)}{(cd^2 - e^2)\sqrt{-c}} \right) - \frac{a}{(ex + d)e}$$

input `integrate((a+b*arctanh(c*x^2))/(e*x+d)^2,x, algorithm="giac")`

output

$$\begin{aligned} & \frac{1}{2} * (c * d * \log(c - 2 * c * d / (e * x + d) + c * d^2 / (e * x + d)^2 + e^2 / (e * x + d)^2) / (c \\ & * d^2 * e + e^3) - c * d * \log(c - 2 * c * d / (e * x + d) + c * d^2 / (e * x + d)^2 - e^2 / (e * x \\ & + d)^2) / (c * d^2 * e - e^3) + 2 * c * \arctan((c * d - c * d^2 / (e * x + d) + e^2 / (e * x + \\ & d)) / (\sqrt{-c} * e)) / ((c * d^2 - e^2) * \sqrt{-c}) + 2 * \sqrt{c} * \arctan((c * d - c * d^2 \\ & / (e * x + d) - e^2 / (e * x + d)) / (\sqrt{c} * e)) / (c * d^2 + e^2) - \log(-(c * x^2 + 1) / \\ & (c * x^2 - 1)) / ((e * x + d) * e) * b - a / ((e * x + d) * e) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 5.03 (sec) , antiderivative size = 727, normalized size of antiderivative = 4.38

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^2} dx$$

$$\ln \left(\frac{16b^4 c^{10} x}{e} - \frac{(b\sqrt{-ce+bcd}) \left(\frac{8x(4bc^{11}d^5e^2+52bc^9de^6)}{e^2} - \frac{(b\sqrt{-ce+bcd}) \left(\frac{384c^8de^6 + \frac{8x(4bc^{11}d^5e^2+52bc^9de^6)}{e^2}}{2(cd^2e+e^3)} \right)}{2(cd^2e+e^3)} \right)}{8b^3c^9e} - \frac{(b\sqrt{-ce+bcd}) \left(\frac{384c^8de^6 + \frac{8x(4bc^{11}d^5e^2+52bc^9de^6)}{e^2}}{2(cd^2e+e^3)} \right)}{2(cd^2e+e^3)} \right)}{2(cd^2e+e^3)}$$

$$= \frac{a}{x^2 + de} - \ln \left(\frac{(b\sqrt{-ce-bcd}) \left(\frac{8x(4bc^{11}d^5e^2+52bc^9de^6)}{e^2} + \frac{(b\sqrt{-ce-bcd}) \left(\frac{384c^8de^6 + \frac{8x(24c^{10}d^4e^5 + \frac{8x(4bc^{11}d^5e^2+52bc^9de^6)}{e^2}}{2(cd^2e+e^3)} \right)}{2(cd^2e+e^3)} \right)}{8b^3c^9e} + \frac{(b\sqrt{-ce-bcd}) \left(\frac{384c^8de^6 + \frac{8x(24c^{10}d^4e^5 + \frac{8x(4bc^{11}d^5e^2+52bc^9de^6)}{e^2}}{2(cd^2e+e^3)} \right)}{2(cd^2e+e^3)} \right)}{2(cd^2e+e^3)} \right)}{2(cd^2e+e^3)}$$

input `int((a + b*atanh(c*x^2))/(d + e*x)^2,x)`

output

$$\begin{aligned} & (\log((16*b^4*c^{10}*x)/e - ((b*(-c)^{(1/2)}*e + b*c*d)*(8*b^3*c^9*e - ((b*(-c)^{(1/2)}*e + b*c*d)*((b*(-c)^{(1/2)}*e + b*c*d)*((8*x*(4*b*c^{11}*d^5*e^2 + 52*b*c^9*d*e^6))/e^2 - ((b*(-c)^{(1/2)}*e + b*c*d)*(384*c^8*d*e^6 + (8*x*(40*c^8*e^9 + 24*c^{10}*d^4*e^5))/e^2 + 128*c^{10}*d^5*e^2))/(2*(e^3 + c*d^2*e)) + 320*b*c^9*d^2*e^3))/(2*(e^3 + c*d^2*e)) - 64*b^2*c^{10}*d^2*e*x))/(2*(e^3 + c*d^2*e)) + (8*b^3*c^{11}*d^3*x)/e^2))/(2*(e^3 + c*d^2*e)))*(b*(-c)^{(1/2)}*e + b*c*d))/(2*(e^3 + c*d^2*e)) - a/(d*e + e^2*x) - (\log(((b*(-c)^{(1/2)}*e - b*c*d)*(8*b^3*c^9*e - ((b*(-c)^{(1/2)}*e - b*c*d)*((b*(-c)^{(1/2)}*e - b*c*d)*((8*x*(4*b*c^{11}*d^5*e^2 + 52*b*c^9*d*e^6))/e^2 + ((b*(-c)^{(1/2)}*e - b*c*d)*(384*c^8*d*e^6 + (8*x*(40*c^8*e^9 + 24*c^{10}*d^4*e^5))/e^2 + 128*c^{10}*d^5*e^2))/(2*(e^3 + c*d^2*e)) + 320*b*c^9*d^2*e^3))/(2*(e^3 + c*d^2*e)) + 64*b^2*c^{10}*d^2*e*x))/(2*(e^3 + c*d^2*e)) + (8*b^3*c^{11}*d^3*x)/e^2))/(2*(e^3 + c*d^2*e)) + (16*b^4*c^{10}*x)/e*(b*(-c)^{(1/2)}*e - b*c*d))/(2*(e^3 + c*d^2*e)) - (b*log(c*x^2 + 1))/(2*e*(d + e*x)) - (b*c^{(1/2)}*log(c^{(1/2)}*x - 1))/(2*(e^2 + c^{(1/2)}*d*e)) + (b*c^{(1/2)}*log(c^{(1/2)}*x + 1))/(2*(e^2 - c^{(1/2)}*d*e)) + (b*log(1 - c*x^2))/(e*(2*d + 2*e*x)) - (2*b*c*d*e*log(d + e*x))/(e^4 - c^2*d^4) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 506, normalized size of antiderivative = 3.05

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^2} dx$$

$$= \frac{-\log(\sqrt{c}x - 1) b e^4 x - \log(\sqrt{c}x + 1) b d e^3 - \log(\sqrt{c}x + 1) b e^4 x + \log(c x^2 + 1) b d e^3 + \log(c x^2 + 1) b}{e^4 - c^2 d^4}$$

input `int((a+b*atanh(c*x^2))/(e*x+d)^2,x)`

output

```
(2*sqrt(c)*atan((c*x)/sqrt(c))*b*c*d**4 + 2*sqrt(c)*atan((c*x)/sqrt(c))*b*
c*d**3*e*x - 2*sqrt(c)*atan((c*x)/sqrt(c))*b*d**2*e**2 - 2*sqrt(c)*atan((c
*x)/sqrt(c))*b*d*e**3*x + 2*atanh(c*x**2)*b*c**2*d**4*x - 2*atanh(c*x**2)*
b*e**4*x + sqrt(c)*log(sqrt(c)*x - 1)*b*c*d**4 + sqrt(c)*log(sqrt(c)*x - 1
)*b*c*d**3*e*x + sqrt(c)*log(sqrt(c)*x - 1)*b*d**2*e**2 + sqrt(c)*log(sqrt
(c)*x - 1)*b*d*e**3*x - sqrt(c)*log(sqrt(c)*x + 1)*b*c*d**4 - sqrt(c)*log(
sqrt(c)*x + 1)*b*c*d**3*e*x - sqrt(c)*log(sqrt(c)*x + 1)*b*d**2*e**2 - sqr
t(c)*log(sqrt(c)*x + 1)*b*d*e**3*x - log(sqrt(c)*x - 1)*b*c*d**3*e - log(s
qrt(c)*x - 1)*b*c*d**2*e**2*x - log(sqrt(c)*x - 1)*b*d*e**3 - log(sqrt(c)*
x - 1)*b*e**4*x - log(sqrt(c)*x + 1)*b*c*d**3*e - log(sqrt(c)*x + 1)*b*c*d
**2*e**2*x - log(sqrt(c)*x + 1)*b*d*e**3 - log(sqrt(c)*x + 1)*b*e**4*x - l
og(c*x**2 + 1)*b*c*d**3*e - log(c*x**2 + 1)*b*c*d**2*e**2*x + log(c*x**2 +
1)*b*d*e**3 + log(c*x**2 + 1)*b*e**4*x + 4*log(d + e*x)*b*c*d**3*e + 4*lo
g(d + e*x)*b*c*d**2*e**2*x + 2*a*c**2*d**4*x - 2*a*e**4*x)/(2*d*(c**2*d**5
+ c**2*d**4*e*x - d*e**4 - e**5*x))
```

$$3.29 \quad \int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^3} dx$$

Optimal result	265
Mathematica [A] (verified)	266
Rubi [A] (verified)	267
Maple [A] (verified)	268
Fricas [B] (verification not implemented)	269
Sympy [F(-1)]	270
Maxima [A] (verification not implemented)	270
Giac [A] (verification not implemented)	271
Mupad [B] (verification not implemented)	271
Reduce [B] (verification not implemented)	272

Optimal result

Integrand size = 18, antiderivative size = 226

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^3} dx = -\frac{bcde}{(c^2d^4 - e^4)(d + ex)} + \frac{bc^{3/2}d \arctan(\sqrt{cx})}{(cd^2 + e^2)^2}$$

$$- \frac{bc^{3/2}d \operatorname{arctanh}(\sqrt{cx})}{(cd^2 - e^2)^2} - \frac{a + b \operatorname{arctanh}(cx^2)}{2e(d + ex)^2}$$

$$+ \frac{bce(3c^2d^4 + e^4) \log(d + ex)}{(c^2d^4 - e^4)^2}$$

$$- \frac{bc(cd^2 + e^2) \log(1 - cx^2)}{4e(cd^2 - e^2)^2} + \frac{bc(cd^2 - e^2) \log(1 + cx^2)}{4e(cd^2 + e^2)^2}$$

output

```
-b*c*d*e/(c^2*d^4-e^4)/(e*x+d)+b*c^(3/2)*d*arctan(c^(1/2)*x)/(c*d^2+e^2)^2
-b*c^(3/2)*d*arctanh(c^(1/2)*x)/(c*d^2-e^2)^2-1/2*(a+b*arctanh(c*x^2))/e/(
e*x+d)^2+b*c*e*(3*c^2*d^4+e^4)*ln(e*x+d)/(c^2*d^4-e^4)^2-1/4*b*c*(c*d^2+e^
2)*ln(-c*x^2+1)/e/(c*d^2-e^2)^2+1/4*b*c*(c*d^2-e^2)*ln(c*x^2+1)/e/(c*d^2+e
^2)^2
```

Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 379, normalized size of antiderivative = 1.68

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^3} dx$$

$$= \frac{1}{4} \left(-\frac{2a}{e(d + ex)^2} - \frac{4bcde}{(c^2d^4 - e^4)(d + ex)} + \frac{4bc^{3/2}d \arctan(\sqrt{cx})}{(cd^2 + e^2)^2} - \frac{2b \operatorname{arctanh}(cx^2)}{e(d + ex)^2} \right.$$

$$- \frac{bc^{3/2}d(c^{5/2}d^5 - 2c^2d^4e - 4cd^2e^3 + 3\sqrt{c}de^4 - 2e^5) \log(1 - \sqrt{cx})}{e(-c^2d^4 + e^4)^2}$$

$$- \frac{bc^{3/2}d(c^{5/2}d^5 + 2c^2d^4e + 4cd^2e^3 + 3\sqrt{c}de^4 + 2e^5) \log(1 + \sqrt{cx})}{e(-c^2d^4 + e^4)^2}$$

$$+ \frac{4bce(3c^2d^4 + e^4) \log(d + ex)}{(-c^2d^4 + e^4)^2} + \frac{bc^2(c^2d^6 + 3d^2e^4) \log(1 + cx^2)}{e(-c^2d^4 + e^4)^2}$$

$$\left. - \frac{bce(3c^2d^4 + e^4) \log(1 - c^2x^4)}{(-c^2d^4 + e^4)^2} \right)$$

input `Integrate[(a + b*ArcTanh[c*x^2])/(d + e*x)^3,x]`output `((-2*a)/(e*(d + e*x)^2) - (4*b*c*d*e)/((c^2*d^4 - e^4)*(d + e*x)) + (4*b*c^(3/2)*d*ArcTan[Sqrt[c]*x])/(c*d^2 + e^2)^2 - (2*b*ArcTanh[c*x^2])/(e*(d + e*x)^2) - (b*c^(3/2)*d*(c^(5/2)*d^5 - 2*c^2*d^4*e - 4*c*d^2*e^3 + 3*Sqrt[c]*d*e^4 - 2*e^5)*Log[1 - Sqrt[c]*x])/(e*(-(c^2*d^4) + e^4)^2) - (b*c^(3/2)*d*(c^(5/2)*d^5 + 2*c^2*d^4*e + 4*c*d^2*e^3 + 3*Sqrt[c]*d*e^4 + 2*e^5)*Log[1 + Sqrt[c]*x])/(e*(-(c^2*d^4) + e^4)^2) + (4*b*c*e*(3*c^2*d^4 + e^4)*Log[d + e*x])/(-(c^2*d^4) + e^4)^2 + (b*c^2*(c^2*d^6 + 3*d^2*e^4)*Log[1 + c*x^2])/(e*(-(c^2*d^4) + e^4)^2) - (b*c*e*(3*c^2*d^4 + e^4)*Log[1 - c^2*x^4])/(e*(-(c^2*d^4) + e^4)^2)/4`

Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 223, normalized size of antiderivative = 0.99, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6486, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{arctanh}(cx^2)}{(d + ex)^3} dx$$

↓ 6486

$$\frac{bc \int \frac{x}{(d+ex)^2(1-c^2x^4)} dx}{e} - \frac{a + \operatorname{arctanh}(cx^2)}{2e(d + ex)^2}$$

↓ 7276

$$\frac{bc \int \left(-\frac{de^3}{(e^4 - c^2d^4)(d+ex)^2} + \frac{e^7 + 3c^2d^4e^3}{(c^2d^4 - e^4)^2(d+ex)} + \frac{c((cd^2 + e^2)x - 2de)}{2(cd^2 - e^2)^2(1 - cx^2)} + \frac{c(2de + (cd^2 - e^2)x)}{2(cd^2 + e^2)^2(cx^2 + 1)} \right) dx}{e} - \frac{a + \operatorname{arctanh}(cx^2)}{2e(d + ex)^2}$$

↓ 2009

$$\frac{bc \left(\frac{\sqrt{cde} \arctan(\sqrt{cx})}{(cd^2 + e^2)^2} - \frac{\sqrt{cde} \operatorname{arctanh}(\sqrt{cx})}{(cd^2 - e^2)^2} - \frac{de^2}{(c^2d^4 - e^4)(d+ex)} + \frac{e^2(3c^2d^4 + e^4) \log(d+ex)}{(c^2d^4 - e^4)^2} - \frac{(cd^2 + e^2) \log(1 - cx^2)}{4(cd^2 - e^2)^2} + \frac{(cd^2 - e^2) \log(1 + cx^2)}{4(cd^2 + e^2)^2} \right)}{e} - \frac{a + \operatorname{arctanh}(cx^2)}{2e(d + ex)^2}$$

input

```
Int[(a + b*ArcTanh[c*x^2])/(d + e*x)^3,x]
```

output

```
-1/2*(a + b*ArcTanh[c*x^2])/(e*(d + e*x)^2) + (b*c*(-((d*e^2)/((c^2*d^4 - e^4)*(d + e*x))) + (Sqrt[c]*d*e*ArcTan[Sqrt[c]*x])/(c*d^2 + e^2)^2 - (Sqrt[c]*d*e*ArcTanh[Sqrt[c]*x])/(c*d^2 - e^2)^2 + (e^2*(3*c^2*d^4 + e^4)*Log[d + e*x])/(c^2*d^4 - e^4)^2 - ((c*d^2 + e^2)*Log[1 - c*x^2])/(4*(c*d^2 - e^2)^2) + ((c*d^2 - e^2)*Log[1 + c*x^2])/(4*(c*d^2 + e^2)^2))/e
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6486 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.)*((d_.) + (e_.)*(x_)^(m_.)), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(e*(m + 1))), x] - Simp[b*c*(n/(e*(m + 1))) Int[x^(n - 1)*((d + e*x)^(m + 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 0.74 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.01

method	result
default	$-\frac{a}{2(ex+d)^2e} + b \left(-\frac{\operatorname{arctanh}(cx^2)}{2(ex+d)^2e} + \frac{c \left(-\frac{c \left(\frac{(cd^2+e^2)\ln(cx^2-1)}{2c} + \frac{2de \operatorname{arctanh}(\sqrt{c}x)}{\sqrt{c}} \right)}{2(cd^2-e^2)^2} + \frac{c \left(\frac{(cd^2-e^2)\ln(cx^2+1)}{2c} + \frac{2de \operatorname{arctanh}(\sqrt{c}x)}{\sqrt{c}} \right)}{2(cd^2+e^2)^2} \right)}{e}$
parts	$-\frac{a}{2(ex+d)^2e} + b \left(-\frac{\operatorname{arctanh}(cx^2)}{2(ex+d)^2e} + \frac{c \left(-\frac{c \left(\frac{(cd^2+e^2)\ln(cx^2-1)}{2c} + \frac{2de \operatorname{arctanh}(\sqrt{c}x)}{\sqrt{c}} \right)}{2(cd^2-e^2)^2} + \frac{c \left(\frac{(cd^2-e^2)\ln(cx^2+1)}{2c} + \frac{2de \operatorname{arctanh}(\sqrt{c}x)}{\sqrt{c}} \right)}{2(cd^2+e^2)^2} \right)}{e}$

input `int((a+b*arctanh(c*x^2))/(e*x+d)^3,x,method=_RETURNVERBOSE)`

output

```
-1/2*a/(e*x+d)^2/e+b*(-1/2/(e*x+d)^2/e*arctanh(c*x^2)+1/e*c*(-1/2*c/(c*d^2
-e^2)^2*(1/2*(c*d^2+e^2)/c*ln(c*x^2-1)+2*d*e/c^(1/2)*arctanh(c^(1/2)*x))+1
/2*c/(c*d^2+e^2)^2*(1/2*(c*d^2-e^2)/c*ln(c*x^2+1)+2*d*e/c^(1/2)*arctan(c^(
1/2)*x))-d*e^2/(c*d^2-e^2)/(c*d^2+e^2)/(e*x+d)+e^2*(3*c^2*d^4+e^4)/(c*d^2-
e^2)^2/(c*d^2+e^2)^2*ln(e*x+d))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 815 vs. $2(212) = 424$.

Time = 22.24 (sec) , antiderivative size = 1639, normalized size of antiderivative = 7.25

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^3} dx = \text{Too large to display}$$

input

```
integrate((a+b*arctanh(c*x^2))/(e*x+d)^3,x, algorithm="fricas")
```

output

```
[-1/4*(2*a*c^4*d^8 + 4*b*c^3*d^6*e^2 - 4*a*c^2*d^4*e^4 - 4*b*c*d^2*e^6 + 2
*a*e^8 - 4*(b*c^3*d^7*e + 2*b*c^2*d^5*e^3 + b*c*d^3*e^5 + (b*c^3*d^5*e^3 +
2*b*c^2*d^3*e^5 + b*c*d*e^7)*x^2 + 2*(b*c^3*d^6*e^2 + 2*b*c^2*d^4*e^4 + b
*c*d^2*e^6)*x)*sqrt(-c)*arctan(sqrt(-c)*x) - 2*(b*c^3*d^7*e - 2*b*c^2*d^5*
e^3 + b*c*d^3*e^5 + (b*c^3*d^5*e^3 - 2*b*c^2*d^3*e^5 + b*c*d*e^7)*x^2 + 2*
(b*c^3*d^6*e^2 - 2*b*c^2*d^4*e^4 + b*c*d^2*e^6)*x)*sqrt(-c)*log((c*x^2 + 2
*sqrt(-c)*x - 1)/(c*x^2 + 1)) + 4*(b*c^3*d^5*e^3 - b*c*d*e^7)*x - (b*c^4*d
^8 - 3*b*c^3*d^6*e^2 + 3*b*c^2*d^4*e^4 - b*c*d^2*e^6 + (b*c^4*d^6*e^2 - 3*
b*c^3*d^4*e^4 + 3*b*c^2*d^2*e^6 - b*c*e^8)*x^2 + 2*(b*c^4*d^7*e - 3*b*c^3*
d^5*e^3 + 3*b*c^2*d^3*e^5 - b*c*d*e^7)*x)*log(c*x^2 + 1) + (b*c^4*d^8 + 3*
b*c^3*d^6*e^2 + 3*b*c^2*d^4*e^4 + b*c*d^2*e^6 + (b*c^4*d^6*e^2 + 3*b*c^3*d
^4*e^4 + 3*b*c^2*d^2*e^6 + b*c*e^8)*x^2 + 2*(b*c^4*d^7*e + 3*b*c^3*d^5*e^3
+ 3*b*c^2*d^3*e^5 + b*c*d*e^7)*x)*log(c*x^2 - 1) - 4*(3*b*c^3*d^6*e^2 + b
*c*d^2*e^6 + (3*b*c^3*d^4*e^4 + b*c*e^8)*x^2 + 2*(3*b*c^3*d^5*e^3 + b*c*d*
e^7)*x)*log(e*x + d) + (b*c^4*d^8 - 2*b*c^2*d^4*e^4 + b*e^8)*log(-(c*x^2 +
1)/(c*x^2 - 1)))/(c^4*d^10*e - 2*c^2*d^6*e^5 + d^2*e^9 + (c^4*d^8*e^3 - 2
*c^2*d^4*e^7 + e^11)*x^2 + 2*(c^4*d^9*e^2 - 2*c^2*d^5*e^6 + d*e^10)*x), -1
/4*(2*a*c^4*d^8 + 4*b*c^3*d^6*e^2 - 4*a*c^2*d^4*e^4 - 4*b*c*d^2*e^6 + 2*a*
e^8 - 4*(b*c^3*d^7*e - 2*b*c^2*d^5*e^3 + b*c*d^3*e^5 + (b*c^3*d^5*e^3 - 2*
b*c^2*d^3*e^5 + b*c*d*e^7)*x^2 + 2*(b*c^3*d^6*e^2 - 2*b*c^2*d^4*e^4 + b...
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^3} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**2))/(e*x+d)**3,x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 311, normalized size of antiderivative = 1.38

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^3} dx$$

$$= \frac{1}{4} \left(\left(\frac{4\sqrt{cd} \arctan(\sqrt{cx})}{c^2d^4 + 2cd^2e^2 + e^4} + \frac{2\sqrt{cd} \log\left(\frac{cx-\sqrt{c}}{cx+\sqrt{c}}\right)}{c^2d^4 - 2cd^2e^2 + e^4} - \frac{4de}{c^2d^5 - de^4 + (c^2d^4e - e^5)x} + \frac{(cd^2 - e^2) \log(cx^2 + 1)}{c^2d^4e + 2cd^2e^3 + e^5} \right) - \frac{a}{2(e^3x^2 + 2de^2x + d^2e)} \right)$$

input `integrate((a+b*arctanh(c*x^2))/(e*x+d)^3,x, algorithm="maxima")`

output `1/4*((4*sqrt(c)*d*arctan(sqrt(c)*x)/(c^2*d^4 + 2*c*d^2*e^2 + e^4) + 2*sqrt(c)*d*log((c*x - sqrt(c))/(c*x + sqrt(c)))/(c^2*d^4 - 2*c*d^2*e^2 + e^4) - 4*d*e/(c^2*d^5 - d*e^4 + (c^2*d^4*e - e^5)*x) + (c*d^2 - e^2)*log(c*x^2 + 1)/(c^2*d^4*e + 2*c*d^2*e^3 + e^5) - (c*d^2 + e^2)*log(c*x^2 - 1)/(c^2*d^4*e - 2*c*d^2*e^3 + e^5) + 4*(3*c^2*d^4*e + e^5)*log(e*x + d)/(c^4*d^8 - 2*c^2*d^4*e^4 + e^8))*c - 2*arctanh(c*x^2)/(e^3*x^2 + 2*d*e^2*x + d^2*e) - 1/2*a/(e^3*x^2 + 2*d*e^2*x + d^2*e)`

Giac [A] (verification not implemented)

Time = 24.16 (sec) , antiderivative size = 362, normalized size of antiderivative = 1.60

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^3} dx = \frac{bc^{\frac{3}{2}}d \arctan(\sqrt{cx})}{c^2d^4 + 2cd^2e^2 + e^4} + \frac{bc^2d \arctan\left(\frac{cx}{\sqrt{-c}}\right)}{(c^2d^4 - 2cd^2e^2 + e^4)\sqrt{-c}}$$

$$+ \frac{(bc^2d^2 - bce^2) \log(cx^2 + 1)}{4(c^2d^4e + 2cd^2e^3 + e^5)}$$

$$- \frac{(bc^2d^2 + bce^2) \log(-cx^2 + 1)}{4(c^2d^4e - 2cd^2e^3 + e^5)}$$

$$+ \frac{(3bc^3d^4e + bce^5) \log(ex + d)}{c^4d^8 - 2c^2d^4e^4 + e^8} - \frac{b \log\left(-\frac{cx^2+1}{cx^2-1}\right)}{4(e^3x^2 + 2de^2x + d^2e)}$$

$$- \frac{ac^2d^4 + 2bcde^3x + 2bcd^2e^2 - ae^4}{2(c^2d^4e^3x^2 + 2c^2d^5e^2x + c^2d^6e - e^7x^2 - 2de^6x - d^2e^5)}$$

input `integrate((a+b*arctanh(c*x^2))/(e*x+d)^3,x, algorithm="giac")`

output `b*c^(3/2)*d*arctan(sqrt(c)*x)/(c^2*d^4 + 2*c*d^2*e^2 + e^4) + b*c^2*d*arctan(c*x/sqrt(-c))/((c^2*d^4 - 2*c*d^2*e^2 + e^4)*sqrt(-c)) + 1/4*(b*c^2*d^2 - b*c*e^2)*log(c*x^2 + 1)/(c^2*d^4*e + 2*c*d^2*e^3 + e^5) - 1/4*(b*c^2*d^2 + b*c*e^2)*log(-c*x^2 + 1)/(c^2*d^4*e - 2*c*d^2*e^3 + e^5) + (3*b*c^3*d^4*e + b*c*e^5)*log(e*x + d)/(c^4*d^8 - 2*c^2*d^4*e^4 + e^8) - 1/4*b*log(-(c*x^2 + 1)/(c*x^2 - 1))/(e^3*x^2 + 2*d*e^2*x + d^2*e) - 1/2*(a*c^2*d^4 + 2*b*c*d*e^3*x + 2*b*c*d^2*e^2 - a*e^4)/(c^2*d^4*e^3*x^2 + 2*c^2*d^5*e^2*x + c^2*d^6*e - e^7*x^2 - 2*d*e^6*x - d^2*e^5)`

Mupad [B] (verification not implemented)

Time = 7.64 (sec) , antiderivative size = 2016, normalized size of antiderivative = 8.92

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^3} dx = \text{Too large to display}$$

input `int((a + b*atanh(c*x^2))/(d + e*x)^3,x)`

output

```

((a*c^2*d^4 - a*e^4 + 2*b*c*d^2*e^2)/(2*(e^4 - c^2*d^4)) + (b*c*d*e^3*x)/(
e^4 - c^2*d^4))/(d^2*e + e^3*x^2 + 2*d*e^2*x) + (log(c*d^28*(-c^3)^(13/2)
+ c^21*d^28*x + 100*c^7*e^28*x - 100*c^2*e^28*(-c^3)^(3/2) + 496*d^2*e^26*
(-c^3)^(5/2) - 7398*d^8*e^20*(-c^3)^(7/2) + 17176*d^14*e^14*(-c^3)^(9/2) -
2703*d^20*e^8*(-c^3)^(11/2) - 20*d^26*e^2*(-c^3)^(13/2) + 496*c^8*d^2*e^2
6*x + 1473*c^9*d^4*e^24*x + 3692*c^10*d^6*e^22*x + 7398*c^11*d^8*e^20*x +
11868*c^12*d^10*e^18*x + 16015*c^13*d^12*e^16*x + 17176*c^14*d^14*e^14*x +
13192*c^15*d^16*e^12*x + 6984*c^16*d^18*e^10*x + 2703*c^17*d^20*e^8*x + 7
64*c^18*d^22*e^6*x + 78*c^19*d^24*e^4*x - 20*c^20*d^26*e^2*x + 3692*c^2*d^
6*e^22*(-c^3)^(5/2) - 16015*c^2*d^12*e^16*(-c^3)^(7/2) + 6984*c^2*d^18*e^1
0*(-c^3)^(9/2) - 78*c^2*d^24*e^4*(-c^3)^(11/2) + 1473*c*d^4*e^24*(-c^3)^(5
/2) - 11868*c*d^10*e^18*(-c^3)^(7/2) + 13192*c*d^16*e^12*(-c^3)^(9/2) - 76
4*c*d^22*e^6*(-c^3)^(11/2))*(b*c^2*d^2 - b*c*e^2 + 2*b*d*e*(-c^3)^(1/2)))/
(4*(e^5 + 2*c*d^2*e^3 + c^2*d^4*e)) - (log(c^21*d^28*x - c*d^28*(-c^3)^(13
/2) + 100*c^7*e^28*x + 100*c^2*e^28*(-c^3)^(3/2) - 496*d^2*e^26*(-c^3)^(5/
2) + 7398*d^8*e^20*(-c^3)^(7/2) - 17176*d^14*e^14*(-c^3)^(9/2) + 2703*d^20
*e^8*(-c^3)^(11/2) + 20*d^26*e^2*(-c^3)^(13/2) + 496*c^8*d^2*e^26*x + 1473
*c^9*d^4*e^24*x + 3692*c^10*d^6*e^22*x + 7398*c^11*d^8*e^20*x + 11868*c^12
*d^10*e^18*x + 16015*c^13*d^12*e^16*x + 17176*c^14*d^14*e^14*x + 13192*c^1
5*d^16*e^12*x + 6984*c^16*d^18*e^10*x + 2703*c^17*d^20*e^8*x + 764*c^18...

```

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 1597, normalized size of antiderivative = 7.07

$$\int \frac{a + b \operatorname{arctanh}(cx^2)}{(d + ex)^3} dx = \text{Too large to display}$$

input

```
int((a+b*atanh(c*x^2))/(e*x+d)^3,x)
```

output

```
(4*sqrt(c)*atan((c*x)/sqrt(c))*b*c**3*d**7*e + 8*sqrt(c)*atan((c*x)/sqrt(c))
)*b*c**3*d**6*e**2*x + 4*sqrt(c)*atan((c*x)/sqrt(c))*b*c**3*d**5*e**3*x**
2 - 8*sqrt(c)*atan((c*x)/sqrt(c))*b*c**2*d**5*e**3 - 16*sqrt(c)*atan((c*x)
/sqrt(c))*b*c**2*d**4*e**4*x - 8*sqrt(c)*atan((c*x)/sqrt(c))*b*c**2*d**3*e
**5*x**2 + 4*sqrt(c)*atan((c*x)/sqrt(c))*b*c*d**3*e**5 + 8*sqrt(c)*atan((c
*x)/sqrt(c))*b*c*d**2*e**6*x + 4*sqrt(c)*atan((c*x)/sqrt(c))*b*c*d*e**7*x*
*2 - 2*atanh(c*x**2)*b*c**4*d**8 + 4*atanh(c*x**2)*b*c**2*d**4*e**4 - 2*at
anh(c*x**2)*b*e**8 + 2*sqrt(c)*log(sqrt(c)*x - 1)*b*c**3*d**7*e + 4*sqrt(c)
)*log(sqrt(c)*x - 1)*b*c**3*d**6*e**2*x + 2*sqrt(c)*log(sqrt(c)*x - 1)*b*c
**3*d**5*e**3*x**2 + 4*sqrt(c)*log(sqrt(c)*x - 1)*b*c**2*d**5*e**3 + 8*sqr
t(c)*log(sqrt(c)*x - 1)*b*c**2*d**4*e**4*x + 4*sqrt(c)*log(sqrt(c)*x - 1)*
b*c**2*d**3*e**5*x**2 + 2*sqrt(c)*log(sqrt(c)*x - 1)*b*c*d**3*e**5 + 4*sqr
t(c)*log(sqrt(c)*x - 1)*b*c*d**2*e**6*x + 2*sqrt(c)*log(sqrt(c)*x - 1)*b*c
*d*e**7*x**2 - 2*sqrt(c)*log(sqrt(c)*x + 1)*b*c**3*d**7*e - 4*sqrt(c)*log(
sqrt(c)*x + 1)*b*c**3*d**6*e**2*x - 2*sqrt(c)*log(sqrt(c)*x + 1)*b*c**3*d*
*5*e**3*x**2 - 4*sqrt(c)*log(sqrt(c)*x + 1)*b*c**2*d**5*e**3 - 8*sqrt(c)*l
og(sqrt(c)*x + 1)*b*c**2*d**4*e**4*x - 4*sqrt(c)*log(sqrt(c)*x + 1)*b*c**2
*d**3*e**5*x**2 - 2*sqrt(c)*log(sqrt(c)*x + 1)*b*c*d**3*e**5 - 4*sqrt(c)*l
og(sqrt(c)*x + 1)*b*c*d**2*e**6*x - 2*sqrt(c)*log(sqrt(c)*x + 1)*b*c*d*e**
7*x**2 - log(sqrt(c)*x - 1)*b*c**4*d**8 - 2*log(sqrt(c)*x - 1)*b*c**4*d...
```

3.30 $\int (d + ex) (a + \operatorname{barctanh}(cx^2))^2 dx$

Optimal result	274
Mathematica [A] (warning: unable to verify)	275
Rubi [A] (verified)	276
Maple [F]	278
Fricas [F]	279
Sympy [F]	279
Maxima [F]	279
Giac [F]	280
Mupad [F(-1)]	280
Reduce [F]	281

Optimal result

Integrand size = 18, antiderivative size = 1085

$$\int (d + ex) (a + \operatorname{barctanh}(cx^2))^2 dx = \text{Too large to display}$$

output

```

-1/2*b^2*d*polylog(2,1-2*c^(1/2)*(1+(-c)^(1/2)*x)/((-c)^(1/2)+c^(1/2))/(1+
c^(1/2)*x))/c^(1/2)-1/2*b^2*d*polylog(2,1+2*c^(1/2)*(1-(-c)^(1/2)*x)/((-c)
^(1/2)-c^(1/2))/(1+c^(1/2)*x))/c^(1/2)+b^2*d*polylog(2,1-2/(1+c^(1/2)*x))/
c^(1/2)+b^2*d*polylog(2,1-2/(1-c^(1/2)*x))/c^(1/2)-b^2*d*arctanh(c^(1/2)*x
)^2/c^(1/2)-1/2*b^2*d*e*polylog(2,1-2/(-c*x^2+1))/c+1/4*b^2*d*x*ln(c*x^2+1)^
2+1/4*b^2*d*x*ln(-c*x^2+1)^2-1/2*b^2*d*x*ln(-c*x^2+1)*ln(c*x^2+1)-2*b^2*d*
arctanh(c^(1/2)*x)*ln(2/(1+c^(1/2)*x))/c^(1/2)+2*b^2*d*arctanh(c^(1/2)*x)*
ln(2/(1-c^(1/2)*x))/c^(1/2)-2*a*b*d*arctanh(c^(1/2)*x)/c^(1/2)+a^2*d*x+2*b
^2*d*arctan(c^(1/2)*x)*ln(2/(1+I*c^(1/2)*x))/c^(1/2)-2*b^2*d*arctan(c^(1/2
)*x)*ln(2/(1-I*c^(1/2)*x))/c^(1/2)+2*a*b*d*arctan(c^(1/2)*x)/c^(1/2)-1/2*I
*b^2*d*polylog(2,1+(-1+I)*(1+c^(1/2)*x)/(1-I*c^(1/2)*x))/c^(1/2)-1/2*I*b^2
*d*polylog(2,1-(1+I)*(1-c^(1/2)*x)/(1-I*c^(1/2)*x))/c^(1/2)+b^2*d*arctan(c
^(1/2)*x)*ln(c*x^2+1)/c^(1/2)+b^2*d*arctanh(c^(1/2)*x)*ln(-c*x^2+1)/c^(1/2
)-b^2*d*arctan(c^(1/2)*x)*ln(-c*x^2+1)/c^(1/2)+I*b^2*d*polylog(2,1-2/(1+I*
c^(1/2)*x))/c^(1/2)+I*b^2*d*polylog(2,1-2/(1-I*c^(1/2)*x))/c^(1/2)+b^2*d*a
rctanh(c^(1/2)*x)*ln(2*c^(1/2)*(1+(-c)^(1/2)*x)/((-c)^(1/2)+c^(1/2))/(1+c^
(1/2)*x))/c^(1/2)+b^2*d*arctanh(c^(1/2)*x)*ln(-2*c^(1/2)*(1-(-c)^(1/2)*x)/
((-c)^(1/2)-c^(1/2))/(1+c^(1/2)*x))/c^(1/2)+b^2*d*arctan(c^(1/2)*x)*ln((1+
I)*(1-c^(1/2)*x)/(1-I*c^(1/2)*x))/c^(1/2)+b^2*d*arctan(c^(1/2)*x)*ln((1-I
)*(1+c^(1/2)*x)/(1-I*c^(1/2)*x))/c^(1/2)+I*b^2*d*arctan(c^(1/2)*x)^2/c^(...

```

Mathematica [A] (warning: unable to verify)

Time = 2.08 (sec) , antiderivative size = 684, normalized size of antiderivative = 0.63

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^2))^2 dx = \text{Too large to display}$$

input

```
Integrate[(d + e*x)*(a + b*ArcTanh[c*x^2])^2,x]
```

output

```
(2*a^2*c*d*x^2 + a^2*c*e*x^3 + 4*a*b*c*d*x^2*ArcTanh[c*x^2] + 4*a*b*d*Sqrt
[c*x^2]*(ArcTan[Sqrt[c*x^2]] - ArcTanh[Sqrt[c*x^2]]) + b^2*e*x*ArcTanh[c*x
^2]*((-1 + c*x^2)*ArcTanh[c*x^2] - 2*Log[1 + E^(-2*ArcTanh[c*x^2])]) + a*b
*e*x*(2*c*x^2*ArcTanh[c*x^2] + Log[1 - c^2*x^4]) + b^2*e*x*PolyLog[2, -E^(-
2*ArcTanh[c*x^2])] - b^2*d*Sqrt[c*x^2]*((2*I)*ArcTan[Sqrt[c*x^2]]^2 - 4*A
rcTan[Sqrt[c*x^2]]*ArcTanh[c*x^2] - 2*Sqrt[c*x^2]*ArcTanh[c*x^2]^2 - 2*Arc
Tan[Sqrt[c*x^2]]*Log[1 + E^((4*I)*ArcTan[Sqrt[c*x^2]])] - 2*ArcTanh[c*x^2]
*Log[1 - Sqrt[c*x^2]] + Log[2]*Log[1 - Sqrt[c*x^2]] - Log[1 - Sqrt[c*x^2]]
^2/2 + Log[1 - Sqrt[c*x^2]]*Log[(1/2 + I/2)*(-I + Sqrt[c*x^2])] + 2*ArcTan
h[c*x^2]*Log[1 + Sqrt[c*x^2]] - Log[2]*Log[1 + Sqrt[c*x^2]] - Log[((1 + I)
- (1 - I)*Sqrt[c*x^2])/2]*Log[1 + Sqrt[c*x^2]] - Log[(-1/2 - I/2)*(I + Sq
rt[c*x^2])]*Log[1 + Sqrt[c*x^2]] + Log[1 + Sqrt[c*x^2]]^2/2 + Log[1 - Sqrt
[c*x^2]]*Log[((1 + I) + (1 - I)*Sqrt[c*x^2])/2] + (I/2)*PolyLog[2, -E^((4*I
)*ArcTan[Sqrt[c*x^2]])] - PolyLog[2, (1 - Sqrt[c*x^2])/2] + PolyLog[2, (-
1/2 - I/2)*(-1 + Sqrt[c*x^2])] + PolyLog[2, (-1/2 + I/2)*(-1 + Sqrt[c*x^2]
)] + PolyLog[2, (1 + Sqrt[c*x^2])/2] - PolyLog[2, (1/2 - I/2)*(1 + Sqrt[c*
x^2])] - PolyLog[2, (1/2 + I/2)*(1 + Sqrt[c*x^2])])/(2*c*x)
```

Rubi [A] (verified)

Time = 2.09 (sec) , antiderivative size = 1085, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {6488, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex) (a + \operatorname{barctanh}(cx^2))^2 dx$$

$$\downarrow 6488$$

$$\int \left(d(a + \operatorname{barctanh}(cx^2))^2 + ex(a + \operatorname{barctanh}(cx^2))^2 \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& dx a^2 + \frac{2bd \arctan(\sqrt{cx}) a}{\sqrt{c}} - \frac{2bd \operatorname{darctanh}(\sqrt{cx}) a}{\sqrt{c}} - bdx \log(1 - cx^2) a + bdx \log(cx^2 + 1) a + \\
& \frac{ib^2 d \arctan(\sqrt{cx})^2}{\sqrt{c}} - \frac{b^2 \operatorname{darctanh}(\sqrt{cx})^2}{\sqrt{c}} + \frac{1}{2} e x^2 (a + b \operatorname{arctanh}(cx^2))^2 + \\
& \frac{e(a + b \operatorname{arctanh}(cx^2))^2}{2c} + \frac{1}{4} b^2 dx \log^2(1 - cx^2) + \frac{1}{4} b^2 dx \log^2(cx^2 + 1) + \\
& \frac{2b^2 \operatorname{darctanh}(\sqrt{cx}) \log\left(\frac{2}{1-\sqrt{cx}}\right)}{\sqrt{c}} - \frac{2b^2 d \arctan(\sqrt{cx}) \log\left(\frac{2}{1-i\sqrt{cx}}\right)}{\sqrt{c}} + \\
& \frac{b^2 d \arctan(\sqrt{cx}) \log\left(\frac{(1+i)(1-\sqrt{cx})}{1-i\sqrt{cx}}\right)}{\sqrt{c}} + \frac{2b^2 d \arctan(\sqrt{cx}) \log\left(\frac{2}{i\sqrt{cx}+1}\right)}{\sqrt{c}} - \\
& \frac{2b^2 \operatorname{darctanh}(\sqrt{cx}) \log\left(\frac{2}{\sqrt{cx}+1}\right)}{\sqrt{c}} + \frac{b^2 \operatorname{darctanh}(\sqrt{cx}) \log\left(-\frac{2\sqrt{c}(1-\sqrt{-cx})}{(\sqrt{-c}-\sqrt{c})(\sqrt{cx}+1)}\right)}{\sqrt{c}} + \\
& \frac{b^2 \operatorname{darctanh}(\sqrt{cx}) \log\left(\frac{2\sqrt{c}(\sqrt{-cx}+1)}{(\sqrt{-c}+\sqrt{c})(\sqrt{cx}+1)}\right)}{\sqrt{c}} + \frac{b^2 d \arctan(\sqrt{cx}) \log\left(\frac{(1-i)(\sqrt{cx}+1)}{1-i\sqrt{cx}}\right)}{\sqrt{c}} - \\
& \frac{be(a + b \operatorname{arctanh}(cx^2)) \log\left(\frac{2}{1-cx^2}\right)}{c} - \frac{b^2 d \arctan(\sqrt{cx}) \log(1 - cx^2)}{\sqrt{c}} + \\
& \frac{b^2 \operatorname{darctanh}(\sqrt{cx}) \log(1 - cx^2)}{\sqrt{c}} + \frac{b^2 d \arctan(\sqrt{cx}) \log(cx^2 + 1)}{\sqrt{c}} - \\
& \frac{b^2 \operatorname{darctanh}(\sqrt{cx}) \log(cx^2 + 1)}{\sqrt{c}} - \frac{1}{2} b^2 dx \log(1 - cx^2) \log(cx^2 + 1) + \\
& \frac{b^2 d \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-\sqrt{cx}}\right)}{\sqrt{c}} + \frac{ib^2 d \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i\sqrt{cx}}\right)}{\sqrt{c}} - \\
& \frac{ib^2 d \operatorname{PolyLog}\left(2, 1 - \frac{(1+i)(1-\sqrt{cx})}{1-i\sqrt{cx}}\right)}{2\sqrt{c}} + \frac{ib^2 d \operatorname{PolyLog}\left(2, 1 - \frac{2}{i\sqrt{cx}+1}\right)}{\sqrt{c}} + \\
& \frac{b^2 d \operatorname{PolyLog}\left(2, 1 - \frac{2}{\sqrt{cx}+1}\right)}{\sqrt{c}} - \frac{b^2 d \operatorname{PolyLog}\left(2, \frac{2\sqrt{c}(1-\sqrt{-cx})}{(\sqrt{-c}-\sqrt{c})(\sqrt{cx}+1)} + 1\right)}{2\sqrt{c}} - \\
& \frac{b^2 d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{c}(\sqrt{-cx}+1)}{(\sqrt{-c}+\sqrt{c})(\sqrt{cx}+1)}\right)}{2\sqrt{c}} - \frac{ib^2 d \operatorname{PolyLog}\left(2, 1 - \frac{(1-i)(\sqrt{cx}+1)}{1-i\sqrt{cx}}\right)}{2\sqrt{c}} - \\
& \frac{b^2 e \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-cx^2}\right)}{2c}
\end{aligned}$$

input `Int[(d + e*x)*(a + b*ArcTanh[c*x^2])^2,x]`

output

```
a^2*d*x + (2*a*b*d*ArcTan[Sqrt[c]*x])/Sqrt[c] + (I*b^2*d*ArcTan[Sqrt[c]*x]^2)/Sqrt[c] - (2*a*b*d*ArcTanh[Sqrt[c]*x])/Sqrt[c] - (b^2*d*ArcTanh[Sqrt[c]*x]^2)/Sqrt[c] + (e*(a + b*ArcTanh[c*x^2])^2)/(2*c) + (e*x^2*(a + b*ArcTanh[c*x^2])^2)/2 + (2*b^2*d*ArcTanh[Sqrt[c]*x]*Log[2/(1 - Sqrt[c]*x)])/Sqrt[c] - (2*b^2*d*ArcTan[Sqrt[c]*x]*Log[2/(1 - I*Sqrt[c]*x)])/Sqrt[c] + (b^2*d*ArcTan[Sqrt[c]*x]*Log[((1 + I)*(1 - Sqrt[c]*x))/(1 - I*Sqrt[c]*x)])/Sqrt[c] + (2*b^2*d*ArcTan[Sqrt[c]*x]*Log[2/(1 + I*Sqrt[c]*x)])/Sqrt[c] - (2*b^2*d*ArcTanh[Sqrt[c]*x]*Log[2/(1 + Sqrt[c]*x)])/Sqrt[c] + (b^2*d*ArcTanh[Sqrt[c]*x]*Log[(-2*Sqrt[c]*(1 - Sqrt[-c]*x))/((Sqrt[-c] - Sqrt[c])*(1 + Sqrt[c]*x))])/Sqrt[c] + (b^2*d*ArcTanh[Sqrt[c]*x]*Log[(2*Sqrt[c]*(1 + Sqrt[-c]*x))/((Sqrt[-c] + Sqrt[c])*(1 + Sqrt[c]*x))])/Sqrt[c] + (b^2*d*ArcTan[Sqrt[c]*x]*Log[((1 - I)*(1 + Sqrt[c]*x))/(1 - I*Sqrt[c]*x)])/Sqrt[c] - (b*e*(a + b*ArcTanh[c*x^2])*Log[2/(1 - c*x^2)])/c - a*b*d*x*Log[1 - c*x^2] - (b^2*d*ArcTan[Sqrt[c]*x]*Log[1 - c*x^2])/Sqrt[c] + (b^2*d*ArcTanh[Sqrt[c]*x]*Log[1 - c*x^2])/Sqrt[c] + (b^2*d*x*Log[1 - c*x^2]^2)/4 + a*b*d*x*Log[1 + c*x^2] + (b^2*d*ArcTan[Sqrt[c]*x]*Log[1 + c*x^2])/Sqrt[c] - (b^2*d*ArcTanh[Sqrt[c]*x]*Log[1 + c*x^2])/Sqrt[c] - (b^2*d*x*Log[1 - c*x^2]*Log[1 + c*x^2])/2 + (b^2*d*x*Log[1 + c*x^2]^2)/4 + (b^2*d*PolyLog[2, 1 - 2/(1 - Sqrt[c]*x)])/Sqrt[c] + (I*b^2*d*PolyLog[2, 1 - 2/(1 - I*Sqrt[c]*x)])/Sqrt[c] - ((I/2)*b^2*d*PolyLog[2, 1 - ((1 + I)*(1 - Sqrt[c]*x))/(1 - I*Sqrt[c]*x)])/...
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6488

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.))^p*((d_) + (e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcTanh[c*x^n])^p, (d + e*x)^m, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[p, 1] && IGtQ[m, 0]
```

Maple [F]

$$\int (ex + d) (a + b \operatorname{arctanh}(cx^2))^2 dx$$

input

```
int((e*x+d)*(a+b*arctanh(c*x^2))^2,x)
```

output `int((e*x+d)*(a+b*arctanh(c*x^2))^2,x)`

Fricas [F]

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (ex + d) (b \operatorname{artanh}(cx^2) + a)^2 dx$$

input `integrate((e*x+d)*(a+b*arctanh(c*x^2))^2,x, algorithm="fricas")`

output `integral(a^2*e*x + a^2*d + (b^2*e*x + b^2*d)*arctanh(c*x^2)^2 + 2*(a*b*e*x + a*b*d)*arctanh(c*x^2), x)`

Sympy [F]

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (a + b \operatorname{atanh}(cx^2))^2 (d + ex) dx$$

input `integrate((e*x+d)*(a+b*atanh(c*x**2))**2,x)`

output `Integral((a + b*atanh(c*x**2))**2*(d + e*x), x)`

Maxima [F]

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (ex + d) (b \operatorname{artanh}(cx^2) + a)^2 dx$$

input `integrate((e*x+d)*(a+b*arctanh(c*x^2))^2,x, algorithm="maxima")`

output

```
1/2*a^2*e*x^2 + (c*(2*arctan(sqrt(c)*x)/c^(3/2) + log((c*x - sqrt(c))/(c*x
+ sqrt(c)))/c^(3/2)) + 2*x*arctanh(c*x^2))*a*b*d + a^2*d*x + 1/2*(2*c*x^2
*arctanh(c*x^2) + log(-c^2*x^4 + 1))*a*b*e/c + 1/8*(b^2*e*x^2 + 2*b^2*d*x)
*log(-c*x^2 + 1)^2 - integrate(-1/4*((b^2*c*e*x^3 + b^2*c*d*x^2 - b^2*e*x
- b^2*d)*log(c*x^2 + 1)^2 - 2*(b^2*c*e*x^3 + 2*b^2*c*d*x^2 + (b^2*c*e*x^3
+ b^2*c*d*x^2 - b^2*e*x - b^2*d)*log(c*x^2 + 1))*log(-c*x^2 + 1))/(c*x^2 -
1), x)
```

Giac [F]

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (ex + d) (b \operatorname{arctanh}(cx^2) + a)^2 dx$$

input

```
integrate((e*x+d)*(a+b*arctanh(c*x^2))^2,x, algorithm="giac")
```

output

```
integrate((e*x + d)*(b*arctanh(c*x^2) + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^2))^2 dx = \int (a + b \operatorname{atanh}(cx^2))^2 (d + ex) dx$$

input

```
int((a + b*atanh(c*x^2))^2*(d + e*x),x)
```

output

```
int((a + b*atanh(c*x^2))^2*(d + e*x), x)
```

Reduce [F]

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^2))^2 dx$$

$$= \frac{4\sqrt{c} \operatorname{atan}\left(\frac{cx}{\sqrt{c}}\right) abd + 4\sqrt{c} \operatorname{atanh}(cx^2) abd + 4 \operatorname{atanh}(cx^2) abcdx + 2 \operatorname{atanh}(cx^2) abce x^2 - 2 \operatorname{atanh}(cx^2) a}{}$$

input `int((e*x+d)*(a+b*atanh(c*x^2))^2,x)`

output `(4*sqrt(c)*atan((c*x)/sqrt(c))*a*b*d + 4*sqrt(c)*atanh(c*x**2)*a*b*d + 4*a
tanh(c*x**2)*a*b*c*d*x + 2*atanh(c*x**2)*a*b*c*e*x**2 - 2*atanh(c*x**2)*a*
b*e + 4*sqrt(c)*log(sqrt(c)*x - 1)*a*b*d - 2*sqrt(c)*log(c*x**2 + 1)*a*b*d
+ 2*int(atanh(c*x**2)**2,x)*b**2*c*d + 2*int(atanh(c*x**2)**2*x,x)*b**2*c
*e + 2*log(c*x**2 + 1)*a*b*e + 2*a**2*c*d*x + a**2*c*e*x**2)/(2*c)`

$$3.31 \quad \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{d + ex} dx$$

Optimal result	282
Mathematica [N/A]	282
Rubi [N/A]	283
Maple [N/A]	283
Fricas [N/A]	284
Sympy [F(-1)]	284
Maxima [N/A]	284
Giac [N/A]	285
Mupad [N/A]	285
Reduce [N/A]	285

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + \operatorname{arctanh}(cx^2))^2}{d + ex} dx = \operatorname{Int}\left(\frac{(a + \operatorname{arctanh}(cx^2))^2}{d + ex}, x\right)$$

output `Defer(Int)((a+b*arctanh(c*x^2))^2/(e*x+d), x)`

Mathematica [N/A]

Not integrable

Time = 48.76 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + \operatorname{arctanh}(cx^2))^2}{d + ex} dx = \int \frac{(a + \operatorname{arctanh}(cx^2))^2}{d + ex} dx$$

input `Integrate[(a + b*ArcTanh[c*x^2])^2/(d + e*x), x]`

output `Integrate[(a + b*ArcTanh[c*x^2])^2/(d + e*x), x]`

Rubi [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{d + ex} dx$$

↓ 6490

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{d + ex} dx$$

input `Int[(a + b*ArcTanh[c*x^2])^2/(d + e*x), x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{ex + d} dx$$

input `int((a+b*arctanh(c*x^2))^2/(e*x+d), x)`

output `int((a+b*arctanh(c*x^2))^2/(e*x+d), x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.80

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{d + ex} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^2}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/(e*x+d),x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x^2)^2 + 2*a*b*arctanh(c*x^2) + a^2)/(e*x + d), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{d + ex} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**2))**2/(e*x+d),x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 81, normalized size of antiderivative = 4.05

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{d + ex} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^2}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/(e*x+d),x, algorithm="maxima")`

output `a^2*log(e*x + d)/e + integrate(1/4*b^2*(log(c*x^2 + 1) - log(-c*x^2 + 1))^2/(e*x + d) + a*b*(log(c*x^2 + 1) - log(-c*x^2 + 1))/(e*x + d), x)`

Giac [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{d + ex} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/(e*x+d),x, algorithm="giac")`

output `integrate((b*arctanh(c*x^2) + a)^2/(e*x + d), x)`

Mupad [N/A]

Not integrable

Time = 3.36 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{d + ex} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{d + ex} dx$$

input `int((a + b*atanh(c*x^2))^2/(d + e*x),x)`

output `int((a + b*atanh(c*x^2))^2/(d + e*x), x)`

Reduce [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 59, normalized size of antiderivative = 2.95

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{d + ex} dx = \frac{2 \left(\int \frac{\operatorname{atanh}(cx^2)}{ex+d} dx \right) a b e + \left(\int \frac{\operatorname{atanh}(cx^2)^2}{ex+d} dx \right) b^2 e + \log(ex + d) a^2}{e}$$

input `int((a+b*atanh(c*x^2))^2/(e*x+d),x)`

output

```
(2*int(atanh(c*x**2)/(d + e*x),x)*a*b*e + int(atanh(c*x**2)**2/(d + e*x),x)
)*b**2*e + log(d + e*x)*a**2)/e
```

$$3.32 \quad \int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(d + ex)^2} dx$$

Optimal result	287
Mathematica [N/A]	287
Rubi [N/A]	288
Maple [N/A]	288
Fricas [N/A]	289
Sympy [F(-1)]	289
Maxima [N/A]	289
Giac [N/A]	290
Mupad [N/A]	290
Reduce [N/A]	291

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + \operatorname{arctanh}(cx^2))^2}{(d + ex)^2} dx = \operatorname{Int} \left(\frac{(a + \operatorname{arctanh}(cx^2))^2}{(d + ex)^2}, x \right)$$

output `Defer(Int)((a+b*arctanh(c*x^2))^2/(e*x+d)^2,x)`

Mathematica [N/A]

Not integrable

Time = 85.62 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + \operatorname{arctanh}(cx^2))^2}{(d + ex)^2} dx = \int \frac{(a + \operatorname{arctanh}(cx^2))^2}{(d + ex)^2} dx$$

input `Integrate[(a + b*ArcTanh[c*x^2])^2/(d + e*x)^2,x]`

output `Integrate[(a + b*ArcTanh[c*x^2])^2/(d + e*x)^2, x]`

Rubi [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(d + ex)^2} dx$$

↓ 6490

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(d + ex)^2} dx$$

input `Int[(a + b*ArcTanh[c*x^2])^2/(d + e*x)^2,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(ex + d)^2} dx$$

input `int((a+b*arctanh(c*x^2))^2/(e*x+d)^2,x)`

output `int((a+b*arctanh(c*x^2))^2/(e*x+d)^2,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.35

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(d + ex)^2} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{(ex + d)^2} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/(e*x+d)^2,x, algorithm="fricas")`

output `integral((b^2*arctanh(c*x^2)^2 + 2*a*b*arctanh(c*x^2) + a^2)/(e^2*x^2 + 2*d*e*x + d^2), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(d + ex)^2} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**2))**2/(e*x+d)**2,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 1.28 (sec) , antiderivative size = 323, normalized size of antiderivative = 16.15

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(d + ex)^2} dx = \int \frac{(b \operatorname{artanh}(cx^2) + a)^2}{(ex + d)^2} dx$$

input `integrate((a+b*arctanh(c*x^2))^2/(e*x+d)^2,x, algorithm="maxima")`

output

```
((4*d*e*log(e*x + d)/(c^2*d^4 - e^4) + d*log(c*x^2 + 1)/(c*d^2*e + e^3) -
d*log(c*x^2 - 1)/(c*d^2*e - e^3) + 2*arctan(sqrt(c)*x)/((c*d^2 + e^2)*sqrt
(c)) + log((c*x - sqrt(c))/(c*x + sqrt(c)))/((c*d^2 - e^2)*sqrt(c))*c - 2
*arctanh(c*x^2)/(e^2*x + d*e))*a*b - 1/4*b^2*(log(-c*x^2 + 1)^2/(e^2*x + d
*e) + integrate(-(c*e*x^2 - e)*log(c*x^2 + 1)^2 + 2*(2*c*e*x^2 + 2*c*d*x
- (c*e*x^2 - e)*log(c*x^2 + 1))*log(-c*x^2 + 1))/(c*e^3*x^4 + 2*c*d*e^2*x^
3 - 2*d*e^2*x - d^2*e + (c*d^2*e - e^3)*x^2), x) - a^2/(e^2*x + d*e)
```

Giac [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(d + ex)^2} dx = \int \frac{(b \operatorname{arctanh}(cx^2) + a)^2}{(ex + d)^2} dx$$

input

```
integrate((a+b*arctanh(c*x^2))^2/(e*x+d)^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(c*x^2) + a)^2/(e*x + d)^2, x)
```

Mupad [N/A]

Not integrable

Time = 3.49 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(d + ex)^2} dx = \int \frac{(a + b \operatorname{atanh}(cx^2))^2}{(d + ex)^2} dx$$

input

```
int((a + b*atanh(c*x^2))^2/(d + e*x)^2,x)
```

output

```
int((a + b*atanh(c*x^2))^2/(d + e*x)^2, x)
```

Reduce [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 695, normalized size of antiderivative = 34.75

$$\int \frac{(a + b \operatorname{arctanh}(cx^2))^2}{(d + ex)^2} dx = \text{Too large to display}$$

input `int((a+b*atanh(c*x^2))^2/(e*x+d)^2,x)`

output

```
(2*sqrt(c)*atan((c*x)/sqrt(c))*a*b*c*d**4 + 2*sqrt(c)*atan((c*x)/sqrt(c))*
a*b*c*d**3*e*x - 2*sqrt(c)*atan((c*x)/sqrt(c))*a*b*d**2*e**2 - 2*sqrt(c)*a
tan((c*x)/sqrt(c))*a*b*d*e**3*x + 2*atanh(c*x**2)*a*b*c**2*d**4*x - 2*atan
h(c*x**2)*a*b*e**4*x + sqrt(c)*log(sqrt(c)*x - 1)*a*b*c*d**4 + sqrt(c)*log
(sqrt(c)*x - 1)*a*b*c*d**3*e*x + sqrt(c)*log(sqrt(c)*x - 1)*a*b*d**2*e**2
+ sqrt(c)*log(sqrt(c)*x - 1)*a*b*d*e**3*x - sqrt(c)*log(sqrt(c)*x + 1)*a*b
*c*d**4 - sqrt(c)*log(sqrt(c)*x + 1)*a*b*c*d**3*e*x - sqrt(c)*log(sqrt(c)*
x + 1)*a*b*d**2*e**2 - sqrt(c)*log(sqrt(c)*x + 1)*a*b*d*e**3*x + int(atanh
(c*x**2)**2/(d**2 + 2*d*e*x + e**2*x**2),x)*b**2*c**2*d**6 + int(atanh(c*x
**2)**2/(d**2 + 2*d*e*x + e**2*x**2),x)*b**2*c**2*d**5*e*x - int(atanh(c*x
**2)**2/(d**2 + 2*d*e*x + e**2*x**2),x)*b**2*d**2*e**4 - int(atanh(c*x**2)
**2/(d**2 + 2*d*e*x + e**2*x**2),x)*b**2*d*e**5*x - log(sqrt(c)*x - 1)*a*b
*c*d**3*e - log(sqrt(c)*x - 1)*a*b*c*d**2*e**2*x - log(sqrt(c)*x - 1)*a*b*
d*e**3 - log(sqrt(c)*x - 1)*a*b*e**4*x - log(sqrt(c)*x + 1)*a*b*c*d**3*e -
log(sqrt(c)*x + 1)*a*b*c*d**2*e**2*x - log(sqrt(c)*x + 1)*a*b*d*e**3 - lo
g(sqrt(c)*x + 1)*a*b*e**4*x - log(c*x**2 + 1)*a*b*c*d**3*e - log(c*x**2 +
1)*a*b*c*d**2*e**2*x + log(c*x**2 + 1)*a*b*d*e**3 + log(c*x**2 + 1)*a*b*e
**4*x + 4*log(d + e*x)*a*b*c*d**3*e + 4*log(d + e*x)*a*b*c*d**2*e**2*x + a
**2*c**2*d**4*x - a**2*e**4*x)/(d*(c**2*d**5 + c**2*d**4*e*x - d*e**4 - e**
5*x))
```

3.33 $\int (d + ex)^2 (a + \operatorname{barctanh}(cx^3)) dx$

Optimal result	292
Mathematica [A] (verified)	293
Rubi [A] (verified)	293
Maple [A] (verified)	295
Fricas [C] (verification not implemented)	297
Sympy [F(-1)]	297
Maxima [A] (verification not implemented)	298
Giac [A] (verification not implemented)	299
Mupad [B] (verification not implemented)	300
Reduce [B] (verification not implemented)	300

Optimal result

Integrand size = 18, antiderivative size = 300

$$\begin{aligned}
 \int (d + ex)^2 (a + \operatorname{barctanh}(cx^3)) dx = & -\frac{\sqrt{3}bde \arctan\left(\frac{1-2\sqrt[3]{c}x}{\sqrt{3}}\right)}{2c^{2/3}} \\
 & + \frac{\sqrt{3}bde \arctan\left(\frac{1+2\sqrt[3]{c}x}{\sqrt{3}}\right)}{2c^{2/3}} \\
 & + \frac{\sqrt{3}bd^2 \arctan\left(\frac{1+2c^{2/3}x^2}{\sqrt{3}}\right)}{2\sqrt[3]{c}} - \frac{bde \operatorname{arctanh}(\sqrt[3]{c}x)}{c^{2/3}} \\
 & + \frac{(d + ex)^3 (a + \operatorname{barctanh}(cx^3))}{3e} \\
 & - \frac{bde \operatorname{arctanh}\left(\frac{\sqrt[3]{c}x}{1+c^{2/3}x^2}\right)}{2c^{2/3}} + \frac{bd^2 \log(1 - c^{2/3}x^2)}{2\sqrt[3]{c}} \\
 & + \frac{b(cd^3 + e^3) \log(1 - cx^3)}{6ce} \\
 & - \frac{b(cd^3 - e^3) \log(1 + cx^3)}{6ce} \\
 & - \frac{bd^2 \log(1 + c^{2/3}x^2 + c^{4/3}x^4)}{4\sqrt[3]{c}}
 \end{aligned}$$

output

$$\begin{aligned}
& -1/2*3^{(1/2)}*b*d*e*\arctan(1/3*(1-2*c^{(1/3)}*x)*3^{(1/2)})/c^{(2/3)}+1/2*3^{(1/2)} \\
& *b*d*e*\arctan(1/3*(1+2*c^{(1/3)}*x)*3^{(1/2)})/c^{(2/3)}+1/2*3^{(1/2)}*b*d^2*\arctan \\
& n(1/3*(1+2*c^{(2/3)}*x^2)*3^{(1/2)})/c^{(1/3)}-b*d*e*\operatorname{arctanh}(c^{(1/3)}*x)/c^{(2/3)}+ \\
& 1/3*(e*x+d)^3*(a+b*\operatorname{arctanh}(c*x^3))/e-1/2*b*d*e*\operatorname{arctanh}(c^{(1/3)}*x/(1+c^{(2/3)} \\
&)*x^2))/c^{(2/3)}+1/2*b*d^2*\ln(1-c^{(2/3)}*x^2)/c^{(1/3)}+1/6*b*(c*d^3+e^3)*\ln(- \\
& c*x^3+1)/c/e-1/6*b*(c*d^3-e^3)*\ln(c*x^3+1)/c/e-1/4*b*d^2*\ln(1+c^{(2/3)}*x^2+ \\
& c^{(4/3)}*x^4)/c^{(1/3)}
\end{aligned}$$
Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 299, normalized size of antiderivative = 1.00

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx^3)) dx$$

$$\begin{aligned}
& 12acd^2x + 12acdex^2 + 4ace^2x^3 + 6\sqrt{3}b\sqrt[3]{cd}(\sqrt[3]{cd} + e) \arctan\left(\frac{-1+2\sqrt[3]{Cx}}{\sqrt{3}}\right) - 6\sqrt{3}b\sqrt[3]{cd}(\sqrt[3]{cd} - e) \arctan \\
& = \frac{\quad}{\quad}
\end{aligned}$$

input

$$\text{Integrate}[(d + e*x)^2*(a + b*\text{ArcTanh}[c*x^3]), x]$$

output

$$\begin{aligned}
& (12*a*c*d^2*x + 12*a*c*d*e*x^2 + 4*a*c*e^2*x^3 + 6*\text{Sqrt}[3]*b*c^{(1/3)}*d*(c^{(1/3)}*d + e)*\text{ArcTan}[(-1 + 2*c^{(1/3)}*x)/\text{Sqrt}[3]] - 6*\text{Sqrt}[3]*b*c^{(1/3)}*d*(c^{(1/3)}*d - e)*\text{ArcTan}[(1 + 2*c^{(1/3)}*x)/\text{Sqrt}[3]] + 4*b*c*x*(3*d^2 + 3*d*e*x + e^2*x^2)*\text{ArcTanh}[c*x^3] + 6*b*c^{(1/3)}*d*(c^{(1/3)}*d + e)*\text{Log}[1 - c^{(1/3)}*x] + 6*b*c^{(1/3)}*d*(c^{(1/3)}*d - e)*\text{Log}[1 + c^{(1/3)}*x] - 3*b*c^{(1/3)}*d*(c^{(1/3)}*d - e)*\text{Log}[1 - c^{(1/3)}*x + c^{(2/3)}*x^2] - 3*b*c^{(1/3)}*d*(c^{(1/3)}*d + e)*\text{Log}[1 + c^{(1/3)}*x + c^{(2/3)}*x^2] + 2*b*e^2*\text{Log}[1 - c^2*x^6])/(12*c)
\end{aligned}$$
Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 326, normalized size of antiderivative = 1.09, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6486, 2370, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int (d + ex)^2 (a + \operatorname{barctanh}(cx^3)) dx \\
& \quad \downarrow \text{6486} \\
& \frac{(d + ex)^3 (a + \operatorname{barctanh}(cx^3))}{3e} - \frac{bc \int \frac{x^2(d+ex)^3}{1-c^2x^6} dx}{e} \\
& \quad \downarrow \text{2370} \\
& \frac{(d + ex)^3 (a + \operatorname{barctanh}(cx^3))}{3e} - \frac{bc \int \left(\frac{3de^2x^4}{1-c^2x^6} + \frac{3d^2ex^3}{1-c^2x^6} + \frac{(d^3+e^3x^3)x^2}{1-c^2x^6} \right) dx}{e} \\
& \quad \downarrow \text{2009} \\
& \frac{(d + ex)^3 (a + \operatorname{barctanh}(cx^3))}{3e} - \\
& bc \left(-\frac{\sqrt{3}d^2e \arctan\left(\frac{2c^{2/3}x^2+1}{\sqrt{3}}\right)}{2c^{4/3}} + \frac{\sqrt{3}de^2 \arctan\left(\frac{1}{\sqrt{3}} - \frac{2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{2c^{5/3}} - \frac{\sqrt{3}de^2 \arctan\left(\frac{2\sqrt[3]{Cx}}{\sqrt{3}} + \frac{1}{\sqrt{3}}\right)}{2c^{5/3}} + \frac{de^2 \operatorname{arctanh}\left(\sqrt[3]{Cx}\right)}{c^{5/3}} + \frac{d^3 \operatorname{arctanh}\left(\sqrt[3]{Cx}\right)}{3c} \right)
\end{aligned}$$

input `Int[(d + e*x)^2*(a + b*ArcTanh[c*x^3]),x]`

output `((d + e*x)^3*(a + b*ArcTanh[c*x^3]))/(3*e) - (b*c*((Sqrt[3]*d*e^2*ArcTan[1/Sqrt[3] - (2*c^(1/3)*x)/Sqrt[3]])/(2*c^(5/3)) - (Sqrt[3]*d*e^2*ArcTan[1/Sqrt[3] + (2*c^(1/3)*x)/Sqrt[3]])/(2*c^(5/3)) - (Sqrt[3]*d^2*e*ArcTan[(1 + 2*c^(2/3)*x^2)/Sqrt[3]])/(2*c^(4/3)) + (d*e^2*ArcTanh[c^(1/3)*x])/c^(5/3) + (d^3*ArcTanh[c*x^3])/(3*c) - (d^2*e*Log[1 - c^(2/3)*x^2])/(2*c^(4/3)) - (d*e^2*Log[1 - c^(1/3)*x + c^(2/3)*x^2])/(4*c^(5/3)) + (d*e^2*Log[1 + c^(1/3)*x + c^(2/3)*x^2])/(4*c^(5/3)) + (d^2*e*Log[1 + c^(2/3)*x^2 + c^(4/3)*x^4])/(4*c^(4/3)) - (e^3*Log[1 - c^2*x^6])/(6*c^2))/e`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2370 `Int[((Pq_)*((c_)*(x_)^(m_)))/((a_) + (b_)*(x_)^(n_)), x_Symbol] := With[
 {v = Sum[(c*x)^(m + ii)*((Coeff[Pq, x, ii] + Coeff[Pq, x, n/2 + ii]*x^(n/2)
)/(c^ii*(a + b*x^n))), {ii, 0, n/2 - 1}], Int[v, x] /; SumQ[v]] /; FreeQ[{
 a, b, c, m}, x] && PolyQ[Pq, x] && IGtQ[n/2, 0] && Expon[Pq, x] < n`

rule 6486 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)]*(b_))*((d_) + (e_)*(x_)^(m_)), x_
 Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(e*(m + 1))), x]
 - Simp[b*c*(n/(e*(m + 1))) Int[x^(n - 1)*((d + e*x)^(m + 1)/(1 - c^2*x^(2
 *n))), x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 467, normalized size of antiderivative = 1.56

method	result
default	$\frac{a(ex+d)^3}{3e} + b \left(\frac{e^2 \operatorname{arctanh}(cx^3)x^3}{3} + e \operatorname{arctanh}(cx^3)x^2d + \operatorname{arctanh}(cx^3)xd^2 + \frac{\operatorname{arctanh}(cx^3)d^3}{3e} - \frac{c}{3e} \right)$

input `int((e*x+d)^2*(a+b*arctanh(c*x^3)),x,method=_RETURNVERBOSE)`

output `1/3*a*(e*x+d)^3/e+b*(1/3*e^2*arctanh(c*x^3)*x^3+e*arctanh(c*x^3)*x^2*d+arctanh(c*x^3)*x*d^2+1/3/e*arctanh(c*x^3)*d^3-1/e*c*(1/2*(-3*d^2*e*(1/3/c/(1/c)^(2/3)*ln(x-(1/c)^(1/3))-1/6/c/(1/c)^(2/3)*ln(x^2+(1/c)^(1/3)*x+(1/c)^(2/3))-1/3/c/(1/c)^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x+1))))-3*d*e^2*(1/3/c/(1/c)^(1/3)*ln(x-(1/c)^(1/3))-1/6/c/(1/c)^(1/3)*ln(x^2+(1/c)^(1/3)*x+(1/c)^(2/3))+1/3*3^(1/2)/c/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x+1)))+1/3*(-c*d^3-e^3)/c*ln(c*x^3-1))/c+1/2*(-3*d^2*e*(1/3/c/(1/c)^(2/3)*ln(x+(1/c)^(1/3))-1/6/c/(1/c)^(2/3)*ln(x^2-(1/c)^(1/3)*x+(1/c)^(2/3)))+1/3/c/(1/c)^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x-1)))-3*d*e^2*(-1/3/c/(1/c)^(1/3)*ln(x+(1/c)^(1/3))+1/6/c/(1/c)^(1/3)*ln(x^2-(1/c)^(1/3)*x+(1/c)^(2/3))+1/3*3^(1/2)/c/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x-1)))+1/3*(c*d^3-e^3)/c*ln(c*x^3+1))/c)`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.99 (sec) , antiderivative size = 9282, normalized size of antiderivative = 30.94

$$\int (d + ex)^2 (a + \operatorname{arctanh}(cx^3)) dx = \text{Too large to display}$$

input `integrate((e*x+d)^2*(a+b*arctanh(c*x^3)),x, algorithm="fricas")`

output Too large to include

Sympy [F(-1)]

Timed out.

$$\int (d + ex)^2 (a + \operatorname{arctanh}(cx^3)) dx = \text{Timed out}$$

input `integrate((e*x+d)**2*(a+b*atanh(c*x**3)),x)`

output Timed out

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 295, normalized size of antiderivative = 0.98

$$\int (d + ex)^2 (a + b \operatorname{arctanh}(cx^3)) dx = \frac{1}{3} ae^2 x^3 + adex^2 + \frac{1}{4} \left(c \left(\frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{\frac{4}{3}}x^2 + c^{\frac{2}{3}})}{3c^{\frac{2}{3}}}\right)}{c^{\frac{4}{3}}} - \frac{\log\left(c^{\frac{4}{3}}x^4 + c^{\frac{2}{3}}x^2 + 1\right)}{c^{\frac{4}{3}}} + \frac{2\log\left(\frac{c^{\frac{2}{3}}x^2 - 1}{c^{\frac{2}{3}}}\right)}{c^{\frac{4}{3}}} \right) + 4x \operatorname{arctanh}(cx^3) + \frac{1}{4} \left(4x^2 \operatorname{arctanh}(cx^3) + c \left(\frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{\frac{2}{3}}x + c^{\frac{1}{3}})}{3c^{\frac{1}{3}}}\right)}{c^{\frac{5}{3}}} + \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{\frac{2}{3}}x - c^{\frac{1}{3}})}{3c^{\frac{1}{3}}}\right)}{c^{\frac{5}{3}}} - \frac{\log\left(c^{\frac{2}{3}}x^2 + 1\right)}{c^{\frac{5}{3}}} \right) + ad^2x + \frac{(2cx^3 \operatorname{arctanh}(cx^3) + \log(-c^2x^6 + 1))be^2}{6c} \right.$$

input `integrate((e*x+d)^2*(a+b*arctanh(c*x^3)),x, algorithm="maxima")`

output `1/3*a*e^2*x^3 + a*d*e*x^2 + 1/4*(c*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(4/3)*x^2 + c^(2/3)))/c^(2/3))/c^(4/3) - log(c^(4/3)*x^4 + c^(2/3)*x^2 + 1)/c^(4/3) + 2*log((c^(2/3)*x^2 - 1)/c^(2/3))/c^(4/3) + 4*x*arctanh(c*x^3))*b*d^2 + 1/4*(4*x^2*arctanh(c*x^3) + c*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*x + c^(1/3)))/c^(1/3))/c^(5/3) + 2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*x - c^(1/3)))/c^(1/3))/c^(5/3) - log(c^(2/3)*x^2 + c^(1/3)*x + 1)/c^(5/3) + log(c^(2/3)*x^2 - c^(1/3)*x + 1)/c^(5/3) - 2*log((c^(1/3)*x + 1)/c^(1/3))/c^(5/3) + 2*log((c^(1/3)*x - 1)/c^(1/3))/c^(5/3))*b*d*e + a*d^2*x + 1/6*(2*c*x^3*arctanh(c*x^3) + log(-c^2*x^6 + 1))*b*e^2/c`

Giac [A] (verification not implemented)

Time = 11.46 (sec) , antiderivative size = 346, normalized size of antiderivative = 1.15

$$\begin{aligned}
& \int (d + ex)^2 (a + b \operatorname{arctanh}(cx^3)) dx \\
&= \frac{1}{3} ae^2 x^3 + adex^2 + ad^2 x + \frac{1}{6} (be^2 x^3 + 3 bde x^2 + 3 bd^2 x) \log \left(-\frac{cx^3 + 1}{cx^3 - 1} \right) \\
&\quad - \frac{\sqrt{3} (bcd^2 |c|^{\frac{2}{3}} - bcde |c|^{\frac{1}{3}}) \arctan \left(\frac{1}{3} \sqrt{3} \left(2x + \frac{1}{|c|^{\frac{1}{3}}} \right) |c|^{\frac{1}{3}} \right)}{2c^2} \\
&\quad + \frac{\sqrt{3} (bcd^2 |c|^{\frac{2}{3}} + bcde |c|^{\frac{1}{3}}) \arctan \left(\frac{1}{3} \sqrt{3} \left(2x - \frac{1}{|c|^{\frac{1}{3}}} \right) |c|^{\frac{1}{3}} \right)}{2c^2} \\
&\quad - \frac{(3bcd^2 |c|^{\frac{2}{3}} + 3bcde |c|^{\frac{1}{3}} - 2bce^2) \log \left(x^2 + \frac{x}{|c|^{\frac{1}{3}}} + \frac{1}{|c|^{\frac{2}{3}}} \right)}{12c^2} \\
&\quad - \frac{(3bcd^2 |c|^{\frac{2}{3}} - 3bcde |c|^{\frac{1}{3}} - 2bce^2) \log \left(x^2 - \frac{x}{|c|^{\frac{1}{3}}} + \frac{1}{|c|^{\frac{2}{3}}} \right)}{12c^2} \\
&\quad + \frac{(3bcd^2 |c|^{\frac{2}{3}} - 3bcde |c|^{\frac{1}{3}} + bce^2) \log \left(\left| x + \frac{1}{|c|^{\frac{1}{3}}} \right| \right)}{6c^2} \\
&\quad + \frac{(3bcd^2 |c|^{\frac{2}{3}} + 3bcde |c|^{\frac{1}{3}} + bce^2) \log \left(\left| x - \frac{1}{|c|^{\frac{1}{3}}} \right| \right)}{6c^2}
\end{aligned}$$

```
input integrate((e*x+d)^2*(a+b*arctanh(c*x^3)),x, algorithm="giac")
```

```
output 1/3*a*e^2*x^3 + a*d*e*x^2 + a*d^2*x + 1/6*(b*e^2*x^3 + 3*b*d*e*x^2 + 3*b*d
^2*x)*log(-(c*x^3 + 1)/(c*x^3 - 1)) - 1/2*sqrt(3)*(b*c*d^2*abs(c)^(2/3) -
b*c*d*e*abs(c)^(1/3))*arctan(1/3*sqrt(3)*(2*x + 1/abs(c)^(1/3))*abs(c)^(1/
3))/c^2 + 1/2*sqrt(3)*(b*c*d^2*abs(c)^(2/3) + b*c*d*e*abs(c)^(1/3))*arctan
(1/3*sqrt(3)*(2*x - 1/abs(c)^(1/3))*abs(c)^(1/3))/c^2 - 1/12*(3*b*c*d^2*ab
s(c)^(2/3) + 3*b*c*d*e*abs(c)^(1/3) - 2*b*c*e^2)*log(x^2 + x/abs(c)^(1/3)
+ 1/abs(c)^(2/3))/c^2 - 1/12*(3*b*c*d^2*abs(c)^(2/3) - 3*b*c*d*e*abs(c)^(1
/3) - 2*b*c*e^2)*log(x^2 - x/abs(c)^(1/3) + 1/abs(c)^(2/3))/c^2 + 1/6*(3*b
*c*d^2*abs(c)^(2/3) - 3*b*c*d*e*abs(c)^(1/3) + b*c*e^2)*log(abs(x + 1/abs(
c)^(1/3)))/c^2 + 1/6*(3*b*c*d^2*abs(c)^(2/3) + 3*b*c*d*e*abs(c)^(1/3) + b*
c*e^2)*log(abs(x - 1/abs(c)^(1/3)))/c^2
```

Mupad [B] (verification not implemented)

Time = 3.63 (sec) , antiderivative size = 1081, normalized size of antiderivative = 3.60

$$\int (d + ex)^2 (a + \operatorname{barctanh}(cx^3)) dx = \text{Too large to display}$$

input `int((a + b*atanh(c*x^3))*(d + e*x)^2,x)`

output `symsum(log(x*(6*b^5*c^7*d^2*e^8 + 162*b^5*c^9*d^8*e^2) + root(216*c^3*z^3 - 108*b*c^2*e^2*z^2 - 162*b^2*c^2*d^3*e*z + 18*b^2*c*e^4*z - 27*b^3*c^2*d^6 - b^3*e^6, z, k)*(x*(486*b^4*c^10*d^8 - 90*b^4*c^8*d^2*e^6) + root(216*c^3*z^3 - 108*b*c^2*e^2*z^2 - 162*b^2*c^2*d^3*e*z + 18*b^2*c*e^4*z - 27*b^3*c^2*d^6 - b^3*e^6, z, k)*(root(216*c^3*z^3 - 108*b*c^2*e^2*z^2 - 162*b^2*c^2*d^3*e*z + 18*b^2*c*e^4*z - 27*b^3*c^2*d^6 - b^3*e^6, z, k)*(3888*b^2*c^10*d^3*e - 3888*root(216*c^3*z^3 - 108*b*c^2*e^2*z^2 - 162*b^2*c^2*d^3*e*z + 18*b^2*c*e^4*z - 27*b^3*c^2*d^6 - b^3*e^6, z, k)*b*c^11*d^2*x + 648*b^2*c^10*d^2*e^2*x) - 972*b^3*c^9*d^3*e^3 + 324*b^3*c^9*d^2*e^4*x)) + 243*b^5*c^9*d^9*e + 9*b^5*c^7*d^3*e^7)*root(216*c^3*z^3 - 108*b*c^2*e^2*z^2 - 162*b^2*c^2*d^3*e*z + 18*b^2*c*e^4*z - 27*b^3*c^2*d^6 - b^3*e^6, z, k), k, 1, 3) + symsum(log(x*(6*b^5*c^7*d^2*e^8 + 162*b^5*c^9*d^8*e^2) + root(216*c^3*z^3 - 108*b*c^2*e^2*z^2 + 162*b^2*c^2*d^3*e*z + 18*b^2*c*e^4*z - 27*b^3*c^2*d^6 - b^3*e^6, z, k)*(x*(486*b^4*c^10*d^8 - 90*b^4*c^8*d^2*e^6) + root(216*c^3*z^3 - 108*b*c^2*e^2*z^2 + 162*b^2*c^2*d^3*e*z + 18*b^2*c*e^4*z - 27*b^3*c^2*d^6 - b^3*e^6, z, k)*(root(216*c^3*z^3 - 108*b*c^2*e^2*z^2 + 162*b^2*c^2*d^3*e*z + 18*b^2*c*e^4*z - 27*b^3*c^2*d^6 - b^3*e^6, z, k)*(3888*b^2*c^10*d^3*e - 3888*root(216*c^3*z^3 - 108*b*c^2*e^2*z^2 + 162*b^2*c^2*d^3*e*z + 18*b^2*c*e^4*z - 27*b^3*c^2*d^6 - b^3*e^6, z, k)*b*c^11*d^2*x + 648*b^2*c^10*d^2*e^2*x) - 972*b^3*c^9*d^3*e^3 + 324*b^3*c^9*d^2*e^4*x))...`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 369, normalized size of antiderivative = 1.23

$$\int (d + ex)^2 (a + \operatorname{barctanh}(cx^3)) dx$$

$$= \frac{6c^{\frac{4}{3}}\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x-1}{\sqrt{3}}\right) b d^2 + 6\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x-1}{\sqrt{3}}\right) bcde - 6c^{\frac{4}{3}}\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x+1}{\sqrt{3}}\right) b d^2 + 6\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x+1}{\sqrt{3}}\right) bc}{}$$

input `int((e*x+d)^2*(a+b*atanh(c*x^3)),x)`

output

```
(6*c**(1/3)*sqrt(3)*atan((2*c**(1/3)*x - 1)/sqrt(3))*b*c*d**2 + 6*sqrt(3)*
atan((2*c**(1/3)*x - 1)/sqrt(3))*b*c*d*e - 6*c**(1/3)*sqrt(3)*atan((2*c**(
1/3)*x + 1)/sqrt(3))*b*c*d**2 + 6*sqrt(3)*atan((2*c**(1/3)*x + 1)/sqrt(3))
*b*c*d*e + 12*c**(2/3)*atanh(c*x**3)*b*c*d**2*x + 12*c**(2/3)*atanh(c*x**3
)*b*c*d*e*x**2 + 4*c**(2/3)*atanh(c*x**3)*b*c*e**2*x**3 - 4*c**(2/3)*atanh
(c*x**3)*b*e**2 + 6*c**(1/3)*atanh(c*x**3)*b*c*d**2 + 6*atanh(c*x**3)*b*c*
d*e + 4*c**(2/3)*log(c**(2/3)*x**2 - c**(1/3)*x + 1)*b*e**2 + 4*c**(2/3)*l
og(c**(2/3)*x + c**(1/3))*b*e**2 + 12*c**(2/3)*a*c*d**2*x + 12*c**(2/3)*a*
c*d*e*x**2 + 4*c**(2/3)*a*c*e**2*x**3 - 6*c**(1/3)*log(c**(2/3)*x**2 - c**
(1/3)*x + 1)*b*c*d**2 + 3*c**(1/3)*log(c**(2/3)*x + c**(1/3))*b*c*d**2 + 9
*c**(1/3)*log(c**(2/3)*x - c**(1/3))*b*c*d**2 - 9*log(c**(2/3)*x + c**(1/3
))*b*c*d*e + 9*log(c**(2/3)*x - c**(1/3))*b*c*d*e)/(12*c**(2/3)*c)
```

3.34 $\int (d + ex) (a + b \operatorname{arctanh}(cx^3)) dx$

Optimal result	302
Mathematica [A] (verified)	303
Rubi [A] (verified)	304
Maple [A] (verified)	305
Fricas [C] (verification not implemented)	306
Sympy [F(-1)]	306
Maxima [A] (verification not implemented)	307
Giac [A] (verification not implemented)	308
Mupad [B] (verification not implemented)	309
Reduce [B] (verification not implemented)	310

Optimal result

Integrand size = 16, antiderivative size = 250

$$\begin{aligned}
 \int (d + ex) (a + b \operatorname{arctanh}(cx^3)) dx = & -\frac{\sqrt{3}be \arctan\left(\frac{1-2\sqrt[3]{cx}}{\sqrt{3}}\right)}{4c^{2/3}} \\
 & + \frac{\sqrt{3}be \arctan\left(\frac{1+2\sqrt[3]{cx}}{\sqrt{3}}\right)}{4c^{2/3}} \\
 & + \frac{\sqrt{3}bd \arctan\left(\frac{1+2c^{2/3}x^2}{\sqrt{3}}\right)}{2\sqrt[3]{c}} \\
 & - \frac{b \operatorname{arctanh}(\sqrt[3]{cx})}{2c^{2/3}} - \frac{bd^2 \operatorname{arctanh}(cx^3)}{2e} \\
 & + \frac{(d + ex)^2 (a + b \operatorname{arctanh}(cx^3))}{2e} \\
 & - \frac{b \operatorname{arctanh}\left(\frac{\sqrt[3]{cx}}{1+c^{2/3}x^2}\right)}{4c^{2/3}} + \frac{bd \log(1 - c^{2/3}x^2)}{2\sqrt[3]{c}} \\
 & - \frac{bd \log(1 + c^{2/3}x^2 + c^{4/3}x^4)}{4\sqrt[3]{c}}
 \end{aligned}$$

output

```
-1/4*3^(1/2)*b*e*arctan(1/3*(1-2*c^(1/3)*x)*3^(1/2))/c^(2/3)+1/4*3^(1/2)*b
*e*arctan(1/3*(1+2*c^(1/3)*x)*3^(1/2))/c^(2/3)+1/2*3^(1/2)*b*d*arctan(1/3*
(1+2*c^(2/3)*x^2)*3^(1/2))/c^(1/3)-1/2*b*e*arctanh(c^(1/3)*x)/c^(2/3)-1/2*
b*d^2*arctanh(c*x^3)/e+1/2*(e*x+d)^2*(a+b*arctanh(c*x^3))/e-1/4*b*e*arctan
h(c^(1/3)*x/(1+c^(2/3)*x^2))/c^(2/3)+1/2*b*d*ln(1-c^(2/3)*x^2)/c^(1/3)-1/4
*b*d*ln(1+c^(2/3)*x^2+c^(4/3)*x^4)/c^(1/3)
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 333, normalized size of antiderivative = 1.33

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^3)) dx = adx + \frac{1}{2} aex^2 + \frac{\sqrt{3}be \operatorname{arctan}\left(\frac{-1+2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{4c^{2/3}}$$

$$+ \frac{\sqrt{3}be \operatorname{arctan}\left(\frac{1+2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{4c^{2/3}} + bdx \operatorname{arctanh}(cx^3) + \frac{1}{2} bex^2 \operatorname{arctanh}(cx^3)$$

$$+ \frac{be \log(1 - \sqrt[3]{cx})}{4c^{2/3}} - \frac{be \log(1 + \sqrt[3]{cx})}{4c^{2/3}} + \frac{be \log(1 - \sqrt[3]{cx} + c^{2/3}x^2)}{8c^{2/3}}$$

$$- \frac{be \log(1 + \sqrt[3]{cx} + c^{2/3}x^2)}{8c^{2/3}} - \frac{bd\left(-2\sqrt{3} \operatorname{arctan}\left(\frac{-1+2\sqrt[3]{Cx}}{\sqrt{3}}\right) + 2\sqrt{3} \operatorname{arctan}\left(\frac{1+2\sqrt[3]{Cx}}{\sqrt{3}}\right) - 2 \log(1 - \sqrt[3]{Cx})\right)}{4}$$

input

```
Integrate[(d + e*x)*(a + b*ArcTanh[c*x^3]),x]
```

output

```
a*d*x + (a*e*x^2)/2 + (Sqrt[3]*b*e*ArcTan[(-1 + 2*c^(1/3)*x)/Sqrt[3]])/(4*
c^(2/3)) + (Sqrt[3]*b*e*ArcTan[(1 + 2*c^(1/3)*x)/Sqrt[3]])/(4*c^(2/3)) + b
*d*x*ArcTanh[c*x^3] + (b*e*x^2*ArcTanh[c*x^3])/2 + (b*e*Log[1 - c^(1/3)*x]
)/(4*c^(2/3)) - (b*e*Log[1 + c^(1/3)*x])/(4*c^(2/3)) + (b*e*Log[1 - c^(1/3)
]*x + c^(2/3)*x^2])/(8*c^(2/3)) - (b*e*Log[1 + c^(1/3)*x + c^(2/3)*x^2])/(
8*c^(2/3)) - (b*d*(-2*Sqrt[3]*ArcTan[(-1 + 2*c^(1/3)*x)/Sqrt[3]] + 2*Sqrt[
3]*ArcTan[(1 + 2*c^(1/3)*x)/Sqrt[3]] - 2*Log[1 - c^(1/3)*x] - 2*Log[1 + c
^(1/3)*x] + Log[1 - c^(1/3)*x + c^(2/3)*x^2] + Log[1 + c^(1/3)*x + c^(2/3)*
x^2]))/(4*c^(1/3))
```


Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.19, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6486, 2370, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (d + ex) (a + \operatorname{arctanh}(cx^3)) dx \\
 & \quad \downarrow 6486 \\
 & \frac{(d + ex)^2 (a + \operatorname{arctanh}(cx^3))}{2e} - \frac{3bc \int \frac{x^2(d+ex)^2}{1-c^2x^6} dx}{2e} \\
 & \quad \downarrow 2370 \\
 & \frac{(d + ex)^2 (a + \operatorname{arctanh}(cx^3))}{2e} - \frac{3bc \int \left(\frac{e^2 x^4}{1-c^2x^6} + \frac{2dex^3}{1-c^2x^6} + \frac{d^2 x^2}{1-c^2x^6} \right) dx}{2e} \\
 & \quad \downarrow 2009 \\
 & \frac{(d + ex)^2 (a + \operatorname{arctanh}(cx^3))}{2e} - \\
 & 3bc \left(-\frac{de \operatorname{arctan}\left(\frac{2c^{2/3}x^2+1}{\sqrt{3}}\right)}{\sqrt{3}c^{4/3}} + \frac{e^2 \operatorname{arctan}\left(\frac{1}{\sqrt{3}} - \frac{2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{2\sqrt{3}c^{5/3}} - \frac{e^2 \operatorname{arctan}\left(\frac{2\sqrt[3]{Cx}}{\sqrt{3}} + \frac{1}{\sqrt{3}}\right)}{2\sqrt{3}c^{5/3}} + \frac{e^2 \operatorname{arctanh}\left(\sqrt[3]{Cx}\right)}{3c^{5/3}} + \frac{d^2 \operatorname{arctanh}(cx^3)}{3c} \right)
 \end{aligned}$$

2e

input `Int[(d + e*x)*(a + b*ArcTanh[c*x^3]), x]`

output `((d + e*x)^2*(a + b*ArcTanh[c*x^3])/(2*e) - (3*b*c*((e^2*ArcTan[1/Sqrt[3] - (2*c^(1/3)*x)/Sqrt[3]])/(2*Sqrt[3]*c^(5/3)) - (e^2*ArcTan[1/Sqrt[3] + (2*c^(1/3)*x)/Sqrt[3]])/(2*Sqrt[3]*c^(5/3)) - (d*e*ArcTan[(1 + 2*c^(2/3)*x^2)/Sqrt[3]])/(Sqrt[3]*c^(4/3)) + (e^2*ArcTanh[c^(1/3)*x])/(3*c^(5/3)) + (d^2*ArcTanh[c*x^3])/(3*c) - (d*e*Log[1 - c^(2/3)*x^2])/(3*c^(4/3)) - (e^2*Log[1 - c^(1/3)*x + c^(2/3)*x^2])/(12*c^(5/3)) + (e^2*Log[1 + c^(1/3)*x + c^(2/3)*x^2])/(12*c^(5/3)) + (d*e*Log[1 + c^(2/3)*x^2 + c^(4/3)*x^4])/(6*c^(4/3)))/(2*e)`

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2370 Int[((Pq_)*((c_)*(x_)^(m_)))/((a_) + (b_)*(x_)^(n_)), x_Symbol] := With[
{v = Sum[(c*x)^(m + ii)*((Coeff[Pq, x, ii] + Coeff[Pq, x, n/2 + ii])*x^(n/2)
)/(c^ii*(a + b*x^n))}, {ii, 0, n/2 - 1}]], Int[v, x] /; SumQ[v]] /; FreeQ[{
a, b, c, m}, x] && PolyQ[Pq, x] && IGtQ[n/2, 0] && Expon[Pq, x] < n
```

```
rule 6486 Int[((a_) + ArcTanh[(c_)*(x_)^(n_)]*(b_))*((d_) + (e_)*(x_)^(m_)), x_
Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(e*(m + 1))), x]
- Simp[b*c*(n/(e*(m + 1))) Int[x^(n - 1)*((d + e*x)^(m + 1)/(1 - c^2*x^(2
*n))), x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 363, normalized size of antiderivative = 1.45

method	result
default	$a\left(\frac{1}{2}ex^2 + dx\right) + \frac{b \operatorname{arctanh}(cx^3)x^2e}{2} + b \operatorname{arctanh}(cx^3) dx + \frac{bd \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{2c\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{bd \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{4c\left(\frac{1}{c}\right)^{\frac{2}{3}}}$
parts	$a\left(\frac{1}{2}ex^2 + dx\right) + \frac{b \operatorname{arctanh}(cx^3)x^2e}{2} + b \operatorname{arctanh}(cx^3) dx + \frac{bd \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{2c\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{bd \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{4c\left(\frac{1}{c}\right)^{\frac{2}{3}}}$
risch	$\frac{b\left(\frac{1}{2}ex^2 + dx\right) \ln(cx^3 + 1)}{2} + \frac{ae x^2}{2} + adx - \frac{bdx \ln(-cx^3 + 1)}{2} + \frac{bd \ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right)}{2c\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{bd \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{4c\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{bd\sqrt{3} \operatorname{arctanh}\left(\frac{x - \left(\frac{1}{c}\right)^{\frac{1}{3}}}{x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}}\right)}{2c\left(\frac{1}{c}\right)^{\frac{2}{3}}}$

```
input int((e*x+d)*(a+b*arctanh(c*x^3)),x,method=_RETURNVERBOSE)
```

output

```
a*(1/2*e*x^2+d*x)+1/2*b*arctanh(c*x^3)*x^2*e+b*arctanh(c*x^3)*d*x+1/2*b*d/c/(1/c)^(2/3)*ln(x-(1/c)^(1/3))-1/4*b*d/c/(1/c)^(2/3)*ln(x^2+(1/c)^(1/3)*x+(1/c)^(2/3))-1/2*b*d/c/(1/c)^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x+1))+1/4*b*e/c/(1/c)^(1/3)*ln(x-(1/c)^(1/3))-1/8*b*e/c/(1/c)^(1/3)*ln(x^2+(1/c)^(1/3)*x+(1/c)^(2/3))+1/4*b*e*3^(1/2)/c/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x+1))+1/2*b*d/c/(1/c)^(2/3)*ln(x+(1/c)^(1/3))-1/4*b*d/c/(1/c)^(2/3)*ln(x^2-(1/c)^(1/3)*x+(1/c)^(2/3))+1/2*b*d/c/(1/c)^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x-1))-1/4*b*e/c/(1/c)^(1/3)*ln(x+(1/c)^(1/3))+1/8*b*e/c/(1/c)^(1/3)*ln(x^2-(1/c)^(1/3)*x+(1/c)^(2/3))+1/4*b*e*3^(1/2)/c/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x-1))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.90 (sec) , antiderivative size = 3928, normalized size of antiderivative = 15.71

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^3)) dx = \text{Too large to display}$$

input

```
integrate((e*x+d)*(a+b*arctanh(c*x^3)),x, algorithm="fricas")
```

output

Too large to include

Sympy [F(-1)]

Timed out.

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^3)) dx = \text{Timed out}$$

input

```
integrate((e*x+d)*(a+b*atanh(c*x**3)),x)
```

output

Timed out

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 248, normalized size of antiderivative = 0.99

$$\int (d + ex) (a + b \operatorname{arctanh}(cx^3)) dx = \frac{1}{2} aex^2 + \frac{1}{4} \left(c \left(\frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{4/3}x^2 + c^{2/3})}{3c^{2/3}}\right)}{c^{4/3}} - \frac{\log\left(c^{4/3}x^4 + c^{2/3}x^2 + 1\right)}{c^{4/3}} + \frac{2\log\left(\frac{c^{2/3}x^2 - 1}{c^{2/3}}\right)}{c^{4/3}} \right) + 4x \operatorname{arctanh}(cx^3) \right) + \frac{1}{8} \left(4x^2 \operatorname{arctanh}(cx^3) + c \left(\frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{2/3}x + c^{1/3})}{3c^{1/3}}\right)}{c^{5/3}} + \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2c^{2/3}x - c^{1/3})}{3c^{1/3}}\right)}{c^{5/3}} - \frac{\log\left(c^{2/3}x^2 + c^{1/3}x + 1\right)}{c^{5/3}} - \frac{\log\left(c^{2/3}x^2 - c^{1/3}x + 1\right)}{c^{5/3}} - 2\log\left(\frac{c^{1/3}x + 1}{c^{1/3}}\right)/c^{5/3} + 2\log\left(\frac{c^{1/3}x - 1}{c^{1/3}}\right)/c^{5/3} \right) \right) + adx$$

input `integrate((e*x+d)*(a+b*arctanh(c*x^3)),x, algorithm="maxima")`

output

```
1/2*a*e*x^2 + 1/4*(c*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(4/3)*x^2 + c^(2/3)))/c^(2/3))/c^(4/3) - log(c^(4/3)*x^4 + c^(2/3)*x^2 + 1)/c^(4/3) + 2*log((c^(2/3)*x^2 - 1)/c^(2/3))/c^(4/3) + 4*x*arctanh(c*x^3))*b*d + 1/8*(4*x^2*arctanh(c*x^3) + c*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*x + c^(1/3))/c^(1/3))/c^(5/3) + 2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(2/3)*x - c^(1/3))/c^(1/3))/c^(5/3) - log(c^(2/3)*x^2 + c^(1/3)*x + 1)/c^(5/3) + log(c^(2/3)*x^2 - c^(1/3)*x + 1)/c^(5/3) - 2*log((c^(1/3)*x + 1)/c^(1/3))/c^(5/3) + 2*log((c^(1/3)*x - 1)/c^(1/3))/c^(5/3)))*b*e + a*d*x
```

Giac [A] (verification not implemented)

Time = 0.84 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.07

$$\begin{aligned}
& \int (d + ex) (a + b \operatorname{arctanh}(cx^3)) dx \\
&= \frac{1}{2} aex^2 + adx + \frac{1}{4} (bex^2 + 2bdx) \log\left(-\frac{cx^3 + 1}{cx^3 - 1}\right) \\
&\quad - \frac{(2\sqrt{3}bcd|c|^{\frac{1}{3}} - \sqrt{3}bce) \arctan\left(\frac{1}{3}\sqrt{3}\left(2x + \frac{1}{|c|^{\frac{1}{3}}}\right)|c|^{\frac{1}{3}}\right)}{4|c|^{\frac{5}{3}}} \\
&\quad + \frac{(2\sqrt{3}bcd|c|^{\frac{1}{3}} + \sqrt{3}bce) \arctan\left(\frac{1}{3}\sqrt{3}\left(2x - \frac{1}{|c|^{\frac{1}{3}}}\right)|c|^{\frac{1}{3}}\right)}{4|c|^{\frac{5}{3}}} \\
&\quad - \frac{(2bcd|c|^{\frac{1}{3}} + bce) \log\left(x^2 + \frac{x}{|c|^{\frac{1}{3}}} + \frac{1}{|c|^{\frac{2}{3}}}\right)}{8|c|^{\frac{5}{3}}} \\
&\quad - \frac{(2bcd|c|^{\frac{1}{3}} - bce) \log\left(x^2 - \frac{x}{|c|^{\frac{1}{3}}} + \frac{1}{|c|^{\frac{2}{3}}}\right)}{8|c|^{\frac{5}{3}}} \\
&\quad + \frac{(2bcd|c|^{\frac{1}{3}} - bce) \log\left(\left|x + \frac{1}{|c|^{\frac{1}{3}}}\right|\right)}{4|c|^{\frac{5}{3}}} + \frac{(2bcd|c|^{\frac{1}{3}} + bce) \log\left(\left|x - \frac{1}{|c|^{\frac{1}{3}}}\right|\right)}{4|c|^{\frac{5}{3}}}
\end{aligned}$$

input `integrate((e*x+d)*(a+b*arctanh(c*x^3)),x, algorithm="giac")`

output `1/2*a*e*x^2 + a*d*x + 1/4*(b*e*x^2 + 2*b*d*x)*log(-(c*x^3 + 1)/(c*x^3 - 1)) - 1/4*(2*sqrt(3)*b*c*d*abs(c)^(1/3) - sqrt(3)*b*c*e)*arctan(1/3*sqrt(3)*(2*x + 1/abs(c)^(1/3))*abs(c)^(1/3))/abs(c)^(5/3) + 1/4*(2*sqrt(3)*b*c*d*abs(c)^(1/3) + sqrt(3)*b*c*e)*arctan(1/3*sqrt(3)*(2*x - 1/abs(c)^(1/3))*abs(c)^(1/3))/abs(c)^(5/3) - 1/8*(2*b*c*d*abs(c)^(1/3) + b*c*e)*log(x^2 + x/abs(c)^(1/3) + 1/abs(c)^(2/3))/abs(c)^(5/3) - 1/8*(2*b*c*d*abs(c)^(1/3) - b*c*e)*log(x^2 - x/abs(c)^(1/3) + 1/abs(c)^(2/3))/abs(c)^(5/3) + 1/4*(2*b*c*d*abs(c)^(1/3) - b*c*e)*log(abs(x + 1/abs(c)^(1/3)))/abs(c)^(5/3) + 1/4*(2*b*c*d*abs(c)^(1/3) + b*c*e)*log(abs(x - 1/abs(c)^(1/3)))/abs(c)^(5/3)`

Mupad [B] (verification not implemented)

Time = 3.53 (sec) , antiderivative size = 621, normalized size of antiderivative = 2.48

$$\begin{aligned}
& \int (d + ex) (a + \operatorname{arctanh}(cx^3)) dx \\
&= \left(\sum_{k=1}^3 \ln \left(-\operatorname{root}(64c^2z^3 + 24b^2cdez - 8b^3cd^3 + b^3e^3, z, k) \left(\operatorname{root}(64c^2z^3 + 24b^2cdez - 8b^3cd^3 + b^3e^3, z, k) \right. \right. \right. \\
&\quad \left. \left. \left. + \frac{243b^5c^9d^4e}{2} + \frac{243b^5c^9d^3e^2x}{4} \right) \operatorname{root}(64c^2z^3 + 24b^2cdez - 8b^3cd^3 + b^3e^3, z, k) \right) \right) \\
&\quad + \left(\sum_{k=1}^3 \ln \left(-\operatorname{root}(64c^2z^3 - 24b^2cdez - 8b^3cd^3 - b^3e^3, z, k) \left(\operatorname{root}(64c^2z^3 - 24b^2cdez - 8b^3cd^3 - b^3e^3, z, k) \right. \right. \right. \\
&\quad \left. \left. \left. + \frac{243b^5c^9d^4e}{2} + \frac{243b^5c^9d^3e^2x}{4} \right) \operatorname{root}(64c^2z^3 - 24b^2cdez - 8b^3cd^3 - b^3e^3, z, k) \right) \right) \\
&\quad + \ln(cx^3 + 1) \left(\frac{bex^2}{4} + \frac{bdx}{2} \right) - \ln(1 - cx^3) \left(\frac{bex^2}{4} + \frac{bdx}{2} \right) + adx + \frac{aex^2}{2}
\end{aligned}$$

input `int((a + b*atanh(c*x^3))*(d + e*x),x)`

output

```

symsum(log((243*b^5*c^9*d^4*e)/2 - root(64*c^2*z^3 + 24*b^2*c*d*e*z - 8*b^3*c*d^3 + b^3*e^3, z, k)*(root(64*c^2*z^3 + 24*b^2*c*d*e*z - 8*b^3*c*d^3 + b^3*e^3, z, k)*(root(64*c^2*z^3 + 24*b^2*c*d*e*z - 8*b^3*c*d^3 + b^3*e^3, z, k)*(486*b^2*c^10*e^2*x - 1944*b^2*c^10*d*e + 3888*root(64*c^2*z^3 + 24*b^2*c*d*e*z - 8*b^3*c*d^3 + b^3*e^3, z, k)*b*c^11*d*x) - (243*b^3*c^9*e^3)/2) - 486*b^4*c^10*d^4*x) + (243*b^5*c^9*d^3*e^2*x)/4)*root(64*c^2*z^3 + 24*b^2*c*d*e*z - 8*b^3*c*d^3 + b^3*e^3, z, k), k, 1, 3) + symsum(log((243*b^5*c^9*d^4*e)/2 - root(64*c^2*z^3 - 24*b^2*c*d*e*z - 8*b^3*c*d^3 - b^3*e^3, z, k)*(root(64*c^2*z^3 - 24*b^2*c*d*e*z - 8*b^3*c*d^3 - b^3*e^3, z, k)*(root(64*c^2*z^3 - 24*b^2*c*d*e*z - 8*b^3*c*d^3 - b^3*e^3, z, k)*(486*b^2*c^10*e^2*x - 1944*b^2*c^10*d*e + 3888*root(64*c^2*z^3 - 24*b^2*c*d*e*z - 8*b^3*c*d^3 - b^3*e^3, z, k)*b*c^11*d*x) - (243*b^3*c^9*e^3)/2) - 486*b^4*c^10*d^4*x) + (243*b^5*c^9*d^3*e^2*x)/4)*root(64*c^2*z^3 - 24*b^2*c*d*e*z - 8*b^3*c*d^3 - b^3*e^3, z, k), k, 1, 3) + log(c*x^3 + 1)*((b*d*x)/2 + (b*e*x^2)/4) - log(1 - c*x^3)*((b*d*x)/2 + (b*e*x^2)/4) + a*d*x + (a*e*x^2)/2

```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 252, normalized size of antiderivative = 1.01

$$\int (d + ex) (a + \operatorname{barctanh}(cx^3)) dx$$

$$= \frac{4c^{\frac{1}{3}}\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x-1}{\sqrt{3}}\right) bd + 2\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x-1}{\sqrt{3}}\right) be - 4c^{\frac{1}{3}}\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x+1}{\sqrt{3}}\right) bd + 2\sqrt{3} \operatorname{atan}\left(\frac{2c^{\frac{1}{3}}x+1}{\sqrt{3}}\right) be + 8c^{\frac{2}{3}} \operatorname{atanh}(cx^3) b d x + 4c^{\frac{2}{3}} \operatorname{atanh}(cx^3) b e x^2 + 4c^{\frac{1}{3}} a \operatorname{tanh}(cx^3) b d + 2 \operatorname{atanh}(cx^3) b e + 8c^{\frac{2}{3}} a d x + 4c^{\frac{2}{3}} a e x^2 - 4c^{\frac{1}{3}} \log(c^{\frac{2}{3}} x^2 - c^{\frac{1}{3}} x + 1) b d + 2c^{\frac{1}{3}} \log(c^{\frac{2}{3}} x + c^{\frac{1}{3}}) b d + 6c^{\frac{1}{3}} \log(c^{\frac{2}{3}} x - c^{\frac{1}{3}}) b d - 3 \log(c^{\frac{2}{3}} x + c^{\frac{1}{3}}) b e + 3 \log(c^{\frac{2}{3}} x - c^{\frac{1}{3}}) b e}{(8c^{\frac{2}{3}})}$$

input `int((e*x+d)*(a+b*atanh(c*x^3)),x)`

output

```
(4*c**(1/3)*sqrt(3)*atan((2*c**(1/3)*x - 1)/sqrt(3))*b*d + 2*sqrt(3)*atan(
(2*c**(1/3)*x - 1)/sqrt(3))*b*e - 4*c**(1/3)*sqrt(3)*atan((2*c**(1/3)*x +
1)/sqrt(3))*b*d + 2*sqrt(3)*atan((2*c**(1/3)*x + 1)/sqrt(3))*b*e + 8*c**(2
/3)*atanh(c*x**3)*b*d*x + 4*c**(2/3)*atanh(c*x**3)*b*e*x**2 + 4*c**(1/3)*a
tanh(c*x**3)*b*d + 2*atanh(c*x**3)*b*e + 8*c**(2/3)*a*d*x + 4*c**(2/3)*a*e
*x**2 - 4*c**(1/3)*log(c**(2/3)*x**2 - c**(1/3)*x + 1)*b*d + 2*c**(1/3)*lo
g(c**(2/3)*x + c**(1/3))*b*d + 6*c**(1/3)*log(c**(2/3)*x - c**(1/3))*b*d -
3*log(c**(2/3)*x + c**(1/3))*b*e + 3*log(c**(2/3)*x - c**(1/3))*b*e)/(8*c
**(2/3))
```

3.35 $\int \frac{a+b\operatorname{arctanh}(cx^3)}{d+ex} dx$

Optimal result	312
Mathematica [C] (verified)	313
Rubi [A] (verified)	314
Maple [C] (verified)	316
Fricas [F]	317
Sympy [F(-1)]	317
Maxima [F]	317
Giac [F]	318
Mupad [F(-1)]	318
Reduce [F]	318

Optimal result

Integrand size = 18, antiderivative size = 523

$$\begin{aligned}
\int \frac{a + \operatorname{barctanh}(cx^3)}{d + ex} dx &= \frac{(a + \operatorname{barctanh}(cx^3)) \log(d + ex)}{e} \\
&+ \frac{b \log\left(\frac{e(1 - \sqrt[3]{Cx})}{\sqrt[3]{Cd+e}}\right) \log(d + ex)}{2e} \\
&- \frac{b \log\left(-\frac{e(1 + \sqrt[3]{Cx})}{\sqrt[3]{Cd-e}}\right) \log(d + ex)}{2e} \\
&+ \frac{b \log\left(-\frac{e(\sqrt[3]{-1} + \sqrt[3]{Cx})}{\sqrt[3]{Cd - \sqrt[3]{-1}e}}\right) \log(d + ex)}{2e} \\
&- \frac{b \log\left(-\frac{e((-1)^{2/3} + \sqrt[3]{Cx})}{\sqrt[3]{Cd - (-1)^{2/3}e}}\right) \log(d + ex)}{2e} \\
&+ \frac{b \log\left(\frac{(-1)^{2/3}e(1 + \sqrt[3]{-1}\sqrt[3]{Cx})}{\sqrt[3]{Cd + (-1)^{2/3}e}}\right) \log(d + ex)}{2e} \\
&- \frac{b \log\left(\frac{\sqrt[3]{-1}e(1 + (-1)^{2/3}\sqrt[3]{Cx})}{\sqrt[3]{Cd + \sqrt[3]{-1}e}}\right) \log(d + ex)}{2e} \\
&- \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt[3]{C(d+ex)}}{\sqrt[3]{Cd-e}}\right)}{2e} + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt[3]{C(d+ex)}}{\sqrt[3]{Cd+e}}\right)}{2e} \\
&+ \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt[3]{C(d+ex)}}{\sqrt[3]{Cd - \sqrt[3]{-1}e}}\right)}{2e} \\
&- \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt[3]{C(d+ex)}}{\sqrt[3]{Cd + \sqrt[3]{-1}e}}\right)}{2e} \\
&- \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt[3]{C(d+ex)}}{\sqrt[3]{Cd - (-1)^{2/3}e}}\right)}{2e} \\
&+ \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt[3]{C(d+ex)}}{\sqrt[3]{Cd + (-1)^{2/3}e}}\right)}{2e}
\end{aligned}$$

output

```
(a+b*arctanh(c*x^3))*ln(e*x+d)/e+1/2*b*ln(e*(1-c^(1/3)*x)/(c^(1/3)*d+e))*ln(e*x+d)/e-1/2*b*ln(-e*(1+c^(1/3)*x)/(c^(1/3)*d-e))*ln(e*x+d)/e+1/2*b*ln(-e*((-1)^(1/3)+c^(1/3)*x)/(c^(1/3)*d-(-1)^(1/3)*e))*ln(e*x+d)/e-1/2*b*ln(-e*((-1)^(2/3)+c^(1/3)*x)/(c^(1/3)*d-(-1)^(2/3)*e))*ln(e*x+d)/e+1/2*b*ln((-1)^(2/3)*e*(1+(-1)^(1/3)*c^(1/3)*x)/(c^(1/3)*d+(-1)^(2/3)*e))*ln(e*x+d)/e-1/2*b*ln((-1)^(1/3)*e*(1+(-1)^(2/3)*c^(1/3)*x)/(c^(1/3)*d+(-1)^(1/3)*e))*ln(e*x+d)/e-1/2*b*polylog(2,c^(1/3)*(e*x+d)/(c^(1/3)*d-e))/e+1/2*b*polylog(2,c^(1/3)*(e*x+d)/(c^(1/3)*d+e))/e+1/2*b*polylog(2,c^(1/3)*(e*x+d)/(c^(1/3)*d-(-1)^(1/3)*e))/e-1/2*b*polylog(2,c^(1/3)*(e*x+d)/(c^(1/3)*d+(-1)^(1/3)*e))/e-1/2*b*polylog(2,c^(1/3)*(e*x+d)/(c^(1/3)*d-(-1)^(2/3)*e))/e+1/2*b*polylog(2,c^(1/3)*(e*x+d)/(c^(1/3)*d+(-1)^(2/3)*e))/e
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.78 (sec) , antiderivative size = 515, normalized size of antiderivative = 0.98

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{d + ex} dx = \frac{a \log(d + ex)}{e} + \frac{b \left(2 \operatorname{arctanh}(cx^3) \log(d + ex) - \log \left(\frac{e(1 - i\sqrt{3} - 2\sqrt[3]{Cx})}{2\sqrt[3]{Cd + e - i\sqrt{3}e}} \right) \log(d + ex) + \log \left(\frac{e(-i + \sqrt{3} - 2i\sqrt[3]{Cx})}{2i\sqrt[3]{Cd + (-i + \sqrt{3})e}} \right) \log(d + \right.}{+ \left. \right)}$$

input

```
Integrate[(a + b*ArcTanh[c*x^3])/(d + e*x),x]
```

output

```
(a*Log[d + e*x])/e + (b*(2*ArcTanh[c*x^3]*Log[d + e*x] - Log[(e*(1 - I*Sqr
t[3] - 2*c^(1/3)*x))/(2*c^(1/3)*d + e - I*Sqrt[3]*e)]*Log[d + e*x] + Log[(
e*(-I + Sqrt[3] - (2*I)*c^(1/3)*x))/((2*I)*c^(1/3)*d + (-I + Sqrt[3])*e)]*
Log[d + e*x] + Log[(e*(I + Sqrt[3] + (2*I)*c^(1/3)*x))/((-2*I)*c^(1/3)*d +
(I + Sqrt[3])*e)]*Log[d + e*x] - Log[-((e*(1 + c^(1/3)*x))/(c^(1/3)*d - e
))]*Log[d + e*x] - Log[-((e*(-1 - I*Sqrt[3] + 2*c^(1/3)*x))/(2*c^(1/3)*d +
e + I*Sqrt[3]*e))]*Log[d + e*x] + Log[d + e*x]*Log[(e - c^(1/3)*e*x)/(c^(
1/3)*d + e)] - PolyLog[2, (c^(1/3)*(d + e*x))/(c^(1/3)*d - e)] + PolyLog[2
, (c^(1/3)*(d + e*x))/(c^(1/3)*d + e)] + PolyLog[2, (2*c^(1/3)*(d + e*x))/
(2*c^(1/3)*d - e - I*Sqrt[3]*e)] - PolyLog[2, (2*c^(1/3)*(d + e*x))/(2*c^(
1/3)*d + e - I*Sqrt[3]*e)] + PolyLog[2, (2*c^(1/3)*(d + e*x))/(2*c^(1/3)*d
- e + I*Sqrt[3]*e)] - PolyLog[2, (2*c^(1/3)*(d + e*x))/(2*c^(1/3)*d + e +
I*Sqrt[3]*e)]))/(2*e)
```

Rubi [A] (verified)

Time = 1.24 (sec) , antiderivative size = 519, normalized size of antiderivative = 0.99, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6482, 2863, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}(cx^3)}{d + ex} dx \\
 & \quad \downarrow \text{6482} \\
 & \frac{\log(d + ex) (a + b \operatorname{arctanh}(cx^3))}{e} - \frac{3bc \int \frac{x^2 \log(d+ex)}{1-c^2x^6} dx}{e} \\
 & \quad \downarrow \text{2863} \\
 & \frac{\log(d + ex) (a + b \operatorname{arctanh}(cx^3))}{e} - \frac{3bc \int \left(\frac{c \log(d+ex)x^2}{2(c-c^2x^3)} + \frac{c \log(d+ex)x^2}{2(c^2x^3+c)} \right) dx}{e} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\frac{\log(d+ex)(a + \operatorname{arctanh}(cx^3))}{e} - 3bc \left(\frac{\operatorname{PolyLog}\left(2, \frac{\sqrt[3]{C(d+ex)}}{\sqrt[3]{Cd-e}}\right)}{6c} - \frac{\operatorname{PolyLog}\left(2, \frac{\sqrt[3]{C(d+ex)}}{\sqrt[3]{Cd+e}}\right)}{6c} - \frac{\operatorname{PolyLog}\left(2, \frac{\sqrt[3]{C(d+ex)}}{\sqrt[3]{Cd-\sqrt{-1}e}}\right)}{6c} + \frac{\operatorname{PolyLog}\left(2, \frac{\sqrt[3]{C(d+ex)}}{\sqrt[3]{Cd+\sqrt{-1}e}}\right)}{6c} + \frac{\operatorname{PolyLog}\left(2, \frac{\sqrt[3]{C(d+ex)}}{\sqrt[3]{Cd-\sqrt{-1}e}}\right)}{6c} + \frac{\operatorname{PolyLog}\left(2, \frac{\sqrt[3]{C(d+ex)}}{\sqrt[3]{Cd+\sqrt{-1}e}}\right)}{6c} \right)$$

input `Int[(a + b*ArcTanh[c*x^3])/(d + e*x),x]`

output `((a + b*ArcTanh[c*x^3])*Log[d + e*x])/e - (3*b*c*(-1/6*(Log[(e*(1 - c^(1/3)*x))/(c^(1/3)*d + e)]*Log[d + e*x])/c + (Log[-((e*(1 + c^(1/3)*x))/(c^(1/3)*d - e)])*Log[d + e*x])/(6*c) - (Log[-((e*((-1)^(1/3) + c^(1/3)*x))/(c^(1/3)*d - (-1)^(1/3)*e)])*Log[d + e*x])/(6*c) + (Log[-((e*((-1)^(2/3) + c^(1/3)*x))/(c^(1/3)*d - (-1)^(2/3)*e)])*Log[d + e*x])/(6*c) - (Log[(-1)^(2/3)*e*(1 + (-1)^(1/3)*c^(1/3)*x)/(c^(1/3)*d + (-1)^(2/3)*e])*Log[d + e*x])/(6*c) + (Log[(-1)^(1/3)*e*(1 + (-1)^(2/3)*c^(1/3)*x)/(c^(1/3)*d + (-1)^(1/3)*e])*Log[d + e*x])/(6*c) + PolyLog[2, (c^(1/3)*(d + e*x))/(c^(1/3)*d - e])/(6*c) - PolyLog[2, (c^(1/3)*(d + e*x))/(c^(1/3)*d + e])/(6*c) - PolyLog[2, (c^(1/3)*(d + e*x))/(c^(1/3)*d - (-1)^(1/3)*e])/(6*c) + PolyLog[2, (c^(1/3)*(d + e*x))/(c^(1/3)*d + (-1)^(1/3)*e])/(6*c) + PolyLog[2, (c^(1/3)*(d + e*x))/(c^(1/3)*d - (-1)^(2/3)*e])/(6*c) - PolyLog[2, (c^(1/3)*(d + e*x))/(c^(1/3)*d + (-1)^(2/3)*e])/(6*c)))/e`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2863 `Int[((a_) + Log[(c_)*((d_) + (e_)*(x_))^(n_)]*(b_))^(p_)*((h_)*(x_))^(m_)*((f_) + (g_)*(x_))^(r_)]^(q_), x_Symbol] := Int[ExpandIntegrand[(a + b*Log[c*(d + e*x)^n])^p, (h*x)^m*(f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n, p, q, r}, x] && IntegerQ[m] && IntegerQ[q]`

rule 6482

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.)/((d_.) + (e_.)*(x_)), x_Symbol
] := Simp[Log[d + e*x]*((a + b*ArcTanh[c*x^n])/e), x] - Simp[b*c*(n/e) In
t[x^(n - 1)*(Log[d + e*x]/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, d,
e, n}, x] && IntegerQ[n]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.33 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.35

method	result
default	$\frac{a \ln(ex+d)}{e} + \frac{b \ln(ex+d) \operatorname{arctanh}(cx^3)}{e} + \frac{b \left(\sum_{-R1=\operatorname{RootOf}(_Z^3 c-3_Z^2 cd+3_Zc d^2-c d^3-e^3)} \left(\ln(ex+d) \ln\left(\frac{-ex+_R1-d}{_R1}\right) \right) \right)}{2e}$
parts	$\frac{a \ln(ex+d)}{e} + \frac{b \ln(ex+d) \operatorname{arctanh}(cx^3)}{e} + \frac{b \left(\sum_{-R1=\operatorname{RootOf}(_Z^3 c-3_Z^2 cd+3_Zc d^2-c d^3-e^3)} \left(\ln(ex+d) \ln\left(\frac{-ex+_R1-d}{_R1}\right) \right) \right)}{2e}$
risch	$\frac{a \ln(ex+d)}{e} - \frac{b \ln(ex+d) \ln(-cx^3+1)}{2e} + \frac{b \left(\sum_{-R1=\operatorname{RootOf}(_Z^3 c-3_Z^2 cd+3_Zc d^2-c d^3-e^3)} \left(\ln(ex+d) \ln\left(\frac{-ex+_R1-d}{_R1}\right) \right) \right)}{2e}$

input

```
int((a+b*arctanh(c*x^3))/(e*x+d),x,method=_RETURNVERBOSE)
```

output

```
a*ln(e*x+d)/e+b*ln(e*x+d)/e*arctanh(c*x^3)+1/2*b/e*sum(ln(e*x+d)*ln((-e*x+
_R1-d)/_R1)+dilog((-e*x+_R1-d)/_R1),_R1=RootOf(_Z^3*c-3*_Z^2*c*d+3*_Z*c*d^
2-c*d^3-e^3))-1/2*b/e*sum(ln(e*x+d)*ln((-e*x+_R1-d)/_R1)+dilog((-e*x+_R1-d
)/_R1),_R1=RootOf(_Z^3*c-3*_Z^2*c*d+3*_Z*c*d^2-c*d^3+e^3))
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{d + ex} dx = \int \frac{b \operatorname{artanh}(cx^3) + a}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x^3))/(e*x+d),x, algorithm="fricas")`

output `integral((b*arctanh(c*x^3) + a)/(e*x + d), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{d + ex} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**3))/(e*x+d),x)`

output `Timed out`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{d + ex} dx = \int \frac{b \operatorname{artanh}(cx^3) + a}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x^3))/(e*x+d),x, algorithm="maxima")`

output `1/2*b*integrate((log(c*x^3 + 1) - log(-c*x^3 + 1))/(e*x + d), x) + a*log(e*x + d)/e`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{d + ex} dx = \int \frac{b \operatorname{artanh}(cx^3) + a}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x^3))/(e*x+d),x, algorithm="giac")`

output `integrate((b*arctanh(c*x^3) + a)/(e*x + d), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{d + ex} dx = \int \frac{a + b \operatorname{atanh}(cx^3)}{d + ex} dx$$

input `int((a + b*atanh(c*x^3))/(d + e*x),x)`

output `int((a + b*atanh(c*x^3))/(d + e*x), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{d + ex} dx = \frac{\left(\int \frac{\operatorname{atanh}(cx^3)}{ex+d} dx\right) be + \log(ex + d) a}{e}$$

input `int((a+b*atanh(c*x^3))/(e*x+d),x)`

output `(int(atanh(c*x**3)/(d + e*x),x)*b*e + log(d + e*x)*a)/e`

3.36 $\int \frac{a+b\operatorname{arctanh}(cx^3)}{(d+ex)^2} dx$

Optimal result	319
Mathematica [A] (verified)	320
Rubi [A] (verified)	321
Maple [A] (verified)	323
Fricas [C] (verification not implemented)	325
Sympy [F(-1)]	325
Maxima [A] (verification not implemented)	326
Giac [A] (verification not implemented)	326
Mupad [B] (verification not implemented)	327
Reduce [F]	328

Optimal result

Integrand size = 18, antiderivative size = 414

$$\int \frac{a + b\operatorname{arctanh}(cx^3)}{(d + ex)^2} dx = -\frac{\sqrt{3}b\sqrt[3]{c} \arctan\left(\frac{1-2\sqrt[3]{cx}}{\sqrt{3}}\right)}{2(c^{2/3}d^2 + \sqrt[3]{cde} + e^2)} - \frac{\sqrt{3}b\sqrt[3]{c}(\sqrt[3]{cd} + e) \arctan\left(\frac{1+2\sqrt[3]{cx}}{\sqrt{3}}\right)}{2(cd^3 + e^3)} - \frac{a + b\operatorname{arctanh}(cx^3)}{e(d + ex)} + \frac{b\sqrt[3]{c}(\sqrt[3]{cd} - e) \log(1 - \sqrt[3]{cx})}{2(cd^3 + e^3)} + \frac{b\sqrt[3]{c}(\sqrt[3]{cd} + e) \log(1 + \sqrt[3]{cx})}{2(cd^3 - e^3)} - \frac{3bcd^2e^2 \log(d + ex)}{c^2d^6 - e^6} - \frac{b\sqrt[3]{c}(\sqrt[3]{cd} + e) \log(1 - \sqrt[3]{cx} + c^{2/3}x^2)}{4(cd^3 - e^3)} - \frac{b\sqrt[3]{c}(\sqrt[3]{cd} - e) \log(1 + \sqrt[3]{cx} + c^{2/3}x^2)}{4(cd^3 + e^3)} - \frac{bcd^2 \log(1 - cx^3)}{2e(cd^3 + e^3)} + \frac{bcd^2 \log(1 + cx^3)}{2e(cd^3 - e^3)}$$

output

```

-1/2*3^(1/2)*b*c^(1/3)*arctan(1/3*(1-2*c^(1/3)*x)*3^(1/2))/(c^(2/3)*d^2+c^(1/3)*d*e+e^2)-3^(1/2)*b*c^(1/3)*(c^(1/3)*d+e)*arctan(1/3*(1+2*c^(1/3)*x)*3^(1/2))/(2*c*d^3+2*e^3)-(a+b*arctanh(c*x^3))/e/(e*x+d)+b*c^(1/3)*(c^(1/3)*d-e)*ln(1-c^(1/3)*x)/(2*c*d^3+2*e^3)+b*c^(1/3)*(c^(1/3)*d+e)*ln(1+c^(1/3)*x)/(2*c*d^3-2*e^3)-3*b*c*d^2*e^2*ln(e*x+d)/(c^2*d^6-e^6)-b*c^(1/3)*(c^(1/3)*d+e)*ln(1-c^(1/3)*x+c^(2/3)*x^2)/(4*c*d^3-4*e^3)-b*c^(1/3)*(c^(1/3)*d-e)*ln(1+c^(1/3)*x+c^(2/3)*x^2)/(4*c*d^3+4*e^3)-1/2*b*c*d^2*ln(-c*x^3+1)/e/(c*d^3+e^3)+1/2*b*c*d^2*ln(c*x^3+1)/e/(c*d^3-e^3)

```

Mathematica [A] (verified)

Time = 0.34 (sec) , antiderivative size = 534, normalized size of antiderivative = 1.29

$$\begin{aligned}
& \int \frac{a + b \operatorname{arctanh}(cx^3)}{(d + ex)^2} dx \\
&= \frac{1}{4} \left(-\frac{4a}{e(d + ex)} + \frac{2\sqrt{3}b\sqrt[3]{c} \operatorname{arctan}\left(\frac{-1+2\sqrt[3]{cx}}{\sqrt{3}}\right)}{c^{2/3}d^2 + \sqrt[3]{cde} + e^2} \right. \\
&\quad - \frac{2\sqrt{3}b\sqrt[3]{c}(\sqrt[3]{cd} + e) \operatorname{arctan}\left(\frac{1+2\sqrt[3]{cx}}{\sqrt{3}}\right)}{cd^3 + e^3} - \frac{4b \operatorname{arctanh}(cx^3)}{e(d + ex)} \\
&\quad + \frac{2b\sqrt[3]{c}(c^{5/3}d^5 - c^{4/3}d^4e + cd^3e^2 + \sqrt[3]{cde}^4 - e^5) \log(1 - \sqrt[3]{cx})}{-c^2d^6e + e^7} \\
&\quad - \frac{2b\sqrt[3]{c}(c^{5/3}d^5 + c^{4/3}d^4e + cd^3e^2 + \sqrt[3]{cde}^4 + e^5) \log(1 + \sqrt[3]{cx})}{-c^2d^6e + e^7} \\
&\quad - \frac{12bcd^2e^2 \log(d + ex)}{c^2d^6 - e^6} \\
&\quad + \frac{b\sqrt[3]{c}(2c^{5/3}d^5 - c^{4/3}d^4e - cd^3e^2 - \sqrt[3]{cde}^4 - e^5) \log(1 - \sqrt[3]{cx} + c^{2/3}x^2)}{c^2d^6e - e^7} \\
&\quad + \frac{b\sqrt[3]{c}(2c^{5/3}d^5 + c^{4/3}d^4e - cd^3e^2 - \sqrt[3]{cde}^4 + e^5) \log(1 + \sqrt[3]{cx} + c^{2/3}x^2)}{-c^2d^6e + e^7} \\
&\quad \left. + \frac{2bcd^2e^2 \log(1 - c^2x^6)}{c^2d^6 - e^6} \right)
\end{aligned}$$

input

```
Integrate[(a + b*ArcTanh[c*x^3])/(d + e*x)^2,x]
```

output

```

((-4*a)/(e*(d + e*x)) + (2*Sqrt[3]*b*c^(1/3)*ArcTan[(-1 + 2*c^(1/3)*x)/Sqr
t[3]])/(c^(2/3)*d^2 + c^(1/3)*d*e + e^2) - (2*Sqrt[3]*b*c^(1/3)*(c^(1/3)*d
+ e)*ArcTan[(1 + 2*c^(1/3)*x)/Sqrt[3]])/(c*d^3 + e^3) - (4*b*ArcTanh[c*x^
3])/(e*(d + e*x)) + (2*b*c^(1/3)*(c^(5/3)*d^5 - c^(4/3)*d^4*e + c*d^3*e^2
+ c^(1/3)*d*e^4 - e^5)*Log[1 - c^(1/3)*x]/(-(c^2*d^6*e) + e^7) - (2*b*c^(
1/3)*(c^(5/3)*d^5 + c^(4/3)*d^4*e + c*d^3*e^2 + c^(1/3)*d*e^4 + e^5)*Log[1
+ c^(1/3)*x])/(-(c^2*d^6*e) + e^7) - (12*b*c*d^2*e^2*Log[d + e*x])/(c^2*d
^6 - e^6) + (b*c^(1/3)*(2*c^(5/3)*d^5 - c^(4/3)*d^4*e - c*d^3*e^2 - c^(1/3
)*d*e^4 - e^5)*Log[1 - c^(1/3)*x + c^(2/3)*x^2])/(c^2*d^6*e - e^7) + (b*c
^(1/3)*(2*c^(5/3)*d^5 + c^(4/3)*d^4*e - c*d^3*e^2 - c^(1/3)*d*e^4 + e^5)*Lo
g[1 + c^(1/3)*x + c^(2/3)*x^2])/(-(c^2*d^6*e) + e^7) + (2*b*c*d^2*e^2*Log[
1 - c^2*x^6])/(c^2*d^6 - e^6))/4

```

Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 410, normalized size of antiderivative = 0.99, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6486, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}(cx^3)}{(d + ex)^2} dx \\
 & \quad \downarrow \text{6486} \\
 & \frac{3bc \int \frac{x^2}{(d+ex)(1-c^2x^6)} dx}{e} - \frac{a + b \operatorname{arctanh}(cx^3)}{e(d + ex)} \\
 & \quad \downarrow \text{7276} \\
 & \frac{3bc \int \left(\frac{d^2 e^4}{(e^3 - cd^3)(cd^3 + e^3)(d + ex)} + \frac{-xe^2 + de - cd^2 x^2}{2(cd^3 + e^3)(cx^3 - 1)} + \frac{-xe^2 + de + cd^2 x^2}{2(cd^3 - e^3)(cx^3 + 1)} \right) dx}{e} - \frac{a + b \operatorname{arctanh}(cx^3)}{e(d + ex)} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$3bc \left(-\frac{e \arctan\left(\frac{2\sqrt[3]{Cx+1}}{\sqrt{3}}\right) \left(\sqrt[3]{Cd+e}\right)}{2\sqrt{3}c^{2/3}(cd^3+e^3)} - \frac{e \arctan\left(\frac{1-2\sqrt[3]{Cx}}{\sqrt{3}}\right)}{2\sqrt{3}c^{2/3}\left(c^{2/3}d^2+\sqrt[3]{Cde+e^2}\right)} - \frac{e\left(\sqrt[3]{Cd+e}\right) \log\left(c^{2/3}x^2-\sqrt[3]{Cx+1}\right)}{12c^{2/3}(cd^3-e^3)} - \frac{e\left(\sqrt[3]{Cd-e}\right) \log\left(c^{2/3}x^2+\sqrt[3]{Cx+1}\right)}{12c^{2/3}(cd^3+e^3)} \right) \\ \frac{a + b \operatorname{arctanh}(cx^3)}{e(d+ex)}$$

input `Int[(a + b*ArcTanh[c*x^3])/(d + e*x)^2, x]`

output `-((a + b*ArcTanh[c*x^3])/(e*(d + e*x))) + (3*b*c*(-1/2*(e*ArcTan[(1 - 2*c^(1/3)*x)/Sqrt[3]])/(Sqrt[3]*c^(2/3)*(c^(2/3)*d^2 + c^(1/3)*d*e + e^2)) - (e*(c^(1/3)*d + e)*ArcTan[(1 + 2*c^(1/3)*x)/Sqrt[3]])/(2*Sqrt[3]*c^(2/3)*(c*d^3 + e^3)) + ((c^(1/3)*d - e)*e*Log[1 - c^(1/3)*x])/(6*c^(2/3)*(c*d^3 + e^3)) + (e*(c^(1/3)*d + e)*Log[1 + c^(1/3)*x])/(6*c^(2/3)*(c*d^3 - e^3)) - (d^2*e^3*Log[d + e*x])/(c^2*d^6 - e^6) - (e*(c^(1/3)*d + e)*Log[1 - c^(1/3)*x + c^(2/3)*x^2])/(12*c^(2/3)*(c*d^3 - e^3)) - ((c^(1/3)*d - e)*e*Log[1 + c^(1/3)*x + c^(2/3)*x^2])/(12*c^(2/3)*(c*d^3 + e^3)) - (d^2*Log[1 - c*x^3])/(6*(c*d^3 + e^3)) + (d^2*Log[1 + c*x^3])/(6*(c*d^3 - e^3)))/e`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6486 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)*((d_) + (e_)*(x_)^(m_)), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*ArcTanh[c*x^n])/(e*(m + 1))), x] - Simp[b*c*(n/(e*(m + 1))) Int[x^(n - 1)*((d + e*x)^(m + 1)/(1 - c^2*x^(2*n)))]], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]`

rule 7276 `Int[(u_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 0.81 (sec) , antiderivative size = 461, normalized size of antiderivative = 1.11

method	result
default	$-\frac{a}{(ex+d)e} + b - \frac{\operatorname{arctanh}(cx^3)}{(ex+d)e} + \frac{de}{3c} \left(\frac{\ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right) - \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{3c\left(\frac{1}{c}\right)^{\frac{2}{3}} - \frac{\ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{6c\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{-2x}{\left(\frac{1}{c}\right)^{\frac{1}{3}} + 1\right)}{\left(\frac{1}{c}\right)^{\frac{1}{3}}}\right)}{3c\left(\frac{1}{c}\right)^{\frac{2}{3}}}\right) - e^2 \frac{\ln\left(\dots\right)}{2cd^3}$
	$-\frac{a}{(ex+d)e} + b - \frac{\operatorname{arctanh}(cx^3)}{(ex+d)e} + \frac{de}{3c} \left(\frac{\ln\left(x - \left(\frac{1}{c}\right)^{\frac{1}{3}}\right) - \ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{3c\left(\frac{1}{c}\right)^{\frac{2}{3}} - \frac{\ln\left(x^2 + \left(\frac{1}{c}\right)^{\frac{1}{3}}x + \left(\frac{1}{c}\right)^{\frac{2}{3}}\right)}{6c\left(\frac{1}{c}\right)^{\frac{2}{3}}} - \frac{\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3}\left(\frac{-2x}{\left(\frac{1}{c}\right)^{\frac{1}{3}} + 1\right)}{\left(\frac{1}{c}\right)^{\frac{1}{3}}}\right)}{3c\left(\frac{1}{c}\right)^{\frac{2}{3}}}\right) - e^2 \frac{\ln\left(\dots\right)}{2cd^3}$

input `int((a+b*arctanh(c*x^3))/(e*x+d)^2,x,method=_RETURNVERBOSE)`

output `-a/(e*x+d)/e+b*(-1/(e*x+d)/e*arctanh(c*x^3)+3/e*c*((d*e*(1/3/c/(1/c)^(2/3))*ln(x-(1/c)^(1/3))-1/6/c/(1/c)^(2/3)*ln(x^2+(1/c)^(1/3)*x+(1/c)^(2/3))-1/3/c/(1/c)^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x+1)))-e^2*(1/3/c/(1/c)^(1/3)*ln(x-(1/c)^(1/3))-1/6/c/(1/c)^(1/3)*ln(x^2+(1/c)^(1/3)*x+(1/c)^(2/3))+1/3*3^(1/2)/c/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x+1)))-1/3*d^2*ln(c*x^3-1)/(2*c*d^3+2*e^3)-d^2*e^3/(c*d^3+e^3)/(c*d^3-e^3)*ln(e*x+d)+(d*e*(1/3/c/(1/c)^(2/3)*ln(x+(1/c)^(1/3))-1/6/c/(1/c)^(2/3)*ln(x^2-(1/c)^(1/3)*x+(1/c)^(2/3))+1/3/c/(1/c)^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x-1)))-e^2*(-1/3/c/(1/c)^(1/3)*ln(x+(1/c)^(1/3))+1/6/c/(1/c)^(1/3)*ln(x^2-(1/c)^(1/3)*x+(1/c)^(2/3))+1/3*3^(1/2)/c/(1/c)^(1/3)*arctan(1/3*3^(1/2)*(2/(1/c)^(1/3)*x-1)))+1/3*d^2*ln(c*x^3+1)/(2*c*d^3-2*e^3))`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 137.99 (sec) , antiderivative size = 12984, normalized size of antiderivative = 31.36

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{(d + ex)^2} dx = \text{Too large to display}$$

input `integrate((a+b*arctanh(c*x^3))/(e*x+d)^2,x, algorithm="fricas")`

output Too large to include

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{(d + ex)^2} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**3))/(e*x+d)**2,x)`

output Timed out

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 394, normalized size of antiderivative = 0.95

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{(d + ex)^2} dx =$$

$$-\frac{1}{4} \left(\left(\frac{12 d^2 e^2 \log(ex + d)}{c^2 d^6 - e^6} + \frac{2 \sqrt{3} (cde + c^{\frac{2}{3}} e^2) \arctan\left(\frac{\sqrt{3}(2c^{\frac{2}{3}}x + c^{\frac{1}{3}})}{3c^{\frac{1}{3}}}\right)}{(c^2 d^3 e + ce^4)c^{\frac{1}{3}}} \right) - \frac{2 \sqrt{3} (cde - c^{\frac{2}{3}} e^2) \arctan\left(\frac{\sqrt{3}(2c^{\frac{2}{3}}x + c^{\frac{1}{3}})}{3c^{\frac{1}{3}}}\right)}{(c^2 d^3 e - ce^4)c^{\frac{1}{3}}} \right) - \frac{a}{e^2 x + de}$$

input `integrate((a+b*arctanh(c*x^3))/(e*x+d)^2,x, algorithm="maxima")`

output

```
-1/4*((12*d^2*e^2*log(e*x + d)/(c^2*d^6 - e^6) + 2*sqrt(3)*(c*d*e + c^(2/3)*e^2)*arctan(1/3*sqrt(3)*(2*c^(2/3)*x + c^(1/3))/c^(1/3))/((c^2*d^3*e + c*e^4)*c^(1/3)) - 2*sqrt(3)*(c*d*e - c^(2/3)*e^2)*arctan(1/3*sqrt(3)*(2*c^(2/3)*x - c^(1/3))/c^(1/3))/((c^2*d^3*e - c*e^4)*c^(1/3)) + (2*c*d^2 + c^(2/3)*d*e - c^(1/3)*e^2)*log(c^(2/3)*x^2 + c^(1/3)*x + 1)/(c^2*d^3*e + c*e^4) - (2*c*d^2 - c^(2/3)*d*e - c^(1/3)*e^2)*log(c^(2/3)*x^2 - c^(1/3)*x + 1)/(c^2*d^3*e - c*e^4) - 2*(c*d^2 + c^(2/3)*d*e + c^(1/3)*e^2)*log((c^(1/3)*x + 1)/c^(1/3))/(c^2*d^3*e - c*e^4) + 2*(c*d^2 - c^(2/3)*d*e + c^(1/3)*e^2)*log((c^(1/3)*x - 1)/c^(1/3))/(c^2*d^3*e + c*e^4))*c + 4*arctanh(c*x^3)/(e^2*x + d*e))*b - a/(e^2*x + d*e)
```

Giac [A] (verification not implemented)

Time = 15.72 (sec) , antiderivative size = 554, normalized size of antiderivative = 1.34

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{(d + ex)^2} dx = \text{Too large to display}$$

input `integrate((a+b*arctanh(c*x^3))/(e*x+d)^2,x, algorithm="giac")`

output

```

-3*b*c*d^2*e^2*log(e*x + d)/(c^2*d^6 - e^6) + 1/2*b*c*d^2*log(abs(c*x^3 +
1))/(c*d^3*e - e^4) - 1/2*b*c*d^2*log(abs(-c*x^3 + 1))/(c*d^3*e + e^4) - 1
/2*sqrt(3)*b*c*abs(c)^(2/3)*arctan(1/3*sqrt(3)*c^(1/3)*(2*x + 1/c^(1/3)))/
(c^2*d^2 - c*d*e*abs(c)^(2/3) + e^2*abs(c)^(4/3)) - 1/2*sqrt(3)*b*c*arctan
(1/3*sqrt(3)*(2*x + (-1/c)^(1/3))/(-1/c)^(1/3))/(c*d*e + (-c^2)^(2/3)*d^2
- (-c^2)^(1/3)*e^2) + 1/2*(b*c^3*d^3*e^3*(-1/c)^(1/3) - b*c^3*d^4*e^2 - b*
c^2*e^6*(-1/c)^(1/3) + b*c^2*d*e^5)*(-1/c)^(1/3)*log(abs(x - (-1/c)^(1/3))
)/(c^3*d^6*e^2 - 2*c^2*d^3*e^5 + c*e^8) + 1/4*((-c^2)^(1/3)*b*c*d - (-c^2)
^(2/3)*b*e)*log(x^2 + x*(-1/c)^(1/3) + (-1/c)^(2/3))/(c^2*d^3 - c*e^3) - 1
/4*(b*c*d*abs(c)^(2/3) - b*e*abs(c)^(4/3))*log(x^2 + x/c^(1/3) + 1/c^(2/3)
)/(c^2*d^3 + c*e^3) - 1/2*b*log(-(c*x^3 + 1)/(c*x^3 - 1))/(e^2*x + d*e) +
1/2*(b*c^3*d^4*e^2 - b*c^(8/3)*d^3*e^3 + b*c^2*d*e^5 - b*c^(5/3)*e^6)*log(
abs(x - 1/c^(1/3)))/((c^3*d^6*e^2 + 2*c^2*d^3*e^5 + c*e^8)*c^(1/3)) - a/(e
^2*x + d*e)

```

Mupad [B] (verification not implemented)

Time = 3.92 (sec) , antiderivative size = 2638, normalized size of antiderivative = 6.37

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{(d + ex)^2} dx = \text{Too large to display}$$

input

```
int((a + b*atanh(c*x^3))/(d + e*x)^2,x)
```


output

```

symsum(log(-(729*b^6*c^14*d*e^2 + 54432*root(8*c*d^3*e^3*z^3 - 8*e^6*z^3 -
12*b*c*d^2*e^2*z^2 + 6*b^2*c*d*e*z - b^3*c, z, k)^6*c^12*e^15*x + 729*b^6
*c^14*e^3*x + 31104*root(8*c*d^3*e^3*z^3 - 8*e^6*z^3 - 12*b*c*d^2*e^2*z^2
+ 6*b^2*c*d*e*z - b^3*c, z, k)^6*c^14*d^7*e^8 + 243*root(8*c*d^3*e^3*z^3 -
8*e^6*z^3 - 12*b*c*d^2*e^2*z^2 + 6*b^2*c*d*e*z - b^3*c, z, k)*b^5*c^15*d^
5 + 62208*root(8*c*d^3*e^3*z^3 - 8*e^6*z^3 - 12*b*c*d^2*e^2*z^2 + 6*b^2*c*
d*e*z - b^3*c, z, k)^6*c^12*d*e^14 - 5832*root(8*c*d^3*e^3*z^3 - 8*e^6*z^3
- 12*b*c*d^2*e^2*z^2 + 6*b^2*c*d*e*z - b^3*c, z, k)^2*b^4*c^14*d^3*e^4 -
1944*root(8*c*d^3*e^3*z^3 - 8*e^6*z^3 - 12*b*c*d^2*e^2*z^2 + 6*b^2*c*d*e*z
- b^3*c, z, k)^3*b^3*c^15*d^7*e^2 + 15552*root(8*c*d^3*e^3*z^3 - 8*e^6*z^
3 - 12*b*c*d^2*e^2*z^2 + 6*b^2*c*d*e*z - b^3*c, z, k)^4*b^2*c^14*d^5*e^6 +
10692*root(8*c*d^3*e^3*z^3 - 8*e^6*z^3 - 12*b*c*d^2*e^2*z^2 + 6*b^2*c*d*e
*z - b^3*c, z, k)^3*b^3*c^13*d*e^8 + 101088*root(8*c*d^3*e^3*z^3 - 8*e^6*z
^3 - 12*b*c*d^2*e^2*z^2 + 6*b^2*c*d*e*z - b^3*c, z, k)^5*b*c^13*d^3*e^10 +
3888*root(8*c*d^3*e^3*z^3 - 8*e^6*z^3 - 12*b*c*d^2*e^2*z^2 + 6*b^2*c*d*e*
z - b^3*c, z, k)^5*b*c^15*d^9*e^4 + 12636*root(8*c*d^3*e^3*z^3 - 8*e^6*z^3
- 12*b*c*d^2*e^2*z^2 + 6*b^2*c*d*e*z - b^3*c, z, k)^3*b^3*c^13*e^9*x + 38
880*root(8*c*d^3*e^3*z^3 - 8*e^6*z^3 - 12*b*c*d^2*e^2*z^2 + 6*b^2*c*d*e*z
- b^3*c, z, k)^6*c^14*d^6*e^9*x + 116640*root(8*c*d^3*e^3*z^3 - 8*e^6*z^3
- 12*b*c*d^2*e^2*z^2 + 6*b^2*c*d*e*z - b^3*c, z, k)^5*b*c^13*d^2*e^11*x...

```

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(cx^3)}{(d + ex)^2} dx = \int \frac{\operatorname{atanh}(cx^3) b + a}{(ex + d)^2} dx$$

input

```
int((a+b*atanh(c*x^3))/(e*x+d)^2,x)
```

output

```
int((a+b*atanh(c*x^3))/(e*x+d)^2,x)
```

3.37 $\int \frac{x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} dx$

Optimal result	329
Mathematica [A] (verified)	330
Rubi [A] (verified)	330
Maple [A] (verified)	336
Fricas [F]	336
Sympy [F]	337
Maxima [A] (verification not implemented)	337
Giac [F]	338
Mupad [F(-1)]	338
Reduce [F]	338

Optimal result

Integrand size = 26, antiderivative size = 195

$$\int \frac{x^3 (a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} dx = -\frac{11b\sqrt{x}}{6c^7} - \frac{5bx^{3/2}}{18c^5} - \frac{bx^{5/2}}{15c^3} + \frac{11b \operatorname{arctanh}(c\sqrt{x})}{6c^8}$$

$$- \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{c^6} - \frac{x^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{2c^4}$$

$$- \frac{x^3(a + b \operatorname{arctanh}(c\sqrt{x}))}{3c^2} - \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{bc^8}$$

$$+ \frac{2(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1-c\sqrt{x}}\right)}{c^8}$$

$$+ \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-c\sqrt{x}}\right)}{c^8}$$

output

```
-11/6*b*x^(1/2)/c^7-5/18*b*x^(3/2)/c^5-1/15*b*x^(5/2)/c^3+11/6*b*arctanh(c
*x^(1/2))/c^8-x*(a+b*arctanh(c*x^(1/2)))/c^6-1/2*x^2*(a+b*arctanh(c*x^(1/2)
))/c^4-1/3*x^3*(a+b*arctanh(c*x^(1/2)))/c^2-(a+b*arctanh(c*x^(1/2)))^2/b/
c^8+2*(a+b*arctanh(c*x^(1/2)))*ln(2/(1-c*x^(1/2)))/c^8+b*polylog(2,1-2/(1-
c*x^(1/2)))/c^8
```

Mathematica [A] (verified)

Time = 0.48 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.82

$$\int \frac{x^3(a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2x} dx = \frac{165bc\sqrt{x} + 90ac^2x + 25bc^3x^{3/2} + 45ac^4x^2 + 6bc^5x^{5/2} + 30ac^6x^3 - 90\operatorname{barctanh}(c\sqrt{x})^2 + 15\operatorname{barctanh}(c\sqrt{x})}{c^8}$$

input

```
Integrate[(x^3*(a + b*ArcTanh[c*Sqrt[x]]))/(1 - c^2*x),x]
```

output

```
-1/90*(165*b*c*Sqrt[x] + 90*a*c^2*x + 25*b*c^3*x^(3/2) + 45*a*c^4*x^2 + 6*
b*c^5*x^(5/2) + 30*a*c^6*x^3 - 90*b*ArcTanh[c*Sqrt[x]]^2 + 15*b*ArcTanh[c*
Sqrt[x]]*(-11 + 6*c^2*x + 3*c^4*x^2 + 2*c^6*x^3 - 12*Log[1 + E^(-2*ArcTanh
[c*Sqrt[x]]))]) + 90*a*Log[1 - c^2*x] + 90*b*PolyLog[2, -E^(-2*ArcTanh[c*Sq
rt[x]])))/c^8
```

Rubi [A] (verified)

Time = 3.17 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.54, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.654$, Rules used = {7267, 6542, 6452, 254, 2009, 6542, 6452, 254, 2009, 6542, 6452, 262, 219, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2x} dx$$

↓ 7267

$$2 \int \frac{x^{7/2}(a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2x} d\sqrt{x}$$

↓ 6542

$$2 \left(\frac{\int \frac{x^{5/2}(a + \operatorname{barctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\int x^{5/2}(a + \operatorname{barctanh}(c\sqrt{x})) d\sqrt{x}}{c^2} \right)$$

↓ 6452

$$2 \left(\frac{\int \frac{x^{5/2}(a + \operatorname{barctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{6}x^3(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{6}bc \int \frac{x^3}{1-c^2x} d\sqrt{x}}{c^2} \right)$$

↓ 254

$$2 \left(\frac{\int \frac{x^{5/2}(a + \operatorname{barctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{6}x^3(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{6}bc \int \left(-\frac{x^2}{c^2} - \frac{x}{c^4} + \frac{1}{c^6(1-c^2x)} - \frac{1}{c^6} \right) d\sqrt{x}}{c^2} \right)$$

↓ 2009

$$2 \left(\frac{\int \frac{x^{5/2}(a + \operatorname{barctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{6}x^3(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{6}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^7} - \frac{\sqrt{x}}{c^6} - \frac{x^{3/2}}{3c^4} - \frac{x^{5/2}}{5c^2} \right)}{c^2} \right)$$

↓ 6542

$$2 \left(\frac{\frac{\int \frac{x^{3/2}(a + \operatorname{barctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\int x^{3/2}(a + \operatorname{barctanh}(c\sqrt{x})) d\sqrt{x}}{c^2}}{c^2} - \frac{\frac{1}{6}x^3(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{6}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^7} - \frac{\sqrt{x}}{c^6} - \frac{x^{3/2}}{3c^4} - \frac{x^{5/2}}{5c^2} \right)}{c^2}}{c^2} \right)$$

↓ 6452

$$2 \left(\frac{\frac{\int \frac{x^{3/2}(a + \operatorname{barctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{4}x^2(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{4}bc \int \frac{x^2}{1-c^2x} d\sqrt{x}}{c^2}}{c^2} - \frac{\frac{1}{6}x^3(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{6}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^7} - \frac{\sqrt{x}}{c^6} - \frac{x^{3/2}}{3c^4} - \frac{x^{5/2}}{5c^2} \right)}{c^2}}{c^2} \right)$$

↓ 254

$$2 \left(\frac{\int \frac{x^{3/2}(a+b\operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{4}x^2(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \int \left(-\frac{x}{c^2} + \frac{1}{c^4(1-c^2x)} - \frac{1}{c^4}\right) d\sqrt{x}}{c^2}}{c^2} - \frac{\frac{1}{6}x^3(a+b\operatorname{arctanh}(c\sqrt{x}))}{c^2} \right)$$

↓ 2009

$$2 \left(\frac{\int \frac{x^{3/2}(a+b\operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{4}x^2(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^5} - \frac{\sqrt{x}}{c^4} - \frac{x^{3/2}}{3c^2}\right)}{c^2}}{c^2} - \frac{\frac{1}{6}x^3(a+b\operatorname{arctanh}(c\sqrt{x}))}{c^2} \right)$$

↓ 6542

$$2 \left(\frac{\frac{\int \frac{\sqrt{x}(a+b\operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\int \sqrt{x}(a+b\operatorname{arctanh}(c\sqrt{x})) d\sqrt{x}}{c^2}}{c^2} - \frac{\frac{1}{4}x^2(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^5} - \frac{\sqrt{x}}{c^4} - \frac{x^{3/2}}{3c^2}\right)}{c^2}}{c^2} - \frac{\frac{1}{6}x^3(a+b\operatorname{arctanh}(c\sqrt{x}))}{c^2} \right)$$

↓ 6452

$$2 \left(\frac{\frac{\int \frac{\sqrt{x}(a+b\operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \int \frac{x}{1-c^2x} d\sqrt{x}}{c^2}}{c^2} - \frac{\frac{1}{4}x^2(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^5} - \frac{\sqrt{x}}{c^4}\right)}{c^2}}{c^2} \right)$$

↓ 262

$$2 \left(\frac{\frac{\int \frac{\sqrt{x}(a+b\operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\int \frac{1}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\sqrt{x}}{c^2}\right)}{c^2}}{c^2} - \frac{\frac{1}{4}x^2(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^5}\right)}{c^2}}{c^2} \right)$$

↓ 219

$$2 \left(\frac{\int \frac{\sqrt{x}(a+b\operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2}\right)}{c^2}}{c^2}}{c^2} - \frac{\frac{1}{4}x^2(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2}\right)}{c^2}}{c^2} \right)$$

↓ 6546

$$2 \left(\frac{\frac{\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{1-c\sqrt{x}} d\sqrt{x}}{c} - \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2}\right)}{c^2}}{c^2} - \frac{\frac{1}{4}x^2(a+b\operatorname{arctanh}(c\sqrt{x}))}{c^2}}{c^2} \right)$$

↓ 6470

$$2 \left(\frac{\frac{\frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)(a+b\operatorname{arctanh}(c\sqrt{x}))}{c} - b\int \frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)}{1-c^2x} d\sqrt{x}}{c} - \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2}\right)}{c^2}}{c^2} \right)$$

↓ 2849

$$2 \left(\frac{\frac{b\int \frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)}{1-\frac{2}{1-c\sqrt{x}}} d\frac{1}{1-c\sqrt{x}} + \frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)(a+b\operatorname{arctanh}(c\sqrt{x}))}{c}}{c^2} - \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2}\right)}{c^2}}{c^2} \right)$$

↓ 2752

$$2 \left(\frac{\frac{\frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)(a+b\operatorname{arctanh}(c\sqrt{x}))}{c} + \frac{b\operatorname{PolyLog}\left(2,1-\frac{2}{1-c\sqrt{x}}\right)}{2c}}{c^2} - \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2}\right)}{c^2}}{c^2} \right)$$

input `Int[(x^3*(a + b*ArcTanh[c*Sqrt[x]]))/(1 - c^2*x),x]`

output `2*(-(((x^3*(a + b*ArcTanh[c*Sqrt[x]]))/6 - (b*c*(-(Sqrt[x]/c^6) - x^(3/2)/(3*c^4) - x^(5/2)/(5*c^2) + ArcTanh[c*Sqrt[x]]/c^7))/6)/c^2) + (-(((x^2*(a + b*ArcTanh[c*Sqrt[x]]))/4 - (b*c*(-(Sqrt[x]/c^4) - x^(3/2)/(3*c^2) + ArcTanh[c*Sqrt[x]]/c^5))/4)/c^2) + (-(((x*(a + b*ArcTanh[c*Sqrt[x]]))/2 - (b*c*(-(Sqrt[x]/c^2) + ArcTanh[c*Sqrt[x]]/c^3))/2)/c^2) + (-1/2*(a + b*ArcTanh[c*Sqrt[x]])^2/(b*c^2) + (((a + b*ArcTanh[c*Sqrt[x]])*Log[2/(1 - c*Sqrt[x]])]/c + (b*PolyLog[2, 1 - 2/(1 - c*Sqrt[x]])/(2*c))/c)/c^2)/c^2)`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 254 `Int[(x_)^(m_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^2, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 3]`

rule 262 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 $\text{Int}[\text{Log}[(c_)/(d_ + (e_)(x_))]/((f_ + (g_)(x_)^2), x_Symbol] \text{ :> } \text{Simp}[-e/g \text{ Subst}[\text{Int}[\text{Log}[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] \text{ /; } \text{FreeQ}[\{c, d, e, f, g\}, x] \ \&\& \ \text{EqQ}[c, 2*d] \ \&\& \ \text{EqQ}[e^2*f + d^2*g, 0]$

rule 6452 $\text{Int}[(a_ + \text{ArcTanh}[(c_)(x_)]*(b_))^{(p_)}(x_)^{(m_)}, x_Symbol] \text{ :> } \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)})/(1 - c^2*x^{(2*n)}), x], x] \text{ /; } \text{FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 6470 $\text{Int}[(a_ + \text{ArcTanh}[(c_)(x_)]*(b_))^{(p_)}((d_ + (e_)(x_)), x_Symbol] \text{ :> } \text{Simp}[(-a + b*\text{ArcTanh}[c*x])^p*(\text{Log}[2/(1 + e*(x/d))]/e), x] + \text{Simp}[b*c*(p/e) \text{ Int}[(a + b*\text{ArcTanh}[c*x])^{(p-1)}*(\text{Log}[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] \text{ /; } \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d^2 - e^2, 0]$

rule 6542 $\text{Int}[(a_ + \text{ArcTanh}[(c_)(x_)]*(b_))^{(p_)}((f_)(x_))^{(m_)}((d_ + (e_)(x_)^2), x_Symbol] \text{ :> } \text{Simp}[f^2/e \text{ Int}[(f*x)^{(m-2)}*(a + b*\text{ArcTanh}[c*x])^p, x], x] - \text{Simp}[d*(f^2/e) \text{ Int}[(f*x)^{(m-2)}*(a + b*\text{ArcTanh}[c*x])^p/(d + e*x^2), x], x] \text{ /; } \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{GtQ}[m, 1]$

rule 6546 $\text{Int}[(a_ + \text{ArcTanh}[(c_)(x_)]*(b_))^{(p_)}(x_)/((d_ + (e_)(x_)^2), x_Symbol] \text{ :> } \text{Simp}[(a + b*\text{ArcTanh}[c*x])^{(p+1)}/(b*e*(p+1)), x] + \text{Simp}[1/(c*d) \text{ Int}[(a + b*\text{ArcTanh}[c*x])^p/(1 - c*x), x], x] \text{ /; } \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{IGtQ}[p, 0]$

rule 7267 $\text{Int}[u_, x_Symbol] \text{ :> } \text{With}[\{lst = \text{SubstForFractionalPowerOfLinear}[u, x]\}, \text{Simp}[lst[[2]]*lst[[4]] \text{ Subst}[\text{Int}[lst[[1]], x], x, lst[[3]]^{(1/lst[[2]])}], x] \text{ /; } \text{!FalseQ}[lst] \ \&\& \ \text{SubstForFractionalPowerQ}[u, lst[[3]], x]$

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.27

method	result
parts	$-\frac{ax^3}{3c^2} - \frac{ax^2}{2c^4} - \frac{ax}{c^6} - \frac{a \ln(c^2x-1)}{c^8} - 2b \left(\frac{\operatorname{arctanh}(c\sqrt{x})c^6x^3}{6} + \frac{\operatorname{arctanh}(c\sqrt{x})c^4x^2}{4} + \frac{\operatorname{arctanh}(c\sqrt{x})c^2x}{2} + \frac{\operatorname{arctanh}(c\sqrt{x})}{2} \right)$
derivativedivides	$2 \left(a \left(\frac{c^6x^3}{6} + \frac{c^4x^2}{4} + \frac{c^2x}{2} + \frac{\ln(c\sqrt{x}-1)}{2} + \frac{\ln(1+c\sqrt{x})}{2} \right) + b \left(\frac{\operatorname{arctanh}(c\sqrt{x})c^6x^3}{6} + \frac{\operatorname{arctanh}(c\sqrt{x})c^4x^2}{4} + \frac{\operatorname{arctanh}(c\sqrt{x})c^2x}{2} + \frac{\operatorname{arctanh}(c\sqrt{x})}{2} \right) \right)$
default	$2 \left(a \left(\frac{c^6x^3}{6} + \frac{c^4x^2}{4} + \frac{c^2x}{2} + \frac{\ln(c\sqrt{x}-1)}{2} + \frac{\ln(1+c\sqrt{x})}{2} \right) + b \left(\frac{\operatorname{arctanh}(c\sqrt{x})c^6x^3}{6} + \frac{\operatorname{arctanh}(c\sqrt{x})c^4x^2}{4} + \frac{\operatorname{arctanh}(c\sqrt{x})c^2x}{2} + \frac{\operatorname{arctanh}(c\sqrt{x})}{2} \right) \right)$

```
input int(x^3*(a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x,method=_RETURNVERBOSE)
```

```
output -1/3*a/c^2*x^3-1/2*a/c^4*x^2-a/c^6*x-a/c^8*ln(c^2*x-1)-2*b/c^8*(1/6*arctanh(c*x^(1/2))*c^6*x^3+1/4*arctanh(c*x^(1/2))*c^4*x^2+1/2*arctanh(c*x^(1/2))*c^2*x+1/2*arctanh(c*x^(1/2))*ln(c*x^(1/2)-1)+1/2*arctanh(c*x^(1/2))*ln(1+c*x^(1/2))-1/2*dilog(1/2*c*x^(1/2)+1/2)-1/4*ln(c*x^(1/2)-1)*ln(1/2*c*x^(1/2)+1/2)+1/8*ln(c*x^(1/2)-1)^2-1/8*ln(1+c*x^(1/2))^2+1/4*(ln(1+c*x^(1/2))-ln(1/2*c*x^(1/2)+1/2))*ln(-1/2*c*x^(1/2)+1/2)+1/30*c^5*x^(5/2)+5/36*c^3*x^(3/2)+11/12*c*x^(1/2)+11/24*ln(c*x^(1/2)-1)-11/24*ln(1+c*x^(1/2)))
```

Fricas [F]

$$\int \frac{x^3(a + b\operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx = \int -\frac{(b\operatorname{arctanh}(c\sqrt{x}) + a)x^3}{c^2x - 1} dx$$

```
input integrate(x^3*(a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x, algorithm="fricas")
```

```
output integral(-(b*x^3*arctanh(c*sqrt(x)) + a*x^3)/(c^2*x - 1), x)
```

Sympy [F]

$$\int \frac{x^3(a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2x} dx = - \int \frac{ax^3}{c^2x - 1} dx - \int \frac{bx^3 \operatorname{atanh}(c\sqrt{x})}{c^2x - 1} dx$$

input `integrate(x**3*(a+b*atanh(c*x**(1/2)))/(-c**2*x+1),x)`

output `-Integral(a*x**3/(c**2*x - 1), x) - Integral(b*x**3*atanh(c*sqrt(x))/(c**2*x - 1), x)`

Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.26

$$\int \frac{x^3(a + \operatorname{barctanh}(c\sqrt{x}))}{1 - c^2x} dx = -\frac{1}{6}a \left(\frac{2c^4x^3 + 3c^2x^2 + 6x}{c^6} + \frac{6 \log(c^2x - 1)}{c^8} \right) - \frac{(\log(c\sqrt{x} + 1) \log(-\frac{1}{2}c\sqrt{x} + \frac{1}{2}) + \operatorname{Li}_2(\frac{1}{2}c\sqrt{x} + \frac{1}{2}))b}{c^8} + \frac{11b \log(c\sqrt{x} + 1)}{12c^8} - \frac{11b \log(c\sqrt{x} - 1)}{12c^8} - \frac{12bc^5x^{\frac{5}{2}} + 50bc^3x^{\frac{3}{2}} + 45b \log(c\sqrt{x} + 1)^2 - 45b \log(-c\sqrt{x} + 1)^2 + 330bc\sqrt{x} + 15(2bc^6x^3 + 3bc^4x^2)}{180c^8}$$

input `integrate(x^3*(a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x, algorithm="maxima")`

output `-1/6*a*((2*c^4*x^3 + 3*c^2*x^2 + 6*x)/c^6 + 6*log(c^2*x - 1)/c^8) - (log(c*sqrt(x) + 1)*log(-1/2*c*sqrt(x) + 1/2) + dilog(1/2*c*sqrt(x) + 1/2))*b/c^8 + 11/12*b*log(c*sqrt(x) + 1)/c^8 - 11/12*b*log(c*sqrt(x) - 1)/c^8 - 1/180*(12*b*c^5*x^(5/2) + 50*b*c^3*x^(3/2) + 45*b*log(c*sqrt(x) + 1)^2 - 45*b*log(-c*sqrt(x) + 1)^2 + 330*b*c*sqrt(x) + 15*(2*b*c^6*x^3 + 3*b*c^4*x^2 + 6*b*c^2*x)*log(c*sqrt(x) + 1) - 15*(2*b*c^6*x^3 + 3*b*c^4*x^2 + 6*b*c^2*x + 6*b*log(c*sqrt(x) + 1))*log(-c*sqrt(x) + 1))/c^8`

Giac [F]

$$\int \frac{x^3(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx = \int -\frac{(b \operatorname{artanh}(c\sqrt{x}) + a)x^3}{c^2x - 1} dx$$

input `integrate(x^3*(a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x, algorithm="giac")`

output `integrate(-(b*arctanh(c*sqrt(x)) + a)*x^3/(c^2*x - 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x^3(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx = \int -\frac{x^3(a + b \operatorname{atanh}(c\sqrt{x}))}{c^2x - 1} dx$$

input `int(-(x^3*(a + b*atanh(c*x^(1/2))))/(c^2*x - 1),x)`

output `int(-(x^3*(a + b*atanh(c*x^(1/2))))/(c^2*x - 1), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{x^3(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx \\ &= \frac{-6 \left(\int \frac{\operatorname{atanh}(\sqrt{x}c)x^3}{c^2x-1} dx \right) b c^8 - 6 \log(c^2x - 1) a - 2a c^6 x^3 - 3a c^4 x^2 - 6a c^2 x}{6c^8} \end{aligned}$$

input `int(x^3*(a+b*atanh(c*x^(1/2)))/(-c^2*x+1),x)`

output `(- 6*int((atanh(sqrt(x)*c)*x**3)/(c**2*x - 1),x)*b*c**8 - 6*log(c**2*x - 1)*a - 2*a*c**6*x**3 - 3*a*c**4*x**2 - 6*a*c**2*x)/(6*c**8)`

$$3.38 \quad \int \frac{x^2 \left(a + b \operatorname{arctanh}(c\sqrt{x}) \right)}{1 - c^2 x} dx$$

Optimal result	339
Mathematica [A] (verified)	340
Rubi [A] (verified)	340
Maple [A] (verified)	345
Fricas [F]	345
Sympy [F]	346
Maxima [A] (verification not implemented)	346
Giac [F]	347
Mupad [F(-1)]	347
Reduce [F]	347

Optimal result

Integrand size = 26, antiderivative size = 160

$$\int \frac{x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} dx = -\frac{3b\sqrt{x}}{2c^5} - \frac{bx^{3/2}}{6c^3} + \frac{3b \operatorname{arctanh}(c\sqrt{x})}{2c^6} - \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{c^4} - \frac{x^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{2c^2} - \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{bc^6} + \frac{2(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1 - c\sqrt{x}}\right)}{c^6} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - c\sqrt{x}}\right)}{c^6}$$

output

```
-3/2*b*x^(1/2)/c^5-1/6*b*x^(3/2)/c^3+3/2*b*arctanh(c*x^(1/2))/c^6-x*(a+b*arctanh(c*x^(1/2)))/c^4-1/2*x^2*(a+b*arctanh(c*x^(1/2)))/c^2-(a+b*arctanh(c*x^(1/2)))^2/b/c^6+2*(a+b*arctanh(c*x^(1/2)))*ln(2/(1-c*x^(1/2)))/c^6+b*polylog(2,1-2/(1-c*x^(1/2)))/c^6
```

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.81

$$\int \frac{x^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx = \frac{9bc\sqrt{x} + 6ac^2x + bc^3x^{3/2} + 3ac^4x^2 - 6b \operatorname{arctanh}(c\sqrt{x})^2 + 3b \operatorname{arctanh}(c\sqrt{x}) \left(-3 + 2c^2x + c^4x^2 - 4 \log\right)}{6c^6}$$

input

```
Integrate[(x^2*(a + b*ArcTanh[c*Sqrt[x]]))/(1 - c^2*x),x]
```

output

```
-1/6*(9*b*c*Sqrt[x] + 6*a*c^2*x + b*c^3*x^(3/2) + 3*a*c^4*x^2 - 6*b*ArcTan
h[c*Sqrt[x]]^2 + 3*b*ArcTanh[c*Sqrt[x]]*(-3 + 2*c^2*x + c^4*x^2 - 4*Log[1
+ E^(-2*ArcTanh[c*Sqrt[x]])]) + 6*a*Log[1 - c^2*x] + 6*b*PolyLog[2, -E^(-2
*ArcTanh[c*Sqrt[x]])])/c^6
```

Rubi [A] (verified)

Time = 1.39 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.36, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {7267, 6542, 6452, 254, 2009, 6542, 6452, 262, 219, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx \\ & \quad \downarrow \text{7267} \\ & 2 \int \frac{x^{5/2}(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} d\sqrt{x} \\ & \quad \downarrow \text{6542} \\ & 2 \left(\frac{\int \frac{x^{3/2}(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} d\sqrt{x}}{c^2} - \frac{\int x^{3/2}(a + b \operatorname{arctanh}(c\sqrt{x})) d\sqrt{x}}{c^2} \right) \end{aligned}$$

↓ 6452

$$2 \left(\frac{\int \frac{x^{3/2}(a + b \operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{4}x^2(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \int \frac{x^2}{1-c^2x} d\sqrt{x}}{c^2} \right)$$

↓ 254

$$2 \left(\frac{\int \frac{x^{3/2}(a + b \operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{4}x^2(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \int \left(-\frac{x}{c^2} + \frac{1}{c^4(1-c^2x)} - \frac{1}{c^4} \right) d\sqrt{x}}{c^2} \right)$$

↓ 2009

$$2 \left(\frac{\int \frac{x^{3/2}(a + b \operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{4}x^2(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^5} - \frac{\sqrt{x}}{c^4} - \frac{x^{3/2}}{3c^2} \right)}{c^2} \right)$$

↓ 6542

$$2 \left(\frac{\int \frac{\sqrt{x}(a + b \operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\int \frac{\sqrt{x}(a + b \operatorname{arctanh}(c\sqrt{x}))}{c^2} d\sqrt{x}}{c^2} - \frac{\frac{1}{4}x^2(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^5} - \frac{\sqrt{x}}{c^4} - \frac{x^{3/2}}{3c^2} \right)}{c^2} \right)$$

↓ 6452

$$2 \left(\frac{\int \frac{\sqrt{x}(a + b \operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{2}x(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \int \frac{x}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{4}x^2(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^5} - \frac{\sqrt{x}}{c^4} - \frac{x^{3/2}}{3c^2} \right)}{c^2} \right)$$

↓ 262

$$2 \left(\frac{\int \frac{\sqrt{x}(a + b \operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{2}x(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\int \frac{1}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\sqrt{x}}{c^2} \right)}{c^2} - \frac{\frac{1}{4}x^2(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^5} - \frac{\sqrt{x}}{c^4} - \frac{x^{3/2}}{3c^2} \right)}{c^2} \right)$$

↓ 219

$$2 \left(\frac{\int \frac{\sqrt{x}(a+b\operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2}\right)}{c^2}}{c^2} - \frac{\frac{1}{4}x^2(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc}{c^2} \right)$$

↓ 6546

$$2 \left(\frac{\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{1-c\sqrt{x}} d\sqrt{x}}{c} - \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2}\right)}{c^2}}{c^2} - \frac{\frac{1}{4}x^2(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{4}bc}{c^2} \right)$$

↓ 6470

$$2 \left(\frac{\frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)(a+b\operatorname{arctanh}(c\sqrt{x}))}{c} - b \int \frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)}{1-c^2x} d\sqrt{x}}{c} - \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2}\right)}{c^2}}{c^2} \right)$$

↓ 2849

$$2 \left(\frac{\frac{b \int \frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)}{1-\frac{2}{1-c\sqrt{x}}} d\frac{1}{1-c\sqrt{x}}}{c} + \frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)(a+b\operatorname{arctanh}(c\sqrt{x}))}{c}}{c} - \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2}\right)}{c^2}}{c^2} \right)$$

↓ 2752

$$2 \left(\frac{\frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)(a+b\operatorname{arctanh}(c\sqrt{x}))}{c} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-c\sqrt{x}}\right)}{2c}}{c} - \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2}\right)}{c^2}}{c^2} \right)$$

input

```
Int[(x^2*(a + b*ArcTanh[c*Sqrt[x]])/(1 - c^2*x), x]
```

output

$$2 * (- (((x^2 * (a + b * \operatorname{ArcTanh}[c * \sqrt{x}])) / 4 - (b * c * (- (\sqrt{x} / c^4) - x^{3/2}) / (3 * c^2) + \operatorname{ArcTanh}[c * \sqrt{x}] / c^5)) / 4) / c^2 + (- ((x * (a + b * \operatorname{ArcTanh}[c * \sqrt{x}])) / 2 - (b * c * (- (\sqrt{x} / c^2) + \operatorname{ArcTanh}[c * \sqrt{x}] / c^3)) / 2) / c^2 + (- 1 / 2 * (a + b * \operatorname{ArcTanh}[c * \sqrt{x}])^2 / (b * c^2) + (((a + b * \operatorname{ArcTanh}[c * \sqrt{x}]) * \operatorname{Log}[2 / (1 - c * \sqrt{x})]) / c + (b * \operatorname{PolyLog}[2, 1 - 2 / (1 - c * \sqrt{x})]) / (2 * c)) / c) / c^2) / c^2$$

Defintions of rubi rules used

rule 219

$$\operatorname{Int}[(a_.) + (b_.) * (x_.)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1 / (\operatorname{Rt}[a, 2] * \operatorname{Rt}[-b, 2])) * \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2] * (x / \operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$$

rule 254

$$\operatorname{Int}[(x_.)^{(m_.)} / ((a_.) + (b_.) * (x_.)^2), x_Symbol] \rightarrow \operatorname{Int}[\operatorname{PolynomialDivide}[x^m, a + b * x^2, x], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{IGtQ}[m, 3]$$

rule 262

$$\operatorname{Int}[(c_.) * (x_.)^{(m_.)} * ((a_.) + (b_.) * (x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[c * (c * x)^{(m-1)} * ((a + b * x^2)^{(p+1}) / (b * (m + 2 * p + 1))), x] - \operatorname{Simp}[a * c^2 * ((m-1) / (b * (m + 2 * p + 1))) \operatorname{Int}[(c * x)^{(m-2)} * (a + b * x^2)^p, x], x] /; \operatorname{FreeQ}\{a, b, c, p\}, x \ \&\& \operatorname{GtQ}[m, 2 - 1] \ \&\& \operatorname{NeQ}[m + 2 * p + 1, 0] \ \&\& \operatorname{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 2009

$$\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$$

rule 2752

$$\operatorname{Int}[\operatorname{Log}[(c_.) * (x_.)] / ((d_.) + (e_.) * (x_)), x_Symbol] \rightarrow \operatorname{Simp}[(-e^{-1}) * \operatorname{PolyLog}[2, 1 - c * x], x] /; \operatorname{FreeQ}\{c, d, e\}, x \ \&\& \operatorname{EqQ}[e + c * d, 0]$$

rule 2849

$$\operatorname{Int}[\operatorname{Log}[(c_.) / ((d_.) + (e_.) * (x_))] / ((f_.) + (g_.) * (x_.)^2), x_Symbol] \rightarrow \operatorname{Simp}[-e/g \operatorname{Subst}[\operatorname{Int}[\operatorname{Log}[2 * d * x] / (1 - 2 * d * x), x], x, 1 / (d + e * x)], x] /; \operatorname{FreeQ}\{c, d, e, f, g\}, x \ \&\& \operatorname{EqQ}[c, 2 * d] \ \&\& \operatorname{EqQ}[e^2 * f + d^2 * g, 0]$$

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6470 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol] :> Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6542 `Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_))/((d_) + (e_.)*(x_)^2), x_Symbol] :> Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

rule 6546 `Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*(x_)/((d_) + (e_.)*(x_)^2), x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 7267 `Int[u_, x_Symbol] :> With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Simp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.35

method	result
parts	$-\frac{ax^2}{2c^2} - \frac{ax}{c^4} - \frac{a \ln(c^2x-1)}{c^6} - \frac{2b \left(\frac{\operatorname{arctanh}(c\sqrt{x})c^4x^2}{4} + \frac{\operatorname{arctanh}(c\sqrt{x})c^2x}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x})}{2} \right)}{c^6}$
derivativedivides	$2 \left(a \left(\frac{c^4x^2}{4} + \frac{c^2x}{2} + \frac{\ln(c\sqrt{x}-1)}{2} + \frac{\ln(1+c\sqrt{x})}{2} \right) + b \left(\frac{\operatorname{arctanh}(c\sqrt{x})c^4x^2}{4} + \frac{\operatorname{arctanh}(c\sqrt{x})c^2x}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x})}{2} \right) \right)$
default	$2 \left(a \left(\frac{c^4x^2}{4} + \frac{c^2x}{2} + \frac{\ln(c\sqrt{x}-1)}{2} + \frac{\ln(1+c\sqrt{x})}{2} \right) + b \left(\frac{\operatorname{arctanh}(c\sqrt{x})c^4x^2}{4} + \frac{\operatorname{arctanh}(c\sqrt{x})c^2x}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x})}{2} \right) \right)$

```
input int(x^2*(a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x,method=_RETURNVERBOSE)
```

```
output -1/2*a/c^2*x^2-a/c^4*x-a/c^6*ln(c^2*x-1)-2*b/c^6*(1/4*arctanh(c*x^(1/2))*c^4*x^2+1/2*arctanh(c*x^(1/2))*c^2*x+1/2*arctanh(c*x^(1/2))*ln(c*x^(1/2)-1)+1/2*arctanh(c*x^(1/2))*ln(1+c*x^(1/2))-1/2*dilog(1/2*c*x^(1/2)+1/2)-1/4*ln(c*x^(1/2)-1)*ln(1/2*c*x^(1/2)+1/2)+1/8*ln(c*x^(1/2)-1)^2-1/8*ln(1+c*x^(1/2))^2+1/4*(ln(1+c*x^(1/2))-ln(1/2*c*x^(1/2)+1/2))*ln(-1/2*c*x^(1/2)+1/2)+1/12*c^3*x^(3/2)+3/4*c*x^(1/2)+3/8*ln(c*x^(1/2)-1)-3/8*ln(1+c*x^(1/2)))
```

Fricas [F]

$$\int \frac{x^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx = \int -\frac{(b \operatorname{arctanh}(c\sqrt{x}) + a)x^2}{c^2x - 1} dx$$

```
input integrate(x^2*(a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x, algorithm="fricas")
```

```
output integral(-(b*x^2*arctanh(c*sqrt(x)) + a*x^2)/(c^2*x - 1), x)
```

Sympy [F]

$$\int \frac{x^2(a + \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx = - \int \frac{ax^2}{c^2x - 1} dx - \int \frac{bx^2 \operatorname{atanh}(c\sqrt{x})}{c^2x - 1} dx$$

input `integrate(x**2*(a+b*atanh(c*x**(1/2)))/(-c**2*x+1),x)`

output `-Integral(a*x**2/(c**2*x - 1), x) - Integral(b*x**2*atanh(c*sqrt(x))/(c**2*x - 1), x)`

Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.30

$$\int \frac{x^2(a + \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx = -\frac{1}{2}a \left(\frac{c^2x^2 + 2x}{c^4} + \frac{2 \log(c^2x - 1)}{c^6} \right) - \frac{(\log(c\sqrt{x} + 1) \log(-\frac{1}{2}c\sqrt{x} + \frac{1}{2}) + \operatorname{Li}_2(\frac{1}{2}c\sqrt{x} + \frac{1}{2}))b}{c^6} + \frac{3b \log(c\sqrt{x} + 1)}{4c^6} - \frac{3b \log(c\sqrt{x} - 1)}{4c^6} - \frac{2bc^3x^{\frac{3}{2}} + 3b \log(c\sqrt{x} + 1)^2 - 3b \log(-c\sqrt{x} + 1)^2 + 18bc\sqrt{x} + 3(bc^4x^2 + 2bc^2x) \log(c\sqrt{x} + 1) - 3}{12c^6}$$

input `integrate(x^2*(a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x, algorithm="maxima")`

output `-1/2*a*((c^2*x^2 + 2*x)/c^4 + 2*log(c^2*x - 1)/c^6) - (log(c*sqrt(x) + 1)*log(-1/2*c*sqrt(x) + 1/2) + dilog(1/2*c*sqrt(x) + 1/2))*b/c^6 + 3/4*b*log(c*sqrt(x) + 1)/c^6 - 3/4*b*log(c*sqrt(x) - 1)/c^6 - 1/12*(2*b*c^3*x^(3/2) + 3*b*log(c*sqrt(x) + 1)^2 - 3*b*log(-c*sqrt(x) + 1)^2 + 18*b*c*sqrt(x) + 3*(b*c^4*x^2 + 2*b*c^2*x)*log(c*sqrt(x) + 1) - 3*(b*c^4*x^2 + 2*b*c^2*x + 2*b*log(c*sqrt(x) + 1))*log(-c*sqrt(x) + 1))/c^6`

Giac [F]

$$\int \frac{x^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx = \int -\frac{(b \operatorname{artanh}(c\sqrt{x}) + a)x^2}{c^2x - 1} dx$$

input `integrate(x^2*(a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x, algorithm="giac")`

output `integrate(-(b*arctanh(c*sqrt(x)) + a)*x^2/(c^2*x - 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx = \int -\frac{x^2(a + b \operatorname{atanh}(c\sqrt{x}))}{c^2x - 1} dx$$

input `int(-(x^2*(a + b*atanh(c*x^(1/2))))/(c^2*x - 1),x)`

output `int(-(x^2*(a + b*atanh(c*x^(1/2))))/(c^2*x - 1), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{x^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx \\ &= \frac{-2 \left(\int \frac{\operatorname{atanh}(\sqrt{x}c)x^2}{c^2x-1} dx \right) b c^6 - 2 \log(c^2x - 1) a - a c^4 x^2 - 2 a c^2 x}{2c^6} \end{aligned}$$

input `int(x^2*(a+b*atanh(c*x^(1/2)))/(-c^2*x+1),x)`

output `(- 2*int((atanh(sqrt(x)*c)*x**2)/(c**2*x - 1),x)*b*c**6 - 2*log(c**2*x - 1)*a - a*c**4*x**2 - 2*a*c**2*x)/(2*c**6)`

3.39 $\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx$

Optimal result	348
Mathematica [A] (verified)	349
Rubi [A] (verified)	349
Maple [A] (verified)	353
Fricas [F]	353
Sympy [F]	354
Maxima [A] (verification not implemented)	354
Giac [F]	355
Mupad [F(-1)]	355
Reduce [F]	355

Optimal result

Integrand size = 24, antiderivative size = 120

$$\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx = -\frac{b\sqrt{x}}{c^3} + \frac{b \operatorname{arctanh}(c\sqrt{x})}{c^4} - \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{c^2} - \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{bc^4} + \frac{2(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1 - c\sqrt{x}}\right)}{c^4} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - c\sqrt{x}}\right)}{c^4}$$

output

```
-b*x^(1/2)/c^3+b*arctanh(c*x^(1/2))/c^4-x*(a+b*arctanh(c*x^(1/2)))/c^2-(a+b*arctanh(c*x^(1/2)))^2/b/c^4+2*(a+b*arctanh(c*x^(1/2)))*ln(2/(1-c*x^(1/2)))/c^4+b*polylog(2,1-2/(1-c*x^(1/2)))/c^4
```

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.80

$$\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} dx = \frac{bc\sqrt{x} + ac^2x - b \operatorname{arctanh}(c\sqrt{x})^2 + b \operatorname{arctanh}(c\sqrt{x}) \left(-1 + c^2x - 2 \log\left(1 + e^{-2 \operatorname{arctanh}(c\sqrt{x})}\right)\right) + a \log\left(1 - c^2x\right)}{c^4}$$

input

```
Integrate[(x*(a + b*ArcTanh[c*Sqrt[x]]))/(1 - c^2*x),x]
```

output

```
-((b*c*Sqrt[x] + a*c^2*x - b*ArcTanh[c*Sqrt[x]]^2 + b*ArcTanh[c*Sqrt[x]]*(-1 + c^2*x - 2*Log[1 + E^(-2*ArcTanh[c*Sqrt[x]])])) + a*Log[1 - c^2*x] + b*PolyLog[2, -E^(-2*ArcTanh[c*Sqrt[x]])])/c^4
```

Rubi [A] (verified)Time = 0.85 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.22, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {7267, 6542, 6452, 262, 219, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} dx \\ & \quad \downarrow \text{7267} \\ & 2 \int \frac{x^{3/2}(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x} \\ & \quad \downarrow \text{6542} \\ & 2 \left(\frac{\int \frac{\sqrt{x}(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2 x} d\sqrt{x}}{c^2} - \frac{\int \sqrt{x}(a + b \operatorname{arctanh}(c\sqrt{x})) d\sqrt{x}}{c^2} \right) \\ & \quad \downarrow \text{6452} \end{aligned}$$

$$2 \left(\frac{\int \frac{\sqrt{x}(a+b\operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \int \frac{x}{1-c^2x} d\sqrt{x}}{c^2} \right)$$

↓ 262

$$2 \left(\frac{\int \frac{\sqrt{x}(a+b\operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\int \frac{1}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\sqrt{x}}{c^2} \right)}{c^2} \right)$$

↓ 219

$$2 \left(\frac{\int \frac{\sqrt{x}(a+b\operatorname{arctanh}(c\sqrt{x}))}{1-c^2x} d\sqrt{x}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2} \right)}{c^2} \right)$$

↓ 6546

$$2 \left(\frac{\frac{\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{1-c\sqrt{x}} d\sqrt{x}}{c} - \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2} \right)}{c^2} \right)$$

↓ 6470

$$2 \left(\frac{\frac{\frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)(a+b\operatorname{arctanh}(c\sqrt{x}))}{c} - b \int \frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)}{1-c^2x} d\sqrt{x}}{c} - \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2} \right)}{c^2} \right)$$

↓ 2849

$$2 \left(\frac{\frac{b \int \frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)}{1-c\sqrt{x}} d\frac{1}{1-c\sqrt{x}} + \frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)(a+b\operatorname{arctanh}(c\sqrt{x}))}{c}}{c} - \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2}}{c^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2} \right)}{c^2} \right)$$

↓ 2752

$$2 \left(\frac{\frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)(a+b\operatorname{arctanh}(c\sqrt{x}))}{c} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-c\sqrt{x}}\right)}{2c}}{c^2} - \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2} - \frac{\frac{1}{2}x(a+b\operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c}\right)}{c^2} \right)$$

input `Int[(x*(a + b*ArcTanh[c*Sqrt[x]]))/(1 - c^2*x),x]`

output `2*(-(((x*(a + b*ArcTanh[c*Sqrt[x]]))/2 - (b*c*(-(Sqrt[x]/c^2) + ArcTanh[c*Sqrt[x]]/c^3))/2)/c^2) + (-1/2*(a + b*ArcTanh[c*Sqrt[x]])^2/(b*c^2) + (((a + b*ArcTanh[c*Sqrt[x]])*Log[2/(1 - c*Sqrt[x]])/c + (b*PolyLog[2, 1 - 2/(1 - c*Sqrt[x]])/(2*c))/c)/c^2)`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 262 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m-1)*((a + b*x^2)^(p+1)/(b*(m+2*p+1))), x] - Simp[a*c^2*((m-1)/(b*(m+2*p+1))) Int[(c*x)^(m-2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2-1] && NeQ[m+2*p+1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2752 `Int[Log[(c_)*(x_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_)/((d_) + (e_)*(x_))]/((f_) + (g_)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6470 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol] :> Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6542 `Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_))/((d_) + (e_.)*(x_)^2), x_Symbol] :> Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

rule 6546 `Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*(x_)/((d_) + (e_.)*(x_)^2), x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 7267 `Int[u_, x_Symbol] :> With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Simp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.54

method	result
parts	$-\frac{ax}{c^2} - \frac{a \ln(c^2x-1)}{c^4} - \frac{2b \left(\frac{\operatorname{arctanh}(c\sqrt{x})c^2x}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1+c\sqrt{x})}{2} + \frac{c\sqrt{x}}{2} + \frac{\ln(c\sqrt{x}-1)}{4} \right)}{c^4}$
derivativedivides	$2 \left(a \left(\frac{c^2x}{2} + \frac{\ln(c\sqrt{x}-1)}{2} + \frac{\ln(1+c\sqrt{x})}{2} \right) + b \left(\frac{\operatorname{arctanh}(c\sqrt{x})c^2x}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1+c\sqrt{x})}{2} + \frac{c\sqrt{x}}{2} + \frac{\ln(c\sqrt{x}-1)}{4} \right) \right)$
default	$2 \left(a \left(\frac{c^2x}{2} + \frac{\ln(c\sqrt{x}-1)}{2} + \frac{\ln(1+c\sqrt{x})}{2} \right) + b \left(\frac{\operatorname{arctanh}(c\sqrt{x})c^2x}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}-1)}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1+c\sqrt{x})}{2} + \frac{c\sqrt{x}}{2} + \frac{\ln(c\sqrt{x}-1)}{4} \right) \right)$

input

```
int(x*(a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x,method=_RETURNVERBOSE)
```

output

```
-a/c^2*x-a/c^4*ln(c^2*x-1)-2*b/c^4*(1/2*arctanh(c*x^(1/2))*c^2*x+1/2*arctanh(c*x^(1/2))*ln(c*x^(1/2)-1)+1/2*arctanh(c*x^(1/2))*ln(1+c*x^(1/2))+1/2*c*x^(1/2)+1/4*ln(c*x^(1/2)-1)-1/4*ln(1+c*x^(1/2))-1/2*dilog(1/2*c*x^(1/2)+1/2)-1/4*ln(c*x^(1/2)-1)*ln(1/2*c*x^(1/2)+1/2)+1/8*ln(c*x^(1/2)-1)^2-1/8*ln(1+c*x^(1/2))^2+1/4*(ln(1+c*x^(1/2))-ln(1/2*c*x^(1/2)+1/2))*ln(-1/2*c*x^(1/2)+1/2))
```

Fricas [F]

$$\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx = \int -\frac{(b \operatorname{arctanh}(c\sqrt{x}) + a)x}{c^2x - 1} dx$$

input

```
integrate(x*(a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x, algorithm="fricas")
```

output

```
integral(-(b*x*arctanh(c*sqrt(x)) + a*x)/(c^2*x - 1), x)
```

Sympy [F]

$$\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx = - \int \frac{ax}{c^2x - 1} dx - \int \frac{bx \operatorname{atanh}(c\sqrt{x})}{c^2x - 1} dx$$

input `integrate(x*(a+b*atanh(c*x**(1/2)))/(-c**2*x+1),x)`

output `-Integral(a*x/(c**2*x - 1), x) - Integral(b*x*atanh(c*sqrt(x))/(c**2*x - 1), x)`

Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.38

$$\begin{aligned} & \int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx \\ &= -a \left(\frac{x}{c^2} + \frac{\log(c^2x - 1)}{c^4} \right) - \frac{(\log(c\sqrt{x} + 1) \log(-\frac{1}{2}c\sqrt{x} + \frac{1}{2}) + \operatorname{Li}_2(\frac{1}{2}c\sqrt{x} + \frac{1}{2}))b}{c^4} \\ & \quad + \frac{b \log(c\sqrt{x} + 1)}{2c^4} - \frac{b \log(c\sqrt{x} - 1)}{2c^4} \\ & \quad - \frac{2bc^2x \log(c\sqrt{x} + 1) + b \log(c\sqrt{x} + 1)^2 - b \log(-c\sqrt{x} + 1)^2 + 4bc\sqrt{x} - 2(bc^2x + b \log(c\sqrt{x} + 1))}{4c^4} \end{aligned}$$

input `integrate(x*(a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x, algorithm="maxima")`

output `-a*(x/c^2 + log(c^2*x - 1)/c^4) - (log(c*sqrt(x) + 1)*log(-1/2*c*sqrt(x) + 1/2) + dilog(1/2*c*sqrt(x) + 1/2))*b/c^4 + 1/2*b*log(c*sqrt(x) + 1)/c^4 - 1/2*b*log(c*sqrt(x) - 1)/c^4 - 1/4*(2*b*c^2*x*log(c*sqrt(x) + 1) + b*log(c*sqrt(x) + 1)^2 - b*log(-c*sqrt(x) + 1)^2 + 4*b*c*sqrt(x) - 2*(b*c^2*x + b*log(c*sqrt(x) + 1))*log(-c*sqrt(x) + 1))/c^4`

Giac [F]

$$\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx = \int -\frac{(b \operatorname{artanh}(c\sqrt{x}) + a)x}{c^2x - 1} dx$$

input `integrate(x*(a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x, algorithm="giac")`

output `integrate(-(b*arctanh(c*sqrt(x)) + a)*x/(c^2*x - 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx = \int -\frac{x(a + b \operatorname{atanh}(c\sqrt{x}))}{c^2x - 1} dx$$

input `int(-(x*(a + b*atanh(c*x^(1/2))))/(c^2*x - 1),x)`

output `int(-(x*(a + b*atanh(c*x^(1/2))))/(c^2*x - 1), x)`

Reduce [F]

$$\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} dx = \frac{-\left(\int \frac{\operatorname{atanh}(\sqrt{x}c)x}{c^2x-1} dx\right) b c^4 - \log(c^2x - 1) a - a c^2x}{c^4}$$

input `int(x*(a+b*atanh(c*x^(1/2)))/(-c^2*x+1),x)`

output `(- (int((atanh(sqrt(x)*c)*x)/(c**2*x - 1),x)*b*c**4 + log(c**2*x - 1)*a + a*c**2*x))/c**4`

3.40 $\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{1-c^2x} dx$

Optimal result	356
Mathematica [A] (verified)	356
Rubi [A] (verified)	357
Maple [A] (verified)	359
Fricas [F]	359
Sympy [F]	360
Maxima [A] (verification not implemented)	360
Giac [F]	361
Mupad [F(-1)]	361
Reduce [F]	361

Optimal result

Integrand size = 23, antiderivative size = 78

$$\int \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{1 - c^2x} dx = -\frac{(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{bc^2} + \frac{2(a + b\operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1-c\sqrt{x}}\right)}{c^2} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-c\sqrt{x}}\right)}{c^2}$$

output

```
-(a+b*arctanh(c*x^(1/2)))^2/b/c^2+2*(a+b*arctanh(c*x^(1/2)))*ln(2/(1-c*x^(1/2)))/c^2+b*polylog(2,1-2/(1-c*x^(1/2)))/c^2
```

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.96

$$\int \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{1 - c^2x} dx = -\frac{a \log(1 - c^2x)}{c^2} - \frac{b\left(-\operatorname{arctanh}(c\sqrt{x})\left(\operatorname{arctanh}(c\sqrt{x}) + 2 \log\left(1 + e^{-2\operatorname{arctanh}(c\sqrt{x})}\right)\right) + \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arctanh}(c\sqrt{x})}\right)\right)}{c^2}$$

input `Integrate[(a + b*ArcTanh[c*Sqrt[x]])/(1 - c^2*x),x]`

output `-((a*Log[1 - c^2*x])/c^2) - (b*(-(ArcTanh[c*Sqrt[x]]*(ArcTanh[c*Sqrt[x]] + 2*Log[1 + E^(-2*ArcTanh[c*Sqrt[x]])]))) + PolyLog[2, -E^(-2*ArcTanh[c*Sqrt[x]])])/c^2`

Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.14, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {6484, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{1 - c^2x} dx \\
 & \quad \downarrow 6484 \\
 & 2 \int \frac{\sqrt{x}(a + b \operatorname{arctanh}(c\sqrt{x}))}{1 - c^2x} d\sqrt{x} \\
 & \quad \downarrow 6546 \\
 & 2 \left(\frac{\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{1 - c\sqrt{x}} d\sqrt{x}}{c} - \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2} \right) \\
 & \quad \downarrow 6470 \\
 & 2 \left(\frac{\log\left(\frac{2}{1 - c\sqrt{x}}\right)(a + b \operatorname{arctanh}(c\sqrt{x}))}{c} - b \int \frac{\log\left(\frac{2}{1 - c\sqrt{x}}\right)}{1 - c^2x} d\sqrt{x} - \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2} \right) \\
 & \quad \downarrow 2849 \\
 & 2 \left(\frac{b \int \frac{\log\left(\frac{2}{1 - c\sqrt{x}}\right)}{1 - c\sqrt{x}} d\frac{1}{1 - c\sqrt{x}} + \frac{\log\left(\frac{2}{1 - c\sqrt{x}}\right)(a + b \operatorname{arctanh}(c\sqrt{x}))}{c}}{c} - \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2} \right)
 \end{aligned}$$

$$2 \left(\frac{\log\left(\frac{2}{1-c\sqrt{x}}\right)(a+b\operatorname{arctanh}(c\sqrt{x}))}{c} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-c\sqrt{x}}\right)}{2c} - \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))^2}{2bc^2} \right)$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])/(1 - c^2*x), x]`

output `2*(-1/2*(a + b*ArcTanh[c*Sqrt[x]])^2/(b*c^2) + ((a + b*ArcTanh[c*Sqrt[x]])*Log[2/(1 - c*Sqrt[x])])/c + (b*PolyLog[2, 1 - 2/(1 - c*Sqrt[x])])/(2*c))/c)`

Defintions of rubi rules used

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 6470 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^p/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))])/(1 - c^2*x^2)], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6484 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.)/((d_) + (e_.)*(x_)), x_Symbol] := With[{k = Denominator[n]}, Simp[k Subst[Int[x^(k - 1)*((a + b*ArcTanh[c*x^(k*n)])/(d + e*x^k)), x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, e}, x] && FractionQ[n]`

rule 6546

```
Int[(((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)*(x_.))/((d_.) + (e_.)*(x_)^2),
  x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/
(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.79

method	result
parts	$-\frac{a \ln(c^2 x - 1)}{c^2} - \frac{2b \left(\frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x} - 1)}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1 + c\sqrt{x})}{2} - \frac{\operatorname{dilog}\left(\frac{c\sqrt{x}}{2} + \frac{1}{2}\right)}{2} - \frac{\ln(c\sqrt{x} - 1) \ln\left(\frac{c\sqrt{x}}{2} + \frac{1}{2}\right)}{4} \right)}{c^2} + \dots$
derivativedivides	$2 \left(a \left(\frac{\ln(c\sqrt{x} - 1)}{2} + \frac{\ln(1 + c\sqrt{x})}{2} \right) + b \left(\frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x} - 1)}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1 + c\sqrt{x})}{2} - \frac{\operatorname{dilog}\left(\frac{c\sqrt{x}}{2} + \frac{1}{2}\right)}{2} - \frac{\ln(c\sqrt{x} - 1) \ln\left(\frac{c\sqrt{x}}{2} + \frac{1}{2}\right)}{4} \right) \right) / c^2$
default	$2 \left(a \left(\frac{\ln(c\sqrt{x} - 1)}{2} + \frac{\ln(1 + c\sqrt{x})}{2} \right) + b \left(\frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x} - 1)}{2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(1 + c\sqrt{x})}{2} - \frac{\operatorname{dilog}\left(\frac{c\sqrt{x}}{2} + \frac{1}{2}\right)}{2} - \frac{\ln(c\sqrt{x} - 1) \ln\left(\frac{c\sqrt{x}}{2} + \frac{1}{2}\right)}{4} \right) \right) / c^2$

input

```
int((a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x,method=_RETURNVERBOSE)
```

output

```
-a*ln(c^2*x-1)/c^2-2*b/c^2*(1/2*arctanh(c*x^(1/2))*ln(c*x^(1/2)-1)+1/2*arc
tanh(c*x^(1/2))*ln(1+c*x^(1/2))-1/2*dilog(1/2*c*x^(1/2)+1/2)-1/4*ln(c*x^(1
/2)-1)*ln(1/2*c*x^(1/2)+1/2)+1/8*ln(c*x^(1/2)-1)^2-1/8*ln(1+c*x^(1/2))^2+1
/4*(ln(1+c*x^(1/2))-ln(1/2*c*x^(1/2)+1/2))*ln(-1/2*c*x^(1/2)+1/2))
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{1 - c^2 x} dx = \int -\frac{b \operatorname{arctanh}(c\sqrt{x}) + a}{c^2 x - 1} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x, algorithm="fricas")
```

output

```
integral(-(b*arctanh(c*sqrt(x)) + a)/(c^2*x - 1), x)
```


Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{1 - c^2x} dx = - \int \frac{a}{c^2x - 1} dx - \int \frac{b \operatorname{atanh}(c\sqrt{x})}{c^2x - 1} dx$$

input `integrate((a+b*atanh(c*x**(1/2)))/(-c**2*x+1),x)`

output `-Integral(a/(c**2*x - 1), x) - Integral(b*atanh(c*sqrt(x))/(c**2*x - 1), x)`

Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.29

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{1 - c^2x} dx \\ &= - \frac{(\log(c\sqrt{x} + 1) \log(-\frac{1}{2}c\sqrt{x} + \frac{1}{2}) + \operatorname{Li}_2(\frac{1}{2}c\sqrt{x} + \frac{1}{2}))b}{c^2} - \frac{a \log(c^2x - 1)}{c^2} \\ & \quad - \frac{b \log(c\sqrt{x} + 1)^2 - 2b \log(c\sqrt{x} + 1) \log(-c\sqrt{x} + 1) - b \log(-c\sqrt{x} + 1)^2}{4c^2} \end{aligned}$$

input `integrate((a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x, algorithm="maxima")`

output `-(log(c*sqrt(x) + 1)*log(-1/2*c*sqrt(x) + 1/2) + dilog(1/2*c*sqrt(x) + 1/2)) * b/c^2 - a*log(c^2*x - 1)/c^2 - 1/4*(b*log(c*sqrt(x) + 1)^2 - 2*b*log(c*sqrt(x) + 1)*log(-c*sqrt(x) + 1) - b*log(-c*sqrt(x) + 1)^2)/c^2`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{1 - c^2 x} dx = \int -\frac{b \operatorname{artanh}(c\sqrt{x}) + a}{c^2 x - 1} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/(-c^2*x+1),x, algorithm="giac")`

output `integrate(-(b*arctanh(c*sqrt(x)) + a)/(c^2*x - 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{1 - c^2 x} dx = \int -\frac{a + b \operatorname{atanh}(c\sqrt{x})}{c^2 x - 1} dx$$

input `int(-(a + b*atanh(c*x^(1/2)))/(c^2*x - 1),x)`

output `int(-(a + b*atanh(c*x^(1/2)))/(c^2*x - 1), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{1 - c^2 x} dx = \frac{-\left(\int \frac{\operatorname{atanh}(\sqrt{x}c)}{c^2 x - 1} dx\right) b c^2 - \log(c^2 x - 1) a}{c^2}$$

input `int((a+b*atanh(c*x^(1/2)))/(-c^2*x+1),x)`

output `(- (int(atanh(sqrt(x)*c)/(c**2*x - 1),x)*b*c**2 + log(c**2*x - 1)*a))/c**2`

$$3.41 \quad \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(1-c^2x)} dx$$

Optimal result	362
Mathematica [A] (verified)	362
Rubi [A] (verified)	363
Maple [B] (verified)	365
Fricas [F]	365
Sympy [F]	366
Maxima [B] (verification not implemented)	366
Giac [F]	367
Mupad [F(-1)]	367
Reduce [F]	367

Optimal result

Integrand size = 26, antiderivative size = 69

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(1-c^2x)} dx = \frac{(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{b} + 2(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(2 - \frac{2}{1 + c\sqrt{x}}\right) - b \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + c\sqrt{x}}\right)$$

output

```
(a+b*arctanh(c*x^(1/2)))^2/b+2*(a+b*arctanh(c*x^(1/2)))*ln(2-2/(1+c*x^(1/2)))-b*polylog(2,-1+2/(1+c*x^(1/2)))
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.07

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(1-c^2x)} dx = -b \operatorname{arctanh}(c\sqrt{x})^2 + 2b \operatorname{arctanh}(c\sqrt{x}) \log\left(1 - e^{2 \operatorname{arctanh}(c\sqrt{x})}\right) + a(\log(x) - \log(1 - c^2x)) + b \operatorname{PolyLog}\left(2, e^{2 \operatorname{arctanh}(c\sqrt{x})}\right)$$

input `Integrate[(a + b*ArcTanh[c*Sqrt[x]])/(x*(1 - c^2*x)),x]`

output `-(b*ArcTanh[c*Sqrt[x]]^2) + 2*b*ArcTanh[c*Sqrt[x]]*Log[1 - E^(2*ArcTanh[c*Sqrt[x]])] + a*(Log[x] - Log[1 - c^2*x]) + b*PolyLog[2, E^(2*ArcTanh[c*Sqrt[x]])]`

Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.09, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$, Rules used = {7267, 2026, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x(1 - c^2x)} dx \\
 & \quad \downarrow \text{7267} \\
 & 2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x} - c^2x^{3/2}} d\sqrt{x} \\
 & \quad \downarrow \text{2026} \\
 & 2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1 - c^2x)} d\sqrt{x} \\
 & \quad \downarrow \text{6550} \\
 & 2 \left(\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{(\sqrt{xc} + 1)\sqrt{x}} d\sqrt{x} + \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} \right) \\
 & \quad \downarrow \text{6494} \\
 & 2 \left(-bc \int \frac{\log\left(2 - \frac{2}{\sqrt{xc}+1}\right)}{1 - c^2x} d\sqrt{x} + \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} + \log\left(2 - \frac{2}{c\sqrt{x}+1}\right) (a + \operatorname{barctanh}(c\sqrt{x})) \right) \\
 & \quad \downarrow \text{2897}
 \end{aligned}$$

$$2 \left(\frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} + \log \left(2 - \frac{2}{c\sqrt{x} + 1} \right) (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{2} b \operatorname{PolyLog} \left(2, \frac{2}{\sqrt{xc} + 1} - 1 \right) \right)$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])/(x*(1 - c^2*x)),x]`

output `2*((a + b*ArcTanh[c*Sqrt[x]])^2/(2*b) + (a + b*ArcTanh[c*Sqrt[x]])*Log[2 - 2/(1 + c*Sqrt[x])] - (b*PolyLog[2, -1 + 2/(1 + c*Sqrt[x])]))/2`

Defintions of rubi rules used

rule 2026 `Int[(Fx_)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2897 `Int[Log[u_]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6494 `Int[((a_) + ArcTanh[(c_)*(x_)]*(b_))^(p_)/((x_)*((d_) + (e_)*(x_))), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6550 `Int[((a_) + ArcTanh[(c_)*(x_)]*(b_))^(p_)/((x_)*((d_) + (e_)*(x_)^2)), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]`

rule 7267

```
Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2]])], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 179 vs. $2(61) = 122$.

Time = 0.09 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.61

method	result
parts	$-a(-\ln(x) + \ln(c^2x - 1)) - b\left(-2 \operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}) + \operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x})\right)$
derivativedivides	$-2a\left(-\ln(c\sqrt{x}) + \frac{\ln(c\sqrt{x}-1)}{2} + \frac{\ln(1+c\sqrt{x})}{2}\right) - 2b\left(-\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}) + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x})}{2}\right)$
default	$-2a\left(-\ln(c\sqrt{x}) + \frac{\ln(c\sqrt{x}-1)}{2} + \frac{\ln(1+c\sqrt{x})}{2}\right) - 2b\left(-\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}) + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x})}{2}\right)$

input

```
int((a+b*arctanh(c*x^(1/2)))/x/(-c^2*x+1),x,method=_RETURNVERBOSE)
```

output

```
-a*(-ln(x)+ln(c^2*x-1))-b*(-2*arctanh(c*x^(1/2))*ln(c*x^(1/2))+arctanh(c*x
^(1/2))*ln(c*x^(1/2)-1)+arctanh(c*x^(1/2))*ln(1+c*x^(1/2))+dilog(c*x^(1/2)
)+dilog(1+c*x^(1/2))+ln(c*x^(1/2))*ln(1+c*x^(1/2))+1/4*ln(c*x^(1/2)-1)^2-d
ilog(1/2*c*x^(1/2)+1/2)-1/2*ln(c*x^(1/2)-1)*ln(1/2*c*x^(1/2)+1/2)-1/4*ln(1
+c*x^(1/2))^2+1/2*(ln(1+c*x^(1/2))-ln(1/2*c*x^(1/2)+1/2))*ln(-1/2*c*x^(1/2
)+1/2))
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(1 - c^2x)} dx = \int -\frac{b \operatorname{arctanh}(c\sqrt{x}) + a}{(c^2x - 1)x} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))/x/(-c^2*x+1),x, algorithm="fricas")
```

output `integral(-(b*arctanh(c*sqrt(x)) + a)/(c^2*x^2 - x), x)`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(1 - c^2x)} dx = - \int \frac{a}{c^2x^2 - x} dx - \int \frac{b \operatorname{atanh}(c\sqrt{x})}{c^2x^2 - x} dx$$

input `integrate((a+b*atanh(c*x**(1/2)))/x/(-c**2*x+1), x)`

output `-Integral(a/(c**2*x**2 - x), x) - Integral(b*atanh(c*sqrt(x))/(c**2*x**2 - x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 159 vs. $2(60) = 120$.

Time = 0.17 (sec) , antiderivative size = 159, normalized size of antiderivative = 2.30

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(1 - c^2x)} dx \\ &= -\frac{1}{4} b \log(c\sqrt{x} + 1)^2 + \frac{1}{2} b \log(c\sqrt{x} + 1) \log(-c\sqrt{x} + 1) + \frac{1}{4} b \log(-c\sqrt{x} + 1)^2 \\ & \quad - \left(\log(c\sqrt{x} + 1) \log\left(-\frac{1}{2}c\sqrt{x} + \frac{1}{2}\right) + \operatorname{Li}_2\left(\frac{1}{2}c\sqrt{x} + \frac{1}{2}\right) \right) b \\ & \quad - (\log(c\sqrt{x}) \log(-c\sqrt{x} + 1) + \operatorname{Li}_2(-c\sqrt{x} + 1)) b \\ & \quad + (\log(c\sqrt{x} + 1) \log(-c\sqrt{x}) + \operatorname{Li}_2(c\sqrt{x} + 1)) b \\ & \quad - a(\log(c\sqrt{x} + 1) + \log(c\sqrt{x} - 1) - \log(x)) \end{aligned}$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x/(-c^2*x+1), x, algorithm="maxima")`

output `-1/4*b*log(c*sqrt(x) + 1)^2 + 1/2*b*log(c*sqrt(x) + 1)*log(-c*sqrt(x) + 1) + 1/4*b*log(-c*sqrt(x) + 1)^2 - (log(c*sqrt(x) + 1)*log(-1/2*c*sqrt(x) + 1/2) + dilog(1/2*c*sqrt(x) + 1/2))*b - (log(c*sqrt(x))*log(-c*sqrt(x) + 1) + dilog(-c*sqrt(x) + 1))*b + (log(c*sqrt(x) + 1)*log(-c*sqrt(x)) + dilog(c*sqrt(x) + 1))*b - a*(log(c*sqrt(x) + 1) + log(c*sqrt(x) - 1) - log(x))`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(1 - c^2x)} dx = \int -\frac{b \operatorname{artanh}(c\sqrt{x}) + a}{(c^2x - 1)x} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x/(-c^2*x+1),x, algorithm="giac")`

output `integrate(-(b*arctanh(c*sqrt(x)) + a)/((c^2*x - 1)*x), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(1 - c^2x)} dx = - \int \frac{a + b \operatorname{atanh}(c\sqrt{x})}{x(c^2x - 1)} dx$$

input `int(-(a + b*atanh(c*x^(1/2)))/(x*(c^2*x - 1)),x)`

output `-int((a + b*atanh(c*x^(1/2)))/(x*(c^2*x - 1)), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(1 - c^2x)} dx = - \left(\int \frac{\operatorname{atanh}(\sqrt{x}c)}{c^2x^2 - x} dx \right) b - \log(c^2x - 1) a + \log(x) a$$

input `int((a+b*atanh(c*x^(1/2)))/x/(-c^2*x+1),x)`

output `- int(atanh(sqrt(x)*c)/(c**2*x**2 - x),x)*b - log(c**2*x - 1)*a + log(x)*a`

3.42 $\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x^2(1-c^2x)} dx$

Optimal result	368
Mathematica [A] (verified)	369
Rubi [A] (verified)	369
Maple [B] (verified)	372
Fricas [F]	373
Sympy [F]	373
Maxima [B] (verification not implemented)	374
Giac [F]	374
Mupad [F(-1)]	375
Reduce [F]	375

Optimal result

Integrand size = 26, antiderivative size = 117

$$\int \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{x^2(1 - c^2x)} dx = -\frac{bc}{\sqrt{x}} + bc^2\operatorname{arctanh}(c\sqrt{x}) - \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{x} + \frac{c^2(a + b\operatorname{arctanh}(c\sqrt{x}))^2}{b} + 2c^2(a + b\operatorname{arctanh}(c\sqrt{x})) \log\left(2 - \frac{2}{1 + c\sqrt{x}}\right) - bc^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + c\sqrt{x}}\right)$$

output

```
-b*c/x^(1/2)+b*c^2*arctanh(c*x^(1/2))-(a+b*arctanh(c*x^(1/2)))/x+c^2*(a+b*arctanh(c*x^(1/2)))^2/b+2*c^2*(a+b*arctanh(c*x^(1/2)))*ln(2-2/(1+c*x^(1/2)))-b*c^2*polylog(2,-1+2/(1+c*x^(1/2)))
```

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.01

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(1 - c^2x)} dx = -\frac{a}{x} + 2ac^2 \log(\sqrt{x}) - ac^2 \log(1 - c^2x) - bc^2 \left(\frac{1}{c\sqrt{x}} - \operatorname{arctanh}(c\sqrt{x}) \left(-\frac{1 - c^2x}{c^2x} + \operatorname{arctanh}(c\sqrt{x}) + 2 \log(1 - e^{-2\operatorname{arctanh}(c\sqrt{x})}) \right) + \operatorname{PolyLog}\left(2, e^{-2\operatorname{arctanh}(c\sqrt{x})}\right) \right)$$

input `Integrate[(a + b*ArcTanh[c*Sqrt[x]])/(x^2*(1 - c^2*x)),x]`

output `-(a/x) + 2*a*c^2*Log[Sqrt[x]] - a*c^2*Log[1 - c^2*x] - b*c^2*(1/(c*Sqrt[x]) - ArcTanh[c*Sqrt[x]]*(-((1 - c^2*x)/(c^2*x)) + ArcTanh[c*Sqrt[x]] + 2*Log[1 - E^(-2*ArcTanh[c*Sqrt[x]])]) + PolyLog[2, E^(-2*ArcTanh[c*Sqrt[x]])])`

Rubi [A] (verified)

Time = 0.90 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.05, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.346$, Rules used = {7267, 2026, 6544, 6452, 264, 219, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(1 - c^2x)} dx$$

↓ 7267

$$2 \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^{3/2} - c^2x^{5/2}} d\sqrt{x}$$

↓ 2026

$$\begin{aligned}
& 2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(1-c^2x)} d\sqrt{x} \\
& \quad \downarrow 6544 \\
& 2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} + \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}} d\sqrt{x} \right) \\
& \quad \downarrow 6452 \\
& 2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} + \frac{1}{2}bc \int \frac{1}{x(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{2x} \right) \\
& \quad \downarrow 264 \\
& 2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} + \frac{1}{2}bc \left(c^2 \int \frac{1}{1-c^2x} d\sqrt{x} - \frac{1}{\sqrt{x}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{2x} \right) \\
& \quad \downarrow 219 \\
& 2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{2x} + \frac{1}{2}bc \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) \right) \\
& \quad \downarrow 6550 \\
& 2 \left(c^2 \left(\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{(\sqrt{xc}+1)\sqrt{x}} d\sqrt{x} + \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{2x} + \frac{1}{2}bc \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) \right) \\
& \quad \downarrow 6494 \\
& 2 \left(c^2 \left(-bc \int \frac{\log \left(2 - \frac{2}{\sqrt{xc}+1} \right)}{1-c^2x} d\sqrt{x} + \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} + \log \left(2 - \frac{2}{c\sqrt{x}+1} \right) (a + \operatorname{barctanh}(c\sqrt{x})) \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{2x} + \frac{1}{2}bc \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) \right) \\
& \quad \downarrow 2897 \\
& 2 \left(c^2 \left(\frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} + \log \left(2 - \frac{2}{c\sqrt{x}+1} \right) (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{2}b \operatorname{PolyLog} \left(2, \frac{2}{\sqrt{xc}+1} - 1 \right) \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{2x} + \frac{1}{2}bc \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) \right)
\end{aligned}$$

input `Int[(a + b*ArcTanh[c*sqrt[x]])/(x^2*(1 - c^2*x)),x]`

output

```
2*(-1/2*(a + b*ArcTanh[c*Sqrt[x]])/x + (b*c*(-1/Sqrt[x]) + c*ArcTanh[c*Sqrt[x]))/2 + c^2*((a + b*ArcTanh[c*Sqrt[x]])^2/(2*b) + (a + b*ArcTanh[c*Sqrt[x]])*Log[2 - 2/(1 + c*Sqrt[x])] - (b*PolyLog[2, -1 + 2/(1 + c*Sqrt[x])])/2)
```

Defintions of rubi rules used

rule 219

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

rule 264

```
Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*((m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]
```

rule 2026

```
Int[(Fx_)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])
```

rule 2897

```
Int[Log[u]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]
```

rule 6452

```
Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]
```

rule 6494 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_))), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6544 `Int((((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*((f_.)*(x_))^(m_))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]`

rule 6550 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_)^2)), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]`

rule 7267 `Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Simp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 242 vs. $2(103) = 206$.

Time = 0.10 (sec) , antiderivative size = 243, normalized size of antiderivative = 2.08

method	result
parts	$-\frac{a}{x} + a c^2 \ln(x) - a c^2 \ln(c^2 x - 1) - 2b c^2 \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{2c^2 x} - \operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x}) + \dots \right)$
derivativedivides	$-2c^2 \left(a \left(\frac{1}{2c^2 x} - \ln(c\sqrt{x}) + \frac{\ln(c\sqrt{x}-1)}{2} + \frac{\ln(1+c\sqrt{x})}{2} \right) + b \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{2c^2 x} - \operatorname{arctanh}(c\sqrt{x}) \right) \right)$
default	$-2c^2 \left(a \left(\frac{1}{2c^2 x} - \ln(c\sqrt{x}) + \frac{\ln(c\sqrt{x}-1)}{2} + \frac{\ln(1+c\sqrt{x})}{2} \right) + b \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{2c^2 x} - \operatorname{arctanh}(c\sqrt{x}) \right) \right)$

input `int((a+b*arctanh(c*x^(1/2)))/x^2/(-c^2*x+1), x, method=_RETURNVERBOSE)`

output

```
-a/x+a*c^2*ln(x)-a*c^2*ln(c^2*x-1)-2*b*c^2*(1/2*arctanh(c*x^(1/2))/c^2/x-a
rctanh(c*x^(1/2))*ln(c*x^(1/2))+1/2*arctanh(c*x^(1/2))*ln(c*x^(1/2)-1)+1/2
*arctanh(c*x^(1/2))*ln(1+c*x^(1/2))+1/2/c/x^(1/2)+1/4*ln(c*x^(1/2)-1)-1/4*
ln(1+c*x^(1/2))+1/2*dilog(c*x^(1/2))+1/2*dilog(1+c*x^(1/2))+1/2*ln(c*x^(1/
2))*ln(1+c*x^(1/2))-1/2*dilog(1/2*c*x^(1/2)+1/2)-1/4*ln(c*x^(1/2)-1)*ln(1/
2*c*x^(1/2)+1/2)+1/8*ln(c*x^(1/2)-1)^2-1/8*ln(1+c*x^(1/2))^2+1/4*(ln(1+c*x
^(1/2))-ln(1/2*c*x^(1/2)+1/2))*ln(-1/2*c*x^(1/2)+1/2))
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(1 - c^2x)} dx = \int -\frac{b \operatorname{atanh}(c\sqrt{x}) + a}{(c^2x - 1)x^2} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))/x^2/(-c^2*x+1),x, algorithm="fricas")
```

output

```
integral(-(b*arctanh(c*sqrt(x)) + a)/(c^2*x^3 - x^2), x)
```

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(1 - c^2x)} dx = -\int \frac{a}{c^2x^3 - x^2} dx - \int \frac{b \operatorname{atanh}(c\sqrt{x})}{c^2x^3 - x^2} dx$$

input

```
integrate((a+b*atanh(c*x**(1/2)))/x**2/(-c**2*x+1),x)
```

output

```
-Integral(a/(c**2*x**3 - x**2), x) - Integral(b*atanh(c*sqrt(x))/(c**2*x**
3 - x**2), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 248 vs. $2(102) = 204$.

Time = 0.19 (sec) , antiderivative size = 248, normalized size of antiderivative = 2.12

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(1-c^2x)} dx$$

$$= -\left(\log(c\sqrt{x}+1)\log\left(-\frac{1}{2}c\sqrt{x}+\frac{1}{2}\right) + \operatorname{Li}_2\left(\frac{1}{2}c\sqrt{x}+\frac{1}{2}\right)\right)bc^2$$

$$- (\log(c\sqrt{x})\log(-c\sqrt{x}+1) + \operatorname{Li}_2(-c\sqrt{x}+1))bc^2$$

$$+ (\log(c\sqrt{x}+1)\log(-c\sqrt{x}) + \operatorname{Li}_2(c\sqrt{x}+1))bc^2 + \frac{1}{2}bc^2\log(c\sqrt{x}+1)$$

$$- \frac{1}{2}bc^2\log(c\sqrt{x}-1) - \left(c^2\log(c\sqrt{x}+1) + c^2\log(c\sqrt{x}-1) - c^2\log(x) + \frac{1}{x}\right)a$$

$$- \frac{bc^2x\log(c\sqrt{x}+1)^2 - bc^2x\log(-c\sqrt{x}+1)^2 + 4bc\sqrt{x} + 2b\log(c\sqrt{x}+1) - 2(bc^2x\log(c\sqrt{x}+1) + 1)}{4x}$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^2/(-c^2*x+1),x, algorithm="maxima")`

output `-(log(c*sqrt(x) + 1)*log(-1/2*c*sqrt(x) + 1/2) + dilog(1/2*c*sqrt(x) + 1/2))*b*c^2 - (log(c*sqrt(x))*log(-c*sqrt(x) + 1) + dilog(-c*sqrt(x) + 1))*b*c^2 + (log(c*sqrt(x) + 1)*log(-c*sqrt(x)) + dilog(c*sqrt(x) + 1))*b*c^2 + 1/2*b*c^2*log(c*sqrt(x) + 1) - 1/2*b*c^2*log(c*sqrt(x) - 1) - (c^2*log(c*sqrt(x) + 1) + c^2*log(c*sqrt(x) - 1) - c^2*log(x) + 1/x)*a - 1/4*(b*c^2*x*log(c*sqrt(x) + 1)^2 - b*c^2*x*log(-c*sqrt(x) + 1)^2 + 4*b*c*sqrt(x) + 2*b*log(c*sqrt(x) + 1) - 2*(b*c^2*x*log(c*sqrt(x) + 1) + b)*log(-c*sqrt(x) + 1))/x`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(1-c^2x)} dx = \int -\frac{b \operatorname{artanh}(c\sqrt{x}) + a}{(c^2x-1)x^2} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^2/(-c^2*x+1),x, algorithm="giac")`

output `integrate(-(b*arctanh(c*sqrt(x)) + a)/((c^2*x - 1)*x^2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2 (1 - c^2 x)} dx = - \int \frac{a + b \operatorname{atanh}(c\sqrt{x})}{x^2 (c^2 x - 1)} dx$$

input `int(-(a + b*atanh(c*x^(1/2)))/(x^2*(c^2*x - 1)),x)`

output `-int((a + b*atanh(c*x^(1/2)))/(x^2*(c^2*x - 1)), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2 (1 - c^2 x)} dx \\ &= \frac{- \left(\int \frac{\operatorname{atanh}(\sqrt{x} c)}{c^2 x^3 - x^2} dx \right) b x - \log(c^2 x - 1) a c^2 x + \log(x) a c^2 x - a}{x} \end{aligned}$$

input `int((a+b*atanh(c*x^(1/2)))/x^2/(-c^2*x+1),x)`

output `(- int(atanh(sqrt(x)*c)/(c**2*x**3 - x**2),x)*b*x - log(c**2*x - 1)*a*c**2*x + log(x)*a*c**2*x - a)/x`

$$3.43 \quad \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(1-c^2x)} dx$$

Optimal result	376
Mathematica [A] (verified)	377
Rubi [A] (verified)	377
Maple [B] (verified)	381
Fricas [F]	382
Sympy [F]	382
Maxima [B] (verification not implemented)	383
Giac [F]	384
Mupad [F(-1)]	384
Reduce [F]	384

Optimal result

Integrand size = 26, antiderivative size = 157

$$\begin{aligned} \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(1-c^2x)} dx = & -\frac{bc}{6x^{3/2}} - \frac{3bc^3}{2\sqrt{x}} + \frac{3}{2}bc^4 \operatorname{arctanh}(c\sqrt{x}) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{2x^2} \\ & - \frac{c^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{x} + \frac{c^4(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{b} \\ & + 2c^4(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(2 - \frac{2}{1 + c\sqrt{x}}\right) \\ & - bc^4 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + c\sqrt{x}}\right) \end{aligned}$$

output

```
-1/6*b*c/x^(3/2)-3/2*b*c^3/x^(1/2)+3/2*b*c^4*arctanh(c*x^(1/2))-1/2*(a+b*arctanh(c*x^(1/2)))/x^2-c^2*(a+b*arctanh(c*x^(1/2)))/x+c^4*(a+b*arctanh(c*x^(1/2)))^2/b+2*c^4*(a+b*arctanh(c*x^(1/2)))*ln(2-2/(1+c*x^(1/2)))-b*c^4*polylog(2,-1+2/(1+c*x^(1/2)))
```

Mathematica [A] (verified)

Time = 0.45 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.01

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(1-c^2x)} dx = \frac{3a + bc\sqrt{x} + 6ac^2x + 9bc^3x^{3/2} - 6bc^4x^2 \operatorname{arctanh}(c\sqrt{x})^2 - 3b \operatorname{arctanh}(c\sqrt{x}) \left(-1 - 2c^2x + 3c^4x^2 + 4c^4x^3\right)}{6x^2} - bc^4 \operatorname{PolyLog}\left(2, e^{-2\operatorname{arctanh}(c\sqrt{x})}\right)$$

input `Integrate[(a + b*ArcTanh[c*Sqrt[x]])/(x^3*(1 - c^2*x)),x]`

output `-1/6*(3*a + b*c*Sqrt[x] + 6*a*c^2*x + 9*b*c^3*x^(3/2) - 6*b*c^4*x^2*ArcTanh[c*Sqrt[x]]^2 - 3*b*ArcTanh[c*Sqrt[x]]*(-1 - 2*c^2*x + 3*c^4*x^2 + 4*c^4*x^3*Log[1 - E^(-2*ArcTanh[c*Sqrt[x]])]) - 6*a*c^4*x^2*Log[x] + 6*a*c^4*x^3*Log[1 - c^2*x])/x^2 - b*c^4*PolyLog[2, E^(-2*ArcTanh[c*Sqrt[x]])]`

Rubi [A] (verified)

Time = 1.27 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.18, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$, Rules used = {7267, 2026, 6544, 6452, 264, 264, 219, 6544, 6452, 264, 219, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(1-c^2x)} dx \\ & \quad \downarrow \text{7267} \\ & 2 \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^{5/2} - c^2x^{7/2}} d\sqrt{x} \\ & \quad \downarrow \text{2026} \\ & 2 \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^{5/2}(1-c^2x)} d\sqrt{x} \end{aligned}$$

$$\begin{aligned}
& \downarrow 6544 \\
& 2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(1-c^2x)} d\sqrt{x} + \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{5/2}} d\sqrt{x} \right) \\
& \downarrow 6452 \\
& 2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(1-c^2x)} d\sqrt{x} + \frac{1}{4}bc \int \frac{1}{x^2(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right) \\
& \downarrow 264 \\
& 2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(1-c^2x)} d\sqrt{x} + \frac{1}{4}bc \left(c^2 \int \frac{1}{x(1-c^2x)} d\sqrt{x} - \frac{1}{3x^{3/2}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right) \\
& \downarrow 264 \\
& 2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(1-c^2x)} d\sqrt{x} + \frac{1}{4}bc \left(c^2 \left(c^2 \int \frac{1}{1-c^2x} d\sqrt{x} - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right) \\
& \downarrow 219 \\
& 2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} + \frac{1}{4}bc \left(c^2 \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) \right) \\
& \downarrow 6544 \\
& 2 \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} + \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}} d\sqrt{x} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} + \frac{1}{4}bc \left(c^2 \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) \right) \\
& \downarrow 6452 \\
& 2 \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} + \frac{1}{2}bc \int \frac{1}{x(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{2x} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} + \frac{1}{4}bc \left(c^2 \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) \right) \\
& \downarrow 264 \\
& 2 \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} + \frac{1}{2}bc \left(c^2 \int \frac{1}{1-c^2x} d\sqrt{x} - \frac{1}{\sqrt{x}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{2x} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} + \frac{1}{4}bc \left(c^2 \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) \right) \\
& \downarrow 219
\end{aligned}$$

$$2 \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{2x} + \frac{1}{2}bc \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right)$$

↓ 6550

$$2 \left(c^2 \left(c^2 \left(\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{(\sqrt{xc}+1)\sqrt{x}} d\sqrt{x} + \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{2x} + \frac{1}{2}bc \left(\operatorname{carctanh}(c\sqrt{x}) \right) \right) \right)$$

↓ 6494

$$2 \left(c^2 \left(c^2 \left(-bc \int \frac{\log \left(2 - \frac{2}{\sqrt{xc}+1} \right)}{1-c^2x} d\sqrt{x} + \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} + \log \left(2 - \frac{2}{c\sqrt{x}+1} \right) (a + \operatorname{barctanh}(c\sqrt{x})) \right) \right) \right)$$

↓ 2897

$$2 \left(c^2 \left(c^2 \left(\frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} + \log \left(2 - \frac{2}{c\sqrt{x}+1} \right) (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{2}b \operatorname{PolyLog} \left(2, \frac{2}{\sqrt{xc}+1} - 1 \right) \right) \right) \right)$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])/(x^3*(1 - c^2*x)),x]`

output `2*(-1/4*(a + b*ArcTanh[c*Sqrt[x]])/x^2 + (b*c*(-1/3*1/x^(3/2) + c^2*(-(1/Sqrt[x]) + c*ArcTanh[c*Sqrt[x]])))/4 + c^2*(-1/2*(a + b*ArcTanh[c*Sqrt[x]])/x + (b*c*(-(1/Sqrt[x]) + c*ArcTanh[c*Sqrt[x]]))/2 + c^2*((a + b*ArcTanh[c*Sqrt[x]])^2/(2*b) + (a + b*ArcTanh[c*Sqrt[x]])*Log[2 - 2/(1 + c*Sqrt[x])] - (b*PolyLog[2, -1 + 2/(1 + c*Sqrt[x])])/2))`

Definitions of rubi rules used

rule 219 $\text{Int}[(a + b(x^2)^{-1}), x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))* \text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 264 $\text{Int}[(c(x))^m * (a + b(x^2)^p), x_Symbol] \rightarrow \text{Simp}[(c*x)^{(m+1)} * ((a + b*x^2)^{(p+1}) / (a*c*(m+1))), x] - \text{Simp}[b*(m+2*p+3)/(a*c^{2*(m+1)}) \ \text{Int}[(c*x)^{(m+2)} * (a + b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p, x\} \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 2026 $\text{Int}[(F(x))*(P(x))^p, x_Symbol] \rightarrow \text{With}\{r = \text{Expon}[P, x, \text{Min}]\}, \text{Int}[x^{(p*r)} * \text{ExpandToSum}[P/x^r, x]^p * F, x] /; \text{IGtQ}[r, 0] /; \text{PolyQ}[P, x] \ \&\& \ \text{IntegerQ}[p] \ \&\& \ \text{!MonomialQ}[P, x] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{!PolyQ}[u, x])$

rule 2897 $\text{Int}[\text{Log}[u] * (P(x))^m, x_Symbol] \rightarrow \text{With}\{C = \text{FullSimplify}[P^m * ((1 - u)/D[u, x])]\}, \text{Simp}[C * \text{PolyLog}[2, 1 - u], x] /; \text{FreeQ}[C, x] /; \text{IntegerQ}[m] \ \&\& \ \text{PolyQ}[P, x] \ \&\& \ \text{RationalFunctionQ}[u, x] \ \&\& \ \text{LeQ}[\text{RationalFunctionExponents}[u, x][[2]], \text{Expon}[P, x]]$

rule 6452 $\text{Int}[(a + \text{ArcTanh}[c(x)] * (b))^p * (x)^m, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)} * ((a + b * \text{ArcTanh}[c*x^n])^p / (m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \ \text{Int}[x^{(m+n)} * ((a + b * \text{ArcTanh}[c*x^n])^{(p-1)}) / (1 - c^2*x^{(2*n)}), x], x] /; \text{FreeQ}\{a, b, c, m, n, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 6494 $\text{Int}[(a + \text{ArcTanh}[c(x)] * (b))^p / ((x) * ((d) + (e) * (x))), x_Symbol] \rightarrow \text{Simp}[(a + b * \text{ArcTanh}[c*x])^p * (\text{Log}[2 - 2/(1 + e*(x/d))]/d), x] - \text{Simp}[b*c*(p/d) \ \text{Int}[(a + b * \text{ArcTanh}[c*x])^{(p-1)} * (\text{Log}[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; \text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d^2 - e^2, 0]$

rule 6544

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)*((f_.)*(x_.))^(m_.)/((d_.) + (
e_.)*(x_.^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x
], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x
^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

rule 6550

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)/((x_.)*((d_.) + (e_.)*(x_.^2)),
x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/
d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]
```

rule 7267

```
Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 272 vs. $2(131) = 262$.

Time = 0.10 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.74

method	result
parts	$-a\left(\frac{1}{2x^2} + \frac{c^2}{x} - c^4 \ln(x) + c^4 \ln(c^2x - 1)\right) - 2bc^4\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{4c^4x^2} + \frac{\operatorname{arctanh}(c\sqrt{x})}{2c^2x} - \operatorname{arctan}\right)$
derivativedivides	$-2c^4\left(a\left(\frac{1}{4c^4x^2} + \frac{1}{2c^2x} - \ln(c\sqrt{x}) + \frac{\ln(c\sqrt{x}-1)}{2} + \frac{\ln(1+c\sqrt{x})}{2}\right) + b\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{4c^4x^2} + \frac{\operatorname{arctanh}(c\sqrt{x})}{2c^2x}\right)\right)$
default	$-2c^4\left(a\left(\frac{1}{4c^4x^2} + \frac{1}{2c^2x} - \ln(c\sqrt{x}) + \frac{\ln(c\sqrt{x}-1)}{2} + \frac{\ln(1+c\sqrt{x})}{2}\right) + b\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{4c^4x^2} + \frac{\operatorname{arctanh}(c\sqrt{x})}{2c^2x}\right)\right)$

input

```
int((a+b*arctanh(c*x^(1/2)))/x^3/(-c^2*x+1), x, method=_RETURNVERBOSE)
```

output

```
-a*(1/2/x^2+c^2/x-c^4*ln(x)+c^4*ln(c^2*x-1))-2*b*c^4*(1/4*arctanh(c*x^(1/2)))/c^4/x^2+1/2*arctanh(c*x^(1/2))/c^2/x-arctanh(c*x^(1/2))*ln(c*x^(1/2))+1/2*arctanh(c*x^(1/2))*ln(c*x^(1/2)-1)+1/2*arctanh(c*x^(1/2))*ln(1+c*x^(1/2))+1/2*dilog(c*x^(1/2))+1/2*dilog(1+c*x^(1/2))+1/2*ln(c*x^(1/2))*ln(1+c*x^(1/2))-1/2*dilog(1/2*c*x^(1/2)+1/2)-1/4*ln(c*x^(1/2)-1)*ln(1/2*c*x^(1/2)+1/2)+1/8*ln(c*x^(1/2)-1)^2-1/8*ln(1+c*x^(1/2))^2+1/4*(ln(1+c*x^(1/2))-ln(1/2*c*x^(1/2)+1/2))*ln(-1/2*c*x^(1/2)+1/2)+1/12/c^3/x^(3/2)+3/4/c/x^(1/2)+3/8*ln(c*x^(1/2)-1)-3/8*ln(1+c*x^(1/2))
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(1-c^2x)} dx = \int -\frac{b \operatorname{artanh}(c\sqrt{x}) + a}{(c^2x-1)x^3} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))/x^3/(-c^2*x+1),x, algorithm="fricas")
```

output

```
integral(-(b*arctanh(c*sqrt(x)) + a)/(c^2*x^4 - x^3), x)
```

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(1-c^2x)} dx = -\int \frac{a}{c^2x^4 - x^3} dx - \int \frac{b \operatorname{atanh}(c\sqrt{x})}{c^2x^4 - x^3} dx$$

input

```
integrate((a+b*atanh(c*x**(1/2)))/x**3/(-c**2*x+1),x)
```

output

```
-Integral(a/(c**2*x**4 - x**3), x) - Integral(b*atanh(c*sqrt(x))/(c**2*x**4 - x**3), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 291 vs. $2(130) = 260$.

Time = 0.20 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.85

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(1 - c^2x)} dx$$

$$= - \left(\log(c\sqrt{x} + 1) \log\left(-\frac{1}{2}c\sqrt{x} + \frac{1}{2}\right) + \operatorname{Li}_2\left(\frac{1}{2}c\sqrt{x} + \frac{1}{2}\right) \right) bc^4$$

$$- (\log(c\sqrt{x}) \log(-c\sqrt{x} + 1) + \operatorname{Li}_2(-c\sqrt{x} + 1)) bc^4$$

$$+ (\log(c\sqrt{x} + 1) \log(-c\sqrt{x}) + \operatorname{Li}_2(c\sqrt{x} + 1)) bc^4$$

$$+ \frac{3}{4} bc^4 \log(c\sqrt{x} + 1) - \frac{3}{4} bc^4 \log(c\sqrt{x} - 1)$$

$$- \frac{1}{2} \left(2c^4 \log(c\sqrt{x} + 1) + 2c^4 \log(c\sqrt{x} - 1) - 2c^4 \log(x) + \frac{2c^2x + 1}{x^2} \right) a$$

$$- \frac{3bc^4x^2 \log(c\sqrt{x} + 1)^2 - 3bc^4x^2 \log(-c\sqrt{x} + 1)^2 + 18bc^3x^{\frac{3}{2}} + 2bc\sqrt{x} + 3(2bc^2x + b) \log(c\sqrt{x} + 1)}{12x^2}$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^3/(-c^2*x+1),x, algorithm="maxima")`

output

```

-(log(c*sqrt(x) + 1)*log(-1/2*c*sqrt(x) + 1/2) + dilog(1/2*c*sqrt(x) + 1/2
))*b*c^4 - (log(c*sqrt(x))*log(-c*sqrt(x) + 1) + dilog(-c*sqrt(x) + 1))*b*
c^4 + (log(c*sqrt(x) + 1)*log(-c*sqrt(x)) + dilog(c*sqrt(x) + 1))*b*c^4 +
3/4*b*c^4*log(c*sqrt(x) + 1) - 3/4*b*c^4*log(c*sqrt(x) - 1) - 1/2*(2*c^4*1
og(c*sqrt(x) + 1) + 2*c^4*log(c*sqrt(x) - 1) - 2*c^4*log(x) + (2*c^2*x + 1
)/x^2)*a - 1/12*(3*b*c^4*x^2*log(c*sqrt(x) + 1)^2 - 3*b*c^4*x^2*log(-c*sqr
t(x) + 1)^2 + 18*b*c^3*x^(3/2) + 2*b*c*sqrt(x) + 3*(2*b*c^2*x + b)*log(c*s
qrt(x) + 1) - 3*(2*b*c^4*x^2*log(c*sqrt(x) + 1) + 2*b*c^2*x + b)*log(-c*sqr
t(x) + 1))/x^2

```


Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(1-c^2x)} dx = \int -\frac{b \operatorname{artanh}(c\sqrt{x}) + a}{(c^2x-1)x^3} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^3/(-c^2*x+1),x, algorithm="giac")`

output `integrate(-(b*arctanh(c*sqrt(x)) + a)/((c^2*x - 1)*x^3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(1-c^2x)} dx = - \int \frac{a + b \operatorname{atanh}(c\sqrt{x})}{x^3(c^2x-1)} dx$$

input `int(-(a + b*atanh(c*x^(1/2)))/(x^3*(c^2*x - 1)),x)`

output `-int((a + b*atanh(c*x^(1/2)))/(x^3*(c^2*x - 1)), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(1-c^2x)} dx$$

$$= \frac{9 \operatorname{atanh}(\sqrt{x}c) b c^4 x^2 - 6 \operatorname{atanh}(\sqrt{x}c) b c^2 x - 3 \operatorname{atanh}(\sqrt{x}c) b - 9 \sqrt{x} b c^3 x - \sqrt{x} b c - 6 \left(\int \frac{\operatorname{atanh}(\sqrt{x}c)}{c^2 x^2 - x} dx \right)}{6x^2}$$

input `int((a+b*atanh(c*x^(1/2)))/x^3/(-c^2*x+1),x)`

output `(9*atanh(sqrt(x)*c)*b*c**4*x**2 - 6*atanh(sqrt(x)*c)*b*c**2*x - 3*atanh(sqrt(x)*c)*b - 9*sqrt(x)*b*c**3*x - sqrt(x)*b*c - 6*int(atanh(sqrt(x)*c)/(c**2*x**2 - x),x)*b*c**4*x**2 - 6*log(c**2*x - 1)*a*c**4*x**2 + 6*log(x)*a*c**4*x**2 - 6*a*c**2*x - 3*a)/(6*x**2)`

$$3.44 \quad \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4(1-c^2x)} dx$$

Optimal result	385
Mathematica [A] (verified)	386
Rubi [A] (verified)	386
Maple [A] (verified)	391
Fricas [F]	391
Sympy [F]	392
Maxima [B] (verification not implemented)	392
Giac [F]	393
Mupad [F(-1)]	393
Reduce [F]	394

Optimal result

Integrand size = 26, antiderivative size = 192

$$\begin{aligned} \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4(1-c^2x)} dx = & -\frac{bc}{15x^{5/2}} - \frac{5bc^3}{18x^{3/2}} - \frac{11bc^5}{6\sqrt{x}} + \frac{11}{6}bc^6 \operatorname{arctanh}(c\sqrt{x}) \\ & - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{3x^3} - \frac{c^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{2x^2} \\ & - \frac{c^4(a + b \operatorname{arctanh}(c\sqrt{x}))}{x} + \frac{c^6(a + b \operatorname{arctanh}(c\sqrt{x}))^2}{b} \\ & + 2c^6(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(2 - \frac{2}{1 + c\sqrt{x}}\right) \\ & - bc^6 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + c\sqrt{x}}\right) \end{aligned}$$

output

```
-1/15*b*c/x^(5/2)-5/18*b*c^3/x^(3/2)-11/6*b*c^5/x^(1/2)+11/6*b*c^6*arctanh
(c*x^(1/2))-1/3*(a+b*arctanh(c*x^(1/2)))/x^3-1/2*c^2*(a+b*arctanh(c*x^(1/2)
))/x^2-c^4*(a+b*arctanh(c*x^(1/2)))/x+c^6*(a+b*arctanh(c*x^(1/2)))^2/b+2*
c^6*(a+b*arctanh(c*x^(1/2)))*ln(2-2/(1+c*x^(1/2)))-b*c^6*polylog(2,-1+2/(1
+c*x^(1/2)))
```

Mathematica [A] (verified)

Time = 0.60 (sec) , antiderivative size = 187, normalized size of antiderivative = 0.97

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4(1 - c^2x)} dx =$$

$$\frac{30a + 6bc\sqrt{x} + 45ac^2x + 25bc^3x^{3/2} + 90ac^4x^2 + 165bc^5x^{5/2} - 90bc^6x^3 \operatorname{arctanh}(c\sqrt{x})^2 - 15b \operatorname{arctanh}(c\sqrt{x}) - bc^6 \operatorname{PolyLog}\left(2, e^{-2 \operatorname{arctanh}(c\sqrt{x})}\right)}{x^3}$$

input `Integrate[(a + b*ArcTanh[c*Sqrt[x]])/(x^4*(1 - c^2*x)),x]`

output `-1/90*(30*a + 6*b*c*Sqrt[x] + 45*a*c^2*x + 25*b*c^3*x^(3/2) + 90*a*c^4*x^2 + 165*b*c^5*x^(5/2) - 90*b*c^6*x^3*ArcTanh[c*Sqrt[x]]^2 - 15*b*ArcTanh[c*Sqrt[x]]*(-2 - 3*c^2*x - 6*c^4*x^2 + 11*c^6*x^3 + 12*c^6*x^3*Log[1 - E^(-2*ArcTanh[c*Sqrt[x]])]) - 90*a*c^6*x^3*Log[x] + 90*a*c^6*x^3*Log[1 - c^2*x])/x^3 - b*c^6*PolyLog[2, E^(-2*ArcTanh[c*Sqrt[x]])]`

Rubi [A] (verified)

Time = 1.73 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.36, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.769$, Rules used = {7267, 2026, 6544, 6452, 264, 264, 264, 219, 6544, 6452, 264, 264, 219, 6544, 6452, 264, 219, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4(1 - c^2x)} dx$$

$$\downarrow 7267$$

$$2 \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^{7/2} - c^2x^{9/2}} d\sqrt{x}$$

$$\downarrow 2026$$

$$\begin{aligned}
& 2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{7/2}(1-c^2x)} d\sqrt{x} \\
& \quad \downarrow \text{6544} \\
& 2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{5/2}(1-c^2x)} d\sqrt{x} + \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{7/2}} d\sqrt{x} \right) \\
& \quad \downarrow \text{6452} \\
& 2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{5/2}(1-c^2x)} d\sqrt{x} + \frac{1}{6}bc \int \frac{1}{x^3(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{6x^3} \right) \\
& \quad \downarrow \text{264} \\
& 2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{5/2}(1-c^2x)} d\sqrt{x} + \frac{1}{6}bc \left(c^2 \int \frac{1}{x^2(1-c^2x)} d\sqrt{x} - \frac{1}{5x^{5/2}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{6x^3} \right) \\
& \quad \downarrow \text{264} \\
& 2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{5/2}(1-c^2x)} d\sqrt{x} + \frac{1}{6}bc \left(c^2 \left(c^2 \int \frac{1}{x(1-c^2x)} d\sqrt{x} - \frac{1}{3x^{3/2}} \right) - \frac{1}{5x^{5/2}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{6x^3} \right) \\
& \quad \downarrow \text{264} \\
& 2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{5/2}(1-c^2x)} d\sqrt{x} + \frac{1}{6}bc \left(c^2 \left(c^2 \left(c^2 \int \frac{1}{1-c^2x} d\sqrt{x} - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) - \frac{1}{5x^{5/2}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{6x^3} \right) \\
& \quad \downarrow \text{219} \\
& 2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{5/2}(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{6x^3} + \frac{1}{6}bc \left(c^2 \left(c^2 \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) - \frac{1}{5x^{5/2}} \right) \right) \\
& \quad \downarrow \text{6544} \\
& 2 \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(1-c^2x)} d\sqrt{x} + \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{5/2}} d\sqrt{x} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{6x^3} + \frac{1}{6}bc \left(c^2 \left(c^2 \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) - \frac{1}{5x^{5/2}} \right) \right) \\
& \quad \downarrow \text{6452} \\
& 2 \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(1-c^2x)} d\sqrt{x} + \frac{1}{4}bc \int \frac{1}{x^2(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{6x^3} \right)
\end{aligned}$$

↓ 264

$$2 \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(1-c^2x)} d\sqrt{x} + \frac{1}{4}bc \left(c^2 \int \frac{1}{x(1-c^2x)} d\sqrt{x} - \frac{1}{3x^{3/2}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right)$$

↓ 264

$$2 \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(1-c^2x)} d\sqrt{x} + \frac{1}{4}bc \left(c^2 \left(c^2 \int \frac{1}{1-c^2x} d\sqrt{x} - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right)$$

↓ 219

$$2 \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} + \frac{1}{4}bc \left(c^2 \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right)$$

↓ 6544

$$2 \left(c^2 \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} + \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}} d\sqrt{x} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} + \frac{1}{4}bc \left(c^2 \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right)$$

↓ 6452

$$2 \left(c^2 \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} + \frac{1}{2}bc \int \frac{1}{x(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{2x} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right)$$

↓ 264

$$2 \left(c^2 \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} + \frac{1}{2}bc \left(c^2 \int \frac{1}{1-c^2x} d\sqrt{x} - \frac{1}{\sqrt{x}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{2x} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right)$$

↓ 219

$$2 \left(c^2 \left(c^2 \left(c^2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{2x} + \frac{1}{2}bc \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right)$$

↓ 6550

$$2 \left(c^2 \left(c^2 \left(c^2 \left(\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{(\sqrt{xc}+1)\sqrt{x}} d\sqrt{x} + \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{2x} + \frac{1}{2}bc \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2} \right)$$

↓ 6494

$$2 \left(c^2 \left(c^2 \left(-bc \int \frac{\log \left(2 - \frac{2}{\sqrt{xc+1}} \right)}{1 - c^2x} d\sqrt{x} + \frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} + \log \left(2 - \frac{2}{c\sqrt{x+1}} \right) (a + \operatorname{barctanh}(c\sqrt{x})) \right) \right) \right)$$

↓ 2897

$$2 \left(c^2 \left(c^2 \left(\frac{(a + \operatorname{barctanh}(c\sqrt{x}))^2}{2b} + \log \left(2 - \frac{2}{c\sqrt{x+1}} \right) (a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{2}b \operatorname{PolyLog} \left(2, \frac{2}{\sqrt{xc+1}} \right) \right) \right) \right)$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])/(x^4*(1 - c^2*x)),x]`

output `2*(-1/6*(a + b*ArcTanh[c*Sqrt[x]])/x^3 + (b*c*(-1/5*1/x^(5/2) + c^2*(-1/3*1/x^(3/2) + c^2*(-1/Sqrt[x]) + c*ArcTanh[c*Sqrt[x]])))/6 + c^2*(-1/4*(a + b*ArcTanh[c*Sqrt[x]])/x^2 + (b*c*(-1/3*1/x^(3/2) + c^2*(-1/Sqrt[x]) + c*ArcTanh[c*Sqrt[x]]))/4 + c^2*(-1/2*(a + b*ArcTanh[c*Sqrt[x]])/x + (b*c*(-1/Sqrt[x]) + c*ArcTanh[c*Sqrt[x]]))/2 + c^2*((a + b*ArcTanh[c*Sqrt[x]])^2/(2*b) + (a + b*ArcTanh[c*Sqrt[x]])*Log[2 - 2/(1 + c*Sqrt[x])] - (b*PolyLog[2, -1 + 2/(1 + c*Sqrt[x])])/2))`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 264 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*((m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2897 `Int[Log[u]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6494 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((x_)*((d_) + (e_)*(x_))), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6544 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)*((f_)*(x_)^(m_))/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]`

rule 6550 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((x_)*((d_) + (e_)*(x_)^2)), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]`

rule 7267 `Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Simp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 303, normalized size of antiderivative = 1.58

method	result
parts	$-a\left(\frac{1}{3x^3} + \frac{c^2}{2x^2} + \frac{c^4}{x} - c^6 \ln(x) + c^6 \ln(c^2x - 1)\right) - 2bc^6\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{6c^6x^3} + \frac{\operatorname{arctanh}(c\sqrt{x})}{4c^4x^2} + \dots\right)$
derivativedivides	$-2c^6\left(a\left(\frac{1}{6c^6x^3} + \frac{1}{4c^4x^2} + \frac{1}{2c^2x} - \ln(c\sqrt{x}) + \frac{\ln(c\sqrt{x}-1)}{2} + \frac{\ln(1+c\sqrt{x})}{2}\right) + b\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{6c^6x^3} + \dots\right)\right)$
default	$-2c^6\left(a\left(\frac{1}{6c^6x^3} + \frac{1}{4c^4x^2} + \frac{1}{2c^2x} - \ln(c\sqrt{x}) + \frac{\ln(c\sqrt{x}-1)}{2} + \frac{\ln(1+c\sqrt{x})}{2}\right) + b\left(\frac{\operatorname{arctanh}(c\sqrt{x})}{6c^6x^3} + \dots\right)\right)$

input `int((a+b*arctanh(c*x^(1/2)))/x^4/(-c^2*x+1),x,method=_RETURNVERBOSE)`

output `-a*(1/3/x^3+1/2*c^2/x^2+c^4/x-c^6*ln(x)+c^6*ln(c^2*x-1))-2*b*c^6*(1/6*arctanh(c*x^(1/2))/c^6/x^3+1/4*arctanh(c*x^(1/2))/c^4/x^2+1/2*arctanh(c*x^(1/2))/c^2/x-arctanh(c*x^(1/2))*ln(c*x^(1/2))+1/2*arctanh(c*x^(1/2))*ln(c*x^(1/2)-1)+1/2*arctanh(c*x^(1/2))*ln(1+c*x^(1/2))+1/2*dilog(c*x^(1/2))+1/2*dilog(1+c*x^(1/2))+1/2*ln(c*x^(1/2))*ln(1+c*x^(1/2))-1/2*dilog(1/2*c*x^(1/2)+1/2)-1/4*ln(c*x^(1/2)-1)*ln(1/2*c*x^(1/2)+1/2)+1/8*ln(c*x^(1/2)-1)^2-1/8*ln(1+c*x^(1/2))^2+1/4*(ln(1+c*x^(1/2))-ln(1/2*c*x^(1/2)+1/2))*ln(-1/2*c*x^(1/2)+1/2)+1/30/c^5/x^(5/2)+5/36/c^3/x^(3/2)+11/12/c/x^(1/2)+11/24*ln(c*x^(1/2)-1)-11/24*ln(1+c*x^(1/2)))`

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4(1 - c^2x)} dx = \int -\frac{b \operatorname{arctanh}(c\sqrt{x}) + a}{(c^2x - 1)x^4} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^4/(-c^2*x+1),x, algorithm="fricas")`

output `integral(-(b*arctanh(c*sqrt(x)) + a)/(c^2*x^5 - x^4), x)`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4(1-c^2x)} dx = - \int \frac{a}{c^2x^5 - x^4} dx - \int \frac{b \operatorname{atanh}(c\sqrt{x})}{c^2x^5 - x^4} dx$$

input `integrate((a+b*atanh(c*x**(1/2)))/x**4/(-c**2*x+1),x)`

output `-Integral(a/(c**2*x**5 - x**4), x) - Integral(b*atanh(c*sqrt(x))/(c**2*x**5 - x**4), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 330 vs. $2(157) = 314$.

Time = 0.20 (sec) , antiderivative size = 330, normalized size of antiderivative = 1.72

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4(1-c^2x)} dx \\ &= - \left(\log(c\sqrt{x} + 1) \log\left(-\frac{1}{2}c\sqrt{x} + \frac{1}{2}\right) + \operatorname{Li}_2\left(\frac{1}{2}c\sqrt{x} + \frac{1}{2}\right) \right) bc^6 \\ & \quad - \left(\log(c\sqrt{x}) \log(-c\sqrt{x} + 1) + \operatorname{Li}_2(-c\sqrt{x} + 1) \right) bc^6 \\ & \quad + \left(\log(c\sqrt{x} + 1) \log(-c\sqrt{x}) + \operatorname{Li}_2(c\sqrt{x} + 1) \right) bc^6 \\ & \quad + \frac{11}{12} bc^6 \log(c\sqrt{x} + 1) - \frac{11}{12} bc^6 \log(c\sqrt{x} - 1) \\ & \quad - \frac{1}{6} \left(6c^6 \log(c\sqrt{x} + 1) + 6c^6 \log(c\sqrt{x} - 1) - 6c^6 \log(x) + \frac{6c^4x^2 + 3c^2x + 2}{x^3} \right) a \\ & \quad - \frac{45bc^6x^3 \log(c\sqrt{x} + 1)^2 - 45bc^6x^3 \log(-c\sqrt{x} + 1)^2 + 330bc^5x^{\frac{5}{2}} + 50bc^3x^{\frac{3}{2}} + 12bc\sqrt{x} + 15(6bc^4x^2 - 180x)}{180x^3} \end{aligned}$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^4/(-c^2*x+1),x, algorithm="maxima")`

output

```

-(log(c*sqrt(x) + 1)*log(-1/2*c*sqrt(x) + 1/2) + dilog(1/2*c*sqrt(x) + 1/2
))*b*c^6 - (log(c*sqrt(x))*log(-c*sqrt(x) + 1) + dilog(-c*sqrt(x) + 1))*b*
c^6 + (log(c*sqrt(x) + 1)*log(-c*sqrt(x)) + dilog(c*sqrt(x) + 1))*b*c^6 +
11/12*b*c^6*log(c*sqrt(x) + 1) - 11/12*b*c^6*log(c*sqrt(x) - 1) - 1/6*(6*c
^6*log(c*sqrt(x) + 1) + 6*c^6*log(c*sqrt(x) - 1) - 6*c^6*log(x) + (6*c^4*x
^2 + 3*c^2*x + 2)/x^3)*a - 1/180*(45*b*c^6*x^3*log(c*sqrt(x) + 1)^2 - 45*b
*c^6*x^3*log(-c*sqrt(x) + 1)^2 + 330*b*c^5*x^(5/2) + 50*b*c^3*x^(3/2) + 12
*b*c*sqrt(x) + 15*(6*b*c^4*x^2 + 3*b*c^2*x + 2*b)*log(c*sqrt(x) + 1) - 15*
(6*b*c^6*x^3*log(c*sqrt(x) + 1) + 6*b*c^4*x^2 + 3*b*c^2*x + 2*b)*log(-c*sq
rt(x) + 1))/x^3

```

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4(1 - c^2x)} dx = \int -\frac{b \operatorname{artanh}(c\sqrt{x}) + a}{(c^2x - 1)x^4} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))/x^4/(-c^2*x+1),x, algorithm="giac")
```

output

```
integrate(-(b*arctanh(c*sqrt(x)) + a)/((c^2*x - 1)*x^4), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4(1 - c^2x)} dx = -\int \frac{a + b \operatorname{atanh}(c\sqrt{x})}{x^4(c^2x - 1)} dx$$

input

```
int(-(a + b*atanh(c*x^(1/2)))/(x^4*(c^2*x - 1)),x)
```

output

```
-int((a + b*atanh(c*x^(1/2)))/(x^4*(c^2*x - 1)), x)
```

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^4(1 - c^2x)} dx$$

$$= \frac{165 \operatorname{atanh}(\sqrt{x}c) b c^6 x^3 - 90 \operatorname{atanh}(\sqrt{x}c) b c^4 x^2 - 45 \operatorname{atanh}(\sqrt{x}c) b c^2 x - 30 \operatorname{atanh}(\sqrt{x}c) b - 165 \sqrt{x} b c^5}{90 x^3}$$

input `int((a+b*atanh(c*x^(1/2)))/x^4/(-c^2*x+1),x)`

output `(165*atanh(sqrt(x)*c)*b*c**6*x**3 - 90*atanh(sqrt(x)*c)*b*c**4*x**2 - 45*atanh(sqrt(x)*c)*b*c**2*x - 30*atanh(sqrt(x)*c)*b - 165*sqrt(x)*b*c**5*x**2 - 25*sqrt(x)*b*c**3*x - 6*sqrt(x)*b*c - 90*int(atanh(sqrt(x)*c)/(c**2*x**2 - x),x)*b*c**6*x**3 - 90*log(c**2*x - 1)*a*c**6*x**3 + 90*log(x)*a*c**6*x**3 - 90*a*c**4*x**2 - 45*a*c**2*x - 30*a)/(90*x**3)`

3.45 $\int \frac{x^2 (a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx$

Optimal result	395
Mathematica [C] (warning: unable to verify)	396
Rubi [A] (verified)	397
Maple [A] (verified)	401
Fricas [F]	403
Sympy [F(-1)]	403
Maxima [F]	403
Giac [F]	404
Mupad [F(-1)]	404
Reduce [F]	404

Optimal result

Integrand size = 23, antiderivative size = 460

$$\int \frac{x^2 (a + \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx = -\frac{bd\sqrt{x}}{ce^2} + \frac{b\sqrt{x}}{2c^3e} + \frac{bx^{3/2}}{6ce} + \frac{b\operatorname{arctanh}(c\sqrt{x})}{c^2e^2}$$

$$- \frac{\operatorname{arctanh}(c\sqrt{x})}{2c^4e} - \frac{dx(a + \operatorname{arctanh}(c\sqrt{x}))}{e^2}$$

$$+ \frac{x^2(a + \operatorname{arctanh}(c\sqrt{x}))}{2e}$$

$$- \frac{2d^2(a + \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1+c\sqrt{x}}\right)}{e^3}$$

$$+ \frac{d^2(a + \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2c(\sqrt{-d}-\sqrt{e}\sqrt{x})}{(c\sqrt{-d}-\sqrt{e})(1+c\sqrt{x})}\right)}{e^3}$$

$$+ \frac{d^2(a + \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2c(\sqrt{-d}+\sqrt{e}\sqrt{x})}{(c\sqrt{-d}+\sqrt{e})(1+c\sqrt{x})}\right)}{e^3}$$

$$+ \frac{bd^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c\sqrt{x}}\right)}{e^3}$$

$$- \frac{bd^2 \operatorname{PolyLog}\left(2, 1 - \frac{2c(\sqrt{-d}-\sqrt{e}\sqrt{x})}{(c\sqrt{-d}-\sqrt{e})(1+c\sqrt{x})}\right)}{2e^3}$$

$$- \frac{bd^2 \operatorname{PolyLog}\left(2, 1 - \frac{2c(\sqrt{-d}+\sqrt{e}\sqrt{x})}{(c\sqrt{-d}+\sqrt{e})(1+c\sqrt{x})}\right)}{2e^3}$$

output

```

-b*d*x^(1/2)/c/e^2+1/2*b*x^(1/2)/c^3/e+1/6*b*x^(3/2)/c/e+b*d*arctanh(c*x^(
1/2))/c^2/e^2-1/2*b*arctanh(c*x^(1/2))/c^4/e-d*x*(a+b*arctanh(c*x^(1/2)))/
e^2+1/2*x^2*(a+b*arctanh(c*x^(1/2)))/e-2*d^2*(a+b*arctanh(c*x^(1/2)))*ln(2
/(1+c*x^(1/2)))/e^3+d^2*(a+b*arctanh(c*x^(1/2)))*ln(2*c*((-d)^(1/2)-e^(1/2
)*x^(1/2))/(c*(-d)^(1/2)-e^(1/2))/(1+c*x^(1/2)))/e^3+d^2*(a+b*arctanh(c*x^
(1/2)))*ln(2*c*((-d)^(1/2)+e^(1/2)*x^(1/2))/(c*(-d)^(1/2)+e^(1/2))/(1+c*x^
(1/2)))/e^3+b*d^2*polylog(2,1-2/(1+c*x^(1/2)))/e^3-1/2*b*d^2*polylog(2,1-2
*c*((-d)^(1/2)-e^(1/2)*x^(1/2))/(c*(-d)^(1/2)-e^(1/2))/(1+c*x^(1/2)))/e^3-
1/2*b*d^2*polylog(2,1-2*c*((-d)^(1/2)+e^(1/2)*x^(1/2))/(c*(-d)^(1/2)+e^(1/
2))/(1+c*x^(1/2)))/e^3

```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 1.86 (sec) , antiderivative size = 558, normalized size of antiderivative = 1.21

$$\int \frac{x^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx$$

$$= \frac{-6adex + 3ae^2x^2 + 6ad^2 \log(d + ex) + b \left(2ce(-3c^2d+2e)\sqrt{x} + ce^2\sqrt{x}(-1+c^2x) - 6(c^2d-e)e(-1+c^2x)\operatorname{arctanh}(c\sqrt{x}) + 3e^2 \right)}{e^3}$$

input

```
Integrate[(x^2*(a + b*ArcTanh[c*Sqrt[x]]))/(d + e*x),x]
```

output

```
(-6*a*d*e*x + 3*a*e^2*x^2 + 6*a*d^2*Log[d + e*x] + (b*(2*c*e*(-3*c^2*d + 2
*e)*Sqrt[x] + c*e^2*Sqrt[x]*(-1 + c^2*x) - 6*(c^2*d - e)*e*(-1 + c^2*x)*Ar
cTanh[c*Sqrt[x]] + 3*e^2*(-1 + c^2*x)^2*ArcTanh[c*Sqrt[x]] - 6*c^4*d^2*(Ar
cTanh[c*Sqrt[x]]*(ArcTanh[c*Sqrt[x]] + 2*Log[1 + E^(-2*ArcTanh[c*Sqrt[x]])
]) - PolyLog[2, -E^(-2*ArcTanh[c*Sqrt[x]])]) + 3*c^4*d^2*(2*ArcTanh[c*Sqrt
[x]]^2 - (4*I)*ArcSin[Sqrt[(c^2*d)/(c^2*d + e)]]*ArcTanh[(c*e*Sqrt[x])/Sqr
t[-(c^2*d*e)]] + 2*((-I)*ArcSin[Sqrt[(c^2*d)/(c^2*d + e)]] + ArcTanh[c*Sqr
t[x]])*Log[(-2*Sqrt[-(c^2*d*e)]] + e*(-1 + E^(2*ArcTanh[c*Sqrt[x]]))) + c^2*
d*(1 + E^(2*ArcTanh[c*Sqrt[x]])))/((c^2*d + e)*E^(2*ArcTanh[c*Sqrt[x]]))
+ 2*(I*ArcSin[Sqrt[(c^2*d)/(c^2*d + e)]] + ArcTanh[c*Sqrt[x]])*Log[(2*Sqrt
[-(c^2*d*e)]] + e*(-1 + E^(2*ArcTanh[c*Sqrt[x]])) + c^2*d*(1 + E^(2*ArcTanh
[c*Sqrt[x]])))/((c^2*d + e)*E^(2*ArcTanh[c*Sqrt[x]])) - PolyLog[2, -(c^2
*d) + e - 2*Sqrt[-(c^2*d*e)]]/((c^2*d + e)*E^(2*ArcTanh[c*Sqrt[x]])) - Po
lyLog[2, -(c^2*d) + e + 2*Sqrt[-(c^2*d*e)]]/((c^2*d + e)*E^(2*ArcTanh[c*S
qrt[x]])))]/c^4)/(6*e^3)
```

Rubi [A] (verified)

Time = 1.46 (sec) , antiderivative size = 459, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.478$, Rules used = {7267, 6542, 6452, 254, 2009, 6542, 6452, 262, 219, 6606, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2(a + \operatorname{barctanh}(c\sqrt{x}))}{d + ex} dx \\
 & \quad \downarrow \text{7267} \\
 & 2 \int \frac{x^{5/2}(a + \operatorname{barctanh}(c\sqrt{x}))}{d + ex} d\sqrt{x} \\
 & \quad \downarrow \text{6542} \\
 & 2 \left(\frac{\int x^{3/2}(a + \operatorname{barctanh}(c\sqrt{x})) d\sqrt{x}}{e} - \frac{d \int \frac{x^{3/2}(a + \operatorname{barctanh}(c\sqrt{x}))}{d + ex} d\sqrt{x}}{e} \right) \\
 & \quad \downarrow \text{6452}
 \end{aligned}$$

$$2 \left(\frac{\frac{1}{4}x^2(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{4}bc \int \frac{x^2}{1-c^2x} d\sqrt{x}}{e} - \frac{d \int \frac{x^{3/2}(a + \operatorname{barctanh}(c\sqrt{x}))}{d+ex} d\sqrt{x}}{e} \right)$$

↓ 254

$$2 \left(\frac{\frac{1}{4}x^2(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{4}bc \int \left(-\frac{x}{c^2} + \frac{1}{c^4(1-c^2x)} - \frac{1}{c^4} \right) d\sqrt{x}}{e} - \frac{d \int \frac{x^{3/2}(a + \operatorname{barctanh}(c\sqrt{x}))}{d+ex} d\sqrt{x}}{e} \right)$$

↓ 2009

$$2 \left(\frac{\frac{1}{4}x^2(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^5} - \frac{\sqrt{x}}{c^4} - \frac{x^{3/2}}{3c^2} \right)}{e} - \frac{d \int \frac{x^{3/2}(a + \operatorname{barctanh}(c\sqrt{x}))}{d+ex} d\sqrt{x}}{e} \right)$$

↓ 6542

$$2 \left(\frac{\frac{1}{4}x^2(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^5} - \frac{\sqrt{x}}{c^4} - \frac{x^{3/2}}{3c^2} \right)}{e} - \frac{d \left(\frac{\int \sqrt{x}(a + \operatorname{barctanh}(c\sqrt{x})) d\sqrt{x}}{e} - \frac{d \int \frac{\sqrt{x}(a + \operatorname{barctanh}(c\sqrt{x}))}{d+ex} d\sqrt{x}}{e} \right)}{e} \right)$$

↓ 6452

$$2 \left(\frac{\frac{1}{4}x^2(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^5} - \frac{\sqrt{x}}{c^4} - \frac{x^{3/2}}{3c^2} \right)}{e} - \frac{d \left(\frac{\frac{1}{2}x(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{2}bc \int \frac{x}{1-c^2x} d\sqrt{x}}{e} - \frac{d \int \frac{\sqrt{x}(a + \operatorname{barctanh}(c\sqrt{x}))}{d+ex} d\sqrt{x}}{e} \right)}{e} \right)$$

↓ 262

$$2 \left(\frac{\frac{1}{4}x^2(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^5} - \frac{\sqrt{x}}{c^4} - \frac{x^{3/2}}{3c^2} \right)}{e} \right) - d \left(\frac{\frac{1}{2}x(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\int \frac{1}{1-c^2x} d\sqrt{x} - \frac{\sqrt{x}}{c^2}}{c^3} \right)}{e} \right)$$

↓ 219

$$2 \left(\frac{\frac{1}{4}x^2(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^5} - \frac{\sqrt{x}}{c^4} - \frac{x^{3/2}}{3c^2} \right)}{e} \right) - d \left(\frac{\frac{1}{2}x(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} \right)}{e} \right)$$

↓ 6606

$$2 \left(\frac{\frac{1}{4}x^2(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^5} - \frac{\sqrt{x}}{c^4} - \frac{x^{3/2}}{3c^2} \right)}{e} \right) - d \left(\frac{\frac{1}{2}x(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} \right)}{e} \right)$$

↓ 2009

$$2 \left(\frac{\frac{1}{4}x^2(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{4}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^5} - \frac{\sqrt{x}}{c^4} - \frac{x^{3/2}}{3c^2} \right)}{e} \right) - d \left(\frac{\frac{1}{2}x(a + \operatorname{barctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} \right)}{e} \right)$$

input `Int[(x^2*(a + b*ArcTanh[c*Sqrt[x]])/(d + e*x), x]`

output

```

2*(((x^2*(a + b*ArcTanh[c*Sqrt[x]]))/4 - (b*c*(-(Sqrt[x]/c^4) - x^(3/2)/(3
*c^2) + ArcTanh[c*Sqrt[x]]/c^5))/4)/e - (d*(((x*(a + b*ArcTanh[c*Sqrt[x]]
)/2 - (b*c*(-(Sqrt[x]/c^2) + ArcTanh[c*Sqrt[x]]/c^3))/2)/e - (d*(-((a + b
*ArcTanh[c*Sqrt[x]])*Log[2/(1 + c*Sqrt[x]))]/e) + ((a + b*ArcTanh[c*Sqrt[x
]])*Log[(2*c*(Sqrt[-d] - Sqrt[e]*Sqrt[x]))/(c*Sqrt[-d] - Sqrt[e]*(1 + c*
Sqrt[x]))]/(2*e) + ((a + b*ArcTanh[c*Sqrt[x]])*Log[(2*c*(Sqrt[-d] + Sqrt[
e]*Sqrt[x]))/(c*Sqrt[-d] + Sqrt[e]*(1 + c*Sqrt[x]))]/(2*e) + (b*PolyLog
[2, 1 - 2/(1 + c*Sqrt[x]))]/(2*e) - (b*PolyLog[2, 1 - (2*c*(Sqrt[-d] - Sqr
t[e]*Sqrt[x]))/(c*Sqrt[-d] - Sqrt[e]*(1 + c*Sqrt[x]))]/(4*e) - (b*PolyL
og[2, 1 - (2*c*(Sqrt[-d] + Sqrt[e]*Sqrt[x]))/(c*Sqrt[-d] + Sqrt[e]*(1 +
c*Sqrt[x]))]/(4*e))/e)/e)

```

Defintions of rubi rules used

rule 219

```

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])

```

rule 254

```

Int[(x_)^(m_)/((a_) + (b_)*(x_)^2), x_Symbol] := Int[PolynomialDivide[x^m,
a + b*x^2, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 3]

```

rule 262

```

Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)
^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/
(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b
, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c
, 2, m, p, x]

```

rule 2009

```

Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]

```

rule 6452

```

Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] :=
Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1
] && IntegerQ[m])) && NeQ[m, -1]

```

rule 6542

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^ (p_.)*((f_.)*(x_))^(m_)/((d_) + (
e_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*
x])^p, x], x] - Simp[d*(f^2/e Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/
(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m,
1]
```

rule 6606

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^ (p_.)*((f_.)*(x_))^(m_.)*((d_) + (e
_.)*(x_)^2)^(q_.), x_Symbol] := With[{u = ExpandIntegrand[(a + b*ArcTanh[c*
x])^p, (f*x)^m*(d + e*x^2)^q, x]}, Int[u, x] /; SumQ[u] /; FreeQ[{a, b, c,
d, e, f, m}, x] && IntegerQ[q] && IGtQ[p, 0] && (GtQ[q, 0] || IntegerQ[m])
```

rule 7267

```
Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]
```

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 564, normalized size of antiderivative = 1.23

method	result
parts	$\frac{ax^2}{2e} - \frac{adx}{e^2} + \frac{ad^2 \ln(ex+d)}{e^3} + \left(-\frac{c^6 \operatorname{arctanh}(c\sqrt{x})xd}{2e^2} + \frac{c^6 \operatorname{arctanh}(c\sqrt{x})x^2}{4e} + \frac{c^6 \operatorname{arctanh}(c\sqrt{x})d^2 \ln(c^2 ex+c^2 d)}{2e^3} - \dots \right)$
derivativedivides	$-\frac{ac^6 dx}{e^2} + \frac{ac^6 x^2}{2e} + \frac{ac^6 d^2 \ln(c^2 ex+c^2 d)}{e^3} + 2bc^2 \left(-\frac{\operatorname{arctanh}(c\sqrt{x})c^4 dx}{2e^2} + \frac{\operatorname{arctanh}(c\sqrt{x})c^4 x^2}{4e} + \frac{\operatorname{arctanh}(c\sqrt{x})c^4 d^2 \ln(c^2 ex+c^2 d)}{2e^3} \right)$
default	$-\frac{ac^6 dx}{e^2} + \frac{ac^6 x^2}{2e} + \frac{ac^6 d^2 \ln(c^2 ex+c^2 d)}{e^3} + 2bc^2 \left(-\frac{\operatorname{arctanh}(c\sqrt{x})c^4 dx}{2e^2} + \frac{\operatorname{arctanh}(c\sqrt{x})c^4 x^2}{4e} + \frac{\operatorname{arctanh}(c\sqrt{x})c^4 d^2 \ln(c^2 ex+c^2 d)}{2e^3} \right)$

input

```
int(x^2*(a+b*arctanh(c*x^(1/2)))/(e*x+d), x, method=_RETURNVERBOSE)
```

output

```
1/2*a/e*x^2-a/e^2*d*x+a*d^2/e^3*ln(e*x+d)+2*b/c^6*(-1/2*c^6*arctanh(c*x^(1/2))/e^2*x*d+1/4*c^6*arctanh(c*x^(1/2))/e*x^2+1/2*c^6*arctanh(c*x^(1/2))*d^2/e^3*ln(c^2*e*x+c^2*d)-1/2*c^2*(-1/2/e^2*(1/3*e*c^3*x^(3/2)-2*c^3*d*x^(1/2)+e*c*x^(1/2)+1/2*(-2*c^2*d+e)*ln(c*x^(1/2)-1)-1/2*(-2*c^2*d+e)*ln(1+c*x^(1/2)))-c^4*d^2/e^3*(1/2*ln(c*x^(1/2)-1)*ln(c^2*e*x+c^2*d)-e*(1/2*ln(c*x^(1/2)-1)*(ln((c*(-d*e)^(1/2)-e*(c*x^(1/2)-1)-e)/(c*(-d*e)^(1/2)-e))+ln((c*(-d*e)^(1/2)+e*(c*x^(1/2)-1)+e)/(c*(-d*e)^(1/2)+e)))/e+1/2*(dilog((c*(-d*e)^(1/2)-e*(c*x^(1/2)-1)-e)/(c*(-d*e)^(1/2)-e))+dilog((c*(-d*e)^(1/2)+e*(c*x^(1/2)-1)+e)/(c*(-d*e)^(1/2)+e)))/e)-1/2*ln(1+c*x^(1/2))*ln(c^2*e*x+c^2*d)+e*(1/2*ln(1+c*x^(1/2))*(ln((c*(-d*e)^(1/2)-e*(1+c*x^(1/2)))+e)/(c*(-d*e)^(1/2)+e))+ln((c*(-d*e)^(1/2)+e*(1+c*x^(1/2))-e)/(c*(-d*e)^(1/2)-e)))/e+1/2*(dilog((c*(-d*e)^(1/2)-e*(1+c*x^(1/2)))+e)/(c*(-d*e)^(1/2)+e))+dilog((c*(-d*e)^(1/2)+e*(1+c*x^(1/2))-e)/(c*(-d*e)^(1/2)-e)))/e))))
```

Fricas [F]

$$\int \frac{x^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx = \int \frac{(b \operatorname{arctanh}(c\sqrt{x}) + a)x^2}{ex + d} dx$$

input `integrate(x^2*(a+b*arctanh(c*x^(1/2)))/(e*x+d),x, algorithm="fricas")`

output `integral((b*x^2*arctanh(c*sqrt(x)) + a*x^2)/(e*x + d), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{x^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx = \text{Timed out}$$

input `integrate(x**2*(a+b*atanh(c*x**(1/2)))/(e*x+d),x)`

output `Timed out`

Maxima [F]

$$\int \frac{x^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx = \int \frac{(b \operatorname{arctanh}(c\sqrt{x}) + a)x^2}{ex + d} dx$$

input `integrate(x^2*(a+b*arctanh(c*x^(1/2)))/(e*x+d),x, algorithm="maxima")`

output `1/2*a*(2*d^2*log(e*x + d)/e^3 + (e*x^2 - 2*d*x)/e^2) + b*integrate(1/2*x^2*log(c*sqrt(x) + 1)/(e*x + d), x) - b*integrate(1/2*x^2*log(-c*sqrt(x) + 1)/(e*x + d), x)`

Giac [F]

$$\int \frac{x^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx = \int \frac{(b \operatorname{artanh}(c\sqrt{x}) + a)x^2}{ex + d} dx$$

input `integrate(x^2*(a+b*arctanh(c*x^(1/2)))/(e*x+d),x, algorithm="giac")`

output `integrate((b*arctanh(c*sqrt(x)) + a)*x^2/(e*x + d), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx = \int \frac{x^2(a + b \operatorname{atanh}(c\sqrt{x}))}{d + ex} dx$$

input `int((x^2*(a + b*atanh(c*x^(1/2))))/(d + e*x),x)`

output `int((x^2*(a + b*atanh(c*x^(1/2))))/(d + e*x), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{x^2(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx \\ &= \frac{2 \left(\int \frac{\operatorname{atanh}(\sqrt{x}c)x^2}{ex+d} dx \right) b e^3 + 2 \log(ex + d) a d^2 - 2 a d e x + a e^2 x^2}{2e^3} \end{aligned}$$

input `int(x^2*(a+b*atanh(c*x^(1/2)))/(e*x+d),x)`

output `(2*int((atanh(sqrt(x)*c)*x**2)/(d + e*x),x)*b*e**3 + 2*log(d + e*x)*a*d**2 - 2*a*d*e*x + a*e**2*x**2)/(2*e**3)`

3.46 $\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx$

Optimal result	405
Mathematica [C] (verified)	406
Rubi [A] (verified)	407
Maple [A] (verified)	409
Fricas [F]	411
Sympy [F]	411
Maxima [F]	411
Giac [F]	412
Mupad [F(-1)]	412
Reduce [F]	412

Optimal result

Integrand size = 21, antiderivative size = 374

$$\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx = \frac{b\sqrt{x}}{ce} - \frac{b \operatorname{arctanh}(c\sqrt{x})}{c^2e} + \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{e}$$

$$+ \frac{2d(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1+c\sqrt{x}}\right)}{e^2}$$

$$- \frac{d(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2c(\sqrt{-d}-\sqrt{e}\sqrt{x})}{(c\sqrt{-d}-\sqrt{e})(1+c\sqrt{x})}\right)}{e^2}$$

$$- \frac{d(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2c(\sqrt{-d}+\sqrt{e}\sqrt{x})}{(c\sqrt{-d}+\sqrt{e})(1+c\sqrt{x})}\right)}{e^2}$$

$$- \frac{bd \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c\sqrt{x}}\right)}{e^2}$$

$$+ \frac{bd \operatorname{PolyLog}\left(2, 1 - \frac{2c(\sqrt{-d}-\sqrt{e}\sqrt{x})}{(c\sqrt{-d}-\sqrt{e})(1+c\sqrt{x})}\right)}{2e^2}$$

$$+ \frac{bd \operatorname{PolyLog}\left(2, 1 - \frac{2c(\sqrt{-d}+\sqrt{e}\sqrt{x})}{(c\sqrt{-d}+\sqrt{e})(1+c\sqrt{x})}\right)}{2e^2}$$

output

```

b*x^(1/2)/c/e-b*arctanh(c*x^(1/2))/c^2/e+x*(a+b*arctanh(c*x^(1/2)))/e+2*d*
(a+b*arctanh(c*x^(1/2)))*ln(2/(1+c*x^(1/2)))/e^2-d*(a+b*arctanh(c*x^(1/2))
)*ln(2*c*((-d)^(1/2)-e^(1/2)*x^(1/2))/(c*(-d)^(1/2)-e^(1/2))/(1+c*x^(1/2))
)/e^2-d*(a+b*arctanh(c*x^(1/2)))*ln(2*c*((-d)^(1/2)+e^(1/2)*x^(1/2))/(c*(-
d)^(1/2)+e^(1/2))/(1+c*x^(1/2)))/e^2-b*d*polylog(2,1-2/(1+c*x^(1/2)))/e^2+
1/2*b*d*polylog(2,1-2*c*((-d)^(1/2)-e^(1/2)*x^(1/2))/(c*(-d)^(1/2)-e^(1/2)
)/(1+c*x^(1/2)))/e^2+1/2*b*d*polylog(2,1-2*c*((-d)^(1/2)+e^(1/2)*x^(1/2))/
(c*(-d)^(1/2)+e^(1/2))/(1+c*x^(1/2)))/e^2

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.65 (sec) , antiderivative size = 463, normalized size of antiderivative = 1.24

$$\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx$$

$$= \frac{\frac{2be\sqrt{x}}{c} + 2aex + \frac{2be(-1+c^2x)\operatorname{arctanh}(c\sqrt{x})}{c^2} + 4ibd \arcsin\left(\sqrt{\frac{c^2d}{c^2d+e}}\right) \operatorname{arctanh}\left(\frac{ce\sqrt{x}}{\sqrt{-c^2de}}\right) + 4bd \operatorname{arctanh}(c\sqrt{x}) \operatorname{arctanh}\left(\frac{ce\sqrt{x}}{\sqrt{-c^2de}}\right)}{e^2}$$

input

```
Integrate[(x*(a + b*ArcTanh[c*Sqrt[x]]))/(d + e*x),x]
```

output

```

((2*b*e*Sqrt[x])/c + 2*a*e*x + (2*b*e*(-1 + c^2*x)*ArcTanh[c*Sqrt[x]])/c^2
+ (4*I)*b*d*ArcSin[Sqrt[(c^2*d)/(c^2*d + e)]]*ArcTanh[(c*e*Sqrt[x])/Sqrt[
-(c^2*d*e)]] + 4*b*d*ArcTanh[c*Sqrt[x]]*Log[1 + E^(-2*ArcTanh[c*Sqrt[x]])]
+ (2*I)*b*d*(ArcSin[Sqrt[(c^2*d)/(c^2*d + e)]] + I*ArcTanh[c*Sqrt[x]])*Lo
g[(-2*Sqrt[-(c^2*d*e)] + e*(-1 + E^(2*ArcTanh[c*Sqrt[x]])) + c^2*d*(1 + E^
(2*ArcTanh[c*Sqrt[x]])))/((c^2*d + e)*E^(2*ArcTanh[c*Sqrt[x]]))] - 2*b*d*(
I*ArcSin[Sqrt[(c^2*d)/(c^2*d + e)]] + ArcTanh[c*Sqrt[x]])*Log[(2*Sqrt[-(c^
2*d*e)] + e*(-1 + E^(2*ArcTanh[c*Sqrt[x]])) + c^2*d*(1 + E^(2*ArcTanh[c*Sq
rt[x]])))/((c^2*d + e)*E^(2*ArcTanh[c*Sqrt[x]]))] - 2*a*d*Log[d + e*x] - 2
*b*d*PolyLog[2, -E^(-2*ArcTanh[c*Sqrt[x]])] + b*d*PolyLog[2, -(c^2*d) + e
- 2*Sqrt[-(c^2*d*e)]]/((c^2*d + e)*E^(2*ArcTanh[c*Sqrt[x]]))] + b*d*PolyL
og[2, -(c^2*d) + e + 2*Sqrt[-(c^2*d*e)]]/((c^2*d + e)*E^(2*ArcTanh[c*Sqrt
[x]])))]/(2*e^2)

```

Rubi [A] (verified)

Time = 0.92 (sec) , antiderivative size = 387, normalized size of antiderivative = 1.03, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {7267, 6542, 6452, 262, 219, 6606, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx \\
 & \quad \downarrow \text{7267} \\
 & 2 \int \frac{x^{3/2}(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} d\sqrt{x} \\
 & \quad \downarrow \text{6542} \\
 & 2 \left(\frac{\int \sqrt{x}(a + b \operatorname{arctanh}(c\sqrt{x})) d\sqrt{x}}{e} - \frac{d \int \frac{\sqrt{x}(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} d\sqrt{x}}{e} \right) \\
 & \quad \downarrow \text{6452} \\
 & 2 \left(\frac{\frac{1}{2}x(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \int \frac{x}{1 - c^2x} d\sqrt{x}}{e} - \frac{d \int \frac{\sqrt{x}(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} d\sqrt{x}}{e} \right) \\
 & \quad \downarrow \text{262} \\
 & 2 \left(\frac{\frac{1}{2}x(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\int \frac{1}{1 - c^2x} d\sqrt{x}}{c^2} - \frac{\sqrt{x}}{c^2} \right)}{e} - \frac{d \int \frac{\sqrt{x}(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} d\sqrt{x}}{e} \right) \\
 & \quad \downarrow \text{219} \\
 & 2 \left(\frac{\frac{1}{2}x(a + b \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2} \right)}{e} - \frac{d \int \frac{\sqrt{x}(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} d\sqrt{x}}{e} \right) \\
 & \quad \downarrow \text{6606}
 \end{aligned}$$

$$2 \left(\frac{\frac{1}{2}x(a + \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2} \right)}{e} - \frac{d \int \left(\frac{a + \operatorname{arctanh}(c\sqrt{x})}{2\sqrt{e}(\sqrt{-d} + \sqrt{e}\sqrt{x})} - \frac{a + \operatorname{arctanh}(c\sqrt{x})}{2\sqrt{e}(\sqrt{-d} - \sqrt{e}\sqrt{x})} \right) d\sqrt{x}}{e} \right)$$

↓ 2009

$$2 \left(\frac{\frac{1}{2}x(a + \operatorname{arctanh}(c\sqrt{x})) - \frac{1}{2}bc \left(\frac{\operatorname{arctanh}(c\sqrt{x})}{c^3} - \frac{\sqrt{x}}{c^2} \right)}{e} - \frac{d \left(\frac{(a + \operatorname{arctanh}(c\sqrt{x})) \log \left(\frac{2c(\sqrt{-d} - \sqrt{e}\sqrt{x})}{(c\sqrt{x} + 1)(c\sqrt{-d} - \sqrt{e})} \right)}{2e} + \frac{(a + \operatorname{arctanh}(c\sqrt{x})) \log \left(\frac{2c(\sqrt{-d} + \sqrt{e}\sqrt{x})}{(c\sqrt{x} + 1)(c\sqrt{-d} + \sqrt{e})} \right)}{2e} \right)}{e} \right)$$

input `Int[(x*(a + b*ArcTanh[c*Sqrt[x]])/(d + e*x), x]`

output `2*((((x*(a + b*ArcTanh[c*Sqrt[x]]))/2 - (b*c*(-(Sqrt[x]/c^2) + ArcTanh[c*Sqrt[x]]/c^3))/2)/e - (d*(-(((a + b*ArcTanh[c*Sqrt[x]])*Log[2/(1 + c*Sqrt[x]])/e) + ((a + b*ArcTanh[c*Sqrt[x]])*Log[(2*c*(Sqrt[-d] - Sqrt[e]*Sqrt[x])/((c*Sqrt[-d] - Sqrt[e])*(1 + c*Sqrt[x]))])/(2*e) + ((a + b*ArcTanh[c*Sqrt[x]])*Log[(2*c*(Sqrt[-d] + Sqrt[e]*Sqrt[x])/((c*Sqrt[-d] + Sqrt[e])*(1 + c*Sqrt[x]))])/(2*e) + (b*PolyLog[2, 1 - 2/(1 + c*Sqrt[x])])/(2*e) - (b*PolyLog[2, 1 - (2*c*(Sqrt[-d] - Sqrt[e]*Sqrt[x])/((c*Sqrt[-d] - Sqrt[e])*(1 + c*Sqrt[x]))])/(4*e) - (b*PolyLog[2, 1 - (2*c*(Sqrt[-d] + Sqrt[e]*Sqrt[x])/((c*Sqrt[-d] + Sqrt[e])*(1 + c*Sqrt[x]))])/(4*e)))))/e`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 262 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m-1)*((a + b*x^2)^(p+1)/(b*(m+2*p+1))), x] - Simp[a*c^2*((m-1)/(b*(m+2*p+1)) Int[(c*x)^(m-2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2-1] && NeQ[m+2*p+1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6542 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((f_.)*(x_)^(m_.))/((d_.) + (e_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x^n])^p, x], x] - Simp[d*(f^2/e Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x^n])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

rule 6606 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((f_.)*(x_)^(m_.))*((d_.) + (e_.)*(x_)^2)^(q_.), x_Symbol] := With[{u = ExpandIntegrand[(a + b*ArcTanh[c*x^n])^p, (f*x)^m*(d + e*x^2)^q, x]}, Int[u, x] /; SumQ[u]] /; FreeQ[{a, b, c, d, e, f, m}, x] && IntegerQ[q] && IGtQ[p, 0] && (GtQ[q, 0] || IntegerQ[m])`

rule 7267 `Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Simp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 494, normalized size of antiderivative = 1.32

method	result
parts	$\frac{ax}{e} - \frac{ad \ln(ex+d)}{e^2} + 2b \left(\frac{c^4 \operatorname{arctanh}(c\sqrt{x})x}{2e} - \frac{c^4 \operatorname{arctanh}(c\sqrt{x})d \ln(c^2 ex+c^2 d)}{2e^2} - \frac{d c^2 \left(\frac{\ln(c\sqrt{x}-1) \ln(c^2 ex+c^2 d)}{2} - e \left(\frac{1}{2} \right) \right)}{e^2} \right)$
derivativedivides	$\frac{a c^4 x}{e} - \frac{a c^4 d \ln(c^2 ex+c^2 d)}{e^2} + 2b c^2 \left(\frac{\operatorname{arctanh}(c\sqrt{x})c^2 x}{2e} - \frac{\operatorname{arctanh}(c\sqrt{x})d c^2 \ln(c^2 ex+c^2 d)}{2e^2} - \frac{d c^2 \left(\frac{\ln(c\sqrt{x}-1) \ln(c^2 ex+c^2 d)}{2} - e \left(\frac{1}{2} \right) \right)}{e^2} \right)$
default	$\frac{a c^4 x}{e} - \frac{a c^4 d \ln(c^2 ex+c^2 d)}{e^2} + 2b c^2 \left(\frac{\operatorname{arctanh}(c\sqrt{x})c^2 x}{2e} - \frac{\operatorname{arctanh}(c\sqrt{x})d c^2 \ln(c^2 ex+c^2 d)}{2e^2} - \frac{d c^2 \left(\frac{\ln(c\sqrt{x}-1) \ln(c^2 ex+c^2 d)}{2} - e \left(\frac{1}{2} \right) \right)}{e^2} \right)$

```
input int(x*(a+b*arctanh(c*x^(1/2)))/(e*x+d), x, method=_RETURNVERBOSE)
```

```
output a/e*x-a*d/e^2*ln(e*x+d)+2*b/c^4*(1/2*c^4*arctanh(c*x^(1/2))/e*x-1/2*c^4*arctanh(c*x^(1/2))*d/e^2*ln(c^2*e*x+c^2*d)-1/2*c^2*(d*c^2/e^2*(1/2*ln(c*x^(1/2)-1)*ln(c^2*e*x+c^2*d)-e*(1/2*ln(c*x^(1/2)-1)*(ln((c*(-d*e)^(1/2)-e*(c*x^(1/2)-1)-e)/(c*(-d*e)^(1/2)-e))+ln((c*(-d*e)^(1/2)+e*(c*x^(1/2)-1)+e)/(c*(-d*e)^(1/2)+e)))/e+1/2*(dilog((c*(-d*e)^(1/2)-e*(c*x^(1/2)-1)-e)/(c*(-d*e)^(1/2)-e))+dilog((c*(-d*e)^(1/2)+e*(c*x^(1/2)-1)+e)/(c*(-d*e)^(1/2)+e)))/e)-1/2*ln(1+c*x^(1/2))*ln(c^2*e*x+c^2*d)+e*(1/2*ln(1+c*x^(1/2))*(ln((c*(-d*e)^(1/2)-e*(1+c*x^(1/2))+e)/(c*(-d*e)^(1/2)+e))+ln((c*(-d*e)^(1/2)+e*(1+c*x^(1/2))-e)/(c*(-d*e)^(1/2)-e)))/e+1/2*(dilog((c*(-d*e)^(1/2)-e*(1+c*x^(1/2))+e)/(c*(-d*e)^(1/2)+e))+dilog((c*(-d*e)^(1/2)+e*(1+c*x^(1/2))-e)/(c*(-d*e)^(1/2)-e)))/e)-1/e*(c*x^(1/2)+1/2*ln(c*x^(1/2)-1)-1/2*ln(1+c*x^(1/2))))
```

Fricas [F]

$$\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx = \int \frac{(b \operatorname{artanh}(c\sqrt{x}) + a)x}{ex + d} dx$$

input `integrate(x*(a+b*arctanh(c*x^(1/2)))/(e*x+d),x, algorithm="fricas")`

output `integral((b*x*arctanh(c*sqrt(x)) + a*x)/(e*x + d), x)`

Sympy [F]

$$\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx = \int \frac{x(a + b \operatorname{atanh}(c\sqrt{x}))}{d + ex} dx$$

input `integrate(x*(a+b*atanh(c*x**(1/2)))/(e*x+d),x)`

output `Integral(x*(a + b*atanh(c*sqrt(x)))/(d + e*x), x)`

Maxima [F]

$$\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx = \int \frac{(b \operatorname{artanh}(c\sqrt{x}) + a)x}{ex + d} dx$$

input `integrate(x*(a+b*arctanh(c*x^(1/2)))/(e*x+d),x, algorithm="maxima")`

output `a*(x/e - d*log(e*x + d)/e^2) + b*integrate(1/2*x*log(c*sqrt(x) + 1)/(e*x + d), x) - b*integrate(1/2*x*log(-c*sqrt(x) + 1)/(e*x + d), x)`

Giac [F]

$$\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx = \int \frac{(b \operatorname{artanh}(c\sqrt{x}) + a)x}{ex + d} dx$$

input `integrate(x*(a+b*arctanh(c*x^(1/2)))/(e*x+d),x, algorithm="giac")`

output `integrate((b*arctanh(c*sqrt(x)) + a)*x/(e*x + d), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx = \int \frac{x(a + b \operatorname{atanh}(c\sqrt{x}))}{d + ex} dx$$

input `int((x*(a + b*atanh(c*x^(1/2))))/(d + e*x),x)`

output `int((x*(a + b*atanh(c*x^(1/2))))/(d + e*x), x)`

Reduce [F]

$$\int \frac{x(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} dx = \frac{\left(\int \frac{\operatorname{atanh}(\sqrt{x}c)x}{ex+d} dx\right) b e^2 - \log(ex + d) ad + aex}{e^2}$$

input `int(x*(a+b*atanh(c*x^(1/2)))/(e*x+d),x)`

output `(int((atanh(sqrt(x)*c)*x)/(d + e*x),x)*b*e**2 - log(d + e*x)*a*d + a*e*x)/e**2`

3.47 $\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{d+ex} dx$

Optimal result	413
Mathematica [C] (warning: unable to verify)	414
Rubi [A] (verified)	415
Maple [A] (verified)	416
Fricas [F]	417
Sympy [F]	417
Maxima [F]	418
Giac [F]	418
Mupad [F(-1)]	418
Reduce [F]	419

Optimal result

Integrand size = 20, antiderivative size = 318

$$\int \frac{a + b\operatorname{arctanh}(c\sqrt{x})}{d + ex} dx = -\frac{2(a + b\operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1+c\sqrt{x}}\right)}{e} + \frac{(a + b\operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2c(\sqrt{-d}-\sqrt{e}\sqrt{x})}{(c\sqrt{-d}-\sqrt{e})(1+c\sqrt{x})}\right)}{e} + \frac{(a + b\operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2c(\sqrt{-d}+\sqrt{e}\sqrt{x})}{(c\sqrt{-d}+\sqrt{e})(1+c\sqrt{x})}\right)}{e} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c\sqrt{x}}\right)}{e} - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2c(\sqrt{-d}-\sqrt{e}\sqrt{x})}{(c\sqrt{-d}-\sqrt{e})(1+c\sqrt{x})}\right)}{2e} - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2c(\sqrt{-d}+\sqrt{e}\sqrt{x})}{(c\sqrt{-d}+\sqrt{e})(1+c\sqrt{x})}\right)}{2e}$$

output

```
-2*(a+b*arctanh(c*x^(1/2)))*ln(2/(1+c*x^(1/2)))/e+(a+b*arctanh(c*x^(1/2))
*ln(2*c*((-d)^(1/2)-e^(1/2)*x^(1/2))/(c*(-d)^(1/2)-e^(1/2))/(1+c*x^(1/2)))
/e+(a+b*arctanh(c*x^(1/2)))*ln(2*c*((-d)^(1/2)+e^(1/2)*x^(1/2))/(c*(-d)^(1
/2)+e^(1/2))/(1+c*x^(1/2)))/e+b*polylog(2,1-2/(1+c*x^(1/2)))/e-1/2*b*polyl
og(2,1-2*c*((-d)^(1/2)-e^(1/2)*x^(1/2))/(c*(-d)^(1/2)-e^(1/2))/(1+c*x^(1/2
)))/e-1/2*b*polylog(2,1-2*c*((-d)^(1/2)+e^(1/2)*x^(1/2))/(c*(-d)^(1/2)+e^(
1/2))/(1+c*x^(1/2)))/e
```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 1.16 (sec) , antiderivative size = 432, normalized size of antiderivative = 1.36

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{d + ex} dx = \frac{a \log(d + ex)}{e} + b \left(-2 \operatorname{arctanh}(c\sqrt{x})^2 + 4i \arcsin \left(\sqrt{\frac{c^2 d}{c^2 d + e}} \right) \operatorname{arctanh} \left(\frac{ce\sqrt{x}}{\sqrt{-c^2 de}} \right) + 2 \operatorname{arctanh}(c\sqrt{x}) \left(\operatorname{arctanh}(c\sqrt{x}) + 2 \log \left(\frac{d + ex}{d} \right) \right) \right)$$

input

```
Integrate[(a + b*ArcTanh[c*Sqrt[x]])/(d + e*x), x]
```

output

```
(a*Log[d + e*x])/e - (b*(-2*ArcTanh[c*Sqrt[x]]^2 + (4*I)*ArcSin[Sqrt[(c^2*d
d)/(c^2*d + e)]]*ArcTanh[(c*e*Sqrt[x])/Sqrt[-(c^2*d*e)]] + 2*ArcTanh[c*Sqr
t[x]]*(ArcTanh[c*Sqrt[x]] + 2*Log[1 + E^(-2*ArcTanh[c*Sqrt[x]])]) - 2*((-I
)*ArcSin[Sqrt[(c^2*d)/(c^2*d + e)]] + ArcTanh[c*Sqrt[x]]*Log[(-2*Sqrt[-(c
^2*d*e)] + e*(-1 + E^(2*ArcTanh[c*Sqrt[x]])) + c^2*d*(1 + E^(2*ArcTanh[c*S
qrt[x])])])]/((c^2*d + e)*E^(2*ArcTanh[c*Sqrt[x]])) - 2*(I*ArcSin[Sqrt[(c^2
*d)/(c^2*d + e)]] + ArcTanh[c*Sqrt[x]]*Log[(2*Sqrt[-(c^2*d*e)] + e*(-1 +
E^(2*ArcTanh[c*Sqrt[x]]) + c^2*d*(1 + E^(2*ArcTanh[c*Sqrt[x])])])]/((c^2*d
+ e)*E^(2*ArcTanh[c*Sqrt[x]])) - 2*PolyLog[2, -E^(-2*ArcTanh[c*Sqrt[x]])]
+ PolyLog[2, -(c^2*d) + e - 2*Sqrt[-(c^2*d*e)])/((c^2*d + e)*E^(2*ArcTan
h[c*Sqrt[x]])]) + PolyLog[2, -(c^2*d) + e + 2*Sqrt[-(c^2*d*e)])/((c^2*d +
e)*E^(2*ArcTanh[c*Sqrt[x]])])])]/(2*e)
```

Rubi [A] (verified)

Time = 0.84 (sec) , antiderivative size = 329, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {6484, 6606, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{d + ex} dx$$

↓ 6484

$$2 \int \frac{\sqrt{x}(a + b \operatorname{arctanh}(c\sqrt{x}))}{d + ex} d\sqrt{x}$$

↓ 6606

$$2 \int \left(\frac{a + b \operatorname{arctanh}(c\sqrt{x})}{2\sqrt{e}(\sqrt{-d} + \sqrt{e}\sqrt{x})} - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{2\sqrt{e}(\sqrt{-d} - \sqrt{e}\sqrt{x})} \right) d\sqrt{x}$$

↓ 2009

$$2 \left(\frac{(a + b \operatorname{arctanh}(c\sqrt{x})) \log \left(\frac{2c(\sqrt{-d} - \sqrt{e}\sqrt{x})}{(c\sqrt{x}+1)(c\sqrt{-d} - \sqrt{e})} \right)}{2e} + \frac{(a + b \operatorname{arctanh}(c\sqrt{x})) \log \left(\frac{2c(\sqrt{-d} + \sqrt{e}\sqrt{x})}{(c\sqrt{x}+1)(c\sqrt{-d} + \sqrt{e})} \right)}{2e} - \frac{\log \left(\frac{2}{c\sqrt{x}} \right)}{2e} \right)$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])/(d + e*x),x]`

output `2*(-((a + b*ArcTanh[c*Sqrt[x]])*Log[2/(1 + c*Sqrt[x]])/e) + ((a + b*ArcTanh[c*Sqrt[x]])*Log[(2*c*(Sqrt[-d] - Sqrt[e]*Sqrt[x]))/((c*Sqrt[-d] - Sqrt[e])*(1 + c*Sqrt[x]))]/(2*e) + ((a + b*ArcTanh[c*Sqrt[x]])*Log[(2*c*(Sqrt[-d] + Sqrt[e]*Sqrt[x]))/((c*Sqrt[-d] + Sqrt[e])*(1 + c*Sqrt[x]))]/(2*e) + (b*PolyLog[2, 1 - 2/(1 + c*Sqrt[x])]/(2*e) - (b*PolyLog[2, 1 - (2*c*(Sqrt[-d] - Sqrt[e]*Sqrt[x]))/((c*Sqrt[-d] - Sqrt[e])*(1 + c*Sqrt[x]))]/(4*e) - (b*PolyLog[2, 1 - (2*c*(Sqrt[-d] + Sqrt[e]*Sqrt[x]))/((c*Sqrt[-d] + Sqrt[e])*(1 + c*Sqrt[x]))]/(4*e))`

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6484 Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_)])*(b_.)/((d_) + (e_.)*(x_)), x_Symbol]
:= With[{k = Denominator[n]}, Simp[k Subst[Int[x^(k - 1)*((a + b*ArcTanh
[c*x^(k*n)])/(d + e*x^k)), x], x, x^(1/k)], x]] /; FreeQ[{a, b, c, d, e}, x]
&& FractionQ[n]
```

```
rule 6606 Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_))^(m_.)*((d_) + (e
_.)*(x_)^2)^(q_.), x_Symbol] := With[{u = ExpandIntegrand[(a + b*ArcTanh[c*
x])^p, (f*x)^m*(d + e*x^2)^q, x]}, Int[u, x] /; SumQ[u]] /; FreeQ[{a, b, c,
d, e, f, m}, x] && IntegerQ[q] && IGtQ[p, 0] && (GtQ[q, 0] || IntegerQ[m])
```

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 433, normalized size of antiderivative = 1.36

method	result
parts	$\frac{a \ln(ex+d)}{e} + \frac{2b \left(\frac{c^2 \ln(c^2 ex+c^2 d)}{2e} \operatorname{arctanh}(c\sqrt{x}) - \frac{\ln(c\sqrt{x}-1) \ln(c^2 ex+c^2 d)}{2} + e \left(\frac{\ln(c\sqrt{x}-1) \left(\ln\left(\frac{c\sqrt{-de}-e(c\sqrt{x}-1)}{c\sqrt{-de}-e}\right) - \frac{\ln(c\sqrt{-de}-e(c\sqrt{x}-1)}{c\sqrt{-de}-e}\right)}{2e} \right) \right)}{e}$
derivativedivides	$\frac{a c^2 \ln(c^2 ex+c^2 d)}{e} + 2b c^2 \left(\frac{\ln(c^2 ex+c^2 d) \operatorname{arctanh}(c\sqrt{x})}{2e} - \frac{\ln(c\sqrt{x}-1) \ln(c^2 ex+c^2 d)}{2} + e \left(\frac{\ln(c\sqrt{x}-1) \left(\ln\left(\frac{c\sqrt{-de}-e(c\sqrt{x}-1)}{c\sqrt{-de}-e}\right) - \frac{\ln(c\sqrt{-de}-e(c\sqrt{x}-1)}{c\sqrt{-de}-e}\right)}{2e} \right) \right)$
default	$\frac{a c^2 \ln(c^2 ex+c^2 d)}{e} + 2b c^2 \left(\frac{\ln(c^2 ex+c^2 d) \operatorname{arctanh}(c\sqrt{x})}{2e} - \frac{\ln(c\sqrt{x}-1) \ln(c^2 ex+c^2 d)}{2} + e \left(\frac{\ln(c\sqrt{x}-1) \left(\ln\left(\frac{c\sqrt{-de}-e(c\sqrt{x}-1)}{c\sqrt{-de}-e}\right) - \frac{\ln(c\sqrt{-de}-e(c\sqrt{x}-1)}{c\sqrt{-de}-e}\right)}{2e} \right) \right)$

```
input int((a+b*arctanh(c*x^(1/2)))/(e*x+d), x, method=_RETURNVERBOSE)
```

output

```
a*ln(e*x+d)/e+2*b/c^2*(1/2*c^2/e*ln(c^2*e*x+c^2*d)*arctanh(c*x^(1/2))-1/2*
c^2/e*(-1/2*ln(c*x^(1/2)-1)*ln(c^2*e*x+c^2*d)+e*(1/2*ln(c*x^(1/2)-1)*(ln((
c*(-d*e)^(1/2)-e*(c*x^(1/2)-1)-e)/(c*(-d*e)^(1/2)-e))+ln((c*(-d*e)^(1/2)+e
*(c*x^(1/2)-1)+e)/(c*(-d*e)^(1/2)+e)))/e+1/2*(dilog((c*(-d*e)^(1/2)-e*(c*x
^(1/2)-1)-e)/(c*(-d*e)^(1/2)-e))+dilog((c*(-d*e)^(1/2)+e*(c*x^(1/2)-1)+e)/
(c*(-d*e)^(1/2)+e)))/e)+1/2*ln(1+c*x^(1/2))*ln(c^2*e*x+c^2*d)-e*(1/2*ln(1+
c*x^(1/2))*(ln((c*(-d*e)^(1/2)-e*(1+c*x^(1/2))+e)/(c*(-d*e)^(1/2)+e))+ln((
c*(-d*e)^(1/2)+e*(1+c*x^(1/2))-e)/(c*(-d*e)^(1/2)-e)))/e+1/2*(dilog((c*(-d
*e)^(1/2)-e*(1+c*x^(1/2))+e)/(c*(-d*e)^(1/2)+e))+dilog((c*(-d*e)^(1/2)+e*(
1+c*x^(1/2))-e)/(c*(-d*e)^(1/2)-e)))/e))
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{d + ex} dx = \int \frac{b \operatorname{artanh}(c\sqrt{x}) + a}{ex + d} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))/(e*x+d),x, algorithm="fricas")
```

output

```
integral((b*arctanh(c*sqrt(x)) + a)/(e*x + d), x)
```

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{d + ex} dx = \int \frac{a + b \operatorname{atanh}(c\sqrt{x})}{d + ex} dx$$

input

```
integrate((a+b*atanh(c*x**(1/2)))/(e*x+d),x)
```

output

```
Integral((a + b*atanh(c*sqrt(x)))/(d + e*x), x)
```

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{d + ex} dx = \int \frac{b \operatorname{artanh}(c\sqrt{x}) + a}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/(e*x+d),x, algorithm="maxima")`

output `b*integrate(1/2*log(c*sqrt(x) + 1)/(e*x + d), x) - b*integrate(1/2*log(-c*sqrt(x) + 1)/(e*x + d), x) + a*log(e*x + d)/e`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{d + ex} dx = \int \frac{b \operatorname{artanh}(c\sqrt{x}) + a}{ex + d} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/(e*x+d),x, algorithm="giac")`

output `integrate((b*arctanh(c*sqrt(x)) + a)/(e*x + d), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{d + ex} dx = \int \frac{a + b \operatorname{atanh}(c\sqrt{x})}{d + ex} dx$$

input `int((a + b*atanh(c*x^(1/2)))/(d + e*x),x)`

output `int((a + b*atanh(c*x^(1/2)))/(d + e*x), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{d + ex} dx = \frac{\left(\int \frac{\operatorname{atanh}(\sqrt{x}c)}{ex+d} dx \right) be + \log(ex + d) a}{e}$$

input `int((a+b*atanh(c*x^(1/2)))/(e*x+d),x)`

output `(int(atanh(sqrt(x)*c)/(d + e*x),x)*b*e + log(d + e*x)*a)/e`

3.48 $\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(d+ex)} dx$

Optimal result	420
Mathematica [C] (warning: unable to verify)	421
Rubi [A] (verified)	422
Maple [A] (verified)	424
Fricas [F]	425
Sympy [F(-1)]	425
Maxima [F]	425
Giac [F]	426
Mupad [F(-1)]	426
Reduce [F]	426

Optimal result

Integrand size = 23, antiderivative size = 358

$$\begin{aligned}
 \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(d+ex)} dx = & \frac{2(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1+c\sqrt{x}}\right)}{d} \\
 & - \frac{(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2c(\sqrt{-d}-\sqrt{e}\sqrt{x})}{(c\sqrt{-d}-\sqrt{e})(1+c\sqrt{x})}\right)}{d} \\
 & - \frac{(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2c(\sqrt{-d}+\sqrt{e}\sqrt{x})}{(c\sqrt{-d}+\sqrt{e})(1+c\sqrt{x})}\right)}{d} \\
 & + \frac{a \log(x)}{d} - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c\sqrt{x}}\right)}{d} \\
 & + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2c(\sqrt{-d}-\sqrt{e}\sqrt{x})}{(c\sqrt{-d}-\sqrt{e})(1+c\sqrt{x})}\right)}{2d} \\
 & + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2c(\sqrt{-d}+\sqrt{e}\sqrt{x})}{(c\sqrt{-d}+\sqrt{e})(1+c\sqrt{x})}\right)}{2d} \\
 & - \frac{b \operatorname{PolyLog}(2, -c\sqrt{x})}{d} + \frac{b \operatorname{PolyLog}(2, c\sqrt{x})}{d}
 \end{aligned}$$

output

```
2*(a+b*arctanh(c*x^(1/2)))*ln(2/(1+c*x^(1/2)))/d-(a+b*arctanh(c*x^(1/2)))*
ln(2*c*((-d)^(1/2)-e^(1/2)*x^(1/2))/(c*(-d)^(1/2)-e^(1/2))/(1+c*x^(1/2)))/
d-(a+b*arctanh(c*x^(1/2)))*ln(2*c*((-d)^(1/2)+e^(1/2)*x^(1/2))/(c*(-d)^(1/2)+e^(1/2))/(1+c*x^(1/2)))/d+a*ln(x)/d-b*polylog(2,1-2/(1+c*x^(1/2)))/d+1/
2*b*polylog(2,1-2*c*((-d)^(1/2)-e^(1/2)*x^(1/2))/(c*(-d)^(1/2)-e^(1/2))/(1+c*x^(1/2)))/d+1/2*b*polylog(2,1-2*c*((-d)^(1/2)+e^(1/2)*x^(1/2))/(c*(-d)^(1/2)+e^(1/2))/(1+c*x^(1/2)))/d-b*polylog(2,-c*x^(1/2))/d+b*polylog(2,c*x^(1/2))/d
```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 1.50 (sec) , antiderivative size = 445, normalized size of antiderivative = 1.24

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(d + ex)} dx$$

$$= \frac{-2b \operatorname{arctanh}(c\sqrt{x})^2 + 4ib \arcsin\left(\sqrt{\frac{c^2 d}{c^2 d + e}}\right) \operatorname{arctanh}\left(\frac{ce\sqrt{x}}{\sqrt{-c^2 de}}\right) + 2b \operatorname{arctanh}(c\sqrt{x}) \left(\operatorname{arctanh}(c\sqrt{x}) + 2 \log\left(\frac{d + ex}{d}\right)\right)}{d(d + ex)}$$

input

```
Integrate[(a + b*ArcTanh[c*Sqrt[x]])/(x*(d + e*x)),x]
```

output

```
(-2*b*ArcTanh[c*Sqrt[x]]^2 + (4*I)*b*ArcSin[Sqrt[(c^2*d)/(c^2*d + e)]]*Arc
Tanh[(c*e*Sqrt[x])/Sqrt[-(c^2*d*e)]] + 2*b*ArcTanh[c*Sqrt[x]]*(ArcTanh[c*S
qrt[x]] + 2*Log[1 - E^(-2*ArcTanh[c*Sqrt[x]])]) + (2*I)*b*(ArcSin[Sqrt[(c^
2*d)/(c^2*d + e)]] + I*ArcTanh[c*Sqrt[x]])*Log[(-2*Sqrt[-(c^2*d*e)] + e*(-
1 + E^(2*ArcTanh[c*Sqrt[x]])) + c^2*d*(1 + E^(2*ArcTanh[c*Sqrt[x]]))]/((c^
2*d + e)*E^(2*ArcTanh[c*Sqrt[x]]))] - 2*b*(I*ArcSin[Sqrt[(c^2*d)/(c^2*d +
e)]] + ArcTanh[c*Sqrt[x]])*Log[(2*Sqrt[-(c^2*d*e)] + e*(-1 + E^(2*ArcTanh[
c*Sqrt[x]])) + c^2*d*(1 + E^(2*ArcTanh[c*Sqrt[x]]))]/((c^2*d + e)*E^(2*Arc
Tanh[c*Sqrt[x]]))] + 2*a*Log[x] - 2*a*Log[d + e*x] - 2*b*PolyLog[2, E^(-2*
ArcTanh[c*Sqrt[x]])] + b*PolyLog[2, (-c^2*d + e - 2*Sqrt[-(c^2*d*e)])/((c^
2*d + e)*E^(2*ArcTanh[c*Sqrt[x]])] + b*PolyLog[2, (-c^2*d + e + 2*Sqr
t[-(c^2*d*e)])/((c^2*d + e)*E^(2*ArcTanh[c*Sqrt[x]]))]/(2*d)
```

Rubi [A] (verified)

Time = 1.36 (sec) , antiderivative size = 374, normalized size of antiderivative = 1.04, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {7267, 2026, 6554, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x(d + ex)} dx \\
 & \quad \downarrow \text{7267} \\
 & 2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{ex^{3/2} + d\sqrt{x}} d\sqrt{x} \\
 & \quad \downarrow \text{2026} \\
 & 2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{\sqrt{x}(d + ex)} d\sqrt{x} \\
 & \quad \downarrow \text{6554} \\
 & 2 \int \left(\frac{a + \operatorname{barctanh}(c\sqrt{x})}{d\sqrt{x}} - \frac{e\sqrt{x}(a + \operatorname{barctanh}(c\sqrt{x}))}{d(d + ex)} \right) d\sqrt{x} \\
 & \quad \downarrow \text{2009} \\
 & 2 \left(-\frac{(a + \operatorname{barctanh}(c\sqrt{x})) \log\left(\frac{2c(\sqrt{-d} - \sqrt{e}\sqrt{x})}{(c\sqrt{x} + 1)(c\sqrt{-d} - \sqrt{e})}\right)}{2d} - \frac{(a + \operatorname{barctanh}(c\sqrt{x})) \log\left(\frac{2c(\sqrt{-d} + \sqrt{e}\sqrt{x})}{(c\sqrt{x} + 1)(c\sqrt{-d} + \sqrt{e})}\right)}{2d} + \frac{\log\left(\frac{c\sqrt{x}}{c\sqrt{x} + 1}\right)}{2d} \right)
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c*sqrt[x]])/(x*(d + e*x)),x]`

output

```
2*(((a + b*ArcTanh[c*Sqrt[x]])*Log[2/(1 + c*Sqrt[x])])/d - ((a + b*ArcTanh
[c*Sqrt[x]])*Log[(2*c*(Sqrt[-d] - Sqrt[e]*Sqrt[x])/((c*Sqrt[-d] - Sqrt[e]
)*(1 + c*Sqrt[x]))])/(2*d) - ((a + b*ArcTanh[c*Sqrt[x]])*Log[(2*c*(Sqrt[-d]
+ Sqrt[e]*Sqrt[x])/((c*Sqrt[-d] + Sqrt[e])*(1 + c*Sqrt[x]))])/(2*d) + (
a*Log[Sqrt[x])/d - (b*PolyLog[2, 1 - 2/(1 + c*Sqrt[x])])/(2*d) + (b*PolyL
og[2, 1 - (2*c*(Sqrt[-d] - Sqrt[e]*Sqrt[x])/((c*Sqrt[-d] - Sqrt[e])*(1 +
c*Sqrt[x]))])/(4*d) + (b*PolyLog[2, 1 - (2*c*(Sqrt[-d] + Sqrt[e]*Sqrt[x])
)/((c*Sqrt[-d] + Sqrt[e])*(1 + c*Sqrt[x]))])/(4*d) - (b*PolyLog[2, -(c*Sqrt
[x])])/(2*d) + (b*PolyLog[2, c*Sqrt[x])/(2*d))
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2026

```
Int[(Fx_)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p
*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0] /; PolyQ[Px, x] && Integ
erQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])
```

rule 6554

```
Int[(((a_) + ArcTanh[(c_)*(x_)]*(b_))*(x_)^(m_))/((d_) + (e_)*(x_)^2),
x_Symbol] := Int[ExpandIntegrand[a + b*ArcTanh[c*x], x^m/(d + e*x^2), x],
x] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[m] && !(EqQ[m, 1] && NeQ[a, 0]
)
```

rule 7267

```
Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]]
```


Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 499, normalized size of antiderivative = 1.39

method	result
parts	$\frac{a \ln(x)}{d} - \frac{a \ln(ex+d)}{d} + b \left(\frac{2 \operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x})}{d} - \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c^2ex+c^2d)}{d} - c^2 \left(\frac{\ln(c\sqrt{x}-1) \ln(c^2e...}{2} \right. \right.$
derivativedivides	$- \frac{a \ln(c^2ex+c^2d)}{d} + \frac{2a \ln(c\sqrt{x})}{d} + 2b c^2 \left(- \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c^2ex+c^2d)}{2d c^2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x})}{d c^2} - \frac{\ln(c\sqrt{x})}{c^2} \right.$
default	$- \frac{a \ln(c^2ex+c^2d)}{d} + \frac{2a \ln(c\sqrt{x})}{d} + 2b c^2 \left(- \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c^2ex+c^2d)}{2d c^2} + \frac{\operatorname{arctanh}(c\sqrt{x}) \ln(c\sqrt{x})}{d c^2} - \frac{\ln(c\sqrt{x})}{c^2} \right.$

input `int((a+b*arctanh(c*x^(1/2)))/x/(e*x+d),x,method=_RETURNVERBOSE)`

output `a*ln(x)/d-a/d*ln(e*x+d)+b*(2*arctanh(c*x^(1/2))/d*ln(c*x^(1/2))-arctanh(c*x^(1/2))/d*ln(c^2*e*x+c^2*d)-c^2*(1/d/c^2*(1/2*ln(c*x^(1/2))-1)*ln(c^2*e*x+c^2*d)-e*(1/2*ln(c*x^(1/2))-1)*(ln((c*(-d*e)^(1/2)-e*(c*x^(1/2))-1)-e)/(c*(-d*e)^(1/2)-e))+ln((c*(-d*e)^(1/2)+e*(c*x^(1/2))-1)+e)/(c*(-d*e)^(1/2)+e)))/e+1/2*(dilog((c*(-d*e)^(1/2)-e*(c*x^(1/2))-1)-e)/(c*(-d*e)^(1/2)-e))+dilog((c*(-d*e)^(1/2)+e*(c*x^(1/2))-1)+e)/(c*(-d*e)^(1/2)+e))/e-1/2*ln(1+c*x^(1/2))*ln(c^2*e*x+c^2*d)+e*(1/2*ln(1+c*x^(1/2))*(ln((c*(-d*e)^(1/2)-e*(1+c*x^(1/2))+e)/(c*(-d*e)^(1/2)+e))+ln((c*(-d*e)^(1/2)+e*(1+c*x^(1/2))-e)/(c*(-d*e)^(1/2)-e)))/e+1/2*(dilog((c*(-d*e)^(1/2)-e*(1+c*x^(1/2))+e)/(c*(-d*e)^(1/2)+e))+dilog((c*(-d*e)^(1/2)+e*(1+c*x^(1/2))-e)/(c*(-d*e)^(1/2)-e)))/e)-2/d/c^2*(-1/2*dilog(c*x^(1/2))-1/2*dilog(1+c*x^(1/2))-1/2*ln(c*x^(1/2))*ln(1+c*x^(1/2))))`

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(d + ex)} dx = \int \frac{b \operatorname{artanh}(c\sqrt{x}) + a}{(ex + d)x} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x/(e*x+d),x, algorithm="fricas")`

output `integral((b*arctanh(c*sqrt(x)) + a)/(e*x^2 + d*x), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(d + ex)} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**(1/2)))/x/(e*x+d),x)`

output `Timed out`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(d + ex)} dx = \int \frac{b \operatorname{artanh}(c\sqrt{x}) + a}{(ex + d)x} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x/(e*x+d),x, algorithm="maxima")`

output `-a*(log(e*x + d)/d - log(x)/d) + b*integrate(1/2*log(c*sqrt(x) + 1)/((e*x^(3/2) + d*sqrt(x))*sqrt(x)), x) - b*integrate(1/2*log(-c*sqrt(x) + 1)/((e*x^(3/2) + d*sqrt(x))*sqrt(x)), x)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(d + ex)} dx = \int \frac{b \operatorname{arctanh}(c\sqrt{x}) + a}{(ex + d)x} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x/(e*x+d),x, algorithm="giac")`

output `integrate((b*arctanh(c*sqrt(x)) + a)/((e*x + d)*x), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(d + ex)} dx = \int \frac{a + b \operatorname{atanh}(c\sqrt{x})}{x(d + ex)} dx$$

input `int((a + b*atanh(c*x^(1/2)))/(x*(d + e*x)),x)`

output `int((a + b*atanh(c*x^(1/2)))/(x*(d + e*x)), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x(d + ex)} dx = \frac{\left(\int \frac{\operatorname{atanh}(\sqrt{x}c)}{e x^2 + dx} dx \right) b d - \log(ex + d) a + \log(x) a}{d}$$

input `int((a+b*atanh(c*x^(1/2)))/x/(e*x+d),x)`

output `(int(atanh(sqrt(x)*c)/(d*x + e*x**2),x)*b*d - log(d + e*x)*a + log(x)*a)/d`

3.49 $\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(d+ex)} dx$

Optimal result	427
Mathematica [C] (verified)	428
Rubi [A] (verified)	429
Maple [A] (verified)	432
Fricas [F]	433
Sympy [F(-1)]	433
Maxima [F]	434
Giac [F]	434
Mupad [F(-1)]	434
Reduce [F]	435

Optimal result

Integrand size = 23, antiderivative size = 413

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(d+ex)} dx = -\frac{bc}{d\sqrt{x}} + \frac{bc^2 \operatorname{arctanh}(c\sqrt{x})}{d} - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{dx}$$

$$- \frac{2e(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1+c\sqrt{x}}\right)}{d^2}$$

$$+ \frac{e(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2c(\sqrt{-d}-\sqrt{e}\sqrt{x})}{(c\sqrt{-d}-\sqrt{e})(1+c\sqrt{x})}\right)}{d^2}$$

$$+ \frac{e(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2c(\sqrt{-d}+\sqrt{e}\sqrt{x})}{(c\sqrt{-d}+\sqrt{e})(1+c\sqrt{x})}\right)}{d^2}$$

$$- \frac{ae \log(x)}{d^2} + \frac{be \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c\sqrt{x}}\right)}{d^2}$$

$$- \frac{be \operatorname{PolyLog}\left(2, 1 - \frac{2c(\sqrt{-d}-\sqrt{e}\sqrt{x})}{(c\sqrt{-d}-\sqrt{e})(1+c\sqrt{x})}\right)}{2d^2}$$

$$- \frac{be \operatorname{PolyLog}\left(2, 1 - \frac{2c(\sqrt{-d}+\sqrt{e}\sqrt{x})}{(c\sqrt{-d}+\sqrt{e})(1+c\sqrt{x})}\right)}{2d^2}$$

$$+ \frac{be \operatorname{PolyLog}(2, -c\sqrt{x})}{d^2} - \frac{be \operatorname{PolyLog}(2, c\sqrt{x})}{d^2}$$

output

```
-b*c/d/x^(1/2)+b*c^2*arctanh(c*x^(1/2))/d-(a+b*arctanh(c*x^(1/2)))/d/x-2*
e*(a+b*arctanh(c*x^(1/2)))*ln(2/(1+c*x^(1/2)))/d^2+e*(a+b*arctanh(c*x^(1/2)
))*ln(2*c*((-d)^(1/2)-e^(1/2)*x^(1/2))/(c*(-d)^(1/2)-e^(1/2))/(1+c*x^(1/2)
))/d^2+e*(a+b*arctanh(c*x^(1/2)))*ln(2*c*((-d)^(1/2)+e^(1/2)*x^(1/2))/(c*(
-d)^(1/2)+e^(1/2))/(1+c*x^(1/2)))/d^2-a*e*ln(x)/d^2+b*e*polylog(2,1-2/(1+c
*x^(1/2)))/d^2-1/2*b*e*polylog(2,1-2*c*((-d)^(1/2)-e^(1/2)*x^(1/2))/(c*(-d
)^(1/2)-e^(1/2))/(1+c*x^(1/2)))/d^2-1/2*b*e*polylog(2,1-2*c*((-d)^(1/2)+e
^(1/2)*x^(1/2))/(c*(-d)^(1/2)+e^(1/2))/(1+c*x^(1/2)))/d^2+b*e*polylog(2,-c*
x^(1/2))/d^2-b*e*polylog(2,c*x^(1/2))/d^2
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.94 (sec) , antiderivative size = 479, normalized size of antiderivative = 1.16

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(d + ex)} dx$$

$$= \frac{-2ad - 2aex \log(x) + 2aex \log(d + ex) - 2b \left(cd\sqrt{x} + \operatorname{arctanh}(c\sqrt{x}) \right) \left(d - c^2dx + ex \operatorname{arctanh}(c\sqrt{x}) + 2 \right)}{2d^2x^2 + 2dex + e^2x^2}$$

input

```
Integrate[(a + b*ArcTanh[c*Sqrt[x]])/(x^2*(d + e*x)),x]
```

output

```
(-2*a*d - 2*a*e*x*Log[x] + 2*a*e*x*Log[d + e*x] - 2*b*(c*d*Sqrt[x] + ArcTan
h[c*Sqrt[x]]*(d - c^2*d*x + e*x*ArcTanh[c*Sqrt[x]] + 2*e*x*Log[1 - E^(-2*
ArcTanh[c*Sqrt[x]])]) - e*x*PolyLog[2, E^(-2*ArcTanh[c*Sqrt[x]])]) + b*e*x
*(2*ArcTanh[c*Sqrt[x]]^2 - (4*I)*ArcSin[Sqrt[(c^2*d)/(c^2*d + e)]]*ArcTanh
[(c*e*Sqrt[x])/Sqrt[-(c^2*d*e)]] + 2*((-I)*ArcSin[Sqrt[(c^2*d)/(c^2*d + e)
]]) + ArcTanh[c*Sqrt[x]]*Log[(-2*Sqrt[-(c^2*d*e)] + e*(-1 + E^(2*ArcTanh[c
*Sqrt[x]])) + c^2*d*(1 + E^(2*ArcTanh[c*Sqrt[x]])))/((c^2*d + e)*E^(2*ArcT
anh[c*Sqrt[x]])]) + 2*(I*ArcSin[Sqrt[(c^2*d)/(c^2*d + e)]] + ArcTanh[c*Sqr
t[x]]*Log[(2*Sqrt[-(c^2*d*e)] + e*(-1 + E^(2*ArcTanh[c*Sqrt[x]])) + c^2*d
*(1 + E^(2*ArcTanh[c*Sqrt[x]])))/((c^2*d + e)*E^(2*ArcTanh[c*Sqrt[x]])]) -
PolyLog[2, (-(c^2*d) + e - 2*Sqrt[-(c^2*d*e)])/((c^2*d + e)*E^(2*ArcTanh[
c*Sqrt[x]])]) - PolyLog[2, (-(c^2*d) + e + 2*Sqrt[-(c^2*d*e)])/((c^2*d + e
)*E^(2*ArcTanh[c*Sqrt[x]])))]/(2*d^2*x)
```

Rubi [A] (verified)

Time = 1.67 (sec) , antiderivative size = 429, normalized size of antiderivative = 1.04, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.348$, Rules used = {7267, 2026, 6544, 6452, 264, 219, 6554, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(d + ex)} dx \\
 & \quad \downarrow \text{7267} \\
 & 2 \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{ex^{5/2} + dx^{3/2}} d\sqrt{x} \\
 & \quad \downarrow \text{2026} \\
 & 2 \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^{3/2}(d + ex)} d\sqrt{x} \\
 & \quad \downarrow \text{6544} \\
 & 2 \left(\frac{\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^{3/2}} d\sqrt{x}}{d} - \frac{e \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}(d + ex)} d\sqrt{x}}{d} \right) \\
 & \quad \downarrow \text{6452} \\
 & 2 \left(\frac{\frac{1}{2}bc \int \frac{1}{x(1 - c^2x)} d\sqrt{x} - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{2x}}{d} - \frac{e \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}(d + ex)} d\sqrt{x}}{d} \right) \\
 & \quad \downarrow \text{264} \\
 & 2 \left(\frac{\frac{1}{2}bc \left(c^2 \int \frac{1}{1 - c^2x} d\sqrt{x} - \frac{1}{\sqrt{x}} \right) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{2x}}{d} - \frac{e \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}(d + ex)} d\sqrt{x}}{d} \right) \\
 & \quad \downarrow \text{219} \\
 & 2 \left(\frac{\frac{1}{2}bc \left(\operatorname{arctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{2x}}{d} - \frac{e \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}(d + ex)} d\sqrt{x}}{d} \right) \\
 & \quad \downarrow \text{6554}
 \end{aligned}$$

$$2 \left(\frac{\frac{1}{2}bc \left(\operatorname{arctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{2x}}{d} - \frac{e \int \left(\frac{a+b\operatorname{arctanh}(c\sqrt{x})}{d\sqrt{x}} - \frac{e\sqrt{x}(a+b\operatorname{arctanh}(c\sqrt{x}))}{d(d+ex)} \right) d\sqrt{x}}{d} \right)$$

↓ 2009

$$2 \left(\frac{\frac{1}{2}bc \left(\operatorname{arctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{2x}}{d} - \frac{e \left(-\frac{(a+b\operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2c(\sqrt{-d}-\sqrt{e}\sqrt{x})}{(c\sqrt{x}+1)(c\sqrt{-d}-\sqrt{e})}\right)}{2d} - \frac{(a+b\operatorname{arctanh}(c\sqrt{x}))}{d} \right)}{d} \right)$$

input `Int[(a + b*ArcTanh[c*Sqrt[x]])/(x^2*(d + e*x)),x]`

output `2*((-1/2*(a + b*ArcTanh[c*Sqrt[x]])/x + (b*c*(-1/Sqrt[x]) + c*ArcTanh[c*Sqrt[x]]))/2)/d - (e*((a + b*ArcTanh[c*Sqrt[x]])*Log[2/(1 + c*Sqrt[x])])/d - ((a + b*ArcTanh[c*Sqrt[x]])*Log[(2*c*(Sqrt[-d] - Sqrt[e]*Sqrt[x])]/((c*Sqrt[-d] - Sqrt[e])*(1 + c*Sqrt[x]))])/(2*d) - ((a + b*ArcTanh[c*Sqrt[x]])*Log[(2*c*(Sqrt[-d] + Sqrt[e]*Sqrt[x])]/((c*Sqrt[-d] + Sqrt[e])*(1 + c*Sqrt[x]))])/(2*d) + (a*Log[Sqrt[x]])/d - (b*PolyLog[2, 1 - 2/(1 + c*Sqrt[x])])/(2*d) + (b*PolyLog[2, 1 - (2*c*(Sqrt[-d] - Sqrt[e]*Sqrt[x])]/((c*Sqrt[-d] - Sqrt[e])*(1 + c*Sqrt[x]))])/(4*d) + (b*PolyLog[2, 1 - (2*c*(Sqrt[-d] + Sqrt[e]*Sqrt[x])]/((c*Sqrt[-d] + Sqrt[e])*(1 + c*Sqrt[x]))])/(4*d) - (b*PolyLog[2, -(c*Sqrt[x])])/(2*d) + (b*PolyLog[2, c*Sqrt[x]])/(2*d))/d`

Definitions of rubi rules used

rule 219 $\text{Int}[(a_.) + (b_.) \cdot (x_.)^2]^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[-b, 2])) \cdot \text{ArcTanh}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 264 $\text{Int}[(c_.) \cdot (x_.)^m] \cdot [(a_.) + (b_.) \cdot (x_.)^2]^{p_.), x_Symbol] \rightarrow \text{Simp}[(c \cdot x)^{m+1} \cdot ((a + b \cdot x^2)^{p+1} / (a \cdot c^{m+1}))], x] - \text{Simp}[b \cdot ((m+2 \cdot p+3) / (a \cdot c^{2 \cdot (m+1)})) \text{Int}[(c \cdot x)^{m+2} \cdot (a + b \cdot x^2)^p], x] /; \text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 2026 $\text{Int}[(F x_.) \cdot (P x_.)^{p_.), x_Symbol] \rightarrow \text{With}\{r = \text{Expon}[P x, x, \text{Min}]\}, \text{Int}[x^{(p \cdot r)} \cdot \text{ExpandToSum}[P x / x^r, x]^{p \cdot F x}, x] /; \text{IGtQ}[r, 0] /; \text{PolyQ}[P x, x] \ \&\& \ \text{IntegerQ}[p] \ \&\& \ \text{!MonomialQ}[P x, x] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{!PolyQ}[u, x])$

rule 6452 $\text{Int}[(a_.) + \text{ArcTanh}[(c_.) \cdot (x_.)^{n_.)}] \cdot (b_.)^{p_.)} \cdot (x_.)^{m_.), x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot ((a + b \cdot \text{ArcTanh}[c \cdot x^n])^p / (m+1)), x] - \text{Simp}[b \cdot c \cdot n \cdot (p / (m+1)) \text{Int}[x^{m+n} \cdot ((a + b \cdot \text{ArcTanh}[c \cdot x^n])^{p-1} / (1 - c^2 \cdot x^{2 \cdot n}))], x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 6544 $\text{Int}[(a_.) + \text{ArcTanh}[(c_.) \cdot (x_.)] \cdot (b_.)]^{p_.)} \cdot ((f_.) \cdot (x_.)^m) / ((d_.) + (e_.) \cdot (x_.)^2), x_Symbol] \rightarrow \text{Simp}[1/d \text{Int}[(f \cdot x)^m \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p], x], x] - \text{Simp}[e / (d \cdot f^2) \text{Int}[(f \cdot x)^{m+2} \cdot ((a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2))], x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{LtQ}[m, -1]$

rule 6554 $\text{Int}[(a_.) + \text{ArcTanh}[(c_.) \cdot (x_.)] \cdot (b_.)] \cdot (x_.)^{m_.)} / ((d_.) + (e_.) \cdot (x_.)^2), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[a + b \cdot \text{ArcTanh}[c \cdot x], x^m / (d + e \cdot x^2), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{!(EqQ}[m, 1] \ \&\& \ \text{NeQ}[a, 0])$

rule 7267

```
Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]]
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 577, normalized size of antiderivative = 1.40

method	result
parts	$a \left(-\frac{1}{dx} - \frac{e \ln(x)}{d^2} + \frac{e \ln(ex+d)}{d^2} \right) + 2b c^2 \left(-\frac{\operatorname{arctanh}(c\sqrt{x})}{2d c^2 x} - \frac{\operatorname{arctanh}(c\sqrt{x}) e \ln(c\sqrt{x})}{c^2 d^2} + \frac{\operatorname{arctanh}(c\sqrt{x}) e}{2c^2} \right)$
derivativedivides	$2c^2 \left(\frac{ae \ln(c^2 ex+c^2 d)}{2c^2 d^2} - \frac{a}{2d c^2 x} - \frac{ae \ln(c\sqrt{x})}{c^2 d^2} \right) + b c^2 \left(\frac{\operatorname{arctanh}(c\sqrt{x}) e \ln(c^2 ex+c^2 d)}{2d^2 c^4} - \frac{\operatorname{arctanh}(c\sqrt{x})}{2d c^4 x} - \dots \right)$
default	$2c^2 \left(\frac{ae \ln(c^2 ex+c^2 d)}{2c^2 d^2} - \frac{a}{2d c^2 x} - \frac{ae \ln(c\sqrt{x})}{c^2 d^2} \right) + b c^2 \left(\frac{\operatorname{arctanh}(c\sqrt{x}) e \ln(c^2 ex+c^2 d)}{2d^2 c^4} - \frac{\operatorname{arctanh}(c\sqrt{x})}{2d c^4 x} - \dots \right)$

input

```
int((a+b*arctanh(c*x^(1/2)))/x^2/(e*x+d), x, method=_RETURNVERBOSE)
```

output

```
a*(-1/d/x-e/d^2*ln(x)+e/d^2*ln(e*x+d))+2*b*c^2*(-1/2*arctanh(c*x^(1/2))/d/
c^2/x-1/c^2*arctanh(c*x^(1/2))/d^2*e*ln(c*x^(1/2))+1/2/c^2*arctanh(c*x^(1/
2))*e/d^2*ln(c^2*e*x+c^2*d)-1/2*c^2*(1/d/c^2*(1/c/x^(1/2)+1/2*ln(c*x^(1/2)
-1)-1/2*ln(1+c*x^(1/2)))+2/d^2/c^4*e*(-1/2*dilog(c*x^(1/2))-1/2*dilog(1+c*
x^(1/2))-1/2*ln(c*x^(1/2))*ln(1+c*x^(1/2)))-1/d^2/c^4*e*(1/2*ln(c*x^(1/2)-
1)*ln(c^2*e*x+c^2*d)-e*(1/2*ln(c*x^(1/2)-1)*(ln((c*(-d*e)^(1/2)-e*(c*x^(1/
2)-1)-e)/(c*(-d*e)^(1/2)-e))+ln((c*(-d*e)^(1/2)+e*(c*x^(1/2)-1)+e)/(c*(-d*
e)^(1/2)+e)))/e+1/2*(dilog((c*(-d*e)^(1/2)-e*(c*x^(1/2)-1)-e)/(c*(-d*e)^(1
/2)-e))+dilog((c*(-d*e)^(1/2)+e*(c*x^(1/2)-1)+e)/(c*(-d*e)^(1/2)+e)))/e-1
/2*ln(1+c*x^(1/2))*ln(c^2*e*x+c^2*d)+e*(1/2*ln(1+c*x^(1/2))*(ln((c*(-d*e)^(
1/2)-e*(1+c*x^(1/2))+e)/(c*(-d*e)^(1/2)+e))+ln((c*(-d*e)^(1/2)+e*(1+c*x^(
1/2))-e)/(c*(-d*e)^(1/2)-e)))/e+1/2*(dilog((c*(-d*e)^(1/2)-e*(1+c*x^(1/2)
+e)/(c*(-d*e)^(1/2)+e))+dilog((c*(-d*e)^(1/2)+e*(1+c*x^(1/2))-e)/(c*(-d*e)
^(1/2)-e)))/e))))
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(d + ex)} dx = \int \frac{b \operatorname{artanh}(c\sqrt{x}) + a}{(ex + d)x^2} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))/x^2/(e*x+d),x, algorithm="fricas")
```

output

```
integral((b*arctanh(c*sqrt(x)) + a)/(e*x^3 + d*x^2), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(d + ex)} dx = \text{Timed out}$$

input

```
integrate((a+b*atanh(c*x**(1/2)))/x**2/(e*x+d),x)
```

output

```
Timed out
```

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(d + ex)} dx = \int \frac{b \operatorname{arctanh}(c\sqrt{x}) + a}{(ex + d)x^2} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^2/(e*x+d),x, algorithm="maxima")`

output `a*(e*log(e*x + d)/d^2 - e*log(x)/d^2 - 1/(d*x)) + b*integrate(1/2*log(c*sqrt(x) + 1)/((e*x^(5/2) + d*x^(3/2))*sqrt(x)), x) - b*integrate(1/2*log(-c*sqrt(x) + 1)/((e*x^(5/2) + d*x^(3/2))*sqrt(x)), x)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(d + ex)} dx = \int \frac{b \operatorname{arctanh}(c\sqrt{x}) + a}{(ex + d)x^2} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^2/(e*x+d),x, algorithm="giac")`

output `integrate((b*arctanh(c*sqrt(x)) + a)/((e*x + d)*x^2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(d + ex)} dx = \int \frac{a + b \operatorname{atanh}(c\sqrt{x})}{x^2(d + ex)} dx$$

input `int((a + b*atanh(c*x^(1/2)))/(x^2*(d + e*x)),x)`

output `int((a + b*atanh(c*x^(1/2)))/(x^2*(d + e*x)), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^2(d + ex)} dx = \frac{\left(\int \frac{\operatorname{atanh}(\sqrt{x}c)}{e x^3 + d x^2} dx \right) b d^2 x + \log(ex + d) a e x - \log(x) a e x - a d}{d^2 x}$$

input `int((a+b*atanh(c*x^(1/2)))/x^2/(e*x+d),x)`

output `(int(atanh(sqrt(x)*c)/(d*x**2 + e*x**3),x)*b*d**2*x + log(d + e*x)*a*e*x - log(x)*a*e*x - a*d)/(d**2*x)`

$$3.50 \quad \int \frac{a+b \operatorname{arctanh}(c\sqrt{x})}{x^3(d+ex)} dx$$

Optimal result	437
Mathematica [C] (verified)	438
Rubi [A] (verified)	438
Maple [A] (verified)	443
Fricas [F]	444
Sympy [F(-1)]	444
Maxima [F]	445
Giac [F]	445
Mupad [F(-1)]	445
Reduce [F]	446

Optimal result

Integrand size = 23, antiderivative size = 506

$$\begin{aligned}
 \int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(d + ex)} dx = & -\frac{bc}{6dx^{3/2}} - \frac{bc^3}{2d\sqrt{x}} + \frac{bce}{d^2\sqrt{x}} \\
 & + \frac{bc^4 \operatorname{arctanh}(c\sqrt{x})}{2d} - \frac{bc^2 e \operatorname{arctanh}(c\sqrt{x})}{d^2} \\
 & - \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{2dx^2} + \frac{e(a + b \operatorname{arctanh}(c\sqrt{x}))}{d^2 x} \\
 & + \frac{2e^2(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2}{1+c\sqrt{x}}\right)}{d^3} \\
 & - \frac{e^2(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2c(\sqrt{-d}-\sqrt{e}\sqrt{x})}{(c\sqrt{-d}-\sqrt{e})(1+c\sqrt{x})}\right)}{d^3} \\
 & - \frac{e^2(a + b \operatorname{arctanh}(c\sqrt{x})) \log\left(\frac{2c(\sqrt{-d}+\sqrt{e}\sqrt{x})}{(c\sqrt{-d}+\sqrt{e})(1+c\sqrt{x})}\right)}{d^3} \\
 & + \frac{ae^2 \log(x)}{d^3} - \frac{be^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c\sqrt{x}}\right)}{d^3} \\
 & + \frac{be^2 \operatorname{PolyLog}\left(2, 1 - \frac{2c(\sqrt{-d}-\sqrt{e}\sqrt{x})}{(c\sqrt{-d}-\sqrt{e})(1+c\sqrt{x})}\right)}{2d^3} \\
 & + \frac{be^2 \operatorname{PolyLog}\left(2, 1 - \frac{2c(\sqrt{-d}+\sqrt{e}\sqrt{x})}{(c\sqrt{-d}+\sqrt{e})(1+c\sqrt{x})}\right)}{2d^3} \\
 & - \frac{be^2 \operatorname{PolyLog}(2, -c\sqrt{x})}{d^3} + \frac{be^2 \operatorname{PolyLog}(2, c\sqrt{x})}{d^3}
 \end{aligned}$$

output

```

-1/6*b*c/d/x^(3/2)-1/2*b*c^3/d/x^(1/2)+b*c*e/d^2/x^(1/2)+1/2*b*c^4*arctanh
(c*x^(1/2))/d-b*c^2*e*arctanh(c*x^(1/2))/d^2-1/2*(a+b*arctanh(c*x^(1/2)))/
d/x^2+e*(a+b*arctanh(c*x^(1/2)))/d^2/x+2*e^2*(a+b*arctanh(c*x^(1/2)))*ln(2
/(1+c*x^(1/2)))/d^3-e^2*(a+b*arctanh(c*x^(1/2)))*ln(2*c*((-d)^(1/2)-e^(1/2)
)*x^(1/2))/(c*(-d)^(1/2)-e^(1/2))/(1+c*x^(1/2))/d^3-e^2*(a+b*arctanh(c*x^
(1/2)))*ln(2*c*((-d)^(1/2)+e^(1/2))*x^(1/2))/(c*(-d)^(1/2)+e^(1/2))/(1+c*x^
(1/2))/d^3+a*e^2*ln(x)/d^3-b*e^2*polylog(2,1-2/(1+c*x^(1/2)))/d^3+1/2*b*e
^2*polylog(2,1-2*c*((-d)^(1/2)-e^(1/2))*x^(1/2))/(c*(-d)^(1/2)-e^(1/2))/(1+
c*x^(1/2))/d^3+1/2*b*e^2*polylog(2,1-2*c*((-d)^(1/2)+e^(1/2))*x^(1/2))/(c*
(-d)^(1/2)+e^(1/2))/(1+c*x^(1/2))/d^3-b*e^2*polylog(2,-c*x^(1/2))/d^3+b*e
^2*polylog(2,c*x^(1/2))/d^3

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.62 (sec) , antiderivative size = 538, normalized size of antiderivative = 1.06

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(d + ex)} dx = \frac{3ad^2 - 6adex - 6ae^2x^2 \log(x) + 6ae^2x^2 \log(d + ex) + b \left(cd\sqrt{x}(d + 3c^2dx - 6ex) - 3 \operatorname{arctanh}(c\sqrt{x}) \right)}{d^3x^2}$$

input `Integrate[(a + b*ArcTanh[c*Sqrt[x]])/(x^3*(d + e*x)),x]`

output

```
-1/6*(3*a*d^2 - 6*a*d*e*x - 6*a*e^2*x^2*Log[x] + 6*a*e^2*x^2*Log[d + e*x]
+ b*(c*d*Sqrt[x]*(d + 3*c^2*d*x - 6*e*x) - 3*ArcTanh[c*Sqrt[x]]*(d*(-1 + c
^2*x)*(d + c^2*d*x - 2*e*x) + 2*e^2*x^2*ArcTanh[c*Sqrt[x]] + 4*e^2*x^2*Log
[1 - E^(-2*ArcTanh[c*Sqrt[x]])]) + 6*e^2*x^2*PolyLog[2, E^(-2*ArcTanh[c*Sq
rt[x]])] + 3*e^2*x^2*(2*ArcTanh[c*Sqrt[x]]^2 - (4*I)*ArcSin[Sqrt[(c^2*d)/(
c^2*d + e)]]*ArcTanh[(c*e*Sqrt[x])/Sqrt[-(c^2*d*e)]] + 2*((-I)*ArcSin[Sqrt
[(c^2*d)/(c^2*d + e)]] + ArcTanh[c*Sqrt[x]])*Log[(-2*Sqrt[-(c^2*d*e)] + e
(-1 + E^(2*ArcTanh[c*Sqrt[x]])) + c^2*d*(1 + E^(2*ArcTanh[c*Sqrt[x]]))]/((
c^2*d + e)*E^(2*ArcTanh[c*Sqrt[x]])) + 2*(I*ArcSin[Sqrt[(c^2*d)/(c^2*d +
e)]] + ArcTanh[c*Sqrt[x]])*Log[(2*Sqrt[-(c^2*d*e)] + e*(-1 + E^(2*ArcTanh[
c*Sqrt[x]])) + c^2*d*(1 + E^(2*ArcTanh[c*Sqrt[x]]))]/((c^2*d + e)*E^(2*Arc
Tanh[c*Sqrt[x]])) - PolyLog[2, (-c^2*d + e - 2*Sqrt[-(c^2*d*e)])/((c^2*
d + e)*E^(2*ArcTanh[c*Sqrt[x]])] - PolyLog[2, (-c^2*d + e + 2*Sqrt[-(c^
2*d*e)])/((c^2*d + e)*E^(2*ArcTanh[c*Sqrt[x]])])))/(d^3*x^2)
```

Rubi [A] (verified)

Time = 1.56 (sec) , antiderivative size = 498, normalized size of antiderivative = 0.98, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.565$, Rules used = {7267, 2026, 6544, 6452, 264, 264, 219, 6544, 6452, 264, 219, 6554, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^3(d+ex)} dx \\
& \quad \downarrow \text{7267} \\
& 2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{ex^{7/2} + dx^{5/2}} d\sqrt{x} \\
& \quad \downarrow \text{2026} \\
& 2 \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{5/2}(d+ex)} d\sqrt{x} \\
& \quad \downarrow \text{6544} \\
& 2 \left(\frac{\int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{5/2}} d\sqrt{x}}{d} - \frac{e \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(d+ex)} d\sqrt{x}}{d} \right) \\
& \quad \downarrow \text{6452} \\
& 2 \left(\frac{\frac{1}{4}bc \int \frac{1}{x^2(1-c^2x)} d\sqrt{x} - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2}}{d} - \frac{e \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(d+ex)} d\sqrt{x}}{d} \right) \\
& \quad \downarrow \text{264} \\
& 2 \left(\frac{\frac{1}{4}bc \left(c^2 \int \frac{1}{x(1-c^2x)} d\sqrt{x} - \frac{1}{3x^{3/2}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2}}{d} - \frac{e \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(d+ex)} d\sqrt{x}}{d} \right) \\
& \quad \downarrow \text{264} \\
& 2 \left(\frac{\frac{1}{4}bc \left(c^2 \left(c^2 \int \frac{1}{1-c^2x} d\sqrt{x} - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2}}{d} - \frac{e \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(d+ex)} d\sqrt{x}}{d} \right) \\
& \quad \downarrow \text{219} \\
& 2 \left(\frac{\frac{1}{4}bc \left(c^2 \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) - \frac{a + \operatorname{barctanh}(c\sqrt{x})}{4x^2}}{d} - \frac{e \int \frac{a + \operatorname{barctanh}(c\sqrt{x})}{x^{3/2}(d+ex)} d\sqrt{x}}{d} \right) \\
& \quad \downarrow \text{6544}
\end{aligned}$$

$$2 \left(\frac{\frac{1}{4}bc \left(c^2 \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) - \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{4x^2}}{d} - \frac{e \left(\int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{x^{3/2}} d\sqrt{x} - \frac{e \int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}(d+ex)} dx}{d} \right)}{d} \right)$$

↓ 6452

$$2 \left(\frac{\frac{1}{4}bc \left(c^2 \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) - \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{4x^2}}{d} - \frac{e \left(\frac{\frac{1}{2}bc \int \frac{1}{x(1-c^2x)} d\sqrt{x} - \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{2x}}{d} - \frac{e \int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}(d+ex)} dx}{d} \right)}{d} \right)$$

↓ 264

$$2 \left(\frac{\frac{1}{4}bc \left(c^2 \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) - \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{4x^2}}{d} - \frac{e \left(\frac{\frac{1}{2}bc \left(c^2 \int \frac{1}{1-c^2x} d\sqrt{x} - \frac{1}{\sqrt{x}} \right) - \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{2x}}{d} - \frac{e \int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}(d+ex)} dx}{d} \right)}{d} \right)$$

↓ 219

$$2 \left(\frac{\frac{1}{4}bc \left(c^2 \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) - \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{4x^2}}{d} - \frac{e \left(\frac{\frac{1}{2}bc \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{2x}}{d} - \frac{e \int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}(d+ex)} dx}{d} \right)}{d} \right)$$

↓ 6554

$$2 \left(\frac{\frac{1}{4}bc \left(c^2 \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) - \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{4x^2}}{d} - \frac{e \left(\frac{\frac{1}{2}bc \left(\operatorname{carctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{2x}}{d} - \frac{e \int \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{\sqrt{x}(d+ex)} dx}{d} \right)}{d} \right)$$

↓ 2009

$$2 \left(\frac{\frac{1}{4}bc \left(c^2 \left(\operatorname{arctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{1}{3x^{3/2}} \right) - \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{4x^2}}{d} - \frac{e \left(\frac{\frac{1}{2}bc \left(\operatorname{arctanh}(c\sqrt{x}) - \frac{1}{\sqrt{x}} \right) - \frac{a+b\operatorname{arctanh}(c\sqrt{x})}{2x}}{d} \right)}{d} \right)$$

```
input Int[(a + b*ArcTanh[c*Sqrt[x]])/(x^3*(d + e*x)),x]
```

```
output 2*((-1/4*(a + b*ArcTanh[c*Sqrt[x]])/x^2 + (b*c*(-1/3*1/x^(3/2) + c^2*(-1/Sqrt[x]) + c*ArcTanh[c*Sqrt[x]]))/4)/d - (e*((-1/2*(a + b*ArcTanh[c*Sqrt[x]])/x + (b*c*(-1/Sqrt[x]) + c*ArcTanh[c*Sqrt[x]]))/2)/d - (e*(((a + b*ArcTanh[c*Sqrt[x]])*Log[2/(1 + c*Sqrt[x])])/d - ((a + b*ArcTanh[c*Sqrt[x]])*Log[(2*c*(Sqrt[-d] - Sqrt[e]*Sqrt[x]))/((c*Sqrt[-d] - Sqrt[e])*(1 + c*Sqrt[x]))])/d - ((a + b*ArcTanh[c*Sqrt[x]])*Log[(2*c*(Sqrt[-d] + Sqrt[e]*Sqrt[x]))/((c*Sqrt[-d] + Sqrt[e])*(1 + c*Sqrt[x]))])/d + (a*Log[Sqrt[x]])/d - (b*PolyLog[2, 1 - 2/(1 + c*Sqrt[x])])/d + (b*PolyLog[2, 1 - (2*c*(Sqrt[-d] - Sqrt[e]*Sqrt[x]))/((c*Sqrt[-d] - Sqrt[e])*(1 + c*Sqrt[x]))])/d + (b*PolyLog[2, 1 - (2*c*(Sqrt[-d] + Sqrt[e]*Sqrt[x]))/((c*Sqrt[-d] + Sqrt[e])*(1 + c*Sqrt[x]))])/d - (b*PolyLog[2, -(c*Sqrt[x])])/d + (b*PolyLog[2, c*Sqrt[x])/d))/d)
```

Definitions of rubi rules used

rule 219 $\text{Int}[\{(a_)+ (b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))* \text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 264 $\text{Int}[\{(c_)*(x_)\}^{(m_)}*\{(a_)+ (b_)*(x_)^2\}^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(c*x)^{(m+1)}*((a+b*x^2)^{(p+1)}/(a*c*(m+1))), x] - \text{Simp}[b*((m+2*p+3)/(a*c^2*(m+1))) \ \text{Int}[(c*x)^{(m+2)}*(a+b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 2026 $\text{Int}[(F_x_)*(P_x_)^{(p_)}, x_Symbol] \rightarrow \text{With}\{r = \text{Expon}[P_x, x, \text{Min}]\}, \text{Int}[x^{(p*r)}*\text{ExpandToSum}[P_x/x^r, x]^p*F_x, x] /; \text{IGtQ}[r, 0] /; \text{PolyQ}[P_x, x] \ \&\& \ \text{IntegerQ}[p] \ \&\& \ !\text{MonomialQ}[P_x, x] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ !\text{PolyQ}[u, x])$

rule 6452 $\text{Int}[\{(a_)+ \text{ArcTanh}[(c_)*(x_)]\}^{(p_)}*(b_)^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a+b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \ \text{Int}[x^{(m+n)}*((a+b*\text{ArcTanh}[c*x^n])^{(p-1)})/(1-c^2*x^{(2*n)}), x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 6544 $\text{Int}[\{(a_)+ \text{ArcTanh}[(c_)*(x_)]\}^{(p_)}*(b_)^{(p_)}*((f_)*(x_))^{(m_)} / ((d_)+ (e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[1/d \ \text{Int}[(f*x)^m*(a+b*\text{ArcTanh}[c*x])^p, x], x] - \text{Simp}[e/(d*f^2) \ \text{Int}[(f*x)^{(m+2)}*((a+b*\text{ArcTanh}[c*x])^p/(d+e*x^2)), x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{LtQ}[m, -1]$

rule 6554 $\text{Int}[\{(a_)+ \text{ArcTanh}[(c_)*(x_)]\}^{(p_)}*(b_)^{(p_)}*(x_)^{(m_)} / ((d_)+ (e_)*(x_)^2), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[a+b*\text{ArcTanh}[c*x], x^m/(d+e*x^2), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x \ \&\& \ \text{IntegerQ}[m] \ \&\& \ !(\text{EqQ}[m, 1] \ \&\& \ \text{NeQ}[a, 0])$

rule 7267

```
Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]]
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 650, normalized size of antiderivative = 1.28

method	result
parts	$a \left(-\frac{1}{2dx^2} + \frac{e^2 \ln(x)}{d^3} + \frac{e}{d^2x} - \frac{e^2 \ln(ex+d)}{d^3} \right) + 2bc^4 \left(-\frac{\operatorname{arctanh}(c\sqrt{x})}{4dc^4x^2} + \frac{\operatorname{arctanh}(c\sqrt{x})e^2 \ln(c\sqrt{x})}{c^4d^3} + \dots \right)$
derivativedivides	$2c^4 \left(-\frac{ae^2 \ln(c^2ex+c^2d)}{2c^4d^3} - \frac{a}{4dc^4x^2} + \frac{ae^2 \ln(c\sqrt{x})}{c^4d^3} + \frac{ae}{2c^4d^2x} + bc^2 \left(-\frac{\operatorname{arctanh}(c\sqrt{x})e^2 \ln(c^2ex+c^2d)}{2d^3c^6} \dots \right) \right)$
default	$2c^4 \left(-\frac{ae^2 \ln(c^2ex+c^2d)}{2c^4d^3} - \frac{a}{4dc^4x^2} + \frac{ae^2 \ln(c\sqrt{x})}{c^4d^3} + \frac{ae}{2c^4d^2x} + bc^2 \left(-\frac{\operatorname{arctanh}(c\sqrt{x})e^2 \ln(c^2ex+c^2d)}{2d^3c^6} \dots \right) \right)$

input

```
int((a+b*arctanh(c*x^(1/2)))/x^3/(e*x+d), x, method=_RETURNVERBOSE)
```

output

```
a*(-1/2/d/x^2+e^2/d^3*ln(x)+e/d^2/x-e^2/d^3*ln(e*x+d))+2*b*c^4*(-1/4*arctanh(c*x^(1/2))/d/c^4/x^2+1/c^4*arctanh(c*x^(1/2))/d^3*e^2*ln(c*x^(1/2))+1/2/c^4*arctanh(c*x^(1/2))/d^2*e/x-1/2/c^4*arctanh(c*x^(1/2))*e^2/d^3*ln(c^2*e*x+c^2*d)-1/4*c^2*(-4/d^3/c^6*e^2*(-1/2*dilog(c*x^(1/2))-1/2*dilog(1+c*x^(1/2))-1/2*ln(c*x^(1/2))*ln(1+c*x^(1/2)))+2/d^3/c^6*e^2*(1/2*ln(c*x^(1/2))-1)*ln(c^2*e*x+c^2*d)-e*(1/2*ln(c*x^(1/2))-1)*(ln((c*(-d*e)^(1/2)-e*(c*x^(1/2)-1)-e)/(c*(-d*e)^(1/2)-e))+ln((c*(-d*e)^(1/2)+e*(c*x^(1/2)-1)+e)/(c*(-d*e)^(1/2)+e)))/e+1/2*(dilog((c*(-d*e)^(1/2)-e*(c*x^(1/2)-1)-e)/(c*(-d*e)^(1/2)-e))+dilog((c*(-d*e)^(1/2)+e*(c*x^(1/2)-1)+e)/(c*(-d*e)^(1/2)+e)))/e-1/2*ln(1+c*x^(1/2))*ln(c^2*e*x+c^2*d)+e*(1/2*ln(1+c*x^(1/2))*(ln((c*(-d*e)^(1/2)-e*(1+c*x^(1/2))+e)/(c*(-d*e)^(1/2)+e))+ln((c*(-d*e)^(1/2)+e*(1+c*x^(1/2))-e)/(c*(-d*e)^(1/2)-e)))/e+1/2*(dilog((c*(-d*e)^(1/2)-e*(1+c*x^(1/2))+e)/(c*(-d*e)^(1/2)+e))+dilog((c*(-d*e)^(1/2)+e*(1+c*x^(1/2))-e)/(c*(-d*e)^(1/2)-e)))/e)-1/c^4/d^2*(-(c^2*d-2*e)/c/x^(1/2)-1/3*d/c/x^(3/2)+(-1/2*c^2*d+e)*ln(c*x^(1/2)-1)+(1/2*c^2*d-e)*ln(1+c*x^(1/2))))
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(d + ex)} dx = \int \frac{b \operatorname{arctanh}(c\sqrt{x}) + a}{(ex + d)x^3} dx$$

input

```
integrate((a+b*arctanh(c*x^(1/2)))/x^3/(e*x+d),x, algorithm="fricas")
```

output

```
integral((b*arctanh(c*sqrt(x)) + a)/(e*x^4 + d*x^3), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(d + ex)} dx = \text{Timed out}$$

input

```
integrate((a+b*atanh(c*x**(1/2)))/x**3/(e*x+d),x)
```

output

```
Timed out
```

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(d + ex)} dx = \int \frac{b \operatorname{artanh}(c\sqrt{x}) + a}{(ex + d)x^3} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^3/(e*x+d),x, algorithm="maxima")`

output `-1/2*a*(2*e^2*log(e*x + d)/d^3 - 2*e^2*log(x)/d^3 - (2*e*x - d)/(d^2*x^2)) + b*integrate(1/2*log(c*sqrt(x) + 1)/((e*x^(7/2) + d*x^(5/2))*sqrt(x)), x) - b*integrate(1/2*log(-c*sqrt(x) + 1)/((e*x^(7/2) + d*x^(5/2))*sqrt(x)), x)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(d + ex)} dx = \int \frac{b \operatorname{artanh}(c\sqrt{x}) + a}{(ex + d)x^3} dx$$

input `integrate((a+b*arctanh(c*x^(1/2)))/x^3/(e*x+d),x, algorithm="giac")`

output `integrate((b*arctanh(c*sqrt(x)) + a)/((e*x + d)*x^3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c\sqrt{x})}{x^3(d + ex)} dx = \int \frac{a + b \operatorname{atanh}(c\sqrt{x})}{x^3(d + ex)} dx$$

input `int((a + b*atanh(c*x^(1/2)))/(x^3*(d + e*x)),x)`

output `int((a + b*atanh(c*x^(1/2)))/(x^3*(d + e*x)), x)`

Reduce [F]

$$\int \frac{a + \operatorname{arctanh}(c\sqrt{x})}{x^3(d + ex)} dx = \text{Too large to display}$$

input `int((a+b*atanh(c*x^(1/2)))/x^3/(e*x+d),x)`

output

```
(3*atanh(sqrt(x)*c)*b*c**8*d**4*x**2 - 6*atanh(sqrt(x)*c)*b*c**6*d**3*e*x*
*2 - 3*atanh(sqrt(x)*c)*b*c**4*d**4 + 6*atanh(sqrt(x)*c)*b*c**2*d**3*e - 3
*sqrt(x)*b*c**7*d**4*x - sqrt(x)*b*c**5*d**4 + 6*sqrt(x)*b*c**5*d**3*e*x +
2*sqrt(x)*b*c**3*d**3*e + 6*int(atanh(sqrt(x)*c)/(c**4*d**3*x**3 + c**4*d
**2*e*x**4 - 2*c**2*d**2*e*x**3 - 2*c**2*d*e**2*x**4 + d*e**2*x**3 + e**3*
x**4),x)*b*c**4*d**5*e**2*x**2 - 12*int(atanh(sqrt(x)*c)/(c**4*d**3*x**3 +
c**4*d**2*e*x**4 - 2*c**2*d**2*e*x**3 - 2*c**2*d*e**2*x**4 + d*e**2*x**3
+ e**3*x**4),x)*b*c**2*d**4*e**3*x**2 + 6*int(atanh(sqrt(x)*c)/(c**4*d**3*
x**3 + c**4*d**2*e*x**4 - 2*c**2*d**2*e*x**3 - 2*c**2*d*e**2*x**4 + d*e**2
*x**3 + e**3*x**4),x)*b*d**3*e**4*x**2 + 6*int(atanh(sqrt(x)*c)/(c**4*d**3
*x**2 + c**4*d**2*e*x**3 - 2*c**2*d**2*e*x**2 - 2*c**2*d*e**2*x**3 + d*e**
2*x**2 + e**3*x**3),x)*b*c**6*d**5*e**2*x**2 - 12*int(atanh(sqrt(x)*c)/(c
**4*d**3*x**2 + c**4*d**2*e*x**3 - 2*c**2*d**2*e*x**2 - 2*c**2*d*e**2*x**3
+ d*e**2*x**2 + e**3*x**3),x)*b*c**4*d**4*e**3*x**2 + 6*int(atanh(sqrt(x)*
c)/(c**4*d**3*x**2 + c**4*d**2*e*x**3 - 2*c**2*d**2*e*x**2 - 2*c**2*d*e**2
*x**3 + d*e**2*x**2 + e**3*x**3),x)*b*c**2*d**3*e**4*x**2 - 6*int(atanh(sq
rt(x)*c)/(c**2*d**2*x**2 + c**2*d*e*x**3 - d*e*x**2 - e**2*x**3),x)*b*c**6
*d**5*e*x**2 + 12*int(atanh(sqrt(x)*c)/(c**2*d**2*x**2 + c**2*d*e*x**3 - d
*e*x**2 - e**2*x**3),x)*b*c**4*d**4*e**2*x**2 - 6*int(atanh(sqrt(x)*c)/(c
**2*d**2*x**2 + c**2*d*e*x**3 - d*e*x**2 - e**2*x**3),x)*b*c**2*d**3*e**...
```

CHAPTER 4

APPENDIX

4.1	Listing of Grading functions	447
4.2	Links to plain text integration problems used in this report for each CAS .	465

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```



```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
          finalresult={"C","Result contains complex when optimal does not."}
        ]
      ,(*ELSE*)(*result does not contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
          ]
        ]
      ,(*ELSE*)(*expnResult>expnOptimal*)
        If[FreeQ[result,Integrate] && FreeQ[result,Int],
          finalresult={"C","Result contains higher order function than in optimal. Order "
          ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```
  Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 5]],
  If [AppellFunctionQ [Head [expn]],
    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 6]],
  If [Head [expn] == RootSum,
    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 7]],
  If [Head [expn] == Integrate || Head [expn] == Int,
    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 8]],
  9]]]]]]]]]]]]]
```

```
ElementaryFunctionQ [func_] :=
  MemberQ [{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]
```

```
SpecialFunctionQ [func_] :=
  MemberQ [{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]
```

```
HypergeometricFunctionQ [func_] :=
  MemberQ [{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]
```

```
AppellFunctionQ [func_] :=
  MemberQ [{AppellF1}, func]
```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#                    if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#                    see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co

        fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```



```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```



```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file