

Computer Algebra Independent Integration Tests

Summer 2024

7-Inverse-hyperbolic-functions/7.3-Inverse-hyperbolic-
tangent/340-7.3.5

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 9:23am

Contents

| | | |
|----------|---|-----------|
| 1 | Introduction | 5 |
| 1.1 | Listing of CAS systems tested | 6 |
| 1.2 | Results | 7 |
| 1.3 | Time and leaf size Performance | 11 |
| 1.4 | Performance based on number of rules Rubi used | 13 |
| 1.5 | Performance based on number of steps Rubi used | 14 |
| 1.6 | Solved integrals histogram based on leaf size of result | 15 |
| 1.7 | Solved integrals histogram based on CPU time used | 16 |
| 1.8 | Leaf size vs. CPU time used | 17 |
| 1.9 | list of integrals with no known antiderivative | 18 |
| 1.10 | List of integrals solved by CAS but has no known antiderivative | 18 |
| 1.11 | list of integrals solved by CAS but failed verification | 18 |
| 1.12 | Timing | 19 |
| 1.13 | Verification | 19 |
| 1.14 | Important notes about some of the results | 20 |
| 1.15 | Current tree layout of integration tests | 23 |
| 1.16 | Design of the test system | 24 |
| 2 | detailed summary tables of results | 25 |
| 2.1 | List of integrals sorted by grade for each CAS | 26 |
| 2.2 | Detailed conclusion table per each integral for all CAS systems | 30 |
| 2.3 | Detailed conclusion table specific for Rubi results | 50 |
| 3 | Listing of integrals | 53 |
| 3.1 | $\int x^3 \operatorname{arctanh}(a + bx)^2 dx$ | 56 |
| 3.2 | $\int x^2 \operatorname{arctanh}(a + bx)^2 dx$ | 63 |
| 3.3 | $\int x \operatorname{arctanh}(a + bx)^2 dx$ | 70 |
| 3.4 | $\int \operatorname{arctanh}(a + bx)^2 dx$ | 77 |
| 3.5 | $\int \frac{\operatorname{arctanh}(a+bx)^2}{x} dx$ | 83 |
| 3.6 | $\int \frac{\operatorname{arctanh}(a+bx)^2}{x^2} dx$ | 90 |

| | | |
|------|--|-----|
| 3.7 | $\int \frac{\operatorname{arctanh}(a+bx)^2}{x^3} dx$ | 98 |
| 3.8 | $\int \frac{\operatorname{arctanh}(1+bx)^2}{x} dx$ | 106 |
| 3.9 | $\int (ce + dex)^3 (a + \operatorname{barctanh}(c + dx)) dx$ | 112 |
| 3.10 | $\int (ce + dex)^2 (a + \operatorname{barctanh}(c + dx)) dx$ | 121 |
| 3.11 | $\int (ce + dex)(a + \operatorname{barctanh}(c + dx)) dx$ | 129 |
| 3.12 | $\int \frac{a+\operatorname{barctanh}(c+dx)}{ce+dex} dx$ | 136 |
| 3.13 | $\int \frac{a+\operatorname{barctanh}(c+dx)}{(ce+dex)^2} dx$ | 141 |
| 3.14 | $\int \frac{a+\operatorname{barctanh}(c+dx)}{(ce+dex)^3} dx$ | 148 |
| 3.15 | $\int (ce + dex)^3 (a + \operatorname{barctanh}(c + dx))^2 dx$ | 155 |
| 3.16 | $\int (ce + dex)^2 (a + \operatorname{barctanh}(c + dx))^2 dx$ | 166 |
| 3.17 | $\int (ce + dex)(a + \operatorname{barctanh}(c + dx))^2 dx$ | 175 |
| 3.18 | $\int \frac{(a+\operatorname{barctanh}(c+dx))^2}{ce+dex} dx$ | 184 |
| 3.19 | $\int \frac{(a+\operatorname{barctanh}(c+dx))^2}{(ce+dex)^2} dx$ | 192 |
| 3.20 | $\int \frac{(a+\operatorname{barctanh}(c+dx))^2}{(ce+dex)^3} dx$ | 199 |
| 3.21 | $\int \frac{(a+\operatorname{barctanh}(c+dx))^2}{(ce+dex)^4} dx$ | 210 |
| 3.22 | $\int \frac{(a+\operatorname{barctanh}(c+dx))^2}{(ce+dex)^5} dx$ | 219 |
| 3.23 | $\int (ce + dex)^2 (a + \operatorname{barctanh}(c + dx))^3 dx$ | 231 |
| 3.24 | $\int (ce + dex)(a + \operatorname{barctanh}(c + dx))^3 dx$ | 242 |
| 3.25 | $\int \frac{(a+\operatorname{barctanh}(c+dx))^3}{ce+dex} dx$ | 251 |
| 3.26 | $\int \frac{(a+\operatorname{barctanh}(c+dx))^3}{(ce+dex)^2} dx$ | 259 |
| 3.27 | $\int \frac{(a+\operatorname{barctanh}(c+dx))^3}{(ce+dex)^3} dx$ | 267 |
| 3.28 | $\int \frac{(a+\operatorname{barctanh}(c+dx))^3}{(ce+dex)^4} dx$ | 276 |
| 3.29 | $\int \frac{\operatorname{arctanh}(1+x)}{2+2x} dx$ | 287 |
| 3.30 | $\int \frac{\operatorname{arctanh}(a+bx)}{\frac{a^d}{b}+dx} dx$ | 292 |
| 3.31 | $\int (e + fx)^3 (a + \operatorname{barctanh}(c + dx)) dx$ | 297 |
| 3.32 | $\int (e + fx)^2 (a + \operatorname{barctanh}(c + dx)) dx$ | 308 |
| 3.33 | $\int (e + fx)(a + \operatorname{barctanh}(c + dx)) dx$ | 317 |
| 3.34 | $\int (a + \operatorname{barctanh}(c + dx)) dx$ | 324 |
| 3.35 | $\int \frac{a+\operatorname{barctanh}(c+dx)}{e+fx} dx$ | 330 |
| 3.36 | $\int \frac{a+\operatorname{barctanh}(c+dx)}{(e+fx)^2} dx$ | 337 |
| 3.37 | $\int \frac{a+\operatorname{barctanh}(c+dx)}{(e+fx)^3} dx$ | 345 |
| 3.38 | $\int (e + fx)^3 (a + \operatorname{barctanh}(c + dx))^2 dx$ | 355 |
| 3.39 | $\int (e + fx)^2 (a + \operatorname{barctanh}(c + dx))^2 dx$ | 365 |
| 3.40 | $\int (e + fx)(a + \operatorname{barctanh}(c + dx))^2 dx$ | 375 |

| | | |
|------|---|-----|
| 3.41 | $\int (a + b \operatorname{arctanh}(c + dx))^2 dx$ | 383 |
| 3.42 | $\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{e + fx} dx$ | 390 |
| 3.43 | $\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^2} dx$ | 397 |
| 3.44 | $\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^3} dx$ | 406 |
| 3.45 | $\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx$ | 417 |
| 3.46 | $\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^3 dx$ | 427 |
| 3.47 | $\int (a + b \operatorname{arctanh}(c + dx))^3 dx$ | 435 |
| 3.48 | $\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{e + fx} dx$ | 442 |
| 3.49 | $\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(e + fx)^2} dx$ | 449 |
| 3.50 | $\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^3 dx$ | 460 |
| 3.51 | $\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^2 dx$ | 465 |
| 3.52 | $\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx)) dx$ | 470 |
| 3.53 | $\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^3} dx$ | 476 |
| 3.54 | $\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^2} dx$ | 486 |
| 3.55 | $\int \frac{\operatorname{arctanh}(a + bx)}{c + dx} dx$ | 493 |
| 3.56 | $\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x}} dx$ | 500 |
| 3.57 | $\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^2}} dx$ | 507 |
| 3.58 | $\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^3}} dx$ | 515 |
| 3.59 | $\int \frac{\operatorname{arctanh}(a + bx)}{1 - x^2} dx$ | 524 |
| 3.60 | $\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + f\sqrt{x}} dx$ | 530 |
| 3.61 | $\int \frac{\operatorname{arctanh}(a + bx)}{c + d\sqrt{x}} dx$ | 538 |
| 3.62 | $\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{\sqrt{x}}} dx$ | 546 |
| 3.63 | $\int \frac{x^3 (a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx$ | 557 |
| 3.64 | $\int \frac{x^2 (a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx$ | 565 |
| 3.65 | $\int \frac{x (a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx$ | 572 |
| 3.66 | $\int \frac{a + b \operatorname{arctanh}(c + dx)}{e - fx^2} dx$ | 579 |
| 3.67 | $\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e - fx^2)} dx$ | 585 |
| 3.68 | $\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2(e - fx^2)} dx$ | 592 |
| 3.69 | $\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3(e - fx^2)} dx$ | 599 |
| 3.70 | $\int \frac{x^3 (a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx$ | 607 |
| 3.71 | $\int \frac{x^2 (a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx$ | 617 |

| | | |
|----------|--|------------|
| 3.72 | $\int \frac{x(a+b\operatorname{arctanh}(c+dx))}{e+fx^3} dx$ | 627 |
| 3.73 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{e+fx^3} dx$ | 636 |
| 3.74 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{x(e+fx^3)} dx$ | 646 |
| 3.75 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{x^2(e+fx^3)} dx$ | 655 |
| 3.76 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{x^3(e+fx^3)} dx$ | 664 |
| 3.77 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{e+fx+gx^2} dx$ | 674 |
| 3.78 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{e+fx^2+gx^4} dx$ | 682 |
| 3.79 | $\int \frac{(ce+dex)(a+b\operatorname{arctanh}(c+dx))}{1-(c+dx)^2} dx$ | 691 |
| 4 | Appendix | 698 |
| 4.1 | Listing of Grading functions | 698 |
| 4.2 | Links to plain text integration problems used in this report for each CAS716 | |

CHAPTER 1

INTRODUCTION

| | | |
|------|---|----|
| 1.1 | Listing of CAS systems tested | 6 |
| 1.2 | Results | 7 |
| 1.3 | Time and leaf size Performance | 11 |
| 1.4 | Performance based on number of rules Rubi used | 13 |
| 1.5 | Performance based on number of steps Rubi used | 14 |
| 1.6 | Solved integrals histogram based on leaf size of result | 15 |
| 1.7 | Solved integrals histogram based on CPU time used | 16 |
| 1.8 | Leaf size vs. CPU time used | 17 |
| 1.9 | list of integrals with no known antiderivative | 18 |
| 1.10 | List of integrals solved by CAS but has no known antiderivative | 18 |
| 1.11 | list of integrals solved by CAS but failed verification | 18 |
| 1.12 | Timing | 19 |
| 1.13 | Verification | 19 |
| 1.14 | Important notes about some of the results | 20 |
| 1.15 | Current tree layout of integration tests | 23 |
| 1.16 | Design of the test system | 24 |

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [79]. This is test number [340].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

| System | % solved | % Failed |
|-------------|--------------|--------------|
| Rubi | 98.73 (78) | 1.27 (1) |
| Maple | 98.73 (78) | 1.27 (1) |
| Mathematica | 96.20 (76) | 3.80 (3) |
| Maxima | 44.30 (35) | 55.70 (44) |
| Fricas | 21.52 (17) | 78.48 (62) |
| Mupad | 21.52 (17) | 78.48 (62) |
| Giac | 21.52 (17) | 78.48 (62) |
| Reduce | 21.52 (17) | 78.48 (62) |
| Sympy | 18.99 (15) | 81.01 (64) |

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

| grade | description |
|-------|---|
| A | Integral was solved and antiderivative is optimal in quality and leaf size. |
| B | Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size. |
| C | Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not. |
| F | Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised. |


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


| System | % A grade | % B grade | % C grade | % F grade |
|-------------|-----------|-----------|-----------|-----------|
| Rubi | 96.203 | 0.000 | 0.000 | 3.797 |
| Mathematica | 68.354 | 3.797 | 21.519 | 6.329 |
| Maple | 48.101 | 18.987 | 29.114 | 3.797 |
| Maxima | 18.987 | 20.253 | 2.532 | 58.228 |
| Fricas | 6.329 | 12.658 | 0.000 | 81.013 |
| Sympy | 1.266 | 17.722 | 0.000 | 81.013 |
| Giac | 0.000 | 18.987 | 0.000 | 81.013 |
| Mupad | 0.000 | 18.987 | 0.000 | 81.013 |
| Reduce | 0.000 | 18.987 | 0.000 | 81.013 |

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

| System | Number failed | Percentage normal failure | Percentage time-out failure | Percentage exception failure |
|-------------|---------------|---------------------------|-----------------------------|------------------------------|
| Rubi | 1 | 100.00 | 0.00 | 0.00 |
| Maple | 1 | 100.00 | 0.00 | 0.00 |
| Mathematica | 3 | 100.00 | 0.00 | 0.00 |
| Maxima | 44 | 79.55 | 0.00 | 20.45 |
| Fricas | 62 | 100.00 | 0.00 | 0.00 |
| Mupad | 62 | 0.00 | 100.00 | 0.00 |
| Giac | 62 | 100.00 | 0.00 | 0.00 |
| Reduce | 62 | 100.00 | 0.00 | 0.00 |
| Sympy | 64 | 62.50 | 37.50 | 0.00 |

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

| System | Mean time (sec) |
|-------------|-----------------|
| Fricas | 0.14 |
| Giac | 0.15 |
| Reduce | 0.19 |
| Maxima | 0.25 |
| Rubi | 0.96 |
| Sympy | 2.34 |
| Maple | 3.06 |
| Mupad | 4.41 |
| Mathematica | 4.98 |

Table 1.5: Time performance for each CAS

| System | Mean size | Normalized mean | Median size | Normalized median |
|-------------|-----------|-----------------|-------------|-------------------|
| Fricas | 232.24 | 2.20 | 159.00 | 2.09 |
| Rubi | 330.92 | 1.07 | 216.50 | 1.00 |
| Maxima | 347.63 | 3.08 | 262.00 | 1.98 |
| Mupad | 501.76 | 4.00 | 237.00 | 2.50 |
| Mathematica | 542.18 | 1.80 | 303.00 | 1.36 |
| Giac | 608.12 | 5.11 | 351.00 | 4.16 |
| Maple | 887.64 | 3.16 | 375.50 | 1.53 |
| Sympy | 1944.73 | 13.00 | 313.00 | 3.48 |
| Reduce | 4968.82 | 232.50 | 304.00 | 2.65 |

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{50, 51}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {10, 13, 15, 20, 22, 28, 54, 70, 76, 78}

Mathematica {2, 5, 6, 7, 18, 38, 39, 42, 43, 44, 45, 49, 54, 56, 64, 68, 71, 72, 73, 74, 75, 76, 78}

Maple {5, 8, 18, 23, 25, 26, 27, 28, 42, 45, 46, 48, 49, 53, 58, 70, 71, 72, 73, 74, 75, 76, 78}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

| | | |
|-----|---|----|
| 2.1 | List of integrals sorted by grade for each CAS | 26 |
| 2.2 | Detailed conclusion table per each integral for all CAS systems | 30 |
| 2.3 | Detailed conclusion table specific for Rubi results | 50 |

2.1 List of integrals sorted by grade for each CAS

| | |
|------------------|----|
| Rubi | 26 |
| Mma | 26 |
| Maple | 27 |
| Fricas | 27 |
| Maxima | 27 |
| Giac | 28 |
| Mupad | 28 |
| Sympy | 28 |
| Reduce | 29 |

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 53, 54, 55, 56, 57, 58, 59, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79 }
}

B grade { }

C grade { }

F normal fail { 60 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 3, 4, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 24, 26, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 46, 47, 53, 54, 55, 57, 58, 59, 60, 61, 62, 63, 65, 66, 67, 69, 71, 73, 74, 77, 78, 79 }
}

B grade { 2, 39, 45 }

C grade { 5, 6, 7, 18, 25, 27, 28, 42, 43, 44, 49, 56, 64, 68, 72, 75, 76 }

F normal fail { 48, 52, 70 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 6, 7, 9, 10, 11, 12, 13, 14, 15, 16, 21, 22, 29, 30, 33, 34, 35, 36, 37, 41, 43, 44, 54, 55, 56, 57, 59, 60, 61, 62, 63, 65, 69, 79 }

B grade { 17, 19, 20, 24, 31, 32, 38, 39, 40, 47, 64, 66, 67, 68, 77 }

C grade { 5, 8, 18, 23, 25, 26, 27, 28, 42, 45, 46, 48, 49, 53, 58, 70, 71, 72, 73, 74, 75, 76, 78 }

F normal fail { 52 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 11, 13, 14, 33, 34 }

B grade { 9, 10, 15, 17, 20, 22, 31, 32, 36, 37 }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 12, 16, 18, 19, 21, 23, 24, 25, 26, 27, 28, 29, 30, 35, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maxima

A grade { 1, 2, 3, 4, 6, 7, 13, 32, 33, 34, 36, 37, 55, 56, 59 }

B grade { 9, 10, 11, 14, 15, 16, 17, 20, 22, 24, 29, 30, 31, 38, 39, 40 }

C grade { 54, 57 }

F normal fail { 5, 8, 12, 18, 19, 21, 23, 25, 26, 27, 28, 35, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 53, 58, 60, 61, 62, 63, 65, 67, 69, 71, 74, 78, 79 }

F(-1) timedout fail { }

F(-2) exception fail { 64, 66, 68, 70, 72, 73, 75, 76, 77 }

Giac

A grade { }

B grade { 9, 10, 11, 13, 14, 15, 17, 20, 22, 31, 32, 33, 34, 36, 37 }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 12, 16, 18, 19, 21, 23, 24, 25, 26, 27, 28, 29, 30, 35, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 9, 10, 11, 13, 14, 15, 17, 20, 22, 31, 32, 33, 34, 36, 37 }

C grade { }

F normal fail { }

F(-1) timedout fail { 1, 2, 3, 4, 5, 6, 7, 8, 12, 16, 18, 19, 21, 23, 24, 25, 26, 27, 28, 29, 30, 35, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79 }

F(-2) exception fail { }

Sympy

A grade { 34 }

B grade { 9, 10, 11, 13, 14, 15, 17, 20, 22, 31, 32, 33, 36, 37 }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 12, 16, 18, 19, 21, 23, 24, 25, 26, 27, 28, 29, 30, 35, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 55, 56, 59, 67, 79 }

F(-1) timedout fail { 50, 51, 53, 54, 57, 58, 60, 61, 62, 63, 64, 65, 66, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78 }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 9, 10, 11, 13, 14, 15, 17, 20, 22, 31, 32, 33, 34, 36, 37 }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 12, 16, 18, 19, 21, 23, 24, 25, 26, 27, 28, 29, 30, 35, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

| Problem 1 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|----------|----------|----------|----------|--------------|
| grade | N/A | A | A | A | A | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 263 | 244 | 187 | 449 | 320 | 0 | 0 | 0 | 257 | 0 |
| N.S. | 1 | 0.93 | 0.71 | 1.71 | 1.22 | 0.00 | 0.00 | 0.00 | 0.98 | 0.00 |
| time (sec) | N/A | 0.538 | 0.971 | 1.289 | 0.035 | 0.000 | 0.000 | 0.000 | 0.179 | 0.000 |

| Problem 2 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-----------|-------|--------|----------|----------|----------|----------|--------------|
| grade | N/A | A | B | A | A | F | F | F | F | F(-1) |
| verified | N/A | Yes | No | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 204 | 193 | 463 | 350 | 259 | 0 | 0 | 0 | 231 | 0 |
| N.S. | 1 | 0.95 | 2.27 | 1.72 | 1.27 | 0.00 | 0.00 | 0.00 | 1.13 | 0.00 |
| time (sec) | N/A | 0.499 | 1.464 | 0.256 | 0.034 | 0.000 | 0.000 | 0.000 | 0.172 | 0.000 |

| Problem 3 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|----------|----------|----------|----------|--------------|
| grade | N/A | A | A | A | A | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 136 | 127 | 98 | 244 | 202 | 0 | 0 | 0 | 47 | 0 |
| N.S. | 1 | 0.93 | 0.72 | 1.79 | 1.49 | 0.00 | 0.00 | 0.00 | 0.35 | 0.00 |
| time (sec) | N/A | 0.436 | 0.248 | 0.210 | 0.035 | 0.000 | 0.000 | 0.000 | 0.170 | 0.000 |

| Problem 4 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|----------|----------|----------|----------|--------------|
| grade | N/A | A | A | A | A | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 81 | 76 | 55 | 91 | 139 | 0 | 0 | 0 | 116 | 0 |
| N.S. | 1 | 0.94 | 0.68 | 1.12 | 1.72 | 0.00 | 0.00 | 0.00 | 1.43 | 0.00 |
| time (sec) | N/A | 0.470 | 0.061 | 0.549 | 0.034 | 0.000 | 0.000 | 0.000 | 0.162 | 0.000 |

| Problem 5 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-----------|-----------|----------|----------|----------|----------|----------|--------------|
| grade | N/A | A | C | C | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | No | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 148 | 148 | 753 | 901 | 0 | 0 | 0 | 0 | 14 | 0 |
| N.S. | 1 | 1.00 | 5.09 | 6.09 | 0.00 | 0.00 | 0.00 | 0.00 | 0.09 | 0.00 |
| time (sec) | N/A | 0.341 | 2.260 | 32.319 | 0.000 | 0.000 | 0.000 | 0.000 | 0.158 | 0.000 |

| Problem 6 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-----------|-------|--------|----------|----------|----------|----------|--------------|
| grade | N/A | A | C | A | A | F | F | F | F | F(-1) |
| verified | N/A | Yes | No | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 251 | 257 | 208 | 301 | 244 | 0 | 0 | 0 | 252 | 0 |
| N.S. | 1 | 1.02 | 0.83 | 1.20 | 0.97 | 0.00 | 0.00 | 0.00 | 1.00 | 0.00 |
| time (sec) | N/A | 1.047 | 0.934 | 1.596 | 0.039 | 0.000 | 0.000 | 0.000 | 0.163 | 0.000 |

| Problem 7 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-----------|-------|--------|----------|----------|----------|----------|--------------|
| grade | N/A | A | C | A | A | F | F | F | F | F(-1) |
| verified | N/A | Yes | No | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 370 | 347 | 271 | 385 | 360 | 0 | 0 | 0 | 605 | 0 |
| N.S. | 1 | 0.94 | 0.73 | 1.04 | 0.97 | 0.00 | 0.00 | 0.00 | 1.64 | 0.00 |
| time (sec) | N/A | 1.107 | 1.591 | 0.273 | 0.047 | 0.000 | 0.000 | 0.000 | 0.174 | 0.000 |

| Problem 8 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | C | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 56 | 61 | 75 | 158 | 0 | 0 | 0 | 0 | 14 | 0 |
| N.S. | 1 | 1.09 | 1.34 | 2.82 | 0.00 | 0.00 | 0.00 | 0.00 | 0.25 | 0.00 |
| time (sec) | N/A | 0.521 | 0.111 | 9.164 | 0.000 | 0.000 | 0.000 | 0.000 | 0.171 | 0.000 |

| Problem 9 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | B | B | B | B | B | B |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 72 | 59 | 78 | 74 | 357 | 159 | 231 | 363 | 163 | 414 |
| N.S. | 1 | 0.82 | 1.08 | 1.03 | 4.96 | 2.21 | 3.21 | 5.04 | 2.26 | 5.75 |
| time (sec) | N/A | 0.302 | 0.080 | 0.826 | 0.035 | 0.083 | 1.473 | 0.144 | 0.159 | 4.629 |

| Problem 10 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | B | B | B | B | B | B |
| verified | N/A | No | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 69 | 54 | 59 | 67 | 225 | 144 | 180 | 322 | 126 | 237 |
| N.S. | 1 | 0.78 | 0.86 | 0.97 | 3.26 | 2.09 | 2.61 | 4.67 | 1.83 | 3.43 |
| time (sec) | N/A | 0.305 | 0.067 | 0.368 | 0.033 | 0.095 | 1.216 | 0.136 | 0.154 | 0.643 |

| Problem 11 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | B | A | B | B | B | B |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 48 | 46 | 77 | 61 | 113 | 73 | 95 | 180 | 71 | 73 |
| N.S. | 1 | 0.96 | 1.60 | 1.27 | 2.35 | 1.52 | 1.98 | 3.75 | 1.48 | 1.52 |
| time (sec) | N/A | 0.261 | 0.029 | 0.191 | 0.026 | 0.069 | 0.649 | 0.133 | 0.153 | 5.202 |

| Problem 12 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 54 | 43 | 54 | 54 | 0 | 0 | 0 | 0 | 35 | 0 |
| N.S. | 1 | 0.80 | 1.00 | 1.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.65 | 0.00 |
| time (sec) | N/A | 0.287 | 0.021 | 0.460 | 0.000 | 0.000 | 0.000 | 0.000 | 0.165 | 0.000 |

| Problem 13 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | A | A | B | B | B | B |
| verified | N/A | No | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 63 | 53 | 69 | 64 | 95 | 85 | 219 | 152 | 155 | 122 |
| N.S. | 1 | 0.84 | 1.10 | 1.02 | 1.51 | 1.35 | 3.48 | 2.41 | 2.46 | 1.94 |
| time (sec) | N/A | 0.292 | 0.071 | 0.529 | 0.030 | 0.094 | 1.100 | 0.122 | 0.179 | 4.679 |

| Problem 14 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | B | A | B | B | B | B |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 63 | 50 | 100 | 67 | 131 | 88 | 313 | 194 | 252 | 67 |
| N.S. | 1 | 0.79 | 1.59 | 1.06 | 2.08 | 1.40 | 4.97 | 3.08 | 4.00 | 1.06 |
| time (sec) | N/A | 0.290 | 0.061 | 0.759 | 0.031 | 0.089 | 1.451 | 0.125 | 0.168 | 5.043 |

| Problem 15 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | B | B | B | B | B | B |
| verified | N/A | No | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 159 | 141 | 148 | 261 | 827 | 383 | 581 | 733 | 421 | 1730 |
| N.S. | 1 | 0.89 | 0.93 | 1.64 | 5.20 | 2.41 | 3.65 | 4.61 | 2.65 | 10.88 |
| time (sec) | N/A | 0.917 | 0.121 | 1.451 | 0.256 | 0.109 | 2.406 | 0.169 | 0.168 | 5.601 |

| Problem 16 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|----------|----------|----------|----------|--------------|
| grade | N/A | A | A | A | B | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 179 | 143 | 150 | 251 | 619 | 0 | 0 | 0 | 321 | 0 |
| N.S. | 1 | 0.80 | 0.84 | 1.40 | 3.46 | 0.00 | 0.00 | 0.00 | 1.79 | 0.00 |
| time (sec) | N/A | 0.871 | 0.316 | 1.989 | 0.236 | 0.000 | 0.000 | 0.000 | 0.164 | 0.000 |

| Problem 17 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | B | B | B | B | B | B | B |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 95 | 91 | 134 | 221 | 316 | 191 | 238 | 351 | 193 | 432 |
| N.S. | 1 | 0.96 | 1.41 | 2.33 | 3.33 | 2.01 | 2.51 | 3.69 | 2.03 | 4.55 |
| time (sec) | N/A | 0.521 | 0.089 | 0.314 | 0.230 | 0.095 | 1.100 | 0.145 | 0.162 | 0.823 |

| Problem 18 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-----------|-----------|----------|----------|----------|----------|----------|--------------|
| grade | N/A | A | C | C | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | No | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 168 | 158 | 424 | 705 | 0 | 0 | 0 | 0 | 62 | 0 |
| N.S. | 1 | 0.94 | 2.52 | 4.20 | 0.00 | 0.00 | 0.00 | 0.00 | 0.37 | 0.00 |
| time (sec) | N/A | 0.839 | 0.230 | 12.455 | 0.000 | 0.000 | 0.000 | 0.000 | 0.164 | 0.000 |

| Problem 19 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|----------|----------|----------|----------|----------|--------------|
| grade | N/A | A | A | B | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 104 | 95 | 126 | 252 | 0 | 0 | 0 | 0 | 554 | 0 |
| N.S. | 1 | 0.91 | 1.21 | 2.42 | 0.00 | 0.00 | 0.00 | 0.00 | 5.33 | 0.00 |
| time (sec) | N/A | 0.574 | 0.237 | 1.154 | 0.000 | 0.000 | 0.000 | 0.000 | 0.168 | 0.000 |

| Problem 20 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | B | B | B | B | B | B | B |
| verified | N/A | No | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 119 | 98 | 136 | 238 | 329 | 270 | 1102 | 375 | 503 | 776 |
| N.S. | 1 | 0.82 | 1.14 | 2.00 | 2.76 | 2.27 | 9.26 | 3.15 | 4.23 | 6.52 |
| time (sec) | N/A | 0.628 | 0.179 | 0.902 | 0.045 | 0.118 | 1.536 | 0.140 | 0.174 | 5.740 |

| Problem 21 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 180 | 141 | 218 | 307 | 0 | 0 | 0 | 0 | 0 | 0 |
| N.S. | 1 | 0.78 | 1.21 | 1.71 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| time (sec) | N/A | 0.917 | 0.408 | 2.253 | 0.000 | 0.000 | 0.000 | 0.000 | 0.187 | 0.000 |

| Problem 22 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | B | B | B | B | B | B |
| verified | N/A | No | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 172 | 157 | 218 | 273 | 613 | 547 | 3516 | 730 | 989 | 2746 |
| N.S. | 1 | 0.91 | 1.27 | 1.59 | 3.56 | 3.18 | 20.44 | 4.24 | 5.75 | 15.97 |
| time (sec) | N/A | 1.023 | 0.232 | 2.063 | 0.070 | 0.112 | 3.458 | 0.145 | 0.171 | 7.052 |

| Problem 23 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|--------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | C | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 263 | 222 | 336 | 1099 | 0 | 0 | 0 | 0 | 593 | 0 |
| N.S. | 1 | 0.84 | 1.28 | 4.18 | 0.00 | 0.00 | 0.00 | 0.00 | 2.25 | 0.00 |
| time (sec) | N/A | 1.632 | 0.474 | 11.551 | 0.000 | 0.000 | 0.000 | 0.000 | 0.170 | 0.000 |

| Problem 24 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|----------|----------|----------|----------|--------------|
| grade | N/A | A | A | B | B | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 160 | 143 | 220 | 931 | 629 | 0 | 0 | 0 | 327 | 0 |
| N.S. | 1 | 0.89 | 1.38 | 5.82 | 3.93 | 0.00 | 0.00 | 0.00 | 2.04 | 0.00 |
| time (sec) | N/A | 1.007 | 0.168 | 3.312 | 0.245 | 0.000 | 0.000 | 0.000 | 0.160 | 0.000 |

| Problem 25 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-----------|----------|----------|----------|----------|----------|--------------|
| grade | N/A | A | C | C | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 257 | 231 | 581 | 1442 | 0 | 0 | 0 | 0 | 89 | 0 |
| N.S. | 1 | 0.90 | 2.26 | 5.61 | 0.00 | 0.00 | 0.00 | 0.00 | 0.35 | 0.00 |
| time (sec) | N/A | 1.187 | 0.435 | 2.527 | 0.000 | 0.000 | 0.000 | 0.000 | 0.176 | 0.000 |

| Problem 26 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-----------|----------|----------|----------|----------|----------|--------------|
| grade | N/A | A | A | C | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 143 | 129 | 232 | 1466 | 0 | 0 | 0 | 0 | 1156 | 0 |
| N.S. | 1 | 0.90 | 1.62 | 10.25 | 0.00 | 0.00 | 0.00 | 0.00 | 8.08 | 0.00 |
| time (sec) | N/A | 0.860 | 0.451 | 2.493 | 0.000 | 0.000 | 0.000 | 0.000 | 0.192 | 0.000 |

| Problem 27 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-----------|----------|----------|----------|----------|----------|--------------|
| grade | N/A | A | C | C | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 166 | 143 | 335 | 4949 | 0 | 0 | 0 | 0 | 1236 | 0 |
| N.S. | 1 | 0.86 | 2.02 | 29.81 | 0.00 | 0.00 | 0.00 | 0.00 | 7.45 | 0.00 |
| time (sec) | N/A | 1.030 | 0.933 | 3.484 | 0.000 | 0.000 | 0.000 | 0.000 | 0.177 | 0.000 |

| Problem 28 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | C | C | F | F | F | F | F | F(-1) |
| verified | N/A | No | Yes | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 269 | 220 | 393 | 1768 | 0 | 0 | 0 | 0 | 0 | 0 |
| N.S. | 1 | 0.82 | 1.46 | 6.57 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| time (sec) | N/A | 1.894 | 0.952 | 7.735 | 0.000 | 0.000 | 0.000 | 0.000 | 0.201 | 0.000 |

| Problem 29 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | B | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 21 | 25 | 31 | 14 | 58 | 0 | 0 | 0 | 14 | 0 |
| N.S. | 1 | 1.19 | 1.48 | 0.67 | 2.76 | 0.00 | 0.00 | 0.00 | 0.67 | 0.00 |
| time (sec) | N/A | 0.237 | 0.004 | 0.934 | 0.025 | 0.000 | 0.000 | 0.000 | 0.153 | 0.000 |

| Problem 30 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | B | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 32 | 30 | 52 | 29 | 132 | 0 | 0 | 0 | 21 | 0 |
| N.S. | 1 | 0.94 | 1.62 | 0.91 | 4.12 | 0.00 | 0.00 | 0.00 | 0.66 | 0.00 |
| time (sec) | N/A | 0.258 | 0.007 | 0.632 | 0.032 | 0.000 | 0.000 | 0.000 | 0.161 | 0.000 |

| Problem 31 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | B | B | B | B | B | B | B |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 168 | 167 | 270 | 599 | 333 | 386 | 644 | 2336 | 518 | 737 |
| N.S. | 1 | 0.99 | 1.61 | 3.57 | 1.98 | 2.30 | 3.83 | 13.90 | 3.08 | 4.39 |
| time (sec) | N/A | 0.492 | 0.136 | 0.760 | 0.032 | 0.101 | 2.022 | 0.195 | 0.154 | 4.696 |

| Problem 32 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | B | A | B | B | B | B | B |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 120 | 128 | 174 | 351 | 207 | 242 | 369 | 976 | 304 | 381 |
| N.S. | 1 | 1.07 | 1.45 | 2.92 | 1.72 | 2.02 | 3.08 | 8.13 | 2.53 | 3.18 |
| time (sec) | N/A | 0.415 | 0.084 | 0.530 | 0.033 | 0.104 | 1.843 | 0.152 | 0.163 | 4.207 |

| Problem 33 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | A | A | B | B | B | B |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 97 | 106 | 138 | 122 | 109 | 134 | 173 | 341 | 138 | 136 |
| N.S. | 1 | 1.09 | 1.42 | 1.26 | 1.12 | 1.38 | 1.78 | 3.52 | 1.42 | 1.40 |
| time (sec) | N/A | 0.365 | 0.039 | 0.263 | 0.025 | 0.104 | 0.887 | 0.133 | 0.165 | 4.740 |

| Problem 34 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | A | A | A | B | B | B |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 40 | 40 | 48 | 37 | 36 | 61 | 46 | 200 | 45 | 48 |
| N.S. | 1 | 1.00 | 1.20 | 0.92 | 0.90 | 1.52 | 1.15 | 5.00 | 1.12 | 1.20 |
| time (sec) | N/A | 0.187 | 0.010 | 0.287 | 0.026 | 0.086 | 0.253 | 0.118 | 0.157 | 4.281 |

| Problem 35 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|----------|----------|----------|----------|----------|--------------|
| grade | N/A | A | A | A | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 130 | 148 | 148 | 192 | 0 | 0 | 0 | 0 | 32 | 0 |
| N.S. | 1 | 1.14 | 1.14 | 1.48 | 0.00 | 0.00 | 0.00 | 0.00 | 0.25 | 0.00 |
| time (sec) | N/A | 0.565 | 0.038 | 6.214 | 0.000 | 0.000 | 0.000 | 0.000 | 0.169 | 0.000 |

| Problem 36 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | A | B | B | B | B | B |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 114 | 121 | 125 | 137 | 121 | 263 | 1605 | 474 | 412 | 170 |
| N.S. | 1 | 1.06 | 1.10 | 1.20 | 1.06 | 2.31 | 14.08 | 4.16 | 3.61 | 1.49 |
| time (sec) | N/A | 0.419 | 0.115 | 0.757 | 0.039 | 0.204 | 4.566 | 0.131 | 0.157 | 4.511 |

| Problem 37 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|--------|-------|--------|-------|
| grade | N/A | A | A | A | A | B | B | B | B | B |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 167 | 170 | 174 | 197 | 291 | 834 | 19859 | 2567 | 2159 | 417 |
| N.S. | 1 | 1.02 | 1.04 | 1.18 | 1.74 | 4.99 | 118.92 | 15.37 | 12.93 | 2.50 |
| time (sec) | N/A | 0.498 | 0.201 | 1.286 | 0.042 | 0.692 | 11.076 | 0.192 | 0.166 | 5.818 |

| Problem 38 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-----------|-------|--------|----------|----------|----------|----------|--------------|
| grade | N/A | A | A | B | B | F | F | F | F | F(-1) |
| verified | N/A | Yes | No | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 562 | 547 | 1082 | 2318 | 1363 | 0 | 0 | 0 | 1429 | 0 |
| N.S. | 1 | 0.97 | 1.93 | 4.12 | 2.43 | 0.00 | 0.00 | 0.00 | 2.54 | 0.00 |
| time (sec) | N/A | 1.138 | 4.812 | 1.188 | 0.252 | 0.000 | 0.000 | 0.000 | 0.167 | 0.000 |

| Problem 39 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-----------|-------|--------|----------|----------|----------|----------|--------------|
| grade | N/A | A | B | B | B | F | F | F | F | F(-1) |
| verified | N/A | Yes | No | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 374 | 374 | 795 | 1411 | 806 | 0 | 0 | 0 | 792 | 0 |
| N.S. | 1 | 1.00 | 2.13 | 3.77 | 2.16 | 0.00 | 0.00 | 0.00 | 2.12 | 0.00 |
| time (sec) | N/A | 0.808 | 2.255 | 0.608 | 0.250 | 0.000 | 0.000 | 0.000 | 0.179 | 0.000 |

| Problem 40 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|----------|----------|----------|----------|--------------|
| grade | N/A | A | A | B | B | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 221 | 231 | 271 | 453 | 415 | 0 | 0 | 0 | 350 | 0 |
| N.S. | 1 | 1.05 | 1.23 | 2.05 | 1.88 | 0.00 | 0.00 | 0.00 | 1.58 | 0.00 |
| time (sec) | N/A | 0.663 | 0.837 | 0.426 | 0.233 | 0.000 | 0.000 | 0.000 | 0.176 | 0.000 |

| Problem 41 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|----------|----------|----------|----------|----------|--------------|
| grade | N/A | A | A | A | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 97 | 93 | 103 | 134 | 0 | 0 | 0 | 0 | 106 | 0 |
| N.S. | 1 | 0.96 | 1.06 | 1.38 | 0.00 | 0.00 | 0.00 | 0.00 | 1.09 | 0.00 |
| time (sec) | N/A | 0.552 | 0.133 | 0.715 | 0.000 | 0.000 | 0.000 | 0.000 | 0.175 | 0.000 |

| Problem 42 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-----------|-----------|----------|----------|----------|----------|----------|--------------|
| grade | N/A | A | C | C | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | No | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 214 | 241 | 3808 | 1700 | 0 | 0 | 0 | 0 | 59 | 0 |
| N.S. | 1 | 1.13 | 17.79 | 7.94 | 0.00 | 0.00 | 0.00 | 0.00 | 0.28 | 0.00 |
| time (sec) | N/A | 0.436 | 90.239 | 39.939 | 0.000 | 0.000 | 0.000 | 0.000 | 0.172 | 0.000 |

| Problem 43 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-----------|-------|----------|----------|----------|----------|----------|--------------|
| grade | N/A | A | C | A | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | No | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 401 | 491 | 419 | 592 | 0 | 0 | 0 | 0 | 0 | 0 |
| N.S. | 1 | 1.22 | 1.04 | 1.48 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| time (sec) | N/A | 1.906 | 5.058 | 3.580 | 0.000 | 0.000 | 0.000 | 0.000 | 0.223 | 0.000 |

| Problem 44 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|--------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | C | A | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | No | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 621 | 739 | 1318 | 870 | 0 | 0 | 0 | 0 | 0 | 0 |
| N.S. | 1 | 1.19 | 2.12 | 1.40 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| time (sec) | N/A | 2.192 | 11.299 | 2.013 | 0.000 | 0.000 | 0.000 | 0.000 | 0.417 | 0.000 |

| Problem 45 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|--------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | B | C | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | No | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 546 | 533 | 1646 | 10013 | 0 | 0 | 0 | 0 | 1397 | 0 |
| N.S. | 1 | 0.98 | 3.01 | 18.34 | 0.00 | 0.00 | 0.00 | 0.00 | 2.56 | 0.00 |
| time (sec) | N/A | 1.221 | 6.606 | 42.682 | 0.000 | 0.000 | 0.000 | 0.000 | 0.376 | 0.000 |

| Problem 46 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | C | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 326 | 324 | 566 | 10425 | 0 | 0 | 0 | 0 | 584 | 0 |
| N.S. | 1 | 0.99 | 1.74 | 31.98 | 0.00 | 0.00 | 0.00 | 0.00 | 1.79 | 0.00 |
| time (sec) | N/A | 0.947 | 3.188 | 5.994 | 0.000 | 0.000 | 0.000 | 0.000 | 0.176 | 0.000 |

| Problem 47 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | B | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 132 | 130 | 194 | 265 | 0 | 0 | 0 | 0 | 170 | 0 |
| N.S. | 1 | 0.98 | 1.47 | 2.01 | 0.00 | 0.00 | 0.00 | 0.00 | 1.29 | 0.00 |
| time (sec) | N/A | 0.742 | 0.198 | 1.374 | 0.000 | 0.000 | 0.000 | 0.000 | 0.173 | 0.000 |

| Problem 48 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | F | C | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | N/A | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 308 | 344 | 0 | 3441 | 0 | 0 | 0 | 0 | 86 | 0 |
| N.S. | 1 | 1.12 | 0.00 | 11.17 | 0.00 | 0.00 | 0.00 | 0.00 | 0.28 | 0.00 |
| time (sec) | N/A | 0.479 | 0.000 | 5.136 | 0.000 | 0.000 | 0.000 | 0.000 | 0.201 | 0.000 |

| Problem 49 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|--------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | C | C | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | No | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 634 | 1085 | 3878 | 5109 | 0 | 0 | 0 | 0 | 0 | 0 |
| N.S. | 1 | 1.71 | 6.12 | 8.06 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| time (sec) | N/A | 2.796 | 90.093 | 3.066 | 0.000 | 0.000 | 0.000 | 0.000 | 0.678 | 0.000 |

| Problem 50 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|---------|-------|
| grade | N/A | N/A | N/A | N/A | N/A | N/A | F(-1) | N/A | N/A | N/A |
| verified | N/A | N/A | N/A | N/A | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 20 | 20 | 22 | 20 | 432 | 52 | 0 | 22 | 55220 | 22 |
| N.S. | 1 | 1.00 | 1.10 | 1.00 | 21.60 | 2.60 | 0.00 | 1.10 | 2761.00 | 1.10 |
| time (sec) | N/A | 0.314 | 3.405 | 0.247 | 3.614 | 0.098 | 0.000 | 0.232 | 0.438 | 3.688 |

| Problem 51 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|---------|-------|
| grade | N/A | N/A | N/A | N/A | N/A | N/A | F(-1) | N/A | N/A | N/A |
| verified | N/A | N/A | N/A | N/A | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 20 | 20 | 22 | 20 | 262 | 36 | 0 | 22 | 22801 | 22 |
| N.S. | 1 | 1.00 | 1.10 | 1.00 | 13.10 | 1.80 | 0.00 | 1.10 | 1140.05 | 1.10 |
| time (sec) | N/A | 0.318 | 0.275 | 0.232 | 2.183 | 0.087 | 0.000 | 0.169 | 0.276 | 3.628 |

| Problem 52 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|----------|----------|----------|----------|----------|----------|----------|--------------|
| grade | N/A | A | F | F | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | N/A | N/A | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 162 | 222 | 0 | 0 | 0 | 0 | 0 | 0 | 1358 | 0 |
| N.S. | 1 | 1.37 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 8.38 | 0.00 |
| time (sec) | N/A | 0.494 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.201 | 0.000 |

| Problem 53 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-----------|----------|----------|--------------|----------|----------|--------------|
| grade | N/A | A | A | C | F | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | Yes | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 649 | 800 | 623 | 259 | 0 | 0 | 0 | 0 | 18 | 0 |
| N.S. | 1 | 1.23 | 0.96 | 0.40 | 0.00 | 0.00 | 0.00 | 0.00 | 0.03 | 0.00 |
| time (sec) | N/A | 1.556 | 0.484 | 0.536 | 0.000 | 0.000 | 0.000 | 0.000 | 41.956 | 0.000 |

| Problem 54 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-----------|-----------|-------|--------|----------|--------------|----------|----------|--------------|
| grade | N/A | A | A | A | C | F | F(-1) | F | F | F(-1) |
| verified | N/A | No | No | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 287 | 491 | 365 | 444 | 591 | 0 | 0 | 0 | 18 | 0 |
| N.S. | 1 | 1.71 | 1.27 | 1.55 | 2.06 | 0.00 | 0.00 | 0.00 | 0.06 | 0.00 |
| time (sec) | N/A | 0.915 | 0.196 | 0.471 | 0.238 | 0.000 | 0.000 | 0.000 | 0.176 | 0.000 |

| Problem 55 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|----------|----------|----------|----------|--------------|
| grade | N/A | A | A | A | A | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 120 | 138 | 138 | 181 | 192 | 0 | 0 | 0 | 16 | 0 |
| N.S. | 1 | 1.15 | 1.15 | 1.51 | 1.60 | 0.00 | 0.00 | 0.00 | 0.13 | 0.00 |
| time (sec) | N/A | 0.536 | 0.016 | 0.449 | 0.033 | 0.000 | 0.000 | 0.000 | 0.186 | 0.000 |

| Problem 56 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | C | A | A | F | F | F | F | F(-1) |
| verified | N/A | Yes | No | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 186 | 191 | 394 | 259 | 192 | 0 | 0 | 0 | 17 | 0 |
| N.S. | 1 | 1.03 | 2.12 | 1.39 | 1.03 | 0.00 | 0.00 | 0.00 | 0.09 | 0.00 |
| time (sec) | N/A | 0.517 | 8.765 | 0.483 | 0.035 | 0.000 | 0.000 | 0.000 | 0.192 | 0.000 |

| Problem 57 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | C | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 545 | 560 | 555 | 581 | 651 | 0 | 0 | 0 | 21 | 0 |
| N.S. | 1 | 1.03 | 1.02 | 1.07 | 1.19 | 0.00 | 0.00 | 0.00 | 0.04 | 0.00 |
| time (sec) | N/A | 1.158 | 0.374 | 1.063 | 0.215 | 0.000 | 0.000 | 0.000 | 0.186 | 0.000 |

| Problem 58 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | C | F | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | Yes | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 832 | 847 | 791 | 366 | 0 | 0 | 0 | 0 | 21 | 0 |
| N.S. | 1 | 1.02 | 0.95 | 0.44 | 0.00 | 0.00 | 0.00 | 0.00 | 0.03 | 0.00 |
| time (sec) | N/A | 1.573 | 0.668 | 0.561 | 0.000 | 0.000 | 0.000 | 0.000 | 0.201 | 0.000 |

| Problem 59 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | A | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 135 | 213 | 203 | 184 | 198 | 0 | 0 | 0 | 18 | 0 |
| N.S. | 1 | 1.58 | 1.50 | 1.36 | 1.47 | 0.00 | 0.00 | 0.00 | 0.13 | 0.00 |
| time (sec) | N/A | 0.538 | 0.030 | 0.480 | 0.035 | 0.000 | 0.000 | 0.000 | 0.181 | 0.000 |

| Problem 60 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|----------|--------|-------|----------|----------|--------------|----------|----------|--------------|
| grade | N/A | F | A | A | F | F | F(-1) | F | F | F(-1) |
| verified | N/A | N/A | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 618 | 0 | 576 | 675 | 0 | 0 | 0 | 0 | 44 | 0 |
| N.S. | 1 | 0.00 | 0.93 | 1.09 | 0.00 | 0.00 | 0.00 | 0.00 | 0.07 | 0.00 |
| time (sec) | N/A | 0.000 | 12.028 | 0.175 | 0.000 | 0.000 | 0.000 | 0.000 | 0.199 | 0.000 |

| Problem 61 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|----------|----------|--------------|----------|----------|--------------|
| grade | N/A | A | A | A | F | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 585 | 585 | 549 | 648 | 0 | 0 | 0 | 0 | 17 | 0 |
| N.S. | 1 | 1.00 | 0.94 | 1.11 | 0.00 | 0.00 | 0.00 | 0.00 | 0.03 | 0.00 |
| time (sec) | N/A | 1.187 | 0.343 | 0.092 | 0.000 | 0.000 | 0.000 | 0.000 | 0.176 | 0.000 |

| Problem 62 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|----------|----------|--------------|----------|----------|--------------|
| grade | N/A | A | A | A | F | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 661 | 661 | 668 | 751 | 0 | 0 | 0 | 0 | 19 | 0 |
| N.S. | 1 | 1.00 | 1.01 | 1.14 | 0.00 | 0.00 | 0.00 | 0.00 | 0.03 | 0.00 |
| time (sec) | N/A | 1.385 | 0.306 | 0.096 | 0.000 | 0.000 | 0.000 | 0.000 | 0.183 | 0.000 |

| Problem 63 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|--------|-------|----------|----------|--------------|----------|----------|--------------|
| grade | N/A | A | A | A | F | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 381 | 398 | 694 | 615 | 0 | 0 | 0 | 0 | 725 | 0 |
| N.S. | 1 | 1.04 | 1.82 | 1.61 | 0.00 | 0.00 | 0.00 | 0.00 | 1.90 | 0.00 |
| time (sec) | N/A | 1.263 | 25.802 | 0.438 | 0.000 | 0.000 | 0.000 | 0.000 | 0.229 | 0.000 |

| Problem 64 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|--------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | C | B | F(-2) | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | No | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 323 | 524 | 1198 | 639 | 0 | 0 | 0 | 0 | 75 | 0 |
| N.S. | 1 | 1.62 | 3.71 | 1.98 | 0.00 | 0.00 | 0.00 | 0.00 | 0.23 | 0.00 |
| time (sec) | N/A | 1.164 | 11.079 | 1.987 | 0.000 | 0.000 | 0.000 | 0.000 | 0.188 | 0.000 |

| Problem 65 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | F | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 292 | 300 | 410 | 474 | 0 | 0 | 0 | 0 | 59 | 0 |
| N.S. | 1 | 1.03 | 1.40 | 1.62 | 0.00 | 0.00 | 0.00 | 0.00 | 0.20 | 0.00 |
| time (sec) | N/A | 0.874 | 0.145 | 0.275 | 0.000 | 0.000 | 0.000 | 0.000 | 0.174 | 0.000 |

| Problem 66 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | B | F(-2) | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 273 | 474 | 365 | 467 | 0 | 0 | 0 | 0 | 69 | 0 |
| N.S. | 1 | 1.74 | 1.34 | 1.71 | 0.00 | 0.00 | 0.00 | 0.00 | 0.25 | 0.00 |
| time (sec) | N/A | 0.831 | 0.162 | 0.615 | 0.000 | 0.000 | 0.000 | 0.000 | 0.195 | 0.000 |

| Problem 67 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | B | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 310 | 325 | 594 | 567 | 0 | 0 | 0 | 0 | 65 | 0 |
| N.S. | 1 | 1.05 | 1.92 | 1.83 | 0.00 | 0.00 | 0.00 | 0.00 | 0.21 | 0.00 |
| time (sec) | N/A | 1.175 | 6.534 | 0.355 | 0.000 | 0.000 | 0.000 | 0.000 | 0.214 | 0.000 |

| Problem 68 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-----------|-------|--------------|----------|--------------|----------|----------|--------------|
| grade | N/A | A | C | B | F(-2) | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | No | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 355 | 562 | 1221 | 678 | 0 | 0 | 0 | 0 | 81 | 0 |
| N.S. | 1 | 1.58 | 3.44 | 1.91 | 0.00 | 0.00 | 0.00 | 0.00 | 0.23 | 0.00 |
| time (sec) | N/A | 1.214 | 7.804 | 2.176 | 0.000 | 0.000 | 0.000 | 0.000 | 0.185 | 0.000 |

| Problem 69 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|--------|-------|----------|----------|--------------|----------|----------|--------------|
| grade | N/A | A | A | A | F | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 428 | 453 | 608 | 730 | 0 | 0 | 0 | 0 | 91 | 0 |
| N.S. | 1 | 1.06 | 1.42 | 1.71 | 0.00 | 0.00 | 0.00 | 0.00 | 0.21 | 0.00 |
| time (sec) | N/A | 1.413 | 33.031 | 0.468 | 0.000 | 0.000 | 0.000 | 0.000 | 0.189 | 0.000 |

| Problem 70 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-----------|----------|-----------|--------------|----------|--------------|----------|----------|--------------|
| grade | N/A | A | F | C | F(-2) | F | F(-1) | F | F | F(-1) |
| verified | N/A | No | N/A | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 727 | 956 | 0 | 505 | 0 | 0 | 0 | 0 | 109 | 0 |
| N.S. | 1 | 1.31 | 0.00 | 0.69 | 0.00 | 0.00 | 0.00 | 0.00 | 0.15 | 0.00 |
| time (sec) | N/A | 2.170 | 0.000 | 0.987 | 0.000 | 0.000 | 0.000 | 0.000 | 0.215 | 0.000 |

| Problem 71 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-----------|-----------|----------|----------|--------------|----------|----------|--------------|
| grade | N/A | A | A | C | F | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | No | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 463 | 467 | 679 | 327 | 0 | 0 | 0 | 0 | 66 | 0 |
| N.S. | 1 | 1.01 | 1.47 | 0.71 | 0.00 | 0.00 | 0.00 | 0.00 | 0.14 | 0.00 |
| time (sec) | N/A | 1.248 | 0.380 | 0.329 | 0.000 | 0.000 | 0.000 | 0.000 | 0.208 | 0.000 |

| Problem 72 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | C | C | F(-2) | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | No | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 679 | 778 | 652 | 338 | 0 | 0 | 0 | 0 | 97 | 0 |
| N.S. | 1 | 1.15 | 0.96 | 0.50 | 0.00 | 0.00 | 0.00 | 0.00 | 0.14 | 0.00 |
| time (sec) | N/A | 1.508 | 0.736 | 0.721 | 0.000 | 0.000 | 0.000 | 0.000 | 0.205 | 0.000 |

| Problem 73 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | C | F(-2) | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | No | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 679 | 919 | 714 | 366 | 0 | 0 | 0 | 0 | 104 | 0 |
| N.S. | 1 | 1.35 | 1.05 | 0.54 | 0.00 | 0.00 | 0.00 | 0.00 | 0.15 | 0.00 |
| time (sec) | N/A | 1.553 | 0.541 | 0.315 | 0.000 | 0.000 | 0.000 | 0.000 | 82.602 | 0.000 |

| Problem 74 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|--------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | C | F | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | No | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 480 | 491 | 915 | 420 | 0 | 0 | 0 | 0 | 72 | 0 |
| N.S. | 1 | 1.02 | 1.91 | 0.88 | 0.00 | 0.00 | 0.00 | 0.00 | 0.15 | 0.00 |
| time (sec) | N/A | 1.527 | 16.802 | 0.310 | 0.000 | 0.000 | 0.000 | 0.000 | 54.092 | 0.000 |

| Problem 75 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | C | C | F(-2) | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | No | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 761 | 865 | 2550 | 513 | 0 | 0 | 0 | 0 | 120 | 0 |
| N.S. | 1 | 1.14 | 3.35 | 0.67 | 0.00 | 0.00 | 0.00 | 0.00 | 0.16 | 0.00 |
| time (sec) | N/A | 1.920 | 8.740 | 0.463 | 0.000 | 0.000 | 0.000 | 0.000 | 0.272 | 0.000 |

| Problem 76 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | C | C | F(-2) | F | F(-1) | F | F | F(-1) |
| verified | N/A | No | No | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 791 | 1028 | 1564 | 662 | 0 | 0 | 0 | 0 | 0 | 0 |
| N.S. | 1 | 1.30 | 1.98 | 0.84 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| time (sec) | N/A | 2.168 | 7.370 | 0.533 | 0.000 | 0.000 | 0.000 | 0.000 | 83.256 | 0.000 |

| Problem 77 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | B | F(-2) | F | F(-1) | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 366 | 395 | 437 | 842 | 0 | 0 | 0 | 0 | 1076 | 0 |
| N.S. | 1 | 1.08 | 1.19 | 2.30 | 0.00 | 0.00 | 0.00 | 0.00 | 2.94 | 0.00 |
| time (sec) | N/A | 1.116 | 0.422 | 1.046 | 0.000 | 0.000 | 0.000 | 0.000 | 0.195 | 0.000 |

| Problem 78 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | C | F | F | F(-1) | F | F | F(-1) |
| verified | N/A | No | No | No | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 1135 | 2120 | 2111 | 718 | 0 | 0 | 0 | 0 | 419 | 0 |
| N.S. | 1 | 1.87 | 1.86 | 0.63 | 0.00 | 0.00 | 0.00 | 0.00 | 0.37 | 0.00 |
| time (sec) | N/A | 4.667 | 3.211 | 1.545 | 0.000 | 0.000 | 0.000 | 0.000 | 0.280 | 0.000 |

| Problem 79 | Optimal | Rubi | MMA | Maple | Maxima | Fricas | Sympy | Giac | Reduce | Mupad |
|------------|---------|-------|-------|-------|--------|--------|-------|-------|--------|-------|
| grade | N/A | A | A | A | F | F | F | F | F | F(-1) |
| verified | N/A | Yes | Yes | Yes | TBD | TBD | TBD | TBD | TBD | TBD |
| size | 83 | 72 | 150 | 149 | 0 | 0 | 0 | 0 | 73 | 0 |
| N.S. | 1 | 0.87 | 1.81 | 1.80 | 0.00 | 0.00 | 0.00 | 0.00 | 0.88 | 0.00 |
| time (sec) | N/A | 0.600 | 0.065 | 0.314 | 0.000 | 0.000 | 0.000 | 0.000 | 0.169 | 0.000 |

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [4] had the largest ratio of [.750000000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

| # | grade | number of steps used | number of unique rules | normalized antiderivative leaf size | integrand leaf size | $\frac{\text{number of rules}}{\text{integrand leaf size}}$ |
|----|-------|----------------------|------------------------|-------------------------------------|---------------------|---|
| 1 | A | 6 | 5 | 0.93 | 12 | 0.417 |
| 2 | A | 5 | 4 | 0.95 | 12 | 0.333 |
| 3 | A | 6 | 5 | 0.93 | 10 | 0.500 |
| 4 | A | 7 | 6 | 0.94 | 8 | 0.750 |
| 5 | A | 5 | 4 | 1.00 | 12 | 0.333 |
| 6 | A | 8 | 7 | 1.02 | 12 | 0.583 |
| 7 | A | 7 | 6 | 0.94 | 12 | 0.500 |
| 8 | A | 7 | 6 | 1.09 | 12 | 0.500 |
| 9 | A | 6 | 5 | 0.82 | 21 | 0.238 |
| 10 | A | 7 | 6 | 0.78 | 21 | 0.286 |
| 11 | A | 6 | 5 | 0.96 | 19 | 0.263 |
| 12 | A | 4 | 3 | 0.80 | 21 | 0.143 |
| 13 | A | 8 | 7 | 0.84 | 21 | 0.333 |
| 14 | A | 6 | 5 | 0.79 | 21 | 0.238 |
| 15 | A | 12 | 11 | 0.89 | 23 | 0.478 |
| 16 | A | 12 | 11 | 0.80 | 23 | 0.478 |
| 17 | A | 7 | 6 | 0.96 | 21 | 0.286 |
| 18 | A | 7 | 6 | 0.94 | 23 | 0.261 |
| 19 | A | 7 | 6 | 0.91 | 23 | 0.261 |
| 20 | A | 11 | 10 | 0.82 | 23 | 0.435 |
| 21 | A | 11 | 10 | 0.78 | 23 | 0.435 |

Continued on next page

Table 2.1 – continued from previous page

| # | grade | number of steps used | number of unique rules | normalized antiderivative leaf size | integrand leaf size | $\frac{\text{number of rules}}{\text{integrand leaf size}}$ |
|----|-------|----------------------|------------------------|-------------------------------------|---------------------|---|
| 22 | A | 16 | 15 | 0.91 | 23 | 0.652 |
| 23 | A | 13 | 12 | 0.84 | 23 | 0.522 |
| 24 | A | 11 | 10 | 0.89 | 21 | 0.476 |
| 25 | A | 8 | 7 | 0.90 | 23 | 0.304 |
| 26 | A | 8 | 7 | 0.90 | 23 | 0.304 |
| 27 | A | 10 | 9 | 0.86 | 23 | 0.391 |
| 28 | A | 17 | 16 | 0.82 | 23 | 0.696 |
| 29 | A | 4 | 3 | 1.19 | 12 | 0.250 |
| 30 | A | 4 | 3 | 0.94 | 19 | 0.158 |
| 31 | A | 6 | 5 | 0.99 | 18 | 0.278 |
| 32 | A | 6 | 5 | 1.07 | 18 | 0.278 |
| 33 | A | 6 | 5 | 1.09 | 16 | 0.312 |
| 34 | A | 1 | 1 | 1.00 | 10 | 0.100 |
| 35 | A | 7 | 6 | 1.14 | 18 | 0.333 |
| 36 | A | 4 | 4 | 1.06 | 18 | 0.222 |
| 37 | A | 4 | 4 | 1.02 | 18 | 0.222 |
| 38 | A | 5 | 4 | 0.97 | 20 | 0.200 |
| 39 | A | 5 | 4 | 1.00 | 20 | 0.200 |
| 40 | A | 5 | 4 | 1.05 | 18 | 0.222 |
| 41 | A | 7 | 6 | 0.96 | 12 | 0.500 |
| 42 | A | 4 | 3 | 1.13 | 20 | 0.150 |
| 43 | A | 7 | 6 | 1.22 | 20 | 0.300 |
| 44 | A | 7 | 6 | 1.19 | 20 | 0.300 |
| 45 | A | 5 | 4 | 0.98 | 20 | 0.200 |
| 46 | A | 5 | 4 | 0.99 | 18 | 0.222 |
| 47 | A | 7 | 6 | 0.98 | 12 | 0.500 |
| 48 | A | 4 | 3 | 1.12 | 20 | 0.150 |
| 49 | A | 7 | 6 | 1.71 | 20 | 0.300 |
| 50 | N/A | 3 | 0 | 1.00 | 20 | 0.000 |
| 51 | N/A | 3 | 0 | 1.00 | 20 | 0.000 |
| 52 | A | 5 | 4 | 1.37 | 18 | 0.222 |
| 53 | A | 3 | 3 | 1.23 | 16 | 0.188 |

Continued on next page

Table 2.1 – continued from previous page

| # | grade | number of steps used | number of unique rules | normalized antiderivative leaf size | integrand leaf size | $\frac{\text{number of rules}}{\text{integrand leaf size}}$ |
|----|-------|----------------------|------------------------|-------------------------------------|---------------------|---|
| 54 | A | 3 | 3 | 1.71 | 16 | 0.188 |
| 55 | A | 7 | 6 | 1.15 | 14 | 0.429 |
| 56 | A | 3 | 3 | 1.03 | 16 | 0.188 |
| 57 | A | 3 | 3 | 1.03 | 16 | 0.188 |
| 58 | A | 3 | 3 | 1.02 | 16 | 0.188 |
| 59 | A | 3 | 3 | 1.58 | 16 | 0.188 |
| 60 | F | 0 | 0 | N/A | 0.000 | N/A |
| 61 | A | 5 | 4 | 1.00 | 18 | 0.222 |
| 62 | A | 6 | 5 | 1.00 | 18 | 0.278 |
| 63 | A | 2 | 2 | 1.04 | 24 | 0.083 |
| 64 | A | 2 | 2 | 1.62 | 24 | 0.083 |
| 65 | A | 2 | 2 | 1.03 | 22 | 0.091 |
| 66 | A | 2 | 2 | 1.74 | 21 | 0.095 |
| 67 | A | 2 | 2 | 1.05 | 24 | 0.083 |
| 68 | A | 2 | 2 | 1.58 | 24 | 0.083 |
| 69 | A | 2 | 2 | 1.06 | 24 | 0.083 |
| 70 | A | 2 | 2 | 1.31 | 23 | 0.087 |
| 71 | A | 2 | 2 | 1.01 | 23 | 0.087 |
| 72 | A | 2 | 2 | 1.15 | 21 | 0.095 |
| 73 | A | 2 | 2 | 1.35 | 20 | 0.100 |
| 74 | A | 2 | 2 | 1.02 | 23 | 0.087 |
| 75 | A | 2 | 2 | 1.14 | 23 | 0.087 |
| 76 | A | 2 | 2 | 1.30 | 23 | 0.087 |
| 77 | A | 2 | 2 | 1.08 | 23 | 0.087 |
| 78 | A | 2 | 2 | 1.87 | 25 | 0.080 |
| 79 | A | 7 | 6 | 0.87 | 32 | 0.188 |

CHAPTER 3

LISTING OF INTEGRALS

| | | |
|------|---|-----|
| 3.1 | $\int x^3 \operatorname{arctanh}(a + bx)^2 dx$ | 56 |
| 3.2 | $\int x^2 \operatorname{arctanh}(a + bx)^2 dx$ | 63 |
| 3.3 | $\int x \operatorname{arctanh}(a + bx)^2 dx$ | 70 |
| 3.4 | $\int \operatorname{arctanh}(a + bx)^2 dx$ | 77 |
| 3.5 | $\int \frac{\operatorname{arctanh}(a+bx)^2}{x} dx$ | 83 |
| 3.6 | $\int \frac{\operatorname{arctanh}(a+bx)^2}{x^2} dx$ | 90 |
| 3.7 | $\int \frac{\operatorname{arctanh}(a+bx)^2}{x^3} dx$ | 98 |
| 3.8 | $\int \frac{\operatorname{arctanh}(1+bx)^2}{x} dx$ | 106 |
| 3.9 | $\int (ce + dex)^3 (a + b \operatorname{arctanh}(c + dx)) dx$ | 112 |
| 3.10 | $\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx)) dx$ | 121 |
| 3.11 | $\int (ce + dex) (a + b \operatorname{arctanh}(c + dx)) dx$ | 129 |
| 3.12 | $\int \frac{a+b \operatorname{arctanh}(c+dx)}{ce+dex} dx$ | 136 |
| 3.13 | $\int \frac{a+b \operatorname{arctanh}(c+dx)}{(ce+dex)^2} dx$ | 141 |
| 3.14 | $\int \frac{a+b \operatorname{arctanh}(c+dx)}{(ce+dex)^3} dx$ | 148 |
| 3.15 | $\int (ce + dex)^3 (a + b \operatorname{arctanh}(c + dx))^2 dx$ | 155 |
| 3.16 | $\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx$ | 166 |
| 3.17 | $\int (ce + dex) (a + b \operatorname{arctanh}(c + dx))^2 dx$ | 175 |
| 3.18 | $\int \frac{(a+b \operatorname{arctanh}(c+dx))^2}{ce+dex} dx$ | 184 |
| 3.19 | $\int \frac{(a+b \operatorname{arctanh}(c+dx))^2}{(ce+dex)^2} dx$ | 192 |
| 3.20 | $\int \frac{(a+b \operatorname{arctanh}(c+dx))^2}{(ce+dex)^3} dx$ | 199 |
| 3.21 | $\int \frac{(a+b \operatorname{arctanh}(c+dx))^2}{(ce+dex)^4} dx$ | 210 |
| 3.22 | $\int \frac{(a+b \operatorname{arctanh}(c+dx))^2}{(ce+dex)^5} dx$ | 219 |
| 3.23 | $\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx$ | 231 |
| 3.24 | $\int (ce + dex) (a + b \operatorname{arctanh}(c + dx))^3 dx$ | 242 |

| | | |
|------|--|-----|
| 3.25 | $\int \frac{(a+b\operatorname{arctanh}(c+dx))^3}{ce+dex} dx$ | 251 |
| 3.26 | $\int \frac{(a+b\operatorname{arctanh}(c+dx))^3}{(ce+dex)^2} dx$ | 259 |
| 3.27 | $\int \frac{(a+b\operatorname{arctanh}(c+dx))^3}{(ce+dex)^3} dx$ | 267 |
| 3.28 | $\int \frac{(a+b\operatorname{arctanh}(c+dx))^3}{(ce+dex)^4} dx$ | 276 |
| 3.29 | $\int \frac{\operatorname{arctanh}(1+x)}{2+2x} dx$ | 287 |
| 3.30 | $\int \frac{\operatorname{arctanh}(a+bx)}{\frac{ad}{b}+dx} dx$ | 292 |
| 3.31 | $\int (e+fx)^3(a+b\operatorname{arctanh}(c+dx)) dx$ | 297 |
| 3.32 | $\int (e+fx)^2(a+b\operatorname{arctanh}(c+dx)) dx$ | 308 |
| 3.33 | $\int (e+fx)(a+b\operatorname{arctanh}(c+dx)) dx$ | 317 |
| 3.34 | $\int (a+b\operatorname{arctanh}(c+dx)) dx$ | 324 |
| 3.35 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{e+fx} dx$ | 330 |
| 3.36 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{(e+fx)^2} dx$ | 337 |
| 3.37 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{(e+fx)^3} dx$ | 345 |
| 3.38 | $\int (e+fx)^3(a+b\operatorname{arctanh}(c+dx))^2 dx$ | 355 |
| 3.39 | $\int (e+fx)^2(a+b\operatorname{arctanh}(c+dx))^2 dx$ | 365 |
| 3.40 | $\int (e+fx)(a+b\operatorname{arctanh}(c+dx))^2 dx$ | 375 |
| 3.41 | $\int (a+b\operatorname{arctanh}(c+dx))^2 dx$ | 383 |
| 3.42 | $\int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{e+fx} dx$ | 390 |
| 3.43 | $\int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{(e+fx)^2} dx$ | 397 |
| 3.44 | $\int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{(e+fx)^3} dx$ | 406 |
| 3.45 | $\int (e+fx)^2(a+b\operatorname{arctanh}(c+dx))^3 dx$ | 417 |
| 3.46 | $\int (e+fx)(a+b\operatorname{arctanh}(c+dx))^3 dx$ | 427 |
| 3.47 | $\int (a+b\operatorname{arctanh}(c+dx))^3 dx$ | 435 |
| 3.48 | $\int \frac{(a+b\operatorname{arctanh}(c+dx))^3}{e+fx} dx$ | 442 |
| 3.49 | $\int \frac{(a+b\operatorname{arctanh}(c+dx))^3}{(e+fx)^2} dx$ | 449 |
| 3.50 | $\int (e+fx)^m(a+b\operatorname{arctanh}(c+dx))^3 dx$ | 460 |
| 3.51 | $\int (e+fx)^m(a+b\operatorname{arctanh}(c+dx))^2 dx$ | 465 |
| 3.52 | $\int (e+fx)^m(a+b\operatorname{arctanh}(c+dx)) dx$ | 470 |
| 3.53 | $\int \frac{\operatorname{arctanh}(a+bx)}{c+dx^3} dx$ | 476 |
| 3.54 | $\int \frac{\operatorname{arctanh}(a+bx)}{c+dx^2} dx$ | 486 |
| 3.55 | $\int \frac{\operatorname{arctanh}(a+bx)}{c+dx} dx$ | 493 |
| 3.56 | $\int \frac{\operatorname{arctanh}(a+bx)}{c+\frac{d}{x}} dx$ | 500 |
| 3.57 | $\int \frac{\operatorname{arctanh}(a+bx)}{c+\frac{d}{x^2}} dx$ | 507 |

| | | |
|------|--|-----|
| 3.58 | $\int \frac{\operatorname{arctanh}(a+bx)}{c+\frac{d}{x^3}} dx$ | 515 |
| 3.59 | $\int \frac{\operatorname{arctanh}(a+bx)}{1-x^2} dx$ | 524 |
| 3.60 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{e+f\sqrt{x}} dx$ | 530 |
| 3.61 | $\int \frac{\operatorname{arctanh}(a+bx)}{c+d\sqrt{x}} dx$ | 538 |
| 3.62 | $\int \frac{\operatorname{arctanh}(a+bx)}{c+\frac{d}{\sqrt{x}}} dx$ | 546 |
| 3.63 | $\int \frac{x^3(a+b\operatorname{arctanh}(c+dx))}{e-fx^2} dx$ | 557 |
| 3.64 | $\int \frac{x^2(a+b\operatorname{arctanh}(c+dx))}{e-fx^2} dx$ | 565 |
| 3.65 | $\int \frac{x(a+b\operatorname{arctanh}(c+dx))}{e-fx^2} dx$ | 572 |
| 3.66 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{e-fx^2} dx$ | 579 |
| 3.67 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{x(e-fx^2)} dx$ | 585 |
| 3.68 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{x^2(e-fx^2)} dx$ | 592 |
| 3.69 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{x^3(e-fx^2)} dx$ | 599 |
| 3.70 | $\int \frac{x^3(a+b\operatorname{arctanh}(c+dx))}{e+fx^3} dx$ | 607 |
| 3.71 | $\int \frac{x^2(a+b\operatorname{arctanh}(c+dx))}{e+fx^3} dx$ | 617 |
| 3.72 | $\int \frac{x(a+b\operatorname{arctanh}(c+dx))}{e+fx^3} dx$ | 627 |
| 3.73 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{e+fx^3} dx$ | 636 |
| 3.74 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{x(e+fx^3)} dx$ | 646 |
| 3.75 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{x^2(e+fx^3)} dx$ | 655 |
| 3.76 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{x^3(e+fx^3)} dx$ | 664 |
| 3.77 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{e+fx+gx^2} dx$ | 674 |
| 3.78 | $\int \frac{a+b\operatorname{arctanh}(c+dx)}{e+fx^2+gx^4} dx$ | 682 |
| 3.79 | $\int \frac{(ce+dex)(a+b\operatorname{arctanh}(c+dx))}{1-(c+dx)^2} dx$ | 691 |

3.1 $\int x^3 \operatorname{arctanh}(a + bx)^2 dx$

| | |
|---|----|
| Optimal result | 56 |
| Mathematica [A] (verified) | 57 |
| Rubi [A] (verified) | 57 |
| Maple [A] (verified) | 59 |
| Fricas [F] | 60 |
| Sympy [F] | 60 |
| Maxima [A] (verification not implemented) | 61 |
| Giac [F] | 61 |
| Mupad [F(-1)] | 62 |
| Reduce [F] | 62 |

Optimal result

Integrand size = 12, antiderivative size = 263

$$\begin{aligned}
 \int x^3 \operatorname{arctanh}(a + bx)^2 dx = & -\frac{ax}{b^3} + \frac{(a + bx)^2}{12b^4} + \frac{a \operatorname{arctanh}(a + bx)}{b^4} \\
 & + \frac{(1 + 6a^2)(a + bx) \operatorname{arctanh}(a + bx)}{2b^4} \\
 & - \frac{a(a + bx)^2 \operatorname{arctanh}(a + bx)}{b^4} \\
 & + \frac{(a + bx)^3 \operatorname{arctanh}(a + bx)}{6b^4} - \frac{a(1 + a^2) \operatorname{arctanh}(a + bx)^2}{b^4} \\
 & - \frac{(1 + 6a^2 + a^4) \operatorname{arctanh}(a + bx)^2}{4b^4} + \frac{1}{4} x^4 \operatorname{arctanh}(a + bx)^2 \\
 & + \frac{2a(1 + a^2) \operatorname{arctanh}(a + bx) \log\left(\frac{2}{1 - a - bx}\right)}{b^4} \\
 & + \frac{\log(1 - (a + bx)^2)}{12b^4} + \frac{(1 + 6a^2) \log(1 - (a + bx)^2)}{4b^4} \\
 & + \frac{a(1 + a^2) \operatorname{PolyLog}\left(2, -\frac{1 + a + bx}{1 - a - bx}\right)}{b^4}
 \end{aligned}$$

output

```
-a*x/b^3+1/12*(b*x+a)^2/b^4+a*arctanh(b*x+a)/b^4+1/2*(6*a^2+1)*(b*x+a)*arc
tanh(b*x+a)/b^4-a*(b*x+a)^2*arctanh(b*x+a)/b^4+1/6*(b*x+a)^3*arctanh(b*x+a
)/b^4-a*(a^2+1)*arctanh(b*x+a)^2/b^4-1/4*(a^4+6*a^2+1)*arctanh(b*x+a)^2/b^
4+1/4*x^4*arctanh(b*x+a)^2+2*a*(a^2+1)*arctanh(b*x+a)*ln(2/(-b*x-a+1))/b^4
+1/12*ln(1-(b*x+a)^2)/b^4+1/4*(6*a^2+1)*ln(1-(b*x+a)^2)/b^4+a*(a^2+1)*poly
log(2,-(b*x+a+1)/(-b*x-a+1))/b^4
```

Mathematica [A] (verified)

Time = 0.97 (sec) , antiderivative size = 187, normalized size of antiderivative = 0.71

$$\int x^3 \operatorname{arctanh}(a + bx)^2 dx =$$

$$\frac{1 + 11a^2 + 10abx - b^2x^2 + 3(1 - 4a + 6a^2 - 4a^3 + a^4 - b^4x^4) \operatorname{arctanh}(a + bx)^2 - 2\operatorname{arctanh}(a + bx)}{b^4}$$

input

```
Integrate[x^3*ArcTanh[a + b*x]^2,x]
```

output

```
-1/12*(1 + 11*a^2 + 10*a*b*x - b^2*x^2 + 3*(1 - 4*a + 6*a^2 - 4*a^3 + a^4
- b^4*x^4)*ArcTanh[a + b*x]^2 - 2*ArcTanh[a + b*x]*(9*a + 13*a^3 + 3*b*x +
9*a^2*b*x - 3*a*b^2*x^2 + b^3*x^3 + 12*(a + a^3)*Log[1 + E^(-2*ArcTanh[a
+ b*x]])) + 8*Log[1/Sqrt[1 - (a + b*x)^2]] + 36*a^2*Log[1/Sqrt[1 - (a + b*
x)^2]] + 12*(a + a^3)*PolyLog[2, -E^(-2*ArcTanh[a + b*x]]))/b^4
```

Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 244, normalized size of antiderivative = 0.93, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {6661, 25, 27, 6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \operatorname{arctanh}(a + bx)^2 dx$$

$$\begin{array}{c} \downarrow 6661 \\ \frac{\int x^3 \operatorname{arctanh}(a + bx)^2 d(a + bx)}{b} \\ \downarrow 25 \\ \frac{\int -x^3 \operatorname{arctanh}(a + bx)^2 d(a + bx)}{b} \\ \downarrow 27 \\ \frac{\int -b^3 x^3 \operatorname{arctanh}(a + bx)^2 d(a + bx)}{b^4} \\ \downarrow 6480 \end{array}$$

$$\frac{1}{2} \int \left(-\operatorname{arctanh}(a + bx)(a + bx)^2 + 4a \operatorname{arctanh}(a + bx)(a + bx) - (6a^2 + 1) \operatorname{arctanh}(a + bx) + \frac{(a^4 + 6a^2 - 4)(a^2 + 1)}{1} \right) dx$$

$$b^4$$

$$\downarrow 2009$$

$$\frac{1}{2} \left(-(6a^2 + 1)(a + bx) \operatorname{arctanh}(a + bx) + 2a(a^2 + 1) \operatorname{arctanh}(a + bx)^2 - 4a(a^2 + 1) \operatorname{arctanh}(a + bx) \log \left(\frac{a + bx}{-a - bx} \right) \right)$$

input `Int[x^3*ArcTanh[a + b*x]^2,x]`

output

```

-((-1/4*(b^4*x^4*ArcTanh[a + b*x]^2) + (2*a*(a + b*x) - (a + b*x)^2/6 - 2*
a*ArcTanh[a + b*x] - (1 + 6*a^2)*(a + b*x)*ArcTanh[a + b*x] + 2*a*(a + b*x)
)^2*ArcTanh[a + b*x] - ((a + b*x)^3*ArcTanh[a + b*x])/3 + 2*a*(1 + a^2)*Ar
cTanh[a + b*x]^2 + ((1 + 6*a^2 + a^4)*ArcTanh[a + b*x]^2)/2 - 4*a*(1 + a^2
)*ArcTanh[a + b*x]*Log[2/(1 - a - b*x)] - Log[1 - (a + b*x)^2]/6 - ((1 + 6
*a^2)*Log[1 - (a + b*x)^2])/2 - 2*a*(1 + a^2)*PolyLog[2, -((1 + a + b*x)/(
1 - a - b*x))])/2)/b^4

```

Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6480 `Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^ (p_) * ((d_) + (e_.)*(x_.))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1) * ((a + b*ArcTanh[c*x])^p / (e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]`
- rule 6661 `Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_.)]*(b_.))^ (p_.) * ((e_.) + (f_.)*(x_.))^(m_.), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 1.29 (sec) , antiderivative size = 449, normalized size of antiderivative = 1.71

| method | result |
|-------------------|---|
| parts | $\frac{x^4 \operatorname{arctanh}(bx+a)^2}{4} - \frac{-6 \operatorname{arctanh}(bx+a)a^2(bx+a)+2 \operatorname{arctanh}(bx+a)(bx+a)^2 a - \frac{\operatorname{arctanh}(bx+a)(bx+a)^3}{3} - \operatorname{arctanh}(bx+a)}{3}$ |
| derivativedivides | $\frac{-\operatorname{arctanh}(bx+a) \ln(bx+a-1)a + \frac{3 \operatorname{arctanh}(bx+a) \ln(bx+a-1)a^2}{2} + \frac{\operatorname{arctanh}(bx+a) \ln(bx+a-1)a^4}{4} - \operatorname{arctanh}(bx+a) \ln(bx+a)}{4}$ |
| default | $\frac{-\operatorname{arctanh}(bx+a) \ln(bx+a-1)a + \frac{3 \operatorname{arctanh}(bx+a) \ln(bx+a-1)a^2}{2} + \frac{\operatorname{arctanh}(bx+a) \ln(bx+a-1)a^4}{4} - \operatorname{arctanh}(bx+a) \ln(bx+a)}{4}$ |
| risch | $-\frac{1}{12b^4} - \frac{5ax}{6b^3} - \frac{\ln\left(-\frac{bx}{2} - \frac{a}{2} + \frac{1}{2}\right) \ln\left(\frac{bx}{2} + \frac{a}{2} + \frac{1}{2}\right) a^3}{b^4} + \frac{\ln\left(\frac{bx}{2} + \frac{a}{2} + \frac{1}{2}\right) \ln(-bx-a+1) a^3}{b^4} - \frac{\ln\left(-\frac{bx}{2} - \frac{a}{2} + \frac{1}{2}\right) \ln\left(\frac{bx}{2} + \frac{a}{2} + \frac{1}{2}\right)}{b^4}$ |

input `int(x^3*arctanh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `1/4*x^4*arctanh(b*x+a)^2-1/2/b^4*(-6*arctanh(b*x+a)*a^2*(b*x+a)+2*arctanh(b*x+a)*(b*x+a)^2*a-1/3*arctanh(b*x+a)*(b*x+a)^3-arctanh(b*x+a)*(b*x+a)-1/2*arctanh(b*x+a)*ln(b*x+a-1)*a^4+2*arctanh(b*x+a)*ln(b*x+a-1)*a^3-3*arctanh(b*x+a)*ln(b*x+a-1)*a^2+2*arctanh(b*x+a)*ln(b*x+a-1)*a-1/2*arctanh(b*x+a)*ln(b*x+a-1)+1/2*arctanh(b*x+a)*ln(b*x+a+1)*a^4+2*arctanh(b*x+a)*ln(b*x+a+1)*a^3+3*arctanh(b*x+a)*ln(b*x+a+1)*a^2+2*arctanh(b*x+a)*ln(b*x+a+1)*a+1/2*arctanh(b*x+a)*ln(b*x+a+1)+2*(b*x+a)*a-1/6*(b*x+a)^2-1/6*(18*a^2-6*a+4)*ln(b*x+a-1)+1/6*(-18*a^2-6*a-4)*ln(b*x+a+1)-1/6*(3*a^4-12*a^3+18*a^2-12*a+3)*(1/4*ln(b*x+a-1)^2-1/2*dilog(1/2*b*x+1/2*a+1/2)-1/2*ln(b*x+a-1)*ln(1/2*b*x+1/2*a+1/2))-1/6*(-3*a^4-12*a^3-18*a^2-12*a-3)*(-1/4*ln(b*x+a+1)^2+1/2*(ln(b*x+a+1)-ln(1/2*b*x+1/2*a+1/2))*ln(-1/2*b*x-1/2*a+1/2)-1/2*dilog(1/2*b*x+1/2*a+1/2)))`

Fricas [F]

$$\int x^3 \operatorname{arctanh}(a + bx)^2 dx = \int x^3 \operatorname{artanh}(bx + a)^2 dx$$

input `integrate(x^3*arctanh(b*x+a)^2,x, algorithm="fricas")`

output `integral(x^3*arctanh(b*x + a)^2, x)`

Sympy [F]

$$\int x^3 \operatorname{arctanh}(a + bx)^2 dx = \int x^3 \operatorname{atanh}^2(a + bx) dx$$

input `integrate(x**3*atanh(b*x+a)**2,x)`

output `Integral(x**3*atanh(a + b*x)**2, x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.22

$$\int x^3 \operatorname{arctanh}(a + bx)^2 dx = \frac{1}{4} x^4 \operatorname{artanh}(bx + a)^2 + \frac{1}{48} b^2 \left(\frac{48(a^3 + a) \left(\log(bx + a - 1) \log\left(\frac{1}{2}bx + \frac{1}{2}a + \frac{1}{2}\right) + \operatorname{Li}_2\left(-\frac{1}{2}bx - \frac{1}{2}a + \frac{1}{2}\right)\right)}{b^6} + \frac{4(13a^3 + 18a^2 + 9a + 4) \log(bx + a + 1)}{b^6} + \frac{4(b^2x^2 - 40abx + 3(a^4 + 4a^3 + 6a^2 + 4a + 1) \log(bx + a + 1)^2 - 6(a^4 + 4a^3 + 6a^2 + 4a + 1) \log(bx + a + 1) \log(bx + a - 1) + 3(a^4 - 4a^3 + 6a^2 - 4a + 1) \log(bx + a - 1)^2 - 4(13a^3 - 18a^2 + 9a - 4) \log(bx + a - 1))}{b^6} + \frac{1}{12} b \left(\frac{2(b^2x^3 - 3abx^2 + 3(3a^2 + 1)x)}{b^4} - \frac{3(a^4 + 4a^3 + 6a^2 + 4a + 1) \log(bx + a + 1)}{b^5} + \frac{3(a^4 - 4a^3 + 6a^2 - 4a + 1) \log(bx + a - 1)}{b^5} \right) \operatorname{arctanh}(bx + a) \right)$$

input `integrate(x^3*arctanh(b*x+a)^2,x, algorithm="maxima")`output `1/4*x^4*arctanh(b*x + a)^2 + 1/48*b^2*(48*(a^3 + a)*(log(b*x + a - 1)*log(1/2*b*x + 1/2*a + 1/2) + dilog(-1/2*b*x - 1/2*a + 1/2))/b^6 + 4*(13*a^3 + 18*a^2 + 9*a + 4)*log(b*x + a + 1)/b^6 + (4*b^2*x^2 - 40*a*b*x + 3*(a^4 + 4*a^3 + 6*a^2 + 4*a + 1)*log(b*x + a + 1)^2 - 6*(a^4 + 4*a^3 + 6*a^2 + 4*a + 1)*log(b*x + a + 1)*log(b*x + a - 1) + 3*(a^4 - 4*a^3 + 6*a^2 - 4*a + 1)*log(b*x + a - 1)^2 - 4*(13*a^3 - 18*a^2 + 9*a - 4)*log(b*x + a - 1))/b^6) + 1/12*b*(2*(b^2*x^3 - 3*a*b*x^2 + 3*(3*a^2 + 1)*x)/b^4 - 3*(a^4 + 4*a^3 + 6*a^2 + 4*a + 1)*log(b*x + a + 1)/b^5 + 3*(a^4 - 4*a^3 + 6*a^2 - 4*a + 1)*log(b*x + a - 1)/b^5)*arctanh(b*x + a)`**Giac [F]**

$$\int x^3 \operatorname{arctanh}(a + bx)^2 dx = \int x^3 \operatorname{artanh}(bx + a)^2 dx$$

input `integrate(x^3*arctanh(b*x+a)^2,x, algorithm="giac")`output `integrate(x^3*arctanh(b*x + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^3 \operatorname{arctanh}(a + bx)^2 dx = \int x^3 \operatorname{atanh}(a + bx)^2 dx$$

input `int(x^3*atanh(a + b*x)^2,x)`output `int(x^3*atanh(a + b*x)^2, x)`**Reduce [F]**

$$\int x^3 \operatorname{arctanh}(a + bx)^2 dx$$

$$= \frac{3 \operatorname{atanh}(bx + a)^2 a^4 - 6 \operatorname{atanh}(bx + a)^2 a^2 + 3 \operatorname{atanh}(bx + a)^2 b^4 x^4 + 3 \operatorname{atanh}(bx + a)^2 + 14 \operatorname{atanh}(bx + a)}$$

input `int(x^3*atanh(b*x+a)^2,x)`

output

```
(3*atanh(a + b*x)**2*a**4 - 6*atanh(a + b*x)**2*a**2 + 3*atanh(a + b*x)**2
*b**4*x**4 + 3*atanh(a + b*x)**2 + 14*atanh(a + b*x)*a**3 + 6*atanh(a + b*
x)*a**2*b*x + 24*atanh(a + b*x)*a**2 - 6*atanh(a + b*x)*a*b**2*x**2 + 6*at
anh(a + b*x)*a + 2*atanh(a + b*x)*b**3*x**3 - 6*atanh(a + b*x)*b*x - 4*ata
nh(a + b*x) + 12*int((atanh(a + b*x)*x**2)/(a**2 + 2*a*b*x + b**2*x**2 - 1
),x)*a**2*b**3 + 12*int((atanh(a + b*x)*x**2)/(a**2 + 2*a*b*x + b**2*x**2
- 1),x)*b**3 + 24*log(a + b*x - 1)*a**2 - 4*log(a + b*x - 1) - 10*a*b*x +
b**2*x**2)/(12*b**4)
```

3.2 $\int x^2 \operatorname{arctanh}(a + bx)^2 dx$

| | |
|---|----|
| Optimal result | 63 |
| Mathematica [B] (warning: unable to verify) | 64 |
| Rubi [A] (verified) | 64 |
| Maple [A] (verified) | 66 |
| Fricas [F] | 67 |
| Sympy [F] | 67 |
| Maxima [A] (verification not implemented) | 68 |
| Giac [F] | 68 |
| Mupad [F(-1)] | 69 |
| Reduce [F] | 69 |

Optimal result

Integrand size = 12, antiderivative size = 204

$$\begin{aligned} \int x^2 \operatorname{arctanh}(a + bx)^2 dx = & \frac{x}{3b^2} - \frac{\operatorname{arctanh}(a + bx)}{3b^3} - \frac{2a(a + bx)\operatorname{arctanh}(a + bx)}{b^3} \\ & + \frac{(a + bx)^2 \operatorname{arctanh}(a + bx)}{3b^3} + \frac{a(3 + a^2) \operatorname{arctanh}(a + bx)^2}{3b^3} \\ & + \frac{(1 + 3a^2) \operatorname{arctanh}(a + bx)^2}{3b^3} + \frac{1}{3} x^3 \operatorname{arctanh}(a + bx)^2 \\ & - \frac{2(1 + 3a^2) \operatorname{arctanh}(a + bx) \log\left(\frac{2}{1 - a - bx}\right)}{3b^3} \\ & - \frac{a \log(1 - (a + bx)^2)}{b^3} - \frac{(1 + 3a^2) \operatorname{PolyLog}\left(2, -\frac{1 + a + bx}{1 - a - bx}\right)}{3b^3} \end{aligned}$$

output

```
1/3*x/b^2-1/3*arctanh(b*x+a)/b^3-2*a*(b*x+a)*arctanh(b*x+a)/b^3+1/3*(b*x+a)^2*arctanh(b*x+a)/b^3+1/3*a*(a^2+3)*arctanh(b*x+a)^2/b^3+1/3*(3*a^2+1)*arctanh(b*x+a)^2/b^3+1/3*x^3*arctanh(b*x+a)^2-2/3*(3*a^2+1)*arctanh(b*x+a)*ln(2/(-b*x-a+1))/b^3-a*ln(1-(b*x+a)^2)/b^3-1/3*(3*a^2+1)*polylog(2,-(b*x+a+1)/(-b*x-a+1))/b^3
```


Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 463 vs. $2(204) = 408$.

Time = 1.46 (sec) , antiderivative size = 463, normalized size of antiderivative = 2.27

$$\int x^2 \operatorname{arctanh}(a + bx)^2 dx =$$

$$(1 - (a + bx)^2)^{3/2} \left(-\frac{a+bx}{\sqrt{1-(a+bx)^2}} + \frac{6a(a+bx)\operatorname{arctanh}(a+bx)}{\sqrt{1-(a+bx)^2}} + \frac{3(a+bx)\operatorname{arctanh}(a+bx)^2}{\sqrt{1-(a+bx)^2}} - \frac{3a^2(a+bx)\operatorname{arctanh}(a+bx)}{\sqrt{1-(a+bx)^2}} \right)$$

input `Integrate[x^2*ArcTanh[a + b*x]^2,x]`

output

```
-1/12*((1 - (a + b*x)^2)^(3/2)*(-(a + b*x)/Sqrt[1 - (a + b*x)^2]) + (6*a*(a + b*x)*ArcTanh[a + b*x])/Sqrt[1 - (a + b*x)^2] + (3*(a + b*x)*ArcTanh[a + b*x]^2)/Sqrt[1 - (a + b*x)^2] - (3*a^2*(a + b*x)*ArcTanh[a + b*x]^2)/Sqrt[1 - (a + b*x)^2] + ArcTanh[a + b*x]^2*Cosh[3*ArcTanh[a + b*x]] + 3*a^2*ArcTanh[a + b*x]^2*Cosh[3*ArcTanh[a + b*x]] + 2*ArcTanh[a + b*x]*Cosh[3*ArcTanh[a + b*x]]*Log[1 + E^(-2*ArcTanh[a + b*x])] + 6*a^2*ArcTanh[a + b*x]*Cosh[3*ArcTanh[a + b*x]]*Log[1 + E^(-2*ArcTanh[a + b*x])] - 6*a*Cosh[3*ArcTanh[a + b*x]]*Log[1/Sqrt[1 - (a + b*x)^2]] + (3*(1 - 4*a + 3*a^2)*ArcTanh[a + b*x]^2 + 2*ArcTanh[a + b*x]*(2 + (3 + 9*a^2)*Log[1 + E^(-2*ArcTanh[a + b*x])]) - 18*a*Log[1/Sqrt[1 - (a + b*x)^2]])/Sqrt[1 - (a + b*x)^2] - (4*(1 + 3*a^2)*PolyLog[2, -E^(-2*ArcTanh[a + b*x])])/(1 - (a + b*x)^2)^(3/2) - Sinh[3*ArcTanh[a + b*x]] + 6*a*ArcTanh[a + b*x]*Sinh[3*ArcTanh[a + b*x]] - ArcTanh[a + b*x]^2*Sinh[3*ArcTanh[a + b*x]] - 3*a^2*ArcTanh[a + b*x]^2*Sinh[3*ArcTanh[a + b*x]]))/b^3
```

Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 193, normalized size of antiderivative = 0.95, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6661, 27, 6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int x^2 \operatorname{arctanh}(a + bx)^2 dx \\
& \quad \downarrow \text{6661} \\
& \frac{\int x^2 \operatorname{arctanh}(a + bx)^2 d(a + bx)}{b} \\
& \quad \downarrow \text{27} \\
& \frac{\int b^2 x^2 \operatorname{arctanh}(a + bx)^2 d(a + bx)}{b^3} \\
& \quad \downarrow \text{6480} \\
& \frac{\frac{2}{3} \int \left(-3a \operatorname{arctanh}(a + bx) + (a + bx) \operatorname{arctanh}(a + bx) + \frac{(a(a^2+3) - (3a^2+1)(a+bx)) \operatorname{arctanh}(a+bx)}{1-(a+bx)^2} \right) d(a + bx) + \frac{1}{3} b^3 x}{b^3} \\
& \quad \downarrow \text{2009} \\
& \frac{\frac{2}{3} \left(\frac{1}{2} a (a^2 + 3) \operatorname{arctanh}(a + bx)^2 + \frac{1}{2} (3a^2 + 1) \operatorname{arctanh}(a + bx)^2 - (3a^2 + 1) \operatorname{arctanh}(a + bx) \log \left(\frac{2}{-a - bx + 1} \right) - \frac{1}{2} \right)}{b^3}
\end{aligned}$$

input `Int[x^2*ArcTanh[a + b*x]^2,x]`

output `((b^3*x^3*ArcTanh[a + b*x]^2)/3 + (2*((a + b*x)/2 - ArcTanh[a + b*x])/2 - 3*a*(a + b*x)*ArcTanh[a + b*x] + ((a + b*x)^2*ArcTanh[a + b*x])/2 + (a*(3 + a^2)*ArcTanh[a + b*x]^2)/2 + ((1 + 3*a^2)*ArcTanh[a + b*x]^2)/2 - (1 + 3*a^2)*ArcTanh[a + b*x]*Log[2/(1 - a - b*x)] - (3*a*Log[1 - (a + b*x)^2])/2 - ((1 + 3*a^2)*PolyLog[2, -((1 + a + b*x)/(1 - a - b*x))])/2))/3)/b^3`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6480

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)*((d_.) + (e_.)*(x_.))^(q_.), x_Symbol]
:> Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] -
Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1),
(d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x]
&& IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

rule 6661

```
Int[((a_.) + ArcTanh[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol]
:> Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 350, normalized size of antiderivative = 1.72

| method | result |
|-------------------|--|
| parts | $\frac{x^3 \operatorname{arctanh}(bx+a)^2}{3} - \frac{2 \left(3 \operatorname{arctanh}(bx+a)(bx+a)a - \frac{\operatorname{arctanh}(bx+a)(bx+a)^2}{2} + \frac{\operatorname{arctanh}(bx+a) \ln(bx+a-1)a^3}{2} - 3 \operatorname{arctanh}(bx+a) \right)}{3}$ |
| derivativedivides | $\frac{-\operatorname{arctanh}(bx+a)^2 a^3 + \operatorname{arctanh}(bx+a)^2 a^2 (bx+a) - \operatorname{arctanh}(bx+a)^2 a (bx+a)^2 + \frac{\operatorname{arctanh}(bx+a)^2 (bx+a)^3}{3} - 2 \operatorname{arctanh}(bx+a)}{3}$ |
| default | $\frac{-\operatorname{arctanh}(bx+a)^2 a^3 + \operatorname{arctanh}(bx+a)^2 a^2 (bx+a) - \operatorname{arctanh}(bx+a)^2 a (bx+a)^2 + \frac{\operatorname{arctanh}(bx+a)^2 (bx+a)^3}{3} - 2 \operatorname{arctanh}(bx+a)}{3}$ |
| risch | $-\frac{1}{3b^3} + \frac{x}{3b^2} + \frac{a}{3b^3} + \frac{x^3 \ln(-bx-a+1)^2}{12} - \frac{\ln\left(\frac{bx}{2} + \frac{a}{2} + \frac{1}{2}\right) \ln(-bx-a+1)}{3b^3} + \frac{\ln\left(\frac{bx}{2} + \frac{a}{2} + \frac{1}{2}\right) \ln\left(-\frac{bx}{2} - \frac{a}{2} + \frac{1}{2}\right)}{3b^3}$ |

input

```
int(x^2*arctanh(b*x+a)^2,x,method=_RETURNVERBOSE)
```

output

```
1/3*x^3*arctanh(b*x+a)^2-2/3/b^3*(3*arctanh(b*x+a)*(b*x+a)*a-1/2*arctanh(b
*x+a)*(b*x+a)^2+1/2*arctanh(b*x+a)*ln(b*x+a-1)*a^3-3/2*arctanh(b*x+a)*ln(b
*x+a-1)*a^2+3/2*arctanh(b*x+a)*ln(b*x+a-1)*a-1/2*arctanh(b*x+a)*ln(b*x+a-1
)-1/2*arctanh(b*x+a)*ln(b*x+a+1)*a^3-3/2*arctanh(b*x+a)*ln(b*x+a+1)*a^2-3/
2*arctanh(b*x+a)*ln(b*x+a+1)*a-1/2*arctanh(b*x+a)*ln(b*x+a+1)-1/2*(a^3+3*a
^2+3*a+1)*(-1/4*ln(b*x+a+1)^2+1/2*(ln(b*x+a+1)-ln(1/2*b*x+1/2*a+1/2))*ln(-
1/2*b*x-1/2*a+1/2)-1/2*dilog(1/2*b*x+1/2*a+1/2))-1/2*b*x-1/2*a+1/4*(6*a-1)
*ln(b*x+a-1)-1/4*(-6*a-1)*ln(b*x+a+1)-1/2*(-a^3+3*a^2-3*a+1)*(1/4*ln(b*x+a
-1)^2-1/2*dilog(1/2*b*x+1/2*a+1/2)-1/2*ln(b*x+a-1)*ln(1/2*b*x+1/2*a+1/2)))
```

Fricas [F]

$$\int x^2 \operatorname{arctanh}(a + bx)^2 dx = \int x^2 \operatorname{artanh}(bx + a)^2 dx$$

input

```
integrate(x^2*arctanh(b*x+a)^2,x, algorithm="fricas")
```

output

```
integral(x^2*arctanh(b*x + a)^2, x)
```

Sympy [F]

$$\int x^2 \operatorname{arctanh}(a + bx)^2 dx = \int x^2 \operatorname{atanh}^2(a + bx) dx$$

input

```
integrate(x**2*atanh(b*x+a)**2,x)
```

output

```
Integral(x**2*atanh(a + b*x)**2, x)
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 259, normalized size of antiderivative = 1.27

$$\int x^2 \operatorname{arctanh}(a + bx)^2 dx = \frac{1}{3} x^3 \operatorname{artanh}(bx + a)^2 - \frac{1}{12} b^2 \left(\frac{4(3a^2 + 1)(\log(bx + a - 1) \log(\frac{1}{2}bx + \frac{1}{2}a + \frac{1}{2})) + \operatorname{Li}_2(-\frac{1}{2}bx - \frac{1}{2}a + \frac{1}{2})}{b^5} + \frac{2(5a^2 + 6a + 1)}{b} + \frac{1}{3} b \left(\frac{bx^2 - 4ax}{b^3} + \frac{(a^3 + 3a^2 + 3a + 1) \log(bx + a + 1)}{b^4} - \frac{(a^3 - 3a^2 + 3a - 1) \log(bx + a - 1)}{b^4} \right) \operatorname{art} \right. \\ \left. + a \right)$$

input `integrate(x^2*arctanh(b*x+a)^2,x, algorithm="maxima")`output `1/3*x^3*arctanh(b*x + a)^2 - 1/12*b^2*(4*(3*a^2 + 1)*(log(b*x + a - 1)*log(1/2*b*x + 1/2*a + 1/2) + dilog(-1/2*b*x - 1/2*a + 1/2))/b^5 + 2*(5*a^2 + 6*a + 1)*log(b*x + a + 1)/b^5 + ((a^3 + 3*a^2 + 3*a + 1)*log(b*x + a + 1)^2 - 2*(a^3 + 3*a^2 + 3*a + 1)*log(b*x + a + 1)*log(b*x + a - 1) + (a^3 - 3*a^2 + 3*a - 1)*log(b*x + a - 1)^2 - 4*b*x - 2*(5*a^2 - 6*a + 1)*log(b*x + a - 1))/b^5) + 1/3*b*((b*x^2 - 4*a*x)/b^3 + (a^3 + 3*a^2 + 3*a + 1)*log(b*x + a + 1)/b^4 - (a^3 - 3*a^2 + 3*a - 1)*log(b*x + a - 1)/b^4)*arctanh(b*x + a)`**Giac [F]**

$$\int x^2 \operatorname{arctanh}(a + bx)^2 dx = \int x^2 \operatorname{artanh}(bx + a)^2 dx$$

input `integrate(x^2*arctanh(b*x+a)^2,x, algorithm="giac")`output `integrate(x^2*arctanh(b*x + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{arctanh}(a + bx)^2 dx = \int x^2 \operatorname{atanh}(a + bx)^2 dx$$

input `int(x^2*atanh(a + b*x)^2,x)`output `int(x^2*atanh(a + b*x)^2, x)`**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(a + bx)^2 dx$$

$$= \frac{-\operatorname{atanh}(bx + a)^2 a^4 + 2\operatorname{atanh}(bx + a)^2 a^2 + 2\operatorname{atanh}(bx + a)^2 a b^3 x^3 - \operatorname{atanh}(bx + a)^2 - 4\operatorname{atanh}(bx + a)}{}$$

input `int(x^2*atanh(b*x+a)^2,x)`output `(- atanh(a + b*x)**2*a**4 + 2*atanh(a + b*x)**2*a**2 + 2*atanh(a + b*x)**2*a*b**3*x**3 - atanh(a + b*x)**2 - 4*atanh(a + b*x)*a**3 - 2*atanh(a + b*x)*a**2*b*x - 6*atanh(a + b*x)*a**2 + 2*atanh(a + b*x)*a*b**2*x**2 + 2*atanh(a + b*x)*b*x + 2*atanh(a + b*x) - 6*int((atanh(a + b*x)*x**2)/(a**2 + 2*a*b*x + b**2*x**2 - 1),x)*a**2*b**3 - 2*int((atanh(a + b*x)*x**2)/(a**2 + 2*a*b*x + b**2*x**2 - 1),x)*b**3 - 6*log(a + b*x - 1)*a**2 + 2*log(a + b*x - 1) + 2*a*b*x)/(6*a*b**3)`

3.3 $\int x \operatorname{arctanh}(a + bx)^2 dx$

| | |
|---|----|
| Optimal result | 70 |
| Mathematica [A] (verified) | 71 |
| Rubi [A] (verified) | 71 |
| Maple [A] (verified) | 73 |
| Fricas [F] | 74 |
| Sympy [F] | 74 |
| Maxima [A] (verification not implemented) | 74 |
| Giac [F] | 75 |
| Mupad [F(-1)] | 75 |
| Reduce [F] | 76 |

Optimal result

Integrand size = 10, antiderivative size = 136

$$\int x \operatorname{arctanh}(a + bx)^2 dx = \frac{(a + bx) \operatorname{arctanh}(a + bx)}{b^2} - \frac{a \operatorname{arctanh}(a + bx)^2}{b^2} - \frac{(1 + a^2) \operatorname{arctanh}(a + bx)^2}{2b^2} + \frac{1}{2} x^2 \operatorname{arctanh}(a + bx)^2 + \frac{2a \operatorname{arctanh}(a + bx) \log\left(\frac{2}{1-a-bx}\right)}{b^2} + \frac{\log(1 - (a + bx)^2)}{2b^2} + \frac{a \operatorname{PolyLog}\left(2, -\frac{1+a+bx}{1-a-bx}\right)}{b^2}$$

output

```
(b*x+a)*arctanh(b*x+a)/b^2-a*arctanh(b*x+a)^2/b^2-1/2*(a^2+1)*arctanh(b*x+a)^2/b^2+1/2*x^2*arctanh(b*x+a)^2+2*a*arctanh(b*x+a)*ln(2/(-b*x-a+1))/b^2+1/2*ln(1-(b*x+a)^2)/b^2+a*polylog(2,-(b*x+a+1)/(-b*x-a+1))/b^2
```

Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.72

$$\int x \operatorname{arctanh}(a + bx)^2 dx$$

$$= \frac{(-1 + 2a - a^2 + b^2 x^2) \operatorname{arctanh}(a + bx)^2 + 2 \operatorname{arctanh}(a + bx) (a + bx + 2a \log(1 + e^{-2 \operatorname{arctanh}(a + bx)})) - 2 \log(1 + e^{-2 \operatorname{arctanh}(a + bx)})}{2b^2}$$

input

```
Integrate[x*ArcTanh[a + b*x]^2,x]
```

output

```
((-1 + 2*a - a^2 + b^2*x^2)*ArcTanh[a + b*x]^2 + 2*ArcTanh[a + b*x]*(a + b*x + 2*a*Log[1 + E^(-2*ArcTanh[a + b*x])]) - 2*Log[1/Sqrt[1 - (a + b*x)^2]] - 2*a*PolyLog[2, -E^(-2*ArcTanh[a + b*x])])/(2*b^2)
```

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.93, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6661, 25, 27, 6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(a + bx)^2 dx$$

$$\downarrow 6661$$

$$\frac{\int x \operatorname{arctanh}(a + bx)^2 d(a + bx)}{b}$$

$$\downarrow 25$$

$$-\frac{\int -x \operatorname{arctanh}(a + bx)^2 d(a + bx)}{b}$$

$$\downarrow 27$$

$$-\frac{\int -bx \operatorname{arctanh}(a + bx)^2 d(a + bx)}{b^2}$$

$$\int \left(\frac{(a^2 - 2(a+bx)a + 1) \operatorname{arctanh}(a+bx)}{1 - (a+bx)^2} - \operatorname{arctanh}(a+bx) \right) d(a+bx) - \frac{1}{2} b^2 x^2 \operatorname{arctanh}(a+bx)^2$$

↓ 6480

$$\frac{1}{b^2}$$

↓ 2009

$$\frac{\frac{1}{2}(a^2 + 1) \operatorname{arctanh}(a+bx)^2 - \frac{1}{2} b^2 x^2 \operatorname{arctanh}(a+bx)^2 + a \operatorname{arctanh}(a+bx)^2 - (a+bx) \operatorname{arctanh}(a+bx) - 2a \operatorname{arctanh}(a+bx)}{b^2}$$

input `Int[x*ArcTanh[a + b*x]^2,x]`

output `-((-((a + b*x)*ArcTanh[a + b*x]) + a*ArcTanh[a + b*x]^2 + ((1 + a^2)*ArcTanh[a + b*x]^2)/2 - (b^2*x^2*ArcTanh[a + b*x]^2)/2 - 2*a*ArcTanh[a + b*x]*Log[2/(1 - a - b*x)] - Log[1 - (a + b*x)^2]/2 - a*PolyLog[2, -((1 + a + b*x)/(1 - a - b*x))])/b^2)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6480 `Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)*((d_.) + (e_.)*(x_.))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]`

rule 6661

```
Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] :> Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*
ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IG
tQ[p, 0]
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.79

| method | result |
|-------------------|--|
| derivativedivides | $\frac{\operatorname{arctanh}(bx+a)^2(bx+a)^2}{2} - \operatorname{arctanh}(bx+a)^2(bx+a)a + \operatorname{arctanh}(bx+a)(bx+a) - \operatorname{arctanh}(bx+a) \ln(bx+a-1)a + \frac{\operatorname{arctanh}(bx+a)}{2}$ |
| default | $\frac{\operatorname{arctanh}(bx+a)^2(bx+a)^2}{2} - \operatorname{arctanh}(bx+a)^2(bx+a)a + \operatorname{arctanh}(bx+a)(bx+a) - \operatorname{arctanh}(bx+a) \ln(bx+a-1)a + \frac{\operatorname{arctanh}(bx+a)}{2}$ |
| parts | $\frac{x^2 \operatorname{arctanh}(bx+a)^2}{2} - \operatorname{arctanh}(bx+a)(bx+a) - \frac{\operatorname{arctanh}(bx+a) \ln(bx+a-1)a^2}{2} + \operatorname{arctanh}(bx+a) \ln(bx+a-1)a - \frac{\operatorname{arctanh}(bx+a)}{2}$ |
| risch | $\frac{x^2 \ln(-bx-a+1)^2}{8} + \frac{\ln(-bx-a-1)}{2b^2} + \frac{\ln(-bx-a+1)}{2b^2} - \frac{\ln(-bx-a+1)^2}{8b^2} + \left(-\frac{\ln(-bx-a+1)x^2}{4} + \frac{\ln(-bx-a-1)}{2} \right)$ |

input

```
int(x*arctanh(b*x+a)^2,x,method=_RETURNVERBOSE)
```

output

```
1/b^2*(1/2*arctanh(b*x+a)^2*(b*x+a)^2-arctanh(b*x+a)^2*(b*x+a)*a+arctanh(b
*x+a)*(b*x+a)-arctanh(b*x+a)*ln(b*x+a-1)*a+1/2*arctanh(b*x+a)*ln(b*x+a-1)-
arctanh(b*x+a)*ln(b*x+a+1)*a-1/2*arctanh(b*x+a)*ln(b*x+a+1)+1/2*ln(b*x+a-1
)+1/2*ln(b*x+a+1)+1/2*(-2*a+1)*(1/4*ln(b*x+a-1)^2-1/2*dilog(1/2*b*x+1/2*a
+1/2)-1/2*ln(b*x+a-1)*ln(1/2*b*x+1/2*a+1/2))+1/2*(-2*a-1)*(-1/4*ln(b*x+a+1
)^2+1/2*(ln(b*x+a+1)-ln(1/2*b*x+1/2*a+1/2))*ln(-1/2*b*x-1/2*a+1/2)-1/2*dilo
g(1/2*b*x+1/2*a+1/2))
```

Fricas [F]

$$\int x \operatorname{arctanh}(a + bx)^2 dx = \int x \operatorname{artanh}(bx + a)^2 dx$$

input `integrate(x*arctanh(b*x+a)^2,x, algorithm="fricas")`

output `integral(x*arctanh(b*x + a)^2, x)`

Sympy [F]

$$\int x \operatorname{arctanh}(a + bx)^2 dx = \int x \operatorname{atanh}^2(a + bx) dx$$

input `integrate(x*atanh(b*x+a)**2,x)`

output `Integral(x*atanh(a + b*x)**2, x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.49

$$\begin{aligned} \int x \operatorname{arctanh}(a + bx)^2 dx &= \frac{1}{2} x^2 \operatorname{artanh}(bx + a)^2 \\ &+ \frac{1}{8} b^2 \left(\frac{8 \left(\log(bx + a - 1) \log\left(\frac{1}{2}bx + \frac{1}{2}a + \frac{1}{2}\right) + \operatorname{Li}_2\left(-\frac{1}{2}bx - \frac{1}{2}a + \frac{1}{2}\right)\right) a}{b^4} + \frac{4(a + 1) \log(bx + a + 1)}{b^4} \right. \\ &+ \left. \frac{1}{2} b \left(\frac{2x}{b^2} - \frac{(a^2 + 2a + 1) \log(bx + a + 1)}{b^3} + \frac{(a^2 - 2a + 1) \log(bx + a - 1)}{b^3} \right) \operatorname{artanh}(bx \right. \\ &\quad \left. + a) \end{aligned}$$

input `integrate(x*arctanh(b*x+a)^2,x, algorithm="maxima")`

output

```
1/2*x^2*arctanh(b*x + a)^2 + 1/8*b^2*(8*(log(b*x + a - 1)*log(1/2*b*x + 1/
2*a + 1/2) + dilog(-1/2*b*x - 1/2*a + 1/2))*a/b^4 + 4*(a + 1)*log(b*x + a
+ 1)/b^4 + ((a^2 + 2*a + 1)*log(b*x + a + 1)^2 - 2*(a^2 + 2*a + 1)*log(b*x
+ a + 1)*log(b*x + a - 1) + (a^2 - 2*a + 1)*log(b*x + a - 1)^2 - 4*(a - 1
)*log(b*x + a - 1))/b^4) + 1/2*b*(2*x/b^2 - (a^2 + 2*a + 1)*log(b*x + a +
1)/b^3 + (a^2 - 2*a + 1)*log(b*x + a - 1)/b^3)*arctanh(b*x + a)
```

Giac [F]

$$\int x \operatorname{arctanh}(a + bx)^2 dx = \int x \operatorname{artanh}(bx + a)^2 dx$$

input

```
integrate(x*arctanh(b*x+a)^2,x, algorithm="giac")
```

output

```
integrate(x*arctanh(b*x + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{arctanh}(a + bx)^2 dx = \int x \operatorname{atanh}(a + bx)^2 dx$$

input

```
int(x*atanh(a + b*x)^2,x)
```

output

```
int(x*atanh(a + b*x)^2, x)
```

Reduce [F]

$$\int x \operatorname{arctanh}(a + bx)^2 dx = \frac{\operatorname{atanh}(bx + a)^2 x^2}{2} + \left(\int \frac{\operatorname{atanh}(bx + a) x^2}{b^2 x^2 + 2abx + a^2 - 1} dx \right) b$$

input

```
int(x*atanh(b*x+a)^2,x)
```

output

```
(atanh(a + b*x)**2*x**2 + 2*int((atanh(a + b*x)*x**2)/(a**2 + 2*a*b*x + b*
*2*x**2 - 1),x)*b)/2
```

3.4 $\int \operatorname{arctanh}(a + bx)^2 dx$

| | |
|---|----|
| Optimal result | 77 |
| Mathematica [A] (verified) | 77 |
| Rubi [A] (verified) | 78 |
| Maple [A] (verified) | 80 |
| Fricas [F] | 80 |
| Sympy [F] | 81 |
| Maxima [A] (verification not implemented) | 81 |
| Giac [F] | 82 |
| Mupad [F(-1)] | 82 |
| Reduce [F] | 82 |

Optimal result

Integrand size = 8, antiderivative size = 81

$$\int \operatorname{arctanh}(a + bx)^2 dx = \frac{\operatorname{arctanh}(a + bx)^2}{b} + \frac{(a + bx)\operatorname{arctanh}(a + bx)^2}{b} - \frac{2\operatorname{arctanh}(a + bx) \log\left(\frac{2}{1-a-bx}\right)}{b} - \frac{\operatorname{PolyLog}\left(2, -\frac{1+a+bx}{1-a-bx}\right)}{b}$$

output

```
arctanh(b*x+a)^2/b+(b*x+a)*arctanh(b*x+a)^2/b-2*arctanh(b*x+a)*ln(2/(-b*x-a+1))/b-polylog(2,-(b*x+a+1)/(-b*x-a+1))/b
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.68

$$\int \operatorname{arctanh}(a + bx)^2 dx = \frac{\operatorname{arctanh}(a + bx) \left((-1 + a + bx)\operatorname{arctanh}(a + bx) - 2 \log(1 + e^{-2\operatorname{arctanh}(a+bx)}) \right) + \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arctanh}(a+bx)}\right)}{b}$$

input

```
Integrate[ArcTanh[a + b*x]^2,x]
```

output

```
(ArcTanh[a + b*x]*((-1 + a + b*x)*ArcTanh[a + b*x] - 2*Log[1 + E^(-2*ArcTanh[a + b*x])]) + PolyLog[2, -E^(-2*ArcTanh[a + b*x])])/b
```

Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6653, 6436, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(a + bx)^2 dx$$

$$\downarrow 6653$$

$$\frac{\int \operatorname{arctanh}(a + bx)^2 d(a + bx)}{b}$$

$$\downarrow 6436$$

$$\frac{(a + bx)\operatorname{arctanh}(a + bx)^2 - 2 \int \frac{(a+bx)\operatorname{arctanh}(a+bx)}{1-(a+bx)^2} d(a + bx)}{b}$$

$$\downarrow 6546$$

$$\frac{(a + bx)\operatorname{arctanh}(a + bx)^2 - 2 \left(\int \frac{\operatorname{arctanh}(a+bx)}{-a-bx+1} d(a + bx) - \frac{1}{2}\operatorname{arctanh}(a + bx)^2 \right)}{b}$$

$$\downarrow 6470$$

$$\frac{(a + bx)\operatorname{arctanh}(a + bx)^2 - 2 \left(- \int \frac{\log\left(\frac{2}{-a-bx+1}\right)}{1-(a+bx)^2} d(a + bx) - \frac{1}{2}\operatorname{arctanh}(a + bx)^2 + \operatorname{arctanh}(a + bx) \log\left(\frac{2}{-a-bx+1}\right) \right)}{b}$$

$$\downarrow 2849$$

$$\frac{(a + bx)\operatorname{arctanh}(a + bx)^2 - 2 \left(\int \frac{\log\left(\frac{2}{-a-bx+1}\right)}{1-\frac{2}{-a-bx+1}} d\frac{1}{-a-bx+1} - \frac{1}{2}\operatorname{arctanh}(a + bx)^2 + \operatorname{arctanh}(a + bx) \log\left(\frac{2}{-a-bx+1}\right) \right)}{b}$$

$$\downarrow 2752$$

$$\frac{(a + bx)\operatorname{arctanh}(a + bx)^2 - 2\left(-\frac{1}{2}\operatorname{arctanh}(a + bx)^2 + \operatorname{arctanh}(a + bx)\log\left(\frac{2}{-a-bx+1}\right) + \frac{1}{2}\operatorname{PolyLog}\left(2, 1 - \frac{2}{-a-bx}\right)\right)}{b}$$

input `Int[ArcTanh[a + b*x]^2,x]`

output `((a + b*x)*ArcTanh[a + b*x]^2 - 2*(-1/2*ArcTanh[a + b*x]^2 + ArcTanh[a + b*x]*Log[2/(1 - a - b*x)] + PolyLog[2, 1 - 2/(1 - a - b*x)]/2))/b`

Defintions of rubi rules used

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 6436 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.), x_Symbol] := Simp[x*(a + b*ArcTanh[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`

rule 6470 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6546 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6653

```
Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.), x_Symbol] := Simp[1/d
  Subst[Int[(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d
}, x] && IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.12

| method | result |
|-------------------|---|
| derivativedivides | $\frac{\operatorname{arctanh}(bx+a)^2(bx+a-1)+2\operatorname{arctanh}(bx+a)^2-2\operatorname{arctanh}(bx+a)\ln\left(1+\frac{(bx+a+1)^2}{1-(bx+a)^2}\right)-\operatorname{polylog}\left(2,-\frac{(bx+a+1)^2}{1-(bx+a)^2}\right)}{b}$ |
| default | $\frac{\operatorname{arctanh}(bx+a)^2(bx+a-1)+2\operatorname{arctanh}(bx+a)^2-2\operatorname{arctanh}(bx+a)\ln\left(1+\frac{(bx+a+1)^2}{1-(bx+a)^2}\right)-\operatorname{polylog}\left(2,-\frac{(bx+a+1)^2}{1-(bx+a)^2}\right)}{b}$ |
| risch | $\frac{(bx+a+1)\ln(bx+a+1)^2}{4b} + \left(-\frac{\ln(-bx-a+1)x}{2} + \frac{-\ln(-bx-a+1)a+\ln(-bx-a+1)}{2b}\right)\ln(bx+a+1) + \frac{x\ln(bx+a+1)}{2b}$ |

input

```
int(arctanh(b*x+a)^2,x,method=_RETURNVERBOSE)
```

output

```
1/b*(arctanh(b*x+a)^2*(b*x+a-1)+2*arctanh(b*x+a)^2-2*arctanh(b*x+a)*ln(1+(
b*x+a+1)^2/(1-(b*x+a)^2))-polylog(2,-(b*x+a+1)^2/(1-(b*x+a)^2)))
```

Fricas [F]

$$\int \operatorname{arctanh}(a + bx)^2 dx = \int \operatorname{artanh}(bx + a)^2 dx$$

input

```
integrate(arctanh(b*x+a)^2,x, algorithm="fricas")
```

output

```
integral(arctanh(b*x + a)^2, x)
```

Sympy [F]

$$\int \operatorname{arctanh}(a + bx)^2 dx = \int \operatorname{atanh}^2(a + bx) dx$$

input `integrate(atanh(b*x+a)**2,x)`

output `Integral(atanh(a + b*x)**2, x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.72

$$\begin{aligned} \int \operatorname{arctanh}(a + bx)^2 dx = & \\ & -\frac{1}{4} b^2 \left(\frac{(a + 1) \log(bx + a + 1)^2 - 2(a + 1) \log(bx + a + 1) \log(bx + a - 1) + (a - 1) \log(bx + a - 1)}{b^3} \right. \\ & + b \left(\frac{(a + 1) \log(bx + a + 1)}{b^2} - \frac{(a - 1) \log(bx + a - 1)}{b^2} \right) \operatorname{artanh}(bx + a) \\ & \left. + x \operatorname{artanh}(bx + a)^2 \right) \end{aligned}$$

input `integrate(arctanh(b*x+a)^2,x, algorithm="maxima")`

output `-1/4*b^2*(((a + 1)*log(b*x + a + 1)^2 - 2*(a + 1)*log(b*x + a + 1)*log(b*x + a - 1) + (a - 1)*log(b*x + a - 1)^2)/b^3 + 4*(log(b*x + a - 1)*log(1/2*b*x + 1/2*a + 1/2) + dilog(-1/2*b*x - 1/2*a + 1/2))/b^3) + b*((a + 1)*log(b*x + a + 1)/b^2 - (a - 1)*log(b*x + a - 1)/b^2)*arctanh(b*x + a) + x*arctanh(b*x + a)^2`

Giac [F]

$$\int \operatorname{arctanh}(a + bx)^2 dx = \int \operatorname{artanh}(bx + a)^2 dx$$

input `integrate(arctanh(b*x+a)^2,x, algorithm="giac")`

output `integrate(arctanh(b*x + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \operatorname{arctanh}(a + bx)^2 dx = \int \operatorname{atanh}(a + bx)^2 dx$$

input `int(atanh(a + b*x)^2,x)`

output `int(atanh(a + b*x)^2, x)`

Reduce [F]

$$\int \operatorname{arctanh}(a + bx)^2 dx$$

$$= \frac{\operatorname{atanh}(bx + a)^2 a^2 + 2 \operatorname{atanh}(bx + a)^2 abx - \operatorname{atanh}(bx + a)^2 + 2 \operatorname{atanh}(bx + a) a + 2 \operatorname{atanh}(bx + a) bx + 2}{2ab}$$

input `int(atanh(b*x+a)^2,x)`

output `(atanh(a + b*x)**2*a**2 + 2*atanh(a + b*x)**2*a*b*x - atanh(a + b*x)**2 + 2*atanh(a + b*x)*a + 2*atanh(a + b*x)*b*x + 2*atanh(a + b*x) - 2*int((atanh(a + b*x)*x**2)/(a**2 + 2*a*b*x + b**2*x**2 - 1),x)*b**3 + 2*log(a + b*x - 1))/(2*a*b)`

3.5 $\int \frac{\operatorname{arctanh}(a+bx)^2}{x} dx$

| | |
|---|----|
| Optimal result | 83 |
| Mathematica [C] (warning: unable to verify) | 84 |
| Rubi [A] (verified) | 85 |
| Maple [C] (warning: unable to verify) | 87 |
| Fricas [F] | 88 |
| Sympy [F] | 88 |
| Maxima [F] | 88 |
| Giac [F] | 89 |
| Mupad [F(-1)] | 89 |
| Reduce [F] | 89 |

Optimal result

Integrand size = 12, antiderivative size = 148

$$\int \frac{\operatorname{arctanh}(a+bx)^2}{x} dx = -\operatorname{arctanh}(a+bx)^2 \log\left(\frac{2}{1+a+bx}\right) + \operatorname{arctanh}(a+bx)^2 \log\left(\frac{2bx}{(1-a)(1+a+bx)}\right) + \operatorname{arctanh}(a+bx) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+a+bx}\right) - \operatorname{arctanh}(a+bx) \operatorname{PolyLog}\left(2, 1 - \frac{2bx}{(1-a)(1+a+bx)}\right) + \frac{1}{2} \operatorname{PolyLog}\left(3, 1 - \frac{2}{1+a+bx}\right) - \frac{1}{2} \operatorname{PolyLog}\left(3, 1 - \frac{2bx}{(1-a)(1+a+bx)}\right)$$

output

```
-arctanh(b*x+a)^2*ln(2/(b*x+a+1))+arctanh(b*x+a)^2*ln(2*b*x/(1-a)/(b*x+a+1))+arctanh(b*x+a)*polylog(2,1-2/(b*x+a+1))-arctanh(b*x+a)*polylog(2,1-2*b*x/(1-a)/(b*x+a+1))+1/2*polylog(3,1-2/(b*x+a+1))-1/2*polylog(3,1-2*b*x/(1-a)/(b*x+a+1))
```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 2.26 (sec) , antiderivative size = 753, normalized size of antiderivative = 5.09

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x} dx = \text{Too large to display}$$

input `Integrate[ArcTanh[a + b*x]^2/x,x]`

output

```
(-4*ArcTanh[a + b*x]^3)/3 - (2*ArcTanh[a + b*x]^3)/(3*a) + (2*Sqrt[1 - a^2]
)*E^ArcTanh[a]*ArcTanh[a + b*x]^3)/(3*a) - ArcTanh[a + b*x]^2*Log[1 + E^(-
2*ArcTanh[a + b*x])] - I*Pi*ArcTanh[a + b*x]*Log[(E^(-ArcTanh[a + b*x])) +
E^ArcTanh[a + b*x])/2] - ArcTanh[a + b*x]^2*Log[1 - (Sqrt[-1 + a]*E^ArcTan
h[a + b*x])/Sqrt[-1 - a]] - ArcTanh[a + b*x]^2*Log[1 + (Sqrt[-1 + a]*E^Arc
Tanh[a + b*x])/Sqrt[-1 - a]] + ArcTanh[a + b*x]^2*Log[(1 + a - E^(2*ArcTan
h[a + b*x])) + a*E^(2*ArcTanh[a + b*x])]/(2*E^ArcTanh[a + b*x])] + ArcTanh[
a + b*x]^2*Log[1 - E^(-ArcTanh[a] + ArcTanh[a + b*x])] + ArcTanh[a + b*x]^
2*Log[1 + E^(-ArcTanh[a] + ArcTanh[a + b*x])] - 2*ArcTanh[a]*ArcTanh[a + b
*x]*Log[(I/2)*(-E^(ArcTanh[a] - ArcTanh[a + b*x])) + E^(-ArcTanh[a] + ArcTa
nh[a + b*x])] + ArcTanh[a + b*x]^2*Log[1 - E^(-2*ArcTanh[a] + 2*ArcTanh[a
+ b*x])] + I*Pi*ArcTanh[a + b*x]*Log[1/Sqrt[1 - (a + b*x)^2]] - ArcTanh[a
+ b*x]^2*Log[-((b*x)/Sqrt[1 - (a + b*x)^2])] + 2*ArcTanh[a]*ArcTanh[a + b
*x]*Log[(-I)*Sinh[ArcTanh[a] - ArcTanh[a + b*x]]] + ArcTanh[a + b*x]*PolyL
og[2, -E^(-2*ArcTanh[a + b*x])] - 2*ArcTanh[a + b*x]*PolyLog[2, -((Sqrt[-1
+ a]*E^ArcTanh[a + b*x])/Sqrt[-1 - a])] - 2*ArcTanh[a + b*x]*PolyLog[2, (
Sqrt[-1 + a]*E^ArcTanh[a + b*x])/Sqrt[-1 - a]] + 2*ArcTanh[a + b*x]*PolyLo
g[2, -E^(-ArcTanh[a] + ArcTanh[a + b*x])] + 2*ArcTanh[a + b*x]*PolyLog[2,
E^(-ArcTanh[a] + ArcTanh[a + b*x])] + ArcTanh[a + b*x]*PolyLog[2, E^(-2*Ar
cTanh[a] + 2*ArcTanh[a + b*x])] + PolyLog[3, -E^(-2*ArcTanh[a + b*x])]/...
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6661, 25, 27, 6474}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arctanh}(a+bx)^2}{x} dx \\
 & \quad \downarrow \text{6661} \\
 & \frac{\int \frac{\operatorname{arctanh}(a+bx)^2}{x} d(a+bx)}{b} \\
 & \quad \downarrow \text{25} \\
 & -\frac{\int -\frac{\operatorname{arctanh}(a+bx)^2}{x} d(a+bx)}{b} \\
 & \quad \downarrow \text{27} \\
 & -\int -\frac{\operatorname{arctanh}(a+bx)^2}{bx} d(a+bx) \\
 & \quad \downarrow \text{6474} \\
 & \operatorname{arctanh}(a+bx) \operatorname{PolyLog}\left(2, 1 - \frac{2}{a+bx+1}\right) - \operatorname{arctanh}(a+bx) \operatorname{PolyLog}\left(2, 1 - \frac{2bx}{(1-a)(a+bx+1)}\right) + \operatorname{arctanh}(a+bx)^2 \left(-\log\left(\frac{2}{a+bx+1}\right)\right) + \\
 & \operatorname{arctanh}(a+bx)^2 \log\left(\frac{2bx}{(1-a)(a+bx+1)}\right) + \frac{1}{2} \operatorname{PolyLog}\left(3, 1 - \frac{2}{a+bx+1}\right) - \\
 & \frac{1}{2} \operatorname{PolyLog}\left(3, 1 - \frac{2bx}{(1-a)(a+bx+1)}\right)
 \end{aligned}$$

input

```
Int[ArcTanh[a + b*x]^2/x,x]
```

output

```
-(ArcTanh[a + b*x]^2*Log[2/(1 + a + b*x)]) + ArcTanh[a + b*x]^2*Log[(2*b*x)
)/((1 - a)*(1 + a + b*x))] + ArcTanh[a + b*x]*PolyLog[2, 1 - 2/(1 + a + b*
x)] - ArcTanh[a + b*x]*PolyLog[2, 1 - (2*b*x)/((1 - a)*(1 + a + b*x))] + P
olyLog[3, 1 - 2/(1 + a + b*x)]/2 - PolyLog[3, 1 - (2*b*x)/((1 - a)*(1 + a
+ b*x))]/2
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

rule 6474

```
Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^2/((d_) + (e_)*(x_)), x_Symbol] :=
Simp[(-(a + b*ArcTanh[c*x])^2)*(Log[2/(1 + c*x)]/e), x] + (Simp[(a + b*Arc
Tanh[c*x])^2*(Log[2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/e), x] + Simp[b*(a
+ b*ArcTanh[c*x])*(PolyLog[2, 1 - 2/(1 + c*x)]/e), x] - Simp[b*(a + b*ArcT
anh[c*x])*(PolyLog[2, 1 - 2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/e), x] + S
imp[b^2*(PolyLog[3, 1 - 2/(1 + c*x)]/(2*e)), x] - Simp[b^2*(PolyLog[3, 1 -
2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/(2*e)), x]) /; FreeQ[{a, b, c, d, e}
, x] && NeQ[c^2*d^2 - e^2, 0]
```

rule 6661

```
Int[((a_) + ArcTanh[(c_) + (d_)*(x_)])*(b_)^(p_)*((e_) + (f_)*(x_))^(
m_), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*
ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IG
tQ[p, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 32.32 (sec) , antiderivative size = 901, normalized size of antiderivative = 6.09

| method | result | size |
|--------------------|---------------------------------|------|
| derivativeldivides | Expression too large to display | 901 |
| default | Expression too large to display | 901 |
| parts | Expression too large to display | 1544 |

input `int(arctanh(b*x+a)^2/x,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & \ln(-b*x)*\operatorname{arctanh}(b*x+a)^2 - \operatorname{arctanh}(b*x+a)^2 * \ln\left(-\frac{(b*x+a+1)^2}{(1-(b*x+a)^2)} + 1 + a * \left(1 + \frac{(b*x+a+1)^2}{(1-(b*x+a)^2)}\right)\right) \\ & + \frac{1}{2} * I * \pi * \operatorname{csgn}\left(I * \frac{(b*x+a+1)^2}{(b*x+a)^2 - 1} + 1 + a * \frac{(1-(b*x+a+1)^2)}{(b*x+a)^2 - 1}\right) / \left(1 - \frac{(b*x+a+1)^2}{(b*x+a)^2 - 1}\right) * \operatorname{csgn}\left(I * \frac{(b*x+a+1)^2}{(b*x+a)^2 - 1} + 1 + a * \frac{(1-(b*x+a+1)^2)}{(b*x+a)^2 - 1}\right) * \operatorname{csgn}\left(I / \left(1 - \frac{(b*x+a+1)^2}{(b*x+a)^2 - 1}\right) - \operatorname{csgn}\left(I * \frac{(b*x+a+1)^2}{(b*x+a)^2 - 1} + 1 + a * \frac{(1-(b*x+a+1)^2}{(b*x+a)^2 - 1}\right) / \left(1 - \frac{(b*x+a+1)^2}{(b*x+a)^2 - 1}\right) * \operatorname{csgn}\left(I / \left(1 - \frac{(b*x+a+1)^2}{(b*x+a)^2 - 1}\right) - \operatorname{csgn}\left(I * \frac{(b*x+a+1)^2}{(b*x+a)^2 - 1} + 1 + a * \frac{(1-(b*x+a+1)^2}{(b*x+a)^2 - 1}\right) / \left(1 - \frac{(b*x+a+1)^2}{(b*x+a)^2 - 1}\right) * \operatorname{csgn}\left(I * \frac{(b*x+a+1)^2}{(b*x+a)^2 - 1} + 1 + a * \frac{(1-(b*x+a+1)^2}{(b*x+a)^2 - 1}\right) / \left(1 - \frac{(b*x+a+1)^2}{(b*x+a)^2 - 1}\right) + \operatorname{csgn}\left(I * \frac{(b*x+a+1)^2}{(b*x+a)^2 - 1} + 1 + a * \frac{(1-(b*x+a+1)^2}{(b*x+a)^2 - 1}\right) / \left(1 - \frac{(b*x+a+1)^2}{(b*x+a)^2 - 1}\right)\right)^2 * \operatorname{arctanh}(b*x+a)^2 - \operatorname{arctanh}(b*x+a) * \operatorname{polylog}\left(2, -\frac{(b*x+a+1)^2}{(1-(b*x+a)^2)} + 1\right) / 2 * \operatorname{polylog}\left(3, -\frac{(b*x+a+1)^2}{(1-(b*x+a)^2)} + a / (a-1) * \operatorname{arctanh}(b*x+a)^2 * \ln(1-(a-1) * \frac{(b*x+a+1)^2}{(1-(b*x+a)^2)} / (-1-a)) + a / (a-1) * \operatorname{arctanh}(b*x+a) * \operatorname{polylog}\left(2, (a-1) * \frac{(b*x+a+1)^2}{(1-(b*x+a)^2)} / (-1-a) - 1/2 * a / (a-1) * \operatorname{polylog}\left(3, (a-1) * \frac{(b*x+a+1)^2}{(1-(b*x+a)^2)} / (-1-a) - 1 / (a-1) * \operatorname{arctanh}(b*x+a)^2 * \ln(1-(a-1) * \frac{(b*x+a+1)^2}{(1-(b*x+a)^2)} / (-1-a) - 1 / (a-1) * \operatorname{arctanh}(b*x+a) * \operatorname{polylog}\left(2, (a-1) * \frac{(b*x+a+1)^2}{(1-(b*x+a)^2)} / (-1-a) + 1/2 / (a-1) * \operatorname{polylog}\left(3, (a-1) * \frac{(b*x+a+1)^2}{(1-(b*x+a)^2)} / (-1-a)\right)\right)\right) \end{aligned}$$

Fricas [F]

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x} dx = \int \frac{\operatorname{artanh}(bx + a)^2}{x} dx$$

input `integrate(arctanh(b*x+a)^2/x,x, algorithm="fricas")`

output `integral(arctanh(b*x + a)^2/x, x)`

Sympy [F]

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x} dx = \int \frac{\operatorname{atanh}^2(a + bx)}{x} dx$$

input `integrate(atanh(b*x+a)**2/x,x)`

output `Integral(atanh(a + b*x)**2/x, x)`

Maxima [F]

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x} dx = \int \frac{\operatorname{artanh}(bx + a)^2}{x} dx$$

input `integrate(arctanh(b*x+a)^2/x,x, algorithm="maxima")`

output `integrate(arctanh(b*x + a)^2/x, x)`

Giac [F]

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x} dx = \int \frac{\operatorname{artanh}(bx + a)^2}{x} dx$$

input `integrate(arctanh(b*x+a)^2/x,x, algorithm="giac")`

output `integrate(arctanh(b*x + a)^2/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x} dx = \int \frac{\operatorname{atanh}(a + bx)^2}{x} dx$$

input `int(atanh(a + b*x)^2/x,x)`

output `int(atanh(a + b*x)^2/x, x)`

Reduce [F]

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x} dx = \int \frac{\operatorname{atanh}(bx + a)^2}{x} dx$$

input `int(atanh(b*x+a)^2/x,x)`

output `int(atanh(a + b*x)**2/x,x)`

3.6 $\int \frac{\operatorname{arctanh}(a+bx)^2}{x^2} dx$

| | |
|---|----|
| Optimal result | 90 |
| Mathematica [C] (warning: unable to verify) | 91 |
| Rubi [A] (verified) | 91 |
| Maple [A] (verified) | 94 |
| Fricas [F] | 95 |
| Sympy [F] | 95 |
| Maxima [A] (verification not implemented) | 95 |
| Giac [F] | 96 |
| Mupad [F(-1)] | 96 |
| Reduce [F] | 97 |

Optimal result

Integrand size = 12, antiderivative size = 251

$$\int \frac{\operatorname{arctanh}(a+bx)^2}{x^2} dx = -\frac{\operatorname{arctanh}(a+bx)^2}{x} + \frac{b \operatorname{arctanh}(a+bx) \log\left(\frac{2}{1-a-bx}\right)}{1-a}$$

$$+ \frac{b \operatorname{arctanh}(a+bx) \log\left(\frac{2}{1+a+bx}\right)}{1+a}$$

$$- \frac{2b \operatorname{arctanh}(a+bx) \log\left(\frac{2}{1+a+bx}\right)}{1-a^2}$$

$$+ \frac{2b \operatorname{arctanh}(a+bx) \log\left(\frac{2bx}{(1-a)(1+a+bx)}\right)}{1-a^2}$$

$$+ \frac{b \operatorname{PolyLog}\left(2, -\frac{1+a+bx}{1-a-bx}\right)}{2(1-a)} - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+a+bx}\right)}{2(1+a)}$$

$$+ \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+a+bx}\right)}{1-a^2}$$

$$- \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2bx}{(1-a)(1+a+bx)}\right)}{1-a^2}$$

output

```
-arctanh(b*x+a)^2/x+b*arctanh(b*x+a)*ln(2/(-b*x-a+1))/(1-a)+b*arctanh(b*x+a)*ln(2/(b*x+a+1))/(1+a)-2*b*arctanh(b*x+a)*ln(2/(b*x+a+1))/(-a^2+1)+2*b*arctanh(b*x+a)*ln(2*b*x/(1-a)/(b*x+a+1))/(-a^2+1)+b*polylog(2,-(b*x+a+1)/(-b*x-a+1))/(2-2*a)-b*polylog(2,1-2/(b*x+a+1))/(2+2*a)+b*polylog(2,1-2/(b*x+a+1))/(-a^2+1)-b*polylog(2,1-2*b*x/(1-a)/(b*x+a+1))/(-a^2+1)
```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.93 (sec) , antiderivative size = 208, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x^2} dx$$

$$= \frac{-((-a + a^3 + a^2bx + b(-1 + \sqrt{1 - a^2}e^{\operatorname{arctanh}(a)})x) \operatorname{arctanh}(a + bx)^2) + abx \operatorname{arctanh}(a + bx) (-i\pi + 2i \operatorname{arctanh}(a))}{x^2}$$

input

```
Integrate[ArcTanh[a + b*x]^2/x^2,x]
```

output

```
((-((-a + a^3 + a^2*b*x + b*(-1 + Sqrt[1 - a^2])*E^ArcTanh[a])*x)*ArcTanh[a + b*x]^2) + a*b*x*ArcTanh[a + b*x]*((-I)*Pi + 2*ArcTanh[a] - 2*Log[1 - E^(2*ArcTanh[a] - 2*ArcTanh[a + b*x])]) + a*b*x*(I*Pi*(Log[1 + E^(2*ArcTanh[a + b*x])]) - Log[1/Sqrt[1 - (a + b*x)^2]]) + 2*ArcTanh[a]*(Log[1 - E^(2*ArcTanh[a] - 2*ArcTanh[a + b*x])]) - Log[(-I)*Sinh[ArcTanh[a] - ArcTanh[a + b*x]]])) + a*b*x*PolyLog[2, E^(2*ArcTanh[a] - 2*ArcTanh[a + b*x])]/(a*(-1 + a^2)*x)
```

Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.02, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {6659, 7292, 6671, 25, 27, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{\operatorname{arctanh}(a + bx)^2}{x^2} dx \\
& \quad \downarrow \text{6659} \\
& 2b \int \frac{\operatorname{arctanh}(a + bx)}{x(1 - (a + bx)^2)} dx - \frac{\operatorname{arctanh}(a + bx)^2}{x} \\
& \quad \downarrow \text{7292} \\
& 2b \int \frac{\operatorname{arctanh}(a + bx)}{x(-a^2 - 2bxa - b^2x^2 + 1)} dx - \frac{\operatorname{arctanh}(a + bx)^2}{x} \\
& \quad \downarrow \text{6671} \\
& 2 \int \frac{\operatorname{arctanh}(a + bx)}{x(1 - (a + bx)^2)} d(a + bx) - \frac{\operatorname{arctanh}(a + bx)^2}{x} \\
& \quad \downarrow \text{25} \\
& -2 \int -\frac{\operatorname{arctanh}(a + bx)}{x(1 - (a + bx)^2)} d(a + bx) - \frac{\operatorname{arctanh}(a + bx)^2}{x} \\
& \quad \downarrow \text{27} \\
& -2b \int -\frac{\operatorname{arctanh}(a + bx)}{bx(1 - (a + bx)^2)} d(a + bx) - \frac{\operatorname{arctanh}(a + bx)^2}{x} \\
& \quad \downarrow \text{7276} \\
& -2b \int \left(\frac{\operatorname{arctanh}(a + bx)}{(a^2 - 1)bx} - \frac{\operatorname{arctanh}(a + bx)}{2(a - 1)(a + bx - 1)} + \frac{\operatorname{arctanh}(a + bx)}{2(a + 1)(a + bx + 1)} \right) d(a + bx) - \\
& \quad \frac{\operatorname{arctanh}(a + bx)^2}{x} \\
& \quad \downarrow \text{2009} \\
& -2b \left(\frac{\operatorname{arctanh}(a + bx) \log\left(\frac{2}{a + bx + 1}\right)}{1 - a^2} - \frac{\operatorname{arctanh}(a + bx) \log\left(\frac{2bx}{(1-a)(a + bx + 1)}\right)}{1 - a^2} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{a + bx + 1}\right)}{2(1 - a^2)} + \frac{\operatorname{PolyLog}\left(2, \frac{2bx}{(1-a)(a + bx + 1)}\right)}{2(1 - a^2)} \right) \\
& \quad \frac{\operatorname{arctanh}(a + bx)^2}{x}
\end{aligned}$$

input

Int[ArcTanh[a + b*x]^2/x^2,x]

output
$$\begin{aligned} & -(\text{ArcTanh}[a + b*x]^2/x) - 2*b*(-1/2*(\text{ArcTanh}[a + b*x]*\text{Log}[2/(1 - a - b*x)] \\ &)/(1 - a) - (\text{ArcTanh}[a + b*x]*\text{Log}[2/(1 + a + b*x)])/(2*(1 + a)) + (\text{ArcTanh} \\ & [a + b*x]*\text{Log}[2/(1 + a + b*x)]/(1 - a^2) - (\text{ArcTanh}[a + b*x]*\text{Log}[(2*b*x)/ \\ & ((1 - a)*(1 + a + b*x))]/(1 - a^2) - \text{PolyLog}[2, -((1 + a + b*x)/(1 - a - \\ & b*x)]/(4*(1 - a)) + \text{PolyLog}[2, 1 - 2/(1 + a + b*x)]/(4*(1 + a)) - \text{PolyLog} \\ & [2, 1 - 2/(1 + a + b*x)]/(2*(1 - a^2)) + \text{PolyLog}[2, 1 - (2*b*x)/((1 - a)*(\\ & 1 + a + b*x)]/(2*(1 - a^2))) \end{aligned}$$

Defintions of rubi rules used

rule 25
$$\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \text{ :> } \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$$

rule 27
$$\text{Int}[(a_)*(\text{Fx}_), \text{x_Symbol}] \text{ :> } \text{Simp}[a \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ /; } \text{FreeQ}[a, \text{x}] \ \&\& \ \text{!Ma} \\ \text{tchQ}[\text{Fx}, (b_)*(\text{Gx}_)] \text{ /; } \text{FreeQ}[b, \text{x}]$$

rule 2009
$$\text{Int}[u_, \text{x_Symbol}] \text{ :> } \text{Simp}[\text{IntSum}[u, \text{x}], \text{x}] \text{ /; } \text{SumQ}[u]$$

rule 6659
$$\text{Int}[((a_.) + \text{ArcTanh}[(c_) + (d_.)*(x_)]*(b_.))^{\text{(p_.)*((e_.) + (f_.)*(x_))^{\text{(m_.)}}}, \text{x_Symbol}] \text{ :> } \text{Simp}[(e + f*x)^{\text{(m + 1)}}*(a + b*\text{ArcTanh}[c + d*x])^{\text{p}}/(f*(m \\ + 1))), \text{x}] - \text{Simp}[b*d*(p/(f*(m + 1))) \quad \text{Int}[(e + f*x)^{\text{(m + 1)}}*(a + b*\text{ArcTa} \\ \text{nh}[c + d*x])^{\text{(p - 1)}}/(1 - (c + d*x)^2)), \text{x}], \text{x}] \text{ /; } \text{FreeQ}[\{a, b, c, d, e, f\} \\ , \text{x}] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{ILtQ}[m, -1]$$

rule 6671
$$\text{Int}[((a_.) + \text{ArcTanh}[(c_) + (d_.)*(x_)]*(b_.))^{\text{(p_.)*((e_.) + (f_.)*(x_))^{\text{(m_.)}}*((A_.) + (B_.)*(x_) + (C_.)*(x_)^2)^{\text{(q_.)}}}, \text{x_Symbol}] \text{ :> } \text{Simp}[1/d \quad \text{Sub} \\ \text{st}[\text{Int}[(d*e - c*f)/d + f*(x/d)]^{\text{m}}*(-C/d^2 + (C/d^2)*x^2)^{\text{q}}*(a + b*\text{ArcTanh}[\\ x])^{\text{p}}, \text{x}], \text{x}, c + d*x], \text{x}] \text{ /; } \text{FreeQ}[\{a, b, c, d, e, f, A, B, C, m, p, q\}, \text{x} \\] \ \&\& \ \text{EqQ}[B*(1 - c^2) + 2*A*c*d, 0] \ \&\& \ \text{EqQ}[2*c*C - B*d, 0]$$

rule 7276
$$\text{Int}[(u_)/((a_) + (b_.)*(x_)^{\text{(n)}}), \text{x_Symbol}] \text{ :> } \text{With}[\{v = \text{RationalFunctionE} \\ \text{xpend}[u/(a + b*x^{\text{n}}), \text{x}]\}, \text{Int}[v, \text{x}] \text{ /; } \text{SumQ}[v]] \text{ /; } \text{FreeQ}[\{a, b\}, \text{x}] \ \&\& \ \text{IGtQ} \\ [n, 0]$$

rule 7292

```
Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v !=
= u]
```

Maple [A] (verified)

Time = 1.60 (sec) , antiderivative size = 301, normalized size of antiderivative = 1.20

| method | result |
|-------------------|--|
| parts | $-\frac{\operatorname{arctanh}(bx+a)^2}{x} + 2b \left(-\frac{\operatorname{arctanh}(bx+a) \ln(-bx)}{(a-1)(a+1)} - \frac{\operatorname{arctanh}(bx+a) \ln(bx+a+1)}{2a+2} + \frac{\operatorname{arctanh}(bx+a) \ln(bx+a-1)}{2a-2} \right)$ |
| derivativedivides | $b \left(-\frac{\operatorname{arctanh}(bx+a)^2}{bx} - \frac{2 \operatorname{arctanh}(bx+a) \ln(-bx)}{(a-1)(a+1)} - \frac{2 \operatorname{arctanh}(bx+a) \ln(bx+a+1)}{2a+2} + \frac{2 \operatorname{arctanh}(bx+a) \ln(bx+a-1)}{2a-2} \right)$ |
| default | $b \left(-\frac{\operatorname{arctanh}(bx+a)^2}{bx} - \frac{2 \operatorname{arctanh}(bx+a) \ln(-bx)}{(a-1)(a+1)} - \frac{2 \operatorname{arctanh}(bx+a) \ln(bx+a+1)}{2a+2} + \frac{2 \operatorname{arctanh}(bx+a) \ln(bx+a-1)}{2a-2} \right)$ |

input

```
int(arctanh(b*x+a)^2/x^2,x,method=_RETURNVERBOSE)
```

output

```
-arctanh(b*x+a)^2/x+2*b*(-arctanh(b*x+a)/(a-1)/(a+1)*ln(-b*x)-arctanh(b*x+a)/(2*a+2)*ln(b*x+a+1)+arctanh(b*x+a)/(2*a-2)*ln(b*x+a-1)-1/(a-1)/(a+1)*(-1/2*dilog((-b*x-a-1)/(-1-a))-1/2*ln(-b*x)*ln((-b*x-a-1)/(-1-a))+1/2*dilog(1/(1-a)*(-b*x-a+1))+1/2*ln(-b*x)*ln(1/(1-a)*(-b*x-a+1)))+1/2/(a-1)*(1/4*ln(b*x+a-1)^2-1/2*dilog(1/2*b*x+1/2*a+1/2)-1/2*ln(b*x+a-1)*ln(1/2*b*x+1/2*a+1/2))-1/2/(a+1)*(-1/4*ln(b*x+a+1)^2+1/2*(ln(b*x+a+1)-ln(1/2*b*x+1/2*a+1/2))*ln(-1/2*b*x-1/2*a+1/2)-1/2*dilog(1/2*b*x+1/2*a+1/2)))
```

Fricas [F]

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{artanh}(bx + a)^2}{x^2} dx$$

input `integrate(arctanh(b*x+a)^2/x^2,x, algorithm="fricas")`

output `integral(arctanh(b*x + a)^2/x^2, x)`

Sympy [F]

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{atanh}^2(a + bx)}{x^2} dx$$

input `integrate(atanh(b*x+a)**2/x**2,x)`

output `Integral(atanh(a + b*x)**2/x**2, x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 244, normalized size of antiderivative = 0.97

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(a + bx)^2}{x^2} dx \\ &= \frac{1}{4} b^2 \left(\frac{(a - 1) \log(bx + a + 1)^2 - 2(a - 1) \log(bx + a + 1) \log(bx + a - 1) + (a + 1) \log(bx + a - 1)^2}{a^2 b - b} \right. \\ & \quad \left. - b \left(\frac{\log(bx + a + 1)}{a + 1} - \frac{\log(bx + a - 1)}{a - 1} + \frac{2 \log(x)}{a^2 - 1} \right) \operatorname{artanh}(bx + a) \right. \\ & \quad \left. - \frac{\operatorname{artanh}(bx + a)^2}{x} \right) \end{aligned}$$

input `integrate(arctanh(b*x+a)^2/x^2,x, algorithm="maxima")`

output

```
1/4*b^2*((a - 1)*log(b*x + a + 1)^2 - 2*(a - 1)*log(b*x + a + 1)*log(b*x
+ a - 1) + (a + 1)*log(b*x + a - 1)^2)/(a^2*b - b) - 4*(log(b*x + a - 1)*l
og(1/2*b*x + 1/2*a + 1/2) + dilog(-1/2*b*x - 1/2*a + 1/2))/(a^2*b - b) + 4
*(log(b*x/(a + 1) + 1)*log(x) + dilog(-b*x/(a + 1)))/(a^2*b - b) - 4*(log(
b*x/(a - 1) + 1)*log(x) + dilog(-b*x/(a - 1)))/(a^2*b - b) - b*(log(b*x +
a + 1)/(a + 1) - log(b*x + a - 1)/(a - 1) + 2*log(x)/(a^2 - 1))*arctanh(b
*x + a) - arctanh(b*x + a)^2/x
```

Giac [F]

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{artanh}(bx + a)^2}{x^2} dx$$

input

```
integrate(arctanh(b*x+a)^2/x^2,x, algorithm="giac")
```

output

```
integrate(arctanh(b*x + a)^2/x^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{atanh}(a + bx)^2}{x^2} dx$$

input

```
int(atanh(a + b*x)^2/x^2,x)
```

output

```
int(atanh(a + b*x)^2/x^2, x)
```

Reduce [F]

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x^2} dx$$

$$= \frac{-2\operatorname{atanh}(bx + a)^2 a^3 - \operatorname{atanh}(bx + a)^2 a^2 bx + 2\operatorname{atanh}(bx + a)^2 a + \operatorname{atanh}(bx + a)^2 bx + 2\operatorname{atanh}(bx + a)}$$

input `int(atanh(b*x+a)^2/x^2,x)`

output `(- 2*atanh(a + b*x)**2*a**3 - atanh(a + b*x)**2*a**2*b*x + 2*atanh(a + b*x)**2*a + atanh(a + b*x)**2*b*x + 2*atanh(a + b*x)*a**2 + 2*atanh(a + b*x)*a*b*x - 2*atanh(a + b*x)*b*x - 2*atanh(a + b*x) + 2*int(atanh(a + b*x)/(a**2*x**2 + 2*a*b*x**3 + b**2*x**4 - x**2),x)*a**4*x - 4*int(atanh(a + b*x)/(a**2*x**2 + 2*a*b*x**3 + b**2*x**4 - x**2),x)*a**2*x + 2*int(atanh(a + b*x)/(a**2*x**2 + 2*a*b*x**3 + b**2*x**4 - x**2),x)*x - 2*log(a + b*x - 1)*b*x + 2*log(x)*b*x)/(2*a*x*(a**2 - 1))`

3.7 $\int \frac{\operatorname{arctanh}(a+bx)^2}{x^3} dx$

| | |
|---|-----|
| Optimal result | 98 |
| Mathematica [C] (warning: unable to verify) | 99 |
| Rubi [A] (verified) | 100 |
| Maple [A] (verified) | 102 |
| Fricas [F] | 103 |
| Sympy [F] | 103 |
| Maxima [A] (verification not implemented) | 103 |
| Giac [F] | 104 |
| Mupad [F(-1)] | 104 |
| Reduce [F] | 105 |

Optimal result

Integrand size = 12, antiderivative size = 370

$$\begin{aligned}
 \int \frac{\operatorname{arctanh}(a+bx)^2}{x^3} dx = & -\frac{b\operatorname{arctanh}(a+bx)}{(1-a^2)x} - \frac{\operatorname{arctanh}(a+bx)^2}{2x^2} \\
 & + \frac{b^2 \log(x)}{(1-a^2)^2} + \frac{b^2 \operatorname{arctanh}(a+bx) \log\left(\frac{2}{1-a-bx}\right)}{2(1-a)^2} \\
 & - \frac{b^2 \log(1-a-bx)}{2(1-a)^2(1+a)} - \frac{b^2 \operatorname{arctanh}(a+bx) \log\left(\frac{2}{1+a+bx}\right)}{2(1+a)^2} \\
 & - \frac{2ab^2 \operatorname{arctanh}(a+bx) \log\left(\frac{2}{1+a+bx}\right)}{(1-a^2)^2} \\
 & + \frac{2ab^2 \operatorname{arctanh}(a+bx) \log\left(\frac{2bx}{(1-a)(1+a+bx)}\right)}{(1-a^2)^2} \\
 & - \frac{b^2 \log(1+a+bx)}{2(1-a)(1+a)^2} + \frac{b^2 \operatorname{PolyLog}\left(2, -\frac{1+a+bx}{1-a-bx}\right)}{4(1-a)^2} \\
 & + \frac{b^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+a+bx}\right)}{4(1+a)^2} + \frac{ab^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+a+bx}\right)}{(1-a^2)^2} \\
 & - \frac{ab^2 \operatorname{PolyLog}\left(2, 1 - \frac{2bx}{(1-a)(1+a+bx)}\right)}{(1-a^2)^2}
 \end{aligned}$$

output

```
-b*arctanh(b*x+a)/(-a^2+1)/x-1/2*arctanh(b*x+a)^2/x^2+b^2*ln(x)/(-a^2+1)^2
+1/2*b^2*arctanh(b*x+a)*ln(2/(-b*x-a+1))/(1-a)^2-1/2*b^2*ln(-b*x-a+1)/(1-a
)^2/(1+a)-1/2*b^2*arctanh(b*x+a)*ln(2/(b*x+a+1))/(1+a)^2-2*a*b^2*arctanh(b
*x+a)*ln(2/(b*x+a+1))/(-a^2+1)^2+2*a*b^2*arctanh(b*x+a)*ln(2*b*x/(1-a)/(b*
x+a+1))/(-a^2+1)^2-1/2*b^2*ln(b*x+a+1)/(1-a)/(1+a)^2+1/4*b^2*polylog(2,-(b
*x+a+1)/(-b*x-a+1))/(1-a)^2+1/4*b^2*polylog(2,1-2/(b*x+a+1))/(1+a)^2+a*b^2
*polylog(2,1-2/(b*x+a+1))/(-a^2+1)^2-a*b^2*polylog(2,1-2*b*x/(1-a)/(b*x+a
1))/(-a^2+1)^2
```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 1.59 (sec) , antiderivative size = 271, normalized size of antiderivative = 0.73

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x^3} dx$$

$$= \frac{-((1 + a^4 - b^2(-1 + 2\sqrt{1 - a^2}e^{\operatorname{arctanh}(a)})x^2 - a^2(2 + b^2x^2)) \operatorname{arctanh}(a + bx)^2) + 2bx \operatorname{arctanh}(a + bx)}{x^3}$$

input

```
Integrate[ArcTanh[a + b*x]^2/x^3,x]
```

output

```
(-((1 + a^4 - b^2*(-1 + 2*Sqrt[1 - a^2]*E^ArcTanh[a])*x^2 - a^2*(2 + b^2*x
^2))*ArcTanh[a + b*x]^2) + 2*b*x*ArcTanh[a + b*x]*(-1 + a^2 + a*b*x + I*a*
b*Pi*x - 2*a*b*x*ArcTanh[a] + 2*a*b*x*Log[1 - E^(2*ArcTanh[a] - 2*ArcTanh[
a + b*x]])) + 2*b^2*x^2*((-I)*a*Pi*Log[1 + E^(2*ArcTanh[a + b*x])] + I*a*P
i*Log[1/Sqrt[1 - (a + b*x)^2]] + Log[-(b*x)/Sqrt[1 - (a + b*x)^2]]) - 2*a
*ArcTanh[a]*(Log[1 - E^(2*ArcTanh[a] - 2*ArcTanh[a + b*x])] - Log[(-I)*Sin
h[ArcTanh[a] - ArcTanh[a + b*x]]])) - 2*a*b^2*x^2*PolyLog[2, E^(2*ArcTanh[
a] - 2*ArcTanh[a + b*x])])/(2*(-1 + a^2)^2*x^2)
```

Rubi [A] (verified)

Time = 1.11 (sec) , antiderivative size = 347, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6659, 7292, 6671, 27, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arctanh}(a+bx)^2}{x^3} dx \\
 & \quad \downarrow \text{6659} \\
 & b \int \frac{\operatorname{arctanh}(a+bx)}{x^2(1-(a+bx)^2)} dx - \frac{\operatorname{arctanh}(a+bx)^2}{2x^2} \\
 & \quad \downarrow \text{7292} \\
 & b \int \frac{\operatorname{arctanh}(a+bx)}{x^2(-a^2-2bxa-b^2x^2+1)} dx - \frac{\operatorname{arctanh}(a+bx)^2}{2x^2} \\
 & \quad \downarrow \text{6671} \\
 & \int \frac{\operatorname{arctanh}(a+bx)}{x^2(1-(a+bx)^2)} d(a+bx) - \frac{\operatorname{arctanh}(a+bx)^2}{2x^2} \\
 & \quad \downarrow \text{27} \\
 & b^2 \int \frac{\operatorname{arctanh}(a+bx)}{b^2x^2(1-(a+bx)^2)} d(a+bx) - \frac{\operatorname{arctanh}(a+bx)^2}{2x^2} \\
 & \quad \downarrow \text{7276} \\
 & b^2 \int \left(\frac{2a\operatorname{arctanh}(a+bx)}{(a^2-1)^2 bx} - \frac{\operatorname{arctanh}(a+bx)}{2(a-1)^2(a+bx-1)} + \frac{\operatorname{arctanh}(a+bx)}{2(a+1)^2(a+bx+1)} - \frac{\operatorname{arctanh}(a+bx)}{(a^2-1)b^2x^2} \right) d(a+bx) - \frac{\operatorname{arctanh}(a+bx)^2}{2x^2} \\
 & \quad \downarrow \text{2009} \\
 & b^2 \left(-\frac{\operatorname{arctanh}(a+bx)}{(1-a^2)bx} - \frac{2a\operatorname{arctanh}(a+bx) \log\left(\frac{2}{a+bx+1}\right)}{(1-a^2)^2} + \frac{2a\operatorname{arctanh}(a+bx) \log\left(\frac{2bx}{(1-a)(a+bx+1)}\right)}{(1-a^2)^2} + \frac{a \operatorname{PolyLog}\left(\frac{2}{a+bx+1}\right)}{(1-a^2)^2} - \frac{\operatorname{arctanh}(a+bx)^2}{2x^2} \right)
 \end{aligned}$$

input `Int[ArcTanh[a + b*x]^2/x^3,x]`

output
$$\begin{aligned} & -1/2*\text{ArcTanh}[a + b*x]^2/x^2 + b^2*(-(\text{ArcTanh}[a + b*x]/((1 - a^2)*b*x)) + \text{Log}[-(b*x)]/(1 - a^2)^2 + (\text{ArcTanh}[a + b*x]*\text{Log}[2/(1 - a - b*x)])/(2*(1 - a)^2) - \text{Log}[1 - a - b*x]/(2*(1 - a)^2*(1 + a)) - (\text{ArcTanh}[a + b*x]*\text{Log}[2/(1 + a + b*x)])/(2*(1 + a)^2) - (2*a*\text{ArcTanh}[a + b*x]*\text{Log}[2/(1 + a + b*x)])/(1 - a^2)^2 + (2*a*\text{ArcTanh}[a + b*x]*\text{Log}[(2*b*x)/((1 - a)*(1 + a + b*x))])/(1 - a^2)^2 - \text{Log}[1 + a + b*x]/(2*(1 - a)*(1 + a)^2) + \text{PolyLog}[2, -((1 + a + b*x)/(1 - a - b*x))]/(4*(1 - a)^2) + \text{PolyLog}[2, 1 - 2/(1 + a + b*x)]/(4*(1 + a)^2) + (a*\text{PolyLog}[2, 1 - 2/(1 + a + b*x)])/(1 - a^2)^2 - (a*\text{PolyLog}[2, 1 - (2*b*x)/((1 - a)*(1 + a + b*x))])/(1 - a^2)^2 \end{aligned}$$

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6659 `Int[((a_) + ArcTanh[(c_) + (d_)*(x_)]*(b_.))^ (p_.)*((e_) + (f_.)*(x_))^(m_), x_Symbol] := Simp[(e + f*x)^(m + 1)*((a + b*ArcTanh[c + d*x])^p/(f*(m + 1))), x] - Simp[b*d*(p/(f*(m + 1))) Int[(e + f*x)^(m + 1)*((a + b*ArcTanh[c + d*x])^(p - 1)/(1 - (c + d*x)^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && ILtQ[m, -1]`

rule 6671 `Int[((a_) + ArcTanh[(c_) + (d_)*(x_)]*(b_.))^ (p_.)*((e_) + (f_.)*(x_))^(m_.)*((A_) + (B_.)*(x_) + (C_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(-C/d^2 + (C/d^2)*x^2)^q*(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, p, q}, x] && EqQ[B*(1 - c^2) + 2*A*c*d, 0] && EqQ[2*c*C - B*d, 0]`

rule 7276

```
Int[(u_)/((a_)+(b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpanse[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

rule 7292

```
Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v !=
= u]
```

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 385, normalized size of antiderivative = 1.04

| method | result |
|-------------------|--|
| parts | $-\frac{\operatorname{arctanh}(bx+a)^2}{2x^2} + b^2 \left(\frac{\operatorname{arctanh}(bx+a)}{(a-1)(a+1)bx} + \frac{2 \operatorname{arctanh}(bx+a)a \ln(-bx)}{(a-1)^2(a+1)^2} + \frac{\operatorname{arctanh}(bx+a) \ln(bx+a+1)}{2(a+1)^2} - \frac{\operatorname{arctanh}(bx+a) \ln(bx+a-1)}{2(a+1)^2} \right)$ |
| derivativedivides | $b^2 \left(-\frac{\operatorname{arctanh}(bx+a)^2}{2b^2x^2} + \frac{\operatorname{arctanh}(bx+a)}{(a-1)(a+1)bx} + \frac{2 \operatorname{arctanh}(bx+a)a \ln(-bx)}{(a-1)^2(a+1)^2} + \frac{\operatorname{arctanh}(bx+a) \ln(bx+a+1)}{2(a+1)^2} - \frac{\operatorname{arctanh}(bx+a) \ln(bx+a-1)}{2(a+1)^2} \right)$ |
| default | $b^2 \left(-\frac{\operatorname{arctanh}(bx+a)^2}{2b^2x^2} + \frac{\operatorname{arctanh}(bx+a)}{(a-1)(a+1)bx} + \frac{2 \operatorname{arctanh}(bx+a)a \ln(-bx)}{(a-1)^2(a+1)^2} + \frac{\operatorname{arctanh}(bx+a) \ln(bx+a+1)}{2(a+1)^2} - \frac{\operatorname{arctanh}(bx+a) \ln(bx+a-1)}{2(a+1)^2} \right)$ |

input

```
int(arctanh(b*x+a)^2/x^3,x,method=_RETURNVERBOSE)
```

output

```
-1/2*arctanh(b*x+a)^2/x^2+b^2*(arctanh(b*x+a)/(a-1)/(a+1)/b/x+2*arctanh(b*
x+a)*a/(a-1)^2/(a+1)^2*ln(-b*x)+1/2*arctanh(b*x+a)/(a+1)^2*ln(b*x+a+1)-1/2
*arctanh(b*x+a)/(a-1)^2*ln(b*x+a-1)-1/2/(a-1)^2*(1/4*ln(b*x+a-1)^2-1/2*dil
og(1/2*b*x+1/2*a+1/2)-1/2*ln(b*x+a-1)*ln(1/2*b*x+1/2*a+1/2))+1/2/(a+1)^2*(
-1/4*ln(b*x+a+1)^2+1/2*(ln(b*x+a+1)-ln(1/2*b*x+1/2*a+1/2))*ln(-1/2*b*x-1/2
*a+1/2)-1/2*dilog(1/2*b*x+1/2*a+1/2))-1/(a-1)/(a+1)*(-1/(a-1)/(a+1)*ln(-b*
x)-1/(2*a+2)*ln(b*x+a+1)+1/(2*a-2)*ln(b*x+a-1))+2*a/(a-1)^2/(a+1)^2*(-1/2*
dilog((-b*x-a-1)/(-1-a))-1/2*ln(-b*x)*ln((-b*x-a-1)/(-1-a))+1/2*dilog(1/(1
-a)*(-b*x-a+1))+1/2*ln(-b*x)*ln(1/(1-a)*(-b*x-a+1)))
```

Fricas [F]

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{artanh}(bx + a)^2}{x^3} dx$$

input `integrate(arctanh(b*x+a)^2/x^3,x, algorithm="fricas")`

output `integral(arctanh(b*x + a)^2/x^3, x)`

Sympy [F]

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{atanh}^2(a + bx)}{x^3} dx$$

input `integrate(atanh(b*x+a)**2/x**3,x)`

output `Integral(atanh(a + b*x)**2/x**3, x)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 360, normalized size of antiderivative = 0.97

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(a + bx)^2}{x^3} dx \\ &= \frac{1}{8} \left(\frac{8 \left(\log(bx + a - 1) \log\left(\frac{1}{2}bx + \frac{1}{2}a + \frac{1}{2}\right) + \operatorname{Li}_2\left(-\frac{1}{2}bx - \frac{1}{2}a + \frac{1}{2}\right)\right)a}{a^4 - 2a^2 + 1} - \frac{8 \left(\log\left(\frac{bx}{a+1} + 1\right) \log(x) + \operatorname{Li}_2\left(\frac{bx}{a+1}\right)\right)}{a^4 - 2a^2 + 1} \right. \\ & \quad \left. + \frac{1}{2} \left(\frac{4ab \log(x)}{a^4 - 2a^2 + 1} + \frac{b \log(bx + a + 1)}{a^2 + 2a + 1} - \frac{b \log(bx + a - 1)}{a^2 - 2a + 1} + \frac{2}{(a^2 - 1)x} \right) b \operatorname{artanh}(bx \right. \\ & \quad \left. + a) - \frac{\operatorname{artanh}(bx + a)^2}{2x^2} \right) \end{aligned}$$

input `integrate(arctanh(b*x+a)^2/x^3,x, algorithm="maxima")`

output

```
1/8*(8*(log(b*x + a - 1)*log(1/2*b*x + 1/2*a + 1/2) + dilog(-1/2*b*x - 1/2
*a + 1/2))*a/(a^4 - 2*a^2 + 1) - 8*(log(b*x/(a + 1) + 1)*log(x) + dilog(-b
*x/(a + 1)))*a/(a^4 - 2*a^2 + 1) + 8*(log(b*x/(a - 1) + 1)*log(x) + dilog(
-b*x/(a - 1)))*a/(a^4 - 2*a^2 + 1) - ((a^2 - 2*a + 1)*log(b*x + a + 1)^2 -
2*(a^2 - 2*a + 1)*log(b*x + a + 1)*log(b*x + a - 1) + (a^2 + 2*a + 1)*log
(b*x + a - 1)^2)/(a^4 - 2*a^2 + 1) + 4*log(b*x + a + 1)/(a^3 + a^2 - a - 1
) - 4*log(b*x + a - 1)/(a^3 - a^2 - a + 1) + 8*log(x)/(a^4 - 2*a^2 + 1))*b
^2 + 1/2*(4*a*b*log(x)/(a^4 - 2*a^2 + 1) + b*log(b*x + a + 1)/(a^2 + 2*a +
1) - b*log(b*x + a - 1)/(a^2 - 2*a + 1) + 2/((a^2 - 1)*x))*b*arctanh(b*x
+ a) - 1/2*arctanh(b*x + a)^2/x^2
```

Giac [F]

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{artanh}(bx + a)^2}{x^3} dx$$

input

```
integrate(arctanh(b*x+a)^2/x^3,x, algorithm="giac")
```

output

```
integrate(arctanh(b*x + a)^2/x^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{atanh}(a + bx)^2}{x^3} dx$$

input

```
int(atanh(a + b*x)^2/x^3,x)
```

output

```
int(atanh(a + b*x)^2/x^3, x)
```

Reduce [F]

$$\int \frac{\operatorname{arctanh}(a + bx)^2}{x^3} dx$$

$$= \frac{5 \operatorname{atanh}(bx + a)^2 a^4 - \operatorname{atanh}(bx + a)^2 a^2 - 4 \operatorname{atanh}(bx + a) a^3 + 2 \operatorname{atanh}(bx + a) a - 3 \operatorname{atanh}(bx + a)^2 a^6 -$$

input `int(atanh(b*x+a)^2/x^3,x)`

output

```
( - 3*atanh(a + b*x)**2*a**6 + atanh(a + b*x)**2*a**4*b**2*x**2 + 5*atanh(a + b*x)**2*a**4 - 2*atanh(a + b*x)**2*a**2*b**2*x**2 - atanh(a + b*x)**2*a**2 + atanh(a + b*x)**2*b**2*x**2 - atanh(a + b*x)**2 + 2*atanh(a + b*x)*a**5 - 2*atanh(a + b*x)*a**4*b*x - 4*atanh(a + b*x)*a**3*b**2*x**2 - 4*atanh(a + b*x)*a**3 + 6*atanh(a + b*x)*a**2*b**2*x**2 + 4*atanh(a + b*x)*a**2*b*x + 2*atanh(a + b*x)*a - 2*atanh(a + b*x)*b**2*x**2 - 2*atanh(a + b*x)*b*x + 12*int(atanh(a + b*x)/(3*a**4*x**3 + 6*a**3*b*x**4 + 3*a**2*b**2*x**5 - 2*a**2*x**3 + 2*a*b*x**4 + b**2*x**5 - x**3),x)*a**9*x**2 - 32*int(atanh(a + b*x)/(3*a**4*x**3 + 6*a**3*b*x**4 + 3*a**2*b**2*x**5 - 2*a**2*x**3 + 2*a*b*x**4 + b**2*x**5 - x**3),x)*a**7*x**2 + 24*int(atanh(a + b*x)/(3*a**4*x**3 + 6*a**3*b*x**4 + 3*a**2*b**2*x**5 - 2*a**2*x**3 + 2*a*b*x**4 + b**2*x**5 - x**3),x)*a**5*x**2 - 4*int(atanh(a + b*x)/(3*a**4*x**3 + 6*a**3*b*x**4 + 3*a**2*b**2*x**5 - 2*a**2*x**3 + 2*a*b*x**4 + b**2*x**5 - x**3),x)*a*x**2 + 6*log(a + b*x - 1)*a**2*b**2*x**2 - 2*log(a + b*x - 1)*b**2*x**2 - 6*log(x)*a**2*b**2*x**2 + 2*log(x)*b**2*x**2 - 2*a**3*b*x + 2*a*b*x)/(2*x**2*(3*a**6 - 5*a**4 + a**2 + 1))
```

3.8 $\int \frac{\operatorname{arctanh}(1+bx)^2}{x} dx$

| | |
|---------------------------------------|-----|
| Optimal result | 106 |
| Mathematica [A] (verified) | 106 |
| Rubi [A] (verified) | 107 |
| Maple [C] (warning: unable to verify) | 109 |
| Fricas [F] | 110 |
| Sympy [F] | 110 |
| Maxima [F] | 110 |
| Giac [F] | 111 |
| Mupad [F(-1)] | 111 |
| Reduce [F] | 111 |

Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \frac{\operatorname{arctanh}(1+bx)^2}{x} dx = -\operatorname{arctanh}(1+bx)^2 \log\left(-\frac{2}{bx}\right) - \operatorname{arctanh}(1+bx) \operatorname{PolyLog}\left(2, 1 + \frac{2}{bx}\right) + \frac{1}{2} \operatorname{PolyLog}\left(3, 1 + \frac{2}{bx}\right)$$

output

```
-arctanh(b*x+1)^2*ln(-2/b/x)-arctanh(b*x+1)*polylog(2,1+2/b/x)+1/2*polylog(3,1+2/b/x)
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.34

$$\int \frac{\operatorname{arctanh}(1+bx)^2}{x} dx = -\frac{2}{3} \operatorname{arctanh}(1+bx)^3 - \operatorname{arctanh}(1+bx)^2 \log(1+e^{-2\operatorname{arctanh}(1+bx)}) + \operatorname{arctanh}(1+bx) \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arctanh}(1+bx)}\right) + \frac{1}{2} \operatorname{PolyLog}\left(3, -e^{-2\operatorname{arctanh}(1+bx)}\right)$$

input `Integrate[ArcTanh[1 + b*x]^2/x,x]`

output `(-2*ArcTanh[1 + b*x]^3)/3 - ArcTanh[1 + b*x]^2*Log[1 + E^(-2*ArcTanh[1 + b*x])] + ArcTanh[1 + b*x]*PolyLog[2, -E^(-2*ArcTanh[1 + b*x])] + PolyLog[3, -E^(-2*ArcTanh[1 + b*x])]/2`

Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.09, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6661, 25, 27, 6470, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arctanh}(bx+1)^2}{x} dx \\
 & \quad \downarrow \text{6661} \\
 & \int \frac{\operatorname{arctanh}(bx+1)^2}{x} d(bx+1) \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{\operatorname{arctanh}(bx+1)^2}{x} d(bx+1) \\
 & \quad \downarrow \text{27} \\
 & - \int - \frac{\operatorname{arctanh}(bx+1)^2}{bx} d(bx+1) \\
 & \quad \downarrow \text{6470} \\
 & 2 \int \frac{\operatorname{arctanh}(bx+1) \log\left(-\frac{2}{bx}\right)}{1-(bx+1)^2} d(bx+1) - \operatorname{arctanh}(bx+1)^2 \log\left(-\frac{2}{bx}\right) \\
 & \quad \downarrow \text{6620}
 \end{aligned}$$

$$2 \left(\frac{1}{2} \int \frac{\text{PolyLog} \left(2, 1 + \frac{2}{bx} \right) d(bx + 1)}{1 - (bx + 1)^2} - \frac{1}{2} \text{arctanh}(bx + 1) \text{PolyLog} \left(2, 1 + \frac{2}{bx} \right) \right) - \text{arctanh}(bx + 1)^2 \log \left(-\frac{2}{bx} \right)$$

↓ 7164

$$2 \left(\frac{1}{4} \text{PolyLog} \left(3, 1 + \frac{2}{bx} \right) - \frac{1}{2} \text{arctanh}(bx + 1) \text{PolyLog} \left(2, 1 + \frac{2}{bx} \right) \right) - \text{arctanh}(bx + 1)^2 \log \left(-\frac{2}{bx} \right)$$

input `Int[ArcTanh[1 + b*x]^2/x, x]`

output `-(ArcTanh[1 + b*x]^2*Log[-2/(b*x)]) + 2*(-1/2*(ArcTanh[1 + b*x]*PolyLog[2, 1 + 2/(b*x)]) + PolyLog[3, 1 + 2/(b*x)]/4)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 6470 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^((p_.))/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6620 `Int[(Log[u_]*)((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^((p_.))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(-(a + b*ArcTanh[c*x])^p)*(PolyLog[2, 1 - u]/(2*c*d)), x] + Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]`

rule 6661

```
Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*
ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IG
tQ[p, 0]
```

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 9.16 (sec) , antiderivative size = 158, normalized size of antiderivative = 2.82

| method | result |
|-------------------|--|
| derivativedivides | $\ln(bx) \operatorname{arctanh}(bx+1)^2 - \operatorname{arctanh}(bx+1) \operatorname{polylog}\left(2, -\frac{(bx+2)^2}{-(bx+1)^2+1}\right) + \frac{\operatorname{polylog}\left(3, -\frac{(bx+2)^2}{-(bx+1)^2+1}\right)}{2}$ |
| default | $\ln(bx) \operatorname{arctanh}(bx+1)^2 - \operatorname{arctanh}(bx+1) \operatorname{polylog}\left(2, -\frac{(bx+2)^2}{-(bx+1)^2+1}\right) + \frac{\operatorname{polylog}\left(3, -\frac{(bx+2)^2}{-(bx+1)^2+1}\right)}{2}$ |
| parts | $\ln(x) \operatorname{arctanh}(bx+1)^2 - 2b \left(\frac{\operatorname{arctanh}(bx+1) \operatorname{polylog}\left(2, -\frac{(bx+2)^2}{-(bx+1)^2+1}\right)}{2b} - \frac{\operatorname{polylog}\left(3, -\frac{(bx+2)^2}{-(bx+1)^2+1}\right)}{4b} \right)$ |
| risch | Expression too large to display |

input

```
int(arctanh(b*x+1)^2/x,x,method=_RETURNVERBOSE)
```

output

```
ln(b*x)*arctanh(b*x+1)^2-arctanh(b*x+1)*polylog(2,-(b*x+2)^2/(-(b*x+1)^2+1
))+1/2*polylog(3,-(b*x+2)^2/(-(b*x+1)^2+1))-(-I*Pi*csgn(I/(1-(b*x+2)^2/((b
*x+1)^2-1)))^2+I*Pi*csgn(I/(1-(b*x+2)^2/((b*x+1)^2-1)))^3+I*Pi+ln(2))*arct
anh(b*x+1)^2
```

Fricas [F]

$$\int \frac{\operatorname{arctanh}(1 + bx)^2}{x} dx = \int \frac{\operatorname{artanh}(bx + 1)^2}{x} dx$$

input `integrate(arctanh(b*x+1)^2/x,x, algorithm="fricas")`

output `integral(arctanh(b*x + 1)^2/x, x)`

Sympy [F]

$$\int \frac{\operatorname{arctanh}(1 + bx)^2}{x} dx = \int \frac{\operatorname{atanh}^2(bx + 1)}{x} dx$$

input `integrate(atanh(b*x+1)**2/x,x)`

output `Integral(atanh(b*x + 1)**2/x, x)`

Maxima [F]

$$\int \frac{\operatorname{arctanh}(1 + bx)^2}{x} dx = \int \frac{\operatorname{artanh}(bx + 1)^2}{x} dx$$

input `integrate(arctanh(b*x+1)^2/x,x, algorithm="maxima")`

output `1/12*log(-b*x)^3 + 1/4*log(b*x + 2)^2*log(-x) - 1/4*integrate(2*(b*x*log(b) + 2*(b*x + 1)*log(-x) + 2*log(b))*log(b*x + 2)/(b*x^2 + 2*x), x)`

Giac [F]

$$\int \frac{\operatorname{arctanh}(1 + bx)^2}{x} dx = \int \frac{\operatorname{artanh}(bx + 1)^2}{x} dx$$

input `integrate(arctanh(b*x+1)^2/x,x, algorithm="giac")`

output `integrate(arctanh(b*x + 1)^2/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(1 + bx)^2}{x} dx = \int \frac{\operatorname{atanh}(bx + 1)^2}{x} dx$$

input `int(atanh(b*x + 1)^2/x,x)`

output `int(atanh(b*x + 1)^2/x, x)`

Reduce [F]

$$\int \frac{\operatorname{arctanh}(1 + bx)^2}{x} dx = \int \frac{\operatorname{atanh}(bx + 1)^2}{x} dx$$

input `int(atanh(b*x+1)^2/x,x)`

output `int(atanh(b*x + 1)**2/x,x)`

3.9 $\int (ce + dex)^3(a + b\operatorname{arctanh}(c + dx)) dx$

| | |
|---|-----|
| Optimal result | 112 |
| Mathematica [A] (verified) | 112 |
| Rubi [A] (verified) | 113 |
| Maple [A] (verified) | 115 |
| Fricas [B] (verification not implemented) | 115 |
| Sympy [B] (verification not implemented) | 116 |
| Maxima [B] (verification not implemented) | 117 |
| Giac [B] (verification not implemented) | 117 |
| Mupad [B] (verification not implemented) | 119 |
| Reduce [B] (verification not implemented) | 120 |

Optimal result

Integrand size = 21, antiderivative size = 72

$$\int (ce + dex)^3(a + b\operatorname{arctanh}(c + dx)) dx = \frac{1}{4}be^3x + \frac{be^3(c + dx)^3}{12d} - \frac{be^3\operatorname{arctanh}(c + dx)}{4d} + \frac{e^3(c + dx)^4(a + b\operatorname{arctanh}(c + dx))}{4d}$$

output $\frac{1}{4}b^3e^{3x} + \frac{1}{12}b^3e^{3(d*x+c)^3/d} - \frac{1}{4}b^3e^{3\operatorname{arctanh}(d*x+c)}/d + \frac{1}{4}e^{3(d*x+c)^4(a+b\operatorname{arctanh}(d*x+c))}/d$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.08

$$\int (ce + dex)^3(a + b\operatorname{arctanh}(c + dx)) dx = \frac{e^3(6b(c + dx) + 2b(c + dx)^3 + 6a(c + dx)^4 + 6b(c + dx)^4\operatorname{arctanh}(c + dx) + 3b\log(1 - c - dx) - 3b\log(1 + c + dx))}{24d}$$

input `Integrate[(c*e + d*e*x)^3*(a + b*ArcTanh[c + d*x]),x]`

output

$$(e^{3*(6*b*(c + d*x) + 2*b*(c + d*x)^3 + 6*a*(c + d*x)^4 + 6*b*(c + d*x)^4*ArcTanh[c + d*x] + 3*b*Log[1 - c - d*x] - 3*b*Log[1 + c + d*x])})/(24*d)$$
Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.82, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {6657, 27, 6452, 254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ce + dex)^3(a + b \operatorname{arctanh}(c + dx)) dx$$

$$\downarrow 6657$$

$$\frac{\int e^3(c + dx)^3(a + b \operatorname{arctanh}(c + dx))d(c + dx)}{d}$$

$$\downarrow 27$$

$$\frac{e^3 \int (c + dx)^3(a + b \operatorname{arctanh}(c + dx))d(c + dx)}{d}$$

$$\downarrow 6452$$

$$\frac{e^3 \left(\frac{1}{4}(c + dx)^4(a + b \operatorname{arctanh}(c + dx)) - \frac{1}{4}b \int \frac{(c+dx)^4}{1-(c+dx)^2} d(c + dx) \right)}{d}$$

$$\downarrow 254$$

$$\frac{e^3 \left(\frac{1}{4}(c + dx)^4(a + b \operatorname{arctanh}(c + dx)) - \frac{1}{4}b \int \left(-(c + dx)^2 + \frac{1}{1-(c+dx)^2} - 1 \right) d(c + dx) \right)}{d}$$

$$\downarrow 2009$$

$$\frac{e^3 \left(\frac{1}{4}(c + dx)^4(a + b \operatorname{arctanh}(c + dx)) - \frac{1}{4}b(\operatorname{arctanh}(c + dx) - \frac{1}{3}(c + dx)^3 - c - dx) \right)}{d}$$

input

$$\text{Int}[(c*e + d*e*x)^3*(a + b*ArcTanh[c + d*x]), x]$$

output $(e^{3*(-1/4*(b*(-c - d*x - (c + d*x)^{3/3} + \text{ArcTanh}[c + d*x])) + ((c + d*x)^{4*(a + b*\text{ArcTanh}[c + d*x]))/4))/d$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_)] /; \text{FreeQ}[b, x]$

rule 254 $\text{Int}[(x_)^{(m)}/((a_) + (b_)*(x_)^2), x_Symbol] \rightarrow \text{Int}[\text{PolynomialDivide}[x^m, a + b*x^2, x], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{IGtQ}[m, 3]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 6452 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)})/(1 - c^2*x^{(2*n)}), x], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 6657 $\text{Int}[(a_ + \text{ArcTanh}[(c_ + (d_)*(x_)]*(b_))^{(p_)}*((e_ + (f_)*(x_))^{(m_)}, x_Symbol] \rightarrow \text{Simp}[1/d \ \text{Subst}[\text{Int}[(f*(x/d))^{m*(a + b*\text{ArcTanh}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{EqQ}[d*e - c*f, 0] \ \&\& \ \text{IGtQ}[p, 0]$

Maple [A] (verified)

Time = 0.83 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.03

| method | result |
|-------------------|---|
| derivativedivides | $\frac{e^3 a (dx+c)^4 + b e^3 \left(\frac{(dx+c)^4 \operatorname{arctanh}(dx+c)}{4} + \frac{(dx+c)^3}{12} + \frac{dx}{4} + \frac{c}{4} + \frac{\ln(dx+c-1)}{8} - \frac{\ln(dx+c+1)}{8} \right)}{d}$ |
| default | $\frac{e^3 a (dx+c)^4 + b e^3 \left(\frac{(dx+c)^4 \operatorname{arctanh}(dx+c)}{4} + \frac{(dx+c)^3}{12} + \frac{dx}{4} + \frac{c}{4} + \frac{\ln(dx+c-1)}{8} - \frac{\ln(dx+c+1)}{8} \right)}{d}$ |
| parts | $\frac{e^3 a (dx+c)^4}{4d} + \frac{b e^3 \left(\frac{(dx+c)^4 \operatorname{arctanh}(dx+c)}{4} + \frac{(dx+c)^3}{12} + \frac{dx}{4} + \frac{c}{4} + \frac{\ln(dx+c-1)}{8} - \frac{\ln(dx+c+1)}{8} \right)}{d}$ |
| orering | $\frac{(2x^5 d^5 + 10x^4 d^4 c + 20x^3 d^3 c^2 + 19x^2 d^2 c^3 + 8c^4 dx + 2d^3 x^3 + c^5 + 3c d^2 x^2 - 4dx - c)(dex+ce)^3 (a+b \operatorname{arctanh}(dx+c))}{4d(dx+c)^4} - \dots$ |
| parallelrisc | $- \frac{3b d^5 e^3 \operatorname{arctanh}(dx+c) x^4 - 3x^4 a d^5 e^3 - 12bc d^4 e^3 \operatorname{arctanh}(dx+c) x^3 - 12x^3 ac d^4 e^3 - 18x^2 \operatorname{arctanh}(dx+c) b c^2 d^3 e^3 - \dots}{4d(dx+c)^4}$ |
| risc | $\frac{e^3 (dx+c)^4 b \ln(dx+c+1)}{8d} - \frac{e^3 d^3 b x^4 \ln(-dx-c+1)}{8} - \frac{e^3 d^2 bc x^3 \ln(-dx-c+1)}{2} + \frac{e^3 d^3 a x^4}{4} - \frac{3e^3 db c^2 x^2 \ln(-dx-c+1)}{4}$ |

input `int((d*e*x+c*e)^3*(a+b*arctanh(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*(1/4*e^3*a*(d*x+c)^4+b*e^3*(1/4*(d*x+c)^4*arctanh(d*x+c)+1/12*(d*x+c)^3+1/4*d*x+1/4*c+1/8*ln(d*x+c-1)-1/8*ln(d*x+c+1)))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 159 vs. 2(64) = 128.

Time = 0.08 (sec) , antiderivative size = 159, normalized size of antiderivative = 2.21

$$\int (ce + dex)^3 (a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \frac{6ad^4e^3x^4 + 2(12ac + b)d^3e^3x^3 + 6(6ac^2 + bc)d^2e^3x^2 + 6(4ac^3 + bc^2 + b)de^3x + 3(bd^4e^3x^4 + 4bcd^3e^3x^3 + 3c^2d^2e^3x^2 + 3cd^2e^3x + 3c^3e^3)}{24d}$$

input `integrate((d*e*x+c*e)^3*(a+b*arctanh(d*x+c)),x, algorithm="fricas")`

output

```
1/24*(6*a*d^4*e^3*x^4 + 2*(12*a*c + b)*d^3*e^3*x^3 + 6*(6*a*c^2 + b*c)*d^2
*e^3*x^2 + 6*(4*a*c^3 + b*c^2 + b)*d*e^3*x + 3*(b*d^4*e^3*x^4 + 4*b*c*d^3*
e^3*x^3 + 6*b*c^2*d^2*e^3*x^2 + 4*b*c^3*d*e^3*x + (b*c^4 - b)*e^3)*log(-(d
*x + c + 1)/(d*x + c - 1))/d
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 231 vs. $2(61) = 122$.

Time = 1.47 (sec) , antiderivative size = 231, normalized size of antiderivative = 3.21

$$\int (ce + dex)^3 (a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \begin{cases} ac^3 e^3 x + \frac{3ac^2 de^3 x^2}{2} + acd^2 e^3 x^3 + \frac{ad^3 e^3 x^4}{4} + \frac{bc^4 e^3 \operatorname{atanh}(c+dx)}{4d} + bc^3 e^3 x \operatorname{atanh}(c + dx) + \frac{3bc^2 de^3 x^2 \operatorname{atanh}(c+dx)}{2} \\ c^3 e^3 x (a + b \operatorname{atanh}(c)) \end{cases}$$

input

```
integrate((d*e*x+c*e)**3*(a+b*atanh(d*x+c)),x)
```

output

```
Piecewise((a*c**3*e**3*x + 3*a*c**2*d*e**3*x**2/2 + a*c*d**2*e**3*x**3 + a
*d**3*e**3*x**4/4 + b*c**4*e**3*atanh(c + d*x)/(4*d) + b*c**3*e**3*x*atanh
(c + d*x) + 3*b*c**2*d*e**3*x**2*atanh(c + d*x)/2 + b*c**2*e**3*x/4 + b*c*
d**2*e**3*x**3*atanh(c + d*x) + b*c*d*e**3*x**2/4 + b*d**3*e**3*x**4*atanh
(c + d*x)/4 + b*d**2*e**3*x**3/12 + b*e**3*x/4 - b*e**3*atanh(c + d*x)/(4*
d), Ne(d, 0)), (c**3*e**3*x*(a + b*atanh(c)), True))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 357 vs. $2(64) = 128$.

Time = 0.04 (sec) , antiderivative size = 357, normalized size of antiderivative = 4.96

$$\int (ce + dex)^3(a + \operatorname{barctanh}(c + dx)) dx = \frac{1}{4} ad^3 e^3 x^4 + acd^2 e^3 x^3 + \frac{3}{2} ac^2 de^3 x^2 + \frac{3}{4} \left(2x^2 \operatorname{artanh}(dx + c) + d \left(\frac{2x}{d^2} - \frac{(c^2 + 2c + 1) \log(dx + c + 1)}{d^3} + \frac{(c^2 - 2c + 1) \log(dx + c - 1)}{d^3} \right) \right) + \frac{1}{2} \left(2x^3 \operatorname{artanh}(dx + c) + d \left(\frac{dx^2 - 4cx}{d^3} + \frac{(c^3 + 3c^2 + 3c + 1) \log(dx + c + 1)}{d^4} - \frac{(c^3 - 3c^2 + 3c - 1) \log(dx + c - 1)}{d^4} \right) \right) + \frac{1}{24} \left(6x^4 \operatorname{artanh}(dx + c) + d \left(\frac{2(d^2 x^3 - 3cdx^2 + 3(3c^2 + 1)x)}{d^4} - \frac{3(c^4 + 4c^3 + 6c^2 + 4c + 1) \log(dx + c + 1)}{d^5} + \frac{3(c^4 - 4c^3 + 6c^2 + 4c + 1) \log(dx + c - 1)}{d^5} \right) \right) + ac^3 e^3 x + \frac{(2(dx + c) \operatorname{artanh}(dx + c) + \log(-(dx + c)^2 + 1)) bc^3 e^3}{2d}$$

input `integrate((d*e*x+c*e)^3*(a+b*arctanh(d*x+c)),x, algorithm="maxima")`

output `1/4*a*d^3*e^3*x^4 + a*c*d^2*e^3*x^3 + 3/2*a*c^2*d*e^3*x^2 + 3/4*(2*x^2*arctanh(d*x + c) + d*(2*x/d^2 - (c^2 + 2*c + 1)*log(d*x + c + 1)/d^3 + (c^2 - 2*c + 1)*log(d*x + c - 1)/d^3))*b*c^2*d*e^3 + 1/2*(2*x^3*arctanh(d*x + c) + d*((d*x^2 - 4*c*x)/d^3 + (c^3 + 3*c^2 + 3*c + 1)*log(d*x + c + 1)/d^4 - (c^3 - 3*c^2 + 3*c - 1)*log(d*x + c - 1)/d^4))*b*c*d^2*e^3 + 1/24*(6*x^4*arctanh(d*x + c) + d*(2*(d^2*x^3 - 3*c*d*x^2 + 3*(3*c^2 + 1)*x)/d^4 - 3*(c^4 + 4*c^3 + 6*c^2 + 4*c + 1)*log(d*x + c + 1)/d^5 + 3*(c^4 - 4*c^3 + 6*c^2 - 4*c + 1)*log(d*x + c - 1)/d^5))*b*d^3*e^3 + a*c^3*e^3*x + 1/2*(2*(d*x + c)*arctanh(d*x + c) + log(-(d*x + c)^2 + 1))*b*c^3*e^3/d`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 363 vs. $2(64) = 128$.

Time = 0.14 (sec) , antiderivative size = 363, normalized size of antiderivative = 5.04

$$\int (ce + dex)^3(a + \operatorname{barctanh}(c + dx)) dx = \frac{1}{6} ((c + 1)d - (c - 1)d) \left(\frac{3 \left(\frac{(dx+c+1)^3 be^3}{(dx+c-1)^3} + \frac{(dx+c+1) be^3}{dx+c-1} \right) \log\left(-\frac{dx+c+1}{dx+c-1}\right)}{\frac{(dx+c+1)^4 d^2}{(dx+c-1)^4} - \frac{4(dx+c+1)^3 d^2}{(dx+c-1)^3} + \frac{6(dx+c+1)^2 d^2}{(dx+c-1)^2} - \frac{4(dx+c+1) d^2}{dx+c-1} + d^2} + \frac{6(dx+c+1)^3 a e^3 + 6(dx+c+1)^2 a e^3}{(dx+c-1)^3} + \frac{6(dx+c+1) a e^3}{(dx+c-1)^2} + \frac{6 a e^3}{dx+c-1} \right)$$

input `integrate((d*e*x+c*e)^3*(a+b*arctanh(d*x+c)),x, algorithm="giac")`

output
$$\frac{1}{6}((c + 1)d - (c - 1)d) \cdot (3 \cdot ((d*x + c + 1)^3 * b * e^3 / (d*x + c - 1)^3 + (d*x + c + 1) * b * e^3 / (d*x + c - 1)) * \log(-(d*x + c + 1) / (d*x + c - 1)) / ((d*x + c + 1)^4 * d^2 / (d*x + c - 1)^4 - 4 * (d*x + c + 1)^3 * d^2 / (d*x + c - 1)^3 + 6 * (d*x + c + 1)^2 * d^2 / (d*x + c - 1)^2 - 4 * (d*x + c + 1) * d^2 / (d*x + c - 1) + d^2) + (6 * (d*x + c + 1)^3 * a * e^3 / (d*x + c - 1)^3 + 6 * (d*x + c + 1) * a * e^3 / (d*x + c - 1) + 3 * (d*x + c + 1)^3 * b * e^3 / (d*x + c - 1)^3 - 6 * (d*x + c + 1)^2 * b * e^3 / (d*x + c - 1)^2 + 5 * (d*x + c + 1) * b * e^3 / (d*x + c - 1) - 2 * b * e^3) / ((d*x + c + 1)^4 * d^2 / (d*x + c - 1)^4 - 4 * (d*x + c + 1)^3 * d^2 / (d*x + c - 1)^3 + 6 * (d*x + c + 1)^2 * d^2 / (d*x + c - 1)^2 - 4 * (d*x + c + 1) * d^2 / (d*x + c - 1) + d^2))$$

Mupad [B] (verification not implemented)

Time = 4.63 (sec) , antiderivative size = 414, normalized size of antiderivative = 5.75

$$\begin{aligned}
& \int (ce + dex)^3 (a + b \operatorname{arctanh}(c + dx)) dx \\
&= x^3 \left(\frac{d^2 e^3 (b + 20ac)}{12} - \frac{2acd^2 e^3}{3} \right) \\
&+ \ln(c + dx + 1) \left(\frac{bc^3 e^3 x}{2} + \frac{3bc^2 d e^3 x^2}{4} + \frac{bcd^2 e^3 x^3}{2} + \frac{bd^3 e^3 x^4}{8} \right) \\
&- \ln(1 - dx - c) \left(\frac{bc^3 e^3 x}{2} + \frac{3bc^2 d e^3 x^2}{4} + \frac{bcd^2 e^3 x^3}{2} + \frac{bd^3 e^3 x^4}{8} \right) \\
&- x^2 \left(\frac{c \left(\frac{d^2 e^3 (b+20ac)}{4} - 2acd^2 e^3 \right)}{d} - \frac{de^3 (10ac^2 + bc - a)}{2} + \frac{ade^3 (4c^2 - 4)}{8} \right) \\
&+ x \left(\frac{ce^3 (20ac^2 + 3bc - 6a)}{2} - \frac{(4c^2 - 4) \left(\frac{d^2 e^3 (b+20ac)}{4} - 2acd^2 e^3 \right)}{4d^2} \right. \\
&\quad \left. + \frac{2c \left(\frac{d^2 e^3 (b+20ac)}{4} - 2acd^2 e^3 \right) - de^3 (10ac^2 + bc - a) + \frac{ade^3 (4c^2 - 4)}{4}}{d} \right) \\
&+ \frac{\ln(c + dx - 1) (be^3 - bc^4 e^3)}{8d} + \frac{ad^3 e^3 x^4}{4} \\
&+ \frac{be^3 \ln(c + dx + 1) (c^2 + 1) (c - 1) (c + 1)}{8d}
\end{aligned}$$

input

```
int((c*e + d*e*x)^3*(a + b*atanh(c + d*x)),x)
```


output

```
x^3*((d^2*e^3*(b + 20*a*c))/12 - (2*a*c*d^2*e^3)/3) + log(c + d*x + 1)*((b
*d^3*e^3*x^4)/8 + (b*c^3*e^3*x)/2 + (3*b*c^2*d*e^3*x^2)/4 + (b*c*d^2*e^3*x
^3)/2) - log(1 - d*x - c)*((b*d^3*e^3*x^4)/8 + (b*c^3*e^3*x)/2 + (3*b*c^2*
d*e^3*x^2)/4 + (b*c*d^2*e^3*x^3)/2) - x^2*((c*((d^2*e^3*(b + 20*a*c))/4 -
2*a*c*d^2*e^3))/d - (d*e^3*(b*c - a + 10*a*c^2))/2 + (a*d*e^3*(4*c^2 - 4))
/8) + x*((c*e^3*(3*b*c - 6*a + 20*a*c^2))/2 - ((4*c^2 - 4)*((d^2*e^3*(b +
20*a*c))/4 - 2*a*c*d^2*e^3))/(4*d^2) + (2*c*((2*c*((d^2*e^3*(b + 20*a*c))/
4 - 2*a*c*d^2*e^3))/d - d*e^3*(b*c - a + 10*a*c^2) + (a*d*e^3*(4*c^2 - 4))
/4))/d) + (log(c + d*x - 1)*(b*e^3 - b*c^4*e^3))/(8*d) + (a*d^3*e^3*x^4)/4
+ (b*e^3*log(c + d*x + 1)*(c^2 + 1)*(c - 1)*(c + 1))/(8*d)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 163, normalized size of antiderivative = 2.26

$$\int (ce + dex)^3 (a + \operatorname{barctanh}(c + dx)) dx$$

$$= \frac{e^3 (3 \operatorname{atanh}(dx + c) b c^4 + 12 \operatorname{atanh}(dx + c) b c^3 dx + 18 \operatorname{atanh}(dx + c) b c^2 d^2 x^2 + 12 \operatorname{atanh}(dx + c) b c d^3 x^3 -$$

input

```
int((d*e*x+c*e)^3*(a+b*atanh(d*x+c)),x)
```

output

```
(e**3*(3*atanh(c + d*x)*b*c**4 + 12*atanh(c + d*x)*b*c**3*d*x + 18*atanh(c
+ d*x)*b*c**2*d**2*x**2 + 12*atanh(c + d*x)*b*c*d**3*x**3 + 3*atanh(c + d
*x)*b*d**4*x**4 - 3*atanh(c + d*x)*b + 12*a*c**3*d*x + 18*a*c**2*d**2*x**2
+ 12*a*c*d**3*x**3 + 3*a*d**4*x**4 + 3*b*c**2*d*x + 3*b*c*d**2*x**2 + b*d
**3*x**3 + 3*b*d*x))/(12*d)
```

3.10 $\int (ce + dex)^2(a + \operatorname{barctanh}(c + dx)) dx$

| | |
|---|-----|
| Optimal result | 121 |
| Mathematica [A] (verified) | 121 |
| Rubi [A] (warning: unable to verify) | 122 |
| Maple [A] (verified) | 124 |
| Fricas [B] (verification not implemented) | 124 |
| Sympy [B] (verification not implemented) | 125 |
| Maxima [B] (verification not implemented) | 125 |
| Giac [B] (verification not implemented) | 126 |
| Mupad [B] (verification not implemented) | 127 |
| Reduce [B] (verification not implemented) | 127 |

Optimal result

Integrand size = 21, antiderivative size = 69

$$\int (ce + dex)^2(a + \operatorname{barctanh}(c + dx)) dx = \frac{be^2(c + dx)^2}{6d} + \frac{e^2(c + dx)^3(a + \operatorname{barctanh}(c + dx))}{3d} + \frac{be^2 \log(1 - (c + dx)^2)}{6d}$$

output

```
1/6*b*e^2*(d*x+c)^2/d+1/3*e^2*(d*x+c)^3*(a+b*arctanh(d*x+c))/d+1/6*b*e^2*ln(1-(d*x+c)^2)/d
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.86

$$\int (ce + dex)^2(a + \operatorname{barctanh}(c + dx)) dx = \frac{e^2((c + dx)^2(b + 2a(c + dx)) + 2b(c + dx)^3 \operatorname{arctanh}(c + dx) + b \log(1 - (c + dx)^2))}{6d}$$

input

```
Integrate[(c*e + d*e*x)^2*(a + b*ArcTanh[c + d*x]),x]
```

output

$$\frac{(e^{2((c+dx)^2(b+2a(c+dx)) + 2b(c+dx)^3 \operatorname{ArcTanh}[c+dx] + b \operatorname{Log}[1-(c+dx)^2])})}{(6*d)}$$

Rubi [A] (warning: unable to verify)

Time = 0.30 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.78, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6657, 27, 6452, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (ce + dex)^2(a + \operatorname{barctanh}(c + dx)) dx \\ & \quad \downarrow \text{6657} \\ & \frac{\int e^2(c + dx)^2(a + \operatorname{barctanh}(c + dx))d(c + dx)}{d} \\ & \quad \downarrow \text{27} \\ & \frac{e^2 \int (c + dx)^2(a + \operatorname{barctanh}(c + dx))d(c + dx)}{d} \\ & \quad \downarrow \text{6452} \\ & \frac{e^2 \left(\frac{1}{3}(c + dx)^3(a + \operatorname{barctanh}(c + dx)) - \frac{1}{3}b \int \frac{(c+dx)^3}{1-(c+dx)^2} d(c + dx) \right)}{d} \\ & \quad \downarrow \text{243} \\ & \frac{e^2 \left(\frac{1}{3}(c + dx)^3(a + \operatorname{barctanh}(c + dx)) - \frac{1}{6}b \int \frac{(c+dx)^2}{-c-dx+1} d(c + dx)^2 \right)}{d} \\ & \quad \downarrow \text{49} \\ & \frac{e^2 \left(\frac{1}{3}(c + dx)^3(a + \operatorname{barctanh}(c + dx)) - \frac{1}{6}b \int \left(\frac{1}{-c-dx+1} - 1 \right) d(c + dx)^2 \right)}{d} \\ & \quad \downarrow \text{2009} \\ & \frac{e^2 \left(\frac{1}{3}(c + dx)^3(a + \operatorname{barctanh}(c + dx)) - \frac{1}{6}b(-\log(-c - dx + 1) - c - dx) \right)}{d} \end{aligned}$$

input `Int[(c*e + d*e*x)^2*(a + b*ArcTanh[c + d*x]),x]`

output `(e^2*(((c + d*x)^3*(a + b*ArcTanh[c + d*x]))/3 - (b*(-c - d*x - Log[1 - c - d*x]))/6))/d`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6657 `Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.97

| method | result |
|-------------------|--|
| derivativedivides | $\frac{e^2 a (dx+c)^3 + b e^2 \left(\frac{(dx+c)^3 \operatorname{arctanh}(dx+c)}{3} + \frac{(dx+c)^2}{6} + \frac{\ln(dx+c-1)}{6} + \frac{\ln(dx+c+1)}{6} \right)}{d}$ |
| default | $\frac{e^2 a (dx+c)^3 + b e^2 \left(\frac{(dx+c)^3 \operatorname{arctanh}(dx+c)}{3} + \frac{(dx+c)^2}{6} + \frac{\ln(dx+c-1)}{6} + \frac{\ln(dx+c+1)}{6} \right)}{d}$ |
| parts | $\frac{e^2 a (dx+c)^3}{3d} + \frac{b e^2 \left(\frac{(dx+c)^3 \operatorname{arctanh}(dx+c)}{3} + \frac{(dx+c)^2}{6} + \frac{\ln(dx+c-1)}{6} + \frac{\ln(dx+c+1)}{6} \right)}{d}$ |
| risch | $\frac{e^2 (dx+c)^3 b \ln(dx+c+1)}{6d} - \frac{e^2 d^2 b x^3 \ln(-dx-c+1)}{6} - \frac{e^2 d b c x^2 \ln(-dx-c+1)}{2} + \frac{e^2 d^2 a x^3}{3} - \frac{e^2 b c^2 x \ln(-dx-c+1)}{2}$ |
| parallelrisch | $\frac{2b d^4 e^2 \operatorname{arctanh}(dx+c) x^3 + 2x^3 a d^4 e^2 + 6x^2 \operatorname{arctanh}(dx+c) b c d^3 e^2 + 6x^2 a c d^3 e^2 + 6x \operatorname{arctanh}(dx+c) b c^2 d^2 e^2 + x^2 b d^3 e^2}{6d}$ |

input `int((d*e*x+c*e)^2*(a+b*arctanh(d*x+c)),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{d} \left(\frac{1}{3} e^{2a} (d*x+c)^3 + b e^{2 \left(\frac{1}{3} (d*x+c)^3 \operatorname{arctanh}(d*x+c) + \frac{1}{6} (d*x+c)^2 + \frac{1}{6} \ln(d*x+c-1) + \frac{1}{6} \ln(d*x+c+1) \right)} \right)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 144 vs. 2(63) = 126.

Time = 0.10 (sec) , antiderivative size = 144, normalized size of antiderivative = 2.09

$$\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \frac{2ad^3e^2x^3 + (6ac + b)d^2e^2x^2 + 2(3ac^2 + bc)de^2x + (bc^3 + b)e^2 \log(dx + c + 1) - (bc^3 - b)e^2 \log(dx + c - 1)}{6d}$$

input `integrate((d*e*x+c*e)^2*(a+b*arctanh(d*x+c)),x, algorithm="fricas")`

output
$$\frac{1}{6} \left(2a*d^3*e^2*x^3 + (6*a*c + b)*d^2*e^2*x^2 + 2*(3*a*c^2 + b*c)*d*e^2*x + (b*c^3 + b)*e^2*\log(d*x + c + 1) - (b*c^3 - b)*e^2*\log(d*x + c - 1) + (b*d^3*e^2*x^3 + 3*b*c*d^2*e^2*x^2 + 3*b*c^2*d*e^2*x)*\log(-(d*x + c + 1)/(d*x + c - 1)) \right) / d$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 180 vs. $2(56) = 112$.

Time = 1.22 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.61

$$\int (ce + dex)^2(a + \operatorname{barctanh}(c + dx)) dx$$

$$= \begin{cases} ac^2e^2x + acde^2x^2 + \frac{ad^2e^2x^3}{3} + \frac{bc^3e^2 \operatorname{atanh}(c+dx)}{3d} + bc^2e^2x \operatorname{atanh}(c + dx) + bcde^2x^2 \operatorname{atanh}(c + dx) + \frac{bce^2x}{3} \\ c^2e^2x(a + b \operatorname{atanh}(c)) \end{cases}$$

input `integrate((d*e*x+c*e)**2*(a+b*atanh(d*x+c)),x)`

output `Piecewise((a*c**2*e**2*x + a*c*d*e**2*x**2 + a*d**2*e**2*x**3/3 + b*c**3*e**2*atanh(c + d*x)/(3*d) + b*c**2*e**2*x*atanh(c + d*x) + b*c*d*e**2*x**2*atanh(c + d*x) + b*c*e**2*x/3 + b*d**2*e**2*x**3*atanh(c + d*x)/3 + b*d*e**2*x**2/6 + b*e**2*log(c/d + x + 1/d)/(3*d) - b*e**2*atanh(c + d*x)/(3*d), Ne(d, 0)), (c**2*e**2*x*(a + b*atanh(c)), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 225 vs. $2(63) = 126$.

Time = 0.03 (sec) , antiderivative size = 225, normalized size of antiderivative = 3.26

$$\int (ce + dex)^2(a + \operatorname{barctanh}(c + dx)) dx = \frac{1}{3} ad^2e^2x^3 + acde^2x^2$$

$$+ \frac{1}{2} \left(2x^2 \operatorname{artanh}(dx + c) + d \left(\frac{2x}{d^2} - \frac{(c^2 + 2c + 1) \log(dx + c + 1)}{d^3} + \frac{(c^2 - 2c + 1) \log(dx + c - 1)}{d^3} \right) \right)$$

$$+ \frac{1}{6} \left(2x^3 \operatorname{artanh}(dx + c) + d \left(\frac{dx^2 - 4cx}{d^3} + \frac{(c^3 + 3c^2 + 3c + 1) \log(dx + c + 1)}{d^4} - \frac{(c^3 - 3c^2 + 3c - 1) \log(dx + c - 1)}{d^4} \right) \right)$$

$$+ ac^2e^2x + \frac{(2(dx + c) \operatorname{artanh}(dx + c) + \log(-(dx + c)^2 + 1))bc^2e^2}{2d}$$

input `integrate((d*e*x+c*e)^2*(a+b*arctanh(d*x+c)),x, algorithm="maxima")`

output

```
1/3*a*d^2*e^2*x^3 + a*c*d*e^2*x^2 + 1/2*(2*x^2*arctanh(d*x + c) + d*(2*x/d
^2 - (c^2 + 2*c + 1)*log(d*x + c + 1)/d^3 + (c^2 - 2*c + 1)*log(d*x + c -
1)/d^3))*b*c*d*e^2 + 1/6*(2*x^3*arctanh(d*x + c) + d*((d*x^2 - 4*c*x)/d^3
+ (c^3 + 3*c^2 + 3*c + 1)*log(d*x + c + 1)/d^4 - (c^3 - 3*c^2 + 3*c - 1)*l
og(d*x + c - 1)/d^4))*b*d^2*e^2 + a*c^2*e^2*x + 1/2*(2*(d*x + c)*arctanh(d
*x + c) + log(-(d*x + c)^2 + 1))*b*c^2*e^2/d
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 322 vs. $2(63) = 126$.

Time = 0.14 (sec) , antiderivative size = 322, normalized size of antiderivative = 4.67

$$\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx)) dx =$$

$$-\frac{1}{6} ((c+1)d - (c-1)d) \left(\frac{be^2 \log\left(-\frac{dx+c+1}{dx+c-1} + 1\right)}{d^2} - \frac{be^2 \log\left(-\frac{dx+c+1}{dx+c-1}\right)}{d^2} - \frac{\left(\frac{3(dx+c+1)^2 be^2}{(dx+c-1)^2} + be^2\right) \log\left(\frac{3(dx+c+1)^2 be^2}{(dx+c-1)^2} + be^2\right)}{\frac{(dx+c+1)^3 d^2}{(dx+c-1)^3} - \frac{3(dx+c+1)^2 d^2}{(dx+c-1)^2} + 3d^2} \right)$$

input

```
integrate((d*e*x+c*e)^2*(a+b*arctanh(d*x+c)),x, algorithm="giac")
```

output

```
-1/6*((c + 1)*d - (c - 1)*d)*(b*e^2*log(-(d*x + c + 1)/(d*x + c - 1) + 1)/
d^2 - b*e^2*log(-(d*x + c + 1)/(d*x + c - 1))/d^2 - (3*(d*x + c + 1)^2*b*e
^2/(d*x + c - 1)^2 + b*e^2)*log(-(d*x + c + 1)/(d*x + c - 1))/((d*x + c +
1)^3*d^2/(d*x + c - 1)^3 - 3*(d*x + c + 1)^2*d^2/(d*x + c - 1)^2 + 3*(d*x
+ c + 1)*d^2/(d*x + c - 1) - d^2) - 2*(3*(d*x + c + 1)^2*a*e^2/(d*x + c -
1)^2 + a*e^2 + (d*x + c + 1)^2*b*e^2/(d*x + c - 1)^2 - (d*x + c + 1)*b*e^2
/(d*x + c - 1))/((d*x + c + 1)^3*d^2/(d*x + c - 1)^3 - 3*(d*x + c + 1)^2*d
^2/(d*x + c - 1)^2 + 3*(d*x + c + 1)*d^2/(d*x + c - 1) - d^2))
```

Mupad [B] (verification not implemented)

Time = 0.64 (sec) , antiderivative size = 237, normalized size of antiderivative = 3.43

$$\int (ce + dex)^2 (a + \operatorname{barctanh}(c + dx)) dx$$

$$= \frac{ad^2 e^2 x^3}{3} + \frac{bce^2 x}{3} + \frac{be^2 \ln(c + dx - 1)}{6d} + \frac{be^2 \ln(c + dx + 1)}{6d} + ac^2 e^2 x$$

$$+ \frac{bd^2 e^2 x^2}{6} + acde^2 x^2 + \frac{bc^2 e^2 x \ln(c + dx + 1)}{2} - \frac{bc^3 e^2 \ln(c + dx - 1)}{6d}$$

$$+ \frac{bc^3 e^2 \ln(c + dx + 1)}{6d} - \frac{bc^2 e^2 x \ln(1 - dx - c)}{2} + \frac{bd^2 e^2 x^3 \ln(c + dx + 1)}{6}$$

$$- \frac{bd^2 e^2 x^3 \ln(1 - dx - c)}{6} + \frac{bcd e^2 x^2 \ln(c + dx + 1)}{2} - \frac{bcd e^2 x^2 \ln(1 - dx - c)}{2}$$

input `int((c*e + d*e*x)^2*(a + b*atanh(c + d*x)),x)`output `(a*d^2*e^2*x^3)/3 + (b*c*e^2*x)/3 + (b*e^2*log(c + d*x - 1))/(6*d) + (b*e^2*log(c + d*x + 1))/(6*d) + a*c^2*e^2*x + (b*d*e^2*x^2)/6 + a*c*d*e^2*x^2 + (b*c^2*e^2*x*log(c + d*x + 1))/2 - (b*c^3*e^2*log(c + d*x - 1))/(6*d) + (b*c^3*e^2*log(c + d*x + 1))/(6*d) - (b*c^2*e^2*x*log(1 - d*x - c))/2 + (b*d^2*e^2*x^3*log(c + d*x + 1))/6 - (b*d^2*e^2*x^3*log(1 - d*x - c))/6 + (b*c*d*e^2*x^2*log(c + d*x + 1))/2 - (b*c*d*e^2*x^2*log(1 - d*x - c))/2`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.83

$$\int (ce + dex)^2 (a + \operatorname{barctanh}(c + dx)) dx$$

$$= \frac{e^2(2a \operatorname{atanh}(dx + c) b c^3 + 6a \operatorname{atanh}(dx + c) b c^2 dx + 6a \operatorname{atanh}(dx + c) b c d^2 x^2 + 2a \operatorname{atanh}(dx + c) b d^3 x^3 + 2a \operatorname{atanh}(dx + c) b c^2 d x^2 + 2a \operatorname{atanh}(dx + c) b c^3 d x^2 + 2a \operatorname{atanh}(dx + c) b c^2 d^2 x^2 + 2a \operatorname{atanh}(dx + c) b c^3 d^2 x^2 + 2a \operatorname{atanh}(dx + c) b c^2 d^3 x^2 + 2a \operatorname{atanh}(dx + c) b c^3 d^3 x^2)}{6d}$$

input `int((d*e*x+c*e)^2*(a+b*atanh(d*x+c)),x)`

output

```
(e**2*(2*atanh(c + d*x)*b*c**3 + 6*atanh(c + d*x)*b*c**2*d*x + 6*atanh(c +
d*x)*b*c*d**2*x**2 + 2*atanh(c + d*x)*b*d**3*x**3 + 2*atanh(c + d*x)*b +
2*log(c + d*x - 1)*b + 6*a*c**2*d*x + 6*a*c*d**2*x**2 + 2*a*d**3*x**3 + 2*
b*c*d*x + b*d**2*x**2))/(6*d)
```

3.11 $\int (ce + dex)(a + \operatorname{arctanh}(c + dx)) dx$

| | |
|---|-----|
| Optimal result | 129 |
| Mathematica [A] (verified) | 129 |
| Rubi [A] (verified) | 130 |
| Maple [A] (verified) | 132 |
| Fricas [A] (verification not implemented) | 132 |
| Sympy [B] (verification not implemented) | 133 |
| Maxima [B] (verification not implemented) | 133 |
| Giac [B] (verification not implemented) | 134 |
| Mupad [B] (verification not implemented) | 134 |
| Reduce [B] (verification not implemented) | 135 |

Optimal result

Integrand size = 19, antiderivative size = 48

$$\int (ce + dex)(a + \operatorname{arctanh}(c + dx)) dx = \frac{bex}{2} - \frac{b \operatorname{arctanh}(c + dx)}{2d} + \frac{e(c + dx)^2(a + \operatorname{arctanh}(c + dx))}{2d}$$

output $1/2*b*e*x - 1/2*b*e*\operatorname{arctanh}(d*x+c)/d + 1/2*e*(d*x+c)^2*(a+b*\operatorname{arctanh}(d*x+c))/d$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.60

$$\int (ce + dex)(a + \operatorname{arctanh}(c + dx)) dx = \frac{e(2bc + 2ac^2 + 2bdx + 4acdx + 2ad^2x^2 + 2b(c + dx)^2 \operatorname{arctanh}(c + dx) + b \log(1 - c - dx) - b \log(1 + c + dx))}{4d}$$

input $\operatorname{Integrate}[(c*e + d*e*x)*(a + b*\operatorname{ArcTanh}[c + d*x]),x]$

output $(e*(2*b*c + 2*a*c^2 + 2*b*d*x + 4*a*c*d*x + 2*a*d^2*x^2 + 2*b*(c + d*x)^2*\operatorname{ArcTanh}[c + d*x] + b*\operatorname{Log}[1 - c - d*x] - b*\operatorname{Log}[1 + c + d*x]))/(4*d)$

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.96, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {6657, 27, 6452, 262, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (ce + dex)(a + \operatorname{barctanh}(c + dx)) dx \\
 & \quad \downarrow \text{6657} \\
 & \frac{\int e(c + dx)(a + \operatorname{barctanh}(c + dx))d(c + dx)}{d} \\
 & \quad \downarrow \text{27} \\
 & \frac{e \int (c + dx)(a + \operatorname{barctanh}(c + dx))d(c + dx)}{d} \\
 & \quad \downarrow \text{6452} \\
 & \frac{e\left(\frac{1}{2}(c + dx)^2(a + \operatorname{barctanh}(c + dx)) - \frac{1}{2}b \int \frac{(c+dx)^2}{1-(c+dx)^2} d(c + dx)\right)}{d} \\
 & \quad \downarrow \text{262} \\
 & \frac{e\left(\frac{1}{2}(c + dx)^2(a + \operatorname{barctanh}(c + dx)) - \frac{1}{2}b\left(\int \frac{1}{1-(c+dx)^2} d(c + dx) - c - dx\right)\right)}{d} \\
 & \quad \downarrow \text{219} \\
 & \frac{e\left(\frac{1}{2}(c + dx)^2(a + \operatorname{barctanh}(c + dx)) - \frac{1}{2}b(\operatorname{arctanh}(c + dx) - c - dx)\right)}{d}
 \end{aligned}$$

input

```
Int[(c*e + d*e*x)*(a + b*ArcTanh[c + d*x]),x]
```

output

```
(e*(-1/2*(b*(-c - d*x + ArcTanh[c + d*x])) + ((c + d*x)^2*(a + b*ArcTanh[c + d*x]))/2)/d)
```

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 262 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m-1)*((a + b*x^2)^(p+1)/(b*(m+2*p+1))), x] - Simp[a*c^2*((m-1)/(b*(m+2*p+1)) Int[(c*x)^(m-2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2-1] && NeQ[m+2*p+1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m+1)*((a + b*ArcTanh[c*x^n])^p/(m+1)), x] - Simp[b*c*n*(p/(m+1)) Int[x^(m+n)*((a + b*ArcTanh[c*x^n])^(p-1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6657 `Int[((a_) + ArcTanh[(c_) + (d_)*(x_)])*(b_)^(p_)*((e_) + (f_)*(x_)^(m_)), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.27

| method | result |
|-------------------|---|
| derivativedivides | $\frac{ae(dx+c)^2 + be \left(\frac{(dx+c)^2 \operatorname{arctanh}(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} + \frac{\ln(dx+c-1)}{4} - \frac{\ln(dx+c+1)}{4} \right)}{d}$ |
| default | $\frac{ae(dx+c)^2 + be \left(\frac{(dx+c)^2 \operatorname{arctanh}(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} + \frac{\ln(dx+c-1)}{4} - \frac{\ln(dx+c+1)}{4} \right)}{d}$ |
| parts | $ae \left(\frac{1}{2} dx^2 + cx \right) + \frac{be \left(\frac{(dx+c)^2 \operatorname{arctanh}(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} + \frac{\ln(dx+c-1)}{4} - \frac{\ln(dx+c+1)}{4} \right)}{d}$ |
| parallelrisch | $\frac{-x^2 \operatorname{arctanh}(dx+c) b d^3 e - x^2 a d^3 e - 2x \operatorname{arctanh}(dx+c) b c d^2 e - 2x a c d^2 e - \operatorname{arctanh}(dx+c) b c^2 d e - x b d^2 e + 5a c^2 d e + a r}{2d^2}$ |
| orering | $\frac{(2d^3 x^3 + 5c d^2 x^2 + 4c^2 d x + c^3 - 2dx - c)(dex + ce)(a + b \operatorname{arctanh}(dx+c))}{2d(dx+c)^2} - \frac{x(dx+c+1)(dx+c-1)(de(a+b \operatorname{arctanh}(dx+c)))}{2d(dx+c)}$ |
| risch | $\frac{ebx(dx+2c)\ln(dx+c+1)}{4} - \frac{edb x^2 \ln(-dx-c+1)}{4} - \frac{ebx \ln(-dx-c+1)c}{2} + \frac{ead x^2}{2} - \frac{e \ln(dx+c-1) b c^2}{4d} + \frac{e \ln(-dx-c+1) b c^2}{4d}$ |

input `int((d*e*x+c*e)*(a+b*arctanh(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*(1/2*a*e*(d*x+c)^2+b*e*(1/2*(d*x+c)^2*arctanh(d*x+c)+1/2*d*x+1/2*c+1/4*ln(d*x+c-1)-1/4*ln(d*x+c+1)))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.52

$$\int (ce + dex)(a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \frac{2ad^2ex^2 + 2(2ac + b)dex + (bd^2ex^2 + 2bcdex + (bc^2 - b)e) \log\left(-\frac{dx+c+1}{dx+c-1}\right)}{4d}$$

input `integrate((d*e*x+c*e)*(a+b*arctanh(d*x+c)),x, algorithm="fricas")`

output `1/4*(2*a*d^2*e*x^2 + 2*(2*a*c + b)*d*e*x + (b*d^2*e*x^2 + 2*b*c*d*e*x + (b*c^2 - b)*e)*log(-(d*x + c + 1)/(d*x + c - 1)))/d`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs. $2(41) = 82$.

Time = 0.65 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.98

$$\int (ce + dex)(a + \operatorname{arctanh}(c + dx)) dx$$

$$= \begin{cases} acex + \frac{adex^2}{2} + \frac{bc^2e \operatorname{arctanh}(c+dx)}{2d} + bcex \operatorname{arctanh}(c + dx) + \frac{bdex^2 \operatorname{arctanh}(c+dx)}{2} + \frac{bex}{2} - \frac{be \operatorname{arctanh}(c+dx)}{2d} & \text{for } d \neq 0 \\ cex(a + b \operatorname{arctanh}(c)) & \text{otherwise} \end{cases}$$

input `integrate((d*e*x+c*e)*(a+b*atanh(d*x+c)), x)`

output `Piecewise((a*c*e*x + a*d*e*x**2/2 + b*c**2*e*atanh(c + d*x)/(2*d) + b*c*e*x*atanh(c + d*x) + b*d*e*x**2*atanh(c + d*x)/2 + b*e*x/2 - b*e*atanh(c + d*x)/(2*d), Ne(d, 0)), (c*e*x*(a + b*atanh(c)), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 113 vs. $2(42) = 84$.

Time = 0.03 (sec) , antiderivative size = 113, normalized size of antiderivative = 2.35

$$\int (ce + dex)(a + \operatorname{arctanh}(c + dx)) dx = \frac{1}{2} adex^2$$

$$+ \frac{1}{4} \left(2x^2 \operatorname{arctanh}(dx + c) + d \left(\frac{2x}{d^2} - \frac{(c^2 + 2c + 1) \log(dx + c + 1)}{d^3} + \frac{(c^2 - 2c + 1) \log(dx + c - 1)}{d^3} \right) \right)$$

$$+ acex + \frac{(2(dx + c) \operatorname{arctanh}(dx + c) + \log(-(dx + c)^2 + 1)) bce}{2d}$$

input `integrate((d*e*x+c*e)*(a+b*arctanh(d*x+c)), x, algorithm="maxima")`

output `1/2*a*d*e*x^2 + 1/4*(2*x^2*arctanh(d*x + c) + d*(2*x/d^2 - (c^2 + 2*c + 1)*log(d*x + c + 1)/d^3 + (c^2 - 2*c + 1)*log(d*x + c - 1)/d^3))*b*d*e + a*c*e*x + 1/2*(2*(d*x + c)*arctanh(d*x + c) + log(-(d*x + c)^2 + 1))*b*c*e/d`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 180 vs. $2(42) = 84$.

Time = 0.13 (sec) , antiderivative size = 180, normalized size of antiderivative = 3.75

$$\int (ce + dex)(a + \operatorname{arctanh}(c + dx)) dx$$

$$= \frac{1}{2} ((c + 1)d - (c - 1)d) \left(\frac{(dx + c + 1)be \log\left(-\frac{dx+c+1}{dx+c-1}\right)}{\left(\frac{(dx+c+1)^2 d^2}{(dx+c-1)^2} - \frac{2(dx+c+1)d^2}{dx+c-1} + d^2\right)(dx + c - 1)} + \frac{\frac{2(dx+c+1)ae}{dx+c-1} + \frac{(dx+c+1)be}{dx+c-1} - be}{\frac{(dx+c+1)^2 d^2}{(dx+c-1)^2} - \frac{2(dx+c+1)d^2}{dx+c-1} + d^2} \right)$$

input `integrate((d*e*x+c*e)*(a+b*arctanh(d*x+c)),x, algorithm="giac")`

output `1/2*((c + 1)*d - (c - 1)*d)*((d*x + c + 1)*b*e*log(-(d*x + c + 1)/(d*x + c - 1))/(((d*x + c + 1)^2*d^2/(d*x + c - 1)^2 - 2*(d*x + c + 1)*d^2/(d*x + c - 1) + d^2)*(d*x + c - 1)) + (2*(d*x + c + 1)*a*e/(d*x + c - 1) + (d*x + c + 1)*b*e/(d*x + c - 1) - b*e)/((d*x + c + 1)^2*d^2/(d*x + c - 1)^2 - 2*(d*x + c + 1)*d^2/(d*x + c - 1) + d^2))`

Mupad [B] (verification not implemented)

Time = 5.20 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.52

$$\int (ce + dex)(a + \operatorname{arctanh}(c + dx)) dx = \frac{bex}{2} + acex - \frac{be \operatorname{atanh}(c + dx)}{2d} + \frac{adex^2}{2}$$

$$+ \frac{bc^2e \operatorname{atanh}(c + dx)}{2d} + bcex \operatorname{atanh}(c + dx)$$

$$+ \frac{bdex^2 \operatorname{atanh}(c + dx)}{2}$$

input `int((c*e + d*e*x)*(a + b*atanh(c + d*x)),x)`

output `(b*e*x)/2 + a*c*e*x - (b*e*atanh(c + d*x))/(2*d) + (a*d*e*x^2)/2 + (b*c^2*e*atanh(c + d*x))/(2*d) + b*c*e*x*atanh(c + d*x) + (b*d*e*x^2*atanh(c + d*x))/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.48

$$\int (ce + dex)(a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \frac{e(\operatorname{atanh}(dx + c) b c^2 + 2 \operatorname{atanh}(dx + c) bcdx + \operatorname{atanh}(dx + c) b d^2 x^2 - \operatorname{atanh}(dx + c) b + 2acdx + a d^2 x^2)}{2d}$$

input `int((d*e*x+c*e)*(a+b*atanh(d*x+c)),x)`output `(e*(atanh(c + d*x)*b*c**2 + 2*atanh(c + d*x)*b*c*d*x + atanh(c + d*x)*b*d*
*2*x**2 - atanh(c + d*x)*b + 2*a*c*d*x + a*d**2*x**2 + b*d*x))/(2*d)`

3.12 $\int \frac{a+b\operatorname{arctanh}(c+dx)}{ce+dex} dx$

| | |
|--------------------------------------|-----|
| Optimal result | 136 |
| Mathematica [A] (verified) | 136 |
| Rubi [A] (verified) | 137 |
| Maple [A] (verified) | 138 |
| Fricas [F] | 138 |
| Sympy [F] | 139 |
| Maxima [F] | 139 |
| Giac [F] | 139 |
| Mupad [F(-1)] | 140 |
| Reduce [F] | 140 |

Optimal result

Integrand size = 21, antiderivative size = 54

$$\int \frac{a + b\operatorname{arctanh}(c + dx)}{ce + dex} dx = \frac{a \log(c + dx)}{de} - \frac{b \operatorname{PolyLog}(2, -c - dx)}{2de} + \frac{b \operatorname{PolyLog}(2, c + dx)}{2de}$$

output `a*ln(d*x+c)/d/e-1/2*b*polylog(2,-d*x-c)/d/e+1/2*b*polylog(2,d*x+c)/d/e`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00

$$\int \frac{a + b\operatorname{arctanh}(c + dx)}{ce + dex} dx = \frac{a \log(c + dx)}{de} - \frac{b \operatorname{PolyLog}(2, -c - dx)}{2de} + \frac{b \operatorname{PolyLog}(2, c + dx)}{2de}$$

input `Integrate[(a + b*ArcTanh[c + d*x])/(c*e + d*e*x),x]`

output `(a*Log[c + d*x])/(d*e) - (b*PolyLog[2, -c - d*x])/(2*d*e) + (b*PolyLog[2, c + d*x])/(2*d*e)`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.80, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6657, 27, 6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{a + b \operatorname{arctanh}(c + dx)}{ce + dex} dx \\
 \downarrow 6657 \\
 \int \frac{a + b \operatorname{arctanh}(c + dx)}{e(c + dx)} d(c + dx) \\
 \downarrow 27 \\
 \int \frac{a + b \operatorname{arctanh}(c + dx)}{c + dx} d(c + dx) \\
 \downarrow 6446 \\
 \frac{a \log(c + dx) - \frac{1}{2} b \operatorname{PolyLog}(2, -c - dx) + \frac{1}{2} b \operatorname{PolyLog}(2, c + dx)}{de}
 \end{array}$$

input `Int[(a + b*ArcTanh[c + d*x])/(c*e + d*e*x), x]`

output `(a*Log[c + d*x] - (b*PolyLog[2, -c - d*x])/2 + (b*PolyLog[2, c + d*x])/2)/(d*e)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] :> Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 6446

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))/(x_), x_Symbol] := Simp[a*Log[x], x]
+ (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) /
; FreeQ[{a, b, c}, x]
```

rule 6657

```
Int[((a_.) + ArcTanh[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(
m_.), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x]
, x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0]
&& IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00

| method | result | size |
|-------------------|---|------|
| risch | $\frac{a \ln(-dx-c)}{de} + \frac{b \operatorname{dilog}(-dx-c+1)}{2de} - \frac{b \operatorname{dilog}(dx+c+1)}{2ed}$ | 54 |
| derivativedivides | $\frac{a \ln(dx+c)}{e} + \frac{b \left(\ln(dx+c) \operatorname{arctanh}(dx+c) - \frac{\operatorname{dilog}(dx+c)}{2} - \frac{\operatorname{dilog}(dx+c+1)}{2} - \frac{\ln(dx+c) \ln(dx+c+1)}{2} \right)}{d}$ | 68 |
| default | $\frac{a \ln(dx+c)}{e} + \frac{b \left(\ln(dx+c) \operatorname{arctanh}(dx+c) - \frac{\operatorname{dilog}(dx+c)}{2} - \frac{\operatorname{dilog}(dx+c+1)}{2} - \frac{\ln(dx+c) \ln(dx+c+1)}{2} \right)}{d}$ | 68 |
| parts | $\frac{a \ln(dx+c)}{de} + \frac{b \left(\ln(dx+c) \operatorname{arctanh}(dx+c) - \frac{\operatorname{dilog}(dx+c)}{2} - \frac{\operatorname{dilog}(dx+c+1)}{2} - \frac{\ln(dx+c) \ln(dx+c+1)}{2} \right)}{ed}$ | 70 |

input

```
int((a+b*arctanh(d*x+c))/(d*e*x+c*e),x,method=_RETURNVERBOSE)
```

output

```
1/d/e*a*ln(-d*x-c)+1/2/d/e*b*dilog(-d*x-c+1)-1/2*b/e/d*dilog(d*x+c+1)
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{ce + dex} dx = \int \frac{b \operatorname{arctanh}(dx + c) + a}{dex + ce} dx$$

input

```
integrate((a+b*arctanh(d*x+c))/(d*e*x+c*e),x, algorithm="fricas")
```

output `integral((b*arctanh(d*x + c) + a)/(d*e*x + c*e), x)`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{ce + dex} dx = \frac{\int \frac{a}{c+dx} dx + \int \frac{b \operatorname{atanh}(c+dx)}{c+dx} dx}{e}$$

input `integrate((a+b*atanh(d*x+c))/(d*e*x+c*e), x)`

output `(Integral(a/(c + d*x), x) + Integral(b*atanh(c + d*x)/(c + d*x), x))/e`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{ce + dex} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{dex + ce} dx$$

input `integrate((a+b*arctanh(d*x+c))/(d*e*x+c*e), x, algorithm="maxima")`

output `1/2*b*integrate((log(d*x + c + 1) - log(-d*x - c + 1))/(d*e*x + c*e), x) + a*log(d*e*x + c*e)/(d*e)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{ce + dex} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{dex + ce} dx$$

input `integrate((a+b*arctanh(d*x+c))/(d*e*x+c*e), x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)/(d*e*x + c*e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{ce + dex} dx = \int \frac{a + b \operatorname{atanh}(c + dx)}{ce + dex} dx$$

input `int((a + b*atanh(c + d*x))/(c*e + d*e*x), x)`output `int((a + b*atanh(c + d*x))/(c*e + d*e*x), x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{ce + dex} dx = \frac{\left(\int \frac{\operatorname{atanh}(dx+c)}{dx+c} dx \right) bd + \log(dx + c) a}{de}$$

input `int((a+b*atanh(d*x+c))/(d*e*x+c*e), x)`output `(int(atanh(c + d*x)/(c + d*x), x)*b*d + log(c + d*x)*a)/(d*e)`

3.13 $\int \frac{a+b\operatorname{arctanh}(c+dx)}{(ce+dex)^2} dx$

| | |
|---|-----|
| Optimal result | 141 |
| Mathematica [A] (verified) | 141 |
| Rubi [A] (warning: unable to verify) | 142 |
| Maple [A] (verified) | 144 |
| Fricas [A] (verification not implemented) | 145 |
| Sympy [B] (verification not implemented) | 145 |
| Maxima [A] (verification not implemented) | 146 |
| Giac [B] (verification not implemented) | 146 |
| Mupad [B] (verification not implemented) | 147 |
| Reduce [B] (verification not implemented) | 147 |

Optimal result

Integrand size = 21, antiderivative size = 63

$$\int \frac{a + b\operatorname{arctanh}(c + dx)}{(ce + dex)^2} dx = -\frac{a + b\operatorname{arctanh}(c + dx)}{de^2(c + dx)} + \frac{b \log(c + dx)}{de^2} - \frac{b \log(1 - (c + dx)^2)}{2de^2}$$

output

```
-(a+b*arctanh(d*x+c))/d/e^2/(d*x+c)+b*ln(d*x+c)/d/e^2-1/2*b*ln(1-(d*x+c)^2)/d/e^2
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.10

$$\int \frac{a + b\operatorname{arctanh}(c + dx)}{(ce + dex)^2} dx = -\frac{\frac{2a}{c+dx} + \frac{2b\operatorname{arctanh}(c+dx)}{c+dx} - 2b \log(c + dx) + b \log(1 - c^2 - 2cdx - d^2x^2)}{2de^2}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])/(c*e + d*e*x)^2,x]
```

output

```
-1/2*((2*a)/(c + d*x) + (2*b*ArcTanh[c + d*x])/(c + d*x) - 2*b*Log[c + d*x]
] + b*Log[1 - c^2 - 2*c*d*x - d^2*x^2])/(d*e^2)
```

Rubi [A] (warning: unable to verify)

Time = 0.29 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.84, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6657, 27, 6452, 243, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}(c + dx)}{(c + dx)^2} dx \\
 & \quad \downarrow 6657 \\
 & \frac{\int \frac{a + b \operatorname{arctanh}(c + dx)}{e^{2(c + dx)}} d(c + dx)}{d} \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{a + b \operatorname{arctanh}(c + dx)}{(c + dx)^2} d(c + dx)}{de^2} \\
 & \quad \downarrow 6452 \\
 & \frac{b \int \frac{1}{(c + dx)(1 - (c + dx)^2)} d(c + dx) - \frac{a + b \operatorname{arctanh}(c + dx)}{c + dx}}{de^2} \\
 & \quad \downarrow 243 \\
 & \frac{\frac{1}{2} b \int \frac{1}{(-c - dx + 1)(c + dx)^2} d(c + dx)^2 - \frac{a + b \operatorname{arctanh}(c + dx)}{c + dx}}{de^2} \\
 & \quad \downarrow 47 \\
 & \frac{\frac{1}{2} b \left(\int \frac{1}{-c - dx + 1} d(c + dx)^2 + \int \frac{1}{(c + dx)^2} d(c + dx)^2 \right) - \frac{a + b \operatorname{arctanh}(c + dx)}{c + dx}}{de^2} \\
 & \quad \downarrow 14 \\
 & \frac{\frac{1}{2} b \left(\int \frac{1}{-c - dx + 1} d(c + dx)^2 + \log((c + dx)^2) \right) - \frac{a + b \operatorname{arctanh}(c + dx)}{c + dx}}{de^2}
 \end{aligned}$$

↓ 16

$$\frac{\frac{1}{2}b(\log((c+dx)^2) - \log(-c-dx+1)) - \frac{a+b\operatorname{arctanh}(c+dx)}{c+dx}}{de^2}$$

input `Int[(a + b*ArcTanh[c + d*x])/(c*e + d*e*x)^2,x]`

output `((-(a + b*ArcTanh[c + d*x])/(c + d*x)) + (b*(-Log[1 - c - d*x] + Log[(c + d*x)^2]))/2)/(d*e^2)`

Defintions of rubi rules used

rule 14 `Int[(a_)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_)/((a_) + (b_)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 47 `Int[1/(((a_) + (b_)*(x_))*((c_) + (d_)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

rule 243 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 6452

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :
> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m
+ 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x
], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1]
&& IntegerQ[m])) && NeQ[m, -1]
```

rule 6657

```
Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] :=> Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x
], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0]
&& IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.53 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.02

| method | result |
|-------------------|---|
| derivativedivides | $-\frac{a}{e^2(dx+c)} + \frac{b \left(-\frac{\operatorname{arctanh}(dx+c)}{dx+c} - \frac{\ln(dx+c-1)}{2} + \ln(dx+c) - \frac{\ln(dx+c+1)}{2} \right)}{e^2 d}$ |
| default | $-\frac{a}{e^2(dx+c)} + \frac{b \left(-\frac{\operatorname{arctanh}(dx+c)}{dx+c} - \frac{\ln(dx+c-1)}{2} + \ln(dx+c) - \frac{\ln(dx+c+1)}{2} \right)}{e^2 d}$ |
| parts | $-\frac{a}{d(dx+c)e^2} + \frac{b \left(-\frac{\operatorname{arctanh}(dx+c)}{dx+c} - \frac{\ln(dx+c-1)}{2} + \ln(dx+c) - \frac{\ln(dx+c+1)}{2} \right)}{e^2 d}$ |
| parallelrisch | $-\frac{3 \ln(dx+c-1) x b c d^2 - 3 \ln(dx+c) x b c d^2 + 3 x \operatorname{arctanh}(dx+c) b c d^2 + 3 \ln(dx+c-1) b c^2 d - 3 \ln(dx+c) b c^2 d + 3 \operatorname{arctanh}(dx+c) b c^2 d}{3(dx+c)d^2 e^2 c}$ |
| risch | $-\frac{b \ln(dx+c+1)}{2d(dx+c)e^2} - \frac{\ln(d^2 x^2 + 2cdx + c^2 - 1) b dx - 2 \ln(-dx-c) b dx + \ln(d^2 x^2 + 2cdx + c^2 - 1) b c - 2 \ln(-dx-c) b c - b \ln(-dx-c)}{2e^2(dx+c)d}$ |

input

```
int((a+b*arctanh(d*x+c))/(d*e*x+c*e)^2,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-a/e^2/(d*x+c)+b/e^2*(-1/(d*x+c)*arctanh(d*x+c)-1/2*ln(d*x+c-1)+ln(d*
x+c)-1/2*ln(d*x+c+1)))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.35

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(ce + dex)^2} dx = \frac{(bdx + bc) \log(d^2x^2 + 2cdx + c^2 - 1) - 2(bdx + bc) \log(dx + c) + b \log\left(-\frac{dx+c+1}{dx+c-1}\right) + 2a}{2(d^2e^2x + cde^2)}$$

input `integrate((a+b*arctanh(d*x+c))/(d*e*x+c*e)^2,x, algorithm="fricas")`

output `-1/2*((b*d*x + b*c)*log(d^2*x^2 + 2*c*d*x + c^2 - 1) - 2*(b*d*x + b*c)*log(d*x + c) + b*log(-(d*x + c + 1)/(d*x + c - 1)) + 2*a)/(d^2*e^2*x + c*d*e^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 219 vs. 2(53) = 106.

Time = 1.10 (sec) , antiderivative size = 219, normalized size of antiderivative = 3.48

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(ce + dex)^2} dx = \begin{cases} -\frac{a}{cde^2 + d^2e^2x} + \frac{bc \log\left(\frac{c}{d} + x\right)}{cde^2 + d^2e^2x} - \frac{bc \log\left(\frac{c}{d} + x + \frac{1}{d}\right)}{cde^2 + d^2e^2x} + \frac{bc \operatorname{atanh}(c + dx)}{cde^2 + d^2e^2x} + \frac{bdx \log\left(\frac{c}{d} + x\right)}{cde^2 + d^2e^2x} - \frac{bdx \log\left(\frac{c}{d} + x + \frac{1}{d}\right)}{cde^2 + d^2e^2x} + \frac{bdx \operatorname{atanh}(c + dx)}{cde^2 + d^2e^2x} \\ \frac{x(a + b \operatorname{atanh}(c))}{c^2e^2} \end{cases}$$

input `integrate((a+b*atanh(d*x+c))/(d*e*x+c*e)**2,x)`

output `Piecewise((-a/(c*d*e**2 + d**2*e**2*x) + b*c*log(c/d + x)/(c*d*e**2 + d**2*e**2*x) - b*c*log(c/d + x + 1/d)/(c*d*e**2 + d**2*e**2*x) + b*c*atanh(c + d*x)/(c*d*e**2 + d**2*e**2*x) + b*d*x*log(c/d + x)/(c*d*e**2 + d**2*e**2*x) - b*d*x*log(c/d + x + 1/d)/(c*d*e**2 + d**2*e**2*x) + b*d*x*atanh(c + d*x)/(c*d*e**2 + d**2*e**2*x) - b*atanh(c + d*x)/(c*d*e**2 + d**2*e**2*x), Ne(d, 0)), (x*(a + b*atanh(c))/(c**2*e**2), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.51

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(ce + dex)^2} dx =$$

$$-\frac{1}{2} \left(d \left(\frac{\log(dx + c + 1)}{d^2 e^2} - \frac{2 \log(dx + c)}{d^2 e^2} + \frac{\log(dx + c - 1)}{d^2 e^2} \right) + \frac{2 \operatorname{artanh}(dx + c)}{d^2 e^2 x + c d e^2} \right) b$$

$$- \frac{a}{d^2 e^2 x + c d e^2}$$

input `integrate((a+b*arctanh(d*x+c))/(d*e*x+c*e)^2,x, algorithm="maxima")`

output `-1/2*(d*(log(d*x + c + 1)/(d^2*e^2) - 2*log(d*x + c)/(d^2*e^2) + log(d*x + c - 1)/(d^2*e^2)) + 2*arctanh(d*x + c)/(d^2*e^2*x + c*d*e^2))*b - a/(d^2*e^2*x + c*d*e^2)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 152 vs. 2(61) = 122.

Time = 0.12 (sec) , antiderivative size = 152, normalized size of antiderivative = 2.41

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(ce + dex)^2} dx$$

$$= \frac{1}{2} ((c + 1)d - (c - 1)d) \left(\frac{b \log\left(-\frac{dx+c+1}{dx+c-1}\right)}{\frac{(dx+c+1)d^2 e^2}{dx+c-1} + d^2 e^2} + \frac{2a}{\frac{(dx+c+1)d^2 e^2}{dx+c-1} + d^2 e^2} + \frac{b \log\left(-\frac{dx+c+1}{dx+c-1} - 1\right)}{d^2 e^2} - \frac{b \log\left(-\frac{dx+c+1}{dx+c-1}\right)}{d^2 e^2} \right)$$

input `integrate((a+b*arctanh(d*x+c))/(d*e*x+c*e)^2,x, algorithm="giac")`

output `1/2*((c + 1)*d - (c - 1)*d)*(b*log(-(d*x + c + 1)/(d*x + c - 1))/((d*x + c + 1)*d^2*e^2/(d*x + c - 1) + d^2*e^2) + 2*a/((d*x + c + 1)*d^2*e^2/(d*x + c - 1) + d^2*e^2) + b*log(-(d*x + c + 1)/(d*x + c - 1) - 1)/(d^2*e^2) - b*log(-(d*x + c + 1)/(d*x + c - 1))/(d^2*e^2))`

Mupad [B] (verification not implemented)

Time = 4.68 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.94

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(ce + dex)^2} dx = \frac{b \ln(1 - dx - c)}{2x d^2 e^2 + 2c d e^2} - \frac{b \ln(c + dx + 1)}{2(x d^2 e^2 + c d e^2)} - \frac{a}{x d^2 e^2 + c d e^2} - \frac{b \ln(c^2 + 2c dx + d^2 x^2 - 1)}{2d e^2} + \frac{b \ln(c + dx)}{d e^2}$$

input `int((a + b*atanh(c + d*x))/(c*e + d*e*x)^2,x)`output `(b*log(1 - d*x - c))/(2*d^2*e^2*x + 2*c*d*e^2) - (b*log(c + d*x + 1))/(2*(d^2*e^2*x + c*d*e^2)) - a/(d^2*e^2*x + c*d*e^2) - (b*log(c^2 + d^2*x^2 + 2*c*d*x - 1))/(2*d*e^2) + (b*log(c + d*x))/(d*e^2)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 155, normalized size of antiderivative = 2.46

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(ce + dex)^2} dx = \frac{2a \operatorname{atanh}(dx + c) b dx - \log(dx + c - 1) b c^2 - \log(dx + c - 1) b c dx + \log(dx + c - 1) b c + \log(dx + c - 1)}{2(c + dx)^2}$$

input `int((a+b*atanh(d*x+c))/(d*e*x+c*e)^2,x)`output `(2*atanh(c + d*x)*b*d*x - log(c + d*x - 1)*b*c**2 - log(c + d*x - 1)*b*c*d*x + log(c + d*x - 1)*b*c + log(c + d*x - 1)*b*d*x - log(c + d*x + 1)*b*c**2 - log(c + d*x + 1)*b*c*d*x - log(c + d*x + 1)*b*c - log(c + d*x + 1)*b*d*x + 2*log(c + d*x)*b*c**2 + 2*log(c + d*x)*b*c*d*x + 2*a*d*x)/(2*c*d*e**2*(c + d*x))`

3.14 $\int \frac{a+b\operatorname{arctanh}(c+dx)}{(ce+dex)^3} dx$

| | |
|---|-----|
| Optimal result | 148 |
| Mathematica [A] (verified) | 148 |
| Rubi [A] (verified) | 149 |
| Maple [A] (verified) | 151 |
| Fricas [A] (verification not implemented) | 151 |
| Sympy [B] (verification not implemented) | 152 |
| Maxima [B] (verification not implemented) | 152 |
| Giac [B] (verification not implemented) | 153 |
| Mupad [B] (verification not implemented) | 153 |
| Reduce [B] (verification not implemented) | 154 |

Optimal result

Integrand size = 21, antiderivative size = 63

$$\int \frac{a + b\operatorname{arctanh}(c + dx)}{(ce + dex)^3} dx = -\frac{b}{2de^3(c + dx)} + \frac{b\operatorname{arctanh}(c + dx)}{2de^3} - \frac{a + b\operatorname{arctanh}(c + dx)}{2de^3(c + dx)^2}$$

output

$-1/2*b/d/e^3/(d*x+c)+1/2*b*\operatorname{arctanh}(d*x+c)/d/e^3-1/2*(a+b*\operatorname{arctanh}(d*x+c))/d/e^3/(d*x+c)^2$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.59

$$\int \frac{a + b\operatorname{arctanh}(c + dx)}{(ce + dex)^3} dx = -\frac{a}{2de^3(c + dx)^2} - \frac{b}{2de^3(c + dx)} - \frac{b\operatorname{arctanh}(c + dx)}{2de^3(c + dx)^2} - \frac{b\log(1 - c - dx)}{4de^3} + \frac{b\log(1 + c + dx)}{4de^3}$$

input

`Integrate[(a + b*ArcTanh[c + d*x])/(c*e + d*e*x)^3,x]`

output

$$\frac{-1/2*a/(d*e^3*(c + d*x)^2) - b/(2*d*e^3*(c + d*x)) - (b*ArcTanh[c + d*x])/(2*d*e^3*(c + d*x)^2) - (b*Log[1 - c - d*x])/(4*d*e^3) + (b*Log[1 + c + d*x])/(4*d*e^3)}$$

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.79, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {6657, 27, 6452, 264, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}(c + dx)}{(ce + dex)^3} dx \\ & \quad \downarrow \text{6657} \\ & \int \frac{a + b \operatorname{arctanh}(c + dx)}{e^3(c + dx)^3} d(c + dx) \\ & \quad \downarrow \text{27} \\ & \int \frac{a + b \operatorname{arctanh}(c + dx)}{(c + dx)^3} d(c + dx) \\ & \quad \downarrow \text{6452} \\ & \frac{\frac{1}{2}b \int \frac{1}{(c + dx)^2(1 - (c + dx)^2)} d(c + dx) - \frac{a + b \operatorname{arctanh}(c + dx)}{2(c + dx)^2}}{de^3} \\ & \quad \downarrow \text{264} \\ & \frac{\frac{1}{2}b \left(\int \frac{1}{1 - (c + dx)^2} d(c + dx) - \frac{1}{c + dx} \right) - \frac{a + b \operatorname{arctanh}(c + dx)}{2(c + dx)^2}}{de^3} \\ & \quad \downarrow \text{219} \\ & \frac{\frac{1}{2}b \left(\operatorname{arctanh}(c + dx) - \frac{1}{c + dx} \right) - \frac{a + b \operatorname{arctanh}(c + dx)}{2(c + dx)^2}}{de^3} \end{aligned}$$

input

$$\text{Int}[(a + b*ArcTanh[c + d*x])/(c*e + d*e*x)^3,x]$$

output
$$\frac{((b*(-(c + d*x)^{-1}) + \text{ArcTanh}[c + d*x]))/2 - (a + b*\text{ArcTanh}[c + d*x])/(2*(c + d*x)^2)}{(d*e^3)}$$

Defintions of rubi rules used

rule 27
$$\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_)] /; \text{FreeQ}[b, x]$$

rule 219
$$\text{Int}[(a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 264
$$\text{Int}[(c_)*(x_)^m*((a_) + (b_)*(x_)^2)^p], x_Symbol] \rightarrow \text{Simp}[(c*x)^{(m+1)}*((a + b*x^2)^{(p+1})/(a*c*(m+1))), x] - \text{Simp}[b*((m+2*p+3)/(a*c^2*(m+1))) \text{Int}[(c*x)^{(m+2)}*(a + b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 6452
$$\text{Int}[(a_) + \text{ArcTanh}[(c_)*(x_)^{n_}]]*(b_)^p*(x_)^m], x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{p-1}/(1 - c^2*x^{(2*n)})), x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$$

rule 6657
$$\text{Int}[(a_) + \text{ArcTanh}[(c_) + (d_)*(x_)]*(b_)^p*((e_) + (f_)*(x_)^m], x_Symbol] \rightarrow \text{Simp}[1/d \ \text{Subst}[\text{Int}[(f*(x/d))^m*(a + b*\text{ArcTanh}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m\}, x \ \&\& \ \text{EqQ}[d*e - c*f, 0] \ \&\& \ \text{IGtQ}[p, 0]$$

Maple [A] (verified)

Time = 0.76 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.06

| method | result |
|-------------------|--|
| derivativedivides | $-\frac{a}{2e^3(dx+c)^2} + \frac{b\left(-\frac{\operatorname{arctanh}(dx+c)}{2(dx+c)^2} - \frac{\ln(dx+c-1)}{4} + \frac{\ln(dx+c+1)}{4} - \frac{1}{2(dx+c)}\right)}{e^3 d}$ |
| default | $-\frac{a}{2e^3(dx+c)^2} + \frac{b\left(-\frac{\operatorname{arctanh}(dx+c)}{2(dx+c)^2} - \frac{\ln(dx+c-1)}{4} + \frac{\ln(dx+c+1)}{4} - \frac{1}{2(dx+c)}\right)}{e^3 d}$ |
| parts | $-\frac{a}{2e^3(dx+c)^2 d} + \frac{b\left(-\frac{\operatorname{arctanh}(dx+c)}{2(dx+c)^2} - \frac{\ln(dx+c-1)}{4} + \frac{\ln(dx+c+1)}{4} - \frac{1}{2(dx+c)}\right)}{e^3 d}$ |
| parallelrisc | $-\frac{4x^2 \operatorname{arctanh}(dx+c)bc d^4 - 8x \operatorname{arctanh}(dx+c)bc^2 d^3 - d^4 b x^2 - 4 \operatorname{arctanh}(dx+c)bc^3 d^2 + 2bc d^3 x + 4 \operatorname{arctanh}(dx+c)bc^3}{8(dx+c)^2 c d^3 e^3}$ |
| oring | $\frac{2(d^3 x^3 + 3c d^2 x^2 + 3c^2 dx + c^3 - dx - c)(a + b \operatorname{arctanh}(dx+c))}{d(dx+ce)^3} + \frac{(dx+c)^2(dx+c-1)(dx+c+1)\left(\frac{bd}{(1-(dx+c)^2)(dx+ce)^3} - \frac{1}{2d^2}\right)}{d(dx+ce)^3}$ |
| risc | $-\frac{b \ln(dx+c+1)}{4d(dx+c)^2 e^3} + \frac{\ln(-dx-c-1)bd^2 x^2 - bd^2 x^2 \ln(-dx-c+1) + 2 \ln(-dx-c-1)bcdx - 2bdx \ln(-dx-c+1)c + \ln(-dx-c-1)c^2}{4e^3(dx+c)^2 d}$ |

input `int((a+b*arctanh(d*x+c))/(d*e*x+c*e)^3,x,method=_RETURNVERBOSE)`

output `1/d*(-1/2*a/e^3/(d*x+c)^2+b/e^3*(-1/2/(d*x+c)^2*arctanh(d*x+c)-1/4*ln(d*x+c-1)+1/4*ln(d*x+c+1)-1/2/(d*x+c)))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.40

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(ce + dex)^3} dx$$

$$= -\frac{2 b d x + 2 b c - (b d^2 x^2 + 2 b c d x + b c^2 - b) \log\left(-\frac{d x + c + 1}{d x + c - 1}\right) + 2 a}{4 (d^3 e^3 x^2 + 2 c d^2 e^3 x + c^2 d e^3)}$$

input `integrate((a+b*arctanh(d*x+c))/(d*e*x+c*e)^3,x, algorithm="fricas")`

output `-1/4*(2*b*d*x + 2*b*c - (b*d^2*x^2 + 2*b*c*d*x + b*c^2 - b)*log(-(d*x + c + 1)/(d*x + c - 1)) + 2*a)/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 313 vs. $2(53) = 106$.

Time = 1.45 (sec) , antiderivative size = 313, normalized size of antiderivative = 4.97

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(ce + dex)^3} dx$$

$$= \begin{cases} -\frac{a}{2c^2de^3 + 4cd^2e^3x + 2d^3e^3x^2} + \frac{bc^2 \operatorname{atanh}(c+dx)}{2c^2de^3 + 4cd^2e^3x + 2d^3e^3x^2} + \frac{2bcdx \operatorname{atanh}(c+dx)}{2c^2de^3 + 4cd^2e^3x + 2d^3e^3x^2} - \frac{bc}{2c^2de^3 + 4cd^2e^3x + 2d^3e^3x^2} + \frac{bd^2x^2 a}{2c^2de^3 + 4cd^2e^3x + 2d^3e^3x^2} \\ \frac{x(a+b \operatorname{atanh}(c))}{c^3e^3} \end{cases}$$

input `integrate((a+b*atanh(d*x+c))/(d*e*x+c*e)**3,x)`

output

```
Piecewise((-a/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) + b*c**2*atanh(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) + 2*b*c*d*x*atanh(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) - b*c/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) + b*d**2*x**2*atanh(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) - b*d*x/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) - b*atanh(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2), Ne(d, 0)), (x*(a + b*atanh(c))/(c**3*e**3), True))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 131 vs. $2(57) = 114$.

Time = 0.03 (sec) , antiderivative size = 131, normalized size of antiderivative = 2.08

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(ce + dex)^3} dx =$$

$$-\frac{1}{4} \left(d \left(\frac{2}{d^3e^3x + cd^2e^3} - \frac{\log(dx + c + 1)}{d^2e^3} + \frac{\log(dx + c - 1)}{d^2e^3} \right) + \frac{2 \operatorname{artanh}(dx + c)}{d^3e^3x^2 + 2cd^2e^3x + c^2de^3} \right) b$$

$$- \frac{a}{2(d^3e^3x^2 + 2cd^2e^3x + c^2de^3)}$$

input `integrate((a+b*arctanh(d*x+c))/(d*e*x+c*e)^3,x, algorithm="maxima")`

output

```
-1/4*(d*(2/(d^3*e^3*x + c*d^2*e^3) - log(d*x + c + 1)/(d^2*e^3) + log(d*x
+ c - 1)/(d^2*e^3)) + 2*arctanh(d*x + c)/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^
2*d*e^3))*b - 1/2*a/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 194 vs. $2(57) = 114$.

Time = 0.12 (sec) , antiderivative size = 194, normalized size of antiderivative = 3.08

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(ce + dex)^3} dx$$

$$= \frac{1}{2} ((c + 1)d - (c - 1)d) \left(\frac{(dx + c + 1)b \log\left(-\frac{dx+c+1}{dx+c-1}\right)}{\left(\frac{(dx+c+1)^2 d^2 e^3}{(dx+c-1)^2} + \frac{2(dx+c+1)d^2 e^3}{dx+c-1} + d^2 e^3\right)(dx + c - 1)} + \frac{\frac{2(dx+c+1)a}{dx+c-1} + \frac{(dx+c+1)}{dx+c-1}}{\frac{(dx+c+1)^2 d^2 e^3}{(dx+c-1)^2} + \frac{2(dx+c+1)d^2 e^3}{dx+c-1}} \right)$$

input

```
integrate((a+b*arctanh(d*x+c))/(d*e*x+c*e)^3,x, algorithm="giac")
```

output

```
1/2*((c + 1)*d - (c - 1)*d)*((d*x + c + 1)*b*log(-(d*x + c + 1)/(d*x + c -
1))/(((d*x + c + 1)^2*d^2*e^3/(d*x + c - 1)^2 + 2*(d*x + c + 1)*d^2*e^3/(
d*x + c - 1) + d^2*e^3)*(d*x + c - 1)) + (2*(d*x + c + 1)*a/(d*x + c - 1)
+ (d*x + c + 1)*b/(d*x + c - 1) + b)/((d*x + c + 1)^2*d^2*e^3/(d*x + c - 1)
)^2 + 2*(d*x + c + 1)*d^2*e^3/(d*x + c - 1) + d^2*e^3))
```

Mupad [B] (verification not implemented)

Time = 5.04 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.06

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(ce + dex)^3} dx$$

$$= \frac{b \operatorname{atanh}(c + dx)}{2de^3} - \frac{\frac{a}{2} + \frac{bc}{2} + \frac{b \ln(c+dx+1)}{4} - \frac{b \ln(1-dx-c)}{4}}{de^3(c + dx)^2} + \frac{bdx}{2}$$

input

```
int((a + b*atanh(c + d*x))/(c*e + d*e*x)^3,x)
```

output

```
(b*atanh(c + d*x))/(2*d*e^3) - (a/2 + (b*c)/2 + (b*log(c + d*x + 1))/4 - (
b*log(1 - d*x - c))/4 + (b*d*x)/2)/(d*e^3*(c + d*x)^2)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 252, normalized size of antiderivative = 4.00

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(ce + dex)^3} dx$$

$$= \frac{4 \operatorname{atanh}(dx + c) bcdx + 2 \operatorname{atanh}(dx + c) b d^2 x^2 - \log(dx + c - 1) b c^4 - 2 \log(dx + c - 1) b c^3 dx - \log(dx + c + 1) b c^4 + 2 \log(dx + c + 1) b c^3 dx + \log(dx + c + 1) b d^2 x^2}{(d^3 e^3 (c + dx)^2)}$$

input

```
int((a+b*atanh(d*x+c))/(d*e*x+c*e)^3,x)
```

output

```
(4*atanh(c + d*x)*b*c*d*x + 2*atanh(c + d*x)*b*d**2*x**2 - log(c + d*x - 1)
)*b*c**4 - 2*log(c + d*x - 1)*b*c**3*d*x - log(c + d*x - 1)*b*c**2*d**2*x*
**2 + log(c + d*x - 1)*b*c**2 + 2*log(c + d*x - 1)*b*c*d*x + log(c + d*x -
1)*b*d**2*x**2 + log(c + d*x + 1)*b*c**4 + 2*log(c + d*x + 1)*b*c**3*d*x +
log(c + d*x + 1)*b*c**2*d**2*x**2 - log(c + d*x + 1)*b*c**2 - 2*log(c + d
*x + 1)*b*c*d*x - log(c + d*x + 1)*b*d**2*x**2 - 2*a*c**2 - b*c**3 + b*c*d
**2*x**2)/(4*c**2*d*e**3*(c**2 + 2*c*d*x + d**2*x**2))
```

3.15 $\int (ce + dex)^3 (a + \operatorname{arctanh}(c + dx))^2 dx$

| | |
|---|-----|
| Optimal result | 155 |
| Mathematica [A] (verified) | 156 |
| Rubi [A] (warning: unable to verify) | 156 |
| Maple [A] (verified) | 159 |
| Fricas [B] (verification not implemented) | 160 |
| Sympy [B] (verification not implemented) | 160 |
| Maxima [B] (verification not implemented) | 161 |
| Giac [B] (verification not implemented) | 162 |
| Mupad [B] (verification not implemented) | 163 |
| Reduce [B] (verification not implemented) | 164 |

Optimal result

Integrand size = 23, antiderivative size = 159

$$\int (ce + dex)^3 (a + \operatorname{arctanh}(c + dx))^2 dx = \frac{1}{2}abe^3x + \frac{b^2e^3(c + dx)^2}{12d} + \frac{b^2e^3(c + dx)\operatorname{arctanh}(c + dx)}{2d} + \frac{be^3(c + dx)^3(a + \operatorname{arctanh}(c + dx))}{6d} - \frac{e^3(a + \operatorname{arctanh}(c + dx))^2}{4d} + \frac{e^3(c + dx)^4(a + \operatorname{arctanh}(c + dx))^2}{4d} + \frac{b^2e^3 \log(1 - (c + dx)^2)}{3d}$$

output

```
1/2*a*b*e^3*x+1/12*b^2*e^3*(d*x+c)^2/d+1/2*b^2*e^3*(d*x+c)*arctanh(d*x+c)/
d+1/6*b*e^3*(d*x+c)^3*(a+b*arctanh(d*x+c))/d-1/4*e^3*(a+b*arctanh(d*x+c))^
2/d+1/4*e^3*(d*x+c)^4*(a+b*arctanh(d*x+c))^2/d+1/3*b^2*e^3*ln(1-(d*x+c)^2)
/d
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.93

$$\int (ce + dex)^3 (a + \operatorname{barctanh}(c + dx))^2 dx$$

$$= \frac{e^3(6ab(c + dx) + b^2(c + dx)^2 + 2ab(c + dx)^3 + 3a^2(c + dx)^4 + 2b(c + dx)(3b + b(c + dx)^2 + 3a(c + dx))}{12d}$$

input `Integrate[(c*e + d*e*x)^3*(a + b*ArcTanh[c + d*x])^2,x]`

output $(e^3(6a*b*(c + d*x) + b^2*(c + d*x)^2 + 2*a*b*(c + d*x)^3 + 3*a^2*(c + d*x)^4 + 2*b*(c + d*x)*(3*b + b*(c + d*x)^2 + 3*a*(c + d*x)^3)*\operatorname{ArcTanh}[c + d*x] + 3*b^2*(-1 + (c + d*x)^4)*\operatorname{ArcTanh}[c + d*x]^2 + b*(3*a + 4*b)*\operatorname{Log}[1 - c - d*x] + b*(-3*a + 4*b)*\operatorname{Log}[1 + c + d*x]))/(12*d)$

Rubi [A] (warning: unable to verify)

Time = 0.92 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.89, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.478$, Rules used = {6657, 27, 6452, 6542, 6452, 243, 49, 2009, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ce + dex)^3 (a + \operatorname{barctanh}(c + dx))^2 dx$$

$$\downarrow 6657$$

$$\frac{\int e^3(c + dx)^3 (a + \operatorname{barctanh}(c + dx))^2 d(c + dx)}{d}$$

$$\downarrow 27$$

$$\frac{e^3 \int (c + dx)^3 (a + \operatorname{barctanh}(c + dx))^2 d(c + dx)}{d}$$

$$\downarrow 6452$$

$$\frac{e^3 \left(\frac{1}{4}(c + dx)^4 (a + \operatorname{barctanh}(c + dx))^2 - \frac{1}{2}b \int \frac{(c+dx)^4 (a + \operatorname{barctanh}(c+dx))}{1-(c+dx)^2} d(c + dx) \right)}{d}$$

↓ 6542

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4 (a + \operatorname{barctanh}(c+dx))^2 - \frac{1}{2}b \left(\int \frac{(c+dx)^2 (a + \operatorname{barctanh}(c+dx))}{1-(c+dx)^2} d(c+dx) - \int (c+dx)^2 (a + \operatorname{barctanh}(c+dx)) d(c+dx) \right) \right)}{d}$$

↓ 6452

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4 (a + \operatorname{barctanh}(c+dx))^2 - \frac{1}{2}b \left(\int \frac{(c+dx)^2 (a + \operatorname{barctanh}(c+dx))}{1-(c+dx)^2} d(c+dx) + \frac{1}{3}b \int \frac{(c+dx)^3}{1-(c+dx)^2} d(c+dx) - \int (c+dx)^2 (a + \operatorname{barctanh}(c+dx)) d(c+dx) \right) \right)}{d}$$

↓ 243

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4 (a + \operatorname{barctanh}(c+dx))^2 - \frac{1}{2}b \left(\int \frac{(c+dx)^2 (a + \operatorname{barctanh}(c+dx))}{1-(c+dx)^2} d(c+dx) + \frac{1}{6}b \int \frac{(c+dx)^2}{-c-dx+1} d(c+dx)^2 - \int (c+dx)^2 (a + \operatorname{barctanh}(c+dx)) d(c+dx) \right) \right)}{d}$$

↓ 49

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4 (a + \operatorname{barctanh}(c+dx))^2 - \frac{1}{2}b \left(\int \frac{(c+dx)^2 (a + \operatorname{barctanh}(c+dx))}{1-(c+dx)^2} d(c+dx) + \frac{1}{6}b \int \left(\frac{1}{-c-dx+1} - 1 \right) d(c+dx) - \int (c+dx)^2 (a + \operatorname{barctanh}(c+dx)) d(c+dx) \right) \right)}{d}$$

↓ 2009

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4 (a + \operatorname{barctanh}(c+dx))^2 - \frac{1}{2}b \left(\int \frac{(c+dx)^2 (a + \operatorname{barctanh}(c+dx))}{1-(c+dx)^2} d(c+dx) - \frac{1}{3}(c+dx)^3 (a + \operatorname{barctanh}(c+dx)) - \int (c+dx)^2 (a + \operatorname{barctanh}(c+dx)) d(c+dx) \right) \right)}{d}$$

↓ 6542

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4 (a + \operatorname{barctanh}(c+dx))^2 - \frac{1}{2}b \left(- \int (a + \operatorname{barctanh}(c+dx)) d(c+dx) + \int \frac{a + \operatorname{barctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) - \int (c+dx)^2 (a + \operatorname{barctanh}(c+dx)) d(c+dx) \right) \right)}{d}$$

↓ 2009

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4 (a + \operatorname{barctanh}(c+dx))^2 - \frac{1}{2}b \left(\int \frac{a + \operatorname{barctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) - \frac{1}{3}(c+dx)^3 (a + \operatorname{barctanh}(c+dx)) - \int (c+dx)^2 (a + \operatorname{barctanh}(c+dx)) d(c+dx) \right) \right)}{d}$$

↓ 6510

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4 (a + \operatorname{barctanh}(c+dx))^2 - \frac{1}{2}b \left(-\frac{1}{3}(c+dx)^3 (a + \operatorname{barctanh}(c+dx)) + \frac{(a + \operatorname{barctanh}(c+dx))^2}{2b} - a(c+dx) - \int (c+dx)^2 (a + \operatorname{barctanh}(c+dx)) d(c+dx) \right) \right)}{d}$$

input `Int[(c*e + d*e*x)^3*(a + b*ArcTanh[c + d*x])^2,x]`

output `(e^3*(((c + d*x)^4*(a + b*ArcTanh[c + d*x])^2)/4 - (b*(-(a*(c + d*x)) - b*(c + d*x)*ArcTanh[c + d*x] - ((c + d*x)^3*(a + b*ArcTanh[c + d*x]))/3 + (a + b*ArcTanh[c + d*x])^2/(2*b) + (b*(-c - d*x - Log[1 - c - d*x]))/6 - (b*Log[1 - (c + d*x)^2]/2))/2))/d`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6510 `Int[((a_.) + ArcTanh[(c_.)*(x_)*(b_.))^(p_.)/((d_.) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6542

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)*((f_.)*(x_.))^(m_.)/((d_.) + (
e_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x]
x)]^p, x], x] - Simp[d*(f^2/e Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/
(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m,
1]
```

rule 6657

```
Int[((a_.) + ArcTanh[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(
m_.), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x
], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0]
&& IGtQ[p, 0]
```

Maple [A] (verified)

Time = 1.45 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.64

| method | result |
|-------------------|---|
| derivativedivides | $\frac{e^3 a^2 (dx+c)^4 + e^3 b^2 \left(\frac{(dx+c)^4 \operatorname{arctanh}(dx+c)^2}{4} + \frac{(dx+c)^3 \operatorname{arctanh}(dx+c)}{6} + \frac{(dx+c) \operatorname{arctanh}(dx+c)}{2} + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{4} \right)}{d}$ |
| default | $\frac{e^3 a^2 (dx+c)^4 + e^3 b^2 \left(\frac{(dx+c)^4 \operatorname{arctanh}(dx+c)^2}{4} + \frac{(dx+c)^3 \operatorname{arctanh}(dx+c)}{6} + \frac{(dx+c) \operatorname{arctanh}(dx+c)}{2} + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{4} \right)}{d}$ |
| parts | $\frac{e^3 a^2 (dx+c)^4}{4d} + \frac{e^3 b^2 \left(\frac{(dx+c)^4 \operatorname{arctanh}(dx+c)^2}{4} + \frac{(dx+c)^3 \operatorname{arctanh}(dx+c)}{6} + \frac{(dx+c) \operatorname{arctanh}(dx+c)}{2} + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{4} \right)}{d}$ |
| parallelrisch | $\frac{e^3 b^2 d + 18 d e^3 c^2 a^2 - 5 d e^3 c^2 b^2 + 24 x \operatorname{arctanh}(dx+c) a b c^3 d^2 e^3 + 36 x^2 \operatorname{arctanh}(dx+c) a b c^2 d^3 e^3 + 24 x^3 \operatorname{arctanh}(dx+c) a b c d^4 e^3}{d^2}$ |
| risch | $\frac{a b e^3 x}{2} - e^3 a b c^3 x \ln(-dx - c + 1) + \frac{e^3 d c a b x^2}{2} + \frac{e^3 a b c^2 x}{2} + \frac{e^3 d^2 b^2 c x^3 \ln(-dx - c + 1)^2}{4} - \frac{e^3 d^3 a b x^4}{4}$ |

input

```
int((d*e*x+c*e)^3*(a+b*arctanh(d*x+c))^2,x,method=_RETURNVERBOSE)
```

output

```
1/d*(1/4*e^3*a^2*(d*x+c)^4+e^3*b^2*(1/4*(d*x+c)^4*arctanh(d*x+c)^2+1/6*(d*
x+c)^3*arctanh(d*x+c)+1/2*(d*x+c)*arctanh(d*x+c)+1/4*arctanh(d*x+c)*ln(d*x
+c-1)-1/4*arctanh(d*x+c)*ln(d*x+c+1)-1/8*ln(d*x+c-1)*ln(1/2*d*x+1/2*c+1/2)
+1/16*ln(d*x+c-1)^2+1/16*ln(d*x+c+1)^2-1/8*(ln(d*x+c+1)-ln(1/2*d*x+1/2*c+1
/2))*ln(-1/2*d*x-1/2*c+1/2)+1/12*(d*x+c)^2+1/3*ln(d*x+c-1)+1/3*ln(d*x+c+1)
)+2*e^3*b*a*(1/4*(d*x+c)^4*arctanh(d*x+c)+1/12*(d*x+c)^3+1/4*d*x+1/4*c+1/8
*ln(d*x+c-1)-1/8*ln(d*x+c+1)))
```


output

```
Piecewise((a**2*c**3*e**3*x + 3*a**2*c**2*d*e**3*x**2/2 + a**2*c*d**2*e**3*x**3 + a**2*d**3*e**3*x**4/4 + a*b*c**4*e**3*atanh(c + d*x)/(2*d) + 2*a*b*c**3*e**3*x*atanh(c + d*x) + 3*a*b*c**2*d*e**3*x**2*atanh(c + d*x) + a*b*c**2*e**3*x/2 + 2*a*b*c*d**2*e**3*x**3*atanh(c + d*x) + a*b*c*d*e**3*x**2/2 + a*b*d**3*e**3*x**4*atanh(c + d*x)/2 + a*b*d**2*e**3*x**3/6 + a*b*e**3*x/2 - a*b*e**3*atanh(c + d*x)/(2*d) + b**2*c**4*e**3*atanh(c + d*x)**2/(4*d) + b**2*c**3*e**3*x*atanh(c + d*x)**2 + b**2*c**3*e**3*atanh(c + d*x)/(6*d) + 3*b**2*c**2*d*e**3*x**2*atanh(c + d*x)**2/2 + b**2*c**2*e**3*x*atanh(c + d*x)/2 + b**2*c*d**2*e**3*x**3*atanh(c + d*x)**2 + b**2*c*d*e**3*x**2*atanh(c + d*x)/2 + b**2*c*e**3*x/6 + b**2*c*e**3*atanh(c + d*x)/(2*d) + b**2*d**3*e**3*x**4*atanh(c + d*x)**2/4 + b**2*d**2*e**3*x**3*atanh(c + d*x)/6 + b**2*d*e**3*x**2/12 + b**2*e**3*x*atanh(c + d*x)/2 + 2*b**2*e**3*log(c/d + x + 1/d)/(3*d) - b**2*e**3*atanh(c + d*x)**2/(4*d) - 2*b**2*e**3*atanh(c + d*x)/(3*d), Ne(d, 0)), (c**3*e**3*x*(a + b*atanh(c))**2, True))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 827 vs. $2(145) = 290$.

Time = 0.26 (sec) , antiderivative size = 827, normalized size of antiderivative = 5.20

$$\int (ce + dex)^3 (a + b \operatorname{arctanh}(c + dx))^2 dx = \text{Too large to display}$$

input

```
integrate((d*e*x+c*e)^3*(a+b*arctanh(d*x+c))^2,x, algorithm="maxima")
```

output

```

1/4*a^2*d^3*e^3*x^4 + a^2*c*d^2*e^3*x^3 + 3/2*a^2*c^2*d*e^3*x^2 + 3/2*(2*x
^2*arctanh(d*x + c) + d*(2*x/d^2 - (c^2 + 2*c + 1)*log(d*x + c + 1)/d^3 +
(c^2 - 2*c + 1)*log(d*x + c - 1)/d^3))*a*b*c^2*d*e^3 + (2*x^3*arctanh(d*x
+ c) + d*((d*x^2 - 4*c*x)/d^3 + (c^3 + 3*c^2 + 3*c + 1)*log(d*x + c + 1)/d
^4 - (c^3 - 3*c^2 + 3*c - 1)*log(d*x + c - 1)/d^4))*a*b*c*d^2*e^3 + 1/12*(
6*x^4*arctanh(d*x + c) + d*(2*(d^2*x^3 - 3*c*d*x^2 + 3*(3*c^2 + 1)*x)/d^4
- 3*(c^4 + 4*c^3 + 6*c^2 + 4*c + 1)*log(d*x + c + 1)/d^5 + 3*(c^4 - 4*c^3
+ 6*c^2 - 4*c + 1)*log(d*x + c - 1)/d^5))*a*b*d^3*e^3 + a^2*c^3*e^3*x + (2
*(d*x + c)*arctanh(d*x + c) + log(-(d*x + c)^2 + 1))*a*b*c^3*e^3/d + 1/48*
(4*b^2*d^2*e^3*x^2 + 8*b^2*c*d*e^3*x + 3*(b^2*d^4*e^3*x^4 + 4*b^2*c*d^3*e^
3*x^3 + 6*b^2*c^2*d^2*e^3*x^2 + 4*b^2*c^3*d*e^3*x + (c^4*e^3 - e^3)*b^2)*l
og(d*x + c + 1)^2 + 3*(b^2*d^4*e^3*x^4 + 4*b^2*c*d^3*e^3*x^3 + 6*b^2*c^2*d
^2*e^3*x^2 + 4*b^2*c^3*d*e^3*x + (c^4*e^3 - e^3)*b^2)*log(-d*x - c + 1)^2
+ 4*(b^2*d^3*e^3*x^3 + 3*b^2*c*d^2*e^3*x^2 + 3*(c^2*d*e^3 + d*e^3)*b^2*x +
(c^3*e^3 + 3*c*e^3 + 4*e^3)*b^2)*log(d*x + c + 1) - 2*(2*b^2*d^3*e^3*x^3
+ 6*b^2*c*d^2*e^3*x^2 + 6*(c^2*d*e^3 + d*e^3)*b^2*x + 2*(c^3*e^3 + 3*c*e^3
- 4*e^3)*b^2 + 3*(b^2*d^4*e^3*x^4 + 4*b^2*c*d^3*e^3*x^3 + 6*b^2*c^2*d^2*e
^3*x^2 + 4*b^2*c^3*d*e^3*x + (c^4*e^3 - e^3)*b^2)*log(d*x + c + 1))*log(-d
*x - c + 1))/d

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 733 vs. $2(145) = 290$.

Time = 0.17 (sec) , antiderivative size = 733, normalized size of antiderivative = 4.61

$$\int (ce + dex)^3 (a + b \operatorname{arctanh}(c + dx))^2 dx = \text{Too large to display}$$

input

```
integrate((d*e*x+c*e)^3*(a+b*arctanh(d*x+c))^2,x, algorithm="giac")
```

output

```

-1/12*(4*b^2*e^3*log(-(d*x + c + 1)/(d*x + c - 1) + 1)/d^2 - 4*b^2*e^3*log
(-(d*x + c + 1)/(d*x + c - 1))/d^2 - 3*((d*x + c + 1)^3*b^2*e^3/(d*x + c -
1)^3 + (d*x + c + 1)*b^2*e^3/(d*x + c - 1))*log(-(d*x + c + 1)/(d*x + c -
1))^2/((d*x + c + 1)^4*d^2/(d*x + c - 1)^4 - 4*(d*x + c + 1)^3*d^2/(d*x +
c - 1)^3 + 6*(d*x + c + 1)^2*d^2/(d*x + c - 1)^2 - 4*(d*x + c + 1)*d^2/(d
*x + c - 1) + d^2) - 2*(6*(d*x + c + 1)^3*a*b*e^3/(d*x + c - 1)^3 + 6*(d*x
+ c + 1)*a*b*e^3/(d*x + c - 1) + 3*(d*x + c + 1)^3*b^2*e^3/(d*x + c - 1)^
3 - 6*(d*x + c + 1)^2*b^2*e^3/(d*x + c - 1)^2 + 5*(d*x + c + 1)*b^2*e^3/(d
*x + c - 1) - 2*b^2*e^3)*log(-(d*x + c + 1)/(d*x + c - 1))/((d*x + c + 1)^
4*d^2/(d*x + c - 1)^4 - 4*(d*x + c + 1)^3*d^2/(d*x + c - 1)^3 + 6*(d*x + c
+ 1)^2*d^2/(d*x + c - 1)^2 - 4*(d*x + c + 1)*d^2/(d*x + c - 1) + d^2) - 2
*(6*(d*x + c + 1)^3*a^2*e^3/(d*x + c - 1)^3 + 6*(d*x + c + 1)*a^2*e^3/(d*x
+ c - 1) + 6*(d*x + c + 1)^3*a*b*e^3/(d*x + c - 1)^3 - 12*(d*x + c + 1)^2
*a*b*e^3/(d*x + c - 1)^2 + 10*(d*x + c + 1)*a*b*e^3/(d*x + c - 1) - 4*a*b*
e^3 + (d*x + c + 1)^3*b^2*e^3/(d*x + c - 1)^3 - 2*(d*x + c + 1)^2*b^2*e^3/
(d*x + c - 1)^2 + (d*x + c + 1)*b^2*e^3/(d*x + c - 1))/((d*x + c + 1)^4*d^
2/(d*x + c - 1)^4 - 4*(d*x + c + 1)^3*d^2/(d*x + c - 1)^3 + 6*(d*x + c + 1
)^2*d^2/(d*x + c - 1)^2 - 4*(d*x + c + 1)*d^2/(d*x + c - 1) + d^2))*((c +
1)*d - (c - 1)*d)

```

Mupad [B] (verification not implemented)

Time = 5.60 (sec) , antiderivative size = 1730, normalized size of antiderivative = 10.88

$$\int (ce + dex)^3 (a + b \operatorname{arctanh}(c + dx))^2 dx = \text{Too large to display}$$

input

```
int((c*e + d*e*x)^3*(a + b*atanh(c + d*x))^2,x)
```


output

```
(e**3*(3*atanh(c + d*x)**2*b**2*c**4 + 12*atanh(c + d*x)**2*b**2*c**3*d*x
+ 18*atanh(c + d*x)**2*b**2*c**2*d**2*x**2 + 12*atanh(c + d*x)**2*b**2*c*d
**3*x**3 + 3*atanh(c + d*x)**2*b**2*d**4*x**4 - 3*atanh(c + d*x)**2*b**2 +
6*atanh(c + d*x)*a*b*c**4 + 24*atanh(c + d*x)*a*b*c**3*d*x + 36*atanh(c +
d*x)*a*b*c**2*d**2*x**2 + 24*atanh(c + d*x)*a*b*c*d**3*x**3 + 6*atanh(c +
d*x)*a*b*d**4*x**4 - 6*atanh(c + d*x)*a*b + 2*atanh(c + d*x)*b**2*c**3 +
6*atanh(c + d*x)*b**2*c**2*d*x + 6*atanh(c + d*x)*b**2*c*d**2*x**2 + 6*ata
nh(c + d*x)*b**2*c + 2*atanh(c + d*x)*b**2*d**3*x**3 + 6*atanh(c + d*x)*b*
**2*d*x + 8*atanh(c + d*x)*b**2 + 8*log(c + d*x - 1)*b**2 + 12*a**2*c**3*d*
x + 18*a**2*c**2*d**2*x**2 + 12*a**2*c*d**3*x**3 + 3*a**2*d**4*x**4 + 6*a*
b*c**2*d*x + 6*a*b*c*d**2*x**2 + 2*a*b*d**3*x**3 + 6*a*b*d*x + 2*b**2*c*d*
x + b**2*d**2*x**2))/(12*d)
```

3.16 $\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx$

| | |
|---|-----|
| Optimal result | 166 |
| Mathematica [A] (verified) | 167 |
| Rubi [A] (verified) | 167 |
| Maple [A] (verified) | 171 |
| Fricas [F] | 171 |
| Sympy [F] | 172 |
| Maxima [B] (verification not implemented) | 172 |
| Giac [F] | 173 |
| Mupad [F(-1)] | 173 |
| Reduce [F] | 174 |

Optimal result

Integrand size = 23, antiderivative size = 179

$$\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx = \frac{1}{3} b^2 e^2 x - \frac{b^2 e^2 \operatorname{arctanh}(c + dx)}{3d} + \frac{b e^2 (c + dx)^2 (a + b \operatorname{arctanh}(c + dx))}{3d} + \frac{e^2 (a + b \operatorname{arctanh}(c + dx))^2}{3d} + \frac{e^2 (c + dx)^3 (a + b \operatorname{arctanh}(c + dx))^2}{3d} - \frac{2 b e^2 (a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2}{1 - c - dx}\right)}{3d} - \frac{b^2 e^2 \operatorname{PolyLog}\left(2, -\frac{1 + c + dx}{1 - c - dx}\right)}{3d}$$

output

```
1/3*b^2*e^2*x-1/3*b^2*e^2*arctanh(d*x+c)/d+1/3*b*e^2*(d*x+c)^2*(a+b*arctanh(d*x+c))/d+1/3*e^2*(a+b*arctanh(d*x+c))^2/d+1/3*e^2*(d*x+c)^3*(a+b*arctanh(d*x+c))^2/d-2/3*b*e^2*(a+b*arctanh(d*x+c))*ln(2/(-d*x-c+1))/d-1/3*b^2*e^2*polylog(2,-(d*x+c+1)/(-d*x-c+1))/d
```

Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.84

$$\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx$$

$$= \frac{e^2 (a^2 (c + dx)^3 + ab((c + dx)^2 + 2(c + dx)^3 \operatorname{arctanh}(c + dx) + \log(-1 + (c + dx)^2)) + b^2 (c + dx - \operatorname{arctanh}(c + dx)))}{3d}$$

input

```
Integrate[(c*e + d*e*x)^2*(a + b*ArcTanh[c + d*x])^2,x]
```

output

```
(e^2*(a^2*(c + d*x)^3 + a*b*((c + d*x)^2 + 2*(c + d*x)^3*ArcTanh[c + d*x]
+ Log[-1 + (c + d*x)^2]) + b^2*(c + d*x - ArcTanh[c + d*x] + (c + d*x)^2*ArcTanh[c + d*x] - ArcTanh[c + d*x]^2 + (c + d*x)^3*ArcTanh[c + d*x]^2 - 2*ArcTanh[c + d*x]*Log[1 + E^(-2*ArcTanh[c + d*x])]) + PolyLog[2, -E^(-2*ArcTanh[c + d*x])])))/(3*d)
```

Rubi [A] (verified)

Time = 0.87 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.80, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.478$, Rules used = {6657, 27, 6452, 6542, 6452, 262, 219, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx$$

$$\downarrow 6657$$

$$\frac{\int e^2 (c + dx)^2 (a + b \operatorname{arctanh}(c + dx))^2 d(c + dx)}{d}$$

$$\downarrow 27$$

$$\frac{e^2 \int (c + dx)^2 (a + b \operatorname{arctanh}(c + dx))^2 d(c + dx)}{d}$$

$$\downarrow 6452$$

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a + \operatorname{barctanh}(c+dx))^2 - \frac{2}{3}b \int \frac{(c+dx)^3(a + \operatorname{barctanh}(c+dx))}{1-(c+dx)^2} d(c+dx) \right)}{d}$$

↓ 6542

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a + \operatorname{barctanh}(c+dx))^2 - \frac{2}{3}b \left(\int \frac{(c+dx)(a + \operatorname{barctanh}(c+dx))}{1-(c+dx)^2} d(c+dx) - \int (c+dx)(a + \operatorname{barctanh}(c+dx)) \right) \right)}{d}$$

↓ 6452

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a + \operatorname{barctanh}(c+dx))^2 - \frac{2}{3}b \left(\int \frac{(c+dx)(a + \operatorname{barctanh}(c+dx))}{1-(c+dx)^2} d(c+dx) + \frac{1}{2}b \int \frac{(c+dx)^2}{1-(c+dx)^2} d(c+dx) - \frac{1}{2} \right) \right)}{d}$$

↓ 262

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a + \operatorname{barctanh}(c+dx))^2 - \frac{2}{3}b \left(\int \frac{(c+dx)(a + \operatorname{barctanh}(c+dx))}{1-(c+dx)^2} d(c+dx) + \frac{1}{2}b \left(\int \frac{1}{1-(c+dx)^2} d(c+dx) - \int \frac{1}{1+(c+dx)^2} d(c+dx) \right) \right) \right)}{d}$$

↓ 219

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a + \operatorname{barctanh}(c+dx))^2 - \frac{2}{3}b \left(\int \frac{(c+dx)(a + \operatorname{barctanh}(c+dx))}{1-(c+dx)^2} d(c+dx) - \frac{1}{2}(c+dx)^2(a + \operatorname{barctanh}(c+dx)) \right) \right)}{d}$$

↓ 6546

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a + \operatorname{barctanh}(c+dx))^2 - \frac{2}{3}b \left(\int \frac{a + \operatorname{barctanh}(c+dx)}{-c-dx+1} d(c+dx) - \frac{1}{2}(c+dx)^2(a + \operatorname{barctanh}(c+dx)) \right) \right)}{d}$$

↓ 6470

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a + \operatorname{barctanh}(c+dx))^2 - \frac{2}{3}b \left(-b \int \frac{\log\left(\frac{-c-dx+1}{1-(c+dx)^2}\right)}{1-(c+dx)^2} d(c+dx) - \frac{1}{2}(c+dx)^2(a + \operatorname{barctanh}(c+dx)) \right) \right)}{d}$$

↓ 2849

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a + \operatorname{barctanh}(c+dx))^2 - \frac{2}{3}b \left(b \int \frac{\log\left(\frac{-c-dx+1}{1-\frac{2}{-c-dx+1}}\right)}{1-\frac{2}{-c-dx+1}} d\frac{1}{-c-dx+1} - \frac{1}{2}(c+dx)^2(a + \operatorname{barctanh}(c+dx)) \right) \right)}{d}$$

↓ 2752

$$\frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + \operatorname{arctanh}(c + dx))^2 - \frac{2}{3} b \left(-\frac{1}{2} (c + dx)^2 (a + \operatorname{arctanh}(c + dx)) - \frac{(a + \operatorname{arctanh}(c + dx))^2}{2b} + \log \left(\frac{\dots}{d} \right) \right) \right)}{d}$$

input `Int[(c*e + d*e*x)^2*(a + b*ArcTanh[c + d*x])^2,x]`

output `(e^2*(((c + d*x)^3*(a + b*ArcTanh[c + d*x])^2)/3 - (2*b*((b*(-c - d*x + ArcTanh[c + d*x]))/2 - ((c + d*x)^2*(a + b*ArcTanh[c + d*x]))/2 - (a + b*ArcTanh[c + d*x])^2/(2*b) + (a + b*ArcTanh[c + d*x])*Log[2/(1 - c - d*x)] + (b*PolyLog[2, 1 - 2/(1 - c - d*x)]/2))/3))/d`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 262 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m-1)*((a + b*x^2)^(p+1)/(b*(m+2*p+1))), x] - Simp[a*c^2*((m-1)/(b*(m+2*p+1)) Int[(c*x)^(m-2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2-1] && NeQ[m+2*p+1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6470 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6542 `Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_))/((d_.) + (e_.)*(x_)^2), x_Symbol] :> Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

rule 6546 `Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*(x_)/((d_.) + (e_.)*(x_)^2), x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/(c*d) Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6657 `Int[((a_.) + ArcTanh[(c_.) + (d_.)*(x_)])*(b_.))^(p_.)*((e_.) + (f_.)*(x_)^(m_.), x_Symbol] :> Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 1.99 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.40

| method | result |
|-------------------|--|
| derivativedivides | $\frac{a^2 e^2 (dx+c)^3}{3} + b^2 e^2 \left(\frac{(dx+c)^3 \operatorname{arctanh}(dx+c)^2}{3} + \frac{(dx+c)^2 \operatorname{arctanh}(dx+c)}{3} + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{3} + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c+1)}{3} \right)$ |
| default | $\frac{a^2 e^2 (dx+c)^3}{3} + b^2 e^2 \left(\frac{(dx+c)^3 \operatorname{arctanh}(dx+c)^2}{3} + \frac{(dx+c)^2 \operatorname{arctanh}(dx+c)}{3} + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{3} + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c+1)}{3} \right)$ |
| parts | $\frac{a^2 e^2 (dx+c)^3}{3d} + \frac{b^2 e^2 \left(\frac{(dx+c)^3 \operatorname{arctanh}(dx+c)^2}{3} + \frac{(dx+c)^2 \operatorname{arctanh}(dx+c)}{3} + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{3} + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c+1)}{3} \right)}{d}$ |
| risch | $\frac{2abx e^2 c}{3} + \frac{b^2 e^2 x}{3} - \frac{17b^2 e^2}{54d} - \frac{11e^2 ba}{9d} + \frac{e^2 b^2 c}{3d} + \frac{e^2 c^3 a^2}{3d} - e^2 dba \ln(-dx - c + 1) c x^2 - \frac{b^2 e^2 cd}{3}$ |

input `int((d*e*x+c*e)^2*(a+b*arctanh(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(1/3*a^2*e^2*(d*x+c)^3+b^2*e^2*(1/3*(d*x+c)^3*arctanh(d*x+c)^2+1/3*(d*x+c)^2*arctanh(d*x+c)+1/3*arctanh(d*x+c)*ln(d*x+c-1)+1/3*arctanh(d*x+c)*ln(d*x+c+1)+1/3*d*x+1/3*c+1/6*ln(d*x+c-1)-1/6*ln(d*x+c+1)-1/3*dilog(1/2*d*x+1/2*c+1/2)-1/6*ln(d*x+c-1)*ln(1/2*d*x+1/2*c+1/2)+1/12*ln(d*x+c-1)^2-1/12*ln(d*x+c+1)^2+1/6*(ln(d*x+c+1)-ln(1/2*d*x+1/2*c+1/2))*ln(-1/2*d*x-1/2*c+1/2))+2*a*b*e^2*(1/3*(d*x+c)^3*arctanh(d*x+c)+1/6*(d*x+c)^2+1/6*ln(d*x+c-1)+1/6*ln(d*x+c+1)))`

Fricas [F]

$$\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (dex + ce)^2 (b \operatorname{arctanh}(dx + c) + a)^2 dx$$

input `integrate((d*e*x+c*e)^2*(a+b*arctanh(d*x+c))^2,x, algorithm="fricas")`

output `integral(a^2*d^2*e^2*x^2 + 2*a^2*c*d*e^2*x + a^2*c^2*e^2 + (b^2*d^2*e^2*x^2 + 2*b^2*c*d*e^2*x + b^2*c^2*e^2)*arctanh(d*x + c)^2 + 2*(a*b*d^2*e^2*x^2 + 2*a*b*c*d*e^2*x + a*b*c^2*e^2)*arctanh(d*x + c), x)`

Sympy [F]

$$\begin{aligned}
& \int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx \\
&= e^2 \left(\int a^2 c^2 dx + \int a^2 d^2 x^2 dx + \int b^2 c^2 \operatorname{atanh}^2(c + dx) dx \right. \\
&\quad + \int 2abc^2 \operatorname{atanh}(c + dx) dx + \int 2a^2 cdx dx + \int b^2 d^2 x^2 \operatorname{atanh}^2(c + dx) dx \\
&\quad + \int 2abd^2 x^2 \operatorname{atanh}(c + dx) dx + \int 2b^2 cdx \operatorname{atanh}^2(c + dx) dx \\
&\quad \left. + \int 4abcdx \operatorname{atanh}(c + dx) dx \right)
\end{aligned}$$

input

```
integrate((d*e*x+c*e)**2*(a+b*atanh(d*x+c))**2,x)
```

output

```
e**2*(Integral(a**2*c**2, x) + Integral(a**2*d**2*x**2, x) + Integral(b**2*c**2*atanh(c + d*x)**2, x) + Integral(2*a*b*c**2*atanh(c + d*x), x) + Integral(2*a**2*c*d*x, x) + Integral(b**2*d**2*x**2*atanh(c + d*x)**2, x) + Integral(2*a*b*d**2*x**2*atanh(c + d*x), x) + Integral(2*b**2*c*d*x*atanh(c + d*x)**2, x) + Integral(4*a*b*c*d*x*atanh(c + d*x), x))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 619 vs. $2(157) = 314$.

Time = 0.24 (sec) , antiderivative size = 619, normalized size of antiderivative = 3.46

$$\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx = \text{Too large to display}$$

input

```
integrate((d*e*x+c*e)^2*(a+b*arctanh(d*x+c))^2,x, algorithm="maxima")
```

output

```

1/3*a^2*d^2*e^2*x^3 + a^2*c*d*e^2*x^2 + (2*x^2*arctanh(d*x + c) + d*(2*x/d
^2 - (c^2 + 2*c + 1)*log(d*x + c + 1)/d^3 + (c^2 - 2*c + 1)*log(d*x + c -
1)/d^3))*a*b*c*d*e^2 + 1/3*(2*x^3*arctanh(d*x + c) + d*((d*x^2 - 4*c*x)/d^
3 + (c^3 + 3*c^2 + 3*c + 1)*log(d*x + c + 1)/d^4 - (c^3 - 3*c^2 + 3*c - 1)
*log(d*x + c - 1)/d^4))*a*b*d^2*e^2 + a^2*c^2*e^2*x + (2*(d*x + c)*arctanh
(d*x + c) + log(-(d*x + c)^2 + 1))*a*b*c^2*e^2/d + 1/3*(log(d*x + c + 1)*l
og(-1/2*d*x - 1/2*c + 1/2) + dilog(1/2*d*x + 1/2*c + 1/2))*b^2*e^2/d + 1/6
*(c^2*e^2 - e^2)*b^2*log(d*x + c + 1)/d - 1/6*(c^2*e^2 - e^2)*b^2*log(d*x
+ c - 1)/d + 1/12*(4*b^2*d*e^2*x + (b^2*d^3*e^2*x^3 + 3*b^2*c*d^2*e^2*x^2
+ 3*b^2*c^2*d*e^2*x + (c^3*e^2 + e^2)*b^2)*log(d*x + c + 1)^2 + (b^2*d^3*e
^2*x^3 + 3*b^2*c*d^2*e^2*x^2 + 3*b^2*c^2*d*e^2*x + (c^3*e^2 - e^2)*b^2)*lo
g(-d*x - c + 1)^2 + 2*(b^2*d^2*e^2*x^2 + 2*b^2*c*d*e^2*x)*log(d*x + c + 1)
- 2*(b^2*d^2*e^2*x^2 + 2*b^2*c*d*e^2*x + (b^2*d^3*e^2*x^3 + 3*b^2*c*d^2*e
^2*x^2 + 3*b^2*c^2*d*e^2*x + (c^3*e^2 + e^2)*b^2)*log(d*x + c + 1))*log(-d
*x - c + 1))/d

```

Giac [F]

$$\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (dex + ce)^2 (b \operatorname{artanh}(dx + c) + a)^2 dx$$

input

```
integrate((d*e*x+c*e)^2*(a+b*arctanh(d*x+c))^2,x, algorithm="giac")
```

output

```
integrate((d*e*x + c*e)^2*(b*arctanh(d*x + c) + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (ce + dex)^2 (a + b \operatorname{atanh}(c + dx))^2 dx$$

input

```
int((c*e + d*e*x)^2*(a + b*atanh(c + d*x))^2,x)
```

output

```
int((c*e + d*e*x)^2*(a + b*atanh(c + d*x))^2, x)
```

Reduce [F]

$$\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx$$

$$= \frac{e^2 \left(\operatorname{atanh}(dx + c)^2 b^2 c^3 + 3 \operatorname{atanh}(dx + c)^2 b^2 c^2 dx + 3 \operatorname{atanh}(dx + c)^2 b^2 c d^2 x^2 - \operatorname{atanh}(dx + c)^2 b^2 c + \operatorname{atanh}(dx + c)^2 b^2 c^2 dx + 3 \operatorname{atanh}(dx + c)^2 b^2 c^2 dx^2 - \operatorname{atanh}(dx + c)^2 b^2 c + \operatorname{atanh}(dx + c)^2 b^2 c^2 dx^2 \right)}{(3d)}$$

input `int((d*e*x+c*e)^2*(a+b*atanh(d*x+c))^2,x)`

output `(e**2*(atanh(c + d*x)**2*b**2*c**3 + 3*atanh(c + d*x)**2*b**2*c**2*d*x + 3*atanh(c + d*x)**2*b**2*c*d**2*x**2 - atanh(c + d*x)**2*b**2*c + atanh(c + d*x)**2*b**2*d**3*x**3 + 2*atanh(c + d*x)*a*b*c**3 + 6*atanh(c + d*x)*a*b*c**2*d*x + 6*atanh(c + d*x)*a*b*c*d**2*x**2 + 2*atanh(c + d*x)*a*b*d**3*x**3 + 2*atanh(c + d*x)*a*b + atanh(c + d*x)*b**2*c**2 + 2*atanh(c + d*x)*b**2*c*d*x + atanh(c + d*x)*b**2*d**2*x**2 - atanh(c + d*x)*b**2 + 2*int((atanh(c + d*x)*x)/(c**2 + 2*c*d*x + d**2*x**2 - 1),x)*b**2*d**2 + 2*log(c + d*x - 1)*a*b + 3*a**2*c**2*d*x + 3*a**2*c*d**2*x**2 + a**2*d**3*x**3 + 2*a*b*c*d*x + a*b*d**2*x**2 + b**2*d*x))/(3*d)`

3.17 $\int (ce + dex)(a + b\operatorname{arctanh}(c + dx))^2 dx$

| | |
|---|-----|
| Optimal result | 175 |
| Mathematica [A] (verified) | 176 |
| Rubi [A] (verified) | 176 |
| Maple [B] (verified) | 178 |
| Fricas [B] (verification not implemented) | 179 |
| Sympy [B] (verification not implemented) | 180 |
| Maxima [B] (verification not implemented) | 180 |
| Giac [B] (verification not implemented) | 181 |
| Mupad [B] (verification not implemented) | 182 |
| Reduce [B] (verification not implemented) | 183 |

Optimal result

Integrand size = 21, antiderivative size = 95

$$\int (ce + dex)(a + b\operatorname{arctanh}(c + dx))^2 dx = abex + \frac{b^2e(c + dx)\operatorname{arctanh}(c + dx)}{d} - \frac{e(a + b\operatorname{arctanh}(c + dx))^2}{2d} + \frac{e(c + dx)^2(a + b\operatorname{arctanh}(c + dx))^2}{2d} + \frac{b^2e \log(1 - (c + dx)^2)}{2d}$$

output

```
a*b*e*x+b^2*e*(d*x+c)*arctanh(d*x+c)/d-1/2*e*(a+b*arctanh(d*x+c))^2/d+1/2*
e*(d*x+c)^2*(a+b*arctanh(d*x+c))^2/d+1/2*b^2*e*ln(1-(d*x+c)^2)/d
```


Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.41

$$\int (ce + dex)(a + b \operatorname{arctanh}(c + dx))^2 dx = e \left(\frac{ab(c + dx)}{d} + \frac{a^2(c + dx)^2}{2d} + \frac{b(c + dx)(b + a(c + dx)) \operatorname{arctanh}(c + dx)}{d} + \frac{(-b^2 + b^2(c + dx)^2) \operatorname{arctanh}(c + dx)^2}{2d} + \frac{(ab + b^2) \log(1 - c - dx)}{2d} + \frac{(-ab + b^2) \log(1 + c + dx)}{2d} \right)$$

input

```
Integrate[(c*e + d*e*x)*(a + b*ArcTanh[c + d*x])^2,x]
```

output

```
e*((a*b*(c + d*x))/d + (a^2*(c + d*x)^2)/(2*d) + (b*(c + d*x)*(b + a*(c + d*x))*ArcTanh[c + d*x])/d + ((-b^2 + b^2*(c + d*x)^2)*ArcTanh[c + d*x]^2)/(2*d) + ((a*b + b^2)*Log[1 - c - d*x])/(2*d) + ((-(a*b) + b^2)*Log[1 + c + d*x])/(2*d))
```

Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.96, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6657, 27, 6452, 6542, 2009, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ce + dex)(a + b \operatorname{arctanh}(c + dx))^2 dx$$

$$\downarrow 6657$$

$$\int \frac{e(c + dx)(a + b \operatorname{arctanh}(c + dx))^2 d(c + dx)}{d}$$

$$\downarrow 27$$

$$\frac{e \int (c + dx)(a + b \operatorname{arctanh}(c + dx))^2 d(c + dx)}{d}$$

↓ 6452

$$\frac{e \left(\frac{1}{2}(c + dx)^2(a + b \operatorname{arctanh}(c + dx))^2 - b \int \frac{(c+dx)^2(a+b \operatorname{arctanh}(c+dx))}{1-(c+dx)^2} d(c + dx) \right)}{d}$$

↓ 6542

$$\frac{e \left(\frac{1}{2}(c + dx)^2(a + b \operatorname{arctanh}(c + dx))^2 - b \left(\int \frac{a+b \operatorname{arctanh}(c+dx)}{1-(c+dx)^2} d(c + dx) - \int (a + b \operatorname{arctanh}(c + dx)) d(c + dx) \right) \right)}{d}$$

↓ 2009

$$\frac{e \left(\frac{1}{2}(c + dx)^2(a + b \operatorname{arctanh}(c + dx))^2 - b \left(\int \frac{a+b \operatorname{arctanh}(c+dx)}{1-(c+dx)^2} d(c + dx) - a(c + dx) - b(c + dx) \operatorname{arctanh}(c + dx) \right) \right)}{d}$$

↓ 6510

$$\frac{e \left(\frac{1}{2}(c + dx)^2(a + b \operatorname{arctanh}(c + dx))^2 - b \left(\frac{(a+b \operatorname{arctanh}(c+dx))^2}{2b} - a(c + dx) - b(c + dx) \operatorname{arctanh}(c + dx) - \frac{1}{2} b \log \right) \right)}{d}$$

input

```
Int[(c*e + d*e*x)*(a + b*ArcTanh[c + d*x])^2,x]
```

output

```
(e*(((c + d*x)^2*(a + b*ArcTanh[c + d*x])^2)/2 - b*(-(a*(c + d*x)) - b*(c + d*x)*ArcTanh[c + d*x] + (a + b*ArcTanh[c + d*x])^2/(2*b) - (b*Log[1 - (c + d*x)^2])/2))/d
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] :> Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6510 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)/((d_.) + (e_.)*(x_)^2), x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6542 `Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_.))/((d_.) + (e_.)*(x_)^2), x_Symbol] :> Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTanh[c*x])^p, x], x] - Simp[d*(f^2/e Int[(f*x)^(m - 2)*((a + b*ArcTanh[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

rule 6657 `Int[((a_.) + ArcTanh[(c_.) + (d_.)*(x_)])*(b_.))^(p_.)*((e_.) + (f_.)*(x_)^(m_.), x_Symbol] :> Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 220 vs. $2(89) = 178$.

Time = 0.31 (sec) , antiderivative size = 221, normalized size of antiderivative = 2.33

| method | result |
|------------------|---|
| derivativdivides | $\frac{a^2 e^{(dx+c)^2}}{2} + e b^2 \left(\frac{(dx+c)^2 \operatorname{arctanh}(dx+c)^2}{2} + (dx+c) \operatorname{arctanh}(dx+c) + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{2} - \frac{\operatorname{arctanh}(dx+c) \ln(dx+c+1)}{2} \right)$ |
| default | $\frac{a^2 e^{(dx+c)^2}}{2} + e b^2 \left(\frac{(dx+c)^2 \operatorname{arctanh}(dx+c)^2}{2} + (dx+c) \operatorname{arctanh}(dx+c) + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{2} - \frac{\operatorname{arctanh}(dx+c) \ln(dx+c+1)}{2} \right)$ |
| parts | $a^2 e \left(\frac{1}{2} d x^2 + c x \right) + \frac{e b^2 \left(\frac{(dx+c)^2 \operatorname{arctanh}(dx+c)^2}{2} + (dx+c) \operatorname{arctanh}(dx+c) + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{2} - \frac{\operatorname{arctanh}(dx+c) \ln(dx+c+1)}{2} \right)}{1}$ |
| parallelrisc | $\frac{b^2 d^3 e \operatorname{arctanh}(dx+c)^2 x^2 + 2x^2 \operatorname{arctanh}(dx+c) a b d^3 e + 2c e b^2 \operatorname{arctanh}(dx+c)^2 x d^2 + x^2 a^2 d^3 e + 4x \operatorname{arctanh}(dx+c) a b c d^2}{8d}$ |
| risc | $\frac{e b^2 (d^2 x^2 + 2cdx + c^2 - 1) \ln(dx+c+1)^2}{8d} + \frac{b e (-b d^2 x^2 \ln(-dx-c+1) + 2d^2 a x^2 - 2bdx \ln(-dx-c+1) c + 4adxc - \ln(-dx-c+1))}{4d}$ |

```
input int((d*e*x+c*e)*(a+b*arctanh(d*x+c))^2,x,method=_RETURNVERBOSE)
```

```
output 1/d*(1/2*a^2*e*(d*x+c)^2+e*b^2*(1/2*(d*x+c)^2*arctanh(d*x+c)^2+(d*x+c)*arctanh(d*x+c)+1/2*arctanh(d*x+c)*ln(d*x+c-1)-1/2*arctanh(d*x+c)*ln(d*x+c+1)-1/4*ln(d*x+c-1)*ln(1/2*d*x+1/2*c+1/2)+1/8*ln(d*x+c-1)^2+1/2*ln(d*x+c-1)+1/2*ln(d*x+c+1)+1/8*ln(d*x+c+1)^2-1/4*(ln(d*x+c+1)-ln(1/2*d*x+1/2*c+1/2))*ln(-1/2*d*x-1/2*c+1/2))+2*b*a*e*(1/2*(d*x+c)^2*arctanh(d*x+c)+1/2*d*x+1/2*c+1/4*ln(d*x+c-1)-1/4*ln(d*x+c+1)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 191 vs. 2(89) = 178.

Time = 0.09 (sec) , antiderivative size = 191, normalized size of antiderivative = 2.01

$$\int (ce + dex)(a + b \operatorname{arctanh}(c + dx))^2 dx$$

$$= \frac{4 a^2 d^2 e x^2 + 8 (a^2 c + a b) d e x + 4 (a b c^2 + b^2 c - a b + b^2) e \log (d x + c + 1) - 4 (a b c^2 + b^2 c - a b - b^2) e \log (d x + c - 1)}{8 d}$$

```
input integrate((d*e*x+c*e)*(a+b*arctanh(d*x+c))^2,x, algorithm="fricas")
```

output

```
1/8*(4*a^2*d^2*e*x^2 + 8*(a^2*c + a*b)*d*e*x + 4*(a*b*c^2 + b^2*c - a*b +
b^2)*e*log(d*x + c + 1) - 4*(a*b*c^2 + b^2*c - a*b - b^2)*e*log(d*x + c -
1) + (b^2*d^2*e*x^2 + 2*b^2*c*d*e*x + (b^2*c^2 - b^2)*e)*log(-(d*x + c + 1
)/(d*x + c - 1))^2 + 4*(a*b*d^2*e*x^2 + (2*a*b*c + b^2)*d*e*x)*log(-(d*x +
c + 1)/(d*x + c - 1)))/d
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 238 vs. $2(83) = 166$.

Time = 1.10 (sec) , antiderivative size = 238, normalized size of antiderivative = 2.51

$$\int (ce + dex)(a + b \operatorname{arctanh}(c + dx))^2 dx$$

$$= \begin{cases} a^2 c e x + \frac{a^2 d e x^2}{2} + \frac{a b c^2 e \operatorname{atanh}(c + dx)}{d} + 2 a b c e x \operatorname{atanh}(c + dx) + a b d e x^2 \operatorname{atanh}(c + dx) + a b e x - \frac{a b e \operatorname{atanh}(c + dx)}{d} \\ c e x (a + b \operatorname{atanh}(c))^2 \end{cases}$$

input

```
integrate((d*e*x+c*e)*(a+b*atanh(d*x+c))**2,x)
```

output

```
Piecewise((a**2*c*e*x + a**2*d*e*x**2/2 + a*b*c**2*e*atanh(c + d*x)/d + 2*
a*b*c*e*x*atanh(c + d*x) + a*b*d*e*x**2*atanh(c + d*x) + a*b*e*x - a*b*e*a
tanh(c + d*x)/d + b**2*c**2*e*atanh(c + d*x)**2/(2*d) + b**2*c*e*x*atanh(c
+ d*x)**2 + b**2*c*e*atanh(c + d*x)/d + b**2*d*e*x**2*atanh(c + d*x)**2/2
+ b**2*e*x*atanh(c + d*x) + b**2*e*log(c/d + x + 1/d)/d - b**2*e*atanh(c
+ d*x)**2/(2*d) - b**2*e*atanh(c + d*x)/d, Ne(d, 0)), (c*e*x*(a + b*atanh(
c))**2, True))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 316 vs. $2(89) = 178$.

Time = 0.23 (sec) , antiderivative size = 316, normalized size of antiderivative = 3.33

$$\int (ce + dex)(a + b \operatorname{arctanh}(c + dx))^2 dx = \frac{1}{2} a^2 dex^2 + \frac{1}{2} \left(2x^2 \operatorname{arctanh}(dx + c) + d \left(\frac{2x}{d^2} - \frac{(c^2 + 2c + 1) \log(dx + c + 1)}{d^3} + \frac{(c^2 - 2c + 1) \log(dx + c - 1)}{d^3} \right) \right) + a^2 cex + \frac{(2(dx + c) \operatorname{arctanh}(dx + c) + \log(-(dx + c)^2 + 1)) abce}{d} + \frac{(b^2 d^2 ex^2 + 2b^2 c dex + (c^2 e - e)b^2) \log(dx + c + 1)^2 + (b^2 d^2 ex^2 + 2b^2 c dex + (c^2 e - e)b^2) \log(-dx - c + 1)^2}{d}$$

input `integrate((d*e*x+c*e)*(a+b*arctanh(d*x+c))^2,x, algorithm="maxima")`

output `1/2*a^2*d*e*x^2 + 1/2*(2*x^2*arctanh(d*x + c) + d*(2*x/d^2 - (c^2 + 2*c + 1)*log(d*x + c + 1)/d^3 + (c^2 - 2*c + 1)*log(d*x + c - 1)/d^3))*a*b*d*e + a^2*c*e*x + (2*(d*x + c)*arctanh(d*x + c) + log(-(d*x + c)^2 + 1))*a*b*c*e/d + 1/8*((b^2*d^2*e*x^2 + 2*b^2*c*d*e*x + (c^2*e - e)*b^2)*log(d*x + c + 1)^2 + (b^2*d^2*e*x^2 + 2*b^2*c*d*e*x + (c^2*e - e)*b^2)*log(-d*x - c + 1)^2 + 4*(b^2*d*e*x + (c*e + e)*b^2)*log(d*x + c + 1) - 2*(2*b^2*d*e*x + 2*(c*e - e)*b^2 + (b^2*d^2*e*x^2 + 2*b^2*c*d*e*x + (c^2*e - e)*b^2)*log(d*x + c + 1))*log(-d*x - c + 1))/d`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 351 vs. 2(89) = 178.

Time = 0.14 (sec) , antiderivative size = 351, normalized size of antiderivative = 3.69

$$\int (ce + dex)(a + b \operatorname{arctanh}(c + dx))^2 dx = \frac{1}{4} \left(\frac{(dx + c + 1)b^2 e \log\left(-\frac{dx+c+1}{dx+c-1}\right)^2}{\left(\frac{(dx+c+1)^2 d^2}{(dx+c-1)^2} - \frac{2(dx+c+1)d^2}{dx+c-1} + d^2\right)(dx + c - 1)} - \frac{2b^2 e \log\left(-\frac{dx+c+1}{dx+c-1} + 1\right)}{d^2} + \frac{2b^2 e \log\left(-\frac{dx+c+1}{dx+c-1}\right)}{d^2} + \dots \right)$$

input `integrate((d*e*x+c*e)*(a+b*arctanh(d*x+c))^2,x, algorithm="giac")`

output

```

1/4*((d*x + c + 1)*b^2*e*log(-(d*x + c + 1)/(d*x + c - 1))^2/(((d*x + c +
1)^2*d^2/(d*x + c - 1)^2 - 2*(d*x + c + 1)*d^2/(d*x + c - 1) + d^2)*(d*x +
c - 1)) - 2*b^2*e*log(-(d*x + c + 1)/(d*x + c - 1) + 1)/d^2 + 2*b^2*e*log
(-(d*x + c + 1)/(d*x + c - 1))/d^2 + 2*(2*(d*x + c + 1)*a*b*e/(d*x + c - 1
) + (d*x + c + 1)*b^2*e/(d*x + c - 1) - b^2*e)*log(-(d*x + c + 1)/(d*x + c
- 1))/((d*x + c + 1)^2*d^2/(d*x + c - 1)^2 - 2*(d*x + c + 1)*d^2/(d*x + c
- 1) + d^2) + 4*((d*x + c + 1)*a^2*e/(d*x + c - 1) + (d*x + c + 1)*a*b*e/
(d*x + c - 1) - a*b*e)/((d*x + c + 1)^2*d^2/(d*x + c - 1)^2 - 2*(d*x + c +
1)*d^2/(d*x + c - 1) + d^2))*((c + 1)*d - (c - 1)*d)

```

Mupad [B] (verification not implemented)

Time = 0.82 (sec) , antiderivative size = 432, normalized size of antiderivative = 4.55

$$\begin{aligned}
& \int (ce + dex)(a + b \operatorname{arctanh}(c + dx))^2 dx \\
&= x (ae(b + 3ac) - 2a^2 ce) + \ln(1 - dx - c)^2 \left(\frac{b^2 cex}{4} - \frac{b^2 e - b^2 c^2 e}{8d} + \frac{b^2 dex^2}{8} \right) \\
&\quad - \ln(1 - dx - c) \left(\ln(c + dx + 1) \left(\frac{b^2 cex}{2} - \frac{\frac{b^2 e}{2} - \frac{b^2 c^2 e}{2}}{2d} + \frac{b^2 dex^2}{4} \right) \right. \\
&\quad\quad \left. - \frac{x(4b^2 d^2 e(c - 1) - 4b^2 de(d(c - 1) + d(c + 1)) + 8b^2 cd^2 e)}{16d^2} \right) \\
&\quad + \frac{x(8bd^2 e(4ac - 2a + bc) + 4bd^2 e(4a + b)(c + 1) - 4bde(d(c - 1) + d(c + 1))(4a + b))}{16d^2} \\
&\quad - \frac{b^2 dex^2}{8} + \frac{bdex^2(4a + b)}{8} \Big) + \ln(c + dx + 1)^2 \left(\frac{b^2 cex}{4} - \frac{b^2 e - b^2 c^2 e}{8d} + \frac{b^2 dex^2}{8} \right) \\
&\quad + \frac{\ln(c + dx + 1)(eb^2 c + eb^2 + aebc^2 - aeb)}{2d} \\
&\quad + \frac{\ln(c + dx - 1)(-eb^2 c + eb^2 - aebc^2 + aeb)}{2d} \\
&\quad + d \ln(c + dx + 1) \left(\frac{x(eb^2 + 2aceb)}{2d} + \frac{abex^2}{2} \right) + \frac{a^2 dex^2}{2}
\end{aligned}$$

input

```
int((c*e + d*e*x)*(a + b*atanh(c + d*x))^2,x)
```

output

```
x*(a*e*(b + 3*a*c) - 2*a^2*c*e) + log(1 - d*x - c)^2*((b^2*c*e*x)/4 - (b^2
*e - b^2*c^2*e)/(8*d) + (b^2*d*e*x^2)/8) - log(1 - d*x - c)*(log(c + d*x +
1)*((b^2*c*e*x)/2 - ((b^2*e)/2 - (b^2*c^2*e)/2)/(2*d) + (b^2*d*e*x^2)/4)
- (x*(4*b^2*d^2*e*(c - 1) - 4*b^2*d*e*(d*(c - 1) + d*(c + 1)) + 8*b^2*c*d^
2*e))/(16*d^2) + (x*(8*b*d^2*e*(4*a*c - 2*a + b*c) + 4*b*d^2*e*(4*a + b)*(
c + 1) - 4*b*d*e*(d*(c - 1) + d*(c + 1))*(4*a + b)))/(16*d^2) - (b^2*d*e*x
^2)/8 + (b*d*e*x^2*(4*a + b))/8) + log(c + d*x + 1)^2*((b^2*c*e*x)/4 - (b^
2*e - b^2*c^2*e)/(8*d) + (b^2*d*e*x^2)/8) + (log(c + d*x + 1)*(b^2*e - a*b
*e + b^2*c*e + a*b*c^2*e))/(2*d) + (log(c + d*x - 1)*(b^2*e + a*b*e - b^2*
c*e - a*b*c^2*e))/(2*d) + d*log(c + d*x + 1)*((x*(b^2*e + 2*a*b*c*e))/(2*d
) + (a*b*e*x^2)/2) + (a^2*d*e*x^2)/2
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 193, normalized size of antiderivative = 2.03

$$\int (ce + dex)(a + b \operatorname{arctanh}(c + dx))^2 dx$$

$$= \frac{e(\operatorname{atanh}(dx + c)^2 b^2 c^2 + 2 \operatorname{atanh}(dx + c)^2 b^2 c dx + \operatorname{atanh}(dx + c)^2 b^2 d^2 x^2 - \operatorname{atanh}(dx + c)^2 b^2 + 2 \operatorname{atanh}(c + dx) \operatorname{atanh}(dx + c) b^2 c dx + \operatorname{atanh}(c + dx) \operatorname{atanh}(dx + c) b^2 d^2 x^2 - \operatorname{atanh}(c + dx) \operatorname{atanh}(dx + c) b^2 + 2 \operatorname{atanh}(c + dx) a b c^2 + 4 \operatorname{atanh}(c + dx) a b c d x + 2 \operatorname{atanh}(c + dx) a b d^2 x^2 - 2 \operatorname{atanh}(c + dx) a b + 2 \operatorname{atanh}(c + dx) b^2 c + 2 \operatorname{atanh}(c + dx) b^2 d x + 2 \operatorname{atanh}(c + dx) b^2 + 2 \log(c + d x - 1) b^2 + 2 a^2 c d x + a^2 d^2 x^2 + 2 a b d x)}{2 d}$$

input

```
int((d*e*x+c*e)*(a+b*atanh(d*x+c))^2,x)
```

output

```
(e*(atanh(c + d*x)**2*b**2*c**2 + 2*atanh(c + d*x)**2*b**2*c*d*x + atanh(c
+ d*x)**2*b**2*d**2*x**2 - atanh(c + d*x)**2*b**2 + 2*atanh(c + d*x)*a*b*
c**2 + 4*atanh(c + d*x)*a*b*c*d*x + 2*atanh(c + d*x)*a*b*d**2*x**2 - 2*ata
nh(c + d*x)*a*b + 2*atanh(c + d*x)*b**2*c + 2*atanh(c + d*x)*b**2*d*x + 2*
atanh(c + d*x)*b**2 + 2*log(c + d*x - 1)*b**2 + 2*a**2*c*d*x + a**2*d**2*x
**2 + 2*a*b*d*x))/(2*d)
```


3.18 $\int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{ce+dex} dx$

| | |
|---|-----|
| Optimal result | 184 |
| Mathematica [C] (warning: unable to verify) | 185 |
| Rubi [A] (verified) | 185 |
| Maple [C] (warning: unable to verify) | 188 |
| Fricas [F] | 189 |
| Sympy [F] | 189 |
| Maxima [F] | 190 |
| Giac [F] | 190 |
| Mupad [F(-1)] | 190 |
| Reduce [F] | 191 |

Optimal result

Integrand size = 23, antiderivative size = 168

$$\int \frac{(a + b\operatorname{arctanh}(c + dx))^2}{ce + dex} dx = \frac{2(a + b\operatorname{arctanh}(c + dx))^2 \operatorname{arctanh}\left(1 - \frac{2}{1-c-dx}\right)}{de} - \frac{b(a + b\operatorname{arctanh}(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-c-dx}\right)}{de} + \frac{b(a + b\operatorname{arctanh}(c + dx)) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1-c-dx}\right)}{de} + \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-c-dx}\right)}{2de} - \frac{b^2 \operatorname{PolyLog}\left(3, -1 + \frac{2}{1-c-dx}\right)}{2de}$$

output

```
-2*(a+b*arctanh(d*x+c))^2*arctanh(-1+2/(-d*x-c+1))/d/e-b*(a+b*arctanh(d*x+c))*polylog(2,1-2/(-d*x-c+1))/d/e+b*(a+b*arctanh(d*x+c))*polylog(2,-1+2/(-d*x-c+1))/d/e+1/2*b^2*polylog(3,1-2/(-d*x-c+1))/d/e-1/2*b^2*polylog(3,-1+2/(-d*x-c+1))/d/e
```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.23 (sec) , antiderivative size = 424, normalized size of antiderivative = 2.52

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{ce + dex} dx = \frac{a^2 \log(c + dx)}{de} - \frac{2iab \left(\operatorname{iarctanh}(c + dx) \left(-\log\left(\frac{1}{\sqrt{1-(c+dx)^2}}\right) + \log\left(\frac{i(c+dx)}{\sqrt{1-(c+dx)^2}}\right) \right) + \frac{1}{2} \left(-\frac{1}{4}i(\pi - 2\operatorname{iarctanh}(c + dx))^2 + b^2(\operatorname{arctanh}(c + dx))^2 \log(1 - e^{-2\operatorname{arctanh}(c+dx)}) - \operatorname{arctanh}(c + dx)^2 \log(1 + e^{-2\operatorname{arctanh}(c+dx)}) + \operatorname{arctanh}(c + dx) \right) \right)}{d^2}$$

input `Integrate[(a + b*ArcTanh[c + d*x])^2/(c*e + d*e*x),x]`

output `(a^2*Log[c + d*x])/(d*e) - ((2*I)*a*b*(I*ArcTanh[c + d*x]*(-Log[1/Sqrt[1 - (c + d*x)^2]] + Log[(I*(c + d*x))/Sqrt[1 - (c + d*x)^2]]) + ((-1/4*I)*(Pi - (2*I)*ArcTanh[c + d*x])^2 + I*ArcTanh[c + d*x]^2 + (Pi - (2*I)*ArcTanh[c + d*x])*Log[1 - E^(I*(Pi - (2*I)*ArcTanh[c + d*x])]) + (2*I)*ArcTanh[c + d*x]*Log[1 - E^(-2*ArcTanh[c + d*x])] - (2*I)*ArcTanh[c + d*x]*Log[((2*I)*(c + d*x))/Sqrt[1 - (c + d*x)^2]] - (Pi - (2*I)*ArcTanh[c + d*x])*Log[2*Sin[(Pi - (2*I)*ArcTanh[c + d*x])/2]] - I*PolyLog[2, E^(I*(Pi - (2*I)*ArcTanh[c + d*x])]) - I*PolyLog[2, E^(-2*ArcTanh[c + d*x])])]/(2))/(d*e) + (b^2*(ArcTanh[c + d*x]^2*Log[1 - E^(-2*ArcTanh[c + d*x])] - ArcTanh[c + d*x]^2*Log[1 + E^(-2*ArcTanh[c + d*x])]) + ArcTanh[c + d*x]*PolyLog[2, -E^(-2*ArcTanh[c + d*x])] - ArcTanh[c + d*x]*PolyLog[2, E^(-2*ArcTanh[c + d*x])] + PolyLog[3, -E^(-2*ArcTanh[c + d*x])]/2 - PolyLog[3, E^(-2*ArcTanh[c + d*x])]/2))/(d*e)`

Rubi [A] (verified)

Time = 0.84 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {6657, 27, 6448, 6614, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{ce + dex} dx$$

↓ 6657

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{e(c + dx)} d(c + dx)$$

d

↓ 27

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{c + dx} d(c + dx)$$

de

↓ 6448

$$\frac{2 \operatorname{arctanh}\left(1 - \frac{2}{-c - dx + 1}\right) (a + b \operatorname{arctanh}(c + dx))^2 - 4b \int \frac{(a + b \operatorname{arctanh}(c + dx)) \operatorname{arctanh}\left(1 - \frac{2}{-c - dx + 1}\right)}{1 - (c + dx)^2} d(c + dx)}{de}$$

↓ 6614

$$\frac{2 \operatorname{arctanh}\left(1 - \frac{2}{-c - dx + 1}\right) (a + b \operatorname{arctanh}(c + dx))^2 - 4b \left(\frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(c + dx)) \log\left(2 - \frac{2}{-c - dx + 1}\right)}{1 - (c + dx)^2} d(c + dx) - \frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(c + dx))}{1 - (c + dx)^2} d(c + dx) \right)}{de}$$

↓ 6620

$$\frac{2 \operatorname{arctanh}\left(1 - \frac{2}{-c - dx + 1}\right) (a + b \operatorname{arctanh}(c + dx))^2 - 4b \left(\frac{1}{2} \left(\frac{1}{2} \operatorname{PolyLog}\left(2, 1 - \frac{2}{-c - dx + 1}\right) (a + b \operatorname{arctanh}(c + dx)) - \frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(c + dx))}{1 - (c + dx)^2} d(c + dx) \right) \right)}{de}$$

↓ 7164

$$\frac{2 \operatorname{arctanh}\left(1 - \frac{2}{-c - dx + 1}\right) (a + b \operatorname{arctanh}(c + dx))^2 - 4b \left(\frac{1}{2} \left(\frac{1}{2} \operatorname{PolyLog}\left(2, 1 - \frac{2}{-c - dx + 1}\right) (a + b \operatorname{arctanh}(c + dx)) - \frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(c + dx))}{1 - (c + dx)^2} d(c + dx) \right) \right)}{de}$$

input

`Int[(a + b*ArcTanh[c + d*x])^2/(c*e + d*e*x),x]`

output

```
(2*(a + b*ArcTanh[c + d*x])^2*ArcTanh[1 - 2/(1 - c - d*x)] - 4*b*(((a + b
*ArcTanh[c + d*x])*PolyLog[2, 1 - 2/(1 - c - d*x)])/2 - (b*PolyLog[3, 1 -
2/(1 - c - d*x)]/4)/2 + (-1/2*(a + b*ArcTanh[c + d*x])*PolyLog[2, -1 + 2
/(1 - c - d*x)]) + (b*PolyLog[3, -1 + 2/(1 - c - d*x)]/4)/2))/(d*e)
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

rule 6448

```
Int[((a_) + ArcTanh[(c_)*(x_)]*(b_))^(p_)/(x_), x_Symbol] := Simp[2*(a +
b*ArcTanh[c*x])^p*ArcTanh[1 - 2/(1 - c*x)], x] - Simp[2*b*c*p Int[(a + b
*ArcTanh[c*x])^(p - 1)*(ArcTanh[1 - 2/(1 - c*x)]/(1 - c^2*x^2)), x], x] /;
FreeQ[{a, b, c}, x] && IGtQ[p, 1]
```

rule 6614

```
Int[(ArcTanh[u_]*((a_) + ArcTanh[(c_)*(x_)]*(b_))^(p_))/((d_) + (e_)*(
x_)^2), x_Symbol] := Simp[1/2 Int[Log[1 + u]*((a + b*ArcTanh[c*x])^p/(d +
e*x^2)), x], x] - Simp[1/2 Int[Log[1 - u]*((a + b*ArcTanh[c*x])^p/(d + e
*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e,
0] && EqQ[u^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 6620

```
Int[(Log[u_]*((a_) + ArcTanh[(c_)*(x_)]*(b_))^(p_))/((d_) + (e_)*(x_)^
2), x_Symbol] := Simp[(-a + b*ArcTanh[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d))
, x] + Simp[b*(p/2 Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(
d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d
+ e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 6657

```
Int[((a_) + ArcTanh[(c_) + (d_)*(x_)]*(b_))^(p_)*((e_) + (f_)*(x_))^(
m_), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x
], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0]
&& IGtQ[p, 0]
```

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] :> With[{w = DerivativeDivides[v, u*v, x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 12.46 (sec) , antiderivative size = 705, normalized size of antiderivative = 4.20

| method | result |
|-------------------|---|
| derivativedivides | $\frac{a^2 \ln(dx+c)}{e} + \frac{b^2 \left(\ln(dx+c) \operatorname{arctanh}(dx+c)^2 - \operatorname{arctanh}(dx+c) \operatorname{polylog}\left(2, -\frac{(dx+c+1)^2}{1-(dx+c)^2}\right) + \frac{\operatorname{polylog}\left(3, -\frac{(dx+c+1)^2}{1-(dx+c)^2}\right)}{2} - \operatorname{arctanh}(dx+c) \right)}{e}$ |
| default | $\frac{a^2 \ln(dx+c)}{e} + \frac{b^2 \left(\ln(dx+c) \operatorname{arctanh}(dx+c)^2 - \operatorname{arctanh}(dx+c) \operatorname{polylog}\left(2, -\frac{(dx+c+1)^2}{1-(dx+c)^2}\right) + \frac{\operatorname{polylog}\left(3, -\frac{(dx+c+1)^2}{1-(dx+c)^2}\right)}{2} - \operatorname{arctanh}(dx+c) \right)}{e}$ |
| parts | $\frac{a^2 \ln(dx+c)}{ed} + \frac{b^2 \left(\ln(dx+c) \operatorname{arctanh}(dx+c)^2 - \operatorname{arctanh}(dx+c) \operatorname{polylog}\left(2, -\frac{(dx+c+1)^2}{1-(dx+c)^2}\right) + \frac{\operatorname{polylog}\left(3, -\frac{(dx+c+1)^2}{1-(dx+c)^2}\right)}{2} - \operatorname{arctanh}(dx+c) \right)}{ed}$ |

input

```
int((a+b*arctanh(d*x+c))^2/(d*e*x+c*e),x,method=_RETURNVERBOSE)
```

output

```

1/d*(a^2/e*ln(d*x+c)+b^2/e*(ln(d*x+c)*arctanh(d*x+c)^2-arctanh(d*x+c)*poly
log(2,-(d*x+c+1)^2/(1-(d*x+c)^2))+1/2*polylog(3,-(d*x+c+1)^2/(1-(d*x+c)^2)
)-arctanh(d*x+c)^2*ln((d*x+c+1)^2/(1-(d*x+c)^2)-1)+arctanh(d*x+c)^2*ln(1+(
d*x+c+1)/(1-(d*x+c)^2)^(1/2))+2*arctanh(d*x+c)*polylog(2,-(d*x+c+1)/(1-(d*
x+c)^2)^(1/2))-2*polylog(3,-(d*x+c+1)/(1-(d*x+c)^2)^(1/2))+arctanh(d*x+c)^
2*ln(1-(d*x+c+1)/(1-(d*x+c)^2)^(1/2))+2*arctanh(d*x+c)*polylog(2,(d*x+c+1)
/(1-(d*x+c)^2)^(1/2))-2*polylog(3,(d*x+c+1)/(1-(d*x+c)^2)^(1/2))+1/2*I*Pi*
csgn(I*(-(d*x+c+1)^2/((d*x+c)^2-1)-1)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))*(csgn
(I*(-(d*x+c+1)^2/((d*x+c)^2-1)-1))*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))-c
sgn(I*(-(d*x+c+1)^2/((d*x+c)^2-1)-1))*csgn(I*(-(d*x+c+1)^2/((d*x+c)^2-1)-1
)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))-csgn(I*(-(d*x+c+1)^2/((d*x+c)^2-1)-1)/(1-
(d*x+c+1)^2/((d*x+c)^2-1)))*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))+csgn(I*(
-(d*x+c+1)^2/((d*x+c)^2-1)-1)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))^2)*arctanh(d*
x+c)^2)+2*b*a/e*(ln(d*x+c)*arctanh(d*x+c)-1/2*dilog(d*x+c)-1/2*dilog(d*x+c
+1)-1/2*ln(d*x+c)*ln(d*x+c+1)))

```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{ce + dex} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^2}{dex + ce} dx$$

input

```
integrate((a+b*arctanh(d*x+c))^2/(d*e*x+c*e),x, algorithm="fricas")
```

output

```
integral((b^2*arctanh(d*x + c)^2 + 2*a*b*arctanh(d*x + c) + a^2)/(d*e*x +
c*e), x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{ce + dex} dx = \frac{\int \frac{a^2}{c+dx} dx + \int \frac{b^2 \operatorname{atanh}^2(c+dx)}{c+dx} dx + \int \frac{2ab \operatorname{atanh}(c+dx)}{c+dx} dx}{e}$$

input

```
integrate((a+b*atanh(d*x+c))**2/(d*e*x+c*e),x)
```

output $(\text{Integral}(a^{**2}/(c + d*x), x) + \text{Integral}(b^{**2}*\text{atanh}(c + d*x)**2/(c + d*x), x) + \text{Integral}(2*a*b*\text{atanh}(c + d*x)/(c + d*x), x))/e$

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{ce + dex} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^2}{dex + ce} dx$$

input `integrate((a+b*arctanh(d*x+c))^2/(d*e*x+c*e),x, algorithm="maxima")`

output $a^2*\log(d*e*x + c*e)/(d*e) + \text{integrate}(1/4*b^2*(\log(d*x + c + 1) - \log(-d*x - c + 1))^2/(d*e*x + c*e) + a*b*(\log(d*x + c + 1) - \log(-d*x - c + 1))/(d*e*x + c*e), x)$

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{ce + dex} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^2}{dex + ce} dx$$

input `integrate((a+b*arctanh(d*x+c))^2/(d*e*x+c*e),x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)^2/(d*e*x + c*e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{ce + dex} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^2}{ce + dex} dx$$

input `int((a + b*atanh(c + d*x))^2/(c*e + d*e*x),x)`

output `int((a + b*atanh(c + d*x))^2/(c*e + d*e*x), x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{ce + dex} dx$$

$$= \frac{2 \left(\int \frac{\operatorname{atanh}(dx+c)}{dx+c} dx \right) abd + \left(\int \frac{\operatorname{atanh}(dx+c)^2}{dx+c} dx \right) b^2 d + \log(dx + c) a^2}{de}$$

input `int((a+b*atanh(d*x+c))^2/(d*e*x+c*e), x)`

output `(2*int(atanh(c + d*x)/(c + d*x), x)*a*b*d + int(atanh(c + d*x)**2/(c + d*x), x)*b**2*d + log(c + d*x)*a**2)/(d*e)`

3.19 $\int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{(ce+dex)^2} dx$

| | |
|----------------------------|-----|
| Optimal result | 192 |
| Mathematica [A] (verified) | 192 |
| Rubi [A] (verified) | 193 |
| Maple [B] (verified) | 195 |
| Fricas [F] | 196 |
| Sympy [F] | 196 |
| Maxima [F] | 197 |
| Giac [F] | 197 |
| Mupad [F(-1)] | 197 |
| Reduce [F] | 198 |

Optimal result

Integrand size = 23, antiderivative size = 104

$$\int \frac{(a + b\operatorname{arctanh}(c + dx))^2}{(ce + dex)^2} dx = \frac{(a + b\operatorname{arctanh}(c + dx))^2}{de^2} - \frac{(a + b\operatorname{arctanh}(c + dx))^2}{de^2(c + dx)} + \frac{2b(a + b\operatorname{arctanh}(c + dx)) \log\left(2 - \frac{2}{1+c+dx}\right)}{de^2} - \frac{b^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1+c+dx}\right)}{de^2}$$

output

```
(a+b*arctanh(d*x+c))^2/d/e^2-(a+b*arctanh(d*x+c))^2/d/e^2/(d*x+c)+2*b*(a+b*arctanh(d*x+c))*ln(2-2/(d*x+c+1))/d/e^2-b^2*polylog(2,-1+2/(d*x+c+1))/d/e^2
```

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.21

$$\int \frac{(a + b\operatorname{arctanh}(c + dx))^2}{(ce + dex)^2} dx = \frac{b^2(-1 + c + dx)\operatorname{arctanh}(c + dx)^2 + 2b\operatorname{arctanh}(c + dx) (-a + b(c + dx) \log(1 - e^{-2\operatorname{arctanh}(c+dx)})) + a(-1 + c + dx)}{de^2(c + dx)}$$

input `Integrate[(a + b*ArcTanh[c + d*x])^2/(c*e + d*e*x)^2,x]`

output `(b^2*(-1 + c + d*x)*ArcTanh[c + d*x]^2 + 2*b*ArcTanh[c + d*x]*(-a + b*(c + d*x)*Log[1 - E^(-2*ArcTanh[c + d*x])]) + a*(-a + 2*b*(c + d*x)*Log[(c + d*x)/Sqrt[1 - (c + d*x)^2]]) - b^2*(c + d*x)*PolyLog[2, E^(-2*ArcTanh[c + d*x])])/(d*e^2*(c + d*x))`

Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.91, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {6657, 27, 6452, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^2} dx \\
 & \quad \downarrow \text{6657} \\
 & \frac{\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{e^2(c + dx)^2} d(c + dx)}{d} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(c + dx)^2} d(c + dx)}{de^2} \\
 & \quad \downarrow \text{6452} \\
 & \frac{2b \int \frac{a + b \operatorname{arctanh}(c + dx)}{(c + dx)(1 - (c + dx)^2)} d(c + dx) - \frac{(a + b \operatorname{arctanh}(c + dx))^2}{c + dx}}{de^2} \\
 & \quad \downarrow \text{6550} \\
 & \frac{2b \left(\int \frac{a + b \operatorname{arctanh}(c + dx)}{(c + dx)(c + dx + 1)} d(c + dx) + \frac{(a + b \operatorname{arctanh}(c + dx))^2}{2b} \right) - \frac{(a + b \operatorname{arctanh}(c + dx))^2}{c + dx}}{de^2} \\
 & \quad \downarrow \text{6494}
 \end{aligned}$$

$$2b \left(-b \int \frac{\log\left(2 - \frac{2}{c+dx+1}\right)}{1-(c+dx)^2} d(c+dx) + \frac{(a + \operatorname{arctanh}(c+dx))^2}{2b} + \log\left(2 - \frac{2}{c+dx+1}\right) (a + \operatorname{arctanh}(c+dx)) \right) - \frac{(a + \operatorname{arctanh}(c+dx))^2}{c+dx+1} \Bigg/ de^2$$

↓ 2897

$$2b \left(\frac{(a + \operatorname{arctanh}(c+dx))^2}{2b} + \log\left(2 - \frac{2}{c+dx+1}\right) (a + \operatorname{arctanh}(c+dx)) - \frac{1}{2} b \operatorname{PolyLog}\left(2, \frac{2}{c+dx+1} - 1\right) \right) - \frac{(a + \operatorname{arctanh}(c+dx))^2}{c+dx+1} \Bigg/ de^2$$

input `Int[(a + b*ArcTanh[c + d*x])^2/(c*e + d*e*x)^2,x]`

output `(-((a + b*ArcTanh[c + d*x])^2/(c + d*x)) + 2*b*((a + b*ArcTanh[c + d*x])^2/(2*b) + (a + b*ArcTanh[c + d*x])*Log[2 - 2/(1 + c + d*x)] - (b*PolyLog[2, -1 + 2/(1 + c + d*x)]/2))/(d*e^2)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2897 `Int[Log[u_]*(P_q_)^(m_), x_Symbol] := With[{C = FullSimplify[P_q^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[P_q, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[P_q, x]]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6494 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((x) \cdot (d + (e) \cdot x))$, x_Symbol] $\rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{Log}[2 - 2/(1 + e \cdot (x/d))]/d)$, x] - $\text{Simp}[b \cdot c \cdot (p/d) \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{Log}[2 - 2/(1 + e \cdot (x/d))]) / (1 - c^2 \cdot x^2)$, x], x] /; $\text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d^2 - e^2, 0]$

rule 6550 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((x) \cdot (d + (e) \cdot x^2))$, x_Symbol] $\rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot d \cdot (p+1))$, x] + $\text{Simp}[1/d \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p / (x \cdot (1 + c \cdot x))$, x], x] /; $\text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{GtQ}[p, 0]$

rule 6657 $\text{Int}[(a + \text{ArcTanh}[c + (d) \cdot x] \cdot b)^p \cdot ((e) + (f) \cdot x)^m$, x_Symbol] $\rightarrow \text{Simp}[1/d \text{Subst}[\text{Int}[(f \cdot (x/d))^m \cdot (a + b \cdot \text{ArcTanh}[x])^p$, x], x, c + d \cdot x], x] /; $\text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{EqQ}[d \cdot e - c \cdot f, 0] \ \&\& \ \text{IGtQ}[p, 0]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 251 vs. $2(104) = 208$.

Time = 1.15 (sec) , antiderivative size = 252, normalized size of antiderivative = 2.42

| method | result |
|-------------------|---|
| derivativedivides | $-\frac{a^2}{e^2(dx+c)} + \frac{b^2 \left(-\frac{\text{arctanh}(dx+c)^2}{dx+c} - \text{arctanh}(dx+c) \ln(dx+c-1) + 2 \ln(dx+c) \text{arctanh}(dx+c) - \text{arctanh}(dx+c) \ln(dx+c+1) + \text{dilog} \left(\frac{dx+c-1}{dx+c+1} \right) \right)}{e^2(dx+c)}$ |
| default | $-\frac{a^2}{e^2(dx+c)} + \frac{b^2 \left(-\frac{\text{arctanh}(dx+c)^2}{dx+c} - \text{arctanh}(dx+c) \ln(dx+c-1) + 2 \ln(dx+c) \text{arctanh}(dx+c) - \text{arctanh}(dx+c) \ln(dx+c+1) + \text{dilog} \left(\frac{dx+c-1}{dx+c+1} \right) \right)}{e^2(dx+c)}$ |
| parts | $-\frac{a^2}{e^2(dx+c)d} + \frac{b^2 \left(-\frac{\text{arctanh}(dx+c)^2}{dx+c} - \text{arctanh}(dx+c) \ln(dx+c-1) + 2 \ln(dx+c) \text{arctanh}(dx+c) - \text{arctanh}(dx+c) \ln(dx+c+1) + \text{dilog} \left(\frac{dx+c-1}{dx+c+1} \right) \right)}{e^2(dx+c)d}$ |

input $\text{int}((a+b \cdot \text{arctanh}(d \cdot x+c))^2 / (d \cdot e \cdot x+c \cdot e)^2, x, \text{method}=_RETURNVERBOSE)$

output

```
1/d*(-a^2/e^2/(d*x+c)+b^2/e^2*(-1/(d*x+c)*arctanh(d*x+c)^2-arctanh(d*x+c)*
ln(d*x+c-1)+2*ln(d*x+c)*arctanh(d*x+c)-arctanh(d*x+c)*ln(d*x+c+1)+dilog(1/
2*d*x+1/2*c+1/2)+1/2*ln(d*x+c-1)*ln(1/2*d*x+1/2*c+1/2)-1/4*ln(d*x+c-1)^2+1
/4*ln(d*x+c+1)^2-1/2*(ln(d*x+c+1)-ln(1/2*d*x+1/2*c+1/2))*ln(-1/2*d*x-1/2*c
+1/2)-dilog(d*x+c)-dilog(d*x+c+1)-ln(d*x+c)*ln(d*x+c+1))+2*b*a/e^2*(-1/(d*
x+c)*arctanh(d*x+c)-1/2*ln(d*x+c-1)+ln(d*x+c)-1/2*ln(d*x+c+1)))
```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^2} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^2}{(dex + ce)^2} dx$$

input

```
integrate((a+b*arctanh(d*x+c))^2/(d*e*x+c*e)^2,x, algorithm="fricas")
```

output

```
integral((b^2*arctanh(d*x + c)^2 + 2*a*b*arctanh(d*x + c) + a^2)/(d^2*e^2*
x^2 + 2*c*d*e^2*x + c^2*e^2), x)
```

Sympy [F]

$$\begin{aligned} & \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^2} dx \\ &= \frac{\int \frac{a^2}{c^2 + 2cdx + d^2x^2} dx + \int \frac{b^2 \operatorname{atanh}^2(c + dx)}{c^2 + 2cdx + d^2x^2} dx + \int \frac{2ab \operatorname{atanh}(c + dx)}{c^2 + 2cdx + d^2x^2} dx}{e^2} \end{aligned}$$

input

```
integrate((a+b*atanh(d*x+c))**2/(d*e*x+c*e)**2,x)
```

output

```
(Integral(a**2/(c**2 + 2*c*d*x + d**2*x**2), x) + Integral(b**2*atanh(c +
d*x)**2/(c**2 + 2*c*d*x + d**2*x**2), x) + Integral(2*a*b*atanh(c + d*x)/(
c**2 + 2*c*d*x + d**2*x**2), x))/e**2
```

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^2} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^2}{(dex + ce)^2} dx$$

input `integrate((a+b*arctanh(d*x+c))^2/(d*e*x+c*e)^2,x, algorithm="maxima")`

output `-(d*(log(d*x + c + 1)/(d^2*e^2) - 2*log(d*x + c)/(d^2*e^2) + log(d*x + c - 1)/(d^2*e^2)) + 2*arctanh(d*x + c)/(d^2*e^2*x + c*d*e^2))*a*b - 1/4*b^2*(log(-d*x - c + 1)^2/(d^2*e^2*x + c*d*e^2) + integrate(-((d*x + c - 1)*log(d*x + c + 1)^2 + 2*(d*x - (d*x + c - 1)*log(d*x + c + 1) + c)*log(-d*x - c + 1))/(d^3*e^2*x^3 + c^3*e^2 - c^2*e^2 + (3*c*d^2*e^2 - d^2*e^2)*x^2 + (3*c^2*d*e^2 - 2*c*d*e^2)*x), x)) - a^2/(d^2*e^2*x + c*d*e^2)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^2} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^2}{(dex + ce)^2} dx$$

input `integrate((a+b*arctanh(d*x+c))^2/(d*e*x+c*e)^2,x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)^2/(d*e*x + c*e)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^2} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^2}{(ce + dex)^2} dx$$

input `int((a + b*atanh(c + d*x))^2/(c*e + d*e*x)^2,x)`

output `int((a + b*atanh(c + d*x))^2/(c*e + d*e*x)^2, x)`

3.20 $\int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{(ce+dex)^3} dx$

| | |
|---|-----|
| Optimal result | 199 |
| Mathematica [A] (verified) | 199 |
| Rubi [A] (warning: unable to verify) | 200 |
| Maple [B] (verified) | 203 |
| Fricas [B] (verification not implemented) | 204 |
| Sympy [B] (verification not implemented) | 204 |
| Maxima [B] (verification not implemented) | 205 |
| Giac [B] (verification not implemented) | 206 |
| Mupad [B] (verification not implemented) | 207 |
| Reduce [B] (verification not implemented) | 208 |

Optimal result

Integrand size = 23, antiderivative size = 119

$$\int \frac{(a + b\operatorname{arctanh}(c + dx))^2}{(ce + dex)^3} dx = -\frac{b(a + b\operatorname{arctanh}(c + dx))}{de^3(c + dx)} + \frac{(a + b\operatorname{arctanh}(c + dx))^2}{2de^3} - \frac{(a + b\operatorname{arctanh}(c + dx))^2}{2de^3(c + dx)^2} + \frac{b^2 \log(c + dx)}{de^3} - \frac{b^2 \log(1 - (c + dx)^2)}{2de^3}$$

output

```
-b*(a+b*arctanh(d*x+c))/d/e^3/(d*x+c)+1/2*(a+b*arctanh(d*x+c))^2/d/e^3-1/2
*(a+b*arctanh(d*x+c))^2/d/e^3/(d*x+c)^2+b^2*ln(d*x+c)/d/e^3-1/2*b^2*ln(1-(
d*x+c)^2)/d/e^3
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.14

$$\int \frac{(a + b\operatorname{arctanh}(c + dx))^2}{(ce + dex)^3} dx = -\frac{a^2}{(c+dx)^2} - \frac{2ab}{c+dx} - \frac{2b(a+b(c+dx))\operatorname{arctanh}(c+dx)}{(c+dx)^2} + \frac{b^2(-1+c^2+2cdx+d^2x^2)\operatorname{arctanh}(c+dx)^2}{(c+dx)^2} - \frac{b(a+b)\log(1-c-dx)}{2de^3}$$

input `Integrate[(a + b*ArcTanh[c + d*x])^2/(c*e + d*e*x)^3,x]`

output
$$\frac{-(a^2/(c + d*x)^2) - (2*a*b)/(c + d*x) - (2*b*(a + b*(c + d*x))*ArcTanh[c + d*x])/(c + d*x)^2 + (b^2*(-1 + c^2 + 2*c*d*x + d^2*x^2)*ArcTanh[c + d*x]^2)/(c + d*x)^2 - b*(a + b)*Log[1 - c - d*x] + 2*b^2*Log[c + d*x] + (a - b)*b*Log[1 + c + d*x]}{(2*d*e^3)}$$

Rubi [A] (warning: unable to verify)

Time = 0.63 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.82, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.435$, Rules used = {6657, 27, 6452, 6544, 6452, 243, 47, 14, 16, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^3} dx \\ & \quad \downarrow 6657 \\ & \frac{\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{e^3(c + dx)^3} d(c + dx)}{d} \\ & \quad \downarrow 27 \\ & \frac{\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(c + dx)^3} d(c + dx)}{de^3} \\ & \quad \downarrow 6452 \\ & \frac{b \int \frac{a + b \operatorname{arctanh}(c + dx)}{(c + dx)^2(1 - (c + dx)^2)} d(c + dx) - \frac{(a + b \operatorname{arctanh}(c + dx))^2}{2(c + dx)^2}}{de^3} \\ & \quad \downarrow 6544 \\ & \frac{b \left(\int \frac{a + b \operatorname{arctanh}(c + dx)}{(c + dx)^2} d(c + dx) + \int \frac{a + b \operatorname{arctanh}(c + dx)}{1 - (c + dx)^2} d(c + dx) \right) - \frac{(a + b \operatorname{arctanh}(c + dx))^2}{2(c + dx)^2}}{de^3} \\ & \quad \downarrow 6452 \end{aligned}$$

$$\frac{b \left(\int \frac{a + b \operatorname{arctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) + b \int \frac{1}{(c+dx)(1-(c+dx)^2)} d(c+dx) - \frac{a + b \operatorname{arctanh}(c+dx)}{c+dx} \right) - \frac{(a + b \operatorname{arctanh}(c+dx))^2}{2(c+dx)^2}}{de^3}$$

↓ 243

$$\frac{b \left(\int \frac{a + b \operatorname{arctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) + \frac{1}{2} b \int \frac{1}{(-c-dx+1)(c+dx)^2} d(c+dx)^2 - \frac{a + b \operatorname{arctanh}(c+dx)}{c+dx} \right) - \frac{(a + b \operatorname{arctanh}(c+dx))^2}{2(c+dx)^2}}{de^3}$$

↓ 47

$$\frac{b \left(\int \frac{a + b \operatorname{arctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) + \frac{1}{2} b \left(\int \frac{1}{-c-dx+1} d(c+dx)^2 + \int \frac{1}{(c+dx)^2} d(c+dx)^2 \right) - \frac{a + b \operatorname{arctanh}(c+dx)}{c+dx} \right) - \frac{(a + b \operatorname{arctanh}(c+dx))^2}{2(c+dx)^2}}{de^3}$$

↓ 14

$$\frac{b \left(\int \frac{a + b \operatorname{arctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) + \frac{1}{2} b \left(\int \frac{1}{-c-dx+1} d(c+dx)^2 + \log((c+dx)^2) \right) - \frac{a + b \operatorname{arctanh}(c+dx)}{c+dx} \right) - \frac{(a + b \operatorname{arctanh}(c+dx))^2}{2(c+dx)^2}}{de^3}$$

↓ 16

$$\frac{b \left(\int \frac{a + b \operatorname{arctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) - \frac{a + b \operatorname{arctanh}(c+dx)}{c+dx} + \frac{1}{2} b (\log((c+dx)^2) - \log(-c-dx+1)) \right) - \frac{(a + b \operatorname{arctanh}(c+dx))^2}{2(c+dx)^2}}{de^3}$$

↓ 6510

$$\frac{b \left(\frac{(a + b \operatorname{arctanh}(c+dx))^2}{2b} - \frac{a + b \operatorname{arctanh}(c+dx)}{c+dx} + \frac{1}{2} b (\log((c+dx)^2) - \log(-c-dx+1)) \right) - \frac{(a + b \operatorname{arctanh}(c+dx))^2}{2(c+dx)^2}}{de^3}$$

input `Int[(a + b*ArcTanh[c + d*x])^2/(c*e + d*e*x)^3,x]`

output `(-1/2*(a + b*ArcTanh[c + d*x])^2/(c + d*x)^2 + b*(-((a + b*ArcTanh[c + d*x])/ (c + d*x)) + (a + b*ArcTanh[c + d*x])^2/(2*b) + (b*(-Log[1 - c - d*x] + Log[(c + d*x)^2])/2))/(d*e^3)`

Definitions of rubi rules used

- rule 14 $\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ ; FreeQ}[a, x]$
- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$
- rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] \text{ ; FreeQ}[b, x]$
- rule 47 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x]$
- rule 243 $\text{Int}[(x_)^{(m_)}*((a_) + (b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{((m - 1)/2)*(a + b*x)^p}, x], x, x^2], x] \text{ ; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m - 1)/2]$
- rule 6452 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_))^{(p_)}*(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m + 1)), x] - \text{Simp}[b*c*n*(p/(m + 1)) \text{ Int}[x^{(m + n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p - 1)/(1 - c^2*x^{(2*n)})}), x], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$
- rule 6510 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)]*(b_))^{(p_)} / ((d_) + (e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^{(p + 1)} / (b*c*d*(p + 1)), x] \text{ ; FreeQ}[\{a, b, c, d, e, p\}, x] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{NeQ}[p, -1]$
- rule 6544 $\text{Int}[(a_ + \text{ArcTanh}[(c_)*(x_)]*(b_))^{(p_)}*((f_)*(x_)^{(m_)}) / ((d_) + (e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[1/d \text{ Int}[(f*x)^m*(a + b*\text{ArcTanh}[c*x])^p, x], x] - \text{Simp}[e/(d*f^2) \text{ Int}[(f*x)^{(m + 2)}*((a + b*\text{ArcTanh}[c*x])^p/(d + e*x^2)), x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{LtQ}[m, -1]$

rule 6657

```
Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(
m_.), x_Symbol] :> Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x
], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0]
&& IGtQ[p, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 237 vs. 2(113) = 226.

Time = 0.90 (sec) , antiderivative size = 238, normalized size of antiderivative = 2.00

| method | result |
|------------------|--|
| derivativdivides | $-\frac{a^2}{2e^3(dx+c)^2} + \frac{b^2 \left(-\frac{\operatorname{arctanh}(dx+c)^2}{2(dx+c)^2} - \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{2} + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c+1)}{2} - \frac{\operatorname{arctanh}(dx+c)}{dx+c} + \frac{\ln(dx+c-1)}{dx+c} \right)}{2e^3(dx+c)^2}$ |
| default | $-\frac{a^2}{2e^3(dx+c)^2} + \frac{b^2 \left(-\frac{\operatorname{arctanh}(dx+c)^2}{2(dx+c)^2} - \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{2} + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c+1)}{2} - \frac{\operatorname{arctanh}(dx+c)}{dx+c} + \frac{\ln(dx+c-1)}{dx+c} \right)}{2e^3(dx+c)^2}$ |
| parts | $-\frac{a^2}{2e^3(dx+c)^2 d} + \frac{b^2 \left(-\frac{\operatorname{arctanh}(dx+c)^2}{2(dx+c)^2} - \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{2} + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c+1)}{2} - \frac{\operatorname{arctanh}(dx+c)}{dx+c} + \frac{\ln(dx+c-1)}{dx+c} \right)}{2e^3(dx+c)^2 d}$ |
| parallelrisc | $-\frac{2abc d^3 x + 2a^2 d^2 c + 3ab c^2 d^2 - ab d^4 x^2 - 8x \operatorname{arctanh}(dx+c) ab c^2 d^3 - 4x^2 \operatorname{arctanh}(dx+c) abc d^4 + 4 \operatorname{arctanh}(dx+c) b^2 c^3}{8e^3(dx+c)^2 d}$ |
| risc | $\frac{b^2(d^2 x^2 + 2cdx + c^2 - 1) \ln(dx+c+1)^2}{8e^3(dx+c)^2 d} - \frac{b(b d^2 x^2 \ln(-dx-c+1) + 2bdx \ln(-dx-c+1)c + \ln(-dx-c+1) b c^2 + 2bdx + 2bc)}{4e^3(dx+c)^2 d}$ |

```
input int((a+b*arctanh(d*x+c))^2/(d*e*x+c*e)^3,x,method=_RETURNVERBOSE)
```

```
output 1/d*(-1/2*a^2/e^3/(d*x+c)^2+b^2/e^3*(-1/2/(d*x+c)^2*arctanh(d*x+c)^2-1/2*a
rctanh(d*x+c)*ln(d*x+c-1)+1/2*arctanh(d*x+c)*ln(d*x+c+1)-1/(d*x+c)*arctanh
(d*x+c)+1/4*ln(d*x+c-1)*ln(1/2*d*x+1/2*c+1/2)-1/8*ln(d*x+c-1)^2-1/2*ln(d*x
+c-1)+ln(d*x+c)-1/2*ln(d*x+c+1)-1/8*ln(d*x+c+1)^2+1/4*(ln(d*x+c+1)-ln(1/2*
d*x+1/2*c+1/2))*ln(-1/2*d*x-1/2*c+1/2))+2*b*a/e^3*(-1/2/(d*x+c)^2*arctanh(
d*x+c)-1/4*ln(d*x+c-1)+1/4*ln(d*x+c+1)-1/2/(d*x+c)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 270 vs. 2(113) = 226.

Time = 0.12 (sec) , antiderivative size = 270, normalized size of antiderivative = 2.27

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^3} dx = \frac{8 abdx + 8 abc - (b^2 d^2 x^2 + 2 b^2 c dx + b^2 c^2 - b^2) \log\left(-\frac{dx+c+1}{dx+c-1}\right)^2 + 4 a^2 - 4((ab - b^2)d^2 x^2 + 2(ab - b^2)dx + b^2 c^2)}{(d^3 e^3 x^2 + 2 c d^2 e^3 x + c^2 d e^3)}$$

input `integrate((a+b*arctanh(d*x+c))^2/(d*e*x+c*e)^3,x, algorithm="fricas")`

output `-1/8*(8*a*b*d*x + 8*a*b*c - (b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2 - b^2)*log(-(d*x + c + 1)/(d*x + c - 1))^2 + 4*a^2 - 4*((a*b - b^2)*d^2*x^2 + 2*(a*b - b^2)*c*d*x + (a*b - b^2)*c^2)*log(d*x + c + 1) - 8*(b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2)*log(d*x + c) + 4*((a*b + b^2)*d^2*x^2 + 2*(a*b + b^2)*c*d*x + (a*b + b^2)*c^2)*log(d*x + c - 1) + 4*(b^2*d*x + b^2*c + a*b)*log(-(d*x + c + 1)/(d*x + c - 1)))/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1102 vs. 2(100) = 200.

Time = 1.54 (sec) , antiderivative size = 1102, normalized size of antiderivative = 9.26

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^3} dx = \text{Too large to display}$$

input `integrate((a+b*atanh(d*x+c))**2/(d*e*x+c*e)**3,x)`

output

```
Piecewise((-a**2/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) + 2*
a*b*c**2*atanh(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**
2) + 4*a*b*c*d*x*atanh(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*
e**3*x**2) - 2*a*b*c/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2)
+ 2*a*b*d**2*x**2*atanh(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3
*e**3*x**2) - 2*a*b*d*x/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**
2) - 2*a*b*atanh(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x
**2) + 2*b**2*c**2*log(c/d + x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*
e**3*x**2) - 2*b**2*c**2*log(c/d + x + 1/d)/(2*c**2*d*e**3 + 4*c*d**2*e**3
*x + 2*d**3*e**3*x**2) + b**2*c**2*atanh(c + d*x)**2/(2*c**2*d*e**3 + 4*c*
d**2*e**3*x + 2*d**3*e**3*x**2) + 2*b**2*c**2*atanh(c + d*x)/(2*c**2*d*e**
3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) + 4*b**2*c*d*x*log(c/d + x)/(2*c**
2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) - 4*b**2*c*d*x*log(c/d + x
+ 1/d)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) + 2*b**2*c*d*x
*atanh(c + d*x)**2/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) +
4*b**2*c*d*x*atanh(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3
*x**2) - 2*b**2*c*atanh(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3
*e**3*x**2) + 2*b**2*d**2*x**2*log(c/d + x)/(2*c**2*d*e**3 + 4*c*d**2*e**3
*x + 2*d**3*e**3*x**2) - 2*b**2*d**2*x**2*log(c/d + x + 1/d)/(2*c**2*d*e**
3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) + b**2*d**2*x**2*atanh(c + d*x)...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 329 vs. $2(113) = 226$.

Time = 0.04 (sec) , antiderivative size = 329, normalized size of antiderivative = 2.76

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^3} dx =$$

$$-\frac{1}{2} \left(d \left(\frac{2}{d^3 e^3 x + cd^2 e^3} - \frac{\log(dx + c + 1)}{d^2 e^3} + \frac{\log(dx + c - 1)}{d^2 e^3} \right) + \frac{2 \operatorname{artanh}(dx + c)}{d^3 e^3 x^2 + 2cd^2 e^3 x + c^2 de^3} \right) ab$$

$$-\frac{1}{8} \left(d^2 \left(\frac{\log(dx + c + 1)^2 - 2 \log(dx + c + 1) \log(dx + c - 1) + \log(dx + c - 1)^2 + 4 \log(dx + c - 1)}{d^3 e^3} \right) \right.$$

$$\left. - \frac{b^2 \operatorname{artanh}(dx + c)^2}{2(d^3 e^3 x^2 + 2cd^2 e^3 x + c^2 de^3)} - \frac{a^2}{2(d^3 e^3 x^2 + 2cd^2 e^3 x + c^2 de^3)} \right)$$

input

```
integrate((a+b*arctanh(d*x+c))^2/(d*e*x+c*e)^3,x, algorithm="maxima")
```

output

```
-1/2*(d*(2/(d^3*e^3*x + c*d^2*e^3) - log(d*x + c + 1)/(d^2*e^3) + log(d*x
+ c - 1)/(d^2*e^3)) + 2*arctanh(d*x + c)/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^
2*d*e^3))*a*b - 1/8*(d^2*((log(d*x + c + 1)^2 - 2*log(d*x + c + 1)*log(d*x
+ c - 1) + log(d*x + c - 1)^2 + 4*log(d*x + c - 1)))/(d^3*e^3) + 4*log(d*x
+ c + 1)/(d^3*e^3) - 8*log(d*x + c)/(d^3*e^3)) + 4*d*(2/(d^3*e^3*x + c*d^
2*e^3) - log(d*x + c + 1)/(d^2*e^3) + log(d*x + c - 1)/(d^2*e^3))*arctanh(
d*x + c))*b^2 - 1/2*b^2*arctanh(d*x + c)^2/(d^3*e^3*x^2 + 2*c*d^2*e^3*x +
c^2*d*e^3) - 1/2*a^2/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 375 vs. $2(113) = 226$.

Time = 0.14 (sec) , antiderivative size = 375, normalized size of antiderivative = 3.15

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^3} dx$$

$$= \frac{1}{4} \left(\frac{(dx + c + 1)b^2 \log\left(-\frac{dx+c+1}{dx+c-1}\right)^2}{\left(\frac{(dx+c+1)^2 d^2 e^3}{(dx+c-1)^2} + \frac{2(dx+c+1)d^2 e^3}{dx+c-1} + d^2 e^3\right)} (dx + c - 1) + \frac{2 \left(\frac{2(dx+c+1)ab}{dx+c-1} + \frac{(dx+c+1)b^2}{dx+c-1} + b^2\right) \log\left(-\frac{dx+c+1}{dx+c-1}\right)}{\frac{(dx+c+1)^2 d^2 e^3}{(dx+c-1)^2} + \frac{2(dx+c+1)d^2 e^3}{dx+c-1} + d^2 e^3} \right)$$

input

```
integrate((a+b*arctanh(d*x+c))^2/(d*e*x+c*e)^3,x, algorithm="giac")
```

output

```
1/4*((d*x + c + 1)*b^2*log(-(d*x + c + 1)/(d*x + c - 1))^2/(((d*x + c + 1)
^2*d^2*e^3/(d*x + c - 1)^2 + 2*(d*x + c + 1)*d^2*e^3/(d*x + c - 1) + d^2*e
^3)*(d*x + c - 1)) + 2*(2*(d*x + c + 1)*a*b/(d*x + c - 1) + (d*x + c + 1)*
b^2/(d*x + c - 1) + b^2)*log(-(d*x + c + 1)/(d*x + c - 1))/((d*x + c + 1)^
2*d^2*e^3/(d*x + c - 1)^2 + 2*(d*x + c + 1)*d^2*e^3/(d*x + c - 1) + d^2*e^
3) + 4*((d*x + c + 1)*a^2/(d*x + c - 1) + (d*x + c + 1)*a*b/(d*x + c - 1)
+ a*b)/(((d*x + c + 1)^2*d^2*e^3/(d*x + c - 1)^2 + 2*(d*x + c + 1)*d^2*e^3/
(d*x + c - 1) + d^2*e^3) + 2*b^2*log(-(d*x + c + 1)/(d*x + c - 1) - 1)/(d^
2*e^3) - 2*b^2*log(-(d*x + c + 1)/(d*x + c - 1))/(d^2*e^3))*((c + 1)*d - (
c - 1)*d)
```

Mupad [B] (verification not implemented)

Time = 5.74 (sec) , antiderivative size = 776, normalized size of antiderivative = 6.52

$$\begin{aligned}
& \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^3} dx \\
&= \ln(1 - dx - c)^2 \left(\frac{b^2}{8de^3} - \frac{b^2}{2d(4c^2e^3 + 8cde^3x + 4d^2e^3x^2)} \right) \\
&+ \ln(c + dx + 1)^2 \left(\frac{b^2}{8de^3} - \frac{b^2}{8d^2e^3(2cx + dx^2 + \frac{c^2}{d})} \right) + \ln(1 - dx - c) \left(\ln(c + dx \right. \\
&+ 1) \left(\frac{b^2}{2d(2c^2e^3 + 4cde^3x + 2d^2e^3x^2)} - \frac{b^2(c^2 + 2cdx + d^2x^2)}{2d(2c^2e^3 + 4cde^3x + 2d^2e^3x^2)} \right) \\
&+ \frac{b^2}{2d(4c^2e^3 + 8cde^3x + 4d^2e^3x^2)} + \frac{b(4a - b)}{2d(4c^2e^3 + 8cde^3x + 4d^2e^3x^2)} \\
&- \frac{b^2(x(4cd - d + d(2c - 1)) - c + c^2 + c(2c - 1) + 3d^2x^2 + 1)}{2d(4c^2e^3 + 8cde^3x + 4d^2e^3x^2)} \\
&+ \frac{b^2(x(2de^3 + d(4ce^3 + 2e^3) + 8cde^3) + 2ce^3 + 2e^3 + c(4ce^3 + 2e^3) + 2c^2e^3 + 6d^2e^3x^2)}{4de^3(4c^2e^3 + 8cde^3x + 4d^2e^3x^2)} \\
&- \frac{\frac{a^2 + 2bca}{2d} + abx}{c^2e^3 + 2cde^3x + d^2e^3x^2} \\
&\ln(c + dx + 1) \left(x \left(\frac{2b^2c + b^2}{4de^3} + \frac{b^2c}{4de^3} - \frac{b^2(3c-1)}{4de^3} \right) + \frac{b^2c^2 + b^2c + b^2 + 4ab}{8d^2e^3} - \frac{b^2 \left(\frac{c^2 - c + 1}{2d} + \frac{c(2c-1)}{2d} \right)}{4de^3} + \frac{c(2b^2c + b^2)}{8d^2e^3} \right) \\
&- \frac{2cx + dx^2 + \frac{c^2}{d}}{2de^3} \\
&+ \frac{b^2 \ln(c + dx)}{de^3} - \frac{\ln(c + dx - 1)(b^2 + ab)}{2de^3} + \frac{\ln(c + dx + 1)(ab - b^2)}{2de^3}
\end{aligned}$$

input `int((a + b*atanh(c + d*x))^2/(c*e + d*e*x)^3,x)`

output

```

log(1 - d*x - c)^2*(b^2/(8*d*e^3) - b^2/(2*d*(4*c^2*e^3 + 4*d^2*e^3*x^2 +
8*c*d*e^3*x))) + log(c + d*x + 1)^2*(b^2/(8*d*e^3) - b^2/(8*d^2*e^3*(2*c*x
+ d*x^2 + c^2/d))) + log(1 - d*x - c)*(log(c + d*x + 1)*(b^2/(2*d*(2*c^2*
e^3 + 2*d^2*e^3*x^2 + 4*c*d*e^3*x)) - (b^2*(c^2 + d^2*x^2 + 2*c*d*x))/(2*d
*(2*c^2*e^3 + 2*d^2*e^3*x^2 + 4*c*d*e^3*x))) + b^2/(2*d*(4*c^2*e^3 + 4*d^2
*e^3*x^2 + 8*c*d*e^3*x)) + (b*(4*a - b))/(2*d*(4*c^2*e^3 + 4*d^2*e^3*x^2 +
8*c*d*e^3*x)) - (b^2*(x*(4*c*d - d + d*(2*c - 1)) - c + c^2 + c*(2*c - 1)
+ 3*d^2*x^2 + 1))/(2*d*(4*c^2*e^3 + 4*d^2*e^3*x^2 + 8*c*d*e^3*x)) + (b^2*
(x*(2*d*e^3 + d*(4*c*e^3 + 2*e^3) + 8*c*d*e^3) + 2*c*e^3 + 2*e^3 + c*(4*c*
e^3 + 2*e^3) + 2*c^2*e^3 + 6*d^2*e^3*x^2))/(4*d*e^3*(4*c^2*e^3 + 4*d^2*e^3
*x^2 + 8*c*d*e^3*x)) - ((a^2 + 2*a*b*c)/(2*d) + a*b*x)/(c^2*e^3 + d^2*e^3
*x^2 + 2*c*d*e^3*x) - (log(c + d*x + 1)*(x*((2*b^2*c + b^2)/(4*d*e^3) + (b
^2*c)/(4*d*e^3) - (b^2*(3*c - 1))/(4*d*e^3)) + (4*a*b + b^2*c + b^2 + b^2*
c^2)/(8*d^2*e^3) - (b^2*((c^2 - c + 1)/(2*d) + (c*(2*c - 1))/(2*d)))/(4*d*
e^3) + (c*(2*b^2*c + b^2))/(8*d^2*e^3)))/(2*c*x + d*x^2 + c^2/d) + (b^2*log
(c + d*x))/(d*e^3) - (log(c + d*x - 1)*(a*b + b^2))/(2*d*e^3) + (log(c +
d*x + 1)*(a*b - b^2))/(2*d*e^3)

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 503, normalized size of antiderivative = 4.23

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^3} dx$$

$$= \frac{-2abc^2 + 2abd^2x^2 - 2a^2c - 2 \log(dx + c + 1) b^2c d^2x^2 + 4 \log(dx + c) b^2c d^2x^2 + 2 \operatorname{atanh}(dx + c)^2 b^2c^3 - \dots}{(ce + dex)^3}$$

input

```
int((a+b*atanh(d*x+c))^2/(d*e*x+c*e)^3,x)
```

output

```
(2*atanh(c + d*x)**2*b**2*c**3 + 4*atanh(c + d*x)**2*b**2*c**2*d*x + 2*atanh(c + d*x)**2*b**2*c*d**2*x**2 - 2*atanh(c + d*x)**2*b**2*c - 4*atanh(c + d*x)*a*b*c - 2*atanh(c + d*x)*b**2*c**2 + 2*atanh(c + d*x)*b**2*d**2*x**2 - 2*log(c + d*x - 1)*a*b*c**3 - 4*log(c + d*x - 1)*a*b*c**2*d*x - 2*log(c + d*x - 1)*a*b*c*d**2*x**2 - 2*log(c + d*x - 1)*b**2*c**3 - 4*log(c + d*x - 1)*b**2*c**2*d*x + log(c + d*x - 1)*b**2*c**2 - 2*log(c + d*x - 1)*b**2*c*d**2*x**2 + 2*log(c + d*x - 1)*b**2*c*d*x + log(c + d*x - 1)*b**2*d**2*x**2 + 2*log(c + d*x + 1)*a*b*c**3 + 4*log(c + d*x + 1)*a*b*c**2*d*x + 2*log(c + d*x + 1)*a*b*c*d**2*x**2 - 2*log(c + d*x + 1)*b**2*c**3 - 4*log(c + d*x + 1)*b**2*c**2*d*x - log(c + d*x + 1)*b**2*c**2 - 2*log(c + d*x + 1)*b**2*c*d**2*x**2 - 2*log(c + d*x + 1)*b**2*c*d*x - log(c + d*x + 1)*b**2*d**2*x**2 + 4*log(c + d*x)*b**2*c**3 + 8*log(c + d*x)*b**2*c**2*d*x + 4*log(c + d*x)*b**2*c*d**2*x**2 - 2*a**2*c - 2*a*b*c**2 + 2*a*b*d**2*x**2)/(4*c*d*e**3*(c**2 + 2*c*d*x + d**2*x**2))
```

$$3.21 \quad \int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{(ce+dex)^4} dx$$

| | |
|----------------------------|-----|
| Optimal result | 210 |
| Mathematica [A] (verified) | 211 |
| Rubi [A] (verified) | 211 |
| Maple [A] (verified) | 214 |
| Fricas [F] | 215 |
| Sympy [F] | 215 |
| Maxima [F] | 216 |
| Giac [F] | 216 |
| Mupad [F(-1)] | 217 |
| Reduce [F] | 217 |

Optimal result

Integrand size = 23, antiderivative size = 180

$$\begin{aligned} \int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{(ce+dex)^4} dx = & -\frac{b^2}{3de^4(c+dx)} + \frac{b^2\operatorname{arctanh}(c+dx)}{3de^4} \\ & - \frac{b(a+b\operatorname{arctanh}(c+dx))}{3de^4(c+dx)^2} \\ & + \frac{(a+b\operatorname{arctanh}(c+dx))^2}{3de^4} - \frac{(a+b\operatorname{arctanh}(c+dx))^2}{3de^4(c+dx)^3} \\ & + \frac{2b(a+b\operatorname{arctanh}(c+dx)) \log\left(2 - \frac{2}{1+c+dx}\right)}{3de^4} \\ & - \frac{b^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1+c+dx}\right)}{3de^4} \end{aligned}$$

output

```
-1/3*b^2/d/e^4/(d*x+c)+1/3*b^2*arctanh(d*x+c)/d/e^4-1/3*b*(a+b*arctanh(d*x+c))/d/e^4/(d*x+c)^2+1/3*(a+b*arctanh(d*x+c))^2/d/e^4-1/3*(a+b*arctanh(d*x+c))^2/d/e^4/(d*x+c)^3+2/3*b*(a+b*arctanh(d*x+c))*ln(2-2/(d*x+c+1))/d/e^4-1/3*b^2*polylog(2,-1+2/(d*x+c+1))/d/e^4
```

Mathematica [A] (verified)

Time = 0.41 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.21

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^4} dx = \frac{a^2 - ab \left(-2 \operatorname{arctanh}(c + dx) + (c + dx) \left(-1 + c^2 + 2cdx + d^2x^2 + 2(c + dx)^2 \log \left(\frac{c+dx}{\sqrt{1-(c+dx)^2}} \right) \right) \right)}{d^4}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])^2/(c*e + d*e*x)^4,x]
```

output

```
-1/3*(a^2 - a*b*(-2*ArcTanh[c + d*x] + (c + d*x)*(-1 + c^2 + 2*c*d*x + d^2*x^2 + 2*(c + d*x)^2*Log[(c + d*x)/Sqrt[1 - (c + d*x)^2]])) + b^2*((c + d*x)^2 + (c + d*x)^2*ArcTanh[c + d*x]^2 + (1 - (c + d*x)^2)*ArcTanh[c + d*x]^2 + (c + d*x)*ArcTanh[c + d*x]*(1 - (c + d*x)^2 - (c + d*x)^2*ArcTanh[c + d*x] - 2*(c + d*x)^2*Log[1 - E^(-2*ArcTanh[c + d*x])])) + (c + d*x)^3*PolyLog[2, E^(-2*ArcTanh[c + d*x])])/(d*e^4*(c + d*x)^3)
```

Rubi [A] (verified)

Time = 0.92 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.78, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.435$, Rules used = {6657, 27, 6452, 6544, 6452, 264, 219, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^4} dx \\ & \quad \downarrow \text{6657} \\ & \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{e^4(c + dx)^4} d(c + dx) \\ & \quad \downarrow \text{27} \\ & \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(c + dx)^4} d(c + dx) \\ & \quad \downarrow \\ & \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{de^4} d(c + dx) \end{aligned}$$

$$\begin{aligned}
& \downarrow 6452 \\
& \frac{\frac{2}{3}b \int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)^3(1-(c+dx)^2)} d(c+dx) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{3(c+dx)^3}}{de^4} \\
& \downarrow 6544 \\
& \frac{\frac{2}{3}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)^3} d(c+dx) + \int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)(1-(c+dx)^2)} d(c+dx) \right) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{3(c+dx)^3}}{de^4} \\
& \downarrow 6452 \\
& \frac{\frac{2}{3}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)(1-(c+dx)^2)} d(c+dx) + \frac{1}{2}b \int \frac{1}{(c+dx)^2(1-(c+dx)^2)} d(c+dx) - \frac{a+\operatorname{barctanh}(c+dx)}{2(c+dx)^2} \right) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{3(c+dx)^3}}{de^4} \\
& \downarrow 264 \\
& \frac{\frac{2}{3}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)(1-(c+dx)^2)} d(c+dx) + \frac{1}{2}b \left(\int \frac{1}{1-(c+dx)^2} d(c+dx) - \frac{1}{c+dx} \right) - \frac{a+\operatorname{barctanh}(c+dx)}{2(c+dx)^2} \right) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{3(c+dx)^3}}{de^4} \\
& \downarrow 219 \\
& \frac{\frac{2}{3}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)(1-(c+dx)^2)} d(c+dx) - \frac{a+\operatorname{barctanh}(c+dx)}{2(c+dx)^2} + \frac{1}{2}b \left(\operatorname{arctanh}(c+dx) - \frac{1}{c+dx} \right) \right) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{3(c+dx)^3}}{de^4} \\
& \downarrow 6550 \\
& \frac{\frac{2}{3}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)(c+dx+1)} d(c+dx) + \frac{(a+\operatorname{barctanh}(c+dx))^2}{2b} - \frac{a+\operatorname{barctanh}(c+dx)}{2(c+dx)^2} + \frac{1}{2}b \left(\operatorname{arctanh}(c+dx) - \frac{1}{c+dx} \right) \right) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{3(c+dx)^3}}{de^4} \\
& \downarrow 6494 \\
& \frac{\frac{2}{3}b \left(-b \int \frac{\log\left(2-\frac{2}{c+dx+1}\right)}{1-(c+dx)^2} d(c+dx) + \frac{(a+\operatorname{barctanh}(c+dx))^2}{2b} - \frac{a+\operatorname{barctanh}(c+dx)}{2(c+dx)^2} + \log\left(2-\frac{2}{c+dx+1}\right) (a+\operatorname{barctanh}(c+dx)) \right) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{3(c+dx)^3}}{de^4} \\
& \downarrow 2897 \\
& \frac{\frac{2}{3}b \left(\frac{(a+\operatorname{barctanh}(c+dx))^2}{2b} - \frac{a+\operatorname{barctanh}(c+dx)}{2(c+dx)^2} + \log\left(2-\frac{2}{c+dx+1}\right) (a+\operatorname{barctanh}(c+dx)) + \frac{1}{2}b \left(\operatorname{arctanh}(c+dx) - \frac{1}{c+dx} \right) \right) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{3(c+dx)^3}}{de^4}
\end{aligned}$$

input `Int[(a + b*ArcTanh[c + d*x])^2/(c*e + d*e*x)^4,x]`

output `(-1/3*(a + b*ArcTanh[c + d*x])^2/(c + d*x)^3 + (2*b*((b*(-(c + d*x)^(-1) + ArcTanh[c + d*x]))/2 - (a + b*ArcTanh[c + d*x])/(2*(c + d*x)^2) + (a + b*ArcTanh[c + d*x])^2/(2*b) + (a + b*ArcTanh[c + d*x])*Log[2 - 2/(1 + c + d*x)] - (b*PolyLog[2, -1 + 2/(1 + c + d*x)]/2))/3)/(d*e^4)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 264 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2897 `Int[Log[u_]*(Pq_)^(m_.), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

```
rule 6494 Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_))), x
_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] -
Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]]
/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c
^2*d^2 - e^2, 0]
```

```
rule 6544 Int((((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*((f_.)*(x_))^(m_))/((d_) + (
e_.)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x
], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x
^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

```
rule 6550 Int((((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_)^2)),
x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/
d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]
```

```
rule 6657 Int((((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x
], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0]
&& IGtQ[p, 0]
```

Maple [A] (verified)

Time = 2.25 (sec) , antiderivative size = 307, normalized size of antiderivative = 1.71

| method | result |
|-------------------|--|
| derivativedivides | $-\frac{a^2}{3e^4(dx+c)^3} + \frac{b^2}{3(dx+c)^3} \left(-\frac{\operatorname{arctanh}(dx+c)^2}{3} - \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{3} - \frac{\operatorname{arctanh}(dx+c)}{3(dx+c)^2} + \frac{2 \ln(dx+c) \operatorname{arctanh}(dx+c)}{3} - \frac{\operatorname{arctanh}(dx+c)}{3} \right)$ |
| default | $-\frac{a^2}{3e^4(dx+c)^3} + \frac{b^2}{3(dx+c)^3} \left(-\frac{\operatorname{arctanh}(dx+c)^2}{3} - \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{3} - \frac{\operatorname{arctanh}(dx+c)}{3(dx+c)^2} + \frac{2 \ln(dx+c) \operatorname{arctanh}(dx+c)}{3} - \frac{\operatorname{arctanh}(dx+c)}{3} \right)$ |
| parts | $-\frac{a^2}{3e^4(dx+c)^3 d} + \frac{b^2}{3(dx+c)^3} \left(-\frac{\operatorname{arctanh}(dx+c)^2}{3} - \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{3} - \frac{\operatorname{arctanh}(dx+c)}{3(dx+c)^2} + \frac{2 \ln(dx+c) \operatorname{arctanh}(dx+c)}{3} - \frac{\operatorname{arctanh}(dx+c)}{3} \right)$ |

input `int((a+b*arctanh(d*x+c))^2/(d*e*x+c*e)^4,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{d} \left(-\frac{1}{3} \frac{a^2}{e^4} (d*x+c)^3 + \frac{b^2}{e^4} \left(-\frac{1}{3} (d*x+c)^3 \operatorname{arctanh}(d*x+c)^2 - \frac{1}{3} \operatorname{arctanh}(d*x+c) \ln(d*x+c-1) - \frac{1}{3} (d*x+c)^2 \operatorname{arctanh}(d*x+c) + \frac{2}{3} \ln(d*x+c) \operatorname{arctanh}(d*x+c) - \frac{1}{3} \operatorname{arctanh}(d*x+c) \ln(d*x+c+1) - \frac{1}{6} \ln(d*x+c-1) + \frac{1}{6} \ln(d*x+c+1) - \frac{1}{3} (d*x+c) + \frac{1}{3} \operatorname{dilog}\left(\frac{1}{2}d*x + \frac{1}{2}c + \frac{1}{2}\right) + \frac{1}{6} \ln(d*x+c-1) \ln\left(\frac{1}{2}d*x + \frac{1}{2}c + \frac{1}{2}\right) - \frac{1}{12} \ln(d*x+c-1)^2 + \frac{1}{12} \ln(d*x+c+1)^2 - \frac{1}{6} (\ln(d*x+c+1) - \ln\left(\frac{1}{2}d*x + \frac{1}{2}c + \frac{1}{2}\right)) \ln\left(-\frac{1}{2}d*x - \frac{1}{2}c + \frac{1}{2}\right) - \frac{1}{3} \operatorname{dilog}(d*x+c) - \frac{1}{3} \operatorname{dilog}(d*x+c+1) - \frac{1}{3} \ln(d*x+c) \ln(d*x+c+1) \right) + 2 \frac{b*a}{e^4} \left(-\frac{1}{3} (d*x+c)^3 \operatorname{arctanh}(d*x+c) - \frac{1}{6} \ln(d*x+c-1) - \frac{1}{6} (d*x+c)^2 + \frac{1}{3} \ln(d*x+c) - \frac{1}{6} \ln(d*x+c+1) \right) \right)$$

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^4} dx = \int \frac{(b \operatorname{arctanh}(dx + c) + a)^2}{(dex + ce)^4} dx$$

input `integrate((a+b*arctanh(d*x+c))^2/(d*e*x+c*e)^4,x, algorithm="fricas")`

output `integral((b^2*arctanh(d*x + c)^2 + 2*a*b*arctanh(d*x + c) + a^2)/(d^4*e^4*x^4 + 4*c*d^3*e^4*x^3 + 6*c^2*d^2*e^4*x^2 + 4*c^3*d*e^4*x + c^4*e^4), x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^4} dx = \frac{\int \frac{a^2}{c^4 + 4c^3 dx + 6c^2 d^2 x^2 + 4cd^3 x^3 + d^4 x^4} dx + \int \frac{b^2 \operatorname{atanh}^2(c+dx)}{c^4 + 4c^3 dx + 6c^2 d^2 x^2 + 4cd^3 x^3 + d^4 x^4} dx + \int \frac{2ab \operatorname{atanh}(c+dx)}{c^4 + 4c^3 dx + 6c^2 d^2 x^2 + 4cd^3 x^3 + d^4 x^4} dx}{e^4}$$

input `integrate((a+b*atanh(d*x+c))**2/(d*e*x+c*e)**4,x)`

output

```
(Integral(a**2/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + 4*c*d**3*x**3 + d**4*x**4), x) + Integral(b**2*atanh(c + d*x)**2/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + 4*c*d**3*x**3 + d**4*x**4), x) + Integral(2*a*b*atanh(c + d*x)/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + 4*c*d**3*x**3 + d**4*x**4), x))/e**4
```

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^4} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^2}{(dex + ce)^4} dx$$

input

```
integrate((a+b*arctanh(d*x+c))^2/(d*e*x+c*e)^4,x, algorithm="maxima")
```

output

```
-1/3*(d*(1/(d^4*e^4*x^2 + 2*c*d^3*e^4*x + c^2*d^2*e^4) + log(d*x + c + 1)/(d^2*e^4) - 2*log(d*x + c)/(d^2*e^4) + log(d*x + c - 1)/(d^2*e^4)) + 2*arctanh(d*x + c)/(d^4*e^4*x^3 + 3*c*d^3*e^4*x^2 + 3*c^2*d^2*e^4*x + c^3*d*e^4))*a*b - 1/12*b^2*(log(-d*x - c + 1)^2/(d^4*e^4*x^3 + 3*c*d^3*e^4*x^2 + 3*c^2*d^2*e^4*x + c^3*d*e^4) + 3*integrate(-1/3*(3*(d*x + c - 1)*log(d*x + c + 1)^2 + 2*(d*x - 3*(d*x + c - 1)*log(d*x + c + 1) + c)*log(-d*x - c + 1))/(d^5*e^4*x^5 + c^5*e^4 - c^4*e^4 + (5*c*d^4*e^4 - d^4*e^4)*x^4 + 2*(5*c^2*d^3*e^4 - 2*c*d^3*e^4)*x^3 + 2*(5*c^3*d^2*e^4 - 3*c^2*d^2*e^4)*x^2 + (5*c^4*d*e^4 - 4*c^3*d*e^4)*x), x)) - 1/3*a^2/(d^4*e^4*x^3 + 3*c*d^3*e^4*x^2 + 3*c^2*d^2*e^4*x + c^3*d*e^4)
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^4} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^2}{(dex + ce)^4} dx$$

input

```
integrate((a+b*arctanh(d*x+c))^2/(d*e*x+c*e)^4,x, algorithm="giac")
```

output

```
integrate((b*arctanh(d*x + c) + a)^2/(d*e*x + c*e)^4, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^4} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^2}{(ce + dex)^4} dx$$

input `int((a + b*atanh(c + d*x))^2/(c*e + d*e*x)^4,x)`output `int((a + b*atanh(c + d*x))^2/(c*e + d*e*x)^4, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^4} dx = \text{too large to display}$$

input `int((a+b*atanh(d*x+c))^2/(d*e*x+c*e)^4,x)`

output

```
(3*atanh(c + d*x)**2*b**2*c**5 + 9*atanh(c + d*x)**2*b**2*c**4*d*x + 9*ata
nh(c + d*x)**2*b**2*c**3*d**2*x**2 + 3*atanh(c + d*x)**2*b**2*c**3 + 3*ata
nh(c + d*x)**2*b**2*c**2*d**3*x**3 + 9*atanh(c + d*x)**2*b**2*c**2*d*x + 9
*atanh(c + d*x)**2*b**2*c*d**2*x**2 - 6*atanh(c + d*x)**2*b**2*c + 3*atanh
(c + d*x)**2*b**2*d**3*x**3 - 12*atanh(c + d*x)*a*b*c - 2*atanh(c + d*x)*b
**2*c**4 + 6*atanh(c + d*x)*b**2*c**2*d**2*x**2 - 8*atanh(c + d*x)*b**2*c*
*2 + 4*atanh(c + d*x)*b**2*c*d**3*x**3 - 12*atanh(c + d*x)*b**2*c*d*x - 6*
atanh(c + d*x)*b**2*d**2*x**2 + 6*int((atanh(c + d*x)*x**2)/(c**6 + 6*c**5
*d*x + 15*c**4*d**2*x**2 - c**4 + 20*c**3*d**3*x**3 - 4*c**3*d*x + 15*c**2
*d**4*x**4 - 6*c**2*d**2*x**2 + 6*c*d**5*x**5 - 4*c*d**3*x**3 + d**6*x**6
- d**4*x**4),x)*b**2*c**3*d**3 + 18*int((atanh(c + d*x)*x**2)/(c**6 + 6*c*
*5*d*x + 15*c**4*d**2*x**2 - c**4 + 20*c**3*d**3*x**3 - 4*c**3*d*x + 15*c*
*2*d**4*x**4 - 6*c**2*d**2*x**2 + 6*c*d**5*x**5 - 4*c*d**3*x**3 + d**6*x**
6 - d**4*x**4),x)*b**2*c**2*d**4*x + 18*int((atanh(c + d*x)*x**2)/(c**6 +
6*c**5*d*x + 15*c**4*d**2*x**2 - c**4 + 20*c**3*d**3*x**3 - 4*c**3*d*x + 1
5*c**2*d**4*x**4 - 6*c**2*d**2*x**2 + 6*c*d**5*x**5 - 4*c*d**3*x**3 + d**6
*x**6 - d**4*x**4),x)*b**2*c*d**5*x**2 + 6*int((atanh(c + d*x)*x**2)/(c**6
+ 6*c**5*d*x + 15*c**4*d**2*x**2 - c**4 + 20*c**3*d**3*x**3 - 4*c**3*d*x
+ 15*c**2*d**4*x**4 - 6*c**2*d**2*x**2 + 6*c*d**5*x**5 - 4*c*d**3*x**3 + d
**6*x**6 - d**4*x**4),x)*b**2*d**6*x**3 - 6*log(c + d*x - 1)*a*b*c**4 - ...
```

3.22 $\int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{(ce+dex)^5} dx$

| | |
|---|-----|
| Optimal result | 219 |
| Mathematica [A] (verified) | 220 |
| Rubi [A] (warning: unable to verify) | 220 |
| Maple [A] (verified) | 224 |
| Fricas [B] (verification not implemented) | 225 |
| Sympy [B] (verification not implemented) | 226 |
| Maxima [B] (verification not implemented) | 227 |
| Giac [B] (verification not implemented) | 228 |
| Mupad [B] (verification not implemented) | 228 |
| Reduce [B] (verification not implemented) | 229 |

Optimal result

Integrand size = 23, antiderivative size = 172

$$\int \frac{(a + b\operatorname{arctanh}(c + dx))^2}{(ce + dex)^5} dx = -\frac{b^2}{12de^5(c + dx)^2} - \frac{b(a + b\operatorname{arctanh}(c + dx))}{6de^5(c + dx)^3} - \frac{b(a + b\operatorname{arctanh}(c + dx))}{2de^5(c + dx)} + \frac{(a + b\operatorname{arctanh}(c + dx))^2}{4de^5} - \frac{(a + b\operatorname{arctanh}(c + dx))^2}{4de^5(c + dx)^4} + \frac{2b^2 \log(c + dx)}{3de^5} - \frac{b^2 \log(1 - (c + dx)^2)}{3de^5}$$

output

```
-1/12*b^2/d/e^5/(d*x+c)^2-1/6*b*(a+b*arctanh(d*x+c))/d/e^5/(d*x+c)^3-1/2*b
*(a+b*arctanh(d*x+c))/d/e^5/(d*x+c)+1/4*(a+b*arctanh(d*x+c))^2/d/e^5-1/4*(
a+b*arctanh(d*x+c))^2/d/e^5/(d*x+c)^4+2/3*b^2*ln(d*x+c)/d/e^5-1/3*b^2*ln(1
-(d*x+c)^2)/d/e^5
```

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.27

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^5} dx =$$

$$-\frac{3a^2}{(c+dx)^4} + \frac{2ab}{(c+dx)^3} + \frac{b^2}{(c+dx)^2} + \frac{6ab}{c+dx} + \frac{2b(3a+b(c+3c^3+dx+9c^2dx+9cd^2x^2+3d^3x^3))\operatorname{arctanh}(c+dx)}{(c+dx)^4} - \frac{3b^2(-1+c^4+4c^3dx+4c^2d^2x^2+4cd^3x^3+d^4x^4)}{(c+dx)^5}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])^2/(c*e + d*e*x)^5,x]
```

output

```
-1/12*((3*a^2)/(c + d*x)^4 + (2*a*b)/(c + d*x)^3 + b^2/(c + d*x)^2 + (6*a*b)/(c + d*x) + (2*b*(3*a + b*(c + 3*c^3 + d*x + 9*c^2*d*x + 9*c*d^2*x^2 + 3*d^3*x^3))*ArcTanh[c + d*x])/(c + d*x)^4 - (3*b^2*(-1 + c^4 + 4*c^3*d*x + 6*c^2*d^2*x^2 + 4*c*d^3*x^3 + d^4*x^4))*ArcTanh[c + d*x]^2/(c + d*x)^4 + b*(3*a + 4*b)*Log[1 - c - d*x] - 8*b^2*Log[c + d*x] - (3*a - 4*b)*b*Log[1 + c + d*x])/(d*e^5)
```

Rubi [A] (warning: unable to verify)Time = 1.02 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.91, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.652$, Rules used = {6657, 27, 6452, 6544, 6452, 243, 54, 2009, 6544, 6452, 243, 47, 14, 16, 6510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^5} dx$$

$$\downarrow 6657$$

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{e^5 (c + dx)^5} d(c + dx)$$

$$\frac{d}{d}$$

$$\downarrow 27$$

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(c + dx)^5} d(c + dx)$$

$$\frac{d}{de^5}$$

$$\begin{aligned} & \downarrow 6452 \\ & \frac{\frac{1}{2}b \int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)^4(1-(c+dx)^2)} d(c+dx) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 6544 \\ & \frac{\frac{1}{2}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)^4} d(c+dx) + \int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)^2(1-(c+dx)^2)} d(c+dx) \right) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 6452 \\ & \frac{\frac{1}{2}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)^2(1-(c+dx)^2)} d(c+dx) + \frac{1}{3}b \int \frac{1}{(c+dx)^3(1-(c+dx)^2)} d(c+dx) - \frac{a+\operatorname{barctanh}(c+dx)}{3(c+dx)^3} \right) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 243 \\ & \frac{\frac{1}{2}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)^2(1-(c+dx)^2)} d(c+dx) + \frac{1}{6}b \int \frac{1}{(-c-dx+1)(c+dx)^4} d(c+dx)^2 - \frac{a+\operatorname{barctanh}(c+dx)}{3(c+dx)^3} \right) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 54 \\ & \frac{\frac{1}{2}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)^2(1-(c+dx)^2)} d(c+dx) + \frac{1}{6}b \int \left(\frac{1}{(c+dx)^2} + \frac{1}{(c+dx)^4} + \frac{1}{-c-dx+1} \right) d(c+dx)^2 - \frac{a+\operatorname{barctanh}(c+dx)}{3(c+dx)^3} \right) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 2009 \\ & \frac{\frac{1}{2}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)^2(1-(c+dx)^2)} d(c+dx) - \frac{a+\operatorname{barctanh}(c+dx)}{3(c+dx)^3} + \frac{1}{6}b \left(-\frac{1}{(c+dx)^2} - \log(-c-dx+1) + \log((c+dx)^2) \right) \right) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 6544 \\ & \frac{\frac{1}{2}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)^2} d(c+dx) + \int \frac{a+\operatorname{barctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) - \frac{a+\operatorname{barctanh}(c+dx)}{3(c+dx)^3} + \frac{1}{6}b \left(-\frac{1}{(c+dx)^2} - \log(-c-dx+1) + \log((c+dx)^2) \right) \right) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 6452 \\ & \frac{\frac{1}{2}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) + b \int \frac{1}{(c+dx)(1-(c+dx)^2)} d(c+dx) - \frac{a+\operatorname{barctanh}(c+dx)}{c+dx} - \frac{a+\operatorname{barctanh}(c+dx)}{3(c+dx)^3} + \frac{1}{6}b \left(-\frac{1}{(c+dx)^2} - \log(-c-dx+1) + \log((c+dx)^2) \right) \right) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 243 \end{aligned}$$

$$\frac{\frac{1}{2}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) + \frac{1}{2}b \int \frac{1}{(-c-dx+1)(c+dx)^2} d(c+dx)^2 - \frac{a+\operatorname{barctanh}(c+dx)}{c+dx} - \frac{a+\operatorname{barctanh}(c+dx)}{3(c+dx)^3} + \frac{1}{6} \right)}{de^5}$$

↓ 47

$$\frac{\frac{1}{2}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) + \frac{1}{2}b \left(\int \frac{1}{-c-dx+1} d(c+dx)^2 + \int \frac{1}{(c+dx)^2} d(c+dx)^2 \right) - \frac{a+\operatorname{barctanh}(c+dx)}{c+dx} - \frac{a+\operatorname{barctanh}(c+dx)}{3(c+dx)^3} \right)}{de^5}$$

↓ 14

$$\frac{\frac{1}{2}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) + \frac{1}{2}b \left(\int \frac{1}{-c-dx+1} d(c+dx)^2 + \log((c+dx)^2) \right) - \frac{a+\operatorname{barctanh}(c+dx)}{c+dx} - \frac{a+\operatorname{barctanh}(c+dx)}{3(c+dx)^3} \right)}{de^5}$$

↓ 16

$$\frac{\frac{1}{2}b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) - \frac{a+\operatorname{barctanh}(c+dx)}{c+dx} - \frac{a+\operatorname{barctanh}(c+dx)}{3(c+dx)^3} + \frac{1}{2}b (\log((c+dx)^2) - \log(-c-dx+1)) \right)}{de^5}$$

↓ 6510

$$\frac{\frac{1}{2}b \left(\frac{(a+\operatorname{barctanh}(c+dx))^2}{2b} - \frac{a+\operatorname{barctanh}(c+dx)}{c+dx} - \frac{a+\operatorname{barctanh}(c+dx)}{3(c+dx)^3} + \frac{1}{2}b (\log((c+dx)^2) - \log(-c-dx+1)) + \frac{1}{6}b \right)}{de^5}$$

input `Int[(a + b*ArcTanh[c + d*x])^2/(c*e + d*e*x)^5,x]`

output `(-1/4*(a + b*ArcTanh[c + d*x])^2/(c + d*x)^4 + (b*(-1/3*(a + b*ArcTanh[c + d*x]))/(c + d*x)^3 - (a + b*ArcTanh[c + d*x])/(c + d*x) + (a + b*ArcTanh[c + d*x])^2/(2*b) + (b*(-Log[1 - c - d*x] + Log[(c + d*x)^2]))/2 + (b*(-(c + d*x)^(-2) - Log[1 - c - d*x] + Log[(c + d*x)^2]))/6)/2)/(d*e^5)`

Definitions of rubi rules used

- rule 14 $\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ ; FreeQ}[a, x]$
- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$
- rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] \text{ ; FreeQ}[b, x]$
- rule 47 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x]$
- rule 54 $\text{Int}[(a_)+(b_)*(x_)^{(m_)}*((c_)+(d_)*(x_)^{(n_)}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{ILtQ}[m, 0] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ !(\text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[m + n + 2, 0])$
- rule 243 $\text{Int}[(x_)^{(m_)}*((a_)+(b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] \text{ ; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$
- rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ ; SumQ}[u]$
- rule 6452 $\text{Int}[(a_)+\text{ArcTanh}[(c_)*(x_)^{(n_)}]*(b_)^{(p_)}*(x_)^{(m_)}, x_Symbol] : > \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{(2*n)})}), x], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

- rule 6510 $\text{Int}[\text{((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))}^{\text{(p_.)}}/\text{((d_.) + (e_.)*(x_.)^2)}, \text{x_Symbol}] \text{:> Simp}[\text{(a + b*ArcTanh[c*x])}^{\text{(p + 1)}}/\text{(b*c*d*(p + 1))}, \text{x}] \text{/; FreeQ}\{\text{a, b, c, d, e, p}\}, \text{x}\} \ \&\& \ \text{EqQ}[\text{c}^2\text{*d + e, 0}] \ \&\& \ \text{NeQ}[\text{p, -1}]$
- rule 6544 $\text{Int}[\text{((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))}^{\text{(p_.)}}*\text{((f_.)*(x_.))}^{\text{(m_.)}}/\text{((d_.) + (e_.)*(x_.)^2)}, \text{x_Symbol}] \text{:> Simp}[\text{1/d Int}[\text{(f*x)}^{\text{m*(a + b*ArcTanh[c*x])}^{\text{p}}, \text{x}], \text{x}] - \text{Simp}[\text{e/(d*f}^2) Int}[\text{(f*x)}^{\text{(m + 2)}}*\text{((a + b*ArcTanh[c*x])}^{\text{p}}/\text{(d + e*x}^2)), \text{x}], \text{x}] \text{/; FreeQ}\{\text{a, b, c, d, e, f}\}, \text{x}\} \ \&\& \ \text{GtQ}[\text{p, 0}] \ \&\& \ \text{LtQ}[\text{m, -1}]$
- rule 6657 $\text{Int}[\text{((a_.) + ArcTanh[(c_.) + (d_.)*(x_.)]*(b_.))}^{\text{(p_.)}}*\text{((e_.) + (f_.)*(x_.))}^{\text{(m_.)}}, \text{x_Symbol}] \text{:> Simp}[\text{1/d Subst}[\text{Int}[\text{(f*(x/d))}^{\text{m*(a + b*ArcTanh[x])}^{\text{p}}, \text{x}], \text{x, c + d*x}], \text{x}] \text{/; FreeQ}\{\text{a, b, c, d, e, f, m}\}, \text{x}\} \ \&\& \ \text{EqQ}[\text{d*e - c*f, 0}] \ \&\& \ \text{IGtQ}[\text{p, 0}]$

Maple [A] (verified)

Time = 2.06 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.59

| method | result |
|------------------|---|
| derivativdivides | $-\frac{a^2}{4e^5(dx+c)^4} + \frac{b^2 \left(-\frac{\text{arctanh}(dx+c)^2}{4(dx+c)^4} - \frac{\text{arctanh}(dx+c) \ln(dx+c-1)}{4} - \frac{\text{arctanh}(dx+c)}{6(dx+c)^3} - \frac{\text{arctanh}(dx+c)}{2(dx+c)} + \frac{\text{arctanh}(dx+c) \ln(dx+c)}{4} \right)}{4e^5(dx+c)^4}$ |
| default | $-\frac{a^2}{4e^5(dx+c)^4} + \frac{b^2 \left(-\frac{\text{arctanh}(dx+c)^2}{4(dx+c)^4} - \frac{\text{arctanh}(dx+c) \ln(dx+c-1)}{4} - \frac{\text{arctanh}(dx+c)}{6(dx+c)^3} - \frac{\text{arctanh}(dx+c)}{2(dx+c)} + \frac{\text{arctanh}(dx+c) \ln(dx+c)}{4} \right)}{4e^5(dx+c)^4}$ |
| parts | $-\frac{a^2}{4e^5(dx+c)^4 d} + \frac{b^2 \left(-\frac{\text{arctanh}(dx+c)^2}{4(dx+c)^4} - \frac{\text{arctanh}(dx+c) \ln(dx+c-1)}{4} - \frac{\text{arctanh}(dx+c)}{6(dx+c)^3} - \frac{\text{arctanh}(dx+c)}{2(dx+c)} + \frac{\text{arctanh}(dx+c) \ln(dx+c)}{4} \right)}{4e^5(dx+c)^4 d}$ |
| parallelrisc | $-\frac{3a^2 d^5 + x^2 b^2 d^7 + 6ab c^3 d^5 + b^2 c^2 d^5 + 2cba d^5 + 3b^2 \text{arctanh}(dx+c)^2 d^5 + 32 \ln(dx+c-1) x^3 b^2 c d^8 - 32 \ln(dx+c) x^3 b^2 c d^8}{4e^5(dx+c)^4 d}$ |
| risc | Expression too large to display |

input $\text{int}(\text{(a+b*arctanh(d*x+c))}^2/\text{(d*e*x+c*e)}^5, \text{x, method}=\text{_RETURNVERBOSE})$

output

```
1/d*(-1/4*a^2/e^5/(d*x+c)^4+b^2/e^5*(-1/4/(d*x+c)^4*arctanh(d*x+c)^2-1/4*a
rctanh(d*x+c)*ln(d*x+c-1)-1/6/(d*x+c)^3*arctanh(d*x+c)-1/2/(d*x+c)*arctanh
(d*x+c)+1/4*arctanh(d*x+c)*ln(d*x+c+1)+1/8*ln(d*x+c-1)*ln(1/2*d*x+1/2*c+1/
2)-1/16*ln(d*x+c-1)^2-1/16*ln(d*x+c+1)^2+1/8*(ln(d*x+c+1)-ln(1/2*d*x+1/2*c
+1/2))*ln(-1/2*d*x-1/2*c+1/2)-1/3*ln(d*x+c-1)-1/12/(d*x+c)^2+2/3*ln(d*x+c)
-1/3*ln(d*x+c+1))+2*b*a/e^5*(-1/4/(d*x+c)^4*arctanh(d*x+c)-1/8*ln(d*x+c-1)
-1/12/(d*x+c)^3-1/4/(d*x+c)+1/8*ln(d*x+c+1)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 547 vs. $2(158) = 316$.

Time = 0.11 (sec) , antiderivative size = 547, normalized size of antiderivative = 3.18

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^5} dx =$$

$$\frac{24abd^3x^3 + 24abc^3 + 4(18abc + b^2)d^2x^2 + 4b^2c^2 + 8abc + 8(9abc^2 + b^2c + ab)dx - 3(b^2d^4x^4 + 4b^2$$

input

```
integrate((a+b*arctanh(d*x+c))^2/(d*e*x+c*e)^5,x, algorithm="fricas")
```

output

```
-1/48*(24*a*b*d^3*x^3 + 24*a*b*c^3 + 4*(18*a*b*c + b^2)*d^2*x^2 + 4*b^2*c^
2 + 8*a*b*c + 8*(9*a*b*c^2 + b^2*c + a*b)*d*x - 3*(b^2*d^4*x^4 + 4*b^2*c*d
^3*x^3 + 6*b^2*c^2*d^2*x^2 + 4*b^2*c^3*d*x + b^2*c^4 - b^2)*log(-(d*x + c
+ 1)/(d*x + c - 1))^2 + 12*a^2 - 4*((3*a*b - 4*b^2)*d^4*x^4 + 4*(3*a*b - 4
*b^2)*c*d^3*x^3 + 6*(3*a*b - 4*b^2)*c^2*d^2*x^2 + 4*(3*a*b - 4*b^2)*c^3*d*
x + (3*a*b - 4*b^2)*c^4)*log(d*x + c + 1) - 32*(b^2*d^4*x^4 + 4*b^2*c*d^3*
x^3 + 6*b^2*c^2*d^2*x^2 + 4*b^2*c^3*d*x + b^2*c^4)*log(d*x + c) + 4*((3*a*
b + 4*b^2)*d^4*x^4 + 4*(3*a*b + 4*b^2)*c*d^3*x^3 + 6*(3*a*b + 4*b^2)*c^2*d
^2*x^2 + 4*(3*a*b + 4*b^2)*c^3*d*x + (3*a*b + 4*b^2)*c^4)*log(d*x + c - 1)
+ 4*(3*b^2*d^3*x^3 + 9*b^2*c*d^2*x^2 + 3*b^2*c^3 + b^2*c + (9*b^2*c^2 + b
^2)*d*x + 3*a*b)*log(-(d*x + c + 1)/(d*x + c - 1)))/(d^5*e^5*x^4 + 4*c*d^4
*e^5*x^3 + 6*c^2*d^3*e^5*x^2 + 4*c^3*d^2*e^5*x + c^4*d*e^5)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3516 vs. $2(148) = 296$.

Time = 3.46 (sec) , antiderivative size = 3516, normalized size of antiderivative = 20.44

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^5} dx = \text{Too large to display}$$

input `integrate((a+b*atanh(d*x+c))**2/(d*e*x+c*e)**5,x)`

output

```
Piecewise((-3*a**2/(12*c**4*d*e**5 + 48*c**3*d**2*e**5*x + 72*c**2*d**3*e**5*x**2 + 48*c*d**4*e**5*x**3 + 12*d**5*e**5*x**4) + 6*a*b*c**4*atanh(c + d*x)/(12*c**4*d*e**5 + 48*c**3*d**2*e**5*x + 72*c**2*d**3*e**5*x**2 + 48*c*d**4*e**5*x**3 + 12*d**5*e**5*x**4) + 24*a*b*c**3*d*x*atanh(c + d*x)/(12*c**4*d*e**5 + 48*c**3*d**2*e**5*x + 72*c**2*d**3*e**5*x**2 + 48*c*d**4*e**5*x**3 + 12*d**5*e**5*x**4) - 6*a*b*c**3/(12*c**4*d*e**5 + 48*c**3*d**2*e**5*x + 72*c**2*d**3*e**5*x**2 + 48*c*d**4*e**5*x**3 + 12*d**5*e**5*x**4) + 36*a*b*c**2*d**2*x**2*atanh(c + d*x)/(12*c**4*d*e**5 + 48*c**3*d**2*e**5*x + 72*c**2*d**3*e**5*x**2 + 48*c*d**4*e**5*x**3 + 12*d**5*e**5*x**4) - 18*a*b*c**2*d*x/(12*c**4*d*e**5 + 48*c**3*d**2*e**5*x + 72*c**2*d**3*e**5*x**2 + 48*c*d**4*e**5*x**3 + 12*d**5*e**5*x**4) + 24*a*b*c*d**3*x**3*atanh(c + d*x)/(12*c**4*d*e**5 + 48*c**3*d**2*e**5*x + 72*c**2*d**3*e**5*x**2 + 48*c*d**4*e**5*x**3 + 12*d**5*e**5*x**4) - 18*a*b*c*d**2*x**2/(12*c**4*d*e**5 + 48*c**3*d**2*e**5*x + 72*c**2*d**3*e**5*x**2 + 48*c*d**4*e**5*x**3 + 12*d**5*e**5*x**4) - 2*a*b*c/(12*c**4*d*e**5 + 48*c**3*d**2*e**5*x + 72*c**2*d**3*e**5*x**2 + 48*c*d**4*e**5*x**3 + 12*d**5*e**5*x**4) + 6*a*b*d**4*x**4*atanh(c + d*x)/(12*c**4*d*e**5 + 48*c**3*d**2*e**5*x + 72*c**2*d**3*e**5*x**2 + 48*c*d**4*e**5*x**3 + 12*d**5*e**5*x**4) - 6*a*b*d**3*x**3/(12*c**4*d*e**5 + 48*c**3*d**2*e**5*x + 72*c**2*d**3*e**5*x**2 + 48*c*d**4*e**5*x**3 + 12*d**5*e**5*x**4) - 2*a*b*d*x/(12*c**4*d*e**5 + 48*c**3*d**2*...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 613 vs. $2(158) = 316$.

Time = 0.07 (sec) , antiderivative size = 613, normalized size of antiderivative = 3.56

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^5} dx =$$

$$-\frac{1}{12} \left(d \left(\frac{2(3d^2x^2 + 6cdx + 3c^2 + 1)}{d^5e^5x^3 + 3cd^4e^5x^2 + 3c^2d^3e^5x + c^3d^2e^5} - \frac{3 \log(dx + c + 1)}{d^2e^5} + \frac{3 \log(dx + c - 1)}{d^2e^5} \right) + \frac{3 \log(dx + c + 1)}{d^2e^5} + \frac{3 \log(dx + c - 1)}{d^2e^5} \right) + \frac{3 \log(dx + c + 1)}{d^2e^5} + \frac{3 \log(dx + c - 1)}{d^2e^5}$$

$$-\frac{1}{48} \left(d^2 \left(\frac{3(d^2x^2 + 2cdx + c^2) \log(dx + c + 1)^2 + 3(d^2x^2 + 2cdx + c^2) \log(dx + c - 1)^2 + 2(8d^2x^2 + 16cdx + 8c^2 - 3(d^2x^2 + 2cdx + c^2) \log(dx + c - 1)) \log(dx + c + 1) + 16(d^2x^2 + 2cdx + c^2) \log(dx + c - 1) + 4}{d^5e^5x^4 + 4cd^4e^5x^3 + 6c^2d^3e^5x^2 + 4c^3d^2e^5x + c^4de^5} - 32 \log(dx + c) / (d^3e^5) \right) + 4d \left(\frac{2(3d^2x^2 + 6cdx + 3c^2 + 1)}{d^5e^5x^3 + 3cd^4e^5x^2 + 3c^2d^3e^5x + c^3d^2e^5} - \frac{3 \log(dx + c + 1)}{d^2e^5} + \frac{3 \log(dx + c - 1)}{d^2e^5} \right) \operatorname{arctanh}(dx + c) \right) b^2 - \frac{1}{4} b^2 \operatorname{arctanh}(dx + c)^2 / (d^5e^5x^4 + 4cd^4e^5x^3 + 6c^2d^3e^5x^2 + 4c^3d^2e^5x + c^4de^5) - \frac{1}{4} a^2 / (d^5e^5x^4 + 4cd^4e^5x^3 + 6c^2d^3e^5x^2 + 4c^3d^2e^5x + c^4de^5)$$

input `integrate((a+b*arctanh(d*x+c))^2/(d*e*x+c*e)^5,x, algorithm="maxima")`

output

```
-1/12*(d*(2*(3*d^2*x^2 + 6*c*d*x + 3*c^2 + 1)/(d^5*e^5*x^3 + 3*c*d^4*e^5*x^2 + 3*c^2*d^3*e^5*x + c^3*d^2*e^5) - 3*log(d*x + c + 1)/(d^2*e^5) + 3*log(d*x + c - 1)/(d^2*e^5)) + 6*arctanh(d*x + c)/(d^5*e^5*x^4 + 4*c*d^4*e^5*x^3 + 6*c^2*d^3*e^5*x^2 + 4*c^3*d^2*e^5*x + c^4*d*e^5))*a*b - 1/48*(d^2*((3*(d^2*x^2 + 2*c*d*x + c^2)*log(d*x + c + 1)^2 + 3*(d^2*x^2 + 2*c*d*x + c^2)*log(d*x + c - 1)^2 + 2*(8*d^2*x^2 + 16*c*d*x + 8*c^2 - 3*(d^2*x^2 + 2*c*d*x + c^2)*log(d*x + c - 1))*log(d*x + c + 1) + 16*(d^2*x^2 + 2*c*d*x + c^2)*log(d*x + c - 1) + 4)/(d^5*e^5*x^4 + 2*c*d^4*e^5*x^3 + c^2*d^3*e^5) - 32*log(d*x + c)/(d^3*e^5)) + 4*d*(2*(3*d^2*x^2 + 6*c*d*x + 3*c^2 + 1)/(d^5*e^5*x^3 + 3*c*d^4*e^5*x^2 + 3*c^2*d^3*e^5*x + c^3*d^2*e^5) - 3*log(d*x + c + 1)/(d^2*e^5) + 3*log(d*x + c - 1)/(d^2*e^5))*arctanh(d*x + c))*b^2 - 1/4*b^2*arctanh(d*x + c)^2/(d^5*e^5*x^4 + 4*c*d^4*e^5*x^3 + 6*c^2*d^3*e^5*x^2 + 4*c^3*d^2*e^5*x + c^4*d*e^5) - 1/4*a^2/(d^5*e^5*x^4 + 4*c*d^4*e^5*x^3 + 6*c^2*d^3*e^5*x^2 + 4*c^3*d^2*e^5*x + c^4*d*e^5)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 730 vs. $2(158) = 316$.

Time = 0.14 (sec) , antiderivative size = 730, normalized size of antiderivative = 4.24

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^5} dx = \text{Too large to display}$$

input `integrate((a+b*arctanh(d*x+c))^2/(d*e*x+c*e)^5,x, algorithm="giac")`

output

```
1/12*((c + 1)*d - (c - 1)*d)*(3*((d*x + c + 1)^3*b^2/(d*x + c - 1)^3 + (d*x + c + 1)*b^2/(d*x + c - 1))*log(-(d*x + c + 1)/(d*x + c - 1))^2/((d*x + c + 1)^4*d^2*e^5/(d*x + c - 1)^4 + 4*(d*x + c + 1)^3*d^2*e^5/(d*x + c - 1)^3 + 6*(d*x + c + 1)^2*d^2*e^5/(d*x + c - 1)^2 + 4*(d*x + c + 1)*d^2*e^5/(d*x + c - 1) + d^2*e^5) + 2*(6*(d*x + c + 1)^3*a*b/(d*x + c - 1)^3 + 6*(d*x + c + 1)*a*b/(d*x + c - 1) + 3*(d*x + c + 1)^3*b^2/(d*x + c - 1)^3 + 6*(d*x + c + 1)^2*b^2/(d*x + c - 1)^2 + 5*(d*x + c + 1)*b^2/(d*x + c - 1) + 2*b^2)*log(-(d*x + c + 1)/(d*x + c - 1))/((d*x + c + 1)^4*d^2*e^5/(d*x + c - 1)^4 + 4*(d*x + c + 1)^3*d^2*e^5/(d*x + c - 1)^3 + 6*(d*x + c + 1)^2*d^2*e^5/(d*x + c - 1)^2 + 4*(d*x + c + 1)*d^2*e^5/(d*x + c - 1) + d^2*e^5) + 2*(6*(d*x + c + 1)^3*a^2/(d*x + c - 1)^3 + 6*(d*x + c + 1)*a^2/(d*x + c - 1) + 6*(d*x + c + 1)^3*a*b/(d*x + c - 1)^3 + 12*(d*x + c + 1)^2*a*b/(d*x + c - 1)^2 + 10*(d*x + c + 1)*a*b/(d*x + c - 1) + 4*a*b + (d*x + c + 1)^3*b^2/(d*x + c - 1)^3 + 2*(d*x + c + 1)^2*b^2/(d*x + c - 1)^2 + (d*x + c + 1)*b^2/(d*x + c - 1))/((d*x + c + 1)^4*d^2*e^5/(d*x + c - 1)^4 + 4*(d*x + c + 1)^3*d^2*e^5/(d*x + c - 1)^3 + 6*(d*x + c + 1)^2*d^2*e^5/(d*x + c - 1)^2 + 4*(d*x + c + 1)*d^2*e^5/(d*x + c - 1) + d^2*e^5) + 4*b^2*log(-(d*x + c + 1)/(d*x + c - 1) - 1)/(d^2*e^5) - 4*b^2*log(-(d*x + c + 1)/(d*x + c - 1))/(d^2*e^5))
```

Mupad [B] (verification not implemented)

Time = 7.05 (sec) , antiderivative size = 2746, normalized size of antiderivative = 15.97

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^5} dx = \text{Too large to display}$$

input `int((a + b*atanh(c + d*x))^2/(c*e + d*e*x)^5,x)`

output

```

log(1 - d*x - c)^2*(b^2/(16*d*e^5) - b^2/(4*d*(4*c^4*e^5 + 4*d^4*e^5*x^4 +
16*c*d^3*e^5*x^3 + 24*c^2*d^2*e^5*x^2 + 16*c^3*d*e^5*x))) + log(c + d*x +
1)^2*(b^2/(16*d*e^5) - b^2/(16*d^2*e^5*(4*c^3*x + c^4/d + d^3*x^4 + 6*c^2
*d*x^2 + 4*c*d^2*x^3))) + log(1 - d*x - c)*(log(c + d*x + 1)*(b^2/(4*d*(2*
c^4*e^5 + 2*d^4*e^5*x^4 + 8*c*d^3*e^5*x^3 + 12*c^2*d^2*e^5*x^2 + 8*c^3*d*e
^5*x)) - (b^2*(c^4 + d^4*x^4 + 4*c*d^3*x^3 + 6*c^2*d^2*x^2 + 4*c^3*d*x))/(
4*d*(2*c^4*e^5 + 2*d^4*e^5*x^4 + 8*c*d^3*e^5*x^3 + 12*c^2*d^2*e^5*x^2 + 8*
c^3*d*e^5*x))) + (3*b^2)/(4*d*(24*c^4*e^5 + 24*d^4*e^5*x^4 + 96*c*d^3*e^5*
x^3 + 144*c^2*d^2*e^5*x^2 + 96*c^3*d*e^5*x)) + (3*b*(8*a - b))/(4*d*(24*c^
4*e^5 + 24*d^4*e^5*x^4 + 96*c*d^3*e^5*x^3 + 144*c^2*d^2*e^5*x^2 + 96*c^3*d
*e^5*x)) - (b^2*(c*(2*c - 3*c^2 + 4*c^3 + c*(6*c^2 - 3*c + c*(12*c - 3) +
1) - 1) - 3*c + x^2*(d*(2*d - 6*c*d + 12*c^2*d + d*(6*c^2 - 3*c + c*(12*c
- 3) + 1) + c*(24*c*d - 3*d + d*(12*c - 3))) - 9*c*d^2 + c*(30*c*d^2 - 3*d
^2 + d*(24*c*d - 3*d + d*(12*c - 3))) + 3*d^2 + 18*c^2*d^2) + x*(d*(2*c -
3*c^2 + 4*c^3 + c*(6*c^2 - 3*c + c*(12*c - 3) + 1) - 1) - 3*d + 6*c*d + c*
(2*d - 6*c*d + 12*c^2*d + d*(6*c^2 - 3*c + c*(12*c - 3) + 1) + c*(24*c*d -
3*d + d*(12*c - 3))) - 9*c^2*d + 12*c^3*d) + 3*c^2 - 3*c^3 + 3*c^4 + 25*d
^4*x^4 + x^3*(34*c*d^3 + d*(30*c*d^2 - 3*d^2 + d*(24*c*d - 3*d + d*(12*c -
3))) - 3*d^3) + 3))/(4*d*(24*c^4*e^5 + 24*d^4*e^5*x^4 + 96*c*d^3*e^5*x^3
+ 144*c^2*d^2*e^5*x^2 + 96*c^3*d*e^5*x)) + (b^2*(c*(c*(6*c*e^5 + 2*e^5 ...

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 989, normalized size of antiderivative = 5.75

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(ce + dex)^5} dx = \text{Too large to display}$$

input

```
int((a+b*atanh(d*x+c))^2/(d*e*x+c*e)^5,x)
```

output

```
(12*atanh(c + d*x)**2*b**2*c**5 + 48*atanh(c + d*x)**2*b**2*c**4*d*x + 72*
atanh(c + d*x)**2*b**2*c**3*d**2*x**2 + 48*atanh(c + d*x)**2*b**2*c**2*d**
3*x**3 + 12*atanh(c + d*x)**2*b**2*c*d**4*x**4 - 12*atanh(c + d*x)**2*b**2
*c - 24*atanh(c + d*x)*a*b*c - 18*atanh(c + d*x)*b**2*c**4 - 48*atanh(c +
d*x)*b**2*c**3*d*x - 36*atanh(c + d*x)*b**2*c**2*d**2*x**2 - 8*atanh(c + d
*x)*b**2*c**2 - 8*atanh(c + d*x)*b**2*c*d*x + 6*atanh(c + d*x)*b**2*d**4*x
**4 - 12*log(c + d*x - 1)*a*b*c**5 - 48*log(c + d*x - 1)*a*b*c**4*d*x - 72
*log(c + d*x - 1)*a*b*c**3*d**2*x**2 - 48*log(c + d*x - 1)*a*b*c**2*d**3*x
**3 - 12*log(c + d*x - 1)*a*b*c*d**4*x**4 - 16*log(c + d*x - 1)*b**2*c**5
- 64*log(c + d*x - 1)*b**2*c**4*d*x + 3*log(c + d*x - 1)*b**2*c**4 - 96*lo
g(c + d*x - 1)*b**2*c**3*d**2*x**2 + 12*log(c + d*x - 1)*b**2*c**3*d*x - 6
4*log(c + d*x - 1)*b**2*c**2*d**3*x**3 + 18*log(c + d*x - 1)*b**2*c**2*d**
2*x**2 - 16*log(c + d*x - 1)*b**2*c*d**4*x**4 + 12*log(c + d*x - 1)*b**2*c
*d**3*x**3 + 3*log(c + d*x - 1)*b**2*d**4*x**4 + 12*log(c + d*x + 1)*a*b*c
**5 + 48*log(c + d*x + 1)*a*b*c**4*d*x + 72*log(c + d*x + 1)*a*b*c**3*d**2
*x**2 + 48*log(c + d*x + 1)*a*b*c**2*d**3*x**3 + 12*log(c + d*x + 1)*a*b*c
*d**4*x**4 - 16*log(c + d*x + 1)*b**2*c**5 - 64*log(c + d*x + 1)*b**2*c**4
*d*x - 3*log(c + d*x + 1)*b**2*c**4 - 96*log(c + d*x + 1)*b**2*c**3*d**2*x
**2 - 12*log(c + d*x + 1)*b**2*c**3*d*x - 64*log(c + d*x + 1)*b**2*c**2*d*
*3*x**3 - 18*log(c + d*x + 1)*b**2*c**2*d**2*x**2 - 16*log(c + d*x + 1)...
```

3.23 $\int (ce + dex)^2 (a + \operatorname{barctanh}(c + dx))^3 dx$

| | |
|---------------------------------------|-----|
| Optimal result | 231 |
| Mathematica [A] (verified) | 232 |
| Rubi [A] (verified) | 232 |
| Maple [C] (warning: unable to verify) | 236 |
| Fricas [F] | 237 |
| Sympy [F] | 238 |
| Maxima [F] | 238 |
| Giac [F] | 239 |
| Mupad [F(-1)] | 240 |
| Reduce [F] | 240 |

Optimal result

Integrand size = 23, antiderivative size = 263

$$\begin{aligned}
 & \int (ce + dex)^2 (a + \operatorname{barctanh}(c + dx))^3 dx \\
 &= ab^2 e^2 x + \frac{b^3 e^2 (c + dx) \operatorname{arctanh}(c + dx)}{d} - \frac{be^2 (a + \operatorname{barctanh}(c + dx))^2}{2d} \\
 &+ \frac{be^2 (c + dx)^2 (a + \operatorname{barctanh}(c + dx))^2}{2d} + \frac{e^2 (a + \operatorname{barctanh}(c + dx))^3}{3d} \\
 &+ \frac{e^2 (c + dx)^3 (a + \operatorname{barctanh}(c + dx))^3}{3d} - \frac{be^2 (a + \operatorname{barctanh}(c + dx))^2 \log\left(\frac{2}{1-c-dx}\right)}{d} \\
 &+ \frac{b^3 e^2 \log(1 - (c + dx)^2)}{2d} - \frac{b^2 e^2 (a + \operatorname{barctanh}(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-c-dx}\right)}{d} \\
 &+ \frac{b^3 e^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-c-dx}\right)}{2d}
 \end{aligned}$$

output

```

a*b^2*e^2*x+b^3*e^2*(d*x+c)*arctanh(d*x+c)/d-1/2*b*e^2*(a+b*arctanh(d*x+c)
)^2/d+1/2*b*e^2*(d*x+c)^2*(a+b*arctanh(d*x+c))^2/d+1/3*e^2*(a+b*arctanh(d*
x+c))^3/d+1/3*e^2*(d*x+c)^3*(a+b*arctanh(d*x+c))^3/d-b*e^2*(a+b*arctanh(d*
x+c))^2*ln(2/(-d*x-c+1))/d+1/2*b^3*e^2*ln(1-(d*x+c)^2)/d-b^2*e^2*(a+b*arct
anh(d*x+c))*polylog(2,1-2/(-d*x-c+1))/d+1/2*b^3*e^2*polylog(3,1-2/(-d*x-c+
1))/d

```


Mathematica [A] (verified)

Time = 0.47 (sec) , antiderivative size = 336, normalized size of antiderivative = 1.28

$$\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx$$

$$= \frac{e^2 \left(3a^2 b (c + dx)^2 + 2a^3 (c + dx)^3 + 6a^2 b (c + dx)^3 \operatorname{arctanh}(c + dx) + 3a^2 b \log(1 - (c + dx)^2) + 6ab^2 (c + \right.$$

input

```
Integrate[(c*e + d*e*x)^2*(a + b*ArcTanh[c + d*x])^3,x]
```

output

```
(e^2*(3*a^2*b*(c + d*x)^2 + 2*a^3*(c + d*x)^3 + 6*a^2*b*(c + d*x)^3*ArcTan
h[c + d*x] + 3*a^2*b*Log[1 - (c + d*x)^2] + 6*a*b^2*(c + d*x - ArcTanh[c +
d*x] + (c + d*x)^2*ArcTanh[c + d*x] - ArcTanh[c + d*x]^2 + (c + d*x)^3*Ar
cTanh[c + d*x]^2 - 2*ArcTanh[c + d*x]*Log[1 + E^(-2*ArcTanh[c + d*x]])] + P
olyLog[2, -E^(-2*ArcTanh[c + d*x])]) + b^3*(6*(c + d*x)*ArcTanh[c + d*x] -
3*(1 - (c + d*x)^2)*ArcTanh[c + d*x]^2 - 2*ArcTanh[c + d*x]^3 + 2*(c + d*
x)*ArcTanh[c + d*x]^3 - 2*(c + d*x)*(1 - (c + d*x)^2)*ArcTanh[c + d*x]^3 -
6*ArcTanh[c + d*x]^2*Log[1 + E^(-2*ArcTanh[c + d*x]])] - 6*Log[1/Sqrt[1 -
(c + d*x)^2]] + 6*ArcTanh[c + d*x]*PolyLog[2, -E^(-2*ArcTanh[c + d*x])] +
3*PolyLog[3, -E^(-2*ArcTanh[c + d*x])]))/(6*d)
```

Rubi [A] (verified)

Time = 1.63 (sec) , antiderivative size = 222, normalized size of antiderivative = 0.84, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.522$, Rules used = {6657, 27, 6452, 6542, 6542, 6542, 2009, 6510, 6546, 6470, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx$$

$$\downarrow 6657$$

$$\frac{\int e^2 (c + dx)^2 (a + b \operatorname{arctanh}(c + dx))^3 d(c + dx)}{d}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{e^2 \int (c + dx)^2 (a + \operatorname{barctanh}(c + dx))^3 d(c + dx)}{d} \\ & \downarrow 6452 \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + \operatorname{barctanh}(c + dx))^3 - b \int \frac{(c+dx)^3 (a+\operatorname{barctanh}(c+dx))^2}{1-(c+dx)^2} d(c + dx) \right)}{d} \\ & \downarrow 6542 \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + \operatorname{barctanh}(c + dx))^3 - b \left(\int \frac{(c+dx)(a+\operatorname{barctanh}(c+dx))^2}{1-(c+dx)^2} d(c + dx) - \int (c + dx)(a + \operatorname{barctanh}(c + dx)) \right) \right)}{d} \\ & \downarrow 6452 \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + \operatorname{barctanh}(c + dx))^3 - b \left(b \int \frac{(c+dx)^2 (a+\operatorname{barctanh}(c+dx))}{1-(c+dx)^2} d(c + dx) + \int \frac{(c+dx)(a+\operatorname{barctanh}(c+dx))^2}{1-(c+dx)^2} \right) \right)}{d} \\ & \downarrow 6542 \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + \operatorname{barctanh}(c + dx))^3 - b \left(b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{1-(c+dx)^2} d(c + dx) - \int (a + \operatorname{barctanh}(c + dx)) \right) \right) \right)}{d} \\ & \downarrow 2009 \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + \operatorname{barctanh}(c + dx))^3 - b \left(\int \frac{(c+dx)(a+\operatorname{barctanh}(c+dx))^2}{1-(c+dx)^2} d(c + dx) + b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{1-(c+dx)^2} d(c + dx) \right) \right) \right)}{d} \\ & \downarrow 6510 \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + \operatorname{barctanh}(c + dx))^3 - b \left(\int \frac{(c+dx)(a+\operatorname{barctanh}(c+dx))^2}{1-(c+dx)^2} d(c + dx) - \frac{1}{2} (c + dx)^2 (a + \operatorname{barctanh}(c + dx)) \right) \right)}{d} \\ & \downarrow 6546 \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + \operatorname{barctanh}(c + dx))^3 - b \left(\int \frac{(a+\operatorname{barctanh}(c+dx))^2}{-c-dx+1} d(c + dx) - \frac{(a+\operatorname{barctanh}(c+dx))^3}{3b} - \frac{1}{2} (c + dx)^2 \right) \right)}{d} \\ & \downarrow 6470 \end{aligned}$$

$$e^2 \left(\frac{1}{3}(c+dx)^3(a + \operatorname{barctanh}(c+dx))^3 - b \left(-2b \int \frac{(a + \operatorname{barctanh}(c+dx)) \log\left(\frac{2}{-c-dx+1}\right)}{1-(c+dx)^2} d(c+dx) - \frac{(a + \operatorname{barctanh}(c+dx))}{3b} \right) \right)$$

↓ 6620

$$e^2 \left(\frac{1}{3}(c+dx)^3(a + \operatorname{barctanh}(c+dx))^3 - b \left(-2b \left(\frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{-c-dx+1}\right)}{1-(c+dx)^2} d(c+dx) - \frac{1}{2} \operatorname{PolyLog}\left(2, 1 - \frac{2}{-c-dx+1}\right) \right) \right)$$

↓ 7164

$$e^2 \left(\frac{1}{3}(c+dx)^3(a + \operatorname{barctanh}(c+dx))^3 - b \left(-2b \left(\frac{1}{4} b \operatorname{PolyLog}\left(3, 1 - \frac{2}{-c-dx+1}\right) - \frac{1}{2} \operatorname{PolyLog}\left(2, 1 - \frac{2}{-c-dx+1}\right) \right) \right) (a$$

input `Int[(c*e + d*e*x)^2*(a + b*ArcTanh[c + d*x])^3,x]`

output `(e^2*(((c + d*x)^3*(a + b*ArcTanh[c + d*x])^3)/3 - b*(-1/2*((c + d*x)^2*(a + b*ArcTanh[c + d*x])^2) - (a + b*ArcTanh[c + d*x])^3/(3*b) + (a + b*ArcTanh[c + d*x])^2*Log[2/(1 - c - d*x)] + b*(-(a*(c + d*x)) - b*(c + d*x)*ArcTanh[c + d*x] + (a + b*ArcTanh[c + d*x])^2/(2*b) - (b*Log[1 - (c + d*x)^2])/2) - 2*b*(-1/2*((a + b*ArcTanh[c + d*x])*PolyLog[2, 1 - 2/(1 - c - d*x)]) + (b*PolyLog[3, 1 - 2/(1 - c - d*x)]/4))))/d`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6452 $\text{Int}[(a + \text{ArcTanh}[c \cdot x]^n] \cdot (b \cdot x)^m, x_Symbol] :$
 $> \text{Simp}[x^{m+1} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x]^n)^p / (m+1), x] - \text{Simp}[b \cdot c \cdot n \cdot (p/(m+1)) \text{Int}[x^{m+n} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x]^n)^{p-1} / (1 - c^2 \cdot x^{2n})], x], x] /;$
 $\text{FreeQ}\{a, b, c, m, n\}, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 6470 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x))^p / ((d + e \cdot x)), x_Symbol] :$
 $> \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{Log}[2/(1 + e \cdot (x/d))]) / e, x] + \text{Simp}[b \cdot c \cdot (p/e) \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{Log}[2/(1 + e \cdot (x/d))]) / (1 - c^2 \cdot x^2)], x], x] /;$
 $\text{FreeQ}\{a, b, c, d, e\}, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d^2 - e^2, 0]$

rule 6510 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x))^p / ((d + e \cdot x)^2), x_Symbol] :$
 $> \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot c \cdot d \cdot (p+1)), x] /;$
 $\text{FreeQ}\{a, b, c, d, e, p\}, x\} \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{NeQ}[p, -1]$

rule 6542 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x))^p \cdot (f \cdot x)^m / ((d + e \cdot x)^2), x_Symbol] :$
 $> \text{Simp}[f^2/e \text{Int}[(f \cdot x)^{m-2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x] - \text{Simp}[d \cdot (f^2/e) \text{Int}[(f \cdot x)^{m-2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2)], x], x] /;$
 $\text{FreeQ}\{a, b, c, d, e, f\}, x\} \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{GtQ}[m, 1]$

rule 6546 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x))^p \cdot (x) / ((d + e \cdot x)^2), x_Symbol] :$
 $> \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot e \cdot (p+1)), x] + \text{Simp}[1 / (c \cdot d) \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p / (1 - c \cdot x)], x], x] /;$
 $\text{FreeQ}\{a, b, c, d, e\}, x\} \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{IGtQ}[p, 0]$

rule 6620 $\text{Int}[(\text{Log}[u] \cdot (a + \text{ArcTanh}[c \cdot x] \cdot (b \cdot x))^p) / ((d + e \cdot x)^2), x_Symbol] :$
 $> \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{PolyLog}[2, 1 - u] / (2 \cdot c \cdot d)), x] + \text{Simp}[b \cdot (p/2) \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{PolyLog}[2, 1 - u]) / (d + e \cdot x^2)], x], x] /;$
 $\text{FreeQ}\{a, b, c, d, e\}, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{EqQ}[(1 - u)^2 - (1 - 2/(1 - c \cdot x))^2, 0]$

rule 6657

```
Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x
], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0]
&& IGtQ[p, 0]
```

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 11.55 (sec) , antiderivative size = 1099, normalized size of antiderivative = 4.18

| method | result | size |
|-------------------|---------------------------------|------|
| derivativedivides | Expression too large to display | 1099 |
| default | Expression too large to display | 1099 |
| parts | Expression too large to display | 1107 |

input

```
int((d*e*x+c*e)^2*(a+b*arctanh(d*x+c))^3,x,method=_RETURNVERBOSE)
```

output

```

1/d*(1/3*a^3*e^2*(d*x+c)^3+e^2*b^3*(1/3*(d*x+c)^3*arctanh(d*x+c)^3+1/2*(d*
x+c)^2*arctanh(d*x+c)^2+1/2*arctanh(d*x+c)^2*ln(d*x+c-1)+1/2*arctanh(d*x+c
)^2*ln(d*x+c+1)-arctanh(d*x+c)*polylog(2,-(d*x+c+1)^2/(1-(d*x+c)^2))+1/2*p
olylog(3,-(d*x+c+1)^2/(1-(d*x+c)^2))-arctanh(d*x+c)^2*ln((d*x+c+1)/(1-(d*x
+c)^2)^(1/2))+1/3*arctanh(d*x+c)^3-ln(1+(d*x+c+1)^2/(1-(d*x+c)^2))-1/2*arc
tanh(d*x+c)^2+(d*x+c+1)*arctanh(d*x+c)+1/4*I*Pi*csgn(I/(1-(d*x+c+1)^2/((d*
x+c)^2-1)))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1))*csgn(I*(d*x+c+1)^2/((d*x+c)
^2-1)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))*arctanh(d*x+c)^2+1/4*I*Pi*csgn(I*(d*x+
c+1)^2/((d*x+c)^2-1))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1)/(1-(d*x+c+1)^2/((d*
x+c)^2-1)))^2*arctanh(d*x+c)^2-1/4*I*Pi*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1))
^3*arctanh(d*x+c)^2+1/2*I*Pi*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))^2*arctan
h(d*x+c)^2-1/2*I*Pi*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))^3*arctanh(d*x+c)
^2-1/4*I*Pi*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))*csgn(I*(d*x+c+1)^2/((d*x
+c)^2-1)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))^2*arctanh(d*x+c)^2-1/4*I*Pi*csgn(I
*(d*x+c+1)^2/((d*x+c)^2-1)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))^3*arctanh(d*x+c)
^2-ln(2)*arctanh(d*x+c)^2-1/4*I*Pi*csgn(I*(d*x+c+1)/(1-(d*x+c)^2)^(1/2))^2
*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1))*arctanh(d*x+c)^2-1/2*I*Pi*csgn(I*(d*x+c
+1)/(1-(d*x+c)^2)^(1/2))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1))^2*arctanh(d*x+c
)^2-1/2*I*Pi*arctanh(d*x+c)^2)+3*e^2*a*b^2*(1/3*(d*x+c)^3*arctanh(d*x+c)^2
+1/3*(d*x+c)^2*arctanh(d*x+c)+1/3*arctanh(d*x+c)*ln(d*x+c-1)+1/3*arctan...

```

Fricas [F]

$$\int (ce + dex)^2(a + b \operatorname{arctanh}(c + dx))^3 dx = \int (dex + ce)^2(b \operatorname{arctanh}(dx + c) + a)^3 dx$$

input

```
integrate((d*e*x+c*e)^2*(a+b*arctanh(d*x+c))^3,x, algorithm="fricas")
```

output

```

integral(a^3*d^2*e^2*x^2 + 2*a^3*c*d*e^2*x + a^3*c^2*e^2 + (b^3*d^2*e^2*x^
2 + 2*b^3*c*d*e^2*x + b^3*c^2*e^2)*arctanh(d*x + c)^3 + 3*(a*b^2*d^2*e^2*x
^2 + 2*a*b^2*c*d*e^2*x + a*b^2*c^2*e^2)*arctanh(d*x + c)^2 + 3*(a^2*b*d^2*
e^2*x^2 + 2*a^2*b*c*d*e^2*x + a^2*b*c^2*e^2)*arctanh(d*x + c), x)

```

Sympy [F]

$$\begin{aligned}
& \int (ce + dex)^2 (a + \operatorname{arctanh}(c + dx))^3 dx \\
&= e^2 \left(\int a^3 c^2 dx + \int a^3 d^2 x^2 dx + \int b^3 c^2 \operatorname{atanh}^3(c + dx) dx \right. \\
&\quad + \int 3ab^2 c^2 \operatorname{atanh}^2(c + dx) dx + \int 3a^2 bc^2 \operatorname{atanh}(c + dx) dx + \int 2a^3 cdx dx \\
&\quad + \int b^3 d^2 x^2 \operatorname{atanh}^3(c + dx) dx + \int 3ab^2 d^2 x^2 \operatorname{atanh}^2(c + dx) dx \\
&\quad + \int 3a^2 bd^2 x^2 \operatorname{atanh}(c + dx) dx + \int 2b^3 cdx \operatorname{atanh}^3(c + dx) dx \\
&\quad \left. + \int 6ab^2 cdx \operatorname{atanh}^2(c + dx) dx + \int 6a^2 bcdx \operatorname{atanh}(c + dx) dx \right)
\end{aligned}$$

input `integrate((d*e*x+c*e)**2*(a+b*atanh(d*x+c))**3,x)`

output `e**2*(Integral(a**3*c**2, x) + Integral(a**3*d**2*x**2, x) + Integral(b**3*c**2*atanh(c + d*x)**3, x) + Integral(3*a*b**2*c**2*atanh(c + d*x)**2, x) + Integral(3*a**2*b*c**2*atanh(c + d*x), x) + Integral(2*a**3*c*d*x, x) + Integral(b**3*d**2*x**2*atanh(c + d*x)**3, x) + Integral(3*a*b**2*d**2*x**2*atanh(c + d*x)**2, x) + Integral(3*a**2*b*d**2*x**2*atanh(c + d*x), x) + Integral(2*b**3*c*d*x*atanh(c + d*x)**3, x) + Integral(6*a*b**2*c*d*x*atanh(c + d*x)**2, x) + Integral(6*a**2*b*c*d*x*atanh(c + d*x), x))`

Maxima [F]

$$\int (ce + dex)^2 (a + \operatorname{arctanh}(c + dx))^3 dx = \int (dex + ce)^2 (\operatorname{arctanh}(dx + c) + a)^3 dx$$

input `integrate((d*e*x+c*e)^2*(a+b*arctanh(d*x+c))^3,x, algorithm="maxima")`

output

```

1/3*a^3*d^2*e^2*x^3 + a^3*c*d*e^2*x^2 + 3/2*(2*x^2*arctanh(d*x + c) + d*(2
*x/d^2 - (c^2 + 2*c + 1)*log(d*x + c + 1)/d^3 + (c^2 - 2*c + 1)*log(d*x +
c - 1)/d^3))*a^2*b*c*d*e^2 + 1/2*(2*x^3*arctanh(d*x + c) + d*((d*x^2 - 4*c
*x)/d^3 + (c^3 + 3*c^2 + 3*c + 1)*log(d*x + c + 1)/d^4 - (c^3 - 3*c^2 + 3*
c - 1)*log(d*x + c - 1)/d^4))*a^2*b*d^2*e^2 + a^3*c^2*e^2*x + 3/2*(2*(d*x
+ c)*arctanh(d*x + c) + log(-(d*x + c)^2 + 1))*a^2*b*c^2*e^2/d - 1/24*((b^
3*d^3*e^2*x^3 + 3*b^3*c*d^2*e^2*x^2 + 3*b^3*c^2*d*e^2*x + (c^3*e^2 - e^2)*
b^3)*log(-d*x - c + 1)^3 - 3*(2*a*b^2*d^3*e^2*x^3 + (6*a*b^2*c*d^2*e^2 + b
^3*d^2*e^2)*x^2 + 2*(3*a*b^2*c^2*d*e^2 + b^3*c*d*e^2)*x + (b^3*d^3*e^2*x^3
+ 3*b^3*c*d^2*e^2*x^2 + 3*b^3*c^2*d*e^2*x + (c^3*e^2 + e^2)*b^3)*log(d*x
+ c + 1))*log(-d*x - c + 1)^2)/d - integrate(-1/8*((b^3*d^3*e^2*x^3 + (3*c
*d^2*e^2 - d^2*e^2)*b^3*x^2 + (3*c^2*d*e^2 - 2*c*d*e^2)*b^3*x + (c^3*e^2 -
c^2*e^2)*b^3)*log(d*x + c + 1)^3 + 6*(a*b^2*d^3*e^2*x^3 + (3*c*d^2*e^2 -
d^2*e^2)*a*b^2*x^2 + (3*c^2*d*e^2 - 2*c*d*e^2)*a*b^2*x + (c^3*e^2 - c^2*e^
2)*a*b^2)*log(d*x + c + 1)^2 - (4*a*b^2*d^3*e^2*x^3 + 2*(6*a*b^2*c*d^2*e^2
+ b^3*d^2*e^2)*x^2 + 3*(b^3*d^3*e^2*x^3 + (3*c*d^2*e^2 - d^2*e^2)*b^3*x^2
+ (3*c^2*d*e^2 - 2*c*d*e^2)*b^3*x + (c^3*e^2 - c^2*e^2)*b^3)*log(d*x + c
+ 1)^2 + 4*(3*a*b^2*c^2*d*e^2 + b^3*c*d*e^2)*x + 2*(6*(c^3*e^2 - c^2*e^2)*
a*b^2 + (c^3*e^2 + e^2)*b^3 + (6*a*b^2*d^3*e^2 + b^3*d^3*e^2)*x^3 + 3*(b^3
*c*d^2*e^2 + 2*(3*c*d^2*e^2 - d^2*e^2)*a*b^2)*x^2 + 3*(b^3*c^2*d*e^2 + ...

```

Giac [F]

$$\int (ce + dex)^2(a + b \operatorname{arctanh}(c + dx))^3 dx = \int (dex + ce)^2(b \operatorname{arctanh}(dx + c) + a)^3 dx$$

input

```
integrate((d*e*x+c*e)^2*(a+b*arctanh(d*x+c))^3,x, algorithm="giac")
```

output

```
integrate((d*e*x + c*e)^2*(b*arctanh(d*x + c) + a)^3, x)
```


Mupad [F(-1)]

Timed out.

$$\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx = \int (ce + dex)^2 (a + b \operatorname{atanh}(c + dx))^3 dx$$

input `int((c*e + d*e*x)^2*(a + b*atanh(c + d*x))^3,x)`output `int((c*e + d*e*x)^2*(a + b*atanh(c + d*x))^3, x)`**Reduce [F]**

$$\int (ce + dex)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx$$

$$= \frac{e^2 \left(2 \operatorname{atanh}(dx + c)^3 b^3 c^3 - 2 \operatorname{atanh}(dx + c)^3 b^3 c + 3 \operatorname{atanh}(dx + c)^2 b^3 c^2 + 6 \operatorname{atanh}(dx + c) a^2 b - 6 \operatorname{atanh}(dx + c) a b^2 \right)}{e^2}$$

input `int((d*e*x+c*e)^2*(a+b*atanh(d*x+c))^3,x)`

output

```
(e**2*(2*atanh(c + d*x)**3*b**3*c**3 + 6*atanh(c + d*x)**3*b**3*c**2*d*x +
6*atanh(c + d*x)**3*b**3*c*d**2*x**2 - 2*atanh(c + d*x)**3*b**3*c + 2*ata
nh(c + d*x)**3*b**3*d**3*x**3 + 6*atanh(c + d*x)**2*a*b**2*c**3 + 18*atanh
(c + d*x)**2*a*b**2*c**2*d*x + 18*atanh(c + d*x)**2*a*b**2*c*d**2*x**2 - 6
*atanh(c + d*x)**2*a*b**2*c + 6*atanh(c + d*x)**2*a*b**2*d**3*x**3 + 3*ata
nh(c + d*x)**2*b**3*c**2 + 6*atanh(c + d*x)**2*b**3*c*d*x + 3*atanh(c + d*
x)**2*b**3*d**2*x**2 - 3*atanh(c + d*x)**2*b**3 + 6*atanh(c + d*x)*a**2*b*
c**3 + 18*atanh(c + d*x)*a**2*b*c**2*d*x + 18*atanh(c + d*x)*a**2*b*c*d**2
*x**2 + 6*atanh(c + d*x)*a**2*b*d**3*x**3 + 6*atanh(c + d*x)*a**2*b + 6*at
anh(c + d*x)*a*b**2*c**2 + 12*atanh(c + d*x)*a*b**2*c*d*x + 6*atanh(c + d*
x)*a*b**2*d**2*x**2 - 6*atanh(c + d*x)*a*b**2 + 6*atanh(c + d*x)*b**3*c +
6*atanh(c + d*x)*b**3*d*x + 6*atanh(c + d*x)*b**3 + 12*int((atanh(c + d*x)
*x)/(c**2 + 2*c*d*x + d**2*x**2 - 1),x)*a*b**2*d**2 + 6*int((atanh(c + d*x)
)**2*x)/(c**2 + 2*c*d*x + d**2*x**2 - 1),x)*b**3*d**2 + 6*log(c + d*x - 1)
*a**2*b + 6*log(c + d*x - 1)*b**3 + 6*a**3*c**2*d*x + 6*a**3*c*d**2*x**2 +
2*a**3*d**3*x**3 + 6*a**2*b*c*d*x + 3*a**2*b*d**2*x**2 + 6*a*b**2*d*x))/(
6*d)
```

3.24 $\int (ce + dex)(a + \operatorname{barctanh}(c + dx))^3 dx$

| | |
|---|-----|
| Optimal result | 242 |
| Mathematica [A] (verified) | 243 |
| Rubi [A] (verified) | 243 |
| Maple [B] (verified) | 247 |
| Fricas [F] | 248 |
| Sympy [F] | 248 |
| Maxima [B] (verification not implemented) | 249 |
| Giac [F] | 249 |
| Mupad [F(-1)] | 250 |
| Reduce [F] | 250 |

Optimal result

Integrand size = 21, antiderivative size = 160

$$\int (ce + dex)(a + \operatorname{barctanh}(c + dx))^3 dx = \frac{3be(a + \operatorname{barctanh}(c + dx))^2}{2d} + \frac{3be(c + dx)(a + \operatorname{barctanh}(c + dx))^2}{2d} - \frac{e(a + \operatorname{barctanh}(c + dx))^3}{2d} + \frac{e(c + dx)^2(a + \operatorname{barctanh}(c + dx))^3}{2d} - \frac{3b^2e(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1-c-dx}\right)}{d} - \frac{3b^3e \operatorname{PolyLog}\left(2, -\frac{1+c+dx}{1-c-dx}\right)}{2d}$$

output

```
3/2*b*e*(a+b*arctanh(d*x+c))^2/d+3/2*b*e*(d*x+c)*(a+b*arctanh(d*x+c))^2/d-
1/2*e*(a+b*arctanh(d*x+c))^3/d+1/2*e*(d*x+c)^2*(a+b*arctanh(d*x+c))^3/d-3*
b^2*e*(a+b*arctanh(d*x+c))*ln(2/(-d*x-c+1))/d-3/2*b^3*e*polylog(2,-(d*x+c
+1)/(-d*x-c+1))/d
```

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.38

$$\int (ce + dex)(a + b \operatorname{arctanh}(c + dx))^3 dx$$

$$= \frac{e \left(6b^2(-1 + c + dx)(b + a(1 + c + dx)) \operatorname{arctanh}(c + dx)^2 + 2b^3(-1 + c^2 + 2cdx + d^2x^2) \operatorname{arctanh}(c + dx) \right)}{4d}$$

input

```
Integrate[(c*e + d*e*x)*(a + b*ArcTanh[c + d*x])^3,x]
```

output

```
(e*(6*b^2*(-1 + c + d*x)*(b + a*(1 + c + d*x))*ArcTanh[c + d*x]^2 + 2*b^3*(-1 + c^2 + 2*c*d*x + d^2*x^2)*ArcTanh[c + d*x]^3 + 6*b*ArcTanh[c + d*x]*(a*(c + d*x)*(2*b + a*c + a*d*x) - 2*b^2*Log[1 + E^(-2*ArcTanh[c + d*x])])) + a*(6*a*b*c + 2*a^2*c^2 + 6*a*b*d*x + 4*a^2*c*d*x + 2*a^2*d^2*x^2 + 3*a*b*Log[1 - c - d*x] - 3*a*b*Log[1 + c + d*x] - 12*b^2*Log[1/Sqrt[1 - (c + d*x)^2]]) + 6*b^3*PolyLog[2, -E^(-2*ArcTanh[c + d*x])]))/(4*d)
```

Rubi [A] (verified)

Time = 1.01 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.89, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.476$, Rules used = {6657, 27, 6452, 6542, 6436, 6510, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ce + dex)(a + b \operatorname{arctanh}(c + dx))^3 dx$$

$$\downarrow \text{6657}$$

$$\frac{\int e(c + dx)(a + b \operatorname{arctanh}(c + dx))^3 d(c + dx)}{d}$$

$$\downarrow \text{27}$$

$$\frac{e \int (c + dx)(a + b \operatorname{arctanh}(c + dx))^3 d(c + dx)}{d}$$

$$\begin{aligned} & \downarrow 6452 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+\operatorname{barctanh}(c+dx))^3 - \frac{3}{2}b \int \frac{(c+dx)^2(a+\operatorname{barctanh}(c+dx))^2}{1-(c+dx)^2} d(c+dx)\right)}{d} \\ & \downarrow 6542 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+\operatorname{barctanh}(c+dx))^3 - \frac{3}{2}b\left(\int \frac{(a+\operatorname{barctanh}(c+dx))^2}{1-(c+dx)^2} d(c+dx) - \int (a+\operatorname{barctanh}(c+dx))^2 d(c+dx)\right)\right)}{d} \\ & \downarrow 6436 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+\operatorname{barctanh}(c+dx))^3 - \frac{3}{2}b\left(2b \int \frac{(c+dx)(a+\operatorname{barctanh}(c+dx))}{1-(c+dx)^2} d(c+dx) + \int \frac{(a+\operatorname{barctanh}(c+dx))^2}{1-(c+dx)^2} d(c+dx)\right)\right)}{d} \\ & \downarrow 6510 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+\operatorname{barctanh}(c+dx))^3 - \frac{3}{2}b\left(2b \int \frac{(c+dx)(a+\operatorname{barctanh}(c+dx))}{1-(c+dx)^2} d(c+dx) + \frac{(a+\operatorname{barctanh}(c+dx))^3}{3b} - (c+dx)\right)\right)}{d} \\ & \downarrow 6546 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+\operatorname{barctanh}(c+dx))^3 - \frac{3}{2}b\left(2b\left(\int \frac{a+\operatorname{barctanh}(c+dx)}{-c-dx+1} d(c+dx) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{2b}\right) + \frac{(a+\operatorname{barctanh}(c+dx))^3}{3b}\right)\right)}{d} \\ & \downarrow 6470 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+\operatorname{barctanh}(c+dx))^3 - \frac{3}{2}b\left(2b\left(-b \int \frac{\log\left(\frac{2}{-c-dx+1}\right)}{1-(c+dx)^2} d(c+dx) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{2b} + \log\left(\frac{2}{-c-dx+1}\right)\right)\right)\right)}{d} \\ & \downarrow 2849 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+\operatorname{barctanh}(c+dx))^3 - \frac{3}{2}b\left(2b\left(b \int \frac{\log\left(\frac{2}{-c-dx+1}\right)}{1-\frac{2}{-c-dx+1}} d\frac{1}{-c-dx+1} - \frac{(a+\operatorname{barctanh}(c+dx))^2}{2b} + \log\left(\frac{2}{-c-dx+1}\right)\right)\right)\right)}{d} \\ & \downarrow 2752 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+\operatorname{barctanh}(c+dx))^3 - \frac{3}{2}b\left(2b\left(-\frac{(a+\operatorname{barctanh}(c+dx))^2}{2b} + \log\left(\frac{2}{-c-dx+1}\right)\right)(a+\operatorname{barctanh}(c+dx)) + \log\left(\frac{2}{-c-dx+1}\right)\right)\right)}{d} \end{aligned}$$

input `Int[(c*e + d*e*x)*(a + b*ArcTanh[c + d*x])^3,x]`

output `(e*(((c + d*x)^2*(a + b*ArcTanh[c + d*x])^3)/2 - (3*b*(-((c + d*x)*(a + b*ArcTanh[c + d*x])^2) + (a + b*ArcTanh[c + d*x])^3/(3*b) + 2*b*(-1/2*(a + b*ArcTanh[c + d*x])^2/b + (a + b*ArcTanh[c + d*x])*Log[2/(1 - c - d*x)] + (b*PolyLog[2, 1 - 2/(1 - c - d*x)]))/2))/2))/d`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 6436 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^p, x_Symbol] := Simp[x*(a + b*ArcTanh[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`

rule 6452 `Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^p*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6470 $\text{Int}[\{(a_.) + \text{ArcTanh}[(c_.)(x_)](b_.)\}^{(p_.)}/\{(d_.) + (e_.)(x_)\}, x_Symbol] \rightarrow \text{Simp}[(-a + b*\text{ArcTanh}[c*x])^p*(\text{Log}[2/(1 + e*(x/d))]/e), x] + \text{Simp}[b*c*(p/e) \text{Int}[(a + b*\text{ArcTanh}[c*x])^{(p-1)}*(\text{Log}[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{IGtQ}[p, 0] \&\& \text{EqQ}[c^2*d^2 - e^2, 0]$

rule 6510 $\text{Int}[\{(a_.) + \text{ArcTanh}[(c_.)(x_)](b_.)\}^{(p_.)}/\{(d_.) + (e_.)(x_)^2\}, x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^{(p+1)}/(b*c*d*(p+1)), x] /; \text{FreeQ}\{a, b, c, d, e, p\}, x] \&\& \text{EqQ}[c^2*d + e, 0] \&\& \text{NeQ}[p, -1]$

rule 6542 $\text{Int}[\{((a_.) + \text{ArcTanh}[(c_.)(x_)](b_.))^{(p_.)}*((f_.)(x_)^m)/\{(d_.) + (e_.)(x_)^2\}, x_Symbol] \rightarrow \text{Simp}[f^2/e \text{Int}[(f*x)^{(m-2)}*(a + b*\text{ArcTanh}[c*x])^p, x], x] - \text{Simp}[d*(f^2/e) \text{Int}[(f*x)^{(m-2)}*((a + b*\text{ArcTanh}[c*x])^p/(d + e*x^2)), x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x] \&\& \text{GtQ}[p, 0] \&\& \text{GtQ}[m, 1]$

rule 6546 $\text{Int}[\{((a_.) + \text{ArcTanh}[(c_.)(x_)](b_.))^{(p_.)}*(x_)/\{(d_.) + (e_.)(x_)^2\}, x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^{(p+1)}/(b*e*(p+1)), x] + \text{Simp}[1/(c*d) \text{Int}[(a + b*\text{ArcTanh}[c*x])^p/(1 - c*x), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{EqQ}[c^2*d + e, 0] \&\& \text{IGtQ}[p, 0]$

rule 6657 $\text{Int}[\{(a_.) + \text{ArcTanh}[(c_.) + (d_.)(x_)](b_.)\}^{(p_.)}*((e_.) + (f_.)(x_)^m), x_Symbol] \rightarrow \text{Simp}[1/d \text{Subst}[\text{Int}[(f*(x/d))^m*(a + b*\text{ArcTanh}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m\}, x] \&\& \text{EqQ}[d*e - c*f, 0] \&\& \text{IGtQ}[p, 0]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 930 vs. $2(150) = 300$.

Time = 3.31 (sec) , antiderivative size = 931, normalized size of antiderivative = 5.82

| method | result |
|-------------------|--|
| risch | $\frac{eda^3x^2}{2} + \frac{eb^3\ln(-dx-c+1)^3}{16d} - \frac{3e\ln(-dx-c+1)^2b^3}{8d} + \frac{3eb^3\ln(-dx-c+1)^2x}{8} + \frac{3eb^3\operatorname{dilog}\left(-\frac{dx}{2}-\frac{c}{2}+\frac{1}{2}\right)}{2d} -$ |
| derivatividivides | Expression too large to display |
| default | Expression too large to display |
| parts | Expression too large to display |

input `int((d*e*x+c*e)*(a+b*arctanh(d*x+c))^3,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/2*e*d*a^3*x^2+1/16*e/d*b^3*\ln(-d*x-c+1)^3-3/8*e/d*\ln(-d*x-c+1)^2*b^3+3/8 \\ & *e*b^3*\ln(-d*x-c+1)^2*x+3/2*e*b^3/d*\operatorname{dilog}(-1/2*d*x-1/2*c+1/2)-3/4*e*d*\ln(- \\ & d*x-c+1)*a^2*b*x^2-3/2*e/d*a*b^2*\ln(-d*x-c+1)*c+3/4*e*\ln(-d*x-c+1)^2*a*b^2 \\ & *c*x-3/2*e*\ln(-d*x-c+1)*a^2*b*c*x+3/4*b*a^2*e/d*\ln(-d*x-c+1)*c^2+3/2*b^2*a \\ & *e/d*\ln(-d*x-c+1)*c+3/8*e/d*\ln(-d*x-c+1)^2*a*b^2*c^2-3/4*e/d*\ln(-d*x-c+1)* \\ & a^2*b*c^2+3/8*e*d*\ln(-d*x-c+1)^2*a*b^2*x^2-3/4*b*a^2*e/d*\ln(-d*x-c+1)+3/2* \\ & b^2*a*e/d*\ln(-d*x-c+1)-3/2*e*b^3/d*\ln(-d*x-c+1)*\ln(1/2*d*x+1/2*c+1/2)+3/2* \\ & e*b^3/d*\ln(1/2*d*x+1/2*c+1/2)*\ln(-1/2*d*x-1/2*c+1/2)-1/8*e*\ln(-d*x-c+1)^3* \\ & b^3*c*x-3/2*e*a*b^2*\ln(-d*x-c+1)*x+3/2*e/d*\ln(-d*x-c+1)*a*b^2-1/16*e*d*\ln(\\ & -d*x-c+1)^3*b^3*x^2-1/16*e/d*\ln(-d*x-c+1)^3*b^3*c^2+3/8*e/d*b^3*\ln(-d*x-c+ \\ & 1)^2*c-3/8*e/d*a*b^2*\ln(-d*x-c+1)^2+3/4*e/d*a^2*b*\ln(-d*x-c+1)+e*a^3*c*x+3 \\ & /2*e*a^2*b*x+1/16*b^3*e*(d^2*x^2+2*c*d*x+c^2-1)/d*\ln(d*x+c+1)^3+3/16*e*b^2 \\ & *(-b*d^2*x^2*\ln(-d*x-c+1)+2*d^2*a*x^2-2*b*d*x*\ln(-d*x-c+1)*c+4*a*d*x*c-\ln(\\ & -d*x-c+1)*b*c^2+2*a*c^2+2*b*d*x+2*b*c+b*\ln(-d*x-c+1)-2*a+2*b)/d*\ln(d*x+c+1 \\ &)^2+(3/16*b^3*e*(d^2*x^2+2*c*d*x+c^2-1)/d*\ln(-d*x-c+1)^2-3/4*e*b^2*x*(a*d* \\ & x+2*a*c+b)*\ln(-d*x-c+1)+3/4*b*e*(d^2*x^2*a^2+2*a^2*c*d*x-b*a*\ln(-d*x-c+1)* \\ & c^2+2*a*b*d*x-b^2*\ln(-d*x-c+1)*c+b*a*\ln(-d*x-c+1)+b^2*\ln(-d*x-c+1))/d)*\ln(\\ & d*x+c+1)-1/2*e/d*a^3-3/2*e/d*a^2*b+1/2*e/d*c^2*a^3+3/2*e/d*a^2*b*c \end{aligned}$$

Fricas [F]

$$\int (ce + dex)(a + b \operatorname{arctanh}(c + dx))^3 dx = \int (dex + ce)(b \operatorname{artanh}(dx + c) + a)^3 dx$$

input `integrate((d*e*x+c*e)*(a+b*arctanh(d*x+c))^3,x, algorithm="fricas")`

output `integral(a^3*d*e*x + a^3*c*e + (b^3*d*e*x + b^3*c*e)*arctanh(d*x + c)^3 + 3*(a*b^2*d*e*x + a*b^2*c*e)*arctanh(d*x + c)^2 + 3*(a^2*b*d*e*x + a^2*b*c*e)*arctanh(d*x + c), x)`

Sympy [F]

$$\begin{aligned} \int (ce + dex)(a + b \operatorname{arctanh}(c + dx))^3 dx = e & \left(\int a^3 c dx + \int a^3 dx dx \right. \\ & + \int b^3 c \operatorname{atanh}^3(c + dx) dx \\ & + \int 3ab^2 c \operatorname{atanh}^2(c + dx) dx \\ & + \int 3a^2 bc \operatorname{atanh}(c + dx) dx \\ & + \int b^3 dx \operatorname{atanh}^3(c + dx) dx \\ & + \int 3ab^2 dx \operatorname{atanh}^2(c + dx) dx \\ & \left. + \int 3a^2 b dx \operatorname{atanh}(c + dx) dx \right) \end{aligned}$$

input `integrate((d*e*x+c*e)*(a+b*atanh(d*x+c))**3,x)`

output `e*(Integral(a**3*c, x) + Integral(a**3*d*x, x) + Integral(b**3*c*atanh(c + d*x)**3, x) + Integral(3*a*b**2*c*atanh(c + d*x)**2, x) + Integral(3*a**2*b*c*atanh(c + d*x), x) + Integral(b**3*d*x*atanh(c + d*x)**3, x) + Integral(3*a*b**2*d*x*atanh(c + d*x)**2, x) + Integral(3*a**2*b*d*x*atanh(c + d*x), x))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 629 vs. $2(142) = 284$.

Time = 0.25 (sec) , antiderivative size = 629, normalized size of antiderivative = 3.93

$$\int (ce + dex)(a + \operatorname{arctanh}(c + dx))^3 dx = \text{Too large to display}$$

input `integrate((d*e*x+c*e)*(a+b*arctanh(d*x+c))^3,x, algorithm="maxima")`

output

$$\begin{aligned} & 1/2*a^3*d*e*x^2 + 3/4*(2*x^2*arctanh(d*x + c) + d*(2*x/d^2 - (c^2 + 2*c + 1)*\log(d*x + c + 1)/d^3 + (c^2 - 2*c + 1)*\log(d*x + c - 1)/d^3))*a^2*b*d*e \\ & + a^3*c*e*x + 3/2*(2*(d*x + c)*arctanh(d*x + c) + \log(-(d*x + c)^2 + 1))* \\ & a^2*b*c*e/d + 3/2*(\log(d*x + c + 1)*\log(-1/2*d*x - 1/2*c + 1/2) + \operatorname{dilog}(1/ \\ & 2*d*x + 1/2*c + 1/2))*b^3*e/d + 3/2*(c*e + e)*a*b^2*\log(d*x + c + 1)/d - 3 \\ & /2*(c*e - e)*a*b^2*\log(d*x + c - 1)/d + 1/16*(24*a*b^2*d*e*x*\log(d*x + c + \\ & 1) + (b^3*d^2*e*x^2 + 2*b^3*c*d*e*x + (c^2*e - e)*b^3)*\log(d*x + c + 1)^3 \\ & - (b^3*d^2*e*x^2 + 2*b^3*c*d*e*x + (c^2*e - e)*b^3)*\log(-d*x - c + 1)^3 + \\ & 6*(a*b^2*d^2*e*x^2 + (c^2*e - e)*a*b^2 + (c*e + e)*b^3 + (2*a*b^2*c*d*e + \\ & b^3*d*e)*x)*\log(d*x + c + 1)^2 + 3*(2*a*b^2*d^2*e*x^2 + 2*(c^2*e - e)*a*b \\ & ^2 + 2*(c*e - e)*b^3 + 2*(2*a*b^2*c*d*e + b^3*d*e)*x + (b^3*d^2*e*x^2 + 2* \\ & b^3*c*d*e*x + (c^2*e - e)*b^3)*\log(d*x + c + 1))*\log(-d*x - c + 1)^2 - 3*(\\ & 8*a*b^2*d*e*x + (b^3*d^2*e*x^2 + 2*b^3*c*d*e*x + (c^2*e - e)*b^3)*\log(d*x \\ & + c + 1)^2 + 4*(a*b^2*d^2*e*x^2 + (c^2*e - e)*a*b^2 + (c*e + e)*b^3 + (2*a \\ & *b^2*c*d*e + b^3*d*e)*x)*\log(d*x + c + 1))*\log(-d*x - c + 1))/d \end{aligned}$$
Giac [F]

$$\int (ce + dex)(a + \operatorname{arctanh}(c + dx))^3 dx = \int (dex + ce)(\operatorname{arctanh}(dx + c) + a)^3 dx$$

input `integrate((d*e*x+c*e)*(a+b*arctanh(d*x+c))^3,x, algorithm="giac")`

output `integrate((d*e*x + c*e)*(b*arctanh(d*x + c) + a)^3, x)`

3.25 $\int \frac{(a+b\operatorname{arctanh}(c+dx))^3}{ce+dex} dx$

| | |
|---------------------------------------|-----|
| Optimal result | 251 |
| Mathematica [C] (verified) | 252 |
| Rubi [A] (verified) | 253 |
| Maple [C] (warning: unable to verify) | 255 |
| Fricas [F] | 256 |
| Sympy [F] | 257 |
| Maxima [F] | 257 |
| Giac [F] | 257 |
| Mupad [F(-1)] | 258 |
| Reduce [F] | 258 |

Optimal result

Integrand size = 23, antiderivative size = 257

$$\int \frac{(a + b\operatorname{arctanh}(c + dx))^3}{ce + dex} dx = \frac{2(a + b\operatorname{arctanh}(c + dx))^3 \operatorname{arctanh}\left(1 - \frac{2}{1-c-dx}\right)}{de} - \frac{3b(a + b\operatorname{arctanh}(c + dx))^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-c-dx}\right)}{2de} + \frac{3b(a + b\operatorname{arctanh}(c + dx))^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1-c-dx}\right)}{2de} + \frac{3b^2(a + b\operatorname{arctanh}(c + dx)) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-c-dx}\right)}{2de} - \frac{3b^2(a + b\operatorname{arctanh}(c + dx)) \operatorname{PolyLog}\left(3, -1 + \frac{2}{1-c-dx}\right)}{2de} - \frac{3b^3 \operatorname{PolyLog}\left(4, 1 - \frac{2}{1-c-dx}\right)}{4de} + \frac{3b^3 \operatorname{PolyLog}\left(4, -1 + \frac{2}{1-c-dx}\right)}{4de}$$

output

```
-2*(a+b*arctanh(d*x+c))^3*arctanh(-1+2/(-d*x-c+1))/d/e-3/2*b*(a+b*arctanh(d*x+c))^2*polylog(2,1-2/(-d*x-c+1))/d/e+3/2*b*(a+b*arctanh(d*x+c))^2*polylog(2,-1+2/(-d*x-c+1))/d/e+3/2*b^2*(a+b*arctanh(d*x+c))*polylog(3,1-2/(-d*x-c+1))/d/e-3/2*b^2*(a+b*arctanh(d*x+c))*polylog(3,-1+2/(-d*x-c+1))/d/e-3/4*b^3*polylog(4,1-2/(-d*x-c+1))/d/e+3/4*b^3*polylog(4,-1+2/(-d*x-c+1))/d/e
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.44 (sec) , antiderivative size = 581, normalized size of antiderivative = 2.26

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{ce + dex} dx$$

$$= \frac{4a^3 \log(c + dx) + 12a^2 b \operatorname{arctanh}(c + dx) \left(-\log\left(\frac{1}{\sqrt{1-(c+dx)^2}}\right) + \log\left(\frac{i(c+dx)}{\sqrt{1-(c+dx)^2}}\right) \right) - \frac{3}{2}a^2 b \left(\pi^2 - 4i \operatorname{arctanh}(c + dx) \right)}{d}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])^3/(c*e + d*e*x),x]
```

output

```
(4*a^3*Log[c + d*x] + 12*a^2*b*ArcTanh[c + d*x]*(-Log[1/Sqrt[1 - (c + d*x)^2]] + Log[(I*(c + d*x))/Sqrt[1 - (c + d*x)^2]]) - (3*a^2*b*(Pi^2 - (4*I)*Pi*ArcTanh[c + d*x] - 8*ArcTanh[c + d*x]^2 - 8*ArcTanh[c + d*x]*Log[1 - E^(-2*ArcTanh[c + d*x])]) + (4*I)*Pi*Log[1 + E^(2*ArcTanh[c + d*x])]) + 8*ArcTanh[c + d*x]*Log[1 + E^(2*ArcTanh[c + d*x])] - (4*I)*Pi*Log[2/Sqrt[1 - (c + d*x)^2]] - 8*ArcTanh[c + d*x]*Log[2/Sqrt[1 - (c + d*x)^2]] + 8*ArcTanh[c + d*x]*Log[((2*I)*(c + d*x))/Sqrt[1 - (c + d*x)^2]] + 4*PolyLog[2, E^(-2*ArcTanh[c + d*x])] + 4*PolyLog[2, -E^(2*ArcTanh[c + d*x])])/2 + 6*a*b^2*(2*ArcTanh[c + d*x]^2*Log[1 - E^(-2*ArcTanh[c + d*x])] - 2*ArcTanh[c + d*x]^2*Log[1 + E^(-2*ArcTanh[c + d*x])] + 2*ArcTanh[c + d*x]*PolyLog[2, -E^(-2*ArcTanh[c + d*x])] - 2*ArcTanh[c + d*x]*PolyLog[2, E^(-2*ArcTanh[c + d*x])] + PolyLog[3, -E^(-2*ArcTanh[c + d*x])] - PolyLog[3, E^(-2*ArcTanh[c + d*x])]) + b^3*(4*ArcTanh[c + d*x]^3*Log[1 - E^(-2*ArcTanh[c + d*x])] - 4*ArcTanh[c + d*x]^3*Log[1 + E^(-2*ArcTanh[c + d*x])] + 6*ArcTanh[c + d*x]^2*PolyLog[2, -E^(-2*ArcTanh[c + d*x])] - 6*ArcTanh[c + d*x]^2*PolyLog[2, E^(-2*ArcTanh[c + d*x])] + 6*ArcTanh[c + d*x]*PolyLog[3, -E^(-2*ArcTanh[c + d*x])] - 6*ArcTanh[c + d*x]*PolyLog[3, E^(-2*ArcTanh[c + d*x])] + 3*PolyLog[4, -E^(-2*ArcTanh[c + d*x])] - 3*PolyLog[4, E^(-2*ArcTanh[c + d*x])]))/(4*d*e)
```

Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 231, normalized size of antiderivative = 0.90, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$, Rules used = {6657, 27, 6448, 6614, 6620, 6624, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{ce + dex} dx$$

$$\downarrow \text{6657}$$

$$\frac{\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{e(c + dx)} d(c + dx)}{d}$$

$$\downarrow \text{27}$$

$$\frac{\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{c + dx} d(c + dx)}{de}$$

$$\downarrow \text{6448}$$

$$\frac{2 \operatorname{arctanh}\left(1 - \frac{2}{-c - dx + 1}\right) (a + b \operatorname{arctanh}(c + dx))^3 - 6b \int \frac{(a + b \operatorname{arctanh}(c + dx))^2 \operatorname{arctanh}\left(1 - \frac{2}{-c - dx + 1}\right)}{1 - (c + dx)^2} d(c + dx)}{de}$$

$$\downarrow \text{6614}$$

$$\frac{2 \operatorname{arctanh}\left(1 - \frac{2}{-c - dx + 1}\right) (a + b \operatorname{arctanh}(c + dx))^3 - 6b \left(\frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(c + dx))^2 \log\left(2 - \frac{2}{-c - dx + 1}\right)}{1 - (c + dx)^2} d(c + dx) - \frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{1 - (c + dx)^2} d(c + dx) \right)}{de}$$

$$\downarrow \text{6620}$$

$$\frac{2 \operatorname{arctanh}\left(1 - \frac{2}{-c - dx + 1}\right) (a + b \operatorname{arctanh}(c + dx))^3 - 6b \left(\frac{1}{2} \operatorname{PolyLog}\left(2, 1 - \frac{2}{-c - dx + 1}\right) (a + b \operatorname{arctanh}(c + dx))^2 - \frac{1}{2} \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{1 - (c + dx)^2} d(c + dx) \right)}{de}$$

$$\downarrow \text{6624}$$

$$2\operatorname{arctanh}\left(1 - \frac{2}{-c-dx+1}\right) (a + \operatorname{barctanh}(c + dx))^3 - 6b \left(\frac{1}{2} \left(\frac{1}{2} \operatorname{PolyLog}\left(2, 1 - \frac{2}{-c-dx+1}\right) (a + \operatorname{barctanh}(c + dx))^2\right)\right)$$

↓ 7164

$$2\operatorname{arctanh}\left(1 - \frac{2}{-c-dx+1}\right) (a + \operatorname{barctanh}(c + dx))^3 - 6b \left(\frac{1}{2} \left(\frac{1}{2} \operatorname{PolyLog}\left(2, 1 - \frac{2}{-c-dx+1}\right) (a + \operatorname{barctanh}(c + dx))^2\right)\right)$$

input

```
Int[(a + b*ArcTanh[c + d*x])^3/(c*e + d*e*x), x]
```

output

```
(2*(a + b*ArcTanh[c + d*x])^3*ArcTanh[1 - 2/(1 - c - d*x)] - 6*b*(((a + b
*ArcTanh[c + d*x])^2*PolyLog[2, 1 - 2/(1 - c - d*x)]/2 - b*(((a + b*ArcTa
nh[c + d*x])*PolyLog[3, 1 - 2/(1 - c - d*x)]/2 - (b*PolyLog[4, 1 - 2/(1 -
c - d*x]))/4))/2 + (-1/2*((a + b*ArcTanh[c + d*x])^2*PolyLog[2, -1 + 2/(1
- c - d*x)]) + b*(((a + b*ArcTanh[c + d*x])*PolyLog[3, -1 + 2/(1 - c - d*
x]))/2 - (b*PolyLog[4, -1 + 2/(1 - c - d*x)]/4))/2))/(d*e)
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

rule 6448

```
Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_)/(x_), x_Symbol] := Simp[2*(a +
b*ArcTanh[c*x])^p*ArcTanh[1 - 2/(1 - c*x)], x] - Simp[2*b*c*p Int[(a + b
*ArcTanh[c*x])^(p - 1)*(ArcTanh[1 - 2/(1 - c*x)]/(1 - c^2*x^2)), x], x] /;
FreeQ[{a, b, c}, x] && IGtQ[p, 1]
```

rule 6614

```
Int[(ArcTanh[u_] * ((a_.) + ArcTanh[(c_.)*(x_)]) * (b_.))^(p_.) / ((d_.) + (e_.)*(
x_)^2), x_Symbol] := Simp[1/2 Int[Log[1 + u] * ((a + b*ArcTanh[c*x])^p / (d +
e*x^2)), x], x] - Simp[1/2 Int[Log[1 - u] * ((a + b*ArcTanh[c*x])^p / (d + e
*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e,
0] && EqQ[u^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 6620

```
Int[(Log[u_]*((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.))/((d_) + (e_.)*(x_)^
2), x_Symbol] := Simp[(- (a + b*ArcTanh[c*x])^p)*(PolyLog[2, 1 - u]/(2*c*d))
, x] + Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(
d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d
+ e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 6624

```
Int[(((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^(p_.)*PolyLog[k_, u_]/((d_) + (e_
.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(PolyLog[k + 1, u]/(2*
c*d)), x] - Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[k + 1,
u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, k}, x] && IGtQ[p, 0] && E
qQ[c^2*d + e, 0] && EqQ[u^2 - (1 - 2/(1 - c*x))^2, 0]
```

rule 6657

```
Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)])*(b_.))^(p_.)*((e_.) + (f_.)*(x_)^(
m_.), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x
], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0]
&& IGtQ[p, 0]
```

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.53 (sec) , antiderivative size = 1442, normalized size of antiderivative = 5.61

| method | result | size |
|-------------------|---------------------------------|------|
| derivativedivides | Expression too large to display | 1442 |
| default | Expression too large to display | 1442 |
| parts | Expression too large to display | 1452 |

input

```
int((a+b*arctanh(d*x+c))^3/(d*e*x+c*e),x,method=_RETURNVERBOSE)
```


output

```

1/d*(a^3/e*ln(d*x+c)+b^3/e*(ln(d*x+c)*arctanh(d*x+c)^3-arctanh(d*x+c)^3*ln
((d*x+c+1)^2/(1-(d*x+c)^2)-1)+arctanh(d*x+c)^3*ln(1+(d*x+c+1)/(1-(d*x+c)^2
)^(1/2))+3*arctanh(d*x+c)^2*polylog(2,-(d*x+c+1)/(1-(d*x+c)^2)^(1/2))-6*ar
ctanh(d*x+c)*polylog(3,-(d*x+c+1)/(1-(d*x+c)^2)^(1/2))+6*polylog(4,-(d*x+c
+1)/(1-(d*x+c)^2)^(1/2))+arctanh(d*x+c)^3*ln(1-(d*x+c+1)/(1-(d*x+c)^2)^(1/
2))+3*arctanh(d*x+c)^2*polylog(2,(d*x+c+1)/(1-(d*x+c)^2)^(1/2))-6*arctanh(
d*x+c)*polylog(3,(d*x+c+1)/(1-(d*x+c)^2)^(1/2))+6*polylog(4,(d*x+c+1)/(1-(
d*x+c)^2)^(1/2))+1/2*I*Pi*csgn(I*(-(d*x+c+1)^2/((d*x+c)^2-1)-1)/(1-(d*x+c+
1)^2/((d*x+c)^2-1)))*(csgn(I*(-(d*x+c+1)^2/((d*x+c)^2-1)-1))*csgn(I/(1-(d*
x+c+1)^2/((d*x+c)^2-1)))-csgn(I*(-(d*x+c+1)^2/((d*x+c)^2-1)-1))*csgn(I*(-(
d*x+c+1)^2/((d*x+c)^2-1)-1)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))-csgn(I*(-(d*x+c
+1)^2/((d*x+c)^2-1)-1)/(1-(d*x+c+1)^2/((d*x+c)^2-1))*csgn(I/(1-(d*x+c+1)^
2/((d*x+c)^2-1)))+csgn(I*(-(d*x+c+1)^2/((d*x+c)^2-1)-1)/(1-(d*x+c+1)^2/((d
*x+c)^2-1)))^2)*arctanh(d*x+c)^3-3/2*arctanh(d*x+c)^2*polylog(2,-(d*x+c+1)
^2/(1-(d*x+c)^2))+3/2*arctanh(d*x+c)*polylog(3,-(d*x+c+1)^2/(1-(d*x+c)^2))
-3/4*polylog(4,-(d*x+c+1)^2/(1-(d*x+c)^2))+3*a*b^2/e*(ln(d*x+c)*arctanh(d
*x+c)^2-arctanh(d*x+c)*polylog(2,-(d*x+c+1)^2/(1-(d*x+c)^2))+1/2*polylog(3
,-(d*x+c+1)^2/(1-(d*x+c)^2))-arctanh(d*x+c)^2*ln((d*x+c+1)^2/(1-(d*x+c)^2
)-1)+arctanh(d*x+c)^2*ln(1+(d*x+c+1)/(1-(d*x+c)^2)^(1/2))+2*arctanh(d*x+c)*
polylog(2,-(d*x+c+1)/(1-(d*x+c)^2)^(1/2))-2*polylog(3,-(d*x+c+1)/(1-(d*...

```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{ce + dex} dx = \int \frac{(b \operatorname{arctanh}(dx + c) + a)^3}{dex + ce} dx$$

input

```
integrate((a+b*arctanh(d*x+c))^3/(d*e*x+c*e),x, algorithm="fricas")
```

output

```
integral((b^3*arctanh(d*x + c)^3 + 3*a*b^2*arctanh(d*x + c)^2 + 3*a^2*b*ar
ctanh(d*x + c) + a^3)/(d*e*x + c*e), x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{ce + dex} dx$$

$$= \frac{\int \frac{a^3}{c+dx} dx + \int \frac{b^3 \operatorname{atanh}^3(c+dx)}{c+dx} dx + \int \frac{3ab^2 \operatorname{atanh}^2(c+dx)}{c+dx} dx + \int \frac{3a^2b \operatorname{atanh}(c+dx)}{c+dx} dx}{e}$$

input `integrate((a+b*atanh(d*x+c))**3/(d*e*x+c*e), x)`

output `(Integral(a**3/(c + d*x), x) + Integral(b**3*atanh(c + d*x)**3/(c + d*x), x) + Integral(3*a*b**2*atanh(c + d*x)**2/(c + d*x), x) + Integral(3*a**2*b*atanh(c + d*x)/(c + d*x), x))/e`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{ce + dex} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^3}{dex + ce} dx$$

input `integrate((a+b*arctanh(d*x+c))^3/(d*e*x+c*e), x, algorithm="maxima")`

output `a^3*log(d*e*x + c*e)/(d*e) + integrate(1/8*b^3*(log(d*x + c + 1) - log(-d*x - c + 1))^3/(d*e*x + c*e) + 3/4*a*b^2*(log(d*x + c + 1) - log(-d*x - c + 1))^2/(d*e*x + c*e) + 3/2*a^2*b*(log(d*x + c + 1) - log(-d*x - c + 1))/(d*e*x + c*e), x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{ce + dex} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^3}{dex + ce} dx$$

input `integrate((a+b*arctanh(d*x+c))^3/(d*e*x+c*e), x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)^3/(d*e*x + c*e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{ce + dex} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^3}{ce + dex} dx$$

input `int((a + b*atanh(c + d*x))^3/(c*e + d*e*x), x)`

output `int((a + b*atanh(c + d*x))^3/(c*e + d*e*x), x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{ce + dex} dx$$

$$= \frac{3 \left(\int \frac{\operatorname{atanh}(dx+c)}{dx+c} dx \right) a^2 b d + \left(\int \frac{\operatorname{atanh}(dx+c)^3}{dx+c} dx \right) b^3 d + 3 \left(\int \frac{\operatorname{atanh}(dx+c)^2}{dx+c} dx \right) a b^2 d + \log(dx + c) a^3}{de}$$

input `int((a+b*atanh(d*x+c))^3/(d*e*x+c*e), x)`

output `(3*int(atanh(c + d*x)/(c + d*x), x)*a**2*b*d + int(atanh(c + d*x)**3/(c + d*x), x)*b**3*d + 3*int(atanh(c + d*x)**2/(c + d*x), x)*a*b**2*d + log(c + d*x)*a**3)/(d*e)`

3.26 $\int \frac{(a+b\operatorname{arctanh}(c+dx))^3}{(ce+dex)^2} dx$

| | |
|---------------------------------------|-----|
| Optimal result | 259 |
| Mathematica [A] (verified) | 260 |
| Rubi [A] (verified) | 260 |
| Maple [C] (warning: unable to verify) | 263 |
| Fricas [F] | 264 |
| Sympy [F] | 264 |
| Maxima [F] | 264 |
| Giac [F] | 265 |
| Mupad [F(-1)] | 265 |
| Reduce [F] | 266 |

Optimal result

Integrand size = 23, antiderivative size = 143

$$\int \frac{(a + b\operatorname{arctanh}(c + dx))^3}{(ce + dex)^2} dx = \frac{(a + b\operatorname{arctanh}(c + dx))^3}{de^2} - \frac{(a + b\operatorname{arctanh}(c + dx))^3}{de^2(c + dx)} + \frac{3b(a + b\operatorname{arctanh}(c + dx))^2 \log\left(2 - \frac{2}{1+c+dx}\right)}{de^2} - \frac{3b^2(a + b\operatorname{arctanh}(c + dx)) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1+c+dx}\right)}{de^2} - \frac{3b^3 \operatorname{PolyLog}\left(3, -1 + \frac{2}{1+c+dx}\right)}{2de^2}$$

output

```
(a+b*arctanh(d*x+c))^3/d/e^2-(a+b*arctanh(d*x+c))^3/d/e^2/(d*x+c)+3*b*(a+b
*arctanh(d*x+c))^2*ln(2-2/(d*x+c+1))/d/e^2-3*b^2*(a+b*arctanh(d*x+c))*poly
log(2,-1+2/(d*x+c+1))/d/e^2-3/2*b^3*polylog(3,-1+2/(d*x+c+1))/d/e^2
```

Mathematica [A] (verified)

Time = 0.45 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.62

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^2} dx$$

$$= -\frac{2a^3}{c+dx} - \frac{6a^2 b \operatorname{arctanh}(c+dx)}{c+dx} + 6a^2 b \log(c + dx) - 3a^2 b \log(1 - c^2 - 2cdx - d^2 x^2) + 6ab^2 (\operatorname{arctanh}(c + dx))$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])^3/(c*e + d*e*x)^2,x]
```

output

```
((-2*a^3)/(c + d*x) - (6*a^2*b*ArcTanh[c + d*x])/(c + d*x) + 6*a^2*b*Log[c + d*x] - 3*a^2*b*Log[1 - c^2 - 2*c*d*x - d^2*x^2] + 6*a*b^2*(ArcTanh[c + d*x]*((1 - (c + d*x)^(-1))*ArcTanh[c + d*x] + 2*Log[1 - E^(-2*ArcTanh[c + d*x])])) - PolyLog[2, E^(-2*ArcTanh[c + d*x])]) + 2*b^3*(ArcTanh[c + d*x]^2*((1 - (c + d*x)^(-1))*ArcTanh[c + d*x] + 3*Log[1 - E^(-2*ArcTanh[c + d*x])])) - 3*ArcTanh[c + d*x]*PolyLog[2, E^(-2*ArcTanh[c + d*x])]) - (3*PolyLog[3, E^(-2*ArcTanh[c + d*x])])]/(2*d*e^2)
```

Rubi [A] (verified)

Time = 0.86 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.90, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$, Rules used = {6657, 27, 6452, 6550, 6494, 6618, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^2} dx$$

$$\downarrow 6657$$

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{e^2(c + dx)^2} d(c + dx)$$

$$\downarrow 27$$

$$\begin{aligned}
 & \frac{\int \frac{(a+b\operatorname{arctanh}(c+dx))^3}{(c+dx)^2} d(c+dx)}{de^2} \\
 & \quad \downarrow \text{6452} \\
 & \frac{3b \int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{(c+dx)(1-(c+dx)^2)} d(c+dx) - \frac{(a+b\operatorname{arctanh}(c+dx))^3}{c+dx}}{de^2} \\
 & \quad \downarrow \text{6550} \\
 & \frac{3b \left(\int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{(c+dx)(c+dx+1)} d(c+dx) + \frac{(a+b\operatorname{arctanh}(c+dx))^3}{3b} \right) - \frac{(a+b\operatorname{arctanh}(c+dx))^3}{c+dx}}{de^2} \\
 & \quad \downarrow \text{6494} \\
 & \frac{3b \left(-2b \int \frac{(a+b\operatorname{arctanh}(c+dx)) \log\left(2 - \frac{2}{c+dx+1}\right)}{1-(c+dx)^2} d(c+dx) + \frac{(a+b\operatorname{arctanh}(c+dx))^3}{3b} + \log\left(2 - \frac{2}{c+dx+1}\right) (a+b\operatorname{arctanh}(c+dx)) \right)}{de^2} \\
 & \quad \downarrow \text{6618} \\
 & \frac{3b \left(-2b \left(\frac{1}{2} \operatorname{PolyLog}\left(2, \frac{2}{c+dx+1} - 1\right) (a+b\operatorname{arctanh}(c+dx)) - \frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, \frac{2}{c+dx+1} - 1\right)}{1-(c+dx)^2} d(c+dx) \right) + \frac{(a+b\operatorname{arctanh}(c+dx))^3}{3b} \right)}{de^2} \\
 & \quad \downarrow \text{7164} \\
 & \frac{3b \left(-2b \left(\frac{1}{2} \operatorname{PolyLog}\left(2, \frac{2}{c+dx+1} - 1\right) (a+b\operatorname{arctanh}(c+dx)) + \frac{1}{4} b \operatorname{PolyLog}\left(3, \frac{2}{c+dx+1} - 1\right) \right) + \frac{(a+b\operatorname{arctanh}(c+dx))^3}{3b} \right)}{de^2}
 \end{aligned}$$

input

```
Int[(a + b*ArcTanh[c + d*x])^3/(c*e + d*e*x)^2,x]
```

output

```
(-((a + b*ArcTanh[c + d*x])^3/(c + d*x)) + 3*b*((a + b*ArcTanh[c + d*x])^3/(3*b) + (a + b*ArcTanh[c + d*x])^2*Log[2 - 2/(1 + c + d*x)] - 2*b*((a + b*ArcTanh[c + d*x])*PolyLog[2, -1 + 2/(1 + c + d*x)]/2 + (b*PolyLog[3, -1 + 2/(1 + c + d*x)]/4)))/(d*e^2)
```

Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`
- rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`
- rule 6494 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((x_)*((d_) + (e_)*(x_))), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`
- rule 6550 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((x_)*((d_) + (e_)*(x_)^2)), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]`
- rule 6618 `Int[(Log[u_]*)((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)), x] - Simp[b*(p/2) Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 + c*x))^2, 0]`
- rule 6657 `Int[((a_) + ArcTanh[(c_) + (d_)*(x_)])*(b_)^(p_)*((e_) + (f_)*(x_)^(m_)), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]`

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.49 (sec) , antiderivative size = 1466, normalized size of antiderivative = 10.25

| method | result | size |
|-------------------|---------------------------------|------|
| derivativedivides | Expression too large to display | 1466 |
| default | Expression too large to display | 1466 |
| parts | Expression too large to display | 1474 |

input

```
int((a+b*arctanh(d*x+c))^3/(d*e*x+c*e)^2,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-a^3/e^2/(d*x+c)+b^3/e^2*(-1/(d*x+c)*arctanh(d*x+c)^3-3/2*arctanh(d*x+c)^2*ln(d*x+c-1)+3*ln(d*x+c)*arctanh(d*x+c)^2-3/2*arctanh(d*x+c)^2*ln(d*x+c+1)+3*arctanh(d*x+c)^2*ln((d*x+c+1)/(1-(d*x+c)^2)^(1/2))-arctanh(d*x+c)^3-3*arctanh(d*x+c)^2*ln((d*x+c+1)^2/(1-(d*x+c)^2)-1)+3*arctanh(d*x+c)^2*ln(1+(d*x+c+1)/(1-(d*x+c)^2)^(1/2))+6*arctanh(d*x+c)*polylog(2,-(d*x+c+1)/(1-(d*x+c)^2)^(1/2))-6*polylog(3,-(d*x+c+1)/(1-(d*x+c)^2)^(1/2))+3*arctanh(d*x+c)^2*ln(1-(d*x+c+1)/(1-(d*x+c)^2)^(1/2))+6*arctanh(d*x+c)*polylog(2,(d*x+c+1)/(1-(d*x+c)^2)^(1/2))-6*polylog(3,(d*x+c+1)/(1-(d*x+c)^2)^(1/2))+3/4*(-I*Pi*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))+I*Pi*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))^3+2*I*Pi+2*I*Pi*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))^3+2*I*Pi*csgn(I*(-(d*x+c+1)^2/((d*x+c)^2-1)-1))*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))*csgn(I*(-(d*x+c+1)^2/((d*x+c)^2-1)-1)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))-2*I*Pi*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))^2+2*I*Pi*csgn(I*(d*x+c+1)/(1-(d*x+c)^2)^(1/2))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1))^2+I*Pi*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))^2+I*Pi*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1))^3+I*Pi*csgn(I*(d*x+c+1)/(1-(d*x+c)^2)^(1/2))^2*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1))-2*I*Pi*csgn(I*(-(d*x+c+1)^2/((d*x+c)^2-1)-1))*csgn(I*(-(d*x+c+1)^2/((d*x+c)^2-1)-1)/(1-(d*x+c+1)^2/((d*x+c...
```


Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^2} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^3}{(dex + ce)^2} dx$$

input `integrate((a+b*arctanh(d*x+c))^3/(d*e*x+c*e)^2,x, algorithm="fricas")`

output `integral((b^3*arctanh(d*x + c)^3 + 3*a*b^2*arctanh(d*x + c)^2 + 3*a^2*b*arctanh(d*x + c) + a^3)/(d^2*e^2*x^2 + 2*c*d*e^2*x + c^2*e^2), x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^2} dx$$

$$= \frac{\int \frac{a^3}{c^2+2cdx+d^2x^2} dx + \int \frac{b^3 \operatorname{atanh}^3(c+dx)}{c^2+2cdx+d^2x^2} dx + \int \frac{3ab^2 \operatorname{atanh}^2(c+dx)}{c^2+2cdx+d^2x^2} dx + \int \frac{3a^2b \operatorname{atanh}(c+dx)}{c^2+2cdx+d^2x^2} dx}{e^2}$$

input `integrate((a+b*atanh(d*x+c))**3/(d*e*x+c*e)**2,x)`

output `(Integral(a**3/(c**2 + 2*c*d*x + d**2*x**2), x) + Integral(b**3*atanh(c + d*x)**3/(c**2 + 2*c*d*x + d**2*x**2), x) + Integral(3*a*b**2*atanh(c + d*x)**2/(c**2 + 2*c*d*x + d**2*x**2), x) + Integral(3*a**2*b*atanh(c + d*x)/(c**2 + 2*c*d*x + d**2*x**2), x))/e**2`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^2} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^3}{(dex + ce)^2} dx$$

input `integrate((a+b*arctanh(d*x+c))^3/(d*e*x+c*e)^2,x, algorithm="maxima")`

output

```
-3/2*(d*(log(d*x + c + 1)/(d^2*e^2) - 2*log(d*x + c)/(d^2*e^2) + log(d*x +
c - 1)/(d^2*e^2)) + 2*arctanh(d*x + c)/(d^2*e^2*x + c*d*e^2))*a^2*b - a^3
/(d^2*e^2*x + c*d*e^2) - 1/8*((b^3*d*x + b^3*(c - 1))*log(-d*x - c + 1)^3
+ 3*(2*a*b^2 + (b^3*d*x + b^3*(c + 1))*log(d*x + c + 1))*log(-d*x - c + 1)
^2)/(d^2*e^2*x + c*d*e^2) - integrate(-1/8*((b^3*d*x + b^3*(c - 1))*log(d*
x + c + 1)^3 + 6*(a*b^2*d*x + a*b^2*(c - 1))*log(d*x + c + 1)^2 + 3*(4*a*b
^2*d*x + 4*a*b^2*c - (b^3*d*x + b^3*(c - 1))*log(d*x + c + 1)^2 + 2*(b^3*d
^2*x^2 + (c^2 + c)*b^3 - 2*a*b^2*(c - 1) + ((2*c*d + d)*b^3 - 2*a*b^2*d)*x
)*log(d*x + c + 1))*log(-d*x - c + 1))/(d^3*e^2*x^3 + c^3*e^2 - c^2*e^2 +
(3*c*d^2*e^2 - d^2*e^2)*x^2 + (3*c^2*d*e^2 - 2*c*d*e^2)*x), x)
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^2} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^3}{(dex + ce)^2} dx$$

input

```
integrate((a+b*arctanh(d*x+c))^3/(d*e*x+c*e)^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(d*x + c) + a)^3/(d*e*x + c*e)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^2} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^3}{(ce + dex)^2} dx$$

input

```
int((a + b*atanh(c + d*x))^3/(c*e + d*e*x)^2,x)
```

output

```
int((a + b*atanh(c + d*x))^3/(c*e + d*e*x)^2, x)
```


$$3.27 \quad \int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^3} dx$$

| | |
|---------------------------------------|-----|
| Optimal result | 267 |
| Mathematica [C] (verified) | 268 |
| Rubi [A] (verified) | 268 |
| Maple [C] (warning: unable to verify) | 271 |
| Fricas [F] | 272 |
| Sympy [F] | 273 |
| Maxima [F] | 273 |
| Giac [F] | 274 |
| Mupad [F(-1)] | 274 |
| Reduce [F] | 275 |

Optimal result

Integrand size = 23, antiderivative size = 166

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^3} dx &= \frac{3b(a + b \operatorname{arctanh}(c + dx))^2}{2de^3} \\ &\quad - \frac{3b(a + b \operatorname{arctanh}(c + dx))^2}{2de^3(c + dx)} \\ &\quad + \frac{(a + b \operatorname{arctanh}(c + dx))^3}{2de^3} - \frac{(a + b \operatorname{arctanh}(c + dx))^3}{2de^3(c + dx)^2} \\ &\quad + \frac{3b^2(a + b \operatorname{arctanh}(c + dx)) \log\left(2 - \frac{2}{1+c+dx}\right)}{de^3} \\ &\quad - \frac{3b^3 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1+c+dx}\right)}{2de^3} \end{aligned}$$

output

```
3/2*b*(a+b*arctanh(d*x+c))^2/d/e^3-3/2*b*(a+b*arctanh(d*x+c))^2/d/e^3/(d*x
+c)+1/2*(a+b*arctanh(d*x+c))^3/d/e^3-1/2*(a+b*arctanh(d*x+c))^3/d/e^3/(d*x
+c)^2+3*b^2*(a+b*arctanh(d*x+c))*ln(2-2/(d*x+c+1))/d/e^3-3/2*b^3*polylog(2
,-1+2/(d*x+c+1))/d/e^3
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.93 (sec) , antiderivative size = 335, normalized size of antiderivative = 2.02

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^3} dx$$

$$= \frac{-4a^3 - 12a^2bc + ib^3c^3\pi^3 - 12a^2bdx + 2ib^3c^2d\pi^3x + ib^3cd^2\pi^3x^2 + 12b^2(-1 + c + dx)(b(c + dx) + a(1 +$$

input `Integrate[(a + b*ArcTanh[c + d*x])^3/(c*e + d*e*x)^3,x]`

output

```
(-4*a^3 - 12*a^2*b*c + I*b^3*c^3*Pi^3 - 12*a^2*b*d*x + (2*I)*b^3*c^2*d*Pi^3*x + I*b^3*c*d^2*Pi^3*x^2 + 12*b^2*(-1 + c + d*x)*(b*(c + d*x) + a*(1 + c + d*x))*ArcTanh[c + d*x]^2 + 4*b^3*(-1 + c^2 + 2*c*d*x + d^2*x^2)*ArcTanh[c + d*x]^3 + 12*b*ArcTanh[c + d*x]*(a*(-2*b*(c + d*x) + a*(-1 + c^2 + 2*c*d*x + d^2*x^2)) + 2*b^2*(c + d*x)^2*Log[1 - E^(-2*ArcTanh[c + d*x])]) + 2*4*a*b^2*c^2*Log[(c + d*x)/Sqrt[1 - (c + d*x)^2]] + 48*a*b^2*c*d*x*Log[(c + d*x)/Sqrt[1 - (c + d*x)^2]] + 24*a*b^2*d^2*x^2*Log[(c + d*x)/Sqrt[1 - (c + d*x)^2]] - 12*b^3*(c + d*x)^2*PolyLog[2, E^(-2*ArcTanh[c + d*x])])/(8*d*e^3*(c + d*x)^2)
```

Rubi [A] (verified)

Time = 1.03 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.86, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.391$, Rules used = {6657, 27, 6452, 6544, 6452, 6510, 6550, 6494, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^3} dx$$

↓ 6657

$$\frac{\int \frac{(a+b\operatorname{arctanh}(c+dx))^3}{e^3(c+dx)^3} d(c+dx)}{d}$$

↓ 27

$$\frac{\int \frac{(a+b\operatorname{arctanh}(c+dx))^3}{(c+dx)^3} d(c+dx)}{de^3}$$

↓ 6452

$$\frac{\frac{3}{2}b \int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{(c+dx)^2(1-(c+dx)^2)} d(c+dx) - \frac{(a+b\operatorname{arctanh}(c+dx))^3}{2(c+dx)^2}}{de^3}$$

↓ 6544

$$\frac{\frac{3}{2}b \left(\int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{(c+dx)^2} d(c+dx) + \int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{1-(c+dx)^2} d(c+dx) \right) - \frac{(a+b\operatorname{arctanh}(c+dx))^3}{2(c+dx)^2}}{de^3}$$

↓ 6452

$$\frac{\frac{3}{2}b \left(2b \int \frac{a+b\operatorname{arctanh}(c+dx)}{(c+dx)(1-(c+dx)^2)} d(c+dx) + \int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{1-(c+dx)^2} d(c+dx) - \frac{(a+b\operatorname{arctanh}(c+dx))^2}{c+dx} \right) - \frac{(a+b\operatorname{arctanh}(c+dx))^3}{2(c+dx)^2}}{de^3}$$

↓ 6510

$$\frac{\frac{3}{2}b \left(2b \int \frac{a+b\operatorname{arctanh}(c+dx)}{(c+dx)(1-(c+dx)^2)} d(c+dx) + \frac{(a+b\operatorname{arctanh}(c+dx))^3}{3b} - \frac{(a+b\operatorname{arctanh}(c+dx))^2}{c+dx} \right) - \frac{(a+b\operatorname{arctanh}(c+dx))^3}{2(c+dx)^2}}{de^3}$$

↓ 6550

$$\frac{\frac{3}{2}b \left(2b \left(\int \frac{a+b\operatorname{arctanh}(c+dx)}{(c+dx)(c+dx+1)} d(c+dx) + \frac{(a+b\operatorname{arctanh}(c+dx))^2}{2b} \right) + \frac{(a+b\operatorname{arctanh}(c+dx))^3}{3b} - \frac{(a+b\operatorname{arctanh}(c+dx))^2}{c+dx} \right) - (a+b\operatorname{arctanh}(c+dx))^3}{de^3}$$

↓ 6494

$$\frac{\frac{3}{2}b \left(2b \left(-b \int \frac{\log\left(2 - \frac{2}{c+dx+1}\right)}{1-(c+dx)^2} d(c+dx) + \frac{(a+b\operatorname{arctanh}(c+dx))^2}{2b} + \log\left(2 - \frac{2}{c+dx+1}\right) (a+b\operatorname{arctanh}(c+dx)) \right) + \frac{(a+b\operatorname{arctanh}(c+dx))^3}{2(c+dx)^2} \right) - \frac{(a+b\operatorname{arctanh}(c+dx))^3}{2(c+dx)^2}}{de^3}$$

↓ 2897

$$\frac{\frac{3}{2}b \left(2b \left(\frac{(a+b\operatorname{arctanh}(c+dx))^2}{2b} + \log\left(2 - \frac{2}{c+dx+1}\right) (a+b\operatorname{arctanh}(c+dx)) - \frac{1}{2}b \operatorname{PolyLog}\left(2, \frac{2}{c+dx+1} - 1\right) \right) + \frac{(a+b\operatorname{arctanh}(c+dx))^3}{2(c+dx)^2} \right) - \frac{(a+b\operatorname{arctanh}(c+dx))^3}{2(c+dx)^2}}{de^3}$$

input `Int[(a + b*ArcTanh[c + d*x])^3/(c*e + d*e*x)^3,x]`

output `(-1/2*(a + b*ArcTanh[c + d*x])^3/(c + d*x)^2 + (3*b*(-(a + b*ArcTanh[c + d*x])^2/(c + d*x)) + (a + b*ArcTanh[c + d*x])^3/(3*b) + 2*b*((a + b*ArcTanh[c + d*x])^2/(2*b) + (a + b*ArcTanh[c + d*x])*Log[2 - 2/(1 + c + d*x)] - (b*PolyLog[2, -1 + 2/(1 + c + d*x)]/2)))/2)/(d*e^3)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2897 `Int[Log[u_]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 6452 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTanh[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 6494 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((x_)*((d_) + (e_)*(x_))), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Simp[b*c*(p/d) Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6510 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_)^(p_)/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]`

rule 6544

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*((f_.)*(x_))^(m_)/((d_) + (
e_.)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTanh[c*x])^p, x
], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTanh[c*x])^p/(d + e*x
^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

rule 6550

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_)^2)),
x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*d*(p + 1)), x] + Simp[1/
d Int[(a + b*ArcTanh[c*x])^p/(x*(1 + c*x)), x], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]
```

rule 6657

```
Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x
], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0]
&& IGtQ[p, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.48 (sec) , antiderivative size = 4949, normalized size of antiderivative = 29.81

| method | result | size |
|-------------------|---------------------------------|------|
| derivativedivides | Expression too large to display | 4949 |
| default | Expression too large to display | 4949 |
| parts | Expression too large to display | 4957 |

input

```
int((a+b*arctanh(d*x+c))^3/(d*e*x+c*e)^3,x,method=_RETURNVERBOSE)
```


output

```

1/d*(-1/2*a^3/e^3/(d*x+c)^2+b^3/e^3*(-3/2*arctanh(d*x+c)^2+3/2*polylog(2,-
(d*x+c+1)/(1-(d*x+c)^2)^(1/2))+3/2*polylog(2,(d*x+c+1)/(1-(d*x+c)^2)^(1/2)
)-3/2/(d*x+c)*arctanh(d*x+c)^2+1/2*arctanh(d*x+c)^3-3/4*arctanh(d*x+c)^2*ln
n(d*x+c-1)+3/4*arctanh(d*x+c)^2*ln(d*x+c+1)-3/2*arctanh(d*x+c)^2*ln((d*x+c
+1)/(1-(d*x+c)^2)^(1/2))+3/8*I*Pi*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))*csgn
(I*(d*x+c+1)^2/((d*x+c)^2-1))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1)/(1-(d*x+c
+1)^2/((d*x+c)^2-1)))*arctanh(d*x+c)^2+3/4*I*Pi*arctanh(d*x+c)^2-3/2*dilog
((d*x+c+1)/(1-(d*x+c)^2)^(1/2))+3/2*dilog(1+(d*x+c+1)/(1-(d*x+c)^2)^(1/2))
-3/8*I*Pi*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-
1)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))^2*dilog((d*x+c+1)/(1-(d*x+c)^2)^(1/2))-3
/8*I*Pi*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1)
/(1-(d*x+c+1)^2/((d*x+c)^2-1)))^2*polylog(2,(d*x+c+1)/(1-(d*x+c)^2)^(1/2))
-3/8*I*Pi*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-
1)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))^2*polylog(2,-(d*x+c+1)/(1-(d*x+c)^2)^(1/
2))+3/8*I*Pi*csgn(I*(d*x+c+1)/(1-(d*x+c)^2)^(1/2))^2*csgn(I*(d*x+c+1)^2/((
d*x+c)^2-1))*dilog((d*x+c+1)/(1-(d*x+c)^2)^(1/2))+3/8*I*Pi*csgn(I/(1-(d*x+
c+1)^2/((d*x+c)^2-1))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1)/(1-(d*x+c+1)^2/((d
*x+c)^2-1)))^2*polylog(2,(d*x+c+1)/(1-(d*x+c)^2)^(1/2))+3/8*I*Pi*csgn(I/(1-
(d*x+c+1)^2/((d*x+c)^2-1))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1)/(1-(d*x+c+1)
^2/((d*x+c)^2-1)))^2*polylog(2,-(d*x+c+1)/(1-(d*x+c)^2)^(1/2))+3/4*I*Pi...

```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^3} dx = \int \frac{(b \operatorname{arctanh}(dx + c) + a)^3}{(dex + ce)^3} dx$$

input

```
integrate((a+b*arctanh(d*x+c))^3/(d*e*x+c*e)^3,x, algorithm="fricas")
```

output

```

integral((b^3*arctanh(d*x + c)^3 + 3*a*b^2*arctanh(d*x + c)^2 + 3*a^2*b*ar
ctanh(d*x + c) + a^3)/(d^3*e^3*x^3 + 3*c*d^2*e^3*x^2 + 3*c^2*d*e^3*x + c^3
*e^3), x)

```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^3} dx$$

$$= \frac{\int \frac{a^3}{c^3 + 3c^2 dx + 3cd^2 x^2 + d^3 x^3} dx + \int \frac{b^3 \operatorname{atanh}^3(c + dx)}{c^3 + 3c^2 dx + 3cd^2 x^2 + d^3 x^3} dx + \int \frac{3ab^2 \operatorname{atanh}^2(c + dx)}{c^3 + 3c^2 dx + 3cd^2 x^2 + d^3 x^3} dx + \int \frac{3a^2 b \operatorname{atanh}(c + dx)}{c^3 + 3c^2 dx + 3cd^2 x^2 + d^3 x^3} dx}{e^3}$$

input `integrate((a+b*atanh(d*x+c))**3/(d*e*x+c*e)**3,x)`

output `(Integral(a**3/(c**3 + 3*c**2*d*x + 3*c*d**2*x**2 + d**3*x**3), x) + Integral(b**3*atanh(c + d*x)**3/(c**3 + 3*c**2*d*x + 3*c*d**2*x**2 + d**3*x**3), x) + Integral(3*a*b**2*atanh(c + d*x)**2/(c**3 + 3*c**2*d*x + 3*c*d**2*x**2 + d**3*x**3), x) + Integral(3*a**2*b*atanh(c + d*x)/(c**3 + 3*c**2*d*x + 3*c*d**2*x**2 + d**3*x**3), x))/e**3`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^3} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^3}{(dex + ce)^3} dx$$

input `integrate((a+b*arctanh(d*x+c))^3/(d*e*x+c*e)^3,x, algorithm="maxima")`

output

```
-3/4*(d*(2/(d^3*e^3*x + c*d^2*e^3) - log(d*x + c + 1)/(d^2*e^3) + log(d*x
+ c - 1)/(d^2*e^3)) + 2*arctanh(d*x + c)/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^
2*d*e^3))*a^2*b - 3/8*(d^2*((log(d*x + c + 1)^2 - 2*log(d*x + c + 1)*log(d
*x + c - 1) + log(d*x + c - 1)^2 + 4*log(d*x + c - 1))/(d^3*e^3) + 4*log(d
*x + c + 1)/(d^3*e^3) - 8*log(d*x + c)/(d^3*e^3)) + 4*d*(2/(d^3*e^3*x + c*
d^2*e^3) - log(d*x + c + 1)/(d^2*e^3) + log(d*x + c - 1)/(d^2*e^3))*arctan
h(d*x + c))*a*b^2 - 1/16*b^3*((d^2*x^2 + 2*c*d*x + c^2 - 1)*log(-d*x - c
+ 1)^3 + 3*(2*d*x - (d^2*x^2 + 2*c*d*x + c^2 - 1)*log(d*x + c + 1) + 2*c)*
log(-d*x - c + 1)^2)/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3) + 2*integra
te(-((d*x + c - 1)*log(d*x + c + 1)^3 + 3*(2*d^2*x^2 + 4*c*d*x - (d*x + c
- 1)*log(d*x + c + 1)^2 + 2*c^2 - (d^3*x^3 + 3*c*d^2*x^2 + c^3 + (3*c^2*d
- d)*x - c)*log(d*x + c + 1))*log(-d*x - c + 1))/(d^4*e^3*x^4 + c^4*e^3 -
c^3*e^3 + (4*c*d^3*e^3 - d^3*e^3)*x^3 + 3*(2*c^2*d^2*e^3 - c*d^2*e^3)*x^2
+ (4*c^3*d*e^3 - 3*c^2*d*e^3)*x), x) - 3/2*a*b^2*arctanh(d*x + c)^2/(d^3*
e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3) - 1/2*a^3/(d^3*e^3*x^2 + 2*c*d^2*e^3*
x + c^2*d*e^3)
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^3} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^3}{(dex + ce)^3} dx$$

input

```
integrate((a+b*arctanh(d*x+c))^3/(d*e*x+c*e)^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(d*x + c) + a)^3/(d*e*x + c*e)^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^3} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^3}{(ce + dex)^3} dx$$

input

```
int((a + b*atanh(c + d*x))^3/(c*e + d*e*x)^3,x)
```

output `int((a + b*atanh(c + d*x))^3/(c*e + d*e*x)^3, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^3} dx = \text{Too large to display}$$

input `int((a+b*atanh(d*x+c))^3/(d*e*x+c*e)^3,x)`

output

```
(2*atanh(c + d*x)**3*b**3*c**3 + 4*atanh(c + d*x)**3*b**3*c**2*d*x + 2*atanh(c + d*x)**3*b**3*c*d**2*x**2 - 2*atanh(c + d*x)**3*b**3*c + 6*atanh(c + d*x)**2*a*b**2*c**3 + 12*atanh(c + d*x)**2*a*b**2*c**2*d*x + 6*atanh(c + d*x)**2*a*b**2*c*d**2*x**2 - 6*atanh(c + d*x)**2*a*b**2*c + 6*atanh(c + d*x)**2*b**3*c*d*x + 6*atanh(c + d*x)**2*b**3*d**2*x**2 - 6*atanh(c + d*x)*a**2*b*c - 6*atanh(c + d*x)*a*b**2*c**2 + 6*atanh(c + d*x)*a*b**2*d**2*x**2 - 6*atanh(c + d*x)*b**3*c - 12*atanh(c + d*x)*b**3*d*x + 12*int((atanh(c + d*x)*x)/(c**5 + 5*c**4*d*x + 10*c**3*d**2*x**2 - c**3 + 10*c**2*d**3*x**3 - 3*c**2*d*x + 5*c*d**4*x**4 - 3*c*d**2*x**2 + d**5*x**5 - d**3*x**3),x)*b**3*c**2*d**2 + 24*int((atanh(c + d*x)*x)/(c**5 + 5*c**4*d*x + 10*c**3*d**2*x**2 - c**3 + 10*c**2*d**3*x**3 - 3*c**2*d*x + 5*c*d**4*x**4 - 3*c*d**2*x**2 + d**5*x**5 - d**3*x**3),x)*b**3*c*d**3*x + 12*int((atanh(c + d*x)*x)/(c**5 + 5*c**4*d*x + 10*c**3*d**2*x**2 - c**3 + 10*c**2*d**3*x**3 - 3*c**2*d*x + 5*c*d**4*x**4 - 3*c*d**2*x**2 + d**5*x**5 - d**3*x**3),x)*b**3*d**4*x**2 - 3*log(c + d*x - 1)*a**2*b*c**3 - 6*log(c + d*x - 1)*a**2*b*c**2*d*x - 3*log(c + d*x - 1)*a**2*b*c*d**2*x**2 - 6*log(c + d*x - 1)*a*b**2*c**3 - 12*log(c + d*x - 1)*a*b**2*c**2*d*x + 3*log(c + d*x - 1)*a*b**2*c**2 - 6*log(c + d*x - 1)*a*b**2*c*d**2*x**2 + 6*log(c + d*x - 1)*a*b**2*c*d*x + 3*log(c + d*x - 1)*a*b**2*d**2*x**2 + 3*log(c + d*x - 1)*b**3*c**3 + 6*log(c + d*x - 1)*b**3*c**2*d*x - 6*log(c + d*x - 1)*b**3*c**2 + 3*log(c...
```

3.28 $\int \frac{(a+b\operatorname{arctanh}(c+dx))^3}{(ce+dex)^4} dx$

| | |
|---------------------------------------|-----|
| Optimal result | 276 |
| Mathematica [C] (verified) | 277 |
| Rubi [A] (warning: unable to verify) | 277 |
| Maple [C] (warning: unable to verify) | 282 |
| Fricas [F] | 283 |
| Sympy [F] | 283 |
| Maxima [F] | 284 |
| Giac [F] | 284 |
| Mupad [F(-1)] | 285 |
| Reduce [F] | 285 |

Optimal result

Integrand size = 23, antiderivative size = 269

$$\int \frac{(a + b\operatorname{arctanh}(c + dx))^3}{(ce + dex)^4} dx = -\frac{b^2(a + b\operatorname{arctanh}(c + dx))}{de^4(c + dx)} + \frac{b(a + b\operatorname{arctanh}(c + dx))^2}{2de^4} - \frac{b(a + b\operatorname{arctanh}(c + dx))^2}{2de^4(c + dx)^2} + \frac{(a + b\operatorname{arctanh}(c + dx))^3}{3de^4} - \frac{(a + b\operatorname{arctanh}(c + dx))^3}{3de^4(c + dx)^3} + \frac{b^3 \log(c + dx)}{de^4} - \frac{b^3 \log(1 - (c + dx)^2)}{2de^4} + \frac{b(a + b\operatorname{arctanh}(c + dx))^2 \log\left(2 - \frac{2}{1+c+dx}\right)}{de^4} - \frac{b^2(a + b\operatorname{arctanh}(c + dx)) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1+c+dx}\right)}{de^4} - \frac{b^3 \operatorname{PolyLog}\left(3, -1 + \frac{2}{1+c+dx}\right)}{2de^4}$$

output

```
-b^2*(a+b*arctanh(d*x+c))/d/e^4/(d*x+c)+1/2*b*(a+b*arctanh(d*x+c))^2/d/e^4
-1/2*b*(a+b*arctanh(d*x+c))^2/d/e^4/(d*x+c)^2+1/3*(a+b*arctanh(d*x+c))^3/d
/e^4-1/3*(a+b*arctanh(d*x+c))^3/d/e^4/(d*x+c)^3+b^3*ln(d*x+c)/d/e^4-1/2*b^
3*ln(1-(d*x+c)^2)/d/e^4+b*(a+b*arctanh(d*x+c))^2*ln(2-2/(d*x+c+1))/d/e^4-b
^2*(a+b*arctanh(d*x+c))*polylog(2,-1+2/(d*x+c+1))/d/e^4-1/2*b^3*polylog(3,
-1+2/(d*x+c+1))/d/e^4
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.95 (sec) , antiderivative size = 393, normalized size of antiderivative = 1.46

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^4} dx$$

$$= -\frac{2a^3}{(c+dx)^3} - \frac{3a^2b}{(c+dx)^2} - \frac{6a^2b \operatorname{arctanh}(c+dx)}{(c+dx)^3} + 6a^2b \log(c + dx) - 3a^2b \log(1 - c^2 - 2cdx - d^2x^2) + 6ab^2 \left(-\frac{c}{c+dx} \right)$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])^3/(c*e + d*e*x)^4,x]
```

output

```
((-2*a^3)/(c + d*x)^3 - (3*a^2*b)/(c + d*x)^2 - (6*a^2*b*ArcTanh[c + d*x])
/(c + d*x)^3 + 6*a^2*b*Log[c + d*x] - 3*a^2*b*Log[1 - c^2 - 2*c*d*x - d^2*
x^2] + 6*a*b^2*(-((c + d*x)^2 + ArcTanh[c + d*x]^2)/(c + d*x)^3) + ArcTan
h[c + d*x]*(-((1 - (c + d*x)^2)/(c + d*x)^2) + ArcTanh[c + d*x] + 2*Log[1
- E^(-2*ArcTanh[c + d*x])]) - PolyLog[2, E^(-2*ArcTanh[c + d*x])]) + 6*b^3
*((I/24)*Pi^3 - ArcTanh[c + d*x]/(c + d*x) - ((1 - (c + d*x)^2)*ArcTanh[c
+ d*x]^2)/(2*(c + d*x)^2) - ArcTanh[c + d*x]^3/3 - ArcTanh[c + d*x]^3/(3*(
c + d*x)) - ((1 - (c + d*x)^2)*ArcTanh[c + d*x]^3)/(3*(c + d*x)^3) + ArcTa
nh[c + d*x]^2*Log[1 - E^(2*ArcTanh[c + d*x])] + Log[(c + d*x)/Sqrt[1 - (c
+ d*x)^2]] + ArcTanh[c + d*x]*PolyLog[2, E^(2*ArcTanh[c + d*x])] - PolyLog
[3, E^(2*ArcTanh[c + d*x])]/2))/(6*d*e^4)
```

Rubi [A] (warning: unable to verify)

Time = 1.89 (sec) , antiderivative size = 220, normalized size of antiderivative = 0.82, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.696$, Rules used = {6657, 27, 6452, 6544, 6452, 6544, 6452, 243, 47, 14, 16, 6510, 6550, 6494, 6618, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^4} dx$$

$$\begin{aligned}
& \downarrow 6657 \\
& \frac{\int \frac{(a+\operatorname{barctanh}(c+dx))^3}{e^4(c+dx)^4} d(c+dx)}{d} \\
& \downarrow 27 \\
& \frac{\int \frac{(a+\operatorname{barctanh}(c+dx))^3}{(c+dx)^4} d(c+dx)}{de^4} \\
& \downarrow 6452 \\
& \frac{b \int \frac{(a+\operatorname{barctanh}(c+dx))^2}{(c+dx)^3(1-(c+dx)^2)} d(c+dx) - \frac{(a+\operatorname{barctanh}(c+dx))^3}{3(c+dx)^3}}{de^4} \\
& \downarrow 6544 \\
& \frac{b \left(\int \frac{(a+\operatorname{barctanh}(c+dx))^2}{(c+dx)^3} d(c+dx) + \int \frac{(a+\operatorname{barctanh}(c+dx))^2}{(c+dx)(1-(c+dx)^2)} d(c+dx) \right) - \frac{(a+\operatorname{barctanh}(c+dx))^3}{3(c+dx)^3}}{de^4} \\
& \downarrow 6452 \\
& \frac{b \left(b \int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)^2(1-(c+dx)^2)} d(c+dx) + \int \frac{(a+\operatorname{barctanh}(c+dx))^2}{(c+dx)(1-(c+dx)^2)} d(c+dx) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{2(c+dx)^2} \right) - \frac{(a+\operatorname{barctanh}(c+dx))^3}{3(c+dx)^3}}{de^4} \\
& \downarrow 6544 \\
& \frac{b \left(b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{(c+dx)^2} d(c+dx) + \int \frac{a+\operatorname{barctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) \right) + \int \frac{(a+\operatorname{barctanh}(c+dx))^2}{(c+dx)(1-(c+dx)^2)} d(c+dx) - \frac{(a+\operatorname{barctanh}(c+dx))^2}{2(c+dx)^2} \right)}{de^4} \\
& \downarrow 6452 \\
& \frac{b \left(b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) + b \int \frac{1}{(c+dx)(1-(c+dx)^2)} d(c+dx) - \frac{a+\operatorname{barctanh}(c+dx)}{c+dx} \right) + \int \frac{(a+\operatorname{barctanh}(c+dx))^2}{(c+dx)(1-(c+dx)^2)} d(c+dx) \right)}{de^4} \\
& \downarrow 243 \\
& \frac{b \left(b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) + \frac{1}{2} b \int \frac{1}{(-c-dx+1)(c+dx)^2} d(c+dx)^2 - \frac{a+\operatorname{barctanh}(c+dx)}{c+dx} \right) + \int \frac{(a+\operatorname{barctanh}(c+dx))^2}{(c+dx)(1-(c+dx)^2)} d(c+dx) \right)}{de^4} \\
& \downarrow 47 \\
& \frac{b \left(b \left(\int \frac{a+\operatorname{barctanh}(c+dx)}{1-(c+dx)^2} d(c+dx) + \frac{1}{2} b \left(\int \frac{1}{-c-dx+1} d(c+dx)^2 + \int \frac{1}{(c+dx)^2} d(c+dx)^2 \right) - \frac{a+\operatorname{barctanh}(c+dx)}{c+dx} \right) + \int \frac{(a+\operatorname{barctanh}(c+dx))^2}{(c+dx)(1-(c+dx)^2)} d(c+dx) \right)}{de^4}
\end{aligned}$$

↓ 14

$$\frac{b \left(\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(c + dx)(1 - (c + dx)^2)} d(c + dx) + b \left(\int \frac{a + b \operatorname{arctanh}(c + dx)}{1 - (c + dx)^2} d(c + dx) + \frac{1}{2} b \left(\int \frac{1}{-c - dx + 1} d(c + dx)^2 + \log((c + dx)^2) \right) \right) \right)}{de^4}$$

↓ 16

$$\frac{b \left(\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(c + dx)(1 - (c + dx)^2)} d(c + dx) + b \left(\int \frac{a + b \operatorname{arctanh}(c + dx)}{1 - (c + dx)^2} d(c + dx) - \frac{a + b \operatorname{arctanh}(c + dx)}{c + dx} + \frac{1}{2} b (\log((c + dx)^2) - 1) \right) \right)}{de^4}$$

↓ 6510

$$\frac{b \left(\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(c + dx)(1 - (c + dx)^2)} d(c + dx) - \frac{(a + b \operatorname{arctanh}(c + dx))^2}{2(c + dx)^2} + b \left(\frac{(a + b \operatorname{arctanh}(c + dx))^2}{2b} - \frac{a + b \operatorname{arctanh}(c + dx)}{c + dx} + \frac{1}{2} b (\log((c + dx)^2) - 1) \right) \right)}{de^4}$$

↓ 6550

$$\frac{b \left(\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(c + dx)(c + dx + 1)} d(c + dx) + \frac{(a + b \operatorname{arctanh}(c + dx))^3}{3b} - \frac{(a + b \operatorname{arctanh}(c + dx))^2}{2(c + dx)^2} + b \left(\frac{(a + b \operatorname{arctanh}(c + dx))^2}{2b} - \frac{a + b \operatorname{arctanh}(c + dx)}{c + dx} \right) \right)}{de^4}$$

↓ 6494

$$\frac{b \left(-2b \int \frac{(a + b \operatorname{arctanh}(c + dx)) \log\left(2 - \frac{2}{c + dx + 1}\right)}{1 - (c + dx)^2} d(c + dx) + \frac{(a + b \operatorname{arctanh}(c + dx))^3}{3b} - \frac{(a + b \operatorname{arctanh}(c + dx))^2}{2(c + dx)^2} + \log\left(2 - \frac{2}{c + dx + 1}\right) \right)}{de^4}$$

↓ 6618

$$\frac{b \left(-2b \left(\frac{1}{2} \operatorname{PolyLog}\left(2, \frac{2}{c + dx + 1} - 1\right) (a + b \operatorname{arctanh}(c + dx)) - \frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, \frac{2}{c + dx + 1} - 1\right)}{1 - (c + dx)^2} d(c + dx) \right) + \frac{(a + b \operatorname{arctanh}(c + dx))^3}{3b} \right)}{de^4}$$

↓ 7164

$$\frac{b \left(-2b \left(\frac{1}{2} \operatorname{PolyLog}\left(2, \frac{2}{c + dx + 1} - 1\right) (a + b \operatorname{arctanh}(c + dx)) + \frac{1}{4} b \operatorname{PolyLog}\left(3, \frac{2}{c + dx + 1} - 1\right) \right) + \frac{(a + b \operatorname{arctanh}(c + dx))^3}{3b} \right)}{de^4}$$

input

```
Int[(a + b*ArcTanh[c + d*x])^3/(c*e + d*e*x)^4,x]
```


output

$$\begin{aligned} & (-1/3*(a + b*\text{ArcTanh}[c + d*x])^3/(c + d*x)^3 + b*(-1/2*(a + b*\text{ArcTanh}[c + \\ & d*x])^2/(c + d*x)^2 + (a + b*\text{ArcTanh}[c + d*x])^3/(3*b) + b*(-((a + b*\text{ArcTa} \\ & \text{nh}[c + d*x])/(c + d*x)) + (a + b*\text{ArcTanh}[c + d*x])^2/(2*b) + (b*(-\text{Log}[1 - \\ & c - d*x] + \text{Log}[(c + d*x)^2]))/2) + (a + b*\text{ArcTanh}[c + d*x])^2*\text{Log}[2 - 2/(1 \\ & + c + d*x)] - 2*b*((a + b*\text{ArcTanh}[c + d*x])* \text{PolyLog}[2, -1 + 2/(1 + c + d \\ & *x)])/2 + (b*\text{PolyLog}[3, -1 + 2/(1 + c + d*x)])/4)))/(d*e^4) \end{aligned}$$

Defintions of rubi rules used

rule 14

$$\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ /; FreeQ}[a, x]$$

rule 16

$$\text{Int}[(c_)/((a_.) + (b_.)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ /; FreeQ}[\{a, b, c\}, x]$$

rule 27

$$\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] \text{ /; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] \text{ /; FreeQ}[b, x]$$

rule 47

$$\text{Int}[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ /; FreeQ}[\{a, b, c, d\}, x]$$

rule 243

$$\text{Int}[(x_)^{(m_)}*((a_.) + (b_.)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{((m-1)/2)*(a + b*x)^p}, x], x, x^2], x] \text{ /; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$$

rule 6452

$$\begin{aligned} & \text{Int}[(a_.) + \text{ArcTanh}[(c_.)*(x_)^{(n_)}]*(b_.)^{(p_)}*(x_)^{(m_)}, x_Symbol] : \\ & > \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTanh}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m \\ & + 1)) \text{ Int}[x^{(m+n)}*((a + b*\text{ArcTanh}[c*x^n])^{(p-1)/(1-c^2*x^{2*n})}), x \\ &], x] \text{ /; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \\ &] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1] \end{aligned}$$

rule 6494 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((x) \cdot (d + (e) \cdot (x))), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{Log}[2 - 2/(1 + e \cdot (x/d))]/d), x] - \text{Simp}[b \cdot c \cdot (p/d) \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{Log}[2 - 2/(1 + e \cdot (x/d))]/(1 - c^2 \cdot x^2)), x], x] /;$ FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]

rule 6510 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((d) + (e) \cdot (x)^2), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot c \cdot d \cdot (p+1)), x] /;$ FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && NeQ[p, -1]

rule 6544 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p \cdot (f \cdot (x))^m / ((d) + (e) \cdot (x)^2), x_Symbol] \rightarrow \text{Simp}[1/d \text{Int}[(f \cdot x)^m \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p, x], x] - \text{Simp}[e/(d \cdot f^2) \text{Int}[(f \cdot x)^{m+2} \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2)), x], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]

rule 6550 $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / ((x) \cdot (d + (e) \cdot (x)^2)), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p+1} / (b \cdot d \cdot (p+1)), x] + \text{Simp}[1/d \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p / (x \cdot (1 + c \cdot x)), x], x] /;$ FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[p, 0]

rule 6618 $\text{Int}[(\text{Log}[u] \cdot (a + \text{ArcTanh}[c \cdot x] \cdot b)^p) / ((d) + (e) \cdot (x)^2), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{PolyLog}[2, 1 - u]/(2 \cdot c \cdot d)), x] - \text{Simp}[b \cdot (p/2) \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{PolyLog}[2, 1 - u]/(d + e \cdot x^2)), x], x] /;$ FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 + c \cdot x))^2, 0]

rule 6657 $\text{Int}[(a + \text{ArcTanh}[c] + (d) \cdot (x)) \cdot b)^p \cdot (e + (f) \cdot (x))^m, x_Symbol] \rightarrow \text{Simp}[1/d \text{Subst}[\text{Int}[(f \cdot (x/d))^m \cdot (a + b \cdot \text{ArcTanh}[x])^p, x], x, c + d \cdot x], x] /;$ FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d \cdot e - c \cdot f, 0] && IGtQ[p, 0]

rule 7164 $\text{Int}[(u) \cdot \text{PolyLog}[n, v], x_Symbol] \rightarrow \text{With}[\{w = \text{DerivativeDivides}[v, u \cdot v, x]\}, \text{Simp}[w \cdot \text{PolyLog}[n + 1, v], x] /;$!FalseQ[w]] /;

 FreeQ[n, x]

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 7.74 (sec) , antiderivative size = 1768, normalized size of antiderivative = 6.57

| method | result | size |
|--------------------|---------------------------------|------|
| derivativeldivides | Expression too large to display | 1768 |
| default | Expression too large to display | 1768 |
| parts | Expression too large to display | 1776 |

input `int((a+b*arctanh(d*x+c))^3/(d*e*x+c*e)^4,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & 1/d*(-1/3*a^3/e^4/(d*x+c)^3+b^3/e^4*(1/2*arctanh(d*x+c)^2*polylog(3,(d*x+c+1)/(1-(d*x+c)^2)^{(1/2)}+\ln(1+(d*x+c+1)/(1-(d*x+c)^2)^{(1/2)})-2*polylog(3, \\
 & -(d*x+c+1)/(1-(d*x+c)^2)^{(1/2)})+2*arctanh(d*x+c)*polylog(2,(d*x+c+1)/(1-(d*x+c)^2)^{(1/2)} \\
 & +\ln(d*x+c)*arctanh(d*x+c)^2-arctanh(d*x+c)^2*\ln((d*x+c+1)^2/(1-(d*x+c)^2)-1) \\
 & +arctanh(d*x+c)^2*\ln(1+(d*x+c+1)/(1-(d*x+c)^2)^{(1/2)})+2*arctanh(d*x+c)*polylog(2, \\
 & -(d*x+c+1)/(1-(d*x+c)^2)^{(1/2)})+arctanh(d*x+c)^2*\ln(1-(d*x+c+1)/(1-(d*x+c)^2)^{(1/2)}) \\
 & -1/2/(d*x+c)^2*arctanh(d*x+c)^2+\ln(2)*arctanh(d*x+c)^2-1/3*arctanh(d*x+c)^3-1/2*arctanh(d*x+c)^2 \\
 & *\ln(d*x+c-1)-1/2*arctanh(d*x+c)^2*\ln(d*x+c+1)+arctanh(d*x+c)^2*\ln((d*x+c+1)/(1-(d*x+c)^2)^{(1/2)}) \\
 & -1/4*I*Pi*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1)/(1-(d*x+c+1)^2/((d*x+c)^2-1))) \\
 & *arctanh(d*x+c)^2+\ln((d*x+c+1)/(1-(d*x+c)^2)^{(1/2)})-1/3/(d*x+c)^3*arctanh(d*x+c)^3-1/2*(d*x+c+1-(1-(d*x+c)^2)^{(1/2)})/(d*x+c)*arctanh(d*x+c)-1/2*arctanh(d*x+c)*(d*x+(1-(d*x+c)^2)^{(1/2)}+c+1)/(d*x+c)+1/2*I*Pi*arctanh(d*x+c)^2+1/2*I*Pi*csgn(I*(-(d*x+c+1)^2/((d*x+c)^2-1)-1))*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))*csgn(I*(-(d*x+c+1)^2/((d*x+c)^2-1)-1)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))*arctanh(d*x+c)^2+1/4*I*Pi*csgn(I*(d*x+c+1)/(1-(d*x+c)^2)^{(1/2)})^2*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1))*arctanh(d*x+c)^2+1/4*I*Pi*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))^2*arctanh(d*x+c)^2-1/4*I*Pi*csgn(I*(d*x+c+1)^2/((d*x+c)^2-1))
 \end{aligned}$$

Fricas [F]

$$\int \frac{(a + \operatorname{arctanh}(c + dx))^3}{(ce + dex)^4} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^3}{(dex + ce)^4} dx$$

input `integrate((a+b*arctanh(d*x+c))^3/(d*e*x+c*e)^4,x, algorithm="fricas")`

output `integral((b^3*arctanh(d*x + c)^3 + 3*a*b^2*arctanh(d*x + c)^2 + 3*a^2*b*arctanh(d*x + c) + a^3)/(d^4*e^4*x^4 + 4*c*d^3*e^4*x^3 + 6*c^2*d^2*e^4*x^2 + 4*c^3*d*e^4*x + c^4*e^4), x)`

Sympy [F]

$$\int \frac{(a + \operatorname{arctanh}(c + dx))^3}{(ce + dex)^4} dx = \int \frac{a^3}{c^4 + 4c^3dx + 6c^2d^2x^2 + 4cd^3x^3 + d^4x^4} dx + \int \frac{b^3 \operatorname{atanh}^3(c+dx)}{c^4 + 4c^3dx + 6c^2d^2x^2 + 4cd^3x^3 + d^4x^4} dx + \int \frac{3ab^2 \operatorname{atanh}^2(c+dx)}{c^4 + 4c^3dx + 6c^2d^2x^2 + 4cd^3x^3 + d^4x^4} dx + \int \frac{3ab \operatorname{atanh}(c+dx)}{c^4 + 4c^3dx + 6c^2d^2x^2 + 4cd^3x^3 + d^4x^4} dx + \int \frac{a^2}{c^4 + 4c^3dx + 6c^2d^2x^2 + 4cd^3x^3 + d^4x^4} dx$$

input `integrate((a+b*atanh(d*x+c))**3/(d*e*x+c*e)**4,x)`

output `(Integral(a**3/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + 4*c*d**3*x**3 + d**4*x**4), x) + Integral(b**3*atanh(c + d*x)**3/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + 4*c*d**3*x**3 + d**4*x**4), x) + Integral(3*a*b**2*atanh(c + d*x)**2/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + 4*c*d**3*x**3 + d**4*x**4), x) + Integral(3*a**2*b*atanh(c + d*x)/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + 4*c*d**3*x**3 + d**4*x**4), x))/e**4`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^4} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^3}{(dex + ce)^4} dx$$

input `integrate((a+b*arctanh(d*x+c))^3/(d*e*x+c*e)^4,x, algorithm="maxima")`

output

```
-1/2*(d*(1/(d^4*e^4*x^2 + 2*c*d^3*e^4*x + c^2*d^2*e^4) + log(d*x + c + 1)/
(d^2*e^4) - 2*log(d*x + c)/(d^2*e^4) + log(d*x + c - 1)/(d^2*e^4)) + 2*arc
tanh(d*x + c)/(d^4*e^4*x^3 + 3*c*d^3*e^4*x^2 + 3*c^2*d^2*e^4*x + c^3*d*e^4
))*a^2*b - 1/3*a^3/(d^4*e^4*x^3 + 3*c*d^3*e^4*x^2 + 3*c^2*d^2*e^4*x + c^3*
d*e^4) - 1/24*((b^3*d^3*x^3 + 3*b^3*c*d^2*x^2 + 3*b^3*c^2*d*x + (c^3 - 1)*
b^3)*log(-d*x - c + 1)^3 + 3*(b^3*d*x + b^3*c + 2*a*b^2 + (b^3*d^3*x^3 + 3
*b^3*c*d^2*x^2 + 3*b^3*c^2*d*x + (c^3 + 1)*b^3)*log(d*x + c + 1))*log(-d*x
- c + 1)^2)/(d^4*e^4*x^3 + 3*c*d^3*e^4*x^2 + 3*c^2*d^2*e^4*x + c^3*d*e^4)
- integrate(-1/8*((b^3*d*x + b^3*(c - 1))*log(d*x + c + 1)^3 + 6*(a*b^2*d
*x + a*b^2*(c - 1))*log(d*x + c + 1)^2 + (2*b^3*d^2*x^2 + 2*b^3*c^2 + 4*a*
b^2*c - 3*(b^3*d*x + b^3*(c - 1))*log(d*x + c + 1)^2 + 4*(b^3*c*d + a*b^2*
d)*x + 2*(b^3*d^4*x^4 + 4*b^3*c*d^3*x^3 + 6*b^3*c^2*d^2*x^2 + (c^4 + c)*b^
3 - 6*a*b^2*(c - 1) + ((4*c^3*d + d)*b^3 - 6*a*b^2*d)*x)*log(d*x + c + 1))
*log(-d*x - c + 1))/(d^5*e^4*x^5 + c^5*e^4 - c^4*e^4 + (5*c*d^4*e^4 - d^4*
e^4)*x^4 + 2*(5*c^2*d^3*e^4 - 2*c*d^3*e^4)*x^3 + 2*(5*c^3*d^2*e^4 - 3*c^2*
d^2*e^4)*x^2 + (5*c^4*d*e^4 - 4*c^3*d*e^4)*x), x)
```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^4} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^3}{(dex + ce)^4} dx$$

input `integrate((a+b*arctanh(d*x+c))^3/(d*e*x+c*e)^4,x, algorithm="giac")`

output

```
integrate((b*arctanh(d*x + c) + a)^3/(d*e*x + c*e)^4, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^4} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^3}{(ce + dex)^4} dx$$

input `int((a + b*atanh(c + d*x))^3/(c*e + d*e*x)^4,x)`output `int((a + b*atanh(c + d*x))^3/(c*e + d*e*x)^4, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(ce + dex)^4} dx = \text{too large to display}$$

input `int((a+b*atanh(d*x+c))^3/(d*e*x+c*e)^4,x)`

output

```
(6*atanh(c + d*x)**3*b**3*c**6 + 18*atanh(c + d*x)**3*b**3*c**5*d*x + 18*atanh(c + d*x)**3*b**3*c**4*d**2*x**2 + 6*atanh(c + d*x)**3*b**3*c**4 + 6*atanh(c + d*x)**3*b**3*c**3*d**3*x**3 + 18*atanh(c + d*x)**3*b**3*c**3*d*x + 18*atanh(c + d*x)**3*b**3*c**2*d**2*x**2 - 12*atanh(c + d*x)**3*b**3*c**2 + 6*atanh(c + d*x)**3*b**3*c*d**3*x**3 + 18*atanh(c + d*x)**2*a*b**2*c**6 + 54*atanh(c + d*x)**2*a*b**2*c**5*d*x + 54*atanh(c + d*x)**2*a*b**2*c**4*d**2*x**2 + 18*atanh(c + d*x)**2*a*b**2*c**4 + 18*atanh(c + d*x)**2*a*b**2*c**3*d**3*x**3 + 54*atanh(c + d*x)**2*a*b**2*c**3*d*x + 54*atanh(c + d*x)**2*a*b**2*c**2*d**2*x**2 - 36*atanh(c + d*x)**2*a*b**2*c**2 + 18*atanh(c + d*x)**2*a*b**2*c*d**3*x**3 + 24*atanh(c + d*x)**2*b**3*c**5 + 90*atanh(c + d*x)**2*b**3*c**4*d*x + 108*atanh(c + d*x)**2*b**3*c**3*d**2*x**2 - 24*atanh(c + d*x)**2*b**3*c**3 + 42*atanh(c + d*x)**2*b**3*c**2*d**3*x**3 - 36*atanh(c + d*x)**2*b**3*c**2*d*x - 18*atanh(c + d*x)**2*b**3*c*d**2*x**2 - 36*atanh(c + d*x)*a**2*b*c**2 - 12*atanh(c + d*x)*a*b**2*c**5 + 36*atanh(c + d*x)*a*b**2*c**3*d**2*x**2 - 48*atanh(c + d*x)*a*b**2*c**3 + 24*atanh(c + d*x)*a*b**2*c**2*d**3*x**3 - 72*atanh(c + d*x)*a*b**2*c**2*d*x - 36*atanh(c + d*x)*a*b**2*c*d**2*x**2 + 16*atanh(c + d*x)*b**3*c**6 + 48*atanh(c + d*x)*b**3*c**5*d*x + 48*atanh(c + d*x)*b**3*c**4*d**2*x**2 - 54*atanh(c + d*x)*b**3*c**4 + 16*atanh(c + d*x)*b**3*c**3*d**3*x**3 - 114*atanh(c + d*x)*b**3*c**3*d*x - 48*atanh(c + d*x)*b**3*c**2*d**2*x**2 + 6*atanh...
```

3.29 $\int \frac{\operatorname{arctanh}(1+x)}{2+2x} dx$

| | |
|---|-----|
| Optimal result | 287 |
| Mathematica [A] (verified) | 287 |
| Rubi [A] (verified) | 288 |
| Maple [A] (verified) | 289 |
| Fricas [F] | 289 |
| Sympy [F] | 290 |
| Maxima [B] (verification not implemented) | 290 |
| Giac [F] | 291 |
| Mupad [F(-1)] | 291 |
| Reduce [F] | 291 |

Optimal result

Integrand size = 12, antiderivative size = 21

$$\int \frac{\operatorname{arctanh}(1+x)}{2+2x} dx = -\frac{1}{4} \operatorname{PolyLog}(2, -1-x) + \frac{\operatorname{PolyLog}(2, 1+x)}{4}$$

output `-1/4*polylog(2,-1-x)+1/4*polylog(2,1+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.48

$$\int \frac{\operatorname{arctanh}(1+x)}{2+2x} dx = -\frac{1}{4} \operatorname{PolyLog}\left(2, \frac{1}{2}(-2-2x)\right) + \frac{1}{4} \operatorname{PolyLog}\left(2, \frac{1}{2}(2+2x)\right)$$

input `Integrate[ArcTanh[1 + x]/(2 + 2*x), x]`

output `-1/4*PolyLog[2, (-2 - 2*x)/2] + PolyLog[2, (2 + 2*x)/2]/4`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.19, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6657, 27, 6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(x+1)}{2x+2} dx \\ & \quad \downarrow 6657 \\ & \int \frac{\operatorname{arctanh}(x+1)}{2(x+1)} d(x+1) \\ & \quad \downarrow 27 \\ & \frac{1}{2} \int \frac{\operatorname{arctanh}(x+1)}{x+1} d(x+1) \\ & \quad \downarrow 6446 \\ & \frac{1}{2} \left(\frac{\operatorname{PolyLog}(2, x+1)}{2} - \frac{\operatorname{PolyLog}(2, -x-1)}{2} \right) \end{aligned}$$

input `Int[ArcTanh[1 + x]/(2 + 2*x),x]`

output `(-1/2*PolyLog[2, -1 - x] + PolyLog[2, 1 + x]/2)/2`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 6446 `Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))/(x_), x_Symbol] := Simp[a*Log[x], x] + (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) /; FreeQ[{a, b, c}, x]`

rule 6657

```
Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] :> Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x
], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0]
&& IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.93 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

| method | result | size |
|-------------------|--|------|
| risch | $\frac{\operatorname{dilog}(-x)}{4} - \frac{\operatorname{dilog}(x+2)}{4}$ | 14 |
| derivativedivides | $\frac{\ln(1+x) \operatorname{arctanh}(1+x)}{2} - \frac{\operatorname{dilog}(x+2)}{4} - \frac{\ln(1+x) \ln(x+2)}{4} - \frac{\operatorname{dilog}(1+x)}{4}$ | 34 |
| default | $\frac{\ln(1+x) \operatorname{arctanh}(1+x)}{2} - \frac{\operatorname{dilog}(x+2)}{4} - \frac{\ln(1+x) \ln(x+2)}{4} - \frac{\operatorname{dilog}(1+x)}{4}$ | 34 |
| parts | $\frac{\ln(1+x) \operatorname{arctanh}(1+x)}{2} - \frac{\operatorname{dilog}(x+2)}{4} - \frac{\ln(1+x) \ln(x+2)}{4} - \frac{\operatorname{dilog}(1+x)}{4}$ | 34 |

input

```
int(arctanh(1+x)/(2*x+2),x,method=_RETURNVERBOSE)
```

output

```
1/4*dilog(-x)-1/4*dilog(x+2)
```

Fricas [F]

$$\int \frac{\operatorname{arctanh}(1+x)}{2+2x} dx = \int \frac{\operatorname{artanh}(x+1)}{2(x+1)} dx$$

input

```
integrate(arctanh(1+x)/(2+2*x),x, algorithm="fricas")
```

output

```
integral(1/2*arctanh(x + 1)/(x + 1), x)
```

Sympy [F]

$$\int \frac{\operatorname{arctanh}(1+x)}{2+2x} dx = \int \frac{\operatorname{atanh}(x+1)}{x+1} dx$$

input `integrate(atanh(1+x)/(2+2*x),x)`

output `Integral(atanh(x + 1)/(x + 1), x)/2`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 58 vs. $2(15) = 30$.

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.76

$$\begin{aligned} \int \frac{\operatorname{arctanh}(1+x)}{2+2x} dx &= -\frac{1}{4} (\log(x+2) - \log(x)) \log(x+1) \\ &\quad + \frac{1}{2} \operatorname{artanh}(x+1) \log(x+1) - \frac{1}{4} \log(x+1) \log(x) \\ &\quad + \frac{1}{4} \log(x+2) \log(-x-1) - \frac{1}{4} \operatorname{Li}_2(-x) + \frac{1}{4} \operatorname{Li}_2(x+2) \end{aligned}$$

input `integrate(arctanh(1+x)/(2+2*x),x, algorithm="maxima")`

output `-1/4*(log(x + 2) - log(x))*log(x + 1) + 1/2*arctanh(x + 1)*log(x + 1) - 1/4*log(x + 1)*log(x) + 1/4*log(x + 2)*log(-x - 1) - 1/4*dilog(-x) + 1/4*dilog(x + 2)`

Giac [F]

$$\int \frac{\operatorname{arctanh}(1+x)}{2+2x} dx = \int \frac{\operatorname{artanh}(x+1)}{2(x+1)} dx$$

input `integrate(arctanh(1+x)/(2+2*x),x, algorithm="giac")`

output `integrate(1/2*arctanh(x + 1)/(x + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(1+x)}{2+2x} dx = \int \frac{\operatorname{atanh}(x+1)}{2x+2} dx$$

input `int(atanh(x + 1)/(2*x + 2),x)`

output `int(atanh(x + 1)/(2*x + 2), x)`

Reduce [F]

$$\int \frac{\operatorname{arctanh}(1+x)}{2+2x} dx = \frac{\left(\int \frac{\operatorname{atanh}(x+1)}{x+1} dx \right)}{2}$$

input `int(atanh(1+x)/(2+2*x),x)`

output `int(atanh(x + 1)/(x + 1),x)/2`

3.30 $\int \frac{\operatorname{arctanh}(a+bx)}{\frac{ad}{b}+dx} dx$

| | |
|---|-----|
| Optimal result | 292 |
| Mathematica [A] (verified) | 292 |
| Rubi [A] (verified) | 293 |
| Maple [A] (verified) | 294 |
| Fricas [F] | 294 |
| Sympy [F] | 295 |
| Maxima [B] (verification not implemented) | 295 |
| Giac [F] | 296 |
| Mupad [F(-1)] | 296 |
| Reduce [F] | 296 |

Optimal result

Integrand size = 19, antiderivative size = 32

$$\int \frac{\operatorname{arctanh}(a+bx)}{\frac{ad}{b}+dx} dx = -\frac{\operatorname{PolyLog}(2, -a-bx)}{2d} + \frac{\operatorname{PolyLog}(2, a+bx)}{2d}$$

output `-1/2*polylog(2,-b*x-a)/d+1/2*polylog(2,b*x+a)/d`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.62

$$\int \frac{\operatorname{arctanh}(a+bx)}{\frac{ad}{b}+dx} dx = b \left(-\frac{\operatorname{PolyLog}\left(2, -\frac{ad+bdx}{d}\right)}{2bd} + \frac{\operatorname{PolyLog}\left(2, \frac{ad+bdx}{d}\right)}{2bd} \right)$$

input `Integrate[ArcTanh[a + b*x]/((a*d)/b + d*x), x]`

output `b*(-1/2*PolyLog[2, -((a*d + b*d*x)/d)]/(b*d) + PolyLog[2, (a*d + b*d*x)/d]/(2*b*d))`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.94, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {6657, 27, 6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(a + bx)}{\frac{ad}{b} + dx} dx$$

↓ 6657

$$\int \frac{b \operatorname{arctanh}(a + bx)}{d(a + bx)} d(a + bx)$$

↓ 27

$$\int \frac{\operatorname{arctanh}(a + bx)}{a + bx} d(a + bx)$$

↓ 6446

$$\frac{\frac{1}{2} \operatorname{PolyLog}(2, a + bx) - \frac{1}{2} \operatorname{PolyLog}(2, -a - bx)}{d}$$

input `Int[ArcTanh[a + b*x]/((a*d)/b + d*x), x]`

output `(-1/2*PolyLog[2, -a - b*x] + PolyLog[2, a + b*x]/2)/d`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 6446 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.)/(x_), x_Symbol] := Simp[a*Log[x], x] + (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) /; FreeQ[{a, b, c}, x]`

rule 6657

```
Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] :> Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTanh[x])^p, x
], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0]
&& IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.63 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91

| method | result | size |
|-------------------|---|------|
| risch | $-\frac{\operatorname{dilog}(bx+a+1)}{2d} + \frac{\operatorname{dilog}(-bx-a+1)}{2d}$ | 29 |
| parts | $\frac{\ln(bx+a) \operatorname{arctanh}(bx+a)}{d} - \frac{\frac{\operatorname{dilog}(bx+a+1)}{2} + \frac{\ln(bx+a) \ln(bx+a+1)}{2} + \frac{\operatorname{dilog}(bx+a)}{2}}{d}$ | 56 |
| derivativedivides | $\frac{\frac{b \ln(bx+a) \operatorname{arctanh}(bx+a)}{d} - b \left(\frac{\operatorname{dilog}(bx+a+1)}{2} + \frac{\ln(bx+a) \ln(bx+a+1)}{2} + \frac{\operatorname{dilog}(bx+a)}{2} \right)}{b}$ | 62 |
| default | $\frac{\frac{b \ln(bx+a) \operatorname{arctanh}(bx+a)}{d} - b \left(\frac{\operatorname{dilog}(bx+a+1)}{2} + \frac{\ln(bx+a) \ln(bx+a+1)}{2} + \frac{\operatorname{dilog}(bx+a)}{2} \right)}{b}$ | 62 |

input

```
int(arctanh(b*x+a)/(a*d/b+d*x), x, method=_RETURNVERBOSE)
```

output

```
-1/2/d*dilog(b*x+a+1)+1/2/d*dilog(-b*x-a+1)
```

Fricas [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{artanh}(bx + a)}{dx + \frac{ad}{b}} dx$$

input

```
integrate(arctanh(b*x+a)/(a*d/b+d*x), x, algorithm="fricas")
```

output

```
integral(b*arctanh(b*x + a)/(b*d*x + a*d), x)
```

Sympy [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{\frac{ad}{b} + dx} dx = \frac{b \int \frac{\operatorname{atanh}\left(\frac{a+bx}{a+bx}\right) dx}{d}}$$

input `integrate(atanh(b*x+a)/(a*d/b+d*x), x)`

output `b*Integral(atanh(a + b*x)/(a + b*x), x)/d`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 132 vs. $2(26) = 52$.

Time = 0.03 (sec) , antiderivative size = 132, normalized size of antiderivative = 4.12

$$\int \frac{\operatorname{arctanh}(a + bx)}{\frac{ad}{b} + dx} dx =$$

$$-\frac{1}{2} b \left(\frac{\log(bx + a) \log(bx + a - 1) + \operatorname{Li}_2(-bx - a + 1)}{bd} - \frac{\log(bx + a + 1) \log(-bx - a) + \operatorname{Li}_2(bx + a - 1)}{bd} \right)$$

$$- \frac{b \left(\frac{\log(bx+a+1)}{b} - \frac{\log(bx+a-1)}{b} \right) \log\left(dx + \frac{ad}{b}\right)}{2d} + \frac{\operatorname{arctanh}(bx + a) \log\left(dx + \frac{ad}{b}\right)}{d}$$

input `integrate(arctanh(b*x+a)/(a*d/b+d*x), x, algorithm="maxima")`

output `-1/2*b*((log(b*x + a)*log(b*x + a - 1) + dilog(-b*x - a + 1))/(b*d) - (log(b*x + a + 1)*log(-b*x - a) + dilog(b*x + a + 1))/(b*d)) - 1/2*b*(log(b*x + a + 1)/b - log(b*x + a - 1)/b)*log(d*x + a*d/b)/d + arctanh(b*x + a)*log(d*x + a*d/b)/d`

Giac [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{artanh}(bx + a)}{dx + \frac{ad}{b}} dx$$

input `integrate(arctanh(b*x+a)/(a*d/b+d*x),x, algorithm="giac")`

output `integrate(arctanh(b*x + a)/(d*x + a*d/b), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{atanh}(a + bx)}{dx + \frac{ad}{b}} dx$$

input `int(atanh(a + b*x)/(d*x + (a*d)/b),x)`

output `int(atanh(a + b*x)/(d*x + (a*d)/b), x)`

Reduce [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{\frac{ad}{b} + dx} dx = \frac{\left(\int \frac{\operatorname{atanh}(bx+a)}{bx+a} dx\right) b}{d}$$

input `int(atanh(b*x+a)/(a*d/b+d*x),x)`

output `(int(atanh(a + b*x)/(a + b*x),x)*b)/d`

3.31 $\int (e + fx)^3 (a + \operatorname{barctanh}(c + dx)) dx$

| | |
|---|-----|
| Optimal result | 297 |
| Mathematica [A] (verified) | 298 |
| Rubi [A] (verified) | 298 |
| Maple [B] (verified) | 300 |
| Fricas [B] (verification not implemented) | 301 |
| Sympy [B] (verification not implemented) | 302 |
| Maxima [B] (verification not implemented) | 303 |
| Giac [B] (verification not implemented) | 303 |
| Mupad [B] (verification not implemented) | 305 |
| Reduce [B] (verification not implemented) | 306 |

Optimal result

Integrand size = 18, antiderivative size = 168

$$\int (e + fx)^3 (a + \operatorname{barctanh}(c + dx)) dx = \frac{bf(6d^2e^2 - 12cdef + (1 + 6c^2)f^2)x}{4d^3} + \frac{bf^2(de - cf)(c + dx)^2}{2d^4} + \frac{bf^3(c + dx)^3}{12d^4} + \frac{(e + fx)^4(a + \operatorname{barctanh}(c + dx))}{4f} + \frac{b(de + f - cf)^4 \log(1 - c - dx)}{8d^4 f} - \frac{b(de - f - cf)^4 \log(1 + c + dx)}{8d^4 f}$$

output

```
1/4*b*f*(6*d^2*e^2-12*c*d*e*f+(6*c^2+1)*f^2)*x/d^3+1/2*b*f^2*(-c*f+d*e)*(d*x+c)^2/d^4+1/12*b*f^3*(d*x+c)^3/d^4+1/4*(f*x+e)^4*(a+b*arctanh(d*x+c))/f+1/8*b*(-c*f+d*e+f)^4*ln(-d*x-c+1)/d^4/f-1/8*b*(-c*f+d*e-f)^4*ln(d*x+c+1)/d^4/f
```

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.61

$$\int (e + fx)^3 (a + \operatorname{barctanh}(c + dx)) dx$$

$$= \frac{6d(4ad^3e^3 + bf(6d^2e^2 - 8cdef + (1 + 3c^2)f^2))x + 6d^2f(6ad^2e^2 + bf(2de - cf))x^2 + 2d^3f^2(12ade + b$$

input `Integrate[(e + f*x)^3*(a + b*ArcTanh[c + d*x]),x]`

output $(6*d*(4*a*d^3*e^3 + b*f*(6*d^2*e^2 - 8*c*d*e*f + (1 + 3*c^2)*f^2))*x + 6*d^2*f*(6*a*d^2*e^2 + b*f*(2*d*e - c*f))*x^2 + 2*d^3*f^2*(12*a*d*e + b*f)*x^3 + 6*a*d^4*f^3*x^4 + 6*b*d^4*x*(4*e^3 + 6*e^2*f*x + 4*e*f^2*x^2 + f^3*x^3)*\operatorname{ArcTanh}[c + d*x] - 3*b*(-1 + c)*(4*d^3*e^3 - 6*(-1 + c)*d^2*e^2*f + 4*(-1 + c)^2*d*e*f^2 - (-1 + c)^3*f^3)*\operatorname{Log}[1 - c - d*x] - 3*b*(1 + c)*(-4*d^3*e^3 + 6*(1 + c)*d^2*e^2*f - 4*(1 + c)^2*d*e*f^2 + (1 + c)^3*f^3)*\operatorname{Log}[1 + c + d*x])/(24*d^4)$

Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.99, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {6661, 27, 6478, 477, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)^3 (a + \operatorname{barctanh}(c + dx)) dx$$

$$\downarrow \text{6661}$$

$$\int \frac{\left(\frac{d\left(e - \frac{cf}{d}\right) + f(c + dx)}{d^3}\right)^3 (a + \operatorname{barctanh}(c + dx))}{d} d(c + dx)$$

$$\downarrow \text{27}$$

$$\int \frac{(de - cf + f(c + dx))^3 (a + \operatorname{barctanh}(c + dx))}{d^4} d(c + dx)$$

$$\begin{array}{c}
 \downarrow 6478 \\
 \frac{(f(c+dx)-cf+de)^4(a+b\operatorname{arctanh}(c+dx))}{4f} - \frac{b \int \frac{(de-cf+f(c+dx))^4}{1-(c+dx)^2} d(c+dx)}{4f} \\
 \hline
 d^4 \\
 \downarrow 477 \\
 \frac{(f(c+dx)-cf+de)^4(a+b\operatorname{arctanh}(c+dx))}{4f} - \frac{b \int \left(-(c+dx)^2 f^4 - 4(de-cf)(c+dx)f^3 - (6d^2e^2 - 12cdf e + (6c^2+1)f^2)f^2 + \frac{(de-cf+f)^4}{2(-c-dx+1)} + \frac{(de-cf-f)^4}{2(c+dx+1)} \right) d(c+dx)}{4f} \\
 \hline
 d^4 \\
 \downarrow 2009 \\
 \frac{(f(c+dx)-cf+de)^4(a+b\operatorname{arctanh}(c+dx))}{4f} - \frac{b(-f^2(c+dx)((6c^2+1)f^2 - 12cdf e + 6d^2e^2) - 2f^3(c+dx)^2(de-cf) - \frac{1}{2}(-cf+de+f)^4 \log(-c-dx+1))}{4f} \\
 \hline
 d^4
 \end{array}$$

input `Int[(e + f*x)^3*(a + b*ArcTanh[c + d*x]),x]`

output `((((d*e - c*f + f*(c + d*x))^4*(a + b*ArcTanh[c + d*x]))/(4*f) - (b*(-(f^2*(6*d^2*e^2 - 12*c*d*e*f + (1 + 6*c^2)*f^2)*(c + d*x)) - 2*f^3*(d*e - c*f)*(c + d*x)^2 - (f^4*(c + d*x)^3)/3 - ((d*e + f - c*f)^4*Log[1 - c - d*x])/2 + ((d*e - f - c*f)^4*Log[1 + c + d*x])/2))/(4*f))/d^4`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 477 `Int[((c_) + (d_.)*(x_))^(n_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p Int[ExpandIntegrand[(c + d*x)^n*(1 - Rt[-b/a, 2]*x)^p*(1 + Rt[-b/a, 2]*x)^p, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IntegerQ[n] && NiceSqrtQ[-b/a] && !FractionalPowerFactorQ[Rt[-b/a, 2]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6478

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))*((d_) + (e_.)*(x_.))^(q_.), x_Symbol]
  := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])/(e*(q + 1))), x] - Simp[b
  *(c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 - c^2*x^2), x], x] /; FreeQ[{a,
  b, c, d, e, q}, x] && NeQ[q, -1]
```

rule 6661

```
Int[((a_.) + ArcTanh[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(
m_.), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*
ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IG
tQ[p, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 598 vs. 2(156) = 312.

Time = 0.76 (sec) , antiderivative size = 599, normalized size of antiderivative = 3.57

| method | result |
|-------------------|---|
| parallelrisc | $-\frac{24ac d^3 e^3 - 18f e^2 a d^2 - 6be f^2 d + 9bc f^3 + 18 \operatorname{arctanh}(dx+c) b c^2 f^3 + 12 \operatorname{arctanh}(dx+c) bc f^3 + 12 \operatorname{arctanh}(dx+c) b c^3}{4d^3 f}$ |
| derivativedivides | $\frac{a(cf - de - f(dx+c))^4}{4d^3 f} - \frac{b \left(-\frac{f^3 \operatorname{arctanh}(dx+c)c^4}{4} + f^2 \operatorname{arctanh}(dx+c)c^3 de + f^3 \operatorname{arctanh}(dx+c)c^3(dx+c) - \frac{3f \operatorname{arctanh}(dx+c)c^2 d^2 e^2}{2} \right)}{4d^3 f}$ |
| default | $\frac{a(cf - de - f(dx+c))^4}{4d^3 f} - \frac{b \left(-\frac{f^3 \operatorname{arctanh}(dx+c)c^4}{4} + f^2 \operatorname{arctanh}(dx+c)c^3 de + f^3 \operatorname{arctanh}(dx+c)c^3(dx+c) - \frac{3f \operatorname{arctanh}(dx+c)c^2 d^2 e^2}{2} \right)}{4d^3 f}$ |
| parts | $\frac{a(fx+e)^4}{4f} + \frac{b \left(\frac{f^3 \operatorname{arctanh}(dx+c)(dx+c)^4}{4d^3} - \frac{f^3 \operatorname{arctanh}(dx+c)(dx+c)^3 c}{d^3} + \frac{f^2 \operatorname{arctanh}(dx+c)(dx+c)^3 e}{d^2} + \frac{3f^3 \operatorname{arctanh}(dx+c)(dx+c)^2 d^2 e^2}{2d^3} \right)}{4d^3 f}$ |
| risc | $\frac{b e^3 \ln(dx+c+1)}{2d} + \frac{b e^3 \ln(-dx-c+1)}{2d} - \frac{2f^2 bcex}{d^2} + \frac{f^2 \ln(dx+c+1) b c^3 e}{2d^3} - \frac{3f \ln(dx+c+1) b c^2 e^2}{4d^2} - \frac{f^2 \ln(-dx-c+1) b c^3 e}{2d^3}$ |

input

```
int((f*x+e)^3*(a+b*arctanh(d*x+c)),x,method=_RETURNVERBOSE)
```

output

```
-1/12*(24*a*c*d^3*e^3-18*f*e^2*a*d^2-6*b*e*f^2*d+9*b*c*f^3+18*arctanh(d*x+c)*b*c^2*f^3+12*arctanh(d*x+c)*b*c*f^3+12*arctanh(d*x+c)*b*c^3*f^3+12*ln(d*x+c-1)*b*c^3*f^3-12*ln(d*x+c-1)*b*d^3*e^3+12*ln(d*x+c-1)*b*c*f^3-3*x^4*a*d^4*f^3-12*x*a*d^4*e^3-3*x*b*d*f^3-x^3*b*d^3*f^3+3*arctanh(d*x+c)*b*c^4*f^3-12*arctanh(d*x+c)*b*d^3*e^3-3*x^4*arctanh(d*x+c)*b*d^4*f^3-18*x^2*a*d^4*e^2*f-12*ln(d*x+c-1)*b*d*e*f^2-9*x*b*c^2*d*f^3-18*x*b*d^3*e^2*f-12*x^3*a*d^4*e*f^2-12*x*arctanh(d*x+c)*b*d^4*e^3-12*arctanh(d*x+c)*b*c*d^3*e^3+18*arctanh(d*x+c)*b*d^2*e^2*f-12*arctanh(d*x+c)*b*d*e*f^2+3*x^2*b*c*d^2*f^3-6*x^2*b*d^3*e*f^2+3*arctanh(d*x+c)*b*f^3+18*a*c^2*d^2*e^2*f+18*arctanh(d*x+c)*b*c^2*d^2*e^2*f-36*ln(d*x+c-1)*b*c^2*d*e*f^2+36*ln(d*x+c-1)*b*c*d^2*e^2*f-12*arctanh(d*x+c)*b*c^3*d*e*f^2-36*arctanh(d*x+c)*b*c^2*d*e*f^2+36*arctanh(d*x+c)*b*c*d^2*e^2*f-36*arctanh(d*x+c)*b*c*d*e*f^2-12*x^3*arctanh(d*x+c)*b*d^4*e*f^2+24*x*b*c*d^2*e*f^2-18*x^2*arctanh(d*x+c)*b*d^4*e^2*f-42*b*c^2*d*e*f^2+36*b*c*d^2*e^2*f+15*b*c^3*f^3)/d^4
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 386 vs. $2(156) = 312$.

Time = 0.10 (sec) , antiderivative size = 386, normalized size of antiderivative = 2.30

$$\int (e + fx)^3 (a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \frac{6 ad^4 f^3 x^4 + 2(12 ad^4 e f^2 + bd^3 f^3) x^3 + 6(6 ad^4 e^2 f + 2 bd^3 e f^2 - bcd^2 f^3) x^2 + 6(4 ad^4 e^3 + 6 bd^3 e^2 f - 8 bcd^2 e f^2 + 4 ad^4 e^2 f^2 + 2 bd^3 e f^3) x + 6(4 ad^4 e^3 + 6 bd^3 e^2 f - 8 bcd^2 e f^2 + 4 ad^4 e^2 f^2 + 2 bd^3 e f^3)}{d^4}$$

input

```
integrate((f*x+e)^3*(a+b*arctanh(d*x+c)),x, algorithm="fricas")
```

output

```
1/24*(6*a*d^4*f^3*x^4 + 2*(12*a*d^4*e*f^2 + b*d^3*f^3)*x^3 + 6*(6*a*d^4*e^2*f + 2*b*d^3*e*f^2 - b*c*d^2*f^3)*x^2 + 6*(4*a*d^4*e^3 + 6*b*d^3*e^2*f - 8*b*c*d^2*e*f^2 + (3*b*c^2 + b)*d*f^3)*x + 3*(4*(b*c + b)*d^3*e^3 - 6*(b*c^2 + 2*b*c + b)*d^2*e^2*f + 4*(b*c^3 + 3*b*c^2 + 3*b*c + b)*d*e*f^2 - (b*c^4 + 4*b*c^3 + 6*b*c^2 + 4*b*c + b)*f^3)*log(d*x + c + 1) - 3*(4*(b*c - b)*d^3*e^3 - 6*(b*c^2 - 2*b*c + b)*d^2*e^2*f + 4*(b*c^3 - 3*b*c^2 + 3*b*c - b)*d*e*f^2 - (b*c^4 - 4*b*c^3 + 6*b*c^2 - 4*b*c + b)*f^3)*log(d*x + c - 1) + 3*(b*d^4*f^3*x^4 + 4*b*d^4*e*f^2*x^3 + 6*b*d^4*e^2*f*x^2 + 4*b*d^4*e^3*x)*log(-(d*x + c + 1)/(d*x + c - 1))/d^4
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 644 vs. $2(151) = 302$.

Time = 2.02 (sec) , antiderivative size = 644, normalized size of antiderivative = 3.83

$$\int (e + fx)^3 (a + b \operatorname{arctanh}(c + dx)) dx = \text{Too large to display}$$

input `integrate((f*x+e)**3*(a+b*atanh(d*x+c)),x)`

output

```
Piecewise((a*e**3*x + 3*a*e**2*f*x**2/2 + a*e*f**2*x**3 + a*f**3*x**4/4 -
b*c**4*f**3*atanh(c + d*x)/(4*d**4) + b*c**3*e*f**2*atanh(c + d*x)/d**3 -
b*c**3*f**3*log(c/d + x + 1/d)/d**4 + b*c**3*f**3*atanh(c + d*x)/d**4 - 3*
b*c**2*e**2*f*atanh(c + d*x)/(2*d**2) + 3*b*c**2*e*f**2*log(c/d + x + 1/d)
/d**3 - 3*b*c**2*e*f**2*atanh(c + d*x)/d**3 + 3*b*c**2*f**3*x/(4*d**3) - 3
*b*c**2*f**3*atanh(c + d*x)/(2*d**4) + b*c*e**3*atanh(c + d*x)/d - 3*b*c*e
**2*f*log(c/d + x + 1/d)/d**2 + 3*b*c*e**2*f*atanh(c + d*x)/d**2 - 2*b*c*e
*f**2*x/d**2 - b*c*f**3*x**2/(4*d**2) + 3*b*c*e*f**2*atanh(c + d*x)/d**3 -
b*c*f**3*log(c/d + x + 1/d)/d**4 + b*c*f**3*atanh(c + d*x)/d**4 + b*e**3*
x*atanh(c + d*x) + 3*b*e**2*f*x**2*atanh(c + d*x)/2 + b*e*f**2*x**3*atanh(
c + d*x) + b*f**3*x**4*atanh(c + d*x)/4 + b*e**3*log(c/d + x + 1/d)/d - b*
e**3*atanh(c + d*x)/d + 3*b*e**2*f*x/(2*d) + b*e*f**2*x**2/(2*d) + b*f**3*
x**3/(12*d) - 3*b*e**2*f*atanh(c + d*x)/(2*d**2) + b*e*f**2*log(c/d + x +
1/d)/d**3 - b*e*f**2*atanh(c + d*x)/d**3 + b*f**3*x/(4*d**3) - b*f**3*atan
h(c + d*x)/(4*d**4), Ne(d, 0)), ((a + b*atanh(c))*(e**3*x + 3*e**2*f*x**2/
2 + e*f**2*x**3 + f**3*x**4/4), True))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 333 vs. $2(156) = 312$.

Time = 0.03 (sec) , antiderivative size = 333, normalized size of antiderivative = 1.98

$$\int (e + fx)^3 (a + b \operatorname{arctanh}(c + dx)) dx = \frac{1}{4} a f^3 x^4 + a e f^2 x^3 + \frac{3}{2} a e^2 f x^2 + \frac{3}{4} \left(2 x^2 \operatorname{arctanh}(dx + c) + d \left(\frac{2x}{d^2} - \frac{(c^2 + 2c + 1) \log(dx + c + 1)}{d^3} + \frac{(c^2 - 2c + 1) \log(dx + c - 1)}{d^3} \right) \right) + \frac{1}{2} \left(2 x^3 \operatorname{arctanh}(dx + c) + d \left(\frac{dx^2 - 4cx}{d^3} + \frac{(c^3 + 3c^2 + 3c + 1) \log(dx + c + 1)}{d^4} - \frac{(c^3 - 3c^2 + 3c - 1) \log(dx + c - 1)}{d^4} \right) \right) + \frac{1}{24} \left(6 x^4 \operatorname{arctanh}(dx + c) + d \left(\frac{2(d^2 x^3 - 3cdx^2 + 3(3c^2 + 1)x)}{d^4} - \frac{3(c^4 + 4c^3 + 6c^2 + 4c + 1) \log(dx + c + 1)}{d^5} + \frac{3(c^4 - 4c^3 + 6c^2 - 4c + 1) \log(dx + c - 1)}{d^5} \right) \right) + a e^3 x + \frac{(2(dx + c) \operatorname{arctanh}(dx + c) + \log(-(dx + c)^2 + 1)) b e^3}{2d}$$

input `integrate((f*x+e)^3*(a+b*arctanh(d*x+c)),x, algorithm="maxima")`

output `1/4*a*f^3*x^4 + a*e*f^2*x^3 + 3/2*a*e^2*f*x^2 + 3/4*(2*x^2*arctanh(d*x + c) + d*(2*x/d^2 - (c^2 + 2*c + 1)*log(d*x + c + 1)/d^3 + (c^2 - 2*c + 1)*log(d*x + c - 1)/d^3))*b*e^2*f + 1/2*(2*x^3*arctanh(d*x + c) + d*((d*x^2 - 4*c*x)/d^3 + (c^3 + 3*c^2 + 3*c + 1)*log(d*x + c + 1)/d^4 - (c^3 - 3*c^2 + 3*c - 1)*log(d*x + c - 1)/d^4))*b*e*f^2 + 1/24*(6*x^4*arctanh(d*x + c) + d*(2*(d^2*x^3 - 3*c*d*x^2 + 3*(3*c^2 + 1)*x)/d^4 - 3*(c^4 + 4*c^3 + 6*c^2 + 4*c + 1)*log(d*x + c + 1)/d^5 + 3*(c^4 - 4*c^3 + 6*c^2 - 4*c + 1)*log(d*x + c - 1)/d^5))*b*f^3 + a*e^3*x + 1/2*(2*(d*x + c)*arctanh(d*x + c) + log(-(d*x + c)^2 + 1))*b*e^3/d`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2336 vs. $2(156) = 312$.

Time = 0.19 (sec) , antiderivative size = 2336, normalized size of antiderivative = 13.90

$$\int (e + fx)^3 (a + b \operatorname{arctanh}(c + dx)) dx = \text{Too large to display}$$

input `integrate((f*x+e)^3*(a+b*arctanh(d*x+c)),x, algorithm="giac")`

output

```

1/6*((c + 1)*d - (c - 1)*d)*(3*((d*x + c + 1)^3*b*d^3*e^3/(d*x + c - 1)^3
- 3*(d*x + c + 1)^2*b*d^3*e^3/(d*x + c - 1)^2 + 3*(d*x + c + 1)*b*d^3*e^3/
(d*x + c - 1) - b*d^3*e^3 - 3*(d*x + c + 1)^3*b*c*d^2*e^2*f/(d*x + c - 1)^
3 + 9*(d*x + c + 1)^2*b*c*d^2*e^2*f/(d*x + c - 1)^2 - 9*(d*x + c + 1)*b*c*
d^2*e^2*f/(d*x + c - 1) + 3*b*c*d^2*e^2*f + 3*(d*x + c + 1)^3*b*c^2*d*e*f^
2/(d*x + c - 1)^3 - 9*(d*x + c + 1)^2*b*c^2*d*e*f^2/(d*x + c - 1)^2 + 9*(d
*x + c + 1)*b*c^2*d*e*f^2/(d*x + c - 1) - 3*b*c^2*d*e*f^2 - (d*x + c + 1)^
3*b*c^3*f^3/(d*x + c - 1)^3 + 3*(d*x + c + 1)^2*b*c^3*f^3/(d*x + c - 1)^2
- 3*(d*x + c + 1)*b*c^3*f^3/(d*x + c - 1) + b*c^3*f^3 + 3*(d*x + c + 1)^3*
b*d^2*e^2*f/(d*x + c - 1)^3 - 6*(d*x + c + 1)^2*b*d^2*e^2*f/(d*x + c - 1)^
2 + 3*(d*x + c + 1)*b*d^2*e^2*f/(d*x + c - 1) - 6*(d*x + c + 1)^3*b*c*d*e*
f^2/(d*x + c - 1)^3 + 12*(d*x + c + 1)^2*b*c*d*e*f^2/(d*x + c - 1)^2 - 6*(
d*x + c + 1)*b*c*d*e*f^2/(d*x + c - 1) + 3*(d*x + c + 1)^3*b*c^2*f^3/(d*x
+ c - 1)^3 - 6*(d*x + c + 1)^2*b*c^2*f^3/(d*x + c - 1)^2 + 3*(d*x + c + 1)
*b*c^2*f^3/(d*x + c - 1) + 3*(d*x + c + 1)^3*b*d*e*f^2/(d*x + c - 1)^3 - 3
*(d*x + c + 1)^2*b*d*e*f^2/(d*x + c - 1)^2 + (d*x + c + 1)*b*d*e*f^2/(d*x
+ c - 1) - b*d*e*f^2 - 3*(d*x + c + 1)^3*b*c*f^3/(d*x + c - 1)^3 + 3*(d*x
+ c + 1)^2*b*c*f^3/(d*x + c - 1)^2 - (d*x + c + 1)*b*c*f^3/(d*x + c - 1) +
b*c*f^3 + (d*x + c + 1)^3*b*f^3/(d*x + c - 1)^3 + (d*x + c + 1)*b*f^3/(d*
x + c - 1))*log(-(d*x + c + 1)/(d*x + c - 1))/((d*x + c + 1)^4*d^5/(d*x...

```

Mupad [B] (verification not implemented)

Time = 4.70 (sec) , antiderivative size = 737, normalized size of antiderivative = 4.39

$$\begin{aligned}
& \int (e + fx)^3 (a + \operatorname{barctanh}(c + dx)) dx \\
&= \ln(c + dx + 1) \left(\frac{be^3 x}{2} + \frac{3be^2 f x^2}{4} + \frac{be f^2 x^3}{2} + \frac{b f^3 x^4}{8} \right) \\
&\quad - \ln(1 - dx - c) \left(\frac{be^3 x}{2} + \frac{3be^2 f x^2}{4} + \frac{be f^2 x^3}{2} + \frac{b f^3 x^4}{8} \right) \\
&\quad + x \left(\frac{e(6ac^2 f^2 + 12acdef + 2ad^2 e^2 + 3bdef - 6a f^2)}{2d^2} \right. \\
&\quad\quad\quad \left. - \frac{(4c^2 - 4) \left(\frac{f^2(bf + 8acf + 12ade)}{4d} - \frac{2acf^3}{d} \right)}{4d^2} \right. \\
&\quad\quad\quad \left. + \frac{2c \left(\frac{2c \left(\frac{f^2(bf + 8acf + 12ade)}{4d} - \frac{2acf^3}{d} \right)}{d} - \frac{4ac^2 f^3 + 24acdef^2 + 12ad^2 e^2 f + 4bdef^2 - 4af^3}{4d^2} + \frac{af^3(4c^2 - 4)}{4d^2} \right)}{d} \right) \\
&\quad - x^2 \left(\frac{c \left(\frac{f^2(bf + 8acf + 12ade)}{4d} - \frac{2acf^3}{d} \right)}{d} \right. \\
&\quad\quad\quad \left. - \frac{4ac^2 f^3 + 24acdef^2 + 12ad^2 e^2 f + 4bdef^2 - 4af^3}{8d^2} + \frac{af^3(4c^2 - 4)}{8d^2} \right) \\
&\quad + x^3 \left(\frac{f^2(bf + 8acf + 12ade)}{12d} - \frac{2acf^3}{3d} \right) + \frac{af^3 x^4}{4} \\
&\quad + \frac{\ln(c + dx - 1) (bc^4 f^3 - 4bc^3 def^2 - 4bc^3 f^3 + 6bc^2 d^2 e^2 f + 12bc^2 def^2 + 6bc^2 f^3 - 4bcd^3 e^3 - \frac{8d^4}{8d^4})}{8d^4} \\
&\quad - \frac{\ln(c + dx + 1) (bc^4 f^3 - 4bc^3 def^2 + 4bc^3 f^3 + 6bc^2 d^2 e^2 f - 12bc^2 def^2 + 6bc^2 f^3 - 4bcd^3 e^3 - \frac{8d^4}{8d^4})}{8d^4}
\end{aligned}$$

input `int((e + f*x)^3*(a + b*atanh(c + d*x)),x)`

output

```
( - 3*atanh(c + d*x)*b*c**4*f**3 + 12*atanh(c + d*x)*b*c**3*d*e*f**2 - 12*
atanh(c + d*x)*b*c**3*f**3 - 18*atanh(c + d*x)*b*c**2*d**2*e**2*f + 36*ata
anh(c + d*x)*b*c**2*d*e*f**2 - 18*atanh(c + d*x)*b*c**2*f**3 + 12*atanh(c +
d*x)*b*c*d**3*e**3 - 36*atanh(c + d*x)*b*c*d**2*e**2*f + 36*atanh(c + d*x
)*b*c*d*e*f**2 - 12*atanh(c + d*x)*b*c*f**3 + 12*atanh(c + d*x)*b*d**4*e**
3*x + 18*atanh(c + d*x)*b*d**4*e**2*f*x**2 + 12*atanh(c + d*x)*b*d**4*e*f*
*2*x**3 + 3*atanh(c + d*x)*b*d**4*f**3*x**4 + 12*atanh(c + d*x)*b*d**3*e**
3 - 18*atanh(c + d*x)*b*d**2*e**2*f + 12*atanh(c + d*x)*b*d*e*f**2 - 3*ata
anh(c + d*x)*b*f**3 - 12*log(c + d*x - 1)*b*c**3*f**3 + 36*log(c + d*x - 1)
*b*c**2*d*e*f**2 - 36*log(c + d*x - 1)*b*c*d**2*e**2*f - 12*log(c + d*x -
1)*b*c*f**3 + 12*log(c + d*x - 1)*b*d**3*e**3 + 12*log(c + d*x - 1)*b*d*e*
f**2 + 12*a*d**4*e**3*x + 18*a*d**4*e**2*f*x**2 + 12*a*d**4*e*f**2*x**3 +
3*a*d**4*f**3*x**4 + 9*b*c**2*d*f**3*x - 24*b*c*d**2*e*f**2*x - 3*b*c*d**2
*f**3*x**2 + 18*b*d**3*e**2*f*x + 6*b*d**3*e*f**2*x**2 + b*d**3*f**3*x**3
+ 3*b*d*f**3*x)/(12*d**4)
```

3.32 $\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx)) dx$

| | |
|---|-----|
| Optimal result | 308 |
| Mathematica [A] (verified) | 309 |
| Rubi [A] (verified) | 309 |
| Maple [B] (verified) | 311 |
| Fricas [B] (verification not implemented) | 312 |
| Sympy [B] (verification not implemented) | 312 |
| Maxima [A] (verification not implemented) | 313 |
| Giac [B] (verification not implemented) | 314 |
| Mupad [B] (verification not implemented) | 315 |
| Reduce [B] (verification not implemented) | 316 |

Optimal result

Integrand size = 18, antiderivative size = 120

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx)) dx = \frac{bf(de - cf)x}{d^2} + \frac{bf^2(c + dx)^2}{6d^3} + \frac{(e + fx)^3(a + b \operatorname{arctanh}(c + dx))}{3f} + \frac{b(de + f - cf)^3 \log(1 - c - dx)}{6d^3 f} - \frac{b(de - (1 + c)f)^3 \log(1 + c + dx)}{6d^3 f}$$

output

```
b*f*(-c*f+d*e)*x/d^2+1/6*b*f^2*(d*x+c)^2/d^3+1/3*(f*x+e)^3*(a+b*arctanh(d*x+c))/f+1/6*b*(-c*f+d*e+f)^3*ln(-d*x-c+1)/d^3/f-1/6*b*(d*e-(1+c)*f)^3*ln(d*x+c+1)/d^3/f
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.45

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \frac{2d(3ad^2e^2 + bf(3de - 2cf))x + d^2f(6ade + bf)x^2 + 2ad^3f^2x^3 + 2bd^3x(3e^2 + 3efx + f^2x^2) \operatorname{arctanh}(c + dx) - 2bd^3x(3e^2 + 3efx + f^2x^2) \operatorname{arctanh}(c - dx)}{6d^3}$$

input `Integrate[(e + f*x)^2*(a + b*ArcTanh[c + d*x]),x]`

output $(2*d*(3*a*d^2*e^2 + b*f*(3*d*e - 2*c*f))*x + d^2*f*(6*a*d*e + b*f)*x^2 + 2*a*d^3*f^2*x^3 + 2*b*d^3*x*(3*e^2 + 3*e*f*x + f^2*x^2)*\operatorname{ArcTanh}[c + d*x] - b*(-1 + c)*(3*d^2*e^2 - 3*(-1 + c)*d*e*f + (-1 + c)^2*f^2)*\operatorname{Log}[1 - c - d*x] + b*(1 + c)*(3*d^2*e^2 - 3*(1 + c)*d*e*f + (1 + c)^2*f^2)*\operatorname{Log}[1 + c + d*x])/(6*d^3)$

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.07, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {6661, 27, 6478, 477, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx)) dx$$

$$\downarrow \text{6661}$$

$$\int \frac{\left(\frac{d\left(e - \frac{cf}{d}\right) + f(c + dx)}{d^2}\right)^2 (a + b \operatorname{arctanh}(c + dx))}{d} d(c + dx)$$

$$\downarrow \text{27}$$

$$\int \frac{(de - cf + f(c + dx))^2 (a + b \operatorname{arctanh}(c + dx))}{d^3} d(c + dx)$$

$$\downarrow \text{6478}$$

$$\frac{\frac{(f(c+dx)-cf+de)^3(a+b\operatorname{arctanh}(c+dx))}{3f} - \frac{b \int \frac{(de-cf+f(c+dx))^3 d(c+dx)}{1-(c+dx)^2}}{3f}}{d^3} \xrightarrow{477} \frac{\frac{(f(c+dx)-cf+de)^3(a+b\operatorname{arctanh}(c+dx))}{3f} - \frac{b \int \left(-((c+dx)f^3) - 3(de-cf)f^2 + \frac{(de-cf+f)^3}{2(-c-dx+1)} + \frac{(de-(c+1)f)^3}{2(c+dx+1)} \right) d(c+dx)}{3f}}{d^3} \xrightarrow{2009} \frac{\frac{(f(c+dx)-cf+de)^3(a+b\operatorname{arctanh}(c+dx))}{3f} - \frac{b(-3f^2(c+dx)(de-cf) - \frac{1}{2}(-cf+de+f)^3 \log(-c-dx+1) + \frac{1}{2}(de-(c+1)f)^3 \log(c+dx+1) - \frac{1}{2}f^3(c+dx+1))}{3f}}{d^3}$$

input `Int[(e + f*x)^2*(a + b*ArcTanh[c + d*x]),x]`

output `((((d*e - c*f + f*(c + d*x))^3*(a + b*ArcTanh[c + d*x]))/(3*f) - (b*(-3*f^2*(d*e - c*f)*(c + d*x) - (f^3*(c + d*x)^2)/2 - ((d*e + f - c*f)^3*Log[1 - c - d*x])/2 + ((d*e - (1 + c)*f)^3*Log[1 + c + d*x])/2))/(3*f))/d^3`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 477 `Int[((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p Int[ExpandIntegrand[(c + d*x)^n*(1 - Rt[-b/a, 2]*x)^p*(1 + Rt[-b/a, 2]*x)^p, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IntegerQ[n] && NiceSqrtQ[-b/a] && !FractionalPowerFactorQ[Rt[-b/a, 2]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6478

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))*((d_) + (e_.)*(x_.))^(q_.), x_Symbol]
-> Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])/(e*(q + 1))), x] - Simp[b
*(c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 - c^2*x^2), x], x] /; FreeQ[{a,
b, c, d, e, q}, x] && NeQ[q, -1]
```

rule 6661

```
Int[((a_.) + ArcTanh[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(
m_.), x_Symbol] :> Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*
ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IG
tQ[p, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 350 vs. 2(112) = 224.

Time = 0.53 (sec) , antiderivative size = 351, normalized size of antiderivative = 2.92

| method | result |
|-------------------|---|
| parallelrisch | $b f^2 - 12ac d^2 e^2 + 6aef d + 6 \operatorname{arctanh}(dx+c)bc d^2 e^2 - 6 \operatorname{arctanh}(dx+c)bde f - 4x bcd f^2 + 6xb d^2 e f + 6x^2 a d^3 e f + 2x^3 \operatorname{arctanh}(dx+c)bc d^2 e^2$ |
| parts | $\frac{a(fx+e)^3}{3f} + \frac{b \left(\frac{f^2 \operatorname{arctanh}(dx+c)(dx+c)^3}{3d^2} - \frac{f^2 \operatorname{arctanh}(dx+c)(dx+c)^2 c}{d^2} + \frac{f \operatorname{arctanh}(dx+c)(dx+c)^2 e}{d} + \frac{f^2 \operatorname{arctanh}(dx+c)(dx+c)}{d^2} \right)}{3d^2 f}$ |
| derivativedivides | $-\frac{a(cf-de-f(dx+c))^3}{3d^2 f} + \frac{b \left(-\frac{f^2 \operatorname{arctanh}(dx+c)c^3}{3} + f \operatorname{arctanh}(dx+c)c^2 de + f^2 \operatorname{arctanh}(dx+c)c^2(dx+c) - \operatorname{arctanh}(dx+c)c d^2 e^2 - 2f \operatorname{arctanh}(dx+c)c d^2 e \right)}{3d^2 f}$ |
| default | $-\frac{a(cf-de-f(dx+c))^3}{3d^2 f} + \frac{b \left(-\frac{f^2 \operatorname{arctanh}(dx+c)c^3}{3} + f \operatorname{arctanh}(dx+c)c^2 de + f^2 \operatorname{arctanh}(dx+c)c^2(dx+c) - \operatorname{arctanh}(dx+c)c d^2 e^2 - 2f \operatorname{arctanh}(dx+c)c d^2 e \right)}{3d^2 f}$ |
| risch | $-\frac{f \ln(-dx-c-1)bc^2 e}{2d^2} + \frac{f \ln(dx+c-1)bc^2 e}{2d^2} - \frac{f \ln(-dx-c-1)bce}{d^2} - \frac{f \ln(dx+c-1)bce}{d^2} + fae x^2 + ae^2 x$ |

input

```
int((f*x+e)^2*(a+b*arctanh(d*x+c)),x,method=_RETURNVERBOSE)
```


output

```
1/6*(b*f^2-12*a*c*d^2*e^2+6*a*e*f*d+6*arctanh(d*x+c)*b*c*d^2*e^2-6*arctanh
(d*x+c)*b*d*e*f-4*x*b*c*d*f^2+6*x*b*d^2*e*f+6*x^2*a*d^3*e*f+2*x^3*arctanh(
d*x+c)*b*d^3*f^2+6*x*arctanh(d*x+c)*b*d^3*e^2+2*arctanh(d*x+c)*b*f^2+2*ln(
d*x+c-1)*b*f^2-6*arctanh(d*x+c)*b*c^2*d*e*f-12*ln(d*x+c-1)*b*c*d*e*f+6*x^2
*arctanh(d*x+c)*b*d^3*e*f-12*arctanh(d*x+c)*b*c*d*e*f+x^2*b*d^2*f^2+6*ln(d
*x+c-1)*b*c^2*f^2+6*ln(d*x+c-1)*b*d^2*e^2+2*arctanh(d*x+c)*b*c^3*f^2+6*arc
tanh(d*x+c)*b*c^2*f^2+6*arctanh(d*x+c)*b*d^2*e^2+6*arctanh(d*x+c)*b*c*f^2+
6*x*a*d^3*e^2+2*x^3*a*d^3*f^2-6*a*c^2*e*f*d+7*b*c^2*f^2-12*b*c*d*e*f)/d^3
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 242 vs. $2(112) = 224$.

Time = 0.10 (sec) , antiderivative size = 242, normalized size of antiderivative = 2.02

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \frac{2ad^3f^2x^3 + (6ad^3ef + bd^2f^2)x^2 + 2(3ad^3e^2 + 3bd^2ef - 2bcd^2f^2)x + (3(bc + b)d^2e^2 - 3(bc^2 + 2bc + b)d^2e^2 - 3(b^2c + b^2d^2e^2 - 2b^2c^2 - 2b^2d^2e^2 + b^2d^2e^2))x + (b^2c^3 + 3b^2c^2 + 3b^2c + b^2d^2e^2)f^2 \log(d*x + c + 1) - (3(b^2c - b^2d^2e^2 - 3(b^2c^2 - 2b^2c + b^2d^2e^2))d*e*f + (b^2c^3 + 3b^2c^2 + 3b^2c + b^2d^2e^2)*f^2)*\log(d*x + c - 1) + (b*d^3*f^2*x^3 + 3*b*d^3*e*f*x^2 + 3*b*d^3*e^2*x)*\log(-(d*x + c + 1)/(d*x + c - 1))}{d^3}$$

input

```
integrate((f*x+e)^2*(a+b*arctanh(d*x+c)),x, algorithm="fricas")
```

output

```
1/6*(2*a*d^3*f^2*x^3 + (6*a*d^3*e*f + b*d^2*f^2)*x^2 + 2*(3*a*d^3*e^2 + 3*
b*d^2*e*f - 2*b*c*d*f^2)*x + (3*(b*c + b)*d^2*e^2 - 3*(b*c^2 + 2*b*c + b)*
d*e*f + (b*c^3 + 3*b*c^2 + 3*b*c + b)*f^2)*log(d*x + c + 1) - (3*(b*c - b)
*d^2*e^2 - 3*(b*c^2 - 2*b*c + b)*d*e*f + (b*c^3 - 3*b*c^2 + 3*b*c - b)*f^2
)*log(d*x + c - 1) + (b*d^3*f^2*x^3 + 3*b*d^3*e*f*x^2 + 3*b*d^3*e^2*x)*log
(-(d*x + c + 1)/(d*x + c - 1))/d^3
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 369 vs. $2(105) = 210$.

Time = 1.84 (sec) , antiderivative size = 369, normalized size of antiderivative = 3.08

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \begin{cases} ae^2x + aefx^2 + \frac{af^2x^3}{3} + \frac{bc^3f^2 \operatorname{atanh}(c+dx)}{3d^3} - \frac{bc^2ef \operatorname{atanh}(c+dx)}{d^2} + \frac{bc^2f^2 \log\left(\frac{c}{d} + x + \frac{1}{d}\right)}{d^3} - \frac{bc^2f^2 \operatorname{atanh}(c+dx)}{d^3} + \frac{bce^2 \operatorname{atanh}(c+dx)}{d} \\ (a + b \operatorname{atanh}(c)) \left(e^2x + efx^2 + \frac{f^2x^3}{3} \right) \end{cases}$$

input `integrate((f*x+e)**2*(a+b*atanh(d*x+c)),x)`

output `Piecewise((a*e**2*x + a*e*f*x**2 + a*f**2*x**3/3 + b*c**3*f**2*atanh(c + d*x)/(3*d**3) - b*c**2*e*f*atanh(c + d*x)/d**2 + b*c**2*f**2*log(c/d + x + 1/d)/d**3 - b*c**2*f**2*atanh(c + d*x)/d**3 + b*c*e**2*atanh(c + d*x)/d - 2*b*c*e*f*log(c/d + x + 1/d)/d**2 + 2*b*c*e*f*atanh(c + d*x)/d**2 - 2*b*c*f**2*x/(3*d**2) + b*c*f**2*atanh(c + d*x)/d**3 + b*e**2*x*atanh(c + d*x) + b*e*f*x**2*atanh(c + d*x) + b*f**2*x**3*atanh(c + d*x)/3 + b*e**2*log(c/d + x + 1/d)/d - b*e**2*atanh(c + d*x)/d + b*e*f*x/d + b*f**2*x**2/(6*d) - b*e*f*atanh(c + d*x)/d**2 + b*f**2*log(c/d + x + 1/d)/(3*d**3) - b*f**2*atanh(c + d*x)/(3*d**3), Ne(d, 0)), ((a + b*atanh(c))*(e**2*x + e*f*x**2 + f**2*x**3/3), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.72

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx)) dx = \frac{1}{3} af^2 x^3 + aefx^2$$

$$+ \frac{1}{2} \left(2x^2 \operatorname{artanh}(dx + c) + d \left(\frac{2x}{d^2} - \frac{(c^2 + 2c + 1) \log(dx + c + 1)}{d^3} + \frac{(c^2 - 2c + 1) \log(dx + c - 1)}{d^3} \right) \right)$$

$$+ \frac{1}{6} \left(2x^3 \operatorname{artanh}(dx + c) + d \left(\frac{dx^2 - 4cx}{d^3} + \frac{(c^3 + 3c^2 + 3c + 1) \log(dx + c + 1)}{d^4} - \frac{(c^3 - 3c^2 + 3c - 1) \log(dx + c - 1)}{d^4} \right) \right)$$

$$+ ae^2x + \frac{(2(dx + c) \operatorname{artanh}(dx + c) + \log(-(dx + c)^2 + 1)) be^2}{2d}$$

input `integrate((f*x+e)^2*(a+b*arctanh(d*x+c)),x, algorithm="maxima")`

output

```
1/3*a*f^2*x^3 + a*e*f*x^2 + 1/2*(2*x^2*arctanh(d*x + c) + d*(2*x/d^2 - (c^
2 + 2*c + 1)*log(d*x + c + 1)/d^3 + (c^2 - 2*c + 1)*log(d*x + c - 1)/d^3))
*b*e*f + 1/6*(2*x^3*arctanh(d*x + c) + d*((d*x^2 - 4*c*x)/d^3 + (c^3 + 3*c
^2 + 3*c + 1)*log(d*x + c + 1)/d^4 - (c^3 - 3*c^2 + 3*c - 1)*log(d*x + c -
1)/d^4))*b*f^2 + a*e^2*x + 1/2*(2*(d*x + c)*arctanh(d*x + c) + log(-(d*x
+ c)^2 + 1))*b*e^2/d
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 976 vs. $2(112) = 224$.

Time = 0.15 (sec) , antiderivative size = 976, normalized size of antiderivative = 8.13

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx)) dx = \text{Too large to display}$$

input

```
integrate((f*x+e)^2*(a+b*arctanh(d*x+c)),x, algorithm="giac")
```

output

```
1/6*((c + 1)*d - (c - 1)*d)*((3*(d*x + c + 1)^2*b*d^2*e^2/(d*x + c - 1)^2
- 6*(d*x + c + 1)*b*d^2*e^2/(d*x + c - 1) + 3*b*d^2*e^2 - 6*(d*x + c + 1)^
2*b*c*d*e*f/(d*x + c - 1)^2 + 12*(d*x + c + 1)*b*c*d*e*f/(d*x + c - 1) - 6
*b*c*d*e*f + 3*(d*x + c + 1)^2*b*c^2*f^2/(d*x + c - 1)^2 - 6*(d*x + c + 1)
*b*c^2*f^2/(d*x + c - 1) + 3*b*c^2*f^2 + 6*(d*x + c + 1)^2*b*d*e*f/(d*x +
c - 1)^2 - 6*(d*x + c + 1)*b*d*e*f/(d*x + c - 1) - 6*(d*x + c + 1)^2*b*c*f
^2/(d*x + c - 1)^2 + 6*(d*x + c + 1)*b*c*f^2/(d*x + c - 1) + 3*(d*x + c +
1)^2*b*f^2/(d*x + c - 1)^2 + b*f^2)*log(-(d*x + c + 1)/(d*x + c - 1))/((d*
x + c + 1)^3*d^4/(d*x + c - 1)^3 - 3*(d*x + c + 1)^2*d^4/(d*x + c - 1)^2 +
3*(d*x + c + 1)*d^4/(d*x + c - 1) - d^4) + 2*(3*(d*x + c + 1)^2*a*d^2*e^2
/(d*x + c - 1)^2 - 6*(d*x + c + 1)*a*d^2*e^2/(d*x + c - 1) + 3*a*d^2*e^2 -
6*(d*x + c + 1)^2*a*c*d*e*f/(d*x + c - 1)^2 + 12*(d*x + c + 1)*a*c*d*e*f/
(d*x + c - 1) - 6*a*c*d*e*f + 3*(d*x + c + 1)^2*a*c^2*f^2/(d*x + c - 1)^2
- 6*(d*x + c + 1)*a*c^2*f^2/(d*x + c - 1) + 3*a*c^2*f^2 + 6*(d*x + c + 1)^
2*a*d*e*f/(d*x + c - 1)^2 - 6*(d*x + c + 1)*a*d*e*f/(d*x + c - 1) + 3*(d*x
+ c + 1)^2*b*d*e*f/(d*x + c - 1)^2 - 6*(d*x + c + 1)*b*d*e*f/(d*x + c - 1
) + 3*b*d*e*f - 6*(d*x + c + 1)^2*a*c*f^2/(d*x + c - 1)^2 + 6*(d*x + c + 1
)*a*c*f^2/(d*x + c - 1) - 3*(d*x + c + 1)^2*b*c*f^2/(d*x + c - 1)^2 + 6*(d
*x + c + 1)*b*c*f^2/(d*x + c - 1) - 3*b*c*f^2 + 3*(d*x + c + 1)^2*a*f^2/(d
*x + c - 1)^2 + a*f^2 + (d*x + c + 1)^2*b*f^2/(d*x + c - 1)^2 - (d*x + ...
```

Mupad [B] (verification not implemented)

Time = 4.21 (sec) , antiderivative size = 381, normalized size of antiderivative = 3.18

$$\begin{aligned}
\int (e + fx)^2 (a + \operatorname{barctanh}(c + dx)) dx &= x^2 \left(\frac{f(bf + 6acf + 6ade)}{6d} - \frac{acf^2}{d} \right) \\
&- \ln(1 - dx - c) \left(\frac{be^2x}{2} + \frac{befx^2}{2} + \frac{bf^2x^3}{6} \right) - x \left(\frac{2c \left(\frac{f(bf + 6acf + 6ade)}{3d} - \frac{2acf^2}{d} \right)}{d} \right. \\
&\quad \left. - \frac{3ac^2f^2 + 12acdef + 3ad^2e^2 + 3bdef - 3af^2}{3d^2} + \frac{af^2(3c^2 - 3)}{3d^2} \right) \\
&+ \ln(c + dx + 1) \left(\frac{be^2x}{2} + \frac{befx^2}{2} + \frac{bf^2x^3}{6} \right) + \frac{af^2x^3}{3} \\
&+ \frac{\ln(c + dx - 1) \left(\frac{bf^2}{6} + d \left(\frac{befc^2}{2} - bef c + \frac{bef}{2} \right) + d^2 \left(\frac{be^2}{2} - \frac{bce^2}{2} \right) + \frac{bc^2f^2}{2} - \frac{bc^3f^2}{6} - \frac{bcf^2}{2} \right)}{d^3} \\
&+ \frac{\ln(c + dx + 1) \left(\frac{bf^2}{6} - d \left(\frac{befc^2}{2} + bef c + \frac{bef}{2} \right) + d^2 \left(\frac{be^2}{2} + \frac{bce^2}{2} \right) + \frac{bc^2f^2}{2} + \frac{bc^3f^2}{6} + \frac{bcf^2}{2} \right)}{d^3}
\end{aligned}$$

input `int((e + f*x)^2*(a + b*atanh(c + d*x)),x)`output

```

x^2*((f*(b*f + 6*a*c*f + 6*a*d*e))/(6*d) - (a*c*f^2)/d) - log(1 - d*x - c)
*((b*f^2*x^3)/6 + (b*e^2*x)/2 + (b*e*f*x^2)/2) - x*((2*c*(f*(b*f + 6*a*c*
f + 6*a*d*e))/(3*d) - (2*a*c*f^2)/d))/d - (3*a*c^2*f^2 - 3*a*f^2 + 3*a*d^2
*e^2 + 3*b*d*e*f + 12*a*c*d*e*f)/(3*d^2) + (a*f^2*(3*c^2 - 3))/(3*d^2)) +
log(c + d*x + 1)*((b*f^2*x^3)/6 + (b*e^2*x)/2 + (b*e*f*x^2)/2) + (a*f^2*x^
3)/3 + (log(c + d*x - 1)*((b*f^2)/6 + d*((b*e*f)/2 + (b*c^2*e*f)/2 - b*c*e
*f) + d^2*((b*e^2)/2 - (b*c*e^2)/2) + (b*c^2*f^2)/2 - (b*c^3*f^2)/6 - (b*c
*f^2)/2))/d^3 + (log(c + d*x + 1)*((b*f^2)/6 - d*((b*e*f)/2 + (b*c^2*e*f)/
2 + b*c*e*f) + d^2*((b*e^2)/2 + (b*c*e^2)/2) + (b*c^2*f^2)/2 + (b*c^3*f^2)
/6 + (b*c*f^2)/2))/d^3

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 304, normalized size of antiderivative = 2.53

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \frac{2a \operatorname{atanh}(dx + c) b c^3 f^2 - 6a \operatorname{atanh}(dx + c) b c^2 d e f + 6a \operatorname{atanh}(dx + c) b c^2 f^2 + 6a \operatorname{atanh}(dx + c) b c d^2 e^2 - 12a$$

input `int((f*x+e)^2*(a+b*atanh(d*x+c)),x)`output

```
(2*atanh(c + d*x)*b*c**3*f**2 - 6*atanh(c + d*x)*b*c**2*d*e*f + 6*atanh(c
+ d*x)*b*c**2*f**2 + 6*atanh(c + d*x)*b*c*d**2*e**2 - 12*atanh(c + d*x)*b*
c*d*e*f + 6*atanh(c + d*x)*b*c*f**2 + 6*atanh(c + d*x)*b*d**3*e**2*x + 6*a
tanh(c + d*x)*b*d**3*e*f*x**2 + 2*atanh(c + d*x)*b*d**3*f**2*x**3 + 6*atan
h(c + d*x)*b*d**2*e**2 - 6*atanh(c + d*x)*b*d*e*f + 2*atanh(c + d*x)*b*f**
2 + 6*log(c + d*x - 1)*b*c**2*f**2 - 12*log(c + d*x - 1)*b*c*d*e*f + 6*log
(c + d*x - 1)*b*d**2*e**2 + 2*log(c + d*x - 1)*b*f**2 + 6*a*d**3*e**2*x +
6*a*d**3*e*f*x**2 + 2*a*d**3*f**2*x**3 - 4*b*c*d*f**2*x + 6*b*d**2*e*f*x +
b*d**2*f**2*x**2)/(6*d**3)
```

3.33 $\int (e + fx)(a + b \operatorname{arctanh}(c + dx)) dx$

| | |
|---|-----|
| Optimal result | 317 |
| Mathematica [A] (verified) | 317 |
| Rubi [A] (verified) | 318 |
| Maple [A] (verified) | 320 |
| Fricas [A] (verification not implemented) | 320 |
| Sympy [B] (verification not implemented) | 321 |
| Maxima [A] (verification not implemented) | 321 |
| Giac [B] (verification not implemented) | 322 |
| Mupad [B] (verification not implemented) | 322 |
| Reduce [B] (verification not implemented) | 323 |

Optimal result

Integrand size = 16, antiderivative size = 97

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx)) dx = \frac{bf x}{2d} + \frac{(e + fx)^2(a + b \operatorname{arctanh}(c + dx))}{2f} + \frac{b(de + f - cf)^2 \log(1 - c - dx)}{4d^2 f} - \frac{b(de - (1 + c)f)^2 \log(1 + c + dx)}{4d^2 f}$$

output

```
1/2*b*f*x/d+1/2*(f*x+e)^2*(a+b*arctanh(d*x+c))/f+1/4*b*(-c*f+d*e+f)^2*ln(-d*x-c+1)/d^2/f-1/4*b*(d*e-(1+c)*f)^2*ln(d*x+c+1)/d^2/f
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.42

$$\begin{aligned} & \int (e + fx)(a + b \operatorname{arctanh}(c + dx)) dx \\ &= aex + \frac{bf x}{2d} + \frac{1}{2} a f x^2 + bex \operatorname{arctanh}(c + dx) + \frac{1}{2} b f x^2 \operatorname{arctanh}(c + dx) \\ &+ \frac{b(1 - 2c + c^2) f \log(1 - c - dx)}{4d^2} + \frac{b(-1 - 2c - c^2) f \log(1 + c + dx)}{4d^2} \\ &+ \frac{be(-((-1 + c) \log(1 - c - dx)) + (1 + c) \log(1 + c + dx))}{2d} \end{aligned}$$

input `Integrate[(e + f*x)*(a + b*ArcTanh[c + d*x]),x]`

output `a*e*x + (b*f*x)/(2*d) + (a*f*x^2)/2 + b*e*x*ArcTanh[c + d*x] + (b*f*x^2*ArcTanh[c + d*x])/2 + (b*(1 - 2*c + c^2)*f*Log[1 - c - d*x])/(4*d^2) + (b*(-1 - 2*c - c^2)*f*Log[1 + c + d*x])/(4*d^2) + (b*e*(-((-1 + c)*Log[1 - c - d*x]) + (1 + c)*Log[1 + c + d*x]))/(2*d)`

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.09, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {6661, 27, 6478, 477, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (e + fx)(a + \text{barctanh}(c + dx)) dx \\
 & \quad \downarrow \text{6661} \\
 & \int \frac{(d(e - \frac{cf}{d}) + f(c + dx))(a + \text{barctanh}(c + dx))}{d} d(c + dx) \\
 & \quad \downarrow \text{27} \\
 & \int \frac{(de - cf + f(c + dx))(a + \text{barctanh}(c + dx))d(c + dx)}{d^2} \\
 & \quad \downarrow \text{6478} \\
 & \frac{(f(c + dx) - cf + de)^2(a + \text{barctanh}(c + dx))}{2f} - \frac{b \int \frac{(de - cf + f(c + dx))^2}{1 - (c + dx)^2} d(c + dx)}{2f} \\
 & \quad \downarrow \text{477} \\
 & \frac{(f(c + dx) - cf + de)^2(a + \text{barctanh}(c + dx))}{2f} - \frac{b \int \left(-f^2 + \frac{(de - cf + f)^2}{2(-c - dx + 1)} + \frac{(de - (c + 1)f)^2}{2(c + dx + 1)} \right) d(c + dx)}{2f} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\frac{\frac{(f(c+dx)-cf+de)^2(a+b\operatorname{arctanh}(c+dx))}{2f} - \frac{b(-\frac{1}{2}(-cf+de+f)^2 \log(-c-dx+1) + \frac{1}{2}(de-(c+1)f)^2 \log(c+dx+1) - (f^2(c+dx)))}{2f}}{d^2}$$

input `Int[(e + f*x)*(a + b*ArcTanh[c + d*x]),x]`

output `((((d*e - c*f + f*(c + d*x))^2*(a + b*ArcTanh[c + d*x]))/(2*f) - (b*(-(f^2*(c + d*x)) - ((d*e + f - c*f)^2*Log[1 - c - d*x])/2 + ((d*e - (1 + c)*f)^2*Log[1 + c + d*x])/2))/(2*f))/d^2`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 477 `Int[((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p Int[ExpandIntegrand[(c + d*x)^n*(1 - Rt[-b/a, 2]*x)^p*(1 + Rt[-b/a, 2]*x)^p, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IntegerQ[n] && NiceSqrtQ[-b/a] && !FractionalPowerFactorQ[Rt[-b/a, 2]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6478 `Int[((a_) + ArcTanh[(c_)*(x_)])*(b_))*((d_) + (e_)*(x_))^(q_), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])/(e*(q + 1))), x] - Simp[b*(c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 - c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[q, -1]`

rule 6661 `Int[((a_) + ArcTanh[(c_) + (d_)*(x_)])*(b_))^(p_)*((e_) + (f_)*(x_))^(m_), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.26

| method | result |
|-------------------|---|
| parts | $a\left(\frac{1}{2}fx^2 + ex\right) + \frac{b\left(\frac{\operatorname{arctanh}(dx+c)(dx+c)^2f}{2d} - \frac{\operatorname{arctanh}(dx+c)(dx+c)cf}{d} + \operatorname{arctanh}(dx+c)e(dx+c) - \frac{-f(dx+c) - (-2cf)}{d}\right)}{d}$ |
| derivativedivides | $\frac{a\left(\frac{fc(dx+c) - e(dx+c)d - \frac{f(dx+c)^2}{2}}{d}\right) - b\left(\frac{\operatorname{arctanh}(dx+c)fc(dx+c) - \operatorname{arctanh}(dx+c)e(dx+c)d - \frac{\operatorname{arctanh}(dx+c)f(dx+c)^2}{2} - \frac{f(dx+c)}{2}}{d}\right)}{d}$ |
| default | $\frac{a\left(\frac{fc(dx+c) - e(dx+c)d - \frac{f(dx+c)^2}{2}}{d}\right) - b\left(\frac{\operatorname{arctanh}(dx+c)fc(dx+c) - \operatorname{arctanh}(dx+c)e(dx+c)d - \frac{\operatorname{arctanh}(dx+c)f(dx+c)^2}{2} - \frac{f(dx+c)}{2}}{d}\right)}{d}$ |
| parallelrisc | $-\frac{\operatorname{arctanh}(dx+c)b d^2 f x^2 - a d^2 f x^2 - 2x \operatorname{arctanh}(dx+c)b d^2 e - 2a d^2 e x + \operatorname{arctanh}(dx+c)b c^2 f - 2 \operatorname{arctanh}(dx+c)b c d e}{d^2}$ |
| risc | $\frac{bx(fx+2e)\ln(dx+c+1)}{4} - \frac{bf x^2 \ln(-dx-c+1)}{4} - \frac{bex \ln(-dx-c+1)}{2} + \frac{af x^2}{2} - \frac{\ln(dx+c+1)bc^2 f}{4d^2} + \frac{\ln(dx+c+1)}{2d}$ |

```
input int((f*x+e)*(a+b*arctanh(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output a*(1/2*f*x^2+e*x)+b/d*(1/2/d*arctanh(d*x+c)*(d*x+c)^2*f-1/d*arctanh(d*x+c)
*(d*x+c)*c*f+arctanh(d*x+c)*e*(d*x+c)-1/2/d*(-f*(d*x+c)-1/2*(-2*c*f+2*d*e+
f)*ln(d*x+c-1)+1/2*(2*c*f-2*d*e+f)*ln(d*x+c+1)))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.38

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \frac{2ad^2fx^2 + 2(2ad^2e + bdf)x + (2(bc + b)de - (bc^2 + 2bc + b)f)\log(dx + c + 1) - (2(bc - b)de - (bc^2 - 2bc + b)f)\log(dx + c - 1) + (b*d^2*f*x^2 + 2*b*d^2*e*x)*\log(-(d*x + c + 1)/(d*x + c - 1))}{4d^2}$$

```
input integrate((f*x+e)*(a+b*arctanh(d*x+c)),x, algorithm="fricas")
```

```
output 1/4*(2*a*d^2*f*x^2 + 2*(2*a*d^2*e + b*d*f)*x + (2*(b*c + b)*d*e - (b*c^2 +
2*b*c + b)*f)*log(d*x + c + 1) - (2*(b*c - b)*d*e - (b*c^2 - 2*b*c + b)*f)
*log(d*x + c - 1) + (b*d^2*f*x^2 + 2*b*d^2*e*x)*log(-(d*x + c + 1)/(d*x +
c - 1)))/d^2
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 173 vs. $2(82) = 164$.

Time = 0.89 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.78

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \begin{cases} aex + \frac{afx^2}{2} - \frac{bc^2 f \operatorname{atanh}(c+dx)}{2d^2} + \frac{bce \operatorname{atanh}(c+dx)}{d} - \frac{bcf \log\left(\frac{c}{d} + x + \frac{1}{d}\right)}{d^2} + \frac{bcf \operatorname{atanh}(c+dx)}{d^2} + bex \operatorname{atanh}(c + dx) + \frac{bf}{d} \\ (a + b \operatorname{atanh}(c)) \left(ex + \frac{fx^2}{2} \right) \end{cases}$$

input `integrate((f*x+e)*(a+b*atanh(d*x+c)),x)`

output `Piecewise((a*e*x + a*f*x**2/2 - b*c**2*f*atanh(c + d*x)/(2*d**2) + b*c*e*atanh(c + d*x)/d - b*c*f*log(c/d + x + 1/d)/d**2 + b*c*f*atanh(c + d*x)/d**2 + b*e*x*atanh(c + d*x) + b*f*x**2*atanh(c + d*x)/2 + b*e*log(c/d + x + 1/d)/d - b*e*atanh(c + d*x)/d + b*f*x/(2*d) - b*f*atanh(c + d*x)/(2*d**2), Ne(d, 0)), ((a + b*atanh(c))*(e*x + f*x**2/2), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.12

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx)) dx = \frac{1}{2} a f x^2$$

$$+ \frac{1}{4} \left(2 x^2 \operatorname{artanh}(dx + c) + d \left(\frac{2x}{d^2} - \frac{(c^2 + 2c + 1) \log(dx + c + 1)}{d^3} + \frac{(c^2 - 2c + 1) \log(dx + c - 1)}{d^3} \right) \right)$$

$$+ aex + \frac{(2(dx + c) \operatorname{artanh}(dx + c) + \log(-(dx + c)^2 + 1))be}{2d}$$

input `integrate((f*x+e)*(a+b*arctanh(d*x+c)),x, algorithm="maxima")`

output `1/2*a*f*x^2 + 1/4*(2*x^2*arctanh(d*x + c) + d*(2*x/d^2 - (c^2 + 2*c + 1)*log(d*x + c + 1)/d^3 + (c^2 - 2*c + 1)*log(d*x + c - 1)/d^3))*b*f + a*e*x + 1/2*(2*(d*x + c)*arctanh(d*x + c) + log(-(d*x + c)^2 + 1))*b*e/d`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 341 vs. $2(89) = 178$.

Time = 0.13 (sec) , antiderivative size = 341, normalized size of antiderivative = 3.52

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \frac{1}{2} ((c + 1)d - (c - 1)d) \left(\frac{\left(\frac{(dx+c+1)bde}{dx+c-1} - bde - \frac{(dx+c+1)bcf}{dx+c-1} + bcf + \frac{(dx+c+1)bf}{dx+c-1} \right) \log\left(-\frac{dx+c+1}{dx+c-1}\right) + \frac{2(dx+c+1)}{dx+c-1}}{\frac{(dx+c+1)^2 d^3}{(dx+c-1)^2} - \frac{2(dx+c+1)d^3}{dx+c-1} + d^3} \right)$$

input `integrate((f*x+e)*(a+b*arctanh(d*x+c)),x, algorithm="giac")`

output

```
1/2*((c + 1)*d - (c - 1)*d)*(((d*x + c + 1)*b*d*e/(d*x + c - 1) - b*d*e -
(d*x + c + 1)*b*c*f/(d*x + c - 1) + b*c*f + (d*x + c + 1)*b*f/(d*x + c - 1
))*log(-(d*x + c + 1)/(d*x + c - 1))/((d*x + c + 1)^2*d^3/(d*x + c - 1)^2
- 2*(d*x + c + 1)*d^3/(d*x + c - 1) + d^3) + (2*(d*x + c + 1)*a*d*e/(d*x +
c - 1) - 2*a*d*e - 2*(d*x + c + 1)*a*c*f/(d*x + c - 1) + 2*a*c*f + 2*(d*x
+ c + 1)*a*f/(d*x + c - 1) + (d*x + c + 1)*b*f/(d*x + c - 1) - b*f)/((d*x
+ c + 1)^2*d^3/(d*x + c - 1)^2 - 2*(d*x + c + 1)*d^3/(d*x + c - 1) + d^3)
- (b*d*e - b*c*f)*log(-(d*x + c + 1)/(d*x + c - 1) + 1)/d^3 + (b*d*e - b*
c*f)*log(-(d*x + c + 1)/(d*x + c - 1))/d^3)
```

Mupad [B] (verification not implemented)

Time = 4.74 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.40

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx)) dx = a e x + \frac{a f x^2}{2} + \frac{b e \ln(c^2 + 2 c d x + d^2 x^2 - 1)}{2 d}$$

$$- \frac{b f \operatorname{atanh}(c + dx)}{2 d^2} + \frac{b f x^2 \operatorname{atanh}(c + dx)}{2}$$

$$+ \frac{b f x}{2 d} + b e x \operatorname{atanh}(c + dx)$$

$$- \frac{b c^2 f \operatorname{atanh}(c + dx)}{2 d^2}$$

$$- \frac{b c f \ln(c^2 + 2 c d x + d^2 x^2 - 1)}{2 d^2}$$

$$+ \frac{b c e \operatorname{atanh}(c + dx)}{d}$$

input `int((e + f*x)*(a + b*atanh(c + d*x)),x)`

output `a*e*x + (a*f*x^2)/2 + (b*e*log(c^2 + d^2*x^2 + 2*c*d*x - 1))/(2*d) - (b*f*atanh(c + d*x))/(2*d^2) + (b*f*x^2*atanh(c + d*x))/2 + (b*f*x)/(2*d) + b*e*x*atanh(c + d*x) - (b*c^2*f*atanh(c + d*x))/(2*d^2) - (b*c*f*log(c^2 + d^2*x^2 + 2*c*d*x - 1))/(2*d^2) + (b*c*e*atanh(c + d*x))/d`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.42

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \frac{-\operatorname{atanh}(dx + c) b c^2 f + 2 \operatorname{atanh}(dx + c) bcde - 2 \operatorname{atanh}(dx + c) bcf + 2 \operatorname{atanh}(dx + c) b d^2 ex + \operatorname{atanh}(dx$$

input `int((f*x+e)*(a+b*atanh(d*x+c)),x)`

output `(- atanh(c + d*x)*b*c**2*f + 2*atanh(c + d*x)*b*c*d*e - 2*atanh(c + d*x)*b*c*f + 2*atanh(c + d*x)*b*d**2*e*x + atanh(c + d*x)*b*d**2*f*x**2 + 2*atanh(c + d*x)*b*d*e - atanh(c + d*x)*b*f - 2*log(c + d*x - 1)*b*c*f + 2*log(c + d*x - 1)*b*d*e + 2*a*d**2*e*x + a*d**2*f*x**2 + b*d*f*x)/(2*d**2)`

3.34 $\int (a + b \operatorname{arctanh}(c + dx)) dx$

| | |
|---|-----|
| Optimal result | 324 |
| Mathematica [A] (verified) | 324 |
| Rubi [A] (verified) | 325 |
| Maple [A] (verified) | 326 |
| Fricas [A] (verification not implemented) | 326 |
| Sympy [A] (verification not implemented) | 327 |
| Maxima [A] (verification not implemented) | 327 |
| Giac [B] (verification not implemented) | 327 |
| Mupad [B] (verification not implemented) | 328 |
| Reduce [B] (verification not implemented) | 329 |

Optimal result

Integrand size = 10, antiderivative size = 40

$$\int (a + b \operatorname{arctanh}(c + dx)) dx = ax + \frac{b(c + dx) \operatorname{arctanh}(c + dx)}{d} + \frac{b \log(1 - (c + dx)^2)}{2d}$$

output

```
a*x+b*(d*x+c)*arctanh(d*x+c)/d+1/2*b*ln(1-(d*x+c)^2)/d
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.20

$$\int (a + b \operatorname{arctanh}(c + dx)) dx$$

$$= ax + b \operatorname{arctanh}(c + dx) + \frac{b(-((-1 + c) \log(1 - c - dx)) + (1 + c) \log(1 + c + dx))}{2d}$$

input

```
Integrate[a + b*ArcTanh[c + d*x], x]
```

output

```
a*x + b*x*ArcTanh[c + d*x] + (b*(-((-1 + c)*Log[1 - c - d*x]) + (1 + c)*Log[1 + c + d*x]))/(2*d)
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{arctanh}(c + dx)) dx$$

↓ 2009

$$ax + \frac{b(c + dx)\operatorname{arctanh}(c + dx)}{d} + \frac{b \log(1 - (c + dx)^2)}{2d}$$

input `Int[a + b*ArcTanh[c + d*x],x]`

output `a*x + (b*(c + d*x)*ArcTanh[c + d*x])/d + (b*Log[1 - (c + d*x)^2]/(2*d)`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.92

| method | result |
|-------------------|---|
| default | $ax + \frac{b \left((dx+c) \operatorname{arctanh}(dx+c) + \frac{\ln(1-(dx+c)^2)}{2} \right)}{d}$ |
| parts | $ax + \frac{b \left((dx+c) \operatorname{arctanh}(dx+c) + \frac{\ln(1-(dx+c)^2)}{2} \right)}{d}$ |
| derivativedivides | $\frac{(dx+c)a+b \left((dx+c) \operatorname{arctanh}(dx+c) + \frac{\ln(1-(dx+c)^2)}{2} \right)}{d}$ |
| parallelrisc | $-\frac{b(-\operatorname{arctanh}(dx+c)x d^2 - \operatorname{arctanh}(dx+c)cd - d \ln(dx+c-1) - d \operatorname{arctanh}(dx+c))}{d^2} + ax$ |
| risc | $ax + \frac{b \ln(dx+c+1)x}{2} - \frac{b \ln(-dx-c+1)x}{2} - \frac{b \ln(dx+c-1)c}{2d} + \frac{b \ln(-dx-c-1)c}{2d} + \frac{b \ln(dx+c-1)}{2d} + \frac{b \ln(-dx-c-1)}{2d}$ |

input `int(a+b*arctanh(d*x+c),x,method=_RETURNVERBOSE)`output `a*x+b/d*((d*x+c)*arctanh(d*x+c)+1/2*ln(1-(d*x+c)^2))`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.52

$$\int (a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \frac{bdx \log\left(-\frac{dx+c+1}{dx+c-1}\right) + 2adx + (bc+b) \log(dx+c+1) - (bc-b) \log(dx+c-1)}{2d}$$

input `integrate(a+b*arctanh(d*x+c),x, algorithm="fricas")`output `1/2*(b*d*x*log(-(d*x + c + 1)/(d*x + c - 1)) + 2*a*d*x + (b*c + b)*log(d*x + c + 1) - (b*c - b)*log(d*x + c - 1))/d`

Sympy [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.15

$$\int (a + b \operatorname{arctanh}(c + dx)) dx$$

$$= ax + b \left(\begin{cases} \frac{c \operatorname{atanh}(c+dx)}{d} + x \operatorname{atanh}(c + dx) + \frac{\log(c+dx+1)}{d} - \frac{\operatorname{atanh}(c+dx)}{d} & \text{for } d \neq 0 \\ x \operatorname{atanh}(c) & \text{otherwise} \end{cases} \right)$$

input `integrate(a+b*atanh(d*x+c),x)`

output `a*x + b*Piecewise((c*atanh(c + d*x)/d + x*atanh(c + d*x) + log(c + d*x + 1)/d - atanh(c + d*x)/d, Ne(d, 0)), (x*atanh(c), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.90

$$\int (a + b \operatorname{arctanh}(c + dx)) dx = ax + \frac{(2(dx + c) \operatorname{artanh}(dx + c) + \log(-(dx + c)^2 + 1))b}{2d}$$

input `integrate(a+b*arctanh(d*x+c),x, algorithm="maxima")`

output `a*x + 1/2*(2*(d*x + c)*arctanh(d*x + c) + log(-(d*x + c)^2 + 1))*b/d`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 200 vs. $2(38) = 76$.

Time = 0.12 (sec) , antiderivative size = 200, normalized size of antiderivative = 5.00

$$\int (a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \frac{1}{2} ((c + 1)d - (c - 1)d)b \left(\frac{\log\left(\frac{|-dx - c - 1|}{|dx + c - 1|}\right)}{d^2} - \frac{\log\left(\left| -\frac{dx + c + 1}{dx + c - 1} + 1 \right|\right)}{d^2} + \frac{\log\left(-\frac{c - \frac{((dx + c + 1)(c - 1) - c - 1)d}{dx + c - 1} - d}{\frac{((dx + c + 1)(c - 1) - c - 1)d}{dx + c - 1} - d} + 1 \right)}{d^2 \left(\frac{dx + c + 1}{dx + c - 1} - 1 \right)} \right) + ax$$

input `integrate(a+b*arctanh(d*x+c),x, algorithm="giac")`

output `1/2*((c + 1)*d - (c - 1)*d)*b*(log(abs(-d*x - c - 1)/abs(d*x + c - 1))/d^2 - log(abs(-(d*x + c + 1)/(d*x + c - 1) + 1))/d^2 + log(-(c - ((d*x + c + 1)*(c - 1)/(d*x + c - 1) - c - 1)*d/((d*x + c + 1)*d/(d*x + c - 1) - d) + 1)/(c - ((d*x + c + 1)*(c - 1)/(d*x + c - 1) - c - 1)*d/((d*x + c + 1)*d/(d*x + c - 1) - d) - 1))/d^2*((d*x + c + 1)/(d*x + c - 1) - 1))) + a*x`

Mupad [B] (verification not implemented)

Time = 4.28 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.20

$$\int (a + b \operatorname{arctanh}(c + dx)) dx = ax + \frac{b \ln(c^2 + 2cdx + d^2x^2 - 1)}{2d} + bc \operatorname{atanh}(c + dx) + bx \operatorname{atanh}(c + dx)$$

input `int(a + b*atanh(c + d*x),x)`

output `a*x + ((b*log(c^2 + d^2*x^2 + 2*c*d*x - 1))/2 + b*c*atanh(c + d*x))/d + b*x*atanh(c + d*x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.12

$$\int (a + b \operatorname{arctanh}(c + dx)) dx$$

$$= \frac{\operatorname{atanh}(dx + c)bc + \operatorname{atanh}(dx + c)bdx + \operatorname{atanh}(dx + c)b + \log(dx + c - 1)b + adx}{d}$$

input `int(a+b*atanh(d*x+c),x)`

output `(atanh(c + d*x)*b*c + atanh(c + d*x)*b*d*x + atanh(c + d*x)*b + log(c + d*x - 1)*b + a*d*x)/d`

3.35 $\int \frac{a+b\operatorname{arctanh}(c+dx)}{e+fx} dx$

| | |
|----------------------------|-----|
| Optimal result | 330 |
| Mathematica [A] (verified) | 331 |
| Rubi [A] (verified) | 331 |
| Maple [A] (verified) | 334 |
| Fricas [F] | 334 |
| Sympy [F] | 335 |
| Maxima [F] | 335 |
| Giac [F] | 335 |
| Mupad [F(-1)] | 336 |
| Reduce [F] | 336 |

Optimal result

Integrand size = 18, antiderivative size = 130

$$\int \frac{a + b\operatorname{arctanh}(c + dx)}{e + fx} dx = -\frac{(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{f} + \frac{(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2d(e+fx)}{(de+f-cf)(1+c+dx)}\right)}{f} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{2f} - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d(e+fx)}{(de+f-cf)(1+c+dx)}\right)}{2f}$$

output

```
-(a+b*arctanh(d*x+c))*ln(2/(d*x+c+1))/f+(a+b*arctanh(d*x+c))*ln(2*d*(f*x+e)/(-c*f+d*e+f)/(d*x+c+1))/f+1/2*b*polylog(2,1-2/(d*x+c+1))/f-1/2*b*polylog(2,1-2*d*(f*x+e)/(-c*f+d*e+f)/(d*x+c+1))/f
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.14

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx} dx = \frac{a \log(e + fx)}{f} - \frac{b \log(1 - c - dx) \log\left(\frac{d(e+fx)}{de+f-cf}\right)}{2f}$$

$$+ \frac{b \log(1 + c + dx) \log\left(\frac{d(e+fx)}{de-(1+c)f}\right)}{2f}$$

$$- \frac{b \operatorname{PolyLog}\left(2, \frac{f(1-c-dx)}{de+f-cf}\right)}{2f} + \frac{b \operatorname{PolyLog}\left(2, -\frac{f(1+c+dx)}{de-f-cf}\right)}{2f}$$

input `Integrate[(a + b*ArcTanh[c + d*x])/(e + f*x),x]`

output `(a*Log[e + f*x])/f - (b*Log[1 - c - d*x]*Log[(d*(e + f*x))/(d*e + f - c*f]
)/(2*f) + (b*Log[1 + c + d*x]*Log[(d*(e + f*x))/(d*e - (1 + c)*f]
)/(2*f) - (b*PolyLog[2, (f*(1 - c - d*x))/(d*e + f - c*f]
)/(2*f) + (b*PolyLog[2, -((f*(1 + c + d*x))/(d*e - f - c*f)
)
)/(2*f)`

Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.14, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6661, 27, 6472, 2849, 2752, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx} dx$$

$$\downarrow \text{6661}$$

$$\int \frac{d(a + b \operatorname{arctanh}(c + dx))}{d\left(e - \frac{cf}{d}\right) + f(c + dx)} d(c + dx)$$

$$\hline d$$

$$\downarrow \text{27}$$

$$\begin{aligned}
& \int \frac{a + b \operatorname{arctanh}(c + dx)}{f(c + dx) - cf + de} d(c + dx) \\
& \quad \downarrow 6472 \\
& \frac{b \int \frac{\log\left(\frac{2(de - cf + f(c + dx))}{(de - cf + f)(c + dx + 1)}\right)}{1 - (c + dx)^2} d(c + dx)}{f} + \frac{b \int \frac{\log\left(\frac{2}{c + dx + 1}\right)}{1 - (c + dx)^2} d(c + dx)}{f} + \\
& \frac{(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2(f(c + dx) - cf + de)}{(c + dx + 1)(-cf + de + f)}\right)}{f} - \frac{\log\left(\frac{2}{c + dx + 1}\right) (a + b \operatorname{arctanh}(c + dx))}{f} \\
& \quad \downarrow 2849 \\
& \frac{b \int \frac{\log\left(\frac{2(de - cf + f(c + dx))}{(de - cf + f)(c + dx + 1)}\right)}{1 - (c + dx)^2} d(c + dx)}{f} + \frac{b \int \frac{\log\left(\frac{2}{c + dx + 1}\right)}{1 - \frac{2}{c + dx + 1}} d \frac{1}{c + dx + 1}}{f} + \\
& \frac{(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2(f(c + dx) - cf + de)}{(c + dx + 1)(-cf + de + f)}\right)}{f} - \frac{\log\left(\frac{2}{c + dx + 1}\right) (a + b \operatorname{arctanh}(c + dx))}{f} \\
& \quad \downarrow 2752 \\
& \frac{b \int \frac{\log\left(\frac{2(de - cf + f(c + dx))}{(de - cf + f)(c + dx + 1)}\right)}{1 - (c + dx)^2} d(c + dx)}{f} + \frac{(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2(f(c + dx) - cf + de)}{(c + dx + 1)(-cf + de + f)}\right)}{f} - \\
& \frac{\log\left(\frac{2}{c + dx + 1}\right) (a + b \operatorname{arctanh}(c + dx))}{f} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{c + dx + 1}\right)}{2f} \\
& \quad \downarrow 2897 \\
& \frac{(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2(f(c + dx) - cf + de)}{(c + dx + 1)(-cf + de + f)}\right)}{f} - \frac{\log\left(\frac{2}{c + dx + 1}\right) (a + b \operatorname{arctanh}(c + dx))}{f} - \\
& \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2(de - cf + f(c + dx))}{(de - cf + f)(c + dx + 1)}\right)}{2f} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{c + dx + 1}\right)}{2f}
\end{aligned}$$

input `Int[(a + b*ArcTanh[c + d*x])/(e + f*x),x]`

output `-(((a + b*ArcTanh[c + d*x])*Log[2/(1 + c + d*x)])/f) + ((a + b*ArcTanh[c + d*x])*Log[(2*(d*e - c*f + f*(c + d*x))]/((d*e + f - c*f)*(1 + c + d*x)))]/f + (b*PolyLog[2, 1 - 2/(1 + c + d*x)])/(2*f) - (b*PolyLog[2, 1 - (2*(d*e - c*f + f*(c + d*x))]/((d*e + f - c*f)*(1 + c + d*x)))]/(2*f)`

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 2752 $\text{Int}[\text{Log}[(c_*)(x_)]/((d_) + (e_*)(x_)), x_Symbol] \rightarrow \text{Simp}[(-e^{(-1)})*\text{PolyLog}[2, 1 - c*x], x] /; \text{FreeQ}[\{c, d, e\}, x] \ \&\& \ \text{EqQ}[e + c*d, 0]$
- rule 2849 $\text{Int}[\text{Log}[(c_)/((d_) + (e_*)(x_))]/((f_) + (g_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[-e/g \text{ Subst}[\text{Int}[\text{Log}[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; \text{FreeQ}[\{c, d, e, f, g\}, x] \ \&\& \ \text{EqQ}[c, 2*d] \ \&\& \ \text{EqQ}[e^2*f + d^2*g, 0]$
- rule 2897 $\text{Int}[\text{Log}[u_]*(Pq_)^{(m_.)}, x_Symbol] \rightarrow \text{With}[\{C = \text{FullSimplify}[Pq^m*((1 - u)/D[u, x])]\}, \text{Simp}[C*\text{PolyLog}[2, 1 - u], x] /; \text{FreeQ}[C, x] /; \text{IntegerQ}[m] \ \&\& \ \text{PolyQ}[Pq, x] \ \&\& \ \text{RationalFunctionQ}[u, x] \ \&\& \ \text{LeQ}[\text{RationalFunctionExponents}[u, x][[2]], \text{Expon}[Pq, x]]]$
- rule 6472 $\text{Int}[(a_.) + \text{ArcTanh}[(c_*)(x_)]*(b_.)/((d_) + (e_*)(x_)), x_Symbol] \rightarrow \text{Simp}[(-a + b*\text{ArcTanh}[c*x])*(\text{Log}[2/(1 + c*x)]/e), x] + (\text{Simp}[(a + b*\text{ArcTanh}[c*x])*(\text{Log}[2*c*((d + e*x)/((c*d + e)*(1 + c*x))])]/e), x] + \text{Simp}[b*(c/e) \text{Int}[\text{Log}[2/(1 + c*x)]/(1 - c^2*x^2), x], x] - \text{Simp}[b*(c/e) \text{Int}[\text{Log}[2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/(1 - c^2*x^2), x], x)] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[c^2*d^2 - e^2, 0]$
- rule 6661 $\text{Int}[(a_.) + \text{ArcTanh}[(c_) + (d_*)(x_)]*(b_.))^{(p_.)*((e_.) + (f_*)(x_))^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[1/d \text{ Subst}[\text{Int}[(d*e - c*f)/d + f*(x/d)]^m*(a + b*\text{ArcTanh}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[p, 0]$

Maple [A] (verified)

Time = 6.21 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.48

| method | result |
|-------------------|---|
| parts | $\frac{a \ln(fx+e)}{f} + \frac{b \ln(f(dx+c)-cf+de) \operatorname{arctanh}(dx+c)}{f} + \frac{b \ln(f(dx+c)-cf+de) \ln\left(\frac{f(dx+c)-f}{cf-de-f}\right)}{2f} + \frac{b \operatorname{dilog}\left(\frac{f(dx+c)}{cf-de-f}\right)}{2f}$ |
| derivativedivides | $\frac{ad \ln(cf-de-f(dx+c))}{f} - bd \left(-\frac{\ln(cf-de-f(dx+c)) \operatorname{arctanh}(dx+c)}{f} - \frac{f \left(\operatorname{dilog}\left(\frac{-f(dx+c)-f}{-cf+de-f}\right) + \ln(cf-de-f(dx+c)) \ln\left(\frac{-f(dx+c)}{-cf+de-f}\right) \right)}{2} \right)$ |
| default | $\frac{ad \ln(cf-de-f(dx+c))}{f} - bd \left(-\frac{\ln(cf-de-f(dx+c)) \operatorname{arctanh}(dx+c)}{f} - \frac{f \left(\operatorname{dilog}\left(\frac{-f(dx+c)-f}{-cf+de-f}\right) + \ln(cf-de-f(dx+c)) \ln\left(\frac{-f(dx+c)}{-cf+de-f}\right) \right)}{2} \right)$ |
| risch | $-\frac{b \operatorname{dilog}\left(\frac{(-dx-c+1)f+cf-de-f}{cf-de-f}\right)}{2f} - \frac{b \ln(-dx-c+1) \ln\left(\frac{(-dx-c+1)f+cf-de-f}{cf-de-f}\right)}{2f} + \frac{a \ln((-dx-c+1)f+cf-de-f)}{f}$ |

input `int((a+b*arctanh(d*x+c))/(f*x+e),x,method=_RETURNVERBOSE)`

output `a*ln(f*x+e)/f+b*ln(f*(d*x+c)-c*f+d*e)/f*arctanh(d*x+c)+1/2*b/f*ln(f*(d*x+c)-c*f+d*e)*ln((f*(d*x+c)-f)/(c*f-d*e-f))+1/2*b/f*dilog((f*(d*x+c)-f)/(c*f-d*e-f))-1/2*b/f*ln(f*(d*x+c)-c*f+d*e)*ln((f*(d*x+c)+f)/(c*f-d*e+f))-1/2*b/f*dilog((f*(d*x+c)+f)/(c*f-d*e+f))`

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{fx + e} dx$$

input `integrate((a+b*arctanh(d*x+c))/(f*x+e),x, algorithm="fricas")`

output `integral((b*arctanh(d*x + c) + a)/(f*x + e), x)`

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx} dx = \int \frac{a + b \operatorname{atanh}(c + dx)}{e + fx} dx$$

input `integrate((a+b*atanh(d*x+c))/(f*x+e),x)`

output `Integral((a + b*atanh(c + d*x))/(e + f*x), x)`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{fx + e} dx$$

input `integrate((a+b*arctanh(d*x+c))/(f*x+e),x, algorithm="maxima")`

output `1/2*b*integrate((log(d*x + c + 1) - log(-d*x - c + 1))/(f*x + e), x) + a*log(f*x + e)/f`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{fx + e} dx$$

input `integrate((a+b*arctanh(d*x+c))/(f*x+e),x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)/(f*x + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx} dx = \int \frac{a + b \operatorname{atanh}(c + dx)}{e + fx} dx$$

input `int((a + b*atanh(c + d*x))/(e + f*x),x)`output `int((a + b*atanh(c + d*x))/(e + f*x), x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx} dx = \frac{\left(\int \frac{\operatorname{atanh}(dx+c)}{fx+e} dx \right) bf + \log(fx + e) a}{f}$$

input `int((a+b*atanh(d*x+c))/(f*x+e),x)`output `(int(atanh(c + d*x)/(e + f*x),x)*b*f + log(e + f*x)*a)/f`

3.36 $\int \frac{a+b\operatorname{arctanh}(c+dx)}{(e+fx)^2} dx$

| | |
|---|-----|
| Optimal result | 337 |
| Mathematica [A] (verified) | 337 |
| Rubi [A] (verified) | 338 |
| Maple [A] (verified) | 340 |
| Fricas [B] (verification not implemented) | 340 |
| Sympy [B] (verification not implemented) | 341 |
| Maxima [A] (verification not implemented) | 342 |
| Giac [B] (verification not implemented) | 342 |
| Mupad [B] (verification not implemented) | 343 |
| Reduce [B] (verification not implemented) | 344 |

Optimal result

Integrand size = 18, antiderivative size = 114

$$\int \frac{a + b\operatorname{arctanh}(c + dx)}{(e + fx)^2} dx = -\frac{a + b\operatorname{arctanh}(c + dx)}{f(e + fx)} - \frac{bd \log(1 - c - dx)}{2f(de + f - cf)} + \frac{bd \log(1 + c + dx)}{2f(de - (1 + c)f)} - \frac{bd \log(e + fx)}{(de + f - cf)(de - (1 + c)f)}$$

output

```
-(a+b*arctanh(d*x+c))/f/(f*x+e)-1/2*b*d*ln(-d*x-c+1)/f/(-c*f+d*e+f)+1/2*b*d*ln(d*x+c+1)/f/(d*e-(1+c)*f)-b*d*ln(f*x+e)/(-c*f+d*e+f)/(d*e-(1+c)*f)
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.10

$$\int \frac{a + b\operatorname{arctanh}(c + dx)}{(e + fx)^2} dx = \frac{1}{2} \left(-\frac{2a}{f(e + fx)} - \frac{2b\operatorname{arctanh}(c + dx)}{f(e + fx)} + \frac{bd \log(1 - c - dx)}{f(-de + (-1 + c)f)} - \frac{bd \log(1 + c + dx)}{f(-de + f + cf)} - \frac{2bd \log(e + fx)}{d^2e^2 - 2cdef + (-1 + c^2)f^2} \right)$$

input `Integrate[(a + b*ArcTanh[c + d*x])/(e + f*x)^2,x]`

output
$$\frac{((-2*a)/(f*(e + f*x)) - (2*b*ArcTanh[c + d*x])/(f*(e + f*x)) + (b*d*Log[1 - c - d*x])/(f*(-(d*e) + (-1 + c)*f)) - (b*d*Log[1 + c + d*x])/(f*(-(d*e) + f + c*f)) - (2*b*d*Log[e + f*x])/(d^2*e^2 - 2*c*d*e*f + (-1 + c^2)*f^2))}{2}$$

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.06, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {6659, 2081, 1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + \text{barctanh}(c + dx)}{(e + fx)^2} dx \\ & \quad \downarrow \text{6659} \\ & \frac{bd \int \frac{1}{(e+fx)(1-(c+dx)^2)} dx}{f} - \frac{a + \text{barctanh}(c + dx)}{f(e + fx)} \\ & \quad \downarrow \text{2081} \\ & \frac{bd \int \frac{1}{(e+fx)(-c^2-2dxc-d^2x^2+1)} dx}{f} - \frac{a + \text{barctanh}(c + dx)}{f(e + fx)} \\ & \quad \downarrow \text{1141} \\ & \frac{bd^3 \int \left(\frac{f^2}{d^2(de-cf+f)(de-(c+1)f)(e+fx)} - \frac{1}{2d(de-cf+f)(-c-dx+1)} - \frac{1}{2d(de-cf-f)(c+dx+1)} \right) dx}{f} - \frac{a + \text{barctanh}(c + dx)}{f(e + fx)} \\ & \quad \downarrow \text{2009} \\ & -\frac{a + \text{barctanh}(c + dx)}{f(e + fx)} - \frac{bd^3 \left(\frac{\log(-c-dx+1)}{2d^2(-cf+de+f)} - \frac{\log(c+dx+1)}{2d^2(de-(c+1)f)} + \frac{f \log(e+fx)}{d^2(-cf+de+f)(de-(c+1)f)} \right)}{f} \end{aligned}$$

input `Int[(a + b*ArcTanh[c + d*x])/(e + f*x)^2,x]`

output `-((a + b*ArcTanh[c + d*x])/(f*(e + f*x))) - (b*d^3*(Log[1 - c - d*x]/(2*d^2*(d*e + f - c*f)) - Log[1 + c + d*x]/(2*d^2*(d*e - (1 + c)*f)) + (f*Log[e + f*x])/(d^2*(d*e + f - c*f)*(d*e - (1 + c)*f))))/f`

Defintions of rubi rules used

rule 1141 `Int[((d_.) + (e_.)*(x_)^(m_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x] /; EqQ[p, -1] || !FractionalPowerFactorQ[q]] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p, 0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2081 `Int[(u_)^(m_.)*(v_)^(p_.), x_Symbol] := Int[ExpandToSum[u, x]^m*ExpandToSum[v, x]^p, x] /; FreeQ[{m, p}, x] && LinearQ[u, x] && QuadraticQ[v, x] && !(LinearMatchQ[u, x] && QuadraticMatchQ[v, x])`

rule 6659 `Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)])*(b_.))^(p_.)*((e_.) + (f_.)*(x_)^(m_), x_Symbol] := Simp[(e + f*x)^(m + 1)*((a + b*ArcTanh[c + d*x])^p/(f*(m + 1))), x] - Simp[b*d*(p/(f*(m + 1))) Int[(e + f*x)^(m + 1)*((a + b*ArcTanh[c + d*x])^(p - 1)/(1 - (c + d*x)^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && ILtQ[m, -1]`

Maple [A] (verified)

Time = 0.76 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.20

| method | result |
|-------------------|---|
| parts | $-\frac{a}{(fx+e)f} - \frac{bd \operatorname{arctanh}(dx+c)}{(dfx+de)f} - \frac{bd \ln(f(dx+c)-cf+de)}{(cf-de+f)(cf-de-f)} + \frac{bd \ln(dx+c-1)}{f(2cf-2de-2f)} - \frac{bd \ln(dx+c+1)}{f(2cf-2de+2f)}$ |
| derivativedivides | $\frac{\frac{a d^2}{(cf-de-f(dx+c))f} + b d^2 \left(\frac{\operatorname{arctanh}(dx+c)}{(cf-de-f(dx+c))f} - \frac{\ln(dx+c-1)}{2cf-2de-2f} + \frac{f \ln(cf-de-f(dx+c))}{(cf-de-f)(cf-de+f)} + \frac{\ln(dx+c+1)}{2cf-2de+2f} \right)}{d}$ |
| default | $\frac{\frac{a d^2}{(cf-de-f(dx+c))f} + b d^2 \left(\frac{\operatorname{arctanh}(dx+c)}{(cf-de-f(dx+c))f} - \frac{\ln(dx+c-1)}{2cf-2de-2f} + \frac{f \ln(cf-de-f(dx+c))}{(cf-de-f)(cf-de+f)} + \frac{\ln(dx+c+1)}{2cf-2de+2f} \right)}{d}$ |
| parallelrisch | $-a d^4 e^2 - a c^2 d^2 f^2 + a d^2 f^2 + 2ac d^3 e f + \operatorname{arctanh}(dx+c) b d^3 e f - x \operatorname{arctanh}(dx+c) b c d^3 f^2 + x \operatorname{arctanh}(dx+c) b d^4 e f + \dots$ |
| risch | $-\frac{b \ln(dx+c+1)}{2f(fx+e)} - \frac{\ln(dx+c+1) b c d f^2 x - \ln(dx+c+1) b d^2 e f x - \ln(-dx-c+1) b c d f^2 x + \ln(-dx-c+1) b d^2 e f x + \ln(dx+c-1) b c d f^2 x - \ln(dx+c-1) b d^2 e f x}{2f(fx+e)}$ |

```
input int((a+b*arctanh(d*x+c))/(f*x+e)^2,x,method=_RETURNVERBOSE)
```

```
output -a/(f*x+e)/f-b*d/(d*f*x+d*e)/f*arctanh(d*x+c)-b*d/(c*f-d*e+f)/(c*f-d*e-f)*
ln(f*(d*x+c)-c*f+d*e)+b*d/f/(2*c*f-2*d*e-2*f)*ln(d*x+c-1)-b*d/f/(2*c*f-2*d
*e+2*f)*ln(d*x+c+1)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 263 vs. 2(110) = 220.

Time = 0.20 (sec) , antiderivative size = 263, normalized size of antiderivative = 2.31

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(e + fx)^2} dx = \frac{2 a d^2 e^2 - 4 a c d e f + 2 (a c^2 - a) f^2 - (b d^2 e^2 - (b c - b) d e f + (b d^2 e f - (b c - b) d f^2) x) \log(dx + c + 1)}{2 (d^2 e^3 f - 2 d e^2 f^2)}$$

```
input integrate((a+b*arctanh(d*x+c))/(f*x+e)^2,x, algorithm="fricas")
```

output

```
-1/2*(2*a*d^2*e^2 - 4*a*c*d*e*f + 2*(a*c^2 - a)*f^2 - (b*d^2*e^2 - (b*c -
b)*d*e*f + (b*d^2*e*f - (b*c - b)*d*f^2)*x)*log(d*x + c + 1) + (b*d^2*e^2
- (b*c + b)*d*e*f + (b*d^2*e*f - (b*c + b)*d*f^2)*x)*log(d*x + c - 1) + 2*
(b*d*f^2*x + b*d*e*f)*log(f*x + e) + (b*d^2*e^2 - 2*b*c*d*e*f + (b*c^2 - b
)*f^2)*log(-(d*x + c + 1)/(d*x + c - 1))/(d^2*e^3*f - 2*c*d*e^2*f^2 + (c^
2 - 1)*e*f^3 + (d^2*e^2*f^2 - 2*c*d*e*f^3 + (c^2 - 1)*f^4)*x)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1605 vs. $2(92) = 184$.

Time = 4.57 (sec) , antiderivative size = 1605, normalized size of antiderivative = 14.08

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(e + fx)^2} dx = \text{Too large to display}$$

input

```
integrate((a+b*atanh(d*x+c))/(f*x+e)**2,x)
```

output

```
Piecewise((- (a + b*atanh(c))/(e*f + f**2*x), Eq(d, 0)), ((a*x + b*c*atanh(
c + d*x)/d + b*x*atanh(c + d*x) + b*log(c/d + x + 1/d)/d - b*atanh(c + d*x
)/d)/e**2, Eq(f, 0)), (-2*a*f/(2*e*f**2 + 2*f**3*x) + b*d*e*atanh(d*e/f +
d*x - 1)/(2*e*f**2 + 2*f**3*x) + b*d*f*x*atanh(d*e/f + d*x - 1)/(2*e*f**2
+ 2*f**3*x) - 2*b*f*atanh(d*e/f + d*x - 1)/(2*e*f**2 + 2*f**3*x) - b*f/(2*
e*f**2 + 2*f**3*x), Eq(c, (d*e - f)/f)), (-2*a*f/(2*e*f**2 + 2*f**3*x) - b
*d*e*atanh(d*e/f + d*x + 1)/(2*e*f**2 + 2*f**3*x) - b*d*f*x*atanh(d*e/f +
d*x + 1)/(2*e*f**2 + 2*f**3*x) - 2*b*f*atanh(d*e/f + d*x + 1)/(2*e*f**2 +
2*f**3*x) + b*f/(2*e*f**2 + 2*f**3*x), Eq(c, (d*e + f)/f)), (-a*c**2*f**2/
(c**2*e*f**3 + c**2*f**4*x - 2*c*d*e**2*f**2 - 2*c*d*e*f**3*x + d**2*e**3*
f + d**2*e**2*f**2*x - e*f**3 - f**4*x) + 2*a*c*d*e*f/(c**2*e*f**3 + c**2*
f**4*x - 2*c*d*e**2*f**2 - 2*c*d*e*f**3*x + d**2*e**3*f + d**2*e**2*f**2*x
- e*f**3 - f**4*x) - a*d**2*e**2/(c**2*e*f**3 + c**2*f**4*x - 2*c*d*e**2*
f**2 - 2*c*d*e*f**3*x + d**2*e**3*f + d**2*e**2*f**2*x - e*f**3 - f**4*x)
+ a*f**2/(c**2*e*f**3 + c**2*f**4*x - 2*c*d*e**2*f**2 - 2*c*d*e*f**3*x + d
**2*e**3*f + d**2*e**2*f**2*x - e*f**3 - f**4*x) - b*c**2*f**2*atanh(c + d
*x)/(c**2*e*f**3 + c**2*f**4*x - 2*c*d*e**2*f**2 - 2*c*d*e*f**3*x + d**2*
e**3*f + d**2*e**2*f**2*x - e*f**3 - f**4*x) + b*c*d*e*f*atanh(c + d*x)/(c
**2*e*f**3 + c**2*f**4*x - 2*c*d*e**2*f**2 - 2*c*d*e*f**3*x + d**2*e**3*f +
d**2*e**2*f**2*x - e*f**3 - f**4*x) - b*c*d*f**2*x*atanh(c + d*x)/(c**...
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.06

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(e + fx)^2} dx$$

$$= \frac{1}{2} \left(d \left(\frac{\log(dx + c + 1)}{def - (c + 1)f^2} - \frac{\log(dx + c - 1)}{def - (c - 1)f^2} - \frac{2 \log(fx + e)}{d^2e^2 - 2cdef + (c^2 - 1)f^2} \right) - \frac{2 \operatorname{artanh}(dx + c)}{f^2x + ef} \right) b - \frac{a}{f^2x + ef}$$

input `integrate((a+b*arctanh(d*x+c))/(f*x+e)^2,x, algorithm="maxima")`

output `1/2*(d*(log(d*x + c + 1)/(d*e*f - (c + 1)*f^2) - log(d*x + c - 1)/(d*e*f - (c - 1)*f^2) - 2*log(f*x + e)/(d^2*e^2 - 2*c*d*e*f + (c^2 - 1)*f^2)) - 2*arctanh(d*x + c)/(f^2*x + e*f))*b - a/(f^2*x + e*f)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 474 vs. 2(110) = 220.

Time = 0.13 (sec) , antiderivative size = 474, normalized size of antiderivative = 4.16

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(e + fx)^2} dx =$$

$$-\frac{1}{2} ((c + 1)d - (c - 1)d) \left(\frac{b \log \left(-\frac{(dx+c+1)de}{dx+c-1} + de + \frac{(dx+c+1)cf}{dx+c-1} - cf - \frac{(dx+c+1)f}{dx+c-1} - f \right)}{d^2e^2 - 2cdef + c^2f^2 - f^2} - \frac{(dx+c+1)d^2e^2}{dx+c-1} \right)$$

input `integrate((a+b*arctanh(d*x+c))/(f*x+e)^2,x, algorithm="giac")`

output

```
-1/2*((c + 1)*d - (c - 1)*d)*(b*log(-(d*x + c + 1)*d*e/(d*x + c - 1) + d*e
+ (d*x + c + 1)*c*f/(d*x + c - 1) - c*f - (d*x + c + 1)*f/(d*x + c - 1) -
f)/(d^2*e^2 - 2*c*d*e*f + c^2*f^2 - f^2) - b*log(-(d*x + c + 1)/(d*x + c
- 1))/((d*x + c + 1)*d^2*e^2/(d*x + c - 1) - d^2*e^2 - 2*(d*x + c + 1)*c*d
*e*f/(d*x + c - 1) + 2*c*d*e*f + (d*x + c + 1)*c^2*f^2/(d*x + c - 1) - c^2
*f^2 + 2*(d*x + c + 1)*d*e*f/(d*x + c - 1) - 2*(d*x + c + 1)*c*f^2/(d*x +
c - 1) + (d*x + c + 1)*f^2/(d*x + c - 1) + f^2) - b*log(-(d*x + c + 1)/(d*
x + c - 1))/(d^2*e^2 - 2*c*d*e*f + c^2*f^2 - f^2) - 2*a/((d*x + c + 1)*d^2
*e^2/(d*x + c - 1) - d^2*e^2 - 2*(d*x + c + 1)*c*d*e*f/(d*x + c - 1) + 2*c
*d*e*f + (d*x + c + 1)*c^2*f^2/(d*x + c - 1) - c^2*f^2 + 2*(d*x + c + 1)*d
*e*f/(d*x + c - 1) - 2*(d*x + c + 1)*c*f^2/(d*x + c - 1) + (d*x + c + 1)*f
^2/(d*x + c - 1) + f^2))
```

Mupad [B] (verification not implemented)

Time = 4.51 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.49

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(e + fx)^2} dx = \ln(e + fx) \left(\frac{b(c-1)}{2e(de - f(c-1))} - \frac{b(c+1)}{2e(de - f(c+1))} \right) - \frac{a}{xf^2 + ef} + \frac{b \ln(1 - dx - c)}{f(2e + 2fx)} - \frac{b \ln(c + dx + 1)}{2f(e + fx)} - \frac{bd \ln(c + dx - 1)}{2f^2 - 2cf^2 + 2def} - \frac{bd \ln(c + dx + 1)}{2cf^2 + 2f^2 - 2def}$$

input

```
int((a + b*atanh(c + d*x))/(e + f*x)^2,x)
```

output

```
log(e + f*x)*((b*(c - 1))/(2*e*(d*e - f*(c - 1))) - (b*(c + 1))/(2*e*(d*e
- f*(c + 1)))) - a/(e*f + f^2*x) + (b*log(1 - d*x - c))/(f*(2*e + 2*f*x))
- (b*log(c + d*x + 1))/(2*f*(e + f*x)) - (b*d*log(c + d*x - 1))/(2*f^2 - 2
*c*f^2 + 2*d*e*f) - (b*d*log(c + d*x + 1))/(2*c*f^2 + 2*f^2 - 2*d*e*f)
```


Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 412, normalized size of antiderivative = 3.61

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(e + fx)^2} dx$$

$$= \frac{-\log(dx + c - 1) bcdefx + \log(dx + c + 1) bcdefx + \log(dx + c + 1) bde^2 + \log(dx + c + 1) bef + \log(dx + c + 1) bcf + \log(dx + c + 1) cde}{(e + fx)^2}$$

input `int((a+b*atanh(d*x+c))/(f*x+e)^2,x)`

output

```
(2*atanh(c + d*x)*b*c**2*f**2*x - 4*atanh(c + d*x)*b*c*d*e*f*x + 2*atanh(c + d*x)*b*d**2*e**2*x - 2*atanh(c + d*x)*b*f**2*x + log(c + d*x - 1)*b*c**2*e*f + log(c + d*x - 1)*b*c**2*f**2*x - log(c + d*x - 1)*b*c*d*e**2 - log(c + d*x - 1)*b*c*d*e*f*x + log(c + d*x - 1)*b*d*e**2 + log(c + d*x - 1)*b*d*e*f*x - log(c + d*x - 1)*b*e*f - log(c + d*x - 1)*b*f**2*x - log(c + d*x + 1)*b*c**2*e*f - log(c + d*x + 1)*b*c**2*f**2*x + log(c + d*x + 1)*b*c*d*e**2 + log(c + d*x + 1)*b*c*d*e*f*x + log(c + d*x + 1)*b*d*e**2 + log(c + d*x + 1)*b*d*e*f*x + log(c + d*x + 1)*b*e*f + log(c + d*x + 1)*b*f**2*x - 2*log(e + f*x)*b*d*e**2 - 2*log(e + f*x)*b*d*e*f*x + 2*a*c**2*f**2*x - 4*a*c*d*e*f*x + 2*a*d**2*e**2*x - 2*a*f**2*x)/(2*e*(c**2*e*f**2 + c**2*f**3*x - 2*c*d*e**2*f - 2*c*d*e*f**2*x + d**2*e**3 + d**2*e**2*f*x - e*f**2 - f**3*x))
```

3.37 $\int \frac{a+b\operatorname{arctanh}(c+dx)}{(e+fx)^3} dx$

| | |
|---|-----|
| Optimal result | 345 |
| Mathematica [A] (verified) | 346 |
| Rubi [A] (verified) | 346 |
| Maple [A] (verified) | 348 |
| Fricas [B] (verification not implemented) | 349 |
| Sympy [B] (verification not implemented) | 349 |
| Maxima [A] (verification not implemented) | 350 |
| Giac [B] (verification not implemented) | 351 |
| Mupad [B] (verification not implemented) | 352 |
| Reduce [B] (verification not implemented) | 353 |

Optimal result

Integrand size = 18, antiderivative size = 167

$$\int \frac{a + b\operatorname{arctanh}(c + dx)}{(e + fx)^3} dx = \frac{bd}{2(de + f - cf)(de - (1 + c)f)(e + fx)} - \frac{a + b\operatorname{arctanh}(c + dx)}{2f(e + fx)^2} - \frac{bd^2 \log(1 - c - dx)}{4f(de + f - cf)^2} + \frac{bd^2 \log(1 + c + dx)}{4f(de - f - cf)^2} - \frac{bd^2(de - cf) \log(e + fx)}{(de + f - cf)^2(de - (1 + c)f)^2}$$

output

```
1/2*b*d/(-c*f+d*e+f)/(d*e-(1+c)*f)/(f*x+e)-1/2*(a+b*arctanh(d*x+c))/f/(f*x
+e)^2-1/4*b*d^2*ln(-d*x-c+1)/f/(-c*f+d*e+f)^2+1/4*b*d^2*ln(d*x+c+1)/f/(-c*
f+d*e-f)^2-b*d^2*(-c*f+d*e)*ln(f*x+e)/(-c*f+d*e+f)^2/(d*e-(1+c)*f)^2
```

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.04

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(e + fx)^3} dx = \frac{1}{4} \left(-\frac{2a}{f(e + fx)^2} + \frac{2bd}{(d^2e^2 - 2cdef + (-1 + c^2)f^2)(e + fx)} - \frac{2b \operatorname{arctanh}(c + dx)}{f(e + fx)^2} - \frac{bd^2 \log(1 - c - dx)}{f(de + f - cf)^2} + \frac{bd^2 \log(1 + c + dx)}{f(-de + f + cf)^2} - \frac{4bd^2(de - cf) \log(e + fx)}{(d^2e^2 - 2cdef + (-1 + c^2)f^2)^2} \right)$$

input `Integrate[(a + b*ArcTanh[c + d*x])/(e + f*x)^3,x]`

output `((-2*a)/(f*(e + f*x)^2) + (2*b*d)/((d^2*e^2 - 2*c*d*e*f + (-1 + c^2)*f^2)*(e + f*x)) - (2*b*ArcTanh[c + d*x])/(f*(e + f*x)^2) - (b*d^2*Log[1 - c - d*x])/(f*(d*e + f - c*f)^2) + (b*d^2*Log[1 + c + d*x])/(f*(-(d*e) + f + c*f)^2) - (4*b*d^2*(d*e - c*f)*Log[e + f*x])/(d^2*e^2 - 2*c*d*e*f + (-1 + c^2)*f^2)^2)/4`

Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.02, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {6659, 2081, 1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(e + fx)^3} dx$$

↓ 6659

$$\frac{bd \int \frac{1}{(e+fx)^2(1-(c+dx)^2)} dx}{2f} - \frac{a + b \operatorname{arctanh}(c + dx)}{2f(e + fx)^2}$$

↓ 2081

$$\frac{bd \int \frac{1}{(e+fx)^2(-c^2-2dxc-d^2x^2+1)} dx}{2f} - \frac{a + \operatorname{arctanh}(c + dx)}{2f(e + fx)^2}$$

↓ 1141

$$\frac{bd^3 \int \left(\frac{2(de-cf)f^2}{d(de-cf+f)^2(de-(c+1)f)^2(e+fx)} + \frac{f^2}{d^2(de-cf+f)(de-(c+1)f)(e+fx)^2} - \frac{1}{2(de-cf+f)^2(-c-dx+1)} - \frac{1}{2(de-(c+1)f)^2(c+dx)} \right)}{2f} - \frac{a + \operatorname{arctanh}(c + dx)}{2f(e + fx)^2}$$

↓ 2009

$$\frac{bd^3 \left(-\frac{f}{d^2(e+fx)(-cf+de+f)(de-(c+1)f)} + \frac{2f(de-cf)\log(e+fx)}{d(-cf+de+f)^2(de-(c+1)f)^2} + \frac{\log(-c-dx+1)}{2d(-cf+de+f)^2} - \frac{\log(c+dx+1)}{2d(de-(c+1)f)^2} \right)}{2f} - \frac{a + \operatorname{arctanh}(c + dx)}{2f(e + fx)^2}$$

input `Int[(a + b*ArcTanh[c + d*x])/(e + f*x)^3,x]`

output `-1/2*(a + b*ArcTanh[c + d*x])/(f*(e + f*x)^2) - (b*d^3*(-(f/(d^2*(d*e + f - c*f)*(d*e - (1 + c)*f)*(e + f*x))) + Log[1 - c - d*x]/(2*d*(d*e + f - c*f)^2) - Log[1 + c + d*x]/(2*d*(d*e - (1 + c)*f)^2) + (2*f*(d*e - c*f)*Log[e + f*x])/(d*(d*e + f - c*f)^2*(d*e - (1 + c)*f)^2))/(2*f)`

Defintions of rubi rules used

rule 1141 `Int[((d_.) + (e_.)*(x_))^(m_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x], x] /; EqQ[p, -1] || !FractionalPowerFactorQ[q] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p, 0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 2081 Int[(u_)^(m_.)*(v_)^(p_.), x_Symbol] := Int[ExpandToSum[u, x]^m*ExpandToSum
[v, x]^p, x] /; FreeQ[{m, p}, x] && LinearQ[u, x] && QuadraticQ[v, x] && !
(LinearMatchQ[u, x] && QuadraticMatchQ[v, x])
```

```
rule 6659 Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)])*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_), x_Symbol] := Simp[(e + f*x)^(m + 1)*((a + b*ArcTanh[c + d*x])^p/(f*(m
+ 1))), x] - Simp[b*d*(p/(f*(m + 1))) Int[(e + f*x)^(m + 1)*((a + b*ArcTa
nh[c + d*x])^(p - 1)/(1 - (c + d*x)^2)), x], x] /; FreeQ[{a, b, c, d, e, f}
, x] && IGtQ[p, 0] && ILtQ[m, -1]
```

Maple [A] (verified)

Time = 1.29 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.18

| method | result |
|-------------------|---|
| parts | $-\frac{a}{2(fx+e)^2f} + \frac{b \left(-\frac{d^3 \operatorname{arctanh}(dx+c)}{2(f(dx+c)-cf+de)^2f} + \frac{d^3 \left(\frac{f}{(cf-de+f)(cf-de-f)(f(dx+c)-cf+de)} + \frac{2f(cf-de) \ln(f(dx+c)-cf+de)}{(cf-de+f)^2(cf-de-f)^2} \right)}{d} \right)}{d}$ |
| derivativedivides | $-\frac{a d^3}{2(cf-de-f(dx+c))^2 f} - b d^3 \left(\frac{\operatorname{arctanh}(dx+c)}{2(cf-de-f(dx+c))^2 f} - \frac{\ln(dx+c-1)}{2(cf-de-f)^2} + \frac{\ln(dx+c+1)}{2(cf-de+f)^2} - \frac{f}{(cf-de-f)(cf-de+f)(cf-de-f(dx+c))} \right)$ |
| default | $-\frac{a d^3}{2(cf-de-f(dx+c))^2 f} - b d^3 \left(\frac{\operatorname{arctanh}(dx+c)}{2(cf-de-f(dx+c))^2 f} - \frac{\ln(dx+c-1)}{2(cf-de-f)^2} + \frac{\ln(dx+c+1)}{2(cf-de+f)^2} - \frac{f}{(cf-de-f)(cf-de+f)(cf-de-f(dx+c))} \right)$ |
| parallelrisch | $-\frac{-b d^5 e^3 f^2 + b d^3 e f^4 + a d^6 e^4 f - 2a d^4 e^2 f^3 - 4a c^3 d^3 e f^4 + 6a c^2 d^4 e^2 f^3 - 4ac d^5 e^3 f^2 + 4ac d^3 e f^4 - b c^2 d^3 e f^4 + 2bc d^4 e^3}{d}$ |
| risch | Expression too large to display |

```
input int((a+b*arctanh(d*x+c))/(f*x+e)^3,x,method=_RETURNVERBOSE)
```

```
output -1/2*a/(f*x+e)^2/f+b/d*(-1/2*d^3/(f*(d*x+c)-c*f+d*e)^2/f*arctanh(d*x+c)+1/
2*d^3/f*(f/(c*f-d*e+f)/(c*f-d*e-f)/(f*(d*x+c)-c*f+d*e)+2*f*(c*f-d*e)/(c*f-
d*e+f)^2/(c*f-d*e-f)^2*ln(f*(d*x+c)-c*f+d*e)-1/2/(c*f-d*e-f)^2*ln(d*x+c-1)
+1/2/(c*f-d*e+f)^2*ln(d*x+c+1))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 834 vs. $2(159) = 318$.

Time = 0.69 (sec) , antiderivative size = 834, normalized size of antiderivative = 4.99

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(e + fx)^3} dx = \text{Too large to display}$$

input `integrate((a+b*arctanh(d*x+c))/(f*x+e)^3,x, algorithm="fricas")`

output

```
-1/4*(2*a*d^4*e^4 - 2*(4*a*c + b)*d^3*e^3*f + 4*(3*a*c^2 + b*c - a)*d^2*e^2*f^2 - 2*(4*a*c^3 + b*c^2 - 4*a*c - b)*d*e*f^3 + 2*(a*c^4 - 2*a*c^2 + a)*f^4 - 2*(b*d^3*e^2*f^2 - 2*b*c*d^2*e*f^3 + (b*c^2 - b)*d*f^4)*x - (b*d^4*e^4 - 2*(b*c - b)*d^3*e^3*f + (b*c^2 - 2*b*c + b)*d^2*e^2*f^2 + (b*d^4*e^2*f^2 - 2*(b*c - b)*d^3*e*f^3 + (b*c^2 - 2*b*c + b)*d^2*f^4)*x^2 + 2*(b*d^4*e^3*f - 2*(b*c - b)*d^3*e^2*f^2 + (b*c^2 - 2*b*c + b)*d^2*e*f^3)*x*log(d*x + c + 1) + (b*d^4*e^4 - 2*(b*c + b)*d^3*e^3*f + (b*c^2 + 2*b*c + b)*d^2*e^2*f^2 + (b*d^4*e^2*f^2 - 2*(b*c + b)*d^3*e*f^3 + (b*c^2 + 2*b*c + b)*d^2*f^4)*x^2 + 2*(b*d^4*e^3*f - 2*(b*c + b)*d^3*e^2*f^2 + (b*c^2 + 2*b*c + b)*d^2*e*f^3)*x*log(d*x + c - 1) + 4*(b*d^3*e^3*f - b*c*d^2*e^2*f^2 + (b*d^3*e*f^3 - b*c*d^2*f^4)*x^2 + 2*(b*d^3*e^2*f^2 - b*c*d^2*e*f^3)*x)*log(f*x + e) + (b*d^4*e^4 - 4*b*c*d^3*e^3*f + 2*(3*b*c^2 - b)*d^2*e^2*f^2 - 4*(b*c^3 - b*c)*d*e*f^3 + (b*c^4 - 2*b*c^2 + b)*f^4)*log(-(d*x + c + 1)/(d*x + c - 1)))/(d^4*e^6*f - 4*c*d^3*e^5*f^2 + 2*(3*c^2 - 1)*d^2*e^4*f^3 - 4*(c^3 - c)*d*e^3*f^4 + (c^4 - 2*c^2 + 1)*e^2*f^5 + (d^4*e^4*f^3 - 4*c*d^3*e^3*f^4 + 2*(3*c^2 - 1)*d^2*e^2*f^5 - 4*(c^3 - c)*d*e*f^6 + (c^4 - 2*c^2 + 1)*f^7)*x^2 + 2*(d^4*e^5*f^2 - 4*c*d^3*e^4*f^3 + 2*(3*c^2 - 1)*d^2*e^3*f^4 - 4*(c^3 - c)*d*e^2*f^5 + (c^4 - 2*c^2 + 1)*e*f^6)*x)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19859 vs. $2(141) = 282$.

Time = 11.08 (sec) , antiderivative size = 19859, normalized size of antiderivative = 118.92

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(e + fx)^3} dx = \text{Too large to display}$$

input `integrate((a+b*atanh(d*x+c))/(f*x+e)**3,x)`

output `Piecewise((- (a + b*atanh(c))/(2*e**2*f + 4*e*f**2*x + 2*f**3*x**2), Eq(d, 0)), ((a*x + b*c*atanh(c + d*x)/d + b*x*atanh(c + d*x) + b*log(c/d + x + 1/d)/d - b*atanh(c + d*x)/d)/e**3, Eq(f, 0)), (-4*a*f**2/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2) + b*d**2*e**2*atanh(d*e/f + d*x - 1)/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2) + 2*b*d**2*e*f*x*atanh(d*e/f + d*x - 1)/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2) + b*d**2*f**2*x**2*atanh(d*e/f + d*x - 1)/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2) - b*d*e*f/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2) - b*d*f**2*x/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2) - 4*b*f**2*atanh(d*e/f + d*x - 1)/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2) - b*f**2/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2), Eq(c, (d*e - f)/f)), (-4*a*f**2/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2) + b*d**2*e**2*atanh(d*e/f + d*x + 1)/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2) + 2*b*d**2*e*f*x*atanh(d*e/f + d*x + 1)/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2) + b*d**2*f**2*x**2*atanh(d*e/f + d*x + 1)/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2) - b*d*e*f/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2) - b*d*f**2*x/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2) - 4*b*f**2*atanh(d*e/f + d*x + 1)/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2) + b*f**2/(8*e**2*f**3 + 16*e*f**4*x + 8*f**5*x**2), Eq(c, (d*e + f)/f)), (-a*c**4*f**4/(2*c**4*e**2*f**5 + 4*c**4*e*f**6*x + 2*c**4*f**7*x**2 - 8*c**3*d*e**3*f**4 - 16*c**3*d*e**2*f**5*x - 8*c**3*d*e*f**6*x**2 + 12*c**2*d**2*e**4*f**3 + 24*c**2*d**2...`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.74

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(e + fx)^3} dx$$

$$= \frac{1}{4} \left(d \left(\frac{d \log(dx + c + 1)}{d^2 e^2 f - 2(c + 1) d e f^2 + (c^2 + 2c + 1) f^3} - \frac{d \log(dx + c - 1)}{d^2 e^2 f - 2(c - 1) d e f^2 + (c^2 - 2c + 1) f^3} - \frac{1}{d^4 e^4 - 4} \right) - \frac{1}{2(f^3 x^2 + 2 e f^2 x + e^2 f)} \right)$$

input `integrate((a+b*arctanh(d*x+c))/(f*x+e)^3,x, algorithm="maxima")`

output

```
1/4*(d*(d*log(d*x + c + 1)/(d^2*e^2*f - 2*(c + 1)*d*e*f^2 + (c^2 + 2*c + 1)*f^3) - d*log(d*x + c - 1)/(d^2*e^2*f - 2*(c - 1)*d*e*f^2 + (c^2 - 2*c + 1)*f^3) - 4*(d^2*e - c*d*f)*log(f*x + e)/(d^4*e^4 - 4*c*d^3*e^3*f + 2*(3*c^2 - 1)*d^2*e^2*f^2 - 4*(c^3 - c)*d*e*f^3 + (c^4 - 2*c^2 + 1)*f^4) + 2/(d^2*e^3 - 2*c*d*e^2*f + (c^2 - 1)*e*f^2 + (d^2*e^2*f - 2*c*d*e*f^2 + (c^2 - 1)*f^3)*x)) - 2*arctanh(d*x + c)/(f^3*x^2 + 2*e*f^2*x + e^2*f))*b - 1/2*a/(f^3*x^2 + 2*e*f^2*x + e^2*f)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2567 vs. $2(159) = 318$.

Time = 0.19 (sec) , antiderivative size = 2567, normalized size of antiderivative = 15.37

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(e + fx)^3} dx = \text{Too large to display}$$

input

```
integrate((a+b*arctanh(d*x+c))/(f*x+e)^3,x, algorithm="giac")
```


output

```

-1/2*((c + 1)*d - (c - 1)*d)*((b*d^2*e - b*c*d*f)*log(-(d*x + c + 1)*d*e/(
d*x + c - 1) + d*e + (d*x + c + 1)*c*f/(d*x + c - 1) - c*f - (d*x + c + 1)
*f/(d*x + c - 1) - f)/(d^4*e^4 - 4*c*d^3*e^3*f + 6*c^2*d^2*e^2*f^2 - 4*c^3
*d*e*f^3 + c^4*f^4 - 2*d^2*e^2*f^2 + 4*c*d*e*f^3 - 2*c^2*f^4 + f^4) - ((d*
x + c + 1)*b*d^2*e/(d*x + c - 1) - b*d^2*e - (d*x + c + 1)*b*c*d*f/(d*x +
c - 1) + b*c*d*f + (d*x + c + 1)*b*d*f/(d*x + c - 1))*log(-(d*x + c + 1)/(
d*x + c - 1))/((d*x + c + 1)^2*d^4*e^4/(d*x + c - 1)^2 - 2*(d*x + c + 1)*d
^4*e^4/(d*x + c - 1) + d^4*e^4 - 4*(d*x + c + 1)^2*c*d^3*e^3*f/(d*x + c -
1)^2 + 8*(d*x + c + 1)*c*d^3*e^3*f/(d*x + c - 1) - 4*c*d^3*e^3*f + 6*(d*x
+ c + 1)^2*c^2*d^2*e^2*f^2/(d*x + c - 1)^2 - 12*(d*x + c + 1)*c^2*d^2*e^2*
f^2/(d*x + c - 1) + 6*c^2*d^2*e^2*f^2 - 4*(d*x + c + 1)^2*c^3*d*e*f^3/(d*x
+ c - 1)^2 + 8*(d*x + c + 1)*c^3*d*e*f^3/(d*x + c - 1) - 4*c^3*d*e*f^3 +
(d*x + c + 1)^2*c^4*f^4/(d*x + c - 1)^2 - 2*(d*x + c + 1)*c^4*f^4/(d*x + c
- 1) + c^4*f^4 + 4*(d*x + c + 1)^2*d^3*e^3*f/(d*x + c - 1)^2 - 4*(d*x + c
+ 1)*d^3*e^3*f/(d*x + c - 1) - 12*(d*x + c + 1)^2*c*d^2*e^2*f^2/(d*x + c
- 1)^2 + 12*(d*x + c + 1)*c*d^2*e^2*f^2/(d*x + c - 1) + 12*(d*x + c + 1)^2
*c^2*d*e*f^3/(d*x + c - 1)^2 - 12*(d*x + c + 1)*c^2*d*e*f^3/(d*x + c - 1)
- 4*(d*x + c + 1)^2*c^3*f^4/(d*x + c - 1)^2 + 4*(d*x + c + 1)*c^3*f^4/(d*x
+ c - 1) + 6*(d*x + c + 1)^2*d^2*e^2*f^2/(d*x + c - 1)^2 - 2*d^2*e^2*f^2
- 12*(d*x + c + 1)^2*c*d*e*f^3/(d*x + c - 1)^2 + 4*c*d*e*f^3 + 6*(d*x +...

```

Mupad [B] (verification not implemented)

Time = 5.82 (sec) , antiderivative size = 417, normalized size of antiderivative = 2.50

$$\begin{aligned}
& \int \frac{a + b \operatorname{arctanh}(c + dx)}{(e + fx)^3} dx = \frac{bd^2 \ln(c + dx + 1)}{4c^2 f^3 - 8cde f^2 + 8cf^3 + 4d^2 e^2 f - 8def^2 + 4f^3} \\
& - \frac{\ln(e + fx) (bd^3 e - bcd^2 f)}{c^4 f^4 - 4c^3 def^3 + 6c^2 d^2 e^2 f^2 - 2c^2 f^4 - 4cd^3 e^3 f + 4cde f^3 + d^4 e^4 - 2d^2 e^2 f^2 + f^4} \\
& - \frac{b \ln(c + dx + 1)}{4f(e^2 + 2efx + f^2 x^2)} - \frac{bd^2 \ln(c + dx - 1)}{4c^2 f^3 - 8cde f^2 - 8cf^3 + 4d^2 e^2 f + 8def^2 + 4f^3} \\
& - \frac{-ac^2 f^2 + 2acdef - ad^2 e^2 + bdef + af^2}{-c^2 f^2 + 2cdef - d^2 e^2 + f^2} + \frac{bdf^2 x}{-c^2 f^2 + 2cdef - d^2 e^2 + f^2} \\
& - \frac{2e^2 f + 4ef^2 x + 2f^3 x^2}{2ef(2e^2 + 4efx + 2f^2 x^2)} \\
& + \frac{b \ln(1 - dx - c)}{2f(2e^2 + 4efx + 2f^2 x^2)}
\end{aligned}$$

input

```
int((a + b*atanh(c + d*x))/(e + f*x)^3,x)
```

output

```
(b*d^2*log(c + d*x + 1))/(8*c*f^3 + 4*f^3 + 4*c^2*f^3 + 4*d^2*e^2*f - 8*d*
e*f^2 - 8*c*d*e*f^2) - (log(e + f*x)*(b*d^3*e - b*c*d^2*f))/(f^4 - 2*c^2*f
^4 + c^4*f^4 + d^4*e^4 - 2*d^2*e^2*f^2 + 4*c*d*e*f^3 + 6*c^2*d^2*e^2*f^2 -
4*c*d^3*e^3*f - 4*c^3*d*e*f^3) - (b*log(c + d*x + 1))/(4*f*(e^2 + f^2*x^2
+ 2*e*f*x)) - (b*d^2*log(c + d*x - 1))/(4*f^3 - 8*c*f^3 + 4*c^2*f^3 + 4*d
^2*e^2*f + 8*d*e*f^2 - 8*c*d*e*f^2) - ((a*f^2 - a*c^2*f^2 - a*d^2*e^2 + b*
d*e*f + 2*a*c*d*e*f)/(f^2 - c^2*f^2 - d^2*e^2 + 2*c*d*e*f) + (b*d*f^2*x)/(
f^2 - c^2*f^2 - d^2*e^2 + 2*c*d*e*f))/(2*e^2*f + 2*f^3*x^2 + 4*e*f^2*x) +
(b*log(1 - d*x - c))/(2*f*(2*e^2 + 2*f^2*x^2 + 4*e*f*x))
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 2159, normalized size of antiderivative = 12.93

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{(e + fx)^3} dx = \text{Too large to display}$$

input

```
int((a+b*atanh(d*x+c))/(f*x+e)^3,x)
```

output

```
(4*atanh(c + d*x)*b*c**4*e*f**5*x + 2*atanh(c + d*x)*b*c**4*f**6*x**2 - 16
*atanh(c + d*x)*b*c**3*d*e**2*f**4*x - 8*atanh(c + d*x)*b*c**3*d*e*f**5*x*
*2 + 24*atanh(c + d*x)*b*c**2*d**2*e**3*f**3*x + 12*atanh(c + d*x)*b*c**2*
d**2*e**2*f**4*x**2 - 8*atanh(c + d*x)*b*c**2*e*f**5*x - 4*atanh(c + d*x)*
b*c**2*f**6*x**2 - 16*atanh(c + d*x)*b*c*d**3*e**4*f**2*x - 8*atanh(c + d*
x)*b*c*d**3*e**3*f**3*x**2 + 16*atanh(c + d*x)*b*c*d*e**2*f**4*x + 8*atanh
(c + d*x)*b*c*d*e*f**5*x**2 + 4*atanh(c + d*x)*b*d**4*e**5*f*x + 2*atanh(c
+ d*x)*b*d**4*e**4*f**2*x**2 - 8*atanh(c + d*x)*b*d**2*e**3*f**3*x - 4*at
anh(c + d*x)*b*d**2*e**2*f**4*x**2 + 4*atanh(c + d*x)*b*e*f**5*x + 2*atanh
(c + d*x)*b*f**6*x**2 + log(c + d*x - 1)*b*c**4*e**2*f**4 + 2*log(c + d*x
- 1)*b*c**4*e*f**5*x + log(c + d*x - 1)*b*c**4*f**6*x**2 - 4*log(c + d*x -
1)*b*c**3*d*e**3*f**3 - 8*log(c + d*x - 1)*b*c**3*d*e**2*f**4*x - 4*log(c
+ d*x - 1)*b*c**3*d*e*f**5*x**2 + 5*log(c + d*x - 1)*b*c**2*d**2*e**4*f**
2 + 10*log(c + d*x - 1)*b*c**2*d**2*e**3*f**3*x + 5*log(c + d*x - 1)*b*c**
2*d**2*e**2*f**4*x**2 - 2*log(c + d*x - 1)*b*c**2*e**2*f**4 - 4*log(c + d*
x - 1)*b*c**2*e*f**5*x - 2*log(c + d*x - 1)*b*c**2*f**6*x**2 - 2*log(c + d
*x - 1)*b*c*d**3*e**5*f - 4*log(c + d*x - 1)*b*c*d**3*e**4*f**2*x - 2*log(
c + d*x - 1)*b*c*d**3*e**3*f**3*x**2 - 2*log(c + d*x - 1)*b*c*d**2*e**4*f*
*2 - 4*log(c + d*x - 1)*b*c*d**2*e**3*f**3*x - 2*log(c + d*x - 1)*b*c*d**2
*e**2*f**4*x**2 + 4*log(c + d*x - 1)*b*c*d*e**3*f**3 + 8*log(c + d*x - ...
```

3.38 $\int (e + fx)^3 (a + b \operatorname{arctanh}(c + dx))^2 dx$

| | |
|---|-----|
| Optimal result | 356 |
| Mathematica [A] (warning: unable to verify) | 357 |
| Rubi [A] (verified) | 357 |
| Maple [B] (verified) | 359 |
| Fricas [F] | 360 |
| Sympy [F] | 361 |
| Maxima [B] (verification not implemented) | 361 |
| Giac [F] | 362 |
| Mupad [F(-1)] | 363 |
| Reduce [F] | 363 |

Optimal result

Integrand size = 20, antiderivative size = 562

$$\begin{aligned}
 & \int (e + fx)^3 (a + \operatorname{barctanh}(c + dx))^2 dx \\
 = & \frac{b^2 f^2 (de - cf)x}{d^3} + \frac{abf(6d^2 e^2 - 12cdef + (1 + 6c^2) f^2) x}{2d^3} \\
 & + \frac{b^2 f^3 (c + dx)^2}{12d^4} - \frac{b^2 f^2 (de - cf) \operatorname{arctanh}(c + dx)}{d^4} \\
 & + \frac{b^2 f(6d^2 e^2 - 12cdef + (1 + 6c^2) f^2) (c + dx) \operatorname{arctanh}(c + dx)}{2d^4} \\
 & + \frac{bf^2 (de - cf)(c + dx)^2 (a + \operatorname{barctanh}(c + dx))}{d^4} \\
 & + \frac{bf^3 (c + dx)^3 (a + \operatorname{barctanh}(c + dx))}{6d^4} \\
 & + \frac{(de - cf)(d^2 e^2 - 2cdef + (1 + c^2) f^2) (a + \operatorname{barctanh}(c + dx))^2}{d^4} \\
 & - \frac{(d^4 e^4 - 4cd^3 e^3 f + 6(1 + c^2) d^2 e^2 f^2 - 4c(3 + c^2) def^3 + (1 + 6c^2 + c^4) f^4) (a + \operatorname{barctanh}(c + dx))^2}{4d^4 f} \\
 & + \frac{(e + fx)^4 (a + \operatorname{barctanh}(c + dx))^2}{4f} \\
 & - \frac{2b(de - cf)(d^2 e^2 - 2cdef + (1 + c^2) f^2) (a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1 - c - dx}\right)}{d^4} \\
 & + \frac{b^2 f^3 \log(1 - (c + dx)^2)}{12d^4} + \frac{b^2 f(6d^2 e^2 - 12cdef + (1 + 6c^2) f^2) \log(1 - (c + dx)^2)}{4d^4} \\
 & - \frac{b^2 (de - cf)(d^2 e^2 - 2cdef + (1 + c^2) f^2) \operatorname{PolyLog}\left(2, -\frac{1+c+dx}{1-c-dx}\right)}{d^4}
 \end{aligned}$$

output

```

b^2*f^2*(-c*f+d*e)*x/d^3+1/2*a*b*f*(6*d^2*e^2-12*c*d*e*f+(6*c^2+1)*f^2)*x/
d^3+1/12*b^2*f^3*(d*x+c)^2/d^4-b^2*f^2*(-c*f+d*e)*arctanh(d*x+c)/d^4+1/2*b
^2*f*(6*d^2*e^2-12*c*d*e*f+(6*c^2+1)*f^2)*(d*x+c)*arctanh(d*x+c)/d^4+b*f^2
*(-c*f+d*e)*(d*x+c)^2*(a+b*arctanh(d*x+c))/d^4+1/6*b*f^3*(d*x+c)^3*(a+b*ar
ctanh(d*x+c))/d^4+(-c*f+d*e)*(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)*(a+b*arctanh(
d*x+c))^2/d^4-1/4*(d^4*e^4-4*c*d^3*e^3*f+6*(c^2+1)*d^2*e^2*f^2-4*c*(c^2+3)
*d*e*f^3+(c^4+6*c^2+1)*f^4)*(a+b*arctanh(d*x+c))^2/d^4/f+1/4*(f*x+e)^4*(a+
b*arctanh(d*x+c))^2/f-2*b*(-c*f+d*e)*(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)*(a+b*
arctanh(d*x+c))*ln(2/(-d*x-c+1))/d^4+1/12*b^2*f^3*ln(1-(d*x+c)^2)/d^4+1/4*
b^2*f*(6*d^2*e^2-12*c*d*e*f+(6*c^2+1)*f^2)*ln(1-(d*x+c)^2)/d^4-b^2*(-c*f+d
e)*(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)*polylog(2,-(d*x+c+1)/(-d*x-c+1))/d^4

```

Mathematica [A] (warning: unable to verify)

Time = 4.81 (sec) , antiderivative size = 1082, normalized size of antiderivative = 1.93

$$\int (e + fx)^3 (a + b \operatorname{arctanh}(c + dx))^2 dx = \text{Too large to display}$$

input `Integrate[(e + f*x)^3*(a + b*ArcTanh[c + d*x])^2,x]`

output

```
(12*a^2*e^3*x + 18*a^2*e^2*f*x^2 + 12*a^2*e*f^2*x^3 + 3*a^2*f^3*x^4 + a*b*
(6*x*(4*e^3 + 6*e^2*f*x + 4*e*f^2*x^2 + f^3*x^3)*ArcTanh[c + d*x] - (-2*d*
f*x*(3*(1 + 3*c^2)*f^2 - 3*c*d*f*(8*e + f*x) + d^2*(18*e^2 + 6*e*f*x + f^2
*x^2)) + 3*(-1 + c)*(4*d^3*e^3 - 6*(-1 + c)*d^2*e^2*f + 4*(-1 + c)^2*d*e*f
^2 - (-1 + c)^3*f^3)*Log[1 - c - d*x] + 3*(1 + c)*(-4*d^3*e^3 + 6*(1 + c)*
d^2*e^2*f - 4*(1 + c)^2*d*e*f^2 + (1 + c)^3*f^3)*Log[1 + c + d*x])/d^4 +
(12*b^2*e^3*(ArcTanh[c + d*x]*((-1 + c + d*x)*ArcTanh[c + d*x] - 2*Log[1 +
E^(-2*ArcTanh[c + d*x])])) + PolyLog[2, -E^(-2*ArcTanh[c + d*x])]))/d - (1
8*b^2*e^2*f*((1 - 2*c + c^2 - d^2*x^2)*ArcTanh[c + d*x]^2 - 2*ArcTanh[c +
d*x]*(c + d*x + 2*c*Log[1 + E^(-2*ArcTanh[c + d*x])])) + 2*Log[1/Sqrt[1 - (
c + d*x)^2]] + 2*c*PolyLog[2, -E^(-2*ArcTanh[c + d*x])]))/d^2 + (b^2*f^3*(
-1 - 11*c^2 - 10*c*d*x + d^2*x^2 - 3*(1 - 4*c + 6*c^2 - 4*c^3 + c^4 - d^4*
x^4)*ArcTanh[c + d*x]^2 + 2*ArcTanh[c + d*x]*(9*c + 13*c^3 + 3*d*x + 9*c^2
*d*x - 3*c*d^2*x^2 + d^3*x^3 + 12*(c + c^3)*Log[1 + E^(-2*ArcTanh[c + d*x]
)])) - 8*Log[1/Sqrt[1 - (c + d*x)^2]] - 36*c^2*Log[1/Sqrt[1 - (c + d*x)^2]]
- 12*(c + c^3)*PolyLog[2, -E^(-2*ArcTanh[c + d*x])]))/d^4 - (3*b^2*e*f^2*
(1 - (c + d*x)^2)^(3/2)*(-((c + d*x)/Sqrt[1 - (c + d*x)^2]) + (6*c*(c + d*
x)*ArcTanh[c + d*x])/Sqrt[1 - (c + d*x)^2] + (3*(c + d*x)*ArcTanh[c + d*x]
^2)/Sqrt[1 - (c + d*x)^2] - (3*c^2*(c + d*x)*ArcTanh[c + d*x]^2)/Sqrt[1 -
(c + d*x)^2] + ArcTanh[c + d*x]^2*Cosh[3*ArcTanh[c + d*x]] + 3*c^2*ArcT...
```

Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 547, normalized size of antiderivative = 0.97, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {6661, 27, 6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int (e + fx)^3 (a + b \operatorname{arctanh}(c + dx))^2 dx \\
& \quad \downarrow \text{6661} \\
& \int \frac{\left(d\left(e - \frac{cf}{d}\right) + f(c + dx)\right)^3 (a + b \operatorname{arctanh}(c + dx))^2}{d^3} d(c + dx) \\
& \quad \downarrow \text{27} \\
& \frac{\int (de - cf + f(c + dx))^3 (a + b \operatorname{arctanh}(c + dx))^2 d(c + dx)}{d^4} \\
& \quad \downarrow \text{6480} \\
& \frac{\frac{(f(c + dx) - cf + de)^4 (a + b \operatorname{arctanh}(c + dx))^2}{4f} - b \int \left(-(c + dx)^2 (a + b \operatorname{arctanh}(c + dx))^4 - 4(de - cf)(c + dx)(a + b \operatorname{arctanh}(c + dx))^3 - (6d^2 e^2 - \dots) \right)}{4f}}{4f} \\
& \quad \downarrow \text{2009} \\
& \frac{\frac{(f(c + dx) - cf + de)^4 (a + b \operatorname{arctanh}(c + dx))^2}{4f} - b \left(\frac{-2f(de - cf)((c^2 + 1)f^2 - 2cdef + d^2 e^2)}{b} (a + b \operatorname{arctanh}(c + dx))^2 + 4f(de - cf)((c^2 + 1)f^2 - 2cdef + \dots) \right)}{4f}}{4f}
\end{aligned}$$

input `Int[(e + f*x)^3*(a + b*ArcTanh[c + d*x])^2,x]`

output

```

((((d*e - c*f + f*(c + d*x))^4*(a + b*ArcTanh[c + d*x])^2)/(4*f) - (b*(-2*b*f^3*(d*e - c*f)*(c + d*x) - a*f^2*(6*d^2*e^2 - 12*c*d*e*f + (1 + 6*c^2)*f^2)*(c + d*x) - (b*f^4*(c + d*x)^2)/6 + 2*b*f^3*(d*e - c*f)*ArcTanh[c + d*x] - b*f^2*(6*d^2*e^2 - 12*c*d*e*f + (1 + 6*c^2)*f^2)*(c + d*x)*ArcTanh[c + d*x] - 2*f^3*(d*e - c*f)*(c + d*x)^2*(a + b*ArcTanh[c + d*x]) - (f^4*(c + d*x)^3*(a + b*ArcTanh[c + d*x]))/3 - (2*f*(d*e - c*f)*(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2)*(a + b*ArcTanh[c + d*x])^2)/b + ((d^4*e^4 - 4*c*d^3*e^3*f + 6*(1 + c^2)*d^2*e^2*f^2 - 4*c*(3 + c^2)*d*e*f^3 + (1 + 6*c^2 + c^4)*f^4)*(a + b*ArcTanh[c + d*x])^2)/(2*b) + 4*f*(d*e - c*f)*(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2)*(a + b*ArcTanh[c + d*x])*Log[2/(1 - c - d*x)] - (b*f^4*Log[1 - (c + d*x)^2])/6 - (b*f^2*(6*d^2*e^2 - 12*c*d*e*f + (1 + 6*c^2)*f^2)*Log[1 - (c + d*x)^2])/2 + 2*b*f*(d*e - c*f)*(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2)*PolyLog[2, -((1 + c + d*x)/(1 - c - d*x))]))/(2*f))/d^4

```

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6480 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*((d_) + (e_.)*(x_))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]`

rule 6661 `Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[p, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2317 vs. 2(546) = 1092.

Time = 1.19 (sec) , antiderivative size = 2318, normalized size of antiderivative = 4.12

| method | result | size |
|--------------------|---------------------------------|------|
| derivativeldivides | Expression too large to display | 2318 |
| default | Expression too large to display | 2318 |
| parts | Expression too large to display | 2339 |
| risch | Expression too large to display | 3412 |

input `int((f*x+e)^3*(a+b*arctanh(d*x+c))^2,x,method=_RETURNVERBOSE)`

output

```

1/d*(1/4*a^2/d^3*(c*f-d*e-f*(d*x+c))^4/f-b^2/d^3*(-1/4*f^3*arctanh(d*x+c)^
2*c^4+f^2*arctanh(d*x+c)^2*c^3*d*e+f^3*arctanh(d*x+c)^2*c^3*(d*x+c)-3/2*f*
arctanh(d*x+c)^2*c^2*d^2*e^2-3*f^2*arctanh(d*x+c)^2*c^2*d*e*(d*x+c)-3/2*f^
3*arctanh(d*x+c)^2*c^2*(d*x+c)^2+arctanh(d*x+c)^2*c*d^3*e^3+3*f*arctanh(d*
x+c)^2*c*d^2*e^2*(d*x+c)+3*f^2*arctanh(d*x+c)^2*c*d*e*(d*x+c)^2+f^3*arctan
h(d*x+c)^2*c*(d*x+c)^3-1/4/f*arctanh(d*x+c)^2*d^4*e^4-arctanh(d*x+c)^2*d^3
*e^3*(d*x+c)-3/2*f*arctanh(d*x+c)^2*d^2*e^2*(d*x+c)^2-f^2*arctanh(d*x+c)^2
*d*e*(d*x+c)^3-1/4*f^3*arctanh(d*x+c)^2*(d*x+c)^4+1/2/f*(1/2*arctanh(d*x+c
))*ln(d*x+c+1)*f^4-1/2*arctanh(d*x+c)*ln(d*x+c-1)*f^4-arctanh(d*x+c)*f^4*(d
*x+c)-1/3*arctanh(d*x+c)*f^4*(d*x+c)^3+2*arctanh(d*x+c)*ln(d*x+c-1)*c^3*d*
e*f^3-3*arctanh(d*x+c)*ln(d*x+c-1)*c^2*d^2*e^2*f^2+2*arctanh(d*x+c)*ln(d*x
+c-1)*c*d^3*e^3*f-6*arctanh(d*x+c)*ln(d*x+c-1)*c^2*d*e*f^3+6*arctanh(d*x+c
)*ln(d*x+c-1)*c*d^2*e^2*f^2+6*arctanh(d*x+c)*ln(d*x+c-1)*c*d*e*f^3-2*arcta
nh(d*x+c)*ln(d*x+c+1)*c^3*d*e*f^3+3*arctanh(d*x+c)*ln(d*x+c+1)*c^2*d^2*e^2
*f^2-2*arctanh(d*x+c)*ln(d*x+c+1)*c*d^3*e^3*f-6*arctanh(d*x+c)*ln(d*x+c+1)
*c^2*d*e*f^3+6*arctanh(d*x+c)*ln(d*x+c+1)*c*d^2*e^2*f^2-6*arctanh(d*x+c)*l
n(d*x+c+1)*c*d*e*f^3+12*arctanh(d*x+c)*c*d*e*f^3*(d*x+c)-2*arctanh(d*x+c)*
ln(d*x+c-1)*d*e*f^3-6*arctanh(d*x+c)*d^2*e^2*f^2*(d*x+c)-2*arctanh(d*x+c)*
d*e*f^3*(d*x+c)^2-2*arctanh(d*x+c)*ln(d*x+c+1)*d^3*e^3*f+3*arctanh(d*x+c)*
ln(d*x+c+1)*d^2*e^2*f^2-2*arctanh(d*x+c)*ln(d*x+c+1)*d*e*f^3-2*arctanh(...

```

Fricas [F]

$$\int (e + fx)^3 (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (fx + e)^3 (b \operatorname{arctanh}(dx + c) + a)^2 dx$$

input

```
integrate((f*x+e)^3*(a+b*arctanh(d*x+c))^2,x, algorithm="fricas")
```

output

```

integral(a^2*f^3*x^3 + 3*a^2*e*f^2*x^2 + 3*a^2*e^2*f*x + a^2*e^3 + (b^2*f^
3*x^3 + 3*b^2*e*f^2*x^2 + 3*b^2*e^2*f*x + b^2*e^3)*arctanh(d*x + c)^2 + 2*
(a*b*f^3*x^3 + 3*a*b*e*f^2*x^2 + 3*a*b*e^2*f*x + a*b*e^3)*arctanh(d*x + c)
, x)

```

Sympy [F]

$$\int (e + fx)^3 (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (a + b \operatorname{atanh}(c + dx))^2 (e + fx)^3 dx$$

input `integrate((f*x+e)**3*(a+b*atanh(d*x+c))**2,x)`

output `Integral((a + b*atanh(c + d*x))**2*(e + f*x)**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1363 vs. 2(538) = 1076.

Time = 0.25 (sec) , antiderivative size = 1363, normalized size of antiderivative = 2.43

$$\int (e + fx)^3 (a + b \operatorname{arctanh}(c + dx))^2 dx = \text{Too large to display}$$

input `integrate((f*x+e)^3*(a+b*arctanh(d*x+c))^2,x, algorithm="maxima")`

output

```

1/4*a^2*f^3*x^4 + a^2*e*f^2*x^3 + 3/2*a^2*e^2*f*x^2 + 3/2*(2*x^2*arctanh(d
*x + c) + d*(2*x/d^2 - (c^2 + 2*c + 1)*log(d*x + c + 1)/d^3 + (c^2 - 2*c +
1)*log(d*x + c - 1)/d^3))*a*b*e^2*f + (2*x^3*arctanh(d*x + c) + d*((d*x^2
- 4*c*x)/d^3 + (c^3 + 3*c^2 + 3*c + 1)*log(d*x + c + 1)/d^4 - (c^3 - 3*c^
2 + 3*c - 1)*log(d*x + c - 1)/d^4))*a*b*e*f^2 + 1/12*(6*x^4*arctanh(d*x +
c) + d*(2*(d^2*x^3 - 3*c*d*x^2 + 3*(3*c^2 + 1)*x)/d^4 - 3*(c^4 + 4*c^3 + 6
*c^2 + 4*c + 1)*log(d*x + c + 1)/d^5 + 3*(c^4 - 4*c^3 + 6*c^2 - 4*c + 1)*l
og(d*x + c - 1)/d^5))*a*b*f^3 + a^2*e^3*x + (2*(d*x + c)*arctanh(d*x + c)
+ log(-(d*x + c)^2 + 1))*a*b*e^3/d + (d^3*e^3 + 3*c^2*d*e*f^2 - c^3*f^3 +
d*e*f^2 - (3*d^2*e^2*f + f^3)*c)*(log(d*x + c + 1)*log(-1/2*d*x - 1/2*c +
1/2) + dilog(1/2*d*x + 1/2*c + 1/2))*b^2/d^4 + 1/12*(13*c^3*f^3 + 18*d^2*e
^2*f - 6*d*e*f^2 - 6*(5*d*e*f^2 - 3*f^3)*c^2 + 4*f^3 + 9*(2*d^2*e^2*f - 4*
d*e*f^2 + f^3)*c)*b^2*log(d*x + c + 1)/d^4 - 1/12*(13*c^3*f^3 - 18*d^2*e^
2*f - 6*d*e*f^2 - 6*(5*d*e*f^2 + 3*f^3)*c^2 - 4*f^3 + 9*(2*d^2*e^2*f + 4*d*
e*f^2 + f^3)*c)*b^2*log(d*x + c - 1)/d^4 + 1/48*(4*b^2*d^2*f^3*x^2 + 8*(6*
d^2*e*f^2 - 5*c*d*f^3)*b^2*x + 3*(b^2*d^4*f^3*x^4 + 4*b^2*d^4*e*f^2*x^3 +
6*b^2*d^4*e^2*f*x^2 + 4*b^2*d^4*e^3*x - (c^4*f^3 - 4*d^3*e^3 + 6*d^2*e^2*f
- 4*(d*e*f^2 - f^3)*c^3 - 4*d*e*f^2 + 6*(d^2*e^2*f - 2*d*e*f^2 + f^3)*c^2
+ f^3 - 4*(d^3*e^3 - 3*d^2*e^2*f + 3*d*e*f^2 - f^3)*c)*b^2)*log(d*x + c +
1)^2 + 3*(b^2*d^4*f^3*x^4 + 4*b^2*d^4*e*f^2*x^3 + 6*b^2*d^4*e^2*f*x^2 ...

```

Giac [F]

$$\int (e + fx)^3 (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (fx + e)^3 (b \operatorname{arctanh}(dx + c) + a)^2 dx$$

input

```
integrate((f*x+e)^3*(a+b*arctanh(d*x+c))^2,x, algorithm="giac")
```

output

```
integrate((f*x + e)^3*(b*arctanh(d*x + c) + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int (e + fx)^3 (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (e + fx)^3 (a + b \operatorname{atanh}(c + dx))^2 dx$$

input `int((e + f*x)^3*(a + b*atanh(c + d*x))^2,x)`output `int((e + f*x)^3*(a + b*atanh(c + d*x))^2, x)`**Reduce [F]**

$$\int (e + fx)^3 (a + b \operatorname{arctanh}(c + dx))^2 dx = \text{Too large to display}$$

input `int((f*x+e)^3*(a+b*atanh(d*x+c))^2,x)`

output

```
(9*atanh(c + d*x)**2*b**2*c**4*f**3 - 24*atanh(c + d*x)**2*b**2*c**3*d*e*f
**2 + 18*atanh(c + d*x)**2*b**2*c**2*d**2*e**2*f - 6*atanh(c + d*x)**2*b**
2*c**2*f**3 + 24*atanh(c + d*x)**2*b**2*c*d*e*f**2 + 12*atanh(c + d*x)**2*
b**2*d**4*e**3*x + 18*atanh(c + d*x)**2*b**2*d**4*e**2*f*x**2 + 12*atanh(c
+ d*x)**2*b**2*d**4*e*f**2*x**3 + 3*atanh(c + d*x)**2*b**2*d**4*f**3*x**4
- 18*atanh(c + d*x)**2*b**2*d**2*e**2*f - 3*atanh(c + d*x)**2*b**2*f**3 -
6*atanh(c + d*x)*a*b*c**4*f**3 + 24*atanh(c + d*x)*a*b*c**3*d*e*f**2 - 24
*atanh(c + d*x)*a*b*c**3*f**3 - 36*atanh(c + d*x)*a*b*c**2*d**2*e**2*f + 7
2*atanh(c + d*x)*a*b*c**2*d*e*f**2 - 36*atanh(c + d*x)*a*b*c**2*f**3 + 24*
atanh(c + d*x)*a*b*c*d**3*e**3 - 72*atanh(c + d*x)*a*b*c*d**2*e**2*f + 72*
atanh(c + d*x)*a*b*c*d*e*f**2 - 24*atanh(c + d*x)*a*b*c*f**3 + 24*atanh(c
+ d*x)*a*b*d**4*e**3*x + 36*atanh(c + d*x)*a*b*d**4*e**2*f*x**2 + 24*atanh
(c + d*x)*a*b*d**4*e*f**2*x**3 + 6*atanh(c + d*x)*a*b*d**4*f**3*x**4 + 24*
atanh(c + d*x)*a*b*d**3*e**3 - 36*atanh(c + d*x)*a*b*d**2*e**2*f + 24*atan
h(c + d*x)*a*b*d*e*f**2 - 6*atanh(c + d*x)*a*b*f**3 + 26*atanh(c + d*x)*b*
*2*c**3*f**3 - 60*atanh(c + d*x)*b**2*c**2*d*e*f**2 + 18*atanh(c + d*x)*b*
*2*c**2*d*f**3*x + 36*atanh(c + d*x)*b**2*c**2*f**3 + 36*atanh(c + d*x)*b*
*2*c*d**2*e**2*f - 48*atanh(c + d*x)*b**2*c*d**2*e*f**2*x - 6*atanh(c + d*
x)*b**2*c*d**2*f**3*x**2 - 72*atanh(c + d*x)*b**2*c*d*e*f**2 + 18*atanh(c
+ d*x)*b**2*c*f**3 + 36*atanh(c + d*x)*b**2*d**3*e**2*f*x + 12*atanh(c ...
```

3.39 $\int (e + fx)^2 (a + \operatorname{barctanh}(c + dx))^2 dx$

| | |
|---|-----|
| Optimal result | 365 |
| Mathematica [B] (warning: unable to verify) | 366 |
| Rubi [A] (verified) | 367 |
| Maple [B] (verified) | 369 |
| Fricas [F] | 370 |
| Sympy [F] | 371 |
| Maxima [B] (verification not implemented) | 371 |
| Giac [F] | 372 |
| Mupad [F(-1)] | 373 |
| Reduce [F] | 373 |

Optimal result

Integrand size = 20, antiderivative size = 374

$$\begin{aligned}
 & \int (e + fx)^2 (a + \operatorname{barctanh}(c + dx))^2 dx \\
 &= \frac{b^2 f^2 x}{3d^2} + \frac{2abf(de - cf)x}{d^2} - \frac{b^2 f^2 \operatorname{arctanh}(c + dx)}{3d^3} \\
 &+ \frac{2b^2 f(de - cf)(c + dx) \operatorname{arctanh}(c + dx)}{d^3} + \frac{bf^2(c + dx)^2 (a + \operatorname{barctanh}(c + dx))}{3d^3} \\
 &- \frac{(de - cf)(d^2 e^2 - 2cdef + (3 + c^2)f^2)(a + \operatorname{barctanh}(c + dx))^2}{3d^3 f} \\
 &+ \frac{(3d^2 e^2 - 6cdef + (1 + 3c^2)f^2)(a + \operatorname{barctanh}(c + dx))^2}{3d^3} \\
 &+ \frac{(e + fx)^3 (a + \operatorname{barctanh}(c + dx))^2}{3f} \\
 &- \frac{2b(3d^2 e^2 - 6cdef + (1 + 3c^2)f^2)(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1 - c - dx}\right)}{3d^3} \\
 &+ \frac{b^2 f(de - cf) \log(1 - (c + dx)^2)}{d^3} \\
 &- \frac{b^2(3d^2 e^2 - 6cdef + (1 + 3c^2)f^2) \operatorname{PolyLog}\left(2, -\frac{1 + c + dx}{1 - c - dx}\right)}{3d^3}
 \end{aligned}$$

output

```
1/3*b^2*f^2*x/d^2+2*a*b*f*(-c*f+d*e)*x/d^2-1/3*b^2*f^2*arctanh(d*x+c)/d^3+
2*b^2*f*(-c*f+d*e)*(d*x+c)*arctanh(d*x+c)/d^3+1/3*b*f^2*(d*x+c)^2*(a+b*arc
tanh(d*x+c))/d^3-1/3*(-c*f+d*e)*(d^2*e^2-2*c*d*e*f+(c^2+3)*f^2)*(a+b*arcta
nhd(x+c))^2/d^3/f+1/3*(3*d^2*e^2-6*c*d*e*f+(3*c^2+1)*f^2)*(a+b*arctanh(d*x
+c))^2/d^3+1/3*(f*x+e)^3*(a+b*arctanh(d*x+c))^2/f-2/3*b*(3*d^2*e^2-6*c*d*
e*f+(3*c^2+1)*f^2)*(a+b*arctanh(d*x+c))*ln(2/(-d*x-c+1))/d^3+b^2*f*(-c*f+d
*e)*ln(1-(d*x+c)^2)/d^3-1/3*b^2*(3*d^2*e^2-6*c*d*e*f+(3*c^2+1)*f^2)*polylo
g(2,-(d*x+c+1)/(-d*x-c+1))/d^3
```

Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 795 vs. $2(374) = 748$.

Time = 2.25 (sec) , antiderivative size = 795, normalized size of antiderivative = 2.13

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx = \text{Too large to display}$$

input

```
Integrate[(e + f*x)^2*(a + b*ArcTanh[c + d*x])^2,x]
```

output

```

a^2*e^2*x + a^2*e*f*x^2 + (a^2*f^2*x^3)/3 + (a*b*(2*x*(3*e^2 + 3*e*f*x + f
^2*x^2)*ArcTanh[c + d*x] + (d*f*x*(6*d*e - 4*c*f + d*f*x) - (-1 + c)*(3*d^
2*e^2 - 3*(-1 + c)*d*e*f + (-1 + c)^2*f^2)*Log[1 - c - d*x] + (1 + c)*(3*d
^2*e^2 - 3*(1 + c)*d*e*f + (1 + c)^2*f^2)*Log[1 + c + d*x])/d^3 + (b^2
*e^2*(ArcTanh[c + d*x]*((-1 + c + d*x)*ArcTanh[c + d*x] - 2*Log[1 + E^(-2*
ArcTanh[c + d*x])) + PolyLog[2, -E^(-2*ArcTanh[c + d*x])]))/d + (b^2*e*f*
((-1 + 2*c - c^2 + d^2*x^2)*ArcTanh[c + d*x]^2 + 2*ArcTanh[c + d*x]*(c + d
*x + 2*c*Log[1 + E^(-2*ArcTanh[c + d*x])]) - 2*Log[1/Sqrt[1 - (c + d*x)^2]
] - 2*c*PolyLog[2, -E^(-2*ArcTanh[c + d*x])]))/d^2 - (b^2*f^2*(1 - (c + d*
x)^2)^(3/2)*(-(c + d*x)/Sqrt[1 - (c + d*x)^2]) + (6*c*(c + d*x)*ArcTanh[c
+ d*x])/Sqrt[1 - (c + d*x)^2] + (3*(c + d*x)*ArcTanh[c + d*x]^2)/Sqrt[1 -
(c + d*x)^2] - (3*c^2*(c + d*x)*ArcTanh[c + d*x]^2)/Sqrt[1 - (c + d*x)^2]
+ ArcTanh[c + d*x]^2*Cosh[3*ArcTanh[c + d*x]] + 3*c^2*ArcTanh[c + d*x]^2*
Cosh[3*ArcTanh[c + d*x]] + 2*ArcTanh[c + d*x]*Cosh[3*ArcTanh[c + d*x]]*Log
[1 + E^(-2*ArcTanh[c + d*x])] + 6*c^2*ArcTanh[c + d*x]*Cosh[3*ArcTanh[c +
d*x]]*Log[1 + E^(-2*ArcTanh[c + d*x])] - 6*c*Cosh[3*ArcTanh[c + d*x]]*Log[
1/Sqrt[1 - (c + d*x)^2]] + (3*(1 - 4*c + 3*c^2)*ArcTanh[c + d*x]^2 + 2*Arc
Tanh[c + d*x]*(2 + (3 + 9*c^2)*Log[1 + E^(-2*ArcTanh[c + d*x])]) - 18*c*Lo
g[1/Sqrt[1 - (c + d*x)^2]])/Sqrt[1 - (c + d*x)^2] - (4*(1 + 3*c^2)*PolyLog
[2, -E^(-2*ArcTanh[c + d*x])])/(1 - (c + d*x)^2)^(3/2) - Sinh[3*ArcTanh...

```

Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 374, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {6661, 27, 6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)^2 (a + \text{barctanh}(c + dx))^2 dx$$

$$\downarrow 6661$$

$$\int \frac{\left(\frac{d(e - \frac{ef}{d}) + f(c + dx)}{d^2}\right)^2 (a + \text{barctanh}(c + dx))^2}{d} d(c + dx)$$

$$\downarrow 27$$

$$\frac{\int (de - cf + f(c + dx))^2 (a + \operatorname{arctanh}(c + dx))^2 d(c + dx)}{d^3}$$

↓ 6480

$$\frac{\frac{(f(c+dx)-cf+de)^3(a+b\operatorname{arctanh}(c+dx))^2}{3f} - \frac{2b \int \left(-((c+dx)(a+b\operatorname{arctanh}(c+dx))f^3) - 3(de-cf)(a+b\operatorname{arctanh}(c+dx))f^2 + \frac{(de-cf)(d^2)}{3f} \right)}{d^3}}{d^3}$$

↓ 2009

$$\frac{\frac{(f(c+dx)-cf+de)^3(a+b\operatorname{arctanh}(c+dx))^2}{3f} - \frac{2b \left(-\frac{f((3c^2+1)f^2-6cdef+3d^2e^2)(a+b\operatorname{arctanh}(c+dx))^2}{2b} + \frac{(de-cf)((c^2+3)f^2-2cdef+d^2e^2)(a+b)}{2b} \right)}{d^3}}{d^3}$$

input `Int[(e + f*x)^2*(a + b*ArcTanh[c + d*x])^2,x]`

output `((((d*e - c*f + f*(c + d*x))^3*(a + b*ArcTanh[c + d*x])^2)/(3*f) - (2*b*(-1/2*(b*f^3*(c + d*x)) - 3*a*f^2*(d*e - c*f)*(c + d*x) + (b*f^3*ArcTanh[c + d*x])/2 - 3*b*f^2*(d*e - c*f)*(c + d*x)*ArcTanh[c + d*x] - (f^3*(c + d*x)^2*(a + b*ArcTanh[c + d*x]))/2 + ((d*e - c*f)*(d^2*e^2 - 2*c*d*e*f + (3 + c^2)*f^2)*(a + b*ArcTanh[c + d*x])^2)/(2*b) - (f*(3*d^2*e^2 - 6*c*d*e*f + (1 + 3*c^2)*f^2)*(a + b*ArcTanh[c + d*x])^2)/(2*b) + f*(3*d^2*e^2 - 6*c*d*e*f + (1 + 3*c^2)*f^2)*(a + b*ArcTanh[c + d*x])*Log[2/(1 - c - d*x)] - (3*b*f^2*(d*e - c*f)*Log[1 - (c + d*x)^2])/2 + (b*f*(3*d^2*e^2 - 6*c*d*e*f + (1 + 3*c^2)*f^2)*PolyLog[2, -((1 + c + d*x)/(1 - c - d*x))])/2))/(3*f))/d^3`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6480

```
Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)*((d_.) + (e_.)*(x_.))^(q_.), x_Symbol]
:> Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] -
Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1),
(d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x]
&& IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

rule 6661

```
Int[((a_.) + ArcTanh[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol]
:> Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x]
&& IGtQ[p, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1410 vs. $2(358) = 716$.

Time = 0.61 (sec) , antiderivative size = 1411, normalized size of antiderivative = 3.77

| method | result | size |
|-------------------|---------------------------------|------|
| parts | Expression too large to display | 1411 |
| derivativedivides | Expression too large to display | 1412 |
| default | Expression too large to display | 1412 |
| risch | Expression too large to display | 1993 |

input

```
int((f*x+e)^2*(a+b*arctanh(d*x+c))^2,x,method=_RETURNVERBOSE)
```

output

```

1/3*a^2*(f*x+e)^3/f+b^2/d*(1/3/d^2*f^2*arctanh(d*x+c)^2*(d*x+c)^3-1/d^2*f^
2*arctanh(d*x+c)^2*(d*x+c)^2*c+1/d*f*arctanh(d*x+c)^2*(d*x+c)^2*e+1/d^2*f^
2*arctanh(d*x+c)^2*(d*x+c)*c^2-2/d*f*arctanh(d*x+c)^2*(d*x+c)*c*e+arctanh(
d*x+c)^2*(d*x+c)*e^2-1/3/d^2*f^2*arctanh(d*x+c)^2*c^3+1/d*f*arctanh(d*x+c)
^2*c^2*e-arctanh(d*x+c)^2*c*e^2+1/3*d/f*arctanh(d*x+c)^2*e^3-2/3/d^2/f*(1/
2*arctanh(d*x+c)*ln(d*x+c+1)*d^3*e^3-1/2*arctanh(d*x+c)*ln(d*x+c-1)*d^3*e^
3-1/2*f^3*arctanh(d*x+c)*ln(d*x+c+1)*c^3-3/2*f^3*arctanh(d*x+c)*ln(d*x+c+1
)*c^2-3/2*f^3*arctanh(d*x+c)*ln(d*x+c+1)*c+1/2*f^3*arctanh(d*x+c)*ln(d*x+c
-1)*c^3-3/2*f^3*arctanh(d*x+c)*ln(d*x+c-1)*c^2+3/2*f^3*arctanh(d*x+c)*ln(d
*x+c-1)*c+3*arctanh(d*x+c)*f^3*c*(d*x+c)-3/2*f*arctanh(d*x+c)*ln(d*x+c-1)*
d^2*e^2-3*arctanh(d*x+c)*f^2*d*e*(d*x+c)-3/2*f^2*arctanh(d*x+c)*ln(d*x+c-1
)*d*e-3/2*f*arctanh(d*x+c)*ln(d*x+c+1)*d^2*e^2+3/2*f^2*arctanh(d*x+c)*ln(d
*x+c+1)*d*e-1/2*f^3*arctanh(d*x+c)*ln(d*x+c+1)-1/2*f^3*arctanh(d*x+c)*ln(d
*x+c-1)-1/2*arctanh(d*x+c)*f^3*(d*x+c)^2-3/2*f*arctanh(d*x+c)*ln(d*x+c+1)*
c*d^2*e^2+3*f^2*arctanh(d*x+c)*ln(d*x+c+1)*c*d*e-3/2*f^2*arctanh(d*x+c)*ln
(d*x+c-1)*c^2*d*e+3/2*f*arctanh(d*x+c)*ln(d*x+c-1)*c*d^2*e^2+3*f^2*arctanh
(d*x+c)*ln(d*x+c-1)*c*d*e+3/2*f^2*arctanh(d*x+c)*ln(d*x+c+1)*c^2*d*e-1/2*(
-c^3*f^3+3*c^2*d*e*f^2-3*c*d^2*e^2*f+d^3*e^3+3*c^2*f^3-6*c*d*e*f^2+3*d^2*e
^2*f-3*c*f^3+3*d*e*f^2+f^3)*(-1/2*dilog(1/2*d*x+1/2*c+1/2)-1/2*ln(d*x+c-1)
*ln(1/2*d*x+1/2*c+1/2)+1/4*ln(d*x+c-1)^2)-1/2*f^2*(f*(d*x+c)+1/2*(-6*c*...

```

Fricas [F]

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (fx + e)^2 (b \operatorname{arctanh}(dx + c) + a)^2 dx$$

input

```
integrate((f*x+e)^2*(a+b*arctanh(d*x+c))^2,x, algorithm="fricas")
```

output

```

integral(a^2*f^2*x^2 + 2*a^2*e*f*x + a^2*e^2 + (b^2*f^2*x^2 + 2*b^2*e*f*x
+ b^2*e^2)*arctanh(d*x + c)^2 + 2*(a*b*f^2*x^2 + 2*a*b*e*f*x + a*b*e^2)*ar
ctanh(d*x + c), x)

```

Sympy [F]

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (a + b \operatorname{atanh}(c + dx))^2 (e + fx)^2 dx$$

input `integrate((f*x+e)**2*(a+b*atanh(d*x+c))**2,x)`

output `Integral((a + b*atanh(c + d*x))**2*(e + f*x)**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 806 vs. $2(350) = 700$.

Time = 0.25 (sec) , antiderivative size = 806, normalized size of antiderivative = 2.16

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx = \text{Too large to display}$$

input `integrate((f*x+e)^2*(a+b*arctanh(d*x+c))^2,x, algorithm="maxima")`

output

```

1/3*a^2*f^2*x^3 + a^2*e*f*x^2 + (2*x^2*arctanh(d*x + c) + d*(2*x/d^2 - (c^
2 + 2*c + 1)*log(d*x + c + 1)/d^3 + (c^2 - 2*c + 1)*log(d*x + c - 1)/d^3))
*a*b*e*f + 1/3*(2*x^3*arctanh(d*x + c) + d*((d*x^2 - 4*c*x)/d^3 + (c^3 + 3
*c^2 + 3*c + 1)*log(d*x + c + 1)/d^4 - (c^3 - 3*c^2 + 3*c - 1)*log(d*x + c
- 1)/d^4))*a*b*f^2 + a^2*e^2*x + (2*(d*x + c)*arctanh(d*x + c) + log(-(d*
x + c)^2 + 1))*a*b*e^2/d + 1/3*(3*d^2*e^2 - 6*c*d*e*f + 3*c^2*f^2 + f^2)*(
log(d*x + c + 1)*log(-1/2*d*x - 1/2*c + 1/2) + dilog(1/2*d*x + 1/2*c + 1/2
))*b^2/d^3 - 1/6*(5*c^2*f^2 - 6*d*e*f - 6*(d*e*f - f^2)*c + f^2)*b^2*log(d
*x + c + 1)/d^3 + 1/6*(5*c^2*f^2 + 6*d*e*f - 6*(d*e*f + f^2)*c + f^2)*b^2*
log(d*x + c - 1)/d^3 + 1/12*(4*b^2*d*f^2*x + (b^2*d^3*f^2*x^3 + 3*b^2*d^3*
e*f*x^2 + 3*b^2*d^3*e^2*x + (c^3*f^2 + 3*d^2*e^2 - 3*(d*e*f - f^2)*c^2 - 3
*d*e*f + 3*(d^2*e^2 - 2*d*e*f + f^2)*c + f^2)*b^2)*log(d*x + c + 1)^2 + (b
^2*d^3*f^2*x^3 + 3*b^2*d^3*e*f*x^2 + 3*b^2*d^3*e^2*x + (c^3*f^2 - 3*d^2*e^
2 - 3*(d*e*f + f^2)*c^2 - 3*d*e*f + 3*(d^2*e^2 + 2*d*e*f + f^2)*c - f^2)*b
^2)*log(-d*x - c + 1)^2 + 2*(b^2*d^2*f^2*x^2 + 2*(3*d^2*e*f - 2*c*d*f^2)*b
^2*x)*log(d*x + c + 1) - 2*(b^2*d^2*f^2*x^2 + 2*(3*d^2*e*f - 2*c*d*f^2)*b
^2*x + (b^2*d^3*f^2*x^3 + 3*b^2*d^3*e*f*x^2 + 3*b^2*d^3*e^2*x + (c^3*f^2 +
3*d^2*e^2 - 3*(d*e*f - f^2)*c^2 - 3*d*e*f + 3*(d^2*e^2 - 2*d*e*f + f^2)*c
+ f^2)*b^2)*log(d*x + c + 1))*log(-d*x - c + 1))/d^3

```

Giac [F]

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (fx + e)^2 (b \operatorname{arctanh}(dx + c) + a)^2 dx$$

input

```
integrate((f*x+e)^2*(a+b*arctanh(d*x+c))^2,x, algorithm="giac")
```

output

```
integrate((f*x + e)^2*(b*arctanh(d*x + c) + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (e + fx)^2 (a + b \operatorname{atanh}(c + dx))^2 dx$$

input `int((e + f*x)^2*(a + b*atanh(c + d*x))^2,x)`output `int((e + f*x)^2*(a + b*atanh(c + d*x))^2, x)`**Reduce [F]**

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^2 dx = \text{Too large to display}$$

input `int((f*x+e)^2*(a+b*atanh(d*x+c))^2,x)`

output

```
( - 2*atanh(c + d*x)**2*b**2*c**3*f**2 + 3*atanh(c + d*x)**2*b**2*c**2*d*e
*f + 2*atanh(c + d*x)**2*b**2*c*f**2 + 3*atanh(c + d*x)**2*b**2*d**3*e**2*
x + 3*atanh(c + d*x)**2*b**2*d**3*e*f*x**2 + atanh(c + d*x)**2*b**2*d**3*f
**2*x**3 - 3*atanh(c + d*x)**2*b**2*d*e*f + 2*atanh(c + d*x)*a*b*c**3*f**2
- 6*atanh(c + d*x)*a*b*c**2*d*e*f + 6*atanh(c + d*x)*a*b*c**2*f**2 + 6*at
anh(c + d*x)*a*b*c*d**2*e**2 - 12*atanh(c + d*x)*a*b*c*d*e*f + 6*atanh(c +
d*x)*a*b*c*f**2 + 6*atanh(c + d*x)*a*b*d**3*e**2*x + 6*atanh(c + d*x)*a*b
*d**3*e*f*x**2 + 2*atanh(c + d*x)*a*b*d**3*f**2*x**3 + 6*atanh(c + d*x)*a*
b*d**2*e**2 - 6*atanh(c + d*x)*a*b*d*e*f + 2*atanh(c + d*x)*a*b*f**2 - 5*a
tanh(c + d*x)*b**2*c**2*f**2 + 6*atanh(c + d*x)*b**2*c*d*e*f - 4*atanh(c +
d*x)*b**2*c*d*f**2*x - 6*atanh(c + d*x)*b**2*c*f**2 + 6*atanh(c + d*x)*b*
**2*d**2*e*f*x + atanh(c + d*x)*b**2*d**2*f**2*x**2 + 6*atanh(c + d*x)*b**2
*d*e*f - atanh(c + d*x)*b**2*f**2 + 6*int((atanh(c + d*x)*x)/(c**2 + 2*c*d
*x + d**2*x**2 - 1),x)*b**2*c**2*d**2*f**2 - 12*int((atanh(c + d*x)*x)/(c*
**2 + 2*c*d*x + d**2*x**2 - 1),x)*b**2*c*d**3*e*f + 6*int((atanh(c + d*x)*x
)/(c**2 + 2*c*d*x + d**2*x**2 - 1),x)*b**2*d**4*e**2 + 2*int((atanh(c + d*
x)*x)/(c**2 + 2*c*d*x + d**2*x**2 - 1),x)*b**2*d**2*f**2 + 6*log(c + d*x -
1)*a*b*c**2*f**2 - 12*log(c + d*x - 1)*a*b*c*d*e*f + 6*log(c + d*x - 1)*a
*b*d**2*e**2 + 2*log(c + d*x - 1)*a*b*f**2 - 6*log(c + d*x - 1)*b**2*c*f**
2 + 6*log(c + d*x - 1)*b**2*d*e*f + 3*a**2*d**3*e**2*x + 3*a**2*d**3*e...
```

3.40 $\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^2 dx$

| | |
|---|-----|
| Optimal result | 375 |
| Mathematica [A] (verified) | 376 |
| Rubi [A] (verified) | 376 |
| Maple [B] (verified) | 378 |
| Fricas [F] | 379 |
| Sympy [F] | 379 |
| Maxima [B] (verification not implemented) | 380 |
| Giac [F] | 381 |
| Mupad [F(-1)] | 381 |
| Reduce [F] | 381 |

Optimal result

Integrand size = 18, antiderivative size = 221

$$\begin{aligned} & \int (e + fx)(a + b \operatorname{arctanh}(c + dx))^2 dx \\ &= \frac{abfx}{d} + \frac{b^2 f(c + dx) \operatorname{arctanh}(c + dx)}{d^2} + \frac{(de - cf)(a + b \operatorname{arctanh}(c + dx))^2}{d^2} \\ & \quad - \frac{(d^2 e^2 - 2cdef + (1 + c^2) f^2)(a + b \operatorname{arctanh}(c + dx))^2}{2d^2 f} \\ & \quad + \frac{(e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^2}{2f} \\ & \quad - \frac{2b(de - cf)(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2}{1 - c - dx}\right)}{d^2} \\ & \quad + \frac{b^2 f \log(1 - (c + dx)^2)}{2d^2} - \frac{b^2 (de - cf) \operatorname{PolyLog}\left(2, -\frac{1 + c + dx}{1 - c - dx}\right)}{d^2} \end{aligned}$$

output

```
a*b*f*x/d+b^2*f*(d*x+c)*arctanh(d*x+c)/d^2+(-c*f+d*e)*(a+b*arctanh(d*x+c))
^2/d^2-1/2*(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)*(a+b*arctanh(d*x+c))^2/d^2/f+1/
2*(f*x+e)^2*(a+b*arctanh(d*x+c))^2/f-2*b*(-c*f+d*e)*(a+b*arctanh(d*x+c))*l
n(2/(-d*x-c+1))/d^2+1/2*b^2*f*ln(1-(d*x+c)^2)/d^2-b^2*(-c*f+d*e)*polylog(2
,-(d*x+c+1)/(-d*x-c+1))/d^2
```


Mathematica [A] (verified)

Time = 0.84 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.23

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^2 dx$$

$$= \frac{2a^2cde + 2abcf - a^2c^2f + 2a^2d^2ex + 2abdfx + a^2d^2fx^2 + b^2(-1 + c + dx)(2de + f - cf + dfx) \operatorname{arctanh}(c + dx)}{d^2}$$

input

```
Integrate[(e + f*x)*(a + b*ArcTanh[c + d*x])^2,x]
```

output

```
(2*a^2*c*d*e + 2*a*b*c*f - a^2*c^2*f + 2*a^2*d^2*e*x + 2*a*b*d*f*x + a^2*d^2*f*x^2 + b^2*(-1 + c + d*x)*(2*d*e + f - c*f + d*f*x)*ArcTanh[c + d*x]^2 + 2*b*ArcTanh[c + d*x]*(-((c + d*x)*(-(b*f) + a*c*f - a*d*(2*e + f*x))) - 2*b*(d*e - c*f)*Log[1 + E^(-2*ArcTanh[c + d*x])]) + a*b*f*Log[1 - c - d*x] - a*b*f*Log[1 + c + d*x] - 4*a*b*d*e*Log[1/Sqrt[1 - (c + d*x)^2]] - 2*b^2*f*Log[1/Sqrt[1 - (c + d*x)^2]] + 4*a*b*c*f*Log[1/Sqrt[1 - (c + d*x)^2]] + 2*b^2*(d*e - c*f)*PolyLog[2, -E^(-2*ArcTanh[c + d*x])])/(2*d^2)
```

Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.05, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {6661, 27, 6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^2 dx$$

$$\downarrow \text{6661}$$

$$\int \frac{(d(e - \frac{cf}{d}) + f(c + dx))(a + b \operatorname{arctanh}(c + dx))^2}{d} d(c + dx)$$

$$\downarrow \text{27}$$

$$\int \frac{(de - cf + f(c + dx))(a + b \operatorname{arctanh}(c + dx))^2}{d^2} d(c + dx)$$

$$\frac{\frac{(f(c+dx)-cf+de)^2(a+b\operatorname{arctanh}(c+dx))^2}{2f} - \frac{b \int \left(\frac{(d^2e^2 - 2cdf e + (c^2+1)f^2 + 2f(de-cf)(c+dx))^{(a+b\operatorname{arctanh}(c+dx))}}{1-(c+dx)^2} - f^2(a+b\operatorname{arctanh}(c+dx)) \right)}{d^2}}{f}}{d^2}$$

$$\frac{\frac{(f(c+dx)-cf+de)^2(a+b\operatorname{arctanh}(c+dx))^2}{2f} - \frac{b \left(\frac{((c^2+1)f^2 - 2cdf e + d^2e^2)^{(a+b\operatorname{arctanh}(c+dx))^2}}{2b} - \frac{f(de-cf)(a+b\operatorname{arctanh}(c+dx))^2}{b} + 2f(de-c) \right)}{d^2}}{f}}$$

input `Int[(e + f*x)*(a + b*ArcTanh[c + d*x])^2,x]`

output `((d*e - c*f + f*(c + d*x))^2*(a + b*ArcTanh[c + d*x])^2)/(2*f) - (b*(-(a*f^2*(c + d*x)) - b*f^2*(c + d*x)*ArcTanh[c + d*x] - (f*(d*e - c*f)*(a + b*ArcTanh[c + d*x])^2)/b + ((d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2)*(a + b*ArcTanh[c + d*x])^2)/(2*b) + 2*f*(d*e - c*f)*(a + b*ArcTanh[c + d*x])*Log[2/(1 - c - d*x)] - (b*f^2*Log[1 - (c + d*x)^2])/2 + b*f*(d*e - c*f)*PolyLog[2, -((1 + c + d*x)/(1 - c - d*x))])/f)/d^2`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6480 `Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^p_)*((d_.) + (e_.)*(x_.))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]`

rule 6661

```
Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] :> Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*
ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IG
tQ[p, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 452 vs. 2(215) = 430.

Time = 0.43 (sec) , antiderivative size = 453, normalized size of antiderivative = 2.05

| method | result |
|-------------------|---|
| parts | $a^2 \left(\frac{1}{2} f x^2 + e x \right) + \frac{b^2 \left(\frac{\operatorname{arctanh}(dx+c)^2 (dx+c)^2 f}{2d} - \frac{\operatorname{arctanh}(dx+c)^2 (dx+c) c f}{d} + \operatorname{arctanh}(dx+c)^2 e (dx+c) - \operatorname{arctanh}(dx+c) \right)}{d}$ |
| derivativedivides | $\frac{a^2 \left(f c (dx+c) - e (dx+c) d - \frac{f (dx+c)^2}{2} \right)}{d} - \frac{b^2 \left(\operatorname{arctanh}(dx+c)^2 f c (dx+c) - \operatorname{arctanh}(dx+c)^2 e (dx+c) d - \frac{\operatorname{arctanh}(dx+c)^2 f (dx+c)^2}{2} \right)}{d}$ |
| default | $\frac{a^2 \left(f c (dx+c) - e (dx+c) d - \frac{f (dx+c)^2}{2} \right)}{d} - \frac{b^2 \left(\operatorname{arctanh}(dx+c)^2 f c (dx+c) - \operatorname{arctanh}(dx+c)^2 e (dx+c) d - \frac{\operatorname{arctanh}(dx+c)^2 f (dx+c)^2}{2} \right)}{d}$ |
| risch | $a^2 e x + \frac{a b f x}{d} + \frac{e \ln(-dx-c-1) a b}{d} - \frac{e b^2 \ln(-dx-c+1)^2}{4d} + \frac{b a c f}{d^2} + \frac{b^2 \operatorname{dilog}\left(-\frac{dx}{2} - \frac{c}{2} + \frac{1}{2}\right) e}{d} + \frac{b^2 \ln(-dx-c)}{2d^2}$ |

input

```
int((f*x+e)*(a+b*arctanh(d*x+c))^2,x,method=_RETURNVERBOSE)
```

output

```
a^2*(1/2*f*x^2+e*x)+b^2/d*(1/2/d*arctanh(d*x+c)^2*(d*x+c)^2*f-1/d*arctanh(d*x+c)^2*(d*x+c)*c*f+arctanh(d*x+c)^2*e*(d*x+c)-1/d*(-arctanh(d*x+c)*(d*x+c)*f+arctanh(d*x+c)*ln(d*x+c-1)*f*c-arctanh(d*x+c)*ln(d*x+c-1)*d*e-1/2*arctanh(d*x+c)*ln(d*x+c-1)*f+arctanh(d*x+c)*ln(d*x+c+1)*f*c-arctanh(d*x+c)*ln(d*x+c+1)*d*e+1/2*arctanh(d*x+c)*ln(d*x+c+1)*f-1/2*ln(d*x+c-1)*f-1/2*ln(d*x+c+1)*f-1/2*(-2*c*f+2*d*e+f)*(-1/2*dilog(1/2*d*x+1/2*c+1/2)-1/2*ln(d*x+c-1)*ln(1/2*d*x+1/2*c+1/2)+1/4*ln(d*x+c-1)^2)-1/2*(-2*c*f+2*d*e-f)*(-1/4*ln(d*x+c+1)^2+1/2*(ln(d*x+c+1)-ln(1/2*d*x+1/2*c+1/2))*ln(-1/2*d*x-1/2*c+1/2)-1/2*dilog(1/2*d*x+1/2*c+1/2))))+2*b*a/d*(1/2/d*arctanh(d*x+c)*(d*x+c)^2*f-1/d*arctanh(d*x+c)*(d*x+c)*c*f+arctanh(d*x+c)*e*(d*x+c)-1/2/d*(-f*(d*x+c)-1/2*(-2*c*f+2*d*e+f)*ln(d*x+c-1)+1/2*(2*c*f-2*d*e+f)*ln(d*x+c+1)))
```

Fricas [F]

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^2 dx = \int (fx + e)(b \operatorname{artanh}(dx + c) + a)^2 dx$$

input

```
integrate((f*x+e)*(a+b*arctanh(d*x+c))^2,x, algorithm="fricas")
```

output

```
integral(a^2*f*x + a^2*e + (b^2*f*x + b^2*e)*arctanh(d*x + c)^2 + 2*(a*b*f*x + a*b*e)*arctanh(d*x + c), x)
```

Sympy [F]

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^2 dx = \int (a + b \operatorname{atanh}(c + dx))^2 (e + fx) dx$$

input

```
integrate((f*x+e)*(a+b*atanh(d*x+c))**2,x)
```

output

```
Integral((a + b*atanh(c + d*x))**2*(e + f*x), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 415 vs. $2(207) = 414$.

Time = 0.23 (sec) , antiderivative size = 415, normalized size of antiderivative = 1.88

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^2 dx = \frac{1}{2} a^2 f x^2 + \frac{1}{2} \left(2x^2 \operatorname{artanh}(dx + c) + d \left(\frac{2x}{d^2} - \frac{(c^2 + 2c + 1) \log(dx + c + 1)}{d^3} + \frac{(c^2 - 2c + 1) \log(dx + c - 1)}{d^3} \right) \right) + a^2 e x + \frac{(2(dx + c) \operatorname{artanh}(dx + c) + \log(-(dx + c)^2 + 1)) a b e}{d} + \frac{(de - cf) (\log(dx + c + 1) \log(-\frac{1}{2} dx - \frac{1}{2} c + \frac{1}{2}) + \operatorname{Li}_2(\frac{1}{2} dx + \frac{1}{2} c + \frac{1}{2})) b^2}{d^2} + \frac{(cf + f) b^2 \log(dx + c + 1)}{2 d^2} - \frac{(cf - f) b^2 \log(dx + c - 1)}{2 d^2} + \frac{4 b^2 d f x \log(dx + c + 1) + (b^2 d^2 f x^2 + 2 b^2 d^2 e x - (c^2 f - 2(de - f)c - 2de + f) b^2) \log(dx + c + 1)^2}{d^2}$$

input `integrate((f*x+e)*(a+b*arctanh(d*x+c))^2,x, algorithm="maxima")`

output

```
1/2*a^2*f*x^2 + 1/2*(2*x^2*arctanh(d*x + c) + d*(2*x/d^2 - (c^2 + 2*c + 1)
*log(d*x + c + 1)/d^3 + (c^2 - 2*c + 1)*log(d*x + c - 1)/d^3))*a*b*f + a^2
*e*x + (2*(d*x + c)*arctanh(d*x + c) + log(-(d*x + c)^2 + 1))*a*b*e/d + (d
*e - c*f)*(log(d*x + c + 1)*log(-1/2*d*x - 1/2*c + 1/2) + dilog(1/2*d*x +
1/2*c + 1/2))*b^2/d^2 + 1/2*(c*f + f)*b^2*log(d*x + c + 1)/d^2 - 1/2*(c*f
- f)*b^2*log(d*x + c - 1)/d^2 + 1/8*(4*b^2*d*f*x*log(d*x + c + 1) + (b^2*d
^2*f*x^2 + 2*b^2*d^2*e*x - (c^2*f - 2*(d*e - f)*c - 2*d*e + f)*b^2)*log(d*
x + c + 1)^2 + (b^2*d^2*f*x^2 + 2*b^2*d^2*e*x - (c^2*f - 2*(d*e + f)*c + 2
*d*e + f)*b^2)*log(-d*x - c + 1)^2 - 2*(2*b^2*d*f*x + (b^2*d^2*f*x^2 + 2*b
^2*d^2*e*x - (c^2*f - 2*(d*e - f)*c - 2*d*e + f)*b^2)*log(d*x + c + 1))*lo
g(-d*x - c + 1))/d^2
```

Giac [F]

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^2 dx = \int (fx + e)(b \operatorname{artanh}(dx + c) + a)^2 dx$$

input `integrate((f*x+e)*(a+b*arctanh(d*x+c))^2,x, algorithm="giac")`

output `integrate((f*x + e)*(b*arctanh(d*x + c) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^2 dx = \int (e + fx) (a + b \operatorname{atanh}(c + dx))^2 dx$$

input `int((e + f*x)*(a + b*atanh(c + d*x))^2,x)`

output `int((e + f*x)*(a + b*atanh(c + d*x))^2, x)`

Reduce [F]

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^2 dx$$

$$= \frac{\operatorname{atanh}(dx + c)^2 b^2 c^2 f + 2 \operatorname{atanh}(dx + c)^2 b^2 d^2 e x + \operatorname{atanh}(dx + c)^2 b^2 d^2 f x^2 - \operatorname{atanh}(dx + c)^2 b^2 f - 2 \operatorname{atanh}(dx + c) b^2 d^2 e x - 2 \operatorname{atanh}(dx + c) b^2 d^2 f x^2 + \operatorname{atanh}(dx + c) b^2 f + a^2 b^2 d^2 e x + a^2 b^2 d^2 f x^2 - a^2 b^2 f}{2}$$

input `int((f*x+e)*(a+b*atanh(d*x+c))^2,x)`

output

```
(atanh(c + d*x)**2*b**2*c**2*f + 2*atanh(c + d*x)**2*b**2*d**2*e*x + atanh
(c + d*x)**2*b**2*d**2*f*x**2 - atanh(c + d*x)**2*b**2*f - 2*atanh(c + d*x
)*a*b*c**2*f + 4*atanh(c + d*x)*a*b*c*d*e - 4*atanh(c + d*x)*a*b*c*f + 4*a
tanh(c + d*x)*a*b*d**2*e*x + 2*atanh(c + d*x)*a*b*d**2*f*x**2 + 4*atanh(c
+ d*x)*a*b*d*e - 2*atanh(c + d*x)*a*b*f + 2*atanh(c + d*x)*b**2*c*f + 2*at
anh(c + d*x)*b**2*d*f*x + 2*atanh(c + d*x)*b**2*f - 4*int((atanh(c + d*x)*
x)/(c**2 + 2*c*d*x + d**2*x**2 - 1),x)*b**2*c*d**2*f + 4*int((atanh(c + d*
x)*x)/(c**2 + 2*c*d*x + d**2*x**2 - 1),x)*b**2*d**3*e - 4*log(c + d*x - 1)
*a*b*c*f + 4*log(c + d*x - 1)*a*b*d*e + 2*log(c + d*x - 1)*b**2*f + 2*a**2
*d**2*e*x + a**2*d**2*f*x**2 + 2*a*b*d*f*x)/(2*d**2)
```

3.41 $\int (a + b \operatorname{arctanh}(c + dx))^2 dx$

| | |
|----------------------------|-----|
| Optimal result | 383 |
| Mathematica [A] (verified) | 383 |
| Rubi [A] (verified) | 384 |
| Maple [A] (verified) | 386 |
| Fricas [F] | 387 |
| Sympy [F] | 387 |
| Maxima [F] | 387 |
| Giac [F] | 388 |
| Mupad [F(-1)] | 388 |
| Reduce [F] | 389 |

Optimal result

Integrand size = 12, antiderivative size = 97

$$\int (a + b \operatorname{arctanh}(c + dx))^2 dx = \frac{(a + b \operatorname{arctanh}(c + dx))^2}{d} + \frac{(c + dx)(a + b \operatorname{arctanh}(c + dx))^2}{d} - \frac{2b(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2}{1-c-dx}\right)}{d} - \frac{b^2 \operatorname{PolyLog}\left(2, -\frac{1+c+dx}{1-c-dx}\right)}{d}$$

output

$(a+b*\operatorname{arctanh}(d*x+c))^2/d+(d*x+c)*(a+b*\operatorname{arctanh}(d*x+c))^2/d-2*b*(a+b*\operatorname{arctanh}(d*x+c))*\ln(2/(-d*x-c+1))/d-b^2*\operatorname{polylog}(2,-(d*x+c+1)/(-d*x-c+1))/d$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.06

$$\int (a + b \operatorname{arctanh}(c + dx))^2 dx$$

$$= \frac{b^2(-1 + c + dx)\operatorname{arctanh}(c + dx)^2 + 2b\operatorname{arctanh}(c + dx) (ac + adx - b \log(1 + e^{-2\operatorname{arctanh}(c+dx)})) + a(ac + adx - b \log(1 + e^{-2\operatorname{arctanh}(c+dx)}))}{d}$$

input `Integrate[(a + b*ArcTanh[c + d*x])^2,x]`

output `(b^2*(-1 + c + d*x)*ArcTanh[c + d*x]^2 + 2*b*ArcTanh[c + d*x]*(a*c + a*d*x - b*Log[1 + E^(-2*ArcTanh[c + d*x])]) + a*(a*c + a*d*x - 2*b*Log[1/Sqrt[1 - (c + d*x)^2]]) + b^2*PolyLog[2, -E^(-2*ArcTanh[c + d*x])])/d`

Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.96, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6653, 6436, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + b \operatorname{arctanh}(c + dx))^2 dx \\
 & \quad \downarrow \text{6653} \\
 & \frac{\int (a + b \operatorname{arctanh}(c + dx))^2 d(c + dx)}{d} \\
 & \quad \downarrow \text{6436} \\
 & \frac{(c + dx)(a + b \operatorname{arctanh}(c + dx))^2 - 2b \int \frac{(c+dx)(a+b \operatorname{arctanh}(c+dx))}{1-(c+dx)^2} d(c + dx)}{d} \\
 & \quad \downarrow \text{6546} \\
 & \frac{(c + dx)(a + b \operatorname{arctanh}(c + dx))^2 - 2b \left(\int \frac{a+b \operatorname{arctanh}(c+dx)}{-c-dx+1} d(c + dx) - \frac{(a+b \operatorname{arctanh}(c+dx))^2}{2b} \right)}{d} \\
 & \quad \downarrow \text{6470} \\
 & \frac{(c + dx)(a + b \operatorname{arctanh}(c + dx))^2 - 2b \left(-b \int \frac{\log\left(\frac{2}{-c-dx+1}\right)}{1-(c+dx)^2} d(c + dx) - \frac{(a+b \operatorname{arctanh}(c+dx))^2}{2b} + \log\left(\frac{2}{-c-dx+1}\right) (a + \right)}{d} \\
 & \quad \downarrow \text{2849}
 \end{aligned}$$

$$(c + dx)(a + \operatorname{arctanh}(c + dx))^2 - 2b \left(b \int \frac{\log\left(\frac{2}{-c-dx+1}\right)}{1-\frac{2}{-c-dx+1}} d \frac{1}{-c-dx+1} - \frac{(a+\operatorname{arctanh}(c+dx))^2}{2b} + \log\left(\frac{2}{-c-dx+1}\right) (a + b \operatorname{arctanh}(c + dx)) \right) / d$$

↓ 2752

$$(c + dx)(a + \operatorname{arctanh}(c + dx))^2 - 2b \left(-\frac{(a+\operatorname{arctanh}(c+dx))^2}{2b} + \log\left(\frac{2}{-c-dx+1}\right) (a + \operatorname{arctanh}(c + dx)) + \frac{1}{2} b \operatorname{PolyLog}[2, 1 - 2/(1 - c - dx)] \right) / d$$

input

```
Int[(a + b*ArcTanh[c + d*x])^2,x]
```

output

```
((c + d*x)*(a + b*ArcTanh[c + d*x])^2 - 2*b*(-1/2*(a + b*ArcTanh[c + d*x])^2/b + (a + b*ArcTanh[c + d*x])*Log[2/(1 - c - d*x)] + (b*PolyLog[2, 1 - 2/(1 - c - d*x)])/2))/d
```

Defintions of rubi rules used

rule 2752

```
Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]
```

rule 2849

```
Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]
```

rule 6436

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^p, x_Symbol] := Simp[x*(a + b*ArcTanh[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])
```

rule 6470 `Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)/((d_.) + (e_.)*(x_.)), x_Symbol]
-> Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e)
Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x]
/; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6546 `Int((((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)*(x_.))/((d_.) + (e_.)*(x_.)^2),
x_Symbol]
-> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/(c*d)
Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x]
/; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6653 `Int((((a_.) + ArcTanh[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.), x_Symbol]
-> Simp[1/d
Subst[Int[(a + b*ArcTanh[x])^p, x], x, c + d*x], x]
/; FreeQ[{a, b, c, d}, x] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.38

| method | result |
|-------------------|--|
| parts | $a^2x + \frac{b^2 \left(\operatorname{arctanh}(dx+c)^2(dx+c-1) + 2 \operatorname{arctanh}(dx+c)^2 - 2 \operatorname{arctanh}(dx+c) \ln \left(1 + \frac{(dx+c+1)^2}{1-(dx+c)^2} \right) - \operatorname{polylog} \left(2, -\frac{dx+c+1}{1-(dx+c)} \right) \right)}{d}$ |
| derivativedivides | $\frac{(dx+c)a^2+b^2 \left(\operatorname{arctanh}(dx+c)^2(dx+c-1) + 2 \operatorname{arctanh}(dx+c)^2 - 2 \operatorname{arctanh}(dx+c) \ln \left(1 + \frac{(dx+c+1)^2}{1-(dx+c)^2} \right) - \operatorname{polylog} \left(2, -\frac{dx+c+1}{1-(dx+c)} \right) \right)}{d}$ |
| default | $\frac{(dx+c)a^2+b^2 \left(\operatorname{arctanh}(dx+c)^2(dx+c-1) + 2 \operatorname{arctanh}(dx+c)^2 - 2 \operatorname{arctanh}(dx+c) \ln \left(1 + \frac{(dx+c+1)^2}{1-(dx+c)^2} \right) - \operatorname{polylog} \left(2, -\frac{dx+c+1}{1-(dx+c)} \right) \right)}{d}$ |
| risch | $a^2x + \frac{b^2 \ln(-dx-c+1)^2 c}{4d} - ba \ln(-dx - c + 1) x + \frac{ba \ln(-dx-c+1)}{d} + \frac{ba \ln(-dx-c-1)}{d} - \frac{b^2 \ln\left(\frac{dx}{2}\right)}{d}$ |

input `int((a+b*arctanh(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `a^2*x+b^2/d*(arctanh(d*x+c)^2*(d*x+c-1)+2*arctanh(d*x+c)^2-2*arctanh(d*x+c)*ln(1+(d*x+c+1)^2/(1-(d*x+c)^2))-polylog(2,-(d*x+c+1)^2/(1-(d*x+c)^2)))+2*b*a/d*((d*x+c)*arctanh(d*x+c)+1/2*ln(1-(d*x+c)^2))`

Fricas [F]

$$\int (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (b \operatorname{artanh}(dx + c) + a)^2 dx$$

input `integrate((a+b*arctanh(d*x+c))^2,x, algorithm="fricas")`

output `integral(b^2*arctanh(d*x + c)^2 + 2*a*b*arctanh(d*x + c) + a^2, x)`

Sympy [F]

$$\int (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (a + b \operatorname{atanh}(c + dx))^2 dx$$

input `integrate((a+b*atanh(d*x+c))**2,x)`

output `Integral((a + b*atanh(c + d*x))**2, x)`

Maxima [F]

$$\int (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (b \operatorname{artanh}(dx + c) + a)^2 dx$$

input `integrate((a+b*arctanh(d*x+c))^2,x, algorithm="maxima")`

output

```
-1/4*(c*d*((c + 1)*log(d*x + c + 1)/d^2 - (c - 1)*log(d*x + c - 1)/d^2) +
d^2*(2*x/d^2 - (c^2 + 2*c + 1)*log(d*x + c + 1)/d^3 + (c^2 - 2*c + 1)*log(
d*x + c - 1)/d^3) - 2*c*d*integrate(x*log(d*x + c + 1)/(d^2*x^2 + 2*c*d*x
+ c^2 - 1), x) - 2*c^2*integrate(log(d*x + c + 1)/(d^2*x^2 + 2*c*d*x + c^2
- 1), x) + d*((c + 1)*log(d*x + c + 1)/d^2 - (c - 1)*log(d*x + c - 1)/d^2
) - 6*d*integrate(x*log(d*x + c + 1)/(d^2*x^2 + 2*c*d*x + c^2 - 1), x) - 4
*c*integrate(log(d*x + c + 1)/(d^2*x^2 + 2*c*d*x + c^2 - 1), x) - (d*x + c
- 1)*(log(-d*x - c + 1)^2 - 2*log(-d*x - c + 1) + 2)/d - (d*x*log(d*x + c
+ 1)^2 + 2*(d*x - (d*x + c + 1)*log(d*x + c + 1))*log(-d*x - c + 1))/d -
2*integrate(log(d*x + c + 1)/(d^2*x^2 + 2*c*d*x + c^2 - 1), x))*b^2 + a^2*
x + (2*(d*x + c)*arctanh(d*x + c) + log(-(d*x + c)^2 + 1))*a*b/d
```

Giac [F]

$$\int (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (b \operatorname{artanh}(dx + c) + a)^2 dx$$

input

```
integrate((a+b*arctanh(d*x+c))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(d*x + c) + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (a + b \operatorname{atanh}(c + dx))^2 dx$$

input

```
int((a + b*atanh(c + d*x))^2,x)
```

output

```
int((a + b*atanh(c + d*x))^2, x)
```

Reduce [F]

$$\int (a + b \operatorname{arctanh}(c + dx))^2 dx$$

$$= \frac{\operatorname{atanh}(dx + c)^2 b^2 dx + 2 \operatorname{atanh}(dx + c) abc + 2 \operatorname{atanh}(dx + c) abdx + 2 \operatorname{atanh}(dx + c) ab + 2 \left(\int \frac{\operatorname{atanh}(dx + c)}{d^2 x^2 + 2cdx + c^2} dx \right)}{d}$$

input

```
int((a+b*atanh(d*x+c))^2,x)
```

output

```
(atanh(c + d*x)**2*b**2*d*x + 2*atanh(c + d*x)*a*b*c + 2*atanh(c + d*x)*a*
b*d*x + 2*atanh(c + d*x)*a*b + 2*int((atanh(c + d*x)*x)/(c**2 + 2*c*d*x +
d**2*x**2 - 1),x)*b**2*d**2 + 2*log(c + d*x - 1)*a*b + a**2*d*x)/d
```

$$3.42 \quad \int \frac{(a+b\operatorname{arctanh}(c+dx))^2}{e+fx} dx$$

| | |
|---|-----|
| Optimal result | 390 |
| Mathematica [C] (warning: unable to verify) | 391 |
| Rubi [A] (verified) | 392 |
| Maple [C] (warning: unable to verify) | 393 |
| Fricas [F] | 394 |
| Sympy [F] | 395 |
| Maxima [F] | 395 |
| Giac [F] | 395 |
| Mupad [F(-1)] | 396 |
| Reduce [F] | 396 |

Optimal result

Integrand size = 20, antiderivative size = 214

$$\begin{aligned} & \int \frac{(a + b\operatorname{arctanh}(c + dx))^2}{e + fx} dx \\ &= -\frac{(a + b\operatorname{arctanh}(c + dx))^2 \log\left(\frac{2}{1+c+dx}\right)}{f} \\ & \quad + \frac{(a + b\operatorname{arctanh}(c + dx))^2 \log\left(\frac{2d(e+fx)}{(de+f-cf)(1+c+dx)}\right)}{f} \\ & \quad + \frac{b(a + b\operatorname{arctanh}(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{f} \\ & \quad - \frac{b(a + b\operatorname{arctanh}(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2d(e+fx)}{(de+f-cf)(1+c+dx)}\right)}{f} \\ & \quad + \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1+c+dx}\right)}{2f} - \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2d(e+fx)}{(de+f-cf)(1+c+dx)}\right)}{2f} \end{aligned}$$

output

```
-(a+b*arctanh(d*x+c))^2*ln(2/(d*x+c+1))/f+(a+b*arctanh(d*x+c))^2*ln(2*d*(f
*x+e)/(-c*f+d*e+f)/(d*x+c+1))/f+b*(a+b*arctanh(d*x+c))*polylog(2,1-2/(d*x+
c+1))/f-b*(a+b*arctanh(d*x+c))*polylog(2,1-2*d*(f*x+e)/(-c*f+d*e+f)/(d*x+c
+1))/f+1/2*b^2*polylog(3,1-2/(d*x+c+1))/f-1/2*b^2*polylog(3,1-2*d*(f*x+e)/
(-c*f+d*e+f)/(d*x+c+1))/f
```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 90.24 (sec) , antiderivative size = 3808, normalized size of antiderivative = 17.79

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{e + fx} dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])^2/(e + f*x),x]
```

output

```
(a^2*Log[e + f*x])/f - ((2*I)*a*b*(I*ArcTanh[c + d*x]*(-Log[1/Sqrt[1 - (c + d*x)^2]] + Log[I*Sinh[ArcTanh[(d*e - c*f)/f] + ArcTanh[c + d*x]]]) + ((-I)*(I*ArcTanh[(d*e - c*f)/f] + I*ArcTanh[c + d*x])^2 - (I/4)*(Pi - (2*I)*ArcTanh[c + d*x])^2 + 2*(I*ArcTanh[(d*e - c*f)/f] + I*ArcTanh[c + d*x])*Log[1 - E^((2*I)*(I*ArcTanh[(d*e - c*f)/f] + I*ArcTanh[c + d*x]))] + (Pi - (2*I)*ArcTanh[c + d*x])*Log[1 - E^(I*(Pi - (2*I)*ArcTanh[c + d*x]))] - (Pi - (2*I)*ArcTanh[c + d*x])*Log[2*Sin[(Pi - (2*I)*ArcTanh[c + d*x])/2]] - 2*(I*ArcTanh[(d*e - c*f)/f] + I*ArcTanh[c + d*x])*Log[(2*I)*Sinh[ArcTanh[(d*e - c*f)/f] + ArcTanh[c + d*x]]) - I*PolyLog[2, E^((2*I)*(I*ArcTanh[(d*e - c*f)/f] + I*ArcTanh[c + d*x]))] - I*PolyLog[2, E^(I*(Pi - (2*I)*ArcTanh[c + d*x]))]/2))/f + (b^2*(d*e - c*f + f*(c + d*x))*((Sqrt[1 - (c + d*x)^2]*((d*e)/Sqrt[1 - (c + d*x)^2] - (c*f)/Sqrt[1 - (c + d*x)^2] + (f*(c + d*x))/Sqrt[1 - (c + d*x)^2])*ArcTanh[c + d*x]^3)/(3*(d*e - c*f)*(d*e - c*f + f*(c + d*x))) - (Sqrt[1 - (c + d*x)^2]*((d*e)/Sqrt[1 - (c + d*x)^2] - (c*f)/Sqrt[1 - (c + d*x)^2] + (f*(c + d*x))/Sqrt[1 - (c + d*x)^2])*ArcTanh[c + d*x]^3/3 + ArcTanh[c + d*x]^2*Log[1 + E^(-2*ArcTanh[c + d*x])] - ArcTanh[c + d*x]*PolyLog[2, -E^(-2*ArcTanh[c + d*x])] - PolyLog[3, -E^(-2*ArcTanh[c + d*x])]/2))/(f*(d*e - c*f + f*(c + d*x))) - (-6*d*e*ArcTanh[c + d*x]^3 + 2*f*ArcTanh[c + d*x]^3 + 6*c*f*ArcTanh[c + d*x]^3 - 4*E^ArcTanh[c - (d*e)/f]*Sqrt[1 - c^2 - (d^2*e^2)/f^2 + (2*c*d*e)/f]*f*ArcTanh[c + d*x]^3 - ...
```


Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.13, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {6661, 27, 6474}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{e + fx} dx \\
 & \quad \downarrow \text{6661} \\
 & \int \frac{d(a + b \operatorname{arctanh}(c + dx))^2}{d(e - \frac{cf}{d}) + f(c + dx)} d(c + dx) \\
 & \quad \downarrow \text{27} \\
 & \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{f(c + dx) - cf + de} d(c + dx) \\
 & \quad \downarrow \text{6474} \\
 & -\frac{b(a + b \operatorname{arctanh}(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2(de - cf + f(c + dx))}{(de - cf + f)(c + dx + 1)}\right)}{f} + \\
 & \quad \frac{(a + b \operatorname{arctanh}(c + dx))^2 \log\left(\frac{2(f(c + dx) - cf + de)}{(c + dx + 1)(-cf + de + f)}\right)}{f} + \\
 & \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{c + dx + 1}\right) (a + b \operatorname{arctanh}(c + dx))}{f} - \frac{\log\left(\frac{2}{c + dx + 1}\right) (a + b \operatorname{arctanh}(c + dx))^2}{f} \\
 & \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2(de - cf + f(c + dx))}{(de - cf + f)(c + dx + 1)}\right)}{2f} + \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{c + dx + 1}\right)}{2f}
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c + d*x])^2/(e + f*x),x]`

output

```

-(((a + b*ArcTanh[c + d*x])^2*Log[2/(1 + c + d*x)]/f) + ((a + b*ArcTanh[c
+ d*x])^2*Log[(2*(d*e - c*f + f*(c + d*x))]/((d*e + f - c*f)*(1 + c + d*x
)))]/f + (b*(a + b*ArcTanh[c + d*x])*PolyLog[2, 1 - 2/(1 + c + d*x)]/f -
(b*(a + b*ArcTanh[c + d*x])*PolyLog[2, 1 - (2*(d*e - c*f + f*(c + d*x))]/(
(d*e + f - c*f)*(1 + c + d*x)))]/f + (b^2*PolyLog[3, 1 - 2/(1 + c + d*x)]
)/(2*f) - (b^2*PolyLog[3, 1 - (2*(d*e - c*f + f*(c + d*x))]/((d*e + f - c*f
)*(1 + c + d*x)))]/(2*f)

```

Defintions of rubi rules used

rule 27

```

Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]

```

rule 6474

```

Int[((a_) + ArcTanh[(c_)*(x_)]*(b_))^2/((d_) + (e_)*(x_)), x_Symbol] :=
Simp[(-(a + b*ArcTanh[c*x])^2)*(Log[2/(1 + c*x)]/e), x] + (Simp[(a + b*Arc
Tanh[c*x])^2*(Log[2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/e), x] + Simp[b*(a
+ b*ArcTanh[c*x])*(PolyLog[2, 1 - 2/(1 + c*x)]/e), x] - Simp[b*(a + b*ArcT
anh[c*x])*(PolyLog[2, 1 - 2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/e), x] + S
imp[b^2*(PolyLog[3, 1 - 2/(1 + c*x)]/(2*e)), x] - Simp[b^2*(PolyLog[3, 1 -
2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/(2*e)), x]) /; FreeQ[{a, b, c, d, e}
, x] && NeQ[c^2*d^2 - e^2, 0]

```

rule 6661

```

Int[((a_) + ArcTanh[(c_) + (d_)*(x_)]*(b_))^(p_)*((e_) + (f_)*(x_))^(
m_), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*
ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IG
tQ[p, 0]

```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 39.94 (sec) , antiderivative size = 1700, normalized size of antiderivative = 7.94

| method | result | size |
|-------------------|---------------------------------|------|
| derivativedivides | Expression too large to display | 1700 |
| default | Expression too large to display | 1700 |
| parts | Expression too large to display | 1800 |

input `int((a+b*arctanh(d*x+c))^2/(f*x+e),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/d*(a^2*d*\ln(c*f-d*e-f*(d*x+c))/f-b^2*d*(-\ln(c*f-d*e-f*(d*x+c))/f*arctanh \\ & (d*x+c)^2+2/f*(1/2*arctanh(d*x+c)^2*\ln(f*c*(1+(d*x+c+1)^2/(1-(d*x+c)^2))+ \\ & (-1-(d*x+c+1)^2/(1-(d*x+c)^2))*e*d+(-(d*x+c+1)^2/(1-(d*x+c)^2)+1)*f)-1/4*I* \\ & Pi*csgn(I*(f*c*(1-(d*x+c+1)^2/((d*x+c)^2-1))+(-1+(d*x+c+1)^2/((d*x+c)^2-1) \\ &)*e*d+((d*x+c+1)^2/((d*x+c)^2-1)+1)*f)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))*(csg \\ & n(I*(f*c*(1-(d*x+c+1)^2/((d*x+c)^2-1))+(-1+(d*x+c+1)^2/((d*x+c)^2-1))*e*d+ \\ & ((d*x+c+1)^2/((d*x+c)^2-1)+1)*f))*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))-csg \\ & n(I*(f*c*(1-(d*x+c+1)^2/((d*x+c)^2-1))+(-1+(d*x+c+1)^2/((d*x+c)^2-1))*e*d \\ & +((d*x+c+1)^2/((d*x+c)^2-1)+1)*f)/(1-(d*x+c+1)^2/((d*x+c)^2-1))*csgn(I/(1 \\ & -(d*x+c+1)^2/((d*x+c)^2-1)))-csgn(I*(f*c*(1-(d*x+c+1)^2/((d*x+c)^2-1))+(-1 \\ & +(d*x+c+1)^2/((d*x+c)^2-1))*e*d+((d*x+c+1)^2/((d*x+c)^2-1)+1)*f))*csgn(I*(\\ & f*c*(1-(d*x+c+1)^2/((d*x+c)^2-1))+(-1+(d*x+c+1)^2/((d*x+c)^2-1))*e*d+((d*x \\ & +c+1)^2/((d*x+c)^2-1)+1)*f)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))+csgn(I*(f*c*(1- \\ & (d*x+c+1)^2/((d*x+c)^2-1))+(-1+(d*x+c+1)^2/((d*x+c)^2-1))*e*d+((d*x+c+1)^2 \\ & /((d*x+c)^2-1)+1)*f)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))^2*arctanh(d*x+c)^2+1/ \\ & 2*arctanh(d*x+c)*polylog(2,-(d*x+c+1)^2/(1-(d*x+c)^2))-1/4*polylog(3,-(d*x \\ & +c+1)^2/(1-(d*x+c)^2))-1/2*f*c/(c*f-d*e-f)*arctanh(d*x+c)^2*\ln(1-(c*f-d*e- \\ & f)*(d*x+c+1)^2/(1-(d*x+c)^2)/(-c*f+d*e-f))-1/2*f*c/(c*f-d*e-f)*arctanh(d*x \\ & +c)*polylog(2,(c*f-d*e-f)*(d*x+c+1)^2/(1-(d*x+c)^2)/(-c*f+d*e-f))+1/4*f*c/ \\ & (c*f-d*e-f)*polylog(3,(c*f-d*e-f)*(d*x+c+1)^2/(1-(d*x+c)^2)/(-c*f+d*e-f)... \end{aligned}$$

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{e + fx} dx = \int \frac{(b \operatorname{arctanh}(dx + c) + a)^2}{fx + e} dx$$

input `integrate((a+b*arctanh(d*x+c))^2/(f*x+e),x, algorithm="fricas")`

output `integral((b^2*arctanh(d*x + c)^2 + 2*a*b*arctanh(d*x + c) + a^2)/(f*x + e), x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{e + fx} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^2}{e + fx} dx$$

input `integrate((a+b*atanh(d*x+c))**2/(f*x+e),x)`

output `Integral((a + b*atanh(c + d*x))**2/(e + f*x), x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{e + fx} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^2}{fx + e} dx$$

input `integrate((a+b*arctanh(d*x+c))^2/(f*x+e),x, algorithm="maxima")`

output `a^2*log(f*x + e)/f + integrate(1/4*b^2*(log(d*x + c + 1) - log(-d*x - c + 1))^2/(f*x + e) + a*b*(log(d*x + c + 1) - log(-d*x - c + 1))/(f*x + e), x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{e + fx} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^2}{fx + e} dx$$

input `integrate((a+b*arctanh(d*x+c))^2/(f*x+e),x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)^2/(f*x + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{e + fx} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^2}{e + fx} dx$$

input `int((a + b*atanh(c + d*x))^2/(e + f*x),x)`output `int((a + b*atanh(c + d*x))^2/(e + f*x), x)`**Reduce [F]**

$$\begin{aligned} & \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{e + fx} dx \\ &= \frac{2 \left(\int \frac{\operatorname{atanh}(dx+c)}{fx+e} dx \right) abf + \left(\int \frac{\operatorname{atanh}(dx+c)^2}{fx+e} dx \right) b^2 f + \log(fx + e) a^2}{f} \end{aligned}$$

input `int((a+b*atanh(d*x+c))^2/(f*x+e),x)`output `(2*int(atanh(c + d*x)/(e + f*x),x)*a*b*f + int(atanh(c + d*x)**2/(e + f*x),x)*b**2*f + log(e + f*x)*a**2)/f`

$$3.43 \quad \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^2} dx$$

| | |
|---|-----|
| Optimal result | 397 |
| Mathematica [C] (warning: unable to verify) | 398 |
| Rubi [A] (verified) | 399 |
| Maple [A] (verified) | 401 |
| Fricas [F] | 402 |
| Sympy [F] | 402 |
| Maxima [F] | 403 |
| Giac [F] | 403 |
| Mupad [F(-1)] | 404 |
| Reduce [F] | 404 |

Optimal result

Integrand size = 20, antiderivative size = 401

$$\begin{aligned} \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^2} dx = & -\frac{(a + b \operatorname{arctanh}(c + dx))^2}{f(e + fx)} \\ & + \frac{bd(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2}{1-c-dx}\right)}{f(de + f - cf)} \\ & - \frac{bd(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{f(de - (1+c)f)} \\ & + \frac{2bd(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{(de + f - cf)(de - (1+c)f)} \\ & - \frac{2bd(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2d(e+fx)}{(de+f-cf)(1+c+dx)}\right)}{(de + f - cf)(de - (1+c)f)} \\ & + \frac{b^2 d \operatorname{PolyLog}\left(2, -\frac{1+c+dx}{1-c-dx}\right)}{2f(de + f - cf)} \\ & + \frac{b^2 d \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{2f(de - (1+c)f)} \\ & - \frac{b^2 d \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{(de + f - cf)(de - (1+c)f)} \\ & + \frac{b^2 d \operatorname{PolyLog}\left(2, 1 - \frac{2d(e+fx)}{(de+f-cf)(1+c+dx)}\right)}{(de + f - cf)(de - (1+c)f)} \end{aligned}$$

output

$$\begin{aligned}
& -(a+b*\operatorname{arctanh}(d*x+c))^2/f/(f*x+e)+b*d*(a+b*\operatorname{arctanh}(d*x+c))*\ln(2/(-d*x-c+1)) \\
& /f/(-c*f+d*e+f)-b*d*(a+b*\operatorname{arctanh}(d*x+c))*\ln(2/(d*x+c+1))/f/(d*e-(1+c)*f)+ \\
& 2*b*d*(a+b*\operatorname{arctanh}(d*x+c))*\ln(2/(d*x+c+1))/(-c*f+d*e+f)/(d*e-(1+c)*f)-2*b* \\
& d*(a+b*\operatorname{arctanh}(d*x+c))*\ln(2*d*(f*x+e)/(-c*f+d*e+f)/(d*x+c+1))/(-c*f+d*e+f) \\
& /f/(d*e-(1+c)*f)+1/2*b^2*d*\operatorname{polylog}(2,-(d*x+c+1)/(-d*x-c+1))/f/(-c*f+d*e+f)+1 \\
& /2*b^2*d*\operatorname{polylog}(2,1-2/(d*x+c+1))/f/(d*e-(1+c)*f)-b^2*d*\operatorname{polylog}(2,1-2/(d*x \\
& +c+1))/(-c*f+d*e+f)/(d*e-(1+c)*f)+b^2*d*\operatorname{polylog}(2,1-2*d*(f*x+e)/(-c*f+d*e+f) \\
& /f/(d*x+c+1))/(-c*f+d*e+f)/(d*e-(1+c)*f)
\end{aligned}$$

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 5.06 (sec) , antiderivative size = 419, normalized size of antiderivative = 1.04

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^2} dx$$

$$\begin{aligned}
& = \frac{-\frac{a^2}{f} + \frac{2ab \left((f-c^2f+d^2ex+cd(e-fx)) \operatorname{arctanh}(c+dx) - d(e+fx) \log \left(\frac{d(e+fx)}{\sqrt{1-(c+dx)^2}} \right) \right)}{(de+f-cf)(de-(1+c)f)}}{b^2 d(e+fx)} \left(-\frac{e^{-\operatorname{arctanh}\left(\frac{de-cf}{f}\right)} \operatorname{arctanh}\left(\frac{de-cf}{f}\right)}{f \sqrt{1-\frac{(de-cf)^2}{f^2}}} \right)
\end{aligned}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])^2/(e + f*x)^2,x]
```

output

$$\begin{aligned}
& (-\frac{a^2}{f}) + (2*a*b*((f - c^2*f + d^2*e*x + c*d*(e - f*x))*\operatorname{ArcTanh}[c + d*x] \\
& - d*(e + f*x)*\operatorname{Log}[(d*(e + f*x))/\operatorname{Sqrt}[1 - (c + d*x)^2]])/((d*e + f - c*f) \\
& *(d*e - (1 + c)*f)) + (b^2*d*(e + f*x)*(-(\operatorname{ArcTanh}[c + d*x]^2/(E^{\operatorname{ArcTanh}[(d \\
& *e - c*f)/f]*f*\operatorname{Sqrt}[1 - (d*e - c*f)^2/f^2])) + ((c + d*x)*\operatorname{ArcTanh}[c + d*x] \\
& ^2)/(d*(e + f*x)) + ((d*e - c*f)*(-2*\operatorname{ArcTanh}[c + d*x]*\operatorname{Log}[1 - E^{(2*(\operatorname{ArcTan} \\
& h[c - (d*e)/f] - \operatorname{ArcTanh}[c + d*x])]) + I*\operatorname{Pi}*\operatorname{Log}[1 + E^{(2*\operatorname{ArcTanh}[c + d*x])} \\
&] - I*\operatorname{Pi}*(\operatorname{ArcTanh}[c + d*x] + \operatorname{Log}[1/\operatorname{Sqrt}[1 - (c + d*x)^2]]) + 2*\operatorname{ArcTanh}[c - \\
& (d*e)/f]*(\operatorname{ArcTanh}[c + d*x] + \operatorname{Log}[1 - E^{(2*(\operatorname{ArcTanh}[c - (d*e)/f] - \operatorname{ArcTanh} \\
& [c + d*x])]) - \operatorname{Log}[I*\operatorname{Sinh}[\operatorname{ArcTanh}[(d*e - c*f)/f] + \operatorname{ArcTanh}[c + d*x]]) + \operatorname{Poly} \\
& \operatorname{Log}[2, E^{(2*(\operatorname{ArcTanh}[c - (d*e)/f] - \operatorname{ArcTanh}[c + d*x])})])])/(d^2*e^2 - 2* \\
& c*d*e*f + (-1 + c^2)*f^2))/(d*e - c*f)/(e + f*x)
\end{aligned}$$

Rubi [A] (verified)

Time = 1.91 (sec) , antiderivative size = 491, normalized size of antiderivative = 1.22, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {6659, 7292, 6671, 27, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + \operatorname{barctanh}(c + dx))^2}{(e + fx)^2} dx \\
 & \quad \downarrow \text{6659} \\
 & \frac{2bd \int \frac{a + \operatorname{barctanh}(c + dx)}{(e + fx)(1 - (c + dx)^2)} dx}{f} - \frac{(a + \operatorname{barctanh}(c + dx))^2}{f(e + fx)} \\
 & \quad \downarrow \text{7292} \\
 & \frac{2bd \int \frac{a + \operatorname{barctanh}(c + dx)}{(e + fx)(-c^2 - 2dxc - d^2x^2 + 1)} dx}{f} - \frac{(a + \operatorname{barctanh}(c + dx))^2}{f(e + fx)} \\
 & \quad \downarrow \text{6671} \\
 & \frac{2b \int \frac{d(a + \operatorname{barctanh}(c + dx))}{(d(e - \frac{cf}{d}) + f(c + dx))(1 - (c + dx)^2)} d(c + dx)}{f} - \frac{(a + \operatorname{barctanh}(c + dx))^2}{f(e + fx)} \\
 & \quad \downarrow \text{27} \\
 & \frac{2bd \int \frac{a + \operatorname{barctanh}(c + dx)}{(de - cf + f(c + dx))(1 - (c + dx)^2)} d(c + dx)}{f} - \frac{(a + \operatorname{barctanh}(c + dx))^2}{f(e + fx)} \\
 & \quad \downarrow \text{7276} \\
 & \frac{2bd \int \left(-\frac{a}{(c + dx - 1)(c + dx + 1)(de - cf + f(c + dx))} - \frac{\operatorname{barctanh}(c + dx)}{(c + dx - 1)(c + dx + 1)(de - cf + f(c + dx))} \right) d(c + dx)}{f} - \\
 & \quad \frac{(a + \operatorname{barctanh}(c + dx))^2}{f(e + fx)} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$2bd \left(-\frac{a \log(-c-dx+1)}{2(-cf+de+f)} + \frac{a \log(c+dx+1)}{2(de-(c+1)f)} - \frac{af \log(f(c+dx)-cf+de)}{(-cf+de+f)(de-(c+1)f)} + \frac{\operatorname{barctanh}(c+dx) \log\left(\frac{2}{-c-dx+1}\right)}{2(-cf+de+f)} - \frac{\operatorname{barctanh}(c+dx) \log\left(\frac{2}{c-dx+1}\right)}{2(-cf+de-f)} \right) \\ \frac{(a + \operatorname{barctanh}(c + dx))^2}{f(e + fx)}$$

input `Int[(a + b*ArcTanh[c + d*x])^2/(e + f*x)^2,x]`

output `-((a + b*ArcTanh[c + d*x])^2/(f*(e + f*x))) + (2*b*d*((b*ArcTanh[c + d*x]*Log[2/(1 - c - d*x)])/(2*(d*e + f - c*f)) - (a*Log[1 - c - d*x]/(2*(d*e + f - c*f)) - (b*ArcTanh[c + d*x]*Log[2/(1 + c + d*x)])/(2*(d*e - f - c*f)) + (b*f*ArcTanh[c + d*x]*Log[2/(1 + c + d*x)])/((d*e + f - c*f)*(d*e - (1 + c)*f)) + (a*Log[1 + c + d*x]/(2*(d*e - (1 + c)*f)) - (a*f*Log[d*e - c*f + f*(c + d*x)])/((d*e + f - c*f)*(d*e - (1 + c)*f)) - (b*f*ArcTanh[c + d*x]*Log[(2*(d*e - c*f + f*(c + d*x))]/((d*e + f - c*f)*(1 + c + d*x)))/((d*e + f - c*f)*(d*e - (1 + c)*f)) + (b*PolyLog[2, -((1 + c + d*x)/(1 - c - d*x))]/(4*(d*e + f - c*f)) + (b*PolyLog[2, 1 - 2/(1 + c + d*x)]/(4*(d*e - f - c*f)) - (b*f*PolyLog[2, 1 - 2/(1 + c + d*x)]/(2*(d*e + f - c*f)*(d*e - (1 + c)*f)) + (b*f*PolyLog[2, 1 - (2*(d*e - c*f + f*(c + d*x))]/((d*e + f - c*f)*(1 + c + d*x)))/((2*(d*e + f - c*f)*(d*e - (1 + c)*f)))))/f`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6659 `Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)])*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_), x_Symbol] := Simp[(e + f*x)^(m + 1)*((a + b*ArcTanh[c + d*x])^p/(f*(m + 1))), x] - Simp[b*d*(p/(f*(m + 1))) Int[(e + f*x)^(m + 1)*((a + b*ArcTanh[c + d*x])^(p - 1)/(1 - (c + d*x)^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && ILtQ[m, -1]`

```
rule 6671 Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.)*((A_.) + (B_.)*(x_) + (C_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/d Sub
st[Int[((d*e - c*f)/d + f*(x/d))^m*(-C/d^2 + (C/d^2)*x^2)^q*(a + b*ArcTanh[
x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, p, q}, x
] && EqQ[B*(1 - c^2) + 2*A*c*d, 0] && EqQ[2*c*C - B*d, 0]
```

```
rule 7276 Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

```
rule 7292 Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v !=
= u]
```

Maple [A] (verified)

Time = 3.58 (sec) , antiderivative size = 592, normalized size of antiderivative = 1.48

| method | result |
|-------------------|---|
| parts | $-\frac{a^2}{(fx+e)f} + b^2 \left(-\frac{d^2 \operatorname{arctanh}(dx+c)^2}{(f(dx+c)-cf+de)f} + \frac{2d^2}{(cf-de+f)(cf-de-f)} \left(-\frac{\operatorname{arctanh}(dx+c)f \ln(f(dx+c)-cf+de)}{(cf-de+f)(cf-de-f)} + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{2cf-2de-2f} - \frac{\operatorname{arctanh}(dx+c)}{cf-de+f} \right) \right)$ |
| derivativedivides | $\frac{a^2 d^2}{(cf-de-f(dx+c))f} + b^2 d^2 \left(\frac{\operatorname{arctanh}(dx+c)^2}{(cf-de-f(dx+c))f} - 2 \left(-\frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{2cf-2de-2f} + \frac{\operatorname{arctanh}(dx+c)f \ln(cf-de-f(dx+c))}{(cf-de-f)(cf-de+f)} + \frac{\operatorname{arctanh}(dx+c)}{cf-de+f} \right) \right)$ |
| default | $\frac{a^2 d^2}{(cf-de-f(dx+c))f} + b^2 d^2 \left(\frac{\operatorname{arctanh}(dx+c)^2}{(cf-de-f(dx+c))f} - 2 \left(-\frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{2cf-2de-2f} + \frac{\operatorname{arctanh}(dx+c)f \ln(cf-de-f(dx+c))}{(cf-de-f)(cf-de+f)} + \frac{\operatorname{arctanh}(dx+c)}{cf-de+f} \right) \right)$ |

input `int((a+b*arctanh(d*x+c))^2/(f*x+e)^2,x,method=_RETURNVERBOSE)`

output `-a^2/(f*x+e)/f+b^2/d*(-d^2/(f*(d*x+c)-c*f+d*e)/f*arctanh(d*x+c)^2+2*d^2/f*(-arctanh(d*x+c)*f/(c*f-d*e+f)/(c*f-d*e-f)*ln(f*(d*x+c)-c*f+d*e)+arctanh(d*x+c)/(2*c*f-2*d*e-2*f)*ln(d*x+c-1)-arctanh(d*x+c)/(2*c*f-2*d*e+2*f)*ln(d*x+c+1)-1/(c*f-d*e+f)/(c*f-d*e-f)*(1/2*f*(dilog((f*(d*x+c)-f)/(c*f-d*e-f))+ln(f*(d*x+c)-c*f+d*e)*ln((f*(d*x+c)-f)/(c*f-d*e-f)))-1/2*f*(dilog((f*(d*x+c)+f)/(c*f-d*e+f))+ln(f*(d*x+c)-c*f+d*e)*ln((f*(d*x+c)+f)/(c*f-d*e+f))))+1/2/(c*f-d*e+f)*(-1/2*dilog(1/2*d*x+1/2*c+1/2)-1/2*ln(d*x+c-1)*ln(1/2*d*x+1/2*c+1/2)+1/4*ln(d*x+c-1)^2)-1/2/(c*f-d*e+f)*(-1/4*ln(d*x+c+1)^2+1/2*(ln(d*x+c+1)-ln(1/2*d*x+1/2*c+1/2))*ln(-1/2*d*x-1/2*c+1/2)-1/2*dilog(1/2*d*x+1/2*c+1/2))))-2*b*a*d/(d*f*x+d*e)/f*arctanh(d*x+c)-2*b*a*d/(c*f-d*e+f)/(c*f-d*e-f)*ln(f*(d*x+c)-c*f+d*e)+2*b*a*d/f/(2*c*f-2*d*e-2*f)*ln(d*x+c-1)-2*b*a*d/f/(2*c*f-2*d*e+2*f)*ln(d*x+c+1)`

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^2} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^2}{(fx + e)^2} dx$$

input `integrate((a+b*arctanh(d*x+c))^2/(f*x+e)^2,x, algorithm="fricas")`

output `integral((b^2*arctanh(d*x + c)^2 + 2*a*b*arctanh(d*x + c) + a^2)/(f^2*x^2 + 2*e*f*x + e^2), x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^2} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^2}{(e + fx)^2} dx$$

input `integrate((a+b*atanh(d*x+c))**2/(f*x+e)**2,x)`

output `Integral((a + b*atanh(c + d*x))**2/(e + f*x)**2, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^2} dx = \int \frac{(b \operatorname{arctanh}(dx + c) + a)^2}{(fx + e)^2} dx$$

input `integrate((a+b*arctanh(d*x+c))^2/(f*x+e)^2,x, algorithm="maxima")`

output `(d*(log(d*x + c + 1)/(d*e*f - (c + 1)*f^2) - log(d*x + c - 1)/(d*e*f - (c - 1)*f^2) - 2*log(f*x + e)/(d^2*e^2 - 2*c*d*e*f + (c^2 - 1)*f^2)) - 2*arctanh(d*x + c)/(f^2*x + e*f))*a*b - 1/4*b^2*(log(-d*x - c + 1)^2/(f^2*x + e*f) + integrate(-((d*f*x + c*f - f)*log(d*x + c + 1)^2 + 2*(d*f*x + d*e - (d*f*x + c*f - f)*log(d*x + c + 1))*log(-d*x - c + 1))/(d*f^3*x^3 + c*e^2*f - e^2*f + (2*d*e*f^2 + c*f^3 - f^3)*x^2 + (d*e^2*f + 2*c*e*f^2 - 2*e*f^2)*x), x)) - a^2/(f^2*x + e*f)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^2} dx = \int \frac{(b \operatorname{arctanh}(dx + c) + a)^2}{(fx + e)^2} dx$$

input `integrate((a+b*arctanh(d*x+c))^2/(f*x+e)^2,x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)^2/(f*x + e)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^2} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^2}{(e + fx)^2} dx$$

input `int((a + b*atanh(c + d*x))^2/(e + f*x)^2,x)`output `int((a + b*atanh(c + d*x))^2/(e + f*x)^2, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^2} dx = \text{too large to display}$$

input `int((a+b*atanh(d*x+c))^2/(f*x+e)^2,x)`

output

```
( - atanh(c + d*x)**2*b**2*c**4*e*f**3 + 2*atanh(c + d*x)**2*b**2*c**3*d*e
**2*f**2 - atanh(c + d*x)**2*b**2*c**2*d**2*e**3*f - atanh(c + d*x)**2*b**
2*c**2*d**2*e**2*f**2*x + 2*atanh(c + d*x)**2*b**2*c**2*e*f**3 + 2*atanh(c
+ d*x)**2*b**2*c*d**3*e**3*f*x - 2*atanh(c + d*x)**2*b**2*c*d*e**2*f**2 -
atanh(c + d*x)**2*b**2*d**4*e**4*x + atanh(c + d*x)**2*b**2*d**2*e**3*f +
atanh(c + d*x)**2*b**2*d**2*e**2*f**2*x - atanh(c + d*x)**2*b**2*e*f**3 +
2*atanh(c + d*x)*a*b*c**4*f**4*x - 4*atanh(c + d*x)*a*b*c**3*d*e*f**3*x -
4*atanh(c + d*x)*a*b*c**2*f**4*x + 4*atanh(c + d*x)*a*b*c*d**3*e**3*f*x +
4*atanh(c + d*x)*a*b*c*d*e*f**3*x - 2*atanh(c + d*x)*a*b*d**4*e**4*x + 2*
atanh(c + d*x)*a*b*f**4*x - 2*atanh(c + d*x)*b**2*c**2*d*e*f**3*x + 4*atan
h(c + d*x)*b**2*c*d**2*e**2*f**2*x - 2*atanh(c + d*x)*b**2*d**3*e**3*f*x +
2*atanh(c + d*x)*b**2*d*e*f**3*x - 2*int((atanh(c + d*x)*x)/(c**4*e**2*f*
*2 + 2*c**4*e*f**3*x + c**4*f**4*x**2 + 2*c**3*d*e**2*f**2*x + 4*c**3*d*e*
f**3*x**2 + 2*c**3*d*f**4*x**3 - c**2*d**2*e**4 - 2*c**2*d**2*e**3*f*x + 2
*c**2*d**2*e*f**3*x**3 + c**2*d**2*f**4*x**4 - 2*c**2*e**2*f**2 - 4*c**2*e
*f**3*x - 2*c**2*f**4*x**2 - 2*c*d**3*e**4*x - 4*c*d**3*e**3*f*x**2 - 2*c*
d**3*e**2*f**2*x**3 - 2*c*d*e**2*f**2*x - 4*c*d*e*f**3*x**2 - 2*c*d*f**4*x
**3 - d**4*e**4*x**2 - 2*d**4*e**3*f*x**3 - d**4*e**2*f**2*x**4 + d**2*e**
4 + 2*d**2*e**3*f*x - 2*d**2*e*f**3*x**3 - d**2*f**4*x**4 + e**2*f**2 + 2*
e*f**3*x + f**4*x**2),x)*b**2*c**6*d*e**2*f**6 - 2*int((atanh(c + d*x)*...
```

$$3.44 \quad \int \frac{(a+b \operatorname{arctanh}(c+dx))^2}{(e+fx)^3} dx$$

| | |
|---|-----|
| Optimal result | 407 |
| Mathematica [C] (warning: unable to verify) | 408 |
| Rubi [A] (verified) | 409 |
| Maple [A] (verified) | 411 |
| Fricas [F] | 413 |
| Sympy [F] | 414 |
| Maxima [F] | 414 |
| Giac [F] | 415 |
| Mupad [F(-1)] | 415 |
| Reduce [F] | 415 |

Optimal result

Integrand size = 20, antiderivative size = 621

$$\begin{aligned}
& \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^3} dx \\
&= \frac{bd(a + b \operatorname{arctanh}(c + dx))}{(de + f - cf)(de - (1 + c)f)(e + fx)} - \frac{(a + b \operatorname{arctanh}(c + dx))^2}{2f(e + fx)^2} \\
&+ \frac{bd^2(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2}{1 - c - dx}\right)}{2f(de + f - cf)^2} \\
&+ \frac{b^2 d^2 \log(1 - c - dx)}{2(de + f - cf)^2(de - (1 + c)f)} - \frac{bd^2(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2}{1 + c + dx}\right)}{2f(de - f - cf)^2} \\
&+ \frac{2bd^2(de - cf)(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2}{1 + c + dx}\right)}{(de + f - cf)^2(de - (1 + c)f)^2} \\
&- \frac{b^2 d^2 \log(1 + c + dx)}{2(de + f - cf)(de - (1 + c)f)^2} + \frac{b^2 d^2 f \log(e + fx)}{(de + f - cf)^2(de - (1 + c)f)^2} \\
&- \frac{2bd^2(de - cf)(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2d(e + fx)}{(de + f - cf)(1 + c + dx)}\right)}{(de + f - cf)^2(de - (1 + c)f)^2} \\
&+ \frac{b^2 d^2 \operatorname{PolyLog}\left(2, -\frac{1 + c + dx}{1 - c - dx}\right)}{4f(de + f - cf)^2} + \frac{b^2 d^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 + c + dx}\right)}{4f(de - f - cf)^2} \\
&- \frac{b^2 d^2(de - cf) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 + c + dx}\right)}{(de + f - cf)^2(de - (1 + c)f)^2} \\
&+ \frac{b^2 d^2(de - cf) \operatorname{PolyLog}\left(2, 1 - \frac{2d(e + fx)}{(de + f - cf)(1 + c + dx)}\right)}{(de + f - cf)^2(de - (1 + c)f)^2}
\end{aligned}$$

output

```

b*d*(a+b*arctanh(d*x+c))/(-c*f+d*e+f)/(d*e-(1+c)*f)/(f*x+e)-1/2*(a+b*arctanh(d*x+c))^2/f/(f*x+e)^2+1/2*b*d^2*(a+b*arctanh(d*x+c))*ln(2/(-d*x-c+1))/f/(-c*f+d*e+f)^2+1/2*b^2*d^2*ln(-d*x-c+1)/(-c*f+d*e+f)^2/(d*e-(1+c)*f)-1/2*b*d^2*(a+b*arctanh(d*x+c))*ln(2/(d*x+c+1))/f/(-c*f+d*e-f)^2+2*b*d^2*(-c*f+d*e)*(a+b*arctanh(d*x+c))*ln(2/(d*x+c+1))/(-c*f+d*e+f)^2/(d*e-(1+c)*f)^2-1/2*b^2*d^2*ln(d*x+c+1)/(-c*f+d*e+f)/(d*e-(1+c)*f)^2+b^2*d^2*f*ln(f*x+e)/(-c*f+d*e+f)^2/(d*e-(1+c)*f)^2-2*b*d^2*(-c*f+d*e)*(a+b*arctanh(d*x+c))*ln(2*d*(f*x+e)/(-c*f+d*e+f)/(d*x+c+1))/(-c*f+d*e+f)^2/(d*e-(1+c)*f)^2+1/4*b^2*d^2*polylog(2,-(d*x+c+1)/(-d*x-c+1))/f/(-c*f+d*e+f)^2+1/4*b^2*d^2*polylog(2,1-2/(d*x+c+1))/f/(-c*f+d*e-f)^2-b^2*d^2*(-c*f+d*e)*polylog(2,1-2/(d*x+c+1))/(-c*f+d*e+f)^2/(d*e-(1+c)*f)^2+b^2*d^2*(-c*f+d*e)*polylog(2,1-2*d*(f*x+e)/(-c*f+d*e+f)/(d*x+c+1))/(-c*f+d*e+f)^2/(d*e-(1+c)*f)^2

```


Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 11.30 (sec) , antiderivative size = 1318, normalized size of antiderivative = 2.12

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^3} dx = \text{Too large to display}$$

input `Integrate[(a + b*ArcTanh[c + d*x])^2/(e + f*x)^3,x]`

output

```
-1/2*a^2/(f*(e + f*x)^2) + (a*b*(d*e - c*f + f*(c + d*x))^3*((f*(2 + ((d*e
+ f - c*f)*(d*e - (1 + c)*f))/((d*e - c*f)/Sqrt[1 - (c + d*x)^2] + (f*(c
+ d*x))/Sqrt[1 - (c + d*x)^2])^2)*ArcTanh[c + d*x])/((d*e + f - c*f)^2*(-(
d*e) + f + c*f)^2) - ((c + d*x)*(f - 2*d*e*ArcTanh[c + d*x] + 2*c*f*ArcTan
h[c + d*x]))/((d*e - c*f)*(d*e + f - c*f)*(d*e - (1 + c)*f)*Sqrt[1 - (c +
d*x)^2]*((d*e - c*f)/Sqrt[1 - (c + d*x)^2] + (f*(c + d*x))/Sqrt[1 - (c + d
*x)^2])) - (2*(d*e - c*f)*Log[(d*e)/Sqrt[1 - (c + d*x)^2] - (c*f)/Sqrt[1 -
(c + d*x)^2] + (f*(c + d*x))/Sqrt[1 - (c + d*x)^2]])/(d^2*e^2 - 2*c*d*e*f
+ (-1 + c^2)*f^2)^2)/(d*(e + f*x)^3) + (b^2*(d*e - c*f + f*(c + d*x))^3*
((d^2*(-(f^2*ArcTanh[c + d*x]) - I*d^2*e^2*Pi*ArcTanh[c + d*x] + (2*I)*c*d
*e*f*Pi*ArcTanh[c + d*x] - I*c^2*f^2*Pi*ArcTanh[c + d*x] + d*e*E^ArcTanh[c
- (d*e)/f]*Sqrt[1 - c^2 - (d^2*e^2)/f^2 + (2*c*d*e)/f]*f*ArcTanh[c + d*x]
^2 - c*E^ArcTanh[c - (d*e)/f]*Sqrt[1 - c^2 - (d^2*e^2)/f^2 + (2*c*d*e)/f]*
f^2*ArcTanh[c + d*x]^2 - 2*d^2*e^2*ArcTanh[c + d*x]*Log[1 - E^(2*ArcTanh[c
- (d*e)/f] - 2*ArcTanh[c + d*x]]) + 4*c*d*e*f*ArcTanh[c + d*x]*Log[1 - E
(2*ArcTanh[c - (d*e)/f] - 2*ArcTanh[c + d*x]]) - 2*c^2*f^2*ArcTanh[c + d*x
]*Log[1 - E^(2*ArcTanh[c - (d*e)/f] - 2*ArcTanh[c + d*x]]) + I*d^2*e^2*Pi*
Log[1 + E^(2*ArcTanh[c + d*x]]) - (2*I)*c*d*e*f*Pi*Log[1 + E^(2*ArcTanh[c
+ d*x]]) + I*c^2*f^2*Pi*Log[1 + E^(2*ArcTanh[c + d*x]]) - I*d^2*e^2*Pi*Log
[1/Sqrt[1 - (c + d*x)^2]] + (2*I)*c*d*e*f*Pi*Log[1/Sqrt[1 - (c + d*x)^2...)
```

Rubi [A] (verified)

Time = 2.19 (sec) , antiderivative size = 739, normalized size of antiderivative = 1.19, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {6659, 7292, 6671, 27, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + \operatorname{barctanh}(c + dx))^2}{(e + fx)^3} dx \\
 & \quad \downarrow \text{6659} \\
 & \frac{bd \int \frac{a + \operatorname{barctanh}(c + dx)}{(e + fx)^2(1 - (c + dx)^2)} dx}{f} - \frac{(a + \operatorname{barctanh}(c + dx))^2}{2f(e + fx)^2} \\
 & \quad \downarrow \text{7292} \\
 & \frac{bd \int \frac{a + \operatorname{barctanh}(c + dx)}{(e + fx)^2(-c^2 - 2dxc - d^2x^2 + 1)} dx}{f} - \frac{(a + \operatorname{barctanh}(c + dx))^2}{2f(e + fx)^2} \\
 & \quad \downarrow \text{6671} \\
 & \frac{b \int \frac{d^2(a + \operatorname{barctanh}(c + dx))}{\left(d\left(e - \frac{cf}{d}\right) + f(c + dx)\right)^2(1 - (c + dx)^2)} d(c + dx)}{f} - \frac{(a + \operatorname{barctanh}(c + dx))^2}{2f(e + fx)^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{bd^2 \int \frac{a + \operatorname{barctanh}(c + dx)}{(de - cf + f(c + dx))^2(1 - (c + dx)^2)} d(c + dx)}{f} - \frac{(a + \operatorname{barctanh}(c + dx))^2}{2f(e + fx)^2} \\
 & \quad \downarrow \text{7276} \\
 & \frac{bd^2 \int \left(-\frac{a}{(de - cf + f(c + dx))^2((c + dx)^2 - 1)} - \frac{\operatorname{barctanh}(c + dx)}{(de - cf + f(c + dx))^2((c + dx)^2 - 1)} \right) d(c + dx)}{f} - \\
 & \quad \frac{(a + \operatorname{barctanh}(c + dx))^2}{2f(e + fx)^2} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$bd^2 \left(\frac{af}{(-cf+de+f)(de-(c+1)f)(f(c+dx)-cf+de)} - \frac{2af(de-cf) \log(f(c+dx)-cf+de)}{(-cf+de+f)^2(de-(c+1)f)^2} - \frac{a \log(-c-dx+1)}{2(-cf+de+f)^2} + \frac{a \log(c+dx+1)}{2(de-(c+1)f)^2} + \frac{1}{(-cf+de+f)} \right)$$

$$\frac{(a + b \operatorname{arctanh}(c + dx))^2}{2f(e + fx)^2}$$

input `Int[(a + b*ArcTanh[c + d*x])^2/(e + f*x)^3,x]`

output

```
-1/2*(a + b*ArcTanh[c + d*x])^2/(f*(e + f*x)^2) + (b*d^2*((a*f)/((d*e + f
- c*f)*(d*e - (1 + c)*f)*(d*e - c*f + f*(c + d*x))) + (b*f*ArcTanh[c + d*x
])/((d*e + f - c*f)*(d*e - (1 + c)*f)*(d*e - c*f + f*(c + d*x))) + (b*ArcT
anh[c + d*x]*Log[2/(1 - c - d*x)])/(2*(d*e + f - c*f)^2) - (a*Log[1 - c -
d*x])/(2*(d*e + f - c*f)^2) + (b*f*Log[1 - c - d*x])/(2*(d*e + f - c*f)^2*
(d*e - (1 + c)*f)) - (b*ArcTanh[c + d*x]*Log[2/(1 + c + d*x)])/(2*(d*e - (
1 + c)*f)^2) + (2*b*f*(d*e - c*f)*ArcTanh[c + d*x]*Log[2/(1 + c + d*x)])/(
(d*e + f - c*f)^2*(d*e - (1 + c)*f)^2) + (a*Log[1 + c + d*x])/(2*(d*e - (1
+ c)*f)^2) - (b*f*Log[1 + c + d*x])/(2*(d*e + f - c*f)*(d*e - (1 + c)*f)^
2) + (b*f^2*Log[d*e - c*f + f*(c + d*x)])/((d*e + f - c*f)^2*(d*e - (1 + c
)*f)^2) - (2*a*f*(d*e - c*f)*Log[d*e - c*f + f*(c + d*x)])/((d*e + f - c*f
)^2*(d*e - (1 + c)*f)^2) - (2*b*f*(d*e - c*f)*ArcTanh[c + d*x]*Log[(2*(d*e
- c*f + f*(c + d*x))]/((d*e + f - c*f)*(1 + c + d*x)))]/((d*e + f - c*f)^
2*(d*e - (1 + c)*f)^2) + (b*PolyLog[2, -(1 + c + d*x)/(1 - c - d*x)])/(4
*(d*e + f - c*f)^2) + (b*PolyLog[2, 1 - 2/(1 + c + d*x)])/(4*(d*e - (1 + c
)*f)^2) - (b*f*(d*e - c*f)*PolyLog[2, 1 - 2/(1 + c + d*x)])/((d*e + f - c*
f)^2*(d*e - (1 + c)*f)^2) + (b*f*(d*e - c*f)*PolyLog[2, 1 - (2*(d*e - c*f
+ f*(c + d*x))]/((d*e + f - c*f)*(1 + c + d*x)))]/((d*e + f - c*f)^2*(d*e
- (1 + c)*f)^2))/f
```

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6659

```
Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_), x_Symbol] := Simp[(e + f*x)^(m + 1)*((a + b*ArcTanh[c + d*x])^p/(f*(m
+ 1))), x] - Simp[b*d*(p/(f*(m + 1))) Int[(e + f*x)^(m + 1)*((a + b*ArcTa
nh[c + d*x])^(p - 1)/(1 - (c + d*x)^2)), x], x] /; FreeQ[{a, b, c, d, e, f}
, x] && IGtQ[p, 0] && ILtQ[m, -1]
```

rule 6671

```
Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.)*((A_.) + (B_.)*(x_) + (C_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/d Sub
st[Int[((d*e - c*f)/d + f*(x/d))^m*(-C/d^2 + (C/d^2)*x^2)^q*(a + b*ArcTanh[
x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, p, q}, x
] && EqQ[B*(1 - c^2) + 2*A*c*d, 0] && EqQ[2*c*C - B*d, 0]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

rule 7292

```
Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v !=
= u]
```

Maple [A] (verified)

Time = 2.01 (sec) , antiderivative size = 870, normalized size of antiderivative = 1.40

| method | result |
|-------------------|---|
| parts | $-\frac{a^2}{2(fx+e)^2 f} + b^2 \left[-\frac{d^3 \operatorname{arctanh}(dx+c)^2}{2(f(dx+c)-cf+de)^2 f} + d^3 \left(\frac{\operatorname{arctanh}(dx+c)f}{(cf-de+f)(cf-de-f)(f(dx+c)-cf+de)} + \frac{2 \operatorname{arctanh}(dx+c)f^2 \ln(f(dx+c)-cf+de)}{(cf-de+f)^2(cf-de-f)} \right) \right]$ |
| derivativedivides | $-\frac{a^2 d^3}{2(cf-de-f(dx+c))^2 f} - b^2 d^3 \left(\frac{\operatorname{arctanh}(dx+c)^2}{2(cf-de-f(dx+c))^2 f} - \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{2(cf-de-f)^2} + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c+1)}{2(cf-de+f)^2} - \frac{1}{(cf-de-f)} \right)$ |
| default | $-\frac{a^2 d^3}{2(cf-de-f(dx+c))^2 f} - b^2 d^3 \left(\frac{\operatorname{arctanh}(dx+c)^2}{2(cf-de-f(dx+c))^2 f} - \frac{\operatorname{arctanh}(dx+c) \ln(dx+c-1)}{2(cf-de-f)^2} + \frac{\operatorname{arctanh}(dx+c) \ln(dx+c+1)}{2(cf-de+f)^2} - \frac{1}{(cf-de-f)} \right)$ |

input `int((a+b*arctanh(d*x+c))^2/(f*x+e)^3,x,method=_RETURNVERBOSE)`

output

```

-1/2*a^2/(f*x+e)^2/f+b^2/d*(-1/2*d^3/(f*(d*x+c)-c*f+d*e)^2/f*arctanh(d*x+c)
)^2+d^3/f*(arctanh(d*x+c)*f/(c*f-d*e+f)/(c*f-d*e-f)/(f*(d*x+c)-c*f+d*e)+2*
arctanh(d*x+c)*f^2/(c*f-d*e+f)^2/(c*f-d*e-f)^2*ln(f*(d*x+c)-c*f+d*e)*c-2*a
rctanh(d*x+c)*f/(c*f-d*e+f)^2/(c*f-d*e-f)^2*ln(f*(d*x+c)-c*f+d*e)*d*e-1/2*
arctanh(d*x+c)/(c*f-d*e-f)^2*ln(d*x+c-1)+1/2*arctanh(d*x+c)/(c*f-d*e+f)^2*
ln(d*x+c+1)-1/2/(c*f-d*e-f)^2*(-1/2*dilog(1/2*d*x+1/2*c+1/2)-1/2*ln(d*x+c-
1)*ln(1/2*d*x+1/2*c+1/2)+1/4*ln(d*x+c-1)^2)+1/2/(c*f-d*e+f)^2*(-1/4*ln(d*x
+c+1)^2+1/2*(ln(d*x+c+1)-ln(1/2*d*x+1/2*c+1/2))*ln(-1/2*d*x-1/2*c+1/2)-1/2
*dilog(1/2*d*x+1/2*c+1/2))+f/(c*f-d*e+f)/(c*f-d*e-f)*(f/(c*f-d*e+f)/(c*f-d
*e-f)*ln(f*(d*x+c)-c*f+d*e)-1/(2*c*f-2*d*e-2*f)*ln(d*x+c-1)+1/(2*c*f-2*d*e
+2*f)*ln(d*x+c+1))+2*(c*f-d*e)/(c*f-d*e+f)^2/(c*f-d*e-f)^2*(1/2*f*(dilog((
f*(d*x+c)-f)/(c*f-d*e-f))+ln(f*(d*x+c)-c*f+d*e)*ln((f*(d*x+c)-f)/(c*f-d*e-
f)))-1/2*f*(dilog((f*(d*x+c)+f)/(c*f-d*e+f))+ln(f*(d*x+c)-c*f+d*e)*ln((f*(
d*x+c)+f)/(c*f-d*e+f)))))+2*b*a/d*(-1/2*d^3/(f*(d*x+c)-c*f+d*e)^2/f*arcta
nh(d*x+c)+1/2*d^3/f*(f/(c*f-d*e+f)/(c*f-d*e-f)/(f*(d*x+c)-c*f+d*e)+2*f*(c*
f-d*e)/(c*f-d*e+f)^2/(c*f-d*e-f)^2*ln(f*(d*x+c)-c*f+d*e)-1/2/(c*f-d*e-f)^2
*ln(d*x+c-1)+1/2/(c*f-d*e+f)^2*ln(d*x+c+1)))

```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^3} dx = \int \frac{(b \operatorname{arctanh}(dx + c) + a)^2}{(fx + e)^3} dx$$

input

```
integrate((a+b*arctanh(d*x+c))^2/(f*x+e)^3,x, algorithm="fricas")
```

output

```
integral((b^2*arctanh(d*x + c)^2 + 2*a*b*arctanh(d*x + c) + a^2)/(f^3*x^3
+ 3*e*f^2*x^2 + 3*e^2*f*x + e^3), x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^3} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^2}{(e + fx)^3} dx$$

input `integrate((a+b*atanh(d*x+c))**2/(f*x+e)**3,x)`

output `Integral((a + b*atanh(c + d*x))**2/(e + f*x)**3, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^3} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^2}{(fx + e)^3} dx$$

input `integrate((a+b*arctanh(d*x+c))^2/(f*x+e)^3,x, algorithm="maxima")`

output `1/2*(d*(d*log(d*x + c + 1)/(d^2*e^2*f - 2*(c + 1)*d*e*f^2 + (c^2 + 2*c + 1)*f^3) - d*log(d*x + c - 1)/(d^2*e^2*f - 2*(c - 1)*d*e*f^2 + (c^2 - 2*c + 1)*f^3) - 4*(d^2*e - c*d*f)*log(f*x + e)/(d^4*e^4 - 4*c*d^3*e^3*f + 2*(3*c^2 - 1)*d^2*e^2*f^2 - 4*(c^3 - c)*d*e*f^3 + (c^4 - 2*c^2 + 1)*f^4) + 2/(d^2*e^3 - 2*c*d*e^2*f + (c^2 - 1)*e*f^2 + (d^2*e^2*f - 2*c*d*e*f^2 + (c^2 - 1)*f^3)*x)) - 2*arctanh(d*x + c)/(f^3*x^2 + 2*e*f^2*x + e^2*f))*a*b - 1/8*b^2*(log(-d*x - c + 1)^2/(f^3*x^2 + 2*e*f^2*x + e^2*f) + 2*integrate(-((d*f*x + c*f - f)*log(d*x + c + 1)^2 + (d*f*x + d*e - 2*(d*f*x + c*f - f)*log(d*x + c + 1))*log(-d*x - c + 1))/(d*f^4*x^4 + c*e^3*f - e^3*f + (3*d*e*f^3 + c*f^4 - f^4)*x^3 + 3*(d*e^2*f^2 + c*e*f^3 - e*f^3)*x^2 + (d*e^3*f + 3*c*e^2*f^2 - 3*e^2*f^2)*x), x)) - 1/2*a^2/(f^3*x^2 + 2*e*f^2*x + e^2*f)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^3} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^2}{(fx + e)^3} dx$$

input `integrate((a+b*arctanh(d*x+c))^2/(f*x+e)^3,x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)^2/(f*x + e)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^3} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^2}{(e + fx)^3} dx$$

input `int((a + b*atanh(c + d*x))^2/(e + f*x)^3,x)`

output `int((a + b*atanh(c + d*x))^2/(e + f*x)^3, x)`

Reduce [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)^3} dx = \text{too large to display}$$

input `int((a+b*atanh(d*x+c))^2/(f*x+e)^3,x)`

output

```
( - 12*atanh(c + d*x)**2*b**2*c**9*e*f**9 + 90*atanh(c + d*x)**2*b**2*c**8
*d**e**2*f**8 - 288*atanh(c + d*x)**2*b**2*c**7*d**2*e**3*f**7 + 32*atanh(c
+ d*x)**2*b**2*c**7*e*f**9 + 510*atanh(c + d*x)**2*b**2*c**6*d**3*e**4*f*
*6 + 12*atanh(c + d*x)**2*b**2*c**6*d**3*e**3*f**7*x + 6*atanh(c + d*x)**2
*b**2*c**6*d**3*e**2*f**8*x**2 - 200*atanh(c + d*x)**2*b**2*c**6*d**e**2*f*
*8 - 540*atanh(c + d*x)**2*b**2*c**5*d**4*e**5*f**5 - 72*atanh(c + d*x)**2
*b**2*c**5*d**4*e**4*f**6*x - 36*atanh(c + d*x)**2*b**2*c**5*d**4*e**3*f**
7*x**2 + 528*atanh(c + d*x)**2*b**2*c**5*d**2*e**3*f**7 - 24*atanh(c + d*x
)**2*b**2*c**5*e*f**9 + 342*atanh(c + d*x)**2*b**2*c**4*d**5*e**6*f**4 + 1
80*atanh(c + d*x)**2*b**2*c**4*d**5*e**5*f**5*x + 90*atanh(c + d*x)**2*b**
2*c**4*d**5*e**4*f**6*x**2 - 770*atanh(c + d*x)**2*b**2*c**4*d**3*e**4*f**
6 - 20*atanh(c + d*x)**2*b**2*c**4*d**3*e**3*f**7*x - 10*atanh(c + d*x)**2
*b**2*c**4*d**3*e**2*f**8*x**2 + 124*atanh(c + d*x)**2*b**2*c**4*d**e**2*f*
*8 - 120*atanh(c + d*x)**2*b**2*c**3*d**6*e**7*f**3 - 240*atanh(c + d*x)**
2*b**2*c**3*d**6*e**6*f**4*x - 120*atanh(c + d*x)**2*b**2*c**3*d**6*e**5*f
**5*x**2 + 680*atanh(c + d*x)**2*b**2*c**3*d**4*e**5*f**5 + 80*atanh(c + d
*x)**2*b**2*c**3*d**4*e**4*f**6*x + 40*atanh(c + d*x)**2*b**2*c**3*d**4*e
**3*f**7*x**2 - 256*atanh(c + d*x)**2*b**2*c**3*d**2*e**3*f**7 + 18*atanh(c
+ d*x)**2*b**2*c**2*d**7*e**8*f**2 + 180*atanh(c + d*x)**2*b**2*c**2*d**7
*e**7*f**3*x + 90*atanh(c + d*x)**2*b**2*c**2*d**7*e**6*f**4*x**2 - 372...
```

3.45 $\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx$

| | |
|---|-----|
| Optimal result | 418 |
| Mathematica [B] (warning: unable to verify) | 419 |
| Rubi [A] (verified) | 420 |
| Maple [C] (warning: unable to verify) | 422 |
| Fricas [F] | 423 |
| Sympy [F] | 423 |
| Maxima [F] | 423 |
| Giac [F] | 424 |
| Mupad [F(-1)] | 425 |
| Reduce [F] | 425 |

Optimal result

Integrand size = 20, antiderivative size = 546

$$\begin{aligned}
& \int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx \\
&= \frac{ab^2 f^2 x}{d^2} + \frac{b^3 f^2 (c + dx) \operatorname{arctanh}(c + dx)}{d^3} - \frac{bf^2 (a + b \operatorname{arctanh}(c + dx))^2}{2d^3} \\
&+ \frac{3bf(de - cf)(a + b \operatorname{arctanh}(c + dx))^2}{d^3} \\
&+ \frac{3bf(de - cf)(c + dx)(a + b \operatorname{arctanh}(c + dx))^2}{d^3} \\
&+ \frac{bf^2(c + dx)^2(a + b \operatorname{arctanh}(c + dx))^2}{2d^3} \\
&- \frac{(de - cf)(d^2 e^2 - 2cdef + (3 + c^2)f^2)(a + b \operatorname{arctanh}(c + dx))^3}{3d^3 f} \\
&+ \frac{(3d^2 e^2 - 6cdef + (1 + 3c^2)f^2)(a + b \operatorname{arctanh}(c + dx))^3}{3d^3} \\
&+ \frac{(e + fx)^3 (a + b \operatorname{arctanh}(c + dx))^3}{3f} \\
&- \frac{6b^2 f(de - cf)(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2}{1 - c - dx}\right)}{d^3} \\
&- \frac{b(3d^2 e^2 - 6cdef + (1 + 3c^2)f^2)(a + b \operatorname{arctanh}(c + dx))^2 \log\left(\frac{2}{1 - c - dx}\right)}{d^3} \\
&+ \frac{b^3 f^2 \log(1 - (c + dx)^2)}{2d^3} - \frac{3b^3 f(de - cf) \operatorname{PolyLog}\left(2, -\frac{1 + c + dx}{1 - c - dx}\right)}{d^3} \\
&- \frac{b^2(3d^2 e^2 - 6cdef + (1 + 3c^2)f^2)(a + b \operatorname{arctanh}(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - c - dx}\right)}{d^3} \\
&+ \frac{b^3(3d^2 e^2 - 6cdef + (1 + 3c^2)f^2) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - c - dx}\right)}{2d^3}
\end{aligned}$$

output

```
a*b^2*f^2*x/d^2+b^3*f^2*(d*x+c)*arctanh(d*x+c)/d^3-1/2*b*f^2*(a+b*arctanh(
d*x+c))^2/d^3+3*b*f*(-c*f+d*e)*(a+b*arctanh(d*x+c))^2/d^3+3*b*f*(-c*f+d*e)
*(d*x+c)*(a+b*arctanh(d*x+c))^2/d^3+1/2*b*f^2*(d*x+c)^2*(a+b*arctanh(d*x+c
))^2/d^3-1/3*(-c*f+d*e)*(d^2*e^2-2*c*d*e*f+(c^2+3)*f^2)*(a+b*arctanh(d*x+c
))^3/d^3/f+1/3*(3*d^2*e^2-6*c*d*e*f+(3*c^2+1)*f^2)*(a+b*arctanh(d*x+c))^3/
d^3+1/3*(f*x+e)^3*(a+b*arctanh(d*x+c))^3/f-6*b^2*f*(-c*f+d*e)*(a+b*arctanh
(d*x+c))*ln(2/(-d*x-c+1))/d^3-b*(3*d^2*e^2-6*c*d*e*f+(3*c^2+1)*f^2)*(a+b*a
rctanh(d*x+c))^2*ln(2/(-d*x-c+1))/d^3+1/2*b^3*f^2*ln(1-(d*x+c)^2)/d^3-3*b^
3*f*(-c*f+d*e)*polylog(2,-(d*x+c+1)/(-d*x-c+1))/d^3-b^2*(3*d^2*e^2-6*c*d*e
*f+(3*c^2+1)*f^2)*(a+b*arctanh(d*x+c))*polylog(2,1-2/(-d*x-c+1))/d^3+1/2*b
^3*(3*d^2*e^2-6*c*d*e*f+(3*c^2+1)*f^2)*polylog(3,1-2/(-d*x-c+1))/d^3
```

Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 1646 vs. $2(546) = 1092$.

Time = 6.61 (sec) , antiderivative size = 1646, normalized size of antiderivative = 3.01

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx = \text{Too large to display}$$

input

```
Integrate[(e + f*x)^2*(a + b*ArcTanh[c + d*x])^3,x]
```

output

```

a^2*(a*e^2 + (b*f*(3*d*e - 2*c*f))/d^2)*x + (a^2*f*(2*a*d*e + b*f)*x^2)/(2
*d) + (a^3*f^2*x^3)/3 + a^2*b*x*(3*e^2 + 3*e*f*x + f^2*x^2)*ArcTanh[c + d*
x] - (a^2*b*(-1 + c)*(3*d^2*e^2 - 3*(-1 + c)*d*e*f + (-1 + c)^2*f^2)*Log[1
- c - d*x])/(2*d^3) + (a^2*b*(1 + c)*(3*d^2*e^2 - 3*(1 + c)*d*e*f + (1 +
c)^2*f^2)*Log[1 + c + d*x])/(2*d^3) + (3*a*b^2*e^2*(ArcTanh[c + d*x]*((-1
+ c + d*x)*ArcTanh[c + d*x] - 2*Log[1 + E^(-2*ArcTanh[c + d*x])]) + PolyLo
g[2, -E^(-2*ArcTanh[c + d*x])]))/d - (3*a*b^2*e*f*((1 - 2*c + c^2 - d^2*x^
2)*ArcTanh[c + d*x]^2 - 2*ArcTanh[c + d*x]*(c + d*x + 2*c*Log[1 + E^(-2*Ar
cTanh[c + d*x])]) + 2*Log[1/Sqrt[1 - (c + d*x)^2]] + 2*c*PolyLog[2, -E^(-2
*ArcTanh[c + d*x])]))/d^2 + (b^3*e^2*(2*ArcTanh[c + d*x]^2*((-1 + c + d*x)
*ArcTanh[c + d*x] - 3*Log[1 + E^(-2*ArcTanh[c + d*x])]) + 6*ArcTanh[c + d*
x]*PolyLog[2, -E^(-2*ArcTanh[c + d*x])]) + 3*PolyLog[3, -E^(-2*ArcTanh[c +
d*x])]))/(2*d) - (b^3*e*f*(ArcTanh[c + d*x]*((1 - 2*c + c^2 - d^2*x^2)*Arc
Tanh[c + d*x]^2 + 6*Log[1 + E^(-2*ArcTanh[c + d*x])]) - 3*ArcTanh[c + d*x]*
(-1 + c + d*x + 2*c*Log[1 + E^(-2*ArcTanh[c + d*x])])) + (-3 + 6*c*ArcTanh
[c + d*x])*PolyLog[2, -E^(-2*ArcTanh[c + d*x])]) + 3*c*PolyLog[3, -E^(-2*Ar
cTanh[c + d*x])]))/d^2 - (a*b^2*f^2*(1 - (c + d*x)^2)^(3/2)*(-(c + d*x)/S
qrt[1 - (c + d*x)^2]) + (6*c*(c + d*x)*ArcTanh[c + d*x])/Sqrt[1 - (c + d*x
)^2] + (3*(c + d*x)*ArcTanh[c + d*x]^2)/Sqrt[1 - (c + d*x)^2] - (3*c^2*(c
+ d*x)*ArcTanh[c + d*x]^2)/Sqrt[1 - (c + d*x)^2] + ArcTanh[c + d*x]^2*C...

```

Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 533, normalized size of antiderivative = 0.98, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {6661, 27, 6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx$$

$$\downarrow 6661$$

$$\int \frac{\left(d\left(e - \frac{ef}{d}\right) + f(c + dx)\right)^2 (a + b \operatorname{arctanh}(c + dx))^3}{d^2} d(c + dx)$$

$$\downarrow 27$$

$$\frac{\int (de - cf + f(c + dx))^2 (a + \operatorname{barctanh}(c + dx))^3 d(c + dx)}{d^3}$$

↓ 6480

$$\frac{\frac{(f(c+dx)-cf+de)^3(a+\operatorname{barctanh}(c+dx))^3}{3f} - \frac{bf\left(-((c+dx)(a+\operatorname{barctanh}(c+dx))^2f^3) - 3(de-cf)(a+\operatorname{barctanh}(c+dx))^2f^2 + \frac{(de-cf)(d^2e^2 - 2cde f + (1+3c^2)f^2)}{f}\right)}{d^3}}{d^3}$$

↓ 2009

$$\frac{\frac{(f(c+dx)-cf+de)^3(a+\operatorname{barctanh}(c+dx))^3}{3f} - \frac{b\left(bf((3c^2+1)f^2 - 6cde f + 3d^2e^2) \operatorname{PolyLog}\left(2, 1 - \frac{2}{-c-dx+1}\right) (a+\operatorname{barctanh}(c+dx)) - \frac{f((3e^2+1)d^2e^2 - 2cde f + (1+3c^2)f^2)}{f}\right)}{d^3}}{d^3}$$

input

```
Int[(e + f*x)^2*(a + b*ArcTanh[c + d*x])^3,x]
```

output

```
((d*e - c*f + f*(c + d*x))^3*(a + b*ArcTanh[c + d*x])^3)/(3*f) - (b*(-(a*b*f^3*(c + d*x)) - b^2*f^3*(c + d*x)*ArcTanh[c + d*x] + (f^3*(a + b*ArcTanh[c + d*x])^2)/2 - 3*f^2*(d*e - c*f)*(a + b*ArcTanh[c + d*x])^2 - 3*f^2*(d*e - c*f)*(c + d*x)*(a + b*ArcTanh[c + d*x])^2 - (f^3*(c + d*x)^2*(a + b*ArcTanh[c + d*x])^2)/2 + ((d*e - c*f)*(d^2*e^2 - 2*c*d*e*f + (3 + c^2)*f^2)*(a + b*ArcTanh[c + d*x])^3)/(3*b) - (f*(3*d^2*e^2 - 6*c*d*e*f + (1 + 3*c^2)*f^2)*(a + b*ArcTanh[c + d*x])^3)/(3*b) + 6*b*f^2*(d*e - c*f)*(a + b*ArcTanh[c + d*x])*Log[2/(1 - c - d*x)] + f*(3*d^2*e^2 - 6*c*d*e*f + (1 + 3*c^2)*f^2)*(a + b*ArcTanh[c + d*x])^2*Log[2/(1 - c - d*x)] - (b^2*f^3*Log[1 - (c + d*x)^2])/2 + 3*b^2*f^2*(d*e - c*f)*PolyLog[2, -((1 + c + d*x)/(1 - c - d*x))] + b*f*(3*d^2*e^2 - 6*c*d*e*f + (1 + 3*c^2)*f^2)*(a + b*ArcTanh[c + d*x])*PolyLog[2, 1 - 2/(1 - c - d*x)] - (b^2*f*(3*d^2*e^2 - 6*c*d*e*f + (1 + 3*c^2)*f^2)*PolyLog[3, 1 - 2/(1 - c - d*x)]/2)/f)/d^3
```

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6480 `Int[((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)*((d_.) + (e_.)*(x_.))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])^p/(e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTanh[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 - c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]`

rule 6661 `Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[p, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 42.68 (sec) , antiderivative size = 10013, normalized size of antiderivative = 18.34

| method | result | size |
|-------------------|---------------------------------|-------|
| derivativedivides | Expression too large to display | 10013 |
| default | Expression too large to display | 10013 |
| parts | Expression too large to display | 10025 |

input `int((f*x+e)^2*(a+b*arctanh(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F]

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx = \int (fx + e)^2 (b \operatorname{artanh}(dx + c) + a)^3 dx$$

input `integrate((f*x+e)^2*(a+b*arctanh(d*x+c))^3,x, algorithm="fricas")`

output `integral(a^3*f^2*x^2 + 2*a^3*e*f*x + a^3*e^2 + (b^3*f^2*x^2 + 2*b^3*e*f*x + b^3*e^2)*arctanh(d*x + c)^3 + 3*(a*b^2*f^2*x^2 + 2*a*b^2*e*f*x + a*b^2*e^2)*arctanh(d*x + c)^2 + 3*(a^2*b*f^2*x^2 + 2*a^2*b*e*f*x + a^2*b*e^2)*arctanh(d*x + c), x)`

Sympy [F]

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx = \int (a + b \operatorname{atanh}(c + dx))^3 (e + fx)^2 dx$$

input `integrate((f*x+e)**2*(a+b*atanh(d*x+c))**3,x)`

output `Integral((a + b*atanh(c + d*x))**3*(e + f*x)**2, x)`

Maxima [F]

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx = \int (fx + e)^2 (b \operatorname{artanh}(dx + c) + a)^3 dx$$

input `integrate((f*x+e)^2*(a+b*arctanh(d*x+c))^3,x, algorithm="maxima")`

output

```

1/3*a^3*f^2*x^3 + a^3*e*f*x^2 + 3/2*(2*x^2*arctanh(d*x + c) + d*(2*x/d^2 -
(c^2 + 2*c + 1)*log(d*x + c + 1)/d^3 + (c^2 - 2*c + 1)*log(d*x + c - 1)/d
^3))*a^2*b*e*f + 1/2*(2*x^3*arctanh(d*x + c) + d*((d*x^2 - 4*c*x)/d^3 + (c
^3 + 3*c^2 + 3*c + 1)*log(d*x + c + 1)/d^4 - (c^3 - 3*c^2 + 3*c - 1)*log(d
*x + c - 1)/d^4))*a^2*b*f^2 + a^3*e^2*x + 3/2*(2*(d*x + c)*arctanh(d*x + c
) + log(-(d*x + c)^2 + 1))*a^2*b*e^2/d - 1/24*((b^3*d^3*f^2*x^3 + 3*b^3*d
^3*e*f*x^2 + 3*b^3*d^3*e^2*x + (c^3*f^2 - 3*d^2*e^2 - 3*(d*e*f + f^2)*c^2 -
3*d*e*f + 3*(d^2*e^2 + 2*d*e*f + f^2)*c - f^2)*b^3)*log(-d*x - c + 1)^3 -
3*(2*a*b^2*d^3*f^2*x^3 + (6*a*b^2*d^3*e*f + b^3*d^2*f^2)*x^2 + 2*(3*a*b^2
*d^3*e^2 + (3*d^2*e*f - 2*c*d*f^2)*b^3)*x + (b^3*d^3*f^2*x^3 + 3*b^3*d^3*e
*f*x^2 + 3*b^3*d^3*e^2*x + (c^3*f^2 + 3*d^2*e^2 - 3*(d*e*f - f^2)*c^2 - 3*
d*e*f + 3*(d^2*e^2 - 2*d*e*f + f^2)*c + f^2)*b^3)*log(d*x + c + 1))*log(-d
*x - c + 1)^2)/d^3 - integrate(-1/8*((b^3*d^3*f^2*x^3 + (2*d^3*e*f + c*d^2
*f^2 - d^2*f^2)*b^3*x^2 + (d^3*e^2 + 2*c*d^2*e*f - 2*d^2*e*f)*b^3*x + (c*d
^2*e^2 - d^2*e^2)*b^3)*log(d*x + c + 1)^3 + 6*(a*b^2*d^3*f^2*x^3 + (2*d^3*
e*f + c*d^2*f^2 - d^2*f^2)*a*b^2*x^2 + (d^3*e^2 + 2*c*d^2*e*f - 2*d^2*e*f)
*a*b^2*x + (c*d^2*e^2 - d^2*e^2)*a*b^2)*log(d*x + c + 1)^2 - (4*a*b^2*d^3*
f^2*x^3 + 2*(6*a*b^2*d^3*e*f + b^3*d^2*f^2)*x^2 + 3*(b^3*d^3*f^2*x^3 + (2*
d^3*e*f + c*d^2*f^2 - d^2*f^2)*b^3*x^2 + (d^3*e^2 + 2*c*d^2*e*f - 2*d^2*e*
f)*b^3*x + (c*d^2*e^2 - d^2*e^2)*b^3)*log(d*x + c + 1)^2 + 4*(3*a*b^2*d...

```

Giac [F]

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx = \int (fx + e)^2 (b \operatorname{arctanh}(dx + c) + a)^3 dx$$

input

```
integrate((f*x+e)^2*(a+b*arctanh(d*x+c))^3,x, algorithm="giac")
```

output

```
integrate((f*x + e)^2*(b*arctanh(d*x + c) + a)^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx = \int (e + fx)^2 (a + b \operatorname{atanh}(c + dx))^3 dx$$

input `int((e + f*x)^2*(a + b*atanh(c + d*x))^3,x)`output `int((e + f*x)^2*(a + b*atanh(c + d*x))^3, x)`**Reduce [F]**

$$\int (e + fx)^2 (a + b \operatorname{arctanh}(c + dx))^3 dx = \text{Too large to display}$$

input `int((f*x+e)^2*(a+b*atanh(d*x+c))^3,x)`

output

```
( - 4*atanh(c + d*x)**3*b**3*c**3*f**2 + 6*atanh(c + d*x)**3*b**3*c**2*d*e
*f + 4*atanh(c + d*x)**3*b**3*c*f**2 + 6*atanh(c + d*x)**3*b**3*d**3*e**2*
x + 6*atanh(c + d*x)**3*b**3*d**3*e*f*x**2 + 2*atanh(c + d*x)**3*b**3*d**3
*f**2*x**3 - 6*atanh(c + d*x)**3*b**3*d*e*f - 12*atanh(c + d*x)**2*a*b**2*
c**3*f**2 + 18*atanh(c + d*x)**2*a*b**2*c**2*d*e*f + 12*atanh(c + d*x)**2*
a*b**2*c*f**2 + 18*atanh(c + d*x)**2*a*b**2*d**3*e**2*x + 18*atanh(c + d*x
)**2*a*b**2*d**3*e*f*x**2 + 6*atanh(c + d*x)**2*a*b**2*d**3*f**2*x**3 - 18
*atanh(c + d*x)**2*a*b**2*d*e*f + 3*atanh(c + d*x)**2*b**3*c**2*f**2 - 12*
atanh(c + d*x)**2*b**3*c*d*f**2*x + 18*atanh(c + d*x)**2*b**3*d**2*e*f*x +
3*atanh(c + d*x)**2*b**3*d**2*f**2*x**2 - 3*atanh(c + d*x)**2*b**3*f**2 +
6*atanh(c + d*x)*a**2*b*c**3*f**2 - 18*atanh(c + d*x)*a**2*b*c**2*d*e*f +
18*atanh(c + d*x)*a**2*b*c**2*f**2 + 18*atanh(c + d*x)*a**2*b*c*d**2*e**2
- 36*atanh(c + d*x)*a**2*b*c*d*e*f + 18*atanh(c + d*x)*a**2*b*c*f**2 + 18
*atanh(c + d*x)*a**2*b*d**3*e**2*x + 18*atanh(c + d*x)*a**2*b*d**3*e*f*x**
2 + 6*atanh(c + d*x)*a**2*b*d**3*f**2*x**3 + 18*atanh(c + d*x)*a**2*b*d**2
*e**2 - 18*atanh(c + d*x)*a**2*b*d*e*f + 6*atanh(c + d*x)*a**2*b*f**2 - 30
*atanh(c + d*x)*a*b**2*c**2*f**2 + 36*atanh(c + d*x)*a*b**2*c*d*e*f - 24*a
tanh(c + d*x)*a*b**2*c*d*f**2*x - 36*atanh(c + d*x)*a*b**2*c*f**2 + 36*ata
nh(c + d*x)*a*b**2*d**2*e*f*x + 6*atanh(c + d*x)*a*b**2*d**2*f**2*x**2 + 3
6*atanh(c + d*x)*a*b**2*d*e*f - 6*atanh(c + d*x)*a*b**2*f**2 + 6*atanh(...
```

3.46 $\int (e + fx)(a + \operatorname{barctanh}(c + dx))^3 dx$

| | |
|---------------------------------------|-----|
| Optimal result | 427 |
| Mathematica [A] (verified) | 428 |
| Rubi [A] (verified) | 429 |
| Maple [C] (warning: unable to verify) | 431 |
| Fricas [F] | 431 |
| Sympy [F] | 432 |
| Maxima [F] | 432 |
| Giac [F] | 433 |
| Mupad [F(-1)] | 433 |
| Reduce [F] | 433 |

Optimal result

Integrand size = 18, antiderivative size = 326

$$\begin{aligned}
 & \int (e + fx)(a + \operatorname{barctanh}(c + dx))^3 dx \\
 &= \frac{3bf(a + \operatorname{barctanh}(c + dx))^2}{2d^2} + \frac{3bf(c + dx)(a + \operatorname{barctanh}(c + dx))^2}{2d^2} \\
 &+ \frac{(de - cf)(a + \operatorname{barctanh}(c + dx))^3}{d^2} \\
 &- \frac{(d^2e^2 - 2cdef + (1 + c^2)f^2)(a + \operatorname{barctanh}(c + dx))^3}{2d^2f} \\
 &+ \frac{(e + fx)^2(a + \operatorname{barctanh}(c + dx))^3}{2f} - \frac{3b^2f(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1-c-dx}\right)}{d^2} \\
 &- \frac{3b(de - cf)(a + \operatorname{barctanh}(c + dx))^2 \log\left(\frac{2}{1-c-dx}\right)}{d^2} - \frac{3b^3f \operatorname{PolyLog}\left(2, -\frac{1+c+dx}{1-c-dx}\right)}{2d^2} \\
 &- \frac{3b^2(de - cf)(a + \operatorname{barctanh}(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-c-dx}\right)}{d^2} \\
 &+ \frac{3b^3(de - cf) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-c-dx}\right)}{2d^2}
 \end{aligned}$$

output

```

3/2*b*f*(a+b*arctanh(d*x+c))^2/d^2+3/2*b*f*(d*x+c)*(a+b*arctanh(d*x+c))^2/
d^2+(-c*f+d*e)*(a+b*arctanh(d*x+c))^3/d^2-1/2*(d^2*e^2-2*c*d*e*f+(c^2+1)*f
^2)*(a+b*arctanh(d*x+c))^3/d^2/f+1/2*(f*x+e)^2*(a+b*arctanh(d*x+c))^3/f-3*
b^2*f*(a+b*arctanh(d*x+c))*ln(2/(-d*x-c+1))/d^2-3*b*(-c*f+d*e)*(a+b*arctan
h(d*x+c))^2*ln(2/(-d*x-c+1))/d^2-3/2*b^3*f*polylog(2,-(d*x+c+1)/(-d*x-c+1)
)/d^2-3*b^2*(-c*f+d*e)*(a+b*arctanh(d*x+c))*polylog(2,1-2/(-d*x-c+1))/d^2+
3/2*b^3*(-c*f+d*e)*polylog(3,1-2/(-d*x-c+1))/d^2

```

Mathematica [A] (verified)

Time = 3.19 (sec) , antiderivative size = 566, normalized size of antiderivative = 1.74

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^3 dx$$

$$= \frac{2a^2(2ade + 3bf - 2acf)(c + dx) + 2a^3f(c + dx)^2 - 6a^2b(c + dx)(cf - d(2e + fx))\operatorname{arctanh}(c + dx) + 3b^2f(c + dx)^2 \operatorname{arctanh}(c + dx) + 3b^3 \operatorname{arctanh}(c + dx)^3}{4d^2}$$

input

```
Integrate[(e + f*x)*(a + b*ArcTanh[c + d*x])^3,x]
```

output

```

(2*a^2*(2*a*d*e + 3*b*f - 2*a*c*f)*(c + d*x) + 2*a^3*f*(c + d*x)^2 - 6*a^2
*b*(c + d*x)*(c*f - d*(2*e + f*x))*ArcTanh[c + d*x] + 3*a^2*b*(2*d*e + f -
2*c*f)*Log[1 - c - d*x] + 3*a^2*b*(2*d*e - (1 + 2*c)*f)*Log[1 + c + d*x]
+ 12*a*b^2*f*((c + d*x)*ArcTanh[c + d*x] - ((1 - (c + d*x)^2)*ArcTanh[c +
d*x]^2)/2 - Log[1/Sqrt[1 - (c + d*x)^2]]) + 12*a*b^2*d*e*(ArcTanh[c + d*x]
*((-1 + c + d*x)*ArcTanh[c + d*x] - 2*Log[1 + E^(-2*ArcTanh[c + d*x])]) +
PolyLog[2, -E^(-2*ArcTanh[c + d*x])]) - 12*a*b^2*c*f*(ArcTanh[c + d*x]*((-
1 + c + d*x)*ArcTanh[c + d*x] - 2*Log[1 + E^(-2*ArcTanh[c + d*x])]) + Poly
Log[2, -E^(-2*ArcTanh[c + d*x])]) + 2*b^3*f*(ArcTanh[c + d*x]*(3*(-1 + c +
d*x)*ArcTanh[c + d*x] + (-1 + c^2 + 2*c*d*x + d^2*x^2)*ArcTanh[c + d*x]^2
- 6*Log[1 + E^(-2*ArcTanh[c + d*x])]) + 3*PolyLog[2, -E^(-2*ArcTanh[c + d
*x])]) + 4*b^3*d*e*(ArcTanh[c + d*x]^2*((-1 + c + d*x)*ArcTanh[c + d*x] -
3*Log[1 + E^(-2*ArcTanh[c + d*x])]) + 3*ArcTanh[c + d*x]*PolyLog[2, -E^(-2
*ArcTanh[c + d*x])]) + (3*PolyLog[3, -E^(-2*ArcTanh[c + d*x])])/2) - 4*b^3*
c*f*(ArcTanh[c + d*x]^2*((-1 + c + d*x)*ArcTanh[c + d*x] - 3*Log[1 + E^(-2
*ArcTanh[c + d*x])]) + 3*ArcTanh[c + d*x]*PolyLog[2, -E^(-2*ArcTanh[c + d
*x])]) + (3*PolyLog[3, -E^(-2*ArcTanh[c + d*x])])/2))/(4*d^2)

```

Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 324, normalized size of antiderivative = 0.99, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {6661, 27, 6480, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (e + fx)(a + b \operatorname{arctanh}(c + dx))^3 dx \\
 & \quad \downarrow \text{6661} \\
 & \int \frac{(d(e - \frac{cf}{d}) + f(c + dx))(a + b \operatorname{arctanh}(c + dx))^3}{d} d(c + dx) \\
 & \quad \downarrow \text{27} \\
 & \int \frac{(de - cf + f(c + dx))(a + b \operatorname{arctanh}(c + dx))^3}{d^2} d(c + dx) \\
 & \quad \downarrow \text{6480} \\
 & \frac{(f(c + dx) - cf + de)^2 (a + b \operatorname{arctanh}(c + dx))^3}{2f} - \frac{3b \int \left(\frac{(d^2 e^2 - 2cdf e + (c^2 + 1)f^2 + 2f(de - cf)(c + dx))(a + b \operatorname{arctanh}(c + dx))^2}{1 - (c + dx)^2} - f^2 (a + b \operatorname{arctanh}(c + dx))^2 \right)}{d^2} dx}{2f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{(f(c + dx) - cf + de)^2 (a + b \operatorname{arctanh}(c + dx))^3}{2f} - \frac{3b \left(\frac{((c^2 + 1)f^2 - 2cdf + d^2 e^2)(a + b \operatorname{arctanh}(c + dx))^3}{3b} + 2bf(de - cf) \operatorname{PolyLog}\left(2, 1 - \frac{2}{-c - dx + 1}\right) \right) (a + b \operatorname{arctanh}(c + dx))^2}{d^2}
 \end{aligned}$$

input

```
Int[(e + f*x)*(a + b*ArcTanh[c + d*x])^3,x]
```

output

$$\begin{aligned} & \left(\frac{((d*e - c*f + f*(c + d*x))^2*(a + b*\text{ArcTanh}[c + d*x])^3)/(2*f) - (3*b*(-f^2*(a + b*\text{ArcTanh}[c + d*x])^2) - f^2*(c + d*x)*(a + b*\text{ArcTanh}[c + d*x])^2}{(2*f*(d*e - c*f)*(a + b*\text{ArcTanh}[c + d*x])^3)/(3*b) + ((d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2)*(a + b*\text{ArcTanh}[c + d*x])^3)/(3*b) + 2*b*f^2*(a + b*\text{ArcTanh}[c + d*x])*Log[2/(1 - c - d*x)] + 2*f*(d*e - c*f)*(a + b*\text{ArcTanh}[c + d*x])^2*Log[2/(1 - c - d*x)] + b^2*f^2*PolyLog[2, -((1 + c + d*x)/(1 - c - d*x))] + 2*b*f*(d*e - c*f)*(a + b*\text{ArcTanh}[c + d*x])*PolyLog[2, 1 - 2/(1 - c - d*x)] - b^2*f*(d*e - c*f)*PolyLog[3, 1 - 2/(1 - c - d*x)]}{(2*f)} \right) / d^2 \end{aligned}$$

Defintions of rubi rules used

rule 27

$$\text{Int}[(a_*)(F_x), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \&\& \text{!MatchQ}[F_x, (b_*)(G_x)] \text{ ; FreeQ}[b, x]$$

rule 2009

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ ; SumQ}[u]$$

rule 6480

$$\begin{aligned} & \text{Int}[(a_.) + \text{ArcTanh}[(c_.)*(x_)]*(b_.)]^{(p_.)}*((d_.) + (e_.)*(x_))^{(q_.)}, x_Symbol] \rightarrow \text{Simp}[(d + e*x)^{(q + 1)}*((a + b*\text{ArcTanh}[c*x])^p/(e*(q + 1))), x] - \\ & \text{Simp}[b*c*(p/(e*(q + 1))) \quad \text{Int}[\text{ExpandIntegrand}[(a + b*\text{ArcTanh}[c*x])^{(p - 1)}, (d + e*x)^{(q + 1)}/(1 - c^2*x^2), x], x], x] \text{ ; FreeQ}[\{a, b, c, d, e\}, x] \\ & \&\& \text{IGtQ}[p, 1] \&\& \text{IntegerQ}[q] \&\& \text{NeQ}[q, -1] \end{aligned}$$

rule 6661

$$\text{Int}[(a_.) + \text{ArcTanh}[(c_.) + (d_.)*(x_)]*(b_.)]^{(p_.)}*((e_.) + (f_.)*(x_))^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[1/d \quad \text{Subst}[\text{Int}[(d*e - c*f)/d + f*(x/d)]^{m*(a + b*\text{ArcTanh}[x])^p}, x], x, c + d*x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, m\}, x] \&\& \text{IGtQ}[p, 0]$$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 5.99 (sec) , antiderivative size = 10425, normalized size of antiderivative = 31.98

| method | result | size |
|--------------------|---------------------------------|-------|
| parts | Expression too large to display | 10425 |
| derivativeldivides | Expression too large to display | 10431 |
| default | Expression too large to display | 10431 |

input `int((f*x+e)*(a+b*arctanh(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F]

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^3 dx = \int (fx + e)(b \operatorname{arctanh}(dx + c) + a)^3 dx$$

input `integrate((f*x+e)*(a+b*arctanh(d*x+c))^3,x, algorithm="fricas")`

output `integral(a^3*f*x + a^3*e + (b^3*f*x + b^3*e)*arctanh(d*x + c)^3 + 3*(a*b^2*f*x + a*b^2*e)*arctanh(d*x + c)^2 + 3*(a^2*b*f*x + a^2*b*e)*arctanh(d*x + c), x)`

Sympy [F]

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^3 dx = \int (a + b \operatorname{atanh}(c + dx))^3 (e + fx) dx$$

input `integrate((f*x+e)*(a+b*atanh(d*x+c))**3,x)`

output `Integral((a + b*atanh(c + d*x))**3*(e + f*x), x)`

Maxima [F]

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^3 dx = \int (fx + e)(b \operatorname{artanh}(dx + c) + a)^3 dx$$

input `integrate((f*x+e)*(a+b*arctanh(d*x+c))^3,x, algorithm="maxima")`

output `1/2*a^3*f*x^2 + 3/4*(2*x^2*arctanh(d*x + c) + d*(2*x/d^2 - (c^2 + 2*c + 1)*log(d*x + c + 1)/d^3 + (c^2 - 2*c + 1)*log(d*x + c - 1)/d^3))*a^2*b*f + a^3*e*x + 3/2*(2*(d*x + c)*arctanh(d*x + c) + log(-(d*x + c)^2 + 1))*a^2*b*e/d - 1/16*((b^3*d^2*f*x^2 + 2*b^3*d^2*e*x - (c^2*f - 2*(d*e + f)*c + 2*d*e + f)*b^3)*log(-d*x - c + 1)^3 - 3*(2*a*b^2*d^2*f*x^2 + 2*(2*a*b^2*d^2*e + b^3*d*f)*x + (b^3*d^2*f*x^2 + 2*b^3*d^2*e*x - (c^2*f - 2*(d*e - f)*c - 2*d*e + f)*b^3)*log(d*x + c + 1))*log(-d*x - c + 1)^2/d^2 - integrate(-1/8*((b^3*d^2*f*x^2 + (d^2*e + c*d*f - d*f)*b^3*x + (c*d*e - d*e)*b^3)*log(d*x + c + 1)^3 + 6*(a*b^2*d^2*f*x^2 + (d^2*e + c*d*f - d*f)*a*b^2*x + (c*d*e - d*e)*a*b^2)*log(d*x + c + 1)^2 - 3*(2*a*b^2*d^2*f*x^2 + (b^3*d^2*f*x^2 + (d^2*e + c*d*f - d*f)*b^3*x + (c*d*e - d*e)*b^3)*log(d*x + c + 1)^2 + 2*(2*a*b^2*d^2*e + b^3*d*f)*x + (4*(c*d*e - d*e)*a*b^2 - (c^2*f - 2*(d*e - f)*c - 2*d*e + f)*b^3 + (4*a*b^2*d^2*f + b^3*d^2*f)*x^2 + 2*(b^3*d^2*e + 2*(d^2*e + c*d*f - d*f)*a*b^2)*x)*log(d*x + c + 1))*log(-d*x - c + 1)/(d^2*x + c*d - d), x)`

Giac [F]

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^3 dx = \int (fx + e)(b \operatorname{artanh}(dx + c) + a)^3 dx$$

input `integrate((f*x+e)*(a+b*arctanh(d*x+c))^3,x, algorithm="giac")`

output `integrate((f*x + e)*(b*arctanh(d*x + c) + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^3 dx = \int (e + fx) (a + b \operatorname{atanh}(c + dx))^3 dx$$

input `int((e + f*x)*(a + b*atanh(c + d*x))^3,x)`

output `int((e + f*x)*(a + b*atanh(c + d*x))^3, x)`

Reduce [F]

$$\int (e + fx)(a + b \operatorname{arctanh}(c + dx))^3 dx$$

$$= \frac{\operatorname{atanh}(dx + c)^3 b^3 c^2 f - \operatorname{atanh}(dx + c)^3 b^3 f + a^3 d^2 f x^2 - 3 \operatorname{atanh}(dx + c)^2 a b^2 f - 3 \operatorname{atanh}(dx + c) a^2 b f + \dots}{\dots}$$

input `int((f*x+e)*(a+b*atanh(d*x+c))^3,x)`

output

```
(atanh(c + d*x)**3*b**3*c**2*f + 2*atanh(c + d*x)**3*b**3*d**2*e*x + atanh
(c + d*x)**3*b**3*d**2*f*x**2 - atanh(c + d*x)**3*b**3*f + 3*atanh(c + d*x
)**2*a*b**2*c**2*f + 6*atanh(c + d*x)**2*a*b**2*d**2*e*x + 3*atanh(c + d*x
)**2*a*b**2*d**2*f*x**2 - 3*atanh(c + d*x)**2*a*b**2*f + 3*atanh(c + d*x)*
*2*b**3*d*f*x - 3*atanh(c + d*x)*a**2*b*c**2*f + 6*atanh(c + d*x)*a**2*b*c
*d*e - 6*atanh(c + d*x)*a**2*b*c*f + 6*atanh(c + d*x)*a**2*b*d**2*e*x + 3*
atanh(c + d*x)*a**2*b*d**2*f*x**2 + 6*atanh(c + d*x)*a**2*b*d*e - 3*atanh(
c + d*x)*a**2*b*f + 6*atanh(c + d*x)*a*b**2*c*f + 6*atanh(c + d*x)*a*b**2*
d*f*x + 6*atanh(c + d*x)*a*b**2*f - 12*int((atanh(c + d*x)*x)/(c**2 + 2*c*
d*x + d**2*x**2 - 1),x)*a*b**2*c*d**2*f + 12*int((atanh(c + d*x)*x)/(c**2
+ 2*c*d*x + d**2*x**2 - 1),x)*a*b**2*d**3*e + 6*int((atanh(c + d*x)*x)/(c*
**2 + 2*c*d*x + d**2*x**2 - 1),x)*b**3*d**2*f - 6*int((atanh(c + d*x)**2*x)
/(c**2 + 2*c*d*x + d**2*x**2 - 1),x)*b**3*c*d**2*f + 6*int((atanh(c + d*x)
**2*x)/(c**2 + 2*c*d*x + d**2*x**2 - 1),x)*b**3*d**3*e - 6*log(c + d*x - 1
)*a**2*b*c*f + 6*log(c + d*x - 1)*a**2*b*d*e + 6*log(c + d*x - 1)*a*b**2*f
+ 2*a**3*d**2*e*x + a**3*d**2*f*x**2 + 3*a**2*b*d*f*x)/(2*d**2)
```

3.47 $\int (a + b \operatorname{arctanh}(c + dx))^3 dx$

| | |
|----------------------------|-----|
| Optimal result | 435 |
| Mathematica [A] (verified) | 436 |
| Rubi [A] (verified) | 436 |
| Maple [B] (verified) | 438 |
| Fricas [F] | 439 |
| Sympy [F] | 440 |
| Maxima [F] | 440 |
| Giac [F] | 440 |
| Mupad [F(-1)] | 441 |
| Reduce [F] | 441 |

Optimal result

Integrand size = 12, antiderivative size = 132

$$\int (a + b \operatorname{arctanh}(c + dx))^3 dx = \frac{(a + b \operatorname{arctanh}(c + dx))^3}{d} + \frac{(c + dx)(a + b \operatorname{arctanh}(c + dx))^3}{d} - \frac{3b(a + b \operatorname{arctanh}(c + dx))^2 \log\left(\frac{2}{1-c-dx}\right)}{d} - \frac{3b^2(a + b \operatorname{arctanh}(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-c-dx}\right)}{d} + \frac{3b^3 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-c-dx}\right)}{2d}$$

output

```
(a+b*arctanh(d*x+c))^3/d+(d*x+c)*(a+b*arctanh(d*x+c))^3/d-3*b*(a+b*arctanh
(d*x+c))^2*ln(2/(-d*x-c+1))/d-3*b^2*(a+b*arctanh(d*x+c))*polylog(2,1-2/(-d
*x-c+1))/d+3/2*b^3*polylog(3,1-2/(-d*x-c+1))/d
```

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.47

$$\int (a + b \operatorname{arctanh}(c + dx))^3 dx$$

$$= \frac{2a^3(c + dx) + 6a^2b(c + dx)\operatorname{arctanh}(c + dx) + 3a^2b \log(1 - (c + dx)^2) + 6ab^2(\operatorname{arctanh}(c + dx))((-1 + c + dx)) + 2b^3(\operatorname{arctanh}(c + dx))^2((-1 + c + dx)\operatorname{arctanh}(c + dx) - 3\log(1 + E^{-2\operatorname{arctanh}(c + dx)})) + 3\operatorname{arctanh}(c + dx)\operatorname{PolyLog}[2, -E^{-2\operatorname{arctanh}(c + dx)}] + (3\operatorname{PolyLog}[3, -E^{-2\operatorname{arctanh}(c + dx)}])/2)}{2d}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])^3,x]
```

output

```
(2*a^3*(c + d*x) + 6*a^2*b*(c + d*x)*ArcTanh[c + d*x] + 3*a^2*b*Log[1 - (c + d*x)^2] + 6*a*b^2*(ArcTanh[c + d*x]*((-1 + c + d*x)*ArcTanh[c + d*x] - 2*Log[1 + E^(-2*ArcTanh[c + d*x])]) + PolyLog[2, -E^(-2*ArcTanh[c + d*x])]) + 2*b^3*(ArcTanh[c + d*x]^2*(-1 + c + d*x)*ArcTanh[c + d*x] - 3*Log[1 + E^(-2*ArcTanh[c + d*x])]) + 3*ArcTanh[c + d*x]*PolyLog[2, -E^(-2*ArcTanh[c + d*x])]) + (3*PolyLog[3, -E^(-2*ArcTanh[c + d*x])])/2)/(2*d)
```

Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.98, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6653, 6436, 6546, 6470, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{arctanh}(c + dx))^3 dx$$

$$\downarrow \text{6653}$$

$$\frac{\int (a + b \operatorname{arctanh}(c + dx))^3 d(c + dx)}{d}$$

$$\downarrow \text{6436}$$

$$\frac{(c + dx)(a + b \operatorname{arctanh}(c + dx))^3 - 3b \int \frac{(c + dx)(a + b \operatorname{arctanh}(c + dx))^2}{1 - (c + dx)^2} d(c + dx)}{d}$$

$$\frac{(c + dx)(a + \operatorname{arctanh}(c + dx))^3 - 3b \left(\int \frac{(a + \operatorname{arctanh}(c + dx))^2}{-c - dx + 1} d(c + dx) - \frac{(a + \operatorname{arctanh}(c + dx))^3}{3b} \right)}{d}$$

↓ 6546

$$\frac{(c + dx)(a + \operatorname{arctanh}(c + dx))^3 - 3b \left(-2b \int \frac{(a + \operatorname{arctanh}(c + dx)) \log\left(\frac{2}{-c - dx + 1}\right)}{1 - (c + dx)^2} d(c + dx) - \frac{(a + \operatorname{arctanh}(c + dx))^3}{3b} \right)}{d}$$

↓ 6470

↓ 6620

$$\frac{(c + dx)(a + \operatorname{arctanh}(c + dx))^3 - 3b \left(-2b \left(\frac{1}{2} b \int \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{-c - dx + 1}\right)}{1 - (c + dx)^2} d(c + dx) - \frac{1}{2} \operatorname{PolyLog}\left(2, 1 - \frac{2}{-c - dx + 1}\right) \right) \right)}{d}$$

↓ 7164

$$\frac{(c + dx)(a + \operatorname{arctanh}(c + dx))^3 - 3b \left(-2b \left(\frac{1}{4} b \operatorname{PolyLog}\left(3, 1 - \frac{2}{-c - dx + 1}\right) - \frac{1}{2} \operatorname{PolyLog}\left(2, 1 - \frac{2}{-c - dx + 1}\right) \right) (a + b \operatorname{arctanh}(c + dx)) \right)}{d}$$

input `Int[(a + b*ArcTanh[c + d*x])^3,x]`

output `((c + d*x)*(a + b*ArcTanh[c + d*x])^3 - 3*b*(-1/3*(a + b*ArcTanh[c + d*x])^3/b + (a + b*ArcTanh[c + d*x])^2*Log[2/(1 - c - d*x)] - 2*b*(-1/2*((a + b*ArcTanh[c + d*x])*PolyLog[2, 1 - 2/(1 - c - d*x)]) + (b*PolyLog[3, 1 - 2/(1 - c - d*x]))/4))/d`

Defintions of rubi rules used

rule 6436

```
Int[((a_.) + ArcTanh[(c_.)*(x_)^(n_.)]*(b_.))^(p_.), x_Symbol] := Simp[x*(a + b*ArcTanh[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])
```

rule 6470 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol]
-> Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e)
Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /;
FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]`

rule 6546 `Int((((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)*(x_))/((d_) + (e_.)*(x_)^2),
x_Symbol]
-> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Simp[1/(c*d)
Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0]
&& IGtQ[p, 0]`

rule 6620 `Int[(Log[u_]*)((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2),
x_Symbol]
-> Simp[(-(a + b*ArcTanh[c*x])^p)*(PolyLog[2, 1 - u]/(2*c*d)), x] + Simp[b*(p/2)
Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /;
FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]`

rule 6653 `Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.), x_Symbol]
-> Simp[1/d Subst[Int[(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[p, 0]`

rule 7164 `Int[(u_)*PolyLog[n_, v_], x_Symbol]
-> With[{w = DerivativeDivides[v, u*v, x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /;
FreeQ[n, x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 264 vs. $2(130) = 260$.

Time = 1.37 (sec) , antiderivative size = 265, normalized size of antiderivative = 2.01

| method | result |
|-------------------|---|
| derivativedivides | $\frac{a^3(dx+c)+b^3 \left(\operatorname{arctanh}(dx+c)^3(dx+c-1)+2 \operatorname{arctanh}(dx+c)^3-3 \operatorname{arctanh}(dx+c)^2 \ln \left(1+\frac{(dx+c+1)^2}{1-(dx+c)^2} \right) \right)-3 \operatorname{arctanh}(dx+c)}{d}$ |
| default | $\frac{a^3(dx+c)+b^3 \left(\operatorname{arctanh}(dx+c)^3(dx+c-1)+2 \operatorname{arctanh}(dx+c)^3-3 \operatorname{arctanh}(dx+c)^2 \ln \left(1+\frac{(dx+c+1)^2}{1-(dx+c)^2} \right) \right)-3 \operatorname{arctanh}(dx+c)}{d}$ |
| parts | $a^3x + \frac{b^3 \left(\operatorname{arctanh}(dx+c)^3(dx+c-1)+2 \operatorname{arctanh}(dx+c)^3-3 \operatorname{arctanh}(dx+c)^2 \ln \left(1+\frac{(dx+c+1)^2}{1-(dx+c)^2} \right) \right)-3 \operatorname{arctanh}(dx+c)}{d}$ |

input `int((a+b*arctanh(d*x+c))^3,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{d} \left(a^3(dx+c) + b^3 \left(\operatorname{arctanh}(dx+c)^3(dx+c-1) + 2 \operatorname{arctanh}(dx+c)^3 - 3 \operatorname{arctanh}(dx+c)^2 \ln \left(1 + \frac{(dx+c+1)^2}{1-(dx+c)^2} \right) \right) - 3 \operatorname{arctanh}(dx+c) \right) \operatorname{polylog}(2, -\frac{dx+c+1}{1-(dx+c)^2}) + \frac{3}{2} \operatorname{polylog}(3, -\frac{dx+c+1}{1-(dx+c)^2}) + 3a^2b \left(\operatorname{arctanh}(dx+c)^2(dx+c-1) + 2 \operatorname{arctanh}(dx+c)^2 - 2 \operatorname{arctanh}(dx+c) \ln \left(1 + \frac{dx+c+1}{1-(dx+c)^2} \right) \right) - \operatorname{polylog}(2, -\frac{dx+c+1}{1-(dx+c)^2}) + 3a^2b \left(\operatorname{arctanh}(dx+c) + \frac{1}{2} \ln(1-(dx+c)^2) \right)$$

Fricas [F]

$$\int (a + b \operatorname{arctanh}(c + dx))^3 dx = \int (b \operatorname{arctanh}(dx + c) + a)^3 dx$$

input `integrate((a+b*arctanh(d*x+c))^3,x, algorithm="fricas")`

output `integral(b^3*arctanh(d*x + c)^3 + 3*a*b^2*arctanh(d*x + c)^2 + 3*a^2*b*arctanh(d*x + c) + a^3, x)`

Sympy [F]

$$\int (a + b \operatorname{arctanh}(c + dx))^3 dx = \int (a + b \operatorname{atanh}(c + dx))^3 dx$$

input `integrate((a+b*atanh(d*x+c))**3,x)`

output `Integral((a + b*atanh(c + d*x))**3, x)`

Maxima [F]

$$\int (a + b \operatorname{arctanh}(c + dx))^3 dx = \int (b \operatorname{artanh}(dx + c) + a)^3 dx$$

input `integrate((a+b*arctanh(d*x+c))^3,x, algorithm="maxima")`

output `a^3*x + 3/2*(2*(d*x + c)*arctanh(d*x + c) + log(-(d*x + c)^2 + 1))*a^2*b/d - 1/8*((b^3*d*x + b^3*(c - 1))*log(-d*x - c + 1)^3 - 3*(2*a*b^2*d*x + (b^3*d*x + b^3*(c + 1))*log(d*x + c + 1))*log(-d*x - c + 1)^2)/d - integrate(-1/8*((b^3*d*x + b^3*(c - 1))*log(d*x + c + 1)^3 + 6*(a*b^2*d*x + a*b^2*(c - 1))*log(d*x + c + 1)^2 - 3*(4*a*b^2*d*x + (b^3*d*x + b^3*(c - 1))*log(d*x + c + 1)^2 + 2*(b^3*(c + 1) + 2*a*b^2*(c - 1) + (2*a*b^2*d + b^3*d)*x)*log(d*x + c + 1))*log(-d*x - c + 1))/(d*x + c - 1), x)`

Giac [F]

$$\int (a + b \operatorname{arctanh}(c + dx))^3 dx = \int (b \operatorname{artanh}(dx + c) + a)^3 dx$$

input `integrate((a+b*arctanh(d*x+c))^3,x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)^3, x)`

$$3.48 \quad \int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{e + fx} dx$$

| | |
|---------------------------------------|-----|
| Optimal result | 442 |
| Mathematica [F] | 443 |
| Rubi [A] (verified) | 443 |
| Maple [C] (warning: unable to verify) | 445 |
| Fricas [F] | 446 |
| Sympy [F] | 447 |
| Maxima [F] | 447 |
| Giac [F] | 447 |
| Mupad [F(-1)] | 448 |
| Reduce [F] | 448 |

Optimal result

Integrand size = 20, antiderivative size = 308

$$\begin{aligned} & \int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{e + fx} dx \\ &= - \frac{(a + b \operatorname{arctanh}(c + dx))^3 \log\left(\frac{2}{1+c+dx}\right)}{f} \\ & \quad + \frac{(a + b \operatorname{arctanh}(c + dx))^3 \log\left(\frac{2d(e+fx)}{(de+f-cf)(1+c+dx)}\right)}{f} \\ & \quad + \frac{3b(a + b \operatorname{arctanh}(c + dx))^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{2f} \\ & \quad - \frac{3b(a + b \operatorname{arctanh}(c + dx))^2 \operatorname{PolyLog}\left(2, 1 - \frac{2d(e+fx)}{(de+f-cf)(1+c+dx)}\right)}{2f} \\ & \quad + \frac{3b^2(a + b \operatorname{arctanh}(c + dx)) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1+c+dx}\right)}{2f} \\ & \quad - \frac{3b^2(a + b \operatorname{arctanh}(c + dx)) \operatorname{PolyLog}\left(3, 1 - \frac{2d(e+fx)}{(de+f-cf)(1+c+dx)}\right)}{2f} \\ & \quad + \frac{3b^3 \operatorname{PolyLog}\left(4, 1 - \frac{2}{1+c+dx}\right)}{4f} - \frac{3b^3 \operatorname{PolyLog}\left(4, 1 - \frac{2d(e+fx)}{(de+f-cf)(1+c+dx)}\right)}{4f} \end{aligned}$$

output

```

-(a+b*arctanh(d*x+c))^3*ln(2/(d*x+c+1))/f+(a+b*arctanh(d*x+c))^3*ln(2*d*(f
*x+e)/(-c*f+d*e+f)/(d*x+c+1))/f+3/2*b*(a+b*arctanh(d*x+c))^2*polylog(2,1-2
/(d*x+c+1))/f-3/2*b*(a+b*arctanh(d*x+c))^2*polylog(2,1-2*d*(f*x+e)/(-c*f+d
*e+f)/(d*x+c+1))/f+3/2*b^2*(a+b*arctanh(d*x+c))*polylog(3,1-2/(d*x+c+1))/f
-3/2*b^2*(a+b*arctanh(d*x+c))*polylog(3,1-2*d*(f*x+e)/(-c*f+d*e+f)/(d*x+c+
1))/f+3/4*b^3*polylog(4,1-2/(d*x+c+1))/f-3/4*b^3*polylog(4,1-2*d*(f*x+e)/(-
c*f+d*e+f)/(d*x+c+1))/f

```

Mathematica [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{e + fx} dx = \int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{e + fx} dx$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])^3/(e + f*x), x]
```

output

```
Integrate[(a + b*ArcTanh[c + d*x])^3/(e + f*x), x]
```

Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 344, normalized size of antiderivative = 1.12, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {6661, 27, 6476}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{e + fx} dx \\
\downarrow \text{6661} \\
\int \frac{d(a + b \operatorname{arctanh}(c + dx))^3}{d(e - \frac{cf}{d}) + f(c + dx)} d(c + dx) \\
\hline
d \\
\downarrow \text{27}
\end{array}$$

$$\begin{aligned}
& \int \frac{(a + \operatorname{barctanh}(c + dx))^3}{f(c + dx) - cf + de} d(c + dx) \\
& \quad \downarrow \text{6476} \\
& -\frac{3b^2(a + \operatorname{barctanh}(c + dx)) \operatorname{PolyLog}\left(3, 1 - \frac{2(de - cf + f(c + dx))}{(de - cf + f)(c + dx + 1)}\right)}{2f} + \\
& \quad \frac{3b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{c + dx + 1}\right) (a + \operatorname{barctanh}(c + dx))}{2f} - \\
& \quad \frac{3b(a + \operatorname{barctanh}(c + dx))^2 \operatorname{PolyLog}\left(2, 1 - \frac{2(de - cf + f(c + dx))}{(de - cf + f)(c + dx + 1)}\right)}{2f} + \\
& \quad \frac{(a + \operatorname{barctanh}(c + dx))^3 \log\left(\frac{2(f(c + dx) - cf + de)}{(c + dx + 1)(-cf + de + f)}\right)}{f} + \\
& \frac{3b \operatorname{PolyLog}\left(2, 1 - \frac{2}{c + dx + 1}\right) (a + \operatorname{barctanh}(c + dx))^2 \log\left(\frac{2}{c + dx + 1}\right) (a + \operatorname{barctanh}(c + dx))^3}{2f} - \\
& \quad \frac{3b^3 \operatorname{PolyLog}\left(4, 1 - \frac{2(de - cf + f(c + dx))}{(de - cf + f)(c + dx + 1)}\right)}{4f} + \frac{3b^3 \operatorname{PolyLog}\left(4, 1 - \frac{2}{c + dx + 1}\right)}{4f}
\end{aligned}$$

input `Int[(a + b*ArcTanh[c + d*x])^3/(e + f*x),x]`

output `-(((a + b*ArcTanh[c + d*x])^3*Log[2/(1 + c + d*x)])/f) + ((a + b*ArcTanh[c + d*x])^3*Log[(2*(d*e - c*f + f*(c + d*x))]/((d*e + f - c*f)*(1 + c + d*x)))]/f + (3*b*(a + b*ArcTanh[c + d*x])^2*PolyLog[2, 1 - 2/(1 + c + d*x)])/(2*f) - (3*b*(a + b*ArcTanh[c + d*x])^2*PolyLog[2, 1 - (2*(d*e - c*f + f*(c + d*x))]/((d*e + f - c*f)*(1 + c + d*x)))]/(2*f) + (3*b^2*(a + b*ArcTanh[c + d*x])*PolyLog[3, 1 - 2/(1 + c + d*x)])/(2*f) - (3*b^2*(a + b*ArcTanh[c + d*x])*PolyLog[3, 1 - (2*(d*e - c*f + f*(c + d*x))]/((d*e + f - c*f)*(1 + c + d*x)))]/(2*f) + (3*b^3*PolyLog[4, 1 - 2/(1 + c + d*x)])/(4*f) - (3*b^3*PolyLog[4, 1 - (2*(d*e - c*f + f*(c + d*x))]/((d*e + f - c*f)*(1 + c + d*x)))]/(4*f)`

Definitions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

rule 6476

```
Int[((a_) + ArcTanh[(c_)*(x_)]*(b_))^3/((d_) + (e_)*(x_)), x_Symbol] :=
  Simp[(-(a + b*ArcTanh[c*x])^3)*(Log[2/(1 + c*x)]/e), x] + (Simp[(a + b*ArcTanh[c*x])^3*(Log[2*c*((d + e*x)/((c*d + e)*(1 + c*x))])/e), x] + Simp[3*b*(a + b*ArcTanh[c*x])^2*(PolyLog[2, 1 - 2/(1 + c*x)]/(2*e)), x] - Simp[3*b*(a + b*ArcTanh[c*x])^2*(PolyLog[2, 1 - 2*c*((d + e*x)/((c*d + e)*(1 + c*x))])/((2*e)), x] + Simp[3*b^2*(a + b*ArcTanh[c*x])*(PolyLog[3, 1 - 2/(1 + c*x)]/(2*e)), x] - Simp[3*b^2*(a + b*ArcTanh[c*x])*(PolyLog[3, 1 - 2*c*((d + e*x)/((c*d + e)*(1 + c*x))])/((2*e)), x] + Simp[3*b^3*(PolyLog[4, 1 - 2/(1 + c*x)]/(4*e)), x] - Simp[3*b^3*(PolyLog[4, 1 - 2*c*((d + e*x)/((c*d + e)*(1 + c*x))])/((4*e)), x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 - e^2, 0]
```

rule 6661

```
Int[((a_) + ArcTanh[(c_) + (d_)*(x_)]*(b_))^(p_)*((e_) + (f_)*(x_))^(m_), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IntegerQ[p, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 5.14 (sec) , antiderivative size = 3441, normalized size of antiderivative = 11.17

| method | result | size |
|-------------------|---------------------------------|------|
| derivativedivides | Expression too large to display | 3441 |
| default | Expression too large to display | 3441 |
| parts | Expression too large to display | 3650 |

input

```
int((a+b*arctanh(d*x+c))^3/(f*x+e),x,method=_RETURNVERBOSE)
```

output

```

1/d*(a^3*d*ln(f*c-d*e-f*(d*x+c))/f-b^3*d*(-ln(f*c-d*e-f*(d*x+c))/f*arctanh
(d*x+c)^3+3/f*(1/3*arctanh(d*x+c)^3*ln(f*c*(1+(d*x+c+1)^2/(1-(d*x+c)^2)))+(
-(d*x+c+1)^2/(1-(d*x+c)^2)-1)*e*d+(-(d*x+c+1)^2/(1-(d*x+c)^2)+1)*f)-1/6*I*
Pi*csgn(I*(f*c*(1-(d*x+c+1)^2/((d*x+c)^2-1))+((d*x+c+1)^2/((d*x+c)^2-1)-1)
*e*d+((d*x+c+1)^2/((d*x+c)^2-1)+1)*f)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))*csgn
(I*(f*c*(1-(d*x+c+1)^2/((d*x+c)^2-1))+((d*x+c+1)^2/((d*x+c)^2-1)-1)*e*d+((
d*x+c+1)^2/((d*x+c)^2-1)+1)*f))*csgn(I/(1-(d*x+c+1)^2/((d*x+c)^2-1)))-csgn
(I*(f*c*(1-(d*x+c+1)^2/((d*x+c)^2-1))+((d*x+c+1)^2/((d*x+c)^2-1)-1)*e*d+((
d*x+c+1)^2/((d*x+c)^2-1)+1)*f)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))*csgn(I/(1-(d
*x+c+1)^2/((d*x+c)^2-1)))-csgn(I*(f*c*(1-(d*x+c+1)^2/((d*x+c)^2-1))+((d*x+
c+1)^2/((d*x+c)^2-1)-1)*e*d+((d*x+c+1)^2/((d*x+c)^2-1)+1)*f))*csgn(I*(f*c*
(1-(d*x+c+1)^2/((d*x+c)^2-1))+((d*x+c+1)^2/((d*x+c)^2-1)-1)*e*d+((d*x+c+1)
^2/((d*x+c)^2-1)+1)*f)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))+csgn(I*(f*c*(1-(d*x+
c+1)^2/((d*x+c)^2-1))+((d*x+c+1)^2/((d*x+c)^2-1)-1)*e*d+((d*x+c+1)^2/((d*x
+c)^2-1)+1)*f)/(1-(d*x+c+1)^2/((d*x+c)^2-1)))^2)*arctanh(d*x+c)^3+1/2*arct
anh(d*x+c)^2*polylog(2,-(d*x+c+1)^2/(1-(d*x+c)^2))-1/2*arctanh(d*x+c)*poly
log(3,-(d*x+c+1)^2/(1-(d*x+c)^2))+1/4*polylog(4,-(d*x+c+1)^2/(1-(d*x+c)^2)
)-1/3/(c*f-d*e-f)*f*c*arctanh(d*x+c)^3*ln(1-(c*f-d*e-f)*(d*x+c+1)^2/(1-(d
*x+c)^2)/(-c*f+d*e-f))-1/2/(c*f-d*e-f)*f*c*arctanh(d*x+c)^2*polylog(2,(c*f-
d*e-f)*(d*x+c+1)^2/(1-(d*x+c)^2)/(-c*f+d*e-f))+1/2/(c*f-d*e-f)*f*c*arct...

```

Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{e + fx} dx = \int \frac{(b \operatorname{arctanh}(dx + c) + a)^3}{fx + e} dx$$

input

```
integrate((a+b*arctanh(d*x+c))^3/(f*x+e),x, algorithm="fricas")
```

output

```
integral((b^3*arctanh(d*x + c)^3 + 3*a*b^2*arctanh(d*x + c)^2 + 3*a^2*b*ar
ctanh(d*x + c) + a^3)/(f*x + e), x)
```

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{e + fx} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^3}{e + fx} dx$$

input `integrate((a+b*atanh(d*x+c))**3/(f*x+e),x)`

output `Integral((a + b*atanh(c + d*x))**3/(e + f*x), x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{e + fx} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^3}{fx + e} dx$$

input `integrate((a+b*arctanh(d*x+c))^3/(f*x+e),x, algorithm="maxima")`

output `a^3*log(f*x + e)/f + integrate(1/8*b^3*(log(d*x + c + 1) - log(-d*x - c + 1))^3/(f*x + e) + 3/4*a*b^2*(log(d*x + c + 1) - log(-d*x - c + 1))^2/(f*x + e) + 3/2*a^2*b*(log(d*x + c + 1) - log(-d*x - c + 1))/(f*x + e), x)`

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{e + fx} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^3}{fx + e} dx$$

input `integrate((a+b*arctanh(d*x+c))^3/(f*x+e),x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)^3/(f*x + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{e + fx} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^3}{e + fx} dx$$

input `int((a + b*atanh(c + d*x))^3/(e + f*x),x)`output `int((a + b*atanh(c + d*x))^3/(e + f*x), x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{e + fx} dx$$

$$= \frac{3 \left(\int \frac{\operatorname{atanh}(dx+c)}{fx+e} dx \right) a^2 b f + \left(\int \frac{\operatorname{atanh}(dx+c)^3}{fx+e} dx \right) b^3 f + 3 \left(\int \frac{\operatorname{atanh}(dx+c)^2}{fx+e} dx \right) a b^2 f + \log(fx + e) a^3}{f}$$

input `int((a+b*atanh(d*x+c))^3/(f*x+e),x)`output `(3*int(atanh(c + d*x)/(e + f*x),x)*a**2*b*f + int(atanh(c + d*x)**3/(e + f*x),x)*b**3*f + 3*int(atanh(c + d*x)**2/(e + f*x),x)*a*b**2*f + log(e + f*x)*a**3)/f`

$$3.49 \quad \int \frac{(a+b \operatorname{arctanh}(c+dx))^3}{(e+fx)^2} dx$$

| | |
|---|-----|
| Optimal result | 450 |
| Mathematica [C] (warning: unable to verify) | 451 |
| Rubi [A] (verified) | 452 |
| Maple [C] (warning: unable to verify) | 455 |
| Fricas [F] | 456 |
| Sympy [F] | 456 |
| Maxima [F] | 456 |
| Giac [F] | 457 |
| Mupad [F(-1)] | 458 |
| Reduce [F] | 458 |

Optimal result

Integrand size = 20, antiderivative size = 634

$$\begin{aligned}
& \int \frac{(a + \operatorname{barctanh}(c + dx))^3}{(e + fx)^2} dx \\
&= -\frac{(a + \operatorname{barctanh}(c + dx))^3}{f(e + fx)} + \frac{3bd(a + \operatorname{barctanh}(c + dx))^2 \log\left(\frac{2}{1-c-dx}\right)}{2f(de + f - cf)} \\
&\quad - \frac{3bd(a + \operatorname{barctanh}(c + dx))^2 \log\left(\frac{2}{1+c+dx}\right)}{2f(de - (1 + c)f)} \\
&\quad + \frac{3bd(a + \operatorname{barctanh}(c + dx))^2 \log\left(\frac{2}{1+c+dx}\right)}{(de + f - cf)(de - (1 + c)f)} \\
&\quad - \frac{3bd(a + \operatorname{barctanh}(c + dx))^2 \log\left(\frac{2d(e+fx)}{(de+f-cf)(1+c+dx)}\right)}{(de + f - cf)(de - (1 + c)f)} \\
&\quad + \frac{3b^2d(a + \operatorname{barctanh}(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-c-dx}\right)}{2f(de + f - cf)} \\
&\quad + \frac{3b^2d(a + \operatorname{barctanh}(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{2f(de - (1 + c)f)} \\
&\quad - \frac{3b^2d(a + \operatorname{barctanh}(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{(de + f - cf)(de - (1 + c)f)} \\
&\quad + \frac{3b^2d(a + \operatorname{barctanh}(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2d(e+fx)}{(de+f-cf)(1+c+dx)}\right)}{(de + f - cf)(de - (1 + c)f)} \\
&\quad - \frac{3b^3d \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-c-dx}\right)}{4f(de + f - cf)} + \frac{3b^3d \operatorname{PolyLog}\left(3, 1 - \frac{2}{1+c+dx}\right)}{4f(de - (1 + c)f)} \\
&\quad - \frac{3b^3d \operatorname{PolyLog}\left(3, 1 - \frac{2}{1+c+dx}\right)}{2(de + f - cf)(de - (1 + c)f)} + \frac{3b^3d \operatorname{PolyLog}\left(3, 1 - \frac{2d(e+fx)}{(de+f-cf)(1+c+dx)}\right)}{2(de + f - cf)(de - (1 + c)f)}
\end{aligned}$$

output

```

-(a+b*arctanh(d*x+c))^3/f/(f*x+e)+3/2*b*d*(a+b*arctanh(d*x+c))^2*ln(2/(-d*
x-c+1))/f/(-c*f+d*e+f)-3/2*b*d*(a+b*arctanh(d*x+c))^2*ln(2/(d*x+c+1))/f/(d
*e-(1+c)*f)+3*b*d*(a+b*arctanh(d*x+c))^2*ln(2/(d*x+c+1))/(-c*f+d*e+f)/(d*e
-(1+c)*f)-3*b*d*(a+b*arctanh(d*x+c))^2*ln(2*d*(f*x+e)/(-c*f+d*e+f)/(d*x+c+
1))/(-c*f+d*e+f)/(d*e-(1+c)*f)+3/2*b^2*d*(a+b*arctanh(d*x+c))*polylog(2,1-
2/(-d*x-c+1))/f/(-c*f+d*e+f)+3/2*b^2*d*(a+b*arctanh(d*x+c))*polylog(2,1-2/
(d*x+c+1))/f/(d*e-(1+c)*f)-3*b^2*d*(a+b*arctanh(d*x+c))*polylog(2,1-2/(d*x
+c+1))/(-c*f+d*e+f)/(d*e-(1+c)*f)+3*b^2*d*(a+b*arctanh(d*x+c))*polylog(2,1
-2*d*(f*x+e)/(-c*f+d*e+f)/(d*x+c+1))/(-c*f+d*e+f)/(d*e-(1+c)*f)-3/4*b^3*d*
polylog(3,1-2/(-d*x-c+1))/f/(-c*f+d*e+f)+3/4*b^3*d*polylog(3,1-2/(d*x+c+1)
)/f/(d*e-(1+c)*f)-3/2*b^3*d*polylog(3,1-2/(d*x+c+1))/(-c*f+d*e+f)/(d*e-(1+
c)*f)+3/2*b^3*d*polylog(3,1-2*d*(f*x+e)/(-c*f+d*e+f)/(d*x+c+1))/(-c*f+d*e+
f)/(d*e-(1+c)*f)

```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 90.09 (sec) , antiderivative size = 3878, normalized size of antiderivative = 6.12

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(e + fx)^2} dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])^3/(e + f*x)^2,x]
```

output

```

-(a^3/(f*(e + f*x))) - (3*a^2*b*ArcTanh[c + d*x])/(f*(e + f*x)) + (3*a^2*b
*d*Log[1 - c - d*x])/(2*f*(-(d*e) - f + c*f)) - (3*a^2*b*d*Log[1 + c + d*x
])/ (2*f*(-(d*e) + f + c*f)) - (3*a^2*b*d*Log[e + f*x])/(d^2*e^2 - 2*c*d*e*
f - f^2 + c^2*f^2) + (3*a*b^2*(1 - (c + d*x)^2)*((d*e - c*f)/Sqrt[1 - (c +
d*x)^2] + (f*(c + d*x))/Sqrt[1 - (c + d*x)^2])^2*(-((E^ArcTanh[c - (d*e)/
f]*ArcTanh[c + d*x]^2)/(f*Sqrt[1 - (d*e - c*f)^2/f^2])) + ((c + d*x)*ArcTa
nh[c + d*x]^2)/(Sqrt[1 - (c + d*x)^2]*((d*e)/Sqrt[1 - (c + d*x)^2] - (c*f)
/Sqrt[1 - (c + d*x)^2] + (f*(c + d*x))/Sqrt[1 - (c + d*x)^2])) + ((d*e - c
*f)*(-2*ArcTanh[c + d*x]*Log[1 - E^(2*ArcTanh[c - (d*e)/f] - 2*ArcTanh[c +
d*x])] + I*Pi*Log[1 + E^(2*ArcTanh[c + d*x])]) - I*Pi*(ArcTanh[c + d*x] +
Log[1/Sqrt[1 - (c + d*x)^2]]) + 2*ArcTanh[c - (d*e)/f]*(ArcTanh[c + d*x] +
Log[1 - E^(2*ArcTanh[c - (d*e)/f] - 2*ArcTanh[c + d*x]]) - Log[I*Sinh[Arc
Tanh[(d*e - c*f)/f] + ArcTanh[c + d*x]]) + PolyLog[2, E^(2*ArcTanh[c - (d
e)/f] - 2*ArcTanh[c + d*x])])/(d^2*e^2 - 2*c*d*e*f + (-1 + c^2)*f^2))/(
d*(d*e - c*f)*(e + f*x)^2) + (b^3*(1 - (c + d*x)^2)*((d*e - c*f)/Sqrt[1 -
(c + d*x)^2] + (f*(c + d*x))/Sqrt[1 - (c + d*x)^2])^2*((d*(c + d*x)*ArcTan
h[c + d*x]^3)/((d*e - c*f)*Sqrt[1 - (c + d*x)^2]*((d*e)/Sqrt[1 - (c + d*x)
^2] - (c*f)/Sqrt[1 - (c + d*x)^2] + (f*(c + d*x))/Sqrt[1 - (c + d*x)^2]))
- (d*(-6*d*e*ArcTanh[c + d*x]^3 + 2*f*ArcTanh[c + d*x]^3 + 6*c*f*ArcTanh[c
+ d*x]^3 - 4*E^ArcTanh[c - (d*e)/f]*Sqrt[1 - c^2 - (d^2*e^2)/f^2 + (2*...

```

Rubi [A] (verified)

Time = 2.80 (sec) , antiderivative size = 1085, normalized size of antiderivative = 1.71, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {6659, 7292, 6671, 27, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(e + fx)^2} dx$$

$$\downarrow \text{6659}$$

$$\frac{3bd \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)(1 - (c + dx)^2)} dx}{f} - \frac{(a + b \operatorname{arctanh}(c + dx))^3}{f(e + fx)}$$

$$\downarrow \text{7292}$$

$$\frac{3bd \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(e + fx)(-c^2 - 2dxc - d^2x^2 + 1)} dx}{f} - \frac{(a + b \operatorname{arctanh}(c + dx))^3}{f(e + fx)}$$

↓ 6671

$$\frac{3b \int \frac{d(a + b \operatorname{arctanh}(c + dx))^2}{\left(d\left(e - \frac{cf}{d}\right) + f(c + dx)\right)(1 - (c + dx)^2)} d(c + dx)}{f} - \frac{(a + b \operatorname{arctanh}(c + dx))^3}{f(e + fx)}$$

↓ 27

$$\frac{3bd \int \frac{(a + b \operatorname{arctanh}(c + dx))^2}{(de - cf + f(c + dx))(1 - (c + dx)^2)} d(c + dx)}{f} - \frac{(a + b \operatorname{arctanh}(c + dx))^3}{f(e + fx)}$$

↓ 7276

$$\frac{3bd \int \left(-\frac{a^2}{(c + dx - 1)(c + dx + 1)(de - cf + f(c + dx))} - \frac{2b \operatorname{arctanh}(c + dx)a}{(c + dx - 1)(c + dx + 1)(de - cf + f(c + dx))} - \frac{b^2 \operatorname{arctanh}(c + dx)^2}{(c + dx - 1)(c + dx + 1)(de - cf + f(c + dx))} \right) d(c + dx)}{(a + b \operatorname{arctanh}(c + dx))^3}{f(e + fx)}$$

↓ 2009

$$\frac{3bd \left(-\frac{\log(-c - dx + 1)a^2}{2(de - cf + f)} + \frac{\log(c + dx + 1)a^2}{2(de - (c + 1)f)} - \frac{f \log(de - cf + f(c + dx))a^2}{(de - cf + f)(de - (c + 1)f)} + \frac{b \operatorname{arctanh}(c + dx) \log\left(\frac{2}{-c - dx + 1}\right)a}{de - cf + f} - \frac{b \operatorname{arctanh}(c + dx) \log\left(\frac{2}{-c - dx + 1}\right)a}{de - cf - f} \right)}{(a + b \operatorname{arctanh}(c + dx))^3}{f(e + fx)}$$

input `Int[(a + b*ArcTanh[c + d*x])^3/(e + f*x)^2,x]`

output

```

-((a + b*ArcTanh[c + d*x])^3/(f*(e + f*x))) + (3*b*d*((a*b*ArcTanh[c + d*x]
]*Log[2/(1 - c - d*x)])/(d*e + f - c*f) + (b^2*ArcTanh[c + d*x]^2*Log[2/(1
- c - d*x)])/(2*(d*e + f - c*f)) - (a^2*Log[1 - c - d*x])/(2*(d*e + f - c
*f)) - (a*b*ArcTanh[c + d*x]*Log[2/(1 + c + d*x)])/(d*e - f - c*f) + (2*a*
b*f*ArcTanh[c + d*x]*Log[2/(1 + c + d*x)])/((d*e + f - c*f)*(d*e - (1 + c)
*f)) - (b^2*ArcTanh[c + d*x]^2*Log[2/(1 + c + d*x)])/(2*(d*e - f - c*f)) +
(b^2*f*ArcTanh[c + d*x]^2*Log[2/(1 + c + d*x)])/((d*e + f - c*f)*(d*e - (
1 + c)*f)) + (a^2*Log[1 + c + d*x])/(2*(d*e - (1 + c)*f)) - (a^2*f*Log[d*e
- c*f + f*(c + d*x)])/((d*e + f - c*f)*(d*e - (1 + c)*f)) - (2*a*b*f*ArcT
anh[c + d*x]*Log[(2*(d*e - c*f + f*(c + d*x))]/((d*e + f - c*f)*(1 + c + d
*x)))]/((d*e + f - c*f)*(d*e - (1 + c)*f)) - (b^2*f*ArcTanh[c + d*x]^2*Log
[(2*(d*e - c*f + f*(c + d*x))]/((d*e + f - c*f)*(1 + c + d*x)))]/((d*e + f
- c*f)*(d*e - (1 + c)*f)) + (a*b*PolyLog[2, -((1 + c + d*x)/(1 - c - d*x)
)])/(2*(d*e + f - c*f)) + (b^2*ArcTanh[c + d*x]*PolyLog[2, 1 - 2/(1 - c -
d*x)])/(2*(d*e + f - c*f)) + (a*b*PolyLog[2, 1 - 2/(1 + c + d*x)])/(2*(d*e
- f - c*f)) - (a*b*f*PolyLog[2, 1 - 2/(1 + c + d*x)])/((d*e + f - c*f)*(d
*e - (1 + c)*f)) + (b^2*ArcTanh[c + d*x]*PolyLog[2, 1 - 2/(1 + c + d*x)])/
(2*(d*e - f - c*f)) - (b^2*f*ArcTanh[c + d*x]*PolyLog[2, 1 - 2/(1 + c + d*
x)])/((d*e + f - c*f)*(d*e - (1 + c)*f)) + (a*b*f*PolyLog[2, 1 - (2*(d*e -
c*f + f*(c + d*x))]/((d*e + f - c*f)*(1 + c + d*x)))]/((d*e + f - c*f)...

```

Defintions of rubi rules used

rule 27

```

Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]

```

rule 2009

```

Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]

```

rule 6659

```

Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)])*(b_.))^ (p_.)*((e_.) + (f_.)*(x_))^(
m_), x_Symbol] := Simp[(e + f*x)^(m + 1)*((a + b*ArcTanh[c + d*x])^p/(f*(m
+ 1))), x] - Simp[b*d*(p/(f*(m + 1))) Int[(e + f*x)^(m + 1)*((a + b*ArcTa
nh[c + d*x])^(p - 1)/(1 - (c + d*x)^2)), x], x] /; FreeQ[{a, b, c, d, e, f}
, x] && IGtQ[p, 0] && ILtQ[m, -1]

```

rule 6671

```
Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.)*((A_.) + (B_.)*(x_) + (C_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/d Sub
st[Int[((d*e - c*f)/d + f*(x/d))^m*(-C/d^2 + (C/d^2)*x^2)^q*(a + b*ArcTanh[
x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, p, q}, x
] && EqQ[B*(1 - c^2) + 2*A*c*d, 0] && EqQ[2*c*C - B*d, 0]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

rule 7292

```
Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v !=
= u]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.07 (sec) , antiderivative size = 5109, normalized size of antiderivative = 8.06

| method | result | size |
|-------------------|---------------------------------|------|
| derivativedivides | Expression too large to display | 5109 |
| default | Expression too large to display | 5109 |
| parts | Expression too large to display | 5242 |

input

```
int((a+b*arctanh(d*x+c))^3/(f*x+e)^2,x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```


Fricas [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(e + fx)^2} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^3}{(fx + e)^2} dx$$

input `integrate((a+b*arctanh(d*x+c))^3/(f*x+e)^2,x, algorithm="fricas")`

output `integral((b^3*arctanh(d*x + c)^3 + 3*a*b^2*arctanh(d*x + c)^2 + 3*a^2*b*arctanh(d*x + c) + a^3)/(f^2*x^2 + 2*e*f*x + e^2), x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(e + fx)^2} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^3}{(e + fx)^2} dx$$

input `integrate((a+b*atanh(d*x+c))**3/(f*x+e)**2,x)`

output `Integral((a + b*atanh(c + d*x))**3/(e + f*x)**2, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(e + fx)^2} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)^3}{(fx + e)^2} dx$$

input `integrate((a+b*arctanh(d*x+c))^3/(f*x+e)^2,x, algorithm="maxima")`

output

```

3/2*(d*(log(d*x + c + 1)/(d*e*f - (c + 1)*f^2) - log(d*x + c - 1)/(d*e*f -
(c - 1)*f^2) - 2*log(f*x + e)/(d^2*e^2 - 2*c*d*e*f + (c^2 - 1)*f^2)) - 2*
arctanh(d*x + c)/(f^2*x + e*f))*a^2*b - a^3/(f^2*x + e*f) - 1/8*(((d^2*e*f
- c*d*f^2 - d*f^2)*b^3*x + (c*d*e*f - c^2*f^2 - d*e*f + f^2)*b^3)*log(-d*
x - c + 1)^3 + 3*(2*(d^2*e^2 - 2*c*d*e*f + c^2*f^2 - f^2)*a*b^2 - ((d^2*e*
f - c*d*f^2 + d*f^2)*b^3*x + (c*d*e*f - c^2*f^2 + d*e*f + f^2)*b^3)*log(d*
x + c + 1))*log(-d*x - c + 1)^2)/(d^2*e^3*f - 2*c*d*e^2*f^2 + c^2*e*f^3 -
e*f^3 + (d^2*e^2*f^2 - 2*c*d*e*f^3 + c^2*f^4 - f^4)*x) - integrate(-1/8*((
d^2*e*f - c*d*f^2 - d*f^2)*b^3*x + (c*d*e*f - c^2*f^2 - d*e*f + f^2)*b^3)
*log(d*x + c + 1)^3 + 6*((d^2*e*f - c*d*f^2 - d*f^2)*a*b^2*x + (c*d*e*f -
c^2*f^2 - d*e*f + f^2)*a*b^2)*log(d*x + c + 1)^2 + 3*(4*(d^2*e*f - c*d*f^2
- d*f^2)*a*b^2*x + 4*(d^2*e^2 - c*d*e*f - d*e*f)*a*b^2 - ((d^2*e*f - c*d*
f^2 - d*f^2)*b^3*x + (c*d*e*f - c^2*f^2 - d*e*f + f^2)*b^3)*log(d*x + c +
1)^2 - 2*(b^3*d^2*f^2*x^2 + 2*(c*d*e*f - c^2*f^2 - d*e*f + f^2)*a*b^2 + (c
*d*e*f + d*e*f)*b^3 + (2*(d^2*e*f - c*d*f^2 - d*f^2)*a*b^2 + (d^2*e*f + c*
d*f^2 + d*f^2)*b^3)*x)*log(d*x + c + 1))*log(-d*x - c + 1))/(c*d*e^3*f - c
^2*e^2*f^2 - d*e^3*f + e^2*f^2 + (d^2*e*f^3 - c*d*f^4 - d*f^4)*x^3 + (2*d^
2*e^2*f^2 - c*d*e*f^3 - c^2*f^4 - 3*d*e*f^3 + f^4)*x^2 + (d^2*e^3*f + c*d*
e^2*f^2 - 2*c^2*e*f^3 - 3*d*e^2*f^2 + 2*e*f^3)*x), x)

```

Giac [F]

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(e + fx)^2} dx = \int \frac{(b \operatorname{arctanh}(dx + c) + a)^3}{(fx + e)^2} dx$$

input

```
integrate((a+b*arctanh(d*x+c))^3/(f*x+e)^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(d*x + c) + a)^3/(f*x + e)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(e + fx)^2} dx = \int \frac{(a + b \operatorname{atanh}(c + dx))^3}{(e + fx)^2} dx$$

input `int((a + b*atanh(c + d*x))^3/(e + f*x)^2,x)`output `int((a + b*atanh(c + d*x))^3/(e + f*x)^2, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{arctanh}(c + dx))^3}{(e + fx)^2} dx = \text{too large to display}$$

input `int((a+b*atanh(d*x+c))^3/(f*x+e)^2,x)`

output

```
( - 4*atanh(c + d*x)**3*b**3*c**6*e*f**5 + 16*atanh(c + d*x)**3*b**3*c**5*
d**2*f**4 - 24*atanh(c + d*x)**3*b**3*c**4*d**2*e**3*f**3 - 4*atanh(c +
d*x)**3*b**3*c**4*d**2*e**2*f**4*x + 8*atanh(c + d*x)**3*b**3*c**4*e*f**5
+ 16*atanh(c + d*x)**3*b**3*c**3*d**3*e**4*f**2 + 16*atanh(c + d*x)**3*b**
3*c**3*d**3*e**3*f**3*x - 24*atanh(c + d*x)**3*b**3*c**3*d**2*f**4 - 4*a
tanh(c + d*x)**3*b**3*c**2*d**4*e**5*f - 24*atanh(c + d*x)**3*b**3*c**2*d
**4*e**4*f**2*x + 28*atanh(c + d*x)**3*b**3*c**2*d**2*e**3*f**3 + 4*atanh(c
+ d*x)**3*b**3*c**2*d**2*e**2*f**4*x - 4*atanh(c + d*x)**3*b**3*c**2*e*f
**5 + 16*atanh(c + d*x)**3*b**3*c*d**5*e**5*f*x - 16*atanh(c + d*x)**3*b**3
*c*d**3*e**4*f**2 - 8*atanh(c + d*x)**3*b**3*c*d**3*e**3*f**3*x + 8*atanh(
c + d*x)**3*b**3*c*d**2*f**4 - 4*atanh(c + d*x)**3*b**3*d**6*e**6*x + 4*
atanh(c + d*x)**3*b**3*d**4*e**5*f + 4*atanh(c + d*x)**3*b**3*d**4*e**4*f
**2*x - 4*atanh(c + d*x)**3*b**3*d**2*e**3*f**3 - 12*atanh(c + d*x)**2*a*b
**2*c**6*e*f**5 + 48*atanh(c + d*x)**2*a*b**2*c**5*d**2*f**4 - 72*atanh(c
+ d*x)**2*a*b**2*c**4*d**2*e**3*f**3 - 12*atanh(c + d*x)**2*a*b**2*c**4*d
**2*e**2*f**4*x + 24*atanh(c + d*x)**2*a*b**2*c**4*e*f**5 + 48*atanh(c + d
*x)**2*a*b**2*c**3*d**3*e**4*f**2 + 48*atanh(c + d*x)**2*a*b**2*c**3*d**3*
e**3*f**3*x - 72*atanh(c + d*x)**2*a*b**2*c**3*d**2*f**4 - 12*atanh(c +
d*x)**2*a*b**2*c**2*d**4*e**5*f - 72*atanh(c + d*x)**2*a*b**2*c**2*d**4*e
**4*f**2*x + 84*atanh(c + d*x)**2*a*b**2*c**2*d**2*e**3*f**3 + 12*atanh(...
```

3.50 $\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^3 dx$

| | |
|-------------------|-----|
| Optimal result | 460 |
| Mathematica [N/A] | 460 |
| Rubi [N/A] | 461 |
| Maple [N/A] | 461 |
| Fricas [N/A] | 462 |
| Sympy [F(-1)] | 462 |
| Maxima [N/A] | 462 |
| Giac [N/A] | 463 |
| Mupad [N/A] | 463 |
| Reduce [N/A] | 464 |

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^3 dx = \operatorname{Int}((e + fx)^m (a + b \operatorname{arctanh}(c + dx))^3, x)$$

output `Defer(Int)((f*x+e)^m*(a+b*arctanh(d*x+c))^3,x)`

Mathematica [N/A]

Not integrable

Time = 3.40 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^3 dx = \int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^3 dx$$

input `Integrate[(e + f*x)^m*(a + b*ArcTanh[c + d*x])^3,x]`

output `Integrate[(e + f*x)^m*(a + b*ArcTanh[c + d*x])^3, x]`

Rubi [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^3 dx$$

$$\downarrow 6661$$

$$\frac{\int \left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^m (a + b \operatorname{arctanh}(c + dx))^3 d(c + dx)}{d}$$

$$\downarrow 6651$$

$$\frac{\int \left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^m (a + b \operatorname{arctanh}(c + dx))^3 d(c + dx)}{d}$$

input `Int[(e + f*x)^m*(a + b*ArcTanh[c + d*x])^3,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (fx + e)^m (a + b \operatorname{arctanh}(dx + c))^3 dx$$

input `int((f*x+e)^m*(a+b*arctanh(d*x+c))^3,x)`

output `int((f*x+e)^m*(a+b*arctanh(d*x+c))^3,x)`

Fricas [N/A]

Not integrable

Time = 0.10 (sec) , antiderivative size = 52, normalized size of antiderivative = 2.60

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^3 dx = \int (b \operatorname{artanh}(dx + c) + a)^3 (fx + e)^m dx$$

input `integrate((f*x+e)^m*(a+b*arctanh(d*x+c))^3,x, algorithm="fricas")`

output `integral((b^3*arctanh(d*x + c)^3 + 3*a*b^2*arctanh(d*x + c)^2 + 3*a^2*b*arctanh(d*x + c) + a^3)*(f*x + e)^m, x)`

Sympy [F(-1)]

Timed out.

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^3 dx = \text{Timed out}$$

input `integrate((f*x+e)**m*(a+b*atanh(d*x+c))**3,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 3.61 (sec) , antiderivative size = 432, normalized size of antiderivative = 21.60

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^3 dx = \int (b \operatorname{artanh}(dx + c) + a)^3 (fx + e)^m dx$$

input `integrate((f*x+e)^m*(a+b*arctanh(d*x+c))^3,x, algorithm="maxima")`

output

```
-1/8*(b^3*f*x + b^3*e)*(f*x + e)^m*log(-d*x - c + 1)^3/(f*(m + 1)) + (f*x
+ e)^(m + 1)*a^3/(f*(m + 1)) + integrate(1/8*((b^3*d*f*(m + 1)*x + (c*f*(m
+ 1) - f*(m + 1))*b^3)*log(d*x + c + 1)^3 + 6*(a*b^2*d*f*(m + 1)*x + (c*f
*(m + 1) - f*(m + 1))*a*b^2)*log(d*x + c + 1)^2 + 3*(b^3*d*e + 2*(c*f*(m +
1) - f*(m + 1))*a*b^2 + (2*a*b^2*d*f*(m + 1) + b^3*d*f)*x + (b^3*d*f*(m +
1)*x + (c*f*(m + 1) - f*(m + 1))*b^3)*log(d*x + c + 1))*log(-d*x - c + 1)
^2 + 12*(a^2*b*d*f*(m + 1)*x + (c*f*(m + 1) - f*(m + 1))*a^2*b)*log(d*x +
c + 1) - 3*(4*a^2*b*d*f*(m + 1)*x + 4*(c*f*(m + 1) - f*(m + 1))*a^2*b + (b
^3*d*f*(m + 1)*x + (c*f*(m + 1) - f*(m + 1))*b^3)*log(d*x + c + 1)^2 + 4*(
a*b^2*d*f*(m + 1)*x + (c*f*(m + 1) - f*(m + 1))*a*b^2)*log(d*x + c + 1))*l
og(-d*x - c + 1))*(f*x + e)^m/(d*f*(m + 1)*x + c*f*(m + 1) - f*(m + 1)), x
)
```

Giac [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^3 dx = \int (b \operatorname{artanh}(dx + c) + a)^3 (fx + e)^m dx$$

input

```
integrate((f*x+e)^m*(a+b*arctanh(d*x+c))^3,x, algorithm="giac")
```

output

```
integrate((b*arctanh(d*x + c) + a)^3*(f*x + e)^m, x)
```

Mupad [N/A]

Not integrable

Time = 3.69 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^3 dx = \int (e + fx)^m (a + b \operatorname{atanh}(c + dx))^3 dx$$

input

```
int((e + f*x)^m*(a + b*atanh(c + d*x))^3,x)
```


output `int((e + f*x)^m*(a + b*atanh(c + d*x))^3, x)`

Reduce [N/A]

Not integrable

Time = 0.44 (sec) , antiderivative size = 55220, normalized size of antiderivative = 2761.00

$$\int (e + fx)^m (a + \operatorname{barctanh}(c + dx))^3 dx = \text{Too large to display}$$

input `int((f*x+e)^m*(a+b*atanh(d*x+c))^3,x)`

output

```
(2*(e + f*x)**m*atanh(c + d*x)**3*b**3*c**5*d*e*f**m**3 + 2*(e + f*x)**m*atanh(c + d*x)**3*b**3*c**5*d*f**2*m**3*x - 2*(e + f*x)**m*atanh(c + d*x)**3*b**3*c**3*d*e*f**m**3 - 2*(e + f*x)**m*atanh(c + d*x)**3*b**3*c**3*d*f**2*m**3*x + 6*(e + f*x)**m*atanh(c + d*x)**2*a*b**2*c**5*d*e*f**m**3 + 6*(e + f*x)**m*atanh(c + d*x)**2*a*b**2*c**5*d*f**2*m**3*x - 6*(e + f*x)**m*atanh(c + d*x)**2*a*b**2*c**3*d*e*f**m**3 - 6*(e + f*x)**m*atanh(c + d*x)**2*a*b**2*c**3*d*f**2*m**3*x - 6*(e + f*x)**m*atanh(c + d*x)**2*b**3*c**5*f**2*m**2 + 6*(e + f*x)**m*atanh(c + d*x)**2*b**3*c**4*d*e*f**m**2 + 6*(e + f*x)**m*atanh(c + d*x)**2*b**3*c**3*d**2*e**2*m**2 + 6*(e + f*x)**m*atanh(c + d*x)**2*b**3*c**3*f**2*m**2 - 6*(e + f*x)**m*atanh(c + d*x)**2*b**3*c**2*d*e*f**m**2 + 6*(e + f*x)**m*atanh(c + d*x)*a**2*b*c**5*d*e*f**m**3 + 6*(e + f*x)**m*atanh(c + d*x)*a**2*b*c**5*d*f**2*m**3*x - 6*(e + f*x)**m*atanh(c + d*x)*a**2*b*c**3*d*e*f**m**3 - 6*(e + f*x)**m*atanh(c + d*x)*a**2*b*c**3*d*f**2*m**3*x - 12*(e + f*x)**m*atanh(c + d*x)*a*b**2*c**5*f**2*m**2 + 12*(e + f*x)**m*atanh(c + d*x)*a*b**2*c**4*d*e*f**m**2 + 12*(e + f*x)**m*atanh(c + d*x)*a*b**2*c**3*d**2*e**2*m**2 + 12*(e + f*x)**m*atanh(c + d*x)*a*b**2*c**3*f**2*m**2 - 12*(e + f*x)**m*atanh(c + d*x)*a*b**2*c**2*d*e*f**m**2 - 6*(e + f*x)**m*atanh(c + d*x)*b**3*c**4*f**2*m + 18*(e + f*x)**m*atanh(c + d*x)*b**3*c**3*d*e*f**m + 6*(e + f*x)**m*atanh(c + d*x)*b**3*c**2*d**2*e**2*m + 6*(e + f*x)**m*atanh(c + d*x)*b**3*c**2*f**2*m - 6*(e + f*x)**m...
```

3.51 $\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^2 dx$

| | |
|-------------------|-----|
| Optimal result | 465 |
| Mathematica [N/A] | 465 |
| Rubi [N/A] | 466 |
| Maple [N/A] | 466 |
| Fricas [N/A] | 467 |
| Sympy [F(-1)] | 467 |
| Maxima [N/A] | 467 |
| Giac [N/A] | 468 |
| Mupad [N/A] | 468 |
| Reduce [N/A] | 469 |

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^2 dx = \operatorname{Int}((e + fx)^m (a + b \operatorname{arctanh}(c + dx))^2, x)$$

output `Defer(Int)((f*x+e)^m*(a+b*arctanh(d*x+c))^2,x)`

Mathematica [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^2 dx$$

input `Integrate[(e + f*x)^m*(a + b*ArcTanh[c + d*x])^2,x]`

output `Integrate[(e + f*x)^m*(a + b*ArcTanh[c + d*x])^2, x]`

Rubi [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^2 dx$$

$$\downarrow 6661$$

$$\frac{\int \left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^m (a + b \operatorname{arctanh}(c + dx))^2 d(c + dx)}{d}$$

$$\downarrow 6651$$

$$\frac{\int \left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^m (a + b \operatorname{arctanh}(c + dx))^2 d(c + dx)}{d}$$

input `Int[(e + f*x)^m*(a + b*ArcTanh[c + d*x])^2,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (fx + e)^m (a + b \operatorname{arctanh}(dx + c))^2 dx$$

input `int((f*x+e)^m*(a+b*arctanh(d*x+c))^2,x)`

output `int((f*x+e)^m*(a+b*arctanh(d*x+c))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.80

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (b \operatorname{artanh}(dx + c) + a)^2 (fx + e)^m dx$$

input `integrate((f*x+e)^m*(a+b*arctanh(d*x+c))^2,x, algorithm="fricas")`

output `integral((b^2*arctanh(d*x + c)^2 + 2*a*b*arctanh(d*x + c) + a^2)*(f*x + e)^m, x)`

Sympy [F(-1)]

Timed out.

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^2 dx = \text{Timed out}$$

input `integrate((f*x+e)**m*(a+b*atanh(d*x+c))**2,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 2.18 (sec) , antiderivative size = 262, normalized size of antiderivative = 13.10

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (b \operatorname{artanh}(dx + c) + a)^2 (fx + e)^m dx$$

input `integrate((f*x+e)^m*(a+b*arctanh(d*x+c))^2,x, algorithm="maxima")`

output

```
1/4*(b^2*f*x + b^2*e)*(f*x + e)^m*log(-d*x - c + 1)^2/(f*(m + 1)) + (f*x +
e)^(m + 1)*a^2/(f*(m + 1)) - integrate(-1/4*((b^2*d*f*(m + 1)*x + (c*f*(m
+ 1) - f*(m + 1))*b^2)*log(d*x + c + 1)^2 + 4*(a*b*d*f*(m + 1)*x + (c*f*(
m + 1) - f*(m + 1))*a*b)*log(d*x + c + 1) - 2*(b^2*d*e + 2*(c*f*(m + 1) -
f*(m + 1))*a*b + (2*a*b*d*f*(m + 1) + b^2*d*f)*x + (b^2*d*f*(m + 1)*x + (c
*f*(m + 1) - f*(m + 1))*b^2)*log(d*x + c + 1))*log(-d*x - c + 1)*(f*x + e
)^m/(d*f*(m + 1)*x + c*f*(m + 1) - f*(m + 1)), x)
```

Giac [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (b \operatorname{artanh}(dx + c) + a)^2 (fx + e)^m dx$$

input

```
integrate((f*x+e)^m*(a+b*arctanh(d*x+c))^2,x, algorithm="giac")
```

output

```
integrate((b*arctanh(d*x + c) + a)^2*(f*x + e)^m, x)
```

Mupad [N/A]

Not integrable

Time = 3.63 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^2 dx = \int (e + fx)^m (a + b \operatorname{atanh}(c + dx))^2 dx$$

input

```
int((e + f*x)^m*(a + b*atanh(c + d*x))^2,x)
```

output

```
int((e + f*x)^m*(a + b*atanh(c + d*x))^2, x)
```

Reduce [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 22801, normalized size of antiderivative = 1140.05

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx))^2 dx = \text{Too large to display}$$

input `int((f*x+e)^m*(a+b*atanh(d*x+c))^2,x)`

output

```
((e + f*x)**m*atanh(c + d*x)**2*b**2*c**4*d*e*f*m**2 + (e + f*x)**m*atanh(c + d*x)**2*b**2*c**4*d*f**2*m**2*x - (e + f*x)**m*atanh(c + d*x)**2*b**2*c**2*d*e*f*m**2 - (e + f*x)**m*atanh(c + d*x)**2*b**2*c**2*d*f**2*m**2*x + 2*(e + f*x)**m*atanh(c + d*x)*a*b*c**4*d*e*f*m**2 + 2*(e + f*x)**m*atanh(c + d*x)*a*b*c**4*d*f**2*m**2*x - 2*(e + f*x)**m*atanh(c + d*x)*a*b*c**2*d*e*f*m**2 - 2*(e + f*x)**m*atanh(c + d*x)*a*b*c**2*d*f**2*m**2*x - 2*(e + f*x)**m*atanh(c + d*x)*b**2*c**4*f**2*m + 2*(e + f*x)**m*atanh(c + d*x)*b**2*c**3*d*e*f*m + 2*(e + f*x)**m*atanh(c + d*x)*b**2*c**2*d**2*e**2*m + 2*(e + f*x)**m*atanh(c + d*x)*b**2*c**2*f**2*m - 2*(e + f*x)**m*atanh(c + d*x)*b**2*c*d*e*f*m + (e + f*x)**m*a**2*c**4*d*e*f*m**2 + (e + f*x)**m*a**2*c**4*d*f**2*m**2*x - (e + f*x)**m*a**2*c**2*d*e*f*m**2 - (e + f*x)**m*a**2*c**2*d*f**2*m**2*x + 2*(e + f*x)**m*a*b*c**4*f**2*m - 2*(e + f*x)**m*a*b*c**2*f**2*m - 2*(e + f*x)**m*b**2*c**3*f**2 + (e + f*x)**m*b**2*c**2*d*e*f + (e + f*x)**m*b**2*c*d**2*e**2 + 2*(e + f*x)**m*b**2*c*f**2 - (e + f*x)*m*b**2*d*e*f + int((e + f*x)**m/(c**4*e*m + c**4*e + c**4*f*m*x + c**4*f*x + 2*c**3*d*e*m*x + 2*c**3*d*e*x + 2*c**3*d*f*m*x**2 + 2*c**3*d*f*x**2 + c**2*d**2*e*m*x**2 + c**2*d**2*e*x**2 + c**2*d**2*f*m*x**3 + c**2*d**2*f*x**3 - 2*c**2*e*m - 2*c**2*e - 2*c**2*f*m*x - 2*c**2*f*x - 2*c*d*e*m*x - 2*c*d*e*x - 2*c*d*f*m*x**2 - 2*c*d*f*x**2 - d**2*e*m*x**2 - d**2*e*x**2 - d**2*f*m*x**3 - d**2*f*x**3 + e*m + e + f*m*x + f*x),x)*b**2*c**7*f**3*m...
```

3.52 $\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx)) dx$

| | |
|---------------------|-----|
| Optimal result | 470 |
| Mathematica [F] | 471 |
| Rubi [A] (verified) | 471 |
| Maple [F] | 473 |
| Fricas [F] | 473 |
| Sympy [F] | 473 |
| Maxima [F] | 474 |
| Giac [F] | 474 |
| Mupad [F(-1)] | 474 |
| Reduce [F] | 475 |

Optimal result

Integrand size = 18, antiderivative size = 162

$$\begin{aligned} & \int (e + fx)^m (a + b \operatorname{arctanh}(c + dx)) dx \\ &= \frac{(e + fx)^{1+m} (a + b \operatorname{arctanh}(c + dx))}{f(1+m)} \\ & \quad + \frac{bd(e + fx)^{2+m} \operatorname{Hypergeometric2F1}\left(1, 2+m, 3+m, \frac{d(e+fx)}{de-f-cf}\right)}{2f(de - (1+c)f)(1+m)(2+m)} \\ & \quad - \frac{bd(e + fx)^{2+m} \operatorname{Hypergeometric2F1}\left(1, 2+m, 3+m, \frac{d(e+fx)}{de+f-cf}\right)}{2f(de + f - cf)(1+m)(2+m)} \end{aligned}$$

output

```
(f*x+e)^(1+m)*(a+b*arctanh(d*x+c))/f/(1+m)+1/2*b*d*(f*x+e)^(2+m)*hypergeom
([1, 2+m],[3+m],d*(f*x+e)/(-c*f+d*e-f))/f/(d*e-(1+c)*f)/(1+m)/(2+m)-1/2*b*
d*(f*x+e)^(2+m)*hypergeom([1, 2+m],[3+m],d*(f*x+e)/(-c*f+d*e+f))/f/(-c*f+d
*e+f)/(1+m)/(2+m)
```

Mathematica [F]

$$\int (e + fx)^m (a + \operatorname{barctanh}(c + dx)) dx = \int (e + fx)^m (a + \operatorname{barctanh}(c + dx)) dx$$

input `Integrate[(e + f*x)^m*(a + b*ArcTanh[c + d*x]),x]`

output `Integrate[(e + f*x)^m*(a + b*ArcTanh[c + d*x]), x]`

Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.37, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {6661, 6478, 485, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (e + fx)^m (a + \operatorname{barctanh}(c + dx)) dx \\ & \quad \downarrow \text{6661} \\ & \frac{\int \left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^m (a + \operatorname{barctanh}(c + dx)) d(c + dx)}{d} \\ & \quad \downarrow \text{6478} \\ & \frac{d(a + \operatorname{barctanh}(c + dx)) \left(\frac{f(c+dx)}{d} - \frac{cf}{d} + e \right)^{m+1}}{f(m+1)} - \frac{bd \int \frac{\left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^{m+1}}{1 - (c+dx)^2} d(c+dx)}{f(m+1)} \\ & \quad \downarrow \text{485} \\ & \frac{d(a + \operatorname{barctanh}(c + dx)) \left(\frac{f(c+dx)}{d} - \frac{cf}{d} + e \right)^{m+1}}{f(m+1)} - \frac{bd \int \left(\frac{\left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^{m+1}}{2(-c-dx+1)} + \frac{\left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^{m+1}}{2(c+dx+1)} \right) d(c+dx)}{f(m+1)} \\ & \quad \downarrow \text{2009} \end{aligned}$$

$$\frac{d(a+b\operatorname{arctanh}(c+dx))\left(\frac{f(c+dx)}{d}-\frac{cf}{d}+e\right)^{m+1}}{f^{(m+1)}} - \frac{bd\left(\frac{d\left(\frac{f(c+dx)}{d}-\frac{cf}{d}+e\right)^{m+2}\operatorname{Hypergeometric2F1}\left(1,m+2,m+3,\frac{de-cf+f(c+dx)}{de-cf+f}\right)}{2(m+2)(-cf+de+f)} - \frac{d\left(\frac{f(c+dx)}{d}-\frac{cf}{d}+e\right)^{m+2}}{2(m+2)(-cf+de+f)}\right)}{d f^{(m+1)}}$$

input `Int[(e + f*x)^m*(a + b*ArcTanh[c + d*x]),x]`

output `((d*(e - (c*f)/d + (f*(c + d*x))/d)^(1 + m)*(a + b*ArcTanh[c + d*x]))/(f*(1 + m)) - (b*d*(-1/2*(d*(e - (c*f)/d + (f*(c + d*x))/d)^(2 + m)*Hypergeometric2F1[1, 2 + m, 3 + m, (d*e - c*f + f*(c + d*x))/(d*e - f - c*f)]/(d*e - (1 + c)*f)*(2 + m)) + (d*(e - (c*f)/d + (f*(c + d*x))/d)^(2 + m)*Hypergeometric2F1[1, 2 + m, 3 + m, (d*e - c*f + f*(c + d*x))/(d*e + f - c*f)]/(2*(d*e + f - c*f)*(2 + m))))/(f*(1 + m))/d`

Defintions of rubi rules used

rule 485 `Int[((c_) + (d_)*(x_))^(n_)/((a_) + (b_)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^n, 1/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, n}, x] && !IntegerQ[2*n]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6478 `Int[((a_) + ArcTanh[(c_)*(x_)]*(b_.))*((d_) + (e_)*(x_))^(q_), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTanh[c*x])/(e*(q + 1))), x] - Simp[b*(c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 - c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[q, -1]`

rule 6661 `Int[((a_) + ArcTanh[(c_) + (d_)*(x_)]*(b_.))^(p_)*((e_) + (f_)*(x_))^(m_), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[p, 0]`

Maple [F]

$$\int (fx + e)^m (a + b \operatorname{arctanh}(dx + c)) dx$$

input `int((f*x+e)^m*(a+b*arctanh(d*x+c)),x)`

output `int((f*x+e)^m*(a+b*arctanh(d*x+c)),x)`

Fricas [F]

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx)) dx = \int (b \operatorname{artanh}(dx + c) + a)(fx + e)^m dx$$

input `integrate((f*x+e)^m*(a+b*arctanh(d*x+c)),x, algorithm="fricas")`

output `integral((b*arctanh(d*x + c) + a)*(f*x + e)^m, x)`

Sympy [F]

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx)) dx = \int (a + b \operatorname{atanh}(c + dx))(e + fx)^m dx$$

input `integrate((f*x+e)**m*(a+b*atanh(d*x+c)),x)`

output `Integral((a + b*atanh(c + d*x))*(e + f*x)**m, x)`

Maxima [F]

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx)) dx = \int (b \operatorname{artanh}(dx + c) + a)(fx + e)^m dx$$

input `integrate((f*x+e)^m*(a+b*arctanh(d*x+c)),x, algorithm="maxima")`

output `-1/2*b*((f*x + e)*(f*x + e)^m*log(-d*x - c + 1)/(f*(m + 1)) - integrate((d*f*x + d*e + (d*f*(m + 1)*x + c*f*(m + 1) - f*(m + 1))*log(d*x + c + 1))*(f*x + e)^m/(d*f*(m + 1)*x + c*f*(m + 1) - f*(m + 1)), x)) + (f*x + e)^(m + 1)*a/(f*(m + 1))`

Giac [F]

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx)) dx = \int (b \operatorname{artanh}(dx + c) + a)(fx + e)^m dx$$

input `integrate((f*x+e)^m*(a+b*arctanh(d*x+c)),x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)*(f*x + e)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx)) dx = \int (e + fx)^m (a + b \operatorname{atanh}(c + dx)) dx$$

input `int((e + f*x)^m*(a + b*atanh(c + d*x)),x)`

output `int((e + f*x)^m*(a + b*atanh(c + d*x)), x)`

Reduce [F]

$$\int (e + fx)^m (a + b \operatorname{arctanh}(c + dx)) dx = \text{Too large to display}$$

input `int((f*x+e)^m*(a+b*atanh(d*x+c)),x)`

output

```
((e + f*x)**m*atanh(c + d*x)*b*c*e**m + (e + f*x)**m*atanh(c + d*x)*b*c*f*m
*x + (e + f*x)**m*a*c*e**m + (e + f*x)**m*a*c*f*m*x + (e + f*x)**m*b*e - in
t((e + f*x)**m/(c**2*e**m + c**2*e + c**2*f*m*x + c**2*f*x + 2*c*d*e**m*x +
2*c*d*e*x + 2*c*d*f*m*x**2 + 2*c*d*f*x**2 + d**2*e**m*x**2 + d**2*e*x**2 +
d**2*f*m*x**3 + d**2*f*x**3 - e*m - e - f*m*x - f*x),x)*b*c**2*e*f*m**2 -
int((e + f*x)**m/(c**2*e**m + c**2*e + c**2*f*m*x + c**2*f*x + 2*c*d*e**m*x
+ 2*c*d*e*x + 2*c*d*f*m*x**2 + 2*c*d*f*x**2 + d**2*e**m*x**2 + d**2*e*x**2
+ d**2*f*m*x**3 + d**2*f*x**3 - e*m - e - f*m*x - f*x),x)*b*c**2*e*f*m + i
nt((e + f*x)**m/(c**2*e**m + c**2*e + c**2*f*m*x + c**2*f*x + 2*c*d*e**m*x +
2*c*d*e*x + 2*c*d*f*m*x**2 + 2*c*d*f*x**2 + d**2*e**m*x**2 + d**2*e*x**2 +
d**2*f*m*x**3 + d**2*f*x**3 - e*m - e - f*m*x - f*x),x)*b*c*d*e**2*m**2 +
int((e + f*x)**m/(c**2*e**m + c**2*e + c**2*f*m*x + c**2*f*x + 2*c*d*e**m*x
+ 2*c*d*e*x + 2*c*d*f*m*x**2 + 2*c*d*f*x**2 + d**2*e**m*x**2 + d**2*e*x**2
+ d**2*f*m*x**3 + d**2*f*x**3 - e*m - e - f*m*x - f*x),x)*b*c*d*e**2*m +
int((e + f*x)**m/(c**2*e**m + c**2*e + c**2*f*m*x + c**2*f*x + 2*c*d*e**m*x
+ 2*c*d*e*x + 2*c*d*f*m*x**2 + 2*c*d*f*x**2 + d**2*e**m*x**2 + d**2*e*x**2
+ d**2*f*m*x**3 + d**2*f*x**3 - e*m - e - f*m*x - f*x),x)*b*e*f*m**2 + int
((e + f*x)**m/(c**2*e**m + c**2*e + c**2*f*m*x + c**2*f*x + 2*c*d*e**m*x + 2
*c*d*e*x + 2*c*d*f*m*x**2 + 2*c*d*f*x**2 + d**2*e**m*x**2 + d**2*e*x**2 + d
**2*f*m*x**3 + d**2*f*x**3 - e*m - e - f*m*x - f*x),x)*b*e*f*m + int(((...
```

3.53 $\int \frac{\operatorname{arctanh}(a+bx)}{c+dx^3} dx$

| | |
|---------------------------------------|-----|
| Optimal result | 477 |
| Mathematica [A] (verified) | 478 |
| Rubi [A] (verified) | 479 |
| Maple [C] (warning: unable to verify) | 482 |
| Fricas [F] | 483 |
| Sympy [F(-1)] | 483 |
| Maxima [F] | 484 |
| Giac [F] | 484 |
| Mupad [F(-1)] | 484 |
| Reduce [F] | 485 |

Optimal result

Integrand size = 16, antiderivative size = 649

$$\begin{aligned}
\int \frac{\operatorname{arctanh}(a+bx)}{c+dx^3} dx = & -\frac{\operatorname{arctanh}(a+bx) \log\left(\frac{2}{1+a+bx}\right)}{3c^{2/3}\sqrt[3]{d}} \\
& + \frac{\sqrt[3]{-1}\operatorname{arctanh}(a+bx) \log\left(\frac{2}{1+a+bx}\right)}{3c^{2/3}\sqrt[3]{d}} \\
& - \frac{(-1)^{2/3}\operatorname{arctanh}(a+bx) \log\left(\frac{2}{1+a+bx}\right)}{3c^{2/3}\sqrt[3]{d}} \\
& + \frac{\operatorname{arctanh}(a+bx) \log\left(\frac{2b\left(\sqrt[3]{c}+\sqrt[3]{dx}\right)}{\left(b\sqrt[3]{c}+(1-a)\sqrt[3]{d}\right)(1+a+bx)}\right)}{3c^{2/3}\sqrt[3]{d}} \\
& + \frac{(-1)^{2/3}\operatorname{arctanh}(a+bx) \log\left(\frac{2b\left(\sqrt[3]{c}-\sqrt[3]{-1}\sqrt[3]{dx}\right)}{\left(b\sqrt[3]{c}-\sqrt[3]{-1}(1-a)\sqrt[3]{d}\right)(1+a+bx)}\right)}{3c^{2/3}\sqrt[3]{d}} \\
& - \frac{\sqrt[3]{-1}\operatorname{arctanh}(a+bx) \log\left(\frac{2b\left(\sqrt[3]{c}+(-1)^{2/3}\sqrt[3]{dx}\right)}{\left(b\sqrt[3]{c}+(-1)^{2/3}(1-a)\sqrt[3]{d}\right)(1+a+bx)}\right)}{3c^{2/3}\sqrt[3]{d}} \\
& + \frac{\operatorname{PolyLog}\left(2, 1-\frac{2}{1+a+bx}\right)}{6c^{2/3}\sqrt[3]{d}} - \frac{\sqrt[3]{-1}\operatorname{PolyLog}\left(2, 1-\frac{2}{1+a+bx}\right)}{6c^{2/3}\sqrt[3]{d}} \\
& + \frac{(-1)^{2/3}\operatorname{PolyLog}\left(2, 1-\frac{2}{1+a+bx}\right)}{6c^{2/3}\sqrt[3]{d}} \\
& - \frac{\operatorname{PolyLog}\left(2, 1-\frac{2b\left(\sqrt[3]{c}+\sqrt[3]{dx}\right)}{\left(b\sqrt[3]{c}+(1-a)\sqrt[3]{d}\right)(1+a+bx)}\right)}{6c^{2/3}\sqrt[3]{d}} \\
& - \frac{(-1)^{2/3}\operatorname{PolyLog}\left(2, 1-\frac{2b\left(\sqrt[3]{c}-\sqrt[3]{-1}\sqrt[3]{dx}\right)}{\left(b\sqrt[3]{c}-\sqrt[3]{-1}(1-a)\sqrt[3]{d}\right)(1+a+bx)}\right)}{6c^{2/3}\sqrt[3]{d}} \\
& - \frac{\sqrt[3]{-1}\operatorname{PolyLog}\left(2, 1-\frac{2b\left(\sqrt[3]{c}+(-1)^{2/3}\sqrt[3]{dx}\right)}{\left(b\sqrt[3]{c}+(-1)^{2/3}(1-a)\sqrt[3]{d}\right)(1+a+bx)}\right)}{6c^{2/3}\sqrt[3]{d}}
\end{aligned}$$

output

```

-1/3*arctanh(b*x+a)*ln(2/(b*x+a+1))/c^(2/3)/d^(1/3)+1/3*(-1)^(1/3)*arctanh
(b*x+a)*ln(2/(b*x+a+1))/c^(2/3)/d^(1/3)-1/3*(-1)^(2/3)*arctanh(b*x+a)*ln(2
/(b*x+a+1))/c^(2/3)/d^(1/3)+1/3*arctanh(b*x+a)*ln(2*b*(c^(1/3)+d^(1/3)*x)/
(b*c^(1/3)+(1-a)*d^(1/3)))/(b*x+a+1))/c^(2/3)/d^(1/3)+1/3*(-1)^(2/3)*arctan
h(b*x+a)*ln(2*b*(c^(1/3)-(-1)^(1/3)*d^(1/3)*x)/(b*c^(1/3)-(-1)^(1/3)*(1-a)
*d^(1/3)))/(b*x+a+1))/c^(2/3)/d^(1/3)-1/3*(-1)^(1/3)*arctanh(b*x+a)*ln(2*b*
(c^(1/3)+(-1)^(2/3)*d^(1/3)*x)/(b*c^(1/3)+(-1)^(2/3)*(1-a)*d^(1/3)))/(b*x+a
+1))/c^(2/3)/d^(1/3)+1/6*polylog(2,1-2/(b*x+a+1))/c^(2/3)/d^(1/3)-1/6*(-1)
^(1/3)*polylog(2,1-2/(b*x+a+1))/c^(2/3)/d^(1/3)+1/6*(-1)^(2/3)*polylog(2,1
-2/(b*x+a+1))/c^(2/3)/d^(1/3)-1/6*polylog(2,1-2*b*(c^(1/3)+d^(1/3)*x)/(b*c
^(1/3)+(1-a)*d^(1/3)))/(b*x+a+1))/c^(2/3)/d^(1/3)-1/6*(-1)^(2/3)*polylog(2,
1-2*b*(c^(1/3)-(-1)^(1/3)*d^(1/3)*x)/(b*c^(1/3)-(-1)^(1/3)*(1-a)*d^(1/3)))/
(b*x+a+1))/c^(2/3)/d^(1/3)+1/6*(-1)^(1/3)*polylog(2,1-2*b*(c^(1/3)+(-1)^(2
/3)*d^(1/3)*x)/(b*c^(1/3)+(-1)^(2/3)*(1-a)*d^(1/3)))/(b*x+a+1))/c^(2/3)/d^(
1/3)

```

Mathematica [A] (verified)

Time = 0.48 (sec) , antiderivative size = 623, normalized size of antiderivative = 0.96

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^3} dx$$

$$= \frac{-\log(1 - a - bx) \log\left(\frac{b(\sqrt[3]{c} + \sqrt[3]{dx})}{b\sqrt[3]{c} - (-1+a)\sqrt[3]{d}}\right) + \log(1 + a + bx) \log\left(\frac{b(\sqrt[3]{c} + \sqrt[3]{dx})}{b\sqrt[3]{c} - (-1+a)\sqrt[3]{d}}\right) - (-1)^{2/3} \log(1 - a - bx) \log\left(\frac{b(\sqrt[3]{c} + \sqrt[3]{dx})}{b\sqrt[3]{c} - (-1+a)\sqrt[3]{d}}\right) + (-1)^{1/3} \log(1 + a + bx) \log\left(\frac{b(\sqrt[3]{c} + \sqrt[3]{dx})}{b\sqrt[3]{c} - (-1+a)\sqrt[3]{d}}\right)}{c + dx^3}$$

input

```
Integrate[ArcTanh[a + b*x]/(c + d*x^3), x]
```

output

```
(-Log[1 - a - b*x]*Log[(b*(c^(1/3) + d^(1/3)*x))/(b*c^(1/3) - (-1 + a)*d^(1/3))] + Log[1 + a + b*x]*Log[(b*(c^(1/3) + d^(1/3)*x))/(b*c^(1/3) - (1 + a)*d^(1/3))] - (-1)^(2/3)*Log[1 - a - b*x]*Log[(b*(c^(1/3) - (-1)^(1/3)*d^(1/3)*x))/(b*c^(1/3) + (-1)^(1/3)*(-1 + a)*d^(1/3))] + (-1)^(2/3)*Log[1 + a + b*x]*Log[(b*(c^(1/3) - (-1)^(1/3)*d^(1/3)*x))/(b*c^(1/3) + (-1)^(1/3)*(1 + a)*d^(1/3))] + (-1)^(1/3)*Log[1 - a - b*x]*Log[(b*(c^(1/3) + (-1)^(2/3)*d^(1/3)*x))/(b*c^(1/3) - (-1)^(2/3)*d^(1/3)*x)] - (-1)^(1/3)*Log[1 + a + b*x]*Log[(b*(c^(1/3) + (-1)^(2/3)*d^(1/3)*x))/(b*c^(1/3) - (-1)^(2/3)*(1 + a)*d^(1/3))] - PolyLog[2, -((d^(1/3)*(-1 + a + b*x))/(b*c^(1/3) - (-1 + a)*d^(1/3))] - (-1)^(2/3)*PolyLog[2, ((-1)^(1/3)*d^(1/3)*(-1 + a + b*x))/(b*c^(1/3) + (-1)^(1/3)*(-1 + a)*d^(1/3))] + (-1)^(1/3)*PolyLog[2, ((-1)^(2/3)*d^(1/3)*(-1 + a + b*x))/(-b*c^(1/3) + (-1)^(2/3)*(-1 + a)*d^(1/3))] + PolyLog[2, -((d^(1/3)*(1 + a + b*x))/(b*c^(1/3) - (1 + a)*d^(1/3))] + (-1)^(2/3)*PolyLog[2, ((-1)^(1/3)*d^(1/3)*(1 + a + b*x))/(b*c^(1/3) + (-1)^(1/3)*(1 + a)*d^(1/3))] - (-1)^(1/3)*PolyLog[2, ((-1)^(2/3)*d^(1/3)*(1 + a + b*x))/(-b*c^(1/3) + (-1)^(2/3)*(1 + a)*d^(1/3))]/(6*c^(2/3)*d^(1/3))
```

Rubi [A] (verified)

Time = 1.56 (sec) , antiderivative size = 800, normalized size of antiderivative = 1.23, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6665, 2856, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^3} dx$$

$$\downarrow \text{6665}$$

$$\frac{1}{2} \int \frac{\log(a + bx + 1)}{dx^3 + c} dx - \frac{1}{2} \int \frac{\log(-a - bx + 1)}{dx^3 + c} dx$$

$$\downarrow \text{2856}$$

$$\frac{1}{2} \int \left(-\frac{\log(a + bx + 1)}{3c^{2/3}(-\sqrt[3]{dx} - \sqrt[3]{c})} - \frac{\log(a + bx + 1)}{3c^{2/3}(\sqrt[3]{-1}\sqrt[3]{dx} - \sqrt[3]{c})} - \frac{\log(a + bx + 1)}{3c^{2/3}(-(-1)^{2/3}\sqrt[3]{dx} - \sqrt[3]{c})} \right) dx -$$

$$\frac{1}{2} \int \left(-\frac{\log(-a - bx + 1)}{3c^{2/3}(-\sqrt[3]{dx} - \sqrt[3]{c})} - \frac{\log(-a - bx + 1)}{3c^{2/3}(\sqrt[3]{-1}\sqrt[3]{dx} - \sqrt[3]{c})} - \frac{\log(-a - bx + 1)}{3c^{2/3}(-(-1)^{2/3}\sqrt[3]{dx} - \sqrt[3]{c})} \right) dx$$

↓ 2009

$$\frac{1}{2} \left(-\frac{\log(-a - bx + 1) \log\left(\frac{b(\sqrt[3]{dx} + \sqrt[3]{c})}{\sqrt[3]{d}(1-a) + b\sqrt[3]{c}}\right)}{3c^{2/3}\sqrt[3]{d}} - \frac{(-1)^{2/3} \log(-a - bx + 1) \log\left(\frac{b(\sqrt[3]{c} - \sqrt[3]{-1}\sqrt[3]{dx})}{b\sqrt[3]{c} - \sqrt[3]{-1}(1-a)\sqrt[3]{d}}\right)}{3c^{2/3}\sqrt[3]{d}} + \frac{\sqrt[3]{-1} \log(-a - bx + 1) \log\left(\frac{b(\sqrt[3]{c} - \sqrt[3]{-1}\sqrt[3]{dx})}{b\sqrt[3]{c} - \sqrt[3]{-1}(1-a)\sqrt[3]{d}}\right)}{3c^{2/3}\sqrt[3]{d}} \right) +$$

$$\frac{1}{2} \left(\frac{\log(a + bx + 1) \log\left(\frac{b(\sqrt[3]{dx} + \sqrt[3]{c})}{b\sqrt[3]{c} - (a+1)\sqrt[3]{d}}\right)}{3c^{2/3}\sqrt[3]{d}} + \frac{(-1)^{2/3} \log(a + bx + 1) \log\left(\frac{b(\sqrt[3]{c} - \sqrt[3]{-1}\sqrt[3]{dx})}{\sqrt[3]{-1}\sqrt[3]{d}(a+1) + b\sqrt[3]{c}}\right)}{3c^{2/3}\sqrt[3]{d}} - \frac{\sqrt[3]{-1} \log(a + bx + 1) \log\left(\frac{b(\sqrt[3]{c} - \sqrt[3]{-1}\sqrt[3]{dx})}{\sqrt[3]{-1}\sqrt[3]{d}(a+1) + b\sqrt[3]{c}}\right)}{3c^{2/3}\sqrt[3]{d}} \right)$$

input Int[ArcTanh[a + b*x]/(c + d*x^3),x]

output

$$\begin{aligned} & (-1/3 * (\text{Log}[1 - a - b*x] * \text{Log}[(b*(c^{1/3}) + d^{1/3}*x))/(b*c^{1/3} + (1 - a) \\ & * d^{1/3}])) / (c^{2/3} * d^{1/3}) - ((-1)^{2/3} * \text{Log}[1 - a - b*x] * \text{Log}[(b*(c^{1/3}) \\ & - (-1)^{1/3} * d^{1/3} * x]) / (b*c^{1/3} - (-1)^{1/3} * (1 - a) * d^{1/3}])) / (3 * \\ & c^{2/3} * d^{1/3}) + ((-1)^{1/3} * \text{Log}[1 - a - b*x] * \text{Log}[(b*(c^{1/3}) + (-1)^{2/3} \\ & * d^{1/3} * x]) / (b*c^{1/3} + (-1)^{2/3} * (1 - a) * d^{1/3}])) / (3 * c^{2/3} * d^{1/3}) \\ & - \text{PolyLog}[2, (d^{1/3} * (1 - a - b*x)) / (b*c^{1/3} + (1 - a) * d^{1/3})] / (3 * \\ & c^{2/3} * d^{1/3}) - ((-1)^{2/3} * \text{PolyLog}[2, -(((-1)^{1/3} * d^{1/3} * (1 - a - \\ & b*x)) / (b*c^{1/3} - (-1)^{1/3} * (1 - a) * d^{1/3}))]) / (3 * c^{2/3} * d^{1/3}) + ((\\ & -1)^{1/3} * \text{PolyLog}[2, ((-1)^{2/3} * d^{1/3} * (1 - a - b*x)) / (b*c^{1/3} + (-1)^{ \\ & 2/3} * (1 - a) * d^{1/3})]) / (3 * c^{2/3} * d^{1/3})) / 2 + ((\text{Log}[1 + a + b*x] * \text{Log}[(\\ & b*(c^{1/3}) + d^{1/3} * x]) / (b*c^{1/3} - (1 + a) * d^{1/3})]) / (3 * c^{2/3} * d^{1/3} \\ &)) + ((-1)^{2/3} * \text{Log}[1 + a + b*x] * \text{Log}[(b*(c^{1/3}) - (-1)^{1/3} * d^{1/3} * x)) \\ & / (b*c^{1/3} + (-1)^{1/3} * (1 + a) * d^{1/3})]) / (3 * c^{2/3} * d^{1/3}) - ((-1)^{1/3} * \\ & \text{Log}[1 + a + b*x] * \text{Log}[(b*(c^{1/3}) + (-1)^{2/3} * d^{1/3} * x]) / (b*c^{1/3} - \\ & (-1)^{2/3} * (1 + a) * d^{1/3})]) / (3 * c^{2/3} * d^{1/3}) + \text{PolyLog}[2, -((d^{1/3}) \\ & * (1 + a + b*x)) / (b*c^{1/3} - (1 + a) * d^{1/3})]) / (3 * c^{2/3} * d^{1/3}) + ((-1) \\ & ^{2/3} * \text{PolyLog}[2, ((-1)^{1/3} * d^{1/3} * (1 + a + b*x)) / (b*c^{1/3} + (-1)^{1/3} * \\ & (1 + a) * d^{1/3})]) / (3 * c^{2/3} * d^{1/3}) - ((-1)^{1/3} * \text{PolyLog}[2, -((d^{1/3}) \\ & * ((-1)^{2/3} + (-1)^{2/3} * a + (-1)^{2/3} * b*x)) / (b*c^{1/3} - (-1)^{2/3} * \\ & (1 + a) * d^{1/3})]) / (3 * c^{2/3} * d^{1/3})) / 2 \end{aligned}$$

Defintions of rubi rules used

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 2856 $\text{Int}[(a_.) + \text{Log}[(c_.) * ((d_.) + (e_.) * (x_.)^{n_.}) * (b_.)^{p_.} * ((f_.) + (g_.) * (x_.)^{r_.})^{q_.}], x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b * \text{Log}[c * (d + e * x)^n])^p, (f + g * x^r)^q, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, g, n, r\}, x\} \&\& \text{I GtQ}[p, 0] \&\& \text{IntegerQ}[q] \&\& (\text{GtQ}[q, 0] \|\| (\text{IntegerQ}[r] \&\& \text{NeQ}[r, 1]))$

rule 6665 $\text{Int}[\text{ArcTanh}[(c_.) + (d_.) * (x_.)] / ((e_.) + (f_.) * (x_.)^{n_.}), x_Symbol] \rightarrow \text{Simp}[1/2 \text{Int}[\text{Log}[1 + c + d * x] / (e + f * x^n), x], x] - \text{Simp}[1/2 \text{Int}[\text{Log}[1 - c - d * x] / (e + f * x^n), x], x] /; \text{FreeQ}\{c, d, e, f\}, x\} \&\& \text{RationalQ}[n]$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.54 (sec) , antiderivative size = 259, normalized size of antiderivative = 0.40

| method | result |
|-------------------|---|
| risch | $\frac{b^2 \left(\frac{\ln(-bx-a+1) \ln\left(\frac{bx+R1+a}{R1}\right)}{R1^2+2R1a+} \right)}{6d}$ |
| derivativedivides | $\frac{b^3 \left(\frac{\ln\left(\frac{bx-R+a}{R^2+2Ra-a^2}\right) \operatorname{arctanh}(bx+a)}{3d} \right)}{+ \left(\frac{b^3 \operatorname{arctanh}(bx+a)}{R=1} \right)}$ |
| default | $\frac{b^3 \left(\frac{\ln\left(\frac{bx-R+a}{R^2+2Ra-a^2}\right) \operatorname{arctanh}(bx+a)}{3d} \right)}{+ \left(\frac{b^3 \operatorname{arctanh}(bx+a)}{R=1} \right)}$ |

```
input int(arctanh(b*x+a)/(d*x^3+c),x,method=_RETURNVERBOSE)
```

output

```
-1/6*b^2/d*sum(1/(_R1^2+2*_R1*a+a^2-2*_R1-2*a+1)*(ln(-b*x-a+1)*ln((b*x+_R1+a-1)/_R1)+dilog((b*x+_R1+a-1)/_R1)),_R1=RootOf(d*_Z^3+(3*a*d-3*d)*_Z^2+(3*a^2*d-6*a*d+3*d)*_Z+a^3*d-c*b^3-3*a^2*d+3*a*d-d))+1/6*b^2/d*sum(1/(_R1^2-2*_R1*a+a^2-2*_R1+2*a+1)*(ln(b*x+a+1)*ln((-b*x+_R1-a-1)/_R1)+dilog((-b*x+_R1-a-1)/_R1)),_R1=RootOf(d*_Z^3+(-3*a*d-3*d)*_Z^2+(3*a^2*d+6*a*d+3*d)*_Z-a^3*d+c*b^3-3*a^2*d-3*a*d-d))
```

Fricas [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^3} dx = \int \frac{\operatorname{artanh}(bx + a)}{dx^3 + c} dx$$

input

```
integrate(arctanh(b*x+a)/(d*x^3+c),x, algorithm="fricas")
```

output

```
integral(arctanh(b*x + a)/(d*x^3 + c), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^3} dx = \text{Timed out}$$

input

```
integrate(atanh(b*x+a)/(d*x**3+c),x)
```

output

```
Timed out
```

Maxima [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^3} dx = \int \frac{\operatorname{artanh}(bx + a)}{dx^3 + c} dx$$

input `integrate(arctanh(b*x+a)/(d*x^3+c),x, algorithm="maxima")`

output `integrate(arctanh(b*x + a)/(d*x^3 + c), x)`

Giac [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^3} dx = \int \frac{\operatorname{artanh}(bx + a)}{dx^3 + c} dx$$

input `integrate(arctanh(b*x+a)/(d*x^3+c),x, algorithm="giac")`

output `sage0*x`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^3} dx = \int \frac{\operatorname{atanh}(a + bx)}{dx^3 + c} dx$$

input `int(atanh(a + b*x)/(c + d*x^3),x)`

output `int(atanh(a + b*x)/(c + d*x^3), x)`

Reduce [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^3} dx = \int \frac{\operatorname{atanh}(bx + a)}{dx^3 + c} dx$$

input `int(atanh(b*x+a)/(d*x^3+c),x)`

output `int(atanh(a + b*x)/(c + d*x**3),x)`

3.54 $\int \frac{\operatorname{arctanh}(a+bx)}{c+dx^2} dx$

| | |
|---|-----|
| Optimal result | 486 |
| Mathematica [A] (warning: unable to verify) | 487 |
| Rubi [A] (warning: unable to verify) | 487 |
| Maple [A] (verified) | 489 |
| Fricas [F] | 490 |
| Sympy [F(-1)] | 490 |
| Maxima [C] (verification not implemented) | 490 |
| Giac [F] | 491 |
| Mupad [F(-1)] | 491 |
| Reduce [F] | 492 |

Optimal result

Integrand size = 16, antiderivative size = 287

$$\int \frac{\operatorname{arctanh}(a+bx)}{c+dx^2} dx = \frac{\operatorname{arctanh}(a+bx) \log\left(\frac{2b(\sqrt{-c}-\sqrt{dx})}{(b\sqrt{-c}-(1-a)\sqrt{d})(1+a+bx)}\right)}{2\sqrt{-c}\sqrt{d}} - \frac{\operatorname{arctanh}(a+bx) \log\left(\frac{2b(\sqrt{-c}+\sqrt{dx})}{(b\sqrt{-c}+(1-a)\sqrt{d})(1+a+bx)}\right)}{2\sqrt{-c}\sqrt{d}} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2b(\sqrt{-c}-\sqrt{dx})}{(b\sqrt{-c}-(1-a)\sqrt{d})(1+a+bx)}\right)}{4\sqrt{-c}\sqrt{d}} + \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2b(\sqrt{-c}+\sqrt{dx})}{(b\sqrt{-c}+(1-a)\sqrt{d})(1+a+bx)}\right)}{4\sqrt{-c}\sqrt{d}}$$

output

```
1/2*arctanh(b*x+a)*ln(2*b*((-c)^(1/2)-d^(1/2)*x)/(b*(-c)^(1/2)-(1-a)*d^(1/2)))/(b*x+a+1))/(-c)^(1/2)/d^(1/2)-1/2*arctanh(b*x+a)*ln(2*b*((-c)^(1/2)+d^(1/2)*x)/(b*(-c)^(1/2)+(1-a)*d^(1/2)))/(b*x+a+1))/(-c)^(1/2)/d^(1/2)-1/4*polylog(2,1-2*b*((-c)^(1/2)-d^(1/2)*x)/(b*(-c)^(1/2)-(1-a)*d^(1/2)))/(b*x+a+1))/(-c)^(1/2)/d^(1/2)+1/4*polylog(2,1-2*b*((-c)^(1/2)+d^(1/2)*x)/(b*(-c)^(1/2)+(1-a)*d^(1/2)))/(b*x+a+1))/(-c)^(1/2)/d^(1/2)
```

Mathematica [A] (warning: unable to verify)

Time = 0.20 (sec) , antiderivative size = 365, normalized size of antiderivative = 1.27

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^2} dx$$

$$= \frac{-\log(1 - a - bx) \log\left(\frac{b(\sqrt{-c} - \sqrt{dx})}{b\sqrt{-c} + (-1+a)\sqrt{d}}\right) + \log(1 + a + bx) \log\left(\frac{b(\sqrt{-c} - \sqrt{dx})}{b\sqrt{-c} + (-1+a)\sqrt{d}}\right) + \log(1 - a - bx) \log\left(\frac{b(\sqrt{-c} + \sqrt{dx})}{b\sqrt{-c} - (-1+a)\sqrt{d}}\right) - \log(1 + a + bx) \log\left(\frac{b(\sqrt{-c} + \sqrt{dx})}{b\sqrt{-c} - (-1+a)\sqrt{d}}\right) + \operatorname{PolyLog}[2, -((\sqrt{d}*(-1 + a + b*x)) / (b*\sqrt{-c} - (-1 + a)*\sqrt{d}))] - \operatorname{PolyLog}[2, (\sqrt{d}*(-1 + a + b*x)) / (b*\sqrt{-c} + (-1 + a)*\sqrt{d})] - \operatorname{PolyLog}[2, -((\sqrt{d}*(1 + a + b*x)) / (b*\sqrt{-c} - (1 + a)*\sqrt{d}))] + \operatorname{PolyLog}[2, (\sqrt{d}*(1 + a + b*x)) / (b*\sqrt{-c} + (1 + a)*\sqrt{d})]}{4*\sqrt{-c}*\sqrt{d}}$$

input `Integrate[ArcTanh[a + b*x]/(c + d*x^2),x]`

output

```
(- (Log[1 - a - b*x]*Log[(b*(Sqrt[-c] - Sqrt[d]*x))/(b*Sqrt[-c] + (-1 + a)*Sqrt[d])]) + Log[1 + a + b*x]*Log[(b*(Sqrt[-c] - Sqrt[d]*x))/(b*Sqrt[-c] + (1 + a)*Sqrt[d])]) + Log[1 - a - b*x]*Log[(b*(Sqrt[-c] + Sqrt[d]*x))/(b*Sqrt[-c] - (-1 + a)*Sqrt[d])] - Log[1 + a + b*x]*Log[(b*(Sqrt[-c] + Sqrt[d]*x))/(b*Sqrt[-c] - (1 + a)*Sqrt[d])] + PolyLog[2, -((Sqrt[d]*(-1 + a + b*x))/(b*Sqrt[-c] - (-1 + a)*Sqrt[d]))] - PolyLog[2, (Sqrt[d]*(-1 + a + b*x))/(b*Sqrt[-c] + (-1 + a)*Sqrt[d])] - PolyLog[2, -((Sqrt[d]*(1 + a + b*x))/(b*Sqrt[-c] - (1 + a)*Sqrt[d]))] + PolyLog[2, (Sqrt[d]*(1 + a + b*x))/(b*Sqrt[-c] + (1 + a)*Sqrt[d])])/(4*Sqrt[-c]*Sqrt[d])
```

Rubi [A] (warning: unable to verify)Time = 0.92 (sec) , antiderivative size = 491, normalized size of antiderivative = 1.71, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6665, 2856, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^2} dx$$

$$\downarrow \text{6665}$$

$$\frac{1}{2} \int \frac{\log(a + bx + 1)}{dx^2 + c} dx - \frac{1}{2} \int \frac{\log(-a - bx + 1)}{dx^2 + c} dx$$

$$\begin{aligned} & \downarrow 2856 \\ & \frac{1}{2} \int \left(\frac{\sqrt{-c} \log(a + bx + 1)}{2c(\sqrt{-c} - \sqrt{dx})} + \frac{\sqrt{-c} \log(a + bx + 1)}{2c(\sqrt{dx} + \sqrt{-c})} \right) dx - \\ & \frac{1}{2} \int \left(\frac{\sqrt{-c} \log(-a - bx + 1)}{2c(\sqrt{-c} - \sqrt{dx})} + \frac{\sqrt{-c} \log(-a - bx + 1)}{2c(\sqrt{dx} + \sqrt{-c})} \right) dx \\ & \downarrow 2009 \end{aligned}$$

$$\begin{aligned} & \frac{1}{2} \left(-\frac{\text{PolyLog}\left(2, -\frac{\sqrt{d}(-a-bx+1)}{b\sqrt{-c}-(1-a)\sqrt{d}}\right)}{2\sqrt{-c}\sqrt{d}} + \frac{\text{PolyLog}\left(2, \frac{\sqrt{d}(-a-bx+1)}{\sqrt{d}(1-a)+b\sqrt{-c}}\right)}{2\sqrt{-c}\sqrt{d}} - \frac{\log(-a-bx+1) \log\left(\frac{b(\sqrt{-c}-\sqrt{dx})}{b\sqrt{-c}-(1-a)\sqrt{d}}\right)}{2\sqrt{-c}\sqrt{d}} + \dots \right) \\ & \frac{1}{2} \left(-\frac{\text{PolyLog}\left(2, -\frac{\sqrt{d}(a+bx+1)}{b\sqrt{-c}-(a+1)\sqrt{d}}\right)}{2\sqrt{-c}\sqrt{d}} + \frac{\text{PolyLog}\left(2, \frac{\sqrt{d}(a+bx+1)}{\sqrt{d}(a+1)+b\sqrt{-c}}\right)}{2\sqrt{-c}\sqrt{d}} + \frac{\log(a+bx+1) \log\left(\frac{b(\sqrt{-c}-\sqrt{dx})}{(a+1)\sqrt{d}+b\sqrt{-c}}\right)}{2\sqrt{-c}\sqrt{d}} - \dots \right) \end{aligned}$$

input `Int[ArcTanh[a + b*x]/(c + d*x^2), x]`

output

```
(-1/2*(Log[1 - a - b*x]*Log[(b*(Sqrt[-c] - Sqrt[d]*x))/(b*Sqrt[-c] - (1 - a)*Sqrt[d])])/(Sqrt[-c]*Sqrt[d]) + (Log[1 - a - b*x]*Log[(b*(Sqrt[-c] + Sqrt[d]*x))/(b*Sqrt[-c] + (1 - a)*Sqrt[d])])/(2*Sqrt[-c]*Sqrt[d]) - PolyLog[2, -((Sqrt[d]*(1 - a - b*x))/(b*Sqrt[-c] - (1 - a)*Sqrt[d]))]/(2*Sqrt[-c]*Sqrt[d]) + PolyLog[2, (Sqrt[d]*(1 - a - b*x))/(b*Sqrt[-c] + (1 - a)*Sqrt[d])]/(2*Sqrt[-c]*Sqrt[d]))/2 + ((Log[1 + a + b*x]*Log[(b*(Sqrt[-c] - Sqrt[d]*x))/(b*Sqrt[-c] + (1 + a)*Sqrt[d])])/(2*Sqrt[-c]*Sqrt[d]) - (Log[1 + a + b*x]*Log[(b*(Sqrt[-c] + Sqrt[d]*x))/(b*Sqrt[-c] - (1 + a)*Sqrt[d])])/(2*Sqrt[-c]*Sqrt[d]) - PolyLog[2, -((Sqrt[d]*(1 + a + b*x))/(b*Sqrt[-c] - (1 + a)*Sqrt[d]))]/(2*Sqrt[-c]*Sqrt[d]) + PolyLog[2, (Sqrt[d]*(1 + a + b*x))/(b*Sqrt[-c] + (1 + a)*Sqrt[d])]/(2*Sqrt[-c]*Sqrt[d]))/2
```

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2856 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]*(b_.))^(p_.)*((f_) + (g_.)*(x_)^(r_))^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log[c*(d + e*x)^n])^p, (f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, r}, x] && I GtQ[p, 0] && IntegerQ[q] && (GtQ[q, 0] || (IntegerQ[r] && NeQ[r, 1]))`

rule 6665 `Int[ArcTanh[(c_) + (d_.)*(x_)]/((e_) + (f_.)*(x_)^(n_.)), x_Symbol] := Simp[1/2 Int[Log[1 + c + d*x]/(e + f*x^n), x], x] - Simp[1/2 Int[Log[1 - c - d*x]/(e + f*x^n), x], x] /; FreeQ[{c, d, e, f}, x] && RationalQ[n]`

Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 444, normalized size of antiderivative = 1.55

| method | result |
|-------------------|---|
| risch | $\frac{\ln(-bx-a+1) \ln\left(\frac{b\sqrt{-cd} - (-bx-a+1)d - ad+d}{b\sqrt{-cd} - ad+d}\right)}{4\sqrt{-cd}} - \frac{\ln(-bx-a+1) \ln\left(\frac{b\sqrt{-cd} + (-bx-a+1)d + ad-d}{b\sqrt{-cd} + ad-d}\right)}{4\sqrt{-cd}} + \frac{\operatorname{dilog}\left(\frac{b\sqrt{-cd} - (-bx-a+1)d - ad+d}{b\sqrt{-cd} - ad+d}\right)}{4\sqrt{-cd}}$ |
| derivativedivides | Expression too large to display |
| default | Expression too large to display |

input `int(arctanh(b*x+a)/(d*x^2+c), x, method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 1/4*\ln(-b*x-a+1)/(-c*d)^{(1/2)}*\ln((b*(-c*d)^{(1/2)}-(-b*x-a+1)*d-a*d+d)/(b*(-c*d)^{(1/2)}-a*d+d))-1/4*\ln(-b*x-a+1)/(-c*d)^{(1/2)}*\ln((b*(-c*d)^{(1/2)}+(-b*x-a+1)*d+a*d-d)/(b*(-c*d)^{(1/2)}+a*d-d))+1/4/(-c*d)^{(1/2)}*dilog((b*(-c*d)^{(1/2)}-(-b*x-a+1)*d-a*d+d)/(b*(-c*d)^{(1/2)}-a*d+d))-1/4/(-c*d)^{(1/2)}*dilog((b*(-c*d)^{(1/2)}+(-b*x-a+1)*d+a*d-d)/(b*(-c*d)^{(1/2)}+a*d-d))+1/4*\ln(b*x+a+1)/(-c*d)^{(1/2)}*\ln((b*(-c*d)^{(1/2)}-(b*x+a+1)*d+a*d+d)/(b*(-c*d)^{(1/2)}+a*d+d))-1/4*\ln(b*x+a+1)/(-c*d)^{(1/2)}*\ln((b*(-c*d)^{(1/2)}+(b*x+a+1)*d-a*d-d)/(b*(-c*d)^{(1/2)}-a*d+d))+1/4/(-c*d)^{(1/2)}*dilog((b*(-c*d)^{(1/2)}-(b*x+a+1)*d+a*d+d)/(b*(-c*d)^{(1/2)}+a*d+d))-1/4/(-c*d)^{(1/2)}*dilog((b*(-c*d)^{(1/2)}+(b*x+a+1)*d-a*d-d)/(b*(-c*d)^{(1/2)}-a*d-d)) \end{aligned}$$

Fricas [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^2} dx = \int \frac{\operatorname{artanh}(bx + a)}{dx^2 + c} dx$$

input `integrate(arctanh(b*x+a)/(d*x^2+c),x, algorithm="fricas")`

output `integral(arctanh(b*x + a)/(d*x^2 + c), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^2} dx = \text{Timed out}$$

input `integrate(atanh(b*x+a)/(d*x**2+c),x)`

output `Timed out`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.24 (sec) , antiderivative size = 591, normalized size of antiderivative = 2.06

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^2} dx = \text{Too large to display}$$

input `integrate(arctanh(b*x+a)/(d*x^2+c),x, algorithm="maxima")`

output

```

arctan(d*x/sqrt(c*d))*arctanh(b*x + a)/sqrt(c*d) + 1/4*((arctan2((b^2*x +
(a + 1)*b)*sqrt(c)*sqrt(d)/(b^2*c + (a^2 + 2*a + 1)*d), ((a + 1)*b*d*x + (
a^2 + 2*a + 1)*d)/(b^2*c + (a^2 + 2*a + 1)*d)) - arctan2((b^2*x + (a - 1)*
b)*sqrt(c)*sqrt(d)/(b^2*c + (a^2 - 2*a + 1)*d), ((a - 1)*b*d*x + (a^2 - 2*
a + 1)*d)/(b^2*c + (a^2 - 2*a + 1)*d))*log(d*x^2 + c) - arctan(sqrt(d)*x/
sqrt(c))*log((b^2*d*x^2 + 2*(a + 1)*b*d*x + (a^2 + 2*a + 1)*d)/(b^2*c + (a
^2 + 2*a + 1)*d)) + arctan(sqrt(d)*x/sqrt(c))*log((b^2*d*x^2 + 2*(a - 1)*b
*d*x + (a^2 - 2*a + 1)*d)/(b^2*c + (a^2 - 2*a + 1)*d)) - I*dilog(((a - 1)*
b*d*x + b^2*c + (I*b^2*x + (-I*a + I)*b)*sqrt(c)*sqrt(d))/(b^2*c + 2*(-I*a
+ I)*b*sqrt(c)*sqrt(d) - (a^2 - 2*a + 1)*d)) + I*dilog(((a - 1)*b*d*x + b
^2*c - (I*b^2*x + (-I*a + I)*b)*sqrt(c)*sqrt(d))/(b^2*c - 2*(-I*a + I)*b*s
qrt(c)*sqrt(d) - (a^2 - 2*a + 1)*d)) + I*dilog(((a + 1)*b*d*x + b^2*c + (I
*b^2*x + (-I*a - I)*b)*sqrt(c)*sqrt(d))/(b^2*c + 2*(-I*a - I)*b*sqrt(c)*sq
rt(d) - (a^2 + 2*a + 1)*d)) - I*dilog(((a + 1)*b*d*x + b^2*c - (I*b^2*x +
(-I*a - I)*b)*sqrt(c)*sqrt(d))/(b^2*c - 2*(-I*a - I)*b*sqrt(c)*sqrt(d) - (
a^2 + 2*a + 1)*d)))/sqrt(c*d)

```

Giac [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^2} dx = \int \frac{\operatorname{artanh}(bx + a)}{dx^2 + c} dx$$

input

```
integrate(arctanh(b*x+a)/(d*x^2+c),x, algorithm="giac")
```

output

```
integrate(arctanh(b*x + a)/(d*x^2 + c), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^2} dx = \int \frac{\operatorname{atanh}(a + bx)}{dx^2 + c} dx$$

input

```
int(atanh(a + b*x)/(c + d*x^2),x)
```

output `int(atanh(a + b*x)/(c + d*x^2), x)`

Reduce [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx^2} dx = \int \frac{\operatorname{atanh}(bx + a)}{dx^2 + c} dx$$

input `int(atanh(b*x+a)/(d*x^2+c), x)`

output `int(atanh(a + b*x)/(c + d*x**2), x)`

3.55 $\int \frac{\operatorname{arctanh}(a+bx)}{c+dx} dx$

| | |
|---|-----|
| Optimal result | 493 |
| Mathematica [A] (verified) | 494 |
| Rubi [A] (verified) | 494 |
| Maple [A] (verified) | 497 |
| Fricas [F] | 497 |
| Sympy [F] | 498 |
| Maxima [A] (verification not implemented) | 498 |
| Giac [F] | 499 |
| Mupad [F(-1)] | 499 |
| Reduce [F] | 499 |

Optimal result

Integrand size = 14, antiderivative size = 120

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx} dx = -\frac{\operatorname{arctanh}(a + bx) \log\left(\frac{2}{1+a+bx}\right)}{d} + \frac{\operatorname{arctanh}(a + bx) \log\left(\frac{2b(c+dx)}{(bc+d-ad)(1+a+bx)}\right)}{d} + \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1+a+bx}\right)}{2d} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2b(c+dx)}{(bc+d-ad)(1+a+bx)}\right)}{2d}$$

output

```
-arctanh(b*x+a)*ln(2/(b*x+a+1))/d+arctanh(b*x+a)*ln(2*b*(d*x+c)/(-a*d+b*c+d)/(b*x+a+1))/d+1/2*polylog(2,1-2/(b*x+a+1))/d-1/2*polylog(2,1-2*b*(d*x+c)/(-a*d+b*c+d)/(b*x+a+1))/d
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.15

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx} dx = -\frac{\log(1 - a - bx) \log\left(-\frac{b(c+dx)}{-bc-(1-a)d}\right)}{2d} + \frac{\log(1 + a + bx) \log\left(\frac{b(c+dx)}{bc-(1+a)d}\right)}{2d} - \frac{\operatorname{PolyLog}\left(2, -\frac{d(1-a-bx)}{-bc-d+ad}\right)}{2d} + \frac{\operatorname{PolyLog}\left(2, \frac{d(1+a+bx)}{-bc+d+ad}\right)}{2d}$$

input

```
Integrate[ArcTanh[a + b*x]/(c + d*x), x]
```

output

```
-1/2*(Log[1 - a - b*x]*Log[-((b*(c + d*x))/(-b*c) - (1 - a)*d))]/d + (Log[1 + a + b*x]*Log[(b*(c + d*x))/(b*c - (1 + a)*d)]/(2*d) - PolyLog[2, -(d*(1 - a - b*x))/(-b*c) - d + a*d)]/(2*d) + PolyLog[2, (d*(1 + a + b*x))/(-b*c) + d + a*d)]/(2*d)
```

Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.15, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {6661, 27, 6472, 2849, 2752, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx} dx$$

↓ 6661

$$\int \frac{b \operatorname{arctanh}(a+bx)}{b\left(c-\frac{ad}{b}\right)+d(a+bx)} d(a + bx)$$

↓ 27

$$\begin{aligned}
& \int \frac{\operatorname{arctanh}(a+bx)}{d(a+bx)-ad+bc} d(a+bx) \\
& \quad \downarrow 6472 \\
& -\frac{\int \frac{\log\left(\frac{2(bc-ad+d(a+bx))}{(bc-ad+d)(a+bx+1)}\right) d(a+bx)}{1-(a+bx)^2}}{d} + \frac{\int \frac{\log\left(\frac{2}{a+bx+1}\right) d(a+bx)}{1-(a+bx)^2}}{d} + \\
& \frac{\operatorname{arctanh}(a+bx) \log\left(\frac{2(d(a+bx)-ad+bc)}{(a+bx+1)(-ad+bc+d)}\right)}{d} - \frac{\operatorname{arctanh}(a+bx) \log\left(\frac{2}{a+bx+1}\right)}{d} \\
& \quad \downarrow 2849 \\
& -\frac{\int \frac{\log\left(\frac{2(bc-ad+d(a+bx))}{(bc-ad+d)(a+bx+1)}\right) d(a+bx)}{1-(a+bx)^2}}{d} + \frac{\int \frac{\log\left(\frac{2}{a+bx+1}\right) d \frac{1}{a+bx+1}}{1-\frac{2}{a+bx+1}}}{d} + \\
& \frac{\operatorname{arctanh}(a+bx) \log\left(\frac{2(d(a+bx)-ad+bc)}{(a+bx+1)(-ad+bc+d)}\right)}{d} - \frac{\operatorname{arctanh}(a+bx) \log\left(\frac{2}{a+bx+1}\right)}{d} \\
& \quad \downarrow 2752 \\
& -\frac{\int \frac{\log\left(\frac{2(bc-ad+d(a+bx))}{(bc-ad+d)(a+bx+1)}\right) d(a+bx)}{1-(a+bx)^2}}{d} + \frac{\operatorname{arctanh}(a+bx) \log\left(\frac{2(d(a+bx)-ad+bc)}{(a+bx+1)(-ad+bc+d)}\right)}{d} - \\
& \frac{\operatorname{arctanh}(a+bx) \log\left(\frac{2}{a+bx+1}\right)}{d} + \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{a+bx+1}\right)}{2d} \\
& \quad \downarrow 2897 \\
& \frac{\operatorname{arctanh}(a+bx) \log\left(\frac{2(d(a+bx)-ad+bc)}{(a+bx+1)(-ad+bc+d)}\right)}{d} - \frac{\operatorname{arctanh}(a+bx) \log\left(\frac{2}{a+bx+1}\right)}{d} - \\
& \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2(bc-ad+d(a+bx))}{(bc-ad+d)(a+bx+1)}\right)}{2d} + \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{a+bx+1}\right)}{2d}
\end{aligned}$$

input `Int[ArcTanh[a + b*x]/(c + d*x), x]`

output `-((ArcTanh[a + b*x]*Log[2/(1 + a + b*x)])/d) + (ArcTanh[a + b*x]*Log[(2*(b*c - a*d + d*(a + b*x)))/((b*c + d - a*d)*(1 + a + b*x))])/d + PolyLog[2, 1 - 2/(1 + a + b*x)]/(2*d) - PolyLog[2, 1 - (2*(b*c - a*d + d*(a + b*x)))/((b*c + d - a*d)*(1 + a + b*x))]/(2*d)`

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 2752 $\text{Int}[\text{Log}[(c_*)(x_)]/((d_) + (e_*)(x_)), x_Symbol] \rightarrow \text{Simp}[(-e^{(-1)})*\text{PolyLog}[2, 1 - c*x], x] /; \text{FreeQ}[\{c, d, e\}, x] \ \&\& \ \text{EqQ}[e + c*d, 0]$
- rule 2849 $\text{Int}[\text{Log}[(c_)/((d_) + (e_*)(x_))]/((f_) + (g_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[-e/g \text{ Subst}[\text{Int}[\text{Log}[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; \text{FreeQ}[\{c, d, e, f, g\}, x] \ \&\& \ \text{EqQ}[c, 2*d] \ \&\& \ \text{EqQ}[e^2*f + d^2*g, 0]$
- rule 2897 $\text{Int}[\text{Log}[u_]*(Pq_)^{(m_.)}, x_Symbol] \rightarrow \text{With}[\{C = \text{FullSimplify}[Pq^m*((1 - u)/D[u, x])]\}, \text{Simp}[C*\text{PolyLog}[2, 1 - u], x] /; \text{FreeQ}[C, x] /; \text{IntegerQ}[m] \ \&\& \ \text{PolyQ}[Pq, x] \ \&\& \ \text{RationalFunctionQ}[u, x] \ \&\& \ \text{LeQ}[\text{RationalFunctionExponents}[u, x][[2]], \text{Expon}[Pq, x]]]$
- rule 6472 $\text{Int}[(a_.) + \text{ArcTanh}[(c_*)(x_)]*(b_.))/((d_) + (e_*)(x_)), x_Symbol] \rightarrow \text{Simp}[(-a + b*\text{ArcTanh}[c*x])*(\text{Log}[2/(1 + c*x)]/e), x] + (\text{Simp}[(a + b*\text{ArcTanh}[c*x])*(\text{Log}[2*c*((d + e*x)/((c*d + e)*(1 + c*x))])]/e), x] + \text{Simp}[b*(c/e) \text{Int}[\text{Log}[2/(1 + c*x)]/(1 - c^2*x^2), x], x] - \text{Simp}[b*(c/e) \text{Int}[\text{Log}[2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/(1 - c^2*x^2), x], x)] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[c^2*d^2 - e^2, 0]$
- rule 6661 $\text{Int}[(a_.) + \text{ArcTanh}[(c_) + (d_*)(x_)]*(b_.))^{(p_.)*((e_.) + (f_*)(x_))^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[1/d \text{ Subst}[\text{Int}[(d*e - c*f)/d + f*(x/d)]^m*(a + b*\text{ArcTanh}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[p, 0]$

Maple [A] (verified)

Time = 0.45 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.51

| method | result |
|-------------------|--|
| risch | $-\frac{\operatorname{dilog}\left(\frac{(-bx-a+1)d+ad-bc-d}{ad-bc-d}\right)}{2d} - \frac{\ln(-bx-a+1)\ln\left(\frac{(-bx-a+1)d+ad-bc-d}{ad-bc-d}\right)}{2d} + \frac{\operatorname{dilog}\left(\frac{(bx+a+1)d-ad+bc-d}{-ad+bc-d}\right)}{2d} + \frac{\ln(bx+a+1)\ln\left(\frac{(bx+a+1)d-ad+bc-d}{-ad+bc-d}\right)}{2d}$ |
| derivativedivides | $\frac{\frac{b \ln(ad-bc-d(bx+a)) \operatorname{arctanh}(bx+a)}{d} + \frac{b \left(\frac{d \left(\operatorname{dilog}\left(\frac{-d(bx+a)-d}{-ad+bc-d}\right) + \ln(ad-bc-d(bx+a)) \ln\left(\frac{-d(bx+a)-d}{-ad+bc-d}\right) \right)}{2} + \frac{d \left(\operatorname{dilog}\left(\frac{-d(bx+a)-d}{-ad+bc-d}\right) \right)}{d} \right)}{d^2}}{b}$ |
| default | $\frac{\frac{b \ln(ad-bc-d(bx+a)) \operatorname{arctanh}(bx+a)}{d} + \frac{b \left(\frac{d \left(\operatorname{dilog}\left(\frac{-d(bx+a)-d}{-ad+bc-d}\right) + \ln(ad-bc-d(bx+a)) \ln\left(\frac{-d(bx+a)-d}{-ad+bc-d}\right) \right)}{2} + \frac{d \left(\operatorname{dilog}\left(\frac{-d(bx+a)-d}{-ad+bc-d}\right) \right)}{d} \right)}{d^2}}{b}$ |
| parts | $\frac{\ln(dx+c) \operatorname{arctanh}(bx+a)}{d} - \frac{b \left(\frac{d \left(\frac{\operatorname{dilog}\left(\frac{ad-bc+b(dx+c)-d}{ad-bc-d}\right)}{b} + \frac{\ln(dx+c)\ln\left(\frac{ad-bc+b(dx+c)-d}{ad-bc-d}\right)}{b} \right)}{2} + \frac{d \left(\frac{\operatorname{dilog}\left(\frac{ad-bc+b(dx+c)-d}{ad-bc-d}\right)}{b} \right)}{d} \right)}{d^2}$ |

input `int(arctanh(b*x+a)/(d*x+c),x,method=_RETURNVERBOSE)`

output `-1/2*dilog(((b*x-a+1)*d+a*d-b*c-d)/(a*d-b*c-d))/d-1/2*ln(-b*x-a+1)*ln(((b*x-a+1)*d+a*d-b*c-d)/(a*d-b*c-d))/d+1/2*dilog(((b*x+a+1)*d-a*d+b*c-d)/(-a*d+b*c-d))/d+1/2*ln(b*x+a+1)*ln(((b*x+a+1)*d-a*d+b*c-d)/(-a*d+b*c-d))/d`

Fricas [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx} dx = \int \frac{\operatorname{arctanh}(bx + a)}{dx + c} dx$$

input `integrate(arctanh(b*x+a)/(d*x+c),x, algorithm="fricas")`

output `integral(arctanh(b*x + a)/(d*x + c), x)`

Sympy [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx} dx = \int \frac{\operatorname{atanh}(a + bx)}{c + dx} dx$$

input `integrate(atanh(b*x+a)/(d*x+c), x)`

output `Integral(atanh(a + b*x)/(c + d*x), x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.60

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx} dx =$$

$$-\frac{1}{2} b \left(\frac{\log(bx + a - 1) \log\left(\frac{bdx + ad - d}{bc - ad + d} + 1\right) + \operatorname{Li}_2\left(-\frac{bdx + ad - d}{bc - ad + d}\right)}{bd} - \frac{\log(bx + a + 1) \log\left(\frac{bdx + ad + d}{bc - ad - d} + 1\right) + \operatorname{Li}_2\left(-\frac{bdx + ad + d}{bc - ad - d}\right)}{bd} \right)$$

$$- \frac{b \left(\frac{\log(bx + a + 1)}{b} - \frac{\log(bx + a - 1)}{b} \right) \log(dx + c)}{2d} + \frac{\operatorname{artanh}(bx + a) \log(dx + c)}{d}$$

input `integrate(arctanh(b*x+a)/(d*x+c), x, algorithm="maxima")`

output `-1/2*b*((log(b*x + a - 1)*log((b*d*x + a*d - d)/(b*c - a*d + d) + 1) + dilog(-(b*d*x + a*d - d)/(b*c - a*d + d)))/(b*d) - (log(b*x + a + 1)*log((b*d*x + a*d + d)/(b*c - a*d - d) + 1) + dilog(-(b*d*x + a*d + d)/(b*c - a*d - d)))/(b*d)) - 1/2*b*(log(b*x + a + 1)/b - log(b*x + a - 1)/b)*log(d*x + c)/d + arctanh(b*x + a)*log(d*x + c)/d`

Giac [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx} dx = \int \frac{\operatorname{artanh}(bx + a)}{dx + c} dx$$

input `integrate(arctanh(b*x+a)/(d*x+c),x, algorithm="giac")`

output `integrate(arctanh(b*x + a)/(d*x + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx} dx = \int \frac{\operatorname{atanh}(a + bx)}{c + dx} dx$$

input `int(atanh(a + b*x)/(c + d*x),x)`

output `int(atanh(a + b*x)/(c + d*x), x)`

Reduce [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + dx} dx = \int \frac{\operatorname{atanh}(bx + a)}{dx + c} dx$$

input `int(atanh(b*x+a)/(d*x+c),x)`

output `int(atanh(a + b*x)/(c + d*x),x)`

3.56 $\int \frac{\operatorname{arctanh}(a+bx)}{c+\frac{d}{x}} dx$

| | |
|---|-----|
| Optimal result | 500 |
| Mathematica [C] (warning: unable to verify) | 501 |
| Rubi [A] (verified) | 501 |
| Maple [A] (verified) | 503 |
| Fricas [F] | 504 |
| Sympy [F] | 504 |
| Maxima [A] (verification not implemented) | 504 |
| Giac [F] | 505 |
| Mupad [F(-1)] | 505 |
| Reduce [F] | 506 |

Optimal result

Integrand size = 16, antiderivative size = 186

$$\int \frac{\operatorname{arctanh}(a+bx)}{c+\frac{d}{x}} dx = \frac{(1-a-bx)\log(1-a-bx)}{2bc} + \frac{(1+a+bx)\log(1+a+bx)}{2bc} - \frac{d \log(1+a+bx) \log\left(-\frac{b(d+cx)}{c+ac-bd}\right)}{2c^2} + \frac{d \log(1-a-bx) \log\left(\frac{b(d+cx)}{c-ac+bd}\right)}{2c^2} + \frac{d \operatorname{PolyLog}\left(2, \frac{c(1-a-bx)}{c-ac+bd}\right)}{2c^2} - \frac{d \operatorname{PolyLog}\left(2, \frac{c(1+a+bx)}{c+ac-bd}\right)}{2c^2}$$

output

```
1/2*(-b*x-a+1)*ln(-b*x-a+1)/b/c+1/2*(b*x+a+1)*ln(b*x+a+1)/b/c-1/2*d*ln(b*x+a+1)*ln(-b*(c*x+d)/(a*c-b*d+c))/c^2+1/2*d*ln(-b*x-a+1)*ln(b*(c*x+d)/(-a*c+b*d+c))/c^2+1/2*d*polylog(2,c*(-b*x-a+1)/(-a*c+b*d+c))/c^2-1/2*d*polylog(2,c*(b*x+a+1)/(a*c-b*d+c))/c^2
```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 8.77 (sec) , antiderivative size = 394, normalized size of antiderivative = 2.12

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x}} dx$$

$$= \frac{2c(a + bx)\operatorname{arctanh}(a + bx) + \frac{bcd\operatorname{arctanh}(a+bx)^2}{ac-bd} - 2c \log\left(\frac{1}{\sqrt{1-(a+bx)^2}}\right) + \frac{bd\left(c\sqrt{1-a^2+\frac{2abd}{c}-\frac{b^2d^2}{c^2}}e^{\operatorname{arctanh}\left(a-\frac{bd}{c}\right)}\right)}{c^2}}$$

input `Integrate[ArcTanh[a + b*x]/(c + d/x),x]`

output

```
(2*c*(a + b*x)*ArcTanh[a + b*x] + (b*c*d*ArcTanh[a + b*x]^2)/(a*c - b*d) -
  2*c*Log[1/Sqrt[1 - (a + b*x)^2]] + (b*d*(c*Sqrt[1 - a^2 + (2*a*b*d)/c - (
  b^2*d^2)/c^2]*E^ArcTanh[a - (b*d)/c]*ArcTanh[a + b*x]^2 + (a*c - b*d)*ArcT
  anh[a + b*x]*(I*Pi - 2*ArcTanh[a - (b*d)/c] + 2*Log[1 - E^(2*(ArcTanh[a -
  (b*d)/c] - ArcTanh[a + b*x]))]) - (a*c - b*d)*(I*Pi*(Log[1 + E^(2*ArcTanh[
  a + b*x]]) - Log[1/Sqrt[1 - (a + b*x)^2]]) + 2*ArcTanh[a - (b*d)/c]*(Log[1
  - E^(2*(ArcTanh[a - (b*d)/c] - ArcTanh[a + b*x]))] - Log[(-I)*Sinh[ArcTan
  h[a - (b*d)/c] - ArcTanh[a + b*x]])) + (-a*c) + b*d)*PolyLog[2, E^(2*(Ar
  cTanh[a - (b*d)/c] - ArcTanh[a + b*x]))])/(-a*c) + b*d*(ArcTanh[a
  + b*x]*(ArcTanh[a + b*x] + 2*Log[1 + E^(-2*ArcTanh[a + b*x]))] - PolyLog[
  2, -E^(-2*ArcTanh[a + b*x]))]))/(2*b*c^2)
```

Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.03, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6665, 2856, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x}} dx$$

$$\begin{aligned}
& \downarrow \text{6665} \\
& \frac{1}{2} \int \frac{\log(a + bx + 1)}{c + \frac{d}{x}} dx - \frac{1}{2} \int \frac{\log(-a - bx + 1)}{c + \frac{d}{x}} dx \\
& \downarrow \text{2856} \\
& \frac{1}{2} \int \left(\frac{\log(a + bx + 1)}{c} - \frac{d \log(a + bx + 1)}{c(d + cx)} \right) dx - \\
& \frac{1}{2} \int \left(\frac{\log(-a - bx + 1)}{c} - \frac{d \log(-a - bx + 1)}{c(d + cx)} \right) dx \\
& \downarrow \text{2009} \\
& \frac{1}{2} \left(\frac{d \operatorname{PolyLog} \left(2, \frac{c(-a-bx+1)}{-ac+c+bd} \right)}{c^2} + \frac{d \log(-a - bx + 1) \log \left(\frac{b(cx+d)}{-ac+bd+c} \right)}{c^2} + \frac{(-a - bx + 1) \log(-a - bx + 1)}{bc} + \frac{x}{c} \right) + \\
& \frac{1}{2} \left(-\frac{d \operatorname{PolyLog} \left(2, \frac{c(a+bx+1)}{ac+c-bd} \right)}{c^2} - \frac{d \log(a + bx + 1) \log \left(-\frac{b(cx+d)}{ac-bd+c} \right)}{c^2} + \frac{(a + bx + 1) \log(a + bx + 1)}{bc} - \frac{x}{c} \right)
\end{aligned}$$

input `Int[ArcTanh[a + b*x]/(c + d/x),x]`

output `(x/c + ((1 - a - b*x)*Log[1 - a - b*x])/(b*c) + (d*Log[1 - a - b*x]*Log[(b*(d + c*x))/(c - a*c + b*d]))/c^2 + (d*PolyLog[2, (c*(1 - a - b*x))/(c - a*c + b*d)]/c^2)/2 + (-x/c) + ((1 + a + b*x)*Log[1 + a + b*x])/(b*c) - (d*Log[1 + a + b*x]*Log[-((b*(d + c*x))/(c + a*c - b*d))])/c^2 - (d*PolyLog[2, (c*(1 + a + b*x))/(c + a*c - b*d)]/c^2)/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2856 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_))^(n_.)]*(b_.))^(p_.)*((f_) + (g_.)*(x_)^(r_))^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log[c*(d + e*x)^n])^p, (f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, r}, x] && ! GtQ[p, 0] && IntegerQ[q] && (GtQ[q, 0] || (IntegerQ[r] && NeQ[r, 1]))`

rule 6665

```
Int[ArcTanh[(c_) + (d_.)*(x_)]/((e_) + (f_.)*(x_)^(n_.)), x_Symbol] := Simp
[1/2 Int[Log[1 + c + d*x]/(e + f*x^n), x], x] - Simp[1/2 Int[Log[1 - c
- d*x]/(e + f*x^n), x], x] /; FreeQ[{c, d, e, f}, x] && RationalQ[n]
```

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 259, normalized size of antiderivative = 1.39

| method | result |
|-------------------|---|
| parts | $\frac{\operatorname{arctanh}(bx+a)x}{c} - \frac{\operatorname{arctanh}(bx+a)d \ln(cx+d)}{c^2} - \frac{b \left(\frac{(a-1) \ln(ac-bd+b(cx+d)-c)}{2b^2} + \frac{(-1-a) \ln(ac-bd+b(cx+d)+c)}{2b^2} \right) + d \left(\frac{\operatorname{arctanh}(bx+a)(bx+a)}{c} - \frac{\operatorname{arctanh}(bx+a)db \ln(ac-bd-c(bx+a))}{c^2} \right)}{c^2}$ |
| risch | $\frac{\ln(bx+a+1)x}{2c} + \frac{\ln(bx+a+1)a}{2bc} + \frac{\ln(bx+a+1)}{2bc} - \frac{1}{bc} - \frac{d \operatorname{dilog}\left(\frac{(bx+a+1)c-ac+bd-c}{-ac+bd-c}\right)}{2c^2} - \frac{d \ln(bx+a+1) \ln\left(\frac{bx+a+1}{2c^2}\right)}{2c^2}$ |
| derivativedivides | $\frac{\operatorname{arctanh}(bx+a)(bx+a)}{c} - \frac{\operatorname{arctanh}(bx+a)db \ln(ac-bd-c(bx+a))}{c^2} + \frac{bd \left(-\frac{\operatorname{dilog}\left(\frac{-c(bx+a)+c}{-ac+bd+c}\right) + \ln(ac-bd-c(bx+a)) \ln\left(\frac{-c(bx+a)+c}{-ac+bd+c}\right)}{2c} \right)}{c^2}$ |
| default | $\frac{\operatorname{arctanh}(bx+a)(bx+a)}{c} - \frac{\operatorname{arctanh}(bx+a)db \ln(ac-bd-c(bx+a))}{c^2} + \frac{bd \left(-\frac{\operatorname{dilog}\left(\frac{-c(bx+a)+c}{-ac+bd+c}\right) + \ln(ac-bd-c(bx+a)) \ln\left(\frac{-c(bx+a)+c}{-ac+bd+c}\right)}{2c} \right)}{c^2}$ |

input

```
int(arctanh(b*x+a)/(c+d/x), x, method=_RETURNVERBOSE)
```

output

```
arctanh(b*x+a)*x/c-arctanh(b*x+a)/c^2*d*ln(c*x+d)-b/c*(1/2*(a-1)/b^2*ln(a*
c-b*d+b*(c*x+d)-c)+1/2*(-1-a)/b^2*ln(a*c-b*d+b*(c*x+d)+c)+d*(1/2/c*(dilog(
(a*c-b*d+b*(c*x+d)-c)/(a*c-b*d-c))/b+ln(c*x+d)*ln((a*c-b*d+b*(c*x+d)-c)/(a
*c-b*d-c))/b)-1/2/c*(dilog((a*c-b*d+b*(c*x+d)+c)/(a*c-b*d+c))/b+ln(c*x+d)*
ln((a*c-b*d+b*(c*x+d)+c)/(a*c-b*d+c))/b))
```


Fricas [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x}} dx = \int \frac{\operatorname{artanh}(bx + a)}{c + \frac{d}{x}} dx$$

input `integrate(arctanh(b*x+a)/(c+d/x),x, algorithm="fricas")`

output `integral(x*arctanh(b*x + a)/(c*x + d), x)`

Sympy [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x}} dx = \int \frac{x \operatorname{atanh}(a + bx)}{cx + d} dx$$

input `integrate(atanh(b*x+a)/(c+d/x),x)`

output `Integral(x*atanh(a + b*x)/(c*x + d), x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.03

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x}} dx \\ &= \frac{1}{2} b \left(\frac{(\log(cx + d) \log\left(\frac{bcx+bd}{ac-bd+c} + 1\right) + \operatorname{Li}_2\left(-\frac{bcx+bd}{ac-bd+c}\right))d}{bc^2} - \frac{(\log(cx + d) \log\left(\frac{bcx+bd}{ac-bd-c} + 1\right) + \operatorname{Li}_2\left(-\frac{bcx+bd}{ac-bd-c}\right))d}{bc^2} \right. \\ & \quad \left. + \left(\frac{x}{c} - \frac{d \log(cx + d)}{c^2}\right) \operatorname{artanh}(bx + a) \right) \end{aligned}$$

input `integrate(arctanh(b*x+a)/(c+d/x),x, algorithm="maxima")`

output

```
1/2*b*((log(c*x + d)*log((b*c*x + b*d)/(a*c - b*d + c) + 1) + dilog(-(b*c*x + b*d)/(a*c - b*d + c)))*d/(b*c^2) - (log(c*x + d)*log((b*c*x + b*d)/(a*c - b*d - c) + 1) + dilog(-(b*c*x + b*d)/(a*c - b*d - c)))*d/(b*c^2) + (a + 1)*log(b*x + a + 1)/(b^2*c) - (a - 1)*log(b*x + a - 1)/(b^2*c)) + (x/c - d*log(c*x + d)/c^2)*arctanh(b*x + a)
```

Giac [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x}} dx = \int \frac{\operatorname{artanh}(bx + a)}{c + \frac{d}{x}} dx$$

input

```
integrate(arctanh(b*x+a)/(c+d/x),x, algorithm="giac")
```

output

```
integrate(arctanh(b*x + a)/(c + d/x), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x}} dx = \int \frac{\operatorname{atanh}(a + bx)}{c + \frac{d}{x}} dx$$

input

```
int(atanh(a + b*x)/(c + d/x),x)
```

output

```
int(atanh(a + b*x)/(c + d/x), x)
```

Reduce [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x}} dx = \int \frac{\operatorname{atanh}(bx + a) x}{cx + d} dx$$

input `int(atanh(b*x+a)/(c+d/x),x)`

output `int((atanh(a + b*x)*x)/(c*x + d),x)`

$$3.57 \quad \int \frac{\operatorname{arctanh}(a+bx)}{c+\frac{d}{x^2}} dx$$

| | |
|---|-----|
| Optimal result | 508 |
| Mathematica [A] (verified) | 509 |
| Rubi [A] (verified) | 510 |
| Maple [A] (verified) | 511 |
| Fricas [F] | 512 |
| Sympy [F(-1)] | 512 |
| Maxima [C] (verification not implemented) | 513 |
| Giac [F] | 513 |
| Mupad [F(-1)] | 514 |
| Reduce [F] | 514 |

Optimal result

Integrand size = 16, antiderivative size = 545

$$\begin{aligned}
\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^2}} dx &= \frac{(1 - a - bx) \log(1 - a - bx)}{2bc} + \frac{(1 + a + bx) \log(1 + a + bx)}{2bc} \\
&+ \frac{\sqrt{d} \log(1 - a - bx) \log\left(-\frac{b(\sqrt{d} - \sqrt{-cx})}{(1-a)\sqrt{-c} - b\sqrt{d}}\right)}{4(-c)^{3/2}} \\
&- \frac{\sqrt{d} \log(1 + a + bx) \log\left(\frac{b(\sqrt{d} - \sqrt{-cx})}{(1+a)\sqrt{-c} + b\sqrt{d}}\right)}{4(-c)^{3/2}} \\
&+ \frac{\sqrt{d} \log(1 + a + bx) \log\left(-\frac{b(\sqrt{d} + \sqrt{-cx})}{(1+a)\sqrt{-c} - b\sqrt{d}}\right)}{4(-c)^{3/2}} \\
&- \frac{\sqrt{d} \log(1 - a - bx) \log\left(\frac{b(\sqrt{d} + \sqrt{-cx})}{(1-a)\sqrt{-c} + b\sqrt{d}}\right)}{4(-c)^{3/2}} \\
&+ \frac{\sqrt{d} \operatorname{PolyLog}\left(2, \frac{\sqrt{-c}(1-a-bx)}{\sqrt{-c} - a\sqrt{-c} - b\sqrt{d}}\right)}{4(-c)^{3/2}} \\
&- \frac{\sqrt{d} \operatorname{PolyLog}\left(2, \frac{\sqrt{-c}(1-a-bx)}{(1-a)\sqrt{-c} + b\sqrt{d}}\right)}{4(-c)^{3/2}} \\
&+ \frac{\sqrt{d} \operatorname{PolyLog}\left(2, \frac{\sqrt{-c}(1+a+bx)}{(1+a)\sqrt{-c} - b\sqrt{d}}\right)}{4(-c)^{3/2}} \\
&- \frac{\sqrt{d} \operatorname{PolyLog}\left(2, \frac{\sqrt{-c}(1+a+bx)}{(1+a)\sqrt{-c} + b\sqrt{d}}\right)}{4(-c)^{3/2}}
\end{aligned}$$

output

```

1/2*(-b*x-a+1)*ln(-b*x-a+1)/b/c+1/2*(b*x+a+1)*ln(b*x+a+1)/b/c+1/4*d^(1/2)*
ln(-b*x-a+1)*ln(-b*(d^(1/2)-(-c)^(1/2)*x)/((1-a)*(-c)^(1/2)-b*d^(1/2)))/(-
c)^(3/2)-1/4*d^(1/2)*ln(b*x+a+1)*ln(b*(d^(1/2)-(-c)^(1/2)*x)/((1+a)*(-c)^(
1/2)+b*d^(1/2)))/(-c)^(3/2)+1/4*d^(1/2)*ln(b*x+a+1)*ln(-b*(d^(1/2)+(-c)^(1
/2)*x)/((1+a)*(-c)^(1/2)-b*d^(1/2)))/(-c)^(3/2)-1/4*d^(1/2)*ln(-b*x-a+1)*l
n(b*(d^(1/2)+(-c)^(1/2)*x)/((1-a)*(-c)^(1/2)+b*d^(1/2)))/(-c)^(3/2)+1/4*d^
(1/2)*polylog(2, (-c)^(1/2)*(-b*x-a+1)/((-c)^(1/2)-a*(-c)^(1/2)-b*d^(1/2)))
/(-c)^(3/2)-1/4*d^(1/2)*polylog(2, (-c)^(1/2)*(-b*x-a+1)/((1-a)*(-c)^(1/2)+
b*d^(1/2)))/(-c)^(3/2)+1/4*d^(1/2)*polylog(2, (-c)^(1/2)*(b*x+a+1)/((1+a)*(-
c)^(1/2)-b*d^(1/2)))/(-c)^(3/2)-1/4*d^(1/2)*polylog(2, (-c)^(1/2)*(b*x+a+1
)/((1+a)*(-c)^(1/2)+b*d^(1/2)))/(-c)^(3/2)

```

Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 555, normalized size of antiderivative = 1.02

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^2}} dx =$$

$$\frac{2\sqrt{-c} \log(1 - a - bx) - 2a\sqrt{-c} \log(1 - a - bx) - 2b\sqrt{-c} \log(1 - a - bx) + 2\sqrt{-c} \log(1 + a + bx)}{c + \frac{d}{x^2}}$$

input

```
Integrate[ArcTanh[a + b*x]/(c + d/x^2), x]
```

output

```

-1/4*(2*Sqrt[-c]*Log[1 - a - b*x] - 2*a*Sqrt[-c]*Log[1 - a - b*x] - 2*b*Sq
rt[-c]*x*Log[1 - a - b*x] + 2*Sqrt[-c]*Log[1 + a + b*x] + 2*a*Sqrt[-c]*Log
[1 + a + b*x] + 2*b*Sqrt[-c]*x*Log[1 + a + b*x] - b*Sqrt[d]*Log[1 - a - b*
x]*Log[(b*(Sqrt[d] - Sqrt[-c]*x))/((-1 + a)*Sqrt[-c] + b*Sqrt[d])] + b*Sqr
t[d]*Log[1 + a + b*x]*Log[(b*(Sqrt[d] - Sqrt[-c]*x))/((1 + a)*Sqrt[-c] + b
*Sqrt[d])] - b*Sqrt[d]*Log[1 + a + b*x]*Log[-((b*(Sqrt[d] + Sqrt[-c]*x))/
(1 + a)*Sqrt[-c] - b*Sqrt[d])] + b*Sqrt[d]*Log[1 - a - b*x]*Log[(b*(Sqrt[
d] + Sqrt[-c]*x))/(-((-1 + a)*Sqrt[-c]) + b*Sqrt[d])] + b*Sqrt[d]*PolyLog[
2, (Sqrt[-c]*(-1 + a + b*x))/(-Sqrt[-c] + a*Sqrt[-c] - b*Sqrt[d])] - b*Sqr
t[d]*PolyLog[2, (Sqrt[-c]*(-1 + a + b*x))/(-Sqrt[-c] + a*Sqrt[-c] + b*Sqr
t[d])] - b*Sqrt[d]*PolyLog[2, (Sqrt[-c]*(1 + a + b*x))/(Sqrt[-c] + a*Sqrt[-
c] - b*Sqrt[d])] + b*Sqrt[d]*PolyLog[2, (Sqrt[-c]*(1 + a + b*x))/(Sqrt[-c]
+ a*Sqrt[-c] + b*Sqrt[d])])]/(b*(-c)^(3/2))

```

Rubi [A] (verified)

Time = 1.16 (sec) , antiderivative size = 560, normalized size of antiderivative = 1.03, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6665, 2856, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^2}} dx$$

↓ 6665

$$\frac{1}{2} \int \frac{\log(a + bx + 1)}{c + \frac{d}{x^2}} dx - \frac{1}{2} \int \frac{\log(-a - bx + 1)}{c + \frac{d}{x^2}} dx$$

↓ 2856

$$\frac{1}{2} \int \left(\frac{\log(a + bx + 1)}{c} - \frac{d \log(a + bx + 1)}{c(cx^2 + d)} \right) dx - \frac{1}{2} \int \left(\frac{\log(-a - bx + 1)}{c} - \frac{d \log(-a - bx + 1)}{c(cx^2 + d)} \right) dx$$

↓ 2009

$$\frac{1}{2} \left(\frac{\sqrt{d} \operatorname{PolyLog} \left(2, \frac{\sqrt{-c}(-a-bx+1)}{-\sqrt{-c}a + \sqrt{-c}b\sqrt{d}} \right)}{2(-c)^{3/2}} - \frac{\sqrt{d} \operatorname{PolyLog} \left(2, \frac{\sqrt{-c}(-a-bx+1)}{\sqrt{-c}(1-a) + b\sqrt{d}} \right)}{2(-c)^{3/2}} + \frac{\sqrt{d} \log(-a - bx + 1) \log \left(-\frac{b(\sqrt{d} - \sqrt{-c})}{(1-a)\sqrt{-c}} \right)}{2(-c)^{3/2}} \right)$$

$$\frac{1}{2} \left(\frac{\sqrt{d} \operatorname{PolyLog} \left(2, \frac{\sqrt{-c}(a+bx+1)}{(a+1)\sqrt{-c} - b\sqrt{d}} \right)}{2(-c)^{3/2}} - \frac{\sqrt{d} \operatorname{PolyLog} \left(2, \frac{\sqrt{-c}(a+bx+1)}{\sqrt{-c}(a+1) + b\sqrt{d}} \right)}{2(-c)^{3/2}} - \frac{\sqrt{d} \log(a + bx + 1) \log \left(\frac{b(\sqrt{d} - \sqrt{-c})}{(a+1)\sqrt{-c} + b\sqrt{d}} \right)}{2(-c)^{3/2}} \right)$$

input

```
Int[ArcTanh[a + b*x]/(c + d/x^2), x]
```

output

$$\begin{aligned} & (x/c + ((1 - a - b*x)*\text{Log}[1 - a - b*x])/(b*c) + (\text{Sqrt}[d]*\text{Log}[1 - a - b*x]* \\ & \text{Log}[-(b*(\text{Sqrt}[d] - \text{Sqrt}[-c]*x))/((1 - a)*\text{Sqrt}[-c] - b*\text{Sqrt}[d])])/(2*(-c) \\ & ^{(3/2)}) - (\text{Sqrt}[d]*\text{Log}[1 - a - b*x]*\text{Log}[(b*(\text{Sqrt}[d] + \text{Sqrt}[-c]*x))/((1 - a) \\ &)*\text{Sqrt}[-c] + b*\text{Sqrt}[d])])/(2*(-c)^{(3/2)}) + (\text{Sqrt}[d]*\text{PolyLog}[2, (\text{Sqrt}[-c]*(\\ & 1 - a - b*x))/(\text{Sqrt}[-c] - a*\text{Sqrt}[-c] - b*\text{Sqrt}[d])])/(2*(-c)^{(3/2)}) - (\text{Sqrt} \\ & [d]*\text{PolyLog}[2, (\text{Sqrt}[-c]*(1 - a - b*x))/((1 - a)*\text{Sqrt}[-c] + b*\text{Sqrt}[d])])/(\\ & 2*(-c)^{(3/2)}))/2 + (-(x/c) + ((1 + a + b*x)*\text{Log}[1 + a + b*x])/(b*c) - (\text{Sqr} \\ & \text{t}[d]*\text{Log}[1 + a + b*x]*\text{Log}[(b*(\text{Sqrt}[d] - \text{Sqrt}[-c]*x))/((1 + a)*\text{Sqrt}[-c] + b \\ & * \text{Sqrt}[d])])/(2*(-c)^{(3/2)}) + (\text{Sqrt}[d]*\text{Log}[1 + a + b*x]*\text{Log}[-(b*(\text{Sqrt}[d] + \\ & \text{Sqrt}[-c]*x))/((1 + a)*\text{Sqrt}[-c] - b*\text{Sqrt}[d])])/(2*(-c)^{(3/2)}) + (\text{Sqrt}[d]* \\ & \text{PolyLog}[2, (\text{Sqrt}[-c]*(1 + a + b*x))/((1 + a)*\text{Sqrt}[-c] - b*\text{Sqrt}[d])])/(2*(- \\ & c)^{(3/2)}) - (\text{Sqrt}[d]*\text{PolyLog}[2, (\text{Sqrt}[-c]*(1 + a + b*x))/((1 + a)*\text{Sqrt}[-c] \\ & + b*\text{Sqrt}[d])])/(2*(-c)^{(3/2)}))/2 \end{aligned}$$

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2856

```
Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]*(b_.))^(p_.)*((f_) + (g_.)
)*(x_)^(r_)^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log[c*(d + e*x)
^ n])^p, (f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, r}, x] && I
GtQ[p, 0] && IntegerQ[q] && (GtQ[q, 0] || (IntegerQ[r] && NeQ[r, 1]))
```

rule 6665

```
Int[ArcTanh[(c_) + (d_.)*(x_)]/((e_) + (f_.)*(x_)^(n_.)), x_Symbol] := Simp
[1/2 Int[Log[1 + c + d*x]/(e + f*x^n), x], x] - Simp[1/2 Int[Log[1 - c
- d*x]/(e + f*x^n), x], x] /; FreeQ[{c, d, e, f}, x] && RationalQ[n]
```

Maple [A] (verified)

Time = 1.06 (sec) , antiderivative size = 581, normalized size of antiderivative = 1.07

| method | result |
|------------------|---|
| risch | $\frac{\ln(bx+a+1)x}{2c} + \frac{\ln(bx+a+1)a}{2bc} + \frac{\ln(bx+a+1)}{2bc} - \frac{1}{bc} - \frac{d \ln(bx+a+1) \ln\left(\frac{b\sqrt{-cd} - (bx+a+1)c + ac+c}{b\sqrt{-cd} + ac+c}\right)}{4c\sqrt{-cd}} + \frac{d \ln(bx+a+1)}{4c\sqrt{-cd}}$ |
| derivativdivides | Expression too large to display |
| default | Expression too large to display |

input `int(arctanh(b*x+a)/(c+d/x^2),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 1/2/c*\ln(b*x+a+1)*x+1/2/b/c*\ln(b*x+a+1)*a+1/2/b/c*\ln(b*x+a+1)-1/b/c-1/4*d/ \\ & c*\ln(b*x+a+1)/(-c*d)^{(1/2)}*\ln((b*(-c*d)^{(1/2)}-(b*x+a+1)*c+a*c+c)/(b*(-c*d) \\ & ^{(1/2)}+a*c+c))+1/4*d/c*\ln(b*x+a+1)/(-c*d)^{(1/2)}*\ln((b*(-c*d)^{(1/2)}+(b*x+a+ \\ & 1)*c-a*c-c)/(b*(-c*d)^{(1/2)}-a*c-c))-1/4*d/c/(-c*d)^{(1/2)}*dilog((b*(-c*d)^{(\\ & 1/2)}-(b*x+a+1)*c+a*c+c)/(b*(-c*d)^{(1/2)}+a*c+c))+1/4*d/c/(-c*d)^{(1/2)}*dilog \\ & ((b*(-c*d)^{(1/2)}+(b*x+a+1)*c-a*c-c)/(b*(-c*d)^{(1/2)}-a*c-c))-1/2/c*\ln(-b*x- \\ & a+1)*x-1/2/b/c*\ln(-b*x-a+1)*a+1/2/b/c*\ln(-b*x-a+1)-1/4*d/c*\ln(-b*x-a+1)/(- \\ & c*d)^{(1/2)}*\ln((b*(-c*d)^{(1/2)}-(-b*x-a+1)*c-a*c+c)/(b*(-c*d)^{(1/2)}-a*c+c))+ \\ & 1/4*d/c*\ln(-b*x-a+1)/(-c*d)^{(1/2)}*\ln((b*(-c*d)^{(1/2)}+(-b*x-a+1)*c+a*c-c)/(\\ & b*(-c*d)^{(1/2)}+a*c-c))-1/4*d/c/(-c*d)^{(1/2)}*dilog((b*(-c*d)^{(1/2)}-(-b*x-a+ \\ & 1)*c-a*c+c)/(b*(-c*d)^{(1/2)}-a*c+c))+1/4*d/c/(-c*d)^{(1/2)}*dilog((b*(-c*d)^{(\\ & 1/2)}+(-b*x-a+1)*c+a*c-c)/(b*(-c*d)^{(1/2)}+a*c-c)) \end{aligned}$$

Fricas [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^2}} dx = \int \frac{\operatorname{artanh}(bx + a)}{c + \frac{d}{x^2}} dx$$

input `integrate(arctanh(b*x+a)/(c+d/x^2),x, algorithm="fricas")`

output `integral(x^2*arctanh(b*x + a)/(c*x^2 + d), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^2}} dx = \text{Timed out}$$

input `integrate(atanh(b*x+a)/(c+d/x**2),x)`

output `Timed out`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.21 (sec) , antiderivative size = 651, normalized size of antiderivative = 1.19

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^2}} dx = \text{Too large to display}$$

input `integrate(arctanh(b*x+a)/(c+d/x^2),x, algorithm="maxima")`

output

```

-(d*arctan(c*x/sqrt(c*d))/(sqrt(c*d)*c) - x/c)*arctanh(b*x + a) + 1/4*(2*(
a + 1)*c*log(b*x + a + 1) - 2*(a - 1)*c*log(b*x + a - 1) + (b*arctan(sqrt(
c)*x/sqrt(d))*log((b^2*c*x^2 + 2*(a + 1)*b*c*x + (a^2 + 2*a + 1)*c)/(b^2*d
+ (a^2 + 2*a + 1)*c)) - b*arctan(sqrt(c)*x/sqrt(d))*log((b^2*c*x^2 + 2*(a
- 1)*b*c*x + (a^2 - 2*a + 1)*c)/(b^2*d + (a^2 - 2*a + 1)*c)) + I*b*dilog(
((a - 1)*b*c*x + b^2*d + (I*b^2*x + (-I*a + I)*b)*sqrt(c)*sqrt(d))/(2*(-I*
a + I)*b*sqrt(c)*sqrt(d) + b^2*d - (a^2 - 2*a + 1)*c)) - I*b*dilog(-((a -
1)*b*c*x + b^2*d - (I*b^2*x + (-I*a + I)*b)*sqrt(c)*sqrt(d))/(2*(-I*a + I)
*b*sqrt(c)*sqrt(d) - b^2*d + (a^2 - 2*a + 1)*c)) - I*b*dilog(((a + 1)*b*c*
x + b^2*d + (I*b^2*x + (-I*a - I)*b)*sqrt(c)*sqrt(d))/(2*(-I*a - I)*b*sqrt
(c)*sqrt(d) + b^2*d - (a^2 + 2*a + 1)*c)) + I*b*dilog(-((a + 1)*b*c*x + b^
2*d - (I*b^2*x + (-I*a - I)*b)*sqrt(c)*sqrt(d))/(2*(-I*a - I)*b*sqrt(c)*sq
rt(d) - b^2*d + (a^2 + 2*a + 1)*c)) - (b*arctan2((b^2*x + (a + 1)*b)*sqrt(
c)*sqrt(d)/(b^2*d + (a^2 + 2*a + 1)*c), ((a + 1)*b*c*x + (a^2 + 2*a + 1)*c
)/(b^2*d + (a^2 + 2*a + 1)*c)) - b*arctan2((b^2*x + (a - 1)*b)*sqrt(c)*sq
rt(d)/(b^2*d + (a^2 - 2*a + 1)*c), ((a - 1)*b*c*x + (a^2 - 2*a + 1)*c)/(b^2
*d + (a^2 - 2*a + 1)*c))) * log(c*x^2 + d) * sqrt(c) * sqrt(d) / (b*c^2)

```

Giac [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^2}} dx = \int \frac{\operatorname{artanh}(bx + a)}{c + \frac{d}{x^2}} dx$$

input `integrate(arctanh(b*x+a)/(c+d/x^2),x, algorithm="giac")`

output `integrate(arctanh(b*x + a)/(c + d/x^2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^2}} dx = \int \frac{\operatorname{atanh}(a + bx)}{c + \frac{d}{x^2}} dx$$

input `int(atanh(a + b*x)/(c + d/x^2), x)`output `int(atanh(a + b*x)/(c + d/x^2), x)`**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^2}} dx = \int \frac{\operatorname{atanh}(bx + a) x^2}{c x^2 + d} dx$$

input `int(atanh(b*x+a)/(c+d/x^2), x)`output `int((atanh(a + b*x)*x**2)/(c*x**2 + d), x)`

$$3.58 \quad \int \frac{\operatorname{arctanh}(a+bx)}{c+\frac{d}{x^3}} dx$$

| | |
|---------------------------------------|-----|
| Optimal result | 515 |
| Mathematica [A] (verified) | 516 |
| Rubi [A] (verified) | 517 |
| Maple [C] (warning: unable to verify) | 520 |
| Fricas [F] | 521 |
| Sympy [F(-1)] | 521 |
| Maxima [F] | 522 |
| Giac [F] | 522 |
| Mupad [F(-1)] | 522 |
| Reduce [F] | 523 |

Optimal result

Integrand size = 16, antiderivative size = 832

$$\int \frac{\operatorname{arctanh}(a+bx)}{c+\frac{d}{x^3}} dx = \text{Too large to display}$$

output

```

1/2*(-b*x-a+1)*ln(-b*x-a+1)/b/c+1/2*(b*x+a+1)*ln(b*x+a+1)/b/c-1/6*d^(1/3)*
ln(b*x+a+1)*ln(-b*(d^(1/3)+c^(1/3)*x)/((1+a)*c^(1/3)-b*d^(1/3)))/c^(4/3)+1
/6*d^(1/3)*ln(-b*x-a+1)*ln(b*(d^(1/3)+c^(1/3)*x)/((1-a)*c^(1/3)+b*d^(1/3))
)/c^(4/3)+1/6*(-1)^(2/3)*d^(1/3)*ln(-b*x-a+1)*ln(-b*(d^(1/3)-(-1)^(1/3)*c^
(1/3)*x)/((-1)^(1/3)*(1-a)*c^(1/3)-b*d^(1/3)))/c^(4/3)-1/6*(-1)^(2/3)*d^(1
/3)*ln(b*x+a+1)*ln(b*(d^(1/3)-(-1)^(1/3)*c^(1/3)*x)/((-1)^(1/3)*(1+a)*c^(1
/3)+b*d^(1/3)))/c^(4/3)+1/6*(-1)^(1/3)*d^(1/3)*ln(b*x+a+1)*ln(-b*(d^(1/3)+
(-1)^(2/3)*c^(1/3)*x)/((-1)^(2/3)*(1+a)*c^(1/3)-b*d^(1/3)))/c^(4/3)-1/6*(-
1)^(1/3)*d^(1/3)*ln(-b*x-a+1)*ln(b*(d^(1/3)+(-1)^(2/3)*c^(1/3)*x)/((-1)^(2
/3)*(1-a)*c^(1/3)+b*d^(1/3)))/c^(4/3)+1/6*(-1)^(2/3)*d^(1/3)*polylog(2,(-1
)^(1/3)*c^(1/3)*(-b*x-a+1)/((-1)^(1/3)*(1-a)*c^(1/3)-b*d^(1/3)))/c^(4/3)+1
/6*d^(1/3)*polylog(2,c^(1/3)*(-b*x-a+1)/((1-a)*c^(1/3)+b*d^(1/3)))/c^(4/3)
-1/6*(-1)^(1/3)*d^(1/3)*polylog(2,(-1)^(2/3)*c^(1/3)*(-b*x-a+1)/((-1)^(2/3
)*c^(1/3)+b*d^(1/3)))/c^(4/3)-1/6*d^(1/3)*polylog(2,c^(1/3)*(b*x+a+1)
)/((1+a)*c^(1/3)-b*d^(1/3)))/c^(4/3)+1/6*(-1)^(1/3)*d^(1/3)*polylog(2,(-1)
^(2/3)*c^(1/3)*(b*x+a+1)/((-1)^(2/3)*(1+a)*c^(1/3)-b*d^(1/3)))/c^(4/3)-1/6
*(-1)^(2/3)*d^(1/3)*polylog(2,(-1)^(1/3)*c^(1/3)*(b*x+a+1)/((-1)^(1/3)*(1+
a)*c^(1/3)+b*d^(1/3)))/c^(4/3)

```

Mathematica [A] (verified)

Time = 0.67 (sec) , antiderivative size = 791, normalized size of antiderivative = 0.95

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^3}} dx$$

$$3\sqrt[3]{c} \log(1 - a - bx) - 3a\sqrt[3]{c} \log(1 - a - bx) - 3b\sqrt[3]{cx} \log(1 - a - bx) + 3\sqrt[3]{c} \log(1 + a + bx) + 3a\sqrt[3]{c} \log(1 + a + bx)$$

input

```
Integrate[ArcTanh[a + b*x]/(c + d/x^3),x]
```

output

```
(3*c^(1/3)*Log[1 - a - b*x] - 3*a*c^(1/3)*Log[1 - a - b*x] - 3*b*c^(1/3)*x
*Log[1 - a - b*x] + 3*c^(1/3)*Log[1 + a + b*x] + 3*a*c^(1/3)*Log[1 + a + b
*x] + 3*b*c^(1/3)*x*Log[1 + a + b*x] + b*d^(1/3)*Log[1 - a - b*x]*Log[(b*(
d^(1/3) + c^(1/3)*x))/(-((-1 + a)*c^(1/3)) + b*d^(1/3))] - b*d^(1/3)*Log[1
+ a + b*x]*Log[(b*(d^(1/3) + c^(1/3)*x))/(-((1 + a)*c^(1/3)) + b*d^(1/3))
] + (-1)^(2/3)*b*d^(1/3)*Log[1 - a - b*x]*Log[(b*(d^(1/3) - (-1)^(1/3)*c^(
1/3)*x))/((-1)^(1/3)*(-1 + a)*c^(1/3) + b*d^(1/3))] - (-1)^(2/3)*b*d^(1/3)
*Log[1 + a + b*x]*Log[(b*(d^(1/3) - (-1)^(1/3)*c^(1/3)*x))/((-1)^(1/3)*(1
+ a)*c^(1/3) + b*d^(1/3))] - (-1)^(1/3)*b*d^(1/3)*Log[1 - a - b*x]*Log[(b*
(d^(1/3) + (-1)^(2/3)*c^(1/3)*x))/(-((-1)^(2/3)*(-1 + a)*c^(1/3)) + b*d^(1
/3))] + (-1)^(1/3)*b*d^(1/3)*Log[1 + a + b*x]*Log[(b*(d^(1/3) + (-1)^(2/3)
*c^(1/3)*x))/(-((-1)^(2/3)*(1 + a)*c^(1/3)) + b*d^(1/3))] + b*d^(1/3)*Poly
Log[2, (c^(1/3)*(-1 + a + b*x))/((-1 + a)*c^(1/3) - b*d^(1/3))] - (-1)^(1/
3)*b*d^(1/3)*PolyLog[2, ((-1)^(2/3)*c^(1/3)*(-1 + a + b*x))/((-1)^(2/3)*(-
1 + a)*c^(1/3) - b*d^(1/3))] + (-1)^(2/3)*b*d^(1/3)*PolyLog[2, ((-1)^(1/3)
*c^(1/3)*(-1 + a + b*x))/((-1)^(1/3)*(-1 + a)*c^(1/3) + b*d^(1/3))] - b*d^(
1/3)*PolyLog[2, (c^(1/3)*(1 + a + b*x))/((1 + a)*c^(1/3) - b*d^(1/3))] +
(-1)^(1/3)*b*d^(1/3)*PolyLog[2, ((-1)^(2/3)*c^(1/3)*(1 + a + b*x))/((-1)^(
2/3)*(1 + a)*c^(1/3) - b*d^(1/3))] - (-1)^(2/3)*b*d^(1/3)*PolyLog[2, ((-1)
^(1/3)*c^(1/3)*(1 + a + b*x))/((-1)^(1/3)*(1 + a)*c^(1/3) + b*d^(1/3))]...
```

Rubi [A] (verified)

Time = 1.57 (sec) , antiderivative size = 847, normalized size of antiderivative = 1.02, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6665, 2856, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^3}} dx$$

$$\downarrow \text{6665}$$

$$\frac{1}{2} \int \frac{\log(a + bx + 1)}{c + \frac{d}{x^3}} dx - \frac{1}{2} \int \frac{\log(-a - bx + 1)}{c + \frac{d}{x^3}} dx$$

$$\downarrow \text{2856}$$

$$\frac{1}{2} \int \left(\frac{\log(a + bx + 1)}{c} - \frac{d \log(a + bx + 1)}{c(cx^3 + d)} \right) dx -$$

$$\frac{1}{2} \int \left(\frac{\log(-a - bx + 1)}{c} - \frac{d \log(-a - bx + 1)}{c(cx^3 + d)} \right) dx$$

↓ 2009

$$\frac{1}{2} \left(\frac{x}{c} + \frac{(-a - bx + 1) \log(-a - bx + 1)}{bc} + \frac{\sqrt[3]{d} \log(-a - bx + 1) \log \left(\frac{b(\sqrt[3]{cx} + \sqrt[3]{d})}{\sqrt[3]{c(1-a)} + b\sqrt[3]{d}} \right)}{3c^{4/3}} + \frac{(-1)^{2/3} \sqrt[3]{d} \log(-a - bx + 1)}{3c^{4/3}} \right)$$

$$\frac{1}{2} \left(-\frac{x}{c} + \frac{(a + bx + 1) \log(a + bx + 1)}{bc} - \frac{\sqrt[3]{d} \log(a + bx + 1) \log \left(-\frac{b(\sqrt[3]{cx} + \sqrt[3]{d})}{(a+1)\sqrt[3]{c} - b\sqrt[3]{d}} \right)}{3c^{4/3}} - \frac{(-1)^{2/3} \sqrt[3]{d} \log(a + bx + 1)}{3c^{4/3}} \right)$$

input Int[ArcTanh[a + b*x]/(c + d/x^3), x]

output

```
(x/c + ((1 - a - b*x)*Log[1 - a - b*x])/(b*c) + (d^(1/3)*Log[1 - a - b*x]*
Log[(b*(d^(1/3) + c^(1/3)*x))/((1 - a)*c^(1/3) + b*d^(1/3))])/(3*c^(4/3))
+ ((-1)^(2/3)*d^(1/3)*Log[1 - a - b*x]*Log[-(b*(d^(1/3) - (-1)^(1/3)*c^(1
/3)*x))/((-1)^(1/3)*(1 - a)*c^(1/3) - b*d^(1/3))])/(3*c^(4/3)) - ((-1)^(1
/3)*d^(1/3)*Log[1 - a - b*x]*Log[(b*(d^(1/3) + (-1)^(2/3)*c^(1/3)*x))/((-1
)^(2/3)*(1 - a)*c^(1/3) + b*d^(1/3))])/(3*c^(4/3)) + ((-1)^(2/3)*d^(1/3)*P
olyLog[2, ((-1)^(1/3)*c^(1/3)*(1 - a - b*x))/((-1)^(1/3)*(1 - a)*c^(1/3) -
b*d^(1/3))])/(3*c^(4/3)) + (d^(1/3)*PolyLog[2, (c^(1/3)*(1 - a - b*x))/((
1 - a)*c^(1/3) + b*d^(1/3))])/(3*c^(4/3)) - ((-1)^(1/3)*d^(1/3)*PolyLog[2,
((-1)^(2/3)*c^(1/3)*(1 - a - b*x))/((-1)^(2/3)*(1 - a)*c^(1/3) + b*d^(1/3
))])/(3*c^(4/3)))/2 + (-x/c) + ((1 + a + b*x)*Log[1 + a + b*x])/(b*c) - (
d^(1/3)*Log[1 + a + b*x]*Log[-(b*(d^(1/3) + c^(1/3)*x))/((1 + a)*c^(1/3)
- b*d^(1/3))])/(3*c^(4/3)) - ((-1)^(2/3)*d^(1/3)*Log[1 + a + b*x]*Log[(b*
(d^(1/3) - (-1)^(1/3)*c^(1/3)*x))/((-1)^(1/3)*(1 + a)*c^(1/3) + b*d^(1/3)
])/(3*c^(4/3)) + ((-1)^(1/3)*d^(1/3)*Log[1 + a + b*x]*Log[-(b*(d^(1/3) +
(-1)^(2/3)*c^(1/3)*x))/((-1)^(2/3)*(1 + a)*c^(1/3) - b*d^(1/3))])/(3*c^(4
/3)) - (d^(1/3)*PolyLog[2, (c^(1/3)*(1 + a + b*x))/((1 + a)*c^(1/3) - b*d^(
1/3))])/(3*c^(4/3)) + ((-1)^(1/3)*d^(1/3)*PolyLog[2, ((-1)^(2/3)*c^(1/3)*
(1 + a + b*x))/((-1)^(2/3)*(1 + a)*c^(1/3) - b*d^(1/3))])/(3*c^(4/3)) - ((
-1)^(2/3)*d^(1/3)*PolyLog[2, ((-1)^(1/3)*c^(1/3)*(1 + a + b*x))/((-1)^(...
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2856 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]*(b_.))^(p_.)*((f_) + (g_.
)*(x_)^(r_))^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log[c*(d + e*x)
^n])^p, (f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, r}, x] && I
GtQ[p, 0] && IntegerQ[q] && (GtQ[q, 0] || (IntegerQ[r] && NeQ[r, 1]))`

rule 6665 `Int[ArcTanh[(c_) + (d_.)*(x_)]/((e_) + (f_.)*(x_)^(n_.)), x_Symbol] := Simp
[1/2 Int[Log[1 + c + d*x]/(e + f*x^n), x], x] - Simp[1/2 Int[Log[1 - c
- d*x]/(e + f*x^n), x], x] /; FreeQ[{c, d, e, f}, x] && RationalQ[n]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.56 (sec) , antiderivative size = 366, normalized size of antiderivative = 0.44

| method | result |
|-------------------|--|
| risch | $\frac{\ln(bx+a+1)x}{2c} + \frac{\ln(bx+a+1)a}{2bc} + \frac{\ln(bx+a+1)}{2bc} - \frac{1}{bc} - \frac{b^2 d}{\sqrt{-R \text{I} = \text{RootOf}(c_Z^3 + (-3ac-3c)_Z^2 + (3a^2c+6ac+3c)}}$ |
| derivativedivides | $\frac{\arctanh\left(\frac{bx+a}{c}\right) + \frac{\arctanh(bx+a) \left(\sqrt{-R = \text{RootOf}(c_Z^3 - 3ac_Z^2 + 3a^2c_Z - a^3c + b^3d)} - \frac{\ln\left(\frac{bx - R + a}{R^2 + 2_R a - a^2}\right)}{d} \right) d b^3}{3c^2} + \frac{\ln(bx+a)}{c}$ |
| default | $\frac{\arctanh\left(\frac{bx+a}{c}\right) + \frac{\arctanh(bx+a) \left(\sqrt{-R = \text{RootOf}(c_Z^3 - 3ac_Z^2 + 3a^2c_Z - a^3c + b^3d)} - \frac{\ln\left(\frac{bx - R + a}{R^2 + 2_R a - a^2}\right)}{d} \right) d b^3}{3c^2} + \frac{\ln(bx+a)}{c}$ |

input `int(arctanh(b*x+a)/(c+d/x^3), x, method=_RETURNVERBOSE)`

output

```
1/2/c*ln(b*x+a+1)*x+1/2/b/c*ln(b*x+a+1)*a+1/2/b/c*ln(b*x+a+1)-1/b/c-1/6*b^
2*d/c^2*sum(1/(_R1^2-2*_R1*a+a^2-2*_R1+2*a+1)*(ln(b*x+a+1)*ln((-b*x+_R1-a-
1)/_R1)+dilog((-b*x+_R1-a-1)/_R1)),_R1=RootOf(c*_Z^3+(-3*a*c-3*c)*_Z^2+(3*
a^2*c+6*a*c+3*c)*_Z-a^3*c+b^3*d-3*a^2*c-3*a*c-c))-1/2/c*ln(-b*x-a+1)*x-1/2
/b/c*ln(-b*x-a+1)*a+1/2/b/c*ln(-b*x-a+1)+1/6*b^2*d/c^2*sum(1/(_R1^2+2*_R1*
a+a^2-2*_R1-2*a+1)*(ln(-b*x-a+1)*ln((b*x+_R1+a-1)/_R1)+dilog((b*x+_R1+a-1)
/_R1)),_R1=RootOf(c*_Z^3+(3*a*c-3*c)*_Z^2+(3*a^2*c-6*a*c+3*c)*_Z+a^3*c-b^3
*d-3*a^2*c+3*a*c-c))
```

Fricas [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^3}} dx = \int \frac{\operatorname{artanh}(bx + a)}{c + \frac{d}{x^3}} dx$$

input

```
integrate(arctanh(b*x+a)/(c+d/x^3),x, algorithm="fricas")
```

output

```
integral(x^3*arctanh(b*x + a)/(c*x^3 + d), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^3}} dx = \text{Timed out}$$

input

```
integrate(atanh(b*x+a)/(c+d/x**3),x)
```

output

```
Timed out
```

Maxima [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^3}} dx = \int \frac{\operatorname{artanh}(bx + a)}{c + \frac{d}{x^3}} dx$$

input `integrate(arctanh(b*x+a)/(c+d/x^3),x, algorithm="maxima")`

output `integrate(arctanh(b*x + a)/(c + d/x^3), x)`

Giac [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^3}} dx = \int \frac{\operatorname{artanh}(bx + a)}{c + \frac{d}{x^3}} dx$$

input `integrate(arctanh(b*x+a)/(c+d/x^3),x, algorithm="giac")`

output `sage0*x`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^3}} dx = \int \frac{\operatorname{atanh}(a + bx)}{c + \frac{d}{x^3}} dx$$

input `int(atanh(a + b*x)/(c + d/x^3),x)`

output `int(atanh(a + b*x)/(c + d/x^3), x)`

Reduce [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{x^3}} dx = \int \frac{\operatorname{atanh}(bx + a) x^3}{c x^3 + d} dx$$

input `int(atanh(b*x+a)/(c+d/x^3),x)`

output `int((atanh(a + b*x)*x**3)/(c*x**3 + d),x)`

3.59 $\int \frac{\operatorname{arctanh}(a+bx)}{1-x^2} dx$

| | |
|---|-----|
| Optimal result | 524 |
| Mathematica [A] (verified) | 525 |
| Rubi [A] (verified) | 525 |
| Maple [A] (verified) | 527 |
| Fricas [F] | 527 |
| Sympy [F] | 528 |
| Maxima [A] (verification not implemented) | 528 |
| Giac [F] | 529 |
| Mupad [F(-1)] | 529 |
| Reduce [F] | 529 |

Optimal result

Integrand size = 16, antiderivative size = 135

$$\int \frac{\operatorname{arctanh}(a+bx)}{1-x^2} dx = -\frac{1}{2}\operatorname{arctanh}(a+bx) \log\left(-\frac{2b(1-x)}{(1-a-b)(1+a+bx)}\right) + \frac{1}{2}\operatorname{arctanh}(a+bx) \log\left(\frac{2b(1+x)}{(1-a+b)(1+a+bx)}\right) + \frac{1}{4}\operatorname{PolyLog}\left(2, 1 + \frac{2b(1-x)}{(1-a-b)(1+a+bx)}\right) - \frac{1}{4}\operatorname{PolyLog}\left(2, 1 - \frac{2b(1+x)}{(1-a+b)(1+a+bx)}\right)$$

output

```
-1/2*arctanh(b*x+a)*ln(-2*b*(1-x)/(1-a-b)/(b*x+a+1))+1/2*arctanh(b*x+a)*ln
(2*b*(1+x)/(1-a+b)/(b*x+a+1))+1/4*polylog(2,1+2*b*(1-x)/(1-a-b)/(b*x+a+1))
-1/4*polylog(2,1-2*b*(1+x)/(1-a+b)/(b*x+a+1))
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.50

$$\int \frac{\operatorname{arctanh}(a + bx)}{1 - x^2} dx = \frac{1}{4} \log \left(-\frac{b(1-x)}{1-a-b} \right) \log(1-a-bx) - \frac{1}{4} \log \left(\frac{b(1+x)}{1-a+b} \right) \log(1-a-bx) - \frac{1}{4} \log \left(\frac{b(1-x)}{1+a+b} \right) \log(1+a+bx) + \frac{1}{4} \log \left(-\frac{b(1+x)}{1+a-b} \right) \log(1+a+bx) + \frac{1}{4} \operatorname{PolyLog} \left(2, \frac{1-a-bx}{1-a-b} \right) - \frac{1}{4} \operatorname{PolyLog} \left(2, \frac{1-a-bx}{1-a+b} \right) + \frac{1}{4} \operatorname{PolyLog} \left(2, \frac{1+a+bx}{1+a-b} \right) - \frac{1}{4} \operatorname{PolyLog} \left(2, \frac{1+a+bx}{1+a+b} \right)$$

input `Integrate[ArcTanh[a + b*x]/(1 - x^2), x]`

output `(Log[-((b*(1 - x))/(1 - a - b))]*Log[1 - a - b*x])/4 - (Log[(b*(1 + x))/(1 - a + b)]*Log[1 - a - b*x])/4 - (Log[(b*(1 - x))/(1 + a + b)]*Log[1 + a + b*x])/4 + (Log[-((b*(1 + x))/(1 + a - b))]*Log[1 + a + b*x])/4 + PolyLog[2, (1 - a - b*x)/(1 - a - b)]/4 - PolyLog[2, (1 - a - b*x)/(1 - a + b)]/4 + PolyLog[2, (1 + a + b*x)/(1 + a - b)]/4 - PolyLog[2, (1 + a + b*x)/(1 + a + b)]/4`

Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.58, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {6665, 2856, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(a + bx)}{1 - x^2} dx$$

$$\frac{1}{2} \int \frac{\log(a + bx + 1)}{1 - x^2} dx - \frac{1}{2} \int \frac{\log(-a - bx + 1)}{1 - x^2} dx$$

$$\frac{1}{2} \int \left(\frac{\log(a + bx + 1)}{2(1 - x)} + \frac{\log(a + bx + 1)}{2(x + 1)} \right) dx - \frac{1}{2} \int \left(\frac{\log(-a - bx + 1)}{2(1 - x)} + \frac{\log(-a - bx + 1)}{2(x + 1)} \right) dx$$

$$\frac{1}{2} \left(\frac{1}{2} \text{PolyLog} \left(2, \frac{-a - bx + 1}{-a - b + 1} \right) - \frac{1}{2} \text{PolyLog} \left(2, \frac{-a - bx + 1}{-a + b + 1} \right) + \frac{1}{2} \log \left(\frac{b(1 - x)}{-a - b + 1} \right) \log(-a - bx + 1) - \frac{1}{2} \log \left(\frac{b(1 - x)}{-a + b + 1} \right) \log(-a - bx + 1) \right) - \frac{1}{2} \left(\frac{1}{2} \text{PolyLog} \left(2, \frac{a + bx + 1}{a - b + 1} \right) - \frac{1}{2} \text{PolyLog} \left(2, \frac{a + bx + 1}{a + b + 1} \right) - \frac{1}{2} \log \left(\frac{b(1 - x)}{a + b + 1} \right) \log(a + bx + 1) + \frac{1}{2} \log \left(\frac{b(1 - x)}{a - b + 1} \right) \log(a + bx + 1) \right)$$

input `Int[ArcTanh[a + b*x]/(1 - x^2), x]`

output `((Log[-((b*(1 - x))/(1 - a - b))]*Log[1 - a - b*x])/2 - (Log[(b*(1 + x))/(1 - a + b)]*Log[1 - a - b*x])/2 + PolyLog[2, (1 - a - b*x)/(1 - a - b)]/2 - PolyLog[2, (1 - a - b*x)/(1 - a + b)]/2)/2 + (-1/2*(Log[(b*(1 - x))/(1 + a + b)]*Log[1 + a + b*x]) + (Log[-((b*(1 + x))/(1 + a - b))]*Log[1 + a + b*x])/2 + PolyLog[2, (1 + a + b*x)/(1 + a - b)]/2 - PolyLog[2, (1 + a + b*x)/(1 + a + b)]/2)/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2856 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]*(b_.))^(p_.)*((f_) + (g_.)*(x_)^(r_.))^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log[c*(d + e*x)^n])^p, (f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, r}, x] && I GtQ[p, 0] && IntegerQ[q] && (GtQ[q, 0] || (IntegerQ[r] && NeQ[r, 1]))`

rule 6665

```
Int[ArcTanh[(c_) + (d_)*(x_)]/((e_) + (f_)*(x_)^(n_)), x_Symbol] := Simp
[1/2 Int[Log[1 + c + d*x]/(e + f*x^n), x], x] - Simp[1/2 Int[Log[1 - c
- d*x]/(e + f*x^n), x], x] /; FreeQ[{c, d, e, f}, x] && RationalQ[n]
```

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.36

| method | result |
|-------------------|---|
| risch | $-\frac{\ln(-bx-a+1)\ln\left(\frac{-bx-b}{-b-1+a}\right)}{4} - \frac{\operatorname{dilog}\left(\frac{-bx-b}{-b-1+a}\right)}{4} + \frac{\ln(-bx-a+1)\ln\left(\frac{-bx+b}{b-1+a}\right)}{4} + \frac{\operatorname{dilog}\left(\frac{-bx+b}{b-1+a}\right)}{4} + \frac{\ln(bx+a+1)}{4}$ |
| parts | $\operatorname{arctanh}(x)\operatorname{arctanh}(bx+a) - b\left(\frac{\operatorname{arctanh}(x)\ln(bx+a+1)}{2b} - \frac{\operatorname{arctanh}(x)\ln(bx+a-1)}{2b} - \frac{\ln(bx+a-1)}{4b}\right)$ |
| derivativedivides | $\frac{\operatorname{arctanh}(bx+a)b\ln(-bx-b)}{2} - \frac{\operatorname{arctanh}(bx+a)b\ln(-bx+b)}{2} + b^2\left(\frac{\operatorname{dilog}\left(\frac{-bx-a-1}{-1+b-a}\right)}{2} - \frac{\ln(-bx-b)\ln\left(\frac{-bx-a-1}{-1+b-a}\right)}{2} + \frac{\operatorname{dilog}\left(\frac{-bx-a+1}{1-a+b}\right)}{2}\right)$ |
| default | $\frac{\operatorname{arctanh}(bx+a)b\ln(-bx-b)}{2} - \frac{\operatorname{arctanh}(bx+a)b\ln(-bx+b)}{2} + b^2\left(\frac{\operatorname{dilog}\left(\frac{-bx-a-1}{-1+b-a}\right)}{2} - \frac{\ln(-bx-b)\ln\left(\frac{-bx-a-1}{-1+b-a}\right)}{2} + \frac{\operatorname{dilog}\left(\frac{-bx-a+1}{1-a+b}\right)}{2}\right)$ |

input

```
int(arctanh(b*x+a)/(-x^2+1),x,method=_RETURNVERBOSE)
```

output

```
-1/4*ln(-b*x-a+1)*ln((-b*x-b)/(-b-1+a))-1/4*dilog((-b*x-b)/(-b-1+a))+1/4*ln
(-b*x-a+1)*ln((-b*x+b)/(b-1+a))+1/4*dilog((-b*x+b)/(b-1+a))+1/4*ln(b*x+a+
1)*ln((b*x+b)/(-1+b-a))+1/4*dilog((b*x+b)/(-1+b-a))-1/4*ln(b*x+a+1)*ln((b*
x-b)/(-1-b-a))-1/4*dilog((b*x-b)/(-1-b-a))
```

Fricas [F]

$$\int \frac{\operatorname{arctanh}(a+bx)}{1-x^2} dx = \int -\frac{\operatorname{arctanh}(bx+a)}{x^2-1} dx$$

input

```
integrate(arctanh(b*x+a)/(-x^2+1),x, algorithm="fricas")
```


output `integral(-arctanh(b*x + a)/(x^2 - 1), x)`

Sympy [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{1 - x^2} dx = - \int \frac{\operatorname{atanh}(a + bx)}{x^2 - 1} dx$$

input `integrate(atanh(b*x+a)/(-x**2+1), x)`

output `-Integral(atanh(a + b*x)/(x**2 - 1), x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.47

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(a + bx)}{1 - x^2} dx \\ &= \frac{1}{4} b \left(\frac{\log(x - 1) \log\left(\frac{bx-b}{a+b+1} + 1\right) + \operatorname{Li}_2\left(-\frac{bx-b}{a+b+1}\right)}{b} - \frac{\log(x - 1) \log\left(\frac{bx-b}{a+b-1} + 1\right) + \operatorname{Li}_2\left(-\frac{bx-b}{a+b-1}\right)}{b} - \frac{\log(x + 1) \log\left(\frac{bx+b}{a-b+1} + 1\right) + \operatorname{Li}_2\left(-\frac{bx+b}{a-b+1}\right)}{b} \right. \\ & \quad \left. + \frac{1}{2} (\log(x + 1) - \log(x - 1)) \operatorname{artanh}(bx + a) \right) \end{aligned}$$

input `integrate(arctanh(b*x+a)/(-x^2+1), x, algorithm="maxima")`

output `1/4*b*((log(x - 1)*log((b*x - b)/(a + b + 1) + 1) + dilog(-(b*x - b)/(a + b + 1)))/b - (log(x - 1)*log((b*x - b)/(a + b - 1) + 1) + dilog(-(b*x - b)/(a + b - 1)))/b - (log(x + 1)*log((b*x + b)/(a - b + 1) + 1) + dilog(-(b*x + b)/(a - b + 1)))/b + (log(x + 1)*log((b*x + b)/(a - b - 1) + 1) + dilog(-(b*x + b)/(a - b - 1)))/b) + 1/2*(log(x + 1) - log(x - 1))*arctanh(b*x + a)`

Giac [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{1 - x^2} dx = \int -\frac{\operatorname{artanh}(bx + a)}{x^2 - 1} dx$$

input `integrate(arctanh(b*x+a)/(-x^2+1),x, algorithm="giac")`

output `integrate(-arctanh(b*x + a)/(x^2 - 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{1 - x^2} dx = - \int \frac{\operatorname{atanh}(a + bx)}{x^2 - 1} dx$$

input `int(-atanh(a + b*x)/(x^2 - 1),x)`

output `-int(atanh(a + b*x)/(x^2 - 1), x)`

Reduce [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{1 - x^2} dx = - \left(\int \frac{\operatorname{atanh}(bx + a)}{x^2 - 1} dx \right)$$

input `int(atanh(b*x+a)/(-x^2+1),x)`

output `- int(atanh(a + b*x)/(x**2 - 1),x)`

$$3.60 \quad \int \frac{a+b \operatorname{arctanh}(c+dx)}{e+f\sqrt{x}} dx$$

| | |
|--------------------------------------|-----|
| Optimal result | 531 |
| Mathematica [A] (verified) | 532 |
| Rubi [F] | 533 |
| Maple [A] (verified) | 534 |
| Fricas [F] | 535 |
| Sympy [F(-1)] | 535 |
| Maxima [F] | 536 |
| Giac [F] | 536 |
| Mupad [F(-1)] | 536 |
| Reduce [F] | 537 |

Optimal result

Integrand size = 22, antiderivative size = 618

$$\begin{aligned}
\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + f\sqrt{x}} dx = & \frac{2a\sqrt{x}}{f} + \frac{2b\sqrt{1+c} \operatorname{arctan}\left(\frac{\sqrt{d}\sqrt{x}}{\sqrt{1+c}}\right)}{\sqrt{d}f} \\
& - \frac{2b\sqrt{1-c} \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{x}}{\sqrt{1-c}}\right)}{\sqrt{d}f} \\
& + \frac{be \log\left(\frac{f(\sqrt{-1-c}-\sqrt{d}\sqrt{x})}{\sqrt{de}+\sqrt{-1-c}f}\right) \log(e + f\sqrt{x})}{f^2} \\
& - \frac{be \log\left(\frac{f(\sqrt{1-c}-\sqrt{d}\sqrt{x})}{\sqrt{de}+\sqrt{1-c}f}\right) \log(e + f\sqrt{x})}{f^2} \\
& + \frac{be \log\left(-\frac{f(\sqrt{-1-c}+\sqrt{d}\sqrt{x})}{\sqrt{de}-\sqrt{-1-c}f}\right) \log(e + f\sqrt{x})}{f^2} \\
& - \frac{be \log\left(-\frac{f(\sqrt{1-c}+\sqrt{d}\sqrt{x})}{\sqrt{de}-\sqrt{1-c}f}\right) \log(e + f\sqrt{x})}{f^2} \\
& - \frac{b\sqrt{x} \log(1-c-dx)}{f} \\
& - \frac{e \log(e + f\sqrt{x}) (a - b \log(1-c-dx))}{f^2} \\
& + \frac{b\sqrt{x} \log(1+c+dx)}{f} \\
& - \frac{e \log(e + f\sqrt{x}) (a + b \log(1+c+dx))}{f^2} \\
& + \frac{be \operatorname{PolyLog}\left(2, \frac{\sqrt{d}(e+f\sqrt{x})}{\sqrt{de}-\sqrt{-1-c}f}\right)}{f^2} \\
& + \frac{be \operatorname{PolyLog}\left(2, \frac{\sqrt{d}(e+f\sqrt{x})}{\sqrt{de}+\sqrt{-1-c}f}\right)}{f^2} \\
& - \frac{be \operatorname{PolyLog}\left(2, \frac{\sqrt{d}(e+f\sqrt{x})}{\sqrt{de}-\sqrt{1-c}f}\right)}{f^2} \\
& - \frac{be \operatorname{PolyLog}\left(2, \frac{\sqrt{d}(e+f\sqrt{x})}{\sqrt{de}+\sqrt{1-c}f}\right)}{f^2}
\end{aligned}$$

output

```

2*a*x^(1/2)/f+2*b*(1+c)^(1/2)*arctan(d^(1/2)*x^(1/2)/(1+c)^(1/2))/d^(1/2)/
f-2*b*(1-c)^(1/2)*arctanh(d^(1/2)*x^(1/2)/(1-c)^(1/2))/d^(1/2)/f+b*e*ln(f*
((-1-c)^(1/2)-d^(1/2)*x^(1/2))/(d^(1/2)*e+(-1-c)^(1/2)*f))*ln(e+f*x^(1/2))
/f^2-b*e*ln(f*((1-c)^(1/2)-d^(1/2)*x^(1/2))/(d^(1/2)*e+(1-c)^(1/2)*f))*ln(
e+f*x^(1/2))/f^2+b*e*ln(-f*((-1-c)^(1/2)+d^(1/2)*x^(1/2))/(d^(1/2)*e-(-1-c)
^(1/2)*f))*ln(e+f*x^(1/2))/f^2-b*e*ln(-f*((1-c)^(1/2)+d^(1/2)*x^(1/2))/(d
^(1/2)*e-(1-c)^(1/2)*f))*ln(e+f*x^(1/2))/f^2-b*x^(1/2)*ln(-d*x-c+1)/f-e*ln
(e+f*x^(1/2))*(a-b*ln(-d*x-c+1))/f^2+b*x^(1/2)*ln(d*x+c+1)/f-e*ln(e+f*x^(1
/2))*(a+b*ln(d*x+c+1))/f^2+b*e*polylog(2,d^(1/2)*(e+f*x^(1/2))/(d^(1/2)*e-
(-1-c)^(1/2)*f))/f^2+b*e*polylog(2,d^(1/2)*(e+f*x^(1/2))/(d^(1/2)*e+(-1-c)
^(1/2)*f))/f^2-b*e*polylog(2,d^(1/2)*(e+f*x^(1/2))/(d^(1/2)*e-(1-c)^(1/2)*
f))/f^2-b*e*polylog(2,d^(1/2)*(e+f*x^(1/2))/(d^(1/2)*e+(1-c)^(1/2)*f))/f^2

```

Mathematica [A] (verified)

Time = 12.03 (sec) , antiderivative size = 576, normalized size of antiderivative = 0.93

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + f\sqrt{x}} dx$$

$$= \frac{2a(f\sqrt{x} - e \log(e + f\sqrt{x})) + b \left(\frac{2\sqrt{1+cf} \operatorname{arctan}\left(\frac{\sqrt{d}\sqrt{x}}{\sqrt{1+c}}\right)}{\sqrt{d}} - \frac{2\sqrt{1-cf} \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{x}}{\sqrt{1-c}}\right)}{\sqrt{d}} + e \log\left(\frac{f(\sqrt{-1-c}-\sqrt{d}\sqrt{x})}{\sqrt{de+\sqrt{-1-c}f}}\right) \right)}{1}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])/(e + f*Sqrt[x]),x]
```

output

```
(2*a*(f*Sqrt[x] - e*Log[e + f*Sqrt[x]]) + b*((2*Sqrt[1 + c]*f*ArcTan[(Sqrt[d]*Sqrt[x])/Sqrt[1 + c]])/Sqrt[d] - (2*Sqrt[1 - c]*f*ArcTanh[(Sqrt[d]*Sqrt[x])/Sqrt[1 - c]])/Sqrt[d] + e*Log[(f*(Sqrt[-1 - c] - Sqrt[d]*Sqrt[x]))/(Sqrt[d]*e + Sqrt[-1 - c]*f)]*Log[e + f*Sqrt[x]] - e*Log[(f*(Sqrt[1 - c] - Sqrt[d]*Sqrt[x]))/(Sqrt[d]*e + Sqrt[1 - c]*f)]*Log[e + f*Sqrt[x]] + e*Log[(f*(Sqrt[-1 - c] + Sqrt[d]*Sqrt[x]))/(-(Sqrt[d]*e) + Sqrt[-1 - c]*f)]*Log[e + f*Sqrt[x]] - e*Log[(f*(Sqrt[1 - c] + Sqrt[d]*Sqrt[x]))/(-(Sqrt[d]*e) + Sqrt[1 - c]*f)]*Log[e + f*Sqrt[x]] - f*Sqrt[x]*Log[1 - c - d*x] + e*Log[e + f*Sqrt[x]]*Log[1 - c - d*x] + f*Sqrt[x]*Log[1 + c + d*x] - e*Log[e + f*Sqrt[x]]*Log[1 + c + d*x] + e*PolyLog[2, (Sqrt[d]*(e + f*Sqrt[x]))/(Sqrt[d]*e - Sqrt[-1 - c]*f)] + e*PolyLog[2, (Sqrt[d]*(e + f*Sqrt[x]))/(Sqrt[d]*e + Sqrt[-1 - c]*f)] - e*PolyLog[2, (Sqrt[d]*(e + f*Sqrt[x]))/(Sqrt[d]*e - Sqrt[1 - c]*f)] - e*PolyLog[2, (Sqrt[d]*(e + f*Sqrt[x]))/(Sqrt[d]*e + Sqrt[1 - c]*f)))/f^2
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}(c + dx)}{e + f\sqrt{x}} dx \\
 & \quad \downarrow 7267 \\
 & 2 \int \frac{\sqrt{x}(a + b \operatorname{arctanh}(c + dx))}{e + f\sqrt{x}} d\sqrt{x} \\
 & \quad \downarrow 7293 \\
 & 2 \int \left(\frac{\sqrt{x}a}{e + f\sqrt{x}} + \frac{b\sqrt{x} \operatorname{arctanh}(c + dx)}{e + f\sqrt{x}} \right) d\sqrt{x} \\
 & \quad \downarrow 2009 \\
 & 2 \left(-\frac{be \int \frac{\operatorname{arctanh}(c+dx)}{e+f\sqrt{x}} d\sqrt{x}}{f} - \frac{ae \log(e + f\sqrt{x})}{f^2} + \frac{a\sqrt{x}}{f} + \frac{b\sqrt{c+1} \arctan\left(\frac{\sqrt{d}\sqrt{x}}{\sqrt{c+1}}\right)}{\sqrt{df}} - \frac{b\sqrt{1-c} \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{x}}{\sqrt{1-c}}\right)}{\sqrt{df}} \right)
 \end{aligned}$$

input

```
Int[(a + b*ArcTanh[c + d*x])/(e + f*Sqrt[x]),x]
```


output

```
2*a*x^(1/2)/f-2*a*e/f^2*ln(e+f*x^(1/2))+2*b*(arctanh(d*x+c)/f*x^(1/2)-arctanh(d*x+c)*e/f^2*ln(e+f*x^(1/2))-2*d/f^2*(-f^2*(1/2*(1+c)/d/(c*d*f^2+d*f^2)^(1/2)*arctan(1/2*(-2*d*e+2*(e+f*x^(1/2))*d)/(c*d*f^2+d*f^2)^(1/2))+1/2*(1-c)/d/(c*d*f^2-d*f^2)^(1/2)*arctan(1/2*(-2*d*e+2*(e+f*x^(1/2))*d)/(c*d*f^2-d*f^2)^(1/2)))-e*f^2*(1/2/f^2*(1/2*ln(e+f*x^(1/2))*(ln((d*e-(e+f*x^(1/2))*d+(-c*d*f^2-d*f^2)^(1/2))/(d*e+(-c*d*f^2-d*f^2)^(1/2))))+ln((-d*e+(e+f*x^(1/2))*d+(-c*d*f^2-d*f^2)^(1/2))/(-d*e+(-c*d*f^2-d*f^2)^(1/2))))/d+1/2*(dilog((d*e-(e+f*x^(1/2))*d+(-c*d*f^2-d*f^2)^(1/2))/(d*e+(-c*d*f^2-d*f^2)^(1/2)))+dilog((-d*e+(e+f*x^(1/2))*d+(-c*d*f^2-d*f^2)^(1/2))/(-d*e+(-c*d*f^2-d*f^2)^(1/2))))/d)+1/2/f^2*(-1/2*ln(e+f*x^(1/2))*(ln((d*e-(e+f*x^(1/2))*d+(-c*d*f^2+d*f^2)^(1/2))/(d*e+(-c*d*f^2+d*f^2)^(1/2))))+ln((-d*e+(e+f*x^(1/2))*d+(-c*d*f^2+d*f^2)^(1/2))/(-d*e+(-c*d*f^2+d*f^2)^(1/2))))/d-1/2*(dilog((d*e-(e+f*x^(1/2))*d+(-c*d*f^2+d*f^2)^(1/2))/(d*e+(-c*d*f^2+d*f^2)^(1/2)))+dilog((-d*e+(e+f*x^(1/2))*d+(-c*d*f^2+d*f^2)^(1/2))/(-d*e+(-c*d*f^2+d*f^2)^(1/2))))/d))))
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + f\sqrt{x}} dx = \int \frac{b \operatorname{arctanh}(dx + c) + a}{f\sqrt{x} + e} dx$$

input

```
integrate((a+b*arctanh(d*x+c))/(e+f*x^(1/2)),x, algorithm="fricas")
```

output

```
integral(-(b*e*arctanh(d*x + c) + a*e - (b*f*arctanh(d*x + c) + a*f)*sqrt(x))/(f^2*x - e^2), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + f\sqrt{x}} dx = \text{Timed out}$$

input

```
integrate((a+b*atanh(d*x+c))/(e+f*x**(1/2)),x)
```


output Timed out

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + f\sqrt{x}} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{f\sqrt{x} + e} dx$$

input `integrate((a+b*arctanh(d*x+c))/(e+f*x^(1/2)),x, algorithm="maxima")`

output `-2*a*(e*log(f*sqrt(x) + e)/f^2 - sqrt(x)/f) + b*integrate(1/2*log(d*x + c + 1)/(f*sqrt(x) + e), x) - b*integrate(1/2*log(-d*x - c + 1)/(f*sqrt(x) + e), x)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + f\sqrt{x}} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{f\sqrt{x} + e} dx$$

input `integrate((a+b*arctanh(d*x+c))/(e+f*x^(1/2)),x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)/(f*sqrt(x) + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + f\sqrt{x}} dx = \int \frac{a + b \operatorname{atanh}(c + dx)}{e + f\sqrt{x}} dx$$

input `int((a + b*atanh(c + d*x))/(e + f*x^(1/2)),x)`

output `int((a + b*atanh(c + d*x))/(e + f*x^(1/2)), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + f\sqrt{x}} dx = \frac{2\sqrt{x} af + \left(\int \frac{\operatorname{atanh}(dx+c)}{\sqrt{x}f+e} dx \right) b f^2 - 2 \log(\sqrt{x} f + e) ae}{f^2}$$

input `int((a+b*atanh(d*x+c))/(e+f*x^(1/2)),x)`

output `(2*sqrt(x)*a*f + int(atanh(c + d*x)/(sqrt(x)*f + e),x)*b*f**2 - 2*log(sqrt(x)*f + e)*a*e)/f**2`

3.61 $\int \frac{\operatorname{arctanh}(a+bx)}{c+d\sqrt{x}} dx$

| | |
|--------------------------------------|-----|
| Optimal result | 539 |
| Mathematica [A] (verified) | 540 |
| Rubi [A] (verified) | 540 |
| Maple [A] (verified) | 543 |
| Fricas [F] | 544 |
| Sympy [F(-1)] | 544 |
| Maxima [F] | 544 |
| Giac [F] | 545 |
| Mupad [F(-1)] | 545 |
| Reduce [F] | 545 |

Optimal result

Integrand size = 18, antiderivative size = 585

$$\begin{aligned}
 \int \frac{\operatorname{arctanh}(a + bx)}{c + d\sqrt{x}} dx &= \frac{2\sqrt{1+a} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{1+a}}\right)}{\sqrt{bd}} - \frac{2\sqrt{1-a} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{1-a}}\right)}{\sqrt{bd}} \\
 &+ \frac{c \log\left(\frac{d(\sqrt{-1-a}-\sqrt{b}\sqrt{x})}{\sqrt{bc}+\sqrt{-1-ad}}\right) \log(c + d\sqrt{x})}{d^2} \\
 &- \frac{c \log\left(\frac{d(\sqrt{1-a}-\sqrt{b}\sqrt{x})}{\sqrt{bc}+\sqrt{1-ad}}\right) \log(c + d\sqrt{x})}{d^2} \\
 &+ \frac{c \log\left(-\frac{d(\sqrt{-1-a}+\sqrt{b}\sqrt{x})}{\sqrt{bc}-\sqrt{-1-ad}}\right) \log(c + d\sqrt{x})}{d^2} \\
 &- \frac{c \log\left(-\frac{d(\sqrt{1-a}+\sqrt{b}\sqrt{x})}{\sqrt{bc}-\sqrt{1-ad}}\right) \log(c + d\sqrt{x})}{d^2} \\
 &- \frac{\sqrt{x} \log(1-a-bx)}{d} + \frac{c \log(c + d\sqrt{x}) \log(1-a-bx)}{d^2} \\
 &+ \frac{\sqrt{x} \log(1+a+bx)}{d} - \frac{c \log(c + d\sqrt{x}) \log(1+a+bx)}{d^2} \\
 &+ \frac{c \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}-\sqrt{-1-ad}}\right)}{d^2} + \frac{c \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}+\sqrt{-1-ad}}\right)}{d^2} \\
 &- \frac{c \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}-\sqrt{1-ad}}\right)}{d^2} - \frac{c \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}+\sqrt{1-ad}}\right)}{d^2}
 \end{aligned}$$

output

```

2*(1+a)^(1/2)*arctan(b^(1/2)*x^(1/2)/(1+a)^(1/2))/b^(1/2)/d-2*(1-a)^(1/2)*
arctanh(b^(1/2)*x^(1/2)/(1-a)^(1/2))/b^(1/2)/d+c*ln(d*((-1-a)^(1/2)-b^(1/2)
)*x^(1/2))/(b^(1/2)*c+(-1-a)^(1/2)*d)*ln(c+d*x^(1/2))/d^2-c*ln(d*((1-a)^(
1/2)-b^(1/2)*x^(1/2))/(b^(1/2)*c+(1-a)^(1/2)*d))*ln(c+d*x^(1/2))/d^2+c*ln(
-d*((-1-a)^(1/2)+b^(1/2)*x^(1/2))/(b^(1/2)*c-(-1-a)^(1/2)*d))*ln(c+d*x^(1/
2))/d^2-c*ln(-d*((1-a)^(1/2)+b^(1/2)*x^(1/2))/(b^(1/2)*c-(1-a)^(1/2)*d))*l
n(c+d*x^(1/2))/d^2-x^(1/2)*ln(-b*x-a+1)/d+c*ln(c+d*x^(1/2))*ln(-b*x-a+1)/d
^2+x^(1/2)*ln(b*x+a+1)/d-c*ln(c+d*x^(1/2))*ln(b*x+a+1)/d^2+c*polylog(2,b^(
1/2)*(c+d*x^(1/2))/(b^(1/2)*c-(-1-a)^(1/2)*d))/d^2+c*polylog(2,b^(1/2)*(c+
d*x^(1/2))/(b^(1/2)*c+(-1-a)^(1/2)*d))/d^2-c*polylog(2,b^(1/2)*(c+d*x^(1/2)
))/(b^(1/2)*c-(1-a)^(1/2)*d))/d^2-c*polylog(2,b^(1/2)*(c+d*x^(1/2))/(b^(1/
2)*c+(1-a)^(1/2)*d))/d^2

```

Mathematica [A] (verified)

Time = 0.34 (sec) , antiderivative size = 549, normalized size of antiderivative = 0.94

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + d\sqrt{x}} dx$$

$$= \frac{2\sqrt{1+ad} \operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{1+a}}\right)}{\sqrt{b}} - \frac{2\sqrt{1-ad} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{1-a}}\right)}{\sqrt{b}} + c \log\left(\frac{d(\sqrt{-1-a}-\sqrt{b}\sqrt{x})}{\sqrt{bc}+\sqrt{-1-ad}}\right) \log(c + d\sqrt{x}) - c \log\left(\frac{d(\sqrt{1-a}-\sqrt{b}\sqrt{x})}{\sqrt{bc}+\sqrt{1-ad}}\right)$$

input `Integrate[ArcTanh[a + b*x]/(c + d*Sqrt[x]), x]`

output

```
((2*Sqrt[1 + a]*d*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[1 + a]])/Sqrt[b] - (2*Sqrt[1 - a]*d*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[1 - a]])/Sqrt[b] + c*Log[(d*(Sqrt[-1 - a] - Sqrt[b]*Sqrt[x]))/(Sqrt[b]*c + Sqrt[-1 - a]*d)]*Log[c + d*Sqrt[x]] - c*Log[(d*(Sqrt[1 - a] - Sqrt[b]*Sqrt[x]))/(Sqrt[b]*c + Sqrt[1 - a]*d)]*Log[c + d*Sqrt[x]] + c*Log[(d*(Sqrt[-1 - a] + Sqrt[b]*Sqrt[x]))/(-(Sqrt[b]*c) + Sqrt[-1 - a]*d)]*Log[c + d*Sqrt[x]] - c*Log[(d*(Sqrt[1 - a] + Sqrt[b]*Sqrt[x]))/(-(Sqrt[b]*c) + Sqrt[1 - a]*d)]*Log[c + d*Sqrt[x]] - d*Sqrt[x]*Log[1 - a - b*x] + c*Log[c + d*Sqrt[x]]*Log[1 - a - b*x] + d*Sqrt[x]*Log[1 + a + b*x] - c*Log[c + d*Sqrt[x]]*Log[1 + a + b*x] + c*PolyLog[2, (Sqrt[b]*(c + d*Sqrt[x]))/(Sqrt[b]*c - Sqrt[-1 - a]*d)] + c*PolyLog[2, (Sqrt[b]*(c + d*Sqrt[x]))/(Sqrt[b]*c + Sqrt[-1 - a]*d)] - c*PolyLog[2, (Sqrt[b]*(c + d*Sqrt[x]))/(Sqrt[b]*c - Sqrt[1 - a]*d)] - c*PolyLog[2, (Sqrt[b]*(c + d*Sqrt[x]))/(Sqrt[b]*c + Sqrt[1 - a]*d)]/d^2
```

Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 585, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {6665, 2855, 2916, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + d\sqrt{x}} dx$$

$$\begin{aligned}
& \downarrow \text{6665} \\
& \frac{1}{2} \int \frac{\log(a + bx + 1)}{c + d\sqrt{x}} dx - \frac{1}{2} \int \frac{\log(-a - bx + 1)}{c + d\sqrt{x}} dx \\
& \downarrow \text{2855} \\
& \int \frac{\sqrt{x} \log(a + bx + 1)}{c + d\sqrt{x}} d\sqrt{x} - \int \frac{\sqrt{x} \log(-a - bx + 1)}{c + d\sqrt{x}} d\sqrt{x} \\
& \downarrow \text{2916} \\
& \int \left(\frac{\log(a + bx + 1)}{d} - \frac{c \log(a + bx + 1)}{d(c + d\sqrt{x})} \right) d\sqrt{x} - \\
& \int \left(\frac{\log(-a - bx + 1)}{d} - \frac{c \log(-a - bx + 1)}{d(c + d\sqrt{x})} \right) d\sqrt{x} \\
& \downarrow \text{2009} \\
& \frac{2\sqrt{a+1} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a+1}}\right)}{\sqrt{bd}} - \frac{2\sqrt{1-a} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{1-a}}\right)}{\sqrt{bd}} + \frac{c \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}-\sqrt{-a-1d}}\right)}{d^2} + \\
& \frac{c \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}+\sqrt{-a-1d}}\right)}{d^2} - \frac{c \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}-\sqrt{1-ad}}\right)}{d^2} - \frac{c \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}+\sqrt{1-ad}}\right)}{d^2} + \\
& \frac{c \log(c + d\sqrt{x}) \log\left(\frac{d(\sqrt{-a-1}-\sqrt{b}\sqrt{x})}{\sqrt{-a-1d}+\sqrt{bc}}\right)}{d^2} - \frac{c \log(c + d\sqrt{x}) \log\left(\frac{d(\sqrt{1-a}-\sqrt{b}\sqrt{x})}{\sqrt{1-ad}+\sqrt{bc}}\right)}{d^2} + \\
& \frac{c \log(c + d\sqrt{x}) \log\left(-\frac{d(\sqrt{-a-1}+\sqrt{b}\sqrt{x})}{\sqrt{bc}-\sqrt{-a-1d}}\right)}{d^2} - \frac{c \log(c + d\sqrt{x}) \log\left(-\frac{d(\sqrt{1-a}+\sqrt{b}\sqrt{x})}{\sqrt{bc}-\sqrt{1-ad}}\right)}{d^2} + \\
& \frac{c \log(-a - bx + 1) \log(c + d\sqrt{x})}{d^2} - \frac{c \log(a + bx + 1) \log(c + d\sqrt{x})}{d^2} - \frac{\sqrt{x} \log(-a - bx + 1)}{d} + \\
& \frac{\sqrt{x} \log(a + bx + 1)}{d}
\end{aligned}$$

input

```
Int[ArcTanh[a + b*x]/(c + d*Sqrt[x]), x]
```

output

$$\begin{aligned}
& (2\sqrt{1+a}\operatorname{ArcTan}[\sqrt{b}\sqrt{x}]/\sqrt{1+a}]/(\sqrt{b}d) - (2\sqrt{1-a}\operatorname{ArcTanh}[\sqrt{b}\sqrt{x}]/\sqrt{1-a}]/(\sqrt{b}d) + (c\operatorname{Log}[(d(\sqrt{-1-a} - \sqrt{b}\sqrt{x}))/(\sqrt{b}c + \sqrt{-1-a}d)]\operatorname{Log}[c + d\sqrt{x}])/d^2 - (c\operatorname{Log}[(d(\sqrt{1-a} - \sqrt{b}\sqrt{x}))/(\sqrt{b}c + \sqrt{1-a}d)]\operatorname{Log}[c + d\sqrt{x}])/d^2 + (c\operatorname{Log}[-(d(\sqrt{-1-a} + \sqrt{b}\sqrt{x}))/(\sqrt{b}c - \sqrt{-1-a}d)]\operatorname{Log}[c + d\sqrt{x}])/d^2 - (c\operatorname{Log}[-(d(\sqrt{1-a} + \sqrt{b}\sqrt{x}))/(\sqrt{b}c - \sqrt{1-a}d)]\operatorname{Log}[c + d\sqrt{x}])/d^2 - (\sqrt{x}\operatorname{Log}[1-a-bx])/d + (c\operatorname{Log}[c + d\sqrt{x}]\operatorname{Log}[1-a-bx])/d^2 + (\sqrt{x}\operatorname{Log}[1+a+bx])/d - (c\operatorname{Log}[c + d\sqrt{x}]\operatorname{Log}[1+a+bx])/d^2 + (c\operatorname{PolyLog}[2, (\sqrt{b}(c + d\sqrt{x}))/(\sqrt{b}c - \sqrt{-1-a}d)])/d^2 + (c\operatorname{PolyLog}[2, (\sqrt{b}(c + d\sqrt{x}))/(\sqrt{b}c + \sqrt{-1-a}d)])/d^2 - (c\operatorname{PolyLog}[2, (\sqrt{b}(c + d\sqrt{x}))/(\sqrt{b}c - \sqrt{1-a}d)])/d^2 - (c\operatorname{PolyLog}[2, (\sqrt{b}(c + d\sqrt{x}))/(\sqrt{b}c + \sqrt{1-a}d)])/d^2
\end{aligned}$$

Defintions of rubi rules used

rule 2009

$$\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] \;/; \operatorname{SumQ}[u]$$

rule 2855

$$\operatorname{Int}[(a_.) + \operatorname{Log}[(c_.) * ((d_.) + (e_.) * (x_.)^{(n_.)}) * (b_.)^{(p_.)} * ((f_.) + (g_.) * (x_.)^{(r_.)})^{(q_.)}], x_Symbol] \rightarrow \operatorname{With}[\{k = \operatorname{Denominator}[r]\}, \operatorname{Simp}[k \operatorname{Subst}[\operatorname{Int}[x^{(k-1)} * (f + g * x^{(k*r)})^q * (a + b * \operatorname{Log}[c * (d + e * x^k)^n])^p, x], x, x^{(1/k)}], x]] \;/; \operatorname{FreeQ}\{a, b, c, d, e, f, g, n, p, q\}, x] \ \&\& \operatorname{FractionQ}[r] \ \&\& \operatorname{IntegerQ}[p, 0]$$

rule 2916

$$\operatorname{Int}[(a_.) + \operatorname{Log}[(c_.) * ((d_.) + (e_.) * (x_.)^{(n_.)})^{(p_.)} * (b_.)^{(q_.)} * (x_.)^{(m_.)} * ((f_.) + (g_.) * (x_.)^{(r_.)})], x_Symbol] \rightarrow \operatorname{Int}[\operatorname{ExpandIntegrand}[(a + b * \operatorname{Log}[c * (d + e * x^n)^p])^q, x^m * (f + g * x)^r, x], x] \;/; \operatorname{FreeQ}\{a, b, c, d, e, f, g, n, p, q\}, x] \ \&\& \operatorname{IntegerQ}[m] \ \&\& \operatorname{IntegerQ}[r]$$

rule 6665

$$\operatorname{Int}[\operatorname{ArcTanh}[(c_.) + (d_.) * (x_.)]/((e_.) + (f_.) * (x_.)^{(n_.)}), x_Symbol] \rightarrow \operatorname{Simp}[1/2 \operatorname{Int}[\operatorname{Log}[1 + c + d * x]/(e + f * x^n), x], x] - \operatorname{Simp}[1/2 \operatorname{Int}[\operatorname{Log}[1 - c - d * x]/(e + f * x^n), x], x] \;/; \operatorname{FreeQ}\{c, d, e, f\}, x] \ \&\& \operatorname{RationalQ}[n]$$

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 648, normalized size of antiderivative = 1.11

| method | result |
|-------------------|---|
| derivativedivides | $\frac{2 \operatorname{arctanh}(bx+a)\sqrt{x}}{d} - \frac{2 \operatorname{arctanh}(bx+a)c \ln(c+d\sqrt{x})}{d^2} - \frac{4b \left(-d^2 \left(\frac{(1+a) \operatorname{arctan}\left(\frac{-2bc+2b(c+d\sqrt{x})}{2\sqrt{ab}d^2+b d^2}\right)}{2b\sqrt{ab}d^2+b d^2} \right) + \frac{(1-a) \operatorname{arctan}\left(\frac{-2bc+2b(c+d\sqrt{x})}{2\sqrt{ab}d^2+b d^2}\right)}{2b\sqrt{ab}d^2+b d^2} \right)}{d^2}$ |
| default | $\frac{2 \operatorname{arctanh}(bx+a)\sqrt{x}}{d} - \frac{2 \operatorname{arctanh}(bx+a)c \ln(c+d\sqrt{x})}{d^2} - \frac{4b \left(-d^2 \left(\frac{(1+a) \operatorname{arctan}\left(\frac{-2bc+2b(c+d\sqrt{x})}{2\sqrt{ab}d^2+b d^2}\right)}{2b\sqrt{ab}d^2+b d^2} \right) + \frac{(1-a) \operatorname{arctan}\left(\frac{-2bc+2b(c+d\sqrt{x})}{2\sqrt{ab}d^2+b d^2}\right)}{2b\sqrt{ab}d^2+b d^2} \right)}{d^2}$ |

input

```
int(arctanh(b*x+a)/(c+d*x^(1/2)),x,method=_RETURNVERBOSE)
```

output

```
2*arctanh(b*x+a)/d*x^(1/2)-2*arctanh(b*x+a)*c/d^2*ln(c+d*x^(1/2))-4*b/d^2*
(-d^2*(1/2*(1+a)/b/(a*b*d^2+b*d^2)^(1/2)*arctan(1/2*(-2*b*c+2*b*(c+d*x^(1/2)))/
(a*b*d^2+b*d^2)^(1/2))+1/2*(1-a)/b/(a*b*d^2-b*d^2)^(1/2)*arctan(1/2*(
-2*b*c+2*b*(c+d*x^(1/2)))/
(a*b*d^2-b*d^2)^(1/2)))-c*d^2*(1/2/d^2*(1/2*ln(c
+d*x^(1/2))*(ln((b*c-b*(c+d*x^(1/2))+(-a*b*d^2-b*d^2)^(1/2))/(b*c+(-a*b*d^
2-b*d^2)^(1/2)))+ln((-b*c+b*(c+d*x^(1/2))+(-a*b*d^2-b*d^2)^(1/2))/(-b*c+(-
a*b*d^2-b*d^2)^(1/2))))/b+1/2*(dilog((b*c-b*(c+d*x^(1/2))+(-a*b*d^2-b*d^2)
^(1/2))/(b*c+(-a*b*d^2-b*d^2)^(1/2)))+dilog((-b*c+b*(c+d*x^(1/2))+(-a*b*d^
2-b*d^2)^(1/2))/(-b*c+(-a*b*d^2-b*d^2)^(1/2))))/b)+1/2/d^2*(-1/2*ln(c+d*x^
(1/2))*(ln((b*c-b*(c+d*x^(1/2))+(-a*b*d^2+b*d^2)^(1/2))/(b*c+(-a*b*d^2+b*d
^2)^(1/2)))+ln((-b*c+b*(c+d*x^(1/2))+(-a*b*d^2+b*d^2)^(1/2))/(-b*c+(-a*b*d
^2+b*d^2)^(1/2))))/b-1/2*(dilog((b*c-b*(c+d*x^(1/2))+(-a*b*d^2+b*d^2)^(1/2)
))/(b*c+(-a*b*d^2+b*d^2)^(1/2)))+dilog((-b*c+b*(c+d*x^(1/2))+(-a*b*d^2+b*d
^2)^(1/2))/(-b*c+(-a*b*d^2+b*d^2)^(1/2))))/b))
```


Fricas [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + d\sqrt{x}} dx = \int \frac{\operatorname{artanh}(bx + a)}{d\sqrt{x} + c} dx$$

input `integrate(arctanh(b*x+a)/(c+d*x^(1/2)),x, algorithm="fricas")`

output `integral((d*sqrt(x)*arctanh(b*x + a) - c*arctanh(b*x + a))/(d^2*x - c^2), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + d\sqrt{x}} dx = \text{Timed out}$$

input `integrate(atanh(b*x+a)/(c+d*x**(1/2)),x)`

output `Timed out`

Maxima [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + d\sqrt{x}} dx = \int \frac{\operatorname{artanh}(bx + a)}{d\sqrt{x} + c} dx$$

input `integrate(arctanh(b*x+a)/(c+d*x^(1/2)),x, algorithm="maxima")`

output `integrate(arctanh(b*x + a)/(d*sqrt(x) + c), x)`

Giac [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + d\sqrt{x}} dx = \int \frac{\operatorname{artanh}(bx + a)}{d\sqrt{x} + c} dx$$

input `integrate(arctanh(b*x+a)/(c+d*x^(1/2)),x, algorithm="giac")`

output `integrate(arctanh(b*x + a)/(d*sqrt(x) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + d\sqrt{x}} dx = \int \frac{\operatorname{atanh}(a + bx)}{c + d\sqrt{x}} dx$$

input `int(atanh(a + b*x)/(c + d*x^(1/2)),x)`

output `int(atanh(a + b*x)/(c + d*x^(1/2)), x)`

Reduce [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + d\sqrt{x}} dx = \int \frac{\operatorname{atanh}(bx + a)}{\sqrt{x}d + c} dx$$

input `int(atanh(b*x+a)/(c+d*x^(1/2)),x)`

output `int(atanh(a + b*x)/(sqrt(x)*d + c),x)`

$$3.62 \quad \int \frac{\operatorname{arctanh}(a+bx)}{c+\frac{d}{\sqrt{x}}} dx$$

| | |
|----------------------------|-----|
| Optimal result | 547 |
| Mathematica [A] (verified) | 549 |
| Rubi [A] (verified) | 550 |
| Maple [A] (verified) | 553 |
| Fricas [F] | 554 |
| Sympy [F(-1)] | 554 |
| Maxima [F] | 555 |
| Giac [F] | 555 |
| Mupad [F(-1)] | 555 |
| Reduce [F] | 556 |

Optimal result

Integrand size = 18, antiderivative size = 661

$$\begin{aligned}
\int \frac{\operatorname{arctanh}(a+bx)}{c+\frac{d}{\sqrt{x}}} dx = & -\frac{2\sqrt{1+ad} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{1+a}}\right)}{\sqrt{bc^2}} + \frac{2\sqrt{1-ad} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{1-a}}\right)}{\sqrt{bc^2}} \\
& - \frac{d^2 \log\left(\frac{c(\sqrt{-1-a}-\sqrt{b}\sqrt{x})}{\sqrt{-1-ac}+\sqrt{bd}}\right) \log(d+c\sqrt{x})}{c^3} \\
& + \frac{d^2 \log\left(\frac{c(\sqrt{1-a}-\sqrt{b}\sqrt{x})}{\sqrt{1-ac}+\sqrt{bd}}\right) \log(d+c\sqrt{x})}{c^3} \\
& - \frac{d^2 \log\left(\frac{c(\sqrt{-1-a}+\sqrt{b}\sqrt{x})}{\sqrt{-1-ac}-\sqrt{bd}}\right) \log(d+c\sqrt{x})}{c^3} \\
& + \frac{d^2 \log\left(\frac{c(\sqrt{1-a}+\sqrt{b}\sqrt{x})}{\sqrt{1-ac}-\sqrt{bd}}\right) \log(d+c\sqrt{x})}{c^3} \\
& + \frac{d\sqrt{x} \log(1-a-bx)}{c^2} + \frac{(1-a-bx) \log(1-a-bx)}{2bc} \\
& - \frac{d^2 \log(d+c\sqrt{x}) \log(1-a-bx)}{c^3} \\
& - \frac{d\sqrt{x} \log(1+a+bx)}{c^2} + \frac{(1+a+bx) \log(1+a+bx)}{2bc} \\
& + \frac{d^2 \log(d+c\sqrt{x}) \log(1+a+bx)}{c^3} \\
& - \frac{d^2 \operatorname{PolyLog}\left(2, -\frac{\sqrt{b}(d+c\sqrt{x})}{\sqrt{-1-ac}-\sqrt{bd}}\right)}{c^3} \\
& + \frac{d^2 \operatorname{PolyLog}\left(2, -\frac{\sqrt{b}(d+c\sqrt{x})}{\sqrt{1-ac}-\sqrt{bd}}\right)}{c^3} \\
& - \frac{d^2 \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(d+c\sqrt{x})}{\sqrt{-1-ac}+\sqrt{bd}}\right)}{c^3} + \frac{d^2 \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(d+c\sqrt{x})}{\sqrt{1-ac}+\sqrt{bd}}\right)}{c^3}
\end{aligned}$$

output

```

-2*(1+a)^(1/2)*d*arctan(b^(1/2)*x^(1/2)/(1+a)^(1/2))/b^(1/2)/c^2+2*(1-a)^(
1/2)*d*arctanh(b^(1/2)*x^(1/2)/(1-a)^(1/2))/b^(1/2)/c^2-d^2*ln(c*((-1-a)^(
1/2)-b^(1/2)*x^(1/2))/((-1-a)^(1/2)*c+b^(1/2)*d))*ln(d+c*x^(1/2))/c^3+d^2*
ln(c*((-1-a)^(1/2)-b^(1/2)*x^(1/2))/((-1-a)^(1/2)*c+b^(1/2)*d))*ln(d+c*x^(1/
2))/c^3-d^2*ln(c*((-1-a)^(1/2)+b^(1/2)*x^(1/2))/((-1-a)^(1/2)*c-b^(1/2)*d)
)*ln(d+c*x^(1/2))/c^3+d^2*ln(c*((-1-a)^(1/2)+b^(1/2)*x^(1/2))/((-1-a)^(1/2)*
c-b^(1/2)*d))*ln(d+c*x^(1/2))/c^3+d*x^(1/2)*ln(-b*x-a+1)/c^2+1/2*(-b*x-a+1
)*ln(-b*x-a+1)/b/c-d^2*ln(d+c*x^(1/2))*ln(-b*x-a+1)/c^3-d*x^(1/2)*ln(b*x+a
+1)/c^2+1/2*(b*x+a+1)*ln(b*x+a+1)/b/c+d^2*ln(d+c*x^(1/2))*ln(b*x+a+1)/c^3-
d^2*polylog(2,-b^(1/2)*(d+c*x^(1/2))/((-1-a)^(1/2)*c-b^(1/2)*d))/c^3+d^2*p
olylog(2,-b^(1/2)*(d+c*x^(1/2))/((-1-a)^(1/2)*c-b^(1/2)*d))/c^3-d^2*polylog
(2,b^(1/2)*(d+c*x^(1/2))/((-1-a)^(1/2)*c+b^(1/2)*d))/c^3+d^2*polylog(2,b^(
1/2)*(d+c*x^(1/2))/((-1-a)^(1/2)*c+b^(1/2)*d))/c^3

```

Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 668, normalized size of antiderivative = 1.01

$$\begin{aligned}
\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{\sqrt{x}}} dx = & -\frac{2\sqrt{1+ad} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{1+a}}\right)}{\sqrt{bc^2}} + \frac{2\sqrt{1-ad} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{1-a}}\right)}{\sqrt{bc^2}} \\
& - \frac{d^2 \log\left(\frac{c(\sqrt{-1-a}-\sqrt{b}\sqrt{x})}{\sqrt{-1-ac}+\sqrt{bd}}\right) \log(d + c\sqrt{x})}{c^3} \\
& + \frac{d^2 \log\left(\frac{c(\sqrt{1-a}-\sqrt{b}\sqrt{x})}{\sqrt{1-ac}+\sqrt{bd}}\right) \log(d + c\sqrt{x})}{c^3} \\
& - \frac{d^2 \log\left(\frac{c(\sqrt{-1-a}+\sqrt{b}\sqrt{x})}{\sqrt{-1-ac}-\sqrt{bd}}\right) \log(d + c\sqrt{x})}{c^3} \\
& + \frac{d^2 \log\left(\frac{c(\sqrt{1-a}+\sqrt{b}\sqrt{x})}{\sqrt{1-ac}-\sqrt{bd}}\right) \log(d + c\sqrt{x})}{c^3} \\
& + \frac{d\sqrt{x} \log(1-a-bx)}{c^2} - \frac{d^2 \log(d + c\sqrt{x}) \log(1-a-bx)}{c^3} \\
& + \frac{x + \frac{(1-a-bx) \log(1-a-bx)}{b}}{2c} - \frac{d\sqrt{x} \log(1+a+bx)}{c^2} \\
& + \frac{d^2 \log(d + c\sqrt{x}) \log(1+a+bx)}{c^3} \\
& - \frac{x - \frac{(1+a+bx) \log(1+a+bx)}{b}}{2c} - \frac{d^2 \operatorname{PolyLog}\left(2, -\frac{\sqrt{b}(d+c\sqrt{x})}{\sqrt{-1-ac}-\sqrt{bd}}\right)}{c^3} \\
& + \frac{d^2 \operatorname{PolyLog}\left(2, -\frac{\sqrt{b}(d+c\sqrt{x})}{\sqrt{1-ac}-\sqrt{bd}}\right)}{c^3} \\
& - \frac{d^2 \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(d+c\sqrt{x})}{\sqrt{-1-ac}+\sqrt{bd}}\right)}{c^3} + \frac{d^2 \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(d+c\sqrt{x})}{\sqrt{1-ac}+\sqrt{bd}}\right)}{c^3}
\end{aligned}$$

input

```
Integrate[ArcTanh[a + b*x]/(c + d/Sqrt[x]), x]
```

output

```
(-2*Sqrt[1 + a]*d*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[1 + a]]/(Sqrt[b]*c^2) + (
2*Sqrt[1 - a]*d*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[1 - a]]/(Sqrt[b]*c^2) - (d
^2*Log[(c*(Sqrt[-1 - a] - Sqrt[b]*Sqrt[x]))/(Sqrt[-1 - a]*c + Sqrt[b]*d)]*
Log[d + c*Sqrt[x]])/c^3 + (d^2*Log[(c*(Sqrt[1 - a] - Sqrt[b]*Sqrt[x]))/(Sq
rt[1 - a]*c + Sqrt[b]*d)]*Log[d + c*Sqrt[x]])/c^3 - (d^2*Log[(c*(Sqrt[-1 -
a] + Sqrt[b]*Sqrt[x]))/(Sqrt[-1 - a]*c - Sqrt[b]*d)]*Log[d + c*Sqrt[x]])/
c^3 + (d^2*Log[(c*(Sqrt[1 - a] + Sqrt[b]*Sqrt[x]))/(Sqrt[1 - a]*c - Sqrt[b
]*d)]*Log[d + c*Sqrt[x]])/c^3 + (d*Sqrt[x]*Log[1 - a - b*x])/c^2 - (d^2*Lo
g[d + c*Sqrt[x]*Log[1 - a - b*x])/c^3 + (x + ((1 - a - b*x)*Log[1 - a - b
*x])/b)/(2*c) - (d*Sqrt[x]*Log[1 + a + b*x])/c^2 + (d^2*Log[d + c*Sqrt[x]]
*Log[1 + a + b*x])/c^3 - (x - ((1 + a + b*x)*Log[1 + a + b*x])/b)/(2*c) -
(d^2*PolyLog[2, -((Sqrt[b]*(d + c*Sqrt[x]))/(Sqrt[-1 - a]*c - Sqrt[b]*d))
]/c^3 + (d^2*PolyLog[2, -((Sqrt[b]*(d + c*Sqrt[x]))/(Sqrt[1 - a]*c - Sqrt[
b]*d))])/c^3 - (d^2*PolyLog[2, (Sqrt[b]*(d + c*Sqrt[x]))/(Sqrt[-1 - a]*c +
Sqrt[b]*d))]/c^3 + (d^2*PolyLog[2, (Sqrt[b]*(d + c*Sqrt[x]))/(Sqrt[1 - a]
*c + Sqrt[b]*d))])/c^3
```

Rubi [A] (verified)

Time = 1.38 (sec) , antiderivative size = 661, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {6665, 2855, 2005, 2916, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{\sqrt{x}}} dx$$

$$\downarrow \text{6665}$$

$$\frac{1}{2} \int \frac{\log(a + bx + 1)}{c + \frac{d}{\sqrt{x}}} dx - \frac{1}{2} \int \frac{\log(-a - bx + 1)}{c + \frac{d}{\sqrt{x}}} dx$$

$$\downarrow \text{2855}$$

$$\int \frac{\sqrt{x} \log(a + bx + 1)}{c + \frac{d}{\sqrt{x}}} d\sqrt{x} - \int \frac{\sqrt{x} \log(-a - bx + 1)}{c + \frac{d}{\sqrt{x}}} d\sqrt{x}$$

$$\downarrow \text{2005}$$

$$\begin{aligned}
& \int \frac{x \log(a + bx + 1)}{\sqrt{xc + d}} d\sqrt{x} - \int \frac{x \log(-a - bx + 1)}{\sqrt{xc + d}} d\sqrt{x} \\
& \quad \downarrow \text{2916} \\
& \int \left(\frac{\log(a + bx + 1)d^2}{c^2 (\sqrt{xc + d})} - \frac{\log(a + bx + 1)d}{c^2} + \frac{\sqrt{x} \log(a + bx + 1)}{c} \right) d\sqrt{x} - \\
& \int \left(\frac{\log(-a - bx + 1)d^2}{c^2 (\sqrt{xc + d})} - \frac{\log(-a - bx + 1)d}{c^2} + \frac{\sqrt{x} \log(-a - bx + 1)}{c} \right) d\sqrt{x} \\
& \quad \downarrow \text{2009} \\
& -\frac{2\sqrt{a+1}d \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a+1}}\right)}{\sqrt{bc^2}} + \frac{2\sqrt{1-a}d \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{1-a}}\right)}{\sqrt{bc^2}} - \frac{d^2 \operatorname{PolyLog}\left(2, -\frac{\sqrt{b}(\sqrt{xc+d})}{\sqrt{-a-1c-\sqrt{bd}}}\right)}{c^3} + \\
& \frac{d^2 \operatorname{PolyLog}\left(2, -\frac{\sqrt{b}(\sqrt{xc+d})}{\sqrt{1-ac-\sqrt{bd}}}\right)}{c^3} - \frac{d^2 \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(\sqrt{xc+d})}{\sqrt{-a-1c+\sqrt{bd}}}\right)}{c^3} + \\
& \frac{d^2 \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(\sqrt{xc+d})}{\sqrt{1-ac+\sqrt{bd}}}\right)}{c^3} - \frac{d^2 \log(c\sqrt{x} + d) \log\left(\frac{c(\sqrt{-a-1}-\sqrt{b}\sqrt{x})}{\sqrt{-a-1c+\sqrt{bd}}}\right)}{c^3} + \\
& \frac{d^2 \log(c\sqrt{x} + d) \log\left(\frac{c(\sqrt{1-a}-\sqrt{b}\sqrt{x})}{\sqrt{1-ac+\sqrt{bd}}}\right)}{c^3} - \frac{d^2 \log(c\sqrt{x} + d) \log\left(\frac{c(\sqrt{-a-1}+\sqrt{b}\sqrt{x})}{\sqrt{-a-1c-\sqrt{bd}}}\right)}{c^3} + \\
& \frac{d^2 \log(c\sqrt{x} + d) \log\left(\frac{c(\sqrt{1-a}+\sqrt{b}\sqrt{x})}{\sqrt{1-ac-\sqrt{bd}}}\right)}{c^3} - \frac{d^2 \log(-a - bx + 1) \log(c\sqrt{x} + d)}{c^3} + \\
& \frac{d^2 \log(a + bx + 1) \log(c\sqrt{x} + d)}{c^3} + \frac{d\sqrt{x} \log(-a - bx + 1)}{c^2} - \frac{d\sqrt{x} \log(a + bx + 1)}{c^2} + \\
& \frac{(-a - bx + 1) \log(-a - bx + 1)}{2bc} + \frac{(a + bx + 1) \log(a + bx + 1)}{2bc}
\end{aligned}$$

input `Int[ArcTanh[a + b*x]/(c + d/Sqrt[x]),x]`

output

$$\begin{aligned}
& (-2\sqrt{1+a} \operatorname{ArcTan}[\sqrt{b}\sqrt{x}]/\sqrt{1+a}]/(\sqrt{b}c^2) + (2\sqrt{1-a} \operatorname{ArcTanh}[\sqrt{b}\sqrt{x}]/\sqrt{1-a}]/(\sqrt{b}c^2) - (d^2 \operatorname{Log}[(c(\sqrt{-1-a} - \sqrt{b}\sqrt{x}))/(\sqrt{-1-a}c + \sqrt{b}d)] * \operatorname{Log}[d + c\sqrt{x}])/c^3 + (d^2 \operatorname{Log}[(c(\sqrt{1-a} - \sqrt{b}\sqrt{x}))/(\sqrt{1-a}c + \sqrt{b}d)] * \operatorname{Log}[d + c\sqrt{x}])/c^3 - (d^2 \operatorname{Log}[(c(\sqrt{-1-a} + \sqrt{b}\sqrt{x}))/(\sqrt{-1-a}c - \sqrt{b}d)] * \operatorname{Log}[d + c\sqrt{x}])/c^3 + (d^2 \operatorname{Log}[(c(\sqrt{1-a} + \sqrt{b}\sqrt{x}))/(\sqrt{1-a}c - \sqrt{b}d)] * \operatorname{Log}[d + c\sqrt{x}])/c^3 + (d\sqrt{x} \operatorname{Log}[1-a-bx])/c^2 + ((1-a-bx) \operatorname{Log}[1-a-bx])/(2bc) - (d^2 \operatorname{Log}[d + c\sqrt{x}] \operatorname{Log}[1-a-bx])/c^3 - (d\sqrt{x} \operatorname{Log}[1+a+bx])/c^2 + ((1+a+bx) \operatorname{Log}[1+a+bx])/(2bc) + (d^2 \operatorname{Log}[d + c\sqrt{x}] \operatorname{Log}[1+a+bx])/c^3 - (d^2 \operatorname{PolyLog}[2, -((\sqrt{b}(d + c\sqrt{x}))/(\sqrt{-1-a}c - \sqrt{b}d))])/c^3 + (d^2 \operatorname{PolyLog}[2, -((\sqrt{b}(d + c\sqrt{x}))/(\sqrt{1-a}c - \sqrt{b}d))])/c^3 - (d^2 \operatorname{PolyLog}[2, (\sqrt{b}(d + c\sqrt{x}))/(\sqrt{-1-a}c + \sqrt{b}d)])/c^3 + (d^2 \operatorname{PolyLog}[2, (\sqrt{b}(d + c\sqrt{x}))/(\sqrt{1-a}c + \sqrt{b}d)])/c^3
\end{aligned}$$

Defintions of rubi rules used

rule 2005

$$\operatorname{Int}[(F x_{-}) (x_{-})^{(m_{-})} ((a_{-}) + (b_{-}) (x_{-})^{(n_{-})})^{(p_{-})}, x_{-} \operatorname{Symbol}] \rightarrow \operatorname{Int}[x^{(m+n p)} (b + a/x^n)^p F x, x] \text{ ; FreeQ}\{a, b, m, n\}, x \text{ \&\& IntegerQ}\{p\} \text{ \&\& NegQ}\{n\}$$

rule 2009

$$\operatorname{Int}[u_{-}, x_{-} \operatorname{Symbol}] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] \text{ ; SumQ}[u]$$

rule 2855

$$\operatorname{Int}[(a_{-}) + \operatorname{Log}[(c_{-}) ((d_{-}) + (e_{-}) (x_{-})^{(n_{-})})] (b_{-})]^{(p_{-})} ((f_{-}) + (g_{-}) (x_{-})^{(r_{-})})^{(q_{-})}, x_{-} \operatorname{Symbol}] \rightarrow \operatorname{With}[\{k = \operatorname{Denominator}[r]\}, \operatorname{Simp}[k \operatorname{Subst}[\operatorname{Int}[x^{(k-1)} (f + g x^{(k r)})^q (a + b \operatorname{Log}[c(d + e x^k)^n])^p, x], x, x^{(1/k)}], x]] \text{ ; FreeQ}\{a, b, c, d, e, f, g, n, p, q\}, x \text{ \&\& FractionQ}\{r\} \text{ \&\& IGtQ}\{p, 0\}$$

rule 2916

$$\operatorname{Int}[(a_{-}) + \operatorname{Log}[(c_{-}) ((d_{-}) + (e_{-}) (x_{-})^{(n_{-})})] (b_{-})]^{(q_{-})} (x_{-})^{(m_{-})} ((f_{-}) + (g_{-}) (x_{-})^{(r_{-})}), x_{-} \operatorname{Symbol}] \rightarrow \operatorname{Int}[\operatorname{ExpandIntegrand}[(a + b \operatorname{Log}[c(d + e x^n)^p])^q, x^m (f + g x)^r, x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, g, n, p, q\}, x \text{ \&\& IntegerQ}\{m\} \text{ \&\& IntegerQ}\{r\}$$

rule 6665

```
Int[ArcTanh[(c_) + (d_.)*(x_)]/((e_) + (f_.)*(x_)^(n_.)), x_Symbol] := Simp
[1/2 Int[Log[1 + c + d*x]/(e + f*x^n), x], x] - Simp[1/2 Int[Log[1 - c
- d*x]/(e + f*x^n), x], x] /; FreeQ[{c, d, e, f}, x] && RationalQ[n]
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 751, normalized size of antiderivative = 1.14

| method | result |
|-------------------|--|
| derivativedivides | $\frac{\operatorname{arctanh}(bx+a)x}{c} - \frac{2 \operatorname{arctanh}(bx+a)d\sqrt{x}}{c^2} + \frac{2 \operatorname{arctanh}(bx+a)d^2 \ln(d+c\sqrt{x})}{c^3} - \frac{4b}{c d^2} \left(\frac{\ln(d+c\sqrt{x}) \left(\ln\left(\frac{bd-b(d+)}{bd}\right) \right)}{\dots} \right)$ |
| default | $\frac{\operatorname{arctanh}(bx+a)x}{c} - \frac{2 \operatorname{arctanh}(bx+a)d\sqrt{x}}{c^2} + \frac{2 \operatorname{arctanh}(bx+a)d^2 \ln(d+c\sqrt{x})}{c^3} - \frac{4b}{c d^2} \left(\frac{\ln(d+c\sqrt{x}) \left(\ln\left(\frac{bd-b(d+)}{bd}\right) \right)}{\dots} \right)$ |

input

```
int(arctanh(b*x+a)/(c+d/x^(1/2)),x,method=_RETURNVERBOSE)
```

output

```

arctanh(b*x+a)*x/c-2*arctanh(b*x+a)/c^2*d*x^(1/2)+2*arctanh(b*x+a)*d^2/c^3
*ln(d+c*x^(1/2))-4*b/c^2*(c*d^2*(1/2/c^2*(1/2*ln(d+c*x^(1/2)))*(ln((b*d-b*(
d+c*x^(1/2))+(-a*b*c^2-b*c^2)^(1/2))/(b*d+(-a*b*c^2-b*c^2)^(1/2)))+ln((-b*
d+b*(d+c*x^(1/2))+(-a*b*c^2-b*c^2)^(1/2))/(-b*d+(-a*b*c^2-b*c^2)^(1/2))))/
b+1/2*(dilog((b*d-b*(d+c*x^(1/2))+(-a*b*c^2-b*c^2)^(1/2))/(b*d+(-a*b*c^2-b
*c^2)^(1/2)))+dilog((-b*d+b*(d+c*x^(1/2))+(-a*b*c^2-b*c^2)^(1/2))/(-b*d+(-
a*b*c^2-b*c^2)^(1/2))))/b)+1/2/c^2*(-1/2*ln(d+c*x^(1/2))*(ln((b*d-b*(d+c*x
^(1/2))+(-a*b*c^2+b*c^2)^(1/2))/(b*d+(-a*b*c^2+b*c^2)^(1/2)))+ln((-b*d+b*(
d+c*x^(1/2))+(-a*b*c^2+b*c^2)^(1/2))/(-b*d+(-a*b*c^2+b*c^2)^(1/2))))/b-1/2
*(dilog((b*d-b*(d+c*x^(1/2))+(-a*b*c^2+b*c^2)^(1/2))/(b*d+(-a*b*c^2+b*c^2)
^(1/2)))+dilog((-b*d+b*(d+c*x^(1/2))+(-a*b*c^2+b*c^2)^(1/2))/(-b*d+(-a*b*c
^2+b*c^2)^(1/2))))/b)+1/2*c*(1/2*(1+a)/b*(-1/2/b*ln(a*c^2+b*d^2-2*b*(d+c*
x^(1/2))*d+b*(d+c*x^(1/2))^2+c^2)+2*d/(a*b*c^2+b*c^2)^(1/2)*arctan(1/2*(-2
*b*d+2*b*(d+c*x^(1/2)))/(a*b*c^2+b*c^2)^(1/2))-1/2*(a-1)/b*(-1/2/b*ln(a*c
^2+b*d^2-2*b*(d+c*x^(1/2))*d+b*(d+c*x^(1/2))^2-c^2)+2*d/(a*b*c^2-b*c^2)^(1
/2)*arctan(1/2*(-2*b*d+2*b*(d+c*x^(1/2)))/(a*b*c^2-b*c^2)^(1/2))))

```

Fricas [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{\sqrt{x}}} dx = \int \frac{\operatorname{arctanh}(bx + a)}{c + \frac{d}{\sqrt{x}}} dx$$

input

```
integrate(arctanh(b*x+a)/(c+d/x^(1/2)),x, algorithm="fricas")
```

output

```
integral((c*x*arctanh(b*x + a) - d*sqrt(x)*arctanh(b*x + a))/(c^2*x - d^2)
, x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{\sqrt{x}}} dx = \text{Timed out}$$

input

```
integrate(atanh(b*x+a)/(c+d/x**(1/2)),x)
```

output Timed out

Maxima [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{\sqrt{x}}} dx = \int \frac{\operatorname{artanh}(bx + a)}{c + \frac{d}{\sqrt{x}}} dx$$

input `integrate(arctanh(b*x+a)/(c+d/x^(1/2)),x, algorithm="maxima")`

output `integrate(arctanh(b*x + a)/(c + d/sqrt(x)), x)`

Giac [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{\sqrt{x}}} dx = \int \frac{\operatorname{artanh}(bx + a)}{c + \frac{d}{\sqrt{x}}} dx$$

input `integrate(arctanh(b*x+a)/(c+d/x^(1/2)),x, algorithm="giac")`

output `integrate(arctanh(b*x + a)/(c + d/sqrt(x)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{\sqrt{x}}} dx = \int \frac{\operatorname{atanh}(a + bx)}{c + \frac{d}{\sqrt{x}}} dx$$

input `int(atanh(a + b*x)/(c + d/x^(1/2)),x)`

output `int(atanh(a + b*x)/(c + d/x^(1/2)), x)`

Reduce [F]

$$\int \frac{\operatorname{arctanh}(a + bx)}{c + \frac{d}{\sqrt{x}}} dx = \int \frac{\sqrt{x} \operatorname{atanh}(bx + a)}{\sqrt{x} c + d} dx$$

input `int(atanh(b*x+a)/(c+d/x^(1/2)),x)`

output `int((sqrt(x)*atanh(a + b*x))/(sqrt(x)*c + d),x)`

3.63 $\int \frac{x^3(a+b\operatorname{arctanh}(c+dx))}{e-fx^2} dx$

| | |
|----------------------------|-----|
| Optimal result | 557 |
| Mathematica [A] (verified) | 558 |
| Rubi [A] (verified) | 559 |
| Maple [A] (verified) | 560 |
| Fricas [F] | 562 |
| Sympy [F(-1)] | 562 |
| Maxima [F] | 562 |
| Giac [F] | 563 |
| Mupad [F(-1)] | 563 |
| Reduce [F] | 563 |

Optimal result

Integrand size = 24, antiderivative size = 381

$$\int \frac{x^3(a+b\operatorname{arctanh}(c+dx))}{e-fx^2} dx = -\frac{bx}{2df} - \frac{x^2(a+b\operatorname{arctanh}(c+dx))}{2f}$$

$$- \frac{b(1-c)^2 \log(1-c-dx)}{4d^2 f}$$

$$+ \frac{e(a+b\operatorname{arctanh}(c+dx)) \log\left(\frac{2}{1+c+dx}\right)}{f^2}$$

$$+ \frac{b(1+c)^2 \log(1+c+dx)}{4d^2 f}$$

$$- \frac{e(a+b\operatorname{arctanh}(c+dx)) \log\left(\frac{2d(\sqrt{e}-\sqrt{fx})}{(d\sqrt{e}-(1-c)\sqrt{f})(1+c+dx)}\right)}{2f^2}$$

$$- \frac{e(a+b\operatorname{arctanh}(c+dx)) \log\left(\frac{2d(\sqrt{e}+\sqrt{fx})}{(d\sqrt{e}+(1-c)\sqrt{f})(1+c+dx)}\right)}{2f^2}$$

$$- \frac{be \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{2f^2}$$

$$+ \frac{be \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e}-\sqrt{fx})}{(d\sqrt{e}-(1-c)\sqrt{f})(1+c+dx)}\right)}{4f^2}$$

$$+ \frac{be \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e}+\sqrt{fx})}{(d\sqrt{e}+(1-c)\sqrt{f})(1+c+dx)}\right)}{4f^2}$$

output

$$\begin{aligned}
& -1/2*b*x/d/f-1/2*x^2*(a+b*\operatorname{arctanh}(d*x+c))/f-1/4*b*(1-c)^2*\ln(-d*x-c+1)/d^2 \\
& /f+e*(a+b*\operatorname{arctanh}(d*x+c))*\ln(2/(d*x+c+1))/f^2+1/4*b*(1+c)^2*\ln(d*x+c+1)/d^2 \\
& /f-1/2*e*(a+b*\operatorname{arctanh}(d*x+c))*\ln(2*d*(e^{1/2}-f^{1/2})*x)/(d*e^{1/2}-(1-c)*f^{1/2}) \\
& /f^2-1/2*e*(a+b*\operatorname{arctanh}(d*x+c))*\ln(2*d*(e^{1/2}+f^{1/2})*x)/(d*e^{1/2}+(1-c)*f^{1/2}) \\
& /f^2-1/2*b*e*\operatorname{polylog}(2,1-2/(d*x+c+1))/f^2+1/4*b*e*\operatorname{polylog}(2,1-2*d*(e^{1/2}-f^{1/2})*x)/(d*e^{1/2}-(1-c)*f^{1/2}) \\
& /f^2+1/4*b*e*\operatorname{polylog}(2,1-2*d*(e^{1/2}+f^{1/2})*x)/(d*e^{1/2}+(1-c)*f^{1/2})/f^2
\end{aligned}$$

Mathematica [A] (verified)

Time = 25.80 (sec) , antiderivative size = 694, normalized size of antiderivative = 1.82

$$\int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \frac{2ad^2(fx^2 + e \log(e - fx^2)) + b \left(2dfx + 2d^2fx^2 \operatorname{arctanh}(c + dx) + f \log(1 - c - dx) - 2cf \log(1 - c - dx) \right)}{e - fx^2}$$

input

```
Integrate[(x^3*(a + b*ArcTanh[c + d*x]))/(e - f*x^2),x]
```

output

$$\begin{aligned}
& -1/4*(2*a*d^2*(f*x^2 + e*\operatorname{Log}[e - f*x^2]) + b*(2*d*f*x + 2*d^2*f*x^2*\operatorname{ArcTan} \\
& h[c + d*x] + f*\operatorname{Log}[1 - c - d*x] - 2*c*f*\operatorname{Log}[1 - c - d*x] + c^2*f*\operatorname{Log}[1 - c \\
& - d*x] - d^2*e*\operatorname{Log}[-(\operatorname{Sqrt}[e]/\operatorname{Sqrt}[f]) + x]*\operatorname{Log}[1 - c - d*x] - d^2*e*\operatorname{Log}[\operatorname{S} \\
& \operatorname{qrt}[e]/\operatorname{Sqrt}[f] + x]*\operatorname{Log}[1 - c - d*x] + d^2*e*\operatorname{Log}[\operatorname{Sqrt}[e]/\operatorname{Sqrt}[f] + x]*\operatorname{Log}[- \\
& ((\operatorname{Sqrt}[f]*(-1 + c + d*x))/(d*\operatorname{Sqrt}[e] - (-1 + c)*\operatorname{Sqrt}[f]))] + d^2*e*\operatorname{Log}[-(\\
& \operatorname{Sqrt}[e]/\operatorname{Sqrt}[f] + x)*\operatorname{Log}[(\operatorname{Sqrt}[f]*(-1 + c + d*x))/(d*\operatorname{Sqrt}[e] + (-1 + c)*\operatorname{S} \\
& \operatorname{qrt}[f])] - f*\operatorname{Log}[1 + c + d*x] - 2*c*f*\operatorname{Log}[1 + c + d*x] - c^2*f*\operatorname{Log}[1 + c + \\
& d*x] + d^2*e*\operatorname{Log}[-(\operatorname{Sqrt}[e]/\operatorname{Sqrt}[f]) + x]*\operatorname{Log}[1 + c + d*x] + d^2*e*\operatorname{Log}[\operatorname{Sqr} \\
& t[e]/\operatorname{Sqrt}[f] + x]*\operatorname{Log}[1 + c + d*x] - d^2*e*\operatorname{Log}[\operatorname{Sqrt}[e]/\operatorname{Sqrt}[f] + x]*\operatorname{Log}[-(\\
& (\operatorname{Sqrt}[f]*(1 + c + d*x))/(d*\operatorname{Sqrt}[e] - (1 + c)*\operatorname{Sqrt}[f]))] - d^2*e*\operatorname{Log}[-(\operatorname{Sqrt} \\
& [e]/\operatorname{Sqrt}[f]) + x]*\operatorname{Log}[(\operatorname{Sqrt}[f]*(1 + c + d*x))/(d*\operatorname{Sqrt}[e] + (1 + c)*\operatorname{Sqrt}[f] \\
&)] + 2*d^2*e*\operatorname{ArcTanh}[c + d*x]*\operatorname{Log}[e - f*x^2] + d^2*e*\operatorname{Log}[1 - c - d*x]*\operatorname{Log}[\\
& e - f*x^2] - d^2*e*\operatorname{Log}[1 + c + d*x]*\operatorname{Log}[e - f*x^2] + d^2*e*\operatorname{PolyLog}[2, (d* \\
& \operatorname{Sqrt}[e]/\operatorname{Sqrt}[f] + x)/(1 - c + (d*\operatorname{Sqrt}[e])/ \operatorname{Sqrt}[f])] + d^2*e*\operatorname{PolyLog}[2, (d \\
& *(\operatorname{Sqrt}[e] - \operatorname{Sqrt}[f]*x))/(d*\operatorname{Sqrt}[e] + (-1 + c)*\operatorname{Sqrt}[f])] - d^2*e*\operatorname{PolyLog}[2, \\
& (d*(\operatorname{Sqrt}[e] - \operatorname{Sqrt}[f]*x))/(d*\operatorname{Sqrt}[e] + (1 + c)*\operatorname{Sqrt}[f])] - d^2*e*\operatorname{PolyLog}[\\
& 2, (d*(\operatorname{Sqrt}[e] + \operatorname{Sqrt}[f]*x))/(d*\operatorname{Sqrt}[e] - (1 + c)*\operatorname{Sqrt}[f])])]/(d^2*f^2)
\end{aligned}$$

Rubi [A] (verified)

Time = 1.26 (sec) , antiderivative size = 398, normalized size of antiderivative = 1.04, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx \\
 & \quad \downarrow \text{7276} \\
 & \int \left(-\frac{ax^3}{fx^2 - e} - \frac{bx^3 \operatorname{arctanh}(c + dx)}{fx^2 - e} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & -\frac{ae \log(e - fx^2)}{2f^2} - \frac{ax^2}{2f} + \frac{b \operatorname{arctanh}(c + dx) \log\left(\frac{2}{c+dx+1}\right)}{f^2} - \\
 & \quad \frac{b \operatorname{arctanh}(c + dx) \log\left(\frac{2d(\sqrt{e}-\sqrt{fx})}{(c+dx+1)(d\sqrt{e}-(1-c)\sqrt{f})}\right)}{2f^2} - \\
 & \quad \frac{b \operatorname{arctanh}(c + dx) \log\left(\frac{2d(\sqrt{e}+\sqrt{fx})}{(c+dx+1)((1-c)\sqrt{f}+d\sqrt{e})}\right)}{2f^2} - \frac{bx^2 \operatorname{arctanh}(c + dx)}{2f} - \\
 & \quad \frac{b(1-c)^2 \log(-c - dx + 1)}{4d^2 f} + \frac{b(c+1)^2 \log(c + dx + 1)}{4d^2 f} - \frac{be \operatorname{PolyLog}\left(2, 1 - \frac{2}{c+dx+1}\right)}{2f^2} + \\
 & \quad \frac{be \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e}-\sqrt{fx})}{(d\sqrt{e}-(1-c)\sqrt{f})(c+dx+1)}\right)}{4f^2} + \frac{be \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{fx}+\sqrt{e})}{(\sqrt{f}(1-c)+d\sqrt{e})(c+dx+1)}\right)}{4f^2} - \frac{bx}{2df}
 \end{aligned}$$

input

```
Int[(x^3*(a + b*ArcTanh[c + d*x]))/(e - f*x^2),x]
```


output

```

-1/2*(b*x)/(d*f) - (a*x^2)/(2*f) - (b*x^2*ArcTanh[c + d*x])/(2*f) - (b*(1
- c)^2*Log[1 - c - d*x])/(4*d^2*f) + (b*e*ArcTanh[c + d*x]*Log[2/(1 + c +
d*x)])/f^2 + (b*(1 + c)^2*Log[1 + c + d*x])/(4*d^2*f) - (b*e*ArcTanh[c + d
*x]*Log[(2*d*(Sqrt[e] - Sqrt[f]*x))/((d*Sqrt[e] - (1 - c)*Sqrt[f])*(1 + c
+ d*x))])/(2*f^2) - (b*e*ArcTanh[c + d*x]*Log[(2*d*(Sqrt[e] + Sqrt[f]*x))/
((d*Sqrt[e] + (1 - c)*Sqrt[f])*(1 + c + d*x))])/(2*f^2) - (a*e*Log[e - f*x
^2])/(2*f^2) - (b*e*PolyLog[2, 1 - 2/(1 + c + d*x)])/(2*f^2) + (b*e*PolyLo
g[2, 1 - (2*d*(Sqrt[e] - Sqrt[f]*x))/((d*Sqrt[e] - (1 - c)*Sqrt[f])*(1 + c
+ d*x))])/(4*f^2) + (b*e*PolyLog[2, 1 - (2*d*(Sqrt[e] + Sqrt[f]*x))/((d*S
qrt[e] + (1 - c)*Sqrt[f])*(1 + c + d*x))])/(4*f^2)

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 615, normalized size of antiderivative = 1.61

| method | result |
|-----------------|---|
| parts | $-\frac{ax^2}{2f} - \frac{ae \ln(fx^2 - e)}{2f^2} + b \left(-\frac{d^2 \operatorname{arctanh}\left(\frac{dx+c}{2f}\right)(dx+c)^2}{2f} + \frac{d^2 \operatorname{arctanh}\left(\frac{dx+c}{f}\right)(dx+c)c}{f} - \frac{d^4 \operatorname{arctanh}(dx+c)e \ln(c^2 f - 2cf(dx+c))}{2f^2} \right)$ |
| derivativelimit | $ad^2 \left(\frac{c(dx+c) - \frac{(dx+c)^2}{2}}{f} - \frac{e d^2 \ln(c^2 f - 2cf(dx+c) - e d^2 + f(dx+c)^2)}{2f^2} \right) + b d^2 \left(\frac{\operatorname{arctanh}\left(\frac{dx+c}{f}\right)c(dx+c)}{f} - \frac{\operatorname{arctanh}\left(\frac{dx+c}{2f}\right)(dx+c)^2}{2f} \right)$ |
| default | $ad^2 \left(\frac{c(dx+c) - \frac{(dx+c)^2}{2}}{f} - \frac{e d^2 \ln(c^2 f - 2cf(dx+c) - e d^2 + f(dx+c)^2)}{2f^2} \right) + b d^2 \left(\frac{\operatorname{arctanh}\left(\frac{dx+c}{f}\right)c(dx+c)}{f} - \frac{\operatorname{arctanh}\left(\frac{dx+c}{2f}\right)(dx+c)^2}{2f} \right)$ |
| risch | $-\frac{ax^2}{2f} + \frac{b \ln(-dx-c+1)x^2}{4f} - \frac{b \ln(-dx-c+1)}{4d^2 f} + \frac{be \operatorname{dilog}\left(\frac{d\sqrt{ef} - (-dx-c+1)f - fc + f}{d\sqrt{ef} - fc + f}\right)}{4f^2} + \frac{be \operatorname{dilog}\left(\frac{d\sqrt{ef} + (-dx-c+1)f - fc + f}{d\sqrt{ef} + fc + f}\right)}{4f^2}$ |

```
input int(x^3*(a+b*arctanh(d*x+c))/(-f*x^2+e),x,method=_RETURNVERBOSE)
```

```
output -1/2*a/f*x^2-1/2*a*e/f^2*ln(f*x^2-e)+b/d^4*(-1/2*d^2*arctanh(d*x+c)/f*(d*x+c)^2+d^2*arctanh(d*x+c)/f*(d*x+c)*c-1/2*d^4*arctanh(d*x+c)*e/f^2*ln(c^2*f-2*c*f*(d*x+c)-e*d^2+f*(d*x+c)^2)+1/2*d^2*(-1/f*(d*x+c+1/2*(-2*c+1)*ln(d*x+c-1)-1/2*(2*c+1)*ln(d*x+c+1))-e*d^2/f^2*(-1/2*ln(d*x+c+1)*ln(c^2*f-2*c*f*(d*x+c)-e*d^2+f*(d*x+c)^2)+f*(1/2*ln(d*x+c+1)*(ln((d*(e*f)^(1/2)+f*c-f*(d*x+c+1)+f)/(d*(e*f)^(1/2)+f*c+f))+ln((d*(e*f)^(1/2)-f*c+f*(d*x+c+1)-f)/(d*(e*f)^(1/2)-f*c-f)))/f+1/2*(dilog((d*(e*f)^(1/2)+f*c-f*(d*x+c+1)+f)/(d*(e*f)^(1/2)+f*c+f))+dilog((d*(e*f)^(1/2)-f*c+f*(d*x+c+1)-f)/(d*(e*f)^(1/2)-f*c-f)))/f)+1/2*ln(d*x+c-1)*ln(c^2*f-2*c*f*(d*x+c)-e*d^2+f*(d*x+c)^2)-f*(1/2*ln(d*x+c-1)*(ln((d*(e*f)^(1/2)+f*c-f*(d*x+c-1)-f)/(d*(e*f)^(1/2)+f*c-f))+ln((d*(e*f)^(1/2)-f*c+f*(d*x+c-1)+f)/(d*(e*f)^(1/2)-f*c+f)))/f+1/2*(dilog((d*(e*f)^(1/2)+f*c-f*(d*x+c-1)-f)/(d*(e*f)^(1/2)+f*c+f))+dilog((d*(e*f)^(1/2)-f*c+f*(d*x+c-1)+f)/(d*(e*f)^(1/2)-f*c+f)))/f))))
```

Fricas [F]

$$\int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \int -\frac{(b \operatorname{artanh}(dx + c) + a)x^3}{fx^2 - e} dx$$

input `integrate(x^3*(a+b*arctanh(d*x+c))/(-f*x^2+e),x, algorithm="fricas")`

output `integral(-(b*x^3*arctanh(d*x + c) + a*x^3)/(f*x^2 - e), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \text{Timed out}$$

input `integrate(x**3*(a+b*atanh(d*x+c))/(-f*x**2+e),x)`

output `Timed out`

Maxima [F]

$$\int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \int -\frac{(b \operatorname{artanh}(dx + c) + a)x^3}{fx^2 - e} dx$$

input `integrate(x^3*(a+b*arctanh(d*x+c))/(-f*x^2+e),x, algorithm="maxima")`

output `-1/2*a*(x^2/f + e*log(f*x^2 - e)/f^2) - 1/2*b*integrate(x^3*(log(d*x + c + 1) - log(-d*x - c + 1))/(f*x^2 - e), x)`

Giac [F]

$$\int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \int -\frac{(b \operatorname{artanh}(dx + c) + a)x^3}{fx^2 - e} dx$$

input `integrate(x^3*(a+b*arctanh(d*x+c))/(-f*x^2+e),x, algorithm="giac")`

output `integrate(-(b*arctanh(d*x + c) + a)*x^3/(f*x^2 - e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \int \frac{x^3(a + b \operatorname{atanh}(c + dx))}{e - fx^2} dx$$

input `int((x^3*(a + b*atanh(c + d*x)))/(e - f*x^2),x)`

output `int((x^3*(a + b*atanh(c + d*x)))/(e - f*x^2), x)`

Reduce [F]

$$\int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \text{Too large to display}$$

input `int(x^3*(a+b*atanh(d*x+c))/(-f*x^2+e),x)`

output

```
(3*atanh(c + d*x)**2*b*c**2*d**2*e*f + atanh(c + d*x)**2*b*d**4*e**2 + atanh(c + d*x)**2*b*d**2*e*f + 2*atanh(c + d*x)*b*c**3*f**2 + 4*atanh(c + d*x)*b*c**2*f**2 - 2*atanh(c + d*x)*b*c*d**2*e*f - 2*atanh(c + d*x)*b*c*d**2*f**2*x**2 + 2*atanh(c + d*x)*b*c*f**2 - 2*atanh(c + d*x)*b*d**3*e*f*x - 2*atanh(c + d*x)*b*d**2*e*f + 8*int(atanh(c + d*x)/(c**2*e - c**2*f*x**2 + 2*c*d*e*x - 2*c*d*f*x**3 + d**2*e*x**2 - d**2*f*x**4 - e + f*x**2),x)*b*c**2*d**3*e**2*f + 2*int(atanh(c + d*x)/(c**2*e - c**2*f*x**2 + 2*c*d*e*x - 2*c*d*f*x**3 + d**2*e*x**2 - d**2*f*x**4 - e + f*x**2),x)*b*d**5*e**3 - 2*int((atanh(c + d*x)*x**4)/(c**2*e - c**2*f*x**2 + 2*c*d*e*x - 2*c*d*f*x**3 + d**2*e*x**2 - d**2*f*x**4 - e + f*x**2),x)*b*d**5*e*f**2 + 4*int((atanh(c + d*x)*x)/(c**2*e - c**2*f*x**2 + 2*c*d*e*x - 2*c*d*f*x**3 + d**2*e*x**2 - d**2*f*x**4 - e + f*x**2),x)*b*c**3*d**2*e*f**2 + 4*int((atanh(c + d*x)*x)/(c**2*e - c**2*f*x**2 + 2*c*d*e*x - 2*c*d*f*x**3 + d**2*e*x**2 - d**2*f*x**4 - e + f*x**2),x)*b*c*d**4*e**2*f - 4*int((atanh(c + d*x)*x)/(c**2*e - c**2*f*x**2 + 2*c*d*e*x - 2*c*d*f*x**3 + d**2*e*x**2 - d**2*f*x**4 - e + f*x**2),x)*b*c*d**2*e*f**2 - 2*log(-sqrt(f)*sqrt(e) - f*x)*a*c*d**2*e*f - 2*log(sqrt(f)*sqrt(e) - f*x)*a*c*d**2*e*f + 4*log(c + d*x - 1)*b*c**2*f**2 - 2*log(c + d*x - 1)*b*d**2*e*f - 2*a*c*d**2*f**2*x**2 - 2*b*c*d*f**2*x)/(4*c*d**2*f**3)
```

3.64 $\int \frac{x^2(a+b\operatorname{arctanh}(c+dx))}{e-fx^2} dx$

| | |
|---|-----|
| Optimal result | 565 |
| Mathematica [C] (warning: unable to verify) | 566 |
| Rubi [A] (verified) | 567 |
| Maple [B] (verified) | 569 |
| Fricas [F] | 569 |
| Sympy [F(-1)] | 570 |
| Maxima [F(-2)] | 570 |
| Giac [F] | 571 |
| Mupad [F(-1)] | 571 |
| Reduce [F] | 571 |

Optimal result

Integrand size = 24, antiderivative size = 323

$$\begin{aligned}
 & \int \frac{x^2(a + b\operatorname{arctanh}(c + dx))}{e - fx^2} dx \\
 &= -\frac{ax}{f} - \frac{b(c + dx)\operatorname{arctanh}(c + dx)}{df} \\
 &\quad - \frac{\sqrt{e}(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2d(\sqrt{e} - \sqrt{fx})}{(d\sqrt{e} - (1-c)\sqrt{f})(1+c+dx)}\right)}{2f^{3/2}} \\
 &\quad + \frac{\sqrt{e}(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2d(\sqrt{e} + \sqrt{fx})}{(d\sqrt{e} + (1-c)\sqrt{f})(1+c+dx)}\right)}{2f^{3/2}} \\
 &\quad - \frac{b \log(1 - (c + dx)^2)}{2df} + \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e} - \sqrt{fx})}{(d\sqrt{e} - (1-c)\sqrt{f})(1+c+dx)}\right)}{4f^{3/2}} \\
 &\quad - \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e} + \sqrt{fx})}{(d\sqrt{e} + (1-c)\sqrt{f})(1+c+dx)}\right)}{4f^{3/2}}
 \end{aligned}$$

output

```
-a*x/f-b*(d*x+c)*arctanh(d*x+c)/d/f-1/2*e^(1/2)*(a+b*arctanh(d*x+c))*ln(2*
d*(e^(1/2)-f^(1/2)*x)/(d*e^(1/2)-(1-c)*f^(1/2))/(d*x+c+1))/f^(3/2)+1/2*e^(
1/2)*(a+b*arctanh(d*x+c))*ln(2*d*(e^(1/2)+f^(1/2)*x)/(d*e^(1/2)+(1-c)*f^(1
/2))/(d*x+c+1))/f^(3/2)-1/2*b*ln(1-(d*x+c)^2)/d/f+1/4*b*e^(1/2)*polylog(2,
1-2*d*(e^(1/2)-f^(1/2)*x)/(d*e^(1/2)-(1-c)*f^(1/2))/(d*x+c+1))/f^(3/2)-1/4
*b*e^(1/2)*polylog(2,1-2*d*(e^(1/2)+f^(1/2)*x)/(d*e^(1/2)+(1-c)*f^(1/2))/(
d*x+c+1))/f^(3/2)
```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 11.08 (sec) , antiderivative size = 1198, normalized size of antiderivative = 3.71

$$\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \text{Too large to display}$$

input

```
Integrate[(x^2*(a + b*ArcTanh[c + d*x]))/(e - f*x^2),x]
```

output

```

-((a*x)/f) + (a*Sqrt[e]*ArcTanh[(Sqrt[f]*x)/Sqrt[e]])/f^(3/2) + (b*(-((c +
d*x)*ArcTanh[c + d*x]) + Log[1/Sqrt[1 - (c + d*x)^2]]))/(d*f) + (b*Sqrt[e
]*(2*d^2*e*ArcTanh[c - (d*Sqrt[e])/Sqrt[f]]*ArcTanh[c + d*x] - 2*c^2*f*Arc
Tanh[c - (d*Sqrt[e])/Sqrt[f]]*ArcTanh[c + d*x] - 2*d^2*e*ArcTanh[c + (d*Sq
rt[e])/Sqrt[f]]*ArcTanh[c + d*x] + 2*c^2*f*ArcTanh[c + (d*Sqrt[e])/Sqrt[f]
]*ArcTanh[c + d*x] - 2*d*Sqrt[e]*Sqrt[f]*ArcTanh[c + d*x]^2 + d*Sqrt[e]*E^
ArcTanh[c + (d*Sqrt[e])/Sqrt[f]]*Sqrt[1 - c^2 - (d^2*e)/f - (2*c*d*Sqrt[e]
)/Sqrt[f]]*Sqrt[f]*ArcTanh[c + d*x]^2 + d*Sqrt[e]*E^ArcTanh[c - (d*Sqrt[e]
)/Sqrt[f]]*Sqrt[1 - c^2 - (d^2*e)/f + (2*c*d*Sqrt[e])/Sqrt[f]]*Sqrt[f]*Arc
Tanh[c + d*x]^2 - c*E^ArcTanh[c + (d*Sqrt[e])/Sqrt[f]]*Sqrt[1 - c^2 - (d^2
*e)/f - (2*c*d*Sqrt[e])/Sqrt[f]]*f*ArcTanh[c + d*x]^2 + c*E^ArcTanh[c - (d
*Sqrt[e])/Sqrt[f]]*Sqrt[1 - c^2 - (d^2*e)/f + (2*c*d*Sqrt[e])/Sqrt[f]]*f*A
rcTanh[c + d*x]^2 + 2*d^2*e*ArcTanh[c - (d*Sqrt[e])/Sqrt[f]]*Log[1 - E^(2*
ArcTanh[c - (d*Sqrt[e])/Sqrt[f]] - 2*ArcTanh[c + d*x])] - 2*c^2*f*ArcTanh[
c - (d*Sqrt[e])/Sqrt[f]]*Log[1 - E^(2*ArcTanh[c - (d*Sqrt[e])/Sqrt[f]] - 2
*ArcTanh[c + d*x])] - 2*d^2*e*ArcTanh[c + d*x]*Log[1 - E^(2*ArcTanh[c - (d
*Sqrt[e])/Sqrt[f]] - 2*ArcTanh[c + d*x])] + 2*c^2*f*ArcTanh[c + d*x]*Log[1
- E^(2*ArcTanh[c - (d*Sqrt[e])/Sqrt[f]] - 2*ArcTanh[c + d*x])] - 2*d^2*e*
ArcTanh[c + (d*Sqrt[e])/Sqrt[f]]*Log[1 - E^(2*ArcTanh[c + (d*Sqrt[e])/Sqrt
[f]] - 2*ArcTanh[c + d*x])] + 2*c^2*f*ArcTanh[c + (d*Sqrt[e])/Sqrt[f]]*...

```

Rubi [A] (verified)

Time = 1.16 (sec) , antiderivative size = 524, normalized size of antiderivative = 1.62, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx$$

$$\downarrow 7276$$

$$\int \left(-\frac{ax^2}{fx^2 - e} - \frac{bx^2 \operatorname{arctanh}(c + dx)}{fx^2 - e} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& \frac{a\sqrt{e}\operatorname{arctanh}\left(\frac{\sqrt{fx}}{\sqrt{e}}\right)}{f^{3/2}} - \frac{ax}{f} - \frac{b(c+dx)\operatorname{arctanh}(c+dx)}{df} + \frac{b\sqrt{e}\operatorname{PolyLog}\left(2, -\frac{\sqrt{f}(-c-dx+1)}{d\sqrt{e-(1-c)\sqrt{f}}}\right)}{4f^{3/2}} - \\
& \frac{b\sqrt{e}\operatorname{PolyLog}\left(2, \frac{\sqrt{f}(-c-dx+1)}{\sqrt{f}(1-c)+d\sqrt{e}}\right)}{4f^{3/2}} + \frac{b\sqrt{e}\operatorname{PolyLog}\left(2, -\frac{\sqrt{f}(c+dx+1)}{d\sqrt{e-(c+1)\sqrt{f}}}\right)}{4f^{3/2}} - \\
& \frac{b\sqrt{e}\operatorname{PolyLog}\left(2, \frac{\sqrt{f}(c+dx+1)}{\sqrt{f}(c+1)+d\sqrt{e}}\right)}{4f^{3/2}} + \frac{b\sqrt{e}\log(-c-dx+1)\log\left(\frac{d(\sqrt{e}-\sqrt{fx})}{d\sqrt{e-(1-c)\sqrt{f}}}\right)}{4f^{3/2}} - \\
& \frac{b\sqrt{e}\log(c+dx+1)\log\left(\frac{d(\sqrt{e}-\sqrt{fx})}{(c+1)\sqrt{f}+d\sqrt{e}}\right)}{4f^{3/2}} - \frac{b\sqrt{e}\log(-c-dx+1)\log\left(\frac{d(\sqrt{e}+\sqrt{fx})}{(1-c)\sqrt{f}+d\sqrt{e}}\right)}{4f^{3/2}} + \\
& \frac{b\sqrt{e}\log(c+dx+1)\log\left(\frac{d(\sqrt{e}+\sqrt{fx})}{d\sqrt{e-(c+1)\sqrt{f}}}\right)}{4f^{3/2}} - \frac{b\log(1-(c+dx)^2)}{2df}
\end{aligned}$$

input `Int[(x^2*(a + b*ArcTanh[c + d*x]))/(e - f*x^2), x]`

output `-((a*x)/f) + (a*Sqrt[e]*ArcTanh[(Sqrt[f]*x)/Sqrt[e]])/f^(3/2) - (b*(c + d*x)*ArcTanh[c + d*x])/(d*f) + (b*Sqrt[e]*Log[1 - c - d*x]*Log[(d*(Sqrt[e] - Sqrt[f]*x))/(d*Sqrt[e] - (1 - c)*Sqrt[f])])/(4*f^(3/2)) - (b*Sqrt[e]*Log[1 + c + d*x]*Log[(d*(Sqrt[e] - Sqrt[f]*x))/(d*Sqrt[e] + (1 + c)*Sqrt[f])])/(4*f^(3/2)) - (b*Sqrt[e]*Log[1 - c - d*x]*Log[(d*(Sqrt[e] + Sqrt[f]*x))/(d*Sqrt[e] + (1 - c)*Sqrt[f])])/(4*f^(3/2)) + (b*Sqrt[e]*Log[1 + c + d*x]*Log[(d*(Sqrt[e] + Sqrt[f]*x))/(d*Sqrt[e] - (1 + c)*Sqrt[f])])/(4*f^(3/2)) - (b*Log[1 - (c + d*x)^2])/(2*d*f) + (b*Sqrt[e]*PolyLog[2, -((Sqrt[f]*(1 - c - d*x))/(d*Sqrt[e] - (1 - c)*Sqrt[f])])/(4*f^(3/2)) - (b*Sqrt[e]*PolyLog[2, (Sqrt[f]*(1 - c - d*x))/(d*Sqrt[e] + (1 - c)*Sqrt[f])])/(4*f^(3/2)) + (b*Sqrt[e]*PolyLog[2, -((Sqrt[f]*(1 + c + d*x))/(d*Sqrt[e] - (1 + c)*Sqrt[f])])/(4*f^(3/2)) - (b*Sqrt[e]*PolyLog[2, (Sqrt[f]*(1 + c + d*x))/(d*Sqrt[e] + (1 + c)*Sqrt[f])])/(4*f^(3/2))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 638 vs. $2(265) = 530$.

Time = 1.99 (sec) , antiderivative size = 639, normalized size of antiderivative = 1.98

| method | result |
|------------------|---|
| risch | $\frac{b \ln(-dx-c+1)x}{2f} + \frac{b \ln(-dx-c+1)c}{2df} - \frac{b \ln(-dx-c+1)}{2df} + \frac{b}{df} - \frac{be \ln(-dx-c+1) \ln\left(\frac{d\sqrt{ef}-(-dx-c+1)f-fc+f}{d\sqrt{ef}-fc+f}\right)}{4f\sqrt{ef}}$ |
| derivativdivides | Expression too large to display |
| default | Expression too large to display |
| parts | Expression too large to display |

input `int(x^2*(a+b*arctanh(d*x+c))/(-f*x^2+e),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/2*b/f*\ln(-d*x-c+1)*x+1/2/d*b/f*\ln(-d*x-c+1)*c-1/2/d*b/f*\ln(-d*x-c+1)+1/d \\ & *b/f-1/4*b*e/f*\ln(-d*x-c+1)/(e*f)^{(1/2)}*\ln((d*(e*f)^{(1/2)}-(-d*x-c+1)*f-f*c \\ & +f)/(d*(e*f)^{(1/2)}-f*c+f))+1/4*b*e/f*\ln(-d*x-c+1)/(e*f)^{(1/2)}*\ln((d*(e*f)^{(1/2)} \\ & +(-d*x-c+1)*f+f*c-f)/(d*(e*f)^{(1/2)}+f*c-f))-1/4*b*e/f/(e*f)^{(1/2)}*di \\ & log((d*(e*f)^{(1/2)}-(-d*x-c+1)*f-f*c+f)/(d*(e*f)^{(1/2)}-f*c+f))+1/4*b*e/f/(e* \\ & f)^{(1/2)}*dilog((d*(e*f)^{(1/2)}+(-d*x-c+1)*f+f*c-f)/(d*(e*f)^{(1/2)}+f*c-f))-a \\ & *x/f-1/d*a/f*c+1/d*a/f-a*e/f/(e*f)^{(1/2)}*arctanh(1/2*(2*(-d*x-c+1)*f+2*f*c \\ & -2*f)/d/(e*f)^{(1/2)})-1/2*b/f*\ln(d*x+c+1)*x-1/2*b/d/f*\ln(d*x+c+1)*c-1/2*b/d \\ & /f*\ln(d*x+c+1)-1/4*b*e/f*\ln(d*x+c+1)/(e*f)^{(1/2)}*\ln((d*(e*f)^{(1/2)}+f*c-f*(\\ & d*x+c+1)+f)/(d*(e*f)^{(1/2)}+f*c+f))+1/4*b*e/f*\ln(d*x+c+1)/(e*f)^{(1/2)}*\ln((d \\ & *(e*f)^{(1/2)}-f*c+f*(d*x+c+1)-f)/(d*(e*f)^{(1/2)}-f*c-f))-1/4*b*e/f/(e*f)^{(1/2)} \\ & *dilog((d*(e*f)^{(1/2)}+f*c-f*(d*x+c+1)+f)/(d*(e*f)^{(1/2)}+f*c+f))+1/4*b*e/ \\ & f/(e*f)^{(1/2)}*dilog((d*(e*f)^{(1/2)}-f*c+f*(d*x+c+1)-f)/(d*(e*f)^{(1/2)}-f*c-f) \\ &)) \end{aligned}$$
Fricas [F]

$$\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \int -\frac{(b \operatorname{arctanh}(dx + c) + a)x^2}{fx^2 - e} dx$$

input `integrate(x^2*(a+b*arctanh(d*x+c))/(-f*x^2+e),x, algorithm="fricas")`

output `integral(-(b*x^2*arctanh(d*x + c) + a*x^2)/(f*x^2 - e), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \text{Timed out}$$

input `integrate(x**2*(a+b*atanh(d*x+c))/(-f*x**2+e),x)`

output `Timed out`

Maxima [F(-2)]

Exception generated.

$$\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \text{Exception raised: ValueError}$$

input `integrate(x^2*(a+b*arctanh(d*x+c))/(-f*x^2+e),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

Giac [F]

$$\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \int -\frac{(b \operatorname{artanh}(dx + c) + a)x^2}{fx^2 - e} dx$$

input `integrate(x^2*(a+b*arctanh(d*x+c))/(-f*x^2+e),x, algorithm="giac")`

output `integrate(-(b*arctanh(d*x + c) + a)*x^2/(f*x^2 - e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \int \frac{x^2(a + b \operatorname{atanh}(c + dx))}{e - fx^2} dx$$

input `int((x^2*(a + b*atanh(c + d*x)))/(e - f*x^2),x)`

output `int((x^2*(a + b*atanh(c + d*x)))/(e - f*x^2), x)`

Reduce [F]

$$\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \frac{\sqrt{f} \sqrt{e} \log(-\sqrt{f} \sqrt{e} - fx) a - \sqrt{f} \sqrt{e} \log(\sqrt{f} \sqrt{e} - fx) a + 2 \left(\int \frac{\operatorname{atanh}(dx+c)x^2}{-fx^2+e} dx \right) b f^2 - 2afx}{2f^2}$$

input `int(x^2*(a+b*atanh(d*x+c))/(-f*x^2+e),x)`

output `(sqrt(f)*sqrt(e)*log(-sqrt(f)*sqrt(e) - f*x)*a - sqrt(f)*sqrt(e)*log(sqrt(f)*sqrt(e) - f*x)*a + 2*int((atanh(c + d*x)*x**2)/(e - f*x**2),x)*b*f**2 - 2*a*f*x)/(2*f**2)`

3.65 $\int \frac{x(a+b\operatorname{arctanh}(c+dx))}{e-fx^2} dx$

| | |
|----------------------------|-----|
| Optimal result | 572 |
| Mathematica [A] (verified) | 573 |
| Rubi [A] (verified) | 574 |
| Maple [A] (verified) | 575 |
| Fricas [F] | 577 |
| Sympy [F(-1)] | 577 |
| Maxima [F] | 577 |
| Giac [F] | 578 |
| Mupad [F(-1)] | 578 |
| Reduce [F] | 578 |

Optimal result

Integrand size = 22, antiderivative size = 292

$$\int \frac{x(a + b\operatorname{arctanh}(c + dx))}{e - fx^2} dx = \frac{(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{f} - \frac{(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2d(\sqrt{e}-\sqrt{fx})}{(d\sqrt{e}-(1-c)\sqrt{f})(1+c+dx)}\right)}{2f} - \frac{(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2d(\sqrt{e}+\sqrt{fx})}{(d\sqrt{e}+(1-c)\sqrt{f})(1+c+dx)}\right)}{2f} - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{2f} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e}-\sqrt{fx})}{(d\sqrt{e}-(1-c)\sqrt{f})(1+c+dx)}\right)}{4f} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e}+\sqrt{fx})}{(d\sqrt{e}+(1-c)\sqrt{f})(1+c+dx)}\right)}{4f}$$

output

```
(a+b*arctanh(d*x+c))*ln(2/(d*x+c+1))/f-1/2*(a+b*arctanh(d*x+c))*ln(2*d*(e^(1/2)-f^(1/2)*x)/(d*e^(1/2)-(1-c)*f^(1/2))/(d*x+c+1))/f-1/2*(a+b*arctanh(d*x+c))*ln(2*d*(e^(1/2)+f^(1/2)*x)/(d*e^(1/2)+(1-c)*f^(1/2))/(d*x+c+1))/f-1/2*b*polylog(2,1-2/(d*x+c+1))/f+1/4*b*polylog(2,1-2*d*(e^(1/2)-f^(1/2)*x)/(d*e^(1/2)-(1-c)*f^(1/2))/(d*x+c+1))/f+1/4*b*polylog(2,1-2*d*(e^(1/2)+f^(1/2)*x)/(d*e^(1/2)+(1-c)*f^(1/2))/(d*x+c+1))/f
```

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 410, normalized size of antiderivative = 1.40

$$\int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \frac{b \log(1 - c - dx) \log\left(\frac{d(\sqrt{e} - \sqrt{fx})}{d\sqrt{e} - (1-c)\sqrt{f}}\right)}{4f} - \frac{b \log(1 + c + dx) \log\left(\frac{d(\sqrt{e} - \sqrt{fx})}{d\sqrt{e} + (1+c)\sqrt{f}}\right)}{4f} + \frac{b \log(1 - c - dx) \log\left(\frac{d(\sqrt{e} + \sqrt{fx})}{d\sqrt{e} + (1-c)\sqrt{f}}\right)}{4f} - \frac{b \log(1 + c + dx) \log\left(\frac{d(\sqrt{e} + \sqrt{fx})}{d\sqrt{e} - (1+c)\sqrt{f}}\right)}{4f} - \frac{a \log(e - fx^2)}{2f} + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{f}(1-c-dx)}{d\sqrt{e} - (1-c)\sqrt{f}}\right)}{4f} + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{f}(1-c-dx)}{d\sqrt{e} + (1-c)\sqrt{f}}\right)}{4f} - \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{f}(1+c+dx)}{d\sqrt{e} - (1+c)\sqrt{f}}\right)}{4f} - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{f}(1+c+dx)}{d\sqrt{e} + (1+c)\sqrt{f}}\right)}{4f}$$

input

```
Integrate[(x*(a + b*ArcTanh[c + d*x]))/(e - f*x^2),x]
```

output

```
(b*Log[1 - c - d*x]*Log[(d*(Sqrt[e] - Sqrt[f]*x))/(d*Sqrt[e] - (1 - c)*Sqrt[f])])/(4*f) - (b*Log[1 + c + d*x]*Log[(d*(Sqrt[e] - Sqrt[f]*x))/(d*Sqrt[e] + (1 + c)*Sqrt[f])])/(4*f) + (b*Log[1 - c - d*x]*Log[(d*(Sqrt[e] + Sqrt[f]*x))/(d*Sqrt[e] + (1 - c)*Sqrt[f])])/(4*f) - (b*Log[1 + c + d*x]*Log[(d*(Sqrt[e] + Sqrt[f]*x))/(d*Sqrt[e] - (1 + c)*Sqrt[f])])/(4*f) - (a*Log[e - f*x^2])/(2*f) + (b*PolyLog[2, -((Sqrt[f]*(1 - c - d*x))/(d*Sqrt[e] - (1 - c)*Sqrt[f]))])/(4*f) + (b*PolyLog[2, (Sqrt[f]*(1 - c - d*x))/(d*Sqrt[e] + (1 - c)*Sqrt[f])])/(4*f) - (b*PolyLog[2, -((Sqrt[f]*(1 + c + d*x))/(d*Sqrt[e] - (1 + c)*Sqrt[f])])/(4*f) - (b*PolyLog[2, (Sqrt[f]*(1 + c + d*x))/(d*Sqrt[e] + (1 + c)*Sqrt[f])])/(4*f)
```

Rubi [A] (verified)

Time = 0.87 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.03, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx$$

$$\downarrow 7276$$

$$\int \left(-\frac{ax}{fx^2 - e} - \frac{b \operatorname{arctanh}(c + dx)}{fx^2 - e} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned} & -\frac{a \log(e - fx^2)}{2f} - \frac{b \operatorname{arctanh}(c + dx) \log\left(\frac{2d(\sqrt{e} - \sqrt{fx})}{(c + dx + 1)(d\sqrt{e} - (1 - c)\sqrt{f})}\right)}{2f} - \\ & \frac{b \operatorname{arctanh}(c + dx) \log\left(\frac{2d(\sqrt{e} + \sqrt{fx})}{(c + dx + 1)((1 - c)\sqrt{f} + d\sqrt{e})}\right)}{2f} + \frac{b \operatorname{arctanh}(c + dx) \log\left(\frac{2}{c + dx + 1}\right)}{f} + \\ & \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e} - \sqrt{fx})}{(d\sqrt{e} - (1 - c)\sqrt{f})(c + dx + 1)}\right)}{4f} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{fx} + \sqrt{e})}{(\sqrt{f}(1 - c) + d\sqrt{e})(c + dx + 1)}\right)}{4f} - \\ & \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{c + dx + 1}\right)}{2f} \end{aligned}$$

input `Int[(x*(a + b*ArcTanh[c + d*x]))/(e - f*x^2),x]`

output `(b*ArcTanh[c + d*x]*Log[2/(1 + c + d*x)]/f - (b*ArcTanh[c + d*x]*Log[(2*d*(Sqrt[e] - Sqrt[f]*x))/((d*Sqrt[e] - (1 - c)*Sqrt[f])*(1 + c + d*x))]/(2*f) - (b*ArcTanh[c + d*x]*Log[(2*d*(Sqrt[e] + Sqrt[f]*x))/((d*Sqrt[e] + (1 - c)*Sqrt[f])*(1 + c + d*x))]/(2*f) - (a*Log[e - f*x^2])/(2*f) - (b*PolyLog[2, 1 - 2/(1 + c + d*x)]/(2*f) + (b*PolyLog[2, 1 - (2*d*(Sqrt[e] - Sqrt[f]*x))/((d*Sqrt[e] - (1 - c)*Sqrt[f])*(1 + c + d*x))]/(4*f) + (b*PolyLog[2, 1 - (2*d*(Sqrt[e] + Sqrt[f]*x))/((d*Sqrt[e] + (1 - c)*Sqrt[f])*(1 + c + d*x))]/(4*f)`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 474, normalized size of antiderivative = 1.62

| method | result |
|------------------|--|
| risch | $\frac{b \ln(-dx-c+1) \ln\left(\frac{d\sqrt{ef}-(-dx-c+1)f-fc+f}{d\sqrt{ef}-fc+f}\right)}{4f} + \frac{b \ln(-dx-c+1) \ln\left(\frac{d\sqrt{ef}+(-dx-c+1)f+fc-f}{d\sqrt{ef}+fc-f}\right)}{4f} + b \operatorname{dilog}\left(\frac{d\sqrt{ef}-(-dx-c+1)f-fc+f}{d\sqrt{ef}-fc+f}\right) + \frac{b \operatorname{dilog}\left(\frac{d\sqrt{ef}+(-dx-c+1)f+fc-f}{d\sqrt{ef}+fc-f}\right)}{4f}$ |
| parts | $-\frac{a \ln(fx^2-e)}{2f} + b \left(-\frac{d^2 \ln(c^2 f - 2cf(dx+c) - e d^2 + f(dx+c)^2) \operatorname{arctanh}(dx+c)}{2f} + \frac{d^2 \left(\frac{\ln(dx+c+1) \ln(c^2 f - 2cf(dx+c) - e d^2 + f(dx+c)^2)}{2} \right)}{2f} \right)$ |
| derivativdivides | $-\frac{a d^2 \ln(c^2 f - 2cf(dx+c) - e d^2 + f(dx+c)^2)}{2f} + b d^2 \left(-\frac{\ln(c^2 f - 2cf(dx+c) - e d^2 + f(dx+c)^2) \operatorname{arctanh}(dx+c)}{2f} + \frac{\ln(dx+c-1) \ln(c^2 f - 2cf(dx+c) - e d^2 + f(dx+c)^2)}{2} \right)$ |
| default | $-\frac{a d^2 \ln(c^2 f - 2cf(dx+c) - e d^2 + f(dx+c)^2)}{2f} + b d^2 \left(-\frac{\ln(c^2 f - 2cf(dx+c) - e d^2 + f(dx+c)^2) \operatorname{arctanh}(dx+c)}{2f} + \frac{\ln(dx+c-1) \ln(c^2 f - 2cf(dx+c) - e d^2 + f(dx+c)^2)}{2} \right)$ |

input `int(x*(a+b*arctanh(d*x+c))/(-f*x^2+e),x,method=_RETURNVERBOSE)`

output

```

1/4*b*ln(-d*x-c+1)/f*ln((d*(e*f)^(1/2)-(-d*x-c+1)*f-f*c+f)/(d*(e*f)^(1/2)-f*c+f))+1/4*b*ln(-d*x-c+1)/f*ln((d*(e*f)^(1/2)+(-d*x-c+1)*f+f*c-f)/(d*(e*f)^(1/2)+f*c-f))+1/4*b/f*dilog((d*(e*f)^(1/2)-(-d*x-c+1)*f-f*c+f)/(d*(e*f)^(1/2)-f*c+f))+1/4*b/f*dilog((d*(e*f)^(1/2)+(-d*x-c+1)*f+f*c-f)/(d*(e*f)^(1/2)+f*c-f))-1/2*a/f*ln(f*(-d*x-c+1)^2+2*(-d*x-c+1)*c*f+c^2*f-e*d^2-2*(-d*x-c+1)*f-2*f*c+f)-1/4*b*ln(d*x+c+1)/f*ln((d*(e*f)^(1/2)+f*c-f*(d*x+c+1)+f)/(d*(e*f)^(1/2)+f*c+f))-1/4*b*ln(d*x+c+1)/f*ln((d*(e*f)^(1/2)-f*c+f*(d*x+c+1)-f)/(d*(e*f)^(1/2)-f*c+f))-1/4*b/f*dilog((d*(e*f)^(1/2)+f*c-f*(d*x+c+1)+f)/(d*(e*f)^(1/2)+f*c+f))-1/4*b/f*dilog((d*(e*f)^(1/2)-f*c+f*(d*x+c+1)-f)/(d*(e*f)^(1/2)-f*c+f))
    
```

Fricas [F]

$$\int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \int -\frac{(b \operatorname{artanh}(dx + c) + a)x}{fx^2 - e} dx$$

input `integrate(x*(a+b*arctanh(d*x+c))/(-f*x^2+e),x, algorithm="fricas")`

output `integral(-(b*x*arctanh(d*x + c) + a*x)/(f*x^2 - e), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \text{Timed out}$$

input `integrate(x*(a+b*atanh(d*x+c))/(-f*x**2+e),x)`

output `Timed out`

Maxima [F]

$$\int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \int -\frac{(b \operatorname{artanh}(dx + c) + a)x}{fx^2 - e} dx$$

input `integrate(x*(a+b*arctanh(d*x+c))/(-f*x^2+e),x, algorithm="maxima")`

output `-1/2*b*integrate(x*(log(d*x + c + 1) - log(-d*x - c + 1))/(f*x^2 - e), x)
- 1/2*a*log(f*x^2 - e)/f`

Giac [F]

$$\int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \int -\frac{(b \operatorname{artanh}(dx + c) + a)x}{fx^2 - e} dx$$

input `integrate(x*(a+b*arctanh(d*x+c))/(-f*x^2+e),x, algorithm="giac")`

output `integrate(-(b*arctanh(d*x + c) + a)*x/(f*x^2 - e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx = \int \frac{x(a + b \operatorname{atanh}(c + dx))}{e - fx^2} dx$$

input `int((x*(a + b*atanh(c + d*x)))/(e - f*x^2),x)`

output `int((x*(a + b*atanh(c + d*x)))/(e - f*x^2), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e - fx^2} dx \\ &= \frac{2 \left(\int \frac{\operatorname{atanh}(dx+c)x}{-fx^2+e} dx \right) bf - \log(-\sqrt{f}\sqrt{e} - fx) a - \log(\sqrt{f}\sqrt{e} - fx) a}{2f} \end{aligned}$$

input `int(x*(a+b*atanh(d*x+c))/(-f*x^2+e),x)`

output `(2*int((atanh(c + d*x)*x)/(e - f*x**2),x)*b*f - log(-sqrt(f)*sqrt(e) - f*x)*a - log(sqrt(f)*sqrt(e) - f*x)*a)/(2*f)`

3.66 $\int \frac{a+b\operatorname{arctanh}(c+dx)}{e-fx^2} dx$

| | |
|----------------------------|-----|
| Optimal result | 579 |
| Mathematica [A] (verified) | 580 |
| Rubi [A] (verified) | 580 |
| Maple [B] (verified) | 582 |
| Fricas [F] | 582 |
| Sympy [F(-1)] | 583 |
| Maxima [F(-2)] | 583 |
| Giac [F] | 583 |
| Mupad [F(-1)] | 584 |
| Reduce [F] | 584 |

Optimal result

Integrand size = 21, antiderivative size = 273

$$\int \frac{a + b\operatorname{arctanh}(c + dx)}{e - fx^2} dx = -\frac{(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2d(\sqrt{e}-\sqrt{fx})}{(d\sqrt{e}-(1-c)\sqrt{f})(1+c+dx)}\right)}{2\sqrt{e}\sqrt{f}} + \frac{(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2d(\sqrt{e}+\sqrt{fx})}{(d\sqrt{e}+(1-c)\sqrt{f})(1+c+dx)}\right)}{2\sqrt{e}\sqrt{f}} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e}-\sqrt{fx})}{(d\sqrt{e}-(1-c)\sqrt{f})(1+c+dx)}\right)}{4\sqrt{e}\sqrt{f}} - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e}+\sqrt{fx})}{(d\sqrt{e}+(1-c)\sqrt{f})(1+c+dx)}\right)}{4\sqrt{e}\sqrt{f}}$$

output

```
-1/2*(a+b*arctanh(d*x+c))*ln(2*d*(e^(1/2)-f^(1/2)*x)/(d*e^(1/2)-(1-c)*f^(1/2)))/(d*x+c+1)/e^(1/2)/f^(1/2)+1/2*(a+b*arctanh(d*x+c))*ln(2*d*(e^(1/2)+f^(1/2)*x)/(d*e^(1/2)+(1-c)*f^(1/2)))/(d*x+c+1)/e^(1/2)/f^(1/2)+1/4*b*polylog(2,1-2*d*(e^(1/2)-f^(1/2)*x)/(d*e^(1/2)-(1-c)*f^(1/2)))/(d*x+c+1)/e^(1/2)/f^(1/2)-1/4*b*polylog(2,1-2*d*(e^(1/2)+f^(1/2)*x)/(d*e^(1/2)+(1-c)*f^(1/2)))/(d*x+c+1)/e^(1/2)/f^(1/2)
```

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 365, normalized size of antiderivative = 1.34

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e - fx^2} dx$$

$$= \frac{4a \operatorname{arctanh}\left(\frac{\sqrt{fx}}{\sqrt{e}}\right) + b \log(1 - c - dx) \log\left(\frac{d(\sqrt{e} - \sqrt{fx})}{d\sqrt{e} + (-1+c)\sqrt{f}}\right) - b \log(1 + c + dx) \log\left(\frac{d(\sqrt{e} - \sqrt{fx})}{d\sqrt{e} + (1+c)\sqrt{f}}\right) - b \log(1 + c + dx) \log\left(\frac{d(\sqrt{e} + \sqrt{fx})}{d\sqrt{e} - (-1+c)\sqrt{f}}\right) + b \log(1 - c - dx) \log\left(\frac{d(\sqrt{e} + \sqrt{fx})}{d\sqrt{e} - (1+c)\sqrt{f}}\right) - b \operatorname{PolyLog}[2, -((\sqrt{fx})(-1 + c + dx))/(d\sqrt{e} - (-1 + c)\sqrt{f})] + b \operatorname{PolyLog}[2, (\sqrt{fx})(-1 + c + dx)/(d\sqrt{e} + (-1 + c)\sqrt{f})] + b \operatorname{PolyLog}[2, -((\sqrt{fx})(1 + c + dx))/(d\sqrt{e} - (1 + c)\sqrt{f})] - b \operatorname{PolyLog}[2, (\sqrt{fx})(1 + c + dx)/(d\sqrt{e} + (1 + c)\sqrt{f})]}{4\sqrt{e}\sqrt{f}}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])/(e - f*x^2),x]
```

output

```
(4*a*ArcTanh[(Sqrt[f]*x)/Sqrt[e]] + b*Log[1 - c - d*x]*Log[(d*(Sqrt[e] - Sqrt[f]*x))/(d*Sqrt[e] + (-1 + c)*Sqrt[f])] - b*Log[1 + c + d*x]*Log[(d*(Sqrt[e] - Sqrt[f]*x))/(d*Sqrt[e] + (1 + c)*Sqrt[f])] - b*Log[1 - c - d*x]*Log[(d*(Sqrt[e] + Sqrt[f]*x))/(d*Sqrt[e] - (-1 + c)*Sqrt[f])] + b*Log[1 + c + d*x]*Log[(d*(Sqrt[e] + Sqrt[f]*x))/(d*Sqrt[e] - (1 + c)*Sqrt[f])] - b*PolyLog[2, -((Sqrt[f]*(-1 + c + d*x))/(d*Sqrt[e] - (-1 + c)*Sqrt[f]))] + b*PolyLog[2, (Sqrt[f]*(-1 + c + d*x))/(d*Sqrt[e] + (-1 + c)*Sqrt[f])] + b*PolyLog[2, -((Sqrt[f]*(1 + c + d*x))/(d*Sqrt[e] - (1 + c)*Sqrt[f]))] - b*PolyLog[2, (Sqrt[f]*(1 + c + d*x))/(d*Sqrt[e] + (1 + c)*Sqrt[f])])/(4*Sqrt[e]*Sqrt[f])
```

Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 474, normalized size of antiderivative = 1.74, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e - fx^2} dx$$

$$\downarrow 7276$$

$$\int \left(\frac{a}{e - fx^2} + \frac{b \operatorname{arctanh}(c + dx)}{e - fx^2} \right) dx$$

$$\begin{aligned}
& \downarrow 2009 \\
& \frac{a \operatorname{arctanh}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)}{\sqrt{e}\sqrt{f}} + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{f}(-c-dx+1)}{d\sqrt{e}-(1-c)\sqrt{f}}\right)}{4\sqrt{e}\sqrt{f}} - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{f}(-c-dx+1)}{\sqrt{f}(1-c)+d\sqrt{e}}\right)}{4\sqrt{e}\sqrt{f}} + \\
& \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{f}(c+dx+1)}{d\sqrt{e}-(c+1)\sqrt{f}}\right)}{4\sqrt{e}\sqrt{f}} - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{f}(c+dx+1)}{\sqrt{f}(c+1)+d\sqrt{e}}\right)}{4\sqrt{e}\sqrt{f}} + \\
& \frac{b \log(-c-dx+1) \log\left(\frac{d(\sqrt{e}-\sqrt{f}x)}{d\sqrt{e}-(1-c)\sqrt{f}}\right)}{4\sqrt{e}\sqrt{f}} - \frac{b \log(c+dx+1) \log\left(\frac{d(\sqrt{e}-\sqrt{f}x)}{(c+1)\sqrt{f}+d\sqrt{e}}\right)}{4\sqrt{e}\sqrt{f}} - \\
& \frac{b \log(-c-dx+1) \log\left(\frac{d(\sqrt{e}+\sqrt{f}x)}{(1-c)\sqrt{f}+d\sqrt{e}}\right)}{4\sqrt{e}\sqrt{f}} + \frac{b \log(c+dx+1) \log\left(\frac{d(\sqrt{e}+\sqrt{f}x)}{d\sqrt{e}-(c+1)\sqrt{f}}\right)}{4\sqrt{e}\sqrt{f}}
\end{aligned}$$

input `Int[(a + b*ArcTanh[c + d*x])/(e - f*x^2), x]`

output `(a*ArcTanh[(Sqrt[f]*x)/Sqrt[e]]/(Sqrt[e]*Sqrt[f]) + (b*Log[1 - c - d*x]*Log[(d*(Sqrt[e] - Sqrt[f]*x))/(d*Sqrt[e] - (1 - c)*Sqrt[f])])/(4*Sqrt[e]*Sqrt[f]) - (b*Log[1 + c + d*x]*Log[(d*(Sqrt[e] - Sqrt[f]*x))/(d*Sqrt[e] + (1 + c)*Sqrt[f])])/(4*Sqrt[e]*Sqrt[f]) - (b*Log[1 - c - d*x]*Log[(d*(Sqrt[e] + Sqrt[f]*x))/(d*Sqrt[e] + (1 - c)*Sqrt[f])])/(4*Sqrt[e]*Sqrt[f]) + (b*Log[1 + c + d*x]*Log[(d*(Sqrt[e] + Sqrt[f]*x))/(d*Sqrt[e] - (1 + c)*Sqrt[f])])/(4*Sqrt[e]*Sqrt[f]) + (b*PolyLog[2, -(Sqrt[f]*(1 - c - d*x))/(d*Sqrt[e] - (1 - c)*Sqrt[f])])/(4*Sqrt[e]*Sqrt[f]) - (b*PolyLog[2, (Sqrt[f]*(1 - c - d*x))/(d*Sqrt[e] + (1 - c)*Sqrt[f])])/(4*Sqrt[e]*Sqrt[f]) + (b*PolyLog[2, -(Sqrt[f]*(1 + c + d*x))/(d*Sqrt[e] - (1 + c)*Sqrt[f])])/(4*Sqrt[e]*Sqrt[f]) - (b*PolyLog[2, (Sqrt[f]*(1 + c + d*x))/(d*Sqrt[e] + (1 + c)*Sqrt[f])])/(4*Sqrt[e]*Sqrt[f]))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 466 vs. $2(217) = 434$.

Time = 0.62 (sec) , antiderivative size = 467, normalized size of antiderivative = 1.71

| method | result |
|-------------------|---|
| risch | $-\frac{b \ln(-dx-c+1) \ln\left(\frac{d\sqrt{ef}-(-dx-c+1)f-fc+f}{d\sqrt{ef}-fc+f}\right)}{4\sqrt{ef}} + \frac{b \ln(-dx-c+1) \ln\left(\frac{d\sqrt{ef}+(-dx-c+1)f+fc-f}{d\sqrt{ef}+fc-f}\right)}{4\sqrt{ef}} - \frac{b \operatorname{dilog}\left(\frac{d\sqrt{ef}}{d\sqrt{ef}+fc-f}\right)}{4\sqrt{ef}}$ |
| parts | Expression too large to display |
| derivativedivides | Expression too large to display |
| default | Expression too large to display |

input `int((a+b*arctanh(d*x+c))/(-f*x^2+e),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & -1/4*b*\ln(-d*x-c+1)/(e*f)^{(1/2)}*\ln((d*(e*f)^{(1/2)}-(-d*x-c+1)*f-f*c+f)/(d*(e*f)^{(1/2)}-f*c+f))+1/4*b*\ln(-d*x-c+1)/(e*f)^{(1/2)}*\ln((d*(e*f)^{(1/2)}+(-d*x-c+1)*f+f*c-f)/(d*(e*f)^{(1/2)}+f*c-f))-1/4*b/(e*f)^{(1/2)}*\operatorname{dilog}((d*(e*f)^{(1/2)}-(-d*x-c+1)*f-f*c+f)/(d*(e*f)^{(1/2)}-f*c+f))+1/4*b/(e*f)^{(1/2)}*\operatorname{dilog}((d*(e*f)^{(1/2)}+(-d*x-c+1)*f+f*c-f)/(d*(e*f)^{(1/2)}+f*c-f))-a/(e*f)^{(1/2)}*\operatorname{arctanh}(1/2*(2*(-d*x-c+1)*f+2*f*c-2*f)/d/(e*f)^{(1/2)}))-1/4*b*\ln(d*x+c+1)/(e*f)^{(1/2)}*\ln((d*(e*f)^{(1/2)}+f*c-f*(d*x+c+1)+f)/(d*(e*f)^{(1/2)}+f*c+f))+1/4*b*\ln(d*x+c+1)/(e*f)^{(1/2)}*\ln((d*(e*f)^{(1/2)}-f*c+f*(d*x+c+1)-f)/(d*(e*f)^{(1/2)}-f*c-f))-1/4*b/(e*f)^{(1/2)}*\operatorname{dilog}((d*(e*f)^{(1/2)}+f*c-f*(d*x+c+1)+f)/(d*(e*f)^{(1/2)}+f*c+f))+1/4*b/(e*f)^{(1/2)}*\operatorname{dilog}((d*(e*f)^{(1/2)}-f*c+f*(d*x+c+1)-f)/(d*(e*f)^{(1/2)}-f*c-f)) \end{aligned}$$
Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e - fx^2} dx = \int -\frac{b \operatorname{artanh}(dx + c) + a}{fx^2 - e} dx$$

input `integrate((a+b*arctanh(d*x+c))/(-f*x^2+e),x, algorithm="fricas")`

output `integral(-(b*arctanh(d*x + c) + a)/(f*x^2 - e), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e - fx^2} dx = \text{Timed out}$$

input `integrate((a+b*atanh(d*x+c))/(-f*x**2+e),x)`

output Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e - fx^2} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arctanh(d*x+c))/(-f*x^2+e),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e - fx^2} dx = \int -\frac{b \operatorname{arctanh}(dx + c) + a}{fx^2 - e} dx$$

input `integrate((a+b*arctanh(d*x+c))/(-f*x^2+e),x, algorithm="giac")`

output `integrate(-(b*arctanh(d*x + c) + a)/(f*x^2 - e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e - fx^2} dx = \int \frac{a + b \operatorname{atanh}(c + dx)}{e - fx^2} dx$$

input `int((a + b*atanh(c + d*x))/(e - f*x^2),x)`output `int((a + b*atanh(c + d*x))/(e - f*x^2), x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e - fx^2} dx$$

$$= \frac{\sqrt{f} \sqrt{e} \log(-\sqrt{f} \sqrt{e} - fx) a - \sqrt{f} \sqrt{e} \log(\sqrt{f} \sqrt{e} - fx) a + 2 \left(\int \frac{\operatorname{atanh}(dx+c)}{-fx^2+e} dx \right) b e f}{2ef}$$

input `int((a+b*atanh(d*x+c))/(-f*x^2+e),x)`output `(sqrt(f)*sqrt(e)*log(-sqrt(f)*sqrt(e) - f*x)*a - sqrt(f)*sqrt(e)*log(sqrt(f)*sqrt(e) - f*x)*a + 2*int(atanh(c + d*x)/(e - f*x**2),x)*b*e*f)/(2*e*f)`

3.67 $\int \frac{a+b\operatorname{arctanh}(c+dx)}{x(e-fx^2)} dx$

| | |
|----------------------------|-----|
| Optimal result | 585 |
| Mathematica [A] (verified) | 586 |
| Rubi [A] (verified) | 587 |
| Maple [B] (verified) | 588 |
| Fricas [F] | 589 |
| Sympy [F] | 589 |
| Maxima [F] | 590 |
| Giac [F] | 590 |
| Mupad [F(-1)] | 590 |
| Reduce [F] | 591 |

Optimal result

Integrand size = 24, antiderivative size = 310

$$\int \frac{a + b\operatorname{arctanh}(c + dx)}{x(e - fx^2)} dx = \frac{(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2dx}{(1-c)(1+c+dx)}\right)}{e} - \frac{(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2d(\sqrt{e}-\sqrt{fx})}{(d\sqrt{e}-(1-c)\sqrt{f})(1+c+dx)}\right)}{2e} - \frac{(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2d(\sqrt{e}+\sqrt{fx})}{(d\sqrt{e}+(1-c)\sqrt{f})(1+c+dx)}\right)}{2e} - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2dx}{(1-c)(1+c+dx)}\right)}{2e} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e}-\sqrt{fx})}{(d\sqrt{e}-(1-c)\sqrt{f})(1+c+dx)}\right)}{4e} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e}+\sqrt{fx})}{(d\sqrt{e}+(1-c)\sqrt{f})(1+c+dx)}\right)}{4e}$$

output

```
(a+b*arctanh(d*x+c))*ln(2*d*x/(1-c)/(d*x+c+1))/e-1/2*(a+b*arctanh(d*x+c))*
ln(2*d*(e^(1/2)-f^(1/2)*x)/(d*e^(1/2)-(1-c)*f^(1/2))/(d*x+c+1))/e-1/2*(a+b
*arctanh(d*x+c))*ln(2*d*(e^(1/2)+f^(1/2)*x)/(d*e^(1/2)+(1-c)*f^(1/2))/(d*x
+c+1))/e-1/2*b*polylog(2,1-2*d*x/(1-c)/(d*x+c+1))/e+1/4*b*polylog(2,1-2*d*
(e^(1/2)-f^(1/2)*x)/(d*e^(1/2)-(1-c)*f^(1/2))/(d*x+c+1))/e+1/4*b*polylog(2
,1-2*d*(e^(1/2)+f^(1/2)*x)/(d*e^(1/2)+(1-c)*f^(1/2))/(d*x+c+1))/e
```

Mathematica [A] (verified)

Time = 6.53 (sec) , antiderivative size = 594, normalized size of antiderivative = 1.92

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e - fx^2)} dx$$

$$= \frac{4a \log(x) - 2a \log(e - fx^2) + b \left(4 \operatorname{arctanh}(c + dx) \log(x) + \log\left(-\frac{\sqrt{e}}{\sqrt{f}} + x\right) \log(1 - c - dx) + \log\left(\frac{\sqrt{e}}{\sqrt{f}} - x\right) \log(1 - c + dx) \right)}{e}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])/(x*(e - f*x^2)),x]
```

output

```
(4*a*Log[x] - 2*a*Log[e - f*x^2] + b*(4*ArcTanh[c + d*x]*Log[x] + Log[-(Sqrt[e]/Sqrt[f]) + x]*Log[1 - c - d*x] + Log[Sqrt[e]/Sqrt[f] + x]*Log[1 - c - d*x] - Log[Sqrt[e]/Sqrt[f] + x]*Log[-((Sqrt[f]*(-1 + c + d*x))/(d*Sqrt[e] - (-1 + c)*Sqrt[f]))] - Log[-(Sqrt[e]/Sqrt[f]) + x]*Log[(Sqrt[f]*(-1 + c + d*x))/(d*Sqrt[e] + (-1 + c)*Sqrt[f])] - Log[-(Sqrt[e]/Sqrt[f]) + x]*Log[1 + c + d*x] - Log[Sqrt[e]/Sqrt[f] + x]*Log[1 + c + d*x] + Log[Sqrt[e]/Sqrt[f] + x]*Log[-((Sqrt[f]*(1 + c + d*x))/(d*Sqrt[e] - (1 + c)*Sqrt[f]))] + Log[-(Sqrt[e]/Sqrt[f]) + x]*Log[(Sqrt[f]*(1 + c + d*x))/(d*Sqrt[e] + (1 + c)*Sqrt[f])] + 2*Log[x]*Log[1 + (d*x)/(-1 + c)] - 2*Log[x]*Log[1 + (d*x)/(1 + c)] - 2*ArcTanh[c + d*x]*Log[e - f*x^2] - Log[1 - c - d*x]*Log[e - f*x^2] + Log[1 + c + d*x]*Log[e - f*x^2] + 2*PolyLog[2, -((d*x)/(-1 + c))] - 2*PolyLog[2, -((d*x)/(1 + c))] - PolyLog[2, (d*(Sqrt[e]/Sqrt[f] + x))/(1 - c + (d*Sqrt[e])/Sqrt[f])] - PolyLog[2, (d*(Sqrt[e] - Sqrt[f]*x))/(d*Sqrt[e] + (-1 + c)*Sqrt[f])] + PolyLog[2, (d*(Sqrt[e] - Sqrt[f]*x))/(d*Sqrt[e] + (1 + c)*Sqrt[f])] + PolyLog[2, (d*(Sqrt[e] + Sqrt[f]*x))/(d*Sqrt[e] - (1 + c)*Sqrt[f])]))/(4*e)
```

Rubi [A] (verified)

Time = 1.18 (sec) , antiderivative size = 325, normalized size of antiderivative = 1.05, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{arctanh}(c + dx)}{x(e - fx^2)} dx$$

$$\downarrow 7276$$

$$\int \left(\frac{a}{x(e - fx^2)} + \frac{\operatorname{arctanh}(c + dx)}{x(e - fx^2)} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned} & -\frac{a \log(e - fx^2)}{2e} + \frac{a \log(x)}{2e} - \frac{\operatorname{arctanh}(c + dx) \log\left(\frac{2d(\sqrt{e} - \sqrt{fx})}{(c + dx + 1)(d\sqrt{e} - (1 - c)\sqrt{f})}\right)}{2e} - \\ & \frac{\operatorname{arctanh}(c + dx) \log\left(\frac{e}{(c + dx + 1)((1 - c)\sqrt{f} + d\sqrt{e})}\right)}{2e} + \frac{\operatorname{arctanh}(c + dx) \log\left(\frac{2dx}{(1 - c)(c + dx + 1)}\right)}{2e} + \\ & \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e} - \sqrt{fx})}{(d\sqrt{e} - (1 - c)\sqrt{f})(c + dx + 1)}\right)}{4e} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{e}{(\sqrt{f}(1 - c) + d\sqrt{e})(c + dx + 1)}\right)}{4e} - \\ & \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2dx}{(1 - c)(c + dx + 1)}\right)}{2e} \end{aligned}$$

input `Int[(a + b*ArcTanh[c + d*x])/(x*(e - f*x^2)),x]`

output `(a*Log[x])/e + (b*ArcTanh[c + d*x]*Log[(2*d*x)/((1 - c)*(1 + c + d*x))])/e - (b*ArcTanh[c + d*x]*Log[(2*d*(Sqrt[e] - Sqrt[f]*x))/((d*Sqrt[e] - (1 - c)*Sqrt[f])*(1 + c + d*x))])/(2*e) - (b*ArcTanh[c + d*x]*Log[(2*d*(Sqrt[e] + Sqrt[f]*x))/((d*Sqrt[e] + (1 - c)*Sqrt[f])*(1 + c + d*x))])/(2*e) - (a*Log[e - f*x^2])/(2*e) - (b*PolyLog[2, 1 - (2*d*x)/((1 - c)*(1 + c + d*x))])/(2*e) + (b*PolyLog[2, 1 - (2*d*(Sqrt[e] - Sqrt[f]*x))/((d*Sqrt[e] - (1 - c)*Sqrt[f])*(1 + c + d*x))])/(4*e) + (b*PolyLog[2, 1 - (2*d*(Sqrt[e] + Sqrt[f]*x))/((d*Sqrt[e] + (1 - c)*Sqrt[f])*(1 + c + d*x))])/(4*e)`

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 7276 Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xprand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 566 vs. 2(268) = 536.

Time = 0.36 (sec) , antiderivative size = 567, normalized size of antiderivative = 1.83

| method | result |
|-------------------|---|
| risch | $-\frac{b \ln(-dx-c+1) \ln\left(-\frac{xd}{-1+c}\right)}{2e} - \frac{b \operatorname{dilog}\left(-\frac{xd}{-1+c}\right)}{2e} + \frac{b \ln(-dx-c+1) \ln\left(\frac{d\sqrt{ef}-(-dx-c+1)f-fc+f}{d\sqrt{ef}-fc+f}\right)}{4e} + \frac{b \ln(-dx-c+1)}{2e}$ |
| parts | $\frac{a \ln(x)}{e} - \frac{a \ln(fx^2-e)}{2e} + b \left(-\frac{\operatorname{arctanh}(dx+c) \ln\left(c^2f-2cf(dx+c)-e d^2+f(dx+c)^2\right)}{2e} + \frac{\operatorname{arctanh}(dx+c) \ln(dx)}{e} \right)$ |
| derivativedivides | $\frac{a \ln(-dx)}{e} - \frac{a \ln\left(c^2f-2cf(dx+c)-e d^2+f(dx+c)^2\right)}{2e} + b d^2 \left(\frac{\operatorname{arctanh}(dx+c) \ln(-dx)}{e d^2} - \frac{\operatorname{arctanh}(dx+c) \ln\left(c^2f-2cf(dx+c)-e d^2+f(dx+c)^2\right)}{e d^2} \right)$ |
| default | $\frac{a \ln(-dx)}{e} - \frac{a \ln\left(c^2f-2cf(dx+c)-e d^2+f(dx+c)^2\right)}{2e} + b d^2 \left(\frac{\operatorname{arctanh}(dx+c) \ln(-dx)}{e d^2} - \frac{\operatorname{arctanh}(dx+c) \ln\left(c^2f-2cf(dx+c)-e d^2+f(dx+c)^2\right)}{e d^2} \right)$ |

```
input int((a+b*arctanh(d*x+c))/x/(-f*x^2+e), x, method=_RETURNVERBOSE)
```

output

```
-1/2*b/e*ln(-d*x-c+1)*ln(-x*d/(-1+c))-1/2*b/e*dilog(-x*d/(-1+c))+1/4*b/e*ln(-d*x-c+1)*ln((d*(e*f)^(1/2)-(-d*x-c+1)*f-f*c+f)/(d*(e*f)^(1/2)-f*c+f))+1/4*b/e*ln(-d*x-c+1)*ln((d*(e*f)^(1/2)+(-d*x-c+1)*f+f*c-f)/(d*(e*f)^(1/2)+f*c-f))+1/4*b/e*dilog((d*(e*f)^(1/2)-(-d*x-c+1)*f-f*c+f)/(d*(e*f)^(1/2)-f*c+f))+1/4*b/e*dilog((d*(e*f)^(1/2)+(-d*x-c+1)*f+f*c-f)/(d*(e*f)^(1/2)+f*c-f))+a/e*ln(-d*x)-1/2*a/e*ln(f*(-d*x-c+1)^2+2*(-d*x-c+1)*c*f+c^2*f-e*d^2-2*(-d*x-c+1)*f-2*f*c+f)+1/2*b/e*ln(d*x+c+1)*ln(x*d/(-1-c))+1/2*b/e*dilog(x*d/(-1-c))-1/4*b/e*ln(d*x+c+1)*ln((d*(e*f)^(1/2)+f*c-f*(d*x+c+1)+f)/(d*(e*f)^(1/2)+f*c+f))-1/4*b/e*ln(d*x+c+1)*ln((d*(e*f)^(1/2)-f*c+f*(d*x+c+1)-f)/(d*(e*f)^(1/2)-f*c-f))-1/4*b/e*dilog((d*(e*f)^(1/2)+f*c-f*(d*x+c+1)+f)/(d*(e*f)^(1/2)+f*c+f))-1/4*b/e*dilog((d*(e*f)^(1/2)-f*c+f*(d*x+c+1)-f)/(d*(e*f)^(1/2)-f*c-f))
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e - fx^2)} dx = \int -\frac{b \operatorname{artanh}(dx + c) + a}{(fx^2 - e)x} dx$$

input

```
integrate((a+b*arctanh(d*x+c))/x/(-f*x^2+e),x, algorithm="fricas")
```

output

```
integral(-(b*arctanh(d*x + c) + a)/(f*x^3 - e*x), x)
```

Sympy [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e - fx^2)} dx = -\int \frac{a}{-ex + fx^3} dx - \int \frac{b \operatorname{atanh}(c + dx)}{-ex + fx^3} dx$$

input

```
integrate((a+b*atanh(d*x+c))/x/(-f*x**2+e),x)
```

output

```
-Integral(a/(-e*x + f*x**3), x) - Integral(b*atanh(c + d*x)/(-e*x + f*x**3), x)
```

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e - fx^2)} dx = \int -\frac{b \operatorname{artanh}(dx + c) + a}{(fx^2 - e)x} dx$$

input `integrate((a+b*arctanh(d*x+c))/x/(-f*x^2+e),x, algorithm="maxima")`

output `-1/2*a*(log(f*x^2 - e)/e - 2*log(x)/e) - 1/2*b*integrate((log(d*x + c + 1) - log(-d*x - c + 1))/(f*x^3 - e*x), x)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e - fx^2)} dx = \int -\frac{b \operatorname{artanh}(dx + c) + a}{(fx^2 - e)x} dx$$

input `integrate((a+b*arctanh(d*x+c))/x/(-f*x^2+e),x, algorithm="giac")`

output `integrate(-(b*arctanh(d*x + c) + a)/((f*x^2 - e)*x), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e - fx^2)} dx = \int \frac{a + b \operatorname{atanh}(c + dx)}{x(e - fx^2)} dx$$

input `int((a + b*atanh(c + d*x))/(x*(e - f*x^2)),x)`

output `int((a + b*atanh(c + d*x))/(x*(e - f*x^2)), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e - fx^2)} dx$$

$$= \frac{2 \left(\int \frac{a \operatorname{atanh}(dx+c)}{-fx^3+ex} dx \right) b e - \log(-\sqrt{f} \sqrt{e} - fx) a - \log(\sqrt{f} \sqrt{e} - fx) a + 2 \log(x) a}{2e}$$

input `int((a+b*atanh(d*x+c))/x/(-f*x^2+e),x)`

output `(2*int(atanh(c + d*x)/(e*x - f*x**3),x)*b*e - log(-sqrt(f)*sqrt(e) - f*x)*a - log(sqrt(f)*sqrt(e) - f*x)*a + 2*log(x)*a)/(2*e)`

3.68 $\int \frac{a+b\operatorname{arctanh}(c+dx)}{x^2(e-fx^2)} dx$

| | |
|---|-----|
| Optimal result | 592 |
| Mathematica [C] (warning: unable to verify) | 593 |
| Rubi [A] (verified) | 594 |
| Maple [B] (verified) | 596 |
| Fricas [F] | 596 |
| Sympy [F(-1)] | 597 |
| Maxima [F(-2)] | 597 |
| Giac [F] | 598 |
| Mupad [F(-1)] | 598 |
| Reduce [F] | 598 |

Optimal result

Integrand size = 24, antiderivative size = 355

$$\int \frac{a + b\operatorname{arctanh}(c + dx)}{x^2(e - fx^2)} dx = -\frac{a + b\operatorname{arctanh}(c + dx)}{ex} + \frac{bd \log(x)}{(1 - c^2)e}$$

$$- \frac{bd \log(1 - c - dx)}{2(1 - c)e} - \frac{bd \log(1 + c + dx)}{2(1 + c)e}$$

$$- \frac{\sqrt{f}(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2d(\sqrt{e} - \sqrt{fx})}{(d\sqrt{e} - (1 - c)\sqrt{f})(1 + c + dx)}\right)}{2e^{3/2}}$$

$$+ \frac{\sqrt{f}(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2d(\sqrt{e} + \sqrt{fx})}{(d\sqrt{e} + (1 - c)\sqrt{f})(1 + c + dx)}\right)}{2e^{3/2}}$$

$$+ \frac{b\sqrt{f} \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e} - \sqrt{fx})}{(d\sqrt{e} - (1 - c)\sqrt{f})(1 + c + dx)}\right)}{4e^{3/2}}$$

$$- \frac{b\sqrt{f} \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e} + \sqrt{fx})}{(d\sqrt{e} + (1 - c)\sqrt{f})(1 + c + dx)}\right)}{4e^{3/2}}$$

output

```

-(a+b*arctanh(d*x+c))/e/x+b*d*ln(x)/(-c^2+1)/e-1/2*b*d*ln(-d*x-c+1)/(1-c)/
e-1/2*b*d*ln(d*x+c+1)/(1+c)/e-1/2*f^(1/2)*(a+b*arctanh(d*x+c))*ln(2*d*(e^(
1/2)-f^(1/2)*x)/(d*e^(1/2)-(1-c)*f^(1/2)))/(d*x+c+1))/e^(3/2)+1/2*f^(1/2)*(
a+b*arctanh(d*x+c))*ln(2*d*(e^(1/2)+f^(1/2)*x)/(d*e^(1/2)+(1-c)*f^(1/2)))/(
d*x+c+1))/e^(3/2)+1/4*b*f^(1/2)*polylog(2,1-2*d*(e^(1/2)-f^(1/2)*x)/(d*e^(
1/2)-(1-c)*f^(1/2)))/(d*x+c+1))/e^(3/2)-1/4*b*f^(1/2)*polylog(2,1-2*d*(e^(1
/2)+f^(1/2)*x)/(d*e^(1/2)+(1-c)*f^(1/2)))/(d*x+c+1))/e^(3/2)

```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 7.80 (sec) , antiderivative size = 1221, normalized size of antiderivative = 3.44

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e - fx^2)} dx = \text{Too large to display}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])/(x^2*(e - f*x^2)),x]
```

output

```

-(a/(e*x)) + (a*Sqrt[f]*ArcTanh[(Sqrt[f]*x)/Sqrt[e]])/e^(3/2) - (b*((-1 +
c^2 + c*d*x)*ArcTanh[c + d*x] + d*x*Log[-((d*x)/Sqrt[1 - (c + d*x)^2])))/
((-1 + c)*(1 + c)*e*x) - (b*Sqrt[f]*(2*d^2*e*ArcTanh[c - (d*Sqrt[e])/Sqrt[
f]]*ArcTanh[c + d*x] - 2*c^2*f*ArcTanh[c - (d*Sqrt[e])/Sqrt[f]]*ArcTanh[c
+ d*x] - 2*d^2*e*ArcTanh[c + (d*Sqrt[e])/Sqrt[f]]*ArcTanh[c + d*x] + 2*c^2
*f*ArcTanh[c + (d*Sqrt[e])/Sqrt[f]]*ArcTanh[c + d*x] - 2*d*Sqrt[e]*Sqrt[f]
*ArcTanh[c + d*x]^2 + d*Sqrt[e]*E^ArcTanh[c + (d*Sqrt[e])/Sqrt[f]]*Sqrt[1
- c^2 - (d^2*e)/f - (2*c*d*Sqrt[e])/Sqrt[f]]*Sqrt[f]*ArcTanh[c + d*x]^2 +
d*Sqrt[e]*E^ArcTanh[c - (d*Sqrt[e])/Sqrt[f]]*Sqrt[1 - c^2 - (d^2*e)/f + (2
*c*d*Sqrt[e])/Sqrt[f]]*Sqrt[f]*ArcTanh[c + d*x]^2 - c*E^ArcTanh[c + (d*Sqr
t[e])/Sqrt[f]]*Sqrt[1 - c^2 - (d^2*e)/f - (2*c*d*Sqrt[e])/Sqrt[f]]*f*ArcTa
nh[c + d*x]^2 + c*E^ArcTanh[c - (d*Sqrt[e])/Sqrt[f]]*Sqrt[1 - c^2 - (d^2*e
)/f + (2*c*d*Sqrt[e])/Sqrt[f]]*f*ArcTanh[c + d*x]^2 + 2*d^2*e*ArcTanh[c -
(d*Sqrt[e])/Sqrt[f]]*Log[1 - E^(2*ArcTanh[c - (d*Sqrt[e])/Sqrt[f]] - 2*Arc
Tanh[c + d*x])] - 2*c^2*f*ArcTanh[c - (d*Sqrt[e])/Sqrt[f]]*Log[1 - E^(2*Arc
Tanh[c - (d*Sqrt[e])/Sqrt[f]] - 2*ArcTanh[c + d*x])] - 2*d^2*e*ArcTanh[c
+ d*x]*Log[1 - E^(2*ArcTanh[c - (d*Sqrt[e])/Sqrt[f]] - 2*ArcTanh[c + d*x])
] + 2*c^2*f*ArcTanh[c + d*x]*Log[1 - E^(2*ArcTanh[c - (d*Sqrt[e])/Sqrt[f]]
- 2*ArcTanh[c + d*x])] - 2*d^2*e*ArcTanh[c + (d*Sqrt[e])/Sqrt[f]]*Log[1 -
E^(2*ArcTanh[c + (d*Sqrt[e])/Sqrt[f]] - 2*ArcTanh[c + d*x])] + 2*c^2*f...

```

Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 562, normalized size of antiderivative = 1.58, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e - fx^2)} dx$$

$$\downarrow 7276$$

$$\int \left(\frac{a}{x^2 (e - fx^2)} + \frac{b \operatorname{arctanh}(c + dx)}{x^2 (e - fx^2)} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned} & \frac{a\sqrt{f}\operatorname{arctanh}\left(\frac{\sqrt{fx}}{\sqrt{e}}\right)}{e^{3/2}} - \frac{a}{ex} - \frac{b\operatorname{arctanh}(c+dx)}{ex} + \frac{bd\log(x)}{(1-c^2)e} + \frac{b\sqrt{f}\operatorname{PolyLog}\left(2, -\frac{\sqrt{f}(-c-dx+1)}{d\sqrt{e-(1-c)\sqrt{f}}}\right)}{4e^{3/2}} - \\ & \frac{b\sqrt{f}\operatorname{PolyLog}\left(2, \frac{\sqrt{f}(-c-dx+1)}{\sqrt{f}(1-c)+d\sqrt{e}}\right)}{4e^{3/2}} + \frac{b\sqrt{f}\operatorname{PolyLog}\left(2, -\frac{\sqrt{f}(c+dx+1)}{d\sqrt{e}-(c+1)\sqrt{f}}\right)}{4e^{3/2}} - \\ & \frac{b\sqrt{f}\operatorname{PolyLog}\left(2, \frac{\sqrt{f}(c+dx+1)}{\sqrt{f}(c+1)+d\sqrt{e}}\right)}{4e^{3/2}} + \frac{b\sqrt{f}\log(-c-dx+1)\log\left(\frac{d(\sqrt{e}-\sqrt{fx})}{d\sqrt{e}-(1-c)\sqrt{f}}\right)}{4e^{3/2}} - \\ & \frac{b\sqrt{f}\log(c+dx+1)\log\left(\frac{d(\sqrt{e}-\sqrt{fx})}{(c+1)\sqrt{f}+d\sqrt{e}}\right)}{4e^{3/2}} - \frac{b\sqrt{f}\log(-c-dx+1)\log\left(\frac{d(\sqrt{e}+\sqrt{fx})}{(1-c)\sqrt{f}+d\sqrt{e}}\right)}{4e^{3/2}} + \\ & \frac{b\sqrt{f}\log(c+dx+1)\log\left(\frac{d(\sqrt{e}+\sqrt{fx})}{d\sqrt{e}-(c+1)\sqrt{f}}\right)}{4e^{3/2}} - \frac{bd\log(-c-dx+1)}{2(1-c)e} - \frac{bd\log(c+dx+1)}{2(c+1)e} \end{aligned}$$

input

```
Int[(a + b*ArcTanh[c + d*x])/(x^2*(e - f*x^2)),x]
```

output

```
-(a/(e*x)) + (a*Sqrt[f]*ArcTanh[(Sqrt[f]*x)/Sqrt[e]])/e^(3/2) - (b*ArcTanh
[c + d*x])/(e*x) + (b*d*Log[x])/((1 - c^2)*e) - (b*d*Log[1 - c - d*x])/(2*
(1 - c)*e) - (b*d*Log[1 + c + d*x])/(2*(1 + c)*e) + (b*Sqrt[f]*Log[1 - c -
d*x]*Log[(d*(Sqrt[e] - Sqrt[f]*x))/(d*Sqrt[e] - (1 - c)*Sqrt[f])])/(4*e^(
3/2)) - (b*Sqrt[f]*Log[1 + c + d*x]*Log[(d*(Sqrt[e] - Sqrt[f]*x))/(d*Sqrt[
e] + (1 + c)*Sqrt[f])])/(4*e^(3/2)) - (b*Sqrt[f]*Log[1 - c - d*x]*Log[(d*(
Sqrt[e] + Sqrt[f]*x))/(d*Sqrt[e] + (1 - c)*Sqrt[f])])/(4*e^(3/2)) + (b*Sqr
t[f]*Log[1 + c + d*x]*Log[(d*(Sqrt[e] + Sqrt[f]*x))/(d*Sqrt[e] - (1 + c)*S
qrt[f])])/(4*e^(3/2)) + (b*Sqrt[f]*PolyLog[2, -((Sqrt[f]*(1 - c - d*x))/(d
*Sqrt[e] - (1 - c)*Sqrt[f]))])/(4*e^(3/2)) - (b*Sqrt[f]*PolyLog[2, (Sqrt[f]
*(1 - c - d*x))/(d*Sqrt[e] + (1 - c)*Sqrt[f])])/(4*e^(3/2)) + (b*Sqrt[f]*
PolyLog[2, -((Sqrt[f]*(1 + c + d*x))/(d*Sqrt[e] - (1 + c)*Sqrt[f]))])/(4*e
^(3/2)) - (b*Sqrt[f]*PolyLog[2, (Sqrt[f]*(1 + c + d*x))/(d*Sqrt[e] + (1 +
c)*Sqrt[f])])/(4*e^(3/2))
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 677 vs. $2(295) = 590$.

Time = 2.18 (sec) , antiderivative size = 678, normalized size of antiderivative = 1.91

| method | result |
|-------------------|---|
| risch | $-\frac{db \ln(-dx)}{2e(-1+c)} + \frac{db \ln(-dx-c+1)}{2e(-1+c)} + \frac{b \ln(-dx-c+1)c}{2e(-1+c)x} - \frac{b \ln(-dx-c+1)}{2e(-1+c)} - \frac{bf \ln(-dx-c+1) \ln\left(\frac{d\sqrt{ef}-(-dx-c+1)}{d\sqrt{ef}-fc}\right)}{4e\sqrt{ef}}$ |
| parts | Expression too large to display |
| derivativedivides | Expression too large to display |
| default | Expression too large to display |

input `int((a+b*arctanh(d*x+c))/x^2/(-f*x^2+e),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & -1/2*d*b/e/(-1+c)*\ln(-d*x)+1/2*d*b/e*\ln(-d*x-c+1)/(-1+c)+1/2*b/e*\ln(-d*x-c+1)/(-1+c)/x*c-1/2*b/e*\ln(-d*x-c+1)/(-1+c)/x-1/4*b*f/e*\ln(-d*x-c+1)/(e*f)^{(1/2)}*\ln((d*(e*f)^{(1/2)}-(-d*x-c+1)*f-f*c+f)/(d*(e*f)^{(1/2)}-f*c+f))+1/4*b*f/e*\ln(-d*x-c+1)/(e*f)^{(1/2)}*\ln((d*(e*f)^{(1/2)}+(-d*x-c+1)*f+f*c-f)/(d*(e*f)^{(1/2)}+f*c-f))-1/4*b*f/e/(e*f)^{(1/2)}*dilog((d*(e*f)^{(1/2)}-(-d*x-c+1)*f-f*c+f)/(d*(e*f)^{(1/2)}-f*c+f))+1/4*b*f/e/(e*f)^{(1/2)}*dilog((d*(e*f)^{(1/2)}+(-d*x-c+1)*f+f*c-f)/(d*(e*f)^{(1/2)}+f*c-f))-a/e/x-a*f/e/(e*f)^{(1/2)}*arctanh(1/2*(2*(-d*x-c+1)*f+2*f*c-2*f)/d/(e*f)^{(1/2)}+1/2*b*d/e/(1+c)*\ln(d*x)-1/2*b*d*\ln(d*x+c+1)/(1+c)/e-1/2*b/e*\ln(d*x+c+1)/(1+c)/x*c-1/2*b/e*\ln(d*x+c+1)/(1+c)/x-1/4*b*f/e*\ln(d*x+c+1)/(e*f)^{(1/2)}*\ln((d*(e*f)^{(1/2)}+f*c-f*(d*x+c+1)+f)/(d*(e*f)^{(1/2)}+f*c+f))+1/4*b*f/e*\ln(d*x+c+1)/(e*f)^{(1/2)}*\ln((d*(e*f)^{(1/2)}-f*c+f*(d*x+c+1)-f)/(d*(e*f)^{(1/2)}-f*c-f))-1/4*b*f/e/(e*f)^{(1/2)}*dilog((d*(e*f)^{(1/2)}+f*c-f*(d*x+c+1)+f)/(d*(e*f)^{(1/2)}+f*c+f))+1/4*b*f/e/(e*f)^{(1/2)}*dilog((d*(e*f)^{(1/2)}-f*c+f*(d*x+c+1)-f)/(d*(e*f)^{(1/2)}-f*c-f))
 \end{aligned}$$
Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e - fx^2)} dx = \int -\frac{b \operatorname{artanh}(dx + c) + a}{(fx^2 - e)x^2} dx$$

input `integrate((a+b*arctanh(d*x+c))/x^2/(-f*x^2+e),x, algorithm="fricas")`

output `integral(-(b*arctanh(d*x + c) + a)/(f*x^4 - e*x^2), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e - fx^2)} dx = \text{Timed out}$$

input `integrate((a+b*atanh(d*x+c))/x**2/(-f*x**2+e),x)`

output `Timed out`

Maxima [F(-2)]

Exception generated.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e - fx^2)} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arctanh(d*x+c))/x^2/(-f*x^2+e),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e - fx^2)} dx = \int -\frac{b \operatorname{artanh}(dx + c) + a}{(fx^2 - e)x^2} dx$$

input `integrate((a+b*arctanh(d*x+c))/x^2/(-f*x^2+e),x, algorithm="giac")`

output `integrate(-(b*arctanh(d*x + c) + a)/((f*x^2 - e)*x^2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e - fx^2)} dx = \int \frac{a + b \operatorname{atanh}(c + dx)}{x^2 (e - fx^2)} dx$$

input `int((a + b*atanh(c + d*x))/(x^2*(e - f*x^2)),x)`

output `int((a + b*atanh(c + d*x))/(x^2*(e - f*x^2)), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e - fx^2)} dx = \frac{\sqrt{f} \sqrt{e} \log(-\sqrt{f} \sqrt{e} - fx) ax - \sqrt{f} \sqrt{e} \log(\sqrt{f} \sqrt{e} - fx) ax + 2 \left(\int \frac{\operatorname{atanh}(dx+c)}{-fx^2+ex^2} dx \right) b e^2 x - 2ae}{2e^2 x}$$

input `int((a+b*atanh(d*x+c))/x^2/(-f*x^2+e),x)`

output `(sqrt(f)*sqrt(e)*log(-sqrt(f)*sqrt(e)-f*x)*a*x - sqrt(f)*sqrt(e)*log(sqrt(f)*sqrt(e)-f*x)*a*x + 2*int(atanh(c+d*x)/(e*x**2-f*x**4),x)*b*e**2*x - 2*a*e)/(2*e**2*x)`

3.69 $\int \frac{a+b\operatorname{arctanh}(c+dx)}{x^3(e-fx^2)} dx$

| | |
|----------------------------|-----|
| Optimal result | 599 |
| Mathematica [A] (verified) | 600 |
| Rubi [A] (verified) | 601 |
| Maple [A] (verified) | 602 |
| Fricas [F] | 604 |
| Sympy [F(-1)] | 604 |
| Maxima [F] | 605 |
| Giac [F] | 605 |
| Mupad [F(-1)] | 605 |
| Reduce [F] | 606 |

Optimal result

Integrand size = 24, antiderivative size = 428

$$\int \frac{a + b\operatorname{arctanh}(c + dx)}{x^3(e - fx^2)} dx = -\frac{bd}{2(1 - c^2)ex} - \frac{a + b\operatorname{arctanh}(c + dx)}{2ex^2}$$

$$+ \frac{bcd^2 \log(x)}{(1 - c^2)^2 e} - \frac{bd^2 \log(1 - c - dx)}{4(1 - c)^2 e}$$

$$+ \frac{f(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2dx}{(1 - c)(1 + c + dx)}\right)}{e^2}$$

$$+ \frac{bd^2 \log(1 + c + dx)}{4(1 + c)^2 e}$$

$$- \frac{f(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2d(\sqrt{e} - \sqrt{fx})}{(d\sqrt{e} - (1 - c)\sqrt{f})(1 + c + dx)}\right)}{2e^2}$$

$$- \frac{f(a + b\operatorname{arctanh}(c + dx)) \log\left(\frac{2d(\sqrt{e} + \sqrt{fx})}{(d\sqrt{e} + (1 - c)\sqrt{f})(1 + c + dx)}\right)}{2e^2}$$

$$- \frac{bf \operatorname{PolyLog}\left(2, 1 - \frac{2dx}{(1 - c)(1 + c + dx)}\right)}{2e^2}$$

$$+ \frac{bf \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e} - \sqrt{fx})}{(d\sqrt{e} - (1 - c)\sqrt{f})(1 + c + dx)}\right)}{4e^2}$$

$$+ \frac{bf \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e} + \sqrt{fx})}{(d\sqrt{e} + (1 - c)\sqrt{f})(1 + c + dx)}\right)}{4e^2}$$

output

$$\begin{aligned}
& -1/2*b*d/(-c^2+1)/e/x-1/2*(a+b*\operatorname{arctanh}(d*x+c))/e/x^2+b*c*d^2*\ln(x)/(-c^2+1) \\
&)^2/e-1/4*b*d^2*\ln(-d*x-c+1)/(1-c)^2/e+f*(a+b*\operatorname{arctanh}(d*x+c))*\ln(2*d*x/(1- \\
& c)/(d*x+c+1))/e^2+1/4*b*d^2*\ln(d*x+c+1)/(1+c)^2/e-1/2*f*(a+b*\operatorname{arctanh}(d*x+c) \\
&))*\ln(2*d*(e^{(1/2)}-f^{(1/2)}*x)/(d*e^{(1/2)}-(1-c)*f^{(1/2)})/(d*x+c+1))/e^2-1/2 \\
& *f*(a+b*\operatorname{arctanh}(d*x+c))*\ln(2*d*(e^{(1/2)}+f^{(1/2)}*x)/(d*e^{(1/2)}+(1-c)*f^{(1/2)} \\
&))/(d*x+c+1))/e^2-1/2*b*f*\operatorname{polylog}(2,1-2*d*x/(1-c)/(d*x+c+1))/e^2+1/4*b*f*p \\
& \operatorname{olylog}(2,1-2*d*(e^{(1/2)}-f^{(1/2)}*x)/(d*e^{(1/2)}-(1-c)*f^{(1/2)})/(d*x+c+1))/e^ \\
& 2+1/4*b*f*\operatorname{polylog}(2,1-2*d*(e^{(1/2)}+f^{(1/2)}*x)/(d*e^{(1/2)}+(1-c)*f^{(1/2)})/(d \\
& *x+c+1))/e^2
\end{aligned}$$

Mathematica [A] (verified)

Time = 33.03 (sec) , antiderivative size = 608, normalized size of antiderivative = 1.42

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e - fx^2)} dx$$

$$\begin{aligned}
& -\frac{2ae}{x^2} + 4af \log(x) - 2af \log(e - fx^2) + b \left(-\frac{2de}{x-c^2x} + \frac{4cd^2e \log(x)}{(-1+c^2)^2} - \frac{d^2e \log(1-c-dx)}{(-1+c)^2} + \frac{d^2e \log(1+c+dx)}{(1+c)^2} - \frac{2a \operatorname{arctanh}(c+dx)}{x} \right) \\
& = \dots
\end{aligned}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])/(x^3*(e - f*x^2)),x]
```

output

$$\begin{aligned}
& ((-2*a*e)/x^2 + 4*a*f*\operatorname{Log}[x] - 2*a*f*\operatorname{Log}[e - f*x^2] + b*((-2*d*e)/(x - c^2 \\
& *x) + (4*c*d^2*e*\operatorname{Log}[x])/(-1 + c^2)^2 - (d^2*e*\operatorname{Log}[1 - c - d*x])/(-1 + c)^ \\
& 2 + (d^2*e*\operatorname{Log}[1 + c + d*x])/(1 + c)^2 - (2*\operatorname{ArcTanh}[c + d*x]*(e - 2*f*x^2* \\
& \operatorname{Log}[x] + f*x^2*\operatorname{Log}[e - f*x^2]))/x^2 + 2*f*(\operatorname{Log}[x]*(\operatorname{Log}[1 + (d*x)/(-1 + c)] \\
& - \operatorname{Log}[1 + (d*x)/(1 + c)]) + \operatorname{PolyLog}[2, -((d*x)/(-1 + c))] - \operatorname{PolyLog}[2, -(\\
& (d*x)/(1 + c)]) + f*(-(\operatorname{Log}[\operatorname{Sqrt}[e]/\operatorname{Sqrt}[f] + x]*\operatorname{Log}[-((\operatorname{Sqrt}[f]*(-1 + c + \\
& d*x))/(d*\operatorname{Sqrt}[e] - (-1 + c)*\operatorname{Sqrt}[f]))]) - \operatorname{Log}[-(\operatorname{Sqrt}[e]/\operatorname{Sqrt}[f] + x)*\operatorname{Log}[\\
& (\operatorname{Sqrt}[f]*(-1 + c + d*x))/(d*\operatorname{Sqrt}[e] + (-1 + c)*\operatorname{Sqrt}[f])] + \operatorname{Log}[\operatorname{Sqrt}[e]/\operatorname{Sqr \\
& t}[f] + x]*\operatorname{Log}[-((\operatorname{Sqrt}[f]*(1 + c + d*x))/(d*\operatorname{Sqrt}[e] - (1 + c)*\operatorname{Sqrt}[f]))] + \\
& \operatorname{Log}[-(\operatorname{Sqrt}[e]/\operatorname{Sqrt}[f] + x)*\operatorname{Log}[(\operatorname{Sqrt}[f]*(1 + c + d*x))/(d*\operatorname{Sqrt}[e] + (1 + \\
& c)*\operatorname{Sqrt}[f])] + (\operatorname{Log}[1 - c - d*x] - \operatorname{Log}[1 + c + d*x])*(\operatorname{Log}[-(\operatorname{Sqrt}[e]/\operatorname{Sqrt}[f] \\
&) + x] + \operatorname{Log}[\operatorname{Sqrt}[e]/\operatorname{Sqrt}[f] + x] - \operatorname{Log}[e - f*x^2]) - \operatorname{PolyLog}[2, (d*(\operatorname{Sqrt} \\
& [e]/\operatorname{Sqrt}[f] + x))/(1 - c + (d*\operatorname{Sqrt}[e])/ \operatorname{Sqrt}[f])] - \operatorname{PolyLog}[2, (d*(\operatorname{Sqrt}[e] \\
& - \operatorname{Sqrt}[f]*x))/(d*\operatorname{Sqrt}[e] + (-1 + c)*\operatorname{Sqrt}[f])] + \operatorname{PolyLog}[2, (d*(\operatorname{Sqrt}[e] - \operatorname{S} \\
& \operatorname{qrt}[f]*x))/(d*\operatorname{Sqrt}[e] + (1 + c)*\operatorname{Sqrt}[f])] + \operatorname{PolyLog}[2, (d*(\operatorname{Sqrt}[e] + \operatorname{Sqrt}[\\
& f]*x))/(d*\operatorname{Sqrt}[e] - (1 + c)*\operatorname{Sqrt}[f])])))/(4*e^2)
\end{aligned}$$

Rubi [A] (verified)

Time = 1.41 (sec) , antiderivative size = 453, normalized size of antiderivative = 1.06, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e - fx^2)} dx \\
 & \quad \downarrow \text{7276} \\
 & \int \left(\frac{a}{x^3 (e - fx^2)} + \frac{b \operatorname{arctanh}(c + dx)}{x^3 (e - fx^2)} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & -\frac{af \log(e - fx^2)}{2e^2} + \frac{af \log(x)}{e^2} - \frac{a}{2ex^2} + \frac{b \operatorname{arctanh}(c + dx) \log\left(\frac{2dx}{(1-c)(c+dx+1)}\right)}{e^2} - \\
 & \quad \frac{b \operatorname{arctanh}(c + dx) \log\left(\frac{2d(\sqrt{e}-\sqrt{fx})}{(c+dx+1)(d\sqrt{e}-(1-c)\sqrt{f})}\right)}{2e^2} - \\
 & \quad \frac{b \operatorname{arctanh}(c + dx) \log\left(\frac{2d(\sqrt{e}+\sqrt{fx})}{(c+dx+1)((1-c)\sqrt{f}+d\sqrt{e})}\right)}{2e^2} - \frac{b \operatorname{arctanh}(c + dx)}{2ex^2} + \frac{bcd^2 \log(x)}{(1-c^2)^2 e} - \\
 & \quad \frac{bd}{2(1-c^2)ex} - \frac{bd^2 \log(-c - dx + 1)}{4(1-c)^2 e} + \frac{bd^2 \log(c + dx + 1)}{4(c+1)^2 e} - \\
 & \quad \frac{bf \operatorname{PolyLog}\left(2, 1 - \frac{2dx}{(1-c)(c+dx+1)}\right)}{2e^2} + \frac{bf \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{e}-\sqrt{fx})}{(d\sqrt{e}-(1-c)\sqrt{f})(c+dx+1)}\right)}{4e^2} + \\
 & \quad \frac{bf \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt{fx}+\sqrt{e})}{(\sqrt{f}(1-c)+d\sqrt{e})(c+dx+1)}\right)}{4e^2}
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c + d*x])/(x^3*(e - f*x^2)),x]`

output

$$\begin{aligned}
& -1/2*a/(e*x^2) - (b*d)/(2*(1 - c^2)*e*x) - (b*ArcTanh[c + d*x])/(2*e*x^2) \\
& + (b*c*d^2*Log[x])/((1 - c^2)^2*e) + (a*f*Log[x])/e^2 - (b*d^2*Log[1 - c - \\
& d*x])/(4*(1 - c)^2*e) + (b*f*ArcTanh[c + d*x]*Log[(2*d*x)/((1 - c)*(1 + c \\
& + d*x))])/e^2 + (b*d^2*Log[1 + c + d*x])/(4*(1 + c)^2*e) - (b*f*ArcTanh[c \\
& + d*x]*Log[(2*d*(Sqrt[e] - Sqrt[f]*x))/((d*Sqrt[e] - (1 - c)*Sqrt[f])*(1 \\
& + c + d*x))])/ (2*e^2) - (b*f*ArcTanh[c + d*x]*Log[(2*d*(Sqrt[e] + Sqrt[f]* \\
& x))/((d*Sqrt[e] + (1 - c)*Sqrt[f])*(1 + c + d*x))])/ (2*e^2) - (a*f*Log[e - \\
& f*x^2])/ (2*e^2) - (b*f*PolyLog[2, 1 - (2*d*x)/((1 - c)*(1 + c + d*x))])/ (\\
& 2*e^2) + (b*f*PolyLog[2, 1 - (2*d*(Sqrt[e] - Sqrt[f]*x))/((d*Sqrt[e] - (1 \\
& - c)*Sqrt[f])*(1 + c + d*x))])/ (4*e^2) + (b*f*PolyLog[2, 1 - (2*d*(Sqrt[e] \\
& + Sqrt[f]*x))/((d*Sqrt[e] + (1 - c)*Sqrt[f])*(1 + c + d*x))])/ (4*e^2)
\end{aligned}$$

Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \text{ :> Simp}[\text{IntSum}[u, x], x] \text{ /; SumQ}[u]$$

rule 7276

$$\text{Int}[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] \text{ :> With}[\{v = \text{RationalFunctionE}$$

$$\text{xpend}[u/(a + b*x^n), x]\}, \text{Int}[v, x] \text{ /; SumQ}[v]] \text{ /; FreeQ}[\{a, b\}, x] \ \&\& \ \text{IGtQ}$$

$$[n, 0]$$

Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 730, normalized size of antiderivative = 1.71

| method | result |
|-------------------|---|
| parts | $a \left(-\frac{1}{2e x^2} + \frac{f \ln(x)}{e^2} - \frac{f \ln(f x^2 - e)}{2e^2} \right) + b d^2 \left(-\frac{\operatorname{arctanh}(dx+c) f \ln(c^2 f - 2cf(dx+c) - e d^2 + f(dx+c)^2)}{2d^2 e^2} - \dots \right)$ |
| derivativedivides | $d^2 \left(-\frac{a}{2e d^2 x^2} + \frac{af \ln(-dx)}{d^2 e^2} - \frac{af \ln(c^2 f - 2cf(dx+c) - e d^2 + f(dx+c)^2)}{2d^2 e^2} \right) + b d^2 \left(-\frac{\operatorname{arctanh}(dx+c)}{2e d^4 x^2} + \operatorname{arct} \dots \right)$ |
| default | $d^2 \left(-\frac{a}{2e d^2 x^2} + \frac{af \ln(-dx)}{d^2 e^2} - \frac{af \ln(c^2 f - 2cf(dx+c) - e d^2 + f(dx+c)^2)}{2d^2 e^2} \right) + b d^2 \left(-\frac{\operatorname{arctanh}(dx+c)}{2e d^4 x^2} + \operatorname{arct} \dots \right)$ |
| risch | $-\frac{a}{2e x^2} - \frac{db}{4e(-1+c)^2 x} + \frac{bf \operatorname{dilog}\left(\frac{-xd}{-1-c}\right)}{2e^2} - \frac{bf \operatorname{dilog}\left(\frac{d\sqrt{ef}+fc-f(dx+c+1)+f}{d\sqrt{ef}+fc+f}\right)}{4e^2} - \frac{bf \operatorname{dilog}\left(\frac{d\sqrt{ef}-fc+f(dx+c)}{d\sqrt{ef}-fc-f}\right)}{4e^2}$ |

input `int((a+b*arctanh(d*x+c))/x^3/(-f*x^2+e),x,method=_RETURNVERBOSE)`

output

```
a*(-1/2/e/x^2+f/e^2*ln(x)-1/2*f/e^2*ln(f*x^2-e))+b*d^2*(-1/2/d^2*arctanh(d
*x+c)*f/e^2*ln(c^2*f-2*c*f*(d*x+c)-e*d^2+f*(d*x+c)^2)-1/2*arctanh(d*x+c)/e
/x^2/d^2+1/d^2*arctanh(d*x+c)*f/e^2*ln(d*x)+1/2*d^2*(-1/e/d^2*(-1/2/(1+c)^
2*ln(d*x+c+1)-1/(-1+c)/(1+c)/d/x-2*c/(-1+c)^2/(1+c)^2*ln(d*x)+1/2/(-1+c)^2
*ln(d*x+c-1))+2*f/d^4/e^2*(1/2*dilog((d*x+c-1)/(-1+c))+1/2*ln(d*x)*ln((d*x
+c-1)/(-1+c))-1/2*dilog((d*x+c+1)/(1+c))-1/2*ln(d*x)*ln((d*x+c+1)/(1+c)))-
f/d^4/e^2*(-1/2*ln(d*x+c+1)*ln(c^2*f-2*c*f*(d*x+c)-e*d^2+f*(d*x+c)^2)+f*(1
/2*ln(d*x+c+1)*(ln((d*(e*f)^(1/2)+f*c-f*(d*x+c+1)+f)/(d*(e*f)^(1/2)+f*c+f)
)+ln((d*(e*f)^(1/2)-f*c+f*(d*x+c+1)-f)/(d*(e*f)^(1/2)-f*c-f)))/f+1/2*(dilo
g((d*(e*f)^(1/2)+f*c-f*(d*x+c+1)+f)/(d*(e*f)^(1/2)+f*c+f))+dilog((d*(e*f)^(
1/2)-f*c+f*(d*x+c+1)-f)/(d*(e*f)^(1/2)-f*c-f)))/f)+1/2*ln(d*x+c-1)*ln(c^2
*f-2*c*f*(d*x+c)-e*d^2+f*(d*x+c)^2)-f*(1/2*ln(d*x+c-1)*(ln((d*(e*f)^(1/2)+
f*c-f*(d*x+c-1)-f)/(d*(e*f)^(1/2)+f*c-f))+ln((d*(e*f)^(1/2)-f*c+f*(d*x+c-1
)+f)/(d*(e*f)^(1/2)-f*c+f)))/f+1/2*(dilog((d*(e*f)^(1/2)+f*c-f*(d*x+c-1)-f
)/(d*(e*f)^(1/2)+f*c-f))+dilog((d*(e*f)^(1/2)-f*c+f*(d*x+c-1)+f)/(d*(e*f)^(
1/2)-f*c+f)))/f))))
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e - fx^2)} dx = \int -\frac{b \operatorname{arctanh}(dx + c) + a}{(fx^2 - e)x^3} dx$$

input

```
integrate((a+b*arctanh(d*x+c))/x^3/(-f*x^2+e),x, algorithm="fricas")
```

output

```
integral(-(b*arctanh(d*x + c) + a)/(f*x^5 - e*x^3), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e - fx^2)} dx = \text{Timed out}$$

input

```
integrate((a+b*atanh(d*x+c))/x**3/(-f*x**2+e),x)
```

output Timed out

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e - fx^2)} dx = \int -\frac{b \operatorname{artanh}(dx + c) + a}{(fx^2 - e)x^3} dx$$

input `integrate((a+b*arctanh(d*x+c))/x^3/(-f*x^2+e),x, algorithm="maxima")`

output `-1/2*a*(f*log(f*x^2 - e)/e^2 - 2*f*log(x)/e^2 + 1/(e*x^2)) - 1/2*b*integrate((log(d*x + c + 1) - log(-d*x - c + 1))/(f*x^5 - e*x^3), x)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e - fx^2)} dx = \int -\frac{b \operatorname{artanh}(dx + c) + a}{(fx^2 - e)x^3} dx$$

input `integrate((a+b*arctanh(d*x+c))/x^3/(-f*x^2+e),x, algorithm="giac")`

output `integrate(-(b*arctanh(d*x + c) + a)/((f*x^2 - e)*x^3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e - fx^2)} dx = \int \frac{a + b \operatorname{atanh}(c + dx)}{x^3 (e - fx^2)} dx$$

input `int((a + b*atanh(c + d*x))/(x^3*(e - f*x^2)),x)`

output `int((a + b*atanh(c + d*x))/(x^3*(e - f*x^2)), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e - fx^2)} dx$$

$$= \frac{2 \left(\int \frac{\operatorname{atanh}(dx+c)}{-fx^5+ex^3} dx \right) b e^2 x^2 - \log(-\sqrt{f} \sqrt{e} - fx) a f x^2 - \log(\sqrt{f} \sqrt{e} - fx) a f x^2 + 2 \log(x) a f x^2 - a e}{2e^2 x^2}$$

input `int((a+b*atanh(d*x+c))/x^3/(-f*x^2+e),x)`

output `(2*int(atanh(c + d*x)/(e*x**3 - f*x**5),x)*b*e**2*x**2 - log(- sqrt(f)*sqrt(e) - f*x)*a*f*x**2 - log(sqrt(f)*sqrt(e) - f*x)*a*f*x**2 + 2*log(x)*a*f*x**2 - a*e)/(2*e**2*x**2)`

$$3.70 \quad \int \frac{x^3(a+b \operatorname{arctanh}(c+dx))}{e+fx^3} dx$$

| | |
|---------------------------------------|-----|
| Optimal result | 608 |
| Mathematica [F] | 609 |
| Rubi [A] (warning: unable to verify) | 609 |
| Maple [C] (warning: unable to verify) | 612 |
| Fricas [F] | 614 |
| Sympy [F(-1)] | 614 |
| Maxima [F(-2)] | 615 |
| Giac [F] | 615 |
| Mupad [F(-1)] | 615 |
| Reduce [F] | 616 |

Optimal result

Integrand size = 23, antiderivative size = 727

$$\begin{aligned}
& \int \frac{x^3(a + \operatorname{barctanh}(c + dx))}{e + fx^3} dx \\
&= \frac{ax}{f} + \frac{b(c + dx)\operatorname{arctanh}(c + dx)}{df} + \frac{\sqrt[3]{e}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{3f^{4/3}} \\
&\quad - \frac{\sqrt[3]{-1}\sqrt[3]{e}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{3f^{4/3}} \\
&\quad + \frac{(-1)^{2/3}\sqrt[3]{e}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{3f^{4/3}} \\
&\quad - \frac{\sqrt[3]{e}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d(\sqrt[3]{e} + \sqrt[3]{fx})}{(d\sqrt[3]{e} + (1-c)\sqrt[3]{f})(1+c+dx)}\right)}{3f^{4/3}} \\
&\quad - \frac{(-1)^{2/3}\sqrt[3]{e}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx})}{(d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f})(1+c+dx)}\right)}{3f^{4/3}} \\
&\quad + \frac{\sqrt[3]{-1}\sqrt[3]{e}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx})}{(d\sqrt[3]{e} + (-1)^{2/3}(1-c)\sqrt[3]{f})(1+c+dx)}\right)}{3f^{4/3}} \\
&\quad + \frac{b \log(1 - (c + dx)^2)}{2df} - \frac{b\sqrt[3]{e} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{6f^{4/3}} \\
&\quad + \frac{\sqrt[3]{-1}b\sqrt[3]{e} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{6f^{4/3}} - \frac{(-1)^{2/3}b\sqrt[3]{e} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{6f^{4/3}} \\
&\quad + \frac{b\sqrt[3]{e} \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt[3]{e} + \sqrt[3]{fx})}{(d\sqrt[3]{e} + (1-c)\sqrt[3]{f})(1+c+dx)}\right)}{6f^{4/3}} \\
&\quad + \frac{(-1)^{2/3}b\sqrt[3]{e} \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx})}{(d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f})(1+c+dx)}\right)}{6f^{4/3}} \\
&\quad - \frac{\sqrt[3]{-1}b\sqrt[3]{e} \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx})}{(d\sqrt[3]{e} + (-1)^{2/3}(1-c)\sqrt[3]{f})(1+c+dx)}\right)}{6f^{4/3}}
\end{aligned}$$

output

```
a*x/f+b*(d*x+c)*arctanh(d*x+c)/d/f+1/3*e^(1/3)*(a+b*arctanh(d*x+c))*ln(2/(
d*x+c+1))/f^(4/3)-1/3*(-1)^(1/3)*e^(1/3)*(a+b*arctanh(d*x+c))*ln(2/(d*x+c+
1))/f^(4/3)+1/3*(-1)^(2/3)*e^(1/3)*(a+b*arctanh(d*x+c))*ln(2/(d*x+c+1))/f^
(4/3)-1/3*e^(1/3)*(a+b*arctanh(d*x+c))*ln(2*d*(e^(1/3)+f^(1/3)*x)/(d*e^(1/
3)+(1-c)*f^(1/3))/(d*x+c+1))/f^(4/3)-1/3*(-1)^(2/3)*e^(1/3)*(a+b*arctanh(d
*x+c))*ln(2*d*(e^(1/3)-(-1)^(1/3)*f^(1/3)*x)/(d*e^(1/3)-(-1)^(1/3)*(1-c)*f
^(1/3))/(d*x+c+1))/f^(4/3)+1/3*(-1)^(1/3)*e^(1/3)*(a+b*arctanh(d*x+c))*ln(
2*d*(e^(1/3)+(-1)^(2/3)*f^(1/3)*x)/(d*e^(1/3)+(-1)^(2/3)*(1-c)*f^(1/3))/(d
*x+c+1))/f^(4/3)+1/2*b*ln(1-(d*x+c)^2)/d/f-1/6*b*e^(1/3)*polylog(2,1-2/(d*
x+c+1))/f^(4/3)+1/6*(-1)^(1/3)*b*e^(1/3)*polylog(2,1-2/(d*x+c+1))/f^(4/3)-
1/6*(-1)^(2/3)*b*e^(1/3)*polylog(2,1-2/(d*x+c+1))/f^(4/3)+1/6*b*e^(1/3)*po
lylog(2,1-2*d*(e^(1/3)+f^(1/3)*x)/(d*e^(1/3)+(1-c)*f^(1/3))/(d*x+c+1))/f^(
4/3)+1/6*(-1)^(2/3)*b*e^(1/3)*polylog(2,1-2*d*(e^(1/3)-(-1)^(1/3)*f^(1/3)*
x)/(d*e^(1/3)-(-1)^(1/3)*(1-c)*f^(1/3))/(d*x+c+1))/f^(4/3)-1/6*(-1)^(1/3)*
b*e^(1/3)*polylog(2,1-2*d*(e^(1/3)+(-1)^(2/3)*f^(1/3)*x)/(d*e^(1/3)+(-1)^(
2/3)*(1-c)*f^(1/3))/(d*x+c+1))/f^(4/3)
```

Mathematica [F]

$$\int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx$$

input

```
Integrate[(x^3*(a + b*ArcTanh[c + d*x]))/(e + f*x^3),x]
```

output

```
Integrate[(x^3*(a + b*ArcTanh[c + d*x]))/(e + f*x^3), x]
```

Rubi [A] (warning: unable to verify)

Time = 2.17 (sec) , antiderivative size = 956, normalized size of antiderivative = 1.31, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx$$

↓ 7276

$$\int \left(\frac{ax^3}{e + fx^3} + \frac{bx^3 \operatorname{arctanh}(c + dx)}{e + fx^3} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{ax}{f} + \frac{a\sqrt[3]{e} \arctan\left(\frac{\sqrt[3]{e}-2\sqrt[3]{fx}}{\sqrt{3}\sqrt[3]{e}}\right)}{\sqrt{3}f^{4/3}} + \frac{b(c+dx)\operatorname{arctanh}(c+dx)}{df} - \frac{a\sqrt[3]{e} \log\left(\sqrt[3]{fx} + \sqrt[3]{e}\right)}{3f^{4/3}} + \\
& \frac{b\sqrt[3]{e} \log(-c-dx+1) \log\left(\frac{d\left(\sqrt[3]{fx} + \sqrt[3]{e}\right)}{\sqrt[3]{f(1-c)} + d\sqrt[3]{e}}\right)}{6f^{4/3}} - \frac{b\sqrt[3]{e} \log(c+dx+1) \log\left(\frac{d\left(\sqrt[3]{fx} + \sqrt[3]{e}\right)}{d\sqrt[3]{e} - (c+1)\sqrt[3]{f}}\right)}{6f^{4/3}} + \\
& \frac{(-1)^{2/3} b\sqrt[3]{e} \log(-c-dx+1) \log\left(\frac{d\left(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx}\right)}{d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f}}\right)}{6f^{4/3}} - \\
& \frac{(-1)^{2/3} b\sqrt[3]{e} \log(c+dx+1) \log\left(\frac{d\left(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx}\right)}{\sqrt[3]{-1}\sqrt[3]{f(c+1)} + d\sqrt[3]{e}}\right)}{6f^{4/3}} - \\
& \frac{\sqrt[3]{-1} b\sqrt[3]{e} \log(-c-dx+1) \log\left(\frac{d\left((-1)^{2/3}\sqrt[3]{fx} + \sqrt[3]{e}\right)}{(-1)^{2/3}\sqrt[3]{f(1-c)} + d\sqrt[3]{e}}\right)}{6f^{4/3}} + \\
& \frac{\sqrt[3]{-1} b\sqrt[3]{e} \log(c+dx+1) \log\left(\frac{d\left((-1)^{2/3}\sqrt[3]{fx} + \sqrt[3]{e}\right)}{d\sqrt[3]{e} - (-1)^{2/3}(c+1)\sqrt[3]{f}}\right)}{6f^{4/3}} + \\
& \frac{a\sqrt[3]{e} \log\left(f^{2/3}x^2 - \sqrt[3]{e}\sqrt[3]{fx} + e^{2/3}\right)}{6f^{4/3}} + \frac{b \log(1 - (c+dx)^2)}{2df} + \\
& \frac{b\sqrt[3]{e} \operatorname{PolyLog}\left(2, \frac{\sqrt[3]{f}(-c-dx+1)}{\sqrt[3]{f(1-c)} + d\sqrt[3]{e}}\right)}{6f^{4/3}} + \frac{(-1)^{2/3} b\sqrt[3]{e} \operatorname{PolyLog}\left(2, -\frac{\sqrt[3]{-1}\sqrt[3]{f}(-c-dx+1)}{d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f}}\right)}{6f^{4/3}} - \\
& \frac{\sqrt[3]{-1} b\sqrt[3]{e} \operatorname{PolyLog}\left(2, \frac{(-1)^{2/3}\sqrt[3]{f}(-c-dx+1)}{(-1)^{2/3}\sqrt[3]{f(1-c)} + d\sqrt[3]{e}}\right)}{6f^{4/3}} - \frac{b\sqrt[3]{e} \operatorname{PolyLog}\left(2, -\frac{\sqrt[3]{f}(c+dx+1)}{d\sqrt[3]{e} - (c+1)\sqrt[3]{f}}\right)}{6f^{4/3}} - \\
& \frac{(-1)^{2/3} b\sqrt[3]{e} \operatorname{PolyLog}\left(2, \frac{\sqrt[3]{-1}\sqrt[3]{f}(c+dx+1)}{\sqrt[3]{-1}\sqrt[3]{f(c+1)} + d\sqrt[3]{e}}\right)}{6f^{4/3}} + \\
& \frac{\sqrt[3]{-1} b\sqrt[3]{e} \operatorname{PolyLog}\left(2, -\frac{(-1)^{2/3}\sqrt[3]{f}(c+dx+1)}{d\sqrt[3]{e} - (-1)^{2/3}(c+1)\sqrt[3]{f}}\right)}{6f^{4/3}}
\end{aligned}$$

input

```
Int[(x^3*(a + b*ArcTanh[c + d*x]))/(e + f*x^3),x]
```

output

$$\begin{aligned} & (a*x)/f + (a*e^{1/3}*ArcTan[(e^{1/3} - 2*f^{1/3}*x)/(Sqrt[3]*e^{1/3})])/(Sqrt[3]*f^{4/3}) + (b*(c + d*x)*ArcTanh[c + d*x])/(d*f) - (a*e^{1/3}*Log[e^{1/3} + f^{1/3}*x])/(3*f^{4/3}) + (b*e^{1/3}*Log[1 - c - d*x]*Log[(d*(e^{1/3} + f^{1/3}*x))/(d*e^{1/3} + (1 - c)*f^{1/3})])/(6*f^{4/3}) - (b*e^{1/3} \\ & *Log[1 + c + d*x]*Log[(d*(e^{1/3} + f^{1/3}*x))/(d*e^{1/3} - (1 + c)*f^{1/3})])/(6*f^{4/3}) + ((-1)^{2/3}*b*e^{1/3}*Log[1 - c - d*x]*Log[(d*(e^{1/3} - (-1)^{1/3}*f^{1/3}*x))/(d*e^{1/3} - (-1)^{1/3}*(1 - c)*f^{1/3})])/(6*f^{4/3}) - ((-1)^{2/3}*b*e^{1/3}*Log[1 + c + d*x]*Log[(d*(e^{1/3} - (-1)^{1/3}*f^{1/3}*x))/(d*e^{1/3} + (-1)^{1/3}*(1 + c)*f^{1/3})])/(6*f^{4/3}) - ((-1)^{1/3}*b*e^{1/3}*Log[1 - c - d*x]*Log[(d*(e^{1/3} + (-1)^{2/3}*f^{1/3}*x))/(d*e^{1/3} + (-1)^{2/3}*(1 - c)*f^{1/3})])/(6*f^{4/3}) + ((-1)^{1/3}*b \\ & *e^{1/3}*Log[1 + c + d*x]*Log[(d*(e^{1/3} + (-1)^{2/3}*f^{1/3}*x))/(d*e^{1/3} - (-1)^{2/3}*(1 + c)*f^{1/3})])/(6*f^{4/3}) + (a*e^{1/3}*Log[e^{2/3} - e^{1/3}*f^{1/3}*x + f^{2/3}*x^2])/(6*f^{4/3}) + (b*Log[1 - (c + d*x)^2])/(2*d*f) + (b*e^{1/3}*PolyLog[2, (f^{1/3}*(1 - c - d*x))/(d*e^{1/3} + (1 - c)*f^{1/3})])/(6*f^{4/3}) + ((-1)^{2/3}*b*e^{1/3}*PolyLog[2, -(((-1)^{1/3})*f^{1/3}*(1 - c - d*x))/(d*e^{1/3} - (-1)^{1/3}*(1 - c)*f^{1/3})])/(6*f^{4/3}) - ((-1)^{1/3}*b*e^{1/3}*PolyLog[2, ((-1)^{2/3}*f^{1/3}*(1 - c - d*x))/(d*e^{1/3} + (-1)^{2/3}*(1 - c)*f^{1/3})])/(6*f^{4/3}) - (b*e^{1/3}*PolyLog[2, -((f^{1/3}*(1 + c + d*x))/(d*e^{1/3} - (1 + c)*f^{1/3})])/(6*f^{4/3}) + \dots \end{aligned}$$

Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

rule 7276

$$\text{Int}[(u_)/((a_) + (b_)*(x_)^{(n_)}), x_Symbol] \rightarrow \text{With}[\{v = \text{RationalFunctionExpand}[u/(a + b*x^n), x]\}, \text{Int}[v, x] /; \text{SumQ}[v]] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{IGtQ}[n, 0]$$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.99 (sec) , antiderivative size = 505, normalized size of antiderivative = 0.69

output

```
-1/2*b/f*ln(-d*x-c+1)*x-1/2/d*b/f*ln(-d*x-c+1)*c+1/2/d*b/f*ln(-d*x-c+1)-1/
d*b/f+1/6*d^2*b*e/f^2*sum(1/(_R1^2+2*_R1*c+c^2-2*_R1-2*c+1)*(ln(-d*x-c+1)*
ln((d*x+_R1+c-1)/_R1)+dilog((d*x+_R1+c-1)/_R1)),_R1=RootOf(f*_Z^3+(3*c*f-3
*f)*_Z^2+(3*c^2*f-6*c*f+3*f)*_Z+c^3*f-d^3*e-3*c^2*f+3*f*c-f))+a*x/f+1/d*a/
f*c-1/d*a/f-1/3*d^2*a/f^2*sum(1/(_R^2+2*_R*c+c^2-2*_R-2*c+1)*ln(-d*x-_R-c+
1),_R=RootOf(f*_Z^3+(3*c*f-3*f)*_Z^2+(3*c^2*f-6*c*f+3*f)*_Z+c^3*f-d^3*e-3*
c^2*f+3*f*c-f))*e+1/2*b/f*ln(d*x+c+1)*x+1/2*b/d/f*ln(d*x+c+1)*c+1/2*b/d/f*
ln(d*x+c+1)-1/6*b*d^2*e/f^2*sum(1/(_R1^2-2*_R1*c+c^2-2*_R1+2*c+1)*(ln(d*x+
c+1)*ln((-d*x+_R1-c-1)/_R1)+dilog((-d*x+_R1-c-1)/_R1)),_R1=RootOf(f*_Z^3+(
-3*c*f-3*f)*_Z^2+(3*c^2*f+6*c*f+3*f)*_Z-c^3*f+d^3*e-3*c^2*f-3*f*c-f))
```

Fricas [F]

$$\int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)x^3}{fx^3 + e} dx$$

input

```
integrate(x^3*(a+b*arctanh(d*x+c))/(f*x^3+e),x, algorithm="fricas")
```

output

```
integral((b*x^3*arctanh(d*x + c) + a*x^3)/(f*x^3 + e), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \text{Timed out}$$

input

```
integrate(x**3*(a+b*atanh(d*x+c))/(f*x**3+e),x)
```

output

```
Timed out
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \text{Exception raised: ValueError}$$

input `integrate(x^3*(a+b*arctanh(d*x+c))/(f*x^3+e),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

Giac [F]

$$\int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \int \frac{(b \operatorname{arctanh}(dx + c) + a)x^3}{fx^3 + e} dx$$

input `integrate(x^3*(a+b*arctanh(d*x+c))/(f*x^3+e),x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)*x^3/(f*x^3 + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x^3(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \int \frac{x^3(a + b \operatorname{atanh}(c + dx))}{fx^3 + e} dx$$

input `int((x^3*(a + b*atanh(c + d*x)))/(e + f*x^3),x)`

output `int((x^3*(a + b*atanh(c + d*x)))/(e + f*x^3), x)`

Reduce [F]

$$\int \frac{x^3(a + \operatorname{arctanh}(c + dx))}{e + fx^3} dx$$

$$= \frac{2e^{\frac{1}{3}}\sqrt{3} \operatorname{atan}\left(\frac{e^{\frac{1}{3}} - 2f^{\frac{1}{3}}x}{e^{\frac{1}{3}}\sqrt{3}}\right) a + e^{\frac{1}{3}}\log\left(e^{\frac{2}{3}} - f^{\frac{1}{3}}e^{\frac{1}{3}}x + f^{\frac{2}{3}}x^2\right) a - 2e^{\frac{1}{3}}\log\left(e^{\frac{1}{3}} + f^{\frac{1}{3}}x\right) a + 6f^{\frac{4}{3}}\left(\int \frac{\operatorname{atanh}(dx+c)x^3}{fx^3+e}\right)}{6f^{\frac{4}{3}}}$$

input

```
int(x^3*(a+b*atanh(d*x+c))/(f*x^3+e),x)
```

output

```
(2*e**(1/3)*sqrt(3)*atan((e**(1/3) - 2*f**(1/3)*x)/(e**(1/3)*sqrt(3)))*a +
e**(1/3)*log(e**(2/3) - f**(1/3)*e**(1/3)*x + f**(2/3)*x**2)*a - 2*e**(1/
3)*log(e**(1/3) + f**(1/3)*x)*a + 6*f**(1/3)*int((atanh(c + d*x)*x**3)/(e
+ f*x**3),x)*b*f + 6*f**(1/3)*a*x)/(6*f**(1/3)*f)
```

$$3.71 \quad \int \frac{x^2(a+b \operatorname{arctanh}(c+dx))}{e+fx^3} dx$$

| | |
|---|-----|
| Optimal result | 618 |
| Mathematica [A] (warning: unable to verify) | 620 |
| Rubi [A] (verified) | 621 |
| Maple [C] (warning: unable to verify) | 623 |
| Fricas [F] | 624 |
| Sympy [F(-1)] | 624 |
| Maxima [F] | 625 |
| Giac [F] | 625 |
| Mupad [F(-1)] | 625 |
| Reduce [F] | 626 |

Optimal result

Integrand size = 23, antiderivative size = 463

$$\begin{aligned}
& \int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx \\
&= -\frac{(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{f} \\
&\quad + \frac{(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2d\left(\sqrt[3]{e} + \sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} + (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{3f} \\
&\quad + \frac{(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2d\left((-1)^{2/3}\sqrt[3]{e} + \sqrt[3]{fx}\right)}{\left((-1)^{2/3}d\sqrt[3]{e} + (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{3f} \\
&\quad + \frac{(a + b \operatorname{arctanh}(c + dx)) \log\left(\frac{2\sqrt[3]{-1}d\left(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx}\right)}{\left(\sqrt[3]{-1}d\sqrt[3]{e} - (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{3f} \\
&\quad + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{2f} - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d\left(\sqrt[3]{e} + \sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} + (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{6f} \\
&\quad - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d\left((-1)^{2/3}\sqrt[3]{e} + \sqrt[3]{fx}\right)}{\left((-1)^{2/3}d\sqrt[3]{e} + (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{6f} \\
&\quad - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt[3]{-1}d\left(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx}\right)}{\left(\sqrt[3]{-1}d\sqrt[3]{e} - (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{6f}
\end{aligned}$$

output

```

-(a+b*arctanh(d*x+c))*ln(2/(d*x+c+1))/f+1/3*(a+b*arctanh(d*x+c))*ln(2*d*(e
^(1/3)+f^(1/3)*x)/(d*e^(1/3)+(1-c)*f^(1/3))/(d*x+c+1))/f+1/3*(a+b*arctanh(
d*x+c))*ln(2*d*((-1)^(2/3)*e^(1/3)+f^(1/3)*x)/((-1)^(2/3)*d*e^(1/3)+(1-c)*
f^(1/3))/(d*x+c+1))/f+1/3*(a+b*arctanh(d*x+c))*ln(2*(-1)^(1/3)*d*(e^(1/3)+
(-1)^(2/3)*f^(1/3)*x)/((-1)^(1/3)*d*e^(1/3)-(1-c)*f^(1/3))/(d*x+c+1))/f+1/
2*b*polylog(2,1-2/(d*x+c+1))/f-1/6*b*polylog(2,1-2*d*(e^(1/3)+f^(1/3)*x)/(
d*e^(1/3)+(1-c)*f^(1/3))/(d*x+c+1))/f-1/6*b*polylog(2,1-2*d*((-1)^(2/3)*e^(
1/3)+f^(1/3)*x)/((-1)^(2/3)*d*e^(1/3)+(1-c)*f^(1/3))/(d*x+c+1))/f-1/6*b*p
olylog(2,1-2*(-1)^(1/3)*d*(e^(1/3)+(-1)^(2/3)*f^(1/3)*x)/((-1)^(1/3)*d*e^(
1/3)-(1-c)*f^(1/3))/(d*x+c+1))/f

```

Mathematica [A] (warning: unable to verify)

Time = 0.38 (sec) , antiderivative size = 679, normalized size of antiderivative = 1.47

$$\begin{aligned}
\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = & - \frac{b \log(1 - c - dx) \log\left(\frac{d(\sqrt[3]{e} + \sqrt[3]{fx})}{d\sqrt[3]{e} + (1-c)\sqrt[3]{f}}\right)}{6f} \\
& + \frac{b \log(1 + c + dx) \log\left(\frac{d(\sqrt[3]{e} + \sqrt[3]{fx})}{d\sqrt[3]{e} - (1+c)\sqrt[3]{f}}\right)}{6f} \\
& - \frac{b \log(1 - c - dx) \log\left(\frac{d((-1)^{2/3}\sqrt[3]{e} + \sqrt[3]{fx})}{(-1)^{2/3}d\sqrt[3]{e} + (1-c)\sqrt[3]{f}}\right)}{6f} \\
& + \frac{b \log(1 + c + dx) \log\left(\frac{(-1)^{2/3}d(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx})}{(-1)^{2/3}d\sqrt[3]{e} - (1+c)\sqrt[3]{f}}\right)}{6f} \\
& - \frac{b \log(1 - c - dx) \log\left(\frac{\sqrt[3]{-1}d(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx})}{\sqrt[3]{-1}d\sqrt[3]{e} - (1-c)\sqrt[3]{f}}\right)}{6f} \\
& + \frac{b \log(1 + c + dx) \log\left(\frac{\sqrt[3]{-1}d(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx})}{\sqrt[3]{-1}d\sqrt[3]{e} + (1+c)\sqrt[3]{f}}\right)}{6f} \\
& + \frac{a \log(e + fx^3)}{3f} \\
& - \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt[3]{f}(1-c-dx)}{\sqrt[3]{-1}d\sqrt[3]{e} - (1-c)\sqrt[3]{f}}\right)}{6f} \\
& - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt[3]{f}(1-c-dx)}{d\sqrt[3]{e} + (1-c)\sqrt[3]{f}}\right)}{6f} \\
& - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt[3]{f}(1-c-dx)}{(-1)^{2/3}d\sqrt[3]{e} + (1-c)\sqrt[3]{f}}\right)}{6f} \\
& + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt[3]{f}(1+c+dx)}{d\sqrt[3]{e} - (1+c)\sqrt[3]{f}}\right)}{6f} \\
& + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt[3]{f}(1+c+dx)}{(-1)^{2/3}d\sqrt[3]{e} - (1+c)\sqrt[3]{f}}\right)}{6f}
\end{aligned}$$

input `Integrate[(x^2*(a + b*ArcTanh[c + d*x]))/(e + f*x^3),x]`

output

```
-1/6*(b*Log[1 - c - d*x]*Log[(d*(e^(1/3) + f^(1/3)*x))/(d*e^(1/3) + (1 - c)*f^(1/3))]/f + (b*Log[1 + c + d*x]*Log[(d*(e^(1/3) + f^(1/3)*x))/(d*e^(1/3) - (1 + c)*f^(1/3))]/(6*f) - (b*Log[1 - c - d*x]*Log[(d*(-1)^(2/3)*e^(1/3) + f^(1/3)*x)/((-1)^(2/3)*d*e^(1/3) + (1 - c)*f^(1/3))]/(6*f) + (b*Log[1 + c + d*x]*Log[(-1)^(2/3)*d*(e^(1/3) - (-1)^(1/3)*f^(1/3)*x)/((-1)^(2/3)*d*e^(1/3) - (1 + c)*f^(1/3))]/(6*f) - (b*Log[1 - c - d*x]*Log[(-1)^(1/3)*d*(e^(1/3) + (-1)^(2/3)*f^(1/3)*x)/((-1)^(1/3)*d*e^(1/3) - (1 - c)*f^(1/3))]/(6*f) + (b*Log[1 + c + d*x]*Log[(-1)^(1/3)*d*(e^(1/3) + (-1)^(2/3)*f^(1/3)*x)/((-1)^(1/3)*d*e^(1/3) + (1 + c)*f^(1/3))]/(6*f) + (a*Log[e + f*x^3]/(3*f) - (b*PolyLog[2, -(f^(1/3)*(1 - c - d*x))/((-1)^(1/3)*d*e^(1/3) - (1 - c)*f^(1/3))]/(6*f) - (b*PolyLog[2, (f^(1/3)*(1 - c - d*x))/(d*e^(1/3) + (1 - c)*f^(1/3))]/(6*f) - (b*PolyLog[2, (f^(1/3)*(1 - c - d*x))/((-1)^(2/3)*d*e^(1/3) + (1 - c)*f^(1/3))]/(6*f) + (b*PolyLog[2, -(f^(1/3)*(1 + c + d*x))/(d*e^(1/3) - (1 + c)*f^(1/3))]/(6*f) + (b*PolyLog[2, -(f^(1/3)*(1 + c + d*x))/((-1)^(2/3)*d*e^(1/3) - (1 + c)*f^(1/3))]/(6*f) + (b*PolyLog[2, (f^(1/3)*(1 + c + d*x))/((-1)^(1/3)*d*e^(1/3) + (1 + c)*f^(1/3))]/(6*f))
```

Rubi [A] (verified)

Time = 1.25 (sec) , antiderivative size = 467, normalized size of antiderivative = 1.01, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx$$

$$\downarrow 7276$$

$$\int \left(\frac{ax^2}{e + fx^3} + \frac{bx^2 \operatorname{arctanh}(c + dx)}{e + fx^3} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& \frac{a \log(e + fx^3)}{3f} + \frac{\operatorname{barctanh}(c + dx) \log\left(\frac{2d(\sqrt[3]{e} + \sqrt[3]{fx})}{(c+dx+1)((1-c)\sqrt[3]{f+d\sqrt[3]{e}})}\right)}{3f} + \\
& \frac{\operatorname{barctanh}(c + dx) \log\left(\frac{2d((-1)^{2/3}\sqrt[3]{e} + \sqrt[3]{fx})}{(c+dx+1)((1-c)\sqrt[3]{f+(-1)^{2/3}d\sqrt[3]{e}})}\right)}{3f} + \\
& \frac{\operatorname{barctanh}(c + dx) \log\left(\frac{2\sqrt[3]{-1}d(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx})}{(c+dx+1)(\sqrt[3]{-1}d\sqrt[3]{e} - (1-c)\sqrt[3]{f})}\right)}{3f} - \frac{\operatorname{barctanh}(c + dx) \log\left(\frac{2}{c+dx+1}\right)}{f} - \\
& \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt[3]{fx} + \sqrt[3]{e})}{(\sqrt[3]{f}(1-c) + d\sqrt[3]{e})(c+dx+1)}\right)}{6f} - \\
& \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt[3]{fx} + (-1)^{2/3}\sqrt[3]{e})}{(\sqrt[3]{f}(1-c) + (-1)^{2/3}d\sqrt[3]{e})(c+dx+1)}\right)}{6f} - \\
& \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt[3]{-1}d((-1)^{2/3}\sqrt[3]{fx} + \sqrt[3]{e})}{(\sqrt[3]{-1}d\sqrt[3]{e} - (1-c)\sqrt[3]{f})(c+dx+1)}\right)}{6f} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{c+dx+1}\right)}{2f}
\end{aligned}$$

input `Int[(x^2*(a + b*ArcTanh[c + d*x]))/(e + f*x^3),x]`

output `-((b*ArcTanh[c + d*x]*Log[2/(1 + c + d*x)])/f) + (b*ArcTanh[c + d*x]*Log[(2*d*(e^(1/3) + f^(1/3)*x))/((d*e^(1/3) + (1 - c)*f^(1/3))*(1 + c + d*x))])/(3*f) + (b*ArcTanh[c + d*x]*Log[(2*d*((-1)^(2/3)*e^(1/3) + f^(1/3)*x))/(((-1)^(2/3)*d*e^(1/3) + (1 - c)*f^(1/3))*(1 + c + d*x))])/(3*f) + (b*ArcTanh[c + d*x]*Log[(2*(-1)^(1/3)*d*(e^(1/3) + (-1)^(2/3)*f^(1/3)*x))/((-1)^(1/3)*d*e^(1/3) - (1 - c)*f^(1/3))*(1 + c + d*x))]/(3*f) + (a*Log[e + f*x^3])/(3*f) + (b*PolyLog[2, 1 - 2/(1 + c + d*x)])/(2*f) - (b*PolyLog[2, 1 - (2*d*(e^(1/3) + f^(1/3)*x))/((d*e^(1/3) + (1 - c)*f^(1/3))*(1 + c + d*x))])/(6*f) - (b*PolyLog[2, 1 - (2*d*((-1)^(2/3)*e^(1/3) + f^(1/3)*x))/((-1)^(2/3)*d*e^(1/3) + (1 - c)*f^(1/3))*(1 + c + d*x))]/(6*f) - (b*PolyLog[2, 1 - (2*(-1)^(1/3)*d*(e^(1/3) + (-1)^(2/3)*f^(1/3)*x))/((-1)^(1/3)*d*e^(1/3) - (1 - c)*f^(1/3))*(1 + c + d*x))]/(6*f)`

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 7276 Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xprand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.33 (sec) , antiderivative size = 327, normalized size of antiderivative = 0.71

| method | result |
|-------------------|--|
| risch | $\frac{b \left(\sum_{-R1=RootOf(fZ^3+(3fc-3f)Z^2+(3c^2f-6fc+3f)Z+c^3f-d^3e-3c^2f+3fc-f)} \ln(-dx-c+1) \ln\left(\frac{dx+R1+c}{-R1}\right) \right)}{6f}$ |
| parts | $\frac{a \ln(fx^3+e)}{3f} + b \left(\frac{d^3 \ln(f(dx+c)^3-3cf(dx+c)^2+3c^2f(dx+c)-c^3f+d^3e) \operatorname{arctanh}(dx+c)}{3f} - d^3 \frac{\ln(dx+c+1) \ln(f(dx+c)^3-3cf(dx+c)+c^3f-d^3e)}{3f} \right)$ |
| derivativedivides | $\frac{a d^3 \ln(c^3 f-3c^2 f(dx+c)+3cf(dx+c)^2-d^3 e-f(dx+c)^3)}{3f} - b d^3 \left(-\frac{\ln(c^3 f-3c^2 f(dx+c)+3cf(dx+c)^2-d^3 e-f(dx+c)^3) \operatorname{arctanh}(dx+c)}{3f} \right)$ |
| default | $\frac{a d^3 \ln(c^3 f-3c^2 f(dx+c)+3cf(dx+c)^2-d^3 e-f(dx+c)^3)}{3f} - b d^3 \left(-\frac{\ln(c^3 f-3c^2 f(dx+c)+3cf(dx+c)^2-d^3 e-f(dx+c)^3) \operatorname{arctanh}(dx+c)}{3f} \right)$ |

input `int(x^2*(a+b*arctanh(d*x+c))/(f*x^3+e),x,method=_RETURNVERBOSE)`

output `-1/6*b/f*sum(ln(-d*x-c+1)*ln((d*x+_R1+c-1)/_R1)+dilog((d*x+_R1+c-1)/_R1),_R1=RootOf(f*_Z^3+(3*c*f-3*f)*_Z^2+(3*c^2*f-6*c*f+3*f)*_Z+c^3*f-d^3*e-3*c^2*f+3*f*c-f))+1/3*a/f*ln((-d*x-c+1)^3*f+3*(-d*x-c+1)^2*c*f+3*(-d*x-c+1)*c^2*f+c^3*f-d^3*e-3*f*(-d*x-c+1)^2-6*(-d*x-c+1)*c*f-3*c^2*f+3*(-d*x-c+1)*f+3*f*c-f)+1/6*b/f*sum(ln(d*x+c+1)*ln((-d*x+_R1-c-1)/_R1)+dilog((-d*x+_R1-c-1)/_R1),_R1=RootOf(f*_Z^3+(-3*c*f-3*f)*_Z^2+(3*c^2*f+6*c*f+3*f)*_Z-c^3*f+d^3*e-3*c^2*f-3*f*c-f))`

Fricas [F]

$$\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)x^2}{fx^3 + e} dx$$

input `integrate(x^2*(a+b*arctanh(d*x+c))/(f*x^3+e),x, algorithm="fricas")`

output `integral((b*x^2*arctanh(d*x + c) + a*x^2)/(f*x^3 + e), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \text{Timed out}$$

input `integrate(x**2*(a+b*atanh(d*x+c))/(f*x**3+e),x)`

output `Timed out`

Maxima [F]

$$\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)x^2}{fx^3 + e} dx$$

input `integrate(x^2*(a+b*arctanh(d*x+c))/(f*x^3+e),x, algorithm="maxima")`

output `1/2*b*integrate(x^2*(log(d*x + c + 1) - log(-d*x - c + 1))/(f*x^3 + e), x) + 1/3*a*log(f*x^3 + e)/f`

Giac [F]

$$\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \int \frac{(b \operatorname{artanh}(dx + c) + a)x^2}{fx^3 + e} dx$$

input `integrate(x^2*(a+b*arctanh(d*x+c))/(f*x^3+e),x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)*x^2/(f*x^3 + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \int \frac{x^2(a + b \operatorname{atanh}(c + dx))}{fx^3 + e} dx$$

input `int((x^2*(a + b*atanh(c + d*x)))/(e + f*x^3),x)`

output `int((x^2*(a + b*atanh(c + d*x)))/(e + f*x^3), x)`

Reduce [F]

$$\int \frac{x^2(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx$$

$$= \frac{3 \left(\int \frac{\operatorname{atanh}(dx+c)x^2}{fx^3+e} dx \right) bf + \log\left(e^{\frac{2}{3}} - f^{\frac{1}{3}}e^{\frac{1}{3}}x + f^{\frac{2}{3}}x^2\right) a + \log\left(e^{\frac{1}{3}} + f^{\frac{1}{3}}x\right) a}{3f}$$

input `int(x^2*(a+b*atanh(d*x+c))/(f*x^3+e),x)`

output `(3*int((atanh(c + d*x)*x**2)/(e + f*x**3),x)*b*f + log(e**(2/3) - f**(1/3)*e**(1/3)*x + f**(2/3)*x**2)*a + log(e**(1/3) + f**(1/3)*x)*a)/(3*f)`

$$3.72 \quad \int \frac{x(a+b \operatorname{arctanh}(c+dx))}{e+fx^3} dx$$

| | |
|---|-----|
| Optimal result | 628 |
| Mathematica [C] (warning: unable to verify) | 629 |
| Rubi [A] (verified) | 630 |
| Maple [C] (warning: unable to verify) | 632 |
| Fricas [F] | 633 |
| Sympy [F(-1)] | 634 |
| Maxima [F(-2)] | 634 |
| Giac [F] | 634 |
| Mupad [F(-1)] | 635 |
| Reduce [F] | 635 |

Optimal result

Integrand size = 21, antiderivative size = 679

$$\begin{aligned}
& \int \frac{x(a + \operatorname{barctanh}(c + dx))}{e + fx^3} dx \\
&= \frac{(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{3\sqrt[3]{e}f^{2/3}} - \frac{\sqrt[3]{-1}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{3\sqrt[3]{e}f^{2/3}} \\
&+ \frac{(-1)^{2/3}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{3\sqrt[3]{e}f^{2/3}} \\
&- \frac{(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d\left(\sqrt[3]{e} + \sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} + (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{3\sqrt[3]{e}f^{2/3}} \\
&+ \frac{\sqrt[3]{-1}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d\left(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{3\sqrt[3]{e}f^{2/3}} \\
&- \frac{(-1)^{2/3}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d\left(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} + (-1)^{2/3}(1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{3\sqrt[3]{e}f^{2/3}} \\
&- \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{6\sqrt[3]{e}f^{2/3}} + \frac{\sqrt[3]{-1}b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{6\sqrt[3]{e}f^{2/3}} \\
&- \frac{(-1)^{2/3}b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{6\sqrt[3]{e}f^{2/3}} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d\left(\sqrt[3]{e} + \sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} + (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{6\sqrt[3]{e}f^{2/3}} \\
&- \frac{\sqrt[3]{-1}b \operatorname{PolyLog}\left(2, 1 - \frac{2d\left(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{6\sqrt[3]{e}f^{2/3}} \\
&+ \frac{(-1)^{2/3}b \operatorname{PolyLog}\left(2, 1 - \frac{2d\left(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} + (-1)^{2/3}(1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{6\sqrt[3]{e}f^{2/3}}
\end{aligned}$$

output

```

1/3*(a+b*arctanh(d*x+c))*ln(2/(d*x+c+1))/e^(1/3)/f^(2/3)-1/3*(-1)^(1/3)*(a
+b*arctanh(d*x+c))*ln(2/(d*x+c+1))/e^(1/3)/f^(2/3)+1/3*(-1)^(2/3)*(a+b*arc
tanh(d*x+c))*ln(2/(d*x+c+1))/e^(1/3)/f^(2/3)-1/3*(a+b*arctanh(d*x+c))*ln(2
*d*(e^(1/3)+f^(1/3)*x)/(d*e^(1/3)+(1-c)*f^(1/3))/(d*x+c+1))/e^(1/3)/f^(2/3
)+1/3*(-1)^(1/3)*(a+b*arctanh(d*x+c))*ln(2*d*(e^(1/3)-(-1)^(1/3)*f^(1/3)*x
)/(d*e^(1/3)-(-1)^(1/3)*(1-c)*f^(1/3))/(d*x+c+1))/e^(1/3)/f^(2/3)-1/3*(-1)
^(2/3)*(a+b*arctanh(d*x+c))*ln(2*d*(e^(1/3)+(-1)^(2/3)*f^(1/3)*x)/(d*e^(1/
3)+(-1)^(2/3)*(1-c)*f^(1/3))/(d*x+c+1))/e^(1/3)/f^(2/3)-1/6*b*polylog(2,1-
2/(d*x+c+1))/e^(1/3)/f^(2/3)+1/6*(-1)^(1/3)*b*polylog(2,1-2/(d*x+c+1))/e^(
1/3)/f^(2/3)-1/6*(-1)^(2/3)*b*polylog(2,1-2/(d*x+c+1))/e^(1/3)/f^(2/3)+1/6
*b*polylog(2,1-2*d*(e^(1/3)+f^(1/3)*x)/(d*e^(1/3)+(1-c)*f^(1/3))/(d*x+c+1)
)/e^(1/3)/f^(2/3)-1/6*(-1)^(1/3)*b*polylog(2,1-2*d*(e^(1/3)-(-1)^(1/3)*f^(
1/3)*x)/(d*e^(1/3)-(-1)^(1/3)*(1-c)*f^(1/3))/(d*x+c+1))/e^(1/3)/f^(2/3)+1/
6*(-1)^(2/3)*b*polylog(2,1-2*d*(e^(1/3)+(-1)^(2/3)*f^(1/3)*x)/(d*e^(1/3)+(-
1)^(2/3)*(1-c)*f^(1/3))/(d*x+c+1))/e^(1/3)/f^(2/3)

```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.74 (sec) , antiderivative size = 652, normalized size of antiderivative = 0.96

$$\int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx$$

$$= \frac{3ax^2 \operatorname{Hypergeometric2F1}\left(\frac{2}{3}, 1, \frac{5}{3}, -\frac{fx^3}{e}\right) + \frac{be^{2/3} \left(\log(1-c-dx) \log\left(\frac{d\left(\sqrt[3]{e} + \sqrt[3]{fx}\right)}{d\sqrt[3]{e} - (-1+c)\sqrt[3]{f}}\right) - \log(1+c+dx) \log\left(\frac{d\left(\sqrt[3]{e} + \sqrt[3]{fx}\right)}{d\sqrt[3]{e} - (1+c)}{d\sqrt[3]{e} - (1+c)}\right)}{\right)}{e^{2/3}}}{e^{2/3}}$$

input

```
Integrate[(x*(a + b*ArcTanh[c + d*x]))/(e + f*x^3),x]
```

output

```
(3*a*x^2*Hypergeometric2F1[2/3, 1, 5/3, -((f*x^3)/e)] + (b*e^(2/3)*(Log[1
- c - d*x]*Log[(d*(e^(1/3) + f^(1/3)*x))/(d*e^(1/3) - (-1 + c)*f^(1/3))] -
Log[1 + c + d*x]*Log[(d*(e^(1/3) + f^(1/3)*x))/(d*e^(1/3) - (1 + c)*f^(1/3)
3))]) - (-1)^(1/3)*Log[1 - c - d*x]*Log[(d*(e^(1/3) - (-1)^(1/3)*f^(1/3)*x)
)/(d*e^(1/3) + (-1)^(1/3)*(-1 + c)*f^(1/3))] + (-1)^(1/3)*Log[1 + c + d*x]
*Log[(d*(e^(1/3) - (-1)^(1/3)*f^(1/3)*x))/(d*e^(1/3) + (-1)^(1/3)*(1 + c)*
f^(1/3))] + (-1)^(2/3)*Log[1 - c - d*x]*Log[(d*(e^(1/3) + (-1)^(2/3)*f^(1/3)
*x))/(d*e^(1/3) - (-1)^(2/3)*(-1 + c)*f^(1/3))] - (-1)^(2/3)*Log[1 + c +
d*x]*Log[(d*(e^(1/3) + (-1)^(2/3)*f^(1/3)*x))/(d*e^(1/3) - (-1)^(2/3)*(1
+ c)*f^(1/3))] + PolyLog[2, -((f^(1/3)*(-1 + c + d*x))/(d*e^(1/3) - (-1 +
c)*f^(1/3)))] - (-1)^(1/3)*PolyLog[2, ((-1)^(1/3)*f^(1/3)*(-1 + c + d*x))/
(d*e^(1/3) + (-1)^(1/3)*(-1 + c)*f^(1/3))] + (-1)^(2/3)*PolyLog[2, ((-1)^(
2/3)*f^(1/3)*(-1 + c + d*x))/(-d*e^(1/3) + (-1)^(2/3)*(-1 + c)*f^(1/3))]
- PolyLog[2, -((f^(1/3)*(1 + c + d*x))/(d*e^(1/3) - (1 + c)*f^(1/3)))] +
(-1)^(1/3)*PolyLog[2, ((-1)^(1/3)*f^(1/3)*(1 + c + d*x))/(d*e^(1/3) + (-1)
^(1/3)*(1 + c)*f^(1/3))] - (-1)^(2/3)*PolyLog[2, ((-1)^(2/3)*f^(1/3)*(1 +
c + d*x))/(-d*e^(1/3) + (-1)^(2/3)*(1 + c)*f^(1/3)))]/f^(2/3))/(6*e)
```

Rubi [A] (verified)

Time = 1.51 (sec) , antiderivative size = 778, normalized size of antiderivative = 1.15, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx$$

$$\downarrow \text{7276}$$

$$\int \left(\frac{ax}{e + fx^3} + \frac{bx \operatorname{arctanh}(c + dx)}{e + fx^3} \right) dx$$

$$\downarrow \text{2009}$$

$$\begin{aligned}
& -\frac{a \arctan\left(\frac{\sqrt[3]{e}-2\sqrt[3]{fx}}{\sqrt{3}\sqrt[3]{e}}\right)}{\sqrt{3}\sqrt[3]{ef^{2/3}}} + \frac{a \log\left(e^{2/3} - \sqrt[3]{e}\sqrt[3]{fx} + f^{2/3}x^2\right)}{6\sqrt[3]{ef^{2/3}}} - \frac{a \log\left(\sqrt[3]{e} + \sqrt[3]{fx}\right)}{3\sqrt[3]{ef^{2/3}}} + \\
& \frac{(-1)^{2/3}b \operatorname{arctanh}(c+dx) \log\left(\frac{2}{c+dx+1}\right)}{3\sqrt[3]{ef^{2/3}}} - \frac{\sqrt[3]{-1}b \operatorname{arctanh}(c+dx) \log\left(\frac{2}{c+dx+1}\right)}{3\sqrt[3]{ef^{2/3}}} + \\
& \frac{b \operatorname{arctanh}(c+dx) \log\left(\frac{2}{c+dx+1}\right)}{3\sqrt[3]{ef^{2/3}}} - \frac{b \operatorname{arctanh}(c+dx) \log\left(\frac{2d\left(\sqrt[3]{e} + \sqrt[3]{fx}\right)}{(c+dx+1)\left((1-c)\sqrt[3]{f} + d\sqrt[3]{e}\right)}\right)}{3\sqrt[3]{ef^{2/3}}} + \\
& \frac{\sqrt[3]{-1}b \operatorname{arctanh}(c+dx) \log\left(\frac{2d\left(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx}\right)}{(c+dx+1)\left(d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f}\right)}\right)}{3\sqrt[3]{ef^{2/3}}} - \\
& \frac{(-1)^{2/3}b \operatorname{arctanh}(c+dx) \log\left(\frac{2d\left(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx}\right)}{(c+dx+1)\left((-1)^{2/3}(1-c)\sqrt[3]{f} + d\sqrt[3]{e}\right)}\right)}{3\sqrt[3]{ef^{2/3}}} - \\
& \frac{(-1)^{2/3}b \operatorname{PolyLog}\left(2, 1 - \frac{2}{c+dx+1}\right)}{6\sqrt[3]{ef^{2/3}}} + \frac{\sqrt[3]{-1}b \operatorname{PolyLog}\left(2, 1 - \frac{2}{c+dx+1}\right)}{6\sqrt[3]{ef^{2/3}}} - \\
& \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{c+dx+1}\right)}{6\sqrt[3]{ef^{2/3}}} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d\left(\sqrt[3]{fx} + \sqrt[3]{e}\right)}{\left(\sqrt[3]{f}(1-c) + d\sqrt[3]{e}\right)(c+dx+1)}\right)}{6\sqrt[3]{ef^{2/3}}} - \\
& \frac{\sqrt[3]{-1}b \operatorname{PolyLog}\left(2, 1 - \frac{2d\left(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f}\right)(c+dx+1)}\right)}{6\sqrt[3]{ef^{2/3}}} + \\
& \frac{(-1)^{2/3}b \operatorname{PolyLog}\left(2, 1 - \frac{2d\left((-1)^{2/3}\sqrt[3]{fx} + \sqrt[3]{e}\right)}{\left((-1)^{2/3}\sqrt[3]{f}(1-c) + d\sqrt[3]{e}\right)(c+dx+1)}\right)}{6\sqrt[3]{ef^{2/3}}}
\end{aligned}$$

input

```
Int[(x*(a + b*ArcTanh[c + d*x]))/(e + f*x^3),x]
```


output

$$\begin{aligned}
& -((a*\text{ArcTan}[(e^{(1/3)} - 2*f^{(1/3)}*x)/(\text{Sqrt}[3]*e^{(1/3)})]) / (\text{Sqrt}[3]*e^{(1/3)}*f^{(2/3)})) + (b*\text{ArcTanh}[c + d*x]*\text{Log}[2/(1 + c + d*x)]) / (3*e^{(1/3)}*f^{(2/3)}) - \\
& ((-1)^{(1/3)}*b*\text{ArcTanh}[c + d*x]*\text{Log}[2/(1 + c + d*x)]) / (3*e^{(1/3)}*f^{(2/3)}) \\
& + ((-1)^{(2/3)}*b*\text{ArcTanh}[c + d*x]*\text{Log}[2/(1 + c + d*x)]) / (3*e^{(1/3)}*f^{(2/3)}) \\
& - (a*\text{Log}[e^{(1/3)} + f^{(1/3)}*x]) / (3*e^{(1/3)}*f^{(2/3)}) - (b*\text{ArcTanh}[c + d*x]* \\
& \text{Log}[(2*d*(e^{(1/3)} + f^{(1/3)}*x)) / ((d*e^{(1/3)} + (1 - c)*f^{(1/3)})*(1 + c + d* \\
& x))]) / (3*e^{(1/3)}*f^{(2/3)}) + ((-1)^{(1/3)}*b*\text{ArcTanh}[c + d*x]*\text{Log}[(2*d*(e^{(1/3)} \\
& - (-1)^{(1/3)}*f^{(1/3)}*x)) / ((d*e^{(1/3)} - (-1)^{(1/3)}*(1 - c)*f^{(1/3)})*(1 + \\
& c + d*x))]) / (3*e^{(1/3)}*f^{(2/3)}) - ((-1)^{(2/3)}*b*\text{ArcTanh}[c + d*x]*\text{Log}[(2*d* \\
& *(e^{(1/3)} + (-1)^{(2/3)}*f^{(1/3)}*x)) / ((d*e^{(1/3)} + (-1)^{(2/3)}*(1 - c)*f^{(1/3)}) \\
& *(1 + c + d*x))]) / (3*e^{(1/3)}*f^{(2/3)}) + (a*\text{Log}[e^{(2/3)} - e^{(1/3)}*f^{(1/3)} \\
& *x + f^{(2/3)}*x^2]) / (6*e^{(1/3)}*f^{(2/3)}) - (b*\text{PolyLog}[2, 1 - 2/(1 + c + d*x)] \\
&) / (6*e^{(1/3)}*f^{(2/3)}) + ((-1)^{(1/3)}*b*\text{PolyLog}[2, 1 - 2/(1 + c + d*x)]) / (6 \\
& *e^{(1/3)}*f^{(2/3)}) - ((-1)^{(2/3)}*b*\text{PolyLog}[2, 1 - 2/(1 + c + d*x)]) / (6*e^{(1 \\
& /3)}*f^{(2/3)}) + (b*\text{PolyLog}[2, 1 - (2*d*(e^{(1/3)} + f^{(1/3)}*x)) / ((d*e^{(1/3)} + \\
& (1 - c)*f^{(1/3)})*(1 + c + d*x))]) / (6*e^{(1/3)}*f^{(2/3)}) - ((-1)^{(1/3)}*b*\text{Poly} \\
& \text{Log}[2, 1 - (2*d*(e^{(1/3)} - (-1)^{(1/3)}*f^{(1/3)}*x)) / ((d*e^{(1/3)} - (-1)^{(1/3)} \\
&)*(1 - c)*f^{(1/3)})*(1 + c + d*x))]) / (6*e^{(1/3)}*f^{(2/3)}) + ((-1)^{(2/3)}*b*\text{Poly} \\
& \text{Log}[2, 1 - (2*d*(e^{(1/3)} + (-1)^{(2/3)}*f^{(1/3)}*x)) / ((d*e^{(1/3)} + (-1)^{(2/3)} \\
&)*(1 - c)*f^{(1/3)})*(1 + c + d*x))]) / (6*e^{(1/3)}*f^{(2/3)})
\end{aligned}$$

Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

rule 7276

$$\text{Int}[(u_)/((a_) + (b_.)*(x_)^{(n_)}), x_Symbol] \rightarrow \text{With}[\{v = \text{RationalFunctionE} \\ \text{xpend}[u/(a + b*x^n), x]\}, \text{Int}[v, x] /; \text{SumQ}[v]] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{IGtQ} \\ [n, 0]$$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.72 (sec) , antiderivative size = 338, normalized size of antiderivative = 0.50

| method | result |
|------------------|---|
| risch | $\frac{db \left(\sum_{R1=\text{RootOf}(f_Z^3+(3fc-3f)_Z^2+(3c^2f-6fc+3f)_Z+c^3f-d^3e-3c^2f+3fc-f)} \frac{\ln(-dx-c+1) \ln\left(\frac{dx+R1+c-1}{R1}\right)}{R1-1} \right)}{6f}$ |
| derivativdivides | Expression too large to display |
| default | Expression too large to display |
| parts | Expression too large to display |

input

```
int(x*(a+b*arctanh(d*x+c))/(f*x^3+e),x,method=_RETURNVERBOSE)
```

output

```
1/6*d*b/f*sum(1/(_R1-1+c)*(ln(-d*x-c+1)*ln((d*x+_R1+c-1)/_R1)+dilog((d*x+_R1+c-1)/_R1)),_R1=RootOf(f*_Z^3+(3*c*f-3*f)*_Z^2+(3*c^2*f-6*c*f+3*f)*_Z+c^3*f-d^3*e-3*c^2*f+3*f*c-f))-1/3*d*a/f*sum((c+_R-1)/(_R^2+2*_R*c+c^2-2*_R-2*c+1)*ln(-d*x-_R-c+1),_R=RootOf(f*_Z^3+(3*c*f-3*f)*_Z^2+(3*c^2*f-6*c*f+3*f)*_Z+c^3*f-d^3*e-3*c^2*f+3*f*c-f))+1/6*b*d/f*sum(1/(_R1-1-c)*(ln(d*x+c+1)*ln((-d*x+_R1-c-1)/_R1)+dilog((-d*x+_R1-c-1)/_R1)),_R1=RootOf(f*_Z^3+(-3*c*f-3*f)*_Z^2+(3*c^2*f+6*c*f+3*f)*_Z-c^3*f+d^3*e-3*c^2*f-3*f*c-f))
```

Fricas [F]

$$\int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \int \frac{(b \operatorname{arctanh}(dx + c) + a)x}{fx^3 + e} dx$$

input

```
integrate(x*(a+b*arctanh(d*x+c))/(f*x^3+e),x, algorithm="fricas")
```

output

```
integral((b*x*arctanh(d*x + c) + a*x)/(f*x^3 + e), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \text{Timed out}$$

input `integrate(x*(a+b*atanh(d*x+c))/(f*x**3+e),x)`

output Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \text{Exception raised: ValueError}$$

input `integrate(x*(a+b*arctanh(d*x+c))/(f*x^3+e),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e

Giac [F]

$$\int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \int \frac{(b \operatorname{arctanh}(dx + c) + a)x}{fx^3 + e} dx$$

input `integrate(x*(a+b*arctanh(d*x+c))/(f*x^3+e),x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)*x/(f*x^3 + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx = \int \frac{x(a + b \operatorname{atanh}(c + dx))}{fx^3 + e} dx$$

input `int((x*(a + b*atanh(c + d*x)))/(e + f*x^3),x)`

output `int((x*(a + b*atanh(c + d*x)))/(e + f*x^3), x)`

Reduce [F]

$$\int \frac{x(a + b \operatorname{arctanh}(c + dx))}{e + fx^3} dx$$

$$= \frac{-2\sqrt{3} \operatorname{atan}\left(\frac{e^{\frac{1}{3}} - 2f^{\frac{1}{3}}x}{e^{\frac{1}{3}}\sqrt{3}}\right) a + 6f^{\frac{2}{3}}e^{\frac{1}{3}} \left(\int \frac{\operatorname{atanh}(dx+c)x}{fx^3+e} dx\right) b + \log\left(e^{\frac{2}{3}} - f^{\frac{1}{3}}e^{\frac{1}{3}}x + f^{\frac{2}{3}}x^2\right) a - 2\log\left(e^{\frac{1}{3}} + f^{\frac{1}{3}}x\right)}{6f^{\frac{2}{3}}e^{\frac{1}{3}}}$$

input `int(x*(a+b*atanh(d*x+c))/(f*x^3+e),x)`

output `(- 2*sqrt(3)*atan((e**(1/3) - 2*f**(1/3)*x)/(e**(1/3)*sqrt(3)))*a + 6*f**(2/3)*e**(1/3)*int((atanh(c + d*x)*x)/(e + f*x**3),x)*b + log(e**(2/3) - f**(1/3)*e**(1/3)*x + f**(2/3)*x**2)*a - 2*log(e**(1/3) + f**(1/3)*x)*a)/(6*f**(2/3)*e**(1/3))`

3.73 $\int \frac{a+b\operatorname{arctanh}(c+dx)}{e+fx^3} dx$

| | |
|---|-----|
| Optimal result | 637 |
| Mathematica [A] (warning: unable to verify) | 638 |
| Rubi [A] (verified) | 639 |
| Maple [C] (warning: unable to verify) | 641 |
| Fricas [F] | 643 |
| Sympy [F(-1)] | 643 |
| Maxima [F(-2)] | 644 |
| Giac [F] | 644 |
| Mupad [F(-1)] | 644 |
| Reduce [F] | 645 |

Optimal result

Integrand size = 20, antiderivative size = 679

$$\begin{aligned}
& \int \frac{a + \operatorname{barctanh}(c + dx)}{e + fx^3} dx \\
&= -\frac{(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{3e^{2/3} \sqrt[3]{f}} + \frac{\sqrt[3]{-1}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{3e^{2/3} \sqrt[3]{f}} \\
&\quad - \frac{(-1)^{2/3}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{3e^{2/3} \sqrt[3]{f}} \\
&\quad + \frac{(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d\left(\sqrt[3]{e} + \sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} + (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{3e^{2/3} \sqrt[3]{f}} \\
&\quad + \frac{(-1)^{2/3}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d\left(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{3e^{2/3} \sqrt[3]{f}} \\
&\quad - \frac{\sqrt[3]{-1}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d\left(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} + (-1)^{2/3}(1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{3e^{2/3} \sqrt[3]{f}} \\
&\quad + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{6e^{2/3} \sqrt[3]{f}} - \frac{\sqrt[3]{-1}b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{6e^{2/3} \sqrt[3]{f}} \\
&\quad + \frac{(-1)^{2/3}b \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{6e^{2/3} \sqrt[3]{f}} - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d\left(\sqrt[3]{e} + \sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} + (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{6e^{2/3} \sqrt[3]{f}} \\
&\quad - \frac{(-1)^{2/3}b \operatorname{PolyLog}\left(2, 1 - \frac{2d\left(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{6e^{2/3} \sqrt[3]{f}} \\
&\quad + \frac{\sqrt[3]{-1}b \operatorname{PolyLog}\left(2, 1 - \frac{2d\left(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} + (-1)^{2/3}(1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{6e^{2/3} \sqrt[3]{f}}
\end{aligned}$$

output

```
-1/3*(a+b*arctanh(d*x+c))*ln(2/(d*x+c+1))/e^(2/3)/f^(1/3)+1/3*(-1)^(1/3)*
(a+b*arctanh(d*x+c))*ln(2/(d*x+c+1))/e^(2/3)/f^(1/3)-1/3*(-1)^(2/3)*(a+b*ar
ctanh(d*x+c))*ln(2/(d*x+c+1))/e^(2/3)/f^(1/3)+1/3*(a+b*arctanh(d*x+c))*ln(
2*d*(e^(1/3)+f^(1/3)*x)/(d*e^(1/3)+(1-c)*f^(1/3))/(d*x+c+1))/e^(2/3)/f^(1/
3)+1/3*(-1)^(2/3)*(a+b*arctanh(d*x+c))*ln(2*d*(e^(1/3)-(-1)^(1/3)*f^(1/3)*
x)/(d*e^(1/3)-(-1)^(1/3)*(1-c)*f^(1/3))/(d*x+c+1))/e^(2/3)/f^(1/3)-1/3*(-1
)^(1/3)*(a+b*arctanh(d*x+c))*ln(2*d*(e^(1/3)+(-1)^(2/3)*f^(1/3)*x)/(d*e^(1
/3)+(-1)^(2/3)*(1-c)*f^(1/3))/(d*x+c+1))/e^(2/3)/f^(1/3)+1/6*b*polylog(2,1
-2/(d*x+c+1))/e^(2/3)/f^(1/3)-1/6*(-1)^(1/3)*b*polylog(2,1-2/(d*x+c+1))/e^
(2/3)/f^(1/3)+1/6*(-1)^(2/3)*b*polylog(2,1-2/(d*x+c+1))/e^(2/3)/f^(1/3)-1/
6*b*polylog(2,1-2*d*(e^(1/3)+f^(1/3)*x)/(d*e^(1/3)+(1-c)*f^(1/3))/(d*x+c+1
))/e^(2/3)/f^(1/3)-1/6*(-1)^(2/3)*b*polylog(2,1-2*d*(e^(1/3)-(-1)^(1/3)*f^
(1/3)*x)/(d*e^(1/3)-(-1)^(1/3)*(1-c)*f^(1/3))/(d*x+c+1))/e^(2/3)/f^(1/3)+1
/6*(-1)^(1/3)*b*polylog(2,1-2*d*(e^(1/3)+(-1)^(2/3)*f^(1/3)*x)/(d*e^(1/3)+
(-1)^(2/3)*(1-c)*f^(1/3))/(d*x+c+1))/e^(2/3)/f^(1/3)
```

Mathematica [A] (warning: unable to verify)

Time = 0.54 (sec) , antiderivative size = 714, normalized size of antiderivative = 1.05

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^3} dx =$$

$$\frac{2\sqrt{3}a \arctan\left(\frac{1 - \frac{2\sqrt[3]{fx}}{\sqrt[3]{e}}}{\sqrt[3]{3}}\right) - 2a \log\left(\sqrt[3]{e} + \sqrt[3]{fx}\right) + b \log(1 - c - dx) \log\left(\frac{d\left(\sqrt[3]{e} + \sqrt[3]{fx}\right)}{d\sqrt[3]{e} - (-1+c)\sqrt[3]{f}}\right) - b \log\left(\frac{d\left(\sqrt[3]{e} - \sqrt[3]{fx}\right)}{d\sqrt[3]{e} - (-1+c)\sqrt[3]{f}}\right)}{\dots}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])/(e + f*x^3),x]
```

output

```

-1/6*(2*Sqrt[3]*a*ArcTan[(1 - (2*f^(1/3)*x)/e^(1/3))/Sqrt[3]] - 2*a*Log[e^(
(1/3) + f^(1/3)*x] + b*Log[1 - c - d*x]*Log[(d*(e^(1/3) + f^(1/3)*x))/(d*e
^(1/3) - (-1 + c)*f^(1/3))] - b*Log[1 + c + d*x]*Log[(d*(e^(1/3) + f^(1/3)
*x))/(d*e^(1/3) - (1 + c)*f^(1/3))] + (-1)^(2/3)*b*Log[1 - c - d*x]*Log[(d
*(e^(1/3) - (-1)^(1/3)*f^(1/3)*x))/(d*e^(1/3) + (-1)^(1/3)*(-1 + c)*f^(1/3
))] - (-1)^(2/3)*b*Log[1 + c + d*x]*Log[(d*(e^(1/3) - (-1)^(1/3)*f^(1/3)*x
))/(d*e^(1/3) + (-1)^(1/3)*(1 + c)*f^(1/3))] - (-1)^(1/3)*b*Log[1 - c - d*
x]*Log[(d*(e^(1/3) + (-1)^(2/3)*f^(1/3)*x))/(d*e^(1/3) - (-1)^(2/3)*(-1 +
c)*f^(1/3))] + (-1)^(1/3)*b*Log[1 + c + d*x]*Log[(d*(e^(1/3) + (-1)^(2/3)*
f^(1/3)*x))/(d*e^(1/3) - (-1)^(2/3)*(1 + c)*f^(1/3))] + a*Log[e^(2/3) - e^(
1/3)*f^(1/3)*x + f^(2/3)*x^2] + b*PolyLog[2, -((f^(1/3)*(-1 + c + d*x))/(
d*e^(1/3) - (-1 + c)*f^(1/3))] + (-1)^(2/3)*b*PolyLog[2, ((-1)^(1/3)*f^(1
/3)*(-1 + c + d*x))/(d*e^(1/3) + (-1)^(1/3)*(-1 + c)*f^(1/3))] - (-1)^(1/3
)*b*PolyLog[2, ((-1)^(2/3)*f^(1/3)*(-1 + c + d*x))/(-d*e^(1/3)) + (-1)^(2
/3)*(-1 + c)*f^(1/3))] - b*PolyLog[2, -((f^(1/3)*(1 + c + d*x))/(d*e^(1/3)
- (1 + c)*f^(1/3))] - (-1)^(2/3)*b*PolyLog[2, ((-1)^(1/3)*f^(1/3)*(1 + c
+ d*x))/(d*e^(1/3) + (-1)^(1/3)*(1 + c)*f^(1/3))] + (-1)^(1/3)*b*PolyLog[
2, ((-1)^(2/3)*f^(1/3)*(1 + c + d*x))/(-d*e^(1/3)) + (-1)^(2/3)*(1 + c)*f
^(1/3)]]/(e^(2/3)*f^(1/3))

```

Rubi [A] (verified)

Time = 1.55 (sec) , antiderivative size = 919, normalized size of antiderivative = 1.35, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^3} dx$$

$$\downarrow 7276$$

$$\int \left(\frac{a}{e + fx^3} + \frac{b \operatorname{arctanh}(c + dx)}{e + fx^3} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& - \frac{a \arctan\left(\frac{\sqrt[3]{e-2}\sqrt[3]{fx}}{\sqrt{3}\sqrt[3]{e}}\right)}{\sqrt{3}e^{2/3}\sqrt[3]{f}} + \frac{a \log\left(\sqrt[3]{fx} + \sqrt[3]{e}\right)}{3e^{2/3}\sqrt[3]{f}} - \frac{b \log(-c-dx+1) \log\left(\frac{d\left(\sqrt[3]{fx} + \sqrt[3]{e}\right)}{\sqrt[3]{f}(1-c)+d\sqrt[3]{e}}\right)}{6e^{2/3}\sqrt[3]{f}} + \\
& \frac{b \log(c+dx+1) \log\left(\frac{d\left(\sqrt[3]{fx} + \sqrt[3]{e}\right)}{d\sqrt[3]{e}-(c+1)\sqrt[3]{f}}\right)}{6e^{2/3}\sqrt[3]{f}} - \\
& \frac{(-1)^{2/3}b \log(-c-dx+1) \log\left(\frac{d\left(\sqrt[3]{e}-\sqrt[3]{-1}\sqrt[3]{fx}\right)}{d\sqrt[3]{e}-\sqrt[3]{-1}(1-c)\sqrt[3]{f}}\right)}{6e^{2/3}\sqrt[3]{f}} + \\
& \frac{(-1)^{2/3}b \log(c+dx+1) \log\left(\frac{d\left(\sqrt[3]{e}-\sqrt[3]{-1}\sqrt[3]{fx}\right)}{\sqrt[3]{-1}\sqrt[3]{f}(c+1)+d\sqrt[3]{e}}\right)}{6e^{2/3}\sqrt[3]{f}} + \\
& \frac{\sqrt[3]{-1}b \log(-c-dx+1) \log\left(\frac{d\left((-1)^{2/3}\sqrt[3]{fx} + \sqrt[3]{e}\right)}{(-1)^{2/3}\sqrt[3]{f}(1-c)+d\sqrt[3]{e}}\right)}{6e^{2/3}\sqrt[3]{f}} - \\
& \frac{\sqrt[3]{-1}b \log(c+dx+1) \log\left(\frac{d\left((-1)^{2/3}\sqrt[3]{fx} + \sqrt[3]{e}\right)}{d\sqrt[3]{e}-(-1)^{2/3}(c+1)\sqrt[3]{f}}\right)}{6e^{2/3}\sqrt[3]{f}} - \frac{a \log\left(f^{2/3}x^2 - \sqrt[3]{e}\sqrt[3]{fx} + e^{2/3}\right)}{6e^{2/3}\sqrt[3]{f}} - \\
& \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt[3]{f}(-c-dx+1)}{\sqrt[3]{f}(1-c)+d\sqrt[3]{e}}\right)}{6e^{2/3}\sqrt[3]{f}} - \frac{(-1)^{2/3}b \operatorname{PolyLog}\left(2, -\frac{\sqrt[3]{-1}\sqrt[3]{f}(-c-dx+1)}{d\sqrt[3]{e}-\sqrt[3]{-1}(1-c)\sqrt[3]{f}}\right)}{6e^{2/3}\sqrt[3]{f}} + \\
& \frac{\sqrt[3]{-1}b \operatorname{PolyLog}\left(2, \frac{(-1)^{2/3}\sqrt[3]{f}(-c-dx+1)}{(-1)^{2/3}\sqrt[3]{f}(1-c)+d\sqrt[3]{e}}\right)}{6e^{2/3}\sqrt[3]{f}} + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt[3]{f}(c+dx+1)}{d\sqrt[3]{e}-(c+1)\sqrt[3]{f}}\right)}{6e^{2/3}\sqrt[3]{f}} + \\
& \frac{(-1)^{2/3}b \operatorname{PolyLog}\left(2, \frac{\sqrt[3]{-1}\sqrt[3]{f}(c+dx+1)}{\sqrt[3]{-1}\sqrt[3]{f}(c+1)+d\sqrt[3]{e}}\right)}{6e^{2/3}\sqrt[3]{f}} - \\
& \frac{\sqrt[3]{-1}b \operatorname{PolyLog}\left(2, -\frac{\sqrt[3]{f}\left((-1)^{2/3}c+(-1)^{2/3}dx+(-1)^{2/3}\right)}{d\sqrt[3]{e}-(-1)^{2/3}(c+1)\sqrt[3]{f}}\right)}{6e^{2/3}\sqrt[3]{f}}
\end{aligned}$$

input

```
Int[(a + b*ArcTanh[c + d*x])/(e + f*x^3), x]
```

output

$$\begin{aligned}
& -((a*\text{ArcTan}[(e^{1/3} - 2*f^{1/3}*x)/(\text{Sqrt}[3]*e^{1/3})]) / (\text{Sqrt}[3]*e^{2/3}*f^{1/3})) + (a*\text{Log}[e^{1/3} + f^{1/3}*x]) / (3*e^{2/3}*f^{1/3}) - (b*\text{Log}[1 - c - d*x]*\text{Log}[(d*(e^{1/3} + f^{1/3}*x)) / (d*e^{1/3} + (1 - c)*f^{1/3})]) / (6*e^{2/3}*f^{1/3}) + (b*\text{Log}[1 + c + d*x]*\text{Log}[(d*(e^{1/3} + f^{1/3}*x)) / (d*e^{1/3} - (1 + c)*f^{1/3})]) / (6*e^{2/3}*f^{1/3}) - ((-1)^{2/3}*b*\text{Log}[1 - c - d*x]*\text{Log}[(d*(e^{1/3} - (-1)^{1/3}*f^{1/3}*x)) / (d*e^{1/3} - (-1)^{1/3}*(1 - c)*f^{1/3})]) / (6*e^{2/3}*f^{1/3}) + ((-1)^{2/3}*b*\text{Log}[1 + c + d*x]*\text{Log}[(d*(e^{1/3} - (-1)^{1/3}*f^{1/3}*x)) / (d*e^{1/3} + (-1)^{1/3}*(1 + c)*f^{1/3})]) / (6*e^{2/3}*f^{1/3}) + ((-1)^{1/3}*b*\text{Log}[1 - c - d*x]*\text{Log}[(d*(e^{1/3} + (-1)^{2/3}*f^{1/3}*x)) / (d*e^{1/3} + (-1)^{2/3}*(1 - c)*f^{1/3})]) / (6*e^{2/3}*f^{1/3}) - ((-1)^{1/3}*b*\text{Log}[1 + c + d*x]*\text{Log}[(d*(e^{1/3} + (-1)^{2/3}*f^{1/3}*x)) / (d*e^{1/3} - (-1)^{2/3}*(1 + c)*f^{1/3})]) / (6*e^{2/3}*f^{1/3}) - (a*\text{Log}[e^{2/3} - e^{1/3}*f^{1/3}*x + f^{2/3}*x^2]) / (6*e^{2/3}*f^{1/3}) - (b*\text{PolyLog}[2, (f^{1/3}*(1 - c - d*x)) / (d*e^{1/3} + (1 - c)*f^{1/3})]) / (6*e^{2/3}*f^{1/3}) - ((-1)^{2/3}*b*\text{PolyLog}[2, -(((1 - c - d*x)) / (d*e^{1/3} - (-1)^{1/3}*(1 - c)*f^{1/3}))]) / (6*e^{2/3}*f^{1/3}) + ((-1)^{1/3}*b*\text{PolyLog}[2, ((1 - c - d*x)) / (d*e^{1/3} + (-1)^{2/3}*(1 - c)*f^{1/3})]) / (6*e^{2/3}*f^{1/3}) + (b*\text{PolyLog}[2, -((f^{1/3}*(1 + c + d*x)) / (d*e^{1/3} - (1 + c)*f^{1/3}))]) / (6*e^{2/3}*f^{1/3}) + ((-1)^{2/3}*b*\text{PolyLog}[2, ((1 + c + d*x)) / (d*e^{1/3} + ...
\end{aligned}$$

Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \text{ :> } \text{Simp}[\text{IntSum}[u, x], x] \text{ /; } \text{SumQ}[u]$$

rule 7276

$$\text{Int}[(u_)/((a_) + (b_.)*(x_)^{(n_)}), x_Symbol] \text{ :> } \text{With}[\{v = \text{RationalFunctionExpand}[u/(a + b*x^n), x]\}, \text{Int}[v, x] \text{ /; } \text{SumQ}[v]] \text{ /; } \text{FreeQ}[\{a, b\}, x] \text{ \&\& } \text{IGtQ}[n, 0]$$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.32 (sec) , antiderivative size = 366, normalized size of antiderivative = 0.54

| method | result |
|------------------|---|
| risch | $\frac{d^2 b \left(\frac{\sum_{R1=\text{RootOf}(fZ^3+(3fc-3f)Z^2+(3c^2f-6fc+3f)Z+c^3f-d^3e-3c^2f+3fc-f)} \ln(-dx-c+1) \ln\left(\frac{dx-R1+c}{R1}\right)}{-R1^2+2R1c} \right)}{6f}$ |
| derivativdivides | $\frac{a d^3 \left(\frac{\sum_{R=\text{RootOf}(fZ^3-3fcZ^2+3c^2fZ-c^3f+d^3e)} \ln(dx-R+c)}{3f} \right)}{-R^2+2Rc-c^2} - b d^3$ |
| default | $\frac{a d^3 \left(\frac{\sum_{R=\text{RootOf}(fZ^3-3fcZ^2+3c^2fZ-c^3f+d^3e)} \ln(dx-R+c)}{3f} \right)}{-R^2+2Rc-c^2} - b d^3$ |
| parts | $a \left(\frac{\ln\left(x+\left(\frac{e}{f}\right)^{\frac{1}{3}}\right)}{3f\left(\frac{e}{f}\right)^{\frac{2}{3}}} - \frac{\ln\left(x^2-\left(\frac{e}{f}\right)^{\frac{1}{3}}x+\left(\frac{e}{f}\right)^{\frac{2}{3}}\right)}{6f\left(\frac{e}{f}\right)^{\frac{2}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{e}{f}\right)^{\frac{1}{3}}}-1\right)}{3}\right)}{3f\left(\frac{e}{f}\right)^{\frac{2}{3}}} \right) - \frac{2b d^2}{-R1=\text{RootOf}(c^3f-d^3e-3c^2f+3fc-f)}$ |

```
input int((a+b*arctanh(d*x+c))/(f*x^3+e),x,method=_RETURNVERBOSE)
```

output

```
-1/6*d^2*b/f*sum(1/(_R1^2+2*_R1*c+c^2-2*_R1-2*c+1)*(ln(-d*x-c+1)*ln((d*x+_R1+c-1)/_R1)+dilog((d*x+_R1+c-1)/_R1)),_R1=RootOf(f*_Z^3+(3*c*f-3*f)*_Z^2+(3*c^2*f-6*c*f+3*f)*_Z+c^3*f-d^3*e-3*c^2*f+3*f*c-f))+1/3*d^2*a/f*sum(1/(_R1^2+2*_R1*c+c^2-2*_R1-2*c+1)*ln(-d*x-_R1-c+1),_R1=RootOf(f*_Z^3+(3*c*f-3*f)*_Z^2+(3*c^2*f-6*c*f+3*f)*_Z+c^3*f-d^3*e-3*c^2*f+3*f*c-f))+1/6*b*d^2/f*sum(1/(_R1^2-2*_R1*c+c^2-2*_R1+2*c+1)*(ln(d*x+c+1)*ln((-d*x+_R1-c-1)/_R1)+dilog((-d*x+_R1-c-1)/_R1)),_R1=RootOf(f*_Z^3+(-3*c*f-3*f)*_Z^2+(3*c^2*f+6*c*f+3*f)*_Z-c^3*f+d^3*e-3*c^2*f-3*f*c-f))
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^3} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{fx^3 + e} dx$$

input

```
integrate((a+b*arctanh(d*x+c))/(f*x^3+e),x, algorithm="fricas")
```

output

```
integral((b*arctanh(d*x + c) + a)/(f*x^3 + e), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^3} dx = \text{Timed out}$$

input

```
integrate((a+b*atanh(d*x+c))/(f*x**3+e),x)
```

output

```
Timed out
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^3} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arctanh(d*x+c))/(f*x^3+e),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^3} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{fx^3 + e} dx$$

input `integrate((a+b*arctanh(d*x+c))/(f*x^3+e),x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)/(f*x^3 + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^3} dx = \int \frac{a + b \operatorname{atanh}(c + dx)}{fx^3 + e} dx$$

input `int((a + b*atanh(c + d*x))/(e + f*x^3),x)`

output `int((a + b*atanh(c + d*x))/(e + f*x^3), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^3} dx$$

$$= \frac{-2e^{\frac{1}{3}}\sqrt{3} \operatorname{atan}\left(\frac{e^{\frac{1}{3}} - 2f^{\frac{1}{3}}x}{e^{\frac{1}{3}}\sqrt{3}}\right) a - e^{\frac{1}{3}}\log\left(e^{\frac{2}{3}} - f^{\frac{1}{3}}e^{\frac{1}{3}}x + f^{\frac{2}{3}}x^2\right) a + 2e^{\frac{1}{3}}\log\left(e^{\frac{1}{3}} + f^{\frac{1}{3}}x\right) a + 6f^{\frac{1}{3}}\left(\int \frac{\operatorname{atanh}(dx+c)}{fx^3+e}\right)}{6f^{\frac{1}{3}}e}$$

input

```
int((a+b*atanh(d*x+c))/(f*x^3+e),x)
```

output

```
( - 2*e**(1/3)*sqrt(3)*atan((e**(1/3) - 2*f**(1/3)*x)/(e**(1/3)*sqrt(3)))*
a - e**(1/3)*log(e**(2/3) - f**(1/3)*e**(1/3)*x + f**(2/3)*x**2)*a + 2*e**
(1/3)*log(e**(1/3) + f**(1/3)*x)*a + 6*f**(1/3)*int(atanh(c + d*x)/(e + f*
x**3),x)*b*e)/(6*f**(1/3)*e)
```

$$3.74 \quad \int \frac{a+b \operatorname{arctanh}(c+dx)}{x(e+fx^3)} dx$$

| | |
|---|-----|
| Optimal result | 647 |
| Mathematica [A] (warning: unable to verify) | 648 |
| Rubi [A] (verified) | 649 |
| Maple [C] (warning: unable to verify) | 651 |
| Fricas [F] | 652 |
| Sympy [F(-1)] | 652 |
| Maxima [F] | 653 |
| Giac [F] | 653 |
| Mupad [F(-1)] | 653 |
| Reduce [F] | 654 |

Optimal result

Integrand size = 23, antiderivative size = 480

$$\begin{aligned}
& \int \frac{a + \operatorname{barctanh}(c + dx)}{x(e + fx^3)} dx \\
&= \frac{(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2dx}{(1-c)(1+c+dx)}\right)}{e} \\
&\quad - \frac{(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d\left(\sqrt[3]{e} + \sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} + (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{3e} \\
&\quad - \frac{(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d\left((-1)^{2/3}\sqrt[3]{e} + \sqrt[3]{fx}\right)}{\left((-1)^{2/3}d\sqrt[3]{e} + (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{3e} \\
&\quad - \frac{(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2\sqrt[3]{-1}d\left(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx}\right)}{\left(\sqrt[3]{-1}d\sqrt[3]{e} - (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{3e} \\
&\quad - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2dx}{(1-c)(1+c+dx)}\right)}{2e} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d\left(\sqrt[3]{e} + \sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} + (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{6e} \\
&\quad + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d\left((-1)^{2/3}\sqrt[3]{e} + \sqrt[3]{fx}\right)}{\left((-1)^{2/3}d\sqrt[3]{e} + (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{6e} \\
&\quad + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt[3]{-1}d\left(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx}\right)}{\left(\sqrt[3]{-1}d\sqrt[3]{e} - (1-c)\sqrt[3]{f}\right)(1+c+dx)}\right)}{6e}
\end{aligned}$$

output

```
(a+b*arctanh(d*x+c))*ln(2*d*x/(1-c)/(d*x+c+1))/e-1/3*(a+b*arctanh(d*x+c))*
ln(2*d*(e^(1/3)+f^(1/3)*x)/(d*e^(1/3)+(1-c)*f^(1/3))/(d*x+c+1))/e-1/3*(a+b
*arctanh(d*x+c))*ln(2*d*((-1)^(2/3)*e^(1/3)+f^(1/3)*x)/((-1)^(2/3)*d*e^(1/
3)+(1-c)*f^(1/3))/(d*x+c+1))/e-1/3*(a+b*arctanh(d*x+c))*ln(2*(-1)^(1/3)*d*
(e^(1/3)+(-1)^(2/3)*f^(1/3)*x)/((-1)^(1/3)*d*e^(1/3)-(1-c)*f^(1/3))/(d*x+c
+1))/e-1/2*b*polylog(2,1-2*d*x/(1-c)/(d*x+c+1))/e+1/6*b*polylog(2,1-2*d*(e
^(1/3)+f^(1/3)*x)/(d*e^(1/3)+(1-c)*f^(1/3))/(d*x+c+1))/e+1/6*b*polylog(2,1
-2*d*((-1)^(2/3)*e^(1/3)+f^(1/3)*x)/((-1)^(2/3)*d*e^(1/3)+(1-c)*f^(1/3))/(
d*x+c+1))/e+1/6*b*polylog(2,1-2*(-1)^(1/3)*d*(e^(1/3)+(-1)^(2/3)*f^(1/3)*x
)/((-1)^(1/3)*d*e^(1/3)-(1-c)*f^(1/3))/(d*x+c+1))/e
```

Mathematica [A] (warning: unable to verify)

Time = 16.80 (sec) , antiderivative size = 915, normalized size of antiderivative = 1.91

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e + fx^3)} dx = \text{Too large to display}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])/(x*(e + f*x^3)),x]
```

output

```
(6*a*Log[x] - 2*a*Log[e + f*x^3] + b*(6*ArcTanh[c + d*x]*Log[x] + Log[e^(1/3)/f^(1/3) + x]*Log[1 - c - d*x] + Log[-(((-1)^(1/3)*e^(1/3))/f^(1/3)) + x]*Log[1 - c - d*x] + Log[(-1)^(2/3)*e^(1/3)/f^(1/3) + x]*Log[1 - c - d*x] - Log[e^(1/3)/f^(1/3) + x]*Log[-((f^(1/3)*(-1 + c + d*x))/(d*e^(1/3) - (-1 + c)*f^(1/3)))] - Log[(-1)^(2/3)*e^(1/3)/f^(1/3) + x]*Log[-((f^(1/3)*(-1 + c + d*x))/((-1)^(2/3)*d*e^(1/3) - (-1 + c)*f^(1/3)))] - Log[-(((-1)^(1/3)*e^(1/3))/f^(1/3)) + x]*Log[(f^(1/3)*(-1 + c + d*x))/((-1)^(1/3)*d*e^(1/3) + (-1 + c)*f^(1/3))] - Log[e^(1/3)/f^(1/3) + x]*Log[1 + c + d*x] - Log[-(((-1)^(1/3)*e^(1/3))/f^(1/3)) + x]*Log[1 + c + d*x] - Log[(-1)^(2/3)*e^(1/3)/f^(1/3) + x]*Log[1 + c + d*x] + Log[e^(1/3)/f^(1/3) + x]*Log[-((f^(1/3)*(1 + c + d*x))/(d*e^(1/3) - (1 + c)*f^(1/3)))] + Log[(-1)^(2/3)*e^(1/3)/f^(1/3) + x]*Log[-((f^(1/3)*(1 + c + d*x))/((-1)^(2/3)*d*e^(1/3) - (1 + c)*f^(1/3)))] + Log[-(((-1)^(1/3)*e^(1/3))/f^(1/3)) + x]*Log[(f^(1/3)*(1 + c + d*x))/((-1)^(1/3)*d*e^(1/3) + (1 + c)*f^(1/3))] + 3*Log[x]*Log[1 + (d*x)/(-1 + c)] - 3*Log[x]*Log[1 + (d*x)/(1 + c)] - 2*ArcTanh[c + d*x]*Log[e + f*x^3] - Log[1 - c - d*x]*Log[e + f*x^3] + Log[1 + c + d*x]*Log[e + f*x^3] + 3*PolyLog[2, -((d*x)/(-1 + c))] - 3*PolyLog[2, -((d*x)/(1 + c))] - PolyLog[2, (d*(e^(1/3)/f^(1/3) + x))/(1 - c + (d*e^(1/3))/f^(1/3))] - PolyLog[2, (d*(((-1)^(2/3)*e^(1/3))/f^(1/3) + x))/(1 - c + ((-1)^(2/3)*d*e^(1/3))/f^(1/3))] - PolyLog[2, (d*((-1)^(1/3)*e^(1/3) - f^(1/3)*x))/(...
```

Rubi [A] (verified)

Time = 1.53 (sec) , antiderivative size = 491, normalized size of antiderivative = 1.02, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e + fx^3)} dx$$

$$\downarrow 7276$$

$$\int \left(\frac{a}{x(e + fx^3)} + \frac{b \operatorname{arctanh}(c + dx)}{x(e + fx^3)} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& \frac{a \log(e + fx^3)}{3e} + \frac{a \log(x)}{e} - \frac{\operatorname{barctanh}(c + dx) \log\left(\frac{2d(\sqrt[3]{e} + \sqrt[3]{fx})}{(c+dx+1)((1-c)\sqrt[3]{f} + d\sqrt[3]{e})}\right)}{3e} \\
& \frac{\operatorname{barctanh}(c + dx) \log\left(\frac{2d(-1)^{2/3}\sqrt[3]{e} + \sqrt[3]{fx}}{(c+dx+1)((1-c)\sqrt[3]{f} + (-1)^{2/3}d\sqrt[3]{e})}\right)}{3e} \\
& \frac{\operatorname{barctanh}(c + dx) \log\left(\frac{2\sqrt[3]{-1}d(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx})}{(c+dx+1)(\sqrt[3]{-1}d\sqrt[3]{e} - (1-c)\sqrt[3]{f})}\right)}{3e} + \\
& \frac{\operatorname{barctanh}(c + dx) \log\left(\frac{2dx}{(1-c)(c+dx+1)}\right)}{e} + \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt[3]{fx} + \sqrt[3]{e})}{(\sqrt[3]{f}(1-c) + d\sqrt[3]{e})(c+dx+1)}\right)}{6e} + \\
& \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt[3]{fx} + (-1)^{2/3}\sqrt[3]{e})}{(\sqrt[3]{f}(1-c) + (-1)^{2/3}d\sqrt[3]{e})(c+dx+1)}\right)}{6e} + \\
& \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt[3]{-1}d((-1)^{2/3}\sqrt[3]{fx} + \sqrt[3]{e})}{(\sqrt[3]{-1}d\sqrt[3]{e} - (1-c)\sqrt[3]{f})(c+dx+1)}\right)}{6e} - \frac{b \operatorname{PolyLog}\left(2, 1 - \frac{2dx}{(1-c)(c+dx+1)}\right)}{2e}
\end{aligned}$$

input `Int[(a + b*ArcTanh[c + d*x])/(x*(e + f*x^3)),x]`

output `(a*Log[x])/e + (b*ArcTanh[c + d*x]*Log[(2*d*x)/((1 - c)*(1 + c + d*x))])/e - (b*ArcTanh[c + d*x]*Log[(2*d*(e^(1/3) + f^(1/3)*x))/((d*e^(1/3) + (1 - c)*f^(1/3))*(1 + c + d*x))])/ (3*e) - (b*ArcTanh[c + d*x]*Log[(2*d*((-1)^(2/3)*e^(1/3) + f^(1/3)*x))/(((-1)^(2/3)*d*e^(1/3) + (1 - c)*f^(1/3))*(1 + c + d*x))])/ (3*e) - (b*ArcTanh[c + d*x]*Log[(2*(-1)^(1/3)*d*(e^(1/3) + (-1)^(2/3)*f^(1/3)*x))/(((-1)^(1/3)*d*e^(1/3) - (1 - c)*f^(1/3))*(1 + c + d*x))])/ (3*e) - (a*Log[e + f*x^3])/ (3*e) - (b*PolyLog[2, 1 - (2*d*x)/((1 - c)*(1 + c + d*x))])/ (2*e) + (b*PolyLog[2, 1 - (2*d*(e^(1/3) + f^(1/3)*x))/((d*e^(1/3) + (1 - c)*f^(1/3))*(1 + c + d*x))])/ (6*e) + (b*PolyLog[2, 1 - (2*d*((-1)^(2/3)*e^(1/3) + f^(1/3)*x))/(((-1)^(2/3)*d*e^(1/3) + (1 - c)*f^(1/3))*(1 + c + d*x))])/ (6*e) + (b*PolyLog[2, 1 - (2*(-1)^(1/3)*d*(e^(1/3) + (-1)^(2/3)*f^(1/3)*x))/(((-1)^(1/3)*d*e^(1/3) - (1 - c)*f^(1/3))*(1 + c + d*x))])/ (6*e)`

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 7276 Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xprand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.31 (sec) , antiderivative size = 420, normalized size of antiderivative = 0.88

| method | result |
|-------------------|---|
| risch | $-\frac{b \ln(-dx-c+1) \ln\left(-\frac{xd}{-1+c}\right)}{2e} - \frac{b \operatorname{dilog}\left(-\frac{xd}{-1+c}\right)}{2e} + \frac{b \left(\operatorname{RootOf}\left(f Z^3 + (3fc-3f) Z^2 + (3c^2f-6fc+3f) Z + c^3\right) \right)}{e}$ |
| parts | $\frac{a \ln(x)}{e} - \frac{a \ln(fx^3+e)}{3e} + b \left(-\frac{\operatorname{arctanh}(dx+c) \ln\left(f(dx+c)^3 - 3cf(dx+c)^2 + 3c^2f(dx+c) - c^3f + d^3e\right)}{3e} + \operatorname{arctanh}(dx+c) \right)$ |
| derivativedivides | $\frac{a \ln(-dx)}{e} - \frac{a \ln\left(c^3f - 3c^2f(dx+c) + 3cf(dx+c)^2 - d^3e - f(dx+c)^3\right)}{3e} + b d^3 \left(\frac{\operatorname{arctanh}(dx+c) \ln(-dx)}{e d^3} - \operatorname{arctanh}(dx+c) \right)$ |
| default | $\frac{a \ln(-dx)}{e} - \frac{a \ln\left(c^3f - 3c^2f(dx+c) + 3cf(dx+c)^2 - d^3e - f(dx+c)^3\right)}{3e} + b d^3 \left(\frac{\operatorname{arctanh}(dx+c) \ln(-dx)}{e d^3} - \operatorname{arctanh}(dx+c) \right)$ |

input `int((a+b*arctanh(d*x+c))/x/(f*x^3+e),x,method=_RETURNVERBOSE)`

output `-1/2*b/e*ln(-d*x-c+1)*ln(-x*d/(-1+c))-1/2*b/e*dilog(-x*d/(-1+c))+1/6*b/e*sum(ln(-d*x-c+1)*ln((d*x+_R1+c-1)/_R1)+dilog((d*x+_R1+c-1)/_R1),_R1=RootOf(f*_Z^3+(3*c*f-3*f)*_Z^2+(3*c^2*f-6*c*f+3*f)*_Z+c^3*f-d^3*e-3*c^2*f+3*f*c-f))+a/e*ln(-d*x)-1/3*a/e*ln((-d*x-c+1)^3*f+3*(-d*x-c+1)^2*c*f+3*(-d*x-c+1)*c^2*f+c^3*f-d^3*e-3*f*(-d*x-c+1)^2-6*(-d*x-c+1)*c*f-3*c^2*f+3*(-d*x-c+1)*f+3*f*c-f)+1/2*b/e*ln(d*x+c+1)*ln(x*d/(-1-c))+1/2*b/e*dilog(x*d/(-1-c))-1/6*b/e*sum(ln(d*x+c+1)*ln((-d*x+_R1-c-1)/_R1)+dilog((-d*x+_R1-c-1)/_R1),_R1=RootOf(f*_Z^3+(-3*c*f-3*f)*_Z^2+(3*c^2*f+6*c*f+3*f)*_Z-c^3*f+d^3*e-3*c^2*f-3*f*c-f))`

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e + fx^3)} dx = \int \frac{b \operatorname{arctanh}(dx + c) + a}{(fx^3 + e)x} dx$$

input `integrate((a+b*arctanh(d*x+c))/x/(f*x^3+e),x, algorithm="fricas")`

output `integral((b*arctanh(d*x + c) + a)/(f*x^4 + e*x), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e + fx^3)} dx = \text{Timed out}$$

input `integrate((a+b*atanh(d*x+c))/x/(f*x**3+e),x)`

output `Timed out`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e + fx^3)} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{(fx^3 + e)x} dx$$

input `integrate((a+b*arctanh(d*x+c))/x/(f*x^3+e),x, algorithm="maxima")`

output `-1/3*a*(log(f*x^3 + e)/e - 3*log(x)/e) + 1/2*b*integrate((log(d*x + c + 1) - log(-d*x - c + 1))/(f*x^4 + e*x), x)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e + fx^3)} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{(fx^3 + e)x} dx$$

input `integrate((a+b*arctanh(d*x+c))/x/(f*x^3+e),x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)/((f*x^3 + e)*x), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e + fx^3)} dx = \int \frac{a + b \operatorname{atanh}(c + dx)}{x(fx^3 + e)} dx$$

input `int((a + b*atanh(c + d*x))/(x*(e + f*x^3)),x)`

output `int((a + b*atanh(c + d*x))/(x*(e + f*x^3)), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x(e + fx^3)} dx$$

$$= \frac{3 \left(\int \frac{\operatorname{atanh}(dx+c)}{fx^4+ex} dx \right) b e - \log\left(e^{\frac{2}{3}} - f^{\frac{1}{3}} e^{\frac{1}{3}} x + f^{\frac{2}{3}} x^2\right) a - \log\left(e^{\frac{1}{3}} + f^{\frac{1}{3}} x\right) a + 3 \log(x) a}{3e}$$

input `int((a+b*atanh(d*x+c))/x/(f*x^3+e),x)`

output `(3*int(atanh(c + d*x)/(e*x + f*x**4),x)*b*e - log(e**(2/3) - f**(1/3)*e**(1/3)*x + f**(2/3)*x**2)*a - log(e**(1/3) + f**(1/3)*x)*a + 3*log(x)*a)/(3*e)`

$$3.75 \quad \int \frac{a+b \operatorname{arctanh}(c+dx)}{x^2(e+fx^3)} dx$$

| | |
|---|-----|
| Optimal result | 656 |
| Mathematica [C] (warning: unable to verify) | 657 |
| Rubi [A] (verified) | 658 |
| Maple [C] (warning: unable to verify) | 660 |
| Fricas [F] | 661 |
| Sympy [F(-1)] | 662 |
| Maxima [F(-2)] | 662 |
| Giac [F] | 662 |
| Mupad [F(-1)] | 663 |
| Reduce [F] | 663 |

Optimal result

Integrand size = 23, antiderivative size = 761

$$\begin{aligned}
& \int \frac{a + \operatorname{barctanh}(c + dx)}{x^2 (e + fx^3)} dx \\
&= -\frac{a + \operatorname{barctanh}(c + dx)}{ex} + \frac{bd \log(x)}{(1 - c^2)e} - \frac{bd \log(1 - c - dx)}{2(1 - c)e} \\
&\quad - \frac{\sqrt[3]{f}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{3e^{4/3}} \\
&\quad + \frac{\sqrt[3]{-1} \sqrt[3]{f}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{3e^{4/3}} \\
&\quad - \frac{(-1)^{2/3} \sqrt[3]{f}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{3e^{4/3}} - \frac{bd \log(1 + c + dx)}{2(1 + c)e} \\
&\quad + \frac{\sqrt[3]{f}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d(\sqrt[3]{e} + \sqrt[3]{fx})}{(d\sqrt[3]{e} + (1-c)\sqrt[3]{f})(1+c+dx)}\right)}{3e^{4/3}} \\
&\quad - \frac{\sqrt[3]{-1} \sqrt[3]{f}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d(\sqrt[3]{e} - \sqrt[3]{-1} \sqrt[3]{fx})}{(d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f})(1+c+dx)}\right)}{3e^{4/3}} \\
&\quad + \frac{(-1)^{2/3} \sqrt[3]{f}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d(\sqrt[3]{e} + (-1)^{2/3} \sqrt[3]{fx})}{(d\sqrt[3]{e} + (-1)^{2/3}(1-c)\sqrt[3]{f})(1+c+dx)}\right)}{3e^{4/3}} \\
&\quad + \frac{b\sqrt[3]{f} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{6e^{4/3}} - \frac{\sqrt[3]{-1} b\sqrt[3]{f} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{6e^{4/3}} \\
&\quad + \frac{(-1)^{2/3} b\sqrt[3]{f} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{6e^{4/3}} \\
&\quad - \frac{b\sqrt[3]{f} \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt[3]{e} + \sqrt[3]{fx})}{(d\sqrt[3]{e} + (1-c)\sqrt[3]{f})(1+c+dx)}\right)}{6e^{4/3}} \\
&\quad + \frac{\sqrt[3]{-1} b\sqrt[3]{f} \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt[3]{e} - \sqrt[3]{-1} \sqrt[3]{fx})}{(d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f})(1+c+dx)}\right)}{6e^{4/3}} \\
&\quad - \frac{(-1)^{2/3} b\sqrt[3]{f} \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt[3]{e} + (-1)^{2/3} \sqrt[3]{fx})}{(d\sqrt[3]{e} + (-1)^{2/3}(1-c)\sqrt[3]{f})(1+c+dx)}\right)}{6e^{4/3}}
\end{aligned}$$

output

```

-(a+b*arctanh(d*x+c))/e/x+b*d*ln(x)/(-c^2+1)/e-1/2*b*d*ln(-d*x-c+1)/(1-c)/
e-1/3*f^(1/3)*(a+b*arctanh(d*x+c))*ln(2/(d*x+c+1))/e^(4/3)+1/3*(-1)^(1/3)*
f^(1/3)*(a+b*arctanh(d*x+c))*ln(2/(d*x+c+1))/e^(4/3)-1/3*(-1)^(2/3)*f^(1/3)
*(a+b*arctanh(d*x+c))*ln(2/(d*x+c+1))/e^(4/3)-1/2*b*d*ln(d*x+c+1)/(1+c)/e
+1/3*f^(1/3)*(a+b*arctanh(d*x+c))*ln(2*d*(e^(1/3)+f^(1/3)*x)/(d*e^(1/3)+(1
-c)*f^(1/3)))/(d*x+c+1))/e^(4/3)-1/3*(-1)^(1/3)*f^(1/3)*(a+b*arctanh(d*x+c)
)*ln(2*d*(e^(1/3)-(-1)^(1/3)*f^(1/3)*x)/(d*e^(1/3)-(-1)^(1/3)*(1-c)*f^(1/3
)))/(d*x+c+1))/e^(4/3)+1/3*(-1)^(2/3)*f^(1/3)*(a+b*arctanh(d*x+c))*ln(2*d*(
e^(1/3)+(-1)^(2/3)*f^(1/3)*x)/(d*e^(1/3)+(-1)^(2/3)*(1-c)*f^(1/3)))/(d*x+c+
1))/e^(4/3)+1/6*b*f^(1/3)*polylog(2,1-2/(d*x+c+1))/e^(4/3)-1/6*(-1)^(1/3)*
b*f^(1/3)*polylog(2,1-2/(d*x+c+1))/e^(4/3)+1/6*(-1)^(2/3)*b*f^(1/3)*polylo
g(2,1-2/(d*x+c+1))/e^(4/3)-1/6*b*f^(1/3)*polylog(2,1-2*d*(e^(1/3)+f^(1/3)*
x)/(d*e^(1/3)+(1-c)*f^(1/3)))/(d*x+c+1))/e^(4/3)+1/6*(-1)^(1/3)*b*f^(1/3)*p
olylog(2,1-2*d*(e^(1/3)-(-1)^(1/3)*f^(1/3)*x)/(d*e^(1/3)-(-1)^(1/3)*(1-c)*
f^(1/3)))/(d*x+c+1))/e^(4/3)-1/6*(-1)^(2/3)*b*f^(1/3)*polylog(2,1-2*d*(e^(1
/3)+(-1)^(2/3)*f^(1/3)*x)/(d*e^(1/3)+(-1)^(2/3)*(1-c)*f^(1/3)))/(d*x+c+1))/
e^(4/3)

```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4 in optimal.

Time = 8.74 (sec) , antiderivative size = 2550, normalized size of antiderivative = 3.35

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e + fx^3)} dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])/(x^2*(e + f*x^3)),x]
```

output

```

((-6*a*e^(1/3))/x + 2*Sqrt[3]*a*f^(1/3)*ArcTan[(1 - (2*f^(1/3)*x)/e^(1/3))
/Sqrt[3]] + 2*a*f^(1/3)*Log[e^(1/3) + f^(1/3)*x] - a*f^(1/3)*Log[e^(2/3) -
e^(1/3)*f^(1/3)*x + f^(2/3)*x^2] - (6*b*e^(1/3)*((-1 + c^2 + c*d*x)*ArcTa
nh[c + d*x] + d*x*Log[-((d*x)/Sqrt[1 - (c + d*x)^2])]))/((-1 + c)*(1 + c)*
x) + b*d*e^(1/3)*f*(-4*ArcTanh[c + d*x]*RootSum[d^3*e - f - 3*c*f - 3*c^2*
f - c^3*f + 3*d^3*e**#1 + 3*f**#1 + 3*c*f**#1 - 3*c^2*f**#1 - 3*c^3*f**#1 + 3*d
^3*e**#1^2 - 3*f**#1^2 + 3*c*f**#1^2 + 3*c^2*f**#1^2 - 3*c^3*f**#1^2 + d^3*e**#1
^3 + f**#1^3 - 3*c*f**#1^3 + 3*c^2*f**#1^3 - c^3*f**#1^3 & , (ArcTanh[c + d*x]
+ c*ArcTanh[c + d*x] + Log[(-1 - c - d*x + #1 - c**#1 - d*x**#1)/Sqrt[1 - (
c + d*x)^2]] + c*Log[(-1 - c - d*x + #1 - c**#1 - d*x**#1)/Sqrt[1 - (c + d*x
)^2]] - ArcTanh[c + d*x]**#1 + c*ArcTanh[c + d*x]**#1 - Log[(-1 - c - d*x +
#1 - c**#1 - d*x**#1)/Sqrt[1 - (c + d*x)^2]]**#1 + c*Log[(-1 - c - d*x + #1 -
c**#1 - d*x**#1)/Sqrt[1 - (c + d*x)^2]]**#1)/(-(d^3*e) - f - c*f + c^2*f + c
^3*f - 2*d^3*e**#1 + 2*f**#1 - 2*c*f**#1 - 2*c^2*f**#1 + 2*c^3*f**#1 - d^3*e**#1
^2 - f**#1^2 + 3*c*f**#1^2 - 3*c^2*f**#1^2 + c^3*f**#1^2) & ] + 2*RootSum[d^3*
e - f - 3*c*f - 3*c^2*f - c^3*f + 3*d^3*e**#1 + 3*f**#1 + 3*c*f**#1 - 3*c^2*f
**#1 - 3*c^3*f**#1 + 3*d^3*e**#1^2 - 3*f**#1^2 + 3*c*f**#1^2 + 3*c^2*f**#1^2 - 3
*c^3*f**#1^2 + d^3*e**#1^3 + f**#1^3 - 3*c*f**#1^3 + 3*c^2*f**#1^3 - c^3*f**#1^3
& , (I*Pi*ArcTanh[c + d*x] + 2*ArcTanh[c + d*x]*ArcTanh[(1 - #1)/(1 + #1)
] - I*Pi*Log[1 + E^(2*ArcTanh[c + d*x])]) + 2*ArcTanh[c + d*x]*Log[1 - E...

```

Rubi [A] (verified)

Time = 1.92 (sec) , antiderivative size = 865, normalized size of antiderivative = 1.14, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e + fx^3)} dx$$

$$\downarrow 7276$$

$$\int \left(\frac{a}{x^2 (e + fx^3)} + \frac{b \operatorname{arctanh}(c + dx)}{x^2 (e + fx^3)} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& \frac{\sqrt[3]{f} \arctan\left(\frac{\sqrt[3]{e-2\sqrt[3]{fx}}}{\sqrt{3}\sqrt[3]{e}}\right) a}{\sqrt{3e^{4/3}}} + \frac{\sqrt[3]{f} \log\left(\sqrt[3]{fx} + \sqrt[3]{e}\right) a}{3e^{4/3}} - \\
& \frac{\sqrt[3]{f} \log\left(f^{2/3}x^2 - \sqrt[3]{e}\sqrt[3]{fx} + e^{2/3}\right) a}{6e^{4/3}} - \frac{a}{ex} - \frac{b \operatorname{arctanh}(c+dx)}{ex} + \frac{bd \log(x)}{(1-c^2)e} - \\
& \frac{bd \log(-c-dx+1)}{2(1-c)e} - \frac{(-1)^{2/3} b \sqrt[3]{f} \operatorname{arctanh}(c+dx) \log\left(\frac{2}{c+dx+1}\right)}{3e^{4/3}} + \\
& \frac{\sqrt[3]{-1} b \sqrt[3]{f} \operatorname{arctanh}(c+dx) \log\left(\frac{2}{c+dx+1}\right)}{3e^{4/3}} - \frac{b \sqrt[3]{f} \operatorname{arctanh}(c+dx) \log\left(\frac{2}{c+dx+1}\right)}{3e^{4/3}} - \\
& \frac{bd \log(c+dx+1)}{2(c+1)e} + \frac{b \sqrt[3]{f} \operatorname{arctanh}(c+dx) \log\left(\frac{2d\left(\sqrt[3]{fx} + \sqrt[3]{e}\right)}{\left(\sqrt[3]{f}(1-c) + d\sqrt[3]{e}\right)(c+dx+1)}\right)}{3e^{4/3}} - \\
& \frac{\sqrt[3]{-1} b \sqrt[3]{f} \operatorname{arctanh}(c+dx) \log\left(\frac{2d\left(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f}\right)(c+dx+1)}\right)}{3e^{4/3}} + \\
& \frac{(-1)^{2/3} b \sqrt[3]{f} \operatorname{arctanh}(c+dx) \log\left(\frac{2d\left((-1)^{2/3}\sqrt[3]{fx} + \sqrt[3]{e}\right)}{\left((-1)^{2/3}\sqrt[3]{f}(1-c) + d\sqrt[3]{e}\right)(c+dx+1)}\right)}{3e^{4/3}} + \\
& \frac{(-1)^{2/3} b \sqrt[3]{f} \operatorname{PolyLog}\left(2, 1 - \frac{2}{c+dx+1}\right)}{6e^{4/3}} - \frac{\sqrt[3]{-1} b \sqrt[3]{f} \operatorname{PolyLog}\left(2, 1 - \frac{2}{c+dx+1}\right)}{6e^{4/3}} + \\
& \frac{b \sqrt[3]{f} \operatorname{PolyLog}\left(2, 1 - \frac{2}{c+dx+1}\right)}{6e^{4/3}} - \frac{b \sqrt[3]{f} \operatorname{PolyLog}\left(2, 1 - \frac{2d\left(\sqrt[3]{fx} + \sqrt[3]{e}\right)}{\left(\sqrt[3]{f}(1-c) + d\sqrt[3]{e}\right)(c+dx+1)}\right)}{6e^{4/3}} + \\
& \frac{\sqrt[3]{-1} b \sqrt[3]{f} \operatorname{PolyLog}\left(2, 1 - \frac{2d\left(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx}\right)}{\left(d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f}\right)(c+dx+1)}\right)}{6e^{4/3}} - \\
& \frac{(-1)^{2/3} b \sqrt[3]{f} \operatorname{PolyLog}\left(2, 1 - \frac{2d\left((-1)^{2/3}\sqrt[3]{fx} + \sqrt[3]{e}\right)}{\left((-1)^{2/3}\sqrt[3]{f}(1-c) + d\sqrt[3]{e}\right)(c+dx+1)}\right)}{6e^{4/3}}
\end{aligned}$$

input

```
Int[(a + b*ArcTanh[c + d*x])/(x^2*(e + f*x^3)),x]
```

output

```

-(a/(e*x)) + (a*f^(1/3)*ArcTan[(e^(1/3) - 2*f^(1/3)*x)/(Sqrt[3]*e^(1/3))])
/(Sqrt[3]*e^(4/3)) - (b*ArcTanh[c + d*x]/(e*x) + (b*d*Log[x])/((1 - c^2)*
e) - (b*d*Log[1 - c - d*x]/(2*(1 - c)*e) - (b*f^(1/3)*ArcTanh[c + d*x]*Lo
g[2/(1 + c + d*x)])/(3*e^(4/3)) + ((-1)^(1/3)*b*f^(1/3)*ArcTanh[c + d*x]*L
og[2/(1 + c + d*x)])/(3*e^(4/3)) - ((-1)^(2/3)*b*f^(1/3)*ArcTanh[c + d*x]*
Log[2/(1 + c + d*x)])/(3*e^(4/3)) - (b*d*Log[1 + c + d*x]/(2*(1 + c)*e) +
(a*f^(1/3)*Log[e^(1/3) + f^(1/3)*x])/(3*e^(4/3)) + (b*f^(1/3)*ArcTanh[c +
d*x]*Log[(2*d*(e^(1/3) + f^(1/3)*x))/((d*e^(1/3) + (1 - c)*f^(1/3))*(1 +
c + d*x))])/(3*e^(4/3)) - ((-1)^(1/3)*b*f^(1/3)*ArcTanh[c + d*x]*Log[(2*d*
(e^(1/3) - (-1)^(1/3)*f^(1/3)*x))/((d*e^(1/3) - (-1)^(1/3)*(1 - c)*f^(1/3)
)*(1 + c + d*x))])/(3*e^(4/3)) + ((-1)^(2/3)*b*f^(1/3)*ArcTanh[c + d*x]*Lo
g[(2*d*(e^(1/3) + (-1)^(2/3)*f^(1/3)*x))/((d*e^(1/3) + (-1)^(2/3)*(1 - c)*
f^(1/3))*(1 + c + d*x))])/(3*e^(4/3)) - (a*f^(1/3)*Log[e^(2/3) - e^(1/3)*f
^(1/3)*x + f^(2/3)*x^2]/(6*e^(4/3)) + (b*f^(1/3)*PolyLog[2, 1 - 2/(1 + c
+ d*x)])/(6*e^(4/3)) - ((-1)^(1/3)*b*f^(1/3)*PolyLog[2, 1 - 2/(1 + c + d*x
)])/ (6*e^(4/3)) + ((-1)^(2/3)*b*f^(1/3)*PolyLog[2, 1 - 2/(1 + c + d*x)])/(
6*e^(4/3)) - (b*f^(1/3)*PolyLog[2, 1 - (2*d*(e^(1/3) + f^(1/3)*x))/((d*e^(
1/3) + (1 - c)*f^(1/3))*(1 + c + d*x))])/ (6*e^(4/3)) + ((-1)^(1/3)*b*f^(1/
3)*PolyLog[2, 1 - (2*d*(e^(1/3) - (-1)^(1/3)*f^(1/3)*x))/((d*e^(1/3) - (-1
)^(1/3)*(1 - c)*f^(1/3))*(1 + c + d*x))])/ (6*e^(4/3)) - ((-1)^(2/3)*b*f...

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.46 (sec) , antiderivative size = 513, normalized size of antiderivative = 0.67

| method | result |
|------------------|---|
| risch | $-\frac{db \ln(-dx)}{2e(-1+c)} + \frac{db \ln(-dx-c+1)}{2e(-1+c)} + \frac{b \ln(-dx-c+1)c}{2e(-1+c)x} - \frac{b \ln(-dx-c+1)}{2e(-1+c)x} - \frac{db}{\sqrt{-R1=\text{RootOf}(f_Z^3+(3fc-3f)_}}$ |
| derivativdivides | Expression too large to display |
| default | Expression too large to display |
| parts | Expression too large to display |

input `int((a+b*arctanh(d*x+c))/x^2/(f*x^3+e),x,method=_RETURNVERBOSE)`

output

```
-1/2*d*b/e/(-1+c)*ln(-d*x)+1/2*d*b/e*ln(-d*x-c+1)/(-1+c)+1/2*b/e*ln(-d*x-c+1)/(-1+c)/x*c-1/2*b/e*ln(-d*x-c+1)/(-1+c)/x-1/6*d*b/e*sum(1/(_R1-1+c)*(ln(-d*x-c+1)*ln((d*x+_R1+c-1)/_R1)+dilog((d*x+_R1+c-1)/_R1)),_R1=RootOf(f*_Z^3+(3*c*f-3*f)*_Z^2+(3*c^2*f-6*c*f+3*f)*_Z+c^3*f-d^3*e-3*c^2*f+3*f*c-f))-a/e/x+1/3*d*a*sum((c+_R-1)/(_R^2+2*_R*c+c^2-2*_R-2*c+1)*ln(-d*x-_R-c+1),_R=RootOf(f*_Z^3+(3*c*f-3*f)*_Z^2+(3*c^2*f-6*c*f+3*f)*_Z+c^3*f-d^3*e-3*c^2*f+3*f*c-f))/e+1/2*b*d/e/(1+c)*ln(d*x)-1/2*b*d*ln(d*x+c+1)/(1+c)/e-1/2*b/e*ln(d*x+c+1)/(1+c)/x*c-1/2*b/e*ln(d*x+c+1)/(1+c)/x-1/6*b*d/e*sum(1/(_R1-1-c)*(ln(d*x+c+1)*ln((-d*x+_R1-c-1)/_R1)+dilog((-d*x+_R1-c-1)/_R1)),_R1=RootOf(f*_Z^3+(-3*c*f-3*f)*_Z^2+(3*c^2*f+6*c*f+3*f)*_Z-c^3*f+d^3*e-3*c^2*f-3*f*c-f))
```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e + fx^3)} dx = \int \frac{b \operatorname{arctanh}(dx + c) + a}{(fx^3 + e)x^2} dx$$

input `integrate((a+b*arctanh(d*x+c))/x^2/(f*x^3+e),x, algorithm="fricas")`

output

```
integral((b*arctanh(d*x + c) + a)/(f*x^5 + e*x^2), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e + fx^3)} dx = \text{Timed out}$$

input `integrate((a+b*atanh(d*x+c))/x**2/(f*x**3+e),x)`

output Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e + fx^3)} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arctanh(d*x+c))/x^2/(f*x^3+e),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e + fx^3)} dx = \int \frac{b \operatorname{arctanh}(dx + c) + a}{(fx^3 + e)x^2} dx$$

input `integrate((a+b*arctanh(d*x+c))/x^2/(f*x^3+e),x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)/((f*x^3 + e)*x^2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e + fx^3)} dx = \int \frac{a + b \operatorname{atanh}(c + dx)}{x^2 (fx^3 + e)} dx$$

input `int((a + b*atanh(c + d*x))/(x^2*(e + f*x^3)),x)`

output `int((a + b*atanh(c + d*x))/(x^2*(e + f*x^3)), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^2 (e + fx^3)} dx$$

$$= \frac{2\sqrt{3} \operatorname{atan}\left(\frac{e^{\frac{1}{3}} - 2f^{\frac{1}{3}}x}{e^{\frac{1}{3}}\sqrt{3}}\right) a f x + 6f^{\frac{2}{3}}e^{\frac{4}{3}}\left(\int \frac{\operatorname{atanh}(dx+c)}{fx^5+ex^2} dx\right) b x - 6f^{\frac{2}{3}}e^{\frac{1}{3}}a - \log\left(e^{\frac{2}{3}} - f^{\frac{1}{3}}e^{\frac{1}{3}}x + f^{\frac{2}{3}}x^2\right) a f x + 21}{6f^{\frac{2}{3}}e^{\frac{4}{3}}x}$$

input `int((a+b*atanh(d*x+c))/x^2/(f*x^3+e),x)`

output `(2*sqrt(3)*atan((e**(1/3) - 2*f**(1/3)*x)/(e**(1/3)*sqrt(3)))*a*f*x + 6*f**
*(2/3)*e**(1/3)*int(atanh(c + d*x)/(e*x**2 + f*x**5),x)*b*e*x - 6*f**(2/3)
*e**(1/3)*a - log(e**(2/3) - f**(1/3)*e**(1/3)*x + f**(2/3)*x**2)*a*f*x +
2*log(e**(1/3) + f**(1/3)*x)*a*f*x)/(6*f**(2/3)*e**(1/3)*e*x)`

$$3.76 \quad \int \frac{a+b \operatorname{arctanh}(c+dx)}{x^3(e+fx^3)} dx$$

| | |
|---|-----|
| Optimal result | 665 |
| Mathematica [C] (warning: unable to verify) | 666 |
| Rubi [A] (warning: unable to verify) | 667 |
| Maple [C] (warning: unable to verify) | 669 |
| Fricas [F] | 671 |
| Sympy [F(-1)] | 671 |
| Maxima [F(-2)] | 672 |
| Giac [F] | 672 |
| Mupad [F(-1)] | 672 |
| Reduce [F] | 673 |

Optimal result

Integrand size = 23, antiderivative size = 791

$$\begin{aligned}
& \int \frac{a + \operatorname{barctanh}(c + dx)}{x^3 (e + fx^3)} dx \\
&= -\frac{bd}{2(1-c^2)ex} - \frac{a + \operatorname{barctanh}(c + dx)}{2ex^2} + \frac{bcd^2 \log(x)}{(1-c^2)^2 e} \\
&\quad - \frac{bd^2 \log(1-c-dx)}{4(1-c)^2 e} + \frac{f^{2/3}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{3e^{5/3}} \\
&\quad - \frac{\sqrt[3]{-1} f^{2/3}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{3e^{5/3}} \\
&\quad + \frac{(-1)^{2/3} f^{2/3}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2}{1+c+dx}\right)}{3e^{5/3}} + \frac{bd^2 \log(1+c+dx)}{4(1+c)^2 e} \\
&\quad - \frac{f^{2/3}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d(\sqrt[3]{e} + \sqrt[3]{fx})}{(d\sqrt[3]{e} + (1-c)\sqrt[3]{f})(1+c+dx)}\right)}{3e^{5/3}} \\
&\quad - \frac{(-1)^{2/3} f^{2/3}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx})}{(d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f})(1+c+dx)}\right)}{3e^{5/3}} \\
&\quad + \frac{\sqrt[3]{-1} f^{2/3}(a + \operatorname{barctanh}(c + dx)) \log\left(\frac{2d(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx})}{(d\sqrt[3]{e} + (-1)^{2/3}(1-c)\sqrt[3]{f})(1+c+dx)}\right)}{3e^{5/3}} \\
&\quad - \frac{bf^{2/3} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{6e^{5/3}} + \frac{\sqrt[3]{-1} bf^{2/3} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{6e^{5/3}} \\
&\quad - \frac{(-1)^{2/3} bf^{2/3} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+c+dx}\right)}{6e^{5/3}} \\
&\quad + \frac{bf^{2/3} \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt[3]{e} + \sqrt[3]{fx})}{(d\sqrt[3]{e} + (1-c)\sqrt[3]{f})(1+c+dx)}\right)}{6e^{5/3}} \\
&\quad + \frac{(-1)^{2/3} bf^{2/3} \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx})}{(d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f})(1+c+dx)}\right)}{6e^{5/3}} \\
&\quad + \frac{\sqrt[3]{-1} bf^{2/3} \operatorname{PolyLog}\left(2, 1 - \frac{2d(\sqrt[3]{e} + (-1)^{2/3}\sqrt[3]{fx})}{(d\sqrt[3]{e} + (-1)^{2/3}(1-c)\sqrt[3]{f})(1+c+dx)}\right)}{6e^{5/3}}
\end{aligned}$$

output

```

-1/2*b*d/(-c^2+1)/e/x-1/2*(a+b*arctanh(d*x+c))/e/x^2+b*c*d^2*ln(x)/(-c^2+1
)^2/e-1/4*b*d^2*ln(-d*x-c+1)/(1-c)^2/e+1/3*f^(2/3)*(a+b*arctanh(d*x+c))*ln
(2/(d*x+c+1))/e^(5/3)-1/3*(-1)^(1/3)*f^(2/3)*(a+b*arctanh(d*x+c))*ln(2/(d*
x+c+1))/e^(5/3)+1/3*(-1)^(2/3)*f^(2/3)*(a+b*arctanh(d*x+c))*ln(2/(d*x+c+1)
)/e^(5/3)+1/4*b*d^2*ln(d*x+c+1)/(1+c)^2/e-1/3*f^(2/3)*(a+b*arctanh(d*x+c))
*ln(2*d*(e^(1/3)+f^(1/3)*x)/(d*e^(1/3)+(1-c)*f^(1/3))/(d*x+c+1))/e^(5/3)-1
/3*(-1)^(2/3)*f^(2/3)*(a+b*arctanh(d*x+c))*ln(2*d*(e^(1/3)-(-1)^(1/3)*f^(1
/3)*x)/(d*e^(1/3)-(-1)^(1/3)*(1-c)*f^(1/3))/(d*x+c+1))/e^(5/3)+1/3*(-1)^(1
/3)*f^(2/3)*(a+b*arctanh(d*x+c))*ln(2*d*(e^(1/3)+(-1)^(2/3)*f^(1/3)*x)/(d*
e^(1/3)+(-1)^(2/3)*(1-c)*f^(1/3))/(d*x+c+1))/e^(5/3)-1/6*b*f^(2/3)*polylog
(2,1-2/(d*x+c+1))/e^(5/3)+1/6*(-1)^(1/3)*b*f^(2/3)*polylog(2,1-2/(d*x+c+1)
)/e^(5/3)-1/6*(-1)^(2/3)*b*f^(2/3)*polylog(2,1-2/(d*x+c+1))/e^(5/3)+1/6*b*
f^(2/3)*polylog(2,1-2*d*(e^(1/3)+f^(1/3)*x)/(d*e^(1/3)+(1-c)*f^(1/3))/(d*x
+c+1))/e^(5/3)+1/6*(-1)^(2/3)*b*f^(2/3)*polylog(2,1-2*d*(e^(1/3)-(-1)^(1/3)
)*f^(1/3)*x)/(d*e^(1/3)-(-1)^(1/3)*(1-c)*f^(1/3))/(d*x+c+1))/e^(5/3)-1/6*(
-1)^(1/3)*b*f^(2/3)*polylog(2,1-2*d*(e^(1/3)+(-1)^(2/3)*f^(1/3)*x)/(d*e^(1
/3)+(-1)^(2/3)*(1-c)*f^(1/3))/(d*x+c+1))/e^(5/3)

```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4 in optimal.

Time = 7.37 (sec) , antiderivative size = 1564, normalized size of antiderivative = 1.98

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e + fx^3)} dx = \text{Too large to display}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])/(x^3*(e + f*x^3)),x]
```

output

```

((-3*a*e^(2/3))/x^2 + 2*Sqrt[3]*a*f^(2/3)*ArcTan[(1 - (2*f^(1/3)*x)/e^(1/3
))/Sqrt[3]] - 2*a*f^(2/3)*Log[e^(1/3) + f^(1/3)*x] + a*f^(2/3)*Log[e^(2/3)
- e^(1/3)*f^(1/3)*x + f^(2/3)*x^2] + (3*b*e^(2/3)*(c*(-1 - c^4 + d^2*x^2
+ c^2*(2 + d^2*x^2))*ArcTanh[c + d*x] + d*x*((-1 + c^2)*(c + d*x) + 2*c^2*
d*x*Log[-((d*x)/Sqrt[1 - (c + d*x)^2])])))/((-1 + c)^2*c*(1 + c)^2*x^2) -
b*d^2*e^(2/3)*f*(4*ArcTanh[c + d*x]*RootSum[d^3*e - f - 3*c*f - 3*c^2*f -
c^3*f + 3*d^3*e**#1 + 3*f**#1 + 3*c*f**#1 - 3*c^2*f**#1 - 3*c^3*f**#1 + 3*d^3*e
**#1^2 - 3*f**#1^2 + 3*c*f**#1^2 + 3*c^2*f**#1^2 - 3*c^3*f**#1^2 + d^3*e**#1^3 +
f**#1^3 - 3*c*f**#1^3 + 3*c^2*f**#1^3 - c^3*f**#1^3 & , (ArcTanh[c + d*x] + L
og[(-1 - c - d*x + #1 - c**#1 - d*x**#1)/Sqrt[1 - (c + d*x)^2]] + ArcTanh[c
+ d*x]**#1 + Log[(-1 - c - d*x + #1 - c**#1 - d*x**#1)/Sqrt[1 - (c + d*x)^2]]
**#1)/(d^3*e + f + c*f - c^2*f - c^3*f + 2*d^3*e**#1 - 2*f**#1 + 2*c*f**#1 + 2
*c^2*f**#1 - 2*c^3*f**#1 + d^3*e**#1^2 + f**#1^2 - 3*c*f**#1^2 + 3*c^2*f**#1^2 -
c^3*f**#1^2) & ] + RootSum[d^3*e - f - 3*c*f - 3*c^2*f - c^3*f + 3*d^3*e**#
1 + 3*f**#1 + 3*c*f**#1 - 3*c^2*f**#1 - 3*c^3*f**#1 + 3*d^3*e**#1^2 - 3*f**#1^2
+ 3*c*f**#1^2 + 3*c^2*f**#1^2 - 3*c^3*f**#1^2 + d^3*e**#1^3 + f**#1^3 - 3*c*f**#
1^3 + 3*c^2*f**#1^3 - c^3*f**#1^3 & , (I*Pi*ArcTanh[c + d*x] + 2*ArcTanh[c +
d*x]*ArcTanh[(1 - #1)/(1 + #1)] - I*Pi*Log[1 + E^(2*ArcTanh[c + d*x])]) +
2*ArcTanh[c + d*x]*Log[1 - E^(-2*(ArcTanh[c + d*x] + ArcTanh[(1 - #1)/(1 +
#1])))] + 2*ArcTanh[(1 - #1)/(1 + #1)]*Log[1 - E^(-2*(ArcTanh[c + d*x]...

```

Rubi [A] (warning: unable to verify)

Time = 2.17 (sec) , antiderivative size = 1028, normalized size of antiderivative = 1.30, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e + fx^3)} dx$$

$$\downarrow 7276$$

$$\int \left(\frac{a}{x^3 (e + fx^3)} + \frac{b \operatorname{arctanh}(c + dx)}{x^3 (e + fx^3)} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& \frac{bc \log(x)d^2}{(1-c^2)^2 e} - \frac{b \log(-c-dx+1)d^2}{4(1-c)^2 e} + \frac{b \log(c+dx+1)d^2}{4(c+1)^2 e} - \frac{bd}{2(1-c^2)ex} + \\
& \frac{af^{2/3} \arctan\left(\frac{\sqrt[3]{e-2}\sqrt[3]{fx}}{\sqrt{3}\sqrt[3]{e}}\right)}{\sqrt{3}e^{5/3}} - \frac{b \operatorname{arctanh}(c+dx)}{2ex^2} - \frac{af^{2/3} \log\left(\sqrt[3]{fx} + \sqrt[3]{e}\right)}{3e^{5/3}} + \\
& \frac{bf^{2/3} \log(-c-dx+1) \log\left(\frac{d\left(\sqrt[3]{fx} + \sqrt[3]{e}\right)}{\sqrt[3]{f(1-c)} + d\sqrt[3]{e}}\right)}{6e^{5/3}} - \frac{bf^{2/3} \log(c+dx+1) \log\left(\frac{d\left(\sqrt[3]{fx} + \sqrt[3]{e}\right)}{d\sqrt[3]{e} - (c+1)\sqrt[3]{f}}\right)}{6e^{5/3}} + \\
& \frac{(-1)^{2/3} bf^{2/3} \log(-c-dx+1) \log\left(\frac{d\left(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx}\right)}{d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f}}\right)}{6e^{5/3}} - \\
& \frac{(-1)^{2/3} bf^{2/3} \log(c+dx+1) \log\left(\frac{d\left(\sqrt[3]{e} - \sqrt[3]{-1}\sqrt[3]{fx}\right)}{\sqrt[3]{-1}\sqrt[3]{f(c+1)} + d\sqrt[3]{e}}\right)}{6e^{5/3}} - \\
& \frac{\sqrt[3]{-1} bf^{2/3} \log(-c-dx+1) \log\left(\frac{d\left((-1)^{2/3}\sqrt[3]{fx} + \sqrt[3]{e}\right)}{(-1)^{2/3}\sqrt[3]{f(1-c)} + d\sqrt[3]{e}}\right)}{6e^{5/3}} + \\
& \frac{\sqrt[3]{-1} bf^{2/3} \log(c+dx+1) \log\left(\frac{d\left((-1)^{2/3}\sqrt[3]{fx} + \sqrt[3]{e}\right)}{d\sqrt[3]{e} - (-1)^{2/3}(c+1)\sqrt[3]{f}}\right)}{6e^{5/3}} + \\
& \frac{af^{2/3} \log\left(f^{2/3}x^2 - \sqrt[3]{e}\sqrt[3]{fx} + e^{2/3}\right)}{6e^{5/3}} + \frac{bf^{2/3} \operatorname{PolyLog}\left(2, \frac{\sqrt[3]{f(-c-dx+1)}}{\sqrt[3]{f(1-c)} + d\sqrt[3]{e}}\right)}{6e^{5/3}} + \\
& \frac{(-1)^{2/3} bf^{2/3} \operatorname{PolyLog}\left(2, -\frac{\sqrt[3]{-1}\sqrt[3]{f(-c-dx+1)}}{d\sqrt[3]{e} - \sqrt[3]{-1}(1-c)\sqrt[3]{f}}\right)}{6e^{5/3}} - \\
& \frac{\sqrt[3]{-1} bf^{2/3} \operatorname{PolyLog}\left(2, \frac{(-1)^{2/3}\sqrt[3]{f(-c-dx+1)}}{(-1)^{2/3}\sqrt[3]{f(1-c)} + d\sqrt[3]{e}}\right)}{6e^{5/3}} - \frac{bf^{2/3} \operatorname{PolyLog}\left(2, -\frac{\sqrt[3]{f(c+dx+1)}}{d\sqrt[3]{e} - (c+1)\sqrt[3]{f}}\right)}{6e^{5/3}} - \\
& \frac{(-1)^{2/3} bf^{2/3} \operatorname{PolyLog}\left(2, \frac{\sqrt[3]{-1}\sqrt[3]{f(c+dx+1)}}{\sqrt[3]{-1}\sqrt[3]{f(c+1)} + d\sqrt[3]{e}}\right)}{6e^{5/3}} + \\
& \frac{\sqrt[3]{-1} bf^{2/3} \operatorname{PolyLog}\left(2, -\frac{(-1)^{2/3}\sqrt[3]{f(c+dx+1)}}{d\sqrt[3]{e} - (-1)^{2/3}(c+1)\sqrt[3]{f}}\right)}{6e^{5/3}} - \frac{a}{2ex^2}
\end{aligned}$$

input

```
Int[(a + b*ArcTanh[c + d*x])/(x^3*(e + f*x^3)),x]
```

output

```

-1/2*a/(e*x^2) - (b*d)/(2*(1 - c^2)*e*x) + (a*f^(2/3)*ArcTan[(e^(1/3) - 2*
f^(1/3)*x)/(Sqrt[3]*e^(1/3))]/(Sqrt[3]*e^(5/3)) - (b*ArcTanh[c + d*x]/(2
*e*x^2) + (b*c*d^2*Log[x])/((1 - c^2)^2*e) - (b*d^2*Log[1 - c - d*x]/(4*(
1 - c)^2*e) + (b*d^2*Log[1 + c + d*x]/(4*(1 + c)^2*e) - (a*f^(2/3)*Log[e^
(1/3) + f^(1/3)*x]/(3*e^(5/3)) + (b*f^(2/3)*Log[1 - c - d*x]*Log[(d*(e^(1
/3) + f^(1/3)*x))/(d*e^(1/3) + (1 - c)*f^(1/3))]/(6*e^(5/3)) - (b*f^(2/3)
*Log[1 + c + d*x]*Log[(d*(e^(1/3) + f^(1/3)*x))/(d*e^(1/3) - (1 + c)*f^(1
/3))]/(6*e^(5/3)) + ((-1)^(2/3)*b*f^(2/3)*Log[1 - c - d*x]*Log[(d*(e^(1/3)
- (-1)^(1/3)*f^(1/3)*x))/(d*e^(1/3) - (-1)^(1/3)*(1 - c)*f^(1/3))]/(6*e^
(5/3)) - ((-1)^(2/3)*b*f^(2/3)*Log[1 + c + d*x]*Log[(d*(e^(1/3) - (-1)^(1
/3)*f^(1/3)*x))/(d*e^(1/3) + (-1)^(1/3)*(1 + c)*f^(1/3))]/(6*e^(5/3)) - ((
-1)^(1/3)*b*f^(2/3)*Log[1 - c - d*x]*Log[(d*(e^(1/3) + (-1)^(2/3)*f^(1/3)*
x))/(d*e^(1/3) + (-1)^(2/3)*(1 - c)*f^(1/3))]/(6*e^(5/3)) + ((-1)^(1/3)*b
*f^(2/3)*Log[1 + c + d*x]*Log[(d*(e^(1/3) + (-1)^(2/3)*f^(1/3)*x))/(d*e^(1
/3) - (-1)^(2/3)*(1 + c)*f^(1/3))]/(6*e^(5/3)) + (a*f^(2/3)*Log[e^(2/3) -
e^(1/3)*f^(1/3)*x + f^(2/3)*x^2]/(6*e^(5/3)) + (b*f^(2/3)*PolyLog[2, (f^
(1/3)*(1 - c - d*x))/(d*e^(1/3) + (1 - c)*f^(1/3))]/(6*e^(5/3)) + ((-1)^(
2/3)*b*f^(2/3)*PolyLog[2, -(((-1)^(1/3)*f^(1/3)*(1 - c - d*x))/(d*e^(1/3)
- (-1)^(1/3)*(1 - c)*f^(1/3)))]/(6*e^(5/3)) - ((-1)^(1/3)*b*f^(2/3)*PolyL
og[2, ((-1)^(2/3)*f^(1/3)*(1 - c - d*x))/(d*e^(1/3) + (-1)^(2/3)*(1 - c...

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.53 (sec) , antiderivative size = 662, normalized size of antiderivative = 0.84

| method | result |
|-------------------|--|
| risch | $\frac{d^2 b \ln(-dx)}{4e(-1+c)^2} - \frac{db}{4e(-1+c)^2 x} + \frac{dbc}{4e(-1+c)^2 x} - \frac{d^2 b \ln(-dx-c+1)}{4e(-1+c)^2} + \frac{b \ln(-dx-c+1)c^2}{4e x^2(-1+c)^2} - \frac{b \ln(-dx-c+1)c}{2e x^2(-1+c)^2} +$ |
| derivativedivides | $d^2 \left(\frac{a \left(\frac{\sum_{-R=\text{RootOf}(f_Z^3 - 3fc_Z^2 + 3c^2 f_Z - c^3 f + d^3 e)} \ln(dx - R + c)}{-R^2 + 2Rc - c^2} \right)}{3e} - \frac{a}{2e d^2 x^2} - \frac{b \operatorname{arctanh}(dx+c)}{2e d^2 x^2} \right)$ |
| default | $d^2 \left(\frac{a \left(\frac{\sum_{-R=\text{RootOf}(f_Z^3 - 3fc_Z^2 + 3c^2 f_Z - c^3 f + d^3 e)} \ln(dx - R + c)}{-R^2 + 2Rc - c^2} \right)}{3e} - \frac{a}{2e d^2 x^2} - \frac{b \operatorname{arctanh}(dx+c)}{2e d^2 x^2} \right)$ |
| parts | $-\frac{a \ln\left(x + \left(\frac{e}{f}\right)^{\frac{1}{3}}\right)}{3e \left(\frac{e}{f}\right)^{\frac{2}{3}}} + \frac{a \ln\left(x^2 - \left(\frac{e}{f}\right)^{\frac{1}{3}} x + \left(\frac{e}{f}\right)^{\frac{2}{3}}\right)}{6e \left(\frac{e}{f}\right)^{\frac{2}{3}}} - \frac{a\sqrt{3} \operatorname{arctan}\left(\frac{\sqrt{3} \left(\frac{2x}{\left(\frac{e}{f}\right)^{\frac{1}{3}} - 1\right)}\right)}{3e \left(\frac{e}{f}\right)^{\frac{2}{3}}} - \frac{a}{2e x^2} - \frac{b \operatorname{arctanh}(dx+c)}{2e x^2}$ |

input `int((a+b*arctanh(d*x+c))/x^3/(f*x^3+e),x,method=_RETURNVERBOSE)`

output

```

1/4*d^2*b/e/(-1+c)^2*ln(-d*x)-1/4*d*b/e/(-1+c)^2/x+1/4*d*b/e/(-1+c)^2/x*c-
1/4*d^2*b/e*ln(-d*x-c+1)/(-1+c)^2+1/4*b/e*ln(-d*x-c+1)/x^2/(-1+c)^2*c^2-1/
2*b/e*ln(-d*x-c+1)/x^2/(-1+c)^2*c+1/4*b/e*ln(-d*x-c+1)/x^2/(-1+c)^2+1/6*d^
2*b/e*sum(1/(_R1^2+2*_R1*c+c^2-2*_R1-2*c+1)*(ln(-d*x-c+1)*ln((d*x+_R1+c-1)
/_R1)+dilog((d*x+_R1+c-1)/_R1)),_R1=RootOf(f*_Z^3+(3*c*f-3*f)*_Z^2+(3*c^2*
f-6*c*f+3*f)*_Z+c^3*f-d^3*e-3*c^2*f+3*f*c-f))-1/2*a/e/x^2-1/3*d^2*a*sum(1/
(_R^2+2*_R*c+c^2-2*_R-2*c+1)*ln(-d*x-_R-c+1),_R=RootOf(f*_Z^3+(3*c*f-3*f)*
_Z^2+(3*c^2*f-6*c*f+3*f)*_Z+c^3*f-d^3*e-3*c^2*f+3*f*c-f))/e-1/4*b*d^2/e/(1
+c)^2*ln(d*x)-1/4*b*d/e/(1+c)^2/x-1/4*b*d/e/(1+c)^2/x*c+1/4*b*d^2*ln(d*x+c
+1)/(1+c)^2/e-1/4*b/e*ln(d*x+c+1)/x^2/(1+c)^2*c^2-1/2*b/e*ln(d*x+c+1)/x^2/
(1+c)^2*c-1/4*b/e*ln(d*x+c+1)/x^2/(1+c)^2-1/6*b*d^2/e*sum(1/(_R1^2-2*_R1*c
+c^2-2*_R1+2*c+1)*(ln(d*x+c+1)*ln((-d*x+_R1-c-1)/_R1)+dilog((-d*x+_R1-c-1)
/_R1)),_R1=RootOf(f*_Z^3+(-3*c*f-3*f)*_Z^2+(3*c^2*f+6*c*f+3*f)*_Z-c^3*f+d^
3*e-3*c^2*f-3*f*c-f))

```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e + fx^3)} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{(fx^3 + e)x^3} dx$$

input

```
integrate((a+b*arctanh(d*x+c))/x^3/(f*x^3+e),x, algorithm="fricas")
```

output

```
integral((b*arctanh(d*x + c) + a)/(f*x^6 + e*x^3), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e + fx^3)} dx = \text{Timed out}$$

input

```
integrate((a+b*atanh(d*x+c))/x**3/(f*x**3+e),x)
```

output

```
Timed out
```


Maxima [F(-2)]

Exception generated.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e + fx^3)} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arctanh(d*x+c))/x^3/(f*x^3+e),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e + fx^3)} dx = \int \frac{b \operatorname{arctanh}(dx + c) + a}{(fx^3 + e)x^3} dx$$

input `integrate((a+b*arctanh(d*x+c))/x^3/(f*x^3+e),x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)/((f*x^3 + e)*x^3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e + fx^3)} dx = \int \frac{a + b \operatorname{atanh}(c + dx)}{x^3 (fx^3 + e)} dx$$

input `int((a + b*atanh(c + d*x))/(x^3*(e + f*x^3)),x)`

output `int((a + b*atanh(c + d*x))/(x^3*(e + f*x^3)), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{x^3 (e + fx^3)} dx = \text{too large to display}$$

input `int((a+b*atanh(d*x+c))/x^3/(f*x^3+e),x)`

output `(6*e**(1/3)*sqrt(3)*atan((e**(1/3) - 2*f**(1/3)*x)/(e**(1/3)*sqrt(3)))*a*c**4*f*x**2 - 4*e**(1/3)*sqrt(3)*atan((e**(1/3) - 2*f**(1/3)*x)/(e**(1/3)*sqrt(3)))*a*c**2*f*x**2 - 2*e**(1/3)*sqrt(3)*atan((e**(1/3) - 2*f**(1/3)*x)/(e**(1/3)*sqrt(3)))*a*f*x**2 + 6*f**(1/3)*atanh(c + d*x)**2*b*c**3*d**2*e*x**2 - 6*f**(1/3)*atanh(c + d*x)**2*b*c*d**2*e*x**2 + 3*f**(1/3)*atanh(c + d*x)*b*c**4*e - 12*f**(1/3)*atanh(c + d*x)*b*c**3*d*e*x - 15*f**(1/3)*atanh(c + d*x)*b*c**2*d**2*e*x**2 - 6*f**(1/3)*atanh(c + d*x)*b*c**2*e + 18*f**(1/3)*atanh(c + d*x)*b*c*d**2*e*x**2 + 12*f**(1/3)*atanh(c + d*x)*b*c*d*e*x - 3*f**(1/3)*atanh(c + d*x)*b*d**2*e*x**2 + 3*f**(1/3)*atanh(c + d*x)*b*e + 3*e**(1/3)*log(e**(2/3) - f**(1/3)*e**(1/3)*x + f**(2/3)*x**2)*a*c**4*f*x**2 - 2*e**(1/3)*log(e**(2/3) - f**(1/3)*e**(1/3)*x + f**(2/3)*x**2)*a*c**2*f*x**2 - e**(1/3)*log(e**(2/3) - f**(1/3)*e**(1/3)*x + f**(2/3)*x**2)*a*f*x**2 - 6*e**(1/3)*log(e**(1/3) + f**(1/3)*x)*a*c**4*f*x**2 + 4*e**(1/3)*log(e**(1/3) + f**(1/3)*x)*a*c**2*f*x**2 + 2*e**(1/3)*log(e**(1/3) + f**(1/3)*x)*a*f*x**2 + 72*f**(1/3)*int(atanh(c + d*x)/(3*c**4*e*x**3 + 3*c**4*f*x**6 + 6*c**3*d*e*x**4 + 6*c**3*d*f*x**7 + 3*c**2*d**2*e*x**5 + 3*c**2*d**2*f*x**8 - 2*c**2*e*x**3 - 2*c**2*f*x**6 + 2*c*d*e*x**4 + 2*c*d*f*x**7 + d**2*e*x**5 + d**2*f*x**8 - e*x**3 - f*x**6),x)*b*c**8*e**2*x**2 - 120*f**(1/3)*int(atanh(c + d*x)/(3*c**4*e*x**3 + 3*c**4*f*x**6 + 6*c**3*d*e*x**4 + 6*c**3*d*f*x**7 + 3*c**2*d**2*e*x**5 + 3*c**2*d**2*f*x**8 - 2*c...`

3.77 $\int \frac{a+b\operatorname{arctanh}(c+dx)}{e+fx+gx^2} dx$

| | |
|----------------------------|-----|
| Optimal result | 674 |
| Mathematica [A] (verified) | 675 |
| Rubi [A] (verified) | 676 |
| Maple [B] (verified) | 677 |
| Fricas [F] | 678 |
| Sympy [F(-1)] | 679 |
| Maxima [F(-2)] | 679 |
| Giac [F] | 679 |
| Mupad [F(-1)] | 680 |
| Reduce [F] | 680 |

Optimal result

Integrand size = 23, antiderivative size = 366

$$\begin{aligned}
 & \int \frac{a + b\operatorname{arctanh}(c + dx)}{e + fx + gx^2} dx \\
 &= \frac{(a + b\operatorname{arctanh}(c + dx)) \log\left(-\frac{2(2cg-d(f-\sqrt{f^2-4eg})-2g(c+dx))}{(df+2g-2cg-d\sqrt{f^2-4eg})(1+c+dx)}\right)}{\sqrt{f^2-4eg}} \\
 &\quad - \frac{(a + b\operatorname{arctanh}(c + dx)) \log\left(-\frac{2(2cg-d(f+\sqrt{f^2-4eg})-2g(c+dx))}{(2(1-c)g+d(f+\sqrt{f^2-4eg}))(1+c+dx)}\right)}{\sqrt{f^2-4eg}} \\
 &\quad - \frac{b \operatorname{PolyLog}\left(2, 1 + \frac{2(2cg-d(f-\sqrt{f^2-4eg})-2g(c+dx))}{(df+2g-2cg-d\sqrt{f^2-4eg})(1+c+dx)}\right)}{2\sqrt{f^2-4eg}} \\
 &\quad + \frac{b \operatorname{PolyLog}\left(2, 1 + \frac{2(2cg-d(f+\sqrt{f^2-4eg})-2g(c+dx))}{(2(1-c)g+d(f+\sqrt{f^2-4eg}))(1+c+dx)}\right)}{2\sqrt{f^2-4eg}}
 \end{aligned}$$

output

```
(a+b*arctanh(d*x+c))*ln((-4*c*g+2*d*(f-(-4*e*g+f^2)^(1/2))+4*g*(d*x+c))/(d*f+2*g-2*c*g-d*(-4*e*g+f^2)^(1/2))/(d*x+c+1))/(-4*e*g+f^2)^(1/2)-(a+b*arctanh(d*x+c))*ln((-4*c*g+2*d*(f+(-4*e*g+f^2)^(1/2))+4*g*(d*x+c))/(2*(1-c)*g+d*(f+(-4*e*g+f^2)^(1/2)))/(d*x+c+1))/(-4*e*g+f^2)^(1/2)-1/2*b*polylog(2,1+2*(2*c*g-d*(f-(-4*e*g+f^2)^(1/2))-2*g*(d*x+c))/(d*f+2*g-2*c*g-d*(-4*e*g+f^2)^(1/2))/(d*x+c+1))/(-4*e*g+f^2)^(1/2)+1/2*b*polylog(2,1+2*(2*c*g-d*(f+(-4*e*g+f^2)^(1/2))-2*g*(d*x+c))/(2*(1-c)*g+d*(f+(-4*e*g+f^2)^(1/2)))/(d*x+c+1))/(-4*e*g+f^2)^(1/2)
```

Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 437, normalized size of antiderivative = 1.19

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx + gx^2} dx =$$

$$\frac{4a \operatorname{arctanh}\left(\frac{f+2gx}{\sqrt{f^2-4eg}}\right) - b \log(1+c+dx) \log\left(\frac{d(-f+\sqrt{f^2-4eg}-2gx)}{2(1+c)g+d(-f+\sqrt{f^2-4eg})}\right) + b \log(1-c-dx) \log\left(\frac{d(f-df+2g}{df+2g}\right)}{-}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])/(e + f*x + g*x^2),x]
```

output

```
-1/2*(4*a*ArcTanh[(f + 2*g*x)/Sqrt[f^2 - 4*e*g]] - b*Log[1 + c + d*x]*Log[(d*(-f + Sqrt[f^2 - 4*e*g] - 2*g*x))/(2*(1 + c)*g + d*(-f + Sqrt[f^2 - 4*e*g]))] + b*Log[1 - c - d*x]*Log[(d*(f - Sqrt[f^2 - 4*e*g] + 2*g*x))/(d*f + 2*g - 2*c*g - d*Sqrt[f^2 - 4*e*g])] - b*Log[1 - c - d*x]*Log[(d*(f + Sqrt[f^2 - 4*e*g] + 2*g*x))/(-2*(-1 + c)*g + d*(f + Sqrt[f^2 - 4*e*g]))] + b*Log[1 + c + d*x]*Log[(d*(f + Sqrt[f^2 - 4*e*g] + 2*g*x))/(-2*(1 + c)*g + d*(f + Sqrt[f^2 - 4*e*g]))] + b*PolyLog[2, (2*g*(-1 + c + d*x))/(2*(-1 + c)*g + d*(-f + Sqrt[f^2 - 4*e*g]))] - b*PolyLog[2, (2*g*(-1 + c + d*x))/(2*(-1 + c)*g - d*(f + Sqrt[f^2 - 4*e*g]))] - b*PolyLog[2, (2*g*(1 + c + d*x))/(2*(1 + c)*g + d*(-f + Sqrt[f^2 - 4*e*g]))] + b*PolyLog[2, (2*g*(1 + c + d*x))/(2*(1 + c)*g - d*(f + Sqrt[f^2 - 4*e*g]))])/Sqrt[f^2 - 4*e*g]
```

Rubi [A] (verified)

Time = 1.12 (sec) , antiderivative size = 395, normalized size of antiderivative = 1.08, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {7279, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx + gx^2} dx \\
 & \quad \downarrow \text{7279} \\
 & \int \left(\frac{a}{e + fx + gx^2} + \frac{b \operatorname{arctanh}(c + dx)}{e + fx + gx^2} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{2a \operatorname{arctanh}\left(\frac{f+2gx}{\sqrt{f^2-4eg}}\right)}{\sqrt{f^2-4eg}} + \frac{b \operatorname{arctanh}(c + dx) \log\left(\frac{2(-2g(c+dx)+2cg-d(f-\sqrt{f^2-4eg}))}{(c+dx+1)(-2cg-d\sqrt{f^2-4eg}+df+2g)}\right)}{\sqrt{f^2-4eg}} \\
 & \quad \frac{b \operatorname{arctanh}(c + dx) \log\left(\frac{2(-2g(c+dx)+2cg-d(\sqrt{f^2-4eg}+f))}{(c+dx+1)(2(1-c)g+d(\sqrt{f^2-4eg}+f))}\right)}{\sqrt{f^2-4eg}} \\
 & \quad \frac{b \operatorname{PolyLog}\left(2, \frac{2(2cg-2(c+dx)g-d(f-\sqrt{f^2-4eg}))}{(fd-\sqrt{f^2-4eg}d-2cg+2g)(c+dx+1)} + 1\right)}{2\sqrt{f^2-4eg}} + \\
 & \quad \frac{b \operatorname{PolyLog}\left(2, \frac{2(2cg-2(c+dx)g-d(f+\sqrt{f^2-4eg}))}{(2(1-c)g+d(f+\sqrt{f^2-4eg}))(c+dx+1)} + 1\right)}{2\sqrt{f^2-4eg}}
 \end{aligned}$$

input

```
Int[(a + b*ArcTanh[c + d*x])/(e + f*x + g*x^2),x]
```

output

$$\begin{aligned} & (-2*a*ArcTanh[(f + 2*g*x)/Sqrt[f^2 - 4*e*g])/Sqrt[f^2 - 4*e*g] + (b*ArcTanh[c + d*x]*Log[(-2*(2*c*g - d*(f - Sqrt[f^2 - 4*e*g]) - 2*g*(c + d*x))]/((d*f + 2*g - 2*c*g - d*Sqrt[f^2 - 4*e*g])*(1 + c + d*x)))]/Sqrt[f^2 - 4*e*g] \\ & - (b*ArcTanh[c + d*x]*Log[(-2*(2*c*g - d*(f + Sqrt[f^2 - 4*e*g]) - 2*g*(c + d*x))]/((2*(1 - c)*g + d*(f + Sqrt[f^2 - 4*e*g])*(1 + c + d*x)))]/Sqrt[f^2 - 4*e*g] \\ & - (b*PolyLog[2, 1 + (2*(2*c*g - d*(f - Sqrt[f^2 - 4*e*g]) - 2*g*(c + d*x))]/((d*f + 2*g - 2*c*g - d*Sqrt[f^2 - 4*e*g])*(1 + c + d*x)))]/(2*Sqrt[f^2 - 4*e*g]) + (b*PolyLog[2, 1 + (2*(2*c*g - d*(f + Sqrt[f^2 - 4*e*g]) - 2*g*(c + d*x))]/((2*(1 - c)*g + d*(f + Sqrt[f^2 - 4*e*g])*(1 + c + d*x)))]/(2*Sqrt[f^2 - 4*e*g]) \end{aligned}$$

Defintions of rubi rules used

rule 2009

Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]

rule 7279

Int[(u_)/((a_.) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n + c*x^(2*n)), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b, c}, x] && EqQ[n2, 2*n] && IGtQ[n, 0]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 841 vs. 2(336) = 672.

Time = 1.05 (sec) , antiderivative size = 842, normalized size of antiderivative = 2.30

| method | result |
|------------------|---|
| risch | $-\frac{2da \arctan\left(\frac{2(-dx-c+1)g+2cg-df-2g}{\sqrt{4d^2eg-d^2f^2}}\right)}{\sqrt{4d^2eg-d^2f^2}} + \frac{db \ln(-dx-c+1) \ln\left(\frac{-2(-dx-c+1)g-2cg+df+\sqrt{-4d^2eg+d^2f^2+2g}}{-2cg+df+2g+\sqrt{-4d^2eg+d^2f^2}}\right)}{2\sqrt{-4d^2eg+d^2f^2}}$ |
| parts | Expression too large to display |
| derivativdivides | Expression too large to display |
| default | Expression too large to display |

input

int((a+b*arctanh(d*x+c))/(g*x^2+f*x+e), x, method=_RETURNVERBOSE)

output

```

-2*d*a/(4*d^2*e*g-d^2*f^2)^(1/2)*arctan((2*(-d*x-c+1)*g+2*c*g-d*f-2*g)/(4*
d^2*e*g-d^2*f^2)^(1/2))+1/2*d*b*ln(-d*x-c+1)/(-4*d^2*e*g+d^2*f^2)^(1/2)*ln
((-2*(-d*x-c+1)*g-2*c*g+d*f+(-4*d^2*e*g+d^2*f^2)^(1/2)+2*g)/(-2*c*g+d*f+2*
g+(-4*d^2*e*g+d^2*f^2)^(1/2)))-1/2*d*b*ln(-d*x-c+1)/(-4*d^2*e*g+d^2*f^2)^(
1/2)*ln((2*(-d*x-c+1)*g+2*c*g-d*f+(-4*d^2*e*g+d^2*f^2)^(1/2)-2*g)/(2*c*g-d
*f+(-4*d^2*e*g+d^2*f^2)^(1/2)-2*g))+1/2*d*b/(-4*d^2*e*g+d^2*f^2)^(1/2)*dil
og((-2*(-d*x-c+1)*g-2*c*g+d*f+(-4*d^2*e*g+d^2*f^2)^(1/2)+2*g)/(-2*c*g+d*f+
2*g+(-4*d^2*e*g+d^2*f^2)^(1/2)))-1/2*d*b/(-4*d^2*e*g+d^2*f^2)^(1/2)*dilog(
(2*(-d*x-c+1)*g+2*c*g-d*f+(-4*d^2*e*g+d^2*f^2)^(1/2)-2*g)/(2*c*g-d*f+(-4*d
^2*e*g+d^2*f^2)^(1/2)-2*g))+1/2*b*d*ln(d*x+c+1)/(-4*d^2*e*g+d^2*f^2)^(1/2)
*ln((-2*(d*x+c+1)*g+2*c*g-d*f+(-4*d^2*e*g+d^2*f^2)^(1/2)+2*g)/(2*c*g-d*f+2
*g+(-4*d^2*e*g+d^2*f^2)^(1/2)))-1/2*b*d*ln(d*x+c+1)/(-4*d^2*e*g+d^2*f^2)^(
1/2)*ln((2*(d*x+c+1)*g-2*c*g+d*f+(-4*d^2*e*g+d^2*f^2)^(1/2)-2*g)/(-2*c*g+d
*f+(-4*d^2*e*g+d^2*f^2)^(1/2)-2*g))+1/2*b*d/(-4*d^2*e*g+d^2*f^2)^(1/2)*dil
og((-2*(d*x+c+1)*g+2*c*g-d*f+(-4*d^2*e*g+d^2*f^2)^(1/2)+2*g)/(2*c*g-d*f+2*
g+(-4*d^2*e*g+d^2*f^2)^(1/2)))-1/2*b*d/(-4*d^2*e*g+d^2*f^2)^(1/2)*dilog((2
*(d*x+c+1)*g-2*c*g+d*f+(-4*d^2*e*g+d^2*f^2)^(1/2)-2*g)/(-2*c*g+d*f+(-4*d^2
*e*g+d^2*f^2)^(1/2)-2*g))

```

Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx + gx^2} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{gx^2 + fx + e} dx$$

input

```
integrate((a+b*arctanh(d*x+c))/(g*x^2+f*x+e),x, algorithm="fricas")
```

output

```
integral((b*arctanh(d*x + c) + a)/(g*x^2 + f*x + e), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx + gx^2} dx = \text{Timed out}$$

input `integrate((a+b*atanh(d*x+c))/(g*x**2+f*x+e),x)`

output `Timed out`

Maxima [F(-2)]

Exception generated.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx + gx^2} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arctanh(d*x+c))/(g*x^2+f*x+e),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*e*g-f^2>0)', see `assume?` for more deta`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx + gx^2} dx = \int \frac{b \operatorname{arctanh}(dx + c) + a}{gx^2 + fx + e} dx$$

input `integrate((a+b*arctanh(d*x+c))/(g*x^2+f*x+e),x, algorithm="giac")`

output `integrate((b*arctanh(d*x + c) + a)/(g*x^2 + f*x + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx + gx^2} dx = \int \frac{a + b \operatorname{atanh}(c + dx)}{gx^2 + fx + e} dx$$

input `int((a + b*atanh(c + d*x))/(e + f*x + g*x^2),x)`output `int((a + b*atanh(c + d*x))/(e + f*x + g*x^2), x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx + gx^2} dx = \text{Too large to display}$$

input `int((a+b*atanh(d*x+c))/(g*x^2+f*x+e),x)`

output

```
(2*sqrt(4*e*g - f**2)*atan((f + 2*g*x)/sqrt(4*e*g - f**2))*a*f - 4*atanh(c
+ d*x)**2*b*c*e*g + atanh(c + d*x)**2*b*c*f**2 + 4*int(atanh(c + d*x)/(c*
**2*e + c**2*f*x + c**2*g*x**2 + 2*c*d*e*x + 2*c*d*f*x**2 + 2*c*d*g*x**3 +
d**2*e*x**2 + d**2*f*x**3 + d**2*g*x**4 - e - f*x - g*x**2),x)*b*c**2*e*f*
g - int(atanh(c + d*x)/(c**2*e + c**2*f*x + c**2*g*x**2 + 2*c*d*e*x + 2*c*
d*f*x**2 + 2*c*d*g*x**3 + d**2*e*x**2 + d**2*f*x**3 + d**2*g*x**4 - e - f*
x - g*x**2),x)*b*c**2*f**3 - 8*int(atanh(c + d*x)/(c**2*e + c**2*f*x + c**
2*g*x**2 + 2*c*d*e*x + 2*c*d*f*x**2 + 2*c*d*g*x**3 + d**2*e*x**2 + d**2*f*
x**3 + d**2*g*x**4 - e - f*x - g*x**2),x)*b*c*d*e**2*g + 2*int(atanh(c + d
*x)/(c**2*e + c**2*f*x + c**2*g*x**2 + 2*c*d*e*x + 2*c*d*f*x**2 + 2*c*d*g*
x**3 + d**2*e*x**2 + d**2*f*x**3 + d**2*g*x**4 - e - f*x - g*x**2),x)*b*c*
d*e*f**2 - 4*int(atanh(c + d*x)/(c**2*e + c**2*f*x + c**2*g*x**2 + 2*c*d*e
*x + 2*c*d*f*x**2 + 2*c*d*g*x**3 + d**2*e*x**2 + d**2*f*x**3 + d**2*g*x**4
- e - f*x - g*x**2),x)*b*e*f*g + int(atanh(c + d*x)/(c**2*e + c**2*f*x +
c**2*g*x**2 + 2*c*d*e*x + 2*c*d*f*x**2 + 2*c*d*g*x**3 + d**2*e*x**2 + d**2
*f*x**3 + d**2*g*x**4 - e - f*x - g*x**2),x)*b*f**3 - 8*int((atanh(c + d*x
)*x**2)/(c**2*e + c**2*f*x + c**2*g*x**2 + 2*c*d*e*x + 2*c*d*f*x**2 + 2*c*
d*g*x**3 + d**2*e*x**2 + d**2*f*x**3 + d**2*g*x**4 - e - f*x - g*x**2),x)*
b*c*d*e*g**2 + 2*int((atanh(c + d*x)*x**2)/(c**2*e + c**2*f*x + c**2*g*x**
2 + 2*c*d*e*x + 2*c*d*f*x**2 + 2*c*d*g*x**3 + d**2*e*x**2 + d**2*f*x**3...
```

$$3.78 \quad \int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^2 + gx^4} dx$$

| | |
|---|-----|
| Optimal result | 682 |
| Mathematica [A] (warning: unable to verify) | 683 |
| Rubi [A] (warning: unable to verify) | 684 |
| Maple [C] (warning: unable to verify) | 687 |
| Fricas [F] | 688 |
| Sympy [F(-1)] | 688 |
| Maxima [F] | 688 |
| Giac [F] | 689 |
| Mupad [F(-1)] | 689 |
| Reduce [F] | 689 |

Optimal result

Integrand size = 25, antiderivative size = 1135

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^2 + gx^4} dx = \text{Too large to display}$$

output

```

-1/2*g^(1/2)*(a+b*arctanh(d*x+c))*ln(-2*d*((-f-(-4*e*g+f^2)^(1/2))^(1/2)-g
^(1/2)*x^2^(1/2))/(2^(1/2)*(1-c)*g^(1/2)-d*(-f-(-4*e*g+f^2)^(1/2))^(1/2))/
(d*x+c+1))*2^(1/2)/(-4*e*g+f^2)^(1/2)/(-f-(-4*e*g+f^2)^(1/2))^(1/2)+1/2*g^
(1/2)*(a+b*arctanh(d*x+c))*ln(-2*d*((-f+(-4*e*g+f^2)^(1/2))^(1/2)-g^(1/2)*
x^2^(1/2))/(2^(1/2)*(1-c)*g^(1/2)-d*(-f+(-4*e*g+f^2)^(1/2))^(1/2))/(d*x+c+
1))*2^(1/2)/(-4*e*g+f^2)^(1/2)/(-f+(-4*e*g+f^2)^(1/2))^(1/2)+1/2*g^(1/2)*(
a+b*arctanh(d*x+c))*ln(2*d*((-f-(-4*e*g+f^2)^(1/2))^(1/2)+g^(1/2)*x^2^(1/2)
))/(2^(1/2)*(1-c)*g^(1/2)+d*(-f-(-4*e*g+f^2)^(1/2))^(1/2))/(d*x+c+1))*2^(1
/2)/(-4*e*g+f^2)^(1/2)/(-f-(-4*e*g+f^2)^(1/2))^(1/2)-1/2*g^(1/2)*(a+b*arct
anh(d*x+c))*ln(2*d*((-f+(-4*e*g+f^2)^(1/2))^(1/2)+g^(1/2)*x^2^(1/2))/(2^(1
/2)*(1-c)*g^(1/2)+d*(-f+(-4*e*g+f^2)^(1/2))^(1/2))/(d*x+c+1))*2^(1/2)/(-4*
e*g+f^2)^(1/2)/(-f+(-4*e*g+f^2)^(1/2))^(1/2)+1/4*b*g^(1/2)*polylog(2,1+2*d
*((-f-(-4*e*g+f^2)^(1/2))^(1/2)-g^(1/2)*x^2^(1/2))/(2^(1/2)*(1-c)*g^(1/2)-
d*(-f-(-4*e*g+f^2)^(1/2))^(1/2))/(d*x+c+1))*2^(1/2)/(-4*e*g+f^2)^(1/2)/(-f
-(-4*e*g+f^2)^(1/2))^(1/2)-1/4*b*g^(1/2)*polylog(2,1+2*d*((-f+(-4*e*g+f^2)
^(1/2))^(1/2)-g^(1/2)*x^2^(1/2))/(2^(1/2)*(1-c)*g^(1/2)-d*(-f+(-4*e*g+f^2)
^(1/2))^(1/2))/(d*x+c+1))*2^(1/2)/(-4*e*g+f^2)^(1/2)/(-f+(-4*e*g+f^2)^(1/2)
))^(1/2)-1/4*b*g^(1/2)*polylog(2,1-2*d*((-f-(-4*e*g+f^2)^(1/2))^(1/2)+g^(1
/2)*x^2^(1/2))/(2^(1/2)*(1-c)*g^(1/2)+d*(-f-(-4*e*g+f^2)^(1/2))^(1/2))/(d*
x+c+1))*2^(1/2)/(-4*e*g+f^2)^(1/2)/(-f-(-4*e*g+f^2)^(1/2))^(1/2)+1/4*b*...

```

Mathematica [A] (warning: unable to verify)

Time = 3.21 (sec) , antiderivative size = 2111, normalized size of antiderivative = 1.86

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^2 + gx^4} dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*ArcTanh[c + d*x])/(e + f*x^2 + g*x^4),x]
```

output

```
(Sqrt[g]*(4*a*Sqrt[-f + Sqrt[f^2 - 4*e*g]]*Sqrt[-(f + Sqrt[f^2 - 4*e*g])^2]
]*ArcTan[(Sqrt[2]*Sqrt[g]*x)/Sqrt[f - Sqrt[f^2 - 4*e*g]]] - 4*a*Sqrt[-f -
Sqrt[f^2 - 4*e*g]]*Sqrt[-(f - Sqrt[f^2 - 4*e*g])^2]*ArcTan[(Sqrt[2]*Sqrt[g]
*x)/Sqrt[f + Sqrt[f^2 - 4*e*g]]] + b*Sqrt[-(f - Sqrt[f^2 - 4*e*g])^2]*Sqr
t[f + Sqrt[f^2 - 4*e*g]]*Log[1 - c - d*x]*Log[(d*(Sqrt[-f - Sqrt[f^2 - 4*e
*g]] - Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(-1 + c)*Sqrt[g] + d*Sqrt[-f - Sqrt[f^
2 - 4*e*g]])] - b*Sqrt[-(f - Sqrt[f^2 - 4*e*g])^2]*Sqrt[f + Sqrt[f^2 - 4*e
*g]]*Log[1 + c + d*x]*Log[(d*(Sqrt[-f - Sqrt[f^2 - 4*e*g]] - Sqrt[2]*Sqrt[
g]*x))/(Sqrt[2]*(1 + c)*Sqrt[g] + d*Sqrt[-f - Sqrt[f^2 - 4*e*g]])] - b*Sqr
t[f - Sqrt[f^2 - 4*e*g]]*Sqrt[-(f + Sqrt[f^2 - 4*e*g])^2]*Log[1 - c - d*x]
*Log[(d*(Sqrt[-f + Sqrt[f^2 - 4*e*g]] - Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(-1 +
c)*Sqrt[g] + d*Sqrt[-f + Sqrt[f^2 - 4*e*g]])] + b*Sqrt[f - Sqrt[f^2 - 4*e
*g]]*Sqrt[-(f + Sqrt[f^2 - 4*e*g])^2]*Log[1 + c + d*x]*Log[(d*(Sqrt[-f + S
qrt[f^2 - 4*e*g]] - Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(1 + c)*Sqrt[g] + d*Sqrt[
-f + Sqrt[f^2 - 4*e*g]])] - b*Sqrt[-(f - Sqrt[f^2 - 4*e*g])^2]*Sqrt[f + Sq
rt[f^2 - 4*e*g]]*Log[1 - c - d*x]*Log[(d*(Sqrt[-f - Sqrt[f^2 - 4*e*g]] + S
qrt[2]*Sqrt[g]*x))/(-(Sqrt[2]*(-1 + c)*Sqrt[g]) + d*Sqrt[-f - Sqrt[f^2 - 4
*e*g]])] + b*Sqrt[-(f - Sqrt[f^2 - 4*e*g])^2]*Sqrt[f + Sqrt[f^2 - 4*e*g]]*
Log[1 + c + d*x]*Log[(d*(Sqrt[-f - Sqrt[f^2 - 4*e*g]] + Sqrt[2]*Sqrt[g]*x)
)/(-(Sqrt[2]*(1 + c)*Sqrt[g]) + d*Sqrt[-f - Sqrt[f^2 - 4*e*g]])] + b*Sq...
```

Rubi [A] (warning: unable to verify)

Time = 4.67 (sec) , antiderivative size = 2120, normalized size of antiderivative = 1.87, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {7279, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \text{barctanh}(c + dx)}{e + fx^2 + gx^4} dx$$

$$\downarrow 7279$$

$$\int \left(\frac{a}{e + fx^2 + gx^4} + \frac{\text{barctanh}(c + dx)}{e + fx^2 + gx^4} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
 & \frac{\sqrt{2a}\sqrt{g} \arctan\left(\frac{\sqrt{2}\sqrt{gx}}{\sqrt{f-\sqrt{f^2-4eg}}}\right)}{\sqrt{f^2-4eg}\sqrt{f-\sqrt{f^2-4eg}}} - \frac{\sqrt{2a}\sqrt{g} \arctan\left(\frac{\sqrt{2}\sqrt{gx}}{\sqrt{f+\sqrt{f^2-4eg}}}\right)}{\sqrt{f^2-4eg}\sqrt{f+\sqrt{f^2-4eg}}} + \\
 & \frac{b\sqrt{g} \log(-c-dx+1) \log\left(-\frac{d(\sqrt{-f-\sqrt{f^2-4eg}}-\sqrt{2}\sqrt{gx})}{\sqrt{2}(1-c)\sqrt{g}-d\sqrt{-f-\sqrt{f^2-4eg}}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{-f-\sqrt{f^2-4eg}}} - \\
 & \frac{b\sqrt{g} \log(c+dx+1) \log\left(\frac{d(\sqrt{-f-\sqrt{f^2-4eg}}-\sqrt{2}\sqrt{gx})}{\sqrt{2}\sqrt{g}(c+1)+d\sqrt{-f-\sqrt{f^2-4eg}}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{-f-\sqrt{f^2-4eg}}} - \\
 & \frac{b\sqrt{g} \log(-c-dx+1) \log\left(-\frac{d(\sqrt{\sqrt{f^2-4eg}-f}-\sqrt{2}\sqrt{gx})}{\sqrt{2}(1-c)\sqrt{g}-d\sqrt{\sqrt{f^2-4eg}-f}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{\sqrt{f^2-4eg}-f}} + \\
 & \frac{b\sqrt{g} \log(c+dx+1) \log\left(\frac{d(\sqrt{\sqrt{f^2-4eg}-f}-\sqrt{2}\sqrt{gx})}{\sqrt{2}\sqrt{g}(c+1)+d\sqrt{\sqrt{f^2-4eg}-f}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{\sqrt{f^2-4eg}-f}} + \\
 & \frac{b\sqrt{g} \log(c+dx+1) \log\left(-\frac{d(\sqrt{2}\sqrt{gx}+\sqrt{-f-\sqrt{f^2-4eg}})}{\sqrt{2}(c+1)\sqrt{g}-d\sqrt{-f-\sqrt{f^2-4eg}}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{-f-\sqrt{f^2-4eg}}} - \\
 & \frac{b\sqrt{g} \log(-c-dx+1) \log\left(\frac{d(\sqrt{2}\sqrt{gx}+\sqrt{-f-\sqrt{f^2-4eg}})}{\sqrt{2}\sqrt{g}(1-c)+d\sqrt{-f-\sqrt{f^2-4eg}}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{-f-\sqrt{f^2-4eg}}} - \\
 & \frac{b\sqrt{g} \log(c+dx+1) \log\left(-\frac{d(\sqrt{2}\sqrt{gx}+\sqrt{\sqrt{f^2-4eg}-f})}{\sqrt{2}(c+1)\sqrt{g}-d\sqrt{\sqrt{f^2-4eg}-f}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{\sqrt{f^2-4eg}-f}} + \\
 & \frac{b\sqrt{g} \log(-c-dx+1) \log\left(\frac{d(\sqrt{2}\sqrt{gx}+\sqrt{\sqrt{f^2-4eg}-f})}{\sqrt{2}\sqrt{g}(1-c)+d\sqrt{\sqrt{f^2-4eg}-f}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{\sqrt{f^2-4eg}-f}} + \\
 & \frac{b\sqrt{g} \operatorname{PolyLog}\left(2, \frac{\sqrt{2}\sqrt{g}(-c-dx+1)}{\sqrt{2}(1-c)\sqrt{g}-d\sqrt{-f-\sqrt{f^2-4eg}}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{-f-\sqrt{f^2-4eg}}} - \\
 & \frac{b\sqrt{g} \operatorname{PolyLog}\left(2, \frac{\sqrt{2}\sqrt{g}(-c-dx+1)}{\sqrt{2}\sqrt{g}(1-c)+d\sqrt{-f-\sqrt{f^2-4eg}}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{-f-\sqrt{f^2-4eg}}} - \frac{b\sqrt{g} \operatorname{PolyLog}\left(2, \frac{\sqrt{2}\sqrt{g}(-c-dx+1)}{\sqrt{2}(1-c)\sqrt{g}-d\sqrt{\sqrt{f^2-4eg}-f}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{\sqrt{f^2-4eg}-f}} + \\
 & \frac{b\sqrt{g} \operatorname{PolyLog}\left(2, \frac{\sqrt{2}\sqrt{g}(-c-dx+1)}{\sqrt{2}\sqrt{g}(1-c)+d\sqrt{\sqrt{f^2-4eg}-f}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{\sqrt{f^2-4eg}-f}} +
 \end{aligned}$$

input `Int[(a + b*ArcTanh[c + d*x])/(e + f*x^2 + g*x^4),x]`

output `(Sqrt[2]*a*Sqrt[g]*ArcTan[(Sqrt[2]*Sqrt[g]*x)/Sqrt[f - Sqrt[f^2 - 4*e*g]])/(Sqrt[f^2 - 4*e*g]*Sqrt[f - Sqrt[f^2 - 4*e*g]]) - (Sqrt[2]*a*Sqrt[g]*ArcTan[(Sqrt[2]*Sqrt[g]*x)/Sqrt[f + Sqrt[f^2 - 4*e*g]])/(Sqrt[f^2 - 4*e*g]*Sqrt[f + Sqrt[f^2 - 4*e*g]]) + (b*Sqrt[g]*Log[1 - c - d*x]*Log[-((d*(Sqrt[-f - Sqrt[f^2 - 4*e*g]] - Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(1 - c)*Sqrt[g] - d*Sqrt[-f - Sqrt[f^2 - 4*e*g]]))])/(2*Sqrt[2]*Sqrt[f^2 - 4*e*g]*Sqrt[-f - Sqrt[f^2 - 4*e*g]]) - (b*Sqrt[g]*Log[1 + c + d*x]*Log[(d*(Sqrt[-f - Sqrt[f^2 - 4*e*g]] - Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(1 + c)*Sqrt[g] + d*Sqrt[-f - Sqrt[f^2 - 4*e*g]]))])/(2*Sqrt[2]*Sqrt[f^2 - 4*e*g]*Sqrt[-f - Sqrt[f^2 - 4*e*g]]) - (b*Sqrt[g]*Log[1 - c - d*x]*Log[-((d*(Sqrt[-f + Sqrt[f^2 - 4*e*g]] - Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(1 - c)*Sqrt[g] - d*Sqrt[-f + Sqrt[f^2 - 4*e*g]]))])/(2*Sqrt[2]*Sqrt[f^2 - 4*e*g]*Sqrt[-f + Sqrt[f^2 - 4*e*g]]) + (b*Sqrt[g]*Log[1 + c + d*x]*Log[(d*(Sqrt[-f + Sqrt[f^2 - 4*e*g]] - Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(1 + c)*Sqrt[g] + d*Sqrt[-f + Sqrt[f^2 - 4*e*g]]))])/(2*Sqrt[2]*Sqrt[f^2 - 4*e*g]*Sqrt[-f + Sqrt[f^2 - 4*e*g]]) + (b*Sqrt[g]*Log[1 + c + d*x]*Log[-((d*(Sqrt[-f - Sqrt[f^2 - 4*e*g]] + Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(1 + c)*Sqrt[g] - d*Sqrt[-f - Sqrt[f^2 - 4*e*g]]))])/(2*Sqrt[2]*Sqrt[f^2 - 4*e*g]*Sqrt[-f - Sqrt[f^2 - 4*e*g]]) - (b*Sqrt[g]*Log[1 - c - d*x]*Log[(d*(Sqrt[-f - Sqrt[f^2 - 4*e*g]] + Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(1 - c)*Sqrt[g] + d*Sqrt[-f - Sqrt[f^2 - 4*e*g]]))])/(2*Sqrt[2]*Sqrt[f^2 - 4...`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7279 `Int[(u_)/((a_.) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n + c*x^(2*n)), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b, c}, x] && EqQ[n2, 2*n] && IGtQ[n, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.54 (sec) , antiderivative size = 718, normalized size of antiderivative = 0.63

| method | result |
|-------------------|--|
| risch | $\frac{d^3 a}{\sqrt{-R=\text{RootOf}(g_Z^4+(4cg-4g)_Z^3+(6c^2g+d^2f-12cg+6g)_Z^2+(4c^3g+2cd^2f-12c^2g-2d^2f+12cg-4g)_Z+c^4g+c^2}}$ |
| parts | Expression too large to display |
| derivativedivides | Expression too large to display |
| default | Expression too large to display |

input

```
int((a+b*arctanh(d*x+c))/(g*x^4+f*x^2+e),x,method=_RETURNVERBOSE)
```

output

```
-1/2*d^3*a*sum(1/(2*_R^3*g+6*_R^2*c*g+6*_R*c^2*g+_R*d^2*f+2*c^3*g+c*d^2*f-6*_R^2*g-12*_R*c*g-6*c^2*g-d^2*f+6*_R*g+6*c*g-2*g)*ln(-d*x-_R-c+1),_R=RootOf(g*_Z^4+(4*c*g-4*g)*_Z^3+(6*c^2*g+d^2*f-12*c*g+6*g)*_Z^2+(4*c^3*g+2*c*d^2*f-12*c^2*g-2*d^2*f+12*c*g-4*g)*_Z+c^4*g+c^2*d^2*f+e*d^4-4*c^3*g-2*c*d^2*f+6*c^2*g+d^2*f-4*c*g+g))+1/4*d^3*b*sum(1/(2*_R1^3*g+6*_R1^2*c*g+6*_R1*c^2*g+_R1*d^2*f+2*c^3*g+c*d^2*f-6*_R1^2*g-12*_R1*c*g-6*c^2*g-d^2*f+6*_R1*g+6*c*g-2*g)*(ln(-d*x-c+1)*ln((d*x+_R1+c-1)/_R1)+dilog((d*x+_R1+c-1)/_R1)),_R1=RootOf(g*_Z^4+(4*c*g-4*g)*_Z^3+(6*c^2*g+d^2*f-12*c*g+6*g)*_Z^2+(4*c^3*g+2*c*d^2*f-12*c^2*g-2*d^2*f+12*c*g-4*g)*_Z+c^4*g+c^2*d^2*f+e*d^4-4*c^3*g-2*c*d^2*f+6*c^2*g+d^2*f-4*c*g+g))+1/4*b*d^3*sum(1/(2*_R1^3*g-6*_R1^2*c*g+6*_R1*c^2*g+_R1*d^2*f-2*c^3*g-c*d^2*f-6*_R1^2*g+12*_R1*c*g-6*c^2*g-d^2*f+6*_R1*g-6*c*g-2*g)*(ln(d*x+c+1)*ln((-d*x+_R1-c-1)/_R1)+dilog((-d*x+_R1-c-1)/_R1)),_R1=RootOf(g*_Z^4+(-4*c*g-4*g)*_Z^3+(6*c^2*g+d^2*f+12*c*g+6*g)*_Z^2+(-4*c^3*g-2*c*d^2*f-12*c^2*g-2*d^2*f-12*c*g-4*g)*_Z+c^4*g+c^2*d^2*f+e*d^4+4*c^3*g+2*c*d^2*f+6*c^2*g+d^2*f+4*c*g+g))
```


Fricas [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^2 + gx^4} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{gx^4 + fx^2 + e} dx$$

input `integrate((a+b*arctanh(d*x+c))/(g*x^4+f*x^2+e),x, algorithm="fricas")`

output `integral((b*arctanh(d*x + c) + a)/(g*x^4 + f*x^2 + e), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^2 + gx^4} dx = \text{Timed out}$$

input `integrate((a+b*atanh(d*x+c))/(g*x**4+f*x**2+e),x)`

output `Timed out`

Maxima [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^2 + gx^4} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{gx^4 + fx^2 + e} dx$$

input `integrate((a+b*arctanh(d*x+c))/(g*x^4+f*x^2+e),x, algorithm="maxima")`

output `integrate((b*arctanh(d*x + c) + a)/(g*x^4 + f*x^2 + e), x)`

Giac [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^2 + gx^4} dx = \int \frac{b \operatorname{artanh}(dx + c) + a}{gx^4 + fx^2 + e} dx$$

input `integrate((a+b*arctanh(d*x+c))/(g*x^4+f*x^2+e),x, algorithm="giac")`

output `sage0*x`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^2 + gx^4} dx = \int \frac{a + b \operatorname{atanh}(c + dx)}{gx^4 + fx^2 + e} dx$$

input `int((a + b*atanh(c + d*x))/(e + f*x^2 + g*x^4),x)`

output `int((a + b*atanh(c + d*x))/(e + f*x^2 + g*x^4), x)`

Reduce [F]

$$\int \frac{a + b \operatorname{arctanh}(c + dx)}{e + fx^2 + gx^4} dx$$

$$= \frac{2\sqrt{e} \sqrt{2\sqrt{g}\sqrt{e+f}} \operatorname{atan}\left(\frac{\sqrt{2\sqrt{g}\sqrt{e-f}-2\sqrt{g}x}}{\sqrt{2\sqrt{g}\sqrt{e+f}}}\right) af - 4\sqrt{g} \sqrt{2\sqrt{g}\sqrt{e+f}} \operatorname{atan}\left(\frac{\sqrt{2\sqrt{g}\sqrt{e-f}-2\sqrt{g}x}}{\sqrt{2\sqrt{g}\sqrt{e+f}}}\right) ae - 2\sqrt{e}}{\dots}$$

input `int((a+b*atanh(d*x+c))/(g*x^4+f*x^2+e),x)`

output

```
(2*sqrt(e)*sqrt(2*sqrt(g)*sqrt(e) + f)*atan((sqrt(2*sqrt(g)*sqrt(e) - f) -
2*sqrt(g)*x)/sqrt(2*sqrt(g)*sqrt(e) + f))*a*f - 4*sqrt(g)*sqrt(2*sqrt(g)*
sqrt(e) + f)*atan((sqrt(2*sqrt(g)*sqrt(e) - f) - 2*sqrt(g)*x)/sqrt(2*sqrt(
g)*sqrt(e) + f))*a*e - 2*sqrt(e)*sqrt(2*sqrt(g)*sqrt(e) + f)*atan((sqrt(2*
sqrt(g)*sqrt(e) - f) + 2*sqrt(g)*x)/sqrt(2*sqrt(g)*sqrt(e) + f))*a*f + 4*s
qrt(g)*sqrt(2*sqrt(g)*sqrt(e) + f)*atan((sqrt(2*sqrt(g)*sqrt(e) - f) + 2*s
qrt(g)*x)/sqrt(2*sqrt(g)*sqrt(e) + f))*a*e - sqrt(e)*sqrt(2*sqrt(g)*sqrt(e
) - f)*log( - sqrt(2*sqrt(g)*sqrt(e) - f)*x + sqrt(e) + sqrt(g)*x**2)*a*f
+ sqrt(e)*sqrt(2*sqrt(g)*sqrt(e) - f)*log(sqrt(2*sqrt(g)*sqrt(e) - f)*x +
sqrt(e) + sqrt(g)*x**2)*a*f - 2*sqrt(g)*sqrt(2*sqrt(g)*sqrt(e) - f)*log( -
sqrt(2*sqrt(g)*sqrt(e) - f)*x + sqrt(e) + sqrt(g)*x**2)*a*e + 2*sqrt(g)*s
qrt(2*sqrt(g)*sqrt(e) - f)*log(sqrt(2*sqrt(g)*sqrt(e) - f)*x + sqrt(e) + s
qrt(g)*x**2)*a*e + 16*int(atanh(c + d*x)/(e + f*x**2 + g*x**4),x)*b*e**2*g
- 4*int(atanh(c + d*x)/(e + f*x**2 + g*x**4),x)*b*e*f**2)/(4*e*(4*e*g - f
**2))
```

3.79 $\int \frac{(ce+dex)(a+b\mathbf{arctanh}(c+dx))}{1-(c+dx)^2} dx$

| | |
|----------------------------|-----|
| Optimal result | 691 |
| Mathematica [A] (verified) | 692 |
| Rubi [A] (verified) | 692 |
| Maple [A] (verified) | 695 |
| Fricas [F] | 695 |
| Sympy [F] | 696 |
| Maxima [F] | 696 |
| Giac [F] | 697 |
| Mupad [F(-1)] | 697 |
| Reduce [F] | 697 |

Optimal result

Integrand size = 32, antiderivative size = 83

$$\int \frac{(ce + dex)(a + \mathbf{barctanh}(c + dx))}{1 - (c + dx)^2} dx = -\frac{e(a + \mathbf{barctanh}(c + dx))^2}{2bd} + \frac{e(a + \mathbf{barctanh}(c + dx)) \log\left(\frac{2}{1-c-dx}\right)}{d} + \frac{be \operatorname{PolyLog}\left(2, -\frac{1+c+dx}{1-c-dx}\right)}{2d}$$

output

```
-1/2*e*(a+b*arctanh(d*x+c))^2/b/d+e*(a+b*arctanh(d*x+c))*ln(2/(-d*x-c+1))/d+1/2*b*e*polylog(2,-(d*x+c+1)/(-d*x-c+1))/d
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.81

$$\int \frac{(ce + dex)(a + b \operatorname{arctanh}(c + dx))}{1 - (c + dx)^2} dx = e \left(-\frac{a \log(1 - c - dx)}{2d} + \frac{b \log^2(1 - c - dx)}{8d} - \frac{b \log(2) \log(-1 + c + dx)}{4d} - \frac{a \log(1 + c + dx)}{2d} + \frac{b \log(2) \log(1 + c + dx)}{4d} - \frac{b \log^2(1 + c + dx)}{8d} + \frac{b \operatorname{PolyLog}\left(2, \frac{1}{2}(1 - c - dx)\right)}{4d} - \frac{b \operatorname{PolyLog}\left(2, \frac{1}{2}(1 + c + dx)\right)}{4d} \right)$$

input `Integrate[((c*e + d*e*x)*(a + b*ArcTanh[c + d*x]))/(1 - (c + d*x)^2), x]`

output `e*(-1/2*(a*Log[1 - c - d*x])/d + (b*Log[1 - c - d*x]^2)/(8*d) - (b*Log[2]*Log[-1 + c + d*x])/(4*d) - (a*Log[1 + c + d*x])/(2*d) + (b*Log[2]*Log[1 + c + d*x])/(4*d) - (b*Log[1 + c + d*x]^2)/(8*d) + (b*PolyLog[2, (1 - c - d*x)/2])/(4*d) - (b*PolyLog[2, (1 + c + d*x)/2])/(4*d))`

Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.87, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {7281, 27, 6546, 6470, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(ce + dex)(a + b \operatorname{arctanh}(c + dx))}{1 - (c + dx)^2} dx$$

$$\begin{aligned}
& \int \frac{e^{(c+dx)(a+b\operatorname{arctanh}(c+dx))}}{1-(c+dx)^2} d(c+dx) \\
& \quad \downarrow \text{7281} \\
& \frac{d}{d} \int \frac{e^{(c+dx)(a+b\operatorname{arctanh}(c+dx))}}{1-(c+dx)^2} d(c+dx) \\
& \quad \downarrow \text{27} \\
& \frac{d}{d} \int \frac{e^{\left(\int \frac{a+b\operatorname{arctanh}(c+dx)}{-c-dx+1} d(c+dx) - \frac{(a+b\operatorname{arctanh}(c+dx))^2}{2b}\right)}}{1-(c+dx)^2} d(c+dx) \\
& \quad \downarrow \text{6546} \\
& \frac{d}{d} \left(\int \frac{e^{\left(\int \frac{a+b\operatorname{arctanh}(c+dx)}{-c-dx+1} d(c+dx) - \frac{(a+b\operatorname{arctanh}(c+dx))^2}{2b}\right)}}{1-(c+dx)^2} d(c+dx) - \frac{(a+b\operatorname{arctanh}(c+dx))^2}{2b} \right) \\
& \quad \downarrow \text{6470} \\
& \frac{d}{d} \left(-b \int \frac{\log\left(\frac{2}{-c-dx+1}\right)}{1-(c+dx)^2} d(c+dx) - \frac{(a+b\operatorname{arctanh}(c+dx))^2}{2b} + \log\left(\frac{2}{-c-dx+1}\right) (a+b\operatorname{arctanh}(c+dx)) \right) \\
& \quad \downarrow \text{2849} \\
& \frac{d}{d} \left(b \int \frac{\log\left(\frac{2}{-c-dx+1}\right)}{1-\frac{2}{-c-dx+1}} d\frac{1}{-c-dx+1} - \frac{(a+b\operatorname{arctanh}(c+dx))^2}{2b} + \log\left(\frac{2}{-c-dx+1}\right) (a+b\operatorname{arctanh}(c+dx)) \right) \\
& \quad \downarrow \text{2752} \\
& \frac{d}{d} \left(-\frac{(a+b\operatorname{arctanh}(c+dx))^2}{2b} + \log\left(\frac{2}{-c-dx+1}\right) (a+b\operatorname{arctanh}(c+dx)) + \frac{1}{2}b \operatorname{PolyLog}\left(2, 1 - \frac{2}{-c-dx+1}\right) \right)
\end{aligned}$$

input

```
Int[((c*e + d*e*x)*(a + b*ArcTanh[c + d*x]))/(1 - (c + d*x)^2),x]
```

output

```
(e*(-1/2*(a + b*ArcTanh[c + d*x])^2/b + (a + b*ArcTanh[c + d*x])*Log[2/(1 - c - d*x)] + (b*PolyLog[2, 1 - 2/(1 - c - d*x)]/2))/d
```

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 2752 $\text{Int}[\text{Log}[(c_*)(x_)]/((d_) + (e_*)(x_)), x_Symbol] \rightarrow \text{Simp}[(-e^{(-1)})*\text{PolyLog}[2, 1 - c*x], x] /; \text{FreeQ}\{c, d, e\}, x] \ \&\& \ \text{EqQ}[e + c*d, 0]$
- rule 2849 $\text{Int}[\text{Log}[(c_)/((d_) + (e_*)(x_))]/((f_) + (g_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[-e/g \text{ Subst}[\text{Int}[\text{Log}[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; \text{FreeQ}\{c, d, e, f, g\}, x] \ \&\& \ \text{EqQ}[c, 2*d] \ \&\& \ \text{EqQ}[e^2*f + d^2*g, 0]$
- rule 6470 $\text{Int}[((a_.) + \text{ArcTanh}[(c_*)(x_)]*(b_.))^p/((d_) + (e_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[(-(a + b*\text{ArcTanh}[c*x])^p)*(\text{Log}[2/(1 + e*(x/d))]/e), x] + \text{Simp}[b*c*(p/e) \text{ Int}[(a + b*\text{ArcTanh}[c*x])^{p-1}*(\text{Log}[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d^2 - e^2, 0]$
- rule 6546 $\text{Int}[(((a_.) + \text{ArcTanh}[(c_*)(x_)]*(b_.))^p*(x_))/((d_) + (e_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTanh}[c*x])^{p+1}/(b*e*(p+1)), x] + \text{Simp}[1/(c*d) \text{ Int}[(a + b*\text{ArcTanh}[c*x])^p/(1 - c*x), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{IGtQ}[p, 0]$
- rule 7281 $\text{Int}[u_, x_Symbol] \rightarrow \text{With}\{\{lst = \text{FunctionOfLinear}[u, x]\}, \text{Simp}[1/lst[[3]] \text{ Subst}[\text{Int}[lst[[1]], x], x, lst[[2]] + lst[[3]]*x], x] /; \ !\text{FalseQ}[lst]]$

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.80

| method | result |
|-------------------|--|
| derivativedivides | $-ae\left(\frac{\ln(dx+c-1)}{2} + \frac{\ln(dx+c+1)}{2}\right) - eb\left(\frac{\operatorname{arctanh}(dx+c)\ln(dx+c-1)}{2} + \frac{\operatorname{arctanh}(dx+c)\ln(dx+c+1)}{2} - \frac{\operatorname{dilog}\left(\frac{dx}{2} + \frac{c}{2} + \frac{1}{2}\right)}{2} - \frac{\ln(dx+c)}{d}\right)$ |
| default | $-ae\left(\frac{\ln(dx+c-1)}{2} + \frac{\ln(dx+c+1)}{2}\right) - eb\left(\frac{\operatorname{arctanh}(dx+c)\ln(dx+c-1)}{2} + \frac{\operatorname{arctanh}(dx+c)\ln(dx+c+1)}{2} - \frac{\operatorname{dilog}\left(\frac{dx}{2} + \frac{c}{2} + \frac{1}{2}\right)}{2} - \frac{\ln(dx+c)}{d}\right)$ |
| risch | $\frac{eb\ln(-dx-c+1)^2}{8d} + \frac{eb\ln\left(\frac{dx}{2} + \frac{c}{2} + \frac{1}{2}\right)\ln(-dx-c+1)}{4d} - \frac{eb\operatorname{dilog}\left(-\frac{dx}{2} - \frac{c}{2} + \frac{1}{2}\right)}{4d} - \frac{ea\ln((-dx-c+1)(-dx-c-1))}{2d}$ |
| parts | $-\frac{ae\ln(d^2x^2+2cdx+c^2-1)}{2d} - \frac{eb\left(\frac{\operatorname{arctanh}(dx+c)\ln(dx+c-1)}{2} + \frac{\operatorname{arctanh}(dx+c)\ln(dx+c+1)}{2} - \frac{\operatorname{dilog}\left(\frac{dx}{2} + \frac{c}{2} + \frac{1}{2}\right)}{2} - \frac{\ln(dx+c)}{d}\right)}{d}$ |

input `int((d*e*x+c*e)*(a+b*arctanh(d*x+c))/(1-(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/d*(-a*e*(1/2*ln(d*x+c-1)+1/2*ln(d*x+c+1))-e*b*(1/2*arctanh(d*x+c)*ln(d*x+c-1)+1/2*arctanh(d*x+c)*ln(d*x+c+1)-1/2*dilog(1/2*d*x+1/2*c+1/2)-1/4*ln(d*x+c-1)*ln(1/2*d*x+1/2*c+1/2)+1/8*ln(d*x+c-1)^2+1/4*(ln(d*x+c+1)-ln(1/2*d*x+1/2*c+1/2))*ln(-1/2*d*x-1/2*c+1/2)-1/8*ln(d*x+c+1)^2))`

Fricas [F]

$$\int \frac{(ce + dex)(a + b\operatorname{arctanh}(c + dx))}{1 - (c + dx)^2} dx = \int -\frac{(dex + ce)(b\operatorname{arctanh}(dx + c) + a)}{(dx + c)^2 - 1} dx$$

input `integrate((d*e*x+c*e)*(a+b*arctanh(d*x+c))/(1-(d*x+c)^2),x, algorithm="fricas")`

output `integral(-(a*d*e*x + a*c*e + (b*d*e*x + b*c*e)*arctanh(d*x + c))/(d^2*x^2 + 2*c*d*x + c^2 - 1), x)`

Sympy [F]

$$\int \frac{(ce + dex)(a + b \operatorname{arctanh}(c + dx))}{1 - (c + dx)^2} dx = -e \left(\int \frac{ac}{c^2 + 2cdx + d^2x^2 - 1} dx + \int \frac{adx}{c^2 + 2cdx + d^2x^2 - 1} dx + \int \frac{bc \operatorname{atanh}(c + dx)}{c^2 + 2cdx + d^2x^2 - 1} dx + \int \frac{bdx \operatorname{atanh}(c + dx)}{c^2 + 2cdx + d^2x^2 - 1} dx \right)$$

input `integrate((d*e*x+c*e)*(a+b*atanh(d*x+c))/(1-(d*x+c)**2),x)`

output `-e*(Integral(a*c/(c**2 + 2*c*d*x + d**2*x**2 - 1), x) + Integral(a*d*x/(c**2 + 2*c*d*x + d**2*x**2 - 1), x) + Integral(b*c*atanh(c + d*x)/(c**2 + 2*c*d*x + d**2*x**2 - 1), x) + Integral(b*d*x*atanh(c + d*x)/(c**2 + 2*c*d*x + d**2*x**2 - 1), x))`

Maxima [F]

$$\int \frac{(ce + dex)(a + b \operatorname{arctanh}(c + dx))}{1 - (c + dx)^2} dx = \int -\frac{(dex + ce)(b \operatorname{atanh}(dx + c) + a)}{(dx + c)^2 - 1} dx$$

input `integrate((d*e*x+c*e)*(a+b*arctanh(d*x+c))/(1-(d*x+c)^2),x, algorithm="maxima")`

output `1/2*b*c*e*(log(d*x + c + 1)/d - log(d*x + c - 1)/d)*arctanh(d*x + c) - 1/2*a*d*e*((c + 1)*log(d*x + c + 1)/d^2 - (c - 1)*log(d*x + c - 1)/d^2) + 1/2*a*c*e*(log(d*x + c + 1)/d - log(d*x + c - 1)/d) + 1/8*b*d*e*((2*(c + 1)*log(d*x + c + 1)*log(-d*x - c + 1) - (c - 1)*log(-d*x - c + 1)^2)/d^2 - 4*integrate(1/2*(c^2 + (c*d + 3*d)*x + 2*c + 1)*log(d*x + c + 1)/(d^3*x^2 + 2*c*d^2*x + c^2*d - d), x)) - 1/8*(log(d*x + c + 1)^2 - 2*log(d*x + c + 1)*log(d*x + c - 1) + log(d*x + c - 1)^2)*b*c*e/d`

Giac [F]

$$\int \frac{(ce + dex)(a + b \operatorname{arctanh}(c + dx))}{1 - (c + dx)^2} dx = \int -\frac{(dex + ce)(b \operatorname{artanh}(dx + c) + a)}{(dx + c)^2 - 1} dx$$

input `integrate((d*e*x+c*e)*(a+b*arctanh(d*x+c))/(1-(d*x+c)^2),x, algorithm="giac")`

output `integrate(-(d*e*x + c*e)*(b*arctanh(d*x + c) + a)/((d*x + c)^2 - 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(ce + dex)(a + b \operatorname{arctanh}(c + dx))}{1 - (c + dx)^2} dx = \int -\frac{(ce + dex)(a + b \operatorname{atanh}(c + dx))}{(c + dx)^2 - 1} dx$$

input `int(-((c*e + d*e*x)*(a + b*atanh(c + d*x)))/((c + d*x)^2 - 1),x)`

output `int(-((c*e + d*e*x)*(a + b*atanh(c + d*x)))/((c + d*x)^2 - 1), x)`

Reduce [F]

$$\int \frac{(ce + dex)(a + b \operatorname{arctanh}(c + dx))}{1 - (c + dx)^2} dx$$

$$= \frac{e \left(\operatorname{atanh}(dx + c)^2 bc - 2 \left(\int \frac{\operatorname{atanh}(dx+c)x}{d^2x^2+2cdx+c^2-1} dx \right) b d^2 - \log(dx + c - 1) a - \log(dx + c + 1) a \right)}{2d}$$

input `int((d*e*x+c*e)*(a+b*atanh(d*x+c))/(1-(d*x+c)^2),x)`

output `(e*(atanh(c + d*x)**2*b*c - 2*int((atanh(c + d*x)*x)/(c**2 + 2*c*d*x + d**2*x**2 - 1),x)*b*d**2 - log(c + d*x - 1)*a - log(c + d*x + 1)*a))/(2*d)`

CHAPTER 4

APPENDIX

| | |
|---|-----|
| 4.1 Listing of Grading functions | 698 |
| 4.2 Links to plain text integration problems used in this report for each CAS . | 716 |

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```

    Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
    If[AppellFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
      If[Head[expn]===RootSum,
        Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
        If[Head[expn]===Integrate || Head[expn]===Int,
          Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
          9]]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

```

```

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

```

```

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022  add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result    := ExpnType(result);
      ExpnType_optimal   := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#     is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```



```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```



```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file