

# Computer Algebra Independent Integration Tests

Summer 2024

7-Inverse-hyperbolic-functions/7.5-Inverse-hyperbolic-  
secant/346-7.5

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 10:16am

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
1.1	Listing of CAS systems tested . . . . .	4
1.2	Results . . . . .	5
1.3	Time and leaf size Performance . . . . .	9
1.4	Performance based on number of rules Rubi used . . . . .	11
1.5	Performance based on number of steps Rubi used . . . . .	12
1.6	Solved integrals histogram based on leaf size of result . . . . .	13
1.7	Solved integrals histogram based on CPU time used . . . . .	14
1.8	Leaf size vs. CPU time used . . . . .	15
1.9	list of integrals with no known antiderivative . . . . .	16
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	16
1.11	list of integrals solved by CAS but failed verification . . . . .	16
1.12	Timing . . . . .	17
1.13	Verification . . . . .	17
1.14	Important notes about some of the results . . . . .	18
1.15	Current tree layout of integration tests . . . . .	21
1.16	Design of the test system . . . . .	22
<b>2</b>	<b>detailed summary tables of results</b>	<b>23</b>
2.1	List of integrals sorted by grade for each CAS . . . . .	24
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	28
2.3	Detailed conclusion table specific for Rubi results . . . . .	37
<b>3</b>	<b>Listing of integrals</b>	<b>39</b>
3.1	$\int x^3 \operatorname{sech}^{-1}(a + bx) dx$ . . . . .	41
3.2	$\int x^2 \operatorname{sech}^{-1}(a + bx) dx$ . . . . .	48
3.3	$\int x \operatorname{sech}^{-1}(a + bx) dx$ . . . . .	55
3.4	$\int \operatorname{sech}^{-1}(a + bx) dx$ . . . . .	62
3.5	$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x} dx$ . . . . .	68
3.6	$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^2} dx$ . . . . .	78

3.7	$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^3} dx$	86
3.8	$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^4} dx$	96
3.9	$\int x^2 \operatorname{sech}^{-1}(a+bx)^2 dx$	107
3.10	$\int x \operatorname{sech}^{-1}(a+bx)^2 dx$	114
3.11	$\int \operatorname{sech}^{-1}(a+bx)^2 dx$	121
3.12	$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} dx$	128
3.13	$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx$	138
3.14	$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^3} dx$	145
3.15	$\int x \operatorname{sech}^{-1}(a+bx)^3 dx$	154
3.16	$\int \operatorname{sech}^{-1}(a+bx)^3 dx$	161
3.17	$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx$	168
3.18	$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^2} dx$	181
3.19	$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^3} dx$	188
3.20	$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx$	196
3.21	$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx$	202
3.22	$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx$	208
3.23	$\int \operatorname{sech}^{-1}(\sqrt{x}) dx$	214
3.24	$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx$	219
3.25	$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx$	226
3.26	$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx$	232
3.27	$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx$	238
3.28	$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx$	245
3.29	$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx$	250
3.30	$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx$	257
3.31	$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx$	264
3.32	$\int \frac{\operatorname{sech}^{-1}(a+bx)}{\frac{a}{b}+dx} dx$	271
3.33	$\int x^3 \operatorname{sech}^{-1}(a+bx^4) dx$	278
3.34	$\int x^{-1+n} \operatorname{sech}^{-1}(a+bx^n) dx$	284

**4 Appendix 290**

4.1	Listing of Grading functions	290
4.2	Links to plain text integration problems used in this report for each CAS308	

# CHAPTER 1

## INTRODUCTION

1.1	Listing of CAS systems tested . . . . .	4
1.2	Results . . . . .	5
1.3	Time and leaf size Performance . . . . .	9
1.4	Performance based on number of rules Rubi used . . . . .	11
1.5	Performance based on number of steps Rubi used . . . . .	12
1.6	Solved integrals histogram based on leaf size of result . . . . .	13
1.7	Solved integrals histogram based on CPU time used . . . . .	14
1.8	Leaf size vs. CPU time used . . . . .	15
1.9	list of integrals with no known antiderivative . . . . .	16
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	16
1.11	list of integrals solved by CAS but failed verification . . . . .	16
1.12	Timing . . . . .	17
1.13	Verification . . . . .	17
1.14	Important notes about some of the results . . . . .	18
1.15	Current tree layout of integration tests . . . . .	21
1.16	Design of the test system . . . . .	22

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [ 34 ]. This is test number [ 346 ].

## 1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

## 1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 ( 34 )	0.00 ( 0 )
Mathematica	94.12 ( 32 )	5.88 ( 2 )
Maple	76.47 ( 26 )	23.53 ( 8 )
Fricas	50.00 ( 17 )	50.00 ( 17 )
Maxima	32.35 ( 11 )	67.65 ( 23 )
Mupad	14.71 ( 5 )	85.29 ( 29 )
Giac	0.00 ( 0 )	100.00 ( 34 )
Reduce	0.00 ( 0 )	100.00 ( 34 )
Sympy	0.00 ( 0 )	100.00 ( 34 )

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> <li>1. antiderivative contains a hypergeometric function and the optimal antiderivative does not.</li> <li>2. antiderivative contains a special function and the optimal antiderivative does not.</li> <li>3. antiderivative contains the imaginary unit and the optimal antiderivative does not.</li> </ol>
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.

System	% A grade	% B grade	% C grade	% F grade
Rubi	76.471	0.000	23.529	0.000
Maple	64.706	2.941	8.824	23.529
Mathematica	52.941	23.529	17.647	5.882
Maxima	32.353	0.000	0.000	67.647
Fricas	26.471	23.529	0.000	50.000
Giac	0.000	0.000	0.000	100.000
Mupad	0.000	14.706	0.000	85.294
Reduce	0.000	0.000	0.000	100.000
Sympy	0.000	0.000	0.000	100.000


Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.





The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	2	50.00	50.00	0.00
Maple	8	100.00	0.00	0.00
Fricas	17	88.24	0.00	11.76
Maxima	23	100.00	0.00	0.00
Mupad	29	0.00	100.00	0.00
Giac	34	100.00	0.00	0.00
Reduce	34	100.00	0.00	0.00
Sympy	34	97.06	2.94	0.00

Table 1.4: Failure statistics for each CAS

## 1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.03
Fricas	0.12
Mathematica	0.43
Maple	0.49
Rubi	0.60
Mupad	4.09
Sympy	-nan(ind)
Reduce	-nan(ind)
Giac	-nan(ind)

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	42.00	0.92	43.00	1.08
Maxima	50.45	0.65	40.00	0.73
Rubi	175.03	1.05	104.50	1.01
Mathematica	227.62	1.65	162.50	1.19
Maple	230.96	1.60	111.00	1.32
Fricas	281.06	3.09	253.00	1.70
Sympy	-nan(ind)	-nan(ind)	nan	nan
Reduce	-nan(ind)	-nan(ind)	nan	nan
Giac	-nan(ind)	-nan(ind)	nan	nan

Table 1.6: Leaf size performance for each CAS

# 1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the  $y$  axis is the percentage solved which Rubi itself needed the number of rules given the  $x$  axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

## 1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

## 1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

## 1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

## 1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range



## 1.9 list of integrals with no known antiderivative

{}

## 1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

## 1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {24, 29, 30, 31, 32}

Mathematica {5, 13, 14, 15}

Maple {}

**Maxima** Verification phase not currently implemented.

**Fricas** Verification phase not currently implemented.

**Sympy** Verification phase not currently implemented.

**Giac** Verification phase not currently implemented.

**Reduce** Verification phase not currently implemented.

**Mupad** Verification phase not currently implemented.

## 1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

## 1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

## 1.14 Important notes about some of the results

### Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

## Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

## Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at [https://ask.sagemath.org/question/57123/could-we-have-a-leaf\\_count-function-in-base-sagemath/](https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/)

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

### Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives  $\sin(x)^2/2$


# 1.15 Current tree layout of integration tests



Figure 1.6: CAS integration tests tree

# 1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



**High level overview of the CAS independent integration test build system**

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

*The following fields are present only in Rubi Table file*

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi  
January 13, 2024  
Design note

# CHAPTER 2

## DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS . . . . .	24
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	28
2.3	Detailed conclusion table specific for Rubi results . . . . .	37



## 2.1 List of integrals sorted by grade for each CAS

Rubi . . . . .	24
Mma . . . . .	24
Maple . . . . .	25
Fricas . . . . .	25
Maxima . . . . .	25
Giac . . . . .	26
Mupad . . . . .	26
Sympy . . . . .	26
Reduce . . . . .	27

### Rubi

**A grade** { 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 13, 14, 15, 16, 18, 19, 20, 21, 22, 23, 25, 26, 27, 28, 33, 34 }

**B grade** { }

**C grade** { 5, 12, 17, 24, 29, 30, 31, 32 }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

### Mma

**A grade** { 8, 9, 10, 11, 12, 15, 16, 17, 20, 21, 22, 24, 25, 26, 27, 28, 30, 32 }

**B grade** { 4, 6, 7, 23, 29, 31, 33, 34 }

**C grade** { 1, 2, 3, 5, 13, 14 }

**F normal fail** { 19 }

**F(-1) timedout fail** { 18 }

**F(-2) exception fail** { }

## Maple

**A grade** { 1, 2, 3, 4, 9, 10, 11, 13, 14, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 31, 32, 33 }

**B grade** { 6 }

**C grade** { 5, 7, 8 }

**F normal fail** { 12, 15, 16, 17, 18, 19, 30, 34 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Fricas

**A grade** { 1, 20, 21, 22, 23, 25, 26, 27, 28 }

**B grade** { 2, 3, 4, 6, 7, 8, 33, 34 }

**C grade** { }

**F normal fail** { 5, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 24, 30, 32 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { 29, 31 }

## Maxima

**A grade** { 4, 20, 21, 22, 23, 25, 26, 27, 28, 33, 34 }

**B grade** { }

**C grade** { }

**F normal fail** { 1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 24, 29, 30, 31, 32 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Giac

A grade { }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34 }

F(-1) timedout fail { }

F(-2) exception fail { }

## Mupad

A grade { }

B grade { 4, 25, 28, 33, 34 }

C grade { }

F normal fail { }

F(-1) timedout fail { 1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 26, 27, 29, 30, 31, 32 }

F(-2) exception fail { }

## Sympy

A grade { }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33 }

F(-1) timedout fail { 34 }

F(-2) exception fail { }

## Reduce

**A grade** { }

**B grade** { }

**C grade** { }

**F normal fail** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24,  
25, 26, 27, 28, 29, 30, 31, 32, 33, 34 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## 2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as  $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$ . To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	203	206	225	250	0	345	0	0	12	0
N.S.	1	1.01	1.11	1.23	0.00	1.70	0.00	0.00	0.06	0.00
time (sec)	N/A	0.536	0.314	1.023	0.000	0.136	0.000	0.000	0.310	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	153	153	200	189	0	327	0	0	12	0
N.S.	1	1.00	1.31	1.24	0.00	2.14	0.00	0.00	0.08	0.00
time (sec)	N/A	0.401	0.146	0.914	0.000	0.121	0.000	0.000	0.249	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	106	176	111	0	308	0	0	10	0
N.S.	1	0.99	1.64	1.04	0.00	2.88	0.00	0.00	0.09	0.00
time (sec)	N/A	0.435	0.106	0.819	0.000	0.162	0.000	0.000	0.214	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	44	97	44	31	253	0	0	8	43
N.S.	1	1.16	2.55	1.16	0.82	6.66	0.00	0.00	0.21	1.13
time (sec)	N/A	0.235	0.162	0.575	0.028	0.155	0.000	0.000	0.205	4.334

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	C	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	170	233	332	886	0	0	0	0	12	0
N.S.	1	1.37	1.95	5.21	0.00	0.00	0.00	0.00	0.07	0.00
time (sec)	N/A	1.059	0.188	0.515	0.000	0.000	0.000	0.000	0.204	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	72	244	171	0	651	0	0	12	0
N.S.	1	1.03	3.49	2.44	0.00	9.30	0.00	0.00	0.17	0.00
time (sec)	N/A	0.425	0.167	0.921	0.000	0.144	0.000	0.000	0.207	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	133	159	315	370	0	865	0	0	12	0
N.S.	1	1.20	2.37	2.78	0.00	6.50	0.00	0.00	0.09	0.00
time (sec)	N/A	0.757	0.552	0.956	0.000	0.142	0.000	0.000	0.195	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	197	239	368	608	0	987	0	0	12	0
N.S.	1	1.21	1.87	3.09	0.00	5.01	0.00	0.00	0.06	0.00
time (sec)	N/A	1.066	0.224	0.971	0.000	0.164	0.000	0.000	0.209	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	279	266	305	599	0	0	0	0	14	0
N.S.	1	0.95	1.09	2.15	0.00	0.00	0.00	0.00	0.05	0.00
time (sec)	N/A	0.550	1.358	0.513	0.000	0.000	0.000	0.000	0.348	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	149	140	172	331	0	0	0	0	12	0
N.S.	1	0.94	1.15	2.22	0.00	0.00	0.00	0.00	0.08	0.00
time (sec)	N/A	0.443	0.309	0.422	0.000	0.000	0.000	0.000	0.253	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	77	105	192	0	0	0	0	10	0
N.S.	1	0.96	1.31	2.40	0.00	0.00	0.00	0.00	0.12	0.00
time (sec)	N/A	0.447	0.152	0.292	0.000	0.000	0.000	0.000	0.209	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	274	337	280	0	0	0	0	0	14	0
N.S.	1	1.23	1.02	0.00	0.00	0.00	0.00	0.00	0.05	0.00
time (sec)	N/A	1.583	0.208	0.000	0.000	0.000	0.000	0.000	0.194	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	224	227	678	362	0	0	0	0	14	0
N.S.	1	1.01	3.03	1.62	0.00	0.00	0.00	0.00	0.06	0.00
time (sec)	N/A	0.739	1.717	0.490	0.000	0.000	0.000	0.000	0.199	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	537	517	1439	982	0	0	0	0	14	0
N.S.	1	0.96	2.68	1.83	0.00	0.00	0.00	0.00	0.03	0.00
time (sec)	N/A	1.176	4.843	0.685	0.000	0.000	0.000	0.000	0.198	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	260	234	254	0	0	0	0	0	12	0
N.S.	1	0.90	0.98	0.00	0.00	0.00	0.00	0.00	0.05	0.00
time (sec)	N/A	0.565	0.398	0.000	0.000	0.000	0.000	0.000	0.264	0.000



Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	136	125	153	0	0	0	0	0	10	0
N.S.	1	0.92	1.12	0.00	0.00	0.00	0.00	0.00	0.07	0.00
time (sec)	N/A	0.582	0.086	0.000	0.000	0.000	0.000	0.000	0.197	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	378	452	384	0	0	0	0	0	14	0
N.S.	1	1.20	1.02	0.00	0.00	0.00	0.00	0.00	0.04	0.00
time (sec)	N/A	1.830	0.225	0.000	0.000	0.000	0.000	0.000	0.189	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	330	332	0	0	0	0	0	0	14	0
N.S.	1	1.01	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.00
time (sec)	N/A	0.917	0.000	0.000	0.000	0.000	0.000	0.000	0.204	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	965	911	0	0	0	0	0	0	14	0
N.S.	1	0.94	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00
time (sec)	N/A	1.777	0.000	0.000	0.000	0.000	0.000	0.000	0.204	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	138	103	84	54	58	57	0	0	9	0
N.S.	1	0.75	0.61	0.39	0.42	0.41	0.00	0.00	0.07	0.00
time (sec)	N/A	0.251	0.031	0.237	0.025	0.103	0.000	0.000	0.285	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	92	72	49	46	52	0	0	9	0
N.S.	1	0.86	0.67	0.46	0.43	0.49	0.00	0.00	0.08	0.00
time (sec)	N/A	0.243	0.022	0.192	0.026	0.098	0.000	0.000	0.286	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	79	56	42	34	45	0	0	7	0
N.S.	1	1.04	0.74	0.55	0.45	0.59	0.00	0.00	0.09	0.00
time (sec)	N/A	0.228	0.019	0.178	0.025	0.091	0.000	0.000	0.240	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	43	118	36	19	39	0	0	5	0
N.S.	1	1.13	3.11	0.95	0.50	1.03	0.00	0.00	0.13	0.00
time (sec)	N/A	0.187	0.083	0.168	0.030	0.105	0.000	0.000	0.210	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	F	F	F	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	58	45	65	0	0	0	0	9	0
N.S.	1	1.26	0.98	1.41	0.00	0.00	0.00	0.00	0.20	0.00
time (sec)	N/A	0.474	0.031	0.305	0.000	0.000	0.000	0.000	0.194	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	79	111	64	65	45	0	0	9	40
N.S.	1	0.85	1.19	0.69	0.70	0.48	0.00	0.00	0.10	0.43
time (sec)	N/A	0.219	0.057	0.187	0.025	0.096	0.000	0.000	0.197	4.239

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	102	125	79	92	54	0	0	9	0
N.S.	1	0.81	0.99	0.63	0.73	0.43	0.00	0.00	0.07	0.00
time (sec)	N/A	0.242	0.078	0.192	0.030	0.109	0.000	0.000	0.194	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	123	140	91	116	60	0	0	9	0
N.S.	1	0.78	0.89	0.58	0.74	0.38	0.00	0.00	0.06	0.00
time (sec)	N/A	0.250	0.083	0.197	0.025	0.106	0.000	0.000	0.200	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	25	20	16	22	0	0	6	17
N.S.	1	1.00	1.19	0.95	0.76	1.05	0.00	0.00	0.29	0.81
time (sec)	N/A	0.212	0.010	0.181	0.026	0.109	0.000	0.000	0.180	3.493

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	B	A	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	77	219	111	0	0	0	0	12	0
N.S.	1	1.26	3.59	1.82	0.00	0.00	0.00	0.00	0.20	0.00
time (sec)	N/A	0.526	0.644	0.655	0.000	0.000	0.000	0.000	0.188	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	54	68	49	0	0	0	0	0	12	0
N.S.	1	1.26	0.91	0.00	0.00	0.00	0.00	0.00	0.22	0.00
time (sec)	N/A	0.493	0.037	0.000	0.000	0.000	0.000	0.000	0.178	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	B	A	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	97	249	135	0	0	0	0	12	0
N.S.	1	1.26	3.23	1.75	0.00	0.00	0.00	0.00	0.16	0.00
time (sec)	N/A	0.497	1.125	0.635	0.000	0.000	0.000	0.000	0.208	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	F	F	F	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	67	52	111	0	0	0	0	21	0
N.S.	1	1.10	0.85	1.82	0.00	0.00	0.00	0.00	0.34	0.00
time (sec)	N/A	0.526	0.061	0.359	0.000	0.000	0.000	0.000	0.207	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	B	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	56	106	53	38	283	0	0	14	56
N.S.	1	1.14	2.16	1.08	0.78	5.78	0.00	0.00	0.29	1.14
time (sec)	N/A	0.327	0.109	0.198	0.025	0.116	0.000	0.000	0.267	4.481

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	A	B	F(-1)	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	50	56	106	0	40	385	0	0	17	54
N.S.	1	1.12	2.12	0.00	0.80	7.70	0.00	0.00	0.34	1.08
time (sec)	N/A	0.351	0.139	0.000	0.030	0.141	0.000	0.000	0.180	3.918

## 2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio  $\frac{\text{number of rules}}{\text{integrand size}}$  is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [5] had the largest ratio of [1.5000000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	9	8	1.01	10	0.800
2	A	6	5	1.00	10	0.500
3	A	10	9	0.99	8	1.125
4	A	5	4	1.16	6	0.667
5	C	16	15	1.37	10	1.500
6	A	8	7	1.03	10	0.700
7	A	13	12	1.20	10	1.200
8	A	14	13	1.21	10	1.300
9	A	6	5	0.95	12	0.417
10	A	7	6	0.94	10	0.600
11	A	8	7	0.96	8	0.875
12	C	17	16	1.23	12	1.333
13	A	6	5	1.01	12	0.417
14	A	7	6	0.96	12	0.500
15	A	7	6	0.90	10	0.600
16	A	9	8	0.92	8	1.000
17	C	17	16	1.20	12	1.333
18	A	6	5	1.01	12	0.417
19	A	7	6	0.94	12	0.500
20	A	4	4	0.75	10	0.400
21	A	4	4	0.86	10	0.400

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	4	4	1.04	8	0.500
23	A	2	2	1.13	6	0.333
24	C	10	9	1.26	10	0.900
25	A	6	5	0.85	10	0.500
26	A	7	6	0.81	10	0.600
27	A	8	7	0.78	10	0.700
28	A	3	3	1.00	4	0.750
29	C	10	9	1.26	10	0.900
30	C	10	9	1.26	10	0.900
31	C	10	9	1.26	10	0.900
32	C	11	10	1.10	19	0.526
33	A	6	5	1.14	12	0.417
34	A	6	5	1.12	14	0.357

# CHAPTER 3

## LISTING OF INTEGRALS

3.1	$\int x^3 \operatorname{sech}^{-1}(a + bx) dx$ . . . . .	41
3.2	$\int x^2 \operatorname{sech}^{-1}(a + bx) dx$ . . . . .	48
3.3	$\int x \operatorname{sech}^{-1}(a + bx) dx$ . . . . .	55
3.4	$\int \operatorname{sech}^{-1}(a + bx) dx$ . . . . .	62
3.5	$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x} dx$ . . . . .	68
3.6	$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^2} dx$ . . . . .	78
3.7	$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^3} dx$ . . . . .	86
3.8	$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^4} dx$ . . . . .	96
3.9	$\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx$ . . . . .	107
3.10	$\int x \operatorname{sech}^{-1}(a + bx)^2 dx$ . . . . .	114
3.11	$\int \operatorname{sech}^{-1}(a + bx)^2 dx$ . . . . .	121
3.12	$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} dx$ . . . . .	128
3.13	$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx$ . . . . .	138
3.14	$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^3} dx$ . . . . .	145
3.15	$\int x \operatorname{sech}^{-1}(a + bx)^3 dx$ . . . . .	154
3.16	$\int \operatorname{sech}^{-1}(a + bx)^3 dx$ . . . . .	161
3.17	$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx$ . . . . .	168
3.18	$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^2} dx$ . . . . .	181
3.19	$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^3} dx$ . . . . .	188
3.20	$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx$ . . . . .	196
3.21	$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx$ . . . . .	202
3.22	$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx$ . . . . .	208
3.23	$\int \operatorname{sech}^{-1}(\sqrt{x}) dx$ . . . . .	214
3.24	$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx$ . . . . .	219



---

3.25	$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx$	226
3.26	$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx$	232
3.27	$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx$	238
3.28	$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx$	245
3.29	$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx$	250
3.30	$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx$	257
3.31	$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx$	264
3.32	$\int \frac{\operatorname{sech}^{-1}(a+bx)}{\frac{a}{b}+dx} dx$	271
3.33	$\int x^3 \operatorname{sech}^{-1}(a+bx^4) dx$	278
3.34	$\int x^{-1+n} \operatorname{sech}^{-1}(a+bx^n) dx$	284

### 3.1 $\int x^3 \operatorname{sech}^{-1}(a + bx) dx$

Optimal result	41
Mathematica [C] (verified)	42
Rubi [A] (verified)	42
Maple [A] (verified)	45
Fricas [A] (verification not implemented)	45
Sympy [F]	46
Maxima [F]	46
Giac [F]	47
Mupad [F(-1)]	47
Reduce [F]	47

#### Optimal result

Integrand size = 10, antiderivative size = 203

$$\int x^3 \operatorname{sech}^{-1}(a + bx) dx = -\frac{(2 + 17a^2) \sqrt{\frac{1-a-bx}{1+a+bx}}(1 + a + bx)}{12b^4} - \frac{x^2 \sqrt{\frac{1-a-bx}{1+a+bx}}(1 + a + bx)}{12b^2}$$

$$+ \frac{a(a + bx) \sqrt{\frac{1-a-bx}{1+a+bx}}(1 + a + bx)}{3b^4} - \frac{a^4 \operatorname{sech}^{-1}(a + bx)}{4b^4}$$

$$+ \frac{1}{4} x^4 \operatorname{sech}^{-1}(a + bx) + \frac{a(1 + 2a^2) \arctan\left(\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{a+bx}\right)}{2b^4}$$

output

```
-1/12*(17*a^2+2)*((-b*x-a+1)/(b*x+a+1))^(1/2)*(b*x+a+1)/b^4-1/12*x^2*(-b*x-a+1)/(b*x+a+1)^(1/2)*(b*x+a+1)/b^2+1/3*a*(b*x+a)*((-b*x-a+1)/(b*x+a+1))^(1/2)*(b*x+a+1)/b^4-1/4*a^4*arcsech(b*x+a)/b^4+1/4*x^4*arcsech(b*x+a)+1/2*a*(2*a^2+1)*arctan(((b*x+a+1)/(b*x+a+1))^(1/2)*(b*x+a+1)/(b*x+a))/b^4
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.11

$$\int x^3 \operatorname{sech}^{-1}(a + bx) dx = \frac{\sqrt{-\frac{-1+a+bx}{1+a+bx}}(2 + 2a + 13a^2 + 13a^3 + (2 - 4a + 9a^2)bx + (1 - 3a)b^2x^2 + b^3x^3) - 3b^4x^4 \operatorname{sech}^{-1}(a + bx)}{b^4}$$

input

```
Integrate[x^3*ArcSech[a + b*x],x]
```

output

```
-1/12*(Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(2 + 2*a + 13*a^2 + 13*a^3 +
(2 - 4*a + 9*a^2)*b*x + (1 - 3*a)*b^2*x^2 + b^3*x^3) - 3*b^4*x^4*ArcSech[a
+ b*x] - 3*a^4*Log[a + b*x] + 3*a^4*Log[1 + Sqrt[-((-1 + a + b*x)/(1 + a
+ b*x))] + a*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))] + b*x*Sqrt[-((-1 + a +
b*x)/(1 + a + b*x))]) + (6*I)*a*(1 + 2*a^2)*Log[(-2*I)*(a + b*x) + 2*Sqrt[
-((-1 + a + b*x)/(1 + a + b*x))]*(1 + a + b*x)])/b^4
```

**Rubi [A] (verified)**

Time = 0.54 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.01, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$ , Rules used = {6875, 25, 5991, 3042, 4269, 3042, 4536, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \operatorname{sech}^{-1}(a + bx) dx$$

$$\downarrow \text{6875}$$

$$\frac{\int b^3 x^3 (a + bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a + bx + 1) \operatorname{sech}^{-1}(a + bx) d \operatorname{sech}^{-1}(a + bx)}{b^4}$$

$$\downarrow \text{25}$$

$$\begin{aligned}
& \frac{\int -b^3 x^3 (a + bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a + bx + 1) \operatorname{sech}^{-1}(a + bx) d\operatorname{sech}^{-1}(a + bx)}{b^4} \\
& \quad \downarrow 5991 \\
& \frac{-\frac{1}{4} \int b^4 x^4 d\operatorname{sech}^{-1}(a + bx) - \frac{1}{4} b^4 x^4 \operatorname{sech}^{-1}(a + bx)}{b^4} \\
& \quad \downarrow 3042 \\
& \frac{-\frac{1}{4} b^4 x^4 \operatorname{sech}^{-1}(a + bx) + \frac{1}{4} \int (a - \csc(i\operatorname{sech}^{-1}(a + bx) + \frac{\pi}{2}))^4 d\operatorname{sech}^{-1}(a + bx)}{b^4} \\
& \quad \downarrow 4269 \\
& \frac{\frac{1}{4} \left( \frac{1}{3} \int -bx(3a^3 + 8(a + bx)^2 a - (9a^2 + 2)(a + bx)) d\operatorname{sech}^{-1}(a + bx) + \frac{1}{3} b^2 x^2 \sqrt{\frac{-a-bx+1}{a+bx+1}} (a + bx + 1) \right) - \frac{1}{4} b^4 x^4}{b^4} \\
& \quad \downarrow 3042 \\
& \frac{-\frac{1}{4} b^4 x^4 \operatorname{sech}^{-1}(a + bx) + \frac{1}{4} \left( \frac{1}{3} b^2 x^2 \sqrt{\frac{-a-bx+1}{a+bx+1}} (a + bx + 1) + \frac{1}{3} \int (a - \csc(i\operatorname{sech}^{-1}(a + bx) + \frac{\pi}{2})) (3a^3 + 8 \csc(i\operatorname{sech}^{-1}(a + bx) + \frac{\pi}{2})) d\operatorname{sech}^{-1}(a + bx) \right)}{b^4} \\
& \quad \downarrow 4536 \\
& \frac{\frac{1}{4} \left( \frac{1}{3} \left( \frac{1}{2} \int (6a^4 - 12(2a^2 + 1)(a + bx)a + 2(17a^2 + 2)(a + bx)^2) d\operatorname{sech}^{-1}(a + bx) - 4a(a + bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a + bx + 1) \right) - \frac{1}{4} b^4 x^4 \operatorname{sech}^{-1}(a + bx) \right)}{b^4} \\
& \quad \downarrow 2009 \\
& \frac{\frac{1}{4} \left( \frac{1}{3} \left( \frac{1}{2} \left( 6a^4 \operatorname{sech}^{-1}(a + bx) - 12(2a^2 + 1) a \arctan \left( \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1)}{a+bx} \right) + 2(17a^2 + 2) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a + bx + 1) \right) - \frac{1}{4} b^4 x^4 \operatorname{sech}^{-1}(a + bx) \right) \right)}{b^4}
\end{aligned}$$

input `Int[x^3*ArcSech[a + b*x],x]`

output 
$$\begin{aligned}
& -\left( \frac{-1/4 * (b^4 * x^4 * \operatorname{ArcSech}[a + b*x]) + ((b^2 * x^2 * \sqrt{[(1 - a - b*x)/(1 + a + b*x)] * (1 + a + b*x)})/3 + (-4 * a * (a + b*x) * \sqrt{[(1 - a - b*x)/(1 + a + b*x)] * (1 + a + b*x)} + (2 * (2 + 17 * a^2) * \sqrt{[(1 - a - b*x)/(1 + a + b*x)] * (1 + a + b*x)} + 6 * a^4 * \operatorname{ArcSech}[a + b*x] - 12 * a * (1 + 2 * a^2) * \operatorname{ArcTan}[(\sqrt{[(1 - a - b*x)/(1 + a + b*x)] * (1 + a + b*x)})/(a + b*x)])/2)/3)/4 \right) / b^4
\end{aligned}$$

## Defintions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 2009  $\text{Int}[\text{u}_, \text{x\_Symbol}] \rightarrow \text{Simp}[\text{IntSum}[\text{u}, \text{x}], \text{x}] \text{ /; SumQ}[\text{u}]$
- rule 3042  $\text{Int}[\text{u}_, \text{x\_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ /; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4269  $\text{Int}[(\text{csc}[(\text{c}_.) + (\text{d}_.) * (\text{x}_)] * (\text{b}_.) + (\text{a}_.))^{\text{n}_.}, \text{x\_Symbol}] \rightarrow \text{Simp}[(-\text{b}^2) * \text{Cot}[\text{c} + \text{d} * \text{x}] * ((\text{a} + \text{b} * \text{Csc}[\text{c} + \text{d} * \text{x}])^{\text{n} - 2} / (\text{d} * (\text{n} - 1))), \text{x}] + \text{Simp}[1 / (\text{n} - 1) \text{ Int}[(\text{a} + \text{b} * \text{Csc}[\text{c} + \text{d} * \text{x}])^{\text{n} - 3} * \text{Simp}[\text{a}^3 * (\text{n} - 1) + (\text{b} * (\text{b}^2 * (\text{n} - 2) + 3 * \text{a}^2 * (\text{n} - 1))) * \text{Csc}[\text{c} + \text{d} * \text{x}] + (\text{a} * \text{b}^2 * (3 * \text{n} - 4)) * \text{Csc}[\text{c} + \text{d} * \text{x}]^2, \text{x}], \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \&\& \text{GtQ}[\text{n}, 2] \&\& \text{IntegerQ}[2 * \text{n}]$
- rule 4536  $\text{Int}[(\text{A}_.) + \text{csc}[(\text{e}_.) + (\text{f}_.) * (\text{x}_)] * (\text{B}_.) + \text{csc}[(\text{e}_.) + (\text{f}_.) * (\text{x}_)]^2 * (\text{C}_.) * (\text{csc}[(\text{e}_.) + (\text{f}_.) * (\text{x}_)] * (\text{b}_.) + (\text{a}_.)), \text{x\_Symbol}] \rightarrow \text{Simp}[(-\text{b}) * \text{C} * \text{Csc}[\text{e} + \text{f} * \text{x}] * (\text{Cot}[\text{e} + \text{f} * \text{x}] / (2 * \text{f})), \text{x}] + \text{Simp}[1 / 2 \quad \text{Int}[\text{Simp}[2 * \text{A} * \text{a} + (2 * \text{B} * \text{a} + \text{b} * (2 * \text{A} + \text{C})) * \text{Csc}[\text{e} + \text{f} * \text{x}] + 2 * (\text{a} * \text{C} + \text{B} * \text{b}) * \text{Csc}[\text{e} + \text{f} * \text{x}]^2, \text{x}], \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{e}, \text{f}, \text{A}, \text{B}, \text{C}\}, \text{x}]$
- rule 5991  $\text{Int}[(\text{e}_.) + (\text{f}_.) * (\text{x}_)]^{\text{m}_.} * \text{Sech}[(\text{c}_.) + (\text{d}_.) * (\text{x}_)] * ((\text{a}_.) + (\text{b}_.) * \text{Sech}[(\text{c}_.) + (\text{d}_.) * (\text{x}_)]^{\text{n}_.} * \text{Tanh}[(\text{c}_.) + (\text{d}_.) * (\text{x}_)]), \text{x\_Symbol}] \rightarrow \text{Simp}[(-\text{e} + \text{f} * \text{x})^{\text{m}} * ((\text{a} + \text{b} * \text{Sech}[\text{c} + \text{d} * \text{x}])^{\text{n} + 1} / (\text{b} * \text{d} * (\text{n} + 1))), \text{x}] + \text{Simp}[\text{f} * (\text{m} / (\text{b} * \text{d} * (\text{n} + 1))) \quad \text{Int}[(\text{e} + \text{f} * \text{x})^{\text{m} - 1} * (\text{a} + \text{b} * \text{Sech}[\text{c} + \text{d} * \text{x}])^{\text{n} + 1}, \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{n}\}, \text{x}] \&\& \text{IGtQ}[\text{m}, 0] \&\& \text{NeQ}[\text{n}, -1]$
- rule 6875  $\text{Int}[(\text{a}_.) + \text{ArcSech}[(\text{c}_.) + (\text{d}_.) * (\text{x}_)] * (\text{b}_.)]^{\text{p}_.} * ((\text{e}_.) + (\text{f}_.) * (\text{x}_)]^{\text{m}_.}, \text{x\_Symbol}] \rightarrow \text{Simp}[-(\text{d}^{\text{m} + 1})^{-1} \quad \text{Subst}[\text{Int}[(\text{a} + \text{b} * \text{x})^{\text{p}} * \text{Sech}[\text{x}] * \text{Tanh}[\text{x}] * (\text{d} * \text{e} - \text{c} * \text{f} + \text{f} * \text{Sech}[\text{x}])^{\text{m}}, \text{x}], \text{x}, \text{ArcSech}[\text{c} + \text{d} * \text{x}]], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \&\& \text{IGtQ}[\text{p}, 0] \&\& \text{IntegerQ}[\text{m}]$

**Maple [A] (verified)**

Time = 1.02 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.23

method	result
derivativedivides	$\frac{\frac{\operatorname{arcsech}(bx+a)a^4}{4} - \operatorname{arcsech}(bx+a)a^3(bx+a) + \frac{3 \operatorname{arcsech}(bx+a)a^2(bx+a)^2}{2} - \operatorname{arcsech}(bx+a)a(bx+a)^3 + \frac{\operatorname{arcsech}(bx+a)(bx+a)^4}{4}}{1}$
default	$\frac{\frac{\operatorname{arcsech}(bx+a)a^4}{4} - \operatorname{arcsech}(bx+a)a^3(bx+a) + \frac{3 \operatorname{arcsech}(bx+a)a^2(bx+a)^2}{2} - \operatorname{arcsech}(bx+a)a(bx+a)^3 + \frac{\operatorname{arcsech}(bx+a)(bx+a)^4}{4}}{1}$
parts	$\frac{x^4 \operatorname{arcsech}(bx+a)}{4} - \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \left( 3 \operatorname{csgn}(b) \operatorname{arctanh}\left(\frac{1}{\sqrt{-b^2x^2-2bxa-a^2+1}}\right) a^4 + \operatorname{csgn}(b)b^2x^2\sqrt{-b^2x^2-2bxa-a^2+1} \right)}{4}$

input `int(x^3*arcsech(b*x+a),x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{b^4} \left( \frac{1}{4} \operatorname{arcsech}(bx+a) a^4 - \operatorname{arcsech}(bx+a) a^3 (bx+a) + \frac{3}{2} \operatorname{arcsech}(bx+a) a^2 (bx+a)^2 - \operatorname{arcsech}(bx+a) a (bx+a)^3 + \frac{1}{4} \operatorname{arcsech}(bx+a) (bx+a)^4 - \frac{1}{12} \left( -\frac{bx+a-1}{bx+a} \right)^{1/2} (bx+a) \left( \frac{bx+a+1}{bx+a} \right)^{1/2} \left( 3a^4 \operatorname{arctanh}\left(\frac{1}{\sqrt{1-(bx+a)^2}}\right) + 12a^3 \operatorname{arcsin}(bx+a) + 18a^2 \sqrt{1-(bx+a)^2} - 6a \sqrt{1-(bx+a)^2} + \sqrt{1-(bx+a)^2} (bx+a)^2 + 6a \operatorname{arcsin}(bx+a) + 2 \sqrt{1-(bx+a)^2} \right) / \sqrt{1-(bx+a)^2} \right)$$

**Fricas [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 345, normalized size of antiderivative = 1.70

$$\int x^3 \operatorname{sech}^{-1}(a+bx) dx$$

$$= \frac{6b^4x^4 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}+1}}{bx+a}\right) - 3a^4 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}+1}}{x}\right) + 3a^4 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}+1}}{x}\right)}{1}$$

input `integrate(x^3*arcsech(b*x+a),x, algorithm="fricas")`

output

```
1/24*(6*b^4*x^4*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)) - 3*a^4*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/x) + 3*a^4*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) - 1)/x) + 12*(2*a^3 + a)*arctan((b^2*x^2 + 2*a*b*x + a^2)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)))/(b^2*x^2 + 2*a*b*x + a^2 - 1)) - 2*(b^3*x^3 - 3*a*b^2*x^2 + 13*a^3 + (9*a^2 + 2)*b*x + 2*a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)))/b^4
```

### Sympy [F]

$$\int x^3 \operatorname{sech}^{-1}(a + bx) dx = \int x^3 \operatorname{asech}(a + bx) dx$$

input

```
integrate(x**3*asech(b*x+a), x)
```

output

```
Integral(x**3*asech(a + b*x), x)
```

### Maxima [F]

$$\int x^3 \operatorname{sech}^{-1}(a + bx) dx = \int x^3 \operatorname{arsech}(bx + a) dx$$

input

```
integrate(x^3*arcsech(b*x+a), x, algorithm="maxima")
```

output

```
1/8*(2*b^4*x^4*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a) - 2*b^4*x^4*log(b*x + a) - b^2*x^2 + 6*a*b*x - (a^4 + 4*a^3 + 6*a^2 + 4*a + 1)*log(b*x + a + 1) - 2*(b^4*x^4 - a^4)*log(b*x + a) - (a^4 - 4*a^3 + 6*a^2 - 4*a + 1)*log(-b*x - a + 1))/b^4 + integrate(1/4*(b^2*x^5 + a*b*x^4)/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^(1/2*log(b*x + a + 1) + 1/2*log(-b*x - a + 1)) - 1), x)
```

**Giac [F]**

$$\int x^3 \operatorname{sech}^{-1}(a + bx) dx = \int x^3 \operatorname{arsech}(bx + a) dx$$

input `integrate(x^3*arcsech(b*x+a),x, algorithm="giac")`

output `integrate(x^3*arcsech(b*x + a), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 \operatorname{sech}^{-1}(a + bx) dx = \int x^3 \operatorname{acosh}\left(\frac{1}{a + bx}\right) dx$$

input `int(x^3*acosh(1/(a + b*x)),x)`

output `int(x^3*acosh(1/(a + b*x)), x)`

**Reduce [F]**

$$\int x^3 \operatorname{sech}^{-1}(a + bx) dx = \int \operatorname{asech}(bx + a) x^3 dx$$

input `int(x^3*asech(b*x+a),x)`

output `int(asech(a + b*x)*x**3,x)`



### 3.2 $\int x^2 \operatorname{sech}^{-1}(a + bx) dx$

Optimal result	48
Mathematica [C] (verified)	49
Rubi [A] (verified)	49
Maple [A] (verified)	51
Fricas [B] (verification not implemented)	52
Sympy [F]	53
Maxima [F]	53
Giac [F]	53
Mupad [F(-1)]	54
Reduce [F]	54

#### Optimal result

Integrand size = 10, antiderivative size = 153

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx = \frac{5a\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6b^3} - \frac{x\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6b^2} + \frac{a^3 \operatorname{sech}^{-1}(a+bx)}{3b^3} + \frac{1}{3}x^3 \operatorname{sech}^{-1}(a+bx) - \frac{(1+6a^2) \arctan\left(\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{a+bx}\right)}{6b^3}$$

output `5/6*a*((-b*x-a+1)/(b*x+a+1))^(1/2)*(b*x+a+1)/b^3-1/6*x*((-b*x-a+1)/(b*x+a+1))^(1/2)*(b*x+a+1)/b^2+1/3*a^3*arcsech(b*x+a)/b^3+1/3*x^3*arcsech(b*x+a)-1/6*(6*a^2+1)*arctan(((b*x+a+1)/(-b*x-a+1))^(1/2)*(b*x+a+1)/(b*x+a))/b^3`

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.31

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx$$

$$= \sqrt{-\frac{-1+a+bx}{1+a+bx}} (5a^2 - bx(1 + bx) + a(5 + 4bx)) + 2b^3 x^3 \operatorname{sech}^{-1}(a + bx) - 2a^3 \log(a + bx) + 2a^3 \log\left(1 + \sqrt{-\frac{-1+a+bx}{1+a+bx}}\right)$$

input

```
Integrate[x^2*ArcSech[a + b*x],x]
```

output

```
(Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(5*a^2 - b*x*(1 + b*x) + a*(5 + 4*b*x)) + 2*b^3*x^3*ArcSech[a + b*x] - 2*a^3*Log[a + b*x] + 2*a^3*Log[1 + Sqrt[-((-1 + a + b*x)/(1 + a + b*x))] + a*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))] + b*x*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))] + I*(1 + 6*a^2)*Log[(-2*I)*(a + b*x) + 2*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(1 + a + b*x)])/(6*b^3)
```

**Rubi [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6875, 5991, 3042, 4269, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx$$

$$\downarrow 6875$$

$$\frac{\int b^2 x^2 (a + bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a + bx + 1) \operatorname{sech}^{-1}(a + bx) d \operatorname{sech}^{-1}(a + bx)}{b^3}$$

$$\downarrow 5991$$

$$\begin{aligned}
 & \frac{-\frac{1}{3} \int -b^3 x^3 d\operatorname{sech}^{-1}(a+bx) - \frac{1}{3} b^3 x^3 \operatorname{sech}^{-1}(a+bx)}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\frac{1}{3} b^3 x^3 \operatorname{sech}^{-1}(a+bx) - \frac{1}{3} \int (a - \csc(\operatorname{sech}^{-1}(a+bx) + \frac{\pi}{2}))^3 d\operatorname{sech}^{-1}(a+bx)}{b^3} \\
 & \quad \downarrow \text{4269} \\
 & \frac{\frac{1}{3} \left( \frac{1}{2} b x \sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1) - \frac{1}{2} \int (2a^3 + 5(a+bx)^2 a - (6a^2+1)(a+bx)) d\operatorname{sech}^{-1}(a+bx) \right) - \frac{1}{3} b^3 x^3 \operatorname{sech}^{-1}(a+bx)}{b^3} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\frac{1}{3} \left( \frac{1}{2} \left( -2a^3 \operatorname{sech}^{-1}(a+bx) + (6a^2+1) \arctan \left( \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1)}{a+bx} \right) - 5a \sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1) \right) + \frac{1}{2} b x \sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1) \right)}{b^3}
 \end{aligned}$$

input `Int[x^2*ArcSech[a + b*x], x]`

output `-((-1/3*(b^3*x^3*ArcSech[a + b*x]) + (b*x*Sqrt[(1 - a - b*x)/(1 + a + b*x)])*(1 + a + b*x))/2 + (-5*a*Sqrt[(1 - a - b*x)/(1 + a + b*x)])*(1 + a + b*x) - 2*a^3*ArcSech[a + b*x] + (1 + 6*a^2)*ArcTan[(Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x))/(a + b*x)]/2)/3)/b^3)`

### Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4269

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_), x_Symbol] := Simp[(-b^2)*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n - 2)/(d*(n - 1))), x] + Simp[1/(n - 1) Int[(a + b*Csc[c + d*x])^(n - 3)*Simp[a^3*(n - 1) + (b*(b^2*(n - 2) + 3*a^2*(n - 1)))*Csc[c + d*x] + (a*b^2*(3*n - 4))*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 2] && IntegerQ[2*n]
```

rule 5991

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_) + (b_.)*Sech[(c_.) + (d_.)*(x_)]^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

rule 6875

```
Int[((a_.) + ArcSech[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

### Maple [A] (verified)

Time = 0.91 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.24

method	result
derivativedivides	$\frac{-\frac{\operatorname{arcsech}(bx+a)a^3}{3} + \operatorname{arcsech}(bx+a)a^2(bx+a) - \operatorname{arcsech}(bx+a)a(bx+a)^2 + \frac{\operatorname{arcsech}(bx+a)(bx+a)^3}{3} + \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}}}{b^3}}{b^3}$
default	$\frac{-\frac{\operatorname{arcsech}(bx+a)a^3}{3} + \operatorname{arcsech}(bx+a)a^2(bx+a) - \operatorname{arcsech}(bx+a)a(bx+a)^2 + \frac{\operatorname{arcsech}(bx+a)(bx+a)^3}{3} + \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}}}{b^3}}{b^3}$
parts	$\frac{x^3 \operatorname{arcsech}(bx+a)}{3} + \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}}}{b^3} \left( 2 \operatorname{csign}(b) \operatorname{arctanh}\left(\frac{1}{\sqrt{-b^2x^2 - 2bxa - a^2 + 1}}\right) a^3 - \sqrt{-b^2x^2 - 2bxa - a^2 + 1} \right)$

input

```
int(x^2*arcsech(b*x+a), x, method=_RETURNVERBOSE)
```

output

```
1/b^3*(-1/3*arcsech(b*x+a)*a^3+arcsech(b*x+a)*a^2*(b*x+a)-arcsech(b*x+a)*
*(b*x+a)^2+1/3*arcsech(b*x+a)*(b*x+a)^3+1/6*(-(b*x+a-1)/(b*x+a))^(1/2)*(b*
x+a)*((b*x+a+1)/(b*x+a))^(1/2)*(2*a^3*arctanh(1/(1-(b*x+a)^2)^(1/2))+6*a^2
*arcsin(b*x+a)+6*a*(1-(b*x+a)^2)^(1/2)-(b*x+a)*(1-(b*x+a)^2)^(1/2)+arcsin(
b*x+a))/(1-(b*x+a)^2)^(1/2))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 327 vs.  $2(131) = 262$ .

Time = 0.12 (sec) , antiderivative size = 327, normalized size of antiderivative = 2.14

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx$$

$$= \frac{2b^3x^3 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}+1}{bx+a}\right) + a^3 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}+1}{x}\right) - a^3 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}}{x}\right)}{1}$$

6

input

```
integrate(x^2*arcsech(b*x+a),x, algorithm="fricas")
```

output

```
1/6*(2*b^3*x^3*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 +
2*a*b*x + a^2)) + 1)/(b*x + a)) + a^3*log(((b*x + a)*sqrt(-(b^2*x^2 + 2
*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/x) - a^3*log(((b*x + a)*
sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) - 1)/x) - (
6*a^2 + 1)*arctan((b^2*x^2 + 2*a*b*x + a^2)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2
- 1)/(b^2*x^2 + 2*a*b*x + a^2))/(b^2*x^2 + 2*a*b*x + a^2 - 1)) - (b^2*x^2
- 4*a*b*x - 5*a^2)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x
+ a^2)))/b^3
```

**Sympy [F]**

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx = \int x^2 \operatorname{arsech}(a + bx) dx$$

input `integrate(x**2*asech(b*x+a),x)`

output `Integral(x**2*asech(a + b*x), x)`

**Maxima [F]**

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx = \int x^2 \operatorname{arsech}(bx + a) dx$$

input `integrate(x^2*arcsech(b*x+a),x, algorithm="maxima")`

output `1/6*(2*b^3*x^3*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a) - 2*b^3*x^3*log(b*x + a) - 2*b*x + (a^3 + 3*a^2 + 3*a + 1)*log(b*x + a + 1) - 2*(b^3*x^3 + a^3)*log(b*x + a) + (a^3 - 3*a^2 + 3*a - 1)*log(-b*x - a + 1))/b^3 + integrate(1/3*(b^2*x^4 + a*b*x^3)/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^(1/2*log(b*x + a + 1) + 1/2*log(-b*x - a + 1)) - 1), x)`

**Giac [F]**

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx = \int x^2 \operatorname{arsech}(bx + a) dx$$

input `integrate(x^2*arcsech(b*x+a),x, algorithm="giac")`

output `integrate(x^2*arcsech(b*x + a), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx = \int x^2 \operatorname{acosh}\left(\frac{1}{a + bx}\right) dx$$

input `int(x^2*acosh(1/(a + b*x)),x)`output `int(x^2*acosh(1/(a + b*x)), x)`**Reduce [F]**

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx = \int \operatorname{asech}(bx + a) x^2 dx$$

input `int(x^2*asech(b*x+a),x)`output `int(asech(a + b*x)*x**2,x)`

### 3.3 $\int x \operatorname{sech}^{-1}(a + bx) dx$

Optimal result . . . . .	55
Mathematica [C] (verified) . . . . .	55
Rubi [A] (verified) . . . . .	56
Maple [A] (verified) . . . . .	59
Fricas [B] (verification not implemented) . . . . .	59
Sympy [F] . . . . .	60
Maxima [F] . . . . .	60
Giac [F] . . . . .	61
Mupad [F(-1)] . . . . .	61
Reduce [F] . . . . .	61

#### Optimal result

Integrand size = 8, antiderivative size = 107

$$\int x \operatorname{sech}^{-1}(a + bx) dx = -\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{2b^2} - \frac{a^2 \operatorname{sech}^{-1}(a+bx)}{2b^2} + \frac{1}{2}x^2 \operatorname{sech}^{-1}(a+bx) + \frac{a \arctan\left(\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{a+bx}\right)}{b^2}$$

output

```
-1/2*((-b*x-a+1)/(b*x+a+1))^(1/2)*(b*x+a+1)/b^2-1/2*a^2*arcsech(b*x+a)/b^2
+1/2*x^2*arcsech(b*x+a)+a*arctan(((b*x+a+1)/(b*x+a+1))^(1/2)*(b*x+a+1)/(b
*x+a))/b^2
```

#### Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.64

$$\int x \operatorname{sech}^{-1}(a + bx) dx = \frac{-\sqrt{-\frac{-1+a+bx}{1+a+bx}}(1+a+bx) + b^2 x^2 \operatorname{sech}^{-1}(a+bx) + a^2 \log(a+bx) - a^2 \log\left(1 + \sqrt{-\frac{-1+a+bx}{1+a+bx}} + a \sqrt{-\frac{-1}{1+a+bx}}\right)}{2b^2}$$



input `Integrate[x*ArcSech[a + b*x],x]`

output 
$$\begin{aligned} & (-\sqrt{-((-1 + a + b*x)/(1 + a + b*x))}*(1 + a + b*x)) + b^2*x^2*ArcSech[ \\ & a + b*x] + a^2*Log[a + b*x] - a^2*Log[1 + \sqrt{-((-1 + a + b*x)/(1 + a + b \\ & *x))}] + a*\sqrt{-((-1 + a + b*x)/(1 + a + b*x))} + b*x*\sqrt{-((-1 + a + b*x \\ & )/(1 + a + b*x))}] - (2*I)*a*Log[(-2*I)*(a + b*x) + 2*\sqrt{-((-1 + a + b*x \\ & )/(1 + a + b*x))}*(1 + a + b*x)]/(2*b^2) \end{aligned}$$

### Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.99, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.125$ , Rules used = {6875, 25, 5991, 3042, 4260, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \operatorname{sech}^{-1}(a + bx) dx \\ & \quad \downarrow \text{6875} \\ & \frac{\int bx(a + bx) \sqrt{\frac{-a - bx + 1}{a + bx + 1}} (a + bx + 1) \operatorname{sech}^{-1}(a + bx) d \operatorname{sech}^{-1}(a + bx)}{b^2} \\ & \quad \downarrow \text{25} \\ & \frac{\int -bx(a + bx) \sqrt{\frac{-a - bx + 1}{a + bx + 1}} (a + bx + 1) \operatorname{sech}^{-1}(a + bx) d \operatorname{sech}^{-1}(a + bx)}{b^2} \\ & \quad \downarrow \text{5991} \\ & \frac{\frac{1}{2} \int b^2 x^2 d \operatorname{sech}^{-1}(a + bx) - \frac{1}{2} b^2 x^2 \operatorname{sech}^{-1}(a + bx)}{b^2} \\ & \quad \downarrow \text{3042} \\ & \frac{-\frac{1}{2} b^2 x^2 \operatorname{sech}^{-1}(a + bx) + \frac{1}{2} \int (a - \csc(\operatorname{isech}^{-1}(a + bx) + \frac{\pi}{2}))^2 d \operatorname{sech}^{-1}(a + bx)}{b^2} \\ & \quad \downarrow \text{4260} \end{aligned}$$

$$\frac{\frac{1}{2}(-2a \int (a + bx) d\operatorname{sech}^{-1}(a + bx) + \int (a + bx)^2 d\operatorname{sech}^{-1}(a + bx) + a^2 \operatorname{sech}^{-1}(a + bx)) - \frac{1}{2}b^2 x^2 \operatorname{sech}^{-1}(a + bx)}{b^2}$$

↓ 3042

$$\frac{-\frac{1}{2}b^2 x^2 \operatorname{sech}^{-1}(a + bx) + \frac{1}{2} \left( -2a \int \csc \left( i \operatorname{sech}^{-1}(a + bx) + \frac{\pi}{2} \right) d\operatorname{sech}^{-1}(a + bx) + \int \csc \left( i \operatorname{sech}^{-1}(a + bx) + \frac{\pi}{2} \right)^2 \right)}{b^2}$$

↓ 4254

$$\frac{-\frac{1}{2}b^2 x^2 \operatorname{sech}^{-1}(a + bx) + \frac{1}{2} \left( i \int 1 d \left( -i \sqrt{\frac{-a-bx+1}{a+bx+1}} (a + bx + 1) \right) - 2a \int \csc \left( i \operatorname{sech}^{-1}(a + bx) + \frac{\pi}{2} \right) d\operatorname{sech}^{-1}(a + bx) \right)}{b^2}$$

↓ 24

$$\frac{-\frac{1}{2}b^2 x^2 \operatorname{sech}^{-1}(a + bx) + \frac{1}{2} \left( -2a \int \csc \left( i \operatorname{sech}^{-1}(a + bx) + \frac{\pi}{2} \right) d\operatorname{sech}^{-1}(a + bx) + a^2 \operatorname{sech}^{-1}(a + bx) + \sqrt{\frac{-a-bx+1}{a+bx+1}} \right)}{b^2}$$

↓ 4257

$$\frac{\frac{1}{2} \left( a^2 \operatorname{sech}^{-1}(a + bx) - 2a \arctan \left( \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1)}{a+bx} \right) + \sqrt{\frac{-a-bx+1}{a+bx+1}} (a + bx + 1) \right) - \frac{1}{2}b^2 x^2 \operatorname{sech}^{-1}(a + bx)}{b^2}$$

input `Int[x*ArcSech[a + b*x], x]`

output `-((-1/2*(b^2*x^2*ArcSech[a + b*x]) + (Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x) + a^2*ArcSech[a + b*x] - 2*a*ArcTan[(Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x))/(a + b*x]))/2)/b^2)`

## Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`
- rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4260 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_))^2, x_Symbol] := Simp[a^2*x, x] + (Simp[2*a*b Int[Csc[c + d*x], x], x] + Simp[b^2 Int[Csc[c + d*x]^2, x], x]) /; FreeQ[{a, b, c, d}, x]`
- rule 5991 `Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_) + (b_.)*Sech[(c_.) + (d_.)*(x_)]^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`
- rule 6875 `Int[((a_.) + ArcSech[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

### Maple [A] (verified)

Time = 0.82 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.04

method	result
derivativedivides	$\frac{\frac{\operatorname{arcsech}(bx+a)(bx+a)^2}{2} - \operatorname{arcsech}(bx+a)a(bx+a) - \sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}}(2a \arcsin(bx+a) + \sqrt{1-(bx+a)^2})}{b^2}$
default	$\frac{\frac{\operatorname{arcsech}(bx+a)(bx+a)^2}{2} - \operatorname{arcsech}(bx+a)a(bx+a) - \sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}}(2a \arcsin(bx+a) + \sqrt{1-(bx+a)^2})}{b^2}$
parts	$\frac{x^2 \operatorname{arcsech}(bx+a)}{2} - \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \left( \operatorname{csgn}(b) \operatorname{arctanh}\left(\frac{1}{\sqrt{-b^2x^2-2bxa-a^2+1}}\right) a^2 + \sqrt{-b^2x^2-2bxa-a^2+1} \right)}{2b^2\sqrt{-b^2x^2-2bxa-a^2+1}}$

input `int(x*arcsech(b*x+a), x, method=_RETURNVERBOSE)`

output 
$$\frac{1}{b^2} \left( \frac{1}{2} \operatorname{arcsech}(bx+a) (bx+a)^2 - \operatorname{arcsech}(bx+a) a (bx+a) - \frac{1}{2} \left( -\frac{bx+a-1}{bx+a} \right)^{1/2} (bx+a) \left( \frac{bx+a+1}{bx+a} \right)^{1/2} (2a \arcsin(bx+a) + (1 - \frac{bx+a-1}{bx+a})^{1/2}) \right) / (1 - (bx+a)^2)^{1/2}$$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 308 vs. 2(93) = 186.

Time = 0.16 (sec) , antiderivative size = 308, normalized size of antiderivative = 2.88

$$\int x \operatorname{sech}^{-1}(a + bx) dx$$

$$= \frac{2b^2x^2 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}+1}{bx+a}\right) - a^2 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}+1}{x}\right) + a^2 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}}{x}\right)}{4b^2}$$

input `integrate(x*arcsech(b*x+a), x, algorithm="fricas")`

output

```
1/4*(2*b^2*x^2*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)) - a^2*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/x) + a^2*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) - 1)/x) + 4*a*arctan((b^2*x^2 + 2*a*b*x + a^2)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)))/(b^2*x^2 + 2*a*b*x + a^2 - 1)) - 2*(b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)))/b^2
```

**Sympy [F]**

$$\int x \operatorname{sech}^{-1}(a + bx) dx = \int x \operatorname{asech}(a + bx) dx$$

input

```
integrate(x*asech(b*x+a),x)
```

output

```
Integral(x*asech(a + b*x), x)
```

**Maxima [F]**

$$\int x \operatorname{sech}^{-1}(a + bx) dx = \int x \operatorname{arsech}(bx + a) dx$$

input

```
integrate(x*arcsech(b*x+a),x, algorithm="maxima")
```

output

```
1/4*(2*b^2*x^2*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a) - 2*b^2*x^2*log(b*x + a) - (a^2 + 2*a + 1)*log(b*x + a + 1) - 2*(b^2*x^2 - a^2)*log(b*x + a) - (a^2 - 2*a + 1)*log(-b*x - a + 1))/b^2 + integrate(1/2*(b^2*x^3 + a*b*x^2)/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^(1/2*log(b*x + a + 1) + 1/2*log(-b*x - a + 1)) - 1), x)
```

**Giac [F]**

$$\int x \operatorname{sech}^{-1}(a + bx) dx = \int x \operatorname{arsech}(bx + a) dx$$

input `integrate(x*arcsech(b*x+a),x, algorithm="giac")`

output `integrate(x*arcsech(b*x + a), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{sech}^{-1}(a + bx) dx = \int x \operatorname{acosh}\left(\frac{1}{a + bx}\right) dx$$

input `int(x*acosh(1/(a + b*x)),x)`

output `int(x*acosh(1/(a + b*x)), x)`

**Reduce [F]**

$$\int x \operatorname{sech}^{-1}(a + bx) dx = \int \operatorname{asech}(bx + a) x dx$$

input `int(x*asech(b*x+a),x)`

output `int(asech(a + b*x)*x,x)`

### 3.4 $\int \operatorname{sech}^{-1}(a + bx) dx$

Optimal result	62
Mathematica [B] (verified)	62
Rubi [A] (verified)	63
Maple [A] (verified)	64
Fricas [B] (verification not implemented)	65
Sympy [F]	65
Maxima [A] (verification not implemented)	66
Giac [F]	66
Mupad [B] (verification not implemented)	66
Reduce [F]	67

#### Optimal result

Integrand size = 6, antiderivative size = 38

$$\int \operatorname{sech}^{-1}(a + bx) dx = \frac{(a + bx)\operatorname{sech}^{-1}(a + bx)}{b} - \frac{2 \arctan\left(\sqrt{-1 + \frac{2}{1+a+bx}}\right)}{b}$$

output

```
(b*x+a)*arcsech(b*x+a)/b-2*arctan((-1+2/(b*x+a+1))^(1/2))/b
```

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 97 vs. 2(38) = 76.

Time = 0.16 (sec) , antiderivative size = 97, normalized size of antiderivative = 2.55

$$\int \operatorname{sech}^{-1}(a + bx) dx = x \operatorname{sech}^{-1}(a + bx) - \frac{2\sqrt{-\frac{-1+a+bx}{1+a+bx}} \left( -a \arctan\left(\sqrt{\frac{-1+a+bx}{1+a+bx}}\right) + \operatorname{arctanh}\left(\sqrt{\frac{-1+a+bx}{1+a+bx}}\right) \right)}{b\sqrt{\frac{-1+a+bx}{1+a+bx}}}$$

input

```
Integrate[ArcSech[a + b*x], x]
```

output

```
x*ArcSech[a + b*x] - (2*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(-(a*ArcTan[
Sqrt[(-1 + a + b*x)/(1 + a + b*x)]) + ArcTanh[Sqrt[(-1 + a + b*x)/(1 + a
+ b*x)]])))/(b*Sqrt[(-1 + a + b*x)/(1 + a + b*x)])
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.16, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {6867, 2055, 27, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \operatorname{sech}^{-1}(a + bx) dx \\
 & \quad \downarrow \text{6867} \\
 & \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}}{-a-bx+1} dx + \frac{(a+bx)\operatorname{sech}^{-1}(a+bx)}{b} \\
 & \quad \downarrow \text{2055} \\
 & \frac{(a+bx)\operatorname{sech}^{-1}(a+bx)}{b} - 4b \int \frac{1}{2b^2 \left( \frac{-a-bx+1}{a+bx+1} + 1 \right)} d\sqrt{\frac{-a-bx+1}{a+bx+1}} \\
 & \quad \downarrow \text{27} \\
 & \frac{(a+bx)\operatorname{sech}^{-1}(a+bx)}{b} - \frac{2 \int \frac{1}{\frac{-a-bx+1}{a+bx+1} + 1} d\sqrt{\frac{-a-bx+1}{a+bx+1}}}{b} \\
 & \quad \downarrow \text{216} \\
 & \frac{(a+bx)\operatorname{sech}^{-1}(a+bx)}{b} - \frac{2 \arctan \left( \sqrt{\frac{-a-bx+1}{a+bx+1}} \right)}{b}
 \end{aligned}$$

input

```
Int[ArcSech[a + b*x], x]
```



output  $((a + b*x)*\text{ArcSech}[a + b*x])/b - (2*\text{ArcTan}[\text{Sqrt}[(1 - a - b*x)/(1 + a + b*x)]])/b$

**Defintions of rubi rules used**

rule 27  $\text{Int}[(a_)*(F_x_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_)] /; \text{FreeQ}[b, x]$

rule 216  $\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 2055  $\text{Int}[(u_)^{(r_)*((e_)*((a_ + (b_)*(x_)^{(n_)})))/((c_ + (d_)*(x_)^{(n_)}))^{(p_)}, x\_Symbol] \rightarrow \text{With}\{q = \text{Denominator}[p]\}, \text{Simp}[q*e*((b*c - a*d)/n) \text{ Subst}[\text{Int}[\text{SimplifyIntegrand}[x^{(q*(p + 1) - 1)}*(((-a)*e + c*x^q)^{(1/n - 1)}/(b*e - d*x^q)^{(1/n + 1)})*(u /. x \rightarrow ((-a)*e + c*x^q)^{(1/n)}/(b*e - d*x^q)^{(1/n)})^r, x], x], x, (e*((a + b*x^n)/(c + d*x^n)))^{(1/q)}, x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \ \&\& \ \text{PolynomialQ}[u, x] \ \&\& \ \text{FractionQ}[p] \ \&\& \ \text{IntegerQ}[1/n] \ \&\& \ \text{IntegerQ}[r]$

rule 6867  $\text{Int}[\text{ArcSech}[(c_ + (d_)*(x_)], x\_Symbol] \rightarrow \text{Simp}[(c + d*x)*(\text{ArcSech}[c + d*x]/d), x] + \text{Int}[\text{Sqrt}[(1 - c - d*x)/(1 + c + d*x)]/(1 - c - d*x), x] /; \text{FreeQ}\{c, d\}, x]$

**Maple [A] (verified)**

Time = 0.58 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.16

method	result
derivativedivides	$\frac{(bx+a) \operatorname{arcsech}(bx+a) - \operatorname{arctan}\left(\sqrt{\frac{1}{bx+a}-1} \sqrt{\frac{1}{bx+a}+1}\right)}{b}$
default	$\frac{(bx+a) \operatorname{arcsech}(bx+a) - \operatorname{arctan}\left(\sqrt{\frac{1}{bx+a}-1} \sqrt{\frac{1}{bx+a}+1}\right)}{b}$
parts	$x \operatorname{arcsech}(bx+a) + \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \left(\operatorname{csgn}(b) \operatorname{arctanh}\left(\frac{1}{\sqrt{-b^2x^2-2bxa-a^2+1}}\right) a + \operatorname{arctan}\left(\frac{1}{\sqrt{-b^2x^2-2bxa-a^2+1}}\right)\right)}{b\sqrt{-b^2x^2-2bxa-a^2+1}}$

input `int(arcsech(b*x+a),x,method=_RETURNVERBOSE)`

output `1/b*((b*x+a)*arcsech(b*x+a)-arctan((1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 253 vs.  $2(36) = 72$ .

Time = 0.15 (sec) , antiderivative size = 253, normalized size of antiderivative = 6.66

$$\int \operatorname{sech}^{-1}(a + bx) dx$$

$$= \frac{2bx \log\left(\frac{(bx+a)\sqrt{\frac{-b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}+1}{bx+a}\right) + a \log\left(\frac{(bx+a)\sqrt{\frac{-b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}}{x}\right) - a \log\left(\frac{(bx+a)\sqrt{\frac{-b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}}{x}\right)}{2b}$$

input `integrate(arcsech(b*x+a),x, algorithm="fricas")`

output `1/2*(2*b*x*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)) + a*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/x) - a*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) - 1)/x) - 2*arctan((b^2*x^2 + 2*a*b*x + a^2)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)))/(b^2*x^2 + 2*a*b*x + a^2 - 1))/b`

### Sympy [F]

$$\int \operatorname{sech}^{-1}(a + bx) dx = \int \operatorname{asech}(a + bx) dx$$

input `integrate(asech(b*x+a),x)`

output `Integral(asech(a + b*x), x)`

### Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int \operatorname{sech}^{-1}(a + bx) dx = \frac{(bx + a) \operatorname{ar} \operatorname{sech}(bx + a) - \arctan\left(\sqrt{\frac{1}{(bx+a)^2} - 1}\right)}{b}$$

input `integrate(arcsech(b*x+a),x, algorithm="maxima")`

output `((b*x + a)*arcsech(b*x + a) - arctan(sqrt(1/(b*x + a)^2 - 1)))/b`

### Giac [F]

$$\int \operatorname{sech}^{-1}(a + bx) dx = \int \operatorname{ar} \operatorname{sech}(bx + a) dx$$

input `integrate(arcsech(b*x+a),x, algorithm="giac")`

output `integrate(arcsech(b*x + a), x)`

### Mupad [B] (verification not implemented)

Time = 4.33 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.13

$$\int \operatorname{sech}^{-1}(a + bx) dx = \frac{\operatorname{atan}\left(\frac{1}{\sqrt{\frac{1}{a+bx}-1}\sqrt{\frac{1}{a+bx}+1}}\right) + \operatorname{acosh}\left(\frac{1}{a+bx}\right)(a + bx)}{b}$$

input `int(acosh(1/(a + b*x)),x)`

output

```
(atan(1/((1/(a + b*x) - 1)^(1/2)*(1/(a + b*x) + 1)^(1/2))) + acosh(1/(a + b*x))*(a + b*x))/b
```

**Reduce [F]**

$$\int \operatorname{sech}^{-1}(a + bx) dx = \int \operatorname{asech}(bx + a) dx$$

input

```
int(asech(b*x+a), x)
```

output

```
int(asech(a + b*x), x)
```

### 3.5 $\int \frac{\operatorname{sech}^{-1}(a+bx)}{x} dx$

Optimal result	68
Mathematica [C] (warning: unable to verify)	69
Rubi [C] (verified)	70
Maple [C] (verified)	75
Fricas [F]	76
Sympy [F]	76
Maxima [F]	76
Giac [F]	77
Mupad [F(-1)]	77
Reduce [F]	77

#### Optimal result

Integrand size = 10, antiderivative size = 170

$$\begin{aligned} \int \frac{\operatorname{sech}^{-1}(a+bx)}{x} dx &= \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\ &\quad + \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\ &\quad - \operatorname{sech}^{-1}(a+bx) \log\left(1 + e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\ &\quad + \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) + \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\ &\quad - \frac{1}{2} \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) \end{aligned}$$

output

```
arcsech(b*x+a)*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/
(1-(-a^2+1)^(1/2)))+arcsech(b*x+a)*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*
(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))-arcsech(b*x+a)*ln(1+(1/(b*x+a)+(1/
(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)+polylog(2,a*(1/(b*x+a)+(1/(b*x+a)
-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))+polylog(2,a*(1/(b*x+a)+
(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))-1/2*polylog(2
,-(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)
```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 332, normalized size of antiderivative = 1.95

$$\begin{aligned}
& \int \frac{\operatorname{sech}^{-1}(a+bx)}{x} dx \\
&= -4i \arcsin\left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}}\right) \operatorname{arctanh}\left(\frac{(1+a) \tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a^2}}\right) \\
&\quad - \operatorname{sech}^{-1}(a+bx) \log\left(1 + e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\
&\quad + \operatorname{sech}^{-1}(a+bx) \log\left(1 + \frac{(-1 + \sqrt{1-a^2}) e^{-\operatorname{sech}^{-1}(a+bx)}}{a}\right) \\
&\quad + 2i \arcsin\left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}}\right) \log\left(1 + \frac{(-1 + \sqrt{1-a^2}) e^{-\operatorname{sech}^{-1}(a+bx)}}{a}\right) \\
&\quad + \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{(1 + \sqrt{1-a^2}) e^{-\operatorname{sech}^{-1}(a+bx)}}{a}\right) \\
&\quad - 2i \arcsin\left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}}\right) \log\left(1 - \frac{(1 + \sqrt{1-a^2}) e^{-\operatorname{sech}^{-1}(a+bx)}}{a}\right) \\
&\quad + \frac{1}{2} \operatorname{PolyLog}\left(2, -e^{-2\operatorname{sech}^{-1}(a+bx)}\right) - \operatorname{PolyLog}\left(2, -\frac{(-1 + \sqrt{1-a^2}) e^{-\operatorname{sech}^{-1}(a+bx)}}{a}\right) \\
&\quad - \operatorname{PolyLog}\left(2, \frac{(1 + \sqrt{1-a^2}) e^{-\operatorname{sech}^{-1}(a+bx)}}{a}\right)
\end{aligned}$$

input

```
Integrate[ArcSech[a + b*x]/x, x]
```

output

```
(-4*I)*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*ArcTanh[((1 + a)*Tanh[ArcSech[a +
b*x]/2])/Sqrt[1 - a^2]] - ArcSech[a + b*x]*Log[1 + E^(-2*ArcSech[a + b*x])
] + ArcSech[a + b*x]*Log[1 + (-1 + Sqrt[1 - a^2])/(a*E^ArcSech[a + b*x])]
+ (2*I)*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*Log[1 + (-1 + Sqrt[1 - a^2])/(a*E
^ArcSech[a + b*x])] + ArcSech[a + b*x]*Log[1 - (1 + Sqrt[1 - a^2])/(a*E^Ar
cSech[a + b*x])] - (2*I)*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*Log[1 - (1 + Sqr
t[1 - a^2])/(a*E^ArcSech[a + b*x])] + PolyLog[2, -E^(-2*ArcSech[a + b*x])]
/2 - PolyLog[2, -((-1 + Sqrt[1 - a^2])/(a*E^ArcSech[a + b*x]))] - PolyLog[
2, (1 + Sqrt[1 - a^2])/(a*E^ArcSech[a + b*x])]
```

### Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.06 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.37, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$ , Rules used = {6875, 25, 6129, 6104, 25, 3042, 26, 4201, 2620, 2715, 2838, 6096, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^{-1}(a+bx)}{x} dx \\
 & \quad \downarrow 6875 \\
 & - \int \frac{(a+bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1) \operatorname{sech}^{-1}(a+bx)}{bx} d\operatorname{sech}^{-1}(a+bx) \\
 & \quad \downarrow 25 \\
 & \int - \frac{(a+bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1) \operatorname{sech}^{-1}(a+bx)}{bx} d\operatorname{sech}^{-1}(a+bx) \\
 & \quad \downarrow 6129 \\
 & \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1) \operatorname{sech}^{-1}(a+bx)}{\frac{a}{a+bx} - 1} d\operatorname{sech}^{-1}(a+bx) \\
 & \quad \downarrow 6104
 \end{aligned}$$

$$\begin{aligned}
& a \int -\frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)}{(a+bx)\left(1-\frac{a}{a+bx}\right)}d\operatorname{sech}^{-1}(a+bx) - \int \sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow 25 \\
& - \int \sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)d\operatorname{sech}^{-1}(a+bx) - \\
& \quad a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)}{(a+bx)\left(1-\frac{a}{a+bx}\right)}d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow 3042 \\
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)}{(a+bx)\left(1-\frac{a}{a+bx}\right)}d\operatorname{sech}^{-1}(a+bx) - \int -i\operatorname{sech}^{-1}(a+bx)\tan(i\operatorname{sech}^{-1}(a+bx))d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow 26 \\
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)}{(a+bx)\left(1-\frac{a}{a+bx}\right)}d\operatorname{sech}^{-1}(a+bx) + i \int \operatorname{sech}^{-1}(a+bx)\tan(i\operatorname{sech}^{-1}(a+bx))d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow 4201 \\
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)}{(a+bx)\left(1-\frac{a}{a+bx}\right)}d\operatorname{sech}^{-1}(a+bx) + \\
& \quad i \left( 2i \int \frac{e^{2\operatorname{sech}^{-1}(a+bx)}\operatorname{sech}^{-1}(a+bx)}{1+e^{2\operatorname{sech}^{-1}(a+bx)}}d\operatorname{sech}^{-1}(a+bx) - \frac{1}{2}i\operatorname{sech}^{-1}(a+bx)^2 \right) \\
& \quad \downarrow 2620 \\
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)}{(a+bx)\left(1-\frac{a}{a+bx}\right)}d\operatorname{sech}^{-1}(a+bx) + \\
& \quad i \left( 2i \left( \frac{1}{2}\operatorname{sech}^{-1}(a+bx)\log\left(e^{2\operatorname{sech}^{-1}(a+bx)}+1\right) - \frac{1}{2}\int\log\left(1+e^{2\operatorname{sech}^{-1}(a+bx)}\right)d\operatorname{sech}^{-1}(a+bx) \right) - \frac{1}{2}i\operatorname{sech}^{-1}(a+bx)^2 \right) \\
& \quad \downarrow 2715
\end{aligned}$$



$$\begin{aligned}
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)}{(a+bx)\left(1-\frac{a}{a+bx}\right)} d\operatorname{sech}^{-1}(a+bx) + \\
& i\left(2i\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\log\left(e^{2\operatorname{sech}^{-1}(a+bx)}+1\right)-\frac{1}{4}\int e^{-2\operatorname{sech}^{-1}(a+bx)}\log\left(1+e^{2\operatorname{sech}^{-1}(a+bx)}\right)de^{2\operatorname{sech}^{-1}(a+bx)}\right)\right) -
\end{aligned}$$

↓ 2838

$$\begin{aligned}
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)}{(a+bx)\left(1-\frac{a}{a+bx}\right)} d\operatorname{sech}^{-1}(a+bx) + \\
& i\left(2i\left(\frac{1}{4}\operatorname{PolyLog}\left(2,-e^{2\operatorname{sech}^{-1}(a+bx)}\right)+\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\log\left(e^{2\operatorname{sech}^{-1}(a+bx)}+1\right)\right)-\frac{1}{2}i\operatorname{sech}^{-1}(a+bx)^2\right)
\end{aligned}$$

↓ 6096

$$\begin{aligned}
& -a\left(\int \frac{e^{\operatorname{sech}^{-1}(a+bx)}\operatorname{sech}^{-1}(a+bx)}{-e^{\operatorname{sech}^{-1}(a+bx)}a-\sqrt{1-a^2}+1}d\operatorname{sech}^{-1}(a+bx)+\int \frac{e^{\operatorname{sech}^{-1}(a+bx)}\operatorname{sech}^{-1}(a+bx)}{-e^{\operatorname{sech}^{-1}(a+bx)}a+\sqrt{1-a^2}+1}d\operatorname{sech}^{-1}(a+bx)+\dots\right) \\
& i\left(2i\left(\frac{1}{4}\operatorname{PolyLog}\left(2,-e^{2\operatorname{sech}^{-1}(a+bx)}\right)+\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\log\left(e^{2\operatorname{sech}^{-1}(a+bx)}+1\right)\right)-\frac{1}{2}i\operatorname{sech}^{-1}(a+bx)^2\right)
\end{aligned}$$

↓ 2620

$$\begin{aligned}
& -a\left(\frac{\int \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)d\operatorname{sech}^{-1}(a+bx)}{a}+\frac{\int \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)d\operatorname{sech}^{-1}(a+bx)}{a}-\frac{\operatorname{sech}^{-1}(a+bx)}{a}\right) \\
& i\left(2i\left(\frac{1}{4}\operatorname{PolyLog}\left(2,-e^{2\operatorname{sech}^{-1}(a+bx)}\right)+\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\log\left(e^{2\operatorname{sech}^{-1}(a+bx)}+1\right)\right)-\frac{1}{2}i\operatorname{sech}^{-1}(a+bx)^2\right)
\end{aligned}$$

↓ 2715

$$\begin{aligned}
& -a\left(\frac{\int e^{-\operatorname{sech}^{-1}(a+bx)}\log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)de^{\operatorname{sech}^{-1}(a+bx)}}{a}+\frac{\int e^{-\operatorname{sech}^{-1}(a+bx)}\log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)de^{\operatorname{sech}^{-1}(a+bx)}}{a}\right) \\
& i\left(2i\left(\frac{1}{4}\operatorname{PolyLog}\left(2,-e^{2\operatorname{sech}^{-1}(a+bx)}\right)+\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\log\left(e^{2\operatorname{sech}^{-1}(a+bx)}+1\right)\right)-\frac{1}{2}i\operatorname{sech}^{-1}(a+bx)^2\right)
\end{aligned}$$

↓ 2838

$$-a \left( \frac{\text{PolyLog} \left( 2, \frac{ae^{\text{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a} - \frac{\text{PolyLog} \left( 2, \frac{ae^{\text{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1} \right)}{a} - \frac{\text{sech}^{-1}(a+bx) \log \left( 1 - \frac{ae^{\text{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a} \right) - i \left( 2i \left( \frac{1}{4} \text{PolyLog} \left( 2, -e^{2\text{sech}^{-1}(a+bx)} \right) + \frac{1}{2} \text{sech}^{-1}(a+bx) \log \left( e^{2\text{sech}^{-1}(a+bx)} + 1 \right) \right) - \frac{1}{2} i \text{sech}^{-1}(a+bx)^2 \right)$$

input `Int[ArcSech[a + b*x]/x, x]`

output `-(a*(ArcSech[a + b*x]^2/(2*a) - (ArcSech[a + b*x]*Log[1 - (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2]])]/a - (ArcSech[a + b*x]*Log[1 - (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2]])]/a - PolyLog[2, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2]])/a - PolyLog[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2]])/a)) + I*((-1/2*I)*ArcSech[a + b*x]^2 + (2*I)*((ArcSech[a + b*x]*Log[1 + E^(2*ArcSech[a + b*x])])/2 + PolyLog[2, -E^(2*ArcSech[a + b*x])]/4))`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838  $\text{Int}[\text{Log}[(c\_.) * ((d\_.) + (e\_.) * (x\_.)^{\wedge}(n\_.))] / (x\_.), x\_Symbol] \text{:>} \text{Simp}[-\text{PolyLog}[2, (-c) * e * x^{\wedge}n] / n, x] \text{/; FreeQ}\{c, d, e, n\}, x\} \&\& \text{EqQ}\{c * d, 1\}$

rule 3042  $\text{Int}[u\_., x\_Symbol] \text{:>} \text{Int}[\text{DeactivateTrig}[u, x], x] \text{/; FunctionOfTrigOfLinearQ}[u, x]$

rule 4201  $\text{Int}[((c\_.) + (d\_.) * (x\_.)^{\wedge}(m\_.) * \tan[(e\_.) + (\text{Complex}[0, fz\_]) * (f\_.) * (x\_.)]), x\_Symbol] \text{:>} \text{Simp}[(-I) * ((c + d * x)^{\wedge}(m + 1) / (d * (m + 1))), x] + \text{Simp}[2 * I \text{ Int}[(c + d * x)^{\wedge}m * (E^{\wedge}(2 * ((-I) * e + f * fz * x)) / (1 + E^{\wedge}(2 * ((-I) * e + f * fz * x))))], x], x] \text{/; FreeQ}\{c, d, e, f, fz\}, x\} \&\& \text{IGtQ}[m, 0]$

rule 6096  $\text{Int}[(((e\_.) + (f\_.) * (x\_.)^{\wedge}(m\_.) * \text{Sinh}[(c\_.) + (d\_.) * (x\_.)]) / (\text{Cosh}[(c\_.) + (d\_.) * (x\_.)] * (b\_.) + (a\_))), x\_Symbol] \text{:>} \text{Simp}[-(e + f * x)^{\wedge}(m + 1) / (b * f * (m + 1)), x] + (\text{Int}[(e + f * x)^{\wedge}m * (E^{\wedge}(c + d * x) / (a - \text{Rt}[a^{\wedge}2 - b^{\wedge}2, 2] + b * E^{\wedge}(c + d * x)))] , x] + \text{Int}[(e + f * x)^{\wedge}m * (E^{\wedge}(c + d * x) / (a + \text{Rt}[a^{\wedge}2 - b^{\wedge}2, 2] + b * E^{\wedge}(c + d * x)))] , x]) \text{/; FreeQ}\{a, b, c, d, e, f\}, x\} \&\& \text{IGtQ}[m, 0] \&\& \text{NeQ}[a^{\wedge}2 - b^{\wedge}2, 0]$

rule 6104  $\text{Int}[(((e\_.) + (f\_.) * (x\_.)^{\wedge}(m\_.) * \text{Tanh}[(c\_.) + (d\_.) * (x\_.)]^{\wedge}(n\_.) / (\text{Cosh}[(c\_.) + (d\_.) * (x\_.)] * (b\_.) + (a\_))), x\_Symbol] \text{:>} \text{Simp}[1/a \text{ Int}[(e + f * x)^{\wedge}m * \text{Tanh}[c + d * x]^{\wedge}n, x], x] - \text{Simp}[b/a \text{ Int}[(e + f * x)^{\wedge}m * \text{Sinh}[c + d * x] * (\text{Tanh}[c + d * x]^{\wedge}(n - 1) / (a + b * \text{Cosh}[c + d * x]))], x], x] \text{/; FreeQ}\{a, b, c, d, e, f\}, x\} \&\& \text{IGtQ}[m, 0] \&\& \text{IGtQ}[n, 0]$

rule 6129  $\text{Int}[(((e\_.) + (f\_.) * (x\_.)^{\wedge}(m\_.) * (F\_.)[(c\_.) + (d\_.) * (x\_.)]^{\wedge}(n\_.) * (G\_.)[(c\_.) + (d\_.) * (x\_.)]^{\wedge}(p\_.) / ((a\_.) + (b\_.) * \text{Sech}[(c\_.) + (d\_.) * (x\_.)])), x\_Symbol] \text{:>} \text{Int}[(e + f * x)^{\wedge}m * \text{Cosh}[c + d * x] * F[c + d * x]^{\wedge}n * (G[c + d * x]^{\wedge}p / (b + a * \text{Cosh}[c + d * x])), x] \text{/; FreeQ}\{a, b, c, d, e, f\}, x\} \&\& \text{HyperbolicQ}[F] \&\& \text{HyperbolicQ}[G] \&\& \text{IntegersQ}[m, n, p]$

rule 6875  $\text{Int}[((a\_.) + \text{ArcSech}[(c\_.) + (d\_.) * (x\_.)] * (b\_.)^{\wedge}(p\_.) * ((e\_.) + (f\_.) * (x\_.)^{\wedge}(m\_.)], x\_Symbol] \text{:>} \text{Simp}[-(d^{\wedge}(m + 1))^{\wedge}(-1) \text{ Subst}[\text{Int}[(a + b * x)^{\wedge}p * \text{Sech}[x] * \text{Tanh}[x] * (d * e - c * f + f * \text{Sech}[x])^{\wedge}m, x], x, \text{ArcSech}[c + d * x]], x] \text{/; FreeQ}\{a, b, c, d, e, f\}, x\} \&\& \text{IGtQ}[p, 0] \&\& \text{IntegerQ}[m]$

**Maple [C] (verified)**

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 886, normalized size of antiderivative = 5.21

method	result
derivativedivides	$-\operatorname{arcsech}(bx+a) \ln\left(1+i\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)-\operatorname{arcsech}(bx+a) \ln\left(1-i\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)$
default	$-\operatorname{arcsech}(bx+a) \ln\left(1+i\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)-\operatorname{arcsech}(bx+a) \ln\left(1-i\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)$

input `int(arcsech(b*x+a)/x,x,method=_RETURNVERBOSE)`

output

```
-arcsech(b*x+a)*ln(1+I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))
)-arcsech(b*x+a)*ln(1-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)
))-dilog(1+I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))-dilog(1-
I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))+1/2*arcsech(b*x+a)*
ln((-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)+
1)/(1+(-a^2+1)^(1/2)))+1/2*arcsech(b*x+a)*ln((a*(1/(b*x+a)+(1/(b*x+a)-1)^(
1/2)*(1/(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2)))-1/2*(-a^2
+1)^(1/2)/(a^2-1)*arcsech(b*x+a)*ln((-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/
(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))+1/2*(-a^2+1)^(1/2)
/(a^2-1)*arcsech(b*x+a)*ln((a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)
^(1/2))+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2)))+dilog((a*(1/(b*x+a)+(1/(b*x
+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2)))+d
ilog((-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)
)+1)/(1+(-a^2+1)^(1/2)))+1/2*(a^2-1-(-a^2+1)^(1/2))/a^2/(a^2-1)*arcsech(b*
x+a)*(ln((a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))+(-a^2+1)^(
1/2)-1)/(-1+(-a^2+1)^(1/2)))*a^2+ln((-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/
(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))*a^2-2*ln((-a*(1/(b
*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)+1)/(1+(-a^2+
1)^(1/2)))+2*ln((-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))+(-
a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))*(-a^2+1)^(1/2))
```

**Fricas [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arsech}(bx + a)}{x} dx$$

input `integrate(arcsech(b*x+a)/x,x, algorithm="fricas")`

output `integral(arcsech(b*x + a)/x, x)`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{asech}(a + bx)}{x} dx$$

input `integrate(asech(b*x+a)/x,x)`

output `Integral(asech(a + b*x)/x, x)`

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arsech}(bx + a)}{x} dx$$

input `integrate(arcsech(b*x+a)/x,x, algorithm="maxima")`

output `integrate(arcsech(b*x + a)/x, x)`

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arsech}(bx + a)}{x} dx$$

input `integrate(arcsech(b*x+a)/x,x, algorithm="giac")`

output `integrate(arcsech(b*x + a)/x, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)}{x} dx$$

input `int(acosh(1/(a + b*x)))/x,x)`

output `int(acosh(1/(a + b*x)))/x, x)`

**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{asech}(bx + a)}{x} dx$$

input `int(asech(b*x+a)/x,x)`

output `int(asech(a + b*x)/x,x)`

### 3.6 $\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^2} dx$

Optimal result	78
Mathematica [B] (verified)	78
Rubi [A] (verified)	79
Maple [B] (verified)	81
Fricas [B] (verification not implemented)	82
Sympy [F]	83
Maxima [F]	83
Giac [F]	84
Mupad [F(-1)]	84
Reduce [F]	85

#### Optimal result

Integrand size = 10, antiderivative size = 70

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^2} dx = -\frac{b\operatorname{sech}^{-1}(a+bx)}{a} - \frac{\operatorname{sech}^{-1}(a+bx)}{x} + \frac{2b\operatorname{arctanh}\left(\frac{\sqrt{1+a}\tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{a\sqrt{1-a^2}}$$

output

```
-b*arcsech(b*x+a)/a-arcsech(b*x+a)/x+2*b*arctanh((1+a)^(1/2)*tanh(1/2*arcs
ech(b*x+a))/(1-a)^(1/2))/a/(-a^2+1)^(1/2)
```

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 244 vs. 2(70) = 140.

Time = 0.17 (sec) , antiderivative size = 244, normalized size of antiderivative = 3.49

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^2} dx = -\frac{\operatorname{sech}^{-1}(a+bx)}{x} + \frac{b\left(-\log(x) + \sqrt{1-a^2}\log(a+bx) - \sqrt{1-a^2}\log\left(1 + \sqrt{-\frac{-1+a+bx}{1+a+bx}} + a\sqrt{-\frac{-1+a+bx}{1+a+bx}} + bx\sqrt{-\frac{-1+a+bx}{1+a+bx}}\right)\right)}{a\sqrt{1-a^2}}$$

input `Integrate[ArcSech[a + b*x]/x^2,x]`

output 
$$\begin{aligned} &-(\text{ArcSech}[a + b*x]/x) + (b*(-\text{Log}[x] + \text{Sqrt}[1 - a^2]*\text{Log}[a + b*x] - \text{Sqrt}[1 \\ &- a^2]*\text{Log}[1 + \text{Sqrt}[-( (-1 + a + b*x)/(1 + a + b*x))]] + a*\text{Sqrt}[-( (-1 + a + \\ &b*x)/(1 + a + b*x))]] + b*x*\text{Sqrt}[-( (-1 + a + b*x)/(1 + a + b*x))]] + \text{Log}[1 \\ &- a^2 - a*b*x + \text{Sqrt}[1 - a^2]*\text{Sqrt}[-( (-1 + a + b*x)/(1 + a + b*x))]] + a*\text{Sqrt}[1 - a^2]*\text{Sqrt}[-( (-1 + a + b*x)/(1 + a + b*x))]] + \text{Sqrt}[1 - a^2]*b*x*\text{Sqrt} \\ &[-( (-1 + a + b*x)/(1 + a + b*x))]))/(a*\text{Sqrt}[1 - a^2]) \end{aligned}$$

### Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.03, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$ , Rules used = {6875, 5991, 3042, 4270, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} &\int \frac{\text{sech}^{-1}(a + bx)}{x^2} dx \\ &\quad \downarrow \text{6875} \\ &-b \int \frac{(a + bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a + bx + 1) \text{sech}^{-1}(a + bx)}{b^2 x^2} d\text{sech}^{-1}(a + bx) \\ &\quad \downarrow \text{5991} \\ &-b \left( \int -\frac{1}{bx} d\text{sech}^{-1}(a + bx) + \frac{\text{sech}^{-1}(a + bx)}{bx} \right) \\ &\quad \downarrow \text{3042} \\ &-b \left( \frac{\text{sech}^{-1}(a + bx)}{bx} + \int \frac{1}{a - \csc(i \text{sech}^{-1}(a + bx) + \frac{\pi}{2})} d\text{sech}^{-1}(a + bx) \right) \\ &\quad \downarrow \text{4270} \\ &-b \left( -\frac{\int \frac{1}{1 - \frac{a}{a+bx}} d\text{sech}^{-1}(a + bx)}{a} + \frac{\text{sech}^{-1}(a + bx)}{a} + \frac{\text{sech}^{-1}(a + bx)}{bx} \right) \end{aligned}$$



$$\begin{aligned}
& \downarrow 3042 \\
& -b \left( -\frac{\int \frac{1}{1-a \sin\left(i \operatorname{sech}^{-1}(a+bx) + \frac{\pi}{2}\right)} d \operatorname{sech}^{-1}(a+bx)}{a} + \frac{\operatorname{sech}^{-1}(a+bx)}{a} + \frac{\operatorname{sech}^{-1}(a+bx)}{bx} \right) \\
& \downarrow 3138 \\
& -b \left( -\frac{2 \int \frac{1}{-\left((a+1) \tanh^2\left(\frac{1}{2} \operatorname{sech}^{-1}(a+bx)\right)\right) - a + 1} d \tanh\left(\frac{1}{2} \operatorname{sech}^{-1}(a+bx)\right)}{a} + \frac{\operatorname{sech}^{-1}(a+bx)}{a} + \frac{\operatorname{sech}^{-1}(a+bx)}{bx} \right) \\
& \downarrow 221 \\
& -b \left( -\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{a+1} \tanh\left(\frac{1}{2} \operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{a \sqrt{1-a^2}} + \frac{\operatorname{sech}^{-1}(a+bx)}{a} + \frac{\operatorname{sech}^{-1}(a+bx)}{bx} \right)
\end{aligned}$$

input `Int[ArcSech[a + b*x]/x^2,x]`

output `-(b*(ArcSech[a + b*x]/a + ArcSech[a + b*x]/(b*x) - (2*ArcTanh[(Sqrt[1 + a]*Tanh[ArcSech[a + b*x]/2])/Sqrt[1 - a]])/(a*Sqrt[1 - a^2])))`

### Defintions of rubi rules used

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 4270 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))(-1), x_Symbol] := Simp[x/a, x] - Simp[1/a Int[1/(1 + (a/b)*Sin[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a2 - b2, 0]`

rule 5991 `Int[((e_.) + (f_.)*(x_))(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sech[(c_.) + (d_.)*(x_)])(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-(e + f*x)m)*((a + b*Sech[c + d*x])(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)(m - 1)*(a + b*Sech[c + d*x])(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

rule 6875 `Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)])*(b_.))(p_.)*((e_.) + (f_.)*(x_))(m_.), x_Symbol] := Simp[-(d(m + 1))(-1) Subst[Int[(a + b*x)p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

## Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 170 vs.  $2(62) = 124$ .

Time = 0.92 (sec) , antiderivative size = 171, normalized size of antiderivative = 2.44

method	result
derivativedivides	$b \left( -\frac{\operatorname{arcsech}(bx+a)}{bx} - \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \left( \operatorname{arctanh}\left(\frac{1}{\sqrt{1-(bx+a)^2}}\right) a^2 + \sqrt{-a^2+1} \ln\left(\frac{2\sqrt{-a^2+1}\sqrt{1-(bx+a)^2}}{\sqrt{1-(bx+a)^2} a(a-1)(1+a)}\right) \right)}{\sqrt{1-(bx+a)^2} a(a-1)(1+a)} \right)$
default	$b \left( -\frac{\operatorname{arcsech}(bx+a)}{bx} - \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \left( \operatorname{arctanh}\left(\frac{1}{\sqrt{1-(bx+a)^2}}\right) a^2 + \sqrt{-a^2+1} \ln\left(\frac{2\sqrt{-a^2+1}\sqrt{1-(bx+a)^2}}{\sqrt{1-(bx+a)^2} a(a-1)(1+a)}\right) \right)}{\sqrt{1-(bx+a)^2} a(a-1)(1+a)} \right)$
parts	$-\frac{\operatorname{arcsech}(bx+a)}{x} - \frac{b\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \left( \operatorname{arctanh}\left(\frac{1}{\sqrt{-b^2x^2-2bxa-a^2+1}}\right) a^2 + \sqrt{-a^2+1} \ln\left(\frac{-2a^2+2-2bxa}{\sqrt{-b^2x^2-2bxa-a^2+1}(1-2bxa-a^2)}\right) \right)}{\sqrt{-b^2x^2-2bxa-a^2+1}(1-2bxa-a^2)}$

input `int(arcsech(b*x+a)/x^2,x,method=_RETURNVERBOSE)`

output

```
b*(-1/b/x*arcsech(b*x+a)-(-(b*x+a-1)/(b*x+a))^(1/2)*(b*x+a)*((b*x+a+1)/(b*x+a))^(1/2)*(arctanh(1/(1-(b*x+a)^2)^(1/2))*a^2+(-a^2+1)^(1/2)*ln(2*((-a^2+1)^(1/2)*(1-(b*x+a)^2)^(1/2)-(b*x+a)*a+1)/b/x)-arctanh(1/(1-(b*x+a)^2)^(1/2))))/(1-(b*x+a)^2)^(1/2)/a/(a-1)/(1+a)
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 320 vs.  $2(62) = 124$ .

Time = 0.14 (sec) , antiderivative size = 651, normalized size of antiderivative = 9.30

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^2} dx$$

$$= \frac{(a^2 - 1)bx \log\left(\frac{(bx+a)\sqrt{\frac{-b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}+1}{x}\right) - (a^2 - 1)bx \log\left(\frac{(bx+a)\sqrt{\frac{-b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}-1}{x}\right) + \sqrt{-a^2 + 1}}{2(a^3 - 1)}$$

input

```
integrate(arcsech(b*x+a)/x^2,x, algorithm="fricas")
```

output

```
[-1/2*((a^2 - 1)*b*x*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/x) - (a^2 - 1)*b*x*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) - 1)/x) + sqrt(-a^2 + 1)*b*x*log(((2*a^2 - 1)*b^2*x^2 + 2*a^4 + 4*(a^3 - a)*b*x - 4*a^2 - 2*(a*b^2*x^2 + a^3 + (2*a^2 - 1)*b*x - a)*sqrt(-a^2 + 1)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 2)/x^2) + 2*(a^3 - a)*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)))/((a^3 - a)*x), -1/2*((a^2 - 1)*b*x*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/x) - (a^2 - 1)*b*x*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) - 1)/x) + 2*sqrt(a^2 - 1)*b*x*arctan((a*b^2*x^2 + a^3 + (2*a^2 - 1)*b*x - a)*sqrt(a^2 - 1)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)))/((a^2 - 1)*b^2*x^2 + a^4 + 2*(a^3 - a)*b*x - 2*a^2 + 1)) + 2*(a^3 - a)*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)))/((a^3 - a)*x)]
```

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{asech}(a + bx)}{x^2} dx$$

input

```
integrate(asech(b*x+a)/x**2,x)
```

output

```
Integral(asech(a + b*x)/x**2, x)
```

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{arsech}(bx + a)}{x^2} dx$$

input

```
integrate(arcsech(b*x+a)/x^2,x, algorithm="maxima")
```

output

```
b*log(x)/(a^3 - a) - 1/2*((a^2*b - a*b)*x*log(b*x + a + 1) + (a^2*b + a*b)
*x*log(-b*x - a + 1) + 2*(a^3 - a)*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)
)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a) - 2*(a^3 + (a^2*
b - b)*x - a)*log(b*x + a) - 2*(a^3 - a)*log(b*x + a))/((a^3 - a)*x) - int
egrate((b^2*x + a*b)/(b^2*x^3 + 2*a*b*x^2 + (a^2 - 1)*x + (b^2*x^3 + 2*a*b
*x^2 + (a^2 - 1)*x)*e^(1/2*log(b*x + a + 1) + 1/2*log(-b*x - a + 1))), x)
```

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{ar} \operatorname{sech}(bx + a)}{x^2} dx$$

input

```
integrate(arcsech(b*x+a)/x^2,x, algorithm="giac")
```

output

```
integrate(arcsech(b*x + a)/x^2, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)}{x^2} dx$$

input

```
int(acosh(1/(a + b*x))/x^2,x)
```

output

```
int(acosh(1/(a + b*x))/x^2, x)
```

**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{asech}(bx + a)}{x^2} dx$$

input `int(asech(b*x+a)/x^2,x)`

output `int(asech(a + b*x)/x**2,x)`

### 3.7 $\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^3} dx$

Optimal result	86
Mathematica [B] (verified)	86
Rubi [A] (verified)	87
Maple [C] (verified)	91
Fricas [B] (verification not implemented)	92
Sympy [F]	93
Maxima [F]	94
Giac [F]	94
Mupad [F(-1)]	95
Reduce [F]	95

#### Optimal result

Integrand size = 10, antiderivative size = 133

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^3} dx = \frac{b\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{2a(1-a^2)x} + \frac{b^2\operatorname{sech}^{-1}(a+bx)}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)}{2x^2} - \frac{(1-2a^2)b^2\operatorname{arctanh}\left(\frac{\sqrt{1+a}\tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{a^2(1-a^2)^{3/2}}$$

output

```
1/2*b*((-b*x-a+1)/(b*x+a+1))^(1/2)*(b*x+a+1)/a/(-a^2+1)/x+1/2*b^2*arcsech(b*x+a)/a^2-1/2*arcsech(b*x+a)/x^2-(-2*a^2+1)*b^2*arctanh((1+a)^(1/2)*tanh(1/2*arcsech(b*x+a))/(1-a)^(1/2))/a^2/(-a^2+1)^(3/2)
```

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 315 vs. 2(133) = 266.

Time = 0.55 (sec) , antiderivative size = 315, normalized size of antiderivative = 2.37

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^3} dx$$

$$= \frac{1}{2} \left( -\frac{b\sqrt{-\frac{-1+a+bx}{1+a+bx}}(1+a+bx)}{(-1+a)a(1+a)x} - \frac{\operatorname{sech}^{-1}(a+bx)}{x^2} - \frac{(-1+2a^2)b^2 \log(x)}{a^2(1-a^2)^{3/2}} \right.$$

$$\left. - \frac{b^2 \log(a+bx)}{a^2} + \frac{b^2 \log\left(1 + \sqrt{-\frac{-1+a+bx}{1+a+bx}} + a\sqrt{-\frac{-1+a+bx}{1+a+bx}} + bx\sqrt{-\frac{-1+a+bx}{1+a+bx}}\right)}{a^2} \right.$$

$$\left. + \frac{(-1+2a^2)b^2 \log\left(1 - a^2 - abx + \sqrt{1-a^2}\sqrt{-\frac{-1+a+bx}{1+a+bx}} + a\sqrt{1-a^2}\sqrt{-\frac{-1+a+bx}{1+a+bx}} + \sqrt{1-a^2}bx\sqrt{-\frac{-1+a+bx}{1+a+bx}}\right)}{a^2(1-a^2)^{3/2}} \right)$$

input

```
Integrate[ArcSech[a + b*x]/x^3,x]
```

output

```
(-((b*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(1 + a + b*x))/((-1 + a)*a*(1 + a)*x) - ArcSech[a + b*x]/x^2 - ((-1 + 2*a^2)*b^2*Log[x])/(a^2*(1 - a^2)^(3/2)) - (b^2*Log[a + b*x])/a^2 + (b^2*Log[1 + Sqrt[-((-1 + a + b*x)/(1 + a + b*x))] + a*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))] + b*x*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))])/a^2 + ((-1 + 2*a^2)*b^2*Log[1 - a^2 - a*b*x + Sqrt[1 - a^2]*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))] + a*Sqrt[1 - a^2]*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))] + Sqrt[1 - a^2]*b*x*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))])/a^2*(1 - a^2)^(3/2))/2
```

## Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.20, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.200$ , Rules used = {6875, 25, 5991, 3042, 4272, 3042, 4407, 3042, 4318, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^3} dx$$

↓ 6875



$$\begin{aligned}
& -b^2 \int \frac{(a+bx)\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)}{b^3x^3} d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow 25 \\
& b^2 \int -\frac{(a+bx)\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)}{b^3x^3} d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow 5991 \\
& -b^2 \left( \frac{\operatorname{sech}^{-1}(a+bx)}{2b^2x^2} - \frac{1}{2} \int \frac{1}{b^2x^2} d\operatorname{sech}^{-1}(a+bx) \right) \\
& \quad \downarrow 3042 \\
& -b^2 \left( \frac{\operatorname{sech}^{-1}(a+bx)}{2b^2x^2} - \frac{1}{2} \int \frac{1}{(a - \csc(i\operatorname{sech}^{-1}(a+bx) + \frac{\pi}{2}))^2} d\operatorname{sech}^{-1}(a+bx) \right) \\
& \quad \downarrow 4272 \\
& -b^2 \left( \frac{1}{2} \left( -\frac{\int -\frac{a^2-(a+bx)a+1}{bx} d\operatorname{sech}^{-1}(a+bx)}{a(1-a^2)} - \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a(1-a^2)bx} \right) + \frac{\operatorname{sech}^{-1}(a+bx)}{2b^2x^2} \right) \\
& \quad \downarrow 3042 \\
& -b^2 \left( \frac{\operatorname{sech}^{-1}(a+bx)}{2b^2x^2} + \frac{1}{2} \left( -\frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a(1-a^2)bx} - \frac{\int \frac{-a^2-\csc(i\operatorname{sech}^{-1}(a+bx)+\frac{\pi}{2})^{a+1} d\operatorname{sech}^{-1}(a+bx)}{a-\csc(i\operatorname{sech}^{-1}(a+bx)+\frac{\pi}{2})}}{a(1-a^2)} \right) \right) \\
& \quad \downarrow 4407 \\
& -b^2 \left( \frac{1}{2} \left( -\frac{(1-2a^2)\int -\frac{a+bx}{bx} d\operatorname{sech}^{-1}(a+bx)}{a} + \frac{(1-a^2)\operatorname{sech}^{-1}(a+bx)}{a} - \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a(1-a^2)bx} \right) + \frac{\operatorname{sech}^{-1}(a+bx)}{2b^2x^2} \right) \\
& \quad \downarrow 3042 \\
& -b^2 \left( \frac{\operatorname{sech}^{-1}(a+bx)}{2b^2x^2} + \frac{1}{2} \left( -\frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a(1-a^2)bx} - \frac{(1-a^2)\operatorname{sech}^{-1}(a+bx)}{a} + \frac{(1-2a^2)\int \frac{\csc(i\operatorname{sech}^{-1}(a+bx)+\frac{\pi}{2})}{a-\csc(i\operatorname{sech}^{-1}(a+bx)+\frac{\pi}{2})} d\operatorname{sech}^{-1}(a+bx)}{a(1-a^2)} \right) \right)
\end{aligned}$$

↓ 4318

$$-b^2 \left( \frac{1}{2} \left( -\frac{(1-a^2)\operatorname{sech}^{-1}(a+bx)}{a} - \frac{(1-2a^2) \int \frac{1}{1-\frac{a}{a+bx}} d\operatorname{sech}^{-1}(a+bx)}{a} - \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a(1-a^2)bx} \right) + \frac{\operatorname{sech}^{-1}(a+bx)}{2b^2x^2} \right)$$

↓ 3042

$$-b^2 \left( \frac{\operatorname{sech}^{-1}(a+bx)}{2b^2x^2} + \frac{1}{2} \left( -\frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a(1-a^2)bx} - \frac{(1-a^2)\operatorname{sech}^{-1}(a+bx)}{a} - \frac{(1-2a^2) \int \frac{1}{1-a \sin\left(i\operatorname{sech}^{-1}(a+bx)+\frac{\pi}{2}\right)} d\operatorname{sech}^{-1}(a+bx)}{a} \right) \right)$$

↓ 3138

$$-b^2 \left( \frac{1}{2} \left( -\frac{(1-a^2)\operatorname{sech}^{-1}(a+bx)}{a} - \frac{2(1-2a^2) \int \frac{1}{-\left((a+1)\tanh^2\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)\right)-a+1} d\tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{a} - \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a(1-a^2)} \right) \right)$$

↓ 221

$$-b^2 \left( \frac{1}{2} \left( -\frac{(1-a^2)\operatorname{sech}^{-1}(a+bx)}{a} - \frac{2(1-2a^2)\operatorname{arctanh}\left(\frac{\sqrt{a+1}\tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{a\sqrt{1-a^2}} - \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a(1-a^2)bx} \right) + \frac{\operatorname{sech}^{-1}(a+bx)}{2b^2x^2} \right)$$

input `Int[ArcSech[a + b*x]/x^3,x]`

output `-(b^2*(ArcSech[a + b*x]/(2*b^2*x^2) + (-((Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x))/(a*(1 - a^2)*b*x)) - (((1 - a^2)*ArcSech[a + b*x])/a - (2*(1 - 2*a^2)*ArcTanh[(Sqrt[1 + a]*Tanh[ArcSech[a + b*x]/2])/Sqrt[1 - a]])/(a*Sqrt[1 - a^2]))/(a*(1 - a^2)))/2)`

## Definitions of rubi rules used

- rule 25  $\text{Int}[-(F x_), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F x, x], x]$
- rule 221  $\text{Int}[(a_) + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] \text{ ; FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$
- rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$
- rule 3138  $\text{Int}[(a_) + (b_)*\sin[\text{Pi}/2 + (c_) + (d_)*(x_)]^{-1}, x\_Symbol] \rightarrow \text{With}[\{e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x]\}, \text{Simp}[2*(e/d) \quad \text{Subst}[\text{Int}[1/(a + b + (a - b)*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x]] \text{ ; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$
- rule 4272  $\text{Int}[(\text{csc}[(c_) + (d_)*(x_)]*(b_) + (a_))^{(n)}, x\_Symbol] \rightarrow \text{Simp}[b^2*\text{Cot}[c + d*x]*((a + b*\text{Csc}[c + d*x])^{(n + 1)}/(a*d*(n + 1)*(a^2 - b^2))), x] + \text{Simp}[1/(a*(n + 1)*(a^2 - b^2)) \quad \text{Int}[(a + b*\text{Csc}[c + d*x])^{(n + 1)}*\text{Simp}[(a^2 - b^2)*(n + 1) - a*b*(n + 1)*\text{Csc}[c + d*x] + b^2*(n + 2)*\text{Csc}[c + d*x]^2, x], x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$
- rule 4318  $\text{Int}[\text{csc}[(e_) + (f_)*(x_)]/(\text{csc}[(e_) + (f_)*(x_)]*(b_) + (a_)), x\_Symbol] \rightarrow \text{Simp}[1/b \quad \text{Int}[1/(1 + (a/b)*\text{Sin}[e + f*x]), x], x] \text{ ; FreeQ}[\{a, b, e, f\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$
- rule 4407  $\text{Int}[(\text{csc}[(e_) + (f_)*(x_)]*(d_) + (c_))/(\text{csc}[(e_) + (f_)*(x_)]*(b_) + (a_)), x\_Symbol] \rightarrow \text{Simp}[c*(x/a), x] - \text{Simp}[(b*c - a*d)/a \quad \text{Int}[\text{Csc}[e + f*x]/(a + b*\text{Csc}[e + f*x]), x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0]$

rule 5991

```
Int[((e_.) + (f_.)*(x_.))^(m_.)*Sech[(c_.) + (d_.)*(x_.)]*((a_.) + (b_.)*Sech[
(c_.) + (d_.)*(x_.)]^(n_.)*Tanh[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[(-(e
+ f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b
*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

rule 6875

```
Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(
m_.), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Sech[x]*T
anh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

### Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.96 (sec) , antiderivative size = 370, normalized size of antiderivative = 2.78

method	result
parts	$-\frac{\operatorname{arcsech}(bx+a)}{2x^2} - \frac{b\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \operatorname{csgn}(b)^2 \left(-\operatorname{arctanh}\left(\frac{1}{\sqrt{-b^2x^2-2bxa-a^2+1}}\right)\right) a^4 bx - 2\sqrt{-a^2+1}}{2x^2}$
derivativedivides	$b^2 \left( -\frac{\operatorname{arcsech}(bx+a)}{2b^2x^2} - \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{1-(bx+a)^2}}\right)\right) a^5 - \operatorname{arctanh}\left(\frac{1}{\sqrt{1-(bx+a)^2}}\right) a^4(bx+a)}{2b^2x^2} \right)$
default	$b^2 \left( -\frac{\operatorname{arcsech}(bx+a)}{2b^2x^2} - \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{1-(bx+a)^2}}\right)\right) a^5 - \operatorname{arctanh}\left(\frac{1}{\sqrt{1-(bx+a)^2}}\right) a^4(bx+a)}{2b^2x^2} \right)$

input

```
int(arcsech(b*x+a)/x^3,x,method=_RETURNVERBOSE)
```

output

```
-1/2*arcsech(b*x+a)/x^2-1/2*b*(-(b*x+a-1)/(b*x+a))^(1/2)*(b*x+a)*((b*x+a+1)
)/(b*x+a))^(1/2)*csgn(b)^2*(-arctanh(1/(-b^2*x^2-2*a*b*x-a^2+1)^(1/2))*a^4
*b*x-2*(-a^2+1)^(1/2)*ln(2*(-b*x*a+(-a^2+1)^(1/2)*(-b^2*x^2-2*a*b*x-a^2+1)
^(1/2)-a^2+1)/x)*a^2*b*x+2*arctanh(1/(-b^2*x^2-2*a*b*x-a^2+1)^(1/2))*a^2*b
*x+a^3*(-b^2*x^2-2*a*b*x-a^2+1)^(1/2)+(-a^2+1)^(1/2)*ln(2*(-b*x*a+(-a^2+1)
^(1/2)*(-b^2*x^2-2*a*b*x-a^2+1)^(1/2)-a^2+1)/x)*b*x-arctanh(1/(-b^2*x^2-2*
a*b*x-a^2+1)^(1/2))*b*x-(-b^2*x^2-2*a*b*x-a^2+1)^(1/2)*a)/(-b^2*x^2-2*a*b*
x-a^2+1)^(1/2)/(1+a)/(a-1)/a^2/(a^2-1)/x
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 427 vs.  $2(112) = 224$ .

Time = 0.14 (sec) , antiderivative size = 865, normalized size of antiderivative = 6.50

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^3} dx = \text{Too large to display}$$

input

```
integrate(arcsech(b*x+a)/x^3,x, algorithm="fricas")
```

output

```

[-1/4*((2*a^2 - 1)*sqrt(-a^2 + 1)*b^2*x^2*log(((2*a^2 - 1)*b^2*x^2 + 2*a^4
+ 4*(a^3 - a)*b*x - 4*a^2 + 2*(a*b^2*x^2 + a^3 + (2*a^2 - 1)*b*x - a)*sqrt
(-a^2 + 1)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2))
+ 2)/x^2) - (a^4 - 2*a^2 + 1)*b^2*x^2*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a
*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/x) + (a^4 - 2*a^2 + 1)*b^2
*x^2*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x
+ a^2)) - 1)/x) + 2*(a^6 - 2*a^4 + a^2)*log(((b*x + a)*sqrt(-(b^2*x^2 + 2
*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)) + 2*((a^3 - a
)*b^2*x^2 + (a^4 - a^2)*b*x)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2
+ 2*a*b*x + a^2)))/((a^6 - 2*a^4 + a^2)*x^2), 1/4*(2*(2*a^2 - 1)*sqrt(a^2
- 1)*b^2*x^2*arctan((a*b^2*x^2 + a^3 + (2*a^2 - 1)*b*x - a)*sqrt(a^2 - 1)*
sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)))/((a^2 - 1)*
b^2*x^2 + a^4 + 2*(a^3 - a)*b*x - 2*a^2 + 1)) + (a^4 - 2*a^2 + 1)*b^2*x^2*
log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^
2)) + 1)/x) - (a^4 - 2*a^2 + 1)*b^2*x^2*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*
a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) - 1)/x) - 2*(a^6 - 2*a^4 + a^2
)*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x +
a^2)) + 1)/(b*x + a)) - 2*((a^3 - a)*b^2*x^2 + (a^4 - a^2)*b*x)*sqrt(-(b^2
*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)))/((a^6 - 2*a^4 + a^2)
*x^2)]

```

SymPy [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{asech}(a + bx)}{x^3} dx$$

input

```
integrate(asech(b*x+a)/x**3,x)
```

output

```
Integral(asech(a + b*x)/x**3, x)
```

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{arsech}(bx + a)}{x^3} dx$$

input `integrate(arcsech(b*x+a)/x^3,x, algorithm="maxima")`

output `-1/2*(3*a^2*b^2 - b^2)*log(x)/(a^6 - 2*a^4 + a^2) + 1/4*((a^4*b^2 - 2*a^3*b^2 + a^2*b^2)*x^2*log(b*x + a + 1) + (a^4*b^2 + 2*a^3*b^2 + a^2*b^2)*x^2*log(-b*x - a + 1) - 2*(a^3*b - a*b)*x - 2*(a^6 - 2*a^4 + a^2)*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a) + 2*(a^6 - 2*a^4 - (a^4*b^2 - 2*a^2*b^2 + b^2)*x^2 + a^2)*log(b*x + a) + 2*(a^6 - 2*a^4 + a^2)*log(b*x + a))/((a^6 - 2*a^4 + a^2)*x^2) - integrate(1/2*(b^2*x + a*b)/(b^2*x^4 + 2*a*b*x^3 + (a^2 - 1)*x^2 + (b^2*x^4 + 2*a*b*x^3 + (a^2 - 1)*x^2)*e^(1/2*log(b*x + a + 1) + 1/2*log(-b*x - a + 1))), x)`

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{arsech}(bx + a)}{x^3} dx$$

input `integrate(arcsech(b*x+a)/x^3,x, algorithm="giac")`

output `integrate(arcsech(b*x + a)/x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)}{x^3} dx$$

input `int(acosh(1/(a + b*x))/x^3,x)`output `int(acosh(1/(a + b*x))/x^3, x)`**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{asech}(bx + a)}{x^3} dx$$

input `int(asech(b*x+a)/x^3,x)`output `int(asech(a + b*x)/x**3,x)`



### 3.8 $\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^4} dx$

Optimal result	96
Mathematica [A] (verified)	97
Rubi [A] (verified)	98
Maple [C] (verified)	102
Fricas [B] (verification not implemented)	103
Sympy [F]	104
Maxima [F]	105
Giac [F]	105
Mupad [F(-1)]	106
Reduce [F]	106

#### Optimal result

Integrand size = 10, antiderivative size = 197

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^4} dx = \frac{b\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6a(1-a^2)x^2} - \frac{(2-5a^2)b^2\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6a^2(1-a^2)^2x}$$

$$- \frac{b^3\operatorname{sech}^{-1}(a+bx)}{3a^3} - \frac{\operatorname{sech}^{-1}(a+bx)}{3x^3}$$

$$+ \frac{(2-5a^2+6a^4)b^3\operatorname{arctanh}\left(\frac{\sqrt{1+a}\tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{3a^3(1-a^2)^{5/2}}$$

output

```
1/6*b*((-b*x-a+1)/(b*x+a+1))^(1/2)*(b*x+a+1)/a/(-a^2+1)/x^2-1/6*(-5*a^2+2)
*b^2*((-b*x-a+1)/(b*x+a+1))^(1/2)*(b*x+a+1)/a^2/(-a^2+1)^2/x-1/3*b^3*arcse
ch(b*x+a)/a^3-1/3*arcsech(b*x+a)/x^3+1/3*(6*a^4-5*a^2+2)*b^3*arctanh((1+a)
^(1/2)*tanh(1/2*arcsech(b*x+a))/(1-a)^(1/2))/a^3/(-a^2+1)^(5/2)
```

**Mathematica [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 368, normalized size of antiderivative = 1.87

$$\begin{aligned}
& \int \frac{\operatorname{sech}^{-1}(a + bx)}{x^4} dx \\
&= \frac{1}{6} \left( \frac{b \sqrt{-\frac{-1+a+bx}{1+a+bx}} (a - a^4 - abx - 2bx(1 + bx) + a^3(-1 + 4bx) + a^2(1 + 5bx + 5b^2x^2))}{(-1 + a)^2 a^2 (1 + a)^2 x^2} \right. \\
&\quad - \frac{2 \operatorname{sech}^{-1}(a + bx)}{x^3} - \frac{(2 - 5a^2 + 6a^4) b^3 \log(x)}{a^3 (1 - a^2)^{5/2}} + \frac{2b^3 \log(a + bx)}{a^3} \\
&\quad - \frac{2b^3 \log\left(1 + \sqrt{-\frac{-1+a+bx}{1+a+bx}} + a \sqrt{-\frac{-1+a+bx}{1+a+bx}} + bx \sqrt{-\frac{-1+a+bx}{1+a+bx}}\right)}{a^3} \\
&\quad \left. + \frac{(2 - 5a^2 + 6a^4) b^3 \log\left(1 - a^2 - abx + \sqrt{1 - a^2} \sqrt{-\frac{-1+a+bx}{1+a+bx}} + a \sqrt{1 - a^2} \sqrt{-\frac{-1+a+bx}{1+a+bx}} + \sqrt{1 - a^2} bx \sqrt{-\frac{-1+a+bx}{1+a+bx}}\right)}{a^3 (1 - a^2)^{5/2}} \right)
\end{aligned}$$

input

Integrate[ArcSech[a + b\*x]/x^4,x]

output

```

((b*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(a - a^4 - a*b*x - 2*b*x*(1 + b*
x) + a^3*(-1 + 4*b*x) + a^2*(1 + 5*b*x + 5*b^2*x^2)))/((-1 + a)^2*a^2*(1 +
a)^2*x^2) - (2*ArcSech[a + b*x])/x^3 - ((2 - 5*a^2 + 6*a^4)*b^3*Log[x])/
(a^3*(1 - a^2)^(5/2)) + (2*b^3*Log[a + b*x])/a^3 - (2*b^3*Log[1 + Sqrt[-((-
1 + a + b*x)/(1 + a + b*x))] + a*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))] + b
*x*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))])/a^3 + ((2 - 5*a^2 + 6*a^4)*b^3*
Log[1 - a^2 - a*b*x + Sqrt[1 - a^2]*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]
+ a*Sqrt[1 - a^2]*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))] + Sqrt[1 - a^2]*b*
x*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))])/a^3*(1 - a^2)^(5/2))/6

```

**Rubi [A] (verified)**

Time = 1.07 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.21, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.300$ , Rules used = {6875, 5991, 3042, 4272, 3042, 4548, 3042, 4407, 3042, 4318, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^{-1}(a+bx)}{x^4} dx \\
 & \quad \downarrow \text{6875} \\
 & -b^3 \int \frac{(a+bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1) \operatorname{sech}^{-1}(a+bx)}{b^4 x^4} d\operatorname{sech}^{-1}(a+bx) \\
 & \quad \downarrow \text{5991} \\
 & -b^3 \left( \frac{1}{3} \int -\frac{1}{b^3 x^3} d\operatorname{sech}^{-1}(a+bx) + \frac{\operatorname{sech}^{-1}(a+bx)}{3b^3 x^3} \right) \\
 & \quad \downarrow \text{3042} \\
 & -b^3 \left( \frac{\operatorname{sech}^{-1}(a+bx)}{3b^3 x^3} + \frac{1}{3} \int \frac{1}{(a - \csc(i \operatorname{sech}^{-1}(a+bx) + \frac{\pi}{2}))^3} d\operatorname{sech}^{-1}(a+bx) \right) \\
 & \quad \downarrow \text{4272} \\
 & -b^3 \left( \frac{1}{3} \left( \int \frac{-(a+bx)^2 - 2a(a+bx) + 2(1-a^2)}{b^2 x^2} d\operatorname{sech}^{-1}(a+bx) - \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1)}{2a(1-a^2)b^2 x^2} \right) + \frac{\operatorname{sech}^{-1}(a+bx)}{3b^3 x^3} \right) \\
 & \quad \downarrow \text{3042} \\
 & -b^3 \left( \frac{\operatorname{sech}^{-1}(a+bx)}{3b^3 x^3} + \frac{1}{3} \left( -\frac{\sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1)}{2a(1-a^2)b^2 x^2} + \frac{\int \frac{-\csc(i \operatorname{sech}^{-1}(a+bx) + \frac{\pi}{2})^2 - 2a \csc(i \operatorname{sech}^{-1}(a+bx) + \frac{\pi}{2}) + 2(1-a^2)}{(a - \csc(i \operatorname{sech}^{-1}(a+bx) + \frac{\pi}{2}))^2}}{2a(1-a^2)} \right) \right) \\
 & \quad \downarrow \text{4548}
 \end{aligned}$$

$$-b^3 \left( \frac{1}{3} \left( \frac{\int \frac{-2(1-a^2)^2 - a(1-4a^2)(a+bx)}{bx} d\operatorname{sech}^{-1}(a+bx)}{2a(1-a^2)} + \frac{(2-5a^2)\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a(1-a^2)bx} - \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{2a(1-a^2)b^2x^2} \right) + \frac{\operatorname{sech}^{-1}(a+bx)}{3b^3x^3} \right)$$

↓ 3042

$$-b^3 \left( \frac{\operatorname{sech}^{-1}(a+bx)}{3b^3x^3} + \frac{1}{3} \left( -\frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{2a(1-a^2)b^2x^2} + \frac{(2-5a^2)\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a(1-a^2)bx} + \frac{\int \frac{2(1-a^2)^2 - a(1-4a^2) \operatorname{csc}(i\operatorname{sech}^{-1}(a+bx))}{a - \operatorname{csc}(i\operatorname{sech}^{-1}(a+bx))} dx}{2a(1-a^2)} \right) \right)$$

↓ 4407

$$-b^3 \left( \frac{1}{3} \left( \frac{\frac{(6a^4-5a^2+2) \int \frac{-a+bx}{bx} d\operatorname{sech}^{-1}(a+bx)}{a} + \frac{2(1-a^2)^2 \operatorname{sech}^{-1}(a+bx)}{a}}{2a(1-a^2)} + \frac{(2-5a^2)\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a(1-a^2)bx} - \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{2a(1-a^2)b^2x^2} \right) + \frac{\operatorname{sech}^{-1}(a+bx)}{3b^3x^3} \right)$$

↓ 3042

$$-b^3 \left( \frac{\operatorname{sech}^{-1}(a+bx)}{3b^3x^3} + \frac{1}{3} \left( -\frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{2a(1-a^2)b^2x^2} + \frac{(2-5a^2)\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a(1-a^2)bx} + \frac{\frac{2(1-a^2)^2 \operatorname{sech}^{-1}(a+bx)}{a} + \frac{(6a^4-5a^2+2) \int \frac{-a+bx}{bx} d\operatorname{sech}^{-1}(a+bx)}{a}}{2a(1-a^2)} \right) \right)$$

↓ 4318

$$-b^3 \left( \frac{1}{3} \left( \frac{\frac{2(1-a^2)^2 \operatorname{sech}^{-1}(a+bx)}{a} - \frac{(6a^4-5a^2+2) \int \frac{1}{1-\frac{a}{a+bx}} d\operatorname{sech}^{-1}(a+bx)}{a}}{2a(1-a^2)} + \frac{(2-5a^2)\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a(1-a^2)bx} - \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{2a(1-a^2)b^2x^2} \right) + \frac{\operatorname{sech}^{-1}(a+bx)}{3b^3x^3} \right)$$

↓ 3042

$$-b^3 \left( \frac{\operatorname{sech}^{-1}(a+bx)}{3b^3x^3} + \frac{1}{3} \left( -\frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{2a(1-a^2)b^2x^2} + \frac{(2-5a^2)\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a(1-a^2)bx} + \frac{2(1-a^2)^2\operatorname{sech}^{-1}(a+bx)}{a} - \frac{(6a^4-5a^2)}{2a(1-a^2)} \right) \right)$$

↓ 3138

$$-b^3 \left( \frac{1}{3} \left( \frac{\frac{2(1-a^2)^2\operatorname{sech}^{-1}(a+bx)}{a} - \frac{2(6a^4-5a^2+2) \int \frac{1}{-(a+1)\tanh^2\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)} dx \tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{a(1-a^2)}}{2a(1-a^2)} + \frac{(2-5a^2)\sqrt{\frac{-a-bx+1}{a+bx+1}}}{a(1-a^2)} \right) \right)$$

↓ 221

$$-b^3 \left( \frac{1}{3} \left( \frac{\frac{(2-5a^2)\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a(1-a^2)bx} + \frac{2(1-a^2)^2\operatorname{sech}^{-1}(a+bx)}{a} - \frac{2(6a^4-5a^2+2)\operatorname{arctanh}\left(\frac{\sqrt{a+1}\tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{a\sqrt{1-a^2}}}{2a(1-a^2)} - \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}}{2a} \right) \right)$$

input `Int[ArcSech[a + b*x]/x^4,x]`

output `-(b^3*(ArcSech[a + b*x]/(3*b^3*x^3) + (-1/2*(Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x))/(a*(1 - a^2)*b^2*x^2) + (((2 - 5*a^2)*Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x))/(a*(1 - a^2)*b*x) + ((2*(1 - a^2)^2*ArcSech[a + b*x])/a - (2*(2 - 5*a^2 + 6*a^4)*ArcTanh[(Sqrt[1 + a]*Tanh[ArcSech[a + b*x]/2])/Sqrt[1 - a]])/(a*Sqrt[1 - a^2]))/(a*(1 - a^2)))/(2*a*(1 - a^2)))/3)`

## Definitions of rubi rules used

- rule 221  $\text{Int}[(a_ + (b_ \cdot x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) \cdot \text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] \text{ ; FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b]$
- rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$
- rule 3138  $\text{Int}[(a_ + (b_ \cdot \sin[\text{Pi}/2 + (c_ \cdot x_ ) + (d_ \cdot x_ )])^{-1}, x\_Symbol] \rightarrow \text{With}\{e = \text{FreeFactors}[\text{Tan}[(c + d \cdot x)/2], x]\}, \text{Simp}[2 \cdot (e/d) \text{ Subst}[\text{Int}[1/(a + b + (a - b) \cdot e^2 \cdot x^2), x], x, \text{Tan}[(c + d \cdot x)/2]/e], x]] \text{ ; FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$
- rule 4272  $\text{Int}[(\text{csc}[(c_ \cdot x_ ) + (d_ \cdot x_ )] \cdot (b_ \cdot x_ ) + (a_ ))^n, x\_Symbol] \rightarrow \text{Simp}[b^2 \cdot \text{Cot}[c + d \cdot x] \cdot ((a + b \cdot \text{Csc}[c + d \cdot x])^{n+1} / (a \cdot d \cdot (n+1) \cdot (a^2 - b^2))), x] + \text{Simp}[1 / (a \cdot (n+1) \cdot (a^2 - b^2)) \text{ Int}[(a + b \cdot \text{Csc}[c + d \cdot x])^{n+1} \cdot \text{Simp}[(a^2 - b^2) \cdot (n+1) - a \cdot b \cdot (n+1) \cdot \text{Csc}[c + d \cdot x] + b^2 \cdot (n+2) \cdot \text{Csc}[c + d \cdot x]^2, x], x], x] \text{ ; FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2 \cdot n]$
- rule 4318  $\text{Int}[\text{csc}[(e_ \cdot x_ ) + (f_ \cdot x_ )] / (\text{csc}[(e_ \cdot x_ ) + (f_ \cdot x_ )] \cdot (b_ \cdot x_ ) + (a_ )), x\_Symbol] \rightarrow \text{Simp}[1/b \text{ Int}[1/(1 + (a/b) \cdot \text{Sin}[e + f \cdot x]), x], x] \text{ ; FreeQ}\{a, b, e, f\}, x \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$
- rule 4407  $\text{Int}[(\text{csc}[(e_ \cdot x_ ) + (f_ \cdot x_ )] \cdot (d_ \cdot x_ ) + (c_ )) / (\text{csc}[(e_ \cdot x_ ) + (f_ \cdot x_ )] \cdot (b_ \cdot x_ ) + (a_ )), x\_Symbol] \rightarrow \text{Simp}[c \cdot (x/a), x] - \text{Simp}[(b \cdot c - a \cdot d) / a \text{ Int}[\text{Csc}[e + f \cdot x] / (a + b \cdot \text{Csc}[e + f \cdot x]), x], x] \text{ ; FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0]$

rule 4548

```
Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.
))*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_), x_Symbol] := Simp[(A*b^2 -
a*b*B + a^2*C)*Cot[e + f*x]*((a + b*Csc[e + f*x])^(m + 1)/(a*f*(m + 1)*(a^2
- b^2))), x] + Simp[1/(a*(m + 1)*(a^2 - b^2)) Int[(a + b*Csc[e + f*x])^(
m + 1)*Simp[A*(a^2 - b^2)*(m + 1) - a*(A*b - a*B + b*C)*(m + 1)*Csc[e + f*x
] + (A*b^2 - a*b*B + a^2*C)*(m + 2)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, e, f, A, B, C}, x] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]
```

rule 5991

```
Int[((e_.) + (f_.)*(x_.))^(m_.)*Sech[(c_.) + (d_.)*(x_.)]*((a_.) + (b_.)*Sech[
(c_.) + (d_.)*(x_.)]^(n_.)*Tanh[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[(-e
+ f*x)^m*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b
*d*(n + 1)) Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

rule 6875

```
Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(
m_.), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Sech[x]*T
anh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

## Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.97 (sec) , antiderivative size = 608, normalized size of antiderivative = 3.09

method	result
parts	$-\frac{\operatorname{arcsech}(bx+a)}{3x^3} - \frac{b\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \operatorname{csgn}(b)^2 \left(2 \operatorname{arctanh}\left(\frac{1}{\sqrt{-b^2x^2-2bxa-a^2+1}}\right)\right) a^6 b^2 x^2 + 6\sqrt{-a^2+1}}{3x^3}$
derivativedivides	Expression too large to display
default	Expression too large to display

input

```
int(arcsech(b*x+a)/x^4,x,method=_RETURNVERBOSE)
```

output

```

-1/3*arcsech(b*x+a)/x^3-1/6*b*(-(b*x+a-1)/(b*x+a))^(1/2)*(b*x+a)*((b*x+a+1)
)/(b*x+a))^(1/2)*csgn(b)^2*(2*arctanh(1/(-b^2*x^2-2*a*b*x-a^2+1)^(1/2))*a^
6*b^2*x^2+6*(-a^2+1)^(1/2)*ln(2*(-b*x*a+(-a^2+1)^(1/2)*(-b^2*x^2-2*a*b*x-a
^2+1)^(1/2)-a^2+1)/x)*a^4*b^2*x^2-6*arctanh(1/(-b^2*x^2-2*a*b*x-a^2+1)^(1/
2))*a^4*b^2*x^2-5*(-a^2+1)^(1/2)*ln(2*(-b*x*a+(-a^2+1)^(1/2)*(-b^2*x^2-2*a
*b*x-a^2+1)^(1/2)-a^2+1)/x)*a^2*b^2*x^2-5*a^5*b*x*(-b^2*x^2-2*a*b*x-a^2+1)
^(1/2)+6*arctanh(1/(-b^2*x^2-2*a*b*x-a^2+1)^(1/2))*a^2*b^2*x^2+a^6*(-b^2*x
^2-2*a*b*x-a^2+1)^(1/2)+2*(-a^2+1)^(1/2)*ln(2*(-b*x*a+(-a^2+1)^(1/2)*(-b^2
*x^2-2*a*b*x-a^2+1)^(1/2)-a^2+1)/x)*b^2*x^2+7*(-b^2*x^2-2*a*b*x-a^2+1)^(1/
2)*a^3*b*x-2*arctanh(1/(-b^2*x^2-2*a*b*x-a^2+1)^(1/2))*b^2*x^2-2*(-b^2*x^2
-2*a*b*x-a^2+1)^(1/2)*a^4-2*(-b^2*x^2-2*a*b*x-a^2+1)^(1/2)*a*b*x+(-b^2*x^2
-2*a*b*x-a^2+1)^(1/2)*a^2)/x^2/(a^2-1)^2/(a-1)/(1+a)/a^3/(-b^2*x^2-2*a*b*x
-a^2+1)^(1/2)

```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 487 vs.  $2(167) = 334$ .

Time = 0.16 (sec) , antiderivative size = 987, normalized size of antiderivative = 5.01

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^4} dx = \text{Too large to display}$$

input

```
integrate(arcsech(b*x+a)/x^4,x, algorithm="fricas")
```



output

```

[-1/12*((6*a^4 - 5*a^2 + 2)*sqrt(-a^2 + 1)*b^3*x^3*log(((2*a^2 - 1)*b^2*x^
2 + 2*a^4 + 4*(a^3 - a)*b*x - 4*a^2 - 2*(a*b^2*x^2 + a^3 + (2*a^2 - 1)*b*x
- a)*sqrt(-a^2 + 1)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x
+ a^2))) + 2)/x^2) + 2*(a^6 - 3*a^4 + 3*a^2 - 1)*b^3*x^3*log(((b*x + a)*s
qrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/x) - 2*
(a^6 - 3*a^4 + 3*a^2 - 1)*b^3*x^3*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x
+ a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) - 1)/x) + 4*(a^9 - 3*a^7 + 3*a^5 - a
^3)*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x
+ a^2)) + 1)/(b*x + a)) - 2*((5*a^5 - 7*a^3 + 2*a)*b^3*x^3 + (4*a^6 - 5*a^
4 + a^2)*b^2*x^2 - (a^7 - 2*a^5 + a^3)*b*x)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2
- 1)/(b^2*x^2 + 2*a*b*x + a^2)))/((a^9 - 3*a^7 + 3*a^5 - a^3)*x^3), -1/6*
((6*a^4 - 5*a^2 + 2)*sqrt(a^2 - 1)*b^3*x^3*arctan((a*b^2*x^2 + a^3 + (2*a^
2 - 1)*b*x - a)*sqrt(a^2 - 1)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2
+ 2*a*b*x + a^2)))/((a^2 - 1)*b^2*x^2 + a^4 + 2*(a^3 - a)*b*x - 2*a^2 + 1)
) + (a^6 - 3*a^4 + 3*a^2 - 1)*b^3*x^3*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*
b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/x) - (a^6 - 3*a^4 + 3*a^2 -
1)*b^3*x^3*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 +
2*a*b*x + a^2)) - 1)/x) + 2*(a^9 - 3*a^7 + 3*a^5 - a^3)*log(((b*x + a)*sq
rt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)
) - ((5*a^5 - 7*a^3 + 2*a)*b^3*x^3 + (4*a^6 - 5*a^4 + a^2)*b^2*x^2 - (a...

```

## Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^4} dx = \int \frac{\operatorname{asech}(a + bx)}{x^4} dx$$

input

```
integrate(asech(b*x+a)/x**4,x)
```

output

```
Integral(asech(a + b*x)/x**4, x)
```

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^4} dx = \int \frac{\operatorname{arosech}(bx + a)}{x^4} dx$$

input `integrate(arcsech(b*x+a)/x^4,x, algorithm="maxima")`

output `1/3*(6*a^4*b^3 - 3*a^2*b^3 + b^3)*log(x)/(a^9 - 3*a^7 + 3*a^5 - a^3) - 1/6*((a^6*b^3 - 3*a^5*b^3 + 3*a^4*b^3 - a^3*b^3)*x^3*log(b*x + a + 1) + (a^6*b^3 + 3*a^5*b^3 + 3*a^4*b^3 + a^3*b^3)*x^3*log(-b*x - a + 1) - 2*(3*a^5*b^2 - 4*a^3*b^2 + a*b^2)*x^2 + (a^6*b - 2*a^4*b + a^2*b)*x + 2*(a^9 - 3*a^7 + 3*a^5 - a^3)*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a) - 2*(a^9 - 3*a^7 + 3*a^5 + (a^6*b^3 - 3*a^4*b^3 + 3*a^2*b^3 - b^3)*x^3 - a^3)*log(b*x + a) - 2*(a^9 - 3*a^7 + 3*a^5 - a^3)*log(b*x + a))/((a^9 - 3*a^7 + 3*a^5 - a^3)*x^3) - integrate(1/3*(b^2*x + a*b)/(b^2*x^5 + 2*a*b*x^4 + (a^2 - 1)*x^3 + (b^2*x^5 + 2*a*b*x^4 + (a^2 - 1)*x^3)*e^(1/2*log(b*x + a + 1) + 1/2*log(-b*x - a + 1))), x)`

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^4} dx = \int \frac{\operatorname{arosech}(bx + a)}{x^4} dx$$

input `integrate(arcsech(b*x+a)/x^4,x, algorithm="giac")`

output `integrate(arcsech(b*x + a)/x^4, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^4} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)}{x^4} dx$$

input `int(acosh(1/(a + b*x))/x^4,x)`output `int(acosh(1/(a + b*x))/x^4, x)`**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^4} dx = \int \frac{\operatorname{asech}(bx + a)}{x^4} dx$$

input `int(asech(b*x+a)/x^4,x)`output `int(asech(a + b*x)/x**4,x)`

### 3.9 $\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx$

Optimal result	107
Mathematica [A] (verified)	108
Rubi [A] (verified)	109
Maple [A] (verified)	111
Fricas [F]	111
Sympy [F]	112
Maxima [F]	112
Giac [F]	113
Mupad [F(-1)]	113
Reduce [F]	113

#### Optimal result

Integrand size = 12, antiderivative size = 279

$$\begin{aligned}
 \int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx = & -\frac{x}{3b^2} + \frac{2a\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)}{b^3} \\
 & - \frac{(a+bx)\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)}{3b^3} \\
 & + \frac{a^3 \operatorname{sech}^{-1}(a+bx)^2}{3b^3} + \frac{1}{3}x^3 \operatorname{sech}^{-1}(a+bx)^2 \\
 & - \frac{2\operatorname{sech}^{-1}(a+bx) \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3} \\
 & - \frac{4a^2 \operatorname{sech}^{-1}(a+bx) \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3} \\
 & + \frac{2a \log(a+bx)}{b^3} + \frac{i \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3} \\
 & + \frac{2ia^2 \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3} \\
 & - \frac{i \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3} \\
 & - \frac{2ia^2 \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3}
 \end{aligned}$$

output

```
-1/3*x/b^2+2*a*((-b*x-a+1)/(b*x+a+1))^(1/2)*(b*x+a+1)*arcsech(b*x+a)/b^3-1/3*(b*x+a)*((-b*x-a+1)/(b*x+a+1))^(1/2)*(b*x+a+1)*arcsech(b*x+a)/b^3+1/3*a^3*arcsech(b*x+a)^2/b^3+1/3*x^3*arcsech(b*x+a)^2-2/3*arcsech(b*x+a)*arctan(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/b^3-4*a^2*arcsech(b*x+a)*arctan(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/b^3+2*a*ln(b*x+a)/b^3+1/3*I*polylog(2,-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b^3+2*I*a^2*polylog(2,-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b^3-1/3*I*polylog(2,I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b^3-2*I*a^2*polylog(2,I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b^3
```

**Mathematica [A] (verified)**

Time = 1.36 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.09

$$\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx =$$

$$\frac{2(a + bx) \sqrt{-\frac{-1+a+bx}{1+a+bx}} (1 + a + bx) \operatorname{sech}^{-1}(a + bx) + 6a(a + bx)^2 \operatorname{sech}^{-1}(a + bx)^2 - 2(a + bx)^3 \operatorname{sech}^{-1}(a + bx)}{b^3}$$

input

```
Integrate[x^2*ArcSech[a + b*x]^2,x]
```

output

```
-1/6*(2*(a + b*x)*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(1 + a + b*x)*ArcSech[a + b*x] + 6*a*(a + b*x)^2*ArcSech[a + b*x]^2 - 2*(a + b*x)^3*ArcSech[a + b*x]^2 + 2*(a + b*x - 6*a*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(1 + a + b*x)*ArcSech[a + b*x] - 3*a^2*(a + b*x)*ArcSech[a + b*x]^2) + 12*a*Log[(a + b*x)^(-1)] - (1 + 6*a^2)*(Pi*Log[1 - I*E^ArcSech[a + b*x]] - (2*I)*ArcSech[a + b*x]*Log[1 - I*E^ArcSech[a + b*x]] - Pi*Log[1 + I*E^ArcSech[a + b*x]]) + (2*I)*ArcSech[a + b*x]*Log[1 + I*E^ArcSech[a + b*x]] - Pi*Log[Cot[(Pi + (2*I)*ArcSech[a + b*x])/4]] + (2*I)*PolyLog[2, (-I)*E^ArcSech[a + b*x]] - (2*I)*PolyLog[2, I*E^ArcSech[a + b*x]]))/b^3
```

**Rubi [A] (verified)**

Time = 0.55 (sec) , antiderivative size = 266, normalized size of antiderivative = 0.95, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {6875, 5991, 3042, 4678, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx$$

$$\downarrow 6875$$

$$\frac{\int b^2 x^2 (a + bx) \sqrt{\frac{-a - bx + 1}{a + bx + 1}} (a + bx + 1) \operatorname{sech}^{-1}(a + bx)^2 d \operatorname{sech}^{-1}(a + bx)}{b^3}$$

$$\downarrow 5991$$

$$\frac{-\frac{2}{3} \int -b^3 x^3 \operatorname{sech}^{-1}(a + bx) d \operatorname{sech}^{-1}(a + bx) - \frac{1}{3} b^3 x^3 \operatorname{sech}^{-1}(a + bx)^2}{b^3}$$

$$\downarrow 3042$$

$$\frac{-\frac{1}{3} b^3 x^3 \operatorname{sech}^{-1}(a + bx)^2 - \frac{2}{3} \int \operatorname{sech}^{-1}(a + bx) \left( a - \operatorname{csc} \left( i \operatorname{sech}^{-1}(a + bx) + \frac{\pi}{2} \right) \right)^3 d \operatorname{sech}^{-1}(a + bx)}{b^3}$$

$$\downarrow 4678$$

$$\frac{-\frac{2}{3} \int \left( \operatorname{sech}^{-1}(a + bx) a^3 - 3(a + bx) \operatorname{sech}^{-1}(a + bx) a^2 + 3(a + bx)^2 \operatorname{sech}^{-1}(a + bx) a - (a + bx)^3 \operatorname{sech}^{-1}(a + bx) \right)}{b^3}$$

$$\downarrow 2009$$

$$\frac{-\frac{1}{3} b^3 x^3 \operatorname{sech}^{-1}(a + bx)^2 - \frac{2}{3} \left( \frac{1}{2} a^3 \operatorname{sech}^{-1}(a + bx)^2 - 6a^2 \operatorname{sech}^{-1}(a + bx) \arctan \left( e^{\operatorname{sech}^{-1}(a + bx)} \right) + 3ia^2 \operatorname{PolyLog} \right)}{b^3}$$

input `Int [x^2*ArcSech[a + b*x]^2,x]`

output

```

-((-1/3*(b^3*x^3*ArcSech[a + b*x]^2) - (2*((-a - b*x)/2 + 3*a*Sqrt[(1 - a
- b*x)/(1 + a + b*x)]*(1 + a + b*x)*ArcSech[a + b*x] - ((a + b*x)*Sqrt[(1
- a - b*x)/(1 + a + b*x)]*(1 + a + b*x)*ArcSech[a + b*x])/2 + (a^3*ArcSech
[a + b*x]^2)/2 - ArcSech[a + b*x]*ArcTan[E^ArcSech[a + b*x]] - 6*a^2*ArcSe
ch[a + b*x]*ArcTan[E^ArcSech[a + b*x]] - 3*a*Log[(a + b*x)^(-1)] + (I/2)*P
olyLog[2, (-I)*E^ArcSech[a + b*x]] + (3*I)*a^2*PolyLog[2, (-I)*E^ArcSech[a
+ b*x]] - (I/2)*PolyLog[2, I*E^ArcSech[a + b*x]] - (3*I)*a^2*PolyLog[2, I
*E^ArcSech[a + b*x]]))/3)/b^3)

```

### Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4678

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)
, x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

rule 5991

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sech[
(c_.) + (d_.)*(x_)]^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-(e
+ f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b
*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

rule 6875

```
Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Sech[x]*T
anh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

**Maple [A] (verified)**

Time = 0.51 (sec) , antiderivative size = 599, normalized size of antiderivative = 2.15

method	result
derivativedivides	$\frac{\operatorname{arcsech}(bx+a)^2 a^2 (bx+a) - \operatorname{arcsech}(bx+a)^2 a (bx+a)^2 + \frac{\operatorname{arcsech}(bx+a)^2 (bx+a)^3}{3} + 2 \operatorname{arcsech}(bx+a) \sqrt{\frac{bx+a+1}{bx+a}} \sqrt{-\frac{bx+a-1}{bx+a}}}{\operatorname{arcsech}(bx+a)^2 a^2 (bx+a) - \operatorname{arcsech}(bx+a)^2 a (bx+a)^2 + \frac{\operatorname{arcsech}(bx+a)^2 (bx+a)^3}{3} + 2 \operatorname{arcsech}(bx+a) \sqrt{\frac{bx+a+1}{bx+a}} \sqrt{-\frac{bx+a-1}{bx+a}}}$
default	$\operatorname{arcsech}(bx+a)^2 a^2 (bx+a) - \operatorname{arcsech}(bx+a)^2 a (bx+a)^2 + \frac{\operatorname{arcsech}(bx+a)^2 (bx+a)^3}{3} + 2 \operatorname{arcsech}(bx+a) \sqrt{\frac{bx+a+1}{bx+a}} \sqrt{-\frac{bx+a-1}{bx+a}}$

input `int(x^2*arcsech(b*x+a)^2,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/b^3 * (\operatorname{arcsech}(b*x+a)^2 * a^2 * (b*x+a) - \operatorname{arcsech}(b*x+a)^2 * a * (b*x+a)^2 + 1/3 * \operatorname{arcsech}(b*x+a)^2 * (b*x+a)^3 + 2 * \operatorname{arcsech}(b*x+a) * ((b*x+a+1)/(b*x+a))^{1/2} * (-(b*x+a-1)/(b*x+a))^{1/2} * a * (b*x+a) - 1/3 * \operatorname{arcsech}(b*x+a) * (b*x+a)^2 * (-(b*x+a-1)/(b*x+a))^{1/2} * ((b*x+a+1)/(b*x+a))^{1/2} - 2 * a * \operatorname{arcsech}(b*x+a) - 1/3 * b*x - 1/3 * a + 1/3 * I * \operatorname{dilog}(1 + I * (1/(b*x+a) + (1/(b*x+a) - 1)^{1/2} * (1/(b*x+a) + 1)^{1/2})) - 1/3 * I * \operatorname{arcsch}(b*x+a) * \ln(1 - I * (1/(b*x+a) + (1/(b*x+a) - 1)^{1/2} * (1/(b*x+a) + 1)^{1/2})) + 2 * I * a^2 * \operatorname{dilog}(1 + I * (1/(b*x+a) + (1/(b*x+a) - 1)^{1/2} * (1/(b*x+a) + 1)^{1/2})) - 1/3 * I * \operatorname{dilog}(1 - I * (1/(b*x+a) + (1/(b*x+a) - 1)^{1/2} * (1/(b*x+a) + 1)^{1/2})) - 2 * \ln(1 + (1/(b*x+a) + (1/(b*x+a) - 1)^{1/2} * (1/(b*x+a) + 1)^{1/2})^2) * a + 4 * a * \ln(1/(b*x+a) + (1/(b*x+a) - 1)^{1/2} * (1/(b*x+a) + 1)^{1/2}) - 2 * I * a^2 * \operatorname{dilog}(1 - I * (1/(b*x+a) + (1/(b*x+a) - 1)^{1/2} * (1/(b*x+a) + 1)^{1/2})) + 2 * I * a^2 * \operatorname{arcsech}(b*x+a) * \ln(1 + I * (1/(b*x+a) + (1/(b*x+a) - 1)^{1/2} * (1/(b*x+a) + 1)^{1/2})) + 1/3 * I * \operatorname{arcsech}(b*x+a) * \ln(1 + I * (1/(b*x+a) + (1/(b*x+a) - 1)^{1/2} * (1/(b*x+a) + 1)^{1/2})) - 2 * I * a^2 * \operatorname{arcsech}(b*x+a) * \ln(1 - I * (1/(b*x+a) + (1/(b*x+a) - 1)^{1/2} * (1/(b*x+a) + 1)^{1/2}))) \end{aligned}$$
**Fricas [F]**

$$\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx = \int x^2 \operatorname{arcsch}(bx + a)^2 dx$$

input `integrate(x^2*arcsech(b*x+a)^2,x, algorithm="fricas")`

output `integral(x^2*arcsech(b*x + a)^2, x)`



**Sympy [F]**

$$\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx = \int x^2 \operatorname{asech}^2(a + bx) dx$$

input `integrate(x**2*asech(b*x+a)**2,x)`

output `Integral(x**2*asech(a + b*x)**2, x)`

**Maxima [F]**

$$\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx = \int x^2 \operatorname{arsech}(bx + a)^2 dx$$

input `integrate(x^2*arcsech(b*x+a)^2,x, algorithm="maxima")`

output `1/3*x^3*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^2 - integrate(-2/3*(6*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^2 + 6*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2)*log(b*x + a)^2 - (b^3*x^5 + 2*a*b^2*x^4 + (a^2*b - b)*x^3 + 6*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2)*log(b*x + a) + (3*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2)*sqrt(b*x + a + 1)*log(b*x + a) + (2*b^3*x^5 + 4*a*b^2*x^4 + (2*a^2*b - b)*x^3 + 3*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2)*log(b*x + a))*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1) + (3*a^2*b - b)*x - a), x)`

**Giac [F]**

$$\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx = \int x^2 \operatorname{arsech}(bx + a)^2 dx$$

input `integrate(x^2*arcsech(b*x+a)^2,x, algorithm="giac")`

output `integrate(x^2*arcsech(b*x + a)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx = \int x^2 \operatorname{acosh}\left(\frac{1}{a + bx}\right)^2 dx$$

input `int(x^2*acosh(1/(a + b*x))^2,x)`

output `int(x^2*acosh(1/(a + b*x))^2, x)`

**Reduce [F]**

$$\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx = \int \operatorname{asech}(bx + a)^2 x^2 dx$$

input `int(x^2*asech(b*x+a)^2,x)`

output `int(asech(a + b*x)**2*x**2,x)`

### 3.10 $\int x \operatorname{sech}^{-1}(a + bx)^2 dx$

Optimal result	114
Mathematica [A] (verified)	115
Rubi [A] (verified)	115
Maple [A] (verified)	117
Fricas [F]	118
Sympy [F]	118
Maxima [F]	119
Giac [F]	119
Mupad [F(-1)]	120
Reduce [F]	120

#### Optimal result

Integrand size = 10, antiderivative size = 149

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx = -\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)}{b^2} - \frac{a^2 \operatorname{sech}^{-1}(a+bx)^2}{2b^2} + \frac{1}{2}x^2 \operatorname{sech}^{-1}(a+bx)^2 + \frac{4a \operatorname{sech}^{-1}(a+bx) \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} - \frac{\log(a+bx)}{b^2} - \frac{2ia \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} + \frac{2ia \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2}$$

output

```
-((-b*x-a+1)/(b*x+a+1))^(1/2)*(b*x+a+1)*arcsech(b*x+a)/b^2-1/2*a^2*arcsech
(b*x+a)^2/b^2+1/2*x^2*arcsech(b*x+a)^2+4*a*arcsech(b*x+a)*arctan(1/(b*x+a)
+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/b^2-ln(b*x+a)/b^2-2*I*a*polylog(
2,-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b^2+2*I*a*polylo
g(2,I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b^2
```

**Mathematica [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.15

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx$$

$$= \frac{-2\sqrt{-\frac{-1+a+bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx) - 2a(a+bx)\operatorname{sech}^{-1}(a+bx)^2 + (a+bx)^2\operatorname{sech}^{-1}(a+bx)^2 - 4a(a+bx)\operatorname{sech}^{-1}(a+bx)^3 + 2(a+bx)^2\operatorname{sech}^{-1}(a+bx)^3}{b^2}$$

input

Integrate[x\*ArcSech[a + b\*x]^2,x]

output

```
(-2*sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(1 + a + b*x)*ArcSech[a + b*x] -
2*a*(a + b*x)*ArcSech[a + b*x]^2 + (a + b*x)^2*ArcSech[a + b*x]^2 - (4*I)
*a*ArcSech[a + b*x]*(Log[1 - I/E^ArcSech[a + b*x]] - Log[1 + I/E^ArcSech[a
+ b*x]]) + 2*Log[(a + b*x)^(-1)] - (4*I)*a*(PolyLog[2, (-I)/E^ArcSech[a +
b*x]] - PolyLog[2, I/E^ArcSech[a + b*x]]))/(2*b^2)
```

**Rubi [A] (verified)**Time = 0.44 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {6875, 25, 5991, 3042, 4678, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx$$

$$\downarrow \text{6875}$$

$$\frac{\int bx(a + bx)\sqrt{\frac{-a-bx+1}{a+bx+1}}(a + bx + 1)\operatorname{sech}^{-1}(a + bx)^2 d\operatorname{sech}^{-1}(a + bx)}{b^2}$$

$$\downarrow \text{25}$$

$$\frac{\int -bx(a + bx)\sqrt{\frac{-a-bx+1}{a+bx+1}}(a + bx + 1)\operatorname{sech}^{-1}(a + bx)^2 d\operatorname{sech}^{-1}(a + bx)}{b^2}$$

$$\begin{aligned}
& \int \frac{b^2 x^2 \operatorname{sech}^{-1}(a+bx) d\operatorname{sech}^{-1}(a+bx) - \frac{1}{2} b^2 x^2 \operatorname{sech}^{-1}(a+bx)^2}{b^2} \\
& \quad \downarrow \text{5991} \\
& \int \frac{-\frac{1}{2} b^2 x^2 \operatorname{sech}^{-1}(a+bx)^2 + \int \operatorname{sech}^{-1}(a+bx) \left(a - \csc\left(\operatorname{isech}^{-1}(a+bx) + \frac{\pi}{2}\right)\right)^2 d\operatorname{sech}^{-1}(a+bx)}{b^2} \\
& \quad \downarrow \text{3042} \\
& \int \frac{(\operatorname{sech}^{-1}(a+bx)a^2 - 2(a+bx)\operatorname{sech}^{-1}(a+bx)a + (a+bx)^2 \operatorname{sech}^{-1}(a+bx)) d\operatorname{sech}^{-1}(a+bx) - \frac{1}{2} b^2 x^2 \operatorname{sech}^{-1}(a+bx)}{b^2} \\
& \quad \downarrow \text{4678} \\
& \int \frac{(\operatorname{sech}^{-1}(a+bx)a^2 - 2(a+bx)\operatorname{sech}^{-1}(a+bx)a + (a+bx)^2 \operatorname{sech}^{-1}(a+bx)) d\operatorname{sech}^{-1}(a+bx) - \frac{1}{2} b^2 x^2 \operatorname{sech}^{-1}(a+bx)}{b^2} \\
& \quad \downarrow \text{2009} \\
& \int \frac{\frac{1}{2} a^2 \operatorname{sech}^{-1}(a+bx)^2 - 4a \operatorname{sech}^{-1}(a+bx) \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right) - \frac{1}{2} b^2 x^2 \operatorname{sech}^{-1}(a+bx)^2 + 2ia \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2}
\end{aligned}$$

input `Int[x*ArcSech[a + b*x]^2,x]`

output `-((Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x)*ArcSech[a + b*x] + (a^2 *ArcSech[a + b*x]^2)/2 - (b^2*x^2*ArcSech[a + b*x]^2)/2 - 4*a*ArcSech[a + b*x]*ArcTan[E^ArcSech[a + b*x]] - Log[(a + b*x)^(-1)] + (2*I)*a*PolyLog[2, (-I)*E^ArcSech[a + b*x]] - (2*I)*a*PolyLog[2, I*E^ArcSech[a + b*x]])/b^2)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4678 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

rule 5991 `Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sech[(c_.) + (d_.)*(x_)])^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

rule 6875 `Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)])*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

## Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 331, normalized size of antiderivative = 2.22

method	result
derivativedivides	$-\frac{\operatorname{arcsech}(bx+a)\left(2\sqrt{-\frac{bx+a-1}{bx+a}}\sqrt{\frac{bx+a+1}{bx+a}}(bx+a)+2\operatorname{arcsech}(bx+a)a(bx+a)-\operatorname{arcsech}(bx+a)(bx+a)^2-2\right)}{2}-2\ln\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}}\right)$
default	$-\frac{\operatorname{arcsech}(bx+a)\left(2\sqrt{-\frac{bx+a-1}{bx+a}}\sqrt{\frac{bx+a+1}{bx+a}}(bx+a)+2\operatorname{arcsech}(bx+a)a(bx+a)-\operatorname{arcsech}(bx+a)(bx+a)^2-2\right)}{2}-2\ln\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}}\right)$

input `int(x*arcsech(b*x+a)^2,x,method=_RETURNVERBOSE)`

output

```
1/b^2*(-1/2*arcsech(b*x+a)*(2*(-(b*x+a-1)/(b*x+a))^(1/2)*((b*x+a+1)/(b*x+a))^(1/2)*(b*x+a)+2*arcsech(b*x+a)*a*(b*x+a)-arcsech(b*x+a)*(b*x+a)^2-2*ln(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))+ln(1+(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))^2)-2*I*a*arcsech(b*x+a)*ln(1+I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))))+2*I*a*arcsech(b*x+a)*ln(1-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))))-2*I*a*dilog(1+I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))))+2*I*a*dilog(1-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))))
```

**Fricas [F]**

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx = \int x \operatorname{arsech}(bx + a)^2 dx$$

input

```
integrate(x*arcsech(b*x+a)^2,x, algorithm="fricas")
```

output

```
integral(x*arcsech(b*x + a)^2, x)
```

**Sympy [F]**

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx = \int x \operatorname{asech}^2(a + bx) dx$$

input

```
integrate(x*asech(b*x+a)**2,x)
```

output

```
Integral(x*asech(a + b*x)**2, x)
```

**Maxima [F]**

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx = \int x \operatorname{arsech}(bx + a)^2 dx$$

input `integrate(x*arcsech(b*x+a)^2,x, algorithm="maxima")`

output `1/2*x^2*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^2 - integrate(-(4*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^2 + 4*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*log(b*x + a)^2 - (b^3*x^4 + 2*a*b^2*x^3 + (a^2*b - b)*x^2 + 4*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*log(b*x + a) + (2*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*sqrt(b*x + a + 1)*log(b*x + a) + (2*b^3*x^4 + 4*a*b^2*x^3 + (2*a^2*b - b)*x^2 + 2*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*log(b*x + a))*sqrt(b*x + a + 1))*sqrt(-b*x - a + 1)*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1) + (3*a^2*b - b)*x - a), x)`

**Giac [F]**

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx = \int x \operatorname{arsech}(bx + a)^2 dx$$

input `integrate(x*arcsech(b*x+a)^2,x, algorithm="giac")`

output `integrate(x*arcsech(b*x + a)^2, x)`



**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx = \int x \operatorname{acosh}\left(\frac{1}{a + bx}\right)^2 dx$$

input `int(x*acosh(1/(a + b*x))^2,x)`output `int(x*acosh(1/(a + b*x))^2, x)`**Reduce [F]**

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx = \int \operatorname{asech}(bx + a)^2 x dx$$

input `int(x*asech(b*x+a)^2,x)`output `int(asech(a + b*x)**2*x,x)`

### 3.11 $\int \operatorname{sech}^{-1}(a + bx)^2 dx$

Optimal result	121
Mathematica [A] (verified)	121
Rubi [A] (verified)	122
Maple [A] (verified)	124
Fricas [F]	125
Sympy [F]	125
Maxima [F]	126
Giac [F]	126
Mupad [F(-1)]	127
Reduce [F]	127

#### Optimal result

Integrand size = 8, antiderivative size = 80

$$\int \operatorname{sech}^{-1}(a + bx)^2 dx = \frac{(a + bx)\operatorname{sech}^{-1}(a + bx)^2}{b} - \frac{4\operatorname{sech}^{-1}(a + bx) \arctan\left(e^{\operatorname{sech}^{-1}(a + bx)}\right)}{b} + \frac{2i \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b} - \frac{2i \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b}$$

output

```
(b*x+a)*arcsech(b*x+a)^2/b-4*arcsech(b*x+a)*arctan(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/b+2*I*polylog(2,-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b-2*I*polylog(2,I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b
```

#### Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.31

$$\int \operatorname{sech}^{-1}(a + bx)^2 dx = \frac{i\left(\operatorname{sech}^{-1}(a + bx)\left(-i(a + bx)\operatorname{sech}^{-1}(a + bx) + 2 \log\left(1 - ie^{-\operatorname{sech}^{-1}(a + bx)}\right) - 2 \log\left(1 + ie^{-\operatorname{sech}^{-1}(a + bx)}\right)\right)\right)}{b}$$

input `Integrate[ArcSech[a + b*x]^2,x]`

output  $(I*(ArcSech[a + b*x]*((-I)*(a + b*x)*ArcSech[a + b*x] + 2*Log[1 - I/E^ArcSech[a + b*x]]) - 2*Log[1 + I/E^ArcSech[a + b*x]]) + 2*PolyLog[2, (-I)/E^ArcSech[a + b*x]] - 2*PolyLog[2, I/E^ArcSech[a + b*x]]))/b$

### Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.96, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$ , Rules used = {6869, 6833, 5941, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \operatorname{sech}^{-1}(a + bx)^2 dx \\
 & \quad \downarrow \text{6869} \\
 & \frac{\int \operatorname{sech}^{-1}(a + bx)^2 d(a + bx)}{b} \\
 & \quad \downarrow \text{6833} \\
 & -\frac{\int (a + bx) \sqrt{\frac{-a - bx + 1}{a + bx + 1}} (a + bx + 1) \operatorname{sech}^{-1}(a + bx)^2 d\operatorname{sech}^{-1}(a + bx)}{b} \\
 & \quad \downarrow \text{5941} \\
 & -\frac{2 \int (a + bx) \operatorname{sech}^{-1}(a + bx) d\operatorname{sech}^{-1}(a + bx) - (a + bx) \operatorname{sech}^{-1}(a + bx)^2}{b} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{(a + bx) \operatorname{sech}^{-1}(a + bx)^2 + 2 \int \operatorname{sech}^{-1}(a + bx) \csc\left(\operatorname{sech}^{-1}(a + bx) + \frac{\pi}{2}\right) d\operatorname{sech}^{-1}(a + bx)}{b} \\
 & \quad \downarrow \text{4668} \\
 & -\frac{(a + bx) \operatorname{sech}^{-1}(a + bx)^2 + 2\left(-i \int \log\left(1 - ie^{\operatorname{sech}^{-1}(a + bx)}\right) d\operatorname{sech}^{-1}(a + bx) + i \int \log\left(1 + ie^{\operatorname{sech}^{-1}(a + bx)}\right) d\operatorname{sech}^{-1}(a + bx)\right)}{b}
 \end{aligned}$$

↓ 2715

$$\frac{-(a+bx)\operatorname{sech}^{-1}(a+bx)^2 + 2\left(-i \int e^{-\operatorname{sech}^{-1}(a+bx)} \log\left(1 - ie^{\operatorname{sech}^{-1}(a+bx)}\right) de^{\operatorname{sech}^{-1}(a+bx)} + i \int e^{-\operatorname{sech}^{-1}(a+bx)} \log\left(1 + ie^{\operatorname{sech}^{-1}(a+bx)}\right) de^{\operatorname{sech}^{-1}(a+bx)}\right)}{b}$$

↓ 2838

$$\frac{-(a+bx)\operatorname{sech}^{-1}(a+bx)^2 + 2\left(2\operatorname{sech}^{-1}(a+bx) \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right) - i \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right) + i \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)\right)}{b}$$

input `Int[ArcSech[a + b*x]^2,x]`

output `-(((a + b*x)*ArcSech[a + b*x]^2) + 2*(2*ArcSech[a + b*x]*ArcTan[E^ArcSech[a + b*x]] - I*PolyLog[2, (-I)*E^ArcSech[a + b*x]] + I*PolyLog[2, I*E^ArcSech[a + b*x]]))/b)`

### Defintions of rubi rules used

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668

```
Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

rule 5941

```
Int[(x_)^(m_.)*Sech[(a_.) + (b_.)*(x_)^(n_.)]^(p_.)*Tanh[(a_.) + (b_.)*(x_)^(n_.)]^(q_.), x_Symbol] := Simp[(-x^(m - n + 1))*(Sech[a + b*x^n]^p/(b*n*p)), x] + Simp[(m - n + 1)/(b*n*p) Int[x^(m - n)*Sech[a + b*x^n]^p, x], x] /; FreeQ[{a, b, p}, x] && RationalQ[m] && IntegerQ[n] && GeQ[m - n, 0] && EqQ[q, 1]
```

rule 6833

```
Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[-c^(-1) Subst[Int[(a + b*x)^n*Sech[x]*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[n, 0]
```

rule 6869

```
Int[((a_.) + ArcSech[(c_) + (d_.)*(x_)]*(b_.))^(p_.), x_Symbol] := Simp[1/d Subst[Int[(a + b*ArcSech[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[p, 0]
```

## Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 192, normalized size of antiderivative = 2.40

method	result
derivativedivides	$\frac{\operatorname{arcsech}(bx+a)^2(bx+a)+2i \operatorname{arcsech}(bx+a) \ln\left(1+i\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)-2i \operatorname{arcsech}(bx+a) \ln\left(1-i\left(\frac{1}{bx+a}-\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)}{1}$
default	$\frac{\operatorname{arcsech}(bx+a)^2(bx+a)+2i \operatorname{arcsech}(bx+a) \ln\left(1+i\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)-2i \operatorname{arcsech}(bx+a) \ln\left(1-i\left(\frac{1}{bx+a}-\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)}{1}$

input

```
int(arcsech(b*x+a)^2,x,method=_RETURNVERBOSE)
```

output

```
1/b*(arcsech(b*x+a)^2*(b*x+a)+2*I*arcsech(b*x+a)*ln(1+I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))-2*I*arcsech(b*x+a)*ln(1-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))+2*I*dilog(1+I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))-2*I*dilog(1-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))))
```

**Fricas [F]**

$$\int \operatorname{sech}^{-1}(a + bx)^2 dx = \int \operatorname{arsech}(bx + a)^2 dx$$

input

```
integrate(arcsech(b*x+a)^2,x, algorithm="fricas")
```

output

```
integral(arcsech(b*x + a)^2, x)
```

**Sympy [F]**

$$\int \operatorname{sech}^{-1}(a + bx)^2 dx = \int \operatorname{asech}^2(a + bx) dx$$

input

```
integrate(asech(b*x+a)**2,x)
```

output

```
Integral(asech(a + b*x)**2, x)
```

**Maxima [F]**

$$\int \operatorname{sech}^{-1}(a + bx)^2 dx = \int \operatorname{arsech}(bx + a)^2 dx$$

input `integrate(arcsech(b*x+a)^2,x, algorithm="maxima")`

output `x*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^2 - integrate(-2*(2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^2 + 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a)^2 - (b^3*x^3 + 2*a*b^2*x^2 + (a^2*b - b)*x + 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a) + ((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*log(b*x + a) + (2*b^3*x^3 + 4*a*b^2*x^2 + (2*a^2*b - b)*x + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a))*sqrt(b*x + a + 1))*sqrt(-b*x - a + 1))*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1) + (3*a^2*b - b)*x - a), x)`

**Giac [F]**

$$\int \operatorname{sech}^{-1}(a + bx)^2 dx = \int \operatorname{arsech}(bx + a)^2 dx$$

input `integrate(arcsech(b*x+a)^2,x, algorithm="giac")`

output `integrate(arcsech(b*x + a)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{sech}^{-1}(a + bx)^2 dx = \int \operatorname{acosh}\left(\frac{1}{a + bx}\right)^2 dx$$

input `int(acosh(1/(a + b*x))^2,x)`output `int(acosh(1/(a + b*x))^2, x)`**Reduce [F]**

$$\int \operatorname{sech}^{-1}(a + bx)^2 dx = \int \operatorname{asech}(bx + a)^2 dx$$

input `int(asech(b*x+a)^2,x)`output `int(asech(a + b*x)**2,x)`



### 3.12 $\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} dx$

Optimal result	128
Mathematica [A] (verified)	129
Rubi [C] (verified)	130
Maple [F]	135
Fricas [F]	136
Sympy [F]	136
Maxima [F]	136
Giac [F]	137
Mupad [F(-1)]	137
Reduce [F]	137

#### Optimal result

Integrand size = 12, antiderivative size = 274

$$\begin{aligned}
 \int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} dx &= \operatorname{sech}^{-1}(a+bx)^2 \log \left( 1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
 &\quad + \operatorname{sech}^{-1}(a+bx)^2 \log \left( 1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
 &\quad - \operatorname{sech}^{-1}(a+bx)^2 \log \left( 1 + e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
 &\quad + 2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left( 2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
 &\quad + 2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left( 2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
 &\quad - \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left( 2, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
 &\quad - 2 \operatorname{PolyLog} \left( 3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
 &\quad - 2 \operatorname{PolyLog} \left( 3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
 &\quad + \frac{1}{2} \operatorname{PolyLog} \left( 3, -e^{2\operatorname{sech}^{-1}(a+bx)} \right)
 \end{aligned}$$

output

```
arcsech(b*x+a)^2*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)
)/(1-(-a^2+1)^(1/2)))+arcsech(b*x+a)^2*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)
)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2))-arcsech(b*x+a)^2*ln(1+(1/(b*x+
a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)+2*arcsech(b*x+a)*polylog(2,
a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))+
2*arcsech(b*x+a)*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(
1/2))/(1+(-a^2+1)^(1/2)))-arcsech(b*x+a)*polylog(2,-(1/(b*x+a)+(1/(b*x+a)
-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)-2*polylog(3,a*(1/(b*x+a)+(1/(b*x+a)-1)^(
1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))-2*polylog(3,a*(1/(b*x+a)+(1/
(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))+1/2*polylog(3,-(
1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)
```

**Mathematica [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.02

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} dx = -\frac{2}{3}\operatorname{sech}^{-1}(a+bx)^3 - \operatorname{sech}^{-1}(a+bx)^2 \log\left(1 + e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\ + \operatorname{sech}^{-1}(a+bx)^2 \log\left(1 + \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}}\right) \\ + \operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\ + \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\ + 2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}}\right) \\ + 2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\ + \frac{1}{2} \operatorname{PolyLog}\left(3, -e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\ - 2 \operatorname{PolyLog}\left(3, -\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}}\right) \\ - 2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right)$$

input

Integrate[ArcSech[a + b\*x]^2/x, x]

output

```
(-2*ArcSech[a + b*x]^3)/3 - ArcSech[a + b*x]^2*Log[1 + E^(-2*ArcSech[a + b*x])] + ArcSech[a + b*x]^2*Log[1 + (a*E^ArcSech[a + b*x])/(-1 + Sqrt[1 - a^2])] + ArcSech[a + b*x]^2*Log[1 - (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])] + ArcSech[a + b*x]*PolyLog[2, -E^(-2*ArcSech[a + b*x])] + 2*ArcSech[a + b*x]*PolyLog[2, -((a*E^ArcSech[a + b*x])/(-1 + Sqrt[1 - a^2]))] + 2*ArcSech[a + b*x]*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])] + PolyLog[3, -E^(-2*ArcSech[a + b*x])]/2 - 2*PolyLog[3, -((a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2]))] - 2*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])]
```

### Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.58 (sec) , antiderivative size = 337, normalized size of antiderivative = 1.23, number of steps used = 17, number of rules used = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.333$ , Rules used = {6875, 25, 6129, 6104, 25, 3042, 26, 4201, 2620, 3011, 2720, 6096, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x} dx$$

↓ 6875

$$- \int \frac{(a + bx) \sqrt{\frac{-a - bx + 1}{a + bx + 1}} (a + bx + 1) \operatorname{sech}^{-1}(a + bx)^2}{bx} d\operatorname{sech}^{-1}(a + bx)$$

↓ 25

$$\int - \frac{(a + bx) \sqrt{\frac{-a - bx + 1}{a + bx + 1}} (a + bx + 1) \operatorname{sech}^{-1}(a + bx)^2}{bx} d\operatorname{sech}^{-1}(a + bx)$$

↓ 6129

$$\int \frac{\sqrt{\frac{-a - bx + 1}{a + bx + 1}} (a + bx + 1) \operatorname{sech}^{-1}(a + bx)^2}{\frac{a}{a + bx} - 1} d\operatorname{sech}^{-1}(a + bx)$$

↓ 6104

$$\begin{aligned}
& a \int -\frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^2}{(a+bx)\left(1-\frac{a}{a+bx}\right)}d\operatorname{sech}^{-1}(a+bx) - \int \sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^2d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow 25 \\
& - \int \sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^2d\operatorname{sech}^{-1}(a+bx) - \\
& \quad a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^2}{(a+bx)\left(1-\frac{a}{a+bx}\right)}d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow 3042 \\
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^2}{(a+bx)\left(1-\frac{a}{a+bx}\right)}d\operatorname{sech}^{-1}(a+bx) - \int -i\operatorname{sech}^{-1}(a+bx)^2 \tan(i\operatorname{sech}^{-1}(a+bx))d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow 26 \\
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^2}{(a+bx)\left(1-\frac{a}{a+bx}\right)}d\operatorname{sech}^{-1}(a+bx) + i \int \operatorname{sech}^{-1}(a+bx)^2 \tan(i\operatorname{sech}^{-1}(a+bx))d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow 4201 \\
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^2}{(a+bx)\left(1-\frac{a}{a+bx}\right)}d\operatorname{sech}^{-1}(a+bx) + \\
& \quad i \left( 2i \int \frac{e^{2\operatorname{sech}^{-1}(a+bx)}\operatorname{sech}^{-1}(a+bx)^2}{1+e^{2\operatorname{sech}^{-1}(a+bx)}}d\operatorname{sech}^{-1}(a+bx) - \frac{1}{3}i\operatorname{sech}^{-1}(a+bx)^3 \right) \\
& \quad \downarrow 2620 \\
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^2}{(a+bx)\left(1-\frac{a}{a+bx}\right)}d\operatorname{sech}^{-1}(a+bx) + \\
& \quad i \left( 2i \left( \frac{1}{2}\operatorname{sech}^{-1}(a+bx)^2 \log(e^{2\operatorname{sech}^{-1}(a+bx)}+1) - \int \operatorname{sech}^{-1}(a+bx) \log(1+e^{2\operatorname{sech}^{-1}(a+bx)})d\operatorname{sech}^{-1}(a+bx) \right) \right) \\
& \quad \downarrow 3011
\end{aligned}$$

$$\begin{aligned}
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^2}{(a+bx)\left(1-\frac{a}{a+bx}\right)} d\operatorname{sech}^{-1}(a+bx) + \\
& i\left(2i\left(-\frac{1}{2}\int \operatorname{PolyLog}\left(2,-e^{2\operatorname{sech}^{-1}(a+bx)}\right) d\operatorname{sech}^{-1}(a+bx) + \frac{1}{2}\operatorname{sech}^{-1}(a+bx)\operatorname{PolyLog}\left(2,-e^{2\operatorname{sech}^{-1}(a+bx)}\right) + \frac{1}{2}\right)\right)
\end{aligned}$$

↓ 2720

$$\begin{aligned}
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^2}{(a+bx)\left(1-\frac{a}{a+bx}\right)} d\operatorname{sech}^{-1}(a+bx) + \\
& i\left(2i\left(-\frac{1}{4}\int e^{-2\operatorname{sech}^{-1}(a+bx)}\operatorname{PolyLog}\left(2,-e^{2\operatorname{sech}^{-1}(a+bx)}\right) de^{2\operatorname{sech}^{-1}(a+bx)} + \frac{1}{2}\operatorname{sech}^{-1}(a+bx)\operatorname{PolyLog}\left(2,-e^{2\operatorname{sech}^{-1}(a+bx)}\right)\right)\right)
\end{aligned}$$

↓ 6096

$$\begin{aligned}
& -a\left(\int \frac{e^{\operatorname{sech}^{-1}(a+bx)}\operatorname{sech}^{-1}(a+bx)^2}{-e^{\operatorname{sech}^{-1}(a+bx)}a-\sqrt{1-a^2}+1} d\operatorname{sech}^{-1}(a+bx) + \int \frac{e^{\operatorname{sech}^{-1}(a+bx)}\operatorname{sech}^{-1}(a+bx)^2}{-e^{\operatorname{sech}^{-1}(a+bx)}a+\sqrt{1-a^2}+1} d\operatorname{sech}^{-1}(a+bx) + \dots\right) \\
& i\left(2i\left(-\frac{1}{4}\int e^{-2\operatorname{sech}^{-1}(a+bx)}\operatorname{PolyLog}\left(2,-e^{2\operatorname{sech}^{-1}(a+bx)}\right) de^{2\operatorname{sech}^{-1}(a+bx)} + \frac{1}{2}\operatorname{sech}^{-1}(a+bx)\operatorname{PolyLog}\left(2,-e^{2\operatorname{sech}^{-1}(a+bx)}\right)\right)\right)
\end{aligned}$$

↓ 2620

$$\begin{aligned}
& -a\left(\frac{2\int \operatorname{sech}^{-1}(a+bx)\log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) d\operatorname{sech}^{-1}(a+bx)}{a} + \frac{2\int \operatorname{sech}^{-1}(a+bx)\log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) d\operatorname{sech}^{-1}(a+bx)}{a} + \dots\right) \\
& i\left(2i\left(-\frac{1}{4}\int e^{-2\operatorname{sech}^{-1}(a+bx)}\operatorname{PolyLog}\left(2,-e^{2\operatorname{sech}^{-1}(a+bx)}\right) de^{2\operatorname{sech}^{-1}(a+bx)} + \frac{1}{2}\operatorname{sech}^{-1}(a+bx)\operatorname{PolyLog}\left(2,-e^{2\operatorname{sech}^{-1}(a+bx)}\right)\right)\right)
\end{aligned}$$

↓ 3011

$$\begin{aligned}
& -a\left(\frac{2\left(\int \operatorname{PolyLog}\left(2,\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) d\operatorname{sech}^{-1}(a+bx) - \operatorname{sech}^{-1}(a+bx)\operatorname{PolyLog}\left(2,\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)\right)}{a} + \frac{2\left(\int \operatorname{PolyLog}\left(2,\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) d\operatorname{sech}^{-1}(a+bx) - \operatorname{sech}^{-1}(a+bx)\operatorname{PolyLog}\left(2,\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)\right)}{a} + \dots\right) \\
& i\left(2i\left(-\frac{1}{4}\int e^{-2\operatorname{sech}^{-1}(a+bx)}\operatorname{PolyLog}\left(2,-e^{2\operatorname{sech}^{-1}(a+bx)}\right) de^{2\operatorname{sech}^{-1}(a+bx)} + \frac{1}{2}\operatorname{sech}^{-1}(a+bx)\operatorname{PolyLog}\left(2,-e^{2\operatorname{sech}^{-1}(a+bx)}\right)\right)\right)
\end{aligned}$$

↓ 2720

$$\begin{aligned}
& -a \left( \frac{2 \left( \int e^{-\operatorname{sech}^{-1}(a+bx)} \operatorname{PolyLog} \left( 2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) de^{\operatorname{sech}^{-1}(a+bx)} - \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left( 2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) \right)}{a} \right) \\
& i \left( 2i \left( -\frac{1}{4} \int e^{-2\operatorname{sech}^{-1}(a+bx)} \operatorname{PolyLog} \left( 2, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) de^{2\operatorname{sech}^{-1}(a+bx)} + \frac{1}{2} \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left( 2, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) \right) \right) \\
& \quad \downarrow \text{7143} \\
& -a \left( \frac{2 \left( \operatorname{PolyLog} \left( 3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) - \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left( 2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) \right)}{a} \right) + \frac{2 \left( \operatorname{PolyLog} \left( 3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1} \right) \right)}{a} \\
& i \left( 2i \left( \frac{1}{2} \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left( 2, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) - \frac{1}{4} \operatorname{PolyLog} \left( 3, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) + \frac{1}{2} \operatorname{sech}^{-1}(a+bx)^2 \log \left( \dots \right) \right) \right)
\end{aligned}$$

input `Int[ArcSech[a + b*x]^2/x, x]`

output `-(a*(ArcSech[a + b*x]^3/(3*a) - (ArcSech[a + b*x]^2*Log[1 - (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/a - (ArcSech[a + b*x]^2*Log[1 - (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])/a + (2*(-(ArcSech[a + b*x]*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])]) + PolyLog[3, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/a + (2*(-(ArcSech[a + b*x]*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])]) + PolyLog[3, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])/a)) + I*((-1/3*I)*ArcSech[a + b*x]^3 + (2*I)*((ArcSech[a + b*x]^2*Log[1 + E^(2*ArcSech[a + b*x])])/2 + (ArcSech[a + b*x]*PolyLog[2, -E^(2*ArcSech[a + b*x])])/2 - PolyLog[3, -E^(2*ArcSech[a + b*x])])/4)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)
*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4201

```
Int[(((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_]*(f_)*(x_)]), x
_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[
(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x))), x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

rule 6096

```
Int[(((e_) + (f_)*(x_))^(m_)*Sinh[(c_) + (d_)*(x_)])/(Cosh[(c_) + (d_
)*(x_)]*(b_) + (a_)), x_Symbol] := Simp[-(e + f*x)^(m + 1)/(b*f*(m + 1)),
x] + (Int[(e + f*x)^m*(E^(c + d*x)/(a - Rt[a^2 - b^2, 2] + b*E^(c + d*x))
, x] + Int[(e + f*x)^m*(E^(c + d*x)/(a + Rt[a^2 - b^2, 2] + b*E^(c + d*x))
, x)) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[a^2 - b^2, 0]
```

rule 6104

```
Int[(((e_.) + (f_.)*(x_))^(m_.)*Tanh[(c_.) + (d_.)*(x_)]^(n_.))/(Cosh[(c_.)
+ (d_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[1/a Int[(e + f*x)^m*Tanh[
c + d*x]^n, x], x] - Simp[b/a Int[(e + f*x)^m*Sinh[c + d*x]*(Tanh[c + d*x]
)^(n - 1)/(a + b*Cosh[c + d*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] &&
IGtQ[m, 0] && IGtQ[n, 0]
```

rule 6129

```
Int[(((e_.) + (f_.)*(x_))^(m_.)*(F_)[(c_.) + (d_.)*(x_)]^(n_.)*(G_)[(c_.) +
(d_.)*(x_)]^(p_.))/(a_ + (b_.)*Sech[(c_.) + (d_.)*(x_)]), x_Symbol] := I
nt[(e + f*x)^m*Cosh[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Cosh[c + d*x]
)), x] /; FreeQ[{a, b, c, d, e, f}, x] && HyperbolicQ[F] && HyperbolicQ[G]
&& IntegersQ[m, n, p]
```

rule 6875

```
Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Sech[x]*T
anh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

## Maple [F]

$$\int \frac{\operatorname{arcsech}(bx + a)^2}{x} dx$$

input

```
int(arcsech(b*x+a)^2/x,x)
```

output

```
int(arcsech(b*x+a)^2/x,x)
```



**Fricas [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{arsech}(bx + a)^2}{x} dx$$

input `integrate(arcsech(b*x+a)^2/x,x, algorithm="fricas")`

output `integral(arcsech(b*x + a)^2/x, x)`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{asech}^2(a + bx)}{x} dx$$

input `integrate(asech(b*x+a)**2/x,x)`

output `Integral(asech(a + b*x)**2/x, x)`

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{arsech}(bx + a)^2}{x} dx$$

input `integrate(arcsech(b*x+a)^2/x,x, algorithm="maxima")`

output `integrate(arcsech(b*x + a)^2/x, x)`

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{arsech}(bx + a)^2}{x} dx$$

input `integrate(arcsech(b*x+a)^2/x,x, algorithm="giac")`

output `integrate(arcsech(b*x + a)^2/x, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)^2}{x} dx$$

input `int(acosh(1/(a + b*x))^2/x,x)`

output `int(acosh(1/(a + b*x))^2/x, x)`

**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{asech}(bx + a)^2}{x} dx$$

input `int(asech(b*x+a)^2/x,x)`

output `int(asech(a + b*x)**2/x,x)`

### 3.13 $\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx$

Optimal result	138
Mathematica [C] (warning: unable to verify)	139
Rubi [A] (verified)	140
Maple [A] (verified)	142
Fricas [F]	142
Sympy [F]	143
Maxima [F]	143
Giac [F]	144
Mupad [F(-1)]	144
Reduce [F]	144

#### Optimal result

Integrand size = 12, antiderivative size = 224

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx = -\frac{b\operatorname{sech}^{-1}(a+bx)^2}{a} - \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} + \frac{2b\operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{2b\operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{2b \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{2b \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}}$$

output

```
-b*arcsech(b*x+a)^2/a-arcsech(b*x+a)^2/x+2*b*arcsech(b*x+a)*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-2*b*arcsech(b*x+a)*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+2*b*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-2*b*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)
```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 1.72 (sec) , antiderivative size = 678, normalized size of antiderivative = 3.03

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^2} dx = \text{Too large to display}$$

input `Integrate[ArcSech[a + b*x]^2/x^2,x]`

output

```
(-(((a + b*x)*ArcSech[a + b*x]^2)/x) + (2*b*(2*ArcSech[a + b*x]*ArcTan[(-1 + a)*Coth[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]] - (2*I)*ArcCos[a^(-1)]*ArcTan[((1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]] + (ArcCos[a^(-1)] + 2*(ArcTan[(-1 + a)*Coth[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]] + ArcTan[(1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]))*Log[Sqrt[-1 + a^2]/(Sqrt[2]*Sqrt[a]*E^(ArcSech[a + b*x]/2)*Sqrt[-((b*x)/(a + b*x))]) + (ArcCos[a^(-1)] - 2*(ArcTan[(-1 + a)*Coth[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]] + ArcTan[((1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]))*Log[(Sqrt[-1 + a^2]*E^(ArcSech[a + b*x]/2))/(Sqrt[2]*Sqrt[a]*Sqrt[-((b*x)/(a + b*x))])] - (ArcCos[a^(-1)] + 2*ArcTan[((1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]])*Log[-(((1 + a)*(1 + a - I*Sqrt[-1 + a^2])*(-1 + Tanh[ArcSech[a + b*x]/2]))/(a*(-1 + a + I*Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2])))] - (ArcCos[a^(-1)] - 2*ArcTan[((1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]])*Log[(((1 + a)*(1 + a + I*Sqrt[-1 + a^2])*(1 + Tanh[ArcSech[a + b*x]/2]))/(a*(-1 + a + I*Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2])))] + I*(PolyLog[2, ((-1 - I*Sqrt[-1 + a^2])*(-1 + a - I*Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2]))/(a*(-1 + a + I*Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2])))] - PolyLog[2, ((I + Sqrt[-1 + a^2])*(-1 + a - I*Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2]))/(a*((-I)*(-1 + a) + Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2])))]/Sqrt[-1 + a^2])/a
```

**Rubi [A] (verified)**

Time = 0.74 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.01, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {6875, 5991, 3042, 4679, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx \\
 & \quad \downarrow \text{6875} \\
 & -b \int \frac{(a+bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1) \operatorname{sech}^{-1}(a+bx)^2}{b^2 x^2} d\operatorname{sech}^{-1}(a+bx) \\
 & \quad \downarrow \text{5991} \\
 & -b \left( 2 \int -\frac{\operatorname{sech}^{-1}(a+bx)}{bx} d\operatorname{sech}^{-1}(a+bx) + \frac{\operatorname{sech}^{-1}(a+bx)^2}{bx} \right) \\
 & \quad \downarrow \text{3042} \\
 & -b \left( \frac{\operatorname{sech}^{-1}(a+bx)^2}{bx} + 2 \int \frac{\operatorname{sech}^{-1}(a+bx)}{a - \operatorname{csc}\left(\operatorname{isech}^{-1}(a+bx) + \frac{\pi}{2}\right)} d\operatorname{sech}^{-1}(a+bx) \right) \\
 & \quad \downarrow \text{4679} \\
 & -b \left( 2 \int \left( \frac{\operatorname{sech}^{-1}(a+bx)}{a} + \frac{\operatorname{sech}^{-1}(a+bx)}{a \left(\frac{a}{a+bx} - 1\right)} \right) d\operatorname{sech}^{-1}(a+bx) + \frac{\operatorname{sech}^{-1}(a+bx)^2}{bx} \right) \\
 & \quad \downarrow \text{2009} \\
 & -b \left( 2 \left( -\frac{\operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{\operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a\sqrt{1-a^2}} - \frac{\operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \right) \right)
 \end{aligned}$$

input `Int[ArcSech[a + b*x]^2/x^2,x]`

output

```

-(b*(ArcSech[a + b*x]^2/(b*x) + 2*(ArcSech[a + b*x]^2/(2*a) - (ArcSech[a +
b*x]*Log[1 - (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2
]) + (ArcSech[a + b*x]*Log[1 - (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])
])/(a*Sqrt[1 - a^2]) - PolyLog[2, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2
])]/(a*Sqrt[1 - a^2]) + PolyLog[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2
])]/(a*Sqrt[1 - a^2])))

```

### Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4679

```
Int[(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(n_)*((c_) + (d_)*(x_))^(m_)
, x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Si
n[e + f*x])^n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGt
Q[m, 0]
```

rule 5991

```
Int[((e_) + (f_)*(x_))^(m_)*Sech[(c_) + (d_)*(x_)]*((a_) + (b_)*Sech[
(c_) + (d_)*(x_)]^(n_)*Tanh[(c_) + (d_)*(x_)], x_Symbol] := Simp[(-(e
+ f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b
*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

rule 6875

```
Int[((a_) + ArcSech[(c_) + (d_)*(x_)]*(b_))^(p_)*((e_) + (f_)*(x_))^(
m_), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Sech[x]*T
anh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

**Maple [A] (verified)**

Time = 0.49 (sec) , antiderivative size = 362, normalized size of antiderivative = 1.62

method	result
derivativedivides	$b \left( -\frac{(bx+a) \operatorname{arcsech}(bx+a)^2}{abx} + \frac{2\sqrt{-a^2+1} \operatorname{arcsech}(bx+a) \ln \left( \frac{-a \left( \frac{1}{bx+a} + \sqrt{\frac{1}{bx+a}-1} \sqrt{\frac{1}{bx+a}+1} \right) + \sqrt{-a^2+1}+1}{1+\sqrt{-a^2+1}} \right)}{a(a^2-1)} \right)$
default	$b \left( -\frac{(bx+a) \operatorname{arcsech}(bx+a)^2}{abx} + \frac{2\sqrt{-a^2+1} \operatorname{arcsech}(bx+a) \ln \left( \frac{-a \left( \frac{1}{bx+a} + \sqrt{\frac{1}{bx+a}-1} \sqrt{\frac{1}{bx+a}+1} \right) + \sqrt{-a^2+1}+1}{1+\sqrt{-a^2+1}} \right)}{a(a^2-1)} \right)$

input `int(arcsech(b*x+a)^2/x^2,x,method=_RETURNVERBOSE)`

output `b*(-(b*x+a)*arcsech(b*x+a)^2/a/b/x+2*(-a^2+1)^(1/2)/a/(a^2-1)*arcsech(b*x+a)*ln((-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))-2*(-a^2+1)^(1/2)/a/(a^2-1)*arcsech(b*x+a)*ln((a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2)))+2*(-a^2+1)^(1/2)/a/(a^2-1)*dilog((-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))-2*(-a^2+1)^(1/2)/a/(a^2-1)*dilog((a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2))))`

**Fricas [F]**

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx = \int \frac{\operatorname{arsech}(bx+a)^2}{x^2} dx$$

input `integrate(arcsech(b*x+a)^2/x^2,x, algorithm="fricas")`

output `integral(arcsech(b*x + a)^2/x^2, x)`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{arsech}^2(a + bx)}{x^2} dx$$

input `integrate(asech(b*x+a)**2/x**2,x)`

output `Integral(asech(a + b*x)**2/x**2, x)`

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{arsech}(bx + a)^2}{x^2} dx$$

input `integrate(arcsech(b*x+a)^2/x^2,x, algorithm="maxima")`

output `-log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^2/x - integrate(-2*(2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^2 + 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a)^2 + (b^3*x^3 + 2*a*b^2*x^2 + (a^2*b - b)*x - 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a) - ((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*log(b*x + a) - (2*b^3*x^3 + 4*a*b^2*x^2 + (2*a^2*b - b)*x - (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a))*sqrt(b*x + a + 1))*sqrt(-b*x - a + 1))*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a))/(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2 + (b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)), x)`



**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{ar} \operatorname{sech}(bx + a)^2}{x^2} dx$$

input `integrate(arcsech(b*x+a)^2/x^2,x, algorithm="giac")`

output `integrate(arcsech(b*x + a)^2/x^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)^2}{x^2} dx$$

input `int(acosh(1/(a + b*x))^2/x^2,x)`

output `int(acosh(1/(a + b*x))^2/x^2, x)`

**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{asech}(bx + a)^2}{x^2} dx$$

input `int(asech(b*x+a)^2/x^2,x)`

output `int(asech(a + b*x)**2/x**2,x)`

### 3.14 $\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^3} dx$

Optimal result	145
Mathematica [C] (warning: unable to verify)	146
Rubi [A] (verified)	147
Maple [A] (verified)	150
Fricas [F]	151
Sympy [F]	152
Maxima [F]	152
Giac [F]	153
Mupad [F(-1)]	153
Reduce [F]	153

#### Optimal result

Integrand size = 12, antiderivative size = 537

$$\begin{aligned}
 \int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^3} dx = & \frac{b^2 \sqrt{\frac{1-a-bx}{1+a+bx}} (1+a+bx) \operatorname{sech}^{-1}(a+bx)}{a(1-a^2)(a+bx)\left(1-\frac{a}{a+bx}\right)} + \frac{b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2} \\
 & - \frac{\operatorname{sech}^{-1}(a+bx)^2}{2x^2} + \frac{b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
 & - \frac{2b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
 & - \frac{b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
 & + \frac{2b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} + \frac{b^2 \log\left(\frac{x}{a+bx}\right)}{a^2(1-a^2)} \\
 & + \frac{b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} - \frac{2b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
 & - \frac{b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} + \frac{2b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}}
 \end{aligned}$$

output

```

b^2*((-b*x-a+1)/(b*x+a+1))^(1/2)*(b*x+a+1)*arcsech(b*x+a)/a/(-a^2+1)/(b*x+
a)/(1-a/(b*x+a))+1/2*b^2*arcsech(b*x+a)^2/a^2-1/2*arcsech(b*x+a)^2/x^2+b^2
*arcsech(b*x+a)*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))
/(1-(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(3/2)-2*b^2*arcsech(b*x+a)*ln(1-a*(1/(b*
x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a^2/(-a^
2+1)^(1/2)-b^2*arcsech(b*x+a)*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*
x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(3/2)+2*b^2*arcsech(b*x+a)
*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1
/2)))/a^2/(-a^2+1)^(1/2)+b^2*ln(x/(b*x+a))/a^2/(-a^2+1)+b^2*polylog(2,a*(1
/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a^2/
(-a^2+1)^(3/2)-2*b^2*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)
+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(1/2)-b^2*polylog(2,a*(1/(b*x+
a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a^2/(-a^2+
1)^(3/2)+2*b^2*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1
/2)))/(1+(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(1/2)

```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 4.84 (sec) , antiderivative size = 1439, normalized size of antiderivative = 2.68

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^3} dx = \text{Too large to display}$$

input

```
Integrate[ArcSech[a + b*x]^2/x^3,x]
```

output

```

-1/2*((a + b*x)^2*ArcSech[a + b*x]^2)/(a^2*x^2) + (b*ArcSech[a + b*x]*(-a
*sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(1 + a + b*x)) + (-1 + a^2)*(a + b*
x)*ArcSech[a + b*x])/((-1 + a)*a^2*(1 + a)*x) + (b^2*Log[(b*x)/(a + b*x)]
)/(a^2 - a^4) - (2*b^2*(2*ArcSech[a + b*x]*ArcTan[((-1 + a)*Coth[ArcSech[a
+ b*x]/2)]/sqrt[-1 + a^2]] - (2*I)*ArcCos[a^(-1)]*ArcTan[((1 + a)*Tanh[ArcS
ech[a + b*x]/2)]/sqrt[-1 + a^2]] + (ArcCos[a^(-1)] + 2*(ArcTan[((-1 + a)
*Coth[ArcSech[a + b*x]/2)]/sqrt[-1 + a^2]] + ArcTan[((1 + a)*Tanh[ArcSech[
a + b*x]/2)]/sqrt[-1 + a^2])))*Log[sqrt[-1 + a^2]/(sqrt[2]*sqrt[a]*E^(ArcS
ech[a + b*x]/2)*sqrt[-((b*x)/(a + b*x))]) + (ArcCos[a^(-1)] - 2*(ArcTan[(-
1 + a)*Coth[ArcSech[a + b*x]/2)]/sqrt[-1 + a^2]] + ArcTan[((1 + a)*Tanh[
ArcSech[a + b*x]/2)]/sqrt[-1 + a^2])))*Log[(sqrt[-1 + a^2]*E^(ArcSech[a +
b*x]/2))/(sqrt[2]*sqrt[a]*sqrt[-((b*x)/(a + b*x))])] - (ArcCos[a^(-1)] + 2
*ArcTan[((1 + a)*Tanh[ArcSech[a + b*x]/2)]/sqrt[-1 + a^2])))*Log[-(((-1 + a
)*(1 + a - I*sqrt[-1 + a^2]))*(-1 + Tanh[ArcSech[a + b*x]/2]))/(a*(-1 + a +
I*sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2]))] - (ArcCos[a^(-1)] - 2*ArcTan
[((1 + a)*Tanh[ArcSech[a + b*x]/2)]/sqrt[-1 + a^2])))*Log[((-1 + a)*(1 + a
+ I*sqrt[-1 + a^2]))*(1 + Tanh[ArcSech[a + b*x]/2]))/(a*(-1 + a + I*sqrt[-
1 + a^2]*Tanh[ArcSech[a + b*x]/2]))] + I*(PolyLog[2, ((-1 - I*sqrt[-1 + a^
2]))*(-1 + a - I*sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2]))/(a*(-1 + a + I*S
qrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2]))] - PolyLog[2, ((I + sqrt[-1 + ...

```

## Rubi [A] (verified)

Time = 1.18 (sec) , antiderivative size = 517, normalized size of antiderivative = 0.96, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6875, 25, 5991, 3042, 4679, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^3} dx$$

$$\downarrow \text{6875}$$

$$-b^2 \int \frac{(a + bx) \sqrt{\frac{-a - bx + 1}{a + bx + 1}} (a + bx + 1) \operatorname{sech}^{-1}(a + bx)^2}{b^3 x^3} d \operatorname{sech}^{-1}(a + bx)$$

$$\downarrow \text{25}$$

$$\begin{aligned}
& b^2 \int -\frac{(a+bx)\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^2}{b^3 x^3} d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow \text{5991} \\
& -b^2 \left( \frac{\operatorname{sech}^{-1}(a+bx)^2}{2b^2 x^2} - \int \frac{\operatorname{sech}^{-1}(a+bx)}{b^2 x^2} d\operatorname{sech}^{-1}(a+bx) \right) \\
& \quad \downarrow \text{3042} \\
& -b^2 \left( \frac{\operatorname{sech}^{-1}(a+bx)^2}{2b^2 x^2} - \int \frac{\operatorname{sech}^{-1}(a+bx)}{(a - \csc(i\operatorname{sech}^{-1}(a+bx) + \frac{\pi}{2}))^2} d\operatorname{sech}^{-1}(a+bx) \right) \\
& \quad \downarrow \text{4679} \\
& -b^2 \left( \frac{\operatorname{sech}^{-1}(a+bx)^2}{2b^2 x^2} - \int \left( \frac{2\operatorname{sech}^{-1}(a+bx)}{a^2 \left(\frac{a}{a+bx} - 1\right)} + \frac{\operatorname{sech}^{-1}(a+bx)}{a^2} + \frac{\operatorname{sech}^{-1}(a+bx)}{a^2 \left(\frac{a}{a+bx} - 1\right)^2} \right) d\operatorname{sech}^{-1}(a+bx) \right) \\
& \quad \downarrow \text{2009} \\
& -b^2 \left( \frac{2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} - \frac{\operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 (1-a^2)^{3/2}} - \frac{2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a^2 \sqrt{1-a^2}} + \frac{\operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a^2 (1-a^2)^{3/2}} \right)
\end{aligned}$$

input `Int[ArcSech[a + b*x]^2/x^3,x]`

output

```

-(b^2*(-((Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x)*ArcSech[a + b*x]
)/(a*(1 - a^2)*(a + b*x)*(1 - a/(a + b*x)))) - ArcSech[a + b*x]^2/(2*a^2)
+ ArcSech[a + b*x]^2/(2*b^2*x^2) - (ArcSech[a + b*x]*Log[1 - (a*E^ArcSech[
a + b*x)]/(1 - Sqrt[1 - a^2])])/(a^2*(1 - a^2)^(3/2)) + (2*ArcSech[a + b*x]
]*Log[1 - (a*E^ArcSech[a + b*x)]/(1 - Sqrt[1 - a^2])])/(a^2*Sqrt[1 - a^2])
+ (ArcSech[a + b*x]*Log[1 - (a*E^ArcSech[a + b*x)]/(1 + Sqrt[1 - a^2])])/(
a^2*(1 - a^2)^(3/2)) - (2*ArcSech[a + b*x]*Log[1 - (a*E^ArcSech[a + b*x]
)/(1 + Sqrt[1 - a^2])])/(a^2*Sqrt[1 - a^2]) - Log[1 - a/(a + b*x)]/(a^2*(1
- a^2)) - PolyLog[2, (a*E^ArcSech[a + b*x)]/(1 - Sqrt[1 - a^2])]/(a^2*(1 -
a^2)^(3/2)) + (2*PolyLog[2, (a*E^ArcSech[a + b*x)]/(1 - Sqrt[1 - a^2])])/(
a^2*Sqrt[1 - a^2]) + PolyLog[2, (a*E^ArcSech[a + b*x)]/(1 + Sqrt[1 - a^2]
)]/(a^2*(1 - a^2)^(3/2)) - (2*PolyLog[2, (a*E^ArcSech[a + b*x)]/(1 + Sqrt[
1 - a^2])])/(a^2*Sqrt[1 - a^2]))

```

### Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4679

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)
, x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Si
n[e + f*x])^n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGt
Q[m, 0]
```

rule 5991

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_) + (b_.)*Sech[
(c_.) + (d_.)*(x_)]^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-(e
+ f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b
*d*(n + 1)) Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

rule 6875

```
Int[((a_.) + ArcSech[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] :> Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Sech[x]*T
anh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

**Maple [A] (verified)**

Time = 0.68 (sec) , antiderivative size = 982, normalized size of antiderivative = 1.83

method	result
derivativedivides	$b^2 \left( -\frac{\operatorname{arcsech}(bx+a) \left( -2\sqrt{\frac{bx+a+1}{bx+a}} \sqrt{-\frac{bx+a-1}{bx+a}} a^2 (bx+a) + 2\sqrt{\frac{bx+a+1}{bx+a}} \sqrt{-\frac{bx+a-1}{bx+a}} a (bx+a)^2 + 2 \operatorname{arcsech}(bx+a) a^3 \right)}{2a^2} \right)$
default	$b^2 \left( -\frac{\operatorname{arcsech}(bx+a) \left( -2\sqrt{\frac{bx+a+1}{bx+a}} \sqrt{-\frac{bx+a-1}{bx+a}} a^2 (bx+a) + 2\sqrt{\frac{bx+a+1}{bx+a}} \sqrt{-\frac{bx+a-1}{bx+a}} a (bx+a)^2 + 2 \operatorname{arcsech}(bx+a) a^3 \right)}{2a^2} \right)$

input

```
int(arcsech(b*x+a)^2/x^3,x,method=_RETURNVERBOSE)
```

output

```

b^2*(-1/2*arcsech(b*x+a)*(-2*((b*x+a+1)/(b*x+a))^(1/2)*(-(b*x+a-1)/(b*x+a)
)^(1/2)*a^2*(b*x+a)+2*((b*x+a+1)/(b*x+a))^(1/2)*(-(b*x+a-1)/(b*x+a))^(1/2)
)*a*(b*x+a)^2+2*arcsech(b*x+a)*a^3*(b*x+a)-arcsech(b*x+a)*a^2*(b*x+a)^2-2*a
rcsech(b*x+a)*a*(b*x+a)+arcsech(b*x+a)*(b*x+a)^2+2*a^2-4*(b*x+a)*a+2*(b*x+
a)^2)/a^2/(a^2-1)/b^2/x^2-1/a^2/(a^2-1)*ln(a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)
)*(1/(b*x+a)+1)^(1/2))^2+a-2/(b*x+a)-2*(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(
1/2))+2/a^2/(a^2-1)*ln(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))+
(-a^2+1)^(1/2)/a^2/(a^2-1)^2*arcsech(b*x+a)*ln((-a*(1/(b*x+a)+(1/(b*x+a)-1)
)^(1/2)*(1/(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))-(-a^2+1
)^(1/2)/a^2/(a^2-1)^2*arcsech(b*x+a)*ln((a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*
(1/(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2)))+(-a^2+1)^(1/2)
/a^2/(a^2-1)^2*dilog((-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)
))+(-a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))-(-a^2+1)^(1/2)/a^2/(a^2-1)^2*dilo
g((a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)-1)
/(-1+(-a^2+1)^(1/2)))-2*(-a^2+1)^(1/2)/(a^2-1)^2*arcsech(b*x+a)*ln((-a*(1/
(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)+1)/(1+(-a^
2+1)^(1/2)))+2*(-a^2+1)^(1/2)/(a^2-1)^2*arcsech(b*x+a)*ln((a*(1/(b*x+a)+(1
/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2)
)))-2*(-a^2+1)^(1/2)/(a^2-1)^2*dilog((-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1
/(b*x+a)+1)^(1/2))+(-a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))+2*(-a^2+1)^(1/...

```

**Fricas [F]**

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^3} dx = \int \frac{\operatorname{ar} \operatorname{sech}(bx+a)^2}{x^3} dx$$

input

```
integrate(arcsech(b*x+a)^2/x^3,x, algorithm="fricas")
```

output

```
integral(arcsech(b*x + a)^2/x^3, x)
```



**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{asech}^2(a + bx)}{x^3} dx$$

input `integrate(asech(b*x+a)**2/x**3,x)`

output `Integral(asech(a + b*x)**2/x**3, x)`

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{arsech}(bx + a)^2}{x^3} dx$$

input `integrate(arcsech(b*x+a)^2/x^3,x, algorithm="maxima")`

output `-1/2*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^2/x^2 - integrate(-(4*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^2 + 4*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a)^2 + (b^3*x^3 + 2*a*b^2*x^2 + (a^2*b - b)*x - 4*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a) - (2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*log(b*x + a) - (2*b^3*x^3 + 4*a*b^2*x^2 + (2*a^2*b - b)*x - 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a))*sqrt(b*x + a + 1))*sqrt(-b*x - a + 1))*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a))/(b^3*x^6 + 3*a*b^2*x^5 + (3*a^2*b - b)*x^4 + (a^3 - a)*x^3 + (b^3*x^6 + 3*a*b^2*x^5 + (3*a^2*b - b)*x^4 + (a^3 - a)*x^3)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)), x)`

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{ar} \operatorname{sech}(bx + a)^2}{x^3} dx$$

input `integrate(arcsech(b*x+a)^2/x^3,x, algorithm="giac")`

output `integrate(arcsech(b*x + a)^2/x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)^2}{x^3} dx$$

input `int(acosh(1/(a + b*x))^2/x^3,x)`

output `int(acosh(1/(a + b*x))^2/x^3, x)`

**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{asech}(bx + a)^2}{x^3} dx$$

input `int(asech(b*x+a)^2/x^3,x)`

output `int(asech(a + b*x)**2/x**3,x)`

### 3.15 $\int x \operatorname{sech}^{-1}(a + bx)^3 dx$

Optimal result	154
Mathematica [A] (warning: unable to verify)	155
Rubi [A] (verified)	156
Maple [F]	158
Fricas [F]	158
Sympy [F]	158
Maxima [F]	159
Giac [F]	159
Mupad [F(-1)]	160
Reduce [F]	160

#### Optimal result

Integrand size = 10, antiderivative size = 260

$$\begin{aligned}
 \int x \operatorname{sech}^{-1}(a + bx)^3 dx = & -\frac{3 \operatorname{sech}^{-1}(a + bx)^2}{2b^2} - \frac{3 \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx) \operatorname{sech}^{-1}(a + bx)^2}{2b^2} \\
 & - \frac{a^2 \operatorname{sech}^{-1}(a + bx)^3}{2b^2} + \frac{1}{2} x^2 \operatorname{sech}^{-1}(a + bx)^3 \\
 & + \frac{6a \operatorname{sech}^{-1}(a + bx)^2 \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
 & + \frac{3 \operatorname{sech}^{-1}(a + bx) \log\left(1 + e^{2 \operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
 & - \frac{6ia \operatorname{sech}^{-1}(a + bx) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
 & + \frac{6ia \operatorname{sech}^{-1}(a + bx) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
 & + \frac{3 \operatorname{PolyLog}\left(2, -e^{2 \operatorname{sech}^{-1}(a+bx)}\right)}{2b^2} \\
 & + \frac{6ia \operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
 & - \frac{6ia \operatorname{PolyLog}\left(3, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2}
 \end{aligned}$$

output

```
-3/2*arcsech(b*x+a)^2/b^2-3/2*((-b*x-a+1)/(b*x+a+1))^(1/2)*(b*x+a+1)*arcsech(b*x+a)^2/b^2-1/2*a^2*arcsech(b*x+a)^3/b^2+1/2*x^2*arcsech(b*x+a)^3+6*a*arcsech(b*x+a)^2*arctan(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/b^2+3*arcsech(b*x+a)*ln(1+(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))^2)/b^2-6*I*a*arcsech(b*x+a)*polylog(2,-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b^2+6*I*a*arcsech(b*x+a)*polylog(2,I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b^2+3/2*polylog(2,-(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))^2)/b^2+6*I*a*polylog(3,-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b^2-6*I*a*polylog(3,I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b^2
```

**Mathematica [A] (warning: unable to verify)**

Time = 0.40 (sec) , antiderivative size = 254, normalized size of antiderivative = 0.98

$$\int x \operatorname{sech}^{-1}(a + bx)^3 dx$$

$$= -3\sqrt{-\frac{-1+a+bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)^2 - 2a(a+bx)\operatorname{sech}^{-1}(a+bx)^3 + (a+bx)^2\operatorname{sech}^{-1}(a+bx)^3 +$$

input

```
Integrate[x*ArcSech[a + b*x]^3,x]
```

output

```
(-3*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(1 + a + b*x)*ArcSech[a + b*x]^2 - 2*a*(a + b*x)*ArcSech[a + b*x]^3 + (a + b*x)^2*ArcSech[a + b*x]^3 + 3*ArcSech[a + b*x]*(ArcSech[a + b*x] + 2*Log[1 + E^(-2*ArcSech[a + b*x])]) - 3*PolyLog[2, -E^(-2*ArcSech[a + b*x])] + (6*I)*a*(-(ArcSech[a + b*x]^2*(Log[1 - I/E^ArcSech[a + b*x]] - Log[1 + I/E^ArcSech[a + b*x]]) - 2*ArcSech[a + b*x]*(PolyLog[2, (-I)/E^ArcSech[a + b*x]] - PolyLog[2, I/E^ArcSech[a + b*x]]) - 2*PolyLog[3, (-I)/E^ArcSech[a + b*x]] + 2*PolyLog[3, I/E^ArcSech[a + b*x]]))/(2*b^2)
```

**Rubi [A] (verified)**

Time = 0.57 (sec) , antiderivative size = 234, normalized size of antiderivative = 0.90, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {6875, 25, 5991, 3042, 4678, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \operatorname{sech}^{-1}(a + bx)^3 dx \\
 & \quad \downarrow \text{6875} \\
 & - \frac{\int bx(a + bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a + bx + 1) \operatorname{sech}^{-1}(a + bx)^3 d \operatorname{sech}^{-1}(a + bx)}{b^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int -bx(a + bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a + bx + 1) \operatorname{sech}^{-1}(a + bx)^3 d \operatorname{sech}^{-1}(a + bx)}{b^2} \\
 & \quad \downarrow \text{5991} \\
 & - \frac{\frac{3}{2} \int b^2 x^2 \operatorname{sech}^{-1}(a + bx)^2 d \operatorname{sech}^{-1}(a + bx) - \frac{1}{2} b^2 x^2 \operatorname{sech}^{-1}(a + bx)^3}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{-\frac{1}{2} b^2 x^2 \operatorname{sech}^{-1}(a + bx)^3 + \frac{3}{2} \int \operatorname{sech}^{-1}(a + bx)^2 (a - \csc(i \operatorname{sech}^{-1}(a + bx) + \frac{\pi}{2}))^2 d \operatorname{sech}^{-1}(a + bx)}{b^2} \\
 & \quad \downarrow \text{4678} \\
 & - \frac{\frac{3}{2} \int (a^2 \operatorname{sech}^{-1}(a + bx)^2 + (a + bx)^2 \operatorname{sech}^{-1}(a + bx)^2 - 2a(a + bx) \operatorname{sech}^{-1}(a + bx)^2) d \operatorname{sech}^{-1}(a + bx) - \frac{1}{2} b^2 x^2 \operatorname{sech}^{-1}(a + bx)^3}{b^2} \\
 & \quad \downarrow \text{2009} \\
 & - \frac{-\frac{1}{2} b^2 x^2 \operatorname{sech}^{-1}(a + bx)^3 + \frac{3}{2} \left( \frac{1}{3} a^2 \operatorname{sech}^{-1}(a + bx)^3 - 4a \operatorname{sech}^{-1}(a + bx)^2 \arctan \left( e^{\operatorname{sech}^{-1}(a + bx)} \right) + 4i a \operatorname{sech}^{-1}(a + bx) \right)}{b^2}
 \end{aligned}$$

input

Int[x\*ArcSech[a + b\*x]^3,x]

output

```

-((-1/2*(b^2*x^2*ArcSech[a + b*x]^3) + (3*(ArcSech[a + b*x]^2 + Sqrt[(1 -
a - b*x)/(1 + a + b*x)]*(1 + a + b*x)*ArcSech[a + b*x]^2 + (a^2*ArcSech[a
+ b*x]^3)/3 - 4*a*ArcSech[a + b*x]^2*ArcTan[E^ArcSech[a + b*x]] - 2*ArcSec
h[a + b*x]*Log[1 + E^(2*ArcSech[a + b*x])] + (4*I)*a*ArcSech[a + b*x]*Poly
Log[2, (-I)*E^ArcSech[a + b*x]] - (4*I)*a*ArcSech[a + b*x]*PolyLog[2, I*E^
ArcSech[a + b*x]] - PolyLog[2, -E^(2*ArcSech[a + b*x])] - (4*I)*a*PolyLog[
3, (-I)*E^ArcSech[a + b*x]] + (4*I)*a*PolyLog[3, I*E^ArcSech[a + b*x]]))/2
)/b^2)

```

### Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4678

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)
, x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

rule 5991

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sech[
(c_.) + (d_.)*(x_)]^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-(e
+ f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b
*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

rule 6875

```
Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Sech[x]*T
anh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

**Maple [F]**

$$\int x \operatorname{arcsech}(bx + a)^3 dx$$

input `int(x*arcsech(b*x+a)^3,x)`

output `int(x*arcsech(b*x+a)^3,x)`

**Fricas [F]**

$$\int x \operatorname{sech}^{-1}(a + bx)^3 dx = \int x \operatorname{ar} \operatorname{sech}(bx + a)^3 dx$$

input `integrate(x*arcsech(b*x+a)^3,x, algorithm="fricas")`

output `integral(x*arcsech(b*x + a)^3, x)`

**Sympy [F]**

$$\int x \operatorname{sech}^{-1}(a + bx)^3 dx = \int x \operatorname{asech}^3(a + bx) dx$$

input `integrate(x*asech(b*x+a)**3,x)`

output `Integral(x*asech(a + b*x)**3, x)`

**Maxima [F]**

$$\int x \operatorname{sech}^{-1}(a + bx)^3 dx = \int x \operatorname{arsech}(bx + a)^3 dx$$

input `integrate(x*arcsech(b*x+a)^3,x, algorithm="maxima")`

output

```
1/2*x^2*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*s
qrt(-b*x - a + 1)*a + b*x + a)^3 - integrate(1/2*(16*(b^3*x^4 + 3*a*b^2*x^
3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*
log(b*x + a)^3 + 16*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)
*x)*log(b*x + a)^3 + 3*(b^3*x^4 + 2*a*b^2*x^3 + (a^2*b - b)*x^2 + 4*(b^3*x
^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*log(b*x + a) + (2*(b^3
*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*sqrt(b*x + a + 1)*lo
g(b*x + a) + (2*b^3*x^4 + 4*a*b^2*x^3 + (2*a^2*b - b)*x^2 + 2*(b^3*x^4 + 3
*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*log(b*x + a))*sqrt(b*x + a +
1))*sqrt(-b*x - a + 1))*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sq
rt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^2 - 24*((b^3*x^4 + 3*a*b^2
*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*sqrt(b*x + a + 1)*sqrt(-b*x - a +
1)*log(b*x + a)^2 + (b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)
*x)*log(b*x + a)^2)*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*
x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 +
(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqr
t(-b*x - a + 1) + (3*a^2*b - b)*x - a), x)
```

**Giac [F]**

$$\int x \operatorname{sech}^{-1}(a + bx)^3 dx = \int x \operatorname{arsech}(bx + a)^3 dx$$

input `integrate(x*arcsech(b*x+a)^3,x, algorithm="giac")`

output `integrate(x*arcsech(b*x + a)^3, x)`



**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{sech}^{-1}(a + bx)^3 dx = \int x \operatorname{acosh}\left(\frac{1}{a + bx}\right)^3 dx$$

input `int(x*acosh(1/(a + b*x))^3,x)`output `int(x*acosh(1/(a + b*x))^3, x)`**Reduce [F]**

$$\int x \operatorname{sech}^{-1}(a + bx)^3 dx = \int \operatorname{asech}(bx + a)^3 x dx$$

input `int(x*asech(b*x+a)^3,x)`output `int(asech(a + b*x)**3*x,x)`

### 3.16 $\int \operatorname{sech}^{-1}(a + bx)^3 dx$

Optimal result	161
Mathematica [A] (verified)	162
Rubi [A] (verified)	162
Maple [F]	165
Fricas [F]	165
Sympy [F]	166
Maxima [F]	166
Giac [F]	167
Mupad [F(-1)]	167
Reduce [F]	167

#### Optimal result

Integrand size = 8, antiderivative size = 136

$$\int \operatorname{sech}^{-1}(a + bx)^3 dx = \frac{(a + bx)\operatorname{sech}^{-1}(a + bx)^3}{b} - \frac{6\operatorname{sech}^{-1}(a + bx)^2 \arctan\left(e^{\operatorname{sech}^{-1}(a + bx)}\right)}{b} + \frac{6i\operatorname{sech}^{-1}(a + bx) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b} - \frac{6i\operatorname{sech}^{-1}(a + bx) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b} - \frac{6i \operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b} + \frac{6i \operatorname{PolyLog}\left(3, ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b}$$

output

```
(b*x+a)*arcsech(b*x+a)^3/b-6*arcsech(b*x+a)^2*arctan(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/b+6*I*arcsech(b*x+a)*polylog(2,-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b-6*I*arcsech(b*x+a)*polylog(2,I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b-6*I*polylog(3,-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b+6*I*polylog(3,I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b
```

**Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.12

$$\int \operatorname{sech}^{-1}(a + bx)^3 dx = \frac{(a + bx)\operatorname{sech}^{-1}(a + bx)^3}{b} - 3i \left( -\operatorname{sech}^{-1}(a + bx)^2 \left( \log \left( 1 - ie^{-\operatorname{sech}^{-1}(a+bx)} \right) - \log \left( 1 + ie^{-\operatorname{sech}^{-1}(a+bx)} \right) \right) - 2\operatorname{sech}^{-1}(a + bx) \left( \operatorname{PolyLog} \right) \right)$$

input `Integrate[ArcSech[a + b*x]^3,x]`

output

```
((a + b*x)*ArcSech[a + b*x]^3)/b - ((3*I)*(-(ArcSech[a + b*x]^2*(Log[1 - I/E^ArcSech[a + b*x]] - Log[1 + I/E^ArcSech[a + b*x]])) - 2*ArcSech[a + b*x]*PolyLog[2, (-I)/E^ArcSech[a + b*x]] - PolyLog[2, I/E^ArcSech[a + b*x]])) - 2*(PolyLog[3, (-I)/E^ArcSech[a + b*x]] - PolyLog[3, I/E^ArcSech[a + b*x]])))/b
```

**Rubi [A] (verified)**

Time = 0.58 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.92, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {6869, 6833, 5941, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{sech}^{-1}(a + bx)^3 dx$$

$$\downarrow \text{6869}$$

$$\frac{\int \operatorname{sech}^{-1}(a + bx)^3 d(a + bx)}{b}$$

$$\downarrow \text{6833}$$

$$\frac{\int (a + bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a + bx + 1) \operatorname{sech}^{-1}(a + bx)^3 d\operatorname{sech}^{-1}(a + bx)}{b}$$

$$\downarrow \text{5941}$$

$$\frac{3 \int (a + bx) \operatorname{sech}^{-1}(a + bx)^2 d \operatorname{sech}^{-1}(a + bx) - (a + bx) \operatorname{sech}^{-1}(a + bx)^3}{b}$$

↓ 3042

$$\frac{-(a + bx) \operatorname{sech}^{-1}(a + bx)^3 + 3 \int \operatorname{sech}^{-1}(a + bx)^2 \csc\left(i \operatorname{sech}^{-1}(a + bx) + \frac{\pi}{2}\right) d \operatorname{sech}^{-1}(a + bx)}{b}$$

↓ 4668

$$\frac{-(a + bx) \operatorname{sech}^{-1}(a + bx)^3 + 3 \left( -2i \int \operatorname{sech}^{-1}(a + bx) \log\left(1 - i e^{\operatorname{sech}^{-1}(a + bx)}\right) d \operatorname{sech}^{-1}(a + bx) + 2i \int \operatorname{sech}^{-1}(a + bx) \operatorname{PolyLog}\left(2, -i e^{\operatorname{sech}^{-1}(a + bx)}\right) d \operatorname{sech}^{-1}(a + bx) - \operatorname{sech}^{-1}(a + bx) \operatorname{PolyLog}\left(2, -i e^{\operatorname{sech}^{-1}(a + bx)}\right) \right)}{b}$$

↓ 3011

$$\frac{-(a + bx) \operatorname{sech}^{-1}(a + bx)^3 + 3 \left( 2i \left( \int \operatorname{PolyLog}\left(2, -i e^{\operatorname{sech}^{-1}(a + bx)}\right) d \operatorname{sech}^{-1}(a + bx) - \operatorname{sech}^{-1}(a + bx) \operatorname{PolyLog}\left(2, -i e^{\operatorname{sech}^{-1}(a + bx)}\right) \right) \right)}{b}$$

↓ 2720

$$\frac{-(a + bx) \operatorname{sech}^{-1}(a + bx)^3 + 3 \left( 2i \left( \int e^{-\operatorname{sech}^{-1}(a + bx)} \operatorname{PolyLog}\left(2, -i e^{\operatorname{sech}^{-1}(a + bx)}\right) d e^{\operatorname{sech}^{-1}(a + bx)} - \operatorname{sech}^{-1}(a + bx) \operatorname{PolyLog}\left(2, -i e^{\operatorname{sech}^{-1}(a + bx)}\right) \right) \right)}{b}$$

↓ 7143

$$\frac{-(a + bx) \operatorname{sech}^{-1}(a + bx)^3 + 3 \left( 2 \operatorname{sech}^{-1}(a + bx)^2 \arctan\left(e^{\operatorname{sech}^{-1}(a + bx)}\right) + 2i \left( \operatorname{PolyLog}\left(3, -i e^{\operatorname{sech}^{-1}(a + bx)}\right) - \operatorname{sech}^{-1}(a + bx) \operatorname{PolyLog}\left(2, -i e^{\operatorname{sech}^{-1}(a + bx)}\right) \right) \right)}{b}$$

input `Int[ArcSech[a + b*x]^3, x]`

output `-((-((a + b*x)*ArcSech[a + b*x]^3) + 3*(2*ArcSech[a + b*x]^2*ArcTan[E^ArcSech[a + b*x]] + (2*I)*(-(ArcSech[a + b*x]*PolyLog[2, (-I)*E^ArcSech[a + b*x]]) + PolyLog[3, (-I)*E^ArcSech[a + b*x]]) - (2*I)*(-(ArcSech[a + b*x]*PolyLog[2, I*E^ArcSech[a + b*x]]) + PolyLog[3, I*E^ArcSech[a + b*x]])))/b)`

## Definitions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^(c_)*((a_) + (b_)*(x_)))^(n_)]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 5941 `Int[(x_)^(m_)*Sech[(a_) + (b_)*(x_)^(n_)]^(p_)*Tanh[(a_) + (b_)*(x_)^(n_)]^(q_), x_Symbol] := Simp[(-x^(m - n + 1))*(Sech[a + b*x^n]^p/(b*n*p)), x] + Simp[(m - n + 1)/(b*n*p) Int[x^(m - n)*Sech[a + b*x^n]^p, x], x] /; FreeQ[{a, b, p}, x] && RationalQ[m] && IntegerQ[n] && GeQ[m - n, 0] && EqQ[q, 1]`

rule 6833 `Int[((a_) + ArcSech[(c_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[-c^(-1) Subst[Int[(a + b*x)^n*Sech[x]*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[n, 0]`

rule 6869 `Int[((a_.) + ArcSech[(c_) + (d_.)*(x_)]*(b_.))^(p_.), x_Symbol] := Simp[1/d  
Subst[Int[(a + b*ArcSech[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d  
, x] && IGtQ[p, 0]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S  
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d  
, e, n, p}, x] && EqQ[b*d, a*e]`

### Maple [F]

$$\int \operatorname{arcsech}(bx + a)^3 dx$$

input `int(arcsech(b*x+a)^3,x)`

output `int(arcsech(b*x+a)^3,x)`

### Fricas [F]

$$\int \operatorname{sech}^{-1}(a + bx)^3 dx = \int \operatorname{arsech}(bx + a)^3 dx$$

input `integrate(arcsech(b*x+a)^3,x, algorithm="fricas")`

output `integral(arcsech(b*x + a)^3, x)`

**Sympy [F]**

$$\int \operatorname{sech}^{-1}(a + bx)^3 dx = \int \operatorname{asech}^3(a + bx) dx$$

input `integrate(asech(b*x+a)**3,x)`

output `Integral(asech(a + b*x)**3, x)`

**Maxima [F]**

$$\int \operatorname{sech}^{-1}(a + bx)^3 dx = \int \operatorname{arsech}(bx + a)^3 dx$$

input `integrate(arcsech(b*x+a)^3,x, algorithm="maxima")`

output

```
x*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^3 - integrate((8*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^3 + 8*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a)^3 + 3*(b^3*x^3 + 2*a*b^2*x^2 + (a^2*b - b)*x + 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a) + ((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*log(b*x + a) + (2*b^3*x^3 + 4*a*b^2*x^2 + (2*a^2*b - b)*x + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a))*sqrt(b*x + a + 1))*sqrt(-b*x - a + 1))*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^2 - 12*((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1))*sqrt(-b*x - a + 1)*log(b*x + a)^2 + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a)^2)*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1))*sqrt(-b*x - a + 1) + (3*a^2*b - b)*x - a), x)
```

**Giac [F]**

$$\int \operatorname{sech}^{-1}(a + bx)^3 dx = \int \operatorname{arsech}(bx + a)^3 dx$$

input `integrate(arcsech(b*x+a)^3,x, algorithm="giac")`

output `integrate(arcsech(b*x + a)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{sech}^{-1}(a + bx)^3 dx = \int \operatorname{acosh}\left(\frac{1}{a + bx}\right)^3 dx$$

input `int(acosh(1/(a + b*x))^3,x)`

output `int(acosh(1/(a + b*x))^3, x)`

**Reduce [F]**

$$\int \operatorname{sech}^{-1}(a + bx)^3 dx = \int \operatorname{asech}(bx + a)^3 dx$$

input `int(asech(b*x+a)^3,x)`

output `int(asech(a + b*x)**3,x)`



$$3.17 \quad \int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx$$

Optimal result	169
Mathematica [A] (verified)	171
Rubi [C] (verified)	172
Maple [F]	178
Fricas [F]	178
Sympy [F]	178
Maxima [F]	179
Giac [F]	179
Mupad [F(-1)]	179
Reduce [F]	180

**Optimal result**

Integrand size = 12, antiderivative size = 378

$$\begin{aligned}
\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx &= \operatorname{sech}^{-1}(a+bx)^3 \log \left( 1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
&+ \operatorname{sech}^{-1}(a+bx)^3 \log \left( 1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&- \operatorname{sech}^{-1}(a+bx)^3 \log \left( 1 + e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
&+ 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog} \left( 2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
&+ 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog} \left( 2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&- \frac{3}{2}\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog} \left( 2, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
&- 6\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left( 3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
&- 6\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left( 3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&+ \frac{3}{2}\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left( 3, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
&+ 6 \operatorname{PolyLog} \left( 4, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
&+ 6 \operatorname{PolyLog} \left( 4, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&- \frac{3}{4} \operatorname{PolyLog} \left( 4, -e^{2\operatorname{sech}^{-1}(a+bx)} \right)
\end{aligned}$$

output

```

arcsech(b*x+a)^3*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)
)/(1-(-a^2+1)^(1/2)))+arcsech(b*x+a)^3*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)
)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2))-arcsech(b*x+a)^3*ln(1+(1/(b*x+
a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)+3*arcsech(b*x+a)^2*polylog(
2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2))
)+3*arcsech(b*x+a)^2*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)
+1)^(1/2))/(1+(-a^2+1)^(1/2)))-3/2*arcsech(b*x+a)^2*polylog(2,-(1/(b*x+a)+
(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)-6*arcsech(b*x+a)*polylog(3,a*(
1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))-6*a
rcsech(b*x+a)*polylog(3,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/
2)))/(1+(-a^2+1)^(1/2)))+3/2*arcsech(b*x+a)*polylog(3,-(1/(b*x+a)+(1/(b*x+a)
)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)+6*polylog(4,a*(1/(b*x+a)+(1/(b*x+a)-1)^(
1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))+6*polylog(4,a*(1/(b*x+a)+(1
/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))-3/4*polylog(4,-
(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)

```

**Mathematica [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 384, normalized size of antiderivative = 1.02

$$\begin{aligned}
\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx = & -\frac{1}{2} \operatorname{sech}^{-1}(a+bx)^4 - \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 + e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\
& + \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 + \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}}\right) \\
& + \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
& + \frac{3}{2} \operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, -e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\
& + 3 \operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, -\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}}\right) \\
& + 3 \operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
& + \frac{3}{2} \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, -e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\
& - 6 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, -\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}}\right) \\
& - 6 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
& + \frac{3}{4} \operatorname{PolyLog}\left(4, -e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\
& + 6 \operatorname{PolyLog}\left(4, -\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}}\right) \\
& + 6 \operatorname{PolyLog}\left(4, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right)
\end{aligned}$$

input `Integrate[ArcSech[a + b*x]^3/x, x]`

output

```

-1/2*ArcSech[a + b*x]^4 - ArcSech[a + b*x]^3*Log[1 + E^(-2*ArcSech[a + b*x
])] + ArcSech[a + b*x]^3*Log[1 + (a*E^ArcSech[a + b*x])/(-1 + Sqrt[1 - a^2
])] + ArcSech[a + b*x]^3*Log[1 - (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2
])] + (3*ArcSech[a + b*x]^2*PolyLog[2, -E^(-2*ArcSech[a + b*x])])/2 + 3*Arc
Sech[a + b*x]^2*PolyLog[2, -((a*E^ArcSech[a + b*x])/(-1 + Sqrt[1 - a^2]))]
+ 3*ArcSech[a + b*x]^2*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^
2])] + (3*ArcSech[a + b*x]*PolyLog[3, -E^(-2*ArcSech[a + b*x])])/2 - 6*Arc
Sech[a + b*x]*PolyLog[3, -((a*E^ArcSech[a + b*x])/(-1 + Sqrt[1 - a^2]))] -
6*ArcSech[a + b*x]*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])]
+ (3*PolyLog[4, -E^(-2*ArcSech[a + b*x])])/4 + 6*PolyLog[4, -((a*E^ArcSec
h[a + b*x])/(-1 + Sqrt[1 - a^2]))] + 6*PolyLog[4, (a*E^ArcSech[a + b*x])/(
1 + Sqrt[1 - a^2])]

```

### Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.83 (sec) , antiderivative size = 452, normalized size of antiderivative = 1.20, number of steps used = 17, number of rules used = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.333$ , Rules used = {6875, 25, 6129, 6104, 25, 3042, 26, 4201, 2620, 3011, 6096, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx \\
 & \quad \downarrow 6875 \\
 & - \int \frac{(a+bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1) \operatorname{sech}^{-1}(a+bx)^3}{bx} d\operatorname{sech}^{-1}(a+bx) \\
 & \quad \downarrow 25 \\
 & \int - \frac{(a+bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1) \operatorname{sech}^{-1}(a+bx)^3}{bx} d\operatorname{sech}^{-1}(a+bx) \\
 & \quad \downarrow 6129
 \end{aligned}$$

$$\begin{aligned}
& \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^3}{\frac{a}{a+bx}-1} d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow \text{6104} \\
& a \int -\frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^3}{(a+bx)\left(1-\frac{a}{a+bx}\right)} d\operatorname{sech}^{-1}(a+bx) - \int \sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^3 d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow \text{25} \\
& - \int \sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^3 d\operatorname{sech}^{-1}(a+bx) - \\
& a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^3}{(a+bx)\left(1-\frac{a}{a+bx}\right)} d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow \text{3042} \\
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^3}{(a+bx)\left(1-\frac{a}{a+bx}\right)} d\operatorname{sech}^{-1}(a+bx) - \int -i\operatorname{sech}^{-1}(a+bx)^3 \tan(i\operatorname{sech}^{-1}(a+bx)) d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow \text{26} \\
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^3}{(a+bx)\left(1-\frac{a}{a+bx}\right)} d\operatorname{sech}^{-1}(a+bx) + i \int \operatorname{sech}^{-1}(a+bx)^3 \tan(i\operatorname{sech}^{-1}(a+bx)) d\operatorname{sech}^{-1}(a+bx) \\
& \quad \downarrow \text{4201} \\
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^3}{(a+bx)\left(1-\frac{a}{a+bx}\right)} d\operatorname{sech}^{-1}(a+bx) + \\
& i \left( 2i \int \frac{e^{2\operatorname{sech}^{-1}(a+bx)}\operatorname{sech}^{-1}(a+bx)^3}{1+e^{2\operatorname{sech}^{-1}(a+bx)}} d\operatorname{sech}^{-1}(a+bx) - \frac{1}{4}i\operatorname{sech}^{-1}(a+bx)^4 \right) \\
& \quad \downarrow \text{2620} \\
& -a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^3}{(a+bx)\left(1-\frac{a}{a+bx}\right)} d\operatorname{sech}^{-1}(a+bx) + \\
& i \left( \frac{1}{2}\operatorname{sech}^{-1}(a+bx)^3 \log\left(e^{2\operatorname{sech}^{-1}(a+bx)}+1\right) - \frac{3}{2} \int \operatorname{sech}^{-1}(a+bx)^2 \log\left(1+e^{2\operatorname{sech}^{-1}(a+bx)}\right) d\operatorname{sech}^{-1}(a+bx) \right)
\end{aligned}$$

↓ 3011

$$-a \int \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^3}{(a+bx)\left(1-\frac{a}{a+bx}\right)} d\operatorname{sech}^{-1}(a+bx) +$$

$$i\left(2i\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)^3 \log\left(e^{2\operatorname{sech}^{-1}(a+bx)}+1\right) - \frac{3}{2}\left(\int \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) d\operatorname{sech}^{-1}(a+bx)\right)\right)\right)$$

↓ 6096

$$-a\left(\int \frac{e^{\operatorname{sech}^{-1}(a+bx)}\operatorname{sech}^{-1}(a+bx)^3}{-e^{\operatorname{sech}^{-1}(a+bx)}a-\sqrt{1-a^2}+1} d\operatorname{sech}^{-1}(a+bx) + \int \frac{e^{\operatorname{sech}^{-1}(a+bx)}\operatorname{sech}^{-1}(a+bx)^3}{-e^{\operatorname{sech}^{-1}(a+bx)}a+\sqrt{1-a^2}+1} d\operatorname{sech}^{-1}(a+bx) + \dots\right)$$

$$i\left(2i\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)^3 \log\left(e^{2\operatorname{sech}^{-1}(a+bx)}+1\right) - \frac{3}{2}\left(\int \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) d\operatorname{sech}^{-1}(a+bx)\right)\right)\right)$$

↓ 2620

$$-a\left(\frac{3 \int \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) d\operatorname{sech}^{-1}(a+bx)}{a} + \frac{3 \int \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) d\operatorname{sech}^{-1}(a+bx)}{a}\right)$$

$$i\left(2i\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)^3 \log\left(e^{2\operatorname{sech}^{-1}(a+bx)}+1\right) - \frac{3}{2}\left(\int \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) d\operatorname{sech}^{-1}(a+bx)\right)\right)\right)$$

↓ 3011

$$-a\left(\frac{3\left(2 \int \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) d\operatorname{sech}^{-1}(a+bx) - \operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)\right)}{a}\right)$$

$$i\left(2i\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)^3 \log\left(e^{2\operatorname{sech}^{-1}(a+bx)}+1\right) - \frac{3}{2}\left(\int \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) d\operatorname{sech}^{-1}(a+bx)\right)\right)\right)$$

↓ 7163

$$-a\left(\frac{3\left(2\left(\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)\right) - \int \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) d\operatorname{sech}^{-1}(a+bx)\right) - \operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a}\right)$$

$$i\left(2i\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)^3 \log\left(e^{2\operatorname{sech}^{-1}(a+bx)}+1\right) - \frac{3}{2}\left(-\frac{1}{2} \int \operatorname{PolyLog}\left(3, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) d\operatorname{sech}^{-1}(a+bx) - \frac{1}{2} \dots\right)\right)\right)$$

↓ 2720

$$\begin{aligned}
& -a \left( \frac{3 \left( 2 \left( \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left( 3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) - \int e^{-\operatorname{sech}^{-1}(a+bx)} \operatorname{PolyLog} \left( 3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) de^{\operatorname{sech}^{-1}(a+bx)} \right)}{a} \right. \\
& \left. + i \left( 2i \left( \frac{1}{2} \operatorname{sech}^{-1}(a+bx) \right)^3 \log \left( e^{2\operatorname{sech}^{-1}(a+bx)} + 1 \right) - \frac{3}{2} \left( -\frac{1}{4} \int e^{-2\operatorname{sech}^{-1}(a+bx)} \operatorname{PolyLog} \left( 3, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) de^{2\operatorname{sech}^{-1}(a+bx)} \right) \right) \right) \\
& \quad \downarrow \text{7143} \\
& -a \left( \frac{3 \left( 2 \left( \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left( 3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) - \operatorname{PolyLog} \left( 4, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) \right) - \operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog} \left( 2, -e^{2\operatorname{sech}^{-1}(a+bx)} \right)}{a} \right. \\
& \left. + i \left( 2i \left( \frac{1}{2} \operatorname{sech}^{-1}(a+bx) \right)^3 \log \left( e^{2\operatorname{sech}^{-1}(a+bx)} + 1 \right) - \frac{3}{2} \left( -\frac{1}{2} \operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog} \left( 2, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) + \frac{1}{2} \operatorname{sech}^{-1}(a+bx) \right) \right) \right)
\end{aligned}$$

input `Int[ArcSech[a + b*x]^3/x,x]`

output

```

-(a*(ArcSech[a + b*x]^4/(4*a) - (ArcSech[a + b*x]^3*Log[1 - (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/a - (ArcSech[a + b*x]^3*Log[1 - (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])/a + (3*(-(ArcSech[a + b*x]^2*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])]) + 2*(ArcSech[a + b*x]*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])]) - PolyLog[4, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])))/a + (3*(-(ArcSech[a + b*x]^2*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])]) + 2*(ArcSech[a + b*x]*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])]) - PolyLog[4, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])))/a) + I*((-1/4*I)*ArcSech[a + b*x]^4 + (2*I)*((ArcSech[a + b*x]^3*Log[1 + E^(2*ArcSech[a + b*x])])/2 - (3*(-1/2*(ArcSech[a + b*x]^2*PolyLog[2, -E^(2*ArcSech[a + b*x])]) + (ArcSech[a + b*x]*PolyLog[3, -E^(2*ArcSech[a + b*x])])/2 - PolyLog[4, -E^(2*ArcSech[a + b*x])])/4))/2))

```



## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
- rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4201 `Int[(((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_]*(f_)*(x_)]), x_Symbol] := Simp[(-I)*(c + d*x)^(m + 1)/(d*(m + 1)), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x))), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6096

```
Int[(((e_.) + (f_.)*(x_))^(m_.)*Sinh[(c_.) + (d_.)*(x_)])/(Cosh[(c_.) + (d_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[-(e + f*x)^(m + 1)/(b*f*(m + 1)), x] + (Int[(e + f*x)^m*(E^(c + d*x)/(a - Rt[a^2 - b^2, 2] + b*E^(c + d*x))), x] + Int[(e + f*x)^m*(E^(c + d*x)/(a + Rt[a^2 - b^2, 2] + b*E^(c + d*x))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[a^2 - b^2, 0]
```

rule 6104

```
Int[(((e_.) + (f_.)*(x_))^(m_.)*Tanh[(c_.) + (d_.)*(x_)]^(n_.))/(Cosh[(c_.) + (d_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[1/a Int[(e + f*x)^m*Tanh[c + d*x]^n, x], x] - Simp[b/a Int[(e + f*x)^m*Sinh[c + d*x]*(Tanh[c + d*x])^(n - 1)/(a + b*Cosh[c + d*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

rule 6129

```
Int[(((e_.) + (f_.)*(x_))^(m_.)*(F_)[(c_.) + (d_.)*(x_)]^(n_.)*(G_)[(c_.) + (d_.)*(x_)]^(p_.))/(a_ + (b_.)*Sech[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[(e + f*x)^m*Cosh[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Cosh[c + d*x])), x] /; FreeQ[{a, b, c, d, e, f}, x] && HyperbolicQ[F] && HyperbolicQ[G] && IntegersQ[m, n, p]
```

rule 6875

```
Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[-(d^(m + 1))^(p - 1) Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[(((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^(c_.)*((a_.) + (b_.)*(x_))^(p_.)]), x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]
```

**Maple [F]**

$$\int \frac{\operatorname{arcsech}(bx+a)^3}{x} dx$$

input `int(arcsech(b*x+a)^3/x,x)`

output `int(arcsech(b*x+a)^3/x,x)`

**Fricas [F]**

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx = \int \frac{\operatorname{arsech}(bx+a)^3}{x} dx$$

input `integrate(arcsech(b*x+a)^3/x,x, algorithm="fricas")`

output `integral(arcsech(b*x + a)^3/x, x)`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx = \int \frac{\operatorname{asech}^3(a+bx)}{x} dx$$

input `integrate(asech(b*x+a)**3/x,x)`

output `Integral(asech(a + b*x)**3/x, x)`

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x} dx = \int \frac{\operatorname{arsech}(bx + a)^3}{x} dx$$

input `integrate(arcsech(b*x+a)^3/x,x, algorithm="maxima")`

output `integrate(arcsech(b*x + a)^3/x, x)`

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x} dx = \int \frac{\operatorname{arsech}(bx + a)^3}{x} dx$$

input `integrate(arcsech(b*x+a)^3/x,x, algorithm="giac")`

output `integrate(arcsech(b*x + a)^3/x, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)^3}{x} dx$$

input `int(acosh(1/(a + b*x))^3/x,x)`

output `int(acosh(1/(a + b*x))^3/x, x)`

**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x} dx = \int \frac{\operatorname{asech}(bx + a)^3}{x} dx$$

input `int(asech(b*x+a)^3/x,x)`

output `int(asech(a + b*x)**3/x,x)`

### 3.18 $\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^2} dx$

Optimal result	181
Mathematica [F(-1)]	182
Rubi [A] (verified)	182
Maple [F]	184
Fricas [F]	185
Sympy [F]	185
Maxima [F]	185
Giac [F]	186
Mupad [F(-1)]	186
Reduce [F]	187

#### Optimal result

Integrand size = 12, antiderivative size = 330

$$\begin{aligned}
 \int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^2} dx = & -\frac{b\operatorname{sech}^{-1}(a+bx)^3}{a} - \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} \\
 & + \frac{3b\operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
 & - \frac{3b\operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
 & + \frac{6b\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
 & - \frac{6b\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
 & - \frac{6b \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{6b \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}}
 \end{aligned}$$

output

```
-b*arcsech(b*x+a)^3/a-arcsech(b*x+a)^3/x+3*b*arcsech(b*x+a)^2*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-3*b*arcsech(b*x+a)^2*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+6*b*arcsech(b*x+a)*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-6*b*arcsech(b*x+a)*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-6*b*polylog(3,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+6*b*polylog(3,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)
```

**Mathematica [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^2} dx = \$Aborted$$

input

```
Integrate[ArcSech[a + b*x]^3/x^2,x]
```

output

```
Aborted
```

**Rubi [A] (verified)**

Time = 0.92 (sec) , antiderivative size = 332, normalized size of antiderivative = 1.01, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {6875, 5991, 3042, 4679, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^2} dx$$

↓ 6875

$$-b \int \frac{(a+bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1) \operatorname{sech}^{-1}(a+bx)^3}{b^2 x^2} d \operatorname{sech}^{-1}(a+bx)$$

$$\begin{aligned}
& \downarrow 5991 \\
& -b \left( 3 \int -\frac{\operatorname{sech}^{-1}(a+bx)^2}{bx} d\operatorname{sech}^{-1}(a+bx) + \frac{\operatorname{sech}^{-1}(a+bx)^3}{bx} \right) \\
& \downarrow 3042 \\
& -b \left( \frac{\operatorname{sech}^{-1}(a+bx)^3}{bx} + 3 \int \frac{\operatorname{sech}^{-1}(a+bx)^2}{a - \csc\left(i\operatorname{sech}^{-1}(a+bx) + \frac{\pi}{2}\right)} d\operatorname{sech}^{-1}(a+bx) \right) \\
& \downarrow 4679 \\
& -b \left( 3 \int \left( \frac{\operatorname{sech}^{-1}(a+bx)^2}{a} + \frac{\operatorname{sech}^{-1}(a+bx)^2}{a\left(\frac{a}{a+bx} - 1\right)} \right) d\operatorname{sech}^{-1}(a+bx) + \frac{\operatorname{sech}^{-1}(a+bx)^3}{bx} \right) \\
& \downarrow 2009 \\
& -b \left( 3 \left( -\frac{2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a\sqrt{1-a^2}} + \frac{2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \right) \right)
\end{aligned}$$

input

```
Int[ArcSech[a + b*x]^3/x^2,x]
```

output

```
-(b*(ArcSech[a + b*x]^3/(b*x) + 3*(ArcSech[a + b*x]^3/(3*a) - (ArcSech[a +
b*x]^2*Log[1 - (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 - a
^2]) + (ArcSech[a + b*x]^2*Log[1 - (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^
2])])/(a*Sqrt[1 - a^2]) - (2*ArcSech[a + b*x]*PolyLog[2, (a*E^ArcSech[a +
b*x])/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) + (2*ArcSech[a + b*x]*PolyLo
g[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) + (2*P
olyLog[3, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) -
(2*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])/(a*Sqrt[1 - a^
2])))
```



## Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4679 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Sin[e + f*x]^n)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGtQ[m, 0]`

rule 5991 `Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sech[(c_.) + (d_.)*(x_)]^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

rule 6875 `Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

## Maple [F]

$$\int \frac{\operatorname{arcsech}(bx + a)^3}{x^2} dx$$

input `int(arcsech(b*x+a)^3/x^2,x)`

output `int(arcsech(b*x+a)^3/x^2,x)`

**Fricas [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{arsech}(bx + a)^3}{x^2} dx$$

input `integrate(arcsech(b*x+a)^3/x^2,x, algorithm="fricas")`

output `integral(arcsech(b*x + a)^3/x^2, x)`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{asech}^3(a + bx)}{x^2} dx$$

input `integrate(asech(b*x+a)**3/x**2,x)`

output `Integral(asech(a + b*x)**3/x**2, x)`

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{arsech}(bx + a)^3}{x^2} dx$$

input `integrate(arcsech(b*x+a)^3/x^2,x, algorithm="maxima")`

output

```
-log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^3/x - integrate((8*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^3 + 8*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a)^3 - 3*(b^3*x^3 + 2*a*b^2*x^2 + (a^2*b - b)*x - 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a) - ((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*log(b*x + a) - (2*b^3*x^3 + 4*a*b^2*x^2 + (2*a^2*b - b)*x - (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a))*sqrt(b*x + a + 1))*sqrt(-b*x - a + 1))*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^2 - 12*((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^2 + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a)^2)*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a))/(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2 + (b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)), x)
```

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{ar} \operatorname{sech}(bx + a)^3}{x^2} dx$$

input

```
integrate(arcsech(b*x+a)^3/x^2,x, algorithm="giac")
```

output

```
integrate(arcsech(b*x + a)^3/x^2, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)^3}{x^2} dx$$

input

```
int(acosh(1/(a + b*x)))^3/x^2,x)
```

output `int(acosh(1/(a + b*x))^3/x^2, x)`

### Reduce [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{asech}(bx + a)^3}{x^2} dx$$

input `int(asech(b*x+a)^3/x^2,x)`

output `int(asech(a + b*x)**3/x**2,x)`

$$3.19 \quad \int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^3} dx$$

Optimal result	188
Mathematica [F]	189
Rubi [A] (verified)	190
Maple [F]	192
Fricas [F]	192
Sympy [F]	193
Maxima [F]	193
Giac [F]	194
Mupad [F(-1)]	194
Reduce [F]	195

### Optimal result

Integrand size = 12, antiderivative size = 965

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^3} dx = \text{Too large to display}$$

output

```

-3/2*b^2*arcsech(b*x+a)^2/a^2/(-a^2+1)+3/2*b^2*((-b*x-a+1)/(b*x+a+1))^(1/2)
)*(b*x+a+1)*arcsech(b*x+a)^2/a/(-a^2+1)/(b*x+a)/(1-a/(b*x+a))+1/2*b^2*arcs
ech(b*x+a)^3/a^2-1/2*arcsech(b*x+a)^3/x^2+3*b^2*arcsech(b*x+a)*ln(1-a*(1/(
b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a^2/(-
a^2+1)+3/2*b^2*arcsech(b*x+a)^2*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(
b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(3/2)-3*b^2*arcsech(b*x+
a)^2*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1
)^(1/2)))/a^2/(-a^2+1)^(1/2)+3*b^2*arcsech(b*x+a)*ln(1-a*(1/(b*x+a)+(1/(b*
x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a^2/(-a^2+1)-3/2*b^
2*arcsech(b*x+a)^2*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/
2)))/(1+(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(3/2)+3*b^2*arcsech(b*x+a)^2*ln(1-a*(
1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a^2
/(-a^2+1)^(1/2)+3*b^2*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a
)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a^2/(-a^2+1)+3*b^2*arcsech(b*x+a)*polylog(
2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)
)/a^2/(-a^2+1)^(3/2)-6*b^2*arcsech(b*x+a)*polylog(2,a*(1/(b*x+a)+(1/(b*x+a
)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(1/2)+3*b
^2*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^
2+1)^(1/2)))/a^2/(-a^2+1)-3*b^2*arcsech(b*x+a)*polylog(2,a*(1/(b*x+a)+(1/(
b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(...

```

### Mathematica [F]

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^3} dx = \int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^3} dx$$

input

```
Integrate[ArcSech[a + b*x]^3/x^3,x]
```

output

```
Integrate[ArcSech[a + b*x]^3/x^3, x]
```

**Rubi [A] (verified)**

Time = 1.78 (sec) , antiderivative size = 911, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6875, 25, 5991, 3042, 4679, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^3} dx \\
 & \quad \downarrow \text{6875} \\
 & -b^2 \int \frac{(a+bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1) \operatorname{sech}^{-1}(a+bx)^3}{b^3 x^3} d\operatorname{sech}^{-1}(a+bx) \\
 & \quad \downarrow \text{25} \\
 & b^2 \int -\frac{(a+bx) \sqrt{\frac{-a-bx+1}{a+bx+1}} (a+bx+1) \operatorname{sech}^{-1}(a+bx)^3}{b^3 x^3} d\operatorname{sech}^{-1}(a+bx) \\
 & \quad \downarrow \text{5991} \\
 & -b^2 \left( \frac{\operatorname{sech}^{-1}(a+bx)^3}{2b^2 x^2} - \frac{3}{2} \int \frac{\operatorname{sech}^{-1}(a+bx)^2}{b^2 x^2} d\operatorname{sech}^{-1}(a+bx) \right) \\
 & \quad \downarrow \text{3042} \\
 & -b^2 \left( \frac{\operatorname{sech}^{-1}(a+bx)^3}{2b^2 x^2} - \frac{3}{2} \int \frac{\operatorname{sech}^{-1}(a+bx)^2}{\left(a - \csc\left(\operatorname{isech}^{-1}(a+bx) + \frac{\pi}{2}\right)\right)^2} d\operatorname{sech}^{-1}(a+bx) \right) \\
 & \quad \downarrow \text{4679} \\
 & -b^2 \left( \frac{\operatorname{sech}^{-1}(a+bx)^3}{2b^2 x^2} - \frac{3}{2} \int \left( \frac{2\operatorname{sech}^{-1}(a+bx)^2}{a^2 \left(\frac{a}{a+bx} - 1\right)} + \frac{\operatorname{sech}^{-1}(a+bx)^2}{a^2} + \frac{\operatorname{sech}^{-1}(a+bx)^2}{a^2 \left(\frac{a}{a+bx} - 1\right)^2} \right) d\operatorname{sech}^{-1}(a+bx) \right) \\
 & \quad \downarrow \text{2009} \\
 & -b^2 \left( \frac{\operatorname{sech}^{-1}(a+bx)^3}{2b^2 x^2} - \frac{3}{2} \left( \frac{\operatorname{sech}^{-1}(a+bx)^3}{3a^2} - \frac{2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) \operatorname{sech}^{-1}(a+bx)^2}{a^2 \sqrt{1-a^2}} + \frac{\log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2} \right) \right)
 \end{aligned}$$

input `Int[ArcSech[a + b*x]^3/x^3,x]`

output

$$\begin{aligned}
 & -(b^2*(\text{ArcSech}[a + b*x]^3/(2*b^2*x^2) - (3*(-\text{ArcSech}[a + b*x]^2/(a^2*(1 - a^2))) + (\text{Sqrt}[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x)*\text{ArcSech}[a + b*x]^2)/(a*(1 - a^2)*(a + b*x)*(1 - a/(a + b*x))) + \text{ArcSech}[a + b*x]^3/(3*a^2) \\
 & ) + (2*\text{ArcSech}[a + b*x]*\text{Log}[1 - (a*E^{\text{ArcSech}[a + b*x]})/(1 - \text{Sqrt}[1 - a^2])])/(a^2*(1 - a^2)) + (\text{ArcSech}[a + b*x]^2*\text{Log}[1 - (a*E^{\text{ArcSech}[a + b*x]})/(1 - \text{Sqrt}[1 - a^2])])/(a^2*(1 - a^2)^{(3/2)}) - (2*\text{ArcSech}[a + b*x]^2*\text{Log}[1 - (a*E^{\text{ArcSech}[a + b*x]})/(1 - \text{Sqrt}[1 - a^2])])/(a^2*\text{Sqrt}[1 - a^2]) + (2*\text{ArcSech}[a + b*x]*\text{Log}[1 - (a*E^{\text{ArcSech}[a + b*x]})/(1 + \text{Sqrt}[1 - a^2])])/(a^2*(1 - a^2)) - (\text{ArcSech}[a + b*x]^2*\text{Log}[1 - (a*E^{\text{ArcSech}[a + b*x]})/(1 + \text{Sqrt}[1 - a^2])])/(a^2*(1 - a^2)^{(3/2)}) + (2*\text{ArcSech}[a + b*x]^2*\text{Log}[1 - (a*E^{\text{ArcSech}[a + b*x]})/(1 + \text{Sqrt}[1 - a^2])])/(a^2*\text{Sqrt}[1 - a^2]) + (2*\text{PolyLog}[2, (a*E^{\text{ArcSech}[a + b*x]})/(1 - \text{Sqrt}[1 - a^2])])/(a^2*(1 - a^2)) + (2*\text{ArcSech}[a + b*x]*\text{PolyLog}[2, (a*E^{\text{ArcSech}[a + b*x]})/(1 - \text{Sqrt}[1 - a^2])])/(a^2*(1 - a^2)^{(3/2)}) - (4*\text{ArcSech}[a + b*x]*\text{PolyLog}[2, (a*E^{\text{ArcSech}[a + b*x]})/(1 - \text{Sqrt}[1 - a^2])])/(a^2*\text{Sqrt}[1 - a^2]) + (2*\text{PolyLog}[2, (a*E^{\text{ArcSech}[a + b*x]})/(1 + \text{Sqrt}[1 - a^2])])/(a^2*(1 - a^2)) - (2*\text{ArcSech}[a + b*x]*\text{PolyLog}[2, (a*E^{\text{ArcSech}[a + b*x]})/(1 + \text{Sqrt}[1 - a^2])])/(a^2*(1 - a^2)^{(3/2)}) + (4*\text{ArcSech}[a + b*x]*\text{PolyLog}[2, (a*E^{\text{ArcSech}[a + b*x]})/(1 + \text{Sqrt}[1 - a^2])])/(a^2*\text{Sqrt}[1 - a^2]) - (2*\text{PolyLog}[3, (a*E^{\text{ArcSech}[a + b*x]})/(1 - \text{Sqrt}[1 - a^2])])/(a^2*(1 - a^2)^{(3/2)}) + (4*\text{PolyLog}[3, (a*E^{\text{ArcSech}[a + b*x]})/(1 - \text{Sqrt}[1...
 \end{aligned}$$

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



rule 4679 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Sin[e + f*x])^n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGtQ[m, 0]`

rule 5991 `Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sech[(c_.) + (d_.)*(x_)])^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

rule 6875 `Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)])*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

## Maple [F]

$$\int \frac{\operatorname{arcsech}(bx + a)^3}{x^3} dx$$

input `int(arcsech(b*x+a)^3/x^3,x)`

output `int(arcsech(b*x+a)^3/x^3,x)`

## Fricas [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^3} dx = \int \frac{\operatorname{ar sech}(bx + a)^3}{x^3} dx$$

input `integrate(arcsech(b*x+a)^3/x^3,x, algorithm="fricas")`

output `integral(arcsech(b*x + a)^3/x^3, x)`

### Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^3} dx = \int \frac{\operatorname{arsech}^3(a + bx)}{x^3} dx$$

input `integrate(arsech(b*x+a)**3/x**3,x)`

output `Integral(arsech(a + b*x)**3/x**3, x)`

### Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^3} dx = \int \frac{\operatorname{arsech}(bx + a)^3}{x^3} dx$$

input `integrate(arcsech(b*x+a)^3/x^3,x, algorithm="maxima")`

output

```
-1/2*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt
(-b*x - a + 1)*a + b*x + a)^3/x^2 - integrate(1/2*(16*(b^3*x^3 + 3*a*b^2*x
^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b
*x + a)^3 + 16*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x
+ a)^3 - 3*(b^3*x^3 + 2*a*b^2*x^2 + (a^2*b - b)*x - 4*(b^3*x^3 + 3*a*b^2*x
^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a) - (2*(b^3*x^3 + 3*a*b^2*x^2
+ a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*log(b*x + a) - (2*b^3*x^3 +
4*a*b^2*x^2 + (2*a^2*b - b)*x - 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b
- b)*x - a)*log(b*x + a))*sqrt(b*x + a + 1))*sqrt(-b*x - a + 1))*log(sqrt
(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1
)*a + b*x + a)^2 - 24*((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)
*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^2 + (b^3*x^3 + 3*a*b^2*x
^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a)^2)*log(sqrt(b*x + a + 1)*sqr
t(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a))/
(b^3*x^6 + 3*a*b^2*x^5 + (3*a^2*b - b)*x^4 + (a^3 - a)*x^3 + (b^3*x^6 + 3*a
*b^2*x^5 + (3*a^2*b - b)*x^4 + (a^3 - a)*x^3)*sqrt(b*x + a + 1)*sqrt(-b*x
- a + 1)), x)
```

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^3} dx = \int \frac{\operatorname{ar} \operatorname{sech}(bx + a)^3}{x^3} dx$$

input

```
integrate(arcsech(b*x+a)^3/x^3,x, algorithm="giac")
```

output

```
integrate(arcsech(b*x + a)^3/x^3, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^3} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)^3}{x^3} dx$$

input

```
int(acosh(1/(a + b*x))^3/x^3,x)
```

output `int(acosh(1/(a + b*x))^3/x^3, x)`

### Reduce [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^3} dx = \int \frac{\operatorname{asech}(bx + a)^3}{x^3} dx$$

input `int(asech(b*x+a)^3/x^3,x)`

output `int(asech(a + b*x)**3/x**3,x)`

### 3.20 $\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx$

Optimal result	196
Mathematica [A] (verified)	197
Rubi [A] (verified)	197
Maple [A] (verified)	199
Fricas [A] (verification not implemented)	199
Sympy [F]	200
Maxima [A] (verification not implemented)	200
Giac [F]	200
Mupad [F(-1)]	201
Reduce [F]	201

#### Optimal result

Integrand size = 10, antiderivative size = 138

$$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx$$

$$= -\frac{1}{4} \sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}}} \sqrt{x} + \frac{1}{4} \left(-1 + \frac{1}{\sqrt{x}}\right)^{3/2} \left(1 + \frac{1}{\sqrt{x}}\right)^{3/2} x^{3/2}$$

$$- \frac{3}{20} \left(-1 + \frac{1}{\sqrt{x}}\right)^{5/2} \left(1 + \frac{1}{\sqrt{x}}\right)^{5/2} x^{5/2} + \frac{1}{28} \left(-1 + \frac{1}{\sqrt{x}}\right)^{7/2} \left(1 + \frac{1}{\sqrt{x}}\right)^{7/2} x^{7/2} + \frac{1}{4} x^4 \operatorname{sech}^{-1}(\sqrt{x})$$

output

```
-1/4*(-1+1/x^(1/2))^(1/2)*(1/x^(1/2)+1)^(1/2)*x^(1/2)+1/4*(-1+1/x^(1/2))^(3/2)*(1/x^(1/2)+1)^(3/2)*x^(3/2)-3/20*(-1+1/x^(1/2))^(5/2)*(1/x^(1/2)+1)^(5/2)*x^(5/2)+1/28*(-1+1/x^(1/2))^(7/2)*(1/x^(1/2)+1)^(7/2)*x^(7/2)+1/4*x^4*arcsech(x^(1/2))
```

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.61

$$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{1}{140} \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}} (16 + 16\sqrt{x} + 8x + 8x^{3/2} + 6x^2 + 6x^{5/2} + 5x^3 + 5x^{7/2}) + \frac{1}{4} x^4 \operatorname{sech}^{-1}(\sqrt{x})$$

input

```
Integrate[x^3*ArcSech[Sqrt[x]],x]
```

output

```
-1/140*(Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]*(16 + 16*Sqrt[x] + 8*x + 8*x^(3/2) + 6*x^2 + 6*x^(5/2) + 5*x^3 + 5*x^(7/2))) + (x^4*ArcSech[Sqrt[x]])/4
```

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.75, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {6899, 27, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx \\ & \quad \downarrow \text{6899} \\ & \frac{\sqrt{1-x} \int \frac{x^3}{2\sqrt{1-x}} dx}{4\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} + \frac{1}{4} x^4 \operatorname{sech}^{-1}(\sqrt{x}) \\ & \quad \downarrow \text{27} \\ & \frac{\sqrt{1-x} \int \frac{x^3}{\sqrt{1-x}} dx}{8\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} + \frac{1}{4} x^4 \operatorname{sech}^{-1}(\sqrt{x}) \\ & \quad \downarrow \text{53} \end{aligned}$$

$$\frac{\sqrt{1-x} \int \left( -(1-x)^{5/2} + 3(1-x)^{3/2} - 3\sqrt{1-x} + \frac{1}{\sqrt{1-x}} \right) dx}{8\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} + \frac{1}{4}x^4 \operatorname{sech}^{-1}(\sqrt{x})$$

↓ 2009

$$\frac{1}{4}x^4 \operatorname{sech}^{-1}(\sqrt{x}) + \frac{\left(\frac{2}{7}(1-x)^{7/2} - \frac{6}{5}(1-x)^{5/2} + 2(1-x)^{3/2} - 2\sqrt{1-x}\right)\sqrt{1-x}}{8\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}}$$

input `Int[x^3*ArcSech[Sqrt[x]],x]`

output `((-2*Sqrt[1 - x] + 2*(1 - x)^(3/2) - (6*(1 - x)^(5/2)))/5 + (2*(1 - x)^(7/2)))/7)*Sqrt[1 - x]/(8*Sqrt[-1 + 1/Sqrt[x]]*Sqrt[1 + 1/Sqrt[x]]*Sqrt[x]) + (x^4*ArcSech[Sqrt[x]])/4`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6899 `Int[((a_.) + ArcSech[u]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcSech[u])/(d*(m + 1))), x] + Simp[b*(Sqrt[1 - u^2]/(d*(m + 1)*u*Sqrt[-1 + 1/u]*Sqrt[1 + 1/u])) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[1 - u^2])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]`

**Maple [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.39

method	result	size
derivativedivides	$\frac{x^4 \operatorname{arcsech}(\sqrt{x})}{4} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}}{140} (5x^3+6x^2+8x+16)$	54
default	$\frac{x^4 \operatorname{arcsech}(\sqrt{x})}{4} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}}{140} (5x^3+6x^2+8x+16)$	54
parts	$\frac{x^4 \operatorname{arcsech}(\sqrt{x})}{4} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}}{140} (5x^3+6x^2+8x+16)$	54

input `int(x^3*arcsech(x^(1/2)),x,method=_RETURNVERBOSE)`

output `1/4*x^4*arcsech(x^(1/2))-1/140*(-(x^(1/2)-1)/x^(1/2))^(1/2)*x^(1/2)*((x^(1/2)+1)/x^(1/2))^(1/2)*(5*x^3+6*x^2+8*x+16)`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.41

$$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx = \frac{1}{4} x^4 \log \left( \frac{x \sqrt{-\frac{x-1}{x}} + \sqrt{x}}{x} \right) - \frac{1}{140} (5x^3 + 6x^2 + 8x + 16) \sqrt{x} \sqrt{-\frac{x-1}{x}}$$

input `integrate(x^3*arcsech(x^(1/2)),x, algorithm="fricas")`

output `1/4*x^4*log((x*sqrt(-(x-1)/x) + sqrt(x))/x) - 1/140*(5*x^3 + 6*x^2 + 8*x + 16)*sqrt(x)*sqrt(-(x-1)/x)`



**Sympy [F]**

$$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x^3 \operatorname{arsech}(\sqrt{x}) dx$$

input `integrate(x**3*asech(x**(1/2)),x)`

output `Integral(x**3*asech(sqrt(x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.42

$$\begin{aligned} \int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx &= \frac{1}{28} x^{\frac{7}{2}} \left(\frac{1}{x} - 1\right)^{\frac{7}{2}} - \frac{3}{20} x^{\frac{5}{2}} \left(\frac{1}{x} - 1\right)^{\frac{5}{2}} \\ &\quad + \frac{1}{4} x^4 \operatorname{arsech}(\sqrt{x}) + \frac{1}{4} x^{\frac{3}{2}} \left(\frac{1}{x} - 1\right)^{\frac{3}{2}} - \frac{1}{4} \sqrt{x} \sqrt{\frac{1}{x} - 1} \end{aligned}$$

input `integrate(x^3*arcsech(x^(1/2)),x, algorithm="maxima")`

output `1/28*x^(7/2)*(1/x - 1)^(7/2) - 3/20*x^(5/2)*(1/x - 1)^(5/2) + 1/4*x^4*arcsech(sqrt(x)) + 1/4*x^(3/2)*(1/x - 1)^(3/2) - 1/4*sqrt(x)*sqrt(1/x - 1)`

**Giac [F]**

$$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x^3 \operatorname{arsech}(\sqrt{x}) dx$$

input `integrate(x^3*arcsech(x^(1/2)),x, algorithm="giac")`

output `integrate(x^3*arcsech(sqrt(x)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x^3 \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right) dx$$

input `int(x^3*acosh(1/x^(1/2)),x)`output `int(x^3*acosh(1/x^(1/2)), x)`**Reduce [F]**

$$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx = \int \operatorname{asech}(\sqrt{x}) x^3 dx$$

input `int(x^3*asech(x^(1/2)),x)`output `int(asech(sqrt(x))*x**3,x)`

### 3.21 $\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx$

Optimal result	202
Mathematica [A] (verified)	202
Rubi [A] (verified)	203
Maple [A] (verified)	204
Fricas [A] (verification not implemented)	205
Sympy [F]	205
Maxima [A] (verification not implemented)	206
Giac [F]	206
Mupad [F(-1)]	206
Reduce [F]	207

#### Optimal result

Integrand size = 10, antiderivative size = 107

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{1}{3} \sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}}} \sqrt{x} + \frac{2}{9} \left(-1 + \frac{1}{\sqrt{x}}\right)^{3/2} \left(1 + \frac{1}{\sqrt{x}}\right)^{3/2} x^{3/2} - \frac{1}{15} \left(-1 + \frac{1}{\sqrt{x}}\right)^{5/2} \left(1 + \frac{1}{\sqrt{x}}\right)^{5/2} x^{5/2} + \frac{1}{3} x^3 \operatorname{sech}^{-1}(\sqrt{x})$$

output

```
-1/3*(-1+1/x^(1/2))^(1/2)*(1/x^(1/2)+1)^(1/2)*x^(1/2)+2/9*(-1+1/x^(1/2))^(3/2)*(1/x^(1/2)+1)^(3/2)*x^(3/2)-1/15*(-1+1/x^(1/2))^(5/2)*(1/x^(1/2)+1)^(5/2)*x^(5/2)+1/3*x^3*arcsech(x^(1/2))
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.67

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{1}{45} \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}} (8 + 8\sqrt{x} + 4x + 4x^{3/2} + 3x^2 + 3x^{5/2}) + \frac{1}{3} x^3 \operatorname{sech}^{-1}(\sqrt{x})$$

input `Integrate[x^2*ArcSech[Sqrt[x]],x]`

output 
$$-1/45*(\text{Sqrt}[(1 - \text{Sqrt}[x])/(1 + \text{Sqrt}[x])]*(8 + 8*\text{Sqrt}[x] + 4*x + 4*x^{(3/2)} + 3*x^2 + 3*x^{(5/2)})) + (x^3*\text{ArcSech}[\text{Sqrt}[x]])/3$$

### Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.86, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {6899, 27, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2 \text{sech}^{-1}(\sqrt{x}) dx \\ & \quad \downarrow \text{6899} \\ & \frac{\sqrt{1-x} \int \frac{x^2}{2\sqrt{1-x}} dx}{3\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} + \frac{1}{3}x^3 \text{sech}^{-1}(\sqrt{x}) \\ & \quad \downarrow \text{27} \\ & \frac{\sqrt{1-x} \int \frac{x^2}{\sqrt{1-x}} dx}{6\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} + \frac{1}{3}x^3 \text{sech}^{-1}(\sqrt{x}) \\ & \quad \downarrow \text{53} \\ & \frac{\sqrt{1-x} \int \left( (1-x)^{3/2} - 2\sqrt{1-x} + \frac{1}{\sqrt{1-x}} \right) dx}{6\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} + \frac{1}{3}x^3 \text{sech}^{-1}(\sqrt{x}) \\ & \quad \downarrow \text{2009} \\ & \frac{1}{3}x^3 \text{sech}^{-1}(\sqrt{x}) + \frac{\left(-\frac{2}{5}(1-x)^{5/2} + \frac{4}{3}(1-x)^{3/2} - 2\sqrt{1-x}\right)\sqrt{1-x}}{6\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} \end{aligned}$$

input `Int[x^2*ArcSech[Sqrt[x]],x]`

output 
$$\frac{((-2\sqrt{1-x} + (4(1-x)^{3/2})/3 - (2(1-x)^{5/2})/5)\sqrt{1-x}) / (6\sqrt{-1 + 1/\sqrt{x}}\sqrt{1 + 1/\sqrt{x}}\sqrt{x}) + (x^3 \operatorname{ArcSech}[\sqrt{x}]))/3$$

**Defintions of rubi rules used**

rule 27 
$$\operatorname{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \operatorname{Simp}[a \operatorname{Int}[Fx, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{MatchQ}[Fx, (b_*)(Gx_)] /; \operatorname{FreeQ}[b, x]$$

rule 53 
$$\operatorname{Int}[((a_.) + (b_.)(x_))^{(m_.)}*((c_.) + (d_.)(x_))^{(n_.)}, x\_Symbol] \rightarrow \operatorname{Int}[\operatorname{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \operatorname{FreeQ}\{a, b, c, d, n\}, x \ \&\& \ \operatorname{IGtQ}[m, 0] \ \&\& \ ( \ !\operatorname{IntegerQ}[n] \ || \ (\operatorname{EqQ}[c, 0] \ \&\& \ \operatorname{LeQ}[7*m + 4*n + 4, 0]) \ || \ \operatorname{LtQ}[9*m + 5*(n + 1), 0] \ || \ \operatorname{GtQ}[m + n + 2, 0])$$

rule 2009 
$$\operatorname{Int}[u_, x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$$

rule 6899 
$$\operatorname{Int}[((a_.) + \operatorname{ArcSech}[u_]*(b_.))*((c_.) + (d_.)(x_))^{(m_.)}, x\_Symbol] \rightarrow \operatorname{Simp}[(c + d*x)^{(m + 1)}*((a + b*\operatorname{ArcSech}[u])/(d*(m + 1))), x] + \operatorname{Simp}[b*(\sqrt{1 - u^2})/(d*(m + 1)*u*\sqrt{-1 + 1/u}*\sqrt{1 + 1/u}) \operatorname{Int}[\operatorname{SimplifyIntegrand}[(c + d*x)^{(m + 1)}*(D[u, x]/(u*\sqrt{1 - u^2}))], x], x], x] /; \operatorname{FreeQ}\{a, b, c, d, m\}, x \ \&\& \ \operatorname{NeQ}[m, -1] \ \&\& \ \operatorname{InverseFunctionFreeQ}[u, x] \ \&\& \ !\operatorname{FunctionOfQ}[(c + d*x)^{(m + 1)}, u, x] \ \&\& \ !\operatorname{FunctionOfExponentialQ}[u, x]$$

**Maple [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.46

method	result	size
derivativedivides	$\frac{x^3 \operatorname{arcsech}(\sqrt{x})}{3} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}}{45} (3x^2+4x+8)$	49
default	$\frac{x^3 \operatorname{arcsech}(\sqrt{x})}{3} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}}{45} (3x^2+4x+8)$	49
parts	$\frac{x^3 \operatorname{arcsech}(\sqrt{x})}{3} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}}{45} (3x^2+4x+8)$	49

input `int(x^2*arcsech(x^(1/2)),x,method=_RETURNVERBOSE)`

output `1/3*x^3*arcsech(x^(1/2))-1/45*(-(x^(1/2)-1)/x^(1/2))^(1/2)*x^(1/2)*((x^(1/2)+1)/x^(1/2))^(1/2)*(3*x^2+4*x+8)`

### Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.49

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = \frac{1}{3} x^3 \log \left( \frac{x \sqrt{-\frac{x-1}{x}} + \sqrt{x}}{x} \right) - \frac{1}{45} (3x^2 + 4x + 8) \sqrt{x} \sqrt{-\frac{x-1}{x}}$$

input `integrate(x^2*arcsech(x^(1/2)),x, algorithm="fricas")`

output `1/3*x^3*log((x*sqrt(-(x - 1)/x) + sqrt(x))/x) - 1/45*(3*x^2 + 4*x + 8)*sqrt(x)*sqrt(-(x - 1)/x)`

### Sympy [F]

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x^2 \operatorname{asech}(\sqrt{x}) dx$$

input `integrate(x**2*asech(x**(1/2)),x)`

output `Integral(x**2*asech(sqrt(x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.43

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{1}{15} x^{\frac{5}{2}} \left(\frac{1}{x} - 1\right)^{\frac{5}{2}} + \frac{1}{3} x^3 \operatorname{ar} \operatorname{sech}(\sqrt{x}) \\ + \frac{2}{9} x^{\frac{3}{2}} \left(\frac{1}{x} - 1\right)^{\frac{3}{2}} - \frac{1}{3} \sqrt{x} \sqrt{\frac{1}{x} - 1}$$

input `integrate(x^2*arcsech(x^(1/2)),x, algorithm="maxima")`output `-1/15*x^(5/2)*(1/x - 1)^(5/2) + 1/3*x^3*arcsech(sqrt(x)) + 2/9*x^(3/2)*(1/x - 1)^(3/2) - 1/3*sqrt(x)*sqrt(1/x - 1)`**Giac [F]**

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x^2 \operatorname{ar} \operatorname{sech}(\sqrt{x}) dx$$

input `integrate(x^2*arcsech(x^(1/2)),x, algorithm="giac")`output `integrate(x^2*arcsech(sqrt(x)), x)`**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x^2 \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right) dx$$

input `int(x^2*acosh(1/x^(1/2)),x)`output `int(x^2*acosh(1/x^(1/2)), x)`

**Reduce [F]**

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = \int \operatorname{asech}(\sqrt{x}) x^2 dx$$

input `int(x^2*asech(x^(1/2)),x)`

output `int(asech(sqrt(x))*x**2,x)`



### 3.22 $\int x \operatorname{sech}^{-1}(\sqrt{x}) dx$

Optimal result	208
Mathematica [A] (verified)	208
Rubi [A] (verified)	209
Maple [A] (verified)	210
Fricas [A] (verification not implemented)	211
Sympy [F]	211
Maxima [A] (verification not implemented)	212
Giac [F]	212
Mupad [F(-1)]	212
Reduce [F]	213

#### Optimal result

Integrand size = 8, antiderivative size = 76

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{1}{2} \sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}}} \sqrt{x} + \frac{1}{6} \left(-1 + \frac{1}{\sqrt{x}}\right)^{3/2} \left(1 + \frac{1}{\sqrt{x}}\right)^{3/2} x^{3/2} + \frac{1}{2} x^2 \operatorname{sech}^{-1}(\sqrt{x})$$

output

$-1/2*(-1+1/x^{(1/2)})^{(1/2)}*(1/x^{(1/2)}+1)^{(1/2)}*x^{(1/2)}+1/6*(-1+1/x^{(1/2)})^{(3/2)}*(1/x^{(1/2)}+1)^{(3/2)}*x^{(3/2)}+1/2*x^2*\operatorname{arcsech}(x^{(1/2)})$

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.74

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{1}{6} \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}} (2 + 2\sqrt{x} + x + x^{3/2}) + \frac{1}{2} x^2 \operatorname{sech}^{-1}(\sqrt{x})$$

input

`Integrate[x*ArcSech[Sqrt[x]],x]`

output

```
-1/6*(Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]*(2 + 2*Sqrt[x] + x + x^(3/2))) + (
x^2*ArcSech[Sqrt[x]])/2
```

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.04, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6899, 27, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx$$

$$\downarrow 6899$$

$$\frac{\sqrt{1-x} \int \frac{x}{2\sqrt{1-x}} dx}{2\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} + \frac{1}{2}x^2 \operatorname{sech}^{-1}(\sqrt{x})$$

$$\downarrow 27$$

$$\frac{\sqrt{1-x} \int \frac{x}{\sqrt{1-x}} dx}{4\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} + \frac{1}{2}x^2 \operatorname{sech}^{-1}(\sqrt{x})$$

$$\downarrow 53$$

$$\frac{\sqrt{1-x} \int \left( \frac{1}{\sqrt{1-x}} - \sqrt{1-x} \right) dx}{4\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} + \frac{1}{2}x^2 \operatorname{sech}^{-1}(\sqrt{x})$$

$$\downarrow 2009$$

$$\frac{1}{2}x^2 \operatorname{sech}^{-1}(\sqrt{x}) + \frac{\left(\frac{2}{3}(1-x)^{3/2} - 2\sqrt{1-x}\right)\sqrt{1-x}}{4\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}}$$

input

```
Int [x*ArcSech[Sqrt [x]] , x]
```

output  $((-2\sqrt{1-x} + (2(1-x)^{3/2})/3)\sqrt{1-x})/(4\sqrt{-1+1/\sqrt{x}})\sqrt{1+1/\sqrt{x}}\sqrt{x} + (x^2\text{ArcSech}[\sqrt{x}])/2$

**Defintions of rubi rules used**

rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$

rule 53  $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}*((c_.) + (d_.)(x_)^{(n_.)}), x\_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ (\ !\text{IntegerQ}[n] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{LeQ}[7*m + 4*n + 4, 0]) \ || \ \text{LtQ}[9*m + 5*(n + 1), 0] \ || \ \text{GtQ}[m + n + 2, 0])$

rule 2009  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 6899  $\text{Int}[(a_.) + \text{ArcSech}[u_]*(b_.))*((c_.) + (d_.)(x_)^{(m_.)}), x\_Symbol] \rightarrow \text{Simp}[(c + d*x)^{(m + 1)}*((a + b*\text{ArcSech}[u])/(d*(m + 1))), x] + \text{Simp}[b*(\text{Sqrt}[1 - u^2]/(d*(m + 1)*u*\text{Sqrt}[-1 + 1/u]*\text{Sqrt}[1 + 1/u])) \ \text{Int}[\text{SimplifyIntegrand}[(c + d*x)^{(m + 1)}*(D[u, x]/(u*\text{Sqrt}[1 - u^2])), x], x], x] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \ \&\& \ \text{NeQ}[m, -1] \ \&\& \ \text{InverseFunctionFreeQ}[u, x] \ \&\& \ !\text{FunctionOfQ}[(c + d*x)^{(m + 1)}, u, x] \ \&\& \ !\text{FunctionOfExponentialQ}[u, x]$

**Maple [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.55

method	result	size
derivativedivides	$\frac{x^2 \text{arcsech}(\sqrt{x})}{2} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}}{6} (x+2)$	42
default	$\frac{x^2 \text{arcsech}(\sqrt{x})}{2} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}}{6} (x+2)$	42
parts	$\frac{x^2 \text{arcsech}(\sqrt{x})}{2} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}}{6} (x+2)$	42

input `int(x*arcsech(x^(1/2)),x,method=_RETURNVERBOSE)`

output `1/2*x^2*arcsech(x^(1/2))-1/6*(-(x^(1/2)-1)/x^(1/2))^(1/2)*x^(1/2)*((x^(1/2)+1)/x^(1/2))^(1/2)*(x+2)`

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.59

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = \frac{1}{2} x^2 \log \left( \frac{x \sqrt{-\frac{x-1}{x}} + \sqrt{x}}{x} \right) - \frac{1}{6} (x+2) \sqrt{x} \sqrt{-\frac{x-1}{x}}$$

input `integrate(x*arcsech(x^(1/2)),x, algorithm="fricas")`

output `1/2*x^2*log((x*sqrt(-(x - 1)/x) + sqrt(x))/x) - 1/6*(x + 2)*sqrt(x)*sqrt(-(x - 1)/x)`

### Sympy [F]

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x \operatorname{asech}(\sqrt{x}) dx$$

input `integrate(x*asech(x**(1/2)),x)`

output `Integral(x*asech(sqrt(x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.45

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = \frac{1}{6} x^{\frac{3}{2}} \left( \frac{1}{x} - 1 \right)^{\frac{3}{2}} + \frac{1}{2} x^2 \operatorname{arsech}(\sqrt{x}) - \frac{1}{2} \sqrt{x} \sqrt{\frac{1}{x} - 1}$$

input `integrate(x*arcsech(x^(1/2)),x, algorithm="maxima")`output `1/6*x^(3/2)*(1/x - 1)^(3/2) + 1/2*x^2*arcsech(sqrt(x)) - 1/2*sqrt(x)*sqrt(1/x - 1)`**Giac [F]**

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x \operatorname{arsech}(\sqrt{x}) dx$$

input `integrate(x*arcsech(x^(1/2)),x, algorithm="giac")`output `integrate(x*arcsech(sqrt(x)), x)`**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right) dx$$

input `int(x*acosh(1/x^(1/2)),x)`output `int(x*acosh(1/x^(1/2)), x)`

Reduce [F]

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = \int \operatorname{asech}(\sqrt{x}) x dx$$

input `int(x*asech(x^(1/2)),x)`

output `int(asech(sqrt(x))*x,x)`

### 3.23 $\int \operatorname{sech}^{-1}(\sqrt{x}) dx$

Optimal result	214
Mathematica [B] (verified)	214
Rubi [A] (verified)	215
Maple [A] (verified)	216
Fricas [A] (verification not implemented)	216
Sympy [F]	217
Maxima [A] (verification not implemented)	217
Giac [F]	217
Mupad [F(-1)]	218
Reduce [F]	218

#### Optimal result

Integrand size = 6, antiderivative size = 38

$$\int \operatorname{sech}^{-1}(\sqrt{x}) dx = -\sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}}} \sqrt{x} + x \operatorname{sech}^{-1}(\sqrt{x})$$

output `-(-1+1/x^(1/2))^(1/2)*(1/x^(1/2)+1)^(1/2)*x^(1/2)+x*arcsech(x^(1/2))`

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 118 vs. 2(38) = 76.

Time = 0.08 (sec) , antiderivative size = 118, normalized size of antiderivative = 3.11

$$\int \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{2(-1 + \sqrt{1 - \sqrt{x}})^2 (-1 + \sqrt{1 + \sqrt{x}})^2 \sqrt{\frac{1 - \sqrt{x}}{1 + \sqrt{x}}} \sqrt{1 + \sqrt{x}}}{(-2 + \sqrt{1 - \sqrt{x}} + \sqrt{1 + \sqrt{x}})^2 \sqrt{1 - \sqrt{x}}} + x \operatorname{sech}^{-1}(\sqrt{x})$$

input `Integrate[ArcSech[Sqrt[x]], x]`

output

```
(-2*(-1 + Sqrt[1 - Sqrt[x]])^2*(-1 + Sqrt[1 + Sqrt[x]])^2*Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]*Sqrt[1 + Sqrt[x]])/((-2 + Sqrt[1 - Sqrt[x]] + Sqrt[1 + Sqrt[x]])^2*Sqrt[1 - Sqrt[x]]) + x*ArcSech[Sqrt[x]]
```

**Rubi [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.13, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6897, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{sech}^{-1}(\sqrt{x}) dx$$

$$\downarrow 6897$$

$$\frac{\sqrt{1-x} \int \frac{1}{2\sqrt{1-x}} dx}{\sqrt{\frac{1}{x}} - 1 \sqrt{\frac{1}{x}} + 1\sqrt{x}} + x \operatorname{sech}^{-1}(\sqrt{x})$$

$$\downarrow 17$$

$$x \operatorname{sech}^{-1}(\sqrt{x}) - \frac{1-x}{\sqrt{\frac{1}{x}} - 1 \sqrt{\frac{1}{x}} + 1\sqrt{x}}$$

input

```
Int[ArcSech[Sqrt[x]], x]
```

output

```
-((1 - x)/(Sqrt[-1 + 1/Sqrt[x]]*Sqrt[1 + 1/Sqrt[x]]*Sqrt[x])) + x*ArcSech[Sqrt[x]]
```



**Defintions of rubi rules used**

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 6897 `Int[ArcSech[u_], x_Symbol] := Simp[x*ArcSech[u], x] + Simp[Sqrt[1 - u^2]/(u*Sqrt[-1 + 1/u]*Sqrt[1 + 1/u]) Int[SimplifyIntegrand[x*(D[u, x]/(u*Sqrt[1 - u^2])), x], x], x] /; InverseFunctionFreeQ[u, x] && !FunctionOfExponentialQ[u, x]`

**Maple [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.95

method	result	size
derivativedivides	$x \operatorname{arcsech}(\sqrt{x}) - \sqrt{x} \sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}$	36
default	$x \operatorname{arcsech}(\sqrt{x}) - \sqrt{x} \sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}$	36
parts	$x \operatorname{arcsech}(\sqrt{x}) - \sqrt{x} \sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}$	36

input `int(arcsech(x^(1/2)),x,method=_RETURNVERBOSE)`

output `x*arcsech(x^(1/2))-x^(1/2)*(-(x^(1/2)-1)/x^(1/2))^(1/2)*((x^(1/2)+1)/x^(1/2))^(1/2)`

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.03

$$\int \operatorname{sech}^{-1}(\sqrt{x}) dx = x \log \left( \frac{x \sqrt{-\frac{x-1}{x}} + \sqrt{x}}{x} \right) - \sqrt{x} \sqrt{-\frac{x-1}{x}}$$

input `integrate(arcsech(x^(1/2)),x, algorithm="fricas")`

output `x*log((x*sqrt(-(x - 1)/x) + sqrt(x))/x) - sqrt(x)*sqrt(-(x - 1)/x)`

### Sympy [F]

$$\int \operatorname{sech}^{-1}(\sqrt{x}) \, dx = \int \operatorname{arsech}(\sqrt{x}) \, dx$$

input `integrate(arsech(x**(1/2)),x)`

output `Integral(arsech(sqrt(x)), x)`

### Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.50

$$\int \operatorname{sech}^{-1}(\sqrt{x}) \, dx = x \operatorname{arsech}(\sqrt{x}) - \sqrt{x} \sqrt{\frac{1}{x} - 1}$$

input `integrate(arcsech(x^(1/2)),x, algorithm="maxima")`

output `x*arcsech(sqrt(x)) - sqrt(x)*sqrt(1/x - 1)`

### Giac [F]

$$\int \operatorname{sech}^{-1}(\sqrt{x}) \, dx = \int \operatorname{arsech}(\sqrt{x}) \, dx$$

input `integrate(arcsech(x^(1/2)),x, algorithm="giac")`

output `integrate(arcsech(sqrt(x)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{sech}^{-1}(\sqrt{x}) \, dx = \int \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right) \, dx$$

input `int(acosh(1/x^(1/2)), x)`output `int(acosh(1/x^(1/2)), x)`**Reduce [F]**

$$\int \operatorname{sech}^{-1}(\sqrt{x}) \, dx = \int \operatorname{asech}(\sqrt{x}) \, dx$$

input `int(asech(x^(1/2)), x)`output `int(asech(sqrt(x)), x)`

### 3.24 $\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx$

Optimal result	219
Mathematica [A] (verified)	219
Rubi [C] (warning: unable to verify)	220
Maple [A] (verified)	223
Fricas [F]	223
Sympy [F]	223
Maxima [F]	224
Giac [F]	224
Mupad [F(-1)]	224
Reduce [F]	225

#### Optimal result

Integrand size = 10, antiderivative size = 46

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx = \operatorname{sech}^{-1}(\sqrt{x})^2 - 2\operatorname{sech}^{-1}(\sqrt{x}) \log\left(1 + e^{2\operatorname{sech}^{-1}(\sqrt{x})}\right) - \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(\sqrt{x})}\right)$$

output

```
arcsech(x^(1/2))^2-2*arcsech(x^(1/2))*ln(1+(1/x^(1/2)+(-1+1/x^(1/2))^(1/2)
*(1/x^(1/2)+1)^(1/2))^2)-polylog(2,-(1/x^(1/2)+(-1+1/x^(1/2))^(1/2)*(1/x^(
1/2)+1)^(1/2))^2)
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.98

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx = -\operatorname{sech}^{-1}(\sqrt{x}) \left( \operatorname{sech}^{-1}(\sqrt{x}) + 2 \log\left(1 + e^{-2\operatorname{sech}^{-1}(\sqrt{x})}\right) \right) + \operatorname{PolyLog}\left(2, -e^{-2\operatorname{sech}^{-1}(\sqrt{x})}\right)$$

input

```
Integrate[ArcSech[Sqrt[x]]/x,x]
```

output

```
-(ArcSech[Sqrt[x]]*(ArcSech[Sqrt[x]] + 2*Log[1 + E^(-2*ArcSech[Sqrt[x]])])
) + PolyLog[2, -E^(-2*ArcSech[Sqrt[x]])]
```

**Rubi [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 0.47 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.26, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$ , Rules used = {7267, 6835, 6297, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx \\ & \quad \downarrow 7267 \\ & 2 \int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{\sqrt{x}} d\sqrt{x} \\ & \quad \downarrow 6835 \\ & -2 \int \frac{\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)}{\sqrt{x}} d\frac{1}{\sqrt{x}} \\ & \quad \downarrow 6297 \\ & -2 \int \sqrt{\frac{\frac{1}{\sqrt{x}} - 1}{1 + \frac{1}{\sqrt{x}}}} \left(1 + \frac{1}{\sqrt{x}}\right) \sqrt{x} \operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right) d\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right) \\ & \quad \downarrow 3042 \\ & -2 \int -i \operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right) \tan\left(i \operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)\right) d\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right) \\ & \quad \downarrow 26 \\ & 2i \int \operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right) \tan\left(i \operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)\right) d\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right) \\ & \quad \downarrow 4201 \end{aligned}$$

$$2i \left( 2i \int \frac{e^{2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)} \operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)}{1 + e^{2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)}} d\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right) - \frac{ix}{2} \right)$$

↓ 2620

$$2i \left( 2i \left( \frac{1}{2} \operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right) \log\left(e^{2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)} + 1\right) - \frac{1}{2} \int \log\left(1 + e^{2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)}\right) d\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right) \right) - \frac{ix}{2} \right)$$

↓ 2715

$$2i \left( 2i \left( \frac{1}{2} \operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right) \log\left(e^{2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)} + 1\right) - \frac{1}{4} \int e^{2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)} \log\left(1 + e^{2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)}\right) de^{2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)} \right) - \frac{ix}{2} \right)$$

↓ 2838

$$2i \left( 2i \left( \frac{1}{4} \operatorname{PolyLog}\left(2, -e^{2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)}\right) + \frac{1}{2} \operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right) \log\left(e^{2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)} + 1\right) \right) - \frac{ix}{2} \right)$$

input `Int[ArcSech[Sqrt[x]]/x, x]`

output `(2*I)*((-1/2*I)*x + (2*I)*((ArcCosh[1/Sqrt[x]]*Log[1 + E^(2*ArcCosh[1/Sqrt[x]])]))/2 + PolyLog[2, -E^(2*ArcCosh[1/Sqrt[x]])]/4)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F x_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] :> Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4201 `Int[((c_.) + (d_.)*(x_)^(m_.))*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x
_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[
(c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x))), x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6297 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(x_), x_Symbol] :> Simp[1/b
Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a
, b, c}, x] && IGtQ[n, 0]`

rule 6835 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))/(x_), x_Symbol] :> -Subst[Int[(a +
b*ArcCosh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]`

rule 7267 `Int[u_, x_Symbol] :> With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]`

**Maple [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.41

method	result
derivativedivides	$\operatorname{arcsech}(\sqrt{x})^2 - 2 \operatorname{arcsech}(\sqrt{x}) \ln \left( 1 + \left( \frac{1}{\sqrt{x}} + \sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}}} \right)^2 \right) - \operatorname{polylog}$
default	$\operatorname{arcsech}(\sqrt{x})^2 - 2 \operatorname{arcsech}(\sqrt{x}) \ln \left( 1 + \left( \frac{1}{\sqrt{x}} + \sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}}} \right)^2 \right) - \operatorname{polylog}$

input `int(arcsech(x^(1/2))/x,x,method=_RETURNVERBOSE)`

output `arcsech(x^(1/2))^2-2*arcsech(x^(1/2))*ln(1+(1/x^(1/2)+(-1+1/x^(1/2))^(1/2)*(1+1/x^(1/2))^(1/2))^2)-polylog(2,-(1/x^(1/2)+(-1+1/x^(1/2))^(1/2)*(1+1/x^(1/2))^(1/2))^2)`

**Fricas [F]**

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arsech}(\sqrt{x})}{x} dx$$

input `integrate(arcsech(x^(1/2))/x,x, algorithm="fricas")`

output `integral(arcsech(sqrt(x))/x, x)`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{asech}(\sqrt{x})}{x} dx$$

input `integrate(asech(x**(1/2))/x,x)`



output `Integral(asech(sqrt(x))/x, x)`

### Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arsech}(\sqrt{x})}{x} dx$$

input `integrate(arcsech(x^(1/2))/x,x, algorithm="maxima")`

output `-1/4*log(x)^2 + log(x)*log(sqrt(sqrt(x) + 1)*sqrt(-sqrt(x) + 1) + 1) - log(sqrt(x) + 1)*log(sqrt(x)) - log(sqrt(x))*log(-sqrt(x) + 1) - dilog(-sqrt(x)) - dilog(sqrt(x)) + integrate(1/2*log(x)/((x - 1)*e^(1/2*log(sqrt(x) + 1) + 1/2*log(-sqrt(x) + 1)) + x - 1), x)`

### Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arsech}(\sqrt{x})}{x} dx$$

input `integrate(arcsech(x^(1/2))/x,x, algorithm="giac")`

output `integrate(arcsech(sqrt(x))/x, x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right)}{x} dx$$

input `int(acosh(1/x^(1/2))/x,x)`

output `int(acosh(1/x^(1/2))/x, x)`

**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{asech}(\sqrt{x})}{x} dx$$

input `int(asech(x^(1/2))/x, x)`

output `int(asech(sqrt(x))/x, x)`

### 3.25 $\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx$

Optimal result	226
Mathematica [A] (verified)	226
Rubi [A] (verified)	227
Maple [A] (verified)	229
Fricas [A] (verification not implemented)	229
Sympy [F]	230
Maxima [A] (verification not implemented)	230
Giac [F]	230
Mupad [B] (verification not implemented)	231
Reduce [F]	231

#### Optimal result

Integrand size = 10, antiderivative size = 93

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}}{2\sqrt{x}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} + \frac{\sqrt{1-x}\operatorname{arctanh}(\sqrt{1-x})}{2\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}}$$

output

```
1/2*(-1+1/x^(1/2))^(1/2)*(1/x^(1/2)+1)^(1/2)/x^(1/2)-arcsech(x^(1/2))/x+1/2*(1-x)^(1/2)*arctanh((1-x)^(1/2))/(-1+1/x^(1/2))^(1/2)/(1/x^(1/2)+1)^(1/2)/x^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.19

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}(1+\sqrt{x}) - 2\operatorname{sech}^{-1}(\sqrt{x}) + x \log\left(1 + \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}\right) + \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}\sqrt{x}}{2x} - \frac{1}{2}x \log(x)$$

input

```
Integrate[ArcSech[Sqrt[x]]/x^2,x]
```

output

```
(Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]*(1 + Sqrt[x]) - 2*ArcSech[Sqrt[x]] + x*
Log[1 + Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]] + Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x]
)])*Sqrt[x]] - (x*Log[x])/2)/(2*x)
```

**Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.85, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6899, 27, 52, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx \\
 & \quad \downarrow \text{6899} \\
 & -\frac{\sqrt{1-x} \int \frac{1}{2\sqrt{1-xx^2}} dx}{\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} \\
 & \quad \downarrow \text{27} \\
 & -\frac{\sqrt{1-x} \int \frac{1}{\sqrt{1-xx^2}} dx}{2\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} \\
 & \quad \downarrow \text{52} \\
 & -\frac{\sqrt{1-x} \left( \frac{1}{2} \int \frac{1}{\sqrt{1-xx}} dx - \frac{\sqrt{1-x}}{x} \right)}{2\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} \\
 & \quad \downarrow \text{73} \\
 & -\frac{\sqrt{1-x} \left( -\int \frac{1}{x} d\sqrt{1-x} - \frac{\sqrt{1-x}}{x} \right)}{2\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} \\
 & \quad \downarrow \text{219} \\
 & -\frac{\sqrt{1-x} \left( -\operatorname{arctanh}(\sqrt{1-x}) - \frac{\sqrt{1-x}}{x} \right)}{2\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x}
 \end{aligned}$$

input `Int[ArcSech[Sqrt[x]]/x^2,x]`

output `-(ArcSech[Sqrt[x]]/x) - (Sqrt[1 - x]*(-(Sqrt[1 - x]/x) - ArcTanh[Sqrt[1 - x]]))/(2*Sqrt[-1 + 1/Sqrt[x]]*Sqrt[1 + 1/Sqrt[x]]*Sqrt[x])`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 52 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntegerQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 6899 `Int[((a_.) + ArcSech[u_]*(b_.))*((c_.) + (d_.)*(x_)^(m_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcSech[u])/(d*(m + 1))), x] + Simp[b*(Sqrt[1 - u^2]/(d*(m + 1)*u*Sqrt[-1 + 1/u]*Sqrt[1 + 1/u])) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[1 - u^2])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]`

**Maple [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.69

method	result	size
derivativedivides	$-\frac{\operatorname{arcsech}(\sqrt{x})}{x} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(\operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x+\sqrt{1-x}\right)}{2\sqrt{x}\sqrt{1-x}}$	64
default	$-\frac{\operatorname{arcsech}(\sqrt{x})}{x} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(\operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x+\sqrt{1-x}\right)}{2\sqrt{x}\sqrt{1-x}}$	64
parts	$-\frac{\operatorname{arcsech}(\sqrt{x})}{x} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(\operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x+\sqrt{1-x}\right)}{2\sqrt{x}\sqrt{1-x}}$	64

input `int(arcsech(x^(1/2))/x^2,x,method=_RETURNVERBOSE)`

output 
$$-\operatorname{arcsech}(x^{1/2})/x+1/2*(-(x^{1/2}-1)/x^{1/2})^{1/2}/x^{1/2}*((x^{1/2}+1)/x^{1/2})^{1/2}*(\operatorname{arctanh}(1/(1-x)^{1/2}))*x+(1-x)^{1/2}/(1-x)^{1/2}$$

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.48

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx = \frac{(x-2) \log\left(\frac{x\sqrt{-\frac{x-1}{x}}+\sqrt{x}}{x}\right) + \sqrt{x}\sqrt{-\frac{x-1}{x}}}{2x}$$

input `integrate(arcsech(x^(1/2))/x^2,x, algorithm="fricas")`

output 
$$1/2*((x-2)*\log((x*\sqrt{-(x-1)/x} + \sqrt{x})/x) + \sqrt{x}*\sqrt{-(x-1)/x})/x$$

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx = \int \frac{\operatorname{arsech}(\sqrt{x})}{x^2} dx$$

input `integrate(asech(x**(1/2))/x**2,x)`

output `Integral(asech(sqrt(x))/x**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.70

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx = -\frac{\sqrt{x}\sqrt{\frac{1}{x}-1}}{2\left(x\left(\frac{1}{x}-1\right)-1\right)} - \frac{\operatorname{arsech}(\sqrt{x})}{x} + \frac{1}{4} \log\left(\sqrt{x}\sqrt{\frac{1}{x}-1}+1\right) - \frac{1}{4} \log\left(\sqrt{x}\sqrt{\frac{1}{x}-1}-1\right)$$

input `integrate(arcsech(x^(1/2))/x^2,x, algorithm="maxima")`

output `-1/2*sqrt(x)*sqrt(1/x - 1)/(x*(1/x - 1) - 1) - arcsech(sqrt(x))/x + 1/4*log(sqrt(x)*sqrt(1/x - 1) + 1) - 1/4*log(sqrt(x)*sqrt(1/x - 1) - 1)`

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx = \int \frac{\operatorname{arsech}(\sqrt{x})}{x^2} dx$$

input `integrate(arcsech(x^(1/2))/x^2,x, algorithm="giac")`

output `integrate(arcsech(sqrt(x))/x^2, x)`

**Mupad [B] (verification not implemented)**

Time = 4.24 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.43

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{\frac{1}{\sqrt{x}} - 1} \sqrt{\frac{1}{\sqrt{x}} + 1}}{2\sqrt{x}} - \frac{2 \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right) \left(\frac{1}{2\sqrt{x}} - \frac{\sqrt{x}}{4}\right)}{\sqrt{x}}$$

input `int(acosh(1/x^(1/2))/x^2,x)`output `((1/x^(1/2) - 1)^(1/2)*(1/x^(1/2) + 1)^(1/2))/(2*x^(1/2)) - (2*acosh(1/x^(1/2))*(1/(2*x^(1/2)) - x^(1/2)/4))/x^(1/2)`**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx = \int \frac{\operatorname{asech}(\sqrt{x})}{x^2} dx$$

input `int(asech(x^(1/2))/x^2,x)`output `int(asech(sqrt(x))/x**2,x)`



### 3.26 $\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx$

Optimal result	232
Mathematica [A] (verified)	232
Rubi [A] (verified)	233
Maple [A] (verified)	235
Fricas [A] (verification not implemented)	235
Sympy [F]	236
Maxima [A] (verification not implemented)	236
Giac [F]	237
Mupad [F(-1)]	237
Reduce [F]	237

#### Optimal result

Integrand size = 10, antiderivative size = 126

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx = \frac{\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}}{8x^{3/2}} + \frac{3\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}}{16\sqrt{x}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{2x^2} + \frac{3\sqrt{1-x}\operatorname{arctanh}(\sqrt{1-x})}{16\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}}$$

output

```
1/8*(-1+1/x^(1/2))^(1/2)*(1/x^(1/2)+1)^(1/2)/x^(3/2)+3/16*(-1+1/x^(1/2))^(1/2)*(1/x^(1/2)+1)^(1/2)/x^(1/2)-1/2*arcsech(x^(1/2))/x^2+3/16*(1-x)^(1/2)*arctanh((1-x)^(1/2))/(-1+1/x^(1/2))^(1/2)/(1/x^(1/2)+1)^(1/2)/x^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.99

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx = \frac{1}{16} \left( \frac{\sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}(2 + 2\sqrt{x} + 3x + 3x^{3/2})}{x^2} - \frac{8\operatorname{sech}^{-1}(\sqrt{x})}{x^2} + 3 \log \left( 1 + \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}} + \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}\sqrt{x} \right) - \frac{3 \log(x)}{2} \right)$$

input `Integrate[ArcSech[Sqrt[x]]/x^3,x]`

output `((Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])])*(2 + 2*Sqrt[x] + 3*x + 3*x^(3/2)))/x^2 - (8*ArcSech[Sqrt[x]])/x^2 + 3*Log[1 + Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]] + Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]*Sqrt[x] - (3*Log[x])/2)/16`

### Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.81, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {6899, 27, 52, 52, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx \\
 & \quad \downarrow 6899 \\
 & -\frac{\sqrt{1-x} \int \frac{1}{2\sqrt{1-xx^3}} dx}{2\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{2x^2} \\
 & \quad \downarrow 27 \\
 & -\frac{\sqrt{1-x} \int \frac{1}{\sqrt{1-xx^3}} dx}{4\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{2x^2} \\
 & \quad \downarrow 52 \\
 & -\frac{\sqrt{1-x} \left( \frac{3}{4} \int \frac{1}{\sqrt{1-xx^2}} dx - \frac{\sqrt{1-x}}{2x^2} \right)}{4\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{2x^2} \\
 & \quad \downarrow 52 \\
 & -\frac{\sqrt{1-x} \left( \frac{3}{4} \left( \frac{1}{2} \int \frac{1}{\sqrt{1-xx}} dx - \frac{\sqrt{1-x}}{x} \right) - \frac{\sqrt{1-x}}{2x^2} \right)}{4\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{2x^2} \\
 & \quad \downarrow 73
 \end{aligned}$$

$$-\frac{\sqrt{1-x}\left(\frac{3}{4}\left(-\int\frac{1}{x}d\sqrt{1-x}-\frac{\sqrt{1-x}}{x}\right)-\frac{\sqrt{1-x}}{2x^2}\right)}{4\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}}-\frac{\operatorname{sech}^{-1}(\sqrt{x})}{2x^2}$$

↓ 219

$$-\frac{\sqrt{1-x}\left(\frac{3}{4}\left(-\operatorname{arctanh}(\sqrt{1-x})-\frac{\sqrt{1-x}}{x}\right)-\frac{\sqrt{1-x}}{2x^2}\right)}{4\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}}-\frac{\operatorname{sech}^{-1}(\sqrt{x})}{2x^2}$$

input

```
Int[ArcSech[Sqrt[x]]/x^3,x]
```

output

```
-1/2*ArcSech[Sqrt[x]]/x^2 - (Sqrt[1 - x]*(-1/2*Sqrt[1 - x]/x^2 + (3*(-Sqrt[1 - x]/x) - ArcTanh[Sqrt[1 - x]]))/4)/(4*Sqrt[-1 + 1/Sqrt[x]]*Sqrt[1 + 1/Sqrt[x]]*Sqrt[x])
```

### Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

rule 52

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1))/((b*c - a*d)*(m + 1)), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]
```

rule 73

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

rule 219

```
Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

rule 6899

```
Int[((a_.) + ArcSech[u_]*(b_.))*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Si
mp[(c + d*x)^(m + 1)*((a + b*ArcSech[u])/(d*(m + 1))), x] + Simp[b*(Sqrt[1
- u^2]/(d*(m + 1)*u*Sqrt[-1 + 1/u]*Sqrt[1 + 1/u])) Int[SimplifyIntegrand[
(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[1 - u^2])), x], x], x] /; FreeQ[{a, b, c
, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c
+ d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]
```

**Maple [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.63

method	result	size
derivativedivides	$-\frac{\operatorname{arcsech}(\sqrt{x})}{2x^2} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(3\operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x^2+3\sqrt{1-x}x+2\sqrt{1-x}\right)}{16x^{\frac{3}{2}}\sqrt{1-x}}$	79
default	$-\frac{\operatorname{arcsech}(\sqrt{x})}{2x^2} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(3\operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x^2+3\sqrt{1-x}x+2\sqrt{1-x}\right)}{16x^{\frac{3}{2}}\sqrt{1-x}}$	79
parts	$-\frac{\operatorname{arcsech}(\sqrt{x})}{2x^2} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(3\operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x^2+3\sqrt{1-x}x+2\sqrt{1-x}\right)}{16x^{\frac{3}{2}}\sqrt{1-x}}$	79

input

```
int(arcsech(x^(1/2))/x^3,x,method=_RETURNVERBOSE)
```

output

```
-1/2*arcsech(x^(1/2))/x^2+1/16*(-(x^(1/2)-1)/x^(1/2))^1/2/x^(3/2)*((x^(1
/2)+1)/x^(1/2))^1/2*(3*arctanh(1/(1-x)^(1/2))*x^2+3*(1-x)^(1/2)*x+2*(1-x
)^(1/2))/(1-x)^(1/2)
```

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.43

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx = \frac{(3x+2)\sqrt{x}\sqrt{-\frac{x-1}{x}} + (3x^2-8)\log\left(\frac{x\sqrt{-\frac{x-1}{x}}+\sqrt{x}}{x}\right)}{16x^2}$$

input

```
integrate(arcsech(x^(1/2))/x^3,x, algorithm="fricas")
```

output

```
1/16*((3*x + 2)*sqrt(x)*sqrt(-(x - 1)/x) + (3*x^2 - 8)*log((x*sqrt(-(x - 1)/x) + sqrt(x))/x))/x^2
```

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\operatorname{asech}(\sqrt{x})}{x^3} dx$$

input

```
integrate(asech(x**(1/2))/x**3,x)
```

output

```
Integral(asech(sqrt(x))/x**3, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.73

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx = -\frac{3x^{\frac{3}{2}}\left(\frac{1}{x}-1\right)^{\frac{3}{2}}-5\sqrt{x}\sqrt{\frac{1}{x}-1}}{16\left(x^2\left(\frac{1}{x}-1\right)^2-2x\left(\frac{1}{x}-1\right)+1\right)}-\frac{\operatorname{arsech}(\sqrt{x})}{2x^2}$$

$$+\frac{3}{32}\log\left(\sqrt{x}\sqrt{\frac{1}{x}-1}+1\right)-\frac{3}{32}\log\left(\sqrt{x}\sqrt{\frac{1}{x}-1}-1\right)$$

input

```
integrate(arcsech(x^(1/2))/x^3,x, algorithm="maxima")
```

output

```
-1/16*(3*x^(3/2)*(1/x - 1)^(3/2) - 5*sqrt(x)*sqrt(1/x - 1))/(x^2*(1/x - 1)^2 - 2*x*(1/x - 1) + 1) - 1/2*arcsech(sqrt(x))/x^2 + 3/32*log(sqrt(x)*sqrt(1/x - 1) + 1) - 3/32*log(sqrt(x)*sqrt(1/x - 1) - 1)
```

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\operatorname{arsech}(\sqrt{x})}{x^3} dx$$

input `integrate(arcsech(x^(1/2))/x^3,x, algorithm="giac")`

output `integrate(arcsech(sqrt(x))/x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right)}{x^3} dx$$

input `int(acosh(1/x^(1/2))/x^3,x)`

output `int(acosh(1/x^(1/2))/x^3, x)`

**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\operatorname{asech}(\sqrt{x})}{x^3} dx$$

input `int(asech(x^(1/2))/x^3,x)`

output `int(asech(sqrt(x))/x**3,x)`

### 3.27 $\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx$

Optimal result	238
Mathematica [A] (verified)	239
Rubi [A] (verified)	239
Maple [A] (verified)	242
Fricas [A] (verification not implemented)	242
Sympy [F]	243
Maxima [A] (verification not implemented)	243
Giac [F]	244
Mupad [F(-1)]	244
Reduce [F]	244

#### Optimal result

Integrand size = 10, antiderivative size = 157

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx = \frac{\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}}{18x^{5/2}} + \frac{5\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}}{72x^{3/2}} + \frac{5\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}}{48\sqrt{x}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3} + \frac{5\sqrt{1-x}\operatorname{arctanh}(\sqrt{1-x})}{48\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}}$$

output

```
1/18*(-1+1/x^(1/2))^(1/2)*(1/x^(1/2)+1)^(1/2)/x^(5/2)+5/72*(-1+1/x^(1/2))^(1/2)*(1/x^(1/2)+1)^(1/2)/x^(3/2)+5/48*(-1+1/x^(1/2))^(1/2)*(1/x^(1/2)+1)^(1/2)/x^(1/2)-1/3*arcsech(x^(1/2))/x^3+5/48*(1-x)^(1/2)*arctanh((1-x)^(1/2))/(-1+1/x^(1/2))^(1/2)/(1/x^(1/2)+1)^(1/2)/x^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.89

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx$$

$$= \frac{\sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}(8 + 8\sqrt{x} + 10x + 10x^{3/2} + 15x^2 + 15x^{5/2}) - 48\operatorname{sech}^{-1}(\sqrt{x}) + 15x^3 \log\left(1 + \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}\right) + \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}}{144x^3}$$

input `Integrate[ArcSech[Sqrt[x]]/x^4,x]`

output `(Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]*(8 + 8*Sqrt[x] + 10*x + 10*x^(3/2) + 15*x^2 + 15*x^(5/2)) - 48*ArcSech[Sqrt[x]] + 15*x^3*Log[1 + Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]] + Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]*Sqrt[x]] - (15*x^3*Log[x])/2)/(144*x^3)`

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.78, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$ , Rules used = {6899, 27, 52, 52, 52, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx$$

$$\downarrow \text{6899}$$

$$-\frac{\sqrt{1-x} \int \frac{1}{2\sqrt{1-xx^4}} dx}{3\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3}$$

$$\downarrow \text{27}$$

$$-\frac{\sqrt{1-x} \int \frac{1}{\sqrt{1-xx^4}} dx}{6\sqrt{\frac{1}{\sqrt{x}} - 1}\sqrt{\frac{1}{\sqrt{x}} + 1}\sqrt{x}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3}$$



$$\begin{aligned}
& \downarrow 52 \\
& -\frac{\sqrt{1-x}\left(\frac{5}{6}\int\frac{1}{\sqrt{1-xx^3}}dx-\frac{\sqrt{1-x}}{3x^3}\right)}{6\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}}-\frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3} \\
& \downarrow 52 \\
& -\frac{\sqrt{1-x}\left(\frac{5}{6}\left(\frac{3}{4}\int\frac{1}{\sqrt{1-xx^2}}dx-\frac{\sqrt{1-x}}{2x^2}\right)-\frac{\sqrt{1-x}}{3x^3}\right)}{6\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}}-\frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3} \\
& \downarrow 52 \\
& -\frac{\sqrt{1-x}\left(\frac{5}{6}\left(\frac{3}{4}\left(\frac{1}{2}\int\frac{1}{\sqrt{1-xx}}dx-\frac{\sqrt{1-x}}{x}\right)-\frac{\sqrt{1-x}}{2x^2}\right)-\frac{\sqrt{1-x}}{3x^3}\right)}{6\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}}-\frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3} \\
& \downarrow 73 \\
& -\frac{\sqrt{1-x}\left(\frac{5}{6}\left(\frac{3}{4}\left(-\int\frac{1}{x}d\sqrt{1-x}-\frac{\sqrt{1-x}}{x}\right)-\frac{\sqrt{1-x}}{2x^2}\right)-\frac{\sqrt{1-x}}{3x^3}\right)}{6\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}}-\frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3} \\
& \downarrow 219 \\
& -\frac{\sqrt{1-x}\left(\frac{5}{6}\left(\frac{3}{4}\left(-\operatorname{arctanh}(\sqrt{1-x})-\frac{\sqrt{1-x}}{x}\right)-\frac{\sqrt{1-x}}{2x^2}\right)-\frac{\sqrt{1-x}}{3x^3}\right)}{6\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}}-\frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3}
\end{aligned}$$

input `Int[ArcSech[Sqrt[x]]/x^4,x]`

output `-1/3*ArcSech[Sqrt[x]]/x^3 - (Sqrt[1-x]*(-1/3*Sqrt[1-x]/x^3 + (5*(-1/2*Sqrt[1-x]/x^2 + (3*(-(Sqrt[1-x]/x) - ArcTanh[Sqrt[1-x]]))/4))/6))/(6*Sqrt[-1 + 1/Sqrt[x]]*Sqrt[1 + 1/Sqrt[x]]*Sqrt[x])`

## Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`
- rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 6899 `Int[((a_.) + ArcSech[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcSech[u])/(d*(m + 1))), x] + Simp[b*(Sqrt[1 - u^2]/(d*(m + 1)*u*Sqrt[-1 + 1/u]*Sqrt[1 + 1/u])) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[1 - u^2])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]`

**Maple [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.58

method	result	size
derivativedivides	$-\frac{\operatorname{arcsech}(\sqrt{x})}{3x^3} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(15\operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x^3+15\sqrt{1-x}x^2+10\sqrt{1-x}x+8\sqrt{1-x}\right)}{144x^{\frac{5}{2}}\sqrt{1-x}}$	91
default	$-\frac{\operatorname{arcsech}(\sqrt{x})}{3x^3} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(15\operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x^3+15\sqrt{1-x}x^2+10\sqrt{1-x}x+8\sqrt{1-x}\right)}{144x^{\frac{5}{2}}\sqrt{1-x}}$	91
parts	$-\frac{\operatorname{arcsech}(\sqrt{x})}{3x^3} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(15\operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x^3+15\sqrt{1-x}x^2+10\sqrt{1-x}x+8\sqrt{1-x}\right)}{144x^{\frac{5}{2}}\sqrt{1-x}}$	91

input `int(arcsech(x^(1/2))/x^4,x,method=_RETURNVERBOSE)`

output `-1/3*arcsech(x^(1/2))/x^3+1/144*(-(x^(1/2)-1)/x^(1/2))^(1/2)/x^(5/2)*((x^(1/2)+1)/x^(1/2))^(1/2)*(15*arctanh(1/(1-x)^(1/2))*x^3+15*(1-x)^(1/2)*x^2+10*(1-x)^(1/2)*x+8*(1-x)^(1/2))/(1-x)^(1/2)`

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.38

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx = \frac{(15x^2 + 10x + 8)\sqrt{x}\sqrt{-\frac{x-1}{x}} + 3(5x^3 - 16)\log\left(\frac{x\sqrt{-\frac{x-1}{x}} + \sqrt{x}}{x}\right)}{144x^3}$$

input `integrate(arcsech(x^(1/2))/x^4,x, algorithm="fricas")`

output `1/144*((15*x^2 + 10*x + 8)*sqrt(x)*sqrt(-(x - 1)/x) + 3*(5*x^3 - 16)*log((x*sqrt(-(x - 1)/x) + sqrt(x))/x))/x^3`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\operatorname{asech}(\sqrt{x})}{x^4} dx$$

input `integrate(asech(x**(1/2))/x**4,x)`

output `Integral(asech(sqrt(x))/x**4, x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.74

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx = -\frac{15x^{\frac{5}{2}}\left(\frac{1}{x}-1\right)^{\frac{5}{2}} - 40x^{\frac{3}{2}}\left(\frac{1}{x}-1\right)^{\frac{3}{2}} + 33\sqrt{x}\sqrt{\frac{1}{x}-1}}{144\left(x^3\left(\frac{1}{x}-1\right)^3 - 3x^2\left(\frac{1}{x}-1\right)^2 + 3x\left(\frac{1}{x}-1\right) - 1\right)} - \frac{\operatorname{arosech}(\sqrt{x})}{3x^3} + \frac{5}{96} \log\left(\sqrt{x}\sqrt{\frac{1}{x}-1} + 1\right) - \frac{5}{96} \log\left(\sqrt{x}\sqrt{\frac{1}{x}-1} - 1\right)$$

input `integrate(arcsech(x^(1/2))/x^4,x, algorithm="maxima")`

output `-1/144*(15*x^(5/2)*(1/x - 1)^(5/2) - 40*x^(3/2)*(1/x - 1)^(3/2) + 33*sqrt(x)*sqrt(1/x - 1))/(x^3*(1/x - 1)^3 - 3*x^2*(1/x - 1)^2 + 3*x*(1/x - 1) - 1) - 1/3*arcsech(sqrt(x))/x^3 + 5/96*log(sqrt(x)*sqrt(1/x - 1) + 1) - 5/96*log(sqrt(x)*sqrt(1/x - 1) - 1)`

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\operatorname{arsech}(\sqrt{x})}{x^4} dx$$

input `integrate(arcsech(x^(1/2))/x^4,x, algorithm="giac")`

output `integrate(arcsech(sqrt(x))/x^4, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right)}{x^4} dx$$

input `int(acosh(1/x^(1/2))/x^4,x)`

output `int(acosh(1/x^(1/2))/x^4, x)`

**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\operatorname{asech}(\sqrt{x})}{x^4} dx$$

input `int(asech(x^(1/2))/x^4,x)`

output `int(asech(sqrt(x))/x**4,x)`

### 3.28 $\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx$

Optimal result	245
Mathematica [A] (verified)	245
Rubi [A] (verified)	246
Maple [A] (verified)	247
Fricas [A] (verification not implemented)	247
Sympy [F]	248
Maxima [A] (verification not implemented)	248
Giac [F]	248
Mupad [B] (verification not implemented)	249
Reduce [F]	249

#### Optimal result

Integrand size = 4, antiderivative size = 21

$$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx = -\sqrt{-1+x}\sqrt{1+x} + x \operatorname{arccosh}(x)$$

output

```
-(-1+x)^(1/2)*(1+x)^(1/2)+x*arccosh(x)
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.19

$$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx = -\sqrt{\frac{-1+x}{1+x}}(1+x) + x \operatorname{sech}^{-1}\left(\frac{1}{x}\right)$$

input

```
Integrate[ArcSech[x^(-1)],x]
```

output

```
-(Sqrt[(-1 + x)/(1 + x)]*(1 + x)) + x*ArcSech[x^(-1)]
```

**Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {6881, 6294, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx \\ & \quad \downarrow \text{6881} \\ & \int \operatorname{arccosh}(x) dx \\ & \quad \downarrow \text{6294} \\ & x \operatorname{arccosh}(x) - \int \frac{x}{\sqrt{x-1}\sqrt{x+1}} dx \\ & \quad \downarrow \text{83} \\ & x \operatorname{arccosh}(x) - \sqrt{x-1}\sqrt{x+1} \end{aligned}$$

input `Int[ArcSech[x^(-1)], x]`

output `-(Sqrt[-1 + x]*Sqrt[1 + x]) + x*ArcCosh[x]`

**Defintions of rubi rules used**

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] :> Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

rule 6294 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.), x_Symbol] :> Simp[x*(a + b*ArcCosh[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcCosh[c*x])^(n - 1))/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

rule 6881

```
Int[ArcSech[(c_.)/((a_.) + (b_.)*(x_)^(n_.))]^(m_.)*(u_.), x_Symbol] :> Int
[u*ArcCosh[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]
```

**Maple [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.95

method	result	size
parts	$x \operatorname{arcsech}\left(\frac{1}{x}\right) - \sqrt{x-1} \sqrt{x+1}$	20
derivativedivides	$x \operatorname{arcsech}\left(\frac{1}{x}\right) - \sqrt{-\left(-1 + \frac{1}{x}\right)x} \sqrt{\left(\frac{1}{x} + 1\right)x}$	29
default	$x \operatorname{arcsech}\left(\frac{1}{x}\right) - \sqrt{-\left(-1 + \frac{1}{x}\right)x} \sqrt{\left(\frac{1}{x} + 1\right)x}$	29

input

```
int(arcsech(1/x), x, method=_RETURNVERBOSE)
```

output

```
x*arcsech(1/x)-(x-1)^(1/2)*(x+1)^(1/2)
```

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx = x \log\left(x + \sqrt{x^2 - 1}\right) - \sqrt{x^2 - 1}$$

input

```
integrate(arcsech(1/x), x, algorithm="fricas")
```

output

```
x*log(x + sqrt(x^2 - 1)) - sqrt(x^2 - 1)
```



**Sympy [F]**

$$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx = \int \operatorname{asech}\left(\frac{1}{x}\right) dx$$

input `integrate(asech(1/x), x)`

output `Integral(asech(1/x), x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.76

$$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx = x \operatorname{arsech}\left(\frac{1}{x}\right) - \sqrt{x^2 - 1}$$

input `integrate(arcsech(1/x), x, algorithm="maxima")`

output `x*arcsech(1/x) - sqrt(x^2 - 1)`

**Giac [F]**

$$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx = \int \operatorname{arsech}\left(\frac{1}{x}\right) dx$$

input `integrate(arcsech(1/x), x, algorithm="giac")`

output `integrate(arcsech(1/x), x)`

**Mupad [B] (verification not implemented)**

Time = 3.49 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx = x \operatorname{acosh}(x) - \sqrt{x-1} \sqrt{x+1}$$

input `int(acosh(x),x)`

output `x*acosh(x) - (x - 1)^(1/2)*(x + 1)^(1/2)`

**Reduce [F]**

$$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx = \int \operatorname{asech}\left(\frac{1}{x}\right) dx$$

input `int(asech(1/x),x)`

output `int(asech(1/x),x)`

### 3.29 $\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx$

Optimal result	250
Mathematica [B] (verified)	250
Rubi [C] (warning: unable to verify)	251
Maple [A] (verified)	254
Fricas [F(-2)]	254
Sympy [F]	255
Maxima [F]	255
Giac [F]	255
Mupad [F(-1)]	256
Reduce [F]	256

#### Optimal result

Integrand size = 10, antiderivative size = 61

$$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx = \frac{\operatorname{sech}^{-1}(ax^n)^2}{2n} - \frac{\operatorname{sech}^{-1}(ax^n) \log\left(1 + e^{2\operatorname{sech}^{-1}(ax^n)}\right)}{n} - \frac{\operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(ax^n)}\right)}{2n}$$

output

```
1/2*arcsech(a*x^n)^2/n-arcsech(a*x^n)*ln(1+(1/a/(x^n)+(1/a/(x^n)-1)^(1/2)*
(1/a/(x^n)+1)^(1/2))^2)/n-1/2*polylog(2,-(1/a/(x^n)+(1/a/(x^n)-1)^(1/2)*(1
/a/(x^n)+1)^(1/2))^2)/n
```

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 219 vs. 2(61) = 122.

Time = 0.64 (sec) , antiderivative size = 219, normalized size of antiderivative = 3.59

$$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx = \operatorname{sech}^{-1}(ax^n) \log(x) + \frac{\sqrt{\frac{1-ax^n}{1+ax^n}} (4\sqrt{-1+a^2x^{2n}} \arctan(\sqrt{-1+a^2x^{2n}}) (2n \log(x) - \log(a^2x^{2n})) + \sqrt{1-a^2x^{2n}} (\log^2(a^2x^{2n}) - 8(n$$

input `Integrate[ArcSech[a*x^n]/x,x]`

output `ArcSech[a*x^n]*Log[x] + (Sqrt[(1 - a*x^n)/(1 + a*x^n)]*(4*Sqrt[-1 + a^2*x^(2*n)]*ArcTan[Sqrt[-1 + a^2*x^(2*n)]]*(2*n*Log[x] - Log[a^2*x^(2*n)]) + Sqrt[1 - a^2*x^(2*n)]*(Log[a^2*x^(2*n)]^2 - 4*Log[a^2*x^(2*n)]*Log[(1 + Sqrt[1 - a^2*x^(2*n)])/2] + 2*Log[(1 + Sqrt[1 - a^2*x^(2*n)])/2]^2 - 4*PolyLog[2, 1/2 - Sqrt[1 - a^2*x^(2*n)])/2])))/(8*(n - a*n*x^n))`

### Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.53 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.26, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$ , Rules used = {7282, 6835, 6297, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx \\
 & \quad \downarrow 7282 \\
 & \int \frac{x^{-n} \operatorname{sech}^{-1}(ax^n) dx^n}{n} \\
 & \quad \downarrow 6835 \\
 & \int \frac{x^{-n} \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right) dx^{-n}}{n} \\
 & \quad \downarrow 6297 \\
 & \int \frac{ax^n \sqrt{\frac{\frac{x^{-n}}{a}-1}{\frac{x^{-n}}{a}+1}} \left(\frac{x^{-n}}{a} + 1\right) \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right) d\operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)}{n} \\
 & \quad \downarrow 3042 \\
 & \int \frac{-i \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right) \tan\left(i \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)\right) d\operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)}{n}
 \end{aligned}$$

$$\begin{array}{c}
\downarrow 26 \\
\frac{i \int \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right) \tan\left(i \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)\right) d \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)}{n} \\
\downarrow 4201 \\
\frac{i \left( 2i \int \frac{e^{2 \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)} \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)}{1+e^{2 \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)}} d \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right) - \frac{1}{2} i x^{2n} \right)}{n} \\
\downarrow 2620 \\
\frac{i \left( 2i \left( \frac{1}{2} \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right) \log\left(e^{2 \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)} + 1\right) - \frac{1}{2} \int \log\left(1 + e^{2 \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)}\right) d \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right) \right) - \frac{1}{2} i x^{2n} \right)}{n} \\
\downarrow 2715 \\
\frac{i \left( 2i \left( \frac{1}{2} \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right) \log\left(e^{2 \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)} + 1\right) - \frac{1}{4} \int e^{2 \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)} \log\left(1 + e^{2 \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)}\right) d e^{2 \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)} \right) \right)}{n} \\
\downarrow 2838 \\
\frac{i \left( 2i \left( \frac{1}{4} \operatorname{PolyLog}\left(2, -e^{2 \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)}\right) + \frac{1}{2} \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right) \log\left(e^{2 \operatorname{arccosh}\left(\frac{x^{-n}}{a}\right)} + 1\right) \right) - \frac{1}{2} i x^{2n} \right)}{n}
\end{array}$$

input `Int[ArcSech[a*x^n]/x,x]`

output `(I*((-1/2*I)*x^(2*n) + (2*I)*((ArcCosh[1/(a*x^n)]*Log[1 + E^(2*ArcCosh[1/(a*x^n)])]))/2 + PolyLog[2, -E^(2*ArcCosh[1/(a*x^n)])]/4))/n`

## Defintions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 2620  $\text{Int}[(((F_)^((g_)*(e_) + (f_)*(x_)))^{(n_)*((c_) + (d_)*(x_))^{(m_))}/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^{(n_)}), x\_Symbol] \rightarrow \text{Simp}[(c + d*x)^m/(b*f*g*n*\text{Log}[F])]*\text{Log}[1 + b*((F^(g*(e + f*x)))^n/a)], x] - \text{Simp}[d*(m/(b*f*g*n*\text{Log}[F])) \text{Int}[(c + d*x)^{m-1}*\text{Log}[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; \text{FreeQ}\{F, a, b, c, d, e, f, g, n\}, x\} \ \&\& \ \text{IGtQ}[m, 0]$
- rule 2715  $\text{Int}[\text{Log}[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^{(n_)}], x\_Symbol] \rightarrow \text{Simp}[1/(d*e*n*\text{Log}[F]) \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; \text{FreeQ}\{F, a, b, c, d, e, n\}, x\} \ \&\& \ \text{GtQ}[a, 0]$
- rule 2838  $\text{Int}[\text{Log}[(c_)*((d_) + (e_)*(x_)^{(n_)})]/(x_), x\_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n/n, x] /; \text{FreeQ}\{c, d, e, n\}, x\} \ \&\& \ \text{EqQ}[c*d, 1]$
- rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4201  $\text{Int}[((c_) + (d_)*(x_))^{(m_)*\tan[(e_) + (\text{Complex}[0, fz_])*(f_)*(x_)], x\_Symbol] \rightarrow \text{Simp}[(-I)*((c + d*x)^{(m+1)}/(d*(m+1))), x] + \text{Simp}[2*I \text{Int}[(c + d*x)^m*(E^{(2*(-I)*e + f*fz*x)})/(1 + E^{(2*(-I)*e + f*fz*x}))], x], x] /; \text{FreeQ}\{c, d, e, f, fz\}, x\} \ \&\& \ \text{IGtQ}[m, 0]$
- rule 6297  $\text{Int}[((a_) + \text{ArcCosh}[(c_)*(x_)]*(b_))^{(n_)} / (x_), x\_Symbol] \rightarrow \text{Simp}[1/b \text{Subst}[\text{Int}[x^n*\text{Tanh}[-a/b + x/b], x], x, a + b*\text{ArcCosh}[c*x]], x] /; \text{FreeQ}\{a, b, c\}, x\} \ \&\& \ \text{IGtQ}[n, 0]$
- rule 6835  $\text{Int}[((a_) + \text{ArcSech}[(c_)*(x_)]*(b_)) / (x_), x\_Symbol] \rightarrow -\text{Subst}[\text{Int}[(a + b*\text{ArcCosh}[x/c])/x, x], x, 1/x] /; \text{FreeQ}\{a, b, c\}, x\}$

rule 7282

```
Int[(u_)/(x_), x_Symbol] := With[{lst = PowerVariableExpn[u, 0, x]}, Simp[1
/lst[[2]] Subst[Int[NormalizeIntegrand[Simplify[lst[[1]]/x], x], x], x, (
lst[[3]]*x)^lst[[2]]], x] /; !FalseQ[lst] && NeQ[lst[[2]], 0] /; NonsumQ[
u] && !RationalFunctionQ[u, x]
```

**Maple [A] (verified)**

Time = 0.66 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.82

method	result
derivativedivides	$\frac{\frac{\operatorname{arcsech}(ax^n)^2}{2} - \operatorname{arcsech}(ax^n) \ln\left(1 + \left(\frac{x^{-n}}{a} + \sqrt{\frac{x^{-n}}{a} - 1} \sqrt{\frac{x^{-n}}{a} + 1}\right)^2\right)}{n} - \frac{\operatorname{polylog}\left(2, -\left(\frac{x^{-n}}{a} + \sqrt{\frac{x^{-n}}{a} - 1} \sqrt{\frac{x^{-n}}{a} + 1}\right)^2\right)}{2}$
default	$\frac{\frac{\operatorname{arcsech}(ax^n)^2}{2} - \operatorname{arcsech}(ax^n) \ln\left(1 + \left(\frac{x^{-n}}{a} + \sqrt{\frac{x^{-n}}{a} - 1} \sqrt{\frac{x^{-n}}{a} + 1}\right)^2\right)}{n} - \frac{\operatorname{polylog}\left(2, -\left(\frac{x^{-n}}{a} + \sqrt{\frac{x^{-n}}{a} - 1} \sqrt{\frac{x^{-n}}{a} + 1}\right)^2\right)}{2}$

input

```
int(arcsech(a*x^n)/x,x,method=_RETURNVERBOSE)
```

output

```
1/n*(1/2*arcsech(a*x^n)^2-arcsech(a*x^n)*ln(1+(1/a/(x^n)+(1/a/(x^n)-1)^(1/2)
/2)*(1/a/(x^n)+1)^(1/2))^2)-1/2*polylog(2,-(1/a/(x^n)+(1/a/(x^n)-1)^(1/2)*
(1/a/(x^n)+1)^(1/2))^2))
```

**Fricas [F(-2)]**

Exception generated.

$$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx = \text{Exception raised: TypeError}$$

input

```
integrate(arcsech(a*x^n)/x,x, algorithm="fricas")
```

output

```
Exception raised: TypeError >> Error detected within library code: inte
grate: implementation incomplete (constant residues)
```

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{arsech}(ax^n)}{x} dx$$

input `integrate(asech(a*x**n)/x,x)`

output `Integral(asech(a*x**n)/x, x)`

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{arsech}(ax^n)}{x} dx$$

input `integrate(arcsech(a*x^n)/x,x, algorithm="maxima")`

output `a^2*n*integrate(x^(2*n)*log(x)/(a^2*x*x^(2*n) + (a^2*x*x^(2*n) - x)*sqrt(a*x^n + 1)*sqrt(-a*x^n + 1) - x), x) + n*integrate(1/2*log(x)/(a*x*x^n + x), x) - n*integrate(1/2*log(x)/(a*x*x^n - x), x) + log(sqrt(a*x^n + 1)*sqrt(-a*x^n + 1) + 1)*log(x) - log(a)*log(x) - log(x)*log(x^n)`

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{arsech}(ax^n)}{x} dx$$

input `integrate(arcsech(a*x^n)/x,x, algorithm="giac")`

output `integrate(arcsech(a*x^n)/x, x)`



**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{ax^n}\right)}{x} dx$$

input `int(acosh(1/(a*x^n))/x,x)`output `int(acosh(1/(a*x^n))/x, x)`**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{asech}(x^na)}{x} dx$$

input `int(asech(a*x^n)/x,x)`output `int(asech(x**n*a)/x,x)`

### 3.30 $\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx$

Optimal result	257
Mathematica [A] (verified)	257
Rubi [C] (warning: unable to verify)	258
Maple [F]	261
Fricas [F]	261
Sympy [F]	261
Maxima [F]	262
Giac [F]	262
Mupad [F(-1)]	262
Reduce [F]	263

#### Optimal result

Integrand size = 10, antiderivative size = 54

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = \frac{1}{10} \operatorname{sech}^{-1}(ax^5)^2 - \frac{1}{5} \operatorname{sech}^{-1}(ax^5) \log\left(1 + e^{2\operatorname{sech}^{-1}(ax^5)}\right) - \frac{1}{10} \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(ax^5)}\right)$$

output

```
1/10*arcsech(a*x^5)^2-1/5*arcsech(a*x^5)*ln(1+(1/a/x^5+(1/a/x^5-1)^(1/2))*(1/a/x^5+1)^(1/2))^2)-1/10*polylog(2,-(1/a/x^5+(1/a/x^5-1)^(1/2)*(1/a/x^5+1)^(1/2))^2)
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.91

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = \frac{1}{10} \left( -\operatorname{sech}^{-1}(ax^5) \left( \operatorname{sech}^{-1}(ax^5) + 2 \log\left(1 + e^{-2\operatorname{sech}^{-1}(ax^5)}\right) \right) + \operatorname{PolyLog}\left(2, -e^{-2\operatorname{sech}^{-1}(ax^5)}\right) \right)$$

input

```
Integrate[ArcSech[a*x^5]/x,x]
```

output

```
(-(ArcSech[a*x^5]*(ArcSech[a*x^5] + 2*Log[1 + E^(-2*ArcSech[a*x^5])])) + PolyLog[2, -E^(-2*ArcSech[a*x^5])])/10
```

**Rubi [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 0.49 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.26, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$ , Rules used = {7282, 6835, 6297, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx \\
 & \quad \downarrow \text{7282} \\
 & \frac{1}{5} \int \frac{\operatorname{sech}^{-1}(ax^5)}{x^5} dx^5 \\
 & \quad \downarrow \text{6835} \\
 & -\frac{1}{5} \int \frac{\operatorname{arccosh}\left(\frac{1}{ax^5}\right)}{x^5} d\frac{1}{x^5} \\
 & \quad \downarrow \text{6297} \\
 & -\frac{1}{5} \int a \sqrt{\frac{\frac{1}{ax^5} - 1}{1 + \frac{1}{x^5 a}}} \left(1 + \frac{1}{x^5 a}\right) x^5 \operatorname{arccosh}\left(\frac{1}{ax^5}\right) d\operatorname{arccosh}\left(\frac{1}{ax^5}\right) \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{5} \int -i \operatorname{arccosh}\left(\frac{1}{ax^5}\right) \tan\left(i \operatorname{arccosh}\left(\frac{1}{ax^5}\right)\right) d\operatorname{arccosh}\left(\frac{1}{ax^5}\right) \\
 & \quad \downarrow \text{26} \\
 & \frac{1}{5} i \int \operatorname{arccosh}\left(\frac{1}{ax^5}\right) \tan\left(i \operatorname{arccosh}\left(\frac{1}{ax^5}\right)\right) d\operatorname{arccosh}\left(\frac{1}{ax^5}\right) \\
 & \quad \downarrow \text{4201}
 \end{aligned}$$

$$\frac{1}{5}i \left( 2i \int \frac{e^{2\operatorname{arccosh}\left(\frac{1}{ax^5}\right)} \operatorname{arccosh}\left(\frac{1}{ax^5}\right)}{1 + e^{2\operatorname{arccosh}\left(\frac{1}{ax^5}\right)}} d\operatorname{arccosh}\left(\frac{1}{ax^5}\right) - \frac{ix^{10}}{2} \right)$$

↓ 2620

$$\frac{1}{5}i \left( 2i \left( \frac{1}{2} \operatorname{arccosh}\left(\frac{1}{ax^5}\right) \log\left(e^{2\operatorname{arccosh}\left(\frac{1}{ax^5}\right)} + 1\right) - \frac{1}{2} \int \log\left(1 + e^{2\operatorname{arccosh}\left(\frac{1}{ax^5}\right)}\right) d\operatorname{arccosh}\left(\frac{1}{ax^5}\right) \right) - \frac{ix^{10}}{2} \right)$$

↓ 2715

$$\frac{1}{5}i \left( 2i \left( \frac{1}{2} \operatorname{arccosh}\left(\frac{1}{ax^5}\right) \log\left(e^{2\operatorname{arccosh}\left(\frac{1}{ax^5}\right)} + 1\right) - \frac{1}{4} \int e^{2\operatorname{arccosh}\left(\frac{1}{ax^5}\right)} \log\left(1 + e^{2\operatorname{arccosh}\left(\frac{1}{ax^5}\right)}\right) de^{2\operatorname{arccosh}\left(\frac{1}{ax^5}\right)} \right) \right)$$

↓ 2838

$$\frac{1}{5}i \left( 2i \left( \frac{1}{4} \operatorname{PolyLog}\left(2, -e^{2\operatorname{arccosh}\left(\frac{1}{ax^5}\right)}\right) + \frac{1}{2} \operatorname{arccosh}\left(\frac{1}{ax^5}\right) \log\left(e^{2\operatorname{arccosh}\left(\frac{1}{ax^5}\right)} + 1\right) \right) - \frac{ix^{10}}{2} \right)$$

input `Int[ArcSech[a*x^5]/x, x]`

output `(I/5)*((-1/2*I)*x^10 + (2*I)*((ArcCosh[1/(a*x^5)]*Log[1 + E^(2*ArcCosh[1/(a*x^5)])]))/2 + PolyLog[2, -E^(2*ArcCosh[1/(a*x^5)])]/4)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F_x), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^((n_) + ((c_) + (d_)*(x_))^(m_)))/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^((n_))), x_Symbol] := Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4201 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x
_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[
(c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x))), x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6297 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(x_), x_Symbol] :> Simp[1/b
Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a
, b, c}, x] && IGtQ[n, 0]`

rule 6835 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))/(x_), x_Symbol] :> -Subst[Int[(a +
b*ArcCosh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]`

rule 7282 `Int[(u_)/(x_), x_Symbol] :> With[{lst = PowerVariableExpn[u, 0, x]}, Simp[1
/lst[[2]] Subst[Int[NormalizeIntegrand[Simplify[lst[[1]]/x], x], x], x, (
lst[[3]]*x)^lst[[2]]], x] /; !FalseQ[lst] && NeQ[lst[[2]], 0] /; NonsumQ[
u] && !RationalFunctionQ[u, x]`

**Maple [F]**

$$\int \frac{\operatorname{arcsech}(ax^5)}{x} dx$$

input `int(arcsech(a*x^5)/x,x)`

output `int(arcsech(a*x^5)/x,x)`

**Fricas [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arsech}(ax^5)}{x} dx$$

input `integrate(arcsech(a*x^5)/x,x, algorithm="fricas")`

output `integral(arcsech(a*x^5)/x, x)`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{asech}(ax^5)}{x} dx$$

input `integrate(asech(a*x**5)/x,x)`

output `Integral(asech(a*x**5)/x, x)`

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arsech}(ax^5)}{x} dx$$

input `integrate(arcsech(a*x^5)/x,x, algorithm="maxima")`

output `integrate(arcsech(a*x^5)/x, x)`

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arsech}(ax^5)}{x} dx$$

input `integrate(arcsech(a*x^5)/x,x, algorithm="giac")`

output `integrate(arcsech(a*x^5)/x, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{ax^5}\right)}{x} dx$$

input `int(acosh(1/(a*x^5))/x,x)`

output `int(acosh(1/(a*x^5))/x, x)`

Reduce [F]

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{asech}(ax^5)}{x} dx$$

input `int(asech(a*x^5)/x,x)`

output `int(asech(a*x**5)/x,x)`



### 3.31 $\int \operatorname{sech}^{-1}(ce^{a+bx}) dx$

Optimal result	264
Mathematica [B] (verified)	264
Rubi [C] (warning: unable to verify)	265
Maple [A] (verified)	268
Fricas [F(-2)]	268
Sympy [F]	269
Maxima [F]	269
Giac [F]	270
Mupad [F(-1)]	270
Reduce [F]	270

#### Optimal result

Integrand size = 10, antiderivative size = 77

$$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx = \frac{\operatorname{sech}^{-1}(ce^{a+bx})^2}{2b} - \frac{\operatorname{sech}^{-1}(ce^{a+bx}) \log\left(1 + e^{2\operatorname{sech}^{-1}(ce^{a+bx})}\right)}{b} - \frac{\operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(ce^{a+bx})}\right)}{2b}$$

output

```
1/2*arcsech(c*exp(b*x+a))^2/b-arcsech(c*exp(b*x+a))*ln(1+(1/c/(exp(1)^(b*x+a)))+(1/c/(exp(1)^(b*x+a))-1)^(1/2)*(1/c/(exp(1)^(b*x+a))+1)^(1/2))^2)/b-1/2*polylog(2,-(1/c/(exp(1)^(b*x+a)))+(1/c/(exp(1)^(b*x+a))-1)^(1/2)*(1/c/(exp(1)^(b*x+a))+1)^(1/2))^2)/b
```

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 249 vs. 2(77) = 154.

Time = 1.12 (sec) , antiderivative size = 249, normalized size of antiderivative = 3.23

$$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx = x\operatorname{sech}^{-1}(ce^{a+bx}) - \frac{\sqrt{\frac{1-ce^{a+bx}}{1+ce^{a+bx}}}\sqrt{1+ce^{a+bx}}\left(\operatorname{arctanh}\left(\sqrt{1-c^2e^{2(a+bx)}}\right)\right)(8bx-4\log(c^2e^{2(a+bx)}))-\log^2(c^2e^{2(a+bx)})+4\log(c^2e^{2(a+bx)})}{2b}$$

input `Integrate[ArcSech[c*E^(a + b*x)],x]`

output `x*ArcSech[c*E^(a + b*x)] - (Sqrt[(1 - c*E^(a + b*x))/(1 + c*E^(a + b*x))]*  
Sqrt[1 + c*E^(a + b*x)]*(ArcTanh[Sqrt[1 - c^2*E^(2*(a + b*x))]]*(8*b*x - 4  
*Log[c^2*E^(2*(a + b*x))]) - Log[c^2*E^(2*(a + b*x))]^2 + 4*Log[c^2*E^(2*(  
a + b*x))]*Log[(1 + Sqrt[1 - c^2*E^(2*(a + b*x))])/2] - 2*Log[(1 + Sqrt[1  
- c^2*E^(2*(a + b*x))])/2]^2 + 4*PolyLog[2, (1 - Sqrt[1 - c^2*E^(2*(a + b*  
x))])/2]))/(8*b*Sqrt[1 - c*E^(a + b*x)])`

### Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.50 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.26,  
number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$ , Rules  
used = {2720, 6835, 6297, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the  
transformation is given above next to the arrow. The rules definitions used are listed  
below.

$$\begin{aligned}
 & \int \operatorname{sech}^{-1}(ce^{a+bx}) dx \\
 & \quad \downarrow \text{2720} \\
 & \frac{\int e^{-a-bx} \operatorname{sech}^{-1}(ce^{a+bx}) de^{a+bx}}{b} \\
 & \quad \downarrow \text{6835} \\
 & -\frac{\int e^{-a-bx} \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right) de^{-a-bx}}{b} \\
 & \quad \downarrow \text{6297} \\
 & -\frac{\int ce^{a+bx} \sqrt{\frac{\frac{e^{-a-bx}}{c}-1}{1+\frac{e^{-a-bx}}{c}}}}{\left(1+\frac{e^{-a-bx}}{c}\right)} \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right) d\operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right)}{b} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{\int -i \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right) \tan\left(i \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right)\right) d \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right)}{b}$$

↓ 26

$$\frac{i \int \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right) \tan\left(i \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right)\right) d \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right)}{b}$$

↓ 4201

$$\frac{i \left( 2i \int \frac{e^{\frac{a+bx+2 \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right)}}}{1+e^{2 \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right)}} d \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right) - \frac{1}{2} i e^{2a+2bx} \right)}{b}$$

↓ 2620

$$\frac{i \left( 2i \left( \frac{1}{2} \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right) \log\left(e^{2 \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right)} + 1\right) - \frac{1}{2} \int \log\left(1 + e^{2 \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right)}\right) d \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right) \right) - \frac{1}{2} i e^{2a+2bx} \right)}{b}$$

↓ 2715

$$\frac{i \left( 2i \left( \frac{1}{2} \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right) \log\left(e^{2 \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right)} + 1\right) - \frac{1}{4} \int e^{2 \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right)} \log\left(1 + e^{2 \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right)}\right) d e^{2 \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right)} \right) - \frac{1}{2} i e^{2a+2bx} \right)}{b}$$

↓ 2838

$$\frac{i \left( 2i \left( \frac{1}{4} \operatorname{PolyLog}\left(2, -e^{2 \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right)}\right) + \frac{1}{2} \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right) \log\left(e^{2 \operatorname{arccosh}\left(\frac{e^{-a-bx}}{c}\right)} + 1\right) \right) - \frac{1}{2} i e^{2a+2bx} \right)}{b}$$

input

```
Int[ArcSech[c*E^(a + b*x)], x]
```

output

```
(I*((-1/2*I)*E^(2*a + 2*b*x) + (2*I)*((ArcCosh[E^(-a - b*x)/c]*Log[1 + E^(2*ArcCosh[E^(-a - b*x)/c]])/2 + PolyLog[2, -E^(2*ArcCosh[E^(-a - b*x)/c]])/4)))/b
```

## Definitions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 2620  $\text{Int}[(((F_)^{((g_)*(e_) + (f_)*(x_))})^{(n_)*((c_) + (d_)*(x_))^{(m_))}/((a_) + (b_)*((F_)^{((g_)*(e_) + (f_)*(x_))})^{(n_)}), x\_Symbol] \rightarrow \text{Simp}[(c + d*x)^m/(b*f*g*n*\text{Log}[F])]*\text{Log}[1 + b*((F^{(g*(e + f*x)))^n/a}], x] - \text{Simp}[d*(m/(b*f*g*n*\text{Log}[F])) \text{Int}[(c + d*x)^{(m-1)}*\text{Log}[1 + b*((F^{(g*(e + f*x)))^n/a}], x], x] /; \text{FreeQ}\{F, a, b, c, d, e, f, g, n\}, x\} \ \&\& \ \text{IGtQ}[m, 0]$
- rule 2715  $\text{Int}[\text{Log}[(a_) + (b_)*((F_)^{((e_)*((c_) + (d_)*(x_))})^{(n_)}], x\_Symbol] \rightarrow \text{Simp}[1/(d*e*n*\text{Log}[F]) \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^{(e*(c + d*x))})^n], x] /; \text{FreeQ}\{F, a, b, c, d, e, n\}, x\} \ \&\& \ \text{GtQ}[a, 0]$
- rule 2720  $\text{Int}[u_, x\_Symbol] \rightarrow \text{With}\{v = \text{FunctionOfExponential}[u, x]\}, \text{Simp}[v/D[v, x] \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x] /; \text{FunctionOfExponentialQ}[u, x] \ \&\& \ !\text{MatchQ}[u, (w_)*((a_)*(v_)^{(n_)})^{(m_)} /; \text{FreeQ}\{a, m, n\}, x\} \ \&\& \ \text{IntegerQ}[m*n] \ \&\& \ !\text{MatchQ}[u, E^{((c_)*((a_) + (b_)*x))* (F_)[v_]} /; \text{FreeQ}\{a, b, c\}, x\} \ \&\& \ \text{InverseFunctionQ}[F[x]]]$
- rule 2838  $\text{Int}[\text{Log}[(c_)*((d_) + (e_)*(x_)^{(n_)}]/(x_), x\_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n]/n, x] /; \text{FreeQ}\{c, d, e, n\}, x\} \ \&\& \ \text{EqQ}[c*d, 1]$
- rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4201  $\text{Int}[((c_) + (d_)*(x_))^{(m_)*\tan[(e_) + (\text{Complex}[0, fz_])*(f_)*(x_)], x\_Symbol] \rightarrow \text{Simp}[(-I)*((c + d*x)^{(m+1)}/(d*(m+1))), x] + \text{Simp}[2*I \text{Int}[(c + d*x)^m*(E^{(2*((-I)*e + f*fz*x))}/(1 + E^{(2*((-I)*e + f*fz*x))})), x], x] /; \text{FreeQ}\{c, d, e, f, fz\}, x\} \ \&\& \ \text{IGtQ}[m, 0]$

rule 6297 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(x_), x_Symbol] := Simp[1/b Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

rule 6835 `Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))/(x_), x_Symbol] := -Subst[Int[(a + b*ArcCosh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]`

### Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.75

method	result
derivativedivides	$\frac{\frac{\operatorname{arcsech}\left(\frac{e^{bx+a}c}{2}\right)^2}{2} - \operatorname{arcsech}(e^{bx+a}c) \ln\left(1 + \left(\frac{e^{-bx-a}}{c} + \sqrt{\frac{e^{-bx-a}}{c} - 1} \sqrt{\frac{e^{-bx-a}}{c} + 1}\right)^2\right)}{b} - \frac{\operatorname{polylog}\left(2, -\left(\frac{e^{-bx-a}}{c} + \sqrt{\frac{e^{-bx-a}}{c} - 1} \sqrt{\frac{e^{-bx-a}}{c} + 1}\right)\right)}{2}$
default	$\frac{\frac{\operatorname{arcsech}\left(\frac{e^{bx+a}c}{2}\right)^2}{2} - \operatorname{arcsech}(e^{bx+a}c) \ln\left(1 + \left(\frac{e^{-bx-a}}{c} + \sqrt{\frac{e^{-bx-a}}{c} - 1} \sqrt{\frac{e^{-bx-a}}{c} + 1}\right)^2\right)}{b} - \frac{\operatorname{polylog}\left(2, -\left(\frac{e^{-bx-a}}{c} + \sqrt{\frac{e^{-bx-a}}{c} - 1} \sqrt{\frac{e^{-bx-a}}{c} + 1}\right)\right)}{2}$

input `int(arcsech(exp(b*x+a)*c), x, method=_RETURNVERBOSE)`

output `1/b*(1/2*arcsech(exp(b*x+a)*c)^2-arcsech(exp(b*x+a)*c)*ln(1+(1/exp(b*x+a)/c+(1/exp(b*x+a)/c-1)^(1/2)*(1/exp(b*x+a)/c+1)^(1/2))^2)-1/2*polylog(2,-(1/exp(b*x+a)/c+(1/exp(b*x+a)/c-1)^(1/2)*(1/exp(b*x+a)/c+1)^(1/2))^2)`

### Fricas [F(-2)]

Exception generated.

$$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx = \text{Exception raised: TypeError}$$

input `integrate(arcsech(c*exp(b*x+a)), x, algorithm="fricas")`

output Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

### Sympy [F]

$$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx = \int \operatorname{asech}(ce^{a+bx}) dx$$

input `integrate(asech(c*exp(b*x+a)), x)`

output `Integral(asech(c*exp(a + b*x)), x)`

### Maxima [F]

$$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx = \int \operatorname{arsech}(ce^{(bx+a)}) dx$$

input `integrate(arcsech(c*exp(b*x+a)), x, algorithm="maxima")`

output `b*c^2*integrate(x*e^(2*b*x + 2*a)/(c^2*e^(2*b*x + 2*a) + (c^2*e^(2*b*x + 2*a) - 1)*e^(1/2*log(c*e^(b*x + a) + 1) + 1/2*log(-c*e^(b*x + a) + 1)) - 1), x) - 1/2*b*x^2 - (a + log(c))*x + x*log(sqrt(c*e^(b*x + a) + 1)*sqrt(-c*e^(b*x + a) + 1) + 1) - 1/2*(b*x*log(c*e^(b*x + a) + 1) + dilog(-c*e^(b*x + a)))/b - 1/2*(b*x*log(-c*e^(b*x + a) + 1) + dilog(c*e^(b*x + a)))/b`

**Giac [F]**

$$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx = \int \operatorname{arsech}(ce^{(bx+a)}) dx$$

input `integrate(arcsech(c*exp(b*x+a)),x, algorithm="giac")`

output `integrate(arcsech(c*e^(b*x + a)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx = \int \operatorname{acosh}\left(\frac{e^{-a-bx}}{c}\right) dx$$

input `int(acosh(exp(- a - b*x)/c),x)`

output `int(acosh(exp(- a - b*x)/c), x)`

**Reduce [F]**

$$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx = \int \operatorname{asech}(e^{bx+a}c) dx$$

input `int(asech(c*exp(b*x+a)),x)`

output `int(asech(e**(a + b*x)*c),x)`

**3.32**  $\int \frac{\operatorname{sech}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx$

Optimal result	271
Mathematica [A] (verified)	271
Rubi [C] (warning: unable to verify)	272
Maple [A] (verified)	275
Fricas [F]	275
Sympy [F]	276
Maxima [F]	276
Giac [F]	276
Mupad [F(-1)]	277
Reduce [F]	277

**Optimal result**

Integrand size = 19, antiderivative size = 61

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \frac{\operatorname{sech}^{-1}(a+bx)^2}{2d} - \frac{\operatorname{sech}^{-1}(a+bx) \log\left(1 + e^{2\operatorname{sech}^{-1}(a+bx)}\right)}{d} - \frac{\operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right)}{2d}$$

output

```
1/2*arcsech(b*x+a)^2/d-arcsech(b*x+a)*ln(1+(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*
(1/(b*x+a)+1)^(1/2))^2)/d-1/2*polylog(2,-(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1
/(b*x+a)+1)^(1/2))^2)/d
```

**Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.85

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \frac{-\operatorname{sech}^{-1}(a+bx) \left( \operatorname{sech}^{-1}(a+bx) + 2 \log\left(1 + e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \right) + \operatorname{PolyLog}\left(2, -e^{-2\operatorname{sech}^{-1}(a+bx)}\right)}{2d}$$



input `Integrate[ArcSech[a + b*x]/((a*d)/b + d*x), x]`

output `(-(ArcSech[a + b*x]*(ArcSech[a + b*x] + 2*Log[1 + E^(-2*ArcSech[a + b*x])])  
) + PolyLog[2, -E^(-2*ArcSech[a + b*x])])/(2*d)`

### Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.53 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.10, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.526$ , Rules used = {6873, 27, 6835, 6297, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx \\
 & \quad \downarrow \text{6873} \\
 & \int \frac{b \operatorname{sech}^{-1}(a + bx) d(a + bx)}{d(a + bx)} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{\operatorname{sech}^{-1}(a + bx) d(a + bx)}{a + bx} \\
 & \quad \downarrow \text{6835} \\
 & \int (a + bx) \operatorname{arccosh}\left(\frac{1}{a + bx}\right) d\frac{1}{a + bx} \\
 & \quad \downarrow \text{6297} \\
 & \int (a + bx) \sqrt{\frac{\frac{1}{a + bx} - 1}{1 + \frac{1}{a + bx}}} \left(1 + \frac{1}{a + bx}\right) \operatorname{arccosh}\left(\frac{1}{a + bx}\right) d\operatorname{arccosh}\left(\frac{1}{a + bx}\right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{\int -i \operatorname{arccosh}\left(\frac{1}{a+bx}\right) \tan\left(i \operatorname{arccosh}\left(\frac{1}{a+bx}\right)\right) d \operatorname{arccosh}\left(\frac{1}{a+bx}\right)}{d} \\
& \quad \downarrow \text{26} \\
& \frac{i \int \operatorname{arccosh}\left(\frac{1}{a+bx}\right) \tan\left(i \operatorname{arccosh}\left(\frac{1}{a+bx}\right)\right) d \operatorname{arccosh}\left(\frac{1}{a+bx}\right)}{d} \\
& \quad \downarrow \text{4201} \\
& \frac{i \left( 2i \int \frac{e^{2 \operatorname{arccosh}\left(\frac{1}{a+bx}\right)} \operatorname{arccosh}\left(\frac{1}{a+bx}\right)}{1+e^{2 \operatorname{arccosh}\left(\frac{1}{a+bx}\right)}} d \operatorname{arccosh}\left(\frac{1}{a+bx}\right) - \frac{i}{2(a+bx)^2} \right)}{d} \\
& \quad \downarrow \text{2620} \\
& \frac{i \left( 2i \left( \frac{1}{2} \operatorname{arccosh}\left(\frac{1}{a+bx}\right) \log\left(e^{2 \operatorname{arccosh}\left(\frac{1}{a+bx}\right)} + 1\right) - \frac{1}{2} \int \log\left(1 + e^{2 \operatorname{arccosh}\left(\frac{1}{a+bx}\right)}\right) d \operatorname{arccosh}\left(\frac{1}{a+bx}\right) \right) - \frac{i}{2(a+bx)^2} \right)}{d} \\
& \quad \downarrow \text{2715} \\
& \frac{i \left( 2i \left( \frac{1}{2} \operatorname{arccosh}\left(\frac{1}{a+bx}\right) \log\left(e^{2 \operatorname{arccosh}\left(\frac{1}{a+bx}\right)} + 1\right) - \frac{1}{4} \int (a+bx) \log\left(1 + e^{2 \operatorname{arccosh}\left(\frac{1}{a+bx}\right)}\right) d e^{2 \operatorname{arccosh}\left(\frac{1}{a+bx}\right)} \right) - \frac{i}{2(a+bx)^2} \right)}{d} \\
& \quad \downarrow \text{2838} \\
& \frac{i \left( 2i \left( \frac{1}{2} \operatorname{arccosh}\left(\frac{1}{a+bx}\right) \log\left(e^{2 \operatorname{arccosh}\left(\frac{1}{a+bx}\right)} + 1\right) + \frac{1}{4} \operatorname{PolyLog}(2, -a - bx) \right) - \frac{i}{2(a+bx)^2} \right)}{d}
\end{aligned}$$

input `Int[ArcSech[a + b*x]/((a*d)/b + d*x), x]`

output `(I*((-1/2*I)/(a + b*x)^2 + (2*I)*((ArcCosh[(a + b*x)^(-1)]*Log[1 + E^(2*ArcCosh[(a + b*x)^(-1)])])/2 + PolyLog[2, -a - b*x]/4)))/d`

## Definitions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 27  $\text{Int}[(a)*(F_x), x\_Symbol] \rightarrow \text{Simp}[a \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b)*(G_x)] /; \text{FreeQ}[b, x]$
- rule 2620  $\text{Int}[(((F_)^\wedge((g_)*(e_) + (f_)*(x_)))^\wedge(n_))*((c_) + (d_)*(x_))^\wedge(m_)] / ((a_) + (b_)*((F_)^\wedge((g_)*(e_) + (f_)*(x_)))^\wedge(n_)), x\_Symbol] \rightarrow \text{Simp}[(c + d*x)^\wedge m / (b*f*g*n*\text{Log}[F])]*\text{Log}[1 + b*((F)^\wedge(g*(e + f*x)))^\wedge n/a], x] - \text{Simp}[d*(m/(b*f*g*n*\text{Log}[F])) \text{Int}[(c + d*x)^\wedge(m - 1)*\text{Log}[1 + b*((F)^\wedge(g*(e + f*x)))^\wedge n/a], x], x] /; \text{FreeQ}\{F, a, b, c, d, e, f, g, n\}, x] \ \&\& \ \text{IGtQ}[m, 0]$
- rule 2715  $\text{Int}[\text{Log}[(a_) + (b_)*((F_)^\wedge((e_)*((c_) + (d_)*(x_)))^\wedge(n_)]], x\_Symbol] \rightarrow \text{Simp}[1/(d*e*n*\text{Log}[F]) \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F)^\wedge(e*(c + d*x))^\wedge n], x] /; \text{FreeQ}\{F, a, b, c, d, e, n\}, x] \ \&\& \ \text{GtQ}[a, 0]$
- rule 2838  $\text{Int}[\text{Log}[(c_)*((d_) + (e_)*(x_))^\wedge(n_)]/(x_), x\_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^\wedge n]/n, x] /; \text{FreeQ}\{c, d, e, n\}, x] \ \&\& \ \text{EqQ}[c*d, 1]$
- rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4201  $\text{Int}[((c_) + (d_)*(x_))^\wedge(m_)*\text{tan}[(e_) + (\text{Complex}[0, fz_])*(f_)*(x_)], x\_Symbol] \rightarrow \text{Simp}[(-I)*((c + d*x)^\wedge(m + 1)/(d*(m + 1))), x] + \text{Simp}[2*I \text{Int}[(c + d*x)^\wedge m*(E^\wedge(2*((-I)*e + f*fz*x)))/(1 + E^\wedge(2*((-I)*e + f*fz*x)))]], x] /; \text{FreeQ}\{c, d, e, f, fz\}, x] \ \&\& \ \text{IGtQ}[m, 0]$
- rule 6297  $\text{Int}[((a_) + \text{ArcCosh}[(c_)*(x_)]*(b_))^\wedge(n_)/(x_), x\_Symbol] \rightarrow \text{Simp}[1/b \text{Subst}[\text{Int}[x^\wedge n*\text{Tanh}[-a/b + x/b], x], x, a + b*\text{ArcCosh}[c*x]], x] /; \text{FreeQ}\{a, b, c\}, x] \ \&\& \ \text{IGtQ}[n, 0]$

rule 6835 `Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))/(x_), x_Symbol] := -Subst[Int[(a + b*ArcCosh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]`

rule 6873 `Int[((a_.) + ArcSech[(c_) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcSech[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]`

## Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.82

method	result
derivativedivides	$\frac{\frac{b \operatorname{arcsech}(bx+a)^2}{2d} - \frac{b \operatorname{arcsech}(bx+a) \ln\left(1 + \left(\frac{1}{bx+a} + \sqrt{\frac{1}{bx+a} - 1} \sqrt{\frac{1}{bx+a} + 1}\right)^2\right)}{d}}{b} - \frac{b \operatorname{polylog}\left(2, -\left(\frac{1}{bx+a} + \sqrt{\frac{1}{bx+a} - 1} \sqrt{\frac{1}{bx+a} + 1}\right)^2\right)}{2d}$
default	$\frac{\frac{b \operatorname{arcsech}(bx+a)^2}{2d} - \frac{b \operatorname{arcsech}(bx+a) \ln\left(1 + \left(\frac{1}{bx+a} + \sqrt{\frac{1}{bx+a} - 1} \sqrt{\frac{1}{bx+a} + 1}\right)^2\right)}{d}}{b} - \frac{b \operatorname{polylog}\left(2, -\left(\frac{1}{bx+a} + \sqrt{\frac{1}{bx+a} - 1} \sqrt{\frac{1}{bx+a} + 1}\right)^2\right)}{2d}$

input `int(arcsech(b*x+a)/(a*d/b+d*x),x,method=_RETURNVERBOSE)`

output `1/b*(1/2/d*b*arcsech(b*x+a)^2-1/d*b*arcsech(b*x+a)*ln(1+(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)-1/2/d*b*polylog(2,-(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2))`

## Fricas [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{ar} \operatorname{sech}(bx + a)}{dx + \frac{ad}{b}} dx$$

input `integrate(arcsech(b*x+a)/(a*d/b+d*x),x, algorithm="fricas")`

output `integral(b*arcsech(b*x + a)/(b*d*x + a*d), x)`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \frac{b \int \frac{\operatorname{asech}\left(\frac{a+bx}{a+bx}\right) dx}{d}}$$

input `integrate(asech(b*x+a)/(a*d/b+d*x), x)`

output `b*Integral(asech(a + b*x)/(a + b*x), x)/d`

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{arsech}\left(\frac{bx + a}{d}\right)}{dx + \frac{ad}{b}} dx$$

input `integrate(arcsech(b*x+a)/(a*d/b+d*x), x, algorithm="maxima")`

output `1/2*(2*log(sqrt(b*x + a + 1))*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)*log(b*x + a) - 3*log(b*x + a)^2)/d - 1/2*(log(b*x + a + 1)*log(b*x + a) + dilog(-b*x - a))/d - 1/2*(log(b*x + a)*log(-b*x - a + 1) + dilog(b*x + a))/d + integrate((b^2*x + a*b)*log(b*x + a)/(b^2*d*x^2 + 2*a*b*d*x + a^2*d + (b^2*d*x^2 + 2*a*b*d*x + a^2*d - d)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1) - d), x)`

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{arsech}\left(\frac{bx + a}{d}\right)}{dx + \frac{ad}{b}} dx$$

input `integrate(arcsech(b*x+a)/(a*d/b+d*x), x, algorithm="giac")`

output `integrate(arcsech(b*x + a)/(d*x + a*d/b), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)}{dx + \frac{ad}{b}} dx$$

input `int(acosh(1/(a + b*x))/(d*x + (a*d)/b), x)`output `int(acosh(1/(a + b*x))/(d*x + (a*d)/b), x)`**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \frac{\left(\int \frac{\operatorname{asech}(bx+a)}{bx+a} dx\right) b}{d}$$

input `int(asech(b*x+a)/(a*d/b+d*x), x)`output `(int(asech(a + b*x)/(a + b*x), x)*b)/d`

### 3.33 $\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx$

Optimal result	278
Mathematica [B] (verified)	278
Rubi [A] (verified)	279
Maple [A] (verified)	281
Fricas [B] (verification not implemented)	281
Sympy [F]	282
Maxima [A] (verification not implemented)	282
Giac [F]	282
Mupad [B] (verification not implemented)	283
Reduce [F]	283

#### Optimal result

Integrand size = 12, antiderivative size = 49

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx = \frac{(a + bx^4) \operatorname{sech}^{-1}(a + bx^4)}{4b} - \frac{\arctan\left(\sqrt{-1 + \frac{2}{1+a+bx^4}}\right)}{2b}$$

output `1/4*(b*x^4+a)*arcsech(b*x^4+a)/b-1/2*arctan((-1+2/(b*x^4+a+1))^(1/2))/b`

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 106 vs. 2(49) = 98.

Time = 0.11 (sec) , antiderivative size = 106, normalized size of antiderivative = 2.16

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx = \frac{(a + bx^4) \operatorname{sech}^{-1}(a + bx^4) + \frac{2\sqrt{-\frac{-1+a+bx^4}{1+a+bx^4}} \sqrt{1-(a+bx^4)^2} \arctan\left(\frac{\sqrt{1-(a+bx^4)^2}}{-1+a+bx^4}\right)}{-1+a+bx^4}}{4b}$$

input `Integrate[x^3*ArcSech[a + b*x^4],x]`

output

```
((a + b*x^4)*ArcSech[a + b*x^4] + (2*Sqrt[-((-1 + a + b*x^4)/(1 + a + b*x^4))]*Sqrt[1 - (a + b*x^4)^2]*ArcTan[Sqrt[1 - (a + b*x^4)^2]/(-1 + a + b*x^4)])/(-1 + a + b*x^4)/(4*b)
```

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.14, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {7266, 6867, 2055, 27, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx$$

$$\downarrow 7266$$

$$\frac{1}{4} \int \operatorname{sech}^{-1}(bx^4 + a) dx^4$$

$$\downarrow 6867$$

$$\frac{1}{4} \left( \int \frac{\sqrt{\frac{-bx^4 - a + 1}{bx^4 + a + 1}}}{-bx^4 - a + 1} dx^4 + \frac{(a + bx^4) \operatorname{sech}^{-1}(a + bx^4)}{b} \right)$$

$$\downarrow 2055$$

$$\frac{1}{4} \left( \frac{(a + bx^4) \operatorname{sech}^{-1}(a + bx^4)}{b} - 4b \int \frac{1}{2b^2 (x^8 + 1)} d\sqrt{\frac{-bx^4 - a + 1}{bx^4 + a + 1}} \right)$$

$$\downarrow 27$$

$$\frac{1}{4} \left( \frac{(a + bx^4) \operatorname{sech}^{-1}(a + bx^4)}{b} - \frac{2 \int \frac{1}{x^8 + 1} d\sqrt{\frac{-bx^4 - a + 1}{bx^4 + a + 1}}}{b} \right)$$

$$\downarrow 216$$

$$\frac{1}{4} \left( \frac{(a + bx^4) \operatorname{sech}^{-1}(a + bx^4)}{b} - \frac{2 \arctan \left( \sqrt{\frac{-a - bx^4 + 1}{a + bx^4 + 1}} \right)}{b} \right)$$



input `Int[x^3*ArcSech[a + b*x^4],x]`

output `((a + b*x^4)*ArcSech[a + b*x^4])/b - (2*ArcTan[Sqrt[(1 - a - b*x^4)/(1 + a + b*x^4)]])/b/4`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 2055 `Int[(u_)^(r_.)*(((e_.)*((a_.) + (b_.)*(x_)^(n_.)))/((c_) + (d_.)*(x_)^(n_.)))^(p_), x_Symbol] := With[{q = Denominator[p]}, Simp[q*e*((b*c - a*d)/n) Subst[Int[SimplifyIntegrand[x^(q*(p + 1) - 1)*((-a)*e + c*x^q)^(1/n - 1)/(b*e - d*x^q)^(1/n + 1)*(u /. x -> ((-a)*e + c*x^q)^(1/n)/(b*e - d*x^q)^(1/n))]^r, x], x], x, (e*((a + b*x^n)/(c + d*x^n)))^(1/q)], x] /; FreeQ[{a, b, c, d, e}, x] && PolynomialQ[u, x] && FractionQ[p] && IntegerQ[1/n] && IntegerQ[r]`

rule 6867 `Int[ArcSech[(c_) + (d_.)*(x_)], x_Symbol] := Simp[(c + d*x)*(ArcSech[c + d*x]/d), x] + Int[Sqrt[(1 - c - d*x)/(1 + c + d*x)]/(1 - c - d*x), x] /; FreeQ[{c, d}, x]`

rule 7266 `Int[(u_)*(x_)^(m_.), x_Symbol] := Simp[1/(m + 1) Subst[Int[SubstFor[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^(m + 1), u, x]`

**Maple [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.08

method	result	size
derivativedivides	$\frac{(bx^4+a) \operatorname{arcsech}(bx^4+a) - \arctan\left(\sqrt{\frac{1}{bx^4+a}-1} \sqrt{\frac{1}{bx^4+a}+1}\right)}{4b}$	53
default	$\frac{(bx^4+a) \operatorname{arcsech}(bx^4+a) - \arctan\left(\sqrt{\frac{1}{bx^4+a}-1} \sqrt{\frac{1}{bx^4+a}+1}\right)}{4b}$	53

input `int(x^3*arcsech(b*x^4+a),x,method=_RETURNVERBOSE)`

output  $\frac{1}{4} \frac{1}{b} \left( (bx^4+a) \operatorname{arcsech}(bx^4+a) - \arctan\left(\sqrt{\frac{1}{bx^4+a}-1} \sqrt{\frac{1}{bx^4+a}+1}\right) \right)$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 283 vs. 2(43) = 86.

Time = 0.12 (sec) , antiderivative size = 283, normalized size of antiderivative = 5.78

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx$$

$$= \frac{2bx^4 \log\left(\frac{(bx^4+a)\sqrt{-\frac{b^2x^8+2abx^4+a^2-1}{b^2x^8+2abx^4+a^2}}+1}{bx^4+a}\right) + a \log\left(\frac{(bx^4+a)\sqrt{-\frac{b^2x^8+2abx^4+a^2-1}{b^2x^8+2abx^4+a^2}}+1}{x^4}\right) - a \log\left(\frac{(bx^4+a)\sqrt{-\frac{b^2x^8+2abx^4+a^2-1}{b^2x^8+2abx^4+a^2}}-1}{x^4}\right)}{8b}$$

input `integrate(x^3*arcsech(b*x^4+a),x, algorithm="fricas")`

output  $\frac{1}{8} \frac{1}{b} \left( 2bx^4 \log\left(\frac{(bx^4+a)\sqrt{-(b^2x^8+2abx^4+a^2-1)/(b^2x^8+2abx^4+a^2)}+1}{bx^4+a}\right) + a \log\left(\frac{(bx^4+a)\sqrt{-(b^2x^8+2abx^4+a^2-1)/(b^2x^8+2abx^4+a^2)}+1}{x^4}\right) - a \log\left(\frac{(bx^4+a)\sqrt{-(b^2x^8+2abx^4+a^2-1)/(b^2x^8+2abx^4+a^2)}-1}{x^4}\right) - 2 \arctan\left(\frac{(b^2x^8+2abx^4+a^2)\sqrt{-(b^2x^8+2abx^4+a^2-1)/(b^2x^8+2abx^4+a^2)}}{(b^2x^8+2abx^4+a^2)}\right) \right) / b$

**Sympy [F]**

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx = \int x^3 \operatorname{asech}(a + bx^4) dx$$

input `integrate(x**3*asech(b*x**4+a), x)`

output `Integral(x**3*asech(a + b*x**4), x)`

**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.78

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx = \frac{(bx^4 + a) \operatorname{arsech}(bx^4 + a) - \arctan\left(\sqrt{\frac{1}{(bx^4 + a)^2} - 1}\right)}{4b}$$

input `integrate(x^3*arcsech(b*x^4+a), x, algorithm="maxima")`

output `1/4*((b*x^4 + a)*arcsech(b*x^4 + a) - arctan(sqrt(1/(b*x^4 + a)^2 - 1)))/b`

**Giac [F]**

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx = \int x^3 \operatorname{arsech}(bx^4 + a) dx$$

input `integrate(x^3*arcsech(b*x^4+a), x, algorithm="giac")`

output `integrate(x^3*arcsech(b*x^4 + a), x)`

**Mupad [B] (verification not implemented)**

Time = 4.48 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.14

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx = \frac{\operatorname{atan}\left(\frac{1}{\sqrt{\frac{1}{bx^4+a}-1}\sqrt{\frac{1}{bx^4+a}+1}}\right)}{4b} + \frac{\operatorname{acosh}\left(\frac{1}{bx^4+a}\right)(bx^4+a)}{4b}$$

input `int(x^3*acosh(1/(a + b*x^4)),x)`output `atan(1/((1/(a + b*x^4) - 1)^(1/2)*(1/(a + b*x^4) + 1)^(1/2)))/(4*b) + (acosh(1/(a + b*x^4))*(a + b*x^4))/(4*b)`**Reduce [F]**

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx = \int \operatorname{asech}(bx^4 + a) x^3 dx$$

input `int(x^3*asech(b*x^4+a),x)`output `int(asech(a + b*x**4)*x**3,x)`

### 3.34 $\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx$

Optimal result	284
Mathematica [B] (verified)	284
Rubi [A] (verified)	285
Maple [F]	287
Fricas [B] (verification not implemented)	287
Sympy [F(-1)]	288
Maxima [A] (verification not implemented)	288
Giac [F]	288
Mupad [B] (verification not implemented)	289
Reduce [F]	289

#### Optimal result

Integrand size = 14, antiderivative size = 50

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx = \frac{(a + bx^n) \operatorname{sech}^{-1}(a + bx^n)}{bn} - \frac{2 \arctan\left(\sqrt{-1 + \frac{2}{1+bx^n}}\right)}{bn}$$

output `(a+b*x^n)*arcsech(a+b*x^n)/b/n-2*arctan((-1+2/(1+a+b*x^n))^(1/2))/b/n`

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 106 vs. 2(50) = 100.

Time = 0.14 (sec) , antiderivative size = 106, normalized size of antiderivative = 2.12

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx = \frac{(a + bx^n) \operatorname{sech}^{-1}(a + bx^n) + \frac{2\sqrt{-\frac{-1+a+bx^n}{1+a+bx^n}} \sqrt{1-(a+bx^n)^2} \arctan\left(\frac{\sqrt{1-(a+bx^n)^2}}{-1+a+bx^n}\right)}{-1+a+bx^n}}{bn}$$

input `Integrate[x^(-1 + n)*ArcSech[a + b*x^n], x]`

output

$$\frac{((a + b*x^n)*ArcSech[a + b*x^n] + (2*sqrt[-((-1 + a + b*x^n)/(1 + a + b*x^n))])*sqrt[1 - (a + b*x^n)^2])*ArcTan[sqrt[1 - (a + b*x^n)^2]/(-1 + a + b*x^n)]}{(-1 + a + b*x^n)/(b*n)}$$
**Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.12, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$ , Rules used = {7266, 6867, 2055, 27, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^{n-1} \operatorname{sech}^{-1}(a + bx^n) dx \\ & \quad \downarrow \text{7266} \\ & \frac{\int \operatorname{sech}^{-1}(bx^n + a) dx^n}{n} \\ & \quad \downarrow \text{6867} \\ & \frac{\int \frac{\sqrt{\frac{-bx^n - a + 1}{bx^n + a + 1}}}{-bx^n - a + 1} dx^n + \frac{(a + bx^n) \operatorname{sech}^{-1}(a + bx^n)}{b}}{n} \\ & \quad \downarrow \text{2055} \\ & \frac{\frac{(a + bx^n) \operatorname{sech}^{-1}(a + bx^n)}{b} - 4b \int \frac{1}{2b^2(x^{2n} + 1)} d\sqrt{\frac{-bx^n - a + 1}{bx^n + a + 1}}}{n} \\ & \quad \downarrow \text{27} \\ & \frac{\frac{(a + bx^n) \operatorname{sech}^{-1}(a + bx^n)}{b} - \frac{2 \int \frac{1}{x^{2n} + 1} d\sqrt{\frac{-bx^n - a + 1}{bx^n + a + 1}}}{b}}{n} \\ & \quad \downarrow \text{216} \\ & \frac{\frac{(a + bx^n) \operatorname{sech}^{-1}(a + bx^n)}{b} - \frac{2 \arctan\left(\sqrt{\frac{-a - bx^n + 1}{a + bx^n + 1}}\right)}{b}}{n} \end{aligned}$$

input

$$\text{Int}[x^{(-1 + n)} * \text{ArcSech}[a + b*x^n], x]$$

output 
$$\frac{((a + b*x^n)*\text{ArcSech}[a + b*x^n])/b - (2*\text{ArcTan}[\text{Sqrt}[(1 - a - b*x^n)/(1 + a + b*x^n)])]/b}{n}$$

### Defintions of rubi rules used

rule 27 
$$\text{Int}[(a_)*(F_x_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_) /; \text{FreeQ}[b, x]]$$

rule 216 
$$\text{Int}[(a_) + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$$

rule 2055 
$$\text{Int}[(u_)^{(r_)*((e_)*((a_.) + (b_.)*(x_)^{(n_.)}))}/((c_) + (d_.)*(x_)^{(n_.)})^{(p_)}, x\_Symbol] \rightarrow \text{With}\{q = \text{Denominator}[p]\}, \text{Simp}[q*e*((b*c - a*d)/n) \text{ Subst}[\text{Int}[\text{SimplifyIntegrand}[x^{(q*(p + 1) - 1)}*(((-a)*e + c*x^q)^{(1/n - 1)}/(b*e - d*x^q)^{(1/n + 1)})*(u /. x \rightarrow ((-a)*e + c*x^q)^{(1/n)}/(b*e - d*x^q)^{(1/n)})^r, x], x], x, (e*((a + b*x^n)/(c + d*x^n)))^{(1/q)}, x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \ \&\& \ \text{PolynomialQ}[u, x] \ \&\& \ \text{FractionQ}[p] \ \&\& \ \text{IntegerQ}[1/n] \ \&\& \ \text{IntegerQ}[r]$$

rule 6867 
$$\text{Int}[\text{ArcSech}[(c_) + (d_.)*(x_)], x\_Symbol] \rightarrow \text{Simp}[(c + d*x)*(\text{ArcSech}[c + d*x]/d), x] + \text{Int}[\text{Sqrt}[(1 - c - d*x)/(1 + c + d*x)]/(1 - c - d*x), x] /; \text{FreeQ}\{c, d\}, x]$$

rule 7266 
$$\text{Int}[(u_)*(x_)^{(m_.)}, x\_Symbol] \rightarrow \text{Simp}[1/(m + 1) \text{ Subst}[\text{Int}[\text{SubstFor}[x^{(m + 1)}, u, x], x], x, x^{(m + 1)}], x] /; \text{FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1] \ \&\& \ \text{FunctionOfQ}[x^{(m + 1)}, u, x]$$

**Maple [F]**

$$\int x^{-1+n} \operatorname{arcsech}(a + b x^n) dx$$

input `int(x^(-1+n)*arcsech(a+b*x^n),x)`

output `int(x^(-1+n)*arcsech(a+b*x^n),x)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 385 vs.  $2(48) = 96$ .

Time = 0.14 (sec) , antiderivative size = 385, normalized size of antiderivative = 7.70

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + b x^n) dx$$

$$= \frac{2(b \cosh(n \log(x)) + b \sinh(n \log(x))) \log\left(\frac{\sqrt{-\frac{2ab + (a^2 + b^2 - 1) \cosh(n \log(x)) - (a^2 - b^2 - 1) \sinh(n \log(x))}{\cosh(n \log(x)) - \sinh(n \log(x))} + 1}}{b \cosh(n \log(x)) + b \sinh(n \log(x)) + a}\right) + a \log\left(\frac{\sqrt{-\frac{2ab + (a^2 + b^2 - 1) \cosh(n \log(x)) - (a^2 - b^2 - 1) \sinh(n \log(x))}{\cosh(n \log(x)) - \sinh(n \log(x))} + 1}}{b \cosh(n \log(x)) + b \sinh(n \log(x)) + a}\right)}{b \cosh(n \log(x)) + b \sinh(n \log(x)) + a} + a \log\left(\frac{\sqrt{-\frac{2ab + (a^2 + b^2 - 1) \cosh(n \log(x)) - (a^2 - b^2 - 1) \sinh(n \log(x))}{\cosh(n \log(x)) - \sinh(n \log(x))} + 1}}{b \cosh(n \log(x)) + b \sinh(n \log(x)) + a}\right)}{b \cosh(n \log(x)) + b \sinh(n \log(x)) + a}$$

input `integrate(x^(-1+n)*arcsech(a+b*x^n),x, algorithm="fricas")`

output `1/2*(2*(b*cosh(n*log(x)) + b*sinh(n*log(x)))*log((sqrt(-(2*a*b + (a^2 + b^2 - 1)*cosh(n*log(x)) - (a^2 - b^2 - 1)*sinh(n*log(x)))/(cosh(n*log(x)) - sinh(n*log(x)))) + 1)/(b*cosh(n*log(x)) + b*sinh(n*log(x)) + a)) + a*log((sqrt(-(2*a*b + (a^2 + b^2 - 1)*cosh(n*log(x)) - (a^2 - b^2 - 1)*sinh(n*log(x)))/(cosh(n*log(x)) - sinh(n*log(x)))) + 1)/(cosh(n*log(x)) + sinh(n*log(x)))) - a*log((sqrt(-(2*a*b + (a^2 + b^2 - 1)*cosh(n*log(x)) - (a^2 - b^2 - 1)*sinh(n*log(x)))/(cosh(n*log(x)) - sinh(n*log(x)))) - 1)/(cosh(n*log(x)) + sinh(n*log(x)))) - 2*arctan((b*cosh(n*log(x)) + b*sinh(n*log(x)) + a)*sqrt(-(2*a*b + (a^2 + b^2 - 1)*cosh(n*log(x)) - (a^2 - b^2 - 1)*sinh(n*log(x)))/(cosh(n*log(x)) - sinh(n*log(x))))/(b^2*cosh(n*log(x))^2 + b^2*sinh(n*log(x))^2 + 2*a*b*cosh(n*log(x)) + a^2 + 2*(b^2*cosh(n*log(x)) + a*b)*sinh(n*log(x)) - 1)))/(b*n)`



**Sympy [F(-1)]**

Timed out.

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx = \text{Timed out}$$

input `integrate(x**(-1+n)*asech(a+b*x**n), x)`

output `Timed out`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.80

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx = \frac{(bx^n + a) \operatorname{ar} \operatorname{sech}(bx^n + a) - \arctan\left(\sqrt{\frac{1}{(bx^n + a)^2} - 1}\right)}{bn}$$

input `integrate(x^(-1+n)*arcsech(a+b*x^n), x, algorithm="maxima")`

output `((b*x^n + a)*arcsech(b*x^n + a) - arctan(sqrt(1/(b*x^n + a)^2 - 1)))/(b*n)`

**Giac [F]**

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx = \int x^{n-1} \operatorname{ar} \operatorname{sech}(bx^n + a) dx$$

input `integrate(x^(-1+n)*arcsech(a+b*x^n), x, algorithm="giac")`

output `integrate(x^(n - 1)*arcsech(b*x^n + a), x)`

**Mupad [B] (verification not implemented)**

Time = 3.92 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.08

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx = \frac{\operatorname{atan}\left(\frac{1}{\sqrt{\frac{1}{a+bx^n}-1}\sqrt{\frac{1}{a+bx^n}+1}}\right) + \operatorname{acosh}\left(\frac{1}{a+bx^n}\right) (a + bx^n)}{bn}$$

input `int(x^(n - 1)*acosh(1/(a + b*x^n)),x)`output `(atan(1/((1/(a + b*x^n) - 1)^(1/2)*(1/(a + b*x^n) + 1)^(1/2))) + acosh(1/(a + b*x^n))*(a + b*x^n))/(b*n)`**Reduce [F]**

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx = \int \frac{x^n \operatorname{asech}(x^n b + a)}{x} dx$$

input `int(x^(-1+n)*asech(a+b*x^n),x)`output `int((x**n*asech(x**n*b + a))/x,x)`

# CHAPTER 4

## APPENDIX

4.1	Listing of Grading functions . . . . .	290
4.2	Links to plain text integration problems used in this report for each CAS .	308

### 4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

#### Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal."}
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal."}
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order of result is higher than in optimal."}
  ]
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```
Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
If[AppellFunctionQ[Head[expn]],
Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
If[Head[expn]===RootSum,
Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
If[Head[expn]===Integrate || Head[expn]===Int,
Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
9]]]]]]]]]]]]
```

```
ElementaryFunctionQ[func_] :=
MemberQ[{
Exp, Log,
Sin, Cos, Tan, Cot, Sec, Csc,
ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
Sinh, Cosh, Tanh, Coth, Sech, Csch,
ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
}, func]
```

```
SpecialFunctionQ[func_] :=
MemberQ[{
Erf, Erfc, Erfi,
FresnelS, FresnelC,
ExpIntegralE, ExpIntegralEi, LogIntegral,
SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
Gamma, LogGamma, PolyGamma,
Zeta, PolyLog, ProductLog,
EllipticF, EllipticE, EllipticPi
}, func]
```

```
HypergeometricFunctionQ[func_] :=
MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]
```

```
AppellFunctionQ[func_] :=
MemberQ[{AppellF1}, func]
```

## Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022  add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result    := ExpnType(result);
      ExpnType_optimal   := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#     is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```



```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
        fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

## Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

## SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```



```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

## 4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file