

# Computer Algebra Independent Integration Tests

Summer 2024

7-Inverse-hyperbolic-functions/7.5-Inverse-hyperbolic-  
secant/347-7.5.1

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 10:22am

# Contents

<b>1</b>	<b>Introduction</b>	<b>9</b>
1.1	Listing of CAS systems tested . . . . .	10
1.2	Results . . . . .	11
1.3	Time and leaf size Performance . . . . .	15
1.4	Performance based on number of rules Rubi used . . . . .	17
1.5	Performance based on number of steps Rubi used . . . . .	18
1.6	Solved integrals histogram based on leaf size of result . . . . .	19
1.7	Solved integrals histogram based on CPU time used . . . . .	20
1.8	Leaf size vs. CPU time used . . . . .	21
1.9	list of integrals with no known antiderivative . . . . .	22
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	22
1.11	list of integrals solved by CAS but failed verification . . . . .	22
1.12	Timing . . . . .	23
1.13	Verification . . . . .	23
1.14	Important notes about some of the results . . . . .	24
1.15	Current tree layout of integration tests . . . . .	27
1.16	Design of the test system . . . . .	28
<b>2</b>	<b>detailed summary tables of results</b>	<b>29</b>
2.1	List of integrals sorted by grade for each CAS . . . . .	30
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	35
2.3	Detailed conclusion table specific for Rubi results . . . . .	83
<b>3</b>	<b>Listing of integrals</b>	<b>90</b>
3.1	$\int x^4 \operatorname{sech}^{-1}(ax)^2 dx$ . . . . .	97
3.2	$\int x^3 \operatorname{sech}^{-1}(ax)^2 dx$ . . . . .	104
3.3	$\int x^2 \operatorname{sech}^{-1}(ax)^2 dx$ . . . . .	111
3.4	$\int x \operatorname{sech}^{-1}(ax)^2 dx$ . . . . .	118
3.5	$\int \operatorname{sech}^{-1}(ax)^2 dx$ . . . . .	124
3.6	$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x} dx$ . . . . .	130

3.7	$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^2} dx$	137
3.8	$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^3} dx$	143
3.9	$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^4} dx$	149
3.10	$\int x^4 \operatorname{sech}^{-1}(ax)^3 dx$	156
3.11	$\int x^3 \operatorname{sech}^{-1}(ax)^3 dx$	165
3.12	$\int x^2 \operatorname{sech}^{-1}(ax)^3 dx$	174
3.13	$\int x \operatorname{sech}^{-1}(ax)^3 dx$	182
3.14	$\int \operatorname{sech}^{-1}(ax)^3 dx$	189
3.15	$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x} dx$	196
3.16	$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^2} dx$	203
3.17	$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^3} dx$	210
3.18	$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^4} dx$	217
3.19	$\int x^6 (a + b \operatorname{sech}^{-1}(cx)) dx$	225
3.20	$\int x^5 (a + b \operatorname{sech}^{-1}(cx)) dx$	233
3.21	$\int x^4 (a + b \operatorname{sech}^{-1}(cx)) dx$	240
3.22	$\int x^3 (a + b \operatorname{sech}^{-1}(cx)) dx$	247
3.23	$\int x^2 (a + b \operatorname{sech}^{-1}(cx)) dx$	253
3.24	$\int x (a + b \operatorname{sech}^{-1}(cx)) dx$	259
3.25	$\int (a + b \operatorname{sech}^{-1}(cx)) dx$	264
3.26	$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x} dx$	269
3.27	$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2} dx$	276
3.28	$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3} dx$	281
3.29	$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4} dx$	288
3.30	$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5} dx$	294
3.31	$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^6} dx$	301
3.32	$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^7} dx$	307
3.33	$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^2 dx$	315
3.34	$\int x^2 (a + b \operatorname{sech}^{-1}(cx))^2 dx$	322
3.35	$\int x (a + b \operatorname{sech}^{-1}(cx))^2 dx$	329
3.36	$\int (a + b \operatorname{sech}^{-1}(cx))^2 dx$	336
3.37	$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x} dx$	343
3.38	$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^2} dx$	351
3.39	$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^3} dx$	358

3.40	$\int \frac{(a+b\operatorname{sech}^{-1}(cx))^2}{x^4} dx$	365
3.41	$\int \frac{(a+b\operatorname{sech}^{-1}(cx))^2}{x^5} dx$	373
3.42	$\int x^3(a+b\operatorname{sech}^{-1}(cx))^3 dx$	380
3.43	$\int x^2(a+b\operatorname{sech}^{-1}(cx))^3 dx$	390
3.44	$\int x(a+b\operatorname{sech}^{-1}(cx))^3 dx$	398
3.45	$\int (a+b\operatorname{sech}^{-1}(cx))^3 dx$	406
3.46	$\int \frac{(a+b\operatorname{sech}^{-1}(cx))^3}{x} dx$	413
3.47	$\int \frac{(a+b\operatorname{sech}^{-1}(cx))^3}{x^2} dx$	421
3.48	$\int \frac{(a+b\operatorname{sech}^{-1}(cx))^3}{x^3} dx$	429
3.49	$\int \frac{(a+b\operatorname{sech}^{-1}(cx))^3}{x^4} dx$	437
3.50	$\int \frac{(a+b\operatorname{sech}^{-1}(cx))^3}{x^5} dx$	446
3.51	$\int \frac{x}{a+b\operatorname{sech}^{-1}(cx)} dx$	455
3.52	$\int \frac{1}{a+b\operatorname{sech}^{-1}(cx)} dx$	460
3.53	$\int \frac{1}{x(a+b\operatorname{sech}^{-1}(cx))} dx$	465
3.54	$\int \frac{1}{x^2(a+b\operatorname{sech}^{-1}(cx))} dx$	470
3.55	$\int \frac{1}{x^3(a+b\operatorname{sech}^{-1}(cx))} dx$	476
3.56	$\int \frac{1}{x^4(a+b\operatorname{sech}^{-1}(cx))} dx$	483
3.57	$\int \frac{x}{(a+b\operatorname{sech}^{-1}(cx))^2} dx$	489
3.58	$\int \frac{1}{(a+b\operatorname{sech}^{-1}(cx))^2} dx$	494
3.59	$\int \frac{1}{x(a+b\operatorname{sech}^{-1}(cx))^2} dx$	499
3.60	$\int \frac{1}{x^2(a+b\operatorname{sech}^{-1}(cx))^2} dx$	504
3.61	$\int \frac{1}{x^3(a+b\operatorname{sech}^{-1}(cx))^2} dx$	512
3.62	$\int \frac{1}{x^4(a+b\operatorname{sech}^{-1}(cx))^2} dx$	520
3.63	$\int \frac{x}{(a+b\operatorname{sech}^{-1}(cx))^3} dx$	526
3.64	$\int \frac{1}{(a+b\operatorname{sech}^{-1}(cx))^3} dx$	531
3.65	$\int \frac{1}{x(a+b\operatorname{sech}^{-1}(cx))^3} dx$	536

3.66	$\int \frac{1}{x^2(a+b\operatorname{sech}^{-1}(cx))^3} dx$	541
3.67	$\int \frac{1}{x^3(a+b\operatorname{sech}^{-1}(cx))^3} dx$	551
3.68	$\int \frac{1}{x^4(a+b\operatorname{sech}^{-1}(cx))^3} dx$	562
3.69	$\int (dx)^m (a + b\operatorname{sech}^{-1}(cx))^3 dx$	569
3.70	$\int (dx)^m (a + b\operatorname{sech}^{-1}(cx))^2 dx$	574
3.71	$\int (dx)^m (a + b\operatorname{sech}^{-1}(cx)) dx$	579
3.72	$\int \frac{(dx)^m}{a+b\operatorname{sech}^{-1}(cx)} dx$	584
3.73	$\int \frac{(dx)^m}{(a+b\operatorname{sech}^{-1}(cx))^2} dx$	589
3.74	$\int (d+ex)^3 (a + b\operatorname{sech}^{-1}(cx)) dx$	594
3.75	$\int (d+ex)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$	604
3.76	$\int (d+ex) (a + b\operatorname{sech}^{-1}(cx)) dx$	613
3.77	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{d+ex} dx$	621
3.78	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex)^2} dx$	628
3.79	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex)^3} dx$	635
3.80	$\int (d+ex)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx$	643
3.81	$\int \sqrt{d+ex} (a + b\operatorname{sech}^{-1}(cx)) dx$	655
3.82	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{\sqrt{d+ex}} dx$	665
3.83	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex)^{3/2}} dx$	672
3.84	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex)^{5/2}} dx$	679
3.85	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex)^{7/2}} dx$	689
3.86	$\int (d+ex)^m (a + b\operatorname{sech}^{-1}(cx)) dx$	701
3.87	$\int x^4 (d+ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$	706
3.88	$\int x^2 (d+ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$	714
3.89	$\int (d+ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$	721
3.90	$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^2} dx$	728
3.91	$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx$	735
3.92	$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx$	742
3.93	$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^8} dx$	749
3.94	$\int x^5 (d+ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$	757
3.95	$\int x^3 (d+ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$	764
3.96	$\int x (d+ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$	771

3.97	$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x} dx$	778
3.98	$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^3} dx$	785
3.99	$\int x^2(d+ex^2)^2(a+b\operatorname{sech}^{-1}(cx)) dx$	792
3.100	$\int (d+ex^2)^2(a+b\operatorname{sech}^{-1}(cx)) dx$	801
3.101	$\int \frac{(d+ex^2)^2(a+b\operatorname{sech}^{-1}(cx))}{x^2} dx$	809
3.102	$\int \frac{(d+ex^2)^2(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx$	817
3.103	$\int \frac{(d+ex^2)^2(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx$	825
3.104	$\int \frac{(d+ex^2)^2(a+b\operatorname{sech}^{-1}(cx))}{x^8} dx$	833
3.105	$\int x^3(d+ex^2)^2(a+b\operatorname{sech}^{-1}(cx)) dx$	841
3.106	$\int x(d+ex^2)^2(a+b\operatorname{sech}^{-1}(cx)) dx$	849
3.107	$\int \frac{(d+ex^2)^2(a+b\operatorname{sech}^{-1}(cx))}{x} dx$	857
3.108	$\int \frac{(d+ex^2)^2(a+b\operatorname{sech}^{-1}(cx))}{x^3} dx$	865
3.109	$\int \frac{x^2(a+b\operatorname{sech}^{-1}(cx))}{d+ex^2} dx$	872
3.110	$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{d+ex^2} dx$	881
3.111	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{d+ex^2} dx$	890
3.112	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x(d+ex^2)} dx$	898
3.113	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^2(d+ex^2)} dx$	906
3.114	$\int \frac{x^5(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^2} dx$	915
3.115	$\int \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^2} dx$	925
3.116	$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^2} dx$	935
3.117	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x(d+ex^2)^2} dx$	943
3.118	$\int \frac{x^4(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^2} dx$	952
3.119	$\int \frac{x^2(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^2} dx$	962
3.120	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex^2)^2} dx$	972
3.121	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^2(d+ex^2)^2} dx$	982
3.122	$\int \frac{x^5(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^3} dx$	992
3.123	$\int \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^3} dx$	1002

3.124	$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^3} dx$	1010
3.125	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x(d+ex^2)^3} dx$	1020
3.126	$\int \frac{x^4(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^3} dx$	1030
3.127	$\int \frac{x^2(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^3} dx$	1039
3.128	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex^2)^3} dx$	1048
3.129	$\int x^5\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1057
3.130	$\int x^3\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1069
3.131	$\int x\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1079
3.132	$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x} dx$	1088
3.133	$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^3} dx$	1093
3.134	$\int x^2\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1098
3.135	$\int \sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1103
3.136	$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^2} dx$	1108
3.137	$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx$	1113
3.138	$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx$	1122
3.139	$\int x^3(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1132
3.140	$\int x(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1143
3.141	$\int \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{x} dx$	1153
3.142	$\int \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{x^3} dx$	1158
3.143	$\int x^2(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1163
3.144	$\int (d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1168
3.145	$\int \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{x^2} dx$	1173
3.146	$\int \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx$	1178
3.147	$\int \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx$	1183
3.148	$\int \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{x^8} dx$	1193
3.149	$\int \frac{x^5(a+b\operatorname{sech}^{-1}(cx))}{\sqrt{d+ex^2}} dx$	1203
3.150	$\int \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{\sqrt{d+ex^2}} dx$	1214
3.151	$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{\sqrt{d+ex^2}} dx$	1223

3.152	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x\sqrt{d+ex^2}} dx$	1231
3.153	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^3\sqrt{d+ex^2}} dx$	1236
3.154	$\int \frac{x^2(a+b\operatorname{sech}^{-1}(cx))}{\sqrt{d+ex^2}} dx$	1241
3.155	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{\sqrt{d+ex^2}} dx$	1246
3.156	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^2\sqrt{d+ex^2}} dx$	1251
3.157	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^4\sqrt{d+ex^2}} dx$	1259
3.158	$\int \frac{x^5(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{3/2}} dx$	1269
3.159	$\int \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{3/2}} dx$	1278
3.160	$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{3/2}} dx$	1286
3.161	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x(d+ex^2)^{3/2}} dx$	1292
3.162	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^3(d+ex^2)^{3/2}} dx$	1297
3.163	$\int \frac{x^4(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{3/2}} dx$	1302
3.164	$\int \frac{x^2(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{3/2}} dx$	1307
3.165	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex^2)^{3/2}} dx$	1312
3.166	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^2(d+ex^2)^{3/2}} dx$	1318
3.167	$\int \frac{x^5(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{5/2}} dx$	1327
3.168	$\int \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{5/2}} dx$	1336
3.169	$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{5/2}} dx$	1344
3.170	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x(d+ex^2)^{5/2}} dx$	1351
3.171	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^3(d+ex^2)^{5/2}} dx$	1356
3.172	$\int \frac{x^6(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{5/2}} dx$	1361
3.173	$\int \frac{x^4(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{5/2}} dx$	1366
3.174	$\int \frac{x^2(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{5/2}} dx$	1371
3.175	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex^2)^{5/2}} dx$	1379



3.176	$\int (fx)^m (d + ex^2)^3 (a + b\operatorname{sech}^{-1}(cx)) dx$	1388
3.177	$\int (fx)^m (d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$	1398
3.178	$\int (fx)^m (d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$	1406
3.179	$\int \frac{(fx)^m (a + b\operatorname{sech}^{-1}(cx))}{d + ex^2} dx$	1413
3.180	$\int \frac{(fx)^m (a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx$	1418
3.181	$\int (fx)^m (d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx$	1423
3.182	$\int (fx)^m \sqrt{d + ex^2} (a + b\operatorname{sech}^{-1}(cx)) dx$	1428
3.183	$\int \frac{(fx)^m (a + b\operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx$	1433
3.184	$\int \frac{(fx)^m (a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$	1438
3.185	$\int \frac{x^{11} (a + b\operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx$	1443
3.186	$\int \frac{x^7 (a + b\operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx$	1451
3.187	$\int \frac{x^3 (a + b\operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx$	1459
3.188	$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x\sqrt{1 - c^4 x^4}} dx$	1466
3.189	$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^5\sqrt{1 - c^4 x^4}} dx$	1471
<b>4</b>	<b>Appendix</b>	<b>1476</b>
4.1	Listing of Grading functions	1476
4.2	Links to plain text integration problems used in this report for each CAS	494

# CHAPTER 1

## INTRODUCTION

1.1	Listing of CAS systems tested . . . . .	10
1.2	Results . . . . .	11
1.3	Time and leaf size Performance . . . . .	15
1.4	Performance based on number of rules Rubi used . . . . .	17
1.5	Performance based on number of steps Rubi used . . . . .	18
1.6	Solved integrals histogram based on leaf size of result . . . . .	19
1.7	Solved integrals histogram based on CPU time used . . . . .	20
1.8	Leaf size vs. CPU time used . . . . .	21
1.9	list of integrals with no known antiderivative . . . . .	22
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	22
1.11	list of integrals solved by CAS but failed verification . . . . .	22
1.12	Timing . . . . .	23
1.13	Verification . . . . .	23
1.14	Important notes about some of the results . . . . .	24
1.15	Current tree layout of integration tests . . . . .	27
1.16	Design of the test system . . . . .	28

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [ 189 ]. This is test number [ 347 ].

## 1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

## 1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 ( 189 )	0.00 ( 0 )
Mathematica	100.00 ( 189 )	0.00 ( 0 )
Maple	80.95 ( 153 )	19.05 ( 36 )
Fricas	68.25 ( 129 )	31.75 ( 60 )
Maxima	32.80 ( 62 )	67.20 ( 127 )
Mupad	26.98 ( 51 )	73.02 ( 138 )
Sympy	25.93 ( 49 )	74.07 ( 140 )
Giac	23.81 ( 45 )	76.19 ( 144 )
Reduce	23.81 ( 45 )	76.19 ( 144 )

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> <li>1. antiderivative contains a hypergeometric function and the optimal antiderivative does not.</li> <li>2. antiderivative contains a special function and the optimal antiderivative does not.</li> <li>3. antiderivative contains the imaginary unit and the optimal antiderivative does not.</li> </ol>
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	65.079	0.000	11.111	23.810
Mathematica	49.206	0.529	26.455	23.810
Maple	35.979	11.640	9.524	42.857
Fricas	26.455	17.989	0.000	55.556
Maxima	21.164	0.000	0.000	78.836
Sympy	5.291	0.000	0.000	94.709
Giac	0.000	0.000	0.000	100.000
Mupad	0.000	3.175	0.000	96.825
Reduce	0.000	0.000	0.000	100.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Maple	36	100.00	0.00	0.00
Fricas	60	95.00	5.00	0.00
Maxima	127	53.54	0.79	45.67
Sympy	140	82.86	17.14	0.00
Mupad	138	0.00	100.00	0.00
Giac	144	98.61	0.00	1.39
Reduce	144	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

### 1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.



System	Mean time (sec)
Giac	0.12
Fricas	0.16
Maxima	0.45
Rubi	0.64
Mupad	3.96
Maple	5.01
Mathematica	6.05
Sympy	11.36
Reduce	117.21

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Giac	20.84	1.05	23.00	1.00
Mupad	29.73	1.22	27.00	1.17
Reduce	31.78	1.73	23.00	1.00
Sympy	45.08	0.97	22.00	0.96
Rubi	196.38	0.98	132.00	1.00
Maxima	295.87	17.39	115.50	0.99
Maple	300.78	1.44	147.00	1.00
Fricas	308.99	2.08	143.00	1.50
Mathematica	379.52	1.59	140.00	1.09

Table 1.6: Leaf size performance for each CAS

## 1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the  $y$  axis is the percentage solved which Rubi itself needed the number of rules given the  $x$  axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

## 1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

## 1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

## 1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

## 1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

## 1.9 list of integrals with no known antiderivative

{51, 52, 53, 57, 58, 59, 63, 64, 65, 69, 70, 72, 73, 86, 132, 133, 134, 135, 136, 141, 142, 143, 144, 145, 146, 152, 153, 154, 155, 161, 162, 163, 164, 170, 171, 172, 173, 179, 180, 181, 182, 183, 184, 188, 189}

## 1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

## 1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {19, 25, 26, 71, 186}

**Mathematica** {77, 80, 81, 82, 83, 84, 85, 109, 110, 111, 112, 113, 114, 115, 117, 118, 119, 120, 121, 122, 125, 126, 127, 128}

**Maple** {109, 110, 112, 114, 115, 117, 118, 119, 120, 121, 122, 125, 126, 127, 128}

**Maxima** Verification phase not currently implemented.

**Fricas** Verification phase not currently implemented.

**Sympy** Verification phase not currently implemented.

**Giac** Verification phase not currently implemented.

**Reduce** Verification phase not currently implemented.

**Mupad** Verification phase not currently implemented.

## 1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

## 1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.



Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

## 1.14 Important notes about some of the results

### Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

### Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

### Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at [https://ask.sagemath.org/question/57123/could-we-have-a-leaf\\_count-function-in-base-sagemath/](https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/)

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

### Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives  $\sin(x)^2/2$

# 1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

# 1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



**High level overview of the CAS independent integration test build system**

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

*The following fields are present only in Rubi Table file*

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi  
January 13, 2024  
Design note

# CHAPTER 2

## DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS . . . . .	30
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	35
2.3	Detailed conclusion table specific for Rubi results . . . . .	83

## 2.1 List of integrals sorted by grade for each CAS

Rubi . . . . .	30
Mma . . . . .	30
Maple . . . . .	31
Fricas . . . . .	31
Maxima . . . . .	32
Giac . . . . .	32
Mupad . . . . .	33
Sympy . . . . .	33
Reduce . . . . .	34

### Rubi

**A grade** { 1, 2, 3, 4, 5, 8, 9, 10, 12, 14, 17, 19, 20, 21, 22, 23, 24, 25, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 39, 40, 41, 43, 45, 48, 50, 56, 62, 68, 71, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 137, 138, 139, 140, 147, 148, 149, 150, 151, 156, 157, 158, 159, 160, 165, 166, 167, 168, 169, 174, 175, 176, 177, 178, 185, 186, 187 }

**B grade** { }

**C grade** { 6, 7, 11, 13, 15, 16, 18, 26, 37, 38, 42, 44, 46, 47, 49, 54, 55, 60, 61, 66, 67 }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

### Mma

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 22, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 46, 47, 48, 49, 50, 54, 55, 56, 60, 61, 62, 66, 67, 68, 71, 76, 78, 90, 91, 92, 93, 94, 95, 96, 97, 98, 103, 104, 105, 106, 107, 108, 129, 130, 131, 139, 140, 149, 150, 151, 158, 159, 160, 167, 168, 169, 176, 177, 178, 185, 186, 187 }

**B grade** { 45 }

**C grade** { 19, 21, 23, 74, 75, 77, 79, 80, 81, 82, 83, 84, 85, 87, 88, 89, 99, 100, 101, 102, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 137, 138, 147, 148, 156, 157, 165, 166, 174, 175 }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Maple

**A grade** { 1, 2, 3, 5, 6, 7, 8, 9, 11, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 34, 36, 37, 39, 40, 42, 44, 48, 54, 55, 56, 60, 74, 75, 76, 78, 81, 82, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108 }

**B grade** { 4, 33, 35, 38, 41, 46, 47, 49, 50, 61, 62, 66, 67, 68, 79, 80, 83, 84, 85, 116, 123, 124 }

**C grade** { 77, 109, 110, 111, 112, 113, 114, 115, 117, 118, 119, 120, 121, 122, 125, 126, 127, 128 }

**F normal fail** { 10, 12, 14, 43, 45, 71, 129, 130, 131, 137, 138, 139, 140, 147, 148, 149, 150, 151, 156, 157, 158, 159, 160, 165, 166, 167, 168, 169, 174, 175, 176, 177, 178, 185, 186, 187 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Fricas

**A grade** { 2, 8, 9, 16, 17, 18, 19, 20, 21, 22, 23, 24, 27, 28, 29, 30, 31, 32, 39, 40, 41, 48, 49, 50, 87, 91, 92, 93, 94, 95, 96, 99, 103, 104, 105, 106, 129, 130, 137, 138, 139, 147, 148, 149, 156, 157, 165, 166, 185, 186 }

**B grade** { 4, 7, 25, 33, 35, 38, 47, 74, 75, 76, 78, 79, 88, 89, 90, 100, 101, 102, 116, 123, 124, 131, 140, 150, 151, 158, 159, 160, 167, 168, 169, 174, 175, 187 }

**C grade** { }

**F normal fail** { 1, 3, 5, 6, 10, 11, 12, 13, 14, 15, 26, 34, 36, 37, 42, 43, 44, 45, 46, 54, 55, 56, 60, 61, 62, 66, 67, 68, 71, 77, 83, 84, 85, 97, 98, 107, 108, 109, 110, 111, 112, 113, 114, 115, 117, 118, 119, 120, 121, 122, 125, 126, 127, 128, 176, 177, 178 }



**F(-1) timedout fail** { 80, 81, 82 }

**F(-2) exception fail** { }

## Maxima

**A grade** { 4, 7, 16, 19, 20, 21, 22, 23, 24, 25, 27, 28, 29, 30, 31, 32, 35, 38, 47, 74, 75, 76, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 99, 100, 101, 102, 103, 104, 105, 106 }

**B grade** { }

**C grade** { }

**F normal fail** { 2, 3, 5, 6, 8, 9, 10, 11, 12, 13, 14, 15, 17, 18, 26, 33, 34, 36, 37, 39, 40, 41, 42, 43, 44, 45, 46, 48, 49, 50, 54, 55, 56, 60, 61, 62, 66, 67, 68, 71, 77, 78, 97, 98, 107, 108, 110, 112, 114, 115, 116, 122, 125, 131, 140, 151, 160, 165, 168, 169, 174, 175, 176, 177, 178, 185, 186, 187 }

**F(-1) timedout fail** { 1 }

**F(-2) exception fail** { 79, 80, 81, 82, 83, 84, 85, 109, 111, 113, 117, 118, 119, 120, 121, 123, 124, 126, 127, 128, 129, 130, 132, 133, 134, 135, 136, 137, 138, 139, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 152, 153, 154, 155, 156, 157, 158, 159, 161, 162, 163, 164, 166, 167, 170, 171, 172, 173 }

## Giac

**A grade** { }

**B grade** { }

**C grade** { }

**F normal fail** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 54, 55, 56, 60, 61, 62, 66, 67, 68, 71, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 137, 138, 139, 140, 147, 148, 149, 150, 151, 156, 157, 158, 159, 160, 165, 166, 167, 168, 169, 174, 175, 176, 177, 178, 187 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { 185, 186 }

## Mupad

**A grade** { }

**B grade** { 24, 25, 27, 28, 76, 90 }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 26, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 54, 55, 56, 60, 61, 62, 66, 67, 68, 71, 74, 75, 77, 78, 79, 80, 81, 82, 83, 84, 85, 87, 88, 89, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 137, 138, 139, 140, 147, 148, 149, 150, 151, 156, 157, 158, 159, 160, 165, 166, 167, 168, 169, 174, 175, 176, 177, 178, 185, 186, 187 }

**F(-2) exception fail** { }

## Sympy

**A grade** { 4, 20, 22, 24, 35, 94, 95, 96, 105, 106 }

**B grade** { }

**C grade** { }

**F normal fail** { 1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 21, 23, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 54, 55, 56, 60, 61, 62, 66, 67, 68, 71, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 87, 88, 89, 90, 91, 92, 93, 97, 98, 99, 100, 101, 102, 103, 104, 107, 108, 109, 110, 111, 112, 113, 115, 116, 117, 118, 119, 120, 129, 130, 131, 137, 138, 140, 147, 149, 150, 151, 156, 157, 158, 159, 160, 165, 166, 176, 177, 178, 186, 187 }

**F(-1) timedout fail** { 85, 114, 121, 122, 123, 124, 125, 126, 127, 128, 139, 148, 167, 168, 169, 170, 171, 172, 173, 174, 175, 180, 181, 185 }

**F(-2) exception fail** { }

## Reduce

**A grade** { }

**B grade** { }

**C grade** { }

**F normal fail** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24,  
25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49,  
50, 54, 55, 56, 60, 61, 62, 66, 67, 68, 71, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 87, 88,  
89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110,  
111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129,  
130, 131, 137, 138, 139, 140, 147, 148, 149, 150, 151, 156, 157, 158, 159, 160, 165, 166, 167,  
168, 169, 174, 175, 176, 177, 178, 185, 186, 187 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

## 2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as  $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$ . To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	164	168	182	280	0	0	0	0	12	0
N.S.	1	1.02	1.11	1.71	0.00	0.00	0.00	0.00	0.07	0.00
time (sec)	N/A	0.602	0.271	0.470	0.000	0.000	0.000	0.000	0.330	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	104	117	77	150	0	125	0	0	12	0
N.S.	1	1.12	0.74	1.44	0.00	1.20	0.00	0.00	0.12	0.00
time (sec)	N/A	0.504	0.072	0.332	0.000	0.091	0.000	0.000	0.296	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	116	138	230	0	0	0	0	12	0
N.S.	1	0.99	1.18	1.97	0.00	0.00	0.00	0.00	0.10	0.00
time (sec)	N/A	0.498	0.174	0.395	0.000	0.000	0.000	0.000	0.261	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	60	53	100	40	106	42	0	10	0
N.S.	1	1.13	1.00	1.89	0.75	2.00	0.79	0.00	0.19	0.00
time (sec)	N/A	0.371	0.041	0.283	0.032	0.099	0.358	0.000	0.241	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	64	90	183	0	0	0	0	8	0
N.S.	1	1.02	1.43	2.90	0.00	0.00	0.00	0.00	0.13	0.00
time (sec)	N/A	0.378	0.139	0.295	0.000	0.000	0.000	0.000	0.226	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	79	63	136	0	0	0	0	12	0
N.S.	1	1.23	0.98	2.12	0.00	0.00	0.00	0.00	0.19	0.00
time (sec)	N/A	0.469	0.026	0.234	0.000	0.000	0.000	0.000	0.214	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	69	42	61	35	97	0	0	12	0
N.S.	1	1.41	0.86	1.24	0.71	1.98	0.00	0.00	0.24	0.00
time (sec)	N/A	0.352	0.069	0.250	0.032	0.086	0.000	0.000	0.199	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	101	54	43	0	106	0	0	12	0
N.S.	1	1.12	0.60	0.48	0.00	1.18	0.00	0.00	0.13	0.00
time (sec)	N/A	0.310	0.032	0.286	0.000	0.084	0.000	0.000	0.188	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	121	73	112	0	116	0	0	12	0
N.S.	1	1.19	0.72	1.10	0.00	1.14	0.00	0.00	0.12	0.00
time (sec)	N/A	0.460	0.069	0.459	0.000	0.082	0.000	0.000	0.201	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	297	319	281	0	0	0	0	0	12	0
N.S.	1	1.07	0.95	0.00	0.00	0.00	0.00	0.00	0.04	0.00
time (sec)	N/A	1.069	0.412	0.000	0.000	0.000	0.000	0.000	0.344	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	184	199	188	236	0	0	0	0	12	0
N.S.	1	1.08	1.02	1.28	0.00	0.00	0.00	0.00	0.07	0.00
time (sec)	N/A	0.752	0.472	0.353	0.000	0.000	0.000	0.000	0.300	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	198	187	199	0	0	0	0	0	12	0
N.S.	1	0.94	1.01	0.00	0.00	0.00	0.00	0.00	0.06	0.00
time (sec)	N/A	0.701	0.327	0.000	0.000	0.000	0.000	0.000	0.265	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	114	101	149	0	0	0	0	10	0
N.S.	1	1.12	0.99	1.46	0.00	0.00	0.00	0.00	0.10	0.00
time (sec)	N/A	0.531	0.289	0.293	0.000	0.000	0.000	0.000	0.233	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	111	104	128	0	0	0	0	0	8	0
N.S.	1	0.94	1.15	0.00	0.00	0.00	0.00	0.00	0.07	0.00
time (sec)	N/A	0.505	0.091	0.000	0.000	0.000	0.000	0.000	0.208	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	106	84	181	0	0	0	0	12	0
N.S.	1	1.20	0.95	2.06	0.00	0.00	0.00	0.00	0.14	0.00
time (sec)	N/A	0.566	0.042	0.243	0.000	0.000	0.000	0.000	0.194	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	115	75	98	55	155	0	0	12	0
N.S.	1	1.39	0.90	1.18	0.66	1.87	0.00	0.00	0.14	0.00
time (sec)	N/A	0.435	0.057	0.243	0.030	0.093	0.000	0.000	0.193	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	137	158	147	58	0	174	0	0	12	0
N.S.	1	1.15	1.07	0.42	0.00	1.27	0.00	0.00	0.09	0.00
time (sec)	N/A	0.403	0.098	0.301	0.000	0.106	0.000	0.000	0.195	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	179	245	120	192	0	186	0	0	12	0
N.S.	1	1.37	0.67	1.07	0.00	1.04	0.00	0.00	0.07	0.00
time (sec)	N/A	0.655	0.078	0.463	0.000	0.088	0.000	0.000	0.194	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	165	154	143	134	135	183	0	0	19	0
N.S.	1	0.93	0.87	0.81	0.82	1.11	0.00	0.00	0.12	0.00
time (sec)	N/A	0.299	0.138	0.401	0.107	0.121	0.000	0.000	0.392	0.000



Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	134	97	77	78	100	94	0	19	0
N.S.	1	1.02	0.74	0.59	0.60	0.76	0.72	0.00	0.15	0.00
time (sec)	N/A	0.298	0.065	0.366	0.037	0.092	0.912	0.000	0.436	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	117	123	114	106	174	0	0	19	0
N.S.	1	0.89	0.94	0.87	0.81	1.33	0.00	0.00	0.15	0.00
time (sec)	N/A	0.263	0.094	0.314	0.118	0.105	0.000	0.000	0.342	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	97	77	68	57	90	68	0	19	0
N.S.	1	1.10	0.88	0.77	0.65	1.02	0.77	0.00	0.22	0.00
time (sec)	N/A	0.249	0.060	0.295	0.031	0.085	0.519	0.000	0.351	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	80	103	92	73	162	0	0	19	0
N.S.	1	0.82	1.06	0.95	0.75	1.67	0.00	0.00	0.20	0.00
time (sec)	N/A	0.243	0.067	0.300	0.104	0.103	0.000	0.000	0.278	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	45	57	59	36	73	46	0	17	50
N.S.	1	0.96	1.21	1.26	0.77	1.55	0.98	0.00	0.36	1.06
time (sec)	N/A	0.224	0.044	0.277	0.026	0.097	0.294	0.000	0.287	3.570

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	40	80	42	31	119	0	0	12	44
N.S.	1	0.70	1.40	0.74	0.54	2.09	0.00	0.00	0.21	0.77
time (sec)	N/A	0.179	0.092	0.070	0.027	0.096	0.000	0.000	0.227	3.411

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	78	47	98	0	0	0	0	17	0
N.S.	1	1.39	0.84	1.75	0.00	0.00	0.00	0.00	0.30	0.00
time (sec)	N/A	0.531	0.041	0.358	0.000	0.000	0.000	0.000	0.197	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	40	42	54	32	66	0	0	21	46
N.S.	1	0.95	1.00	1.29	0.76	1.57	0.00	0.00	0.50	1.10
time (sec)	N/A	0.219	0.050	0.303	0.029	0.100	0.000	0.000	0.206	3.592

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	96	117	108	105	77	0	0	25	61
N.S.	1	0.90	1.09	1.01	0.98	0.72	0.00	0.00	0.23	0.57
time (sec)	N/A	0.256	0.055	0.383	0.026	0.082	0.000	0.000	0.206	3.476

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	97	74	73	56	79	0	0	25	0
N.S.	1	1.20	0.91	0.90	0.69	0.98	0.00	0.00	0.31	0.00
time (sec)	N/A	0.249	0.050	0.361	0.023	0.084	0.000	0.000	0.199	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	141	130	137	131	147	90	0	0	25	0
N.S.	1	0.92	0.97	0.93	1.04	0.64	0.00	0.00	0.18	0.00
time (sec)	N/A	0.293	0.070	0.378	0.033	0.083	0.000	0.000	0.206	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	131	94	81	73	89	0	0	25	0
N.S.	1	1.14	0.82	0.70	0.63	0.77	0.00	0.00	0.22	0.00
time (sec)	N/A	0.285	0.065	0.398	0.028	0.082	0.000	0.000	0.218	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	175	164	157	151	185	100	0	0	25	0
N.S.	1	0.94	0.90	0.86	1.06	0.57	0.00	0.00	0.14	0.00
time (sec)	N/A	0.314	0.105	0.434	0.030	0.089	0.000	0.000	0.217	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	132	212	224	0	244	0	0	39	0
N.S.	1	1.06	1.71	1.81	0.00	1.97	0.00	0.00	0.31	0.00
time (sec)	N/A	0.581	0.216	0.759	0.000	0.110	0.000	0.000	0.676	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	132	241	329	0	0	0	0	39	0
N.S.	1	0.94	1.72	2.35	0.00	0.00	0.00	0.00	0.28	0.00
time (sec)	N/A	0.570	0.936	0.799	0.000	0.000	0.000	0.000	0.565	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	72	112	165	84	205	99	0	35	0
N.S.	1	1.11	1.72	2.54	1.29	3.15	1.52	0.00	0.54	0.00
time (sec)	N/A	0.434	0.259	0.644	0.033	0.100	0.380	0.000	0.482	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	75	126	232	0	0	0	0	28	0
N.S.	1	0.96	1.62	2.97	0.00	0.00	0.00	0.00	0.36	0.00
time (sec)	N/A	0.411	0.215	0.490	0.000	0.000	0.000	0.000	0.318	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	99	116	241	0	0	0	0	37	0
N.S.	1	1.19	1.40	2.90	0.00	0.00	0.00	0.00	0.45	0.00
time (sec)	N/A	0.540	0.113	0.500	0.000	0.000	0.000	0.000	0.231	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	79	87	121	78	143	0	0	42	0
N.S.	1	1.30	1.43	1.98	1.28	2.34	0.00	0.00	0.69	0.00
time (sec)	N/A	0.398	0.158	0.488	0.034	0.089	0.000	0.000	0.237	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	120	183	157	0	165	0	0	48	0
N.S.	1	1.13	1.73	1.48	0.00	1.56	0.00	0.00	0.45	0.00
time (sec)	N/A	0.370	0.112	0.491	0.000	0.091	0.000	0.000	0.246	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	122	136	134	191	0	181	0	0	48	0
N.S.	1	1.11	1.10	1.57	0.00	1.48	0.00	0.00	0.39	0.00
time (sec)	N/A	0.513	0.168	0.733	0.000	0.097	0.000	0.000	0.237	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	139	158	268	263	0	204	0	0	48	0
N.S.	1	1.14	1.93	1.89	0.00	1.47	0.00	0.00	0.35	0.00
time (sec)	N/A	0.462	0.165	0.661	0.000	0.094	0.000	0.000	0.229	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	223	233	337	464	0	0	0	0	59	0
N.S.	1	1.04	1.51	2.08	0.00	0.00	0.00	0.00	0.26	0.00
time (sec)	N/A	0.925	1.353	0.838	0.000	0.000	0.000	0.000	1.005	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	242	224	440	0	0	0	0	0	59	0
N.S.	1	0.93	1.82	0.00	0.00	0.00	0.00	0.00	0.24	0.00
time (sec)	N/A	0.874	0.755	0.000	0.000	0.000	0.000	0.000	0.803	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	136	219	318	0	0	0	0	53	0
N.S.	1	1.08	1.74	2.52	0.00	0.00	0.00	0.00	0.42	0.00
time (sec)	N/A	0.626	0.682	0.688	0.000	0.000	0.000	0.000	0.610	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	140	127	282	0	0	0	0	0	44	0
N.S.	1	0.91	2.01	0.00	0.00	0.00	0.00	0.00	0.31	0.00
time (sec)	N/A	0.589	0.353	0.000	0.000	0.000	0.000	0.000	0.402	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	114	130	182	429	0	0	0	0	57	0
N.S.	1	1.14	1.60	3.76	0.00	0.00	0.00	0.00	0.50	0.00
time (sec)	N/A	0.687	0.157	0.492	0.000	0.000	0.000	0.000	0.321	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	130	165	225	144	228	0	0	63	0
N.S.	1	1.27	1.62	2.21	1.41	2.24	0.00	0.00	0.62	0.00
time (sec)	N/A	0.490	0.219	0.481	0.038	0.085	0.000	0.000	0.298	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	163	182	245	219	0	271	0	0	71	0
N.S.	1	1.12	1.50	1.34	0.00	1.66	0.00	0.00	0.44	0.00
time (sec)	N/A	0.466	0.293	0.484	0.000	0.088	0.000	0.000	0.311	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	213	275	256	387	0	305	0	0	71	0
N.S.	1	1.29	1.20	1.82	0.00	1.43	0.00	0.00	0.33	0.00
time (sec)	N/A	0.781	0.251	0.746	0.000	0.109	0.000	0.000	0.287	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	242	314	332	485	0	351	0	0	71	0
N.S.	1	1.30	1.37	2.00	0.00	1.45	0.00	0.00	0.29	0.00
time (sec)	N/A	0.826	0.441	0.724	0.000	0.109	0.000	0.000	0.305	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	14	14	10	14	14	18
N.S.	1	1.00	1.17	1.00	1.17	1.17	0.83	1.17	1.17	1.50
time (sec)	N/A	0.191	2.623	0.371	0.106	0.076	0.343	0.104	0.216	3.506



Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	10	12	12	16
N.S.	1	1.00	1.20	1.00	1.20	1.20	1.00	1.20	1.20	1.60
time (sec)	N/A	0.173	0.027	0.229	0.104	0.090	0.326	0.107	0.209	3.464

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	16	14	16	15	12	16	15	20
N.S.	1	1.00	1.14	1.00	1.14	1.07	0.86	1.14	1.07	1.43
time (sec)	N/A	0.203	0.218	0.201	0.116	0.091	1.176	0.105	0.203	3.619

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	54	43	54	0	0	0	0	19	0
N.S.	1	1.17	0.93	1.17	0.00	0.00	0.00	0.00	0.41	0.00
time (sec)	N/A	0.461	0.059	0.384	0.000	0.000	0.000	0.000	0.208	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	66	56	60	0	0	0	0	19	0
N.S.	1	1.05	0.89	0.95	0.00	0.00	0.00	0.00	0.30	0.00
time (sec)	N/A	0.574	0.058	0.569	0.000	0.000	0.000	0.000	0.214	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	110	91	110	0	0	0	0	19	0
N.S.	1	0.94	0.78	0.94	0.00	0.00	0.00	0.00	0.16	0.00
time (sec)	N/A	0.503	0.117	0.710	0.000	0.000	0.000	0.000	0.231	0.000

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	546	28	12	14	28	18
N.S.	1	1.00	1.17	1.00	45.50	2.33	1.00	1.17	2.33	1.50
time (sec)	N/A	0.191	12.204	0.272	0.398	0.080	0.698	0.102	0.224	3.531

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	535	26	12	12	26	16
N.S.	1	1.00	1.20	1.00	53.50	2.60	1.20	1.20	2.60	1.60
time (sec)	N/A	0.180	59.169	0.198	0.417	0.100	0.779	0.108	0.210	3.449

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	16	14	544	30	14	16	30	20
N.S.	1	1.00	1.14	1.00	38.86	2.14	1.00	1.14	2.14	1.43
time (sec)	N/A	0.200	4.349	0.158	0.313	0.087	1.473	0.107	0.222	3.481

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	103	82	164	0	0	0	0	36	0
N.S.	1	1.20	0.95	1.91	0.00	0.00	0.00	0.00	0.42	0.00
time (sec)	N/A	0.581	0.286	0.404	0.000	0.000	0.000	0.000	0.228	0.000

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	93	92	186	0	0	0	0	36	0
N.S.	1	1.09	1.08	2.19	0.00	0.00	0.00	0.00	0.42	0.00
time (sec)	N/A	0.675	0.375	0.604	0.000	0.000	0.000	0.000	0.275	0.000

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	190	180	250	420	0	0	0	0	36	0
N.S.	1	0.95	1.32	2.21	0.00	0.00	0.00	0.00	0.19	0.00
time (sec)	N/A	0.575	0.574	0.777	0.000	0.000	0.000	0.000	0.267	0.000

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	2818	42	12	14	42	18
N.S.	1	1.00	1.17	1.00	234.83	3.50	1.00	1.17	3.50	1.50
time (sec)	N/A	0.215	3.812	0.273	1.993	0.091	1.271	0.103	0.247	3.656

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	2771	40	12	12	40	16
N.S.	1	1.00	1.20	1.00	277.10	4.00	1.20	1.20	4.00	1.60
time (sec)	N/A	0.183	88.386	0.200	1.942	0.092	1.312	0.107	0.215	3.642

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	16	14	2638	45	14	16	45	20
N.S.	1	1.00	1.14	1.00	188.43	3.21	1.00	1.14	3.21	1.43
time (sec)	N/A	0.203	1.637	0.159	1.777	0.103	2.467	0.105	0.226	3.684

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	114	141	103	244	0	0	0	0	53	0
N.S.	1	1.24	0.90	2.14	0.00	0.00	0.00	0.00	0.46	0.00
time (sec)	N/A	0.687	0.190	0.427	0.000	0.000	0.000	0.000	0.242	0.000

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	112	130	122	277	0	0	0	0	53	0
N.S.	1	1.16	1.09	2.47	0.00	0.00	0.00	0.00	0.47	0.00
time (sec)	N/A	0.776	0.257	0.608	0.000	0.000	0.000	0.000	0.281	0.000

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	240	227	204	628	0	0	0	0	53	0
N.S.	1	0.95	0.85	2.62	0.00	0.00	0.00	0.00	0.22	0.00
time (sec)	N/A	0.653	0.366	0.868	0.000	0.000	0.000	0.000	0.295	0.000

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	1450	44	15	18	121	22
N.S.	1	1.00	1.12	1.00	90.62	2.75	0.94	1.12	7.56	1.38
time (sec)	N/A	0.202	3.257	0.581	10.373	0.120	6.623	0.122	0.567	3.881

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	704	30	15	18	80	22
N.S.	1	1.00	1.12	1.00	44.00	1.88	0.94	1.12	5.00	1.38
time (sec)	N/A	0.210	1.798	0.395	4.308	0.112	2.607	0.111	0.451	3.949

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	104	87	97	0	0	0	0	0	41	0
N.S.	1	0.84	0.93	0.00	0.00	0.00	0.00	0.00	0.39	0.00
time (sec)	N/A	0.259	0.103	0.000	0.000	0.000	0.000	0.000	0.295	0.000

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	18	18	14	18	20	22
N.S.	1	1.00	1.12	1.00	1.12	1.12	0.88	1.12	1.25	1.38
time (sec)	N/A	0.211	0.287	0.734	0.129	0.090	0.512	0.107	0.222	3.576

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	616	32	15	18	34	22
N.S.	1	1.00	1.12	1.00	38.50	2.00	0.94	1.12	2.12	1.38
time (sec)	N/A	0.218	0.595	0.610	0.764	0.100	1.479	0.109	0.219	3.838

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	264	196	190	264	221	358	0	0	93	0
N.S.	1	0.74	0.72	1.00	0.84	1.36	0.00	0.00	0.35	0.00
time (sec)	N/A	0.751	0.239	0.323	0.112	0.192	0.000	0.000	0.663	0.000

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	201	149	147	199	152	280	0	0	62	0
N.S.	1	0.74	0.73	0.99	0.76	1.39	0.00	0.00	0.31	0.00
time (sec)	N/A	0.482	0.138	0.318	0.108	0.162	0.000	0.000	0.501	0.000

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	142	105	142	107	70	177	0	0	32	99
N.S.	1	0.74	1.00	0.75	0.49	1.25	0.00	0.00	0.23	0.70
time (sec)	N/A	0.314	0.218	0.302	0.034	0.122	0.000	0.000	0.324	3.830

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	211	229	393	511	0	0	0	0	30	0
N.S.	1	1.09	1.86	2.42	0.00	0.00	0.00	0.00	0.14	0.00
time (sec)	N/A	1.373	0.392	0.811	0.000	0.000	0.000	0.000	0.253	0.000

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	147	132	222	207	0	578	0	0	75	0
N.S.	1	0.90	1.51	1.41	0.00	3.93	0.00	0.00	0.51	0.00
time (sec)	N/A	0.380	0.149	1.780	0.000	0.145	0.000	0.000	0.294	0.000

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	<b>F(-2)</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	234	244	342	593	0	1212	0	0	159	0
N.S.	1	1.04	1.46	2.53	0.00	5.18	0.00	0.00	0.68	0.00
time (sec)	N/A	0.472	0.399	1.971	0.000	0.224	0.000	0.000	0.319	0.000

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F(-2)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	343	306	2653	818	0	0	0	0	83	0
N.S.	1	0.89	7.73	2.38	0.00	0.00	0.00	0.00	0.24	0.00
time (sec)	N/A	0.825	19.991	18.740	0.000	0.000	0.000	0.000	1.490	0.000

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F(-2)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	279	256	2938	413	0	0	0	0	44	0
N.S.	1	0.92	10.53	1.48	0.00	0.00	0.00	0.00	0.16	0.00
time (sec)	N/A	0.624	24.883	17.445	0.000	0.000	0.000	0.000	0.821	0.000

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F(-2)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	187	170	1707	286	0	0	0	0	32	0
N.S.	1	0.91	9.13	1.53	0.00	0.00	0.00	0.00	0.17	0.00
time (sec)	N/A	0.478	15.075	17.128	0.000	0.000	0.000	0.000	0.568	0.000

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F(-2)	F	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	105	123	1675	251	0	0	0	0	52	0
N.S.	1	1.17	15.95	2.39	0.00	0.00	0.00	0.00	0.50	0.00
time (sec)	N/A	0.358	14.374	16.984	0.000	0.000	0.000	0.000	0.629	0.000



Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F(-2)	F	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	278	254	4527	890	0	0	0	0	136	0
N.S.	1	0.91	16.28	3.20	0.00	0.00	0.00	0.00	0.49	0.00
time (sec)	N/A	0.581	24.325	20.956	0.000	0.000	0.000	0.000	0.777	0.000

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F(-2)	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	472	433	8675	1612	0	0	0	0	260	0
N.S.	1	0.92	18.38	3.42	0.00	0.00	0.00	0.00	0.55	0.00
time (sec)	N/A	0.900	25.399	25.119	0.000	0.000	0.000	0.000	0.877	0.000

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	222	18	15	18	66	22
N.S.	1	1.00	1.12	1.00	13.88	1.12	0.94	1.12	4.12	1.38
time (sec)	N/A	0.260	14.703	0.394	0.662	0.110	17.638	0.110	0.310	3.887

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	229	164	162	213	244	259	0	0	41	0
N.S.	1	0.72	0.71	0.93	1.07	1.13	0.00	0.00	0.18	0.00
time (sec)	N/A	0.367	0.247	0.507	0.107	0.177	0.000	0.000	0.694	0.000

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	174	132	144	171	182	238	0	0	41	0
N.S.	1	0.76	0.83	0.98	1.05	1.37	0.00	0.00	0.24	0.00
time (sec)	N/A	0.324	0.172	0.484	0.106	0.144	0.000	0.000	0.551	0.000

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	112	96	189	120	107	209	0	0	34	0
N.S.	1	0.86	1.69	1.07	0.96	1.87	0.00	0.00	0.30	0.00
time (sec)	N/A	0.274	0.247	0.318	0.111	0.148	0.000	0.000	0.369	0.000

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	96	80	127	112	66	182	0	0	39	98
N.S.	1	0.83	1.32	1.17	0.69	1.90	0.00	0.00	0.41	1.02
time (sec)	N/A	0.276	0.168	0.325	0.026	0.117	0.000	0.000	0.264	4.336

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	110	76	110	91	106	0	0	51	0
N.S.	1	0.87	0.60	0.87	0.72	0.84	0.00	0.00	0.40	0.00
time (sec)	N/A	0.300	0.094	0.332	0.030	0.094	0.000	0.000	0.257	0.000

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	183	141	101	129	132	128	0	0	51	0
N.S.	1	0.77	0.55	0.70	0.72	0.70	0.00	0.00	0.28	0.00
time (sec)	N/A	0.341	0.127	0.379	0.034	0.106	0.000	0.000	0.269	0.000

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	238	170	117	147	165	149	0	0	51	0
N.S.	1	0.71	0.49	0.62	0.69	0.63	0.00	0.00	0.21	0.00
time (sec)	N/A	0.350	0.152	0.352	0.034	0.110	0.000	0.000	0.294	0.000

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	232	172	126	139	177	168	228	0	41	0
N.S.	1	0.74	0.54	0.60	0.76	0.72	0.98	0.00	0.18	0.00
time (sec)	N/A	0.397	0.160	0.498	0.032	0.114	1.619	0.000	0.800	0.000

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	180	141	106	121	138	147	177	0	41	0
N.S.	1	0.78	0.59	0.67	0.77	0.82	0.98	0.00	0.23	0.00
time (sec)	N/A	0.378	0.139	0.529	0.033	0.102	0.867	0.000	0.659	0.000

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	164	126	85	194	96	125	126	0	39	0
N.S.	1	0.77	0.52	1.18	0.59	0.76	0.77	0.00	0.24	0.00
time (sec)	N/A	0.479	0.105	0.480	0.031	0.102	0.477	0.000	0.481	0.000

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	296	323	108	161	0	0	0	0	37	0
N.S.	1	1.09	0.36	0.54	0.00	0.00	0.00	0.00	0.12	0.00
time (sec)	N/A	1.283	0.154	0.807	0.000	0.000	0.000	0.000	0.268	0.000

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	309	347	170	168	0	0	0	0	53	0
N.S.	1	1.12	0.55	0.54	0.00	0.00	0.00	0.00	0.17	0.00
time (sec)	N/A	1.163	0.168	1.089	0.000	0.000	0.000	0.000	0.251	0.000

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	275	212	207	286	328	341	0	0	72	0
N.S.	1	0.77	0.75	1.04	1.19	1.24	0.00	0.00	0.26	0.00
time (sec)	N/A	0.456	0.292	0.533	0.114	0.247	0.000	0.000	0.783	0.000

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	204	175	174	208	224	305	0	0	65	0
N.S.	1	0.86	0.85	1.02	1.10	1.50	0.00	0.00	0.32	0.00
time (sec)	N/A	0.376	0.182	0.355	0.110	0.206	0.000	0.000	0.563	0.000

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	177	142	158	184	152	287	0	0	74	0
N.S.	1	0.80	0.89	1.04	0.86	1.62	0.00	0.00	0.42	0.00
time (sec)	N/A	0.392	0.156	0.375	0.115	0.150	0.000	0.000	0.335	0.000

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	176	143	149	201	134	267	0	0	81	0
N.S.	1	0.81	0.85	1.14	0.76	1.52	0.00	0.00	0.46	0.00
time (sec)	N/A	0.391	0.175	0.352	0.036	0.135	0.000	0.000	0.334	0.000

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	213	181	134	177	175	167	0	0	85	0
N.S.	1	0.85	0.63	0.83	0.82	0.78	0.00	0.00	0.40	0.00
time (sec)	N/A	0.411	0.181	0.393	0.027	0.109	0.000	0.000	0.319	0.000

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	281	212	160	209	232	199	0	0	85	0
N.S.	1	0.75	0.57	0.74	0.83	0.71	0.00	0.00	0.30	0.00
time (sec)	N/A	0.446	0.222	0.385	0.034	0.098	0.000	0.000	0.318	0.000

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	278	218	168	198	245	227	332	0	72	0
N.S.	1	0.78	0.60	0.71	0.88	0.82	1.19	0.00	0.26	0.00
time (sec)	N/A	0.506	0.177	0.527	0.035	0.124	1.696	0.000	0.956	0.000

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	230	171	139	283	185	192	252	0	70	0
N.S.	1	0.74	0.60	1.23	0.80	0.83	1.10	0.00	0.30	0.00
time (sec)	N/A	0.518	0.167	0.532	0.033	0.119	0.914	0.000	0.755	0.000

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	370	401	176	276	0	0	0	0	67	0
N.S.	1	1.08	0.48	0.75	0.00	0.00	0.00	0.00	0.18	0.00
time (sec)	N/A	1.438	0.269	0.986	0.000	0.000	0.000	0.000	0.355	0.000

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	373	411	223	250	0	0	0	0	84	0
N.S.	1	1.10	0.60	0.67	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	1.365	0.667	1.313	0.000	0.000	0.000	0.000	0.336	0.000

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	519	571	921	411	0	0	0	0	52	0
N.S.	1	1.10	1.77	0.79	0.00	0.00	0.00	0.00	0.10	0.00
time (sec)	N/A	1.578	1.837	36.369	0.000	0.000	0.000	0.000	0.330	0.000

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	441	523	860	506	0	0	0	0	37	0
N.S.	1	1.19	1.95	1.15	0.00	0.00	0.00	0.00	0.08	0.00
time (sec)	N/A	1.523	0.407	1.031	0.000	0.000	0.000	0.000	0.302	0.000

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	469	517	849	302	0	0	0	0	46	0
N.S.	1	1.10	1.81	0.64	0.00	0.00	0.00	0.00	0.10	0.00
time (sec)	N/A	1.215	0.406	14.655	0.000	0.000	0.000	0.000	0.298	0.000

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	417	469	841	2081	0	0	0	0	44	0
N.S.	1	1.12	2.02	4.99	0.00	0.00	0.00	0.00	0.11	0.00
time (sec)	N/A	1.297	0.347	1.419	0.000	0.000	0.000	0.000	0.310	0.000

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	523	575	933	372	0	0	0	0	58	0
N.S.	1	1.10	1.78	0.71	0.00	0.00	0.00	0.00	0.11	0.00
time (sec)	N/A	1.425	2.032	30.194	0.000	0.000	0.000	0.000	0.387	0.000

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	611	703	1278	786	0	0	0	0	137	0
N.S.	1	1.15	2.09	1.29	0.00	0.00	0.00	0.00	0.22	0.00
time (sec)	N/A	1.762	3.231	5.637	0.000	0.000	0.000	0.000	0.661	0.000

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	562	648	1208	644	0	0	0	0	123	0
N.S.	1	1.15	2.15	1.15	0.00	0.00	0.00	0.00	0.22	0.00
time (sec)	N/A	1.759	0.925	1.318	0.000	0.000	0.000	0.000	0.601	0.000



Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	147	127	345	466	0	602	0	0	90	0
N.S.	1	0.86	2.35	3.17	0.00	4.10	0.00	0.00	0.61	0.00
time (sec)	N/A	0.481	0.730	4.224	0.000	0.123	0.000	0.000	0.579	0.000

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	542	598	1189	2226	0	0	0	0	138	0
N.S.	1	1.10	2.19	4.11	0.00	0.00	0.00	0.00	0.25	0.00
time (sec)	N/A	1.591	1.239	5.510	0.000	0.000	0.000	0.000	0.613	0.000

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	840	900	1270	1006	0	0	0	0	149	0
N.S.	1	1.07	1.51	1.20	0.00	0.00	0.00	0.00	0.18	0.00
time (sec)	N/A	2.995	0.904	84.631	0.000	0.000	0.000	0.000	0.710	0.000

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	786	842	1226	910	0	0	0	0	144	0
N.S.	1	1.07	1.56	1.16	0.00	0.00	0.00	0.00	0.18	0.00
time (sec)	N/A	1.768	1.102	30.899	0.000	0.000	0.000	0.000	0.650	0.000

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F(-2)	F	F	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	786	842	1216	898	0	0	0	0	137	0
N.S.	1	1.07	1.55	1.14	0.00	0.00	0.00	0.00	0.17	0.00
time (sec)	N/A	2.623	1.022	39.140	0.000	0.000	0.000	0.000	0.612	0.000

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F(-2)	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	844	904	1305	1007	0	0	0	0	155	0
N.S.	1	1.07	1.55	1.19	0.00	0.00	0.00	0.00	0.18	0.00
time (sec)	N/A	2.750	0.876	78.124	0.000	0.000	0.000	0.000	0.721	0.000

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	749	850	2000	1549	0	0	0	0	238	0
N.S.	1	1.13	2.67	2.07	0.00	0.00	0.00	0.00	0.32	0.00
time (sec)	N/A	2.006	7.267	5.553	0.000	0.000	0.000	0.000	0.893	0.000

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F(-2)	B	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	173	153	486	1362	0	1346	0	0	180	0
N.S.	1	0.88	2.81	7.87	0.00	7.78	0.00	0.00	1.04	0.00
time (sec)	N/A	0.376	1.018	4.156	0.000	0.254	0.000	0.000	0.857	0.000

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	<b>F(-2)</b>	B	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	217	198	486	1318	0	1232	0	0	172	0
N.S.	1	0.91	2.24	6.07	0.00	5.68	0.00	0.00	0.79	0.00
time (sec)	N/A	0.540	0.748	4.005	0.000	0.206	0.000	0.000	0.857	0.000

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	730	801	2054	3727	0	0	0	0	265	0
N.S.	1	1.10	2.81	5.11	0.00	0.00	0.00	0.00	0.36	0.00
time (sec)	N/A	1.860	6.062	2.089	0.000	0.000	0.000	0.000	0.857	0.000

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	1272	1336	1823	1960	0	0	0	0	272	0
N.S.	1	1.05	1.43	1.54	0.00	0.00	0.00	0.00	0.21	0.00
time (sec)	N/A	2.525	5.556	75.241	0.000	0.000	0.000	0.000	0.924	0.000

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	1276	1340	2030	1398	0	0	0	0	271	0
N.S.	1	1.05	1.59	1.10	0.00	0.00	0.00	0.00	0.21	0.00
time (sec)	N/A	3.612	6.071	72.883	0.000	0.000	0.000	0.000	0.872	0.000

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	1272	1336	1813	1950	0	0	0	0	263	0
N.S.	1	1.05	1.43	1.53	0.00	0.00	0.00	0.00	0.21	0.00
time (sec)	N/A	4.261	6.046	94.872	0.000	0.000	0.000	0.000	0.878	0.000

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F(-2)</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	447	380	340	0	0	1995	0	0	23	0
N.S.	1	0.85	0.76	0.00	0.00	4.46	0.00	0.00	0.05	0.00
time (sec)	N/A	1.539	35.618	0.000	0.000	1.422	0.000	0.000	106.761	0.000

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F(-2)</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	329	279	365	0	0	1669	0	0	23	0
N.S.	1	0.85	1.11	0.00	0.00	5.07	0.00	0.00	0.07	0.00
time (sec)	N/A	0.582	22.144	0.000	0.000	0.663	0.000	0.000	95.253	0.000

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	221	190	307	0	0	1382	0	0	21	0
N.S.	1	0.86	1.39	0.00	0.00	6.25	0.00	0.00	0.10	0.00
time (sec)	N/A	0.580	21.745	0.000	0.000	0.329	0.000	0.000	84.260	0.000

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	23	20	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	1.00	0.87	1.00	1.00	1.17
time (sec)	N/A	0.273	5.649	0.193	0.000	0.103	4.767	0.116	200.027	4.136

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	23	22	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	1.00	0.96	1.00	1.00	1.17
time (sec)	N/A	0.278	5.378	0.382	0.000	0.109	4.986	0.125	200.030	4.352

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	27	22	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	1.17	0.96	1.00	1.00	1.17
time (sec)	N/A	0.287	12.873	0.170	0.000	0.105	9.742	0.128	200.027	4.462

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	18	0	20	19	20	20	24
N.S.	1	1.00	1.10	0.90	0.00	1.00	0.95	1.00	1.00	1.20
time (sec)	N/A	0.224	5.201	0.211	0.000	0.106	2.553	0.126	200.038	4.282

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	23	22	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	1.00	0.96	1.00	1.00	1.17
time (sec)	N/A	0.266	1.612	0.222	0.000	0.099	3.092	0.117	200.025	4.394

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F(-2)</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	312	256	576	0	0	242	0	0	23	0
N.S.	1	0.82	1.85	0.00	0.00	0.78	0.00	0.00	0.07	0.00
time (sec)	N/A	0.603	23.721	0.000	0.000	0.107	0.000	0.000	200.027	0.000

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F(-2)</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	446	369	641	0	0	340	0	0	23	0
N.S.	1	0.83	1.44	0.00	0.00	0.76	0.00	0.00	0.05	0.00
time (sec)	N/A	0.802	25.403	0.000	0.000	0.118	0.000	0.000	200.026	0.000

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F(-2)</b>	A	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	418	354	313	0	0	1989	0	0	23	0
N.S.	1	0.85	0.75	0.00	0.00	4.76	0.00	0.00	0.06	0.00
time (sec)	N/A	0.672	35.485	0.000	0.000	1.494	0.000	0.000	109.479	0.000

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	297	254	342	0	0	1667	0	0	21	0
N.S.	1	0.86	1.15	0.00	0.00	5.61	0.00	0.00	0.07	0.00
time (sec)	N/A	0.654	22.116	0.000	0.000	0.601	0.000	0.000	94.842	0.000

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	40	20	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	1.74	0.87	1.00	1.00	1.17
time (sec)	N/A	0.296	6.957	0.183	0.000	0.096	30.754	0.150	200.029	4.117

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	40	22	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	1.74	0.96	1.00	1.00	1.17
time (sec)	N/A	0.302	5.836	0.477	0.000	0.114	22.926	0.128	200.026	4.390

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	43	22	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	1.87	0.96	1.00	1.00	1.17
time (sec)	N/A	0.288	13.368	0.185	0.000	0.103	79.061	0.136	200.032	4.331

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	18	0	37	19	20	20	24
N.S.	1	1.00	1.10	0.90	0.00	1.85	0.95	1.00	1.00	1.20
time (sec)	N/A	0.226	5.997	0.168	0.000	0.107	23.489	0.146	200.202	4.146

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	40	22	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	1.74	0.96	1.00	1.00	1.17
time (sec)	N/A	0.276	10.872	0.234	0.000	0.101	20.383	0.134	200.028	4.496

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	40	22	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	1.74	0.96	1.00	1.00	1.17
time (sec)	N/A	0.280	12.657	0.414	0.000	0.123	23.082	0.130	200.025	4.245

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F(-2)</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	409	338	620	0	0	338	0	0	23	0
N.S.	1	0.83	1.52	0.00	0.00	0.83	0.00	0.00	0.06	0.00
time (sec)	N/A	0.707	25.260	0.000	0.000	0.148	0.000	0.000	200.026	0.000



Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F(-2)</b>	A	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	556	462	728	0	0	447	0	0	23	0
N.S.	1	0.83	1.31	0.00	0.00	0.80	0.00	0.00	0.04	0.00
time (sec)	N/A	0.919	26.739	0.000	0.000	0.131	0.000	0.000	200.028	0.000

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F(-2)</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	356	306	366	0	0	1679	0	0	23	0
N.S.	1	0.86	1.03	0.00	0.00	4.72	0.00	0.00	0.06	0.00
time (sec)	N/A	1.361	22.199	0.000	0.000	0.661	0.000	0.000	136.049	0.000

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F(-2)</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	251	216	406	0	0	1389	0	0	23	0
N.S.	1	0.86	1.62	0.00	0.00	5.53	0.00	0.00	0.09	0.00
time (sec)	N/A	0.505	21.820	0.000	0.000	0.332	0.000	0.000	131.240	0.000

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	153	137	239	0	0	1102	0	0	21	0
N.S.	1	0.90	1.56	0.00	0.00	7.20	0.00	0.00	0.14	0.00
time (sec)	N/A	0.497	18.934	0.000	0.000	0.229	0.000	0.000	129.928	0.000

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	31	20	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	1.35	0.87	1.00	1.00	1.17
time (sec)	N/A	0.277	2.037	0.186	0.000	0.106	2.875	0.123	200.027	4.267

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	33	22	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	1.43	0.96	1.00	1.00	1.17
time (sec)	N/A	0.280	5.158	0.324	0.000	0.095	9.168	0.121	200.027	4.431

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	27	22	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	1.17	0.96	1.00	1.00	1.17
time (sec)	N/A	0.282	10.271	0.163	0.000	0.103	4.436	0.128	200.024	4.415

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	18	0	20	19	20	20	24
N.S.	1	1.00	1.10	0.90	0.00	1.00	0.95	1.00	1.00	1.20
time (sec)	N/A	0.217	0.700	0.157	0.000	0.091	1.353	0.122	67.928	4.283

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F(-2)</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	218	186	501	0	0	154	0	0	23	0
N.S.	1	0.85	2.30	0.00	0.00	0.71	0.00	0.00	0.11	0.00
time (sec)	N/A	0.453	23.807	0.000	0.000	0.125	0.000	0.000	200.024	0.000

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F(-2)</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	346	288	612	0	0	244	0	0	23	0
N.S.	1	0.83	1.77	0.00	0.00	0.71	0.00	0.00	0.07	0.00
time (sec)	N/A	0.638	24.174	0.000	0.000	0.132	0.000	0.000	200.025	0.000

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F(-2)</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	278	243	436	0	0	1771	0	0	23	0
N.S.	1	0.87	1.57	0.00	0.00	6.37	0.00	0.00	0.08	0.00
time (sec)	N/A	1.351	22.077	0.000	0.000	0.333	0.000	0.000	140.492	0.000

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F(-2)</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	177	161	249	0	0	1311	0	0	23	0
N.S.	1	0.91	1.41	0.00	0.00	7.41	0.00	0.00	0.13	0.00
time (sec)	N/A	0.460	21.965	0.000	0.000	0.251	0.000	0.000	136.047	0.000

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	135	0	0	379	0	0	21	0
N.S.	1	1.00	1.55	0.00	0.00	4.36	0.00	0.00	0.24	0.00
time (sec)	N/A	0.461	19.877	0.000	0.000	0.152	0.000	0.000	131.797	0.000

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	42	20	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	1.83	0.87	1.00	1.00	1.17
time (sec)	N/A	0.301	7.969	0.201	0.000	0.116	23.168	0.124	200.024	4.036

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	44	22	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	1.91	0.96	1.00	1.00	1.17
time (sec)	N/A	0.307	10.505	0.375	0.000	0.122	97.817	0.123	200.034	4.269

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	47	22	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	2.04	0.96	1.00	1.00	1.17
time (sec)	N/A	0.287	14.139	0.203	0.000	0.108	47.522	0.133	200.045	4.179

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	47	22	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	2.04	0.96	1.00	1.00	1.17
time (sec)	N/A	0.289	5.708	0.177	0.000	0.123	11.069	0.134	200.047	3.817

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	334	0	0	107	0	0	20	0
N.S.	1	1.00	3.63	0.00	0.00	1.16	0.00	0.00	0.22	0.00
time (sec)	N/A	0.283	48.588	0.000	0.000	0.119	0.000	0.000	200.035	0.000

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F(-2)</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	249	214	501	0	0	238	0	0	23	0
N.S.	1	0.86	2.01	0.00	0.00	0.96	0.00	0.00	0.09	0.00
time (sec)	N/A	0.552	23.958	0.000	0.000	0.129	0.000	0.000	200.036	0.000

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F(-2)</b>	B	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	272	229	348	0	0	2415	0	0	23	0
N.S.	1	0.84	1.28	0.00	0.00	8.88	0.00	0.00	0.08	0.00
time (sec)	N/A	1.449	22.104	0.000	0.000	0.353	0.000	0.000	115.789	0.000

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	B	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	179	157	218	0	0	786	0	0	23	0
N.S.	1	0.88	1.22	0.00	0.00	4.39	0.00	0.00	0.13	0.00
time (sec)	N/A	0.475	0.429	0.000	0.000	0.221	0.000	0.000	102.752	0.000

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	B	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	154	135	204	0	0	692	0	0	21	0
N.S.	1	0.88	1.32	0.00	0.00	4.49	0.00	0.00	0.14	0.00
time (sec)	N/A	0.548	0.373	0.000	0.000	0.195	0.000	0.000	99.027	0.000

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	53	0	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	2.30	0.00	1.00	1.00	1.17
time (sec)	N/A	0.312	13.684	0.181	0.000	0.116	0.000	0.134	200.032	4.076

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	55	0	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	2.39	0.00	1.00	1.00	1.17
time (sec)	N/A	0.318	15.700	0.415	0.000	0.109	0.000	0.129	200.023	4.440

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	<b>F(-1)</b>	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	58	0	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	2.52	0.00	1.00	1.00	1.17
time (sec)	N/A	0.302	15.047	0.188	0.000	0.122	0.000	0.136	200.029	4.204

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	<b>F(-1)</b>	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	0	58	0	23	23	27
N.S.	1	1.00	1.09	0.91	0.00	2.52	0.00	1.00	1.00	1.17
time (sec)	N/A	0.293	14.799	0.188	0.000	0.110	0.000	0.133	200.026	4.083

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	B	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	243	205	488	0	0	335	0	0	23	0
N.S.	1	0.84	2.01	0.00	0.00	1.38	0.00	0.00	0.09	0.00
time (sec)	N/A	0.528	1.977	0.000	0.000	0.148	0.000	0.000	200.028	0.000

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	B	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	266	224	517	0	0	397	0	0	20	0
N.S.	1	0.84	1.94	0.00	0.00	1.49	0.00	0.00	0.08	0.00
time (sec)	N/A	0.468	24.743	0.000	0.000	0.130	0.000	0.000	200.023	0.000

Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	593	527	441	0	0	0	0	0	656	0
N.S.	1	0.89	0.74	0.00	0.00	0.00	0.00	0.00	1.11	0.00
time (sec)	N/A	2.180	2.903	0.000	0.000	0.000	0.000	0.000	1.228	0.000

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	372	328	322	0	0	0	0	0	360	0
N.S.	1	0.88	0.87	0.00	0.00	0.00	0.00	0.00	0.97	0.00
time (sec)	N/A	0.665	0.486	0.000	0.000	0.000	0.000	0.000	0.780	0.000

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	206	184	190	0	0	0	0	0	149	0
N.S.	1	0.89	0.92	0.00	0.00	0.00	0.00	0.00	0.72	0.00
time (sec)	N/A	0.396	0.325	0.000	0.000	0.000	0.000	0.000	0.460	0.000

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	23	25	25	20	25	43	29
N.S.	1	1.00	1.09	1.00	1.09	1.09	0.87	1.09	1.87	1.26
time (sec)	N/A	0.259	2.368	1.081	0.136	0.114	10.460	0.111	0.332	3.788



Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	23	25	36	0	25	65	29
N.S.	1	1.00	1.09	1.00	1.09	1.57	0.00	1.09	2.83	1.26
time (sec)	N/A	0.258	4.944	1.250	0.158	0.128	0.000	0.113	0.911	3.725

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	27	23	25	42	0	25	25	29
N.S.	1	1.00	1.08	0.92	1.00	1.68	0.00	1.00	1.00	1.16
time (sec)	N/A	0.287	0.937	0.495	0.156	0.117	0.000	0.158	200.025	3.676

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	27	23	25	25	24	25	25	29
N.S.	1	1.00	1.08	0.92	1.00	1.00	0.96	1.00	1.00	1.16
time (sec)	N/A	0.278	0.138	0.506	0.142	0.114	10.973	0.141	200.027	3.558

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	27	23	25	25	24	25	25	29
N.S.	1	1.00	1.08	0.92	1.00	1.00	0.96	1.00	1.00	1.16
time (sec)	N/A	0.280	0.975	0.516	0.166	0.104	4.469	0.124	200.024	3.598

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	27	23	25	45	24	25	25	29
N.S.	1	1.00	1.08	0.92	1.00	1.80	0.96	1.00	1.00	1.16
time (sec)	N/A	0.298	1.192	0.533	0.150	0.105	30.176	0.132	200.025	3.676

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F(-1)</b>	<b>F(-2)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	473	239	213	0	0	393	0	0	98	0
N.S.	1	0.51	0.45	0.00	0.00	0.83	0.00	0.00	0.21	0.00
time (sec)	N/A	0.732	0.372	0.000	0.000	0.146	0.000	0.000	5.301	0.000

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	316	168	178	0	0	336	0	0	78	0
N.S.	1	0.53	0.56	0.00	0.00	1.06	0.00	0.00	0.25	0.00
time (sec)	N/A	0.560	0.313	0.000	0.000	0.130	0.000	0.000	3.128	0.000

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	159	113	140	0	0	279	0	0	58	0
N.S.	1	0.71	0.88	0.00	0.00	1.75	0.00	0.00	0.36	0.00
time (sec)	N/A	0.379	0.296	0.000	0.000	0.118	0.000	0.000	1.491	0.000

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	28	24	100	37	34	26	50	30
N.S.	1	1.00	1.08	0.92	3.85	1.42	1.31	1.00	1.92	1.15
time (sec)	N/A	0.263	0.496	0.177	0.570	0.105	3.092	0.117	0.235	4.725

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	28	24	124	39	36	26	79	30
N.S.	1	1.00	1.08	0.92	4.77	1.50	1.38	1.00	3.04	1.15
time (sec)	N/A	0.279	7.132	0.281	0.630	0.108	28.589	0.118	0.326	4.230

## 2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio  $\frac{\text{number of rules}}{\text{integrand size}}$  is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [10] had the largest ratio of [1.5000000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	11	10	1.02	10	1.000
2	A	11	10	1.12	10	1.000
3	A	9	8	0.99	10	0.800
4	A	9	8	1.13	8	1.000
5	A	7	6	1.02	6	1.000
6	C	9	8	1.23	10	0.800
7	C	11	10	1.41	10	1.000
8	A	7	6	1.12	10	0.600
9	A	11	10	1.19	10	1.000
10	A	16	15	1.07	10	1.500
11	C	16	15	1.08	10	1.500
12	A	11	10	0.94	10	1.000
13	C	12	11	1.12	8	1.375
14	A	8	7	0.94	6	1.167
15	C	10	9	1.20	10	0.900
16	C	13	12	1.39	10	1.200
17	A	12	11	1.15	10	1.100
18	C	15	14	1.37	10	1.400
19	A	9	9	0.93	12	0.750
20	A	6	6	1.02	12	0.500
21	A	7	7	0.89	12	0.583

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	4	4	1.10	12	0.333
23	A	5	5	0.82	12	0.417
24	A	2	2	0.96	10	0.200
25	A	1	1	0.70	8	0.125
26	C	10	9	1.39	12	0.750
27	A	2	2	0.95	12	0.167
28	A	7	6	0.90	12	0.500
29	A	4	4	1.20	12	0.333
30	A	9	8	0.92	12	0.667
31	A	6	6	1.14	12	0.500
32	A	11	10	0.94	12	0.833
33	A	11	10	1.06	14	0.714
34	A	9	8	0.94	14	0.571
35	A	9	8	1.11	12	0.667
36	A	7	6	0.96	10	0.600
37	C	9	8	1.19	14	0.571
38	C	11	10	1.30	14	0.714
39	A	7	6	1.13	14	0.429
40	A	11	10	1.11	14	0.714
41	A	8	7	1.14	14	0.500
42	C	16	15	1.04	14	1.071
43	A	11	10	0.93	14	0.714
44	C	12	11	1.08	12	0.917
45	A	8	7	0.91	10	0.700
46	C	10	9	1.14	14	0.643
47	C	13	12	1.27	14	0.857
48	A	12	11	1.12	14	0.786
49	C	15	14	1.29	14	1.000
50	A	15	14	1.30	14	1.000
51	N/A	1	0	1.00	12	0.000
52	N/A	1	0	1.00	10	0.000
53	N/A	1	0	1.00	14	0.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	C	10	9	1.17	14	0.643
55	C	12	11	1.05	14	0.786
56	A	4	3	0.94	14	0.214
57	N/A	1	0	1.00	12	0.000
58	N/A	1	0	1.00	10	0.000
59	N/A	1	0	1.00	14	0.000
60	C	12	11	1.20	14	0.786
61	C	14	13	1.09	14	0.929
62	A	4	3	0.95	14	0.214
63	N/A	1	0	1.00	12	0.000
64	N/A	1	0	1.00	10	0.000
65	N/A	1	0	1.00	14	0.000
66	C	16	15	1.24	14	1.071
67	C	18	17	1.16	14	1.214
68	A	4	3	0.95	14	0.214
69	N/A	1	0	1.00	16	0.000
70	N/A	1	0	1.00	16	0.000
71	A	3	3	0.84	14	0.214
72	N/A	1	0	1.00	16	0.000
73	N/A	1	0	1.00	16	0.000
74	A	13	12	0.74	16	0.750
75	A	12	11	0.74	16	0.688
76	A	10	9	0.74	14	0.643
77	A	2	2	1.09	16	0.125
78	A	3	3	0.90	16	0.188
79	A	3	3	1.04	16	0.188
80	A	14	13	0.89	18	0.722
81	A	12	11	0.92	18	0.611
82	A	3	3	0.91	18	0.167
83	A	6	5	1.17	18	0.278
84	A	13	12	0.91	18	0.667

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
85	A	16	15	0.92	18	0.833
86	N/A	2	0	1.00	16	0.000
87	A	6	6	0.72	19	0.316
88	A	5	5	0.76	19	0.263
89	A	4	4	0.86	16	0.250
90	A	4	4	0.83	19	0.211
91	A	4	4	0.87	19	0.211
92	A	5	5	0.77	19	0.263
93	A	6	6	0.71	19	0.316
94	A	6	5	0.74	19	0.263
95	A	6	5	0.78	19	0.263
96	A	6	5	0.77	17	0.294
97	A	6	5	1.09	19	0.263
98	A	6	5	1.12	19	0.263
99	A	7	7	0.77	21	0.333
100	A	6	6	0.86	18	0.333
101	A	6	6	0.80	21	0.286
102	A	6	6	0.81	21	0.286
103	A	6	6	0.85	21	0.286
104	A	7	7	0.75	21	0.333
105	A	6	5	0.78	21	0.238
106	A	6	5	0.74	19	0.263
107	A	6	5	1.08	21	0.238
108	A	6	5	1.10	21	0.238
109	A	4	3	1.10	21	0.143
110	A	4	3	1.19	19	0.158
111	A	4	3	1.10	18	0.167
112	A	4	3	1.12	21	0.143
113	A	4	3	1.10	21	0.143
114	A	4	3	1.15	21	0.143
115	A	4	3	1.15	21	0.143
116	A	7	6	0.86	19	0.316

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
117	A	4	3	1.10	21	0.143
118	A	4	3	1.07	21	0.143
119	A	4	3	1.07	21	0.143
120	A	4	3	1.07	18	0.167
121	A	4	3	1.07	21	0.143
122	A	4	3	1.13	21	0.143
123	A	7	6	0.88	21	0.286
124	A	9	8	0.91	19	0.421
125	A	4	3	1.10	21	0.143
126	A	4	3	1.05	21	0.143
127	A	4	3	1.05	21	0.143
128	A	4	3	1.05	18	0.167
129	A	15	14	0.85	23	0.609
130	A	13	12	0.85	23	0.522
131	A	11	10	0.86	21	0.476
132	N/A	1	0	1.00	23	0.000
133	N/A	1	0	1.00	23	0.000
134	N/A	1	0	1.00	23	0.000
135	N/A	1	0	1.00	20	0.000
136	N/A	1	0	1.00	23	0.000
137	A	11	11	0.82	23	0.478
138	A	12	12	0.83	23	0.522
139	A	15	14	0.85	23	0.609
140	A	13	12	0.86	21	0.571
141	N/A	1	0	1.00	23	0.000
142	N/A	1	0	1.00	23	0.000
143	N/A	1	0	1.00	23	0.000
144	N/A	1	0	1.00	20	0.000
145	N/A	1	0	1.00	23	0.000
146	N/A	1	0	1.00	23	0.000
147	A	12	12	0.83	23	0.522

Continued on next page



Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
148	A	13	13	0.83	23	0.565
149	A	13	12	0.86	23	0.522
150	A	11	10	0.86	23	0.435
151	A	10	9	0.90	21	0.429
152	N/A	1	0	1.00	23	0.000
153	N/A	1	0	1.00	23	0.000
154	N/A	1	0	1.00	23	0.000
155	N/A	1	0	1.00	20	0.000
156	A	10	10	0.85	23	0.435
157	A	11	11	0.83	23	0.478
158	A	11	10	0.87	23	0.435
159	A	9	8	0.91	23	0.348
160	A	6	5	1.00	21	0.238
161	N/A	1	0	1.00	23	0.000
162	N/A	1	0	1.00	23	0.000
163	N/A	1	0	1.00	23	0.000
164	N/A	1	0	1.00	23	0.000
165	A	4	4	1.00	20	0.200
166	A	11	11	0.86	23	0.478
167	A	11	10	0.84	23	0.435
168	A	8	7	0.88	23	0.304
169	A	7	6	0.88	21	0.286
170	N/A	1	0	1.00	23	0.000
171	N/A	1	0	1.00	23	0.000
172	N/A	1	0	1.00	23	0.000
173	N/A	1	0	1.00	23	0.000
174	A	8	8	0.84	23	0.348
175	A	10	10	0.84	20	0.500
176	A	7	7	0.89	23	0.304
177	A	6	6	0.88	23	0.261
178	A	4	4	0.89	21	0.190

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
179	N/A	1	0	1.00	23	0.000
180	N/A	1	0	1.00	23	0.000
181	N/A	1	0	1.00	25	0.000
182	N/A	1	0	1.00	25	0.000
183	N/A	1	0	1.00	25	0.000
184	N/A	1	0	1.00	25	0.000
185	A	7	6	0.51	26	0.231
186	A	9	8	0.53	26	0.308
187	A	8	7	0.71	26	0.269
188	N/A	1	0	1.00	26	0.000
189	N/A	1	0	1.00	26	0.000

# CHAPTER 3

## LISTING OF INTEGRALS

3.1	$\int x^4 \operatorname{sech}^{-1}(ax)^2 dx$ . . . . .	97
3.2	$\int x^3 \operatorname{sech}^{-1}(ax)^2 dx$ . . . . .	104
3.3	$\int x^2 \operatorname{sech}^{-1}(ax)^2 dx$ . . . . .	111
3.4	$\int x \operatorname{sech}^{-1}(ax)^2 dx$ . . . . .	118
3.5	$\int \operatorname{sech}^{-1}(ax)^2 dx$ . . . . .	124
3.6	$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x} dx$ . . . . .	130
3.7	$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^2} dx$ . . . . .	137
3.8	$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^3} dx$ . . . . .	143
3.9	$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^4} dx$ . . . . .	149
3.10	$\int x^4 \operatorname{sech}^{-1}(ax)^3 dx$ . . . . .	156
3.11	$\int x^3 \operatorname{sech}^{-1}(ax)^3 dx$ . . . . .	165
3.12	$\int x^2 \operatorname{sech}^{-1}(ax)^3 dx$ . . . . .	174
3.13	$\int x \operatorname{sech}^{-1}(ax)^3 dx$ . . . . .	182
3.14	$\int \operatorname{sech}^{-1}(ax)^3 dx$ . . . . .	189
3.15	$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x} dx$ . . . . .	196
3.16	$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^2} dx$ . . . . .	203
3.17	$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^3} dx$ . . . . .	210
3.18	$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^4} dx$ . . . . .	217
3.19	$\int x^6 (a + b \operatorname{sech}^{-1}(cx)) dx$ . . . . .	225
3.20	$\int x^5 (a + b \operatorname{sech}^{-1}(cx)) dx$ . . . . .	233
3.21	$\int x^4 (a + b \operatorname{sech}^{-1}(cx)) dx$ . . . . .	240
3.22	$\int x^3 (a + b \operatorname{sech}^{-1}(cx)) dx$ . . . . .	247
3.23	$\int x^2 (a + b \operatorname{sech}^{-1}(cx)) dx$ . . . . .	253
3.24	$\int x (a + b \operatorname{sech}^{-1}(cx)) dx$ . . . . .	259

3.25	$\int (a + b \operatorname{sech}^{-1}(cx)) dx$	264
3.26	$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x} dx$	269
3.27	$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2} dx$	276
3.28	$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3} dx$	281
3.29	$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4} dx$	288
3.30	$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5} dx$	294
3.31	$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^6} dx$	301
3.32	$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^7} dx$	307
3.33	$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^2 dx$	315
3.34	$\int x^2 (a + b \operatorname{sech}^{-1}(cx))^2 dx$	322
3.35	$\int x (a + b \operatorname{sech}^{-1}(cx))^2 dx$	329
3.36	$\int (a + b \operatorname{sech}^{-1}(cx))^2 dx$	336
3.37	$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x} dx$	343
3.38	$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^2} dx$	351
3.39	$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^3} dx$	358
3.40	$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^4} dx$	365
3.41	$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^5} dx$	373
3.42	$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^3 dx$	380
3.43	$\int x^2 (a + b \operatorname{sech}^{-1}(cx))^3 dx$	390
3.44	$\int x (a + b \operatorname{sech}^{-1}(cx))^3 dx$	398
3.45	$\int (a + b \operatorname{sech}^{-1}(cx))^3 dx$	406
3.46	$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x} dx$	413
3.47	$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^2} dx$	421
3.48	$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^3} dx$	429
3.49	$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^4} dx$	437
3.50	$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^5} dx$	446
3.51	$\int \frac{x}{a + b \operatorname{sech}^{-1}(cx)} dx$	455
3.52	$\int \frac{1}{a + b \operatorname{sech}^{-1}(cx)} dx$	460
3.53	$\int \frac{1}{x(a + b \operatorname{sech}^{-1}(cx))} dx$	465

3.54	$\int \frac{1}{x^2(a+b\operatorname{sech}^{-1}(cx))} dx$	470
3.55	$\int \frac{1}{x^3(a+b\operatorname{sech}^{-1}(cx))} dx$	476
3.56	$\int \frac{1}{x^4(a+b\operatorname{sech}^{-1}(cx))} dx$	483
3.57	$\int \frac{x}{(a+b\operatorname{sech}^{-1}(cx))^2} dx$	489
3.58	$\int \frac{1}{(a+b\operatorname{sech}^{-1}(cx))^2} dx$	494
3.59	$\int \frac{1}{x(a+b\operatorname{sech}^{-1}(cx))^2} dx$	499
3.60	$\int \frac{1}{x^2(a+b\operatorname{sech}^{-1}(cx))^2} dx$	504
3.61	$\int \frac{1}{x^3(a+b\operatorname{sech}^{-1}(cx))^2} dx$	512
3.62	$\int \frac{1}{x^4(a+b\operatorname{sech}^{-1}(cx))^2} dx$	520
3.63	$\int \frac{x}{(a+b\operatorname{sech}^{-1}(cx))^3} dx$	526
3.64	$\int \frac{1}{(a+b\operatorname{sech}^{-1}(cx))^3} dx$	531
3.65	$\int \frac{1}{x(a+b\operatorname{sech}^{-1}(cx))^3} dx$	536
3.66	$\int \frac{1}{x^2(a+b\operatorname{sech}^{-1}(cx))^3} dx$	541
3.67	$\int \frac{1}{x^3(a+b\operatorname{sech}^{-1}(cx))^3} dx$	551
3.68	$\int \frac{1}{x^4(a+b\operatorname{sech}^{-1}(cx))^3} dx$	562
3.69	$\int (dx)^m (a + b\operatorname{sech}^{-1}(cx))^3 dx$	569
3.70	$\int (dx)^m (a + b\operatorname{sech}^{-1}(cx))^2 dx$	574
3.71	$\int (dx)^m (a + b\operatorname{sech}^{-1}(cx)) dx$	579
3.72	$\int \frac{(dx)^m}{a+b\operatorname{sech}^{-1}(cx)} dx$	584
3.73	$\int \frac{(dx)^m}{(a+b\operatorname{sech}^{-1}(cx))^2} dx$	589
3.74	$\int (d + ex)^3 (a + b\operatorname{sech}^{-1}(cx)) dx$	594
3.75	$\int (d + ex)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$	604
3.76	$\int (d + ex) (a + b\operatorname{sech}^{-1}(cx)) dx$	613
3.77	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{d+ex} dx$	621
3.78	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex)^2} dx$	628
3.79	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex)^3} dx$	635
3.80	$\int (d + ex)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx$	643
3.81	$\int \sqrt{d + ex} (a + b\operatorname{sech}^{-1}(cx)) dx$	655

3.82	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{\sqrt{d+ex}} dx$	665
3.83	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex)^{3/2}} dx$	672
3.84	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex)^{5/2}} dx$	679
3.85	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex)^{7/2}} dx$	689
3.86	$\int (d+ex)^m (a+b\operatorname{sech}^{-1}(cx)) dx$	701
3.87	$\int x^4 (d+ex^2) (a+b\operatorname{sech}^{-1}(cx)) dx$	706
3.88	$\int x^2 (d+ex^2) (a+b\operatorname{sech}^{-1}(cx)) dx$	714
3.89	$\int (d+ex^2) (a+b\operatorname{sech}^{-1}(cx)) dx$	721
3.90	$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^2} dx$	728
3.91	$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx$	735
3.92	$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx$	742
3.93	$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^8} dx$	749
3.94	$\int x^5 (d+ex^2) (a+b\operatorname{sech}^{-1}(cx)) dx$	757
3.95	$\int x^3 (d+ex^2) (a+b\operatorname{sech}^{-1}(cx)) dx$	764
3.96	$\int x (d+ex^2) (a+b\operatorname{sech}^{-1}(cx)) dx$	771
3.97	$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x} dx$	778
3.98	$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^3} dx$	785
3.99	$\int x^2 (d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx)) dx$	792
3.100	$\int (d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx)) dx$	801
3.101	$\int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x^2} dx$	809
3.102	$\int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x^4} dx$	817
3.103	$\int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x^6} dx$	825
3.104	$\int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x^8} dx$	833
3.105	$\int x^3 (d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx)) dx$	841
3.106	$\int x (d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx)) dx$	849
3.107	$\int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x} dx$	857
3.108	$\int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x^3} dx$	865
3.109	$\int \frac{x^2 (a+b\operatorname{sech}^{-1}(cx))}{d+ex^2} dx$	872
3.110	$\int \frac{x (a+b\operatorname{sech}^{-1}(cx))}{d+ex^2} dx$	881
3.111	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{d+ex^2} dx$	890

3.112	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x(d+ex^2)} dx$	898
3.113	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^2(d+ex^2)} dx$	906
3.114	$\int \frac{x^5(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^2} dx$	915
3.115	$\int \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^2} dx$	925
3.116	$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^2} dx$	935
3.117	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x(d+ex^2)^2} dx$	943
3.118	$\int \frac{x^4(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^2} dx$	952
3.119	$\int \frac{x^2(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^2} dx$	962
3.120	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex^2)^2} dx$	972
3.121	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^2(d+ex^2)^2} dx$	982
3.122	$\int \frac{x^5(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^3} dx$	992
3.123	$\int \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^3} dx$	1002
3.124	$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^3} dx$	1010
3.125	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x(d+ex^2)^3} dx$	1020
3.126	$\int \frac{x^4(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^3} dx$	1030
3.127	$\int \frac{x^2(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^3} dx$	1039
3.128	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex^2)^3} dx$	1048
3.129	$\int x^5\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1057
3.130	$\int x^3\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1069
3.131	$\int x\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1079
3.132	$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x} dx$	1088
3.133	$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^3} dx$	1093
3.134	$\int x^2\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1098
3.135	$\int \sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1103
3.136	$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^2} dx$	1108
3.137	$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx$	1113

3.138	$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx$	1122
3.139	$\int x^3(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1132
3.140	$\int x(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1143
3.141	$\int \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{x} dx$	1153
3.142	$\int \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{x^3} dx$	1158
3.143	$\int x^2(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1163
3.144	$\int (d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx)) dx$	1168
3.145	$\int \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{x^2} dx$	1173
3.146	$\int \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx$	1178
3.147	$\int \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx$	1183
3.148	$\int \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{x^8} dx$	1193
3.149	$\int \frac{x^5(a+b\operatorname{sech}^{-1}(cx))}{\sqrt{d+ex^2}} dx$	1203
3.150	$\int \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{\sqrt{d+ex^2}} dx$	1214
3.151	$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{\sqrt{d+ex^2}} dx$	1223
3.152	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x\sqrt{d+ex^2}} dx$	1231
3.153	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^3\sqrt{d+ex^2}} dx$	1236
3.154	$\int \frac{x^2(a+b\operatorname{sech}^{-1}(cx))}{\sqrt{d+ex^2}} dx$	1241
3.155	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{\sqrt{d+ex^2}} dx$	1246
3.156	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^2\sqrt{d+ex^2}} dx$	1251
3.157	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^4\sqrt{d+ex^2}} dx$	1259
3.158	$\int \frac{x^5(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{3/2}} dx$	1269
3.159	$\int \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{3/2}} dx$	1278
3.160	$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{3/2}} dx$	1286
3.161	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x(d+ex^2)^{3/2}} dx$	1292
3.162	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^3(d+ex^2)^{3/2}} dx$	1297
3.163	$\int \frac{x^4(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{3/2}} dx$	1302



3.164	$\int \frac{x^2(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{3/2}} dx$	1307
3.165	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex^2)^{3/2}} dx$	1312
3.166	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^2(d+ex^2)^{3/2}} dx$	1318
3.167	$\int \frac{x^5(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{5/2}} dx$	1327
3.168	$\int \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{5/2}} dx$	1336
3.169	$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{5/2}} dx$	1344
3.170	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x(d+ex^2)^{5/2}} dx$	1351
3.171	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^3(d+ex^2)^{5/2}} dx$	1356
3.172	$\int \frac{x^6(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{5/2}} dx$	1361
3.173	$\int \frac{x^4(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{5/2}} dx$	1366
3.174	$\int \frac{x^2(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{5/2}} dx$	1371
3.175	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex^2)^{5/2}} dx$	1379
3.176	$\int (fx)^m (d+ex^2)^3 (a+b\operatorname{sech}^{-1}(cx)) dx$	1388
3.177	$\int (fx)^m (d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx)) dx$	1398
3.178	$\int (fx)^m (d+ex^2) (a+b\operatorname{sech}^{-1}(cx)) dx$	1406
3.179	$\int \frac{(fx)^m (a+b\operatorname{sech}^{-1}(cx))}{d+ex^2} dx$	1413
3.180	$\int \frac{(fx)^m (a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^2} dx$	1418
3.181	$\int (fx)^m (d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx)) dx$	1423
3.182	$\int (fx)^m \sqrt{d+ex^2} (a+b\operatorname{sech}^{-1}(cx)) dx$	1428
3.183	$\int \frac{(fx)^m (a+b\operatorname{sech}^{-1}(cx))}{\sqrt{d+ex^2}} dx$	1433
3.184	$\int \frac{(fx)^m (a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{3/2}} dx$	1438
3.185	$\int \frac{x^{11}(a+b\operatorname{sech}^{-1}(cx))}{\sqrt{1-c^4x^4}} dx$	1443
3.186	$\int \frac{x^7(a+b\operatorname{sech}^{-1}(cx))}{\sqrt{1-c^4x^4}} dx$	1451
3.187	$\int \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{\sqrt{1-c^4x^4}} dx$	1459
3.188	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x\sqrt{1-c^4x^4}} dx$	1466
3.189	$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^5\sqrt{1-c^4x^4}} dx$	1471

### 3.1 $\int x^4 \operatorname{sech}^{-1}(ax)^2 dx$

Optimal result	97
Mathematica [A] (verified)	98
Rubi [A] (verified)	98
Maple [A] (verified)	101
Fricas [F]	102
Sympy [F]	102
Maxima [F(-1)]	102
Giac [F]	103
Mupad [F(-1)]	103
Reduce [F]	103

#### Optimal result

Integrand size = 10, antiderivative size = 164

$$\int x^4 \operatorname{sech}^{-1}(ax)^2 dx = -\frac{3x}{20a^4} - \frac{x^3}{30a^2} - \frac{3x\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)}{20a^4} - \frac{x^3\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)}{10a^2} + \frac{1}{5}x^5\operatorname{sech}^{-1}(ax)^2 - \frac{3\operatorname{sech}^{-1}(ax)\arctan\left(e^{\operatorname{sech}^{-1}(ax)}\right)}{10a^5} + \frac{3i\operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right)}{20a^5} - \frac{3i\operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(ax)}\right)}{20a^5}$$

output

```
-3/20*x/a^4-1/30*x^3/a^2-3/20*x*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*arcsech(a*x)/a^4-1/10*x^3*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*arcsech(a*x)/a^2+1/5*x^5*arcsech(a*x)^2-3/10*arcsech(a*x)*arctan(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2))/a^5+3/20*I*polylog(2,-I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a^5-3/20*I*polylog(2,I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a^5
```

**Mathematica [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.11

$$\int x^4 \operatorname{sech}^{-1}(ax)^2 dx$$

$$= \frac{-9ax - 2a^3x^3 - 9ax\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax) - 6a^3x^3\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax) + 12a^5x^5\operatorname{sech}^{-1}(ax)}{60a^5}$$

input

```
Integrate[x^4*ArcSech[a*x]^2,x]
```

output

```
(-9*a*x - 2*a^3*x^3 - 9*a*x*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x] - 6*a^3*x^3*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x] + 12*a^5*x^5*ArcSech[a*x]^2 + (9*I)*ArcSech[a*x]*Log[1 - I/E^ArcSech[a*x]] - (9*I)*ArcSech[a*x]*Log[1 + I/E^ArcSech[a*x]] + (9*I)*PolyLog[2, (-I)/E^ArcSech[a*x]] - (9*I)*PolyLog[2, I/E^ArcSech[a*x]])/(60*a^5)
```

**Rubi [A] (verified)**

Time = 0.60 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.02, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {6839, 5941, 3042, 4673, 3042, 4673, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4 \operatorname{sech}^{-1}(ax)^2 dx$$

$$\downarrow \text{6839}$$

$$\frac{\int a^5 x^5 \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 d\operatorname{sech}^{-1}(ax)}{a^5}$$

$$\downarrow \text{5941}$$

$$\frac{\frac{2}{5} \int a^5 x^5 \operatorname{sech}^{-1}(ax) d\operatorname{sech}^{-1}(ax) - \frac{1}{5} a^5 x^5 \operatorname{sech}^{-1}(ax)^2}{a^5}$$

$$\downarrow \text{3042}$$

$$\frac{-\frac{1}{5}a^5x^5\operatorname{sech}^{-1}(ax)^2 + \frac{2}{5}\int\operatorname{sech}^{-1}(ax)\csc\left(\operatorname{isech}^{-1}(ax) + \frac{\pi}{2}\right)^5 d\operatorname{sech}^{-1}(ax)}{a^5}$$

↓ 4673

$$\frac{\frac{2}{5}\left(\frac{3}{4}\int a^3x^3\operatorname{sech}^{-1}(ax)d\operatorname{sech}^{-1}(ax) + \frac{a^3x^3}{12} + \frac{1}{4}a^3x^3\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)\right) - \frac{1}{5}a^5x^5\operatorname{sech}^{-1}(ax)^2}{a^5}$$

↓ 3042

$$\frac{-\frac{1}{5}a^5x^5\operatorname{sech}^{-1}(ax)^2 + \frac{2}{5}\left(\frac{3}{4}\int\operatorname{sech}^{-1}(ax)\csc\left(\operatorname{isech}^{-1}(ax) + \frac{\pi}{2}\right)^3 d\operatorname{sech}^{-1}(ax) + \frac{a^3x^3}{12} + \frac{1}{4}a^3x^3\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)\right)}{a^5}$$

↓ 4673

$$\frac{\frac{2}{5}\left(\frac{3}{4}\left(\frac{1}{2}\int ax\operatorname{sech}^{-1}(ax)d\operatorname{sech}^{-1}(ax) + \frac{ax}{2} + \frac{1}{2}ax\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)\right) + \frac{a^3x^3}{12} + \frac{1}{4}a^3x^3\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)\right)}{a^5}$$

↓ 3042

$$\frac{-\frac{1}{5}a^5x^5\operatorname{sech}^{-1}(ax)^2 + \frac{2}{5}\left(\frac{3}{4}\left(\frac{1}{2}\int\operatorname{sech}^{-1}(ax)\csc\left(\operatorname{isech}^{-1}(ax) + \frac{\pi}{2}\right) d\operatorname{sech}^{-1}(ax) + \frac{ax}{2} + \frac{1}{2}ax\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)\right)\right)}{a^5}$$

↓ 4668

$$\frac{-\frac{1}{5}a^5x^5\operatorname{sech}^{-1}(ax)^2 + \frac{2}{5}\left(\frac{3}{4}\left(\frac{1}{2}\left(-i\int\log\left(1 - ie^{\operatorname{sech}^{-1}(ax)}\right) d\operatorname{sech}^{-1}(ax) + i\int\log\left(1 + ie^{\operatorname{sech}^{-1}(ax)}\right) d\operatorname{sech}^{-1}(ax)\right)\right)\right)}{a^5}$$

↓ 2715

$$\frac{-\frac{1}{5}a^5x^5\operatorname{sech}^{-1}(ax)^2 + \frac{2}{5}\left(\frac{3}{4}\left(\frac{1}{2}\left(-i\int e^{-\operatorname{sech}^{-1}(ax)}\log\left(1 - ie^{\operatorname{sech}^{-1}(ax)}\right) de^{\operatorname{sech}^{-1}(ax)} + i\int e^{-\operatorname{sech}^{-1}(ax)}\log\left(1 + ie^{\operatorname{sech}^{-1}(ax)}\right) de^{\operatorname{sech}^{-1}(ax)}\right)\right)\right)}{a^5}$$

↓ 2838

$$\frac{-\frac{1}{5}a^5x^5\operatorname{sech}^{-1}(ax)^2 + \frac{2}{5}\left(\frac{a^3x^3}{12} + \frac{1}{4}a^3x^3\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax) + \frac{3}{4}\left(\frac{1}{2}\left(2\operatorname{sech}^{-1}(ax)\arctan\left(e^{\operatorname{sech}^{-1}(ax)}\right) - \right)\right)\right)}{a^5}$$

input `Int [x^4*ArcSech [a*x]^2, x]`

output

```

-((-1/5*(a^5*x^5*ArcSech[a*x]^2) + (2*((a^3*x^3)/12 + (a^3*x^3*Sqrt[(1 - a
*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x])/4 + (3*((a*x)/2 + (a*x*Sqrt[(1 - a*
x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x])/2 + (2*ArcSech[a*x]*ArcTan[E^ArcSech
[a*x]] - I*PolyLog[2, (-I)*E^ArcSech[a*x]] + I*PolyLog[2, I*E^ArcSech[a*x]
])/2))/4))/5)/a^5)

```

### Defintions of rubi rules used

rule 2715

```

Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

```

rule 2838

```

Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

```

rule 3042

```

Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

rule 4668

```

Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_
))^m, x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(
I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[
1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c
+ d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c
, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]

```

rule 4673

```

Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^n*((c_.) + (d_.)*(x_)), x_Symbol] :>
Simp[(-b^2)*(c + d*x)*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))),
x] + (-Simp[b^2*d*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x] + S
imp[b^2*((n - 2)/(n - 1)) Int[(c + d*x)*(b*Csc[e + f*x])^(n - 2), x], x])
/; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2]

```

rule 5941

```
Int[(x_)^(m_)*Sech[(a_) + (b_)*(x_)^(n_)]^(p_)*Tanh[(a_) + (b_)*(x_)
^(n_)]^(q_), x_Symbol] := Simp[(-x^(m - n + 1))*(Sech[a + b*x^n]^p/(b*n*p
)), x] + Simp[(m - n + 1)/(b*n*p) Int[x^(m - n)*Sech[a + b*x^n]^p, x], x]
/; FreeQ[{a, b, p}, x] && RationalQ[m] && IntegerQ[n] && GeQ[m - n, 0] &&
EqQ[q, 1]
```

rule 6839

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_)^(n_)*(x_)^(m_), x_Symbol] := Simp[
-(c^(m + 1))^( -1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

### Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.71

method	result
derivativedivides	$\frac{(12x^4a^4 \operatorname{arcsech}(xa)^2 - 6 \operatorname{arcsech}(xa) \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} x^3a^3 - 9 \operatorname{arcsech}(xa) \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} ax - 2a^2x^2 - 9)xa}{60} + \frac{3i \operatorname{arcsech}(xa)}{60}$
default	$\frac{(12x^4a^4 \operatorname{arcsech}(xa)^2 - 6 \operatorname{arcsech}(xa) \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} x^3a^3 - 9 \operatorname{arcsech}(xa) \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} ax - 2a^2x^2 - 9)xa}{60} + \frac{3i \operatorname{arcsech}(xa)}{60}$

input

```
int(x^4*arcsech(x*a)^2,x,method=_RETURNVERBOSE)
```

output

```
1/a^5*(1/60*(12*x^4*a^4*arcsech(x*a)^2-6*arcsech(x*a)*(-(a*x-1)/x/a)^(1/2)
*((a*x+1)/x/a)^(1/2)*x^3*a^3-9*arcsech(x*a)*(-(a*x-1)/x/a)^(1/2)*((a*x+1)/
x/a)^(1/2)*a*x-2*a^2*x^2-9)*x*a+3/20*I*arcsech(x*a)*ln(1+I*(1/a/x+(1/a/x-1
)^(1/2)*(1+1/a/x)^(1/2)))-3/20*I*arcsech(x*a)*ln(1-I*(1/a/x+(1/a/x-1)^(1/2
)*(1+1/a/x)^(1/2)))+3/20*I*dilog(1+I*(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2
)))-3/20*I*dilog(1-I*(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))))
```

**Fricas [F]**

$$\int x^4 \operatorname{sech}^{-1}(ax)^2 dx = \int x^4 \operatorname{arsech}(ax)^2 dx$$

input `integrate(x^4*arcsech(a*x)^2,x, algorithm="fricas")`

output `integral(x^4*arcsech(a*x)^2, x)`

**Sympy [F]**

$$\int x^4 \operatorname{sech}^{-1}(ax)^2 dx = \int x^4 \operatorname{asech}^2(ax) dx$$

input `integrate(x**4*asech(a*x)**2,x)`

output `Integral(x**4*asech(a*x)**2, x)`

**Maxima [F(-1)]**

Timed out.

$$\int x^4 \operatorname{sech}^{-1}(ax)^2 dx = \text{Timed out}$$

input `integrate(x^4*arcsech(a*x)^2,x, algorithm="maxima")`

output `Timed out`

**Giac [F]**

$$\int x^4 \operatorname{sech}^{-1}(ax)^2 dx = \int x^4 \operatorname{arsech}(ax)^2 dx$$

input `integrate(x^4*arcsech(a*x)^2,x, algorithm="giac")`

output `integrate(x^4*arcsech(a*x)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^4 \operatorname{sech}^{-1}(ax)^2 dx = \int x^4 \operatorname{acosh}\left(\frac{1}{ax}\right)^2 dx$$

input `int(x^4*acosh(1/(a*x))^2,x)`

output `int(x^4*acosh(1/(a*x))^2, x)`

**Reduce [F]**

$$\int x^4 \operatorname{sech}^{-1}(ax)^2 dx = \int \operatorname{asech}(ax)^2 x^4 dx$$

input `int(x^4*asech(a*x)^2,x)`

output `int(asech(a*x)**2*x**4,x)`



### 3.2 $\int x^3 \operatorname{sech}^{-1}(ax)^2 dx$

Optimal result	104
Mathematica [A] (verified)	104
Rubi [A] (verified)	105
Maple [A] (verified)	108
Fricas [A] (verification not implemented)	108
Sympy [F]	109
Maxima [F]	109
Giac [F]	109
Mupad [F(-1)]	110
Reduce [F]	110

#### Optimal result

Integrand size = 10, antiderivative size = 104

$$\int x^3 \operatorname{sech}^{-1}(ax)^2 dx = -\frac{x^2}{12a^2} - \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)}{3a^4} - \frac{x^2\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)}{6a^2} + \frac{1}{4}x^4\operatorname{sech}^{-1}(ax)^2 - \frac{\log(x)}{3a^4}$$

output

```
-1/12*x^2/a^2-1/3*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*arcsech(a*x)/a^4-1/6*x^2*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*arcsech(a*x)/a^2+1/4*x^4*arcsech(a*x)^2-1/3*ln(x)/a^4
```

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.74

$$\int x^3 \operatorname{sech}^{-1}(ax)^2 dx = \frac{a^2 x^2 + 2\sqrt{\frac{1-ax}{1+ax}}(2 + 2ax + a^2 x^2 + a^3 x^3) \operatorname{sech}^{-1}(ax) - 3a^4 x^4 \operatorname{sech}^{-1}(ax)^2 + 4 \log(x)}{12a^4}$$

input `Integrate[x^3*ArcSech[a*x]^2,x]`

output 
$$\frac{-1/12*(a^2*x^2 + 2*sqrt[(1 - a*x)/(1 + a*x)]*(2 + 2*a*x + a^2*x^2 + a^3*x^3)*ArcSech[a*x] - 3*a^4*x^4*ArcSech[a*x]^2 + 4*Log[x])}{a^4}$$

### Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.12, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {6839, 5941, 3042, 4673, 3042, 4672, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^3 \operatorname{sech}^{-1}(ax)^2 dx \\ & \quad \downarrow 6839 \\ & \frac{\int a^4 x^4 \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 d\operatorname{sech}^{-1}(ax)}{a^4} \\ & \quad \downarrow 5941 \\ & \frac{\frac{1}{2} \int a^4 x^4 \operatorname{sech}^{-1}(ax) d\operatorname{sech}^{-1}(ax) - \frac{1}{4} a^4 x^4 \operatorname{sech}^{-1}(ax)^2}{a^4} \\ & \quad \downarrow 3042 \\ & \frac{-\frac{1}{4} a^4 x^4 \operatorname{sech}^{-1}(ax)^2 + \frac{1}{2} \int \operatorname{sech}^{-1}(ax) \csc\left(i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2}\right)^4 d\operatorname{sech}^{-1}(ax)}{a^4} \\ & \quad \downarrow 4673 \\ & \frac{\frac{1}{2} \left( \frac{2}{3} \int a^2 x^2 \operatorname{sech}^{-1}(ax) d\operatorname{sech}^{-1}(ax) + \frac{a^2 x^2}{6} + \frac{1}{3} a^2 x^2 \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax) \right) - \frac{1}{4} a^4 x^4 \operatorname{sech}^{-1}(ax)^2}{a^4} \\ & \quad \downarrow 3042 \\ & \frac{-\frac{1}{4} a^4 x^4 \operatorname{sech}^{-1}(ax)^2 + \frac{1}{2} \left( \frac{2}{3} \int \operatorname{sech}^{-1}(ax) \csc\left(i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2}\right)^2 d\operatorname{sech}^{-1}(ax) + \frac{a^2 x^2}{6} + \frac{1}{3} a^2 x^2 \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax) \right)}{a^4} \end{aligned}$$

↓ 4672

$$\frac{-\frac{1}{4}a^4x^4\operatorname{sech}^{-1}(ax)^2 + \frac{1}{2}\left(\frac{2}{3}\left(\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax) - i\int -i\sqrt{\frac{1-ax}{ax+1}}(ax+1)d\operatorname{sech}^{-1}(ax)\right) + \frac{a^2x^2}{6} + \frac{1}{3}a^2x^2\right)}{a^4}$$

↓ 26

$$\frac{\frac{1}{2}\left(\frac{2}{3}\left(\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax) - \int \sqrt{\frac{1-ax}{ax+1}}(ax+1)d\operatorname{sech}^{-1}(ax)\right) + \frac{a^2x^2}{6} + \frac{1}{3}a^2x^2\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)\right)}{a^4}$$

↓ 3042

$$\frac{-\frac{1}{4}a^4x^4\operatorname{sech}^{-1}(ax)^2 + \frac{1}{2}\left(\frac{2}{3}\left(\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax) - \int -i\tan(i\operatorname{sech}^{-1}(ax))d\operatorname{sech}^{-1}(ax)\right) + \frac{a^2x^2}{6} + \frac{1}{3}a^2x^2\right)}{a^4}$$

↓ 26

$$\frac{-\frac{1}{4}a^4x^4\operatorname{sech}^{-1}(ax)^2 + \frac{1}{2}\left(\frac{2}{3}\left(\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax) + i\int \tan(i\operatorname{sech}^{-1}(ax))d\operatorname{sech}^{-1}(ax)\right) + \frac{a^2x^2}{6} + \frac{1}{3}a^2x^2\right)}{a^4}$$

↓ 3956

$$\frac{\frac{1}{2}\left(\frac{a^2x^2}{6} + \frac{1}{3}a^2x^2\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax) + \frac{2}{3}\left(\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax) - \log\left(\frac{1}{ax}\right)\right)\right) - \frac{1}{4}a^4x^4\operatorname{sech}^{-1}(ax)^2}{a^4}$$

input

```
Int [x^3*ArcSech[a*x]^2, x]
```

output

```
-((-1/4*(a^4*x^4*ArcSech[a*x]^2) + ((a^2*x^2)/6 + (a^2*x^2*sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x])/3 + (2*(sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x] - Log[1/(a*x)]))/3)/2)/a^4
```

## Definitions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4672 `Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(-c + d*x)^m*(Cot[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`
- rule 4673 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((c_.) + (d_.)*(x_)), x_Symbol] := Simp[(-b^2)*(c + d*x)*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))), x] + (-Simp[b^2*d*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(c + d*x)*(b*Csc[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2]`
- rule 5941 `Int[(x_)^(m_.)*Sech[(a_.) + (b_.)*(x_)^(n_.)]^(p_.)*Tanh[(a_.) + (b_.)*(x_)^(n_.)]^(q_.), x_Symbol] := Simp[(-x^(m - n + 1))*(Sech[a + b*x^n]^p/(b*n*p)), x] + Simp[(m - n + 1)/(b*n*p) Int[x^(m - n)*Sech[a + b*x^n]^p, x], x] /; FreeQ[{a, b, p}, x] && RationalQ[m] && IntegerQ[n] && GeQ[m - n, 0] && EqQ[q, 1]`
- rule 6839 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))^(n_)*(x_)^(m_.), x_Symbol] := Simp[-(c^(m + 1))^( -1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])`

**Maple [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.44

method	result
derivativedivides	$\frac{-\frac{\operatorname{arcsech}(xa)}{3} + \frac{x^4 a^4 \operatorname{arcsech}(xa)^2}{4} - \frac{\operatorname{arcsech}(xa) \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} x^3 a^3}{6} - \frac{\operatorname{arcsech}(xa) \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} ax}{3} - \frac{a^2 x^2}{12} + \ln\left(1 + \left(\frac{1}{ax}\right)^2\right)}{a^4}$
default	$\frac{-\frac{\operatorname{arcsech}(xa)}{3} + \frac{x^4 a^4 \operatorname{arcsech}(xa)^2}{4} - \frac{\operatorname{arcsech}(xa) \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} x^3 a^3}{6} - \frac{\operatorname{arcsech}(xa) \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} ax}{3} - \frac{a^2 x^2}{12} + \ln\left(1 + \left(\frac{1}{ax}\right)^2\right)}{a^4}$

input `int(x^3*arcsech(x*a)^2,x,method=_RETURNVERBOSE)`output 
$$\frac{1}{a^4} \left( -\frac{1}{3} \operatorname{arcsech}(x/a) + \frac{1}{4} x^4 a^4 \operatorname{arcsech}(x/a)^2 - \frac{1}{6} \operatorname{arcsech}(x/a) \left( -\frac{a^2 x^2 - 1}{x/a} \right)^{1/2} \left( \frac{a^2 x^2 + 1}{x/a} \right)^{1/2} x^3 a^3 - \frac{1}{3} \operatorname{arcsech}(x/a) \left( -\frac{a^2 x^2 - 1}{x/a} \right)^{1/2} \left( \frac{a^2 x^2 + 1}{x/a} \right)^{1/2} a x - \frac{a^2 x^2}{12} + \ln\left(1 + \left(\frac{1}{ax}\right)^2\right) \right)$$
**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.20

$$\int x^3 \operatorname{sech}^{-1}(ax)^2 dx$$

$$= \frac{3 a^4 x^4 \log\left(\frac{ax \sqrt{-\frac{a^2 x^2 - 1}{a^2 x^2}} + 1}{ax}\right)^2 - a^2 x^2 - 2(a^3 x^3 + 2ax) \sqrt{-\frac{a^2 x^2 - 1}{a^2 x^2}} \log\left(\frac{ax \sqrt{-\frac{a^2 x^2 - 1}{a^2 x^2}} + 1}{ax}\right) - 4 \log(x)}{12 a^4}$$

input `integrate(x^3*arcsech(a*x)^2,x, algorithm="fricas")`output 
$$\frac{1}{12} \left( 3 a^4 x^4 \log\left(\frac{a x \sqrt{-(a^2 x^2 - 1)/(a^2 x^2)} + 1}{a x}\right)^2 - a^2 x^2 - 2(a^3 x^3 + 2 a x) \sqrt{-(a^2 x^2 - 1)/(a^2 x^2)} \log\left(\frac{a x \sqrt{-(a^2 x^2 - 1)/(a^2 x^2)} + 1}{a x}\right) - 4 \log(x) \right) / a^4$$

**Sympy [F]**

$$\int x^3 \operatorname{sech}^{-1}(ax)^2 dx = \int x^3 \operatorname{asech}^2(ax) dx$$

input `integrate(x**3*asech(a*x)**2,x)`

output `Integral(x**3*asech(a*x)**2, x)`

**Maxima [F]**

$$\int x^3 \operatorname{sech}^{-1}(ax)^2 dx = \int x^3 \operatorname{arsech}(ax)^2 dx$$

input `integrate(x^3*arcsech(a*x)^2,x, algorithm="maxima")`

output `integrate(x^3*arcsech(a*x)^2, x)`

**Giac [F]**

$$\int x^3 \operatorname{sech}^{-1}(ax)^2 dx = \int x^3 \operatorname{arsech}(ax)^2 dx$$

input `integrate(x^3*arcsech(a*x)^2,x, algorithm="giac")`

output `integrate(x^3*arcsech(a*x)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 \operatorname{sech}^{-1}(ax)^2 dx = \int x^3 \operatorname{acosh}\left(\frac{1}{ax}\right)^2 dx$$

input `int(x^3*acosh(1/(a*x))^2,x)`output `int(x^3*acosh(1/(a*x))^2, x)`**Reduce [F]**

$$\int x^3 \operatorname{sech}^{-1}(ax)^2 dx = \int a \operatorname{sech}(ax)^2 x^3 dx$$

input `int(x^3*asech(a*x)^2,x)`output `int(asech(a*x)**2*x**3,x)`

### 3.3 $\int x^2 \operatorname{sech}^{-1}(ax)^2 dx$

Optimal result	111
Mathematica [A] (verified)	112
Rubi [A] (verified)	112
Maple [A] (verified)	115
Fricas [F]	115
Sympy [F]	116
Maxima [F]	116
Giac [F]	116
Mupad [F(-1)]	117
Reduce [F]	117

#### Optimal result

Integrand size = 10, antiderivative size = 117

$$\int x^2 \operatorname{sech}^{-1}(ax)^2 dx = -\frac{x}{3a^2} - \frac{x \sqrt{\frac{1-ax}{1+ax}} (1+ax) \operatorname{sech}^{-1}(ax)}{3a^2} + \frac{1}{3} x^3 \operatorname{sech}^{-1}(ax)^2 - \frac{2 \operatorname{sech}^{-1}(ax) \arctan\left(e^{\operatorname{sech}^{-1}(ax)}\right)}{3a^3} + \frac{i \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right)}{3a^3} - \frac{i \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(ax)}\right)}{3a^3}$$

output

```
-1/3*x/a^2-1/3*x*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*arcsech(a*x)/a^2+1/3*x^3
*arcsech(a*x)^2-2/3*arcsech(a*x)*arctan(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(
1/2))/a^3+1/3*I*polylog(2,-I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a^3
-1/3*I*polylog(2,I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a^3
```



**Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.18

$$\int x^2 \operatorname{sech}^{-1}(ax)^2 dx = \frac{-ax - ax \sqrt{\frac{1-ax}{1+ax}}(1+ax) \operatorname{sech}^{-1}(ax) + a^3 x^3 \operatorname{sech}^{-1}(ax)^2 + i \operatorname{sech}^{-1}(ax) \log(1 - i e^{-\operatorname{sech}^{-1}(ax)}) - i \operatorname{sech}^{-1}(ax)}{3a^3}$$

input

```
Integrate[x^2*ArcSech[a*x]^2,x]
```

output

```
(-(a*x) - a*x*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x] + a^3*x^3*ArcSech[a*x]^2 + I*ArcSech[a*x]*Log[1 - I/E^ArcSech[a*x]] - I*ArcSech[a*x]*Log[1 + I/E^ArcSech[a*x]] + I*PolyLog[2, (-I)/E^ArcSech[a*x]] - I*PolyLog[2, I/E^ArcSech[a*x]])/(3*a^3)
```

**Rubi [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.99, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$ , Rules used = {6839, 5941, 3042, 4673, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2 \operatorname{sech}^{-1}(ax)^2 dx \\ & \quad \downarrow \text{6839} \\ & \frac{\int a^3 x^3 \sqrt{\frac{1-ax}{ax+1}}(ax+1) \operatorname{sech}^{-1}(ax)^2 d \operatorname{sech}^{-1}(ax)}{a^3} \\ & \quad \downarrow \text{5941} \\ & \frac{\frac{2}{3} \int a^3 x^3 \operatorname{sech}^{-1}(ax) d \operatorname{sech}^{-1}(ax) - \frac{1}{3} a^3 x^3 \operatorname{sech}^{-1}(ax)^2}{a^3} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\frac{-\frac{1}{3}a^3x^3\operatorname{sech}^{-1}(ax)^2 + \frac{2}{3}\int \operatorname{sech}^{-1}(ax) \csc\left(\operatorname{isech}^{-1}(ax) + \frac{\pi}{2}\right)^3 d\operatorname{sech}^{-1}(ax)}{a^3}$$

↓ 4673

$$\frac{\frac{2}{3}\left(\frac{1}{2}\int ax\operatorname{sech}^{-1}(ax)d\operatorname{sech}^{-1}(ax) + \frac{ax}{2} + \frac{1}{2}ax\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)\right) - \frac{1}{3}a^3x^3\operatorname{sech}^{-1}(ax)^2}{a^3}$$

↓ 3042

$$\frac{-\frac{1}{3}a^3x^3\operatorname{sech}^{-1}(ax)^2 + \frac{2}{3}\left(\frac{1}{2}\int \operatorname{sech}^{-1}(ax) \csc\left(\operatorname{isech}^{-1}(ax) + \frac{\pi}{2}\right) d\operatorname{sech}^{-1}(ax) + \frac{ax}{2} + \frac{1}{2}ax\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)\right)}{a^3}$$

↓ 4668

$$\frac{-\frac{1}{3}a^3x^3\operatorname{sech}^{-1}(ax)^2 + \frac{2}{3}\left(\frac{1}{2}\left(-i\int \log\left(1 - ie^{\operatorname{sech}^{-1}(ax)}\right) d\operatorname{sech}^{-1}(ax) + i\int \log\left(1 + ie^{\operatorname{sech}^{-1}(ax)}\right) d\operatorname{sech}^{-1}(ax) + \right)}{a^3}$$

↓ 2715

$$\frac{-\frac{1}{3}a^3x^3\operatorname{sech}^{-1}(ax)^2 + \frac{2}{3}\left(\frac{1}{2}\left(-i\int e^{-\operatorname{sech}^{-1}(ax)} \log\left(1 - ie^{\operatorname{sech}^{-1}(ax)}\right) de^{\operatorname{sech}^{-1}(ax)} + i\int e^{-\operatorname{sech}^{-1}(ax)} \log\left(1 + ie^{\operatorname{sech}^{-1}(ax)}\right) de^{\operatorname{sech}^{-1}(ax)} + \right)}{a^3}$$

↓ 2838

$$\frac{-\frac{1}{3}a^3x^3\operatorname{sech}^{-1}(ax)^2 + \frac{2}{3}\left(\frac{1}{2}\left(2\operatorname{sech}^{-1}(ax) \arctan\left(e^{\operatorname{sech}^{-1}(ax)}\right) - i\operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right) + i\operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(ax)}\right)\right)\right)}{a^3}$$

input `Int [x^2*ArcSech[a*x]^2, x]`

output `-((-1/3*(a^3*x^3*ArcSech[a*x]^2) + (2*((a*x)/2 + (a*x*Sqrt[(1 - a*x)/(1 + a*x)])*(1 + a*x)*ArcSech[a*x])/2 + (2*ArcSech[a*x]*ArcTan[E^ArcSech[a*x]] - I*PolyLog[2, (-1)*E^ArcSech[a*x]] + I*PolyLog[2, I*E^ArcSech[a*x]])/2))/3)/a^3)`

## Definitions of rubi rules used

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_
))^m_], x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(
I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[
1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c
+ d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c
, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 4673 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^n_)*((c_.) + (d_.)*(x_)), x_Symbol] :>
Simp[(-b^2)*(c + d*x)*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))),
x] + (-Simp[b^2*d*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x] + S
imp[b^2*((n - 2)/(n - 1)) Int[(c + d*x)*(b*Csc[e + f*x])^(n - 2), x], x])
/; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2]`

rule 5941 `Int[(x_)^(m_.)*Sech[(a_.) + (b_.)*(x_)^(n_.)]^(p_.)*Tanh[(a_.) + (b_.)*(x_
)^(n_.)]^(q_.), x_Symbol] :> Simp[(-x^(m - n + 1))*(Sech[a + b*x^n]^p/(b*n*p
)), x] + Simp[(m - n + 1)/(b*n*p) Int[x^(m - n)*Sech[a + b*x^n]^p, x], x]
/; FreeQ[{a, b, p}, x] && RationalQ[m] && IntegerQ[n] && GeQ[m - n, 0] &&
EqQ[q, 1]`

rule 6839

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_.), x_Symbol] := Simp[
-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

**Maple [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.97

method	result
derivativedivides	$\frac{\left(-\operatorname{arcsech}(xa)\sqrt{-\frac{xa-1}{xa}}\sqrt{\frac{xa+1}{xa}}ax+x^2a^2\operatorname{arcsech}(xa)^2-1\right)xa}{3} + \frac{i\operatorname{arcsech}(xa)\ln\left(1+i\left(\frac{1}{ax}+\sqrt{\frac{1}{ax}-1}\sqrt{1+\frac{1}{ax}}\right)\right)}{3} - \frac{i\operatorname{arcsech}(xa)}{a^3}$
default	$\frac{\left(-\operatorname{arcsech}(xa)\sqrt{-\frac{xa-1}{xa}}\sqrt{\frac{xa+1}{xa}}ax+x^2a^2\operatorname{arcsech}(xa)^2-1\right)xa}{3} + \frac{i\operatorname{arcsech}(xa)\ln\left(1+i\left(\frac{1}{ax}+\sqrt{\frac{1}{ax}-1}\sqrt{1+\frac{1}{ax}}\right)\right)}{3} - \frac{i\operatorname{arcsech}(xa)}{a^3}$

input

```
int(x^2*arcsech(x*a)^2,x,method=_RETURNVERBOSE)
```

output

```
1/a^3*(1/3*(-arcsech(x*a)*(-(a*x-1)/x/a)^(1/2)*((a*x+1)/x/a)^(1/2)*a*x+x^2
*a^2*arcsech(x*a)^2-1)*x*a+1/3*I*arcsech(x*a)*ln(1+I*(1/a/x+(1/a/x-1)^(1/2)
)*(1+1/a/x)^(1/2)))-1/3*I*arcsech(x*a)*ln(1-I*(1/a/x+(1/a/x-1)^(1/2)*(1+1/
a/x)^(1/2)))+1/3*I*dilog(1+I*(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)))-1/3*
I*dilog(1-I*(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))))
```

**Fricas [F]**

$$\int x^2 \operatorname{sech}^{-1}(ax)^2 dx = \int x^2 \operatorname{ar} \operatorname{sech}(ax)^2 dx$$

input

```
integrate(x^2*arcsech(a*x)^2,x, algorithm="fricas")
```

output

```
integral(x^2*arcsech(a*x)^2, x)
```

**Sympy [F]**

$$\int x^2 \operatorname{sech}^{-1}(ax)^2 dx = \int x^2 \operatorname{asech}^2(ax) dx$$

input `integrate(x**2*asech(a*x)**2,x)`

output `Integral(x**2*asech(a*x)**2, x)`

**Maxima [F]**

$$\int x^2 \operatorname{sech}^{-1}(ax)^2 dx = \int x^2 \operatorname{arsech}(ax)^2 dx$$

input `integrate(x^2*arcsech(a*x)^2,x, algorithm="maxima")`

output `integrate(x^2*arcsech(a*x)^2, x)`

**Giac [F]**

$$\int x^2 \operatorname{sech}^{-1}(ax)^2 dx = \int x^2 \operatorname{arsech}(ax)^2 dx$$

input `integrate(x^2*arcsech(a*x)^2,x, algorithm="giac")`

output `integrate(x^2*arcsech(a*x)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{sech}^{-1}(ax)^2 dx = \int x^2 \operatorname{acosh}\left(\frac{1}{ax}\right)^2 dx$$

input `int(x^2*acosh(1/(a*x))^2,x)`output `int(x^2*acosh(1/(a*x))^2, x)`**Reduce [F]**

$$\int x^2 \operatorname{sech}^{-1}(ax)^2 dx = \int \operatorname{asech}(ax)^2 x^2 dx$$

input `int(x^2*asech(a*x)^2,x)`output `int(asech(a*x)**2*x**2,x)`

### 3.4 $\int x \operatorname{sech}^{-1}(ax)^2 dx$

Optimal result	118
Mathematica [A] (verified)	118
Rubi [A] (verified)	119
Maple [B] (verified)	121
Fricas [B] (verification not implemented)	121
Sympy [A] (verification not implemented)	122
Maxima [A] (verification not implemented)	122
Giac [F]	123
Mupad [F(-1)]	123
Reduce [F]	123

#### Optimal result

Integrand size = 8, antiderivative size = 53

$$\int x \operatorname{sech}^{-1}(ax)^2 dx = -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)}{a^2} + \frac{1}{2}x^2\operatorname{sech}^{-1}(ax)^2 - \frac{\log(x)}{a^2}$$

output

```
-((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*arcsech(a*x)/a^2+1/2*x^2*arcsech(a*x)^2-
ln(x)/a^2
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00

$$\int x \operatorname{sech}^{-1}(ax)^2 dx = -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)}{a^2} + \frac{1}{2}x^2\operatorname{sech}^{-1}(ax)^2 - \frac{\log(x)}{a^2}$$

input

```
Integrate[x*ArcSech[a*x]^2,x]
```

output

```
-((Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x])/a^2) + (x^2*ArcSech[a
*x]^2)/2 - Log[x]/a^2
```

**Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.13, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {6839, 5941, 3042, 4672, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \operatorname{sech}^{-1}(ax)^2 dx \\
 & \quad \downarrow \text{6839} \\
 & \frac{\int a^2 x^2 \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 d \operatorname{sech}^{-1}(ax)}{a^2} \\
 & \quad \downarrow \text{5941} \\
 & \frac{\int a^2 x^2 \operatorname{sech}^{-1}(ax) d \operatorname{sech}^{-1}(ax) - \frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^2}{a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^2 + \int \operatorname{sech}^{-1}(ax) \csc \left( i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2} \right)^2 d \operatorname{sech}^{-1}(ax)}{a^2} \\
 & \quad \downarrow \text{4672} \\
 & \frac{-i \int -i \sqrt{\frac{1-ax}{ax+1}} (ax+1) d \operatorname{sech}^{-1}(ax) - \frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^2 + \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)}{a^2} \\
 & \quad \downarrow \text{26} \\
 & \frac{-\int \sqrt{\frac{1-ax}{ax+1}} (ax+1) d \operatorname{sech}^{-1}(ax) - \frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^2 + \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)}{a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\int -i \tan \left( i \operatorname{sech}^{-1}(ax) \right) d \operatorname{sech}^{-1}(ax) - \frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^2 + \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)}{a^2} \\
 & \quad \downarrow \text{26} \\
 & \frac{i \int \tan \left( i \operatorname{sech}^{-1}(ax) \right) d \operatorname{sech}^{-1}(ax) - \frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^2 + \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)}{a^2}
 \end{aligned}$$



↓ 3956

$$\frac{-\frac{1}{2}a^2x^2\operatorname{sech}^{-1}(ax)^2 - \log\left(\frac{1}{ax}\right) + \sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{a^2}$$

input `Int[x*ArcSech[a*x]^2,x]`

output `-((Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x] - (a^2*x^2*ArcSech[a*x]^2)/2 - Log[1/(a*x)])/a^2)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4672 `Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 5941 `Int[(x_)^(m_.)*Sech[(a_.) + (b_.)*(x_)^(n_.)]^(p_.)*Tanh[(a_.) + (b_.)*(x_)^(n_.)]^(q_.), x_Symbol] := Simp[(-x^(m - n + 1))*(Sech[a + b*x^n]^p/(b*n*p)), x] + Simp[(m - n + 1)/(b*n*p) Int[x^(m - n)*Sech[a + b*x^n]^p, x], x] /; FreeQ[{a, b, p}, x] && RationalQ[m] && IntegerQ[n] && GeQ[m - n, 0] && EqQ[q, 1]`

rule 6839

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))^ (n_.)*(x_)^(m_.), x_Symbol] := Simp[
-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 99 vs.  $2(49) = 98$ .

Time = 0.28 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.89

method	result	size
derivativedivides	$\frac{-2 \operatorname{arcsech}(xa) + \frac{\operatorname{arcsech}(xa) \left( \operatorname{arcsech}(xa) x^2 a^2 - 2 \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} xa + 2 \right)}{2}}{a^2} + \ln \left( 1 + \left( \frac{1}{ax} + \sqrt{\frac{1}{ax} - 1} \sqrt{1 + \frac{1}{ax}} \right)^2 \right)$	100
default	$\frac{-2 \operatorname{arcsech}(xa) + \frac{\operatorname{arcsech}(xa) \left( \operatorname{arcsech}(xa) x^2 a^2 - 2 \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} xa + 2 \right)}{2}}{a^2} + \ln \left( 1 + \left( \frac{1}{ax} + \sqrt{\frac{1}{ax} - 1} \sqrt{1 + \frac{1}{ax}} \right)^2 \right)$	100

input

```
int(x*arcsech(x*a)^2,x,method=_RETURNVERBOSE)
```

output

```
1/a^2*(-2*arcsech(x*a)+1/2*arcsech(x*a)*(arcsech(x*a)*x^2*a^2-2*(-(a*x-1)/
x/a)^(1/2)*((a*x+1)/x/a)^(1/2)*x*a+2)+ln(1+(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x
)^(1/2))^2))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 106 vs.  $2(49) = 98$ .

Time = 0.10 (sec) , antiderivative size = 106, normalized size of antiderivative = 2.00

$$\int x \operatorname{sech}^{-1}(ax)^2 dx$$

$$= \frac{a^2 x^2 \log \left( \frac{ax \sqrt{-\frac{a^2 x^2 - 1}{a^2 x^2}} + 1}{ax} \right) - 2 ax \sqrt{-\frac{a^2 x^2 - 1}{a^2 x^2}} \log \left( \frac{ax \sqrt{-\frac{a^2 x^2 - 1}{a^2 x^2}} + 1}{ax} \right) - 2 \log(x)}{2 a^2}$$

input

```
integrate(x*arcsech(a*x)^2,x, algorithm="fricas")
```

output

```
1/2*(a^2*x^2*log((a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)) + 1)/(a*x))^2 - 2*a*x
*sqrt(-(a^2*x^2 - 1)/(a^2*x^2))*log((a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)) +
1)/(a*x)) - 2*log(x))/a^2
```

**Sympy [A] (verification not implemented)**

Time = 0.36 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.79

$$\int x \operatorname{sech}^{-1}(ax)^2 dx = \begin{cases} \frac{x^2 \operatorname{asech}^2(ax)}{2} - \frac{\sqrt{-a^2x^2+1} \operatorname{asech}(ax)}{a^2} - \frac{\log(x)}{a^2} & \text{for } a \neq 0 \\ \infty x^2 & \text{otherwise} \end{cases}$$

input

```
integrate(x*asech(a*x)**2,x)
```

output

```
Piecewise((x**2*asech(a*x)**2/2 - sqrt(-a**2*x**2 + 1)*asech(a*x)/a**2 - 1
og(x)/a**2, Ne(a, 0)), (oo*x**2, True))
```

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.75

$$\int x \operatorname{sech}^{-1}(ax)^2 dx = \frac{1}{2} x^2 \operatorname{arsech}(ax)^2 - \frac{x \sqrt{\frac{1}{a^2x^2} - 1} \operatorname{arsech}(ax)}{a} - \frac{\log(x)}{a^2}$$

input

```
integrate(x*arcsech(a*x)^2,x, algorithm="maxima")
```

output

```
1/2*x^2*arcsech(a*x)^2 - x*sqrt(1/(a^2*x^2) - 1)*arcsech(a*x)/a - log(x)/a
^2
```

**Giac [F]**

$$\int x \operatorname{sech}^{-1}(ax)^2 dx = \int x \operatorname{arsech}(ax)^2 dx$$

input `integrate(x*arcsech(a*x)^2,x, algorithm="giac")`

output `integrate(x*arcsech(a*x)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{sech}^{-1}(ax)^2 dx = \int x \operatorname{acosh}\left(\frac{1}{ax}\right)^2 dx$$

input `int(x*acosh(1/(a*x))^2,x)`

output `int(x*acosh(1/(a*x))^2, x)`

**Reduce [F]**

$$\int x \operatorname{sech}^{-1}(ax)^2 dx = \int \operatorname{asech}(ax)^2 x dx$$

input `int(x*asech(a*x)^2,x)`

output `int(asech(a*x)**2*x,x)`

### 3.5 $\int \operatorname{sech}^{-1}(ax)^2 dx$

Optimal result	124
Mathematica [A] (verified)	124
Rubi [A] (verified)	125
Maple [A] (verified)	127
Fricas [F]	127
Sympy [F]	128
Maxima [F]	128
Giac [F]	128
Mupad [F(-1)]	129
Reduce [F]	129

#### Optimal result

Integrand size = 6, antiderivative size = 63

$$\int \operatorname{sech}^{-1}(ax)^2 dx = x \operatorname{sech}^{-1}(ax)^2 - \frac{4 \operatorname{sech}^{-1}(ax) \arctan\left(e^{\operatorname{sech}^{-1}(ax)}\right)}{a} + \frac{2i \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right)}{a} - \frac{2i \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(ax)}\right)}{a}$$

output

```
x*arcsech(a*x)^2-4*arcsech(a*x)*arctan(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2))/a+2*I*polylog(2,-I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a-2*I*polylog(2,I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a
```

#### Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.43

$$\int \operatorname{sech}^{-1}(ax)^2 dx = \frac{i \left( \operatorname{sech}^{-1}(ax) \left( -iax \operatorname{sech}^{-1}(ax) + 2 \log \left( 1 - ie^{-\operatorname{sech}^{-1}(ax)} \right) - 2 \log \left( 1 + ie^{-\operatorname{sech}^{-1}(ax)} \right) \right) + 2 \operatorname{PolyLog} \left( 2, -ie^{-\operatorname{sech}^{-1}(ax)} \right) - 2 \operatorname{PolyLog} \left( 2, ie^{-\operatorname{sech}^{-1}(ax)} \right) \right)}{a}$$

input

```
Integrate[ArcSech[a*x]^2,x]
```

output

```
(I*(ArcSech[a*x]*((-I)*a*x*ArcSech[a*x] + 2*Log[1 - I/E^ArcSech[a*x]] - 2*
Log[1 + I/E^ArcSech[a*x]]) + 2*PolyLog[2, (-I)/E^ArcSech[a*x]] - 2*PolyLog
[2, I/E^ArcSech[a*x]]))/a
```

**Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.02, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {6833, 5941, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{sech}^{-1}(ax)^2 dx$$

$$\downarrow 6833$$

$$\frac{\int ax \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 d\operatorname{sech}^{-1}(ax)}{a}$$

$$\downarrow 5941$$

$$\frac{2 \int ax \operatorname{sech}^{-1}(ax) d\operatorname{sech}^{-1}(ax) - ax \operatorname{sech}^{-1}(ax)^2}{a}$$

$$\downarrow 3042$$

$$\frac{-ax \operatorname{sech}^{-1}(ax)^2 + 2 \int \operatorname{sech}^{-1}(ax) \csc\left(i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2}\right) d\operatorname{sech}^{-1}(ax)}{a}$$

$$\downarrow 4668$$

$$\frac{-ax \operatorname{sech}^{-1}(ax)^2 + 2 \left( -i \int \log\left(1 - ie^{\operatorname{sech}^{-1}(ax)}\right) d\operatorname{sech}^{-1}(ax) + i \int \log\left(1 + ie^{\operatorname{sech}^{-1}(ax)}\right) d\operatorname{sech}^{-1}(ax) + 2 \operatorname{sech}^{-1}(ax) \right)}{a}$$

$$\downarrow 2715$$

$$\frac{-ax \operatorname{sech}^{-1}(ax)^2 + 2 \left( -i \int e^{-\operatorname{sech}^{-1}(ax)} \log\left(1 - ie^{\operatorname{sech}^{-1}(ax)}\right) de^{\operatorname{sech}^{-1}(ax)} + i \int e^{-\operatorname{sech}^{-1}(ax)} \log\left(1 + ie^{\operatorname{sech}^{-1}(ax)}\right) de^{\operatorname{sech}^{-1}(ax)} \right)}{a}$$

$$\downarrow 2838$$

$$\frac{-ax \operatorname{sech}^{-1}(ax)^2 + 2 \left( 2 \operatorname{sech}^{-1}(ax) \arctan \left( e^{\operatorname{sech}^{-1}(ax)} \right) - i \operatorname{PolyLog} \left( 2, -i e^{\operatorname{sech}^{-1}(ax)} \right) + i \operatorname{PolyLog} \left( 2, i e^{\operatorname{sech}^{-1}(ax)} \right) \right)}{a}$$

input `Int[ArcSech[a*x]^2,x]`

output `-((- (a*x*ArcSech[a*x]^2) + 2*(2*ArcSech[a*x]*ArcTan[E^ArcSech[a*x]] - I*PolyLog[2, (-I)*E^ArcSech[a*x]] + I*PolyLog[2, I*E^ArcSech[a*x]]))/a)`

### Defintions of rubi rules used

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] :> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 5941 `Int[(x_)^(m_)*Sech[(a_) + (b_)*(x_)^(n_)]^(p_)*Tanh[(a_) + (b_)*(x_)^(n_)]^(q_), x_Symbol] :> Simp[(-x^(m - n + 1))*(Sech[a + b*x^n]^p/(b*n*p)), x] + Simp[(m - n + 1)/(b*n*p) Int[x^(m - n)*Sech[a + b*x^n]^p, x], x] /; FreeQ[{a, b, p}, x] && RationalQ[m] && IntegerQ[n] && GeQ[m - n, 0] && EqQ[q, 1]`

rule 6833

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Simp[-c^(-1) S
ubst[Int[(a + b*x)^n*Sech[x]*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a,
b, c, n}, x] && IGtQ[n, 0]
```

**Maple [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 183, normalized size of antiderivative = 2.90

method	result
derivativedivides	$\frac{\operatorname{arcsech}(xa)^2 xa + 2i \operatorname{arcsech}(xa) \ln\left(1+i\left(\frac{1}{ax} + \sqrt{\frac{1}{ax}-1}\sqrt{1+\frac{1}{ax}}\right)\right) - 2i \operatorname{arcsech}(xa) \ln\left(1-i\left(\frac{1}{ax} + \sqrt{\frac{1}{ax}-1}\sqrt{1+\frac{1}{ax}}\right)\right)}{a}$
default	$\frac{\operatorname{arcsech}(xa)^2 xa + 2i \operatorname{arcsech}(xa) \ln\left(1+i\left(\frac{1}{ax} + \sqrt{\frac{1}{ax}-1}\sqrt{1+\frac{1}{ax}}\right)\right) - 2i \operatorname{arcsech}(xa) \ln\left(1-i\left(\frac{1}{ax} + \sqrt{\frac{1}{ax}-1}\sqrt{1+\frac{1}{ax}}\right)\right)}{a}$

input

```
int(arcsech(x*a)^2,x,method=_RETURNVERBOSE)
```

output

```
1/a*(arcsech(x*a)^2*x*a+2*I*arcsech(x*a)*ln(1+I*(1/a/x+(1/a/x-1)^(1/2))*(1+
1/a/x)^(1/2)))-2*I*arcsech(x*a)*ln(1-I*(1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1
/2)))+2*I*dilog(1+I*(1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2)))-2*I*dilog(1-I
*(1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2)))
```

**Fricas [F]**

$$\int \operatorname{sech}^{-1}(ax)^2 dx = \int \operatorname{arsech}(ax)^2 dx$$

input

```
integrate(arcsech(a*x)^2,x, algorithm="fricas")
```

output

```
integral(arcsech(a*x)^2, x)
```



**Sympy [F]**

$$\int \operatorname{sech}^{-1}(ax)^2 dx = \int \operatorname{arsech}^2(ax) dx$$

input `integrate(asech(a*x)**2,x)`

output `Integral(asech(a*x)**2, x)`

**Maxima [F]**

$$\int \operatorname{sech}^{-1}(ax)^2 dx = \int \operatorname{arsech}(ax)^2 dx$$

input `integrate(arcsech(a*x)^2,x, algorithm="maxima")`

output `x*log(sqrt(a*x + 1)*sqrt(-a*x + 1) + 1)^2 - integrate(-(a^2*x^2*log(a)^2 + (a^2*x^2 - 1)*log(x)^2 + (a^2*x^2*log(a)^2 + (a^2*x^2 - 1)*log(x)^2 - log(a)^2 + 2*(a^2*x^2*log(a) - log(a))*log(x))*sqrt(a*x + 1)*sqrt(-a*x + 1) - 2*(a^2*x^2*log(a) + (a^2*x^2*(log(a) + 1) + (a^2*x^2 - 1)*log(x) - log(a)))*sqrt(a*x + 1)*sqrt(-a*x + 1) + (a^2*x^2 - 1)*log(x) - log(a))*sqrt(a*x + 1)*sqrt(-a*x + 1) + (a^2*x^2 - 1)*log(x) - log(a))*log(sqrt(a*x + 1)*sqrt(-a*x + 1) + 1) - log(a)^2 + 2*(a^2*x^2*log(a) - log(a))*log(x))/(a^2*x^2 + (a^2*x^2 - 1)*sqrt(a*x + 1)*sqrt(-a*x + 1) - 1), x)`

**Giac [F]**

$$\int \operatorname{sech}^{-1}(ax)^2 dx = \int \operatorname{arsech}(ax)^2 dx$$

input `integrate(arcsech(a*x)^2,x, algorithm="giac")`

output `integrate(arcsech(a*x)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{sech}^{-1}(ax)^2 dx = \int \operatorname{acosh}\left(\frac{1}{ax}\right)^2 dx$$

input `int(acosh(1/(a*x))^2,x)`output `int(acosh(1/(a*x))^2, x)`**Reduce [F]**

$$\int \operatorname{sech}^{-1}(ax)^2 dx = \int \operatorname{asech}(ax)^2 dx$$

input `int(asech(a*x)^2,x)`output `int(asech(a*x)**2,x)`

### 3.6 $\int \frac{\operatorname{sech}^{-1}(ax)^2}{x} dx$

Optimal result	130
Mathematica [A] (verified)	130
Rubi [C] (verified)	131
Maple [A] (verified)	134
Fricas [F]	134
Sympy [F]	134
Maxima [F]	135
Giac [F]	135
Mupad [F(-1)]	135
Reduce [F]	136

#### Optimal result

Integrand size = 10, antiderivative size = 64

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x} dx = \frac{1}{3} \operatorname{sech}^{-1}(ax)^3 - \operatorname{sech}^{-1}(ax)^2 \log\left(1 + e^{2\operatorname{sech}^{-1}(ax)}\right) - \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(ax)}\right) + \frac{1}{2} \operatorname{PolyLog}\left(3, -e^{2\operatorname{sech}^{-1}(ax)}\right)$$

output `1/3*arcsech(a*x)^3-arcsech(a*x)^2*ln(1+(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2))^2)-arcsech(a*x)*polylog(2,-(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2))^2)+1/2*polylog(3,-(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2))^2)`

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.98

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x} dx = -\frac{1}{3} \operatorname{sech}^{-1}(ax)^3 - \operatorname{sech}^{-1}(ax)^2 \log\left(1 + e^{-2\operatorname{sech}^{-1}(ax)}\right) + \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(2, -e^{-2\operatorname{sech}^{-1}(ax)}\right) + \frac{1}{2} \operatorname{PolyLog}\left(3, -e^{-2\operatorname{sech}^{-1}(ax)}\right)$$

input `Integrate[ArcSech[a*x]^2/x,x]`

output `-1/3*ArcSech[a*x]^3 - ArcSech[a*x]^2*Log[1 + E^(-2*ArcSech[a*x])] + ArcSech[a*x]*PolyLog[2, -E^(-2*ArcSech[a*x])] + PolyLog[3, -E^(-2*ArcSech[a*x])]/2`

### Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.47 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.23, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$ , Rules used = {6839, 3042, 26, 4201, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^{-1}(ax)^2}{x} dx \\
 & \quad \downarrow \text{6839} \\
 & - \int \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 d\operatorname{sech}^{-1}(ax) \\
 & \quad \downarrow \text{3042} \\
 & - \int -i \operatorname{sech}^{-1}(ax)^2 \tan(i \operatorname{sech}^{-1}(ax)) d\operatorname{sech}^{-1}(ax) \\
 & \quad \downarrow \text{26} \\
 & i \int \operatorname{sech}^{-1}(ax)^2 \tan(i \operatorname{sech}^{-1}(ax)) d\operatorname{sech}^{-1}(ax) \\
 & \quad \downarrow \text{4201} \\
 & i \left( 2i \int \frac{e^{2\operatorname{sech}^{-1}(ax)} \operatorname{sech}^{-1}(ax)^2}{1 + e^{2\operatorname{sech}^{-1}(ax)}} d\operatorname{sech}^{-1}(ax) - \frac{1}{3} i \operatorname{sech}^{-1}(ax)^3 \right) \\
 & \quad \downarrow \text{2620}
 \end{aligned}$$

$$i \left( 2i \left( \frac{1}{2} \operatorname{sech}^{-1}(ax)^2 \log \left( e^{2\operatorname{sech}^{-1}(ax)} + 1 \right) - \int \operatorname{sech}^{-1}(ax) \log \left( 1 + e^{2\operatorname{sech}^{-1}(ax)} \right) d\operatorname{sech}^{-1}(ax) \right) - \frac{1}{3} i \operatorname{sech}^{-1}(ax)^3 \right)$$

↓ 3011

$$i \left( 2i \left( -\frac{1}{2} \int \operatorname{PolyLog} \left( 2, -e^{2\operatorname{sech}^{-1}(ax)} \right) d\operatorname{sech}^{-1}(ax) + \frac{1}{2} \operatorname{sech}^{-1}(ax) \operatorname{PolyLog} \left( 2, -e^{2\operatorname{sech}^{-1}(ax)} \right) + \frac{1}{2} \operatorname{sech}^{-1}(ax)^2 \right) \right)$$

↓ 2720

$$i \left( 2i \left( -\frac{1}{4} \int e^{-2\operatorname{sech}^{-1}(ax)} \operatorname{PolyLog} \left( 2, -e^{2\operatorname{sech}^{-1}(ax)} \right) de^{2\operatorname{sech}^{-1}(ax)} + \frac{1}{2} \operatorname{sech}^{-1}(ax) \operatorname{PolyLog} \left( 2, -e^{2\operatorname{sech}^{-1}(ax)} \right) + \frac{1}{2} \right) \right)$$

↓ 7143

$$i \left( 2i \left( \frac{1}{2} \operatorname{sech}^{-1}(ax) \operatorname{PolyLog} \left( 2, -e^{2\operatorname{sech}^{-1}(ax)} \right) - \frac{1}{4} \operatorname{PolyLog} \left( 3, -e^{2\operatorname{sech}^{-1}(ax)} \right) + \frac{1}{2} \operatorname{sech}^{-1}(ax)^2 \log \left( e^{2\operatorname{sech}^{-1}(ax)} \right) \right) \right)$$

input `Int[ArcSech[a*x]^2/x, x]`

output `I*((-1/3*I)*ArcSech[a*x]^3 + (2*I)*((ArcSech[a*x]^2*Log[1 + E^(2*ArcSech[a*x])]))/2 + (ArcSech[a*x]*PolyLog[2, -E^(2*ArcSech[a*x])])/2 - PolyLog[3, -E^(2*ArcSech[a*x])]/4)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4201 `Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_]*(f_)*(x_)]], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x))), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6839 `Int[((a_) + ArcSech[(c_)*(x_)]*(b_))^(n_)*(x_)^m_, x_Symbol] := Simp[-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])`

rule 7143 `Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

**Maple [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 136, normalized size of antiderivative = 2.12

method	result
derivativedivides	$\frac{\operatorname{arcsech}(xa)^3}{3} - \operatorname{arcsech}(xa)^2 \ln \left( 1 + \left( \frac{1}{ax} + \sqrt{\frac{1}{ax} - 1} \sqrt{1 + \frac{1}{ax}} \right)^2 \right) - \operatorname{arcsech}(xa) \operatorname{polylog}$
default	$\frac{\operatorname{arcsech}(xa)^3}{3} - \operatorname{arcsech}(xa)^2 \ln \left( 1 + \left( \frac{1}{ax} + \sqrt{\frac{1}{ax} - 1} \sqrt{1 + \frac{1}{ax}} \right)^2 \right) - \operatorname{arcsech}(xa) \operatorname{polylog}$

input `int(arcsech(x*a)^2/x,x,method=_RETURNVERBOSE)`

output `1/3*arcsech(x*a)^3-arcsech(x*a)^2*ln(1+(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2)-arcsech(x*a)*polylog(2,-(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2)+1/2*polylog(3,-(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2)`

**Fricas [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x} dx = \int \frac{\operatorname{arsech}(ax)^2}{x} dx$$

input `integrate(arcsech(a*x)^2/x,x, algorithm="fricas")`

output `integral(arcsech(a*x)^2/x, x)`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x} dx = \int \frac{\operatorname{asech}^2(ax)}{x} dx$$

input `integrate(asech(a*x)**2/x,x)`

output `Integral(asech(a*x)**2/x, x)`

### Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x} dx = \int \frac{\operatorname{arosech}(ax)^2}{x} dx$$

input `integrate(arcsech(a*x)^2/x,x, algorithm="maxima")`

output `integrate(arcsech(a*x)^2/x, x)`

### Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x} dx = \int \frac{\operatorname{arosech}(ax)^2}{x} dx$$

input `integrate(arcsech(a*x)^2/x,x, algorithm="giac")`

output `integrate(arcsech(a*x)^2/x, x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{ax}\right)^2}{x} dx$$

input `int(acosh(1/(a*x))^2/x,x)`

output `int(acosh(1/(a*x))^2/x, x)`



**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x} dx = \int \frac{a \operatorname{sech}(ax)^2}{x} dx$$

input `int(asech(a*x)^2/x,x)`

output `int(asech(a*x)**2/x,x)`

### 3.7 $\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^2} dx$

Optimal result	137
Mathematica [A] (verified)	137
Rubi [C] (verified)	138
Maple [A] (verified)	140
Fricas [B] (verification not implemented)	140
Sympy [F]	141
Maxima [A] (verification not implemented)	141
Giac [F]	141
Mupad [F(-1)]	142
Reduce [F]	142

#### Optimal result

Integrand size = 10, antiderivative size = 49

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^2} dx = -\frac{2}{x} + \frac{2\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)}{x} - \frac{\operatorname{sech}^{-1}(ax)^2}{x}$$

output

$$-2/x + 2*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*\operatorname{arcsech}(a*x)/x - \operatorname{arcsech}(a*x)^2/x$$

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.86

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^2} dx = -\frac{2 - 2\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax) + \operatorname{sech}^{-1}(ax)^2}{x}$$

input

$$\operatorname{Integrate}[\operatorname{ArcSech}[a*x]^2/x^2, x]$$

output

$$-((2 - 2*\operatorname{Sqrt}[(1 - a*x)/(1 + a*x)]*(1 + a*x)*\operatorname{ArcSech}[a*x] + \operatorname{ArcSech}[a*x]^2)/x)$$

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.41, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {6839, 3042, 26, 3777, 3042, 3777, 26, 3042, 26, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^{-1}(ax)^2}{x^2} dx \\
 & \quad \downarrow \text{6839} \\
 & -a \int \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{ax} d\operatorname{sech}^{-1}(ax) \\
 & \quad \downarrow \text{3042} \\
 & -a \int -i\operatorname{sech}^{-1}(ax)^2 \sin(i\operatorname{sech}^{-1}(ax)) d\operatorname{sech}^{-1}(ax) \\
 & \quad \downarrow \text{26} \\
 & ia \int \operatorname{sech}^{-1}(ax)^2 \sin(i\operatorname{sech}^{-1}(ax)) d\operatorname{sech}^{-1}(ax) \\
 & \quad \downarrow \text{3777} \\
 & ia \left( \frac{i\operatorname{sech}^{-1}(ax)^2}{ax} - 2i \int \frac{\operatorname{sech}^{-1}(ax)}{ax} d\operatorname{sech}^{-1}(ax) \right) \\
 & \quad \downarrow \text{3042} \\
 & ia \left( \frac{i\operatorname{sech}^{-1}(ax)^2}{ax} - 2i \int \operatorname{sech}^{-1}(ax) \sin\left(i\operatorname{sech}^{-1}(ax) + \frac{\pi}{2}\right) d\operatorname{sech}^{-1}(ax) \right) \\
 & \quad \downarrow \text{3777} \\
 & ia \left( \frac{i\operatorname{sech}^{-1}(ax)^2}{ax} - 2i \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{ax} - i \int -\frac{i\sqrt{\frac{1-ax}{ax+1}}(ax+1)}{ax} d\operatorname{sech}^{-1}(ax) \right) \right) \\
 & \quad \downarrow \text{26}
 \end{aligned}$$

$$\begin{aligned}
& ia \left( \frac{\operatorname{isech}^{-1}(ax)^2}{ax} - 2i \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{ax} - \int \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)}{ax} d\operatorname{sech}^{-1}(ax) \right) \right) \\
& \quad \downarrow \text{3042} \\
& ia \left( \frac{\operatorname{isech}^{-1}(ax)^2}{ax} - 2i \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{ax} - \int -i \sin(\operatorname{isech}^{-1}(ax)) d\operatorname{sech}^{-1}(ax) \right) \right) \\
& \quad \downarrow \text{26} \\
& ia \left( \frac{\operatorname{isech}^{-1}(ax)^2}{ax} - 2i \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{ax} + i \int \sin(\operatorname{isech}^{-1}(ax)) d\operatorname{sech}^{-1}(ax) \right) \right) \\
& \quad \downarrow \text{3118} \\
& ia \left( \frac{\operatorname{isech}^{-1}(ax)^2}{ax} - 2i \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{ax} - \frac{1}{ax} \right) \right)
\end{aligned}$$

input `Int[ArcSech[a*x]^2/x^2,x]`

output `I*a*((I*ArcSech[a*x]^2)/(a*x) - (2*I)*(-1/(a*x)) + (Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x])/(a*x))`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_)^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-c + d*x)^m*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 6839 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))^(n_)*(x_)^(m_.), x_Symbol] := Simp[-(c^(m + 1))^(1/2) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])`

### Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.24

method	result	size
derivativedivides	$a \left( -\frac{\operatorname{arcsech}(xa)^2}{xa} + 2\sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} \operatorname{arcsech}(xa) - \frac{2}{ax} \right)$	61
default	$a \left( -\frac{\operatorname{arcsech}(xa)^2}{xa} + 2\sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} \operatorname{arcsech}(xa) - \frac{2}{ax} \right)$	61

input `int(arcsech(x*a)^2/x^2,x,method=_RETURNVERBOSE)`

output `a*(-1/x/a*arcsech(x*a)^2+2*(-(a*x-1)/x/a)^(1/2)*((a*x+1)/x/a)^(1/2)*arcsech(x*a)-2/a/x)`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 97 vs. 2(47) = 94.

Time = 0.09 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.98

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^2} dx = \frac{2ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}} \log\left(\frac{ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}}+1}{ax}\right) - \log\left(\frac{ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}}+1}{ax}\right)^2}{x} - 2$$

input `integrate(arcsech(a*x)^2/x^2,x, algorithm="fricas")`

output  $(2ax\sqrt{-(a^2x^2 - 1)/(a^2x^2)})\log((ax\sqrt{-(a^2x^2 - 1)/(a^2x^2)} + 1)/(ax)) - \log((ax\sqrt{-(a^2x^2 - 1)/(a^2x^2)} + 1)/(ax))^2 - 2)/x$

### Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^2} dx = \int \frac{\operatorname{arsech}^2(ax)}{x^2} dx$$

input `integrate(asech(a*x)**2/x**2,x)`

output `Integral(asech(a*x)**2/x**2, x)`

### Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.71

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^2} dx = 2a\sqrt{\frac{1}{a^2x^2} - 1} \operatorname{arsech}(ax) - \frac{\operatorname{arsech}(ax)^2}{x} - \frac{2}{x}$$

input `integrate(arcsech(a*x)^2/x^2,x, algorithm="maxima")`

output `2*a*sqrt(1/(a^2*x^2) - 1)*arcsech(a*x) - arcsech(a*x)^2/x - 2/x`

### Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^2} dx = \int \frac{\operatorname{arsech}(ax)^2}{x^2} dx$$

input `integrate(arcsech(a*x)^2/x^2,x, algorithm="giac")`

output `integrate(arcsech(a*x)^2/x^2, x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^2} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{ax}\right)^2}{x^2} dx$$

input `int(acosh(1/(a*x))^2/x^2,x)`

output `int(acosh(1/(a*x))^2/x^2, x)`

### Reduce [F]

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^2} dx = \int \frac{\operatorname{asech}(ax)^2}{x^2} dx$$

input `int(asech(a*x)^2/x^2,x)`

output `int(asech(a*x)**2/x**2,x)`

### 3.8 $\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^3} dx$

Optimal result	143
Mathematica [A] (verified)	143
Rubi [A] (verified)	144
Maple [A] (verified)	146
Fricas [A] (verification not implemented)	146
Sympy [F]	147
Maxima [F]	147
Giac [F]	147
Mupad [F(-1)]	148
Reduce [F]	148

#### Optimal result

Integrand size = 10, antiderivative size = 90

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^3} dx = -\frac{(1-ax)(1+ax)}{4x^2} + \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)}{2x^2} - \frac{1}{4}a^2\operatorname{sech}^{-1}(ax)^2 - \frac{(1-ax)(1+ax)\operatorname{sech}^{-1}(ax)^2}{2x^2}$$

output

```
-1/4*(-a*x+1)*(a*x+1)/x^2+1/2*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*arcsech(a*x)/x^2-1/4*a^2*arcsech(a*x)^2-1/2*(-a*x+1)*(a*x+1)*arcsech(a*x)^2/x^2
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.60

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^3} dx = \frac{-1 + 2\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax) + (-2 + a^2x^2)\operatorname{sech}^{-1}(ax)^2}{4x^2}$$

input

```
Integrate[ArcSech[a*x]^2/x^3,x]
```



output

```
(-1 + 2*sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x] + (-2 + a^2*x^2)*
ArcSech[a*x]^2)/(4*x^2)
```

**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {6839, 5895, 3042, 25, 3791, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^3} dx$$

$$\downarrow 6839$$

$$-a^2 \int \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{a^2x^2} d\operatorname{sech}^{-1}(ax)$$

$$\downarrow 5895$$

$$-a^2 \left( \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)^2}{2a^2x^2} - \int \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)}{a^2x^2} d\operatorname{sech}^{-1}(ax) \right)$$

$$\downarrow 3042$$

$$-a^2 \left( \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)^2}{2a^2x^2} - \int -\operatorname{sech}^{-1}(ax) \sin(i\operatorname{sech}^{-1}(ax))^2 d\operatorname{sech}^{-1}(ax) \right)$$

$$\downarrow 25$$

$$-a^2 \left( \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)^2}{2a^2x^2} + \int \operatorname{sech}^{-1}(ax) \sin(i\operatorname{sech}^{-1}(ax))^2 d\operatorname{sech}^{-1}(ax) \right)$$

$$\downarrow 3791$$

$$-a^2 \left( \frac{1}{2} \int \operatorname{sech}^{-1}(ax) d\operatorname{sech}^{-1}(ax) + \frac{(1-ax)(ax+1)}{4a^2x^2} + \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)^2}{2a^2x^2} - \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{2a^2x^2} \right)$$

$$\downarrow 15$$

$$-a^2 \left( \frac{(1-ax)(ax+1)}{4a^2x^2} + \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)^2}{2a^2x^2} - \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{2a^2x^2} + \frac{1}{4}\operatorname{sech}^{-1}(ax)^2 \right)$$

input `Int[ArcSech[a*x]^2/x^3,x]`

output `-(a^2*(((1 - a*x)*(1 + a*x))/(4*a^2*x^2) - (Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x])/(2*a^2*x^2) + ArcSech[a*x]^2/4 + ((1 - a*x)*(1 + a*x)*ArcSech[a*x]^2)/(2*a^2*x^2)))`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3791 `Int[((c_.) + (d_.)*(x_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[d*((b*Sine[e + f*x])^n/(f^2*n^2)), x] + (-Simp[b*(c + d*x)*Cos[e + f*x]*((b*Sine[e + f*x])^(n - 1)/(f*n)), x] + Simp[b^2*((n - 1)/n) Int[(c + d*x)*(b*Sine[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1]`

rule 5895 `Int[Cosh[(a_.) + (b_.)*(x_)^(n_.)]*(x_)^(m_.)*Sinh[(a_.) + (b_.)*(x_)^(n_.)]^(p_.), x_Symbol] := Simp[x^(m - n + 1)*(Sinh[a + b*x^n]^(p + 1)/(b*n*(p + 1))), x] - Simp[(m - n + 1)/(b*n*(p + 1)) Int[x^(m - n)*Sinh[a + b*x^n]^(p + 1), x], x] /; FreeQ[{a, b, p}, x] && LtQ[0, n, m + 1] && NeQ[p, -1]`

rule 6839

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_.), x_Symbol] :> Simp[
-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

**Maple [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.48

method	result	size
derivativedivides	$a^2 \left( -\frac{\cosh(2 \operatorname{arcsech}(xa)) \operatorname{arcsech}(xa)^2}{4} + \frac{\sinh(2 \operatorname{arcsech}(xa)) \operatorname{arcsech}(xa)}{4} - \frac{\cosh(2 \operatorname{arcsech}(xa))}{8} \right)$	43
default	$a^2 \left( -\frac{\cosh(2 \operatorname{arcsech}(xa)) \operatorname{arcsech}(xa)^2}{4} + \frac{\sinh(2 \operatorname{arcsech}(xa)) \operatorname{arcsech}(xa)}{4} - \frac{\cosh(2 \operatorname{arcsech}(xa))}{8} \right)$	43

input

```
int(arcsech(x*a)^2/x^3,x,method=_RETURNVERBOSE)
```

output

```
a^2*(-1/4*cosh(2*arcsech(x*a))*arcsech(x*a)^2+1/4*sinh(2*arcsech(x*a))*arc
sech(x*a)-1/8*cosh(2*arcsech(x*a)))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.18

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^3} dx$$

$$= \frac{2ax \sqrt{-\frac{a^2x^2-1}{a^2x^2}} \log\left(\frac{ax \sqrt{-\frac{a^2x^2-1}{a^2x^2}}+1}{ax}\right) + (a^2x^2-2) \log\left(\frac{ax \sqrt{-\frac{a^2x^2-1}{a^2x^2}}+1}{ax}\right)^2 - 1}{4x^2}$$

input

```
integrate(arcsech(a*x)^2/x^3,x, algorithm="fricas")
```

output

```
1/4*(2*a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2))*log((a*x*sqrt(-(a^2*x^2 - 1)/(a^
2*x^2)) + 1)/(a*x)) + (a^2*x^2 - 2)*log((a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2
) + 1)/(a*x))^2 - 1)/x^2
```

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^3} dx = \int \frac{\operatorname{arsech}^2(ax)}{x^3} dx$$

input `integrate(asech(a*x)**2/x**3,x)`

output `Integral(asech(a*x)**2/x**3, x)`

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^3} dx = \int \frac{\operatorname{arsech}(ax)^2}{x^3} dx$$

input `integrate(arcsech(a*x)^2/x^3,x, algorithm="maxima")`

output `integrate(arcsech(a*x)^2/x^3, x)`

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^3} dx = \int \frac{\operatorname{arsech}(ax)^2}{x^3} dx$$

input `integrate(arcsech(a*x)^2/x^3,x, algorithm="giac")`

output `integrate(arcsech(a*x)^2/x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^3} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{ax}\right)^2}{x^3} dx$$

input `int(acosh(1/(a*x))^2/x^3,x)`output `int(acosh(1/(a*x))^2/x^3, x)`**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^3} dx = \int \frac{\operatorname{asech}(ax)^2}{x^3} dx$$

input `int(asech(a*x)^2/x^3,x)`output `int(asech(a*x)**2/x**3,x)`

### 3.9 $\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^4} dx$

Optimal result	149
Mathematica [A] (verified)	149
Rubi [A] (verified)	150
Maple [A] (verified)	153
Fricas [A] (verification not implemented)	153
Sympy [F]	154
Maxima [F]	154
Giac [F]	154
Mupad [F(-1)]	155
Reduce [F]	155

#### Optimal result

Integrand size = 10, antiderivative size = 102

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^4} dx = -\frac{2}{27x^3} - \frac{4a^2}{9x} + \frac{2\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)}{9x^3} + \frac{4a^2\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)}{9x} - \frac{\operatorname{sech}^{-1}(ax)^2}{3x^3}$$

output

$$-2/27/x^3-4/9*a^2/x+2/9*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*\operatorname{arcsech}(a*x)/x^3+4/9*a^2*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*\operatorname{arcsech}(a*x)/x-1/3*\operatorname{arcsech}(a*x)^2/x^3$$

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.72

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^4} dx = \frac{-2(1+6a^2x^2) + 6\sqrt{\frac{1-ax}{1+ax}}(1+ax+2a^2x^2+2a^3x^3)\operatorname{sech}^{-1}(ax) - 9\operatorname{sech}^{-1}(ax)^2}{27x^3}$$

input `Integrate[ArcSech[a*x]^2/x^4,x]`

output  $(-2*(1 + 6*a^2*x^2) + 6*\text{Sqrt}[(1 - a*x)/(1 + a*x)]*(1 + a*x + 2*a^2*x^2 + 2*a^3*x^3)*\text{ArcSech}[a*x] - 9*\text{ArcSech}[a*x]^2)/(27*x^3)$

### Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.19, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {6839, 5896, 3042, 3791, 3042, 3777, 26, 3042, 26, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\text{sech}^{-1}(ax)^2}{x^4} dx \\
 & \quad \downarrow \text{6839} \\
 & -a^3 \int \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\text{sech}^{-1}(ax)^2}{a^3x^3} d\text{sech}^{-1}(ax) \\
 & \quad \downarrow \text{5896} \\
 & -a^3 \left( \frac{\text{sech}^{-1}(ax)^2}{3a^3x^3} - \frac{2}{3} \int \frac{\text{sech}^{-1}(ax)}{a^3x^3} d\text{sech}^{-1}(ax) \right) \\
 & \quad \downarrow \text{3042} \\
 & -a^3 \left( \frac{\text{sech}^{-1}(ax)^2}{3a^3x^3} - \frac{2}{3} \int \text{sech}^{-1}(ax) \sin \left( i\text{sech}^{-1}(ax) + \frac{\pi}{2} \right)^3 d\text{sech}^{-1}(ax) \right) \\
 & \quad \downarrow \text{3791} \\
 & -a^3 \left( \frac{\text{sech}^{-1}(ax)^2}{3a^3x^3} - \frac{2}{3} \left( \frac{2}{3} \int \frac{\text{sech}^{-1}(ax)}{ax} d\text{sech}^{-1}(ax) - \frac{1}{9a^3x^3} + \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\text{sech}^{-1}(ax)}{3a^3x^3} \right) \right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$-a^3 \left( \frac{\operatorname{sech}^{-1}(ax)^2}{3a^3x^3} - \frac{2}{3} \left( \frac{2}{3} \int \operatorname{sech}^{-1}(ax) \sin \left( i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2} \right) d \operatorname{sech}^{-1}(ax) - \frac{1}{9a^3x^3} + \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{3a^3x^3} \right) \right)$$

↓ 3777

$$-a^3 \left( \frac{\operatorname{sech}^{-1}(ax)^2}{3a^3x^3} - \frac{2}{3} \left( \frac{2}{3} \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{ax} - i \int -\frac{i\sqrt{\frac{1-ax}{ax+1}}(ax+1)}{ax} d \operatorname{sech}^{-1}(ax) \right) - \frac{1}{9a^3x^3} + \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{3a^3x^3} \right) \right)$$

↓ 26

$$-a^3 \left( \frac{\operatorname{sech}^{-1}(ax)^2}{3a^3x^3} - \frac{2}{3} \left( \frac{2}{3} \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{ax} - \int \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)}{ax} d \operatorname{sech}^{-1}(ax) \right) - \frac{1}{9a^3x^3} + \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{3a^3x^3} \right) \right)$$

↓ 3042

$$-a^3 \left( \frac{\operatorname{sech}^{-1}(ax)^2}{3a^3x^3} - \frac{2}{3} \left( \frac{2}{3} \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{ax} - \int -i \sin \left( i \operatorname{sech}^{-1}(ax) \right) d \operatorname{sech}^{-1}(ax) \right) - \frac{1}{9a^3x^3} + \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{3a^3x^3} \right) \right)$$

↓ 26

$$-a^3 \left( \frac{\operatorname{sech}^{-1}(ax)^2}{3a^3x^3} - \frac{2}{3} \left( \frac{2}{3} \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{ax} + i \int \sin \left( i \operatorname{sech}^{-1}(ax) \right) d \operatorname{sech}^{-1}(ax) \right) - \frac{1}{9a^3x^3} + \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{3a^3x^3} \right) \right)$$

↓ 3118

$$-a^3 \left( \frac{\operatorname{sech}^{-1}(ax)^2}{3a^3x^3} - \frac{2}{3} \left( -\frac{1}{9a^3x^3} + \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{3a^3x^3} + \frac{2}{3} \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{ax} - \frac{1}{ax} \right) \right) \right)$$

input `Int[ArcSech[a*x]^2/x^4,x]`

output `-(a^3*(ArcSech[a*x]^2/(3*a^3*x^3) - (2*(-1/9*1/(a^3*x^3) + (Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x]))/(3*a^3*x^3) + (2*(-1/(a*x)) + (Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x])/(a*x)))/3))/3)`



## Definitions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a])*(Fx\_), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3118  $\text{Int}[\sin[(c\_.) + (d\_.)*(x\_)], x\_Symbol] \rightarrow \text{Simp}[-\text{Cos}[c + d*x]/d, x] /; \text{FreeQ}[\{c, d\}, x]$
- rule 3777  $\text{Int}[((c\_.) + (d\_.)*(x\_))^m*\sin[(e\_.) + (f\_.)*(x\_)], x\_Symbol] \rightarrow \text{Simp}[(-c + d*x)^m*(\text{Cos}[e + f*x]/f), x] + \text{Simp}[d*(m/f) \text{Int}[(c + d*x)^{m-1}*\text{Cos}[e + f*x], x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \ \&\& \ \text{GtQ}[m, 0]$
- rule 3791  $\text{Int}[((c\_.) + (d\_.)*(x\_))*((b\_.)*\sin[(e\_.) + (f\_.)*(x\_)])^n, x\_Symbol] \rightarrow \text{Simp}[d*((b*\sin[e + f*x])^n/(f^2*n^2)), x] + (-\text{Simp}[b*(c + d*x)*\text{Cos}[e + f*x]*((b*\sin[e + f*x])^{n-1}/(f*n)), x] + \text{Simp}[b^2*((n-1)/n) \text{Int}[(c + d*x)*(b*\sin[e + f*x])^{n-2}, x], x]) /; \text{FreeQ}[\{b, c, d, e, f\}, x] \ \&\& \ \text{GtQ}[n, 1]$
- rule 5896  $\text{Int}[\text{Cosh}[(a\_.) + (b\_.)*(x\_)]^{(n\_)}]^{(p\_)}*(x\_)^m*\text{Sinh}[(a\_.) + (b\_.)*(x\_)]^{(n\_)}, x\_Symbol] \rightarrow \text{Simp}[x^{(m-n+1)}*(\text{Cosh}[a + b*x^n]^{(p+1)}/(b*n*(p+1))), x] - \text{Simp}[(m-n+1)/(b*n*(p+1)) \text{Int}[x^{(m-n)}*\text{Cosh}[a + b*x^n]^{(p+1)}, x], x] /; \text{FreeQ}[\{a, b, p\}, x] \ \&\& \ \text{LtQ}[0, n, m+1] \ \&\& \ \text{NeQ}[p, -1]$
- rule 6839  $\text{Int}[((a\_.) + \text{ArcSech}[(c\_.)*(x\_)]*(b\_.)^n)*(x\_)^m, x\_Symbol] \rightarrow \text{Simp}[-(c^{(m+1)})^{(-1)} \text{Subst}[\text{Int}[(a + b*x)^n*\text{Sech}[x]^{(m+1)}*\text{Tanh}[x], x], x, \text{ArcSech}[c*x]], x] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ (\text{GtQ}[n, 0] \ || \ \text{LtQ}[m, -1])$

**Maple [A] (verified)**

Time = 0.46 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.10

method	result
derivativedivides	$a^3 \left( -\frac{\operatorname{arcsech}(xa)^2}{3x^3a^3} + \frac{4\sqrt{-\frac{xa-1}{xa}}\sqrt{\frac{xa+1}{xa}}\operatorname{arcsech}(xa)}{9} + \frac{2\sqrt{-\frac{xa-1}{xa}}\sqrt{\frac{xa+1}{xa}}\operatorname{arcsech}(xa)}{9x^2a^2} - \frac{4}{9ax} - \frac{2}{27x^3a^3} \right)$
default	$a^3 \left( -\frac{\operatorname{arcsech}(xa)^2}{3x^3a^3} + \frac{4\sqrt{-\frac{xa-1}{xa}}\sqrt{\frac{xa+1}{xa}}\operatorname{arcsech}(xa)}{9} + \frac{2\sqrt{-\frac{xa-1}{xa}}\sqrt{\frac{xa+1}{xa}}\operatorname{arcsech}(xa)}{9x^2a^2} - \frac{4}{9ax} - \frac{2}{27x^3a^3} \right)$

input `int(arcsech(x*a)^2/x^4,x,method=_RETURNVERBOSE)`

output `a^3*(-1/3/x^3/a^3*arcsech(x*a)^2+4/9*(-(a*x-1)/x/a)^(1/2)*((a*x+1)/x/a)^(1/2)*arcsech(x*a)+2/9*(-(a*x-1)/x/a)^(1/2)*((a*x+1)/x/a)^(1/2)/x^2/a^2*arcsech(x*a)-4/9/a/x-2/27/x^3/a^3)`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.14

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^4} dx = \frac{12a^2x^2 - 6(2a^3x^3 + ax)\sqrt{-\frac{a^2x^2-1}{a^2x^2}} \log\left(\frac{ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}}+1}{ax}\right) + 9 \log\left(\frac{ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}}+1}{ax}\right)^2 + 2}{27x^3}$$

input `integrate(arcsech(a*x)^2/x^4,x, algorithm="fricas")`

output `-1/27*(12*a^2*x^2 - 6*(2*a^3*x^3 + a*x)*sqrt(-(a^2*x^2 - 1)/(a^2*x^2))*log((a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)) + 1)/(a*x)) + 9*log((a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)) + 1)/(a*x))^2 + 2)/x^3`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^4} dx = \int \frac{\operatorname{arsech}^2(ax)}{x^4} dx$$

input `integrate(asech(a*x)**2/x**4,x)`

output `Integral(asech(a*x)**2/x**4, x)`

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^4} dx = \int \frac{\operatorname{arsech}(ax)^2}{x^4} dx$$

input `integrate(arcsech(a*x)^2/x^4,x, algorithm="maxima")`

output `integrate(arcsech(a*x)^2/x^4, x)`

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^4} dx = \int \frac{\operatorname{arsech}(ax)^2}{x^4} dx$$

input `integrate(arcsech(a*x)^2/x^4,x, algorithm="giac")`

output `integrate(arcsech(a*x)^2/x^4, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^4} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{ax}\right)^2}{x^4} dx$$

input `int(acosh(1/(a*x))^2/x^4,x)`output `int(acosh(1/(a*x))^2/x^4, x)`**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^2}{x^4} dx = \int \frac{\operatorname{asech}(ax)^2}{x^4} dx$$

input `int(asech(a*x)^2/x^4,x)`output `int(asech(a*x)**2/x**4,x)`

### 3.10 $\int x^4 \operatorname{sech}^{-1}(ax)^3 dx$

Optimal result	156
Mathematica [A] (verified)	157
Rubi [A] (verified)	158
Maple [F]	162
Fricas [F]	162
Sympy [F]	163
Maxima [F]	163
Giac [F]	163
Mupad [F(-1)]	164
Reduce [F]	164

#### Optimal result

Integrand size = 10, antiderivative size = 297

$$\begin{aligned}
 \int x^4 \operatorname{sech}^{-1}(ax)^3 dx = & \frac{x \sqrt{\frac{1-ax}{1+ax}} (1+ax)}{20a^4} - \frac{9x \operatorname{sech}^{-1}(ax)}{20a^4} \\
 & - \frac{x^3 \operatorname{sech}^{-1}(ax)}{10a^2} - \frac{9x \sqrt{\frac{1-ax}{1+ax}} (1+ax) \operatorname{sech}^{-1}(ax)^2}{40a^4} \\
 & - \frac{3x^3 \sqrt{\frac{1-ax}{1+ax}} (1+ax) \operatorname{sech}^{-1}(ax)^2}{20a^2} + \frac{1}{5} x^5 \operatorname{sech}^{-1}(ax)^3 \\
 & - \frac{9 \operatorname{sech}^{-1}(ax)^2 \arctan\left(e^{\operatorname{sech}^{-1}(ax)}\right)}{20a^5} + \frac{\arctan\left(\frac{\sqrt{\frac{1-ax}{1+ax}} (1+ax)}{ax}\right)}{2a^5} \\
 & + \frac{9i \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right)}{20a^5} \\
 & - \frac{9i \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(ax)}\right)}{20a^5} \\
 & - \frac{9i \operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(ax)}\right)}{20a^5} + \frac{9i \operatorname{PolyLog}\left(3, ie^{\operatorname{sech}^{-1}(ax)}\right)}{20a^5}
 \end{aligned}$$

output

```

1/20*x*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)/a^4-9/20*x*arcsech(a*x)/a^4-1/10*x
^3*arcsech(a*x)/a^2-9/40*x*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*arcsech(a*x)^2
/a^4-3/20*x^3*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*arcsech(a*x)^2/a^2+1/5*x^5*
arcsech(a*x)^3-9/20*arcsech(a*x)^2*arctan(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)
^(1/2))/a^5+1/2*arctan(((a*x+1)/(a*x+1))^(1/2)*(a*x+1)/a/x)/a^5+9/20*I*ar
csech(a*x)*polylog(2,-I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a^5-9/20
*I*arcsech(a*x)*polylog(2,I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a^5-
9/20*I*polylog(3,-I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a^5+9/20*I*p
olylog(3,I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a^5

```

**Mathematica [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 281, normalized size of antiderivative = 0.95

$$\int x^4 \operatorname{sech}^{-1}(ax)^3 dx$$

$$= \frac{2ax \sqrt{\frac{1-ax}{1+ax}}(1+ax) - 18ax \operatorname{sech}^{-1}(ax) - 4a^3 x^3 \operatorname{sech}^{-1}(ax) - 9ax \sqrt{\frac{1-ax}{1+ax}}(1+ax) \operatorname{sech}^{-1}(ax)^2 - 6a^3 x^3 \sqrt{\frac{1-ax}{1+ax}}}{1}$$

input

```
Integrate[x^4*ArcSech[a*x]^3,x]
```

output

```

(2*a*x*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x) - 18*a*x*ArcSech[a*x] - 4*a^3*x
^3*ArcSech[a*x] - 9*a*x*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x]^2
- 6*a^3*x^3*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x]^2 + 8*a^5*x^
5*ArcSech[a*x]^3 + 40*ArcTan[Tanh[ArcSech[a*x]/2]] + (9*I)*ArcSech[a*x]^2*
Log[1 - I/E^ArcSech[a*x]] - (9*I)*ArcSech[a*x]^2*Log[1 + I/E^ArcSech[a*x]]
+ (18*I)*ArcSech[a*x]*PolyLog[2, (-I)/E^ArcSech[a*x]] - (18*I)*ArcSech[a*
x]*PolyLog[2, I/E^ArcSech[a*x]] + (18*I)*PolyLog[3, (-I)/E^ArcSech[a*x]] -
(18*I)*PolyLog[3, I/E^ArcSech[a*x]])/(40*a^5)

```

**Rubi [A] (verified)**

Time = 1.07 (sec) , antiderivative size = 319, normalized size of antiderivative = 1.07, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$ , Rules used = {6839, 5941, 3042, 4674, 3042, 4255, 3042, 4257, 4674, 3042, 4257, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^4 \operatorname{sech}^{-1}(ax)^3 dx \\
 & \quad \downarrow \text{6839} \\
 & \frac{\int a^5 x^5 \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^3 d\operatorname{sech}^{-1}(ax)}{a^5} \\
 & \quad \downarrow \text{5941} \\
 & \frac{\frac{3}{5} \int a^5 x^5 \operatorname{sech}^{-1}(ax)^2 d\operatorname{sech}^{-1}(ax) - \frac{1}{5} a^5 x^5 \operatorname{sech}^{-1}(ax)^3}{a^5} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\frac{1}{5} a^5 x^5 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{5} \int \operatorname{sech}^{-1}(ax)^2 \csc\left(i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2}\right)^5 d\operatorname{sech}^{-1}(ax)}{a^5} \\
 & \quad \downarrow \text{4674} \\
 & \frac{\frac{3}{5} \left( -\frac{1}{6} \int a^3 x^3 d\operatorname{sech}^{-1}(ax) + \frac{3}{4} \int a^3 x^3 \operatorname{sech}^{-1}(ax)^2 d\operatorname{sech}^{-1}(ax) + \frac{1}{4} a^3 x^3 \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 + \frac{1}{6} a^3 x^3 \operatorname{sech}^{-1}(ax) \right)}{a^5} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\frac{1}{5} a^5 x^5 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{5} \left( -\frac{1}{6} \int \csc\left(i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2}\right)^3 d\operatorname{sech}^{-1}(ax) + \frac{3}{4} \int \operatorname{sech}^{-1}(ax)^2 \csc\left(i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2}\right)^3 d\operatorname{sech}^{-1}(ax) \right)}{a^5} \\
 & \quad \downarrow \text{4255} \\
 & \frac{-\frac{1}{5} a^5 x^5 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{5} \left( \frac{1}{6} \left( -\frac{1}{2} \int ax d\operatorname{sech}^{-1}(ax) - \frac{1}{2} ax \sqrt{\frac{1-ax}{ax+1}} (ax+1) \right) + \frac{3}{4} \int \operatorname{sech}^{-1}(ax)^2 \csc\left(i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2}\right)^3 d\operatorname{sech}^{-1}(ax) \right)}{a^5} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{-\frac{1}{5}a^5x^5\operatorname{sech}^{-1}(ax)^3 + \frac{3}{5}\left(\frac{1}{6}\left(-\frac{1}{2}ax\sqrt{\frac{1-ax}{ax+1}}(ax+1) - \frac{1}{2}\int\csc\left(\operatorname{isech}^{-1}(ax) + \frac{\pi}{2}\right)d\operatorname{sech}^{-1}(ax)\right) + \frac{3}{4}\int\operatorname{sech}^{-1}(ax)\right)}{a^5}$$

↓ 4257

$$\frac{-\frac{1}{5}a^5x^5\operatorname{sech}^{-1}(ax)^3 + \frac{3}{5}\left(\frac{3}{4}\int\operatorname{sech}^{-1}(ax)^2\csc\left(\operatorname{isech}^{-1}(ax) + \frac{\pi}{2}\right)^3d\operatorname{sech}^{-1}(ax) + \frac{1}{4}a^3x^3\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)\right)}{a^5}$$

↓ 4674

$$\frac{\frac{3}{5}\left(\frac{3}{4}\left(-\int axd\operatorname{sech}^{-1}(ax) + \frac{1}{2}\int ax\operatorname{sech}^{-1}(ax)^2d\operatorname{sech}^{-1}(ax) + \frac{1}{2}ax\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2 + ax\operatorname{sech}^{-1}(ax)\right)\right)}{a^5}$$

↓ 3042

$$\frac{-\frac{1}{5}a^5x^5\operatorname{sech}^{-1}(ax)^3 + \frac{3}{5}\left(\frac{3}{4}\left(-\int\csc\left(\operatorname{isech}^{-1}(ax) + \frac{\pi}{2}\right)d\operatorname{sech}^{-1}(ax) + \frac{1}{2}\int\operatorname{sech}^{-1}(ax)^2\csc\left(\operatorname{isech}^{-1}(ax) + \frac{\pi}{2}\right)\right)\right)}{a^5}$$

↓ 4257

$$\frac{-\frac{1}{5}a^5x^5\operatorname{sech}^{-1}(ax)^3 + \frac{3}{5}\left(\frac{3}{4}\left(\frac{1}{2}\int\operatorname{sech}^{-1}(ax)^2\csc\left(\operatorname{isech}^{-1}(ax) + \frac{\pi}{2}\right)d\operatorname{sech}^{-1}(ax) - \arctan\left(\frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)}{ax}\right) + \frac{1}{2}\int\operatorname{sech}^{-1}(ax)\right)\right)}{a^5}$$

↓ 4668

$$\frac{-\frac{1}{5}a^5x^5\operatorname{sech}^{-1}(ax)^3 + \frac{3}{5}\left(\frac{3}{4}\left(\frac{1}{2}\left(-2i\int\operatorname{sech}^{-1}(ax)\log\left(1 - ie^{\operatorname{sech}^{-1}(ax)}\right)d\operatorname{sech}^{-1}(ax) + 2i\int\operatorname{sech}^{-1}(ax)\log\left(1 - ie^{\operatorname{sech}^{-1}(ax)}\right)\right)\right)\right)}{a^5}$$

↓ 3011

$$\frac{-\frac{1}{5}a^5x^5\operatorname{sech}^{-1}(ax)^3 + \frac{3}{5}\left(\frac{3}{4}\left(\frac{1}{2}\left(2i\left(\int\operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right)d\operatorname{sech}^{-1}(ax) - \operatorname{sech}^{-1}(ax)\operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right)\right)\right)\right)\right)}{a^5}$$

↓ 2720



$$-\frac{1}{5}a^5x^5\operatorname{sech}^{-1}(ax)^3 + \frac{3}{5}\left(\frac{3}{4}\left(\frac{1}{2}\left(2i\left(\int e^{-\operatorname{sech}^{-1}(ax)}\operatorname{PolyLog}\left(2,-ie^{\operatorname{sech}^{-1}(ax)}\right)de^{\operatorname{sech}^{-1}(ax)} - \operatorname{sech}^{-1}(ax)\operatorname{PolyLog}\right.\right.\right.\right.$$

↓ 7143

$$-\frac{1}{5}a^5x^5\operatorname{sech}^{-1}(ax)^3 + \frac{3}{5}\left(\frac{1}{4}a^3x^3\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2 + \frac{1}{6}a^3x^3\operatorname{sech}^{-1}(ax) + \frac{3}{4}\left(\frac{1}{2}\left(2\operatorname{sech}^{-1}(ax)^2\arctan\right.\right.\right.$$

input

```
Int[x^4*ArcSech[a*x]^3,x]
```

output

```
-((-1/5*(a^5*x^5*ArcSech[a*x]^3) + (3*((a^3*x^3*ArcSech[a*x])/6 + (a^3*x^3*
*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x]^2)/4 + (-1/2*(a*x*Sqrt[(
1 - a*x)/(1 + a*x)]*(1 + a*x)) - ArcTan[(Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*
x))/(a*x])/2)/6 + (3*(a*x*ArcSech[a*x] + (a*x*Sqrt[(1 - a*x)/(1 + a*x)]*(1
+ a*x)*ArcSech[a*x]^2)/2 - ArcTan[(Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x))/(
a*x]) + (2*ArcSech[a*x]^2*ArcTan[E^ArcSech[a*x]]) + (2*I)*(-(ArcSech[a*x]*P
olyLog[2, (-I)*E^ArcSech[a*x]]) + PolyLog[3, (-I)*E^ArcSech[a*x]]) - (2*I)
*(-(ArcSech[a*x]*PolyLog[2, I*E^ArcSech[a*x]]) + PolyLog[3, I*E^ArcSech[a*
x])))/2)/4)/5)/a^5)
```

### Defintions of rubi rules used

rule 2720

```
Int[u_, x_Symbol] :=> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] :=> Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n_, x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^m_, x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 4674 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^n_*(c_.) + (d_.)*(x_)^m_, x_Symbol] := Simp[(-b^2)*(c + d*x)^m*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))), x] + (-Simp[b^2*d*m*(c + d*x)^(m - 1)*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x] + Simp[b^2*d^2*m*((m - 1)/(f^2*(n - 1)*(n - 2))) Int[(c + d*x)^(m - 2)*(b*Csc[e + f*x])^(n - 2), x], x] + Simp[b^2*((n - 2)/(n - 1)) Int[(c + d*x)^m*(b*Csc[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2] && GtQ[m, 1]`

rule 5941 `Int[(x_)^m_*Sech[(a_.) + (b_.)*(x_)^n_]^(p_)*Tanh[(a_.) + (b_.)*(x_)^n_]^(q_), x_Symbol] := Simp[(-x^(m - n + 1))*(Sech[a + b*x^n]^p/(b*n*p)), x] + Simp[(m - n + 1)/(b*n*p) Int[x^(m - n)*Sech[a + b*x^n]^p, x], x] /; FreeQ[{a, b, p}, x] && RationalQ[m] && IntegerQ[n] && GeQ[m - n, 0] && EqQ[q, 1]`

rule 6839

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))^(n_)*(x_)^(m_.), x_Symbol] := Simp[
-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

**Maple [F]**

$$\int x^4 \operatorname{arcsech}(xa)^3 dx$$

input

```
int(x^4*arcsech(x*a)^3,x)
```

output

```
int(x^4*arcsech(x*a)^3,x)
```

**Fricas [F]**

$$\int x^4 \operatorname{sech}^{-1}(ax)^3 dx = \int x^4 \operatorname{arsech}(ax)^3 dx$$

input

```
integrate(x^4*arcsech(a*x)^3,x, algorithm="fricas")
```

output

```
integral(x^4*arcsech(a*x)^3, x)
```

**Sympy [F]**

$$\int x^4 \operatorname{sech}^{-1}(ax)^3 dx = \int x^4 \operatorname{asech}^3(ax) dx$$

input `integrate(x**4*asech(a*x)**3,x)`

output `Integral(x**4*asech(a*x)**3, x)`

**Maxima [F]**

$$\int x^4 \operatorname{sech}^{-1}(ax)^3 dx = \int x^4 \operatorname{arsech}(ax)^3 dx$$

input `integrate(x^4*arcsech(a*x)^3,x, algorithm="maxima")`

output `integrate(x^4*arcsech(a*x)^3, x)`

**Giac [F]**

$$\int x^4 \operatorname{sech}^{-1}(ax)^3 dx = \int x^4 \operatorname{arsech}(ax)^3 dx$$

input `integrate(x^4*arcsech(a*x)^3,x, algorithm="giac")`

output `integrate(x^4*arcsech(a*x)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^4 \operatorname{sech}^{-1}(ax)^3 dx = \int x^4 \operatorname{acosh}\left(\frac{1}{ax}\right)^3 dx$$

input `int(x^4*acosh(1/(a*x))^3,x)`output `int(x^4*acosh(1/(a*x))^3, x)`**Reduce [F]**

$$\int x^4 \operatorname{sech}^{-1}(ax)^3 dx = \int \operatorname{asech}(ax)^3 x^4 dx$$

input `int(x^4*asech(a*x)^3,x)`output `int(asech(a*x)**3*x**4,x)`

### 3.11 $\int x^3 \operatorname{sech}^{-1}(ax)^3 dx$

Optimal result	165
Mathematica [A] (verified)	166
Rubi [C] (verified)	166
Maple [A] (verified)	170
Fricas [F]	171
Sympy [F]	171
Maxima [F]	172
Giac [F]	172
Mupad [F(-1)]	172
Reduce [F]	173

#### Optimal result

Integrand size = 10, antiderivative size = 184

$$\int x^3 \operatorname{sech}^{-1}(ax)^3 dx = \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)}{4a^4} - \frac{x^2 \operatorname{sech}^{-1}(ax)}{4a^2}$$

$$- \frac{\operatorname{sech}^{-1}(ax)^2}{2a^4} - \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax) \operatorname{sech}^{-1}(ax)^2}{2a^4}$$

$$- \frac{x^2 \sqrt{\frac{1-ax}{1+ax}}(1+ax) \operatorname{sech}^{-1}(ax)^2}{4a^2} + \frac{1}{4} x^4 \operatorname{sech}^{-1}(ax)^3$$

$$+ \frac{\operatorname{sech}^{-1}(ax) \log\left(1 + e^{2\operatorname{sech}^{-1}(ax)}\right)}{a^4} + \frac{\operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(ax)}\right)}{2a^4}$$

output

```
1/4*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)/a^4-1/4*x^2*arcsech(a*x)/a^2-1/2*arcs
ech(a*x)^2/a^4-1/2*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*arcsech(a*x)^2/a^4-1/4
*x^2*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*arcsech(a*x)^2/a^2+1/4*x^4*arcsech(a
*x)^3+arcsech(a*x)*ln(1+(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2))^2)/a^4+1/
2*polylog(2,-(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2))^2)/a^4
```

**Mathematica [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.02

$$\int x^3 \operatorname{sech}^{-1}(ax)^3 dx$$

$$= \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax) - \left(-2 + 2\sqrt{\frac{1-ax}{1+ax}} + 2ax\sqrt{\frac{1-ax}{1+ax}} + a^2x^2\sqrt{\frac{1-ax}{1+ax}} + a^3x^3\sqrt{\frac{1-ax}{1+ax}}\right) \operatorname{sech}^{-1}(ax)^2 + a^4x^4 \operatorname{sech}^{-1}(ax)}{4a^4}$$

input

```
Integrate[x^3*ArcSech[a*x]^3,x]
```

output

```
(Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x) - (-2 + 2*Sqrt[(1 - a*x)/(1 + a*x)] +
2*a*x*Sqrt[(1 - a*x)/(1 + a*x)] + a^2*x^2*Sqrt[(1 - a*x)/(1 + a*x)] + a^3
*x^3*Sqrt[(1 - a*x)/(1 + a*x)])*ArcSech[a*x]^2 + a^4*x^4*ArcSech[a*x]^3 +
ArcSech[a*x]*(-a^2*x^2 + 4*Log[1 + E^(-2*ArcSech[a*x])]) - 2*PolyLog[2,
-E^(-2*ArcSech[a*x])])/(4*a^4)
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.75 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.08, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$ , Rules used = {6839, 5941, 3042, 4674, 3042, 4254, 24, 4672, 26, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \operatorname{sech}^{-1}(ax)^3 dx$$

$$\downarrow \text{6839}$$

$$\frac{\int a^4 x^4 \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^3 d\operatorname{sech}^{-1}(ax)}{a^4}$$

$$\downarrow \text{5941}$$

$$\frac{\frac{3}{4} \int a^4 x^4 \operatorname{sech}^{-1}(ax)^2 d\operatorname{sech}^{-1}(ax) - \frac{1}{4} a^4 x^4 \operatorname{sech}^{-1}(ax)^3}{a^4}$$

↓ 3042

$$\frac{-\frac{1}{4} a^4 x^4 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{4} \int \operatorname{sech}^{-1}(ax)^2 \csc \left( i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2} \right)^4 d\operatorname{sech}^{-1}(ax)}{a^4}$$

↓ 4674

$$\frac{\frac{3}{4} \left( -\frac{1}{3} \int a^2 x^2 d\operatorname{sech}^{-1}(ax) + \frac{2}{3} \int a^2 x^2 \operatorname{sech}^{-1}(ax)^2 d\operatorname{sech}^{-1}(ax) + \frac{1}{3} a^2 x^2 \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 + \frac{1}{3} a^2 x^2 \operatorname{sech}^{-1}(ax) \right)}{a^4}$$

↓ 3042

$$\frac{-\frac{1}{4} a^4 x^4 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{4} \left( -\frac{1}{3} \int \csc \left( i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2} \right)^2 d\operatorname{sech}^{-1}(ax) + \frac{2}{3} \int \operatorname{sech}^{-1}(ax)^2 \csc \left( i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2} \right)^2 d\operatorname{sech}^{-1}(ax) \right)}{a^4}$$

↓ 4254

$$\frac{-\frac{1}{4} a^4 x^4 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{4} \left( -\frac{1}{3} i \int 1 d \left( -i \sqrt{\frac{1-ax}{ax+1}} (ax+1) \right) + \frac{2}{3} \int \operatorname{sech}^{-1}(ax)^2 \csc \left( i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2} \right)^2 d\operatorname{sech}^{-1}(ax) \right)}{a^4}$$

↓ 24

$$\frac{-\frac{1}{4} a^4 x^4 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{4} \left( \frac{2}{3} \int \operatorname{sech}^{-1}(ax)^2 \csc \left( i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2} \right)^2 d\operatorname{sech}^{-1}(ax) + \frac{1}{3} a^2 x^2 \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax) \right)}{a^4}$$

↓ 4672

$$\frac{-\frac{1}{4} a^4 x^4 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{4} \left( \frac{2}{3} \left( \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 - 2i \int -i \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax) d\operatorname{sech}^{-1}(ax) \right) \right)}{a^4}$$

↓ 26

$$\frac{\frac{3}{4} \left( \frac{2}{3} \left( \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 - 2 \int \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax) d\operatorname{sech}^{-1}(ax) \right) + \frac{1}{3} a^2 x^2 \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax) \right)}{a^4}$$

↓ 3042

$$\frac{-\frac{1}{4} a^4 x^4 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{4} \left( \frac{2}{3} \left( \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 - 2 \int -i \operatorname{sech}^{-1}(ax) \tan \left( i \operatorname{sech}^{-1}(ax) \right) d\operatorname{sech}^{-1}(ax) \right) \right)}{a^4}$$



↓ 26

$$\frac{-\frac{1}{4}a^4x^4\operatorname{sech}^{-1}(ax)^3 + \frac{3}{4}\left(\frac{2}{3}\left(\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2 + 2i \int \operatorname{sech}^{-1}(ax) \tan(i\operatorname{sech}^{-1}(ax)) d\operatorname{sech}^{-1}(ax)\right) + \dots}{a^4}$$

↓ 4201

$$\frac{-\frac{1}{4}a^4x^4\operatorname{sech}^{-1}(ax)^3 + \frac{3}{4}\left(\frac{2}{3}\left(\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2 + 2i\left(2i \int \frac{e^{2\operatorname{sech}^{-1}(ax)}\operatorname{sech}^{-1}(ax)}{1+e^{2\operatorname{sech}^{-1}(ax)}} d\operatorname{sech}^{-1}(ax) - \frac{1}{2}i\operatorname{sech}^{-1}(ax)\right)\right) + \dots}{a^4}$$

↓ 2620

$$\frac{-\frac{1}{4}a^4x^4\operatorname{sech}^{-1}(ax)^3 + \frac{3}{4}\left(\frac{2}{3}\left(\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2 + 2i\left(2i\left(\frac{1}{2}\operatorname{sech}^{-1}(ax) \log\left(e^{2\operatorname{sech}^{-1}(ax)} + 1\right) - \frac{1}{2} \int \log\right)\right)\right) + \dots}{a^4}$$

↓ 2715

$$\frac{-\frac{1}{4}a^4x^4\operatorname{sech}^{-1}(ax)^3 + \frac{3}{4}\left(\frac{2}{3}\left(\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2 + 2i\left(2i\left(\frac{1}{2}\operatorname{sech}^{-1}(ax) \log\left(e^{2\operatorname{sech}^{-1}(ax)} + 1\right) - \frac{1}{4} \int e^{-2\operatorname{sech}^{-1}(ax)}\right)\right)\right) + \dots}{a^4}$$

↓ 2838

$$\frac{-\frac{1}{4}a^4x^4\operatorname{sech}^{-1}(ax)^3 + \frac{3}{4}\left(\frac{1}{3}a^2x^2\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2 + \frac{1}{3}a^2x^2\operatorname{sech}^{-1}(ax) + \frac{2}{3}\left(\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)\right)\right) + \dots}{a^4}$$

input `Int[x^3*ArcSech[a*x]^3,x]`

output `-((-1/4*(a^4*x^4*ArcSech[a*x]^3) + (3*(-1/3*(Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)) + (a^2*x^2*ArcSech[a*x])/3 + (a^2*x^2*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x]^2)/3 + (2*(Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x]^2 + (2*I)*((-1/2*I)*ArcSech[a*x]^2 + (2*I)*((ArcSech[a*x]*Log[1 + E^(2*ArcSech[a*x]]))/2 + PolyLog[2, -E^(2*ArcSech[a*x]])/4))))/3))/4)/a^4`

## Definitions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4201 `Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_]*(f_)*(x_))], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))]], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 4254 `Int[csc[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

```
rule 4672 Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Simp
[(-c + d*x)^m*(Cot[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)
*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

```
rule 4674 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((c_.) + (d_.)*(x_)^(m_), x_Symbol] := Simp[(-b^2)*(c + d*x)^m*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))), x] + (-Simp[b^2*d*m*(c + d*x)^(m - 1)*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x] + Simp[b^2*d^2*m*((m - 1)/(f^2*(n - 1)*(n - 2))) Int[(c + d*x)^(m - 2)*(b*Csc[e + f*x])^(n - 2), x], x] + Simp[b^2*((n - 2)/(n - 1)) Int[(c + d*x)^m*(b*Csc[e + f*x])^(n - 2), x], x] /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2] && GtQ[m, 1]
```

```
rule 5941 Int[(x_)^(m_)*Sech[(a_.) + (b_.)*(x_)^(n_.)]^(p_)*Tanh[(a_.) + (b_.)*(x_)^(n_.)]^(q_), x_Symbol] := Simp[(-x^(m - n + 1))*(Sech[a + b*x^n]^p/(b*n*p)), x] + Simp[(m - n + 1)/(b*n*p) Int[x^(m - n)*Sech[a + b*x^n]^p, x], x] /; FreeQ[{a, b, p}, x] && RationalQ[m] && IntegerQ[n] && GeQ[m - n, 0] && EqQ[q, 1]
```

```
rule 6839 Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^(n_)*(x_)^(m_), x_Symbol] := Simp[-(c^(m + 1))^( -1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])
```

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.28

method	result
derivativedivides	$\frac{x^4 a^4 \operatorname{arcsech}(xa)^3}{4} - \frac{\operatorname{arcsech}(xa)^2 \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} x^3 a^3}{4} - \frac{\operatorname{arcsech}(xa)^2 \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} xa}{2} - \frac{\operatorname{arcsech}(xa)x^2 a^2}{4} + \frac{\sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}}}{4}$
default	$\frac{x^4 a^4 \operatorname{arcsech}(xa)^3}{4} - \frac{\operatorname{arcsech}(xa)^2 \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} x^3 a^3}{4} - \frac{\operatorname{arcsech}(xa)^2 \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} xa}{2} - \frac{\operatorname{arcsech}(xa)x^2 a^2}{4} + \frac{\sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}}}{4}$

```
input int(x^3*arcsech(x*a)^3,x,method=_RETURNVERBOSE)
```

output

```
1/a^4*(1/4*x^4*a^4*arcsech(x*a)^3-1/4*arcsech(x*a)^2*(-(a*x-1)/x/a)^(1/2)*
((a*x+1)/x/a)^(1/2)*x^3*a^3-1/2*arcsech(x*a)^2*(-(a*x-1)/x/a)^(1/2)*((a*x+
1)/x/a)^(1/2)*x*a-1/4*arcsech(x*a)*x^2*a^2+1/4*(-(a*x-1)/x/a)^(1/2)*((a*x+
1)/x/a)^(1/2)*x*a-1/2*arcsech(x*a)^2-1/4+arcsech(x*a)*ln(1+(1/a/x+(1/a/x-1
)^(1/2)*(1+1/a/x)^(1/2))^2)+1/2*polylog(2,-(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x
)^(1/2))^2))
```

**Fricas [F]**

$$\int x^3 \operatorname{sech}^{-1}(ax)^3 dx = \int x^3 \operatorname{arsech}(ax)^3 dx$$

input

```
integrate(x^3*arcsech(a*x)^3,x, algorithm="fricas")
```

output

```
integral(x^3*arcsech(a*x)^3, x)
```

**Sympy [F]**

$$\int x^3 \operatorname{sech}^{-1}(ax)^3 dx = \int x^3 \operatorname{asech}^3(ax) dx$$

input

```
integrate(x**3*asech(a*x)**3,x)
```

output

```
Integral(x**3*asech(a*x)**3, x)
```

**Maxima [F]**

$$\int x^3 \operatorname{sech}^{-1}(ax)^3 dx = \int x^3 \operatorname{arsech}(ax)^3 dx$$

input `integrate(x^3*arcsech(a*x)^3,x, algorithm="maxima")`

output `integrate(x^3*arcsech(a*x)^3, x)`

**Giac [F]**

$$\int x^3 \operatorname{sech}^{-1}(ax)^3 dx = \int x^3 \operatorname{arsech}(ax)^3 dx$$

input `integrate(x^3*arcsech(a*x)^3,x, algorithm="giac")`

output `integrate(x^3*arcsech(a*x)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 \operatorname{sech}^{-1}(ax)^3 dx = \int x^3 \operatorname{acosh}\left(\frac{1}{ax}\right)^3 dx$$

input `int(x^3*acosh(1/(a*x))^3,x)`

output `int(x^3*acosh(1/(a*x))^3, x)`

Reduce [F]

$$\int x^3 \operatorname{sech}^{-1}(ax)^3 dx = \int \operatorname{asech}(ax)^3 x^3 dx$$

input `int(x^3*asech(a*x)^3,x)`

output `int(asech(a*x)**3*x**3,x)`

### 3.12 $\int x^2 \operatorname{sech}^{-1}(ax)^3 dx$

Optimal result	174
Mathematica [A] (verified)	175
Rubi [A] (verified)	175
Maple [F]	179
Fricas [F]	179
Sympy [F]	179
Maxima [F]	180
Giac [F]	180
Mupad [F(-1)]	180
Reduce [F]	181

#### Optimal result

Integrand size = 10, antiderivative size = 198

$$\int x^2 \operatorname{sech}^{-1}(ax)^3 dx = -\frac{x \operatorname{sech}^{-1}(ax)}{a^2} - \frac{x \sqrt{\frac{1-ax}{1+ax}}(1+ax) \operatorname{sech}^{-1}(ax)^2}{2a^2} + \frac{1}{3}x^3 \operatorname{sech}^{-1}(ax)^3$$

$$- \frac{\operatorname{sech}^{-1}(ax)^2 \arctan\left(e^{\operatorname{sech}^{-1}(ax)}\right)}{a^3} + \frac{\arctan\left(\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)}{ax}\right)}{a^3}$$

$$+ \frac{i \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right)}{a^3}$$

$$- \frac{i \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(ax)}\right)}{a^3}$$

$$- \frac{i \operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(ax)}\right)}{a^3} + \frac{i \operatorname{PolyLog}\left(3, ie^{\operatorname{sech}^{-1}(ax)}\right)}{a^3}$$

output

```
-x*arcsech(a*x)/a^2-1/2*x*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*arcsech(a*x)^2/
a^2+1/3*x^3*arcsech(a*x)^3-arcsech(a*x)^2*arctan(1/a/x+(-1+1/a/x)^(1/2)*(1
+1/a/x)^(1/2))/a^3+arctan(((a*x+1)/(a*x+1))^(1/2)*(a*x+1)/a/x)/a^3+I*arcs
ech(a*x)*polylog(2,-I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a^3-I*arcs
ech(a*x)*polylog(2,I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a^3-I*polyl
og(3,-I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a^3+I*polylog(3,I*(1/a/x
+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a^3
```

**Mathematica [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.01

$$\int x^2 \operatorname{sech}^{-1}(ax)^3 dx$$

$$= \frac{-6ax \operatorname{sech}^{-1}(ax) - 3ax \sqrt{\frac{1-ax}{1+ax}}(1+ax) \operatorname{sech}^{-1}(ax)^2 + 2a^3 x^3 \operatorname{sech}^{-1}(ax)^3 + 3i \left( -4i \arctan \left( \tanh \left( \frac{1}{2} \operatorname{sech}^{-1}(ax) \right) \right) \right)}{a^3}$$

input

```
Integrate[x^2*ArcSech[a*x]^3,x]
```

output

```
(-6*a*x*ArcSech[a*x] - 3*a*x*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x]^2 + 2*a^3*x^3*ArcSech[a*x]^3 + (3*I)*((-4*I)*ArcTan[Tanh[ArcSech[a*x]/2]] + ArcSech[a*x]^2*Log[1 - I/E^ArcSech[a*x]] - ArcSech[a*x]^2*Log[1 + I/E^ArcSech[a*x]] + 2*ArcSech[a*x]*PolyLog[2, (-I)/E^ArcSech[a*x]] - 2*ArcSech[a*x]*PolyLog[2, I/E^ArcSech[a*x]] + 2*PolyLog[3, (-I)/E^ArcSech[a*x]] - 2*PolyLog[3, I/E^ArcSech[a*x]])/(6*a^3)
```

**Rubi [A] (verified)**

Time = 0.70 (sec) , antiderivative size = 187, normalized size of antiderivative = 0.94, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {6839, 5941, 3042, 4674, 3042, 4257, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{sech}^{-1}(ax)^3 dx$$

$$\downarrow \text{6839}$$

$$\frac{\int a^3 x^3 \sqrt{\frac{1-ax}{ax+1}}(ax+1) \operatorname{sech}^{-1}(ax)^3 d \operatorname{sech}^{-1}(ax)}{a^3}$$

$$\downarrow \text{5941}$$

$$\frac{\int a^3 x^3 \operatorname{sech}^{-1}(ax)^2 d \operatorname{sech}^{-1}(ax) - \frac{1}{3} a^3 x^3 \operatorname{sech}^{-1}(ax)^3}{a^3}$$



---

↓ 3042

$$\frac{-\frac{1}{3}a^3x^3\operatorname{sech}^{-1}(ax)^3 + \int \operatorname{sech}^{-1}(ax)^2 \csc\left(\operatorname{isech}^{-1}(ax) + \frac{\pi}{2}\right)^3 d\operatorname{sech}^{-1}(ax)}{a^3}$$

↓ 4674

$$\frac{-\int ax d\operatorname{sech}^{-1}(ax) + \frac{1}{2} \int ax \operatorname{sech}^{-1}(ax)^2 d\operatorname{sech}^{-1}(ax) - \frac{1}{3}a^3x^3\operatorname{sech}^{-1}(ax)^3 + \frac{1}{2}ax\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{a^3}$$

↓ 3042

$$\frac{-\int \csc\left(\operatorname{isech}^{-1}(ax) + \frac{\pi}{2}\right) d\operatorname{sech}^{-1}(ax) + \frac{1}{2} \int \operatorname{sech}^{-1}(ax)^2 \csc\left(\operatorname{isech}^{-1}(ax) + \frac{\pi}{2}\right) d\operatorname{sech}^{-1}(ax) - \frac{1}{3}a^3x^3\operatorname{sech}^{-1}(ax)^3}{a^3}$$

↓ 4257

$$\frac{\frac{1}{2} \int \operatorname{sech}^{-1}(ax)^2 \csc\left(\operatorname{isech}^{-1}(ax) + \frac{\pi}{2}\right) d\operatorname{sech}^{-1}(ax) - \frac{1}{3}a^3x^3\operatorname{sech}^{-1}(ax)^3 - \arctan\left(\frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)}{ax}\right) + \frac{1}{2}ax\sqrt{\frac{1-ax}{ax+1}}}{a^3}$$

↓ 4668

$$\frac{\frac{1}{2}\left(-2i \int \operatorname{sech}^{-1}(ax) \log\left(1 - ie^{\operatorname{sech}^{-1}(ax)}\right) d\operatorname{sech}^{-1}(ax) + 2i \int \operatorname{sech}^{-1}(ax) \log\left(1 + ie^{\operatorname{sech}^{-1}(ax)}\right) d\operatorname{sech}^{-1}(ax) + \dots\right)}{a^3}$$

↓ 3011

$$\frac{\frac{1}{2}\left(2i\left(\int \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right) d\operatorname{sech}^{-1}(ax) - \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right)\right) - 2i\left(\int \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(ax)}\right) d\operatorname{sech}^{-1}(ax) - \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(ax)}\right)\right)\right)}{a^3}$$

↓ 2720

$$\frac{\frac{1}{2}\left(2i\left(\int e^{-\operatorname{sech}^{-1}(ax)} \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right) de^{\operatorname{sech}^{-1}(ax)} - \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right)\right) - 2i\left(\int e^{\operatorname{sech}^{-1}(ax)} \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(ax)}\right) de^{\operatorname{sech}^{-1}(ax)} - \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(ax)}\right)\right)\right)}{a^3}$$

↓ 7143

---

$$-\frac{1}{3}a^3x^3\operatorname{sech}^{-1}(ax)^3 + \frac{1}{2}\left(2\operatorname{sech}^{-1}(ax)^2 \arctan\left(e^{\operatorname{sech}^{-1}(ax)}\right) + 2i\left(\operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(ax)}\right) - \operatorname{sech}^{-1}(ax)\operatorname{Pol}\right)\right)$$

input `Int[x^2*ArcSech[a*x]^3,x]`

output `-((a*x*ArcSech[a*x] + (a*x*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x]^2)/2 - (a^3*x^3*ArcSech[a*x]^3)/3 - ArcTan[(Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x))/(a*x)] + (2*ArcSech[a*x]^2*ArcTan[E^ArcSech[a*x]] + (2*I)*(-(ArcSech[a*x]*PolyLog[2, (-I)*E^ArcSech[a*x]]) + PolyLog[3, (-I)*E^ArcSech[a*x]]) - (2*I)*(-(ArcSech[a*x]*PolyLog[2, I*E^ArcSech[a*x]]) + PolyLog[3, I*E^ArcSech[a*x]]))/2)/a^3)`

### Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x))))^n]/(b*c*n*Log[F]), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x))))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 4674 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((c_.) + (d_.)*(x_))^(m_), x_Symbol] := Simp[(-b^2)*(c + d*x)^m*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))), x] + (-Simp[b^2*d*m*(c + d*x)^(m - 1)*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x] + Simp[b^2*d^2*m*((m - 1)/(f^2*(n - 1)*(n - 2))) Int[(c + d*x)^(m - 2)*(b*Csc[e + f*x])^(n - 2), x], x] + Simp[b^2*((n - 2)/(n - 1)) Int[(c + d*x)^m*(b*Csc[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2] && GtQ[m, 1]`

rule 5941 `Int[(x_)^(m_.)*Sech[(a_.) + (b_.)*(x_)^(n_.)]^(p_.)*Tanh[(a_.) + (b_.)*(x_)^(n_.)]^(q_.), x_Symbol] := Simp[(-x^(m - n + 1))*(Sech[a + b*x^n]^p/(b*n*p)), x] + Simp[(m - n + 1)/(b*n*p) Int[x^(m - n)*Sech[a + b*x^n]^p, x] /; FreeQ[{a, b, p}, x] && RationalQ[m] && IntegerQ[n] && GeQ[m - n, 0] && EqQ[q, 1]`

rule 6839 `Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^(n_)*(x_)^(m_.), x_Symbol] := Simp[-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

**Maple [F]**

$$\int x^2 \operatorname{arcsech}(xa)^3 dx$$

input `int(x^2*arcsech(x*a)^3,x)`

output `int(x^2*arcsech(x*a)^3,x)`

**Fricas [F]**

$$\int x^2 \operatorname{sech}^{-1}(ax)^3 dx = \int x^2 \operatorname{arsech}(ax)^3 dx$$

input `integrate(x^2*arcsech(a*x)^3,x, algorithm="fricas")`

output `integral(x^2*arcsech(a*x)^3, x)`

**Sympy [F]**

$$\int x^2 \operatorname{sech}^{-1}(ax)^3 dx = \int x^2 \operatorname{asech}^3(ax) dx$$

input `integrate(x**2*asech(a*x)**3,x)`

output `Integral(x**2*asech(a*x)**3, x)`

**Maxima [F]**

$$\int x^2 \operatorname{sech}^{-1}(ax)^3 dx = \int x^2 \operatorname{arsech}(ax)^3 dx$$

input `integrate(x^2*arcsech(a*x)^3,x, algorithm="maxima")`

output `integrate(x^2*arcsech(a*x)^3, x)`

**Giac [F]**

$$\int x^2 \operatorname{sech}^{-1}(ax)^3 dx = \int x^2 \operatorname{arsech}(ax)^3 dx$$

input `integrate(x^2*arcsech(a*x)^3,x, algorithm="giac")`

output `integrate(x^2*arcsech(a*x)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{sech}^{-1}(ax)^3 dx = \int x^2 \operatorname{acosh}\left(\frac{1}{ax}\right)^3 dx$$

input `int(x^2*acosh(1/(a*x))^3,x)`

output `int(x^2*acosh(1/(a*x))^3, x)`

**Reduce [F]**

$$\int x^2 \operatorname{sech}^{-1}(ax)^3 dx = \int \operatorname{asech}(ax)^3 x^2 dx$$

input `int(x^2*asech(a*x)^3,x)`

output `int(asech(a*x)**3*x**2,x)`

### 3.13 $\int x \operatorname{sech}^{-1}(ax)^3 dx$

Optimal result	182
Mathematica [A] (verified)	182
Rubi [C] (verified)	183
Maple [A] (verified)	186
Fricas [F]	187
Sympy [F]	187
Maxima [F]	187
Giac [F]	188
Mupad [F(-1)]	188
Reduce [F]	188

#### Optimal result

Integrand size = 8, antiderivative size = 102

$$\int x \operatorname{sech}^{-1}(ax)^3 dx = -\frac{3 \operatorname{sech}^{-1}(ax)^2}{2a^2} - \frac{3 \sqrt{\frac{1-ax}{1+ax}}(1+ax) \operatorname{sech}^{-1}(ax)^2}{2a^2} + \frac{1}{2} x^2 \operatorname{sech}^{-1}(ax)^3$$

$$+ \frac{3 \operatorname{sech}^{-1}(ax) \log\left(1 + e^{2 \operatorname{sech}^{-1}(ax)}\right)}{a^2} + \frac{3 \operatorname{PolyLog}\left(2, -e^{2 \operatorname{sech}^{-1}(ax)}\right)}{2a^2}$$

output

$$\begin{aligned} & -3/2*\operatorname{arcsech}(a*x)^2/a^2-3/2*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*\operatorname{arcsech}(a*x)^2/a^2 \\ & +1/2*x^2*\operatorname{arcsech}(a*x)^3+3*\operatorname{arcsech}(a*x)*\ln(1+(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2))^2)/a^2 \\ & +3/2*\operatorname{polylog}(2,-(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2))^2)/a^2 \end{aligned}$$

#### Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.99

$$\int x \operatorname{sech}^{-1}(ax)^3 dx = \frac{\operatorname{sech}^{-1}(ax) \left( -3 \left( -1 + \sqrt{\frac{1-ax}{1+ax}} + ax \sqrt{\frac{1-ax}{1+ax}} \right) \operatorname{sech}^{-1}(ax) + a^2 x^2 \operatorname{sech}^{-1}(ax)^2 + 6 \log \left( 1 + e^{-2 \operatorname{sech}^{-1}(ax)} \right) \right)}{2a^2}$$

input `Integrate[x*ArcSech[a*x]^3,x]`

output  $(\text{ArcSech}[a*x]*(-3*(-1 + \text{Sqrt}[(1 - a*x)/(1 + a*x)] + a*x*\text{Sqrt}[(1 - a*x)/(1 + a*x)]))*\text{ArcSech}[a*x] + a^2*x^2*\text{ArcSech}[a*x]^2 + 6*\text{Log}[1 + E^{(-2*\text{ArcSech}[a*x])}] - 3*\text{PolyLog}[2, -E^{(-2*\text{ArcSech}[a*x])}])/(2*a^2)$

## Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.53 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.12, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.375$ , Rules used = {6839, 5941, 3042, 4672, 26, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \operatorname{sech}^{-1}(ax)^3 dx \\
 & \quad \downarrow 6839 \\
 & \frac{\int a^2 x^2 \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^3 d\operatorname{sech}^{-1}(ax)}{a^2} \\
 & \quad \downarrow 5941 \\
 & \frac{\frac{3}{2} \int a^2 x^2 \operatorname{sech}^{-1}(ax)^2 d\operatorname{sech}^{-1}(ax) - \frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^3}{a^2} \\
 & \quad \downarrow 3042 \\
 & \frac{-\frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{2} \int \operatorname{sech}^{-1}(ax)^2 \csc\left(i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2}\right)^2 d\operatorname{sech}^{-1}(ax)}{a^2} \\
 & \quad \downarrow 4672 \\
 & \frac{-\frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{2} \left( \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 - 2i \int -i \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax) d\operatorname{sech}^{-1}(ax) \right)}{a^2} \\
 & \quad \downarrow 26
 \end{aligned}$$



$$\frac{\frac{3}{2} \left( \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 - 2 \int \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax) d\operatorname{sech}^{-1}(ax) \right) - \frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^3}{a^2}$$

↓ 3042

$$\frac{-\frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{2} \left( \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 - 2 \int -i \operatorname{sech}^{-1}(ax) \tan(i \operatorname{sech}^{-1}(ax)) d\operatorname{sech}^{-1}(ax) \right)}{a^2}$$

↓ 26

$$\frac{-\frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{2} \left( \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 + 2i \int \operatorname{sech}^{-1}(ax) \tan(i \operatorname{sech}^{-1}(ax)) d\operatorname{sech}^{-1}(ax) \right)}{a^2}$$

↓ 4201

$$\frac{-\frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{2} \left( \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 + 2i \left( 2i \int \frac{e^{2\operatorname{sech}^{-1}(ax)} \operatorname{sech}^{-1}(ax)}{1+e^{2\operatorname{sech}^{-1}(ax)}} d\operatorname{sech}^{-1}(ax) - \frac{1}{2} i \operatorname{sech}^{-1}(ax) \right) \right)}{a^2}$$

↓ 2620

$$\frac{-\frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{2} \left( \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 + 2i \left( 2i \left( \frac{1}{2} \operatorname{sech}^{-1}(ax) \log(e^{2\operatorname{sech}^{-1}(ax)} + 1) \right) - \frac{1}{2} \int \log(1) \right) \right)}{a^2}$$

↓ 2715

$$\frac{-\frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{2} \left( \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 + 2i \left( 2i \left( \frac{1}{2} \operatorname{sech}^{-1}(ax) \log(e^{2\operatorname{sech}^{-1}(ax)} + 1) \right) - \frac{1}{4} \int e^{-2\operatorname{sech}^{-1}(ax)} \right) \right)}{a^2}$$

↓ 2838

$$\frac{-\frac{1}{2} a^2 x^2 \operatorname{sech}^{-1}(ax)^3 + \frac{3}{2} \left( \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^2 + 2i \left( 2i \left( \frac{1}{4} \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(ax)}\right) \right) + \frac{1}{2} \operatorname{sech}^{-1}(ax) \log(1) \right) \right)}{a^2}$$

input `Int [x*ArcSech[a*x]^3, x]`

output

```

-((-1/2*(a^2*x^2*ArcSech[a*x]^3) + (3*(Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)
*ArcSech[a*x]^2 + (2*I)*((-1/2*I)*ArcSech[a*x]^2 + (2*I)*((ArcSech[a*x]*Lo
g[1 + E^(2*ArcSech[a*x]]))/2 + PolyLog[2, -E^(2*ArcSech[a*x]])/4))))/2)/a^
2)

```

### Defintions of rubi rules used

rule 26

```

Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]

```

rule 2620

```

Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*(F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

```

rule 2715

```

Int[Log[(a_) + (b_)*(F_)^((e_)*((c_) + (d_)*(x_)))^(n_)]], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

```

rule 2838

```

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

```

rule 3042

```

Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

rule 4201

```

Int[(((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_])*(f_)*(x_)], x
_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[
(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

```

rule 4672

```
Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Simp
[(-c + d*x)^m*(Cot[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)
*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

rule 5941

```
Int[(x_)^(m_.)*Sech[(a_.) + (b_.)*(x_)^(n_.)]^(p_.)*Tanh[(a_.) + (b_.)*(x_)
^(n_.)]^(q_.), x_Symbol] := Simp[(-x^(m - n + 1))*(Sech[a + b*x^n]^p/(b*n*p
)), x] + Simp[(m - n + 1)/(b*n*p) Int[x^(m - n)*Sech[a + b*x^n]^p, x], x]
/; FreeQ[{a, b, p}, x] && RationalQ[m] && IntegerQ[n] && GeQ[m - n, 0] &&
EqQ[q, 1]
```

rule 6839

```
Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))^(n_.)*(x_)^(m_.), x_Symbol] := Simp[
-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

## Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.46

method	result
derivativedivides	$\frac{\operatorname{arcsech}(xa)^2 \left( \operatorname{arcsech}(xa) x^2 a^2 - 3 \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} xa + 3 \right)}{2} - 3 \operatorname{arcsech}(xa)^2 + 3 \operatorname{arcsech}(xa) \ln \left( 1 + \left( \frac{1}{ax} + \sqrt{\frac{1}{ax} - 1} \sqrt{1 + \frac{1}{ax}} \right) \right)}{a^2}$
default	$\frac{\operatorname{arcsech}(xa)^2 \left( \operatorname{arcsech}(xa) x^2 a^2 - 3 \sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} xa + 3 \right)}{2} - 3 \operatorname{arcsech}(xa)^2 + 3 \operatorname{arcsech}(xa) \ln \left( 1 + \left( \frac{1}{ax} + \sqrt{\frac{1}{ax} - 1} \sqrt{1 + \frac{1}{ax}} \right) \right)}{a^2}$

input

```
int(x*arcsech(x*a)^3,x,method=_RETURNVERBOSE)
```

output

```
1/a^2*(1/2*arcsech(x*a)^2*(arcsech(x*a)*x^2*a^2-3*(-(a*x-1)/x/a)^(1/2)*((a
*x+1)/x/a)^(1/2)*x*a+3)-3*arcsech(x*a)^2+3*arcsech(x*a)*ln(1+(1/a/x+(1/a/x
-1)^(1/2)*(1+1/a/x)^(1/2))^2)+3/2*polylog(2,-(1/a/x+(1/a/x-1)^(1/2)*(1+1/a
/x)^(1/2))^2))
```

**Fricas [F]**

$$\int x \operatorname{sech}^{-1}(ax)^3 dx = \int x \operatorname{arsech}(ax)^3 dx$$

input `integrate(x*arcsech(a*x)^3,x, algorithm="fricas")`

output `integral(x*arcsech(a*x)^3, x)`

**Sympy [F]**

$$\int x \operatorname{sech}^{-1}(ax)^3 dx = \int x \operatorname{asech}^3(ax) dx$$

input `integrate(x*asech(a*x)**3,x)`

output `Integral(x*asech(a*x)**3, x)`

**Maxima [F]**

$$\int x \operatorname{sech}^{-1}(ax)^3 dx = \int x \operatorname{arsech}(ax)^3 dx$$

input `integrate(x*arcsech(a*x)^3,x, algorithm="maxima")`

output `integrate(x*arcsech(a*x)^3, x)`

**Giac [F]**

$$\int x \operatorname{sech}^{-1}(ax)^3 dx = \int x \operatorname{arsech}(ax)^3 dx$$

input `integrate(x*arcsech(a*x)^3,x, algorithm="giac")`

output `integrate(x*arcsech(a*x)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{sech}^{-1}(ax)^3 dx = \int x \operatorname{acosh}\left(\frac{1}{ax}\right)^3 dx$$

input `int(x*acosh(1/(a*x))^3,x)`

output `int(x*acosh(1/(a*x))^3, x)`

**Reduce [F]**

$$\int x \operatorname{sech}^{-1}(ax)^3 dx = \int \operatorname{asech}(ax)^3 x dx$$

input `int(x*asech(a*x)^3,x)`

output `int(asech(a*x)**3*x,x)`

### 3.14 $\int \operatorname{sech}^{-1}(ax)^3 dx$

Optimal result	189
Mathematica [A] (verified)	190
Rubi [A] (verified)	190
Maple [F]	193
Fricas [F]	193
Sympy [F]	193
Maxima [F]	194
Giac [F]	194
Mupad [F(-1)]	195
Reduce [F]	195

#### Optimal result

Integrand size = 6, antiderivative size = 111

$$\int \operatorname{sech}^{-1}(ax)^3 dx = x \operatorname{sech}^{-1}(ax)^3 - \frac{6 \operatorname{sech}^{-1}(ax)^2 \arctan\left(e^{\operatorname{sech}^{-1}(ax)}\right)}{a} + \frac{6i \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right)}{a} - \frac{6i \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(ax)}\right)}{a} - \frac{6i \operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(ax)}\right)}{a} + \frac{6i \operatorname{PolyLog}\left(3, ie^{\operatorname{sech}^{-1}(ax)}\right)}{a}$$

output

```
x*arcsech(a*x)^3-6*arcsech(a*x)^2*arctan(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2))/a+6*I*arcsech(a*x)*polylog(2,-I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a-6*I*arcsech(a*x)*polylog(2,I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a-6*I*polylog(3,-I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a+6*I*polylog(3,I*(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2)))/a
```

**Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.15

$$\int \operatorname{sech}^{-1}(ax)^3 dx = x \operatorname{sech}^{-1}(ax)^3 - \frac{3i \left( -\operatorname{sech}^{-1}(ax)^2 \left( \log \left( 1 - ie^{-\operatorname{sech}^{-1}(ax)} \right) - \log \left( 1 + ie^{-\operatorname{sech}^{-1}(ax)} \right) \right) - 2 \operatorname{sech}^{-1}(ax) \left( \operatorname{PolyLog} \left( 2, -ie^{-\operatorname{sech}^{-1}(ax)} \right) - \operatorname{PolyLog} \left( 2, ie^{-\operatorname{sech}^{-1}(ax)} \right) \right) \right)}{a}$$

input

```
Integrate[ArcSech[a*x]^3,x]
```

output

```
x*ArcSech[a*x]^3 - (((3*I)*(-(ArcSech[a*x]^2*(Log[1 - I/E^ArcSech[a*x]] - Log[1 + I/E^ArcSech[a*x]])) - 2*ArcSech[a*x]*(PolyLog[2, (-I)/E^ArcSech[a*x]] - PolyLog[2, I/E^ArcSech[a*x]]) - 2*(PolyLog[3, (-I)/E^ArcSech[a*x]] - PolyLog[3, I/E^ArcSech[a*x]]))))/a
```

**Rubi [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.94, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.167$ , Rules used = {6833, 5941, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \operatorname{sech}^{-1}(ax)^3 dx \\ & \quad \downarrow \text{6833} \\ & - \frac{\int ax \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^3 d \operatorname{sech}^{-1}(ax)}{a} \\ & \quad \downarrow \text{5941} \\ & - \frac{3 \int ax \operatorname{sech}^{-1}(ax)^2 d \operatorname{sech}^{-1}(ax) - ax \operatorname{sech}^{-1}(ax)^3}{a} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\frac{-ax \operatorname{sech}^{-1}(ax)^3 + 3 \int \operatorname{sech}^{-1}(ax)^2 \csc\left(\operatorname{isech}^{-1}(ax) + \frac{\pi}{2}\right) d\operatorname{sech}^{-1}(ax)}{a}$$

↓ 4668

$$\frac{-ax \operatorname{sech}^{-1}(ax)^3 + 3\left(-2i \int \operatorname{sech}^{-1}(ax) \log\left(1 - ie^{\operatorname{sech}^{-1}(ax)}\right) d\operatorname{sech}^{-1}(ax) + 2i \int \operatorname{sech}^{-1}(ax) \log\left(1 + ie^{\operatorname{sech}^{-1}(ax)}\right) d\operatorname{sech}^{-1}(ax)\right)}{a}$$

↓ 3011

$$\frac{-ax \operatorname{sech}^{-1}(ax)^3 + 3\left(2i \left(\int \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right) d\operatorname{sech}^{-1}(ax) - \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right)\right)\right)}{a}$$

↓ 2720

$$\frac{-ax \operatorname{sech}^{-1}(ax)^3 + 3\left(2i \left(\int e^{-\operatorname{sech}^{-1}(ax)} \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right) de^{\operatorname{sech}^{-1}(ax)} - \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(ax)}\right)\right)\right)}{a}$$

↓ 7143

$$\frac{-ax \operatorname{sech}^{-1}(ax)^3 + 3\left(2 \operatorname{sech}^{-1}(ax)^2 \arctan\left(e^{\operatorname{sech}^{-1}(ax)}\right) + 2i \left(\operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(ax)}\right) - \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(ax)}\right)\right)\right)}{a}$$

input `Int[ArcSech[a*x]^3, x]`

output `-((- (a*x*ArcSech[a*x]^3) + 3*(2*ArcSech[a*x]^2*ArcTan[E^ArcSech[a*x]] + (2*I)*(-(ArcSech[a*x]*PolyLog[2, (-I)*E^ArcSech[a*x]]) + PolyLog[3, (-I)*E^ArcSech[a*x])) - (2*I)*(-(ArcSech[a*x]*PolyLog[2, I*E^ArcSech[a*x]]) + PolyLog[3, I*E^ArcSech[a*x]])))/a)`

### Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`



rule 3011  $\text{Int}[\text{Log}[1 + (e\_.) * ((F\_)^{(c\_.) * (a\_.) + (b\_.) * (x\_))})^{(n\_.)}] * ((f\_.) + (g\_.) * (x\_))^{(m\_.)}, x\_Symbol] \rightarrow \text{Simp}[(-f + g*x)^m * (\text{PolyLog}[2, (-e) * (F^{(c*(a + b*x))})^n]) / (b*c*n*\text{Log}[F]), x] + \text{Simp}[g*(m/(b*c*n*\text{Log}[F])) \text{Int}[(f + g*x)^{(m - 1)} * \text{PolyLog}[2, (-e) * (F^{(c*(a + b*x))})^n], x], x] /; \text{FreeQ}[\{F, a, b, c, e, f, g, n\}, x] \&\& \text{GtQ}[m, 0]$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4668  $\text{Int}[\text{csc}[(e\_.) + \text{Pi} * (k\_.) + (\text{Complex}[0, fz\_]) * (f\_.) * (x\_)] * ((c\_.) + (d\_.) * (x\_))^{(m\_.)}, x\_Symbol] \rightarrow \text{Simp}[-2 * (c + d*x)^m * (\text{ArcTanh}[E^{((-I)*e + f*fz*x)} / E^{(I*k*Pi)}]) / (f*fz*I), x] + (-\text{Simp}[d * (m / (f*fz*I)) \text{Int}[(c + d*x)^{(m - 1)} * \text{Log}[1 - E^{((-I)*e + f*fz*x)} / E^{(I*k*Pi)}], x], x] + \text{Simp}[d * (m / (f*fz*I)) \text{Int}[(c + d*x)^{(m - 1)} * \text{Log}[1 + E^{((-I)*e + f*fz*x)} / E^{(I*k*Pi)}], x], x]) /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \&\& \text{IntegerQ}[2*k] \&\& \text{IGtQ}[m, 0]$

rule 5941  $\text{Int}[(x\_)^{(m\_.)} * \text{Sech}[(a\_.) + (b\_.) * (x\_)]^{(n\_.)}]^{(p\_.)} * \text{Tanh}[(a\_.) + (b\_.) * (x\_)]^{(q\_.)}, x\_Symbol] \rightarrow \text{Simp}[(-x^{(m - n + 1)}) * (\text{Sech}[a + b*x^n]^p / (b^n * p)), x] + \text{Simp}[(m - n + 1) / (b^n * p) \text{Int}[x^{(m - n)} * \text{Sech}[a + b*x^n]^p, x], x] /; \text{FreeQ}[\{a, b, p\}, x] \&\& \text{RationalQ}[m] \&\& \text{IntegerQ}[n] \&\& \text{GeQ}[m - n, 0] \&\& \text{EqQ}[q, 1]$

rule 6833  $\text{Int}[(a\_.) + \text{ArcSech}[(c\_.) * (x\_)] * (b\_.)]^{(n\_.)}, x\_Symbol] \rightarrow \text{Simp}[-c^{(-1)} \text{Subst}[\text{Int}[(a + b*x)^n * \text{Sech}[x] * \text{Tanh}[x], x], x, \text{ArcSech}[c*x]], x] /; \text{FreeQ}[\{a, b, c, n\}, x] \&\& \text{IGtQ}[n, 0]$

rule 7143  $\text{Int}[\text{PolyLog}[n_, (c\_.) * ((a\_.) + (b\_.) * (x\_))^{(p\_.)}] / ((d\_.) + (e\_.) * (x\_)), x\_Symbol] \rightarrow \text{Simp}[\text{PolyLog}[n + 1, c * (a + b*x)^p] / (e*p), x] /; \text{FreeQ}[\{a, b, c, d, e, n, p\}, x] \&\& \text{EqQ}[b*d, a*e]$

**Maple [F]**

$$\int \operatorname{arcsech}(xa)^3 dx$$

input `int(arcsech(x*a)^3,x)`

output `int(arcsech(x*a)^3,x)`

**Fricas [F]**

$$\int \operatorname{sech}^{-1}(ax)^3 dx = \int \operatorname{arsech}(ax)^3 dx$$

input `integrate(arcsech(a*x)^3,x, algorithm="fricas")`

output `integral(arcsech(a*x)^3, x)`

**Sympy [F]**

$$\int \operatorname{sech}^{-1}(ax)^3 dx = \int \operatorname{asech}^3(ax) dx$$

input `integrate(asech(a*x)**3,x)`

output `Integral(asech(a*x)**3, x)`

**Maxima [F]**

$$\int \operatorname{sech}^{-1}(ax)^3 dx = \int \operatorname{arsech}(ax)^3 dx$$

input `integrate(arcsech(a*x)^3,x, algorithm="maxima")`

output `x*log(sqrt(a*x + 1)*sqrt(-a*x + 1) + 1)^3 - integrate((a^2*x^2*log(a)^3 + (a^2*x^2 - 1)*log(x)^3 + 3*(a^2*x^2*log(a) + (a^2*x^2*(log(a) + 1) + (a^2*x^2 - 1)*log(x) - log(a))*sqrt(a*x + 1)*sqrt(-a*x + 1) + (a^2*x^2 - 1)*log(x) - log(a))*log(sqrt(a*x + 1)*sqrt(-a*x + 1) + 1)^2 - log(a)^3 + 3*(a^2*x^2*log(a) - log(a))*log(x)^2 + (a^2*x^2*log(a)^3 + (a^2*x^2 - 1)*log(x)^3 - log(a)^3 + 3*(a^2*x^2*log(a) - log(a))*log(x)^2 + 3*(a^2*x^2*log(a)^2 - log(a)^2)*log(x))*sqrt(a*x + 1)*sqrt(-a*x + 1) - 3*(a^2*x^2*log(a)^2 + (a^2*x^2 - 1)*log(x)^2 + (a^2*x^2*log(a)^2 + (a^2*x^2 - 1)*log(x)^2 - log(a)^2 + 2*(a^2*x^2*log(a) - log(a))*log(x))*sqrt(a*x + 1)*sqrt(-a*x + 1) - log(a)^2 + 2*(a^2*x^2*log(a) - log(a))*log(x))*log(sqrt(a*x + 1)*sqrt(-a*x + 1) + 1) + 1) + 3*(a^2*x^2*log(a)^2 - log(a)^2)*log(x))/(a^2*x^2 + (a^2*x^2 - 1)*sqrt(a*x + 1)*sqrt(-a*x + 1) - 1), x)`

**Giac [F]**

$$\int \operatorname{sech}^{-1}(ax)^3 dx = \int \operatorname{arsech}(ax)^3 dx$$

input `integrate(arcsech(a*x)^3,x, algorithm="giac")`

output `integrate(arcsech(a*x)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{sech}^{-1}(ax)^3 dx = \int \operatorname{acosh}\left(\frac{1}{ax}\right)^3 dx$$

input `int(acosh(1/(a*x))^3,x)`output `int(acosh(1/(a*x))^3, x)`**Reduce [F]**

$$\int \operatorname{sech}^{-1}(ax)^3 dx = \int \operatorname{asech}(ax)^3 dx$$

input `int(asech(a*x)^3,x)`output `int(asech(a*x)**3,x)`

### 3.15 $\int \frac{\operatorname{sech}^{-1}(ax)^3}{x} dx$

Optimal result	196
Mathematica [A] (verified)	197
Rubi [C] (verified)	197
Maple [A] (verified)	200
Fricas [F]	201
Sympy [F]	201
Maxima [F]	201
Giac [F]	202
Mupad [F(-1)]	202
Reduce [F]	202

#### Optimal result

Integrand size = 10, antiderivative size = 88

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x} dx = \frac{1}{4} \operatorname{sech}^{-1}(ax)^4 - \operatorname{sech}^{-1}(ax)^3 \log\left(1 + e^{2\operatorname{sech}^{-1}(ax)}\right) - \frac{3}{2} \operatorname{sech}^{-1}(ax)^2 \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(ax)}\right) + \frac{3}{2} \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}\left(3, -e^{2\operatorname{sech}^{-1}(ax)}\right) - \frac{3}{4} \operatorname{PolyLog}\left(4, -e^{2\operatorname{sech}^{-1}(ax)}\right)$$

output

```
1/4*arcsech(a*x)^4-arcsech(a*x)^3*ln(1+(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2))^2)-3/2*arcsech(a*x)^2*polylog(2,-(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2))^2)+3/2*arcsech(a*x)*polylog(3,-(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2))^2)-3/4*polylog(4,-(1/a/x+(-1+1/a/x)^(1/2)*(1+1/a/x)^(1/2))^2)
```

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.95

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x} dx = \frac{1}{4} \left( -\operatorname{sech}^{-1}(ax)^4 - 4\operatorname{sech}^{-1}(ax)^3 \log \left( 1 + e^{-2\operatorname{sech}^{-1}(ax)} \right) \right. \\ \left. + 6\operatorname{sech}^{-1}(ax)^2 \operatorname{PolyLog} \left( 2, -e^{-2\operatorname{sech}^{-1}(ax)} \right) \right. \\ \left. + 6\operatorname{sech}^{-1}(ax) \operatorname{PolyLog} \left( 3, -e^{-2\operatorname{sech}^{-1}(ax)} \right) \right. \\ \left. + 3 \operatorname{PolyLog} \left( 4, -e^{-2\operatorname{sech}^{-1}(ax)} \right) \right)$$

input `Integrate[ArcSech[a*x]^3/x,x]`

output `(-ArcSech[a*x]^4 - 4*ArcSech[a*x]^3*Log[1 + E^(-2*ArcSech[a*x])] + 6*ArcSech[a*x]^2*PolyLog[2, -E^(-2*ArcSech[a*x])] + 6*ArcSech[a*x]*PolyLog[3, -E^(-2*ArcSech[a*x])] + 3*PolyLog[4, -E^(-2*ArcSech[a*x])])/4`

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.57 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.20, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$ , Rules used = {6839, 3042, 26, 4201, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x} dx \\ \downarrow \text{6839} \\ - \int \sqrt{\frac{1-ax}{ax+1}} (ax+1) \operatorname{sech}^{-1}(ax)^3 d\operatorname{sech}^{-1}(ax) \\ \downarrow \text{3042} \\ - \int -i \operatorname{sech}^{-1}(ax)^3 \tan(i \operatorname{sech}^{-1}(ax)) d\operatorname{sech}^{-1}(ax)$$

↓ 26

$$i \int \operatorname{sech}^{-1}(ax)^3 \tan(i \operatorname{sech}^{-1}(ax)) d \operatorname{sech}^{-1}(ax)$$

↓ 4201

$$i \left( 2i \int \frac{e^{2 \operatorname{sech}^{-1}(ax)} \operatorname{sech}^{-1}(ax)^3}{1 + e^{2 \operatorname{sech}^{-1}(ax)}} d \operatorname{sech}^{-1}(ax) - \frac{1}{4} i \operatorname{sech}^{-1}(ax)^4 \right)$$

↓ 2620

$$i \left( 2i \left( \frac{1}{2} \operatorname{sech}^{-1}(ax)^3 \log(e^{2 \operatorname{sech}^{-1}(ax)} + 1) - \frac{3}{2} \int \operatorname{sech}^{-1}(ax)^2 \log(1 + e^{2 \operatorname{sech}^{-1}(ax)}) d \operatorname{sech}^{-1}(ax) \right) - \frac{1}{4} i \operatorname{sech}^{-1}(ax)^4 \right)$$

↓ 3011

$$i \left( 2i \left( \frac{1}{2} \operatorname{sech}^{-1}(ax)^3 \log(e^{2 \operatorname{sech}^{-1}(ax)} + 1) - \frac{3}{2} \left( \int \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}(2, -e^{2 \operatorname{sech}^{-1}(ax)}) d \operatorname{sech}^{-1}(ax) - \frac{1}{2} \operatorname{sech}^{-1}(ax)^2 \log(e^{2 \operatorname{sech}^{-1}(ax)} + 1) \right) \right) - \frac{1}{4} i \operatorname{sech}^{-1}(ax)^4 \right)$$

↓ 7163

$$i \left( 2i \left( \frac{1}{2} \operatorname{sech}^{-1}(ax)^3 \log(e^{2 \operatorname{sech}^{-1}(ax)} + 1) - \frac{3}{2} \left( -\frac{1}{2} \int \operatorname{PolyLog}(3, -e^{2 \operatorname{sech}^{-1}(ax)}) d \operatorname{sech}^{-1}(ax) - \frac{1}{2} \operatorname{sech}^{-1}(ax)^2 \log(e^{2 \operatorname{sech}^{-1}(ax)} + 1) \right) \right) - \frac{1}{4} i \operatorname{sech}^{-1}(ax)^4 \right)$$

↓ 2720

$$i \left( 2i \left( \frac{1}{2} \operatorname{sech}^{-1}(ax)^3 \log(e^{2 \operatorname{sech}^{-1}(ax)} + 1) - \frac{3}{2} \left( -\frac{1}{4} \int e^{-2 \operatorname{sech}^{-1}(ax)} \operatorname{PolyLog}(3, -e^{2 \operatorname{sech}^{-1}(ax)}) d e^{2 \operatorname{sech}^{-1}(ax)} - \frac{1}{2} \operatorname{sech}^{-1}(ax)^2 \log(e^{2 \operatorname{sech}^{-1}(ax)} + 1) \right) \right) - \frac{1}{4} i \operatorname{sech}^{-1}(ax)^4 \right)$$

↓ 7143

$$i \left( 2i \left( \frac{1}{2} \operatorname{sech}^{-1}(ax)^3 \log(e^{2 \operatorname{sech}^{-1}(ax)} + 1) - \frac{3}{2} \left( -\frac{1}{2} \operatorname{sech}^{-1}(ax)^2 \operatorname{PolyLog}(2, -e^{2 \operatorname{sech}^{-1}(ax)}) + \frac{1}{2} \operatorname{sech}^{-1}(ax) \operatorname{PolyLog}(3, -e^{2 \operatorname{sech}^{-1}(ax)}) \right) \right) - \frac{1}{4} i \operatorname{sech}^{-1}(ax)^4 \right)$$

input

```
Int[ArcSech[a*x]^3/x, x]
```

output

```
I*((-1/4*I)*ArcSech[a*x]^4 + (2*I)*((ArcSech[a*x]^3*Log[1 + E^(2*ArcSech[a*x])])/2 - (3*(-1/2*(ArcSech[a*x]^2*PolyLog[2, -E^(2*ArcSech[a*x])]) + (ArcSech[a*x]*PolyLog[3, -E^(2*ArcSech[a*x])])/2 - PolyLog[4, -E^(2*ArcSech[a*x])]/4))/2))
```

## Definitions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(F x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2620 `Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)] / ((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))* (F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
- rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4201 `Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x))), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`



rule 6839 `Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^(n_)*(x_)^(m_), x_Symbol] := Simp[-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_)^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`

## Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 181, normalized size of antiderivative = 2.06

method	result
derivativedivides	$\frac{\operatorname{arcsech}(xa)^4}{4} - \operatorname{arcsech}(xa)^3 \ln \left( 1 + \left( \frac{1}{ax} + \sqrt{\frac{1}{ax} - 1} \sqrt{1 + \frac{1}{ax}} \right)^2 \right) - \frac{3 \operatorname{arcsech}(xa)^2 \operatorname{polylog} \left( 2, -\left( \frac{1}{ax} + \sqrt{\frac{1}{ax} - 1} \sqrt{1 + \frac{1}{ax}} \right)^2 \right)}{2}$
default	$\frac{\operatorname{arcsech}(xa)^4}{4} - \operatorname{arcsech}(xa)^3 \ln \left( 1 + \left( \frac{1}{ax} + \sqrt{\frac{1}{ax} - 1} \sqrt{1 + \frac{1}{ax}} \right)^2 \right) - \frac{3 \operatorname{arcsech}(xa)^2 \operatorname{polylog} \left( 2, -\left( \frac{1}{ax} + \sqrt{\frac{1}{ax} - 1} \sqrt{1 + \frac{1}{ax}} \right)^2 \right)}{2}$

input `int(arcsech(x*a)^3/x,x,method=_RETURNVERBOSE)`

output `1/4*arcsech(x*a)^4-arcsech(x*a)^3*ln(1+(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2)-3/2*arcsech(x*a)^2*polylog(2,-(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2)+3/2*arcsech(x*a)*polylog(3,-(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2)-3/4*polylog(4,-(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2)`

**Fricas [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x} dx = \int \frac{\operatorname{arsech}(ax)^3}{x} dx$$

input `integrate(arcsech(a*x)^3/x, x, algorithm="fricas")`

output `integral(arcsech(a*x)^3/x, x)`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x} dx = \int \frac{\operatorname{asech}^3(ax)}{x} dx$$

input `integrate(asech(a*x)**3/x, x)`

output `Integral(asech(a*x)**3/x, x)`

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x} dx = \int \frac{\operatorname{arsech}(ax)^3}{x} dx$$

input `integrate(arcsech(a*x)^3/x, x, algorithm="maxima")`

output `integrate(arcsech(a*x)^3/x, x)`

**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x} dx = \int \frac{\operatorname{arsech}(ax)^3}{x} dx$$

input `integrate(arcsech(a*x)^3/x,x, algorithm="giac")`

output `integrate(arcsech(a*x)^3/x, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{ax}\right)^3}{x} dx$$

input `int(acosh(1/(a*x))^3/x,x)`

output `int(acosh(1/(a*x))^3/x, x)`

**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x} dx = \int \frac{\operatorname{asech}(ax)^3}{x} dx$$

input `int(asech(a*x)^3/x,x)`

output `int(asech(a*x)**3/x,x)`

### 3.16 $\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^2} dx$

Optimal result	203
Mathematica [A] (verified)	203
Rubi [C] (verified)	204
Maple [A] (verified)	206
Fricas [A] (verification not implemented)	207
Sympy [F]	207
Maxima [A] (verification not implemented)	208
Giac [F]	208
Mupad [F(-1)]	208
Reduce [F]	209

#### Optimal result

Integrand size = 10, antiderivative size = 83

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^2} dx = \frac{6\sqrt{\frac{1-ax}{1+ax}}(1+ax)}{x} - \frac{6\operatorname{sech}^{-1}(ax)}{x} + \frac{3\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)^2}{x} - \frac{\operatorname{sech}^{-1}(ax)^3}{x}$$

output

```
6*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)/x-6*arcsech(a*x)/x+3*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*arcsech(a*x)^2/x-arcsech(a*x)^3/x
```

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.90

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^2} dx = \frac{6\sqrt{\frac{1-ax}{1+ax}}(1+ax) - 6\operatorname{sech}^{-1}(ax) + 3\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)^2 - \operatorname{sech}^{-1}(ax)^3}{x}$$

input

```
Integrate[ArcSech[a*x]^3/x^2,x]
```

output

```
(6*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x) - 6*ArcSech[a*x] + 3*Sqrt[(1 - a*x)
/(1 + a*x)]*(1 + a*x)*ArcSech[a*x]^2 - ArcSech[a*x]^3)/x
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.44 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.39, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.200$ , Rules used = {6839, 3042, 26, 3777, 3042, 3777, 26, 3042, 26, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^{-1}(ax)^3}{x^2} dx \\
 & \quad \downarrow \text{6839} \\
 & -a \int \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^3}{ax} d\operatorname{sech}^{-1}(ax) \\
 & \quad \downarrow \text{3042} \\
 & -a \int -i\operatorname{sech}^{-1}(ax)^3 \sin(i\operatorname{sech}^{-1}(ax)) d\operatorname{sech}^{-1}(ax) \\
 & \quad \downarrow \text{26} \\
 & ia \int \operatorname{sech}^{-1}(ax)^3 \sin(i\operatorname{sech}^{-1}(ax)) d\operatorname{sech}^{-1}(ax) \\
 & \quad \downarrow \text{3777} \\
 & ia \left( \frac{i\operatorname{sech}^{-1}(ax)^3}{ax} - 3i \int \frac{\operatorname{sech}^{-1}(ax)^2}{ax} d\operatorname{sech}^{-1}(ax) \right) \\
 & \quad \downarrow \text{3042} \\
 & ia \left( \frac{i\operatorname{sech}^{-1}(ax)^3}{ax} - 3i \int \operatorname{sech}^{-1}(ax)^2 \sin\left(i\operatorname{sech}^{-1}(ax) + \frac{\pi}{2}\right) d\operatorname{sech}^{-1}(ax) \right) \\
 & \quad \downarrow \text{3777}
 \end{aligned}$$

$$ia \left( \frac{i \operatorname{sech}^{-1}(ax)^3}{ax} - 3i \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1) \operatorname{sech}^{-1}(ax)^2}{ax} - 2i \int -\frac{i \sqrt{\frac{1-ax}{ax+1}}(ax+1) \operatorname{sech}^{-1}(ax)}{ax} d \operatorname{sech}^{-1}(ax) \right) \right)$$

↓ 26

$$ia \left( \frac{i \operatorname{sech}^{-1}(ax)^3}{ax} - 3i \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1) \operatorname{sech}^{-1}(ax)^2}{ax} - 2 \int \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1) \operatorname{sech}^{-1}(ax)}{ax} d \operatorname{sech}^{-1}(ax) \right) \right)$$

↓ 3042

$$ia \left( \frac{i \operatorname{sech}^{-1}(ax)^3}{ax} - 3i \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1) \operatorname{sech}^{-1}(ax)^2}{ax} - 2 \int -i \operatorname{sech}^{-1}(ax) \sin(i \operatorname{sech}^{-1}(ax)) d \operatorname{sech}^{-1}(ax) \right) \right)$$

↓ 26

$$ia \left( \frac{i \operatorname{sech}^{-1}(ax)^3}{ax} - 3i \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1) \operatorname{sech}^{-1}(ax)^2}{ax} + 2i \int \operatorname{sech}^{-1}(ax) \sin(i \operatorname{sech}^{-1}(ax)) d \operatorname{sech}^{-1}(ax) \right) \right)$$

↓ 3777

$$ia \left( \frac{i \operatorname{sech}^{-1}(ax)^3}{ax} - 3i \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1) \operatorname{sech}^{-1}(ax)^2}{ax} + 2i \left( \frac{i \operatorname{sech}^{-1}(ax)}{ax} - i \int \frac{1}{ax} d \operatorname{sech}^{-1}(ax) \right) \right) \right)$$

↓ 3042

$$ia \left( \frac{i \operatorname{sech}^{-1}(ax)^3}{ax} - 3i \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1) \operatorname{sech}^{-1}(ax)^2}{ax} + 2i \left( \frac{i \operatorname{sech}^{-1}(ax)}{ax} - i \int \sin\left(i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2}\right) d \operatorname{sech}^{-1}(ax) \right) \right) \right)$$

↓ 3117

$$ia \left( \frac{i \operatorname{sech}^{-1}(ax)^3}{ax} - 3i \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1) \operatorname{sech}^{-1}(ax)^2}{ax} + 2i \left( \frac{i \operatorname{sech}^{-1}(ax)}{ax} - \frac{i \sqrt{\frac{1-ax}{ax+1}}(ax+1)}{ax} \right) \right) \right)$$

input `Int[ArcSech[a*x]^3/x^2,x]`

output

```
I*a*((I*ArcSech[a*x]^3)/(a*x) - (3*I)*((Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)
)*ArcSech[a*x]^2)/(a*x) + (2*I)*((-I)*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)
)/(a*x) + (I*ArcSech[a*x])/(a*x)))
```

### Defintions of rubi rules used

rule 26

```
Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3117

```
Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]
```

rule 3777

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(
-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*C
os[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

rule 6839

```
Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))^(n_)*(x_)^(m_.), x_Symbol] := Simp[
-(c^(m + 1))^( -1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

### Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.18

method	result
derivativedivides	$a \left( -\frac{\operatorname{arcsech}(xa)^3}{xa} + 3\sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} \operatorname{arcsech}(xa)^2 - \frac{6 \operatorname{arcsech}(xa)}{xa} + 6\sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} \right)$
default	$a \left( -\frac{\operatorname{arcsech}(xa)^3}{xa} + 3\sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} \operatorname{arcsech}(xa)^2 - \frac{6 \operatorname{arcsech}(xa)}{xa} + 6\sqrt{-\frac{xa-1}{xa}} \sqrt{\frac{xa+1}{xa}} \right)$

input `int(arcsech(x*a)^3/x^2,x,method=_RETURNVERBOSE)`

output `a*(-1/x/a*arcsech(x*a)^3+3*(-(a*x-1)/x/a)^(1/2)*((a*x+1)/x/a)^(1/2)*arcsech(x*a)^2-6/x/a*arcsech(x*a)+6*(-(a*x-1)/x/a)^(1/2)*((a*x+1)/x/a)^(1/2))`

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.87

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^2} dx$$

$$= \frac{3ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}} \log\left(\frac{ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}}+1}{ax}\right)^2 - \log\left(\frac{ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}}+1}{ax}\right)^3 + 6ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}} - 6\log\left(\frac{ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}}+1}{ax}\right)}{x}$$

input `integrate(arcsech(a*x)^3/x^2,x, algorithm="fricas")`

output `(3*a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2))*log((a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)) + 1)/(a*x))^2 - log((a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)) + 1)/(a*x))^3 + 6*a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)) - 6*log((a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)) + 1)/(a*x)))/x`

### Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^2} dx = \int \frac{\operatorname{asech}^3(ax)}{x^2} dx$$

input `integrate(asech(a*x)**3/x**2,x)`

output `Integral(asech(a*x)**3/x**2, x)`



**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.66

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^2} dx = 3a\sqrt{\frac{1}{a^2x^2} - 1} \operatorname{ar} \operatorname{sech}(ax)^2 - \frac{\operatorname{ar} \operatorname{sech}(ax)^3}{x} + 6a\sqrt{\frac{1}{a^2x^2} - 1} - \frac{6 \operatorname{ar} \operatorname{sech}(ax)}{x}$$

input `integrate(arcsech(a*x)^3/x^2,x, algorithm="maxima")`output `3*a*sqrt(1/(a^2*x^2) - 1)*arcsech(a*x)^2 - arcsech(a*x)^3/x + 6*a*sqrt(1/(a^2*x^2) - 1) - 6*arcsech(a*x)/x`**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^2} dx = \int \frac{\operatorname{ar} \operatorname{sech}(ax)^3}{x^2} dx$$

input `integrate(arcsech(a*x)^3/x^2,x, algorithm="giac")`output `integrate(arcsech(a*x)^3/x^2, x)`**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^2} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{ax}\right)^3}{x^2} dx$$

input `int(acosh(1/(a*x))^3/x^2,x)`output `int(acosh(1/(a*x))^3/x^2, x)`

**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^2} dx = \int \frac{a \operatorname{sech}(ax)^3}{x^2} dx$$

input `int(asech(a*x)^3/x^2,x)`

output `int(asech(a*x)**3/x**2,x)`

### 3.17 $\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^3} dx$

Optimal result	210
Mathematica [A] (verified)	211
Rubi [A] (verified)	211
Maple [A] (verified)	214
Fricas [A] (verification not implemented)	215
Sympy [F]	215
Maxima [F]	215
Giac [F]	216
Mupad [F(-1)]	216
Reduce [F]	216

#### Optimal result

Integrand size = 10, antiderivative size = 137

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^3} dx = \frac{3\sqrt{\frac{1-ax}{1+ax}}(1+ax)}{8x^2} - \frac{3}{8}a^2\operatorname{sech}^{-1}(ax) - \frac{3(1-ax)(1+ax)\operatorname{sech}^{-1}(ax)}{4x^2} + \frac{3\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)^2}{4x^2} - \frac{1}{4}a^2\operatorname{sech}^{-1}(ax)^3 - \frac{(1-ax)(1+ax)\operatorname{sech}^{-1}(ax)^3}{2x^2}$$

output

```
3/8*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)/x^2-3/8*a^2*arcsech(a*x)-3/4*(-a*x+1)
*(a*x+1)*arcsech(a*x)/x^2+3/4*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)*arcsech(a*x)
)^2/x^2-1/4*a^2*arcsech(a*x)^3-1/2*(-a*x+1)*(a*x+1)*arcsech(a*x)^3/x^2
```

**Mathematica [A] (verified)**

Time = 0.10 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.07

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^3} dx$$

$$= \frac{3\sqrt{\frac{1-ax}{1+ax}}(1+ax) - 6\operatorname{sech}^{-1}(ax) + 6\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)^2 + 2(-2 + a^2x^2)\operatorname{sech}^{-1}(ax)^3 - 3a^2x^2 \log}{8x^2}$$

input `Integrate[ArcSech[a*x]^3/x^3,x]`

output

```
(3*sqrt((1 - a*x)/(1 + a*x))*(1 + a*x) - 6*ArcSech[a*x] + 6*sqrt((1 - a*x)
/(1 + a*x))*(1 + a*x)*ArcSech[a*x]^2 + 2*(-2 + a^2*x^2)*ArcSech[a*x]^3 - 3
*a^2*x^2*Log[x] + 3*a^2*x^2*Log[1 + Sqrt[(1 - a*x)/(1 + a*x)] + a*x*sqrt[(
1 - a*x)/(1 + a*x)]])/(8*x^2)
```

**Rubi [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.15, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.100$ , Rules used = {6839, 5895, 3042, 25, 3792, 15, 25, 3042, 25, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^3} dx$$

$$\downarrow \text{6839}$$

$$-a^2 \int \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^3}{a^2x^2} d\operatorname{sech}^{-1}(ax)$$

$$\downarrow \text{5895}$$

$$-a^2 \left( \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)^3}{2a^2x^2} - \frac{3}{2} \int \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)^2}{a^2x^2} d\operatorname{sech}^{-1}(ax) \right)$$

$$\downarrow \text{3042}$$

$$-a^2 \left( \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)^3}{2a^2x^2} - \frac{3}{2} \int -\operatorname{sech}^{-1}(ax)^2 \sin(i\operatorname{sech}^{-1}(ax))^2 d\operatorname{sech}^{-1}(ax) \right)$$

↓ 25

$$-a^2 \left( \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)^3}{2a^2x^2} + \frac{3}{2} \int \operatorname{sech}^{-1}(ax)^2 \sin(i\operatorname{sech}^{-1}(ax))^2 d\operatorname{sech}^{-1}(ax) \right)$$

↓ 3792

$$-a^2 \left( \frac{3}{2} \left( \frac{1}{2} \int -\frac{(1-ax)(ax+1)}{a^2x^2} d\operatorname{sech}^{-1}(ax) + \frac{1}{2} \int \operatorname{sech}^{-1}(ax)^2 d\operatorname{sech}^{-1}(ax) - \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{2a^2x^2} + \right. \right.$$

↓ 15

$$\left. -a^2 \left( \frac{3}{2} \left( \frac{1}{2} \int -\frac{(1-ax)(ax+1)}{a^2x^2} d\operatorname{sech}^{-1}(ax) - \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{2a^2x^2} + \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)}{2a^2x^2} + \right. \right. \right.$$

↓ 25

$$\left. -a^2 \left( \frac{3}{2} \left( -\frac{1}{2} \int \frac{(1-ax)(ax+1)}{a^2x^2} d\operatorname{sech}^{-1}(ax) - \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{2a^2x^2} + \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)}{2a^2x^2} + \right. \right. \right.$$

↓ 3042

$$-a^2 \left( \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)^3}{2a^2x^2} + \frac{3}{2} \left( -\frac{1}{2} \int -\sin(i\operatorname{sech}^{-1}(ax))^2 d\operatorname{sech}^{-1}(ax) - \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{2a^2x^2} \right. \right.$$

↓ 25

$$-a^2 \left( \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)^3}{2a^2x^2} + \frac{3}{2} \left( \frac{1}{2} \int \sin(i\operatorname{sech}^{-1}(ax))^2 d\operatorname{sech}^{-1}(ax) - \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{2a^2x^2} + \right. \right.$$

↓ 3115

$$-a^2 \left( \frac{3}{2} \left( \frac{1}{2} \left( \frac{1}{2} \int 1 d\operatorname{sech}^{-1}(ax) - \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)}{2a^2x^2} \right) - \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{2a^2x^2} + \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)}{2a^2x^2} \right. \right.$$

↓ 24

$$-a^2 \left( \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)^3}{2a^2x^2} + \frac{3}{2} \left( -\frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{2a^2x^2} + \frac{(1-ax)(ax+1)\operatorname{sech}^{-1}(ax)}{2a^2x^2} + \frac{1}{2} \left( \frac{1}{2} \right) \right) \right)$$

input `Int[ArcSech[a*x]^3/x^3,x]`

output `-(a^2*((1 - a*x)*(1 + a*x)*ArcSech[a*x]^3)/(2*a^2*x^2) + (3*((-1/2*(Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)))/(a^2*x^2) + ArcSech[a*x]/2)/2 + ((1 - a*x)*(1 + a*x)*ArcSech[a*x])/(2*a^2*x^2) - (Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x]^2)/(2*a^2*x^2) + ArcSech[a*x]^3/6))/2)`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3792

```
Int[((c_.) + (d_.)*(x_)^(m_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol]
:= Simp[d*m*(c + d*x)^(m - 1)*((b*Sine + f*x)^(n)/(f^2*n^2)), x] + (-Simp
p[b*(c + d*x)^m*cos[e + f*x]*((b*Sine + f*x)^(n - 1)/(f*n)), x] + Simp[b^
2*((n - 1)/n) Int[(c + d*x)^m*(b*Sine + f*x)^(n - 2), x], x] - Simp[d^2
*m*((m - 1)/(f^2*n^2)) Int[(c + d*x)^(m - 2)*(b*Sine + f*x)^n, x], x])
/; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 1]
```

rule 5895

```
Int[Cosh[(a_.) + (b_.)*(x_)^(n_.)]*(x_)^(m_.)*Sinh[(a_.) + (b_.)*(x_)^(n_.)
]^(p_.), x_Symbol] := Simp[x^(m - n + 1)*(Sinh[a + b*x^n]^(p + 1)/(b*n*(p +
1))), x] - Simp[(m - n + 1)/(b*n*(p + 1)) Int[x^(m - n)*Sinh[a + b*x^n]^(
p + 1), x], x] /; FreeQ[{a, b, p}, x] && LtQ[0, n, m + 1] && NeQ[p, -1]
```

rule 6839

```
Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))^(n_)*(x_)^(m_), x_Symbol] := Simp[
-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

## Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.42

method	result
derivativedivides	$a^2 \left( -\frac{\cosh(2 \operatorname{arcsech}(xa)) \operatorname{arcsech}(xa)^3}{4} + \frac{3 \sinh(2 \operatorname{arcsech}(xa)) \operatorname{arcsech}(xa)^2}{8} - \frac{3 \cosh(2 \operatorname{arcsech}(xa)) \operatorname{arcsech}(xa)}{8} \right)$
default	$a^2 \left( -\frac{\cosh(2 \operatorname{arcsech}(xa)) \operatorname{arcsech}(xa)^3}{4} + \frac{3 \sinh(2 \operatorname{arcsech}(xa)) \operatorname{arcsech}(xa)^2}{8} - \frac{3 \cosh(2 \operatorname{arcsech}(xa)) \operatorname{arcsech}(xa)}{8} \right)$

input

```
int(arcsech(x*a)^3/x^3,x,method=_RETURNVERBOSE)
```

output

```
a^2*(-1/4*cosh(2*arcsech(x*a))*arcsech(x*a)^3+3/8*sinh(2*arcsech(x*a))*arc
sech(x*a)^2-3/8*cosh(2*arcsech(x*a))*arcsech(x*a)+3/16*sinh(2*arcsech(x*a)
))
```

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.27

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^3} dx$$

$$= \frac{6ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}} \log\left(\frac{ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}}+1}{ax}\right)^2 + 2(a^2x^2-2) \log\left(\frac{ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}}+1}{ax}\right)^3 + 3ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}} + 3(a^2x^2-2) \log\left(\frac{ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}}+1}{ax}\right)}{8x^2}$$

input `integrate(arcsech(a*x)^3/x^3,x, algorithm="fricas")`output `1/8*(6*a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2))*log((a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)) + 1)/(a*x))^2 + 2*(a^2*x^2 - 2)*log((a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)) + 1)/(a*x))^3 + 3*a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)) + 3*(a^2*x^2 - 2)*log((a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)) + 1)/(a*x)))/x^2`**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^3} dx = \int \frac{\operatorname{arsech}^3(ax)}{x^3} dx$$

input `integrate(asech(a*x)**3/x**3,x)`output `Integral(asech(a*x)**3/x**3, x)`**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^3} dx = \int \frac{\operatorname{arsech}(ax)^3}{x^3} dx$$

input `integrate(arcsech(a*x)^3/x^3,x, algorithm="maxima")`



output `integrate(arcsech(a*x)^3/x^3, x)`

### Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^3} dx = \int \frac{\operatorname{arsech}(ax)^3}{x^3} dx$$

input `integrate(arcsech(a*x)^3/x^3,x, algorithm="giac")`

output `integrate(arcsech(a*x)^3/x^3, x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^3} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{ax}\right)^3}{x^3} dx$$

input `int(acosh(1/(a*x))^3/x^3,x)`

output `int(acosh(1/(a*x))^3/x^3, x)`

### Reduce [F]

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^3} dx = \int \frac{\operatorname{asech}(ax)^3}{x^3} dx$$

input `int(asech(a*x)^3/x^3,x)`

output `int(asech(a*x)**3/x**3,x)`

### 3.18 $\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^4} dx$

Optimal result	217
Mathematica [A] (verified)	218
Rubi [C] (verified)	218
Maple [A] (verified)	222
Fricas [A] (verification not implemented)	222
Sympy [F]	223
Maxima [F]	223
Giac [F]	224
Mupad [F(-1)]	224
Reduce [F]	224

#### Optimal result

Integrand size = 10, antiderivative size = 179

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^4} dx = \frac{14a^2 \sqrt{\frac{1-ax}{1+ax}}(1+ax)}{9x} + \frac{2\left(\frac{1-ax}{1+ax}\right)^{3/2}(1+ax)^3}{27x^3} - \frac{2\operatorname{sech}^{-1}(ax)}{9x^3}$$

$$- \frac{4a^2 \operatorname{sech}^{-1}(ax)}{3x} + \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)^2}{3x^3}$$

$$+ \frac{2a^2 \sqrt{\frac{1-ax}{1+ax}}(1+ax)\operatorname{sech}^{-1}(ax)^2}{3x} - \frac{\operatorname{sech}^{-1}(ax)^3}{3x^3}$$

output

```
14/9*a^2*((-a*x+1)/(a*x+1))^(1/2)*(a*x+1)/x+2/27*((-a*x+1)/(a*x+1))^(3/2)*
(a*x+1)^3/x^3-2/9*arcsech(a*x)/x^3-4/3*a^2*arcsech(a*x)/x+1/3*((-a*x+1)/(a
*x+1))^(1/2)*(a*x+1)*arcsech(a*x)^2/x^3+2/3*a^2*((-a*x+1)/(a*x+1))^(1/2)*
(a*x+1)*arcsech(a*x)^2/x-1/3*arcsech(a*x)^3/x^3
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.67

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^4} dx$$

$$= \frac{2\sqrt{\frac{1-ax}{1+ax}}(1+ax+20a^2x^2+20a^3x^3) - 6(1+6a^2x^2)\operatorname{sech}^{-1}(ax) + 9\sqrt{\frac{1-ax}{1+ax}}(1+ax+2a^2x^2+2a^3x^3)\operatorname{sech}^{-1}(ax)}{27x^3}$$

input

```
Integrate[ArcSech[a*x]^3/x^4,x]
```

output

```
(2*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x + 20*a^2*x^2 + 20*a^3*x^3) - 6*(1 + 6*a^2*x^2)*ArcSech[a*x] + 9*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x + 2*a^2*x^2 + 2*a^3*x^3)*ArcSech[a*x]^2 - 9*ArcSech[a*x]^3)/(27*x^3)
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.65 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.37, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.400$ , Rules used = {6839, 5896, 3042, 3792, 3042, 3113, 2009, 3777, 26, 3042, 26, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^4} dx$$

$$\downarrow \text{6839}$$

$$-a^3 \int \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^3}{a^3x^3} d\operatorname{sech}^{-1}(ax)$$

$$\downarrow \text{5896}$$

$$-a^3 \left( \frac{\operatorname{sech}^{-1}(ax)^3}{3a^3x^3} - \int \frac{\operatorname{sech}^{-1}(ax)^2}{a^3x^3} d\operatorname{sech}^{-1}(ax) \right)$$

$$\downarrow \text{3042}$$

$$-a^3 \left( \frac{\operatorname{sech}^{-1}(ax)^3}{3a^3x^3} - \int \operatorname{sech}^{-1}(ax)^2 \sin \left( i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2} \right)^3 d\operatorname{sech}^{-1}(ax) \right)$$

↓ 3792

$$-a^3 \left( -\frac{2}{9} \int \frac{1}{a^3x^3} d\operatorname{sech}^{-1}(ax) - \frac{2}{3} \int \frac{\operatorname{sech}^{-1}(ax)^2}{ax} d\operatorname{sech}^{-1}(ax) + \frac{\operatorname{sech}^{-1}(ax)^3}{3a^3x^3} - \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{3a^3x^3} + \dots \right)$$

↓ 3042

$$-a^3 \left( -\frac{2}{3} \int \operatorname{sech}^{-1}(ax)^2 \sin \left( i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2} \right) d\operatorname{sech}^{-1}(ax) - \frac{2}{9} \int \sin \left( i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2} \right)^3 d\operatorname{sech}^{-1}(ax) + \frac{\operatorname{sech}^{-1}(ax)^3}{3a^3x^3} - \dots \right)$$

↓ 3113

$$-a^3 \left( -\frac{2}{9}i \int \left( \frac{(1-ax)(ax+1)}{a^2x^2} + 1 \right) d \left( -\frac{i\sqrt{\frac{1-ax}{ax+1}}(ax+1)}{ax} \right) - \frac{2}{3} \int \operatorname{sech}^{-1}(ax)^2 \sin \left( i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2} \right) d\operatorname{sech}^{-1}(ax) + \dots \right)$$

↓ 2009

$$-a^3 \left( -\frac{2}{3} \int \operatorname{sech}^{-1}(ax)^2 \sin \left( i \operatorname{sech}^{-1}(ax) + \frac{\pi}{2} \right) d\operatorname{sech}^{-1}(ax) - \frac{2}{9}i \left( -\frac{i\left(\frac{1-ax}{ax+1}\right)^{3/2}(ax+1)^3}{3a^3x^3} - \frac{i\sqrt{\frac{1-ax}{ax+1}}(ax+1)}{ax} \right) + \dots \right)$$

↓ 3777

$$-a^3 \left( -\frac{2}{3} \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{ax} - 2i \int -\frac{i\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{ax} d\operatorname{sech}^{-1}(ax) \right) - \frac{2}{9}i \left( -\frac{i\left(\frac{1-ax}{ax+1}\right)^{3/2}(ax+1)^3}{3a^3x^3} - \frac{i\sqrt{\frac{1-ax}{ax+1}}(ax+1)}{ax} \right) + \dots \right)$$

↓ 26

$$-a^3 \left( -\frac{2}{3} \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{ax} - 2 \int \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)}{ax} d\operatorname{sech}^{-1}(ax) \right) - \frac{2}{9}i \left( -\frac{i\left(\frac{1-ax}{ax+1}\right)^{3/2}(ax+1)^3}{3a^3x^3} - \frac{i\sqrt{\frac{1-ax}{ax+1}}(ax+1)}{ax} \right) + \dots \right)$$

↓ 3042

$$-a^3 \left( -\frac{2}{3} \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{ax} - 2 \int -i\operatorname{sech}^{-1}(ax) \sin(i\operatorname{sech}^{-1}(ax)) d\operatorname{sech}^{-1}(ax) \right) - \frac{2}{9}i \left( -\frac{i\left(\frac{1-ax}{ax+1}\right)^{3/2}}{3a^3x^3} \right) \right)$$

↓ 26

$$-a^3 \left( -\frac{2}{3} \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{ax} + 2i \int \operatorname{sech}^{-1}(ax) \sin(i\operatorname{sech}^{-1}(ax)) d\operatorname{sech}^{-1}(ax) \right) - \frac{2}{9}i \left( -\frac{i\left(\frac{1-ax}{ax+1}\right)^{3/2}}{3a^3x^3} \right) \right)$$

↓ 3777

$$-a^3 \left( -\frac{2}{3} \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{ax} + 2i \left( \frac{i\operatorname{sech}^{-1}(ax)}{ax} - i \int \frac{1}{ax} d\operatorname{sech}^{-1}(ax) \right) \right) - \frac{2}{9}i \left( -\frac{i\left(\frac{1-ax}{ax+1}\right)^{3/2}(ax+1)}{3a^3x^3} \right) \right)$$

↓ 3042

$$-a^3 \left( -\frac{2}{3} \left( \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{ax} + 2i \left( \frac{i\operatorname{sech}^{-1}(ax)}{ax} - i \int \sin\left(i\operatorname{sech}^{-1}(ax) + \frac{\pi}{2}\right) d\operatorname{sech}^{-1}(ax) \right) \right) - \frac{2}{9}i \left( -\frac{i\left(\frac{1-ax}{ax+1}\right)^{3/2}(ax+1)}{3a^3x^3} \right) \right)$$

↓ 3117

$$-a^3 \left( -\frac{2}{9}i \left( -\frac{i\left(\frac{1-ax}{ax+1}\right)^{3/2}(ax+1)^3}{3a^3x^3} - \frac{i\sqrt{\frac{1-ax}{ax+1}}(ax+1)}{ax} \right) + \frac{\operatorname{sech}^{-1}(ax)^3}{3a^3x^3} - \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)\operatorname{sech}^{-1}(ax)^2}{3a^3x^3} + \frac{2\operatorname{sech}^{-1}(ax)}{9} \right)$$

input `Int[ArcSech[a*x]^3/x^4,x]`

output `-(a^3*(((2*I)/9)*(((I)*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x))/(a*x) - ((I/3)*((1 - a*x)/(1 + a*x))^(3/2)*(1 + a*x)^3)/(a^3*x^3)) + (2*ArcSech[a*x])/(9*a^3*x^3) - (Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x]^2)/(3*a^3*x^3) + ArcSech[a*x]^3/(3*a^3*x^3) - (2*((Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)*ArcSech[a*x]^2)/(a*x) + (2*I)*(((I)*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)))/(a*x) + (I*ArcSech[a*x])/(a*x))))/3)`

## Definitions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 2009  $\text{Int}[u, x\_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$
- rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3113  $\text{Int}[\sin[(c.) + (d.)*(x_)]^{(n_)}, x\_Symbol] \rightarrow \text{Simp}[-d^{(-1)} \text{Subst}[\text{Int}[\text{Exp} \text{and}[(1 - x^2)^{(n-1)/2}], x], x], x, \text{Cos}[c + d*x]], x] /; \text{FreeQ}\{c, d\}, x] \ \&\& \ \text{IGtQ}[(n-1)/2, 0]$
- rule 3117  $\text{Int}[\sin[\text{Pi}/2 + (c.) + (d.)*(x_)], x\_Symbol] \rightarrow \text{Simp}[\text{Sin}[c + d*x]/d, x] /; \text{FreeQ}\{c, d\}, x]$
- rule 3777  $\text{Int}[(c.) + (d.)*(x_)]^{(m.)*\sin[(e.) + (f.)*(x_)], x\_Symbol] \rightarrow \text{Simp}[(c + d*x)^m * (\text{Cos}[e + f*x]/f), x] + \text{Simp}[d*(m/f) \text{Int}[(c + d*x)^{(m-1)} * \text{Cos}[e + f*x], x], x] /; \text{FreeQ}\{c, d, e, f\}, x] \ \&\& \ \text{GtQ}[m, 0]$
- rule 3792  $\text{Int}[(c.) + (d.)*(x_)]^{(m.)*((b.)*\sin[(e.) + (f.)*(x_)]^{(n_)}), x\_Symbol] \rightarrow \text{Simp}[d*m*(c + d*x)^{(m-1)}*((b*\text{Sin}[e + f*x])^n/(f^2*n^2)), x] + (-\text{Simp}[b*(c + d*x)^m*\text{Cos}[e + f*x]*((b*\text{Sin}[e + f*x])^{(n-1)})/(f*n), x] + \text{Simp}[b^2*((n-1)/n) \text{Int}[(c + d*x)^m*(b*\text{Sin}[e + f*x])^{(n-2)}, x], x] - \text{Simp}[d^2*m*((m-1)/(f^2*n^2)) \text{Int}[(c + d*x)^{(m-2)}*(b*\text{Sin}[e + f*x])^n, x], x]) /; \text{FreeQ}\{b, c, d, e, f\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{GtQ}[m, 1]$
- rule 5896  $\text{Int}[\text{Cosh}[(a.) + (b.)*(x_)]^{(n.)*\sinh[(a.) + (b.)*(x_)]^{(p.)*x_)]^{(m.)*x_}, x\_Symbol] \rightarrow \text{Simp}[x^{(m-n+1)}*(\text{Cosh}[a + b*x^n]^{(p+1)})/(b*n*(p+1)), x] - \text{Simp}[(m-n+1)/(b*n*(p+1)) \text{Int}[x^{(m-n)}*\text{Cosh}[a + b*x^n]^{(p+1)}, x], x] /; \text{FreeQ}\{a, b, p\}, x] \ \&\& \ \text{LtQ}[0, n, m+1] \ \&\& \ \text{NeQ}[p, -1]$

rule 6839

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))^(n_)*(x_)^(m_.), x_Symbol] := Simp[
-(c^(m + 1))^( -1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

**Maple [A] (verified)**

Time = 0.46 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.07

method	result
derivativedivides	$a^3 \left( -\frac{\operatorname{arcsech}(xa)^3}{3x^3a^3} + \frac{2\sqrt{-\frac{xa-1}{xa}}\sqrt{\frac{xa+1}{xa}}\operatorname{arcsech}(xa)^2}{3} + \frac{\sqrt{-\frac{xa-1}{xa}}\sqrt{\frac{xa+1}{xa}}\operatorname{arcsech}(xa)^2}{3x^2a^2} - \frac{4\operatorname{arcsech}(xa)}{3xa} + \dots \right)$
default	$a^3 \left( -\frac{\operatorname{arcsech}(xa)^3}{3x^3a^3} + \frac{2\sqrt{-\frac{xa-1}{xa}}\sqrt{\frac{xa+1}{xa}}\operatorname{arcsech}(xa)^2}{3} + \frac{\sqrt{-\frac{xa-1}{xa}}\sqrt{\frac{xa+1}{xa}}\operatorname{arcsech}(xa)^2}{3x^2a^2} - \frac{4\operatorname{arcsech}(xa)}{3xa} + \dots \right)$

input

```
int(arcsech(x*a)^3/x^4,x,method=_RETURNVERBOSE)
```

output

```
a^3*(-1/3/x^3/a^3*arcsech(x*a)^3+2/3*(-(a*x-1)/x/a)^(1/2)*((a*x+1)/x/a)^(1
/2)*arcsech(x*a)^2+1/3*(-(a*x-1)/x/a)^(1/2)*((a*x+1)/x/a)^(1/2)/x^2/a^2*ar
csech(x*a)^2-4/3/x/a*arcsech(x*a)+40/27*(-(a*x-1)/x/a)^(1/2)*((a*x+1)/x/a)
^(1/2)-2/9*arcsech(x*a)/x^3/a^3+2/27*(-(a*x-1)/x/a)^(1/2)*((a*x+1)/x/a)^(1
/2)/x^2/a^2)
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.04

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^4} dx$$

$$= \frac{9(2a^3x^3 + ax)\sqrt{-\frac{a^2x^2-1}{a^2x^2}} \log\left(\frac{ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}}+1}{ax}\right)^2 - 9 \log\left(\frac{ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}}+1}{ax}\right)^3 - 6(6a^2x^2 + 1) \log\left(\frac{ax\sqrt{-\frac{a^2x^2-1}{a^2x^2}}}{ax}\right)}{27x^3}$$

input

```
integrate(arcsech(a*x)^3/x^4,x, algorithm="fricas")
```

output

```
1/27*(9*(2*a^3*x^3 + a*x)*sqrt(-(a^2*x^2 - 1)/(a^2*x^2))*log((a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)) + 1)/(a*x))^2 - 9*log((a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)) + 1)/(a*x))^3 - 6*(6*a^2*x^2 + 1)*log((a*x*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)) + 1)/(a*x)) + 2*(20*a^3*x^3 + a*x)*sqrt(-(a^2*x^2 - 1)/(a^2*x^2)))/x^3
```

**Sympy [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^4} dx = \int \frac{\operatorname{arsech}^3(ax)}{x^4} dx$$

input

```
integrate(asech(a*x)**3/x**4,x)
```

output

```
Integral(asech(a*x)**3/x**4, x)
```

**Maxima [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^4} dx = \int \frac{\operatorname{arosech}(ax)^3}{x^4} dx$$

input

```
integrate(arcsech(a*x)^3/x^4,x, algorithm="maxima")
```

output

```
integrate(arcsech(a*x)^3/x^4, x)
```



**Giac [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^4} dx = \int \frac{\operatorname{arsech}(ax)^3}{x^4} dx$$

input `integrate(arcsech(a*x)^3/x^4,x, algorithm="giac")`

output `integrate(arcsech(a*x)^3/x^4, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^4} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{ax}\right)^3}{x^4} dx$$

input `int(acosh(1/(a*x))^3/x^4,x)`

output `int(acosh(1/(a*x))^3/x^4, x)`

**Reduce [F]**

$$\int \frac{\operatorname{sech}^{-1}(ax)^3}{x^4} dx = \int \frac{\operatorname{asech}(ax)^3}{x^4} dx$$

input `int(asech(a*x)^3/x^4,x)`

output `int(asech(a*x)**3/x**4,x)`

### 3.19 $\int x^6 (a + b \operatorname{sech}^{-1}(cx)) dx$

Optimal result	225
Mathematica [C] (verified)	226
Rubi [A] (warning: unable to verify)	226
Maple [A] (verified)	230
Fricas [A] (verification not implemented)	230
Sympy [F]	231
Maxima [A] (verification not implemented)	231
Giac [F]	232
Mupad [F(-1)]	232
Reduce [F]	232

#### Optimal result

Integrand size = 12, antiderivative size = 165

$$\int x^6 (a + b \operatorname{sech}^{-1}(cx)) dx = -\frac{5bx \sqrt{\frac{1-cx}{1+cx}} (1+cx)}{112c^6} - \frac{5bx^3 \sqrt{\frac{1-cx}{1+cx}} (1+cx)}{168c^4}$$

$$- \frac{bx^5 \sqrt{\frac{1-cx}{1+cx}} (1+cx)}{42c^2} + \frac{1}{7} x^7 (a + b \operatorname{sech}^{-1}(cx))$$

$$+ \frac{5b \sqrt{\frac{1-cx}{1+cx}} (1+cx) \arcsin(cx)}{112c^7 \sqrt{1-c^2x^2}}$$

output

```
-5/112*b*x*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/c^6-5/168*b*x^3*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/c^4-1/42*b*x^5*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/c^2+1/7*x^7*(a+b*arcsech(c*x))+5/112*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*arcsin(c*x)/c^7/(-c^2*x^2+1)^(1/2)
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.87

$$\int x^6(a + b\operatorname{sech}^{-1}(cx)) dx = \frac{ax^7}{7} + b\sqrt{\frac{1-cx}{1+cx}} \left( -\frac{5x}{112c^6} - \frac{5x^2}{112c^5} - \frac{5x^3}{168c^4} - \frac{5x^4}{168c^3} - \frac{x^5}{42c^2} - \frac{x^6}{42c} \right) + \frac{1}{7}bx^7\operatorname{sech}^{-1}(cx) + \frac{5ib \log\left(-2icx + 2\sqrt{\frac{1-cx}{1+cx}}(1+cx)\right)}{112c^7}$$

input `Integrate[x^6*(a + b*ArcSech[c*x]),x]`

output `(a*x^7)/7 + b*Sqrt[(1 - c*x)/(1 + c*x)]*((-5*x)/(112*c^6) - (5*x^2)/(112*c^5) - (5*x^3)/(168*c^4) - (5*x^4)/(168*c^3) - x^5/(42*c^2) - x^6/(42*c)) + (b*x^7*ArcSech[c*x])/7 + (((5*I)/112)*b*Log[(-2*I)*c*x + 2*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)])/c^7`

**Rubi [A] (warning: unable to verify)**

Time = 0.30 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.93, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {6837, 111, 27, 111, 27, 101, 25, 39, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^6(a + b\operatorname{sech}^{-1}(cx)) dx$$

$$\downarrow 6837$$

$$\frac{1}{7}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^6}{\sqrt{1-cx}\sqrt{cx+1}} dx + \frac{1}{7}x^7(a + b\operatorname{sech}^{-1}(cx))$$

$$\downarrow 111$$

$$\frac{1}{7}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{\int-\frac{5x^4}{\sqrt{1-cx}\sqrt{cx+1}}dx}{6c^2}-\frac{x^5\sqrt{1-cx}\sqrt{cx+1}}{6c^2}\right)+\frac{1}{7}x^7(a+b\operatorname{sech}^{-1}(cx))$$

↓ 27

$$\frac{1}{7}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{5\int\frac{x^4}{\sqrt{1-cx}\sqrt{cx+1}}dx}{6c^2}-\frac{x^5\sqrt{1-cx}\sqrt{cx+1}}{6c^2}\right)+\frac{1}{7}x^7(a+b\operatorname{sech}^{-1}(cx))$$

↓ 111

$$\frac{1}{7}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{5\left(-\frac{\int-\frac{3x^2}{\sqrt{1-cx}\sqrt{cx+1}}dx}{4c^2}-\frac{x^3\sqrt{1-cx}\sqrt{cx+1}}{4c^2}\right)}{6c^2}-\frac{x^5\sqrt{1-cx}\sqrt{cx+1}}{6c^2}\right)+\frac{1}{7}x^7(a+b\operatorname{sech}^{-1}(cx))$$

↓ 27

$$\frac{1}{7}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{5\left(\frac{3\int\frac{x^2}{\sqrt{1-cx}\sqrt{cx+1}}dx}{4c^2}-\frac{x^3\sqrt{1-cx}\sqrt{cx+1}}{4c^2}\right)}{6c^2}-\frac{x^5\sqrt{1-cx}\sqrt{cx+1}}{6c^2}\right)+\frac{1}{7}x^7(a+b\operatorname{sech}^{-1}(cx))$$

↓ 101

$$\frac{1}{7}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{5\left(3\left(\frac{\int-\frac{1}{\sqrt{1-cx}\sqrt{cx+1}}dx}{2c^2}-\frac{x\sqrt{1-cx}\sqrt{cx+1}}{2c^2}\right)-\frac{x^3\sqrt{1-cx}\sqrt{cx+1}}{4c^2}\right)}{6c^2}-\frac{x^5\sqrt{1-cx}\sqrt{cx+1}}{6c^2}\right)+\frac{1}{7}x^7(a+b\operatorname{sech}^{-1}(cx))$$

↓ 25

$$\frac{1}{7}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{5\left(\frac{3\left(\frac{\int\frac{1}{\sqrt{1-cx}\sqrt{cx+1}}dx}{2c^2}-\frac{x\sqrt{1-cx}\sqrt{cx+1}}{2c^2}\right)}{4c^2}-\frac{x^3\sqrt{1-cx}\sqrt{cx+1}}{4c^2}\right)}{6c^2}-\frac{x^5\sqrt{1-cx}\sqrt{cx+1}}{6c^2}\right)+\frac{1}{7}x^7(a+b\operatorname{sech}^{-1}(cx))$$

↓ 39

$$\frac{1}{7}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{5\left(\frac{3\left(\frac{\int\frac{1}{\sqrt{1-c^2x^2}}dx}{2c^2}-\frac{x\sqrt{1-cx}\sqrt{cx+1}}{2c^2}\right)}{4c^2}-\frac{x^3\sqrt{1-cx}\sqrt{cx+1}}{4c^2}\right)}{6c^2}-\frac{x^5\sqrt{1-cx}\sqrt{cx+1}}{6c^2}\right)+\frac{1}{7}x^7(a+b\operatorname{sech}^{-1}(cx))$$

↓ 223

$$\frac{1}{7}x^7(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{7}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{5\left(\frac{3\left(\frac{\arcsin(cx)}{2c^3}-\frac{x\sqrt{1-cx}\sqrt{cx+1}}{2c^2}\right)}{4c^2}-\frac{x^3\sqrt{1-cx}\sqrt{cx+1}}{4c^2}\right)}{6c^2}-\frac{x^5\sqrt{1-cx}\sqrt{cx+1}}{6c^2}\right)$$

input `Int[x^6*(a + b*ArcSech[c*x]),x]`

output `(x^7*(a + b*ArcSech[c*x]))/7 + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/6*(x^5*Sqrt[1 - c*x]*Sqrt[1 + c*x])/c^2 + (5*(-1/4*(x^3*Sqrt[1 - c*x]*Sqrt[1 + c*x])/c^2 + (3*(-1/2*(x*Sqrt[1 - c*x]*Sqrt[1 + c*x])/c^2 + ArcSin[c*x]/(2*c^3)))/(4*c^2)))/(6*c^2))/7`

## Definitions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27  $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 39  $\text{Int}[(\text{a}_) + (\text{b}_)*(\text{x}_)^{(\text{m}_)}*((\text{c}_) + (\text{d}_)*(\text{x}_)^{(\text{m}_)}), \text{x\_Symbol}] \rightarrow \text{Int}[(\text{a}*c + \text{b}*d*x^2)^m, \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{m}\}, \text{x}] \ \&\& \ \text{EqQ}[\text{b}*c + \text{a}*d, 0] \ \&\& \ (\text{IntegerQ}[\text{m}] \ || \ (\text{GtQ}[\text{a}, 0] \ \&\& \ \text{GtQ}[\text{c}, 0]))$
- rule 101  $\text{Int}[(\text{a}_.) + (\text{b}_.)*(\text{x}_.)^2*((\text{c}_.) + (\text{d}_.)*(\text{x}_.)^{(\text{n}_)}*((\text{e}_.) + (\text{f}_.)*(\text{x}_.)^{(\text{p}_)}), \text{x}_] \rightarrow \text{Simp}[\text{b}*(\text{a} + \text{b}*x)*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)}/(d*f*(n + p + 3))), \text{x}] + \text{Simp}[1/(d*f*(n + p + 3)) \quad \text{Int}[(c + d*x)^n*(e + f*x)^p*\text{Simp}[\text{a}^2*d*f*(n + p + 3) - \text{b}*(\text{b}*c*e + \text{a}*(d*e*(n + 1) + c*f*(p + 1))) + \text{b}*(\text{a}*d*f*(n + p + 4) - \text{b}*(d*e*(n + 2) + c*f*(p + 2)))*x, \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{n}, \text{p}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{n} + \text{p} + 3, 0]$
- rule 111  $\text{Int}[(\text{a}_.) + (\text{b}_.)*(\text{x}_.)^{(\text{m}_)}*((\text{c}_.) + (\text{d}_.)*(\text{x}_.)^{(\text{n}_)}*((\text{e}_.) + (\text{f}_.)*(\text{x}_.)^{(\text{p}_)}), \text{x}_] \rightarrow \text{Simp}[\text{b}*(\text{a} + \text{b}*x)^{(m - 1)}*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)}/(d*f*(m + n + p + 1))), \text{x}] + \text{Simp}[1/(d*f*(m + n + p + 1)) \quad \text{Int}[(\text{a} + \text{b}*x)^{(m - 2)}*(c + d*x)^n*(e + f*x)^p*\text{Simp}[\text{a}^2*d*f*(m + n + p + 1) - \text{b}*(\text{b}*c*e*(m - 1) + \text{a}*(d*e*(n + 1) + c*f*(p + 1))) + \text{b}*(\text{a}*d*f*(2*m + n + p) - \text{b}*(d*e*(m + n) + c*f*(m + p)))*x, \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{n}, \text{p}\}, \text{x}] \ \&\& \ \text{GtQ}[\text{m}, 1] \ \&\& \ \text{NeQ}[\text{m} + \text{n} + \text{p} + 1, 0] \ \&\& \ \text{IntegerQ}[\text{m}]$
- rule 223  $\text{Int}[1/\text{Sqrt}[(\text{a}_) + (\text{b}_.)*(\text{x}_.)^2], \text{x\_Symbol}] \rightarrow \text{Simp}[\text{ArcSin}[\text{Rt}[-\text{b}, 2]*(\text{x}/\text{Sqrt}[\text{a}])]/\text{Rt}[-\text{b}, 2], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{GtQ}[\text{a}, 0] \ \&\& \ \text{NegQ}[\text{b}]$
- rule 6837  $\text{Int}[(\text{a}_.) + \text{ArcSech}[(\text{c}_.)*(\text{x}_.)]*(\text{b}_.))*((\text{d}_.)*(\text{x}_.)^{(\text{m}_)}), \text{x\_Symbol}] \rightarrow \text{Simp}[(d*x)^{(m + 1)}*((a + b*\text{ArcSech}[c*x])/(d*(m + 1))), \text{x}] + \text{Simp}[\text{b}*(\text{Sqrt}[1 + c*x]/(m + 1))*\text{Sqrt}[1/(1 + c*x)] \quad \text{Int}[(d*x)^m/(\text{Sqrt}[1 - c*x]*\text{Sqrt}[1 + c*x]), \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{m}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{m}, -1]$

### Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.81

method	result
parts	$\frac{ax^7}{7} + \frac{b \left( \frac{c^7 x^7 \operatorname{arcsech}(cx)}{7} + \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (-8\sqrt{-c^2x^2+1} c^5 x^5 - 10c^3 x^3 \sqrt{-c^2x^2+1} - 15cx \sqrt{-c^2x^2+1} + 15 \arcsin(cx))}{336\sqrt{-c^2x^2+1}} \right)}{c^7}$
derivativedivides	$\frac{\frac{ac^7x^7}{7} + b \left( \frac{c^7 x^7 \operatorname{arcsech}(cx)}{7} + \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (-8\sqrt{-c^2x^2+1} c^5 x^5 - 10c^3 x^3 \sqrt{-c^2x^2+1} - 15cx \sqrt{-c^2x^2+1} + 15 \arcsin(cx))}{336\sqrt{-c^2x^2+1}} \right)}{c^7}$
default	$\frac{\frac{ac^7x^7}{7} + b \left( \frac{c^7 x^7 \operatorname{arcsech}(cx)}{7} + \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (-8\sqrt{-c^2x^2+1} c^5 x^5 - 10c^3 x^3 \sqrt{-c^2x^2+1} - 15cx \sqrt{-c^2x^2+1} + 15 \arcsin(cx))}{336\sqrt{-c^2x^2+1}} \right)}{c^7}$

input `int(x^6*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{7}ax^7 + \frac{b}{c^7} \left( \frac{1}{7}c^7x^7 \operatorname{arcsech}(cx) + \frac{1}{336} \left( -\frac{cx-1}{cx} \right)^{1/2} c^5 x^5 - 10 \left( -\frac{cx-1}{cx} \right)^{1/2} c^3 x^3 \left( -\frac{cx-1}{cx} \right)^{1/2} - 15 \left( -\frac{cx-1}{cx} \right)^{1/2} c x \left( -\frac{cx-1}{cx} \right)^{1/2} + 15 \arcsin(cx) \right) / \left( -\frac{cx-1}{cx} \right)^{1/2}$$

### Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.11

$$\int x^6 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{48ac^7x^7 - 48bc^7 \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{x}\right) - 30b \arctan\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{cx}\right) + 48(bc^7x^7 - bc^7) \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}+1}{cx}\right)}{336c^7}$$

input `integrate(x^6*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output 
$$\frac{1}{336} (48ac^7x^7 - 48bc^7 \log\left(\frac{cx\sqrt{-(c^2x^2-1)/(c^2x^2)}-1}{x}\right) - 30b \arctan\left(\frac{cx\sqrt{-(c^2x^2-1)/(c^2x^2)}-1}{cx}\right) + 48(bc^7x^7 - bc^7) \log\left(\frac{cx\sqrt{-(c^2x^2-1)/(c^2x^2)}+1}{cx}\right) - (8bc^6x^6 + 10bc^4x^4 + 15bc^2x^2) \sqrt{-(c^2x^2-1)/(c^2x^2)}) / c^7$$

**Sympy [F]**

$$\int x^6(a + b \operatorname{sech}^{-1}(cx)) dx = \int x^6(a + b \operatorname{asech}(cx)) dx$$

input `integrate(x**6*(a+b*asech(c*x)),x)`

output `Integral(x**6*(a + b*asech(c*x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.82

$$\int x^6(a + b \operatorname{sech}^{-1}(cx)) dx = \frac{1}{7} ax^7 + \frac{1}{336} \left( 48 x^7 \operatorname{arsech}(cx) - \frac{15 \left(\frac{1}{c^2 x^2} - 1\right)^{\frac{5}{2}} + 40 \left(\frac{1}{c^2 x^2} - 1\right)^{\frac{3}{2}} + 33 \sqrt{\frac{1}{c^2 x^2} - 1}}{c^6 \left(\frac{1}{c^2 x^2} - 1\right)^3 + 3 c^6 \left(\frac{1}{c^2 x^2} - 1\right)^2 + 3 c^6 \left(\frac{1}{c^2 x^2} - 1\right) + c^6} + \frac{15 \arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right)}{c^6} \right) b$$

input `integrate(x^6*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `1/7*a*x^7 + 1/336*(48*x^7*arcsech(c*x) - ((15*(1/(c^2*x^2) - 1)^(5/2) + 40*(1/(c^2*x^2) - 1)^(3/2) + 33*sqrt(1/(c^2*x^2) - 1))/(c^6*(1/(c^2*x^2) - 1)^3 + 3*c^6*(1/(c^2*x^2) - 1)^2 + 3*c^6*(1/(c^2*x^2) - 1) + c^6) + 15*arctan(sqrt(1/(c^2*x^2) - 1))/c^6)/c)*b`



**Giac [F]**

$$\int x^6(a + b\operatorname{sech}^{-1}(cx)) dx = \int (b \operatorname{arsech}(cx) + a)x^6 dx$$

input `integrate(x^6*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^6, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^6(a + b\operatorname{sech}^{-1}(cx)) dx = \int x^6 \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int(x^6*(a + b*acosh(1/(c*x))),x)`

output `int(x^6*(a + b*acosh(1/(c*x))), x)`

**Reduce [F]**

$$\int x^6(a + b\operatorname{sech}^{-1}(cx)) dx = \left( \int \operatorname{asech}(cx) x^6 dx \right) b + \frac{ax^7}{7}$$

input `int(x^6*(a+b*asech(c*x)),x)`

output `(7*int(asech(c*x)*x**6,x)*b + a*x**7)/7`

### 3.20 $\int x^5(a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	233
Mathematica [A] (verified)	233
Rubi [A] (verified)	234
Maple [A] (verified)	236
Fricas [A] (verification not implemented)	237
Sympy [A] (verification not implemented)	237
Maxima [A] (verification not implemented)	238
Giac [F]	238
Mupad [F(-1)]	238
Reduce [F]	239

#### Optimal result

Integrand size = 12, antiderivative size = 131

$$\int x^5(a + b\operatorname{sech}^{-1}(cx)) dx = -\frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{6c^6} + \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(1-c^2x^2)}{9c^6} - \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(1-c^2x^2)^2}{30c^6} + \frac{1}{6}x^6(a + b\operatorname{sech}^{-1}(cx))$$

output

```
-1/6*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/c^6+1/9*b*((-c*x+1)/(c*x+1))^(1/2)
*(c*x+1)*(-c^2*x^2+1)/c^6-1/30*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(-c^2*x^
2+1)^2/c^6+1/6*x^6*(a+b*arcsech(c*x))
```

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.74

$$\int x^5(a + b\operatorname{sech}^{-1}(cx)) dx = \frac{ax^6}{6} + b\sqrt{\frac{1-cx}{1+cx}}\left(-\frac{4}{45c^6} - \frac{4x}{45c^5} - \frac{2x^2}{45c^4} - \frac{2x^3}{45c^3} - \frac{x^4}{30c^2} - \frac{x^5}{30c}\right) + \frac{1}{6}bx^6\operatorname{sech}^{-1}(cx)$$

input `Integrate[x^5*(a + b*ArcSech[c*x]),x]`

output  $(a*x^6)/6 + b*\text{Sqrt}[(1 - c*x)/(1 + c*x)]*(-4/(45*c^6) - (4*x)/(45*c^5) - (2*x^2)/(45*c^4) - (2*x^3)/(45*c^3) - x^4/(30*c^2) - x^5/(30*c)) + (b*x^6*ArcSech[c*x])/6$

### Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.02, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6837, 111, 27, 111, 27, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^5(a + b\text{sech}^{-1}(cx)) dx$$

$$\downarrow 6837$$

$$\frac{1}{6}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^5}{\sqrt{1-cx}\sqrt{cx+1}} dx + \frac{1}{6}x^6(a + b\text{sech}^{-1}(cx))$$

$$\downarrow 111$$

$$\frac{1}{6}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{\int -\frac{4x^3}{\sqrt{1-cx}\sqrt{cx+1}} dx}{5c^2} - \frac{x^4\sqrt{1-cx}\sqrt{cx+1}}{5c^2} \right) + \frac{1}{6}x^6(a + b\text{sech}^{-1}(cx))$$

$$\downarrow 27$$

$$\frac{1}{6}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{4 \int \frac{x^3}{\sqrt{1-cx}\sqrt{cx+1}} dx}{5c^2} - \frac{x^4\sqrt{1-cx}\sqrt{cx+1}}{5c^2} \right) + \frac{1}{6}x^6(a + b\text{sech}^{-1}(cx))$$

$$\downarrow 111$$

$$\frac{1}{6}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{4 \left( -\frac{\int -\frac{2x}{\sqrt{1-cx}\sqrt{cx+1}} dx}{3c^2} - \frac{x^2\sqrt{1-cx}\sqrt{cx+1}}{3c^2} \right)}{5c^2} - \frac{x^4\sqrt{1-cx}\sqrt{cx+1}}{5c^2} \right) + \frac{1}{6}x^6(a + b\text{sech}^{-1}(cx))$$

$$\begin{aligned}
 & \frac{1}{6}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{4\left(\frac{2\int\frac{x}{\sqrt{1-cx}\sqrt{cx+1}}dx}{3c^2} - \frac{x^2\sqrt{1-cx}\sqrt{cx+1}}{3c^2}\right)}{5c^2} - \frac{x^4\sqrt{1-cx}\sqrt{cx+1}}{5c^2} \right) + \\
 & \frac{1}{6}x^6(a + b\operatorname{sech}^{-1}(cx)) \\
 & \downarrow 83 \\
 & \frac{1}{6}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{4\left(-\frac{2\sqrt{1-cx}\sqrt{cx+1}}{3c^4} - \frac{x^2\sqrt{1-cx}\sqrt{cx+1}}{3c^2}\right)}{5c^2} - \frac{x^4\sqrt{1-cx}\sqrt{cx+1}}{5c^2} \right) + \\
 & \frac{1}{6}x^6(a + b\operatorname{sech}^{-1}(cx)) +
 \end{aligned}$$

input `Int[x^5*(a + b*ArcSech[c*x]),x]`

output `(b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/5*(x^4*Sqrt[1 - c*x]*Sqrt[1 + c*x])/c^2 + (4*((-2*Sqrt[1 - c*x]*Sqrt[1 + c*x])/(3*c^4) - (x^2*Sqrt[1 - c*x]*Sqrt[1 + c*x])/(3*c^2)))/(5*c^2)))/6 + (x^6*(a + b*ArcSech[c*x]))/6`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

rule 111

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/(d*f*(m + n + p + 1)), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]
```

rule 6837

```
Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcSech[c*x])/(d*(m + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(m + 1))*Sqrt[1/(1 + c*x)] Int[(d*x)^m/(Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

### Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.59

method	result	size
parts	$\frac{x^6 a}{6} + \frac{b \left( \frac{c^6 x^6 \operatorname{arcsech}(cx) - \sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (3c^4 x^4 + 4c^2 x^2 + 8)}{6} \right)}{c^6}$	77
derivativedivides	$\frac{\frac{a c^6 x^6}{6} + b \left( \frac{c^6 x^6 \operatorname{arcsech}(cx) - \sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (3c^4 x^4 + 4c^2 x^2 + 8)}{6} \right)}{c^6}$	81
default	$\frac{\frac{a c^6 x^6}{6} + b \left( \frac{c^6 x^6 \operatorname{arcsech}(cx) - \sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (3c^4 x^4 + 4c^2 x^2 + 8)}{6} \right)}{c^6}$	81

input

```
int(x^5*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)
```

output

```
1/6*x^6*a+b/c^6*(1/6*c^6*x^6*arcsech(c*x)-1/90*(-(c*x-1)/c/x)^(1/2)*c*x*((c*x+1)/c/x)^(1/2)*(3*c^4*x^4+4*c^2*x^2+8))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.76

$$\int x^5 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{15 bc^5 x^6 \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx}\right) + 15 ac^5 x^6 - (3 bc^4 x^5 + 4 bc^2 x^3 + 8 bx) \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}}}{90 c^5}$$

input `integrate(x^5*(a+b*arcsech(c*x)),x, algorithm="fricas")`output `1/90*(15*b*c^5*x^6*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 15*a*c^5*x^6 - (3*b*c^4*x^5 + 4*b*c^2*x^3 + 8*b*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/c^5`**Sympy [A] (verification not implemented)**

Time = 0.91 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.72

$$\int x^5 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \begin{cases} \frac{ax^6}{6} + \frac{bx^6 \operatorname{asech}(cx)}{6} - \frac{bx^4 \sqrt{-c^2 x^2 + 1}}{30c^2} - \frac{2bx^2 \sqrt{-c^2 x^2 + 1}}{45c^4} - \frac{4b \sqrt{-c^2 x^2 + 1}}{45c^6} & \text{for } c \neq 0 \\ \frac{x^6(a + \infty b)}{6} & \text{otherwise} \end{cases}$$

input `integrate(x**5*(a+b*asech(c*x)),x)`output `Piecewise((a*x**6/6 + b*x**6*asech(c*x)/6 - b*x**4*sqrt(-c**2*x**2 + 1)/(30*c**2) - 2*b*x**2*sqrt(-c**2*x**2 + 1)/(45*c**4) - 4*b*sqrt(-c**2*x**2 + 1)/(45*c**6), Ne(c, 0)), (x**6*(a + oo*b)/6, True))`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.60

$$\int x^5(a + b \operatorname{sech}^{-1}(cx)) dx = \frac{1}{6} ax^6 + \frac{1}{90} \left( 15x^6 \operatorname{arosech}(cx) - \frac{3c^4x^5 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{5}{2}} - 10c^2x^3 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{3}{2}} + 15x \sqrt{\frac{1}{c^2x^2} - 1}}{c^5} \right) b$$

input `integrate(x^5*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `1/6*a*x^6 + 1/90*(15*x^6*arcsech(c*x) - (3*c^4*x^5*(1/(c^2*x^2) - 1)^(5/2) - 10*c^2*x^3*(1/(c^2*x^2) - 1)^(3/2) + 15*x*sqrt(1/(c^2*x^2) - 1))/c^5)*b`

**Giac [F]**

$$\int x^5(a + b \operatorname{sech}^{-1}(cx)) dx = \int (b \operatorname{arosech}(cx) + a)x^5 dx$$

input `integrate(x^5*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^5, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^5(a + b \operatorname{sech}^{-1}(cx)) dx = \int x^5 \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int(x^5*(a + b*acosh(1/(c*x))),x)`

output `int(x^5*(a + b*acosh(1/(c*x))), x)`

**Reduce [F]**

$$\int x^5 (a + b \operatorname{sech}^{-1}(cx)) dx = \left( \int a \operatorname{sech}(cx) x^5 dx \right) b + \frac{ax^6}{6}$$

input `int(x^5*(a+b*asech(c*x)),x)`

output `(6*int(asech(c*x)*x**5,x)*b + a*x**6)/6`



### 3.21 $\int x^4(a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	240
Mathematica [C] (verified)	240
Rubi [A] (verified)	241
Maple [A] (verified)	244
Fricas [A] (verification not implemented)	244
Sympy [F]	245
Maxima [A] (verification not implemented)	245
Giac [F]	246
Mupad [F(-1)]	246
Reduce [F]	246

#### Optimal result

Integrand size = 12, antiderivative size = 131

$$\int x^4(a + b\operatorname{sech}^{-1}(cx)) dx = -\frac{3bx\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{40c^4} - \frac{bx^3\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{20c^2} + \frac{1}{5}x^5(a + b\operatorname{sech}^{-1}(cx)) + \frac{3b\sqrt{\frac{1-cx}{1+cx}}(1+cx)\arcsin(cx)}{40c^5\sqrt{1-c^2x^2}}$$

output

```
-3/40*b*x*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/c^4-1/20*b*x^3*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/c^2+1/5*x^5*(a+b*arcsech(c*x))+3/40*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*arcsin(c*x)/c^5/(-c^2*x^2+1)^(1/2)
```

#### Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.94

$$\int x^4(a + b\operatorname{sech}^{-1}(cx)) dx = \frac{ax^5}{5} + b\sqrt{\frac{1-cx}{1+cx}}\left(-\frac{3x}{40c^4} - \frac{3x^2}{40c^3} - \frac{x^3}{20c^2} - \frac{x^4}{20c}\right) + \frac{1}{5}bx^5\operatorname{sech}^{-1}(cx) + \frac{3ib\log\left(-2icx + 2\sqrt{\frac{1-cx}{1+cx}}(1+cx)\right)}{40c^5}$$

input `Integrate[x^4*(a + b*ArcSech[c*x]),x]`

output  $(a*x^5)/5 + b*\text{Sqrt}[(1 - c*x)/(1 + c*x)]*((-3*x)/(40*c^4) - (3*x^2)/(40*c^3) - x^3/(20*c^2) - x^4/(20*c)) + (b*x^5*\text{ArcSech}[c*x])/5 + ((3*I)/40)*b*\text{Log}[(-2*I)*c*x + 2*\text{Sqrt}[(1 - c*x)/(1 + c*x)]*(1 + c*x)]/c^5$

### Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.89, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$ , Rules used = {6837, 111, 27, 101, 25, 39, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4(a + b\text{sech}^{-1}(cx)) dx$$

$$\downarrow 6837$$

$$\frac{1}{5}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^4}{\sqrt{1-cx}\sqrt{cx+1}} dx + \frac{1}{5}x^5(a + b\text{sech}^{-1}(cx))$$

$$\downarrow 111$$

$$\frac{1}{5}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{\int -\frac{3x^2}{\sqrt{1-cx}\sqrt{cx+1}} dx}{4c^2} - \frac{x^3\sqrt{1-cx}\sqrt{cx+1}}{4c^2} \right) + \frac{1}{5}x^5(a + b\text{sech}^{-1}(cx))$$

$$\downarrow 27$$

$$\frac{1}{5}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3 \int \frac{x^2}{\sqrt{1-cx}\sqrt{cx+1}} dx}{4c^2} - \frac{x^3\sqrt{1-cx}\sqrt{cx+1}}{4c^2} \right) + \frac{1}{5}x^5(a + b\text{sech}^{-1}(cx))$$

$$\downarrow 101$$

$$\frac{1}{5}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3 \left( -\frac{\int -\frac{1}{\sqrt{1-cx}\sqrt{cx+1}} dx}{2c^2} - \frac{x\sqrt{1-cx}\sqrt{cx+1}}{2c^2} \right)}{4c^2} - \frac{x^3\sqrt{1-cx}\sqrt{cx+1}}{4c^2} \right) + \frac{1}{5}x^5(a + b\text{sech}^{-1}(cx))$$

$$\begin{aligned}
& \frac{1}{5}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3\left(\frac{\int \frac{1}{\sqrt{1-cx}\sqrt{cx+1}} dx}{2c^2} - \frac{x\sqrt{1-cx}\sqrt{cx+1}}{2c^2}\right)}{4c^2} - \frac{x^3\sqrt{1-cx}\sqrt{cx+1}}{4c^2} \right) + \\
& \qquad \qquad \qquad \frac{1}{5}x^5(a + b\operatorname{sech}^{-1}(cx)) \\
& \qquad \qquad \qquad \downarrow 25 \\
& \frac{1}{5}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3\left(\frac{\int \frac{1}{\sqrt{1-c^2x^2}} dx}{2c^2} - \frac{x\sqrt{1-cx}\sqrt{cx+1}}{2c^2}\right)}{4c^2} - \frac{x^3\sqrt{1-cx}\sqrt{cx+1}}{4c^2} \right) + \\
& \qquad \qquad \qquad \frac{1}{5}x^5(a + b\operatorname{sech}^{-1}(cx)) \\
& \qquad \qquad \qquad \downarrow 39 \\
& \frac{1}{5}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3\left(\frac{\arcsin(cx)}{2c^3} - \frac{x\sqrt{1-cx}\sqrt{cx+1}}{2c^2}\right)}{4c^2} - \frac{x^3\sqrt{1-cx}\sqrt{cx+1}}{4c^2} \right) + \\
& \qquad \qquad \qquad \frac{1}{5}x^5(a + b\operatorname{sech}^{-1}(cx)) +
\end{aligned}$$

input `Int[x^4*(a + b*ArcSech[c*x]),x]`

output `(x^5*(a + b*ArcSech[c*x])/5 + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/4*(x^3*Sqrt[1 - c*x]*Sqrt[1 + c*x])/c^2 + (3*(-1/2*(x*Sqrt[1 - c*x]*Sqrt[1 + c*x])/c^2 + ArcSin[c*x]/(2*c^3)))/(4*c^2)))/5`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 39

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(m_), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))
```

rule 101

```
Int[((a_) + (b_)*(x_))^(2)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]
```

rule 111

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]
```

rule 223

```
Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

rule 6837

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_))*((d_)*(x_))^(m_), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcSech[c*x])/(d*(m + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(m + 1))*Sqrt[1/(1 + c*x)] Int[(d*x)^m/(Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

**Maple [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.87

method	result	size
parts	$\frac{ax^5}{5} + \frac{b \left( \frac{c^5 x^5 \operatorname{arcsech}(cx)}{5} + \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (-2c^3 x^3 \sqrt{-c^2 x^2 + 1} - 3cx \sqrt{-c^2 x^2 + 1} + 3 \arcsin(cx))}{40 \sqrt{-c^2 x^2 + 1}} \right)}{c^5}$	114
derivativedivides	$\frac{ac^5 x^5}{5} + b \left( \frac{c^5 x^5 \operatorname{arcsech}(cx)}{5} + \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (-2c^3 x^3 \sqrt{-c^2 x^2 + 1} - 3cx \sqrt{-c^2 x^2 + 1} + 3 \arcsin(cx))}{40 \sqrt{-c^2 x^2 + 1}} \right)$	118
default	$\frac{ac^5 x^5}{5} + b \left( \frac{c^5 x^5 \operatorname{arcsech}(cx)}{5} + \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (-2c^3 x^3 \sqrt{-c^2 x^2 + 1} - 3cx \sqrt{-c^2 x^2 + 1} + 3 \arcsin(cx))}{40 \sqrt{-c^2 x^2 + 1}} \right)$	118

input `int(x^4*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{5}ax^5 + \frac{b}{c^5} \left( \frac{1}{5}c^5 x^5 \operatorname{arcsech}(cx) + \frac{1}{40} \left( -\frac{cx-1}{cx} \right)^{1/2} cx \left( \frac{cx+1}{cx} \right)^{1/2} \left( -2c^3 x^3 \sqrt{-c^2 x^2 + 1} - 3cx \sqrt{-c^2 x^2 + 1} + 3 \arcsin(cx) \right) \right) / \sqrt{-c^2 x^2 + 1}$$

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.33

$$\int x^4 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{8ac^5 x^5 - 8bc^5 \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - 1}{x}\right) - 6b \arctan\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - 1}{cx}\right) + 8(bc^5 x^5 - bc^5) \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx}\right)}{40c^5}$$

input `integrate(x^4*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output 
$$\frac{1}{40} \left( 8ac^5 x^5 - 8bc^5 \log\left(\frac{cx \sqrt{-(c^2 x^2 - 1)/(c^2 x^2)} - 1}{x}\right) - 6b \arctan\left(\frac{cx \sqrt{-(c^2 x^2 - 1)/(c^2 x^2)} - 1}{cx}\right) + 8(bc^5 x^5 - bc^5) \log\left(\frac{cx \sqrt{-(c^2 x^2 - 1)/(c^2 x^2)} + 1}{cx}\right) - (2bc^4 x^4 + 3bc^2 x^2) \sqrt{-(c^2 x^2 - 1)/(c^2 x^2)} \right) / c^5$$

**Sympy [F]**

$$\int x^4(a + b \operatorname{sech}^{-1}(cx)) dx = \int x^4(a + b \operatorname{asech}(cx)) dx$$

input `integrate(x**4*(a+b*asech(c*x)),x)`

output `Integral(x**4*(a + b*asech(c*x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.81

$$\int x^4(a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{1}{5} ax^5 + \frac{1}{40} \left( 8x^5 \operatorname{arsech}(cx) - \frac{3 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{3}{2}} + 5 \sqrt{\frac{1}{c^2 x^2} - 1}}{c^4 \left( \frac{1}{c^2 x^2} - 1 \right)^2 + 2c^4 \left( \frac{1}{c^2 x^2} - 1 \right) + c^4} + \frac{3 \arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right)}{c^4} \right) b$$

input `integrate(x^4*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `1/5*a*x^5 + 1/40*(8*x^5*arcsech(c*x) - ((3*(1/(c^2*x^2) - 1)^(3/2) + 5*sqrt(1/(c^2*x^2) - 1))/(c^4*(1/(c^2*x^2) - 1)^2 + 2*c^4*(1/(c^2*x^2) - 1) + c^4) + 3*arctan(sqrt(1/(c^2*x^2) - 1))/c^4)/c)*b`

**Giac [F]**

$$\int x^4(a + b\operatorname{sech}^{-1}(cx)) dx = \int (b \operatorname{arsech}(cx) + a)x^4 dx$$

input `integrate(x^4*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^4, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^4(a + b\operatorname{sech}^{-1}(cx)) dx = \int x^4 \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int(x^4*(a + b*acosh(1/(c*x))),x)`

output `int(x^4*(a + b*acosh(1/(c*x))), x)`

**Reduce [F]**

$$\int x^4(a + b\operatorname{sech}^{-1}(cx)) dx = \left( \int \operatorname{asech}(cx) x^4 dx \right) b + \frac{ax^5}{5}$$

input `int(x^4*(a+b*asech(c*x)),x)`

output `(5*int(asech(c*x)*x**4,x)*b + a*x**5)/5`

### 3.22 $\int x^3(a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	247
Mathematica [A] (verified)	247
Rubi [A] (verified)	248
Maple [A] (verified)	249
Fricas [A] (verification not implemented)	250
Sympy [A] (verification not implemented)	250
Maxima [A] (verification not implemented)	251
Giac [F]	251
Mupad [F(-1)]	251
Reduce [F]	252

#### Optimal result

Integrand size = 12, antiderivative size = 88

$$\int x^3(a + b\operatorname{sech}^{-1}(cx)) dx = -\frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{4c^4} + \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(1-c^2x^2)}{12c^4} + \frac{1}{4}x^4(a + b\operatorname{sech}^{-1}(cx))$$

output

```
-1/4*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/c^4+1/12*b*((-c*x+1)/(c*x+1))^(1/2)*
(c*x+1)*(-c^2*x^2+1)/c^4+1/4*x^4*(a+b*arcsech(c*x))
```

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.88

$$\int x^3(a + b\operatorname{sech}^{-1}(cx)) dx = \frac{ax^4}{4} + b\sqrt{\frac{1-cx}{1+cx}}\left(-\frac{1}{6c^4} - \frac{x}{6c^3} - \frac{x^2}{12c^2} - \frac{x^3}{12c}\right) + \frac{1}{4}bx^4\operatorname{sech}^{-1}(cx)$$

input

```
Integrate[x^3*(a + b*ArcSech[c*x]),x]
```



output

$$(a*x^4)/4 + b*\text{Sqrt}[(1 - c*x)/(1 + c*x)]*(-1/6*1/c^4 - x/(6*c^3) - x^2/(12*c^2) - x^3/(12*c)) + (b*x^4*\text{ArcSech}[c*x])/4$$

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.10, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6837, 111, 27, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3(a + b\text{sech}^{-1}(cx)) dx$$

$$\downarrow 6837$$

$$\frac{1}{4}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^3}{\sqrt{1-cx}\sqrt{cx+1}} dx + \frac{1}{4}x^4(a + b\text{sech}^{-1}(cx))$$

$$\downarrow 111$$

$$\frac{1}{4}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{\int -\frac{2x}{\sqrt{1-cx}\sqrt{cx+1}} dx}{3c^2} - \frac{x^2\sqrt{1-cx}\sqrt{cx+1}}{3c^2} \right) + \frac{1}{4}x^4(a + b\text{sech}^{-1}(cx))$$

$$\downarrow 27$$

$$\frac{1}{4}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{2 \int \frac{x}{\sqrt{1-cx}\sqrt{cx+1}} dx}{3c^2} - \frac{x^2\sqrt{1-cx}\sqrt{cx+1}}{3c^2} \right) + \frac{1}{4}x^4(a + b\text{sech}^{-1}(cx))$$

$$\downarrow 83$$

$$\frac{1}{4}x^4(a + b\text{sech}^{-1}(cx)) + \frac{1}{4}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{2\sqrt{1-cx}\sqrt{cx+1}}{3c^4} - \frac{x^2\sqrt{1-cx}\sqrt{cx+1}}{3c^2} \right)$$

input

$$\text{Int}[x^3*(a + b*\text{ArcSech}[c*x]),x]$$

output

$$(b*\text{Sqrt}[(1 + c*x)^{-1}]*\text{Sqrt}[1 + c*x]*((-2*\text{Sqrt}[1 - c*x]*\text{Sqrt}[1 + c*x])/(3*c^4) - (x^2*\text{Sqrt}[1 - c*x]*\text{Sqrt}[1 + c*x])/(3*c^2)))/4 + (x^4*(a + b*\text{ArcSech}[c*x]))/4$$

**Defintions of rubi rules used**

rule 27  $\text{Int}[(a\_)(Fx\_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b\_)(Gx\_)] /; \text{FreeQ}[b, x]$

rule 83  $\text{Int}[((a\_.) + (b\_.)*(x\_))*((c\_.) + (d\_.)*(x\_))^n*((e\_.) + (f\_.)*(x\_))^p, x] \rightarrow \text{Simp}[b*(c + d*x)^{n+1}*((e + f*x)^{p+1}/(d*f*(n + p + 2))), x] /; \text{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \text{NeQ}[n + p + 2, 0] \ \&\& \ \text{EqQ}[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]$

rule 111  $\text{Int}[((a\_.) + (b\_.)*(x\_))^m*((c\_.) + (d\_.)*(x\_))^n*((e\_.) + (f\_.)*(x\_))^p, x] \rightarrow \text{Simp}[b*(a + b*x)^{m-1}*(c + d*x)^{n+1}*((e + f*x)^{p+1}/(d*f*(m + n + p + 1))), x] + \text{Simp}[1/(d*f*(m + n + p + 1)) \text{ Int}[(a + b*x)^{m-2}*(c + d*x)^n*(e + f*x)^p*\text{Simp}[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m-1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p))]*x, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{NeQ}[m + n + p + 1, 0] \ \&\& \ \text{IntegerQ}[m]$

rule 6837  $\text{Int}[((a\_.) + \text{ArcSech}[c\_.*(x\_)]*(b\_))*((d\_.)*(x\_))^m, x\_Symbol] \rightarrow \text{Simp}[(d*x)^{m+1}*((a + b*\text{ArcSech}[c*x])/(d*(m + 1))), x] + \text{Simp}[b*(\text{Sqrt}[1 + c*x]/(m + 1))*\text{Sqrt}[1/(1 + c*x)] \text{ Int}[(d*x)^m/(\text{Sqrt}[1 - c*x]*\text{Sqrt}[1 + c*x]), x], x] /; \text{FreeQ}\{a, b, c, d, m\}, x] \ \&\& \ \text{NeQ}[m, -1]$

**Maple [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.77

method	result	size
parts	$\frac{x^4 a}{4} + \frac{b \left( \frac{c^4 x^4 \operatorname{arcsech}(cx) - \sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (c^2 x^2 + 2)}{4} \right)}{c^4}$	68
derivativedivides	$\frac{a c^4 x^4}{4} + b \left( \frac{c^4 x^4 \operatorname{arcsech}(cx) - \sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (c^2 x^2 + 2)}{4} \right)$	72
default	$\frac{a c^4 x^4}{4} + b \left( \frac{c^4 x^4 \operatorname{arcsech}(cx) - \sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (c^2 x^2 + 2)}{4} \right)$	72

input `int(x^3*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)`

output `1/4*x^4*a+b/c^4*(1/4*c^4*x^4*arcsech(c*x)-1/12*(-(c*x-1)/c/x)^(1/2)*c*x*((c*x+1)/c/x)^(1/2)*(c^2*x^2+2))`

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.02

$$\int x^3(a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{3bc^3x^4 \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}+1}{cx}\right) + 3ac^3x^4 - (bc^2x^3 + 2bx)\sqrt{-\frac{c^2x^2-1}{c^2x^2}}}{12c^3}$$

input `integrate(x^3*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `1/12*(3*b*c^3*x^4*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 3*a*c^3*x^4 - (b*c^2*x^3 + 2*b*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/c^3`

### Sympy [A] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.77

$$\int x^3(a + b \operatorname{sech}^{-1}(cx)) dx = \begin{cases} \frac{ax^4}{4} + \frac{bx^4 \operatorname{asech}(cx)}{4} - \frac{bx^2\sqrt{-c^2x^2+1}}{12c^2} - \frac{b\sqrt{-c^2x^2+1}}{6c^4} & \text{for } c \neq 0 \\ \frac{x^4(a+\infty b)}{4} & \text{otherwise} \end{cases}$$

input `integrate(x**3*(a+b*asech(c*x)),x)`

output `Piecewise((a*x**4/4 + b*x**4*asech(c*x)/4 - b*x**2*sqrt(-c**2*x**2 + 1)/(12*c**2) - b*sqrt(-c**2*x**2 + 1)/(6*c**4), Ne(c, 0)), (x**4*(a + oo*b)/4, True))`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.65

$$\int x^3(a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{1}{4} ax^4 + \frac{1}{12} \left( 3x^4 \operatorname{arsech}(cx) + \frac{c^2 x^3 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{3}{2}} - 3x \sqrt{\frac{1}{c^2 x^2} - 1}}{c^3} \right) b$$

input `integrate(x^3*(a+b*arcsech(c*x)),x, algorithm="maxima")`output `1/4*a*x^4 + 1/12*(3*x^4*arcsech(c*x) + (c^2*x^3*(1/(c^2*x^2) - 1)^(3/2) - 3*x*sqrt(1/(c^2*x^2) - 1))/c^3)*b`**Giac [F]**

$$\int x^3(a + b \operatorname{sech}^{-1}(cx)) dx = \int (b \operatorname{arsech}(cx) + a)x^3 dx$$

input `integrate(x^3*(a+b*arcsech(c*x)),x, algorithm="giac")`output `integrate((b*arcsech(c*x) + a)*x^3, x)`**Mupad [F(-1)]**

Timed out.

$$\int x^3(a + b \operatorname{sech}^{-1}(cx)) dx = \int x^3 \left( a + b \operatorname{acosh} \left( \frac{1}{cx} \right) \right) dx$$

input `int(x^3*(a + b*acosh(1/(c*x))),x)`output `int(x^3*(a + b*acosh(1/(c*x))), x)`

**Reduce [F]**

$$\int x^3(a + b\operatorname{sech}^{-1}(cx)) dx = \left( \int a\operatorname{sech}(cx) x^3 dx \right) b + \frac{ax^4}{4}$$

input `int(x^3*(a+b*asech(c*x)),x)`

output `(4*int(asech(c*x)*x**3,x)*b + a*x**4)/4`

### 3.23 $\int x^2(a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	253
Mathematica [C] (verified)	253
Rubi [A] (verified)	254
Maple [A] (verified)	256
Fricas [A] (verification not implemented)	256
Sympy [F]	257
Maxima [A] (verification not implemented)	257
Giac [F]	257
Mupad [F(-1)]	258
Reduce [F]	258

#### Optimal result

Integrand size = 12, antiderivative size = 97

$$\int x^2(a + b\operatorname{sech}^{-1}(cx)) dx = -\frac{bx\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{6c^2} + \frac{1}{3}x^3(a + b\operatorname{sech}^{-1}(cx)) + \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)\arcsin(cx)}{6c^3\sqrt{1-c^2x^2}}$$

output

```
-1/6*b*x*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/c^2+1/3*x^3*(a+b*arcsech(c*x))+1/6*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*arcsin(c*x)/c^3/(-c^2*x^2+1)^(1/2)
```

#### Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.06

$$\int x^2(a + b\operatorname{sech}^{-1}(cx)) dx = \frac{ax^3}{3} + b\sqrt{\frac{1-cx}{1+cx}}\left(-\frac{x}{6c^2} - \frac{x^2}{6c}\right) + \frac{1}{3}bx^3\operatorname{sech}^{-1}(cx) + \frac{ib\log\left(-2icx + 2\sqrt{\frac{1-cx}{1+cx}}(1+cx)\right)}{6c^3}$$

input `Integrate[x^2*(a + b*ArcSech[c*x]),x]`

output  $(a*x^3)/3 + b*\text{Sqrt}[(1 - c*x)/(1 + c*x)]*(-1/6*x/c^2 - x^2/(6*c)) + (b*x^3* \text{ArcSech}[c*x])/3 + ((I/6)*b*\text{Log}[(-2*I)*c*x + 2*\text{Sqrt}[(1 - c*x)/(1 + c*x)]*(1 + c*x)])/c^3$

### Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.82, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {6837, 101, 25, 39, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2(a + b\text{sech}^{-1}(cx)) dx \\
 & \quad \downarrow 6837 \\
 & \frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^2}{\sqrt{1-cx}\sqrt{cx+1}} dx + \frac{1}{3}x^3(a + b\text{sech}^{-1}(cx)) \\
 & \quad \downarrow 101 \\
 & \frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{\int -\frac{1}{\sqrt{1-cx}\sqrt{cx+1}} dx}{2c^2} - \frac{x\sqrt{1-cx}\sqrt{cx+1}}{2c^2} \right) + \frac{1}{3}x^3(a + b\text{sech}^{-1}(cx)) \\
 & \quad \downarrow 25 \\
 & \frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{1}{\sqrt{1-cx}\sqrt{cx+1}} dx}{2c^2} - \frac{x\sqrt{1-cx}\sqrt{cx+1}}{2c^2} \right) + \frac{1}{3}x^3(a + b\text{sech}^{-1}(cx)) \\
 & \quad \downarrow 39 \\
 & \frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{1}{\sqrt{1-c^2x^2}} dx}{2c^2} - \frac{x\sqrt{1-cx}\sqrt{cx+1}}{2c^2} \right) + \frac{1}{3}x^3(a + b\text{sech}^{-1}(cx)) \\
 & \quad \downarrow 223 \\
 & \frac{1}{3}x^3(a + b\text{sech}^{-1}(cx)) + \frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\arcsin(cx)}{2c^3} - \frac{x\sqrt{1-cx}\sqrt{cx+1}}{2c^2} \right)
 \end{aligned}$$

input `Int[x^2*(a + b*ArcSech[c*x]),x]`

output `(x^3*(a + b*ArcSech[c*x]))/3 + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/2 * (x*Sqrt[1 - c*x]*Sqrt[1 + c*x])/c^2 + ArcSin[c*x]/(2*c^3)))/3`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 39 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`

rule 101 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3) Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))] + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`

rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 6837 `Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*((d_)*(x_))^(m_), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcSech[c*x])/(d*(m + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(m + 1))*Sqrt[1/(1 + c*x)] Int[(d*x)^m/(Sqrt[1 - c*x]*Sqrt[1 + c*x]), x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]`



**Maple [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.95

method	result	size
parts	$\frac{x^3 a}{3} + \frac{b \left( \frac{c^3 x^3 \operatorname{arcsech}(cx)}{3} + \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (-cx \sqrt{-c^2 x^2 + 1} + \arcsin(cx))}{6 \sqrt{-c^2 x^2 + 1}} \right)}{c^3}$	92
derivativedivides	$\frac{a c^3 x^3}{3} + b \left( \frac{c^3 x^3 \operatorname{arcsech}(cx)}{3} + \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (-cx \sqrt{-c^2 x^2 + 1} + \arcsin(cx))}{6 \sqrt{-c^2 x^2 + 1}} \right)$	96
default	$\frac{a c^3 x^3}{3} + b \left( \frac{c^3 x^3 \operatorname{arcsech}(cx)}{3} + \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (-cx \sqrt{-c^2 x^2 + 1} + \arcsin(cx))}{6 \sqrt{-c^2 x^2 + 1}} \right)$	96

input `int(x^2*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{3} x^3 a + \frac{b}{c^3} \left( \frac{1}{3} c^3 x^3 \operatorname{arcsech}(cx) + \frac{1}{6} \left( -\frac{cx-1}{cx} \right)^{1/2} cx \left( \frac{cx+1}{cx} \right)^{1/2} (-cx \sqrt{-c^2 x^2 + 1} + \arcsin(cx)) \right) / \sqrt{-c^2 x^2 + 1}$$

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.67

$$\int x^2 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{2 a c^3 x^3 - b c^2 x^2 \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - 2 b c^3 \log \left( \frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - 1}{x} \right) - 2 b \arctan \left( \frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - 1}{cx} \right) + 2 (b c^3 x^3 - b c^3) \log \left( \frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx} \right)}{6 c^3}$$

input `integrate(x^2*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output 
$$\frac{1}{6} (2 a c^3 x^3 - b c^2 x^2 \sqrt{-(c^2 x^2 - 1)/(c^2 x^2)} - 2 b c^3 \log \left( \frac{cx \sqrt{-(c^2 x^2 - 1)/(c^2 x^2)} - 1}{cx} \right) - 2 b \arctan \left( \frac{cx \sqrt{-(c^2 x^2 - 1)/(c^2 x^2)} - 1}{cx} \right) + 2 (b c^3 x^3 - b c^3) \log \left( \frac{cx \sqrt{-(c^2 x^2 - 1)/(c^2 x^2)} + 1}{cx} \right)) / c^3$$

**Sympy [F]**

$$\int x^2(a + b \operatorname{sech}^{-1}(cx)) dx = \int x^2(a + b \operatorname{arsech}(cx)) dx$$

input `integrate(x**2*(a+b*arsech(c*x)),x)`

output `Integral(x**2*(a + b*arsech(c*x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.75

$$\int x^2(a + b \operatorname{sech}^{-1}(cx)) dx = \frac{1}{3} ax^3 + \frac{1}{6} \left( 2x^3 \operatorname{arsech}(cx) - \frac{\frac{\sqrt{\frac{1}{c^2x^2}-1}}{c^2(\frac{1}{c^2x^2}-1)+c^2} + \frac{\arctan(\sqrt{\frac{1}{c^2x^2}-1})}{c^2}}{c} \right) b$$

input `integrate(x^2*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `1/3*a*x^3 + 1/6*(2*x^3*arcsech(c*x) - (sqrt(1/(c^2*x^2) - 1)/(c^2*(1/(c^2*x^2) - 1) + c^2) + arctan(sqrt(1/(c^2*x^2) - 1))/c^2)/c)*b`

**Giac [F]**

$$\int x^2(a + b \operatorname{sech}^{-1}(cx)) dx = \int (b \operatorname{arsech}(cx) + a)x^2 dx$$

input `integrate(x^2*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^2 \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int(x^2*(a + b*acosh(1/(c*x))),x)`output `int(x^2*(a + b*acosh(1/(c*x))), x)`**Reduce [F]**

$$\int x^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \left( \int a \operatorname{sech}(cx) x^2 dx \right) b + \frac{a x^3}{3}$$

input `int(x^2*(a+b*asech(c*x)),x)`output `(3*int(asech(c*x)*x**2,x)*b + a*x**3)/3`

### 3.24 $\int x(a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	259
Mathematica [A] (verified)	259
Rubi [A] (verified)	260
Maple [A] (verified)	261
Fricas [A] (verification not implemented)	261
Sympy [A] (verification not implemented)	262
Maxima [A] (verification not implemented)	262
Giac [F]	262
Mupad [B] (verification not implemented)	263
Reduce [F]	263

#### Optimal result

Integrand size = 10, antiderivative size = 47

$$\int x(a + b\operatorname{sech}^{-1}(cx)) dx = -\frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{2c^2} + \frac{1}{2}x^2(a + b\operatorname{sech}^{-1}(cx))$$

output

```
-1/2*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/c^2+1/2*x^2*(a+b*arcsech(c*x))
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.21

$$\int x(a + b\operatorname{sech}^{-1}(cx)) dx = \frac{ax^2}{2} + b\left(-\frac{1}{2c^2} - \frac{x}{2c}\right)\sqrt{\frac{1-cx}{1+cx}} + \frac{1}{2}bx^2\operatorname{sech}^{-1}(cx)$$

input

```
Integrate[x*(a + b*ArcSech[c*x]),x]
```

output

```
(a*x^2)/2 + b*(-1/2*1/c^2 - x/(2*c))*Sqrt[(1 - c*x)/(1 + c*x)] + (b*x^2*ArcSech[c*x])/2
```

**Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.96, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {6837, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(a + b \operatorname{sech}^{-1}(cx)) dx$$

↓ 6837

$$\frac{1}{2}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x}{\sqrt{1-cx}\sqrt{cx+1}} dx + \frac{1}{2}x^2(a + b \operatorname{sech}^{-1}(cx))$$

↓ 83

$$\frac{1}{2}x^2(a + b \operatorname{sech}^{-1}(cx)) - \frac{b\sqrt{1-cx}}{2c^2\sqrt{\frac{1}{cx+1}}}$$

input `Int[x*(a + b*ArcSech[c*x]),x]`

output `-1/2*(b*Sqrt[1 - c*x])/(c^2*Sqrt[(1 + c*x)^(-1)]) + (x^2*(a + b*ArcSech[c*x]))/2`

**Defintions of rubi rules used**

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

rule 6837 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcSech[c*x])/(d*(m + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(m + 1))*Sqrt[1/(1 + c*x)] Int[(d*x)^m/(Sqrt[1 - c*x]*Sqrt[1 + c*x]), x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]`

**Maple [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.26

method	result	size
parts	$\frac{ax^2}{2} + \frac{b \left( \frac{c^2 x^2 \operatorname{arcsech}(cx)}{2} - \sqrt{-\frac{cx-1}{cx}} \frac{c \sqrt{\frac{cx+1}{cx}} x}{2} \right)}{c^2}$	59
derivativedivides	$\frac{a \frac{c^2 x^2}{2} + b \left( \frac{c^2 x^2 \operatorname{arcsech}(cx)}{2} - \sqrt{-\frac{cx-1}{cx}} \frac{c \sqrt{\frac{cx+1}{cx}} x}{2} \right)}{c^2}$	63
default	$\frac{a \frac{c^2 x^2}{2} + b \left( \frac{c^2 x^2 \operatorname{arcsech}(cx)}{2} - \sqrt{-\frac{cx-1}{cx}} \frac{c \sqrt{\frac{cx+1}{cx}} x}{2} \right)}{c^2}$	63

input `int(x*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)`

output  $\frac{1}{2}ax^2 + \frac{b}{c^2} \left( \frac{1}{2}c^2 x^2 \operatorname{arcsech}(cx) - \frac{1}{2} \left( -\frac{cx-1}{cx} \right)^{1/2} c \left( \frac{cx+1}{cx} \right)^{1/2} x \right)$

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.55

$$\int x(a + b \operatorname{sech}^{-1}(cx)) dx = \frac{bcx^2 \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx}\right) + acx^2 - bx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}}}{2c}$$

input `integrate(x*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output  $\frac{1}{2}(b * c * x^2 * \log((c * x * \sqrt{-(c^2 * x^2 - 1)/(c^2 * x^2)} + 1)/(c * x)) + a * c * x^2 - b * x * \sqrt{-(c^2 * x^2 - 1)/(c^2 * x^2)})/c$

**Sympy [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.98

$$\int x(a + b \operatorname{sech}^{-1}(cx)) dx = \begin{cases} \frac{ax^2}{2} + \frac{bx^2 \operatorname{arsech}(cx)}{2} - \frac{b\sqrt{-c^2x^2+1}}{2c^2} & \text{for } c \neq 0 \\ \frac{x^2(a+\infty b)}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*(a+b*asech(c*x)),x)`output `Piecewise((a*x**2/2 + b*x**2*asech(c*x)/2 - b*sqrt(-c**2*x**2 + 1)/(2*c**2), Ne(c, 0)), (x**2*(a + oo*b)/2, True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.77

$$\int x(a + b \operatorname{sech}^{-1}(cx)) dx = \frac{1}{2} ax^2 + \frac{1}{2} \left( x^2 \operatorname{arsech}(cx) - \frac{x\sqrt{\frac{1}{c^2x^2} - 1}}{c} \right) b$$

input `integrate(x*(a+b*arcsech(c*x)),x, algorithm="maxima")`output `1/2*a*x^2 + 1/2*(x^2*arcsech(c*x) - x*sqrt(1/(c^2*x^2) - 1)/c)*b`**Giac [F]**

$$\int x(a + b \operatorname{sech}^{-1}(cx)) dx = \int (b \operatorname{arsech}(cx) + a)x dx$$

input `integrate(x*(a+b*arcsech(c*x)),x, algorithm="giac")`output `integrate((b*arcsech(c*x) + a)*x, x)`

**Mupad [B] (verification not implemented)**

Time = 3.57 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.06

$$\int x(a + b \operatorname{sech}^{-1}(cx)) dx = \frac{ax^2}{2} + \frac{bx^2 \operatorname{acosh}\left(\frac{1}{cx}\right)}{2} - \frac{bx \sqrt{\frac{1}{cx} - 1} \sqrt{\frac{1}{cx} + 1}}{2c}$$

input `int(x*(a + b*acosh(1/(c*x))),x)`output `(a*x^2)/2 + (b*x^2*acosh(1/(c*x)))/2 - (b*x*(1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2))/(2*c)`**Reduce [F]**

$$\int x(a + b \operatorname{sech}^{-1}(cx)) dx = \left( \int a \operatorname{sech}(cx) x dx \right) b + \frac{ax^2}{2}$$

input `int(x*(a+b*asech(c*x)),x)`output `(2*int(asech(c*x)*x,x)*b + a*x**2)/2`



### 3.25 $\int (a + b \operatorname{sech}^{-1}(cx)) dx$

Optimal result	264
Mathematica [A] (verified)	264
Rubi [A] (warning: unable to verify)	265
Maple [A] (verified)	265
Fricas [B] (verification not implemented)	266
Sympy [F]	266
Maxima [A] (verification not implemented)	267
Giac [F]	267
Mupad [B] (verification not implemented)	267
Reduce [F]	268

#### Optimal result

Integrand size = 8, antiderivative size = 57

$$\int (a + b \operatorname{sech}^{-1}(cx)) dx = ax + bx \operatorname{sech}^{-1}(cx) + \frac{b \sqrt{\frac{1-cx}{1+cx}} (1+cx) \arcsin(cx)}{c \sqrt{1-c^2x^2}}$$

output

```
a*x+b*x*arcsech(c*x)+b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*arcsin(c*x)/c/(-c^2*x^2+1)^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.40

$$\int (a + b \operatorname{sech}^{-1}(cx)) dx = ax + bx \operatorname{sech}^{-1}(cx) + \frac{2b \sqrt{\frac{1-cx}{1+cx}} \sqrt{1-c^2x^2} \arctan\left(\frac{\sqrt{1-c^2x^2}}{1-cx}\right)}{c - c^2x}$$

input

```
Integrate[a + b*ArcSech[c*x], x]
```

output

```
a*x + b*x*ArcSech[c*x] + (2*b*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*ArcTan[Sqrt[1 - c^2*x^2]/(1 - c*x)]/(c - c^2*x)
```

### Rubi [A] (warning: unable to verify)

Time = 0.18 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.70, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{sech}^{-1}(cx)) dx$$

↓ 2009

$$ax + \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \arcsin(cx)}{c} + bx \operatorname{sech}^{-1}(cx)$$

input `Int[a + b*ArcSech[c*x], x]`

output `a*x + b*x*ArcSech[c*x] + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*ArcSin[c*x])/c`

#### Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

### Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.74

method	result	size
default	$xa + bx \operatorname{arcsech}(cx) - \frac{b \arctan\left(\sqrt{-1+\frac{1}{cx}} \sqrt{1+\frac{1}{cx}}\right)}{c}$	42
parts	$xa + bx \operatorname{arcsech}(cx) - \frac{b \arctan\left(\sqrt{-1+\frac{1}{cx}} \sqrt{1+\frac{1}{cx}}\right)}{c}$	42
derivativedivides	$\frac{cxa+b\left(cx \operatorname{arcsech}(cx)-\arctan\left(\sqrt{-1+\frac{1}{cx}} \sqrt{1+\frac{1}{cx}}\right)\right)}{c}$	46

input `int(a+b*arcsech(c*x),x,method=_RETURNVERBOSE)`

output `x*a+b*x*arcsech(c*x)-b/c*arctan((-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs.  $2(53) = 106$ .

Time = 0.10 (sec) , antiderivative size = 119, normalized size of antiderivative = 2.09

$$\int (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{acx - bc \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{x}\right) - 2b \arctan\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{cx}\right) + (bcx - bc) \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}+1}{cx}\right)}{c}$$

input `integrate(a+b*arcsech(c*x),x, algorithm="fricas")`

output `(a*c*x - b*c*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) - 2*b*arctan((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/(c*x)) + (b*c*x - b*c)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)))/c`

### Sympy [F]

$$\int (a + b \operatorname{sech}^{-1}(cx)) dx = \int (a + b \operatorname{asech}(cx)) dx$$

input `integrate(a+b*asech(c*x),x)`

output `Integral(a + b*asech(c*x), x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.54

$$\int (a + b \operatorname{sech}^{-1}(cx)) dx = ax + \frac{(cx \operatorname{ar} \operatorname{sech}(cx) - \arctan(\sqrt{\frac{1}{c^2 x^2} - 1}))b}{c}$$

input `integrate(a+b*arcsech(c*x),x, algorithm="maxima")`

output `a*x + (c*x*arcsech(c*x) - arctan(sqrt(1/(c^2*x^2) - 1)))*b/c`

**Giac [F]**

$$\int (a + b \operatorname{sech}^{-1}(cx)) dx = \int b \operatorname{ar} \operatorname{sech}(cx) + a dx$$

input `integrate(a+b*arcsech(c*x),x, algorithm="giac")`

output `integrate(b*arcsech(c*x) + a, x)`

**Mupad [B] (verification not implemented)**

Time = 3.41 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.77

$$\int (a + b \operatorname{sech}^{-1}(cx)) dx = ax + bx \operatorname{acosh}\left(\frac{1}{cx}\right) + \frac{b \operatorname{atan}\left(\frac{1}{\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}}\right)}{c}$$

input `int(a + b*acosh(1/(c*x)),x)`

output `a*x + b*x*acosh(1/(c*x)) + (b*atan(1/((1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2))))/c`

**Reduce [F]**

$$\int (a + b \operatorname{sech}^{-1}(cx)) dx = \left( \int a \operatorname{sech}(cx) dx \right) b + ax$$

input `int(a+b*asech(c*x),x)`

output `int(asech(c*x),x)*b + a*x`

### 3.26 $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x} dx$

Optimal result	269
Mathematica [A] (verified)	269
Rubi [C] (warning: unable to verify)	270
Maple [A] (verified)	273
Fricas [F]	273
Sympy [F]	274
Maxima [F]	274
Giac [F]	274
Mupad [F(-1)]	275
Reduce [F]	275

#### Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x} dx = \frac{(a + b\operatorname{sech}^{-1}(cx))^2}{2b} - (a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + e^{2\operatorname{sech}^{-1}(cx)}\right) - \frac{1}{2}b \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(cx)}\right)$$

output

```
1/2*(a+b*arcsech(c*x))^2/b-(a+b*arcsech(c*x))*ln(1+(1/c/x+(-1+1/c/x)^(1/2)
*(1+1/c/x)^(1/2))^2)-1/2*b*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1
/2))^2)
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.84

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x} dx = a \log(x) + \frac{1}{2}b \left( -\operatorname{sech}^{-1}(cx) \left( \operatorname{sech}^{-1}(cx) + 2 \log\left(1 + e^{-2\operatorname{sech}^{-1}(cx)}\right) \right) + \operatorname{PolyLog}\left(2, -e^{-2\operatorname{sech}^{-1}(cx)}\right) \right)$$

input `Integrate[(a + b*ArcSech[c*x])/x,x]`

output `a*Log[x] + (b*(-(ArcSech[c*x]*(ArcSech[c*x] + 2*Log[1 + E^(-2*ArcSech[c*x])])) + PolyLog[2, -E^(-2*ArcSech[c*x])]))/2`

### Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.53 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.39, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {6835, 6297, 25, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{x} dx \\
 & \quad \downarrow \text{6835} \\
 & - \int x \left( a + b \operatorname{arccosh}\left(\frac{1}{cx}\right) \right) d\frac{1}{x} \\
 & \quad \downarrow \text{6297} \\
 & \frac{\int - \left( \left( a + b \operatorname{arccosh}\left(\frac{1}{cx}\right) \right) \tanh\left(\frac{a}{b} - \frac{a + b \operatorname{arccosh}\left(\frac{1}{cx}\right)}{b}\right) \right) d\left( a + b \operatorname{arccosh}\left(\frac{1}{cx}\right) \right)}{b} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \left( a + b \operatorname{arccosh}\left(\frac{1}{cx}\right) \right) \tanh\left(\frac{a}{b} - \frac{a + b \operatorname{arccosh}\left(\frac{1}{cx}\right)}{b}\right) d\left( a + b \operatorname{arccosh}\left(\frac{1}{cx}\right) \right)}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int -i \left( a + b \operatorname{arccosh}\left(\frac{1}{cx}\right) \right) \tan\left(\frac{ia}{b} - \frac{i \left( a + b \operatorname{arccosh}\left(\frac{1}{cx}\right) \right)}{b}\right) d\left( a + b \operatorname{arccosh}\left(\frac{1}{cx}\right) \right)}{b} \\
 & \quad \downarrow \text{26}
 \end{aligned}$$

$$\frac{i \int (a + \operatorname{barccosh}(\frac{1}{cx})) \tan \left( \frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(\frac{1}{cx}))}{b} \right) d(a + \operatorname{barccosh}(\frac{1}{cx}))}{b}$$

↓ 4201

$$\frac{i \left( 2i \int \frac{e^{-2\operatorname{arccosh}(\frac{1}{cx})} (a + \operatorname{barccosh}(\frac{1}{cx}))}{1 + e^{-2\operatorname{arccosh}(\frac{1}{cx})}} d(a + \operatorname{barccosh}(\frac{1}{cx})) - \frac{i}{2x^2} \right)}{b}$$

↓ 2620

$$\frac{i \left( 2i \left( \frac{1}{2} b \int \log \left( 1 + e^{-2\operatorname{arccosh}(\frac{1}{cx})} \right) d(a + \operatorname{barccosh}(\frac{1}{cx})) - \frac{1}{2} b \log \left( e^{-2\operatorname{arccosh}(\frac{1}{cx})} + 1 \right) (a + \operatorname{barccosh}(\frac{1}{cx})) \right) - \frac{i}{2x^2} \right)}{b}$$

↓ 2715

$$\frac{i \left( 2i \left( -\frac{1}{4} b^2 \int x \log \left( 1 + e^{-2\operatorname{arccosh}(\frac{1}{cx})} \right) d e^{-2\operatorname{arccosh}(\frac{1}{cx})} - \frac{1}{2} b \log \left( e^{-2\operatorname{arccosh}(\frac{1}{cx})} + 1 \right) (a + \operatorname{barccosh}(\frac{1}{cx})) \right) - \frac{i}{2x^2} \right)}{b}$$

↓ 2838

$$\frac{i \left( 2i \left( \frac{1}{4} b^2 \operatorname{PolyLog} \left( 2, -a - \operatorname{barccosh}(\frac{1}{cx}) \right) - \frac{1}{2} b \log \left( e^{-2\operatorname{arccosh}(\frac{1}{cx})} + 1 \right) (a + \operatorname{barccosh}(\frac{1}{cx})) \right) - \frac{i}{2x^2} \right)}{b}$$

input `Int[(a + b*ArcSech[c*x])/x,x]`

output `((-I)*((-1/2*I)/x^2 + (2*I)*(-1/2*(b*(a + b*ArcCosh[1/(c*x)])*Log[1 + E^(-2*ArcCosh[1/(c*x)]]) + (b^2*PolyLog[2, -a - b*ArcCosh[1/(c*x)]])/4)))/b`

**Defintions of rubi rules used**

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`



rule 2620  $\text{Int}[\frac{((F_{-})^{((g_{-}) * (e_{-}) + (f_{-}) * (x_{-})))^{(n_{-})} * ((c_{-}) + (d_{-}) * (x_{-}))^{(m_{-})})}{((a_{-}) + (b_{-}) * (F_{-})^{((g_{-}) * (e_{-}) + (f_{-}) * (x_{-})))^{(n_{-})})}, x\_Symbol] \rightarrow \text{Simp}[\frac{(c + d*x)^m}{(b*f*g*n*\text{Log}[F])} * \text{Log}[1 + b*((F^{(g*(e + f*x)))^n/a}], x] - \text{Simp}[d*(m/(b*f*g*n*\text{Log}[F])) \text{Int}[(c + d*x)^{m-1} * \text{Log}[1 + b*((F^{(g*(e + f*x)))^n/a}], x], x] /; \text{FreeQ}\{F, a, b, c, d, e, f, g, n\}, x\} \&\& \text{IGtQ}[m, 0]$

rule 2715  $\text{Int}[\text{Log}[(a_{-}) + (b_{-}) * (F_{-})^{((e_{-}) * ((c_{-}) + (d_{-}) * (x_{-})))^{(n_{-})}], x\_Symbol] \rightarrow \text{Simp}[1/(d*e*n*\text{Log}[F]) \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^{(e*(c + d*x))})^n], x] /; \text{FreeQ}\{F, a, b, c, d, e, n\}, x\} \&\& \text{GtQ}[a, 0]$

rule 2838  $\text{Int}[\text{Log}[(c_{-}) * ((d_{-}) + (e_{-}) * (x_{-})^{(n_{-})})]/(x_{-}), x\_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n]/n, x] /; \text{FreeQ}\{c, d, e, n\}, x\} \&\& \text{EqQ}[c*d, 1]$

rule 3042  $\text{Int}[u_{-}, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4201  $\text{Int}[\frac{((c_{-}) + (d_{-}) * (x_{-}))^{(m_{-})} * \tan[(e_{-}) + (\text{Complex}[0, fz_{-}]) * (f_{-}) * (x_{-})], x\_Symbol] \rightarrow \text{Simp}[(-I) * ((c + d*x)^{m+1} / (d*(m+1))), x] + \text{Simp}[2*I \text{Int}[(c + d*x)^m * (E^{(2*((-I)*e + f*fz*x))} / (1 + E^{(2*((-I)*e + f*fz*x))}))], x], x] /; \text{FreeQ}\{c, d, e, f, fz\}, x\} \&\& \text{IGtQ}[m, 0]$

rule 6297  $\text{Int}[\frac{((a_{-}) + \text{ArcCosh}[(c_{-}) * (x_{-})] * (b_{-}))^{(n_{-})}}{(x_{-})}, x\_Symbol] \rightarrow \text{Simp}[1/b \text{Subst}[\text{Int}[x^n * \text{Tanh}[-a/b + x/b], x], x, a + b*\text{ArcCosh}[c*x]], x] /; \text{FreeQ}\{a, b, c\}, x\} \&\& \text{IGtQ}[n, 0]$

rule 6835  $\text{Int}[\frac{((a_{-}) + \text{ArcSech}[(c_{-}) * (x_{-})] * (b_{-}))}{(x_{-})}, x\_Symbol] \rightarrow -\text{Subst}[\text{Int}[(a + b*\text{ArcCosh}[x/c])/x, x], x, 1/x] /; \text{FreeQ}\{a, b, c\}, x\}$

**Maple [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.75

method	result
parts	$a \ln(x) + b \left( \frac{\operatorname{arcsech}(cx)^2}{2} - \operatorname{arcsech}(cx) \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right)^2 \right) - \frac{\operatorname{polylog}}{\dots} \right)$
derivativedivides	$a \ln(cx) + b \left( \frac{\operatorname{arcsech}(cx)^2}{2} - \operatorname{arcsech}(cx) \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right)^2 \right) - \frac{\operatorname{polylog}}{\dots} \right)$
default	$a \ln(cx) + b \left( \frac{\operatorname{arcsech}(cx)^2}{2} - \operatorname{arcsech}(cx) \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right)^2 \right) - \frac{\operatorname{polylog}}{\dots} \right)$

input `int((a+b*arcsech(c*x))/x,x,method=_RETURNVERBOSE)`

output `a*ln(x)+b*(1/2*arcsech(c*x)^2-arcsech(c*x)*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)-1/2*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2))`

**Fricas [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x} dx = \int \frac{b \operatorname{ar} \operatorname{sech}(cx) + a}{x} dx$$

input `integrate((a+b*arcsech(c*x))/x,x, algorithm="fricas")`

output `integral((b*arcsech(c*x) + a)/x, x)`

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x} dx = \int \frac{a + b \operatorname{arsech}(cx)}{x} dx$$

input `integrate((a+b*asech(c*x))/x,x)`

output `Integral((a + b*asech(c*x))/x, x)`

**Maxima [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x} dx = \int \frac{b \operatorname{arsech}(cx) + a}{x} dx$$

input `integrate((a+b*arcsech(c*x))/x,x, algorithm="maxima")`

output `b*integrate(log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/x, x) + a*log(x)`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x} dx = \int \frac{b \operatorname{arsech}(cx) + a}{x} dx$$

input `integrate((a+b*arcsech(c*x))/x,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/x, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x} dx$$

input `int((a + b*acosh(1/(c*x)))/x,x)`output `int((a + b*acosh(1/(c*x)))/x, x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x} dx = \left( \int \frac{a \operatorname{sech}(cx)}{x} dx \right) b + \log(x) a$$

input `int((a+b*asech(c*x))/x,x)`output `int(asech(c*x)/x,x)*b + log(x)*a`

### 3.27 $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^2} dx$

Optimal result	276
Mathematica [A] (verified)	276
Rubi [A] (verified)	277
Maple [A] (verified)	278
Fricas [A] (verification not implemented)	278
Sympy [F]	279
Maxima [A] (verification not implemented)	279
Giac [F]	279
Mupad [B] (verification not implemented)	280
Reduce [F]	280

#### Optimal result

Integrand size = 12, antiderivative size = 42

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^2} dx = \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{x} - \frac{a + b\operatorname{sech}^{-1}(cx)}{x}$$

output `b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/x-(a+b*arcsech(c*x))/x`

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^2} dx = -\frac{a}{x} + b\left(c + \frac{1}{x}\right) \sqrt{\frac{1-cx}{1+cx}} - \frac{b\operatorname{sech}^{-1}(cx)}{x}$$

input `Integrate[(a + b*ArcSech[c*x])/x^2,x]`

output `-(a/x) + b*(c + x^(-1))*Sqrt[(1 - c*x)/(1 + c*x)] - (b*ArcSech[c*x])/x`

**Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.95, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {6837, 106}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2} dx$$

↓ 6837

$$-b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{1}{x^2 \sqrt{1-cx} \sqrt{cx+1}} dx - \frac{a + b \operatorname{sech}^{-1}(cx)}{x}$$

↓ 106

$$\frac{b \sqrt{1-cx}}{x \sqrt{\frac{1}{cx+1}}} - \frac{a + b \operatorname{sech}^{-1}(cx)}{x}$$

input `Int[(a + b*ArcSech[c*x])/x^2,x]`

output `(b*Sqrt[1 - c*x])/(x*Sqrt[(1 + c*x)^(-1)]) - (a + b*ArcSech[c*x])/x`

**Defintions of rubi rules used**

rule 106

```
Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && EqQ[a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1), 0] && NeQ[m, -1]
```

rule 6837

```
Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))*((d_.)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcSech[c*x])/(d*(m + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(m + 1))*Sqrt[1/(1 + c*x)] Int[(d*x)^m/(Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

**Maple [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.29

method	result	size
parts	$-\frac{a}{x} + bc\left(-\frac{\operatorname{arcsech}(cx)}{cx} + \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}\right)$	54
derivativedivides	$c\left(-\frac{a}{cx} + b\left(-\frac{\operatorname{arcsech}(cx)}{cx} + \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}\right)\right)$	58
default	$c\left(-\frac{a}{cx} + b\left(-\frac{\operatorname{arcsech}(cx)}{cx} + \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}\right)\right)$	58

input `int((a+b*arcsech(c*x))/x^2,x,method=_RETURNVERBOSE)`

output `-a/x+b*c*(-1/c/x*arcsech(c*x)+(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.57

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2} dx = \frac{bcx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - b \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx}\right) - a}{x}$$

input `integrate((a+b*arcsech(c*x))/x^2,x, algorithm="fricas")`

output `(b*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - b*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - a)/x`

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2} dx = \int \frac{a + b \operatorname{arsech}(cx)}{x^2} dx$$

input `integrate((a+b*arsech(c*x))/x**2,x)`

output `Integral((a + b*arsech(c*x))/x**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.76

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2} dx = \left( c \sqrt{\frac{1}{c^2 x^2} - 1} - \frac{\operatorname{arsech}(cx)}{x} \right) b - \frac{a}{x}$$

input `integrate((a+b*arcsech(c*x))/x^2,x, algorithm="maxima")`

output `(c*sqrt(1/(c^2*x^2) - 1) - arcsech(c*x)/x)*b - a/x`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2} dx = \int \frac{b \operatorname{arsech}(cx) + a}{x^2} dx$$

input `integrate((a+b*arcsech(c*x))/x^2,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/x^2, x)`



**Mupad [B] (verification not implemented)**

Time = 3.59 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.10

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2} dx = bc \sqrt{\frac{1}{cx} - 1} \sqrt{\frac{1}{cx} + 1} - \frac{b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x} - \frac{a}{x}$$

input `int((a + b*acosh(1/(c*x)))/x^2,x)`output `b*c*(1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2) - (b*acosh(1/(c*x)))/x - a/x`**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2} dx = \frac{\left(\int \frac{a \operatorname{sech}(cx)}{x^2} dx\right) bx - a}{x}$$

input `int((a+b*asech(c*x))/x^2,x)`output `(int(asech(c*x)/x**2,x)*b*x - a)/x`

### 3.28 $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^3} dx$

Optimal result	281
Mathematica [A] (verified)	281
Rubi [A] (verified)	282
Maple [A] (verified)	284
Fricas [A] (verification not implemented)	285
Sympy [F]	285
Maxima [A] (verification not implemented)	286
Giac [F]	286
Mupad [B] (verification not implemented)	287
Reduce [F]	287

#### Optimal result

Integrand size = 12, antiderivative size = 107

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^3} dx = \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{4x^2} - \frac{a + b\operatorname{sech}^{-1}(cx)}{2x^2} + \frac{bc^2\sqrt{\frac{1-cx}{1+cx}}(1+cx)\operatorname{arctanh}(\sqrt{1-c^2x^2})}{4\sqrt{1-c^2x^2}}$$

output `1/4*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/x^2-1/2*(a+b*arcsech(c*x))/x^2+1/4*b*c^2*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*arctanh((-c^2*x^2+1)^(1/2))/(-c^2*x^2+1)^(1/2)`

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.09

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^3} dx = -\frac{a}{2x^2} + b\left(\frac{1}{4x^2} + \frac{c}{4x}\right)\sqrt{\frac{1-cx}{1+cx}} - \frac{b\operatorname{sech}^{-1}(cx)}{2x^2} - \frac{1}{4}bc^2\log(x) + \frac{1}{4}bc^2\log\left(1 + \sqrt{\frac{1-cx}{1+cx}} + cx\sqrt{\frac{1-cx}{1+cx}}\right)$$

input `Integrate[(a + b*ArcSech[c*x])/x^3,x]`

output `-1/2*a/x^2 + b*(1/(4*x^2) + c/(4*x))*Sqrt[(1 - c*x)/(1 + c*x)] - (b*ArcSech[c*x])/(2*x^2) - (b*c^2*Log[x])/4 + (b*c^2*Log[1 + Sqrt[(1 - c*x)/(1 + c*x)]] + c*x*Sqrt[(1 - c*x)/(1 + c*x)])/4`

### Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.90, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6837, 114, 25, 27, 103, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3} dx \\
 & \quad \downarrow 6837 \\
 & -\frac{1}{2}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{x^3\sqrt{1-cx}\sqrt{cx+1}} dx - \frac{a + b \operatorname{sech}^{-1}(cx)}{2x^2} \\
 & \quad \downarrow 114 \\
 & -\frac{1}{2}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{1}{2} \int -\frac{c^2}{x\sqrt{1-cx}\sqrt{cx+1}} dx - \frac{\sqrt{1-cx}\sqrt{cx+1}}{2x^2} \right) - \frac{a + b \operatorname{sech}^{-1}(cx)}{2x^2} \\
 & \quad \downarrow 25 \\
 & -\frac{1}{2}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{2} \int \frac{c^2}{x\sqrt{1-cx}\sqrt{cx+1}} dx - \frac{\sqrt{1-cx}\sqrt{cx+1}}{2x^2} \right) - \frac{a + b \operatorname{sech}^{-1}(cx)}{2x^2} \\
 & \quad \downarrow 27 \\
 & -\frac{1}{2}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{2}c^2 \int \frac{1}{x\sqrt{1-cx}\sqrt{cx+1}} dx - \frac{\sqrt{1-cx}\sqrt{cx+1}}{2x^2} \right) - \frac{a + b \operatorname{sech}^{-1}(cx)}{2x^2} \\
 & \quad \downarrow 103
 \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{2}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{1}{2}c^3\int\frac{1}{c-c(1-cx)(cx+1)}d\left(\sqrt{1-cx}\sqrt{cx+1}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{2x^2}\right)- \\
& \qquad \qquad \qquad \frac{a+b\operatorname{sech}^{-1}(cx)}{2x^2} \\
& \qquad \qquad \qquad \downarrow \text{221} \\
& \qquad \qquad \qquad -\frac{a+b\operatorname{sech}^{-1}(cx)}{2x^2}- \\
& \frac{1}{2}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{1}{2}c^2\operatorname{arctanh}\left(\sqrt{1-cx}\sqrt{cx+1}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{2x^2}\right)
\end{aligned}$$

input `Int[(a + b*ArcSech[c*x])/x^3,x]`

output `-1/2*(a + b*ArcSech[c*x])/x^2 - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/2*(Sqrt[1 - c*x]*Sqrt[1 + c*x])/x^2 - (c^2*ArcTanh[Sqrt[1 - c*x]*Sqrt[1 + c*x]]/2))/2`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 103 `Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))), x_] :> Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]`

```
rule 114 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 6837 Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcSech[c*x])/(d*(m + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(m + 1))*Sqrt[1/(1 + c*x)] Int[(d*x)^m/(Sqrt[1 - c*x]*Sqrt[1 + c*x]), x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

**Maple [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.01

method	result	size
parts	$-\frac{a}{2x^2} + b c^2 \left( -\frac{\operatorname{arcsech}(cx)}{2c^2x^2} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \left( \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right) c^2x^2 + \sqrt{-c^2x^2+1} \right)}{4cx\sqrt{-c^2x^2+1}} \right)$	108
derivativedivides	$c^2 \left( -\frac{a}{2c^2x^2} + b \left( -\frac{\operatorname{arcsech}(cx)}{2c^2x^2} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \left( \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right) c^2x^2 + \sqrt{-c^2x^2+1} \right)}{4cx\sqrt{-c^2x^2+1}} \right) \right)$	112
default	$c^2 \left( -\frac{a}{2c^2x^2} + b \left( -\frac{\operatorname{arcsech}(cx)}{2c^2x^2} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \left( \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right) c^2x^2 + \sqrt{-c^2x^2+1} \right)}{4cx\sqrt{-c^2x^2+1}} \right) \right)$	112

```
input int((a+b*arcsech(c*x))/x^3,x,method=_RETURNVERBOSE)
```

output

```
-1/2*a/x^2+b*c^2*(-1/2/c^2/x^2*arcsech(c*x)+1/4*(-(c*x-1)/c/x)^(1/2)/c/x*(
(c*x+1)/c/x)^(1/2)*(arctanh(1/(-c^2*x^2+1)^(1/2))*c^2*x^2+(-c^2*x^2+1)^(1/
2))/(-c^2*x^2+1)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.72

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3} dx = \frac{bcx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + (bc^2 x^2 - 2b) \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx}\right) - 2a}{4x^2}$$

input

```
integrate((a+b*arcsech(c*x))/x^3,x, algorithm="fricas")
```

output

```
1/4*(b*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + (b*c^2*x^2 - 2*b)*log((c*x*sqrt
(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - 2*a)/x^2
```

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3} dx = \int \frac{a + b \operatorname{asech}(cx)}{x^3} dx$$

input

```
integrate((a+b*asech(c*x))/x**3,x)
```

output

```
Integral((a + b*asech(c*x))/x**3, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.98

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3} dx = -\frac{1}{8} b \left( \frac{2c^4 x \sqrt{\frac{1}{c^2 x^2} - 1}}{c^2 x^2 \left(\frac{1}{c^2 x^2} - 1\right)^{-1}} - c^3 \log \left( cx \sqrt{\frac{1}{c^2 x^2} - 1} + 1 \right) + c^3 \log \left( cx \sqrt{\frac{1}{c^2 x^2} - 1} - 1 \right) \right) + \frac{4 \operatorname{arsech}(cx)}{x^2} - \frac{a}{2x^2}$$

input `integrate((a+b*arcsech(c*x))/x^3,x, algorithm="maxima")`output `-1/8*b*((2*c^4*x*sqrt(1/(c^2*x^2) - 1)/(c^2*x^2*(1/(c^2*x^2) - 1) - 1) - c^3*log(c*x*sqrt(1/(c^2*x^2) - 1) + 1) + c^3*log(c*x*sqrt(1/(c^2*x^2) - 1) - 1))/c + 4*arcsech(c*x)/x^2) - 1/2*a/x^2`**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3} dx = \int \frac{b \operatorname{arsech}(cx) + a}{x^3} dx$$

input `integrate((a+b*arcsech(c*x))/x^3,x, algorithm="giac")`output `integrate((b*arcsech(c*x) + a)/x^3, x)`

**Mupad [B] (verification not implemented)**

Time = 3.48 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.57

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3} dx = \frac{b \operatorname{acosh}\left(\frac{1}{cx}\right) \left(\frac{c^2 x}{4} - \frac{1}{2x}\right)}{x} - \frac{a}{2x^2} + \frac{bc \sqrt{\frac{1}{cx} - 1} \sqrt{\frac{1}{cx} + 1}}{4x}$$

input `int((a + b*acosh(1/(c*x)))/x^3,x)`output `(b*acosh(1/(c*x))*((c^2*x)/4 - 1/(2*x)))/x - a/(2*x^2) + (b*c*(1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2))/(4*x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3} dx = \frac{2 \left( \int \frac{a \operatorname{sech}(cx)}{x^3} dx \right) b x^2 - a}{2x^2}$$

input `int((a+b*asech(c*x))/x^3,x)`output `(2*int(asech(c*x)/x**3,x)*b*x**2 - a)/(2*x**2)`



### 3.29 $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^4} dx$

Optimal result	288
Mathematica [A] (verified)	288
Rubi [A] (verified)	289
Maple [A] (verified)	290
Fricas [A] (verification not implemented)	291
Sympy [F]	291
Maxima [A] (verification not implemented)	292
Giac [F]	292
Mupad [F(-1)]	292
Reduce [F]	293

#### Optimal result

Integrand size = 12, antiderivative size = 81

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^4} dx = \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{9x^3} + \frac{2bc^2\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{9x} - \frac{a + b\operatorname{sech}^{-1}(cx)}{3x^3}$$

output  $\frac{1}{9}b*((-cx+1)/(cx+1))^{(1/2)}*(cx+1)/x^3+2/9*b*c^2*((-cx+1)/(cx+1))^{(1/2)}*(cx+1)/x-1/3*(a+b*\operatorname{arcsech}(cx))/x^3$

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.91

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^4} dx = -\frac{a}{3x^3} + b\left(\frac{2c^3}{9} + \frac{1}{9x^3} + \frac{c}{9x^2} + \frac{2c^2}{9x}\right)\sqrt{\frac{1-cx}{1+cx}} - \frac{b\operatorname{sech}^{-1}(cx)}{3x^3}$$

input `Integrate[(a + b*ArcSech[c*x])/x^4, x]`

output  $-\frac{1}{3}a/x^3 + b*((2*c^3)/9 + 1/(9*x^3) + c/(9*x^2) + (2*c^2)/(9*x))*\operatorname{Sqrt}[(1 - cx)/(1 + cx)] - (b*\operatorname{ArcSech}[c*x])/(3*x^3)$

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.20, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6837, 114, 27, 106}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4} dx \\
 & \quad \downarrow \text{6837} \\
 & -\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{x^4\sqrt{1-cx}\sqrt{cx+1}} dx - \frac{a + b \operatorname{sech}^{-1}(cx)}{3x^3} \\
 & \quad \downarrow \text{114} \\
 & -\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{1}{3} \int -\frac{2c^2}{x^2\sqrt{1-cx}\sqrt{cx+1}} dx - \frac{\sqrt{1-cx}\sqrt{cx+1}}{3x^3} \right) - \\
 & \quad \quad \quad \frac{a + b \operatorname{sech}^{-1}(cx)}{3x^3} \\
 & \quad \downarrow \text{27} \\
 & -\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{2}{3}c^2 \int \frac{1}{x^2\sqrt{1-cx}\sqrt{cx+1}} dx - \frac{\sqrt{1-cx}\sqrt{cx+1}}{3x^3} \right) - \frac{a + b \operatorname{sech}^{-1}(cx)}{3x^3} \\
 & \quad \downarrow \text{106} \\
 & -\frac{a + b \operatorname{sech}^{-1}(cx)}{3x^3} - \frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{2c^2\sqrt{1-cx}\sqrt{cx+1}}{3x} - \frac{\sqrt{1-cx}\sqrt{cx+1}}{3x^3} \right)
 \end{aligned}$$

input

```
Int[(a + b*ArcSech[c*x])/x^4,x]
```

output

```
-1/3*(b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/3*(Sqrt[1 - c*x]*Sqrt[1 + c*x])/x^3 - (2*c^2*Sqrt[1 - c*x]*Sqrt[1 + c*x])/(3*x))) - (a + b*ArcSech[c*x])/(3*x^3)
```

## Definitions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 106 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && EqQ[a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1), 0] && NeQ[m, -1]`

rule 114 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])`

rule 6837 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcSech[c*x])/(d*(m + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(m + 1))*Sqrt[1/(1 + c*x)] Int[(d*x)^m/(Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]`

## Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.90

method	result	size
parts	$-\frac{a}{3x^3} + bc^3 \left( -\frac{\operatorname{arcsech}(cx)}{3c^3x^3} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} (2c^2x^2+1)}{9c^2x^2} \right)$	73
derivativedivides	$c^3 \left( -\frac{a}{3c^3x^3} + b \left( -\frac{\operatorname{arcsech}(cx)}{3c^3x^3} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} (2c^2x^2+1)}{9c^2x^2} \right) \right)$	77
default	$c^3 \left( -\frac{a}{3c^3x^3} + b \left( -\frac{\operatorname{arcsech}(cx)}{3c^3x^3} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} (2c^2x^2+1)}{9c^2x^2} \right) \right)$	77

input `int((a+b*arcsech(c*x))/x^4,x,method=_RETURNVERBOSE)`

output `-1/3*a/x^3+b*c^3*(-1/3/c^3/x^3*arcsech(c*x)+1/9*(-(c*x-1)/c/x)^(1/2)/c^2/x^2*((c*x+1)/c/x)^(1/2)*(2*c^2*x^2+1))`

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.98

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4} dx = -\frac{3b \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}+1}{cx}\right) - (2bc^3x^3 + bcx)\sqrt{-\frac{c^2x^2-1}{c^2x^2}} + 3a}{9x^3}$$

input `integrate((a+b*arcsech(c*x))/x^4,x, algorithm="fricas")`

output `-1/9*(3*b*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (2*b*c^3*x^3 + b*c*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 3*a)/x^3`

### Sympy [F]

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4} dx = \int \frac{a + b \operatorname{asech}(cx)}{x^4} dx$$

input `integrate((a+b*asech(c*x))/x**4,x)`

output `Integral((a + b*asech(c*x))/x**4, x)`

**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.69

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4} dx = \frac{1}{9} b \left( \frac{c^4 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{3}{2}} + 3 c^4 \sqrt{\frac{1}{c^2 x^2} - 1}}{c} - \frac{3 \operatorname{ar} \operatorname{sech}(cx)}{x^3} \right) - \frac{a}{3 x^3}$$

input `integrate((a+b*arcsech(c*x))/x^4,x, algorithm="maxima")`output `1/9*b*((c^4*(1/(c^2*x^2) - 1)^(3/2) + 3*c^4*sqrt(1/(c^2*x^2) - 1))/c - 3*arcsech(c*x)/x^3) - 1/3*a/x^3`**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4} dx = \int \frac{b \operatorname{ar} \operatorname{sech}(cx) + a}{x^4} dx$$

input `integrate((a+b*arcsech(c*x))/x^4,x, algorithm="giac")`output `integrate((b*arcsech(c*x) + a)/x^4, x)`**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x^4} dx$$

input `int((a + b*acosh(1/(c*x)))/x^4,x)`output `int((a + b*acosh(1/(c*x)))/x^4, x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4} dx = \frac{3 \left( \int \frac{\operatorname{asech}(cx)}{x^4} dx \right) b x^3 - a}{3x^3}$$

input `int((a+b*asech(c*x))/x^4,x)`

output `(3*int(asech(c*x)/x**4,x)*b*x**3 - a)/(3*x**3)`

### 3.30 $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^5} dx$

Optimal result	294
Mathematica [A] (verified)	294
Rubi [A] (verified)	295
Maple [A] (verified)	298
Fricas [A] (verification not implemented)	298
Sympy [F]	299
Maxima [A] (verification not implemented)	299
Giac [F]	300
Mupad [F(-1)]	300
Reduce [F]	300

#### Optimal result

Integrand size = 12, antiderivative size = 141

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^5} dx = \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{16x^4} + \frac{3bc^2\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{32x^2} - \frac{a + b\operatorname{sech}^{-1}(cx)}{4x^4} + \frac{3bc^4\sqrt{\frac{1-cx}{1+cx}}(1+cx)\operatorname{arctanh}(\sqrt{1-c^2x^2})}{32\sqrt{1-c^2x^2}}$$

output

```
1/16*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/x^4+3/32*b*c^2*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/x^2-1/4*(a+b*arcsech(c*x))/x^4+3/32*b*c^4*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*arctanh((-c^2*x^2+1)^(1/2))/(-c^2*x^2+1)^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.97

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^5} dx = -\frac{a}{4x^4} + b\left(\frac{1}{16x^4} + \frac{c}{16x^3} + \frac{3c^2}{32x^2} + \frac{3c^3}{32x}\right)\sqrt{\frac{1-cx}{1+cx}} - \frac{b\operatorname{sech}^{-1}(cx)}{4x^4} - \frac{3}{32}bc^4\log(x) + \frac{3}{32}bc^4\log\left(1 + \sqrt{\frac{1-cx}{1+cx}} + cx\sqrt{\frac{1-cx}{1+cx}}\right)$$

input `Integrate[(a + b*ArcSech[c*x])/x^5,x]`

output `-1/4*a/x^4 + b*(1/(16*x^4) + c/(16*x^3) + (3*c^2)/(32*x^2) + (3*c^3)/(32*x))*Sqrt[(1 - c*x)/(1 + c*x)] - (b*ArcSech[c*x])/(4*x^4) - (3*b*c^4*Log[x])/32 + (3*b*c^4*Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]])/32`

### Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.92, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {6837, 114, 27, 114, 25, 27, 103, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5} dx \\
 & \quad \downarrow 6837 \\
 & -\frac{1}{4}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{x^5\sqrt{1-cx}\sqrt{cx+1}} dx - \frac{a + b \operatorname{sech}^{-1}(cx)}{4x^4} \\
 & \quad \downarrow 114 \\
 & -\frac{1}{4}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{1}{4} \int -\frac{3c^2}{x^3\sqrt{1-cx}\sqrt{cx+1}} dx - \frac{\sqrt{1-cx}\sqrt{cx+1}}{4x^4} \right) - \\
 & \quad \quad \quad \frac{a + b \operatorname{sech}^{-1}(cx)}{4x^4} \\
 & \quad \downarrow 27 \\
 & -\frac{1}{4}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3}{4}c^2 \int \frac{1}{x^3\sqrt{1-cx}\sqrt{cx+1}} dx - \frac{\sqrt{1-cx}\sqrt{cx+1}}{4x^4} \right) - \frac{a + b \operatorname{sech}^{-1}(cx)}{4x^4} \\
 & \quad \downarrow 114 \\
 & -\frac{1}{4}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3}{4}c^2 \left( -\frac{1}{2} \int -\frac{c^2}{x\sqrt{1-cx}\sqrt{cx+1}} dx - \frac{\sqrt{1-cx}\sqrt{cx+1}}{2x^2} \right) - \frac{\sqrt{1-cx}\sqrt{cx+1}}{4x^4} \right) -
 \end{aligned}$$



↓ 25

$$-\frac{1}{4}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{3}{4}c^2\left(\frac{1}{2}\int\frac{c^2}{x\sqrt{1-cx}\sqrt{cx+1}}dx-\frac{\sqrt{1-cx}\sqrt{cx+1}}{2x^2}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{4x^4}\right)-$$

$$\frac{a+b\operatorname{sech}^{-1}(cx)}{4x^4}$$

↓ 27

$$-\frac{1}{4}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{3}{4}c^2\left(\frac{1}{2}c^2\int\frac{1}{x\sqrt{1-cx}\sqrt{cx+1}}dx-\frac{\sqrt{1-cx}\sqrt{cx+1}}{2x^2}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{4x^4}\right)-$$

$$\frac{a+b\operatorname{sech}^{-1}(cx)}{4x^4}$$

↓ 103

$$-\frac{1}{4}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{3}{4}c^2\left(-\frac{1}{2}c^3\int\frac{1}{c-c(1-cx)(cx+1)}d(\sqrt{1-cx}\sqrt{cx+1})-\frac{\sqrt{1-cx}\sqrt{cx+1}}{2x^2}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{4x^4}\right)-$$

$$\frac{a+b\operatorname{sech}^{-1}(cx)}{4x^4}$$

↓ 221

$$-\frac{a+b\operatorname{sech}^{-1}(cx)}{4x^4}$$

$$\frac{1}{4}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{3}{4}c^2\left(-\frac{1}{2}c^2\operatorname{arctanh}\left(\sqrt{1-cx}\sqrt{cx+1}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{2x^2}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{4x^4}\right)-$$

input `Int[(a + b*ArcSech[c*x])/x^5,x]`

output `-1/4*(a + b*ArcSech[c*x])/x^4 - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/4*(Sqrt[1 - c*x]*Sqrt[1 + c*x])/x^4 + (3*c^2*(-1/2*(Sqrt[1 - c*x]*Sqrt[1 + c*x])/x^2 - (c^2*ArcTanh[Sqrt[1 - c*x]*Sqrt[1 + c*x]]/2))/4))/4`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 103 `Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]`
- rule 114 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 6837 `Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcSech[c*x])/(d*(m + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(m + 1))*Sqrt[1/(1 + c*x)] Int[(d*x)^m/(Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]`

**Maple [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.93

method	result
parts	$-\frac{a}{4x^4} + bc^4 \left( -\frac{\operatorname{arcsech}(cx)}{4c^4x^4} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \left( 3 \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right) c^4x^4 + 3\sqrt{-c^2x^2+1} c^2x^2 + 2\sqrt{-c^2x^2+1} \right)}{32c^3x^3\sqrt{-c^2x^2+1}} \right)$
derivativedivides	$c^4 \left( -\frac{a}{4c^4x^4} + b \left( -\frac{\operatorname{arcsech}(cx)}{4c^4x^4} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \left( 3 \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right) c^4x^4 + 3\sqrt{-c^2x^2+1} c^2x^2 + 2\sqrt{-c^2x^2+1} \right)}{32c^3x^3\sqrt{-c^2x^2+1}} \right) \right)$
default	$c^4 \left( -\frac{a}{4c^4x^4} + b \left( -\frac{\operatorname{arcsech}(cx)}{4c^4x^4} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \left( 3 \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right) c^4x^4 + 3\sqrt{-c^2x^2+1} c^2x^2 + 2\sqrt{-c^2x^2+1} \right)}{32c^3x^3\sqrt{-c^2x^2+1}} \right) \right)$

input `int((a+b*arcsech(c*x))/x^5,x,method=_RETURNVERBOSE)`

output 
$$-1/4*a/x^4+b*c^4*(-1/4/c^4/x^4*arcsech(c*x)+1/32*(-(c*x-1)/c/x)^(1/2)/c^3/x^3*((c*x+1)/c/x)^(1/2)*(3*arctanh(1/(-c^2*x^2+1)^(1/2))*c^4*x^4+3*(-c^2*x^2+1)^(1/2)*c^2*x^2+2*(-c^2*x^2+1)^(1/2))/(-c^2*x^2+1)^(1/2))$$

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.64

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5} dx$$

$$= \frac{(3bc^4x^4 - 8b) \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}+1}}{cx}\right) + (3bc^3x^3 + 2bcx)\sqrt{-\frac{c^2x^2-1}{c^2x^2}} - 8a}{32x^4}$$

input `integrate((a+b*arcsech(c*x))/x^5,x, algorithm="fricas")`

output 
$$1/32*((3*b*c^4*x^4 - 8*b)*\log((c*x*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} + 1)/(c*x)) + (3*b*c^3*x^3 + 2*b*c*x)*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} - 8*a)/x^4$$

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5} dx = \int \frac{a + b \operatorname{asech}(cx)}{x^5} dx$$

input `integrate((a+b*asech(c*x))/x**5,x)`

output `Integral((a + b*asech(c*x))/x**5, x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.04

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5} dx$$

$$= \frac{1}{64} b \left( \frac{3c^5 \log\left(cx \sqrt{\frac{1}{c^2 x^2} - 1} + 1\right) - 3c^5 \log\left(cx \sqrt{\frac{1}{c^2 x^2} - 1} - 1\right) - \frac{2\left(3c^8 x^3 \left(\frac{1}{c^2 x^2} - 1\right)^{\frac{3}{2}} - 5c^6 x \sqrt{\frac{1}{c^2 x^2} - 1}\right)}{c^4 x^4 \left(\frac{1}{c^2 x^2} - 1\right)^2 - 2c^2 x^2 \left(\frac{1}{c^2 x^2} - 1\right) + 1}}{c} - \frac{a}{4x^4} \right) - 16$$

input `integrate((a+b*arcsech(c*x))/x^5,x, algorithm="maxima")`

output `1/64*b*((3*c^5*log(c*x*sqrt(1/(c^2*x^2) - 1) + 1) - 3*c^5*log(c*x*sqrt(1/(c^2*x^2) - 1) - 1) - 2*(3*c^8*x^3*(1/(c^2*x^2) - 1)^(3/2) - 5*c^6*x*sqrt(1/(c^2*x^2) - 1)))/(c^4*x^4*(1/(c^2*x^2) - 1)^2 - 2*c^2*x^2*(1/(c^2*x^2) - 1) + 1))/c - 16*arcsech(c*x)/x^4) - 1/4*a/x^4`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5} dx = \int \frac{b \operatorname{arsech}(cx) + a}{x^5} dx$$

input `integrate((a+b*arcsech(c*x))/x^5,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/x^5, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x^5} dx$$

input `int((a + b*acosh(1/(c*x)))/x^5,x)`

output `int((a + b*acosh(1/(c*x)))/x^5, x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5} dx = \frac{4 \left( \int \frac{\operatorname{asech}(cx)}{x^5} dx \right) b x^4 - a}{4x^4}$$

input `int((a+b*asech(c*x))/x^5,x)`

output `(4*int(asech(c*x)/x**5,x)*b*x**4 - a)/(4*x**4)`

### 3.31 $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^6} dx$

Optimal result	301
Mathematica [A] (verified)	301
Rubi [A] (verified)	302
Maple [A] (verified)	304
Fricas [A] (verification not implemented)	304
Sympy [F]	305
Maxima [A] (verification not implemented)	305
Giac [F]	306
Mupad [F(-1)]	306
Reduce [F]	306

#### Optimal result

Integrand size = 12, antiderivative size = 115

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^6} dx = \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{25x^5} + \frac{4bc^2\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{75x^3} + \frac{8bc^4\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{75x} - \frac{a + b\operatorname{sech}^{-1}(cx)}{5x^5}$$

output

```
1/25*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/x^5+4/75*b*c^2*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/x^3+8/75*b*c^4*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/x-1/5*(a+b*arcsech(c*x))/x^5
```

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.82

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^6} dx = -\frac{a}{5x^5} + b\left(\frac{8c^5}{75} + \frac{1}{25x^5} + \frac{c}{25x^4} + \frac{4c^2}{75x^3} + \frac{4c^3}{75x^2} + \frac{8c^4}{75x}\right)\sqrt{\frac{1-cx}{1+cx}} - \frac{b\operatorname{sech}^{-1}(cx)}{5x^5}$$

input

```
Integrate[(a + b*ArcSech[c*x])/x^6, x]
```

output

```
-1/5*a/x^5 + b*((8*c^5)/75 + 1/(25*x^5) + c/(25*x^4) + (4*c^2)/(75*x^3) +
(4*c^3)/(75*x^2) + (8*c^4)/(75*x))*Sqrt[(1 - c*x)/(1 + c*x)] - (b*ArcSech[
c*x])/(5*x^5)
```

**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.14, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6837, 114, 27, 114, 27, 106}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^6} dx \\
 & \quad \downarrow 6837 \\
 & -\frac{1}{5}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{x^6\sqrt{1-cx}\sqrt{cx+1}} dx - \frac{a + b \operatorname{sech}^{-1}(cx)}{5x^5} \\
 & \quad \downarrow 114 \\
 & -\frac{1}{5}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{1}{5} \int -\frac{4c^2}{x^4\sqrt{1-cx}\sqrt{cx+1}} dx - \frac{\sqrt{1-cx}\sqrt{cx+1}}{5x^5} \right) - \\
 & \quad \quad \quad \frac{a + b \operatorname{sech}^{-1}(cx)}{5x^5} \\
 & \quad \downarrow 27 \\
 & -\frac{1}{5}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{4}{5}c^2 \int \frac{1}{x^4\sqrt{1-cx}\sqrt{cx+1}} dx - \frac{\sqrt{1-cx}\sqrt{cx+1}}{5x^5} \right) - \frac{a + b \operatorname{sech}^{-1}(cx)}{5x^5} \\
 & \quad \downarrow 114 \\
 & -\frac{1}{5}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{4}{5}c^2 \left( -\frac{1}{3} \int -\frac{2c^2}{x^2\sqrt{1-cx}\sqrt{cx+1}} dx - \frac{\sqrt{1-cx}\sqrt{cx+1}}{3x^3} \right) - \frac{\sqrt{1-cx}\sqrt{cx+1}}{5x^5} \right) - \\
 & \quad \quad \quad \frac{a + b \operatorname{sech}^{-1}(cx)}{5x^5} \\
 & \quad \downarrow 27
 \end{aligned}$$

$$-\frac{1}{5}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{4}{5}c^2\left(\frac{2}{3}c^2\int\frac{1}{x^2\sqrt{1-cx}\sqrt{cx+1}}dx-\frac{\sqrt{1-cx}\sqrt{cx+1}}{3x^3}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{5x^5}\right)-$$

$$\frac{a+b\operatorname{sech}^{-1}(cx)}{5x^5}$$

↓ 106

$$\frac{1}{5}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{4}{5}c^2\left(-\frac{2c^2\sqrt{1-cx}\sqrt{cx+1}}{3x}-\frac{\sqrt{1-cx}\sqrt{cx+1}}{3x^3}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{5x^5}\right)$$

input `Int[(a + b*ArcSech[c*x])/x^6,x]`

output `-1/5*(b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/5*(Sqrt[1 - c*x]*Sqrt[1 + c*x])/x^5 + (4*c^2*(-1/3*(Sqrt[1 - c*x]*Sqrt[1 + c*x])/x^3 - (2*c^2*Sqrt[1 - c*x]*Sqrt[1 + c*x])/(3*x))))/5) - (a + b*ArcSech[c*x])/(5*x^5)`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 106 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && EqQ[a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1), 0] && NeQ[m, -1]`

rule 114 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])`



rule 6837

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))*((d_.)*(x_.))^(m_.), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*ArcSech[c*x])/(d*(m + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(m + 1))*Sqrt[1/(1 + c*x)] Int[(d*x)^m/(Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

**Maple [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.70

method	result	size
parts	$-\frac{a}{5x^5} + bc^5 \left( -\frac{\operatorname{arcsech}(cx)}{5c^5x^5} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} (8c^4x^4 + 4c^2x^2 + 3)}{75c^4x^4} \right)$	81
derivativedivides	$c^5 \left( -\frac{a}{5c^5x^5} + b \left( -\frac{\operatorname{arcsech}(cx)}{5c^5x^5} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} (8c^4x^4 + 4c^2x^2 + 3)}{75c^4x^4} \right) \right)$	85
default	$c^5 \left( -\frac{a}{5c^5x^5} + b \left( -\frac{\operatorname{arcsech}(cx)}{5c^5x^5} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} (8c^4x^4 + 4c^2x^2 + 3)}{75c^4x^4} \right) \right)$	85

input

```
int((a+b*arcsech(c*x))/x^6,x,method=_RETURNVERBOSE)
```

output

```
-1/5*a/x^5+b*c^5*(-1/5/c^5/x^5*arcsech(c*x)+1/75*(-(c*x-1)/c/x)^(1/2)/c^4/x^4*((c*x+1)/c/x)^(1/2)*(8*c^4*x^4+4*c^2*x^2+3))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.77

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^6} dx$$

$$= \frac{15 b \log \left( \frac{cx \sqrt{-\frac{c^2x^2-1}{c^2x^2}} + 1}{cx} \right) - (8bc^5x^5 + 4bc^3x^3 + 3bcx) \sqrt{-\frac{c^2x^2-1}{c^2x^2}} + 15a}{75x^5}$$

input

```
integrate((a+b*arcsech(c*x))/x^6,x, algorithm="fricas")
```

output

```
-1/75*(15*b*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (8*b*c^5
*x^5 + 4*b*c^3*x^3 + 3*b*c*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 15*a)/x^5
```

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^6} dx = \int \frac{a + b \operatorname{asech}(cx)}{x^6} dx$$

input

```
integrate((a+b*asech(c*x))/x**6,x)
```

output

```
Integral((a + b*asech(c*x))/x**6, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.63

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^6} dx$$

$$= \frac{1}{75} b \left( \frac{3 c^6 \left(\frac{1}{c^2 x^2} - 1\right)^{\frac{5}{2}} + 10 c^6 \left(\frac{1}{c^2 x^2} - 1\right)^{\frac{3}{2}} + 15 c^6 \sqrt{\frac{1}{c^2 x^2} - 1}}{c} - \frac{15 \operatorname{arsech}(cx)}{x^5} \right) - \frac{a}{5 x^5}$$

input

```
integrate((a+b*arcsech(c*x))/x^6,x, algorithm="maxima")
```

output

```
1/75*b*((3*c^6*(1/(c^2*x^2) - 1)^(5/2) + 10*c^6*(1/(c^2*x^2) - 1)^(3/2) +
15*c^6*sqrt(1/(c^2*x^2) - 1))/c - 15*arcsech(c*x)/x^5) - 1/5*a/x^5
```

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^6} dx = \int \frac{b \operatorname{arsech}(cx) + a}{x^6} dx$$

input `integrate((a+b*arcsech(c*x))/x^6,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/x^6, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^6} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x^6} dx$$

input `int((a + b*acosh(1/(c*x)))/x^6,x)`

output `int((a + b*acosh(1/(c*x)))/x^6, x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^6} dx = \frac{5 \left( \int \frac{\operatorname{asech}(cx)}{x^6} dx \right) b x^5 - a}{5x^5}$$

input `int((a+b*asech(c*x))/x^6,x)`

output `(5*int(asech(c*x)/x**6,x)*b*x**5 - a)/(5*x**5)`

### 3.32 $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^7} dx$

Optimal result	307
Mathematica [A] (verified)	308
Rubi [A] (verified)	308
Maple [A] (verified)	311
Fricas [A] (verification not implemented)	312
Sympy [F]	312
Maxima [A] (verification not implemented)	313
Giac [F]	313
Mupad [F(-1)]	314
Reduce [F]	314

#### Optimal result

Integrand size = 12, antiderivative size = 175

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^7} dx = \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{36x^6} + \frac{5bc^2\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{144x^4} + \frac{5bc^4\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{96x^2} - \frac{a + b\operatorname{sech}^{-1}(cx)}{6x^6} + \frac{5bc^6\sqrt{\frac{1-cx}{1+cx}}(1+cx)\operatorname{arctanh}(\sqrt{1-c^2x^2})}{96\sqrt{1-c^2x^2}}$$

output `1/36*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/x^6+5/144*b*c^2*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/x^4+5/96*b*c^4*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/x^2-1/6*(a+b*arcsech(c*x))/x^6+5/96*b*c^6*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*arctanh((-c^2*x^2+1)^(1/2))/(-c^2*x^2+1)^(1/2)`

**Mathematica [A] (verified)**

Time = 0.10 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.90

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^7} dx = -\frac{a}{6x^6} + b \left( \frac{1}{36x^6} + \frac{c}{36x^5} + \frac{5c^2}{144x^4} + \frac{5c^3}{144x^3} + \frac{5c^4}{96x^2} + \frac{5c^5}{96x} \right) \sqrt{\frac{1-cx}{1+cx}} - \frac{b \operatorname{sech}^{-1}(cx)}{6x^6} - \frac{5}{96} bc^6 \log(x) + \frac{5}{96} bc^6 \log \left( 1 + \sqrt{\frac{1-cx}{1+cx}} + cx \sqrt{\frac{1-cx}{1+cx}} \right)$$

input `Integrate[(a + b*ArcSech[c*x])/x^7,x]`

output `-1/6*a/x^6 + b*(1/(36*x^6) + c/(36*x^5) + (5*c^2)/(144*x^4) + (5*c^3)/(144*x^3) + (5*c^4)/(96*x^2) + (5*c^5)/(96*x))*Sqrt[(1 - c*x)/(1 + c*x)] - (b*ArcSech[c*x])/(6*x^6) - (5*b*c^6*Log[x])/96 + (5*b*c^6*Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]])/96`

**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.94, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$ , Rules used = {6837, 114, 27, 114, 27, 114, 25, 27, 103, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^7} dx \quad \downarrow \quad 6837$$

$$-\frac{1}{6} b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{1}{x^7 \sqrt{1-cx} \sqrt{cx+1}} dx - \frac{a + b \operatorname{sech}^{-1}(cx)}{6x^6}$$

$$\quad \downarrow \quad 114$$

$$\begin{aligned}
& -\frac{1}{6}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{1}{6}\int\frac{5c^2}{\frac{x^5\sqrt{1-cx}\sqrt{cx+1}}{a+b\operatorname{sech}^{-1}(cx)}}dx-\frac{\sqrt{1-cx}\sqrt{cx+1}}{6x^6}\right)- \\
& \quad \downarrow 27 \\
& -\frac{1}{6}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{5}{6}c^2\int\frac{1}{x^5\sqrt{1-cx}\sqrt{cx+1}}dx-\frac{\sqrt{1-cx}\sqrt{cx+1}}{6x^6}\right)-\frac{a+b\operatorname{sech}^{-1}(cx)}{6x^6} \\
& \quad \downarrow 114 \\
& -\frac{1}{6}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{5}{6}c^2\left(-\frac{1}{4}\int\frac{3c^2}{\frac{x^3\sqrt{1-cx}\sqrt{cx+1}}{a+b\operatorname{sech}^{-1}(cx)}}dx-\frac{\sqrt{1-cx}\sqrt{cx+1}}{4x^4}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{6x^6}\right)- \\
& \quad \downarrow 27 \\
& -\frac{1}{6}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{5}{6}c^2\left(\frac{3}{4}c^2\int\frac{1}{x^3\sqrt{1-cx}\sqrt{cx+1}}dx-\frac{\sqrt{1-cx}\sqrt{cx+1}}{4x^4}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{6x^6}\right)- \\
& \quad \downarrow 114 \\
& -\frac{1}{6}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{5}{6}c^2\left(\frac{3}{4}c^2\left(-\frac{1}{2}\int\frac{c^2}{\frac{x\sqrt{1-cx}\sqrt{cx+1}}{a+b\operatorname{sech}^{-1}(cx)}}dx-\frac{\sqrt{1-cx}\sqrt{cx+1}}{2x^2}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{4x^4}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{6x^6}\right)- \\
& \quad \downarrow 25 \\
& -\frac{1}{6}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{5}{6}c^2\left(\frac{3}{4}c^2\left(\frac{1}{2}\int\frac{c^2}{\frac{x\sqrt{1-cx}\sqrt{cx+1}}{a+b\operatorname{sech}^{-1}(cx)}}dx-\frac{\sqrt{1-cx}\sqrt{cx+1}}{2x^2}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{4x^4}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{6x^6}\right)- \\
& \quad \downarrow 27 \\
& -\frac{1}{6}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{5}{6}c^2\left(\frac{3}{4}c^2\left(\frac{1}{2}c^2\int\frac{1}{x\sqrt{1-cx}\sqrt{cx+1}}dx-\frac{\sqrt{1-cx}\sqrt{cx+1}}{2x^2}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{4x^4}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{6x^6}\right)- \\
& \quad \downarrow 27
\end{aligned}$$

↓ 103

$$-\frac{1}{6}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{5}{6}c^2\left(\frac{3}{4}c^2\left(-\frac{1}{2}c^3\int\frac{1}{c-c(1-cx)(cx+1)}d(\sqrt{1-cx}\sqrt{cx+1})-\frac{\sqrt{1-cx}\sqrt{cx+1}}{2x^2}\right)-\frac{a+b\operatorname{sech}^{-1}(cx)}{6x^6}\right)\right)$$

↓ 221

$$\frac{1}{6}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{5}{6}c^2\left(\frac{3}{4}c^2\left(-\frac{1}{2}c^2\operatorname{arctanh}(\sqrt{1-cx}\sqrt{cx+1})-\frac{\sqrt{1-cx}\sqrt{cx+1}}{2x^2}\right)-\frac{\sqrt{1-cx}\sqrt{cx+1}}{4x^4}\right)-\frac{a+b\operatorname{sech}^{-1}(cx)}{6x^6}\right)$$

input `Int[(a + b*ArcSech[c*x])/x^7,x]`

output `-1/6*(a + b*ArcSech[c*x])/x^6 - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/6*(Sqrt[1 - c*x]*Sqrt[1 + c*x])/x^6 + (5*c^2*(-1/4*(Sqrt[1 - c*x]*Sqrt[1 + c*x])/x^4 + (3*c^2*(-1/2*(Sqrt[1 - c*x]*Sqrt[1 + c*x])/x^2 - (c^2*ArcTanh[Sqrt[1 - c*x]*Sqrt[1 + c*x]]/2))/4))/6)/6`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 103 `Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]`

```
rule 114 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 6837 Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcSech[c*x])/(d*(m + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(m + 1))*Sqrt[1/(1 + c*x)] Int[(d*x)^m/(Sqrt[1 - c*x]*Sqrt[1 + c*x]), x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

### Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.86

method	result
parts	$-\frac{a}{6x^6} + bc^6 \left( -\frac{\operatorname{arcsech}(cx)}{6c^6x^6} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \left( 15 \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right) c^6x^6 + 15\sqrt{-c^2x^2+1} c^4x^4 + 10\sqrt{-c^2x^2+1} c^2x^2 + 5 \right)}{288c^5x^5\sqrt{-c^2x^2+1}} \right)$
derivativedivides	$c^6 \left( -\frac{a}{6c^6x^6} + b \left( -\frac{\operatorname{arcsech}(cx)}{6c^6x^6} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \left( 15 \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right) c^6x^6 + 15\sqrt{-c^2x^2+1} c^4x^4 + 10\sqrt{-c^2x^2+1} c^2x^2 + 5 \right)}{288c^5x^5\sqrt{-c^2x^2+1}} \right) \right)$
default	$c^6 \left( -\frac{a}{6c^6x^6} + b \left( -\frac{\operatorname{arcsech}(cx)}{6c^6x^6} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \left( 15 \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right) c^6x^6 + 15\sqrt{-c^2x^2+1} c^4x^4 + 10\sqrt{-c^2x^2+1} c^2x^2 + 5 \right)}{288c^5x^5\sqrt{-c^2x^2+1}} \right) \right)$

```
input int((a+b*arcsech(c*x))/x^7,x,method=_RETURNVERBOSE)
```



output

```
-1/6*a/x^6+b*c^6*(-1/6/c^6/x^6*arcsech(c*x)+1/288*(-(c*x-1)/c/x)^(1/2)/c^5
/x^5*((c*x+1)/c/x)^(1/2)*(15*arctanh(1/(-c^2*x^2+1)^(1/2))*c^6*x^6+15*(-c^
2*x^2+1)^(1/2)*c^4*x^4+10*(-c^2*x^2+1)^(1/2)*c^2*x^2+8*(-c^2*x^2+1)^(1/2))
/(-c^2*x^2+1)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.57

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^7} dx$$

$$= \frac{3(5bc^6x^6 - 16b) \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}+1}{cx}\right) + (15bc^5x^5 + 10bc^3x^3 + 8bcx)\sqrt{-\frac{c^2x^2-1}{c^2x^2}} - 48a}{288x^6}$$

input

```
integrate((a+b*arcsech(c*x))/x^7,x, algorithm="fricas")
```

output

```
1/288*(3*(5*b*c^6*x^6 - 16*b)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)
/(c*x)) + (15*b*c^5*x^5 + 10*b*c^3*x^3 + 8*b*c*x)*sqrt(-(c^2*x^2 - 1)/(c^2
*x^2)) - 48*a)/x^6
```

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^7} dx = \int \frac{a + b \operatorname{asech}(cx)}{x^7} dx$$

input

```
integrate((a+b*asech(c*x))/x**7,x)
```

output

```
Integral((a + b*asech(c*x))/x**7, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.06

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^7} dx$$

$$= \frac{1}{576} b \left( \frac{15 c^7 \log \left( cx \sqrt{\frac{1}{c^2 x^2} - 1} + 1 \right) - 15 c^7 \log \left( cx \sqrt{\frac{1}{c^2 x^2} - 1} - 1 \right) - \frac{2 \left( 15 c^{12} x^5 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{5}{2}} - 40 c^{10} x^3 \left( \frac{1}{c^2 x^2} - 1 \right) \right)}{c^6 x^6 \left( \frac{1}{c^2 x^2} - 1 \right)^3 - 3 c^4 x^4 \left( \frac{1}{c^2 x^2} - 1 \right)^2 + 3 c^2 x^2 \left( \frac{1}{c^2 x^2} - 1 \right) - 1}}{c} \right) - \frac{a}{6 x^6}$$

input `integrate((a+b*arcsech(c*x))/x^7,x, algorithm="maxima")`output `1/576*b*((15*c^7*log(c*x*sqrt(1/(c^2*x^2) - 1) + 1) - 15*c^7*log(c*x*sqrt(1/(c^2*x^2) - 1) - 1) - 2*(15*c^12*x^5*(1/(c^2*x^2) - 1)^(5/2) - 40*c^10*x^3*(1/(c^2*x^2) - 1)^(3/2) + 33*c^8*x*sqrt(1/(c^2*x^2) - 1)))/(c^6*x^6*(1/(c^2*x^2) - 1)^3 - 3*c^4*x^4*(1/(c^2*x^2) - 1)^2 + 3*c^2*x^2*(1/(c^2*x^2) - 1) - 1))/c - 96*arcsech(c*x)/x^6) - 1/6*a/x^6`**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^7} dx = \int \frac{b \operatorname{ar} \operatorname{sech}(cx) + a}{x^7} dx$$

input `integrate((a+b*arcsech(c*x))/x^7,x, algorithm="giac")`output `integrate((b*arcsech(c*x) + a)/x^7, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^7} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x^7} dx$$

input `int((a + b*acosh(1/(c*x)))/x^7,x)`output `int((a + b*acosh(1/(c*x)))/x^7, x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^7} dx = \frac{6 \left( \int \frac{a \operatorname{sech}(cx)}{x^7} dx \right) b x^6 - a}{6x^6}$$

input `int((a+b*asech(c*x))/x^7,x)`output `(6*int(asech(c*x)/x**7,x)*b*x**6 - a)/(6*x**6)`

### 3.33 $\int x^3 (a + b \operatorname{sech}^{-1}(cx))^2 dx$

Optimal result . . . . .	315
Mathematica [A] (verified) . . . . .	315
Rubi [A] (verified) . . . . .	316
Maple [B] (verified) . . . . .	319
Fricas [B] (verification not implemented) . . . . .	319
Sympy [F] . . . . .	320
Maxima [F] . . . . .	320
Giac [F] . . . . .	321
Mupad [F(-1)] . . . . .	321
Reduce [F] . . . . .	321

#### Optimal result

Integrand size = 14, antiderivative size = 124

$$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^2 dx = -\frac{b^2 x^2}{12c^2} - \frac{b \sqrt{\frac{1-cx}{1+cx}} (1+cx) (a + b \operatorname{sech}^{-1}(cx))}{3c^4}$$

$$- \frac{bx^2 \sqrt{\frac{1-cx}{1+cx}} (1+cx) (a + b \operatorname{sech}^{-1}(cx))}{6c^2}$$

$$+ \frac{1}{4} x^4 (a + b \operatorname{sech}^{-1}(cx))^2 - \frac{b^2 \log(x)}{3c^4}$$

output

```
-1/12*b^2*x^2/c^2-1/3*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+b*arcsech(c*x)
)/c^4-1/6*b*x^2*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+b*arcsech(c*x))/c^2+1/
4*x^4*(a+b*arcsech(c*x))^2-1/3*b^2*ln(x)/c^4
```

#### Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.71

$$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^2 dx =$$

$$\frac{b^2 c^2 x^2 - 3a^2 c^4 x^4 + 4ab \sqrt{\frac{1-cx}{1+cx}} + 4abcx \sqrt{\frac{1-cx}{1+cx}} + 2abc^2 x^2 \sqrt{\frac{1-cx}{1+cx}} + 2abc^3 x^3 \sqrt{\frac{1-cx}{1+cx}} + 2b(-3ac^4 x^4 + b \sqrt{\frac{1-cx}{1+cx}})}{12c^4}$$

input `Integrate[x^3*(a + b*ArcSech[c*x])^2,x]`

output 
$$\frac{-1/12*(b^2*c^2*x^2 - 3*a^2*c^4*x^4 + 4*a*b*\sqrt{(1 - c*x)/(1 + c*x)} + 4*a*b*c*x*\sqrt{(1 - c*x)/(1 + c*x)} + 2*a*b*c^2*x^2*\sqrt{(1 - c*x)/(1 + c*x)} + 2*a*b*c^3*x^3*\sqrt{(1 - c*x)/(1 + c*x)} + 2*b*(-3*a*c^4*x^4 + b*\sqrt{(1 - c*x)/(1 + c*x)})*(2 + 2*c*x + c^2*x^2 + c^3*x^3))*\text{ArcSech}[c*x] - 3*b^2*c^4*x^4*\text{ArcSech}[c*x]^2 + 4*b^2*\text{Log}[x])/c^4$$

### Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.06, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$ , Rules used = {6839, 5974, 3042, 4673, 3042, 4672, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^3 (a + b \operatorname{sech}^{-1}(cx))^2 dx \\ & \quad \downarrow \text{6839} \\ & \frac{\int c^4 x^4 \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx))^2 d \operatorname{sech}^{-1}(cx)}{c^4} \\ & \quad \downarrow \text{5974} \\ & \frac{\frac{1}{2} b \int c^4 x^4 (a + b \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) - \frac{1}{4} c^4 x^4 (a + b \operatorname{sech}^{-1}(cx))^2}{c^4} \\ & \quad \downarrow \text{3042} \\ & \frac{-\frac{1}{4} c^4 x^4 (a + b \operatorname{sech}^{-1}(cx))^2 + \frac{1}{2} b \int (a + b \operatorname{sech}^{-1}(cx)) \csc \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^4 d \operatorname{sech}^{-1}(cx)}{c^4} \\ & \quad \downarrow \text{4673} \\ & \frac{\frac{1}{2} b \left( \frac{2}{3} \int c^2 x^2 (a + b \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) + \frac{1}{3} c^2 x^2 \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{6} b c^2 x^2 \right) - \frac{1}{4} c^4 x^4 (a + b \operatorname{sech}^{-1}(cx))^2}{c^4} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\frac{-\frac{1}{4}c^4x^4(a + b\operatorname{sech}^{-1}(cx))^2 + \frac{1}{2}b\left(\frac{2}{3}\int(a + b\operatorname{sech}^{-1}(cx))\csc\left(\operatorname{isech}^{-1}(cx) + \frac{\pi}{2}\right)^2 d\operatorname{sech}^{-1}(cx) + \frac{1}{3}c^2x^2\sqrt{\frac{1-cx}{cx+1}}(cx)\right)}{c^4}$$

↓ 4672

$$\frac{-\frac{1}{4}c^4x^4(a + b\operatorname{sech}^{-1}(cx))^2 + \frac{1}{2}b\left(\frac{2}{3}\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx)) - ib\int -i\sqrt{\frac{1-cx}{cx+1}}(cx+1)d\operatorname{sech}^{-1}(cx)\right)\right)}{c^4}$$

↓ 26

$$\frac{\frac{1}{2}b\left(\frac{2}{3}\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx)) - b\int\sqrt{\frac{1-cx}{cx+1}}(cx+1)d\operatorname{sech}^{-1}(cx)\right) + \frac{1}{3}c^2x^2\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))\right)}{c^4}$$

↓ 3042

$$\frac{-\frac{1}{4}c^4x^4(a + b\operatorname{sech}^{-1}(cx))^2 + \frac{1}{2}b\left(\frac{2}{3}\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx)) - b\int -i\tan(\operatorname{isech}^{-1}(cx))d\operatorname{sech}^{-1}(cx)\right)\right)}{c^4}$$

↓ 26

$$\frac{-\frac{1}{4}c^4x^4(a + b\operatorname{sech}^{-1}(cx))^2 + \frac{1}{2}b\left(\frac{2}{3}\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx)) + ib\int\tan(\operatorname{isech}^{-1}(cx))d\operatorname{sech}^{-1}(cx)\right)\right)}{c^4}$$

↓ 3956

$$\frac{\frac{1}{2}b\left(\frac{1}{3}c^2x^2\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx)) + \frac{2}{3}\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx)) - b\log\left(\frac{1}{cx}\right)\right) + \frac{1}{6}bc^2x^2\right)}{c^4}$$

input `Int[x^3*(a + b*ArcSech[c*x])^2,x]`

output `-((-1/4*(c^4*x^4*(a + b*ArcSech[c*x])^2) + (b*((b*c^2*x^2)/6 + (c^2*x^2*sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x])))/3 + (2*(sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x]) - b*Log[1/(c*x)]))/3))/2)/c^4)`

## Definitions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3956  $\text{Int}[\tan[(c.) + (d.)*(x)], x\_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$
- rule 4672  $\text{Int}[\text{csc}[(e.) + (f.)*(x)]^2*((c.) + (d.)*(x))^{(m.)}, x\_Symbol] \rightarrow \text{Simp}[(-c + d*x)^m*(\text{Cot}[e + f*x]/f), x] + \text{Simp}[d*(m/f) \text{Int}[(c + d*x)^{(m-1)}*\text{Cot}[e + f*x], x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \ \&\& \ \text{GtQ}[m, 0]$
- rule 4673  $\text{Int}[(\text{csc}[(e.) + (f.)*(x)]*(b.))^{(n)}*((c.) + (d.)*(x)), x\_Symbol] \rightarrow \text{Simp}[(-b^2)*(c + d*x)*\text{Cot}[e + f*x]*((b*\text{Csc}[e + f*x])^{(n-2)}/(f*(n-1))), x] + (-\text{Simp}[b^2*d*((b*\text{Csc}[e + f*x])^{(n-2)}/(f^2*(n-1)*(n-2))), x] + \text{Simp}[b^2*((n-2)/(n-1)) \text{Int}[(c + d*x)*(b*\text{Csc}[e + f*x])^{(n-2)}, x], x]) /; \text{FreeQ}[\{b, c, d, e, f\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{NeQ}[n, 2]$
- rule 5974  $\text{Int}[(c.) + (d.)*(x))^{(m)}*\text{Sech}[(a.) + (b.)*(x)]^{(n)}*\text{Tanh}[(a.) + (b.)*(x)]^{(p)}, x\_Symbol] \rightarrow \text{Simp}[(-c + d*x)^m*(\text{Sech}[a + b*x]^n/(b*n)), x] + \text{Simp}[d*(m/(b*n)) \text{Int}[(c + d*x)^{(m-1)}*\text{Sech}[a + b*x]^n, x], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \ \&\& \ \text{EqQ}[p, 1] \ \&\& \ \text{GtQ}[m, 0]$
- rule 6839  $\text{Int}[(a.) + \text{ArcSech}[(c.)*(x)]*(b.))^{(n)}*(x)^{(m)}, x\_Symbol] \rightarrow \text{Simp}[-(c^{(m+1)})^{(-1)} \text{Subst}[\text{Int}[(a + b*x)^n*\text{Sech}[x]^{(m+1)}*\text{Tanh}[x], x], x, \text{ArcSech}[c*x]], x] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ (\text{GtQ}[n, 0] \ || \ \text{LtQ}[m, -1])$

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 223 vs. 2(110) = 220.

Time = 0.76 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.81

method	result
parts	$\frac{a^2 x^4}{4} + \frac{b^2 \left( -\frac{\operatorname{arcsech}(cx)}{3} + \frac{\operatorname{arcsech}(cx)^2 c^4 x^4}{4} - \frac{\operatorname{arcsech}(cx) \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} c^3 x^3}{6} - \frac{\operatorname{arcsech}(cx) \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} cx}{3} - \frac{c^2 x^2}{12} \right)}{c^4}$
derivativeldivides	$\frac{a^2 c^4 x^4}{4} + b^2 \left( -\frac{\operatorname{arcsech}(cx)}{3} + \frac{\operatorname{arcsech}(cx)^2 c^4 x^4}{4} - \frac{\operatorname{arcsech}(cx) \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} c^3 x^3}{6} - \frac{\operatorname{arcsech}(cx) \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} cx}{3} - \frac{c^2 x^2}{12} \right)$
default	$\frac{a^2 c^4 x^4}{4} + b^2 \left( -\frac{\operatorname{arcsech}(cx)}{3} + \frac{\operatorname{arcsech}(cx)^2 c^4 x^4}{4} - \frac{\operatorname{arcsech}(cx) \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} c^3 x^3}{6} - \frac{\operatorname{arcsech}(cx) \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} cx}{3} - \frac{c^2 x^2}{12} \right)$

input `int(x^3*(a+b*arcsech(c*x))^2,x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{4} a^2 x^4 + \frac{b^2}{c^4} \left( -\frac{1}{3} \operatorname{arcsech}(cx) + \frac{1}{4} \operatorname{arcsech}(cx)^2 c^4 x^4 - \frac{1}{6} \operatorname{arcsech}(cx) \left( -\frac{cx-1}{cx} \right)^{1/2} \left( \frac{cx+1}{cx} \right)^{1/2} c^3 x^3 - \frac{1}{3} \operatorname{arcsech}(cx) \left( \frac{cx+1}{cx} \right)^{1/2} \left( -\frac{cx-1}{cx} \right)^{1/2} c^2 x^2 + \frac{1}{3} \ln \left( 1 + \frac{1}{c/x} + \left( -1 + \frac{1}{c/x} \right)^{1/2} \left( 1 + \frac{1}{c/x} \right)^{1/2} \right)^2 \right) + 2 a b / c^4 \left( \frac{1}{4} c^4 x^4 \operatorname{arcsech}(cx) - \frac{1}{12} \left( -\frac{cx-1}{cx} \right)^{1/2} c^2 x^2 \left( \frac{cx+1}{cx} \right)^{1/2} \left( c^2 x^2 + 2 \right) \right)$$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 244 vs. 2(110) = 220.

Time = 0.11 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.97

$$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^2 dx$$

$$= \frac{3 b^2 c^4 x^4 \log \left( \frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx} \right)^2 + 3 a^2 c^4 x^4 - 6 a b c^4 \log \left( \frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - 1}{x} \right) - b^2 c^2 x^2 - 4 b^2 \log(x) + 2 \left( 3 a b c^4 \right)}{12 c^4}$$

input `integrate(x^3*(a+b*arcsech(c*x))^2,x, algorithm="fricas")`



output

```
1/12*(3*b^2*c^4*x^4*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x))^2
+ 3*a^2*c^4*x^4 - 6*a*b*c^4*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x
) - b^2*c^2*x^2 - 4*b^2*log(x) + 2*(3*a*b*c^4*x^4 - 3*a*b*c^4 - (b^2*c^3*x
^3 + 2*b^2*c*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*log((c*x*sqrt(-(c^2*x^2 -
1)/(c^2*x^2)) + 1)/(c*x)) - 2*(a*b*c^3*x^3 + 2*a*b*c*x)*sqrt(-(c^2*x^2 - 1
)/(c^2*x^2)))/c^4
```

**Sympy [F]**

$$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int x^3 (a + b \operatorname{arsech}(cx))^2 dx$$

input

```
integrate(x**3*(a+b*asech(c*x))**2,x)
```

output

```
Integral(x**3*(a + b*asech(c*x))**2, x)
```

**Maxima [F]**

$$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int (b \operatorname{arsech}(cx) + a)^2 x^3 dx$$

input

```
integrate(x^3*(a+b*arcsech(c*x))^2,x, algorithm="maxima")
```

output

```
1/4*a^2*x^4 + 1/6*(3*x^4*arcsech(c*x) + (c^2*x^3*(1/(c^2*x^2) - 1)^(3/2) -
3*x*sqrt(1/(c^2*x^2) - 1))/c^3)*a*b + b^2*integrate(x^3*log(sqrt(1/(c*x)
+ 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^2, x)
```

**Giac [F]**

$$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int (b \operatorname{arsech}(cx) + a)^2 x^3 dx$$

input `integrate(x^3*(a+b*arcsech(c*x))^2,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)^2*x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int x^3 \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right)^2 dx$$

input `int(x^3*(a + b*acosh(1/(c*x)))^2,x)`

output `int(x^3*(a + b*acosh(1/(c*x)))^2, x)`

**Reduce [F]**

$$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^2 dx = 2 \left( \int \operatorname{asech}(cx) x^3 dx \right) ab + \left( \int \operatorname{asech}(cx)^2 x^3 dx \right) b^2 + \frac{a^2 x^4}{4}$$

input `int(x^3*(a+b*asech(c*x))^2,x)`

output `(8*int(asech(c*x)*x**3,x)*a*b + 4*int(asech(c*x)**2*x**3,x)*b**2 + a**2*x**4)/4`

### 3.34 $\int x^2(a + b\operatorname{sech}^{-1}(cx))^2 dx$

Optimal result	322
Mathematica [A] (verified)	323
Rubi [A] (verified)	323
Maple [A] (verified)	326
Fricas [F]	327
Sympy [F]	327
Maxima [F]	327
Giac [F]	328
Mupad [F(-1)]	328
Reduce [F]	328

#### Optimal result

Integrand size = 14, antiderivative size = 140

$$\int x^2(a + b\operatorname{sech}^{-1}(cx))^2 dx = -\frac{b^2 x}{3c^2} - \frac{bx\sqrt{\frac{1-cx}{1+cx}}(1+cx)(a + b\operatorname{sech}^{-1}(cx))}{3c^2} + \frac{1}{3}x^3(a + b\operatorname{sech}^{-1}(cx))^2 - \frac{2b(a + b\operatorname{sech}^{-1}(cx))\arctan(e^{\operatorname{sech}^{-1}(cx)})}{3c^3} + \frac{ib^2 \operatorname{PolyLog}(2, -ie^{\operatorname{sech}^{-1}(cx)})}{3c^3} - \frac{ib^2 \operatorname{PolyLog}(2, ie^{\operatorname{sech}^{-1}(cx)})}{3c^3}$$

output

```
-1/3*b^2*x/c^2-1/3*b*x*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+b*arcsech(c*x))
/c^2+1/3*x^3*(a+b*arcsech(c*x))^2-2/3*b*(a+b*arcsech(c*x))*arctan(1/c/x+(-
1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/c^3+1/3*I*b^2*polylog(2,-I*(1/c/x+(-1+1/c/
x)^(1/2)*(1+1/c/x)^(1/2)))/c^3-1/3*I*b^2*polylog(2,I*(1/c/x+(-1+1/c/x)^(1/
2)*(1+1/c/x)^(1/2)))/c^3
```

**Mathematica [A] (verified)**

Time = 0.94 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.72

$$\int x^2 (a + b \operatorname{sech}^{-1}(cx))^2 dx$$

$$= \frac{1}{3} \left( a^2 x^3 + ab \left( 2x^3 \operatorname{sech}^{-1}(cx) + \frac{\sqrt{\frac{1-cx}{1+cx}} (cx - c^3 x^3 + 2\sqrt{1-c^2 x^2} \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right))}{c^3(-1+cx)} \right) \right)$$

$$+ \frac{b^2 \left( -cx - cx \sqrt{\frac{1-cx}{1+cx}} (1+cx) \operatorname{sech}^{-1}(cx) + c^3 x^3 \operatorname{sech}^{-1}(cx)^2 + i \operatorname{sech}^{-1}(cx) \log\left(1 - i e^{-\operatorname{sech}^{-1}(cx)}\right) - i \operatorname{sech}^{-1}(cx) \log\left(1 + i e^{-\operatorname{sech}^{-1}(cx)}\right) \right)}{c^3}$$

input `Integrate[x^2*(a + b*ArcSech[c*x])^2,x]`

output

```
(a^2*x^3 + a*b*(2*x^3*ArcSech[c*x] + (Sqrt[(1 - c*x)/(1 + c*x)]*(c*x - c^3*x^3 + 2*Sqrt[1 - c^2*x^2]*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])))/(c^3*(-1 + c*x)) + (b^2*(-(c*x) - c*x*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*ArcSech[c*x] + c^3*x^3*ArcSech[c*x]^2 + I*ArcSech[c*x]*Log[1 - I/E^ArcSech[c*x]] - I*ArcSech[c*x]*Log[1 + I/E^ArcSech[c*x]] + I*PolyLog[2, (-I)/E^ArcSech[c*x]] - I*PolyLog[2, I/E^ArcSech[c*x]]))/c^3)/3
```

**Rubi [A] (verified)**

Time = 0.57 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.94, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$ , Rules used = {6839, 5974, 3042, 4673, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 (a + b \operatorname{sech}^{-1}(cx))^2 dx$$

$$\downarrow \text{6839}$$

$$\frac{\int c^3 x^3 \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx))^2 d \operatorname{sech}^{-1}(cx)}{c^3}$$

$$\begin{aligned} & \downarrow 5974 \\ & \frac{\frac{2}{3}b \int c^3 x^3 (a + b \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) - \frac{1}{3}c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))^2}{c^3} \\ & \downarrow 3042 \\ & \frac{-\frac{1}{3}c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))^2 + \frac{2}{3}b \int (a + b \operatorname{sech}^{-1}(cx)) \csc \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^3 d \operatorname{sech}^{-1}(cx)}{c^3} \\ & \downarrow 4673 \\ & \frac{\frac{2}{3}b \left( \frac{1}{2} \int cx (a + b \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) + \frac{1}{2}cx \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx)) + \frac{bcx}{2} \right) - \frac{1}{3}c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))^2}{c^3} \\ & \downarrow 3042 \\ & \frac{-\frac{1}{3}c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))^2 + \frac{2}{3}b \left( \frac{1}{2} \int (a + b \operatorname{sech}^{-1}(cx)) \csc \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right) d \operatorname{sech}^{-1}(cx) + \frac{1}{2}cx \sqrt{\frac{1-cx}{cx+1}} (cx+1) \right)}{c^3} \\ & \downarrow 4668 \\ & \frac{-\frac{1}{3}c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))^2 + \frac{2}{3}b \left( \frac{1}{2} \left( -ib \int \log \left( 1 - ie^{\operatorname{sech}^{-1}(cx)} \right) d \operatorname{sech}^{-1}(cx) + ib \int \log \left( 1 + ie^{\operatorname{sech}^{-1}(cx)} \right) d \operatorname{sech}^{-1}(cx) \right) \right)}{c^3} \\ & \downarrow 2715 \\ & \frac{-\frac{1}{3}c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))^2 + \frac{2}{3}b \left( \frac{1}{2} \left( -ib \int e^{-\operatorname{sech}^{-1}(cx)} \log \left( 1 - ie^{\operatorname{sech}^{-1}(cx)} \right) de^{\operatorname{sech}^{-1}(cx)} + ib \int e^{-\operatorname{sech}^{-1}(cx)} \log \left( 1 + ie^{\operatorname{sech}^{-1}(cx)} \right) de^{\operatorname{sech}^{-1}(cx)} \right) \right)}{c^3} \\ & \downarrow 2838 \\ & \frac{-\frac{1}{3}c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))^2 + \frac{2}{3}b \left( \frac{1}{2} \left( 2 \arctan \left( e^{\operatorname{sech}^{-1}(cx)} \right) (a + b \operatorname{sech}^{-1}(cx)) - ib \operatorname{PolyLog} \left( 2, -ie^{\operatorname{sech}^{-1}(cx)} \right) + ib \operatorname{PolyLog} \left( 2, ie^{\operatorname{sech}^{-1}(cx)} \right) \right) \right)}{c^3} \end{aligned}$$

input `Int [x^2*(a + b*ArcSech[c*x])^2,x]`

output

```

-((-1/3*(c^3*x^3*(a + b*ArcSech[c*x])^2) + (2*b*((b*c*x)/2 + (c*x*Sqrt[(1
- c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x])))/2 + (2*(a + b*ArcSech[c*
x])*ArcTan[E^ArcSech[c*x]] - I*b*PolyLog[2, (-I)*E^ArcSech[c*x]] + I*b*Pol
yLog[2, I*E^ArcSech[c*x]])/2))/3)/c^3

```

### Defintions of rubi rules used

rule 2715

```

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^ (n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

```

rule 2838

```

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

```

rule 3042

```

Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

rule 4668

```

Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_
))^ (m_), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(
I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[
1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c
+ d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c
, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]

```

rule 4673

```

Int[(csc[(e_) + (f_)*(x_)]*(b_))^ (n_)*((c_) + (d_)*(x_)), x_Symbol] :>
Simp[(-b^2)*(c + d*x)*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))),
x] + (-Simp[b^2*d*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x] + S
imp[b^2*((n - 2)/(n - 1)) Int[(c + d*x)*(b*Csc[e + f*x])^(n - 2), x], x])
/; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2]

```

```
rule 5974 Int[((c_.) + (d_.)*(x_.))^(m_.)*Sech[(a_.) + (b_.)*(x_.)]^(n_.)*Tanh[(a_.) +
(b_.)*(x_.)]^(p_.), x_Symbol] := Simp[(-c + d*x)^(m)*(Sech[a + b*x]^n/(b*n)
, x] + Simp[d*(m/(b*n)) Int[(c + d*x)^(m - 1)*Sech[a + b*x]^n, x], x] /;
FreeQ[{a, b, c, d, n}, x] && EqQ[p, 1] && GtQ[m, 0]
```

```
rule 6839 Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))^(n_.)*(x_.)^(m_.), x_Symbol] := Simp[
-(c^(m + 1))^(n-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

**Maple [A] (verified)**

Time = 0.80 (sec) , antiderivative size = 329, normalized size of antiderivative = 2.35

method	result
parts	$\frac{a^2 x^3}{3} + \frac{b^2 \left( \frac{(-\operatorname{arcsech}(cx) \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} cx + c^2 x^2 \operatorname{arcsech}(cx)^2 - 1) cx}{3} + i \operatorname{arcsech}(cx) \ln \left( 1 + i \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right) \right)}{3} \right)}{3}$
derivativedivides	$\frac{a^2 c^3 x^3}{3} + b^2 \left( \frac{(-\operatorname{arcsech}(cx) \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} cx + c^2 x^2 \operatorname{arcsech}(cx)^2 - 1) cx}{3} + i \operatorname{arcsech}(cx) \ln \left( 1 + i \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right) \right)}{3} \right)$
default	$\frac{a^2 c^3 x^3}{3} + b^2 \left( \frac{(-\operatorname{arcsech}(cx) \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} cx + c^2 x^2 \operatorname{arcsech}(cx)^2 - 1) cx}{3} + i \operatorname{arcsech}(cx) \ln \left( 1 + i \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right) \right)}{3} \right)$

```
input int(x^2*(a+b*arcsech(c*x))^2,x,method=_RETURNVERBOSE)
```

```
output 1/3*a^2*x^3+b^2/c^3*(1/3*(-arcsech(c*x)*((c*x+1)/c/x)^(1/2)*(-(c*x-1)/c/x)
^(1/2)*c*x+c^2*x^2*arcsech(c*x)^2-1)*c*x+1/3*I*arcsech(c*x)*ln(1+I*(1/c/x+
(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))-1/3*I*arcsech(c*x)*ln(1-I*(1/c/x+(-1+1/
c/x)^(1/2)*(1+1/c/x)^(1/2)))+1/3*I*dilog(1+I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/
c/x)^(1/2)))-1/3*I*dilog(1-I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))))+2*
a*b/c^3*(1/3*c^3*x^3*arcsech(c*x)-1/6*(-(c*x-1)/c/x)^(1/2)*c*x*((c*x+1)/c/
x)^(1/2)*(c*x*(-c^2*x^2+1)^(1/2)-arcsin(c*x))/(-c^2*x^2+1)^(1/2))
```

**Fricas [F]**

$$\int x^2(a + b\operatorname{sech}^{-1}(cx))^2 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^2 x^2 dx$$

input `integrate(x^2*(a+b*arcsech(c*x))^2,x, algorithm="fricas")`

output `integral(b^2*x^2*arcsech(c*x)^2 + 2*a*b*x^2*arcsech(c*x) + a^2*x^2, x)`

**Sympy [F]**

$$\int x^2(a + b\operatorname{sech}^{-1}(cx))^2 dx = \int x^2(a + b \operatorname{ar} \operatorname{sech}(cx))^2 dx$$

input `integrate(x**2*(a+b*asech(c*x))**2,x)`

output `Integral(x**2*(a + b*asech(c*x))**2, x)`

**Maxima [F]**

$$\int x^2(a + b\operatorname{sech}^{-1}(cx))^2 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^2 x^2 dx$$

input `integrate(x^2*(a+b*arcsech(c*x))^2,x, algorithm="maxima")`

output `1/3*a^2*x^3 + 1/3*(2*x^3*arcsech(c*x) - (sqrt(1/(c^2*x^2) - 1)/(c^2*(1/(c^2*x^2) - 1) + c^2) + arctan(sqrt(1/(c^2*x^2) - 1))/c^2)/c)*a*b + b^2*integrate(x^2*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^2, x)`



**Giac [F]**

$$\int x^2 (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^2 x^2 dx$$

input `integrate(x^2*(a+b*arcsech(c*x))^2,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)^2*x^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int x^2 \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right)^2 dx$$

input `int(x^2*(a + b*acosh(1/(c*x)))^2,x)`

output `int(x^2*(a + b*acosh(1/(c*x)))^2, x)`

**Reduce [F]**

$$\int x^2 (a + b \operatorname{sech}^{-1}(cx))^2 dx = 2 \left( \int \operatorname{asech}(cx) x^2 dx \right) ab + \left( \int \operatorname{asech}(cx)^2 x^2 dx \right) b^2 + \frac{a^2 x^3}{3}$$

input `int(x^2*(a+b*asech(c*x))^2,x)`

output `(6*int(asech(c*x)*x**2,x)*a*b + 3*int(asech(c*x)**2*x**2,x)*b**2 + a**2*x**3)/3`

### 3.35 $\int x(a + b\operatorname{sech}^{-1}(cx))^2 dx$

Optimal result . . . . .	329
Mathematica [A] (verified) . . . . .	329
Rubi [A] (verified) . . . . .	330
Maple [B] (verified) . . . . .	332
Fricas [B] (verification not implemented) . . . . .	333
Sympy [A] (verification not implemented) . . . . .	333
Maxima [A] (verification not implemented) . . . . .	334
Giac [F] . . . . .	334
Mupad [F(-1)] . . . . .	335
Reduce [F] . . . . .	335

#### Optimal result

Integrand size = 12, antiderivative size = 65

$$\int x(a + b\operatorname{sech}^{-1}(cx))^2 dx = -\frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(a + b\operatorname{sech}^{-1}(cx))}{c^2} + \frac{1}{2}x^2(a + b\operatorname{sech}^{-1}(cx))^2 - \frac{b^2 \log(x)}{c^2}$$

output

```
-b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+b*arcsech(c*x))/c^2+1/2*x^2*(a+b*arcsech(c*x))^2-b^2*ln(x)/c^2
```

#### Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.72

$$\int x(a + b\operatorname{sech}^{-1}(cx))^2 dx = \frac{a(ac^2x^2 - 2b\sqrt{\frac{1-cx}{1+cx}}(1+cx)) - 2b(-ac^2x^2 + b\sqrt{\frac{1-cx}{1+cx}}(1+cx))\operatorname{sech}^{-1}(cx) + b^2c^2x^2\operatorname{sech}^{-1}(cx)^2 - 2b^2\ln(x)}{2c^2}$$

input

```
Integrate[x*(a + b*ArcSech[c*x])^2,x]
```

output

```
(a*(a*c^2*x^2 - 2*b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)) - 2*b*(-(a*c^2*x^2) + b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))*ArcSech[c*x] + b^2*c^2*x^2*ArcSech[c*x]^2 - 2*b^2*Log[c*x])/(2*c^2)
```

**Rubi [A] (verified)**

Time = 0.43 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.11, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {6839, 5974, 3042, 4672, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(a + b \operatorname{sech}^{-1}(cx))^2 dx$$

$$\downarrow 6839$$

$$\frac{\int c^2 x^2 \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx))^2 d \operatorname{sech}^{-1}(cx)}{c^2}$$

$$\downarrow 5974$$

$$\frac{b \int c^2 x^2 (a + b \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) - \frac{1}{2} c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^2}{c^2}$$

$$\downarrow 3042$$

$$\frac{-\frac{1}{2} c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^2 + b \int (a + b \operatorname{sech}^{-1}(cx)) \csc \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right) d \operatorname{sech}^{-1}(cx)}{c^2}$$

$$\downarrow 4672$$

$$\frac{-\frac{1}{2} c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^2 + b \left( \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx)) - i b \int -i \sqrt{\frac{1-cx}{cx+1}} (cx+1) d \operatorname{sech}^{-1}(cx) \right)}{c^2}$$

$$\downarrow 26$$

$$\frac{b \left( \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx)) - b \int \sqrt{\frac{1-cx}{cx+1}} (cx+1) d \operatorname{sech}^{-1}(cx) \right) - \frac{1}{2} c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^2}{c^2}$$

$$\downarrow 3042$$

$$\frac{-\frac{1}{2}c^2x^2(a + b\operatorname{sech}^{-1}(cx))^2 + b\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx)) - b \int -i \tan(i\operatorname{sech}^{-1}(cx)) d\operatorname{sech}^{-1}(cx)\right)}{c^2}$$

↓ 26

$$\frac{-\frac{1}{2}c^2x^2(a + b\operatorname{sech}^{-1}(cx))^2 + b\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx)) + ib \int \tan(i\operatorname{sech}^{-1}(cx)) d\operatorname{sech}^{-1}(cx)\right)}{c^2}$$

↓ 3956

$$\frac{b\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx)) - b \log\left(\frac{1}{cx}\right)\right) - \frac{1}{2}c^2x^2(a + b\operatorname{sech}^{-1}(cx))^2}{c^2}$$

input `Int[x*(a + b*ArcSech[c*x])^2,x]`

output `-((-1/2*(c^2*x^2*(a + b*ArcSech[c*x])^2) + b*(Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x]) - b*Log[1/(c*x)]))/c^2)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4672 `Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1) * Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

```
rule 5974 Int[((c_.) + (d_.)*(x_)^(m_.)*Sech[(a_.) + (b_.)*(x_)]^(n_.)*Tanh[(a_.) +
(b_.)*(x_)]^(p_.), x_Symbol] := Simp[(-c + d*x)^m*(Sech[a + b*x]^n/(b*n))
, x] + Simp[d*(m/(b*n)) Int[(c + d*x)^(m - 1)*Sech[a + b*x]^n, x], x] /;
FreeQ[{a, b, c, d, n}, x] && EqQ[p, 1] && GtQ[m, 0]
```

```
rule 6839 Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^(n_)*(x_)^(m_.), x_Symbol] := Simp[
-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 164 vs. 2(61) = 122.

Time = 0.64 (sec) , antiderivative size = 165, normalized size of antiderivative = 2.54

method	result
parts	$\frac{a^2 x^2}{2} + \frac{b^2 \left( -2 \operatorname{arcsech}(cx) + \frac{\operatorname{arcsech}(cx) \left( c^2 x^2 \operatorname{arcsech}(cx) - 2 \sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x + 2 \right)}{2} \right)}{c^2} + \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right) \right)$
derivativedivides	$\frac{\frac{a^2 c^2 x^2}{2} + b^2 \left( -2 \operatorname{arcsech}(cx) + \frac{\operatorname{arcsech}(cx) \left( c^2 x^2 \operatorname{arcsech}(cx) - 2 \sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x + 2 \right)}{2} \right)}{c^2} + \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right) \right)}{c^2}$
default	$\frac{\frac{a^2 c^2 x^2}{2} + b^2 \left( -2 \operatorname{arcsech}(cx) + \frac{\operatorname{arcsech}(cx) \left( c^2 x^2 \operatorname{arcsech}(cx) - 2 \sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x + 2 \right)}{2} \right)}{c^2} + \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right) \right)}{c^2}$

```
input int(x*(a+b*arcsech(c*x))^2,x,method=_RETURNVERBOSE)
```

```
output 1/2*a^2*x^2+b^2/c^2*(-2*arcsech(c*x)+1/2*arcsech(c*x)*(c^2*x^2*arcsech(c*x)
)-2*(-(c*x-1)/c/x)^(1/2)*c*((c*x+1)/c/x)^(1/2)*x+2)+ln(1+(1/c/x+(-1+1/c/x)
)^(1/2)*(1+1/c/x)^(1/2))^2)+2*a*b/c^2*(1/2*c^2*x^2*arcsech(c*x)-1/2*(-(c*x
-1)/c/x)^(1/2)*c*((c*x+1)/c/x)^(1/2)*x)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 205 vs.  $2(61) = 122$ .

Time = 0.10 (sec) , antiderivative size = 205, normalized size of antiderivative = 3.15

$$\int x(a + b \operatorname{sech}^{-1}(cx))^2 dx$$

$$= \frac{b^2 c^2 x^2 \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx}\right)^2 + a^2 c^2 x^2 - 2 abc^2 \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - 1}{x}\right) - 2 abcx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - 2 b^2 \log(x) + 2}{2 c^2}$$

input `integrate(x*(a+b*arcsech(c*x))^2,x, algorithm="fricas")`

output  $\frac{1}{2}*(b^2*c^2*x^2*\log((c*x*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} + 1)/(c*x))^2 + a^2*c^2*x^2 - 2*a*b*c^2*\log((c*x*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} - 1)/x) - 2*a*b*c*x*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} - 2*b^2*\log(x) + 2*(a*b*c^2*x^2 - b^2*c*x*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} - a*b*c^2)*\log((c*x*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} + 1)/(c*x)))/c^2$

**Sympy [A] (verification not implemented)**

Time = 0.38 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.52

$$\int x(a + b \operatorname{sech}^{-1}(cx))^2 dx$$

$$= \begin{cases} \frac{a^2 x^2}{2} + abx^2 \operatorname{asech}(cx) - \frac{ab\sqrt{-c^2 x^2 + 1}}{c^2} + \frac{b^2 x^2 \operatorname{asech}^2(cx)}{2} - \frac{b^2 \sqrt{-c^2 x^2 + 1} \operatorname{asech}(cx)}{c^2} - \frac{b^2 \log(x)}{c^2} & \text{for } c \neq 0 \\ \frac{x^2(a + \infty b)^2}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*(a+b*asech(c*x))**2,x)`

output `Piecewise((a**2*x**2/2 + a*b*x**2*asech(c*x) - a*b*sqrt(-c**2*x**2 + 1)/c**2 + b**2*x**2*asech(c*x)**2/2 - b**2*sqrt(-c**2*x**2 + 1)*asech(c*x)/c**2 - b**2*log(x)/c**2, Ne(c, 0)), (x**2*(a + oo*b)**2/2, True))`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.29

$$\int x(a + b \operatorname{sech}^{-1}(cx))^2 dx = \frac{1}{2} b^2 x^2 \operatorname{arsech}(cx)^2 + \frac{1}{2} a^2 x^2 + \left( x^2 \operatorname{arsech}(cx) - \frac{x \sqrt{\frac{1}{c^2 x^2} - 1}}{c} \right) ab - \left( \frac{x \sqrt{\frac{1}{c^2 x^2} - 1} \operatorname{arsech}(cx)}{c} + \frac{\log(x)}{c^2} \right) b^2$$

input `integrate(x*(a+b*arcsech(c*x))^2,x, algorithm="maxima")`output `1/2*b^2*x^2*arcsech(c*x)^2 + 1/2*a^2*x^2 + (x^2*arcsech(c*x) - x*sqrt(1/(c^2*x^2) - 1)/c)*a*b - (x*sqrt(1/(c^2*x^2) - 1)*arcsech(c*x)/c + log(x)/c^2)*b^2`**Giac [F]**

$$\int x(a + b \operatorname{sech}^{-1}(cx))^2 dx = \int (b \operatorname{arsech}(cx) + a)^2 x dx$$

input `integrate(x*(a+b*arcsech(c*x))^2,x, algorithm="giac")`output `integrate((b*arcsech(c*x) + a)^2*x, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x(a + b \operatorname{sech}^{-1}(cx))^2 dx = \int x \left( a + b \operatorname{acosh} \left( \frac{1}{cx} \right) \right)^2 dx$$

input `int(x*(a + b*acosh(1/(c*x)))^2,x)`output `int(x*(a + b*acosh(1/(c*x)))^2, x)`**Reduce [F]**

$$\int x(a + b \operatorname{sech}^{-1}(cx))^2 dx = 2 \left( \int \operatorname{asech}(cx) x dx \right) ab + \left( \int \operatorname{asech}(cx)^2 x dx \right) b^2 + \frac{a^2 x^2}{2}$$

input `int(x*(a+b*asech(c*x))^2,x)`output `(4*int(asech(c*x)*x,x)*a*b + 2*int(asech(c*x)**2*x,x)*b**2 + a**2*x**2)/2`



### 3.36 $\int (a + b \operatorname{sech}^{-1}(cx))^2 dx$

Optimal result	336
Mathematica [A] (verified)	337
Rubi [A] (verified)	337
Maple [A] (verified)	339
Fricas [F]	340
Sympy [F]	340
Maxima [F]	340
Giac [F]	341
Mupad [F(-1)]	341
Reduce [F]	342

#### Optimal result

Integrand size = 10, antiderivative size = 78

$$\int (a + b \operatorname{sech}^{-1}(cx))^2 dx = x(a + b \operatorname{sech}^{-1}(cx))^2 - \frac{4b(a + b \operatorname{sech}^{-1}(cx)) \arctan\left(e^{\operatorname{sech}^{-1}(cx)}\right)}{c} + \frac{2ib^2 \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(cx)}\right)}{c} - \frac{2ib^2 \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(cx)}\right)}{c}$$

output

```
x*(a+b*arcsech(c*x))^2-4*b*(a+b*arcsech(c*x))*arctan(1/c/x+(-1+1/c/x)^(1/2)
)*(1+1/c/x)^(1/2))/c+2*I*b^2*polylog(2,-I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)
)^(1/2))/c-2*I*b^2*polylog(2,I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/
c
```

**Mathematica [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.62

$$\int (a + b \operatorname{sech}^{-1}(cx))^2 dx = a^2 x + \frac{2ab(cx \operatorname{sech}^{-1}(cx) - 2 \arctan(\tanh(\frac{1}{2} \operatorname{sech}^{-1}(cx))))}{c} + \frac{ib^2 \left( \operatorname{sech}^{-1}(cx) \left( -icx \operatorname{sech}^{-1}(cx) + 2 \log(1 - ie^{-\operatorname{sech}^{-1}(cx)}) \right) - 2 \log(1 + ie^{-\operatorname{sech}^{-1}(cx)}) \right)}{c} + 2 \operatorname{PolyLog} \left( 2, \frac{-1 + i \operatorname{sech}^{-1}(cx)}{1 + i \operatorname{sech}^{-1}(cx)} \right) + 2 \operatorname{PolyLog} \left( 2, \frac{1 + i \operatorname{sech}^{-1}(cx)}{1 - i \operatorname{sech}^{-1}(cx)} \right) \Big/ c$$

input

```
Integrate[(a + b*ArcSech[c*x])^2,x]
```

output

```
a^2*x + (2*a*b*(c*x*ArcSech[c*x] - 2*ArcTan[Tanh[ArcSech[c*x]/2]]))/c + (I*b^2*(ArcSech[c*x]*((-I)*c*x*ArcSech[c*x] + 2*Log[1 - I/E^ArcSech[c*x]] - 2*Log[1 + I/E^ArcSech[c*x]]) + 2*PolyLog[2, (-I)/E^ArcSech[c*x]] - 2*PolyLog[2, I/E^ArcSech[c*x]]))/c
```

**Rubi [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.96, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {6833, 5974, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{sech}^{-1}(cx))^2 dx$$

$$\downarrow \text{6833}$$

$$\frac{\int cx \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx))^2 d \operatorname{sech}^{-1}(cx)}{c}$$

$$\downarrow \text{5974}$$

$$\frac{2b \int cx (a + b \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) - cx (a + b \operatorname{sech}^{-1}(cx))^2}{c}$$

$$\downarrow \text{3042}$$

$$\frac{-cx(a + b\operatorname{sech}^{-1}(cx))^2 + 2b \int (a + b\operatorname{sech}^{-1}(cx)) \csc\left(\operatorname{isech}^{-1}(cx) + \frac{\pi}{2}\right) d\operatorname{sech}^{-1}(cx)}{c}$$

↓ 4668

$$\frac{-cx(a + b\operatorname{sech}^{-1}(cx))^2 + 2b\left(-ib \int \log\left(1 - ie^{\operatorname{sech}^{-1}(cx)}\right) d\operatorname{sech}^{-1}(cx) + ib \int \log\left(1 + ie^{\operatorname{sech}^{-1}(cx)}\right) d\operatorname{sech}^{-1}(cx)\right)}{c}$$

↓ 2715

$$\frac{-cx(a + b\operatorname{sech}^{-1}(cx))^2 + 2b\left(-ib \int e^{-\operatorname{sech}^{-1}(cx)} \log\left(1 - ie^{\operatorname{sech}^{-1}(cx)}\right) de^{\operatorname{sech}^{-1}(cx)} + ib \int e^{-\operatorname{sech}^{-1}(cx)} \log\left(1 + ie^{\operatorname{sech}^{-1}(cx)}\right) de^{\operatorname{sech}^{-1}(cx)}\right)}{c}$$

↓ 2838

$$\frac{-cx(a + b\operatorname{sech}^{-1}(cx))^2 + 2b\left(2 \arctan\left(e^{\operatorname{sech}^{-1}(cx)}\right) (a + b\operatorname{sech}^{-1}(cx)) - ib \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(cx)}\right) + ib \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(cx)}\right)\right)}{c}$$

input `Int[(a + b*ArcSech[c*x])^2,x]`

output `-((-c*x*(a + b*ArcSech[c*x])^2) + 2*b*(2*(a + b*ArcSech[c*x])*ArcTan[E^ArcSech[c*x]] - I*b*PolyLog[2, (-I)*E^ArcSech[c*x]] + I*b*PolyLog[2, I*E^ArcSech[c*x]]))/c)`

### Defintions of rubi rules used

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] :> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668

```
Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

rule 5974

```
Int[((c_.) + (d_.)*(x_))^(m_.)*Sech[(a_.) + (b_.)*(x_)]^(n_.)*Tanh[(a_.) + (b_.)*(x_)]^(p_.), x_Symbol] := Simp[(-c + d*x)^m*(Sech[a + b*x]^n/(b^n)), x] + Simp[d*(m/(b^n)) Int[(c + d*x)^(m - 1)*Sech[a + b*x]^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[p, 1] && GtQ[m, 0]
```

rule 6833

```
Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[-c^(-1) Subst[Int[(a + b*x)^n*Sech[x]*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[n, 0]
```

## Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 232, normalized size of antiderivative = 2.97

method	result
derivativedivides	$\frac{cx a^2 + b^2 \left( \operatorname{arcsech}(cx)^2 cx + 2i \operatorname{arcsech}(cx) \ln \left( 1 + i \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right) \right) - 2i \operatorname{arcsech}(cx) \ln \left( 1 - i \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right) \right) \right)}{c}$
default	$\frac{cx a^2 + b^2 \left( \operatorname{arcsech}(cx)^2 cx + 2i \operatorname{arcsech}(cx) \ln \left( 1 + i \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right) \right) - 2i \operatorname{arcsech}(cx) \ln \left( 1 - i \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right) \right) \right)}{c}$
parts	$a^2 x + \frac{b^2 \left( \operatorname{arcsech}(cx)^2 cx + 2i \operatorname{arcsech}(cx) \ln \left( 1 + i \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right) \right) - 2i \operatorname{arcsech}(cx) \ln \left( 1 - i \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right) \right) \right)}{c}$

input

```
int((a+b*arcsech(c*x))^2,x,method=_RETURNVERBOSE)
```

output

```
1/c*(c*x*a^2+b^2*(arcsech(c*x)^2*c*x+2*I*arcsech(c*x)*ln(1+I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))))-2*I*arcsech(c*x)*ln(1-I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))+2*I*dilog(1+I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))-2*I*dilog(1-I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))))+2*a*b*(c*x*arcsech(c*x)-arctan((-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))
```

**Fricas [F]**

$$\int (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^2 dx$$

input `integrate((a+b*arcsech(c*x))^2,x, algorithm="fricas")`

output `integral(b^2*arcsech(c*x)^2 + 2*a*b*arcsech(c*x) + a^2, x)`

**Sympy [F]**

$$\int (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int (a + b \operatorname{ar} \operatorname{sech}(cx))^2 dx$$

input `integrate((a+b*asech(c*x))**2,x)`

output `Integral((a + b*asech(c*x))**2, x)`

**Maxima [F]**

$$\int (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^2 dx$$

input `integrate((a+b*arcsech(c*x))^2,x, algorithm="maxima")`

output

```
(x*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)^2 - integrate(-(c^2*x^2*log(c)^2
+ (c^2*x^2 - 1)*log(x)^2 + (c^2*x^2*log(c)^2 + (c^2*x^2 - 1)*log(x)^2 - lo
g(c)^2 + 2*(c^2*x^2*log(c) - log(c))*log(x))*sqrt(c*x + 1)*sqrt(-c*x + 1)
- 2*(c^2*x^2*log(c) + (c^2*x^2*(log(c) + 1) + (c^2*x^2 - 1)*log(x) - log(c)
))*sqrt(c*x + 1)*sqrt(-c*x + 1) + (c^2*x^2 - 1)*log(x) - log(c))*log(sqrt(
c*x + 1)*sqrt(-c*x + 1) + 1) - log(c)^2 + 2*(c^2*x^2*log(c) - log(c))*log(
x))/(c^2*x^2 + (c^2*x^2 - 1)*sqrt(c*x + 1)*sqrt(-c*x + 1) - 1), x))*b^2 +
a^2*x + 2*(c*x*arcsech(c*x) - arctan(sqrt(1/(c^2*x^2) - 1)))*a*b/c
```

**Giac [F]**

$$\int (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^2 dx$$

input

```
integrate((a+b*arcsech(c*x))^2,x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)^2, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right)^2 dx$$

input

```
int((a + b*acosh(1/(c*x)))^2,x)
```

output

```
int((a + b*acosh(1/(c*x)))^2, x)
```

**Reduce [F]**

$$\int (a + b \operatorname{sech}^{-1}(cx))^2 dx = 2 \left( \int \operatorname{asech}(cx) dx \right) ab + \left( \int \operatorname{asech}(cx)^2 dx \right) b^2 + a^2 x$$

input `int((a+b*asech(c*x))^2,x)`

output `2*int(asech(c*x),x)*a*b + int(asech(c*x)**2,x)*b**2 + a**2*x`

**3.37**  $\int \frac{(a+b\operatorname{sech}^{-1}(cx))^2}{x} dx$

Optimal result	343
Mathematica [A] (verified)	344
Rubi [C] (verified)	344
Maple [A] (verified)	347
Fricas [F]	348
Sympy [F]	348
Maxima [F]	349
Giac [F]	349
Mupad [F(-1)]	349
Reduce [F]	350

**Optimal result**

Integrand size = 14, antiderivative size = 83

$$\int \frac{(a + b\operatorname{sech}^{-1}(cx))^2}{x} dx = \frac{(a + b\operatorname{sech}^{-1}(cx))^3}{3b} - (a + b\operatorname{sech}^{-1}(cx))^2 \log\left(1 + e^{2\operatorname{sech}^{-1}(cx)}\right) - b(a + b\operatorname{sech}^{-1}(cx)) \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(cx)}\right) + \frac{1}{2}b^2 \operatorname{PolyLog}\left(3, -e^{2\operatorname{sech}^{-1}(cx)}\right)$$

output

```
1/3*(a+b*arcsech(c*x))^3/b-(a+b*arcsech(c*x))^2*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)-b*(a+b*arcsech(c*x))*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)+1/2*b^2*polylog(3,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)
```



**Mathematica [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.40

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x} dx = a^2 \log(cx) + ab \left( -\operatorname{sech}^{-1}(cx) \left( \operatorname{sech}^{-1}(cx) + 2 \log \left( 1 + e^{-2 \operatorname{sech}^{-1}(cx)} \right) \right) + \operatorname{PolyLog} \left( 2, -e^{-2 \operatorname{sech}^{-1}(cx)} \right) \right) + b^2 \left( -\frac{1}{3} \operatorname{sech}^{-1}(cx)^3 - \operatorname{sech}^{-1}(cx)^2 \log \left( 1 + e^{-2 \operatorname{sech}^{-1}(cx)} \right) + \operatorname{sech}^{-1}(cx) \operatorname{PolyLog} \left( 2, -e^{-2 \operatorname{sech}^{-1}(cx)} \right) + \frac{1}{2} \operatorname{PolyLog} \left( 3, -e^{-2 \operatorname{sech}^{-1}(cx)} \right) \right)$$

input `Integrate[(a + b*ArcSech[c*x])^2/x, x]`

output `a^2*Log[c*x] + a*b*(-(ArcSech[c*x]*(ArcSech[c*x] + 2*Log[1 + E^(-2*ArcSech[c*x])])) + PolyLog[2, -E^(-2*ArcSech[c*x])]) + b^2*(-1/3*ArcSech[c*x]^3 - ArcSech[c*x]^2*Log[1 + E^(-2*ArcSech[c*x])] + ArcSech[c*x]*PolyLog[2, -E^(-2*ArcSech[c*x])] + PolyLog[3, -E^(-2*ArcSech[c*x])])/2)`

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.54 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.19, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$ , Rules used = {6839, 3042, 26, 4201, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x} dx$$

↓ 6839

$$-\int \sqrt{\frac{1-cx}{cx+1}}(cx+1)(a+b\operatorname{sech}^{-1}(cx))^2 d\operatorname{sech}^{-1}(cx)$$

↓ 3042

$$-\int -i(a+b\operatorname{sech}^{-1}(cx))^2 \tan(i\operatorname{sech}^{-1}(cx)) d\operatorname{sech}^{-1}(cx)$$

↓ 26

$$i \int (a+b\operatorname{sech}^{-1}(cx))^2 \tan(i\operatorname{sech}^{-1}(cx)) d\operatorname{sech}^{-1}(cx)$$

↓ 4201

$$i \left( 2i \int \frac{e^{2\operatorname{sech}^{-1}(cx)}(a+b\operatorname{sech}^{-1}(cx))^2}{1+e^{2\operatorname{sech}^{-1}(cx)}} d\operatorname{sech}^{-1}(cx) - \frac{i(a+b\operatorname{sech}^{-1}(cx))^3}{3b} \right)$$

↓ 2620

$$i \left( 2i \left( \frac{1}{2} \log(e^{2\operatorname{sech}^{-1}(cx)} + 1) (a+b\operatorname{sech}^{-1}(cx))^2 - b \int (a+b\operatorname{sech}^{-1}(cx)) \log(1+e^{2\operatorname{sech}^{-1}(cx)}) d\operatorname{sech}^{-1}(cx) \right) - \right)$$

↓ 3011

$$i \left( 2i \left( \frac{1}{2} \log(e^{2\operatorname{sech}^{-1}(cx)} + 1) (a+b\operatorname{sech}^{-1}(cx))^2 - b \left( \frac{1}{2} b \int \operatorname{PolyLog}(2, -e^{2\operatorname{sech}^{-1}(cx)}) d\operatorname{sech}^{-1}(cx) - \frac{1}{2} \operatorname{PolyLog}(2, -e^{2\operatorname{sech}^{-1}(cx)}) \right) \right) \right)$$

↓ 2720

$$i \left( 2i \left( \frac{1}{2} \log(e^{2\operatorname{sech}^{-1}(cx)} + 1) (a+b\operatorname{sech}^{-1}(cx))^2 - b \left( \frac{1}{4} b \int e^{-2\operatorname{sech}^{-1}(cx)} \operatorname{PolyLog}(2, -e^{2\operatorname{sech}^{-1}(cx)}) de^{2\operatorname{sech}^{-1}(cx)} \right) \right) \right)$$

↓ 7143

$$i \left( 2i \left( \frac{1}{2} \log(e^{2\operatorname{sech}^{-1}(cx)} + 1) (a+b\operatorname{sech}^{-1}(cx))^2 - b \left( \frac{1}{4} b \operatorname{PolyLog}(3, -e^{2\operatorname{sech}^{-1}(cx)}) - \frac{1}{2} \operatorname{PolyLog}(2, -e^{2\operatorname{sech}^{-1}(cx)}) \right) \right) \right)$$

input

Int[(a + b\*ArcSech[c\*x])^2/x,x]

output

$$I * \left( \left( -\frac{1}{3} I \right) * (a + b \operatorname{ArcSech}[c*x])^3 / b + (2*I) * \left( (a + b \operatorname{ArcSech}[c*x])^2 \operatorname{Log}[1 + E^{(2*\operatorname{ArcSech}[c*x])}] \right) / 2 - b * \left( -\frac{1}{2} * (a + b \operatorname{ArcSech}[c*x]) * \operatorname{PolyLog}[2, -E^{(2*\operatorname{ArcSech}[c*x])}] \right) + (b * \operatorname{PolyLog}[3, -E^{(2*\operatorname{ArcSech}[c*x])}] / 4) \right)$$

### Defintions of rubi rules used

rule 26

$$\operatorname{Int}[(\operatorname{Complex}[0, a]) * (F x), x\_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Complex}[\operatorname{Identity}[0], a]) \operatorname{Int}[F x, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ \operatorname{EqQ}[a^2, 1]$$

rule 2620

$$\operatorname{Int}[\left( (F)^{(g \cdot (e \cdot) + (f \cdot)(x \cdot))} \right)^{(n \cdot)} * \left( (c \cdot) + (d \cdot)(x \cdot) \right)^{(m \cdot)} / \left( (a \cdot) + (b \cdot) * (F)^{(g \cdot (e \cdot) + (f \cdot)(x \cdot))} \right)^{(n \cdot)}, x\_Symbol] \rightarrow \operatorname{Simp}[\left( (c + d*x)^m / (b*f*g*n*\operatorname{Log}[F]) \right) * \operatorname{Log}[1 + b * (F^{(g*(e + f*x))})^n / a], x] - \operatorname{Simp}[d * (m / (b*f*g*n*\operatorname{Log}[F])) \operatorname{Int}[(c + d*x)^{(m-1)} * \operatorname{Log}[1 + b * (F^{(g*(e + f*x))})^n / a], x], x] /; \operatorname{FreeQ}\{F, a, b, c, d, e, f, g, n\}, x] \ \&\& \ \operatorname{IGtQ}[m, 0]$$

rule 2720

$$\operatorname{Int}[u, x\_Symbol] \rightarrow \operatorname{With}\{v = \operatorname{FunctionOfExponential}[u, x]\}, \operatorname{Simp}[v/D[v, x] \operatorname{Subst}[\operatorname{Int}[\operatorname{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x] /; \operatorname{FunctionOfExponentialQ}[u, x] \ \&\& \ \operatorname{!MatchQ}[u, (w \cdot) * ((a \cdot) * (v \cdot)^{(n \cdot)})^m] /; \operatorname{FreeQ}\{a, m, n\}, x] \ \&\& \ \operatorname{IntegerQ}[m*n] \ \&\& \ \operatorname{!MatchQ}[u, E^{(c \cdot) * ((a \cdot) + (b \cdot) * x)}] * (F \cdot)[v \cdot] /; \operatorname{FreeQ}\{a, b, c\}, x] \ \&\& \ \operatorname{InverseFunctionQ}[F[x]]$$

rule 3011

$$\operatorname{Int}[\operatorname{Log}[1 + (e \cdot) * (F)^{(c \cdot) * ((a \cdot) + (b \cdot) * (x \cdot))}]^n] * ((f \cdot) + (g \cdot) * (x \cdot))^{(m \cdot)}, x\_Symbol] \rightarrow \operatorname{Simp}[(-f + g*x)^m * (\operatorname{PolyLog}[2, (-e) * (F^{(c*(a + b*x))})^n] / (b*c*n*\operatorname{Log}[F])), x] + \operatorname{Simp}[g * (m / (b*c*n*\operatorname{Log}[F])) \operatorname{Int}[(f + g*x)^{(m-1)} * \operatorname{PolyLog}[2, (-e) * (F^{(c*(a + b*x))})^n], x], x] /; \operatorname{FreeQ}\{F, a, b, c, e, f, g, n\}, x] \ \&\& \ \operatorname{GtQ}[m, 0]$$

rule 3042

$$\operatorname{Int}[u, x\_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$$

rule 4201 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))]], x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6839 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))^(n_)*(x_)^(m_.), x_Symbol] := Simp[-(c^(m + 1))^(n-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.))]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

## Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 241, normalized size of antiderivative = 2.90

method	result
parts	$a^2 \ln(x) + b^2 \left( \frac{\operatorname{arcsech}(cx)^3}{3} - \operatorname{arcsech}(cx)^2 \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right)^2 \right) - \operatorname{arcsech}(cx) \ln \left( 1 + \frac{1}{cx} \right) \right)$
derivativedivides	$a^2 \ln(cx) + b^2 \left( \frac{\operatorname{arcsech}(cx)^3}{3} - \operatorname{arcsech}(cx)^2 \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right)^2 \right) - \operatorname{arcsech}(cx) \ln \left( 1 + \frac{1}{cx} \right) \right)$
default	$a^2 \ln(cx) + b^2 \left( \frac{\operatorname{arcsech}(cx)^3}{3} - \operatorname{arcsech}(cx)^2 \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right)^2 \right) - \operatorname{arcsech}(cx) \ln \left( 1 + \frac{1}{cx} \right) \right)$

input `int((a+b*arcsech(c*x))^2/x,x,method=_RETURNVERBOSE)`

output

```
a^2*ln(x)+b^2*(1/3*arcsech(c*x)^3-arcsech(c*x)^2*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)-arcsech(c*x)*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)+1/2*polylog(3,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2))+2*a*b*(1/2*arcsech(c*x)^2-arcsech(c*x)*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2))-1/2*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2))
```

**Fricas [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2}{x} dx$$

input

```
integrate((a+b*arcsech(c*x))^2/x,x, algorithm="fricas")
```

output

```
integral((b^2*arcsech(c*x)^2 + 2*a*b*arcsech(c*x) + a^2)/x, x)
```

**Sympy [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x} dx = \int \frac{(a + b \operatorname{ar} \operatorname{sech}(cx))^2}{x} dx$$

input

```
integrate((a+b*asech(c*x))**2/x,x)
```

output

```
Integral((a + b*asech(c*x))**2/x, x)
```

**Maxima [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2}{x} dx$$

input `integrate((a+b*arcsech(c*x))^2/x,x, algorithm="maxima")`

output `a^2*log(x) + integrate(b^2*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^2/x + 2*a*b*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/x, x)`

**Giac [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2}{x} dx$$

input `integrate((a+b*arcsech(c*x))^2/x,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)^2/x, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x} dx = \int \frac{(a + b \operatorname{acosh}(\frac{1}{cx}))^2}{x} dx$$

input `int((a + b*acosh(1/(c*x)))^2/x,x)`

output `int((a + b*acosh(1/(c*x)))^2/x, x)`

**Reduce [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x} dx = 2 \left( \int \frac{\operatorname{asech}(cx)}{x} dx \right) ab + \left( \int \frac{\operatorname{asech}(cx)^2}{x} dx \right) b^2 + \log(x) a^2$$

input `int((a+b*asech(c*x))^2/x,x)`

output `2*int(asech(c*x)/x,x)*a*b + int(asech(c*x)**2/x,x)*b**2 + log(x)*a**2`

**3.38**  $\int \frac{(a+b\operatorname{sech}^{-1}(cx))^2}{x^2} dx$

Optimal result	351
Mathematica [A] (verified)	351
Rubi [C] (verified)	352
Maple [B] (verified)	354
Fricas [B] (verification not implemented)	355
Sympy [F]	355
Maxima [A] (verification not implemented)	356
Giac [F]	356
Mupad [F(-1)]	356
Reduce [F]	357

**Optimal result**

Integrand size = 14, antiderivative size = 61

$$\int \frac{(a + b\operatorname{sech}^{-1}(cx))^2}{x^2} dx = -\frac{2b^2}{x} + \frac{2b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(a + b\operatorname{sech}^{-1}(cx))}{x} - \frac{(a + b\operatorname{sech}^{-1}(cx))^2}{x}$$

output

```
-2*b^2/x+2*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+b*arcsech(c*x))/x-(a+b*arcsech(c*x))^2/x
```

**Mathematica [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.43

$$\int \frac{(a + b\operatorname{sech}^{-1}(cx))^2}{x^2} dx = \frac{a^2 + 2b^2 - 2ab\sqrt{\frac{1-cx}{1+cx}}(1+cx) - 2b\left(-a + b\sqrt{\frac{1-cx}{1+cx}}(1+cx)\right)\operatorname{sech}^{-1}(cx) + b^2\operatorname{sech}^{-1}(cx)^2}{x}$$

input

```
Integrate[(a + b*ArcSech[c*x])^2/x^2,x]
```



output

```

-((a^2 + 2*b^2 - 2*a*b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x) - 2*b*(-a + b*S
qrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))*ArcSech[c*x] + b^2*ArcSech[c*x]^2)/x)

```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.40 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.30, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$ , Rules used = {6839, 3042, 26, 3777, 3042, 3777, 26, 3042, 26, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^2} dx \\
& \quad \downarrow \text{6839} \\
& -c \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{cx} d \operatorname{sech}^{-1}(cx) \\
& \quad \downarrow \text{3042} \\
& -c \int -i(a + b \operatorname{sech}^{-1}(cx))^2 \sin(i \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) \\
& \quad \downarrow \text{26} \\
& ic \int (a + b \operatorname{sech}^{-1}(cx))^2 \sin(i \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) \\
& \quad \downarrow \text{3777} \\
& ic \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))^2}{cx} - 2ib \int \frac{a + b \operatorname{sech}^{-1}(cx)}{cx} d \operatorname{sech}^{-1}(cx) \right) \\
& \quad \downarrow \text{3042} \\
& ic \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))^2}{cx} - 2ib \int (a + b \operatorname{sech}^{-1}(cx)) \sin\left(i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2}\right) d \operatorname{sech}^{-1}(cx) \right) \\
& \quad \downarrow \text{3777}
\end{aligned}$$

$$ic \left( \frac{i(a + b\operatorname{sech}^{-1}(cx))^2}{cx} - 2ib \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))}{cx} - ib \int -\frac{i\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{cx} d\operatorname{sech}^{-1}(cx) \right) \right)$$

↓ 26

$$ic \left( \frac{i(a + b\operatorname{sech}^{-1}(cx))^2}{cx} - 2ib \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))}{cx} - b \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{cx} d\operatorname{sech}^{-1}(cx) \right) \right)$$

↓ 3042

$$ic \left( \frac{i(a + b\operatorname{sech}^{-1}(cx))^2}{cx} - 2ib \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))}{cx} - b \int -i \sin(\operatorname{sech}^{-1}(cx)) d\operatorname{sech}^{-1}(cx) \right) \right)$$

↓ 26

$$ic \left( \frac{i(a + b\operatorname{sech}^{-1}(cx))^2}{cx} - 2ib \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))}{cx} + ib \int \sin(\operatorname{sech}^{-1}(cx)) d\operatorname{sech}^{-1}(cx) \right) \right)$$

↓ 3118

$$ic \left( \frac{i(a + b\operatorname{sech}^{-1}(cx))^2}{cx} - 2ib \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))}{cx} - \frac{b}{cx} \right) \right)$$

input `Int[(a + b*ArcSech[c*x])^2/x^2,x]`

output `I*c*((I*(a + b*ArcSech[c*x])^2)/(c*x) - (2*I)*b*(-(b/(c*x)) + (Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x]))/(c*x)))`

## Definitions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a_])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3118  $\text{Int}[\sin[(c_.) + (d_.)*(x_)], x\_Symbol] \rightarrow \text{Simp}[-\text{Cos}[c + d*x]/d, x] /; \text{FreeQ}[\{c, d\}, x]$
- rule 3777  $\text{Int}[((c_.) + (d_.)*(x_))^{(m_.)}*\sin[(e_.) + (f_.)*(x_)], x\_Symbol] \rightarrow \text{Simp}[(-c + d*x)^m*(\text{Cos}[e + f*x]/f), x] + \text{Simp}[d*(m/f) \text{Int}[(c + d*x)^{m-1}*\text{Cos}[e + f*x], x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \ \&\& \ \text{GtQ}[m, 0]$
- rule 6839  $\text{Int}[((a_.) + \text{ArcSech}[(c_.)*(x_)]*(b_.) )^{(n_.)}*(x_)^{(m_.)}, x\_Symbol] \rightarrow \text{Simp}[-(c^{m+1})^{-1} \text{Subst}[\text{Int}[(a + b*x)^n*\text{Sech}[x]^{m+1}*\text{Tanh}[x], x], x, \text{ArcSech}[c*x]], x] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ (\text{GtQ}[n, 0] \ || \ \text{LtQ}[m, -1])$

## Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 120 vs.  $2(59) = 118$ .

Time = 0.49 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.98

method	result
parts	$-\frac{a^2}{x} + b^2c \left( -\frac{\text{arcsech}(cx)^2}{cx} + 2 \text{arcsech}(cx) \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} - \frac{2}{cx} \right) + 2abc \left( -\frac{\text{arcsech}(cx)}{cx} + \right.$
derivativedivides	$c \left( -\frac{a^2}{cx} + b^2 \left( -\frac{\text{arcsech}(cx)^2}{cx} + 2 \text{arcsech}(cx) \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} - \frac{2}{cx} \right) + 2ab \left( -\frac{\text{arcsech}(cx)}{cx} + \right.$
default	$c \left( -\frac{a^2}{cx} + b^2 \left( -\frac{\text{arcsech}(cx)^2}{cx} + 2 \text{arcsech}(cx) \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} - \frac{2}{cx} \right) + 2ab \left( -\frac{\text{arcsech}(cx)}{cx} + \right.$

input  $\text{int}((a+b*\text{arcsech}(c*x))^2/x^2, x, \text{method}=\_RETURNVERBOSE)$

output

```
-1/x*a^2+b^2*c*(-1/c/x*arcsech(c*x)^2+2*arcsech(c*x)*(-(c*x-1)/c/x)^(1/2)*
((c*x+1)/c/x)^(1/2)-2/c/x)+2*a*b*c*(-1/c/x*arcsech(c*x)+(-(c*x-1)/c/x)^(1/2)*
2)*((c*x+1)/c/x)^(1/2))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 143 vs.  $2(59) = 118$ .

Time = 0.09 (sec) , antiderivative size = 143, normalized size of antiderivative = 2.34

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^2} dx$$

$$= \frac{2 abcx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - b^2 \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx}\right)^2 - a^2 - 2b^2 + 2\left(b^2 cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - ab\right) \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx}\right)}{x}$$

input

```
integrate((a+b*arcsech(c*x))^2/x^2,x, algorithm="fricas")
```

output

```
(2*a*b*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - b^2*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x))^2 - a^2 - 2*b^2 + 2*(b^2*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - a*b)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)))/x
```

**Sympy [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^2} dx = \int \frac{(a + b \operatorname{asech}(cx))^2}{x^2} dx$$

input

```
integrate((a+b*asech(c*x))**2/x**2,x)
```

output

```
Integral((a + b*asech(c*x))**2/x**2, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.28

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^2} dx = 2 \left( c \sqrt{\frac{1}{c^2 x^2} - 1} - \frac{\operatorname{arsh}(cx)}{x} \right) ab + 2 \left( c \sqrt{\frac{1}{c^2 x^2} - 1} \operatorname{arsh}(cx) - \frac{1}{x} \right) b^2 - \frac{b^2 \operatorname{arsh}(cx)^2}{x} - \frac{a^2}{x}$$

input `integrate((a+b*arcsech(c*x))^2/x^2,x, algorithm="maxima")`output `2*(c*sqrt(1/(c^2*x^2) - 1) - arcsech(c*x)/x)*a*b + 2*(c*sqrt(1/(c^2*x^2) - 1)*arcsech(c*x) - 1/x)*b^2 - b^2*arcsech(c*x)^2/x - a^2/x`**Giac [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^2} dx = \int \frac{(b \operatorname{arsh}(cx) + a)^2}{x^2} dx$$

input `integrate((a+b*arcsech(c*x))^2/x^2,x, algorithm="giac")`output `integrate((b*arcsech(c*x) + a)^2/x^2, x)`**Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(\frac{1}{cx}))^2}{x^2} dx$$

input `int((a + b*acosh(1/(c*x)))^2/x^2,x)`output `int((a + b*acosh(1/(c*x)))^2/x^2, x)`

**Reduce [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^2} dx = \frac{2 \left( \int \frac{\operatorname{asech}(cx)}{x^2} dx \right) abx + \left( \int \frac{\operatorname{asech}(cx)^2}{x^2} dx \right) b^2x - a^2}{x}$$

input `int((a+b*asech(c*x))^2/x^2,x)`

output `(2*int(asech(c*x)/x**2,x)*a*b*x + int(asech(c*x)**2/x**2,x)*b**2*x - a**2)/x`

**3.39** 
$$\int \frac{(a+b\operatorname{sech}^{-1}(cx))^2}{x^3} dx$$

Optimal result	358
Mathematica [A] (verified)	359
Rubi [A] (verified)	359
Maple [A] (verified)	362
Fricas [A] (verification not implemented)	362
Sympy [F]	363
Maxima [F]	363
Giac [F]	364
Mupad [F(-1)]	364
Reduce [F]	364

**Optimal result**

Integrand size = 14, antiderivative size = 106

$$\int \frac{(a + b\operatorname{sech}^{-1}(cx))^2}{x^3} dx = -\frac{b^2(1 - cx)(1 + cx)}{4x^2} + \frac{b\sqrt{\frac{1-cx}{1+cx}}(1 + cx)(a + b\operatorname{sech}^{-1}(cx))}{2x^2} - \frac{1}{4}c^2(a + b\operatorname{sech}^{-1}(cx))^2 - \frac{(1 - cx)(1 + cx)(a + b\operatorname{sech}^{-1}(cx))^2}{2x^2}$$

output

```
-1/4*b^2*(-c*x+1)*(c*x+1)/x^2+1/2*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+b*
arcsech(c*x))/x^2-1/4*c^2*(a+b*arcsech(c*x))^2-1/2*(-c*x+1)*(c*x+1)*(a+b*a
rcsech(c*x))^2/x^2
```

**Mathematica [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.73

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^3} dx$$

$$= \frac{-2a^2 - b^2 + 2ab\sqrt{\frac{1-cx}{1+cx}} + 2abcx\sqrt{\frac{1-cx}{1+cx}} + 2b\left(-2a + b\sqrt{\frac{1-cx}{1+cx}}(1+cx)\right)\operatorname{sech}^{-1}(cx) + b^2(-2 + c^2x^2)\operatorname{sech}^{-1}(cx)}{4x^2}$$

input

```
Integrate[(a + b*ArcSech[c*x])^2/x^3,x]
```

output

```
(-2*a^2 - b^2 + 2*a*b*Sqrt[(1 - c*x)/(1 + c*x)] + 2*a*b*c*x*Sqrt[(1 - c*x)/(1 + c*x)] + 2*b*(-2*a + b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))*ArcSech[c*x] + b^2*(-2 + c^2*x^2)*ArcSech[c*x]^2 - 2*a*b*c^2*x^2*Log[x] + 2*a*b*c^2*x^2*Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]])/(4*x^2)
```

**Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.13, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {6839, 5969, 3042, 25, 3791, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^3} dx$$

$$\downarrow \text{6839}$$

$$-c^2 \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{c^2x^2} d \operatorname{sech}^{-1}(cx)$$

$$\downarrow \text{5969}$$



$$-c^2 \left( \frac{(1-cx)(cx+1)(a+b\operatorname{sech}^{-1}(cx))^2}{2c^2x^2} - b \int \frac{(1-cx)(cx+1)(a+b\operatorname{sech}^{-1}(cx))}{c^2x^2} d\operatorname{sech}^{-1}(cx) \right)$$

↓ 3042

$$-c^2 \left( \frac{(1-cx)(cx+1)(a+b\operatorname{sech}^{-1}(cx))^2}{2c^2x^2} - b \int -((a+b\operatorname{sech}^{-1}(cx)) \sin(\operatorname{isech}^{-1}(cx))^2) d\operatorname{sech}^{-1}(cx) \right)$$

↓ 25

$$-c^2 \left( \frac{(1-cx)(cx+1)(a+b\operatorname{sech}^{-1}(cx))^2}{2c^2x^2} + b \int (a+b\operatorname{sech}^{-1}(cx)) \sin(\operatorname{isech}^{-1}(cx))^2 d\operatorname{sech}^{-1}(cx) \right)$$

↓ 3791

$$-c^2 \left( b \left( \frac{1}{2} \int (a+b\operatorname{sech}^{-1}(cx)) d\operatorname{sech}^{-1}(cx) - \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a+b\operatorname{sech}^{-1}(cx))}{2c^2x^2} + \frac{b(1-cx)(cx+1)}{4c^2x^2} \right) + \frac{(1-cx)(cx+1)(a+b\operatorname{sech}^{-1}(cx))^2}{2c^2x^2} \right)$$

↓ 17

$$-c^2 \left( \frac{(1-cx)(cx+1)(a+b\operatorname{sech}^{-1}(cx))^2}{2c^2x^2} + b \left( -\frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a+b\operatorname{sech}^{-1}(cx))}{2c^2x^2} + \frac{(a+b\operatorname{sech}^{-1}(cx))^2}{4b} + \frac{b(1-cx)(cx+1)}{4c^2x^2} \right) \right)$$

input `Int[(a + b*ArcSech[c*x])^2/x^3,x]`

output `-(c^2*(((1 - c*x)*(1 + c*x)*(a + b*ArcSech[c*x])^2)/(2*c^2*x^2) + b*((b*(1 - c*x)*(1 + c*x))/(4*c^2*x^2) - (Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x]))/(2*c^2*x^2) + (a + b*ArcSech[c*x])^2/(4*b))))`

## Definitions of rubi rules used

- rule 17  $\text{Int}[(c_.)((a_.) + (b_.)(x_))^{(m_.)}, x\_Symbol] \rightarrow \text{Simp}[c*((a + b*x)^{(m + 1)})/(b*(m + 1)), x] \text{ /; FreeQ}\{a, b, c, m\}, x] \ \&\& \ \text{NeQ}[m, -1]$
- rule 25  $\text{Int}[-(Fx_), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \ \text{Int}[Fx, x], x]$
- rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$
- rule 3791  $\text{Int}[((c_.) + (d_.)(x_))*((b_.)\sin[(e_.) + (f_.)(x_)])^{(n_.)}, x\_Symbol] \rightarrow \text{Simp}[d*((b*\sin[e + f*x])^n/(f^2*n^2)), x] + (-\text{Simp}[b*(c + d*x)*\cos[e + f*x] * ((b*\sin[e + f*x])^{(n - 1)})/(f*n), x] + \text{Simp}[b^2*((n - 1)/n) \ \text{Int}[(c + d*x)*(b*\sin[e + f*x])^{(n - 2)}, x], x]) \text{ /; FreeQ}\{b, c, d, e, f\}, x] \ \&\& \ \text{GtQ}[n, 1]$
- rule 5969  $\text{Int}[\text{Cosh}[(a_.) + (b_.)(x_)]*((c_.) + (d_.)(x_))^{(m_.)}\sinh[(a_.) + (b_.)(x_)]^{(n_.)}, x\_Symbol] \rightarrow \text{Simp}[(c + d*x)^m*(\sinh[a + b*x]^{(n + 1)})/(b*(n + 1)), x] - \text{Simp}[d*(m/(b*(n + 1))) \ \text{Int}[(c + d*x)^{(m - 1)}*\sinh[a + b*x]^{(n + 1)}, x], x] \text{ /; FreeQ}\{a, b, c, d, n\}, x] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{NeQ}[n, -1]$
- rule 6839  $\text{Int}[((a_.) + \text{ArcSech}[(c_.)(x_)]*(b_.))^{(n_.)}(x_)^{(m_.)}, x\_Symbol] \rightarrow \text{Simp}[-(c^{(m + 1)})^{(-1)} \ \text{Subst}[\text{Int}[(a + b*x)^n*\text{sech}[x]^{(m + 1)}*\tanh[x], x], x, \text{ArcSech}[c*x]], x] \text{ /; FreeQ}\{a, b, c\}, x] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ (\text{GtQ}[n, 0] \ || \ \text{LtQ}[m, -1])$

**Maple [A] (verified)**

Time = 0.49 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.48

method	result
parts	$-\frac{a^2}{2x^2} + b^2 c^2 \left( -\frac{\cosh(2 \operatorname{arcsech}(cx)) \operatorname{arcsech}(cx)^2}{4} + \frac{\sinh(2 \operatorname{arcsech}(cx)) \operatorname{arcsech}(cx)}{4} - \frac{\cosh(2 \operatorname{arcsech}(cx))}{8} \right)$
derivativedivides	$c^2 \left( -\frac{a^2}{2c^2 x^2} + b^2 \left( -\frac{\cosh(2 \operatorname{arcsech}(cx)) \operatorname{arcsech}(cx)^2}{4} + \frac{\sinh(2 \operatorname{arcsech}(cx)) \operatorname{arcsech}(cx)}{4} - \frac{\cosh(2 \operatorname{arcsech}(cx))}{8} \right) \right)$
default	$c^2 \left( -\frac{a^2}{2c^2 x^2} + b^2 \left( -\frac{\cosh(2 \operatorname{arcsech}(cx)) \operatorname{arcsech}(cx)^2}{4} + \frac{\sinh(2 \operatorname{arcsech}(cx)) \operatorname{arcsech}(cx)}{4} - \frac{\cosh(2 \operatorname{arcsech}(cx))}{8} \right) \right)$

input `int((a+b*arcsech(c*x))^2/x^3,x,method=_RETURNVERBOSE)`

output 
$$-1/2*a^2/x^2+b^2*c^2*(-1/4*\cosh(2*\operatorname{arcsech}(c*x))*\operatorname{arcsech}(c*x)^2+1/4*\sinh(2*\operatorname{arcsech}(c*x))*\operatorname{arcsech}(c*x)-1/8*\cosh(2*\operatorname{arcsech}(c*x)))+2*a*b*c^2*(-1/2/c^2/x^2*\operatorname{arcsech}(c*x)+1/4*(-(c*x-1)/c/x)^{(1/2)}/c/x*((c*x+1)/c/x)^{(1/2)}*(\operatorname{arctanh}(1/(-c^2*x^2+1)^{(1/2)})*c^2*x^2+(-c^2*x^2+1)^{(1/2)})/(-c^2*x^2+1)^{(1/2)})$$

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.56

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^3} dx$$

$$= \frac{2 abcx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + (b^2 c^2 x^2 - 2 b^2) \log \left( \frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx} \right)^2 - 2 a^2 - b^2 + 2 \left( abc^2 x^2 + b^2 cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - 2 ab \right)}{4 x^2}$$

input `integrate((a+b*arcsech(c*x))^2/x^3,x, algorithm="fricas")`

output

```
1/4*(2*a*b*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + (b^2*c^2*x^2 - 2*b^2)*log(
(c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x))^2 - 2*a^2 - b^2 + 2*(a*b*c
^2*x^2 + b^2*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 2*a*b)*log((c*x*sqrt(-(c
^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)))/x^2
```

**Sympy [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^3} dx = \int \frac{(a + b \operatorname{arsech}(cx))^2}{x^3} dx$$

input

```
integrate((a+b*asech(c*x))**2/x**3,x)
```

output

```
Integral((a + b*asech(c*x))**2/x**3, x)
```

**Maxima [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^3} dx = \int \frac{(b \operatorname{arsech}(cx) + a)^2}{x^3} dx$$

input

```
integrate((a+b*arcsech(c*x))^2/x^3,x, algorithm="maxima")
```

output

```
-1/4*a*b*((2*c^4*x*sqrt(1/(c^2*x^2) - 1)/(c^2*x^2*(1/(c^2*x^2) - 1) - 1) -
c^3*log(c*x*sqrt(1/(c^2*x^2) - 1) + 1) + c^3*log(c*x*sqrt(1/(c^2*x^2) - 1
) - 1))/c + 4*arcsech(c*x)/x^2) + b^2*integrate(log(sqrt(1/(c*x) + 1)*sqrt
(1/(c*x) - 1) + 1/(c*x))^2/x^3, x) - 1/2*a^2/x^2
```

**Giac [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^3} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2}{x^3} dx$$

input `integrate((a+b*arcsech(c*x))^2/x^3,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)^2/x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(\frac{1}{cx}))^2}{x^3} dx$$

input `int((a + b*acosh(1/(c*x)))^2/x^3,x)`

output `int((a + b*acosh(1/(c*x)))^2/x^3, x)`

**Reduce [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^3} dx = \frac{4 \left( \int \frac{\operatorname{asech}(cx)}{x^3} dx \right) ab x^2 + 2 \left( \int \frac{\operatorname{asech}(cx)^2}{x^3} dx \right) b^2 x^2 - a^2}{2x^2}$$

input `int((a+b*asech(c*x))^2/x^3,x)`

output `(4*int(asech(c*x)/x**3,x)*a*b*x**2 + 2*int(asech(c*x)**2/x**3,x)*b**2*x**2 - a**2)/(2*x**2)`

**3.40**  $\int \frac{(a+b\operatorname{sech}^{-1}(cx))^2}{x^4} dx$

Optimal result	365
Mathematica [A] (verified)	366
Rubi [A] (verified)	366
Maple [A] (verified)	369
Fricas [A] (verification not implemented)	370
Sympy [F]	370
Maxima [F]	371
Giac [F]	371
Mupad [F(-1)]	371
Reduce [F]	372

**Optimal result**

Integrand size = 14, antiderivative size = 122

$$\int \frac{(a + b\operatorname{sech}^{-1}(cx))^2}{x^4} dx = -\frac{2b^2}{27x^3} - \frac{4b^2c^2}{9x} + \frac{2b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(a + b\operatorname{sech}^{-1}(cx))}{9x^3} + \frac{4bc^2\sqrt{\frac{1-cx}{1+cx}}(1+cx)(a + b\operatorname{sech}^{-1}(cx))}{9x} - \frac{(a + b\operatorname{sech}^{-1}(cx))^2}{3x^3}$$

output

```
-2/27*b^2/x^3-4/9*b^2*c^2/x+2/9*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+b*ar
csech(c*x))/x^3+4/9*b*c^2*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+b*arcsech(c*
x))/x-1/3*(a+b*arcsech(c*x))^2/x^3
```

**Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^4} dx$$

$$= \frac{-9a^2 - 2b^2(1 + 6c^2x^2) + 6ab\sqrt{\frac{1-cx}{1+cx}}(1 + cx + 2c^2x^2 + 2c^3x^3) + 6b\left(-3a + b\sqrt{\frac{1-cx}{1+cx}}(1 + cx + 2c^2x^2 + 2c^3x^3)\right)}{27x^3}$$

input

```
Integrate[(a + b*ArcSech[c*x])^2/x^4,x]
```

output

```
(-9*a^2 - 2*b^2*(1 + 6*c^2*x^2) + 6*a*b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x
+ 2*c^2*x^2 + 2*c^3*x^3) + 6*b*(-3*a + b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c
*x + 2*c^2*x^2 + 2*c^3*x^3))*ArcSech[c*x] - 9*b^2*ArcSech[c*x]^2)/(27*x^3)
```

**Rubi [A] (verified)**

Time = 0.51 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.11, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$ , Rules used = {6839, 5970, 3042, 3791, 3042, 3777, 26, 3042, 26, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^4} dx$$

$$\downarrow \text{6839}$$

$$-c^3 \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{c^3x^3} d \operatorname{sech}^{-1}(cx)$$

$$\downarrow \text{5970}$$

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{3c^3x^3} - \frac{2}{3}b \int \frac{a + b \operatorname{sech}^{-1}(cx)}{c^3x^3} d \operatorname{sech}^{-1}(cx) \right)$$

$$\downarrow \text{3042}$$

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{3c^3x^3} - \frac{2}{3}b \int (a + b \operatorname{sech}^{-1}(cx)) \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^3 d \operatorname{sech}^{-1}(cx) \right)$$

↓ 3791

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{3c^3x^3} - \frac{2}{3}b \left( \frac{2}{3} \int \frac{a + b \operatorname{sech}^{-1}(cx)}{cx} d \operatorname{sech}^{-1}(cx) + \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{3c^3x^3} - \frac{b}{9c^3x^3} \right) \right)$$

↓ 3042

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{3c^3x^3} - \frac{2}{3}b \left( \frac{2}{3} \int (a + b \operatorname{sech}^{-1}(cx)) \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right) d \operatorname{sech}^{-1}(cx) + \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{3c^3x^3} \right) \right)$$

↓ 3777

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{3c^3x^3} - \frac{2}{3}b \left( \frac{2}{3} \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{cx} - ib \int -\frac{i\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{cx} d \operatorname{sech}^{-1}(cx) \right) \right) \right)$$

↓ 26

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{3c^3x^3} - \frac{2}{3}b \left( \frac{2}{3} \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{cx} - b \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{cx} d \operatorname{sech}^{-1}(cx) \right) \right) \right)$$

↓ 3042

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{3c^3x^3} - \frac{2}{3}b \left( \frac{2}{3} \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{cx} - b \int -i \sin(i \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) \right) \right) \right)$$

↓ 26

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{3c^3x^3} - \frac{2}{3}b \left( \frac{2}{3} \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{cx} + ib \int \sin(i \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) \right) \right) \right)$$

↓ 3118



$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{3c^3 x^3} - \frac{2}{3} b \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{3c^3 x^3} + \frac{2}{3} \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{cx} - \dots \right) \right) \right)$$

input `Int[(a + b*ArcSech[c*x])^2/x^4,x]`

output `-(c^3*((a + b*ArcSech[c*x])^2/(3*c^3*x^3) - (2*b*(-1/9*b/(c^3*x^3) + (Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x]))/(3*c^3*x^3) + (2*(-(b/(c*x)) + (Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x]))/(c*x)))/3))/3))`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_)^(m_.))*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3791 `Int[((c_.) + (d_.)*(x_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[d*((b*Sine[e + f*x])^n/(f^2*n^2)), x] + (-Simp[b*(c + d*x)*Cos[e + f*x]*(b*Sine[e + f*x])^(n - 1)/(f*n), x] + Simp[b^2*((n - 1)/n) Int[(c + d*x)*(b*Sine[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1]`

rule 5970

```
Int[Cosh[(a_.) + (b_.)*(x_)^(n_.)*((c_.) + (d_.)*(x_)^(m_.)*Sinh[(a_.) +
(b_.)*(x_)], x_Symbol] := Simp[(c + d*x)^m*(Cosh[a + b*x]^(n + 1)/(b*(n + 1
))), x] - Simp[d*(m/(b*(n + 1))) Int[(c + d*x)^(m - 1)*Cosh[a + b*x]^(n +
1), x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

rule 6839

```
Int[((a_.) + ArcSech[(c_.)*(x_)*(b_.)]^(n_.)*(x_)^(m_.), x_Symbol] := Simp[
-(c^(m + 1))^(n-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

### Maple [A] (verified)

Time = 0.73 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.57

method	result
parts	$-\frac{a^2}{3x^3} + b^2c^3 \left( -\frac{\operatorname{arcsech}(cx)^2}{3c^3x^3} + \frac{4 \operatorname{arcsech}(cx)\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}}{9} + \frac{2\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}} \operatorname{arcsech}(cx)}{9c^2x^2} - \frac{4}{9cx} \right)$
derivativedivides	$c^3 \left( -\frac{a^2}{3c^3x^3} + b^2 \left( -\frac{\operatorname{arcsech}(cx)^2}{3c^3x^3} + \frac{4 \operatorname{arcsech}(cx)\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}}{9} + \frac{2\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}} \operatorname{arcsech}(cx)}{9c^2x^2} - \frac{4}{9cx} \right) \right)$
default	$c^3 \left( -\frac{a^2}{3c^3x^3} + b^2 \left( -\frac{\operatorname{arcsech}(cx)^2}{3c^3x^3} + \frac{4 \operatorname{arcsech}(cx)\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}}{9} + \frac{2\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}} \operatorname{arcsech}(cx)}{9c^2x^2} - \frac{4}{9cx} \right) \right)$

input

```
int((a+b*arcsech(c*x))^2/x^4,x,method=_RETURNVERBOSE)
```

output

```
-1/3*a^2/x^3+b^2*c^3*(-1/3/c^3/x^3*arcsech(c*x)^2+4/9*arcsech(c*x)*(-(c*x-
1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)+2/9*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(
1/2)/c^2/x^2*arcsech(c*x)-4/9/c/x-2/27/c^3/x^3)+2*a*b*c^3*(-1/3/c^3/x^3*ar
csech(c*x)+1/9*(-(c*x-1)/c/x)^(1/2)/c^2/x^2*((c*x+1)/c/x)^(1/2)*(2*c^2*x^2
+1))
```

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.48

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^4} dx = \frac{12 b^2 c^2 x^2 + 9 b^2 \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx}\right)^2 + 9 a^2 + 2 b^2 + 6 \left(3 ab - (2 b^2 c^3 x^3 + b^2 cx) \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}}\right) \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx}\right)}{27 x^3}$$

input `integrate((a+b*arcsech(c*x))^2/x^4,x, algorithm="fricas")`output `-1/27*(12*b^2*c^2*x^2 + 9*b^2*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x))^2 + 9*a^2 + 2*b^2 + 6*(3*a*b - (2*b^2*c^3*x^3 + b^2*c*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - 6*(2*a*b*c^3*x^3 + a*b*c*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/x^3`**Sympy [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^4} dx = \int \frac{(a + b \operatorname{asech}(cx))^2}{x^4} dx$$

input `integrate((a+b*asech(c*x))**2/x**4,x)`output `Integral((a + b*asech(c*x))**2/x**4, x)`

**Maxima [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^4} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2}{x^4} dx$$

input `integrate((a+b*arcsech(c*x))^2/x^4,x, algorithm="maxima")`

output `2/9*a*b*((c^4*(1/(c^2*x^2) - 1)^(3/2) + 3*c^4*sqrt(1/(c^2*x^2) - 1))/c - 3*arcsech(c*x)/x^3) + b^2*integrate(log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^2/x^4, x) - 1/3*a^2/x^3`

**Giac [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^4} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2}{x^4} dx$$

input `integrate((a+b*arcsech(c*x))^2/x^4,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)^2/x^4, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(\frac{1}{cx}))^2}{x^4} dx$$

input `int((a + b*acosh(1/(c*x)))^2/x^4,x)`

output `int((a + b*acosh(1/(c*x)))^2/x^4, x)`

**Reduce [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^4} dx = \frac{6 \left( \int \frac{\operatorname{asech}(cx)}{x^4} dx \right) ab x^3 + 3 \left( \int \frac{\operatorname{asech}(cx)^2}{x^4} dx \right) b^2 x^3 - a^2}{3x^3}$$

input `int((a+b*asech(c*x))^2/x^4,x)`

output `(6*int(asech(c*x)/x**4,x)*a*b*x**3 + 3*int(asech(c*x)**2/x**4,x)*b**2*x**3 - a**2)/(3*x**3)`

**3.41**  $\int \frac{(a+b\operatorname{sech}^{-1}(cx))^2}{x^5} dx$

Optimal result	373
Mathematica [A] (verified)	374
Rubi [A] (verified)	374
Maple [B] (verified)	377
Fricas [A] (verification not implemented)	377
Sympy [F]	378
Maxima [F]	378
Giac [F]	379
Mupad [F(-1)]	379
Reduce [F]	379

**Optimal result**

Integrand size = 14, antiderivative size = 139

$$\int \frac{(a + b\operatorname{sech}^{-1}(cx))^2}{x^5} dx = -\frac{b^2}{32x^4} - \frac{3b^2c^2}{32x^2} + \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(a + b\operatorname{sech}^{-1}(cx))}{8x^4}$$

$$+ \frac{3bc^2\sqrt{\frac{1-cx}{1+cx}}(1+cx)(a + b\operatorname{sech}^{-1}(cx))}{16x^2}$$

$$+ \frac{3}{32}c^4(a + b\operatorname{sech}^{-1}(cx))^2 - \frac{(a + b\operatorname{sech}^{-1}(cx))^2}{4x^4}$$

output

```
-1/32*b^2/x^4-3/32*b^2*c^2/x^2+1/8*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+b
*arcsech(c*x))/x^4+3/16*b*c^2*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+b*arcsec
h(c*x))/x^2+3/32*c^4*(a+b*arcsech(c*x))^2-1/4*(a+b*arcsech(c*x))^2/x^4
```

**Mathematica [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.93

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^5} dx$$

$$= \frac{-8a^2 - b^2 - 3b^2c^2x^2 + 4ab\sqrt{\frac{1-cx}{1+cx}} + 4abcx\sqrt{\frac{1-cx}{1+cx}} + 6abc^2x^2\sqrt{\frac{1-cx}{1+cx}} + 6abc^3x^3\sqrt{\frac{1-cx}{1+cx}} + 2b(-8a + b\sqrt{\frac{1-cx}{1+cx}})}{32x^4}$$

input

Integrate[(a + b\*ArcSech[c\*x])^2/x^5,x]

output

```
(-8*a^2 - b^2 - 3*b^2*c^2*x^2 + 4*a*b*Sqrt[(1 - c*x)/(1 + c*x)] + 4*a*b*c*x*Sqrt[(1 - c*x)/(1 + c*x)] + 6*a*b*c^2*x^2*Sqrt[(1 - c*x)/(1 + c*x)] + 6*a*b*c^3*x^3*Sqrt[(1 - c*x)/(1 + c*x)] + 2*b*(-8*a + b*Sqrt[(1 - c*x)/(1 + c*x)])*(2 + 2*c*x + 3*c^2*x^2 + 3*c^3*x^3))*ArcSech[c*x] + b^2*(-8 + 3*c^4*x^4)*ArcSech[c*x]^2 - 6*a*b*c^4*x^4*Log[x] + 6*a*b*c^4*x^4*Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]])/(32*x^4)
```

**Rubi [A] (verified)**Time = 0.46 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.14, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6839, 5970, 3042, 3791, 3042, 3791, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^5} dx$$

$$\downarrow \text{6839}$$

$$-c^4 \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{c^4 x^4} d \operatorname{sech}^{-1}(cx)$$

$$\downarrow \text{5970}$$

$$\begin{aligned}
 & -c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{4c^4 x^4} - \frac{1}{2} b \int \frac{a + b \operatorname{sech}^{-1}(cx)}{c^4 x^4} d \operatorname{sech}^{-1}(cx) \right) \\
 & \quad \downarrow \text{3042} \\
 & -c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{4c^4 x^4} - \frac{1}{2} b \int (a + b \operatorname{sech}^{-1}(cx)) \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^4 d \operatorname{sech}^{-1}(cx) \right) \\
 & \quad \downarrow \text{3791} \\
 & -c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{4c^4 x^4} - \frac{1}{2} b \left( \frac{3}{4} \int \frac{a + b \operatorname{sech}^{-1}(cx)}{c^2 x^2} d \operatorname{sech}^{-1}(cx) + \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{4c^4 x^4} - \frac{b}{16c^4 x^4} \right) \right) \\
 & \quad \downarrow \text{3042} \\
 & -c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{4c^4 x^4} - \frac{1}{2} b \left( \frac{3}{4} \int (a + b \operatorname{sech}^{-1}(cx)) \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^2 d \operatorname{sech}^{-1}(cx) + \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{4c^4 x^4} \right) \right) \\
 & \quad \downarrow \text{3791} \\
 & -c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{4c^4 x^4} - \frac{1}{2} b \left( \frac{3}{4} \left( \frac{1}{2} \int (a + b \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) + \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{2c^2 x^2} \right) \right) \right) \\
 & \quad \downarrow \text{17} \\
 & -c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{4c^4 x^4} - \frac{1}{2} b \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{4c^4 x^4} + \frac{3}{4} \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{2c^2 x^2} + \dots \right) \right) \right)
 \end{aligned}$$

input

```
Int[(a + b*ArcSech[c*x])^2/x^5,x]
```

output

```
-(c^4*((a + b*ArcSech[c*x])^2/(4*c^4*x^4) - (b*(-1/16*b/(c^4*x^4) + (Sqrt[
(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x]))/(4*c^4*x^4) + (3*(-1/
4*b/(c^2*x^2) + (Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x]))
/(2*c^2*x^2) + (a + b*ArcSech[c*x])^2/(4*b)))/4))/2))
```



## Definitions of rubi rules used

- rule 17  $\text{Int}[(c_.)*(a_.) + (b_.)*(x_)^{\wedge}(m_.), x\_Symbol] \rightarrow \text{Simp}[c*(a + b*x)^{\wedge}(m + 1)/(b*(m + 1)), x] /; \text{FreeQ}[\{a, b, c, m\}, x] \ \&\& \ \text{NeQ}[m, -1]$
- rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3791  $\text{Int}[((c_.) + (d_.)*(x_))*((b_.)*\sin[(e_.) + (f_.)*(x_)])^{\wedge}(n_), x\_Symbol] \rightarrow \text{Simp}[d*((b*\sin[e + f*x])^{\wedge}n/(f^{\wedge}2*n^{\wedge}2)), x] + (-\text{Simp}[b*(c + d*x)*\cos[e + f*x]*((b*\sin[e + f*x])^{\wedge}(n - 1)/(f*n)), x] + \text{Simp}[b^{\wedge}2*((n - 1)/n) \ \text{Int}[(c + d*x)*(b*\sin[e + f*x])^{\wedge}(n - 2), x], x]) /; \text{FreeQ}[\{b, c, d, e, f\}, x] \ \&\& \ \text{GtQ}[n, 1]$
- rule 5970  $\text{Int}[\text{Cosh}[(a_.) + (b_.)*(x_)]^{\wedge}(n_.)*((c_.) + (d_.)*(x_))^{\wedge}(m_.)*\text{Sinh}[(a_.) + (b_.)*(x_)], x\_Symbol] \rightarrow \text{Simp}[(c + d*x)^{\wedge}m*(\text{Cosh}[a + b*x]^{\wedge}(n + 1)/(b*(n + 1))), x] - \text{Simp}[d*(m/(b*(n + 1))) \ \text{Int}[(c + d*x)^{\wedge}(m - 1)*\text{Cosh}[a + b*x]^{\wedge}(n + 1), x], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{NeQ}[n, -1]$
- rule 6839  $\text{Int}[(a_.) + \text{ArcSech}[(c_.)*(x_)]*(b_.))^{\wedge}(n_)*(x_)^{\wedge}(m_.), x\_Symbol] \rightarrow \text{Simp}[-(c^{\wedge}(m + 1))^{\wedge}(-1) \ \text{Subst}[\text{Int}[(a + b*x)^{\wedge}n*\text{sech}[x]^{\wedge}(m + 1)*\text{Tanh}[x], x], x, \text{ArcSech}[c*x]], x] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ (\text{GtQ}[n, 0] \ || \ \text{LtQ}[m, -1])$

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 262 vs. 2(123) = 246.

Time = 0.66 (sec) , antiderivative size = 263, normalized size of antiderivative = 1.89

method	result
parts	$-\frac{a^2}{4x^4} + b^2c^4 \left( -\frac{\operatorname{arcsech}(cx)^2}{4c^4x^4} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}\operatorname{arcsech}(cx)}{8c^3x^3} + \frac{3\operatorname{arcsech}(cx)\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}}{16cx} + \frac{3\operatorname{arcsech}(cx)}{32c^2x^2} \right)$
derivativedivides	$c^4 \left( -\frac{a^2}{4c^4x^4} + b^2 \left( -\frac{\operatorname{arcsech}(cx)^2}{4c^4x^4} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}\operatorname{arcsech}(cx)}{8c^3x^3} + \frac{3\operatorname{arcsech}(cx)\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}}{16cx} + \frac{3\operatorname{arcsech}(cx)}{32c^2x^2} \right) \right)$
default	$c^4 \left( -\frac{a^2}{4c^4x^4} + b^2 \left( -\frac{\operatorname{arcsech}(cx)^2}{4c^4x^4} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}\operatorname{arcsech}(cx)}{8c^3x^3} + \frac{3\operatorname{arcsech}(cx)\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}}{16cx} + \frac{3\operatorname{arcsech}(cx)}{32c^2x^2} \right) \right)$

input `int((a+b*arcsech(c*x))^2/x^5,x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -1/4*a^2/x^4 + b^2*c^4*(-1/4/c^4/x^4*arcsech(c*x)^2 + 1/8*(-(c*x-1)/c/x)^(1/2) \\ & *((c*x+1)/c/x)^(1/2)/c^3/x^3*arcsech(c*x) + 3/16*arcsech(c*x)/c/x*(-(c*x-1)/ \\ & c/x)^(1/2)*((c*x+1)/c/x)^(1/2) + 3/32*arcsech(c*x)^2 - 1/32/c^4/x^4 - 3/32/c^2/x \\ & ^2) + 2*a*b*c^4*(-1/4/c^4/x^4*arcsech(c*x) + 1/32*(-(c*x-1)/c/x)^(1/2)/c^3/x^3 \\ & *((c*x+1)/c/x)^(1/2)*(3*arctanh(1/(-c^2*x^2+1)^(1/2))*c^4*x^4 + 3*(-c^2*x^2+ \\ & 1)^(1/2)*c^2*x^2 + 2*(-c^2*x^2+1)^(1/2))/(-c^2*x^2+1)^(1/2)) \end{aligned}$$

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.47

$$\int \frac{(a + b\operatorname{sech}^{-1}(cx))^2}{x^5} dx = \frac{3b^2c^2x^2 - (3b^2c^4x^4 - 8b^2)\log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}+1}}{cx}\right)^2 + 8a^2 + b^2 - 2\left(3abc^4x^4 - 8ab + (3b^2c^3x^3 + 2b^2cx^2)\right)}{32x^4}$$

input `integrate((a+b*arcsech(c*x))^2/x^5,x, algorithm="fricas")`

output

```
-1/32*(3*b^2*c^2*x^2 - (3*b^2*c^4*x^4 - 8*b^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x))^2 + 8*a^2 + b^2 - 2*(3*a*b*c^4*x^4 - 8*a*b + (3*b^2*c^3*x^3 + 2*b^2*c*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - 2*(3*a*b*c^3*x^3 + 2*a*b*c*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/x^4
```

**Sympy [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^5} dx = \int \frac{(a + b \operatorname{asech}(cx))^2}{x^5} dx$$

input

```
integrate((a+b*asech(c*x))**2/x**5,x)
```

output

```
Integral((a + b*asech(c*x))**2/x**5, x)
```

**Maxima [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^5} dx = \int \frac{(b \operatorname{arsech}(cx) + a)^2}{x^5} dx$$

input

```
integrate((a+b*arcsech(c*x))^2/x^5,x, algorithm="maxima")
```

output

```
1/32*a*b*((3*c^5*log(c*x*sqrt(1/(c^2*x^2) - 1) + 1) - 3*c^5*log(c*x*sqrt(1/(c^2*x^2) - 1) - 1) - 2*(3*c^8*x^3*(1/(c^2*x^2) - 1)^(3/2) - 5*c^6*x*sqrt(1/(c^2*x^2) - 1))/(c^4*x^4*(1/(c^2*x^2) - 1)^2 - 2*c^2*x^2*(1/(c^2*x^2) - 1) + 1))/c - 16*arcsech(c*x)/x^4) + b^2*integrate(log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^2/x^5, x) - 1/4*a^2/x^4
```

**Giac [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^5} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2}{x^5} dx$$

input `integrate((a+b*arcsech(c*x))^2/x^5,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)^2/x^5, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^5} dx = \int \frac{(a + b \operatorname{acosh}(\frac{1}{cx}))^2}{x^5} dx$$

input `int((a + b*acosh(1/(c*x)))^2/x^5,x)`

output `int((a + b*acosh(1/(c*x)))^2/x^5, x)`

**Reduce [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{x^5} dx = \frac{8 \left( \int \frac{\operatorname{asech}(cx)}{x^5} dx \right) ab x^4 + 4 \left( \int \frac{\operatorname{asech}(cx)^2}{x^5} dx \right) b^2 x^4 - a^2}{4x^4}$$

input `int((a+b*asech(c*x))^2/x^5,x)`

output `(8*int(asech(c*x)/x**5,x)*a*b*x**4 + 4*int(asech(c*x)**2/x**5,x)*b**2*x**4 - a**2)/(4*x**4)`

### 3.42 $\int x^3 (a + b \operatorname{sech}^{-1}(cx))^3 dx$

Optimal result	380
Mathematica [A] (verified)	381
Rubi [C] (verified)	382
Maple [A] (verified)	386
Fricas [F]	387
Sympy [F]	387
Maxima [F]	388
Giac [F]	388
Mupad [F(-1)]	388
Reduce [F]	389

#### Optimal result

Integrand size = 14, antiderivative size = 223

$$\begin{aligned}
 \int x^3 (a + b \operatorname{sech}^{-1}(cx))^3 dx = & \frac{b^3 \sqrt{\frac{1-cx}{1+cx}} (1+cx)}{4c^4} - \frac{b^2 x^2 (a + b \operatorname{sech}^{-1}(cx))}{4c^2} \\
 & - \frac{b(a + b \operatorname{sech}^{-1}(cx))^2}{2c^4} \\
 & - \frac{b \sqrt{\frac{1-cx}{1+cx}} (1+cx) (a + b \operatorname{sech}^{-1}(cx))^2}{2c^4} \\
 & - \frac{bx^2 \sqrt{\frac{1-cx}{1+cx}} (1+cx) (a + b \operatorname{sech}^{-1}(cx))^2}{4c^2} \\
 & + \frac{1}{4} x^4 (a + b \operatorname{sech}^{-1}(cx))^3 \\
 & + \frac{b^2 (a + b \operatorname{sech}^{-1}(cx)) \log(1 + e^{2 \operatorname{sech}^{-1}(cx)})}{c^4} \\
 & + \frac{b^3 \operatorname{PolyLog}(2, -e^{2 \operatorname{sech}^{-1}(cx)})}{2c^4}
 \end{aligned}$$

output

```
1/4*b^3*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/c^4-1/4*b^2*x^2*(a+b*arcsech(c*x)
)/c^2-1/2*b*(a+b*arcsech(c*x))^2/c^4-1/2*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1
)*(a+b*arcsech(c*x))^2/c^4-1/4*b*x^2*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+b
*arcsech(c*x))^2/c^2+1/4*x^4*(a+b*arcsech(c*x))^3+b^2*(a+b*arcsech(c*x))*l
n(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)/c^4+1/2*b^3*polylog(2,-(1/
c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)/c^4
```

**Mathematica [A] (verified)**

Time = 1.35 (sec) , antiderivative size = 337, normalized size of antiderivative = 1.51

$$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^3 dx$$

$$= \frac{1}{4} \left( a^3 x^4 + b^3 x^4 \operatorname{sech}^{-1}(cx)^3 + a^2 b \left( -\frac{\sqrt{\frac{1-cx}{1+cx}}(1+cx)(2+c^2x^2)}{c^4} + 3x^4 \operatorname{sech}^{-1}(cx) \right) \right. \\ \left. + \frac{ab^2 \left( -c^2x^2 - 2\sqrt{\frac{1-cx}{1+cx}}(2+2cx+c^2x^2+c^3x^3) \operatorname{sech}^{-1}(cx) + 3c^4x^4 \operatorname{sech}^{-1}(cx)^2 + 4 \log\left(\frac{1}{cx}\right) \right)}{c^4} \right. \\ \left. - \frac{b^3 \left( -\sqrt{\frac{1-cx}{1+cx}}(1+cx) + \left( -2 + 2\sqrt{\frac{1-cx}{1+cx}} + 2cx\sqrt{\frac{1-cx}{1+cx}} + c^2x^2\sqrt{\frac{1-cx}{1+cx}} + c^3x^3\sqrt{\frac{1-cx}{1+cx}} \right) \operatorname{sech}^{-1}(cx)^2 + \operatorname{sech}^{-1}(cx) \right)}{c^4} \right)$$

input

```
Integrate[x^3*(a + b*ArcSech[c*x])^3,x]
```

output

```
(a^3*x^4 + b^3*x^4*ArcSech[c*x]^3 + a^2*b*(-((Sqrt[(1 - c*x)/(1 + c*x)]*(1
+ c*x)*(2 + c^2*x^2))/c^4) + 3*x^4*ArcSech[c*x]) + (a*b^2*(-(c^2*x^2) - 2
*Sqrt[(1 - c*x)/(1 + c*x)]*(2 + 2*c*x + c^2*x^2 + c^3*x^3)*ArcSech[c*x] +
3*c^4*x^4*ArcSech[c*x]^2 + 4*Log[1/(c*x)]))/c^4 - (b^3*(-(Sqrt[(1 - c*x)/(
1 + c*x)]*(1 + c*x)) + (-2 + 2*Sqrt[(1 - c*x)/(1 + c*x)] + 2*c*x*Sqrt[(1 -
c*x)/(1 + c*x)] + c^2*x^2*Sqrt[(1 - c*x)/(1 + c*x)] + c^3*x^3*Sqrt[(1 - c
*x)/(1 + c*x)])*ArcSech[c*x]^2 + ArcSech[c*x]*(c^2*x^2 - 4*Log[1 + E^(-2*A
rcSech[c*x])])) + 2*PolyLog[2, -E^(-2*ArcSech[c*x])])))/c^4)/4
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.93 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.04, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.071$ , Rules used = {6839, 5974, 3042, 4674, 3042, 4254, 24, 4672, 26, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 (a + b \operatorname{sech}^{-1}(cx))^3 dx \\
 & \quad \downarrow \text{6839} \\
 & - \frac{\int c^4 x^4 \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx))^3 d \operatorname{sech}^{-1}(cx)}{c^4} \\
 & \quad \downarrow \text{5974} \\
 & - \frac{\frac{3}{4} b \int c^4 x^4 (a + b \operatorname{sech}^{-1}(cx))^2 d \operatorname{sech}^{-1}(cx) - \frac{1}{4} c^4 x^4 (a + b \operatorname{sech}^{-1}(cx))^3}{c^4} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{-\frac{1}{4} c^4 x^4 (a + b \operatorname{sech}^{-1}(cx))^3 + \frac{3}{4} b \int (a + b \operatorname{sech}^{-1}(cx))^2 \csc \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^4 d \operatorname{sech}^{-1}(cx)}{c^4} \\
 & \quad \downarrow \text{4674} \\
 & - \frac{\frac{3}{4} b \left( \frac{2}{3} \int c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^2 d \operatorname{sech}^{-1}(cx) - \frac{1}{3} b^2 \int c^2 x^2 d \operatorname{sech}^{-1}(cx) + \frac{1}{3} b c^2 x^2 (a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{3} c^2 x^2 \sqrt{\frac{1-cx}{cx}} \right)}{c^4} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{-\frac{1}{4} c^4 x^4 (a + b \operatorname{sech}^{-1}(cx))^3 + \frac{3}{4} b \left( \frac{2}{3} \int (a + b \operatorname{sech}^{-1}(cx))^2 \csc \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^2 d \operatorname{sech}^{-1}(cx) - \frac{1}{3} b^2 \int \csc \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right) d \operatorname{sech}^{-1}(cx) \right)}{c^4} \\
 & \quad \downarrow \text{4254} \\
 & - \frac{-\frac{1}{4} c^4 x^4 (a + b \operatorname{sech}^{-1}(cx))^3 + \frac{3}{4} b \left( \frac{2}{3} \int (a + b \operatorname{sech}^{-1}(cx))^2 \csc \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^2 d \operatorname{sech}^{-1}(cx) - \frac{1}{3} b^2 \int 1 d \left( -i \sqrt{\frac{1-cx}{cx}} \right) \right)}{c^4}
 \end{aligned}$$

---

↓ 24

$$-\frac{1}{4}c^4x^4(a + b\operatorname{sech}^{-1}(cx))^3 + \frac{3}{4}b\left(\frac{2}{3}\int(a + b\operatorname{sech}^{-1}(cx))^2 \csc\left(\operatorname{isech}^{-1}(cx) + \frac{\pi}{2}\right)^2 d\operatorname{sech}^{-1}(cx) + \frac{1}{3}bc^2x^2(a + b\operatorname{sech}^{-1}(cx))\right)$$


---

↓ 4672

$$-\frac{1}{4}c^4x^4(a + b\operatorname{sech}^{-1}(cx))^3 + \frac{3}{4}b\left(\frac{2}{3}\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))^2 - 2ib\int -i\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))\right)\right)$$


---

↓ 26

$$\frac{3}{4}b\left(\frac{2}{3}\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))^2 - 2b\int\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))d\operatorname{sech}^{-1}(cx)\right) + \frac{1}{3}bc^2x^2(a + b\operatorname{sech}^{-1}(cx))\right)$$


---

↓ 3042

$$-\frac{1}{4}c^4x^4(a + b\operatorname{sech}^{-1}(cx))^3 + \frac{3}{4}b\left(\frac{2}{3}\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))^2 - 2b\int -i(a + b\operatorname{sech}^{-1}(cx))\tan(\operatorname{isech}^{-1}(cx))\right)\right)$$


---

↓ 26

$$-\frac{1}{4}c^4x^4(a + b\operatorname{sech}^{-1}(cx))^3 + \frac{3}{4}b\left(\frac{2}{3}\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))^2 + 2ib\int(a + b\operatorname{sech}^{-1}(cx))\tan(\operatorname{isech}^{-1}(cx))\right)\right)$$


---

↓ 4201

$$-\frac{1}{4}c^4x^4(a + b\operatorname{sech}^{-1}(cx))^3 + \frac{3}{4}b\left(\frac{2}{3}\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))^2 + 2ib\left(2i\int\frac{e^{2\operatorname{sech}^{-1}(cx)}(a + b\operatorname{sech}^{-1}(cx))}{1 + e^{2\operatorname{sech}^{-1}(cx)}}\right)\right)\right)$$


---

↓ 2620

$$-\frac{1}{4}c^4x^4(a + b\operatorname{sech}^{-1}(cx))^3 + \frac{3}{4}b\left(\frac{2}{3}\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))^2 + 2ib\left(2i\left(\frac{1}{2}\log(e^{2\operatorname{sech}^{-1}(cx)} + 1)\right)(a + b\operatorname{sech}^{-1}(cx))\right)\right)\right)$$


---

↓ 2715



$$-\frac{1}{4}c^4x^4(a + b\operatorname{sech}^{-1}(cx))^3 + \frac{3}{4}b\left(\frac{2}{3}\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))^2 + 2ib\left(2i\left(\frac{1}{2}\log\left(e^{2\operatorname{sech}^{-1}(cx)} + 1\right)\right)(a + b\operatorname{sech}^{-1}(cx))\right)\right)\right)$$

↓ 2838

$$-\frac{1}{4}c^4x^4(a + b\operatorname{sech}^{-1}(cx))^3 + \frac{3}{4}b\left(\frac{1}{3}bc^2x^2(a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{3}c^2x^2\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))^2 + \frac{2}{3}\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))^2 + 2ib\left(2i\left(\frac{1}{2}\log\left(e^{2\operatorname{sech}^{-1}(cx)} + 1\right)\right)(a + b\operatorname{sech}^{-1}(cx))\right)\right)\right)$$

input `Int[x^3*(a + b*ArcSech[c*x])^3,x]`

output `-((-1/4*(c^4*x^4*(a + b*ArcSech[c*x])^3) + (3*b*(-1/3*(b^2*sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)) + (b*c^2*x^2*(a + b*ArcSech[c*x])))/3 + (c^2*x^2*sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x])^2)/3 + (2*(sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x])^2 + (2*I)*b*((-1/2*I)*(a + b*ArcSech[c*x])^2)/b + (2*I)*(((a + b*ArcSech[c*x])*Log[1 + E^(2*ArcSech[c*x])])))/2 + (b*PolyLog[2, -E^(2*ArcSech[c*x])])/4))))/3)/4)/c^4`

### Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715  $\text{Int}[\text{Log}[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^{(n_)}], x\_Symbol]$   
 $\text{:> Simp}[1/(d*e*n*\text{Log}[F]) \text{ Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^{(e*(c + d*x)})^n], x] /;$   $\text{FreeQ}\{F, a, b, c, d, e, n\}, x\} \ \&\& \ \text{GtQ}\{a, 0\}$

rule 2838  $\text{Int}[\text{Log}[(c_)*((d_) + (e_)*(x_)^{(n_)})]/(x_), x\_Symbol] \text{:> Simp}[-\text{PolyLog}[2,$   
 $(-c)*e*x^n/n, x] /;$   $\text{FreeQ}\{c, d, e, n\}, x\} \ \&\& \ \text{EqQ}\{c*d, 1\}$

rule 3042  $\text{Int}[u_, x\_Symbol] \text{:> Int}[\text{DeactivateTrig}[u, x], x] /;$   $\text{FunctionOfTrigOfLinear}$   
 $\text{Q}[u, x]$

rule 4201  $\text{Int}[(c_) + (d_)*(x_)^{(m_)}*\text{tan}[(e_) + (\text{Complex}[0, fz_])*(f_)*(x_)], x$   
 $\_Symbol] \text{:> Simp}[(-I)*((c + d*x)^{(m + 1)}/(d*(m + 1))), x] + \text{Simp}[2*I \text{ Int}[(c + d*x)^m*(E^{(2*(-I)*e + f*fz*x})/(1 + E^{(2*(-I)*e + f*fz*x}))], x], x]$   
 $/;$   $\text{FreeQ}\{c, d, e, f, fz\}, x\} \ \&\& \ \text{IGtQ}\{m, 0\}$

rule 4254  $\text{Int}[\text{csc}[(c_) + (d_)*(x_)]^{(n_)}, x\_Symbol] \text{:> Simp}[-d^{(-1)} \text{ Subst}[\text{Int}[\text{Exp}$   
 $\text{andIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \text{Cot}[c + d*x], x] /;$   $\text{FreeQ}\{c,$   
 $d\}, x\} \ \&\& \ \text{IGtQ}\{n/2, 0\}$

rule 4672  $\text{Int}[\text{csc}[(e_) + (f_)*(x_)]^2*((c_) + (d_)*(x_))^{(m_)}, x\_Symbol] \text{:> Simp}$   
 $[(-c + d*x)^m*(\text{Cot}[e + f*x]/f), x] + \text{Simp}[d*(m/f) \text{ Int}[(c + d*x)^{(m - 1)}$   
 $*\text{Cot}[e + f*x], x], x] /;$   $\text{FreeQ}\{c, d, e, f\}, x\} \ \&\& \ \text{GtQ}\{m, 0\}$

rule 4674  $\text{Int}[(\text{csc}[(e_) + (f_)*(x_)]*(b_))^{(n_)}*((c_) + (d_)*(x_))^{(m_)}, x\_Symbo$   
 $l] \text{:> Simp}[(-b^2)*(c + d*x)^m*\text{Cot}[e + f*x]*((b*\text{Csc}[e + f*x])^{(n - 2)})/(f*(n$   
 $- 1))), x] + (-\text{Simp}[b^2*d*m*(c + d*x)^{(m - 1)}*((b*\text{Csc}[e + f*x])^{(n - 2)})/(f^$   
 $2*(n - 1)*(n - 2))), x] + \text{Simp}[b^2*d^2*m*((m - 1)/(f^2*(n - 1)*(n - 2)))$   
 $\text{Int}[(c + d*x)^{(m - 2)}*(b*\text{Csc}[e + f*x])^{(n - 2)}, x], x] + \text{Simp}[b^2*((n - 2)/$   
 $(n - 1)) \text{ Int}[(c + d*x)^m*(b*\text{Csc}[e + f*x])^{(n - 2)}, x], x]) /;$   $\text{FreeQ}\{b, c,$   
 $d, e, f\}, x\} \ \&\& \ \text{GtQ}\{n, 1\} \ \&\& \ \text{NeQ}\{n, 2\} \ \&\& \ \text{GtQ}\{m, 1\}$

rule 5974

```
Int[((c_.) + (d_.)*(x_)^(m_.)*Sech[(a_.) + (b_.)*(x_)^(n_.)*Tanh[(a_.) +
(b_.)*(x_)^(p_.)], x_Symbol] := Simp[(-c + d*x)^m*(Sech[a + b*x]^n/(b*n))
, x] + Simp[d*(m/(b*n)) Int[(c + d*x)^(m - 1)*Sech[a + b*x]^n, x], x] /;
FreeQ[{a, b, c, d, n}, x] && EqQ[p, 1] && GtQ[m, 0]
```

rule 6839

```
Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.)^(n_)*(x_)^(m_.), x_Symbol] := Simp[
-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

### Maple [A] (verified)

Time = 0.84 (sec) , antiderivative size = 464, normalized size of antiderivative = 2.08

method	result
derivativedivides	$\frac{a^3 c^4 x^4}{4} + b^3 \left( \frac{\operatorname{arcsech}(cx)^3 c^4 x^4}{4} - \frac{c^3 x^3 \operatorname{arcsech}(cx)^2 \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}}{4} - \frac{\operatorname{arcsech}(cx)^2 \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} cx}{2} - \frac{c^2 x^2 \operatorname{arcsech}(cx)}{4} + \dots \right)$
default	$\frac{a^3 c^4 x^4}{4} + b^3 \left( \frac{\operatorname{arcsech}(cx)^3 c^4 x^4}{4} - \frac{c^3 x^3 \operatorname{arcsech}(cx)^2 \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}}{4} - \frac{\operatorname{arcsech}(cx)^2 \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} cx}{2} - \frac{c^2 x^2 \operatorname{arcsech}(cx)}{4} + \dots \right)$
parts	$\frac{a^3 x^4}{4} + b^3 \left( \frac{\operatorname{arcsech}(cx)^3 c^4 x^4}{4} - \frac{c^3 x^3 \operatorname{arcsech}(cx)^2 \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}}{4} - \frac{\operatorname{arcsech}(cx)^2 \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} cx}{2} - \frac{c^2 x^2 \operatorname{arcsech}(cx)}{4} + \dots \right)$

input

```
int(x^3*(a+b*arcsech(c*x))^3,x,method=_RETURNVERBOSE)
```

output

```
1/c^4*(1/4*a^3*c^4*x^4+b^3*(1/4*arcsech(c*x)^3*c^4*x^4-1/4*c^3*x^3*arcsech
(c*x)^2*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)-1/2*arcsech(c*x)^2*((c*x+
1)/c/x)^(1/2)*(-(c*x-1)/c/x)^(1/2)*c*x-1/4*c^2*x^2*arcsech(c*x)+1/4*(-(c*x
-1)/c/x)^(1/2)*c*((c*x+1)/c/x)^(1/2)*x-1/2*arcsech(c*x)^2-1/4+arcsech(c*x)
*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)+1/2*polylog(2,-(1/c/x+(-
1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2))+3*a*b^2*(-1/3*arcsech(c*x)+1/4*arcsech
(c*x)^2*c^4*x^4-1/6*arcsech(c*x)*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)*
c^3*x^3-1/3*arcsech(c*x)*((c*x+1)/c/x)^(1/2)*(-(c*x-1)/c/x)^(1/2)*c*x-1/12
*c^2*x^2+1/3*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2))+3*a^2*b*(1/
4*c^4*x^4*arcsech(c*x)-1/12*(-(c*x-1)/c/x)^(1/2)*c*x*((c*x+1)/c/x)^(1/2)*
(c^2*x^2+2))
```

**Fricas [F]**

$$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^3 x^3 dx$$

input

```
integrate(x^3*(a+b*arcsech(c*x))^3,x, algorithm="fricas")
```

output

```
integral(b^3*x^3*arcsech(c*x)^3 + 3*a*b^2*x^3*arcsech(c*x)^2 + 3*a^2*b*x^3
*arcsech(c*x) + a^3*x^3, x)
```

**Sympy [F]**

$$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int x^3 (a + b \operatorname{ar} \operatorname{sech}(cx))^3 dx$$

input

```
integrate(x**3*(a+b*asech(c*x))**3,x)
```

output

```
Integral(x**3*(a + b*asech(c*x))**3, x)
```

**Maxima [F]**

$$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^3 x^3 dx$$

input `integrate(x^3*(a+b*arcsech(c*x))^3,x, algorithm="maxima")`

output `1/4*a^3*x^4 + 1/4*(3*x^4*arcsech(c*x) + (c^2*x^3*(1/(c^2*x^2) - 1)^(3/2) - 3*x*sqrt(1/(c^2*x^2) - 1))/c^3)*a^2*b + integrate(b^3*x^3*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^3 + 3*a*b^2*x^3*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^2, x)`

**Giac [F]**

$$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^3 x^3 dx$$

input `integrate(x^3*(a+b*arcsech(c*x))^3,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)^3*x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int x^3 \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right)^3 dx$$

input `int(x^3*(a + b*acosh(1/(c*x)))^3,x)`

output `int(x^3*(a + b*acosh(1/(c*x)))^3, x)`

**Reduce [F]**

$$\int x^3 (a + b \operatorname{sech}^{-1}(cx))^3 dx = 3 \left( \int a \operatorname{sech}(cx) x^3 dx \right) a^2 b + \left( \int a \operatorname{sech}(cx)^3 x^3 dx \right) b^3 \\ + 3 \left( \int a \operatorname{sech}(cx)^2 x^3 dx \right) a b^2 + \frac{a^3 x^4}{4}$$

input `int(x^3*(a+b*asech(c*x))^3,x)`

output `(12*int(asech(c*x)*x**3,x)*a**2*b + 4*int(asech(c*x)**3*x**3,x)*b**3 + 12*int(asech(c*x)**2*x**3,x)*a*b**2 + a**3*x**4)/4`

### 3.43 $\int x^2 (a + b \operatorname{sech}^{-1}(cx))^3 dx$

Optimal result	390
Mathematica [A] (verified)	391
Rubi [A] (verified)	392
Maple [F]	395
Fricas [F]	395
Sympy [F]	396
Maxima [F]	396
Giac [F]	396
Mupad [F(-1)]	397
Reduce [F]	397

#### Optimal result

Integrand size = 14, antiderivative size = 242

$$\begin{aligned}
 \int x^2 (a + b \operatorname{sech}^{-1}(cx))^3 dx = & -\frac{b^2 x (a + b \operatorname{sech}^{-1}(cx))}{c^2} \\
 & - \frac{bx \sqrt{\frac{1-cx}{1+cx}} (1+cx) (a + b \operatorname{sech}^{-1}(cx))^2}{2c^2} \\
 & + \frac{1}{3} x^3 (a + b \operatorname{sech}^{-1}(cx))^3 \\
 & - \frac{b (a + b \operatorname{sech}^{-1}(cx))^2 \arctan(e^{\operatorname{sech}^{-1}(cx)})}{c^3} \\
 & + \frac{b^3 \arctan\left(\frac{\sqrt{\frac{1-cx}{1+cx}} (1+cx)}{cx}\right)}{c^3} \\
 & + \frac{ib^2 (a + b \operatorname{sech}^{-1}(cx)) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(cx)}\right)}{c^3} \\
 & - \frac{ib^2 (a + b \operatorname{sech}^{-1}(cx)) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(cx)}\right)}{c^3} \\
 & - \frac{ib^3 \operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(cx)}\right)}{c^3} \\
 & + \frac{ib^3 \operatorname{PolyLog}\left(3, ie^{\operatorname{sech}^{-1}(cx)}\right)}{c^3}
 \end{aligned}$$

output

```
-b^2*x*(a+b*arcsech(c*x))/c^2-1/2*b*x*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+
b*arcsech(c*x))^2/c^2+1/3*x^3*(a+b*arcsech(c*x))^3-b*(a+b*arcsech(c*x))^2*b
arctan(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/c^3+b^3*arctan(((c*x+1)/(c
*x+1))^(1/2)*(c*x+1)/c/x)/c^3+I*b^2*(a+b*arcsech(c*x))*polylog(2,-I*(1/c/x
+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/c^3-I*b^2*(a+b*arcsech(c*x))*polylog(2
,I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/c^3-I*b^3*polylog(3,-I*(1/c/x
+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/c^3+I*b^3*polylog(3,I*(1/c/x+(-1+1/c/x
)^(1/2)*(1+1/c/x)^(1/2)))/c^3
```

**Mathematica [A] (verified)**

Time = 0.76 (sec) , antiderivative size = 440, normalized size of antiderivative = 1.82

$$\int x^2 (a + b \operatorname{sech}^{-1}(cx))^3 dx$$

$$= \frac{2a^3 c^3 x^3 - 3a^2 b c x \sqrt{\frac{1-cx}{1+cx}} (1+cx) + 6a^2 b c^3 x^3 \operatorname{sech}^{-1}(cx) + 3ia^2 b \log\left(-2icx + 2\sqrt{\frac{1-cx}{1+cx}}(1+cx)\right) - 6ab^2 \left(\dots\right)}{\dots}$$

input

```
Integrate[x^2*(a + b*ArcSech[c*x])^3,x]
```

output

```
(2*a^3*c^3*x^3 - 3*a^2*b*c*x*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x) + 6*a^2*b
*c^3*x^3*ArcSech[c*x] + (3*I)*a^2*b*Log[(-2*I)*c*x + 2*Sqrt[(1 - c*x)/(1 +
c*x)]*(1 + c*x)] - 6*a*b^2*(c*x + c*x*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)
*ArcSech[c*x] - c^3*x^3*ArcSech[c*x]^2 - I*ArcSech[c*x]*Log[1 - I/E^ArcSec
h[c*x]] + I*ArcSech[c*x]*Log[1 + I/E^ArcSech[c*x]] - I*PolyLog[2, (-I)/E^A
rcSech[c*x]] + I*PolyLog[2, I/E^ArcSech[c*x]]) - b^3*(6*c*x*ArcSech[c*x] +
3*c*x*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*ArcSech[c*x]^2 - 2*c^3*x^3*ArcS
ech[c*x]^3 - (3*I)*((-4*I)*ArcTan[Tanh[ArcSech[c*x]/2]] + ArcSech[c*x]^2*L
og[1 - I/E^ArcSech[c*x]] - ArcSech[c*x]^2*Log[1 + I/E^ArcSech[c*x]] + 2*Ar
cSech[c*x]*PolyLog[2, (-I)/E^ArcSech[c*x]] - 2*ArcSech[c*x]*PolyLog[2, I/E
^ArcSech[c*x]] + 2*PolyLog[3, (-I)/E^ArcSech[c*x]] - 2*PolyLog[3, I/E^ArcS
ech[c*x]])))/(6*c^3)
```



**Rubi [A] (verified)**

Time = 0.87 (sec) , antiderivative size = 224, normalized size of antiderivative = 0.93, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$ , Rules used = {6839, 5974, 3042, 4674, 3042, 4257, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 (a + b \operatorname{sech}^{-1}(cx))^3 dx \\
 & \quad \downarrow \text{6839} \\
 & \frac{\int c^3 x^3 \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx))^3 d \operatorname{sech}^{-1}(cx)}{c^3} \\
 & \quad \downarrow \text{5974} \\
 & \frac{b \int c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))^2 d \operatorname{sech}^{-1}(cx) - \frac{1}{3} c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))^3}{c^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\frac{1}{3} c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))^3 + b \int (a + b \operatorname{sech}^{-1}(cx))^2 \csc(\operatorname{isech}^{-1}(cx) + \frac{\pi}{2})^3 d \operatorname{sech}^{-1}(cx)}{c^3} \\
 & \quad \downarrow \text{4674} \\
 & \frac{b \left( \frac{1}{2} \int cx (a + b \operatorname{sech}^{-1}(cx))^2 d \operatorname{sech}^{-1}(cx) + b^2 \left( - \int cx d \operatorname{sech}^{-1}(cx) \right) + bcx (a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{2} cx \sqrt{\frac{1-cx}{cx+1}} (cx + \right)}{c^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\frac{1}{3} c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))^3 + b \left( \frac{1}{2} \int (a + b \operatorname{sech}^{-1}(cx))^2 \csc(\operatorname{isech}^{-1}(cx) + \frac{\pi}{2}) d \operatorname{sech}^{-1}(cx) + b^2 \left( - \int \csc(\operatorname{isech}^{-1}(cx) + \frac{\pi}{2}) d \operatorname{sech}^{-1}(cx) \right) + bcx (a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{2} cx \sqrt{\frac{1-cx}{cx+1}} (cx + \right)}{c^3} \\
 & \quad \downarrow \text{4257} \\
 & \frac{-\frac{1}{3} c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))^3 + b \left( \frac{1}{2} \int (a + b \operatorname{sech}^{-1}(cx))^2 \csc(\operatorname{isech}^{-1}(cx) + \frac{\pi}{2}) d \operatorname{sech}^{-1}(cx) + bcx (a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{2} cx \sqrt{\frac{1-cx}{cx+1}} (cx + \right)}{c^3} \\
 & \quad \downarrow \text{4668}
 \end{aligned}$$

$$-\frac{1}{3}c^3x^3(a + b\operatorname{sech}^{-1}(cx))^3 + b\left(\frac{1}{2}\left(-2ib \int (a + b\operatorname{sech}^{-1}(cx)) \log(1 - ie^{\operatorname{sech}^{-1}(cx)}) d\operatorname{sech}^{-1}(cx) + 2ib \int (a + b\operatorname{sech}^{-1}(cx)) \log(1 + ie^{\operatorname{sech}^{-1}(cx)}) d\operatorname{sech}^{-1}(cx)\right)\right)$$

↓ 3011

$$-\frac{1}{3}c^3x^3(a + b\operatorname{sech}^{-1}(cx))^3 + b\left(\frac{1}{2}\left(2ib\left(b \int \operatorname{PolyLog}(2, -ie^{\operatorname{sech}^{-1}(cx)}) d\operatorname{sech}^{-1}(cx) - \operatorname{PolyLog}(2, -ie^{\operatorname{sech}^{-1}(cx)})\right)\right)\right)$$

↓ 2720

$$-\frac{1}{3}c^3x^3(a + b\operatorname{sech}^{-1}(cx))^3 + b\left(\frac{1}{2}\left(2ib\left(b \int e^{-\operatorname{sech}^{-1}(cx)} \operatorname{PolyLog}(2, -ie^{\operatorname{sech}^{-1}(cx)}) de^{\operatorname{sech}^{-1}(cx)} - \operatorname{PolyLog}(2, -ie^{\operatorname{sech}^{-1}(cx)})\right)\right)\right)$$

↓ 7143

$$-\frac{1}{3}c^3x^3(a + b\operatorname{sech}^{-1}(cx))^3 + b\left(\frac{1}{2}\left(2 \arctan(e^{\operatorname{sech}^{-1}(cx)}) (a + b\operatorname{sech}^{-1}(cx))^2 + 2ib\left(b \operatorname{PolyLog}(3, -ie^{\operatorname{sech}^{-1}(cx)}) - \operatorname{PolyLog}(3, -ie^{\operatorname{sech}^{-1}(cx)})\right)\right)\right)$$

input `Int[x^2*(a + b*ArcSech[c*x])^3,x]`

output `-((-1/3*(c^3*x^3*(a + b*ArcSech[c*x])^3) + b*(b*c*x*(a + b*ArcSech[c*x]) + (c*x*sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x])^2)/2 - b^2*ArcTan[(sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(c*x)] + (2*(a + b*ArcSech[c*x])^2*ArcTan[E^ArcSech[c*x]]) + (2*I)*b*(-((a + b*ArcSech[c*x])*PolyLog[2, (-I)*E^ArcSech[c*x]]) + b*PolyLog[3, (-I)*E^ArcSech[c*x]]) - (2*I)*b*(-((a + b*ArcSech[c*x])*PolyLog[2, I*E^ArcSech[c*x]]) + b*PolyLog[3, I*E^ArcSech[c*x]]))/2)/c^3)`

## Definitions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))] * ((f_) + (g_) * (x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m * (PolyLog[2, (-e)*(F^(c*(a + b*x)))^n] / (b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1) * PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4668 `Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)] * ((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m * (ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)] / (f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1) * Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1) * Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 4674 `Int[(csc[(e_) + (f_)*(x_)]*(b_))^(n_)*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[(-b^2)*(c + d*x)^m * Cot[e + f*x] * ((b*Csc[e + f*x])^(n - 2) / (f*(n - 1))), x] + (-Simp[b^2*d*m*(c + d*x)^(m - 1) * ((b*Csc[e + f*x])^(n - 2) / (f^2*(n - 1)*(n - 2))), x] + Simp[b^2*d^2*m*(m - 1) / (f^2*(n - 1)*(n - 2)) Int[(c + d*x)^(m - 2) * (b*Csc[e + f*x])^(n - 2), x], x] + Simp[b^2*((n - 2) / (n - 1)) Int[(c + d*x)^m * (b*Csc[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2] && GtQ[m, 1]`

rule 5974

```
Int[((c_.) + (d_.)*(x_)^(m_.)*Sech[(a_.) + (b_.)*(x_)^(n_.)*Tanh[(a_.) +
(b_.)*(x_)^(p_.)], x_Symbol] := Simp[(-c + d*x)^m*(Sech[a + b*x]^n/(b*n))
, x] + Simp[d*(m/(b*n)) Int[(c + d*x)^(m - 1)*Sech[a + b*x]^n, x], x] /;
FreeQ[{a, b, c, d, n}, x] && EqQ[p, 1] && GtQ[m, 0]
```

rule 6839

```
Int[((a_.) + ArcSech[(c_.)*(x_)*(b_.)]^(n_.)*(x_)^(m_.), x_Symbol] := Simp[
-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

**Maple [F]**

$$\int x^2(a + b \operatorname{arcsech}(cx))^3 dx$$

input

```
int(x^2*(a+b*arcsech(c*x))^3,x)
```

output

```
int(x^2*(a+b*arcsech(c*x))^3,x)
```

**Fricas [F]**

$$\int x^2(a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (b \operatorname{arsech}(cx) + a)^3 x^2 dx$$

input

```
integrate(x^2*(a+b*arcsech(c*x))^3,x, algorithm="fricas")
```

output

```
integral(b^3*x^2*arcsech(c*x)^3 + 3*a*b^2*x^2*arcsech(c*x)^2 + 3*a^2*b*x^2
*arcsech(c*x) + a^3*x^2, x)
```

**Sympy [F]**

$$\int x^2 (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int x^2 (a + b \operatorname{arosech}(cx))^3 dx$$

input `integrate(x**2*(a+b*asech(c*x))**3,x)`

output `Integral(x**2*(a + b*asech(c*x))**3, x)`

**Maxima [F]**

$$\int x^2 (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (b \operatorname{arosech}(cx) + a)^3 x^2 dx$$

input `integrate(x^2*(a+b*arcsech(c*x))^3,x, algorithm="maxima")`

output `1/3*a^3*x^3 + 1/2*(2*x^3*arcsech(c*x) - (sqrt(1/(c^2*x^2) - 1)/(c^2*(1/(c^2*x^2) - 1) + c^2) + arctan(sqrt(1/(c^2*x^2) - 1)/c^2)/c)*a^2*b + integrate(b^3*x^2*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^3 + 3*a*b^2*x^2*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^2, x)`

**Giac [F]**

$$\int x^2 (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (b \operatorname{arosech}(cx) + a)^3 x^2 dx$$

input `integrate(x^2*(a+b*arcsech(c*x))^3,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)^3*x^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int x^2 \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right)^3 dx$$

input `int(x^2*(a + b*acosh(1/(c*x)))^3,x)`output `int(x^2*(a + b*acosh(1/(c*x)))^3, x)`**Reduce [F]**

$$\int x^2 (a + b \operatorname{sech}^{-1}(cx))^3 dx = 3 \left( \int \operatorname{asech}(cx) x^2 dx \right) a^2 b + \left( \int \operatorname{asech}(cx)^3 x^2 dx \right) b^3 + 3 \left( \int \operatorname{asech}(cx)^2 x^2 dx \right) a b^2 + \frac{a^3 x^3}{3}$$

input `int(x^2*(a+b*asech(c*x))^3,x)`output `(9*int(asech(c*x)*x**2,x)*a**2*b + 3*int(asech(c*x)**3*x**2,x)*b**3 + 9*int(asech(c*x)**2*x**2,x)*a*b**2 + a**3*x**3)/3`

### 3.44 $\int x(a + b\operatorname{sech}^{-1}(cx))^3 dx$

Optimal result	398
Mathematica [A] (verified)	399
Rubi [C] (verified)	399
Maple [A] (verified)	403
Fricas [F]	403
Sympy [F]	404
Maxima [F]	404
Giac [F]	404
Mupad [F(-1)]	405
Reduce [F]	405

#### Optimal result

Integrand size = 12, antiderivative size = 126

$$\int x(a + b\operatorname{sech}^{-1}(cx))^3 dx = -\frac{3b(a + b\operatorname{sech}^{-1}(cx))^2}{2c^2} - \frac{3b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(a + b\operatorname{sech}^{-1}(cx))^2}{2c^2} + \frac{1}{2}x^2(a + b\operatorname{sech}^{-1}(cx))^3 + \frac{3b^2(a + b\operatorname{sech}^{-1}(cx)) \log(1 + e^{2\operatorname{sech}^{-1}(cx)})}{c^2} + \frac{3b^3 \operatorname{PolyLog}(2, -e^{2\operatorname{sech}^{-1}(cx)})}{2c^2}$$

output

```
-3/2*b*(a+b*arcsech(c*x))^2/c^2-3/2*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+b*arcsech(c*x))^2/c^2+1/2*x^2*(a+b*arcsech(c*x))^3+3*b^2*(a+b*arcsech(c*x))*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)/c^2+3/2*b^3*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)/c^2
```

**Mathematica [A] (verified)**

Time = 0.68 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.74

$$\int x(a + b\operatorname{sech}^{-1}(cx))^3 dx$$

$$= \frac{-3b^2\left(-ac^2x^2 + b\left(-1 + \sqrt{\frac{1-cx}{1+cx}} + cx\sqrt{\frac{1-cx}{1+cx}}\right)\right)\operatorname{sech}^{-1}(cx)^2 + b^3c^2x^2\operatorname{sech}^{-1}(cx)^3 + 3b\operatorname{sech}^{-1}(cx)\left(a\left(ac^2\right.\right.$$

input

```
Integrate[x*(a + b*ArcSech[c*x])^3,x]
```

output

```
(-3*b^2*(-(a*c^2*x^2) + b*(-1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]))*ArcSech[c*x]^2 + b^3*c^2*x^2*ArcSech[c*x]^3 + 3*b*ArcSech[c*x]*(a*(a*c^2*x^2 - 2*b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)) + 2*b^2*Log[1 + E^(-2*ArcSech[c*x])]) + a*(a*(a*c^2*x^2 - 3*b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)) + 6*b^2*Log[1/(c*x)]) - 3*b^3*PolyLog[2, -E^(-2*ArcSech[c*x])])/(2*c^2)
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.63 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.08, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$ , Rules used = {6839, 5974, 3042, 4672, 26, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(a + b\operatorname{sech}^{-1}(cx))^3 dx$$

$$\downarrow 6839$$

$$\frac{\int c^2x^2\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))^3 d\operatorname{sech}^{-1}(cx)}{c^2}$$

$$\downarrow 5974$$



$$\begin{aligned}
& - \frac{\frac{3}{2}b \int c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^2 d \operatorname{sech}^{-1}(cx) - \frac{1}{2}c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^3}{c^2} \\
& \quad \downarrow \text{3042} \\
& - \frac{-\frac{1}{2}c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^3 + \frac{3}{2}b \int (a + b \operatorname{sech}^{-1}(cx))^2 \csc \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^2 d \operatorname{sech}^{-1}(cx)}{c^2} \\
& \quad \downarrow \text{4672} \\
& - \frac{-\frac{1}{2}c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^3 + \frac{3}{2}b \left( \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx))^2 - 2ib \int -i \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx)) \right)}{c^2} \\
& \quad \downarrow \text{26} \\
& - \frac{\frac{3}{2}b \left( \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx))^2 - 2b \int \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) \right) - \frac{1}{2}c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^3}{c^2} \\
& \quad \downarrow \text{3042} \\
& - \frac{-\frac{1}{2}c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^3 + \frac{3}{2}b \left( \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx))^2 - 2b \int -i (a + b \operatorname{sech}^{-1}(cx)) \tan \left( i \operatorname{sech}^{-1}(cx) \right) \right)}{c^2} \\
& \quad \downarrow \text{26} \\
& - \frac{-\frac{1}{2}c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^3 + \frac{3}{2}b \left( \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx))^2 + 2ib \int (a + b \operatorname{sech}^{-1}(cx)) \tan \left( i \operatorname{sech}^{-1}(cx) \right) \right)}{c^2} \\
& \quad \downarrow \text{4201} \\
& - \frac{-\frac{1}{2}c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^3 + \frac{3}{2}b \left( \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx))^2 + 2ib \left( 2i \int \frac{e^{2 \operatorname{sech}^{-1}(cx)} (a + b \operatorname{sech}^{-1}(cx))}{1 + e^{2 \operatorname{sech}^{-1}(cx)}} d \operatorname{sech}^{-1}(cx) \right) \right)}{c^2} \\
& \quad \downarrow \text{2620} \\
& - \frac{-\frac{1}{2}c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^3 + \frac{3}{2}b \left( \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx))^2 + 2ib \left( 2i \left( \frac{1}{2} \log \left( e^{2 \operatorname{sech}^{-1}(cx)} + 1 \right) \right) (a + b \operatorname{sech}^{-1}(cx)) \right) \right)}{c^2} \\
& \quad \downarrow \text{2715}
\end{aligned}$$

$$\frac{-\frac{1}{2}c^2x^2(a + b\operatorname{sech}^{-1}(cx))^3 + \frac{3}{2}b\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))^2 + 2ib\left(2i\left(\frac{1}{2}\log\left(e^{2\operatorname{sech}^{-1}(cx)} + 1\right)\right)(a + b\right)\right)}{c^2}$$

↓ 2838

$$\frac{-\frac{1}{2}c^2x^2(a + b\operatorname{sech}^{-1}(cx))^3 + \frac{3}{2}b\left(\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b\operatorname{sech}^{-1}(cx))^2 + 2ib\left(2i\left(\frac{1}{2}\log\left(e^{2\operatorname{sech}^{-1}(cx)} + 1\right)\right)(a + b\right)\right)}{c^2}$$

input `Int[x*(a + b*ArcSech[c*x])^3,x]`

output `-((-1/2*(c^2*x^2*(a + b*ArcSech[c*x])^3) + (3*b*(Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x])^2 + (2*I)*b*((( -1/2*I)*(a + b*ArcSech[c*x])^2)/b + (2*I)*(((a + b*ArcSech[c*x])*Log[1 + E^(2*ArcSech[c*x])])/2 + (b*PolyLog[2, -E^(2*ArcSech[c*x])])/4))))/2)/c^2)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838  $\text{Int}[\text{Log}[(c\_.) * ((d\_.) + (e\_.) * (x\_.)^{(n\_.)})] / (x\_.), x\_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c) * e * x^n / n, x] /; \text{FreeQ}[\{c, d, e, n\}, x] \ \&\& \ \text{EqQ}[c * d, 1]$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4201  $\text{Int}[((c\_.) + (d\_.) * (x\_.)^{(m\_.)}) * \tan[(e\_.) + (\text{Complex}[0, fz\_]) * (f\_.) * (x\_.)], x\_Symbol] \rightarrow \text{Simp}[(-I) * ((c + d * x)^{(m + 1)} / (d * (m + 1))), x] + \text{Simp}[2 * I \ \text{Int}[(c + d * x)^m * (E^{(2 * ((-I) * e + f * fz * x))} / (1 + E^{(2 * ((-I) * e + f * fz * x))}))], x], x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \ \&\& \ \text{IGtQ}[m, 0]$

rule 4672  $\text{Int}[\text{csc}[(e\_.) + (f\_.) * (x\_)]^{2 * ((c\_.) + (d\_.) * (x\_.)^{(m\_.)})}, x\_Symbol] \rightarrow \text{Simp}[(-c + d * x)^m * (\text{Cot}[e + f * x] / f), x] + \text{Simp}[d * (m / f) \ \text{Int}[(c + d * x)^{(m - 1)} * \text{Cot}[e + f * x], x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \ \&\& \ \text{GtQ}[m, 0]$

rule 5974  $\text{Int}[((c\_.) + (d\_.) * (x\_.)^{(m\_.)}) * \text{Sech}[(a\_.) + (b\_.) * (x\_)]^{(n\_.)} * \text{Tanh}[(a\_.) + (b\_.) * (x\_)]^{(p\_.)}, x\_Symbol] \rightarrow \text{Simp}[(-c + d * x)^m * (\text{Sech}[a + b * x]^n / (b * n)), x] + \text{Simp}[d * (m / (b * n)) \ \text{Int}[(c + d * x)^{(m - 1)} * \text{Sech}[a + b * x]^n, x], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \ \&\& \ \text{EqQ}[p, 1] \ \&\& \ \text{GtQ}[m, 0]$

rule 6839  $\text{Int}[((a\_.) + \text{ArcSech}[(c\_.) * (x\_)] * (b\_.)^{(n\_.)}) * (x\_.)^{(m\_.)}, x\_Symbol] \rightarrow \text{Simp}[-(c^{(m + 1)})^{(-1)} \ \text{Subst}[\text{Int}[(a + b * x)^n * \text{Sech}[x]^{(m + 1)} * \text{Tanh}[x], x], x, \text{ArcSech}[c * x]], x] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ (\text{GtQ}[n, 0] \ || \ \text{LtQ}[m, -1])$

### Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 318, normalized size of antiderivative = 2.52

method	result
derivativedivides	$\frac{c^2 x^2 a^3}{2} + b^3 \left( \frac{\operatorname{arcsech}(cx)^2 \left( c^2 x^2 \operatorname{arcsech}(cx) - 3 \sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x + 3 \right)}{2} - 3 \operatorname{arcsech}(cx)^2 + 3 \operatorname{arcsech}(cx) \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1} \right) \right) \right)$
default	$\frac{c^2 x^2 a^3}{2} + b^3 \left( \frac{\operatorname{arcsech}(cx)^2 \left( c^2 x^2 \operatorname{arcsech}(cx) - 3 \sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x + 3 \right)}{2} - 3 \operatorname{arcsech}(cx)^2 + 3 \operatorname{arcsech}(cx) \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1} \right) \right) \right)$
parts	$\frac{x^2 a^3}{2} + \frac{b^3 \left( \frac{\operatorname{arcsech}(cx)^2 \left( c^2 x^2 \operatorname{arcsech}(cx) - 3 \sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x + 3 \right)}{2} - 3 \operatorname{arcsech}(cx)^2 + 3 \operatorname{arcsech}(cx) \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1} \right) \right) \right)}{c^2}$

input `int(x*(a+b*arcsech(c*x))^3,x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{c^2} \left( \frac{1}{2} c^2 x^2 a^3 + b^3 \left( \frac{1}{2} \operatorname{arcsech}(cx)^2 \left( c^2 x^2 \operatorname{arcsech}(cx) - 3 \left( -\frac{cx-1}{c/x} \right)^{1/2} c \left( \frac{cx+1}{c/x} \right)^{1/2} x + 3 \right) - 3 \operatorname{arcsech}(cx)^2 + 3 \operatorname{arcsech}(cx) \ln \left( 1 + \left( \frac{1}{c/x} + (-1+1/c/x)^{1/2} (1+1/c/x)^{1/2} \right)^2 \right) + 3/2 \operatorname{polylog}(2, -\left( \frac{1}{c/x} + (-1+1/c/x)^{1/2} (1+1/c/x)^{1/2} \right)^2) \right) + 3 a b^2 \left( -2 \operatorname{arcsech}(cx) + 1/2 \operatorname{arcsech}(cx) \left( c^2 x^2 \operatorname{arcsech}(cx) - 2 \left( -\frac{cx-1}{c/x} \right)^{1/2} c \left( \frac{cx+1}{c/x} \right)^{1/2} x + 2 \right) + \ln \left( 1 + \left( \frac{1}{c/x} + (-1+1/c/x)^{1/2} (1+1/c/x)^{1/2} \right)^2 \right) \right) + 3 a^2 b \left( \frac{1}{2} c^2 x^2 \operatorname{arcsech}(cx) - 1/2 \left( -\frac{cx-1}{c/x} \right)^{1/2} c \left( \frac{cx+1}{c/x} \right)^{1/2} x \right) \right)$$

### Fricas [F]

$$\int x(a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^3 x dx$$

input `integrate(x*(a+b*arcsech(c*x))^3,x, algorithm="fricas")`

output `integral(b^3*x*arcsech(c*x)^3 + 3*a*b^2*x*arcsech(c*x)^2 + 3*a^2*b*x*arcsech(c*x) + a^3*x, x)`

**Sympy [F]**

$$\int x(a + b \operatorname{sech}^{-1}(cx))^3 dx = \int x(a + b \operatorname{arsech}(cx))^3 dx$$

input `integrate(x*(a+b*asech(c*x))**3,x)`

output `Integral(x*(a + b*asech(c*x))**3, x)`

**Maxima [F]**

$$\int x(a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (b \operatorname{arsech}(cx) + a)^3 x dx$$

input `integrate(x*(a+b*arcsech(c*x))^3,x, algorithm="maxima")`

output `3/2*a*b^2*x^2*arcsech(c*x)^2 + 1/2*a^3*x^2 + 3/2*(x^2*arcsech(c*x) - x*sqrt(1/(c^2*x^2) - 1)/c)*a^2*b - 3*(x*sqrt(1/(c^2*x^2) - 1)*arcsech(c*x)/c + log(x)/c^2)*a*b^2 + b^3*integrate(x*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^3, x)`

**Giac [F]**

$$\int x(a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (b \operatorname{arsech}(cx) + a)^3 x dx$$

input `integrate(x*(a+b*arcsech(c*x))^3,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)^3*x, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x(a + b \operatorname{sech}^{-1}(cx))^3 dx = \int x \left( a + b \operatorname{acosh} \left( \frac{1}{cx} \right) \right)^3 dx$$

input `int(x*(a + b*acosh(1/(c*x)))^3,x)`output `int(x*(a + b*acosh(1/(c*x)))^3, x)`**Reduce [F]**

$$\int x(a + b \operatorname{sech}^{-1}(cx))^3 dx = 3 \left( \int a \operatorname{sech}(cx) x dx \right) a^2 b + \left( \int a \operatorname{sech}(cx)^3 x dx \right) b^3 \\ + 3 \left( \int a \operatorname{sech}(cx)^2 x dx \right) a b^2 + \frac{a^3 x^2}{2}$$

input `int(x*(a+b*asech(c*x))^3,x)`output `(6*int(asech(c*x)*x,x)*a**2*b + 2*int(asech(c*x)**3*x,x)*b**3 + 6*int(asech(c*x)**2*x,x)*a*b**2 + a**3*x**2)/2`

### 3.45 $\int (a + b \operatorname{sech}^{-1}(cx))^3 dx$

Optimal result	406
Mathematica [B] (verified)	407
Rubi [A] (verified)	407
Maple [F]	410
Fricas [F]	410
Sympy [F]	410
Maxima [F]	411
Giac [F]	411
Mupad [F(-1)]	412
Reduce [F]	412

#### Optimal result

Integrand size = 10, antiderivative size = 140

$$\int (a + b \operatorname{sech}^{-1}(cx))^3 dx = x(a + b \operatorname{sech}^{-1}(cx))^3 - \frac{6b(a + b \operatorname{sech}^{-1}(cx))^2 \arctan(e^{\operatorname{sech}^{-1}(cx)})}{c} + \frac{6ib^2(a + b \operatorname{sech}^{-1}(cx)) \operatorname{PolyLog}(2, -ie^{\operatorname{sech}^{-1}(cx)})}{c} - \frac{6ib^2(a + b \operatorname{sech}^{-1}(cx)) \operatorname{PolyLog}(2, ie^{\operatorname{sech}^{-1}(cx)})}{c} - \frac{6ib^3 \operatorname{PolyLog}(3, -ie^{\operatorname{sech}^{-1}(cx)})}{c} + \frac{6ib^3 \operatorname{PolyLog}(3, ie^{\operatorname{sech}^{-1}(cx)})}{c}$$

output

```
x*(a+b*arcsech(c*x))^3-6*b*(a+b*arcsech(c*x))^2*arctan(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/c+6*I*b^2*(a+b*arcsech(c*x))*polylog(2,-I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/c-6*I*b^2*(a+b*arcsech(c*x))*polylog(2,I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/c-6*I*b^3*polylog(3,-I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/c+6*I*b^3*polylog(3,I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/c
```

**Mathematica [B] (verified)**

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 282 vs.  $2(140) = 280$ .

Time = 0.35 (sec) , antiderivative size = 282, normalized size of antiderivative = 2.01

$$\int (a + b \operatorname{sech}^{-1}(cx))^3 dx = a^3 x + 3a^2 b x \operatorname{sech}^{-1}(cx) - \frac{3a^2 b \arctan\left(\frac{cx \sqrt{\frac{1-cx}{1+cx}}}{-1+cx}\right)}{c} + \frac{3iab^2 \left( \operatorname{sech}^{-1}(cx) \left( -icx \operatorname{sech}^{-1}(cx) + 2 \log\left(1 - ie^{-\operatorname{sech}^{-1}(cx)}\right) - 2 \log\left(1 + ie^{-\operatorname{sech}^{-1}(cx)}\right) \right) \right) + 2 \operatorname{PolyLog}}{c} + \frac{b^3 \left( cx \operatorname{sech}^{-1}(cx)^3 - 3i \left( -\operatorname{sech}^{-1}(cx)^2 \left( \log\left(1 - ie^{-\operatorname{sech}^{-1}(cx)}\right) - \log\left(1 + ie^{-\operatorname{sech}^{-1}(cx)}\right) \right) \right) - 2 \operatorname{sech}^{-1}(cx) \right)}{c}$$

input `Integrate[(a + b*ArcSech[c*x])^3,x]`

output `a^3*x + 3*a^2*b*x*ArcSech[c*x] - (3*a^2*b*ArcTan[(c*x*Sqrt[(1 - c*x)/(1 + c*x)])/(-1 + c*x)])/c + ((3*I)*a*b^2*(ArcSech[c*x]*((-I)*c*x*ArcSech[c*x] + 2*Log[1 - I/E^ArcSech[c*x]] - 2*Log[1 + I/E^ArcSech[c*x]]) + 2*PolyLog[2, (-I)/E^ArcSech[c*x]] - 2*PolyLog[2, I/E^ArcSech[c*x]]))/c + (b^3*(c*x*ArcSech[c*x]^3 - (3*I)*(-ArcSech[c*x]^2*(Log[1 - I/E^ArcSech[c*x]] - Log[1 + I/E^ArcSech[c*x]]) - 2*ArcSech[c*x]*(PolyLog[2, (-I)/E^ArcSech[c*x]] - PolyLog[2, I/E^ArcSech[c*x]]) - 2*(PolyLog[3, (-I)/E^ArcSech[c*x]] - PolyLog[3, I/E^ArcSech[c*x]]))))/c`

**Rubi [A] (verified)**

Time = 0.59 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.91, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$ , Rules used = {6833, 5974, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{sech}^{-1}(cx))^3 dx$$



$$\begin{aligned} & \downarrow 6833 \\ & \frac{\int cx \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a+b\operatorname{sech}^{-1}(cx))^3 d\operatorname{sech}^{-1}(cx)}{c} \\ & \downarrow 5974 \\ & \frac{3b \int cx (a+b\operatorname{sech}^{-1}(cx))^2 d\operatorname{sech}^{-1}(cx) - cx (a+b\operatorname{sech}^{-1}(cx))^3}{c} \\ & \downarrow 3042 \\ & \frac{-cx (a+b\operatorname{sech}^{-1}(cx))^3 + 3b \int (a+b\operatorname{sech}^{-1}(cx))^2 \csc\left(\operatorname{isech}^{-1}(cx) + \frac{\pi}{2}\right) d\operatorname{sech}^{-1}(cx)}{c} \\ & \downarrow 4668 \\ & \frac{-cx (a+b\operatorname{sech}^{-1}(cx))^3 + 3b \left( -2ib \int (a+b\operatorname{sech}^{-1}(cx)) \log\left(1 - ie^{\operatorname{sech}^{-1}(cx)}\right) d\operatorname{sech}^{-1}(cx) + 2ib \int (a+b\operatorname{sech}^{-1}(cx)) \right)}{c} \\ & \downarrow 3011 \\ & \frac{-cx (a+b\operatorname{sech}^{-1}(cx))^3 + 3b \left( 2ib \left( b \int \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(cx)}\right) d\operatorname{sech}^{-1}(cx) - \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(cx)}\right) \right) (a+b\operatorname{sech}^{-1}(cx)) \right)}{c} \\ & \downarrow 2720 \\ & \frac{-cx (a+b\operatorname{sech}^{-1}(cx))^3 + 3b \left( 2ib \left( b \int e^{-\operatorname{sech}^{-1}(cx)} \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(cx)}\right) d\operatorname{sech}^{-1}(cx) - \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(cx)}\right) \right) \right)}{c} \\ & \downarrow 7143 \\ & \frac{-cx (a+b\operatorname{sech}^{-1}(cx))^3 + 3b \left( 2 \arctan\left(e^{\operatorname{sech}^{-1}(cx)}\right) (a+b\operatorname{sech}^{-1}(cx))^2 + 2ib \left( b \operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(cx)}\right) - \operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(cx)}\right) \right) \right)}{c} \end{aligned}$$

input `Int[(a + b*ArcSech[c*x])^3,x]`

output `-((-c*x*(a + b*ArcSech[c*x])^3) + 3*b*(2*(a + b*ArcSech[c*x])^2*ArcTan[E^ArcSech[c*x]] + (2*I)*b*(-((a + b*ArcSech[c*x])*PolyLog[2, (-I)*E^ArcSech[c*x]]) + b*PolyLog[3, (-I)*E^ArcSech[c*x]]) - (2*I)*b*(-((a + b*ArcSech[c*x])*PolyLog[2, I*E^ArcSech[c*x]]) + b*PolyLog[3, I*E^ArcSech[c*x]])))/c`

## Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 5974 `Int[((c_) + (d_)*(x_))^(m_)*Sech[(a_) + (b_)*(x_)]^(n_)*Tanh[(a_) + (b_)*(x_)]^(p_), x_Symbol] := Simp[(-c + d*x)^m*(Sech[a + b*x]^n/(b^n)), x] + Simp[d*(m/(b^n)) Int[(c + d*x)^(m - 1)*Sech[a + b*x]^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[p, 1] && GtQ[m, 0]`

rule 6833 `Int[((a_) + ArcSech[(c_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[-c^(-1) Subst[Int[(a + b*x)^n*Sech[x]*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[n, 0]`

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol]
:> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x]
&& EqQ[b*d, a*e]
```

**Maple [F]**

$$\int (a + b \operatorname{arcsech}(cx))^3 dx$$

input

```
int((a+b*arcsech(c*x))^3,x)
```

output

```
int((a+b*arcsech(c*x))^3,x)
```

**Fricas [F]**

$$\int (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (b \operatorname{arsech}(cx) + a)^3 dx$$

input

```
integrate((a+b*arcsech(c*x))^3,x, algorithm="fricas")
```

output

```
integral(b^3*arcsech(c*x)^3 + 3*a*b^2*arcsech(c*x)^2 + 3*a^2*b*arcsech(c*x)
) + a^3, x)
```

**Sympy [F]**

$$\int (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (a + b \operatorname{asech}(cx))^3 dx$$

input

```
integrate((a+b*asech(c*x))**3,x)
```

output

```
Integral((a + b*asech(c*x))**3, x)
```

**Maxima [F]**

$$\int (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^3 dx$$

input `integrate((a+b*arcsech(c*x))^3,x, algorithm="maxima")`

output

```

b^3*x*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)^3 + a^3*x + 3*(c*x*arcsech(c*x)
) - arctan(sqrt(1/(c^2*x^2) - 1)))*a^2*b/c - integrate(-(b^3*log(c)^3 - 3*
a*b^2*log(c)^2 - (b^3*c^2*x^2 - b^3)*log(x)^3 - (b^3*c^2*log(c)^3 - 3*a*b^
2*c^2*log(c)^2)*x^2 + 3*(b^3*log(c) - a*b^2 - (b^3*c^2*log(c) - a*b^2*c^2)
*x^2 + (b^3*log(c) - a*b^2 - (b^3*c^2*(log(c) + 1) - a*b^2*c^2)*x^2 - (b^3
*c^2*x^2 - b^3)*log(x))*sqrt(c*x + 1)*sqrt(-c*x + 1) - (b^3*c^2*x^2 - b^3)
*log(x))*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)^2 + 3*(b^3*log(c) - a*b^2 -
(b^3*c^2*log(c) - a*b^2*c^2)*x^2)*log(x)^2 + (b^3*log(c)^3 - 3*a*b^2*log(
c)^2 - (b^3*c^2*x^2 - b^3)*log(x)^3 - (b^3*c^2*log(c)^3 - 3*a*b^2*c^2*log(
c)^2)*x^2 + 3*(b^3*log(c) - a*b^2 - (b^3*c^2*log(c) - a*b^2*c^2)*x^2)*log(
x)^2 + 3*(b^3*log(c)^2 - 2*a*b^2*log(c) - (b^3*c^2*log(c)^2 - 2*a*b^2*c^2*
log(c))*x^2)*log(x))*sqrt(c*x + 1)*sqrt(-c*x + 1) - 3*(b^3*log(c)^2 - 2*a*
b^2*log(c) - (b^3*c^2*log(c)^2 - 2*a*b^2*c^2*log(c))*x^2 - (b^3*c^2*x^2 -
b^3)*log(x)^2 + (b^3*log(c)^2 - 2*a*b^2*log(c) - (b^3*c^2*log(c)^2 - 2*a*b
^2*c^2*log(c))*x^2 - (b^3*c^2*x^2 - b^3)*log(x)^2 + 2*(b^3*log(c) - a*b^2
- (b^3*c^2*log(c) - a*b^2*c^2)*x^2)*log(x))*sqrt(c*x + 1)*sqrt(-c*x + 1) +
2*(b^3*log(c) - a*b^2 - (b^3*c^2*log(c) - a*b^2*c^2)*x^2)*log(x))*log(sqrt
t(c*x + 1)*sqrt(-c*x + 1) + 1) + 3*(b^3*log(c)^2 - 2*a*b^2*log(c) - (b^3*c
^2*log(c)^2 - 2*a*b^2*c^2*log(c))*x^2)*log(x))/(c^2*x^2 + (c^2*x^2 - 1)*sq
rt(c*x + 1)*sqrt(-c*x + 1) - 1), x)

```

**Giac [F]**

$$\int (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^3 dx$$

input `integrate((a+b*arcsech(c*x))^3,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right)^3 dx$$

input `int((a + b*acosh(1/(c*x)))^3,x)`output `int((a + b*acosh(1/(c*x)))^3, x)`**Reduce [F]**

$$\int (a + b \operatorname{sech}^{-1}(cx))^3 dx = 3 \left( \int \operatorname{asech}(cx) dx \right) a^2 b + \left( \int \operatorname{asech}(cx)^3 dx \right) b^3 \\ + 3 \left( \int \operatorname{asech}(cx)^2 dx \right) a b^2 + a^3 x$$

input `int((a+b*asech(c*x))^3,x)`output `3*int(asech(c*x),x)*a**2*b + int(asech(c*x)**3,x)*b**3 + 3*int(asech(c*x)*  
*2,x)*a*b**2 + a**3*x`

$$3.46 \quad \int \frac{(a+b\operatorname{sech}^{-1}(cx))^3}{x} dx$$

Optimal result	413
Mathematica [A] (verified)	414
Rubi [C] (verified)	414
Maple [B] (verified)	417
Fricas [F]	418
Sympy [F]	419
Maxima [F]	419
Giac [F]	419
Mupad [F(-1)]	420
Reduce [F]	420

### Optimal result

Integrand size = 14, antiderivative size = 114

$$\int \frac{(a + b\operatorname{sech}^{-1}(cx))^3}{x} dx = \frac{(a + b\operatorname{sech}^{-1}(cx))^4}{4b} - (a + b\operatorname{sech}^{-1}(cx))^3 \log\left(1 + e^{2\operatorname{sech}^{-1}(cx)}\right) - \frac{3}{2}b(a + b\operatorname{sech}^{-1}(cx))^2 \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(cx)}\right) + \frac{3}{2}b^2(a + b\operatorname{sech}^{-1}(cx)) \operatorname{PolyLog}\left(3, -e^{2\operatorname{sech}^{-1}(cx)}\right) - \frac{3}{4}b^3 \operatorname{PolyLog}\left(4, -e^{2\operatorname{sech}^{-1}(cx)}\right)$$

output

```
1/4*(a+b*arcsech(c*x))^4/b-(a+b*arcsech(c*x))^3*ln(1+(1/c/x+(-1+1/c/x)^(1/2))*(1+1/c/x)^(1/2))^2)-3/2*b*(a+b*arcsech(c*x))^2*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2))*(1+1/c/x)^(1/2))^2)+3/2*b^2*(a+b*arcsech(c*x))*polylog(3,-(1/c/x+(-1+1/c/x)^(1/2))*(1+1/c/x)^(1/2))^2)-3/4*b^3*polylog(4,-(1/c/x+(-1+1/c/x)^(1/2))*(1+1/c/x)^(1/2))^2)
```

**Mathematica [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.60

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x} dx = \frac{1}{4} \left( -6a^2 b \operatorname{sech}^{-1}(cx)^2 - 4ab^2 \operatorname{sech}^{-1}(cx)^3 - b^3 \operatorname{sech}^{-1}(cx)^4 \right. \\ \left. - 12a^2 b \operatorname{sech}^{-1}(cx) \log \left( 1 + e^{-2 \operatorname{sech}^{-1}(cx)} \right) \right. \\ \left. - 12ab^2 \operatorname{sech}^{-1}(cx)^2 \log \left( 1 + e^{-2 \operatorname{sech}^{-1}(cx)} \right) \right. \\ \left. - 4b^3 \operatorname{sech}^{-1}(cx)^3 \log \left( 1 + e^{-2 \operatorname{sech}^{-1}(cx)} \right) + 4a^3 \log(cx) \right. \\ \left. + 6b(a + b \operatorname{sech}^{-1}(cx))^2 \operatorname{PolyLog} \left( 2, -e^{-2 \operatorname{sech}^{-1}(cx)} \right) \right. \\ \left. + 6b^2(a + b \operatorname{sech}^{-1}(cx)) \operatorname{PolyLog} \left( 3, -e^{-2 \operatorname{sech}^{-1}(cx)} \right) \right. \\ \left. + 3b^3 \operatorname{PolyLog} \left( 4, -e^{-2 \operatorname{sech}^{-1}(cx)} \right) \right)$$

input `Integrate[(a + b*ArcSech[c*x])^3/x,x]`

output `(-6*a^2*b*ArcSech[c*x]^2 - 4*a*b^2*ArcSech[c*x]^3 - b^3*ArcSech[c*x]^4 - 12*a^2*b*ArcSech[c*x]*Log[1 + E^(-2*ArcSech[c*x])] - 12*a*b^2*ArcSech[c*x]^2*Log[1 + E^(-2*ArcSech[c*x])] - 4*b^3*ArcSech[c*x]^3*Log[1 + E^(-2*ArcSech[c*x])] + 4*a^3*Log[c*x] + 6*b*(a + b*ArcSech[c*x])^2*PolyLog[2, -E^(-2*ArcSech[c*x])] + 6*b^2*(a + b*ArcSech[c*x])*PolyLog[3, -E^(-2*ArcSech[c*x])] + 3*b^3*PolyLog[4, -E^(-2*ArcSech[c*x])])/4`

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.69 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.14, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$ , Rules used = {6839, 3042, 26, 4201, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x} dx \\
& \quad \downarrow \text{6839} \\
& - \int \sqrt{\frac{1-cx}{cx+1}} (cx+1) (a + b \operatorname{sech}^{-1}(cx))^3 d \operatorname{sech}^{-1}(cx) \\
& \quad \downarrow \text{3042} \\
& - \int -i(a + b \operatorname{sech}^{-1}(cx))^3 \tan(i \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) \\
& \quad \downarrow \text{26} \\
& i \int (a + b \operatorname{sech}^{-1}(cx))^3 \tan(i \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) \\
& \quad \downarrow \text{4201} \\
& i \left( 2i \int \frac{e^{2 \operatorname{sech}^{-1}(cx)} (a + b \operatorname{sech}^{-1}(cx))^3}{1 + e^{2 \operatorname{sech}^{-1}(cx)}} d \operatorname{sech}^{-1}(cx) - \frac{i(a + b \operatorname{sech}^{-1}(cx))^4}{4b} \right) \\
& \quad \downarrow \text{2620} \\
& i \left( 2i \left( \frac{1}{2} \log(e^{2 \operatorname{sech}^{-1}(cx)} + 1) (a + b \operatorname{sech}^{-1}(cx))^3 - \frac{3}{2} b \int (a + b \operatorname{sech}^{-1}(cx))^2 \log(1 + e^{2 \operatorname{sech}^{-1}(cx)}) d \operatorname{sech}^{-1}(cx) \right) \right) \\
& \quad \downarrow \text{3011} \\
& i \left( 2i \left( \frac{1}{2} \log(e^{2 \operatorname{sech}^{-1}(cx)} + 1) (a + b \operatorname{sech}^{-1}(cx))^3 - \frac{3}{2} b \left( b \int (a + b \operatorname{sech}^{-1}(cx)) \operatorname{PolyLog}(2, -e^{2 \operatorname{sech}^{-1}(cx)}) d \operatorname{sech}^{-1}(cx) \right) \right) \right) \\
& \quad \downarrow \text{7163} \\
& i \left( 2i \left( \frac{1}{2} \log(e^{2 \operatorname{sech}^{-1}(cx)} + 1) (a + b \operatorname{sech}^{-1}(cx))^3 - \frac{3}{2} b \left( b \left( \frac{1}{2} \operatorname{PolyLog}(3, -e^{2 \operatorname{sech}^{-1}(cx)}) (a + b \operatorname{sech}^{-1}(cx)) - \frac{1}{2} b \right) \right) \right) \right) \\
& \quad \downarrow \text{2720} \\
& i \left( 2i \left( \frac{1}{2} \log(e^{2 \operatorname{sech}^{-1}(cx)} + 1) (a + b \operatorname{sech}^{-1}(cx))^3 - \frac{3}{2} b \left( b \left( \frac{1}{2} \operatorname{PolyLog}(3, -e^{2 \operatorname{sech}^{-1}(cx)}) (a + b \operatorname{sech}^{-1}(cx)) - \frac{1}{4} b \right) \right) \right) \right) \\
& \quad \downarrow \text{7143}
\end{aligned}$$



$$i \left( 2i \left( \frac{1}{2} \log \left( e^{2\operatorname{sech}^{-1}(cx)} + 1 \right) (a + b\operatorname{sech}^{-1}(cx))^3 - \frac{3}{2} b \left( \frac{1}{2} \operatorname{PolyLog} \left( 3, -e^{2\operatorname{sech}^{-1}(cx)} \right) (a + b\operatorname{sech}^{-1}(cx)) - \frac{1}{4} b \right. \right. \right.$$

input `Int[(a + b*ArcSech[c*x])^3/x,x]`

output `I*(((−1/4*I)*(a + b*ArcSech[c*x])^4)/b + (2*I)*(((a + b*ArcSech[c*x])^3*Log[1 + E^(2*ArcSech[c*x])])/2 - (3*b*(−1/2*((a + b*ArcSech[c*x])^2*PolyLog[2, −E^(2*ArcSech[c*x])]) + b*((a + b*ArcSech[c*x])*PolyLog[3, −E^(2*ArcSech[c*x])])/2 - (b*PolyLog[4, −E^(2*ArcSech[c*x])])/4))/2))`

### Definitions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4201 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x))))], x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6839 `Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.), x_Symbol] := Simp[-(c^(m + 1))^(n-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`

## Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 428 vs.  $2(206) = 412$ .

Time = 0.49 (sec) , antiderivative size = 429, normalized size of antiderivative = 3.76

method	result
parts	$a^3 \ln(x) + b^3 \left( \frac{\operatorname{arcsech}(cx)^4}{4} - \operatorname{arcsech}(cx)^3 \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right)^2 \right) - \dots \right)$
derivativedivides	$a^3 \ln(cx) + b^3 \left( \frac{\operatorname{arcsech}(cx)^4}{4} - \operatorname{arcsech}(cx)^3 \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right)^2 \right) - \dots \right)$
default	$a^3 \ln(cx) + b^3 \left( \frac{\operatorname{arcsech}(cx)^4}{4} - \operatorname{arcsech}(cx)^3 \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} \right)^2 \right) - \dots \right)$

input `int((a+b*arcsech(c*x))^3/x,x,method=_RETURNVERBOSE)`

output `a^3*ln(x)+b^3*(1/4*arcsech(c*x)^4-arcsech(c*x)^3*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)-3/2*arcsech(c*x)^2*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)+3/2*arcsech(c*x)*polylog(3,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)-3/4*polylog(4,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2))+3*a*b^2*(1/3*arcsech(c*x)^3-arcsech(c*x)^2*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)-arcsech(c*x)*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)+1/2*polylog(3,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2))+3*a^2*b*(1/2*arcsech(c*x)^2-arcsech(c*x)*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)-1/2*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2))`

## Fricas [F]

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3}{x} dx$$

input `integrate((a+b*arcsech(c*x))^3/x,x, algorithm="fricas")`

output `integral((b^3*arcsech(c*x)^3 + 3*a*b^2*arcsech(c*x)^2 + 3*a^2*b*arcsech(c*x) + a^3)/x, x)`

**Sympy [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x} dx = \int \frac{(a + b \operatorname{asech}(cx))^3}{x} dx$$

input `integrate((a+b*asech(c*x))**3/x,x)`

output `Integral((a + b*asech(c*x))**3/x, x)`

**Maxima [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x} dx = \int \frac{(b \operatorname{arsech}(cx) + a)^3}{x} dx$$

input `integrate((a+b*arcsech(c*x))^3/x,x, algorithm="maxima")`

output `a^3*log(x) + integrate(b^3*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^3/x + 3*a*b^2*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^2/x + 3*a^2*b*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/x, x)`

**Giac [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x} dx = \int \frac{(b \operatorname{arsech}(cx) + a)^3}{x} dx$$

input `integrate((a+b*arcsech(c*x))^3/x,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)^3/x, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x} dx = \int \frac{(a + b \operatorname{acosh}(\frac{1}{cx}))^3}{x} dx$$

input `int((a + b*acosh(1/(c*x)))^3/x,x)`output `int((a + b*acosh(1/(c*x)))^3/x, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x} dx = 3 \left( \int \frac{\operatorname{asech}(cx)}{x} dx \right) a^2 b + \left( \int \frac{\operatorname{asech}(cx)^3}{x} dx \right) b^3 \\ + 3 \left( \int \frac{\operatorname{asech}(cx)^2}{x} dx \right) a b^2 + \log(x) a^3$$

input `int((a+b*asech(c*x))^3/x,x)`output `3*int(asech(c*x)/x,x)*a**2*b + int(asech(c*x)**3/x,x)*b**3 + 3*int(asech(c*x)**2/x,x)*a*b**2 + log(x)*a**3`

**3.47**  $\int \frac{(a+b\operatorname{sech}^{-1}(cx))^3}{x^2} dx$

Optimal result	421
Mathematica [A] (verified)	422
Rubi [C] (verified)	422
Maple [B] (verified)	425
Fricas [B] (verification not implemented)	425
Sympy [F]	426
Maxima [A] (verification not implemented)	426
Giac [F]	427
Mupad [F(-1)]	427
Reduce [F]	428

**Optimal result**

Integrand size = 14, antiderivative size = 102

$$\int \frac{(a + b\operatorname{sech}^{-1}(cx))^3}{x^2} dx = \frac{6b^3 \sqrt{\frac{1-cx}{1+cx}}(1+cx)}{x} - \frac{6b^2(a + b\operatorname{sech}^{-1}(cx))}{x} + \frac{3b \sqrt{\frac{1-cx}{1+cx}}(1+cx)(a + b\operatorname{sech}^{-1}(cx))^2}{x} - \frac{(a + b\operatorname{sech}^{-1}(cx))^3}{x}$$

output

```
6*b^3*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/x-6*b^2*(a+b*arcsech(c*x))/x+3*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+b*arcsech(c*x))^2/x-(a+b*arcsech(c*x))^3/x
```

**Mathematica [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.62

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^2} dx = \frac{a^3 + 6ab^2 - 3a^2b\sqrt{\frac{1-cx}{1+cx}}(1+cx) - 6b^3\sqrt{\frac{1-cx}{1+cx}}(1+cx) + 3b(a^2 + 2b^2 - 2ab\sqrt{\frac{1-cx}{1+cx}}(1+cx)) \operatorname{sech}^{-1}(cx)}{x}$$

input

```
Integrate[(a + b*ArcSech[c*x])^3/x^2,x]
```

output

```

-((a^3 + 6*a*b^2 - 3*a^2*b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x) - 6*b^3*Sqr
t[(1 - c*x)/(1 + c*x)]*(1 + c*x) + 3*b*(a^2 + 2*b^2 - 2*a*b*Sqrt[(1 - c*x)
/(1 + c*x)]*(1 + c*x))*ArcSech[c*x] - 3*b^2*(-a + b*Sqrt[(1 - c*x)/(1 + c
*x)]*(1 + c*x))*ArcSech[c*x]^2 + b^3*ArcSech[c*x]^3)/x)

```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.49 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.27, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$ , Rules used = {6839, 3042, 26, 3777, 3042, 3777, 26, 3042, 26, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^2} dx$$

$$\downarrow 6839$$

$$-c \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^3}{cx} d \operatorname{sech}^{-1}(cx)$$

$$\downarrow 3042$$

$$-c \int -i(a + b \operatorname{sech}^{-1}(cx))^3 \sin(i \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx)$$

$$\begin{aligned}
& \downarrow 26 \\
& ic \int (a + b \operatorname{sech}^{-1}(cx))^3 \sin(i \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) \\
& \downarrow 3777 \\
& ic \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))^3}{cx} - 3ib \int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{cx} d \operatorname{sech}^{-1}(cx) \right) \\
& \downarrow 3042 \\
& ic \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))^3}{cx} - 3ib \int (a + b \operatorname{sech}^{-1}(cx))^2 \sin\left(i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2}\right) d \operatorname{sech}^{-1}(cx) \right) \\
& \downarrow 3777 \\
& ic \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))^3}{cx} - 3ib \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{cx} - 2ib \int -\frac{i\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{cx} d \operatorname{sech}^{-1}(cx) \right) \right) \\
& \downarrow 26 \\
& ic \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))^3}{cx} - 3ib \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{cx} - 2b \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{cx} d \operatorname{sech}^{-1}(cx) \right) \right) \\
& \downarrow 3042 \\
& ic \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))^3}{cx} - 3ib \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{cx} - 2b \int -i(a + b \operatorname{sech}^{-1}(cx)) \sin(i \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) \right) \right) \\
& \downarrow 26 \\
& ic \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))^3}{cx} - 3ib \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{cx} + 2ib \int (a + b \operatorname{sech}^{-1}(cx)) \sin(i \operatorname{sech}^{-1}(cx)) d \operatorname{sech}^{-1}(cx) \right) \right) \\
& \downarrow 3777 \\
& ic \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))^3}{cx} - 3ib \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{cx} + 2ib \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))}{cx} - ib \int \frac{1}{cx} d \operatorname{sech}^{-1}(cx) \right) \right) \right)
\end{aligned}$$



↓ 3042

$$ic \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))^3}{cx} - 3ib \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{cx} + 2ib \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))}{cx} - ib \int \sin(i \operatorname{sech}^{-1}(cx)) dx \right) \right) \right)$$

↓ 3117

$$ic \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))^3}{cx} - 3ib \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{cx} + 2ib \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))}{cx} - \frac{ib \sqrt{\frac{1-cx}{cx+1}}(cx+1)}{cx} \right) \right) \right)$$

input `Int[(a + b*ArcSech[c*x])^3/x^2,x]`

output `I*c*((I*(a + b*ArcSech[c*x])^3)/(c*x) - (3*I)*b*((Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x])^2)/(c*x) + (2*I)*b*((-I)*b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(c*x) + (I*(a + b*ArcSech[c*x]))/(c*x)))`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 6839

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_.), x_Symbol] := Simp[
-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 224 vs. 2(98) = 196.

Time = 0.48 (sec) , antiderivative size = 225, normalized size of antiderivative = 2.21

method	result
parts	$-\frac{a^3}{x} + b^3 c \left( -\frac{\operatorname{arcsech}(cx)^3}{cx} + 3 \operatorname{arcsech}(cx)^2 \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} - \frac{6 \operatorname{arcsech}(cx)}{cx} + 6 \sqrt{-\frac{cx-1}{cx}} \right)$
derivativedivides	$c \left( -\frac{a^3}{cx} + b^3 \left( -\frac{\operatorname{arcsech}(cx)^3}{cx} + 3 \operatorname{arcsech}(cx)^2 \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} - \frac{6 \operatorname{arcsech}(cx)}{cx} + 6 \sqrt{-\frac{cx-1}{cx}} \right) \right)$
default	$c \left( -\frac{a^3}{cx} + b^3 \left( -\frac{\operatorname{arcsech}(cx)^3}{cx} + 3 \operatorname{arcsech}(cx)^2 \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} - \frac{6 \operatorname{arcsech}(cx)}{cx} + 6 \sqrt{-\frac{cx-1}{cx}} \right) \right)$

input

```
int((a+b*arcsech(c*x))^3/x^2,x,method=_RETURNVERBOSE)
```

output

```
-a^3/x+b^3*c*(-1/c/x*arcsech(c*x)^3+3*arcsech(c*x)^2*(-(c*x-1)/c/x)^(1/2)*
((c*x+1)/c/x)^(1/2)-6/c/x*arcsech(c*x)+6*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x
)^(1/2))+3*a*b^2*c*(-1/c/x*arcsech(c*x)^2+2*arcsech(c*x)*(-(c*x-1)/c/x)^(1
/2)*((c*x+1)/c/x)^(1/2)-2/c/x)+3*a^2*b*c*(-1/c/x*arcsech(c*x)+(-(c*x-1)/c
x)^(1/2)*((c*x+1)/c/x)^(1/2))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 228 vs. 2(98) = 196.

Time = 0.09 (sec) , antiderivative size = 228, normalized size of antiderivative = 2.24

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^2} dx =$$

$$\frac{b^3 \log \left( \frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx} \right)^3 - 3(a^2 b + 2 b^3) cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + a^3 + 6 ab^2 - 3 \left( b^3 cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - ab^2 \right) \log \left( \frac{cx}{\dots} \right)}{x}$$

input `integrate((a+b*arcsech(c*x))^3/x^2,x, algorithm="fricas")`

output `-(b^3*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x))^3 - 3*(a^2*b + 2*b^3)*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + a^3 + 6*a*b^2 - 3*(b^3*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - a*b^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x))^2 - 3*(2*a*b^2*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - a^2*b - 2*b^3)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)))/x`

## Sympy [F]

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^2} dx = \int \frac{(a + b \operatorname{asech}(cx))^3}{x^2} dx$$

input `integrate((a+b*asech(c*x))**3/x**2,x)`

output `Integral((a + b*asech(c*x))**3/x**2, x)`

## Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.41

$$\begin{aligned} & \int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^2} dx \\ &= -\frac{b^3 \operatorname{arsech}(cx)^3}{x} + 3 \left( c \sqrt{\frac{1}{c^2 x^2} - 1} - \frac{\operatorname{arsech}(cx)}{x} \right) a^2 b \\ &+ 6 \left( c \sqrt{\frac{1}{c^2 x^2} - 1} \operatorname{arsech}(cx) - \frac{1}{x} \right) ab^2 \\ &+ 3 \left( c \sqrt{\frac{1}{c^2 x^2} - 1} \operatorname{arsech}(cx)^2 + 2c \sqrt{\frac{1}{c^2 x^2} - 1} - \frac{2 \operatorname{arsech}(cx)}{x} \right) b^3 \\ &- \frac{3ab^2 \operatorname{arsech}(cx)^2}{x} - \frac{a^3}{x} \end{aligned}$$

input `integrate((a+b*arcsech(c*x))^3/x^2,x, algorithm="maxima")`

output `-b^3*arcsech(c*x)^3/x + 3*(c*sqrt(1/(c^2*x^2) - 1) - arcsech(c*x)/x)*a^2*b + 6*(c*sqrt(1/(c^2*x^2) - 1)*arcsech(c*x) - 1/x)*a*b^2 + 3*(c*sqrt(1/(c^2*x^2) - 1)*arcsech(c*x)^2 + 2*c*sqrt(1/(c^2*x^2) - 1) - 2*arcsech(c*x)/x)*b^3 - 3*a*b^2*arcsech(c*x)^2/x - a^3/x`

### Giac [F]

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3}{x^2} dx$$

input `integrate((a+b*arcsech(c*x))^3/x^2,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)^3/x^2, x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(\frac{1}{cx}))^3}{x^2} dx$$

input `int((a + b*acosh(1/(c*x)))^3/x^2,x)`

output `int((a + b*acosh(1/(c*x)))^3/x^2, x)`

**Reduce [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^2} dx$$

$$= \frac{3 \left( \int \frac{\operatorname{asech}(cx)}{x^2} dx \right) a^2 b x + \left( \int \frac{\operatorname{asech}(cx)^3}{x^2} dx \right) b^3 x + 3 \left( \int \frac{\operatorname{asech}(cx)^2}{x^2} dx \right) a b^2 x - a^3}{x}$$

input `int((a+b*asech(c*x))^3/x^2,x)`

output `(3*int(asech(c*x)/x**2,x)*a**2*b*x + int(asech(c*x)**3/x**2,x)*b**3*x + 3*int(asech(c*x)**2/x**2,x)*a*b**2*x - a**3)/x`

**3.48**  $\int \frac{(a+b\operatorname{sech}^{-1}(cx))^3}{x^3} dx$

Optimal result	429
Mathematica [A] (verified)	430
Rubi [A] (verified)	430
Maple [A] (verified)	433
Fricas [A] (verification not implemented)	434
Sympy [F]	435
Maxima [F]	435
Giac [F]	435
Mupad [F(-1)]	436
Reduce [F]	436

**Optimal result**

Integrand size = 14, antiderivative size = 163

$$\int \frac{(a + b\operatorname{sech}^{-1}(cx))^3}{x^3} dx = \frac{3b^3 \sqrt{\frac{1-cx}{1+cx}}(1+cx)}{8x^2} - \frac{3}{8}b^3c^2\operatorname{sech}^{-1}(cx) - \frac{3b^2(1-cx)(1+cx)(a+b\operatorname{sech}^{-1}(cx))}{4x^2} + \frac{3b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(a+b\operatorname{sech}^{-1}(cx))^2}{4x^2} - \frac{1}{4}c^2(a+b\operatorname{sech}^{-1}(cx))^3 - \frac{(1-cx)(1+cx)(a+b\operatorname{sech}^{-1}(cx))^3}{2x^2}$$

output

```
3/8*b^3*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/x^2-3/8*b^3*c^2*arcsech(c*x)-3/4*
b^2*(-c*x+1)*(c*x+1)*(a+b*arcsech(c*x))/x^2+3/4*b*((-c*x+1)/(c*x+1))^(1/2)
*(c*x+1)*(a+b*arcsech(c*x))^2/x^2-1/4*c^2*(a+b*arcsech(c*x))^3-1/2*(-c*x+1)
*(c*x+1)*(a+b*arcsech(c*x))^3/x^2
```

**Mathematica [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.50

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^3} dx$$

$$= \frac{-4a^3 - 6ab^2 + 3b(2a^2 + b^2) \sqrt{\frac{1-cx}{1+cx}}(1+cx) - 6b(2a^2 + b^2 - 2ab \sqrt{\frac{1-cx}{1+cx}}(1+cx)) \operatorname{sech}^{-1}(cx) + 6b^2 \left( b \sqrt{\frac{1-cx}{1+cx}}(1+cx) \right)}{8x^2}$$

input

Integrate[(a + b\*ArcSech[c\*x])^3/x^3,x]

output

```
(-4*a^3 - 6*a*b^2 + 3*b*(2*a^2 + b^2)*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)
- 6*b*(2*a^2 + b^2 - 2*a*b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))*ArcSech[c*
x] + 6*b^2*(b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x) + a*(-2 + c^2*x^2))*ArcS
ech[c*x]^2 + 2*b^3*(-2 + c^2*x^2)*ArcSech[c*x]^3 - 3*b*(2*a^2 + b^2)*c^2*x
^2*Log[x] + 3*b*(2*a^2 + b^2)*c^2*x^2*Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] +
c*x*Sqrt[(1 - c*x)/(1 + c*x)])]/(8*x^2)
```

**Rubi [A] (verified)**Time = 0.47 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.12, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.786$ , Rules used = {6839, 5969, 3042, 25, 3792, 17, 25, 3042, 25, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^3} dx$$

$$\downarrow \text{6839}$$

$$-c^2 \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^3}{c^2 x^2} d \operatorname{sech}^{-1}(cx)$$

$$\downarrow \text{5969}$$

$$-c^2 \left( \frac{(1-cx)(cx+1)(a+b\operatorname{sech}^{-1}(cx))^3}{2c^2x^2} - \frac{3}{2}b \int \frac{(1-cx)(cx+1)(a+b\operatorname{sech}^{-1}(cx))^2}{c^2x^2} d\operatorname{sech}^{-1}(cx) \right)$$

↓ 3042

$$-c^2 \left( \frac{(1-cx)(cx+1)(a+b\operatorname{sech}^{-1}(cx))^3}{2c^2x^2} - \frac{3}{2}b \int -(a+b\operatorname{sech}^{-1}(cx))^2 \sin(\operatorname{isech}^{-1}(cx))^2 d\operatorname{sech}^{-1}(cx) \right)$$

↓ 25

$$-c^2 \left( \frac{(1-cx)(cx+1)(a+b\operatorname{sech}^{-1}(cx))^3}{2c^2x^2} + \frac{3}{2}b \int (a+b\operatorname{sech}^{-1}(cx))^2 \sin(\operatorname{isech}^{-1}(cx))^2 d\operatorname{sech}^{-1}(cx) \right)$$

↓ 3792

$$-c^2 \left( \frac{3}{2}b \left( \frac{1}{2} \int (a+b\operatorname{sech}^{-1}(cx))^2 d\operatorname{sech}^{-1}(cx) + \frac{1}{2}b^2 \int -\frac{(1-cx)(cx+1)}{c^2x^2} d\operatorname{sech}^{-1}(cx) + \frac{b(1-cx)(cx+1)(a+b\operatorname{sech}^{-1}(cx))^3}{2c^2x^2} \right) \right)$$

↓ 17

$$-c^2 \left( \frac{3}{2}b \left( \frac{1}{2}b^2 \int -\frac{(1-cx)(cx+1)}{c^2x^2} d\operatorname{sech}^{-1}(cx) - \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a+b\operatorname{sech}^{-1}(cx))^2}{2c^2x^2} + \frac{b(1-cx)(cx+1)(a+b\operatorname{sech}^{-1}(cx))^3}{2c^2x^2} \right) \right)$$

↓ 25

$$-c^2 \left( \frac{3}{2}b \left( -\frac{1}{2}b^2 \int \frac{(1-cx)(cx+1)}{c^2x^2} d\operatorname{sech}^{-1}(cx) - \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a+b\operatorname{sech}^{-1}(cx))^2}{2c^2x^2} + \frac{b(1-cx)(cx+1)(a+b\operatorname{sech}^{-1}(cx))^3}{2c^2x^2} \right) \right)$$

↓ 3042

$$-c^2 \left( \frac{(1-cx)(cx+1)(a+b\operatorname{sech}^{-1}(cx))^3}{2c^2x^2} + \frac{3}{2}b \left( -\frac{1}{2}b^2 \int -\sin(\operatorname{isech}^{-1}(cx))^2 d\operatorname{sech}^{-1}(cx) - \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a+b\operatorname{sech}^{-1}(cx))^2}{2c^2x^2} \right) \right)$$

↓ 25

$$-c^2 \left( \frac{(1-cx)(cx+1)(a+b\operatorname{sech}^{-1}(cx))^3}{2c^2x^2} + \frac{3}{2}b \left( \frac{1}{2}b^2 \int \sin(\operatorname{isech}^{-1}(cx))^2 d\operatorname{sech}^{-1}(cx) - \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a+b\operatorname{sech}^{-1}(cx))^2}{2c^2x^2} \right) \right)$$



$$\begin{aligned}
 & \downarrow 3115 \\
 & -c^2 \left( \frac{3}{2} b \left( \frac{1}{2} b^2 \left( \frac{1}{2} \int 1 d \operatorname{sech}^{-1}(cx) - \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{2c^2x^2} \right) - \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a+b \operatorname{sech}^{-1}(cx))^2}{2c^2x^2} + \frac{b(1-cx)(cx+1)}{2c^2x^2} \right) \right. \\
 & \downarrow 24 \\
 & \left. -c^2 \left( \frac{3}{2} b \left( -\frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a+b \operatorname{sech}^{-1}(cx))^2}{2c^2x^2} + \frac{b(1-cx)(cx+1)(a+b \operatorname{sech}^{-1}(cx))}{2c^2x^2} + \frac{(a+b \operatorname{sech}^{-1}(cx))^3}{6b} \right) \right) \right.
 \end{aligned}$$

input `Int[(a + b*ArcSech[c*x])^3/x^3,x]`

output `-(c^2*((1 - c*x)*(1 + c*x)*(a + b*ArcSech[c*x])^3)/(2*c^2*x^2) + (3*b*((b^2*(-1/2*(Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(c^2*x^2) + ArcSech[c*x]/2)/2 + (b*(1 - c*x)*(1 + c*x)*(a + b*ArcSech[c*x]))/(2*c^2*x^2) - (Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x])^2)/(2*c^2*x^2) + (a + b*ArcSech[c*x])^3/(6*b)))/2))`

### Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3792 `Int[((c_.) + (d_.)*(x_))^(m_)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[d*m*(c + d*x)^(m - 1)*((b*Sin[e + f*x])^n/(f^2*n^2)), x] + (-Simp[b*(c + d*x)^m*cos[e + f*x]*((b*Sin[e + f*x])^(n - 1)/(f*n)), x] + Simp[b^2*((n - 1)/n) Int[(c + d*x)^m*(b*Sin[e + f*x])^(n - 2), x], x] - Simp[d^2*m*(m - 1)/(f^2*n^2) Int[(c + d*x)^(m - 2)*(b*Sin[e + f*x])^n, x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 1]`

rule 5969 `Int[Cosh[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_)*Sinh[(a_.) + (b_.)*(x_)]^(n_), x_Symbol] := Simp[(c + d*x)^m*(Sinh[a + b*x]^(n + 1)/(b*(n + 1))), x] - Simp[d*(m/(b*(n + 1))) Int[(c + d*x)^(m - 1)*Sinh[a + b*x]^(n + 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

rule 6839 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))^(n_)*(x_)^(m_), x_Symbol] := Simp[-(c^(m + 1))^( -1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])`

### Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.34

method	result
derivativedivides	$c^2 \left( -\frac{a^3}{2c^2x^2} + b^3 \left( -\frac{\cosh(2 \operatorname{arcsech}(cx)) \operatorname{arcsech}(cx)^3}{4} + \frac{3 \sinh(2 \operatorname{arcsech}(cx)) \operatorname{arcsech}(cx)^2}{8} - \frac{3 \cosh(2 \operatorname{arcsech}(cx)) \operatorname{arcsech}(cx)}{8} \right) \right)$
default	$c^2 \left( -\frac{a^3}{2c^2x^2} + b^3 \left( -\frac{\cosh(2 \operatorname{arcsech}(cx)) \operatorname{arcsech}(cx)^3}{4} + \frac{3 \sinh(2 \operatorname{arcsech}(cx)) \operatorname{arcsech}(cx)^2}{8} - \frac{3 \cosh(2 \operatorname{arcsech}(cx)) \operatorname{arcsech}(cx)}{8} \right) \right)$
parts	$-\frac{a^3}{2x^2} + b^3 c^2 \left( -\frac{\cosh(2 \operatorname{arcsech}(cx)) \operatorname{arcsech}(cx)^3}{4} + \frac{3 \sinh(2 \operatorname{arcsech}(cx)) \operatorname{arcsech}(cx)^2}{8} - \frac{3 \cosh(2 \operatorname{arcsech}(cx)) \operatorname{arcsech}(cx)}{8} \right)$

input `int((a+b*arcsech(c*x))^3/x^3,x,method=_RETURNVERBOSE)`

output 
$$c^2*(-1/2*a^3/c^2/x^2+b^3*(-1/4*cosh(2*arcsech(c*x))*arcsech(c*x)^3+3/8*sinh(2*arcsech(c*x))*arcsech(c*x)^2-3/8*cosh(2*arcsech(c*x))*arcsech(c*x)+3/16*sinh(2*arcsech(c*x)))+3*a*b^2*(-1/4*cosh(2*arcsech(c*x))*arcsech(c*x)^2+1/4*sinh(2*arcsech(c*x))*arcsech(c*x)-1/8*cosh(2*arcsech(c*x))+3*a^2*b*(-1/2/c^2/x^2*arcsech(c*x)+1/4*(-(c*x-1)/c/x)^(1/2)/c/x*((c*x+1)/c/x)^(1/2))*(arctanh(1/(-c^2*x^2+1)^(1/2))*c^2*x^2+(-c^2*x^2+1)^(1/2))/(-c^2*x^2+1)^(1/2))$$

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.66

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^3} dx$$

$$= \frac{2(b^3 c^2 x^2 - 2b^3) \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx}\right)^3 + 3(2a^2 b + b^3) cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - 4a^3 - 6ab^2 + 6(ab^2 c^2 x^2 + b^3 cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}})}{x^3}$$

input `integrate((a+b*arcsech(c*x))^3/x^3,x, algorithm="fricas")`

output 
$$\frac{1}{8}*(2*(b^3*c^2*x^2 - 2*b^3)*\log((c*x*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} + 1)/(c*x))^3 + 3*(2*a^2*b + b^3)*c*x*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} - 4*a^3 - 6*a*b^2 + 6*(a*b^2*c^2*x^2 + b^3*c*x*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} - 2*a*b^2)*\log((c*x*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} + 1)/(c*x))^2 + 3*(4*a*b^2*c*x*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} + (2*a^2*b + b^3)*c^2*x^2 - 4*a^2*b - 2*b^3)*\log((c*x*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} + 1)/(c*x)))/x^2$$

**Sympy [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^3} dx = \int \frac{(a + b \operatorname{arsech}(cx))^3}{x^3} dx$$

input `integrate((a+b*asech(c*x))**3/x**3,x)`

output `Integral((a + b*asech(c*x))**3/x**3, x)`

**Maxima [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^3} dx = \int \frac{(b \operatorname{arsech}(cx) + a)^3}{x^3} dx$$

input `integrate((a+b*arcsech(c*x))^3/x^3,x, algorithm="maxima")`

output `-3/8*a^2*b*((2*c^4*x*sqrt(1/(c^2*x^2) - 1)/(c^2*x^2*(1/(c^2*x^2) - 1) - 1) - c^3*log(c*x*sqrt(1/(c^2*x^2) - 1) + 1) + c^3*log(c*x*sqrt(1/(c^2*x^2) - 1) - 1))/c + 4*arcsech(c*x)/x^2) - 1/2*a^3/x^2 + integrate(b^3*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^3/x^3 + 3*a*b^2*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^2/x^3, x)`

**Giac [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^3} dx = \int \frac{(b \operatorname{arsech}(cx) + a)^3}{x^3} dx$$

input `integrate((a+b*arcsech(c*x))^3/x^3,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)^3/x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(\frac{1}{cx}))^3}{x^3} dx$$

input `int((a + b*acosh(1/(c*x)))^3/x^3,x)`output `int((a + b*acosh(1/(c*x)))^3/x^3, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^3} dx$$

$$= \frac{6 \left( \int \frac{\operatorname{asech}(cx)}{x^3} dx \right) a^2 b x^2 + 2 \left( \int \frac{\operatorname{asech}(cx)^3}{x^3} dx \right) b^3 x^2 + 6 \left( \int \frac{\operatorname{asech}(cx)^2}{x^3} dx \right) a b^2 x^2 - a^3}{2x^2}$$

input `int((a+b*asech(c*x))^3/x^3,x)`output `(6*int(asech(c*x)/x**3,x)*a**2*b*x**2 + 2*int(asech(c*x)**3/x**3,x)*b**3*x**2 + 6*int(asech(c*x)**2/x**3,x)*a*b**2*x**2 - a**3)/(2*x**2)`

**3.49** 
$$\int \frac{(a+b\operatorname{sech}^{-1}(cx))^3}{x^4} dx$$

Optimal result . . . . .	437
Mathematica [A] (verified) . . . . .	438
Rubi [C] (verified) . . . . .	438
Maple [B] (verified) . . . . .	442
Fricas [A] (verification not implemented) . . . . .	443
Sympy [F] . . . . .	444
Maxima [F] . . . . .	444
Giac [F] . . . . .	444
Mupad [F(-1)] . . . . .	445
Reduce [F] . . . . .	445

**Optimal result**

Integrand size = 14, antiderivative size = 213

$$\int \frac{(a + b\operatorname{sech}^{-1}(cx))^3}{x^4} dx = \frac{14b^3c^2\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{9x} + \frac{2b^3\left(\frac{1-cx}{1+cx}\right)^{3/2}(1+cx)^3}{27x^3} - \frac{2b^2(a + b\operatorname{sech}^{-1}(cx))}{9x^3} - \frac{4b^2c^2(a + b\operatorname{sech}^{-1}(cx))}{3x} + \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(a + b\operatorname{sech}^{-1}(cx))^2}{3x^3} + \frac{2bc^2\sqrt{\frac{1-cx}{1+cx}}(1+cx)(a + b\operatorname{sech}^{-1}(cx))^2}{3x} - \frac{(a + b\operatorname{sech}^{-1}(cx))^3}{3x^3}$$

output

```
14/9*b^3*c^2*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/x+2/27*b^3*((-c*x+1)/(c*x+1))^(3/2)*(c*x+1)^3/x^3-2/9*b^2*(a+b*arcsech(c*x))/x^3-4/3*b^2*c^2*(a+b*arcsech(c*x))/x+1/3*b*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+b*arcsech(c*x))^2/x^3+2/3*b*c^2*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*(a+b*arcsech(c*x))^2/x-1/3*(a+b*arcsech(c*x))^3/x^3
```

**Mathematica [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 256, normalized size of antiderivative = 1.20

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^4} dx$$

$$= \frac{-9a^3 - 6ab^2(1 + 6c^2x^2) + 9a^2b\sqrt{\frac{1-cx}{1+cx}}(1 + cx + 2c^2x^2 + 2c^3x^3) + 2b^3\sqrt{\frac{1-cx}{1+cx}}(1 + cx + 20c^2x^2 + 20c^3x^3)}{27x^3}$$

input `Integrate[(a + b*ArcSech[c*x])^3/x^4,x]`

output

```
(-9*a^3 - 6*a*b^2*(1 + 6*c^2*x^2) + 9*a^2*b*sqrt[(1 - c*x)/(1 + c*x)]*(1 +
c*x + 2*c^2*x^2 + 2*c^3*x^3) + 2*b^3*sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x +
20*c^2*x^2 + 20*c^3*x^3) - 3*b*(9*a^2 + 2*b^2*(1 + 6*c^2*x^2) - 6*a*b*sqrt
t[(1 - c*x)/(1 + c*x)]*(1 + c*x + 2*c^2*x^2 + 2*c^3*x^3))*ArcSech[c*x] + 9
*b^2*(-3*a + b*sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x + 2*c^2*x^2 + 2*c^3*x^3)
)*ArcSech[c*x]^2 - 9*b^3*ArcSech[c*x]^3)/(27*x^3)
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.78 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.29, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {6839, 5970, 3042, 3792, 3042, 3113, 2009, 3777, 26, 3042, 26, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^4} dx$$

$$\downarrow \text{6839}$$

$$-c^3 \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^3}{c^3x^3} d \operatorname{sech}^{-1}(cx)$$

$$\downarrow \text{5970}$$

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{3c^3 x^3} - b \int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{c^3 x^3} d \operatorname{sech}^{-1}(cx) \right)$$

↓ 3042

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{3c^3 x^3} - b \int (a + b \operatorname{sech}^{-1}(cx))^2 \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^3 d \operatorname{sech}^{-1}(cx) \right)$$

↓ 3792

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{3c^3 x^3} - b \left( \frac{2}{3} \int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{cx} d \operatorname{sech}^{-1}(cx) + \frac{2}{9} b^2 \int \frac{1}{c^3 x^3} d \operatorname{sech}^{-1}(cx) - \frac{2b(a + b \operatorname{sech}^{-1}(cx))}{9c^3 x^3} \right) \right)$$

↓ 3042

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{3c^3 x^3} - b \left( \frac{2}{3} \int (a + b \operatorname{sech}^{-1}(cx))^2 \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right) d \operatorname{sech}^{-1}(cx) + \frac{2}{9} b^2 \int \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^3 d \operatorname{sech}^{-1}(cx) \right) \right)$$

↓ 3113

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{3c^3 x^3} - b \left( \frac{2}{3} \int (a + b \operatorname{sech}^{-1}(cx))^2 \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right) d \operatorname{sech}^{-1}(cx) + \frac{2}{9} i b^2 \int \left( \frac{1 - cx}{c^2} \right) d \operatorname{sech}^{-1}(cx) \right) \right)$$

↓ 2009

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{3c^3 x^3} - b \left( \frac{2}{3} \int (a + b \operatorname{sech}^{-1}(cx))^2 \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right) d \operatorname{sech}^{-1}(cx) - \frac{2b(a + b \operatorname{sech}^{-1}(cx))}{9c^3 x^3} \right) \right)$$

↓ 3777

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{3c^3 x^3} - b \left( \frac{2}{3} \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a+b \operatorname{sech}^{-1}(cx))^2}{cx} - 2ib \int -\frac{i \sqrt{\frac{1-cx}{cx+1}}(cx+1)(a+b \operatorname{sech}^{-1}(cx))^2}{cx} d \operatorname{sech}^{-1}(cx) \right) \right) \right)$$

↓ 26



$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{3c^3 x^3} - b \left( \frac{2}{3} \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{cx} - 2b \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{cx} \right) \right) \right)$$

↓ 3042

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{3c^3 x^3} - b \left( \frac{2}{3} \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{cx} - 2b \int -i(a + b \operatorname{sech}^{-1}(cx)) \sin(i \operatorname{sech}^{-1}(cx)) \right) \right) \right)$$

↓ 26

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{3c^3 x^3} - b \left( \frac{2}{3} \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{cx} + 2ib \int (a + b \operatorname{sech}^{-1}(cx)) \sin(i \operatorname{sech}^{-1}(cx)) \right) \right) \right)$$

↓ 3777

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{3c^3 x^3} - b \left( \frac{2}{3} \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{cx} + 2ib \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))}{cx} - ib \int \frac{1}{cx} d \operatorname{sech}^{-1}(cx) \right) \right) \right) \right)$$

↓ 3042

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{3c^3 x^3} - b \left( \frac{2}{3} \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{cx} + 2ib \left( \frac{i(a + b \operatorname{sech}^{-1}(cx))}{cx} - ib \int \sin(i \operatorname{sech}^{-1}(cx)) \right) \right) \right) \right)$$

↓ 3117

$$-c^3 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{3c^3 x^3} - b \left( -\frac{2b(a + b \operatorname{sech}^{-1}(cx))}{9c^3 x^3} + \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{3c^3 x^3} + \frac{2}{3} \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))}{cx} \right) \right) \right)$$

input

```
Int[(a + b*ArcSech[c*x])^3/x^4,x]
```

output

```

-(c^3*((a + b*ArcSech[c*x])^3/(3*c^3*x^3) - b*((2*I)/9)*b^2*((-I)*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(c*x) - ((I/3)*((1 - c*x)/(1 + c*x))^(3/2)*(1 + c*x)^3)/(c^3*x^3)) - (2*b*(a + b*ArcSech[c*x]))/(9*c^3*x^3) + (Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x])^2)/(3*c^3*x^3) + (2*(Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x])^2)/(c*x) + (2*I)*b*(((I)*b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(c*x) + (I*(a + b*ArcSech[c*x]))/(c*x))))/3)))

```

**Defintions of rubi rules used**

rule 26

```

Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]

```

rule 2009

```

Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]

```

rule 3042

```

Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]

```

rule 3113

```

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp and[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

```

rule 3117

```

Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]

```

rule 3777

```

Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

```

rule 3792

```
Int[((c_.) + (d_.)*(x_)^(m_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol]
  := Simp[d*m*(c + d*x)^(m - 1)*((b*Sine + f*x)^(n)/(f^2*n^2)), x] + (-Simp
  p[b*(c + d*x)^m*cos[e + f*x]*((b*Sine + f*x)^(n - 1)/(f*n)), x] + Simp[b^
  2*((n - 1)/n) Int[(c + d*x)^m*(b*Sine + f*x)^(n - 2), x], x] - Simp[d^2
  *m*((m - 1)/(f^2*n^2)) Int[(c + d*x)^(m - 2)*(b*Sine + f*x)^n, x], x])
  /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 1]
```

rule 5970

```
Int[Cosh[(a_.) + (b_.)*(x_)^(n_.)]*((c_.) + (d_.)*(x_)^(m_.))*Sinh[(a_.) +
  (b_.)*(x_)], x_Symbol] := Simp[(c + d*x)^m*(Cosh[a + b*x]^(n + 1)/(b*(n + 1
 ))), x] - Simp[d*(m/(b*(n + 1))) Int[(c + d*x)^(m - 1)*Cosh[a + b*x]^(n +
  1), x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

rule 6839

```
Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))^(n_)*(x_)^(m_), x_Symbol] := Simp[
  -(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
  rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
  tQ[n, 0] || LtQ[m, -1])
```

## Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 386 vs.  $2(191) = 382$ .

Time = 0.75 (sec) , antiderivative size = 387, normalized size of antiderivative = 1.82

method	result
derivativedivides	$c^3 \left( -\frac{a^3}{3c^3x^3} + b^3 \left( -\frac{\operatorname{arcsech}(cx)^3}{3c^3x^3} + \frac{2 \operatorname{arcsech}(cx)^2 \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}}{3} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \operatorname{arcsech}(cx)^2}{3c^2x^2} - \frac{4 \operatorname{arcsech}(cx)}{3c^2x^2} \right) \right)$
default	$c^3 \left( -\frac{a^3}{3c^3x^3} + b^3 \left( -\frac{\operatorname{arcsech}(cx)^3}{3c^3x^3} + \frac{2 \operatorname{arcsech}(cx)^2 \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}}{3} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \operatorname{arcsech}(cx)^2}{3c^2x^2} - \frac{4 \operatorname{arcsech}(cx)}{3c^2x^2} \right) \right)$
parts	$-\frac{a^3}{3x^3} + b^3 c^3 \left( -\frac{\operatorname{arcsech}(cx)^3}{3c^3x^3} + \frac{2 \operatorname{arcsech}(cx)^2 \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}}{3} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \operatorname{arcsech}(cx)^2}{3c^2x^2} - \frac{4 \operatorname{arcsech}(cx)}{3c^2x^2} \right)$

input

```
int((a+b*arcsech(c*x))^3/x^4,x,method=_RETURNVERBOSE)
```

output

```
c^3*(-1/3*a^3/c^3/x^3+b^3*(-1/3/c^3/x^3*arcsech(c*x)^3+2/3*arcsech(c*x)^2*
(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)+1/3*(-(c*x-1)/c/x)^(1/2)*((c*x+1)
/c/x)^(1/2)/c^2/x^2*arcsech(c*x)^2-4/3/c/x*arcsech(c*x)+40/27*(-(c*x-1)/c/
x)^(1/2)*((c*x+1)/c/x)^(1/2)-2/9/c^3/x^3*arcsech(c*x)+2/27*(-(c*x-1)/c/x)^(
1/2)*((c*x+1)/c/x)^(1/2)/c^2/x^2)+3*a*b^2*(-1/3/c^3/x^3*arcsech(c*x)^2+4/
9*arcsech(c*x)*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)+2/9*(-(c*x-1)/c/x)
^(1/2)*((c*x+1)/c/x)^(1/2)/c^2/x^2*arcsech(c*x)-4/9/c/x-2/27/c^3/x^3)+3*a^
2*b*(-1/3/c^3/x^3*arcsech(c*x)+1/9*(-(c*x-1)/c/x)^(1/2)/c^2/x^2*((c*x+1)/c
/x)^(1/2)*(2*c^2*x^2+1)))
```

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.43

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^4} dx =$$

$$\frac{36 ab^2 c^2 x^2 + 9 b^3 \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx}\right)^3 + 9 a^3 + 6 ab^2 + 9 \left(3 ab^2 - (2 b^3 c^3 x^3 + b^3 cx) \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}}\right) \log\left(\right)}{x^3}$$

input

```
integrate((a+b*arcsech(c*x))^3/x^4,x, algorithm="fricas")
```

output

```
-1/27*(36*a*b^2*c^2*x^2 + 9*b^3*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) +
1)/(c*x))^3 + 9*a^3 + 6*a*b^2 + 9*(3*a*b^2 - (2*b^3*c^3*x^3 + b^3*c*x)*sqr
t(-(c^2*x^2 - 1)/(c^2*x^2)))*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/
(c*x))^2 + 3*(12*b^3*c^2*x^2 + 9*a^2*b + 2*b^3 - 6*(2*a*b^2*c^3*x^3 + a*b^
2*c*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x
^2)) + 1)/(c*x)) - (2*(9*a^2*b + 20*b^3)*c^3*x^3 + (9*a^2*b + 2*b^3)*c*x)*
sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/x^3
```

**Sympy [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^4} dx = \int \frac{(a + b \operatorname{arsech}(cx))^3}{x^4} dx$$

input `integrate((a+b*asech(c*x))**3/x**4,x)`

output `Integral((a + b*asech(c*x))**3/x**4, x)`

**Maxima [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^4} dx = \int \frac{(b \operatorname{arsech}(cx) + a)^3}{x^4} dx$$

input `integrate((a+b*arcsech(c*x))^3/x^4,x, algorithm="maxima")`

output `1/3*a^2*b*((c^4*(1/(c^2*x^2) - 1)^(3/2) + 3*c^4*sqrt(1/(c^2*x^2) - 1))/c - 3*arcsech(c*x)/x^3) - 1/3*a^3/x^3 + integrate(b^3*log(sqrt(1/(c*x) + 1))*sqrt(1/(c*x) - 1) + 1/(c*x))^3/x^4 + 3*a*b^2*log(sqrt(1/(c*x) + 1))*sqrt(1/(c*x) - 1) + 1/(c*x))^2/x^4, x)`

**Giac [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^4} dx = \int \frac{(b \operatorname{arsech}(cx) + a)^3}{x^4} dx$$

input `integrate((a+b*arcsech(c*x))^3/x^4,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)^3/x^4, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(\frac{1}{cx}))^3}{x^4} dx$$

input `int((a + b*acosh(1/(c*x)))^3/x^4,x)`output `int((a + b*acosh(1/(c*x)))^3/x^4, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^4} dx$$

$$= \frac{9 \left( \int \frac{\operatorname{asech}(cx)}{x^4} dx \right) a^2 b x^3 + 3 \left( \int \frac{\operatorname{asech}(cx)^3}{x^4} dx \right) b^3 x^3 + 9 \left( \int \frac{\operatorname{asech}(cx)^2}{x^4} dx \right) a b^2 x^3 - a^3}{3x^3}$$

input `int((a+b*asech(c*x))^3/x^4,x)`output `(9*int(asech(c*x)/x**4,x)*a**2*b*x**3 + 3*int(asech(c*x)**3/x**4,x)*b**3*x**3 + 9*int(asech(c*x)**2/x**4,x)*a*b**2*x**3 - a**3)/(3*x**3)`

**3.50**  $\int \frac{(a+b\operatorname{sech}^{-1}(cx))^3}{x^5} dx$

Optimal result	446
Mathematica [A] (verified)	447
Rubi [A] (verified)	447
Maple [B] (verified)	451
Fricas [A] (verification not implemented)	451
Sympy [F]	452
Maxima [F]	452
Giac [F]	453
Mupad [F(-1)]	453
Reduce [F]	454

**Optimal result**

Integrand size = 14, antiderivative size = 242

$$\int \frac{(a + b\operatorname{sech}^{-1}(cx))^3}{x^5} dx = \frac{3b^3 \sqrt{\frac{1-cx}{1+cx}}(1+cx)}{128x^4} + \frac{45b^3 c^2 \sqrt{\frac{1-cx}{1+cx}}(1+cx)}{256x^2}$$

$$+ \frac{45}{256} b^3 c^4 \operatorname{sech}^{-1}(cx) - \frac{3b^2 (a + b\operatorname{sech}^{-1}(cx))}{32x^4}$$

$$- \frac{9b^2 c^2 (a + b\operatorname{sech}^{-1}(cx))}{32x^2}$$

$$+ \frac{3b \sqrt{\frac{1-cx}{1+cx}}(1+cx) (a + b\operatorname{sech}^{-1}(cx))^2}{16x^4}$$

$$+ \frac{9bc^2 \sqrt{\frac{1-cx}{1+cx}}(1+cx) (a + b\operatorname{sech}^{-1}(cx))^2}{32x^2}$$

$$+ \frac{3}{32} c^4 (a + b\operatorname{sech}^{-1}(cx))^3 - \frac{(a + b\operatorname{sech}^{-1}(cx))^3}{4x^4}$$

output

```
3/128*b^3*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/x^4+45/256*b^3*c^2*((-c*x+1)/(c
*x+1))^(1/2)*(c*x+1)/x^2+45/256*b^3*c^4*arcsech(c*x)-3/32*b^2*(a+b*arcsech
(c*x))/x^4-9/32*b^2*c^2*(a+b*arcsech(c*x))/x^2+3/16*b*((-c*x+1)/(c*x+1))^(
1/2)*(c*x+1)*(a+b*arcsech(c*x))^2/x^4+9/32*b*c^2*((-c*x+1)/(c*x+1))^(1/2)*
(c*x+1)*(a+b*arcsech(c*x))^2/x^2+3/32*c^4*(a+b*arcsech(c*x))^3-1/4*(a+b*ar
csech(c*x))^3/x^4
```

**Mathematica [A] (verified)**

Time = 0.44 (sec) , antiderivative size = 332, normalized size of antiderivative = 1.37

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^5} dx$$

$$= \frac{-8a(8a^2 + 3b^2) - 72ab^2c^2x^2 + 3b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(8a^2(2+3c^2x^2) + b^2(2+15c^2x^2)) - 24b(8a^2 + b^2(1 +$$

input `Integrate[(a + b*ArcSech[c*x])^3/x^5,x]`

output

```
(-8*a*(8*a^2 + 3*b^2) - 72*a*b^2*c^2*x^2 + 3*b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(8*a^2*(2 + 3*c^2*x^2) + b^2*(2 + 15*c^2*x^2)) - 24*b*(8*a^2 + b^2*(1 + 3*c^2*x^2) - 2*a*b*Sqrt[(1 - c*x)/(1 + c*x)]*(2 + 2*c*x + 3*c^2*x^2 + 3*c^3*x^3))*ArcSech[c*x] + 24*b^2*(b*Sqrt[(1 - c*x)/(1 + c*x)]*(2 + 2*c*x + 3*c^2*x^2 + 3*c^3*x^3) + a*(-8 + 3*c^4*x^4))*ArcSech[c*x]^2 + 8*b^3*(-8 + 3*c^4*x^4)*ArcSech[c*x]^3 - 9*b*(8*a^2 + 5*b^2)*c^4*x^4*Log[x] + 9*b*(8*a^2 + 5*b^2)*c^4*x^4*Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]]/(256*x^4)
```

**Rubi [A] (verified)**

Time = 0.83 (sec) , antiderivative size = 314, normalized size of antiderivative = 1.30, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {6839, 5970, 3042, 3792, 3042, 3115, 3042, 3115, 24, 3792, 17, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^5} dx$$

$$\downarrow \text{6839}$$

$$-c^4 \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^3}{c^4 x^4} d \operatorname{sech}^{-1}(cx)$$

$$\downarrow \text{5970}$$



$$-c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{4c^4 x^4} - \frac{3}{4} b \int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{c^4 x^4} d \operatorname{sech}^{-1}(cx) \right)$$

↓ 3042

$$-c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{4c^4 x^4} - \frac{3}{4} b \int (a + b \operatorname{sech}^{-1}(cx))^2 \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^4 d \operatorname{sech}^{-1}(cx) \right)$$

↓ 3792

$$-c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{4c^4 x^4} - \frac{3}{4} b \left( \frac{3}{4} \int \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{c^2 x^2} d \operatorname{sech}^{-1}(cx) + \frac{1}{8} b^2 \int \frac{1}{c^4 x^4} d \operatorname{sech}^{-1}(cx) - \frac{b(a + b \operatorname{sech}^{-1}(cx))}{8c^4 x^4} \right) \right)$$

↓ 3042

$$-c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{4c^4 x^4} - \frac{3}{4} b \left( \frac{3}{4} \int (a + b \operatorname{sech}^{-1}(cx))^2 \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^2 d \operatorname{sech}^{-1}(cx) + \frac{1}{8} b^2 \int \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^4 d \operatorname{sech}^{-1}(cx) \right) \right)$$

↓ 3115

$$-c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{4c^4 x^4} - \frac{3}{4} b \left( \frac{3}{4} \int (a + b \operatorname{sech}^{-1}(cx))^2 \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^2 d \operatorname{sech}^{-1}(cx) + \frac{1}{8} b^2 \left( \frac{3}{4} \int \frac{1}{c^2 x^2} d \operatorname{sech}^{-1}(cx) - \frac{b(a + b \operatorname{sech}^{-1}(cx))}{8c^4 x^4} \right) \right) \right)$$

↓ 3042

$$-c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{4c^4 x^4} - \frac{3}{4} b \left( \frac{3}{4} \int (a + b \operatorname{sech}^{-1}(cx))^2 \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^2 d \operatorname{sech}^{-1}(cx) + \frac{1}{8} b^2 \left( \frac{\sqrt{1-cx}}{cx+1} \int \frac{1}{c^4 x^4} d \operatorname{sech}^{-1}(cx) - \frac{b(a + b \operatorname{sech}^{-1}(cx))}{8c^4 x^4} \right) \right) \right)$$

↓ 3115

$$-c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{4c^4 x^4} - \frac{3}{4} b \left( \frac{3}{4} \int (a + b \operatorname{sech}^{-1}(cx))^2 \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^2 d \operatorname{sech}^{-1}(cx) + \frac{1}{8} b^2 \left( \frac{3}{4} \left( \frac{1}{2} \int \frac{1}{c^2 x^2} d \operatorname{sech}^{-1}(cx) - \frac{b(a + b \operatorname{sech}^{-1}(cx))}{8c^4 x^4} \right) \right) \right) \right)$$

↓ 24

$$-c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{4c^4 x^4} - \frac{3}{4} b \left( \frac{3}{4} \int (a + b \operatorname{sech}^{-1}(cx))^2 \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^2 d \operatorname{sech}^{-1}(cx) - \frac{b(a + b \operatorname{sech}^{-1}(cx))}{8c^4 x^4} \right) \right)$$

↓ 3792

$$-c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{4c^4 x^4} - \frac{3}{4} b \left( \frac{3}{4} \left( \frac{1}{2} \int (a + b \operatorname{sech}^{-1}(cx))^2 d \operatorname{sech}^{-1}(cx) + \frac{1}{2} b^2 \int \frac{1}{c^2 x^2} d \operatorname{sech}^{-1}(cx) - \frac{b(a + b \operatorname{sech}^{-1}(cx))}{2c^2 x^2} \right) \right) \right)$$

↓ 17

$$-c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{4c^4 x^4} - \frac{3}{4} b \left( \frac{3}{4} \left( \frac{1}{2} b^2 \int \frac{1}{c^2 x^2} d \operatorname{sech}^{-1}(cx) + \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{2c^2 x^2} - \frac{b(a + b \operatorname{sech}^{-1}(cx))}{2c^2 x^2} \right) \right) \right)$$

↓ 3042

$$-c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{4c^4 x^4} - \frac{3}{4} b \left( \frac{3}{4} \left( \frac{1}{2} b^2 \int \sin \left( i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)^2 d \operatorname{sech}^{-1}(cx) + \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{2c^2 x^2} \right) \right) \right)$$

↓ 3115

$$-c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{4c^4 x^4} - \frac{3}{4} b \left( \frac{3}{4} \left( \frac{1}{2} b^2 \left( \frac{1}{2} \int 1 d \operatorname{sech}^{-1}(cx) + \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{2c^2 x^2} \right) \right) + \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{2c^2 x^2} \right) \right)$$

↓ 24

$$-c^4 \left( \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{4c^4 x^4} - \frac{3}{4} b \left( \frac{3}{4} \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)(a + b \operatorname{sech}^{-1}(cx))^2}{2c^2 x^2} - \frac{b(a + b \operatorname{sech}^{-1}(cx))}{2c^2 x^2} + \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{6b} \right) \right) \right)$$

input `Int[(a + b*ArcSech[c*x])^3/x^5,x]`

output `-(c^4*((a + b*ArcSech[c*x])^3/(4*c^4*x^4) - (3*b*((b^2*((Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(4*c^4*x^4) + (3*((Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(2*c^2*x^2) + ArcSech[c*x]/2))/4))/8 - (b*(a + b*ArcSech[c*x]))/(8*c^4*x^4) + (Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x])^2)/(4*c^4*x^4) + (3*((b^2*((Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(2*c^2*x^2) + ArcSech[c*x]/2))/2 - (b*(a + b*ArcSech[c*x]))/(2*c^2*x^2) + (Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcSech[c*x])^2)/(2*c^2*x^2) + (a + b*ArcSech[c*x])^3/(6*b)))/4))/4)`

## Definitions of rubi rules used

- rule 17  $\text{Int}[(c_.)*((a_.) + (b_.)*(x_))^{(m_.)}, x\_Symbol] \rightarrow \text{Simp}[c*((a + b*x)^{(m + 1})/(b*(m + 1))), x] \text{ /; FreeQ}\{a, b, c, m\}, x] \ \&\& \ \text{NeQ}[m, -1]$
- rule 24  $\text{Int}[a_, x\_Symbol] \rightarrow \text{Simp}[a*x, x] \text{ /; FreeQ}[a, x]$
- rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$
- rule 3115  $\text{Int}[(b_.)*\sin[(c_.) + (d_.)*(x_)]^{(n_.)}, x\_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*((b*\sin[c + d*x])^{(n - 1)})/(d*n), x] + \text{Simp}[b^2*((n - 1)/n) \text{ Int}[(b*\sin[c + d*x])^{(n - 2)}, x], x] \text{ /; FreeQ}\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$
- rule 3792  $\text{Int}[(c_.) + (d_.)*(x_)]^{(m_.)*((b_.)*\sin[(e_.) + (f_.)*(x_)]^{(n_.)}, x\_Symbol] \rightarrow \text{Simp}[d*m*(c + d*x)^{(m - 1)}*((b*\sin[e + f*x])^n/(f^2*n^2)), x] + (-\text{Simp}[b*(c + d*x)^m*\text{Cos}[e + f*x]*((b*\sin[e + f*x])^{(n - 1)})/(f*n), x] + \text{Simp}[b^2*((n - 1)/n) \text{ Int}[(c + d*x)^m*(b*\sin[e + f*x])^{(n - 2)}, x], x] - \text{Simp}[d^2*m*((m - 1)/(f^2*n^2)) \text{ Int}[(c + d*x)^{(m - 2)}*(b*\sin[e + f*x])^n, x], x]) \text{ /; FreeQ}\{b, c, d, e, f\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{GtQ}[m, 1]$
- rule 5970  $\text{Int}[\text{Cosh}[(a_.) + (b_.)*(x_)]^{(n_.)*((c_.) + (d_.)*(x_))^{(m_.)*\text{Sinh}[(a_.) + (b_.)*(x_)]}, x\_Symbol] \rightarrow \text{Simp}[(c + d*x)^m*(\text{Cosh}[a + b*x]^{(n + 1)})/(b*(n + 1)), x] - \text{Simp}[d*(m/(b*(n + 1))) \text{ Int}[(c + d*x)^{(m - 1)}*\text{Cosh}[a + b*x]^{(n + 1)}, x], x] \text{ /; FreeQ}\{a, b, c, d, n\}, x] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{NeQ}[n, -1]$
- rule 6839  $\text{Int}[(a_.) + \text{ArcSech}[(c_.)*(x_)]*(b_.)]^{(n_.)*(x_)^{(m_.)}, x\_Symbol] \rightarrow \text{Simp}[-(c^{(m + 1)})^{(-1)} \text{ Subst}[\text{Int}[(a + b*x)^n*\text{Sech}[x]^{(m + 1)}*\text{Tanh}[x], x], x, \text{ArcSech}[c*x]], x] \text{ /; FreeQ}\{a, b, c\}, x] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ (\text{GtQ}[n, 0] \ || \ \text{LtQ}[m, -1])$

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 484 vs. 2(216) = 432.

Time = 0.72 (sec) , antiderivative size = 485, normalized size of antiderivative = 2.00

method	result
derivativedivides	$c^4 \left( -\frac{a^3}{4c^4x^4} + b^3 \left( -\frac{\operatorname{arcsech}(cx)^3}{4c^4x^4} + \frac{3\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}\operatorname{arcsech}(cx)^2}{16c^3x^3} + \frac{9\operatorname{arcsech}(cx)^2\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}}{32cx} + \dots \right) \right)$
default	$c^4 \left( -\frac{a^3}{4c^4x^4} + b^3 \left( -\frac{\operatorname{arcsech}(cx)^3}{4c^4x^4} + \frac{3\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}\operatorname{arcsech}(cx)^2}{16c^3x^3} + \frac{9\operatorname{arcsech}(cx)^2\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}}{32cx} + \dots \right) \right)$
parts	$-\frac{a^3}{4x^4} + b^3c^4 \left( -\frac{\operatorname{arcsech}(cx)^3}{4c^4x^4} + \frac{3\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}\operatorname{arcsech}(cx)^2}{16c^3x^3} + \frac{9\operatorname{arcsech}(cx)^2\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}}{32cx} + \dots \right)$

```
input int((a+b*arcsech(c*x))^3/x^5,x,method=_RETURNVERBOSE)
```

```
output c^4*(-1/4*a^3/c^4/x^4+b^3*(-1/4/c^4/x^4*arcsech(c*x)^3+3/16*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)/c^3/x^3*arcsech(c*x)^2+9/32*arcsech(c*x)^2/c/x*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)+3/32*arcsech(c*x)^3-3/32/c^4/x^4*arcsech(c*x)+3/128*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)/c^3/x^3+45/256*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)/c/x+45/256*arcsech(c*x)-9/32/c^2/x^2*arcsech(c*x))+3*a*b^2*(-1/4/c^4/x^4*arcsech(c*x)^2+1/8*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)/c^3/x^3*arcsech(c*x)+3/16*arcsech(c*x)/c/x*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)+3/32*arcsech(c*x)^2-1/32/c^4/x^4-3/32/c^2/x^2)+3*a^2*b*(-1/4/c^4/x^4*arcsech(c*x)+1/32*(-(c*x-1)/c/x)^(1/2)/c^3/x^3*((c*x+1)/c/x)^(1/2)*(3*arctanh(1/(-c^2*x^2+1)^(1/2))*c^4*x^4+3*(-c^2*x^2+1)^(1/2)*c^2*x^2+2*(-c^2*x^2+1)^(1/2))/(-c^2*x^2+1)^(1/2)))
```

### Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 351, normalized size of antiderivative = 1.45

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^5} dx = \frac{72 ab^2 c^2 x^2 - 8 (3 b^3 c^4 x^4 - 8 b^3) \log \left( \frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{cx} \right)^3 + 64 a^3 + 24 ab^2 - 24 \left( 3 ab^2 c^4 x^4 - 8 ab^2 + (3 b^3 c^4 x^4 - 8 ab^2 + (3 b^3 c^4 x^4 - 8 ab^2 + (3 b^3 c^4 x^4 - 8 ab^2 + \dots \right)}{x^5}$$

input `integrate((a+b*arcsech(c*x))^3/x^5,x, algorithm="fricas")`

output 
$$\begin{aligned} & -1/256*(72*a*b^2*c^2*x^2 - 8*(3*b^3*c^4*x^4 - 8*b^3)*\log((c*x*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} + 1)/(c*x))^3 + 64*a^3 + 24*a*b^2 - 24*(3*a*b^2*c^4*x^4 \\ & - 8*a*b^2 + (3*b^3*c^3*x^3 + 2*b^3*c*x)*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)})) * \\ & \log((c*x*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} + 1)/(c*x))^2 - 3*(3*(8*a^2*b + 5*b^3)*c^4*x^4 - 24*b^3*c^2*x^2 - 64*a^2*b - 8*b^3 + 16*(3*a*b^2*c^3*x^3 + 2* \\ & a*b^2*c*x)*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)})) * \log((c*x*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)} + 1)/(c^2*x^2)) + 1)/(c*x)) - 3*(3*(8*a^2*b + 5*b^3)*c^3*x^3 + 2*(8*a^2*b + b^3)* \\ & c*x)*\sqrt{-(c^2*x^2 - 1)/(c^2*x^2)})/x^4 \end{aligned}$$

### Sympy [F]

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^5} dx = \int \frac{(a + b \operatorname{asech}(cx))^3}{x^5} dx$$

input `integrate((a+b*asech(c*x))**3/x**5,x)`

output `Integral((a + b*asech(c*x))**3/x**5, x)`

### Maxima [F]

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^5} dx = \int \frac{(b \operatorname{arsech}(cx) + a)^3}{x^5} dx$$

input `integrate((a+b*arcsech(c*x))^3/x^5,x, algorithm="maxima")`

output

```
3/64*a^2*b*((3*c^5*log(c*x*sqrt(1/(c^2*x^2) - 1) + 1) - 3*c^5*log(c*x*sqrt(1/(c^2*x^2) - 1) - 1) - 2*(3*c^8*x^3*(1/(c^2*x^2) - 1)^(3/2) - 5*c^6*x*sqrt(1/(c^2*x^2) - 1))/(c^4*x^4*(1/(c^2*x^2) - 1)^2 - 2*c^2*x^2*(1/(c^2*x^2) - 1) + 1))/c - 16*arcsech(c*x)/x^4) - 1/4*a^3/x^4 + integrate(b^3*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^3/x^5 + 3*a*b^2*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))^2/x^5, x)
```

**Giac [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^5} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3}{x^5} dx$$

input

```
integrate((a+b*arcsech(c*x))^3/x^5,x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)^3/x^5, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^5} dx = \int \frac{(a + b \operatorname{acosh}(\frac{1}{cx}))^3}{x^5} dx$$

input

```
int((a + b*acosh(1/(c*x)))^3/x^5,x)
```

output

```
int((a + b*acosh(1/(c*x)))^3/x^5, x)
```

**Reduce [F]**

$$\int \frac{(a + b \operatorname{sech}^{-1}(cx))^3}{x^5} dx$$

$$= \frac{12 \left( \int \frac{\operatorname{asech}(cx)}{x^5} dx \right) a^2 b x^4 + 4 \left( \int \frac{\operatorname{asech}(cx)^3}{x^5} dx \right) b^3 x^4 + 12 \left( \int \frac{\operatorname{asech}(cx)^2}{x^5} dx \right) a b^2 x^4 - a^3}{4x^4}$$

input `int((a+b*asech(c*x))^3/x^5,x)`

output `(12*int(asech(c*x)/x**5,x)*a**2*b*x**4 + 4*int(asech(c*x)**3/x**5,x)*b**3*x**4 + 12*int(asech(c*x)**2/x**5,x)*a*b**2*x**4 - a**3)/(4*x**4)`

### 3.51 $\int \frac{x}{a+b\operatorname{sech}^{-1}(cx)} dx$

Optimal result	455
Mathematica [N/A]	455
Rubi [N/A]	456
Maple [N/A]	456
Fricas [N/A]	457
Sympy [N/A]	457
Maxima [N/A]	457
Giac [N/A]	458
Mupad [N/A]	458
Reduce [N/A]	459

#### Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{x}{a + b\operatorname{sech}^{-1}(cx)} dx = \operatorname{Int}\left(\frac{x}{a + b\operatorname{sech}^{-1}(cx)}, x\right)$$

output `Defer(Int)(x/(a+b*arcsech(c*x)), x)`

#### Mathematica [N/A]

Not integrable

Time = 2.62 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{x}{a + b\operatorname{sech}^{-1}(cx)} dx = \int \frac{x}{a + b\operatorname{sech}^{-1}(cx)} dx$$

input `Integrate[x/(a + b*ArcSech[c*x]), x]`

output `Integrate[x/(a + b*ArcSech[c*x]), x]`



**Rubi [N/A]**

Not integrable

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{a + b \operatorname{sech}^{-1}(cx)} dx$$

↓ 6865

$$\int \frac{x}{a + b \operatorname{sech}^{-1}(cx)} dx$$

input `Int[x/(a + b*ArcSech[c*x]),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.37 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{x}{a + b \operatorname{arcsech}(cx)} dx$$

input `int(x/(a+b*arcsech(c*x)),x)`

output `int(x/(a+b*arcsech(c*x)),x)`

**Fricas [N/A]**

Not integrable

Time = 0.08 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{x}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{x}{b \operatorname{ar} \operatorname{sech}(cx) + a} dx$$

input `integrate(x/(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral(x/(b*arcsech(c*x) + a), x)`

**Sympy [N/A]**

Not integrable

Time = 0.34 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{x}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{x}{a + b \operatorname{asech}(cx)} dx$$

input `integrate(x/(a+b*asech(c*x)),x)`

output `Integral(x/(a + b*asech(c*x)), x)`

**Maxima [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{x}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{x}{b \operatorname{ar} \operatorname{sech}(cx) + a} dx$$

input `integrate(x/(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `integrate(x/(b*arcsech(c*x) + a), x)`

### Giac [N/A]

Not integrable

Time = 0.10 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{x}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{x}{b \operatorname{arsech}(cx) + a} dx$$

input `integrate(x/(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate(x/(b*arcsech(c*x) + a), x)`

### Mupad [N/A]

Not integrable

Time = 3.51 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.50

$$\int \frac{x}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{x}{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)} dx$$

input `int(x/(a + b*acosh(1/(c*x))),x)`

output `int(x/(a + b*acosh(1/(c*x))), x)`

**Reduce [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{x}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{x}{a \operatorname{sech}(cx) b + a} dx$$

input `int(x/(a+b*asech(c*x)),x)`output `int(x/(asech(c*x)*b + a),x)`

$$3.52 \quad \int \frac{1}{a+b\operatorname{sech}^{-1}(cx)} dx$$

Optimal result	460
Mathematica [N/A]	460
Rubi [N/A]	461
Maple [N/A]	461
Fricas [N/A]	462
Sympy [N/A]	462
Maxima [N/A]	462
Giac [N/A]	463
Mupad [N/A]	463
Reduce [N/A]	464

### Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{1}{a + b\operatorname{sech}^{-1}(cx)} dx = \operatorname{Int}\left(\frac{1}{a + b\operatorname{sech}^{-1}(cx)}, x\right)$$

output `Defer(Int)(1/(a+b*arcsech(c*x)),x)`

### Mathematica [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{1}{a + b\operatorname{sech}^{-1}(cx)} dx = \int \frac{1}{a + b\operatorname{sech}^{-1}(cx)} dx$$

input `Integrate[(a + b*ArcSech[c*x])^(-1),x]`

output `Integrate[(a + b*ArcSech[c*x])^(-1), x]`

**Rubi [N/A]**

Not integrable

Time = 0.17 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{a + b \operatorname{sech}^{-1}(cx)} dx$$

↓ 6865

$$\int \frac{1}{a + b \operatorname{sech}^{-1}(cx)} dx$$

input `Int[(a + b*ArcSech[c*x])^(-1),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.23 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{a + b \operatorname{arcsech}(cx)} dx$$

input `int(1/(a+b*arcsech(c*x)),x)`

output `int(1/(a+b*arcsech(c*x)),x)`

**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{1}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{1}{b \operatorname{ar} \operatorname{sech}(cx) + a} dx$$

input `integrate(1/(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral(1/(b*arcsech(c*x) + a), x)`

**Sympy [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{1}{a + b \operatorname{ar} \operatorname{sech}(cx)} dx$$

input `integrate(1/(a+b*asech(c*x)),x)`

output `Integral(1/(a + b*asech(c*x)), x)`

**Maxima [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{1}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{1}{b \operatorname{ar} \operatorname{sech}(cx) + a} dx$$

input `integrate(1/(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `integrate(1/(b*arcsech(c*x) + a), x)`

### Giac [N/A]

Not integrable

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{1}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{1}{b \operatorname{ar} \operatorname{sech}(cx) + a} dx$$

input `integrate(1/(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate(1/(b*arcsech(c*x) + a), x)`

### Mupad [N/A]

Not integrable

Time = 3.46 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.60

$$\int \frac{1}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{1}{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)} dx$$

input `int(1/(a + b*acosh(1/(c*x))),x)`

output `int(1/(a + b*acosh(1/(c*x))), x)`



**Reduce [N/A]**

Not integrable

Time = 0.21 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{1}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{1}{a \operatorname{sech}(cx) b + a} dx$$

input `int(1/(a+b*asech(c*x)),x)`output `int(1/(asech(c*x)*b + a),x)`

$$3.53 \quad \int \frac{1}{x(a+b\operatorname{sech}^{-1}(cx))} dx$$

Optimal result	465
Mathematica [N/A]	465
Rubi [N/A]	466
Maple [N/A]	466
Fricas [N/A]	467
Sympy [N/A]	467
Maxima [N/A]	467
Giac [N/A]	468
Mupad [N/A]	468
Reduce [N/A]	469

### Optimal result

Integrand size = 14, antiderivative size = 14

$$\int \frac{1}{x(a+b\operatorname{sech}^{-1}(cx))} dx = \operatorname{Int}\left(\frac{1}{x(a+b\operatorname{sech}^{-1}(cx))}, x\right)$$

output `Defer(Int)(1/x/(a+b*arcsech(c*x)), x)`

### Mathematica [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{1}{x(a+b\operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{x(a+b\operatorname{sech}^{-1}(cx))} dx$$

input `Integrate[1/(x*(a + b*ArcSech[c*x])), x]`

output `Integrate[1/(x*(a + b*ArcSech[c*x])), x]`

**Rubi [N/A]**

Not integrable

Time = 0.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x (a + b \operatorname{sech}^{-1}(cx))} dx$$

↓ 6865

$$\int \frac{1}{x (a + b \operatorname{sech}^{-1}(cx))} dx$$

input `Int[1/(x*(a + b*ArcSech[c*x])),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{x (a + b \operatorname{arcsech}(cx))} dx$$

input `int(1/x/(a+b*arcsech(c*x)),x)`

output `int(1/x/(a+b*arcsech(c*x)),x)`

**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

$$\int \frac{1}{x(a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)x} dx$$

input `integrate(1/x/(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral(1/(b*x*arcsech(c*x) + a*x), x)`

**Sympy [N/A]**

Not integrable

Time = 1.18 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{x(a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{x(a + b \operatorname{ar} \operatorname{sech}(cx))} dx$$

input `integrate(1/x/(a+b*asech(c*x)),x)`

output `Integral(1/(x*(a + b*asech(c*x))), x)`

**Maxima [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{1}{x(a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)x} dx$$

input `integrate(1/x/(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `integrate(1/((b*arcsech(c*x) + a)*x), x)`

**Giac [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{1}{x(a + b\operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{(b\operatorname{ar}\operatorname{sech}(cx) + a)x} dx$$

input `integrate(1/x/(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate(1/((b*arcsech(c*x) + a)*x), x)`

**Mupad [N/A]**

Not integrable

Time = 3.62 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.43

$$\int \frac{1}{x(a + b\operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{x(a + b\operatorname{acosh}(\frac{1}{cx}))} dx$$

input `int(1/(x*(a + b*acosh(1/(c*x))))),x)`

output `int(1/(x*(a + b*acosh(1/(c*x))))), x)`

**Reduce [N/A]**

Not integrable

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

$$\int \frac{1}{x(a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{a \operatorname{sech}(cx) bx + ax} dx$$

input `int(1/x/(a+b*asech(c*x)),x)`output `int(1/(asech(c*x)*b*x + a*x),x)`

### 3.54 $\int \frac{1}{x^2(a+b\operatorname{sech}^{-1}(cx))} dx$

Optimal result	470
Mathematica [A] (verified)	470
Rubi [C] (verified)	471
Maple [A] (verified)	473
Fricas [F]	474
Sympy [F]	474
Maxima [F]	474
Giac [F]	475
Mupad [F(-1)]	475
Reduce [F]	475

#### Optimal result

Integrand size = 14, antiderivative size = 46

$$\int \frac{1}{x^2(a+b\operatorname{sech}^{-1}(cx))} dx = \frac{c\operatorname{Chi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right) \sinh\left(\frac{a}{b}\right)}{b} - \frac{c \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{b}$$

output

```
c*Chi(a/b+arcsech(c*x))*sinh(a/b)/b-c*cosh(a/b)*Shi(a/b+arcsech(c*x))/b
```

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.93

$$\int \frac{1}{x^2(a+b\operatorname{sech}^{-1}(cx))} dx = \frac{c(\operatorname{Chi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right) \sinh\left(\frac{a}{b}\right) - \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right))}{b}$$

input

```
Integrate[1/(x^2*(a + b*ArcSech[c*x])),x]
```

output

```
(c*(CoshIntegral[a/b + ArcSech[c*x]]*Sinh[a/b] - Cosh[a/b]*SinhIntegral[a/
b + ArcSech[c*x]]))/b
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.46 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.17, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$ , Rules used = {6839, 3042, 26, 3784, 26, 3042, 26, 3779, 3782}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))} dx \\
 & \quad \downarrow \text{6839} \\
 & -c \int \frac{\sqrt{\frac{1-cx}{cx+1}} (cx+1)}{cx (a + b \operatorname{sech}^{-1}(cx))} d \operatorname{sech}^{-1}(cx) \\
 & \quad \downarrow \text{3042} \\
 & -c \int -\frac{i \sin(i \operatorname{sech}^{-1}(cx))}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) \\
 & \quad \downarrow \text{26} \\
 & ic \int \frac{\sin(i \operatorname{sech}^{-1}(cx))}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) \\
 & \quad \downarrow \text{3784} \\
 & ic \left( \cosh\left(\frac{a}{b}\right) \int \frac{i \sinh\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) - i \sinh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) \right) \\
 & \quad \downarrow \text{26} \\
 & ic \left( i \cosh\left(\frac{a}{b}\right) \int \frac{\sinh\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) - i \sinh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) \right)
 \end{aligned}$$



↓ 3042

$$ic \left( i \cosh \left( \frac{a}{b} \right) \int -\frac{i \sin \left( \frac{ia}{b} + i \operatorname{sech}^{-1}(cx) \right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) - i \sinh \left( \frac{a}{b} \right) \int \frac{\sin \left( \frac{ia}{b} + i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) \right)$$

↓ 26

$$ic \left( \cosh \left( \frac{a}{b} \right) \int \frac{\sin \left( \frac{ia}{b} + i \operatorname{sech}^{-1}(cx) \right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) - i \sinh \left( \frac{a}{b} \right) \int \frac{\sin \left( \frac{ia}{b} + i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) \right)$$

↓ 3779

$$ic \left( \frac{i \cosh \left( \frac{a}{b} \right) \operatorname{Shi} \left( \frac{a}{b} + \operatorname{sech}^{-1}(cx) \right)}{b} - i \sinh \left( \frac{a}{b} \right) \int \frac{\sin \left( \frac{ia}{b} + i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2} \right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) \right)$$

↓ 3782

$$ic \left( \frac{i \cosh \left( \frac{a}{b} \right) \operatorname{Shi} \left( \frac{a}{b} + \operatorname{sech}^{-1}(cx) \right)}{b} - \frac{i \sinh \left( \frac{a}{b} \right) \operatorname{Chi} \left( \frac{a}{b} + \operatorname{sech}^{-1}(cx) \right)}{b} \right)$$

input `Int[1/(x^2*(a + b*ArcSech[c*x])),x]`

output `I*c*((( -I)*CoshIntegral[a/b + ArcSech[c*x]]*Sinh[a/b])/b + (I*Cosh[a/b]*SinhIntegral[a/b + ArcSech[c*x]])/b)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F x_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[F x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3779 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`

rule 3782 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`

rule 3784 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`

rule 6839 `Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^n*(x_)^m, x_Symbol] := Simp[-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])`

## Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.17

method	result	size
derivativedivides	$c \left( -\frac{e^{\frac{a}{b}} \exp\text{Integral}_1\left(\frac{a}{b} + \text{arcsech}(cx)\right)}{2b} + \frac{e^{-\frac{a}{b}} \exp\text{Integral}_1\left(-\text{arcsech}(cx) - \frac{a}{b}\right)}{2b} \right)$	54
default	$c \left( -\frac{e^{\frac{a}{b}} \exp\text{Integral}_1\left(\frac{a}{b} + \text{arcsech}(cx)\right)}{2b} + \frac{e^{-\frac{a}{b}} \exp\text{Integral}_1\left(-\text{arcsech}(cx) - \frac{a}{b}\right)}{2b} \right)$	54

input `int(1/x^2/(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)`

output `c*(-1/2/b*exp(1/b*a)*Ei(1,1/b*a+arcsech(c*x))+1/2/b*exp(-1/b*a)*Ei(1,-arcsech(c*x)-1/b*a))`

**Fricas [F]**

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{(b \operatorname{arsech}(cx) + a)x^2} dx$$

input `integrate(1/x^2/(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral(1/(b*x^2*arcsech(c*x) + a*x^2), x)`

**Sympy [F]**

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{x^2 (a + b \operatorname{asech}(cx))} dx$$

input `integrate(1/x**2/(a+b*asech(c*x)),x)`

output `Integral(1/(x**2*(a + b*asech(c*x))), x)`

**Maxima [F]**

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{(b \operatorname{arsech}(cx) + a)x^2} dx$$

input `integrate(1/x^2/(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `integrate(1/((b*arcsech(c*x) + a)*x^2), x)`

**Giac [F]**

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{(b \operatorname{arsech}(cx) + a)x^2} dx$$

input `integrate(1/x^2/(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate(1/((b*arcsech(c*x) + a)*x^2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{x^2 (a + b \operatorname{acosh}(\frac{1}{cx}))} dx$$

input `int(1/(x^2*(a + b*acosh(1/(c*x))))),x)`

output `int(1/(x^2*(a + b*acosh(1/(c*x))))), x)`

**Reduce [F]**

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{\operatorname{asech}(cx) b x^2 + a x^2} dx$$

input `int(1/x^2/(a+b*asech(c*x)),x)`

output `int(1/(asech(c*x)*b*x**2 + a*x**2),x)`

### 3.55 $\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))} dx$

Optimal result	476
Mathematica [A] (verified)	476
Rubi [C] (verified)	477
Maple [A] (verified)	480
Fricas [F]	480
Sympy [F]	480
Maxima [F]	481
Giac [F]	481
Mupad [F(-1)]	481
Reduce [F]	482

#### Optimal result

Integrand size = 14, antiderivative size = 63

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))} dx = \frac{c^2 \operatorname{Chi}\left(\frac{2a}{b} + 2 \operatorname{sech}^{-1}(cx)\right) \sinh\left(\frac{2a}{b}\right)}{2b} - \frac{c^2 \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2a}{b} + 2 \operatorname{sech}^{-1}(cx)\right)}{2b}$$

output

$1/2*c^2*Chi(2*a/b+2*arcsech(c*x))*sinh(2*a/b)/b-1/2*c^2*cosh(2*a/b)*Shi(2*a/b+2*arcsech(c*x))/b$

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.89

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))} dx = \frac{c^2 (\operatorname{Chi}\left(\frac{2a}{b} + 2 \operatorname{sech}^{-1}(cx)\right) \sinh\left(\frac{2a}{b}\right) - \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2a}{b} + 2 \operatorname{sech}^{-1}(cx)\right))}{2b}$$

input

`Integrate[1/(x^3*(a + b*ArcSech[c*x])),x]`

output

```
(c^2*(CoshIntegral[(2*a)/b + 2*ArcSech[c*x]]*Sinh[(2*a)/b] - Cosh[(2*a)/b]
*SinhIntegral[(2*a)/b + 2*ArcSech[c*x]]))/(2*b)
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.57 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.05, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.786$ , Rules used = {6839, 5971, 27, 3042, 26, 3784, 26, 3042, 26, 3779, 3782}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))} dx \\
 & \quad \downarrow \text{6839} \\
 & -c^2 \int \frac{\sqrt{\frac{1-cx}{cx+1}} (cx+1)}{c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))} d \operatorname{sech}^{-1}(cx) \\
 & \quad \downarrow \text{5971} \\
 & -c^2 \int \frac{\sinh(2 \operatorname{sech}^{-1}(cx))}{2 (a + b \operatorname{sech}^{-1}(cx))} d \operatorname{sech}^{-1}(cx) \\
 & \quad \downarrow \text{27} \\
 & -\frac{1}{2} c^2 \int \frac{\sinh(2 \operatorname{sech}^{-1}(cx))}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{2} c^2 \int -\frac{i \sin(2i \operatorname{sech}^{-1}(cx))}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) \\
 & \quad \downarrow \text{26} \\
 & \frac{1}{2} i c^2 \int \frac{\sin(2i \operatorname{sech}^{-1}(cx))}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) \\
 & \quad \downarrow \text{3784}
 \end{aligned}$$

$$\frac{1}{2}ic^2 \left( \cosh\left(\frac{2a}{b}\right) \int \frac{i \sinh\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{a + b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) - i \sinh\left(\frac{2a}{b}\right) \int \frac{\cosh\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{a + b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)$$

↓ 26

$$\frac{1}{2}ic^2 \left( i \cosh\left(\frac{2a}{b}\right) \int \frac{\sinh\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{a + b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) - i \sinh\left(\frac{2a}{b}\right) \int \frac{\cosh\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{a + b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)$$

↓ 3042

$$\frac{1}{2}ic^2 \left( i \cosh\left(\frac{2a}{b}\right) \int -\frac{i \sin\left(\frac{2ia}{b} + 2i\operatorname{sech}^{-1}(cx)\right)}{a + b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) - i \sinh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2ia}{b} + 2i\operatorname{sech}^{-1}(cx) + \frac{\pi}{2}\right)}{a + b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)$$

↓ 26

$$\frac{1}{2}ic^2 \left( \cosh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2ia}{b} + 2i\operatorname{sech}^{-1}(cx)\right)}{a + b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) - i \sinh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2ia}{b} + 2i\operatorname{sech}^{-1}(cx) + \frac{\pi}{2}\right)}{a + b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)$$

↓ 3779

$$\frac{1}{2}ic^2 \left( \frac{i \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{b} - i \sinh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2ia}{b} + 2i\operatorname{sech}^{-1}(cx) + \frac{\pi}{2}\right)}{a + b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)$$

↓ 3782

$$\frac{1}{2}ic^2 \left( \frac{i \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{b} - \frac{i \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{b} \right)$$

input `Int[1/(x^3*(a + b*ArcSech[c*x])),x]`

output `(I/2)*c^2*(((-I)*CoshIntegral[(2*a)/b + 2*ArcSech[c*x]]*Sinh[(2*a)/b])/b + (I*Cosh[(2*a)/b]*SinhIntegral[(2*a)/b + 2*ArcSech[c*x]])/b)`

## Definitions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a_])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 27  $\text{Int}[(a_)*(F_x), x\_Symbol] \rightarrow \text{Simp}[a \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] /; \text{FreeQ}[b, x]$
- rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3779  $\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x\_Symbol] \rightarrow \text{Simp}[I*(\text{SinhIntegral}[c*f*(fz/d) + f*fz*x]/d), x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \ \&\& \ \text{EqQ}[d*e - c*f*fz*I, 0]$
- rule 3782  $\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x\_Symbol] \rightarrow \text{Simp}[\text{CoshIntegral}[c*f*(fz/d) + f*fz*x]/d, x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \ \&\& \ \text{EqQ}[d*(e - \text{Pi}/2) - c*f*fz*I, 0]$
- rule 3784  $\text{Int}[\sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x\_Symbol] \rightarrow \text{Simp}[\text{Cos}[(d*e - c*f)/d] \text{Int}[\text{Sin}[c*(f/d) + f*x]/(c + d*x), x], x] + \text{Simp}[\text{Sin}[(d*e - c*f)/d] \text{Int}[\text{Cos}[c*(f/d) + f*x]/(c + d*x), x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \ \&\& \ \text{NeQ}[d*e - c*f, 0]$
- rule 5971  $\text{Int}[\text{Cosh}[(a_.) + (b_.)*(x_)]^{(p_.)}*((c_.) + (d_.)*(x_))^{(m_.)}*\text{Sinh}[(a_.) + (b_.)*(x_)]^{(n_.)}, x\_Symbol] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(c + d*x)^m, \text{Sinh}[a + b*x]^{n*} \text{Cosh}[a + b*x]^p, x], x] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IGtQ}[p, 0]$
- rule 6839  $\text{Int}[(a_.) + \text{ArcSech}[(c_.)*(x_)]*(b_.)^{(n_.)}*(x_)^{(m_.)}, x\_Symbol] \rightarrow \text{Simp}[-(c^{(m+1)})^{(-1)} \text{Subst}[\text{Int}[(a + b*x)^n*\text{Sech}[x]^{(m+1)}*\text{Tanh}[x], x], x, \text{ArcSech}[c*x]], x] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ (\text{GtQ}[n, 0] \ || \ \text{LtQ}[m, -1])$



**Maple [A] (verified)**

Time = 0.57 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.95

method	result	size
derivativedivides	$c^2 \left( -\frac{e^{\frac{2a}{b}} \operatorname{ExpIntegral}_1\left(\frac{2a}{b} + 2 \operatorname{arcsech}(cx)\right)}{4b} + \frac{e^{-\frac{2a}{b}} \operatorname{ExpIntegral}_1\left(-2 \operatorname{arcsech}(cx) - \frac{2a}{b}\right)}{4b} \right)$	60
default	$c^2 \left( -\frac{e^{\frac{2a}{b}} \operatorname{ExpIntegral}_1\left(\frac{2a}{b} + 2 \operatorname{arcsech}(cx)\right)}{4b} + \frac{e^{-\frac{2a}{b}} \operatorname{ExpIntegral}_1\left(-2 \operatorname{arcsech}(cx) - \frac{2a}{b}\right)}{4b} \right)$	60

input `int(1/x^3/(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)`

output `c^2*(-1/4/b*exp(2/b*a)*Ei(1,2/b*a+2*arcsech(c*x))+1/4/b*exp(-2/b*a)*Ei(1,-2*arcsech(c*x)-2/b*a))`

**Fricas [F]**

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a) x^3} dx$$

input `integrate(1/x^3/(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral(1/(b*x^3*arcsech(c*x) + a*x^3), x)`

**Sympy [F]**

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{x^3 (a + b \operatorname{ar} \operatorname{sech}(cx))} dx$$

input `integrate(1/x**3/(a+b*asech(c*x)),x)`

output `Integral(1/(x**3*(a + b*asech(c*x))), x)`

**Maxima [F]**

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^3} dx$$

input `integrate(1/x^3/(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `integrate(1/((b*arcsech(c*x) + a)*x^3), x)`

**Giac [F]**

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^3} dx$$

input `integrate(1/x^3/(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate(1/((b*arcsech(c*x) + a)*x^3), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{x^3 (a + b \operatorname{acosh}(\frac{1}{cx}))} dx$$

input `int(1/(x^3*(a + b*acosh(1/(c*x))))),x)`

output `int(1/(x^3*(a + b*acosh(1/(c*x))))), x)`

**Reduce [F]**

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{\operatorname{asech}(cx) b x^3 + a x^3} dx$$

input `int(1/x^3/(a+b*asech(c*x)),x)`

output `int(1/(asech(c*x)*b*x**3 + a*x**3),x)`

### 3.56 $\int \frac{1}{x^4 (a+b\operatorname{sech}^{-1}(cx))} dx$

Optimal result	483
Mathematica [A] (verified)	484
Rubi [A] (verified)	484
Maple [A] (verified)	485
Fricas [F]	486
Sympy [F]	486
Maxima [F]	487
Giac [F]	487
Mupad [F(-1)]	487
Reduce [F]	488

#### Optimal result

Integrand size = 14, antiderivative size = 117

$$\int \frac{1}{x^4 (a + b\operatorname{sech}^{-1}(cx))} dx = \frac{c^3 \operatorname{Chi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right) \sinh\left(\frac{a}{b}\right)}{4b} + \frac{c^3 \operatorname{Chi}\left(\frac{3a}{b} + 3\operatorname{sech}^{-1}(cx)\right) \sinh\left(\frac{3a}{b}\right)}{4b} - \frac{c^3 \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{4b} - \frac{c^3 \cosh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3a}{b} + 3\operatorname{sech}^{-1}(cx)\right)}{4b}$$

output

```
1/4*c^3*Chi(a/b+arcsech(c*x))*sinh(a/b)/b+1/4*c^3*Chi(3*a/b+3*arcsech(c*x))
)*sinh(3*a/b)/b-1/4*c^3*cosh(a/b)*Shi(a/b+arcsech(c*x))/b-1/4*c^3*cosh(3*a
/b)*Shi(3*a/b+3*arcsech(c*x))/b
```

**Mathematica [A] (verified)**

Time = 0.12 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.78

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))} dx = \frac{c^3 \left( -\operatorname{Chi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right) \sinh\left(\frac{a}{b}\right) - \operatorname{Chi}\left(3\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)\right) \sinh\left(\frac{3a}{b}\right) + \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right) \right)}{4b}$$

input `Integrate[1/(x^4*(a + b*ArcSech[c*x])),x]`

output

```
-1/4*(c^3*(-(CoshIntegral[a/b + ArcSech[c*x]]*Sinh[a/b]) - CoshIntegral[3*(a/b + ArcSech[c*x]])*Sinh[(3*a)/b] + Cosh[a/b]*SinhIntegral[a/b + ArcSech[c*x]] + Cosh[(3*a)/b]*SinhIntegral[3*(a/b + ArcSech[c*x])]))/b
```

**Rubi [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.94, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {6839, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))} dx \\ & \quad \downarrow \text{6839} \\ & -c^3 \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))} d \operatorname{sech}^{-1}(cx) \\ & \quad \downarrow \text{5971} \\ & -c^3 \int \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{4cx (a + b \operatorname{sech}^{-1}(cx))} + \frac{\sinh(3 \operatorname{sech}^{-1}(cx))}{4 (a + b \operatorname{sech}^{-1}(cx))} \right) d \operatorname{sech}^{-1}(cx) \\ & \quad \downarrow \text{2009} \end{aligned}$$

$$-c^3 \left( -\frac{\sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{4b} - \frac{\sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3a}{b} + 3\operatorname{sech}^{-1}(cx)\right)}{4b} + \frac{\cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{4b} + \dots \right)$$

input `Int[1/(x^4*(a + b*ArcSech[c*x])),x]`

output `-(c^3*(-1/4*(CoshIntegral[a/b + ArcSech[c*x]]*Sinh[a/b])/b - (CoshIntegral[(3*a)/b + 3*ArcSech[c*x]]*Sinh[(3*a)/b])/(4*b) + (Cosh[a/b]*SinhIntegral[a/b + ArcSech[c*x]])/(4*b) + (Cosh[(3*a)/b]*SinhIntegral[(3*a)/b + 3*ArcSech[c*x]])/(4*b)))`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)^(p_.)*((c_.) + (d_.)*(x_)^(m_.)*Sinh[(a_.) + (b_.)*(x_)^(n_.)], x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6839 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))^(n_.)*(x_)^(m_.), x_Symbol] := Simp[-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])`

**Maple [A] (verified)**

Time = 0.71 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.94

method	result
derivativedivides	$c^3 \left( -\frac{e^{\frac{3a}{b}} \operatorname{expIntegral}_1\left(\frac{3a}{b} + 3 \operatorname{arcsech}(cx)\right)}{8b} - \frac{e^{\frac{a}{b}} \operatorname{expIntegral}_1\left(\frac{a}{b} + \operatorname{arcsech}(cx)\right)}{8b} + \frac{e^{-\frac{a}{b}} \operatorname{expIntegral}_1\left(-\operatorname{arcsech}(cx)\right)}{8b} \right)$
default	$c^3 \left( -\frac{e^{\frac{3a}{b}} \operatorname{expIntegral}_1\left(\frac{3a}{b} + 3 \operatorname{arcsech}(cx)\right)}{8b} - \frac{e^{\frac{a}{b}} \operatorname{expIntegral}_1\left(\frac{a}{b} + \operatorname{arcsech}(cx)\right)}{8b} + \frac{e^{-\frac{a}{b}} \operatorname{expIntegral}_1\left(-\operatorname{arcsech}(cx)\right)}{8b} \right)$

input `int(1/x^4/(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)`

output `c^3*(-1/8/b*exp(3/b*a)*Ei(1,3/b*a+3*arcsech(c*x))-1/8/b*exp(1/b*a)*Ei(1,1/b*a+arcsech(c*x))+1/8/b*exp(-1/b*a)*Ei(1,-arcsech(c*x)-1/b*a)+1/8/b*exp(-3/b*a)*Ei(1,-3*arcsech(c*x)-3/b*a))`

### Fricas [F]

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^4} dx$$

input `integrate(1/x^4/(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral(1/(b*x^4*arcsech(c*x) + a*x^4), x)`

### Sympy [F]

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{x^4 (a + b \operatorname{ar} \operatorname{sech}(cx))} dx$$

input `integrate(1/x**4/(a+b*asech(c*x)),x)`

output `Integral(1/(x**4*(a + b*asech(c*x))), x)`

**Maxima [F]**

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^4} dx$$

input `integrate(1/x^4/(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `integrate(1/((b*arcsech(c*x) + a)*x^4), x)`

**Giac [F]**

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^4} dx$$

input `integrate(1/x^4/(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate(1/((b*arcsech(c*x) + a)*x^4), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{x^4 (a + b \operatorname{acosh}(\frac{1}{cx}))} dx$$

input `int(1/(x^4*(a + b*acosh(1/(c*x))))),x)`

output `int(1/(x^4*(a + b*acosh(1/(c*x))))), x)`



Reduce [F]

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))} dx = \int \frac{1}{\operatorname{asech}(cx) b x^4 + a x^4} dx$$

input `int(1/x^4/(a+b*asech(c*x)),x)`

output `int(1/(asech(c*x)*b*x**4 + a*x**4),x)`

$$3.57 \quad \int \frac{x}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^2} dx$$

Optimal result	489
Mathematica [N/A]	489
Rubi [N/A]	490
Maple [N/A]	490
Fricas [N/A]	491
Sympy [N/A]	491
Maxima [N/A]	491
Giac [N/A]	492
Mupad [N/A]	492
Reduce [N/A]	493

### Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{x}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^2} dx = \operatorname{Int}\left(\frac{x}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^2}, x\right)$$

output `Defer(Int)(x/(a+b*arcsech(c*x))^2,x)`

### Mathematica [N/A]

Not integrable

Time = 12.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{x}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^2} dx = \int \frac{x}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^2} dx$$

input `Integrate[x/(a + b*ArcSech[c*x])^2,x]`

output `Integrate[x/(a + b*ArcSech[c*x])^2, x]`

**Rubi [N/A]**

Not integrable

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^2} dx$$

↓ 6865

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^2} dx$$

input `Int[x/(a + b*ArcSech[c*x])^2,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{x}{(a + b \operatorname{arcsech}(cx))^2} dx$$

input `int(x/(a+b*arcsech(c*x))^2,x)`

output `int(x/(a+b*arcsech(c*x))^2,x)`

**Fricas [N/A]**

Not integrable

Time = 0.08 (sec) , antiderivative size = 28, normalized size of antiderivative = 2.33

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{x}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2} dx$$

input `integrate(x/(a+b*arcsech(c*x))^2,x, algorithm="fricas")`

output `integral(x/(b^2*arcsech(c*x)^2 + 2*a*b*arcsech(c*x) + a^2), x)`

**Sympy [N/A]**

Not integrable

Time = 0.70 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{x}{(a + b \operatorname{ar} \operatorname{sech}(cx))^2} dx$$

input `integrate(x/(a+b*asech(c*x))**2,x)`

output `Integral(x/(a + b*asech(c*x))**2, x)`

**Maxima [N/A]**

Not integrable

Time = 0.40 (sec) , antiderivative size = 546, normalized size of antiderivative = 45.50

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{x}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2} dx$$

input `integrate(x/(a+b*arcsech(c*x))^2,x, algorithm="maxima")`

output

```

-((c^2*x^3 - x)*sqrt(c*x + 1)*sqrt(-c*x + 1)*x + (c^2*x^3 - x)*x)/((b^2*c^
2*log(c) - a*b*c^2)*x^2 - b^2*log(c) - (b^2*log(c) + b^2*log(x) - a*b)*sqr
t(c*x + 1)*sqrt(-c*x + 1) + a*b - (b^2*c^2*x^2 - sqrt(c*x + 1)*sqrt(-c*x +
1)*b^2 - b^2)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1) + (b^2*c^2*x^2 - b^2)
*log(x)) + integrate((2*(2*c^2*x^2 - 1)*(c*x + 1)*(c*x - 1)*x + (3*c^4*x^4
- 8*c^2*x^2 + 4)*sqrt(c*x + 1)*sqrt(-c*x + 1)*x + 2*(c^4*x^4 - 2*c^2*x^2
+ 1)*x)/((b^2*c^4*log(c) - a*b*c^4)*x^4 - (b^2*log(c) + b^2*log(x) - a*b)*
(c*x + 1)*(c*x - 1) - 2*(b^2*c^2*log(c) - a*b*c^2)*x^2 + b^2*log(c) - 2*((
b^2*c^2*log(c) - a*b*c^2)*x^2 - b^2*log(c) + a*b + (b^2*c^2*x^2 - b^2)*log
(x))*sqrt(c*x + 1)*sqrt(-c*x + 1) - a*b - (b^2*c^4*x^4 - 2*b^2*c^2*x^2 - (
c*x + 1)*(c*x - 1)*b^2 - 2*(b^2*c^2*x^2 - b^2)*sqrt(c*x + 1)*sqrt(-c*x + 1
) + b^2)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1) + (b^2*c^4*x^4 - 2*b^2*c^2*
x^2 + b^2)*log(x)), x)

```

**Giac [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{x}{(b \operatorname{arsech}(cx) + a)^2} dx$$

input

```
integrate(x/(a+b*arcsech(c*x))^2,x, algorithm="giac")
```

output

```
integrate(x/(b*arcsech(c*x) + a)^2, x)
```

**Mupad [N/A]**

Not integrable

Time = 3.53 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.50

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{x}{(a + b \operatorname{acosh}(\frac{1}{cx}))^2} dx$$

input

```
int(x/(a + b*acosh(1/(c*x)))^2,x)
```

output `int(x/(a + b*acosh(1/(c*x)))^2, x)`

**Reduce [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 28, normalized size of antiderivative = 2.33

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{x}{\operatorname{asech}(cx)^2 b^2 + 2 \operatorname{asech}(cx) ab + a^2} dx$$

input `int(x/(a+b*asech(c*x))^2,x)`

output `int(x/(asech(c*x)**2*b**2 + 2*asech(c*x)*a*b + a**2),x)`

$$3.58 \quad \int \frac{1}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^2} dx$$

Optimal result	494
Mathematica [N/A]	494
Rubi [N/A]	495
Maple [N/A]	495
Fricas [N/A]	496
Sympy [N/A]	496
Maxima [N/A]	496
Giac [N/A]	497
Mupad [N/A]	497
Reduce [N/A]	498

### Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{1}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^2} dx = \operatorname{Int}\left(\frac{1}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^2}, x\right)$$

output `Defer(Int)(1/(a+b*arcsech(c*x))^2,x)`

### Mathematica [N/A]

Not integrable

Time = 59.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{1}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^2} dx = \int \frac{1}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^2} dx$$

input `Integrate[(a + b*ArcSech[c*x])^(-2), x]`

output `Integrate[(a + b*ArcSech[c*x])^(-2), x]`

**Rubi [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^2} dx$$

↓ 6865

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^2} dx$$

input `Int[(a + b*ArcSech[c*x])^(-2),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.20 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a + b \operatorname{arcsech}(cx))^2} dx$$

input `int(1/(a+b*arcsech(c*x))^2,x)`

output `int(1/(a+b*arcsech(c*x))^2,x)`



**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 2.60

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2} dx$$

input `integrate(1/(a+b*arcsech(c*x))^2,x, algorithm="fricas")`

output `integral(1/(b^2*arcsech(c*x)^2 + 2*a*b*arcsech(c*x) + a^2), x)`

**Sympy [N/A]**

Not integrable

Time = 0.78 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(a + b \operatorname{asech}(cx))^2} dx$$

input `integrate(1/(a+b*asech(c*x))**2,x)`

output `Integral((a + b*asech(c*x))**(-2), x)`

**Maxima [N/A]**

Not integrable

Time = 0.42 (sec) , antiderivative size = 535, normalized size of antiderivative = 53.50

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2} dx$$

input `integrate(1/(a+b*arcsech(c*x))^2,x, algorithm="maxima")`

output

```

-(c^2*x^3 + (c^2*x^3 - x)*sqrt(c*x + 1)*sqrt(-c*x + 1) - x)/((b^2*c^2*log(
c) - a*b*c^2)*x^2 - b^2*log(c) - (b^2*log(c) + b^2*log(x) - a*b)*sqrt(c*x
+ 1)*sqrt(-c*x + 1) + a*b - (b^2*c^2*x^2 - sqrt(c*x + 1)*sqrt(-c*x + 1)*b^
2 - b^2)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1) + (b^2*c^2*x^2 - b^2)*log(x
)) + integrate((c^4*x^4 - 2*c^2*x^2 + (3*c^2*x^2 - 1)*(c*x + 1)*(c*x - 1)
+ (2*c^4*x^4 - 5*c^2*x^2 + 2)*sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)/((b^2*c^4*
log(c) - a*b*c^4)*x^4 - (b^2*log(c) + b^2*log(x) - a*b)*(c*x + 1)*(c*x - 1
) - 2*(b^2*c^2*log(c) - a*b*c^2)*x^2 + b^2*log(c) - 2*((b^2*c^2*log(c) - a
*b*c^2)*x^2 - b^2*log(c) + a*b + (b^2*c^2*x^2 - b^2)*log(x))*sqrt(c*x + 1)
*sqrt(-c*x + 1) - a*b - (b^2*c^4*x^4 - 2*b^2*c^2*x^2 - (c*x + 1)*(c*x - 1)
*b^2 - 2*(b^2*c^2*x^2 - b^2)*sqrt(c*x + 1)*sqrt(-c*x + 1) + b^2)*log(sqrt(
c*x + 1)*sqrt(-c*x + 1) + 1) + (b^2*c^4*x^4 - 2*b^2*c^2*x^2 + b^2)*log(x)
, x)

```

**Giac [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(b \operatorname{arsech}(cx) + a)^2} dx$$

input

```
integrate(1/(a+b*arcsech(c*x))^2,x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)^(-2), x)
```

**Mupad [N/A]**

Not integrable

Time = 3.45 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.60

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(a + b \operatorname{acosh}(\frac{1}{cx}))^2} dx$$

input

```
int(1/(a + b*acosh(1/(c*x)))^2,x)
```

output `int(1/(a + b*acosh(1/(c*x)))^2, x)`

**Reduce [N/A]**

Not integrable

Time = 0.21 (sec) , antiderivative size = 26, normalized size of antiderivative = 2.60

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{\operatorname{sech}(cx)^2 b^2 + 2 \operatorname{sech}(cx) ab + a^2} dx$$

input `int(1/(a+b*asech(c*x))^2,x)`

output `int(1/(asech(c*x)**2*b**2 + 2*asech(c*x)*a*b + a**2),x)`

$$3.59 \quad \int \frac{1}{x \left( a + b \operatorname{sech}^{-1}(cx) \right)^2} dx$$

Optimal result	499
Mathematica [N/A]	499
Rubi [N/A]	500
Maple [N/A]	500
Fricas [N/A]	501
Sympy [N/A]	501
Maxima [N/A]	501
Giac [N/A]	502
Mupad [N/A]	502
Reduce [N/A]	503

### Optimal result

Integrand size = 14, antiderivative size = 14

$$\int \frac{1}{x \left( a + b \operatorname{sech}^{-1}(cx) \right)^2} dx = \operatorname{Int} \left( \frac{1}{x \left( a + b \operatorname{sech}^{-1}(cx) \right)^2}, x \right)$$

output `Defer(Int)(1/x/(a+b*arcsech(c*x))^2,x)`

### Mathematica [N/A]

Not integrable

Time = 4.35 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{1}{x \left( a + b \operatorname{sech}^{-1}(cx) \right)^2} dx = \int \frac{1}{x \left( a + b \operatorname{sech}^{-1}(cx) \right)^2} dx$$

input `Integrate[1/(x*(a + b*ArcSech[c*x])^2),x]`

output `Integrate[1/(x*(a + b*ArcSech[c*x])^2), x]`

**Rubi [N/A]**

Not integrable

Time = 0.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x (a + b \operatorname{sech}^{-1}(cx))^2} dx$$

↓ 6865

$$\int \frac{1}{x (a + b \operatorname{sech}^{-1}(cx))^2} dx$$

input `Int[1/(x*(a + b*ArcSech[c*x])^2),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{x (a + b \operatorname{arcsech}(cx))^2} dx$$

input `int(1/x/(a+b*arcsech(c*x))^2,x)`

output `int(1/x/(a+b*arcsech(c*x))^2,x)`

**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 30, normalized size of antiderivative = 2.14

$$\int \frac{1}{x (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2 x} dx$$

input `integrate(1/x/(a+b*arcsech(c*x))^2,x, algorithm="fricas")`

output `integral(1/(b^2*x*arcsech(c*x)^2 + 2*a*b*x*arcsech(c*x) + a^2*x), x)`

**Sympy [N/A]**

Not integrable

Time = 1.47 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{x (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{x (a + b \operatorname{asech}(cx))^2} dx$$

input `integrate(1/x/(a+b*asech(c*x))**2,x)`

output `Integral(1/(x*(a + b*asech(c*x))**2), x)`

**Maxima [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 544, normalized size of antiderivative = 38.86

$$\int \frac{1}{x (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2 x} dx$$

input `integrate(1/x/(a+b*arcsech(c*x))^2,x, algorithm="maxima")`

output

```

-(c^2*x^3 + (c^2*x^3 - x)*sqrt(c*x + 1)*sqrt(-c*x + 1) - x)/((b^2*c^2*x^2
- b^2)*x*log(x) - (b^2*x*log(x) + (b^2*log(c) - a*b)*x)*sqrt(c*x + 1)*sqrt
(-c*x + 1) + ((b^2*c^2*log(c) - a*b*c^2)*x^2 - b^2*log(c) + a*b)*x + (sqrt
(c*x + 1)*sqrt(-c*x + 1)*b^2*x - (b^2*c^2*x^2 - b^2)*x)*log(sqrt(c*x + 1)*
sqrt(-c*x + 1) + 1)) + integrate(-(2*(c*x + 1)*(c*x - 1)*c^2*x^2 + (c^4*x^
4 - 2*c^2*x^2)*sqrt(c*x + 1)*sqrt(-c*x + 1))/((b^2*x*log(x) + (b^2*log(c)
- a*b)*x)*(c*x + 1)*(c*x - 1) - (b^2*c^4*x^4 - 2*b^2*c^2*x^2 + b^2)*x*log(
x) + 2*((b^2*c^2*x^2 - b^2)*x*log(x) + ((b^2*c^2*log(c) - a*b*c^2)*x^2 - b
^2*log(c) + a*b)*x)*sqrt(c*x + 1)*sqrt(-c*x + 1) - ((b^2*c^4*log(c) - a*b*
c^4)*x^4 - 2*(b^2*c^2*log(c) - a*b*c^2)*x^2 + b^2*log(c) - a*b)*x - ((c*x
+ 1)*(c*x - 1)*b^2*x + 2*(b^2*c^2*x^2 - b^2)*sqrt(c*x + 1)*sqrt(-c*x + 1)*
x - (b^2*c^4*x^4 - 2*b^2*c^2*x^2 + b^2)*x)*log(sqrt(c*x + 1)*sqrt(-c*x + 1
) + 1)), x)

```

**Giac [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{1}{x(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2 x} dx$$

input

```
integrate(1/x/(a+b*arcsech(c*x))^2,x, algorithm="giac")
```

output

```
integrate(1/((b*arcsech(c*x) + a)^2*x), x)
```

**Mupad [N/A]**

Not integrable

Time = 3.48 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.43

$$\int \frac{1}{x(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{x(a + b \operatorname{acosh}(\frac{1}{cx}))^2} dx$$

input

```
int(1/(x*(a + b*acosh(1/(c*x)))^2),x)
```

output `int(1/(x*(a + b*acosh(1/(c*x))))^2), x)`

**Reduce [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 30, normalized size of antiderivative = 2.14

$$\int \frac{1}{x (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{\operatorname{asech}(cx)^2 b^2 x + 2 \operatorname{asech}(cx) abx + a^2 x} dx$$

input `int(1/x/(a+b*asech(c*x))^2,x)`

output `int(1/(asech(c*x)**2*b**2*x + 2*asech(c*x)*a*b*x + a**2*x),x)`



**3.60**  $\int \frac{1}{x^2 (a+b\operatorname{sech}^{-1}(cx))^2} dx$

Optimal result . . . . .	504
Mathematica [A] (verified) . . . . .	504
Rubi [C] (verified) . . . . .	505
Maple [A] (verified) . . . . .	508
Fricas [F] . . . . .	509
Sympy [F] . . . . .	509
Maxima [F] . . . . .	509
Giac [F] . . . . .	510
Mupad [F(-1)] . . . . .	510
Reduce [F] . . . . .	511

**Optimal result**

Integrand size = 14, antiderivative size = 86

$$\int \frac{1}{x^2 (a + b\operatorname{sech}^{-1}(cx))^2} dx = \frac{\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{bx (a + b\operatorname{sech}^{-1}(cx))} - \frac{c \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{b^2} + \frac{c \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{b^2}$$

output

```
((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/b/x/(a+b*arcsech(c*x))-c*cosh(a/b)*Chi(a/b+arcsech(c*x))/b^2+c*sinh(a/b)*Shi(a/b+arcsech(c*x))/b^2
```

**Mathematica [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.95

$$\int \frac{1}{x^2 (a + b\operatorname{sech}^{-1}(cx))^2} dx = \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{x(a+b\operatorname{sech}^{-1}(cx))} - c \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right) + c \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{b^2}$$

input `Integrate[1/(x^2*(a + b*ArcSech[c*x])^2),x]`

output `((b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(x*(a + b*ArcSech[c*x])) - c*Cosh[a/b]*CoshIntegral[a/b + ArcSech[c*x]] + c*Sinh[a/b]*SinhIntegral[a/b + ArcSech[c*x]])/b^2`

## Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.58 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.20, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.786$ , Rules used = {6839, 3042, 26, 3778, 3042, 3784, 26, 3042, 26, 3779, 3782}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))^2} dx \\
 & \quad \downarrow \text{6839} \\
 & -c \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{cx (a + b \operatorname{sech}^{-1}(cx))^2} d \operatorname{sech}^{-1}(cx) \\
 & \quad \downarrow \text{3042} \\
 & -c \int -\frac{i \sin(i \operatorname{sech}^{-1}(cx))}{(a + b \operatorname{sech}^{-1}(cx))^2} d \operatorname{sech}^{-1}(cx) \\
 & \quad \downarrow \text{26} \\
 & ic \int \frac{\sin(i \operatorname{sech}^{-1}(cx))}{(a + b \operatorname{sech}^{-1}(cx))^2} d \operatorname{sech}^{-1}(cx) \\
 & \quad \downarrow \text{3778} \\
 & ic \left( \frac{i \int \frac{1}{cx (a + b \operatorname{sech}^{-1}(cx))} d \operatorname{sech}^{-1}(cx)}{b} - \frac{i \sqrt{\frac{1-cx}{cx+1}}(cx+1)}{bcx (a + b \operatorname{sech}^{-1}(cx))} \right)
 \end{aligned}$$

↓ 3042

$$ic \left( \frac{i \int \frac{\sin(i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2})}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx)}{b} - \frac{i \sqrt{\frac{1-cx}{cx+1}} (cx+1)}{bcx (a + b \operatorname{sech}^{-1}(cx))} \right)$$

↓ 3784

$$ic \left( \frac{i \left( \cosh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) + i \sinh\left(\frac{a}{b}\right) \int \frac{i \sinh\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) \right)}{b} - \frac{i \sqrt{\frac{1-cx}{cx+1}} (cx+1)}{bcx (a + b \operatorname{sech}^{-1}(cx))} \right)$$

↓ 26

$$ic \left( \frac{i \left( \cosh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) - \sinh\left(\frac{a}{b}\right) \int \frac{\sinh\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) \right)}{b} - \frac{i \sqrt{\frac{1-cx}{cx+1}} (cx+1)}{bcx (a + b \operatorname{sech}^{-1}(cx))} \right)$$

↓ 3042

$$ic \left( \frac{i \left( \cosh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{ia}{b} + i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2}\right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) - \sinh\left(\frac{a}{b}\right) \int \frac{-i \sin\left(\frac{ia}{b} + i \operatorname{sech}^{-1}(cx)\right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) \right)}{b} - \frac{i \sqrt{\frac{1-cx}{cx+1}} (cx+1)}{bcx (a + b \operatorname{sech}^{-1}(cx))} \right)$$

↓ 26

$$ic \left( \frac{i \left( \cosh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{ia}{b} + i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2}\right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) + i \sinh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{ia}{b} + i \operatorname{sech}^{-1}(cx)\right)}{a + b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) \right)}{b} - \frac{i \sqrt{\frac{1-cx}{cx+1}} (cx+1)}{bcx (a + b \operatorname{sech}^{-1}(cx))} \right)$$

↓ 3779

$$ic \left( \frac{i \left( -\frac{\sinh\left(\frac{a}{b}\right) \text{Shi}\left(\frac{a}{b} + \text{sech}^{-1}(cx)\right)}{b} + \cosh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{ia}{b} + i \text{sech}^{-1}(cx) + \frac{\pi}{2}\right)}{a + b \text{sech}^{-1}(cx)} d \text{sech}^{-1}(cx) \right)}{b} - \frac{i \sqrt{\frac{1-cx}{cx+1}} (cx+1)}{bcx (a + b \text{sech}^{-1}(cx))} \right)$$

↓ 3782

$$ic \left( \frac{i \left( \frac{\cosh\left(\frac{a}{b}\right) \text{Chi}\left(\frac{a}{b} + \text{sech}^{-1}(cx)\right)}{b} - \frac{\sinh\left(\frac{a}{b}\right) \text{Shi}\left(\frac{a}{b} + \text{sech}^{-1}(cx)\right)}{b} \right)}{b} - \frac{i \sqrt{\frac{1-cx}{cx+1}} (cx+1)}{bcx (a + b \text{sech}^{-1}(cx))} \right)$$

input `Int[1/(x^2*(a + b*ArcSech[c*x])^2),x]`

output `I*c*((( -I)*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(b*c*x*(a + b*ArcSech[c*x])) + (I*((Cosh[a/b]*CoshIntegral[a/b + ArcSech[c*x]])/b - (Sinh[a/b]*SinhIntegral[a/b + ArcSech[c*x]])/b))/b)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3778 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Simp[f/(d*(m + 1)) Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]`

rule 3779 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`

rule 3782 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`

rule 3784 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`

rule 6839 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))^(n_)*(x_)^(m_.), x_Symbol] := Simp[-(c^(m + 1))^( -1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])`

## Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.91

method	result
derivativedivides	$c \left( \frac{\sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x-1}{2cxb(a+b \operatorname{arcsech}(cx))} + \frac{e^{\frac{a}{b}} \operatorname{expIntegral}_1\left(\frac{a}{b} + \operatorname{arcsech}(cx)\right)}{2b^2} + \frac{\sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x+1}{2bcx(a+b \operatorname{arcsech}(cx))} + \frac{e^{-\frac{a}{b}} \operatorname{expIntegral}_1\left(\frac{a}{b} - \operatorname{arcsech}(cx)\right)}{2b^2} \right)$
default	$c \left( \frac{\sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x-1}{2cxb(a+b \operatorname{arcsech}(cx))} + \frac{e^{\frac{a}{b}} \operatorname{expIntegral}_1\left(\frac{a}{b} + \operatorname{arcsech}(cx)\right)}{2b^2} + \frac{\sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x+1}{2bcx(a+b \operatorname{arcsech}(cx))} + \frac{e^{-\frac{a}{b}} \operatorname{expIntegral}_1\left(\frac{a}{b} - \operatorname{arcsech}(cx)\right)}{2b^2} \right)$

input `int(1/x^2/(a+b*arcsech(c*x))^2,x,method=_RETURNVERBOSE)`

output

```
c*(1/2*((-(c*x-1)/c/x)^(1/2)*c*((c*x+1)/c/x)^(1/2)*x-1)/c/x/b/(a+b*arcsech(c*x))+1/2/b^2*exp(1/b*a)*Ei(1,1/b*a+arcsech(c*x))+1/2/b*((-(c*x-1)/c/x)^(1/2)*c*((c*x+1)/c/x)^(1/2)*x+1)/c/x/(a+b*arcsech(c*x))+1/2/b^2*exp(-1/b*a)*Ei(1,-arcsech(c*x)-1/b*a))
```

**Fricas [F]**

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2 x^2} dx$$

input

```
integrate(1/x^2/(a+b*arcsech(c*x))^2,x, algorithm="fricas")
```

output

```
integral(1/(b^2*x^2*arcsech(c*x)^2 + 2*a*b*x^2*arcsech(c*x) + a^2*x^2), x)
```

**Sympy [F]**

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{x^2 (a + b \operatorname{ar} \operatorname{sech}(cx))^2} dx$$

input

```
integrate(1/x**2/(a+b*asech(c*x))**2,x)
```

output

```
Integral(1/(x**2*(a + b*asech(c*x))**2), x)
```

**Maxima [F]**

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2 x^2} dx$$

input

```
integrate(1/x^2/(a+b*arcsech(c*x))^2,x, algorithm="maxima")
```

output

```

-(c^2*x^3 + (c^2*x^3 - x)*sqrt(c*x + 1)*sqrt(-c*x + 1) - x)/((b^2*c^2*x^2
- b^2)*x^2*log(x) + ((b^2*c^2*log(c) - a*b*c^2)*x^2 - b^2*log(c) + a*b)*x^
2 - (b^2*x^2*log(x) + (b^2*log(c) - a*b)*x^2)*sqrt(c*x + 1)*sqrt(-c*x + 1)
+ (sqrt(c*x + 1)*sqrt(-c*x + 1)*b^2*x^2 - (b^2*c^2*x^2 - b^2)*x^2)*log(sq
rt(c*x + 1)*sqrt(-c*x + 1) + 1)) + integrate(-(c^4*x^4 - 2*c^2*x^2 - (c^2*
x^2 + 1)*(c*x + 1)*(c*x - 1) - (c^2*x^2 - 2)*sqrt(c*x + 1)*sqrt(-c*x + 1)
+ 1)/((b^2*c^4*x^4 - 2*b^2*c^2*x^2 + b^2)*x^2*log(x) - (b^2*x^2*log(x) + (
b^2*log(c) - a*b)*x^2)*(c*x + 1)*(c*x - 1) + ((b^2*c^4*log(c) - a*b*c^4)*x
^4 - 2*(b^2*c^2*log(c) - a*b*c^2)*x^2 + b^2*log(c) - a*b)*x^2 - 2*((b^2*c
^2*x^2 - b^2)*x^2*log(x) + ((b^2*c^2*log(c) - a*b*c^2)*x^2 - b^2*log(c) + a
*b)*x^2)*sqrt(c*x + 1)*sqrt(-c*x + 1) + ((c*x + 1)*(c*x - 1)*b^2*x^2 + 2*(
b^2*c^2*x^2 - b^2)*sqrt(c*x + 1)*sqrt(-c*x + 1)*x^2 - (b^2*c^4*x^4 - 2*b^2
*c^2*x^2 + b^2)*x^2)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)), x)

```

**Giac [F]**

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2 x^2} dx$$

input

```
integrate(1/x^2/(a+b*arcsech(c*x))^2,x, algorithm="giac")
```

output

```
integrate(1/((b*arcsech(c*x) + a)^2*x^2), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{x^2 (a + b \operatorname{acosh}(\frac{1}{cx}))^2} dx$$

input

```
int(1/(x^2*(a + b*acosh(1/(c*x)))^2), x)
```

output

```
int(1/(x^2*(a + b*acosh(1/(c*x)))^2), x)
```

**Reduce [F]**

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{\operatorname{asech}(cx)^2 b^2 x^2 + 2 \operatorname{asech}(cx) ab x^2 + a^2 x^2} dx$$

input `int(1/x^2/(a+b*asech(c*x))^2,x)`

output `int(1/(asech(c*x)**2*b**2*x**2 + 2*asech(c*x)*a*b*x**2 + a**2*x**2),x)`



**3.61** 
$$\int \frac{1}{x^3 (a+b\operatorname{sech}^{-1}(cx))^2} dx$$

Optimal result	512
Mathematica [A] (verified)	512
Rubi [C] (verified)	513
Maple [B] (verified)	517
Fricas [F]	517
Sympy [F]	518
Maxima [F]	518
Giac [F]	519
Mupad [F(-1)]	519
Reduce [F]	519

**Optimal result**

Integrand size = 14, antiderivative size = 85

$$\int \frac{1}{x^3 (a + b\operatorname{sech}^{-1}(cx))^2} dx = -\frac{c^2 \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{b^2} + \frac{c^2 \sinh\left(2\operatorname{sech}^{-1}(cx)\right)}{2b (a + b\operatorname{sech}^{-1}(cx))} + \frac{c^2 \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{b^2}$$

output

```
-c^2*cosh(2*a/b)*Chi(2*a/b+2*arcsech(c*x))/b^2+1/2*c^2*sinh(2*arcsech(c*x)
)/b/(a+b*arcsech(c*x))+c^2*sinh(2*a/b)*Shi(2*a/b+2*arcsech(c*x))/b^2
```

**Mathematica [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.08

$$\int \frac{1}{x^3 (a + b\operatorname{sech}^{-1}(cx))^2} dx = \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{x^2(a+b\operatorname{sech}^{-1}(cx))} - \frac{c^2 \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(2\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)\right) + c^2 \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(2\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)\right)}{b^2}$$

input `Integrate[1/(x^3*(a + b*ArcSech[c*x])^2),x]`

output `((b*sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(x^2*(a + b*ArcSech[c*x])) - c^2*Cosh[(2*a)/b]*CoshIntegral[2*(a/b + ArcSech[c*x])] + c^2*Sinh[(2*a)/b]*SinhIntegral[2*(a/b + ArcSech[c*x])])/b^2`

## Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.68 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.09, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.929$ , Rules used = {6839, 5971, 27, 3042, 26, 3778, 3042, 3784, 26, 3042, 26, 3779, 3782}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))^2} dx \\
 & \quad \downarrow \text{6839} \\
 & -c^2 \int \frac{\sqrt{\frac{1-cx}{cx+1}} (cx+1)}{c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^2} d \operatorname{sech}^{-1}(cx) \\
 & \quad \downarrow \text{5971} \\
 & -c^2 \int \frac{\sinh(2 \operatorname{sech}^{-1}(cx))}{2 (a + b \operatorname{sech}^{-1}(cx))^2} d \operatorname{sech}^{-1}(cx) \\
 & \quad \downarrow \text{27} \\
 & -\frac{1}{2} c^2 \int \frac{\sinh(2 \operatorname{sech}^{-1}(cx))}{(a + b \operatorname{sech}^{-1}(cx))^2} d \operatorname{sech}^{-1}(cx) \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{2} c^2 \int -\frac{i \sin(2i \operatorname{sech}^{-1}(cx))}{(a + b \operatorname{sech}^{-1}(cx))^2} d \operatorname{sech}^{-1}(cx) \\
 & \quad \downarrow \text{26}
 \end{aligned}$$

$$\frac{1}{2}ic^2 \int \frac{\sin(2i\operatorname{sech}^{-1}(cx))}{(a+b\operatorname{sech}^{-1}(cx))^2} d\operatorname{sech}^{-1}(cx)$$

↓ 3778

$$\frac{1}{2}ic^2 \left( \frac{2i \int \frac{\cosh(2\operatorname{sech}^{-1}(cx))}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx)}{b} - \frac{i \sinh(2\operatorname{sech}^{-1}(cx))}{b(a+b\operatorname{sech}^{-1}(cx))} \right)$$

↓ 3042

$$\frac{1}{2}ic^2 \left( \frac{2i \int \frac{\sin(2i\operatorname{sech}^{-1}(cx) + \frac{\pi}{2})}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx)}{b} - \frac{i \sinh(2\operatorname{sech}^{-1}(cx))}{b(a+b\operatorname{sech}^{-1}(cx))} \right)$$

↓ 3784

$$\frac{1}{2}ic^2 \left( \frac{2i \left( \cosh\left(\frac{2a}{b}\right) \int \frac{\cosh\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) + i \sinh\left(\frac{2a}{b}\right) \int \frac{i \sinh\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)}{b} - \frac{i \sinh(2\operatorname{sech}^{-1}(cx))}{b(a+b\operatorname{sech}^{-1}(cx))} \right)$$

↓ 26

$$\frac{1}{2}ic^2 \left( \frac{2i \left( \cosh\left(\frac{2a}{b}\right) \int \frac{\cosh\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) - \sinh\left(\frac{2a}{b}\right) \int \frac{\sinh\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)}{b} - \frac{i \sinh(2\operatorname{sech}^{-1}(cx))}{b(a+b\operatorname{sech}^{-1}(cx))} \right)$$

↓ 3042

$$\frac{1}{2}ic^2 \left( \frac{2i \left( \cosh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2ia}{b} + 2i\operatorname{sech}^{-1}(cx) + \frac{\pi}{2}\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) - \sinh\left(\frac{2a}{b}\right) \int \frac{i \sin\left(\frac{2ia}{b} + 2i\operatorname{sech}^{-1}(cx)\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)}{b} - \frac{i \sinh(2\operatorname{sech}^{-1}(cx))}{b(a+b\operatorname{sech}^{-1}(cx))} \right)$$

↓ 26

$$\frac{1}{2}ic^2 \left( \frac{2i \left( \cosh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2ia}{b} + 2i\operatorname{sech}^{-1}(cx) + \frac{\pi}{2}\right)}{a + b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) + i \sinh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2ia}{b} + 2i\operatorname{sech}^{-1}(cx)\right)}{a + b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)}{b} \right) - \frac{i}{b}$$

↓ 3779

$$\frac{1}{2}ic^2 \left( \frac{2i \left( -\frac{\sinh\left(\frac{2a}{b}\right)\operatorname{Shi}\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{b} + \cosh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2ia}{b} + 2i\operatorname{sech}^{-1}(cx) + \frac{\pi}{2}\right)}{a + b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)}{b} \right) - \frac{i \sinh\left(2\operatorname{sech}^{-1}(cx)\right)}{b(a + b\operatorname{sech}^{-1}(cx))}$$

↓ 3782

$$\frac{1}{2}ic^2 \left( \frac{2i \left( \frac{\cosh\left(\frac{2a}{b}\right)\operatorname{Chi}\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{b} - \frac{\sinh\left(\frac{2a}{b}\right)\operatorname{Shi}\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{b} \right)}{b} \right) - \frac{i \sinh\left(2\operatorname{sech}^{-1}(cx)\right)}{b(a + b\operatorname{sech}^{-1}(cx))}$$

input `Int[1/(x^3*(a + b*ArcSech[c*x])^2),x]`

output `(I/2)*c^2*(((-I)*Sinh[2*ArcSech[c*x]])/(b*(a + b*ArcSech[c*x])) + ((2*I)*(Cosh[(2*a)/b]*CoshIntegral[(2*a)/b + 2*ArcSech[c*x]])/b - (Sinh[(2*a)/b]*SinhIntegral[(2*a)/b + 2*ArcSech[c*x]]/b))/b)`

**Defintions of rubi rules used**

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3778 `Int[((c_.) + (d_.)*(x_)^(m_))*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Simp[f/(d*(m + 1)) Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]`

rule 3779 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`

rule 3782 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`

rule 3784 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)^(p_.)]*((c_.) + (d_.)*(x_)^(m_.))*Sinh[(a_.) + (b_.)*(x_)^(n_.)], x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6839 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))^(n_)*(x_)^(m_), x_Symbol] := Simp[-(c^(m + 1))^( -1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])`

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 185 vs.  $2(83) = 166$ .

Time = 0.60 (sec) , antiderivative size = 186, normalized size of antiderivative = 2.19

method	result
derivativedivides	$c^2 \left( \frac{2\sqrt{-\frac{cx-1}{cx}} c\sqrt{\frac{cx+1}{cx}} x + c^2 x^2 - 2}{4c^2 x^2 b(a+b \operatorname{arcsech}(cx))} + \frac{e^{\frac{2a}{b}} \operatorname{expIntegral}_1\left(\frac{2a}{b} + 2 \operatorname{arcsech}(cx)\right)}{2b^2} - \frac{c^2 x^2 - 2 - 2\sqrt{-\frac{cx-1}{cx}} c\sqrt{\frac{cx+1}{cx}} x}{4b c^2 x^2 (a+b \operatorname{arcsech}(cx))} + \dots \right)$
default	$c^2 \left( \frac{2\sqrt{-\frac{cx-1}{cx}} c\sqrt{\frac{cx+1}{cx}} x + c^2 x^2 - 2}{4c^2 x^2 b(a+b \operatorname{arcsech}(cx))} + \frac{e^{\frac{2a}{b}} \operatorname{expIntegral}_1\left(\frac{2a}{b} + 2 \operatorname{arcsech}(cx)\right)}{2b^2} - \frac{c^2 x^2 - 2 - 2\sqrt{-\frac{cx-1}{cx}} c\sqrt{\frac{cx+1}{cx}} x}{4b c^2 x^2 (a+b \operatorname{arcsech}(cx))} + \dots \right)$

input `int(1/x^3/(a+b*arcsech(c*x))^2,x,method=_RETURNVERBOSE)`

output  $c^2 * (1/4 * (2 * (-c*x-1)/c/x)^{(1/2)} * c * ((c*x+1)/c/x)^{(1/2)} * x + c^2 * x^2 - 2) / c^2 / x^2 / b / (a + b * \operatorname{arcsech}(c*x)) + 1/2 / b^2 * \exp(2/b*a) * \operatorname{Ei}(1, 2/b*a + 2 * \operatorname{arcsech}(c*x)) - 1/4 / b * (c^2 * x^2 - 2 - 2 * (-c*x-1)/c/x)^{(1/2)} * c * ((c*x+1)/c/x)^{(1/2)} * x / c^2 / x^2 / (a + b * \operatorname{arcsech}(c*x)) + 1/2 / b^2 * \exp(-2/b*a) * \operatorname{Ei}(1, -2 * \operatorname{arcsech}(c*x) - 2/b*a)$

**Fricas [F]**

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2 x^3} dx$$

input `integrate(1/x^3/(a+b*arcsech(c*x))^2,x, algorithm="fricas")`

output `integral(1/(b^2*x^3*arcsech(c*x)^2 + 2*a*b*x^3*arcsech(c*x) + a^2*x^3), x)`

**Sympy [F]**

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{x^3 (a + b \operatorname{arsech}(cx))^2} dx$$

input `integrate(1/x**3/(a+b*asech(c*x))**2,x)`

output `Integral(1/(x**3*(a + b*asech(c*x))**2), x)`

**Maxima [F]**

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(b \operatorname{arsech}(cx) + a)^2 x^3} dx$$

input `integrate(1/x^3/(a+b*arcsech(c*x))^2,x, algorithm="maxima")`

output `-(c^2*x^3 + (c^2*x^3 - x)*sqrt(c*x + 1)*sqrt(-c*x + 1) - x)/((b^2*c^2*x^2 - b^2)*x^3*log(x) + ((b^2*c^2*log(c) - a*b*c^2)*x^2 - b^2*log(c) + a*b)*x^3 - (b^2*x^3*log(x) + (b^2*log(c) - a*b)*x^3)*sqrt(c*x + 1)*sqrt(-c*x + 1) + (sqrt(c*x + 1)*sqrt(-c*x + 1)*b^2*x^3 - (b^2*c^2*x^2 - b^2)*x^3)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)) + integrate(-(2*c^4*x^4 - 4*c^2*x^2 - 2*(c*x + 1)*(c*x - 1) + (c^4*x^4 - 4*c^2*x^2 + 4)*sqrt(c*x + 1)*sqrt(-c*x + 1) + 2)/((b^2*c^4*x^4 - 2*b^2*c^2*x^2 + b^2)*x^3*log(x) + ((b^2*c^4*log(c) - a*b*c^4)*x^4 - 2*(b^2*c^2*log(c) - a*b*c^2)*x^2 + b^2*log(c) - a*b)*x^3 - (b^2*x^3*log(x) + (b^2*log(c) - a*b)*x^3)*(c*x + 1)*(c*x - 1) - 2*((b^2*c^2*x^2 - b^2)*x^3*log(x) + ((b^2*c^2*log(c) - a*b*c^2)*x^2 - b^2*log(c) + a*b)*x^3)*sqrt(c*x + 1)*sqrt(-c*x + 1) + ((c*x + 1)*(c*x - 1)*b^2*x^3 + 2*(b^2*c^2*x^2 - b^2)*sqrt(c*x + 1)*sqrt(-c*x + 1)*x^3 - (b^2*c^4*x^4 - 2*b^2*c^2*x^2 + b^2)*x^3)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)), x)`

**Giac [F]**

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(b \operatorname{arsech}(cx) + a)^2 x^3} dx$$

input `integrate(1/x^3/(a+b*arcsech(c*x))^2,x, algorithm="giac")`

output `integrate(1/((b*arcsech(c*x) + a)^2*x^3), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{x^3 (a + b \operatorname{acosh}(\frac{1}{cx}))^2} dx$$

input `int(1/(x^3*(a + b*acosh(1/(c*x)))^2),x)`

output `int(1/(x^3*(a + b*acosh(1/(c*x)))^2), x)`

**Reduce [F]**

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{\operatorname{asech}(cx)^2 b^2 x^3 + 2 \operatorname{asech}(cx) a b x^3 + a^2 x^3} dx$$

input `int(1/x^3/(a+b*asech(c*x))^2,x)`

output `int(1/(asech(c*x)**2*b**2*x**3 + 2*asech(c*x)*a*b*x**3 + a**2*x**3),x)`



**3.62**  $\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^2} dx$

Optimal result . . . . .	520
Mathematica [A] (verified) . . . . .	521
Rubi [A] (verified) . . . . .	521
Maple [B] (verified) . . . . .	523
Fricas [F] . . . . .	523
Sympy [F] . . . . .	524
Maxima [F] . . . . .	524
Giac [F] . . . . .	525
Mupad [F(-1)] . . . . .	525
Reduce [F] . . . . .	525

**Optimal result**

Integrand size = 14, antiderivative size = 190

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \frac{c^2 \sqrt{\frac{1-cx}{1+cx}} (1+cx)}{4bx (a + b \operatorname{sech}^{-1}(cx))} - \frac{c^3 \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{4b^2}$$

$$- \frac{3c^3 \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3a}{b} + 3\operatorname{sech}^{-1}(cx)\right)}{4b^2}$$

$$+ \frac{c^3 \sinh\left(3\operatorname{sech}^{-1}(cx)\right)}{4b (a + b \operatorname{sech}^{-1}(cx))}$$

$$+ \frac{c^3 \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{4b^2}$$

$$+ \frac{3c^3 \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3a}{b} + 3\operatorname{sech}^{-1}(cx)\right)}{4b^2}$$

output

```
1/4*c^2*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/b/x/(a+b*arcsech(c*x))-1/4*c^3*cosh(a/b)*Chi(a/b+arcsech(c*x))/b^2-3/4*c^3*cosh(3*a/b)*Chi(3*a/b+3*arcsech(c*x))/b^2+1/4*c^3*sinh(3*arcsech(c*x))/b/(a+b*arcsech(c*x))+1/4*c^3*sinh(a/b)*Shi(a/b+arcsech(c*x))/b^2+3/4*c^3*sinh(3*a/b)*Shi(3*a/b+3*arcsech(c*x))/b^2
```

**Mathematica [A] (verified)**

Time = 0.57 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.32

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^2} dx$$

$$= \frac{4b \sqrt{\frac{1-cx}{1+cx}} + 4bcx \sqrt{\frac{1-cx}{1+cx}} - c^3 x^3 (a + b \operatorname{sech}^{-1}(cx)) \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right) - 3c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))}{4b^2 x^3 (a + b \operatorname{sech}^{-1}(cx))^2}$$

input `Integrate[1/(x^4*(a + b*ArcSech[c*x])^2),x]`

output `(4*b*Sqrt[(1 - c*x)/(1 + c*x)] + 4*b*c*x*Sqrt[(1 - c*x)/(1 + c*x)] - c^3*x^3*(a + b*ArcSech[c*x])*Cosh[a/b]*CoshIntegral[a/b + ArcSech[c*x]] - 3*c^3*x^3*(a + b*ArcSech[c*x])*Cosh[(3*a)/b]*CoshIntegral[3*(a/b + ArcSech[c*x])] + a*c^3*x^3*Sinh[a/b]*SinhIntegral[a/b + ArcSech[c*x]] + b*c^3*x^3*ArcSech[c*x]*Sinh[a/b]*SinhIntegral[a/b + ArcSech[c*x]] + 3*a*c^3*x^3*Sinh[(3*a)/b]*SinhIntegral[3*(a/b + ArcSech[c*x])] + 3*b*c^3*x^3*ArcSech[c*x]*Sinh[(3*a)/b]*SinhIntegral[3*(a/b + ArcSech[c*x])])/(4*b^2*x^3*(a + b*ArcSech[c*x]))`

**Rubi [A] (verified)**

Time = 0.57 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {6839, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^2} dx$$

$$\downarrow \text{6839}$$

$$-c^3 \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))^2} d \operatorname{sech}^{-1}(cx)$$

$$-c^3 \int \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{4cx(a+b\operatorname{sech}^{-1}(cx))^2} + \frac{\sinh(3\operatorname{sech}^{-1}(cx))}{4(a+b\operatorname{sech}^{-1}(cx))^2} \right) d\operatorname{sech}^{-1}(cx)$$

↓ 5971

↓ 2009

$$-c^3 \left( \frac{\cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{4b^2} + \frac{3 \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3a}{b} + 3\operatorname{sech}^{-1}(cx)\right)}{4b^2} - \frac{\sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{4b^2} - \frac{3 \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3a}{b} + 3\operatorname{sech}^{-1}(cx)\right)}{4b^2} \right)$$

input `Int[1/(x^4*(a + b*ArcSech[c*x])^2),x]`

output `-(c^3*(-1/4*(Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(b*c*x*(a + b*ArcSech[c*x])) + (Cosh[a/b]*CoshIntegral[a/b + ArcSech[c*x]])/(4*b^2) + (3*Cosh[(3*a)/b]*CoshIntegral[(3*a)/b + 3*ArcSech[c*x]])/(4*b^2) - Sinh[3*ArcSech[c*x]]/(4*b*(a + b*ArcSech[c*x])) - (Sinh[a/b]*SinhIntegral[a/b + ArcSech[c*x]])/(4*b^2) - (3*Sinh[(3*a)/b]*SinhIntegral[(3*a)/b + 3*ArcSech[c*x]])/(4*b^2))`

### Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^(n)*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6839 `Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.), x_Symbol] := Simp[-(c^(m + 1))^(1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])`

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 419 vs.  $2(176) = 352$ .

Time = 0.78 (sec) , antiderivative size = 420, normalized size of antiderivative = 2.21

method	result
derivativedivides	$c^3 \left( -\frac{\sqrt{-\frac{cx-1}{cx}} c^3 x^3 \sqrt{\frac{cx+1}{cx}} - 4\sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x - 3c^2 x^2 + 4}{8c^3 x^3 b(a+b \operatorname{arcsech}(cx))} + \frac{3e^{\frac{3a}{b}} \operatorname{expIntegral}_1\left(\frac{3a}{b} + 3 \operatorname{arcsech}(cx)\right)}{8b^2} + \frac{\sqrt{-\frac{cx-1}{cx}}}{8cx} \right)$
default	$c^3 \left( -\frac{\sqrt{-\frac{cx-1}{cx}} c^3 x^3 \sqrt{\frac{cx+1}{cx}} - 4\sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x - 3c^2 x^2 + 4}{8c^3 x^3 b(a+b \operatorname{arcsech}(cx))} + \frac{3e^{\frac{3a}{b}} \operatorname{expIntegral}_1\left(\frac{3a}{b} + 3 \operatorname{arcsech}(cx)\right)}{8b^2} + \frac{\sqrt{-\frac{cx-1}{cx}}}{8cx} \right)$

input `int(1/x^4/(a+b*arcsech(c*x))^2,x,method=_RETURNVERBOSE)`

output

$$c^3 * (-1/8 * ((- (c*x-1)/c/x)^(1/2) * c^3 * x^3 * ((c*x+1)/c/x)^(1/2) - 4 * (- (c*x-1)/c/x)^(1/2) * c * ((c*x+1)/c/x)^(1/2) * x - 3 * c^2 * x^2 + 4) / c^3 / x^3 / b / (a + b * \operatorname{arcsech}(c*x)) + 3/8 / b^2 * \operatorname{exp}(3/b*a) * \operatorname{Ei}(1, 3/b*a + 3 * \operatorname{arcsech}(c*x)) + 1/8 * ((- (c*x-1)/c/x)^(1/2) * c * ((c*x+1)/c/x)^(1/2) * x - 1) / c/x / b / (a + b * \operatorname{arcsech}(c*x)) + 1/8 / b^2 * \operatorname{exp}(1/b*a) * \operatorname{Ei}(1, 1/b*a + \operatorname{arcsech}(c*x)) + 1/8 / b * ((- (c*x-1)/c/x)^(1/2) * c * ((c*x+1)/c/x)^(1/2) * x + 1) / c/x / (a + b * \operatorname{arcsech}(c*x)) + 1/8 / b^2 * \operatorname{exp}(-1/b*a) * \operatorname{Ei}(1, -\operatorname{arcsech}(c*x) - 1/b*a) - 1/8 / b * ((- (c*x-1)/c/x)^(1/2) * c^3 * x^3 * ((c*x+1)/c/x)^(1/2) - 4 * (- (c*x-1)/c/x)^(1/2) * c * ((c*x+1)/c/x)^(1/2) * x + 3 * c^2 * x^2 - 4) / c^3 / x^3 / (a + b * \operatorname{arcsech}(c*x)) + 3/8 / b^2 * \operatorname{exp}(-3/b*a) * \operatorname{Ei}(1, -3 * \operatorname{arcsech}(c*x) - 3/b*a))$$
**Fricas [F]**

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2 x^4} dx$$

input `integrate(1/x^4/(a+b*arcsech(c*x))^2,x, algorithm="fricas")`

output `integral(1/(b^2*x^4*arcsech(c*x)^2 + 2*a*b*x^4*arcsech(c*x) + a^2*x^4), x)`

**Sympy [F]**

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{x^4 (a + b \operatorname{arsech}(cx))^2} dx$$

input `integrate(1/x**4/(a+b*asech(c*x))**2,x)`

output `Integral(1/(x**4*(a + b*asech(c*x))**2), x)`

**Maxima [F]**

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(b \operatorname{arsech}(cx) + a)^2 x^4} dx$$

input `integrate(1/x^4/(a+b*arcsech(c*x))^2,x, algorithm="maxima")`

output `-(c^2*x^3 + (c^2*x^3 - x)*sqrt(c*x + 1)*sqrt(-c*x + 1) - x)/((b^2*c^2*x^2 - b^2)*x^4*log(x) + ((b^2*c^2*log(c) - a*b*c^2)*x^2 - b^2*log(c) + a*b)*x^4 - (b^2*x^4*log(x) + (b^2*log(c) - a*b)*x^4)*sqrt(c*x + 1)*sqrt(-c*x + 1) + (sqrt(c*x + 1)*sqrt(-c*x + 1)*b^2*x^4 - (b^2*c^2*x^2 - b^2)*x^4)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)) - integrate((3*c^4*x^4 - 6*c^2*x^2 + (c^2*x^2 - 3)*(c*x + 1)*(c*x - 1) + (2*c^4*x^4 - 7*c^2*x^2 + 6)*sqrt(c*x + 1)*sqrt(-c*x + 1) + 3)/((b^2*c^4*x^4 - 2*b^2*c^2*x^2 + b^2)*x^4*log(x) + ((b^2*c^4*log(c) - a*b*c^4)*x^4 - 2*(b^2*c^2*log(c) - a*b*c^2)*x^2 + b^2*log(c) - a*b)*x^4 - (b^2*x^4*log(x) + (b^2*log(c) - a*b)*x^4)*(c*x + 1)*(c*x - 1) - 2*((b^2*c^2*x^2 - b^2)*x^4*log(x) + ((b^2*c^2*log(c) - a*b*c^2)*x^2 - b^2*log(c) + a*b)*x^4)*sqrt(c*x + 1)*sqrt(-c*x + 1) + ((c*x + 1)*(c*x - 1)*b^2*x^4 + 2*(b^2*c^2*x^2 - b^2)*sqrt(c*x + 1)*sqrt(-c*x + 1)*x^4 - (b^2*c^4*x^4 - 2*b^2*c^2*x^2 + b^2)*x^4)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)), x)`

**Giac [F]**

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2 x^4} dx$$

input `integrate(1/x^4/(a+b*arcsech(c*x))^2,x, algorithm="giac")`

output `integrate(1/((b*arcsech(c*x) + a)^2*x^4), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{x^4 (a + b \operatorname{acosh}(\frac{1}{cx}))^2} dx$$

input `int(1/(x^4*(a + b*acosh(1/(c*x)))^2),x)`

output `int(1/(x^4*(a + b*acosh(1/(c*x)))^2), x)`

**Reduce [F]**

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{1}{\operatorname{asech}(cx)^2 b^2 x^4 + 2 \operatorname{asech}(cx) a b x^4 + a^2 x^4} dx$$

input `int(1/x^4/(a+b*asech(c*x))^2,x)`

output `int(1/(asech(c*x)**2*b**2*x**4 + 2*asech(c*x)*a*b*x**4 + a**2*x**4),x)`

$$3.63 \quad \int \frac{x}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^3} dx$$

Optimal result	526
Mathematica [N/A]	526
Rubi [N/A]	527
Maple [N/A]	527
Fricas [N/A]	528
Sympy [N/A]	528
Maxima [N/A]	528
Giac [N/A]	529
Mupad [N/A]	530
Reduce [N/A]	530

### Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{x}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^3} dx = \operatorname{Int}\left(\frac{x}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^3}, x\right)$$

output `Defer(Int)(x/(a+b*arcsech(c*x))^3,x)`

### Mathematica [N/A]

Not integrable

Time = 3.81 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{x}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^3} dx = \int \frac{x}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^3} dx$$

input `Integrate[x/(a + b*ArcSech[c*x])^3,x]`

output `Integrate[x/(a + b*ArcSech[c*x])^3, x]`

**Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^3} dx$$

↓ 6865

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^3} dx$$

input `Int[x/(a + b*ArcSech[c*x])^3,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{x}{(a + b \operatorname{arcsech}(cx))^3} dx$$

input `int(x/(a+b*arcsech(c*x))^3,x)`

output `int(x/(a+b*arcsech(c*x))^3,x)`



**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 42, normalized size of antiderivative = 3.50

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{x}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3} dx$$

input `integrate(x/(a+b*arcsech(c*x))^3,x, algorithm="fricas")`

output `integral(x/(b^3*arcsech(c*x)^3 + 3*a*b^2*arcsech(c*x)^2 + 3*a^2*b*arcsech(c*x) + a^3), x)`

**Sympy [N/A]**

Not integrable

Time = 1.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{x}{(a + b \operatorname{ar} \operatorname{sech}(cx))^3} dx$$

input `integrate(x/(a+b*asech(c*x))**3,x)`

output `Integral(x/(a + b*asech(c*x))**3, x)`

**Maxima [N/A]**

Not integrable

Time = 1.99 (sec) , antiderivative size = 2818, normalized size of antiderivative = 234.83

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{x}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3} dx$$

input `integrate(x/(a+b*arcsech(c*x))^3,x, algorithm="maxima")`

output

$$\begin{aligned}
 & -1/2*((2*(2*b*c^4*x^5 - 3*b*c^2*x^3 + b*x)*x*\log(x) + (4*(b*c^4*\log(c) - a \\
 & *c^4)*x^5 - (b*c^2*(6*\log(c) + 1) - 6*a*c^2)*x^3 + (b*(2*\log(c) + 1) - 2*a \\
 & )*x)*x)*(c*x + 1)^(3/2)*(-c*x + 1)^(3/2) + (3*(b*c^6*x^7 - 5*b*c^4*x^5 + 6 \\
 & *b*c^2*x^3 - 2*b*x)*x*\log(x) + (3*(b*c^6*\log(c) - a*c^6)*x^7 - (b*c^4*(15* \\
 & \log(c) + 2) - 15*a*c^4)*x^5 + (b*c^2*(18*\log(c) + 5) - 18*a*c^2)*x^3 - 3*( \\
 & b*(2*\log(c) + 1) - 2*a)*x)*x)*(c*x + 1)*(c*x - 1) - 2*(b*c^6*x^7 - 3*b*c^4 \\
 & *x^5 + 3*b*c^2*x^3 - b*x)*x*\log(x) - ((5*b*c^6*x^7 - 17*b*c^4*x^5 + 18*b*c \\
 & ^2*x^3 - 6*b*x)*x*\log(x) + ((b*c^6*(5*\log(c) + 1) - 5*a*c^6)*x^7 - (b*c^4* \\
 & (17*\log(c) + 5) - 17*a*c^4)*x^5 + (b*c^2*(18*\log(c) + 7) - 18*a*c^2)*x^3 - \\
 & 3*(b*(2*\log(c) + 1) - 2*a)*x)*x)*\sqrt{c*x + 1}*\sqrt{-c*x + 1} - ((b*c^6*( \\
 & 2*\log(c) + 1) - 2*a*c^6)*x^7 - 3*(b*c^4*(2*\log(c) + 1) - 2*a*c^4)*x^5 + 3* \\
 & (b*c^2*(2*\log(c) + 1) - 2*a*c^2)*x^3 - (b*(2*\log(c) + 1) - 2*a)*x)*x - (2* \\
 & (2*b*c^4*x^5 - 3*b*c^2*x^3 + b*x)*(c*x + 1)^(3/2)*(-c*x + 1)^(3/2)*x + 3*( \\
 & b*c^6*x^7 - 5*b*c^4*x^5 + 6*b*c^2*x^3 - 2*b*x)*(c*x + 1)*(c*x - 1)*x - (5* \\
 & b*c^6*x^7 - 17*b*c^4*x^5 + 18*b*c^2*x^3 - 6*b*x)*\sqrt{c*x + 1}*\sqrt{-c*x + \\
 & 1)*x - 2*(b*c^6*x^7 - 3*b*c^4*x^5 + 3*b*c^2*x^3 - b*x)*x*\log(\sqrt{c*x + \\
 & 1}*\sqrt{-c*x + 1} + 1))/((b^4*c^6*\log(c)^2 - 2*a*b^3*c^6*\log(c) + a^2*b^2*c \\
 & ^6)*x^6 - b^4*\log(c)^2 - 3*(b^4*c^4*\log(c)^2 - 2*a*b^3*c^4*\log(c) + a^2*b \\
 & ^2*c^4)*x^4 + 2*a*b^3*\log(c) - (b^4*\log(c)^2 + b^4*\log(x)^2 - 2*a*b^3*\log( \\
 & c) + a^2*b^2 + 2*(b^4*\log(c) - a*b^3)*\log(x))*(c*x + 1)^(3/2)*(-c*x + 1...
 \end{aligned}$$

**Giac [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{x}{(b \operatorname{arsech}(cx) + a)^3} dx$$

input `integrate(x/(a+b*arcsech(c*x))^3,x, algorithm="giac")`

output `integrate(x/(b*arcsech(c*x) + a)^3, x)`

**Mupad [N/A]**

Not integrable

Time = 3.66 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.50

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{x}{(a + b \operatorname{acosh}(\frac{1}{cx}))^3} dx$$

input `int(x/(a + b*acosh(1/(c*x)))^3,x)`output `int(x/(a + b*acosh(1/(c*x)))^3, x)`**Reduce [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 42, normalized size of antiderivative = 3.50

$$\int \frac{x}{(a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{x}{\operatorname{asech}(cx)^3 b^3 + 3 \operatorname{asech}(cx)^2 a b^2 + 3 \operatorname{asech}(cx) a^2 b + a^3} dx$$

input `int(x/(a+b*asech(c*x))^3,x)`output `int(x/(asech(c*x)**3*b**3 + 3*asech(c*x)**2*a*b**2 + 3*asech(c*x)*a**2*b + a**3),x)`

$$3.64 \quad \int \frac{1}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^3} dx$$

Optimal result	531
Mathematica [N/A]	531
Rubi [N/A]	532
Maple [N/A]	532
Fricas [N/A]	533
Sympy [N/A]	533
Maxima [N/A]	533
Giac [N/A]	534
Mupad [N/A]	535
Reduce [N/A]	535

### Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{1}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^3} dx = \operatorname{Int}\left(\frac{1}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^3}, x\right)$$

output `Defer(Int)(1/(a+b*arcsech(c*x))^3,x)`

### Mathematica [N/A]

Not integrable

Time = 88.39 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{1}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^3} dx = \int \frac{1}{\left(a+b\operatorname{sech}^{-1}(cx)\right)^3} dx$$

input `Integrate[(a + b*ArcSech[c*x])^(-3), x]`

output `Integrate[(a + b*ArcSech[c*x])^(-3), x]`

**Rubi [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^3} dx$$

↓ 6865

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^3} dx$$

input `Int[(a + b*ArcSech[c*x])^(-3),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.20 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a + b \operatorname{arcsech}(cx))^3} dx$$

input `int(1/(a+b*arcsech(c*x))^3,x)`

output `int(1/(a+b*arcsech(c*x))^3,x)`

**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 40, normalized size of antiderivative = 4.00

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3} dx$$

input `integrate(1/(a+b*arcsech(c*x))^3,x, algorithm="fricas")`

output `integral(1/(b^3*arcsech(c*x)^3 + 3*a*b^2*arcsech(c*x)^2 + 3*a^2*b*arcsech(c*x) + a^3), x)`

**Sympy [N/A]**

Not integrable

Time = 1.31 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(a + b \operatorname{ar} \operatorname{sech}(cx))^3} dx$$

input `integrate(1/(a+b*asech(c*x))**3,x)`

output `Integral((a + b*asech(c*x))**(-3), x)`

**Maxima [N/A]**

Not integrable

Time = 1.94 (sec) , antiderivative size = 2771, normalized size of antiderivative = 277.10

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3} dx$$

input `integrate(1/(a+b*arcsech(c*x))^3,x, algorithm="maxima")`

output

```

1/2*((b*c^6*(log(c) + 1) - a*c^6)*x^7 - 3*(b*c^4*(log(c) + 1) - a*c^4)*x^5
- (3*(b*c^4*log(c) - a*c^4)*x^5 - (b*c^2*(4*log(c) + 1) - 4*a*c^2)*x^3 +
(b*(log(c) + 1) - a)*x + (3*b*c^4*x^5 - 4*b*c^2*x^3 + b*x)*log(x))*(c*x +
1)^(3/2)*(-c*x + 1)^(3/2) + 3*(b*c^2*(log(c) + 1) - a*c^2)*x^3 - (2*(b*c^6
*log(c) - a*c^6)*x^7 - 2*(b*c^4*(5*log(c) + 1) - 5*a*c^4)*x^5 + (b*c^2*(11
*log(c) + 5) - 11*a*c^2)*x^3 - 3*(b*(log(c) + 1) - a)*x + (2*b*c^6*x^7 - 1
0*b*c^4*x^5 + 11*b*c^2*x^3 - 3*b*x)*log(x))*(c*x + 1)*(c*x - 1) + ((b*c^6*
(3*log(c) + 1) - 3*a*c^6)*x^7 - 5*(b*c^4*(2*log(c) + 1) - 2*a*c^4)*x^5 + (
b*c^2*(10*log(c) + 7) - 10*a*c^2)*x^3 - 3*(b*(log(c) + 1) - a)*x + (3*b*c^
6*x^7 - 10*b*c^4*x^5 + 10*b*c^2*x^3 - 3*b*x)*log(x))*sqrt(c*x + 1)*sqrt(-c
*x + 1) - (b*(log(c) + 1) - a)*x - (b*c^6*x^7 - 3*b*c^4*x^5 + 3*b*c^2*x^3
- (3*b*c^4*x^5 - 4*b*c^2*x^3 + b*x)*(c*x + 1)^(3/2)*(-c*x + 1)^(3/2) - (2*
b*c^6*x^7 - 10*b*c^4*x^5 + 11*b*c^2*x^3 - 3*b*x)*(c*x + 1)*(c*x - 1) + (3*
b*c^6*x^7 - 10*b*c^4*x^5 + 10*b*c^2*x^3 - 3*b*x)*sqrt(c*x + 1)*sqrt(-c*x +
1) - b*x)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1) + (b*c^6*x^7 - 3*b*c^4*x^
5 + 3*b*c^2*x^3 - b*x)*log(x))/((b^4*c^6*log(c)^2 - 2*a*b^3*c^6*log(c) + a
^2*b^2*c^6)*x^6 - b^4*log(c)^2 - 3*(b^4*c^4*log(c)^2 - 2*a*b^3*c^4*log(c)
+ a^2*b^2*c^4)*x^4 + 2*a*b^3*log(c) - (b^4*log(c)^2 + b^4*log(x)^2 - 2*a*b
^3*log(c) + a^2*b^2 + 2*(b^4*log(c) - a*b^3)*log(x))*(c*x + 1)^(3/2)*(-c*x
+ 1)^(3/2) - a^2*b^2 + 3*(b^4*log(c)^2 - 2*a*b^3*log(c) + a^2*b^2 - (b...

```

**Giac [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(b \operatorname{arsech}(cx) + a)^3} dx$$

input `integrate(1/(a+b*arcsech(c*x))^3,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)^(-3), x)`

**Mupad [N/A]**

Not integrable

Time = 3.64 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.60

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(a + b \operatorname{acosh}(\frac{1}{cx}))^3} dx$$

input `int(1/(a + b*acosh(1/(c*x)))^3,x)`output `int(1/(a + b*acosh(1/(c*x)))^3, x)`**Reduce [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 40, normalized size of antiderivative = 4.00

$$\int \frac{1}{(a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{\operatorname{asech}(cx)^3 b^3 + 3 \operatorname{asech}(cx)^2 a b^2 + 3 \operatorname{asech}(cx) a^2 b + a^3} dx$$

input `int(1/(a+b*asech(c*x))^3,x)`output `int(1/(asech(c*x)**3*b**3 + 3*asech(c*x)**2*a*b**2 + 3*asech(c*x)*a**2*b + a**3),x)`



$$3.65 \quad \int \frac{1}{x \left( a + b \operatorname{sech}^{-1}(cx) \right)^3} dx$$

Optimal result	536
Mathematica [N/A]	536
Rubi [N/A]	537
Maple [N/A]	537
Fricas [N/A]	538
Sympy [N/A]	538
Maxima [N/A]	538
Giac [N/A]	539
Mupad [N/A]	540
Reduce [N/A]	540

### Optimal result

Integrand size = 14, antiderivative size = 14

$$\int \frac{1}{x \left( a + b \operatorname{sech}^{-1}(cx) \right)^3} dx = \operatorname{Int} \left( \frac{1}{x \left( a + b \operatorname{sech}^{-1}(cx) \right)^3}, x \right)$$

output `Defer(Int)(1/x/(a+b*arcsech(c*x))^3,x)`

### Mathematica [N/A]

Not integrable

Time = 1.64 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{1}{x \left( a + b \operatorname{sech}^{-1}(cx) \right)^3} dx = \int \frac{1}{x \left( a + b \operatorname{sech}^{-1}(cx) \right)^3} dx$$

input `Integrate[1/(x*(a + b*ArcSech[c*x])^3),x]`

output `Integrate[1/(x*(a + b*ArcSech[c*x])^3), x]`

**Rubi [N/A]**

Not integrable

Time = 0.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x (a + b \operatorname{sech}^{-1}(cx))^3} dx$$

↓ 6865

$$\int \frac{1}{x (a + b \operatorname{sech}^{-1}(cx))^3} dx$$

input `Int[1/(x*(a + b*ArcSech[c*x])^3),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{x (a + b \operatorname{arcsech}(cx))^3} dx$$

input `int(1/x/(a+b*arcsech(c*x))^3,x)`

output `int(1/x/(a+b*arcsech(c*x))^3,x)`

**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 45, normalized size of antiderivative = 3.21

$$\int \frac{1}{x (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3 x} dx$$

input `integrate(1/x/(a+b*arcsech(c*x))^3,x, algorithm="fricas")`

output `integral(1/(b^3*x*arcsech(c*x)^3 + 3*a*b^2*x*arcsech(c*x)^2 + 3*a^2*b*x*arcsech(c*x) + a^3*x), x)`

**Sympy [N/A]**

Not integrable

Time = 2.47 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{x (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{x (a + b \operatorname{ar} \operatorname{sech}(cx))^3} dx$$

input `integrate(1/x/(a+b*asech(c*x))**3,x)`

output `Integral(1/(x*(a + b*asech(c*x))**3), x)`

**Maxima [N/A]**

Not integrable

Time = 1.78 (sec) , antiderivative size = 2638, normalized size of antiderivative = 188.43

$$\int \frac{1}{x (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3 x} dx$$

input `integrate(1/x/(a+b*arcsech(c*x))^3,x, algorithm="maxima")`

output

$$\begin{aligned} & -1/2*(b*c^6*x^7 - 3*b*c^4*x^5 + 3*b*c^2*x^3 - (2*(b*c^4*\log(c) - a*c^4)*x^5 \\ & - (b*c^2*(2*\log(c) + 1) - 2*a*c^2)*x^3 + b*x + 2*(b*c^4*x^5 - b*c^2*x^3) \\ & * \log(x))*(c*x + 1)^{(3/2)}*(-c*x + 1)^{(3/2)} - ((b*c^6*\log(c) - a*c^6)*x^7 - \\ & (b*c^4*(5*\log(c) + 2) - 5*a*c^4)*x^5 + (b*c^2*(4*\log(c) + 5) - 4*a*c^2)*x^3 \\ & - 3*b*x + (b*c^6*x^7 - 5*b*c^4*x^5 + 4*b*c^2*x^3)*\log(x))*(c*x + 1)*(c*x \\ & - 1) + ((b*c^6*(\log(c) + 1) - a*c^6)*x^7 - (b*c^4*(3*\log(c) + 5) - 3*a*c^4) \\ & *x^5 + (b*c^2*(2*\log(c) + 7) - 2*a*c^2)*x^3 - 3*b*x + (b*c^6*x^7 - 3*b*c^4 \\ & *x^5 + 2*b*c^2*x^3)*\log(x))*\sqrt{c*x + 1}*\sqrt{-c*x + 1} - b*x + (2*(b*c^4 \\ & *x^5 - b*c^2*x^3)*(c*x + 1)^{(3/2)}*(-c*x + 1)^{(3/2)} + (b*c^6*x^7 - 5*b*c^4 \\ & *x^5 + 4*b*c^2*x^3)*(c*x + 1)*(c*x - 1) - (b*c^6*x^7 - 3*b*c^4*x^5 + 2*b*c^2 \\ & *x^3)*\sqrt{c*x + 1}*\sqrt{-c*x + 1})*\log(\sqrt{c*x + 1}*\sqrt{-c*x + 1} + 1))/((b^4*x \\ & *\log(x)^2 + 2*(b^4*\log(c) - a*b^3)*x*\log(x) + (b^4*\log(c)^2 - 2*a*b^3*\log(c) \\ & + a^2*b^2)*x)*(c*x + 1)^{(3/2)}*(-c*x + 1)^{(3/2)} - (b^4*c^6*x^6 - 3*b^4*c^4*x^4 \\ & + 3*b^4*c^2*x^2 - b^4)*x*\log(x)^2 + 3*((b^4*c^2*x^2 - b^4)*x*\log(x)^2 - 2*(b^4 \\ & *\log(c) - a*b^3 - (b^4*c^2*\log(c) - a*b^3*c^2)*x^2)*x*\log(x) - (b^4*\log(c)^2 \\ & - 2*a*b^3*\log(c) + a^2*b^2 - (b^4*c^2*\log(c)^2 - 2*a*b^3*c^2*\log(c) + a^2*b^2*c^2) \\ & *x^2)*x)*(c*x + 1)*(c*x - 1) + ((c*x + 1)^{(3/2)}*(-c*x + 1)^{(3/2)}*b^4*x \\ & + 3*(b^4*c^2*x^2 - b^4)*(c*x + 1)*(c*x - 1)*x + 3*(b^4*c^4*x^4 - 2*b^4*c^2*x^2 \\ & + b^4)*\sqrt{c*x + 1}*\sqrt{-c*x + 1}*x - (b^4*c^6*x^6 - 3*b^4*c^4*x^4 + 3*b^4*c^2*x^2 - b^4)*x) \\ & *\log(\sqrt{c*x + 1}*\sqrt{-c*x + 1}) \end{aligned}$$

**Giac [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{1}{x(a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3 x} dx$$

input `integrate(1/x/(a+b*arcsech(c*x))^3,x, algorithm="giac")`

output `integrate(1/((b*arcsech(c*x) + a)^3*x), x)`

**Mupad [N/A]**

Not integrable

Time = 3.68 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.43

$$\int \frac{1}{x (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{x (a + b \operatorname{acosh}(\frac{1}{cx}))^3} dx$$

input `int(1/(x*(a + b*acosh(1/(c*x)))^3),x)`output `int(1/(x*(a + b*acosh(1/(c*x)))^3), x)`**Reduce [N/A]**

Not integrable

Time = 0.23 (sec) , antiderivative size = 45, normalized size of antiderivative = 3.21

$$\begin{aligned} & \int \frac{1}{x (a + b \operatorname{sech}^{-1}(cx))^3} dx \\ &= \int \frac{1}{a \operatorname{sech}(cx)^3 b^3 x + 3 a \operatorname{sech}(cx)^2 a b^2 x + 3 a \operatorname{sech}(cx) a^2 b x + a^3 x} dx \end{aligned}$$

input `int(1/x/(a+b*asech(c*x))^3,x)`output `int(1/(asech(c*x)**3*b**3*x + 3*asech(c*x)**2*a*b**2*x + 3*asech(c*x)*a**2*b*x + a**3*x),x)`

**3.66**  $\int \frac{1}{x^2 (a+b\operatorname{sech}^{-1}(cx))^3} dx$

Optimal result . . . . .	541
Mathematica [A] (verified) . . . . .	542
Rubi [C] (verified) . . . . .	542
Maple [B] (verified) . . . . .	547
Fricas [F] . . . . .	548
Sympy [F] . . . . .	548
Maxima [F] . . . . .	548
Giac [F] . . . . .	549
Mupad [F(-1)] . . . . .	550
Reduce [F] . . . . .	550

**Optimal result**

Integrand size = 14, antiderivative size = 114

$$\int \frac{1}{x^2 (a+b\operatorname{sech}^{-1}(cx))^3} dx = \frac{\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{2bx (a+b\operatorname{sech}^{-1}(cx))^2} + \frac{1}{2b^2x (a+b\operatorname{sech}^{-1}(cx))} + \frac{c\operatorname{Chi}(\frac{a}{b} + \operatorname{sech}^{-1}(cx)) \sinh(\frac{a}{b})}{2b^3} - \frac{c \cosh(\frac{a}{b}) \operatorname{Shi}(\frac{a}{b} + \operatorname{sech}^{-1}(cx))}{2b^3}$$

output

```
1/2*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/b/x/(a+b*arcsech(c*x))^2+1/2/b^2/x/(a
+b*arcsech(c*x))+1/2*c*Chi(a/b+arcsech(c*x))*sinh(a/b)/b^3-1/2*c*cosh(a/b)
*Shi(a/b+arcsech(c*x))/b^3
```

**Mathematica [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.90

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))^3} dx$$

$$= \frac{\frac{b^2 \sqrt{\frac{1-cx}{1+cx}}(1+cx)}{x(a+b \operatorname{sech}^{-1}(cx))^2} + \frac{b}{ax+b \operatorname{sech}^{-1}(cx)} + c(\operatorname{Chi}(\frac{a}{b} + \operatorname{sech}^{-1}(cx)) \sinh(\frac{a}{b}) - \cosh(\frac{a}{b}) \operatorname{Shi}(\frac{a}{b} + \operatorname{sech}^{-1}(cx)))}{2b^3}$$

input `Integrate[1/(x^2*(a + b*ArcSech[c*x])^3),x]`output `((b^2*sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(x*(a + b*ArcSech[c*x])^2) + b/(a*x + b*x*ArcSech[c*x]) + c*(CoshIntegral[a/b + ArcSech[c*x]]*Sinh[a/b] - Cosh[a/b]*SinhIntegral[a/b + ArcSech[c*x]]))/(2*b^3)`**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.69 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.24, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.071$ , Rules used = {6839, 3042, 26, 3778, 3042, 3778, 26, 3042, 26, 3784, 26, 3042, 26, 3779, 3782}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))^3} dx$$

$$\downarrow \text{6839}$$

$$-c \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{cx(a+b \operatorname{sech}^{-1}(cx))^3} d \operatorname{sech}^{-1}(cx)$$

$$\downarrow \text{3042}$$

$$-c \int -\frac{i \sin(i \operatorname{sech}^{-1}(cx))}{(a+b \operatorname{sech}^{-1}(cx))^3} d \operatorname{sech}^{-1}(cx)$$

$$\begin{aligned}
 & \downarrow 26 \\
 & ic \int \frac{\sin(i \operatorname{sech}^{-1}(cx))}{(a + b \operatorname{sech}^{-1}(cx))^3} d \operatorname{sech}^{-1}(cx) \\
 & \downarrow 3778 \\
 & ic \left( \frac{i \int \frac{1}{cx(a + b \operatorname{sech}^{-1}(cx))^2} d \operatorname{sech}^{-1}(cx)}{2b} - \frac{i \sqrt{\frac{1-cx}{cx+1}}(cx+1)}{2bcx(a + b \operatorname{sech}^{-1}(cx))^2} \right) \\
 & \downarrow 3042 \\
 & ic \left( \frac{i \int \frac{\sin(i \operatorname{sech}^{-1}(cx) + \frac{\pi}{2})}{(a + b \operatorname{sech}^{-1}(cx))^2} d \operatorname{sech}^{-1}(cx)}{2b} - \frac{i \sqrt{\frac{1-cx}{cx+1}}(cx+1)}{2bcx(a + b \operatorname{sech}^{-1}(cx))^2} \right) \\
 & \downarrow 3778 \\
 & ic \left( \frac{i \left( -\frac{1}{bcx(a + b \operatorname{sech}^{-1}(cx))} + \frac{i \int -\frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{cx(a + b \operatorname{sech}^{-1}(cx))} d \operatorname{sech}^{-1}(cx)}{b} \right)}{2b} - \frac{i \sqrt{\frac{1-cx}{cx+1}}(cx+1)}{2bcx(a + b \operatorname{sech}^{-1}(cx))^2} \right) \\
 & \downarrow 26 \\
 & ic \left( \frac{i \left( \frac{\int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{cx(a + b \operatorname{sech}^{-1}(cx))} d \operatorname{sech}^{-1}(cx)}{b} - \frac{1}{bcx(a + b \operatorname{sech}^{-1}(cx))} \right)}{2b} - \frac{i \sqrt{\frac{1-cx}{cx+1}}(cx+1)}{2bcx(a + b \operatorname{sech}^{-1}(cx))^2} \right) \\
 & \downarrow 3042
 \end{aligned}$$



$$ic \left( \frac{i \left( -\frac{1}{bcx(a+b\operatorname{sech}^{-1}(cx))} + \frac{\int -\frac{i \sin(i\operatorname{sech}^{-1}(cx))}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx)}{b} \right)}{2b} - \frac{i\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{2bcx(a+b\operatorname{sech}^{-1}(cx))^2} \right)$$

↓ 26

$$ic \left( \frac{i \left( -\frac{1}{bcx(a+b\operatorname{sech}^{-1}(cx))} - \frac{i \int \frac{\sin(i\operatorname{sech}^{-1}(cx))}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx)}{b} \right)}{2b} - \frac{i\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{2bcx(a+b\operatorname{sech}^{-1}(cx))^2} \right)$$

↓ 3784

$$ic \left( \frac{i \left( -\frac{1}{bcx(a+b\operatorname{sech}^{-1}(cx))} - \frac{i \left( \cosh\left(\frac{a}{b}\right) \int \frac{i \sinh\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) - i \sinh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)}{b} \right)}{2b}$$

↓ 26

$$ic \left( \frac{i \left( -\frac{1}{bcx(a+b\operatorname{sech}^{-1}(cx))} - \frac{i \left( i \cosh\left(\frac{a}{b}\right) \int \frac{\sinh\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) - i \sinh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)}{b} \right)}{2b}$$

↓ 3042

$$ic \left( \frac{i \left( -\frac{1}{bcx(a+b\operatorname{sech}^{-1}(cx))} - \frac{i \left( i \cosh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{ia}{b} + i\operatorname{sech}^{-1}(cx)\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) - i \sinh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{ia}{b} + i\operatorname{sech}^{-1}(cx) + \frac{\pi}{2}\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)}{b} \right)}{2b} \right)$$

↓ 26

$$ic \left( \frac{i \left( -\frac{1}{bcx(a+b\operatorname{sech}^{-1}(cx))} - \frac{i \left( \cosh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{ia}{b} + i\operatorname{sech}^{-1}(cx)\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) - i \sinh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{ia}{b} + i\operatorname{sech}^{-1}(cx) + \frac{\pi}{2}\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)}{b} \right)}{2b} \right)$$

↓ 3779

$$ic \left( \frac{i \left( -\frac{1}{bcx(a+b\operatorname{sech}^{-1}(cx))} - \frac{i \left( \frac{i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{b} - i \sinh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{ia}{b} + i\operatorname{sech}^{-1}(cx) + \frac{\pi}{2}\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)}{b} \right)}{2b} \right) - \frac{i \sqrt{\dots}}{2bcx \left( \dots \right)}$$

↓ 3782

$$ic \left( \frac{i \left( \frac{1}{bcx(a+b\operatorname{sech}^{-1}(cx))} - \frac{i \left( \frac{\cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{b} - \frac{\sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{b} \right)}{b} \right)}{2b} - \frac{i \sqrt{\frac{1-cx}{cx+1}} (cx+1)}{2bcx(a+b\operatorname{sech}^{-1}(cx))} \right)$$

input `Int[1/(x^2*(a + b*ArcSech[c*x])^3),x]`

output `I*c*(((1/2*I)*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(b*c*x*(a + b*ArcSech[c*x])^2) + ((I/2)*(-1/(b*c*x*(a + b*ArcSech[c*x]))) - (I*(((1/2)*CoshIntegral[a/b + ArcSech[c*x]]*Sinh[a/b])/b + (I*Cosh[a/b]*SinhIntegral[a/b + ArcSech[c*x]])/b))/b))`

### Definitions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3778 `Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + (f_)*(x_)], x_Symbol] := Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Simp[f/(d*(m + 1)) Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]`

rule 3779 `Int[sin[(e_) + (Complex[0, fz_]*(f_)*(x_))]/((c_) + (d_)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`

rule 3782 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`

rule 3784 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`

rule 6839 `Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^n*(x_)^m, x_Symbol] := Simp[-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])`

## Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 243 vs.  $2(104) = 208$ .

Time = 0.43 (sec) , antiderivative size = 244, normalized size of antiderivative = 2.14

method	result
derivativedivides	$c \left( -\frac{\left(\sqrt{-\frac{cx-1}{cx}} c\sqrt{\frac{cx+1}{cx}} x-1\right) (b \operatorname{arcsech}(cx)+a-b)}{4cx b^2 (b^2 \operatorname{arcsech}(cx)^2+2ab \operatorname{arcsech}(cx)+a^2)} - \frac{e^{\frac{a}{b}} \operatorname{expIntegral}_1\left(\frac{a}{b}+\operatorname{arcsech}(cx)\right)}{4b^3} + \frac{\sqrt{-\frac{cx-1}{cx}} c\sqrt{\frac{cx+1}{cx}}}{4bcx(a+b \operatorname{arcsech}(cx))} \right)$
default	$c \left( -\frac{\left(\sqrt{-\frac{cx-1}{cx}} c\sqrt{\frac{cx+1}{cx}} x-1\right) (b \operatorname{arcsech}(cx)+a-b)}{4cx b^2 (b^2 \operatorname{arcsech}(cx)^2+2ab \operatorname{arcsech}(cx)+a^2)} - \frac{e^{\frac{a}{b}} \operatorname{expIntegral}_1\left(\frac{a}{b}+\operatorname{arcsech}(cx)\right)}{4b^3} + \frac{\sqrt{-\frac{cx-1}{cx}} c\sqrt{\frac{cx+1}{cx}}}{4bcx(a+b \operatorname{arcsech}(cx))} \right)$

input `int(1/x^2/(a+b*arcsech(c*x))^3,x,method=_RETURNVERBOSE)`

output `c*(-1/4*((-(c*x-1)/c/x)^(1/2)*c*((c*x+1)/c/x)^(1/2)*x-1)*(b*arcsech(c*x)+a-b)/c/x/b^2/(b^2*arcsech(c*x)^2+2*a*b*arcsech(c*x)+a^2)-1/4/b^3*exp(1/b*a)*Ei(1,1/b*a+arcsech(c*x))+1/4/b*((-(c*x-1)/c/x)^(1/2)*c*((c*x+1)/c/x)^(1/2)*x+1)/c/x/(a+b*arcsech(c*x))^2+1/4/b^2*((-(c*x-1)/c/x)^(1/2)*c*((c*x+1)/c/x)^(1/2)*x+1)/c/x/(a+b*arcsech(c*x))+1/4/b^3*exp(-1/b*a)*Ei(1,-arcsech(c*x)-1/b*a))`

**Fricas [F]**

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3 x^2} dx$$

input `integrate(1/x^2/(a+b*arcsech(c*x))^3,x, algorithm="fricas")`

output `integral(1/(b^3*x^2*arcsech(c*x)^3 + 3*a*b^2*x^2*arcsech(c*x)^2 + 3*a^2*b*x^2*arcsech(c*x) + a^3*x^2), x)`

**Sympy [F]**

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{x^2 (a + b \operatorname{asech}(cx))^3} dx$$

input `integrate(1/x**2/(a+b*asech(c*x))**3,x)`

output `Integral(1/(x**2*(a + b*asech(c*x))**3), x)`

**Maxima [F]**

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3 x^2} dx$$

input `integrate(1/x^2/(a+b*arcsech(c*x))^3,x, algorithm="maxima")`

output

```

-1/2*((b*c^6*(log(c) - 1) - a*c^6)*x^7 - 3*(b*c^4*(log(c) - 1) - a*c^4)*x^
5 - (b*c^2*x^3 - (b*c^4*log(c) - a*c^4)*x^5 + (b*(log(c) - 1) - a)*x - (b*
c^4*x^5 - b*x)*log(x))*(c*x + 1)^(3/2)*(-c*x + 1)^(3/2) + 3*(b*c^2*(log(c)
- 1) - a*c^2)*x^3 - (2*b*c^4*x^5 + (b*c^2*(3*log(c) - 5) - 3*a*c^2)*x^3 -
3*(b*(log(c) - 1) - a)*x + 3*(b*c^2*x^3 - b*x)*log(x))*(c*x + 1)*(c*x - 1
) + ((b*c^6*(log(c) - 1) - a*c^6)*x^7 - (b*c^4*(4*log(c) - 5) - 4*a*c^4)*x
^5 + (b*c^2*(6*log(c) - 7) - 6*a*c^2)*x^3 - 3*(b*(log(c) - 1) - a)*x + (b*
c^6*x^7 - 4*b*c^4*x^5 + 6*b*c^2*x^3 - 3*b*x)*log(x))*sqrt(c*x + 1)*sqrt(-c
*x + 1) - (b*(log(c) - 1) - a)*x - (b*c^6*x^7 - 3*b*c^4*x^5 + 3*b*c^2*x^3
+ (b*c^4*x^5 - b*x)*(c*x + 1)^(3/2)*(-c*x + 1)^(3/2) - 3*(b*c^2*x^3 - b*x)
*(c*x + 1)*(c*x - 1) + (b*c^6*x^7 - 4*b*c^4*x^5 + 6*b*c^2*x^3 - 3*b*x)*sqr
t(c*x + 1)*sqrt(-c*x + 1) - b*x)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1) + (
b*c^6*x^7 - 3*b*c^4*x^5 + 3*b*c^2*x^3 - b*x)*log(x))/((b^4*c^6*x^6 - 3*b^4
*c^4*x^4 + 3*b^4*c^2*x^2 - b^4)*x^2*log(x)^2 - (b^4*x^2*log(x)^2 + 2*(b^4*
log(c) - a*b^3)*x^2*log(x) + (b^4*log(c)^2 - 2*a*b^3*log(c) + a^2*b^2)*x^2
)*(c*x + 1)^(3/2)*(-c*x + 1)^(3/2) + 2*((b^4*c^6*log(c) - a*b^3*c^6)*x^6 -
3*(b^4*c^4*log(c) - a*b^3*c^4)*x^4 - b^4*log(c) + a*b^3 + 3*(b^4*c^2*log(
c) - a*b^3*c^2)*x^2)*x^2*log(x) - 3*((b^4*c^2*x^2 - b^4)*x^2*log(x)^2 - 2*
(b^4*log(c) - a*b^3 - (b^4*c^2*log(c) - a*b^3*c^2)*x^2)*x^2*log(x) - (b^4*
log(c)^2 - 2*a*b^3*log(c) + a^2*b^2 - (b^4*c^2*log(c)^2 - 2*a*b^3*c^2*1...

```

**Giac [F]**

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(b \operatorname{arsech}(cx) + a)^3 x^2} dx$$

input

```
integrate(1/x^2/(a+b*arcsech(c*x))^3,x, algorithm="giac")
```

output

```
integrate(1/((b*arcsech(c*x) + a)^3*x^2), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{x^2 (a + b \operatorname{acosh}(\frac{1}{cx}))^3} dx$$

input `int(1/(x^2*(a + b*acosh(1/(c*x)))^3),x)`output `int(1/(x^2*(a + b*acosh(1/(c*x)))^3), x)`**Reduce [F]**

$$\begin{aligned} & \int \frac{1}{x^2 (a + b \operatorname{sech}^{-1}(cx))^3} dx \\ &= \int \frac{1}{\operatorname{asech}(cx)^3 b^3 x^2 + 3 \operatorname{asech}(cx)^2 a b^2 x^2 + 3 \operatorname{asech}(cx) a^2 b x^2 + a^3 x^2} dx \end{aligned}$$

input `int(1/x^2/(a+b*asech(c*x))^3,x)`output `int(1/(asech(c*x)**3*b**3*x**2 + 3*asech(c*x)**2*a*b**2*x**2 + 3*asech(c*x)*a**2*b*x**2 + a**3*x**2),x)`

**3.67**  $\int \frac{1}{x^3 (a+b\operatorname{sech}^{-1}(cx))^3} dx$

Optimal result	551
Mathematica [A] (verified)	552
Rubi [C] (verified)	552
Maple [B] (verified)	558
Fricas [F]	558
Sympy [F]	559
Maxima [F]	559
Giac [F]	560
Mupad [F(-1)]	561
Reduce [F]	561

**Optimal result**

Integrand size = 14, antiderivative size = 112

$$\int \frac{1}{x^3 (a + b\operatorname{sech}^{-1}(cx))^3} dx = \frac{c^2 \cosh(2\operatorname{sech}^{-1}(cx))}{2b^2 (a + b\operatorname{sech}^{-1}(cx))} + \frac{c^2 \operatorname{Chi}(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)) \sinh(\frac{2a}{b})}{b^3} + \frac{c^2 \sinh(2\operatorname{sech}^{-1}(cx))}{4b (a + b\operatorname{sech}^{-1}(cx))^2} - \frac{c^2 \cosh(\frac{2a}{b}) \operatorname{Shi}(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx))}{b^3}$$

output

```
1/2*c^2*cosh(2*arcsech(c*x))/b^2/(a+b*arcsech(c*x))+c^2*Chi(2*a/b+2*arcsec
h(c*x))*sinh(2*a/b)/b^3+1/4*c^2*sinh(2*arcsech(c*x))/b/(a+b*arcsech(c*x))^
2-c^2*cosh(2*a/b)*Shi(2*a/b+2*arcsech(c*x))/b^3
```



**Mathematica [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.09

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))^3} dx$$

$$= \frac{\frac{b^2 \sqrt{\frac{1-cx}{1+cx}} (1+cx)}{x^2 (a+b \operatorname{sech}^{-1}(cx))^2} + \frac{b(2-c^2 x^2)}{x^2 (a+b \operatorname{sech}^{-1}(cx))} + 2c^2 (\operatorname{Chi}(2(\frac{a}{b} + \operatorname{sech}^{-1}(cx))) \sinh(\frac{2a}{b}) - \cosh(\frac{2a}{b}) \operatorname{Shi}(2(\frac{a}{b} + \operatorname{sech}^{-1}(cx))))}{2b^3}}$$

input

```
Integrate[1/(x^3*(a + b*ArcSech[c*x])^3),x]
```

output

```
((b^2*sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(x^2*(a + b*ArcSech[c*x])^2) +
(b*(2 - c^2*x^2))/(x^2*(a + b*ArcSech[c*x])) + 2*c^2*(CoshIntegral[2*(a/b
+ ArcSech[c*x]])*Sinh[(2*a)/b] - Cosh[(2*a)/b]*SinhIntegral[2*(a/b + ArcSe
ch[c*x])]))/(2*b^3)
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.78 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.16, number of steps used = 18, number of rules used = 17,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.214$ , Rules used = {6839, 5971, 27, 3042, 26, 3778, 3042, 3778, 26, 3042, 26, 3784, 26, 3042, 26, 3779, 3782}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))^3} dx$$

$$\downarrow \text{6839}$$

$$-c^2 \int \frac{\sqrt{\frac{1-cx}{cx+1}} (cx+1)}{c^2 x^2 (a + b \operatorname{sech}^{-1}(cx))^3} d \operatorname{sech}^{-1}(cx)$$

$$\downarrow \text{5971}$$

$$\begin{aligned}
& -c^2 \int \frac{\sinh(2\operatorname{sech}^{-1}(cx))}{2(a+b\operatorname{sech}^{-1}(cx))^3} d\operatorname{sech}^{-1}(cx) \\
& \quad \downarrow 27 \\
& -\frac{1}{2}c^2 \int \frac{\sinh(2\operatorname{sech}^{-1}(cx))}{(a+b\operatorname{sech}^{-1}(cx))^3} d\operatorname{sech}^{-1}(cx) \\
& \quad \downarrow 3042 \\
& -\frac{1}{2}c^2 \int -\frac{i \sin(2i\operatorname{sech}^{-1}(cx))}{(a+b\operatorname{sech}^{-1}(cx))^3} d\operatorname{sech}^{-1}(cx) \\
& \quad \downarrow 26 \\
& \frac{1}{2}ic^2 \int \frac{\sin(2i\operatorname{sech}^{-1}(cx))}{(a+b\operatorname{sech}^{-1}(cx))^3} d\operatorname{sech}^{-1}(cx) \\
& \quad \downarrow 3778 \\
& \frac{1}{2}ic^2 \left( \frac{i \int \frac{\cosh(2\operatorname{sech}^{-1}(cx))}{(a+b\operatorname{sech}^{-1}(cx))^2} d\operatorname{sech}^{-1}(cx)}{b} - \frac{i \sinh(2\operatorname{sech}^{-1}(cx))}{2b(a+b\operatorname{sech}^{-1}(cx))^2} \right) \\
& \quad \downarrow 3042 \\
& \frac{1}{2}ic^2 \left( \frac{i \int \frac{\sin(2i\operatorname{sech}^{-1}(cx) + \frac{\pi}{2})}{(a+b\operatorname{sech}^{-1}(cx))^2} d\operatorname{sech}^{-1}(cx)}{b} - \frac{i \sinh(2\operatorname{sech}^{-1}(cx))}{2b(a+b\operatorname{sech}^{-1}(cx))^2} \right) \\
& \quad \downarrow 3778 \\
& \frac{1}{2}ic^2 \left( \frac{i \left( -\frac{\cosh(2\operatorname{sech}^{-1}(cx))}{b(a+b\operatorname{sech}^{-1}(cx))} + \frac{2i \int -\frac{i \sinh(2\operatorname{sech}^{-1}(cx))}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx)}{b} \right)}{b} - \frac{i \sinh(2\operatorname{sech}^{-1}(cx))}{2b(a+b\operatorname{sech}^{-1}(cx))^2} \right) \\
& \quad \downarrow 26
\end{aligned}$$

$$\begin{aligned}
 & \frac{1}{2} i c^2 \left( \frac{i \left( \frac{2 \int \frac{\sinh(2 \operatorname{sech}^{-1}(cx))}{a+b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx)}{b} - \frac{\cosh(2 \operatorname{sech}^{-1}(cx))}{b(a+b \operatorname{sech}^{-1}(cx))} \right)}{b} - \frac{i \sinh(2 \operatorname{sech}^{-1}(cx))}{2b(a+b \operatorname{sech}^{-1}(cx))^2} \right) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} i c^2 \left( \frac{i \left( -\frac{\cosh(2 \operatorname{sech}^{-1}(cx))}{b(a+b \operatorname{sech}^{-1}(cx))} + \frac{2 \int -\frac{i \sin(2i \operatorname{sech}^{-1}(cx))}{a+b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx)}{b} \right)}{b} - \frac{i \sinh(2 \operatorname{sech}^{-1}(cx))}{2b(a+b \operatorname{sech}^{-1}(cx))^2} \right) \\
 & \quad \downarrow \text{26} \\
 & \frac{1}{2} i c^2 \left( \frac{i \left( -\frac{\cosh(2 \operatorname{sech}^{-1}(cx))}{b(a+b \operatorname{sech}^{-1}(cx))} - \frac{2i \int \frac{\sin(2i \operatorname{sech}^{-1}(cx))}{a+b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx)}{b} \right)}{b} - \frac{i \sinh(2 \operatorname{sech}^{-1}(cx))}{2b(a+b \operatorname{sech}^{-1}(cx))^2} \right) \\
 & \quad \downarrow \text{3784} \\
 & \frac{1}{2} i c^2 \left( \frac{i \left( -\frac{\cosh(2 \operatorname{sech}^{-1}(cx))}{b(a+b \operatorname{sech}^{-1}(cx))} - \frac{2i \left( \cosh\left(\frac{2a}{b}\right) \int \frac{i \sinh\left(\frac{2a}{b} + 2 \operatorname{sech}^{-1}(cx)\right)}{a+b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) - i \sinh\left(\frac{2a}{b}\right) \int \frac{\cosh\left(\frac{2a}{b} + 2 \operatorname{sech}^{-1}(cx)\right)}{a+b \operatorname{sech}^{-1}(cx)} d \operatorname{sech}^{-1}(cx) \right)}{b} \right)}{b} \right) \\
 & \quad \downarrow \text{26}
 \end{aligned}$$

$$\frac{1}{2}ic^2 \left( i \left( -\frac{\cosh(2\operatorname{sech}^{-1}(cx))}{b(a+b\operatorname{sech}^{-1}(cx))} - \frac{2i \left( i \cosh\left(\frac{2a}{b}\right) \int \frac{\sinh\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) - i \sinh\left(\frac{2a}{b}\right) \int \frac{\cosh\left(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx)\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)}{b} \right) \right)$$

↓ 3042

$$\frac{1}{2}ic^2 \left( i \left( -\frac{\cosh(2\operatorname{sech}^{-1}(cx))}{b(a+b\operatorname{sech}^{-1}(cx))} - \frac{2i \left( i \cosh\left(\frac{2a}{b}\right) \int -\frac{i \sin\left(\frac{2ia}{b} + 2i\operatorname{sech}^{-1}(cx)\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) - i \sinh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2ia}{b} + 2i\operatorname{sech}^{-1}(cx) + \frac{\pi}{2}\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)}{b} \right) \right)$$

↓ 26

$$\frac{1}{2}ic^2 \left( i \left( -\frac{\cosh(2\operatorname{sech}^{-1}(cx))}{b(a+b\operatorname{sech}^{-1}(cx))} - \frac{2i \left( \cosh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2ia}{b} + 2i\operatorname{sech}^{-1}(cx)\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) - i \sinh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2ia}{b} + 2i\operatorname{sech}^{-1}(cx) + \frac{\pi}{2}\right)}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)}{b} \right) \right)$$

↓ 3779

$$\frac{1}{2}ic^2 \left( \frac{i \left( \frac{\cosh(2\operatorname{sech}^{-1}(cx))}{b(a+b\operatorname{sech}^{-1}(cx))} - \frac{2i \left( \frac{i \cosh(\frac{2a}{b}) \operatorname{Shi}(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx))}{b} - i \sinh(\frac{2a}{b}) \int \frac{\sin(\frac{2ia}{b} + 2i\operatorname{sech}^{-1}(cx) + \frac{\pi}{2})}{a+b\operatorname{sech}^{-1}(cx)} d\operatorname{sech}^{-1}(cx) \right)}{b} \right)}{b} \right)$$

↓ 3782

$$\frac{1}{2}ic^2 \left( \frac{i \left( \frac{\cosh(2\operatorname{sech}^{-1}(cx))}{b(a+b\operatorname{sech}^{-1}(cx))} - \frac{2i \left( \frac{i \cosh(\frac{2a}{b}) \operatorname{Shi}(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx))}{b} - \frac{i \sinh(\frac{2a}{b}) \operatorname{Chi}(\frac{2a}{b} + 2\operatorname{sech}^{-1}(cx))}{b} \right)}{b} \right)}{b} \right) - \frac{i \sinh(2\operatorname{sech}^{-1}(cx))}{2b(a+b\operatorname{sech}^{-1}(cx))}$$

input `Int[1/(x^3*(a + b*ArcSech[c*x])^3),x]`

output `(I/2)*c^2*((( -1/2*I)*Sinh[2*ArcSech[c*x]]/(b*(a + b*ArcSech[c*x])^2) + (I*(-(Cosh[2*ArcSech[c*x]]/(b*(a + b*ArcSech[c*x]))) - ((2*I)*((( -I)*CoshIntegral[(2*a)/b + 2*ArcSech[c*x]]*Sinh[(2*a)/b])/b + (I*Cosh[(2*a)/b]*SinhIntegral[(2*a)/b + 2*ArcSech[c*x]]/b))/b)))/b)`

**Defintions of rubi rules used**

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3778 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Simp[f/(d*(m + 1)) Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]`

rule 3779 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`

rule 3782 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`

rule 3784 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^(n)*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6839 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))^(n_)*(x_)^(m_), x_Symbol] := Simp[-(c^(m + 1))^( -1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, ArcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (GtQ[n, 0] || LtQ[m, -1])`

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 276 vs.  $2(108) = 216$ .

Time = 0.61 (sec) , antiderivative size = 277, normalized size of antiderivative = 2.47

method	result
derivativedivides	$c^2 \left( -\frac{\left(2\sqrt{-\frac{cx-1}{cx}} c\sqrt{\frac{cx+1}{cx}} x + c^2 x^2 - 2\right) (2b \operatorname{arcsech}(cx) + 2a - b)}{8c^2 x^2 b^2 (b^2 \operatorname{arcsech}(cx)^2 + 2ab \operatorname{arcsech}(cx) + a^2)} - \frac{e^{\frac{2a}{b}} \operatorname{expIntegral}_1\left(\frac{2a}{b} + 2 \operatorname{arcsech}(cx)\right)}{2b^3} - \frac{c^2 x^2}{8b} \right)$
default	$c^2 \left( -\frac{\left(2\sqrt{-\frac{cx-1}{cx}} c\sqrt{\frac{cx+1}{cx}} x + c^2 x^2 - 2\right) (2b \operatorname{arcsech}(cx) + 2a - b)}{8c^2 x^2 b^2 (b^2 \operatorname{arcsech}(cx)^2 + 2ab \operatorname{arcsech}(cx) + a^2)} - \frac{e^{\frac{2a}{b}} \operatorname{expIntegral}_1\left(\frac{2a}{b} + 2 \operatorname{arcsech}(cx)\right)}{2b^3} - \frac{c^2 x^2}{8b} \right)$

input `int(1/x^3/(a+b*arcsech(c*x))^3,x,method=_RETURNVERBOSE)`

output 
$$c^2 * (-1/8 * (2 * (-c*x-1)/c/x)^{(1/2)} * c * ((c*x+1)/c/x)^{(1/2)} * x + c^2 * x^2 - 2) * (2 * b * \operatorname{arcsech}(c*x) + 2 * a - b) / c^2 / x^2 / b^2 / (b^2 * \operatorname{arcsech}(c*x)^2 + 2 * a * b * \operatorname{arcsech}(c*x) + a^2) - 1/2 / b^3 * \exp(2/b*a) * \operatorname{Ei}(1, 2/b*a + 2 * \operatorname{arcsech}(c*x)) - 1/8 / b * (c^2 * x^2 - 2 - 2 * (-c*x-1)/c/x)^{(1/2)} * c * ((c*x+1)/c/x)^{(1/2)} * x / c^2 / x^2 / (a + b * \operatorname{arcsech}(c*x))^2 - 1/4 / b^2 * (c^2 * x^2 - 2 - 2 * (-c*x-1)/c/x)^{(1/2)} * c * ((c*x+1)/c/x)^{(1/2)} * x / c^2 / x^2 / (a + b * \operatorname{arcsech}(c*x)) + 1/2 / b^3 * \exp(-2/b*a) * \operatorname{Ei}(1, -2 * \operatorname{arcsech}(c*x) - 2/b*a)$$

**Fricas [F]**

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3 x^3} dx$$

input `integrate(1/x^3/(a+b*arcsech(c*x))^3,x, algorithm="fricas")`

output `integral(1/(b^3*x^3*arcsech(c*x)^3 + 3*a*b^2*x^3*arcsech(c*x)^2 + 3*a^2*b*x^3*arcsech(c*x) + a^3*x^3), x)`

**Sympy [F]**

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{x^3 (a + b \operatorname{asech}(cx))^3} dx$$

input `integrate(1/x**3/(a+b*asech(c*x))**3,x)`

output `Integral(1/(x**3*(a + b*asech(c*x))**3), x)`

**Maxima [F]**

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(b \operatorname{arsech}(cx) + a)^3 x^3} dx$$

input `integrate(1/x^3/(a+b*arcsech(c*x))^3,x, algorithm="maxima")`



output

```

-1/2*((b*c^6*(2*log(c) - 1) - 2*a*c^6)*x^7 - 3*(b*c^4*(2*log(c) - 1) - 2*a
*c^4)*x^5 + ((b*c^2*(2*log(c) - 1) - 2*a*c^2)*x^3 - (b*(2*log(c) - 1) - 2*
a)*x + 2*(b*c^2*x^3 - b*x)*log(x))*(c*x + 1)^(3/2)*(-c*x + 1)^(3/2) + 3*(b
*c^2*(2*log(c) - 1) - 2*a*c^2)*x^3 - ((b*c^6*log(c) - a*c^6)*x^7 - (b*c^4*
(5*log(c) - 2) - 5*a*c^4)*x^5 + 5*(b*c^2*(2*log(c) - 1) - 2*a*c^2)*x^3 - 3
*(b*(2*log(c) - 1) - 2*a)*x + (b*c^6*x^7 - 5*b*c^4*x^5 + 10*b*c^2*x^3 - 6*
b*x)*log(x))*(c*x + 1)*(c*x - 1) + ((b*c^6*(3*log(c) - 1) - 3*a*c^6)*x^7 -
(b*c^4*(11*log(c) - 5) - 11*a*c^4)*x^5 + 7*(b*c^2*(2*log(c) - 1) - 2*a*c^
2)*x^3 - 3*(b*(2*log(c) - 1) - 2*a)*x + (3*b*c^6*x^7 - 11*b*c^4*x^5 + 14*b
*c^2*x^3 - 6*b*x)*log(x))*sqrt(c*x + 1)*sqrt(-c*x + 1) - (b*(2*log(c) - 1)
- 2*a)*x - (2*b*c^6*x^7 - 6*b*c^4*x^5 + 6*b*c^2*x^3 + 2*(b*c^2*x^3 - b*x)
*(c*x + 1)^(3/2)*(-c*x + 1)^(3/2) - (b*c^6*x^7 - 5*b*c^4*x^5 + 10*b*c^2*x^
3 - 6*b*x)*(c*x + 1)*(c*x - 1) + (3*b*c^6*x^7 - 11*b*c^4*x^5 + 14*b*c^2*x^
3 - 6*b*x)*sqrt(c*x + 1)*sqrt(-c*x + 1) - 2*b*x*log(sqrt(c*x + 1)*sqrt(-c
*x + 1) + 1) + 2*(b*c^6*x^7 - 3*b*c^4*x^5 + 3*b*c^2*x^3 - b*x)*log(x))/((b
^4*c^6*x^6 - 3*b^4*c^4*x^4 + 3*b^4*c^2*x^2 - b^4)*x^3*log(x)^2 + 2*((b^4*c
^6*log(c) - a*b^3*c^6)*x^6 - 3*(b^4*c^4*log(c) - a*b^3*c^4)*x^4 - b^4*log(
c) + a*b^3 + 3*(b^4*c^2*log(c) - a*b^3*c^2)*x^2)*x^3*log(x) - (b^4*x^3*log
(x)^2 + 2*(b^4*log(c) - a*b^3)*x^3*log(x) + (b^4*log(c)^2 - 2*a*b^3*log(c)
+ a^2*b^2)*x^3)*(c*x + 1)^(3/2)*(-c*x + 1)^(3/2) + ((b^4*c^6*log(c)^2 ...

```

**Giac** [F]

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3 x^3} dx$$

input

```
integrate(1/x^3/(a+b*arcsech(c*x))^3,x, algorithm="giac")
```

output

```
integrate(1/((b*arcsech(c*x) + a)^3*x^3), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{x^3 (a + b \operatorname{acosh}(\frac{1}{cx}))^3} dx$$

input `int(1/(x^3*(a + b*acosh(1/(c*x)))^3),x)`output `int(1/(x^3*(a + b*acosh(1/(c*x)))^3), x)`**Reduce [F]**

$$\begin{aligned} & \int \frac{1}{x^3 (a + b \operatorname{sech}^{-1}(cx))^3} dx \\ &= \int \frac{1}{\operatorname{asech}(cx)^3 b^3 x^3 + 3 \operatorname{asech}(cx)^2 a b^2 x^3 + 3 \operatorname{asech}(cx) a^2 b x^3 + a^3 x^3} dx \end{aligned}$$

input `int(1/x^3/(a+b*asech(c*x))^3,x)`output `int(1/(asech(c*x)**3*b**3*x**3 + 3*asech(c*x)**2*a*b**2*x**3 + 3*asech(c*x)*a**2*b*x**3 + a**3*x**3),x)`

**3.68** 
$$\int \frac{1}{x^4 (a+b\operatorname{sech}^{-1}(cx))^3} dx$$

Optimal result	562
Mathematica [A] (verified)	563
Rubi [A] (verified)	563
Maple [B] (verified)	565
Fricas [F]	566
Sympy [F]	566
Maxima [F]	566
Giac [F]	567
Mupad [F(-1)]	568
Reduce [F]	568

**Optimal result**

Integrand size = 14, antiderivative size = 240

$$\int \frac{1}{x^4 (a+b\operatorname{sech}^{-1}(cx))^3} dx = \frac{c^2 \sqrt{\frac{1-cx}{1+cx}} (1+cx)}{8bx (a+b\operatorname{sech}^{-1}(cx))^2} + \frac{c^2}{8b^2x (a+b\operatorname{sech}^{-1}(cx))} + \frac{3c^3 \cosh(3\operatorname{sech}^{-1}(cx))}{8b^2 (a+b\operatorname{sech}^{-1}(cx))} + \frac{c^3 \operatorname{Chi}(\frac{a}{b} + \operatorname{sech}^{-1}(cx)) \sinh(\frac{a}{b})}{8b^3} + \frac{9c^3 \operatorname{Chi}(\frac{3a}{b} + 3\operatorname{sech}^{-1}(cx)) \sinh(\frac{3a}{b})}{8b^3} + \frac{c^3 \sinh(3\operatorname{sech}^{-1}(cx))}{8b (a+b\operatorname{sech}^{-1}(cx))^2} - \frac{c^3 \cosh(\frac{a}{b}) \operatorname{Shi}(\frac{a}{b} + \operatorname{sech}^{-1}(cx))}{8b^3} - \frac{9c^3 \cosh(\frac{3a}{b}) \operatorname{Shi}(\frac{3a}{b} + 3\operatorname{sech}^{-1}(cx))}{8b^3}$$

output

$$\begin{aligned} & 1/8*c^2*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)/b/x/(a+b*arcsech(c*x))^2+1/8*c^2/ \\ & b^2/x/(a+b*arcsech(c*x))+3/8*c^3*cosh(3*arcsech(c*x))/b^2/(a+b*arcsech(c*x) \\ & ))+1/8*c^3*Chi(a/b+arcsech(c*x))*sinh(a/b)/b^3+9/8*c^3*Chi(3*a/b+3*arcsech \\ & (c*x))*sinh(3*a/b)/b^3+1/8*c^3*sinh(3*arcsech(c*x))/b/(a+b*arcsech(c*x))^2 \\ & -1/8*c^3*cosh(a/b)*Shi(a/b+arcsech(c*x))/b^3-9/8*c^3*cosh(3*a/b)*Shi(3*a/b \\ & +3*arcsech(c*x))/b^3 \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 204, normalized size of antiderivative = 0.85

$$\begin{aligned} & \int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^3} dx \\ & \frac{4b^2 \sqrt{\frac{1-cx}{1+cx}}(1+cx)}{x^3 (a+b \operatorname{sech}^{-1}(cx))^2} + \frac{4b(3-2c^2x^2)}{x^3 (a+b \operatorname{sech}^{-1}(cx))} - 8c^3 \left( \operatorname{Chi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right) \sinh\left(\frac{a}{b}\right) - \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right) \right) \\ & = \end{aligned}$$

input

Integrate[1/(x^4\*(a + b\*ArcSech[c\*x])^3),x]

output

$$\begin{aligned} & ((4*b^2*sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(x^3*(a + b*ArcSech[c*x])^2) \\ & + (4*b*(3 - 2*c^2*x^2))/(x^3*(a + b*ArcSech[c*x])) - 8*c^3*(CoshIntegral[a \\ & /b + ArcSech[c*x]]*Sinh[a/b] - Cosh[a/b]*SinhIntegral[a/b + ArcSech[c*x]]) \\ & + 9*c^3*(CoshIntegral[a/b + ArcSech[c*x]]*Sinh[a/b] + CoshIntegral[3*(a/b \\ & + ArcSech[c*x]))*Sinh[(3*a)/b] - Cosh[a/b]*SinhIntegral[a/b + ArcSech[c*x] \\ & ]) - Cosh[(3*a)/b]*SinhIntegral[3*(a/b + ArcSech[c*x])]))/(8*b^3) \end{aligned}$$

**Rubi [A] (verified)**Time = 0.65 (sec) , antiderivative size = 227, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {6839, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^3} dx \\
& \quad \downarrow \text{6839} \\
& -c^3 \int \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{c^3 x^3 (a + b \operatorname{sech}^{-1}(cx))^3} d \operatorname{sech}^{-1}(cx) \\
& \quad \downarrow \text{5971} \\
& -c^3 \int \left( \frac{\sqrt{\frac{1-cx}{cx+1}}(cx+1)}{4cx (a + b \operatorname{sech}^{-1}(cx))^3} + \frac{\sinh(3 \operatorname{sech}^{-1}(cx))}{4 (a + b \operatorname{sech}^{-1}(cx))^3} \right) d \operatorname{sech}^{-1}(cx) \\
& \quad \downarrow \text{2009} \\
& -c^3 \left( -\frac{\sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{8b^3} - \frac{9 \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3a}{b} + 3 \operatorname{sech}^{-1}(cx)\right)}{8b^3} + \frac{\cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{sech}^{-1}(cx)\right)}{8b^3} + \dots \right)
\end{aligned}$$

input `Int[1/(x^4*(a + b*ArcSech[c*x])^3),x]`

output `-(c^3*(-1/8*(Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(b*c*x*(a + b*ArcSech[c*x])^2) - 1/(8*b^2*c*x*(a + b*ArcSech[c*x])) - (3*Cosh[3*ArcSech[c*x]])/(8*b^2*(a + b*ArcSech[c*x])) - (CoshIntegral[a/b + ArcSech[c*x]]*Sinh[a/b])/(8*b^3) - (9*CoshIntegral[(3*a)/b + 3*ArcSech[c*x]]*Sinh[(3*a)/b])/(8*b^3) - Sinh[3*ArcSech[c*x]]/(8*b*(a + b*ArcSech[c*x])^2) + (Cosh[a/b]*SinhIntegral[a/b + ArcSech[c*x]])/(8*b^3) + (9*Cosh[(3*a)/b]*SinhIntegral[(3*a)/b + 3*ArcSech[c*x]])/(8*b^3))`

### Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^(n)*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6839

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_.), x_Symbol] := Simp[
-(c^(m + 1))^(-1) Subst[Int[(a + b*x)^n*Sech[x]^(m + 1)*Tanh[x], x], x, A
rcSech[c*x]], x] /; FreeQ[{a, b, c}, x] && IntegerQ[n] && IntegerQ[m] && (G
tQ[n, 0] || LtQ[m, -1])
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 627 vs. 2(222) = 444.

Time = 0.87 (sec) , antiderivative size = 628, normalized size of antiderivative = 2.62

method	result
derivativedivides	$c^3 \left( \frac{\left( \sqrt{-\frac{cx-1}{cx}} c^3 x^3 \sqrt{\frac{cx+1}{cx}} - 4 \sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x - 3c^2 x^2 + 4 \right) (3b \operatorname{arcsech}(cx) + 3a - b)}{16c^3 x^3 b^2 (b^2 \operatorname{arcsech}(cx)^2 + 2ab \operatorname{arcsech}(cx) + a^2)} - \frac{9e^{\frac{3a}{b}} \operatorname{expIntegral}_1\left(\frac{3a}{b}\right)}{16b^3} \right)$
default	$c^3 \left( \frac{\left( \sqrt{-\frac{cx-1}{cx}} c^3 x^3 \sqrt{\frac{cx+1}{cx}} - 4 \sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x - 3c^2 x^2 + 4 \right) (3b \operatorname{arcsech}(cx) + 3a - b)}{16c^3 x^3 b^2 (b^2 \operatorname{arcsech}(cx)^2 + 2ab \operatorname{arcsech}(cx) + a^2)} - \frac{9e^{\frac{3a}{b}} \operatorname{expIntegral}_1\left(\frac{3a}{b}\right)}{16b^3} \right)$

input

```
int(1/x^4/(a+b*arcsech(c*x))^3,x,method=_RETURNVERBOSE)
```

output

```
c^3*(1/16*((-(c*x-1)/c/x)^(1/2)*c^3*x^3*((c*x+1)/c/x)^(1/2)-4*-(c*x-1)/c/
x)^(1/2)*c*((c*x+1)/c/x)^(1/2)*x-3*c^2*x^2+4)*(3*b*arcsech(c*x)+3*a-b)/c^3
/x^3/b^2/(b^2*arcsech(c*x)^2+2*a*b*arcsech(c*x)+a^2)-9/16/b^3*exp(3/b*a)*E
i(1,3/b*a+3*arcsech(c*x))-1/16*((-(c*x-1)/c/x)^(1/2)*c*((c*x+1)/c/x)^(1/2)
*x-1)*(b*arcsech(c*x)+a-b)/c/x/b^2/(b^2*arcsech(c*x)^2+2*a*b*arcsech(c*x)+
a^2)-1/16/b^3*exp(1/b*a)*Ei(1,1/b*a+arcsech(c*x))+1/16/b*((-(c*x-1)/c/x)^(
1/2)*c*((c*x+1)/c/x)^(1/2)*x+1)/c/x/(a+b*arcsech(c*x))^2+1/16/b^2*((-(c*x-
1)/c/x)^(1/2)*c*((c*x+1)/c/x)^(1/2)*x+1)/c/x/(a+b*arcsech(c*x))+1/16/b^3*exp(-1/b*a)*Ei(1,-arcsech(c*x)-1/b*a)-1/16/b*((-(c*x-1)/c/x)^(1/2)*c^3*x^3*
((c*x+1)/c/x)^(1/2)-4*-(c*x-1)/c/x)^(1/2)*c*((c*x+1)/c/x)^(1/2)*x+3*c^2*x
^2-4)/c^3/x^3/(a+b*arcsech(c*x))^2-3/16/b^2*((-(c*x-1)/c/x)^(1/2)*c^3*x^3*
((c*x+1)/c/x)^(1/2)-4*-(c*x-1)/c/x)^(1/2)*c*((c*x+1)/c/x)^(1/2)*x+3*c^2*x
^2-4)/c^3/x^3/(a+b*arcsech(c*x))+9/16/b^3*exp(-3/b*a)*Ei(1,-3*arcsech(c*x)
-3/b*a))
```

**Fricas [F]**

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3 x^4} dx$$

input `integrate(1/x^4/(a+b*arcsech(c*x))^3,x, algorithm="fricas")`

output `integral(1/(b^3*x^4*arcsech(c*x)^3 + 3*a*b^2*x^4*arcsech(c*x)^2 + 3*a^2*b*x^4*arcsech(c*x) + a^3*x^4), x)`

**Sympy [F]**

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{x^4 (a + b \operatorname{ar} \operatorname{sech}(cx))^3} dx$$

input `integrate(1/x**4/(a+b*asech(c*x))**3,x)`

output `Integral(1/(x**4*(a + b*asech(c*x))**3), x)`

**Maxima [F]**

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3 x^4} dx$$

input `integrate(1/x^4/(a+b*arcsech(c*x))^3,x, algorithm="maxima")`

output

```

-1/2*((b*c^6*(3*log(c) - 1) - 3*a*c^6)*x^7 - 3*(b*c^4*(3*log(c) - 1) - 3*a
*c^4)*x^5 - ((b*c^4*log(c) - a*c^4)*x^5 - (b*c^2*(4*log(c) - 1) - 4*a*c^2)
*x^3 + (b*(3*log(c) - 1) - 3*a)*x + (b*c^4*x^5 - 4*b*c^2*x^3 + 3*b*x)*log(
x))*(c*x + 1)^(3/2)*(-c*x + 1)^(3/2) + 3*(b*c^2*(3*log(c) - 1) - 3*a*c^2)*
x^3 - (2*(b*c^6*log(c) - a*c^6)*x^7 - 2*(b*c^4*(5*log(c) - 1) - 5*a*c^4)*x
^5 + (b*c^2*(17*log(c) - 5) - 17*a*c^2)*x^3 - 3*(b*(3*log(c) - 1) - 3*a)*x
+ (2*b*c^6*x^7 - 10*b*c^4*x^5 + 17*b*c^2*x^3 - 9*b*x)*log(x))*(c*x + 1)*(
c*x - 1) + ((b*c^6*(5*log(c) - 1) - 5*a*c^6)*x^7 - (b*c^4*(18*log(c) - 5)
- 18*a*c^4)*x^5 + (b*c^2*(22*log(c) - 7) - 22*a*c^2)*x^3 - 3*(b*(3*log(c)
- 1) - 3*a)*x + (5*b*c^6*x^7 - 18*b*c^4*x^5 + 22*b*c^2*x^3 - 9*b*x)*log(x)
)*sqrt(c*x + 1)*sqrt(-c*x + 1) - (b*(3*log(c) - 1) - 3*a)*x - (3*b*c^6*x^7
- 9*b*c^4*x^5 + 9*b*c^2*x^3 - (b*c^4*x^5 - 4*b*c^2*x^3 + 3*b*x)*(c*x + 1)
^(3/2)*(-c*x + 1)^(3/2) - (2*b*c^6*x^7 - 10*b*c^4*x^5 + 17*b*c^2*x^3 - 9*b
*x)*(c*x + 1)*(c*x - 1) + (5*b*c^6*x^7 - 18*b*c^4*x^5 + 22*b*c^2*x^3 - 9*b
*x)*sqrt(c*x + 1)*sqrt(-c*x + 1) - 3*b*x*log(sqrt(c*x + 1)*sqrt(-c*x + 1)
+ 1) + 3*(b*c^6*x^7 - 3*b*c^4*x^5 + 3*b*c^2*x^3 - b*x)*log(x))/((b^4*c^6*
x^6 - 3*b^4*c^4*x^4 + 3*b^4*c^2*x^2 - b^4)*x^4*log(x)^2 + 2*((b^4*c^6*log(
c) - a*b^3*c^6)*x^6 - 3*(b^4*c^4*log(c) - a*b^3*c^4)*x^4 - b^4*log(c) + a
b^3 + 3*(b^4*c^2*log(c) - a*b^3*c^2)*x^2)*x^4*log(x) + ((b^4*c^6*log(c))^2
- 2*a*b^3*c^6*log(c) + a^2*b^2*c^6)*x^6 - b^4*log(c)^2 - 3*(b^4*c^4*log...

```

**Giac** [F]

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^3 x^4} dx$$

input

```
integrate(1/x^4/(a+b*arcsech(c*x))^3,x, algorithm="giac")
```

output

```
integrate(1/((b*arcsech(c*x) + a)^3*x^4), x)
```



**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^3} dx = \int \frac{1}{x^4 (a + b \operatorname{acosh}(\frac{1}{cx}))^3} dx$$

input `int(1/(x^4*(a + b*acosh(1/(c*x)))^3),x)`output `int(1/(x^4*(a + b*acosh(1/(c*x)))^3), x)`**Reduce [F]**

$$\begin{aligned} & \int \frac{1}{x^4 (a + b \operatorname{sech}^{-1}(cx))^3} dx \\ &= \int \frac{1}{\operatorname{asech}(cx)^3 b^3 x^4 + 3 \operatorname{asech}(cx)^2 a b^2 x^4 + 3 \operatorname{asech}(cx) a^2 b x^4 + a^3 x^4} dx \end{aligned}$$

input `int(1/x^4/(a+b*asech(c*x))^3,x)`output `int(1/(asech(c*x)**3*b**3*x**4 + 3*asech(c*x)**2*a*b**2*x**4 + 3*asech(c*x)*a**2*b*x**4 + a**3*x**4),x)`

### 3.69 $\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^3 dx$

Optimal result	569
Mathematica [N/A]	569
Rubi [N/A]	570
Maple [N/A]	570
Fricas [N/A]	571
Sympy [N/A]	571
Maxima [N/A]	571
Giac [N/A]	572
Mupad [N/A]	573
Reduce [N/A]	573

#### Optimal result

Integrand size = 16, antiderivative size = 16

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^3 dx = \operatorname{Int}\left((dx)^m (a + b \operatorname{sech}^{-1}(cx))^3, x\right)$$

output `Defer(Int)((d*x)^m*(a+b*arcsech(c*x))^3,x)`

#### Mathematica [N/A]

Not integrable

Time = 3.26 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^3 dx$$

input `Integrate[(d*x)^m*(a + b*ArcSech[c*x])^3,x]`

output `Integrate[(d*x)^m*(a + b*ArcSech[c*x])^3, x]`

**Rubi [N/A]**

Not integrable

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^3 dx$$

↓ 6865

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^3 dx$$

input `Int[(d*x)^m*(a + b*ArcSech[c*x])^3,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.58 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int (dx)^m (a + b \operatorname{arcsech}(cx))^3 dx$$

input `int((d*x)^m*(a+b*arcsech(c*x))^3,x)`

output `int((d*x)^m*(a+b*arcsech(c*x))^3,x)`

**Fricas [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.75

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^3 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arcsech(c*x))^3,x, algorithm="fricas")`

output `integral((b^3*arcsech(c*x)^3 + 3*a*b^2*arcsech(c*x)^2 + 3*a^2*b*arcsech(c*x) + a^3)*(d*x)^m, x)`

**Sympy [N/A]**

Not integrable

Time = 6.62 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (dx)^m (a + b \operatorname{asech}(cx))^3 dx$$

input `integrate((d*x)**m*(a+b*asech(c*x))**3,x)`

output `Integral((d*x)**m*(a + b*asech(c*x))**3, x)`

**Maxima [N/A]**

Not integrable

Time = 10.37 (sec) , antiderivative size = 1450, normalized size of antiderivative = 90.62

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^3 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arcsech(c*x))^3,x, algorithm="maxima")`

output

```

b^3*d^m*x^m*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)^3/(m + 1) + (d*x)^(m +
1)*a^3/(d*(m + 1)) - integrate(((b^3*c^2*d^m*(m + 1)*x^2 - b^3*d^m*(m + 1
))*x^m*log(x)^3 - 3*(b^3*d^m*(m + 1)*log(c) - a*b^2*d^m*(m + 1) - (b^3*c^2
*d^m*(m + 1)*log(c) - a*b^2*c^2*d^m*(m + 1))*x^2)*x^m*log(x)^2 + 3*((b^3*c
^2*d^m*(m + 1)*x^2 - b^3*d^m*(m + 1))*x^m*log(x) + ((b^3*c^2*d^m*(m + 1)*x
^2 - b^3*d^m*(m + 1))*x^m*log(x) - (b^3*d^m*(m + 1)*log(c) - a*b^2*d^m*(m
+ 1) + (a*b^2*c^2*d^m*(m + 1) - (d^m*(m + 1)*log(c) + d^m)*b^3*c^2)*x^2)*x
^m)*sqrt(c*x + 1)*sqrt(-c*x + 1) - (b^3*d^m*(m + 1)*log(c) - a*b^2*d^m*(m
+ 1) - (b^3*c^2*d^m*(m + 1)*log(c) - a*b^2*c^2*d^m*(m + 1))*x^2)*x^m*log(
sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)^2 - 3*(b^3*d^m*(m + 1)*log(c)^2 - 2*a*b^
2*d^m*(m + 1)*log(c) + a^2*b*d^m*(m + 1) - (b^3*c^2*d^m*(m + 1)*log(c)^2 -
2*a*b^2*c^2*d^m*(m + 1)*log(c) + a^2*b*c^2*d^m*(m + 1))*x^2)*x^m*log(x) +
((b^3*c^2*d^m*(m + 1)*x^2 - b^3*d^m*(m + 1))*x^m*log(x)^3 - 3*(b^3*d^m*(m
+ 1)*log(c) - a*b^2*d^m*(m + 1) - (b^3*c^2*d^m*(m + 1)*log(c) - a*b^2*c^2
*d^m*(m + 1))*x^2)*x^m*log(x)^2 - 3*(b^3*d^m*(m + 1)*log(c)^2 - 2*a*b^2*d
^m*(m + 1)*log(c) + a^2*b*d^m*(m + 1) - (b^3*c^2*d^m*(m + 1)*log(c)^2 - 2*a
*b^2*c^2*d^m*(m + 1)*log(c) + a^2*b*c^2*d^m*(m + 1))*x^2)*x^m*log(x) - (b^
3*d^m*(m + 1)*log(c)^3 - 3*a*b^2*d^m*(m + 1)*log(c)^2 + 3*a^2*b*d^m*(m + 1
)*log(c) - (b^3*c^2*d^m*(m + 1)*log(c)^3 - 3*a*b^2*c^2*d^m*(m + 1)*log(c)^
2 + 3*a^2*b*c^2*d^m*(m + 1)*log(c))*x^2)*x^m)*sqrt(c*x + 1)*sqrt(-c*x + ...

```

**Giac [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^3 (dx)^m dx$$

input

```
integrate((d*x)^m*(a+b*arcsech(c*x))^3,x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)^3*(d*x)^m, x)
```

**Mupad [N/A]**

Not integrable

Time = 3.88 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^3 dx = \int (dx)^m \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right)^3 dx$$

input `int((d*x)^m*(a + b*acosh(1/(c*x)))^3,x)`output `int((d*x)^m*(a + b*acosh(1/(c*x)))^3, x)`**Reduce [N/A]**

Not integrable

Time = 0.57 (sec) , antiderivative size = 121, normalized size of antiderivative = 7.56

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^3 dx$$

$$= \frac{d^m (x^m a^3 x + 3 \int x^m \operatorname{asech}(cx) dx) a^2 b m + 3 \left( \int x^m \operatorname{asech}(cx) dx \right) a^2 b + \left( \int x^m \operatorname{asech}(cx) dx \right)^3 b^3 m + \left( \int x^m \operatorname{asech}(cx) dx \right)^2 a b m}{m + 1}$$

input `int((d*x)^m*(a+b*asech(c*x))^3,x)`output `(d**m*(x**m*a**3*x + 3*int(x**m*asech(c*x),x)*a**2*b*m + 3*int(x**m*asech(c*x),x)*a**2*b + int(x**m*asech(c*x)**3,x)*b**3*m + int(x**m*asech(c*x)**3,x)*b**3 + 3*int(x**m*asech(c*x)**2,x)*a*b**2*m + 3*int(x**m*asech(c*x)**2,x)*a*b**2))/m + 1)`

### 3.70 $\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^2 dx$

Optimal result	574
Mathematica [N/A]	574
Rubi [N/A]	575
Maple [N/A]	575
Fricas [N/A]	576
Sympy [N/A]	576
Maxima [N/A]	576
Giac [N/A]	577
Mupad [N/A]	578
Reduce [N/A]	578

#### Optimal result

Integrand size = 16, antiderivative size = 16

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^2 dx = \operatorname{Int}\left((dx)^m (a + b \operatorname{sech}^{-1}(cx))^2, x\right)$$

output `Defer(Int)((d*x)^m*(a+b*arcsech(c*x))^2,x)`

#### Mathematica [N/A]

Not integrable

Time = 1.80 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^2 dx$$

input `Integrate[(d*x)^m*(a + b*ArcSech[c*x])^2,x]`

output `Integrate[(d*x)^m*(a + b*ArcSech[c*x])^2, x]`

**Rubi [N/A]**

Not integrable

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^2 dx$$

↓ 6865

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^2 dx$$

input `Int[(d*x)^m*(a + b*ArcSech[c*x])^2,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.40 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int (dx)^m (a + b \operatorname{arcsech}(cx))^2 dx$$

input `int((d*x)^m*(a+b*arcsech(c*x))^2,x)`

output `int((d*x)^m*(a+b*arcsech(c*x))^2,x)`



**Fricas [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.88

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^2 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arcsech(c*x))^2,x, algorithm="fricas")`

output `integral((b^2*arcsech(c*x)^2 + 2*a*b*arcsech(c*x) + a^2)*(d*x)^m, x)`

**Sympy [N/A]**

Not integrable

Time = 2.61 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int (dx)^m (a + b \operatorname{asech}(cx))^2 dx$$

input `integrate((d*x)**m*(a+b*asech(c*x))**2,x)`

output `Integral((d*x)**m*(a + b*asech(c*x))**2, x)`

**Maxima [N/A]**

Not integrable

Time = 4.31 (sec) , antiderivative size = 704, normalized size of antiderivative = 44.00

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^2 (dx)^m dx$$

input `integrate((d*x)^m*(a+b*arcsech(c*x))^2,x, algorithm="maxima")`

output

```

b^2*d^m*x*x^m*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)^(2/(m + 1)) + (d*x)^(m +
1)*a^2/(d*(m + 1)) - integrate(-((b^2*c^2*d^m*(m + 1)*x^2 - b^2*d^m*(m +
1))*x^m*log(x)^2 - 2*(b^2*d^m*(m + 1)*log(c) - a*b*d^m*(m + 1) - (b^2*c^2*
d^m*(m + 1)*log(c) - a*b*c^2*d^m*(m + 1))*x^2)*x^m*log(x) + ((b^2*c^2*d^m*
(m + 1)*x^2 - b^2*d^m*(m + 1))*x^m*log(x)^2 - 2*(b^2*d^m*(m + 1)*log(c) -
a*b*d^m*(m + 1) - (b^2*c^2*d^m*(m + 1)*log(c) - a*b*c^2*d^m*(m + 1))*x^2)*
x^m*log(x) - (b^2*d^m*(m + 1)*log(c)^2 - 2*a*b*d^m*(m + 1)*log(c) - (b^2*c
^2*d^m*(m + 1)*log(c)^2 - 2*a*b*c^2*d^m*(m + 1)*log(c))*x^2)*x^m)*sqrt(c*x
+ 1)*sqrt(-c*x + 1) - (b^2*d^m*(m + 1)*log(c)^2 - 2*a*b*d^m*(m + 1)*log(c)
) - (b^2*c^2*d^m*(m + 1)*log(c)^2 - 2*a*b*c^2*d^m*(m + 1)*log(c))*x^2)*x^m
- 2*((b^2*c^2*d^m*(m + 1)*x^2 - b^2*d^m*(m + 1))*x^m*log(x) + ((b^2*c^2*d
^m*(m + 1)*x^2 - b^2*d^m*(m + 1))*x^m*log(x) - (b^2*d^m*(m + 1)*log(c) - a
*b*d^m*(m + 1) + (a*b*c^2*d^m*(m + 1) - (d^m*(m + 1)*log(c) + d^m)*b^2*c^2
)*x^2)*x^m)*sqrt(c*x + 1)*sqrt(-c*x + 1) - (b^2*d^m*(m + 1)*log(c) - a*b*d
^m*(m + 1) - (b^2*c^2*d^m*(m + 1)*log(c) - a*b*c^2*d^m*(m + 1))*x^2)*x^m)*
log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)/(c^2*(m + 1)*x^2 + (c^2*(m + 1)*x^2
- m - 1)*sqrt(c*x + 1)*sqrt(-c*x + 1) - m - 1), x)

```

**Giac [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)^2 (dx)^m dx$$

input

```
integrate((d*x)^m*(a+b*arcsech(c*x))^2,x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)^2*(d*x)^m, x)
```

**Mupad [N/A]**

Not integrable

Time = 3.95 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^2 dx = \int (dx)^m \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right)^2 dx$$

input `int((d*x)^m*(a + b*acosh(1/(c*x)))^2,x)`output `int((d*x)^m*(a + b*acosh(1/(c*x)))^2, x)`**Reduce [N/A]**

Not integrable

Time = 0.45 (sec) , antiderivative size = 80, normalized size of antiderivative = 5.00

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx))^2 dx$$

$$= \frac{d^m (x^m a^2 x + 2(\int x^m \operatorname{asech}(cx) dx) abm + 2(\int x^m \operatorname{asech}(cx) dx) ab + (\int x^m \operatorname{asech}(cx)^2 dx) b^2 m + (\int x^m \operatorname{asech}(cx) dx) b^2)}{m + 1}$$

input `int((d*x)^m*(a+b*asech(c*x))^2,x)`output `(d**m*(x**m*a**2*x + 2*int(x**m*asech(c*x),x)*a*b*m + 2*int(x**m*asech(c*x),x)*a*b + int(x**m*asech(c*x)**2,x)*b**2*m + int(x**m*asech(c*x)**2,x)*b**2))/m + 1)`

### 3.71 $\int (dx)^m (a + b \operatorname{sech}^{-1}(cx)) dx$

Optimal result	579
Mathematica [A] (verified)	579
Rubi [A] (warning: unable to verify)	580
Maple [F]	581
Fricas [F]	582
Sympy [F]	582
Maxima [F]	582
Giac [F]	583
Mupad [F(-1)]	583
Reduce [F]	583

#### Optimal result

Integrand size = 14, antiderivative size = 104

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{(dx)^{1+m} (a + b \operatorname{sech}^{-1}(cx))}{d(1+m)}$$

$$+ \frac{b(dx)^{1+m} \sqrt{\frac{1-cx}{1+cx}} (1+cx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2 x^2\right)}{d(1+m)^2 \sqrt{1-c^2 x^2}}$$

output

```
(d*x)^(1+m)*(a+b*arcsech(c*x))/d/(1+m)+b*(d*x)^(1+m)*((-c*x+1)/(c*x+1))^(1/2)*(c*x+1)*hypergeom([1/2, 1/2+1/2*m],[3/2+1/2*m],c^2*x^2)/d/(1+m)^2/(-c^2*x^2+1)^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.93

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{x(dx)^m \left( (1+m)(-1+cx) (a + b \operatorname{sech}^{-1}(cx)) - b \sqrt{\frac{1-cx}{1+cx}} \sqrt{1-c^2 x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2 x^2\right) \right)}{(1+m)^2 (-1+cx)}$$

input `Integrate[(d*x)^m*(a + b*ArcSech[c*x]), x]`

output `(x*(d*x)^m*((1 + m)*(-1 + c*x)*(a + b*ArcSech[c*x]) - b*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2]))/((1 + m)^2*(-1 + c*x))`

### Rubi [A] (warning: unable to verify)

Time = 0.26 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.84, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {6837, 135, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (dx)^m (a + b \operatorname{sech}^{-1}(cx)) dx \\
 & \quad \downarrow 6837 \\
 & \frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{(dx)^m}{\sqrt{1-cx}\sqrt{cx+1}} dx}{m+1} + \frac{(dx)^{m+1} (a + b \operatorname{sech}^{-1}(cx))}{d(m+1)} \\
 & \quad \downarrow 135 \\
 & \frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{(dx)^m}{\sqrt{1-c^2x^2}} dx}{m+1} + \frac{(dx)^{m+1} (a + b \operatorname{sech}^{-1}(cx))}{d(m+1)} \\
 & \quad \downarrow 278 \\
 & \frac{(dx)^{m+1} (a + b \operatorname{sech}^{-1}(cx))}{d(m+1)} + \\
 & \frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} (dx)^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, c^2x^2\right)}{d(m+1)^2}
 \end{aligned}$$

input `Int[(d*x)^m*(a + b*ArcSech[c*x]), x]`

output 
$$\frac{(d*x)^{(1+m)}*(a + b*\text{ArcSech}[c*x])}{d*(1+m)} + \frac{b*(d*x)^{(1+m)}*\text{Sqrt}[1 + c*x]^{(-1)}*\text{Sqrt}[1 + c*x]*\text{Hypergeometric2F1}[1/2, (1+m)/2, (3+m)/2, c^2*x^2]}{d*(1+m)^2}$$

### Defintions of rubi rules used

rule 135 
$$\text{Int}[\frac{(f_*)^{(p_*)}*((a_*) + (b_*)^{(x_*)^{(m_*)}*((c_*) + (d_*)^{(x_*)^{(n_*)}, x_]}]{:} \text{Int}[(a*c + b*d*x^2)^m*(f*x)^p, x] \text{ ; FreeQ}\{a, b, c, d, f, m, n, p\}, x] \ \&\& \ \text{EqQ}\{b*c + a*d, 0\} \ \&\& \ \text{EqQ}\{n, m\} \ \&\& \ \text{GtQ}\{a, 0\} \ \&\& \ \text{GtQ}\{c, 0\}$$

rule 278 
$$\text{Int}[\frac{(c_*)^{(x_*)^{(m_*)}*((a_*) + (b_*)^{(x_*)^2})^{(p_*)}, x\_Symbol]}{:} \text{Simp}[a^p*((c*x)^{(m+1)}/(c*(m+1)))*\text{Hypergeometric2F1}[-p, (m+1)/2, (m+1)/2 + 1, (-b)*(x^2/a)], x] \text{ ; FreeQ}\{a, b, c, m, p\}, x] \ \&\& \ !\text{IGtQ}\{p, 0\} \ \&\& \ (\text{ILtQ}\{p, 0\} \ || \ \text{GtQ}\{a, 0\})$$

rule 6837 
$$\text{Int}[\frac{(a_*) + \text{ArcSech}[c_*^{(x_*)}]*b_*}{(d_*)^{(x_*)^{(m_*)}}, x\_Symbol]}{:} \text{Simp}[(d*x)^{(m+1)}*((a + b*\text{ArcSech}[c*x])/(d*(m+1))), x] + \text{Simp}[b*(\text{Sqrt}[1 + c*x]/(m+1))*\text{Sqrt}[1/(1 + c*x)] \ \text{Int}[(d*x)^m/(\text{Sqrt}[1 - c*x]*\text{Sqrt}[1 + c*x]), x], x] \text{ ; FreeQ}\{a, b, c, d, m\}, x] \ \&\& \ \text{NeQ}\{m, -1\}$$

### Maple [F]

$$\int (dx)^m (a + b \operatorname{arcsech}(cx)) dx$$

input 
$$\text{int}((d*x)^m*(a+b*\text{arcsech}(c*x)),x)$$

output 
$$\text{int}((d*x)^m*(a+b*\text{arcsech}(c*x)),x)$$

**Fricas [F]**

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx)) dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)(dx)^m dx$$

input `integrate((d*x)^m*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral((b*arcsech(c*x) + a)*(d*x)^m, x)`

**Sympy [F]**

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx)) dx = \int (dx)^m (a + b \operatorname{asech}(cx)) dx$$

input `integrate((d*x)**m*(a+b*asech(c*x)),x)`

output `Integral((d*x)**m*(a + b*asech(c*x)), x)`

**Maxima [F]**

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx)) dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)(dx)^m dx$$

input `integrate((d*x)^m*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `(c^2*d^m*integrate(x^2*x^m/(c^2*(m + 1)*x^2 + (c^2*(m + 1)*x^2 - m - 1)*sqrt(c*x + 1)*sqrt(-c*x + 1) - m - 1), x) + (d^m*x*x^m*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1) - d^m*x*x^m*log(x))/(m + 1) - integrate((c^2*d^m*(m + 1)*x^2*log(c) - d^m*(m + 1)*log(c) + d^m)*x^m/(c^2*(m + 1)*x^2 - m - 1), x))*b + (d*x)^(m + 1)*a/(d*(m + 1))`

**Giac [F]**

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx)) dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)(dx)^m dx$$

input `integrate((d*x)^m*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*(d*x)^m, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (dx)^m (a + b \operatorname{sech}^{-1}(cx)) dx = \int (dx)^m \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int((d*x)^m*(a + b*acosh(1/(c*x))),x)`

output `int((d*x)^m*(a + b*acosh(1/(c*x))), x)`

**Reduce [F]**

$$\begin{aligned} & \int (dx)^m (a + b \operatorname{sech}^{-1}(cx)) dx \\ &= \frac{d^m (x^m a x + (\int x^m \operatorname{asech}(cx) dx) b m + (\int x^m \operatorname{asech}(cx) dx) b)}{m + 1} \end{aligned}$$

input `int((d*x)^m*(a+b*asech(c*x)),x)`

output `(d**m*(x**m*a*x + int(x**m*asech(c*x),x)*b*m + int(x**m*asech(c*x),x)*b))/  
(m + 1)`



$$3.72 \quad \int \frac{(dx)^m}{a+b\operatorname{sech}^{-1}(cx)} dx$$

Optimal result	584
Mathematica [N/A]	584
Rubi [N/A]	585
Maple [N/A]	585
Fricas [N/A]	586
Sympy [N/A]	586
Maxima [N/A]	586
Giac [N/A]	587
Mupad [N/A]	587
Reduce [N/A]	588

### Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{(dx)^m}{a + b\operatorname{sech}^{-1}(cx)} dx = \operatorname{Int}\left(\frac{(dx)^m}{a + b\operatorname{sech}^{-1}(cx)}, x\right)$$

output `Defer(Int)((d*x)^m/(a+b*arcsech(c*x)), x)`

### Mathematica [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(dx)^m}{a + b\operatorname{sech}^{-1}(cx)} dx = \int \frac{(dx)^m}{a + b\operatorname{sech}^{-1}(cx)} dx$$

input `Integrate[(d*x)^m/(a + b*ArcSech[c*x]), x]`

output `Integrate[(d*x)^m/(a + b*ArcSech[c*x]), x]`

**Rubi [N/A]**

Not integrable

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(dx)^m}{a + b \operatorname{sech}^{-1}(cx)} dx$$

↓ 6865

$$\int \frac{(dx)^m}{a + b \operatorname{sech}^{-1}(cx)} dx$$

input `Int[(d*x)^m/(a + b*ArcSech[c*x]),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.73 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{(dx)^m}{a + b \operatorname{arcsech}(cx)} dx$$

input `int((d*x)^m/(a+b*arcsech(c*x)),x)`

output `int((d*x)^m/(a+b*arcsech(c*x)),x)`

**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(dx)^m}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{(dx)^m}{b \operatorname{ar} \operatorname{sech}(cx) + a} dx$$

input `integrate((d*x)^m/(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral((d*x)^m/(b*arcsech(c*x) + a), x)`

**Sympy [N/A]**

Not integrable

Time = 0.51 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{(dx)^m}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{(dx)^m}{a + b \operatorname{asech}(cx)} dx$$

input `integrate((d*x)**m/(a+b*asech(c*x)),x)`

output `Integral((d*x)**m/(a + b*asech(c*x)), x)`

**Maxima [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(dx)^m}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{(dx)^m}{b \operatorname{ar} \operatorname{sech}(cx) + a} dx$$

input `integrate((d*x)^m/(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `integrate((d*x)^m/(b*arcsech(c*x) + a), x)`

### Giac [N/A]

Not integrable

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(dx)^m}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{(dx)^m}{b \operatorname{ar} \operatorname{sech}(cx) + a} dx$$

input `integrate((d*x)^m/(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((d*x)^m/(b*arcsech(c*x) + a), x)`

### Mupad [N/A]

Not integrable

Time = 3.58 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int \frac{(dx)^m}{a + b \operatorname{sech}^{-1}(cx)} dx = \int \frac{(dx)^m}{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)} dx$$

input `int((d*x)^m/(a + b*acosh(1/(c*x))),x)`

output `int((d*x)^m/(a + b*acosh(1/(c*x))), x)`

**Reduce [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int \frac{(dx)^m}{a + b \operatorname{sech}^{-1}(cx)} dx = d^m \left( \int \frac{x^m}{a \operatorname{sech}(cx) b + a} dx \right)$$

input `int((d*x)^m/(a+b*asech(c*x)),x)`output `d**m*int(x**m/(asech(c*x)*b + a),x)`

$$3.73 \quad \int \frac{(dx)^m}{(a+b\operatorname{sech}^{-1}(cx))^2} dx$$

Optimal result	589
Mathematica [N/A]	589
Rubi [N/A]	590
Maple [N/A]	590
Fricas [N/A]	591
Sympy [N/A]	591
Maxima [N/A]	591
Giac [N/A]	592
Mupad [N/A]	592
Reduce [N/A]	593

### Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{(dx)^m}{(a + b\operatorname{sech}^{-1}(cx))^2} dx = \operatorname{Int}\left(\frac{(dx)^m}{(a + b\operatorname{sech}^{-1}(cx))^2}, x\right)$$

output `Defer(Int)((d*x)^m/(a+b*arcsech(c*x))^2,x)`

### Mathematica [N/A]

Not integrable

Time = 0.60 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(dx)^m}{(a + b\operatorname{sech}^{-1}(cx))^2} dx = \int \frac{(dx)^m}{(a + b\operatorname{sech}^{-1}(cx))^2} dx$$

input `Integrate[(d*x)^m/(a + b*ArcSech[c*x])^2,x]`

output `Integrate[(d*x)^m/(a + b*ArcSech[c*x])^2, x]`

**Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(dx)^m}{(a + b \operatorname{sech}^{-1}(cx))^2} dx$$

↓ 6865

$$\int \frac{(dx)^m}{(a + b \operatorname{sech}^{-1}(cx))^2} dx$$

input `Int[(d*x)^m/(a + b*ArcSech[c*x])^2,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.61 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{(dx)^m}{(a + b \operatorname{arcsech}(cx))^2} dx$$

input `int((d*x)^m/(a+b*arcsech(c*x))^2,x)`

output `int((d*x)^m/(a+b*arcsech(c*x))^2,x)`

**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 32, normalized size of antiderivative = 2.00

$$\int \frac{(dx)^m}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{(dx)^m}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2} dx$$

input `integrate((d*x)^m/(a+b*arcsech(c*x))^2,x, algorithm="fricas")`

output `integral((d*x)^m/(b^2*arcsech(c*x)^2 + 2*a*b*arcsech(c*x) + a^2), x)`

**Sympy [N/A]**

Not integrable

Time = 1.48 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{(dx)^m}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{(dx)^m}{(a + b \operatorname{asech}(cx))^2} dx$$

input `integrate((d*x)**m/(a+b*asech(c*x))**2,x)`

output `Integral((d*x)**m/(a + b*asech(c*x))**2, x)`

**Maxima [N/A]**

Not integrable

Time = 0.76 (sec) , antiderivative size = 616, normalized size of antiderivative = 38.50

$$\int \frac{(dx)^m}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{(dx)^m}{(b \operatorname{ar} \operatorname{sech}(cx) + a)^2} dx$$

input `integrate((d*x)^m/(a+b*arcsech(c*x))^2,x, algorithm="maxima")`



output

```

-((c^2*d^m*x^3 - d^m*x)*sqrt(c*x + 1)*sqrt(-c*x + 1)*x^m + (c^2*d^m*x^3 -
d^m*x)*x^m)/((b^2*c^2*log(c) - a*b*c^2)*x^2 - b^2*log(c) - (b^2*log(c) + b
^2*log(x) - a*b)*sqrt(c*x + 1)*sqrt(-c*x + 1) + a*b - (b^2*c^2*x^2 - sqrt(
c*x + 1)*sqrt(-c*x + 1)*b^2 - b^2)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1) +
(b^2*c^2*x^2 - b^2)*log(x)) + integrate(((c^2*d^m*(m + 3)*x^2 - d^m*(m +
1))*(c*x + 1)*(c*x - 1)*x^m + (c^4*d^m*(m + 2)*x^4 - c^2*d^m*(3*m + 5)*x^2
+ 2*d^m*(m + 1))*sqrt(c*x + 1)*sqrt(-c*x + 1)*x^m + (c^4*d^m*(m + 1)*x^4
- 2*c^2*d^m*(m + 1)*x^2 + d^m*(m + 1))*x^m)/((b^2*c^4*log(c) - a*b*c^4)*x^
4 - (b^2*log(c) + b^2*log(x) - a*b)*(c*x + 1)*(c*x - 1) - 2*(b^2*c^2*log(c
) - a*b*c^2)*x^2 + b^2*log(c) - 2*((b^2*c^2*log(c) - a*b*c^2)*x^2 - b^2*lo
g(c) + a*b + (b^2*c^2*x^2 - b^2)*log(x))*sqrt(c*x + 1)*sqrt(-c*x + 1) - a*
b - (b^2*c^4*x^4 - 2*b^2*c^2*x^2 - (c*x + 1)*(c*x - 1)*b^2 - 2*(b^2*c^2*x^
2 - b^2)*sqrt(c*x + 1)*sqrt(-c*x + 1) + b^2)*log(sqrt(c*x + 1)*sqrt(-c*x +
1) + 1) + (b^2*c^4*x^4 - 2*b^2*c^2*x^2 + b^2)*log(x)), x)

```

**Giac [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{(dx)^m}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{(dx)^m}{(b \operatorname{arsech}(cx) + a)^2} dx$$

input

```
integrate((d*x)^m/(a+b*arcsech(c*x))^2,x, algorithm="giac")
```

output

```
integrate((d*x)^m/(b*arcsech(c*x) + a)^2, x)
```

**Mupad [N/A]**

Not integrable

Time = 3.84 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int \frac{(dx)^m}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = \int \frac{(dx)^m}{(a + b \operatorname{acosh}(\frac{1}{cx}))^2} dx$$

input `int((d*x)^m/(a + b*acosh(1/(c*x)))^2,x)`

output `int((d*x)^m/(a + b*acosh(1/(c*x)))^2, x)`

### Reduce [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.12

$$\int \frac{(dx)^m}{(a + b \operatorname{sech}^{-1}(cx))^2} dx = d^m \left( \int \frac{x^m}{a \operatorname{sech}(cx)^2 b^2 + 2 \operatorname{sech}(cx) ab + a^2} dx \right)$$

input `int((d*x)^m/(a+b*asech(c*x))^2,x)`

output `d**m*int(x**m/(asech(c*x)**2*b**2 + 2*asech(c*x)*a*b + a**2),x)`

### 3.74 $\int (d + ex)^3 (a + b \operatorname{sech}^{-1}(cx)) dx$

Optimal result	594
Mathematica [C] (verified)	595
Rubi [A] (verified)	595
Maple [A] (verified)	600
Fricas [B] (verification not implemented)	601
Sympy [F]	601
Maxima [A] (verification not implemented)	602
Giac [F]	602
Mupad [F(-1)]	603
Reduce [F]	603

#### Optimal result

Integrand size = 16, antiderivative size = 264

$$\int (d + ex)^3 (a + b \operatorname{sech}^{-1}(cx)) dx = -\frac{be(9c^2d^2 + e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{6c^4} - \frac{bde^2x \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{2c^2} - \frac{be^3x^2 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{12c^2} + \frac{(d + ex)^4 (a + b \operatorname{sech}^{-1}(cx))}{4e} + \frac{bd(2c^2d^2 + e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \arcsin(cx)}{2c^3} - \frac{bd^4 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \operatorname{arctanh}(\sqrt{1-c^2x^2})}{4e}$$

output

```
-1/6*b*e*(9*c^2*d^2+e^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^4-1/2*b*d*e^2*x*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^2-1/12*b*e^3*x^2*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^2+1/4*(e*x+d)^4*(a+b*arcsech(c*x))/e+1/2*b*d*(2*c^2*d^2+e^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arcsin(c*x)/c^3-1/4*b*d^4*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh((-c^2*x^2+1)^(1/2))/e
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.24 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.72

$$\int (d + ex)^3 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{1}{4} \left( 4ad^3x + 6ad^2ex^2 + 4ade^2x^3 + ae^3x^4 \right. \\ \left. - \frac{be\sqrt{\frac{1-cx}{1+cx}}(1+cx)(2e^2 + c^2(18d^2 + 6dex + e^2x^2))}{3c^4} \right. \\ \left. + bx(4d^3 + 6d^2ex + 4de^2x^2 + e^3x^3) \operatorname{sech}^{-1}(cx) \right. \\ \left. + \frac{2ibd(2c^2d^2 + e^2) \log\left(-2icx + 2\sqrt{\frac{1-cx}{1+cx}}(1+cx)\right)}{c^3} \right)$$

input

```
Integrate[(d + e*x)^3*(a + b*ArcSech[c*x]), x]
```

output

```
(4*a*d^3*x + 6*a*d^2*e*x^2 + 4*a*d*e^2*x^3 + a*e^3*x^4 - (b*e*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(2*e^2 + c^2*(18*d^2 + 6*d*e*x + e^2*x^2)))/(3*c^4) + b*x*(4*d^3 + 6*d^2*e*x + 4*d*e^2*x^2 + e^3*x^3)*ArcSech[c*x] + ((2*I)*b*d*(2*c^2*d^2 + e^2)*Log[(-2*I)*c*x + 2*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)])/c^3)/4
```

**Rubi [A] (verified)**

Time = 0.75 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.74, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {6842, 541, 25, 2340, 27, 2340, 27, 538, 223, 243, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex)^3 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$\begin{aligned}
 & \downarrow 6842 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(d+ex)^4}{x\sqrt{1-c^2x^2}} dx}{4e} + \frac{(d+ex)^4 (a + b\operatorname{sech}^{-1}(cx))}{4e} \\
 & \downarrow 541 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{\int -\frac{3c^2d^4+12c^2exd^3+12c^2e^3x^3d+2e^2(9c^2d^2+e^2)x^2}{x\sqrt{1-c^2x^2}} dx}{3c^2} - \frac{e^4x^2\sqrt{1-c^2x^2}}{3c^2} \right)}{4e} + \\
 & \frac{(d+ex)^4 (a + b\operatorname{sech}^{-1}(cx))}{4e} \\
 & \downarrow 25 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{3c^2d^4+12c^2exd^3+12c^2e^3x^3d+2e^2(9c^2d^2+e^2)x^2}{x\sqrt{1-c^2x^2}} dx}{3c^2} - \frac{e^4x^2\sqrt{1-c^2x^2}}{3c^2} \right)}{4e} + \\
 & \frac{(d+ex)^4 (a + b\operatorname{sech}^{-1}(cx))}{4e} \\
 & \downarrow 2340 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int -\frac{2(3c^4d^4+6c^2e(2c^2d^2+e^2)xd+2c^2e^2(9c^2d^2+e^2)x^2)}{x\sqrt{1-c^2x^2}} dx}{2c^2} - \frac{6de^3x\sqrt{1-c^2x^2}}{3c^2} - \frac{e^4x^2\sqrt{1-c^2x^2}}{3c^2} \right)}{4e} + \\
 & \frac{(d+ex)^4 (a + b\operatorname{sech}^{-1}(cx))}{4e} \\
 & \downarrow 27 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{3c^4d^4+6c^2e(2c^2d^2+e^2)xd+2c^2e^2(9c^2d^2+e^2)x^2}{x\sqrt{1-c^2x^2}} dx}{c^2} - \frac{6de^3x\sqrt{1-c^2x^2}}{3c^2} - \frac{e^4x^2\sqrt{1-c^2x^2}}{3c^2} \right)}{4e} + \\
 & \frac{(d+ex)^4 (a + b\operatorname{sech}^{-1}(cx))}{4e} \\
 & \downarrow 2340
 \end{aligned}$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int -\frac{3c^4d(c^2d^3+2e(2c^2d^2+e^2)x)}{x\sqrt{1-c^2x^2}} dx}{c^2} - \frac{2e^2\sqrt{1-c^2x^2}(9c^2d^2+e^2)}{3c^2} - \frac{6de^3x\sqrt{1-c^2x^2}}{3c^2} - \frac{e^4x^2\sqrt{1-c^2x^2}}{3c^2} \right) +$$

$$\frac{4e}{(d+ex)^4(a+b\operatorname{sech}^{-1}(cx))}$$

4e  
↓ 27

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3c^2d\int \frac{c^2d^3+2e(2c^2d^2+e^2)x}{x\sqrt{1-c^2x^2}} dx - 2e^2\sqrt{1-c^2x^2}(9c^2d^2+e^2)}{c^2} - \frac{6de^3x\sqrt{1-c^2x^2}}{3c^2} - \frac{e^4x^2\sqrt{1-c^2x^2}}{3c^2} \right) +$$

$$\frac{4e}{(d+ex)^4(a+b\operatorname{sech}^{-1}(cx))}$$

4e  
↓ 538

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3c^2d\left(c^2d^3\int \frac{1}{x\sqrt{1-c^2x^2}} dx + 2e(2c^2d^2+e^2)\int \frac{1}{\sqrt{1-c^2x^2}} dx\right) - 2e^2\sqrt{1-c^2x^2}(9c^2d^2+e^2)}{c^2} - \frac{6de^3x\sqrt{1-c^2x^2}}{3c^2} - \frac{e^4x^2\sqrt{1-c^2x^2}}{3c^2} \right) +$$

$$\frac{4e}{(d+ex)^4(a+b\operatorname{sech}^{-1}(cx))}$$

4e  
↓ 223

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3c^2d\left(c^2d^3\int \frac{1}{x\sqrt{1-c^2x^2}} dx + \frac{2e\arcsin(cx)(2c^2d^2+e^2)}{c}\right) - 2e^2\sqrt{1-c^2x^2}(9c^2d^2+e^2)}{c^2} - \frac{6de^3x\sqrt{1-c^2x^2}}{3c^2} - \frac{e^4x^2\sqrt{1-c^2x^2}}{3c^2} \right) +$$

$$\frac{4e}{(d+ex)^4(a+b\operatorname{sech}^{-1}(cx))}$$

4e  
↓ 243

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{3c^2d\left(\frac{1}{2}c^2d^3\int\frac{1}{x^2\sqrt{1-c^2x^2}}dx^2+\frac{2e\arcsin(cx)(2c^2d^2+e^2)}{c}\right)-2e^2\sqrt{1-c^2x^2}(9c^2d^2+e^2)}{c^2}-\frac{6de^3x\sqrt{1-c^2x^2}}{3c^2}-\frac{e^4x^2\sqrt{1-c^2x^2}}{3c^2}\right)$$

$$\frac{(d+ex)^4(a+b\operatorname{sech}^{-1}(cx))}{4e}$$

73

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{3c^2d\left(\frac{2e\arcsin(cx)(2c^2d^2+e^2)}{c}-d^3\int\frac{1}{c^2}\frac{1-x^4}{c^2}d\sqrt{1-c^2x^2}\right)-2e^2\sqrt{1-c^2x^2}(9c^2d^2+e^2)}{c^2}-\frac{6de^3x\sqrt{1-c^2x^2}}{3c^2}-\frac{e^4x^2\sqrt{1-c^2x^2}}{3c^2}\right)$$

$$\frac{(d+ex)^4(a+b\operatorname{sech}^{-1}(cx))}{4e}$$

221

$$\frac{(d+ex)^4(a+b\operatorname{sech}^{-1}(cx))}{4e} +$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{3c^2d\left(\frac{2e\arcsin(cx)(2c^2d^2+e^2)}{c}-c^2d^3\operatorname{arctanh}(\sqrt{1-c^2x^2})\right)-2e^2\sqrt{1-c^2x^2}(9c^2d^2+e^2)}{c^2}-\frac{6de^3x\sqrt{1-c^2x^2}}{3c^2}-\frac{e^4x^2\sqrt{1-c^2x^2}}{3c^2}\right)$$

4e

input `Int[(d + e*x)^3*(a + b*ArcSech[c*x]),x]`

output `((d + e*x)^4*(a + b*ArcSech[c*x]))/(4*e) + (b*sqrt[(1 + c*x)^(-1)]*sqrt[1 + c*x]*(-1/3*(e^4*x^2*sqrt[1 - c^2*x^2])/c^2 + (-6*d*e^3*x*sqrt[1 - c^2*x^2] + (-2*e^2*(9*c^2*d^2 + e^2)*sqrt[1 - c^2*x^2] + 3*c^2*d*((2*e*(2*c^2*d^2 + e^2)*ArcSin[c*x])/c - c^2*d^3*ArcTanh[Sqrt[1 - c^2*x^2]])))/c^2)/(3*c^2)))/(4*e)`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`
- rule 538 `Int[((c_) + (d_.)*(x_))/((x_)*Sqrt[(a_) + (b_.)*(x_)^2]), x_Symbol] := Simp[c Int[1/(x*Sqrt[a + b*x^2]), x], x] + Simp[d Int[1/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d}, x]`
- rule 541 `Int[(x_)^(m_.)*((c_) + (d_.)*(x_)^(n_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[d^n*x^(m + n - 1)*((a + b*x^2)^(p + 1)/(b*(m + n + 2*p + 1))), x] + Simp[1/(b*(m + n + 2*p + 1)) Int[x^m*(a + b*x^2)^p*ExpandToSum[b*(m + n + 2*p + 1)*(c + d*x)^n - b*d^n*(m + n + 2*p + 1)*x^n - a*d^n*(m + n - 1)*x^(n - 2), x], x] /; FreeQ[{a, b, c, d, m, p}, x] && IGtQ[n, 1] && IGtQ[m, -2] && GtQ[p, -1] && IntegerQ[2*p]`



rule 2340

```
Int[(Pq_)*((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[
{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[f*(c*x)^(m + q - 1)
*((a + b*x^2)^(p + 1)/(b*c^(q - 1)*(m + q + 2*p + 1))), x] + Simp[1/(b*(m
+ q + 2*p + 1)) Int[(c*x)^m*(a + b*x^2)^p*ExpandToSum[b*(m + q + 2*p + 1)
*Pq - b*f*(m + q + 2*p + 1)*x^q - a*f*(m + q - 1)*x^(q - 2), x], x] /;
GtQ[q, 1] && NeQ[m + q + 2*p + 1, 0] /; FreeQ[{a, b, c, m, p}, x] && PolyQ
[Pq, x] && (!IGtQ[m, 0] || IGtQ[p + 1/2, -1])
```

rule 6842

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_))*((d_) + (e_)*(x_)^(m_)), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*ArcSech[c*x])/(e*(m + 1))), x] + Simp[
b*(Sqrt[1 + c*x]/(e*(m + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x)^(m + 1)/(x*
Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[m, -1]
```

### Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.00

method	result
parts	$\frac{a(ex+d)^4}{4e} + \frac{b \left( \frac{c e^3 \operatorname{arcsech}(cx)x^4}{4} + c e^2 \operatorname{arcsech}(cx)x^3 d + \frac{3ce \operatorname{arcsech}(cx)x^2 d^2}{2} + \operatorname{arcsech}(cx)cx d^3 + \frac{c \operatorname{arcsech}(cx)d^4}{4e} - \sqrt{-\frac{cx}{c}} \right)}{4e}$
derivativedivides	$\frac{a(cex+cd)^4}{4c^3e} + \frac{b \left( \frac{\operatorname{arcsech}(cx)c^4 d^4}{4e} + \operatorname{arcsech}(cx)c^4 d^3 x + \frac{3e \operatorname{arcsech}(cx)c^4 d^2 x^2}{2} + e^2 \operatorname{arcsech}(cx)c^4 d x^3 + \frac{e^3 \operatorname{arcsech}(cx)c^4 x^4}{4} + \sqrt{-\frac{cx}{c}} \right)}{4c^3e}$
default	$\frac{a(cex+cd)^4}{4c^3e} + \frac{b \left( \frac{\operatorname{arcsech}(cx)c^4 d^4}{4e} + \operatorname{arcsech}(cx)c^4 d^3 x + \frac{3e \operatorname{arcsech}(cx)c^4 d^2 x^2}{2} + e^2 \operatorname{arcsech}(cx)c^4 d x^3 + \frac{e^3 \operatorname{arcsech}(cx)c^4 x^4}{4} + \sqrt{-\frac{cx}{c}} \right)}{4c^3e}$

```
input int((e*x+d)^3*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)
```

```
output 1/4*a*(e*x+d)^4/e+b/c*(1/4*c*e^3*arcsech(c*x)*x^4+c*e^2*arcsech(c*x)*x^3*d
+3/2*c*e*arcsech(c*x)*x^2*d^2+arcsech(c*x)*c*x*d^3+1/4*c/e*arcsech(c*x)*d^
4-1/12/c^2/e*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(3*c^4*d^4*arctanh
(1/(-c^2*x^2+1)^(1/2))-12*c^3*d^3*e*arcsin(c*x)+e^4*(-c^2*x^2+1)^(1/2)*c^2
*x^2+6*c^2*d*e^3*x*(-c^2*x^2+1)^(1/2)+18*c^2*d^2*e^2*(-c^2*x^2+1)^(1/2)-6*
c*d*e^3*arcsin(c*x)+2*e^4*(-c^2*x^2+1)^(1/2))/(-c^2*x^2+1)^(1/2))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 358 vs.  $2(144) = 288$ .

Time = 0.19 (sec) , antiderivative size = 358, normalized size of antiderivative = 1.36

$$\int (d + ex)^3 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{3ac^3e^3x^4 + 12ac^3de^2x^3 + 18ac^3d^2ex^2 + 12ac^3d^3x - 12(2bc^2d^3 + bde^2) \arctan\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{cx}\right) - 3(4b^2c^3d^3 + 6b^2c^3de^2 + 4b^2c^3e^3x^4 + 4b^2c^3d^2ex^3 + 6b^2c^3d^2e^2x^2 + 4b^2c^3d^3x - 4b^2c^3d^3 - 6b^2c^3d^2e - 4b^2c^3de^2 - b^2c^3e^3) \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{cx}\right) - (b^2c^2e^3x^3 + 6b^2c^2de^2x^2 + 2(9b^2c^2d^2e + b^2e^3)x) \sqrt{-\frac{c^2x^2-1}{c^2x^2}}}{c^3}$$

input `integrate((e*x+d)^3*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `1/12*(3*a*c^3*e^3*x^4 + 12*a*c^3*d*e^2*x^3 + 18*a*c^3*d^2*e*x^2 + 12*a*c^3*d^3*x - 12*(2*b*c^2*d^3 + b*d*e^2)*arctan((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/(c*x)) - 3*(4*b*c^3*d^3 + 6*b*c^3*d^2*e + 4*b*c^3*d*e^2 + b*c^3*e^3)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) + 3*(b*c^3*e^3*x^4 + 4*b*c^3*d*e^2*x^3 + 6*b*c^3*d^2*e*x^2 + 4*b*c^3*d^3*x - 4*b*c^3*d^3 - 6*b*c^3*d^2*e - 4*b*c^3*d*e^2 - b*c^3*e^3)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (b*c^2*e^3*x^3 + 6*b*c^2*d*e^2*x^2 + 2*(9*b*c^2*d^2*e + b*e^3)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/c^3`

**Sympy [F]**

$$\int (d + ex)^3 (a + b \operatorname{asech}(cx)) dx = \int (a + b \operatorname{asech}(cx)) (d + ex)^3 dx$$

input `integrate((e*x+d)**3*(a+b*asech(c*x)),x)`

output `Integral((a + b*asech(c*x))*(d + e*x)**3, x)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.84

$$\begin{aligned}
& \int (d + ex)^3 (a + b \operatorname{sech}^{-1}(cx)) dx \\
&= \frac{1}{4} a e^3 x^4 + a d e^2 x^3 + \frac{3}{2} a d^2 e x^2 + \frac{3}{2} \left( x^2 \operatorname{arsech}(cx) - \frac{x \sqrt{\frac{1}{c^2 x^2} - 1}}{c} \right) b d^2 e \\
&+ \frac{1}{2} \left( 2 x^3 \operatorname{arsech}(cx) - \frac{\frac{\sqrt{\frac{1}{c^2 x^2} - 1}}{c^2 \left( \frac{1}{c^2 x^2} - 1 \right) + c^2} + \frac{\arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right)}{c^2}}{c} \right) b d e^2 \\
&+ \frac{1}{12} \left( 3 x^4 \operatorname{arsech}(cx) + \frac{c^2 x^3 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{3}{2}} - 3 x \sqrt{\frac{1}{c^2 x^2} - 1}}{c^3} \right) b e^3 \\
&+ a d^3 x + \frac{\left( c x \operatorname{arsech}(cx) - \arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right) \right) b d^3}{c}
\end{aligned}$$

input `integrate((e*x+d)^3*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `1/4*a*e^3*x^4 + a*d*e^2*x^3 + 3/2*a*d^2*e*x^2 + 3/2*(x^2*arcsech(c*x) - x*sqrt(1/(c^2*x^2) - 1)/c)*b*d^2*e + 1/2*(2*x^3*arcsech(c*x) - (sqrt(1/(c^2*x^2) - 1)/(c^2*(1/(c^2*x^2) - 1) + c^2) + arctan(sqrt(1/(c^2*x^2) - 1))/c^2)/c)*b*d*e^2 + 1/12*(3*x^4*arcsech(c*x) + (c^2*x^3*(1/(c^2*x^2) - 1)^(3/2) - 3*x*sqrt(1/(c^2*x^2) - 1))/c^3)*b*e^3 + a*d^3*x + (c*x*arcsech(c*x) - arctan(sqrt(1/(c^2*x^2) - 1)))*b*d^3/c`

**Giac [F]**

$$\int (d + ex)^3 (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex + d)^3 (b \operatorname{arsech}(cx) + a) dx$$

input `integrate((e*x+d)^3*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((e*x + d)^3*(b*arcsech(c*x) + a), x)`

### Mupad [F(-1)]

Timed out.

$$\int (d + ex)^3 (a + b \operatorname{sech}^{-1}(cx)) dx = \int \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) (d + ex)^3 dx$$

input `int((a + b*acosh(1/(c*x)))*(d + e*x)^3,x)`

output `int((a + b*acosh(1/(c*x)))*(d + e*x)^3, x)`

### Reduce [F]

$$\begin{aligned} \int (d + ex)^3 (a + b \operatorname{sech}^{-1}(cx)) dx &= \left( \int \operatorname{asech}(cx) dx \right) b d^3 + \left( \int \operatorname{asech}(cx) x^3 dx \right) b e^3 \\ &+ 3 \left( \int \operatorname{asech}(cx) x^2 dx \right) b d e^2 \\ &+ 3 \left( \int \operatorname{asech}(cx) x dx \right) b d^2 e + a d^3 x \\ &+ \frac{3a d^2 e x^2}{2} + a d e^2 x^3 + \frac{a e^3 x^4}{4} \end{aligned}$$

input `int((e*x+d)^3*(a+b*asech(c*x)),x)`

output `(4*int(asech(c*x),x)*b*d**3 + 4*int(asech(c*x)*x**3,x)*b*e**3 + 12*int(asech(c*x)*x**2,x)*b*d*e**2 + 12*int(asech(c*x)*x,x)*b*d**2*e + 4*a*d**3*x + 6*a*d**2*e*x**2 + 4*a*d*e**2*x**3 + a*e**3*x**4)/4`

### 3.75 $\int (d + ex)^2 (a + b \operatorname{sech}^{-1}(cx)) dx$

Optimal result	604
Mathematica [C] (verified)	605
Rubi [A] (verified)	605
Maple [A] (verified)	609
Fricas [B] (verification not implemented)	610
Sympy [F]	610
Maxima [A] (verification not implemented)	611
Giac [F]	611
Mupad [F(-1)]	612
Reduce [F]	612

#### Optimal result

Integrand size = 16, antiderivative size = 201

$$\int (d + ex)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = -\frac{bde\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{c^2} - \frac{be^2x\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{6c^2} + \frac{(d + ex)^3 (a + b \operatorname{sech}^{-1}(cx))}{3e} + \frac{b(6c^2d^2 + e^2)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\arcsin(cx)}{6c^3} - \frac{bd^3\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\operatorname{arctanh}(\sqrt{1-c^2x^2})}{3e}$$

output

```
-b*d*e*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^2-1/6*b*e^2*x*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^2+1/3*(e*x+d)^3*(a+b*arcsech(c*x))/e+1/6*b*(6*c^2*d^2+e^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arcsin(c*x)/c^3-1/3*b*d^3*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh((-c^2*x^2+1)^(1/2))/e
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.73

$$\int (d + ex)^2 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{-bce \sqrt{\frac{1-cx}{1+cx}} (1+cx)(6d+ex) + 2ac^3x(3d^2+3dex+e^2x^2) + 2bc^3x(3d^2+3dex+e^2x^2) \operatorname{sech}^{-1}(cx) + ib}{6c^3}$$

input

```
Integrate[(d + e*x)^2*(a + b*ArcSech[c*x]), x]
```

output

```
(-(b*c*e*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(6*d + e*x)) + 2*a*c^3*x*(3*d^2 + 3*d*e*x + e^2*x^2) + 2*b*c^3*x*(3*d^2 + 3*d*e*x + e^2*x^2)*ArcSech[c*x] + I*b*(6*c^2*d^2 + e^2)*Log[(-2*I)*c*x + 2*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)])/(6*c^3)
```

**Rubi [A] (verified)**

Time = 0.48 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.74, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.688$ , Rules used = {6842, 541, 25, 2340, 25, 27, 538, 223, 243, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex)^2 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$\downarrow \text{6842}$$

$$\frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{(d+ex)^3}{x \sqrt{1-c^2x^2}} dx}{3e} + \frac{(d+ex)^3 (a + b \operatorname{sech}^{-1}(cx))}{3e}$$

$$\downarrow \text{541}$$

$$\begin{aligned}
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{\int -\frac{2c^2d^3+6c^2e^2x^2d+e(6c^2d^2+e^2)x}{x\sqrt{1-c^2x^2}} dx}{2c^2} - \frac{e^3x\sqrt{1-c^2x^2}}{2c^2} \right)}{(d+ex)^3 (a+b\operatorname{sech}^{-1}(cx))} + \\
 & \qquad \qquad \qquad \frac{3e}{3e} \\
 & \qquad \qquad \qquad \downarrow \text{25} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{2c^2d^3+6c^2e^2x^2d+e(6c^2d^2+e^2)x}{x\sqrt{1-c^2x^2}} dx}{2c^2} - \frac{e^3x\sqrt{1-c^2x^2}}{2c^2} \right)}{3e} + \frac{(d+ex)^3 (a+b\operatorname{sech}^{-1}(cx))}{3e} \\
 & \qquad \qquad \qquad \downarrow \text{2340} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{\int -\frac{c^2(2c^2d^3+e(6c^2d^2+e^2)x)}{x\sqrt{1-c^2x^2}} dx}{2c^2} - 6de^2\sqrt{1-c^2x^2} - \frac{e^3x\sqrt{1-c^2x^2}}{2c^2} \right)}{(d+ex)^3 (a+b\operatorname{sech}^{-1}(cx))} + \\
 & \qquad \qquad \qquad \frac{3e}{3e} \\
 & \qquad \qquad \qquad \downarrow \text{25} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{c^2(2c^2d^3+e(6c^2d^2+e^2)x)}{x\sqrt{1-c^2x^2}} dx}{2c^2} - 6de^2\sqrt{1-c^2x^2} - \frac{e^3x\sqrt{1-c^2x^2}}{2c^2} \right)}{(d+ex)^3 (a+b\operatorname{sech}^{-1}(cx))} + \\
 & \qquad \qquad \qquad \frac{3e}{3e} \\
 & \qquad \qquad \qquad \downarrow \text{27} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{2c^2d^3+e(6c^2d^2+e^2)x}{x\sqrt{1-c^2x^2}} dx - 6de^2\sqrt{1-c^2x^2} - \frac{e^3x\sqrt{1-c^2x^2}}{2c^2} \right)}{(d+ex)^3 (a+b\operatorname{sech}^{-1}(cx))} + \\
 & \qquad \qquad \qquad \frac{3e}{3e} \\
 & \qquad \qquad \qquad \downarrow \text{538} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{2c^2d^3 \int \frac{1}{x\sqrt{1-c^2x^2}} dx + e(6c^2d^2+e^2) \int \frac{1}{\sqrt{1-c^2x^2}} dx - 6de^2\sqrt{1-c^2x^2} - \frac{e^3x\sqrt{1-c^2x^2}}{2c^2} \right)}{(d+ex)^3 (a+b\operatorname{sech}^{-1}(cx))} + \\
 & \qquad \qquad \qquad \frac{3e}{3e}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 223 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{2c^2d^3\int\frac{1}{x\sqrt{1-c^2x^2}}dx+\frac{e\arcsin(cx)(6c^2d^2+e^2)}{2c^2}-6de^2\sqrt{1-c^2x^2}}{2c^2}-\frac{e^3x\sqrt{1-c^2x^2}}{2c^2}\right)}{(d+ex)^3(a+b\operatorname{sech}^{-1}(cx))} + \\
& \downarrow 243 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{c^2d^3\int\frac{1}{x^2\sqrt{1-c^2x^2}}dx^2+\frac{e\arcsin(cx)(6c^2d^2+e^2)}{2c^2}-6de^2\sqrt{1-c^2x^2}}{2c^2}-\frac{e^3x\sqrt{1-c^2x^2}}{2c^2}\right)}{(d+ex)^3(a+b\operatorname{sech}^{-1}(cx))} + \\
& \downarrow 73 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{-2d^3\int\frac{1}{\frac{1}{c^2}-x^4}d\sqrt{1-c^2x^2}+\frac{e\arcsin(cx)(6c^2d^2+e^2)}{2c^2}-6de^2\sqrt{1-c^2x^2}}{2c^2}-\frac{e^3x\sqrt{1-c^2x^2}}{2c^2}\right)}{(d+ex)^3(a+b\operatorname{sech}^{-1}(cx))} + \\
& \downarrow 221 \\
& \frac{(d+ex)^3(a+b\operatorname{sech}^{-1}(cx))}{3e} + \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{\frac{e\arcsin(cx)(6c^2d^2+e^2)}{c}-2c^2d^3\operatorname{arctanh}(\sqrt{1-c^2x^2})-6de^2\sqrt{1-c^2x^2}}{2c^2}-\frac{e^3x\sqrt{1-c^2x^2}}{2c^2}\right)}{3e}
\end{aligned}$$

input `Int[(d + e*x)^2*(a + b*ArcSech[c*x]), x]`

output `((d + e*x)^3*(a + b*ArcSech[c*x]))/(3*e) + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/2*(e^3*x*Sqrt[1 - c^2*x^2])/c^2 + (-6*d*e^2*Sqrt[1 - c^2*x^2] + (e*(6*c^2*d^2 + e^2)*ArcSin[c*x])/c - 2*c^2*d^3*ArcTanh[Sqrt[1 - c^2*x^2]])/(2*c^2)))/(3*e)`



## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`
- rule 538 `Int[((c_) + (d_.)*(x_))/((x_)*Sqrt[(a_) + (b_.)*(x_)^2]), x_Symbol] := Simp[c Int[1/(x*Sqrt[a + b*x^2]), x], x] + Simp[d Int[1/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d}, x]`
- rule 541 `Int[(x_)^(m_.)*((c_) + (d_.)*(x_)^(n_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[d^n*x^(m + n - 1)*((a + b*x^2)^(p + 1)/(b*(m + n + 2*p + 1))), x] + Simp[1/(b*(m + n + 2*p + 1)) Int[x^m*(a + b*x^2)^p*ExpandToSum[b*(m + n + 2*p + 1)*(c + d*x)^n - b*d^n*(m + n + 2*p + 1)*x^n - a*d^n*(m + n - 1)*x^(n - 2), x], x] /; FreeQ[{a, b, c, d, m, p}, x] && IGtQ[n, 1] && IGtQ[m, -2] && GtQ[p, -1] && IntegerQ[2*p]`

rule 2340

```
Int[(Pq_)*((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[
{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[f*(c*x)^(m + q - 1)
*((a + b*x^2)^(p + 1)/(b*c^(q - 1)*(m + q + 2*p + 1))), x] + Simp[1/(b*(m
+ q + 2*p + 1)) Int[(c*x)^m*(a + b*x^2)^p*ExpandToSum[b*(m + q + 2*p + 1)
*Pq - b*f*(m + q + 2*p + 1)*x^q - a*f*(m + q - 1)*x^(q - 2), x], x] /;
GtQ[q, 1] && NeQ[m + q + 2*p + 1, 0] /; FreeQ[{a, b, c, m, p}, x] && PolyQ
[Pq, x] && (!IGtQ[m, 0] || IGtQ[p + 1/2, -1])
```

rule 6842

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_))*((d_) + (e_)*(x_)^(m_)), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*ArcSech[c*x])/(e*(m + 1))), x] + Simp[
b*(Sqrt[1 + c*x]/(e*(m + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x)^(m + 1)/(x*
Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[m, -1]
```

### Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 199, normalized size of antiderivative = 0.99

method	result
parts	$\frac{a(ex+d)^3}{3e} + \frac{b \left( \frac{c e^2 \operatorname{arcsech}(cx)x^3}{3} + ce \operatorname{arcsech}(cx)x^2 d + \operatorname{arcsech}(cx)cx d^2 + \frac{c \operatorname{arcsech}(cx)d^3}{3e} + \frac{\sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}} (-2c^3 d^3 a)}{c} \right)}{c^2}$
derivativedivides	$\frac{a(cex+cd)^3}{3c^2 e} + \frac{b \left( \frac{\operatorname{arcsech}(cx)c^3 d^3}{3e} + \operatorname{arcsech}(cx)c^3 d^2 x + e \operatorname{arcsech}(cx)c^3 d x^2 + \frac{e^2 \operatorname{arcsech}(cx)c^3 x^3}{3} + \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (-2c^3 d^3 a)}{c} \right)}{c^2}$
default	$\frac{a(cex+cd)^3}{3c^2 e} + \frac{b \left( \frac{\operatorname{arcsech}(cx)c^3 d^3}{3e} + \operatorname{arcsech}(cx)c^3 d^2 x + e \operatorname{arcsech}(cx)c^3 d x^2 + \frac{e^2 \operatorname{arcsech}(cx)c^3 x^3}{3} + \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (-2c^3 d^3 a)}{c} \right)}{c^2}$

input

```
int((e*x+d)^2*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)
```

output

```
1/3*a*(e*x+d)^3/e+b/c*(1/3*c*e^2*arcsech(c*x)*x^3+c*e*arcsech(c*x)*x^2*d+a
rcsech(c*x)*c*x*d^2+1/3*c/e*arcsech(c*x)*d^3+1/6/c/e*(-(c*x-1)/c/x)^(1/2)*
x*((c*x+1)/c/x)^(1/2)*(-2*c^3*d^3*arctanh(1/(-c^2*x^2+1)^(1/2))+6*c^2*d^2*
e*arcsin(c*x)-6*c*d*e^2*(-c^2*x^2+1)^(1/2)-e^3*c*x*(-c^2*x^2+1)^(1/2)+e^3*
arcsin(c*x))/(-c^2*x^2+1)^(1/2))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 280 vs.  $2(107) = 214$ .

Time = 0.16 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.39

$$\int (d + ex)^2 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{2ac^3e^2x^3 + 6ac^3dex^2 + 6ac^3d^2x - 2(6bc^2d^2 + be^2) \arctan\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{cx}\right) - 2(3bc^3d^2 + 3bc^3de + bc^3}{}$$

input `integrate((e*x+d)^2*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `1/6*(2*a*c^3*e^2*x^3 + 6*a*c^3*d*e*x^2 + 6*a*c^3*d^2*x - 2*(6*b*c^2*d^2 + b*e^2)*arctan((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/(c*x)) - 2*(3*b*c^3*d^2 + 3*b*c^3*d*e + b*c^3*e^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) + 2*(b*c^3*e^2*x^3 + 3*b*c^3*d*e*x^2 + 3*b*c^3*d^2*x - 3*b*c^3*d^2 - 3*b*c^3*d*e - b*c^3*e^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (b*c^2*e^2*x^2 + 6*b*c^2*d*e*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/c^3`

**Sympy [F]**

$$\int (d + ex)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \int (a + b \operatorname{asech}(cx)) (d + ex)^2 dx$$

input `integrate((e*x+d)**2*(a+b*asech(c*x)),x)`

output `Integral((a + b*asech(c*x))*(d + e*x)**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.76

$$\int (d + ex)^2 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{1}{3} a e^2 x^3 + a d e x^2 + \left( x^2 \operatorname{ar} \operatorname{sech}(cx) - \frac{x \sqrt{\frac{1}{c^2 x^2} - 1}}{c} \right) b d e$$

$$+ \frac{1}{6} \left( 2 x^3 \operatorname{ar} \operatorname{sech}(cx) - \frac{\frac{\sqrt{\frac{1}{c^2 x^2} - 1}}{c^2 \left( \frac{1}{c^2 x^2} - 1 \right) + c^2} + \frac{\arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right)}{c^2}}{c} \right) b e^2$$

$$+ a d^2 x + \frac{\left( c x \operatorname{ar} \operatorname{sech}(cx) - \arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right) \right) b d^2}{c}$$

input `integrate((e*x+d)^2*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `1/3*a*e^2*x^3 + a*d*e*x^2 + (x^2*arcsech(c*x) - x*sqrt(1/(c^2*x^2) - 1)/c) *b*d*e + 1/6*(2*x^3*arcsech(c*x) - (sqrt(1/(c^2*x^2) - 1)/(c^2*(1/(c^2*x^2) - 1) + c^2) + arctan(sqrt(1/(c^2*x^2) - 1)/c^2)/c)*b*e^2 + a*d^2*x + (c*x*arcsech(c*x) - arctan(sqrt(1/(c^2*x^2) - 1)))*b*d^2/c`

**Giac [F]**

$$\int (d + ex)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex + d)^2 (b \operatorname{ar} \operatorname{sech}(cx) + a) dx$$

input `integrate((e*x+d)^2*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((e*x + d)^2*(b*arcsech(c*x) + a), x)`

**Mupad [F(-1)]**

Timed out.

$$\int (d + ex)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \int \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) (d + ex)^2 dx$$

input `int((a + b*acosh(1/(c*x)))*(d + e*x)^2,x)`output `int((a + b*acosh(1/(c*x)))*(d + e*x)^2, x)`**Reduce [F]**

$$\int (d + ex)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \left( \int a \operatorname{sech}(cx) dx \right) b d^2 + \left( \int a \operatorname{sech}(cx) x^2 dx \right) b e^2 + 2 \left( \int a \operatorname{sech}(cx) x dx \right) b d e + a d^2 x + a d e x^2 + \frac{a e^2 x^3}{3}$$

input `int((e*x+d)^2*(a+b*asech(c*x)),x)`output `(3*int(asech(c*x),x)*b*d**2 + 3*int(asech(c*x)*x**2,x)*b*e**2 + 6*int(asech(c*x)*x,x)*b*d*e + 3*a*d**2*x + 3*a*d*e*x**2 + a*e**2*x**3)/3`

### 3.76 $\int (d + ex) (a + b \operatorname{sech}^{-1}(cx)) dx$

Optimal result	613
Mathematica [A] (verified)	614
Rubi [A] (verified)	614
Maple [A] (verified)	617
Fricas [B] (verification not implemented)	618
Sympy [F]	618
Maxima [A] (verification not implemented)	619
Giac [F]	619
Mupad [B] (verification not implemented)	619
Reduce [F]	620

#### Optimal result

Integrand size = 14, antiderivative size = 142

$$\int (d + ex) (a + b \operatorname{sech}^{-1}(cx)) dx = -\frac{be\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{2c^2} + \frac{(d+ex)^2(a+b\operatorname{sech}^{-1}(cx))}{2e} + \frac{bd\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\arcsin(cx)}{c} - \frac{bd^2\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\operatorname{arctanh}(\sqrt{1-c^2x^2})}{2e}$$

output

```
-1/2*b*e*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^2+1/2*(e*x+d)^2*(a+b*arcsech(c*x))/e+b*d*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arcsin(c*x)/c-1/2*b*d^2*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh((-c^2*x^2+1)^(1/2))/e
```

**Mathematica [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.00

$$\int (d + ex) (a + b \operatorname{sech}^{-1}(cx)) dx = adx + \frac{1}{2} aex^2 + be \left( -\frac{1}{2c^2} - \frac{x}{2c} \right) \sqrt{\frac{1-cx}{1+cx}} \\ + bdx \operatorname{sech}^{-1}(cx) + \frac{1}{2} bex^2 \operatorname{sech}^{-1}(cx) \\ + \frac{2bd \sqrt{\frac{1-cx}{1+cx}} \sqrt{1-c^2x^2} \arctan \left( \frac{\sqrt{1-c^2x^2}}{1-cx} \right)}{c - c^2x}$$

input `Integrate[(d + e*x)*(a + b*ArcSech[c*x]), x]`

output `a*d*x + (a*e*x^2)/2 + b*e*(-1/2*1/c^2 - x/(2*c))*Sqrt[(1 - c*x)/(1 + c*x)] \\ + b*d*x*ArcSech[c*x] + (b*e*x^2*ArcSech[c*x])/2 + (2*b*d*Sqrt[(1 - c*x)/( \\ 1 + c*x)]*Sqrt[1 - c^2*x^2]*ArcTan[Sqrt[1 - c^2*x^2]/(1 - c*x)]/(c - c^2* \\ x)`

**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.74, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$ , Rules used = {6842, 541, 25, 27, 538, 223, 243, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex) (a + b \operatorname{sech}^{-1}(cx)) dx \\ \downarrow 6842 \\ \frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{(d+ex)^2}{x \sqrt{1-c^2x^2}} dx}{2e} + \frac{(d+ex)^2 (a + b \operatorname{sech}^{-1}(cx))}{2e} \\ \downarrow 541$$

$$\begin{aligned}
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{\int-\frac{c^2d(d+2ex)}{x\sqrt{1-c^2x^2}}dx}{c^2}-\frac{e^2\sqrt{1-c^2x^2}}{c^2}\right)}{2e} + \frac{(d+ex)^2(a+b\operatorname{sech}^{-1}(cx))}{2e} \\
& \quad \downarrow \text{25} \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{\int\frac{c^2d(d+2ex)}{x\sqrt{1-c^2x^2}}dx}{c^2}-\frac{e^2\sqrt{1-c^2x^2}}{c^2}\right)}{2e} + \frac{(d+ex)^2(a+b\operatorname{sech}^{-1}(cx))}{2e} \\
& \quad \downarrow \text{27} \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(d\int\frac{d+2ex}{x\sqrt{1-c^2x^2}}dx-\frac{e^2\sqrt{1-c^2x^2}}{c^2}\right)}{2e} + \frac{(d+ex)^2(a+b\operatorname{sech}^{-1}(cx))}{2e} \\
& \quad \downarrow \text{538} \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(d\left(d\int\frac{1}{x\sqrt{1-c^2x^2}}dx+2e\int\frac{1}{\sqrt{1-c^2x^2}}dx\right)-\frac{e^2\sqrt{1-c^2x^2}}{c^2}\right)}{2e} + \frac{(d+ex)^2(a+b\operatorname{sech}^{-1}(cx))}{2e} \\
& \quad \downarrow \text{223} \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(d\left(d\int\frac{1}{x\sqrt{1-c^2x^2}}dx+\frac{2e\arcsin(cx)}{c}\right)-\frac{e^2\sqrt{1-c^2x^2}}{c^2}\right)}{2e} + \frac{(d+ex)^2(a+b\operatorname{sech}^{-1}(cx))}{2e} \\
& \quad \downarrow \text{243} \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(d\left(\frac{1}{2}d\int\frac{1}{x^2\sqrt{1-c^2x^2}}dx^2+\frac{2e\arcsin(cx)}{c}\right)-\frac{e^2\sqrt{1-c^2x^2}}{c^2}\right)}{2e} + \frac{(d+ex)^2(a+b\operatorname{sech}^{-1}(cx))}{2e} \\
& \quad \downarrow \text{73} \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(d\left(\frac{2e\arcsin(cx)}{c}-\frac{d\int\frac{1}{c^2}-\frac{x^4}{c^2}d\sqrt{1-c^2x^2}}{c^2}\right)-\frac{e^2\sqrt{1-c^2x^2}}{c^2}\right)}{2e} + \frac{(d+ex)^2(a+b\operatorname{sech}^{-1}(cx))}{2e} \\
& \quad \downarrow \text{221}
\end{aligned}$$



$$\frac{(d + ex)^2 (a + b \operatorname{sech}^{-1}(cx))}{2e} + \frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \left( d \left( \frac{2e \arcsin(cx)}{c} - d \operatorname{arctanh}(\sqrt{1-c^2x^2}) \right) - \frac{e^2 \sqrt{1-c^2x^2}}{c^2} \right)}{2e}$$

input `Int[(d + e*x)*(a + b*ArcSech[c*x]),x]`

output `((d + e*x)^2*(a + b*ArcSech[c*x]))/(2*e) + (b*sqrt[(1 + c*x)^(-1)]*sqrt[1 + c*x]*(-(e^2*sqrt[1 - c^2*x^2])/c^2) + d*((2*e*ArcSin[c*x])/c - d*ArcTanh[sqrt[1 - c^2*x^2]])))/(2*e)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 538 `Int[((c_) + (d_.)*(x_))/((x_)*Sqrt[(a_) + (b_.)*(x_)^2]), x_Symbol] := Simp[c Int[1/(x*Sqrt[a + b*x^2]), x], x] + Simp[d Int[1/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d}, x]`

rule 541 `Int[(x_)^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[d^n*x^(m + n - 1)*((a + b*x^2)^(p + 1)/(b*(m + n + 2*p + 1))), x] + Simp[1/(b*(m + n + 2*p + 1)) Int[x^m*(a + b*x^2)^p*ExpandToSum[b*(m + n + 2*p + 1)*(c + d*x)^n - b*d^n*(m + n + 2*p + 1)*x^n - a*d^n*(m + n - 1)*x^(n - 2), x], x] /; FreeQ[{a, b, c, d, m, p}, x] && IGtQ[n, 1] && IGtQ[m, -2] && GtQ[p, -1] && IntegerQ[2*p]`

rule 6842 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*ArcSech[c*x])/(e*(m + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(e*(m + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x)^(m + 1)/(x*Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[m, -1]`

### Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.75

method	result	size
parts	$a\left(\frac{1}{2}x^2e + dx\right) + \frac{b\left(\frac{c \operatorname{arcsech}(cx)x^2e}{2} + \operatorname{arcsech}(cx)dcx + \frac{\sqrt{-\frac{cx-1}{cx}}x\sqrt{\frac{cx+1}{cx}}(2dc \arcsin(cx) - e\sqrt{-c^2x^2+1})}{2\sqrt{-c^2x^2+1}}\right)}{c}$	107
derivativedivides	$\frac{a\left(d c^2 x + \frac{1}{2} e c^2 x^2\right)}{c} + \frac{b\left(\operatorname{arcsech}(cx)d c^2 x + \frac{\operatorname{arcsech}(cx)e c^2 x^2}{2} + \frac{\sqrt{-\frac{cx-1}{cx}}cx\sqrt{\frac{cx+1}{cx}}(2dc \arcsin(cx) - e\sqrt{-c^2x^2+1})}{2\sqrt{-c^2x^2+1}}\right)}{c}$	125
default	$\frac{a\left(d c^2 x + \frac{1}{2} e c^2 x^2\right)}{c} + \frac{b\left(\operatorname{arcsech}(cx)d c^2 x + \frac{\operatorname{arcsech}(cx)e c^2 x^2}{2} + \frac{\sqrt{-\frac{cx-1}{cx}}cx\sqrt{\frac{cx+1}{cx}}(2dc \arcsin(cx) - e\sqrt{-c^2x^2+1})}{2\sqrt{-c^2x^2+1}}\right)}{c}$	125

input `int((e*x+d)*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)`

output

```
a*(1/2*x^2*e+d*x)+b/c*(1/2*c*arcsech(c*x)*x^2*e+arcsech(c*x)*d*c*x+1/2*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(2*d*c*arcsin(c*x)-e*(-c^2*x^2+1)^(1/2)))/(-c^2*x^2+1)^(1/2))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 177 vs.  $2(72) = 144$ .

Time = 0.12 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.25

$$\int (d + ex) (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{acex^2 + 2acdx - bex\sqrt{-\frac{c^2x^2-1}{c^2x^2}} - 4bd \arctan\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{cx}\right) - (2bcd + bce) \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{x}\right) + (bce - 2bcd) \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}+1}{x}\right)}{2c}$$

input

```
integrate((e*x+d)*(a+b*arcsech(c*x)),x, algorithm="fricas")
```

output

```
1/2*(a*c*e*x^2 + 2*a*c*d*x - b*e*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 4*b*d*arctan((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/(c*x)) - (2*b*c*d + b*c*e)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) + (b*c*e*x^2 + 2*b*c*d*x - 2*b*c*d - b*c*e)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)))/c
```

### Sympy [F]

$$\int (d + ex) (a + b \operatorname{sech}^{-1}(cx)) dx = \int (a + b \operatorname{asech}(cx)) (d + ex) dx$$

input

```
integrate((e*x+d)*(a+b*asech(c*x)),x)
```

output

```
Integral((a + b*asech(c*x))*(d + e*x), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.49

$$\int (d + ex) (a + b \operatorname{sech}^{-1}(cx)) dx = \frac{1}{2} aex^2 + \frac{1}{2} \left( x^2 \operatorname{arsech}(cx) - \frac{x \sqrt{\frac{1}{c^2 x^2} - 1}}{c} \right) be$$

$$+ adx + \frac{\left( cx \operatorname{arsech}(cx) - \arctan \left( \sqrt{\frac{1}{c^2 x^2} - 1} \right) \right) bd}{c}$$

input `integrate((e*x+d)*(a+b*arcsech(c*x)),x, algorithm="maxima")`output `1/2*a*e*x^2 + 1/2*(x^2*arcsech(c*x) - x*sqrt(1/(c^2*x^2) - 1)/c)*b*e + a*d*x + (c*x*arcsech(c*x) - arctan(sqrt(1/(c^2*x^2) - 1)))*b*d/c`**Giac [F]**

$$\int (d + ex) (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex + d)(b \operatorname{arsech}(cx) + a) dx$$

input `integrate((e*x+d)*(a+b*arcsech(c*x)),x, algorithm="giac")`output `integrate((e*x + d)*(b*arcsech(c*x) + a), x)`**Mupad [B] (verification not implemented)**

Time = 3.83 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.70

$$\int (d + ex) (a + b \operatorname{sech}^{-1}(cx)) dx = \frac{ax(2d + ex)}{2} + \frac{bd \operatorname{atan} \left( \frac{1}{\sqrt{\frac{1}{cx} - 1} \sqrt{\frac{1}{cx} + 1}} \right)}{c}$$

$$+ \frac{bex^2 \operatorname{acosh} \left( \frac{1}{cx} \right)}{2} + bdx \operatorname{acosh} \left( \frac{1}{cx} \right)$$

$$- \frac{bex \sqrt{\frac{1}{cx} - 1} \sqrt{\frac{1}{cx} + 1}}{2c}$$

input `int((a + b*acosh(1/(c*x)))*(d + e*x),x)`

output `(a*x*(2*d + e*x))/2 + (b*d*atan(1/((1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2))))/c + (b*e*x^2*acosh(1/(c*x)))/2 + b*d*x*acosh(1/(c*x)) - (b*e*x*(1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2)))/(2*c)`

### Reduce [F]

$$\int (d + ex) (a + b \operatorname{sech}^{-1}(cx)) dx = \left( \int \operatorname{asech}(cx) dx \right) bd + \left( \int \operatorname{asech}(cx) x dx \right) be + adx + \frac{ae x^2}{2}$$

input `int((e*x+d)*(a+b*asech(c*x)),x)`

output `(2*int(asech(c*x),x)*b*d + 2*int(asech(c*x)*x,x)*b*e + 2*a*d*x + a*e*x**2)/2`

### 3.77 $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{d+ex} dx$

Optimal result	621
Mathematica [C] (warning: unable to verify)	622
Rubi [A] (verified)	622
Maple [C] (verified)	624
Fricas [F]	625
Sympy [F]	625
Maxima [F]	626
Giac [F]	626
Mupad [F(-1)]	626
Reduce [F]	627

#### Optimal result

Integrand size = 16, antiderivative size = 211

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{d + ex} dx = \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{cde^{\operatorname{sech}^{-1}(cx)}}{e - \sqrt{-c^2d^2 + e^2}}\right)}{e} + \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{cde^{\operatorname{sech}^{-1}(cx)}}{e + \sqrt{-c^2d^2 + e^2}}\right)}{e} - \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + e^{2\operatorname{sech}^{-1}(cx)}\right)}{e} + \frac{b \operatorname{PolyLog}\left(2, -\frac{cde^{\operatorname{sech}^{-1}(cx)}}{e - \sqrt{-c^2d^2 + e^2}}\right)}{e} + \frac{b \operatorname{PolyLog}\left(2, -\frac{cde^{\operatorname{sech}^{-1}(cx)}}{e + \sqrt{-c^2d^2 + e^2}}\right)}{e} - \frac{b \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(cx)}\right)}{2e}$$

output

```
(a+b*arcsech(c*x))*ln(1+c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e-(-c^2*d^2+e^2)^(1/2)))/e+(a+b*arcsech(c*x))*ln(1+c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e+(-c^2*d^2+e^2)^(1/2)))/e-(a+b*arcsech(c*x))*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)/e+b*polylog(2,-c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e-(-c^2*d^2+e^2)^(1/2)))/e+b*polylog(2,-c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e+(-c^2*d^2+e^2)^(1/2)))/e-1/2*b*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)/e
```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 0.39 (sec) , antiderivative size = 393, normalized size of antiderivative = 1.86

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex} dx = \frac{a \log(d + ex)}{e} + \frac{b \left( \operatorname{PolyLog} \left( 2, -e^{-2 \operatorname{sech}^{-1}(cx)} \right) - 2 \left( -4i \arcsin \left( \frac{\sqrt{1 + \frac{e}{cd}}}{\sqrt{2}} \right) \operatorname{arctanh} \left( \frac{(-cd+e) \tanh \left( \frac{1}{2} \operatorname{sech}^{-1}(cx) \right)}{\sqrt{-c^2 d^2 + e^2}} \right) \right) \right)}{2e} + \operatorname{sech}^{-1}(cx)$$

input `Integrate[(a + b*ArcSech[c*x])/(d + e*x), x]`

output

```
(a*Log[d + e*x])/e + (b*(PolyLog[2, -E^(-2*ArcSech[c*x])] - 2*((-4*I)*ArcSin[Sqrt[1 + e/(c*d)]/Sqrt[2]]*ArcTanh[(-c*d + e)*Tanh[ArcSech[c*x]/2]]/Sqrt[-(c^2*d^2) + e^2]] + ArcSech[c*x]*Log[1 + E^(-2*ArcSech[c*x])] - ArcSech[c*x]*Log[1 + (e - Sqrt[-(c^2*d^2) + e^2])/(c*d*E^ArcSech[c*x])] + (2*I)*ArcSin[Sqrt[1 + e/(c*d)]/Sqrt[2]]*Log[1 + (e - Sqrt[-(c^2*d^2) + e^2])/(c*d*E^ArcSech[c*x])] - ArcSech[c*x]*Log[1 + (e + Sqrt[-(c^2*d^2) + e^2])/(c*d*E^ArcSech[c*x])] - (2*I)*ArcSin[Sqrt[1 + e/(c*d)]/Sqrt[2]]*Log[1 + (e + Sqrt[-(c^2*d^2) + e^2])/(c*d*E^ArcSech[c*x])] + PolyLog[2, (-e + Sqrt[-(c^2*d^2) + e^2])/(c*d*E^ArcSech[c*x])] + PolyLog[2, -((e + Sqrt[-(c^2*d^2) + e^2])/(c*d*E^ArcSech[c*x]))])))/(2*e)
```

**Rubi [A] (verified)**

Time = 1.37 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.09, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$ , Rules used = {6841, 2998}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex} dx$$

↓ 6841

$$\begin{aligned}
 & \frac{b \int \frac{\sqrt{\frac{1-cx}{cx+1}} \log\left(\frac{e^{-\operatorname{sech}^{-1}(cx)}(e-\sqrt{e^2-c^2d^2})}{cd} + 1\right)}{x(1-cx)} dx}{e} + \frac{b \int \frac{\sqrt{\frac{1-cx}{cx+1}} \log\left(\frac{e^{-\operatorname{sech}^{-1}(cx)}(e+\sqrt{e^2-c^2d^2})}{cd} + 1\right)}{x(1-cx)} dx}{e} \\
 & \frac{b \int \frac{\sqrt{\frac{1-cx}{cx+1}} \log(1+e^{-2\operatorname{sech}^{-1}(cx)})}{x(1-cx)} dx}{e} + \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(\frac{e^{-\operatorname{sech}^{-1}(cx)}(e-\sqrt{e^2-c^2d^2})}{cd} + 1\right)}{e} + \\
 & \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(\frac{(\sqrt{e^2-c^2d^2}+e)e^{-\operatorname{sech}^{-1}(cx)}}{cd} + 1\right)}{e} - \\
 & \frac{\log\left(e^{-2\operatorname{sech}^{-1}(cx)} + 1\right) (a + b\operatorname{sech}^{-1}(cx))}{e} \\
 & \quad \downarrow \text{2998} \\
 & \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(\frac{(e-\sqrt{e^2-c^2d^2})e^{-\operatorname{sech}^{-1}(cx)}}{cd} + 1\right)}{e} + \\
 & \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(\frac{(\sqrt{e^2-c^2d^2}+e)e^{-\operatorname{sech}^{-1}(cx)}}{cd} + 1\right)}{e} - \\
 & \frac{\log\left(e^{-2\operatorname{sech}^{-1}(cx)} + 1\right) (a + b\operatorname{sech}^{-1}(cx))}{e} - \frac{b \operatorname{PolyLog}\left(2, -\frac{(e-\sqrt{e^2-c^2d^2})e^{-\operatorname{sech}^{-1}(cx)}}{cd}\right)}{e} \\
 & \frac{b \operatorname{PolyLog}\left(2, -\frac{(e+\sqrt{e^2-c^2d^2})e^{-\operatorname{sech}^{-1}(cx)}}{cd}\right)}{e} + \frac{b \operatorname{PolyLog}\left(2, -e^{-2\operatorname{sech}^{-1}(cx)}\right)}{2e}
 \end{aligned}$$

input `Int[(a + b*ArcSech[c*x])/(d + e*x),x]`

output `-(((a + b*ArcSech[c*x])*Log[1 + E^(-2*ArcSech[c*x])])/e) + ((a + b*ArcSech[c*x])*Log[1 + (e - Sqrt[-(c^2*d^2) + e^2])/(c*d*E^ArcSech[c*x])])/e + ((a + b*ArcSech[c*x])*Log[1 + (e + Sqrt[-(c^2*d^2) + e^2])/(c*d*E^ArcSech[c*x])])/e + (b*PolyLog[2, -E^(-2*ArcSech[c*x])])/(2*e) - (b*PolyLog[2, -((e - Sqrt[-(c^2*d^2) + e^2])/(c*d*E^ArcSech[c*x])]))/e - (b*PolyLog[2, -((e + Sqrt[-(c^2*d^2) + e^2])/(c*d*E^ArcSech[c*x])]))/e`



Defintions of rubi rules used

```
rule 2998 Int[Log[v_]*(u_), x_Symbol] := With[{w = DerivativeDivides[v, u*(1 - v), x]
}, Simp[w*PolyLog[2, 1 - v], x] /; !FalseQ[w]]
```

```
rule 6841 Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))/((d_.) + (e_.)*(x_.)), x_Symbol] :=
Simp[(a + b*ArcSech[c*x])*(Log[1 + (e - Sqrt[(-c^2)*d^2 + e^2])/(c*d*E^ArcS
ech[c*x]])]/e), x] + (Simp[(a + b*ArcSech[c*x])*(Log[1 + (e + Sqrt[(-c^2)*d^
2 + e^2])/(c*d*E^ArcSech[c*x]])]/e), x] - Simp[(a + b*ArcSech[c*x])*(Log[1 +
1/E^(2*ArcSech[c*x]])]/e), x] + Simp[b/e Int[(Sqrt[(1 - c*x)/(1 + c*x)]*L
og[1 + (e - Sqrt[(-c^2)*d^2 + e^2])/(c*d*E^ArcSech[c*x]])]/(x*(1 - c*x)), x
], x] + Simp[b/e Int[(Sqrt[(1 - c*x)/(1 + c*x)]*Log[1 + (e + Sqrt[(-c^2)*
d^2 + e^2])/(c*d*E^ArcSech[c*x]])]/(x*(1 - c*x)), x], x] - Simp[b/e Int[(
Sqrt[(1 - c*x)/(1 + c*x)]*Log[1 + 1/E^(2*ArcSech[c*x]])]/(x*(1 - c*x)), x],
x]) /; FreeQ[{a, b, c, d, e}, x]
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.81 (sec) , antiderivative size = 511, normalized size of antiderivative = 2.42

method	result
parts	$\frac{a \ln(ex+d)}{e} + \frac{b \operatorname{arcsech}(cx) \ln\left(\frac{-cd\left(\frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}\right) + \sqrt{-c^2d^2 + e^2} - e}{-e + \sqrt{-c^2d^2 + e^2}}\right)}{e} + \frac{b \operatorname{arcsech}(cx) \ln\left(\frac{cd\left(\frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}\right) + \sqrt{-c^2d^2 + e^2} - e}{e + \sqrt{-c^2d^2 + e^2}}\right)}{e}$
derivativedivides	$\frac{ac \ln(cex+cd)}{e} + bc \left( \frac{\operatorname{arcsech}(cx) \ln\left(\frac{-cd\left(\frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}\right) + \sqrt{-c^2d^2 + e^2} - e}{-e + \sqrt{-c^2d^2 + e^2}}\right)}{e} + \frac{\operatorname{arcsech}(cx) \ln\left(\frac{cd\left(\frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}\right) + \sqrt{-c^2d^2 + e^2} - e}{e + \sqrt{-c^2d^2 + e^2}}\right)}{e} \right)$
default	$\frac{ac \ln(cex+cd)}{e} + bc \left( \frac{\operatorname{arcsech}(cx) \ln\left(\frac{-cd\left(\frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}\right) + \sqrt{-c^2d^2 + e^2} - e}{-e + \sqrt{-c^2d^2 + e^2}}\right)}{e} + \frac{\operatorname{arcsech}(cx) \ln\left(\frac{cd\left(\frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}\right) + \sqrt{-c^2d^2 + e^2} - e}{e + \sqrt{-c^2d^2 + e^2}}\right)}{e} \right)$

```
input int((a+b*arcsech(c*x))/(e*x+d), x, method=_RETURNVERBOSE)
```

output

```
a*ln(e*x+d)/e+b/e*arcsech(c*x)*ln((-c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))+(-c^2*d^2+e^2)^(1/2)-e)/(-e+(-c^2*d^2+e^2)^(1/2)))+b/e*arcsech(c*x)*ln((c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))+(-c^2*d^2+e^2)^(1/2)+e)/(e+(-c^2*d^2+e^2)^(1/2)))+b/e*dilog((-c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))+(-c^2*d^2+e^2)^(1/2)-e)/(-e+(-c^2*d^2+e^2)^(1/2)))+b/e*dilog((c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))+(-c^2*d^2+e^2)^(1/2)+e)/(e+(-c^2*d^2+e^2)^(1/2)))-b/e*arcsech(c*x)*ln(1+I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))-b/e*arcsech(c*x)*ln(1-I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))-b/e*dilog(1+I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))-b/e*dilog(1-I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))
```

**Fricas [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex} dx = \int \frac{b \operatorname{arsech}(cx) + a}{ex + d} dx$$

input

```
integrate((a+b*arcsech(c*x))/(e*x+d),x, algorithm="fricas")
```

output

```
integral((b*arcsech(c*x) + a)/(e*x + d), x)
```

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex} dx = \int \frac{a + b \operatorname{asech}(cx)}{d + ex} dx$$

input

```
integrate((a+b*asech(c*x))/(e*x+d),x)
```

output

```
Integral((a + b*asech(c*x))/(d + e*x), x)
```

**Maxima [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex} dx = \int \frac{b \operatorname{arsech}(cx) + a}{ex + d} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x+d),x, algorithm="maxima")`

output `b*integrate(log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(e*x + d), x) + a*log(e*x + d)/e`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex} dx = \int \frac{b \operatorname{arsech}(cx) + a}{ex + d} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x+d),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/(e*x + d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{d + ex} dx$$

input `int((a + b*acosh(1/(c*x)))/(d + e*x),x)`

output `int((a + b*acosh(1/(c*x)))/(d + e*x), x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex} dx = \frac{\left( \int \frac{a \operatorname{sech}(cx)}{ex+d} dx \right) be + \log(ex + d) a}{e}$$

input `int((a+b*asech(c*x))/(e*x+d),x)`

output `(int(asech(c*x)/(d + e*x),x)*b*e + log(d + e*x)*a)/e`

### 3.78 $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex)^2} dx$

Optimal result	628
Mathematica [A] (verified)	629
Rubi [A] (verified)	629
Maple [A] (verified)	631
Fricas [B] (verification not implemented)	631
Sympy [F]	633
Maxima [F]	633
Giac [F]	633
Mupad [F(-1)]	634
Reduce [F]	634

#### Optimal result

Integrand size = 16, antiderivative size = 147

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{(d + ex)^2} dx = -\frac{a + b\operatorname{sech}^{-1}(cx)}{e(d + ex)} + \frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1 + cx} \arctan\left(\frac{e+c^2dx}{\sqrt{c^2d^2-e^2}\sqrt{1-c^2x^2}}\right)}{d\sqrt{c^2d^2 - e^2}} + \frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1 + cx}\operatorname{arctanh}(\sqrt{1 - c^2x^2})}{de}$$

output

```
-(a+b*arcsech(c*x))/e/(e*x+d)+b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctan((c^2*d*x+e)/(c^2*d^2-e^2)^(1/2)/(-c^2*x^2+1)^(1/2))/d/(c^2*d^2-e^2)^(1/2)+b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh((-c^2*x^2+1)^(1/2))/d/e
```

**Mathematica [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.51

$$\begin{aligned} & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^2} dx \\ &= -\frac{a}{e(d + ex)} - \frac{b \operatorname{sech}^{-1}(cx)}{e(d + ex)} - \frac{b \log(x)}{de} \\ &+ \frac{b \log(d + ex)}{d\sqrt{-c^2d^2 + e^2}} + \frac{b \log\left(1 + \sqrt{\frac{1-cx}{1+cx}} + cx\sqrt{\frac{1-cx}{1+cx}}\right)}{de} \\ &- \frac{b \log\left(e + c^2dx + \sqrt{-c^2d^2 + e^2}\sqrt{\frac{1-cx}{1+cx}} + c\sqrt{-c^2d^2 + e^2}x\sqrt{\frac{1-cx}{1+cx}}\right)}{d\sqrt{-c^2d^2 + e^2}} \end{aligned}$$

input

```
Integrate[(a + b*ArcSech[c*x])/(d + e*x)^2,x]
```

output

```
-(a/(e*(d + e*x))) - (b*ArcSech[c*x])/(e*(d + e*x)) - (b*Log[x])/(d*e) + (
b*Log[d + e*x])/(d*Sqrt[-(c^2*d^2) + e^2]) + (b*Log[1 + Sqrt[(1 - c*x)/(1
+ c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]])/(d*e) - (b*Log[e + c^2*d*x + Sqr
t[-(c^2*d^2) + e^2]*Sqrt[(1 - c*x)/(1 + c*x)] + c*Sqrt[-(c^2*d^2) + e^2]*x
*Sqrt[(1 - c*x)/(1 + c*x)]])/(d*Sqrt[-(c^2*d^2) + e^2])
```

**Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.90, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$ , Rules used = {6842, 617, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^2} dx \\ & \quad \downarrow \text{6842} \\ & -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{x(d+ex)\sqrt{1-c^2x^2}} dx}{e} - \frac{a + b \operatorname{sech}^{-1}(cx)}{e(d + ex)} \end{aligned}$$

$$\begin{array}{c}
 \downarrow 617 \\
 \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \left( \frac{1}{dx\sqrt{1-c^2x^2}} - \frac{e}{d(d+ex)\sqrt{1-c^2x^2}} \right) dx}{e} - \frac{a + b\operatorname{sech}^{-1}(cx)}{e(d+ex)} \\
 \downarrow 2009 \\
 \frac{a + b\operatorname{sech}^{-1}(cx)}{e(d+ex)} - \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{e \arctan\left(\frac{c^2 dx + e}{\sqrt{1-c^2x^2}\sqrt{c^2 d^2 - e^2}}\right)}{d\sqrt{c^2 d^2 - e^2}} - \frac{\operatorname{arctanh}\left(\sqrt{1-c^2x^2}\right)}{d} \right)}{e}
 \end{array}$$

input `Int[(a + b*ArcSech[c*x])/(d + e*x)^2,x]`

output `-((a + b*ArcSech[c*x])/(e*(d + e*x))) - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-(e*ArcTan[(e + c^2*d*x)/(Sqrt[c^2*d^2 - e^2]*Sqrt[1 - c^2*x^2])]))/(d*Sqrt[c^2*d^2 - e^2])) - ArcTanh[Sqrt[1 - c^2*x^2]/d])/e`

### Defintions of rubi rules used

rule 617 `Int[(x_)^(m_.)*((c_) + (d_.)*(x_))^(n_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^2)^p, x^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, p}, x] && ILtQ[n, 0] && IntegerQ[m] && IntegerQ[2*p]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6842 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*ArcSech[c*x])/(e*(m + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(e*(m + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x)^(m + 1)/(x*Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[m, -1]`

### Maple [A] (verified)

Time = 1.78 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.41

method	result
parts	$-\frac{a}{(ex+d)e} + \frac{b \left( -\frac{c^2 \operatorname{arcsech}(cx)}{(cex+cd)e} + \frac{c^2 \sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}} \left( \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right) \sqrt{-\frac{c^2d^2-e^2}{e^2}} - \ln\left(\frac{2\sqrt{-c^2x^2+1} \sqrt{-\frac{c^2d^2-e^2}{e^2}}}{cex+cd}\right) \right)}{e \sqrt{-\frac{c^2d^2-e^2}{e^2}} d \sqrt{-c^2x^2+1}} \right)}{c}$
derivativedivides	$-\frac{a e^2}{(cex+cd)e} + b c^2 \left( -\frac{\operatorname{arcsech}(cx)}{(cex+cd)e} + \frac{\sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}} \left( \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right) \sqrt{-\frac{c^2d^2-e^2}{e^2}} - \ln\left(\frac{2\sqrt{-c^2x^2+1} \sqrt{-\frac{c^2d^2-e^2}{e^2}}}{cex+cd}\right) \right)}{e \sqrt{-\frac{c^2d^2-e^2}{e^2}} d \sqrt{-c^2x^2+1}} \right)$
default	$-\frac{a e^2}{(cex+cd)e} + b c^2 \left( -\frac{\operatorname{arcsech}(cx)}{(cex+cd)e} + \frac{\sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}} \left( \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right) \sqrt{-\frac{c^2d^2-e^2}{e^2}} - \ln\left(\frac{2\sqrt{-c^2x^2+1} \sqrt{-\frac{c^2d^2-e^2}{e^2}}}{cex+cd}\right) \right)}{e \sqrt{-\frac{c^2d^2-e^2}{e^2}} d \sqrt{-c^2x^2+1}} \right)$

```
input int((a+b*arcsech(c*x))/(e*x+d)^2,x,method=_RETURNVERBOSE)
```

```
output -a/(e*x+d)/e+b/c*(-c^2/(c*e*x+c*d)/e*arcsech(c*x)+c^2/e*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(arctanh(1/(-c^2*x^2+1)^(1/2))*(-(c^2*d^2-e^2)/e^2)^(1/2)-ln(2*((-c^2*x^2+1)^(1/2))*(-(c^2*d^2-e^2)/e^2)^(1/2)*e+d*c^2*x+e)/(c*e*x+c*d)))/(-c^2*d^2-e^2)/e^2)^(1/2)/d/(-c^2*x^2+1)^(1/2))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 279 vs. 2(99) = 198.



Time = 0.15 (sec) , antiderivative size = 578, normalized size of antiderivative = 3.93

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^2} dx$$

$$= \frac{\left[ \begin{aligned} & ac^2d^3 - ade^2 + \sqrt{-c^2d^2 + e^2}(be^2x + bde) \log \left( \frac{c^2dex - (c^3d^2 - ce^2)x \sqrt{-\frac{c^2x^2-1}{c^2x^2}} + e^2 - \sqrt{-c^2d^2 + e^2} \left( c^2dx + cex \sqrt{-\frac{c^2x^2-1}{c^2x^2}} \right)}{ex+d} \right)}{c^2d^4e - d^2e^3 +} \right. \\ & \left. ac^2d^3 - ade^2 - 2\sqrt{c^2d^2 - e^2}(be^2x + bde) \arctan \left( -\frac{\sqrt{c^2d^2 - e^2}cdx \sqrt{-\frac{c^2x^2-1}{c^2x^2}} - \sqrt{c^2d^2 - e^2}(ex+d)}{(c^2d^2 - e^2)x} \right) + (bc^2d^3 - b} \right]}{c^2d^4e - d^2e^3 + (c^2d^3e^2 - d^2e^4)} \end{aligned} \right.$$

input `integrate((a+b*arcsech(c*x))/(e*x+d)^2,x, algorithm="fricas")`

output

```
[-(a*c^2*d^3 - a*d*e^2 + sqrt(-c^2*d^2 + e^2)*(b*e^2*x + b*d*e)*log((c^2*d
*e*x - (c^3*d^2 - c*e^2)*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e^2 - sqrt(-c^
2*d^2 + e^2)*(c^2*d*x + c*e*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e))/(e*x +
d)) + (b*c^2*d^3 - b*d*e^2 + (b*c^2*d^2*e - b*e^3)*x)*log((c*x*sqrt(-(c^2*
x^2 - 1)/(c^2*x^2)) - 1)/x) + (b*c^2*d^3 - b*d*e^2)*log((c*x*sqrt(-(c^2*x^
2 - 1)/(c^2*x^2)) + 1)/(c*x)))/(c^2*d^4*e - d^2*e^3 + (c^2*d^3*e^2 - d*e^4
)*x), -(a*c^2*d^3 - a*d*e^2 - 2*sqrt(c^2*d^2 - e^2)*(b*e^2*x + b*d*e)*arct
an(-(sqrt(c^2*d^2 - e^2)*c*d*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - sqrt(c^2*d
^2 - e^2)*(e*x + d))/((c^2*d^2 - e^2)*x)) + (b*c^2*d^3 - b*d*e^2 + (b*c^2*
d^2*e - b*e^3)*x)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) + (b*c^2
*d^3 - b*d*e^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)))/(c^2*
d^4*e - d^2*e^3 + (c^2*d^3*e^2 - d*e^4)*x]
```

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^2} dx = \int \frac{a + b \operatorname{arsech}(cx)}{(d + ex)^2} dx$$

input `integrate((a+b*asech(c*x))/(e*x+d)**2,x)`

output `Integral((a + b*asech(c*x))/(d + e*x)**2, x)`

**Maxima [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^2} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex + d)^2} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x+d)^2,x, algorithm="maxima")`

output `(c^2*integrate(x^2/(c^2*d^2*x^2 + (c^2*d^2*x^2 - d^2 + (c^2*d*e*x^2 - d*e)*x)*sqrt(c*x + 1)*sqrt(-c*x + 1) - d^2 + (c^2*d*e*x^2 - d*e)*x), x) + (x*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1) - x*log(c) - x*log(x))/(d*e*x + d^2) - integrate(1/(c^2*d^2*x^2 - d^2 + (c^2*d*e*x^2 - d*e)*x), x))*b - a/(e^2*x + d*e)`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^2} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex + d)^2} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x+d)^2,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/(e*x + d)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^2} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{(d + ex)^2} dx$$

input `int((a + b*acosh(1/(c*x)))/(d + e*x)^2,x)`output `int((a + b*acosh(1/(c*x)))/(d + e*x)^2, x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^2} dx = \frac{\left(\int \frac{\operatorname{asech}(cx)}{e^2 x^2 + 2dex + d^2} dx\right) b d^2 + \left(\int \frac{\operatorname{asech}(cx)}{e^2 x^2 + 2dex + d^2} dx\right) b dex + ax}{d(ex + d)}$$

input `int((a+b*asech(c*x))/(e*x+d)^2,x)`output `(int(asech(c*x)/(d**2 + 2*d*e*x + e**2*x**2),x)*b*d**2 + int(asech(c*x)/(d**2 + 2*d*e*x + e**2*x**2),x)*b*d*e*x + a*x)/(d*(d + e*x))`

**3.79** 
$$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex)^3} dx$$

Optimal result	635
Mathematica [C] (verified)	636
Rubi [A] (verified)	636
Maple [B] (verified)	638
Fricas [B] (verification not implemented)	639
Sympy [F]	640
Maxima [F(-2)]	641
Giac [F]	641
Mupad [F(-1)]	641
Reduce [F]	642

**Optimal result**

Integrand size = 16, antiderivative size = 234

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{(d + ex)^3} dx = \frac{be\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{2d(c^2d^2 - e^2)(d + ex)} - \frac{a + b\operatorname{sech}^{-1}(cx)}{2e(d + ex)^2} + \frac{b(2c^2d^2 - e^2)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\arctan\left(\frac{e+c^2dx}{\sqrt{c^2d^2 - e^2}\sqrt{1-c^2x^2}}\right)}{2d^2(c^2d^2 - e^2)^{3/2}} + \frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\operatorname{arctanh}(\sqrt{1-c^2x^2})}{2d^2e}$$

output

```
1/2*b*e*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/d/(c^2*d^2-e^2)
/(e*x+d)-1/2*(a+b*arcsech(c*x))/e/(e*x+d)^2+1/2*b*(2*c^2*d^2-e^2)*(1/(c*x+
1))^(1/2)*(c*x+1)^(1/2)*arctan((c^2*d*x+e)/(c^2*d^2-e^2)^(1/2)/(-c^2*x^2+1
)^(1/2))/d^2/(c^2*d^2-e^2)^(3/2)+1/2*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arc
tanh((-c^2*x^2+1)^(1/2))/d^2/e
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.40 (sec) , antiderivative size = 342, normalized size of antiderivative = 1.46

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^3} dx$$

$$= \frac{1}{2} \left( -\frac{a}{e(d + ex)^2} + \frac{b \sqrt{\frac{1-cx}{1+cx}} (e + cex)}{d(cd - e)(cd + e)(d + ex)} - \frac{b \operatorname{sech}^{-1}(cx)}{e(d + ex)^2} - \frac{b \log(x)}{d^2 e} \right. \\ \left. + \frac{b \log \left( 1 + \sqrt{\frac{1-cx}{1+cx}} + cx \sqrt{\frac{1-cx}{1+cx}} \right)}{d^2 e} \right. \\ \left. - \frac{ib(2c^2 d^2 - e^2) \log \left( \frac{4d^2 e \sqrt{c^2 d^2 - e^2} (ie + ic^2 dx + \sqrt{c^2 d^2 - e^2} \sqrt{\frac{1-cx}{1+cx}} + c \sqrt{c^2 d^2 - e^2} x \sqrt{\frac{1-cx}{1+cx}})}{b(2c^2 d^2 - e^2)(d + ex)} \right)}{d^2 (cd - e)(cd + e) \sqrt{c^2 d^2 - e^2}} \right)$$

input

```
Integrate[(a + b*ArcSech[c*x])/(d + e*x)^3,x]
```

output

```
(-(a/(e*(d + e*x)^2)) + (b*Sqrt[(1 - c*x)/(1 + c*x)]*(e + c*e*x))/(d*(c*d - e)*(c*d + e)*(d + e*x)) - (b*ArcSech[c*x])/(e*(d + e*x)^2) - (b*Log[x])/
(d^2*e) + (b*Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c
*x)]])/(d^2*e) - (I*b*(2*c^2*d^2 - e^2)*Log[(4*d^2*e*Sqrt[c^2*d^2 - e^2]*(
I*e + I*c^2*d*x + Sqrt[c^2*d^2 - e^2]*Sqrt[(1 - c*x)/(1 + c*x)] + c*Sqrt[c
^2*d^2 - e^2]*x*Sqrt[(1 - c*x)/(1 + c*x)])]/(b*(2*c^2*d^2 - e^2)*(d + e*x
)))/(d^2*(c*d - e)*(c*d + e)*Sqrt[c^2*d^2 - e^2])/2
```

**Rubi [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.04, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$ , Rules used = {6842, 617, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^3} dx \\
& \quad \downarrow \text{6842} \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{x(d+ex)^2\sqrt{1-c^2x^2}} dx}{2e} - \frac{a + b \operatorname{sech}^{-1}(cx)}{2e(d + ex)^2} \\
& \quad \downarrow \text{617} \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \left( -\frac{e}{d^2(d+ex)\sqrt{1-c^2x^2}} - \frac{e}{d(d+ex)^2\sqrt{1-c^2x^2}} + \frac{1}{d^2x\sqrt{1-c^2x^2}} \right) dx}{2e} - \frac{a + b \operatorname{sech}^{-1}(cx)}{2e(d + ex)^2} \\
& \quad \downarrow \text{2009} \\
& -\frac{a + b \operatorname{sech}^{-1}(cx)}{2e(d + ex)^2} - \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{c^2e \arctan\left(\frac{c^2dx+e}{\sqrt{1-c^2x^2}\sqrt{c^2d^2-e^2}}\right)}{(c^2d^2-e^2)^{3/2}} - \frac{e \arctan\left(\frac{c^2dx+e}{d^2\sqrt{c^2d^2-e^2}}\right)}{d^2\sqrt{c^2d^2-e^2}} - \frac{\operatorname{arctanh}\left(\sqrt{1-c^2x^2}\right)}{d^2} - \frac{e^2\sqrt{1-c^2x^2}}{d(c^2d^2-e^2)(d+ex)} \right)}{2e}
\end{aligned}$$

input `Int[(a + b*ArcSech[c*x])/(d + e*x)^3,x]`

output `-1/2*(a + b*ArcSech[c*x])/(e*(d + e*x)^2) - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-(e^2*Sqrt[1 - c^2*x^2])/(d*(c^2*d^2 - e^2)*(d + e*x))) - (c^2*e*ArcTan[(e + c^2*d*x)/(Sqrt[c^2*d^2 - e^2]*Sqrt[1 - c^2*x^2])])/(c^2*d^2 - e^2)^(3/2) - (e*ArcTan[(e + c^2*d*x)/(Sqrt[c^2*d^2 - e^2]*Sqrt[1 - c^2*x^2])])/(d^2*Sqrt[c^2*d^2 - e^2]) - ArcTanh[Sqrt[1 - c^2*x^2]/d^2])/(2*e)`

### Defintions of rubi rules used

rule 617 `Int[(x_)^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^2)^p, x^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, p}, x] && ILtQ[n, 0] && IntegerQ[m] && IntegerQ[2*p]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6842

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))*((d_.) + (e_.)*(x_.))^(m_.), x_Symbol]
  := Simp[(d + e*x)^(m + 1)*((a + b*ArcSech[c*x])/(e*(m + 1))), x] + Simp[
  b*(Sqrt[1 + c*x]/(e*(m + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x)^(m + 1)/(x*
  Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[m, -1]
```

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 592 vs. 2(204) = 408.

Time = 1.97 (sec) , antiderivative size = 593, normalized size of antiderivative = 2.53

method	result
parts	$-\frac{a}{2(ce x+d)^2 e} + b \left( -\frac{c^3 \operatorname{arcsech}(cx)}{2(ce x+cd)^2 e} + \frac{c^2 \sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}}}{\left( \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2 x^2+1}}\right) c^3 d^3 \sqrt{-\frac{c^2 d^2-e^2}{e^2}} + \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2 x^2+1}}\right) \right)} \right)$
derivativedivides	$-\frac{a c^3}{2(ce x+cd)^2 e} + b c^3 \left( -\frac{\operatorname{arcsech}(cx)}{2(ce x+cd)^2 e} + \frac{\sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}}}{\left( \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2 x^2+1}}\right) c^3 d^3 \sqrt{-\frac{c^2 d^2-e^2}{e^2}} + \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2 x^2+1}}\right) \right)} \right)$
default	$-\frac{a c^3}{2(ce x+cd)^2 e} + b c^3 \left( -\frac{\operatorname{arcsech}(cx)}{2(ce x+cd)^2 e} + \frac{\sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}}}{\left( \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2 x^2+1}}\right) c^3 d^3 \sqrt{-\frac{c^2 d^2-e^2}{e^2}} + \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2 x^2+1}}\right) \right)} \right)$

input

```
int((a+b*arcsech(c*x))/(e*x+d)^3,x,method=_RETURNVERBOSE)
```

output

```

-1/2*a/(e*x+d)^2/e+b/c*(-1/2*c^3/(c*e*x+c*d)^2/e*arcsech(c*x)+1/2*c^2/e*(-
(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(arctanh(1/(-c^2*x^2+1)^(1/2))*c^
3*d^3*(-(c^2*d^2-e^2)/e^2)^(1/2)+arctanh(1/(-c^2*x^2+1)^(1/2))*c^3*d^2*e*(
-(c^2*d^2-e^2)/e^2)^(1/2)*x-2*ln(2*((-c^2*x^2+1)^(1/2)*(-(c^2*d^2-e^2)/e^2
)^(1/2)*e+d*c^2*x+e)/(c*e*x+c*d))*c^3*d^3-2*ln(2*((-c^2*x^2+1)^(1/2)*(-(c^
2*d^2-e^2)/e^2)^(1/2)*e+d*c^2*x+e)/(c*e*x+c*d))*c^3*d^2*e*x-arctanh(1/(-c^
2*x^2+1)^(1/2))*c*d*e^2*(-(c^2*d^2-e^2)/e^2)^(1/2)-arctanh(1/(-c^2*x^2+1)^(
1/2))*e^3*(-(c^2*d^2-e^2)/e^2)^(1/2)*c*x+c*d*e^2*(-(c^2*d^2-e^2)/e^2)^(1/
2)*(-c^2*x^2+1)^(1/2)+ln(2*((-c^2*x^2+1)^(1/2)*(-(c^2*d^2-e^2)/e^2)^(1/2)*
e+d*c^2*x+e)/(c*e*x+c*d))*c*d*e^2+ln(2*((-c^2*x^2+1)^(1/2)*(-(c^2*d^2-e^2)
/e^2)^(1/2)*e+d*c^2*x+e)/(c*e*x+c*d))*e^3*c*x)/(-c^2*x^2+1)^(1/2)/(c*d-e)/
(c*d+e)/d^2/(c*e*x+c*d)/(-c^2*d^2-e^2)/e^2)^(1/2))

```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 596 vs.  $2(156) = 312$ .

Time = 0.22 (sec) , antiderivative size = 1212, normalized size of antiderivative = 5.18

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^3} dx = \text{Too large to display}$$

input

```
integrate((a+b*arcsech(c*x))/(e*x+d)^3,x, algorithm="fricas")
```



output

```

[-1/2*(a*c^4*d^6 - (2*a + b)*c^2*d^4*e^2 + (a + b)*d^2*e^4 - (b*c^2*d^2*e^4 - b*e^6)*x^2 + (2*b*c^2*d^4*e - b*d^2*e^3 + (2*b*c^2*d^2*e^3 - b*e^5)*x^2 + 2*(2*b*c^2*d^3*e^2 - b*d*e^4)*x)*sqrt(-c^2*d^2 + e^2)*log((c^2*d*e*x - (c^3*d^2 - c*e^2)*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e^2 - sqrt(-c^2*d^2 + e^2)*(c^2*d*x + c*e*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e))/(e*x + d)) - 2*(b*c^2*d^3*e^3 - b*d*e^5)*x + (b*c^4*d^6 - 2*b*c^2*d^4*e^2 + b*d^2*e^4 + (b*c^4*d^4*e^2 - 2*b*c^2*d^2*e^4 + b*e^6)*x^2 + 2*(b*c^4*d^5*e - 2*b*c^2*d^3*e^3 + b*d*e^5)*x)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) + (b*c^4*d^6 - 2*b*c^2*d^4*e^2 + b*d^2*e^4)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - ((b*c^3*d^3*e^3 - b*c*d*e^5)*x^2 + (b*c^3*d^4*e^2 - b*c*d^2*e^4)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2))/(c^4*d^8*e - 2*c^2*d^6*e^3 + d^4*e^5 + (c^4*d^6*e^3 - 2*c^2*d^4*e^5 + d^2*e^7)*x^2 + 2*(c^4*d^7*e^2 - 2*c^2*d^5*e^4 + d^3*e^6)*x), -1/2*(a*c^4*d^6 - (2*a + b)*c^2*d^4*e^2 + (a + b)*d^2*e^4 - (b*c^2*d^2*e^4 - b*e^6)*x^2 - 2*(2*b*c^2*d^4*e - b*d^2*e^3 + (2*b*c^2*d^2*e^3 - b*e^5)*x^2 + 2*(2*b*c^2*d^3*e^2 - b*d*e^4)*x)*sqrt(c^2*d^2 - e^2)*arctan(-(sqrt(c^2*d^2 - e^2)*c*d*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - sqrt(c^2*d^2 - e^2)*(e*x + d))/(c^2*d^2 - e^2)*x)) - 2*(b*c^2*d^3*e^3 - b*d*e^5)*x + (b*c^4*d^6 - 2*b*c^2*d^4*e^2 + b*d^2*e^4 + (b*c^4*d^4*e^2 - 2*b*c^2*d^2*e^4 + b*e^6)*x^2 + 2*(b*c^4*d^5*e - 2*b*c^2*d^3*e^3 + b*d*e^5)*x)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) + (b*c^4*d^6 - ...

```

### Sympy [F]

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^3} dx = \int \frac{a + b \operatorname{asech}(cx)}{(d + ex)^3} dx$$

input

```
integrate((a+b*asech(c*x))/(e*x+d)**3,x)
```

output

```
Integral((a + b*asech(c*x))/(d + e*x)**3, x)
```

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+b*arcsech(c*x))/(e*x+d)^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^3} dx = \int \frac{b \operatorname{ar} \operatorname{sech}(cx) + a}{(ex + d)^3} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x+d)^3,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/(e*x + d)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^3} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{(d + ex)^3} dx$$

input `int((a + b*acosh(1/(c*x)))/(d + e*x)^3,x)`

output `int((a + b*acosh(1/(c*x)))/(d + e*x)^3, x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^3} dx$$

$$= \frac{2 \left( \int \frac{\operatorname{asech}(cx)}{e^3 x^3 + 3d e^2 x^2 + 3d^2 ex + d^3} dx \right) b d^2 e + 4 \left( \int \frac{\operatorname{asech}(cx)}{e^3 x^3 + 3d e^2 x^2 + 3d^2 ex + d^3} dx \right) b d e^2 x + 2 \left( \int \frac{\operatorname{asech}(cx)}{e^3 x^3 + 3d e^2 x^2 + 3d^2 ex + d^3} dx \right) b e^3 x^2}{2e(e^2 x^2 + 2dex + d^2)}$$

input `int((a+b*asech(c*x))/(e*x+d)^3,x)`

output `(2*int(asech(c*x)/(d**3 + 3*d**2*e*x + 3*d*e**2*x**2 + e**3*x**3),x)*b*d**2*e + 4*int(asech(c*x)/(d**3 + 3*d**2*e*x + 3*d*e**2*x**2 + e**3*x**3),x)*b*d*e**2*x + 2*int(asech(c*x)/(d**3 + 3*d**2*e*x + 3*d*e**2*x**2 + e**3*x**3),x)*b*e**3*x**2 - a)/(2*e*(d**2 + 2*d*e*x + e**2*x**2))`

### 3.80 $\int (d + ex)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx$

Optimal result	643
Mathematica [C] (warning: unable to verify)	644
Rubi [A] (verified)	645
Maple [B] (verified)	651
Fricas [F(-1)]	652
Sympy [F]	652
Maxima [F(-2)]	653
Giac [F]	653
Mupad [F(-1)]	654
Reduce [F]	654

#### Optimal result

Integrand size = 18, antiderivative size = 343

$$\begin{aligned}
 & \int (d + ex)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \\
 & - \frac{4be \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{d+ex} \sqrt{1-c^2x^2}}{15c^2} + \frac{2(d+ex)^{5/2} (a + b \operatorname{sech}^{-1}(cx))}{5e} \\
 & - \frac{28bd \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{d+ex} E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right) \mid \frac{2e}{cd+e}\right)}{15c \sqrt{\frac{c(d+ex)}{cd+e}}} \\
 & - \frac{4b(2c^2d^2 + e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{\frac{c(d+ex)}{cd+e}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{15c^3 \sqrt{d+ex}} \\
 & - \frac{4bd^3 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{\frac{c(d+ex)}{cd+e}} \operatorname{EllipticPi}\left(2, \arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{5e \sqrt{d+ex}}
 \end{aligned}$$

output

```
-4/15*b*e*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(e*x+d)^(1/2)*(-c^2*x^2+1)^(1/2)
/c^2+2/5*(e*x+d)^(5/2)*(a+b*arcsech(c*x))/e-28/15*b*d*(1/(c*x+1))^(1/2)*(c
*x+1)^(1/2)*(e*x+d)^(1/2)*EllipticE(1/2*(-c*x+1)^(1/2)*2^(1/2),2^(1/2)*(e/
(c*d+e))^(1/2))/c/(c*(e*x+d)/(c*d+e))^(1/2)-4/15*b*(2*c^2*d^2+e^2)*(1/(c*x
+1))^(1/2)*(c*x+1)^(1/2)*(c*(e*x+d)/(c*d+e))^(1/2)*EllipticF(1/2*(-c*x+1)^(
1/2)*2^(1/2),2^(1/2)*(e/(c*d+e))^(1/2))/c^3/(e*x+d)^(1/2)-4/5*b*d^3*(1/(c
*x+1))^(1/2)*(c*x+1)^(1/2)*(c*(e*x+d)/(c*d+e))^(1/2)*EllipticPi(1/2*(-c*x+
1)^(1/2)*2^(1/2),2,2^(1/2)*(e/(c*d+e))^(1/2))/e/(e*x+d)^(1/2)
```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 19.99 (sec) , antiderivative size = 2653, normalized size of antiderivative = 7.73

$$\int (d + ex)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \text{Result too large to show}$$

input

```
Integrate[(d + e*x)^(3/2)*(a + b*ArcSech[c*x]),x]
```

output

```

Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[d + e*x]*((-4*b*e)/(15*c^2) - (4*b*e*x)/(15
*c)) + Sqrt[d + e*x]*((2*a*d^2)/(5*e) + (4*a*d*x)/5 + (2*a*e*x^2)/5) + (2*
b*(d + e*x)^(5/2)*ArcSech[c*x])/(5*e) - (4*b*(7*c*d*e*Sqrt[(1 - c*x)/(1 +
c*x)]*(e - (e*(1 - c*x))/(1 + c*x) + c*d*(1 + (1 - c*x)/(1 + c*x)))) + ((7*
I)*c^2*d^2*e*(c*d + e)*Sqrt[1 + (1 - c*x)/(1 + c*x)]*Sqrt[(e - (e*(1 - c*x)
))/(1 + c*x) + c*d*(1 + (1 - c*x)/(1 + c*x))]/(c*d + e)]*(EllipticE[I*ArcS
inh[Sqrt[(1 - c*x)/(1 + c*x)]], (c*d - e)/(c*d + e)] - EllipticF[I*ArcSinh
[Sqrt[(1 - c*x)/(1 + c*x)]], (c*d - e)/(c*d + e)]))/(c*d - e) - ((7*I)*c*d
*e^2*(c*d + e)*Sqrt[1 + (1 - c*x)/(1 + c*x)]*Sqrt[(e - (e*(1 - c*x))/(1 +
c*x) + c*d*(1 + (1 - c*x)/(1 + c*x)))/(c*d + e)]*(EllipticE[I*ArcSinh[Sqrt
[(1 - c*x)/(1 + c*x)]], (c*d - e)/(c*d + e)] - EllipticF[I*ArcSinh[Sqrt[(1
- c*x)/(1 + c*x)]], (c*d - e)/(c*d + e)]))/(c*d - e) + (3*I)*c^3*d^3*Sqrt
[1 + (1 - c*x)/(1 + c*x)]*Sqrt[(e - (e*(1 - c*x))/(1 + c*x) + c*d*(1 + (1
- c*x)/(1 + c*x)))/(c*d + e)]*EllipticF[I*ArcSinh[Sqrt[(1 - c*x)/(1 + c*x)
]], (c*d - e)/(c*d + e)] - (2*I)*c^2*d^2*e*Sqrt[1 + (1 - c*x)/(1 + c*x)]*S
qrt[(e - (e*(1 - c*x))/(1 + c*x) + c*d*(1 + (1 - c*x)/(1 + c*x)))/(c*d + e
)]*EllipticF[I*ArcSinh[Sqrt[(1 - c*x)/(1 + c*x)]], (c*d - e)/(c*d + e)] -
I*e^3*Sqrt[1 + (1 - c*x)/(1 + c*x)]*Sqrt[(e - (e*(1 - c*x))/(1 + c*x) + c*
d*(1 + (1 - c*x)/(1 + c*x)))/(c*d + e)]*EllipticF[I*ArcSinh[Sqrt[(1 - c*x)
/(1 + c*x)]], (c*d - e)/(c*d + e)] + ((3 + 3*I)*c^3*d^3*(-I + Sqrt[(1 - ...

```

## Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 306, normalized size of antiderivative = 0.89, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.722$ , Rules used = {6842, 634, 632, 186, 413, 412, 2185, 27, 600, 508, 327, 511, 321}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (d + ex)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) \, dx \\
 & \qquad \qquad \qquad \downarrow \text{6842} \\
 & \frac{2b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{(d+ex)^{5/2}}{x\sqrt{1-c^2x^2}} dx}{5e} + \frac{2(d+ex)^{5/2} (a + b \operatorname{sech}^{-1}(cx))}{5e} \\
 & \qquad \qquad \qquad \downarrow \text{634}
 \end{aligned}$$

$$\begin{aligned}
& \frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(d^3\int\frac{1}{x\sqrt{d+ex}\sqrt{1-c^2x^2}}dx - \int\frac{-x^2e^3-3dxe^2-3d^2e}{\sqrt{d+ex}\sqrt{1-c^2x^2}}dx\right)}{2(d+ex)^{5/2}(a+b\operatorname{sech}^{-1}(cx))} + \\
& \quad \frac{5e}{5e} \downarrow \mathbf{632} \\
& \frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(d^3\int\frac{1}{x\sqrt{1-cx}\sqrt{cx+1}\sqrt{d+ex}}dx - \int\frac{-x^2e^3-3dxe^2-3d^2e}{\sqrt{d+ex}\sqrt{1-c^2x^2}}dx\right)}{2(d+ex)^{5/2}(a+b\operatorname{sech}^{-1}(cx))} + \\
& \quad \frac{5e}{5e} \downarrow \mathbf{186} \\
& \frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\int\frac{-x^2e^3-3dxe^2-3d^2e}{\sqrt{d+ex}\sqrt{1-c^2x^2}}dx - 2d^3\int\frac{1}{cx\sqrt{cx+1}\sqrt{d+\frac{e}{c}-\frac{e(1-cx)}{c}}}d\sqrt{1-cx}\right)}{2(d+ex)^{5/2}(a+b\operatorname{sech}^{-1}(cx))} + \\
& \quad \frac{5e}{5e} \downarrow \mathbf{413} \\
& \frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\int\frac{-x^2e^3-3dxe^2-3d^2e}{\sqrt{d+ex}\sqrt{1-c^2x^2}}dx - \frac{2d^3\sqrt{1-\frac{e(1-cx)}{cd+e}}\int\frac{1}{cx\sqrt{cx+1}\sqrt{1-\frac{e(1-cx)}{cd+e}}}d\sqrt{1-cx}}{\sqrt{-\frac{e(1-cx)}{c}+\frac{e}{c}+d}}\right)}{2(d+ex)^{5/2}(a+b\operatorname{sech}^{-1}(cx))} + \\
& \quad \frac{5e}{5e} \downarrow \mathbf{412} \\
& \frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\int\frac{-x^2e^3-3dxe^2-3d^2e}{\sqrt{d+ex}\sqrt{1-c^2x^2}}dx - \frac{2d^3\sqrt{1-\frac{e(1-cx)}{cd+e}}\operatorname{EllipticPi}\left(2,\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right),\frac{2e}{cd+e}\right)}{\sqrt{-\frac{e(1-cx)}{c}+\frac{e}{c}+d}}\right)}{2(d+ex)^{5/2}(a+b\operatorname{sech}^{-1}(cx))} + \\
& \quad \frac{5e}{5e} \downarrow \mathbf{2185} \\
& \frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{2\int\frac{e^3(9d^2c^2+7dexc^2+e^2)}{2\sqrt{d+ex}\sqrt{1-c^2x^2}}dx}{3c^2e^2} - \frac{2d^3\sqrt{1-\frac{e(1-cx)}{cd+e}}\operatorname{EllipticPi}\left(2,\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right),\frac{2e}{cd+e}\right)}{\sqrt{-\frac{e(1-cx)}{c}+\frac{e}{c}+d}} - \frac{2e^2\sqrt{1-c^2x^2}\sqrt{d+ex}}{3c^2}\right)}{2(d+ex)^{5/2}(a+b\operatorname{sech}^{-1}(cx))} + \\
& \quad \frac{5e}{5e} \downarrow \mathbf{27}
\end{aligned}$$

$$2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{e \int \frac{9d^2c^2+7dexc^2+e^2}{\sqrt{d+ex}\sqrt{1-c^2x^2}} dx}{3c^2} - \frac{2d^3\sqrt{1-\frac{e(1-cx)}{cd+e}} \operatorname{EllipticPi}\left(2, \arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{\sqrt{-\frac{e(1-cx)}{c} + \frac{e}{c} + d}} - \frac{2e^2\sqrt{1-c^2x^2}\sqrt{d+ex}}{3c^2} \right) +$$

$$\frac{2(d+ex)^{5/2} (a + b\operatorname{sech}^{-1}(cx))}{5e}$$

↓ 600

$$2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{e \left( (2c^2d^2+e^2) \int \frac{1}{\sqrt{d+ex}\sqrt{1-c^2x^2}} dx + 7c^2d \int \frac{\sqrt{d+ex}}{\sqrt{1-c^2x^2}} dx \right)}{3c^2} - \frac{2d^3\sqrt{1-\frac{e(1-cx)}{cd+e}} \operatorname{EllipticPi}\left(2, \arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{\sqrt{-\frac{e(1-cx)}{c} + \frac{e}{c} + d}} \right) +$$

$$\frac{2(d+ex)^{5/2} (a + b\operatorname{sech}^{-1}(cx))}{5e}$$

↓ 508

$$2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{e \left( (2c^2d^2+e^2) \int \frac{1}{\sqrt{d+ex}\sqrt{1-c^2x^2}} dx - \frac{14cd\sqrt{d+ex} \int \frac{\sqrt{1-\frac{e(1-cx)}{cd+e}} d\sqrt{1-cx}}{\sqrt{\frac{1}{2}(cx-1)+1} \sqrt{2}} dx}{\sqrt{\frac{c(d+ex)}{cd+e}}} \right)}{3c^2} - \frac{2d^3\sqrt{1-\frac{e(1-cx)}{cd+e}} \operatorname{EllipticPi}\left(2, \arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{\sqrt{-\frac{e(1-cx)}{c} + \frac{e}{c} + d}} \right) +$$

$$\frac{2(d+ex)^{5/2} (a + b\operatorname{sech}^{-1}(cx))}{5e}$$

↓ 327

$$2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{e \left( (2c^2d^2+e^2) \int \frac{1}{\sqrt{d+ex}\sqrt{1-c^2x^2}} dx - \frac{14cd\sqrt{d+ex} E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right)\right) \frac{2e}{cd+e}}{\sqrt{\frac{c(d+ex)}{cd+e}}} \right)}{3c^2} - \frac{2d^3\sqrt{1-\frac{e(1-cx)}{cd+e}} \operatorname{EllipticPi}\left(2, \arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{\sqrt{-\frac{e(1-cx)}{c} + \frac{e}{c} + d}} \right) +$$

$$\frac{2(d+ex)^{5/2} (a + b\operatorname{sech}^{-1}(cx))}{5e}$$

↓ 511



$$\begin{aligned}
 & 2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{e \left( \frac{2(2c^2d^2+e^2)\sqrt{\frac{c(d+ex)}{cd+e}} \int \frac{1}{\sqrt{1-\frac{e(1-cx)}{cd+e}} \sqrt{\frac{1}{2}(cx-1)+1}} d\sqrt{\frac{1-cx}{2}}} - \frac{14cd\sqrt{d+ex}E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right)\middle|\frac{2e}{cd+e}\right)}{\sqrt{\frac{c(d+ex)}{cd+e}}} \right)}{3c^2} \right) - \frac{2d^3\sqrt{1-\frac{e(1-cx)}{cd+e}}}{5e} \\
 & \frac{2(d+ex)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e} \\
 & \quad \downarrow \text{321} \\
 & \frac{2(d+ex)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e} + \\
 & 2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{e \left( \frac{2(2c^2d^2+e^2)\sqrt{\frac{c(d+ex)}{cd+e}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{c\sqrt{d+ex}} - \frac{14cd\sqrt{d+ex}E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right)\middle|\frac{2e}{cd+e}\right)}{\sqrt{\frac{c(d+ex)}{cd+e}}} \right)}{3c^2} \right) - \frac{2d^3\sqrt{1-\frac{e(1-cx)}{cd+e}}}{5e}
 \end{aligned}$$

input `Int[(d + e*x)^(3/2)*(a + b*ArcSech[c*x]),x]`

output `(2*(d + e*x)^(5/2)*(a + b*ArcSech[c*x]))/(5*e) + (2*b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-2*e^2*Sqrt[d + e*x]*Sqrt[1 - c^2*x^2])/(3*c^2) + (e*((-1 - 4*c*d*Sqrt[d + e*x]*EllipticE[ArcSin[Sqrt[1 - c*x]/Sqrt[2]], (2*e)/(c*d + e)))/Sqrt[(c*(d + e*x))/(c*d + e)] - (2*(2*c^2*d^2 + e^2)*Sqrt[(c*(d + e*x))/(c*d + e)]*EllipticF[ArcSin[Sqrt[1 - c*x]/Sqrt[2]], (2*e)/(c*d + e)]/(c*Sqrt[d + e*x])))/(3*c^2) - (2*d^3*Sqrt[1 - (e*(1 - c*x))/(c*d + e)]*EllipticPi[2, ArcSin[Sqrt[1 - c*x]/Sqrt[2]], (2*e)/(c*d + e)]/Sqrt[d + e/c - (e*(1 - c*x))/c]))/(5*e)`

## Definitions of rubi rules used

- rule 27 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx) /; FreeQ[b, x]]`
- rule 186 `Int[1/(((a_) + (b_)*(x_))*Sqrt[(c_) + (d_)*(x_)]*Sqrt[(e_) + (f_)*(x_)]*Sqrt[(g_) + (h_)*(x_)]), x_] := Simp[-2 Subst[Int[1/(Simp[b*c - a*d - b*x^2, x]*Sqrt[Simp[(d*e - c*f)/d + f*(x^2/d), x]]*Sqrt[Simp[(d*g - c*h)/d + h*(x^2/d), x]]), x], x, Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x] && GtQ[(d*e - c*f)/d, 0]`
- rule 321 `Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] := Simp[(1/(Sqrt[a]*Sqrt[c]*Rt[-d/c, 2]))*EllipticF[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplifierSqrtQ[-b/a, -d/c])`
- rule 327 `Int[Sqrt[(a_) + (b_)*(x_)^2]/Sqrt[(c_) + (d_)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]/(Sqrt[c]*Rt[-d/c, 2]))*EllipticE[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]`
- rule 412 `Int[1/(((a_) + (b_)*(x_)^2)*Sqrt[(c_) + (d_)*(x_)^2]*Sqrt[(e_) + (f_)*(x_)^2]), x_Symbol] := Simp[(1/(a*Sqrt[c]*Sqrt[e]*Rt[-d/c, 2]))*EllipticPi[b*(c/(a*d)), ArcSin[Rt[-d/c, 2]*x], c*(f/(d*e))], x] /; FreeQ[{a, b, c, d, e, f}, x] && !GtQ[d/c, 0] && GtQ[c, 0] && GtQ[e, 0] && !( !GtQ[f/e, 0] && SimplifierSqrtQ[-f/e, -d/c])`
- rule 413 `Int[1/(((a_) + (b_)*(x_)^2)*Sqrt[(c_) + (d_)*(x_)^2]*Sqrt[(e_) + (f_)*(x_)^2]), x_Symbol] := Simp[Sqrt[1 + (d/c)*x^2]/Sqrt[c + d*x^2] Int[1/((a + b*x^2)*Sqrt[1 + (d/c)*x^2]*Sqrt[e + f*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && !GtQ[c, 0]`

rule 508 `Int[Sqrt[(c_) + (d_)*(x_)]/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := With[{q = Rt[-b/a, 2]}, Simp[-2*(Sqrt[c + d*x]/(Sqrt[a]*q*Sqrt[q*(c + d*x)/(d + c*q)]))] Subst[Int[Sqrt[1 - 2*d*(x^2/(d + c*q))]/Sqrt[1 - x^2], x], x, Sqrt[(1 - q*x)/2]], x] /; FreeQ[{a, b, c, d}, x] && NegQ[b/a] && GtQ[a, 0]`

rule 511 `Int[1/(Sqrt[(c_) + (d_)*(x_)]*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] := With[{q = Rt[-b/a, 2]}, Simp[-2*(Sqrt[q*(c + d*x)/(d + c*q)]/(Sqrt[a]*q*Sqrt[c + d*x])) Subst[Int[1/(Sqrt[1 - 2*d*(x^2/(d + c*q))]*Sqrt[1 - x^2]), x], x, Sqrt[(1 - q*x)/2]], x] /; FreeQ[{a, b, c, d}, x] && NegQ[b/a] && GtQ[a, 0]`

rule 600 `Int[((A_) + (B_)*(x_))/(Sqrt[(c_) + (d_)*(x_)]*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] := Simp[B/d Int[Sqrt[c + d*x]/Sqrt[a + b*x^2], x], x] - Simp[(B*c - A*d)/d Int[1/(Sqrt[c + d*x]*Sqrt[a + b*x^2]), x], x] /; FreeQ[{a, b, c, d, A, B}, x] && NegQ[b/a]`

rule 632 `Int[1/((x_)*Sqrt[(c_) + (d_)*(x_)]*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] := With[{q = Rt[-b/a, 2]}, Simp[1/Sqrt[a] Int[1/(x*Sqrt[c + d*x]*Sqrt[1 - q*x]*Sqrt[1 + q*x]), x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[b/a] && GtQ[a, 0]`

rule 634 `Int[((c_) + (d_)*(x_))^(n_)/((x_)*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] := Simp[c^(n + 1/2) Int[1/(x*Sqrt[c + d*x]*Sqrt[a + b*x^2]), x], x] - Int[(1/(Sqrt[c + d*x]*Sqrt[a + b*x^2]))*ExpandToSum[(c^(n + 1/2) - (c + d*x)^(n + 1/2))/x, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[n - 1/2, 0]`

rule 2185

```
Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] :
> With[{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[f*(d + e*x)
^(m + q - 1)*((a + b*x^2)^(p + 1)/(b*e^(q - 1)*(m + q + 2*p + 1))), x] + Si
mp[1/(b*e^q*(m + q + 2*p + 1)) Int[(d + e*x)^m*(a + b*x^2)^p*ExpandToSum[
b*e^q*(m + q + 2*p + 1)*Pq - b*f*(m + q + 2*p + 1)*(d + e*x)^q - f*(d + e*x
)^(q - 2)*(a*e^2*(m + q - 1) - b*d^2*(m + q + 2*p + 1) - 2*b*d*e*(m + q + p
)*x), x], x], x] /; GtQ[q, 1] && NeQ[m + q + 2*p + 1, 0]] /; FreeQ[{a, b, d
, e, m, p}, x] && PolyQ[Pq, x] && NeQ[b*d^2 + a*e^2, 0] && !(EqQ[d, 0] &&
True) && !(IGtQ[m, 0] && RationalQ[a, b, d, e] && (IntegerQ[p] || ILtQ[p +
1/2, 0]))
```

rule 6842

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*((d_) + (e_)*(x_))^(m_), x_Symbol]
:> Simp[(d + e*x)^(m + 1)*((a + b*ArcSech[c*x])/(e*(m + 1))), x] + Simp[
b*(Sqrt[1 + c*x]/(e*(m + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x)^(m + 1)/(x*
Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[m, -1]
```

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 817 vs. 2(302) = 604.

Time = 18.74 (sec) , antiderivative size = 818, normalized size of antiderivative = 2.38

method	result
derivativedivides	$\frac{2a(ex+d)^{\frac{5}{2}}}{5} + 2b \left( \frac{(ex+d)^{\frac{5}{2}} \operatorname{arcsech}(cx)}{5} - \frac{2e^2 \sqrt{\frac{-c(ex+d)+cd+e}{cex}} x \sqrt{\frac{-c(ex+d)+cd-e}{cex}} \left( \sqrt{\frac{c}{cd+e}} c^2 (ex+d)^{\frac{5}{2}} + 9 \sqrt{\frac{-c(ex+d)+cd}{cd+e}} \right)}{5} \right)$
default	$\frac{2a(ex+d)^{\frac{5}{2}}}{5} + 2b \left( \frac{(ex+d)^{\frac{5}{2}} \operatorname{arcsech}(cx)}{5} - \frac{2e^2 \sqrt{\frac{-c(ex+d)+cd+e}{cex}} x \sqrt{\frac{-c(ex+d)+cd-e}{cex}} \left( \sqrt{\frac{c}{cd+e}} c^2 (ex+d)^{\frac{5}{2}} + 9 \sqrt{\frac{-c(ex+d)+cd}{cd+e}} \right)}{5} \right)$
parts	$\frac{2a(ex+d)^{\frac{5}{2}}}{5e} + 2b \left( \frac{(ex+d)^{\frac{5}{2}} \operatorname{arcsech}(cx)}{5} - \frac{2e^2 \sqrt{\frac{-c(ex+d)-cd-e}{cxe}} x \sqrt{\frac{c(ex+d)-cd+e}{cxe}} \left( \sqrt{\frac{c}{cd+e}} c^2 (ex+d)^{\frac{5}{2}} + 9 \sqrt{\frac{-c(ex+d)-cd}{cd+e}} \right)}{5} \right)$

input

```
int((e*x+d)^(3/2)*(a+b*arcsech(c*x)), x, method=_RETURNVERBOSE)
```

output

```

2/e*(1/5*a*(e*x+d)^(5/2)+b*(1/5*(e*x+d)^(5/2)*arcsech(c*x)-2/15/c*e^2*((-c
*(e*x+d)+c*d+e)/c/e/x)^(1/2)*x*(-(-c*(e*x+d)+c*d-e)/c/e/x)^(1/2)*((c/(c*d+
e))^(1/2)*c^2*(e*x+d)^(5/2)+9*((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x
+d)+c*d-e)/(c*d-e))^(1/2)*EllipticF((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+
e)/(c*d-e))^(1/2))*c^2*d^2-7*((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x+
d)+c*d-e)/(c*d-e))^(1/2)*EllipticE((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e
)/(c*d-e))^(1/2))*c^2*d^2-3*((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x+d
)+c*d-e)/(c*d-e))^(1/2)*EllipticPi((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),1/c*(c*
d+e)/d,(c/(c*d-e))^(1/2)/(c/(c*d+e))^(1/2))*c^2*d^2-2*(c/(c*d+e))^(1/2)*c^
2*d*(e*x+d)^(3/2)-7*((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x+d)+c*d-e)
/(c*d-e))^(1/2)*EllipticF((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e)/(c*d-e)
)^(1/2))*c*d*e+7*((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x+d)+c*d-e)/(c
*d-e))^(1/2)*EllipticE((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e)/(c*d-e))^(
1/2))*c*d*e+(c/(c*d+e))^(1/2)*c^2*d^2*(e*x+d)^(1/2)+((-c*(e*x+d)+c*d+e)/(c
*d+e))^(1/2)*((-c*(e*x+d)+c*d-e)/(c*d-e))^(1/2)*EllipticF((e*x+d)^(1/2)*(c
/(c*d+e))^(1/2),((c*d+e)/(c*d-e))^(1/2))*e^2-(c/(c*d+e))^(1/2)*e^2*(e*x+d)
^(1/2))/(c/(c*d+e))^(1/2)/(c^2*(e*x+d)^2-2*c^2*d*(e*x+d)+c^2*d^2-e^2))

```

**Fricas [F(-1)]**

Timed out.

$$\int (d + ex)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \text{Timed out}$$

input

```
integrate((e*x+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="fricas")
```

output

Timed out

**Sympy [F]**

$$\int (d + ex)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int (a + b \operatorname{asech}(cx)) (d + ex)^{3/2} dx$$

input

```
integrate((e*x+d)**(3/2)*(a+b*asech(c*x)),x)
```

output `Integral((a + b*asech(c*x))*(d + e*x)**(3/2), x)`

### Maxima [F(-2)]

Exception generated.

$$\int (d + ex)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \text{Exception raised: ValueError}$$

input `integrate((e*x+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e+c*d>0)', see `assume?` for more details)`

### Giac [F]

$$\int (d + ex)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex + d)^{\frac{3}{2}} (b\operatorname{ar}\operatorname{sech}(cx) + a) dx$$

input `integrate((e*x+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((e*x + d)^(3/2)*(b*arcsech(c*x) + a), x)`

**Mupad [F(-1)]**

Timed out.

$$\int (d + ex)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) (d + ex)^{3/2} dx$$

input `int((a + b*acosh(1/(c*x)))*(d + e*x)^(3/2), x)`

output `int((a + b*acosh(1/(c*x)))*(d + e*x)^(3/2), x)`

**Reduce [F]**

$$\int (d + ex)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \frac{2\sqrt{ex + d} a d^2 + 4\sqrt{ex + d} a d e x + 2\sqrt{ex + d} a e^2 x^2 + 5 \left( \int \sqrt{ex + d} a \operatorname{sech}(cx) x dx \right) b e}{5e}$$

input `int((e*x+d)^(3/2)*(a+b*asech(c*x)), x)`

output `(2*sqrt(d + e*x)*a*d**2 + 4*sqrt(d + e*x)*a*d*e*x + 2*sqrt(d + e*x)*a*e**2 *x**2 + 5*int(sqrt(d + e*x)*asech(c*x)*x,x)*b*e**2 + 5*int(sqrt(d + e*x)*a sech(c*x), x)*b*d*e)/(5*e)`

### 3.81 $\int \sqrt{d+ex}(a+b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	655
Mathematica [C] (warning: unable to verify)	656
Rubi [A] (verified)	657
Maple [A] (verified)	661
Fricas [F(-1)]	662
Sympy [F]	662
Maxima [F(-2)]	663
Giac [F]	663
Mupad [F(-1)]	663
Reduce [F]	664

#### Optimal result

Integrand size = 18, antiderivative size = 279

$$\begin{aligned}
 & \int \sqrt{d+ex}(a+b\operatorname{sech}^{-1}(cx)) dx \\
 &= \frac{2(d+ex)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e} \\
 & \quad - \frac{4b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{d+ex}E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right)\middle|\frac{2e}{cd+e}\right)}{3c\sqrt{\frac{c(d+ex)}{cd+e}}} \\
 & \quad - \frac{4bd\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{\frac{c(d+ex)}{cd+e}}\operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right),\frac{2e}{cd+e}\right)}{3c\sqrt{d+ex}} \\
 & \quad - \frac{4bd^2\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{\frac{c(d+ex)}{cd+e}}\operatorname{EllipticPi}\left(2,\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right),\frac{2e}{cd+e}\right)}{3e\sqrt{d+ex}}
 \end{aligned}$$

output

```

2/3*(e*x+d)^(3/2)*(a+b*arcsech(c*x))/e-4/3*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/
2)*(e*x+d)^(1/2)*EllipticE(1/2*(-c*x+1)^(1/2)*2^(1/2),2^(1/2)*(e/(c*d+e))^(
1/2))/c/(c*(e*x+d)/(c*d+e))^(1/2)-4/3*b*d*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)
*(c*(e*x+d)/(c*d+e))^(1/2)*EllipticF(1/2*(-c*x+1)^(1/2)*2^(1/2),2^(1/2)*(e
/(c*d+e))^(1/2))/c/(e*x+d)^(1/2)-4/3*b*d^2*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)
*(c*(e*x+d)/(c*d+e))^(1/2)*EllipticPi(1/2*(-c*x+1)^(1/2)*2^(1/2),2,2^(1/2)
*(e/(c*d+e))^(1/2))/e/(e*x+d)^(1/2)

```



**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 24.88 (sec) , antiderivative size = 2938, normalized size of antiderivative = 10.53

$$\int \sqrt{d+ex}(a+b\operatorname{sech}^{-1}(cx)) dx = \text{Result too large to show}$$

input `Integrate[Sqrt[d + e*x]*(a + b*ArcSech[c*x]),x]`

output

```
((2*a*d)/(3*e) + (2*a*x)/3)*Sqrt[d + e*x] + (2*b*(d + e*x)^(3/2)*ArcSech[c*x])/(3*e) + (4*b*(-((e*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[c*(1 + (1 - c*x)/(1 + c*x))])*Sqrt[c + (c*(1 - c*x))/(1 + c*x)]*Sqrt[(c*d + e + (c*d*(1 - c*x))/(1 + c*x) - (e*(1 - c*x))/(1 + c*x))/(c + (c*(1 - c*x))/(1 + c*x))])/(c*(1 + (1 - c*x)/(1 + c*x)))) + (Sqrt[c*(1 + (1 - c*x)/(1 + c*x))]*Sqrt[c + (c*(1 - c*x)/(1 + c*x)]*Sqrt[c*(1 + (1 - c*x)/(1 + c*x))*(c*d + e + (c*d*(1 - c*x))/(1 + c*x) - (e*(1 - c*x))/(1 + c*x)]*Sqrt[(c*d + e + (c*d*(1 - c*x))/(1 + c*x) - (e*(1 - c*x))/(1 + c*x))/(c + (c*(1 - c*x))/(1 + c*x))])*((I*c*d*(-(c*d) - e)*e*Sqrt[1 + (1 - c*x)/(1 + c*x)]*Sqrt[1 - ((c*d - e)*(1 - c*x))/((-c*d) - e)*(1 + c*x)])*(EllipticE[I*ArcSinh[Sqrt[(1 - c*x)/(1 + c*x)]], -(c*d - e)/(-(c*d) - e)] - EllipticF[I*ArcSinh[Sqrt[(1 - c*x)/(1 + c*x)]], -(c*d - e)/(-(c*d) - e)]))/((c*d - e)*Sqrt[c*(1 + (1 - c*x)/(1 + c*x))*(c*d + e + ((c*d - e)*(1 - c*x))/(1 + c*x))]) - (I*(-(c*d) - e)*e^2*Sqrt[1 + (1 - c*x)/(1 + c*x)]*Sqrt[1 - ((c*d - e)*(1 - c*x))/((-c*d) - e)*(1 + c*x)])*(EllipticE[I*ArcSinh[Sqrt[(1 - c*x)/(1 + c*x)]], -(c*d - e)/(-(c*d) - e)] - EllipticF[I*ArcSinh[Sqrt[(1 - c*x)/(1 + c*x)]], -(c*d - e)/(-(c*d) - e)]))/((c*d - e)*Sqrt[c*(1 + (1 - c*x)/(1 + c*x))*(c*d + e + ((c*d - e)*(1 - c*x))/(1 + c*x))]) - (I*c^2*d^2*Sqrt[1 + (1 - c*x)/(1 + c*x)]*Sqrt[1 - ((c*d - e)*(1 - c*x))/((-c*d) - e)*(1 + c*x)])*EllipticF[I*ArcSinh[Sqrt[(1 - c*x)/(1 + c*x)]], -(c*d - e)/(-(c*d) - e)])...
```

**Rubi [A] (verified)**

Time = 0.62 (sec) , antiderivative size = 256, normalized size of antiderivative = 0.92, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.611$ , Rules used = {6842, 634, 600, 508, 327, 511, 321, 632, 186, 413, 412}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{d+ex}(a+b\operatorname{sech}^{-1}(cx)) \, dx \\
 & \quad \downarrow \text{6842} \\
 & \frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(d+ex)^{3/2}}{x\sqrt{1-c^2x^2}} dx}{3e} + \frac{2(d+ex)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e} \\
 & \quad \downarrow \text{634} \\
 & \frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( d^2 \int \frac{1}{x\sqrt{d+ex}\sqrt{1-c^2x^2}} dx - \int \frac{-xe^2-2de}{\sqrt{d+ex}\sqrt{1-c^2x^2}} dx \right)}{3e} + \\
 & \quad \frac{2(d+ex)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e} \\
 & \quad \downarrow \text{600} \\
 & \frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( d^2 \int \frac{1}{x\sqrt{d+ex}\sqrt{1-c^2x^2}} dx + de \int \frac{1}{\sqrt{d+ex}\sqrt{1-c^2x^2}} dx + e \int \frac{\sqrt{d+ex}}{\sqrt{1-c^2x^2}} dx \right)}{3e} + \\
 & \quad \frac{2(d+ex)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e} \\
 & \quad \downarrow \text{508} \\
 & \frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( d^2 \int \frac{1}{x\sqrt{d+ex}\sqrt{1-c^2x^2}} dx + de \int \frac{1}{\sqrt{d+ex}\sqrt{1-c^2x^2}} dx - \frac{2e\sqrt{d+ex} \int \frac{\sqrt{1-\frac{e(1-cx)}{cd+e}} d\sqrt{1-cx}}{\sqrt{\frac{1}{2}(cx-1)+1} \sqrt{2}} dx}{c\sqrt{\frac{c(d+ex)}{cd+e}}} \right)}{3e} + \\
 & \quad \frac{2(d+ex)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e} \\
 & \quad \downarrow \text{327}
 \end{aligned}$$

$$\frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(d^2\int\frac{1}{x\sqrt{d+ex}\sqrt{1-c^2x^2}}dx+de\int\frac{1}{\sqrt{d+ex}\sqrt{1-c^2x^2}}dx-\frac{2e\sqrt{d+ex}E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right)\middle|\frac{2e}{cd+e}\right)}{c\sqrt{\frac{c(d+ex)}{cd+e}}}\right)}{2(d+ex)^{3/2}(a+b\operatorname{sech}^{-1}(cx))} + \frac{3e}{3e}$$

511

$$\frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(d^2\int\frac{1}{x\sqrt{d+ex}\sqrt{1-c^2x^2}}dx-\frac{2de\sqrt{\frac{c(d+ex)}{cd+e}}\int\frac{1}{\sqrt{1-\frac{e(1-cx)}{cd+e}}\sqrt{\frac{1}{2}(cx-1)+1}}\frac{d\sqrt{1-cx}}{\sqrt{2}}}{c\sqrt{d+ex}}-\frac{2e\sqrt{d+ex}E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right)\middle|\frac{2e}{cd+e}\right)}{c\sqrt{\frac{c(d+ex)}{cd+e}}}\right)}{2(d+ex)^{3/2}(a+b\operatorname{sech}^{-1}(cx))} + \frac{3e}{3e}$$

321

$$\frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(d^2\int\frac{1}{x\sqrt{d+ex}\sqrt{1-c^2x^2}}dx-\frac{2de\sqrt{\frac{c(d+ex)}{cd+e}}\operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right),\frac{2e}{cd+e}\right)}{c\sqrt{d+ex}}-\frac{2e\sqrt{d+ex}E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right)\middle|\frac{2e}{cd+e}\right)}{c\sqrt{\frac{c(d+ex)}{cd+e}}}\right)}{2(d+ex)^{3/2}(a+b\operatorname{sech}^{-1}(cx))} + \frac{3e}{3e}$$

632

$$\frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(d^2\int\frac{1}{x\sqrt{1-cx}\sqrt{cx+1}\sqrt{d+ex}}dx-\frac{2de\sqrt{\frac{c(d+ex)}{cd+e}}\operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right),\frac{2e}{cd+e}\right)}{c\sqrt{d+ex}}-\frac{2e\sqrt{d+ex}E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right)\middle|\frac{2e}{cd+e}\right)}{c\sqrt{\frac{c(d+ex)}{cd+e}}}\right)}{2(d+ex)^{3/2}(a+b\operatorname{sech}^{-1}(cx))} + \frac{3e}{3e}$$

186

$$\frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-2d^2\int\frac{1}{cx\sqrt{cx+1}\sqrt{d+\frac{e}{c}-\frac{e(1-cx)}{c}}}d\sqrt{1-cx}-\frac{2de\sqrt{\frac{c(d+ex)}{cd+e}}\operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right),\frac{2e}{cd+e}\right)}{c\sqrt{d+ex}}-\frac{2e\sqrt{d+ex}E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right)\middle|\frac{2e}{cd+e}\right)}{c\sqrt{\frac{c(d+ex)}{cd+e}}}\right)}{2(d+ex)^{3/2}(a+b\operatorname{sech}^{-1}(cx))} + \frac{3e}{3e}$$

413

$$\begin{aligned}
 & 2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{2d^2\sqrt{1-\frac{e(1-cx)}{cd+e}} \int \frac{1}{cx\sqrt{cx+1}\sqrt{1-\frac{e(1-cx)}{cd+e}}} d\sqrt{1-cx}}{\sqrt{-\frac{e(1-cx)}{c}+\frac{e}{c}+d}} - \frac{2de\sqrt{\frac{c(d+ex)}{cd+e}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{c\sqrt{d+ex}} - \frac{2e\sqrt{d+ex}}{3e} \right) \\
 & \frac{2(d+ex)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e} \\
 & \quad \downarrow 412 \\
 & \frac{2(d+ex)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e} + \\
 & 2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{2d^2\sqrt{1-\frac{e(1-cx)}{cd+e}} \operatorname{EllipticPi}\left(2, \arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{\sqrt{-\frac{e(1-cx)}{c}+\frac{e}{c}+d}} - \frac{2de\sqrt{\frac{c(d+ex)}{cd+e}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{c\sqrt{d+ex}} - \frac{2e\sqrt{d+ex}}{3e} \right)
 \end{aligned}$$

input `Int[Sqrt[d + e*x]*(a + b*ArcSech[c*x]), x]`

output `(2*(d + e*x)^(3/2)*(a + b*ArcSech[c*x]))/(3*e) + (2*b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-2*e*Sqrt[d + e*x]*EllipticE[ArcSin[Sqrt[1 - c*x]/Sqrt[2]], (2*e)/(c*d + e)]/(c*Sqrt[(c*(d + e*x))/(c*d + e)]) - (2*d*e*Sqrt[(c*(d + e*x))/(c*d + e)]*EllipticF[ArcSin[Sqrt[1 - c*x]/Sqrt[2]], (2*e)/(c*d + e)]/(c*Sqrt[d + e*x]) - (2*d^2*Sqrt[1 - (e*(1 - c*x))/(c*d + e)]*EllipticPi[2, ArcSin[Sqrt[1 - c*x]/Sqrt[2]], (2*e)/(c*d + e)]/Sqrt[d + e/c - (e*(1 - c*x))/c]))/(3*e)`

### Defintions of rubi rules used

rule 186 `Int[1/(((a_.) + (b_.)*(x_))*Sqrt[(c_.) + (d_.)*(x_)]*Sqrt[(e_.) + (f_.)*(x_)]*Sqrt[(g_.) + (h_.)*(x_)]), x_] := Simp[-2 Subst[Int[1/(Simp[b*c - a*d - b*x^2, x]*Sqrt[Simp[(d*e - c*f)/d + f*(x^2/d), x]]*Sqrt[Simp[(d*g - c*h)/d + h*(x^2/d), x]]), x], x, Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x] && GtQ[(d*e - c*f)/d, 0]`

rule 321 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(1/(Sqrt[a]*Sqrt[c]*Rt[-d/c, 2]))*EllipticF[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-b/a, -d/c])`

rule 327 `Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[  
 (Sqrt[a]/(Sqrt[c]*Rt[-d/c, 2]))*EllipticE[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))  
 ], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]`

rule 412 `Int[1/(((a_) + (b_.)*(x_)^2)*Sqrt[(c_) + (d_.)*(x_)^2]*Sqrt[(e_) + (f_.)*(x  
 _)^2]), x_Symbol] := Simp[(1/(a*Sqrt[c]*Sqrt[e]*Rt[-d/c, 2]))*EllipticPi[b*  
 (c/(a*d)), ArcSin[Rt[-d/c, 2]*x], c*(f/(d*e))], x] /; FreeQ[{a, b, c, d, e,  
 f}, x] && !GtQ[d/c, 0] && GtQ[c, 0] && GtQ[e, 0] && !( !GtQ[f/e, 0] && S  
 implerSqrtQ[-f/e, -d/c])`

rule 413 `Int[1/(((a_) + (b_.)*(x_)^2)*Sqrt[(c_) + (d_.)*(x_)^2]*Sqrt[(e_) + (f_.)*(x  
 _)^2]), x_Symbol] := Simp[Sqrt[1 + (d/c)*x^2]/Sqrt[c + d*x^2] Int[1/((a +  
 b*x^2)*Sqrt[1 + (d/c)*x^2]*Sqrt[e + f*x^2]), x], x] /; FreeQ[{a, b, c, d,  
 e, f}, x] && !GtQ[c, 0]`

rule 508 `Int[Sqrt[(c_) + (d_.)*(x_)]/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := With[{q  
 = Rt[-b/a, 2]}, Simp[-2*(Sqrt[c + d*x]/(Sqrt[a]*q*Sqrt[q*((c + d*x)/(d + c  
 *q))])) Subst[Int[Sqrt[1 - 2*d*(x^2/(d + c*q))]/Sqrt[1 - x^2], x], x, Sqr  
 t[(1 - q*x)/2]], x]] /; FreeQ[{a, b, c, d}, x] && NegQ[b/a] && GtQ[a, 0]`

rule 511 `Int[1/(Sqrt[(c_) + (d_.)*(x_)]*Sqrt[(a_) + (b_.)*(x_)^2]), x_Symbol] := Wit  
 h[{q = Rt[-b/a, 2]}, Simp[-2*(Sqrt[q*((c + d*x)/(d + c*q))]/(Sqrt[a]*q*Sqrt  
 [c + d*x])) Subst[Int[1/(Sqrt[1 - 2*d*(x^2/(d + c*q))])*Sqrt[1 - x^2]), x]  
 , x, Sqrt[(1 - q*x)/2]], x]] /; FreeQ[{a, b, c, d}, x] && NegQ[b/a] && GtQ[  
 a, 0]`

rule 600 `Int[((A_.) + (B_.)*(x_))/(Sqrt[(c_) + (d_.)*(x_)]*Sqrt[(a_) + (b_.)*(x_)^2]  
 ), x_Symbol] := Simp[B/d Int[Sqrt[c + d*x]/Sqrt[a + b*x^2], x], x] - Simp  
 [(B*c - A*d)/d Int[1/(Sqrt[c + d*x]*Sqrt[a + b*x^2]), x], x] /; FreeQ[{a,  
 b, c, d, A, B}, x] && NegQ[b/a]`

```
rule 632 Int[1/((x_)*Sqrt[(c_) + (d_)*(x_)]*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] :
> With[{q = Rt[-b/a, 2]}, Simp[1/Sqrt[a] Int[1/(x*Sqrt[c + d*x]*Sqrt[1 -
q*x]*Sqrt[1 + q*x]), x], x]] /; FreeQ[{a, b, c, d}, x] && NegQ[b/a] && GtQ[
a, 0]
```

```
rule 634 Int[((c_) + (d_)*(x_))^(n_)/((x_)*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] :>
Simp[c^(n + 1/2) Int[1/(x*Sqrt[c + d*x]*Sqrt[a + b*x^2]), x], x] - Int[(
1/(Sqrt[c + d*x]*Sqrt[a + b*x^2]))*ExpandToSum[(c^(n + 1/2) - (c + d*x)^(n
+ 1/2))/x, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[n - 1/2, 0]
```

```
rule 6842 Int[((a_) + ArcSech[(c_)*(x_)]*(b_))*((d_) + (e_)*(x_))^(m_), x_Symbo
l] :> Simp[(d + e*x)^(m + 1)*((a + b*ArcSech[c*x])/(e*(m + 1))), x] + Simp[
b*(Sqrt[1 + c*x]/(e*(m + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x)^(m + 1)/(x*
Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[m, -1]
```

### Maple [A] (verified)

Time = 17.44 (sec) , antiderivative size = 413, normalized size of antiderivative = 1.48

method	result
derivativedivides	$\frac{2(ex+d)^{\frac{3}{2}}a}{3} + 2b \left( \frac{(ex+d)^{\frac{3}{2}} \operatorname{arcsech}(cx)}{3} - \frac{2e^2 \sqrt{-\frac{c(ex+d)+cd+e}{cex}} x \sqrt{-\frac{-c(ex+d)+cd-e}{cex}}}{3} \left( 2 \operatorname{EllipticF} \left( \sqrt{ex+d}, \sqrt{\frac{c}{cd+e}}, \sqrt{\frac{cd+e}{cd-e}} \right) \right) \right)$
default	$\frac{2(ex+d)^{\frac{3}{2}}a}{3} + 2b \left( \frac{(ex+d)^{\frac{3}{2}} \operatorname{arcsech}(cx)}{3} - \frac{2e^2 \sqrt{-\frac{c(ex+d)+cd+e}{cex}} x \sqrt{-\frac{-c(ex+d)+cd-e}{cex}}}{3} \left( 2 \operatorname{EllipticF} \left( \sqrt{ex+d}, \sqrt{\frac{c}{cd+e}}, \sqrt{\frac{cd+e}{cd-e}} \right) \right) \right)$
parts	$\frac{2a(ex+d)^{\frac{3}{2}}}{3e} + 2b \left( \frac{(ex+d)^{\frac{3}{2}} \operatorname{arcsech}(cx)}{3} - \frac{2e^2 \sqrt{-\frac{c(ex+d)-cd-e}{cxe}} x \sqrt{\frac{c(ex+d)-cd+e}{cxe}}}{3} \left( 2 \operatorname{EllipticF} \left( \sqrt{ex+d}, \sqrt{\frac{c}{cd+e}}, \sqrt{\frac{cd+e}{cd-e}} \right) \right) \right)$

```
input int((e*x+d)^(1/2)*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)
```

output

```
2/e*(1/3*(e*x+d)^(3/2)*a+b*(1/3*(e*x+d)^(3/2)*arcsech(c*x)-2/3*e^2*((-c*(e*x+d)+c*d+e)/c/e/x)^(1/2)*x*(-(-c*(e*x+d)+c*d-e)/c/e/x)^(1/2)*(2*EllipticF((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e)/(c*d-e))^(1/2))*c*d-EllipticE((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e)/(c*d-e))^(1/2))*c*d-EllipticPi((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),1/c*(c*d+e)/d,(c/(c*d-e))^(1/2)/(c/(c*d+e))^(1/2))*c*d-EllipticF((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e)/(c*d-e))^(1/2))*e+EllipticE((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e)/(c*d-e))^(1/2))*e)*((-c*(e*x+d)+c*d-e)/(c*d-e))^(1/2)*((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)/(c/(c*d+e))^(1/2)/(c^2*(e*x+d)^2-2*c^2*d*(e*x+d)+c^2*d^2-e^2)))
```

**Fricas [F(-1)]**

Timed out.

$$\int \sqrt{d+ex}(a+b\operatorname{sech}^{-1}(cx)) dx = \text{Timed out}$$

input

```
integrate((e*x+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="fricas")
```

output

Timed out

**Sympy [F]**

$$\int \sqrt{d+ex}(a+b\operatorname{sech}^{-1}(cx)) dx = \int (a+b\operatorname{asech}(cx))\sqrt{d+ex} dx$$

input

```
integrate((e*x+d)**(1/2)*(a+b*asech(c*x)),x)
```

output

```
Integral((a + b*asech(c*x))*sqrt(d + e*x), x)
```

**Maxima [F(-2)]**

Exception generated.

$$\int \sqrt{d+ex}(a+b\operatorname{sech}^{-1}(cx)) dx = \text{Exception raised: ValueError}$$

input `integrate((e*x+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e+c*d>0)', see `assume?` for more details)`

**Giac [F]**

$$\int \sqrt{d+ex}(a+b\operatorname{sech}^{-1}(cx)) dx = \int \sqrt{ex+d}(b\operatorname{ar}sech(cx) + a) dx$$

input `integrate((e*x+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate(sqrt(e*x + d)*(b*arcsech(c*x) + a), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{d+ex}(a+b\operatorname{sech}^{-1}(cx)) dx = \int \left(a + b\operatorname{acosh}\left(\frac{1}{cx}\right)\right) \sqrt{d+ex} dx$$

input `int((a + b*acosh(1/(c*x)))*(d + e*x)^(1/2),x)`

output `int((a + b*acosh(1/(c*x)))*(d + e*x)^(1/2), x)`



**Reduce [F]**

$$\int \sqrt{d+ex}(a+b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{2\sqrt{ex+d}ad + 2\sqrt{ex+d}aex + 3\left(\int \sqrt{ex+d} \operatorname{asech}(cx) dx\right) be}{3e}$$

input `int((e*x+d)^(1/2)*(a+b*asech(c*x)),x)`

output `(2*sqrt(d + e*x)*a*d + 2*sqrt(d + e*x)*a*e*x + 3*int(sqrt(d + e*x)*asech(c*x),x)*b*e)/(3*e)`

**3.82** 
$$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{\sqrt{d+ex}} dx$$

Optimal result	665
Mathematica [C] (warning: unable to verify)	666
Rubi [A] (verified)	667
Maple [A] (verified)	668
Fricas [F(-1)]	669
Sympy [F]	669
Maxima [F(-2)]	670
Giac [F]	670
Mupad [F(-1)]	670
Reduce [F]	671

**Optimal result**

Integrand size = 18, antiderivative size = 187

$$\begin{aligned} & \int \frac{a + b\operatorname{sech}^{-1}(cx)}{\sqrt{d + ex}} dx \\ &= \frac{2\sqrt{d + ex}(a + b\operatorname{sech}^{-1}(cx))}{e} \\ & \quad - \frac{4b\sqrt{\frac{1}{1+cx}}\sqrt{1 + cx}\sqrt{\frac{c(d+ex)}{cd+e}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{c\sqrt{d + ex}} \\ & \quad - \frac{4bd\sqrt{\frac{1}{1+cx}}\sqrt{1 + cx}\sqrt{\frac{c(d+ex)}{cd+e}} \operatorname{EllipticPi}\left(2, \arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{e\sqrt{d + ex}} \end{aligned}$$

output

```
2*(e*x+d)^(1/2)*(a+b*arcsech(c*x))/e-4*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(
c*(e*x+d)/(c*d+e))^(1/2)*EllipticF(1/2*(-c*x+1)^(1/2)*2^(1/2),2^(1/2)*(e/(
c*d+e))^(1/2))/c/(e*x+d)^(1/2)-4*b*d*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(c*(e
*x+d)/(c*d+e))^(1/2)*EllipticPi(1/2*(-c*x+1)^(1/2)*2^(1/2),2,2^(1/2)*(e/(c
*d+e))^(1/2))/e/(e*x+d)^(1/2)
```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 15.08 (sec) , antiderivative size = 1707, normalized size of antiderivative = 9.13

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex}} dx = \text{Too large to display}$$

input `Integrate[(a + b*ArcSech[c*x])/Sqrt[d + e*x],x]`

output

```
(2*a*Sqrt[d + e*x])/e + (2*b*Sqrt[d + e*x]*ArcSech[c*x])/e - ((4*I)*b*Sqrt
[(c*d + e + (c*d*(1 - c*x))/(1 + c*x) - (e*(1 - c*x))/(1 + c*x))/(c + (c*
1 - c*x)/(1 + c*x))]*(2*c*d*Sqrt[(-I)*(Sqrt[-(c*d) - e]*Sqrt[c*d - e] +
c*d*Sqrt[(1 - c*x)/(1 + c*x)] - e*Sqrt[(1 - c*x)/(1 + c*x)])]/(((-I)*c*d +
Sqrt[-(c*d) - e]*Sqrt[c*d - e] + I*e)*(-I + Sqrt[(1 - c*x)/(1 + c*x)])))*
Sqrt[(-I)*(Sqrt[-(c*d) - e]*Sqrt[c*d - e] - c*d*Sqrt[(1 - c*x)/(1 + c*x)]
+ e*Sqrt[(1 - c*x)/(1 + c*x)])]/((I*c*d + Sqrt[-(c*d) - e]*Sqrt[c*d - e]
- I*e)*(-I + Sqrt[(1 - c*x)/(1 + c*x)]))]*(1 + (1 - c*x)/(1 + c*x))*Ellipt
icF[ArcSin[Sqrt[((Sqrt[-(c*d) - e] - I*Sqrt[c*d - e])*(I + Sqrt[(1 - c*x)/
(1 + c*x)])]/((Sqrt[-(c*d) - e] + I*Sqrt[c*d - e])*(-I + Sqrt[(1 - c*x)/(1
+ c*x)]))]], (Sqrt[-(c*d) - e] + I*Sqrt[c*d - e])^2/(Sqrt[-(c*d) - e] - I
*Sqrt[c*d - e])^2] + (c*d - e)*Sqrt[((Sqrt[-(c*d) - e] - I*Sqrt[c*d - e])*(
I + Sqrt[(1 - c*x)/(1 + c*x)]))]/((Sqrt[-(c*d) - e] + I*Sqrt[c*d - e])*(-I
+ Sqrt[(1 - c*x)/(1 + c*x)]))]*Sqrt[1 + (1 - c*x)/(1 + c*x)]*Sqrt[(e - (e
*(1 - c*x))/(1 + c*x) + c*d*(1 + (1 - c*x)/(1 + c*x)))/(c*d + e)]*Elliptic
F[I*ArcSinh[Sqrt[(1 - c*x)/(1 + c*x)]], (c*d - e)/(c*d + e)] + (2*I)*c*d*S
qrt[(-I)*(Sqrt[-(c*d) - e]*Sqrt[c*d - e] + c*d*Sqrt[(1 - c*x)/(1 + c*x)]
- e*Sqrt[(1 - c*x)/(1 + c*x)])]/(((-I)*c*d + Sqrt[-(c*d) - e]*Sqrt[c*d - e
] + I*e)*(-I + Sqrt[(1 - c*x)/(1 + c*x)])))*Sqrt[(-I)*(Sqrt[-(c*d) - e]*S
qrt[c*d - e] - c*d*Sqrt[(1 - c*x)/(1 + c*x)] + e*Sqrt[(1 - c*x)/(1 + c*...
```

**Rubi [A] (verified)**

Time = 0.48 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.91, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {6842, 637, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex}} dx \\
 & \quad \downarrow \text{6842} \\
 & \frac{2b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{\sqrt{d+ex}}{x \sqrt{1-c^2x^2}} dx}{e} + \frac{2\sqrt{d+ex}(a + b \operatorname{sech}^{-1}(cx))}{e} \\
 & \quad \downarrow \text{637} \\
 & \frac{2b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \left( \frac{d}{x \sqrt{d+ex} \sqrt{1-c^2x^2}} + \frac{e}{\sqrt{d+ex} \sqrt{1-c^2x^2}} \right) dx}{e} + \frac{2\sqrt{d+ex}(a + b \operatorname{sech}^{-1}(cx))}{e} \\
 & \quad \downarrow \text{2009} \\
 & \frac{2\sqrt{d+ex}(a + b \operatorname{sech}^{-1}(cx))}{e} + \\
 & \frac{2b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \left( -\frac{2e \sqrt{\frac{c(d+ex)}{cd+e}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{c \sqrt{d+ex}} - \frac{2d \sqrt{\frac{c(d+ex)}{cd+e}} \operatorname{EllipticPi}\left(2, \arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{\sqrt{d+ex}} \right)}{e}
 \end{aligned}$$

input `Int[(a + b*ArcSech[c*x])/Sqrt[d + e*x],x]`

output `(2*Sqrt[d + e*x]*(a + b*ArcSech[c*x]))/e + (2*b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-2*e*Sqrt[(c*(d + e*x))/(c*d + e)]*EllipticF[ArcSin[Sqrt[1 - c*x]/Sqrt[2]], (2*e)/(c*d + e)])/(c*Sqrt[d + e*x]) - (2*d*Sqrt[(c*(d + e*x))/(c*d + e)]*EllipticPi[2, ArcSin[Sqrt[1 - c*x]/Sqrt[2]], (2*e)/(c*d + e)]/Sqrt[d + e*x]))/e`

Defintions of rubi rules used

```
rule 637 Int[(x_)^(m_.)*((c_) + (d_.)*(x_))^(n_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol]
  :=> Int[ExpandIntegrand[(a + b*x^2)^p/Sqrt[c + d*x], x^m*(c + d*x)^(n + 1/2), x], x]
  /; FreeQ[{a, b, c, d, m}, x] && IntegerQ[p + 1/2] && IntegerQ[n + 1/2] && IntegerQ[m]
```

```
rule 2009 Int[u_, x_Symbol] :=> Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6842 Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol]
  :=> Simp[(d + e*x)^(m + 1)*((a + b*ArcSech[c*x])/(e*(m + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(e*(m + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x)^(m + 1)/(x*Sqrt[1 - c^2*x^2]), x], x]
  /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[m, -1]
```

Maple [A] (verified)

Time = 17.13 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.53

method	result
derivativedivides	$2\sqrt{ex+d}a+2b \left( \sqrt{ex+d} \operatorname{arcsech}(cx) - \frac{2ce^2 \sqrt{-\frac{c(ex+d)+cd+e}{cex}} x \sqrt{-\frac{c(ex+d)+cd-e}{cex}} \left( \operatorname{EllipticF}\left(\sqrt{ex+d} \sqrt{\frac{c}{cd+e}}, \sqrt{\frac{cd+e}{cd-e}}\right) - \operatorname{EllipticF}\left(\sqrt{ex+d} \sqrt{\frac{c}{cd+e}}, \sqrt{\frac{cd+e}{cd-e}}\right) \right)}{(c^2(ex+d)^2 - 2c^2d(ex+d) + e^2)}$
default	$2\sqrt{ex+d}a+2b \left( \sqrt{ex+d} \operatorname{arcsech}(cx) - \frac{2ce^2 \sqrt{-\frac{c(ex+d)+cd+e}{cex}} x \sqrt{-\frac{c(ex+d)+cd-e}{cex}} \left( \operatorname{EllipticF}\left(\sqrt{ex+d} \sqrt{\frac{c}{cd+e}}, \sqrt{\frac{cd+e}{cd-e}}\right) - \operatorname{EllipticF}\left(\sqrt{ex+d} \sqrt{\frac{c}{cd+e}}, \sqrt{\frac{cd+e}{cd-e}}\right) \right)}{(c^2(ex+d)^2 - 2c^2d(ex+d) + e^2)} \right)$
parts	$\frac{2a\sqrt{ex+d}}{e} + \frac{2b \left( \sqrt{ex+d} \operatorname{arcsech}(cx) - \frac{2ce^2 \sqrt{-\frac{c(ex+d)-cd-e}{cxe}} x \sqrt{\frac{c(ex+d)-cd+e}{cxe}} \left( \operatorname{EllipticF}\left(\sqrt{ex+d} \sqrt{\frac{c}{cd+e}}, \sqrt{\frac{cd+e}{cd-e}}\right) - \operatorname{EllipticF}\left(\sqrt{ex+d} \sqrt{\frac{c}{cd+e}}, \sqrt{\frac{cd+e}{cd-e}}\right) \right)}{\sqrt{\frac{c}{cd+e}} (c^2(ex+d)^2 - 2c^2d(ex+d) + e^2)} \right)}{e}$

```
input int((a+b*arcsech(c*x))/(e*x+d)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
2/e*((e*x+d)^(1/2)*a+b*((e*x+d)^(1/2)*arcsech(c*x)-2*c*e^2*((-c*(e*x+d)+c*d+e)/c/e/x)^(1/2)*x*(-(-c*(e*x+d)+c*d-e)/c/e/x)^(1/2)*(EllipticF((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e)/(c*d-e))^(1/2))-EllipticPi((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),1/c*(c*d+e)/d,(c/(c*d-e))^(1/2)/(c/(c*d+e))^(1/2)))*((-c*(e*x+d)+c*d-e)/(c*d-e))^(1/2)*((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)/(c^2*(e*x+d)^2-2*c^2*d*(e*x+d)+c^2*d^2-e^2)/(c/(c*d+e))^(1/2)))
```

**Fricas [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex}} dx = \text{Timed out}$$

input

```
integrate((a+b*arcsech(c*x))/(e*x+d)^(1/2),x, algorithm="fricas")
```

output

Timed out

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex}} dx = \int \frac{a + b \operatorname{asech}(cx)}{\sqrt{d + ex}} dx$$

input

```
integrate((a+b*asech(c*x))/(e*x+d)**(1/2),x)
```

output

```
Integral((a + b*asech(c*x))/sqrt(d + e*x), x)
```

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex}} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/(e*x+d)^(1/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation \*may\* help (example of legal syntax is 'assume(c\*d-e>0)', see `assume?` for more details)

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex}} dx = \int \frac{b \operatorname{ar} \operatorname{sech}(cx) + a}{\sqrt{ex + d}} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/sqrt(e*x + d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{\sqrt{d + ex}} dx$$

input `int((a + b*acosh(1/(c*x)))/(d + e*x)^(1/2),x)`

output `int((a + b*acosh(1/(c*x)))/(d + e*x)^(1/2), x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex}} dx = \frac{2\sqrt{ex + d} a + \left( \int \frac{\operatorname{asech}(cx)}{\sqrt{ex+d}} dx \right) b e}{e}$$

input `int((a+b*asech(c*x))/(e*x+d)^(1/2),x)`

output `(2*sqrt(d + e*x)*a + int(asech(c*x)/sqrt(d + e*x),x)*b*e)/e`



**3.83** 
$$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex)^{3/2}} dx$$

Optimal result	672
Mathematica [C] (warning: unable to verify)	672
Rubi [A] (verified)	673
Maple [B] (verified)	675
Fricas [F]	676
Sympy [F]	677
Maxima [F(-2)]	677
Giac [F]	677
Mupad [F(-1)]	678
Reduce [F]	678

**Optimal result**

Integrand size = 18, antiderivative size = 105

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{(d + ex)^{3/2}} dx = -\frac{2(a + b\operatorname{sech}^{-1}(cx))}{e\sqrt{d + ex}} + \frac{4b\sqrt{\frac{1}{1+cx}}\sqrt{1 + cx}\sqrt{\frac{c(d+ex)}{cd+e}} \operatorname{EllipticPi}\left(2, \arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{e\sqrt{d + ex}}$$

output

```
(-2*a-2*b*arcsech(c*x))/e/(e*x+d)^(1/2)+4*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)
)*(c*(e*x+d)/(c*d+e))^(1/2)*EllipticPi(1/2*(-c*x+1)^(1/2)*2^(1/2),2,2^(1/2)
)*(e/(c*d+e))^(1/2))/e/(e*x+d)^(1/2)
```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 14.37 (sec) , antiderivative size = 1675, normalized size of antiderivative = 15.95

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{(d + ex)^{3/2}} dx = \text{Too large to display}$$

input

```
Integrate[(a + b*ArcSech[c*x])/(d + e*x)^(3/2),x]
```

output

```

(-2*a)/(e*Sqrt[d + e*x]) - (2*b*ArcSech[c*x])/(e*Sqrt[d + e*x]) + ((4*I)*b
*(2*Sqrt[(-I)*(Sqrt[-(c*d) - e]*Sqrt[c*d - e] + c*d*Sqrt[(1 - c*x)/(1 + c
*x)] - e*Sqrt[(1 - c*x)/(1 + c*x)])]/((-I)*c*d + Sqrt[-(c*d) - e]*Sqrt[c*d
d - e] + I*e)*(-I + Sqrt[(1 - c*x)/(1 + c*x)])))*Sqrt[(-I)*(Sqrt[-(c*d) -
e]*Sqrt[c*d - e] - c*d*Sqrt[(1 - c*x)/(1 + c*x)] + e*Sqrt[(1 - c*x)/(1 +
c*x)])]/((I*c*d + Sqrt[-(c*d) - e]*Sqrt[c*d - e] - I*e)*(-I + Sqrt[(1 - c*
x)/(1 + c*x)]))]*(1 + (1 - c*x)/(1 + c*x))*EllipticF[ArcSin[Sqrt[((Sqrt[-(
c*d) - e] - I*Sqrt[c*d - e])*(I + Sqrt[(1 - c*x)/(1 + c*x)]))]/((Sqrt[-(c*d
) - e] + I*Sqrt[c*d - e])*(-I + Sqrt[(1 - c*x)/(1 + c*x)]))]], (Sqrt[-(c*d
) - e] + I*Sqrt[c*d - e])^2/(Sqrt[-(c*d) - e] - I*Sqrt[c*d - e])^2 + Sqrt
[((Sqrt[-(c*d) - e] - I*Sqrt[c*d - e])*(I + Sqrt[(1 - c*x)/(1 + c*x)]))]/((
Sqrt[-(c*d) - e] + I*Sqrt[c*d - e])*(-I + Sqrt[(1 - c*x)/(1 + c*x)])))*Sqr
t[1 + (1 - c*x)/(1 + c*x)]*Sqrt[(e - (e*(1 - c*x))/(1 + c*x) + c*d*(1 + (1
- c*x)/(1 + c*x)))/(c*d + e)]*EllipticF[I*ArcSinh[Sqrt[(1 - c*x)/(1 + c*x
)]]], (c*d - e)/(c*d + e)] + (2*I)*Sqrt[(-I)*(Sqrt[-(c*d) - e]*Sqrt[c*d -
e] + c*d*Sqrt[(1 - c*x)/(1 + c*x)] - e*Sqrt[(1 - c*x)/(1 + c*x)])]/((-I)*
c*d + Sqrt[-(c*d) - e]*Sqrt[c*d - e] + I*e)*(-I + Sqrt[(1 - c*x)/(1 + c*x
)]])*Sqrt[(-I)*(Sqrt[-(c*d) - e]*Sqrt[c*d - e] - c*d*Sqrt[(1 - c*x)/(1 +
c*x)] + e*Sqrt[(1 - c*x)/(1 + c*x)])]/((I*c*d + Sqrt[-(c*d) - e]*Sqrt[c*d
- e] - I*e)*(-I + Sqrt[(1 - c*x)/(1 + c*x)]))]*(1 + (1 - c*x)/(1 + c*x)...

```

### Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.17, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$ , Rules used = {6842, 632, 186, 413, 412}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{3/2}} dx \\
 & \quad \downarrow 6842 \\
 & -\frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{x\sqrt{d+ex}\sqrt{1-c^2x^2}} dx}{e} - \frac{2(a + b \operatorname{sech}^{-1}(cx))}{e\sqrt{d+ex}} \\
 & \quad \downarrow 632
 \end{aligned}$$

$$\begin{aligned}
& \frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{x\sqrt{1-cx}\sqrt{cx+1}\sqrt{d+ex}} dx}{e} - \frac{2(a + b\operatorname{sech}^{-1}(cx))}{e\sqrt{d+ex}} \\
& \quad \downarrow 186 \\
& \frac{4b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{cx\sqrt{cx+1}\sqrt{d+\frac{e}{c}-\frac{e(1-cx)}{c}}} d\sqrt{1-cx}}{e} - \frac{2(a + b\operatorname{sech}^{-1}(cx))}{e\sqrt{d+ex}} \\
& \quad \downarrow 413 \\
& \frac{4b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\sqrt{1-\frac{e(1-cx)}{cd+e}} \int \frac{1}{cx\sqrt{cx+1}\sqrt{1-\frac{e(1-cx)}{cd+e}}} d\sqrt{1-cx}}{e\sqrt{-\frac{e(1-cx)}{c} + \frac{e}{c} + d}} - \frac{2(a + b\operatorname{sech}^{-1}(cx))}{e\sqrt{d+ex}} \\
& \quad \downarrow 412 \\
& \frac{4b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\sqrt{1-\frac{e(1-cx)}{cd+e}} \operatorname{EllipticPi}\left(2, \arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{e\sqrt{-\frac{e(1-cx)}{c} + \frac{e}{c} + d}} - \frac{2(a + b\operatorname{sech}^{-1}(cx))}{e\sqrt{d+ex}}
\end{aligned}$$

input `Int[(a + b*ArcSech[c*x])/(d + e*x)^(3/2), x]`

output `(-2*(a + b*ArcSech[c*x]))/(e*Sqrt[d + e*x]) + (4*b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*Sqrt[1 - (e*(1 - c*x))/(c*d + e)]*EllipticPi[2, ArcSin[Sqrt[1 - c*x]/Sqrt[2]], (2*e)/(c*d + e)])/(e*Sqrt[d + e/c - (e*(1 - c*x))/c])`

### Defintions of rubi rules used

rule 186 `Int[1/(((a_.) + (b_.)*(x_))*Sqrt[(c_.) + (d_.)*(x_)]*Sqrt[(e_.) + (f_.)*(x_)]*Sqrt[(g_.) + (h_.)*(x_)]), x_] := Simp[-2 Subst[Int[1/(Simp[b*c - a*d - b*x^2, x]*Sqrt[Simp[(d*e - c*f)/d + f*(x^2/d), x]]*Sqrt[Simp[(d*g - c*h)/d + h*(x^2/d), x]]), x], x, Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x] && GtQ[(d*e - c*f)/d, 0]`

rule 412 `Int[1/(((a_) + (b_)*(x_)^2)*Sqrt[(c_) + (d_)*(x_)^2]*Sqrt[(e_) + (f_)*(x_)^2]), x_Symbol] := Simp[(1/(a*Sqrt[c]*Sqrt[e]*Rt[-d/c, 2]))*EllipticPi[b*(c/(a*d)), ArcSin[Rt[-d/c, 2]*x], c*(f/(d*e))], x] /; FreeQ[{a, b, c, d, e, f}, x] && !GtQ[d/c, 0] && GtQ[c, 0] && GtQ[e, 0] && !( !GtQ[f/e, 0] && SimplerSqrtQ[-f/e, -d/c])`

rule 413 `Int[1/(((a_) + (b_)*(x_)^2)*Sqrt[(c_) + (d_)*(x_)^2]*Sqrt[(e_) + (f_)*(x_)^2]), x_Symbol] := Simp[Sqrt[1 + (d/c)*x^2]/Sqrt[c + d*x^2] Int[1/((a + b*x^2)*Sqrt[1 + (d/c)*x^2]*Sqrt[e + f*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && !GtQ[c, 0]`

rule 632 `Int[1/((x_)*Sqrt[(c_) + (d_)*(x_)]*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] :> With[{q = Rt[-b/a, 2]}, Simp[1/Sqrt[a] Int[1/(x*Sqrt[c + d*x]*Sqrt[1 - q*x]*Sqrt[1 + q*x]), x], x]] /; FreeQ[{a, b, c, d}, x] && NegQ[b/a] && GtQ[a, 0]`

rule 6842 `Int[((a_) + ArcSech[(c_)*(x_)])*(b_))*((d_) + (e_)*(x_)^(m_)), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*ArcSech[c*x])/(e*(m + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(e*(m + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x)^(m + 1)/(x*Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[m, -1]`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 250 vs.  $2(98) = 196$ .

Time = 16.98 (sec) , antiderivative size = 251, normalized size of antiderivative = 2.39

method	result
derivativedivides	$-\frac{2a}{\sqrt{ex+d}} + 2b \left( -\frac{\operatorname{arcsech}(cx)}{\sqrt{ex+d}} - \frac{2ce^2 \sqrt{\frac{-c(ex+d)+cd+e}{ce}} x \sqrt{\frac{-c(ex+d)+cd-e}{ce}} \operatorname{EllipticPi}\left(\sqrt{ex+d} \sqrt{\frac{c}{cd+e}}, \frac{cd+e}{cd}, \sqrt{\frac{c}{cd+e}}\right)}{d \sqrt{\frac{c}{cd+e}} (c^2(ex+d)^2 - 2c^2d(ex+d) + c^2d^2 - e^2)} \right)$
default	$-\frac{2a}{\sqrt{ex+d}} + 2b \left( -\frac{\operatorname{arcsech}(cx)}{\sqrt{ex+d}} - \frac{2ce^2 \sqrt{\frac{-c(ex+d)+cd+e}{ce}} x \sqrt{\frac{-c(ex+d)+cd-e}{ce}} \operatorname{EllipticPi}\left(\sqrt{ex+d} \sqrt{\frac{c}{cd+e}}, \frac{cd+e}{cd}, \sqrt{\frac{c}{cd+e}}\right)}{d \sqrt{\frac{c}{cd+e}} (c^2(ex+d)^2 - 2c^2d(ex+d) + c^2d^2 - e^2)} \right)$
parts	$-\frac{2a}{\sqrt{ex+d}e} + 2b \left( -\frac{\operatorname{arcsech}(cx)}{\sqrt{ex+d}} - \frac{2ce^2 \sqrt{\frac{c(ex+d)-cd-e}{cxe}} x \sqrt{\frac{c(ex+d)-cd+e}{cxe}} \operatorname{EllipticPi}\left(\sqrt{ex+d} \sqrt{\frac{e}{cd+e}}, \frac{cd+e}{cd}, \sqrt{\frac{c}{cd+e}}\right)}{d \sqrt{\frac{c}{cd+e}} (c^2(ex+d)^2 - 2c^2d(ex+d) + c^2d^2 - e^2)} \right)$

```
input int((a+b*arcsech(c*x))/(e*x+d)^(3/2), x, method=_RETURNVERBOSE)
```

```
output 2/e*(-a/(e*x+d)^(1/2)+b*(-1/(e*x+d)^(1/2)*arcsech(c*x)-2*c*e^2*((-c*(e*x+d)+c*d+e)/c/e/x)^(1/2)*x*(-(-c*(e*x+d)+c*d-e)/c/e/x)^(1/2)*EllipticPi((e*x+d)^(1/2)*(c/(c*d+e))^(1/2), 1/c*(c*d+e)/d, (c/(c*d-e))^(1/2)/(c/(c*d+e))^(1/2))*((-c*(e*x+d)+c*d-e)/(c*d-e))^(1/2)*((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)/d/(c/(c*d+e))^(1/2)/(c^2*(e*x+d)^2-2*c^2*d*(e*x+d)+c^2*d^2-e^2)))
```

**Fricas [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{3/2}} dx = \int \frac{b \operatorname{ar} \operatorname{sech}(cx) + a}{(ex + d)^{\frac{3}{2}}} dx$$

```
input integrate((a+b*arcsech(c*x))/(e*x+d)^(3/2), x, algorithm="fricas")
```

```
output integral(sqrt(e*x + d)*(b*arcsech(c*x) + a)/(e^2*x^2 + 2*d*e*x + d^2), x)
```

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{3/2}} dx = \int \frac{a + b \operatorname{arsech}(cx)}{(d + ex)^{\frac{3}{2}}} dx$$

input `integrate((a+b*asech(c*x))/(e*x+d)**(3/2), x)`

output `Integral((a + b*asech(c*x))/(d + e*x)**(3/2), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/(e*x+d)^(3/2), x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e+c*d>0)', see `assume?` for more details)`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{3/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex + d)^{\frac{3}{2}}} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x+d)^(3/2), x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/(e*x + d)^(3/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{(d + ex)^{3/2}} dx$$

input `int((a + b*acosh(1/(c*x)))/(d + e*x)^(3/2), x)`output `int((a + b*acosh(1/(c*x)))/(d + e*x)^(3/2), x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{3/2}} dx = \frac{\sqrt{ex + d} \left( \int \frac{a \operatorname{sech}(cx)}{\sqrt{ex+d}d + \sqrt{ex+d}ex} dx \right) be - 2a}{\sqrt{ex + d} e}$$

input `int((a+b*asech(c*x))/(e*x+d)^(3/2), x)`output `(sqrt(d + e*x)*int(asech(c*x)/(sqrt(d + e*x)*d + sqrt(d + e*x)*e*x), x)*b*e - 2*a)/(sqrt(d + e*x)*e)`

**3.84**  $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex)^{5/2}} dx$

Optimal result	679
Mathematica [C] (warning: unable to verify)	680
Rubi [A] (verified)	681
Maple [B] (verified)	685
Fricas [F]	687
Sympy [F]	687
Maxima [F(-2)]	687
Giac [F]	688
Mupad [F(-1)]	688
Reduce [F]	688

**Optimal result**

Integrand size = 18, antiderivative size = 278

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{(d + ex)^{5/2}} dx = \frac{4be\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{3d(c^2d^2 - e^2)\sqrt{d+ex}} - \frac{2(a + b\operatorname{sech}^{-1}(cx))}{3e(d + ex)^{3/2}}$$

$$- \frac{4bc\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{d+ex}E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right) \mid \frac{2e}{cd+e}\right)}{3d(c^2d^2 - e^2)\sqrt{\frac{c(d+ex)}{cd+e}}}$$

$$+ \frac{4b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{\frac{c(d+ex)}{cd+e}}\operatorname{EllipticPi}\left(2, \arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{3de\sqrt{d+ex}}$$

output

```
4/3*b*e*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/d/(c^2*d^2-e^2)
/(e*x+d)^(1/2)-2/3*(a+b*arcsech(c*x))/e/(e*x+d)^(3/2)-4/3*b*c*(1/(c*x+1))^(
1/2)*(c*x+1)^(1/2)*(e*x+d)^(1/2)*EllipticE(1/2*(-c*x+1)^(1/2)*2^(1/2),2^(
1/2)*(e/(c*d+e))^(1/2))/d/(c^2*d^2-e^2)/(c*(e*x+d)/(c*d+e))^(1/2)+4/3*b*(1
/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(c*(e*x+d)/(c*d+e))^(1/2)*EllipticPi(1/2*(-c
*x+1)^(1/2)*2^(1/2),2^(1/2)*(e/(c*d+e))^(1/2))/d/e/(e*x+d)^(1/2)
```



**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 24.33 (sec) , antiderivative size = 4527, normalized size of antiderivative = 16.28

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{5/2}} dx = \text{Result too large to show}$$

input `Integrate[(a + b*ArcSech[c*x])/(d + e*x)^(5/2),x]`

output

```
(-2*a)/(3*e*(d + e*x)^(3/2)) + Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[d + e*x]*((4
*b*c)/(3*d*(c^2*d^2 - e^2)) - (4*b)/(3*d*(c*d + e)*(d + e*x))) - (2*b*ArcS
ech[c*x])/(3*e*(d + e*x)^(3/2)) - (4*b*((e*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[
c*(1 + (1 - c*x)/(1 + c*x))]*(c*d + e + (c*d*(1 - c*x))/(1 + c*x)) - (e*(1
- c*x))/(1 + c*x)))/((1 + (1 - c*x)/(1 + c*x))*Sqrt[c + (c*(1 - c*x))/(1 +
c*x)]*Sqrt[(c*d + e + (c*d*(1 - c*x))/(1 + c*x)) - (e*(1 - c*x))/(1 + c*x)
])/((c + (c*(1 - c*x))/(1 + c*x))) - ((c*d - e)*Sqrt[c*(1 + (1 - c*x)/(1 +
c*x))]*Sqrt[c*(1 + (1 - c*x)/(1 + c*x))*(c*d + e + (c*d*(1 - c*x))/(1 + c
x)) - (e*(1 - c*x))/(1 + c*x)))*((I*(-(c*d) - e)*e*Sqrt[1 + (1 - c*x)/(1 +
c*x)]*Sqrt[1 - ((c*d - e)*(1 - c*x))/((-c*d) - e)*(1 + c*x)]]*(EllipticE[
I*ArcSinh[Sqrt[(1 - c*x)/(1 + c*x)]], -((c*d - e)/(-(c*d) - e))] - Ellipti
cF[I*ArcSinh[Sqrt[(1 - c*x)/(1 + c*x)]], -((c*d - e)/(-(c*d) - e))])/((c*
d - e)*Sqrt[c*(1 + (1 - c*x)/(1 + c*x))*(c*d + e + ((c*d - e)*(1 - c*x))/(
1 + c*x))]) + (I*c*d*Sqrt[1 + (1 - c*x)/(1 + c*x)]*Sqrt[1 - ((c*d - e)*(1
- c*x))/((-c*d) - e)*(1 + c*x)]]*EllipticF[I*ArcSinh[Sqrt[(1 - c*x)/(1 +
c*x)]], -((c*d - e)/(-(c*d) - e))])/Sqrt[c*(1 + (1 - c*x)/(1 + c*x))*(c*d
+ e + ((c*d - e)*(1 - c*x))/(1 + c*x))] + (I*e*Sqrt[1 + (1 - c*x)/(1 + c*x
)]*Sqrt[1 - ((c*d - e)*(1 - c*x))/((-c*d) - e)*(1 + c*x)]]*EllipticF[I*Ar
cSinh[Sqrt[(1 - c*x)/(1 + c*x)]], -((c*d - e)/(-(c*d) - e))])/Sqrt[c*(1 +
(1 - c*x)/(1 + c*x))*(c*d + e + ((c*d - e)*(1 - c*x))/(1 + c*x))] - (I*...
```

**Rubi [A] (verified)**

Time = 0.58 (sec) , antiderivative size = 254, normalized size of antiderivative = 0.91, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {6842, 635, 25, 27, 498, 27, 508, 327, 632, 186, 413, 412}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{5/2}} dx \\
 & \quad \downarrow \text{6842} \\
 & - \frac{2b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{1}{x(d+ex)^{3/2} \sqrt{1-c^2x^2}} dx}{3e} - \frac{2(a + b \operatorname{sech}^{-1}(cx))}{3e(d+ex)^{3/2}} \\
 & \quad \downarrow \text{635} \\
 & - \frac{2b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \left( \int -\frac{e}{d(d+ex)^{3/2} \sqrt{1-c^2x^2}} dx + \frac{\int \frac{1}{x \sqrt{d+ex} \sqrt{1-c^2x^2}} dx}{d} \right)}{3e} - \frac{2(a + b \operatorname{sech}^{-1}(cx))}{3e(d+ex)^{3/2}} \\
 & \quad \downarrow \text{25} \\
 & - \frac{2b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \left( \frac{\int \frac{1}{x \sqrt{d+ex} \sqrt{1-c^2x^2}} dx}{d} - \int \frac{e}{d(d+ex)^{3/2} \sqrt{1-c^2x^2}} dx \right)}{3e} - \frac{2(a + b \operatorname{sech}^{-1}(cx))}{3e(d+ex)^{3/2}} \\
 & \quad \downarrow \text{27} \\
 & - \frac{2b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \left( \frac{\int \frac{1}{x \sqrt{d+ex} \sqrt{1-c^2x^2}} dx}{d} - \frac{e \int \frac{1}{(d+ex)^{3/2} \sqrt{1-c^2x^2}} dx}{d} \right)}{3e} - \frac{2(a + b \operatorname{sech}^{-1}(cx))}{3e(d+ex)^{3/2}} \\
 & \quad \downarrow \text{498} \\
 & - \frac{2b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \left( \frac{\int \frac{1}{x \sqrt{d+ex} \sqrt{1-c^2x^2}} dx}{d} - \frac{e \left( \frac{2e \sqrt{1-c^2x^2}}{(c^2d^2 - e^2) \sqrt{d+ex}} - \frac{2c^2 \int -\frac{\sqrt{d+ex}}{2\sqrt{1-c^2x^2}} dx}{c^2d^2 - e^2} \right)}{d} \right)}{3e} \\
 & \quad \downarrow \text{27} \\
 & \frac{2(a + b \operatorname{sech}^{-1}(cx))}{3e(d+ex)^{3/2}}
 \end{aligned}$$

$$2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{1}{x\sqrt{d+ex}\sqrt{1-c^2x^2}} dx}{d} - \frac{e \left( \frac{c^2 \int \frac{\sqrt{d+ex}}{\sqrt{1-c^2x^2}} dx}{c^2d^2-e^2} + \frac{2e\sqrt{1-c^2x^2}}{(c^2d^2-e^2)\sqrt{d+ex}} \right)}{d} \right)$$

$$\frac{3e}{2(a+b\operatorname{sech}^{-1}(cx))} \frac{1}{3e(d+ex)^{3/2}}$$

↓ 508

$$2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{1}{x\sqrt{d+ex}\sqrt{1-c^2x^2}} dx}{d} - \frac{e \left( \frac{2e\sqrt{1-c^2x^2}}{(c^2d^2-e^2)\sqrt{d+ex}} - \frac{2c\sqrt{d+ex} \int \frac{\sqrt{1-\frac{e(1-cx)}{cd+e}} d\sqrt{1-cx}}{\sqrt{\frac{1}{2}(cx-1)+1} \sqrt{2}}}{(c^2d^2-e^2)\sqrt{\frac{c(d+ex)}{cd+e}}} \right)}{d} \right)$$

$$\frac{3e}{2(a+b\operatorname{sech}^{-1}(cx))} \frac{1}{3e(d+ex)^{3/2}}$$

↓ 327

$$2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{1}{x\sqrt{d+ex}\sqrt{1-c^2x^2}} dx}{d} - \frac{e \left( \frac{2e\sqrt{1-c^2x^2}}{(c^2d^2-e^2)\sqrt{d+ex}} - \frac{2c\sqrt{d+ex} E \left( \arcsin \left( \frac{\sqrt{1-cx}}{\sqrt{2}} \right) \middle| \frac{2e}{cd+e} \right)}{(c^2d^2-e^2)\sqrt{\frac{c(d+ex)}{cd+e}}} \right)}{d} \right)$$

$$\frac{3e}{2(a+b\operatorname{sech}^{-1}(cx))} \frac{1}{3e(d+ex)^{3/2}}$$

↓ 632

$$2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{1}{x\sqrt{1-cx}\sqrt{cx+1}\sqrt{d+ex}} dx}{d} - \frac{e \left( \frac{2e\sqrt{1-c^2x^2}}{(c^2d^2-e^2)\sqrt{d+ex}} - \frac{2c\sqrt{d+ex} E \left( \arcsin \left( \frac{\sqrt{1-cx}}{\sqrt{2}} \right) \middle| \frac{2e}{cd+e} \right)}{(c^2d^2-e^2)\sqrt{\frac{c(d+ex)}{cd+e}}} \right)}{d} \right)$$

$$\frac{3e}{2(a+b\operatorname{sech}^{-1}(cx))} \frac{1}{3e(d+ex)^{3/2}}$$

↓ 186

$$\begin{aligned}
 & 2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{2\int\frac{1}{cx\sqrt{cx+1}\sqrt{d+\frac{e}{c}-\frac{e(1-cx)}{c}}}d\sqrt{1-cx}}{d} - \frac{e\left(\frac{2e\sqrt{1-c^2x^2}}{(c^2d^2-e^2)\sqrt{d+ex}} - \frac{2c\sqrt{d+ex}E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right)\middle|\frac{2e}{cd+e}\right)}{(c^2d^2-e^2)\sqrt{\frac{c(d+ex)}{cd+e}}}\right)}{d} \right) \\
 & \frac{2(a+b\operatorname{sech}^{-1}(cx))}{3e(d+ex)^{3/2}} \\
 & \quad \downarrow 413 \\
 & 2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{2\sqrt{1-\frac{e(1-cx)}{cd+e}}\int\frac{1}{cx\sqrt{cx+1}\sqrt{1-\frac{e(1-cx)}{cd+e}}}d\sqrt{1-cx}}{d\sqrt{-\frac{e(1-cx)}{c}+\frac{e}{c}+d}} - \frac{e\left(\frac{2e\sqrt{1-c^2x^2}}{(c^2d^2-e^2)\sqrt{d+ex}} - \frac{2c\sqrt{d+ex}E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right)\middle|\frac{2e}{cd+e}\right)}{(c^2d^2-e^2)\sqrt{\frac{c(d+ex)}{cd+e}}}\right)}{d} \right) \\
 & \frac{2(a+b\operatorname{sech}^{-1}(cx))}{3e(d+ex)^{3/2}} \\
 & \quad \downarrow 412 \\
 & \frac{2(a+b\operatorname{sech}^{-1}(cx))}{3e(d+ex)^{3/2}} - \frac{2\sqrt{1-\frac{e(1-cx)}{cd+e}}\operatorname{EllipticPi}\left(2,\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right),\frac{2e}{cd+e}\right)}{d\sqrt{-\frac{e(1-cx)}{c}+\frac{e}{c}+d}} \\
 & \frac{3e}{3e}
 \end{aligned}$$

input `Int[(a + b*ArcSech[c*x])/(d + e*x)^(5/2),x]`

output `(-2*(a + b*ArcSech[c*x]))/(3*e*(d + e*x)^(3/2)) - (2*b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-((e*((2*e*Sqrt[1 - c^2*x^2]))/((c^2*d^2 - e^2)*Sqrt[d + e*x]) - (2*c*Sqrt[d + e*x]*EllipticE[ArcSin[Sqrt[1 - c*x]/Sqrt[2]]], (2*e)/(c*d + e)))/((c^2*d^2 - e^2)*Sqrt[(c*(d + e*x))/(c*d + e)])))/d - (2*Sqrt[1 - (e*(1 - c*x))/(c*d + e)]*EllipticPi[2, ArcSin[Sqrt[1 - c*x]/Sqrt[2]], (2*e)/(c*d + e)]/(d*Sqrt[d + e/c - (e*(1 - c*x))/c])))/(3*e)`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 186 `Int[1/(((a_) + (b_)*(x_))*Sqrt[(c_) + (d_)*(x_)]*Sqrt[(e_) + (f_)*(x_)]*Sqrt[(g_) + (h_)*(x_)]), x_] := Simp[-2 Subst[Int[1/(Simp[b*c - a*d - b*x^2, x]*Sqrt[Simp[(d*e - c*f)/d + f*(x^2/d), x]]*Sqrt[Simp[(d*g - c*h)/d + h*(x^2/d), x]]), x], x, Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x] && GtQ[(d*e - c*f)/d, 0]`
- rule 327 `Int[Sqrt[(a_) + (b_)*(x_)^2]/Sqrt[(c_) + (d_)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]/(Sqrt[c]*Rt[-d/c, 2]))*EllipticE[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]`
- rule 412 `Int[1/(((a_) + (b_)*(x_)^2)*Sqrt[(c_) + (d_)*(x_)^2]*Sqrt[(e_) + (f_)*(x_)^2]), x_Symbol] := Simp[(1/(a*Sqrt[c]*Sqrt[e]*Rt[-d/c, 2]))*EllipticPi[b*(c/(a*d)), ArcSin[Rt[-d/c, 2]*x], c*(f/(d*e))], x] /; FreeQ[{a, b, c, d, e, f}, x] && !GtQ[d/c, 0] && GtQ[c, 0] && GtQ[e, 0] && !( !GtQ[f/e, 0] && SimplifierSqrtQ[-f/e, -d/c])`
- rule 413 `Int[1/(((a_) + (b_)*(x_)^2)*Sqrt[(c_) + (d_)*(x_)^2]*Sqrt[(e_) + (f_)*(x_)^2]), x_Symbol] := Simp[Sqrt[1 + (d/c)*x^2]/Sqrt[c + d*x^2] Int[1/((a + b*x^2)*Sqrt[1 + (d/c)*x^2]*Sqrt[e + f*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && !GtQ[c, 0]`
- rule 498 `Int[((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[d*(c + d*x)^(n + 1)*((a + b*x^2)^(p + 1)/((n + 1)*(b*c^2 + a*d^2))), x] + Simp[b/((n + 1)*(b*c^2 + a*d^2)) Int[(c + d*x)^(n + 1)*(a + b*x^2)^p*(c*(n + 1) - d*(n + 2*p + 3)*x), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[n, -1] && ((LtQ[n, -1] && IntQuadraticQ[a, 0, b, c, d, n, p, x]) || (SumSimplerQ[n, 1] && IntegerQ[p]) || ILtQ[Simplify[n + 2*p + 3], 0])`

rule 508 `Int[Sqrt[(c_) + (d_)*(x_)]/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := With[{q = Rt[-b/a, 2]}, Simp[-2*(Sqrt[c + d*x]/(Sqrt[a]*q*Sqrt[q*(c + d*x)/(d + c*q)]))] Subst[Int[Sqrt[1 - 2*d*(x^2/(d + c*q))]/Sqrt[1 - x^2], x], x, Sqrt[(1 - q*x)/2]], x] /; FreeQ[{a, b, c, d}, x] && NegQ[b/a] && GtQ[a, 0]`

rule 632 `Int[1/((x_)*Sqrt[(c_) + (d_)*(x_)]*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] :> With[{q = Rt[-b/a, 2]}, Simp[1/Sqrt[a] Int[1/(x*Sqrt[c + d*x]*Sqrt[1 - q*x]*Sqrt[1 + q*x]), x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[b/a] && GtQ[a, 0]`

rule 635 `Int[((c_) + (d_)*(x_))^(n_)/((x_)*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] := Simp[c^(n + 1/2) Int[1/(x*Sqrt[c + d*x]*Sqrt[a + b*x^2]), x], x] + Int[(c + d*x)^n/Sqrt[a + b*x^2])*ExpandToSum[(1 - c^(n + 1/2)*(c + d*x)^(-n - 1/2))/x, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[n + 1/2, 0]`

rule 6842 `Int[((a_) + ArcSech[(c_)*(x_)]*(b_))*((d_) + (e_)*(x_))^(m_), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*ArcSech[c*x])/(e*(m + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(e*(m + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x)^(m + 1)/(x*Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[m, -1]`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 889 vs.  $2(246) = 492$ .

Time = 20.96 (sec) , antiderivative size = 890, normalized size of antiderivative = 3.20

method	result
derivativedivides	$-\frac{2a}{3(ex+d)^{\frac{3}{2}}} + 2b \left( -\frac{\operatorname{arcsech}(cx)}{3(ex+d)^{\frac{3}{2}}} + \frac{2ce^2 \sqrt{\frac{-c(ex+d)+cd+e}{cex}} x \sqrt{\frac{-c(ex+d)+cd-e}{cex}} \left( \sqrt{\frac{c}{cd+e}} c^2 d(ex+d)^2 - \sqrt{\frac{-c(ex+d)+cd+e}{cd+e}} \sqrt{\frac{-c(ex+d)+cd-e}{cd+e}} \right)}{3(ex+d)^{\frac{3}{2}}} \right)$
default	$-\frac{2a}{3(ex+d)^{\frac{3}{2}}} + 2b \left( -\frac{\operatorname{arcsech}(cx)}{3(ex+d)^{\frac{3}{2}}} + \frac{2ce^2 \sqrt{\frac{-c(ex+d)+cd+e}{cex}} x \sqrt{\frac{-c(ex+d)+cd-e}{cex}} \left( \sqrt{\frac{c}{cd+e}} c^2 d(ex+d)^2 - \sqrt{\frac{-c(ex+d)+cd+e}{cd+e}} \sqrt{\frac{-c(ex+d)+cd-e}{cd+e}} \right)}{3(ex+d)^{\frac{3}{2}}} \right)$
parts	$-\frac{2a}{3(ex+d)^{\frac{3}{2}} e} + 2b \left( -\frac{\operatorname{arcsech}(cx)}{3(ex+d)^{\frac{3}{2}}} + \frac{2ce^2 \sqrt{\frac{-c(ex+d)-cd-e}{cxe}} x \sqrt{\frac{c(ex+d)-cd+e}{cxe}} \left( \sqrt{\frac{c}{cd+e}} c^2 d(ex+d)^2 - \sqrt{\frac{-c(ex+d)-cd-e}{cd+e}} \sqrt{\frac{-c(ex+d)-cd+e}{cd+e}} \right)}{3(ex+d)^{\frac{3}{2}}} \right)$

```
input int((a+b*arcsech(c*x))/(e*x+d)^(5/2),x,method=_RETURNVERBOSE)
```

```
output 2/e*(-1/3*a/(e*x+d)^(3/2)+b*(-1/3/(e*x+d)^(3/2)*arcsech(c*x)+2/3*c*e^2*((-c*(e*x+d)+c*d+e)/c/e/x)^(1/2)*x*(-(-c*(e*x+d)+c*d-e)/c/e/x)^(1/2)*((c/(c*d+e))^(1/2)*c^2*d*(e*x+d)^2-((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x+d)+c*d-e)/(c*d-e))^(1/2)*EllipticF((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e)/(c*d-e))^(1/2))*c^2*d^2*(e*x+d)^(1/2)+c^2*((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x+d)+c*d-e)/(c*d-e))^(1/2)*EllipticE((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e)/(c*d-e))^(1/2))*d^2*(e*x+d)^(1/2)-((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x+d)+c*d-e)/(c*d-e))^(1/2)*EllipticPi((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),1/c*(c*d+e)/d,(c/(c*d-e))^(1/2)/(c/(c*d+e))^(1/2))*c^2*d^2*(e*x+d)^(1/2)-2*(c/(c*d+e))^(1/2)*c^2*d^2*(e*x+d)+((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x+d)+c*d-e)/(c*d-e))^(1/2)*EllipticF((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e)/(c*d-e))^(1/2))*c*d*e*(e*x+d)^(1/2)-((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x+d)+c*d-e)/(c*d-e))^(1/2)*EllipticE((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e)/(c*d-e))^(1/2))*c*d*e*(e*x+d)^(1/2)+(c/(c*d+e))^(1/2)*c^2*d^3+((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x+d)+c*d-e)/(c*d-e))^(1/2)*EllipticPi((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),1/c*(c*d+e)/d,(c/(c*d-e))^(1/2)/(c/(c*d+e))^(1/2))*e^2*(e*x+d)^(1/2)-(c/(c*d+e))^(1/2)*d*e^2)/(c^2*(e*x+d)^2-2*c^2*d*(e*x+d)+c^2*d^2-e^2)/d^2/(c/(c*d+e))^(1/2)/(c*d+e)/(c*d-e)/(e*x+d)^(1/2)))
```

**Fricas [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{5/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex + d)^{\frac{5}{2}}} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x+d)^(5/2),x, algorithm="fricas")`

output `integral(sqrt(e*x + d)*(b*arcsech(c*x) + a)/(e^3*x^3 + 3*d*e^2*x^2 + 3*d^2*e*x + d^3), x)`

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{5/2}} dx = \int \frac{a + b \operatorname{asech}(cx)}{(d + ex)^{\frac{5}{2}}} dx$$

input `integrate((a+b*asech(c*x))/(e*x+d)**(5/2),x)`

output `Integral((a + b*asech(c*x))/(d + e*x)**(5/2), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/(e*x+d)^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e+c*d>0)', see `assume?` for more details)`



**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{5/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex + d)^{5/2}} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/(e*x + d)^(5/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{5/2}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{(d + ex)^{5/2}} dx$$

input `int((a + b*acosh(1/(c*x)))/(d + e*x)^(5/2),x)`

output `int((a + b*acosh(1/(c*x)))/(d + e*x)^(5/2), x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{5/2}} dx = \frac{3\sqrt{ex + d} \left( \int \frac{\operatorname{asech}(cx)}{\sqrt{ex+d}d^2 + 2\sqrt{ex+d}dex + \sqrt{ex+d}e^2x^2} dx \right) bde + 3\sqrt{ex + d} \left( \int \frac{\operatorname{asech}(cx)}{\sqrt{ex+d}d^2 + 2\sqrt{ex+d}dex + \sqrt{ex+d}e^2x^2} dx \right) d}{3\sqrt{ex + d}e(ex + d)}$$

input `int((a+b*asech(c*x))/(e*x+d)^(5/2),x)`

output `(3*sqrt(d + e*x)*int(asech(c*x)/(sqrt(d + e*x)*d**2 + 2*sqrt(d + e*x)*d*e*x + sqrt(d + e*x)*e**2*x**2),x)*b*d*e + 3*sqrt(d + e*x)*int(asech(c*x)/(sqrt(d + e*x)*d**2 + 2*sqrt(d + e*x)*d*e*x + sqrt(d + e*x)*e**2*x**2),x)*b*e**2*x - 2*a)/(3*sqrt(d + e*x)*e*(d + e*x))`

**3.85** 
$$\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex)^{7/2}} dx$$

Optimal result	689
Mathematica [C] (warning: unable to verify)	690
Rubi [A] (verified)	690
Maple [B] (verified)	697
Fricas [F]	698
Sympy [F(-1)]	699
Maxima [F(-2)]	699
Giac [F]	699
Mupad [F(-1)]	700
Reduce [F]	700

**Optimal result**

Integrand size = 18, antiderivative size = 472

$$\begin{aligned} \int \frac{a + b\operatorname{sech}^{-1}(cx)}{(d + ex)^{7/2}} dx &= \frac{4be\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{15d(c^2d^2 - e^2)(d + ex)^{3/2}} \\ &+ \frac{4be(7c^2d^2 - 3e^2)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{15d^2(c^2d^2 - e^2)^2\sqrt{d + ex}} - \frac{2(a + b\operatorname{sech}^{-1}(cx))}{5e(d + ex)^{5/2}} \\ &- \frac{4bc(7c^2d^2 - 3e^2)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{d + ex}E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right) \mid \frac{2e}{cd+e}\right)}{15d^2(c^2d^2 - e^2)^2\sqrt{\frac{c(d+ex)}{cd+e}}} \\ &+ \frac{4bc\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{\frac{c(d+ex)}{cd+e}}\operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{15d(c^2d^2 - e^2)\sqrt{d + ex}} \\ &+ \frac{4b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{\frac{c(d+ex)}{cd+e}}\operatorname{EllipticPi}\left(2, \arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{5d^2e\sqrt{d + ex}} \end{aligned}$$

output

```

4/15*b*e*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/d/(c^2*d^2-e^2
)/(e*x+d)^(3/2)+4/15*b*e*(7*c^2*d^2-3*e^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)
*(-c^2*x^2+1)^(1/2)/d^2/(c^2*d^2-e^2)^2/(e*x+d)^(1/2)-2/5*(a+b*arcsech(c*x
))/e/(e*x+d)^(5/2)-4/15*b*c*(7*c^2*d^2-3*e^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1
/2)*(e*x+d)^(1/2)*EllipticE(1/2*(-c*x+1)^(1/2)*2^(1/2),2^(1/2)*(e/(c*d+e))
^(1/2))/d^2/(c^2*d^2-e^2)^2/(c*(e*x+d)/(c*d+e))^(1/2)+4/15*b*c*(1/(c*x+1))
^(1/2)*(c*x+1)^(1/2)*(c*(e*x+d)/(c*d+e))^(1/2)*EllipticF(1/2*(-c*x+1)^(1/2)
)*2^(1/2),2^(1/2)*(e/(c*d+e))^(1/2))/d/(c^2*d^2-e^2)/(e*x+d)^(1/2)+4/5*b*(
1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(c*(e*x+d)/(c*d+e))^(1/2)*EllipticPi(1/2*(-
c*x+1)^(1/2)*2^(1/2),2,2^(1/2)*(e/(c*d+e))^(1/2))/d^2/e/(e*x+d)^(1/2)

```

### Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 25.40 (sec) , antiderivative size = 8675, normalized size of antiderivative = 18.38

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{7/2}} dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*ArcSech[c*x])/(d + e*x)^(7/2),x]
```

output

```
Result too large to show
```

### Rubi [A] (verified)

Time = 0.90 (sec) , antiderivative size = 433, normalized size of antiderivative = 0.92, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$ , Rules used = {6842, 635, 632, 186, 413, 412, 688, 27, 688, 27, 600, 508, 327, 511, 321}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{7/2}} dx$$

↓ 6842

$$\frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{x(d+ex)^{5/2}\sqrt{1-c^2x^2}} dx}{5e} - \frac{2(a+b\operatorname{sech}^{-1}(cx))}{5e(d+ex)^{5/2}}$$

↓ 635

$$\frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \int \frac{-\frac{xe^2}{d^2} - \frac{2e}{d}}{(d+ex)^{5/2}\sqrt{1-c^2x^2}} dx + \frac{\int \frac{1}{x\sqrt{d+ex}\sqrt{1-c^2x^2}} dx}{d^2} \right)}{5e} - \frac{2(a+b\operatorname{sech}^{-1}(cx))}{5e(d+ex)^{5/2}}$$

↓ 632

$$\frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \int \frac{-\frac{xe^2}{d^2} - \frac{2e}{d}}{(d+ex)^{5/2}\sqrt{1-c^2x^2}} dx + \frac{\int \frac{1}{x\sqrt{1-cx}\sqrt{cx+1}\sqrt{d+ex}} dx}{d^2} \right)}{5e} - \frac{2(a+b\operatorname{sech}^{-1}(cx))}{5e(d+ex)^{5/2}}$$

↓ 186

$$\frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \int \frac{-\frac{xe^2}{d^2} - \frac{2e}{d}}{(d+ex)^{5/2}\sqrt{1-c^2x^2}} dx - \frac{2 \int \frac{1}{cx\sqrt{cx+1}\sqrt{d+\frac{e}{c}-\frac{e(1-cx)}{c}}} d\sqrt{1-cx}}{d^2} \right)}{5e} - \frac{2(a+b\operatorname{sech}^{-1}(cx))}{5e(d+ex)^{5/2}}$$

↓ 413

$$\frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \int \frac{-\frac{xe^2}{d^2} - \frac{2e}{d}}{(d+ex)^{5/2}\sqrt{1-c^2x^2}} dx - \frac{2\sqrt{1-\frac{e(1-cx)}{cd+e}} \int \frac{1}{cx\sqrt{cx+1}\sqrt{1-\frac{e(1-cx)}{cd+e}}} d\sqrt{1-cx}}{d^2\sqrt{-\frac{e(1-cx)}{c}+\frac{e}{c}+d}} \right)}{5e} - \frac{2(a+b\operatorname{sech}^{-1}(cx))}{5e(d+ex)^{5/2}}$$

↓ 412

$$\frac{2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \int \frac{-\frac{xe^2}{d^2} - \frac{2e}{d}}{(d+ex)^{5/2}\sqrt{1-c^2x^2}} dx - \frac{2\sqrt{1-\frac{e(1-cx)}{cd+e}} \operatorname{EllipticPi}\left(2, \arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right)}{d^2\sqrt{-\frac{e(1-cx)}{c}+\frac{e}{c}+d}} \right)}{5e} - \frac{2(a+b\operatorname{sech}^{-1}(cx))}{5e(d+ex)^{5/2}}$$

↓ 688

$$2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{2\int-\frac{e\left(3d\left(2c^2-\frac{e^2}{d^2}\right)-c^2ex\right)}{2d(d+ex)^{3/2}\sqrt{1-c^2x^2}}dx}{3(c^2d^2-e^2)}-\frac{2\sqrt{1-\frac{e(1-cx)}{cd+e}}\operatorname{EllipticPi}\left(2,\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right),\frac{2e}{cd+e}\right)}{d^2\sqrt{-\frac{e(1-cx)}{c}+\frac{e}{c}+d}}-\frac{2e^2\sqrt{1-c^2x^2}}{3d(c^2d^2-e^2)(d+ex)^{3/2}}\right)$$

$$\frac{2(a+b\operatorname{sech}^{-1}(cx))}{5e(d+ex)^{5/2}}$$

↓ 27

$$2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{e\int\frac{3\left(2c^2d-\frac{e^2}{d}\right)-c^2ex}{(d+ex)^{3/2}\sqrt{1-c^2x^2}}dx}{3d(c^2d^2-e^2)}-\frac{2\sqrt{1-\frac{e(1-cx)}{cd+e}}\operatorname{EllipticPi}\left(2,\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right),\frac{2e}{cd+e}\right)}{d^2\sqrt{-\frac{e(1-cx)}{c}+\frac{e}{c}+d}}-\frac{2e^2\sqrt{1-c^2x^2}}{3d(c^2d^2-e^2)(d+ex)^{3/2}}\right)$$

$$\frac{2(a+b\operatorname{sech}^{-1}(cx))}{5e(d+ex)^{5/2}}$$

↓ 688

$$2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{e\left(\frac{2\int\frac{c^2\left(2d\left(3c^2d^2-e^2\right)+e\left(7c^2d^2-3e^2\right)x\right)}{2d\sqrt{d+ex}\sqrt{1-c^2x^2}}dx}{c^2d^2-e^2}+\frac{2e\sqrt{1-c^2x^2}\left(7c^2d^2-3e^2\right)}{d\left(c^2d^2-e^2\right)\sqrt{d+ex}}\right)}{3d(c^2d^2-e^2)}-\frac{2\sqrt{1-\frac{e(1-cx)}{cd+e}}\operatorname{EllipticPi}\left(2,\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right),\frac{2e}{cd+e}\right)}{d^2\sqrt{-\frac{e(1-cx)}{c}+\frac{e}{c}+d}}\right)$$

$$\frac{2(a+b\operatorname{sech}^{-1}(cx))}{5e(d+ex)^{5/2}}$$

↓ 27

$$2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{e\left(\frac{c^2\int\frac{2d\left(3c^2d^2-e^2\right)+e\left(7c^2d^2-3e^2\right)x}{\sqrt{d+ex}\sqrt{1-c^2x^2}}dx}{d\left(c^2d^2-e^2\right)}+\frac{2e\sqrt{1-c^2x^2}\left(7c^2d^2-3e^2\right)}{d\left(c^2d^2-e^2\right)\sqrt{d+ex}}\right)}{3d(c^2d^2-e^2)}-\frac{2\sqrt{1-\frac{e(1-cx)}{cd+e}}\operatorname{EllipticPi}\left(2,\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right),\frac{2e}{cd+e}\right)}{d^2\sqrt{-\frac{e(1-cx)}{c}+\frac{e}{c}+d}}\right)$$

$$\frac{2(a+b\operatorname{sech}^{-1}(cx))}{5e(d+ex)^{5/2}}$$

↓ 600

$$2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{e \left( \frac{c^2 \left( (7c^2d^2-3e^2) \int \frac{\sqrt{d+ex}}{\sqrt{1-c^2x^2}} dx - d(cd-e)(cd+e) \int \frac{1}{\sqrt{d+ex}\sqrt{1-c^2x^2}} dx \right)}{d(c^2d^2-e^2)} + \frac{2e\sqrt{1-c^2x^2}(7c^2d^2-3e^2)}{d(c^2d^2-e^2)\sqrt{d+ex}} \right)}{3d(c^2d^2-e^2)} - \frac{2\sqrt{1-\frac{e(1-cx)}{cd+e}}}{5e} \right)$$

$$\frac{2(a + b\operatorname{sech}^{-1}(cx))}{5e(d + ex)^{5/2}}$$

508

$$2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{e \left( \frac{c^2 \left( -\frac{2(7c^2d^2-3e^2)\sqrt{d+ex} \int \frac{\sqrt{1-\frac{e(1-cx)}{cd+e}} d\sqrt{1-cx}}{\sqrt{\frac{1}{2}(cx-1)+1}\sqrt{2}} - d(cd-e)(cd+e) \int \frac{1}{\sqrt{d+ex}\sqrt{1-c^2x^2}} dx \right)}{c\sqrt{\frac{c(d+ex)}{cd+e}}} - \frac{2e\sqrt{1-c^2x^2}(7c^2d^2-3e^2)}{d(c^2d^2-e^2)\sqrt{d+ex}} \right)}{3d(c^2d^2-e^2)} - \frac{2\sqrt{1-\frac{e(1-cx)}{cd+e}}}{5e} \right)$$

$$\frac{2(a + b\operatorname{sech}^{-1}(cx))}{5e(d + ex)^{5/2}}$$

327

$$2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{e \left( \frac{c^2 \left( -d(cd-e)(cd+e) \int \frac{1}{\sqrt{d+ex}\sqrt{1-c^2x^2}} dx - \frac{2(7c^2d^2-3e^2)\sqrt{d+ex} E \left( \arcsin \left( \frac{\sqrt{1-cx}}{\sqrt{2}} \right) \middle| -\frac{2e}{cd+e} \right)}{c\sqrt{\frac{c(d+ex)}{cd+e}}} \right)}{d(c^2d^2-e^2)} + \frac{2e\sqrt{1-c^2x^2}(7c^2d^2-3e^2)}{d(c^2d^2-e^2)\sqrt{d+ex}} \right)}{3d(c^2d^2-e^2)} - \frac{2\sqrt{1-\frac{e(1-cx)}{cd+e}}}{5e} \right)$$

$$\frac{2(a + b\operatorname{sech}^{-1}(cx))}{5e(d + ex)^{5/2}}$$

5e

$$\begin{array}{c}
 \downarrow 511 \\
 \left( \begin{array}{l}
 c^2 \left( \frac{2d(cd-e)(cd+e)\sqrt{\frac{c(d+ex)}{cd+e}} \int \frac{1}{\sqrt{1-\frac{e(1-cx)}{cd+e}} \sqrt{\frac{1}{2}(cx-1)+1}} d\frac{\sqrt{1-cx}}{\sqrt{2}}} - \frac{2(7c^2d^2-3e^2)\sqrt{d+ex}E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right)\right) \frac{2e}{cd+e}}{c\sqrt{\frac{c(d+ex)}{cd+e}}} \right) \\
 \hline
 e \frac{\quad}{d(c^2d^2-e^2)} \\
 \hline
 2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} - \frac{\quad}{3d(c^2d^2-e^2)}
 \end{array} \right)
 \end{array}$$

$$\begin{array}{c}
 \frac{2(a + b\operatorname{sech}^{-1}(cx))}{5e(d + ex)^{5/2}} \qquad \qquad \qquad 5e \\
 \downarrow 321 \\
 \frac{2(a + b\operatorname{sech}^{-1}(cx))}{5e(d + ex)^{5/2}} - \\
 \left( \begin{array}{l}
 c^2 \left( \frac{2d(cd-e)(cd+e)\sqrt{\frac{c(d+ex)}{cd+e}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right), \frac{2e}{cd+e}\right) - 2(7c^2d^2-3e^2)\sqrt{d+ex}E\left(\arcsin\left(\frac{\sqrt{1-cx}}{\sqrt{2}}\right)\right) \frac{2e}{cd+e}}{c\sqrt{d+ex}} \right) \\
 \hline
 e \frac{\quad}{d(c^2d^2-e^2)} \\
 \hline
 2b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} - \frac{\quad}{3d(c^2d^2-e^2)}
 \end{array} \right)
 \end{array}$$

5e

input `Int[(a + b*ArcSech[c*x])/(d + e*x)^(7/2),x]`

output

```
(-2*(a + b*ArcSech[c*x]))/(5*e*(d + e*x)^(5/2)) - (2*b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-2*e^2*Sqrt[1 - c^2*x^2])/(3*d*(c^2*d^2 - e^2)*(d + e*x)^(3/2)) - (e*((2*e*(7*c^2*d^2 - 3*e^2)*Sqrt[1 - c^2*x^2])/(d*(c^2*d^2 - e^2)*Sqrt[d + e*x]) + (c^2*((-2*(7*c^2*d^2 - 3*e^2)*Sqrt[d + e*x]*EllipticE[ArcSin[Sqrt[1 - c*x]/Sqrt[2]], (2*e)/(c*d + e)))/(c*Sqrt[(c*(d + e*x))/(c*d + e])) + (2*d*(c*d - e)*(c*d + e)*Sqrt[(c*(d + e*x))/(c*d + e])*EllipticF[ArcSin[Sqrt[1 - c*x]/Sqrt[2]], (2*e)/(c*d + e)))/(c*Sqrt[d + e*x])))/(d*(c^2*d^2 - e^2)))/(3*d*(c^2*d^2 - e^2)) - (2*Sqrt[1 - (e*(1 - c*x))/(c*d + e])*EllipticPi[2, ArcSin[Sqrt[1 - c*x]/Sqrt[2]], (2*e)/(c*d + e)]/(d^2*Sqrt[d + e/c - (e*(1 - c*x))/c])))/(5*e)
```

### Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

rule 186

```
Int[1/(((a_.) + (b_.)*(x_))*Sqrt[(c_.) + (d_.)*(x_)]*Sqrt[(e_.) + (f_.)*(x_)]*Sqrt[(g_.) + (h_.)*(x_)]), x_] := Simp[-2 Subst[Int[1/(Simp[b*c - a*d - b*x^2, x]*Sqrt[Simp[(d*e - c*f)/d + f*(x^2/d), x]]*Sqrt[Simp[(d*g - c*h)/d + h*(x^2/d), x]]), x], x, Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x] && GtQ[(d*e - c*f)/d, 0]
```

rule 321

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(1/(Sqrt[a]*Sqrt[c]*Rt[-d/c, 2]))*EllipticF[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-b/a, -d/c])
```

rule 327

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]/(Sqrt[c]*Rt[-d/c, 2]))*EllipticE[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]
```



rule 412 `Int[1/(((a_) + (b_)*(x_)^2)*Sqrt[(c_) + (d_)*(x_)^2]*Sqrt[(e_) + (f_)*(x_)^2]), x_Symbol] := Simp[(1/(a*Sqrt[c]*Sqrt[e]*Rt[-d/c, 2]))*EllipticPi[b*(c/(a*d)), ArcSin[Rt[-d/c, 2]*x], c*(f/(d*e))], x] /; FreeQ[{a, b, c, d, e, f}, x] && !GtQ[d/c, 0] && GtQ[c, 0] && GtQ[e, 0] && !( !GtQ[f/e, 0] && SimplerSqrtQ[-f/e, -d/c])`

rule 413 `Int[1/(((a_) + (b_)*(x_)^2)*Sqrt[(c_) + (d_)*(x_)^2]*Sqrt[(e_) + (f_)*(x_)^2]), x_Symbol] := Simp[Sqrt[1 + (d/c)*x^2]/Sqrt[c + d*x^2] Int[1/((a + b*x^2)*Sqrt[1 + (d/c)*x^2]*Sqrt[e + f*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && !GtQ[c, 0]`

rule 508 `Int[Sqrt[(c_) + (d_)*(x_)]/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := With[{q = Rt[-b/a, 2]}, Simp[-2*(Sqrt[c + d*x]/(Sqrt[a]*q*Sqrt[q*((c + d*x)/(d + c*q))])) Subst[Int[Sqrt[1 - 2*d*(x^2/(d + c*q))]/Sqrt[1 - x^2], x], x, Sqrt[(1 - q*x)/2]], x] /; FreeQ[{a, b, c, d}, x] && NegQ[b/a] && GtQ[a, 0]`

rule 511 `Int[1/(Sqrt[(c_) + (d_)*(x_)]*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] := With[{q = Rt[-b/a, 2]}, Simp[-2*(Sqrt[q*((c + d*x)/(d + c*q))]/(Sqrt[a]*q*Sqrt[c + d*x])) Subst[Int[1/(Sqrt[1 - 2*d*(x^2/(d + c*q))]*Sqrt[1 - x^2]), x], x, Sqrt[(1 - q*x)/2]], x] /; FreeQ[{a, b, c, d}, x] && NegQ[b/a] && GtQ[a, 0]`

rule 600 `Int[((A_) + (B_)*(x_))/(Sqrt[(c_) + (d_)*(x_)]*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] := Simp[B/d Int[Sqrt[c + d*x]/Sqrt[a + b*x^2], x], x] - Simp[(B*c - A*d)/d Int[1/(Sqrt[c + d*x]*Sqrt[a + b*x^2]), x], x] /; FreeQ[{a, b, c, d, A, B}, x] && NegQ[b/a]`

rule 632 `Int[1/((x_)*Sqrt[(c_) + (d_)*(x_)]*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] := With[{q = Rt[-b/a, 2]}, Simp[1/Sqrt[a] Int[1/(x*Sqrt[c + d*x]*Sqrt[1 - q*x]*Sqrt[1 + q*x]), x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[b/a] && GtQ[a, 0]`

rule 635

```
Int[((c_) + (d_)*(x_)^(n_))/((x_)*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] :=
Simp[c^(n + 1/2) Int[1/(x*Sqrt[c + d*x]*Sqrt[a + b*x^2]), x], x] + Int[(
(c + d*x)^n/Sqrt[a + b*x^2])*ExpandToSum[(1 - c^(n + 1/2)*(c + d*x)^(-n - 1
/2))/x, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[n + 1/2, 0]
```

rule 688

```
Int[((d_) + (e_)*(x_)^(m_))*((f_) + (g_)*(x_))*((a_) + (c_)*(x_)^2)^(p
_), x_Symbol] := Simp[(e*f - d*g)*(d + e*x)^(m + 1)*((a + c*x^2)^(p + 1)/(
(m + 1)*(c*d^2 + a*e^2))), x] + Simp[1/((m + 1)*(c*d^2 + a*e^2)) Int[(d +
e*x)^(m + 1)*(a + c*x^2)^p*Simp[(c*d*f + a*e*g)*(m + 1) - c*(e*f - d*g)*(m
+ 2*p + 3)*x, x], x], x] /; FreeQ[{a, c, d, e, f, g, p}, x] && LtQ[m, -1]
&& (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])
```

rule 6842

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_))*((d_) + (e_)*(x_)^(m_)), x_Symbol] :=
Simp[(d + e*x)^(m + 1)*((a + b*ArcSech[c*x])/(e*(m + 1))), x] + Simp[
b*(Sqrt[1 + c*x]/(e*(m + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x)^(m + 1)/(x*
Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[m, -1]
```

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1611 vs.  $2(421) = 842$ .

Time = 25.12 (sec) , antiderivative size = 1612, normalized size of antiderivative = 3.42

method	result	size
derivativdivides	Expression too large to display	1612
default	Expression too large to display	1612
parts	Expression too large to display	1634

input

```
int((a+b*arcsech(c*x))/(e*x+d)^(7/2),x,method=_RETURNVERBOSE)
```

output

```

2/e*(-1/5*a/(e*x+d)^(5/2)+b*(-1/5/(e*x+d)^(5/2)*arcsech(c*x)-2/15*c*e^2*((
-c*(e*x+d)+c*d+e)/c/e/x)^(1/2)*x*(-(-c*(e*x+d)+c*d-e)/c/e/x)^(1/2)*(6*((-c
*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x+d)+c*d-e)/(c*d-e))^(1/2)*Elliptic
F((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e)/(c*d-e))^(1/2))*c^4*d^4*(e*x+d)
^(3/2)-7*((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x+d)+c*d-e)/(c*d-e))^(
1/2)*EllipticE((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e)/(c*d-e))^(1/2))*c^
4*d^4*(e*x+d)^(3/2)+3*((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x+d)+c*d-
e)/(c*d-e))^(1/2)*EllipticPi((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),1*c*(c*d+e)/d
,(c/(c*d-e))^(1/2)/(c/(c*d+e))^(1/2))*c^4*d^4*(e*x+d)^(3/2)-7*(c/(c*d+e))^(
1/2)*c^4*d^3*(e*x+d)^3-7*((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x+d)+
c*d-e)/(c*d-e))^(1/2)*EllipticF((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e)/(
c*d-e))^(1/2))*c^3*d^3*e*(e*x+d)^(3/2)+7*((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)
)*((-c*(e*x+d)+c*d-e)/(c*d-e))^(1/2)*EllipticE((e*x+d)^(1/2)*(c/(c*d+e))^(
1/2),((c*d+e)/(c*d-e))^(1/2))*c^3*d^3*e*(e*x+d)^(3/2)+13*(c/(c*d+e))^(1/2)
*c^4*d^4*(e*x+d)^2-2*((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*((-c*(e*x+d)+c*d-e
)/(c*d-e))^(1/2)*EllipticF((e*x+d)^(1/2)*(c/(c*d+e))^(1/2),((c*d+e)/(c*d-e
))^(1/2))*c^2*d^2*e^2*(e*x+d)^(3/2)+3*((-c*(e*x+d)+c*d+e)/(c*d+e))^(1/2)*(-
c*(e*x+d)+c*d-e)/(c*d-e))^(1/2)*EllipticE((e*x+d)^(1/2)*(c/(c*d+e))^(1/2)
),((c*d+e)/(c*d-e))^(1/2))*c^2*d^2*e^2*(e*x+d)^(3/2)-6*((-c*(e*x+d)+c*d+e)
/(c*d+e))^(1/2)*((-c*(e*x+d)+c*d-e)/(c*d-e))^(1/2)*EllipticPi((e*x+d)^(...

```

**Fricas [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{7/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex + d)^{7/2}} dx$$

input

```
integrate((a+b*arcsech(c*x))/(e*x+d)^(7/2),x, algorithm="fricas")
```

output

```

integral(sqrt(e*x + d)*(b*arcsech(c*x) + a)/(e^4*x^4 + 4*d*e^3*x^3 + 6*d^2
*e^2*x^2 + 4*d^3*e*x + d^4), x)

```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{7/2}} dx = \text{Timed out}$$

input `integrate((a+b*asech(c*x))/(e*x+d)**(7/2), x)`

output `Timed out`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{7/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/(e*x+d)^(7/2), x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e+c*d>0)', see `assume?` for more details)`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{7/2}} dx = \int \frac{b \operatorname{ar} \operatorname{sech}(cx) + a}{(ex + d)^{\frac{7}{2}}} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x+d)^(7/2), x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/(e*x + d)^(7/2), x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{7/2}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{(d + ex)^{7/2}} dx$$

input `int((a + b*acosh(1/(c*x)))/(d + e*x)^(7/2), x)`

output `int((a + b*acosh(1/(c*x)))/(d + e*x)^(7/2), x)`

### Reduce [F]

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex)^{7/2}} dx = \frac{5\sqrt{ex+d} \left( \int \frac{\operatorname{asech}(cx)}{\sqrt{ex+d} d^3 + 3\sqrt{ex+d} d^2 ex + 3\sqrt{ex+d} d e^2 x^2 + \sqrt{ex+d} e^3 x^3} dx \right) b d^2 e + 10\sqrt{ex+d} \left( \dots \right)}$$

input `int((a+b*asech(c*x))/(e*x+d)^(7/2), x)`

output `(5*sqrt(d + e*x)*int(asech(c*x)/(sqrt(d + e*x)*d**3 + 3*sqrt(d + e*x)*d**2*e*x + 3*sqrt(d + e*x)*d*e**2*x**2 + sqrt(d + e*x)*e**3*x**3), x)*b*d**2*e + 10*sqrt(d + e*x)*int(asech(c*x)/(sqrt(d + e*x)*d**3 + 3*sqrt(d + e*x)*d**2*e*x + 3*sqrt(d + e*x)*d*e**2*x**2 + sqrt(d + e*x)*e**3*x**3), x)*b*d*e**2*x + 5*sqrt(d + e*x)*int(asech(c*x)/(sqrt(d + e*x)*d**3 + 3*sqrt(d + e*x)*d**2*e*x + 3*sqrt(d + e*x)*d*e**2*x**2 + sqrt(d + e*x)*e**3*x**3), x)*b*e**3*x**2 - 2*a)/(5*sqrt(d + e*x)*e*(d**2 + 2*d*e*x + e**2*x**2))`

### 3.86 $\int (d + ex)^m (a + b \operatorname{sech}^{-1}(cx)) dx$

Optimal result	701
Mathematica [N/A]	701
Rubi [N/A]	702
Maple [N/A]	703
Fricas [N/A]	703
Sympy [N/A]	703
Maxima [N/A]	704
Giac [N/A]	704
Mupad [N/A]	705
Reduce [N/A]	705

#### Optimal result

Integrand size = 16, antiderivative size = 16

$$\int (d + ex)^m (a + b \operatorname{sech}^{-1}(cx)) dx = \frac{(d + ex)^{1+m} (a + b \operatorname{sech}^{-1}(cx))}{e(1+m)} + \frac{b \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \operatorname{Int}\left(\frac{(d+ex)^{1+m}}{x\sqrt{1-c^2x^2}}, x\right)}{e(1+m)}$$

output

```
(e*x+d)^(1+m)*(a+b*arcsech(c*x))/e/(1+m)+b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)
*Defer(Int)((e*x+d)^(1+m)/x/(-c^2*x^2+1)^(1/2),x)/e/(1+m)
```

#### Mathematica [N/A]

Not integrable

Time = 14.70 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (d + ex)^m (a + b \operatorname{sech}^{-1}(cx)) dx = \int (d + ex)^m (a + b \operatorname{sech}^{-1}(cx)) dx$$

input

```
Integrate[(d + e*x)^m*(a + b*ArcSech[c*x]),x]
```

output `Integrate[(d + e*x)^m*(a + b*ArcSech[c*x]), x]`

**Rubi [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex)^m (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$\downarrow \text{6842}$$

$$\frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{(d+ex)^{m+1}}{x\sqrt{1-c^2x^2}} dx}{e(m+1)} + \frac{(d+ex)^{m+1} (a + b \operatorname{sech}^{-1}(cx))}{e(m+1)}$$

$$\downarrow \text{638}$$

$$\frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{(d+ex)^{m+1}}{x\sqrt{1-c^2x^2}} dx}{e(m+1)} + \frac{(d+ex)^{m+1} (a + b \operatorname{sech}^{-1}(cx))}{e(m+1)}$$

input `Int[(d + e*x)^m*(a + b*ArcSech[c*x]), x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.39 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int (ex + d)^m (a + b \operatorname{arcsech}(cx)) dx$$

input `int((e*x+d)^m*(a+b*arcsech(c*x)),x)`output `int((e*x+d)^m*(a+b*arcsech(c*x)),x)`**Fricas [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (d + ex)^m (a + b \operatorname{sech}^{-1}(cx)) dx = \int (b \operatorname{arsech}(cx) + a)(ex + d)^m dx$$

input `integrate((e*x+d)^m*(a+b*arcsech(c*x)),x, algorithm="fricas")`output `integral((b*arcsech(c*x) + a)*(e*x + d)^m, x)`**Sympy [N/A]**

Not integrable

Time = 17.64 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int (d + ex)^m (a + b \operatorname{sech}^{-1}(cx)) dx = \int (a + b \operatorname{asech}(cx))(d + ex)^m dx$$

input `integrate((e*x+d)**m*(a+b*asech(c*x)),x)`output `Integral((a + b*asech(c*x))*(d + e*x)**m, x)`



**Maxima [N/A]**

Not integrable

Time = 0.66 (sec) , antiderivative size = 222, normalized size of antiderivative = 13.88

$$\int (d + ex)^m (a + b \operatorname{sech}^{-1}(cx)) dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)(ex + d)^m dx$$

input `integrate((e*x+d)^m*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `b*(((e*x + d)*(e*x + d)^m*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1) - (e*x + d)  
 )*(e*x + d)^m*log(x))/(e*(m + 1)) - integrate((c^2*e*(m + 1)*x^3*log(c) -  
 (e*(m + 1)*log(c) - e)*x + d)*(e*x + d)^m/(c^2*e*(m + 1)*x^3 - e*(m + 1)*x  
 ), x) + integrate((c^2*e*x^2 + c^2*d*x)*(e*x + d)^m/(c^2*e*(m + 1)*x^2 + (c^2*e*(m + 1)*x^2 - e*(m + 1))*sqrt(c*x + 1)*sqrt(-c*x + 1) - e*(m + 1)), x)) + (e*x + d)^(m + 1)*a/(e*(m + 1))`

**Giac [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (d + ex)^m (a + b \operatorname{sech}^{-1}(cx)) dx = \int (b \operatorname{ar} \operatorname{sech}(cx) + a)(ex + d)^m dx$$

input `integrate((e*x+d)^m*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*(e*x + d)^m, x)`

**Mupad [N/A]**

Not integrable

Time = 3.89 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int (d + ex)^m (a + b \operatorname{sech}^{-1}(cx)) dx = \int \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) (d + ex)^m dx$$

input `int((a + b*acosh(1/(c*x)))*(d + e*x)^m,x)`output `int((a + b*acosh(1/(c*x)))*(d + e*x)^m, x)`**Reduce [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 66, normalized size of antiderivative = 4.12

$$\int (d + ex)^m (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{(ex + d)^m ad + (ex + d)^m aex + \left(\int (ex + d)^m \operatorname{asech}(cx) dx\right) bem + \left(\int (ex + d)^m \operatorname{asech}(cx) dx\right) be}{e(m + 1)}$$

input `int((e*x+d)^m*(a+b*asech(c*x)),x)`output `((d + e*x)**m*a*d + (d + e*x)**m*a*e*x + int((d + e*x)**m*asech(c*x),x)*b*e*m + int((d + e*x)**m*asech(c*x),x)*b*e)/(e*(m + 1))`

### 3.87 $\int x^4(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	706
Mathematica [C] (verified)	707
Rubi [A] (verified)	707
Maple [A] (verified)	710
Fricas [A] (verification not implemented)	710
Sympy [F]	711
Maxima [A] (verification not implemented)	711
Giac [F]	712
Mupad [F(-1)]	712
Reduce [F]	713

#### Optimal result

Integrand size = 19, antiderivative size = 229

$$\int x^4(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx = -\frac{b(42c^2d + 25e) x \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{560c^6} - \frac{b(42c^2d + 25e) x^3 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{840c^4} - \frac{bex^5 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{42c^2} + \frac{1}{5} dx^5 (a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{7} ex^7 (a + b\operatorname{sech}^{-1}(cx)) + \frac{b(42c^2d + 25e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \arcsin(cx)}{560c^7}$$

output

```
-1/560*b*(42*c^2*d+25*e)*x*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^6-1/840*b*(42*c^2*d+25*e)*x^3*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^4-1/42*b*e*x^5*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^2+1/5*d*x^5*(a+b*arcsech(c*x))+1/7*e*x^7*(a+b*arcsech(c*x))+1/560*b*(42*c^2*d+25*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arcsin(c*x)/c^7
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.71

$$\int x^4(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{48ac^7x^5(7d + 5ex^2) - bcx\sqrt{\frac{1-cx}{1+cx}}(1 + cx)(75e + 2c^2(63d + 25ex^2) + c^4(84dx^2 + 40ex^4)) + 48bc^7x^5(7d + 5ex^2)}{1680c^7}$$

input

```
Integrate[x^4*(d + e*x^2)*(a + b*ArcSech[c*x]),x]
```

output

```
(48*a*c^7*x^5*(7*d + 5*e*x^2) - b*c*x*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*
(75*e + 2*c^2*(63*d + 25*e*x^2) + c^4*(84*d*x^2 + 40*e*x^4)) + 48*b*c^7*x^
5*(7*d + 5*e*x^2)*ArcSech[c*x] + (3*I)*b*(42*c^2*d + 25*e)*Log[(-2*I)*c*x
+ 2*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)]/(1680*c^7)
```

**Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.72, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$ , Rules used = {6855, 27, 363, 262, 262, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx$$

$$\downarrow \text{6855}$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^4(5ex^2 + 7d)}{35\sqrt{1-c^2x^2}} dx + \frac{1}{5}dx^5(a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{7}ex^7(a + b\operatorname{sech}^{-1}(cx))$$

$$\downarrow \text{27}$$

$$\frac{1}{35}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^4(5ex^2 + 7d)}{\sqrt{1-c^2x^2}} dx + \frac{1}{5}dx^5(a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{7}ex^7(a + b\operatorname{sech}^{-1}(cx))$$

↓ 363

$$\frac{1}{35}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{6}\left(\frac{25e}{c^2}+42d\right)\int\frac{x^4}{\sqrt{1-c^2x^2}}dx-\frac{5ex^5\sqrt{1-c^2x^2}}{6c^2}\right)+\frac{1}{5}dx^5(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{7}ex^7(a+b\operatorname{sech}^{-1}(cx))$$

↓ 262

$$\frac{1}{35}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{6}\left(\frac{25e}{c^2}+42d\right)\left(\frac{3\int\frac{x^2}{\sqrt{1-c^2x^2}}dx}{4c^2}-\frac{x^3\sqrt{1-c^2x^2}}{4c^2}\right)-\frac{5ex^5\sqrt{1-c^2x^2}}{6c^2}\right)+\frac{1}{5}dx^5(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{7}ex^7(a+b\operatorname{sech}^{-1}(cx))$$

↓ 262

$$\frac{1}{35}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{6}\left(\frac{25e}{c^2}+42d\right)\left(\frac{3\left(\frac{\int\frac{1}{\sqrt{1-c^2x^2}}dx}{2c^2}-\frac{x\sqrt{1-c^2x^2}}{2c^2}\right)}{4c^2}-\frac{x^3\sqrt{1-c^2x^2}}{4c^2}\right)-\frac{5ex^5\sqrt{1-c^2x^2}}{6c^2}\right)+\frac{1}{5}dx^5(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{7}ex^7(a+b\operatorname{sech}^{-1}(cx))$$

↓ 223

$$\frac{1}{5}dx^5(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{7}ex^7(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{35}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{6}\left(\frac{3\left(\frac{\arcsin(cx)}{2c^3}-\frac{x\sqrt{1-c^2x^2}}{2c^2}\right)}{4c^2}-\frac{x^3\sqrt{1-c^2x^2}}{4c^2}\right)\left(\frac{25e}{c^2}+42d\right)-\frac{5ex^5\sqrt{1-c^2x^2}}{6c^2}\right)$$

input `Int[x^4*(d + e*x^2)*(a + b*ArcSech[c*x]),x]`

output `(d*x^5*(a + b*ArcSech[c*x]))/5 + (e*x^7*(a + b*ArcSech[c*x]))/7 + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-5*e*x^5*Sqrt[1 - c^2*x^2])/(6*c^2) + ((42*d + (25*e)/c^2)*(-1/4*(x^3*Sqrt[1 - c^2*x^2])/c^2 + (3*(-1/2*(x*Sqrt[1 - c^2*x^2])/c^2 + ArcSin[c*x]/(2*c^3)))/(4*c^2)))/6)/35`

## Defintions of rubi rules used

- rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 223  $\text{Int}[1/\text{Sqrt}[(a_*) + (b_*)(x_)^2], x\_Symbol] \rightarrow \text{Simp}[\text{ArcSin}[\text{Rt}[-b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[-b, 2], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{GtQ}[a, 0] \ \&\& \ \text{NegQ}[b]$
- rule 262  $\text{Int}[((c_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^2)^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[c*(c*x)^{(m-1)}((a + b*x^2)^{(p+1})/(b*(m+2*p+1))), x] - \text{Simp}[a*c^2*((m-1)/(b*(m+2*p+1))) \text{ Int}[(c*x)^{(m-2)}*(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{GtQ}[m, 2-1] \ \&\& \ \text{NeQ}[m+2*p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$
- rule 363  $\text{Int}[((e_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^2)^{(p_*)}((c_*) + (d_*)(x_)^2), x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}((a + b*x^2)^{(p+1})/(b*e*(m+2*p+3))), x] - \text{Simp}[(a*d*(m+1) - b*c*(m+2*p+3))/(b*(m+2*p+3)) \text{ Int}[(e*x)^m*(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, p\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m+2*p+3, 0]$
- rule 6855  $\text{Int}[(a_*) + \text{ArcSech}[(c_*)(x_)]*(b_*)((f_*)(x_))^{(m_*)}((d_*) + (e_*)(x_)^2)^{(p_*)}, x\_Symbol] \rightarrow \text{With}[\{u = \text{IntHide}[(f*x)^m*(d + e*x^2)^p, x]\}, \text{Simp}[(a + b*\text{ArcSech}[c*x]) \ u, x] + \text{Simp}[b*\text{Sqrt}[1 + c*x]*\text{Sqrt}[1/(1 + c*x)] \ \text{Int}[\text{SimplifyIntegrand}[u/(x*\text{Sqrt}[1 - c*x]*\text{Sqrt}[1 + c*x]), x], x], x]] /; \text{FreeQ}[\{a, b, c, d, e, f, m, p\}, x] \ \&\& \ ((\text{IGtQ}[p, 0] \ \&\& \ !(\text{ILtQ}[(m-1)/2, 0] \ \&\& \ \text{GtQ}[m+2*p+3, 0])) \ || \ (\text{IGtQ}[(m+1)/2, 0] \ \&\& \ !(\text{ILtQ}[p, 0] \ \&\& \ \text{GtQ}[m+2*p+3, 0])) \ || \ (\text{ILtQ}[(m+2*p+1)/2, 0] \ \&\& \ !\text{ILtQ}[(m-1)/2, 0]))$

### Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 213, normalized size of antiderivative = 0.93

method	result
parts	$a\left(\frac{1}{7}e x^7 + \frac{1}{5}d x^5\right) + \frac{b\left(\frac{c^5 \operatorname{arcsech}(cx)e x^7}{7} + \frac{\operatorname{arcsech}(cx)d c^5 x^5}{5} - \sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}} (84c^5 d \sqrt{-c^2 x^2+1} x^3 + 40e \sqrt{-c^2 x^2+1} c^5)\right)}{c^5}$
derivativelimit	$\frac{a\left(\frac{1}{5}d c^7 x^5 + \frac{1}{7}e c^7 x^7\right)}{c^2} + \frac{b\left(\frac{\operatorname{arcsech}(cx)d c^7 x^5}{5} + \frac{\operatorname{arcsech}(cx)e c^7 x^7}{7} - \sqrt{-\frac{cx-1}{cx}} c x \sqrt{\frac{cx+1}{cx}} (84c^5 d \sqrt{-c^2 x^2+1} x^3 + 40e \sqrt{-c^2 x^2+1} c^5)\right)}{c^5}$
default	$\frac{a\left(\frac{1}{5}d c^7 x^5 + \frac{1}{7}e c^7 x^7\right)}{c^2} + \frac{b\left(\frac{\operatorname{arcsech}(cx)d c^7 x^5}{5} + \frac{\operatorname{arcsech}(cx)e c^7 x^7}{7} - \sqrt{-\frac{cx-1}{cx}} c x \sqrt{\frac{cx+1}{cx}} (84c^5 d \sqrt{-c^2 x^2+1} x^3 + 40e \sqrt{-c^2 x^2+1} c^5)\right)}{c^5}$

```
input int(x^4*(e*x^2+d)*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)
```

```
output a*(1/7*e*x^7+1/5*d*x^5)+b/c^5*(1/7*c^5*arcsech(c*x)*e*x^7+1/5*arcsech(c*x)*d*c^5*x^5-1/1680/c*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(84*c^5*d*(-c^2*x^2+1)^(1/2)*x^3+40*e*(-c^2*x^2+1)^(1/2)*c^5*x^5+126*(-c^2*x^2+1)^(1/2)*c^3*d*x+50*e*c^3*x^3*(-c^2*x^2+1)^(1/2)-126*arcsin(c*x)*c^2*d+75*(-c^2*x^2+1)^(1/2)*e*c*x-75*arcsin(c*x)*e)/(-c^2*x^2+1)^(1/2))
```

### Fricas [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 259, normalized size of antiderivative = 1.13

$$\int x^4(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{240 ac^7 ex^7 + 336 ac^7 dx^5 - 6(42 bc^2 d + 25 be) \arctan\left(\frac{cx\sqrt{-\frac{c^2 x^2-1}{c^2 x^2}}-1}{cx}\right) - 48(7 bc^7 d + 5 bc^7 e) \log\left(\frac{cx\sqrt{-\frac{c^2 x^2-1}{c^2 x^2}}}{cx}\right)}{c^5}$$

```
input integrate(x^4*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="fricas")
```

output

```
1/1680*(240*a*c^7*e*x^7 + 336*a*c^7*d*x^5 - 6*(42*b*c^2*d + 25*b*e)*arctan
((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/(c*x)) - 48*(7*b*c^7*d + 5*b*c^7
*e)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) + 48*(5*b*c^7*e*x^7 +
7*b*c^7*d*x^5 - 7*b*c^7*d - 5*b*c^7*e)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x
^2)) + 1)/(c*x)) - (40*b*c^6*e*x^6 + 2*(42*b*c^6*d + 25*b*c^4*e)*x^4 + 3*(
42*b*c^4*d + 25*b*c^2*e)*x^2)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2))/c^7
```

**Sympy [F]**

$$\int x^4(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx = \int x^4(a + b\operatorname{asech}(cx))(d + ex^2) dx$$

input

```
integrate(x**4*(e*x**2+d)*(a+b*asech(c*x)), x)
```

output

```
Integral(x**4*(a + b*asech(c*x))*(d + e*x**2), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.07

$$\int x^4(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx = \frac{1}{7} aex^7 + \frac{1}{5} adx^5$$

$$+ \frac{1}{40} \left( 8x^5 \operatorname{arsech}(cx) - \frac{3\left(\frac{1}{c^2x^2}-1\right)^{\frac{3}{2}} + 5\sqrt{\frac{1}{c^2x^2}-1} + \frac{3\arctan\left(\sqrt{\frac{1}{c^2x^2}-1}\right)}{c^4}}{c^4\left(\frac{1}{c^2x^2}-1\right)^2 + 2c^4\left(\frac{1}{c^2x^2}-1\right) + c^4} + \frac{3\arctan\left(\sqrt{\frac{1}{c^2x^2}-1}\right)}{c^4} \right) bd$$

$$+ \frac{1}{336} \left( 48x^7 \operatorname{arsech}(cx) - \frac{15\left(\frac{1}{c^2x^2}-1\right)^{\frac{5}{2}} + 40\left(\frac{1}{c^2x^2}-1\right)^{\frac{3}{2}} + 33\sqrt{\frac{1}{c^2x^2}-1} + \frac{15\arctan\left(\sqrt{\frac{1}{c^2x^2}-1}\right)}{c^6}}{c^6\left(\frac{1}{c^2x^2}-1\right)^3 + 3c^6\left(\frac{1}{c^2x^2}-1\right)^2 + 3c^6\left(\frac{1}{c^2x^2}-1\right) + c^6} + \frac{15\arctan\left(\sqrt{\frac{1}{c^2x^2}-1}\right)}{c^6} \right) be$$

input

```
integrate(x^4*(e*x^2+d)*(a+b*arcsech(c*x)), x, algorithm="maxima")
```



output

```
1/7*a*e*x^7 + 1/5*a*d*x^5 + 1/40*(8*x^5*arcsech(c*x) - ((3*(1/(c^2*x^2) - 1)^(3/2) + 5*sqrt(1/(c^2*x^2) - 1))/(c^4*(1/(c^2*x^2) - 1)^2 + 2*c^4*(1/(c^2*x^2) - 1) + c^4) + 3*arctan(sqrt(1/(c^2*x^2) - 1))/c^4)/c)*b*d + 1/336*(48*x^7*arcsech(c*x) - ((15*(1/(c^2*x^2) - 1)^(5/2) + 40*(1/(c^2*x^2) - 1)^(3/2) + 33*sqrt(1/(c^2*x^2) - 1))/(c^6*(1/(c^2*x^2) - 1)^3 + 3*c^6*(1/(c^2*x^2) - 1)^2 + 3*c^6*(1/(c^2*x^2) - 1) + c^6) + 15*arctan(sqrt(1/(c^2*x^2) - 1))/c^6)/c)*b*e
```

**Giac [F]**

$$\int x^4(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)(b\operatorname{ar}\operatorname{sech}(cx) + a)x^4 dx$$

input

```
integrate(x^4*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)*(b*arcsech(c*x) + a)*x^4, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int x^4(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx = \int x^4(ex^2 + d) \left( a + b\operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input

```
int(x^4*(d + e*x^2)*(a + b*acosh(1/(c*x))),x)
```

output

```
int(x^4*(d + e*x^2)*(a + b*acosh(1/(c*x))), x)
```

**Reduce [F]**

$$\int x^4(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx = \left( \int \operatorname{asech}(cx) x^6 dx \right) be + \left( \int \operatorname{asech}(cx) x^4 dx \right) bd + \frac{adx^5}{5} + \frac{aex^7}{7}$$

input `int(x^4*(e*x^2+d)*(a+b*asech(c*x)),x)`

output `(35*int(asech(c*x)*x**6,x)*b*e + 35*int(asech(c*x)*x**4,x)*b*d + 7*a*d*x**5 + 5*a*e*x**7)/35`

### 3.88 $\int x^2(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	714
Mathematica [C] (verified)	715
Rubi [A] (verified)	715
Maple [A] (verified)	717
Fricas [B] (verification not implemented)	718
Sympy [F]	718
Maxima [A] (verification not implemented)	719
Giac [F]	719
Mupad [F(-1)]	720
Reduce [F]	720

#### Optimal result

Integrand size = 19, antiderivative size = 174

$$\int x^2(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx = -\frac{b(20c^2d + 9e) x \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{120c^4} - \frac{bex^3 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{20c^2} + \frac{1}{3} dx^3 (a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{5} ex^5 (a + b\operatorname{sech}^{-1}(cx)) + \frac{b(20c^2d + 9e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \arcsin(cx)}{120c^5}$$

output

```
-1/120*b*(20*c^2*d+9*e)*x*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^4-1/20*b*e*x^3*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^2+1/3*d*x^3*(a+b*arcsech(c*x))+1/5*e*x^5*(a+b*arcsech(c*x))+1/120*b*(20*c^2*d+9*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arcsin(c*x)/c^5
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.17 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.83

$$\int x^2(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{8ac^5x^3(5d + 3ex^2) - bcx\sqrt{\frac{1-cx}{1+cx}}(1 + cx)(9e + c^2(20d + 6ex^2)) + 8bc^5x^3(5d + 3ex^2)\operatorname{sech}^{-1}(cx) + ib(20c^2d + 9e)\operatorname{Log}\left[\frac{-2Ix + 2\sqrt{(1-cx)/(1+cx)}(1+cx)}{120c^5}\right]}{120c^5}$$

input

```
Integrate[x^2*(d + e*x^2)*(a + b*ArcSech[c*x]),x]
```

output

```
(8*a*c^5*x^3*(5*d + 3*e*x^2) - b*c*x*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(9*e + c^2*(20*d + 6*e*x^2)) + 8*b*c^5*x^3*(5*d + 3*e*x^2)*ArcSech[c*x] + I*b*(20*c^2*d + 9*e)*Log[(-2*I)*c*x + 2*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)])/(120*c^5)
```

**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.76, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {6855, 27, 363, 262, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx$$

$$\downarrow \text{6855}$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^2(3ex^2 + 5d)}{15\sqrt{1-c^2x^2}} dx + \frac{1}{3}dx^3(a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{5}ex^5(a + b\operatorname{sech}^{-1}(cx))$$

$$\downarrow \text{27}$$

$$\frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^2(3ex^2 + 5d)}{\sqrt{1-c^2x^2}} dx + \frac{1}{3}dx^3(a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{5}ex^5(a + b\operatorname{sech}^{-1}(cx))$$

$$\begin{aligned}
& \downarrow 363 \\
& \frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{4}\left(\frac{9e}{c^2}+20d\right)\int\frac{x^2}{\sqrt{1-c^2x^2}}dx-\frac{3ex^3\sqrt{1-c^2x^2}}{4c^2}\right)+ \\
& \quad \frac{1}{3}dx^3(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{5}ex^5(a+b\operatorname{sech}^{-1}(cx)) \\
& \downarrow 262 \\
& \frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{4}\left(\frac{9e}{c^2}+20d\right)\left(\frac{\int\frac{1}{\sqrt{1-c^2x^2}}dx}{2c^2}-\frac{x\sqrt{1-c^2x^2}}{2c^2}\right)-\frac{3ex^3\sqrt{1-c^2x^2}}{4c^2}\right)+ \\
& \quad \frac{1}{3}dx^3(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{5}ex^5(a+b\operatorname{sech}^{-1}(cx)) \\
& \downarrow 223 \\
& \quad \frac{1}{3}dx^3(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{5}ex^5(a+b\operatorname{sech}^{-1}(cx))+ \\
& \frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{4}\left(\frac{\arcsin(cx)}{2c^3}-\frac{x\sqrt{1-c^2x^2}}{2c^2}\right)\left(\frac{9e}{c^2}+20d\right)-\frac{3ex^3\sqrt{1-c^2x^2}}{4c^2}\right)
\end{aligned}$$

input `Int[x^2*(d + e*x^2)*(a + b*ArcSech[c*x]),x]`

output `(d*x^3*(a + b*ArcSech[c*x]))/3 + (e*x^5*(a + b*ArcSech[c*x]))/5 + (b*sqrt[
(1 + c*x)^(-1)]*sqrt[1 + c*x]*((-3*e*x^3*sqrt[1 - c^2*x^2])/(4*c^2) + ((20
*d + (9*e)/c^2)*(-1/2*(x*sqrt[1 - c^2*x^2])/c^2 + ArcSin[c*x]/(2*c^3))))/4
)/15`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt
[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

```
rule 262 Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

```
rule 363 Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(b*e*(m + 2*p + 3))), x] - Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(b*(m + 2*p + 3)) Int[(e*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + 2*p + 3, 0]
```

```
rule 6855 Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

### Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.98

method	result
parts	$a\left(\frac{1}{5}e x^5 + \frac{1}{3}d x^3\right) + \frac{b\left(\frac{c^3 \operatorname{arcsech}(cx)e x^5}{5} + \frac{\operatorname{arcsech}(cx)c^3 x^3 d}{3} - \sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}} (20\sqrt{-c^2 x^2+1} c^3 dx + 6e c^3 x^3 \sqrt{-c^2 x^2+1})}{120c\sqrt{-c^2 x^2+1}}\right)}{c^3}$
derivativedivides	$\frac{a\left(\frac{1}{3}d c^5 x^3 + \frac{1}{5}e c^5 x^5\right)}{c^2} + \frac{b\left(\frac{\operatorname{arcsech}(cx)d c^5 x^3}{3} + \frac{\operatorname{arcsech}(cx)e c^5 x^5}{5} - \sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (20\sqrt{-c^2 x^2+1} c^3 dx + 6e c^3 x^3 \sqrt{-c^2 x^2+1})}{120\sqrt{-c^2 x^2+1}}\right)}{c^3}$
default	$\frac{a\left(\frac{1}{3}d c^5 x^3 + \frac{1}{5}e c^5 x^5\right)}{c^2} + \frac{b\left(\frac{\operatorname{arcsech}(cx)d c^5 x^3}{3} + \frac{\operatorname{arcsech}(cx)e c^5 x^5}{5} - \sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (20\sqrt{-c^2 x^2+1} c^3 dx + 6e c^3 x^3 \sqrt{-c^2 x^2+1})}{120\sqrt{-c^2 x^2+1}}\right)}{c^3}$

```
input int(x^2*(e*x^2+d)*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)
```

output

```
a*(1/5*e*x^5+1/3*d*x^3)+b/c^3*(1/5*c^3*arcsech(c*x)*e*x^5+1/3*arcsech(c*x)
*c^3*x^3*d-1/120/c*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(20*(-c^2*x^
2+1)^(1/2)*c^3*d*x+6*e*c^3*x^3*(-c^2*x^2+1)^(1/2)-20*arcsin(c*x)*c^2*d+9*(
-c^2*x^2+1)^(1/2)*e*c*x-9*arcsin(c*x)*e)/(-c^2*x^2+1)^(1/2))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 238 vs.  $2(100) = 200$ .

Time = 0.14 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.37

$$\int x^2(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{24ac^5ex^5 + 40ac^5dx^3 - 2(20bc^2d + 9be) \arctan\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{cx}\right) - 8(5bc^5d + 3bc^5e) \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{x}\right)}{1}$$

input

```
integrate(x^2*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="fricas")
```

output

```
1/120*(24*a*c^5*e*x^5 + 40*a*c^5*d*x^3 - 2*(20*b*c^2*d + 9*b*e)*arctan((c*
x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/(c*x)) - 8*(5*b*c^5*d + 3*b*c^5*e)*l
og((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) + 8*(3*b*c^5*e*x^5 + 5*b*c^
5*d*x^3 - 5*b*c^5*d - 3*b*c^5*e)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) +
1)/(c*x)) - (6*b*c^4*e*x^4 + (20*b*c^4*d + 9*b*c^2*e)*x^2)*sqrt(-(c^2*x^2
- 1)/(c^2*x^2)))/c^5
```

### Sympy [F]

$$\int x^2(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx = \int x^2(a + b\operatorname{asech}(cx))(d + ex^2) dx$$

input

```
integrate(x**2*(e*x**2+d)*(a+b*asech(c*x)),x)
```

output

```
Integral(x**2*(a + b*asech(c*x))*(d + e*x**2), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.05

$$\int x^2(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{1}{5} aex^5 + \frac{1}{3} adx^3 + \frac{1}{6} \left( 2x^3 \operatorname{ar} \operatorname{sech}(cx) - \frac{\frac{\sqrt{\frac{1}{c^2x^2}-1}}{c^2\left(\frac{1}{c^2x^2}-1\right)+c^2} + \frac{\arctan\left(\sqrt{\frac{1}{c^2x^2}-1}\right)}{c^2}}{c} \right) bd$$

$$+ \frac{1}{40} \left( 8x^5 \operatorname{ar} \operatorname{sech}(cx) - \frac{\frac{3\left(\frac{1}{c^2x^2}-1\right)^{\frac{3}{2}}+5\sqrt{\frac{1}{c^2x^2}-1}}{c^4\left(\frac{1}{c^2x^2}-1\right)^2+2c^4\left(\frac{1}{c^2x^2}-1\right)+c^4} + \frac{3\arctan\left(\sqrt{\frac{1}{c^2x^2}-1}\right)}{c^4}}{c} \right) be$$

input `integrate(x^2*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `1/5*a*e*x^5 + 1/3*a*d*x^3 + 1/6*(2*x^3*arcsech(c*x) - (sqrt(1/(c^2*x^2) - 1)/(c^2*(1/(c^2*x^2) - 1) + c^2) + arctan(sqrt(1/(c^2*x^2) - 1)/c^2)/c)*b*d + 1/40*(8*x^5*arcsech(c*x) - ((3*(1/(c^2*x^2) - 1)^(3/2) + 5*sqrt(1/(c^2*x^2) - 1))/(c^4*(1/(c^2*x^2) - 1)^2 + 2*c^4*(1/(c^2*x^2) - 1) + c^4) + 3*arctan(sqrt(1/(c^2*x^2) - 1)/c^4)/c)*b*e`

**Giac [F]**

$$\int x^2(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)(b \operatorname{ar} \operatorname{sech}(cx) + a)x^2 dx$$

input `integrate(x^2*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((e*x^2 + d)*(b*arcsech(c*x) + a)*x^2, x)`



**Mupad [F(-1)]**

Timed out.

$$\int x^2 (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^2 (ex^2 + d) \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int(x^2*(d + e*x^2)*(a + b*acosh(1/(c*x))),x)`output `int(x^2*(d + e*x^2)*(a + b*acosh(1/(c*x))), x)`**Reduce [F]**

$$\int x^2 (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx = \left( \int a \operatorname{sech}(cx) x^4 dx \right) be + \left( \int a \operatorname{sech}(cx) x^2 dx \right) bd + \frac{ad x^3}{3} + \frac{ae x^5}{5}$$

input `int(x^2*(e*x^2+d)*(a+b*asech(c*x)),x)`output `(15*int(asech(c*x)*x**4,x)*b*e + 15*int(asech(c*x)*x**2,x)*b*d + 5*a*d*x**3 + 3*a*e*x**5)/15`

### 3.89 $\int (d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	721
Mathematica [C] (verified)	721
Rubi [A] (verified)	722
Maple [A] (verified)	724
Fricas [B] (verification not implemented)	724
Sympy [F]	725
Maxima [A] (verification not implemented)	725
Giac [F]	726
Mupad [F(-1)]	726
Reduce [F]	726

#### Optimal result

Integrand size = 16, antiderivative size = 112

$$\int (d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx = -\frac{bex\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{6c^2} + dx(a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{3}ex^3(a + b\operatorname{sech}^{-1}(cx)) + \frac{b(6c^2d + e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\arcsin(cx)}{6c^3}$$

output

```
-1/6*b*e*x*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^2+d*x*(a+b*arcsech(c*x))+1/3*e*x^3*(a+b*arcsech(c*x))+1/6*b*(6*c^2*d+e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arcsin(c*x)/c^3
```

#### Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.69

$$\int (d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx = adx + \frac{1}{3}aex^3 + be\sqrt{\frac{1-cx}{1+cx}} \left( -\frac{x}{6c^2} - \frac{x^2}{6c} \right) + bdx\operatorname{sech}^{-1}(cx) + \frac{1}{3}bex^3\operatorname{sech}^{-1}(cx) + \frac{2bd\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\arctan\left(\frac{\sqrt{1-c^2x^2}}{1-cx}\right)}{c-c^2x} + \frac{ibe\log\left(-2icx + 2\sqrt{\frac{1-cx}{1+cx}}(1+cx)\right)}{6c^3}$$

input `Integrate[(d + e*x^2)*(a + b*ArcSech[c*x]),x]`

output `a*d*x + (a*e*x^3)/3 + b*e*Sqrt[(1 - c*x)/(1 + c*x)]*(-1/6*x/c^2 - x^2/(6*c)) + b*d*x*ArcSech[c*x] + (b*e*x^3*ArcSech[c*x])/3 + (2*b*d*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*ArcTan[Sqrt[1 - c^2*x^2]/(1 - c*x)]/(c - c^2*x) + ((I/6)*b*e*Log[(-2*I)*c*x + 2*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)])/c^3`

### Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.86, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {6845, 27, 299, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$$

↓ 6845

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{ex^2 + 3d}{3\sqrt{1-c^2x^2}} dx + dx(a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{3}ex^3(a + b\operatorname{sech}^{-1}(cx))$$

↓ 27

$$\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\int\frac{ex^2+3d}{\sqrt{1-c^2x^2}}dx+dx(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{3}ex^3(a+b\operatorname{sech}^{-1}(cx))$$

↓ 299

$$\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{(6c^2d+e)\int\frac{1}{\sqrt{1-c^2x^2}}dx}{2c^2}-\frac{ex\sqrt{1-c^2x^2}}{2c^2}\right)+dx(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{3}ex^3(a+b\operatorname{sech}^{-1}(cx))$$

↓ 223

$$dx(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{3}ex^3(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{\arcsin(cx)(6c^2d+e)}{2c^3}-\frac{ex\sqrt{1-c^2x^2}}{2c^2}\right)$$

input `Int[(d + e*x^2)*(a + b*ArcSech[c*x]), x]`

output `d*x*(a + b*ArcSech[c*x]) + (e*x^3*(a + b*ArcSech[c*x]))/3 + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/2*(e*x*Sqrt[1 - c^2*x^2])/c^2 + ((6*c^2*d + e)*ArcSin[c*x])/(2*c^3)))/3`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 299 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[d*x*((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NeQ[2*p + 3, 0]`

rule 6845

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol]
:> With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e}, x] && (IGtQ[p, 0] || ILtQ[p + 1/2, 0])
```

**Maple [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.07

method	result
parts	$a\left(\frac{1}{3}x^3e + dx\right) + \frac{b\left(\frac{c \operatorname{arcsech}(cx)x^3e}{3} + \operatorname{arcsech}(cx)dx + \frac{\sqrt{-\frac{cx-1}{cx}}x\sqrt{\frac{cx+1}{cx}}(6 \arcsin(cx)c^2d - \sqrt{-c^2x^2+1}ecx + \arcsin(cx))}{6c\sqrt{-c^2x^2+1}}\right)}{c}$
derivativedivides	$\frac{\frac{a(d c^3 x + \frac{1}{3} c^3 x^3 e)}{c^2} + \frac{b\left(\operatorname{arcsech}(cx) d c^3 x + \frac{\operatorname{arcsech}(cx) c^3 x^3 e}{3} + \frac{\sqrt{-\frac{cx-1}{cx}} c x \sqrt{\frac{cx+1}{cx}}(6 \arcsin(cx) c^2 d - \sqrt{-c^2 x^2+1} e c x + \arcsin(cx) e)}{6 \sqrt{-c^2 x^2+1}}\right)}{c^2}}{c}$
default	$\frac{\frac{a(d c^3 x + \frac{1}{3} c^3 x^3 e)}{c^2} + \frac{b\left(\operatorname{arcsech}(cx) d c^3 x + \frac{\operatorname{arcsech}(cx) c^3 x^3 e}{3} + \frac{\sqrt{-\frac{cx-1}{cx}} c x \sqrt{\frac{cx+1}{cx}}(6 \arcsin(cx) c^2 d - \sqrt{-c^2 x^2+1} e c x + \arcsin(cx) e)}{6 \sqrt{-c^2 x^2+1}}\right)}{c^2}}{c}$

input

```
int((e*x^2+d)*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)
```

output

```
a*(1/3*x^3*e+d*x)+b/c*(1/3*c*arcsech(c*x)*x^3*e+arcsech(c*x)*d*c*x+1/6/c*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(6*arcsin(c*x)*c^2*d-(-c^2*x^2+1)^(1/2)*e*c*x+arcsin(c*x)*e)/(-c^2*x^2+1)^(1/2))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 209 vs. 2(64) = 128.

Time = 0.15 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.87

$$\int (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{2ac^3ex^3 - bc^2ex^2\sqrt{-\frac{c^2x^2-1}{c^2x^2}} + 6ac^3dx - 2(6bc^2d + be) \arctan\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{cx}\right) - 2(3bc^3d + bc^3e) \log\left(\dots\right)}{6c^3}$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output 
$$\frac{1}{6}(2ac^3ex^3 - bc^2e^2x^2\sqrt{-(c^2x^2 - 1)/(c^2x^2)} + 6a^3c^3d^2x - 2(6b^2c^2d + b^2e)\arctan((cx\sqrt{-(c^2x^2 - 1)/(c^2x^2)} - 1)/(cx)) - 2(3b^2c^3d + b^2c^3e)\log((cx\sqrt{-(c^2x^2 - 1)/(c^2x^2)} - 1)/x) + 2(b^2c^3e^2x^3 + 3b^2c^3d^2x - 3b^2c^3d - b^2c^3e)\log((cx\sqrt{-(c^2x^2 - 1)/(c^2x^2)} + 1)/(cx)))/c^3$$

### Sympy [F]

$$\int (d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx = \int (a + b\operatorname{asech}(cx)) (d + ex^2) dx$$

input `integrate((e*x**2+d)*(a+b*asech(c*x)),x)`

output `Integral((a + b*asech(c*x))*(d + e*x**2), x)`

### Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.96

$$\begin{aligned} & \int (d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx \\ &= \frac{1}{3} aex^3 + \frac{1}{6} \left( 2x^3 \operatorname{arsech}(cx) - \frac{\frac{\sqrt{\frac{1}{c^2x^2} - 1}}{c^2(\frac{1}{c^2x^2} - 1) + c^2} + \frac{\arctan(\sqrt{\frac{1}{c^2x^2} - 1})}{c^2}}{c} \right) be \\ &+ adx + \frac{(cx \operatorname{arsech}(cx) - \arctan(\sqrt{\frac{1}{c^2x^2} - 1}))bd}{c} \end{aligned}$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output

```
1/3*a*e*x^3 + 1/6*(2*x^3*arcsech(c*x) - (sqrt(1/(c^2*x^2) - 1)/(c^2*(1/(c^2*x^2) - 1) + c^2) + arctan(sqrt(1/(c^2*x^2) - 1))/c^2)/c)*b*e + a*d*x + (c*x*arcsech(c*x) - arctan(sqrt(1/(c^2*x^2) - 1)))*b*d/c
```

**Giac [F]**

$$\int (d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)(b\operatorname{ar} \operatorname{sech}(cx) + a) dx$$

input

```
integrate((e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)*(b*arcsech(c*x) + a), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int (d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d) \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input

```
int((d + e*x^2)*(a + b*acosh(1/(c*x))),x)
```

output

```
int((d + e*x^2)*(a + b*acosh(1/(c*x))), x)
```

**Reduce [F]**

$$\begin{aligned} \int (d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx &= \left( \int a\operatorname{sech}(cx) dx \right) bd \\ &+ \left( \int a\operatorname{sech}(cx) x^2 dx \right) be + adx + \frac{ae x^3}{3} \end{aligned}$$

input

```
int((e*x^2+d)*(a+b*asech(c*x)),x)
```

output `(3*int(asech(c*x),x)*b*d + 3*int(asech(c*x)*x**2,x)*b*e + 3*a*d*x + a*e*x*  
*3)/3`



**3.90**  $\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^2} dx$

Optimal result	728
Mathematica [A] (verified)	729
Rubi [A] (verified)	729
Maple [A] (verified)	731
Fricas [B] (verification not implemented)	732
Sympy [F]	732
Maxima [A] (verification not implemented)	733
Giac [F]	733
Mupad [B] (verification not implemented)	733
Reduce [F]	734

**Optimal result**

Integrand size = 19, antiderivative size = 96

$$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^2} dx = \frac{bd\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{x} - \frac{d(a+b\operatorname{sech}^{-1}(cx))}{x} + ex(a+b\operatorname{sech}^{-1}(cx)) + \frac{be\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\arcsin(cx)}{c}$$

output

```
b*d*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/x-d*(a+b*arcsech(c*x))/x+e*x*(a+b*arcsech(c*x))+b*e*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arcsin(c*x)/c
```

**Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.32

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^2} dx = -\frac{ad}{x} + aex + bd\left(c + \frac{1}{x}\right)\sqrt{\frac{1-cx}{1+cx}} - \frac{bd\operatorname{sech}^{-1}(cx)}{x} + bex\operatorname{sech}^{-1}(cx) + \frac{2be\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\arctan\left(\frac{\sqrt{1-c^2x^2}}{1-cx}\right)}{c-c^2x}$$

input `Integrate[((d + e*x^2)*(a + b*ArcSech[c*x]))/x^2,x]`

output `-((a*d)/x) + a*e*x + b*d*(c + x^(-1))*Sqrt[(1 - c*x)/(1 + c*x)] - (b*d*ArcSech[c*x])/x + b*e*x*ArcSech[c*x] + (2*b*e*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*ArcTan[Sqrt[1 - c^2*x^2]/(1 - c*x)])/(c - c^2*x)`

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.83, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {6855, 25, 358, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^2} dx$$

$$\downarrow 6855$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int -\frac{d - ex^2}{x^2\sqrt{1-c^2x^2}} dx - \frac{d(a + b\operatorname{sech}^{-1}(cx))}{x} + ex(a + b\operatorname{sech}^{-1}(cx))$$

$$\downarrow 25$$

$$-b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{d - ex^2}{x^2\sqrt{1-c^2x^2}} dx - \frac{d(a + b\operatorname{sech}^{-1}(cx))}{x} + ex(a + b\operatorname{sech}^{-1}(cx))$$



rule 6855

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))*((f_.)*(x_.))^(m_.)*((d_.) + (e_.)*(
x_)^2)^(p_.), x_Symbol] :=> With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Si
mp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)]
Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x]] /; Fre
eQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] &&
GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2
*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

### Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.17

method	result
parts	$a\left(ex - \frac{d}{x}\right) + bc\left(\frac{\operatorname{arcsech}(cx)xe}{c} - \frac{\operatorname{arcsech}(cx)d}{cx} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}\left(\sqrt{-c^2x^2+1}c^2d + \arcsin(cx)ecx\right)}{c^2\sqrt{-c^2x^2+1}}\right)$
derivativedivides	$c\left(\frac{a\left(cex - \frac{dc}{x}\right)}{c^2} + \frac{b\left(\operatorname{arcsech}(cx)ecx - \frac{\operatorname{arcsech}(cx)dc}{x} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}\left(\sqrt{-c^2x^2+1}c^2d + \arcsin(cx)ecx\right)}{\sqrt{-c^2x^2+1}}\right)}{c^2}\right)$
default	$c\left(\frac{a\left(cex - \frac{dc}{x}\right)}{c^2} + \frac{b\left(\operatorname{arcsech}(cx)ecx - \frac{\operatorname{arcsech}(cx)dc}{x} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}\left(\sqrt{-c^2x^2+1}c^2d + \arcsin(cx)ecx\right)}{\sqrt{-c^2x^2+1}}\right)}{c^2}\right)$

input `int((e*x^2+d)*(a+b*arcsech(c*x))/x^2,x,method=_RETURNVERBOSE)`

output `a*(e*x-d/x)+b*c*(1/c*arcsech(c*x)*x*e-arcsech(c*x)*d/c/x+1/c^2*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)*((-c^2*x^2+1)^(1/2)*c^2*d+arcsin(c*x)*e*c*x)/(-c^2*x^2+1)^(1/2))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 182 vs.  $2(54) = 108$ .

Time = 0.12 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.90

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^2} dx$$

$$= \frac{bc^2 dx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + acex^2 - 2bex \arctan\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - 1}{cx}\right) - acd + (bcd - bce)x \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - 1}{x}\right) + (b}{cx}$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x))/x^2,x, algorithm="fricas")`

output `(b*c^2*d*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + a*c*e*x^2 - 2*b*e*x*arctan((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/(c*x)) - a*c*d + (b*c*d - b*c*e)*x*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) + (b*c*e*x^2 - b*c*d + (b*c*d - b*c*e)*x)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)))/(c*x)`

**Sympy [F]**

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{(a + b\operatorname{asech}(cx))(d + ex^2)}{x^2} dx$$

input `integrate((e*x**2+d)*(a+b*asech(c*x))/x**2,x)`

output `Integral((a + b*asech(c*x))*(d + e*x**2)/x**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.69

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^2} dx = \left( c \sqrt{\frac{1}{c^2 x^2} - 1} - \frac{\operatorname{arsech}(cx)}{x} \right) bd + aex$$

$$+ \frac{\left( cx \operatorname{arsech}(cx) - \arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right) \right) be}{c} - \frac{ad}{x}$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x))/x^2,x, algorithm="maxima")`output `(c*sqrt(1/(c^2*x^2) - 1) - arcsech(c*x)/x)*b*d + a*e*x + (c*x*arcsech(c*x) - arctan(sqrt(1/(c^2*x^2) - 1)))*b*e/c - a*d/x`**Giac [F]**

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{(ex^2 + d)(b \operatorname{arsech}(cx) + a)}{x^2} dx$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x))/x^2,x, algorithm="giac")`output `integrate((e*x^2 + d)*(b*arcsech(c*x) + a)/x^2, x)`**Mupad [B] (verification not implemented)**

Time = 4.34 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.02

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^2} dx = aex - \frac{ad}{x}$$

$$+ bcd \left( \sqrt{\frac{1}{cx} - 1} \sqrt{\frac{1}{cx} + 1} - \frac{\operatorname{acosh}\left(\frac{1}{cx}\right)}{cx} \right)$$

$$+ \frac{be \operatorname{atan}\left(\frac{1}{\sqrt{\frac{1}{cx} - 1} \sqrt{\frac{1}{cx} + 1}}\right)}{c} + bex \operatorname{acosh}\left(\frac{1}{cx}\right)$$

input `int(((d + e*x^2)*(a + b*acosh(1/(c*x))))/x^2,x)`

output `a*e*x - (a*d)/x + b*c*d*((1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2) - acosh(1/(c*x)))/(c*x) + (b*e*atan(1/((1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2))))/c + b*e*x*acosh(1/(c*x))`

### Reduce [F]

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^2} dx$$

$$= \frac{\left(\int a\operatorname{sech}(cx) dx\right) bex + \left(\int \frac{a\operatorname{sech}(cx)}{x^2} dx\right) bdx - ad + aex^2}{x}$$

input `int((e*x^2+d)*(a+b*asech(c*x))/x^2,x)`

output `(int(asech(c*x),x)*b*e*x + int(asech(c*x)/x**2,x)*b*d*x - a*d + a*e*x**2)/x`

**3.91**  $\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx$

Optimal result	735
Mathematica [A] (verified)	736
Rubi [A] (verified)	736
Maple [A] (verified)	738
Fricas [A] (verification not implemented)	739
Sympy [F]	739
Maxima [A] (verification not implemented)	739
Giac [F]	740
Mupad [F(-1)]	740
Reduce [F]	741

**Optimal result**

Integrand size = 19, antiderivative size = 126

$$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx = \frac{bd\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{9x^3} + \frac{b(2c^2d+9e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{9x} - \frac{d(a+b\operatorname{sech}^{-1}(cx))}{3x^3} - \frac{e(a+b\operatorname{sech}^{-1}(cx))}{x}$$

output

```
1/9*b*d*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/x^3+1/9*b*(2*c^2*d+9*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/x-1/3*d*(a+b*arcsech(c*x))/x^3-e*(a+b*arcsech(c*x))/x
```



**Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.60

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^4} dx$$

$$= \frac{-3a(d + 3ex^2) + b\sqrt{\frac{1-cx}{1+cx}}(1 + cx)(d + 2c^2dx^2 + 9ex^2) - 3b(d + 3ex^2)\operatorname{sech}^{-1}(cx)}{9x^3}$$

input

```
Integrate[((d + e*x^2)*(a + b*ArcSech[c*x]))/x^4,x]
```

output

```
(-3*a*(d + 3*e*x^2) + b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(d + 2*c^2*d*x^2 + 9*e*x^2) - 3*b*(d + 3*e*x^2)*ArcSech[c*x])/(9*x^3)
```

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.87, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {6855, 27, 359, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^4} dx$$

$$\downarrow \text{6855}$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int -\frac{3ex^2 + d}{3x^4\sqrt{1-c^2x^2}} dx - \frac{d(a + b\operatorname{sech}^{-1}(cx))}{3x^3} - \frac{e(a + b\operatorname{sech}^{-1}(cx))}{x}$$

$$\downarrow \text{27}$$

$$-\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{3ex^2 + d}{x^4\sqrt{1-c^2x^2}} dx - \frac{d(a + b\operatorname{sech}^{-1}(cx))}{3x^3} - \frac{e(a + b\operatorname{sech}^{-1}(cx))}{x}$$

$$\downarrow \text{359}$$

$$\begin{aligned}
& -\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}(2c^2d+9e)\int\frac{1}{x^2\sqrt{1-c^2x^2}}dx-\frac{d\sqrt{1-c^2x^2}}{3x^3}\right)- \\
& \frac{d(a+b\operatorname{sech}^{-1}(cx))}{3x^3}-\frac{e(a+b\operatorname{sech}^{-1}(cx))}{x} \\
& \quad \downarrow 242 \\
& -\frac{d(a+b\operatorname{sech}^{-1}(cx))}{3x^3}-\frac{e(a+b\operatorname{sech}^{-1}(cx))}{x}- \\
& \frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{\sqrt{1-c^2x^2}(2c^2d+9e)}{3x}-\frac{d\sqrt{1-c^2x^2}}{3x^3}\right)
\end{aligned}$$

input `Int[((d + e*x^2)*(a + b*ArcSech[c*x]))/x^4,x]`

output `-1/3*(b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/3*(d*Sqrt[1 - c^2*x^2])/x^3 - ((2*c^2*d + 9*e)*Sqrt[1 - c^2*x^2])/(3*x))) - (d*(a + b*ArcSech[c*x]))/(3*x^3) - (e*(a + b*ArcSech[c*x]))/x`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 242 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m+1)*((a + b*x^2)^(p+1)/(a*c*(m+1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

rule 359 `Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[c*(e*x)^(m+1)*((a + b*x^2)^(p+1)/(a*e*(m+1))), x] + Simp[(a*d*(m+1) - b*c*(m+2*p+3))/(a*e^2*(m+1)) Int[(e*x)^(m+2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && !ILtQ[p, -1]`

rule 6855

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))*((f_.)*(x_.))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

**Maple [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.87

method	result	size
parts	$a\left(-\frac{d}{3x^3} - \frac{e}{x}\right) + bc^3\left(-\frac{\operatorname{arcsech}(cx)d}{3c^3x^3} - \frac{\operatorname{arcsech}(cx)e}{c^3x} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}(2c^4dx^2+9e^2x^2+c^2d)}{9c^4x^2}\right)$	11
derivativedivides	$c^3\left(\frac{a\left(-\frac{d}{3cx^3}-\frac{e}{cx}\right)}{c^2} + \frac{b\left(-\frac{\operatorname{arcsech}(cx)d}{3cx^3}-\frac{\operatorname{arcsech}(cx)e}{cx} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}(2c^4dx^2+9e^2x^2+c^2d)}{9c^2x^2}\right)}{c^2}\right)$	12
default	$c^3\left(\frac{a\left(-\frac{d}{3cx^3}-\frac{e}{cx}\right)}{c^2} + \frac{b\left(-\frac{\operatorname{arcsech}(cx)d}{3cx^3}-\frac{\operatorname{arcsech}(cx)e}{cx} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}(2c^4dx^2+9e^2x^2+c^2d)}{9c^2x^2}\right)}{c^2}\right)$	12

input

```
int((e*x^2+d)*(a+b*arcsech(c*x))/x^4,x,method=_RETURNVERBOSE)
```

output

```
a*(-1/3*d/x^3-e/x)+b*c^3*(-1/3*arcsech(c*x)*d/c^3/x^3-1/c^3*arcsech(c*x)*e/x+1/9/c^4*(-(c*x-1)/c/x)^(1/2)/x^2*((c*x+1)/c/x)^(1/2)*(2*c^4*d*x^2+9*c^2*e*x^2+c^2*d))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.84

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^4} dx = \frac{9 a e x^2 + 3 a d + 3 (3 b e x^2 + b d) \log\left(\frac{c x \sqrt{-\frac{c^2 x^2}{c^2 x^2} - 1} + 1}{c x}\right) - (b c d x + (2 b c^3 d + 9 b c e) x^3) \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}}}{9 x^3}$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x))/x^4,x, algorithm="fricas")`

output `-1/9*(9*a*e*x^2 + 3*a*d + 3*(3*b*e*x^2 + b*d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (b*c*d*x + (2*b*c^3*d + 9*b*c*e)*x^3)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/x^3`

**Sympy [F]**

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{(a + b \operatorname{asech}(cx))(d + ex^2)}{x^4} dx$$

input `integrate((e*x**2+d)*(a+b*asech(c*x))/x**4,x)`

output `Integral((a + b*asech(c*x))*(d + e*x**2)/x**4, x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.72

$$\begin{aligned} & \int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^4} dx \\ &= \left( c \sqrt{\frac{1}{c^2 x^2} - 1} - \frac{\operatorname{arsech}(cx)}{x} \right) b e \\ &+ \frac{1}{9} b d \left( \frac{c^4 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{3}{2}} + 3 c^4 \sqrt{\frac{1}{c^2 x^2} - 1}}{c} - \frac{3 \operatorname{arsech}(cx)}{x^3} \right) - \frac{a e}{x} - \frac{a d}{3 x^3} \end{aligned}$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x))/x^4,x, algorithm="maxima")`

output `(c*sqrt(1/(c^2*x^2) - 1) - arcsech(c*x)/x)*b*e + 1/9*b*d*((c^4*(1/(c^2*x^2) - 1)^(3/2) + 3*c^4*sqrt(1/(c^2*x^2) - 1))/c - 3*arcsech(c*x)/x^3) - a*e/x - 1/3*a*d/x^3`

### Giac [F]

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{(ex^2 + d)(b \operatorname{ar} \operatorname{sech}(cx) + a)}{x^4} dx$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x))/x^4,x, algorithm="giac")`

output `integrate((e*x^2 + d)*(b*arcsech(c*x) + a)/x^4, x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{(ex^2 + d)(a + b \operatorname{acosh}(\frac{1}{cx}))}{x^4} dx$$

input `int(((d + e*x^2)*(a + b*acosh(1/(c*x))))/x^4,x)`

output `int(((d + e*x^2)*(a + b*acosh(1/(c*x))))/x^4, x)`

**Reduce [F]**

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^4} dx$$

$$= \frac{3\left(\int \frac{\operatorname{asech}(cx)}{x^4} dx\right)bdx^3 + 3\left(\int \frac{\operatorname{asech}(cx)}{x^2} dx\right)be x^3 - ad - 3aex^2}{3x^3}$$

input `int((e*x^2+d)*(a+b*asech(c*x))/x^4,x)`

output `(3*int(asech(c*x)/x**4,x)*b*d*x**3 + 3*int(asech(c*x)/x**2,x)*b*e*x**3 - a*d - 3*a*e*x**2)/(3*x**3)`

**3.92** 
$$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx$$

Optimal result	742
Mathematica [A] (verified)	743
Rubi [A] (verified)	743
Maple [A] (verified)	745
Fricas [A] (verification not implemented)	746
Sympy [F]	746
Maxima [A] (verification not implemented)	747
Giac [F]	747
Mupad [F(-1)]	748
Reduce [F]	748

**Optimal result**

Integrand size = 19, antiderivative size = 183

$$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx = \frac{bd\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{25x^5} + \frac{b(12c^2d+25e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{225x^3} + \frac{2bc^2(12c^2d+25e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{225x} - \frac{d(a+b\operatorname{sech}^{-1}(cx))}{5x^5} - \frac{e(a+b\operatorname{sech}^{-1}(cx))}{3x^3}$$

output

```
1/25*b*d*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/x^5+1/225*b*(1
2*c^2*d+25*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/x^3+2/225
*b*c^2*(12*c^2*d+25*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/
x-1/5*d*(a+b*arcsech(c*x))/x^5-1/3*e*(a+b*arcsech(c*x))/x^3
```

**Mathematica [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.55

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^6} dx$$

$$= \frac{-15a(3d + 5ex^2) + b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(25ex^2(1+2c^2x^2) + 3d(3+4c^2x^2+8c^4x^4)) - 15b(3d+5ex^2) \operatorname{sech}^{-1}(cx)}{225x^5}$$

input

```
Integrate[((d + e*x^2)*(a + b*ArcSech[c*x]))/x^6,x]
```

output

```
(-15*a*(3*d + 5*e*x^2) + b*sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(25*e*x^2*(1 + 2*c^2*x^2) + 3*d*(3 + 4*c^2*x^2 + 8*c^4*x^4)) - 15*b*(3*d + 5*e*x^2)*ArcSech[c*x])/(225*x^5)
```

**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.77, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {6855, 27, 359, 245, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^6} dx$$

$$\downarrow 6855$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int -\frac{5ex^2+3d}{15x^6\sqrt{1-c^2x^2}} dx - \frac{d(a+b\operatorname{sech}^{-1}(cx))}{5x^5} - \frac{e(a+b\operatorname{sech}^{-1}(cx))}{3x^3}$$

$$\downarrow 27$$

$$-\frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{5ex^2+3d}{x^6\sqrt{1-c^2x^2}} dx - \frac{d(a+b\operatorname{sech}^{-1}(cx))}{5x^5} - \frac{e(a+b\operatorname{sech}^{-1}(cx))}{3x^3}$$

$$\downarrow 359$$



$$\begin{aligned}
& -\frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}(12c^2d+25e)\int\frac{1}{x^4\sqrt{1-c^2x^2}}dx-\frac{3d\sqrt{1-c^2x^2}}{5x^5}\right)- \\
& \quad \frac{d(a+b\operatorname{sech}^{-1}(cx))}{5x^5}-\frac{e(a+b\operatorname{sech}^{-1}(cx))}{3x^3} \\
& \quad \downarrow 245 \\
& -\frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}(12c^2d+25e)\left(\frac{2}{3}c^2\int\frac{1}{x^2\sqrt{1-c^2x^2}}dx-\frac{\sqrt{1-c^2x^2}}{3x^3}\right)-\frac{3d\sqrt{1-c^2x^2}}{5x^5}\right)- \\
& \quad \frac{d(a+b\operatorname{sech}^{-1}(cx))}{5x^5}-\frac{e(a+b\operatorname{sech}^{-1}(cx))}{3x^3} \\
& \quad \downarrow 242 \\
& -\frac{d(a+b\operatorname{sech}^{-1}(cx))}{5x^5}-\frac{e(a+b\operatorname{sech}^{-1}(cx))}{3x^3}- \\
& \frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\left(-\frac{2c^2\sqrt{1-c^2x^2}}{3x}-\frac{\sqrt{1-c^2x^2}}{3x^3}\right)(12c^2d+25e)-\frac{3d\sqrt{1-c^2x^2}}{5x^5}\right)
\end{aligned}$$

input `Int[((d + e*x^2)*(a + b*ArcSech[c*x]))/x^6,x]`

output `-1/15*(b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-3*d*Sqrt[1 - c^2*x^2])/(5*x^5) + ((12*c^2*d + 25*e)*(-1/3*Sqrt[1 - c^2*x^2]/x^3 - (2*c^2*Sqrt[1 - c^2*x^2])/(3*x)))/5) - (d*(a + b*ArcSech[c*x]))/(5*x^5) - (e*(a + b*ArcSech[c*x]))/(3*x^3)`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] :> Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 242 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

```
rule 245 Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[x^(m + 1)*((a +
b*x^2)^(p + 1)/(a*(m + 1))), x] - Simp[b*(m + 2*(p + 1) + 1)/(a*(m + 1)))
Int[x^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, m, p}, x] && ILtQ[Simplify[(m + 1)/2 + p + 1], 0] && NeQ[m, -1]
```

```
rule 359 Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2), x
_Symbol] := Simp[c*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*e*(m + 1))), x] +
Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(a*e^2*(m + 1)) Int[(e*x)^(m + 2)*
(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0]
&& LtQ[m, -1] && !ILtQ[p, -1]
```

```
rule 6855 Int[((a_) + ArcSech[(c_)*(x_)]*(b_))*((f_)*(x_))^(m_)*((d_) + (e_)*(
x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Si
mp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)]
Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; Fre
eQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] &&
GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2
*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

### Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.70

method	result
parts	$a\left(-\frac{e}{3x^3} - \frac{d}{5x^5}\right) + bc^5\left(-\frac{\operatorname{arcsech}(cx)e}{3c^5x^3} - \frac{\operatorname{arcsech}(cx)d}{5c^5x^5} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}(24c^6dx^4+50c^4ex^4+12c^4dx^2)}{225c^6x^4}\right)$
derivativedivides	$c^5\left(\frac{a\left(-\frac{d}{5c^3x^5} - \frac{e}{3c^3x^3}\right)}{c^2} + \frac{b\left(-\frac{\operatorname{arcsech}(cx)d}{5c^3x^5} - \frac{\operatorname{arcsech}(cx)e}{3c^3x^3} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}(24c^6dx^4+50c^4ex^4+12c^4dx^2+25e^2x^2)}{225c^4x^4}\right)}{c^2}\right)$
default	$c^5\left(\frac{a\left(-\frac{d}{5c^3x^5} - \frac{e}{3c^3x^3}\right)}{c^2} + \frac{b\left(-\frac{\operatorname{arcsech}(cx)d}{5c^3x^5} - \frac{\operatorname{arcsech}(cx)e}{3c^3x^3} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}(24c^6dx^4+50c^4ex^4+12c^4dx^2+25e^2x^2)}{225c^4x^4}\right)}{c^2}\right)$

```
input int((e*x^2+d)*(a+b*arcsech(c*x))/x^6,x,method=_RETURNVERBOSE)
```

output

```
a*(-1/3*e/x^3-1/5*d/x^5)+b*c^5*(-1/3/c^5*arcsech(c*x)*e/x^3-1/5*arcsech(c*x)*d/c^5/x^5+1/225/c^6*(-(c*x-1)/c/x)^(1/2)/x^4*((c*x+1)/c/x)^(1/2)*(24*c^6*d*x^4+50*c^4*e*x^4+12*c^4*d*x^2+25*c^2*e*x^2+9*c^2*d))
```

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.70

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^6} dx = \frac{75 aex^2 + 45 ad + 15(5 bex^2 + 3 bd) \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}+1}}{cx}\right) - (2(12bc^5d + 25bc^3e)x^5 + 9bcdx + (12bc^3d + 12bc^3e)x^3)\sqrt{-\frac{c^2x^2-1}{c^2x^2}}}{225x^5}$$

input

```
integrate((e*x^2+d)*(a+b*arcsech(c*x))/x^6,x, algorithm="fricas")
```

output

```
-1/225*(75*a*e*x^2 + 45*a*d + 15*(5*b*e*x^2 + 3*b*d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (2*(12*b*c^5*d + 25*b*c^3*e)*x^5 + 9*b*c*d*x + (12*b*c^3*d + 25*b*c^3*e)*x^3)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2))/x^5
```

**Sympy [F]**

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^6} dx = \int \frac{(a + b \operatorname{asech}(cx))(d + ex^2)}{x^6} dx$$

input

```
integrate((e*x**2+d)*(a+b*asech(c*x))/x**6,x)
```

output

```
Integral((a + b*asech(c*x))*(d + e*x**2)/x**6, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.72

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^6} dx$$

$$= \frac{1}{75} bd \left( \frac{3c^6 \left(\frac{1}{c^2 x^2} - 1\right)^{\frac{5}{2}} + 10c^6 \left(\frac{1}{c^2 x^2} - 1\right)^{\frac{3}{2}} + 15c^6 \sqrt{\frac{1}{c^2 x^2} - 1}}{c} - \frac{15 \operatorname{arsech}(cx)}{x^5} \right)$$

$$+ \frac{1}{9} be \left( \frac{c^4 \left(\frac{1}{c^2 x^2} - 1\right)^{\frac{3}{2}} + 3c^4 \sqrt{\frac{1}{c^2 x^2} - 1}}{c} - \frac{3 \operatorname{arsech}(cx)}{x^3} \right) - \frac{ae}{3x^3} - \frac{ad}{5x^5}$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x))/x^6,x, algorithm="maxima")`

output `1/75*b*d*((3*c^6*(1/(c^2*x^2) - 1)^(5/2) + 10*c^6*(1/(c^2*x^2) - 1)^(3/2) + 15*c^6*sqrt(1/(c^2*x^2) - 1))/c - 15*arcsech(c*x)/x^5) + 1/9*b*e*((c^4*(1/(c^2*x^2) - 1)^(3/2) + 3*c^4*sqrt(1/(c^2*x^2) - 1))/c - 3*arcsech(c*x)/x^3) - 1/3*a*e/x^3 - 1/5*a*d/x^5`

**Giac [F]**

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^6} dx = \int \frac{(ex^2 + d)(b \operatorname{arsech}(cx) + a)}{x^6} dx$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x))/x^6,x, algorithm="giac")`

output `integrate((e*x^2 + d)*(b*arcsech(c*x) + a)/x^6, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^6} dx = \int \frac{(ex^2 + d)(a + b\operatorname{acosh}(\frac{1}{cx}))}{x^6} dx$$

input `int(((d + e*x^2)*(a + b*acosh(1/(c*x))))/x^6,x)`output `int(((d + e*x^2)*(a + b*acosh(1/(c*x))))/x^6, x)`**Reduce [F]**

$$\begin{aligned} & \int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^6} dx \\ &= \frac{15\left(\int \frac{\operatorname{asech}(cx)}{x^6} dx\right)bdx^5 + 15\left(\int \frac{\operatorname{asech}(cx)}{x^4} dx\right)be x^5 - 3ad - 5ae x^2}{15x^5} \end{aligned}$$

input `int((e*x^2+d)*(a+b*asech(c*x))/x^6,x)`output `(15*int(asech(c*x)/x**6,x)*b*d*x**5 + 15*int(asech(c*x)/x**4,x)*b*e*x**5 - 3*a*d - 5*a*e*x**2)/(15*x**5)`

**3.93**  $\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^8} dx$

Optimal result	749
Mathematica [A] (verified)	750
Rubi [A] (verified)	750
Maple [A] (verified)	753
Fricas [A] (verification not implemented)	753
Sympy [F]	754
Maxima [A] (verification not implemented)	754
Giac [F]	755
Mupad [F(-1)]	755
Reduce [F]	756

**Optimal result**

Integrand size = 19, antiderivative size = 238

$$\int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^8} dx = \frac{bd\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{49x^7} + \frac{b(30c^2d+49e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{1225x^5} + \frac{4bc^2(30c^2d+49e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{3675x^3} + \frac{8bc^4(30c^2d+49e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{3675x} - \frac{d(a+b\operatorname{sech}^{-1}(cx))}{7x^7} - \frac{e(a+b\operatorname{sech}^{-1}(cx))}{5x^5}$$

output

```
1/49*b*d*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/x^7+1/1225*b*(
30*c^2*d+49*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/x^5+4/36
75*b*c^2*(30*c^2*d+49*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2
)/x^3+8/3675*b*c^4*(30*c^2*d+49*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x
^2+1)^(1/2)/x-1/7*d*(a+b*arcsech(c*x))/x^7-1/5*e*(a+b*arcsech(c*x))/x^5
```

**Mathematica [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.49

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^8} dx$$

$$= \frac{-105a(5d + 7ex^2) + b\sqrt{\frac{1-cx}{1+cx}}(1 + cx)(49ex^2(3 + 4c^2x^2 + 8c^4x^4) + 15d(5 + 6c^2x^2 + 8c^4x^4 + 16c^6x^6))}{3675x^7}$$

input

```
Integrate[((d + e*x^2)*(a + b*ArcSech[c*x]))/x^8,x]
```

output

```
(-105*a*(5*d + 7*e*x^2) + b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(49*e*x^2*(3 + 4*c^2*x^2 + 8*c^4*x^4) + 15*d*(5 + 6*c^2*x^2 + 8*c^4*x^4 + 16*c^6*x^6)) - 105*b*(5*d + 7*e*x^2)*ArcSech[c*x])/(3675*x^7)
```

**Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.71, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$ , Rules used = {6855, 27, 359, 245, 245, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^8} dx$$

$$\downarrow \text{6855}$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int -\frac{7ex^2 + 5d}{35x^8\sqrt{1-c^2x^2}} dx - \frac{d(a + b\operatorname{sech}^{-1}(cx))}{7x^7} - \frac{e(a + b\operatorname{sech}^{-1}(cx))}{5x^5}$$

$$\downarrow \text{27}$$

$$-\frac{1}{35}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{7ex^2 + 5d}{x^8\sqrt{1-c^2x^2}} dx - \frac{d(a + b\operatorname{sech}^{-1}(cx))}{7x^7} - \frac{e(a + b\operatorname{sech}^{-1}(cx))}{5x^5}$$

$$\downarrow \text{359}$$

$$\begin{aligned}
& -\frac{1}{35}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{7}(30c^2d+49e)\int\frac{1}{x^6\sqrt{1-c^2x^2}}dx-\frac{5d\sqrt{1-c^2x^2}}{7x^7}\right)- \\
& \quad \frac{d(a+b\operatorname{sech}^{-1}(cx))}{7x^7}-\frac{e(a+b\operatorname{sech}^{-1}(cx))}{5x^5} \\
& \quad \downarrow 245 \\
& -\frac{1}{35}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{7}(30c^2d+49e)\left(\frac{4}{5}c^2\int\frac{1}{x^4\sqrt{1-c^2x^2}}dx-\frac{\sqrt{1-c^2x^2}}{5x^5}\right)-\frac{5d\sqrt{1-c^2x^2}}{7x^7}\right)- \\
& \quad \frac{d(a+b\operatorname{sech}^{-1}(cx))}{7x^7}-\frac{e(a+b\operatorname{sech}^{-1}(cx))}{5x^5} \\
& \quad \downarrow 245 \\
& -\frac{1}{35}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{7}(30c^2d+49e)\left(\frac{4}{5}c^2\left(\frac{2}{3}c^2\int\frac{1}{x^2\sqrt{1-c^2x^2}}dx-\frac{\sqrt{1-c^2x^2}}{3x^3}\right)-\frac{\sqrt{1-c^2x^2}}{5x^5}\right)-\frac{5d\sqrt{1-c^2x^2}}{7x^7}\right)- \\
& \quad \frac{d(a+b\operatorname{sech}^{-1}(cx))}{7x^7}-\frac{e(a+b\operatorname{sech}^{-1}(cx))}{5x^5} \\
& \quad \downarrow 242 \\
& \quad -\frac{d(a+b\operatorname{sech}^{-1}(cx))}{7x^7}-\frac{e(a+b\operatorname{sech}^{-1}(cx))}{5x^5}- \\
& \frac{1}{35}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{7}\left(\frac{4}{5}c^2\left(-\frac{2c^2\sqrt{1-c^2x^2}}{3x}-\frac{\sqrt{1-c^2x^2}}{3x^3}\right)-\frac{\sqrt{1-c^2x^2}}{5x^5}\right)(30c^2d+49e)-\frac{5d\sqrt{1-c^2x^2}}{7x^7}\right)
\end{aligned}$$

input `Int[((d + e*x^2)*(a + b*ArcSech[c*x]))/x^8,x]`

output `-1/35*(b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-5*d*Sqrt[1 - c^2*x^2])/(7*x^7) + ((30*c^2*d + 49*e)*(-1/5*Sqrt[1 - c^2*x^2]/x^5 + (4*c^2*(-1/3*Sqrt[1 - c^2*x^2]/x^3 - (2*c^2*Sqrt[1 - c^2*x^2])/(3*x)))/5))/7) - (d*(a + b*ArcSech[c*x]))/(7*x^7) - (e*(a + b*ArcSech[c*x]))/(5*x^5)`



## Definitions of rubi rules used

- rule 27 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx) /; FreeQ[b, x]]`
- rule 242 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`
- rule 245 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[x^(m + 1)*((a + b*x^2)^(p + 1)/(a*(m + 1))), x] - Simp[b*(m + 2*(p + 1) + 1)/(a*(m + 1)) Int[x^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, m, p}, x] && ILtQ[Simplify[(m + 1)/2 + p + 1], 0] && NeQ[m, -1]`
- rule 359 `Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[c*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*e*(m + 1))), x] + Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(a*e^2*(m + 1)) Int[(e*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && !ILtQ[p, -1]`
- rule 6855 `Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))`

### Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.62

method	result
parts	$a\left(-\frac{d}{7x^7} - \frac{e}{5x^5}\right) + bc^7\left(-\frac{\operatorname{arcsech}(cx)d}{7c^7x^7} - \frac{\operatorname{arcsech}(cx)e}{5c^7x^5} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}(240c^8dx^6+392c^6ex^6+120c^6d^4+196c^4e^2+75c^2d^2)}{3675c^8x^6}\right)$
derivativedivides	$c^7\left(\frac{a\left(-\frac{d}{7c^5x^7}-\frac{e}{5c^5x^5}\right)}{c^2} + \frac{b\left(-\frac{\operatorname{arcsech}(cx)d}{7c^5x^7}-\frac{\operatorname{arcsech}(cx)e}{5c^5x^5} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}(240c^8dx^6+392c^6ex^6+120c^6d^4+196c^4e^2+75c^2d^2)}{3675c^6x^6}\right)}{c^2}\right)$
default	$c^7\left(\frac{a\left(-\frac{d}{7c^5x^7}-\frac{e}{5c^5x^5}\right)}{c^2} + \frac{b\left(-\frac{\operatorname{arcsech}(cx)d}{7c^5x^7}-\frac{\operatorname{arcsech}(cx)e}{5c^5x^5} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}(240c^8dx^6+392c^6ex^6+120c^6d^4+196c^4e^2+75c^2d^2)}{3675c^6x^6}\right)}{c^2}\right)$

input `int((e*x^2+d)*(a+b*arcsech(c*x))/x^8,x,method=_RETURNVERBOSE)`

output `a*(-1/7*d/x^7-1/5*e/x^5)+b*c^7*(-1/7*arcsech(c*x)*d/c^7/x^7-1/5/c^7*arcsech(c*x)*e/x^5+1/3675/c^8*(-(c*x-1)/c/x)^(1/2)/x^6*((c*x+1)/c/x)^(1/2)*(240*c^8*d*x^6+392*c^6*e*x^6+120*c^6*d*x^4+196*c^4*e*x^4+90*c^4*d*x^2+147*c^2*e*x^2+75*c^2*d))`

### Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.63

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^8} dx = \frac{735 aex^2 + 525 ad + 105(7bex^2 + 5bd) \log\left(\frac{cx\sqrt{-\frac{e^2x^2-1}{c^2x^2}}+1}{cx}\right) - (8(30bc^7d + 49bc^5e)x^7 + 4(30bc^5d + 3675x^7))}{3675x^7}$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x))/x^8,x, algorithm="fricas")`

output

```
-1/3675*(735*a*e*x^2 + 525*a*d + 105*(7*b*e*x^2 + 5*b*d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (8*(30*b*c^7*d + 49*b*c^5*e)*x^7 + 4*(30*b*c^5*d + 49*b*c^3*e)*x^5 + 75*b*c*d*x + 3*(30*b*c^3*d + 49*b*c*e)*x^3)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2))/x^7
```

**Sympy [F]**

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^8} dx = \int \frac{(a + b \operatorname{asech}(cx))(d + ex^2)}{x^8} dx$$

input

```
integrate((e*x**2+d)*(a+b*asech(c*x))/x**8,x)
```

output

```
Integral((a + b*asech(c*x))*(d + e*x**2)/x**8, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.69

$$\begin{aligned} & \int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^8} dx \\ &= \frac{1}{245} bd \left( \frac{5c^8 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{7}{2}} + 21c^8 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{5}{2}} + 35c^8 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{3}{2}} + 35c^8 \sqrt{\frac{1}{c^2x^2} - 1}}{c} - \frac{35 \operatorname{arsech}(cx)}{x^7} \right) \\ &+ \frac{1}{75} be \left( \frac{3c^6 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{5}{2}} + 10c^6 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{3}{2}} + 15c^6 \sqrt{\frac{1}{c^2x^2} - 1}}{c} - \frac{15 \operatorname{arsech}(cx)}{x^5} \right) \\ &- \frac{ae}{5x^5} - \frac{ad}{7x^7} \end{aligned}$$

input

```
integrate((e*x^2+d)*(a+b*arcsech(c*x))/x^8,x, algorithm="maxima")
```

output

```
1/245*b*d*((5*c^8*(1/(c^2*x^2) - 1)^(7/2) + 21*c^8*(1/(c^2*x^2) - 1)^(5/2)
+ 35*c^8*(1/(c^2*x^2) - 1)^(3/2) + 35*c^8*sqrt(1/(c^2*x^2) - 1))/c - 35*a
rcsech(c*x)/x^7) + 1/75*b*e*((3*c^6*(1/(c^2*x^2) - 1)^(5/2) + 10*c^6*(1/(c
^2*x^2) - 1)^(3/2) + 15*c^6*sqrt(1/(c^2*x^2) - 1))/c - 15*arcsech(c*x)/x^5
) - 1/5*a*e/x^5 - 1/7*a*d/x^7
```

**Giac [F]**

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^8} dx = \int \frac{(ex^2 + d)(b \operatorname{ar} \operatorname{sech}(cx) + a)}{x^8} dx$$

input

```
integrate((e*x^2+d)*(a+b*arcsech(c*x))/x^8,x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)*(b*arcsech(c*x) + a)/x^8, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^8} dx = \int \frac{(ex^2 + d)(a + b \operatorname{acosh}(\frac{1}{cx}))}{x^8} dx$$

input

```
int(((d + e*x^2)*(a + b*acosh(1/(c*x))))/x^8,x)
```

output

```
int(((d + e*x^2)*(a + b*acosh(1/(c*x))))/x^8, x)
```

**Reduce [F]**

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^8} dx$$

$$= \frac{35 \left( \int \frac{\operatorname{asech}(cx)}{x^8} dx \right) bd x^7 + 35 \left( \int \frac{\operatorname{asech}(cx)}{x^6} dx \right) be x^7 - 5ad - 7ae x^2}{35x^7}$$

input `int((e*x^2+d)*(a+b*asech(c*x))/x^8,x)`

output `(35*int(asech(c*x)/x**8,x)*b*d*x**7 + 35*int(asech(c*x)/x**6,x)*b*e*x**7 - 5*a*d - 7*a*e*x**2)/(35*x**7)`

### 3.94 $\int x^5(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	757
Mathematica [A] (verified)	758
Rubi [A] (verified)	758
Maple [A] (verified)	760
Fricas [A] (verification not implemented)	761
Sympy [A] (verification not implemented)	761
Maxima [A] (verification not implemented)	762
Giac [F]	762
Mupad [F(-1)]	763
Reduce [F]	763

#### Optimal result

Integrand size = 19, antiderivative size = 232

$$\int x^5(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx = -\frac{b(4c^2d + 3e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{24c^8} + \frac{b(8c^2d + 9e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} (1-c^2x^2)^{3/2}}{72c^8} - \frac{b(4c^2d + 9e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} (1-c^2x^2)^{5/2}}{120c^8} + \frac{be \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} (1-c^2x^2)^{7/2}}{56c^8} + \frac{1}{6} dx^6 (a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{8} ex^8 (a + b\operatorname{sech}^{-1}(cx))$$

output

```
-1/24*b*(4*c^2*d+3*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^8+1/72*b*(8*c^2*d+9*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(3/2)/c^8-1/120*b*(4*c^2*d+9*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(5/2)/c^8+1/56*b*e*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(7/2)/c^8+1/6*d*x^6*(a+b*arcsech(c*x))+1/8*e*x^8*(a+b*arcsech(c*x))
```

**Mathematica [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.54

$$\int x^5 (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx = \frac{1}{24} ax^6 (4d + 3ex^2) - \frac{b \sqrt{\frac{1-cx}{1+cx}} (1+cx) (144e + 8c^2(28d + 9ex^2) + 2c^4(56dx^2 + 27ex^4) + c^6(84dx^4 + 45ex^6))}{2520c^8} + \frac{1}{24} bx^6 (4d + 3ex^2) \operatorname{sech}^{-1}(cx)$$

input

```
Integrate[x^5*(d + e*x^2)*(a + b*ArcSech[c*x]),x]
```

output

```
(a*x^6*(4*d + 3*e*x^2))/24 - (b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(144*e + 8*c^2*(28*d + 9*e*x^2) + 2*c^4*(56*d*x^2 + 27*e*x^4) + c^6*(84*d*x^4 + 45*e*x^6)))/(2520*c^8) + (b*x^6*(4*d + 3*e*x^2)*ArcSech[c*x])/24
```

**Rubi [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.74, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {6855, 27, 354, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^5 (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx$$

↓ 6855

$$b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{x^5 (3ex^2 + 4d)}{24\sqrt{1-c^2x^2}} dx + \frac{1}{6} dx^6 (a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{8} ex^8 (a + b \operatorname{sech}^{-1}(cx))$$

↓ 27

$$\frac{1}{24} b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{x^5 (3ex^2 + 4d)}{\sqrt{1-c^2x^2}} dx + \frac{1}{6} dx^6 (a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{8} ex^8 (a + b \operatorname{sech}^{-1}(cx))$$

↓ 354

$$\frac{1}{48}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\int\frac{x^4(3ex^2+4d)}{\sqrt{1-c^2x^2}}dx^2+\frac{1}{6}dx^6(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{8}ex^8(a+b\operatorname{sech}^{-1}(cx))$$

↓ 86

$$\frac{1}{48}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\int\left(-\frac{3e(1-c^2x^2)^{5/2}}{c^6}+\frac{(4dc^2+9e)(1-c^2x^2)^{3/2}}{c^6}+\frac{(-8dc^2-9e)\sqrt{1-c^2x^2}}{c^6}+\frac{4dc^2+9e}{c^6\sqrt{1-c^2x^2}}\right)\frac{1}{6}dx^6(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{8}ex^8(a+b\operatorname{sech}^{-1}(cx))$$

↓ 2009

$$\frac{1}{48}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{6}dx^6(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{8}ex^8(a+b\operatorname{sech}^{-1}(cx))+\frac{2(1-c^2x^2)^{5/2}(4c^2d+9e)}{5c^8}+\frac{2(1-c^2x^2)^{3/2}(8c^2d+9e)}{3c^8}-\frac{2\sqrt{1-c^2x^2}(4c^2d+3e)}{c^8}+\frac{4dc^2+9e}{c^6\sqrt{1-c^2x^2}}\right)$$

input `Int[x^5*(d + e*x^2)*(a + b*ArcSech[c*x]),x]`

output `(b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-2*(4*c^2*d + 3*e)*Sqrt[1 - c^2*x^2])/c^8 + (2*(8*c^2*d + 9*e)*(1 - c^2*x^2)^(3/2))/(3*c^8) - (2*(4*c^2*d + 9*e)*(1 - c^2*x^2)^(5/2))/(5*c^8) + (6*e*(1 - c^2*x^2)^(7/2))/(7*c^8))/48 + (d*x^6*(a + b*ArcSech[c*x]))/6 + (e*x^8*(a + b*ArcSech[c*x]))/8`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`



```
rule 354 Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol]
:> Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]
```

```
rule 2009 Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6855 Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.)*((f_.)*(x_)^(m_.))*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol]
:> With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

### Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.60

method	result
parts	$a\left(\frac{1}{8}e x^8 + \frac{1}{6}d x^6\right) + \frac{b\left(\frac{c^6 \operatorname{arcsech}(cx)e x^8}{8} + \frac{\operatorname{arcsech}(cx)d c^6 x^6}{6} - \frac{\sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}} (45c^6 e x^6 + 84c^6 d x^4 + 54c^4 e x^4 + 112c^4 d x^2 + 72c^2 e x^2 + 224c^2 d + 144e)}{2520c}\right)}{c^6}$
derivativedivides	$\frac{a\left(\frac{1}{6}c^8 d x^6 + \frac{1}{8}e c^8 x^8\right)}{c^2} + \frac{b\left(\frac{\operatorname{arcsech}(cx)d c^8 x^6}{6} + \frac{\operatorname{arcsech}(cx)e c^8 x^8}{8} - \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (45c^6 e x^6 + 84c^6 d x^4 + 54c^4 e x^4 + 112c^4 d x^2 + 72c^2 e x^2 + 224c^2 d + 144e)}{2520}\right)}{c^2}$
default	$\frac{a\left(\frac{1}{6}c^8 d x^6 + \frac{1}{8}e c^8 x^8\right)}{c^2} + \frac{b\left(\frac{\operatorname{arcsech}(cx)d c^8 x^6}{6} + \frac{\operatorname{arcsech}(cx)e c^8 x^8}{8} - \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} (45c^6 e x^6 + 84c^6 d x^4 + 54c^4 e x^4 + 112c^4 d x^2 + 72c^2 e x^2 + 224c^2 d + 144e)}{2520}\right)}{c^2}$

```
input int(x^5*(e*x^2+d)*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)
```

```
output a*(1/8*e*x^8+1/6*d*x^6)+b/c^6*(1/8*c^6*arcsech(c*x)*e*x^8+1/6*arcsech(c*x)*d*c^6*x^6-1/2520/c*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(45*c^6*e*x^6+84*c^6*d*x^4+54*c^4*e*x^4+112*c^4*d*x^2+72*c^2*e*x^2+224*c^2*d+144*e))
```

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.72

$$\int x^5 (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{315 ac^7 ex^8 + 420 ac^7 dx^6 + 105 (3 bc^7 ex^8 + 4 bc^7 dx^6) \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2} + 1}}{cx}\right) - (45 bc^6 ex^7 + 6 (14 bc^6 d + 9 bc^6 e) x^5 + 8 (14 bc^4 d + 9 bc^4 e) x^3 + 16 (14 bc^2 d + 9 bc^2 e) x) \operatorname{sech}^{-1}(cx)}{2520 c^7}$$

input `integrate(x^5*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `1/2520*(315*a*c^7*e*x^8 + 420*a*c^7*d*x^6 + 105*(3*b*c^7*e*x^8 + 4*b*c^7*d*x^6)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (45*b*c^6*e*x^7 + 6*(14*b*c^6*d + 9*b*c^4*e)*x^5 + 8*(14*b*c^4*d + 9*b*c^2*e)*x^3 + 16*(14*b*c^2*d + 9*b*c^2*e)*x)*sech^-1(c*x)/c^7`

**Sympy [A] (verification not implemented)**

Time = 1.62 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.98

$$\int x^5 (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \begin{cases} \frac{adx^6}{6} + \frac{aex^8}{8} + \frac{bdx^6 \operatorname{asech}(cx)}{6} + \frac{bex^8 \operatorname{asech}(cx)}{8} - \frac{bdx^4 \sqrt{-c^2 x^2 + 1}}{30c^2} - \frac{bex^6 \sqrt{-c^2 x^2 + 1}}{56c^2} - \frac{2bdx^2 \sqrt{-c^2 x^2 + 1}}{45c^4} - \frac{3bex^4 \sqrt{-c^2 x^2 + 1}}{140c^4} \\ (a + \infty b) \left( \frac{dx^6}{6} + \frac{ex^8}{8} \right) \end{cases}$$

input `integrate(x**5*(e*x**2+d)*(a+b*asech(c*x)),x)`

output `Piecewise((a*d*x**6/6 + a*e*x**8/8 + b*d*x**6*asech(c*x)/6 + b*e*x**8*asech(c*x)/8 - b*d*x**4*sqrt(-c**2*x**2 + 1)/(30*c**2) - b*e*x**6*sqrt(-c**2*x**2 + 1)/(56*c**2) - 2*b*d*x**2*sqrt(-c**2*x**2 + 1)/(45*c**4) - 3*b*e*x**4*sqrt(-c**2*x**2 + 1)/(140*c**4) - 4*b*d*sqrt(-c**2*x**2 + 1)/(45*c**6) - b*e*x**2*sqrt(-c**2*x**2 + 1)/(35*c**6) - 2*b*e*sqrt(-c**2*x**2 + 1)/(35*c**8), Ne(c, 0)), ((a + oo*b)*(d*x**6/6 + e*x**8/8), True))`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.76

$$\int x^5 (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx = \frac{1}{8} aex^8 + \frac{1}{6} adx^6 + \frac{1}{90} \left( 15x^6 \operatorname{ar} \operatorname{sech}(cx) - \frac{3c^4x^5 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{5}{2}} - 10c^2x^3 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{3}{2}} + 15x \sqrt{\frac{1}{c^2x^2} - 1}}{c^5} \right) bd + \frac{1}{280} \left( 35x^8 \operatorname{ar} \operatorname{sech}(cx) + \frac{5c^6x^7 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{7}{2}} - 21c^4x^5 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{5}{2}} + 35c^2x^3 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{3}{2}} - 35x \sqrt{\frac{1}{c^2x^2} - 1}}{c^7} \right) bde$$

input `integrate(x^5*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `1/8*a*e*x^8 + 1/6*a*d*x^6 + 1/90*(15*x^6*arcsech(c*x) - (3*c^4*x^5*(1/(c^2*x^2) - 1)^(5/2) - 10*c^2*x^3*(1/(c^2*x^2) - 1)^(3/2) + 15*x*sqrt(1/(c^2*x^2) - 1))/c^5)*b*d + 1/280*(35*x^8*arcsech(c*x) + (5*c^6*x^7*(1/(c^2*x^2) - 1)^(7/2) - 21*c^4*x^5*(1/(c^2*x^2) - 1)^(5/2) + 35*c^2*x^3*(1/(c^2*x^2) - 1)^(3/2) - 35*x*sqrt(1/(c^2*x^2) - 1))/c^7)*b*e`

**Giac [F]**

$$\int x^5 (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)(b \operatorname{ar} \operatorname{sech}(cx) + a)x^5 dx$$

input `integrate(x^5*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((e*x^2 + d)*(b*arcsech(c*x) + a)*x^5, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^5 (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^5 (ex^2 + d) \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int(x^5*(d + e*x^2)*(a + b*acosh(1/(c*x))),x)`output `int(x^5*(d + e*x^2)*(a + b*acosh(1/(c*x))), x)`**Reduce [F]**

$$\int x^5 (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx = \left( \int a \operatorname{sech}(cx) x^7 dx \right) be + \left( \int a \operatorname{sech}(cx) x^5 dx \right) bd + \frac{ad x^6}{6} + \frac{ae x^8}{8}$$

input `int(x^5*(e*x^2+d)*(a+b*asech(c*x)),x)`output `(24*int(asech(c*x)*x**7,x)*b*e + 24*int(asech(c*x)*x**5,x)*b*d + 4*a*d*x**6 + 3*a*e*x**8)/24`

### 3.95 $\int x^3(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	764
Mathematica [A] (verified)	765
Rubi [A] (verified)	765
Maple [A] (verified)	767
Fricas [A] (verification not implemented)	768
Sympy [A] (verification not implemented)	768
Maxima [A] (verification not implemented)	769
Giac [F]	769
Mupad [F(-1)]	770
Reduce [F]	770

#### Optimal result

Integrand size = 19, antiderivative size = 180

$$\int x^3(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx = -\frac{b(3c^2d + 2e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{12c^6} + \frac{b(3c^2d + 4e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} (1-c^2x^2)^{3/2}}{36c^6} - \frac{be \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} (1-c^2x^2)^{5/2}}{30c^6} + \frac{1}{4} dx^4 (a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{6} ex^6 (a + b\operatorname{sech}^{-1}(cx))$$

output

```
-1/12*b*(3*c^2*d+2*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c
^6+1/36*b*(3*c^2*d+4*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(3/2)
/c^6-1/30*b*e*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(5/2)/c^6+1/4*d
*x^4*(a+b*arcsech(c*x))+1/6*e*x^6*(a+b*arcsech(c*x))
```

**Mathematica [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.59

$$\int x^3(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{1}{180} \left( 15ax^4(3d + 2ex^2) - \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(16e + c^2(30d + 8ex^2) + 3c^4(5dx^2 + 2ex^4))}{c^6} + 15bx^4(3d + 2ex^2)\operatorname{sech}^{-1}(cx) \right)$$

input `Integrate[x^3*(d + e*x^2)*(a + b*ArcSech[c*x]),x]`

output `(15*a*x^4*(3*d + 2*e*x^2) - (b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(16*e + c^2*(30*d + 8*e*x^2) + 3*c^4*(5*d*x^2 + 2*e*x^4)))/c^6 + 15*b*x^4*(3*d + 2*e*x^2)*ArcSech[c*x])/180`

**Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.78, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {6855, 27, 354, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx$$

$$\downarrow \text{6855}$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^3(2ex^2 + 3d)}{12\sqrt{1-c^2x^2}} dx + \frac{1}{4}dx^4(a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{6}ex^6(a + b\operatorname{sech}^{-1}(cx))$$

$$\downarrow \text{27}$$

$$\begin{aligned}
& \frac{1}{12} b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{x^3(2ex^2+3d)}{\sqrt{1-c^2x^2}} dx + \frac{1}{4} dx^4(a+b\operatorname{sech}^{-1}(cx)) + \frac{1}{6} ex^6(a+b\operatorname{sech}^{-1}(cx)) \\
& \quad \downarrow \text{354} \\
& \frac{1}{24} b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{x^2(2ex^2+3d)}{\sqrt{1-c^2x^2}} dx^2 + \frac{1}{4} dx^4(a+b\operatorname{sech}^{-1}(cx)) + \frac{1}{6} ex^6(a+b\operatorname{sech}^{-1}(cx)) \\
& \quad \downarrow \text{86} \\
& \frac{1}{24} b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \left( \frac{2e(1-c^2x^2)^{3/2}}{c^4} + \frac{(-3dc^2-4e)\sqrt{1-c^2x^2}}{c^4} + \frac{3dc^2+2e}{c^4\sqrt{1-c^2x^2}} \right) dx^2 + \\
& \quad \frac{1}{4} dx^4(a+b\operatorname{sech}^{-1}(cx)) + \frac{1}{6} ex^6(a+b\operatorname{sech}^{-1}(cx)) \\
& \quad \downarrow \text{2009} \\
& \frac{1}{24} b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \left( \frac{2(1-c^2x^2)^{3/2}(3c^2d+4e)}{3c^6} - \frac{2\sqrt{1-c^2x^2}(3c^2d+2e)}{c^6} - \frac{4e(1-c^2x^2)^{5/2}}{5c^6} \right) \\
& \quad \frac{1}{4} dx^4(a+b\operatorname{sech}^{-1}(cx)) + \frac{1}{6} ex^6(a+b\operatorname{sech}^{-1}(cx)) +
\end{aligned}$$

input `Int[x^3*(d + e*x^2)*(a + b*ArcSech[c*x]),x]`

output `(b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-2*(3*c^2*d + 2*e)*Sqrt[1 - c^2*x^2])/c^6 + (2*(3*c^2*d + 4*e)*(1 - c^2*x^2)^(3/2))/(3*c^6) - (4*e*(1 - c^2*x^2)^(5/2))/(5*c^6))/24 + (d*x^4*(a + b*ArcSech[c*x])/4 + (e*x^6*(a + b*ArcSech[c*x]))/6`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

```
rule 354 Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol]
:> Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]
```

```
rule 2009 Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6855 Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.)*((f_.)*(x_)^(m_.))*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol]
:> With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

### Maple [A] (verified)

Time = 0.53 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.67

method	result
parts	$a\left(\frac{1}{6}ex^6 + \frac{1}{4}dx^4\right) + \frac{b\left(\frac{c^4 \operatorname{arcsech}(cx)e^6}{6} + \frac{\operatorname{arcsech}(cx)d^4x^4}{4} - \frac{\sqrt{-\frac{cx-1}{cx}}x\sqrt{\frac{cx+1}{cx}}(6c^4ex^4 + 15c^4dx^2 + 8e^2c^2x^2 + 30c^2d)}{180c}\right)}{c^4}$
derivativedivides	$-\frac{a\left(\frac{c^2d(e^2x^2+c^2d)^2}{2} - \frac{(e^2x^2+c^2d)^3}{3}\right)}{2c^2e^2} + \frac{b\left(-\frac{\operatorname{arcsech}(cx)c^6d^3}{12e^2} + \frac{\operatorname{arcsech}(cx)c^6dx^4}{4} + \frac{e \operatorname{arcsech}(cx)c^6x^6}{6} - \frac{\sqrt{-\frac{cx-1}{cx}}cx\sqrt{\frac{cx+1}{cx}}}{c^2}\right)}{c^4}$
default	$-\frac{a\left(\frac{c^2d(e^2x^2+c^2d)^2}{2} - \frac{(e^2x^2+c^2d)^3}{3}\right)}{2c^2e^2} + \frac{b\left(-\frac{\operatorname{arcsech}(cx)c^6d^3}{12e^2} + \frac{\operatorname{arcsech}(cx)c^6dx^4}{4} + \frac{e \operatorname{arcsech}(cx)c^6x^6}{6} - \frac{\sqrt{-\frac{cx-1}{cx}}cx\sqrt{\frac{cx+1}{cx}}}{c^2}\right)}{c^4}$

```
input int(x^3*(e*x^2+d)*(a+b*arcsech(c*x)), x, method=_RETURNVERBOSE)
```

```
output a*(1/6*e*x^6+1/4*d*x^4)+b/c^4*(1/6*c^4*arcsech(c*x)*e*x^6+1/4*arcsech(c*x)*d*c^4*x^4-1/180/c*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(6*c^4*e*x^4+15*c^4*d*x^2+8*c^2*e*x^2+30*c^2*d+16*e))
```



**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.82

$$\int x^3(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{30ac^5ex^6 + 45ac^5dx^4 + 15(2bc^5ex^6 + 3bc^5dx^4) \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}+1}}{cx}\right) - (6bc^4ex^5 + (15bc^4d + 8bc^2e)x^3)}{180c^5}$$

input `integrate(x^3*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `1/180*(30*a*c^5*e*x^6 + 45*a*c^5*d*x^4 + 15*(2*b*c^5*e*x^6 + 3*b*c^5*d*x^4)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (6*b*c^4*e*x^5 + (15*b*c^4*d + 8*b*c^2*e)*x^3 + 2*(15*b*c^2*d + 8*b*e)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/c^5`

**Sympy [A] (verification not implemented)**

Time = 0.87 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.98

$$\int x^3(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \begin{cases} \frac{adx^4}{4} + \frac{aex^6}{6} + \frac{bdx^4 \operatorname{asech}(cx)}{4} + \frac{bex^6 \operatorname{asech}(cx)}{6} - \frac{bdx^2\sqrt{-c^2x^2+1}}{12c^2} - \frac{bex^4\sqrt{-c^2x^2+1}}{30c^2} - \frac{bd\sqrt{-c^2x^2+1}}{6c^4} - \frac{2bex^2\sqrt{-c^2x^2+1}}{45c^4} \\ (a + \infty b) \left( \frac{dx^4}{4} + \frac{ex^6}{6} \right) \end{cases}$$

input `integrate(x**3*(e*x**2+d)*(a+b*asech(c*x)),x)`

output `Piecewise((a*d*x**4/4 + a*e*x**6/6 + b*d*x**4*asech(c*x)/4 + b*e*x**6*asech(c*x)/6 - b*d*x**2*sqrt(-c**2*x**2 + 1)/(12*c**2) - b*e*x**4*sqrt(-c**2*x**2 + 1)/(30*c**2) - b*d*sqrt(-c**2*x**2 + 1)/(6*c**4) - 2*b*e*x**2*sqrt(-c**2*x**2 + 1)/(45*c**4) - 4*b*e*sqrt(-c**2*x**2 + 1)/(45*c**6), Ne(c, 0)), ((a + oo*b)*(d*x**4/4 + e*x**6/6), True))`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.77

$$\int x^3(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{1}{6}aex^6 + \frac{1}{4}adx^4 + \frac{1}{12} \left( 3x^4 \operatorname{ar} \operatorname{sech}(cx) + \frac{c^2x^3 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{3}{2}} - 3x\sqrt{\frac{1}{c^2x^2} - 1}}{c^3} \right) bd$$

$$+ \frac{1}{90} \left( 15x^6 \operatorname{ar} \operatorname{sech}(cx) - \frac{3c^4x^5 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{5}{2}} - 10c^2x^3 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{3}{2}} + 15x\sqrt{\frac{1}{c^2x^2} - 1}}{c^5} \right) be$$

input `integrate(x^3*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `1/6*a*e*x^6 + 1/4*a*d*x^4 + 1/12*(3*x^4*arcsech(c*x) + (c^2*x^3*(1/(c^2*x^2) - 1)^(3/2) - 3*x*sqrt(1/(c^2*x^2) - 1))/c^3)*b*d + 1/90*(15*x^6*arcsech(c*x) - (3*c^4*x^5*(1/(c^2*x^2) - 1)^(5/2) - 10*c^2*x^3*(1/(c^2*x^2) - 1)^(3/2) + 15*x*sqrt(1/(c^2*x^2) - 1))/c^5)*b*e`

**Giac [F]**

$$\int x^3(d + ex^2)(a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)(b \operatorname{ar} \operatorname{sech}(cx) + a)x^3 dx$$

input `integrate(x^3*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((e*x^2 + d)*(b*arcsech(c*x) + a)*x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^3 (ex^2 + d) \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int(x^3*(d + e*x^2)*(a + b*acosh(1/(c*x))),x)`output `int(x^3*(d + e*x^2)*(a + b*acosh(1/(c*x))), x)`**Reduce [F]**

$$\int x^3 (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx = \left( \int a \operatorname{sech}(cx) x^5 dx \right) be + \left( \int a \operatorname{sech}(cx) x^3 dx \right) bd + \frac{ad x^4}{4} + \frac{ae x^6}{6}$$

input `int(x^3*(e*x^2+d)*(a+b*asech(c*x)),x)`output `(12*int(asech(c*x)*x**5,x)*b*e + 12*int(asech(c*x)*x**3,x)*b*d + 3*a*d*x**4 + 2*a*e*x**6)/12`

### 3.96 $\int x(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	771
Mathematica [A] (verified)	772
Rubi [A] (verified)	772
Maple [A] (verified)	774
Fricas [A] (verification not implemented)	775
Sympy [A] (verification not implemented)	775
Maxima [A] (verification not implemented)	776
Giac [F]	776
Mupad [F(-1)]	777
Reduce [F]	777

#### Optimal result

Integrand size = 17, antiderivative size = 164

$$\int x(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx = -\frac{b(2c^2d + e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{4c^4} + \frac{be \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} (1-c^2x^2)^{3/2}}{12c^4} + \frac{(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx))}{4e} - \frac{bd^2 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \operatorname{arctanh}(\sqrt{1-c^2x^2})}{4e}$$

output

```
-1/4*b*(2*c^2*d+e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^4+
1/12*b*e*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(3/2)/c^4+1/4*(e*x^2
+d)^2*(a+b*arcsech(c*x))/e-1/4*b*d^2*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arcta
nh((-c^2*x^2+1)^(1/2))/e
```

**Mathematica [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.52

$$\int x(d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx = \frac{1}{12} \left( 3ax^2(2d + ex^2) - \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(2e + c^2(6d + ex^2))}{c^4} + 3bx^2(2d + ex^2) \operatorname{sech}^{-1}(cx) \right)$$

input

```
Integrate[x*(d + e*x^2)*(a + b*ArcSech[c*x]),x]
```

output

```
(3*a*x^2*(2*d + e*x^2) - (b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(2*e + c^2*(6*d + e*x^2)))/c^4 + 3*b*x^2*(2*d + e*x^2)*ArcSech[c*x])/12
```

**Rubi [A] (verified)**

Time = 0.48 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.77, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$ , Rules used = {6853, 2036, 354, 99, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x(d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx \\ & \quad \downarrow \text{6853} \\ & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(ex^2+d)^2}{x\sqrt{1-cx}\sqrt{cx+1}} dx}{4e} + \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{4e} \\ & \quad \downarrow \text{2036} \\ & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(ex^2+d)^2}{x\sqrt{1-c^2x^2}} dx}{4e} + \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{4e} \end{aligned}$$

$$\begin{aligned}
& \downarrow 354 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(ex^2+d)^2}{x^2\sqrt{1-c^2x^2}} dx^2}{8e} + \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{4e} \\
& \downarrow 99 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \left( \frac{d^2}{x^2\sqrt{1-c^2x^2}} - \frac{e^2\sqrt{1-c^2x^2}}{c^2} + \frac{e(2dc^2+e)}{c^2\sqrt{1-c^2x^2}} \right) dx^2}{8e} + \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{4e} \\
& \downarrow 2009 \\
& \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{4e} + \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -2d^2\operatorname{arctanh}\left(\sqrt{1-c^2x^2}\right) - \frac{2e\sqrt{1-c^2x^2}(2c^2d+e)}{c^4} + \frac{2e^2(1-c^2x^2)^{3/2}}{3c^4} \right)}{8e}
\end{aligned}$$

input `Int[x*(d + e*x^2)*(a + b*ArcSech[c*x]),x]`

output `((d + e*x^2)^2*(a + b*ArcSech[c*x]))/(4*e) + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-2*e*(2*c^2*d + e)*Sqrt[1 - c^2*x^2])/c^4 + (2*e^2*(1 - c^2*x^2)^(3/2))/(3*c^4) - 2*d^2*ArcTanh[Sqrt[1 - c^2*x^2]]))/(8*e)`

### Defintions of rubi rules used

rule 99 `Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] | (GtQ[m, 0] && GeQ[n, -1]))]`

rule 354 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2036

```
Int[(u_)*((c_) + (d_)*(x_)^(n_))^(q_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_), x_Symbol] := Int[u*(a1*a2 + b1*b2 *x^n)^p*(c + d*x^n)^q, x] /; FreeQ[{a1, b1, a2, b2, c, d, n, p, q}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))
```

rule 6853

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*(x_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcSech[c*x])/(2*e*(p + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(2*e*(p + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x^2)^(p + 1)/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[p, -1]
```

### Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.18

method	result
parts	$\frac{a(x^2e+d)^2}{4e} + \frac{b \left( \frac{c^2e \operatorname{arcsech}(cx)x^4}{4} + \frac{\operatorname{arcsech}(cx)c^2x^2d}{2} + \frac{c^2 \operatorname{arcsech}(cx)d^2}{4e} - \frac{\sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}} \left( 3c^4d^2 \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right) \right)}{12} \right)}{c^2}$
derivativedivides	$\frac{a(e c^2 x^2 + c^2 d)^2}{4c^2e} + \frac{b \left( \frac{\operatorname{arcsech}(cx)c^4d^2}{4e} + \frac{\operatorname{arcsech}(cx)c^4dx^2}{2} + \frac{e \operatorname{arcsech}(cx)c^4x^4}{4} - \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} \left( 3c^4d^2 \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right) \right)}{12} \right)}{c^2}$
default	$\frac{a(e c^2 x^2 + c^2 d)^2}{4c^2e} + \frac{b \left( \frac{\operatorname{arcsech}(cx)c^4d^2}{4e} + \frac{\operatorname{arcsech}(cx)c^4dx^2}{2} + \frac{e \operatorname{arcsech}(cx)c^4x^4}{4} - \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}} \left( 3c^4d^2 \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right) \right)}{12} \right)}{c^2}$

input

```
int(x*(e*x^2+d)*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)
```

output

```
1/4*a*(e*x^2+d)^2/e+b/c^2*(1/4*c^2*e*arcsech(c*x)*x^4+1/2*arcsech(c*x)*c^2*x^2*d+1/4*c^2/e*arcsech(c*x)*d^2-1/12/c/e*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(3*c^4*d^2*arctanh(1/(-c^2*x^2+1)^(1/2))+6*c^2*d*e*(-c^2*x^2+1)^(1/2)+e^2*(-c^2*x^2+1)^(1/2)*c^2*x^2+2*e^2*(-c^2*x^2+1)^(1/2))/(-c^2*x^2+1)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.76

$$\int x(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{3ac^3ex^4 + 6ac^3dx^2 + 3(bc^3ex^4 + 2bc^3dx^2) \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}+1}{cx}\right) - (bc^2ex^3 + 2(3bc^2d + be)x)\sqrt{-\frac{c^2x^2-1}{c^2x^2}}}{12c^3}$$

input `integrate(x*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="fricas")`output `1/12*(3*a*c^3*e*x^4 + 6*a*c^3*d*x^2 + 3*(b*c^3*e*x^4 + 2*b*c^3*d*x^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (b*c^2*e*x^3 + 2*(3*b*c^2*d + b*e)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/c^3`**Sympy [A] (verification not implemented)**

Time = 0.48 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.77

$$\int x(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \begin{cases} \frac{adx^2}{2} + \frac{aex^4}{4} + \frac{bdx^2 \operatorname{asech}(cx)}{2} + \frac{be x^4 \operatorname{asech}(cx)}{4} - \frac{bd\sqrt{-c^2x^2+1}}{2c^2} - \frac{be x^2 \sqrt{-c^2x^2+1}}{12c^2} - \frac{be\sqrt{-c^2x^2+1}}{6c^4} & \text{for } c \neq 0 \\ (a + \infty b) \left( \frac{dx^2}{2} + \frac{ex^4}{4} \right) & \text{otherwise} \end{cases}$$

input `integrate(x*(e*x**2+d)*(a+b*asech(c*x)),x)`output `Piecewise((a*d*x**2/2 + a*e*x**4/4 + b*d*x**2*asech(c*x)/2 + b*e*x**4*asech(c*x)/4 - b*d*sqrt(-c**2*x**2 + 1)/(2*c**2) - b*e*x**2*sqrt(-c**2*x**2 + 1)/(12*c**2) - b*e*sqrt(-c**2*x**2 + 1)/(6*c**4), Ne(c, 0)), ((a + oo*b)*(d*x**2/2 + e*x**4/4), True))`



**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.59

$$\int x(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{1}{4} aex^4 + \frac{1}{2} adx^2 + \frac{1}{2} \left( x^2 \operatorname{ar} \operatorname{sech}(cx) - \frac{x \sqrt{\frac{1}{c^2 x^2} - 1}}{c} \right) bd$$

$$+ \frac{1}{12} \left( 3x^4 \operatorname{ar} \operatorname{sech}(cx) + \frac{c^2 x^3 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{3}{2}} - 3x \sqrt{\frac{1}{c^2 x^2} - 1}}{c^3} \right) be$$

input `integrate(x*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `1/4*a*e*x^4 + 1/2*a*d*x^2 + 1/2*(x^2*arcsech(c*x) - x*sqrt(1/(c^2*x^2) - 1)/c)*b*d + 1/12*(3*x^4*arcsech(c*x) + (c^2*x^3*(1/(c^2*x^2) - 1)^(3/2) - 3*x*sqrt(1/(c^2*x^2) - 1))/c^3)*b*e`

**Giac [F]**

$$\int x(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)(b \operatorname{ar} \operatorname{sech}(cx) + a)x dx$$

input `integrate(x*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((e*x^2 + d)*(b*arcsech(c*x) + a)*x, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx = \int x (ex^2 + d) \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int(x*(d + e*x^2)*(a + b*acosh(1/(c*x))),x)`output `int(x*(d + e*x^2)*(a + b*acosh(1/(c*x))), x)`**Reduce [F]**

$$\int x(d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx = \left( \int a\operatorname{sech}(cx) x^3 dx \right) be + \left( \int a\operatorname{sech}(cx) x dx \right) bd + \frac{ad x^2}{2} + \frac{ae x^4}{4}$$

input `int(x*(e*x^2+d)*(a+b*asech(c*x)),x)`output `(4*int(asech(c*x)*x**3,x)*b*e + 4*int(asech(c*x)*x,x)*b*d + 2*a*d*x**2 + a*e*x**4)/4`

$$3.97 \quad \int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x} dx$$

Optimal result	778
Mathematica [A] (verified)	779
Rubi [A] (verified)	779
Maple [A] (verified)	782
Fricas [F]	782
Sympy [F]	783
Maxima [F]	783
Giac [F]	783
Mupad [F(-1)]	784
Reduce [F]	784

### Optimal result

Integrand size = 19, antiderivative size = 296

$$\begin{aligned} \int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x} dx = & -\frac{be\sqrt{-1+\frac{1}{cx}}\sqrt{1+\frac{1}{cx}}x}{2c} \\ & + \frac{ibd\sqrt{1-\frac{1}{c^2x^2}}\operatorname{csc}^{-1}(cx)^2}{2\sqrt{-1+\frac{1}{cx}}\sqrt{1+\frac{1}{cx}}} + \frac{1}{2}ex^2(a+b\operatorname{sech}^{-1}(cx)) \\ & - \frac{bd\sqrt{1-\frac{1}{c^2x^2}}\operatorname{csc}^{-1}(cx)\log\left(1-e^{2i\operatorname{csc}^{-1}(cx)}\right)}{\sqrt{-1+\frac{1}{cx}}\sqrt{1+\frac{1}{cx}}} \\ & + \frac{bd\sqrt{1-\frac{1}{c^2x^2}}\operatorname{csc}^{-1}(cx)\log\left(\frac{1}{x}\right)}{\sqrt{-1+\frac{1}{cx}}\sqrt{1+\frac{1}{cx}}} \\ & - d(a+b\operatorname{sech}^{-1}(cx))\log\left(\frac{1}{x}\right) \\ & + \frac{ibd\sqrt{1-\frac{1}{c^2x^2}}\operatorname{PolyLog}\left(2,e^{2i\operatorname{csc}^{-1}(cx)}\right)}{2\sqrt{-1+\frac{1}{cx}}\sqrt{1+\frac{1}{cx}}} \end{aligned}$$

output

```
-1/2*b*e*(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)*x/c+1/2*I*b*d*(1-1/c^2/x^2)^(1/2)
)*arccsc(c*x)^2/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)+1/2*e*x^2*(a+b*arcsech(c*
x))-b*d*(1-1/c^2/x^2)^(1/2)*arccsc(c*x)*ln(1-(1/c/x+(1-1/c^2/x^2)^(1/2))^2
)/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)+b*d*(1-1/c^2/x^2)^(1/2)*arccsc(c*x)*ln(
1/x)/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)-d*(a+b*arcsech(c*x))*ln(1/x)+1/2*I*b
*d*(1-1/c^2/x^2)^(1/2)*polylog(2,(1/c/x+(1-1/c^2/x^2)^(1/2))^2)/(-1+1/c/x)
^(1/2)/(1+1/c/x)^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.36

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x} dx$$

$$= \frac{1}{2} a e x^2 + b e \left( -\frac{1}{2c^2} - \frac{x}{2c} \right) \sqrt{\frac{1 - cx}{1 + cx}} + \frac{1}{2} b e x^2 \operatorname{sech}^{-1}(cx) + a d \log(x)$$

$$+ \frac{1}{2} b d \left( -\operatorname{sech}^{-1}(cx) \left( \operatorname{sech}^{-1}(cx) + 2 \log \left( 1 + e^{-2 \operatorname{sech}^{-1}(cx)} \right) \right) \right. \\ \left. + \operatorname{PolyLog} \left( 2, -e^{-2 \operatorname{sech}^{-1}(cx)} \right) \right)$$

input

```
Integrate[((d + e*x^2)*(a + b*ArcSech[c*x]))/x,x]
```

output

```
(a*e*x^2)/2 + b*e*(-1/2*1/c^2 - x/(2*c))*Sqrt[(1 - c*x)/(1 + c*x)] + (b*e*
x^2*ArcSech[c*x])/2 + a*d*Log[x] + (b*d*(-(ArcSech[c*x]*(ArcSech[c*x] + 2*
Log[1 + E^(-2*ArcSech[c*x]))])) + PolyLog[2, -E^(-2*ArcSech[c*x]))]/2
```

**Rubi [A] (verified)**Time = 1.28 (sec) , antiderivative size = 323, normalized size of antiderivative = 1.09, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {6857, 6373, 27, 7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.



output

```
(e*x^2*(a + b*ArcCosh[1/(c*x)]))/2 - d*(a + b*ArcCosh[1/(c*x)]*Log[x^(-1)] - (b*(e*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]*x - (I*c*d*Sqrt[1 - 1/(c^2*x^2)]*ArcSin[1/(c*x)]^2)/(Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) + (2*c*d*Sqrt[1 - 1/(c^2*x^2)]*ArcSin[1/(c*x)]*Log[1 - E^((2*I)*ArcSin[1/(c*x)])])/(Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) - (2*c*d*Sqrt[1 - 1/(c^2*x^2)]*ArcSin[1/(c*x)]*Log[x^(-1)])/(Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) - (I*c*d*Sqrt[1 - 1/(c^2*x^2)]*PolyLog[2, E^((2*I)*ArcSin[1/(c*x)])])/(Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)])))/(2*c)
```

**Defintions of rubi rules used**

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6373

```
Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))
```

rule 6857

```
Int[((a_) + ArcSech[(c_)*(x_)]*(b_))^(n_)*(x_)^m)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := -Subst[Int[(e + d*x^2)^p*(a + b*ArcCosh[x/c])^n/x^(m + 2*(p + 1)), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0] && IntegersQ[m, p]
```

rule 7293

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]
```

**Maple [A] (verified)**

Time = 0.81 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.54

method	result
parts	$\frac{ax^2e}{2} + ad \ln(x) + b \left( \frac{\operatorname{arcsech}(cx)^2 d}{2} + \frac{e \left( c^2 x^2 \operatorname{arcsech}(cx) - \sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x + 1 \right)}{2c^2} - d \operatorname{arcsech}(cx) \right)$
derivativedivides	$\frac{ax^2e}{2} + ad \ln(cx) + \frac{b \left( \frac{c^2 d \operatorname{arcsech}(cx)^2}{2} + \frac{e \left( c^2 x^2 \operatorname{arcsech}(cx) - \sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x + 1 \right)}{2} - \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \right) \sqrt{\frac{cx+1}{cx}} \right) \right)}{c^2}$
default	$\frac{ax^2e}{2} + ad \ln(cx) + \frac{b \left( \frac{c^2 d \operatorname{arcsech}(cx)^2}{2} + \frac{e \left( c^2 x^2 \operatorname{arcsech}(cx) - \sqrt{-\frac{cx-1}{cx}} c \sqrt{\frac{cx+1}{cx}} x + 1 \right)}{2} - \ln \left( 1 + \left( \frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \right) \sqrt{\frac{cx+1}{cx}} \right) \right)}{c^2}$

input `int((e*x^2+d)*(a+b*arcsech(c*x))/x,x,method=_RETURNVERBOSE)`

output `1/2*a*x^2*e+a*d*ln(x)+b*(1/2*arcsech(c*x)^2*d+1/2*e*(c^2*x^2*arcsech(c*x)-(-c*x-1)/c/x)^(1/2)*c*((c*x+1)/c/x)^(1/2)*x+1)/c^2-d*arcsech(c*x)*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)-1/2*d*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2))`

**Fricas [F]**

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(ex^2 + d)(b \operatorname{ar} \operatorname{sech}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x))/x, x, algorithm="fricas")`

output `integral((a*e*x^2 + a*d + (b*e*x^2 + b*d)*arcsech(c*x))/x, x)`

**Sympy [F]**

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(a + b \operatorname{arsech}(cx))(d + ex^2)}{x} dx$$

input `integrate((e*x**2+d)*(a+b*asech(c*x))/x,x)`

output `Integral((a + b*asech(c*x))*(d + e*x**2)/x, x)`

**Maxima [F]**

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(ex^2 + d)(b \operatorname{arsech}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x))/x,x, algorithm="maxima")`

output `1/2*a*e*x^2 + a*d*log(x) + integrate(b*e*x*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x)) + b*d*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x)))/x, x)`

**Giac [F]**

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(ex^2 + d)(b \operatorname{arsech}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x))/x,x, algorithm="giac")`

output `integrate((e*x^2 + d)*(b*arcsech(c*x) + a)/x, x)`



**Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(ex^2 + d)(a + b\operatorname{acosh}(\frac{1}{cx}))}{x} dx$$

input `int(((d + e*x^2)*(a + b*acosh(1/(c*x))))/x,x)`

output `int(((d + e*x^2)*(a + b*acosh(1/(c*x))))/x, x)`

**Reduce [F]**

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x} dx = \left( \int \frac{a\operatorname{sech}(cx)}{x} dx \right) bd + \left( \int a\operatorname{sech}(cx) x dx \right) be + \log(x) ad + \frac{ae x^2}{2}$$

input `int((e*x^2+d)*(a+b*asech(c*x))/x,x)`

output `(2*int(asech(c*x)/x,x)*b*d + 2*int(asech(c*x)*x,x)*b*e + 2*log(x)*a*d + a*e*x**2)/2`

$$3.98 \quad \int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^3} dx$$

Optimal result	785
Mathematica [A] (verified)	786
Rubi [A] (verified)	787
Maple [A] (verified)	789
Fricas [F]	789
Sympy [F]	790
Maxima [F]	790
Giac [F]	790
Mupad [F(-1)]	791
Reduce [F]	791

### Optimal result

Integrand size = 19, antiderivative size = 309

$$\begin{aligned} \int \frac{(d+ex^2)(a+b\operatorname{sech}^{-1}(cx))}{x^3} dx = & \frac{bcd\sqrt{-1+\frac{1}{cx}}\sqrt{1+\frac{1}{cx}}}{4x} + \frac{ibe\sqrt{1-\frac{1}{c^2x^2}}\operatorname{csc}^{-1}(cx)^2}{2\sqrt{-1+\frac{1}{cx}}\sqrt{1+\frac{1}{cx}}} \\ & + \frac{1}{4}bc^2d\operatorname{sech}^{-1}(cx) - \frac{d(a+b\operatorname{sech}^{-1}(cx))}{2x^2} \\ & - \frac{be\sqrt{1-\frac{1}{c^2x^2}}\operatorname{csc}^{-1}(cx)\log\left(1-e^{2i\operatorname{csc}^{-1}(cx)}\right)}{\sqrt{-1+\frac{1}{cx}}\sqrt{1+\frac{1}{cx}}} \\ & + \frac{be\sqrt{1-\frac{1}{c^2x^2}}\operatorname{csc}^{-1}(cx)\log\left(\frac{1}{x}\right)}{\sqrt{-1+\frac{1}{cx}}\sqrt{1+\frac{1}{cx}}} \\ & - e(a+b\operatorname{sech}^{-1}(cx))\log\left(\frac{1}{x}\right) \\ & + \frac{ibe\sqrt{1-\frac{1}{c^2x^2}}\operatorname{PolyLog}\left(2,e^{2i\operatorname{csc}^{-1}(cx)}\right)}{2\sqrt{-1+\frac{1}{cx}}\sqrt{1+\frac{1}{cx}}} \end{aligned}$$

output

```
1/4*b*c*d*(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)/x+1/2*I*b*e*(1-1/c^2/x^2)^(1/2)
*arccsc(c*x)^2/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)+1/4*b*c^2*d*arcsech(c*x)-1
/2*d*(a+b*arcsech(c*x))/x^2-b*e*(1-1/c^2/x^2)^(1/2)*arccsc(c*x)*ln(1-(I/c/
x+(1-1/c^2/x^2)^(1/2))^2)/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)+b*e*(1-1/c^2/x^
2)^(1/2)*arccsc(c*x)*ln(1/x)/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)-e*(a+b*arcse
ch(c*x))*ln(1/x)+1/2*I*b*e*(1-1/c^2/x^2)^(1/2)*polylog(2,(I/c/x+(1-1/c^2/x
^2)^(1/2))^2)/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.55

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^3} dx = -\frac{ad}{2x^2} + bd\left(\frac{1}{4x^2} + \frac{c}{4x}\right)\sqrt{\frac{1-cx}{1+cx}}$$

$$-\frac{bd\operatorname{sech}^{-1}(cx)}{2x^2} - \frac{1}{4}bc^2d\log(x) + ae\log(x)$$

$$+ \frac{1}{4}bc^2d\log\left(1 + \sqrt{\frac{1-cx}{1+cx}} + cx\sqrt{\frac{1-cx}{1+cx}}\right)$$

$$+ \frac{1}{2}be\left(-\operatorname{sech}^{-1}(cx)\left(\operatorname{sech}^{-1}(cx)\right.\right.$$

$$\left.\left.+ 2\log\left(1 + e^{-2\operatorname{sech}^{-1}(cx)}\right)\right)\right)$$

$$+ \operatorname{PolyLog}\left(2, -e^{-2\operatorname{sech}^{-1}(cx)}\right)$$

input

```
Integrate[((d + e*x^2)*(a + b*ArcSech[c*x]))/x^3,x]
```

output

```
-1/2*(a*d)/x^2 + b*d*(1/(4*x^2) + c/(4*x))*Sqrt[(1 - c*x)/(1 + c*x)] - (b*
d*ArcSech[c*x])/(2*x^2) - (b*c^2*d*Log[x])/4 + a*e*Log[x] + (b*c^2*d*Log[1
+ Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x]])/4 + (b*e*(-
(ArcSech[c*x]*(ArcSech[c*x] + 2*Log[1 + E^(-2*ArcSech[c*x])])) + PolyLog[2
, -E^(-2*ArcSech[c*x])]))/2
```



input `Int[((d + e*x^2)*(a + b*ArcSech[c*x]))/x^3,x]`

output `-1/2*(d*(a + b*ArcCosh[1/(c*x)]))/x^2 - e*(a + b*ArcCosh[1/(c*x)])*Log[x^(-1)] + (b*((c^2*d*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)])/(2*x) + (c^3*d*ArcCosh[1/(c*x)])/2 + (I*c*e*Sqrt[1 - 1/(c^2*x^2)]*ArcSin[1/(c*x)]^2)/(Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) - (2*c*e*Sqrt[1 - 1/(c^2*x^2)]*ArcSin[1/(c*x)]*Log[1 - E^((2*I)*ArcSin[1/(c*x)])])/(Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) + (2*c*e*Sqrt[1 - 1/(c^2*x^2)]*ArcSin[1/(c*x)]*Log[x^(-1)])/(Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) + (I*c*e*Sqrt[1 - 1/(c^2*x^2)]*PolyLog[2, E^((2*I)*ArcSin[1/(c*x)])])/(Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)])))/(2*c)`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6873 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[p[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))`

rule 6857 `Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^((n_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := -Subst[Int[(e + d*x^2)^p*(a + b*ArcCosh[x/c])^n/x^(m + 2*(p + 1)), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0] && IntegersQ[m, p]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

### Maple [A] (verified)

Time = 1.09 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.54

method	result
parts	$-\frac{ad}{2x^2} + ae \ln(x) + \frac{b \operatorname{arcsech}(cx)^2 e}{2} + \frac{bc^2 d \operatorname{arcsech}(cx)}{4} + \frac{bcd \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}}{4cx} - \frac{bd \operatorname{arcsech}(cx)}{2x^2} - be$
derivativeldivides	$c^2 \left( \frac{ae \ln(cx)}{c^2} - \frac{ad}{2c^2 x^2} + \frac{be \operatorname{arcsech}(cx)^2}{2c^2} + \frac{bd \operatorname{arcsech}(cx)}{4} + \frac{bd \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}}{4cx} - \frac{b \operatorname{arcsech}(cx)d}{2c^2 x^2} - \frac{be}{c^2} \right)$
default	$c^2 \left( \frac{ae \ln(cx)}{c^2} - \frac{ad}{2c^2 x^2} + \frac{be \operatorname{arcsech}(cx)^2}{2c^2} + \frac{bd \operatorname{arcsech}(cx)}{4} + \frac{bd \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}}{4cx} - \frac{b \operatorname{arcsech}(cx)d}{2c^2 x^2} - \frac{be}{c^2} \right)$

input `int((e*x^2+d)*(a+b*arcsech(c*x))/x^3,x,method=_RETURNVERBOSE)`

output `-1/2*a*d/x^2+a*e*ln(x)+1/2*b*arcsech(c*x)^2*e+1/4*b*c^2*d*arcsech(c*x)+1/4*b*c*d/x*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)-1/2*b*d/x^2*arcsech(c*x)-b*e*arcsech(c*x)*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)-1/2*b*e*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)`

### Fricas [F]

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(ex^2 + d)(b \operatorname{ar} \operatorname{sech}(cx) + a)}{x^3} dx$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x))/x^3,x, algorithm="fricas")`

output `integral((a*e*x^2 + a*d + (b*e*x^2 + b*d)*arcsech(c*x))/x^3, x)`

**Sympy [F]**

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(a + b \operatorname{arsech}(cx))(d + ex^2)}{x^3} dx$$

input `integrate((e*x**2+d)*(a+b*asech(c*x))/x**3,x)`

output `Integral((a + b*asech(c*x))*(d + e*x**2)/x**3, x)`

**Maxima [F]**

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(ex^2 + d)(b \operatorname{arsech}(cx) + a)}{x^3} dx$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x))/x^3,x, algorithm="maxima")`

output `-1/8*b*d*((2*c^4*x*sqrt(1/(c^2*x^2) - 1)/(c^2*x^2*(1/(c^2*x^2) - 1) - 1) - c^3*log(c*x*sqrt(1/(c^2*x^2) - 1) + 1) + c^3*log(c*x*sqrt(1/(c^2*x^2) - 1) - 1))/c + 4*arcsech(c*x)/x^2) + b*e*integrate(log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/x, x) + a*e*log(x) - 1/2*a*d/x^2`

**Giac [F]**

$$\int \frac{(d + ex^2)(a + b \operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(ex^2 + d)(b \operatorname{arsech}(cx) + a)}{x^3} dx$$

input `integrate((e*x^2+d)*(a+b*arcsech(c*x))/x^3,x, algorithm="giac")`

output `integrate((e*x^2 + d)*(b*arcsech(c*x) + a)/x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(ex^2 + d)(a + b\operatorname{acosh}(\frac{1}{cx}))}{x^3} dx$$

input `int(((d + e*x^2)*(a + b*acosh(1/(c*x))))/x^3,x)`output `int(((d + e*x^2)*(a + b*acosh(1/(c*x))))/x^3, x)`**Reduce [F]**

$$\int \frac{(d + ex^2)(a + b\operatorname{sech}^{-1}(cx))}{x^3} dx$$

$$= \frac{2\left(\int \frac{\operatorname{asech}(cx)}{x^3} dx\right) b d x^2 + 2\left(\int \frac{\operatorname{asech}(cx)}{x} dx\right) b e x^2 + 2 \log(x) a e x^2 - a d}{2x^2}$$

input `int((e*x^2+d)*(a+b*asech(c*x))/x^3,x)`output `(2*int(asech(c*x)/x**3,x)*b*d*x**2 + 2*int(asech(c*x)/x,x)*b*e*x**2 + 2*log(x)*a*e*x**2 - a*d)/(2*x**2)`



### 3.99 $\int x^2(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	792
Mathematica [C] (verified)	793
Rubi [A] (verified)	793
Maple [A] (verified)	796
Fricas [A] (verification not implemented)	797
Sympy [F]	798
Maxima [A] (verification not implemented)	798
Giac [F]	799
Mupad [F(-1)]	799
Reduce [F]	800

#### Optimal result

Integrand size = 21, antiderivative size = 275

$$\int x^2(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= -\frac{b(280c^4d^2 + 252c^2de + 75e^2) x \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{1680c^6}$$

$$- \frac{be(84c^2d + 25e) x^3 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{840c^4} - \frac{be^2x^5 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{42c^2}$$

$$+ \frac{1}{3}d^2x^3(a + b\operatorname{sech}^{-1}(cx)) + \frac{2}{5}dex^5(a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{7}e^2x^7(a + b\operatorname{sech}^{-1}(cx))$$

$$+ \frac{b(280c^4d^2 + 252c^2de + 75e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \arcsin(cx)}{1680c^7}$$

output

```
-1/1680*b*(280*c^4*d^2+252*c^2*d*e+75*e^2)*x*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^6-1/840*b*e*(84*c^2*d+25*e)*x^3*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^4-1/42*b*e^2*x^5*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^2+1/3*d^2*x^3*(a+b*arcsech(c*x))+2/5*d*e*x^5*(a+b*arcsech(c*x))+1/7*e^2*x^7*(a+b*arcsech(c*x))+1/1680*b*(280*c^4*d^2+252*c^2*d*e+75*e^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arcsin(c*x)/c^7
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 207, normalized size of antiderivative = 0.75

$$\int x^2(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{16ac^7x^3(35d^2 + 42dex^2 + 15e^2x^4) - bcx\sqrt{\frac{1-cx}{1+cx}}(1+cx)(75e^2 + 2c^2e(126d + 25ex^2)) + 8c^4(35d^2 + 21dex^2 + 5e^2x^4) + 16b^2c^7x^3(35d^2 + 42dex^2 + 15e^2x^4) \operatorname{ArcSech}[cx] + I b^2(280c^4d^2 + 252c^2de + 75e^2) \operatorname{Log}[(-2I)cx + 2\sqrt{\frac{1-cx}{1+cx}}(1+cx)]}{1680c^7}$$

input

```
Integrate[x^2*(d + e*x^2)^2*(a + b*ArcSech[c*x]),x]
```

output

```
(16*a*c^7*x^3*(35*d^2 + 42*d*e*x^2 + 15*e^2*x^4) - b*c*x*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(75*e^2 + 2*c^2*e*(126*d + 25*e*x^2)) + 8*c^4*(35*d^2 + 21*d*e*x^2 + 5*e^2*x^4) + 16*b*c^7*x^3*(35*d^2 + 42*d*e*x^2 + 15*e^2*x^4) *ArcSech[c*x] + I*b*(280*c^4*d^2 + 252*c^2*d*e + 75*e^2)*Log[(-2*I)*c*x + 2*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)]/(1680*c^7)
```

**Rubi [A] (verified)**

Time = 0.46 (sec) , antiderivative size = 212, normalized size of antiderivative = 0.77, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6855, 27, 1590, 27, 363, 262, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$\downarrow 6855$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^2(15e^2x^4 + 42dex^2 + 35d^2)}{105\sqrt{1-c^2x^2}} dx + \frac{1}{3}d^2x^3(a + b\operatorname{sech}^{-1}(cx)) + \frac{2}{5}dex^5(a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{7}e^2x^7(a + b\operatorname{sech}^{-1}(cx))$$

$$\downarrow 27$$

$$\frac{1}{105}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\int\frac{x^2(15e^2x^4+42dex^2+35d^2)}{\sqrt{1-c^2x^2}}dx+\frac{1}{3}d^2x^3(a+b\operatorname{sech}^{-1}(cx))+\frac{2}{5}dex^5(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{7}e^2x^7(a+b\operatorname{sech}^{-1}(cx))$$

↓ 1590

$$\frac{1}{105}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{\int-\frac{3x^2(70c^2d^2+e(84dc^2+25e)x^2)}{\sqrt{1-c^2x^2}}dx}{6c^2}-\frac{5e^2x^5\sqrt{1-c^2x^2}}{2c^2}\right)+\frac{1}{3}d^2x^3(a+b\operatorname{sech}^{-1}(cx))+\frac{2}{5}dex^5(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{7}e^2x^7(a+b\operatorname{sech}^{-1}(cx))$$

↓ 27

$$\frac{1}{105}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{\int\frac{x^2(70c^2d^2+e(84dc^2+25e)x^2)}{\sqrt{1-c^2x^2}}dx}{2c^2}-\frac{5e^2x^5\sqrt{1-c^2x^2}}{2c^2}\right)+\frac{1}{3}d^2x^3(a+b\operatorname{sech}^{-1}(cx))+\frac{2}{5}dex^5(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{7}e^2x^7(a+b\operatorname{sech}^{-1}(cx))$$

↓ 363

$$\frac{1}{105}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{(280c^4d^2+252c^2de+75e^2)\int\frac{x^2}{\sqrt{1-c^2x^2}}dx}{4c^2}-\frac{ex^3\sqrt{1-c^2x^2}(84c^2d+25e)}{4c^2}-\frac{5e^2x^5\sqrt{1-c^2x^2}}{2c^2}\right)+\frac{1}{3}d^2x^3(a+b\operatorname{sech}^{-1}(cx))+\frac{2}{5}dex^5(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{7}e^2x^7(a+b\operatorname{sech}^{-1}(cx))$$

↓ 262

$$\frac{1}{105}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{(280c^4d^2+252c^2de+75e^2)\left(\frac{\int\frac{1}{\sqrt{1-c^2x^2}}dx}{2c^2}-\frac{x\sqrt{1-c^2x^2}}{2c^2}\right)}{4c^2}-\frac{ex^3\sqrt{1-c^2x^2}(84c^2d+25e)}{4c^2}-\frac{5e^2x^5\sqrt{1-c^2x^2}}{2c^2}\right)+\frac{1}{3}d^2x^3(a+b\operatorname{sech}^{-1}(cx))+\frac{2}{5}dex^5(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{7}e^2x^7(a+b\operatorname{sech}^{-1}(cx))$$

↓ 223

$$\frac{1}{3}d^2x^3(a+b\operatorname{sech}^{-1}(cx))+\frac{2}{5}dex^5(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{7}e^2x^7(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{105}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{\left(\frac{\arcsin(cx)}{2c^3}-\frac{x\sqrt{1-c^2x^2}}{2c^2}\right)(280c^4d^2+252c^2de+75e^2)}{4c^2}-\frac{ex^3\sqrt{1-c^2x^2}(84c^2d+25e)}{4c^2}-\frac{5e^2x^5\sqrt{1-c^2x^2}}{2c^2}\right)$$

input `Int[x^2*(d + e*x^2)^2*(a + b*ArcSech[c*x]),x]`

output `(d^2*x^3*(a + b*ArcSech[c*x]))/3 + (2*d*e*x^5*(a + b*ArcSech[c*x]))/5 + (e^2*x^7*(a + b*ArcSech[c*x]))/7 + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-5*e^2*x^5*Sqrt[1 - c^2*x^2])/(2*c^2) + (-1/4*(e*(84*c^2*d + 25*e)*x^3*Sqrt[1 - c^2*x^2])/c^2 + ((280*c^4*d^2 + 252*c^2*d*e + 75*e^2)*(-1/2*(x*Sqrt[1 - c^2*x^2])/c^2 + ArcSin[c*x]/(2*c^3)))/(4*c^2))/(2*c^2))/105`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 262 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 363 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(b*e*(m + 2*p + 3))), x] - Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(b*(m + 2*p + 3)) Int[(e*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + 2*p + 3, 0]`

rule 1590

```
Int[((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[c^p*(f*x)^(m + 4*p - 1)*((d + e*x^2)^(q + 1)/(e*f^(4*p - 1)*(m + 4*p + 2*q + 1))), x] + Simp[1/(e*(m + 4*p + 2*q + 1)) Int[(f*x)^m*(d + e*x^2)^q*ExpandToSum[e*(m + 4*p + 2*q + 1)*((a + b*x^2 + c*x^4)^p - c^p*x^(4*p)) - d*c^p*(m + 4*p - 1)*x^(4*p - 2), x], x] /; FreeQ[{a, b, c, d, e, f, m, q}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && !IntegerQ[q] && NeQ[m + 4*p + 2*q + 1, 0]
```

rule 6855

```
Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

### Maple [A] (verified)

Time = 0.53 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.04

method	result
parts	$a\left(\frac{1}{7}e^2x^7 + \frac{2}{5}dex^5 + \frac{1}{3}d^2x^3\right) + \frac{b\left(\frac{c^3 \operatorname{arcsech}(cx)e^2x^7}{7} + \frac{2c^3 \operatorname{arcsech}(cx)de x^5}{5} + \frac{\operatorname{arcsech}(cx)d^2c^3x^3}{3} + \sqrt{-\frac{cx-1}{cx}}x\sqrt{\frac{cx+1}{cx}}\right)}{c^4}$
derivativedivides	$\frac{a\left(\frac{1}{3}d^2c^7x^3 + \frac{2}{5}dc^7ex^5 + \frac{1}{7}e^2c^7x^7\right)}{c^4} + \frac{b\left(\frac{\operatorname{arcsech}(cx)d^2c^7x^3}{3} + \frac{2 \operatorname{arcsech}(cx)dc^7ex^5}{5} + \frac{\operatorname{arcsech}(cx)e^2c^7x^7}{7} - \sqrt{-\frac{cx-1}{cx}}cx\sqrt{\frac{cx+1}{cx}}\right)}{c^4}$
default	$\frac{a\left(\frac{1}{3}d^2c^7x^3 + \frac{2}{5}dc^7ex^5 + \frac{1}{7}e^2c^7x^7\right)}{c^4} + \frac{b\left(\frac{\operatorname{arcsech}(cx)d^2c^7x^3}{3} + \frac{2 \operatorname{arcsech}(cx)dc^7ex^5}{5} + \frac{\operatorname{arcsech}(cx)e^2c^7x^7}{7} - \sqrt{-\frac{cx-1}{cx}}cx\sqrt{\frac{cx+1}{cx}}\right)}{c^4}$

input

```
int(x^2*(e*x^2+d)^2*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)
```

output

```
a*(1/7*e^2*x^7+2/5*d*e*x^5+1/3*d^2*x^3)+b/c^3*(1/7*c^3*arcsech(c*x)*e^2*x^7+2/5*c^3*arcsech(c*x)*d*e*x^5+1/3*arcsech(c*x)*d^2*c^3*x^3+1/1680/c^3*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(-40*e^2*(-c^2*x^2+1)^(1/2)*c^5*x^5-168*c^5*d*e*(-c^2*x^2+1)^(1/2)*x^3-280*d^2*c^5*x*(-c^2*x^2+1)^(1/2)+280*d^2*c^4*arcsin(c*x)-50*e^2*c^3*x^3*(-c^2*x^2+1)^(1/2)-252*e*(-c^2*x^2+1)^(1/2)*c^3*d*x+252*e*arcsin(c*x)*c^2*d-75*(-c^2*x^2+1)^(1/2)*e^2*c*x+75*arcsin(c*x)*e^2)/(-c^2*x^2+1)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 341, normalized size of antiderivative = 1.24

$$\int x^2(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{240 ac^7 e^2 x^7 + 672 ac^7 d e x^5 + 560 ac^7 d^2 x^3 - 2(280 bc^4 d^2 + 252 bc^2 d e + 75 be^2) \arctan\left(\frac{cx\sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - 1}{cx}\right) - \dots}{\dots}$$

input

```
integrate(x^2*(e*x^2+d)^2*(a+b*arcsech(c*x)),x, algorithm="fricas")
```

output

```
1/1680*(240*a*c^7*e^2*x^7 + 672*a*c^7*d*e*x^5 + 560*a*c^7*d^2*x^3 - 2*(280*b*c^4*d^2 + 252*b*c^2*d*e + 75*b*e^2)*arctan((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/(c*x)) - 16*(35*b*c^7*d^2 + 42*b*c^7*d*e + 15*b*c^7*e^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) + 16*(15*b*c^7*e^2*x^7 + 42*b*c^7*d*e*x^5 + 35*b*c^7*d^2*x^3 - 35*b*c^7*d^2 - 42*b*c^7*d*e - 15*b*c^7*e^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (40*b*c^6*e^2*x^6 + 2*(84*b*c^6*d*e + 25*b*c^4*e^2)*x^4 + (280*b*c^6*d^2 + 252*b*c^4*d*e + 75*b*c^2*e^2)*x^2)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/c^7
```

**Sympy [F]**

$$\int x^2(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx = \int x^2(a + b\operatorname{asech}(cx)) (d + ex^2)^2 dx$$

input `integrate(x**2*(e*x**2+d)**2*(a+b*asech(c*x)), x)`

output `Integral(x**2*(a + b*asech(c*x))*(d + e*x**2)**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.19

$$\begin{aligned} \int x^2(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx &= \frac{1}{7} ae^2 x^7 + \frac{2}{5} adex^5 + \frac{1}{3} ad^2 x^3 \\ &+ \frac{1}{6} \left( 2x^3 \operatorname{arsech}(cx) - \frac{\frac{\sqrt{\frac{1}{c^2 x^2} - 1}}{c^2 \left(\frac{1}{c^2 x^2} - 1\right) + c^2} + \frac{\arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right)}{c^2}}{c} \right) bd^2 \\ &+ \frac{1}{20} \left( 8x^5 \operatorname{arsech}(cx) - \frac{\frac{3\left(\frac{1}{c^2 x^2} - 1\right)^{\frac{3}{2}} + 5\sqrt{\frac{1}{c^2 x^2} - 1}}{c^4 \left(\frac{1}{c^2 x^2} - 1\right)^2 + 2c^4 \left(\frac{1}{c^2 x^2} - 1\right) + c^4} + \frac{3\arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right)}{c^4}}{c} \right) bde \\ &+ \frac{1}{336} \left( 48x^7 \operatorname{arsech}(cx) - \frac{\frac{15\left(\frac{1}{c^2 x^2} - 1\right)^{\frac{5}{2}} + 40\left(\frac{1}{c^2 x^2} - 1\right)^{\frac{3}{2}} + 33\sqrt{\frac{1}{c^2 x^2} - 1}}{c^6 \left(\frac{1}{c^2 x^2} - 1\right)^3 + 3c^6 \left(\frac{1}{c^2 x^2} - 1\right)^2 + 3c^6 \left(\frac{1}{c^2 x^2} - 1\right) + c^6} + \frac{15\arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right)}{c^6}}{c} \right) be^2 \end{aligned}$$

input `integrate(x^2*(e*x^2+d)^2*(a+b*arcsech(c*x)), x, algorithm="maxima")`

output

```
1/7*a*e^2*x^7 + 2/5*a*d*e*x^5 + 1/3*a*d^2*x^3 + 1/6*(2*x^3*arcsech(c*x) -
(sqrt(1/(c^2*x^2) - 1))/(c^2*(1/(c^2*x^2) - 1) + c^2) + arctan(sqrt(1/(c^2*
x^2) - 1))/c^2)/c)*b*d^2 + 1/20*(8*x^5*arcsech(c*x) - ((3*(1/(c^2*x^2) - 1)
)^(3/2) + 5*sqrt(1/(c^2*x^2) - 1))/(c^4*(1/(c^2*x^2) - 1)^2 + 2*c^4*(1/(c^
2*x^2) - 1) + c^4) + 3*arctan(sqrt(1/(c^2*x^2) - 1))/c^4)/c)*b*d*e + 1/336
*(48*x^7*arcsech(c*x) - ((15*(1/(c^2*x^2) - 1)^(5/2) + 40*(1/(c^2*x^2) - 1)
)^(3/2) + 33*sqrt(1/(c^2*x^2) - 1))/(c^6*(1/(c^2*x^2) - 1)^3 + 3*c^6*(1/(c
^2*x^2) - 1)^2 + 3*c^6*(1/(c^2*x^2) - 1) + c^6) + 15*arctan(sqrt(1/(c^2*x^
2) - 1))/c^6)/c)*b*e^2
```

**Giac [F]**

$$\int x^2(d + ex^2)^2(a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^2(b \operatorname{arsech}(cx) + a)x^2 dx$$

input

```
integrate(x^2*(e*x^2+d)^2*(a+b*arcsech(c*x)),x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)^2*(b*arcsech(c*x) + a)*x^2, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int x^2(d + ex^2)^2(a + b\operatorname{sech}^{-1}(cx)) dx = \int x^2(ex^2 + d)^2 \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input

```
int(x^2*(d + e*x^2)^2*(a + b*acosh(1/(c*x))),x)
```

output

```
int(x^2*(d + e*x^2)^2*(a + b*acosh(1/(c*x))), x)
```



**Reduce [F]**

$$\int x^2 (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \left( \int a \operatorname{sech}(cx) x^6 dx \right) b e^2$$

$$+ 2 \left( \int a \operatorname{sech}(cx) x^4 dx \right) b d e$$

$$+ \left( \int a \operatorname{sech}(cx) x^2 dx \right) b d^2$$

$$+ \frac{a d^2 x^3}{3} + \frac{2 a d e x^5}{5} + \frac{a e^2 x^7}{7}$$

input `int(x^2*(e*x^2+d)^2*(a+b*asech(c*x)),x)`

output `(105*int(asech(c*x)*x**6,x)*b*e**2 + 210*int(asech(c*x)*x**4,x)*b*d*e + 105*int(asech(c*x)*x**2,x)*b*d**2 + 35*a*d**2*x**3 + 42*a*d*e*x**5 + 15*a*e**2*x**7)/105`

### 3.100 $\int (d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	801
Mathematica [C] (verified)	802
Rubi [A] (verified)	802
Maple [A] (verified)	805
Fricas [B] (verification not implemented)	805
Sympy [F]	806
Maxima [A] (verification not implemented)	806
Giac [F]	807
Mupad [F(-1)]	807
Reduce [F]	808

#### Optimal result

Integrand size = 18, antiderivative size = 204

$$\int (d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= -\frac{be(40c^2d + 9e)x\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{120c^4} - \frac{be^2x^3\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{20c^2}$$

$$+ d^2x(a + b\operatorname{sech}^{-1}(cx)) + \frac{2}{3}dex^3(a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{5}e^2x^5(a + b\operatorname{sech}^{-1}(cx))$$

$$+ \frac{b(120c^4d^2 + 40c^2de + 9e^2)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\arcsin(cx)}{120c^5}$$

output

```
-1/120*b*e*(40*c^2*d+9*e)*x*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^4-1/20*b*e^2*x^3*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^2+d^2*x*(a+b*arcsech(c*x))+2/3*d*e*x^3*(a+b*arcsech(c*x))+1/5*e^2*x^5*(a+b*arcsech(c*x))+1/120*b*(120*c^4*d^2+40*c^2*d*e+9*e^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arcsin(c*x)/c^5
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.18 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.85

$$\int (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{8ac^5x(15d^2 + 10dex^2 + 3e^2x^4) - bcex\sqrt{\frac{1-cx}{1+cx}}(1+cx)(9e + c^2(40d + 6ex^2)) + 8bc^5x(15d^2 + 10dex^2 + 3e^2x^4)}{120c^5}$$

input

```
Integrate[(d + e*x^2)^2*(a + b*ArcSech[c*x]),x]
```

output

```
(8*a*c^5*x*(15*d^2 + 10*d*e*x^2 + 3*e^2*x^4) - b*c*e*x*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(9*e + c^2*(40*d + 6*e*x^2)) + 8*b*c^5*x*(15*d^2 + 10*d*e*x^2 + 3*e^2*x^4)*ArcSech[c*x] + I*b*(120*c^4*d^2 + 40*c^2*d*e + 9*e^2)*Log[(-2*I)*c*x + 2*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)])/(120*c^5)
```

**Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.86, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6845, 27, 1473, 25, 299, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$\downarrow 6845$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{3e^2x^4 + 10dex^2 + 15d^2}{15\sqrt{1-c^2x^2}} dx + d^2x(a + b \operatorname{sech}^{-1}(cx)) + \frac{2}{3}dex^3(a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{5}e^2x^5(a + b \operatorname{sech}^{-1}(cx))$$

$$\downarrow 27$$

$$\frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\int\frac{3e^2x^4+10dex^2+15d^2}{\sqrt{1-c^2x^2}}dx+d^2x(a+b\operatorname{sech}^{-1}(cx))+\frac{2}{3}dex^3(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{5}e^2x^5(a+b\operatorname{sech}^{-1}(cx))$$

↓ 1473

$$\frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{\int-\frac{60c^2d^2+e(40dc^2+9e)x^2}{\sqrt{1-c^2x^2}}dx}{4c^2}-\frac{3e^2x^3\sqrt{1-c^2x^2}}{4c^2}\right)+d^2x(a+b\operatorname{sech}^{-1}(cx))+\frac{2}{3}dex^3(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{5}e^2x^5(a+b\operatorname{sech}^{-1}(cx))$$

↓ 25

$$\frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{\int\frac{60c^2d^2+e(40dc^2+9e)x^2}{\sqrt{1-c^2x^2}}dx}{4c^2}-\frac{3e^2x^3\sqrt{1-c^2x^2}}{4c^2}\right)+d^2x(a+b\operatorname{sech}^{-1}(cx))+\frac{2}{3}dex^3(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{5}e^2x^5(a+b\operatorname{sech}^{-1}(cx))$$

↓ 299

$$\frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{\frac{(120c^4d^2+40c^2de+9e^2)\int\frac{1}{\sqrt{1-c^2x^2}}dx}{2c^2}-\frac{ex\sqrt{1-c^2x^2}(40c^2d+9e)}{2c^2}}{4c^2}-\frac{3e^2x^3\sqrt{1-c^2x^2}}{4c^2}\right)+d^2x(a+b\operatorname{sech}^{-1}(cx))+\frac{2}{3}dex^3(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{5}e^2x^5(a+b\operatorname{sech}^{-1}(cx))$$

↓ 223

$$d^2x(a+b\operatorname{sech}^{-1}(cx))+\frac{2}{3}dex^3(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{5}e^2x^5(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{\arcsin(cx)\frac{(120c^4d^2+40c^2de+9e^2)}{2c^3}-\frac{ex\sqrt{1-c^2x^2}(40c^2d+9e)}{2c^2}}{4c^2}-\frac{3e^2x^3\sqrt{1-c^2x^2}}{4c^2}\right)$$

input

```
Int[(d + e*x^2)^2*(a + b*ArcSech[c*x]), x]
```

output

```
d^2*x*(a + b*ArcSech[c*x]) + (2*d*e*x^3*(a + b*ArcSech[c*x]))/3 + (e^2*x^5*(a + b*ArcSech[c*x]))/5 + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-3*e^2*x^3*Sqrt[1 - c^2*x^2])/(4*c^2) + (-1/2*(e*(40*c^2*d + 9*e))*x*Sqrt[1 - c^2*x^2])/c^2 + ((120*c^4*d^2 + 40*c^2*d*e + 9*e^2)*ArcSin[c*x])/(2*c^3))/(4*c^2))/15
```

## Definitions of rubi rules used

- rule 25  $\text{Int}[-(F_x), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[F_x, x], x]$
- rule 27  $\text{Int}[(a_*)(F_x), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_*)(G_x)] /; \text{FreeQ}[b, x]$
- rule 223  $\text{Int}[1/\text{Sqrt}[(a_*) + (b_*)(x_)^2], x\_Symbol] \rightarrow \text{Simp}[\text{ArcSin}[\text{Rt}[-b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[-b, 2], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{GtQ}[a, 0] \ \&\& \ \text{NegQ}[b]$
- rule 299  $\text{Int}[((a_*) + (b_*)(x_)^2)^{(p_*)}*((c_*) + (d_*)(x_)^2), x\_Symbol] \rightarrow \text{Simp}[d*x*((a + b*x^2)^{(p + 1)}/(b*(2*p + 3))), x] - \text{Simp}[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) \text{ Int}[(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[2*p + 3, 0]$
- rule 1473  $\text{Int}[((d_*) + (e_*)(x_)^2)^{(q_*)}*((a_*) + (b_*)(x_)^2 + (c_*)(x_)^4)^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[c^p*x^{(4*p - 1)}*((d + e*x^2)^{(q + 1)}/(e*(4*p + 2*q + 1))), x] + \text{Simp}[1/(e*(4*p + 2*q + 1)) \text{ Int}[(d + e*x^2)^q*\text{ExpandToSum}[e*(4*p + 2*q + 1)*(a + b*x^2 + c*x^4)^p - d*c^p*(4*p - 1)*x^{(4*p - 2)} - e*c^p*(4*p + 2*q + 1)*x^{(4*p)}, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, q\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ !\text{LtQ}[q, -1]$
- rule 6845  $\text{Int}[((a_*) + \text{ArcSech}[(c_*)(x_)])*(b_*)*((d_*) + (e_*)(x_)^2)^{(p_*)}, x\_Symbol] \rightarrow \text{With}[\{u = \text{IntHide}[(d + e*x^2)^p, x]\}, \text{Simp}[(a + b*\text{ArcSech}[c*x]) u, x] + \text{Simp}[b*\text{Sqrt}[1 + c*x]*\text{Sqrt}[1/(1 + c*x)] \text{ Int}[\text{SimplifyIntegrand}[u/(x*\text{Sqrt}[1 - c*x]*\text{Sqrt}[1 + c*x]), x], x], x]] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ (\text{IGtQ}[p, 0] \ || \ \text{ILtQ}[p + 1/2, 0])$

### Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.02

method	result
parts	$a\left(\frac{1}{5}e^2x^5 + \frac{2}{3}dex^3 + d^2x\right) + \frac{b\left(\frac{c \operatorname{arcsech}(cx)e^2x^5}{5} + \frac{2c \operatorname{arcsech}(cx)de x^3}{3} + \operatorname{arcsech}(cx)cx d^2 + \sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}}\right)}{c}$
derivativelimit	$\frac{a\left(d^2c^5x + \frac{2}{3}dc^5ex^3 + \frac{1}{5}e^2c^5x^5\right)}{c^4} + \frac{b\left(\operatorname{arcsech}(cx)d^2c^5x + \frac{2 \operatorname{arcsech}(cx)d c^5ex^3}{3} + \frac{\operatorname{arcsech}(cx)e^2c^5x^5}{5} + \sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}}\right)(120d^2)}{c}$
default	$\frac{a\left(d^2c^5x + \frac{2}{3}dc^5ex^3 + \frac{1}{5}e^2c^5x^5\right)}{c^4} + \frac{b\left(\operatorname{arcsech}(cx)d^2c^5x + \frac{2 \operatorname{arcsech}(cx)d c^5ex^3}{3} + \frac{\operatorname{arcsech}(cx)e^2c^5x^5}{5} + \sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}}\right)(120d^2)}{c}$

```
input int((e*x^2+d)^2*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)
```

```
output a*(1/5*e^2*x^5+2/3*d*e*x^3+d^2*x)+b/c*(1/5*c*arcsech(c*x)*e^2*x^5+2/3*c*arcsech(c*x)*d*e*x^3+arcsech(c*x)*c*x*d^2+1/120/c^3*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(120*d^2*c^4*arcsin(c*x)-40*e*(-c^2*x^2+1)^(1/2)*c^3*d*x-6*e^2*c^3*x^3*(-c^2*x^2+1)^(1/2)+40*e*arcsin(c*x)*c^2*d-9*(-c^2*x^2+1)^(1/2)*e^2*c*x+9*arcsin(c*x)*e^2)/(-c^2*x^2+1)^(1/2))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 305 vs. 2(130) = 260.

Time = 0.21 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.50

$$\int (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{24ac^5e^2x^5 + 80ac^5dex^3 + 120ac^5d^2x - 2(120bc^4d^2 + 40bc^2de + 9be^2) \arctan\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{cx}\right) - 8(15}{1}$$

```
input integrate((e*x^2+d)^2*(a+b*arcsech(c*x)),x, algorithm="fricas")
```

output

```
1/120*(24*a*c^5*e^2*x^5 + 80*a*c^5*d*e*x^3 + 120*a*c^5*d^2*x - 2*(120*b*c^4*d^2 + 40*b*c^2*d*e + 9*b*e^2)*arctan((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/(c*x)) - 8*(15*b*c^5*d^2 + 10*b*c^5*d*e + 3*b*c^5*e^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) + 8*(3*b*c^5*e^2*x^5 + 10*b*c^5*d*e*x^3 + 15*b*c^5*d^2*x - 15*b*c^5*d^2 - 10*b*c^5*d*e - 3*b*c^5*e^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (6*b*c^4*e^2*x^4 + (40*b*c^4*d*e + 9*b*c^2*e^2)*x^2)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/c^5
```

**Sympy [F]**

$$\int (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \int (a + b \operatorname{asech}(cx)) (d + ex^2)^2 dx$$

input

```
integrate((e*x**2+d)**2*(a+b*asech(c*x)),x)
```

output

```
Integral((a + b*asech(c*x))*(d + e*x**2)**2, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.10

$$\begin{aligned} & \int (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx \\ &= \frac{1}{5} ae^2 x^5 + \frac{2}{3} adex^3 + \frac{1}{3} \left( 2x^3 \operatorname{arsech}(cx) - \frac{\frac{\sqrt{\frac{1}{c^2 x^2} - 1}}{c^2} + \frac{\arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right)}{c^2}}{\left(\frac{1}{c^2 x^2} - 1\right) + c^2} + \frac{\arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right)}{c^2} \right) bde \\ &+ \frac{1}{40} \left( 8x^5 \operatorname{arsech}(cx) - \frac{\frac{3\left(\frac{1}{c^2 x^2} - 1\right)^{\frac{3}{2}} + 5\sqrt{\frac{1}{c^2 x^2} - 1}}{c^4} + \frac{3\arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right)}{c^4}}{\left(\frac{1}{c^2 x^2} - 1\right)^2 + 2c^4\left(\frac{1}{c^2 x^2} - 1\right) + c^4} + \frac{3\arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right)}{c^4} \right) be^2 \\ &+ ad^2 x + \frac{\left(cx \operatorname{arsech}(cx) - \arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right)\right) bd^2}{c} \end{aligned}$$

input `integrate((e*x^2+d)^2*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `1/5*a*e^2*x^5 + 2/3*a*d*e*x^3 + 1/3*(2*x^3*arcsech(c*x) - (sqrt(1/(c^2*x^2) - 1)/(c^2*(1/(c^2*x^2) - 1) + c^2) + arctan(sqrt(1/(c^2*x^2) - 1)/c^2)/c)*b*d*e + 1/40*(8*x^5*arcsech(c*x) - ((3*(1/(c^2*x^2) - 1)^(3/2) + 5*sqrt(1/(c^2*x^2) - 1))/(c^4*(1/(c^2*x^2) - 1)^2 + 2*c^4*(1/(c^2*x^2) - 1) + c^4) + 3*arctan(sqrt(1/(c^2*x^2) - 1)/c^4)/c)*b*e^2 + a*d^2*x + (c*x*arcsech(c*x) - arctan(sqrt(1/(c^2*x^2) - 1)))*b*d^2/c`

### Giac [F]

$$\int (d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^2 (b \operatorname{ar} \operatorname{sech}(cx) + a) dx$$

input `integrate((e*x^2+d)^2*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((e*x^2 + d)^2*(b*arcsech(c*x) + a), x)`

### Mupad [F(-1)]

Timed out.

$$\int (d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^2 \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int((d + e*x^2)^2*(a + b*acosh(1/(c*x))),x)`

output `int((d + e*x^2)^2*(a + b*acosh(1/(c*x))), x)`



**Reduce [F]**

$$\int (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \left( \int \operatorname{asech}(cx) dx \right) b d^2 + \left( \int \operatorname{asech}(cx) x^4 dx \right) b e^2$$

$$+ 2 \left( \int \operatorname{asech}(cx) x^2 dx \right) b d e$$

$$+ a d^2 x + \frac{2 a d e x^3}{3} + \frac{a e^2 x^5}{5}$$

input `int((e*x^2+d)^2*(a+b*asech(c*x)),x)`

output `(15*int(asech(c*x),x)*b*d**2 + 15*int(asech(c*x)*x**4,x)*b*e**2 + 30*int(asech(c*x)*x**2,x)*b*d*e + 15*a*d**2*x + 10*a*d*e*x**3 + 3*a*e**2*x**5)/15`

**3.101**  $\int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x^2} dx$

Optimal result	809
Mathematica [C] (verified)	810
Rubi [A] (verified)	810
Maple [A] (verified)	813
Fricas [B] (verification not implemented)	813
Sympy [F]	814
Maxima [A] (verification not implemented)	814
Giac [F]	815
Mupad [F(-1)]	815
Reduce [F]	816

**Optimal result**

Integrand size = 21, antiderivative size = 177

$$\int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x^2} dx = \frac{bd^2 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{x} - \frac{be^2x \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{6c^2} - \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{x} + 2dex(a+b\operatorname{sech}^{-1}(cx)) + \frac{1}{3}e^2x^3(a+b\operatorname{sech}^{-1}(cx)) + \frac{be(12c^2d+e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \arcsin(cx)}{6c^3}$$

output

```
b*d^2*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/x-1/6*b*e^2*x*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^2-d^2*(a+b*arcsech(c*x))/x+2*d*e*x*(a+b*arcsech(c*x))+1/3*e^2*x^3*(a+b*arcsech(c*x))+1/6*b*e*(12*c^2*d+e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arcsin(c*x)/c^3
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.16 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.89

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx$$

$$= \frac{-bc \sqrt{\frac{1-cx}{1+cx}} (1+cx) (-6c^2 d^2 + e^2 x^2) + 2ac^3 (-3d^2 + 6dex^2 + e^2 x^4) + 2bc^3 (-3d^2 + 6dex^2 + e^2 x^4) \operatorname{sech}^{-1}(cx)}{6c^3 x}$$

input

```
Integrate[((d + e*x^2)^2*(a + b*ArcSech[c*x]))/x^2,x]
```

output

```
(-(b*c*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(-6*c^2*d^2 + e^2*x^2)) + 2*a*c^3*(-3*d^2 + 6*d*e*x^2 + e^2*x^4) + 2*b*c^3*(-3*d^2 + 6*d*e*x^2 + e^2*x^4)*ArcSech[c*x] + I*b*e*(12*c^2*d + e)*x*Log[(-2*I)*c*x + 2*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)])/(6*c^3*x)
```

**Rubi [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.80, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {6855, 27, 1588, 27, 299, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx$$

$$\downarrow \text{6855}$$

$$b \sqrt{\frac{1}{cx + 1}} \sqrt{cx + 1} \int -\frac{-e^2 x^4 - 6dex^2 + 3d^2}{3x^2 \sqrt{1 - c^2 x^2}} dx - \frac{d^2 (a + b \operatorname{sech}^{-1}(cx))}{x} + 2dex(a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{3} e^2 x^3 (a + b \operatorname{sech}^{-1}(cx))$$

$$\downarrow \text{27}$$

$$\begin{aligned}
& -\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\int\frac{-e^2x^4-6dex^2+3d^2}{x^2\sqrt{1-c^2x^2}}dx-\frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{x}+ \\
& \quad 2dex(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{3}e^2x^3(a+b\operatorname{sech}^{-1}(cx)) \\
& \quad \downarrow 1588 \\
& -\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\int\frac{e(ex^2+6d)}{\sqrt{1-c^2x^2}}dx-\frac{3d^2\sqrt{1-c^2x^2}}{x}\right)-\frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{x}+ \\
& \quad 2dex(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{3}e^2x^3(a+b\operatorname{sech}^{-1}(cx)) \\
& \quad \downarrow 27 \\
& -\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-e\int\frac{ex^2+6d}{\sqrt{1-c^2x^2}}dx-\frac{3d^2\sqrt{1-c^2x^2}}{x}\right)-\frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{x}+ \\
& \quad 2dex(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{3}e^2x^3(a+b\operatorname{sech}^{-1}(cx)) \\
& \quad \downarrow 299 \\
& -\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-e\left(\frac{(12c^2d+e)\int\frac{1}{\sqrt{1-c^2x^2}}dx}{2c^2}-\frac{ex\sqrt{1-c^2x^2}}{2c^2}\right)-\frac{3d^2\sqrt{1-c^2x^2}}{x}\right)- \\
& \quad \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{x}+2dex(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{3}e^2x^3(a+b\operatorname{sech}^{-1}(cx)) \\
& \quad \downarrow 223 \\
& -\frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{x}+2dex(a+b\operatorname{sech}^{-1}(cx))+\frac{1}{3}e^2x^3(a+b\operatorname{sech}^{-1}(cx))- \\
& \quad \frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-e\left(\frac{\arcsin(cx)(12c^2d+e)}{2c^3}-\frac{ex\sqrt{1-c^2x^2}}{2c^2}\right)-\frac{3d^2\sqrt{1-c^2x^2}}{x}\right)
\end{aligned}$$

input `Int[((d + e*x^2)^2*(a + b*ArcSech[c*x]))/x^2,x]`

output `-((d^2*(a + b*ArcSech[c*x]))/x) + 2*d*e*x*(a + b*ArcSech[c*x]) + (e^2*x^3*(a + b*ArcSech[c*x]))/3 - (b*sqrt[(1 + c*x)^(-1)]*sqrt[1 + c*x]*((-3*d^2*sqrt[1 - c^2*x^2])/x - e*(-1/2*(e*x*sqrt[1 - c^2*x^2])/c^2 + ((12*c^2*d + e)*ArcSin[c*x])/(2*c^3))))/3`

## Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`
- rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 299 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[d*x*((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NeQ[2*p + 3, 0]`
- rule 1588 `Int[((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, f*x, x], R = PolynomialRemainder[(a + b*x^2 + c*x^4)^p, f*x, x]}, Simp[R*(f*x)^(m + 1)*((d + e*x^2)^(q + 1)/(d*f*(m + 1))), x] + Simp[1/(d*f^(2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^q*ExpandToSum[d*f*(m + 1)*(Qx/x) - e*R*(m + 2*q + 3), x], x], x] /; FreeQ[{a, b, c, d, e, f, q}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && LtQ[m, -1]`
- rule 6855 `Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))`

### Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.04

method	result
parts	$a\left(\frac{e^2x^3}{3} + 2dex - \frac{d^2}{x}\right) + bc\left(\frac{\operatorname{arcsech}(cx)e^2x^3}{3c} + \frac{2 \operatorname{arcsech}(cx)dex}{c} - \frac{\operatorname{arcsech}(cx)d^2}{cx} + \sqrt{\frac{-cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}\right)$
derivativedivides	$c\left(\frac{a\left(2c^3dex + \frac{e^2c^3x^3}{3} - \frac{c^3d^2}{x}\right)}{c^4} + \frac{b\left(2 \operatorname{arcsech}(cx)c^3dex + \frac{\operatorname{arcsech}(cx)e^2c^3x^3}{3} - \frac{\operatorname{arcsech}(cx)c^3d^2}{x} + \sqrt{\frac{-cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}\right)}{c^4}\right)$
default	$c\left(\frac{a\left(2c^3dex + \frac{e^2c^3x^3}{3} - \frac{c^3d^2}{x}\right)}{c^4} + \frac{b\left(2 \operatorname{arcsech}(cx)c^3dex + \frac{\operatorname{arcsech}(cx)e^2c^3x^3}{3} - \frac{\operatorname{arcsech}(cx)c^3d^2}{x} + \sqrt{\frac{-cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}\right)}{c^4}\right)$

input `int((e*x^2+d)^2*(a+b*arcsech(c*x))/x^2,x,method=_RETURNVERBOSE)`

output `a*(1/3*e^2*x^3+2*d*e*x-d^2/x)+b*c*(1/3/c*arcsech(c*x)*e^2*x^3+2/c*arcsech(c*x)*d*e*x-arcsech(c*x)*d^2/c/x+1/6/c^4*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)*(6*(-c^2*x^2+1)^(1/2)*c^4*d^2+12*arcsin(c*x)*c^3*d*e*x-e^2*(-c^2*x^2+1)^(1/2)*c^2*x^2+arcsin(c*x)*e^2*c*x)/(-c^2*x^2+1)^(1/2))`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 287 vs. 2(107) = 214.

Time = 0.15 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.62

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx$$

$$= \frac{2ac^3e^2x^4 + 12ac^3dex^2 - 6ac^3d^2 - 2(12bc^2de + be^2)x \arctan\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{cx}\right) + 2(3bc^3d^2 - 6bc^3de - b^2e^2)}{c^4}$$

input `integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x^2,x, algorithm="fricas")`

output

```
1/6*(2*a*c^3*e^2*x^4 + 12*a*c^3*d*e*x^2 - 6*a*c^3*d^2 - 2*(12*b*c^2*d*e +
b*e^2)*x*arctan((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/(c*x)) + 2*(3*b*c
^3*d^2 - 6*b*c^3*d*e - b*c^3*e^2)*x*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)
) - 1)/x) + 2*(b*c^3*e^2*x^4 + 6*b*c^3*d*e*x^2 - 3*b*c^3*d^2 + (3*b*c^3*d^
2 - 6*b*c^3*d*e - b*c^3*e^2)*x)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) +
1)/(c*x)) + (6*b*c^4*d^2*x - b*c^2*e^2*x^3)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2))
)/(c^3*x)
```

**Sympy [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{(a + b \operatorname{asech}(cx)) (d + ex^2)^2}{x^2} dx$$

input

```
integrate((e*x**2+d)**2*(a+b*asech(c*x))/x**2,x)
```

output

```
Integral((a + b*asech(c*x))*(d + e*x**2)**2/x**2, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.86

$$\begin{aligned} & \int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx \\ &= \frac{1}{3} a e^2 x^3 + \left( c \sqrt{\frac{1}{c^2 x^2} - 1} - \frac{\operatorname{arsech}(cx)}{x} \right) b d^2 \\ &+ \frac{1}{6} \left( 2 x^3 \operatorname{arsech}(cx) - \frac{\frac{\sqrt{\frac{1}{c^2 x^2} - 1}}{c^2 \left( \frac{1}{c^2 x^2} - 1 \right) + c^2} + \frac{\arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right)}{c^2}}{c} \right) b e^2 \\ &+ 2 a d e x + \frac{2 \left( c x \operatorname{arsech}(cx) - \arctan\left(\sqrt{\frac{1}{c^2 x^2} - 1}\right) \right) b d e}{c} - \frac{a d^2}{x} \end{aligned}$$

input

```
integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x^2,x, algorithm="maxima")
```

output

```
1/3*a*e^2*x^3 + (c*sqrt(1/(c^2*x^2) - 1) - arcsech(c*x)/x)*b*d^2 + 1/6*(2*
x^3*arcsech(c*x) - (sqrt(1/(c^2*x^2) - 1)/(c^2*(1/(c^2*x^2) - 1) + c^2) +
arctan(sqrt(1/(c^2*x^2) - 1))/c^2)/c)*b*e^2 + 2*a*d*e*x + 2*(c*x*arcsech(c
*x) - arctan(sqrt(1/(c^2*x^2) - 1)))*b*d*e/c - a*d^2/x
```

**Giac [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{ar} \operatorname{sech}(cx) + a)}{x^2} dx$$

input

```
integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x^2,x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)^2*(b*arcsech(c*x) + a)/x^2, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{(ex^2 + d)^2 (a + b \operatorname{acosh}(\frac{1}{cx}))}{x^2} dx$$

input

```
int(((d + e*x^2)^2*(a + b*acosh(1/(c*x))))/x^2,x)
```

output

```
int(((d + e*x^2)^2*(a + b*acosh(1/(c*x))))/x^2, x)
```



**Reduce [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx$$

$$= \frac{6 \left( \int \operatorname{asech}(cx) dx \right) b d e x + 3 \left( \int \frac{\operatorname{asech}(cx)}{x^2} dx \right) b d^2 x + 3 \left( \int \operatorname{asech}(cx) x^2 dx \right) b e^2 x - 3 a d^2 + 6 a d e x^2 + a e^2 x^4}{3x}$$

input `int((e*x^2+d)^2*(a+b*asech(c*x))/x^2,x)`

output `(6*int(asech(c*x),x)*b*d*e*x + 3*int(asech(c*x)/x**2,x)*b*d**2*x + 3*int(asech(c*x)*x**2,x)*b*e**2*x - 3*a*d**2 + 6*a*d*e*x**2 + a*e**2*x**4)/(3*x)`

**3.102**  $\int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x^4} dx$

Optimal result	817
Mathematica [C] (verified)	818
Rubi [A] (verified)	818
Maple [A] (verified)	821
Fricas [B] (verification not implemented)	821
Sympy [F]	822
Maxima [A] (verification not implemented)	822
Giac [F]	823
Mupad [F(-1)]	823
Reduce [F]	824

**Optimal result**

Integrand size = 21, antiderivative size = 176

$$\int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x^4} dx = \frac{bd^2 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{9x^3} + \frac{2bd(c^2d+9e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{9x} - \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{3x^3} - \frac{2de(a+b\operatorname{sech}^{-1}(cx))}{x} + e^2x(a+b\operatorname{sech}^{-1}(cx)) + \frac{be^2 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \arcsin(cx)}{c}$$

output

```
1/9*b*d^2*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/x^3+2/9*b*d*(c^2*d+9*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/x-1/3*d^2*(a+b*arcsech(c*x))/x^3-2*d*e*(a+b*arcsech(c*x))/x+e^2*x*(a+b*arcsech(c*x))+b*e^2*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arcsin(c*x)/c
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.17 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.85

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx$$

$$= \frac{bcd \sqrt{\frac{1-cx}{1+cx}} (1+cx) (d + 2c^2 dx^2 + 18ex^2) - 3ac(d^2 + 6dex^2 - 3e^2 x^4) - 3bc(d^2 + 6dex^2 - 3e^2 x^4) \operatorname{sech}^{-1}(cx)}{9cx^3}$$

input

```
Integrate[((d + e*x^2)^2*(a + b*ArcSech[c*x]))/x^4,x]
```

output

```
(b*c*d*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(d + 2*c^2*d*x^2 + 18*e*x^2) -
3*a*c*(d^2 + 6*d*e*x^2 - 3*e^2*x^4) - 3*b*c*(d^2 + 6*d*e*x^2 - 3*e^2*x^4)*
ArcSech[c*x] + (9*I)*b*e^2*x^3*Log[(-2*I)*c*x + 2*Sqrt[(1 - c*x)/(1 + c*x)]
]*(1 + c*x)]/(9*c*x^3)
```

**Rubi [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.81, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {6855, 27, 1588, 25, 358, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx$$

$$\downarrow \text{6855}$$

$$b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int -\frac{-3e^2 x^4 + 6dex^2 + d^2}{3x^4 \sqrt{1-c^2 x^2}} dx - \frac{d^2 (a + b \operatorname{sech}^{-1}(cx))}{3x^3} -$$

$$\frac{2de(a + b \operatorname{sech}^{-1}(cx))}{x} + e^2 x (a + b \operatorname{sech}^{-1}(cx))$$

$$\downarrow \text{27}$$

$$\begin{aligned}
& -\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\int\frac{-3e^2x^4+6dex^2+d^2}{x^4\sqrt{1-c^2x^2}}dx-\frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{3x^3}- \\
& \quad \frac{2de(a+b\operatorname{sech}^{-1}(cx))}{x}+e^2x(a+b\operatorname{sech}^{-1}(cx)) \\
& \quad \downarrow 1588 \\
& -\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{1}{3}\int-\frac{2d(dc^2+9e)-9e^2x^2}{x^2\sqrt{1-c^2x^2}}dx-\frac{d^2\sqrt{1-c^2x^2}}{3x^3}\right)- \\
& \quad \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{3x^3}-\frac{2de(a+b\operatorname{sech}^{-1}(cx))}{x}+e^2x(a+b\operatorname{sech}^{-1}(cx)) \\
& \quad \downarrow 25 \\
& -\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\int\frac{2d(dc^2+9e)-9e^2x^2}{x^2\sqrt{1-c^2x^2}}dx-\frac{d^2\sqrt{1-c^2x^2}}{3x^3}\right)- \\
& \quad \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{3x^3}-\frac{2de(a+b\operatorname{sech}^{-1}(cx))}{x}+e^2x(a+b\operatorname{sech}^{-1}(cx)) \\
& \quad \downarrow 358 \\
& -\frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\left(-9e^2\int\frac{1}{\sqrt{1-c^2x^2}}dx-\frac{2d\sqrt{1-c^2x^2}(c^2d+9e)}{x}\right)-\frac{d^2\sqrt{1-c^2x^2}}{3x^3}\right)- \\
& \quad \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{3x^3}-\frac{2de(a+b\operatorname{sech}^{-1}(cx))}{x}+e^2x(a+b\operatorname{sech}^{-1}(cx)) \\
& \quad \downarrow 223 \\
& -\frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{3x^3}-\frac{2de(a+b\operatorname{sech}^{-1}(cx))}{x}+e^2x(a+b\operatorname{sech}^{-1}(cx))- \\
& \quad \frac{1}{3}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\left(-\frac{9e^2\arcsin(cx)}{c}-\frac{2d\sqrt{1-c^2x^2}(c^2d+9e)}{x}\right)-\frac{d^2\sqrt{1-c^2x^2}}{3x^3}\right)
\end{aligned}$$

input `Int[((d + e*x^2)^2*(a + b*ArcSech[c*x]))/x^4,x]`

output `-1/3*(d^2*(a + b*ArcSech[c*x]))/x^3 - (2*d*e*(a + b*ArcSech[c*x]))/x + e^2*x*(a + b*ArcSech[c*x]) - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/3*(d^2*Sqrt[1 - c^2*x^2])/x^3 + ((-2*d*(c^2*d + 9*e)*Sqrt[1 - c^2*x^2])/x - (9*e^2*ArcSin[c*x])/c)/3)/3`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 358 `Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[c*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*e*(m + 1))), x] + Simp[d/e^2 Int[(e*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d, 0] && EqQ[Simplify[m + 2*p + 3], 0] && NeQ[m, -1]`
- rule 1588 `Int[((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := With[{qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, f*x, x], R = PolynomialRemainder[(a + b*x^2 + c*x^4)^p, f*x, x]}, Simp[R*(f*x)^(m + 1)*((d + e*x^2)^(q + 1)/(d*f*(m + 1))), x] + Simp[1/(d*f^2*(m + 1)) Int[(f*x)^(m + 2)*(d + e*x^2)^q*ExpandToSum[d*f*(m + 1)*(Qx/x) - e*R*(m + 2*q + 3), x], x], x]] /; FreeQ[{a, b, c, d, e, f, q}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && LtQ[m, -1]`
- rule 6855 `Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))`

### Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.14

method	result
parts	$a\left(e^2x - \frac{d^2}{3x^3} - \frac{2de}{x}\right) + bc^3\left(\frac{\operatorname{arcsech}(cx)e^2x}{c^3} - \frac{\operatorname{arcsech}(cx)d^2}{3c^3x^3} - \frac{2\operatorname{arcsech}(cx)de}{c^3x} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}}{c^3}\right)$
derivativedivides	$c^3\left(\frac{a\left(e^2cx - \frac{2cde}{x} - \frac{cd^2}{3x^3}\right)}{c^4} + \frac{b\left(\operatorname{arcsech}(cx)e^2cx - \frac{2\operatorname{arcsech}(cx)cde}{x} - \frac{\operatorname{arcsech}(cx)cd^2}{3x^3} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}}{c^3}\right)}{c^4}\right)$
default	$c^3\left(\frac{a\left(e^2cx - \frac{2cde}{x} - \frac{cd^2}{3x^3}\right)}{c^4} + \frac{b\left(\operatorname{arcsech}(cx)e^2cx - \frac{2\operatorname{arcsech}(cx)cde}{x} - \frac{\operatorname{arcsech}(cx)cd^2}{3x^3} + \frac{\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}}{c^3}\right)}{c^4}\right)$

input `int((e*x^2+d)^2*(a+b*arcsech(c*x))/x^4,x,method=_RETURNVERBOSE)`

output `a*(e^2*x-1/3*d^2/x^3-2*d*e/x)+b*c^3*(1/c^3*arcsech(c*x)*e^2*x-1/3*arcsech(c*x)*d^2/c^3/x^3-2/c^3*arcsech(c*x)*d*e/x+1/9/c^6*(-(c*x-1)/c/x)^(1/2)/x^2*((c*x+1)/c/x)^(1/2)*(2*(-c^2*x^2+1)^(1/2)*c^6*d^2*x^2+(-c^2*x^2+1)^(1/2)*c^4*d^2+18*(-c^2*x^2+1)^(1/2)*c^4*d*e*x^2+9*arcsin(c*x)*e^2*c^3*x^3)/(-c^2*x^2+1)^(1/2)`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 267 vs. 2(106) = 212.

Time = 0.13 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.52

$$\int \frac{(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx))}{x^4} dx$$

$$= \frac{9ace^2x^4 - 18be^2x^3 \arctan\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{cx}\right) - 18acdex^2 + 3(bcd^2 + 6bcde - 3bce^2)x^3 \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}-1}{x}\right)}{c^4}$$

input `integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x^4,x, algorithm="fricas")`

output

```
1/9*(9*a*c*e^2*x^4 - 18*b*e^2*x^3*arctan((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2
)) - 1)/(c*x)) - 18*a*c*d*e*x^2 + 3*(b*c*d^2 + 6*b*c*d*e - 3*b*c*e^2)*x^3*
log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) - 3*a*c*d^2 + 3*(3*b*c*e^2
*x^4 - 6*b*c*d*e*x^2 - b*c*d^2 + (b*c*d^2 + 6*b*c*d*e - 3*b*c*e^2)*x^3)*lo
g((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + (b*c^2*d^2*x + 2*(b*c^
4*d^2 + 9*b*c^2*d*e)*x^3)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c*x^3)
```

**Sympy [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{(a + b \operatorname{sech}(cx)) (d + ex^2)^2}{x^4} dx$$

input

```
integrate((e*x**2+d)**2*(a+b*asech(c*x))/x**4,x)
```

output

```
Integral((a + b*asech(c*x))*(d + e*x**2)**2/x**4, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.76

$$\begin{aligned} & \int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx \\ &= 2 \left( c \sqrt{\frac{1}{c^2 x^2} - 1} - \frac{\operatorname{arsech}(cx)}{x} \right) bde + ae^2 x \\ &+ \frac{1}{9} bd^2 \left( \frac{c^4 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{3}{2}} + 3c^4 \sqrt{\frac{1}{c^2 x^2} - 1}}{c} - \frac{3 \operatorname{arsech}(cx)}{x^3} \right) \\ &+ \frac{\left( cx \operatorname{arsech}(cx) - \arctan \left( \sqrt{\frac{1}{c^2 x^2} - 1} \right) \right) be^2}{c} - \frac{2ade}{x} - \frac{ad^2}{3x^3} \end{aligned}$$

input

```
integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x^4,x, algorithm="maxima")
```

output

```
2*(c*sqrt(1/(c^2*x^2) - 1) - arcsech(c*x)/x)*b*d*e + a*e^2*x + 1/9*b*d^2*(
(c^4*(1/(c^2*x^2) - 1)^(3/2) + 3*c^4*sqrt(1/(c^2*x^2) - 1))/c - 3*arcsech(
c*x)/x^3) + (c*x*arcsech(c*x) - arctan(sqrt(1/(c^2*x^2) - 1)))*b*e^2/c - 2
*a*d*e/x - 1/3*a*d^2/x^3
```

**Giac [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{ar} \operatorname{sech}(cx) + a)}{x^4} dx$$

input

```
integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x^4,x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)^2*(b*arcsech(c*x) + a)/x^4, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{(ex^2 + d)^2 (a + b \operatorname{acosh}(\frac{1}{cx}))}{x^4} dx$$

input

```
int(((d + e*x^2)^2*(a + b*acosh(1/(c*x))))/x^4,x)
```

output

```
int(((d + e*x^2)^2*(a + b*acosh(1/(c*x))))/x^4, x)
```



**Reduce [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx$$

$$= \frac{3 \left( \int \operatorname{asech}(cx) dx \right) b e^2 x^3 + 3 \left( \int \frac{\operatorname{asech}(cx)}{x^4} dx \right) b d^2 x^3 + 6 \left( \int \frac{\operatorname{asech}(cx)}{x^2} dx \right) b d e x^3 - a d^2 - 6 a d e x^2 + 3 a e^2 x^4}{3x^3}$$

input `int((e*x^2+d)^2*(a+b*asech(c*x))/x^4,x)`

output `(3*int(asech(c*x),x)*b*e**2*x**3 + 3*int(asech(c*x)/x**4,x)*b*d**2*x**3 + 6*int(asech(c*x)/x**2,x)*b*d*e*x**3 - a*d**2 - 6*a*d*e*x**2 + 3*a*e**2*x**4)/(3*x**3)`

**3.103**  $\int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x^6} dx$

Optimal result	825
Mathematica [A] (verified)	826
Rubi [A] (verified)	826
Maple [A] (verified)	829
Fricas [A] (verification not implemented)	829
Sympy [F]	830
Maxima [A] (verification not implemented)	830
Giac [F]	831
Mupad [F(-1)]	831
Reduce [F]	832

**Optimal result**

Integrand size = 21, antiderivative size = 213

$$\int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x^6} dx$$

$$= \frac{bd^2 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{25x^5} + \frac{2bd(6c^2d+25e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{225x^3}$$

$$+ \frac{b(24c^4d^2+100c^2de+225e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{225x}$$

$$- \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{5x^5} - \frac{2de(a+b\operatorname{sech}^{-1}(cx))}{3x^3} - \frac{e^2(a+b\operatorname{sech}^{-1}(cx))}{x}$$

output

```
1/25*b*d^2*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/x^5+2/225*b*
d*(6*c^2*d+25*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/x^3+1/
225*b*(24*c^4*d^2+100*c^2*d*e+225*e^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c
^2*x^2+1)^(1/2)/x-1/5*d^2*(a+b*arcsech(c*x))/x^5-2/3*d*e*(a+b*arcsech(c*x)
)/x^3-e^2*(a+b*arcsech(c*x))/x
```

**Mathematica [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.63

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx$$

$$= \frac{-15a(3d^2 + 10dex^2 + 15e^2x^4) + b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(225e^2x^4 + 50dex^2(1+2c^2x^2) + 3d^2(3+4c^2x^2+8c^4x^4)) - 15b(3d^2 + 10dex^2 + 15e^2x^4)\operatorname{ArcSech}[cx]}{225x^5}$$

input

```
Integrate[((d + e*x^2)^2*(a + b*ArcSech[c*x]))/x^6,x]
```

output

```
(-15*a*(3*d^2 + 10*d*e*x^2 + 15*e^2*x^4) + b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(225*e^2*x^4 + 50*d*e*x^2*(1 + 2*c^2*x^2) + 3*d^2*(3 + 4*c^2*x^2 + 8*c^4*x^4)) - 15*b*(3*d^2 + 10*d*e*x^2 + 15*e^2*x^4)*ArcSech[c*x])/(225*x^5)
```

**Rubi [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 181, normalized size of antiderivative = 0.85, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {6855, 27, 1588, 25, 359, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx$$

$$\downarrow 6855$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int -\frac{15e^2x^4 + 10dex^2 + 3d^2}{15x^6\sqrt{1-c^2x^2}} dx - \frac{d^2(a + b \operatorname{sech}^{-1}(cx))}{5x^5} -$$

$$\frac{2de(a + b \operatorname{sech}^{-1}(cx))}{3x^3} - \frac{e^2(a + b \operatorname{sech}^{-1}(cx))}{x}$$

$$\downarrow 27$$

$$\begin{aligned}
& -\frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\int\frac{15e^2x^4+10dex^2+3d^2}{x^6\sqrt{1-c^2x^2}}dx-\frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{5x^5}- \\
& \quad \frac{2de(a+b\operatorname{sech}^{-1}(cx))}{3x^3}-\frac{e^2(a+b\operatorname{sech}^{-1}(cx))}{x} \\
& \quad \downarrow 1588 \\
& -\frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{1}{5}\int-\frac{75e^2x^2+2d(6dc^2+25e)}{x^4\sqrt{1-c^2x^2}}dx-\frac{3d^2\sqrt{1-c^2x^2}}{5x^5}\right)- \\
& \quad \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{5x^5}-\frac{2de(a+b\operatorname{sech}^{-1}(cx))}{3x^3}-\frac{e^2(a+b\operatorname{sech}^{-1}(cx))}{x} \\
& \quad \downarrow 25 \\
& -\frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\int\frac{75e^2x^2+2d(6dc^2+25e)}{x^4\sqrt{1-c^2x^2}}dx-\frac{3d^2\sqrt{1-c^2x^2}}{5x^5}\right)- \\
& \quad \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{5x^5}-\frac{2de(a+b\operatorname{sech}^{-1}(cx))}{3x^3}-\frac{e^2(a+b\operatorname{sech}^{-1}(cx))}{x} \\
& \quad \downarrow 359 \\
& -\frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\left(\frac{1}{3}(24c^4d^2+100c^2de+225e^2)\int\frac{1}{x^2\sqrt{1-c^2x^2}}dx-\frac{2d\sqrt{1-c^2x^2}(6c^2d+25e)}{3x^3}\right)\right)- \\
& \quad \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{5x^5}-\frac{2de(a+b\operatorname{sech}^{-1}(cx))}{3x^3}-\frac{e^2(a+b\operatorname{sech}^{-1}(cx))}{x} \\
& \quad \downarrow 242 \\
& -\frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{5x^5}-\frac{2de(a+b\operatorname{sech}^{-1}(cx))}{3x^3}-\frac{e^2(a+b\operatorname{sech}^{-1}(cx))}{x}- \\
& \frac{1}{15}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\left(-\frac{2d\sqrt{1-c^2x^2}(6c^2d+25e)}{3x^3}-\frac{\sqrt{1-c^2x^2}(24c^4d^2+100c^2de+225e^2)}{3x}\right)\right)-\frac{3d^2\sqrt{1-c^2x^2}}{5x^5}
\end{aligned}$$

input `Int[((d + e*x^2)^2*(a + b*ArcSech[c*x]))/x^6,x]`

output `-1/15*(b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-3*d^2*Sqrt[1 - c^2*x^2]))/(5*x^5) + ((-2*d*(6*c^2*d + 25*e)*Sqrt[1 - c^2*x^2])/(3*x^3) - ((24*c^4*d^2 + 100*c^2*d*e + 225*e^2)*Sqrt[1 - c^2*x^2])/(3*x))/5) - (d^2*(a + b*ArcSech[c*x]))/(5*x^5) - (2*d*e*(a + b*ArcSech[c*x]))/(3*x^3) - (e^2*(a + b*ArcSech[c*x]))/x`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 242 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`
- rule 359 `Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[c*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*e*(m + 1))), x] + Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(a*e^2*(m + 1)) Int[(e*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && !ILtQ[p, -1]`
- rule 1588 `Int[((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, f*x, x], R = PolynomialRemainder[(a + b*x^2 + c*x^4)^p, f*x, x]}, Simp[R*(f*x)^(m + 1)*((d + e*x^2)^(q + 1)/(d*f*(m + 1))), x] + Simp[1/(d*f^2*(m + 1)) Int[(f*x)^(m + 2)*(d + e*x^2)^q*ExpandToSum[d*f*(m + 1)*(Qx/x) - e*R*(m + 2*q + 3), x], x], x] /; FreeQ[{a, b, c, d, e, f, q}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && LtQ[m, -1]`
- rule 6855 `Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))`

### Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.83

method	result
parts	$a\left(-\frac{2de}{3x^3} - \frac{e^2}{x} - \frac{d^2}{5x^5}\right) + b c^5 \left(-\frac{2 \operatorname{arcsech}(cx)de}{3c^5x^3} - \frac{\operatorname{arcsech}(cx)e^2}{c^5x} - \frac{\operatorname{arcsech}(cx)d^2}{5c^5x^5} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}}{c^5x^5}\right)$
derivativedivides	$c^5 \left( \frac{a\left(-\frac{d^2}{5cx^5} - \frac{2de}{3cx^3} - \frac{e^2}{cx}\right)}{c^4} + \frac{b\left(-\frac{\operatorname{arcsech}(cx)d^2}{5cx^5} - \frac{2 \operatorname{arcsech}(cx)de}{3cx^3} - \frac{\operatorname{arcsech}(cx)e^2}{cx} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} (24c^8d^2x^4 + 100c^6d^2x^2 + 225c^4e^2x^4 + 50c^4de^2x^4 + 9c^4d^2)}{c^4}}{c^4} \right)$
default	$c^5 \left( \frac{a\left(-\frac{d^2}{5cx^5} - \frac{2de}{3cx^3} - \frac{e^2}{cx}\right)}{c^4} + \frac{b\left(-\frac{\operatorname{arcsech}(cx)d^2}{5cx^5} - \frac{2 \operatorname{arcsech}(cx)de}{3cx^3} - \frac{\operatorname{arcsech}(cx)e^2}{cx} + \frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} (24c^8d^2x^4 + 100c^6d^2x^2 + 225c^4e^2x^4 + 50c^4de^2x^4 + 9c^4d^2)}{c^4}}{c^4} \right)$

input `int((e*x^2+d)^2*(a+b*arcsech(c*x))/x^6,x,method=_RETURNVERBOSE)`

output `a*(-2/3*d*e/x^3-e^2/x-1/5*d^2/x^5)+b*c^5*(-2/3/c^5*arcsech(c*x)*d*e/x^3-1/c^5*arcsech(c*x)*e^2/x-1/5*arcsech(c*x)*d^2/c^5/x^5+1/225/c^8*(-(c*x-1)/c/x)^(1/2)/x^4*((c*x+1)/c/x)^(1/2)*(24*c^8*d^2*x^4+100*c^6*d*e*x^4+12*c^6*d^2*x^2+225*c^4*e^2*x^4+50*c^4*d*e*x^2+9*c^4*d^2))`

### Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.78

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx = \frac{225 a e^2 x^4 + 150 a d e x^2 + 45 a d^2 + 15 (15 b e^2 x^4 + 10 b d e x^2 + 3 b d^2) \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2} + 1}}{cx}\right) - ((24 b c^5 d^2 + \dots)}{225 x^5}$$

input `integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x^6,x, algorithm="fricas")`

output

```
-1/225*(225*a*e^2*x^4 + 150*a*d*e*x^2 + 45*a*d^2 + 15*(15*b*e^2*x^4 + 10*b*d*e*x^2 + 3*b*d^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - ((24*b*c^5*d^2 + 100*b*c^3*d*e + 225*b*c*e^2)*x^5 + 9*b*c*d^2*x + 2*(6*b*c^3*d^2 + 25*b*c*d*e)*x^3)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2))/x^5
```

**Sympy [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx = \int \frac{(a + b \operatorname{asech}(cx)) (d + ex^2)^2}{x^6} dx$$

input

```
integrate((e*x**2+d)**2*(a+b*asech(c*x))/x**6,x)
```

output

```
Integral((a + b*asech(c*x))*(d + e*x**2)**2/x**6, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.82

$$\begin{aligned} & \int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx \\ &= \left( c \sqrt{\frac{1}{c^2 x^2} - 1} - \frac{\operatorname{arsech}(cx)}{x} \right) b e^2 \\ &+ \frac{1}{75} b d^2 \left( \frac{3 c^6 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{5}{2}} + 10 c^6 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{3}{2}} + 15 c^6 \sqrt{\frac{1}{c^2 x^2} - 1}}{c} - \frac{15 \operatorname{arsech}(cx)}{x^5} \right) \\ &+ \frac{2}{9} b d e \left( \frac{c^4 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{3}{2}} + 3 c^4 \sqrt{\frac{1}{c^2 x^2} - 1}}{c} - \frac{3 \operatorname{arsech}(cx)}{x^3} \right) - \frac{a e^2}{x} - \frac{2 a d e}{3 x^3} - \frac{a d^2}{5 x^5} \end{aligned}$$

input

```
integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x^6,x, algorithm="maxima")
```

output

```
(c*sqrt(1/(c^2*x^2) - 1) - arcsech(c*x)/x)*b*e^2 + 1/75*b*d^2*((3*c^6*(1/(c^2*x^2) - 1)^(5/2) + 10*c^6*(1/(c^2*x^2) - 1)^(3/2) + 15*c^6*sqrt(1/(c^2*x^2) - 1))/c - 15*arcsech(c*x)/x^5) + 2/9*b*d*e*((c^4*(1/(c^2*x^2) - 1)^(3/2) + 3*c^4*sqrt(1/(c^2*x^2) - 1))/c - 3*arcsech(c*x)/x^3) - a*e^2/x - 2/3*a*d*e/x^3 - 1/5*a*d^2/x^5
```

**Giac [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{arsech}(cx) + a)}{x^6} dx$$

input

```
integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x^6,x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)^2*(b*arcsech(c*x) + a)/x^6, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx = \int \frac{(ex^2 + d)^2 (a + b \operatorname{acosh}(\frac{1}{cx}))}{x^6} dx$$

input

```
int(((d + e*x^2)^2*(a + b*acosh(1/(c*x))))/x^6,x)
```

output

```
int(((d + e*x^2)^2*(a + b*acosh(1/(c*x))))/x^6, x)
```



**Reduce [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx$$

$$= \frac{15 \left( \int \frac{\operatorname{asech}(cx)}{x^6} dx \right) b d^2 x^5 + 30 \left( \int \frac{\operatorname{asech}(cx)}{x^4} dx \right) b d e x^5 + 15 \left( \int \frac{\operatorname{asech}(cx)}{x^2} dx \right) b e^2 x^5 - 3a d^2 - 10ade x^2 - 15a}{15x^5}$$

input `int((e*x^2+d)^2*(a+b*asech(c*x))/x^6,x)`

output `(15*int(asech(c*x)/x**6,x)*b*d**2*x**5 + 30*int(asech(c*x)/x**4,x)*b*d*e*x**5 + 15*int(asech(c*x)/x**2,x)*b*e**2*x**5 - 3*a*d**2 - 10*a*d*e*x**2 - 15*a*e**2*x**4)/(15*x**5)`

**3.104** 
$$\int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x^8} dx$$

Optimal result . . . . .	833
Mathematica [A] (verified) . . . . .	834
Rubi [A] (verified) . . . . .	834
Maple [A] (verified) . . . . .	837
Fricas [A] (verification not implemented) . . . . .	838
Sympy [F] . . . . .	838
Maxima [A] (verification not implemented) . . . . .	839
Giac [F] . . . . .	839
Mupad [F(-1)] . . . . .	840
Reduce [F] . . . . .	840

**Optimal result**

Integrand size = 21, antiderivative size = 281

$$\begin{aligned} & \int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x^8} dx \\ &= \frac{bd^2 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{49x^7} + \frac{2bd(15c^2d+49e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{1225x^5} \\ &+ \frac{b(360c^4d^2+1176c^2de+1225e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{11025x^3} \\ &+ \frac{2bc^2(360c^4d^2+1176c^2de+1225e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{11025x} \\ &- \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{7x^7} - \frac{2de(a+b\operatorname{sech}^{-1}(cx))}{5x^5} - \frac{e^2(a+b\operatorname{sech}^{-1}(cx))}{3x^3} \end{aligned}$$

output

```
1/49*b*d^2*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/x^7+2/1225*b
*d*(15*c^2*d+49*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/x^5+
1/11025*b*(360*c^4*d^2+1176*c^2*d*e+1225*e^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1
/2)*(-c^2*x^2+1)^(1/2)/x^3+2/11025*b*c^2*(360*c^4*d^2+1176*c^2*d*e+1225*e^
2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/x-1/7*d^2*(a+b*arcse
ch(c*x))/x^7-2/5*d*e*(a+b*arcsech(c*x))/x^5-1/3*e^2*(a+b*arcsech(c*x))/x^3
```

**Mathematica [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.57

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^8} dx$$

$$= \frac{-105a(15d^2 + 42dex^2 + 35e^2x^4) + b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(1225e^2x^4(1+2c^2x^2) + 294dex^2(3+4c^2x^2+8c^4x^4))}{11025x^7}$$

input

```
Integrate[((d + e*x^2)^2*(a + b*ArcSech[c*x]))/x^8,x]
```

output

```
(-105*a*(15*d^2 + 42*d*e*x^2 + 35*e^2*x^4) + b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(1225*e^2*x^4*(1 + 2*c^2*x^2) + 294*d*e*x^2*(3 + 4*c^2*x^2 + 8*c^4*x^4) + 45*d^2*(5 + 6*c^2*x^2 + 8*c^4*x^4 + 16*c^6*x^6)) - 105*b*(15*d^2 + 42*d*e*x^2 + 35*e^2*x^4)*ArcSech[c*x])/(11025*x^7)
```

**Rubi [A] (verified)**Time = 0.45 (sec) , antiderivative size = 212, normalized size of antiderivative = 0.75, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6855, 27, 1588, 25, 359, 245, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^8} dx$$

$$\downarrow 6855$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int -\frac{35e^2x^4 + 42dex^2 + 15d^2}{105x^8\sqrt{1-c^2x^2}} dx - \frac{d^2(a + b \operatorname{sech}^{-1}(cx))}{7x^7} -$$

$$\frac{2de(a + b \operatorname{sech}^{-1}(cx))}{5x^5} - \frac{e^2(a + b \operatorname{sech}^{-1}(cx))}{3x^3}$$

$$\downarrow 27$$

$$\begin{aligned}
& -\frac{1}{105}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\int\frac{35e^2x^4+42dex^2+15d^2}{x^8\sqrt{1-c^2x^2}}dx-\frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{7x^7}- \\
& \quad \frac{2de(a+b\operatorname{sech}^{-1}(cx))}{5x^5}-\frac{e^2(a+b\operatorname{sech}^{-1}(cx))}{3x^3} \\
& \quad \downarrow 1588 \\
& -\frac{1}{105}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{1}{7}\int-\frac{245e^2x^2+6d(15dc^2+49e)}{x^6\sqrt{1-c^2x^2}}dx-\frac{15d^2\sqrt{1-c^2x^2}}{7x^7}\right)- \\
& \quad \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{7x^7}-\frac{2de(a+b\operatorname{sech}^{-1}(cx))}{5x^5}-\frac{e^2(a+b\operatorname{sech}^{-1}(cx))}{3x^3} \\
& \quad \downarrow 25 \\
& -\frac{1}{105}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{7}\int\frac{245e^2x^2+6d(15dc^2+49e)}{x^6\sqrt{1-c^2x^2}}dx-\frac{15d^2\sqrt{1-c^2x^2}}{7x^7}\right)- \\
& \quad \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{7x^7}-\frac{2de(a+b\operatorname{sech}^{-1}(cx))}{5x^5}-\frac{e^2(a+b\operatorname{sech}^{-1}(cx))}{3x^3} \\
& \quad \downarrow 359 \\
& -\frac{1}{105}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{7}\left(\frac{1}{5}(360c^4d^2+1176c^2de+1225e^2)\int\frac{1}{x^4\sqrt{1-c^2x^2}}dx-\frac{6d\sqrt{1-c^2x^2}(15c^2d+49e)}{5x^5}\right)\right)- \\
& \quad \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{7x^7}-\frac{2de(a+b\operatorname{sech}^{-1}(cx))}{5x^5}-\frac{e^2(a+b\operatorname{sech}^{-1}(cx))}{3x^3} \\
& \quad \downarrow 245 \\
& -\frac{1}{105}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{7}\left(\frac{1}{5}(360c^4d^2+1176c^2de+1225e^2)\left(\frac{2}{3}c^2\int\frac{1}{x^2\sqrt{1-c^2x^2}}dx-\frac{\sqrt{1-c^2x^2}}{3x^3}\right)\right)\right)-\frac{6d\sqrt{1-c^2x^2}}{5x^5} \\
& \quad \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{7x^7}-\frac{2de(a+b\operatorname{sech}^{-1}(cx))}{5x^5}-\frac{e^2(a+b\operatorname{sech}^{-1}(cx))}{3x^3} \\
& \quad \downarrow 242 \\
& -\frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{7x^7}-\frac{2de(a+b\operatorname{sech}^{-1}(cx))}{5x^5}-\frac{e^2(a+b\operatorname{sech}^{-1}(cx))}{3x^3}- \\
& \quad \frac{1}{105}b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{7}\left(\frac{1}{5}\left(-\frac{2c^2\sqrt{1-c^2x^2}}{3x}-\frac{\sqrt{1-c^2x^2}}{3x^3}\right)(360c^4d^2+1176c^2de+1225e^2)-\frac{6d\sqrt{1-c^2x^2}}{5x^5}\right)\right)
\end{aligned}$$

input

```
Int[((d + e*x^2)^2*(a + b*ArcSech[c*x]))/x^8,x]
```

output

$$\begin{aligned} & -1/105*(b*\text{Sqrt}[(1 + c*x)^{-1}]*\text{Sqrt}[1 + c*x]*((-15*d^2*\text{Sqrt}[1 - c^2*x^2])/ \\ & (7*x^7) + ((-6*d*(15*c^2*d + 49*e)*\text{Sqrt}[1 - c^2*x^2])/(5*x^5) + ((360*c^4* \\ & d^2 + 1176*c^2*d*e + 1225*e^2)*(-1/3*\text{Sqrt}[1 - c^2*x^2]/x^3 - (2*c^2*\text{Sqrt}[1 \\ & - c^2*x^2])/(3*x)))/5)/7) - (d^2*(a + b*\text{ArcSech}[c*x]))/(7*x^7) - (2*d*e* \\ & (a + b*\text{ArcSech}[c*x]))/(5*x^5) - (e^2*(a + b*\text{ArcSech}[c*x]))/(3*x^3) \end{aligned}$$

### Defintions of rubi rules used

rule 25

$$\text{Int}[-(\text{Fx}_), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, x], x]$$

rule 27

$$\text{Int}[(a_)*(\text{Fx}_), x\_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[\text{Fx}, (b_)*(\text{Gx}_) /; \text{FreeQ}[b, x]]$$

rule 242

$$\text{Int}[((c_)*(x_))^{(m_)}*((a_) + (b_)*(x_)^2)^{(p_)}, x\_Symbol] \rightarrow \text{Simp}[(c*x)^{(m+1)}*((a + b*x^2)^{(p+1)})/(a*c*(m+1)), x] /; \text{FreeQ}[\{a, b, c, m, p\}, x] \ \&\& \ \text{EqQ}[m + 2*p + 3, 0] \ \&\& \ \text{NeQ}[m, -1]$$

rule 245

$$\begin{aligned} & \text{Int}[(x_)^{(m_)}*((a_) + (b_)*(x_)^2)^{(p_)}, x\_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + \\ & b*x^2)^{(p+1)})/(a*(m+1)), x] - \text{Simp}[b*((m + 2*(p + 1) + 1)/(a*(m + 1))) \\ & \quad \text{Int}[x^{(m+2)}*(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{ILtQ}[\text{Simplify}[(m + 1)/2 + p + 1], 0] \ \&\& \ \text{NeQ}[m, -1] \end{aligned}$$

rule 359

$$\begin{aligned} & \text{Int}[((e_)*(x_))^{(m_)}*((a_) + (b_)*(x_)^2)^{(p_)}*((c_) + (d_)*(x_)^2), x \\ & \_Symbol] \rightarrow \text{Simp}[c*(e*x)^{(m+1)}*((a + b*x^2)^{(p+1)})/(a*e*(m+1)), x] + \\ & \text{Simp}[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(a*e^2*(m + 1)) \quad \text{Int}[(e*x)^{(m+2)}* \\ & (a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, p\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \\ & \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ !\text{ILtQ}[p, -1] \end{aligned}$$

rule 1588

```
Int[((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, f*x, x], R = PolynomialRemainder[(a + b*x^2 + c*x^4)^p, f*x, x]}, Simp[R*(f*x)^(m + 1)*((d + e*x^2)^(q + 1)/(d*f*(m + 1))), x] + Simp[1/(d*f^2*(m + 1)) Int[(f*x)^(m + 2)*(d + e*x^2)^q*ExpandToSum[d*f*(m + 1)*(Qx/x) - e*R*(m + 2*q + 3), x], x], x] /; FreeQ[{a, b, c, d, e, f, q}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && LtQ[m, -1]
```

rule 6855

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_))*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

### Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 209, normalized size of antiderivative = 0.74

method	result
parts	$a \left( -\frac{d^2}{7x^7} - \frac{e^2}{3x^3} - \frac{2de}{5x^5} \right) + b c^7 \left( -\frac{\operatorname{arcsech}(cx)d^2}{7c^7x^7} - \frac{\operatorname{arcsech}(cx)e^2}{3c^7x^3} - \frac{2 \operatorname{arcsech}(cx)de}{5c^7x^5} + \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \right)$
derivativedivides	$c^7 \left( \frac{a \left( -\frac{e^2}{3c^3x^3} - \frac{d^2}{7c^3x^7} - \frac{2de}{5c^3x^5} \right)}{c^4} + \frac{b \left( -\frac{\operatorname{arcsech}(cx)e^2}{3c^3x^3} - \frac{\operatorname{arcsech}(cx)d^2}{7c^3x^7} - \frac{2 \operatorname{arcsech}(cx)de}{5c^3x^5} + \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} (720c^{10}d^2x^6) \right)}{c^4} \right)$
default	$c^7 \left( \frac{a \left( -\frac{e^2}{3c^3x^3} - \frac{d^2}{7c^3x^7} - \frac{2de}{5c^3x^5} \right)}{c^4} + \frac{b \left( -\frac{\operatorname{arcsech}(cx)e^2}{3c^3x^3} - \frac{\operatorname{arcsech}(cx)d^2}{7c^3x^7} - \frac{2 \operatorname{arcsech}(cx)de}{5c^3x^5} + \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} (720c^{10}d^2x^6) \right)}{c^4} \right)$

input

```
int((e*x^2+d)^2*(a+b*arcsech(c*x))/x^8,x,method=_RETURNVERBOSE)
```

output

```
a*(-1/7*d^2/x^7-1/3*e^2/x^3-2/5*d*e/x^5)+b*c^7*(-1/7*arcsech(c*x)*d^2/c^7/
x^7-1/3/c^7*arcsech(c*x)*e^2/x^3-2/5/c^7*arcsech(c*x)*d*e/x^5+1/11025/c^10
*(-(c*x-1)/c/x)^(1/2)/x^6*((c*x+1)/c/x)^(1/2)*(720*c^10*d^2*x^6+2352*c^8*d
*e*x^6+360*c^8*d^2*x^4+2450*c^6*e^2*x^6+1176*c^6*d*e*x^4+270*c^6*d^2*x^2+1
225*c^4*e^2*x^4+882*c^4*d*e*x^2+225*c^4*d^2))
```

### Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 199, normalized size of antiderivative = 0.71

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^8} dx =$$

$$\frac{3675 ae^2 x^4 + 4410 adex^2 + 1575 ad^2 + 105 (35 be^2 x^4 + 42 bdex^2 + 15 bd^2) \log\left(\frac{cx\sqrt{-\frac{e^2 x^2 - 1}{c^2 x^2}} + 1}{cx}\right) - (2(3$$

input

```
integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x^8,x, algorithm="fricas")
```

output

```
-1/11025*(3675*a*e^2*x^4 + 4410*a*d*e*x^2 + 1575*a*d^2 + 105*(35*b*e^2*x^4
+ 42*b*d*e*x^2 + 15*b*d^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(
c*x)) - (2*(360*b*c^7*d^2 + 1176*b*c^5*d*e + 1225*b*c^3*e^2)*x^7 + (360*b*
c^5*d^2 + 1176*b*c^3*d*e + 1225*b*c*e^2)*x^5 + 225*b*c*d^2*x + 18*(15*b*c^
3*d^2 + 49*b*c*d*e)*x^3)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/x^7
```

### Sympy [F]

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{asech}(cx))}{x^8} dx = \int \frac{(a + b \operatorname{asech}(cx)) (d + ex^2)^2}{x^8} dx$$

input

```
integrate((e*x**2+d)**2*(a+b*asech(c*x))/x**8,x)
```

output

```
Integral((a + b*asech(c*x))*(d + e*x**2)**2/x**8, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 232, normalized size of antiderivative = 0.83

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^8} dx$$

$$= \frac{1}{245} bd^2 \left( \frac{5c^8 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{7}{2}} + 21c^8 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{5}{2}} + 35c^8 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{3}{2}} + 35c^8 \sqrt{\frac{1}{c^2x^2} - 1}}{c} - \frac{35 \operatorname{arsech}(cx)}{x^7} \right)$$

$$+ \frac{2}{75} bde \left( \frac{3c^6 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{5}{2}} + 10c^6 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{3}{2}} + 15c^6 \sqrt{\frac{1}{c^2x^2} - 1}}{c} - \frac{15 \operatorname{arsech}(cx)}{x^5} \right)$$

$$+ \frac{1}{9} be^2 \left( \frac{c^4 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{3}{2}} + 3c^4 \sqrt{\frac{1}{c^2x^2} - 1}}{c} - \frac{3 \operatorname{arsech}(cx)}{x^3} \right) - \frac{ae^2}{3x^3} - \frac{2ade}{5x^5} - \frac{ad^2}{7x^7}$$

input `integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x^8,x, algorithm="maxima")`

output `1/245*b*d^2*((5*c^8*(1/(c^2*x^2) - 1)^(7/2) + 21*c^8*(1/(c^2*x^2) - 1)^(5/2) + 35*c^8*(1/(c^2*x^2) - 1)^(3/2) + 35*c^8*sqrt(1/(c^2*x^2) - 1))/c - 35*arcsech(c*x)/x^7) + 2/75*b*d*e*((3*c^6*(1/(c^2*x^2) - 1)^(5/2) + 10*c^6*(1/(c^2*x^2) - 1)^(3/2) + 15*c^6*sqrt(1/(c^2*x^2) - 1))/c - 15*arcsech(c*x)/x^5) + 1/9*b*e^2*((c^4*(1/(c^2*x^2) - 1)^(3/2) + 3*c^4*sqrt(1/(c^2*x^2) - 1))/c - 3*arcsech(c*x)/x^3) - 1/3*a*e^2/x^3 - 2/5*a*d*e/x^5 - 1/7*a*d^2/x^7`

**Giac [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^8} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{arsech}(cx) + a)}{x^8} dx$$

input `integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x^8,x, algorithm="giac")`

output `integrate((e*x^2 + d)^2*(b*arcsech(c*x) + a)/x^8, x)`



**Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^8} dx = \int \frac{(ex^2 + d)^2 (a + b \operatorname{acosh}(\frac{1}{cx}))}{x^8} dx$$

input `int(((d + e*x^2)^2*(a + b*acosh(1/(c*x))))/x^8,x)`

output `int(((d + e*x^2)^2*(a + b*acosh(1/(c*x))))/x^8, x)`

**Reduce [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^8} dx$$

$$= \frac{105 \left( \int \frac{\operatorname{asech}(cx)}{x^8} dx \right) b d^2 x^7 + 210 \left( \int \frac{\operatorname{asech}(cx)}{x^6} dx \right) b d e x^7 + 105 \left( \int \frac{\operatorname{asech}(cx)}{x^4} dx \right) b e^2 x^7 - 15 a d^2 - 42 a d e x^2 - 35 a e^2 x^4}{105 x^7}$$

input `int((e*x^2+d)^2*(a+b*asech(c*x))/x^8,x)`

output `(105*int(asech(c*x)/x**8,x)*b*d**2*x**7 + 210*int(asech(c*x)/x**6,x)*b*d*e*x**7 + 105*int(asech(c*x)/x**4,x)*b*e**2*x**7 - 15*a*d**2 - 42*a*d*e*x**2 - 35*a*e**2*x**4)/(105*x**7)`

### 3.105 $\int x^3(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	841
Mathematica [A] (verified)	842
Rubi [A] (verified)	842
Maple [A] (verified)	845
Fricas [A] (verification not implemented)	845
Sympy [A] (verification not implemented)	846
Maxima [A] (verification not implemented)	847
Giac [F]	847
Mupad [F(-1)]	848
Reduce [F]	848

#### Optimal result

Integrand size = 21, antiderivative size = 278

$$\int x^3(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= -\frac{b(6c^4d^2 + 8c^2de + 3e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{24c^8}$$

$$+ \frac{b(6c^4d^2 + 16c^2de + 9e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} (1-c^2x^2)^{3/2}}{72c^8}$$

$$- \frac{be(8c^2d + 9e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} (1-c^2x^2)^{5/2}}{120c^8} + \frac{be^2 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} (1-c^2x^2)^{7/2}}{56c^8}$$

$$+ \frac{1}{4}d^2x^4(a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{3}dex^6(a + b\operatorname{sech}^{-1}(cx)) + \frac{1}{8}e^2x^8(a + b\operatorname{sech}^{-1}(cx))$$

output

```
-1/24*b*(6*c^4*d^2+8*c^2*d*e+3*e^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^8+1/72*b*(6*c^4*d^2+16*c^2*d*e+9*e^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(3/2)/c^8-1/120*b*e*(8*c^2*d+9*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(5/2)/c^8+1/56*b*e^2*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(7/2)/c^8+1/4*d^2*x^4*(a+b*arcsech(c*x))+1/3*d*e*x^6*(a+b*arcsech(c*x))+1/8*e^2*x^8*(a+b*arcsech(c*x))
```

**Mathematica [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.60

$$\int x^3 (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \frac{1}{24} \left( 6ad^2x^4 + 8adex^6 + 3ae^2x^8 - \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(144e^2 + 8c^2e(56d + 9ex^2) + c^4(420d^2 + 224dex^2 + 54e^2x^4) + 3c^6(70d^2x^2 + 56dex^4 - 105c^8 + bx^4(6d^2 + 8dex^2 + 3e^2x^4) \operatorname{sech}^{-1}(cx))}{105c^8} \right)$$

input

```
Integrate[x^3*(d + e*x^2)^2*(a + b*ArcSech[c*x]),x]
```

output

```
(6*a*d^2*x^4 + 8*a*d*e*x^6 + 3*a*e^2*x^8 - (b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(144*e^2 + 8*c^2*e*(56*d + 9*e*x^2) + c^4*(420*d^2 + 224*d*e*x^2 + 54*e^2*x^4) + 3*c^6*(70*d^2*x^2 + 56*d*e*x^4 + 15*e^2*x^6)))/(105*c^8) + b*x^4*(6*d^2 + 8*d*e*x^2 + 3*e^2*x^4)*ArcSech[c*x])/24
```

**Rubi [A] (verified)**

Time = 0.51 (sec) , antiderivative size = 218, normalized size of antiderivative = 0.78, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {6855, 27, 1578, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx$$

↓ 6855

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^3(3e^2x^4 + 8dex^2 + 6d^2)}{24\sqrt{1-c^2x^2}} dx + \frac{1}{4}d^2x^4(a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{3}dex^6(a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{8}e^2x^8(a + b \operatorname{sech}^{-1}(cx))$$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{1}{24} b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{x^3(3e^2x^4 + 8dex^2 + 6d^2)}{\sqrt{1-c^2x^2}} dx + \frac{1}{4} d^2 x^4 (a + b \operatorname{sech}^{-1}(cx)) + \\
& \quad \frac{1}{3} dex^6 (a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{8} e^2 x^8 (a + b \operatorname{sech}^{-1}(cx)) \\
& \downarrow 1578 \\
& \frac{1}{48} b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{x^2(3e^2x^4 + 8dex^2 + 6d^2)}{\sqrt{1-c^2x^2}} dx^2 + \frac{1}{4} d^2 x^4 (a + b \operatorname{sech}^{-1}(cx)) + \\
& \quad \frac{1}{3} dex^6 (a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{8} e^2 x^8 (a + b \operatorname{sech}^{-1}(cx)) \\
& \downarrow 1195 \\
& \frac{1}{48} b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \left( -\frac{3e^2(1-c^2x^2)^{5/2}}{c^6} + \frac{e(8dc^2+9e)(1-c^2x^2)^{3/2}}{c^6} + \frac{(-6d^2c^4-16dec^2-9e^2)\sqrt{1-c^2x^2}}{c^6} \right) \\
& \quad \frac{1}{4} d^2 x^4 (a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{3} dex^6 (a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{8} e^2 x^8 (a + b \operatorname{sech}^{-1}(cx)) \\
& \downarrow 2009 \\
& \frac{1}{4} d^2 x^4 (a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{3} dex^6 (a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{8} e^2 x^8 (a + b \operatorname{sech}^{-1}(cx)) + \\
& \frac{1}{48} b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \left( -\frac{2e(1-c^2x^2)^{5/2}(8c^2d+9e)}{5c^8} + \frac{6e^2(1-c^2x^2)^{7/2}}{7c^8} + \frac{2(1-c^2x^2)^{3/2}(6c^4d^2+16c^2de+9e^2)}{3c^8} \right)
\end{aligned}$$

input `Int[x^3*(d + e*x^2)^2*(a + b*ArcSech[c*x]),x]`

output `(b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-2*(6*c^4*d^2 + 8*c^2*d*e + 3*e^2)*Sqrt[1 - c^2*x^2])/c^8 + (2*(6*c^4*d^2 + 16*c^2*d*e + 9*e^2)*(1 - c^2*x^2)^(3/2))/(3*c^8) - (2*e*(8*c^2*d + 9*e)*(1 - c^2*x^2)^(5/2))/(5*c^8) + (6*e^2*(1 - c^2*x^2)^(7/2))/(7*c^8))/48 + (d^2*x^4*(a + b*ArcSech[c*x]))/4 + (d*e*x^6*(a + b*ArcSech[c*x]))/3 + (e^2*x^8*(a + b*ArcSech[c*x]))/8`

## Definitions of rubi rules used

- rule 27 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx) /; FreeQ[b, x]]`
- rule 1195 `Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`
- rule 1578 `Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6855 `Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))`

### Maple [A] (verified)

Time = 0.53 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.71

method	result
parts	$a\left(\frac{1}{8}e^2x^8 + \frac{1}{3}dex^6 + \frac{1}{4}d^2x^4\right) + \frac{b\left(\frac{c^4 \operatorname{arcsech}(cx)e^2x^8}{8} + \frac{c^4 \operatorname{arcsech}(cx)dex^6}{3} + \frac{\operatorname{arcsech}(cx)d^2c^4x^4}{4} - \frac{\sqrt{-\frac{cx-1}{cx}}x\sqrt{c^2x^2-1}}{cx}\right)}{2c^4e^2}$
derivativedivides	$\frac{a\left(\frac{c^2d(e^2x^2+c^2d)^3}{3} - \frac{(e^2x^2+c^2d)^4}{4}\right)}{2c^4e^2} + \frac{b\left(-\frac{\operatorname{arcsech}(cx)c^8d^4}{24e^2} + \frac{\operatorname{arcsech}(cx)c^8d^2x^4}{4} + \frac{e \operatorname{arcsech}(cx)c^8dx^6}{3} + \frac{e^2 \operatorname{arcsech}(cx)c^8}{8}\right)}{2c^4e^2}$
default	$\frac{a\left(\frac{c^2d(e^2x^2+c^2d)^3}{3} - \frac{(e^2x^2+c^2d)^4}{4}\right)}{2c^4e^2} + \frac{b\left(-\frac{\operatorname{arcsech}(cx)c^8d^4}{24e^2} + \frac{\operatorname{arcsech}(cx)c^8d^2x^4}{4} + \frac{e \operatorname{arcsech}(cx)c^8dx^6}{3} + \frac{e^2 \operatorname{arcsech}(cx)c^8}{8}\right)}{2c^4e^2}$

```
input int(x^3*(e*x^2+d)^2*(a+b*arcsech(c*x)),x,method=_RETURNVERBOSE)
```

```
output a*(1/8*e^2*x^8+1/3*d*e*x^6+1/4*d^2*x^4)+b/c^4*(1/8*c^4*arcsech(c*x)*e^2*x^8+1/3*c^4*arcsech(c*x)*d*e*x^6+1/4*arcsech(c*x)*d^2*c^4*x^4-1/2520/c^3*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(45*c^6*e^2*x^6+168*c^6*d*e*x^4+210*c^6*d^2*x^2+54*c^4*e^2*x^4+224*c^4*d*e*x^2+420*c^4*d^2+72*c^2*e^2*x^2+448*c^2*d*e+144*e^2))
```

### Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 227, normalized size of antiderivative = 0.82

$$\int x^3(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{315 ac^7 e^2 x^8 + 840 ac^7 dex^6 + 630 ac^7 d^2 x^4 + 105 (3 bc^7 e^2 x^8 + 8 bc^7 dex^6 + 6 bc^7 d^2 x^4) \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}+1}{cx}\right)}{1}$$

```
input integrate(x^3*(e*x^2+d)^2*(a+b*arcsech(c*x)),x, algorithm="fricas")
```

output

```
1/2520*(315*a*c^7*e^2*x^8 + 840*a*c^7*d*e*x^6 + 630*a*c^7*d^2*x^4 + 105*(3
*b*c^7*e^2*x^8 + 8*b*c^7*d*e*x^6 + 6*b*c^7*d^2*x^4)*log((c*x*sqrt(-(c^2*x^
2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (45*b*c^6*e^2*x^7 + 6*(28*b*c^6*d*e + 9*b*
c^4*e^2)*x^5 + 2*(105*b*c^6*d^2 + 112*b*c^4*d*e + 36*b*c^2*e^2)*x^3 + 4*(1
05*b*c^4*d^2 + 112*b*c^2*d*e + 36*b*e^2)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2))
)/c^7
```

**Sympy [A] (verification not implemented)**

Time = 1.70 (sec) , antiderivative size = 332, normalized size of antiderivative = 1.19

$$\int x^3 (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \begin{cases} \frac{ad^2x^4}{4} + \frac{ade^6x^6}{3} + \frac{ae^2x^8}{8} + \frac{bd^2x^4 \operatorname{asech}(cx)}{4} + \frac{bdex^6 \operatorname{asech}(cx)}{3} + \frac{be^2x^8 \operatorname{asech}(cx)}{8} - \frac{bd^2x^2\sqrt{-c^2x^2+1}}{12c^2} - \frac{bdex^4\sqrt{-c^2x^2+1}}{15c^2} - \frac{be^2x^6\sqrt{-c^2x^2+1}}{15c^2} \\ (a + \infty b) \left( \frac{d^2x^4}{4} + \frac{dex^6}{3} + \frac{e^2x^8}{8} \right) \end{cases}$$

input

```
integrate(x**3*(e*x**2+d)**2*(a+b*asech(c*x)),x)
```

output

```
Piecewise((a*d**2*x**4/4 + a*d*e*x**6/3 + a*e**2*x**8/8 + b*d**2*x**4*asec
h(c*x)/4 + b*d*e*x**6*asech(c*x)/3 + b*e**2*x**8*asech(c*x)/8 - b*d**2*x**
2*sqrt(-c**2*x**2 + 1)/(12*c**2) - b*d*e*x**4*sqrt(-c**2*x**2 + 1)/(15*c**
2) - b*e**2*x**6*sqrt(-c**2*x**2 + 1)/(56*c**2) - b*d**2*sqrt(-c**2*x**2 +
1)/(6*c**4) - 4*b*d*e*x**2*sqrt(-c**2*x**2 + 1)/(45*c**4) - 3*b*e**2*x**4
*sqrt(-c**2*x**2 + 1)/(140*c**4) - 8*b*d*e*sqrt(-c**2*x**2 + 1)/(45*c**6)
- b*e**2*x**2*sqrt(-c**2*x**2 + 1)/(35*c**6) - 2*b*e**2*sqrt(-c**2*x**2 +
1)/(35*c**8), Ne(c, 0)), ((a + oo*b)*(d**2*x**4/4 + d*e*x**6/3 + e**2*x**8
/8), True))
```

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 245, normalized size of antiderivative = 0.88

$$\int x^3 (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \frac{1}{8} ae^2 x^8 + \frac{1}{3} adex^6 + \frac{1}{4} ad^2 x^4 + \frac{1}{12} \left( 3x^4 \operatorname{ar} \operatorname{sech}(cx) + \frac{c^2 x^3 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{3}{2}} - 3x \sqrt{\frac{1}{c^2 x^2} - 1}}{c^3} \right) bd^2 + \frac{1}{45} \left( 15x^6 \operatorname{ar} \operatorname{sech}(cx) - \frac{3c^4 x^5 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{5}{2}} - 10c^2 x^3 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{3}{2}} + 15x \sqrt{\frac{1}{c^2 x^2} - 1}}{c^5} \right) bde + \frac{1}{280} \left( 35x^8 \operatorname{ar} \operatorname{sech}(cx) + \frac{5c^6 x^7 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{7}{2}} - 21c^4 x^5 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{5}{2}} + 35c^2 x^3 \left( \frac{1}{c^2 x^2} - 1 \right)^{\frac{3}{2}} - 35x \sqrt{\frac{1}{c^2 x^2} - 1}}{c^7} \right)$$

input `integrate(x^3*(e*x^2+d)^2*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `1/8*a*e^2*x^8 + 1/3*a*d*e*x^6 + 1/4*a*d^2*x^4 + 1/12*(3*x^4*arcsech(c*x) + (c^2*x^3*(1/(c^2*x^2) - 1)^(3/2) - 3*x*sqrt(1/(c^2*x^2) - 1))/c^3)*b*d^2 + 1/45*(15*x^6*arcsech(c*x) - (3*c^4*x^5*(1/(c^2*x^2) - 1)^(5/2) - 10*c^2*x^3*(1/(c^2*x^2) - 1)^(3/2) + 15*x*sqrt(1/(c^2*x^2) - 1))/c^5)*b*d*e + 1/280*(35*x^8*arcsech(c*x) + (5*c^6*x^7*(1/(c^2*x^2) - 1)^(7/2) - 21*c^4*x^5*(1/(c^2*x^2) - 1)^(5/2) + 35*c^2*x^3*(1/(c^2*x^2) - 1)^(3/2) - 35*x*sqrt(1/(c^2*x^2) - 1))/c^7)*b*e^2`

**Giac [F]**

$$\int x^3 (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^2 (b \operatorname{ar} \operatorname{sech}(cx) + a) x^3 dx$$

input `integrate(x^3*(e*x^2+d)^2*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((e*x^2 + d)^2*(b*arcsech(c*x) + a)*x^3, x)`



**Mupad [F(-1)]**

Timed out.

$$\int x^3 (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^3 (ex^2 + d)^2 \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int(x^3*(d + e*x^2)^2*(a + b*acosh(1/(c*x))),x)`

output `int(x^3*(d + e*x^2)^2*(a + b*acosh(1/(c*x))), x)`

**Reduce [F]**

$$\begin{aligned} \int x^3 (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx &= \left( \int a \operatorname{sech}(cx) x^7 dx \right) b e^2 \\ &+ 2 \left( \int a \operatorname{sech}(cx) x^5 dx \right) b d e \\ &+ \left( \int a \operatorname{sech}(cx) x^3 dx \right) b d^2 \\ &+ \frac{a d^2 x^4}{4} + \frac{a d e x^6}{3} + \frac{a e^2 x^8}{8} \end{aligned}$$

input `int(x^3*(e*x^2+d)^2*(a+b*asech(c*x)),x)`

output `(24*int(asech(c*x)*x**7,x)*b*e**2 + 48*int(asech(c*x)*x**5,x)*b*d*e + 24*int(asech(c*x)*x**3,x)*b*d**2 + 6*a*d**2*x**4 + 8*a*d*e*x**6 + 3*a*e**2*x**8)/24`

### 3.106 $\int x(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	849
Mathematica [A] (verified)	850
Rubi [A] (verified)	850
Maple [A] (verified)	852
Fricas [A] (verification not implemented)	853
Sympy [A] (verification not implemented)	853
Maxima [A] (verification not implemented)	854
Giac [F]	855
Mupad [F(-1)]	855
Reduce [F]	855

#### Optimal result

Integrand size = 19, antiderivative size = 230

$$\int x(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx = -\frac{b(3c^4d^2 + 3c^2de + e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{6c^6} + \frac{be(3c^2d + 2e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} (1-c^2x^2)^{3/2}}{18c^6} - \frac{be^2 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} (1-c^2x^2)^{5/2}}{30c^6} + \frac{(d + ex^2)^3 (a + b\operatorname{sech}^{-1}(cx))}{6e} - \frac{bd^3 \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \operatorname{arctanh}(\sqrt{1-c^2x^2})}{6e}$$

output

```
-1/6*b*(3*c^4*d^2+3*c^2*d*e+e^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^6+1/18*b*e*(3*c^2*d+2*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(3/2)/c^6-1/30*b*e^2*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(5/2)/c^6+1/6*(e*x^2+d)^3*(a+b*arcsech(c*x))/e-1/6*b*d^3*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh((-c^2*x^2+1)^(1/2))/e
```

**Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.60

$$\begin{aligned} & \int x(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx \\ &= \frac{1}{6}ax^2(3d^2 + 3dex^2 + e^2x^4) \\ & \quad - \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(8e^2 + 2c^2e(15d + 2ex^2) + 3c^4(15d^2 + 5dex^2 + e^2x^4))}{90c^6} \\ & \quad + \frac{1}{6}bx^2(3d^2 + 3dex^2 + e^2x^4)\operatorname{sech}^{-1}(cx) \end{aligned}$$

input

```
Integrate[x*(d + e*x^2)^2*(a + b*ArcSech[c*x]),x]
```

output

```
(a*x^2*(3*d^2 + 3*d*e*x^2 + e^2*x^4))/6 - (b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(8*e^2 + 2*c^2*e*(15*d + 2*e*x^2) + 3*c^4*(15*d^2 + 5*d*e*x^2 + e^2*x^4)))/(90*c^6) + (b*x^2*(3*d^2 + 3*d*e*x^2 + e^2*x^4)*ArcSech[c*x])/6
```

**Rubi [A] (verified)**

Time = 0.52 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.74, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {6853, 2036, 354, 99, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx \\ & \quad \downarrow \text{6853} \\ & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(ex^2+d)^3}{x\sqrt{1-cx}\sqrt{cx+1}} dx}{6e} + \frac{(d + ex^2)^3 (a + b\operatorname{sech}^{-1}(cx))}{6e} \\ & \quad \downarrow \text{2036} \end{aligned}$$

$$\begin{aligned}
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(ex^2+d)^3}{x\sqrt{1-c^2x^2}} dx}{6e} + \frac{(d+ex^2)^3 (a+b\operatorname{sech}^{-1}(cx))}{6e} \\
& \quad \downarrow \text{354} \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(ex^2+d)^3}{x^2\sqrt{1-c^2x^2}} dx^2}{12e} + \frac{(d+ex^2)^3 (a+b\operatorname{sech}^{-1}(cx))}{6e} \\
& \quad \downarrow \text{99} \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \left( \frac{d^3}{x^2\sqrt{1-c^2x^2}} + \frac{e^3(1-c^2x^2)^{3/2}}{c^4} - \frac{e^2(3dc^2+2e)\sqrt{1-c^2x^2}}{c^4} + \frac{e(3d^2c^4+3dec^2+e^2)}{c^4\sqrt{1-c^2x^2}} \right) dx^2}{12e} + \\
& \quad \frac{(d+ex^2)^3 (a+b\operatorname{sech}^{-1}(cx))}{6e} \\
& \quad \downarrow \text{2009} \\
& \frac{(d+ex^2)^3 (a+b\operatorname{sech}^{-1}(cx))}{6e} + \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -2d^3\operatorname{arctanh}\left(\sqrt{1-c^2x^2}\right) + \frac{2e^2(1-c^2x^2)^{3/2}(3c^2d+2e)}{3c^6} - \frac{2e^3(1-c^2x^2)^{5/2}}{5c^6} - \frac{2e\sqrt{1-c^2x^2}(3c^4d^2+3c^2de+e^2)}{c^6} \right)}{12e}
\end{aligned}$$

input `Int[x*(d + e*x^2)^2*(a + b*ArcSech[c*x]),x]`

output `((d + e*x^2)^3*(a + b*ArcSech[c*x]))/(6*e) + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-2*e*(3*c^4*d^2 + 3*c^2*d*e + e^2)*Sqrt[1 - c^2*x^2])/c^6 + (2*e^2*(3*c^2*d + 2*e)*(1 - c^2*x^2)^(3/2))/(3*c^6) - (2*e^3*(1 - c^2*x^2)^(5/2))/(5*c^6) - 2*d^3*ArcTanh[Sqrt[1 - c^2*x^2]]))/(12*e)`

### Defintions of rubi rules used

rule 99 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] | (GtQ[m, 0] && GeQ[n, -1]))]`

```
rule 354 Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol]
:> Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]
```

```
rule 2009 Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2036 Int[(u_)*((c_) + (d_)*(x_)^(n_))^(q_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_), x_Symbol]
:> Int[u*(a1*a2 + b1*b2*x^n)^p*(c + d*x^n)^q, x] /; FreeQ[{a1, b1, a2, b2, c, d, n, p, q}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))
```

```
rule 6853 Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*(x_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol]
:> Simp[(d + e*x^2)^(p + 1)*((a + b*ArcSech[c*x])/(2*e*(p + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(2*e*(p + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x^2)^(p + 1)/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[p, -1]
```

### Maple [A] (verified)

Time = 0.53 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.23

method	result
parts	$\frac{a(x^2e+d)^3}{6e} + \frac{b \left( \frac{c^2e^2 \operatorname{arcsech}(cx)x^6}{6} + \frac{c^2e \operatorname{arcsech}(cx)x^4d}{2} + \frac{\operatorname{arcsech}(cx)c^2x^2d^2}{2} + \frac{c^2 \operatorname{arcsech}(cx)d^3}{6e} - \frac{\sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}}}{15} \right)}{6e}$
derivativedivides	$\frac{a(e^2x^2+c^2d)^3}{6c^4e} + \frac{b \left( \frac{\operatorname{arcsech}(cx)c^6d^3}{6e} + \frac{\operatorname{arcsech}(cx)c^6d^2x^2}{2} + e \operatorname{arcsech}(cx)c^6dx^4 + \frac{e^2 \operatorname{arcsech}(cx)c^6x^6}{6} - \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}}}{15} \right)}{6c^4e}$
default	$\frac{a(e^2x^2+c^2d)^3}{6c^4e} + \frac{b \left( \frac{\operatorname{arcsech}(cx)c^6d^3}{6e} + \frac{\operatorname{arcsech}(cx)c^6d^2x^2}{2} + e \operatorname{arcsech}(cx)c^6dx^4 + \frac{e^2 \operatorname{arcsech}(cx)c^6x^6}{6} - \frac{\sqrt{-\frac{cx-1}{cx}} cx \sqrt{\frac{cx+1}{cx}}}{15} \right)}{6c^4e}$

```
input int(x*(e*x^2+d)^2*(a+b*arcsech(c*x)), x, method=_RETURNVERBOSE)
```

output

```
1/6*a*(e*x^2+d)^3/e+b/c^2*(1/6*c^2*e^2*arcsech(c*x)*x^6+1/2*c^2*e*arcsech(c*x)*x^4*d+1/2*arcsech(c*x)*c^2*x^2*d^2+1/6*c^2/e*arcsech(c*x)*d^3-1/90/c^3/e*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(15*c^6*d^3*arctanh(1/(-c^2*x^2+1)^(1/2))+45*c^4*d^2*e*(-c^2*x^2+1)^(1/2)+15*c^4*d*e^2*(-c^2*x^2+1)^(1/2)*x^2+3*e^3*(-c^2*x^2+1)^(1/2)*c^4*x^4+30*c^2*d*e^2*(-c^2*x^2+1)^(1/2)+4*e^3*c^2*x^2*(-c^2*x^2+1)^(1/2)+8*e^3*(-c^2*x^2+1)^(1/2))/(-c^2*x^2+1)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.83

$$\int x(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{15 ac^5 e^2 x^6 + 45 ac^5 dex^4 + 45 ac^5 d^2 x^2 + 15 (bc^5 e^2 x^6 + 3 bc^5 dex^4 + 3 bc^5 d^2 x^2) \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2} + 1}}{cx}\right) - (3 bc^5 e^2 x^6 + 3 bc^5 dex^4 + 3 bc^5 d^2 x^2)}{90 c^5}$$

input

```
integrate(x*(e*x^2+d)^2*(a+b*arcsech(c*x)),x, algorithm="fricas")
```

output

```
1/90*(15*a*c^5*e^2*x^6 + 45*a*c^5*d*e*x^4 + 45*a*c^5*d^2*x^2 + 15*(b*c^5*e^2*x^6 + 3*b*c^5*d*e*x^4 + 3*b*c^5*d^2*x^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (3*b*c^4*e^2*x^5 + (15*b*c^4*d*e + 4*b*c^2*e^2)*x^3 + (45*b*c^4*d^2 + 30*b*c^2*d*e + 8*b*e^2)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/c^5
```

**Sympy [A] (verification not implemented)**

Time = 0.91 (sec) , antiderivative size = 252, normalized size of antiderivative = 1.10

$$\int x(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \begin{cases} \frac{ad^2x^2}{2} + \frac{adex^4}{2} + \frac{ae^2x^6}{6} + \frac{bd^2x^2 \operatorname{asech}(cx)}{2} + \frac{bdex^4 \operatorname{asech}(cx)}{2} + \frac{be^2x^6 \operatorname{asech}(cx)}{6} - \frac{bd^2\sqrt{-c^2x^2+1}}{2c^2} - \frac{bdex^2\sqrt{-c^2x^2+1}}{6c^2} - \frac{be^2\sqrt{-c^2x^2+1}}{6c^2} \\ (a + \infty b) \left( \frac{d^2x^2}{2} + \frac{dex^4}{2} + \frac{e^2x^6}{6} \right) \end{cases}$$

input `integrate(x*(e*x**2+d)**2*(a+b*asech(c*x)),x)`

output `Piecewise((a*d**2*x**2/2 + a*d*e*x**4/2 + a*e**2*x**6/6 + b*d**2*x**2*asech(c*x)/2 + b*d*e*x**4*asech(c*x)/2 + b*e**2*x**6*asech(c*x)/6 - b*d**2*sqrt(-c**2*x**2 + 1)/(2*c**2) - b*d*e*x**2*sqrt(-c**2*x**2 + 1)/(6*c**2) - b*e**2*x**4*sqrt(-c**2*x**2 + 1)/(30*c**2) - b*d*e*sqrt(-c**2*x**2 + 1)/(3*c**4) - 2*b*e**2*x**2*sqrt(-c**2*x**2 + 1)/(45*c**4) - 4*b*e**2*sqrt(-c**2*x**2 + 1)/(45*c**6), Ne(c, 0)), ((a + oo*b)*(d**2*x**2/2 + d*e*x**4/2 + e**2*x**6/6), True))`

### Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.80

$$\int x(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{1}{6} ae^2x^6 + \frac{1}{2} adex^4 + \frac{1}{2} ad^2x^2 + \frac{1}{2} \left( x^2 \operatorname{ar} \operatorname{sech}(cx) - \frac{x\sqrt{\frac{1}{c^2x^2} - 1}}{c} \right) bd^2$$

$$+ \frac{1}{6} \left( 3x^4 \operatorname{ar} \operatorname{sech}(cx) + \frac{c^2x^3 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{3}{2}} - 3x\sqrt{\frac{1}{c^2x^2} - 1}}{c^3} \right) bde$$

$$+ \frac{1}{90} \left( 15x^6 \operatorname{ar} \operatorname{sech}(cx) - \frac{3c^4x^5 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{5}{2}} - 10c^2x^3 \left(\frac{1}{c^2x^2} - 1\right)^{\frac{3}{2}} + 15x\sqrt{\frac{1}{c^2x^2} - 1}}{c^5} \right) be^2$$

input `integrate(x*(e*x^2+d)^2*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `1/6*a*e^2*x^6 + 1/2*a*d*e*x^4 + 1/2*a*d^2*x^2 + 1/2*(x^2*arcsech(c*x) - x*sqrt(1/(c^2*x^2) - 1)/c)*b*d^2 + 1/6*(3*x^4*arcsech(c*x) + (c^2*x^3*(1/(c^2*x^2) - 1)^(3/2) - 3*x*sqrt(1/(c^2*x^2) - 1))/c^3)*b*d*e + 1/90*(15*x^6*arcsech(c*x) - (3*c^4*x^5*(1/(c^2*x^2) - 1)^(5/2) - 10*c^2*x^3*(1/(c^2*x^2) - 1)^(3/2) + 15*x*sqrt(1/(c^2*x^2) - 1))/c^5)*b*e^2`

**Giac [F]**

$$\int x(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^2 (b \operatorname{ar} \operatorname{sech}(cx) + a)x dx$$

input `integrate(x*(e*x^2+d)^2*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((e*x^2 + d)^2*(b*arcsech(c*x) + a)*x, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx = \int x (ex^2 + d)^2 \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int(x*(d + e*x^2)^2*(a + b*acosh(1/(c*x))),x)`

output `int(x*(d + e*x^2)^2*(a + b*acosh(1/(c*x))), x)`

**Reduce [F]**

$$\begin{aligned} \int x(d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx &= \left( \int a \operatorname{sech}(cx) x^5 dx \right) b e^2 \\ &+ 2 \left( \int a \operatorname{sech}(cx) x^3 dx \right) b d e \\ &+ \left( \int a \operatorname{sech}(cx) x dx \right) b d^2 \\ &+ \frac{a d^2 x^2}{2} + \frac{a d e x^4}{2} + \frac{a e^2 x^6}{6} \end{aligned}$$

input `int(x*(e*x^2+d)^2*(a+b*asech(c*x)),x)`



output

```
(6*int(asech(c*x)*x**5,x)*b**2 + 12*int(asech(c*x)*x**3,x)*b*d*e + 6*int(asech(c*x)*x,x)*b*d**2 + 3*a*d**2*x**2 + 3*a*d*e*x**4 + a*e**2*x**6)/6
```

$$3.107 \quad \int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x} dx$$

Optimal result	858
Mathematica [A] (verified)	859
Rubi [A] (verified)	859
Maple [A] (verified)	862
Fricas [F]	862
Sympy [F]	863
Maxima [F]	863
Giac [F]	863
Mupad [F(-1)]	864
Reduce [F]	864

## Optimal result

Integrand size = 21, antiderivative size = 370

$$\begin{aligned}
 \int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x} dx = & -\frac{be(6c^2d + e) \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}}{6c^3} \\
 & - \frac{be^2 \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} x^3}{12c} \\
 & + \frac{ibd^2 \sqrt{1 - \frac{1}{c^2x^2}} \operatorname{csc}^{-1}(cx)^2}{2\sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}} \\
 & + dex^2(a + b \operatorname{sech}^{-1}(cx)) + \frac{1}{4}e^2x^4(a + b \operatorname{sech}^{-1}(cx)) \\
 & - \frac{bd^2 \sqrt{1 - \frac{1}{c^2x^2}} \operatorname{csc}^{-1}(cx) \log\left(1 - e^{2i \operatorname{csc}^{-1}(cx)}\right)}{\sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}} \\
 & + \frac{bd^2 \sqrt{1 - \frac{1}{c^2x^2}} \operatorname{csc}^{-1}(cx) \log\left(\frac{1}{x}\right)}{\sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}} \\
 & - d^2(a + b \operatorname{sech}^{-1}(cx)) \log\left(\frac{1}{x}\right) \\
 & + \frac{ibd^2 \sqrt{1 - \frac{1}{c^2x^2}} \operatorname{PolyLog}\left(2, e^{2i \operatorname{csc}^{-1}(cx)}\right)}{2\sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}}
 \end{aligned}$$

output

```

-1/6*b*e*(6*c^2*d+e)*(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)*x/c^3-1/12*b*e^2*(-1
+1/c/x)^(1/2)*(1+1/c/x)^(1/2)*x^3/c+1/2*I*b*d^2*(1-1/c^2/x^2)^(1/2)*arccsc
(c*x)^2/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)+d*e*x^2*(a+b*arcsech(c*x))+1/4*e^
2*x^4*(a+b*arcsech(c*x))-b*d^2*(1-1/c^2/x^2)^(1/2)*arccsc(c*x)*ln(1-(I/c/x
+(1-1/c^2/x^2)^(1/2))^2)/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)+b*d^2*(1-1/c^2/x
^2)^(1/2)*arccsc(c*x)*ln(1/x)/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)-d^2*(a+b*ar
csech(c*x))*ln(1/x)+1/2*I*b*d^2*(1-1/c^2/x^2)^(1/2)*polylog(2,(I/c/x+(1-1/
c^2/x^2)^(1/2))^2)/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)

```

**Mathematica [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.48

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x} dx = adex^2 + \frac{1}{4}ae^2x^4 - \frac{bde\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{c^2}$$

$$- \frac{be^2\sqrt{\frac{1-cx}{1+cx}}(1+cx)(2+c^2x^2)}{12c^4} + bdex^2\operatorname{sech}^{-1}(cx)$$

$$+ \frac{1}{4}be^2x^4\operatorname{sech}^{-1}(cx) - \frac{1}{2}bd^2\operatorname{sech}^{-1}(cx) \left( \operatorname{sech}^{-1}(cx) \right.$$

$$\left. + 2 \log \left( 1 + e^{-2\operatorname{sech}^{-1}(cx)} \right) \right)$$

$$+ ad^2 \log(x) + \frac{1}{2}bd^2 \operatorname{PolyLog} \left( 2, -e^{-2\operatorname{sech}^{-1}(cx)} \right)$$

input

```
Integrate[((d + e*x^2)^2*(a + b*ArcSech[c*x]))/x,x]
```

output

```
a*d*e*x^2 + (a*e^2*x^4)/4 - (b*d*e*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/c^2 - (b*e^2*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(2 + c^2*x^2))/(12*c^4) + b*d*e*x^2*ArcSech[c*x] + (b*e^2*x^4*ArcSech[c*x])/4 - (b*d^2*ArcSech[c*x]*(ArcSech[c*x] + 2*Log[1 + E^(-2*ArcSech[c*x])]))/2 + a*d^2*Log[x] + (b*d^2*PolyLog[2, -E^(-2*ArcSech[c*x])])/2
```

**Rubi [A] (verified)**

Time = 1.44 (sec) , antiderivative size = 401, normalized size of antiderivative = 1.08, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {6857, 6373, 27, 7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x} dx$$

↓ 6857

$$-\int \left(\frac{d}{x^2} + e\right)^2 x^5 \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) d\frac{1}{x}$$

↓ 6373

$$\frac{b \int -\frac{e\left(\frac{4d}{x^2}+e\right)x^4-4d^2 \log\left(\frac{1}{x}\right)}{4\sqrt{\frac{1}{cx}-1}\sqrt{1+\frac{1}{cx}}} d\frac{1}{x}}{c} - d^2 \log\left(\frac{1}{x}\right) \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) + dex^2 \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) + \frac{1}{4}e^2 x^4 \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right)$$

↓ 27

$$-\frac{b \int \frac{e\left(\frac{4d}{x^2}+e\right)x^4-4d^2 \log\left(\frac{1}{x}\right)}{\sqrt{\frac{1}{cx}-1}\sqrt{1+\frac{1}{cx}}} d\frac{1}{x}}{4c} - d^2 \log\left(\frac{1}{x}\right) \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) + dex^2 \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) + \frac{1}{4}e^2 x^4 \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right)$$

↓ 7293

$$-\frac{b \int \left(\frac{e\left(\frac{4d}{x^2}+e\right)x^4}{\sqrt{\frac{1}{cx}-1}\sqrt{1+\frac{1}{cx}}} - \frac{4d^2 \log\left(\frac{1}{x}\right)}{\sqrt{\frac{1}{cx}-1}\sqrt{1+\frac{1}{cx}}}\right) d\frac{1}{x}}{4c} - d^2 \log\left(\frac{1}{x}\right) \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) + dex^2 \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) + \frac{1}{4}e^2 x^4 \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right)$$

↓ 2009

$$-d^2 \log\left(\frac{1}{x}\right) \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) + dex^2 \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) + \frac{1}{4}e^2 x^4 \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) - b \left( -\frac{2icd^2 \sqrt{1-\frac{1}{c^2x^2}} \operatorname{PolyLog}\left(2, e^{2i \arcsin\left(\frac{1}{cx}\right)}\right)}{\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}} - \frac{2icd^2 \sqrt{1-\frac{1}{c^2x^2}} \arcsin\left(\frac{1}{cx}\right)^2}{\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}} + \frac{4cd^2 \sqrt{1-\frac{1}{c^2x^2}} \arcsin\left(\frac{1}{cx}\right) \log\left(1-e^{2i \arcsin\left(\frac{1}{cx}\right)}\right)}{\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}} - \frac{4cd^2 \sqrt{1-\frac{1}{c^2x^2}} \arcsin\left(\frac{1}{cx}\right)}{\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}} \right)$$

4c

input

Int[((d + e\*x^2)^2\*(a + b\*ArcSech[c\*x]))/x,x]

output

```
d*e*x^2*(a + b*ArcCosh[1/(c*x)]) + (e^2*x^4*(a + b*ArcCosh[1/(c*x)]))/4 -
d^2*(a + b*ArcCosh[1/(c*x)]*Log[x^(-1)] - (b*((2*e*(6*d + e/c^2)*Sqrt[-1
+ 1/(c*x)]*Sqrt[1 + 1/(c*x)]*x)/3 + (e^2*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*
x)]*x^3)/3 - ((2*I)*c*d^2*Sqrt[1 - 1/(c^2*x^2)]*ArcSin[1/(c*x)]^2)/(Sqrt[-
1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) + (4*c*d^2*Sqrt[1 - 1/(c^2*x^2)]*ArcSin[1/
(c*x)]*Log[1 - E^((2*I)*ArcSin[1/(c*x)])])/(Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/
(c*x)]) - (4*c*d^2*Sqrt[1 - 1/(c^2*x^2)]*ArcSin[1/(c*x)]*Log[x^(-1)])/(Sqr
t[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) - ((2*I)*c*d^2*Sqrt[1 - 1/(c^2*x^2)]*Po
lyLog[2, E^((2*I)*ArcSin[1/(c*x)])])/(Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)
]))/(4*c)
```

### Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6373

```
Int[((a_) + ArcCosh[(c_)*(x_)])*(b_))*((f_)*(x_)^(m_))*((d_) + (e_)*(x
_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Sim
p[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] &&
NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && Le
Q[m + p, 0]))
```

rule 6857

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_))^(n_)*(x_)^(m_))*((d_) + (e_)*(x
_)^2)^(p_), x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x
^(m + 2*(p + 1))), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0
] && IntegersQ[m, p]
```

rule 7293

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v
]
```

**Maple [A] (verified)**

Time = 0.99 (sec) , antiderivative size = 276, normalized size of antiderivative = 0.75

method	result
parts	$a\left(\frac{e^2x^4}{4} + dex^2 + d^2 \ln(x)\right) + b\left(\frac{d^2 \operatorname{arcsech}(cx)^2}{2} + \frac{e\left(12 \operatorname{arcsech}(cx)c^4dx^2 + 3e \operatorname{arcsech}(cx)c^4x^4 - 12c^3\right)}{2}\right)$
derivativedivides	$ade x^2 + \frac{ae^2x^4}{4} + ad^2 \ln(cx) + \frac{b\left(\frac{c^4d^2 \operatorname{arcsech}(cx)^2}{2} + \frac{e\left(12 \operatorname{arcsech}(cx)c^4dx^2 + 3e \operatorname{arcsech}(cx)c^4x^4 - 12c^3dx\sqrt{\frac{cx}{c-x}}\right)}{2}\right)}{2}$
default	$ade x^2 + \frac{ae^2x^4}{4} + ad^2 \ln(cx) + \frac{b\left(\frac{c^4d^2 \operatorname{arcsech}(cx)^2}{2} + \frac{e\left(12 \operatorname{arcsech}(cx)c^4dx^2 + 3e \operatorname{arcsech}(cx)c^4x^4 - 12c^3dx\sqrt{\frac{cx}{c-x}}\right)}{2}\right)}{2}$

input `int((e*x^2+d)^2*(a+b*arcsech(c*x))/x,x,method=_RETURNVERBOSE)`

output `a*(1/4*e^2*x^4+d*e*x^2+d^2*ln(x))+b*(1/2*d^2*arcsech(c*x)^2+1/12/c^4*e*(12*arcsech(c*x)*c^4*d*x^2+3*e*arcsech(c*x)*c^4*x^4-12*c^3*d*x*((c*x+1)/c/x)^(1/2)*(-(c*x-1)/c/x)^(1/2)-c^3*e*x^3*((c*x+1)/c/x)^(1/2)*(-(c*x-1)/c/x)^(1/2)-2*c*e*x*((c*x+1)/c/x)^(1/2)*(-(c*x-1)/c/x)^(1/2)+12*c^2*d+2*e)-d^2*arcsech(c*x)*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)-1/2*d^2*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2))`

**Fricas [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{ar} \operatorname{sech}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x,x, algorithm="fricas")`

output `integral((a*e^2*x^4 + 2*a*d*e*x^2 + a*d^2 + (b*e^2*x^4 + 2*b*d*e*x^2 + b*d^2)*arcsech(c*x))/x, x)`

**Sympy [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(a + b \operatorname{arsech}(cx)) (d + ex^2)^2}{x} dx$$

input `integrate((e*x**2+d)**2*(a+b*asech(c*x))/x,x)`

output `Integral((a + b*asech(c*x))*(d + e*x**2)**2/x, x)`

**Maxima [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{arsech}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x,x, algorithm="maxima")`

output `1/4*a*e^2*x^4 + a*d*e*x^2 + a*d^2*log(x) + integrate(b*e^2*x^3*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x)) + 2*b*d*e*x*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x)) + b*d^2*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/x, x)`

**Giac [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{arsech}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x,x, algorithm="giac")`

output `integrate((e*x^2 + d)^2*(b*arcsech(c*x) + a)/x, x)`



**Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(ex^2 + d)^2 (a + b \operatorname{acosh}(\frac{1}{cx}))}{x} dx$$

input `int(((d + e*x^2)^2*(a + b*acosh(1/(c*x))))/x,x)`output `int(((d + e*x^2)^2*(a + b*acosh(1/(c*x))))/x, x)`**Reduce [F]**

$$\begin{aligned} \int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x} dx &= \left( \int \frac{a \operatorname{sech}(cx)}{x} dx \right) b d^2 + \left( \int a \operatorname{sech}(cx) x^3 dx \right) b e^2 \\ &+ 2 \left( \int a \operatorname{sech}(cx) x dx \right) b d e \\ &+ \log(x) a d^2 + a d e x^2 + \frac{a e^2 x^4}{4} \end{aligned}$$

input `int((e*x^2+d)^2*(a+b*asech(c*x))/x,x)`output `(4*int(asech(c*x)/x,x)*b*d**2 + 4*int(asech(c*x)*x**3,x)*b*e**2 + 8*int(asech(c*x)*x,x)*b*d*e + 4*log(x)*a*d**2 + 4*a*d*e*x**2 + a*e**2*x**4)/4`

**3.108**  $\int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x^3} dx$

Optimal result	865
Mathematica [A] (verified)	866
Rubi [A] (verified)	867
Maple [A] (verified)	869
Fricas [F]	870
Sympy [F]	870
Maxima [F]	870
Giac [F]	871
Mupad [F(-1)]	871
Reduce [F]	871

**Optimal result**

Integrand size = 21, antiderivative size = 373

$$\int \frac{(d+ex^2)^2 (a+b\operatorname{sech}^{-1}(cx))}{x^3} dx = \frac{bcd^2 \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}}{4x} - \frac{be^2 \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} x}{2c}$$

$$+ \frac{ibde \sqrt{1 - \frac{1}{c^2 x^2}} \operatorname{csc}^{-1}(cx)^2}{\sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}} + \frac{1}{4} bc^2 d^2 \operatorname{sech}^{-1}(cx)$$

$$- \frac{d^2 (a + b\operatorname{sech}^{-1}(cx))}{2x^2} + \frac{1}{2} e^2 x^2 (a + b\operatorname{sech}^{-1}(cx))$$

$$- \frac{2bde \sqrt{1 - \frac{1}{c^2 x^2}} \operatorname{csc}^{-1}(cx) \log\left(1 - e^{2i \operatorname{csc}^{-1}(cx)}\right)}{\sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}}$$

$$+ \frac{2bde \sqrt{1 - \frac{1}{c^2 x^2}} \operatorname{csc}^{-1}(cx) \log\left(\frac{1}{x}\right)}{\sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}}$$

$$- 2de (a + b\operatorname{sech}^{-1}(cx)) \log\left(\frac{1}{x}\right)$$

$$+ \frac{ibde \sqrt{1 - \frac{1}{c^2 x^2}} \operatorname{PolyLog}\left(2, e^{2i \operatorname{csc}^{-1}(cx)}\right)}{\sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}}$$

output

```

1/4*b*c*d^2*(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)/x-1/2*b*e^2*(-1+1/c/x)^(1/2)*
(1+1/c/x)^(1/2)*x/c+I*b*d*e*(1-1/c^2/x^2)^(1/2)*arccsc(c*x)^2/(-1+1/c/x)^(
1/2)/(1+1/c/x)^(1/2)+1/4*b*c^2*d^2*arcsech(c*x)-1/2*d^2*(a+b*arcsech(c*x))
/x^2+1/2*e^2*x^2*(a+b*arcsech(c*x))-2*b*d*e*(1-1/c^2/x^2)^(1/2)*arccsc(c*x
)*ln(1-(I/c/x+(1-1/c^2/x^2)^(1/2))^2)/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)+2*b
*d*e*(1-1/c^2/x^2)^(1/2)*arccsc(c*x)*ln(1/x)/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1
/2)-2*d*e*(a+b*arcsech(c*x))*ln(1/x)+I*b*d*e*(1-1/c^2/x^2)^(1/2)*polylog(2
,(I/c/x+(1-1/c^2/x^2)^(1/2))^2)/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)

```

**Mathematica [A] (verified)**

Time = 0.67 (sec) , antiderivative size = 223, normalized size of antiderivative = 0.60

$$\begin{aligned}
& \int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx \\
&= \frac{1}{4} \left( -\frac{2ad^2}{x^2} + 2ae^2x^2 - \frac{2be^2 \sqrt{\frac{1-cx}{1+cx}}(1+cx)}{c^2} - \frac{2bd^2 \operatorname{sech}^{-1}(cx)}{x^2} + 2be^2x^2 \operatorname{sech}^{-1}(cx) \right. \\
&\quad \left. + \frac{bd^2 \sqrt{\frac{1-cx}{1+cx}} \left( \sqrt{1-cx}(1+cx) + 2c^2x^2 \sqrt{1+cx} \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)}{x^2 \sqrt{1-cx}} \right. \\
&\quad \left. - 4bd \operatorname{sech}^{-1}(cx) \left( \operatorname{sech}^{-1}(cx) + 2 \log \left( 1 + e^{-2 \operatorname{sech}^{-1}(cx)} \right) \right) + 8ade \log(x) \right. \\
&\quad \left. + 4bde \operatorname{PolyLog} \left( 2, -e^{-2 \operatorname{sech}^{-1}(cx)} \right) \right)
\end{aligned}$$

input

```
Integrate[((d + e*x^2)^2*(a + b*ArcSech[c*x]))/x^3,x]
```

output

```

((-2*a*d^2)/x^2 + 2*a*e^2*x^2 - (2*b*e^2*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*
x))/c^2 - (2*b*d^2*ArcSech[c*x])/x^2 + 2*b*e^2*x^2*ArcSech[c*x] + (b*d^2*S
qrt[(1 - c*x)/(1 + c*x)]*(Sqrt[1 - c*x]*(1 + c*x) + 2*c^2*x^2*Sqrt[1 + c*x
]*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]]))/(x^2*Sqrt[1 - c*x]) - 4*b*d*e*Arc
Sech[c*x]*(ArcSech[c*x] + 2*Log[1 + E^(-2*ArcSech[c*x])]) + 8*a*d*e*Log[x]
+ 4*b*d*e*PolyLog[2, -E^(-2*ArcSech[c*x])])/4

```

**Rubi [A] (verified)**

Time = 1.37 (sec) , antiderivative size = 411, normalized size of antiderivative = 1.10, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {6857, 6373, 27, 7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx$$

$$\downarrow 6857$$

$$- \int \left( \frac{d}{x^2} + e \right) x^3 \left( a + \operatorname{barccosh} \left( \frac{1}{cx} \right) \right) d \frac{1}{x}$$

$$\downarrow 6373$$

$$\frac{b \int -\frac{d^2}{x^2} - 4e \log\left(\frac{1}{x}\right) d + e^2 x^2}{2\sqrt{\frac{1}{cx}-1}\sqrt{1+\frac{1}{cx}}} d \frac{1}{x}}{c} - \frac{d^2 (a + \operatorname{barccosh}(\frac{1}{cx}))}{2x^2} - 2de \log\left(\frac{1}{x}\right) \left( a + \operatorname{barccosh}(\frac{1}{cx}) \right) + \frac{1}{2} e^2 x^2 \left( a + \operatorname{barccosh}(\frac{1}{cx}) \right)$$

$$\downarrow 27$$

$$- \frac{b \int -\frac{d^2}{x^2} - 4e \log\left(\frac{1}{x}\right) d + e^2 x^2}{\sqrt{\frac{1}{cx}-1}\sqrt{1+\frac{1}{cx}}} d \frac{1}{x}}{2c} - \frac{d^2 (a + \operatorname{barccosh}(\frac{1}{cx}))}{2x^2} - 2de \log\left(\frac{1}{x}\right) \left( a + \operatorname{barccosh}(\frac{1}{cx}) \right) + \frac{1}{2} e^2 x^2 \left( a + \operatorname{barccosh}(\frac{1}{cx}) \right)$$

$$\downarrow 7293$$

$$- \frac{b \int \left( -\frac{d^2}{\sqrt{\frac{1}{cx}-1}\sqrt{1+\frac{1}{cx}}} x^2 - \frac{4e \log\left(\frac{1}{x}\right) d}{\sqrt{\frac{1}{cx}-1}\sqrt{1+\frac{1}{cx}}} + \frac{e^2 x^2}{\sqrt{\frac{1}{cx}-1}\sqrt{1+\frac{1}{cx}}} \right) d \frac{1}{x}}{2c} - \frac{d^2 (a + \operatorname{barccosh}(\frac{1}{cx}))}{2x^2} - 2de \log\left(\frac{1}{x}\right) \left( a + \operatorname{barccosh}(\frac{1}{cx}) \right) + \frac{1}{2} e^2 x^2 \left( a + \operatorname{barccosh}(\frac{1}{cx}) \right)$$

$$\downarrow 2009$$

$$\frac{-\frac{d^2(a + \operatorname{barccosh}(\frac{1}{cx}))}{2x^2} - 2de \log\left(\frac{1}{x}\right) \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) + \frac{1}{2}e^2x^2 \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) - b \left( -\frac{1}{2}c^3d^2 \operatorname{arccosh}\left(\frac{1}{cx}\right) - \frac{2icde\sqrt{1-\frac{1}{c^2x^2}} \operatorname{PolyLog}\left(2, e^{2i \operatorname{arcsin}\left(\frac{1}{cx}\right)}\right)}{\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}} - \frac{2icde\sqrt{1-\frac{1}{c^2x^2}} \operatorname{arcsin}\left(\frac{1}{cx}\right)^2}{\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}} + \frac{4cde\sqrt{1-\frac{1}{c^2x^2}} \operatorname{arcsin}\left(\frac{1}{cx}\right) \log\left(\frac{1}{cx}\right)}{\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}} \right)}{2c}$$

input `Int[((d + e*x^2)^2*(a + b*ArcSech[c*x]))/x^3,x]`

output `-1/2*(d^2*(a + b*ArcCosh[1/(c*x)]))/x^2 + (e^2*x^2*(a + b*ArcCosh[1/(c*x)]))/2 - 2*d*e*(a + b*ArcCosh[1/(c*x)]*Log[x^(-1)] - (b*(-1/2*(c^2*d^2*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]))/x + e^2*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]*x - (c^3*d^2*ArcCosh[1/(c*x)])/2 - ((2*I)*c*d*e*Sqrt[1 - 1/(c^2*x^2)]*ArcSin[1/(c*x)]^2)/(Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) + (4*c*d*e*Sqrt[1 - 1/(c^2*x^2)]*ArcSin[1/(c*x)]*Log[1 - E^((2*I)*ArcSin[1/(c*x)])])/(Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) - (4*c*d*e*Sqrt[1 - 1/(c^2*x^2)]*ArcSin[1/(c*x)]*Log[x^(-1)])/(Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) - ((2*I)*c*d*e*Sqrt[1 - 1/(c^2*x^2)]*PolyLog[2, E^((2*I)*ArcSin[1/(c*x)])])/(Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)])))/(2*c)`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6373 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))`

rule 6857

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> -Subst[Int[(e + d*x^2)^p*(a + b*ArcCosh[x/c])^n/x^(m + 2*(p + 1))], x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0] && IntegersQ[m, p]
```

rule 7293

```
Int[u_, x_Symbol] :> With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]
```

### Maple [A] (verified)

Time = 1.31 (sec) , antiderivative size = 250, normalized size of antiderivative = 0.67

method	result
parts	$a \left( \frac{e^2 x^2}{2} - \frac{d^2}{2x^2} + 2de \ln(x) \right) + bde \operatorname{arcsech}(cx)^2 + \frac{bc^2 d^2 \operatorname{arcsech}(cx)}{4} + \frac{bc d^2 \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}}{4x} -$
derivativedivides	$c^2 \left( \frac{a x^2 e^2}{2c^2} - \frac{a d^2}{2c^2 x^2} + \frac{2ade \ln(cx)}{c^2} + \frac{bde \operatorname{arcsech}(cx)^2}{c^2} + \frac{b d^2 \operatorname{arcsech}(cx)}{4} + \frac{b d^2 \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}}{4cx} - \frac{b \operatorname{arcsech}(cx)}{2} \right)$
default	$c^2 \left( \frac{a x^2 e^2}{2c^2} - \frac{a d^2}{2c^2 x^2} + \frac{2ade \ln(cx)}{c^2} + \frac{bde \operatorname{arcsech}(cx)^2}{c^2} + \frac{b d^2 \operatorname{arcsech}(cx)}{4} + \frac{b d^2 \sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}}}{4cx} - \frac{b \operatorname{arcsech}(cx)}{2} \right)$

input

```
int((e*x^2+d)^2*(a+b*arcsech(c*x))/x^3,x,method=_RETURNVERBOSE)
```

output

```
a*(1/2*e^2*x^2-1/2*d^2/x^2+2*d*e*ln(x))+b*d*e*arcsech(c*x)^2+1/4*b*c^2*d^2*arcsech(c*x)+1/4*b*c*d^2/x*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)-1/2*b*d^2/x^2*arcsech(c*x)+1/2*b*e^2*arcsech(c*x)*x^2-1/2*b/c*e^2*(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)*x+1/2*b/c^2*e^2-2*b*e*d*arcsech(c*x)*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)-b*e*d*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)
```

**Fricas [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{ar} \operatorname{sech}(cx) + a)}{x^3} dx$$

input `integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x^3,x, algorithm="fricas")`

output `integral((a*e^2*x^4 + 2*a*d*e*x^2 + a*d^2 + (b*e^2*x^4 + 2*b*d*e*x^2 + b*d^2)*arcsech(c*x))/x^3, x)`

**Sympy [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(a + b \operatorname{asech}(cx)) (d + ex^2)^2}{x^3} dx$$

input `integrate((e*x**2+d)**2*(a+b*asech(c*x))/x**3,x)`

output `Integral((a + b*asech(c*x))*(d + e*x**2)**2/x**3, x)`

**Maxima [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{ar} \operatorname{sech}(cx) + a)}{x^3} dx$$

input `integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x^3,x, algorithm="maxima")`

output `1/2*a*e^2*x^2 - 1/8*b*d^2*((2*c^4*x*sqrt(1/(c^2*x^2) - 1)/(c^2*x^2*(1/(c^2*x^2) - 1) - 1) - c^3*log(c*x*sqrt(1/(c^2*x^2) - 1) + 1) + c^3*log(c*x*sqrt(1/(c^2*x^2) - 1) - 1))/c + 4*arcsech(c*x)/x^2) + 2*a*d*e*log(x) - 1/2*a*d^2/x^2 + integrate(b*e^2*x*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x)) + 2*b*d*e*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/x, x)`

**Giac [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{arsech}(cx) + a)}{x^3} dx$$

input `integrate((e*x^2+d)^2*(a+b*arcsech(c*x))/x^3,x, algorithm="giac")`

output `integrate((e*x^2 + d)^2*(b*arcsech(c*x) + a)/x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(ex^2 + d)^2 (a + b \operatorname{acosh}(\frac{1}{cx}))}{x^3} dx$$

input `int(((d + e*x^2)^2*(a + b*acosh(1/(c*x))))/x^3,x)`

output `int(((d + e*x^2)^2*(a + b*acosh(1/(c*x))))/x^3, x)`

**Reduce [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx$$

$$= \frac{2 \left( \int \frac{\operatorname{asech}(cx)}{x^3} dx \right) b d^2 x^2 + 4 \left( \int \frac{\operatorname{asech}(cx)}{x} dx \right) b d e x^2 + 2 \left( \int \operatorname{asech}(cx) x dx \right) b e^2 x^2 + 4 \log(x) a d e x^2 - a d^2 + a e^2 x^4}{2x^2}$$

input `int((e*x^2+d)^2*(a+b*asech(c*x))/x^3,x)`

output `(2*int(asech(c*x)/x**3,x)*b*d**2*x**2 + 4*int(asech(c*x)/x,x)*b*d*e*x**2 + 2*int(asech(c*x)*x,x)*b*e**2*x**2 + 4*log(x)*a*d*e*x**2 - a*d**2 + a*e**2*x**4)/(2*x**2)`



$$3.109 \quad \int \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx$$

Optimal result	873
Mathematica [C] (warning: unable to verify)	874
Rubi [A] (verified)	875
Maple [C] (warning: unable to verify)	878
Fricas [F]	879
Sympy [F]	879
Maxima [F(-2)]	879
Giac [F]	880
Mupad [F(-1)]	880
Reduce [F]	880

**Optimal result**

Integrand size = 21, antiderivative size = 519

$$\begin{aligned}
\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{d + ex^2} dx = & \frac{x(a + b\operatorname{sech}^{-1}(cx))}{e} - \frac{b \arctan\left(\sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}\right)}{ce} \\
& + \frac{\sqrt{-d}(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-d}e\operatorname{sech}^{-1}(cx)}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2e^{3/2}} \\
& - \frac{\sqrt{-d}(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-d}e\operatorname{sech}^{-1}(cx)}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2e^{3/2}} \\
& + \frac{\sqrt{-d}(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-d}e\operatorname{sech}^{-1}(cx)}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2e^{3/2}} \\
& - \frac{\sqrt{-d}(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-d}e\operatorname{sech}^{-1}(cx)}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2e^{3/2}} \\
& - \frac{b\sqrt{-d} \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-d}e\operatorname{sech}^{-1}(cx)}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2e^{3/2}} \\
& + \frac{b\sqrt{-d} \operatorname{PolyLog}\left(2, \frac{c\sqrt{-d}e\operatorname{sech}^{-1}(cx)}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2e^{3/2}} \\
& - \frac{b\sqrt{-d} \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-d}e\operatorname{sech}^{-1}(cx)}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2e^{3/2}} \\
& + \frac{b\sqrt{-d} \operatorname{PolyLog}\left(2, \frac{c\sqrt{-d}e\operatorname{sech}^{-1}(cx)}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2e^{3/2}}
\end{aligned}$$

output

```
x*(a+b*arcsech(c*x))/e-b*arctan((-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/c/e+1/2*
(-d)^(1/2)*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1
+1/c/x)^(1/2)))/(e^(1/2)-(c^2*d+e)^(1/2))/e^(3/2)-1/2*(-d)^(1/2)*(a+b*arcs
ech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1
/2)-(c^2*d+e)^(1/2))/e^(3/2)+1/2*(-d)^(1/2)*(a+b*arcsech(c*x))*ln(1-c*(-d)
)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(1/2)
)/e^(3/2)-1/2*(-d)^(1/2)*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1
/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(1/2))/e^(3/2)-1/2*b*(-d)
^(1/2)*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e
^(1/2)-(c^2*d+e)^(1/2))/e^(3/2)+1/2*b*(-d)^(1/2)*polylog(2,c*(-d)^(1/2)*(
1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2))/e^(3/2)
-1/2*b*(-d)^(1/2)*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x
)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2))/e^(3/2)+1/2*b*(-d)^(1/2)*polylog(2,c*(
-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2
)))/e^(3/2)
```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 1.84 (sec) , antiderivative size = 921, normalized size of antiderivative = 1.77

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \text{Too large to display}$$

input

```
Integrate[(x^2*(a + b*ArcSech[c*x]))/(d + e*x^2),x]
```

output

```
(4*a*c*Sqrt[e]*x - 4*a*c*Sqrt[d]*ArcTan[(Sqrt[e]*x)/Sqrt[d]] + b*(4*Sqrt[e]
)*(c*x*ArcSech[c*x] - 2*ArcTan[Tanh[ArcSech[c*x]/2]]) - (2*I)*c*Sqrt[d]*((
-4*I)*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*ArcTanh[((I*c*Sqrt
[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])/Sqrt[c^2*d + e]] + ArcSech[c*x]*Log[1
+ E^(-2*ArcSech[c*x])] - ArcSech[c*x]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d +
e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + (2*I)*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sq
rt[d])]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcS
ech[c*x])] - ArcSech[c*x]*Log[1 + (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[
d]*E^ArcSech[c*x])] - (2*I)*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[
2]]*Log[1 + (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] +
PolyLog[2, (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] +
PolyLog[2, ((-I)*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])])
+ (2*I)*c*Sqrt[d]*((-4*I)*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2
]]*ArcTanh[((-I)*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])/Sqrt[c^2*d +
e]] + ArcSech[c*x]*Log[1 + E^(-2*ArcSech[c*x])] - ArcSech[c*x]*Log[1 + (I*
(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + (2*I)*ArcSin[S
qrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*
d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - ArcSech[c*x]*Log[1 - (I*(Sqrt[e] +
Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - (2*I)*ArcSin[Sqrt[1 - (I*S
qrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 - (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(...
```

**Rubi [A] (verified)**

Time = 1.58 (sec) , antiderivative size = 571, normalized size of antiderivative = 1.10, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6857, 6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{d + ex^2} dx$$

$$\downarrow 6857$$

$$- \int \frac{x^2(a + b\operatorname{arccosh}(\frac{1}{cx}))}{\frac{d}{x^2} + e} d\frac{1}{x}$$

$$\downarrow 6374$$

$$\begin{aligned}
& - \int \left( \frac{x^2(a + \operatorname{barccosh}(\frac{1}{cx}))}{e} - \frac{d(a + \operatorname{barccosh}(\frac{1}{cx}))}{e(\frac{d}{x^2} + e)} \right) d\frac{1}{x} \\
& \quad \downarrow \text{2009} \\
& \frac{\sqrt{-d}(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( 1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{c^2d + e}}} \right)}{2e^{3/2}} - \\
& \frac{\sqrt{-d}(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{c^2d + e}}} + 1 \right)}{2e^{3/2}} + \\
& \frac{\sqrt{-d}(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( 1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{c^2d + e + \sqrt{e}}} \right)}{2e^{3/2}} - \\
& \frac{\sqrt{-d}(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{c^2d + e + \sqrt{e}}} + 1 \right)}{2e^{3/2}} + \frac{x(a + \operatorname{barccosh}(\frac{1}{cx}))}{e} - \\
& \frac{b\sqrt{-d} \operatorname{PolyLog} \left( 2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{dc^2 + e}}} \right)}{2e^{3/2}} + \frac{b\sqrt{-d} \operatorname{PolyLog} \left( 2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{dc^2 + e}}} \right)}{2e^{3/2}} \\
& \frac{b\sqrt{-d} \operatorname{PolyLog} \left( 2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}} \right)}{2e^{3/2}} + \frac{b\sqrt{-d} \operatorname{PolyLog} \left( 2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}} \right)}{2e^{3/2}} \\
& \quad \frac{b \arctan \left( \sqrt{\frac{1}{cx}} - 1\sqrt{\frac{1}{cx} + 1} \right)}{ce}
\end{aligned}$$

input `Int[(x^2*(a + b*ArcSech[c*x]))/(d + e*x^2),x]`

output

```
(x*(a + b*ArcCosh[1/(c*x)]))/e - (b*ArcTan[Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x]])/(c*e) + (Sqrt[-d]*(a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])])/(2*e^(3/2)) - (Sqrt[-d]*(a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])])/(2*e^(3/2)) + (Sqrt[-d]*(a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(2*e^(3/2)) - (Sqrt[-d]*(a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(2*e^(3/2)) - (b*Sqrt[-d]*PolyLog[2, -((c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e]))])/(2*e^(3/2)) + (b*Sqrt[-d]*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])])/(2*e^(3/2)) - (b*Sqrt[-d]*PolyLog[2, -((c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e]))])/(2*e^(3/2)) + (b*Sqrt[-d]*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(2*e^(3/2))
```

### Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6374

```
Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

rule 6857

```
Int[((a_) + ArcSech[(c_)*(x_)]*(b_))^(n_)*(x_)^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x^(m + 2*(p + 1))), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0] && IntegersQ[m, p]
```

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 36.37 (sec) , antiderivative size = 411, normalized size of antiderivative = 0.79

method	result
parts	$\frac{ax}{e} - \frac{ad \arctan\left(\frac{xe}{\sqrt{de}}\right)}{e\sqrt{de}} + \frac{b \operatorname{arcsech}(cx)x}{e} - \frac{2b \arctan\left(\frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}\right)}{ce} - \frac{bcd \left( \sum_{-R1=\text{RootOf}(c^2d\_Z^4+}$
derivativedivides	$\frac{ac^3x}{e} - \frac{ac^3d \arctan\left(\frac{xe}{\sqrt{de}}\right)}{e\sqrt{de}} + bc^2 \left( \frac{\operatorname{arcsech}(cx)cx}{e} + \frac{dc^2 \left( \sum_{-R1=\text{RootOf}(c^2d\_Z^4+(2c^2d+4e)\_Z^2+c^2d)} \frac{(-R1^2c^2d+4e)}{\dots} \right)}{\dots} \right)$
default	$\frac{ac^3x}{e} - \frac{ac^3d \arctan\left(\frac{xe}{\sqrt{de}}\right)}{e\sqrt{de}} + bc^2 \left( \frac{\operatorname{arcsech}(cx)cx}{e} + \frac{dc^2 \left( \sum_{-R1=\text{RootOf}(c^2d\_Z^4+(2c^2d+4e)\_Z^2+c^2d)} \frac{(-R1^2c^2d+4e)}{\dots} \right)}{\dots} \right)$

input

```
int(x^2*(a+b*arcsech(c*x))/(e*x^2+d), x, method=_RETURNVERBOSE)
```

output

```
a/e*x-a/e*d/(d*e)^(1/2)*arctan(x*e/(d*e)^(1/2))+b*arcsech(c*x)/e*x-2*b/c/e
*arctan(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))-1/8*b*c/e^2*d*sum((_R1^2*c
^2*d+c^2*d+4*e)/_R1/(_R1^2*c^2*d+c^2*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-
1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)+dilog((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+
1/c/x)^(1/2))/_R1)),_R1=RootOf(c^2*d*_Z^4+(2*c^2*d+4*e)*_Z^2+c^2*d))+1/8*b
*c/e^2*d*sum((_R1^2*c^2*d+4*_R1^2*e+c^2*d)/_R1/(_R1^2*c^2*d+c^2*d+2*e)*(ar
csech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)+dilog((_R1
-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)),_R1=RootOf(c^2*d*_Z^4+(2*c^
2*d+4*e)*_Z^2+c^2*d))
```

**Fricas [F]**

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^2}{ex^2 + d} dx$$

input `integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d),x, algorithm="fricas")`

output `integral((b*x^2*arcsech(c*x) + a*x^2)/(e*x^2 + d), x)`

**Sympy [F]**

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \int \frac{x^2(a + b \operatorname{asech}(cx))}{d + ex^2} dx$$

input `integrate(x**2*(a+b*asech(c*x))/(e*x**2+d),x)`

output `Integral(x**2*(a + b*asech(c*x))/(d + e*x**2), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \text{Exception raised: ValueError}$$

input `integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`



**Giac [F]**

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^2}{ex^2 + d} dx$$

input `integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^2/(e*x^2 + d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \int \frac{x^2(a + b \operatorname{acosh}(\frac{1}{cx}))}{ex^2 + d} dx$$

input `int((x^2*(a + b*acosh(1/(c*x))))/(d + e*x^2),x)`

output `int((x^2*(a + b*acosh(1/(c*x))))/(d + e*x^2), x)`

**Reduce [F]**

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \frac{-\sqrt{e}\sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e}\sqrt{d}}\right) a + \left(\int \frac{a\operatorname{sech}(cx)x^2}{ex^2+d} dx\right) b e^2 + aex}{e^2}$$

input `int(x^2*(a+b*asech(c*x))/(e*x^2+d),x)`

output `( - sqrt(e)*sqrt(d)*atan((e*x)/(sqrt(e)*sqrt(d)))*a + int((asech(c*x)*x**2)/(d + e*x**2),x)*b*e**2 + a*e*x)/e**2`

$$3.110 \quad \int \frac{x \left( a + b \operatorname{sech}^{-1}(cx) \right)}{d + ex^2} dx$$

Optimal result	882
Mathematica [C] (warning: unable to verify)	883
Rubi [A] (verified)	884
Maple [C] (warning: unable to verify)	886
Fricas [F]	887
Sympy [F]	887
Maxima [F]	888
Giac [F]	888
Mupad [F(-1)]	888
Reduce [F]	889

**Optimal result**

Integrand size = 19, antiderivative size = 441

$$\begin{aligned}
\int \frac{x(a + b\operatorname{sech}^{-1}(cx))}{d + ex^2} dx = & \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2e} \\
& + \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2e} \\
& + \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2e} \\
& + \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2e} \\
& - \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + e^{2\operatorname{sech}^{-1}(cx)}\right)}{e} \\
& + \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2e} \\
& + \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2e} \\
& + \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2e} \\
& + \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2e} \\
& - \frac{b \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(cx)}\right)}{2e}
\end{aligned}$$

output

```

1/2*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)
^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/e+1/2*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1
/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/e+
1/2*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)
^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/e+1/2*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1
/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/e-
(a+b*arcsech(c*x))*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)/e+1/2*
b*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)
)-(c^2*d+e)^(1/2)))/e+1/2*b*polylog(2,c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)
*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/e+1/2*b*polylog(2,-c*(-d)^(1/
2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/e+1
/2*b*polylog(2,c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1
/2)+(c^2*d+e)^(1/2)))/e-1/2*b*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)
^(1/2))^2)/e

```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 0.41 (sec) , antiderivative size = 860, normalized size of antiderivative = 1.95

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \text{Too large to display}$$

input

```
Integrate[(x*(a + b*ArcSech[c*x]))/(d + e*x^2),x]
```

output

```

((4*I)*b*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*ArcTanh[(((-I)*
c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])/Sqrt[c^2*d + e]] + (4*I)*b*ArcS
in[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*ArcTanh[((I*c*Sqrt[d] + Sqrt
[e])*Tanh[ArcSech[c*x]/2])/Sqrt[c^2*d + e]] - 2*b*ArcSech[c*x]*Log[1 + E^(
-2*ArcSech[c*x])] + b*ArcSech[c*x]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]))
/(c*Sqrt[d]*E^ArcSech[c*x])] - (2*I)*b*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt
[d])]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSec
h[c*x])] + b*ArcSech[c*x]*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt
[d]*E^ArcSech[c*x])] - (2*I)*b*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]/Sq
rt[2]]*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])
] + b*ArcSech[c*x]*Log[1 - (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^Ar
cSech[c*x])] + (2*I)*b*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*L
og[1 - (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + b*Arc
Sech[c*x]*Log[1 + (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x
])] + (2*I)*b*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 + (I
*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + a*Log[d + e*x^
2] + b*PolyLog[2, -E^(-2*ArcSech[c*x])] - b*PolyLog[2, ((-I)*(-Sqrt[e] + S
qrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - b*PolyLog[2, (I*(-Sqrt[e] +
Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - b*PolyLog[2, ((-I)*(Sqrt[
e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - b*PolyLog[2, (I*(S...

```

### Rubi [A] (verified)

Time = 1.52 (sec) , antiderivative size = 523, normalized size of antiderivative = 1.19, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$ , Rules used = {6857, 6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx \\
 & \quad \downarrow \text{6857} \\
 & - \int \frac{x(a + b \operatorname{arccosh}(\frac{1}{cx}))}{\frac{d}{x^2} + e} d \frac{1}{x} \\
 & \quad \downarrow \text{6374}
 \end{aligned}$$

$$\begin{aligned}
 & - \int \left( \frac{x(a + \operatorname{barccosh}(\frac{1}{cx}))}{e} - \frac{d(a + \operatorname{barccosh}(\frac{1}{cx}))}{e(\frac{d}{x^2} + e)x} \right) d\frac{1}{x} \\
 & \quad \downarrow \text{2009} \\
 & \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( 1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{c^2d + e}} \right)}{2e} + \\
 & \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{c^2d + e}} + 1 \right)}{2e} + \\
 & \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( 1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{c^2d + e} + \sqrt{e}} \right)}{2e} + \\
 & \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{c^2d + e} + \sqrt{e}} + 1 \right)}{2e} - \frac{(a + \operatorname{barccosh}(\frac{1}{cx}))^2}{e} - \\
 & \frac{\log \left( e^{-2\operatorname{arccosh}(\frac{1}{cx})} + 1 \right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{e} + \frac{b \operatorname{PolyLog} \left( 2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{c^2d + e}} \right)}{2e} + \\
 & \frac{b \operatorname{PolyLog} \left( 2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{c^2d + e}} \right)}{2e} + \frac{b \operatorname{PolyLog} \left( 2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{c^2d + e}} \right)}{2e} + \\
 & \frac{b \operatorname{PolyLog} \left( 2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{c^2d + e}} \right)}{2e} + \frac{b \operatorname{PolyLog} \left( 2, -e^{-2\operatorname{arccosh}(\frac{1}{cx})} \right)}{2e}
 \end{aligned}$$

input `Int[(x*(a + b*ArcSech[c*x]))/(d + e*x^2),x]`

output `-((a + b*ArcCosh[1/(c*x)])^2/(b*e)) - ((a + b*ArcCosh[1/(c*x)])*Log[1 + E^(-2*ArcCosh[1/(c*x)])])/e + ((a + b*ArcCosh[1/(c*x)])*Log[1 - (c*sqrt[-d]*E^ArcCosh[1/(c*x)])/(sqrt[e] - sqrt[c^2*d + e])])/(2*e) + ((a + b*ArcCosh[1/(c*x)])*Log[1 + (c*sqrt[-d]*E^ArcCosh[1/(c*x)])/(sqrt[e] - sqrt[c^2*d + e])])/(2*e) + ((a + b*ArcCosh[1/(c*x)])*Log[1 - (c*sqrt[-d]*E^ArcCosh[1/(c*x)])/(sqrt[e] + sqrt[c^2*d + e])])/(2*e) + ((a + b*ArcCosh[1/(c*x)])*Log[1 + (c*sqrt[-d]*E^ArcCosh[1/(c*x)])/(sqrt[e] + sqrt[c^2*d + e])])/(2*e) + (b*PolyLog[2, -E^(-2*ArcCosh[1/(c*x)])])/(2*e) + (b*PolyLog[2, -((c*sqrt[-d]*E^ArcCosh[1/(c*x)])/(sqrt[e] - sqrt[c^2*d + e]))])/(2*e) + (b*PolyLog[2, (c*sqrt[-d]*E^ArcCosh[1/(c*x)])/(sqrt[e] - sqrt[c^2*d + e])])/(2*e) + (b*PolyLog[2, -((c*sqrt[-d]*E^ArcCosh[1/(c*x)])/(sqrt[e] + sqrt[c^2*d + e]))])/(2*e) + (b*PolyLog[2, (c*sqrt[-d]*E^ArcCosh[1/(c*x)])/(sqrt[e] + sqrt[c^2*d + e])])/(2*e)`

**Defintions of rubi rules used**

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6374 Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^n_.*((f_.)*(x_.))^m_.*((d_.) + (e_.)*(x_)^2)^p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

```
rule 6857 Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))^n_.*(x_)^m_.*((d_.) + (e_.)*(x_)^2)^p_.), x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x^(m + 2*(p + 1))), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0] && IntegerQ[m, p]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.03 (sec) , antiderivative size = 506, normalized size of antiderivative = 1.15

method	result
parts	$\frac{a \ln(x^2 e + d)}{2e} - \frac{b \operatorname{arcsech}(cx) \ln\left(1 + i\left(\frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}}\right) \sqrt{1 + \frac{1}{cx}}\right)}{e} - \frac{b \operatorname{arcsech}(cx) \ln\left(1 - i\left(\frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}}\right) \sqrt{1 + \frac{1}{cx}}\right)}{e}$
derivativedivides	$\frac{a c^2 \ln\left(\frac{e c^2 x^2 + c^2 d}{2e}\right) + b c^2}{c^2 d} \left( \frac{\sum_{-R1=\operatorname{RootOf}(c^2 d Z^4 + (2c^2 d + 4e) Z^2 + c^2 d)} \left(-R1^2 + 1\right) \left(\operatorname{arcsech}(cx) \ln\left(\frac{-R1 - \frac{1}{cx}}{\dots}\right)\right)}{4e} \right)$
default	$\frac{a c^2 \ln\left(\frac{e c^2 x^2 + c^2 d}{2e}\right) + b c^2}{c^2 d} \left( \frac{\sum_{-R1=\operatorname{RootOf}(c^2 d Z^4 + (2c^2 d + 4e) Z^2 + c^2 d)} \left(-R1^2 + 1\right) \left(\operatorname{arcsech}(cx) \ln\left(\frac{-R1 - \frac{1}{cx}}{\dots}\right)\right)}{4e} \right)$

input `int(x*(a+b*arcsech(c*x))/(e*x^2+d),x,method=_RETURNVERBOSE)`

output `1/2*a/e*ln(e*x^2+d)-b/e*arcsech(c*x)*ln(1+I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))-b/e*arcsech(c*x)*ln(1-I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))-b/e*dilog(1+I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))-b/e*dilog(1-I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))+1/4*b*c^2*d/e*sum((_R1^2+1)/(_R1^2*c^2*d+c^2*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)+dilog((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)),_R1=RootOf(c^2*d*_Z^4+(2*c^2*d+4*e)*_Z^2+c^2*d))+1/4*b/e*sum((_R1^2*c^2*d+c^2*d+4*e)/(_R1^2*c^2*d+c^2*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)+dilog((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)),_R1=RootOf(c^2*d*_Z^4+(2*c^2*d+4*e)*_Z^2+c^2*d))`

### Fricas [F]

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x}{ex^2 + d} dx$$

input `integrate(x*(a+b*arcsech(c*x))/(e*x^2+d),x, algorithm="fricas")`

output `integral((b*x*arcsech(c*x) + a*x)/(e*x^2 + d), x)`

### Sympy [F]

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \int \frac{x(a + b \operatorname{asech}(cx))}{d + ex^2} dx$$

input `integrate(x*(a+b*asech(c*x))/(e*x**2+d),x)`

output `Integral(x*(a + b*asech(c*x))/(d + e*x**2), x)`



**Maxima [F]**

$$\int \frac{x(a + b\operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x}{ex^2 + d} dx$$

input `integrate(x*(a+b*arcsech(c*x))/(e*x^2+d),x, algorithm="maxima")`

output `b*integrate(x*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(e*x^2 + d), x) + 1/2*a*log(e*x^2 + d)/e`

**Giac [F]**

$$\int \frac{x(a + b\operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x}{ex^2 + d} dx$$

input `integrate(x*(a+b*arcsech(c*x))/(e*x^2+d),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x/(e*x^2 + d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + b\operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \int \frac{x(a + b \operatorname{acosh}(\frac{1}{cx}))}{ex^2 + d} dx$$

input `int((x*(a + b*acosh(1/(c*x))))/(d + e*x^2),x)`

output `int((x*(a + b*acosh(1/(c*x))))/(d + e*x^2), x)`

**Reduce [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \frac{2 \left( \int \frac{a \operatorname{sech}(cx)x}{ex^2+d} dx \right) be + \log(ex^2 + d) a}{2e}$$

input `int(x*(a+b*asech(c*x))/(e*x^2+d),x)`

output `(2*int((asech(c*x)*x)/(d + e*x**2),x)*b*e + log(d + e*x**2)*a)/(2*e)`

### 3.111 $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{d+ex^2} dx$

Optimal result	890
Mathematica [C] (warning: unable to verify)	891
Rubi [A] (verified)	892
Maple [C] (verified)	894
Fricas [F]	895
Sympy [F]	896
Maxima [F(-2)]	896
Giac [F]	897
Mupad [F(-1)]	897
Reduce [F]	897

#### Optimal result

Integrand size = 18, antiderivative size = 469

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{d + ex^2} dx = \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2\sqrt{-d}\sqrt{e}} - \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2\sqrt{-d}\sqrt{e}} + \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2\sqrt{-d}\sqrt{e}} - \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2\sqrt{-d}\sqrt{e}} - \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2\sqrt{-d}\sqrt{e}} + \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2\sqrt{-d}\sqrt{e}} - \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2\sqrt{-d}\sqrt{e}} + \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2\sqrt{-d}\sqrt{e}}$$

output

```

1/2*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)
^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(1/2)/e^(1/2)-1/2*(a+b*arcsech(c*x
))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^
2*d+e)^(1/2)))/(-d)^(1/2)/e^(1/2)+1/2*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)
*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/(-d)^(
1/2)/e^(1/2)-1/2*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(
1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/(-d)^(1/2)/e^(1/2)-1/2*b*
polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-
(c^2*d+e)^(1/2)))/(-d)^(1/2)/e^(1/2)+1/2*b*polylog(2,c*(-d)^(1/2)*(1/c/x+(
-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(1/2)/e^(
1/2)-1/2*b*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)
)/(e^(1/2)+(c^2*d+e)^(1/2)))/(-d)^(1/2)/e^(1/2)+1/2*b*polylog(2,c*(-d)^(1/
2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/(-d
)^(1/2)/e^(1/2)

```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 0.41 (sec) , antiderivative size = 849, normalized size of antiderivative = 1.81

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex^2} dx = \text{Too large to display}$$

input

```
Integrate[(a + b*ArcSech[c*x])/(d + e*x^2),x]
```

output

```
(2*a*ArcTan[(Sqrt[e]*x)/Sqrt[d]] - 4*b*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*ArcTanh[((( -I)*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])/Sqrt[c^2*d + e]] + 4*b*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*ArcTanh[(((I*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])/Sqrt[c^2*d + e]] - I*b*ArcSech[c*x]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - 2*b*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + I*b*ArcSech[c*x]*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + 2*b*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + I*b*ArcSech[c*x]*Log[1 - (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - 2*b*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 - (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - I*b*ArcSech[c*x]*Log[1 + (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + 2*b*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - I*b*PolyLog[2, (( -I)*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + I*b*PolyLog[2, (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + I*b*PolyLog[2, (( -I)*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - I*b*PolyLog[2, (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])])/(2*Sqrt[d]*Sqrt[e])
```

### Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 517, normalized size of antiderivative = 1.10, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {6847, 6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex^2} dx$$

$$\downarrow 6847$$

$$- \int \frac{a + b \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\frac{d}{x^2} + e} d \frac{1}{x}$$

$$\downarrow 6324$$

$$\begin{aligned}
& - \int \left( \frac{a + \operatorname{arccosh}\left(\frac{1}{cx}\right)}{2\sqrt{e}\left(\sqrt{e} - \frac{\sqrt{-d}}{x}\right)} + \frac{a + \operatorname{arccosh}\left(\frac{1}{cx}\right)}{2\sqrt{e}\left(\frac{\sqrt{-d}}{x} + \sqrt{e}\right)} \right) d\frac{1}{x} \\
& \quad \downarrow \text{2009} \\
& \frac{(a + \operatorname{arccosh}\left(\frac{1}{cx}\right)) \log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{c^2d+e}}\right)}{2\sqrt{-d}\sqrt{e}} - \\
& \frac{(a + \operatorname{arccosh}\left(\frac{1}{cx}\right)) \log\left(\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{c^2d+e}} + 1\right)}{2\sqrt{-d}\sqrt{e}} + \\
& \frac{(a + \operatorname{arccosh}\left(\frac{1}{cx}\right)) \log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{c^2d+e} + \sqrt{e}}\right)}{2\sqrt{-d}\sqrt{e}} - \\
& \frac{(a + \operatorname{arccosh}\left(\frac{1}{cx}\right)) \log\left(\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{c^2d+e} + \sqrt{e}} + 1\right)}{2\sqrt{-d}\sqrt{e}} - \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{dc^2+e}}\right)}{2\sqrt{-d}\sqrt{e}} + \\
& \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{dc^2+e}}\right)}{2\sqrt{-d}\sqrt{e}} - \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} + \sqrt{dc^2+e}}\right)}{2\sqrt{-d}\sqrt{e}} + \\
& \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} + \sqrt{dc^2+e}}\right)}{2\sqrt{-d}\sqrt{e}}
\end{aligned}$$

input `Int[(a + b*ArcSech[c*x])/(d + e*x^2), x]`

output `((a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])])/(2*Sqrt[-d]*Sqrt[e]) - ((a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])])/(2*Sqrt[-d]*Sqrt[e]) + ((a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(2*Sqrt[-d]*Sqrt[e]) - ((a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(2*Sqrt[-d]*Sqrt[e]) - (b*PolyLog[2, -((c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e]))])/(2*Sqrt[-d]*Sqrt[e]) + (b*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])])/(2*Sqrt[-d]*Sqrt[e]) - (b*PolyLog[2, -((c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e]))])/(2*Sqrt[-d]*Sqrt[e]) + (b*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(2*Sqrt[-d]*Sqrt[e])`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6324 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((d_) + (e_.)*(x_)^2)^(p_.),  
x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x],  
x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] &&  
(p > 0 || IGtQ[n, 0])`

rule 6847 `Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^n_)*((d_.) + (e_.)*(x_)^2)^(p_.),  
x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x^(2*(p + 1)  
)), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0] && IntegerQ[p  
]`

**Maple [C] (verified)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 14.66 (sec) , antiderivative size = 302, normalized size of antiderivative = 0.64

method	result
parts	$\frac{a \arctan\left(\frac{xe}{\sqrt{de}}\right)}{\sqrt{de}} - \frac{bc \left( \frac{-R1 \left( \operatorname{arcsech}(cx) \ln\left(\frac{-R1 - \frac{1}{cx} - \sqrt{-1 + \frac{1}{cx}}\right)}{-R1} \right)}{-R1^2 c^2 d} \right)}{2}$
derivativedivides	$\frac{ac \arctan\left(\frac{xe}{\sqrt{de}}\right)}{\sqrt{de}} + bc^2 \left( \frac{-R1 \left( \operatorname{arcsech}(cx) \ln\left(\frac{-R1 - \frac{1}{cx} - \sqrt{-1 + \frac{1}{cx}}\right)}{-R1} \right)}{-R1^2 c^2 d} \right) - \frac{-R1 \left( \operatorname{arcsech}(cx) \ln\left(\frac{-R1 - \frac{1}{cx} - \sqrt{-1 + \frac{1}{cx}}\right)}{-R1} \right)}{2}$
default	$\frac{ac \arctan\left(\frac{xe}{\sqrt{de}}\right)}{\sqrt{de}} + bc^2 \left( \frac{-R1 \left( \operatorname{arcsech}(cx) \ln\left(\frac{-R1 - \frac{1}{cx} - \sqrt{-1 + \frac{1}{cx}}\right)}{-R1} \right)}{-R1^2 c^2 d} \right) - \frac{-R1 \left( \operatorname{arcsech}(cx) \ln\left(\frac{-R1 - \frac{1}{cx} - \sqrt{-1 + \frac{1}{cx}}\right)}{-R1} \right)}{2}$

```
input int((a+b*arcsech(c*x))/(e*x^2+d),x,method=_RETURNVERBOSE)
```

```
output a/(d*e)^(1/2)*arctan(xe/(d*e)^(1/2))-1/2*b*c*sum(_R1/(_R1^2*c^2*d+c^2*d+2
*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2))*(1+1/c/x)^(1/2))/_R1)+dilog
og((_R1-1/c/x-(-1+1/c/x)^(1/2))*(1+1/c/x)^(1/2))/_R1)),_R1=RootOf(c^2*d*_Z^
4+(2*c^2*d+4*e)*_Z^2+c^2*d))+1/2*b*c*sum(1/_R1/(_R1^2*c^2*d+c^2*d+2*e)*(ar
csech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2))*(1+1/c/x)^(1/2))/_R1)+dilog((_R1
-1/c/x-(-1+1/c/x)^(1/2))*(1+1/c/x)^(1/2))/_R1)),_R1=RootOf(c^2*d*_Z^4+(2*c^
2*d+4*e)*_Z^2+c^2*d))
```

**Fricas [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex^2} dx = \int \frac{b \operatorname{ar} \operatorname{sech}(cx) + a}{ex^2 + d} dx$$

```
input integrate((a+b*arcsech(c*x))/(e*x^2+d),x, algorithm="fricas")
```



output `integral((b*arcsech(c*x) + a)/(e*x^2 + d), x)`

### Sympy [F]

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex^2} dx = \int \frac{a + b \operatorname{asech}(cx)}{d + ex^2} dx$$

input `integrate((a+b*asech(c*x))/(e*x**2+d),x)`

output `Integral((a + b*asech(c*x))/(d + e*x**2), x)`

### Maxima [F(-2)]

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex^2} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/(e*x^2+d),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex^2} dx = \int \frac{b \operatorname{arsech}(cx) + a}{ex^2 + d} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x^2+d),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/(e*x^2 + d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex^2} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{ex^2 + d} dx$$

input `int((a + b*acosh(1/(c*x)))/(d + e*x^2),x)`

output `int((a + b*acosh(1/(c*x)))/(d + e*x^2), x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{d + ex^2} dx = \frac{\sqrt{e} \sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e} \sqrt{d}}\right) a + \left(\int \frac{a \operatorname{sech}(cx)}{ex^2 + d} dx\right) bde}{de}$$

input `int((a+b*asech(c*x))/(e*x^2+d),x)`

output `(sqrt(e)*sqrt(d)*atan((e*x)/(sqrt(e)*sqrt(d)))*a + int(asech(c*x)/(d + e*x**2),x)*b*d*e)/(d*e)`

$$3.112 \quad \int \frac{a+b\operatorname{sech}^{-1}(cx)}{x(d+ex^2)} dx$$

Optimal result	899
Mathematica [C] (warning: unable to verify)	900
Rubi [A] (verified)	901
Maple [C] (warning: unable to verify)	902
Fricas [F]	903
Sympy [F]	904
Maxima [F]	904
Giac [F]	904
Mupad [F(-1)]	905
Reduce [F]	905

## Optimal result

Integrand size = 21, antiderivative size = 417

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)} dx = \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{2bd} - \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2d} - \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2d} - \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2d} - \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2d} - \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2d} - \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{2d} - \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2d} - \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{2d}$$

output

```
1/2*(a+b*arcsech(c*x))^2/b/d-1/2*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/d-1/2*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/d-1/2*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/d-1/2*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/d-1/2*b*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/d-1/2*b*polylog(2,c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/d-1/2*b*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/d-1/2*b*polylog(2,c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/d
```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 841, normalized size of antiderivative = 2.02

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)} dx = \text{Too large to display}$$

input `Integrate[(a + b*ArcSech[c*x])/(x*(d + e*x^2)),x]`

output

```
-1/2*(b*ArcSech[c*x]^2 + (4*I)*b*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]]/
Sqrt[2]]*ArcTanh[(((-I)*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])/Sqrt[c^
2*d + e]] + (4*I)*b*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*ArcT
anh[(((I)*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])/Sqrt[c^2*d + e]] + b*Ar
cSech[c*x]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*
x])] - (2*I)*b*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 + (
I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + b*ArcSech[c*x
]*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - (
2*I)*b*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 + (I*(-Sqr
t[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + b*ArcSech[c*x]*Log[1
- (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + (2*I)*b*A
rcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 - (I*(Sqrt[e] + Sqr
t[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + b*ArcSech[c*x]*Log[1 + (I*(Sq
rt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + (2*I)*b*ArcSin[Sqr
t[1 + (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] + Sqrt[c^2*d +
e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - 2*a*Log[x] + a*Log[d + e*x^2] - b*Poly
Log[2, ((-I)*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - b
*PolyLog[2, (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] -
b*PolyLog[2, ((-I)*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x]
)] - b*PolyLog[2, (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[...
```

**Rubi [A] (verified)**

Time = 1.30 (sec) , antiderivative size = 469, normalized size of antiderivative = 1.12, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6857, 6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)} dx \\
 & \quad \downarrow \text{6857} \\
 & - \int \frac{a + b \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\left(\frac{d}{x^2} + e\right)x} d \frac{1}{x} \\
 & \quad \downarrow \text{6374} \\
 & - \int \left( \frac{\sqrt{-d}(a + b \operatorname{arccosh}\left(\frac{1}{cx}\right))}{2d\left(\frac{\sqrt{-d}}{x} + \sqrt{e}\right)} - \frac{\sqrt{-d}(a + b \operatorname{arccosh}\left(\frac{1}{cx}\right))}{2d\left(\sqrt{e} - \frac{\sqrt{-d}}{x}\right)} \right) d \frac{1}{x} \\
 & \quad \downarrow \text{2009} \\
 & \frac{(a + b \operatorname{arccosh}\left(\frac{1}{cx}\right)) \log\left(1 - \frac{c\sqrt{-d}e^{\operatorname{arccosh}\left(\frac{1}{cx}\right)}}{\sqrt{e} - \sqrt{c^2d + e}}\right)}{2d} - \\
 & \frac{(a + b \operatorname{arccosh}\left(\frac{1}{cx}\right)) \log\left(\frac{c\sqrt{-d}e^{\operatorname{arccosh}\left(\frac{1}{cx}\right)}}{\sqrt{e} - \sqrt{c^2d + e}} + 1\right)}{2d} - \\
 & \frac{(a + b \operatorname{arccosh}\left(\frac{1}{cx}\right)) \log\left(1 - \frac{c\sqrt{-d}e^{\operatorname{arccosh}\left(\frac{1}{cx}\right)}}{\sqrt{c^2d + e} + \sqrt{e}}\right)}{2d} - \\
 & \frac{(a + b \operatorname{arccosh}\left(\frac{1}{cx}\right)) \log\left(\frac{c\sqrt{-d}e^{\operatorname{arccosh}\left(\frac{1}{cx}\right)}}{\sqrt{c^2d + e} + \sqrt{e}} + 1\right)}{2d} + \frac{(a + b \operatorname{arccosh}\left(\frac{1}{cx}\right))^2}{2d} - \\
 & \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-d}e^{\operatorname{arccosh}\left(\frac{1}{cx}\right)}}{\sqrt{e} - \sqrt{c^2d + e}}\right)}{2d} - \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-d}e^{\operatorname{arccosh}\left(\frac{1}{cx}\right)}}{\sqrt{e} - \sqrt{c^2d + e}}\right)}{2d} - \\
 & \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-d}e^{\operatorname{arccosh}\left(\frac{1}{cx}\right)}}{\sqrt{e} + \sqrt{c^2d + e}}\right)}{2d} - \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-d}e^{\operatorname{arccosh}\left(\frac{1}{cx}\right)}}{\sqrt{e} + \sqrt{c^2d + e}}\right)}{2d}
 \end{aligned}$$

input

```
Int[(a + b*ArcSech[c*x])/(x*(d + e*x^2)),x]
```

output

$$\begin{aligned} & (a + b \operatorname{ArcCosh}[1/(c*x)])^2/(2*b*d) - ((a + b \operatorname{ArcCosh}[1/(c*x)]) * \operatorname{Log}[1 - (c * \\ & \operatorname{Sqrt}[-d] * E^{\operatorname{ArcCosh}[1/(c*x)])} / (\operatorname{Sqrt}[e] - \operatorname{Sqrt}[c^2*d + e])]) / (2*d) - ((a + b \\ & * \operatorname{ArcCosh}[1/(c*x)]) * \operatorname{Log}[1 + (c * \operatorname{Sqrt}[-d] * E^{\operatorname{ArcCosh}[1/(c*x)])} / (\operatorname{Sqrt}[e] - \operatorname{Sqrt} \\ & [c^2*d + e])]) / (2*d) - ((a + b \operatorname{ArcCosh}[1/(c*x)]) * \operatorname{Log}[1 - (c * \operatorname{Sqrt}[-d] * E^{\operatorname{Arc} \\ & \operatorname{Cosh}[1/(c*x)])} / (\operatorname{Sqrt}[e] + \operatorname{Sqrt}[c^2*d + e])]) / (2*d) - ((a + b \operatorname{ArcCosh}[1/(c * \\ & x)]) * \operatorname{Log}[1 + (c * \operatorname{Sqrt}[-d] * E^{\operatorname{ArcCosh}[1/(c*x)])} / (\operatorname{Sqrt}[e] + \operatorname{Sqrt}[c^2*d + e])]) \\ & / (2*d) - (b * \operatorname{PolyLog}[2, -((c * \operatorname{Sqrt}[-d] * E^{\operatorname{ArcCosh}[1/(c*x)])} / (\operatorname{Sqrt}[e] - \operatorname{Sqrt}[c \\ & ^2*d + e])]) / (2*d) - (b * \operatorname{PolyLog}[2, (c * \operatorname{Sqrt}[-d] * E^{\operatorname{ArcCosh}[1/(c*x)])} / (\operatorname{Sqrt}[ \\ & e] - \operatorname{Sqrt}[c^2*d + e])]) / (2*d) - (b * \operatorname{PolyLog}[2, -((c * \operatorname{Sqrt}[-d] * E^{\operatorname{ArcCosh}[1/(c \\ & *x)])} / (\operatorname{Sqrt}[e] + \operatorname{Sqrt}[c^2*d + e])]) / (2*d) - (b * \operatorname{PolyLog}[2, (c * \operatorname{Sqrt}[-d] * E^{\operatorname{A} \\ & rcCosh}[1/(c*x)])} / (\operatorname{Sqrt}[e] + \operatorname{Sqrt}[c^2*d + e])]) / (2*d) \end{aligned}$$

### Defintions of rubi rules used

rule 2009

$$\operatorname{Int}[u_, x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$$

rule 6374

$$\operatorname{Int}[(a_. + \operatorname{ArcCosh}[c_. * (x_.)] * (b_.))^n * ((f_.) * (x_.))^m * ((d_.) + (e_.) * (x_.)^2)^p, x\_Symbol] \rightarrow \operatorname{Int}[\operatorname{ExpandIntegrand}[(a + b * \operatorname{ArcCosh}[c*x])^n, (f*x)^m * (d + e*x^2)^p, x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f\}, x] \&\& \operatorname{NeQ}[c^2*d + e, 0] \&\& \operatorname{IGtQ}[n, 0] \&\& \operatorname{IntegerQ}[p] \&\& \operatorname{IntegerQ}[m]$$

rule 6857

$$\operatorname{Int}[(a_. + \operatorname{ArcSech}[c_. * (x_.)] * (b_.))^n * (x_.)^m * ((d_.) + (e_.) * (x_.)^2)^p, x\_Symbol] \rightarrow -\operatorname{Subst}[\operatorname{Int}[(e + d*x^2)^p * ((a + b * \operatorname{ArcCosh}[x/c])^n / x^{m + 2*(p + 1)})], x, 1/x] /; \operatorname{FreeQ}\{a, b, c, d, e, n\}, x] \&\& \operatorname{IGtQ}[n, 0] \&\& \operatorname{IntegersQ}[m, p]$$

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.42 (sec) , antiderivative size = 2081, normalized size of antiderivative = 4.99

method	result	size
parts	Expression too large to display	2081
derivativedivides	Expression too large to display	2108
default	Expression too large to display	2108

input `int((a+b*arcsech(c*x))/x/(e*x^2+d),x,method=_RETURNVERBOSE)`

output

```
-1/2*a/d*ln(e*x^2+d)+a/d*ln(x)+b*(1/2/d*arcsech(c*x)^2+(-c^2*d*(e*(c^2*d+e))^(1/2)+2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^(1/2)*e)/(c^2*d+e)/c^2/d^2*ln(1-d*c^2*(1/c/x+(-1+1/c/x)^(1/2))*(1+1/c/x)^(1/2))^2/(-c^2*d-2*(e*(c^2*d+e))^(1/2)-2*e)*arcsech(c*x)+1/4*(-c^2*d*(e*(c^2*d+e))^(1/2)+2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^(1/2)*e)/e/(c^2*d+e)/d*ln(1-d*c^2*(1/c/x+(-1+1/c/x)^(1/2))*(1+1/c/x)^(1/2))^2/(-c^2*d-2*(e*(c^2*d+e))^(1/2)-2*e)*arcsech(c*x)-(-c^2*d*(e*(c^2*d+e))^(1/2)+2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^(1/2)*e)/(c^2*d+e)/c^2/d^2*arcsech(c*x)^2+1/2*(-c^2*d*(e*(c^2*d+e))^(1/2)+2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^(1/2)*e)/(c^2*d+e)/c^2/d^2*polylog(2,d*c^2*(1/c/x+(-1+1/c/x)^(1/2))*(1+1/c/x)^(1/2))^2/(-c^2*d-2*(e*(c^2*d+e))^(1/2)-2*e))+1/2*(e*(c^2*d+e))^(1/2)/(c^2*d+e)/d*arcsech(c*x)*ln(1-d*c^2*(1/c/x+(-1+1/c/x)^(1/2))*(1+1/c/x)^(1/2))^2/(-c^2*d+2*(e*(c^2*d+e))^(1/2)-2*e))-(-c^2*d-2*(e*(c^2*d+e))^(1/2)+2*e)/c^4/d^3*e*ln(1-d*c^2*(1/c/x+(-1+1/c/x)^(1/2))*(1+1/c/x)^(1/2))^2/(-c^2*d-2*(e*(c^2*d+e))^(1/2)-2*e)*arcsech(c*x)-(-c^2*d*(e*(c^2*d+e))^(1/2)+2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^(1/2)*e)*e/(c^2*d+e)/d^3/c^4*arcsech(c*x)^2+1/2*(-c^2*d*(e*(c^2*d+e))^(1/2)+2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^(1/2)*e)*e/(c^2*d+e)/d^3/c^4*polylog(2,d*c^2*(1/c/x+(-1+1/c/x)^(1/2))*(1+1/c/x)^(1/2))^2/(-c^2*d-2*(e*(c^2*d+e))^(1/2)-2*e))+1/4*(e*(c^2*d+e))^(1/2)/e/(c^2*d+e)*c^2*arcsech(c*x)*ln(1-d*c^2*(1/c/x+(-1+1/c/x)^(1/2))*(1+1/c/x)^(1/2))^2/(-c^2*d+2*(e*(c^2*d+e))^(1/2)-2*e))-1/4*(-c^2*d*(e*(c^2*d+e))^(1/2)+2*...
```

### Fricas [F]

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)x} dx$$

input `integrate((a+b*arcsech(c*x))/x/(e*x^2+d),x, algorithm="fricas")`

output `integral((b*arcsech(c*x) + a)/(e*x^3 + d*x), x)`



**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)} dx = \int \frac{a + b \operatorname{arsech}(cx)}{x(d + ex^2)} dx$$

input `integrate((a+b*asech(c*x))/x/(e*x**2+d),x)`

output `Integral((a + b*asech(c*x))/(x*(d + e*x**2)), x)`

**Maxima [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)x} dx$$

input `integrate((a+b*arcsech(c*x))/x/(e*x^2+d),x, algorithm="maxima")`

output `-1/2*a*(log(e*x^2 + d)/d - 2*log(x)/d) + b*integrate(log(sqrt(1/(c*x) + 1)  
*sqrt(1/(c*x) - 1) + 1/(c*x))/(e*x^3 + d*x), x)`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)x} dx$$

input `integrate((a+b*arcsech(c*x))/x/(e*x^2+d),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/((e*x^2 + d)*x), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x(ex^2 + d)} dx$$

input `int((a + b*acosh(1/(c*x)))/(x*(d + e*x^2)),x)`

output `int((a + b*acosh(1/(c*x)))/(x*(d + e*x^2)), x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)} dx = \frac{2 \left( \int \frac{a \operatorname{sech}(cx)}{e x^3 + dx} dx \right) bd - \log(ex^2 + d)a + 2 \log(x)a}{2d}$$

input `int((a+b*asech(c*x))/x/(e*x^2+d),x)`

output `(2*int(asech(c*x)/(d*x + e*x**3),x)*b*d - log(d + e*x**2)*a + 2*log(x)*a)/(2*d)`

$$3.113 \quad \int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^2(d+ex^2)} dx$$

Optimal result	907
Mathematica [C] (warning: unable to verify)	908
Rubi [A] (verified)	909
Maple [C] (verified)	912
Fricas [F]	913
Sympy [F]	913
Maxima [F(-2)]	913
Giac [F]	914
Mupad [F(-1)]	914
Reduce [F]	914

## Optimal result

Integrand size = 21, antiderivative size = 523

$$\begin{aligned}
 \int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2(d + ex^2)} dx = & \frac{bc\sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}}{d} - \frac{a}{dx} - \frac{b \operatorname{sech}^{-1}(cx)}{dx} \\
 & + \frac{\sqrt{e}(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-de} \operatorname{sech}^{-1}(cx)}{\sqrt{e} - \sqrt{c^2d + e}}\right)}{2(-d)^{3/2}} \\
 & - \frac{\sqrt{e}(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-de} \operatorname{sech}^{-1}(cx)}{\sqrt{e} - \sqrt{c^2d + e}}\right)}{2(-d)^{3/2}} \\
 & + \frac{\sqrt{e}(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-de} \operatorname{sech}^{-1}(cx)}{\sqrt{e} + \sqrt{c^2d + e}}\right)}{2(-d)^{3/2}} \\
 & - \frac{\sqrt{e}(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-de} \operatorname{sech}^{-1}(cx)}{\sqrt{e} + \sqrt{c^2d + e}}\right)}{2(-d)^{3/2}} \\
 & - \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{sech}^{-1}(cx)}{\sqrt{e} - \sqrt{c^2d + e}}\right)}{2(-d)^{3/2}} \\
 & + \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{sech}^{-1}(cx)}{\sqrt{e} - \sqrt{c^2d + e}}\right)}{2(-d)^{3/2}} \\
 & - \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{sech}^{-1}(cx)}{\sqrt{e} + \sqrt{c^2d + e}}\right)}{2(-d)^{3/2}} \\
 & + \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{sech}^{-1}(cx)}{\sqrt{e} + \sqrt{c^2d + e}}\right)}{2(-d)^{3/2}}
 \end{aligned}$$

output

```

b*c*(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)/d-a/d/x-b*arcsech(c*x)/d/x+1/2*e^(1/2)
)*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(
1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(3/2)-1/2*e^(1/2)*(a+b*arcsech(c*x))
*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*
d+e)^(1/2)))/(-d)^(3/2)+1/2*e^(1/2)*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(
1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/(-d)^(3
/2)-1/2*e^(1/2)*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/
2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/(-d)^(3/2)-1/2*b*e^(1/2)*po
lylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c
^2*d+e)^(1/2)))/(-d)^(3/2)+1/2*b*e^(1/2)*polylog(2,c*(-d)^(1/2)*(1/c/x+(-1
+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(3/2)-1/2*b
*e^(1/2)*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/
(e^(1/2)+(c^2*d+e)^(1/2)))/(-d)^(3/2)+1/2*b*e^(1/2)*polylog(2,c*(-d)^(1/2)
*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/(-d)^(
3/2)

```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 2.03 (sec) , antiderivative size = 933, normalized size of antiderivative = 1.78

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2(d + ex^2)} dx = \text{Too large to display}$$

input

```
Integrate[(a + b*ArcSech[c*x])/(x^2*(d + e*x^2)),x]
```

output

```
(-4*a*Sqrt[d] - 4*a*Sqrt[e]*x*ArcTan[(Sqrt[e]*x)/Sqrt[d]] + b*(4*Sqrt[d]*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x) - 4*Sqrt[d]*ArcSech[c*x] - (2*I)*Sqrt[e]*x*((-4*I)*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*ArcTanh[((I*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])/Sqrt[c^2*d + e]] + ArcSech[c*x]*Log[1 + E^(-2*ArcSech[c*x])] - ArcSech[c*x]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + (2*I)*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - ArcSech[c*x]*Log[1 + (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - (2*I)*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + PolyLog[2, (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + PolyLog[2, ((-I)*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + (2*I)*Sqrt[e]*x*((-4*I)*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*ArcTanh[((-I)*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])/Sqrt[c^2*d + e]] + ArcSech[c*x]*Log[1 + E^(-2*ArcSech[c*x])] - ArcSech[c*x]*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + (2*I)*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - ArcSech[c*x]*Log[1 - (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - (2*I)*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 - (I*(Sqrt[e] + Sqrt[c^2*d + ...
```

### Rubi [A] (verified)

Time = 1.43 (sec) , antiderivative size = 575, normalized size of antiderivative = 1.10, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6857, 6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2(d + ex^2)} dx$$

$$\downarrow 6857$$

$$- \int \frac{a + b \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\left(\frac{d}{x^2} + e\right) x^2} d \frac{1}{x}$$

$$\downarrow 6374$$

$$\begin{aligned}
& - \int \left( \frac{a + \operatorname{barccosh}\left(\frac{1}{cx}\right)}{d} - \frac{e(a + \operatorname{barccosh}\left(\frac{1}{cx}\right))}{d\left(\frac{d}{x^2} + e\right)} \right) d \frac{1}{x} \\
& \quad \downarrow \text{2009} \\
& \frac{\sqrt{e}(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)) \log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{c^2 d + e}}\right)}{2(-d)^{3/2}} - \\
& \frac{\sqrt{e}(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)) \log\left(\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{c^2 d + e}} + 1\right)}{2(-d)^{3/2}} + \\
& \frac{\sqrt{e}(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)) \log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{c^2 d + e} + \sqrt{e}}\right)}{2(-d)^{3/2}} - \\
& \frac{\sqrt{e}(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)) \log\left(\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{c^2 d + e} + \sqrt{e}} + 1\right)}{2(-d)^{3/2}} - \frac{a}{dx} - \\
& \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{2(-d)^{3/2}} + \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{2(-d)^{3/2}} - \\
& \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{2(-d)^{3/2}} + \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{2(-d)^{3/2}} - \frac{\operatorname{barccosh}\left(\frac{1}{cx}\right)}{dx} + \\
& \frac{bc\sqrt{\frac{1}{cx} - 1}\sqrt{\frac{1}{cx} + 1}}{d}
\end{aligned}$$

input

```
Int[(a + b*ArcSech[c*x])/(x^2*(d + e*x^2)),x]
```

output

```
(b*c*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]/d - a/(d*x) - (b*ArcCosh[1/(c*x)
]))/(d*x) + (Sqrt[e]*(a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]*E^ArcCos
h[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])])/(2*(-d)^(3/2)) - (Sqrt[e]*(a + b
*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt
[c^2*d + e])])/(2*(-d)^(3/2)) + (Sqrt[e]*(a + b*ArcCosh[1/(c*x)])*Log[1 -
(c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(2*(-d)^(3/2)
) - (Sqrt[e]*(a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*
x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(2*(-d)^(3/2)) - (b*Sqrt[e]*PolyLog[2,
-((c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e]))])/(2*(-d)^(
3/2)) + (b*Sqrt[e]*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - S
qrt[c^2*d + e])])/(2*(-d)^(3/2)) - (b*Sqrt[e]*PolyLog[2, -((c*Sqrt[-d]*E^A
rcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e]))])/(2*(-d)^(3/2)) + (b*Sqrt[e
]*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])
)/(2*(-d)^(3/2))
```

### Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6374

```
Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((f_)*(x_))^(m_)*((d_) + (e
_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

rule 6857

```
Int[((a_) + ArcSech[(c_)*(x_)]*(b_))^(n_)*(x_)^(m_)*((d_) + (e_)*(x_
)^2)^(p_), x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x
^(m + 2*(p + 1))), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0
] && IntegersQ[m, p]
```



### Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 30.19 (sec) , antiderivative size = 372, normalized size of antiderivative = 0.71

method	result
parts	$-\frac{ae \arctan\left(\frac{xe}{\sqrt{de}}\right)}{d\sqrt{de}} - \frac{a}{dx} + \frac{cb\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}}{d} - \frac{b \operatorname{arcsech}(cx)}{dx} + \frac{cbe \left( \sum_{-R1=\operatorname{RootOf}(c^2d\_Z^4+(2c^2d+4e)\_Z^2)} \right)}{\dots}$
derivativedivides	$c \left( -\frac{ae \arctan\left(\frac{xe}{\sqrt{de}}\right)}{cd\sqrt{de}} - \frac{a}{dcx} + \frac{b\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}}{d} - \frac{b \operatorname{arcsech}(cx)}{dcx} + \frac{be \left( \sum_{-R1=\operatorname{RootOf}(c^2d\_Z^4+(2c^2d+4e)\_Z^2)} \right)}{\dots} \right)$
default	$c \left( -\frac{ae \arctan\left(\frac{xe}{\sqrt{de}}\right)}{cd\sqrt{de}} - \frac{a}{dcx} + \frac{b\sqrt{-\frac{cx-1}{cx}}\sqrt{\frac{cx+1}{cx}}}{d} - \frac{b \operatorname{arcsech}(cx)}{dcx} + \frac{be \left( \sum_{-R1=\operatorname{RootOf}(c^2d\_Z^4+(2c^2d+4e)\_Z^2)} \right)}{\dots} \right)$

```
input int((a+b*arcsech(c*x))/x^2/(e*x^2+d),x,method=_RETURNVERBOSE)
```

```
output -a*e/d/(d*e)^(1/2)*arctan(x*e/(d*e)^(1/2))-a/d/x+c*b/d*(-(c*x-1)/c/x)^(1/2)
*((c*x+1)/c/x)^(1/2)-b*arcsech(c*x)/d/x+1/2*c*b*e/d*sum(_R1/(_R1^2*c^2*d+
c^2*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_
R1)+dilog((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)),_R1=RootOf(c^
2*d*_Z^4+(2*c^2*d+4*e)*_Z^2+c^2*d))-1/2*c*b*e/d*sum(1/_R1/(_R1^2*c^2*d+c^2
*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)
+dilog((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)),_R1=RootOf(c^2*d
*_Z^4+(2*c^2*d+4*e)*_Z^2+c^2*d))
```

**Fricas [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2(d + ex^2)} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)x^2} dx$$

input `integrate((a+b*arcsech(c*x))/x^2/(e*x^2+d),x, algorithm="fricas")`

output `integral((b*arcsech(c*x) + a)/(e*x^4 + d*x^2), x)`

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2(d + ex^2)} dx = \int \frac{a + b \operatorname{asech}(cx)}{x^2(d + ex^2)} dx$$

input `integrate((a+b*asech(c*x))/x**2/(e*x**2+d),x)`

output `Integral((a + b*asech(c*x))/(x**2*(d + e*x**2)), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2(d + ex^2)} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/x^2/(e*x^2+d),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)x^2} dx$$

input `integrate((a+b*arcsech(c*x))/x^2/(e*x^2+d),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/((e*x^2 + d)*x^2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x^2 (ex^2 + d)} dx$$

input `int((a + b*acosh(1/(c*x)))/(x^2*(d + e*x^2)),x)`

output `int((a + b*acosh(1/(c*x)))/(x^2*(d + e*x^2)), x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)} dx = \frac{-\sqrt{e} \sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e} \sqrt{d}}\right) ax + \left(\int \frac{\operatorname{asech}(cx)}{ex^4 + dx^2} dx\right) b d^2 x - ad}{d^2 x}$$

input `int((a+b*asech(c*x))/x^2/(e*x^2+d),x)`

output `( - sqrt(e)*sqrt(d)*atan((e*x)/(sqrt(e)*sqrt(d)))*a*x + int(asech(c*x)/(d*x**2 + e*x**4),x)*b*d**2*x - a*d)/(d**2*x)`

$$3.114 \quad \int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx$$

Optimal result	916
Mathematica [C] (warning: unable to verify)	917
Rubi [A] (verified)	918
Maple [C] (warning: unable to verify)	921
Fricas [F]	922
Sympy [F(-1)]	922
Maxima [F]	923
Giac [F]	923
Mupad [F(-1)]	923
Reduce [F]	924

**Optimal result**

Integrand size = 21, antiderivative size = 611

$$\begin{aligned}
\int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = & -\frac{b\sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}x}{2ce^2} \\
& + \frac{d(a + b\operatorname{sech}^{-1}(cx))}{2e^2(e + \frac{d}{x^2})} + \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{2e^2} \\
& - \frac{bd\sqrt{-1 + \frac{1}{c^2x^2}}\operatorname{arctanh}\left(\frac{\sqrt{c^2d+e}}{c\sqrt{e}\sqrt{-1 + \frac{1}{c^2x^2}}x}\right)}{2e^{5/2}\sqrt{c^2d+e}\sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}} \\
& - \frac{d(a + b\operatorname{sech}^{-1}(cx))\log\left(1 - \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{e^3} \\
& - \frac{d(a + b\operatorname{sech}^{-1}(cx))\log\left(1 + \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{e^3} \\
& - \frac{d(a + b\operatorname{sech}^{-1}(cx))\log\left(1 - \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{e^3} \\
& - \frac{d(a + b\operatorname{sech}^{-1}(cx))\log\left(1 + \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{e^3} \\
& + \frac{2d(a + b\operatorname{sech}^{-1}(cx))\log\left(1 + e^{2\operatorname{sech}^{-1}(cx)}\right)}{e^3} \\
& - \frac{bd\operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{e^3} \\
& - \frac{bd\operatorname{PolyLog}\left(2, \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{e^3} \\
& - \frac{bd\operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{e^3} \\
& - \frac{bd\operatorname{PolyLog}\left(2, \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{e^3} \\
& + \frac{bd\operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(cx)}\right)}{e^3}
\end{aligned}$$

output

```

-1/2*b*(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)*x/c/e^2+1/2*d*(a+b*arcsech(c*x))/e
^2/(e+d/x^2)+1/2*x^2*(a+b*arcsech(c*x))/e^2-1/2*b*d*(-1+1/c^2/x^2)^(1/2)*a
rctanh((c^2*d+e)^(1/2)/c/e^(1/2)/(-1+1/c^2/x^2)^(1/2)/x)/e^(5/2)/(c^2*d+e)
^(1/2)/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)-d*(a+b*arcsech(c*x))*ln(1-c*(-d)^(
1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/e
^3-d*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)
^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/e^3-d*(a+b*arcsech(c*x))*ln(1-c*(-d)^(
1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/e
^3-d*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)
^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/e^3+2*d*(a+b*arcsech(c*x))*ln(1+(1/c/x
+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)/e^3-b*d*polylog(2,-c*(-d)^(1/2)*(1/c
/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/e^3-b*d*po
lylog(2,c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^
2*d+e)^(1/2)))/e^3-b*d*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+
1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/e^3-b*d*polylog(2,c*(-d)^(1/2)*(1
/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/e^3+b*d*
polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)/e^3

```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 3.23 (sec) , antiderivative size = 1278, normalized size of antiderivative = 2.09

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \text{Too large to display}$$

input

```
Integrate[(x^5*(a + b*ArcSech[c*x]))/(d + e*x^2)^2,x]
```

output

```

-1/4*(-2*a*e*x^2 + (2*a*d^2)/(d + e*x^2) + 4*a*d*Log[d + e*x^2] + b*((2*e*
Sqrt[(1 - c*x)/(1 + c*x)]/c^2 + (2*e*x*Sqrt[(1 - c*x)/(1 + c*x)]/c - 2*e
*x^2*ArcSech[c*x] + (d^(3/2)*ArcSech[c*x])/(Sqrt[d] - I*Sqrt[e]*x) + (d^(3
/2)*ArcSech[c*x])/(Sqrt[d] + I*Sqrt[e]*x) + (16*I)*d*ArcSin[Sqrt[1 - (I*Sq
rt[e])/(c*Sqrt[d])]/Sqrt[2]]*ArcTanh[((( -I)*c*Sqrt[d] + Sqrt[e])*Tanh[ArcS
ech[c*x]/2])/Sqrt[c^2*d + e]] + (16*I)*d*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sq
rt[d])]/Sqrt[2]]*ArcTanh[((I*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])/Sq
rt[c^2*d + e]] - 8*d*ArcSech[c*x]*Log[1 + E^(-2*ArcSech[c*x])) + 4*d*ArcSe
ch[c*x]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])
] - (8*I)*d*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 + (I*(
Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])) + 4*d*ArcSech[c*x]
*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])) - (8
*I)*d*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 + (I*(-Sqrt[
e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])) + 4*d*ArcSech[c*x]*Log[
1 - (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])) + (8*I)*d*
ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 - (I*(Sqrt[e] + Sq
rt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])) + 4*d*ArcSech[c*x]*Log[1 + (I*
(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])) + (8*I)*d*ArcSin[
Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] + Sqrt[c^2*
d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])) + 2*d*Log[x] - 2*d*Log[1 + Sqrt[(1...

```

### Rubi [A] (verified)

Time = 1.76 (sec) , antiderivative size = 703, normalized size of antiderivative = 1.15, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6857, 6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx \\
 & \quad \downarrow \text{6857} \\
 & - \int \frac{x^3(a + b\operatorname{arccosh}(\frac{1}{cx}))}{(\frac{d}{x^2} + e)^2} d\frac{1}{x} \\
 & \quad \downarrow \text{6374}
 \end{aligned}$$

$$\begin{aligned}
 & - \int \left( \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) x^3}{e^2} - \frac{2d(a + \operatorname{barccosh}(\frac{1}{cx})) x}{e^3} + \frac{2d^2(a + \operatorname{barccosh}(\frac{1}{cx}))}{e^3 (\frac{d}{x^2} + e) x} + \frac{d^2(a + \operatorname{barccosh}(\frac{1}{cx}))}{e^2 (\frac{d}{x^2} + e)^2 x} \right) dx \\
 & \qquad \qquad \qquad \downarrow \text{2009} \\
 & \frac{d(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( 1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{c^2 d + e}}} \right)}{e^3} - \\
 & \frac{d(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{c^2 d + e}}} + 1 \right)}{e^3} - \\
 & \frac{d(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( 1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{c^2 d + e} + \sqrt{e}} \right)}{e^3} - \\
 & \frac{d(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{c^2 d + e} + \sqrt{e}} + 1 \right)}{e^3} + \frac{2d(a + \operatorname{barccosh}(\frac{1}{cx}))^2}{be^3} + \\
 & \frac{2d \log \left( e^{-2\operatorname{arccosh}(\frac{1}{cx})} + 1 \right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{e^3} + \frac{d(a + \operatorname{barccosh}(\frac{1}{cx}))}{2e^2 (\frac{d}{x^2} + e)} + \\
 & \frac{x^2(a + \operatorname{barccosh}(\frac{1}{cx}))}{2e^2} - \frac{bd \operatorname{PolyLog} \left( 2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{dc^2 + e}}} \right)}{e^3} - \\
 & \frac{bd \operatorname{PolyLog} \left( 2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{dc^2 + e}}} \right)}{e^3} - \frac{bd \operatorname{PolyLog} \left( 2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}} \right)}{e^3} - \\
 & \frac{bd \operatorname{PolyLog} \left( 2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}} \right)}{e^3} - \frac{bd \operatorname{PolyLog} \left( 2, -e^{-2\operatorname{arccosh}(\frac{1}{cx})} \right)}{e^3} - \\
 & \frac{bd \sqrt{\frac{1}{c^2 x^2} - 1} \operatorname{arctanh} \left( \frac{\sqrt{c^2 d + e}}{c\sqrt{ex} \sqrt{\frac{1}{c^2 x^2} - 1}} \right)}{2e^{5/2} \sqrt{\frac{1}{cx} - 1} \sqrt{\frac{1}{cx} + 1} \sqrt{c^2 d + e}} - \frac{bx \sqrt{\frac{1}{cx} - 1} \sqrt{\frac{1}{cx} + 1}}{2ce^2}
 \end{aligned}$$

input

```
Int[(x^5*(a + b*ArcSech[c*x]))/(d + e*x^2)^2,x]
```



output

```

-1/2*(b*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]*x)/(c*e^2) + (d*(a + b*ArcCos
h[1/(c*x)]))/(2*e^2*(e + d/x^2)) + (x^2*(a + b*ArcCosh[1/(c*x)]))/(2*e^2)
+ (2*d*(a + b*ArcCosh[1/(c*x)])^2)/(b*e^3) - (b*d*Sqrt[-1 + 1/(c^2*x^2)]*A
rcTanh[Sqrt[c^2*d + e]/(c*Sqrt[e]*Sqrt[-1 + 1/(c^2*x^2)]*x)]/(2*e^(5/2)*S
qrt[c^2*d + e]*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) + (2*d*(a + b*ArcCosh
[1/(c*x)])*Log[1 + E^(-2*ArcCosh[1/(c*x)])])/e^3 - (d*(a + b*ArcCosh[1/(c*
x)])*Log[1 - (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])])
/e^3 - (d*(a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])
/(Sqrt[e] - Sqrt[c^2*d + e])])/e^3 - (d*(a + b*ArcCosh[1/(c*x)])*Log[1 - (
c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/e^3 - (d*(a +
b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sq
rt[c^2*d + e])])/e^3 - (b*d*PolyLog[2, -E^(-2*ArcCosh[1/(c*x)])])/e^3 - (b
*d*PolyLog[2, -(c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e]
)])/e^3 - (b*d*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt
[c^2*d + e])])/e^3 - (b*d*PolyLog[2, -(c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sq
rt[e] + Sqrt[c^2*d + e])])/e^3 - (b*d*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/
(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/e^3

```

### Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6374

```
Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*((f_.)*(x_))^m_.*((d_.) + (e
_.)*(x_)^2)^p_.], x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

rule 6857

```
Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^n_.*(x_)^m_.*((d_.) + (e_.)*(x_
)^2)^p_.], x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x
^(m + 2*(p + 1))), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0
] && IntegersQ[m, p]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 5.64 (sec) , antiderivative size = 786, normalized size of antiderivative = 1.29

method	result
parts	$\frac{ax^2}{2e^2} - \frac{ad \ln(x^2e+d)}{e^3} - \frac{ad^2}{2e^3(x^2e+d)} + b \left( \frac{c^4 \left( 2 \operatorname{arcsech}(cx)c^4 dx^2 + e \operatorname{arcsech}(cx)c^4 x^4 - c^3 dx \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} - c^3 e x^3 \right)}{2(e c^2 x^2 + c^2 d) e^2} \right)$
derivativedivides	$\frac{ac^6x^2}{2e^2} - \frac{ac^8d^2}{2e^3(e c^2x^2 + c^2d)} - \frac{ac^6d \ln(e c^2x^2 + c^2d)}{e^3} + bc^4 \left( \frac{2 \operatorname{arcsech}(cx)c^4 dx^2 + e \operatorname{arcsech}(cx)c^4 x^4 - c^3 dx \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} - c^3 e}{2(e c^2x^2 + c^2d) e^2} \right)$
default	$\frac{ac^6x^2}{2e^2} - \frac{ac^8d^2}{2e^3(e c^2x^2 + c^2d)} - \frac{ac^6d \ln(e c^2x^2 + c^2d)}{e^3} + bc^4 \left( \frac{2 \operatorname{arcsech}(cx)c^4 dx^2 + e \operatorname{arcsech}(cx)c^4 x^4 - c^3 dx \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} - c^3 e}{2(e c^2x^2 + c^2d) e^2} \right)$

input `int(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^2,x,method=_RETURNVERBOSE)`

output

```

1/2*a*x^2/e^2-a*d/e^3*ln(e*x^2+d)-1/2*a*d^2/e^3/(e*x^2+d)+b/c^6*(1/2*c^4*(
2*arcsech(c*x)*c^4*d*x^2+e*arcsech(c*x)*c^4*x^4-c^3*d*x*((c*x+1)/c/x)^(1/2
))*(-(c*x-1)/c/x)^(1/2)-c^3*e*x^3*((c*x+1)/c/x)^(1/2)*(-(c*x-1)/c/x)^(1/2)+
c^2*d+e*c^2*x^2)/(c^2*e*x^2+c^2*d)/e^2+1/2*(e*(c^2*d+e))^(1/2)/e^3/(c^2*d+
e)*d*c^6*arctanh(1/4*(2*c^2*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2+2
*c^2*d+4*e)/(c^2*d*e+e^2)^(1/2))+2/e^3*d*c^6*arcsech(c*x)*ln(1+I*(1/c/x+(-
1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))+2/e^3*d*c^6*arcsech(c*x)*ln(1-I*(1/c/x+(-
1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))+2/e^3*d*c^6*dilog(1+I*(1/c/x+(-1+1/c/x)^(
1/2)*(1+1/c/x)^(1/2)))+2/e^3*d*c^6*dilog(1-I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/
c/x)^(1/2)))-1/2/e^3*d*c^6*sum((_R1^2*c^2*d+c^2*d+4*e)/(_R1^2*c^2*d+c^2*d+
2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)+di
log((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)),_R1=RootOf(c^2*d*_Z
^4+(2*c^2*d+4*e)*_Z^2+c^2*d))-1/2/e^3*d^2*c^8*sum((_R1^2+1)/(_R1^2*c^2*d+c
^2*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R
1)+dilog((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)),_R1=RootOf(c^2
*d*_Z^4+(2*c^2*d+4*e)*_Z^2+c^2*d))

```

**Fricas [F]**

$$\int \frac{x^5(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^5}{(ex^2 + d)^2} dx$$

input

```
integrate(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="fricas")
```

output

```
integral((b*x^5*arcsech(c*x) + a*x^5)/(e^2*x^4 + 2*d*e*x^2 + d^2), x)
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{x^5(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \text{Timed out}$$

input

```
integrate(x**5*(a+b*asech(c*x))/(e*x**2+d)**2,x)
```

output Timed out

### Maxima [F]

$$\int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^5}{(ex^2 + d)^2} dx$$

input `integrate(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="maxima")`

output `-1/2*a*(d^2/(e^4*x^2 + d*e^3) - x^2/e^2 + 2*d*log(e*x^2 + d)/e^3) + b*integrate(x^5*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(e^2*x^4 + 2*d*e*x^2 + d^2), x)`

### Giac [F]

$$\int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^5}{(ex^2 + d)^2} dx$$

input `integrate(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^5/(e*x^2 + d)^2, x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{x^5(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^2} dx$$

input `int((x^5*(a + b*acosh(1/(c*x))))/(d + e*x^2)^2,x)`

output `int((x^5*(a + b*acosh(1/(c*x))))/(d + e*x^2)^2, x)`

### Reduce [F]

$$\int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx$$

$$= \frac{2\left(\int \frac{\operatorname{asech}(cx)x^5}{e^2x^4 + 2dex^2 + d^2} dx\right) bde^3 + 2\left(\int \frac{\operatorname{asech}(cx)x^5}{e^2x^4 + 2dex^2 + d^2} dx\right) be^4x^2 - 2\log(ex^2 + d)ad^2 - 2\log(ex^2 + d)ade x^2}{2e^3(ex^2 + d)}$$

input `int(x^5*(a+b*asech(c*x))/(e*x^2+d)^2,x)`

output `(2*int((asech(c*x)*x**5)/(d**2 + 2*d*e*x**2 + e**2*x**4),x)*b*d*e**3 + 2*int((asech(c*x)*x**5)/(d**2 + 2*d*e*x**2 + e**2*x**4),x)*b*e**4*x**2 - 2*log(d + e*x**2)*a*d**2 - 2*log(d + e*x**2)*a*d*e*x**2 + 2*a*d*e*x**2 + a*e**2*x**4)/(2*e**3*(d + e*x**2))`

$$3.115 \quad \int \frac{x^3 (a+b \operatorname{sech}^{-1}(cx))}{(d+ex^2)^2} dx$$

Optimal result	926
Mathematica [C] (warning: unable to verify)	927
Rubi [A] (verified)	928
Maple [C] (warning: unable to verify)	931
Fricas [F]	932
Sympy [F]	932
Maxima [F]	933
Giac [F]	933
Mupad [F(-1)]	933
Reduce [F]	934

**Optimal result**

Integrand size = 21, antiderivative size = 562

$$\begin{aligned}
\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = & -\frac{a + b\operatorname{sech}^{-1}(cx)}{2e\left(e + \frac{d}{x^2}\right)} + \frac{b\sqrt{-1 + \frac{1}{c^2x^2}} \operatorname{arctanh}\left(\frac{\sqrt{c^2d+e}}{c\sqrt{e}\sqrt{-1 + \frac{1}{c^2x^2}}}\right)}{2e^{3/2}\sqrt{c^2d+e}\sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}} \\
& + \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2e^2} \\
& + \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2e^2} \\
& + \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2e^2} \\
& + \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2e^2} \\
& - \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + e^{2\operatorname{sech}^{-1}(cx)}\right)}{e^2} \\
& + \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2e^2} \\
& + \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2e^2} \\
& + \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2e^2} \\
& + \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2e^2} \\
& - \frac{b \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(cx)}\right)}{2e^2}
\end{aligned}$$

output

```

-1/2*(a+b*arcsech(c*x))/e/(e+d/x^2)+1/2*b*(-1+1/c^2/x^2)^(1/2)*arctanh((c^
2*d+e)^(1/2)/c/e^(1/2)/(-1+1/c^2/x^2)^(1/2)/x)/e^(3/2)/(c^2*d+e)^(1/2)/(-1
+1/c/x)^(1/2)/(1+1/c/x)^(1/2)+1/2*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/
c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)-(c^2*d+e)^(1/2))/e^2+1/2*(
a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2
)))/(e^(1/2)-(c^2*d+e)^(1/2))/e^2+1/2*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)
*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(1/2))/e^2+1
/2*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(
1/2)))/(e^(1/2)+(c^2*d+e)^(1/2))/e^2-(a+b*arcsech(c*x))*ln(1+(1/c/x+(-1+1
/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)/e^2+1/2*b*polylog(2,-c*(-d)^(1/2)*(1/c/x+(
-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)-(c^2*d+e)^(1/2))/e^2+1/2*b*poly
log(2,c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)-(c^2*
d+e)^(1/2))/e^2+1/2*b*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+
1/c/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(1/2))/e^2+1/2*b*polylog(2,c*(-d)^(1/2)*
(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(1/2))/e^2-1/
2*b*polylog(2,-(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)/e^2

```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 0.93 (sec) , antiderivative size = 1208, normalized size of antiderivative = 2.15

$$\int \frac{x^3 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \text{Too large to display}$$

input

```
Integrate[(x^3*(a + b*ArcSech[c*x]))/(d + e*x^2)^2,x]
```



output

```

((2*a*d)/(d + e*x^2) + (b*Sqrt[d]*ArcSech[c*x])/(Sqrt[d] - I*Sqrt[e]*x) +
(b*Sqrt[d]*ArcSech[c*x])/(Sqrt[d] + I*Sqrt[e]*x) + (8*I)*b*ArcSin[Sqrt[1 -
(I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*ArcTanh[(((-I)*c*Sqrt[d] + Sqrt[e])*Tan
h[ArcSech[c*x]/2])/Sqrt[c^2*d + e]] + (8*I)*b*ArcSin[Sqrt[1 + (I*Sqrt[e])/(
c*Sqrt[d])]/Sqrt[2]]*ArcTanh[((I*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2
])/Sqrt[c^2*d + e]] - 4*b*ArcSech[c*x]*Log[1 + E^(-2*ArcSech[c*x])] + 2*b*
ArcSech[c*x]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[
c*x])] - (4*I)*b*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 +
(I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + 2*b*ArcSech
[c*x]*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])]
- (4*I)*b*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 + (I*(-
Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + 2*b*ArcSech[c*x]
*Log[1 - (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + (4*
I)*b*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 - (I*(Sqrt[e]
+ Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + 2*b*ArcSech[c*x]*Log[1
+ (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + (4*I)*b*Ar
cSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] + Sqrt
[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + 2*b*Log[x] + 2*a*Log[d + e*x^2
] - 2*b*Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]]
+ (b*Sqrt[e]*Log[((2*I)*Sqrt[e]*(Sqrt[d]*Sqrt[(1 - c*x)/(1 + c*x)])*(1 ...

```

### Rubi [A] (verified)

Time = 1.76 (sec) , antiderivative size = 648, normalized size of antiderivative = 1.15, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6857, 6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx \\
 & \quad \downarrow \text{6857} \\
 & - \int \frac{x(a + b\operatorname{arccosh}(\frac{1}{cx}))}{(\frac{d}{x^2} + e)^2} d\frac{1}{x} \\
 & \quad \downarrow \text{6374}
 \end{aligned}$$

$$\begin{aligned}
 & - \int \left( \frac{x(a + \operatorname{barccosh}(\frac{1}{cx}))}{e^2} - \frac{d(a + \operatorname{barccosh}(\frac{1}{cx}))}{e^2 (\frac{d}{x^2} + e)x} - \frac{d(a + \operatorname{barccosh}(\frac{1}{cx}))}{e (\frac{d}{x^2} + e)^2 x} \right) d\frac{1}{x} \\
 & \quad \downarrow \text{2009} \\
 & \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( 1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{c^2 d + e}}} \right)}{2e^2} + \\
 & \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{c^2 d + e}}} + 1 \right)}{2e^2} + \\
 & \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( 1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{c^2 d + e} + \sqrt{e}} \right)}{2e^2} + \\
 & \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{c^2 d + e} + \sqrt{e}} + 1 \right)}{2e^2} - \frac{a + \operatorname{barccosh}(\frac{1}{cx})}{2e (\frac{d}{x^2} + e)} - \frac{(a + \operatorname{barccosh}(\frac{1}{cx}))^2}{be^2} - \\
 & \frac{\log \left( e^{-2\operatorname{arccosh}(\frac{1}{cx})} + 1 \right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{e^2} + \frac{b \operatorname{PolyLog} \left( 2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{c^2 d + e}}} \right)}{2e^2} + \\
 & \frac{b \operatorname{PolyLog} \left( 2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{dc^2 + e}}} \right)}{2e^2} + \frac{b \operatorname{PolyLog} \left( 2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}} \right)}{2e^2} + \\
 & \frac{b \operatorname{PolyLog} \left( 2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}} \right)}{2e^2} + \frac{b \operatorname{PolyLog} \left( 2, -e^{-2\operatorname{arccosh}(\frac{1}{cx})} \right)}{2e^2} + \\
 & \frac{b\sqrt{\frac{1}{c^2 x^2}} - 1 \operatorname{arctanh} \left( \frac{\sqrt{c^2 d + e}}{c\sqrt{ex} \sqrt{\frac{1}{c^2 x^2} - 1}} \right)}{2e^{3/2} \sqrt{\frac{1}{cx}} - 1 \sqrt{\frac{1}{cx}} + 1 \sqrt{c^2 d + e}}
 \end{aligned}$$

input `Int[(x^3*(a + b*ArcSech[c*x]))/(d + e*x^2)^2,x]`

output

$$\begin{aligned}
& -1/2*(a + b*\text{ArcCosh}[1/(c*x)])/(e*(e + d/x^2)) - (a + b*\text{ArcCosh}[1/(c*x)])^2 \\
& / (b*e^2) + (b*\text{Sqrt}[-1 + 1/(c^2*x^2)]*\text{ArcTanh}[\text{Sqrt}[c^2*d + e]/(c*\text{Sqrt}[e]*\text{Sqrt} \\
& [-1 + 1/(c^2*x^2)]*x)]/(2*e^{3/2}*\text{Sqrt}[c^2*d + e]*\text{Sqrt}[-1 + 1/(c*x)]*\text{Sqrt} \\
& [1 + 1/(c*x)]) - ((a + b*\text{ArcCosh}[1/(c*x)])*\text{Log}[1 + E^{(-2*\text{ArcCosh}[1/(c*x) \\
& ])]])/e^2 + ((a + b*\text{ArcCosh}[1/(c*x)])*\text{Log}[1 - (c*\text{Sqrt}[-d]*E^{\text{ArcCosh}[1/(c*x) \\
& ]})]/(\text{Sqrt}[e] - \text{Sqrt}[c^2*d + e]))/(2*e^2) + ((a + b*\text{ArcCosh}[1/(c*x)])*\text{Log}[1 \\
& + (c*\text{Sqrt}[-d]*E^{\text{ArcCosh}[1/(c*x)])/(\text{Sqrt}[e] - \text{Sqrt}[c^2*d + e]))/(2*e^2) + \\
& ((a + b*\text{ArcCosh}[1/(c*x)])*\text{Log}[1 - (c*\text{Sqrt}[-d]*E^{\text{ArcCosh}[1/(c*x)])/(\text{Sqrt}[e] \\
& + \text{Sqrt}[c^2*d + e]))/(2*e^2) + ((a + b*\text{ArcCosh}[1/(c*x)])*\text{Log}[1 + (c*\text{Sqrt} \\
& [-d]*E^{\text{ArcCosh}[1/(c*x)])/(\text{Sqrt}[e] + \text{Sqrt}[c^2*d + e]))/(2*e^2) + (b*\text{PolyLo} \\
& g[2, -E^{(-2*\text{ArcCosh}[1/(c*x)])}]/(2*e^2) + (b*\text{PolyLog}[2, -(c*\text{Sqrt}[-d]*E^{\text{Ar} \\
& c\text{Cosh}[1/(c*x)])/(\text{Sqrt}[e] - \text{Sqrt}[c^2*d + e]))]/(2*e^2) + (b*\text{PolyLog}[2, (c* \\
& \text{Sqrt}[-d]*E^{\text{ArcCosh}[1/(c*x)])/(\text{Sqrt}[e] - \text{Sqrt}[c^2*d + e]))]/(2*e^2) + (b*\text{Po} \\
& ly\text{Log}[2, -(c*\text{Sqrt}[-d]*E^{\text{ArcCosh}[1/(c*x)])/(\text{Sqrt}[e] + \text{Sqrt}[c^2*d + e]))]/ \\
& (2*e^2) + (b*\text{PolyLog}[2, (c*\text{Sqrt}[-d]*E^{\text{ArcCosh}[1/(c*x)])/(\text{Sqrt}[e] + \text{Sqrt}[c^ \\
& 2*d + e]))]/(2*e^2)
\end{aligned}$$

### Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x\_Symbol] \text{ :> } \text{Simp}[\text{IntSum}[u, x], x] \text{ /; } \text{SumQ}[u]$$

rule 6374

$$\begin{aligned}
& \text{Int}[(a + \text{ArcCosh}[c*x])*(b + (f*x)^m)^n*(d + e*x^2)^p, x\_Symbol] \text{ :> } \text{Int}[\text{ExpandIntegrand}[(a + b*\text{ArcCosh}[c*x])^n, \\
& (f*x)^m*(d + e*x^2)^p, x], x] \text{ /; } \text{FreeQ}\{a, b, c, d, e, f\}, x \text{ \&\& } \text{NeQ}[c^2*d \\
& + e, 0] \text{ \&\& } \text{IGtQ}[n, 0] \text{ \&\& } \text{IntegerQ}[p] \text{ \&\& } \text{IntegerQ}[m]
\end{aligned}$$

rule 6857

$$\begin{aligned}
& \text{Int}[(a + \text{ArcSech}[c*x])*(b + (f*x)^m)^n*(d + e*x^2)^p, x\_Symbol] \text{ :> } -\text{Subst}[\text{Int}[(e + d*x^2)^p*(a + b*\text{ArcCosh}[x/c])^n/x \\
& ^{(m + 2*(p + 1))}, x], x, 1/x] \text{ /; } \text{FreeQ}\{a, b, c, d, e, n\}, x \text{ \&\& } \text{IGtQ}[n, 0] \\
& \text{ \&\& } \text{IntegersQ}[m, p]
\end{aligned}$$

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.32 (sec) , antiderivative size = 644, normalized size of antiderivative = 1.15

method	result
parts	$\frac{a \ln(x^2 e + d)}{2e^2} + \frac{ad}{2e^2(x^2 e + d)} - \frac{b c^2 x^2 \operatorname{arcsech}(cx)}{2(e c^2 x^2 + c^2 d)e} - \frac{b \sqrt{e(c^2 d + e)} \operatorname{arctanh}\left(\frac{2c^2 d \left(\frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}\right)^2 + 2c^2 d + 4}{4\sqrt{c^2 d e + e^2}}\right)}{2e^2(c^2 d + e)}$
derivativedivides	$\frac{a c^6 d}{2e^2(e c^2 x^2 + c^2 d)} + \frac{a c^4 \ln(e c^2 x^2 + c^2 d)}{2e^2} + b c^4 \left( -\frac{c^2 x^2 \operatorname{arcsech}(cx)}{2(e c^2 x^2 + c^2 d)e} - \frac{\sqrt{e(c^2 d + e)} \operatorname{arctanh}\left(\frac{2c^2 d \left(\frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}\right)^2 + 2c^2 d + 4}{4\sqrt{c^2 d e + e^2}}\right)}{2e^2(c^2 d + e)} \right)$
default	$\frac{a c^6 d}{2e^2(e c^2 x^2 + c^2 d)} + \frac{a c^4 \ln(e c^2 x^2 + c^2 d)}{2e^2} + b c^4 \left( -\frac{c^2 x^2 \operatorname{arcsech}(cx)}{2(e c^2 x^2 + c^2 d)e} - \frac{\sqrt{e(c^2 d + e)} \operatorname{arctanh}\left(\frac{2c^2 d \left(\frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}\right)^2 + 2c^2 d + 4}{4\sqrt{c^2 d e + e^2}}\right)}{2e^2(c^2 d + e)} \right)$

input `int(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^2,x,method=_RETURNVERBOSE)`

output

```

1/2*a/e^2*ln(e*x^2+d)+1/2*a/e^2*d/(e*x^2+d)-1/2*b*c^2*x^2*arcsech(c*x)/(c^
2*e*x^2+c^2*d)/e-1/2*b*(e*(c^2*d+e))^(1/2)/e^2/(c^2*d+e)*arctanh(1/4*(2*c^
2*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2+2*c^2*d+4*e)/(c^2*d+e+e^2)^(
1/2))-b/e^2*arcsech(c*x)*ln(1+I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))
-b/e^2*arcsech(c*x)*ln(1-I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))-b/e^2
*dilog(1+I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))-b/e^2*dilog(1-I*(1/c/
x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))+1/4*b/e^2*sum((_R1^2*c^2*d+c^2*d+4*e)
/(_R1^2*c^2*d+c^2*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1
/c/x)^(1/2))/_R1)+dilog((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1))
,_R1=RootOf(c^2*d*_Z^4+(2*c^2*d+4*e)*_Z^2+c^2*d))+1/4*b*c^2/e^2*d*sum((_R1
^2+1)/(_R1^2*c^2*d+c^2*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)
*(1+1/c/x)^(1/2))/_R1)+dilog((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/
_R1)),_R1=RootOf(c^2*d*_Z^4+(2*c^2*d+4*e)*_Z^2+c^2*d))

```

**Fricas [F]**

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^3}{(ex^2 + d)^2} dx$$

input

```
integrate(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="fricas")
```

output

```
integral((b*x^3*arcsech(c*x) + a*x^3)/(e^2*x^4 + 2*d*e*x^2 + d^2), x)
```

**Sympy [F]**

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{x^3(a + b \operatorname{asech}(cx))}{(d + ex^2)^2} dx$$

input

```
integrate(x**3*(a+b*asech(c*x))/(e*x**2+d)**2,x)
```

output

```
Integral(x**3*(a + b*asech(c*x))/(d + e*x**2)**2, x)
```

**Maxima [F]**

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^3}{(ex^2 + d)^2} dx$$

input `integrate(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="maxima")`

output `1/2*a*(d/(e^3*x^2 + d*e^2) + log(e*x^2 + d)/e^2) + b*integrate(x^3*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(e^2*x^4 + 2*d*e*x^2 + d^2), x)`

**Giac [F]**

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^3}{(ex^2 + d)^2} dx$$

input `integrate(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^3/(e*x^2 + d)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{x^3(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^2} dx$$

input `int((x^3*(a + b*acosh(1/(c*x))))/(d + e*x^2)^2,x)`

output `int((x^3*(a + b*acosh(1/(c*x))))/(d + e*x^2)^2, x)`

**Reduce [F]**

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx$$

$$= \frac{2 \left( \int \frac{a \operatorname{sech}(cx) x^3}{e^2 x^4 + 2de x^2 + d^2} dx \right) b d e^2 + 2 \left( \int \frac{a \operatorname{sech}(cx) x^3}{e^2 x^4 + 2de x^2 + d^2} dx \right) b e^3 x^2 + \log(e x^2 + d) a d + \log(e x^2 + d) a e x^2 - a e x^2}{2e^2 (e x^2 + d)}$$

input `int(x^3*(a+b*asech(c*x))/(e*x^2+d)^2,x)`

output `(2*int((asech(c*x)*x**3)/(d**2 + 2*d*e*x**2 + e**2*x**4),x)*b*d*e**2 + 2*int((asech(c*x)*x**3)/(d**2 + 2*d*e*x**2 + e**2*x**4),x)*b*e**3*x**2 + log(d + e*x**2)*a*d + log(d + e*x**2)*a*e*x**2 - a*e*x**2)/(2*e**2*(d + e*x**2))`

**3.116**  $\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^2} dx$

Optimal result	935
Mathematica [C] (verified)	935
Rubi [A] (verified)	936
Maple [B] (verified)	939
Fricas [B] (verification not implemented)	940
Sympy [F]	941
Maxima [F]	941
Giac [F]	941
Mupad [F(-1)]	942
Reduce [F]	942

**Optimal result**

Integrand size = 19, antiderivative size = 147

$$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^2} dx = -\frac{a+b\operatorname{sech}^{-1}(cx)}{2e(d+ex^2)} + \frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\operatorname{arctanh}(\sqrt{1-c^2x^2})}{2de} - \frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{\sqrt{c^2d+e}}\right)}{2d\sqrt{e}\sqrt{c^2d+e}}$$

output

```
-1/2*(a+b*arcsech(c*x))/e/(e*x^2+d)+1/2*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*
arctanh((-c^2*x^2+1)^(1/2))/d/e-1/2*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arct
anh(e^(1/2)*(-c^2*x^2+1)^(1/2)/(c^2*d+e)^(1/2))/d/e^(1/2)/(c^2*d+e)^(1/2)
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.



Time = 0.73 (sec) , antiderivative size = 345, normalized size of antiderivative = 2.35

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx =$$

$$-\frac{2a}{d+ex^2} + \frac{2b \operatorname{sech}^{-1}(cx)}{d+ex^2} + \frac{2b \log(x)}{d} - \frac{2b \log\left(1 + \sqrt{\frac{1-cx}{1+cx}} + cx \sqrt{\frac{1-cx}{1+cx}}\right)}{d} + \frac{b\sqrt{e} \log\left(\frac{4\left(\frac{ide+c^2d^{3/2}\sqrt{ex}}{\sqrt{c^2d+e}(\sqrt{d+i\sqrt{ex}})} + \frac{de\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{-i\sqrt{d\sqrt{e+ex}}}\right)}{b}\right)}{d\sqrt{c^2d+e}} + \dots$$

input

```
Integrate[(x*(a + b*ArcSech[c*x]))/(d + e*x^2)^2,x]
```

output

```
-1/4*((2*a)/(d + e*x^2) + (2*b*ArcSech[c*x])/(d + e*x^2) + (2*b*Log[x])/d - (2*b*Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]])/d + (b*Sqrt[e]*Log[(4*((I*d*e + c^2*d^(3/2)*Sqrt[e]*x)/(Sqrt[c^2*d + e]*(Sqrt[d] + I*Sqrt[e]*x)) + (d*e*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/((-I)*Sqrt[d]*Sqrt[e] + e*x))/b])/(d*Sqrt[c^2*d + e]) + (b*Sqrt[e]*Log[(4*((d*e + I*c^2*d^(3/2)*Sqrt[e]*x)/(Sqrt[c^2*d + e]*(I*Sqrt[d] + Sqrt[e]*x)) + (d*e*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(I*Sqrt[d]*Sqrt[e] + e*x))/b])/(d*Sqrt[c^2*d + e]))/e
```

### Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.86, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$ , Rules used = {6853, 2036, 354, 97, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx$$

$$\downarrow \text{6853}$$

$$-\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{x\sqrt{1-cx}\sqrt{cx+1}(ex^2+d)} dx}{2e} - \frac{a + b \operatorname{sech}^{-1}(cx)}{2e(d + ex^2)}$$

$$\begin{aligned}
& \downarrow 2036 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{x\sqrt{1-c^2x^2}(ex^2+d)} dx}{2e} - \frac{a + b\operatorname{sech}^{-1}(cx)}{2e(d+ex^2)} \\
& \downarrow 354 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{x^2\sqrt{1-c^2x^2}(ex^2+d)} dx^2}{4e} - \frac{a + b\operatorname{sech}^{-1}(cx)}{2e(d+ex^2)} \\
& \downarrow 97 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{1}{x^2\sqrt{1-c^2x^2}} dx^2}{d} - \frac{e \int \frac{1}{\sqrt{1-c^2x^2}(ex^2+d)} dx^2}{d} \right)}{4e} - \frac{a + b\operatorname{sech}^{-1}(cx)}{2e(d+ex^2)} \\
& \downarrow 73 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{2e \int \frac{1}{-\frac{ex^4}{c^2} + d + \frac{e}{c^2}} d\sqrt{1-c^2x^2}}{c^2d} - \frac{2 \int \frac{1}{\frac{1}{c^2} - \frac{x^4}{c^2}} d\sqrt{1-c^2x^2}}{c^2d} \right)}{4e} - \frac{a + b\operatorname{sech}^{-1}(cx)}{2e(d+ex^2)} \\
& \downarrow 221 \\
& \frac{a + b\operatorname{sech}^{-1}(cx)}{2e(d+ex^2)} - \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{2\sqrt{e}\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{\sqrt{c^2d+e}}\right)}{d\sqrt{c^2d+e}} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{1-c^2x^2}}{d}\right)}{d} \right)}{4e}
\end{aligned}$$

input `Int[(x*(a + b*ArcSech[c*x]))/(d + e*x^2)^2,x]`

output `-1/2*(a + b*ArcSech[c*x])/(e*(d + e*x^2)) - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-2*ArcTanh[Sqrt[1 - c^2*x^2]])/d + (2*Sqrt[e]*ArcTanh[(Sqrt[e]*Sqrt[1 - c^2*x^2])/Sqrt[c^2*d + e]])/(d*Sqrt[c^2*d + e]))/(4*e)`

## Definitions of rubi rules used

- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[  
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +  
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt  
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL  
 inearQ[a, b, c, d, m, n, x]`
- rule 97 `Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))),  
 x_] := Simp[b/(b*c - a*d) Int[(e + f*x)^p/(a + b*x), x], x] - Simp[d/(b*c  
 - a*d) Int[(e + f*x)^p/(c + d*x), x], x] /; FreeQ[{a, b, c, d, e, f, p},  
 x] && !IntegerQ[p]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x  
 /Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 354 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_S  
 ymbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x  
 , x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ  
 [(m - 1)/2]`
- rule 2036 `Int[(u_.)*((c_) + (d_.)*(x_)^(n_.))^(q_.)*((a1_) + (b1_.)*(x_)^(non2_.))^(p  
 _)*((a2_) + (b2_.)*(x_)^(non2_.))^(p_.), x_Symbol] := Int[u*(a1*a2 + b1*b2  
 *x^n)^p*(c + d*x^n)^q, x] /; FreeQ[{a1, b1, a2, b2, c, d, n, p, q}, x] && E  
 qq[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && Gt  
 Q[a2, 0]))`
- rule 6853 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.)*(x_)*((d_.) + (e_.)*(x_)^2)^(p_.),  
 x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcSech[c*x])/(2*e*(p + 1))),  
 x] + Simp[b*(Sqrt[1 + c*x]/(2*e*(p + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x  
 ^2)^(p + 1)/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x] /; FreeQ[{a, b, c, d, e  
 , p}, x] && NeQ[p, -1]`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 465 vs. 2(121) = 242.

Time = 4.22 (sec) , antiderivative size = 466, normalized size of antiderivative = 3.17

method	result
parts	$-\frac{a}{2e(x^2e+d)} + b \left( -\frac{c^4 \operatorname{arcsech}(cx)}{2e(e c^2 x^2 + c^2 d)} - \frac{c^3 \sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}}}{2e(e c^2 x^2 + c^2 d)} \left( 2 \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2 x^2 + 1}}\right) \sqrt{\frac{c^2 d + e}{e}} c^2 d - \ln\left(-\frac{2(\sqrt{-c^2 x^2 + 1}}{e} - \frac{c^2 d + e}{e}\right) \right) \right)$
derivativedivides	$-\frac{a c^4}{2e(e c^2 x^2 + c^2 d)} + b c^4 \left( -\frac{\operatorname{arcsech}(cx)}{2e(e c^2 x^2 + c^2 d)} - \frac{\sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}}}{2e(e c^2 x^2 + c^2 d)} \left( 2 \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2 x^2 + 1}}\right) \sqrt{\frac{c^2 d + e}{e}} c^2 d - \ln\left(\frac{-2\sqrt{-c^2 x^2 + 1}}{e} - \frac{c^2 d + e}{e}\right) \right) \right)$
default	$-\frac{a c^4}{2e(e c^2 x^2 + c^2 d)} + b c^4 \left( -\frac{\operatorname{arcsech}(cx)}{2e(e c^2 x^2 + c^2 d)} - \frac{\sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}}}{2e(e c^2 x^2 + c^2 d)} \left( 2 \operatorname{arctanh}\left(\frac{1}{\sqrt{-c^2 x^2 + 1}}\right) \sqrt{\frac{c^2 d + e}{e}} c^2 d - \ln\left(\frac{-2\sqrt{-c^2 x^2 + 1}}{e} - \frac{c^2 d + e}{e}\right) \right) \right)$

```
input int(x*(a+b*arcsech(c*x))/(e*x^2+d)^2,x,method=_RETURNVERBOSE)
```

```
output -1/2*a/e/(e*x^2+d)+b/c^2*(-1/2*c^4/e/(c^2*e*x^2+c^2*d)*arcsech(c*x)-1/4*c^3*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(2*arctanh(1/(-c^2*x^2+1)^(1/2))*((c^2*d+e)/e)^(1/2)*c^2*d-ln(-2*((-c^2*x^2+1)^(1/2))*((c^2*d+e)/e)^(1/2))*e-(-c^2*d*e)^(1/2)*c*x+e)/(-c*e*x+(-c^2*d*e)^(1/2)))*c^2*d-ln(2*((-c^2*x^2+1)^(1/2))*((c^2*d+e)/e)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x+e)/(c*e*x+(-c^2*d*e)^(1/2)))*c^2*d+2*arctanh(1/(-c^2*x^2+1)^(1/2))*((c^2*d+e)/e)^(1/2)*e-ln(-2*((-c^2*x^2+1)^(1/2))*((c^2*d+e)/e)^(1/2)*e-(-c^2*d*e)^(1/2)*c*x+e)/(-c*e*x+(-c^2*d*e)^(1/2)))*e-ln(2*((-c^2*x^2+1)^(1/2))*((c^2*d+e)/e)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x+e)/(c*e*x+(-c^2*d*e)^(1/2)))*e)/(-c^2*x^2+1)^(1/2)/d/(e+(-c^2*d*e)^(1/2))/(-e+(-c^2*d*e)^(1/2))/((c^2*d+e)/e)^(1/2))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 265 vs.  $2(89) = 178$ .

Time = 0.12 (sec) , antiderivative size = 602, normalized size of antiderivative = 4.10

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx$$

$$= \frac{2ac^2d^2 + 2ade - \sqrt{c^2de + e^2}(bex^2 + bd) \log\left(\frac{c^4d^2 + 4c^2de - (c^4de + 2c^2e^2)x^2 + 4(c^3de + ce^2)x\sqrt{-\frac{c^2x^2-1}{c^2x^2}} + 4e^2 + 2(c^2e^2x^2 + 4e^2d)}{ex^2 + d}\right)}{2(c^2d^3e + d^2e^2 + (c^2d^2e^2 + d^3e^3))} + \frac{ac^2d^2 + ade + \sqrt{-c^2de - e^2}(bex^2 + bd) \arctan\left(\frac{\sqrt{-c^2de - e^2}cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}} - \sqrt{-c^2de - e^2}(ex^2 + d)}{(c^2de + e^2)x^2}\right) + (bc^2d^2 + bde^2)}{2(c^2d^3e + d^2e^2 + (c^2d^2e^2 + d^3e^3))}$$

input `integrate(x*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="fricas")`

output `[-1/4*(2*a*c^2*d^2 + 2*a*d*e - sqrt(c^2*d*e + e^2)*(b*e*x^2 + b*d)*log((c^4*d^2 + 4*c^2*d*e - (c^4*d*e + 2*c^2*e^2)*x^2 + 4*(c^3*d*e + c*e^2)*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 4*e^2 + 2*(c^2*e*x^2 - c^2*d - (c^3*d + 2*c*e)*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 2*e)*sqrt(c^2*d*e + e^2))/(e*x^2 + d) + 2*(b*c^2*d^2 + b*d*e + (b*c^2*d*e + b*e^2)*x^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) + 2*(b*c^2*d^2 + b*d*e)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)))/(c^2*d^3*e + d^2*e^2 + (c^2*d^2*e^2 + d*e^3)*x^2), -1/2*(a*c^2*d^2 + a*d*e + sqrt(-c^2*d*e - e^2)*(b*e*x^2 + b*d)*arctan((sqrt(-c^2*d*e - e^2)*c*d*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - sqrt(-c^2*d*e - e^2)*(e*x^2 + d))/((c^2*d*e + e^2)*x^2)) + (b*c^2*d^2 + b*d*e + (b*c^2*d*e + b*e^2)*x^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) + (b*c^2*d^2 + b*d*e)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)))/(c^2*d^3*e + d^2*e^2 + (c^2*d^2*e^2 + d*e^3)*x^2]`

**Sympy [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{x(a + b \operatorname{arsech}(cx))}{(d + ex^2)^2} dx$$

input `integrate(x*(a+b*asech(c*x))/(e*x**2+d)**2,x)`

output `Integral(x*(a + b*asech(c*x))/(d + e*x**2)**2, x)`

**Maxima [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x}{(ex^2 + d)^2} dx$$

input `integrate(x*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="maxima")`

output `1/2*(2*c^2*integrate(1/2*x^3/(c^2*d^2*x^2 + (c^2*d*e*x^2 - d*e)*x^2 + (c^2*d^2*x^2 + (c^2*d*e*x^2 - d*e)*x^2 - d^2)*sqrt(c*x + 1)*sqrt(-c*x + 1) - d^2), x) + (x^2*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1) - x^2*log(c) - x^2*log(x))/(d*e*x^2 + d^2) - 2*integrate(1/2*x/(c^2*d^2*x^2 + (c^2*d*e*x^2 - d*e)*x^2 - d^2), x))*b - 1/2*a/(e^2*x^2 + d*e)`

**Giac [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x}{(ex^2 + d)^2} dx$$

input `integrate(x*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x/(e*x^2 + d)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{x(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^2} dx$$

input `int((x*(a + b*acosh(1/(c*x))))/(d + e*x^2)^2,x)`

output `int((x*(a + b*acosh(1/(c*x))))/(d + e*x^2)^2, x)`

**Reduce [F]**

$$\begin{aligned} & \int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx \\ &= \frac{2 \left( \int \frac{a \operatorname{sech}(cx)x}{e^2 x^4 + 2de x^2 + d^2} dx \right) b d^2 + 2 \left( \int \frac{a \operatorname{sech}(cx)x}{e^2 x^4 + 2de x^2 + d^2} dx \right) b d e x^2 + a x^2}{2d(e x^2 + d)} \end{aligned}$$

input `int(x*(a+b*asech(c*x))/(e*x^2+d)^2,x)`

output `(2*int((asech(c*x)*x)/(d**2 + 2*d*e*x**2 + e**2*x**4),x)*b*d**2 + 2*int((a sech(c*x)*x)/(d**2 + 2*d*e*x**2 + e**2*x**4),x)*b*d*e*x**2 + a*x**2)/(2*d*(d + e*x**2))`

$$3.117 \quad \int \frac{a+b\operatorname{sech}^{-1}(cx)}{x(d+ex^2)^2} dx$$

Optimal result . . . . .	944
Mathematica [C] (warning: unable to verify) . . . . .	945
Rubi [A] (verified) . . . . .	946
Maple [C] (warning: unable to verify) . . . . .	949
Fricas [F] . . . . .	950
Sympy [F] . . . . .	950
Maxima [F(-2)] . . . . .	950
Giac [F] . . . . .	951
Mupad [F(-1)] . . . . .	951
Reduce [F] . . . . .	951



## Optimal result

Integrand size = 21, antiderivative size = 542

$$\begin{aligned}
 \int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^2} dx = & -\frac{e(a + b \operatorname{sech}^{-1}(cx))}{2d^2(e + \frac{d}{x^2})} + \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{2bd^2} \\
 & + \frac{b\sqrt{e}\sqrt{-1 + \frac{1}{c^2x^2}} \operatorname{arctanh}\left(\frac{\sqrt{c^2d+e}}{c\sqrt{e}\sqrt{-1 + \frac{1}{c^2x^2}}}\right)}{2d^2\sqrt{c^2d+e}\sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}} \\
 & - \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2d^2} \\
 & - \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2d^2} \\
 & - \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2d^2} \\
 & - \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2d^2} \\
 & - \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2d^2} \\
 & - \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2d^2} \\
 & - \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2d^2} \\
 & - \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2d^2}
 \end{aligned}$$

output

```

-1/2*e*(a+b*arcsech(c*x))/d^2/(e+d/x^2)+1/2*(a+b*arcsech(c*x))^2/b/d^2+1/2
*b*e^(1/2)*(-1+1/c^2/x^2)^(1/2)*arctanh((c^2*d+e)^(1/2)/c/e^(1/2)/(-1+1/c^
2/x^2)^(1/2)/x)/d^2/(c^2*d+e)^(1/2)/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)-1/2*(
a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2
)))/(e^(1/2)-(c^2*d+e)^(1/2))/d^2-1/2*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)
*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)-(c^2*d+e)^(1/2))/d^2-1
/2*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(
1/2)))/(e^(1/2)+(c^2*d+e)^(1/2))/d^2-1/2*(a+b*arcsech(c*x))*ln(1+c*(-d)^(
1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(1/2))/d
^2-1/2*b*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/
(e^(1/2)-(c^2*d+e)^(1/2))/d^2-1/2*b*polylog(2,c*(-d)^(1/2)*(1/c/x+(-1+1/c
/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)-(c^2*d+e)^(1/2))/d^2-1/2*b*polylog(2,
-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(
1/2))/d^2-1/2*b*polylog(2,c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(
1/2)))/(e^(1/2)+(c^2*d+e)^(1/2))/d^2

```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 1.24 (sec) , antiderivative size = 1189, normalized size of antiderivative = 2.19

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^2} dx = \text{Too large to display}$$

input

```
Integrate[(a + b*ArcSech[c*x])/(x*(d + e*x^2)^2), x]
```

output

```

((2*a*d)/(d + e*x^2) + (b*Sqrt[d]*ArcSech[c*x])/(Sqrt[d] - I*Sqrt[e]*x) +
(b*Sqrt[d]*ArcSech[c*x])/(Sqrt[d] + I*Sqrt[e]*x) - 2*b*ArcSech[c*x]^2 - (8
*I)*b*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*ArcTanh[(((-I)*c*S
qrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])/Sqrt[c^2*d + e]] - (8*I)*b*ArcSin[
Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*ArcTanh[((I*c*Sqrt[d] + Sqrt[e]
)*Tanh[ArcSech[c*x]/2])/Sqrt[c^2*d + e]] - 2*b*ArcSech[c*x]*Log[1 + (I*(Sqr
t[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + (4*I)*b*ArcSin[Sqr
t[1 + (I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d +
e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - 2*b*ArcSech[c*x]*Log[1 + (I*(-Sqrt[e]
+ Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + (4*I)*b*ArcSin[Sqrt[1 -
(I*Sqrt[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*d + e]))
/(c*Sqrt[d]*E^ArcSech[c*x])] - 2*b*ArcSech[c*x]*Log[1 - (I*(Sqrt[e] + Sqrt
[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - (4*I)*b*ArcSin[Sqrt[1 - (I*Sqr
t[e])/(c*Sqrt[d])]/Sqrt[2]]*Log[1 - (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqr
t[d]*E^ArcSech[c*x])] - 2*b*ArcSech[c*x]*Log[1 + (I*(Sqrt[e] + Sqrt[c^2*d
+ e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - (4*I)*b*ArcSin[Sqrt[1 + (I*Sqrt[e])/
(c*Sqrt[d])]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^
ArcSech[c*x])] + 4*a*Log[x] + 2*b*Log[x] - 2*a*Log[d + e*x^2] - 2*b*Log[1
+ Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]] + (b*Sqrt[e]*
Log[((2*I)*Sqrt[e]*(Sqrt[d]*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x) + (Sqrt...

```

### Rubi [A] (verified)

Time = 1.59 (sec) , antiderivative size = 598, normalized size of antiderivative = 1.10, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6857, 6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^2} dx \\
 & \quad \downarrow \text{6857} \\
 & - \int \frac{a + b \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\left(\frac{d}{x^2} + e\right)^2 x^3} d \frac{1}{x} \\
 & \quad \downarrow \text{6374}
 \end{aligned}$$

$$\begin{aligned}
& - \int \left( \frac{a + \operatorname{barccosh}\left(\frac{1}{cx}\right)}{d\left(\frac{d}{x^2} + e\right)x} - \frac{e\left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right)}{d\left(\frac{d}{x^2} + e\right)^2 x} \right) d\frac{1}{x} \\
& \quad \downarrow \text{2009} \\
& \frac{(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)) \log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{c^2 d + e}}\right)}{2d^2} \\
& \frac{(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)) \log\left(\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{c^2 d + e}} + 1\right)}{2d^2} \\
& \frac{(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)) \log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{c^2 d + e} + \sqrt{e}}\right)}{2d^2} \\
& \frac{(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)) \log\left(\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{c^2 d + e} + \sqrt{e}} + 1\right)}{2d^2} - \frac{e\left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right)}{2d^2\left(\frac{d}{x^2} + e\right)} + \\
& \frac{(a + \operatorname{barccosh}\left(\frac{1}{cx}\right))^2}{2bd^2} - \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{2d^2} \\
& \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{2d^2} - \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{2d^2} \\
& \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{2d^2} + \frac{b\sqrt{e}\sqrt{\frac{1}{c^2 x^2} - 1} \operatorname{arctanh}\left(\frac{\sqrt{c^2 d + e}}{c\sqrt{ex}\sqrt{\frac{1}{c^2 x^2} - 1}}\right)}{2d^2\sqrt{\frac{1}{cx} - 1}\sqrt{\frac{1}{cx} + 1}\sqrt{c^2 d + e}}
\end{aligned}$$

input `Int[(a + b*ArcSech[c*x])/(x*(d + e*x^2)^2), x]`

output

```

-1/2*(e*(a + b*ArcCosh[1/(c*x)]))/(d^2*(e + d/x^2)) + (a + b*ArcCosh[1/(c*
x)]^2/(2*b*d^2) + (b*Sqrt[e]*Sqrt[-1 + 1/(c^2*x^2)]*ArcTanh[Sqrt[c^2*d +
e]/(c*Sqrt[e]*Sqrt[-1 + 1/(c^2*x^2)]*x)]/(2*d^2*Sqrt[c^2*d + e]*Sqrt[-1 +
1/(c*x)]*Sqrt[1 + 1/(c*x)]) - ((a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-
d]*E^ArcCosh[1/(c*x)]/(Sqrt[e] - Sqrt[c^2*d + e]))]/(2*d^2) - ((a + b*Arc
Cosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)]/(Sqrt[e] - Sqrt[c^2
*d + e]))]/(2*d^2) - ((a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]*E^ArcCo
sh[1/(c*x)]/(Sqrt[e] + Sqrt[c^2*d + e]))]/(2*d^2) - ((a + b*ArcCosh[1/(c*
x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)]/(Sqrt[e] + Sqrt[c^2*d + e]))]
)/(2*d^2) - (b*PolyLog[2, -((c*Sqrt[-d]*E^ArcCosh[1/(c*x)]/(Sqrt[e] - Sqrt
[c^2*d + e])))]/(2*d^2) - (b*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)]/(S
qrt[e] - Sqrt[c^2*d + e])))]/(2*d^2) - (b*PolyLog[2, -((c*Sqrt[-d]*E^ArcCos
h[1/(c*x)]/(Sqrt[e] + Sqrt[c^2*d + e])))]/(2*d^2) - (b*PolyLog[2, (c*Sqrt
[-d]*E^ArcCosh[1/(c*x)]/(Sqrt[e] + Sqrt[c^2*d + e])))]/(2*d^2)

```

### Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6374

```
Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((f_)*(x_))^(m_)*((d_) + (e
_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

rule 6857

```
Int[((a_) + ArcSech[(c_)*(x_)]*(b_))^(n_)*(x_)^(m_)*((d_) + (e_)*(x
_)^2)^(p_), x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x
^(m + 2*(p + 1))), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0
] && IntegersQ[m, p]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 5.51 (sec) , antiderivative size = 2226, normalized size of antiderivative = 4.11

method	result	size
parts	Expression too large to display	2226
derivativeldivides	Expression too large to display	2275
default	Expression too large to display	2275

input `int((a+b*arcsech(c*x))/x/(e*x^2+d)^2,x,method=_RETURNVERBOSE)`

output

```
-1/2*a/d^2*ln(e*x^2+d)+1/2*a/d/(e*x^2+d)+a/d^2*ln(x)+b*(1/4*(e*(c^2*d+e))^(1/2)/d/e/(c^2*d+e)*c^2*arcsech(c*x)*ln(1-d*c^2*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2/(-c^2*d+2*(e*(c^2*d+e))^(1/2)-2*e))+(-c^2*d*(e*(c^2*d+e))^(1/2)+2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^(1/2)*e)/d^4*e/(c^2*d+e)/c^4*ln(1-d*c^2*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2/(-c^2*d-2*(e*(c^2*d+e))^(1/2)-2*e))*arcsech(c*x)-1/2/d^2*sum((_R1^2*c^2*d+2*c^2*d+4*e)/(_R1^2*c^2*d+c^2*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)+dilog((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)),_R1=RootOf(c^2*d*_Z^4+(2*c^2*d+4*e)*_Z^2+c^2*d))+1/2*arcsech(c*x)^2/d^2-(c^2*d-2*(e*(c^2*d+e))^(1/2)+2*e)/d^4/c^4*e*ln(1-d*c^2*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2/(-c^2*d-2*(e*(c^2*d+e))^(1/2)-2*e))*arcsech(c*x)+(-c^2*d*(e*(c^2*d+e))^(1/2)+2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^(1/2)*e)/(c^2*d+e)/d^3/c^2*ln(1-d*c^2*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2/(-c^2*d-2*(e*(c^2*d+e))^(1/2)-2*e))*arcsech(c*x)+1/2*(-c^2*d*(e*(c^2*d+e))^(1/2)+2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^(1/2)*e)/d^4*e/(c^2*d+e)/c^4*polylog(2,d*c^2*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2/(-c^2*d-2*(e*(c^2*d+e))^(1/2)-2*e))-(-c^2*d*(e*(c^2*d+e))^(1/2)+2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^(1/2)*e)/d^4*e/(c^2*d+e)/c^4*arcsech(c*x)^2+1/8*(e*(c^2*d+e))^(1/2)/d/e/(c^2*d+e)*c^2*polylog(2,d*c^2*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2/(-c^2*d+2*(e*(c^2*d+e))^(1/2)-2*e))-1/4*(e*(c^2*d+e))^(1/2)/d/e/(c^2*d+e)*c^2*arcsech(c*x)^2+1/4...
```

**Fricas [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^2} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^2 x} dx$$

input `integrate((a+b*arcsech(c*x))/x/(e*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*arcsech(c*x) + a)/(e^2*x^5 + 2*d*e*x^3 + d^2*x), x)`

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^2} dx = \int \frac{a + b \operatorname{asech}(cx)}{x(d + ex^2)^2} dx$$

input `integrate((a+b*asech(c*x))/x/(e*x**2+d)**2,x)`

output `Integral((a + b*asech(c*x))/(x*(d + e*x**2)**2), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+b*arcsech(c*x))/x/(e*x^2+d)^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^2} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^2 x} dx$$

input `integrate((a+b*arcsech(c*x))/x/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/((e*x^2 + d)^2*x), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^2} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x(e x^2 + d)^2} dx$$

input `int((a + b*acosh(1/(c*x)))/(x*(d + e*x^2)^2),x)`

output `int((a + b*acosh(1/(c*x)))/(x*(d + e*x^2)^2), x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^2} dx$$

$$= \frac{2 \left( \int \frac{\operatorname{asech}(cx)}{e^2 x^5 + 2de x^3 + d^2 x} dx \right) b d^3 + 2 \left( \int \frac{\operatorname{asech}(cx)}{e^2 x^5 + 2de x^3 + d^2 x} dx \right) b d^2 e x^2 - \log(e x^2 + d) a d - \log(e x^2 + d) a e x^2 + 2 \log(x) a d + 2 \log(x) a e x^2 - a e x^2}{2d^2 (e x^2 + d)}$$

input `int((a+b*asech(c*x))/x/(e*x^2+d)^2,x)`

output `(2*int(asech(c*x)/(d**2*x + 2*d*e*x**3 + e**2*x**5),x)*b*d**3 + 2*int(asech(c*x)/(d**2*x + 2*d*e*x**3 + e**2*x**5),x)*b*d**2*e*x**2 - log(d + e*x**2)*a*d - log(d + e*x**2)*a*e*x**2 + 2*log(x)*a*d + 2*log(x)*a*e*x**2 - a*e*x**2)/(2*d**2*(d + e*x**2))`



$$3.118 \quad \int \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx$$

Optimal result	953
Mathematica [C] (warning: unable to verify)	954
Rubi [A] (verified)	955
Maple [C] (warning: unable to verify)	958
Fricas [F]	959
Sympy [F]	959
Maxima [F(-2)]	959
Giac [F]	960
Mupad [F(-1)]	960
Reduce [F]	960

## Optimal result

Integrand size = 21, antiderivative size = 840

$$\begin{aligned}
 \int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = & -\frac{d(a + b\operatorname{sech}^{-1}(cx))}{4e^2(\sqrt{-d}\sqrt{e} - \frac{d}{x})} + \frac{d(a + b\operatorname{sech}^{-1}(cx))}{4e^2(\sqrt{-d}\sqrt{e} + \frac{d}{x})} \\
 & + \frac{x(a + b\operatorname{sech}^{-1}(cx))}{e^2} \\
 & + \frac{bd \arctan\left(\frac{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{-1 + \frac{1}{cx}}}\right)}{2\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{cd + \sqrt{-d}\sqrt{e}}e^2} \\
 & + \frac{bd \arctan\left(\frac{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{-1 + \frac{1}{cx}}}\right)}{2\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{cd + \sqrt{-d}\sqrt{e}}e^2} \\
 & - \frac{b \arctan\left(\sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}\right)}{ce^2} \\
 & + \frac{3\sqrt{-d}(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{4e^{5/2}} \\
 & - \frac{3\sqrt{-d}(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{4e^{5/2}} \\
 & + \frac{3\sqrt{-d}(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{4e^{5/2}} \\
 & - \frac{3\sqrt{-d}(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{4e^{5/2}} \\
 & - \frac{3b\sqrt{-d} \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{4e^{5/2}} \\
 & + \frac{3b\sqrt{-d} \operatorname{PolyLog}\left(2, \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2d + e}}}\right)}{4e^{5/2}} \\
 & - \frac{3b\sqrt{-d} \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{4e^{5/2}} \\
 & + \frac{3b\sqrt{-d} \operatorname{PolyLog}\left(2, \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2d + e}}}\right)}{4e^{5/2}}
 \end{aligned}$$

output

```

-1/4*d*(a+b*arcsech(c*x))/e^2/((-d)^(1/2)*e^(1/2)-d/x)+1/4*d*(a+b*arcsech(
c*x))/e^2/((-d)^(1/2)*e^(1/2)+d/x)+x*(a+b*arcsech(c*x))/e^2+1/2*b*d*arctan
((c*d-(-d)^(1/2)*e^(1/2))^(1/2)*(1+1/c/x)^(1/2)/(c*d+(-d)^(1/2)*e^(1/2))^(
1/2)/(-1+1/c/x)^(1/2))/(c*d-(-d)^(1/2)*e^(1/2))^(1/2)/(c*d+(-d)^(1/2)*e^(1
/2))^(1/2)/e^2+1/2*b*d*arctan((c*d+(-d)^(1/2)*e^(1/2))^(1/2)*(1+1/c/x)^(1/
2)/(c*d-(-d)^(1/2)*e^(1/2))^(1/2)/(-1+1/c/x)^(1/2))/(c*d-(-d)^(1/2)*e^(1/2
))^(1/2)/(c*d+(-d)^(1/2)*e^(1/2))^(1/2)/e^2-b*arctan((-1+1/c/x)^(1/2)*(1+1
/c/x)^(1/2))/c/e^2+3/4*(-d)^(1/2)*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/
c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/e^(5/2)-3
/4*(-d)^(1/2)*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)
*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/e^(5/2)+3/4*(-d)^(1/2)*(a+b*a
rcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e
^(1/2)+(c^2*d+e)^(1/2)))/e^(5/2)-3/4*(-d)^(1/2)*(a+b*arcsech(c*x))*ln(1+c*
(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/
2)))/e^(5/2)-3/4*b*(-d)^(1/2)*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1
/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/e^(5/2)+3/4*b*(-d)^(1/2)*p
olylog(2,c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c
^2*d+e)^(1/2)))/e^(5/2)-3/4*b*(-d)^(1/2)*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-
1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/e^(5/2)+3/4*b*(-
d)^(1/2)*polylog(2,c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)...

```

### Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.90 (sec) , antiderivative size = 1270, normalized size of antiderivative = 1.51

$$\int \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \text{Too large to display}$$

input

```
Integrate[(x^4*(a + b*ArcSech[c*x]))/(d + e*x^2)^2,x]
```

output

```
(4*a*Sqrt[e]*x + (2*a*d*Sqrt[e]*x)/(d + e*x^2) + 4*b*Sqrt[e]*x*ArcSech[c*x]
] + (b*d*ArcSech[c*x])/((-I)*Sqrt[d] + Sqrt[e]*x) + (b*d*ArcSech[c*x])/(I*
Sqrt[d] + Sqrt[e]*x) - 6*a*Sqrt[d]*ArcTan[(Sqrt[e]*x)/Sqrt[d]] - (8*b*Sqrt
[e]*ArcTan[Tanh[ArcSech[c*x]/2]])/c + 12*b*Sqrt[d]*ArcSin[Sqrt[1 - (I*Sqrt
[e])/(c*Sqrt[d])]]/Sqrt[2]]*ArcTanh[(((I)*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSec
h[c*x]/2])/Sqrt[c^2*d + e]] - 12*b*Sqrt[d]*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*
Sqrt[d])]]/Sqrt[2]]*ArcTanh[(((I)*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])/
Sqrt[c^2*d + e]] + (3*I)*b*Sqrt[d]*ArcSech[c*x]*Log[1 + (I*(Sqrt[e] - Sqrt
[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + 6*b*Sqrt[d]*ArcSin[Sqrt[1 + (I
*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c
*Sqrt[d]*E^ArcSech[c*x])] - (3*I)*b*Sqrt[d]*ArcSech[c*x]*Log[1 + (I*(-Sqrt
[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - 6*b*Sqrt[d]*ArcSin[S
qrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*
d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - (3*I)*b*Sqrt[d]*ArcSech[c*x]*Log[1
- (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + 6*b*Sqrt[d]
]*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 - (I*(Sqrt[e] +
Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + (3*I)*b*Sqrt[d]*ArcSech[c*
x]*Log[1 + (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - 6
*b*Sqrt[d]*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 + (I*(S
qrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - (I*b*Sqrt[d]*S...
```

### Rubi [A] (verified)

Time = 3.00 (sec) , antiderivative size = 900, normalized size of antiderivative = 1.07, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6857, 6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx$$

$$\downarrow 6857$$

$$- \int \frac{x^2(a + b\operatorname{arccosh}(\frac{1}{cx}))}{(\frac{d}{x^2} + e)^2} d\frac{1}{x}$$

$$\downarrow 6374$$

$$\begin{aligned}
 & - \int \left( \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) x^2}{e^2} - \frac{d(a + \operatorname{barccosh}(\frac{1}{cx}))}{e^2 (\frac{d}{x^2} + e)} - \frac{d(a + \operatorname{barccosh}(\frac{1}{cx}))}{e (\frac{d}{x^2} + e)^2} \right) d\frac{1}{x} \\
 & \qquad \qquad \qquad \downarrow \text{2009} \\
 & \frac{x(a + \operatorname{barccosh}(\frac{1}{cx}))}{e^2} + \frac{3\sqrt{-d} \log \left( 1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{dc^2 + e}}} \right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{4e^{5/2}} - \\
 & \frac{3\sqrt{-d} \log \left( \frac{\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx}) c}{\sqrt{e - \sqrt{dc^2 + e}}} + 1 \right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{4e^{5/2}} + \\
 & \frac{3\sqrt{-d} \log \left( 1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}} \right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{4e^{5/2}} - \\
 & \frac{3\sqrt{-d} \log \left( \frac{\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx}) c}{\sqrt{e + \sqrt{dc^2 + e}}} + 1 \right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{4e^{5/2}} - \frac{d(a + \operatorname{barccosh}(\frac{1}{cx}))}{4e^2 (\sqrt{-d}\sqrt{e} - \frac{d}{x})} + \\
 & \frac{d(a + \operatorname{barccosh}(\frac{1}{cx}))}{4e^2 (\frac{d}{x} + \sqrt{-d}\sqrt{e})} + \frac{bd \arctan \left( \frac{\sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{1 + \frac{1}{cx}}}{\sqrt{cd + \sqrt{-d}\sqrt{e}} \sqrt{\frac{1}{cx} - 1}} \right)}{2\sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{cd + \sqrt{-d}\sqrt{e}^2}} + \\
 & \frac{bd \arctan \left( \frac{\sqrt{cd + \sqrt{-d}\sqrt{e}} \sqrt{1 + \frac{1}{cx}}}{\sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{\frac{1}{cx} - 1}} \right)}{2\sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{cd + \sqrt{-d}\sqrt{e}^2}} - \frac{b \arctan \left( \sqrt{\frac{1}{cx} - 1} \sqrt{1 + \frac{1}{cx}} \right)}{ce^2} - \\
 & \frac{3b\sqrt{-d} \operatorname{PolyLog} \left( 2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{dc^2 + e}}} \right)}{4e^{5/2}} + \frac{3b\sqrt{-d} \operatorname{PolyLog} \left( 2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{dc^2 + e}}} \right)}{4e^{5/2}} - \\
 & \frac{3b\sqrt{-d} \operatorname{PolyLog} \left( 2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}} \right)}{4e^{5/2}} + \frac{3b\sqrt{-d} \operatorname{PolyLog} \left( 2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}} \right)}{4e^{5/2}}
 \end{aligned}$$

input `Int[(x^4*(a + b*ArcSech[c*x]))/(d + e*x^2)^2,x]`

output

```

-1/4*(d*(a + b*ArcCosh[1/(c*x)]))/(e^2*(Sqrt[-d]*Sqrt[e] - d/x)) + (d*(a +
b*ArcCosh[1/(c*x)]))/(4*e^2*(Sqrt[-d]*Sqrt[e] + d/x)) + (x*(a + b*ArcCosh
[1/(c*x)]))/e^2 + (b*d*ArcTan[(Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[1 + 1/(c*
x)])/(Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*Sqrt[-1 + 1/(c*x)])])/(2*Sqrt[c*d - Sqr
t[-d]*Sqrt[e]]*Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*e^2) + (b*d*ArcTan[(Sqrt[c*d +
Sqrt[-d]*Sqrt[e]]*Sqrt[1 + 1/(c*x)])/(Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[-
1 + 1/(c*x)])])/(2*Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[c*d + Sqrt[-d]*Sqrt[e
]]*e^2) - (b*ArcTan[Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)])]/(c*e^2) + (3*Sq
rt[-d]*(a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(S
qrt[e] - Sqrt[c^2*d + e])])/(4*e^(5/2)) - (3*Sqrt[-d]*(a + b*ArcCosh[1/(c*
x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])])
/(4*e^(5/2)) + (3*Sqrt[-d]*(a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]*E^
ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(4*e^(5/2)) - (3*Sqrt[-d]*
(a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e]
+ Sqrt[c^2*d + e])])/(4*e^(5/2)) - (3*b*Sqrt[-d]*PolyLog[2, -((c*Sqrt[-d]*
E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e]))])/(4*e^(5/2)) + (3*b*Sqrt
[-d]*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e]
)])/ (4*e^(5/2)) - (3*b*Sqrt[-d]*PolyLog[2, -((c*Sqrt[-d]*E^ArcCosh[1/(c*x)
])/ (Sqrt[e] + Sqrt[c^2*d + e]))])/(4*e^(5/2)) + (3*b*Sqrt[-d]*PolyLog[2, (
c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(4*e^(5/2)...

```

### Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6374

```
Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^ (n_.)*((f_.)*(x_))^(m_.)*((d_) + (e
_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

rule 6857

```
Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^ (n_.)*(x_)^(m_.)*((d_.) + (e_.)*(x
_)^2)^(p_.), x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x
^(m + 2*(p + 1))), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0
] && IntegersQ[m, p]
```



**Fricas [F]**

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^4}{(ex^2 + d)^2} dx$$

input `integrate(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*x^4*arcsech(c*x) + a*x^4)/(e^2*x^4 + 2*d*e*x^2 + d^2), x)`

**Sympy [F]**

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{x^4(a + b \operatorname{asech}(cx))}{(d + ex^2)^2} dx$$

input `integrate(x**4*(a+b*asech(c*x))/(e*x**2+d)**2,x)`

output `Integral(x**4*(a + b*asech(c*x))/(d + e*x**2)**2, x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \text{Exception raised: ValueError}$$

input `integrate(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`



**Giac [F]**

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^4}{(ex^2 + d)^2} dx$$

input `integrate(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^4/(e*x^2 + d)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{x^4(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^2} dx$$

input `int((x^4*(a + b*acosh(1/(c*x))))/(d + e*x^2)^2,x)`

output `int((x^4*(a + b*acosh(1/(c*x))))/(d + e*x^2)^2, x)`

**Reduce [F]**

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx$$

$$= \frac{-3\sqrt{e}\sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e}\sqrt{d}}\right) ad - 3\sqrt{e}\sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e}\sqrt{d}}\right) aex^2 + 2\left(\int \frac{a\operatorname{sech}(cx)x^4}{e^2x^4+2dex^2+d^2} dx\right) bde^3 + 2\left(\int \frac{a\operatorname{sech}(cx)x^4}{e^2x^4+2dex^2+d^2} dx\right)}{2e^3(e^2x^2 + d)}$$

input `int(x^4*(a+b*asech(c*x))/(e*x^2+d)^2,x)`

output

```
( - 3*sqrt(e)*sqrt(d)*atan((e*x)/(sqrt(e)*sqrt(d)))*a*d - 3*sqrt(e)*sqrt(d)
)*atan((e*x)/(sqrt(e)*sqrt(d)))*a*e*x**2 + 2*int((asech(c*x)*x**4)/(d**2 +
2*d*e*x**2 + e**2*x**4),x)*b*d*e**3 + 2*int((asech(c*x)*x**4)/(d**2 + 2*d
*e*x**2 + e**2*x**4),x)*b*e**4*x**2 + 3*a*d*e*x + 2*a*e**2*x**3)/(2*e**3*(
d + e*x**2))
```

$$3.119 \quad \int \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx$$

Optimal result	963
Mathematica [C] (warning: unable to verify)	964
Rubi [A] (verified)	965
Maple [C] (warning: unable to verify)	968
Fricas [F]	969
Sympy [F]	970
Maxima [F(-2)]	970
Giac [F]	970
Mupad [F(-1)]	971
Reduce [F]	971

**Optimal result**

Integrand size = 21, antiderivative size = 786

$$\begin{aligned}
\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx &= \frac{a + b\operatorname{sech}^{-1}(cx)}{4e(\sqrt{-d}\sqrt{e} - \frac{d}{x})} - \frac{a + b\operatorname{sech}^{-1}(cx)}{4e(\sqrt{-d}\sqrt{e} + \frac{d}{x})} \\
&\quad - \frac{b \arctan\left(\frac{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{-1 + \frac{1}{cx}}}\right)}{2\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{cd + \sqrt{-d}\sqrt{e}}} \\
&\quad - \frac{b \arctan\left(\frac{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{-1 + \frac{1}{cx}}}\right)}{2\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{cd + \sqrt{-d}\sqrt{e}}} \\
&\quad + \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e} - \sqrt{c^2d + e}}\right)}{4\sqrt{-de}^{3/2}} \\
&\quad - \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e} - \sqrt{c^2d + e}}\right)}{4\sqrt{-de}^{3/2}} \\
&\quad + \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e} + \sqrt{c^2d + e}}\right)}{4\sqrt{-de}^{3/2}} \\
&\quad - \frac{(a + b\operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e} + \sqrt{c^2d + e}}\right)}{4\sqrt{-de}^{3/2}} \\
&\quad - \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e} - \sqrt{c^2d + e}}\right)}{4\sqrt{-de}^{3/2}} \\
&\quad + \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e} - \sqrt{c^2d + e}}\right)}{4\sqrt{-de}^{3/2}} \\
&\quad - \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e} + \sqrt{c^2d + e}}\right)}{4\sqrt{-de}^{3/2}} \\
&\quad + \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e} + \sqrt{c^2d + e}}\right)}{4\sqrt{-de}^{3/2}}
\end{aligned}$$

output

```

1/4*(a+b*arcsech(c*x))/e/((-d)^(1/2)*e^(1/2)-d/x)-1/4*(a+b*arcsech(c*x))/e
/((-d)^(1/2)*e^(1/2)+d/x)-1/2*b*arctan((c*d-(-d)^(1/2)*e^(1/2))^(1/2)*(1+1
/c/x)^(1/2)/(c*d+(-d)^(1/2)*e^(1/2))^(1/2)/(-1+1/c/x)^(1/2))/(c*d-(-d)^(1
/2)*e^(1/2))^(1/2)/(c*d+(-d)^(1/2)*e^(1/2))^(1/2)/e-1/2*b*arctan((c*d+(-d)^(
1/2)*e^(1/2))^(1/2)*(1+1/c/x)^(1/2)/(c*d-(-d)^(1/2)*e^(1/2))^(1/2)/(-1+1/
c/x)^(1/2))/(c*d-(-d)^(1/2)*e^(1/2))^(1/2)/(c*d+(-d)^(1/2)*e^(1/2))^(1/2)/
e+1/4*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/
x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(1/2)/e^(3/2)-1/4*(a+b*arcsech(c
*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(
c^2*d+e)^(1/2)))/(-d)^(1/2)/e^(3/2)+1/4*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/
2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/(-d
)^(1/2)/e^(3/2)-1/4*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)
^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/(-d)^(1/2)/e^(3/2)-1/4*
b*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2
)-(c^2*d+e)^(1/2)))/(-d)^(1/2)/e^(3/2)+1/4*b*polylog(2,c*(-d)^(1/2)*(1/c/x
+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(1/2)/e
^(3/2)-1/4*b*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/
2))/(e^(1/2)+(c^2*d+e)^(1/2)))/(-d)^(1/2)/e^(3/2)+1/4*b*polylog(2,c*(-d)^(
1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/(-
d)^(1/2)/e^(3/2)

```

### Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 1.10 (sec) , antiderivative size = 1226, normalized size of antiderivative = 1.56

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \text{Too large to display}$$

input

```
Integrate[(x^2*(a + b*ArcSech[c*x]))/(d + e*x^2)^2,x]
```

output

```

((-2*a*Sqrt[e]*x)/(d + e*x^2) + (b*ArcSech[c*x])/(I*Sqrt[d] - Sqrt[e]*x) -
(b*ArcSech[c*x])/(I*Sqrt[d] + Sqrt[e]*x) + (2*a*ArcTan[(Sqrt[e]*x)/Sqrt[d
]])/Sqrt[d] - (4*b*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])])/Sqrt[2]]*ArcTa
nh[(((-I)*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])/Sqrt[c^2*d + e]]/Sqr
t[d] + (4*b*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])])/Sqrt[2]]*ArcTanh[((I*
c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])/Sqrt[c^2*d + e]]/Sqrt[d] - (I*
b*ArcSech[c*x]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSec
h[c*x])])/Sqrt[d] - (2*b*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])])/Sqrt[2]]
*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])])/Sqrt
[d] + (I*b*ArcSech[c*x]*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d
]*E^ArcSech[c*x])])/Sqrt[d] + (2*b*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])
])/Sqrt[2]]*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c
*x])])/Sqrt[d] + (I*b*ArcSech[c*x]*Log[1 - (I*(Sqrt[e] + Sqrt[c^2*d + e]))
/(c*Sqrt[d]*E^ArcSech[c*x])])/Sqrt[d] - (2*b*ArcSin[Sqrt[1 - (I*Sqrt[e])/(
c*Sqrt[d])])/Sqrt[2]]*Log[1 - (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^
ArcSech[c*x])])/Sqrt[d] - (I*b*ArcSech[c*x]*Log[1 + (I*(Sqrt[e] + Sqrt[c^2
*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])])/Sqrt[d] + (2*b*ArcSin[Sqrt[1 + (I*S
qrt[e])/(c*Sqrt[d])])/Sqrt[2]]*Log[1 + (I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*S
qrt[d]*E^ArcSech[c*x])])/Sqrt[d] + (I*b*Sqrt[e]*Log[((2*I)*Sqrt[e]*(Sqrt[d
]*Sqrt[(1 - c*x)/(1 + c*x])*(1 + c*x) + (Sqrt[d]*Sqrt[e] + I*c^2*d*x)/S...

```

### Rubi [A] (verified)

Time = 1.77 (sec) , antiderivative size = 842, normalized size of antiderivative = 1.07, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6857, 6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx \\
 & \quad \downarrow 6857 \\
 & - \int \frac{a + b\operatorname{arcosh}\left(\frac{1}{cx}\right)}{\left(\frac{d}{x^2} + e\right)^2} d\frac{1}{x} \\
 & \quad \downarrow 6324
 \end{aligned}$$

$$\begin{aligned}
& - \int \left( -\frac{d(a + \operatorname{barccosh}(\frac{1}{cx}))}{2e \left(-\frac{d^2}{x^2} - ed\right)} - \frac{d(a + \operatorname{barccosh}(\frac{1}{cx}))}{4e \left(\sqrt{-d}\sqrt{e} - \frac{d}{x}\right)^2} - \frac{d(a + \operatorname{barccosh}(\frac{1}{cx}))}{4e \left(\frac{d}{x} + \sqrt{-d}\sqrt{e}\right)^2} \right) d\frac{1}{x} \\
& \qquad \qquad \qquad \downarrow \text{2009} \\
& \frac{\log \left( 1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{dc^2 + e}} \right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{4\sqrt{-de}^{3/2}} - \\
& \frac{\log \left( \frac{\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{dc^2 + e}} c + 1 \right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{4\sqrt{-de}^{3/2}} + \\
& \frac{\log \left( 1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{dc^2 + e}} \right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{4\sqrt{-de}^{3/2}} - \\
& \frac{\log \left( \frac{\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{dc^2 + e}} c + 1 \right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{4\sqrt{-de}^{3/2}} + \frac{a + \operatorname{barccosh}(\frac{1}{cx})}{4e \left(\sqrt{-d}\sqrt{e} - \frac{d}{x}\right)} - \frac{a + \operatorname{barccosh}(\frac{1}{cx})}{4e \left(\frac{d}{x} + \sqrt{-d}\sqrt{e}\right)} - \\
& \frac{b \arctan \left( \frac{\sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{1 + \frac{1}{cx}}}{\sqrt{cd + \sqrt{-d}\sqrt{e}} \sqrt{\frac{1}{cx} - 1}} \right)}{2\sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{cd + \sqrt{-d}\sqrt{e}}} - \frac{b \arctan \left( \frac{\sqrt{cd + \sqrt{-d}\sqrt{e}} \sqrt{1 + \frac{1}{cx}}}{\sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{\frac{1}{cx} - 1}} \right)}{2\sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{cd + \sqrt{-d}\sqrt{e}}} - \\
& \frac{b \operatorname{PolyLog} \left( 2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{dc^2 + e}} \right)}{4\sqrt{-de}^{3/2}} + \frac{b \operatorname{PolyLog} \left( 2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{dc^2 + e}} \right)}{4\sqrt{-de}^{3/2}} - \\
& \frac{b \operatorname{PolyLog} \left( 2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{dc^2 + e}} \right)}{4\sqrt{-de}^{3/2}} + \frac{b \operatorname{PolyLog} \left( 2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{dc^2 + e}} \right)}{4\sqrt{-de}^{3/2}}
\end{aligned}$$

input `Int[(x^2*(a + b*ArcSech[c*x]))/(d + e*x^2)^2,x]`

output

```
(a + b*ArcCosh[1/(c*x)])/(4*e*(Sqrt[-d]*Sqrt[e] - d/x)) - (a + b*ArcCosh[1/(c*x)])/(4*e*(Sqrt[-d]*Sqrt[e] + d/x)) - (b*ArcTan[(Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[1 + 1/(c*x)])]/(Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*Sqrt[-1 + 1/(c*x)])))/(2*Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*e) - (b*ArcTan[(Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*Sqrt[1 + 1/(c*x)])]/(Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[-1 + 1/(c*x)])))/(2*Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*e) + ((a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])]/(Sqrt[e] - Sqrt[c^2*d + e]))/(4*Sqrt[-d]*e^(3/2)) - ((a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])]/(Sqrt[e] - Sqrt[c^2*d + e]))/(4*Sqrt[-d]*e^(3/2)) + ((a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])]/(Sqrt[e] + Sqrt[c^2*d + e]))/(4*Sqrt[-d]*e^(3/2)) - ((a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])]/(Sqrt[e] + Sqrt[c^2*d + e]))/(4*Sqrt[-d]*e^(3/2)) - (b*PolyLog[2, -(c*Sqrt[-d]*E^ArcCosh[1/(c*x)])]/(Sqrt[e] - Sqrt[c^2*d + e]))/(4*Sqrt[-d]*e^(3/2)) + (b*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])]/(Sqrt[e] - Sqrt[c^2*d + e]))/(4*Sqrt[-d]*e^(3/2)) - (b*PolyLog[2, -(c*Sqrt[-d]*E^ArcCosh[1/(c*x)])]/(Sqrt[e] + Sqrt[c^2*d + e]))/(4*Sqrt[-d]*e^(3/2)) + (b*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])]/(Sqrt[e] + Sqrt[c^2*d + e]))/(4*Sqrt[-d]*e^(3/2))
```

### Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6324

```
Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (p > 0 || IGtQ[n, 0])
```

rule 6857

```
Int[((a_) + ArcSech[(c_)*(x_)]*(b_))^(n_)*(x_)^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x^(m + 2*(p + 1))), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0] && IntegersQ[m, p]
```



### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 30.90 (sec) , antiderivative size = 910, normalized size of antiderivative = 1.16

method	result
parts	$-\frac{ax}{2e(x^2e+d)} + \frac{a \arctan\left(\frac{xe}{\sqrt{de}}\right)}{2e\sqrt{de}} + b \left( -\frac{c^5 \operatorname{arcsech}(cx)x}{2e(e c^2 x^2 + c^2 d)} - \frac{c^4 \left( -R1 = \operatorname{RootOf}(c^2 d \_Z^4 + (2c^2 d + 4e) \_Z^2 + c^2 d) \right)}{2e(e c^2 x^2 + c^2 d)} \right)$
derivativeldivides	$-\frac{a c^5 x}{2e(e c^2 x^2 + c^2 d)} + \frac{a c^3 \arctan\left(\frac{xe}{\sqrt{de}}\right)}{2e\sqrt{de}} + b c^4 \left( -\frac{\operatorname{arcsech}(cx)cx}{2e(e c^2 x^2 + c^2 d)} - \frac{-R1 = \operatorname{RootOf}(c^2 d \_Z^4 + (2c^2 d + 4e) \_Z^2 + c^2 d)}{2e(e c^2 x^2 + c^2 d)} \right)$
default	$-\frac{a c^5 x}{2e(e c^2 x^2 + c^2 d)} + \frac{a c^3 \arctan\left(\frac{xe}{\sqrt{de}}\right)}{2e\sqrt{de}} + b c^4 \left( -\frac{\operatorname{arcsech}(cx)cx}{2e(e c^2 x^2 + c^2 d)} - \frac{-R1 = \operatorname{RootOf}(c^2 d \_Z^4 + (2c^2 d + 4e) \_Z^2 + c^2 d)}{2e(e c^2 x^2 + c^2 d)} \right)$

input `int(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^2,x,method=_RETURNVERBOSE)`

output

```

-1/2*a/e*x/(e*x^2+d)+1/2*a/e/(d*e)^(1/2)*arctan(x*e/(d*e)^(1/2))+b/c^3*(-1
/2*c^5*arcsech(c*x)/e*x/(c^2*e*x^2+c^2*d)-1/4/e*c^4*sum(_R1/(_R1^2*c^2*d+c
^2*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R
1)+dilog((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)),_R1=RootOf(c^2
*d*_Z^4+(2*c^2*d+4*e)*_Z^2+c^2*d))+1/2*(-(c^2*d-2*(e*(c^2*d+e))^(1/2)+2*e)
*d)^(1/2)*(c^2*d+2*(e*(c^2*d+e))^(1/2)+2*e)*arctanh(c*d*(1/c/x+(-1+1/c/x)^(
1/2)*(1+1/c/x)^(1/2))/((-c^2*d+2*(e*(c^2*d+e))^(1/2)-2*e)*d)^(1/2))/c/d^3
/e-1/2*(-(c^2*d-2*(e*(c^2*d+e))^(1/2)+2*e)*d)^(1/2)*(c^2*d*(e*(c^2*d+e))^(
1/2)+2*c^2*d*e+2*e^2+2*(e*(c^2*d+e))^(1/2)*e)*arctanh(c*d*(1/c/x+(-1+1/c/x)
)^(1/2)*(1+1/c/x)^(1/2))/((-c^2*d+2*(e*(c^2*d+e))^(1/2)-2*e)*d)^(1/2))/e/(
c^2*d+e)/d^3/c+1/2*((c^2*d+2*(e*(c^2*d+e))^(1/2)+2*e)*d)^(1/2)*(c^2*d-2*(e
*(c^2*d+e))^(1/2)+2*e)*arctan(c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))
/((c^2*d+2*(e*(c^2*d+e))^(1/2)+2*e)*d)^(1/2))/c/d^3/e-1/2*((c^2*d+2*(e*(c^
2*d+e))^(1/2)+2*e)*d)^(1/2)*(-c^2*d*(e*(c^2*d+e))^(1/2)+2*c^2*d*e+2*e^2-2*
(e*(c^2*d+e))^(1/2)*e)*arctan(c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))
/((c^2*d+2*(e*(c^2*d+e))^(1/2)+2*e)*d)^(1/2))/e/(c^2*d+e)/d^3/c+1/4/e*c^4*
sum(1/_R1/(_R1^2*c^2*d+c^2*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(
1/2)*(1+1/c/x)^(1/2))/_R1)+dilog((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/
2))/_R1)),_R1=RootOf(c^2*d*_Z^4+(2*c^2*d+4*e)*_Z^2+c^2*d))

```

**Fricas [F]**

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^2}{(ex^2 + d)^2} dx$$

input

```
integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="fricas")
```

output

```
integral((b*x^2*arcsech(c*x) + a*x^2)/(e^2*x^4 + 2*d*e*x^2 + d^2), x)
```

**Sympy [F]**

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{x^2(a + b \operatorname{asech}(cx))}{(d + ex^2)^2} dx$$

input `integrate(x**2*(a+b*asech(c*x))/(e*x**2+d)**2,x)`

output `Integral(x**2*(a + b*asech(c*x))/(d + e*x**2)**2, x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \text{Exception raised: ValueError}$$

input `integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^2}{(ex^2 + d)^2} dx$$

input `integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^2/(e*x^2 + d)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{x^2(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^2} dx$$

input `int((x^2*(a + b*acosh(1/(c*x))))/(d + e*x^2)^2,x)`output `int((x^2*(a + b*acosh(1/(c*x))))/(d + e*x^2)^2, x)`**Reduce [F]**

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx$$

$$= \frac{\sqrt{e} \sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e} \sqrt{d}}\right) ad + \sqrt{e} \sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e} \sqrt{d}}\right) ae x^2 + 2 \left( \int \frac{\operatorname{asech}(cx)x^2}{e^2 x^4 + 2de x^2 + d^2} dx \right) b d^2 e^2 + 2 \left( \int \frac{\operatorname{asech}(cx)x^2}{e^2 x^4 + 2de x^2 + d^2} dx \right) d}{2d e^2 (e x^2 + d)}$$

input `int(x^2*(a+b*asech(c*x))/(e*x^2+d)^2,x)`output `(sqrt(e)*sqrt(d)*atan((e*x)/(sqrt(e)*sqrt(d)))*a*d + sqrt(e)*sqrt(d)*atan((e*x)/(sqrt(e)*sqrt(d)))*a*e*x**2 + 2*int((asech(c*x)*x**2)/(d**2 + 2*d*e*x**2 + e**2*x**4),x)*b*d**2*e**2 + 2*int((asech(c*x)*x**2)/(d**2 + 2*d*e*x**2 + e**2*x**4),x)*b*d*e**3*x**2 - a*d*e*x)/(2*d*e**2*(d + e*x**2))`

$$3.120 \quad \int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex^2)^2} dx$$

Optimal result	973
Mathematica [C] (warning: unable to verify)	974
Rubi [A] (verified)	975
Maple [C] (warning: unable to verify)	978
Fricas [F]	979
Sympy [F]	980
Maxima [F(-2)]	980
Giac [F]	980
Mupad [F(-1)]	981
Reduce [F]	981

## Optimal result

Integrand size = 18, antiderivative size = 786

$$\begin{aligned}
 \int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^2} dx = & -\frac{a + b \operatorname{sech}^{-1}(cx)}{4d(\sqrt{-d}\sqrt{e} - \frac{d}{x})} + \frac{a + b \operatorname{sech}^{-1}(cx)}{4d(\sqrt{-d}\sqrt{e} + \frac{d}{x})} \\
 & + \frac{b \arctan\left(\frac{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{-1 + \frac{1}{cx}}}\right)}{2d\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{cd + \sqrt{-d}\sqrt{e}}} \\
 & + \frac{b \arctan\left(\frac{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{-1 + \frac{1}{cx}}}\right)}{2d\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{cd + \sqrt{-d}\sqrt{e}}} \\
 & - \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e} - \sqrt{c^2d + e}}\right)}{4(-d)^{3/2}\sqrt{e}} \\
 & + \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e} - \sqrt{c^2d + e}}\right)}{4(-d)^{3/2}\sqrt{e}} \\
 & - \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e} + \sqrt{c^2d + e}}\right)}{4(-d)^{3/2}\sqrt{e}} \\
 & + \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e} + \sqrt{c^2d + e}}\right)}{4(-d)^{3/2}\sqrt{e}} \\
 & + \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e} - \sqrt{c^2d + e}}\right)}{4(-d)^{3/2}\sqrt{e}} \\
 & - \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e} - \sqrt{c^2d + e}}\right)}{4(-d)^{3/2}\sqrt{e}} \\
 & + \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e} + \sqrt{c^2d + e}}\right)}{4(-d)^{3/2}\sqrt{e}} \\
 & - \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e} + \sqrt{c^2d + e}}\right)}{4(-d)^{3/2}\sqrt{e}}
 \end{aligned}$$

output

```

-1/4*(a+b*arcsech(c*x))/d/((-d)^(1/2)*e^(1/2)-d/x)+1/4*(a+b*arcsech(c*x))/
d/((-d)^(1/2)*e^(1/2)+d/x)+1/2*b*arctan((c*d-(-d)^(1/2)*e^(1/2))^(1/2)*(1+
1/c/x)^(1/2)/(c*d+(-d)^(1/2)*e^(1/2))^(1/2)/(-1+1/c/x)^(1/2))/d/(c*d-(-d)^(
1/2)*e^(1/2))^(1/2)/(c*d+(-d)^(1/2)*e^(1/2))^(1/2)+1/2*b*arctan((c*d+(-d)
^(1/2)*e^(1/2))^(1/2)*(1+1/c/x)^(1/2)/(c*d-(-d)^(1/2)*e^(1/2))^(1/2)/(-1+1
/c/x)^(1/2))/d/(c*d-(-d)^(1/2)*e^(1/2))^(1/2)/(c*d+(-d)^(1/2)*e^(1/2))^(1/
2)-1/4*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c
/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(3/2)/e^(1/2)+1/4*(a+b*arcsech(
c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-
(c^2*d+e)^(1/2)))/(-d)^(3/2)/e^(1/2)-1/4*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1
/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/(-
d)^(3/2)/e^(1/2)+1/4*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x
)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/(-d)^(3/2)/e^(1/2)+1/4
*b*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/
2)-(c^2*d+e)^(1/2)))/(-d)^(3/2)/e^(1/2)-1/4*b*polylog(2,c*(-d)^(1/2)*(1/c/
x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(3/2)/
e^(1/2)+1/4*b*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1
/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/(-d)^(3/2)/e^(1/2)-1/4*b*polylog(2,c*(-d)^(
1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)+(c^2*d+e)^(1/2)))/
(-d)^(3/2)/e^(1/2)

```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 1.02 (sec) , antiderivative size = 1216, normalized size of antiderivative = 1.55

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^2} dx = \text{Too large to display}$$

input

```
Integrate[(a + b*ArcSech[c*x])/(d + e*x^2)^2,x]
```

output

```

((2*a*Sqrt[d]*x)/(d + e*x^2) + (b*Sqrt[d]*ArcSech[c*x])/((-1)*Sqrt[d]*Sqrt
[e] + e*x) + (b*Sqrt[d]*ArcSech[c*x])/(I*Sqrt[d]*Sqrt[e] + e*x) + (2*a*Arc
Tan[(Sqrt[e]*x)/Sqrt[d]])/Sqrt[e] - (4*b*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sq
rt[d])]]/Sqrt[2]]*ArcTanh[(((I*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2)
/Sqrt[c^2*d + e])/Sqrt[e] + (4*b*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]]
/Sqrt[2]]*ArcTanh[(((I*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2)/Sqrt[c^2*
d + e])/Sqrt[e] - (I*b*ArcSech[c*x]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]
)))/(c*Sqrt[d]*E^ArcSech[c*x]))]/Sqrt[e] - (2*b*ArcSin[Sqrt[1 + (I*Sqrt[e]
)/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*
E^ArcSech[c*x]))]/Sqrt[e] + (I*b*ArcSech[c*x]*Log[1 + (I*(-Sqrt[e] + Sqrt[
c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x]))]/Sqrt[e] + (2*b*ArcSin[Sqrt[1 - (
I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/
(c*Sqrt[d]*E^ArcSech[c*x]))]/Sqrt[e] + (I*b*ArcSech[c*x]*Log[1 - (I*(Sqrt[
e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x]))]/Sqrt[e] - (2*b*ArcSin[
Sqrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 - (I*(Sqrt[e] + Sqrt[c^2*
d + e]))/(c*Sqrt[d]*E^ArcSech[c*x]))]/Sqrt[e] - (I*b*ArcSech[c*x]*Log[1 +
(I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x]))]/Sqrt[e] + (2*
b*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] +
Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x]))]/Sqrt[e] - (I*b*Log[((2*I)*S
qrt[e]*(Sqrt[d]*Sqrt[(1 - c*x)/(1 + c*x])*(1 + c*x) + (Sqrt[d]*Sqrt[e] ...

```

### Rubi [A] (verified)

Time = 2.62 (sec) , antiderivative size = 842, normalized size of antiderivative = 1.07, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {6847, 6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^2} dx \\
 & \quad \downarrow 6847 \\
 & - \int \frac{a + b \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\left(\frac{d}{x^2} + e\right)^2} d \frac{1}{x} \\
 & \quad \downarrow 6374
 \end{aligned}$$



$$\begin{aligned}
& - \int \left( \frac{a + \operatorname{barccosh}\left(\frac{1}{cx}\right)}{d\left(\frac{d}{x^2} + e\right)} - \frac{e\left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right)}{d\left(\frac{d}{x^2} + e\right)^2} \right) d\frac{1}{x} \\
& \quad \downarrow \text{2009} \\
& \frac{\log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{dc^2 + e}}\right) \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right)}{4(-d)^{3/2}\sqrt{e}} + \\
& \frac{\log\left(\frac{\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)c + 1}{\sqrt{e} - \sqrt{dc^2 + e}}\right) \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right)}{4(-d)^{3/2}\sqrt{e}} - \\
& \frac{\log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} + \sqrt{dc^2 + e}}\right) \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right)}{4(-d)^{3/2}\sqrt{e}} + \\
& \frac{\log\left(\frac{\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)c + 1}{\sqrt{e} + \sqrt{dc^2 + e}}\right) \left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right)}{4(-d)^{3/2}\sqrt{e}} - \frac{a + \operatorname{barccosh}\left(\frac{1}{cx}\right)}{4d\left(\sqrt{-d}\sqrt{e} - \frac{d}{x}\right)} + \frac{a + \operatorname{barccosh}\left(\frac{1}{cx}\right)}{4d\left(\frac{d}{x} + \sqrt{-d}\sqrt{e}\right)} + \\
& \frac{b \arctan\left(\frac{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{\frac{1}{cx} - 1}}\right)}{2d\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{cd + \sqrt{-d}\sqrt{e}}} + \frac{b \arctan\left(\frac{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{\frac{1}{cx} - 1}}\right)}{2d\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{cd + \sqrt{-d}\sqrt{e}}} + \\
& \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{4(-d)^{3/2}\sqrt{e}} - \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{4(-d)^{3/2}\sqrt{e}} + \\
& \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{4(-d)^{3/2}\sqrt{e}} - \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{4(-d)^{3/2}\sqrt{e}}
\end{aligned}$$

input `Int[(a + b*ArcSech[c*x])/(d + e*x^2)^2,x]`

output

```

-1/4*(a + b*ArcCosh[1/(c*x)])/(d*(Sqrt[-d]*Sqrt[e] - d/x)) + (a + b*ArcCos
h[1/(c*x)])/(4*d*(Sqrt[-d]*Sqrt[e] + d/x)) + (b*ArcTan[(Sqrt[c*d - Sqrt[-d
]*Sqrt[e]]*Sqrt[1 + 1/(c*x)])/(Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*Sqrt[-1 + 1/(c
*x)])))/(2*d*Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[c*d + Sqrt[-d]*Sqrt[e]]) +
(b*ArcTan[(Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*Sqrt[1 + 1/(c*x)])/(Sqrt[c*d - Sqr
t[-d]*Sqrt[e]]*Sqrt[-1 + 1/(c*x)])))/(2*d*Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqr
t[c*d + Sqrt[-d]*Sqrt[e]]) - ((a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]
)*E^ArcCosh[1/(c*x)]/(Sqrt[e] - Sqrt[c^2*d + e])])/(4*(-d)^(3/2)*Sqrt[e])
+ ((a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[
e] - Sqrt[c^2*d + e])])/(4*(-d)^(3/2)*Sqrt[e]) - ((a + b*ArcCosh[1/(c*x)])
*Log[1 - (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(4*
(-d)^(3/2)*Sqrt[e]) + ((a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcC
osh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(4*(-d)^(3/2)*Sqrt[e]) + (b*Po
lyLog[2, -((c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e]))])/(
4*(-d)^(3/2)*Sqrt[e]) - (b*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqr
t[e] - Sqrt[c^2*d + e])])/(4*(-d)^(3/2)*Sqrt[e]) + (b*PolyLog[2, -((c*Sqr
t[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e]))])/(4*(-d)^(3/2)*Sqr
t[e]) - (b*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*
d + e]))])/(4*(-d)^(3/2)*Sqrt[e])

```

### Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6374

```
Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((f_.)*(x_))^m_)*((d_) + (e
_)*(x_)^2)^p_, x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

rule 6847

```
Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^n_)*((d_.) + (e_.)*(x_)^2)^p_,
x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x^(2*(p + 1)
)), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0] && IntegerQ[p
]
```

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 39.14 (sec) , antiderivative size = 898, normalized size of antiderivative = 1.14

method	result
parts	$\frac{ax}{2d(x^2e+d)} + \frac{a \arctan\left(\frac{xe}{\sqrt{de}}\right)}{2d\sqrt{de}} + b \left( \frac{c^3 \operatorname{arcsech}(cx)x}{2d(e c^2 x^2 + c^2 d)} - \frac{\sqrt{-(c^2 d - 2\sqrt{e(c^2 d + e)} + 2e)} d (c^2 d + 2\sqrt{e(c^2 d + e)} + 2e) \operatorname{arctanh}\left(\frac{\sqrt{-(c^2 d - 2\sqrt{e(c^2 d + e)} + 2e)} d (c^2 d + 2\sqrt{e(c^2 d + e)} + 2e)}{2d^4 c^3}\right)}{2d^4 c^3} \right)$
derivativedivides	$\frac{a c^3 x}{2d(e c^2 x^2 + c^2 d)} + \frac{ac \arctan\left(\frac{xe}{\sqrt{de}}\right)}{2d\sqrt{de}} + b c^4 \left( \frac{\operatorname{arcsech}(cx)x}{2cd(e c^2 x^2 + c^2 d)} - \frac{\sqrt{-(c^2 d - 2\sqrt{e(c^2 d + e)} + 2e)} d (c^2 d + 2\sqrt{e(c^2 d + e)} + 2e) \operatorname{arctanh}\left(\frac{\sqrt{-(c^2 d - 2\sqrt{e(c^2 d + e)} + 2e)} d (c^2 d + 2\sqrt{e(c^2 d + e)} + 2e)}{2c^7 d^4}\right)}{2c^7 d^4} \right)$
default	$\frac{a c^3 x}{2d(e c^2 x^2 + c^2 d)} + \frac{ac \arctan\left(\frac{xe}{\sqrt{de}}\right)}{2d\sqrt{de}} + b c^4 \left( \frac{\operatorname{arcsech}(cx)x}{2cd(e c^2 x^2 + c^2 d)} - \frac{\sqrt{-(c^2 d - 2\sqrt{e(c^2 d + e)} + 2e)} d (c^2 d + 2\sqrt{e(c^2 d + e)} + 2e) \operatorname{arctanh}\left(\frac{\sqrt{-(c^2 d - 2\sqrt{e(c^2 d + e)} + 2e)} d (c^2 d + 2\sqrt{e(c^2 d + e)} + 2e)}{2c^7 d^4}\right)}{2c^7 d^4} \right)$

input `int((a+b*arcsech(c*x))/(e*x^2+d)^2,x,method=_RETURNVERBOSE)`

output

```

1/2*a*x/d/(e*x^2+d)+1/2*a/d/(d*e)^(1/2)*arctan(x*e/(d*e)^(1/2))+b/c*(1/2*c
^3*arcsech(c*x)*x/d/(c^2*e*x^2+c^2*d)-1/2*(-(c^2*d-2*(e*(c^2*d+e))^(1/2)+2
*e)*d)^(1/2)*(c^2*d+2*(e*(c^2*d+e))^(1/2)+2*e)*arctanh(c*d*(1/c/x+(-1+1/c/
x)^(1/2)*(1+1/c/x)^(1/2)))/((-c^2*d+2*(e*(c^2*d+e))^(1/2)-2*e)*d)^(1/2))/d^
4/c^3+1/2*(-(c^2*d-2*(e*(c^2*d+e))^(1/2)+2*e)*d)^(1/2)*(c^2*d*(e*(c^2*d+e)
)^(1/2)+2*c^2*d*e+2*e^2+2*(e*(c^2*d+e))^(1/2)*e)*arctanh(c*d*(1/c/x+(-1+1/
c/x)^(1/2)*(1+1/c/x)^(1/2)))/((-c^2*d+2*(e*(c^2*d+e))^(1/2)-2*e)*d)^(1/2))/
d^4/(c^2*d+e)/c^3-1/2*((c^2*d+2*(e*(c^2*d+e))^(1/2)+2*e)*d)^(1/2)*(c^2*d-2
*(e*(c^2*d+e))^(1/2)+2*e)*arctan(c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/
2)))/((c^2*d+2*(e*(c^2*d+e))^(1/2)+2*e)*d)^(1/2))/d^4/c^3+1/2*((c^2*d+2*(e
*(c^2*d+e))^(1/2)+2*e)*d)^(1/2)*(-c^2*d*(e*(c^2*d+e))^(1/2)+2*c^2*d*e+2*e^2
-2*(e*(c^2*d+e))^(1/2)*e)*arctan(c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/
2)))/((c^2*d+2*(e*(c^2*d+e))^(1/2)+2*e)*d)^(1/2))/d^4/(c^2*d+e)/c^3-1/4/d*c
^2*sum(_R1/(_R1^2*c^2*d+c^2*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(
1/2)*(1+1/c/x)^(1/2))/_R1)+dilog((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1
/2))/_R1)),_R1=RootOf(c^2*d*_Z^4+(2*c^2*d+4*e)*_Z^2+c^2*d))+1/4/d*c^2*sum(
1/_R1/(_R1^2*c^2*d+c^2*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)
*(1+1/c/x)^(1/2))/_R1)+dilog((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/
_R1)),_R1=RootOf(c^2*d*_Z^4+(2*c^2*d+4*e)*_Z^2+c^2*d))

```

**Fricas [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^2} dx = \int \frac{b \operatorname{ar} \operatorname{sech}(cx) + a}{(ex^2 + d)^2} dx$$

input

```
integrate((a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="fricas")
```

output

```
integral((b*arcsech(c*x) + a)/(e^2*x^4 + 2*d*e*x^2 + d^2), x)
```

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^2} dx = \int \frac{a + b \operatorname{arsech}(cx)}{(d + ex^2)^2} dx$$

input `integrate((a+b*asech(c*x))/(e*x**2+d)**2,x)`

output `Integral((a + b*asech(c*x))/(d + e*x**2)**2, x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^2} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^2} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^2} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/(e*x^2 + d)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^2} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{(ex^2 + d)^2} dx$$

input `int((a + b*acosh(1/(c*x)))/(d + e*x^2)^2,x)`

output `int((a + b*acosh(1/(c*x)))/(d + e*x^2)^2, x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^2} dx$$

$$= \frac{\sqrt{e} \sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e} \sqrt{d}}\right) ad + \sqrt{e} \sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e} \sqrt{d}}\right) aex^2 + 2 \left( \int \frac{a \operatorname{sech}(cx)}{e^2 x^4 + 2de x^2 + d^2} dx \right) b d^3 e + 2 \left( \int \frac{a \operatorname{sech}(cx)}{e^2 x^4 + 2de x^2 + d^2} dx \right)}{2d^2 e (ex^2 + d)}$$

input `int((a+b*asech(c*x))/(e*x^2+d)^2,x)`

output `(sqrt(e)*sqrt(d)*atan((e*x)/(sqrt(e)*sqrt(d)))*a*d + sqrt(e)*sqrt(d)*atan((e*x)/(sqrt(e)*sqrt(d)))*a*e*x**2 + 2*int(asech(c*x)/(d**2 + 2*d*e*x**2 + e**2*x**4),x)*b*d**3*e + 2*int(asech(c*x)/(d**2 + 2*d*e*x**2 + e**2*x**4),x)*b*d**2*e**2*x**2 + a*d*e*x)/(2*d**2*e*(d + e*x**2))`

$$3.121 \quad \int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^2(d+ex^2)^2} dx$$

Optimal result	983
Mathematica [C] (warning: unable to verify)	984
Rubi [A] (verified)	985
Maple [C] (warning: unable to verify)	988
Fricas [F]	989
Sympy [F(-1)]	989
Maxima [F(-2)]	989
Giac [F]	990
Mupad [F(-1)]	990
Reduce [F]	990

## Optimal result

Integrand size = 21, antiderivative size = 844

$$\begin{aligned}
\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^2} dx = & \frac{bc \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}}{d^2} - \frac{a}{d^2 x} - \frac{b \operatorname{sech}^{-1}(cx)}{d^2 x} \\
& + \frac{e(a + b \operatorname{sech}^{-1}(cx))}{4d^2 (\sqrt{-d}\sqrt{e} - \frac{d}{x})} - \frac{e(a + b \operatorname{sech}^{-1}(cx))}{4d^2 (\sqrt{-d}\sqrt{e} + \frac{d}{x})} \\
& - \frac{be \arctan\left(\frac{\sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{1 + \frac{1}{cx}}}{\sqrt{cd + \sqrt{-d}\sqrt{e}} \sqrt{-1 + \frac{1}{cx}}}\right)}{2d^2 \sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{cd + \sqrt{-d}\sqrt{e}}} \\
& - \frac{be \arctan\left(\frac{\sqrt{cd + \sqrt{-d}\sqrt{e}} \sqrt{1 + \frac{1}{cx}}}{\sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{-1 + \frac{1}{cx}}}\right)}{2d^2 \sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{cd + \sqrt{-d}\sqrt{e}}} \\
& - \frac{3\sqrt{e}(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2 d + e}}}\right)}{4(-d)^{5/2}} \\
& + \frac{3\sqrt{e}(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2 d + e}}}\right)}{4(-d)^{5/2}} \\
& - \frac{3\sqrt{e}(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2 d + e}}}\right)}{4(-d)^{5/2}} \\
& + \frac{3\sqrt{e}(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2 d + e}}}\right)}{4(-d)^{5/2}} \\
& + \frac{3b\sqrt{e} \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2 d + e}}}\right)}{4(-d)^{5/2}} \\
& - \frac{3b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e - \sqrt{c^2 d + e}}}\right)}{4(-d)^{5/2}} \\
& + \frac{3b\sqrt{e} \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2 d + e}}}\right)}{4(-d)^{5/2}} \\
& - \frac{3b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{c\sqrt{-d}e^{\operatorname{sech}^{-1}(cx)}}{\sqrt{e + \sqrt{c^2 d + e}}}\right)}{4(-d)^{5/2}}
\end{aligned}$$



output

```

b*c*(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)/d^2-a/d^2/x-b*arcsech(c*x)/d^2/x+1/4*
e*(a+b*arcsech(c*x))/d^2/((-d)^(1/2)*e^(1/2)-d/x)-1/4*e*(a+b*arcsech(c*x))
/d^2/((-d)^(1/2)*e^(1/2)+d/x)-1/2*b*e*arctan((c*d-(-d)^(1/2)*e^(1/2))^(1/2)
)*(1+1/c/x)^(1/2)/(c*d+(-d)^(1/2)*e^(1/2))^(1/2)/(-1+1/c/x)^(1/2))/d^2/(c*
d-(-d)^(1/2)*e^(1/2))^(1/2)/(c*d+(-d)^(1/2)*e^(1/2))^(1/2)-1/2*b*e*arctan(
(c*d+(-d)^(1/2)*e^(1/2))^(1/2)*(1+1/c/x)^(1/2)/(c*d-(-d)^(1/2)*e^(1/2))^(1
/2)/(-1+1/c/x)^(1/2))/d^2/(c*d-(-d)^(1/2)*e^(1/2))^(1/2)/(c*d+(-d)^(1/2)*e
^(1/2))^(1/2)-3/4*e^(1/2)*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+
1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(5/2)+3/4*e^
(1/2)*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/
x)^(1/2)))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(5/2)-3/4*e^(1/2)*(a+b*arcsech(c
*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)+(
c^2*d+e)^(1/2)))/(-d)^(5/2)+3/4*e^(1/2)*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/
2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(1/2)))/(-d
)^(5/2)+3/4*b*e^(1/2)*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1
/c/x)^(1/2)))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(5/2)-3/4*b*e^(1/2)*polylog(2
,c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)-(c^2*d+e)^(
1/2)))/(-d)^(5/2)+3/4*b*e^(1/2)*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)
^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(1/2)))/(-d)^(5/2)-3/4*b*e^(1/2
)*polylog(2,c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1...

```

### Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.88 (sec) , antiderivative size = 1305, normalized size of antiderivative = 1.55

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^2} dx = \text{Too large to display}$$

input

```
Integrate[(a + b*ArcSech[c*x])/(x^2*(d + e*x^2)^2),x]
```

output

```

((-4*a*Sqrt[d])/x + 4*b*c*Sqrt[d]*Sqrt[(1 - c*x)/(1 + c*x)] + (4*b*Sqrt[d]
*Sqrt[(1 - c*x)/(1 + c*x)])/x - (2*a*Sqrt[d]*e*x)/(d + e*x^2) - (4*b*Sqrt[d]
*d)*ArcSech[c*x])/x - (b*Sqrt[d]*e*ArcSech[c*x])/((-I)*Sqrt[d]*Sqrt[e] + e*
x) - (b*Sqrt[d]*e*ArcSech[c*x])/(I*Sqrt[d]*Sqrt[e] + e*x) - 6*a*Sqrt[e]*Ar
cTan[(Sqrt[e]*x)/Sqrt[d]] + 12*b*Sqrt[e]*ArcSin[Sqrt[1 - (I*Sqrt[e])/(c*Sq
rt[d])]]/Sqrt[2]]*ArcTanh[(((-I)*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])
/Sqrt[c^2*d + e]] - 12*b*Sqrt[e]*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]]/
Sqrt[2]]*ArcTanh[(((I*c*Sqrt[d] + Sqrt[e])*Tanh[ArcSech[c*x]/2])/Sqrt[c^2*d
+ e]] + (3*I)*b*Sqrt[e]*ArcSech[c*x]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e
]))/(c*Sqrt[d]*E^ArcSech[c*x])] + 6*b*Sqrt[e]*ArcSin[Sqrt[1 + (I*Sqrt[e])/
(c*Sqrt[d])]]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] - Sqrt[c^2*d + e]))/(c*Sqrt[d]*E
^ArcSech[c*x])] - (3*I)*b*Sqrt[e]*ArcSech[c*x]*Log[1 + (I*(-Sqrt[e] + Sqrt
[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - 6*b*Sqrt[e]*ArcSin[Sqrt[1 - (I
*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 + (I*(-Sqrt[e] + Sqrt[c^2*d + e]))/(
c*Sqrt[d]*E^ArcSech[c*x])] - (3*I)*b*Sqrt[e]*ArcSech[c*x]*Log[1 - (I*(Sqrt
[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + 6*b*Sqrt[e]*ArcSin[S
qrt[1 - (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 - (I*(Sqrt[e] + Sqrt[c^2*d
+ e]))/(c*Sqrt[d]*E^ArcSech[c*x])] + (3*I)*b*Sqrt[e]*ArcSech[c*x]*Log[1 +
(I*(Sqrt[e] + Sqrt[c^2*d + e]))/(c*Sqrt[d]*E^ArcSech[c*x])] - 6*b*Sqrt[e]
*ArcSin[Sqrt[1 + (I*Sqrt[e])/(c*Sqrt[d])]]/Sqrt[2]]*Log[1 + (I*(Sqrt[e] ...

```

### Rubi [A] (verified)

Time = 2.75 (sec) , antiderivative size = 904, normalized size of antiderivative = 1.07, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6857, 6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^2} dx \\
 & \quad \downarrow 6857 \\
 & - \int \frac{a + b \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\left(\frac{d}{x^2} + e\right)^2 x^4} d \frac{1}{x} \\
 & \quad \downarrow 6374
 \end{aligned}$$

$$\begin{aligned}
 & - \int \left( \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) e^2}{d^2 (\frac{d}{x^2} + e)^2} - \frac{2(a + \operatorname{barccosh}(\frac{1}{cx})) e}{d^2 (\frac{d}{x^2} + e)} + \frac{a + \operatorname{barccosh}(\frac{1}{cx})}{d^2} \right) d \frac{1}{x} \\
 & \quad \downarrow \text{2009} \\
 & - \frac{a}{d^2 x} - \frac{\operatorname{barccosh}(\frac{1}{cx})}{d^2 x} + \frac{e(a + \operatorname{barccosh}(\frac{1}{cx}))}{4d^2 (\sqrt{-d}\sqrt{e} - \frac{d}{x})} - \frac{e(a + \operatorname{barccosh}(\frac{1}{cx}))}{4d^2 (\frac{d}{x} + \sqrt{-d}\sqrt{e})} - \\
 & \frac{be \arctan \left( \frac{\sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{1 + \frac{1}{cx}}}{\sqrt{cd + \sqrt{-d}\sqrt{e}} \sqrt{\frac{1}{cx} - 1}} \right)}{2d^2 \sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{cd + \sqrt{-d}\sqrt{e}}} - \frac{be \arctan \left( \frac{\sqrt{cd + \sqrt{-d}\sqrt{e}} \sqrt{1 + \frac{1}{cx}}}{\sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{\frac{1}{cx} - 1}} \right)}{2d^2 \sqrt{cd - \sqrt{-d}\sqrt{e}} \sqrt{cd + \sqrt{-d}\sqrt{e}}} - \\
 & \frac{3\sqrt{e}(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( 1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{dc^2 + e}} \right)}{4(-d)^{5/2}} + \\
 & \frac{3\sqrt{e}(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( \frac{\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{dc^2 + e}} c + 1 \right)}{4(-d)^{5/2}} - \\
 & \frac{3\sqrt{e}(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( 1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{dc^2 + e}} \right)}{4(-d)^{5/2}} + \\
 & \frac{3\sqrt{e}(a + \operatorname{barccosh}(\frac{1}{cx})) \log \left( \frac{\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{dc^2 + e}} c + 1 \right)}{4(-d)^{5/2}} + \frac{3b\sqrt{e} \operatorname{PolyLog} \left( 2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{dc^2 + e}} \right)}{4(-d)^{5/2}} - \\
 & \frac{3b\sqrt{e} \operatorname{PolyLog} \left( 2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{dc^2 + e}} \right)}{4(-d)^{5/2}} + \frac{3b\sqrt{e} \operatorname{PolyLog} \left( 2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{dc^2 + e}} \right)}{4(-d)^{5/2}} - \\
 & \frac{3b\sqrt{e} \operatorname{PolyLog} \left( 2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{dc^2 + e}} \right)}{4(-d)^{5/2}} + \frac{bc\sqrt{\frac{1}{cx} - 1} \sqrt{1 + \frac{1}{cx}}}{d^2}
 \end{aligned}$$

input

`Int[(a + b*ArcSech[c*x])/(x^2*(d + e*x^2)^2), x]`

output

```
(b*c*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]/d^2 - a/(d^2*x) - (b*ArcCosh[1/
(c*x)]/(d^2*x) + (e*(a + b*ArcCosh[1/(c*x)])))/(4*d^2*(Sqrt[-d]*Sqrt[e] -
d/x)) - (e*(a + b*ArcCosh[1/(c*x)])))/(4*d^2*(Sqrt[-d]*Sqrt[e] + d/x)) - (b
*e*ArcTan[(Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[1 + 1/(c*x)])]/(Sqrt[c*d + Sqr
t[-d]*Sqrt[e]]*Sqrt[-1 + 1/(c*x)])))/(2*d^2*Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*S
qrt[c*d + Sqrt[-d]*Sqrt[e]]) - (b*e*ArcTan[(Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*S
qrt[1 + 1/(c*x)])/(Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[-1 + 1/(c*x)])))/(2*d
^2*Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[c*d + Sqrt[-d]*Sqrt[e]]) - (3*Sqrt[e]
*(a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]*E^ArcCosh[1/(c*x)]/(Sqrt[e]
- Sqrt[c^2*d + e]))]/(4*(-d)^(5/2)) + (3*Sqrt[e]*(a + b*ArcCosh[1/(c*x)]))
*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)]/(Sqrt[e] - Sqrt[c^2*d + e]))]/(4*
(-d)^(5/2)) - (3*Sqrt[e]*(a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]*E^Ar
cCosh[1/(c*x)]/(Sqrt[e] + Sqrt[c^2*d + e]))]/(4*(-d)^(5/2)) + (3*Sqrt[e]*
(a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)]/(Sqrt[e]
+ Sqrt[c^2*d + e]))]/(4*(-d)^(5/2)) + (3*b*Sqrt[e]*PolyLog[2, -((c*Sqrt[-d
]*E^ArcCosh[1/(c*x)]/(Sqrt[e] - Sqrt[c^2*d + e])))]/(4*(-d)^(5/2)) - (3*b
*Sqrt[e]*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)]/(Sqrt[e] - Sqrt[c^2*d
+ e]))]/(4*(-d)^(5/2)) + (3*b*Sqrt[e]*PolyLog[2, -((c*Sqrt[-d]*E^ArcCosh[1
/(c*x)]/(Sqrt[e] + Sqrt[c^2*d + e])))]/(4*(-d)^(5/2)) - (3*b*Sqrt[e]*Poly
Log[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)]/(Sqrt[e] + Sqrt[c^2*d + e]))]/(4...
```

### Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6374

```
Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^ (n_.)*((f_.)*(x_))^(m_.)*((d_) + (e
_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

rule 6857

```
Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^ (n_.)*(x_)^(m_.)*((d_.) + (e_.)*(x
_)^2)^(p_.), x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x
^(m + 2*(p + 1))), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0
] && IntegersQ[m, p]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 78.12 (sec) , antiderivative size = 1007, normalized size of antiderivative = 1.19

method	result	size
parts	Expression too large to display	1007
derivativedivides	Expression too large to display	1034
default	Expression too large to display	1034

input `int((a+b*arcsech(c*x))/x^2/(e*x^2+d)^2,x,method=_RETURNVERBOSE)`

output

```
a*(-1/d^2*e*(1/2*x/(e*x^2+d)+3/2/(d*e)^(1/2)*arctan(x*e/(d*e)^(1/2)))-1/d^2/x)+b*c*(-1/2*(-1+arcsech(c*x))/d^2*((-c*x-1)/c/x)^(1/2)*c*((c*x+1)/c/x)^(1/2)*x+1)/c/x+1/2*((-c*x-1)/c/x)^(1/2)*c*((c*x+1)/c/x)^(1/2)*x-1)*(1+arcsech(c*x))/d^2/c/x-1/2*arcsech(c*x)/d^2*e*c*x/(c^2*e*x^2+c^2*d)+1/2*(-(c^2*d-2*(e*(c^2*d+e))^(1/2)+2*e)*d)^(1/2)*(c^2*d+2*(e*(c^2*d+e))^(1/2)+2*e)*e*arctanh(c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/((-c^2*d+2*(e*(c^2*d+e))^(1/2)-2*e)*d)^(1/2))/d^5/c^5-1/2*(-(c^2*d-2*(e*(c^2*d+e))^(1/2)+2*e)*d)^(1/2)*(c^2*d*(e*(c^2*d+e))^(1/2)+2*c^2*d*e+2*e^2+2*(e*(c^2*d+e))^(1/2)*e)*e*arctanh(c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/((-c^2*d+2*(e*(c^2*d+e))^(1/2)-2*e)*d)^(1/2))/d^5/c^5/(c^2*d+e)+1/2*((c^2*d+2*(e*(c^2*d+e))^(1/2)+2*e)*d)^(1/2)*(c^2*d-2*(e*(c^2*d+e))^(1/2)+2*e)*e*arctan(c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/((c^2*d+2*(e*(c^2*d+e))^(1/2)+2*e)*d)^(1/2))/d^5/c^5-1/2*((c^2*d+2*(e*(c^2*d+e))^(1/2)+2*e)*d)^(1/2)*(-c^2*d*(e*(c^2*d+e))^(1/2)+2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^(1/2)*e)*e*arctan(c*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/((c^2*d+2*(e*(c^2*d+e))^(1/2)+2*e)*d)^(1/2))/d^5/c^5/(c^2*d+e)-3/4/d^2*e*sum(1/_R1/(_R1^2*c^2*d+c^2*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)+dilog((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)),_R1=RootOf(c^2*d*_Z^4+(2*c^2*d+4*e)*_Z^2+c^2*d))+3/4/d^2*e*sum(_R1/(_R1^2*c^2*d+c^2*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)+dilog((_R1-1/...
```

**Fricas [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^2} dx = \int \frac{b \operatorname{ar} \operatorname{sech}(cx) + a}{(ex^2 + d)^2 x^2} dx$$

input `integrate((a+b*arcsech(c*x))/x^2/(e*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*arcsech(c*x) + a)/(e^2*x^6 + 2*d*e*x^4 + d^2*x^2), x)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^2} dx = \text{Timed out}$$

input `integrate((a+b*asech(c*x))/x**2/(e*x**2+d)**2,x)`

output `Timed out`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^2} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/x^2/(e*x^2+d)^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^2} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^2 x^2} dx$$

input `integrate((a+b*arcsech(c*x))/x^2/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/((e*x^2 + d)^2*x^2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^2} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x^2 (ex^2 + d)^2} dx$$

input `int((a + b*acosh(1/(c*x)))/(x^2*(d + e*x^2)^2), x)`

output `int((a + b*acosh(1/(c*x)))/(x^2*(d + e*x^2)^2), x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^2} dx = \frac{-3\sqrt{e}\sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e}\sqrt{d}}\right) adx - 3\sqrt{e}\sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e}\sqrt{d}}\right) aex^3 + 2\left(\int \frac{\operatorname{asech}(cx)}{e^2x^6 + 2dex^4 + d^2x^2} dx\right) bd^4x + 2\left(\int \frac{\operatorname{asech}(cx)}{e^2x^6 + 2dex^4 + d^2x^2} dx\right)}{2d^3x(ex^2 + d)}$$

input `int((a+b*asech(c*x))/x^2/(e*x^2+d)^2,x)`

output

```
( - 3*sqrt(e)*sqrt(d)*atan((e*x)/(sqrt(e)*sqrt(d)))*a*d*x - 3*sqrt(e)*sqrt
(d)*atan((e*x)/(sqrt(e)*sqrt(d)))*a*e*x**3 + 2*int(asech(c*x)/(d**2*x**2 +
2*d*e*x**4 + e**2*x**6),x)*b*d**4*x + 2*int(asech(c*x)/(d**2*x**2 + 2*d*e
*x**4 + e**2*x**6),x)*b*d**3*e*x**3 - 2*a*d**2 - 3*a*d*e*x**2)/(2*d**3*x*(
d + e*x**2))
```



$$3.122 \quad \int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx$$

Optimal result	993
Mathematica [C] (warning: unable to verify)	994
Rubi [A] (verified)	995
Maple [C] (warning: unable to verify)	998
Fricas [F]	999
Sympy [F(-1)]	999
Maxima [F]	999
Giac [F]	1000
Mupad [F(-1)]	1000
Reduce [F]	1000

## Optimal result

Integrand size = 21, antiderivative size = 749

$$\begin{aligned}
\int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = & -\frac{bcd\sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}}{8e^2(c^2d + e)\left(e + \frac{d}{x^2}\right)x} \\
& -\frac{a + b\operatorname{sech}^{-1}(cx)}{4e\left(e + \frac{d}{x^2}\right)^2} - \frac{a + b\operatorname{sech}^{-1}(cx)}{2e^2\left(e + \frac{d}{x^2}\right)} \\
& + \frac{b\sqrt{-1 + \frac{1}{c^2x^2}}\operatorname{arctanh}\left(\frac{\sqrt{c^2d+e}}{c\sqrt{e}\sqrt{-1 + \frac{1}{c^2x^2}}x}\right)}{2e^{5/2}\sqrt{c^2d+e}\sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}} \\
& + \frac{b(c^2d + 2e)\sqrt{-1 + \frac{1}{c^2x^2}}\operatorname{arctanh}\left(\frac{\sqrt{c^2d+e}}{c\sqrt{e}\sqrt{-1 + \frac{1}{c^2x^2}}x}\right)}{8e^{5/2}(c^2d + e)^{3/2}\sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}} \\
& + \frac{(a + b\operatorname{sech}^{-1}(cx))\log\left(1 - \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2e^3} \\
& + \frac{(a + b\operatorname{sech}^{-1}(cx))\log\left(1 + \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2e^3} \\
& + \frac{(a + b\operatorname{sech}^{-1}(cx))\log\left(1 - \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2e^3} \\
& + \frac{(a + b\operatorname{sech}^{-1}(cx))\log\left(1 + \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2e^3} \\
& - \frac{(a + b\operatorname{sech}^{-1}(cx))\log\left(1 + e^{2\operatorname{sech}^{-1}(cx)}\right)}{e^3} \\
& + \frac{b\operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2e^3} \\
& + \frac{b\operatorname{PolyLog}\left(2, \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2e^3} \\
& + \frac{b\operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2e^3} \\
& + \frac{b\operatorname{PolyLog}\left(2, \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2e^3} \\
& - \frac{b\operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(cx)}\right)}{2e^3}
\end{aligned}$$

output

```

-1/8*b*c*d*(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)/e^2/(c^2*d+e)/(e+d/x^2)/x-1/4*
(a+b*arcsech(c*x))/e/(e+d/x^2)^2-1/2*(a+b*arcsech(c*x))/e^2/(e+d/x^2)+1/2*
b*(-1+1/c^2/x^2)^(1/2)*arctanh((c^2*d+e)^(1/2)/c/e^(1/2)/(-1+1/c^2/x^2)^(1
/2)/x)/e^(5/2)/(c^2*d+e)^(1/2)/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)+1/8*b*(c^2
*d+2*e)*(-1+1/c^2/x^2)^(1/2)*arctanh((c^2*d+e)^(1/2)/c/e^(1/2)/(-1+1/c^2/x
^2)^(1/2)/x)/e^(5/2)/(c^2*d+e)^(3/2)/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)+1/2*
(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/
2)))/(e^(1/2)-(c^2*d+e)^(1/2)))/e^3+1/2*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2
)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(1/2)))/e^3+
1/2*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)
^(1/2)))/(e^(1/2)+(c^2*d+e)^(1/2)))/e^3+1/2*(a+b*arcsech(c*x))*ln(1+c*(-d)^(
1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(1/2)))/
e^3-(a+b*arcsech(c*x))*ln(1+(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2)/e^
3+1/2*b*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(
e^(1/2)-(c^2*d+e)^(1/2)))/e^3+1/2*b*polylog(2,c*(-d)^(1/2)*(1/c/x+(-1+1/c/
x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)-(c^2*d+e)^(1/2)))/e^3+1/2*b*polylog(2,-
c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(
1/2)))/e^3+1/2*b*polylog(2,c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(
1/2)))/(e^(1/2)+(c^2*d+e)^(1/2)))/e^3-1/2*b*polylog(2,-(1/c/x+(-1+1/c/x)^(
1/2)*(1+1/c/x)^(1/2))^2)/e^3

```

### Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 7.27 (sec) , antiderivative size = 2000, normalized size of antiderivative = 2.67

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Too large to display}$$

input

```
Integrate[(x^5*(a + b*ArcSech[c*x]))/(d + e*x^2)^3,x]
```

output

```

-1/4*(a*d^2)/(e^3*(d + e*x^2)^2) + (a*d)/(e^3*(d + e*x^2)) + (a*Log[d + e*
x^2])/(2*e^3) + b*(-1/16*(d*((-I)*Sqrt[e]*Sqrt[(1 - c*x)/(1 + c*x)]*(1 +
c*x))/(Sqrt[d]*(c^2*d + e)*((-I)*Sqrt[d] + Sqrt[e]*x)) - ArcSech[c*x]/(Sqr
t[e]*((-I)*Sqrt[d] + Sqrt[e]*x)^2) + Log[x]/(d*Sqrt[e]) - Log[1 + Sqrt[(1
- c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]/(d*Sqrt[e]) + ((2*c^2*d
+ e)*Log[(-4*d*Sqrt[e]*Sqrt[c^2*d + e]*(Sqrt[e] - I*c^2*Sqrt[d]*x + Sqrt[
c^2*d + e]*Sqrt[(1 - c*x)/(1 + c*x)] + c*Sqrt[c^2*d + e]*x*Sqrt[(1 - c*x)/
(1 + c*x)])))/((2*c^2*d + e)*((-I)*Sqrt[d] + Sqrt[e]*x)))/(d*(c^2*d + e)^(
3/2))))/e^(5/2) - (d*((I*Sqrt[e]*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(Sqr
t[d]*(c^2*d + e)*(I*Sqrt[d] + Sqrt[e]*x)) - ArcSech[c*x]/(Sqrt[e]*(I*Sqrt[
d] + Sqrt[e]*x)^2) + Log[x]/(d*Sqrt[e]) - Log[1 + Sqrt[(1 - c*x)/(1 + c*x)
] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]/(d*Sqrt[e]) + ((2*c^2*d + e)*Log[(-4*d*
Sqrt[e]*Sqrt[c^2*d + e]*(Sqrt[e] + I*c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[
(1 - c*x)/(1 + c*x)] + c*Sqrt[c^2*d + e]*x*Sqrt[(1 - c*x)/(1 + c*x)])))/((2
*c^2*d + e)*(I*Sqrt[d] + Sqrt[e]*x)))/(d*(c^2*d + e)^(3/2))))/(16*e^(5/2)
) - (((7*I)/16)*Sqrt[d]*(-ArcSech[c*x]/(I*Sqrt[d]*Sqrt[e] + e*x)) + (I*(L
og[x]/Sqrt[e] - Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1
+ c*x)]/Sqrt[e] + Log[((2*I)*Sqrt[e]*(Sqrt[d]*Sqrt[(1 - c*x)/(1 + c*x)]*(
1 + c*x) + (Sqrt[d]*Sqrt[e] + I*c^2*d*x)/Sqrt[c^2*d + e]))/(I*Sqrt[d] + Sq
rt[e]*x])/Sqrt[c^2*d + e]))/Sqrt[d]))/e^(5/2) + (((7*I)/16)*Sqrt[d]*(-...

```

### Rubi [A] (verified)

Time = 2.01 (sec) , antiderivative size = 850, normalized size of antiderivative = 1.13, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6857, 6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx \\
 & \quad \downarrow \text{6857} \\
 & - \int \frac{x (a + b \operatorname{arccosh}(\frac{1}{cx}))}{(\frac{d}{x^2} + e)^3} d \frac{1}{x} \\
 & \quad \downarrow \text{6374}
 \end{aligned}$$

$$\begin{aligned}
 & - \int \left( \frac{x(a + \operatorname{barccosh}(\frac{1}{cx}))}{e^3} - \frac{d(a + \operatorname{barccosh}(\frac{1}{cx}))}{e^3 (\frac{d}{x^2} + e)} - \frac{d(a + \operatorname{barccosh}(\frac{1}{cx}))}{e^2 (\frac{d}{x^2} + e)^2 x} - \frac{d(a + \operatorname{barccosh}(\frac{1}{cx}))}{e (\frac{d}{x^2} + e)^3 x} \right) d\frac{1}{x} \\
 & \qquad \qquad \qquad \downarrow \text{2009} \\
 & - \frac{(a + \operatorname{barccosh}(\frac{1}{cx}))^2}{be^3} - \frac{\log\left(1 + e^{-2\operatorname{arccosh}(\frac{1}{cx})}\right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{e^3} + \\
 & \frac{\log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{dc^2 + e}}}\right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{2e^3} + \\
 & \frac{\log\left(\frac{\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{dc^2 + e}}} + 1\right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{2e^3} + \\
 & \frac{\log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}}\right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{2e^3} + \\
 & \frac{\log\left(\frac{\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}} + 1\right) (a + \operatorname{barccosh}(\frac{1}{cx}))}{2e^3} - \frac{a + \operatorname{barccosh}(\frac{1}{cx})}{2e^2 (\frac{d}{x^2} + e)} - \frac{a + \operatorname{barccosh}(\frac{1}{cx})}{4e (\frac{d}{x^2} + e)^2} + \\
 & \frac{b(dc^2 + 2e) \sqrt{\frac{1}{c^2 x^2} - 1} \operatorname{arctanh}\left(\frac{\sqrt{dc^2 + e}}{c\sqrt{e} \sqrt{\frac{1}{c^2 x^2} - 1}}\right)}{8e^{5/2} (dc^2 + e)^{3/2} \sqrt{\frac{1}{cx} - 1} \sqrt{1 + \frac{1}{cx}}} + \frac{b \sqrt{\frac{1}{c^2 x^2} - 1} \operatorname{arctanh}\left(\frac{\sqrt{dc^2 + e}}{c\sqrt{e} \sqrt{\frac{1}{c^2 x^2} - 1}}\right)}{2e^{5/2} \sqrt{dc^2 + e} \sqrt{\frac{1}{cx} - 1} \sqrt{1 + \frac{1}{cx}}} + \\
 & \frac{b \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arccosh}(\frac{1}{cx})}\right)}{2e^3} + \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{dc^2 + e}}}\right)}{2e^3} + \\
 & \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{dc^2 + e}}}\right)}{2e^3} + \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}}\right)}{2e^3} + \\
 & \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}}\right)}{2e^3} + \frac{bd(c^2 - \frac{1}{x^2})}{8ce^2 (dc^2 + e) (\frac{d}{x^2} + e) \sqrt{\frac{1}{cx} - 1} \sqrt{1 + \frac{1}{cx}}}
 \end{aligned}$$

input

```
Int[(x^5*(a + b*ArcSech[c*x]))/(d + e*x^2)^3,x]
```

output

```
(b*d*(c^2 - x^(-2)))/(8*c*e^2*(c^2*d + e)*(e + d/x^2)*Sqrt[-1 + 1/(c*x)]*S
qrt[1 + 1/(c*x)]*x) - (a + b*ArcCosh[1/(c*x)])/(4*e*(e + d/x^2)^2) - (a +
b*ArcCosh[1/(c*x)])/(2*e^2*(e + d/x^2)) - (a + b*ArcCosh[1/(c*x)]^2/(b*e^
3) + (b*Sqrt[-1 + 1/(c^2*x^2)]*ArcTanh[Sqrt[c^2*d + e]/(c*Sqrt[e]*Sqrt[-1
+ 1/(c^2*x^2)]*x)])/(2*e^(5/2)*Sqrt[c^2*d + e]*Sqrt[-1 + 1/(c*x)]*Sqrt[1 +
1/(c*x)]) + (b*(c^2*d + 2*e)*Sqrt[-1 + 1/(c^2*x^2)]*ArcTanh[Sqrt[c^2*d +
e]/(c*Sqrt[e]*Sqrt[-1 + 1/(c^2*x^2)]*x)])/(8*e^(5/2)*(c^2*d + e)^(3/2)*Sqr
t[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) - ((a + b*ArcCosh[1/(c*x)])*Log[1 + E^(-
2*ArcCosh[1/(c*x)])])/e^3 + ((a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]
*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])])/(2*e^3) + ((a + b*ArcCo
sh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d
+ e])])/(2*e^3) + ((a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]*E^ArcCosh
[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(2*e^3) + ((a + b*ArcCosh[1/(c*x)
])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(
2*e^3) + (b*PolyLog[2, -E^(-2*ArcCosh[1/(c*x)])])/(2*e^3) + (b*PolyLog[2,
-((c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e]))])/(2*e^3) +
(b*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])
])/ (2*e^3) + (b*PolyLog[2, -((c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sq
rt[c^2*d + e]))])/(2*e^3) + (b*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/
(Sqrt[e] + Sqrt[c^2*d + e]))])/(2*e^3)
```

### Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6374 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^ (n_.)*((f_.)*(x_))^(m_.)*((d_) + (e
_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]`

rule 6857 `Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^ (n_.)*(x_)^(m_.)*((d_.) + (e_.)*(x
_)^2)^(p_.), x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x
^(m + 2*(p + 1))), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0
] && IntegersQ[m, p]`

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 5.55 (sec) , antiderivative size = 1549, normalized size of antiderivative = 2.07

method	result	size
parts	Expression too large to display	1549
derivativedivides	Expression too large to display	1562
default	Expression too large to display	1562

input `int(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^3,x,method=_RETURNVERBOSE)`

output

```
a*(1/e^3*d/(e*x^2+d)+1/2/e^3*ln(e*x^2+d)-1/4*d^2/e^3/(e*x^2+d)^2)+b/c^6*(-
1/8*c^6*(4*arcsech(c*x)*c^6*d^2*x^2+6*e*arcsech(c*x)*c^6*d*x^4+(-(c*x-1)/c
/x)^(1/2)*((c*x+1)/c/x)^(1/2)*c^5*d^2*x+(-(c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)
^(1/2)*c^5*d*e*x^3+4*c^4*d*e*arcsech(c*x)*x^2+6*arcsech(c*x)*e^2*c^4*x^4-c
^4*d^2-2*c^4*d*e*x^2-c^4*e^2*x^4)/e^2/(c^2*d+e)/(c^2*e*x^2+c^2*d)^2-3/4*(e
*(c^2*d+e))^(1/2)/(c^2*d+e)^2/e^2*c^6*arctanh(1/4*(2*c^2*d*(1/c/x+(-1+1/c/
x)^(1/2)*(1+1/c/x)^(1/2))^2+2*c^2*d+4*e)/(c^2*d*e+e^2)^(1/2))-1/(c^2*d+e)/
e^2*c^6*arcsech(c*x)*ln(1+I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))-1/(c
^2*d+e)/e^2*c^6*arcsech(c*x)*ln(1-I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)
)))-1/(c^2*d+e)/e^2*c^6*dilog(1+I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)
))-1/(c^2*d+e)/e^2*c^6*dilog(1-I*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))
+1/4/(c^2*d+e)/e^2*c^6*sum((_R1^2*c^2*d+c^2*d+4*e)/(_R1^2*c^2*d+c^2*d+2*e)*
(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)+dilog((
_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)),_R1=RootOf(c^2*d*_Z^4+(2
*c^2*d+4*e)*_Z^2+c^2*d))-5/8*(e*(c^2*d+e))^(1/2)/(c^2*d+e)^2/e^3*c^8*d*arc
tanh(1/4*(2*c^2*d*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))^2+2*c^2*d+4*e)/
(c^2*d*e+e^2)^(1/2))+1/4/(c^2*d+e)/e^2*c^8*d*sum((_R1^2+1)/(_R1^2*c^2*d+c^
2*d+2*e)*(arcsech(c*x)*ln((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1
)+dilog((_R1-1/c/x-(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/_R1)),_R1=RootOf(c^2*
d*_Z^4+(2*c^2*d+4*e)*_Z^2+c^2*d))-1/(c^2*d+e)/e^3*c^8*d*arcsech(c*x)*ln...
```

**Fricas [F]**

$$\int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^5}{(ex^2 + d)^3} dx$$

input `integrate(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="fricas")`

output `integral((b*x^5*arcsech(c*x) + a*x^5)/(e^3*x^6 + 3*d*e^2*x^4 + 3*d^2*e*x^2 + d^3), x)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Timed out}$$

input `integrate(x**5*(a+b*asech(c*x))/(e*x**2+d)**3,x)`

output `Timed out`

**Maxima [F]**

$$\int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^5}{(ex^2 + d)^3} dx$$

input `integrate(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="maxima")`

output `1/4*a*((4*d*e*x^2 + 3*d^2)/(e^5*x^4 + 2*d*e^4*x^2 + d^2*e^3) + 2*log(e*x^2 + d)/e^3) + b*integrate(x^5*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(e^3*x^6 + 3*d*e^2*x^4 + 3*d^2*e*x^2 + d^3), x)`



**Giac [F]**

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^5}{(ex^2 + d)^3} dx$$

input `integrate(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^5/(e*x^2 + d)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \int \frac{x^5 (a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^3} dx$$

input `int((x^5*(a + b*acosh(1/(c*x))))/(d + e*x^2)^3,x)`

output `int((x^5*(a + b*acosh(1/(c*x))))/(d + e*x^2)^3, x)`

**Reduce [F]**

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx$$

$$= \frac{4 \left( \int \frac{\operatorname{asech}(cx)x^5}{e^3x^6 + 3de^2x^4 + 3d^2ex^2 + d^3} dx \right) b d^2 e^3 + 8 \left( \int \frac{\operatorname{asech}(cx)x^5}{e^3x^6 + 3de^2x^4 + 3d^2ex^2 + d^3} dx \right) b d e^4 x^2 + 4 \left( \int \frac{\operatorname{asech}(cx)x^5}{e^3x^6 + 3de^2x^4 + 3d^2ex^2 + d^3} dx \right) b d e^3 (e^2 x^4 + 2de^2 x^2 + d^2)}{4e^3 (e^2 x^4 + 2de^2 x^2 + d^2)}$$

input `int(x^5*(a+b*asech(c*x))/(e*x^2+d)^3,x)`

output

```
(4*int((asech(c*x)*x**5)/(d**3 + 3*d**2*e*x**2 + 3*d*e**2*x**4 + e**3*x**6),x)*b*d**2*e**3 + 8*int((asech(c*x)*x**5)/(d**3 + 3*d**2*e*x**2 + 3*d*e**2*x**4 + e**3*x**6),x)*b*d*e**4*x**2 + 4*int((asech(c*x)*x**5)/(d**3 + 3*d**2*e*x**2 + 3*d*e**2*x**4 + e**3*x**6),x)*b*e**5*x**4 + 2*log(d + e*x**2)*a*d**2 + 4*log(d + e*x**2)*a*d*e*x**2 + 2*log(d + e*x**2)*a*e**2*x**4 + a*d**2 - 2*a*e**2*x**4)/(4*e**3*(d**2 + 2*d*e*x**2 + e**2*x**4))
```

**3.123** 
$$\int \frac{x^3 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx$$

Optimal result	1002
Mathematica [C] (verified)	1003
Rubi [A] (verified)	1003
Maple [B] (verified)	1006
Fricas [B] (verification not implemented)	1007
Sympy [F(-1)]	1008
Maxima [F(-2)]	1008
Giac [F]	1008
Mupad [F(-1)]	1009
Reduce [F]	1009

**Optimal result**

Integrand size = 21, antiderivative size = 173

$$\int \frac{x^3 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \frac{b \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{8e(c^2d+e)(d+ex^2)} + \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{4d(d+ex^2)^2} - \frac{b(c^2d+2e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{\sqrt{c^2d+e}}\right)}{8de^{3/2}(c^2d+e)^{3/2}}$$

output

```
1/8*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/e/(c^2*d+e)/(e*x^2+d)+1/4*x^4*(a+b*arcsech(c*x))/d/(e*x^2+d)^2-1/8*b*(c^2*d+2*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh(e^(1/2)*(-c^2*x^2+1)^(1/2)/(c^2*d+e)^(1/2))/d/e^(3/2)/(c^2*d+e)^(3/2)
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 1.02 (sec) , antiderivative size = 486, normalized size of antiderivative = 2.81

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx =$$

$$-\frac{4ad}{(d+ex^2)^2} + \frac{8a}{d+ex^2} - \frac{2e\sqrt{\frac{1-cx}{1+cx}}(b+bcx)}{(c^2d+e)(d+ex^2)} + \frac{4b(d+2ex^2)\operatorname{sech}^{-1}(cx)}{(d+ex^2)^2} + \frac{4b\log(x)}{d} - \frac{4b\log\left(1+\sqrt{\frac{1-cx}{1+cx}}+cx\sqrt{\frac{1-cx}{1+cx}}\right)}{d} + \frac{b\sqrt{e}(c^2d+e)}{(d+ex^2)^2}$$

input

```
Integrate[(x^3*(a + b*ArcSech[c*x]))/(d + e*x^2)^3,x]
```

output

```
-1/16*((-4*a*d)/(d + e*x^2)^2 + (8*a)/(d + e*x^2) - (2*e*Sqrt[(1 - c*x)/(1 + c*x)]*(b + b*c*x))/((c^2*d + e)*(d + e*x^2)) + (4*b*(d + 2*e*x^2)*ArcSech[c*x])/(d + e*x^2)^2 + (4*b*Log[x])/d - (4*b*Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]])/d + (b*Sqrt[e]*(c^2*d + 2*e)*Log[(16*d*e^(3/2)*Sqrt[c^2*d + e]*(Sqrt[e] - I*c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[(1 - c*x)/(1 + c*x)] + c*Sqrt[c^2*d + e]*x*Sqrt[(1 - c*x)/(1 + c*x)])]/(b*(c^2*d + 2*e)*((-I)*Sqrt[d] + Sqrt[e]*x)))/(d*(c^2*d + e)^(3/2)) + (b*Sqrt[e]*(c^2*d + 2*e)*Log[(16*d*e^(3/2)*Sqrt[c^2*d + e]*(Sqrt[e] + I*c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[(1 - c*x)/(1 + c*x)] + c*Sqrt[c^2*d + e]*x*Sqrt[(1 - c*x)/(1 + c*x)])]/(b*(c^2*d + 2*e)*(I*Sqrt[d] + Sqrt[e]*x)))/(d*(c^2*d + e)^(3/2)))/e^2
```

**Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.88, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {6855, 27, 354, 87, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx$$

$$\begin{aligned}
& \downarrow 6855 \\
& b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^3}{4d\sqrt{1-c^2x^2}(ex^2+d)^2} dx + \frac{x^4(a+b\operatorname{sech}^{-1}(cx))}{4d(d+ex^2)^2} \\
& \downarrow 27 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^3}{\sqrt{1-c^2x^2}(ex^2+d)^2} dx}{4d} + \frac{x^4(a+b\operatorname{sech}^{-1}(cx))}{4d(d+ex^2)^2} \\
& \downarrow 354 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^2}{\sqrt{1-c^2x^2}(ex^2+d)^2} dx^2}{8d} + \frac{x^4(a+b\operatorname{sech}^{-1}(cx))}{4d(d+ex^2)^2} \\
& \downarrow 87 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{(c^2d+2e) \int \frac{1}{\sqrt{1-c^2x^2}(ex^2+d)} dx^2}{2e(c^2d+e)} + \frac{d\sqrt{1-c^2x^2}}{e(c^2d+e)(d+ex^2)} \right)}{8d} + \frac{x^4(a+b\operatorname{sech}^{-1}(cx))}{4d(d+ex^2)^2} \\
& \downarrow 73 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{d\sqrt{1-c^2x^2}}{e(c^2d+e)(d+ex^2)} - \frac{(c^2d+2e) \int \frac{1}{-\frac{ex^4}{c^2}+d+\frac{e}{c^2}} d\sqrt{1-c^2x^2}}{c^2e(c^2d+e)} \right)}{8d} + \frac{x^4(a+b\operatorname{sech}^{-1}(cx))}{4d(d+ex^2)^2} \\
& \downarrow 221 \\
& \frac{x^4(a+b\operatorname{sech}^{-1}(cx))}{4d(d+ex^2)^2} + \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{d\sqrt{1-c^2x^2}}{e(c^2d+e)(d+ex^2)} - \frac{(c^2d+2e)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{\sqrt{c^2d+e}}\right)}{e^{3/2}(c^2d+e)^{3/2}} \right)}{8d}
\end{aligned}$$

input `Int[(x^3*(a + b*ArcSech[c*x]))/(d + e*x^2)^3,x]`

output `(x^4*(a + b*ArcSech[c*x]))/(4*d*(d + e*x^2)^2) + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((d*Sqrt[1 - c^2*x^2])/(e*(c^2*d + e)*(d + e*x^2)) - ((c^2*d + 2*e)*ArcTanh[(Sqrt[e]*Sqrt[1 - c^2*x^2])/Sqrt[c^2*d + e]])/(e^(3/2)*(c^2*d + e)^(3/2))))/(8*d)`

## Definitions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e)), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 354 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]`
- rule 6855 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))`

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 1361 vs.  $2(147) = 294$ .

Time = 4.16 (sec) , antiderivative size = 1362, normalized size of antiderivative = 7.87

method	result	size
parts	Expression too large to display	1362
derivativeldivides	Expression too large to display	1395
default	Expression too large to display	1395

input `int(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^3,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & a * (-1/2/e^2/(e*x^2+d) + 1/4/e^2*d/(e*x^2+d)^2) + b/c^4 * (1/4*c^8*arcsech(c*x)*d \\
 & /e^2/(c^2*e*x^2+c^2*d)^2 - 1/2*c^6*arcsech(c*x)/e^2/(c^2*e*x^2+c^2*d) - 1/16*c \\
 & ^5 * ((-c*x-1)/c/x)^{(1/2)} * x * ((c*x+1)/c/x)^{(1/2)} * e * (4 * ((c^2*d+e)/e)^{(1/2)} * arc \\
 & tanh(1/(-c^2*x^2+1)^{(1/2)}) * c^6*d^2*e*x^2+4 * ((c^2*d+e)/e)^{(1/2)} * arctanh(1/( \\
 & -c^2*x^2+1)^{(1/2)}) * c^6*d^3 - \ln(-2 * ((-c^2*x^2+1)^{(1/2)} * ((c^2*d+e)/e)^{(1/2)} * e \\
 & - (-c^2*d*e)^{(1/2)} * c*x+e) / (-c*e*x+(-c^2*d*e)^{(1/2)})) * c^6*x^2*d^2*e - \ln(-2 * ( \\
 & -c^2*x^2+1)^{(1/2)} * ((c^2*d+e)/e)^{(1/2)} * e - (-c^2*d*e)^{(1/2)} * c*x+e) / (-c*e*x+(- \\
 & c^2*d*e)^{(1/2)})) * c^6*d^3 - \ln(2 * ((-c^2*x^2+1)^{(1/2)} * ((c^2*d+e)/e)^{(1/2)} * e + ( \\
 & -c^2*d*e)^{(1/2)} * c*x+e) / (c*e*x+(-c^2*d*e)^{(1/2)})) * c^6*d^2*e*x^2 - \ln(2 * ((-c^2* \\
 & x^2+1)^{(1/2)} * ((c^2*d+e)/e)^{(1/2)} * e + (-c^2*d*e)^{(1/2)} * c*x+e) / (c*e*x+(-c^2*d* \\
 & e)^{(1/2)})) * c^6*d^3 + 8 * ((c^2*d+e)/e)^{(1/2)} * arctanh(1/(-c^2*x^2+1)^{(1/2)}) * c^4 \\
 & * d * e^2 * x^2 + 8 * ((c^2*d+e)/e)^{(1/2)} * arctanh(1/(-c^2*x^2+1)^{(1/2)}) * c^4 * d^2 * e + 2 \\
 & * (-c^2*x^2+1)^{(1/2)} * ((c^2*d+e)/e)^{(1/2)} * c^4 * d^2 * e - 3 * \ln(-2 * ((-c^2*x^2+1)^{(1 \\
 & /2)} * ((c^2*d+e)/e)^{(1/2)} * e - (-c^2*d*e)^{(1/2)} * c*x+e) / (-c*e*x+(-c^2*d*e)^{(1/2)} \\
 & )) * c^4 * x^2 * d * e^2 - 3 * \ln(-2 * ((-c^2*x^2+1)^{(1/2)} * ((c^2*d+e)/e)^{(1/2)} * e - (-c^2*d \\
 & * e)^{(1/2)} * c*x+e) / (-c*e*x+(-c^2*d*e)^{(1/2)})) * c^4 * d^2 * e - 3 * \ln(2 * ((-c^2*x^2+1) \\
 & ^{(1/2)} * ((c^2*d+e)/e)^{(1/2)} * e + (-c^2*d*e)^{(1/2)} * c*x+e) / (c*e*x+(-c^2*d*e)^{(1/ \\
 & 2)})) * c^4 * d * e^2 * x^2 - 3 * \ln(2 * ((-c^2*x^2+1)^{(1/2)} * ((c^2*d+e)/e)^{(1/2)} * e + (-c^2*d \\
 & * e)^{(1/2)} * c*x+e) / (c*e*x+(-c^2*d*e)^{(1/2)})) * c^4 * d^2 * e + 4 * ((c^2*d+e)/e)^{(1/2)} \\
 & ) * arctanh(1/(-c^2*x^2+1)^{(1/2)}) * e^3 * c^2 * x^2 + 4 * ((c^2*d+e)/e)^{(1/2)} * arcta...
 \end{aligned}$$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 638 vs.  $2(115) = 230$ .

Time = 0.25 (sec) , antiderivative size = 1346, normalized size of antiderivative = 7.78

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Too large to display}$$

input `integrate(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="fricas")`

output

```
[-1/16*(4*a*c^4*d^4 + 2*(4*a - b)*c^2*d^3*e + 2*(2*a - b)*d^2*e^2 - 2*(b*c^2*d*e^3 + b*e^4)*x^4 + 4*(2*a*c^4*d^3*e + (4*a - b)*c^2*d^2*e^2 + (2*a - b)*d*e^3)*x^2 - (b*c^2*d^3 + (b*c^2*d*e^2 + 2*b*e^3)*x^4 + 2*b*d^2*e + 2*(b*c^2*d^2*e + 2*b*d*e^2)*x^2)*sqrt(c^2*d*e + e^2)*log((c^4*d^2 + 4*c^2*d*e - (c^4*d*e + 2*c^2*e^2)*x^2 + 4*(c^3*d*e + c*e^2)*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2))) + 4*e^2 + 2*(c^2*e*x^2 - c^2*d - (c^3*d + 2*c*e)*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 2*e)*sqrt(c^2*d*e + e^2))/(e*x^2 + d)) + 4*(b*c^4*d^4 + 2*b*c^2*d^3*e + b*d^2*e^2 + (b*c^4*d^2*e^2 + 2*b*c^2*d*e^3 + b*e^4)*x^4 + 2*(b*c^4*d^3*e + 2*b*c^2*d^2*e^2 + b*d*e^3)*x^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) + 4*(b*c^4*d^4 + 2*b*c^2*d^3*e + b*d^2*e^2 + 2*(b*c^4*d^3*e + 2*b*c^2*d^2*e^2 + b*d*e^3)*x^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - 2*((b*c^3*d^2*e^2 + b*c*d*e^3)*x^3 + (b*c^3*d^3*e + b*c*d^2*e^2)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^4*d^5*e^2 + 2*c^2*d^4*e^3 + d^3*e^4 + (c^4*d^3*e^4 + 2*c^2*d^2*e^5 + d*e^6)*x^4 + 2*(c^4*d^4*e^3 + 2*c^2*d^3*e^4 + d^2*e^5)*x^2), -1/8*(2*a*c^4*d^4 + (4*a - b)*c^2*d^3*e + (2*a - b)*d^2*e^2 - (b*c^2*d*e^3 + b*e^4)*x^4 + 2*(2*a*c^4*d^3*e + (4*a - b)*c^2*d^2*e^2 + (2*a - b)*d*e^3)*x^2 + (b*c^2*d^3 + (b*c^2*d*e^2 + 2*b*e^3)*x^4 + 2*b*d^2*e + 2*(b*c^2*d^2*e + 2*b*d*e^2)*x^2)*sqrt(-c^2*d*e - e^2)*arctan((sqrt(-c^2*d*e - e^2)*c*d*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - sqrt(-c^2*d*e - e^2)*(e*x^2 + d))/((c^2*d*e + e^2)*x^2)) + 2*(b*c^4*d^4 ...
```



**Sympy [F(-1)]**

Timed out.

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Timed out}$$

input `integrate(x**3*(a+b*asech(c*x))/(e*x**2+d)**3,x)`

output `Timed out`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

**Giac [F]**

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^3}{(ex^2 + d)^3} dx$$

input `integrate(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^3/(e*x^2 + d)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \int \frac{x^3(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^3} dx$$

input `int((x^3*(a + b*acosh(1/(c*x))))/(d + e*x^2)^3,x)`

output `int((x^3*(a + b*acosh(1/(c*x))))/(d + e*x^2)^3, x)`

**Reduce [F]**

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx$$

$$= \frac{4 \left( \int \frac{\operatorname{asech}(cx)x^3}{e^3x^6 + 3de^2x^4 + 3d^2ex^2 + d^3} dx \right) b d^3 + 8 \left( \int \frac{\operatorname{asech}(cx)x^3}{e^3x^6 + 3de^2x^4 + 3d^2ex^2 + d^3} dx \right) b d^2 e x^2 + 4 \left( \int \frac{\operatorname{asech}(cx)x^3}{e^3x^6 + 3de^2x^4 + 3d^2ex^2 + d^3} dx \right)}{4d(e^2x^4 + 2dex^2 + d^2)}$$

input `int(x^3*(a+b*asech(c*x))/(e*x^2+d)^3,x)`

output `(4*int((asech(c*x)*x**3)/(d**3 + 3*d**2*e*x**2 + 3*d*e**2*x**4 + e**3*x**6),x)*b*d**3 + 8*int((asech(c*x)*x**3)/(d**3 + 3*d**2*e*x**2 + 3*d*e**2*x**4 + e**3*x**6),x)*b*d**2*e*x**2 + 4*int((asech(c*x)*x**3)/(d**3 + 3*d**2*e*x**2 + 3*d*e**2*x**4 + e**3*x**6),x)*b*d*e**2*x**4 + a*x**4)/(4*d*(d**2 + 2*d*e*x**2 + e**2*x**4))`

**3.124**  $\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^3} dx$

Optimal result	1010
Mathematica [C] (verified)	1011
Rubi [A] (verified)	1012
Maple [B] (verified)	1015
Fricas [B] (verification not implemented)	1016
Sympy [F(-1)]	1017
Maxima [F(-2)]	1018
Giac [F]	1018
Mupad [F(-1)]	1018
Reduce [F]	1019

**Optimal result**

Integrand size = 19, antiderivative size = 217

$$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^3} dx = -\frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{8d(c^2d+e)(d+ex^2)} - \frac{a+b\operatorname{sech}^{-1}(cx)}{4e(d+ex^2)^2} + \frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\operatorname{arctanh}(\sqrt{1-c^2x^2})}{4d^2e} - \frac{b(3c^2d+2e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{\sqrt{c^2d+e}}\right)}{8d^2\sqrt{e}(c^2d+e)^{3/2}}$$

output

```
-1/8*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/d/(c^2*d+e)/(e*x^2+d)-1/4*(a+b*arcsech(c*x))/e/(e*x^2+d)^2+1/4*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh((-c^2*x^2+1)^(1/2))/d^2/e-1/8*b*(3*c^2*d+2*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh(e^(1/2)*(-c^2*x^2+1)^(1/2)/(c^2*d+e)^(1/2))/d^2/e^(1/2)/(c^2*d+e)^(3/2)
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.75 (sec) , antiderivative size = 486, normalized size of antiderivative = 2.24

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx$$

$$= \frac{1}{16} \left( -\frac{4a}{e(d + ex^2)^2} - \frac{2\sqrt{\frac{1-cx}{1+cx}}(b + bcx)}{d(c^2d + e)(d + ex^2)} - \frac{4b \operatorname{sech}^{-1}(cx)}{e(d + ex^2)^2} - \frac{4b \log(x)}{d^2e} \right.$$

$$+ \frac{4b \log\left(1 + \sqrt{\frac{1-cx}{1+cx}} + cx\sqrt{\frac{1-cx}{1+cx}}\right)}{d^2e}$$

$$- \frac{b(3c^2d + 2e) \log\left(\frac{16d^2\sqrt{e}\sqrt{c^2d+e}(\sqrt{e-ic^2\sqrt{dx}+\sqrt{c^2d+e}}\sqrt{\frac{1-cx}{1+cx}} + c\sqrt{c^2d+ex}\sqrt{\frac{1-cx}{1+cx}})}{b(3c^2d+2e)(-i\sqrt{d}+\sqrt{ex})}\right)}{d^2\sqrt{e}(c^2d + e)^{3/2}}$$

$$\left. - \frac{b(3c^2d + 2e) \log\left(\frac{16d^2\sqrt{e}\sqrt{c^2d+e}(\sqrt{e+ic^2\sqrt{dx}+\sqrt{c^2d+e}}\sqrt{\frac{1-cx}{1+cx}} + c\sqrt{c^2d+ex}\sqrt{\frac{1-cx}{1+cx}})}{b(3c^2d+2e)(i\sqrt{d}+\sqrt{ex})}\right)}{d^2\sqrt{e}(c^2d + e)^{3/2}} \right)$$

input `Integrate[(x*(a + b*ArcSech[c*x]))/(d + e*x^2)^3,x]`

output `((-4*a)/(e*(d + e*x^2)^2) - (2*sqrt[(1 - c*x)/(1 + c*x)]*(b + b*c*x))/(d*(c^2*d + e)*(d + e*x^2)) - (4*b*ArcSech[c*x])/(e*(d + e*x^2)^2) - (4*b*Log[x])/(d^2*e) + (4*b*Log[1 + sqrt[(1 - c*x)/(1 + c*x)] + c*x*sqrt[(1 - c*x)/(1 + c*x])])/(d^2*e) - (b*(3*c^2*d + 2*e)*Log[(16*d^2*sqrt[e]*sqrt[c^2*d + e]*(sqrt[e] - I*c^2*sqrt[d]*x + sqrt[c^2*d + e]*sqrt[(1 - c*x)/(1 + c*x)] + c*sqrt[c^2*d + e]*x*sqrt[(1 - c*x)/(1 + c*x]))]/(b*(3*c^2*d + 2*e)*((-I)*sqrt[d] + sqrt[e]*x)))/(d^2*sqrt[e]*(c^2*d + e)^(3/2)) - (b*(3*c^2*d + 2*e)*Log[(16*d^2*sqrt[e]*sqrt[c^2*d + e]*(sqrt[e] + I*c^2*sqrt[d]*x + sqrt[c^2*d + e]*sqrt[(1 - c*x)/(1 + c*x)] + c*sqrt[c^2*d + e]*x*sqrt[(1 - c*x)/(1 + c*x]))]/(b*(3*c^2*d + 2*e)*(I*sqrt[d] + sqrt[e]*x)))/(d^2*sqrt[e]*(c^2*d + e)^(3/2)))/16`

**Rubi [A] (verified)**

Time = 0.54 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.91, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$ , Rules used = {6853, 2036, 354, 114, 27, 174, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx \\
 & \quad \downarrow \text{6853} \\
 & - \frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{1}{x \sqrt{1-cx} \sqrt{cx+1} (ex^2+d)^2} dx}{4e} - \frac{a + b \operatorname{sech}^{-1}(cx)}{4e (d + ex^2)^2} \\
 & \quad \downarrow \text{2036} \\
 & - \frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{1}{x \sqrt{1-c^2x^2} (ex^2+d)^2} dx}{4e} - \frac{a + b \operatorname{sech}^{-1}(cx)}{4e (d + ex^2)^2} \\
 & \quad \downarrow \text{354} \\
 & - \frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{1}{x^2 \sqrt{1-c^2x^2} (ex^2+d)^2} dx^2}{8e} - \frac{a + b \operatorname{sech}^{-1}(cx)}{4e (d + ex^2)^2} \\
 & \quad \downarrow \text{114} \\
 & - \frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \left( \frac{\int \frac{-ex^2c^2+2dc^2+2e}{2x^2 \sqrt{1-c^2x^2} (ex^2+d)} dx^2}{d(c^2d+e)} + \frac{e \sqrt{1-c^2x^2}}{d(c^2d+e)(d+ex^2)} \right)}{8e} - \frac{a + b \operatorname{sech}^{-1}(cx)}{4e (d + ex^2)^2} \\
 & \quad \downarrow \text{27} \\
 & - \frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \left( \frac{\int \frac{2(dc^2+e)-c^2ex^2}{x^2 \sqrt{1-c^2x^2} (ex^2+d)} dx^2}{2d(c^2d+e)} + \frac{e \sqrt{1-c^2x^2}}{d(c^2d+e)(d+ex^2)} \right)}{8e} - \frac{a + b \operatorname{sech}^{-1}(cx)}{4e (d + ex^2)^2} \\
 & \quad \downarrow \text{174}
 \end{aligned}$$

$$\begin{aligned}
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{2(c^2d+e) \int \frac{1}{x^2\sqrt{1-c^2x^2}} dx^2 - \frac{e(3c^2d+2e) \int \frac{1}{\sqrt{1-c^2x^2}(ex^2+d)} dx^2}{2d(c^2d+e)}}{d} + \frac{e\sqrt{1-c^2x^2}}{d(c^2d+e)(d+ex^2)} \right) \\
 & \frac{8e}{a + b\operatorname{sech}^{-1}(cx)} \\
 & \frac{4e(d+ex^2)^2}{\phantom{a + b\operatorname{sech}^{-1}(cx)}} \\
 & \downarrow 73 \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{2e(3c^2d+2e) \int \frac{1}{-\frac{ex^4}{c^2}+d+\frac{e}{c^2}} d\sqrt{1-c^2x^2} - 4(c^2d+e) \int \frac{1}{\frac{1}{c^2}-\frac{x^4}{c^2}} d\sqrt{1-c^2x^2}}{2d(c^2d+e)} + \frac{e\sqrt{1-c^2x^2}}{d(c^2d+e)(d+ex^2)} \right) \\
 & \frac{8e}{a + b\operatorname{sech}^{-1}(cx)} \\
 & \frac{4e(d+ex^2)^2}{\phantom{a + b\operatorname{sech}^{-1}(cx)}} \\
 & \downarrow 221 \\
 & \frac{a + b\operatorname{sech}^{-1}(cx)}{4e(d+ex^2)^2} \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{2\sqrt{e}(3c^2d+2e)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{\sqrt{c^2d+e}}\right) - 4\operatorname{arctanh}(\sqrt{1-c^2x^2})(c^2d+e)}{d\sqrt{c^2d+e}} + \frac{e\sqrt{1-c^2x^2}}{d(c^2d+e)(d+ex^2)} \right) \\
 & \frac{8e}{\phantom{a + b\operatorname{sech}^{-1}(cx)}}
 \end{aligned}$$

input `Int[(x*(a + b*ArcSech[c*x]))/(d + e*x^2)^3,x]`

output `-1/4*(a + b*ArcSech[c*x])/(e*(d + e*x^2)^2) - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((e*Sqrt[1 - c^2*x^2])/(d*(c^2*d + e)*(d + e*x^2)) + ((-4*(c^2*d + e)*ArcTanh[Sqrt[1 - c^2*x^2]])/d + (2*Sqrt[e]*(3*c^2*d + 2*e)*ArcTanh[(Sqrt[e]*Sqrt[1 - c^2*x^2])/Sqrt[c^2*d + e]]/(d*Sqrt[c^2*d + e])))/(2*d*(c^2*d + e)))/(8*e)`

## Definitions of rubi rules used

- rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 73  $\text{Int}[(a_.) + (b_.)*(x_)^m*((c_.) + (d_.)*(x_))^n, x\_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \text{ Subst}[\text{Int}[x^{p*(m+1)-1}*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^{1/p}], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$
- rule 114  $\text{Int}[(a_.) + (b_.)*(x_)^m*((c_.) + (d_.)*(x_))^n*((e_.) + (f_.)*(x_))^{p_}, x_] \rightarrow \text{Simp}[b*(a + b*x)^{m+1}*(c + d*x)^{n+1}*((e + f*x)^{p+1})/((m+1)*(b*c - a*d)*(b*e - a*f)), x] + \text{Simp}[1/((m+1)*(b*c - a*d)*(b*e - a*f)) \text{ Int}[(a + b*x)^{m+1}*(c + d*x)^n*(e + f*x)^p*\text{Simp}[a*d*f*(m+1) - b*(d*e*(m+n+2) + c*f*(m+p+2)) - b*d*f*(m+n+p+3)*x, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \text{ILtQ}[m, -1] \ \&\& \ (\text{IntegerQ}[n] \ || \ \text{IntegersQ}[2*n, 2*p] \ || \ \text{ILtQ}[m+n+p+3, 0])$
- rule 174  $\text{Int}[(((e_.) + (f_.)*(x_))^{p_})*((g_.) + (h_.)*(x_)))/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_] \rightarrow \text{Simp}[(b*g - a*h)/(b*c - a*d) \text{ Int}[(e + f*x)^p/(a + b*x), x], x] - \text{Simp}[(d*g - c*h)/(b*c - a*d) \text{ Int}[(e + f*x)^p/(c + d*x), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, h\}, x]$
- rule 221  $\text{Int}[(a_.) + (b_.)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$
- rule 354  $\text{Int}[(x_)^m*((a_.) + (b_.)*(x_)^2)^{p_})*((c_.) + (d_.)*(x_)^2)^{q_}, x\_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; \text{FreeQ}[\{a, b, c, d, p, q\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IntegerQ}[(m-1)/2]$

rule 2036

```
Int[(u_)*((c_) + (d_)*(x_)^(n_))^(q_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_), x_Symbol] := Int[u*(a1*a2 + b1*b2*x^n)^p*(c + d*x^n)^q, x] /; FreeQ[{a1, b1, a2, b2, c, d, n, p, q}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))
```

rule 6853

```
Int[((a_) + ArcSech[(c_)*(x_)]*(b_))*(x_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcSech[c*x])/(2*e*(p + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(2*e*(p + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x^2)^(p + 1)/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[p, -1]
```

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1317 vs.  $2(183) = 366$ .

Time = 4.00 (sec) , antiderivative size = 1318, normalized size of antiderivative = 6.07

method	result	size
parts	Expression too large to display	1318
derivativedivides	Expression too large to display	1341
default	Expression too large to display	1341

input

```
int(x*(a+b*arcsech(c*x))/(e*x^2+d)^3,x,method=_RETURNVERBOSE)
```



output

```

-1/4*a/e/(e*x^2+d)^2+b/c^2*(-1/4*c^6/e/(c^2*e*x^2+c^2*d)^2*arcsech(c*x)-1/
16*c^3*e^2*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(4*((c^2*d+e)/e)^(1/
2)*arctanh(1/(-c^2*x^2+1)^(1/2))*c^6*d^2*e*x^2+4*((c^2*d+e)/e)^(1/2)*arcta
nh(1/(-c^2*x^2+1)^(1/2))*c^6*d^3-3*ln(-2*((-c^2*x^2+1)^(1/2))*((c^2*d+e)/e)
^(1/2)*e-(-c^2*d*e)^(1/2)*c*x+e)/(-c*e*x+(-c^2*d*e)^(1/2))) *c^6*x^2*d^2*e-
3*ln(-2*((-c^2*x^2+1)^(1/2))*((c^2*d+e)/e)^(1/2)*e-(-c^2*d*e)^(1/2)*c*x+e)/
(-c*e*x+(-c^2*d*e)^(1/2))) *c^6*d^3-3*ln(2*((-c^2*x^2+1)^(1/2))*((c^2*d+e)/e)
^(1/2)*e+(-c^2*d*e)^(1/2)*c*x+e)/(c*e*x+(-c^2*d*e)^(1/2))) *c^6*d^2*e*x^2-
3*ln(2*((-c^2*x^2+1)^(1/2))*((c^2*d+e)/e)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x+e)/(
c*e*x+(-c^2*d*e)^(1/2))) *c^6*d^3+8*((c^2*d+e)/e)^(1/2)*arctanh(1/(-c^2*x^2
+1)^(1/2))*c^4*d*e^2*x^2+8*((c^2*d+e)/e)^(1/2)*arctanh(1/(-c^2*x^2+1)^(1/2)
))*c^4*d^2*e-2*(-c^2*x^2+1)^(1/2))*((c^2*d+e)/e)^(1/2)*c^4*d^2*e-5*ln(-2*((
-c^2*x^2+1)^(1/2))*((c^2*d+e)/e)^(1/2)*e-(-c^2*d*e)^(1/2)*c*x+e)/(-c*e*x+(-
c^2*d*e)^(1/2))) *c^4*x^2*d*e^2-5*ln(-2*((-c^2*x^2+1)^(1/2))*((c^2*d+e)/e)^(
1/2)*e-(-c^2*d*e)^(1/2)*c*x+e)/(-c*e*x+(-c^2*d*e)^(1/2))) *c^4*d^2*e-5*ln(2
*((-c^2*x^2+1)^(1/2))*((c^2*d+e)/e)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x+e)/(c*e*x+
(-c^2*d*e)^(1/2))) *c^4*d*e^2*x^2-5*ln(2*((-c^2*x^2+1)^(1/2))*((c^2*d+e)/e)^(
1/2)*e+(-c^2*d*e)^(1/2)*c*x+e)/(c*e*x+(-c^2*d*e)^(1/2))) *c^4*d^2*e+4*((c^
2*d+e)/e)^(1/2)*arctanh(1/(-c^2*x^2+1)^(1/2))*e^3*c^2*x^2+4*((c^2*d+e)/e)^(
1/2)*arctanh(1/(-c^2*x^2+1)^(1/2))*c^2*d*e^2-2*(-c^2*x^2+1)^(1/2))*((c^...

```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 580 vs.  $2(135) = 270$ .

Time = 0.21 (sec) , antiderivative size = 1232, normalized size of antiderivative = 5.68

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Too large to display}$$

input

```
integrate(x*(a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="fricas")
```

output

```

[-1/16*(4*a*c^4*d^4 + 2*(4*a + b)*c^2*d^3*e + 2*(2*a + b)*d^2*e^2 + 2*(b*c
^2*d*e^3 + b*e^4)*x^4 + 4*(b*c^2*d^2*e^2 + b*d*e^3)*x^2 - (3*b*c^2*d^3 + (
3*b*c^2*d*e^2 + 2*b*e^3)*x^4 + 2*b*d^2*e + 2*(3*b*c^2*d^2*e + 2*b*d*e^2)*x
^2)*sqrt(c^2*d*e + e^2)*log((c^4*d^2 + 4*c^2*d*e - (c^4*d*e + 2*c^2*e^2)*x
^2 + 4*(c^3*d*e + c*e^2)*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 4*e^2 + 2*(c^2
*e*x^2 - c^2*d - (c^3*d + 2*c*e)*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 2*e)*s
qrt(c^2*d*e + e^2))/(e*x^2 + d)) + 4*(b*c^4*d^4 + 2*b*c^2*d^3*e + b*d^2*e^
2 + (b*c^4*d^2*e^2 + 2*b*c^2*d*e^3 + b*e^4)*x^4 + 2*(b*c^4*d^3*e + 2*b*c^2
*d^2*e^2 + b*d*e^3)*x^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/x) +
4*(b*c^4*d^4 + 2*b*c^2*d^3*e + b*d^2*e^2)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c
^2*x^2)) + 1)/(c*x)) + 2*((b*c^3*d^2*e^2 + b*c*d*e^3)*x^3 + (b*c^3*d^3*e +
b*c*d^2*e^2)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^4*d^6*e + 2*c^2*d^5*e^
2 + d^4*e^3 + (c^4*d^4*e^3 + 2*c^2*d^3*e^4 + d^2*e^5)*x^4 + 2*(c^4*d^5*e^2
+ 2*c^2*d^4*e^3 + d^3*e^4)*x^2), -1/8*(2*a*c^4*d^4 + (4*a + b)*c^2*d^3*e
+ (2*a + b)*d^2*e^2 + (b*c^2*d*e^3 + b*e^4)*x^4 + 2*(b*c^2*d^2*e^2 + b*d*e
^3)*x^2 + (3*b*c^2*d^3 + (3*b*c^2*d*e^2 + 2*b*e^3)*x^4 + 2*b*d^2*e + 2*(3*
b*c^2*d^2*e + 2*b*d*e^2)*x^2)*sqrt(-c^2*d*e - e^2)*arctan((sqrt(-c^2*d*e -
e^2)*c*d*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - sqrt(-c^2*d*e - e^2)*(e*x^2 +
d))/((c^2*d*e + e^2)*x^2)) + 2*(b*c^4*d^4 + 2*b*c^2*d^3*e + b*d^2*e^2 + (
b*c^4*d^2*e^2 + 2*b*c^2*d*e^3 + b*e^4)*x^4 + 2*(b*c^4*d^3*e + 2*b*c^2*d...

```

### Sympy [F(-1)]

Timed out.

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Timed out}$$

input

```
integrate(x*(a+b*asech(c*x))/(e*x**2+d)**3,x)
```

output

Timed out

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate(x*(a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

**Giac [F]**

$$\int \frac{x(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x}{(ex^2 + d)^3} dx$$

input `integrate(x*(a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x/(e*x^2 + d)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \int \frac{x(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^3} dx$$

input `int((x*(a + b*acosh(1/(c*x))))/(d + e*x^2)^3,x)`

output `int((x*(a + b*acosh(1/(c*x))))/(d + e*x^2)^3, x)`

**Reduce [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx$$

$$= \frac{4 \left( \int \frac{a \operatorname{sech}(cx)x}{e^3 x^6 + 3d e^2 x^4 + 3d^2 e x^2 + d^3} dx \right) b d^2 e + 8 \left( \int \frac{a \operatorname{sech}(cx)x}{e^3 x^6 + 3d e^2 x^4 + 3d^2 e x^2 + d^3} dx \right) b d e^2 x^2 + 4 \left( \int \frac{a \operatorname{sech}(cx)x}{e^3 x^6 + 3d e^2 x^4 + 3d^2 e x^2 + d^3} dx \right) b d^2 e}{4e(e^2 x^4 + 2d e x^2 + d^2)}$$

input `int(x*(a+b*asech(c*x))/(e*x^2+d)^3,x)`

output `(4*int((asech(c*x)*x)/(d**3 + 3*d**2*e*x**2 + 3*d*e**2*x**4 + e**3*x**6),x)*b*d**2*e + 8*int((asech(c*x)*x)/(d**3 + 3*d**2*e*x**2 + 3*d*e**2*x**4 + e**3*x**6),x)*b*d*e**2*x**2 + 4*int((asech(c*x)*x)/(d**3 + 3*d**2*e*x**2 + 3*d*e**2*x**4 + e**3*x**6),x)*b*e**3*x**4 - a)/(4*e*(d**2 + 2*d*e*x**2 + e**2*x**4))`

$$3.125 \quad \int \frac{a+b\operatorname{sech}^{-1}(cx)}{x(d+ex^2)^3} dx$$

Optimal result	1021
Mathematica [C] (warning: unable to verify)	1022
Rubi [A] (verified)	1023
Maple [C] (warning: unable to verify)	1026
Fricas [F]	1027
Sympy [F(-1)]	1027
Maxima [F]	1027
Giac [F]	1028
Mupad [F(-1)]	1028
Reduce [F]	1028

**Optimal result**

Integrand size = 21, antiderivative size = 730

$$\begin{aligned}
\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^3} dx &= \frac{bce\sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}}{8d^2(c^2d + e)\left(e + \frac{d}{x^2}\right)x} + \frac{e^2(a + b \operatorname{sech}^{-1}(cx))}{4d^3\left(e + \frac{d}{x^2}\right)^2} \\
&- \frac{e(a + b \operatorname{sech}^{-1}(cx))}{d^3\left(e + \frac{d}{x^2}\right)} + \frac{(a + b \operatorname{sech}^{-1}(cx))^2}{2bd^3} \\
&+ \frac{b\sqrt{e}\sqrt{-1 + \frac{1}{c^2x^2}} \operatorname{arctanh}\left(\frac{\sqrt{c^2d+e}}{c\sqrt{e}\sqrt{-1 + \frac{1}{c^2x^2}}x}\right)}{d^3\sqrt{c^2d+e}\sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}} \\
&- \frac{b\sqrt{e}(c^2d + 2e)\sqrt{-1 + \frac{1}{c^2x^2}} \operatorname{arctanh}\left(\frac{\sqrt{c^2d+e}}{c\sqrt{e}\sqrt{-1 + \frac{1}{c^2x^2}}x}\right)}{8d^3(c^2d + e)^{3/2}\sqrt{-1 + \frac{1}{cx}}\sqrt{1 + \frac{1}{cx}}} \\
&- \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2d^3} \\
&- \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2d^3} \\
&- \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 - \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2d^3} \\
&- \frac{(a + b \operatorname{sech}^{-1}(cx)) \log\left(1 + \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2d^3} \\
&- \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2d^3} \\
&- \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}-\sqrt{c^2d+e}}\right)}{2d^3} \\
&- \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2d^3} \\
&- \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de}\operatorname{sech}^{-1}(cx)}{\sqrt{e}+\sqrt{c^2d+e}}\right)}{2d^3}
\end{aligned}$$

output

```

1/8*b*c*e*(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)/d^2/(c^2*d+e)/(e+d/x^2)/x+1/4*
e^2*(a+b*arcsech(c*x))/d^3/(e+d/x^2)^2-e*(a+b*arcsech(c*x))/d^3/(e+d/x^2)+1
/2*(a+b*arcsech(c*x))^2/b/d^3+b*e^(1/2)*(-1+1/c^2/x^2)^(1/2)*arctanh((c^2*
d+e)^(1/2)/c/e^(1/2)/(-1+1/c^2/x^2)^(1/2)/x)/d^3/(c^2*d+e)^(1/2)/(-1+1/c/x
)^(1/2)/(1+1/c/x)^(1/2)-1/8*b*e^(1/2)*(c^2*d+2*e)*(-1+1/c^2/x^2)^(1/2)*arc
tanh((c^2*d+e)^(1/2)/c/e^(1/2)/(-1+1/c^2/x^2)^(1/2)/x)/d^3/(c^2*d+e)^(3/2)
/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)-1/2*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)
*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)-(c^2*d+e)^(1/2))/d^3-1
/2*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(
1/2)))/(e^(1/2)-(c^2*d+e)^(1/2))/d^3-1/2*(a+b*arcsech(c*x))*ln(1-c*(-d)^(
1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(1/2))/d
^3-1/2*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c
/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(1/2))/d^3-1/2*b*polylog(2,-c*(-d)^(1/2)*(1
/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)-(c^2*d+e)^(1/2))/d^3-1/2*
b*polylog(2,c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)
-(c^2*d+e)^(1/2))/d^3-1/2*b*polylog(2,-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/
2)*(1+1/c/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(1/2))/d^3-1/2*b*polylog(2,c*(-d)^(
1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)))/(e^(1/2)+(c^2*d+e)^(1/2))/
d^3

```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 6.06 (sec) , antiderivative size = 2054, normalized size of antiderivative = 2.81

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^3} dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*ArcSech[c*x])/(x*(d + e*x^2)^3),x]
```

output

```

a/(4*d*(d + e*x^2)^2) + a/(2*d^2*(d + e*x^2)) + (a*Log[x])/d^3 - (a*Log[d
+ e*x^2])/(2*d^3) + b*((Sqrt[e]*((-I)*Sqrt[e]*Sqrt[(1 - c*x)/(1 + c*x)]*(
1 + c*x))/(Sqrt[d]*(c^2*d + e)*((-I)*Sqrt[d] + Sqrt[e]*x)) - ArcSech[c*x]/
(Sqrt[e]*((-I)*Sqrt[d] + Sqrt[e]*x)^2) + Log[x]/(d*Sqrt[e]) - Log[1 + Sqrt
[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]]/(d*Sqrt[e]) + ((2*c
^2*d + e)*Log[(-4*d*Sqrt[e]*Sqrt[c^2*d + e]*(Sqrt[e] - I*c^2*Sqrt[d]*x + S
qrt[c^2*d + e]*Sqrt[(1 - c*x)/(1 + c*x]) + c*Sqrt[c^2*d + e]*x*Sqrt[(1 - c
*x)/(1 + c*x]))]/((2*c^2*d + e)*((-I)*Sqrt[d] + Sqrt[e]*x)))/(d*(c^2*d +
e)^(3/2)))/(16*d^2) + (Sqrt[e]*((I*Sqrt[e]*Sqrt[(1 - c*x)/(1 + c*x)]*(1 +
c*x))/(Sqrt[d]*(c^2*d + e)*(I*Sqrt[d] + Sqrt[e]*x)) - ArcSech[c*x]/(Sqrt[
e]*(I*Sqrt[d] + Sqrt[e]*x)^2) + Log[x]/(d*Sqrt[e]) - Log[1 + Sqrt[(1 - c*x
)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]]/(d*Sqrt[e]) + ((2*c^2*d + e)
*Log[(-4*d*Sqrt[e]*Sqrt[c^2*d + e]*(Sqrt[e] + I*c^2*Sqrt[d]*x + Sqrt[c^2*d
+ e]*Sqrt[(1 - c*x)/(1 + c*x]) + c*Sqrt[c^2*d + e]*x*Sqrt[(1 - c*x)/(1 +
c*x]))]/((2*c^2*d + e)*(I*Sqrt[d] + Sqrt[e]*x)))/(d*(c^2*d + e)^(3/2)))/
(16*d^2) - (((5*I)/16)*Sqrt[e]*(-(ArcSech[c*x]/(I*Sqrt[d]*Sqrt[e] + e*x))
+ (I*(Log[x]/Sqrt[e] - Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c
*x)/(1 + c*x)]]/Sqrt[e] + Log[((2*I)*Sqrt[e]*(Sqrt[d]*Sqrt[(1 - c*x)/(1 +
c*x)]*(1 + c*x) + (Sqrt[d]*Sqrt[e] + I*c^2*d*x)/Sqrt[c^2*d + e]))/(I*Sqrt[
d] + Sqrt[e]*x])/Sqrt[c^2*d + e]))/Sqrt[d])/d^(5/2) + (((5*I)/16)*Sqrt...

```

### Rubi [A] (verified)

Time = 1.86 (sec) , antiderivative size = 801, normalized size of antiderivative = 1.10, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6857, 6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^3} dx \\
 & \quad \downarrow 6857 \\
 & - \int \frac{a + b \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\left(\frac{d}{x^2} + e\right)^3 x^5} d \frac{1}{x} \\
 & \quad \downarrow 6374
 \end{aligned}$$



$$\begin{aligned}
 & - \int \left( \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) e^2}{d^2 (\frac{d}{x^2} + e)^3 x} - \frac{2(a + \operatorname{barccosh}(\frac{1}{cx})) e}{d^2 (\frac{d}{x^2} + e)^2 x} + \frac{a + \operatorname{barccosh}(\frac{1}{cx})}{d^2 (\frac{d}{x^2} + e) x} \right) d \frac{1}{x} \\
 & \qquad \qquad \qquad \downarrow \text{2009} \\
 & \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) e^2}{4d^3 (\frac{d}{x^2} + e)^2} - \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) e}{d^3 (\frac{d}{x^2} + e)} - \frac{b(c^2 - \frac{1}{x^2}) e}{8cd^2 (dc^2 + e) (\frac{d}{x^2} + e) \sqrt{\frac{1}{cx} - 1} \sqrt{1 + \frac{1}{cx} x}} - \\
 & \frac{b(dc^2 + 2e) \sqrt{\frac{1}{c^2 x^2} - 1} \operatorname{arctanh}\left(\frac{\sqrt{dc^2 + e}}{c\sqrt{e}\sqrt{\frac{1}{c^2 x^2} - 1} x}\right) \sqrt{e}}{8d^3 (dc^2 + e)^{3/2} \sqrt{\frac{1}{cx} - 1} \sqrt{1 + \frac{1}{cx}}} + \frac{b\sqrt{\frac{1}{c^2 x^2} - 1} \operatorname{arctanh}\left(\frac{\sqrt{dc^2 + e}}{c\sqrt{e}\sqrt{\frac{1}{c^2 x^2} - 1} x}\right) \sqrt{e}}{d^3 \sqrt{dc^2 + e} \sqrt{\frac{1}{cx} - 1} \sqrt{1 + \frac{1}{cx}}} + \\
 & \frac{(a + \operatorname{barccosh}(\frac{1}{cx}))^2}{2bd^3} - \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) \log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{2d^3} - \\
 & \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) \log\left(\frac{\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx}) c + 1}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{2d^3} - \\
 & \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) \log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{2d^3} - \\
 & \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) \log\left(\frac{\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx}) c + 1}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{2d^3} - \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{2d^3} - \\
 & \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{2d^3} - \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{2d^3} - \\
 & \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{2d^3}
 \end{aligned}$$

input `Int[(a + b*ArcSech[c*x])/(x*(d + e*x^2)^3), x]`

output

```

-1/8*(b*e*(c^2 - x^(-2)))/(c*d^2*(c^2*d + e)*(e + d/x^2)*Sqrt[-1 + 1/(c*x)
]*Sqrt[1 + 1/(c*x)]*x + (e^2*(a + b*ArcCosh[1/(c*x)]))/(4*d^3*(e + d/x^2)
^2) - (e*(a + b*ArcCosh[1/(c*x)]))/(d^3*(e + d/x^2)) + (a + b*ArcCosh[1/(c
*x)])^2/(2*b*d^3) + (b*Sqrt[e]*Sqrt[-1 + 1/(c^2*x^2)]*ArcTanh[Sqrt[c^2*d +
e]/(c*Sqrt[e]*Sqrt[-1 + 1/(c^2*x^2)]*x)]/(d^3*Sqrt[c^2*d + e]*Sqrt[-1 +
1/(c*x)]*Sqrt[1 + 1/(c*x)]) - (b*Sqrt[e]*(c^2*d + 2*e)*Sqrt[-1 + 1/(c^2*x^
2)]*ArcTanh[Sqrt[c^2*d + e]/(c*Sqrt[e]*Sqrt[-1 + 1/(c^2*x^2)]*x)]/(8*d^3*
(c^2*d + e)^(3/2)*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) - ((a + b*ArcCosh[
1/(c*x)])*Log[1 - (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d +
e])])/(2*d^3) - ((a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/
(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])])/(2*d^3) - ((a + b*ArcCosh[1/(c*x)])*
Log[1 - (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(2*d
^3) - ((a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(S
qrt[e] + Sqrt[c^2*d + e])])/(2*d^3) - (b*PolyLog[2, -(c*Sqrt[-d]*E^ArcCos
h[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])])/(2*d^3) - (b*PolyLog[2, (c*Sqrt
[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])])/(2*d^3) - (b*PolyLo
g[2, -(c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d + e])])/(2*d
^3) - (b*PolyLog[2, (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] + Sqrt[c^2*d
+ e])])/(2*d^3)

```

### Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6374

```
Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*((f_.)*(x_))^(m_.)*((d_) + (e
_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

rule 6857

```
Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^n_.*(x_)^(m_.)*((d_.) + (e_.)*(x_
)^2)^(p_.), x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x
^(m + 2*(p + 1))), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0
] && IntegersQ[m, p]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.09 (sec) , antiderivative size = 3727, normalized size of antiderivative = 5.11

method	result	size
parts	Expression too large to display	3727
derivativedivides	Expression too large to display	3801
default	Expression too large to display	3801

input `int((a+b*arcsech(c*x))/x/(e*x^2+d)^3,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & \frac{1}{2}a/d^2/(e*x^2+d) - \frac{1}{2}a/d^3*\ln(e*x^2+d) + \frac{1}{4}a/d/(e*x^2+d)^2 + a/d^3*\ln(x) + \\
 & b*(-2*(-c^2*d*(e*(c^2*d+e))^{(1/2)} + 2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^{(1/2)}*e) \\
 & *e/d^4/(c^4*d^2+2*c^2*d*e+e^2)/c^2*arcsech(c*x)^2 + 3/2*(c^2*d-2*(e*(c^2*d+e) \\
 & ))^{(1/2)}+2*e)/(c^2*d+e)/d^4/c^2*e*arcsech(c*x)^2 + 1/8*(-c^2*d*(e*(c^2*d+e)) \\
 & ^{(1/2)}+2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^{(1/2)}*e)/e/(c^4*d^2+2*c^2*d*e+e^2)* \\
 & c^2/d^2*polylog(2,d*c^2*(1/c/x+(-1+1/c/x)^{(1/2)}*(1+1/c/x)^{(1/2)})^2/(-c^2*d \\
 & -2*(e*(c^2*d+e))^{(1/2)}-2*e))-1/4*(e*(c^2*d+e))^{(1/2)}/(c^2*d+e)^2/d/e*c^4*a \\
 & rcsech(c*x)^2 + 1/2*(-c^2*d*(e*(c^2*d+e))^{(1/2)}+2*c^2*d*e+2*e^2-2*(e*(c^2*d+ \\
 & e))^{(1/2)}*e)/(c^4*d^2+2*c^2*d*e+e^2)*e^2/d^5/c^4*polylog(2,d*c^2*(1/c/x+(- \\
 & 1+1/c/x)^{(1/2)}*(1+1/c/x)^{(1/2)})^2/(-c^2*d-2*(e*(c^2*d+e))^{(1/2)}-2*e))+1/2* \\
 & (e*(c^2*d+e))^{(1/2)}/(c^2*d+e)^2/d^3*e*arcsech(c*x)*\ln(1-d*c^2*(1/c/x+(-1+1 \\
 & /c/x)^{(1/2)}*(1+1/c/x)^{(1/2)})^2/(-c^2*d+2*(e*(c^2*d+e))^{(1/2)}-2*e))+3/4*(e \\
 & (c^2*d+e))^{(1/2)}/(c^2*d+e)^2/d^2*c^2*arcsech(c*x)*\ln(1-d*c^2*(1/c/x+(-1+1/ \\
 & c/x)^{(1/2)}*(1+1/c/x)^{(1/2)})^2/(-c^2*d+2*(e*(c^2*d+e))^{(1/2)}-2*e))+(-c^2*d* \\
 & (e*(c^2*d+e))^{(1/2)}+2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^{(1/2)}*e)*e/d^4/(c^4*d^ \\
 & 2+2*c^2*d*e+e^2)/c^2*polylog(2,d*c^2*(1/c/x+(-1+1/c/x)^{(1/2)}*(1+1/c/x)^{(1/ \\
 & 2)})^2/(-c^2*d-2*(e*(c^2*d+e))^{(1/2)}-2*e))-1/2*(c^2*d-2*(e*(c^2*d+e))^{(1/2)} \\
 & +2*e)/(c^2*d+e)/d^5/c^4*e^2*polylog(2,d*c^2*(1/c/x+(-1+1/c/x)^{(1/2)}*(1+1/c \\
 & /x)^{(1/2)})^2/(-c^2*d-2*(e*(c^2*d+e))^{(1/2)}-2*e))+(-c^2*d*(e*(c^2*d+e))^{(1/ \\
 & 2)}+2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^{(1/2)}*e)/(c^4*d^2+2*c^2*d*e+e^2)*e^2\dots
 \end{aligned}$$

**Fricas [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x (d + ex^2)^3} dx = \int \frac{b \operatorname{ar} \operatorname{sech}(cx) + a}{(ex^2 + d)^3 x} dx$$

input `integrate((a+b*arcsech(c*x))/x/(e*x^2+d)^3,x, algorithm="fricas")`

output `integral((b*arcsech(c*x) + a)/(e^3*x^7 + 3*d*e^2*x^5 + 3*d^2*e*x^3 + d^3*x), x)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x (d + ex^2)^3} dx = \text{Timed out}$$

input `integrate((a+b*asech(c*x))/x/(e*x**2+d)**3,x)`

output `Timed out`

**Maxima [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x (d + ex^2)^3} dx = \int \frac{b \operatorname{ar} \operatorname{sech}(cx) + a}{(ex^2 + d)^3 x} dx$$

input `integrate((a+b*arcsech(c*x))/x/(e*x^2+d)^3,x, algorithm="maxima")`

output `1/4*a*((2*e*x^2 + 3*d)/(d^2*e^2*x^4 + 2*d^3*e*x^2 + d^4) - 2*log(e*x^2 + d)/d^3 + 4*log(x)/d^3) + b*integrate(log(sqrt(1/(c*x) + 1))*sqrt(1/(c*x) - 1) + 1/(c*x))/(e^3*x^7 + 3*d*e^2*x^5 + 3*d^2*e*x^3 + d^3*x), x)`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^3} dx = \int \frac{b \operatorname{ar} \operatorname{sech}(cx) + a}{(ex^2 + d)^3 x} dx$$

input `integrate((a+b*arcsech(c*x))/x/(e*x^2+d)^3,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/((e*x^2 + d)^3*x), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^3} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x(e x^2 + d)^3} dx$$

input `int((a + b*acosh(1/(c*x)))/(x*(d + e*x^2)^3), x)`

output `int((a + b*acosh(1/(c*x)))/(x*(d + e*x^2)^3), x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^3} dx$$

$$= \frac{4 \left( \int \frac{\operatorname{asech}(cx)}{e^3 x^7 + 3d e^2 x^5 + 3d^2 e x^3 + d^3 x} dx \right) b d^5 + 8 \left( \int \frac{\operatorname{asech}(cx)}{e^3 x^7 + 3d e^2 x^5 + 3d^2 e x^3 + d^3 x} dx \right) b d^4 e x^2 + 4 \left( \int \frac{\operatorname{asech}(cx)}{e^3 x^7 + 3d e^2 x^5 + 3d^2 e x^3 + d^3 x} dx \right)}$$

input `int((a+b*asech(c*x))/x/(e*x^2+d)^3,x)`

output

```
(4*int(asech(c*x)/(d**3*x + 3*d**2*e*x**3 + 3*d*e**2*x**5 + e**3*x**7),x)*
b*d**5 + 8*int(asech(c*x)/(d**3*x + 3*d**2*e*x**3 + 3*d*e**2*x**5 + e**3*x
**7),x)*b*d**4*e*x**2 + 4*int(asech(c*x)/(d**3*x + 3*d**2*e*x**3 + 3*d*e**
2*x**5 + e**3*x**7),x)*b*d**3*e**2*x**4 - 2*log(d + e*x**2)*a*d**2 - 4*log
(d + e*x**2)*a*d*e*x**2 - 2*log(d + e*x**2)*a*e**2*x**4 + 4*log(x)*a*d**2
+ 8*log(x)*a*d*e*x**2 + 4*log(x)*a*e**2*x**4 + 2*a*d**2 - a*e**2*x**4)/(4*
d**3*(d**2 + 2*d*e*x**2 + e**2*x**4))
```

**3.126** 
$$\int \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx$$

Optimal result	1030
Mathematica [C] (warning: unable to verify)	1031
Rubi [A] (verified)	1032
Maple [C] (warning: unable to verify)	1035
Fricas [F]	1036
Sympy [F(-1)]	1036
Maxima [F(-2)]	1036
Giac [F]	1037
Mupad [F(-1)]	1037
Reduce [F]	1037

**Optimal result**

Integrand size = 21, antiderivative size = 1272

$$\int \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Too large to display}$$

output

```

1/16*b*c*(-d)^(1/2)*(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)/e^(3/2)/(c^2*d+e)/((-
d)^(1/2)*e^(1/2)-d/x)+1/16*b*c*(-d)^(1/2)*(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)
/e^(3/2)/(c^2*d+e)/((-d)^(1/2)*e^(1/2)+d/x)+1/16*(-d)^(1/2)*(a+b*arcsech(c
*x))/e^(3/2)/((-d)^(1/2)*e^(1/2)-d/x)^2+3/16*(a+b*arcsech(c*x))/e^2/((-d)^(
1/2)*e^(1/2)-d/x)-1/16*(-d)^(1/2)*(a+b*arcsech(c*x))/e^(3/2)/((-d)^(1/2)*
e^(1/2)+d/x)^2-3/16*(a+b*arcsech(c*x))/e^2/((-d)^(1/2)*e^(1/2)+d/x)-3/8*b*
arctan((c*d-(-d)^(1/2)*e^(1/2))^(1/2)*(1+1/c/x)^(1/2)/(c*d+(-d)^(1/2)*e^(1
/2))^(1/2)/(-1+1/c/x)^(1/2))/(c*d-(-d)^(1/2)*e^(1/2))^(1/2)/(c*d+(-d)^(1/2
)*e^(1/2))^(1/2)/e^2-1/8*b*d*arctan((c*d-(-d)^(1/2)*e^(1/2))^(1/2)*(1+1/c/
x)^(1/2)/(c*d+(-d)^(1/2)*e^(1/2))^(1/2)/(-1+1/c/x)^(1/2))/(c*d-(-d)^(1/2)*
e^(1/2))^(3/2)/(c*d+(-d)^(1/2)*e^(1/2))^(3/2)/e-3/8*b*arctan((c*d+(-d)^(1/
2)*e^(1/2))^(1/2)*(1+1/c/x)^(1/2)/(c*d-(-d)^(1/2)*e^(1/2))^(1/2)/(-1+1/c/x
)^(1/2))/(c*d-(-d)^(1/2)*e^(1/2))^(1/2)/(c*d+(-d)^(1/2)*e^(1/2))^(1/2)/e^2
-1/8*b*d*arctan((c*d+(-d)^(1/2)*e^(1/2))^(1/2)*(1+1/c/x)^(1/2)/(c*d-(-d)^(
1/2)*e^(1/2))^(1/2)/(-1+1/c/x)^(1/2))/(c*d-(-d)^(1/2)*e^(1/2))^(3/2)/(c*d+
(-d)^(1/2)*e^(1/2))^(3/2)/e+3/16*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c
/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(1/2)
/e^(5/2)-3/16*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)
*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(1/2)/e^(5/2)+3/16*(a+b*
arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))...

```

### Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 5.56 (sec) , antiderivative size = 1823, normalized size of antiderivative = 1.43

$$\int \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Too large to display}$$

input

```
Integrate[(x^4*(a + b*ArcSech[c*x]))/(d + e*x^2)^3,x]
```



output

```

((b*e*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/((c^2*d + e)*((-I)*Sqrt[d] + Sqrt[e]*x)) + (b*e*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/((c^2*d + e)*(I*Sqrt[d] + Sqrt[e]*x)) + (4*a*d*Sqrt[e]*x)/(d + e*x^2)^2 - (10*a*Sqrt[e]*x)/(d + e*x^2) + (5*b*ArcSech[c*x])/(I*Sqrt[d] - Sqrt[e]*x) + (I*b*Sqrt[d]*ArcSech[c*x])/(Sqrt[d] + I*Sqrt[e]*x)^2 + (I*b*Sqrt[d]*ArcSech[c*x])/(I*Sqrt[d] + Sqrt[e]*x)^2 - (5*b*ArcSech[c*x])/(I*Sqrt[d] + Sqrt[e]*x) + (6*a*ArcTan[(Sqrt[e]*x)/Sqrt[d]])/Sqrt[d] + (I*b*Sqrt[e]*(2*c^2*d + e)*Log[(-4*d*Sqrt[e]*Sqrt[c^2*d + e]*(Sqrt[e] - I*c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[(1 - c*x)/(1 + c*x]) + c*Sqrt[c^2*d + e]*x*Sqrt[(1 - c*x)/(1 + c*x]))]/((2*c^2*d + e)*((-I)*Sqrt[d] + Sqrt[e]*x)))/(Sqrt[d]*(c^2*d + e)^(3/2)) - (I*b*Sqrt[e]*(2*c^2*d + e)*Log[(-4*d*Sqrt[e]*Sqrt[c^2*d + e]*(Sqrt[e] + I*c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[(1 - c*x)/(1 + c*x]) + c*Sqrt[c^2*d + e]*x*Sqrt[(1 - c*x)/(1 + c*x]))]/((2*c^2*d + e)*(I*Sqrt[d] + Sqrt[e]*x)))/(Sqrt[d]*(c^2*d + e)^(3/2)) + ((5*I)*b*Sqrt[e]*(Log[x]/Sqrt[e] - Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]]/Sqrt[e] + Log[(2*I)*Sqrt[e]*(Sqrt[d]*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x) + (Sqrt[d]*Sqrt[e] + I*c^2*d*x)/Sqrt[c^2*d + e]))/(I*Sqrt[d] + Sqrt[e]*x)]/Sqrt[c^2*d + e])/Sqrt[d] - ((5*I)*b*Sqrt[e]*(Log[x]/Sqrt[e] - Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]]/Sqrt[e] + Log[(2*Sqrt[e]*(I*Sqrt[d]*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x) + (I*Sqrt[d]*Sqrt[e] + c^2*d*x)/Sqrt...

```

### Rubi [A] (verified)

Time = 2.52 (sec) , antiderivative size = 1336, normalized size of antiderivative = 1.05, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6857, 6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx \\
 & \quad \downarrow \text{6857} \\
 & - \int \frac{a + b\operatorname{arccosh}\left(\frac{1}{cx}\right)}{\left(\frac{d}{x^2} + e\right)^3} d\frac{1}{x} \\
 & \quad \downarrow \text{6324}
 \end{aligned}$$

$$\begin{aligned}
 & - \int \left( - \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) d^3}{8(-d)^{3/2} e^{3/2} (\sqrt{-d}\sqrt{e} - \frac{d}{x})^3} - \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) d^3}{8(-d)^{3/2} e^{3/2} (\frac{d}{x} + \sqrt{-d}\sqrt{e})^3} - \frac{3(a + \operatorname{barccosh}(\frac{1}{cx})) d}{8e^2 \left(-\frac{d^2}{x^2} - ed\right)} - \frac{3(a + \operatorname{barccosh}(\frac{1}{cx}))}{16e^2 (\sqrt{-d}\sqrt{e} - \frac{d}{x})} \right) \\
 & \quad \downarrow \text{2009} \\
 & \frac{b\sqrt{-d}\sqrt{\frac{1}{cx} - 1}\sqrt{1 + \frac{1}{cx}}c}{16e^{3/2} (dc^2 + e) (\sqrt{-d}\sqrt{e} - \frac{d}{x})} + \frac{b\sqrt{-d}\sqrt{\frac{1}{cx} - 1}\sqrt{1 + \frac{1}{cx}}c}{16e^{3/2} (dc^2 + e) (\frac{d}{x} + \sqrt{-d}\sqrt{e})} + \frac{3(a + \operatorname{barccosh}(\frac{1}{cx}))}{16e^2 (\sqrt{-d}\sqrt{e} - \frac{d}{x})} - \\
 & \frac{3(a + \operatorname{barccosh}(\frac{1}{cx}))}{16e^2 (\frac{d}{x} + \sqrt{-d}\sqrt{e})} + \frac{\sqrt{-d}(a + \operatorname{barccosh}(\frac{1}{cx}))}{16e^{3/2} (\sqrt{-d}\sqrt{e} - \frac{d}{x})^2} - \frac{\sqrt{-d}(a + \operatorname{barccosh}(\frac{1}{cx}))}{16e^{3/2} (\frac{d}{x} + \sqrt{-d}\sqrt{e})^2} - \\
 & \frac{bd \arctan\left(\frac{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{\frac{1}{cx} - 1}}\right)}{8(cd - \sqrt{-d}\sqrt{e})^{3/2} (cd + \sqrt{-d}\sqrt{e})^{3/2} e} - \frac{3b \arctan\left(\frac{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{\frac{1}{cx} - 1}}\right)}{8\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{cd + \sqrt{-d}\sqrt{e}}e^2} - \\
 & \frac{bd \arctan\left(\frac{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{\frac{1}{cx} - 1}}\right)}{8(cd - \sqrt{-d}\sqrt{e})^{3/2} (cd + \sqrt{-d}\sqrt{e})^{3/2} e} - \frac{3b \arctan\left(\frac{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{\frac{1}{cx} - 1}}\right)}{8\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{cd + \sqrt{-d}\sqrt{e}}e^2} + \\
 & \frac{3(a + \operatorname{barccosh}(\frac{1}{cx})) \log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{16\sqrt{-de}e^{5/2}} - \\
 & \frac{3(a + \operatorname{barccosh}(\frac{1}{cx})) \log\left(\frac{\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})c + 1}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{16\sqrt{-de}e^{5/2}} + \\
 & \frac{3(a + \operatorname{barccosh}(\frac{1}{cx})) \log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{16\sqrt{-de}e^{5/2}} - \\
 & \frac{3(a + \operatorname{barccosh}(\frac{1}{cx})) \log\left(\frac{\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})c + 1}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{16\sqrt{-de}e^{5/2}} - \frac{3b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{16\sqrt{-de}e^{5/2}} + \\
 & \frac{3b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{16\sqrt{-de}e^{5/2}} - \frac{3b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{16\sqrt{-de}e^{5/2}} + \\
 & \frac{3b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{16\sqrt{-de}e^{5/2}}
 \end{aligned}$$

input `Int[(x^4*(a + b*ArcSech[c*x]))/(d + e*x^2)^3,x]`

output

```
(b*c*Sqrt[-d]*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]/(16*e^(3/2)*(c^2*d + e)
)*(Sqrt[-d]*Sqrt[e] - d/x)) + (b*c*Sqrt[-d]*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/
(c*x)]/(16*e^(3/2)*(c^2*d + e)*(Sqrt[-d]*Sqrt[e] + d/x)) + (Sqrt[-d]*(a +
b*ArcCosh[1/(c*x)]))/(16*e^(3/2)*(Sqrt[-d]*Sqrt[e] - d/x)^2) + (3*(a + b*
ArcCosh[1/(c*x)]))/(16*e^2*(Sqrt[-d]*Sqrt[e] - d/x)) - (Sqrt[-d]*(a + b*Ar
cCosh[1/(c*x)]))/(16*e^(3/2)*(Sqrt[-d]*Sqrt[e] + d/x)^2) - (3*(a + b*ArcCo
sh[1/(c*x)]))/(16*e^2*(Sqrt[-d]*Sqrt[e] + d/x)) - (3*b*ArcTan[(Sqrt[c*d -
Sqrt[-d]*Sqrt[e]]*Sqrt[1 + 1/(c*x)])/(Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*Sqrt[-1
+ 1/(c*x)]))/(8*Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[c*d + Sqrt[-d]*Sqrt[e]
]*e^2) - (b*d*ArcTan[(Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[1 + 1/(c*x)])/(Sqr
t[c*d + Sqrt[-d]*Sqrt[e]]*Sqrt[-1 + 1/(c*x)]))/(8*(c*d - Sqrt[-d]*Sqrt[e]
)^(3/2)*(c*d + Sqrt[-d]*Sqrt[e])^(3/2)*e) - (3*b*ArcTan[(Sqrt[c*d + Sqrt[-
d]*Sqrt[e]]*Sqrt[1 + 1/(c*x)])/(Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[-1 + 1/(
c*x)]))/(8*Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*e^2)
- (b*d*ArcTan[(Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*Sqrt[1 + 1/(c*x)])/(Sqrt[c*d
- Sqrt[-d]*Sqrt[e]]*Sqrt[-1 + 1/(c*x)]))/(8*(c*d - Sqrt[-d]*Sqrt[e])^(3/2)
)*(c*d + Sqrt[-d]*Sqrt[e])^(3/2)*e) + (3*(a + b*ArcCosh[1/(c*x)])*Log[1 -
(c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])]/(16*Sqrt[-d]
*e^(5/2)) - (3*(a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c
*x)])/(Sqrt[e] - Sqrt[c^2*d + e])]/(16*Sqrt[-d]*e^(5/2)) + (3*(a + b*A...
```

### Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6324

```
Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*((d_) + (e_.)*(x_)^2)^(p_.),
x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x],
x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] &&
(p > 0 || IGtQ[n, 0])
```

rule 6857

```
Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^n_.*(x_)^(m_.)*((d_.) + (e_.)*(x_)
^2)^(p_.), x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x
^(m + 2*(p + 1))), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0]
&& IntegersQ[m, p]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 75.24 (sec) , antiderivative size = 1960, normalized size of antiderivative = 1.54

method	result	size
parts	Expression too large to display	1960
derivativeldivides	Expression too large to display	1983
default	Expression too large to display	1983

input `int(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^3,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & a * \left( \frac{-5/8/e*x^3 - 3/8/e^2*d*x}{(e*x^2+d)^2} + \frac{3/8/e^2/(d*e)^{(1/2)} * \arctan(x*e/(d*e)^{(1/2)})}{(e*x^2+d)^2} \right) + b/c^5 * \left( -\frac{1}{8} * c^7 * x * (3*d^2*c^4 * \operatorname{arcsech}(c*x) + 5*c^4*d*e * \operatorname{arcsech}(c*x)) \right. \\
 & * x^2 - \left. \frac{-(c*x-1)/c/x)^{(1/2)} * ((c*x+1)/c/x)^{(1/2)} * c^3*d*e*x - (-(c*x-1)/c/x)^{(1/2)} * ((c*x+1)/c/x)^{(1/2)} * e^2*c^3*x^3 + 3*c^2*d*e * \operatorname{arcsech}(c*x) + 5*e^2 * \operatorname{arcsech}(c*x)}{e^2/(c^2*e*x^2+c^2*d)^2/(c^2*d+e)} \right. \\
 & - \left. \frac{3/8 * (-(c^2*d-2*(e*(c^2*d+e))^{(1/2)}+2*e)*d)^{(1/2)} * (c^2*d*(e*(c^2*d+e))^{(1/2)}+2*c^2*d*e+2*e^2+2*(e*(c^2*d+e))^{(1/2)}*e)}{c^3 * \operatorname{arctanh}(c*d*(1/c/x+(-1+1/c/x)^{(1/2)}*(1+1/c/x)^{(1/2)})} \right) / \left( \frac{-(c^2*d+2*(e*(c^2*d+e))^{(1/2)}-2*e)*d)^{(1/2)}}{(c^2*d+e)^2/e^2/d^2-3/8 * ((c^2*d+2*(e*(c^2*d+e))^{(1/2)}+2*e)*d)^{(1/2)} * (-c^2*d*(e*(c^2*d+e))^{(1/2)}+2*c^2*d*e+2*e^2-2*(e*(c^2*d+e))^{(1/2)}*e)} \right. \\
 & * \left. \frac{c^3 * \operatorname{arctan}(c*d*(1/c/x+(-1+1/c/x)^{(1/2)}*(1+1/c/x)^{(1/2)})}{((c^2*d+2*(e*(c^2*d+e))^{(1/2)}+2*e)*d)^{(1/2)}} \right) / \left( \frac{c^2*d+e}{e/d^3-1/2 * (-c^2*d-2*(e*(c^2*d+e))^{(1/2)}+2*e)*d)^{(1/2)} * (c^2*d*(e*(c^2*d+e))^{(1/2)}+2*c^2*d*e+2*e^2+2*(e*(c^2*d+e))^{(1/2)}*e)} \right. \\
 & * \left. \frac{c * \operatorname{arctanh}(c*d*(1/c/x+(-1+1/c/x)^{(1/2)}*(1+1/c/x)^{(1/2)})}{((-c^2*d+2*(e*(c^2*d+e))^{(1/2)}-2*e)*d)^{(1/2)}} \right) / \left( \frac{c^2*d+e}{e/d^3-1/2 * (-c^2*d-2*(e*(c^2*d+e))^{(1/2)}+2*e)*d)^{(1/2)} * (c^2*d*(e*(c^2*d+e))^{(1/2)}+2*c^2*d*e+2*e^2+2*(e*(c^2*d+e))^{(1/2)}*e)} \right. \\
 & * \left. \frac{c * \operatorname{arctan}(c*d*(1/c/x+(-1+1/c/x)^{(1/2)}*(1+1/c/x)^{(1/2)})}{((c^2*d+2*(e*(c^2*d+e))^{(1/2)}+2*e)*d)^{(1/2)}} \right) / \left( \frac{c^2*d+e}{e/d^3-1/2 * ((c^2*d+2*(e*(c^2*d+e))^{(1/2)}+2*e)*d)^{(1/2)} * (c^2*d*(e*(c^2*d+e))^{(1/2)}+2*c^2*d*e+2*e^2+2*(e*(c^2*d+e))^{(1/2)}*e)} \right)
 \end{aligned}$$

**Fricas [F]**

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^4}{(ex^2 + d)^3} dx$$

input `integrate(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="fricas")`

output `integral((b*x^4*arcsech(c*x) + a*x^4)/(e^3*x^6 + 3*d*e^2*x^4 + 3*d^2*e*x^2 + d^3), x)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Timed out}$$

input `integrate(x**4*(a+b*asech(c*x))/(e*x**2+d)**3,x)`

output `Timed out`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Exception raised: ValueError}$$

input `integrate(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^4}{(ex^2 + d)^3} dx$$

input `integrate(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^4/(e*x^2 + d)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \int \frac{x^4(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^3} dx$$

input `int((x^4*(a + b*acosh(1/(c*x))))/(d + e*x^2)^3,x)`

output `int((x^4*(a + b*acosh(1/(c*x))))/(d + e*x^2)^3, x)`

**Reduce [F]**

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx$$

$$= \frac{3\sqrt{e}\sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e}\sqrt{d}}\right) a d^2 + 6\sqrt{e}\sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e}\sqrt{d}}\right) a d e x^2 + 3\sqrt{e}\sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e}\sqrt{d}}\right) a e^2 x^4 + 8 \left( \int \frac{a \operatorname{sech}^{-1}(cx)}{e^3 x^6 + 3d e^2 x^4} dx \right)}{8d}$$

input `int(x^4*(a+b*asech(c*x))/(e*x^2+d)^3,x)`

output

```
(3*sqrt(e)*sqrt(d)*atan((e*x)/(sqrt(e)*sqrt(d)))*a*d**2 + 6*sqrt(e)*sqrt(d)
)*atan((e*x)/(sqrt(e)*sqrt(d)))*a*d*e*x**2 + 3*sqrt(e)*sqrt(d)*atan((e*x)/
(sqrt(e)*sqrt(d)))*a*e**2*x**4 + 8*int((asech(c*x)*x**4)/(d**3 + 3*d**2*e*
x**2 + 3*d*e**2*x**4 + e**3*x**6),x)*b*d**3*e**3 + 16*int((asech(c*x)*x**4
)/(d**3 + 3*d**2*e*x**2 + 3*d*e**2*x**4 + e**3*x**6),x)*b*d**2*e**4*x**2 +
8*int((asech(c*x)*x**4)/(d**3 + 3*d**2*e*x**2 + 3*d*e**2*x**4 + e**3*x**6
),x)*b*d*e**5*x**4 - 3*a*d**2*e*x - 5*a*d*e**2*x**3)/(8*d*e**3*(d**2 + 2*d
*e*x**2 + e**2*x**4))
```

$$3.127 \quad \int \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx$$

Optimal result	1039
Mathematica [C] (warning: unable to verify)	1040
Rubi [A] (verified)	1041
Maple [C] (warning: unable to verify)	1044
Fricas [F]	1045
Sympy [F(-1)]	1045
Maxima [F(-2)]	1045
Giac [F]	1046
Mupad [F(-1)]	1046
Reduce [F]	1046

### Optimal result

Integrand size = 21, antiderivative size = 1276

$$\int \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Too large to display}$$



output

```

1/16*b*c*(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)/(-d)^(1/2)/e^(1/2)/(c^2*d+e)/((-
d)^(1/2)*e^(1/2)-d/x)+1/16*b*c*(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)/(-d)^(1/2)
/e^(1/2)/(c^2*d+e)/((-d)^(1/2)*e^(1/2)+d/x)+1/16*(a+b*arcsech(c*x))/(-d)^(
1/2)/e^(1/2)/((-d)^(1/2)*e^(1/2)-d/x)^2+1/16*(a+b*arcsech(c*x))/d/e/((-d)^(
1/2)*e^(1/2)-d/x)-1/16*(a+b*arcsech(c*x))/(-d)^(1/2)/e^(1/2)/((-d)^(1/2)*
e^(1/2)+d/x)^2-1/16*(a+b*arcsech(c*x))/d/e/((-d)^(1/2)*e^(1/2)+d/x)+1/8*b*
arctan((c*d-(-d)^(1/2)*e^(1/2))^(1/2)*(1+1/c/x)^(1/2)/(c*d+(-d)^(1/2)*e^(1
/2))^(1/2)/(-1+1/c/x)^(1/2))/(c*d-(-d)^(1/2)*e^(1/2))^(3/2)/(c*d+(-d)^(1/2)
)*e^(1/2))^(3/2)-1/8*b*arctan((c*d-(-d)^(1/2)*e^(1/2))^(1/2)*(1+1/c/x)^(1/
2)/(c*d+(-d)^(1/2)*e^(1/2))^(1/2)/(-1+1/c/x)^(1/2))/d/(c*d-(-d)^(1/2)*e^(1
/2))^(1/2)/(c*d+(-d)^(1/2)*e^(1/2))^(1/2)/e+1/8*b*arctan((c*d+(-d)^(1/2)*e
^(1/2))^(1/2)*(1+1/c/x)^(1/2)/(c*d-(-d)^(1/2)*e^(1/2))^(1/2)/(-1+1/c/x)^(1
/2))/c*d-(-d)^(1/2)*e^(1/2))^(3/2)/(c*d+(-d)^(1/2)*e^(1/2))^(3/2)-1/8*b*a
rctan((c*d+(-d)^(1/2)*e^(1/2))^(1/2)*(1+1/c/x)^(1/2)/(c*d-(-d)^(1/2)*e^(1/
2))^(1/2)/(-1+1/c/x)^(1/2))/d/(c*d-(-d)^(1/2)*e^(1/2))^(1/2)/(c*d+(-d)^(1/
2)*e^(1/2))^(1/2)/e-1/16*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1
/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(3/2)/e^(3/2)
+1/16*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/
x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(3/2)/e^(3/2)-1/16*(a+b*arcsech(
c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/...

```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 6.07 (sec) , antiderivative size = 2030, normalized size of antiderivative = 1.59

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Result too large to show}$$

input

```
Integrate[(x^2*(a + b*ArcSech[c*x]))/(d + e*x^2)^3,x]
```

output

```

-1/4*(a*x)/(e*(d + e*x^2)^2) + (a*x)/(8*d*e*(d + e*x^2)) + (a*ArcTan[(Sqrt
[e]*x)/Sqrt[d]])/(8*d^(3/2)*e^(3/2)) + b*(((1/16*I)*((-I)*Sqrt[e]*Sqrt[(
1 - c*x)/(1 + c*x)]*(1 + c*x))/(Sqrt[d]*(c^2*d + e)*((-I)*Sqrt[d] + Sqrt[e
]*x)) - ArcSech[c*x]/(Sqrt[e]*((-I)*Sqrt[d] + Sqrt[e]*x)^2) + Log[x]/(d*Sq
rt[e]) - Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)
]]/(d*Sqrt[e]) + ((2*c^2*d + e)*Log[(-4*d*Sqrt[e]*Sqrt[c^2*d + e]*(Sqrt[e]
- I*c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[(1 - c*x)/(1 + c*x]) + c*Sqrt[c^2
*d + e]*x*Sqrt[(1 - c*x)/(1 + c*x]))]/((2*c^2*d + e)*((-I)*Sqrt[d] + Sqrt[
e]*x)))]/(d*(c^2*d + e)^(3/2)))/(Sqrt[d]*e) + ((I/16)*((I*Sqrt[e]*Sqrt[(1
- c*x)/(1 + c*x)]*(1 + c*x))/(Sqrt[d]*(c^2*d + e)*(I*Sqrt[d] + Sqrt[e]*x)
) - ArcSech[c*x]/(Sqrt[e]*(I*Sqrt[d] + Sqrt[e]*x)^2) + Log[x]/(d*Sqrt[e])
- Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]]/(d*Sq
rt[e]) + ((2*c^2*d + e)*Log[(-4*d*Sqrt[e]*Sqrt[c^2*d + e]*(Sqrt[e] + I*c^2
*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[(1 - c*x)/(1 + c*x]) + c*Sqrt[c^2*d + e
]*x*Sqrt[(1 - c*x)/(1 + c*x]))]/((2*c^2*d + e)*(I*Sqrt[d] + Sqrt[e]*x)))]/(
d*(c^2*d + e)^(3/2)))/(Sqrt[d]*e) - (- (ArcSech[c*x]/(I*Sqrt[d]*Sqrt[e] +
e*x)) + (I*(Log[x]/Sqrt[e] - Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[
(1 - c*x)/(1 + c*x)]]/Sqrt[e] + Log[((2*I)*Sqrt[e]*(Sqrt[d]*Sqrt[(1 - c*x)
/(1 + c*x)]*(1 + c*x) + (Sqrt[d]*Sqrt[e] + I*c^2*d*x)/Sqrt[c^2*d + e]))/(I
*Sqrt[d] + Sqrt[e]*x)]/Sqrt[c^2*d + e]))/Sqrt[d])/(16*d*e) - (- (ArcSech...

```

### Rubi [A] (verified)

Time = 3.61 (sec) , antiderivative size = 1340, normalized size of antiderivative = 1.05, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6857, 6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx \\
 & \quad \downarrow \text{6857} \\
 & - \int \frac{a + b \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\left(\frac{d}{x^2} + e\right)^3} d \frac{1}{x} \\
 & \quad \downarrow \text{6374}
 \end{aligned}$$

$$\begin{aligned}
& - \int \left( \frac{a + \operatorname{barccosh}\left(\frac{1}{cx}\right)}{d\left(\frac{d}{x^2} + e\right)^2} - \frac{e\left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right)}{d\left(\frac{d}{x^2} + e\right)^3} \right) d\frac{1}{x} \\
& \quad \downarrow \text{2009} \\
& \frac{b\sqrt{\frac{1}{cx} - 1}\sqrt{1 + \frac{1}{cx}c}}{16\sqrt{-d}\sqrt{e}(dc^2 + e)\left(\sqrt{-d}\sqrt{e} - \frac{d}{x}\right)} + \frac{b\sqrt{\frac{1}{cx} - 1}\sqrt{1 + \frac{1}{cx}c}}{16\sqrt{-d}\sqrt{e}(dc^2 + e)\left(\frac{d}{x} + \sqrt{-d}\sqrt{e}\right)} + \\
& \frac{a + \operatorname{barccosh}\left(\frac{1}{cx}\right)}{16de\left(\sqrt{-d}\sqrt{e} - \frac{d}{x}\right)} - \frac{a + \operatorname{barccosh}\left(\frac{1}{cx}\right)}{16de\left(\frac{d}{x} + \sqrt{-d}\sqrt{e}\right)} + \frac{a + \operatorname{barccosh}\left(\frac{1}{cx}\right)}{16\sqrt{-d}\sqrt{e}\left(\sqrt{-d}\sqrt{e} - \frac{d}{x}\right)^2} - \\
& \frac{a + \operatorname{barccosh}\left(\frac{1}{cx}\right)}{16\sqrt{-d}\sqrt{e}\left(\frac{d}{x} + \sqrt{-d}\sqrt{e}\right)^2} - \frac{b \arctan\left(\frac{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{\frac{1}{cx} - 1}}\right)}{8d\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{cd + \sqrt{-d}\sqrt{e}}} + \\
& \frac{b \arctan\left(\frac{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{\frac{1}{cx} - 1}}\right)}{8\left(cd - \sqrt{-d}\sqrt{e}\right)^{3/2}\left(cd + \sqrt{-d}\sqrt{e}\right)^{3/2}} - \frac{b \arctan\left(\frac{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{\frac{1}{cx} - 1}}\right)}{8d\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{cd + \sqrt{-d}\sqrt{e}}} + \\
& \frac{b \arctan\left(\frac{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{\frac{1}{cx} - 1}}\right)}{8\left(cd - \sqrt{-d}\sqrt{e}\right)^{3/2}\left(cd + \sqrt{-d}\sqrt{e}\right)^{3/2}} - \frac{\left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) \log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{16(-d)^{3/2}e^{3/2}} + \\
& \frac{\left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) \log\left(\frac{\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)c}{\sqrt{e} - \sqrt{dc^2 + e}} + 1\right)}{16(-d)^{3/2}e^{3/2}} - \\
& \frac{\left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) \log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{16(-d)^{3/2}e^{3/2}} + \\
& \frac{\left(a + \operatorname{barccosh}\left(\frac{1}{cx}\right)\right) \log\left(\frac{\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)c}{\sqrt{e} + \sqrt{dc^2 + e}} + 1\right)}{16(-d)^{3/2}e^{3/2}} + \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{16(-d)^{3/2}e^{3/2}} - \\
& \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} - \sqrt{dc^2 + e}}\right)}{16(-d)^{3/2}e^{3/2}} + \frac{b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{16(-d)^{3/2}e^{3/2}} - \\
& \frac{b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\sqrt{e} + \sqrt{dc^2 + e}}\right)}{16(-d)^{3/2}e^{3/2}}
\end{aligned}$$

input

```
Int[(x^2*(a + b*ArcSech[c*x]))/(d + e*x^2)^3,x]
```

output

```
(b*c*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]/(16*Sqrt[-d]*Sqrt[e]*(c^2*d + e)
)*(Sqrt[-d]*Sqrt[e] - d/x)) + (b*c*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]/(
16*Sqrt[-d]*Sqrt[e]*(c^2*d + e)*(Sqrt[-d]*Sqrt[e] + d/x)) + (a + b*ArcCosh
[1/(c*x)]/(16*Sqrt[-d]*Sqrt[e]*(Sqrt[-d]*Sqrt[e] - d/x)^2) + (a + b*ArcCo
sh[1/(c*x)]/(16*d*e*(Sqrt[-d]*Sqrt[e] - d/x)) - (a + b*ArcCosh[1/(c*x)]/
(16*Sqrt[-d]*Sqrt[e]*(Sqrt[-d]*Sqrt[e] + d/x)^2) - (a + b*ArcCosh[1/(c*x)]
)/(16*d*e*(Sqrt[-d]*Sqrt[e] + d/x)) + (b*ArcTan[(Sqrt[c*d - Sqrt[-d]*Sqrt[
e]]*Sqrt[1 + 1/(c*x)]/(Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*Sqrt[-1 + 1/(c*x)])))
/(8*(c*d - Sqrt[-d]*Sqrt[e])^(3/2)*(c*d + Sqrt[-d]*Sqrt[e])^(3/2)) - (b*Ar
cTan[(Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[1 + 1/(c*x)]/(Sqrt[c*d + Sqrt[-d]
*Sqrt[e]]*Sqrt[-1 + 1/(c*x)])))/(8*d*Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[c*d
+ Sqrt[-d]*Sqrt[e]]*e) + (b*ArcTan[(Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*Sqrt[1 +
1/(c*x)]/(Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[-1 + 1/(c*x)])))/(8*(c*d - S
qrt[-d]*Sqrt[e])^(3/2)*(c*d + Sqrt[-d]*Sqrt[e])^(3/2)) - (b*ArcTan[(Sqrt[c
*d + Sqrt[-d]*Sqrt[e]]*Sqrt[1 + 1/(c*x)]/(Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sq
rt[-1 + 1/(c*x)])))/(8*d*Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[c*d + Sqrt[-d]*
Sqrt[e]]*e) - ((a + b*ArcCosh[1/(c*x)])*Log[1 - (c*Sqrt[-d]*E^ArcCosh[1/(c
*x)]/(Sqrt[e] - Sqrt[c^2*d + e]))]/(16*(-d)^(3/2)*e^(3/2)) + ((a + b*ArcC
osh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^ArcCosh[1/(c*x)]/(Sqrt[e] - Sqrt[c^2*d
+ e]))]/(16*(-d)^(3/2)*e^(3/2)) - ((a + b*ArcCosh[1/(c*x)])*Log[1 - (...
```

### Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6374

```
Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^ (n_.)*((f_.)*(x_))^(m_.)*((d_) + (e
_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

rule 6857

```
Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^ (n_.)*(x_)^(m_.)*((d_.) + (e_.)*(x
_)^2)^(p_.), x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x
^(m + 2*(p + 1))), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0
] && IntegersQ[m, p]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 72.88 (sec) , antiderivative size = 1398, normalized size of antiderivative = 1.10

method	result	size
parts	Expression too large to display	1398
derivativeldivides	Expression too large to display	1417
default	Expression too large to display	1417

input `int(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^3,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & a \left( \frac{1}{8} \frac{d^3}{d^3} - \frac{1}{8} \frac{e^3}{e^3} \right) / (e^2 x^2 + d)^2 + \frac{1}{8} \frac{e}{d} \frac{d}{(d e)^{1/2}} \arctan \left( \frac{x e}{(d e)^{1/2}} \right) \\
 & + b/c^3 \left( \frac{1}{8} c^5 x^* (c^4 d e \operatorname{arcsech}(c x) x^2 - d^2 c^4 \operatorname{arcsech}(c x) - ((c x - 1)/c/x)^{1/2} ((c x + 1)/c/x)^{1/2} e^2 c^3 x^3 - ((c x - 1)/c/x)^{1/2} ((c x + 1)/c/x)^{1/2} c^3 d e x + e^2 \operatorname{arcsech}(c x) c^2 x^2 - c^2 d e \operatorname{arcsech}(c x)) / d \right. \\
 & / e / (c^2 d + e) / (c^2 e x^2 + c^2 d)^2 - 1/16 / d / (c^2 d + e) c^4 \operatorname{sum}(\_R1 / (\_R1^2 c^2 d + c^2 d + 2 e) * (\operatorname{arcsech}(c x) * \ln((\_R1 - 1/c/x - (-1 + 1/c/x)^{1/2}) * (1 + 1/c/x)^{1/2})) / \_R1) \\
 & + \operatorname{dilog}((\_R1 - 1/c/x - (-1 + 1/c/x)^{1/2}) * (1 + 1/c/x)^{1/2}) / \_R1), \_R1 = \operatorname{RootOf}(c^2 d * \_Z^4 + (2 c^2 d + 4 e) * \_Z^2 + c^2 d) - 1/16 / (c^2 d + e) / e c^6 \operatorname{sum}(\_R1 / (\_R1^2 c^2 d + c^2 d + 2 e) * (\operatorname{arcsech}(c x) * \ln((\_R1 - 1/c/x - (-1 + 1/c/x)^{1/2}) * (1 + 1/c/x)^{1/2})) / \_R1) \\
 & + \operatorname{dilog}((\_R1 - 1/c/x - (-1 + 1/c/x)^{1/2}) * (1 + 1/c/x)^{1/2}) / \_R1), \_R1 = \operatorname{RootOf}(c^2 d * \_Z^4 + (2 c^2 d + 4 e) * \_Z^2 + c^2 d) + 1/16 / d / (c^2 d + e) c^4 \operatorname{sum}(1 / \_R1 / (\_R1^2 c^2 d + c^2 d + 2 e) * (\operatorname{arcsech}(c x) * \ln((\_R1 - 1/c/x - (-1 + 1/c/x)^{1/2}) * (1 + 1/c/x)^{1/2})) / \_R1) \\
 & + \operatorname{dilog}((\_R1 - 1/c/x - (-1 + 1/c/x)^{1/2}) * (1 + 1/c/x)^{1/2}) / \_R1), \_R1 = \operatorname{RootOf}(c^2 d * \_Z^4 + (2 c^2 d + 4 e) * \_Z^2 + c^2 d) + 1/8 * (- (c^2 d - 2 * (e * (c^2 d + e))^{1/2} + 2 e) * d)^{1/2} * (c^2 d + 2 * (e * (c^2 d + e))^{1/2} + 2 e) * c * \operatorname{arctanh}(c * d * (1/c/x + (-1 + 1/c/x)^{1/2}) * (1 + 1/c/x)^{1/2}) / ((-c^2 d + 2 * (e * (c^2 d + e))^{1/2} - 2 e) * d)^{1/2} \\
 & / (c^2 d + e) / e / d^3 - 1/8 * (- (c^2 d - 2 * (e * (c^2 d + e))^{1/2} + 2 e) * d)^{1/2} * (c^2 d * (e * (c^2 d + e))^{1/2} + 2 c^2 d e + 2 e^2 + 2 * (e * (c^2 d + e))^{1/2} * e) * c * \operatorname{arctanh}(c * d * (1/c/x + (-1 + 1/c/x)^{1/2}) * (1 + 1/c/x)^{1/2}) / ((-c^2 d + 2 * (e * (c^2 d + e))^{1/2} - 2 e) * d)^{1/2} \\
 & / (c^2 d + e)^2 / e / d^3 + 1/8 * ((c^2 d + 2 * (e * (c^2 d + e))^{1/2} \dots
 \end{aligned}$$

**Fricas [F]**

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^2}{(ex^2 + d)^3} dx$$

input `integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="fricas")`

output `integral((b*x^2*arcsech(c*x) + a*x^2)/(e^3*x^6 + 3*d*e^2*x^4 + 3*d^2*e*x^2 + d^3), x)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Timed out}$$

input `integrate(x**2*(a+b*asech(c*x))/(e*x**2+d)**3,x)`

output `Timed out`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \text{Exception raised: ValueError}$$

input `integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^2}{(ex^2 + d)^3} dx$$

input `integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^2/(e*x^2 + d)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx = \int \frac{x^2(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^3} dx$$

input `int((x^2*(a + b*acosh(1/(c*x))))/(d + e*x^2)^3,x)`

output `int((x^2*(a + b*acosh(1/(c*x))))/(d + e*x^2)^3, x)`

**Reduce [F]**

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^3} dx$$

$$= \frac{\sqrt{e} \sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e} \sqrt{d}}\right) a d^2 + 2\sqrt{e} \sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e} \sqrt{d}}\right) a d e x^2 + \sqrt{e} \sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e} \sqrt{d}}\right) a e^2 x^4 + 8 \left( \int \frac{a \operatorname{sech}(cx)}{e^3 x^6 + 3d e^2 x^4 + \dots} dx \right)}{8d^2 e^2}$$

input `int(x^2*(a+b*asech(c*x))/(e*x^2+d)^3,x)`

output

```
(sqrt(e)*sqrt(d)*atan((e*x)/(sqrt(e)*sqrt(d)))*a*d**2 + 2*sqrt(e)*sqrt(d)*
atan((e*x)/(sqrt(e)*sqrt(d)))*a*d*e*x**2 + sqrt(e)*sqrt(d)*atan((e*x)/(sqr
t(e)*sqrt(d)))*a*e**2*x**4 + 8*int((asech(c*x)*x**2)/(d**3 + 3*d**2*e*x**2
+ 3*d*e**2*x**4 + e**3*x**6),x)*b*d**4*e**2 + 16*int((asech(c*x)*x**2)/(d
**3 + 3*d**2*e*x**2 + 3*d*e**2*x**4 + e**3*x**6),x)*b*d**3*e**3*x**2 + 8*i
nt((asech(c*x)*x**2)/(d**3 + 3*d**2*e*x**2 + 3*d*e**2*x**4 + e**3*x**6),x)
*b*d**2*e**4*x**4 - a*d**2*e*x + a*d*e**2*x**3)/(8*d**2*e**2*(d**2 + 2*d*e
*x**2 + e**2*x**4))
```



**3.128**       $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex^2)^3} dx$

Optimal result	1048
Mathematica [C] (warning: unable to verify)	1049
Rubi [A] (verified)	1050
Maple [C] (warning: unable to verify)	1053
Fricas [F]	1054
Sympy [F(-1)]	1054
Maxima [F(-2)]	1054
Giac [F]	1055
Mupad [F(-1)]	1055
Reduce [F]	1055

**Optimal result**

Integrand size = 18, antiderivative size = 1272

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{(d + ex^2)^3} dx = \text{Too large to display}$$

output

```

1/16*b*c*e^(1/2)*(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)/(-d)^(3/2)/(c^2*d+e)/((-
d)^(1/2)*e^(1/2)-d/x)+1/16*b*c*e^(1/2)*(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2)/(-
d)^(3/2)/(c^2*d+e)/((-d)^(1/2)*e^(1/2)+d/x)+1/16*e^(1/2)*(a+b*arcsech(c*x)
)/(-d)^(3/2)/((-d)^(1/2)*e^(1/2)-d/x)^2-5/16*(a+b*arcsech(c*x))/d^2/((-d)^(
1/2)*e^(1/2)-d/x)-1/16*e^(1/2)*(a+b*arcsech(c*x))/(-d)^(3/2)/((-d)^(1/2)*
e^(1/2)+d/x)^2+5/16*(a+b*arcsech(c*x))/d^2/((-d)^(1/2)*e^(1/2)+d/x)+5/8*b*
arctan((c*d-(-d)^(1/2)*e^(1/2))^(1/2)*(1+1/c/x)^(1/2)/(c*d+(-d)^(1/2)*e^(1
/2))^(1/2)/(-1+1/c/x)^(1/2))/d^2/(c*d-(-d)^(1/2)*e^(1/2))^(1/2)/(c*d+(-d)^(
1/2)*e^(1/2))^(1/2)-1/8*b*e*arctan((c*d-(-d)^(1/2)*e^(1/2))^(1/2)*(1+1/c/
x)^(1/2)/(c*d+(-d)^(1/2)*e^(1/2))^(1/2)/(-1+1/c/x)^(1/2))/d/(c*d-(-d)^(1/2
)*e^(1/2))^(3/2)/(c*d+(-d)^(1/2)*e^(1/2))^(3/2)+5/8*b*arctan((c*d+(-d)^(1/
2)*e^(1/2))^(1/2)*(1+1/c/x)^(1/2)/(c*d-(-d)^(1/2)*e^(1/2))^(1/2)/(-1+1/c/x
)^(1/2))/d^2/(c*d-(-d)^(1/2)*e^(1/2))^(1/2)/(c*d+(-d)^(1/2)*e^(1/2))^(1/2)
-1/8*b*e*arctan((c*d+(-d)^(1/2)*e^(1/2))^(1/2)*(1+1/c/x)^(1/2)/(c*d-(-d)^(
1/2)*e^(1/2))^(1/2)/(-1+1/c/x)^(1/2))/d/(c*d-(-d)^(1/2)*e^(1/2))^(3/2)/(c*
d+(-d)^(1/2)*e^(1/2))^(3/2)+3/16*(a+b*arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c
/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(5/2)
/e^(1/2)-3/16*(a+b*arcsech(c*x))*ln(1+c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)
*(1+1/c/x)^(1/2))/(e^(1/2)-(c^2*d+e)^(1/2)))/(-d)^(5/2)/e^(1/2)+3/16*(a+b*
arcsech(c*x))*ln(1-c*(-d)^(1/2)*(1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))...

```

**Mathematica [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 6.05 (sec) , antiderivative size = 1813, normalized size of antiderivative = 1.43

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^3} dx = \text{Too large to display}$$

input

```
Integrate[(a + b*ArcSech[c*x])/(d + e*x^2)^3,x]
```

output

```

((b*Sqrt[d]*Sqrt[e]*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x))/((c^2*d + e)*((-I)
)*Sqrt[d] + Sqrt[e]*x)) + (b*Sqrt[d]*Sqrt[e]*Sqrt[(1 - c*x)/(1 + c*x)]*(1
+ c*x))/((c^2*d + e)*(I*Sqrt[d] + Sqrt[e]*x)) + (4*a*d^(3/2)*x)/(d + e*x^2
)^2 + (6*a*Sqrt[d]*x)/(d + e*x^2) + (I*b*d*ArcSech[c*x])/(Sqrt[e]*(Sqrt[d]
+ I*Sqrt[e]*x)^2) + (I*b*d*ArcSech[c*x])/(Sqrt[e]*(I*Sqrt[d] + Sqrt[e]*x)
^2) + (3*b*Sqrt[d]*ArcSech[c*x])/((-I)*Sqrt[d]*Sqrt[e] + e*x) + (3*b*Sqrt[
d]*ArcSech[c*x])/(I*Sqrt[d]*Sqrt[e] + e*x) + (6*a*ArcTan[(Sqrt[e]*x)/Sqrt[
d]])/Sqrt[e] + (I*b*(2*c^2*d + e)*Log[(-4*d*Sqrt[e]*Sqrt[c^2*d + e]*(Sqrt[
e] - I*c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[(1 - c*x)/(1 + c*x]) + c*Sqrt[
c^2*d + e]*x*Sqrt[(1 - c*x)/(1 + c*x]))]/((2*c^2*d + e)*((-I)*Sqrt[d] + Sq
rt[e]*x)))/(c^2*d + e)^(3/2) - (I*b*(2*c^2*d + e)*Log[(-4*d*Sqrt[e]*Sqrt[
c^2*d + e]*(Sqrt[e] + I*c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[(1 - c*x)/(1
+ c*x]) + c*Sqrt[c^2*d + e]*x*Sqrt[(1 - c*x)/(1 + c*x]))]/((2*c^2*d + e)*
(I*Sqrt[d] + Sqrt[e]*x)))/(c^2*d + e)^(3/2) - (3*I)*b*(Log[x]/Sqrt[e] - Lo
g[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]]/Sqrt[e] +
Log[((2*I)*Sqrt[e]*(Sqrt[d]*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x) + (Sqrt[d]
)*Sqrt[e] + I*c^2*d*x)/Sqrt[c^2*d + e]))/(I*Sqrt[d] + Sqrt[e]*x)/Sqrt[c^2
*d + e] + (3*I)*b*(Log[x]/Sqrt[e] - Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c
*x*Sqrt[(1 - c*x)/(1 + c*x)]]/Sqrt[e] + Log[(2*Sqrt[e]*(I*Sqrt[d]*Sqrt[(1
- c*x)/(1 + c*x)]*(1 + c*x) + (I*Sqrt[d]*Sqrt[e] + c^2*d*x)/Sqrt[c^2*d ...

```

### Rubi [A] (verified)

Time = 4.26 (sec) , antiderivative size = 1336, normalized size of antiderivative = 1.05, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {6847, 6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^3} dx \\
 & \quad \downarrow 6847 \\
 & - \int \frac{a + b \operatorname{arccosh}\left(\frac{1}{cx}\right)}{\left(\frac{d}{x^2} + e\right)^3 x^4} d \frac{1}{x} \\
 & \quad \downarrow 6374
 \end{aligned}$$

$$\begin{aligned}
 & - \int \left( \frac{(a + \operatorname{barccosh}(\frac{1}{cx})) e^2}{d^2 (\frac{d}{x^2} + e)^3} - \frac{2(a + \operatorname{barccosh}(\frac{1}{cx})) e}{d^2 (\frac{d}{x^2} + e)^2} + \frac{a + \operatorname{barccosh}(\frac{1}{cx})}{d^2 (\frac{d}{x^2} + e)} \right) d \frac{1}{x} \\
 & \qquad \qquad \qquad \downarrow \text{2009} \\
 & \frac{b\sqrt{e}\sqrt{\frac{1}{cx} - 1}\sqrt{1 + \frac{1}{cx}c}}{16(-d)^{3/2}(dc^2 + e)(\sqrt{-d}\sqrt{e} - \frac{d}{x})} + \frac{b\sqrt{e}\sqrt{\frac{1}{cx} - 1}\sqrt{1 + \frac{1}{cx}c}}{16(-d)^{3/2}(dc^2 + e)(\frac{d}{x} + \sqrt{-d}\sqrt{e})} - \\
 & \frac{5(a + \operatorname{barccosh}(\frac{1}{cx}))}{16d^2(\sqrt{-d}\sqrt{e} - \frac{d}{x})} + \frac{5(a + \operatorname{barccosh}(\frac{1}{cx}))}{16d^2(\frac{d}{x} + \sqrt{-d}\sqrt{e})} + \frac{\sqrt{e}(a + \operatorname{barccosh}(\frac{1}{cx}))}{16(-d)^{3/2}(\sqrt{-d}\sqrt{e} - \frac{d}{x})^2} - \\
 & \frac{\sqrt{e}(a + \operatorname{barccosh}(\frac{1}{cx}))}{16(-d)^{3/2}(\frac{d}{x} + \sqrt{-d}\sqrt{e})^2} - \frac{be \arctan\left(\frac{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{\frac{1}{cx} - 1}}\right)}{8d(cd - \sqrt{-d}\sqrt{e})^{3/2}(cd + \sqrt{-d}\sqrt{e})^{3/2}} + \\
 & \frac{5b \arctan\left(\frac{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{\frac{1}{cx} - 1}}\right)}{8d^2\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{cd + \sqrt{-d}\sqrt{e}}} - \frac{be \arctan\left(\frac{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{\frac{1}{cx} - 1}}\right)}{8d(cd - \sqrt{-d}\sqrt{e})^{3/2}(cd + \sqrt{-d}\sqrt{e})^{3/2}} + \\
 & \frac{5b \arctan\left(\frac{\sqrt{cd + \sqrt{-d}\sqrt{e}}\sqrt{1 + \frac{1}{cx}}}{\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{\frac{1}{cx} - 1}}\right)}{8d^2\sqrt{cd - \sqrt{-d}\sqrt{e}}\sqrt{cd + \sqrt{-d}\sqrt{e}}} + \frac{3(a + \operatorname{barccosh}(\frac{1}{cx})) \log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{dc^2 + e}}}\right)}{16(-d)^{5/2}\sqrt{e}} - \\
 & \frac{3(a + \operatorname{barccosh}(\frac{1}{cx})) \log\left(\frac{\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})c}{\sqrt{e - \sqrt{dc^2 + e}}} + 1\right)}{16(-d)^{5/2}\sqrt{e}} + \\
 & \frac{3(a + \operatorname{barccosh}(\frac{1}{cx})) \log\left(1 - \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}}\right)}{16(-d)^{5/2}\sqrt{e}} - \\
 & \frac{3(a + \operatorname{barccosh}(\frac{1}{cx})) \log\left(\frac{\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})c}{\sqrt{e + \sqrt{dc^2 + e}}} + 1\right)}{16(-d)^{5/2}\sqrt{e}} - \frac{3b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{dc^2 + e}}}\right)}{16(-d)^{5/2}\sqrt{e}} + \\
 & \frac{3b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e - \sqrt{dc^2 + e}}}\right)}{16(-d)^{5/2}\sqrt{e}} - \frac{3b \operatorname{PolyLog}\left(2, -\frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}}\right)}{16(-d)^{5/2}\sqrt{e}} + \\
 & \frac{3b \operatorname{PolyLog}\left(2, \frac{c\sqrt{-de} \operatorname{arccosh}(\frac{1}{cx})}{\sqrt{e + \sqrt{dc^2 + e}}}\right)}{16(-d)^{5/2}\sqrt{e}}
 \end{aligned}$$

input

Int[(a + b\*ArcSech[c\*x])/(d + e\*x^2)^3,x]

output

```
(b*c*Sqrt[e]*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)])/(16*(-d)^(3/2)*(c^2*d +
e)*(Sqrt[-d]*Sqrt[e] - d/x)) + (b*c*Sqrt[e]*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1
/(c*x)])/(16*(-d)^(3/2)*(c^2*d + e)*(Sqrt[-d]*Sqrt[e] + d/x)) + (Sqrt[e]*(
a + b*ArcCosh[1/(c*x)]))/(16*(-d)^(3/2)*(Sqrt[-d]*Sqrt[e] - d/x)^2) - (5*(
a + b*ArcCosh[1/(c*x)]))/(16*d^2*(Sqrt[-d]*Sqrt[e] - d/x)) - (Sqrt[e]*(a +
b*ArcCosh[1/(c*x)]))/(16*(-d)^(3/2)*(Sqrt[-d]*Sqrt[e] + d/x)^2) + (5*(a +
b*ArcCosh[1/(c*x)]))/(16*d^2*(Sqrt[-d]*Sqrt[e] + d/x)) + (5*b*ArcTan[(Sqr
t[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[1 + 1/(c*x)])/(Sqrt[c*d + Sqrt[-d]*Sqrt[e]]
*Sqrt[-1 + 1/(c*x)])])/(8*d^2*Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[c*d + Sqrt
[-d]*Sqrt[e]]) - (b*e*ArcTan[(Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[1 + 1/(c*x
)])/(Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*Sqrt[-1 + 1/(c*x)])])/(8*d*(c*d - Sqrt[-
d]*Sqrt[e])^(3/2)*(c*d + Sqrt[-d]*Sqrt[e])^(3/2)) + (5*b*ArcTan[(Sqrt[c*d
+ Sqrt[-d]*Sqrt[e]]*Sqrt[1 + 1/(c*x)])/(Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[
-1 + 1/(c*x)])])/(8*d^2*Sqrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[c*d + Sqrt[-d]*S
qrt[e]]) - (b*e*ArcTan[(Sqrt[c*d + Sqrt[-d]*Sqrt[e]]*Sqrt[1 + 1/(c*x)])/(S
qrt[c*d - Sqrt[-d]*Sqrt[e]]*Sqrt[-1 + 1/(c*x)])])/(8*d*(c*d - Sqrt[-d]*Sqr
t[e])^(3/2)*(c*d + Sqrt[-d]*Sqrt[e])^(3/2)) + (3*(a + b*ArcCosh[1/(c*x)])*
Log[1 - (c*Sqrt[-d]*E^ArcCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])])/(16*
(-d)^(5/2)*Sqrt[e]) - (3*(a + b*ArcCosh[1/(c*x)])*Log[1 + (c*Sqrt[-d]*E^Ar
cCosh[1/(c*x)])/(Sqrt[e] - Sqrt[c^2*d + e])])/(16*(-d)^(5/2)*Sqrt[e]) + ...
```

### Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6374

```
Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^ (n_.)*((f_.)*(x_))^(m_.)*((d_) + (e
_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

rule 6847

```
Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))^ (n_.)*((d_.) + (e_.)*(x_)^2)^(p_.),
x_Symbol] := -Subst[Int[(e + d*x^2)^p*((a + b*ArcCosh[x/c])^n/x^(2*(p + 1)
)), x], x, 1/x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[n, 0] && IntegerQ[p
]
```



**Fricas [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^3} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^3} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="fricas")`

output `integral((b*arcsech(c*x) + a)/(e^3*x^6 + 3*d*e^2*x^4 + 3*d^2*e*x^2 + d^3), x)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^3} dx = \text{Timed out}$$

input `integrate((a+b*asech(c*x))/(e*x**2+d)**3,x)`

output `Timed out`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^3} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^3} dx = \int \frac{b \operatorname{ar} \operatorname{sech}(cx) + a}{(ex^2 + d)^3} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x^2+d)^3,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/(e*x^2 + d)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^3} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{(ex^2 + d)^3} dx$$

input `int((a + b*acosh(1/(c*x)))/(d + e*x^2)^3,x)`

output `int((a + b*acosh(1/(c*x)))/(d + e*x^2)^3, x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^3} dx$$

$$= \frac{3\sqrt{e}\sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e}\sqrt{d}}\right) a d^2 + 6\sqrt{e}\sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e}\sqrt{d}}\right) a d e x^2 + 3\sqrt{e}\sqrt{d} \operatorname{atan}\left(\frac{ex}{\sqrt{e}\sqrt{d}}\right) a e^2 x^4 + 8 \left( \int \frac{a \operatorname{sech}^{-1}(cx)}{e^3 x^6 + 3d e^2 x^4 + 3d^2 e x^2 + d^3} dx \right)}{8d^3}$$

input `int((a+b*asech(c*x))/(e*x^2+d)^3,x)`



output

```
(3*sqrt(e)*sqrt(d)*atan((e*x)/(sqrt(e)*sqrt(d)))*a*d**2 + 6*sqrt(e)*sqrt(d)
)*atan((e*x)/(sqrt(e)*sqrt(d)))*a*d*e*x**2 + 3*sqrt(e)*sqrt(d)*atan((e*x)/
(sqrt(e)*sqrt(d)))*a*e**2*x**4 + 8*int(asech(c*x)/(d**3 + 3*d**2*e*x**2 +
3*d*e**2*x**4 + e**3*x**6),x)*b*d**5*e + 16*int(asech(c*x)/(d**3 + 3*d**2*
e*x**2 + 3*d*e**2*x**4 + e**3*x**6),x)*b*d**4*e**2*x**2 + 8*int(asech(c*x)
/(d**3 + 3*d**2*e*x**2 + 3*d*e**2*x**4 + e**3*x**6),x)*b*d**3*e**3*x**4 +
5*a*d**2*e*x + 3*a*d*e**2*x**3)/(8*d**3*e*(d**2 + 2*d*e*x**2 + e**2*x**4))
```

### 3.129 $\int x^5 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx$

Optimal result	1057
Mathematica [A] (verified)	1058
Rubi [A] (verified)	1059
Maple [F]	1065
Fricas [A] (verification not implemented)	1066
Sympy [F]	1066
Maxima [F(-2)]	1067
Giac [F]	1067
Mupad [F(-1)]	1068
Reduce [F]	1068

#### Optimal result

Integrand size = 23, antiderivative size = 447

$$\begin{aligned}
 & \int x^5 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx \\
 = & \frac{b(23c^4d^2 + 12c^2de - 75e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} \sqrt{d+ex^2}}{1680c^6e^2} \\
 & + \frac{b(29c^2d - 25e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} (d+ex^2)^{3/2}}{840c^4e^2} \\
 & - \frac{b \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} (d+ex^2)^{5/2}}{42c^2e^2} + \frac{d^2(d+ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{3e^3} \\
 & - \frac{2d(d+ex^2)^{5/2} (a + b \operatorname{sech}^{-1}(cx))}{5e^3} + \frac{(d+ex^2)^{7/2} (a + b \operatorname{sech}^{-1}(cx))}{7e^3} \\
 & - \frac{b(105c^6d^3 - 35c^4d^2e + 63c^2de^2 + 75e^3) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{1680c^7e^{5/2}} \\
 & - \frac{8bd^{7/2} \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{105e^3}
 \end{aligned}$$

output

$$\frac{1}{1680} b (23c^4 d^2 + 12c^2 d e - 75e^2) (1/(cx+1))^{1/2} (cx+1)^{1/2} (-c^2 x^2 + 1)^{1/2} (e x^2 + d)^{1/2} / c^6 / e^2 + 1/840 b (29c^2 d - 25e) (1/(cx+1))^{1/2} (cx+1)^{1/2} (-c^2 x^2 + 1)^{1/2} (e x^2 + d)^{3/2} / c^4 / e^2 - 1/42 b (1/(cx+1))^{1/2} (cx+1)^{1/2} (-c^2 x^2 + 1)^{1/2} (e x^2 + d)^{5/2} / c^2 / e^2 + 1/3 d^2 (e x^2 + d)^{3/2} (a + b \operatorname{arcsech}(cx)) / e^3 - 2/5 d (e x^2 + d)^{5/2} (a + b \operatorname{arcsech}(cx)) / e^3 + 1/7 (e x^2 + d)^{7/2} (a + b \operatorname{arcsech}(cx)) / e^3 - 1/1680 b (105c^6 d^3 - 35c^4 d^2 e + 63c^2 d e^2 + 75e^3) (1/(cx+1))^{1/2} (cx+1)^{1/2} \arctan(e^{1/2} (-c^2 x^2 + 1)^{1/2} / c (e x^2 + d)^{1/2}) / c^7 / e^{5/2} - 8/105 b d^{7/2} (1/(cx+1))^{1/2} (cx+1)^{1/2} \operatorname{arctanh}((e x^2 + d)^{1/2} / d^{1/2} / (-c^2 x^2 + 1)^{1/2}) / e^3$$
**Mathematica [A] (verified)**

Time = 35.62 (sec) , antiderivative size = 340, normalized size of antiderivative = 0.76

$$\int x^5 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{\sqrt{d + ex^2} \left( 16ac^6(8d^3 - 4d^2 ex^2 + 3de^2 x^4 + 15e^3 x^6) - be \sqrt{\frac{1-cx}{1+cx}} (1+cx) (75e^2 + 2c^2 e(19d + 25ex^2) + c^4(1680c^6 e^3)) \right)}{1680c^6 e^3} - \frac{b \sqrt{\frac{1-cx}{1+cx}} \sqrt{-1 + c^2 x^2} \left( 128c^7 d^{7/2} \arctan\left(\frac{\sqrt{d} \sqrt{-1 + c^2 x^2}}{\sqrt{d + ex^2}}\right) + \sqrt{e} (105c^6 d^3 - 35c^4 d^2 e + 63c^2 d e^2 + 75e^3) \operatorname{arctan}\left(\frac{\sqrt{d} \sqrt{-1 + c^2 x^2}}{\sqrt{d + ex^2}}\right) \right)}{1680c^7 e^3 (-1 + cx)}$$

input

Integrate[x^5\*Sqrt[d + e\*x^2]\*(a + b\*ArcSech[c\*x]),x]

output

$$\frac{(\operatorname{Sqrt}[d + e x^2] (16 a c^6 (8 d^3 - 4 d^2 e x^2 + 3 d e^2 x^4 + 15 e^3 x^6) - b e \operatorname{Sqrt}[(1 - c x)/(1 + c x)] (1 + c x) (75 e^2 + 2 c^2 e (19 d + 25 e x^2) + c^4 (-41 d^2 + 22 d e x^2 + 40 e^2 x^4)) + 16 b c^6 (8 d^3 - 4 d^2 e x^2 + 3 d e^2 x^4 + 15 e^3 x^6) \operatorname{ArcSech}[c x])) / (1680 c^6 e^3) - (b \operatorname{Sqrt}[(1 - c x)/(1 + c x)] \operatorname{Sqrt}[-1 + c^2 x^2] (128 c^7 d^{7/2} \operatorname{ArcTan}[(\operatorname{Sqrt}[d] \operatorname{Sqrt}[-1 + c^2 x^2]) / \operatorname{Sqrt}[d + e x^2]] + \operatorname{Sqrt}[e] (105 c^6 d^3 - 35 c^4 d^2 e + 63 c^2 d e^2 + 75 e^3) \operatorname{ArcTanh}[(\operatorname{Sqrt}[e] \operatorname{Sqrt}[-1 + c^2 x^2]) / (c \operatorname{Sqrt}[d + e x^2])])) / (1680 c^7 e^3 (-1 + c x))$$

**Rubi [A] (verified)**

Time = 1.54 (sec) , antiderivative size = 380, normalized size of antiderivative = 0.85, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.609$ , Rules used = {6855, 27, 7282, 2118, 27, 171, 27, 171, 27, 175, 66, 104, 218, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^5 \sqrt{d+ex^2} (a+b\operatorname{sech}^{-1}(cx)) dx \\
 & \quad \downarrow \text{6855} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(ex^2+d)^{3/2} (15e^2x^4-12dex^2+8d^2)}{105e^3x\sqrt{1-c^2x^2}} dx + \frac{d^2(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{3e^3} + \frac{(d+ex^2)^{7/2} (a+b\operatorname{sech}^{-1}(cx))}{7e^3} - \frac{2d(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{5e^3}}{210e^3} \\
 & \quad \downarrow \text{27} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(ex^2+d)^{3/2} (15e^2x^4-12dex^2+8d^2)}{x\sqrt{1-c^2x^2}} dx + \frac{d^2(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{3e^3} + \frac{(d+ex^2)^{7/2} (a+b\operatorname{sech}^{-1}(cx))}{7e^3} - \frac{2d(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{5e^3}}{210e^3} \\
 & \quad \downarrow \text{7282} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(ex^2+d)^{3/2} (15e^2x^4-12dex^2+8d^2)}{x^2\sqrt{1-c^2x^2}} dx^2 + \frac{d^2(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{3e^3} + \frac{(d+ex^2)^{7/2} (a+b\operatorname{sech}^{-1}(cx))}{7e^3} - \frac{2d(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{5e^3}}{210e^3} \\
 & \quad \downarrow \text{2118} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{\int -\frac{3e(ex^2+d)^{3/2} (16c^2d^2-(29c^2d-25e)ex^2)}{2x^2\sqrt{1-c^2x^2}} dx^2 - \frac{5e\sqrt{1-c^2x^2}(d+ex^2)^{5/2}}{c^2}}{3c^2e} \right) + \frac{d^2(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{3e^3} + \frac{(d+ex^2)^{7/2} (a+b\operatorname{sech}^{-1}(cx))}{7e^3} - \frac{2d(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{5e^3}}{210e^3}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 27 \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{(ex^2+d)^{3/2}(16c^2d^2-(29c^2d-25e)ex^2)}{x^2\sqrt{1-c^2x^2}} dx}{2c^2} - \frac{5e\sqrt{1-c^2x^2}(d+ex^2)^{5/2}}{c^2} \right) \\
 & \hline
 & \frac{210e^3}{3e^3} \frac{d^2(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} + \frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^3} - \\
 & \frac{2d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3} \\
 & \downarrow 171 \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{e\sqrt{1-c^2x^2}(29c^2d-25e)(d+ex^2)^{3/2}}{2c^2} - \frac{\int \frac{\sqrt{ex^2+d}(64c^4d^3-e(23d^2c^4+12dec^2-75e^2)x^2)}{2x^2\sqrt{1-c^2x^2}} dx}{2c^2} - \frac{5e\sqrt{1-c^2x^2}(d+ex^2)^{5/2}}{c^2} \right) \\
 & \hline
 & \frac{210e^3}{3e^3} \frac{d^2(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} + \frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^3} - \\
 & \frac{2d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3} \\
 & \downarrow 27 \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{\sqrt{ex^2+d}(64c^4d^3-e(23d^2c^4+12dec^2-75e^2)x^2)}{x^2\sqrt{1-c^2x^2}} dx}{4c^2} + \frac{e\sqrt{1-c^2x^2}(29c^2d-25e)(d+ex^2)^{3/2}}{2c^2} - \frac{5e\sqrt{1-c^2x^2}(d+ex^2)^{5/2}}{c^2} \right) \\
 & \hline
 & \frac{210e^3}{3e^3} \frac{d^2(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} + \frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^3} - \\
 & \frac{2d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3} \\
 & \downarrow 171
 \end{aligned}$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\frac{e\sqrt{1-c^2x^2}(23c^4d^2+12c^2de-75e^2)\sqrt{d+ex^2}}{c^2} - \int \frac{128d^4c^6+e(105d^3c^6-35d^2ec^4+63de^2c^2+75e^3)x^2}{2x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2}{4c^2} + \frac{e\sqrt{1-c^2x^2}(29c^2d-25e)}{2c^2} \right)$$

$$\frac{d^2(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} + \frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^3} - \frac{2d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3}$$

↓ 27

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{128d^4c^6+e(105d^3c^6-35d^2ec^4+63de^2c^2+75e^3)x^2}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 + \frac{e\sqrt{1-c^2x^2}(23c^4d^2+12c^2de-75e^2)\sqrt{d+ex^2}}{c^2}}{4c^2} + \frac{e\sqrt{1-c^2x^2}(29c^2d-25e)}{2c^2} \right)$$

$$\frac{d^2(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} + \frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^3} - \frac{2d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3}$$

↓ 175

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\frac{128c^6d^4 \int \frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 + e(105c^6d^3-35c^4d^2e+63c^2de^2+75e^3) \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2}{2c^2} + \frac{e\sqrt{1-c^2x^2}(23c^4d^2+12c^2de-75e^2)\sqrt{d+ex^2}}{c^2}}{4c^2} \right)$$

$$\frac{d^2(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} + \frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^3} - \frac{2d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3}$$

↓ 66

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\frac{128c^6d^4 \int \frac{1}{x^2\sqrt{1-c^2x^2}} dx^2 + 2e(105c^6d^3 - 35c^4d^2e + 63c^2de^2 + 75e^3) \int \frac{1}{-ex^4-c^2} d\frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}}}{2c^2}}{\frac{4c^2}{2c^2}} + \frac{e\sqrt{1-c^2x^2}(23c^4d^2 + 12c^2de - 75e^2)}{c^2} \right)$$

---


$$\frac{d^2(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} + \frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^3} - \frac{2d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3}$$

↓ 104

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\frac{256c^6d^4 \int \frac{1}{x^4-d} d\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} + 2e(105c^6d^3 - 35c^4d^2e + 63c^2de^2 + 75e^3) \int \frac{1}{-ex^4-c^2} d\frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}}}{2c^2}}{\frac{4c^2}{2c^2}} + \frac{e\sqrt{1-c^2x^2}(23c^4d^2 + 12c^2de - 75e^2)}{c^2} \right)$$

---


$$\frac{d^2(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} + \frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^3} - \frac{2d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3}$$

↓ 218

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\frac{256c^6d^4 \int \frac{1}{x^4-d} d\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} - \frac{2\sqrt{e}(105c^6d^3 - 35c^4d^2e + 63c^2de^2 + 75e^3) \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c}}{2c^2}}{\frac{4c^2}{2c^2}} + \frac{e\sqrt{1-c^2x^2}(23c^4d^2 + 12c^2de - 75e^2)}{c^2} \right)$$

---


$$\frac{d^2(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} + \frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^3} - \frac{2d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3}$$

↓ 220

$$\begin{aligned}
 & \frac{d^2(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} + \frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^3} - \\
 & \frac{2d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3} + \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \begin{aligned}
 & -\frac{2\sqrt{e}(105c^6d^3-35c^4d^2e+63c^2de^2+75e^3)\arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c} - \frac{256c^6d^{7/2}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{4c^2} + \frac{e\sqrt{1-c^2x^2}(23c^4d^2+12c^2d^2+12c^2d^2)}{2c^2} \right) \\
 & \hline
 & 210e^3
 \end{aligned} \right)
 \end{aligned}$$

input `Int[x^5*sqrt[d + e*x^2]*(a + b*ArcSech[c*x]),x]`

output `(d^2*(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/(3*e^3) - (2*d*(d + e*x^2)^(5/2)*(a + b*ArcSech[c*x]))/(5*e^3) + ((d + e*x^2)^(7/2)*(a + b*ArcSech[c*x]))/(7*e^3) + (b*sqrt[(1 + c*x)^(-1)]*sqrt[1 + c*x]*((-5*e*sqrt[1 - c^2*x^2])*(d + e*x^2)^(5/2))/c^2 + (((29*c^2*d - 25*e)*e*sqrt[1 - c^2*x^2])*(d + e*x^2)^(3/2))/(2*c^2) + ((e*(23*c^4*d^2 + 12*c^2*d*e - 75*e^2)*sqrt[1 - c^2*x^2]*sqrt[d + e*x^2])/c^2 + ((-2*sqrt[e]*(105*c^6*d^3 - 35*c^4*d^2*e + 63*c^2*d*e^2 + 75*e^3)*ArcTan[(sqrt[e]*sqrt[1 - c^2*x^2])/(c*sqrt[d + e*x^2])])/c - 256*c^6*d^(7/2)*ArcTanh[sqrt[d + e*x^2]/(sqrt[d]*sqrt[1 - c^2*x^2])])/(2*c^2))/(4*c^2)/(2*c^2))/(210*e^3)`

**Defintions of rubi rules used**

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 66 `Int[1/(sqrt[(a_) + (b_.)*(x_)]*sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, sqrt[a + b*x]/sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`



rule 104 `Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 171 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[h*(a + b*x)^m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Simp[1/(d*f*(m + n + p + 2)) Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegersQ[2*m, 2*n, 2*p]`

rule 175 `Int[(((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)))/((a_.) + (b_.)*(x_)), x_] := Simp[h/b Int[(c + d*x)^n*(e + f*x)^p, x], x] + Simp[(b*g - a*h)/b Int[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 2118

```
Int[(Px_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*
(x_))^(p_), x_Symbol] := With[{q = Expon[Px, x], k = Coeff[Px, x, Expo
n[Px, x]]}, Simp[k*(a + b*x)^(m + q - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p +
1)/(d*f*b^(q - 1)*(m + n + p + q + 1))), x] + Simp[1/(d*f*b^q*(m + n + p +
q + 1)) Int[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p*ExpandToSum[d*f*b^q*(m +
n + p + q + 1)*Px - d*f*k*(m + n + p + q + 1)*(a + b*x)^q + k*(a + b*x)^(q
- 2)*(a^2*d*f*(m + n + p + q + 1) - b*(b*c*e*(m + q - 1) + a*(d*e*(n + 1) +
c*f*(p + 1))) + b*(a*d*f*(2*(m + q) + n + p) - b*(d*e*(m + q + n) + c*f*(m
+ q + p)))*x), x], x], x] /; NeQ[m + n + p + q + 1, 0] /; FreeQ[{a, b, c,
d, e, f, m, n, p}, x] && PolyQ[Px, x]
```

rule 6855

```
Int[((a_) + ArcSech[(c_)*(x_)]*(b_))*((f_)*(x_))^(m_)*((d_) + (e_)*(
x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Si
mp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)]
Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x]] /; Fre
eQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] &&
GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2
*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

rule 7282

```
Int[(u_)/(x_), x_Symbol] := With[{lst = PowerVariableExpn[u, 0, x]}, Simp[1
/lst[[2]] Subst[Int[NormalizeIntegrand[Simplify[lst[[1]]/x], x], x], x, (
lst[[3]]*x)^lst[[2]]], x] /; !FalseQ[lst] && NeQ[lst[[2]], 0] /; NonsumQ[
u] && !RationalFunctionQ[u, x]
```

## Maple [F]

$$\int x^5 \sqrt{x^2 e + d} (a + b \operatorname{arcsech}(cx)) dx$$

input

```
int(x^5*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x)
```

output

```
int(x^5*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x)
```

**Fricas [A] (verification not implemented)**

Time = 1.42 (sec) , antiderivative size = 1995, normalized size of antiderivative = 4.46

$$\int x^5 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \text{Too large to display}$$

input `integrate(x^5*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `[1/6720*(128*b*c^7*d^(7/2)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 + 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) - (105*b*c^6*d^3 - 35*b*c^4*d^2*e + 63*b*c^2*d*e^2 + 75*b*e^3)*sqrt(-e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e + 8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^4*e*x^3 + (c^4*d - c^2*e)*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e^2) + 64*(15*b*c^7*e^3*x^6 + 3*b*c^7*d*e^2*x^4 - 4*b*c^7*d^2*e*x^2 + 8*b*c^7*d^3)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 4*(240*a*c^7*e^3*x^6 + 48*a*c^7*d*e^2*x^4 - 64*a*c^7*d^2*e*x^2 + 128*a*c^7*d^3 - (40*b*c^6*e^3*x^5 + 2*(11*b*c^6*d*e^2 + 25*b*c^4*e^3)*x^3 - (41*b*c^6*d^2*e - 38*b*c^4*d*e^2 - 75*b*c^2*e^3)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d)/(c^7*e^3), 1/3360*(64*b*c^7*d^(7/2)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 + 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) - (105*b*c^6*d^3 - 35*b*c^4*d^2*e + 63*b*c^2*d*e^2 + 75*b*e^3)*sqrt(e)*arctan(1/2*(2*c^2*e*x^3 + (c^2*d - e)*x)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*e^2*x^4 + (c^2*d*e - e^2)*x^2 - d*e) + 32*(15*b*c^7*e^3*x^6 + 3*b*c^7*d*e^2*x^4 - 4*b*c^7*d^2*e*x^2 + 8*b*c^7*d^3)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 2*(240*a*c^7*e^3*x^6 + 48*a*c^7*d*e^2*x^4 - 64*a*c^7*d^2*e*x^2 + 128*a*c^7*d^3 - (40*b*c^6*e^3*x^...`

**Sympy [F]**

$$\int x^5 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^5 (a + b \operatorname{asech}(cx)) \sqrt{d + ex^2} dx$$

input `integrate(x**5*(e*x**2+d)**(1/2)*(a+b*asech(c*x)),x)`

output `Integral(x**5*(a + b*asech(c*x))*sqrt(d + e*x**2), x)`

### Maxima [F(-2)]

Exception generated.

$$\int x^5 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \text{Exception raised: ValueError}$$

input `integrate(x^5*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

### Giac [F]

$$\int x^5 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int \sqrt{ex^2 + d} (b \operatorname{ar} \operatorname{sech}(cx) + a) x^5 dx$$

input `integrate(x^5*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)*x^5, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^5 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^5 \sqrt{ex^2 + d} \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int(x^5*(d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))),x)`output `int(x^5*(d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))), x)`**Reduce [F]**

$$\int x^5 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^5 \sqrt{ex^2 + d} (a \operatorname{sech}(cx) b + a) dx$$

input `int(x^5*(e*x^2+d)^(1/2)*(a+b*asech(c*x)),x)`output `int(x^5*(e*x^2+d)^(1/2)*(a+b*asech(c*x)),x)`

### 3.130 $\int x^3 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx$

Optimal result	1069
Mathematica [A] (verified)	1070
Rubi [A] (verified)	1071
Maple [F]	1075
Fricas [A] (verification not implemented)	1075
Sympy [F]	1076
Maxima [F(-2)]	1077
Giac [F]	1077
Mupad [F(-1)]	1077
Reduce [F]	1078

#### Optimal result

Integrand size = 23, antiderivative size = 329

$$\begin{aligned}
 & \int x^3 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx \\
 &= -\frac{b(c^2d + 9e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} \sqrt{d+ex^2}}{120c^4e} \\
 &\quad - \frac{b \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} (d+ex^2)^{3/2}}{20c^2e} \\
 &\quad - \frac{d(d+ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{3e^2} + \frac{(d+ex^2)^{5/2} (a + b \operatorname{sech}^{-1}(cx))}{5e^2} \\
 &\quad + \frac{b(15c^4d^2 - 10c^2de - 9e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{120c^5e^{3/2}} \\
 &\quad + \frac{2bd^{5/2} \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{15e^2}
 \end{aligned}$$

output

```
-1/120*b*(c^2*d+9*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*(e
*x^2+d)^(1/2)/c^4/e-1/20*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1
/2)*(e*x^2+d)^(3/2)/c^2/e-1/3*d*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/e^2+1/5
*(e*x^2+d)^(5/2)*(a+b*arcsech(c*x))/e^2+1/120*b*(15*c^4*d^2-10*c^2*d*e-9*e
^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctan(e^(1/2)*(-c^2*x^2+1)^(1/2)/c/(e
*x^2+d)^(1/2))/c^5/e^(3/2)+2/15*b*d^(5/2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*
arctanh((e*x^2+d)^(1/2)/d^(1/2)/(-c^2*x^2+1)^(1/2))/e^2
```

**Mathematica [A] (verified)**

Time = 22.14 (sec) , antiderivative size = 365, normalized size of antiderivative = 1.11

$$\int x^3 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx =$$

$$\frac{\sqrt{d + ex^2} \left( 8ac^4(2d^2 - dex^2 - 3e^2x^4) + be\sqrt{\frac{1-cx}{1+cx}}(1+cx)(9e + c^2(7d + 6ex^2)) + 8bc^4(2d^2 - dex^2 - 3e^2x^4) \right)}{120c^4e^2} +$$

$$\frac{b\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2} \left( \sqrt{-c^2}\sqrt{-c^2d-e}e\sqrt{e}(15c^4d^2 - 10c^2de - 9e^2) \sqrt{\frac{c^2(d+ex^2)}{c^2d+e}} \arcsin\left(\frac{c\sqrt{e}\sqrt{1-c^2x^2}}{\sqrt{-c^2}\sqrt{-c^2d-e}}\right) + 1 \right)}{120c^7e^2(-1+cx)\sqrt{d+ex^2}}$$

input

```
Integrate[x^3*Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]),x]
```

output

```
-1/120*(Sqrt[d + e*x^2]*(8*a*c^4*(2*d^2 - d*e*x^2 - 3*e^2*x^4) + b*e*Sqrt[
(1 - c*x)/(1 + c*x)]*(1 + c*x)*(9*e + c^2*(7*d + 6*e*x^2)) + 8*b*c^4*(2*d^
2 - d*e*x^2 - 3*e^2*x^4)*ArcSech[c*x]))/(c^4*e^2) - (b*Sqrt[(1 - c*x)/(1 +
c*x)]*Sqrt[1 - c^2*x^2]*(Sqrt[-c^2]*Sqrt[-(c^2*d) - e]*Sqrt[e]*(15*c^4*d^
2 - 10*c^2*d*e - 9*e^2)*Sqrt[(c^2*(d + e*x^2))/(c^2*d + e)]*ArcSin[(c*Sqrt
[e]*Sqrt[1 - c^2*x^2])/(Sqrt[-c^2]*Sqrt[-(c^2*d) - e]]) + 16*c^7*d^(5/2)*S
qrt[-d - e*x^2]*ArcTan[(Sqrt[d]*Sqrt[1 - c^2*x^2])/Sqrt[-d - e*x^2]]))/(12
0*c^7*e^2*(-1 + c*x)*Sqrt[d + e*x^2])
```

**Rubi [A] (verified)**

Time = 0.58 (sec) , antiderivative size = 279, normalized size of antiderivative = 0.85, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.522$ , Rules used = {6855, 27, 435, 171, 27, 171, 27, 175, 66, 104, 218, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \sqrt{d+ex^2} (a+b\operatorname{sech}^{-1}(cx)) \, dx \\
 & \quad \downarrow \text{6855} \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int -\frac{(2d-3ex^2)(ex^2+d)^{3/2}}{15e^2x\sqrt{1-c^2x^2}} dx + \frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2} - \\
 & \quad \frac{d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2} \\
 & \quad \downarrow \text{27} \\
 & -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(2d-3ex^2)(ex^2+d)^{3/2}}{x\sqrt{1-c^2x^2}} dx}{15e^2} + \frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2} - \\
 & \quad \frac{d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2} \\
 & \quad \downarrow \text{435} \\
 & -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(2d-3ex^2)(ex^2+d)^{3/2}}{x^2\sqrt{1-c^2x^2}} dx^2}{30e^2} + \frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2} - \\
 & \quad \frac{d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2} \\
 & \quad \downarrow \text{171} \\
 & -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2} - \frac{\int -\frac{\sqrt{ex^2+d}(8c^2d^2-e(dc^2+9e)x^2)}{2x^2\sqrt{1-c^2x^2}} dx^2}{2c^2} \right)}{30e^2} + \\
 & \quad \frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2} - \frac{d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$



$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{\int\frac{\sqrt{ex^2+d}(8c^2d^2-e(dc^2+9e)x^2)}{x^2\sqrt{1-c^2x^2}}dx^2}{4c^2}+\frac{3e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2}\right)$$


---


$$\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2}-\frac{d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2}$$

↓ 171

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{e\sqrt{1-c^2x^2}(c^2d+9e)\sqrt{d+ex^2}}{c^2}-\frac{\int-\frac{16d^3c^4+e(15d^2c^4-10dec^2-9e^2)x^2}{2x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2}{4c^2}+\frac{3e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2}\right)$$


---


$$\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2}-\frac{d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2}$$

↓ 27

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{\int\frac{16d^3c^4+e(15d^2c^4-10dec^2-9e^2)x^2}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2}{4c^2}+\frac{e\sqrt{1-c^2x^2}(c^2d+9e)\sqrt{d+ex^2}}{c^2}+\frac{3e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2}\right)$$


---


$$\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2}-\frac{d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2}$$

↓ 175

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{16c^4d^3\int\frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2+e(15c^4d^2-10c^2de-9e^2)\int\frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2}{2c^2}+\frac{e\sqrt{1-c^2x^2}(c^2d+9e)\sqrt{d+ex^2}}{c^2}+\frac{3e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2}\right)$$


---


$$\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2}-\frac{d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2}$$

↓ 66

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{16c^4d^3\int\frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2+2e(15c^4d^2-10c^2de-9e^2)\int\frac{1}{-ex^4-c^2}\frac{d\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}}}{2c^2}+\frac{e\sqrt{1-c^2x^2}(c^2d+9e)\sqrt{d+ex^2}}{c^2}+\frac{3e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2}\right)$$


---


$$\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2}-\frac{d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2}$$

↓ 104

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{32c^4d^3 \int \frac{1}{x^4-d} d \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} + 2e(15c^4d^2-10c^2de-9e^2) \int \frac{1}{-ex^4-c^2} d \frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}} + \frac{e\sqrt{1-c^2x^2}(c^2d+9e)\sqrt{d+ex^2}}{c^2}}{4c^2} + \frac{3e\sqrt{1-c^2x^2}}{2c} \right)$$


---


$$\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2} - \frac{d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2}$$

↓ 218

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{32c^4d^3 \int \frac{1}{x^4-d} d \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} - \frac{2\sqrt{e}(15c^4d^2-10c^2de-9e^2) \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{2c^2} + \frac{e\sqrt{1-c^2x^2}(c^2d+9e)\sqrt{d+ex^2}}{c^2}}{4c^2} + \frac{3e\sqrt{1-c^2x^2}}{2c} \right)$$


---


$$\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2} - \frac{d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2}$$

↓ 220

$$\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2} - \frac{d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2} -$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\frac{2\sqrt{e}(15c^4d^2-10c^2de-9e^2) \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c} - 32c^4d^{5/2}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{2c^2} + \frac{e\sqrt{1-c^2x^2}(c^2d+9e)\sqrt{d+ex^2}}{c^2}}{4c^2} + \frac{3e\sqrt{1-c^2x^2}}{2c} \right)$$


---


$$30e^2$$

input `Int[x^3*Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]),x]`

output `-1/3*(d*(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/e^2 + ((d + e*x^2)^(5/2)*(a + b*ArcSech[c*x]))/(5*e^2) - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((3*e*Sqrt[1 - c^2*x^2]*(d + e*x^2)^(3/2))/(2*c^2) + ((e*(c^2*d + 9*e)*Sqrt[1 - c^2*x^2]*Sqrt[d + e*x^2])/c^2 + ((-2*Sqrt[e]*(15*c^4*d^2 - 10*c^2*d*e - 9*e^2)*ArcTan[(Sqrt[e]*Sqrt[1 - c^2*x^2])/(c*Sqrt[d + e*x^2])])/c - 32*c^4*d^(5/2)*ArcTanh[Sqrt[d + e*x^2]/(Sqrt[d]*Sqrt[1 - c^2*x^2])])/(2*c^2))/(4*c^2)))/(30*e^2)`

## Defintions of rubi rules used

- rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 66  $\text{Int}[1/(\text{Sqrt}[(a_*) + (b_*)(x_)]*\text{Sqrt}[(c_*) + (d_*)(x_)]), x\_Symbol] \rightarrow \text{Simp}[2 \text{ Subst}[\text{Int}[1/(b - d*x^2), x], x, \text{Sqrt}[a + b*x]/\text{Sqrt}[c + d*x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ !\text{GtQ}[c - a*(d/b), 0]$
- rule 104  $\text{Int}[(((a_*) + (b_*)(x_))^{(m_*)}*((c_*) + (d_*)(x_))^{(n_*)})/((e_*) + (f_*)(x_))), x_] \rightarrow \text{With}[\{q = \text{Denominator}[m]\}, \text{Simp}[q \text{ Subst}[\text{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{EqQ}[m + n + 1, 0] \ \&\& \ \text{RationalQ}[n] \ \&\& \ \text{LTQ}[-1, m, 0] \ \&\& \ \text{SimplerQ}[a + b*x, c + d*x]$
- rule 171  $\text{Int}[(((a_*) + (b_*)(x_))^{(m_*)}*((c_*) + (d_*)(x_))^{(n_*)}*((e_*) + (f_*)(x_))^{(p_*)}*((g_*) + (h_*)(x_))), x_] \rightarrow \text{Simp}[h*(a + b*x)^m*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)}/(d*f*(m+n+p+2))), x] + \text{Simp}[1/(d*f*(m+n+p+2)) \text{ Int}[(a + b*x)^{(m-1)}*(c + d*x)^n*(e + f*x)^p*\text{Simp}[a*d*f*g*(m+n+p+2) - h*(b*c*e*m + a*(d*e*(n+1) + c*f*(p+1))] + (b*d*f*g*(m+n+p+2) + h*(a*d*f*m - b*(d*e*(m+n+1) + c*f*(m+p+1)))*x, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, h, n, p\}, x] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{NeQ}[m+n+p+2, 0] \ \&\& \ \text{IntegersQ}[2*m, 2*n, 2*p]$
- rule 175  $\text{Int}[(((c_*) + (d_*)(x_))^{(n_*)}*((e_*) + (f_*)(x_))^{(p_*)}*((g_*) + (h_*)(x_))))/((a_*) + (b_*)(x_)), x_] \rightarrow \text{Simp}[h/b \text{ Int}[(c + d*x)^n*(e + f*x)^p, x], x] + \text{Simp}[(b*g - a*h)/b \text{ Int}[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, h, n, p\}, x]$
- rule 218  $\text{Int}[((a_*) + (b_*)(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

rule 220 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 435 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2)^(r_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q*(e + f*x)^r, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, f, p, q, r}, x] && IntegerQ[(m - 1)/2]`

rule 6855 `Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*((f_)*(x_)^(m_)*((d_) + (e_)*(x_)^2)^(p_)), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))`

## Maple [F]

$$\int x^3 \sqrt{x^2 e + d} (a + b \operatorname{arcsech}(cx)) dx$$

input `int(x^3*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x)`

output `int(x^3*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x)`

## Fricas [A] (verification not implemented)

Time = 0.66 (sec) , antiderivative size = 1669, normalized size of antiderivative = 5.07

$$\int x^3 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \text{Too large to display}$$

input `integrate(x^3*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output

```
[1/480*(16*b*c^5*d^(5/2)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 - 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) + (15*b*c^4*d^2 - 10*b*c^2*d*e - 9*b*e^2)*sqrt(-e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e + 8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^4*e*x^3 + (c^4*d - c^2*e)*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e^2) + 32*(3*b*c^5*e^2*x^4 + b*c^5*d*e*x^2 - 2*b*c^5*d^2)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 4*(24*a*c^5*e^2*x^4 + 8*a*c^5*d*e*x^2 - 16*a*c^5*d^2 - (6*b*c^4*e^2*x^3 + (7*b*c^4*d*e + 9*b*c^2*e^2)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d))/(c^5*e^2), 1/240*(8*b*c^5*d^(5/2)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 - 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) + (15*b*c^4*d^2 - 10*b*c^2*d*e - 9*b*e^2)*sqrt(e)*arctan(1/2*(2*c^2*e*x^3 + (c^2*d - e)*x)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*e^2*x^4 + (c^2*d*e - e^2)*x^2 - d*e) + 16*(3*b*c^5*e^2*x^4 + b*c^5*d*e*x^2 - 2*b*c^5*d^2)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 2*(24*a*c^5*e^2*x^4 + 8*a*c^5*d*e*x^2 - 16*a*c^5*d^2 - (6*b*c^4*e^2*x^3 + (7*b*c^4*d*e + 9*b*c^2*e^2)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d))/(c^5*e^2), 1/480*(32*b*c^5*sqrt(-d)*d^2*arctan(-1/2*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(-d)*sqrt(-(c^2*x^2 - 1)/...
```

### Sympy [F]

$$\int x^3 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^3 (a + b \operatorname{asech}(cx)) \sqrt{d + ex^2} dx$$

input

```
integrate(x**3*(e*x**2+d)**(1/2)*(a+b*asech(c*x)),x)
```

output

```
Integral(x**3*(a + b*asech(c*x))*sqrt(d + e*x**2), x)
```

**Maxima [F(-2)]**

Exception generated.

$$\int x^3 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \text{Exception raised: ValueError}$$

input `integrate(x^3*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int x^3 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int \sqrt{ex^2 + d} (b \operatorname{ar} \operatorname{sech}(cx) + a) x^3 dx$$

input `integrate(x^3*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)*x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^3 \sqrt{ex^2 + d} \left( a + b \operatorname{acosh} \left( \frac{1}{cx} \right) \right) dx$$

input `int(x^3*(d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))),x)`

output `int(x^3*(d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))), x)`

**Reduce [F]**

$$\int x^3 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^3 \sqrt{ex^2 + d} (a \operatorname{sech}(cx) b + a) dx$$

input `int(x^3*(e*x^2+d)^(1/2)*(a+b*asech(c*x)),x)`

output `int(x^3*(e*x^2+d)^(1/2)*(a+b*asech(c*x)),x)`

### 3.131 $\int x\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	1079
Mathematica [A] (verified)	1080
Rubi [A] (verified)	1080
Maple [F]	1084
Fricas [B] (verification not implemented)	1084
Sympy [F]	1085
Maxima [F]	1086
Giac [F]	1086
Mupad [F(-1)]	1086
Reduce [F]	1087

#### Optimal result

Integrand size = 21, antiderivative size = 221

$$\int x\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx = -\frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{6c^2} + \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e} - \frac{b(3c^2d+e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{6c^3\sqrt{e}} - \frac{bd^{3/2}\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{3e}$$

output

```
-1/6*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*(e*x^2+d)^(1/2)/
c^2+1/3*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/e-1/6*b*(3*c^2*d+e)*(1/(c*x+1))
^(1/2)*(c*x+1)^(1/2)*arctan(e^(1/2)*(-c^2*x^2+1)^(1/2)/c/(e*x^2+d)^(1/2))/
c^3/e^(1/2)-1/3*b*d^(3/2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh((e*x^2+d
)^(1/2)/d^(1/2)/(-c^2*x^2+1)^(1/2))/e
```



**Mathematica [A] (verified)**

Time = 21.75 (sec) , antiderivative size = 307, normalized size of antiderivative = 1.39

$$\int x\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))dx$$

$$= \frac{\sqrt{d+ex^2}\left(-be\sqrt{\frac{1-cx}{1+cx}}(1+cx)+2ac^2(d+ex^2)+2bc^2(d+ex^2)\operatorname{sech}^{-1}(cx)\right)}{6c^2e}$$

$$+ \frac{b\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\left(\sqrt{-c^2}\sqrt{-c^2d-e}\sqrt{e}(3c^2d+e)\sqrt{\frac{c^2(d+ex^2)}{c^2d+e}}\arcsin\left(\frac{c\sqrt{e}\sqrt{1-c^2x^2}}{\sqrt{-c^2}\sqrt{-c^2d-e}}\right)+2c^5d^{3/2}\sqrt{-d-e}\right)}{6c^5e(-1+cx)\sqrt{d+ex^2}}$$

input

```
Integrate[x*Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]),x]
```

output

```
(Sqrt[d + e*x^2]*(-(b*e*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)) + 2*a*c^2*(d + e*x^2) + 2*b*c^2*(d + e*x^2)*ArcSech[c*x]))/(6*c^2*e) + (b*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*(Sqrt[-c^2]*Sqrt[-(c^2*d) - e]*Sqrt[e]*(3*c^2*d + e)*Sqrt[(c^2*(d + e*x^2))/(c^2*d + e)]*ArcSin[(c*Sqrt[e]*Sqrt[1 - c^2*x^2])/(Sqrt[-c^2]*Sqrt[-(c^2*d) - e]]) + 2*c^5*d^(3/2)*Sqrt[-d - e*x^2]*ArcTan[(Sqrt[d]*Sqrt[1 - c^2*x^2])/Sqrt[-d - e*x^2]]))/(6*c^5*e*(-1 + c*x))*Sqrt[d + e*x^2]
```

**Rubi [A] (verified)**

Time = 0.58 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.86, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.476$ , Rules used = {6853, 2036, 354, 113, 27, 175, 66, 104, 218, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))dx$$

$$\downarrow 6853$$

$$\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\int\frac{(ex^2+d)^{3/2}}{x\sqrt{1-cx}\sqrt{cx+1}}dx}{3e} + \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e}$$

$$\begin{aligned}
& \downarrow 2036 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(ex^2+d)^{3/2}}{x\sqrt{1-c^2x^2}} dx}{3e} + \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{3e} \\
& \downarrow 354 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(ex^2+d)^{3/2}}{x^2\sqrt{1-c^2x^2}} dx^2}{6e} + \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{3e} \\
& \downarrow 113 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{\int -\frac{2c^2d^2+e(3dc^2+e)x^2}{2x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2}{c^2} - \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} \right)}{6e} + \\
& \quad \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{3e} \\
& \downarrow 27 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{2c^2d^2+e(3dc^2+e)x^2}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2}{2c^2} - \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} \right)}{6e} + \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{3e} \\
& \downarrow 175 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{2c^2d^2 \int \frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 + e(3c^2d+e) \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2}{2c^2} - \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} \right)}{6e} + \\
& \quad \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{3e} \\
& \downarrow 66 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{2c^2d^2 \int \frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 + 2e(3c^2d+e) \int \frac{1}{-ex^4-c^2} d \frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}}}{2c^2} - \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} \right)}{6e} + \\
& \quad \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{3e} \\
& \downarrow 104
\end{aligned}$$

$$\begin{aligned}
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{4c^2d^2 \int \frac{1}{x^4-d} d \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} + 2e(3c^2d+e) \int \frac{1}{-ex^4-c^2} d \frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}} - \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} \right)}{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))} + \\
 & \qquad \qquad \qquad \frac{6e}{3e} \\
 & \qquad \qquad \qquad \downarrow \text{218} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{4c^2d^2 \int \frac{1}{x^4-d} d \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} - \frac{2\sqrt{e}(3c^2d+e) \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c} - \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} \right)}{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))} + \\
 & \qquad \qquad \qquad \frac{6e}{3e} \\
 & \qquad \qquad \qquad \downarrow \text{220} \\
 & \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{3e} + \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{2\sqrt{e}(3c^2d+e) \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c} - \frac{4c^2d^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{2c^2} - \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} \right)}{6e}
 \end{aligned}$$

input `Int[x*Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]),x]`

output `((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/(3*e) + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-(e*Sqrt[1 - c^2*x^2]*Sqrt[d + e*x^2])/c^2) + ((-2*Sqrt[e]*(3*c^2*d + e)*ArcTan[(Sqrt[e]*Sqrt[1 - c^2*x^2])/(c*Sqrt[d + e*x^2])])/c - 4*c^2*d^(3/2)*ArcTanh[Sqrt[d + e*x^2]/(Sqrt[d]*Sqrt[1 - c^2*x^2])])/(2*c^2)))/(6*e)`

## Definitions of rubi rules used

- rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 66  $\text{Int}[1/(\text{Sqrt}[(a_*) + (b_*)(x_)]*\text{Sqrt}[(c_*) + (d_*)(x_)]), x\_Symbol] \rightarrow \text{Simp}[2 \text{ Subst}[\text{Int}[1/(b - d*x^2), x], x, \text{Sqrt}[a + b*x]/\text{Sqrt}[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ !\text{GtQ}[c - a*(d/b), 0]$
- rule 104  $\text{Int}[(((a_*) + (b_*)(x_))^{(m_*)}*((c_*) + (d_*)(x_))^{(n_*)})/((e_*) + (f_*)(x_))), x_] \rightarrow \text{With}[\{q = \text{Denominator}[m]\}, \text{Simp}[q \text{ Subst}[\text{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{EqQ}[m + n + 1, 0] \ \&\& \ \text{RationalQ}[n] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{SimplerQ}[a + b*x, c + d*x]$
- rule 113  $\text{Int}[(((a_*) + (b_*)(x_))^{(m_*)}*((c_*) + (d_*)(x_))^{(n_*)}*((e_*) + (f_*)(x_))^{(p_*)}), x_] \rightarrow \text{Simp}[b*(a + b*x)^{(m-1)}*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)}/(d*f*(m+n+p+1))), x] + \text{Simp}[1/(d*f*(m+n+p+1)) \text{ Int}[(a + b*x)^{(m-2)}*(c + d*x)^n*(e + f*x)^p*\text{Simp}[a^2*d*f*(m+n+p+1) - b*(b*c*e*(m-1) + a*(d*e*(n+1) + c*f*(p+1))) + b*(a*d*f*(2*m+n+p) - b*(d*e*(m+n) + c*f*(m+p)))*x, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{NeQ}[m+n+p+1, 0] \ \&\& \ \text{IntegersQ}[2*m, 2*n, 2*p]$
- rule 175  $\text{Int}[(((c_*) + (d_*)(x_))^{(n_*)}*((e_*) + (f_*)(x_))^{(p_*)}*((g_*) + (h_*)(x_))))/((a_*) + (b_*)(x_)), x_] \rightarrow \text{Simp}[h/b \text{ Int}[(c + d*x)^n*(e + f*x)^p, x], x] + \text{Simp}[(b*g - a*h)/b \text{ Int}[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, h, n, p\}, x]$
- rule 218  $\text{Int}[((a_*) + (b_*)(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$
- rule 220  $\text{Int}[((a_*) + (b_*)(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[b, 2])^{-1})*\text{ArcTanh}[\text{Rt}[b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 354

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol]
:= Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x]
/; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]
```

rule 2036

```
Int[(u_)*((c_) + (d_)*(x_)^(n_))^(q_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_), x_Symbol]
:= Int[u*(a1*a2 + b1*b2*x^n)^p*(c + d*x^n)^q, x]
/; FreeQ[{a1, b1, a2, b2, c, d, n, p, q}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))
```

rule 6853

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*(x_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol]
:= Simp[(d + e*x^2)^(p + 1)*((a + b*ArcSech[c*x])/(2*e*(p + 1))), x]
+ Simp[b*(Sqrt[1 + c*x]/(2*e*(p + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x^2)^(p + 1)/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x]
/; FreeQ[{a, b, c, d, e, p}, x] && NeQ[p, -1]
```

**Maple [F]**

$$\int x\sqrt{x^2e+d}(a+b \operatorname{arcsech}(cx)) dx$$

input

```
int(x*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x)
```

output

```
int(x*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 328 vs.  $2(131) = 262$ .

Time = 0.33 (sec) , antiderivative size = 1382, normalized size of antiderivative = 6.25

$$\int x\sqrt{d+ex^2}(a+b \operatorname{sech}^{-1}(cx)) dx = \text{Too large to display}$$

input `integrate(x*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `[1/24*(2*b*c^3*d^(3/2)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 + 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) - (3*b*c^2*d + b*e)*sqrt(-e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e + 8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^4*e*x^3 + (c^4*d - c^2*e)*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e^2) + 8*(b*c^3*e*x^2 + b*c^3*d)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 4*(2*a*c^3*e*x^2 - b*c^2*e*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 2*a*c^3*d)*sqrt(e*x^2 + d))/(c^3*e), 1/12*(b*c^3*d^(3/2)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 + 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) - (3*b*c^2*d + b*e)*sqrt(e)*arctan(1/2*(2*c^2*e*x^3 + (c^2*d - e)*x)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*e^2*x^4 + (c^2*d*e - e^2)*x^2 - d*e) + 4*(b*c^3*e*x^2 + b*c^3*d)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 2*(2*a*c^3*e*x^2 - b*c^2*e*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 2*a*c^3*d)*sqrt(e*x^2 + d))/(c^3*e), -1/24*(4*b*c^3*sqrt(-d)*d*arctan(-1/2*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(-d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*d*e*x^4 + (c^2*d^2 - d*e)*x^2 - d^2)) + (3*b*c^2*d + b*e)*sqrt(-e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e + 8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^4*e*x^3 + (c^4*d - c^2*e)*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(-(c^2*x^2 - 1)/...`

## Sympy [F]

$$\int x\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx = \int x(a+b\operatorname{asech}(cx))\sqrt{d+ex^2} dx$$

input `integrate(x*(e*x**2+d)**(1/2)*(a+b*asech(c*x)),x)`

output `Integral(x*(a + b*asech(c*x))*sqrt(d + e*x**2), x)`

**Maxima [F]**

$$\int x\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx = \int \sqrt{ex^2+d}(b\operatorname{ar}\operatorname{sech}(cx)+a)x dx$$

input `integrate(x*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `1/3*((e*x^2 + d)^(3/2)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)/e - 3*integrate(1/3*sqrt(e*x^2 + d)*(6*(c^2*e*x^2 - e)*x*log(sqrt(x)) + 3*(c^2*e*x^2*log(c) - e*log(c))*x + (6*(c^2*e*x^2 - e)*x*log(sqrt(x)) + ((3*e*log(c) + e)*c^2*x^2 + c^2*d - 3*e*log(c))*x)*e^(1/2*log(c*x + 1) + 1/2*log(-c*x + 1)))/(c^2*e*x^2 + (c^2*e*x^2 - e)*e^(1/2*log(c*x + 1) + 1/2*log(-c*x + 1)) - e), x)*b + 1/3*(e*x^2 + d)^(3/2)*a/e`

**Giac [F]**

$$\int x\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx = \int \sqrt{ex^2+d}(b\operatorname{ar}\operatorname{sech}(cx)+a)x dx$$

input `integrate(x*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)*x, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx = \int x\sqrt{ex^2+d}\left(a+b\operatorname{acosh}\left(\frac{1}{cx}\right)\right) dx$$

input `int(x*(d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))),x)`

output `int(x*(d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))), x)`

**Reduce [F]**

$$\int x\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx)) dx = \int x\sqrt{ex^2+d}(a\operatorname{sech}(cx)b+a) dx$$

input `int(x*(e*x^2+d)^(1/2)*(a+b*asech(c*x)),x)`

output `int(x*(e*x^2+d)^(1/2)*(a+b*asech(c*x)),x)`



$$3.132 \quad \int \frac{\sqrt{d+ex^2} \left( a + b \operatorname{sech}^{-1}(cx) \right)}{x} dx$$

Optimal result	1088
Mathematica [N/A]	1088
Rubi [N/A]	1089
Maple [N/A]	1089
Fricas [N/A]	1090
Sympy [N/A]	1090
Maxima [F(-2)]	1090
Giac [N/A]	1091
Mupad [N/A]	1091
Reduce [N/A]	1092

### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{\sqrt{d+ex^2} \left( a + b \operatorname{sech}^{-1}(cx) \right)}{x} dx = \operatorname{Int} \left( \frac{\sqrt{d+ex^2} \left( a + b \operatorname{sech}^{-1}(cx) \right)}{x}, x \right)$$

output `Defer(Int)((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x,x)`

### Mathematica [N/A]

Not integrable

Time = 5.65 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{\sqrt{d+ex^2} \left( a + b \operatorname{sech}^{-1}(cx) \right)}{x} dx = \int \frac{\sqrt{d+ex^2} \left( a + b \operatorname{sech}^{-1}(cx) \right)}{x} dx$$

input `Integrate[(Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/x,x]`

output `Integrate[(Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/x, x]`

**Rubi [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{d + ex^2}(a + b\operatorname{sech}^{-1}(cx))}{x} dx$$

↓ 6865

$$\int \frac{\sqrt{d + ex^2}(a + b\operatorname{sech}^{-1}(cx))}{x} dx$$

input `Int[(Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/x,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{\sqrt{x^2e + d}(a + b \operatorname{arcsech}(cx))}{x} dx$$

input `int((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x,x)`

output `int((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x,x)`

**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{\sqrt{ex^2+d}(b\operatorname{arsech}(cx)+a)}{x} dx$$

input `integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x,x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)/x, x)`

**Sympy [N/A]**

Not integrable

Time = 4.77 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(a+b\operatorname{asech}(cx))\sqrt{d+ex^2}}{x} dx$$

input `integrate((e*x**2+d)**(1/2)*(a+b*asech(c*x))/x,x)`

output `Integral((a + b*asech(c*x))*sqrt(d + e*x**2)/x, x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x} dx = \text{Exception raised: ValueError}$$

input `integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x,x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{\sqrt{ex^2+d}(b\operatorname{ar} \operatorname{sech}(cx)+a)}{x} dx$$

input

```
integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x,x, algorithm="giac")
```

output

```
integrate(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)/x, x)
```

**Mupad [N/A]**

Not integrable

Time = 4.14 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{\sqrt{ex^2+d}(a+b\operatorname{acosh}(\frac{1}{cx}))}{x} dx$$

input

```
int(((d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))))/x,x)
```

output

```
int(((d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))))/x, x)
```

**Reduce [N/A]**

Not integrable

Time = 200.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d + ex^2}(a + b\operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{\sqrt{ex^2 + d}(a\operatorname{sech}(cx) b + a)}{x} dx$$

input `int((e*x^2+d)^(1/2)*(a+b*asech(c*x))/x,x)`output `int((e*x^2+d)^(1/2)*(a+b*asech(c*x))/x,x)`

**3.133** 
$$\int \frac{\sqrt{d+ex^2} \left( a+b\operatorname{sech}^{-1}(cx) \right)}{x^3} dx$$

Optimal result	1093
Mathematica [N/A]	1093
Rubi [N/A]	1094
Maple [N/A]	1094
Fricas [N/A]	1095
Sympy [N/A]	1095
Maxima [F(-2)]	1095
Giac [N/A]	1096
Mupad [N/A]	1096
Reduce [N/A]	1097

**Optimal result**

Integrand size = 23, antiderivative size = 23

$$\int \frac{\sqrt{d+ex^2} \left( a+b\operatorname{sech}^{-1}(cx) \right)}{x^3} dx = \operatorname{Int} \left( \frac{\sqrt{d+ex^2} \left( a+b\operatorname{sech}^{-1}(cx) \right)}{x^3}, x \right)$$

output `Defer(Int)((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^3,x)`

**Mathematica [N/A]**

Not integrable

Time = 5.38 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{\sqrt{d+ex^2} \left( a+b\operatorname{sech}^{-1}(cx) \right)}{x^3} dx = \int \frac{\sqrt{d+ex^2} \left( a+b\operatorname{sech}^{-1}(cx) \right)}{x^3} dx$$

input `Integrate[(Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/x^3,x]`

output `Integrate[(Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/x^3, x]`

**Rubi [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{d + ex^2}(a + b\operatorname{sech}^{-1}(cx))}{x^3} dx$$

↓ 6865

$$\int \frac{\sqrt{d + ex^2}(a + b\operatorname{sech}^{-1}(cx))}{x^3} dx$$

input `Int[(Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/x^3,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.38 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{\sqrt{x^2e + d}(a + b \operatorname{arcsech}(cx))}{x^3} dx$$

input `int((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^3,x)`

output `int((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^3,x)`

**Fricas [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{\sqrt{ex^2+d}(b\operatorname{ar} \operatorname{sech}(cx)+a)}{x^3} dx$$

input `integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^3,x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)/x^3, x)`

**Sympy [N/A]**

Not integrable

Time = 4.99 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(a+b\operatorname{asech}(cx))\sqrt{d+ex^2}}{x^3} dx$$

input `integrate((e*x**2+d)**(1/2)*(a+b*asech(c*x))/x**3,x)`

output `Integral((a + b*asech(c*x))*sqrt(d + e*x**2)/x**3, x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^3} dx = \text{Exception raised: ValueError}$$

input `integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^3,x, algorithm="maxima")`



output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{\sqrt{ex^2+d}(b\operatorname{arsech}(cx)+a)}{x^3} dx$$

input

```
integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^3,x, algorithm="giac")
```

output

```
integrate(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)/x^3, x)
```

**Mupad [N/A]**

Not integrable

Time = 4.35 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{\sqrt{ex^2+d}(a+b\operatorname{acosh}(\frac{1}{cx}))}{x^3} dx$$

input

```
int(((d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))))/x^3,x)
```

output

```
int(((d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))))/x^3, x)
```

**Reduce [N/A]**

Not integrable

Time = 200.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d + ex^2}(a + b\operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{\sqrt{ex^2 + d}(a\operatorname{sech}(cx) b + a)}{x^3} dx$$

input

`int((e*x^2+d)^(1/2)*(a+b*asech(c*x))/x^3,x)`

output

`int((e*x^2+d)^(1/2)*(a+b*asech(c*x))/x^3,x)`

### 3.134 $\int x^2 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx$

Optimal result	1098
Mathematica [N/A]	1098
Rubi [N/A]	1099
Maple [N/A]	1099
Fricas [N/A]	1100
Sympy [N/A]	1100
Maxima [F(-2)]	1100
Giac [N/A]	1101
Mupad [N/A]	1101
Reduce [N/A]	1102

#### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int x^2 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \operatorname{Int}\left(x^2 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)), x\right)$$

output `Defer(Int)(x^2*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x)`

#### Mathematica [N/A]

Not integrable

Time = 12.87 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int x^2 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^2 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx$$

input `Integrate[x^2*Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]),x]`

output `Integrate[x^2*Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]), x]`

**Rubi [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx$$

↓ 6865

$$\int x^2 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx$$

input `Int[x^2*Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.17 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int x^2 \sqrt{x^2 e + d} (a + b \operatorname{arcsech}(cx)) dx$$

input `int(x^2*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x)`

output `int(x^2*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x)`

**Fricas [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int x^2 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int \sqrt{ex^2 + d} (b \operatorname{ar} \operatorname{sech}(cx) + a) x^2 dx$$

input `integrate(x^2*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral((b*x^2*arcsech(c*x) + a*x^2)*sqrt(e*x^2 + d), x)`

**Sympy [N/A]**

Not integrable

Time = 9.74 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int x^2 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^2 (a + b \operatorname{asech}(cx)) \sqrt{d + ex^2} dx$$

input `integrate(x**2*(e*x**2+d)**(1/2)*(a+b*asech(c*x)),x)`

output `Integral(x**2*(a + b*asech(c*x))*sqrt(d + e*x**2), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int x^2 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \text{Exception raised: ValueError}$$

input `integrate(x^2*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int x^2 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int \sqrt{ex^2 + d} (b \operatorname{ar} \operatorname{sech}(cx) + a) x^2 dx$$

input

```
integrate(x^2*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="giac")
```

output

```
integrate(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)*x^2, x)
```

**Mupad [N/A]**

Not integrable

Time = 4.46 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int x^2 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^2 \sqrt{ex^2 + d} \left( a + b \operatorname{acosh} \left( \frac{1}{cx} \right) \right) dx$$

input

```
int(x^2*(d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))),x)
```

output

```
int(x^2*(d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))), x)
```

**Reduce [N/A]**

Not integrable

Time = 200.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int x^2 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^2 \sqrt{ex^2 + d} (a \operatorname{sech}(cx) b + a) dx$$

input `int(x^2*(e*x^2+d)^(1/2)*(a+b*asech(c*x)),x)`output `int(x^2*(e*x^2+d)^(1/2)*(a+b*asech(c*x)),x)`

### 3.135 $\int \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx$

Optimal result	1103
Mathematica [N/A]	1103
Rubi [N/A]	1104
Maple [N/A]	1104
Fricas [N/A]	1105
Sympy [N/A]	1105
Maxima [F(-2)]	1105
Giac [N/A]	1106
Mupad [N/A]	1106
Reduce [N/A]	1107

#### Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \operatorname{Int}\left(\sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)), x\right)$$

output `Defer(Int)((e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x)`

#### Mathematica [N/A]

Not integrable

Time = 5.20 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx$$

input `Integrate[Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]),x]`

output `Integrate[Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]), x]`



**Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{d + ex^2}(a + b\operatorname{sech}^{-1}(cx)) dx$$

↓ 6865

$$\int \sqrt{d + ex^2}(a + b\operatorname{sech}^{-1}(cx)) dx$$

input `Int[Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.21 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \sqrt{x^2e + d}(a + b \operatorname{arcsech}(cx)) dx$$

input `int((e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x)`

output `int((e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x)`

**Fricas [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \sqrt{d + ex^2}(a + b \operatorname{sech}^{-1}(cx)) dx = \int \sqrt{ex^2 + d}(b \operatorname{arsech}(cx) + a) dx$$

input `integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a), x)`

**Sympy [N/A]**

Not integrable

Time = 2.55 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \sqrt{d + ex^2}(a + b \operatorname{sech}^{-1}(cx)) dx = \int (a + b \operatorname{asech}(cx)) \sqrt{d + ex^2} dx$$

input `integrate((e*x**2+d)**(1/2)*(a+b*asech(c*x)),x)`

output `Integral((a + b*asech(c*x))*sqrt(d + e*x**2), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \sqrt{d + ex^2}(a + b \operatorname{sech}^{-1}(cx)) dx = \text{Exception raised: ValueError}$$

input `integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \sqrt{d + ex^2}(a + b \operatorname{sech}^{-1}(cx)) dx = \int \sqrt{ex^2 + d}(b \operatorname{ar} \operatorname{sech}(cx) + a) dx$$

input

```
integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="giac")
```

output

```
integrate(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a), x)
```

**Mupad [N/A]**

Not integrable

Time = 4.28 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.20

$$\int \sqrt{d + ex^2}(a + b \operatorname{sech}^{-1}(cx)) dx = \int \sqrt{ex^2 + d} \left( a + b \operatorname{acosh} \left( \frac{1}{cx} \right) \right) dx$$

input

```
int((d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))),x)
```

output

```
int((d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))), x)
```

**Reduce [N/A]**

Not integrable

Time = 200.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int \sqrt{ex^2 + d} (a \operatorname{sech}(cx) b + a) dx$$

input `int((e*x^2+d)^(1/2)*(a+b*asech(c*x)),x)`output `int((e*x^2+d)^(1/2)*(a+b*asech(c*x)),x)`

**3.136** 
$$\int \frac{\sqrt{d+ex^2} \left( a+b\operatorname{sech}^{-1}(cx) \right)}{x^2} dx$$

Optimal result	1108
Mathematica [N/A]	1108
Rubi [N/A]	1109
Maple [N/A]	1109
Fricas [N/A]	1110
Sympy [N/A]	1110
Maxima [F(-2)]	1110
Giac [N/A]	1111
Mupad [N/A]	1111
Reduce [N/A]	1112

**Optimal result**

Integrand size = 23, antiderivative size = 23

$$\int \frac{\sqrt{d+ex^2} \left( a+b\operatorname{sech}^{-1}(cx) \right)}{x^2} dx = \operatorname{Int} \left( \frac{\sqrt{d+ex^2} \left( a+b\operatorname{sech}^{-1}(cx) \right)}{x^2}, x \right)$$

output

```
Defer(Int)((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^2,x)
```

**Mathematica [N/A]**

Not integrable

Time = 1.61 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{\sqrt{d+ex^2} \left( a+b\operatorname{sech}^{-1}(cx) \right)}{x^2} dx = \int \frac{\sqrt{d+ex^2} \left( a+b\operatorname{sech}^{-1}(cx) \right)}{x^2} dx$$

input

```
Integrate[(Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/x^2,x]
```

output

```
Integrate[(Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/x^2, x]
```

**Rubi [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{d + ex^2}(a + b\operatorname{sech}^{-1}(cx))}{x^2} dx$$

↓ 6865

$$\int \frac{\sqrt{d + ex^2}(a + b\operatorname{sech}^{-1}(cx))}{x^2} dx$$

input `Int[(Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/x^2,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{\sqrt{x^2e + d}(a + b \operatorname{arcsech}(cx))}{x^2} dx$$

input `int((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^2,x)`

output `int((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^2,x)`

**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{\sqrt{ex^2+d}(b\operatorname{ar}\operatorname{sech}(cx)+a)}{x^2} dx$$

input `integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^2,x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)/x^2, x)`

**Sympy [N/A]**

Not integrable

Time = 3.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{(a+b\operatorname{asech}(cx))\sqrt{d+ex^2}}{x^2} dx$$

input `integrate((e*x**2+d)**(1/2)*(a+b*asech(c*x))/x**2,x)`

output `Integral((a + b*asech(c*x))*sqrt(d + e*x**2)/x**2, x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^2} dx = \text{Exception raised: ValueError}$$

input `integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^2,x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{\sqrt{ex^2+d}(b\operatorname{arsech}(cx)+a)}{x^2} dx$$

input

```
integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^2,x, algorithm="giac")
```

output

```
integrate(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)/x^2, x)
```

**Mupad [N/A]**

Not integrable

Time = 4.39 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{\sqrt{ex^2+d}(a+b\operatorname{acosh}(\frac{1}{cx}))}{x^2} dx$$

input

```
int(((d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))))/x^2,x)
```

output

```
int(((d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))))/x^2, x)
```



**Reduce [N/A]**

Not integrable

Time = 200.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d + ex^2}(a + b\operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{\sqrt{ex^2 + d}(a\operatorname{sech}(cx) b + a)}{x^2} dx$$

input

`int((e*x^2+d)^(1/2)*(a+b*asech(c*x))/x^2,x)`

output

`int((e*x^2+d)^(1/2)*(a+b*asech(c*x))/x^2,x)`

**3.137** 
$$\int \frac{\sqrt{d+ex^2} \left( a+b\operatorname{sech}^{-1}(cx) \right)}{x^4} dx$$

Optimal result	1113
Mathematica [C] (verified)	1114
Rubi [A] (verified)	1115
Maple [F]	1119
Fricas [A] (verification not implemented)	1119
Sympy [F]	1120
Maxima [F(-2)]	1120
Giac [F]	1120
Mupad [F(-1)]	1121
Reduce [F]	1121

**Optimal result**

Integrand size = 23, antiderivative size = 312

$$\begin{aligned} & \int \frac{\sqrt{d+ex^2} \left( a+b\operatorname{sech}^{-1}(cx) \right)}{x^4} dx \\ &= \frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{9x^3} \\ &+ \frac{2b(c^2d+2e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{9dx} - \frac{(d+ex^2)^{3/2} \left( a+b\operatorname{sech}^{-1}(cx) \right)}{3dx^3} \\ &+ \frac{2bc(c^2d+2e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{d+ex^2}E\left(\arcsin(cx)\left|-\frac{e}{c^2d}\right.\right)}{9d\sqrt{1+\frac{ex^2}{d}}} \\ &- \frac{b(c^2d+e)(2c^2d+3e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1+\frac{ex^2}{d}}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)}{9cd\sqrt{d+ex^2}} \end{aligned}$$

output

$$\frac{1}{9}b*(1/(c*x+1))^{(1/2)}*(c*x+1)^{(1/2)}*(-c^2*x^2+1)^{(1/2)}*(e*x^2+d)^{(1/2)}/x^3+2/9*b*(c^2*d+2*e)*(1/(c*x+1))^{(1/2)}*(c*x+1)^{(1/2)}*(-c^2*x^2+1)^{(1/2)}*(e*x^2+d)^{(1/2)}/d/x-1/3*(e*x^2+d)^{(3/2)}*(a+b*\operatorname{arcsech}(c*x))/d/x^3+2/9*b*c*(c^2*d+2*e)*(1/(c*x+1))^{(1/2)}*(c*x+1)^{(1/2)}*(e*x^2+d)^{(1/2)}*\operatorname{EllipticE}(c*x,(-e/c^2/d)^{(1/2)})/d/(1+e*x^2/d)^{(1/2)}-1/9*b*(c^2*d+e)*(2*c^2*d+3*e)*(1/(c*x+1))^{(1/2)}*(c*x+1)^{(1/2)}*(1+e*x^2/d)^{(1/2)}*\operatorname{EllipticF}(c*x,(-e/c^2/d)^{(1/2)})/c/d/(e*x^2+d)^{(1/2)}$$
**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 23.72 (sec) , antiderivative size = 576, normalized size of antiderivative = 1.85

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx$$

$$\frac{b\sqrt{\frac{1-cx}{1+cx}}(d+ex^2)}{x^3} + \frac{bc\sqrt{\frac{1-cx}{1+cx}}(d+ex^2)}{x^2} + \frac{2b(c^2d+2e)\sqrt{\frac{1-cx}{1+cx}}(d+ex^2)}{dx} - \frac{3a(d+ex^2)^2}{dx^3} - \frac{3b(d+ex^2)^2\operatorname{sech}^{-1}(cx)}{dx^3} - \frac{2ib(c\sqrt{d-i\sqrt{e}})^2}{dx^3}$$

=

input

$$\operatorname{Integrate}[(\operatorname{Sqrt}[d+e*x^2]*(a+b*\operatorname{ArcSech}[c*x]))/x^4,x]$$

output

$$\begin{aligned} & ((b*\operatorname{Sqrt}[(1-c*x)/(1+c*x)]*(d+e*x^2))/x^3 + (b*c*\operatorname{Sqrt}[(1-c*x)/(1+c*x)]*(d+e*x^2))/x^2 + (2*b*(c^2*d+2*e)*\operatorname{Sqrt}[(1-c*x)/(1+c*x)]*(d+e*x^2))/(d*x) - (3*a*(d+e*x^2)^2)/(d*x^3) - (3*b*(d+e*x^2)^2*\operatorname{ArcSech}[c*x))/(d*x^3) - ((2*I)*b*(c*\operatorname{Sqrt}[d-I*\operatorname{Sqrt}[e]]^2*\operatorname{Sqrt}[(1-c*x)/(1+c*x)]*(1+c*x)*\operatorname{Sqrt}[(c*(\operatorname{Sqrt}[d]-I*\operatorname{Sqrt}[e]*x))/((c*\operatorname{Sqrt}[d]-I*\operatorname{Sqrt}[e])*(1+c*x))])*\operatorname{Sqrt}[(c*(\operatorname{Sqrt}[d]+I*\operatorname{Sqrt}[e]*x))/((c*\operatorname{Sqrt}[d]+I*\operatorname{Sqrt}[e])*(1+c*x))])*((c^2*d+2*e)*\operatorname{EllipticE}[I*\operatorname{ArcSinh}[\operatorname{Sqrt}[(c^2*d+e)*(1-c*x)]/((c*\operatorname{Sqrt}[d]+I*\operatorname{Sqrt}[e])^2*(1+c*x))]]), (c*\operatorname{Sqrt}[d]+I*\operatorname{Sqrt}[e])^2/(c*\operatorname{Sqrt}[d]-I*\operatorname{Sqrt}[e])^2 + ((2*I)*c*\operatorname{Sqrt}[d]-3*\operatorname{Sqrt}[e])* \operatorname{Sqrt}[e]*\operatorname{EllipticF}[I*\operatorname{ArcSinh}[\operatorname{Sqrt}[(c^2*d+e)*(1-c*x)]/((c*\operatorname{Sqrt}[d]+I*\operatorname{Sqrt}[e])^2*(1+c*x))]]), (c*\operatorname{Sqrt}[d]+I*\operatorname{Sqrt}[e])^2/(c*\operatorname{Sqrt}[d]-I*\operatorname{Sqrt}[e])^2))/((c*d*\operatorname{Sqrt}[-((c*\operatorname{Sqrt}[d]-I*\operatorname{Sqrt}[e])*(-1+c*x))/((c*\operatorname{Sqrt}[d]+I*\operatorname{Sqrt}[e])*(1+c*x))])))/(9*\operatorname{Sqrt}[d+e*x^2]) \end{aligned}$$

**Rubi [A] (verified)**

Time = 0.60 (sec) , antiderivative size = 256, normalized size of antiderivative = 0.82, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.478$ , Rules used = {6855, 27, 376, 445, 25, 27, 399, 323, 321, 330, 327}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx \\
 & \quad \downarrow \text{6855} \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int -\frac{(ex^2+d)^{3/2}}{3dx^4\sqrt{1-c^2x^2}} dx - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3dx^3} \\
 & \quad \downarrow \text{27} \\
 & -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(ex^2+d)^{3/2}}{x^4\sqrt{1-c^2x^2}} dx}{3d} - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3dx^3} \\
 & \quad \downarrow \text{376} \\
 & -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{3} \int \frac{e(dc^2+3e)x^2+2d(dc^2+2e)}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx - \frac{d\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{3x^3} \right)}{3d} - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3dx^3} \\
 & \quad \downarrow \text{445} \\
 & -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{3} \left( -\frac{\int -\frac{de(-2(dc^2+2e)x^2c^2+dc^2+3e)}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx}{d} - \frac{2\sqrt{1-c^2x^2}(c^2d+2e)\sqrt{d+ex^2}}{x} \right) - \frac{d\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{3x^3} \right)}{3d} - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3dx^3} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\left(\int\frac{de(-2(dc^2+2e)x^2c^2+dc^2+3e)}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx-\frac{2\sqrt{1-c^2x^2}(c^2d+2e)\sqrt{d+ex^2}}{x}\right)-\frac{d\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{3x^3}\right)$$


---


$$\frac{3d}{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))} \frac{3dx^3}{3dx^3}$$

↓ 27

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\left(e\int\frac{-2(dc^2+2e)x^2c^2+dc^2+3e}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx-\frac{2\sqrt{1-c^2x^2}(c^2d+2e)\sqrt{d+ex^2}}{x}\right)-\frac{d\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{3x^3}\right)$$


---


$$\frac{3d}{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))} \frac{3dx^3}{3dx^3}$$

↓ 399

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\left(e\left(\frac{(c^2d+e)(2c^2d+3e)\int\frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx}{e}-\frac{2c^2(c^2d+2e)\int\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}dx}{e}\right)-\frac{2\sqrt{1-c^2x^2}(c^2d+2e)\sqrt{d+ex^2}}{x}\right)\right)$$


---


$$\frac{3d}{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))} \frac{3dx^3}{3dx^3}$$

↓ 323

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\left(e\left(\frac{(c^2d+e)(2c^2d+3e)\sqrt{\frac{ex^2}{d}+1}\int\frac{1}{\sqrt{1-c^2x^2}\sqrt{\frac{ex^2}{d}+1}}dx}{e\sqrt{d+ex^2}}-\frac{2c^2(c^2d+2e)\int\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}dx}{e}\right)-\frac{2\sqrt{1-c^2x^2}(c^2d+2e)\sqrt{d+ex^2}}{x}\right)\right)$$


---


$$\frac{3d}{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))} \frac{3dx^3}{3dx^3}$$

↓ 321

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\left(e\left(\frac{(c^2d+e)(2c^2d+3e)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}}-\frac{2c^2(c^2d+2e)\int\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}dx}{e}\right)-\frac{2\sqrt{1-c^2x^2}(c^2d+2e)\sqrt{d+ex^2}}{x}\right)\right)$$


---


$$\frac{3d}{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))} \frac{3dx^3}{3dx^3}$$

↓ 330

$$\begin{aligned}
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{3} \left( e \left( \frac{(c^2d+e)(2c^2d+3e)\sqrt{\frac{ex^2}{d}+1} \operatorname{EllipticF}\left(\arcsin(cx), -\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}} - \frac{2c^2(c^2d+2e)\sqrt{d+ex^2} \int \frac{\sqrt{\frac{ex^2}{d}+1}}{\sqrt{1-c^2x^2}} dx}{e\sqrt{\frac{ex^2}{d}+1}} \right) - \frac{2\sqrt{1-c^2x^2}}{3d} \right) \right. \\
 & \qquad \qquad \qquad \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{3dx^3} \\
 & \qquad \qquad \qquad \downarrow \text{327} \\
 & \qquad \qquad \qquad \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{3dx^3} \\
 & \left. - \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{3} \left( e \left( \frac{(c^2d+e)(2c^2d+3e)\sqrt{\frac{ex^2}{d}+1} \operatorname{EllipticF}\left(\arcsin(cx), -\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}} - \frac{2c(c^2d+2e)\sqrt{d+ex^2} E\left(\arcsin(cx) \middle| -\frac{e}{c^2d}\right)}{e\sqrt{\frac{ex^2}{d}+1}} \right) - \frac{2\sqrt{1-c^2x^2}}{3d} \right) \right)}{3d} \right)
 \end{aligned}$$

```
input Int[(Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/x^4,x]
```

```
output -1/3*((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/(d*x^3) - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/3*(d*Sqrt[1 - c^2*x^2]*Sqrt[d + e*x^2])/x^3 + ((-2*(c^2*d + 2*e)*Sqrt[1 - c^2*x^2]*Sqrt[d + e*x^2])/x + e*((-2*c*(c^2*d + 2*e)*Sqrt[d + e*x^2]*EllipticE[ArcSin[c*x], -(e/(c^2*d))]))/(e*Sqrt[1 + (e*x^2)/d]) + ((c^2*d + e)*(2*c^2*d + 3*e)*Sqrt[1 + (e*x^2)/d]*EllipticF[ArcSin[c*x], -(e/(c^2*d))]))/(c*e*Sqrt[d + e*x^2])))/(3*d)
```

**Defintions of rubi rules used**

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 321 Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(1/(Sqrt[a]*Sqrt[c]*Rt[-d/c, 2]))*EllipticF[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-b/a, -d/c])
```

rule 323 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[  
Simp[Sqrt[1 + (d/c)*x^2]/Sqrt[c + d*x^2] Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d/c)*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]`

rule 327 `Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[  
(Sqrt[a]/(Sqrt[c]*Rt[-d/c, 2]))*EllipticE[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]`

rule 330 `Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[  
Sqrt[a + b*x^2]/Sqrt[1 + (b/a)*x^2] Int[Sqrt[1 + (b/a)*x^2]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]`

rule 376 `Int[((e_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[c*(e*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q - 1))/(a*e*(m + 1)), x] - Simp[1/(a*e^2*(m + 1)) Int[(e*x)^(m + 2)*(a + b*x^2)^p*(c + d*x^2)^(q - 2)*Simp[c*(b*c - a*d)*(m + 1) + 2*c*(b*c*(p + 1) + a*d*(q - 1)) + d*((b*c - a*d)*(m + 1) + 2*b*c*(p + q))*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && LtQ[m, -1] && IntBinomialQ[a, b, c, d, e, m, 2, p, q, x]`

rule 399 `Int[((e_) + (f_.)*(x_)^2)/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[f/b Int[Sqrt[a + b*x^2]/Sqrt[c + d*x^2], x], x] + Simp[(b*e - a*f)/b Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && !((PosQ[b/a] && PosQ[d/c]) || (NegQ[b/a] && (PosQ[d/c] || (GtQ[a, 0] && (!GtQ[c, 0] || SimplerSqrtQ[-b/a, -d/c]))))`

rule 445 `Int[((g_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.)*((e_) + (f_.)*(x_)^2), x_Symbol] := Simp[e*(g*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1))/(a*c*g*(m + 1)), x] + Simp[1/(a*c*g^2*(m + 1)) Int[(g*x)^(m + 2)*(a + b*x^2)^p*(c + d*x^2)^q*Simp[a*f*c*(m + 1) - e*(b*c + a*d)*(m + 2 + 1) - e*2*(b*c*p + a*d*q) - b*e*d*(m + 2*(p + q + 2) + 1)*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, p, q}, x] && LtQ[m, -1]`

rule 6855

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))*((f_.)*(x_.))^(m_.)*((d_.) + (e_.)*(
x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Si
mp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)]
Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x]] /; Fre
eQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] &&
GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2
*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

**Maple [F]**

$$\int \frac{\sqrt{x^2 e + d} (a + b \operatorname{arcsech}(cx))}{x^4} dx$$

input `int((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^4,x)`

output `int((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^4,x)`

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 242, normalized size of antiderivative = 0.78

$$\int \frac{\sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx =$$

$$\frac{3(bcde x^2 + bcd^2)\sqrt{ex^2 + d} \log\left(\frac{cx\sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2} + 1}}{cx}\right) + (3acde x^2 + 3acd^2 - (bc^2 d^2 x + 2(bc^4 d^2 + 2bc^2 de))}{-}}$$

input `integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^4,x, algorithm="fricas")`

output `-1/9*(3*(b*c*d*e*x^2 + b*c*d^2)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + (3*a*c*d*e*x^2 + 3*a*c*d^2 - (b*c^2*d^2*x + 2*(b*c^4*d^2 + 2*b*c^2*d*e)*x^3)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2))*sqrt(e*x^2 + d) - (2*(b*c^6*d^2 + 2*b*c^4*d*e)*x^3*elliptic_e(arcsin(c*x), -e/(c^2*d)) - (2*b*c^6*d^2 + (4*b*c^4 + b*c^2)*d*e + 3*b*e^2)*x^3*elliptic_f(arcsin(c*x), -e/(c^2*d)))*sqrt(d))/(c*d^2*x^3)`



**Sympy [F]**

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{(a+b\operatorname{asech}(cx))\sqrt{d+ex^2}}{x^4} dx$$

input `integrate((e*x**2+d)**(1/2)*(a+b*asech(c*x))/x**4,x)`

output `Integral((a + b*asech(c*x))*sqrt(d + e*x**2)/x**4, x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx = \text{Exception raised: ValueError}$$

input `integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^4,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{\sqrt{ex^2+d}(b\operatorname{arsech}(cx) + a)}{x^4} dx$$

input `integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^4,x, algorithm="giac")`

output `integrate(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)/x^4, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{\sqrt{ex^2+d}(a+b\operatorname{acosh}(\frac{1}{cx}))}{x^4} dx$$

input `int(((d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))))/x^4,x)`

output `int(((d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))))/x^4, x)`

**Reduce [F]**

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{\sqrt{ex^2+d}(a+b\operatorname{asech}(cx))}{x^4} dx$$

input `int((e*x^2+d)^(1/2)*(a+b*asech(c*x))/x^4,x)`

output `int((e*x^2+d)^(1/2)*(a+b*asech(c*x))/x^4,x)`

$$3.138 \quad \int \frac{\sqrt{d+ex^2} \left( a + b \operatorname{sech}^{-1}(cx) \right)}{x^6} dx$$

Optimal result	1122
Mathematica [C] (verified)	1123
Rubi [A] (verified)	1124
Maple [F]	1129
Fricas [A] (verification not implemented)	1129
Sympy [F]	1130
Maxima [F(-2)]	1130
Giac [F]	1130
Mupad [F(-1)]	1131
Reduce [F]	1131

### Optimal result

Integrand size = 23, antiderivative size = 446

$$\begin{aligned}
 \int \frac{\sqrt{d+ex^2} \left( a + b \operatorname{sech}^{-1}(cx) \right)}{x^6} dx = & \frac{b(12c^2d - e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} \sqrt{d+ex^2}}{225dx^3} \\
 & + \frac{b(24c^4d^2 + 19c^2de - 31e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} \sqrt{d+ex^2}}{225d^2x} \\
 & + \frac{b \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} (d+ex^2)^{3/2}}{25dx^5} \\
 & - \frac{(d+ex^2)^{3/2} \left( a + b \operatorname{sech}^{-1}(cx) \right)}{5dx^5} + \frac{2e(d+ex^2)^{3/2} \left( a + b \operatorname{sech}^{-1}(cx) \right)}{15d^2x^3} \\
 & + \frac{bc(24c^4d^2 + 19c^2de - 31e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{d+ex^2} E(\arcsin(cx) \mid -\frac{e}{c^2d})}{225d^2 \sqrt{1+\frac{ex^2}{d}}} \\
 & - \frac{b(c^2d + e) (24c^4d^2 + 7c^2de - 30e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1+\frac{ex^2}{d}} \operatorname{EllipticF}(\arcsin(cx), -\frac{e}{c^2d})}{225cd^2 \sqrt{d+ex^2}}
 \end{aligned}$$

output

```

1/225*b*(12*c^2*d-e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*(e
*x^2+d)^(1/2)/d/x^3+1/225*b*(24*c^4*d^2+19*c^2*d*e-31*e^2)*(1/(c*x+1))^(1/
2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*(e*x^2+d)^(1/2)/d^2/x+1/25*b*(1/(c*x+1
))^2*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*(e*x^2+d)^(3/2)/d/x^5-1/5*(e*x^2
+d)^(3/2)*(a+b*arcsech(c*x))/d/x^5+2/15*e*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x
))/d^2/x^3+1/225*b*c*(24*c^4*d^2+19*c^2*d*e-31*e^2)*(1/(c*x+1))^(1/2)*(c*x
+1)^(1/2)*(e*x^2+d)^(1/2)*EllipticE(c*x,(-e/c^2/d)^(1/2))/d^2/(1+e*x^2/d)^(
1/2)-1/225*b*(c^2*d+e)*(24*c^4*d^2+7*c^2*d*e-30*e^2)*(1/(c*x+1))^(1/2)*(c
*x+1)^(1/2)*(1+e*x^2/d)^(1/2)*EllipticF(c*x,(-e/c^2/d)^(1/2))/c/d^2/(e*x^2
+d)^(1/2)

```

### Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 25.40 (sec) , antiderivative size = 641, normalized size of antiderivative = 1.44

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx$$

$$= \frac{15a(d+ex^2)^2(-3d+2ex^2)}{x^5} + \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(d+ex^2)(-31e^2x^4+dex^2(8+19c^2x^2)+3d^2(3+4c^2x^2+8c^4x^4))}{x^5} + \frac{15b(d+ex^2)^2(-3d+2ex^2)}{x^5}$$

input

```
Integrate[(Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/x^6,x]
```

output

```

((15*a*(d + e*x^2)^2*(-3*d + 2*e*x^2))/x^5 + (b*Sqrt[(1 - c*x)/(1 + c*x)]*
(1 + c*x)*(d + e*x^2)*(-31*e^2*x^4 + d*e*x^2*(8 + 19*c^2*x^2) + 3*d^2*(3 +
4*c^2*x^2 + 8*c^4*x^4)))/x^5 + (15*b*(d + e*x^2)^2*(-3*d + 2*e*x^2)*ArcSe
ch[c*x])/x^5 + (b*Sqrt[(1 - c*x)/(1 + c*x)]*(-(c^2*(24*c^4*d^2 + 19*c^2*d*
e - 31*e^2)*(d + e*x^2)) - (I*(c*Sqrt[d] - I*Sqrt[e])^2*(1 + c*x)*Sqrt[(c*
(Sqrt[d] - I*Sqrt[e]*x))/((c*Sqrt[d] - I*Sqrt[e])*(1 + c*x))]*Sqrt[(c*(Sqr
t[d] + I*Sqrt[e]*x))/((c*Sqrt[d] + I*Sqrt[e])*(1 + c*x))]*((24*c^4*d^2 + 1
9*c^2*d*e - 31*e^2)*EllipticE[I*ArcSinh[Sqrt[((c^2*d + e)*(1 - c*x))/((c*S
qrt[d] + I*Sqrt[e])^2*(1 + c*x))]]], (c*Sqrt[d] + I*Sqrt[e])^2/(c*Sqrt[d] -
I*Sqrt[e])^2 + 2*Sqrt[e]*((24*I)*c^3*d^(3/2) - 36*c^2*d*Sqrt[e] - (29*I)
*c*Sqrt[d]*e + 30*e^(3/2))*EllipticF[I*ArcSinh[Sqrt[((c^2*d + e)*(1 - c*x)
)/((c*Sqrt[d] + I*Sqrt[e])^2*(1 + c*x))]]], (c*Sqrt[d] + I*Sqrt[e])^2/(c*Sq
rt[d] - I*Sqrt[e])^2))/Sqrt[-(((c*Sqrt[d] - I*Sqrt[e])*(-1 + c*x))/((c*Sq
rt[d] + I*Sqrt[e])*(1 + c*x)))))/c)/(225*d^2*Sqrt[d + e*x^2])

```

## Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 369, normalized size of antiderivative = 0.83, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.522$ , Rules used = {6855, 27, 442, 442, 445, 25, 27, 399, 323, 321, 330, 327}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx \\
 & \quad \downarrow \text{6855} \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int -\frac{(3d-2ex^2)(ex^2+d)^{3/2}}{15d^2x^6\sqrt{1-c^2x^2}} dx + \frac{2e(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{15d^2x^3} - \\
 & \quad \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5} \\
 & \quad \downarrow \text{27} \\
 & -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(3d-2ex^2)(ex^2+d)^{3/2}}{x^6\sqrt{1-c^2x^2}} dx}{15d^2} + \frac{2e(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{15d^2x^3} - \\
 & \quad \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5}
 \end{aligned}$$

$$\begin{aligned} & \downarrow 442 \\ & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\int\frac{\sqrt{ex^2+d}((3c^2d-10e)ex^2+d(12c^2d-e))}{x^4\sqrt{1-c^2x^2}}dx-\frac{3d\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{5x^5}\right)}{15d^2} + \\ & \frac{2e(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{15d^2x^3} - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5} \\ & \downarrow 442 \\ & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\left(\frac{1}{3}\int\frac{2e(6d^2c^4+4dec^2-15e^2)x^2+d(24d^2c^4+19dec^2-31e^2)}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx-\frac{d\sqrt{1-c^2x^2}(12c^2d-e)\sqrt{d+ex^2}}{3x^3}\right)-\frac{3d\sqrt{1-c^2x^2}}{5}\right)}{15d^2} \\ & \frac{2e(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{15d^2x^3} - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5} \\ & \downarrow 445 \\ & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\left(\frac{1}{3}\left(\int-\frac{de(2(6d^2c^4+4dec^2-15e^2)-c^2(24d^2c^4+19dec^2-31e^2)x^2)}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx-\frac{\sqrt{1-c^2x^2}(24c^4d^2+19c^2de-31e^2)\sqrt{d+ex^2}}{x}\right)\right)}{15d^2} \right)}{15d^2} \\ & \frac{2e(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{15d^2x^3} - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5} \\ & \downarrow 25 \\ & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\left(\frac{1}{3}\left(\int\frac{de(2(6d^2c^4+4dec^2-15e^2)-c^2(24d^2c^4+19dec^2-31e^2)x^2)}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx-\frac{\sqrt{1-c^2x^2}(24c^4d^2+19c^2de-31e^2)\sqrt{d+ex^2}}{x}\right)\right)\right)}{15d^2} \\ & \frac{2e(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{15d^2x^3} - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5} \\ & \downarrow 27 \\ & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\left(\frac{1}{3}\left(e\int\frac{2(6d^2c^4+4dec^2-15e^2)-c^2(24d^2c^4+19dec^2-31e^2)x^2}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx-\frac{\sqrt{1-c^2x^2}(24c^4d^2+19c^2de-31e^2)\sqrt{d+ex^2}}{x}\right)\right)\right)}{15d^2} \\ & \frac{2e(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{15d^2x^3} - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5} \\ & \downarrow 399 \end{aligned}$$

$$\begin{aligned}
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{5} \left( \frac{1}{3} \left( e \left( \frac{(c^2d+e)(24c^4d^2+7c^2de-30e^2) \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx}{e} - \frac{c^2(24c^4d^2+19c^2de-31e^2) \int \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} dx}{e} \right) \right) \right) - \right. \\
 & \left. \frac{2e(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{15d^2x^3} - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5} \right) \quad 15d^2 \\
 & \qquad \qquad \qquad \downarrow \text{323} \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{5} \left( \frac{1}{3} \left( e \left( \frac{(c^2d+e)(24c^4d^2+7c^2de-30e^2)\sqrt{\frac{ex^2}{d}+1} \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{\frac{ex^2}{d}+1}} dx}{e\sqrt{d+ex^2}} - \frac{c^2(24c^4d^2+19c^2de-31e^2) \int \frac{\sqrt{\frac{ex^2}{d}+1}}{\sqrt{1-c^2x^2}} dx}{e} \right) \right) \right) - \right. \\
 & \left. \frac{2e(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{15d^2x^3} - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5} \right) \quad 15d^2 \\
 & \qquad \qquad \qquad \downarrow \text{321} \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{5} \left( \frac{1}{3} \left( e \left( \frac{(c^2d+e)(24c^4d^2+7c^2de-30e^2)\sqrt{\frac{ex^2}{d}+1} \operatorname{EllipticF}\left(\arcsin(cx), -\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}} - \frac{c^2(24c^4d^2+19c^2de-31e^2) \int \frac{\sqrt{\frac{ex^2}{d}+1}}{\sqrt{1-c^2x^2}} dx}{e} \right) \right) \right) - \right. \\
 & \left. \frac{2e(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{15d^2x^3} - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5} \right) \quad 15d^2 \\
 & \qquad \qquad \qquad \downarrow \text{330} \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{5} \left( \frac{1}{3} \left( e \left( \frac{(c^2d+e)(24c^4d^2+7c^2de-30e^2)\sqrt{\frac{ex^2}{d}+1} \operatorname{EllipticF}\left(\arcsin(cx), -\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}} - \frac{c^2(24c^4d^2+19c^2de-31e^2)\sqrt{d+ex^2}}{e\sqrt{\frac{ex^2}{d}+1}} \right) \right) \right) - \right. \\
 & \left. \frac{2e(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{15d^2x^3} - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5} \right) \quad 15d \\
 & \qquad \qquad \qquad \downarrow \text{327} \\
 & \frac{2e(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{15d^2x^3} - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5} - \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{5} \left( \frac{1}{3} \left( e \left( \frac{(c^2d+e)(24c^4d^2+7c^2de-30e^2)\sqrt{\frac{ex^2}{d}+1} \operatorname{EllipticF}\left(\arcsin(cx), -\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}} - \frac{c(24c^4d^2+19c^2de-31e^2)\sqrt{d+ex^2}}{e\sqrt{\frac{ex^2}{d}+1}} \right) \right) \right) - \right. \\
 & \left. \frac{2e(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{15d^2x^3} - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5} \right) \quad 15d
 \end{aligned}$$

input `Int[(Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/x^6,x]`

output `-1/5*((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/(d*x^5) + (2*e*(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/(15*d^2*x^3) - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-3*d*Sqrt[1 - c^2*x^2]*(d + e*x^2)^(3/2))/(5*x^5) + (-1/3*(d*(12*c^2*d - e)*Sqrt[1 - c^2*x^2]*Sqrt[d + e*x^2])/x^3 + (-((24*c^4*d^2 + 19*c^2*d*e - 31*e^2)*Sqrt[1 - c^2*x^2]*Sqrt[d + e*x^2])/x) + e*(-((c*(24*c^4*d^2 + 19*c^2*d*e - 31*e^2)*Sqrt[d + e*x^2]*EllipticE[ArcSin[c*x], -(e/(c^2*d))]))/(e*Sqrt[1 + (e*x^2)/d])) + ((c^2*d + e)*(24*c^4*d^2 + 7*c^2*d*e - 30*e^2)*Sqrt[1 + (e*x^2)/d]*EllipticF[ArcSin[c*x], -(e/(c^2*d))])/(c*e*Sqrt[d + e*x^2])))/3)/5)/(15*d^2)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 321 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(1/(Sqrt[a]*Sqrt[c]*Rt[-d/c, 2]))*EllipticF[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-b/a, -d/c])`

rule 323 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[Sqrt[1 + (d/c)*x^2]/Sqrt[c + d*x^2] Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d/c)*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]`

rule 327 `Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]/(Sqrt[c]*Rt[-d/c, 2]))*EllipticE[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]`



rule 330 `Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[Sqrt[a + b*x^2]/Sqrt[1 + (b/a)*x^2] Int[Sqrt[1 + (b/a)*x^2]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]`

rule 399 `Int[((e_) + (f_.)*(x_)^2)/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[f/b Int[Sqrt[a + b*x^2]/Sqrt[c + d*x^2], x], x] + Simp[(b*e - a*f)/b Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && !((PosQ[b/a] && PosQ[d/c]) || (NegQ[b/a] && (PosQ[d/c] || (GtQ[a, 0] && (!GtQ[c, 0] || SimplerSqrtQ[-b/a, -d/c]))))`

rule 442 `Int[((g_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.)*((e_) + (f_.)*(x_)^2), x_Symbol] := Simp[e*(g*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^q/(a*g*(m + 1))), x] - Simp[1/(a*g^2*(m + 1)) Int[(g*x)^(m + 2)*(a + b*x^2)^p*(c + d*x^2)^(q - 1)*Simp[c*(b*e - a*f)*(m + 1) + e*2*(b*c*(p + 1) + a*d*q) + d*((b*e - a*f)*(m + 1) + b*e*2*(p + q + 1))*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && GtQ[q, 0] && LtQ[m, -1] && !(EqQ[q, 1] && SimplerQ[e + f*x^2, c + d*x^2])`

rule 445 `Int[((g_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.)*((e_) + (f_.)*(x_)^2), x_Symbol] := Simp[e*(g*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(a*c*g*(m + 1))), x] + Simp[1/(a*c*g^2*(m + 1)) Int[(g*x)^(m + 2)*(a + b*x^2)^p*(c + d*x^2)^q*Simp[a*f*c*(m + 1) - e*(b*c + a*d)*(m + 2 + 1) - e*2*(b*c*p + a*d*q) - b*e*d*(m + 2*(p + q + 2) + 1)*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, p, q}, x] && LtQ[m, -1]`

rule 6855 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))`

**Maple [F]**

$$\int \frac{\sqrt{x^2 e + d} (a + b \operatorname{arcsech}(cx))}{x^6} dx$$

input `int((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^6,x)`

output `int((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^6,x)`

**Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 340, normalized size of antiderivative = 0.76

$$\int \frac{\sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx$$

$$= \frac{15(2bcde^2x^4 - bcd^2ex^2 - 3bcd^3)\sqrt{ex^2 + d} \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}+1}}{cx}\right) + (30acde^2x^4 - 15acd^2ex^2 - 45acd^3 + \dots}{\dots}$$

input `integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^6,x, algorithm="fricas")`

output `1/225*(15*(2*b*c*d*e^2*x^4 - b*c*d^2*e*x^2 - 3*b*c*d^3)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + (30*a*c*d*e^2*x^4 - 15*a*c*d^2*e*x^2 - 45*a*c*d^3 + (9*b*c^2*d^3*x + (24*b*c^6*d^3 + 19*b*c^4*d^2*e - 31*b*c^2*d*e^2)*x^5 + 4*(3*b*c^4*d^3 + 2*b*c^2*d^2*e)*x^3)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d) + ((24*b*c^8*d^3 + 19*b*c^6*d^2*e - 31*b*c^4*d*e^2)*x^5*elliptic_e(arcsin(c*x), -e/(c^2*d)) - (24*b*c^8*d^3 + (19*b*c^6 + 12*b*c^4)*d^2*e - (31*b*c^4 - 8*b*c^2)*d*e^2 - 30*b*e^3)*x^5*elliptic_f(arcsin(c*x), -e/(c^2*d))*sqrt(d))/(c*d^3*x^5)`

**Sympy [F]**

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx = \int \frac{(a+b\operatorname{asech}(cx))\sqrt{d+ex^2}}{x^6} dx$$

input `integrate((e*x**2+d)**(1/2)*(a+b*asech(c*x))/x**6,x)`

output `Integral((a + b*asech(c*x))*sqrt(d + e*x**2)/x**6, x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx = \text{Exception raised: ValueError}$$

input `integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^6,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx = \int \frac{\sqrt{ex^2+d}(b\operatorname{arsech}(cx) + a)}{x^6} dx$$

input `integrate((e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/x^6,x, algorithm="giac")`

output `integrate(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)/x^6, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx = \int \frac{\sqrt{ex^2+d}(a+b\operatorname{acosh}(\frac{1}{cx}))}{x^6} dx$$

input `int(((d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))))/x^6,x)`output `int(((d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))))/x^6, x)`**Reduce [F]**

$$\int \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{x^6} dx = \int \frac{\sqrt{ex^2+d}(a+b\operatorname{asech}(cx))}{x^6} dx$$

input `int((e*x^2+d)^(1/2)*(a+b*asech(c*x))/x^6,x)`output `int((e*x^2+d)^(1/2)*(a+b*asech(c*x))/x^6,x)`

### 3.139 $\int x^3(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	1132
Mathematica [A] (verified)	1133
Rubi [A] (verified)	1134
Maple [F]	1139
Fricas [A] (verification not implemented)	1139
Sympy [F(-1)]	1140
Maxima [F(-2)]	1141
Giac [F]	1141
Mupad [F(-1)]	1141
Reduce [F]	1142

#### Optimal result

Integrand size = 23, antiderivative size = 418

$$\begin{aligned}
 & \int x^3(d + ex^2)^{3/2} (a \\
 & + b\operatorname{sech}^{-1}(cx)) dx = \frac{b(3c^4d^2 - 38c^2de - 25e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} \sqrt{d+ex^2}}{560c^6e} \\
 & - \frac{b(13c^2d + 25e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} (d+ex^2)^{3/2}}{840c^4e} \\
 & - \frac{b \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} (d+ex^2)^{5/2}}{42c^2e} \\
 & - \frac{d(d+ex^2)^{5/2} (a + b\operatorname{sech}^{-1}(cx))}{5e^2} + \frac{(d+ex^2)^{7/2} (a + b\operatorname{sech}^{-1}(cx))}{7e^2} \\
 & + \frac{b(35c^6d^3 - 35c^4d^2e - 63c^2de^2 - 25e^3) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{560c^7e^{3/2}} \\
 & + \frac{2bd^{7/2} \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{35e^2}
 \end{aligned}$$

output

$$\frac{1}{560}b(3c^4d^2 - 38c^2de - 25e^2) \frac{1}{(cx+1)^{1/2}} (cx+1)^{1/2} (-c^2x^2+1)^{1/2} (ex^2+d)^{1/2} / c^6/e - 1/840b(13c^2d+25e) \frac{1}{(cx+1)^{1/2}} (cx+1)^{1/2} (-c^2x^2+1)^{1/2} (ex^2+d)^{3/2} / c^4/e - 1/42b \frac{1}{(cx+1)^{1/2}} (cx+1)^{1/2} (-c^2x^2+1)^{1/2} (ex^2+d)^{5/2} / c^2/e - 1/5d(e^{ex^2+d})^{5/2} (a+b \operatorname{arcsech}(cx)) / e^{2+1/7} (ex^2+d)^{7/2} (a+b \operatorname{arcsech}(cx)) / e^{2+1/560} b(35c^6d^3 - 35c^4d^2e - 63c^2de^2 - 25e^3) \frac{1}{(cx+1)^{1/2}} (cx+1)^{1/2} \arctan(e^{1/2}(-c^2x^2+1)^{1/2}/c/(ex^2+d)^{1/2}) / c^7 / e^{3/2} + 2/35bd^{7/2} \frac{1}{(cx+1)^{1/2}} (cx+1)^{1/2} \operatorname{arctanh}((ex^2+d)^{1/2}/d^{1/2}/(-c^2x^2+1)^{1/2}) / e^2$$
**Mathematica [A] (verified)**

Time = 35.49 (sec) , antiderivative size = 313, normalized size of antiderivative = 0.75

$$\int x^3(d+ex^2)^{3/2}(a+b \operatorname{sech}^{-1}(cx)) dx =$$

$$\frac{\sqrt{d+ex^2} \left( 48ac^6(2d-5ex^2)(d+ex^2)^2 + be\sqrt{\frac{1-cx}{1+cx}}(1+cx)(75e^2+2c^2e(82d+25ex^2)) + c^4(57d^2+106d^2+40e^2x^4) \right) + 48b \frac{1-cx}{1+cx} \sqrt{-1+c^2x^2} \left( -32c^7d^{7/2} \arctan\left(\frac{\sqrt{d}\sqrt{-1+c^2x^2}}{\sqrt{d+ex^2}}\right) + \sqrt{e}(-35c^6d^3+35c^4d^2e+63c^2de^2+25e^3) \operatorname{arctan}\left(\frac{\sqrt{d}\sqrt{-1+c^2x^2}}{\sqrt{d+ex^2}}\right) \right)}{1680c^6e^2}$$

$$\frac{b\sqrt{\frac{1-cx}{1+cx}}\sqrt{-1+c^2x^2}\left(-32c^7d^{7/2}\arctan\left(\frac{\sqrt{d}\sqrt{-1+c^2x^2}}{\sqrt{d+ex^2}}\right)+\sqrt{e}(-35c^6d^3+35c^4d^2e+63c^2de^2+25e^3)\arctan\left(\frac{\sqrt{d}\sqrt{-1+c^2x^2}}{\sqrt{d+ex^2}}\right)\right)}{560c^7e^2(-1+cx)}$$

input

`Integrate[x^3*(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]),x]`

output

$$\frac{-1}{1680} \left( \sqrt{d+ex^2} (48ac^6(2d-5ex^2)(d+ex^2)^2 + b e \sqrt{\frac{1-cx}{1+cx}}(1+cx)(75e^2+2c^2e(82d+25ex^2)) + c^4(57d^2+106d^2+40e^2x^4)) + 48b \frac{1-cx}{1+cx} \sqrt{-1+c^2x^2} \left( -32c^7d^{7/2} \operatorname{ArcTan}\left[\frac{\sqrt{d}\sqrt{-1+c^2x^2}}{\sqrt{d+ex^2}}\right] + \sqrt{e}(-35c^6d^3+35c^4d^2e+63c^2de^2+25e^3) \operatorname{ArcTanh}\left[\frac{\sqrt{d}\sqrt{-1+c^2x^2}}{\sqrt{d+ex^2}}\right] \right) \right) / (560c^7e^2(-1+cx))$$

**Rubi [A] (verified)**

Time = 0.67 (sec) , antiderivative size = 354, normalized size of antiderivative = 0.85, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.609$ , Rules used = {6855, 27, 435, 171, 27, 171, 27, 171, 27, 175, 66, 104, 218, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx \\
 & \quad \downarrow \text{6855} \\
 & b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int -\frac{(2d-5ex^2)(ex^2+d)^{5/2}}{35e^2 x \sqrt{1-c^2x^2}} dx + \frac{(d+ex^2)^{7/2} (a+b \operatorname{sech}^{-1}(cx))}{7e^2} - \\
 & \quad \frac{d(d+ex^2)^{5/2} (a+b \operatorname{sech}^{-1}(cx))}{5e^2} \\
 & \quad \downarrow \text{27} \\
 & -\frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{(2d-5ex^2)(ex^2+d)^{5/2}}{x \sqrt{1-c^2x^2}} dx}{35e^2} + \frac{(d+ex^2)^{7/2} (a+b \operatorname{sech}^{-1}(cx))}{7e^2} - \\
 & \quad \frac{d(d+ex^2)^{5/2} (a+b \operatorname{sech}^{-1}(cx))}{5e^2} \\
 & \quad \downarrow \text{435} \\
 & -\frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{(2d-5ex^2)(ex^2+d)^{5/2}}{x^2 \sqrt{1-c^2x^2}} dx^2}{70e^2} + \frac{(d+ex^2)^{7/2} (a+b \operatorname{sech}^{-1}(cx))}{7e^2} - \\
 & \quad \frac{d(d+ex^2)^{5/2} (a+b \operatorname{sech}^{-1}(cx))}{5e^2} \\
 & \quad \downarrow \text{171} \\
 & -\frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \left( \frac{5e \sqrt{1-c^2x^2} (d+ex^2)^{5/2}}{3c^2} - \frac{\int -\frac{(ex^2+d)^{3/2} (12c^2d^2 - e(13dc^2+25e)x^2)}{2x^2 \sqrt{1-c^2x^2}} dx^2}{3c^2} \right)}{70e^2} + \\
 & \quad \frac{(d+ex^2)^{7/2} (a+b \operatorname{sech}^{-1}(cx))}{7e^2} - \frac{d(d+ex^2)^{5/2} (a+b \operatorname{sech}^{-1}(cx))}{5e^2} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{\int\frac{(ex^2+d)^{3/2}(12c^2d^2-e(13dc^2+25e)x^2)}{x^2\sqrt{1-c^2x^2}}dx^2}{6c^2}+\frac{5e\sqrt{1-c^2x^2}(d+ex^2)^{5/2}}{3c^2}\right)$$


---


$$\frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^2}-\frac{70e^2}{5e^2}\frac{d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2}$$

↓ 171

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{e\sqrt{1-c^2x^2}(13c^2d+25e)(d+ex^2)^{3/2}}{2c^2}-\frac{\int-\frac{3\sqrt{ex^2+d}(16d^3c^4+e(3d^2c^4-38dec^2-25e^2)x^2)}{2x^2\sqrt{1-c^2x^2}}dx^2}{6c^2}+\frac{5e\sqrt{1-c^2x^2}(d+ex^2)^{5/2}}{3c^2}\right)$$


---


$$\frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^2}-\frac{70e^2}{5e^2}\frac{d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2}$$

↓ 27

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{3\int\frac{\sqrt{ex^2+d}(16d^3c^4+e(3d^2c^4-38dec^2-25e^2)x^2)}{x^2\sqrt{1-c^2x^2}}dx^2}{4c^2}+\frac{e\sqrt{1-c^2x^2}(13c^2d+25e)(d+ex^2)^{3/2}}{2c^2}+\frac{5e\sqrt{1-c^2x^2}(d+ex^2)^{5/2}}{3c^2}\right)$$


---


$$\frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^2}-\frac{70e^2}{5e^2}\frac{d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2}$$

↓ 171

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{3\left(\frac{\int-\frac{32d^4c^6+e(35d^3c^6-35d^2ec^4-63de^2c^2-25e^3)x^2}{2x^2\sqrt{1-c^2x^2}}dx^2}{c^2}-\frac{e\sqrt{1-c^2x^2}(3c^4d^2-38c^2de-25e^2)\sqrt{d+ex^2}}{c^2}\right)}{4c^2}+\frac{e\sqrt{1-c^2x^2}(13c^2d+25e)(d+ex^2)^{3/2}}{2c^2}+\frac{5e\sqrt{1-c^2x^2}(d+ex^2)^{5/2}}{3c^2}\right)$$


---


$$\frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^2}-\frac{70e^2}{5e^2}\frac{d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2}$$

↓ 27



$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3 \left( \int \frac{32d^4c^6 + e(35d^3c^6 - 35d^2ec^4 - 63de^2c^2 - 25e^3)x^2}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 - \frac{e\sqrt{1-c^2x^2}(3c^4d^2 - 38c^2de - 25e^2)\sqrt{d+ex^2}}{c^2} \right)}{4c^2} + \frac{e\sqrt{1-c^2x^2}(13c^2d+25e^2)}{6c^2} \right)$$

$$\frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^2} - \frac{d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2} \quad 70e^2$$

↓ 175

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3 \left( \frac{32c^6d^4 \int \frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 + e(35c^6d^3 - 35c^4d^2e - 63c^2de^2 - 25e^3) \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 - \frac{e\sqrt{1-c^2x^2}(3c^4d^2 - 38c^2de - 25e^2)}{c^2} \right)}{4c^2} + \frac{e\sqrt{1-c^2x^2}(13c^2d+25e^2)}{6c^2} \right)$$

$$\frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^2} - \frac{d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2} \quad 70e^2$$

↓ 66

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3 \left( \frac{32c^6d^4 \int \frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 + 2e(35c^6d^3 - 35c^4d^2e - 63c^2de^2 - 25e^3) \int \frac{1}{-ex^4-c^2} d \frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}} - \frac{e\sqrt{1-c^2x^2}(3c^4d^2 - 38c^2de - 25e^2)}{c^2} \right)}{4c^2} + \frac{e\sqrt{1-c^2x^2}(13c^2d+25e^2)}{6c^2} \right)$$

$$\frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^2} - \frac{d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2} \quad 70e^2$$

↓ 104

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3 \left( \frac{64c^6d^4 \int \frac{1}{x^4-d} d \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} + 2e(35c^6d^3 - 35c^4d^2e - 63c^2de^2 - 25e^3) \int \frac{1}{-ex^4-c^2} d \frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}} - \frac{e\sqrt{1-c^2x^2}(3c^4d^2 - 38c^2de - 25e^2)}{c^2} \right)}{4c^2} + \frac{e\sqrt{1-c^2x^2}(13c^2d+25e^2)}{6c^2} \right)$$

$$\frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^2} - \frac{d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2} \quad 70e^2$$

218

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3 \left( \frac{64c^6d^4 \int \frac{1}{x^4-d} d \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} - \frac{2\sqrt{e}(35c^6d^3-35c^4d^2e-63c^2de^2-25e^3)}{c} \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{2c^2} - \frac{e\sqrt{1-c^2x^2}(3c^4d^2-38c^2de-25e^2)}{c^2} \right)}{4c^2} - \frac{\phantom{3 \left( \dots \right)}}{6c^2} \right)$$

---


$$\frac{(d+ex^2)^{7/2}(a+b\operatorname{sech}^{-1}(cx))}{7e^2} - \frac{d(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^2} \quad 70e^2$$

220

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{3 \left( \frac{2\sqrt{e}(35c^6d^3-35c^4d^2e-63c^2de^2-25e^3)}{c} \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right) - \frac{5e^2}{2c^2} - 64c^6d^{7/2}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right) - \frac{e\sqrt{1-c^2x^2}(3c^4d^2-38c^2de-25e^2)}{c^2} \right)}{4c^2} - \frac{\phantom{3 \left( \dots \right)}}{6c^2} \right)$$

70e<sup>2</sup>

```
input Int[x^3*(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]),x]
```

```
output -1/5*(d*(d + e*x^2)^(5/2)*(a + b*ArcSech[c*x]))/e^2 + ((d + e*x^2)^(7/2)*(a + b*ArcSech[c*x]))/(7*e^2) - (b*sqrt[(1 + c*x)^(-1)]*sqrt[1 + c*x]*((5*e*sqrt[1 - c^2*x^2]*(d + e*x^2)^(5/2))/(3*c^2) + ((e*(13*c^2*d + 25*e)*sqrt[1 - c^2*x^2]*(d + e*x^2)^(3/2))/(2*c^2) + (3*(-((e*(3*c^4*d^2 - 38*c^2*d*e - 25*e^2)*sqrt[1 - c^2*x^2]*sqrt[d + e*x^2])/c^2) + ((-2*sqrt[e]*(35*c^6*d^3 - 35*c^4*d^2*e - 63*c^2*d*e^2 - 25*e^3)*ArcTan[(sqrt[e]*sqrt[1 - c^2*x^2])/(c*sqrt[d + e*x^2])])/c - 64*c^6*d^(7/2)*ArcTanh[Sqrt[d + e*x^2]/(sqrt[d]*sqrt[1 - c^2*x^2])])/(2*c^2)))/(4*c^2))/(6*c^2))/(70*e^2)
```

## Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx) /; FreeQ[b, x]]`
- rule 66 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`
- rule 104 `Int[(((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 171 `Int[(((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_)*((g_) + (h_)*(x_)), x_] := Simp[h*(a + b*x)^m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Simp[1/(d*f*(m + n + p + 2)) Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegersQ[2*m, 2*n, 2*p]`
- rule 175 `Int[(((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_)*((g_) + (h_)*(x_)))/((a_) + (b_)*(x_)), x_] := Simp[h/b Int[(c + d*x)^n*(e + f*x)^p, x], x] + Simp[(b*g - a*h)/b Int[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x]`
- rule 218 `Int[(((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 220 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 435 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2)^(r_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2) * (a + b*x)^p * (c + d*x)^q * (e + f*x)^r, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, f, p, q, r}, x] && IntegerQ[(m - 1)/2]`

rule 6855 `Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*((f_)*(x_)^(m_))*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))`

## Maple [F]

$$\int x^3 (x^2 e + d)^{\frac{3}{2}} (a + b \operatorname{arcsech}(cx)) dx$$

input `int(x^3*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x)`

output `int(x^3*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x)`

## Fricas [A] (verification not implemented)

Time = 1.49 (sec) , antiderivative size = 1989, normalized size of antiderivative = 4.76

$$\int x^3 (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \text{Too large to display}$$

input `integrate(x^3*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output

```
[1/6720*(96*b*c^7*d^(7/2)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 - 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) + 3*(35*b*c^6*d^3 - 35*b*c^4*d^2*e - 63*b*c^2*d*e^2 - 25*b*e^3)*sqrt(-e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e + 8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^4*e*x^3 + (c^4*d - c^2*e)*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e^2) + 192*(5*b*c^7*e^3*x^6 + 8*b*c^7*d*e^2*x^4 + b*c^7*d^2*e*x^2 - 2*b*c^7*d^3)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 4*(240*a*c^7*e^3*x^6 + 384*a*c^7*d*e^2*x^4 + 48*a*c^7*d^2*e*x^2 - 96*a*c^7*d^3 - (40*b*c^6*e^3*x^5 + 2*(53*b*c^6*d*e^2 + 25*b*c^4*e^3)*x^3 + (57*b*c^6*d^2*e + 164*b*c^4*d*e^2 + 75*b*c^2*e^3)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d))/(c^7*e^2), 1/3360*(48*b*c^7*d^(7/2)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 - 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) + 3*(35*b*c^6*d^3 - 35*b*c^4*d^2*e - 63*b*c^2*d*e^2 - 25*b*e^3)*sqrt(e)*arctan(1/2*(2*c^2*e*x^3 + (c^2*d - e)*x)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*e^2*x^4 + (c^2*d*e - e^2)*x^2 - d*e) + 96*(5*b*c^7*e^3*x^6 + 8*b*c^7*d*e^2*x^4 + b*c^7*d^2*e*x^2 - 2*b*c^7*d^3)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 2*(240*a*c^7*e^3*x^6 + 384*a*c^7*d*e^2*x^4 + 48*a*c^7*d^2*e*x^2 - 96*a*c^7*d^3 - (40*b*c^6*e^3*x^5 + ...
```

**Sympy [F(-1)]**

Timed out.

$$\int x^3(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \text{Timed out}$$

input

```
integrate(x**3*(e*x**2+d)**(3/2)*(a+b*asech(c*x)),x)
```

output

Timed out

**Maxima [F(-2)]**

Exception generated.

$$\int x^3 (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \text{Exception raised: ValueError}$$

input `integrate(x^3*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int x^3 (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^{\frac{3}{2}} (b \operatorname{ar} \operatorname{sech}(cx) + a) x^3 dx$$

input `integrate(x^3*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((e*x^2 + d)^(3/2)*(b*arcsech(c*x) + a)*x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^3 (ex^2 + d)^{3/2} \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int(x^3*(d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))),x)`

output `int(x^3*(d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))), x)`

**Reduce [F]**

$$\int x^3 (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^3 (ex^2 + d)^{\frac{3}{2}} (a \operatorname{sech}(cx) b + a) dx$$

input `int(x^3*(e*x^2+d)^(3/2)*(a+b*asech(c*x)),x)`

output `int(x^3*(e*x^2+d)^(3/2)*(a+b*asech(c*x)),x)`

### 3.140 $\int x(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	1143
Mathematica [A] (verified)	1144
Rubi [A] (verified)	1144
Maple [F]	1149
Fricas [B] (verification not implemented)	1149
Sympy [F]	1150
Maxima [F]	1151
Giac [F]	1151
Mupad [F(-1)]	1152
Reduce [F]	1152

#### Optimal result

Integrand size = 21, antiderivative size = 297

$$\int x(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx =$$

$$\frac{b(7c^2d + 3e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} \sqrt{d+ex^2}}{40c^4}$$

$$- \frac{b\sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} (d+ex^2)^{3/2}}{20c^2} + \frac{(d+ex^2)^{5/2} (a + b\operatorname{sech}^{-1}(cx))}{5e}$$

$$- \frac{b(15c^4d^2 + 10c^2de + 3e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{40c^5\sqrt{e}}$$

$$- \frac{bd^{5/2} \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{5e}$$

output

```
-1/40*b*(7*c^2*d+3*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*(
e*x^2+d)^(1/2)/c^4-1/20*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*
(e*x^2+d)^(3/2)/c^2+1/5*(e*x^2+d)^(5/2)*(a+b*arcsech(c*x))/e-1/40*b*(15
*c^4*d^2+10*c^2*d*e+3*e^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctan(e^(1/2)*
(-c^2*x^2+1)^(1/2)/c/(e*x^2+d)^(1/2))/c^5/e^(1/2)-1/5*b*d^(5/2)*(1/(c*x+1)
)^(1/2)*(c*x+1)^(1/2)*arctanh((e*x^2+d)^(1/2)/d^(1/2)/(-c^2*x^2+1)^(1/2))/
e
```



### Mathematica [A] (verified)

Time = 22.12 (sec) , antiderivative size = 342, normalized size of antiderivative = 1.15

$$\int x(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \frac{\sqrt{d + ex^2} \left( 8ac^4(d + ex^2)^2 - be \sqrt{\frac{1-cx}{1+cx}} (1 + cx) (3e + c^2(9d + 2ex^2)) + 8bc^4(d + ex^2)^2 \right)}{40c^4e} + \frac{b \sqrt{\frac{1-cx}{1+cx}} \sqrt{1 - c^2x^2} \left( \sqrt{-c^2} \sqrt{-c^2d - e} \sqrt{e} (15c^4d^2 + 10c^2de + 3e^2) \sqrt{\frac{c^2(d+ex^2)}{c^2d+e}} \arcsin \left( \frac{c\sqrt{e}\sqrt{1-c^2x^2}}{\sqrt{-c^2}\sqrt{-c^2d-e}} \right) + 8c^7d \right)}{40c^7e(-1 + cx)\sqrt{d + ex^2}}$$

input

```
Integrate[x*(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]),x]
```

output

```
(Sqrt[d + e*x^2]*(8*a*c^4*(d + e*x^2)^2 - b*e*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(3*e + c^2*(9*d + 2*e*x^2)) + 8*b*c^4*(d + e*x^2)^2*ArcSech[c*x]))/(40*c^4*e) + (b*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*(Sqrt[-c^2]*Sqrt[-(c^2*d) - e]*Sqrt[e]*(15*c^4*d^2 + 10*c^2*d*e + 3*e^2)*Sqrt[(c^2*(d + e*x^2))/(c^2*d + e)]*ArcSin[(c*Sqrt[e]*Sqrt[1 - c^2*x^2])/(Sqrt[-c^2]*Sqrt[-(c^2*d) - e]]) + 8*c^7*d^(5/2)*Sqrt[-d - e*x^2]*ArcTan[(Sqrt[d]*Sqrt[1 - c^2*x^2])/Sqrt[-d - e*x^2]]))/(40*c^7*e*(-1 + c*x)*Sqrt[d + e*x^2])
```

### Rubi [A] (verified)

Time = 0.65 (sec) , antiderivative size = 254, normalized size of antiderivative = 0.86, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$ , Rules used = {6853, 2036, 354, 113, 27, 171, 27, 175, 66, 104, 218, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx \quad \downarrow \text{6853}$$

$$\frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx + 1} \int \frac{(ex^2+d)^{5/2}}{x\sqrt{1-cx}\sqrt{cx+1}} dx}{5e} + \frac{(d + ex^2)^{5/2} (a + b \operatorname{sech}^{-1}(cx))}{5e}$$

$$\begin{aligned}
 & \downarrow 2036 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(ex^2+d)^{5/2}}{x\sqrt{1-c^2x^2}} dx}{5e} + \frac{(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{5e} \\
 & \downarrow 354 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(ex^2+d)^{5/2}}{x^2\sqrt{1-c^2x^2}} dx^2}{10e} + \frac{(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{5e} \\
 & \downarrow 113 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{\int -\frac{\sqrt{ex^2+d}(4c^2d^2+e(7dc^2+3e)x^2)}{2x^2\sqrt{1-c^2x^2}} dx^2}{2c^2} - \frac{e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2} \right)}{10e} + \\
 & \frac{(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{5e} \\
 & \downarrow 27 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{\sqrt{ex^2+d}(4c^2d^2+e(7dc^2+3e)x^2)}{x^2\sqrt{1-c^2x^2}} dx^2}{4c^2} - \frac{e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2} \right)}{10e} + \\
 & \frac{(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{5e} \\
 & \downarrow 171 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{\int -\frac{8d^3c^4+e(15d^2c^4+10dec^2+3e^2)x^2}{2x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2}{c^2} - \frac{e\sqrt{1-c^2x^2}(7c^2d+3e)\sqrt{d+ex^2}}{c^2} - \frac{e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2} \right)}{10e} + \\
 & \frac{(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{5e} \\
 & \downarrow 27
 \end{aligned}$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{8d^3c^4+e(15d^2c^4+10dec^2+3e^2)x^2}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2}{2c^2} - \frac{e\sqrt{1-c^2x^2}(7c^2d+3e)\sqrt{d+ex^2}}{c^2} - \frac{e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2} \right)$$


---


$$\frac{10e}{(d+ex^2)^{5/2}} (a + b\operatorname{sech}^{-1}(cx))$$

$$\downarrow 175$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{8c^4d^3 \int \frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 + e(15c^4d^2+10c^2de+3e^2) \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2}{2c^2} - \frac{e\sqrt{1-c^2x^2}(7c^2d+3e)\sqrt{d+ex^2}}{c^2} - \frac{e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2} \right)$$


---


$$\frac{10e}{(d+ex^2)^{5/2}} (a + b\operatorname{sech}^{-1}(cx))$$

$$\downarrow 66$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{8c^4d^3 \int \frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 + 2e(15c^4d^2+10c^2de+3e^2) \int \frac{1}{-ex^4-c^2} d \frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}}}{2c^2} - \frac{e\sqrt{1-c^2x^2}(7c^2d+3e)\sqrt{d+ex^2}}{c^2} - \frac{e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2} \right)$$


---


$$\frac{10e}{(d+ex^2)^{5/2}} (a + b\operatorname{sech}^{-1}(cx))$$

$$\downarrow 104$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{16c^4d^3 \int \frac{1}{x^4-d} d \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} + 2e(15c^4d^2+10c^2de+3e^2) \int \frac{1}{-ex^4-c^2} d \frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}}}{2c^2} - \frac{e\sqrt{1-c^2x^2}(7c^2d+3e)\sqrt{d+ex^2}}{c^2} - \frac{e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2} \right)$$


---


$$\frac{10e}{(d+ex^2)^{5/2}} (a + b\operatorname{sech}^{-1}(cx))$$

$$\downarrow 218$$

$$\begin{aligned}
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{16c^4d^3 \int \frac{1}{x^4-d} d \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} - \frac{2\sqrt{e}(15c^4d^2+10c^2de+3e^2) \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c} - \frac{e\sqrt{1-c^2x^2}(7c^2d+3e)\sqrt{d+ex^2}}{c^2} - \frac{e\sqrt{1-c^2x^2}}{2c} \right) \\
 & \frac{(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{5e} \\
 & \quad \downarrow \text{220} \\
 & \frac{(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{5e} + \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{-\frac{2\sqrt{e}(15c^4d^2+10c^2de+3e^2) \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c} - 16c^4d^{5/2}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right) - \frac{e\sqrt{1-c^2x^2}(7c^2d+3e)\sqrt{d+ex^2}}{c^2} - e}{2c^2} - \frac{e}{4c^2} \right) \\
 & \frac{10e}{10e}
 \end{aligned}$$

input

```
Int[x*(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]), x]
```

output

```
((d + e*x^2)^(5/2)*(a + b*ArcSech[c*x]))/(5*e) + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/2*(e*Sqrt[1 - c^2*x^2]*(d + e*x^2)^(3/2))/c^2 + (-((e*(7*c^2*d + 3*e)*Sqrt[1 - c^2*x^2]*Sqrt[d + e*x^2])/c^2) + ((-2*Sqrt[e]*(15*c^4*d^2 + 10*c^2*d*e + 3*e^2)*ArcTan[(Sqrt[e]*Sqrt[1 - c^2*x^2])/(c*Sqrt[d + e*x^2])])/c - 16*c^4*d^(5/2)*ArcTanh[Sqrt[d + e*x^2]/(Sqrt[d]*Sqrt[1 - c^2*x^2])])/(2*c^2))/(4*c^2))/(10*e)
```

**Defintions of rubi rules used**

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 66

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]
```

rule 104 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 113 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/(d*f*(m + n + p + 1)), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegersQ[2*m, 2*n, 2*p]`

rule 171 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[h*(a + b*x)^(m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Simp[1/(d*f*(m + n + p + 2)) Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegersQ[2*m, 2*n, 2*p]`

rule 175 `Int[((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)))/((a_.) + (b_.)*(x_)), x_] := Simp[h/b Int[(c + d*x)^n*(e + f*x)^p, x], x] + Simp[(b*g - a*h)/b Int[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 354 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]`

rule 2036 `Int[(u_)*((c_) + (d_)*(x_)^(n_))^(q_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_), x_Symbol] := Int[u*(a1*a2 + b1*b2*x^n)^p*(c + d*x^n)^q, x] /; FreeQ[{a1, b1, a2, b2, c, d, n, p, q}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))`

rule 6853 `Int[((a_) + ArcSech[(c_)*(x_)])*(b_)]*(x_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcSech[c*x])/(2*e*(p + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(2*e*(p + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x^2)^(p + 1)/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[p, -1]`

### Maple [F]

$$\int x(x^2e + d)^{\frac{3}{2}}(a + b \operatorname{arcsech}(cx)) dx$$

input `int(x*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x)`

output `int(x*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x)`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 400 vs.  $2(181) = 362$ .

Time = 0.60 (sec) , antiderivative size = 1667, normalized size of antiderivative = 5.61

$$\int x(d + ex^2)^{3/2}(a + b \operatorname{sech}^{-1}(cx)) dx = \text{Too large to display}$$

input `integrate(x*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `[1/160*(8*b*c^5*d^(5/2)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 + 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) - (15*b*c^4*d^2 + 10*b*c^2*d*e + 3*b*e^2)*sqrt(-e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e + 8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^4*e*x^3 + (c^4*d - c^2*e)*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e^2) + 32*(b*c^5*e^2*x^4 + 2*b*c^5*d*e*x^2 + b*c^5*d^2)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 4*(8*a*c^5*e^2*x^4 + 16*a*c^5*d*e*x^2 + 8*a*c^5*d^2 - (2*b*c^4*e^2*x^3 + 3*(3*b*c^4*d*e + b*c^2*e^2)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d)/(c^5*e), 1/80*(4*b*c^5*d^(5/2)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 + 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) - (15*b*c^4*d^2 + 10*b*c^2*d*e + 3*b*e^2)*sqrt(e)*arctan(1/2*(2*c^2*e*x^3 + (c^2*d - e)*x)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2))/(c^2*e^2*x^4 + (c^2*d*e - e^2)*x^2 - d*e)) + 16*(b*c^5*e^2*x^4 + 2*b*c^5*d*e*x^2 + b*c^5*d^2)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 2*(8*a*c^5*e^2*x^4 + 16*a*c^5*d*e*x^2 + 8*a*c^5*d^2 - (2*b*c^4*e^2*x^3 + 3*(3*b*c^4*d*e + b*c^2*e^2)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d)/(c^5*e), -1/160*(16*b*c^5*sqrt(-d)*d^2*arctan(-1/2*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(-d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/...`

## Sympy [F]

$$\int x(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int x(a + b\operatorname{asech}(cx)) (d + ex^2)^{\frac{3}{2}} dx$$

input `integrate(x*(e*x**2+d)**(3/2)*(a+b*asech(c*x)),x)`

output `Integral(x*(a + b*asech(c*x))*(d + e*x**2)**(3/2), x)`

**Maxima [F]**

$$\int x(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^{\frac{3}{2}} (b \operatorname{arsech}(cx) + a)x dx$$

input `integrate(x*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `1/5*(e*x^2 + d)^(5/2)*a/e + 1/15*b*(((3*e^2*x^4 + d*e*x^2 - 2*d^2)*x^3 + 5*(d*e*x^4 + d^2*x^2)*x)*sqrt(e*x^2 + d)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)/(e*x^3) - 15*integrate(1/15*(15*(c^2*e^2*x^4*log(c) - e^2*x^2*log(c))*x^3 + 15*(c^2*d*e*x^4*log(c) - d*e*x^2*log(c))*x + ((3*(5*e^2*log(c) + e^2)*c^2*x^4 - 2*c^2*d^2 + (c^2*d*e - 15*e^2*log(c))*x^2)*x^3 + 5*((3*d*e*log(c) + d*e)*c^2*x^4 + (c^2*d^2 - 3*d*e*log(c))*x^2)*x + 30*((c^2*e^2*x^4 - e^2*x^2)*x^3 + (c^2*d*e*x^4 - d*e*x^2)*x)*log(sqrt(x)))*e^(1/2*log(c*x + 1) + 1/2*log(-c*x + 1)) + 30*((c^2*e^2*x^4 - e^2*x^2)*x^3 + (c^2*d*e*x^4 - d*e*x^2)*x)*log(sqrt(x))*sqrt(e*x^2 + d)/(c^2*e*x^4 - e*x^2 + (c^2*e*x^4 - e*x^2)*e^(1/2*log(c*x + 1) + 1/2*log(-c*x + 1))), x)`

**Giac [F]**

$$\int x(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^{\frac{3}{2}} (b \operatorname{arsech}(cx) + a)x dx$$

input `integrate(x*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((e*x^2 + d)^(3/2)*(b*arcsech(c*x) + a)*x, x)`



**Mupad [F(-1)]**

Timed out.

$$\int x(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int x (ex^2 + d)^{3/2} \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int(x*(d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))),x)`

output `int(x*(d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))), x)`

**Reduce [F]**

$$\int x(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int x (ex^2 + d)^{\frac{3}{2}} (a \operatorname{sech}(cx) b + a) dx$$

input `int(x*(e*x^2+d)^(3/2)*(a+b*asech(c*x)),x)`

output `int(x*(e*x^2+d)^(3/2)*(a+b*asech(c*x)),x)`

$$3.141 \quad \int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x} dx$$

Optimal result	1153
Mathematica [N/A]	1153
Rubi [N/A]	1154
Maple [N/A]	1154
Fricas [N/A]	1155
Sympy [N/A]	1155
Maxima [F(-2)]	1155
Giac [N/A]	1156
Mupad [N/A]	1156
Reduce [N/A]	1157

### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x} dx = \operatorname{Int}\left(\frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x}, x\right)$$

output `Defer(Int)((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x,x)`

### Mathematica [N/A]

Not integrable

Time = 6.96 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x} dx$$

input `Integrate[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x,x]`

output `Integrate[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x, x]`

**Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x} dx$$

↓ 6865

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x} dx$$

input

```
Int[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x,x]
```

output

```
$Aborted
```

**Maple [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{(x^2 e + d)^{3/2} (a + b \operatorname{arcsech}(cx))}{x} dx$$

input

```
int((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x,x)
```

output

```
int((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x,x)
```

**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.74

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(ex^2 + d)^{\frac{3}{2}} (b \operatorname{arsech}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x,x, algorithm="fricas")`

output `integral((a*e*x^2 + a*d + (b*e*x^2 + b*d)*arcsech(c*x))*sqrt(e*x^2 + d)/x, x)`

**Sympy [N/A]**

Not integrable

Time = 30.75 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(a + b \operatorname{asech}(cx)) (d + ex^2)^{\frac{3}{2}}}{x} dx$$

input `integrate((e*x**2+d)**(3/2)*(a+b*asech(c*x))/x,x)`

output `Integral((a + b*asech(c*x))*(d + e*x**2)**(3/2)/x, x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x} dx = \text{Exception raised: ValueError}$$

input `integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x,x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.15 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(ex^2 + d)^{3/2} (b \operatorname{arsech}(cx) + a)}{x} dx$$

input

```
integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x,x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)^(3/2)*(b*arcsech(c*x) + a)/x, x)
```

**Mupad [N/A]**

Not integrable

Time = 4.12 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(ex^2 + d)^{3/2} (a + b \operatorname{acosh}(\frac{1}{cx}))}{x} dx$$

input

```
int(((d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))))/x,x)
```

output

```
int(((d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))))/x, x)
```

**Reduce [N/A]**

Not integrable

Time = 200.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x} dx = \int \frac{(ex^2 + d)^{\frac{3}{2}} (a \operatorname{sech}(cx) b + a)}{x} dx$$

input `int((e*x^2+d)^(3/2)*(a+b*asech(c*x))/x,x)`output `int((e*x^2+d)^(3/2)*(a+b*asech(c*x))/x,x)`

**3.142** 
$$\int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^3} dx$$

Optimal result	1158
Mathematica [N/A]	1158
Rubi [N/A]	1159
Maple [N/A]	1159
Fricas [N/A]	1160
Sympy [N/A]	1160
Maxima [F(-2)]	1160
Giac [N/A]	1161
Mupad [N/A]	1161
Reduce [N/A]	1162

**Optimal result**

Integrand size = 23, antiderivative size = 23

$$\int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^3} dx = \operatorname{Int}\left(\frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^3}, x\right)$$

output

```
Defer(Int)((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^3,x)
```

**Mathematica [N/A]**

Not integrable

Time = 5.84 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^3} dx$$

input

```
Integrate[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x^3,x]
```

output

```
Integrate[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x^3, x]
```

**Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx$$

↓ 6865

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx$$

input

```
Int[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x^3,x]
```

output

```
$Aborted
```

**Maple [N/A]**

Not integrable

Time = 0.48 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{(x^2e + d)^{\frac{3}{2}} (a + b \operatorname{arcsech}(cx))}{x^3} dx$$

input

```
int((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^3,x)
```

output

```
int((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^3,x)
```



**Fricas [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.74

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(ex^2 + d)^{3/2} (b \operatorname{arsech}(cx) + a)}{x^3} dx$$

input `integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^3,x, algorithm="fricas")`

output `integral((a*e*x^2 + a*d + (b*e*x^2 + b*d)*arcsech(c*x))*sqrt(e*x^2 + d)/x^3, x)`

**Sympy [N/A]**

Not integrable

Time = 22.93 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(a + b \operatorname{asech}(cx)) (d + ex^2)^{3/2}}{x^3} dx$$

input `integrate((e*x**2+d)**(3/2)*(a+b*asech(c*x))/x**3,x)`

output `Integral((a + b*asech(c*x))*(d + e*x**2)**(3/2)/x**3, x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx = \text{Exception raised: ValueError}$$

input `integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^3,x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(ex^2 + d)^{3/2} (b \operatorname{arsech}(cx) + a)}{x^3} dx$$

input

```
integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^3,x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)^(3/2)*(b*arcsech(c*x) + a)/x^3, x)
```

**Mupad [N/A]**

Not integrable

Time = 4.39 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(ex^2 + d)^{3/2} (a + b \operatorname{acosh}(\frac{1}{cx}))}{x^3} dx$$

input

```
int(((d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))))/x^3,x)
```

output

```
int(((d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))))/x^3, x)
```

**Reduce [N/A]**

Not integrable

Time = 200.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^3} dx = \int \frac{(ex^2 + d)^{\frac{3}{2}} (a \operatorname{sech}(cx) b + a)}{x^3} dx$$

input `int((e*x^2+d)^(3/2)*(a+b*asech(c*x))/x^3,x)`output `int((e*x^2+d)^(3/2)*(a+b*asech(c*x))/x^3,x)`

### 3.143 $\int x^2(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	1163
Mathematica [N/A]	1163
Rubi [N/A]	1164
Maple [N/A]	1164
Fricas [N/A]	1165
Sympy [N/A]	1165
Maxima [F(-2)]	1165
Giac [N/A]	1166
Mupad [N/A]	1166
Reduce [N/A]	1167

#### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int x^2(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \operatorname{Int}\left(x^2(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)), x\right)$$

output

```
Defer(Int)(x^2*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x)
```

#### Mathematica [N/A]

Not integrable

Time = 13.37 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int x^2(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int x^2(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx$$

input

```
Integrate[x^2*(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]),x]
```

output

```
Integrate[x^2*(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]), x]
```

**Rubi [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx$$

↓ 6865

$$\int x^2 (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx$$

input `Int[x^2*(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int x^2 (x^2 e + d)^{\frac{3}{2}} (a + b \operatorname{arcsech}(cx)) dx$$

input `int(x^2*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x)`

output `int(x^2*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x)`

**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.87

$$\int x^2(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^{\frac{3}{2}}(b \operatorname{ar} \operatorname{sech}(cx) + a)x^2 dx$$

input `integrate(x^2*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral((a*e*x^4 + a*d*x^2 + (b*e*x^4 + b*d*x^2)*arcsech(c*x))*sqrt(e*x^2 + d), x)`

**Sympy [N/A]**

Not integrable

Time = 79.06 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int x^2(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int x^2(a + b \operatorname{asech}(cx)) (d + ex^2)^{\frac{3}{2}} dx$$

input `integrate(x**2*(e*x**2+d)**(3/2)*(a+b*asech(c*x)),x)`

output `Integral(x**2*(a + b*asech(c*x))*(d + e*x**2)**(3/2), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int x^2(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \text{Exception raised: ValueError}$$

input `integrate(x^2*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.14 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int x^2(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^{\frac{3}{2}} (b \operatorname{ar}\operatorname{sech}(cx) + a)x^2 dx$$

input

```
integrate(x^2*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)^(3/2)*(b*arcsech(c*x) + a)*x^2, x)
```

**Mupad [N/A]**

Not integrable

Time = 4.33 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int x^2(d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int x^2 (ex^2 + d)^{3/2} \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input

```
int(x^2*(d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))),x)
```

output

```
int(x^2*(d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))), x)
```

**Reduce [N/A]**

Not integrable

Time = 200.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int x^2 (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int x^2 (ex^2 + d)^{\frac{3}{2}} (a \operatorname{sech}(cx) b + a) dx$$

input `int(x^2*(e*x^2+d)^(3/2)*(a+b*asech(c*x)),x)`output `int(x^2*(e*x^2+d)^(3/2)*(a+b*asech(c*x)),x)`



### 3.144 $\int (d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	1168
Mathematica [N/A]	1168
Rubi [N/A]	1169
Maple [N/A]	1169
Fricas [N/A]	1170
Sympy [N/A]	1170
Maxima [F(-2)]	1170
Giac [N/A]	1171
Mupad [N/A]	1171
Reduce [N/A]	1172

#### Optimal result

Integrand size = 20, antiderivative size = 20

$$\int (d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \operatorname{Int}\left((d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)), x\right)$$

output

```
Defer(Int)((e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x)
```

#### Mathematica [N/A]

Not integrable

Time = 6.00 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int (d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx$$

input

```
Integrate[(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]),x]
```

output

```
Integrate[(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]), x]
```

**Rubi [N/A]**

Not integrable

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx$$

↓ 6865

$$\int (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx$$

input `Int[(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.17 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int (x^2 e + d)^{\frac{3}{2}} (a + b \operatorname{arcsech}(cx)) dx$$

input `int((e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x)`

output `int((e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x)`

**Fricas [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.85

$$\int (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^{\frac{3}{2}} (b \operatorname{ar} \operatorname{sech}(cx) + a) dx$$

input `integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral((a*e*x^2 + a*d + (b*e*x^2 + b*d)*arcsech(c*x))*sqrt(e*x^2 + d), x)`

**Sympy [N/A]**

Not integrable

Time = 23.49 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int (a + b \operatorname{asech}(cx)) (d + ex^2)^{\frac{3}{2}} dx$$

input `integrate((e*x**2+d)**(3/2)*(a+b*asech(c*x)),x)`

output `Integral((a + b*asech(c*x))*(d + e*x**2)**(3/2), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \text{Exception raised: ValueError}$$

input `integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^{\frac{3}{2}} (b \operatorname{ar} \operatorname{sech}(cx) + a) dx$$

input

```
integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)^(3/2)*(b*arcsech(c*x) + a), x)
```

**Mupad [N/A]**

Not integrable

Time = 4.15 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.20

$$\int (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^{3/2} \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input

```
int((d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))),x)
```

output

```
int((d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))), x)
```

**Reduce [N/A]**

Not integrable

Time = 200.20 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^{\frac{3}{2}} (a\operatorname{sech}(cx) b + a) dx$$

input `int((e*x^2+d)^(3/2)*(a+b*asech(c*x)),x)`output `int((e*x^2+d)^(3/2)*(a+b*asech(c*x)),x)`

$$3.145 \quad \int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^2} dx$$

Optimal result	1173
Mathematica [N/A]	1173
Rubi [N/A]	1174
Maple [N/A]	1174
Fricas [N/A]	1175
Sympy [N/A]	1175
Maxima [F(-2)]	1175
Giac [N/A]	1176
Mupad [N/A]	1176
Reduce [N/A]	1177

### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^2} dx = \operatorname{Int}\left(\frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^2}, x\right)$$

output `Defer(Int)((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^2,x)`

### Mathematica [N/A]

Not integrable

Time = 10.87 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^2} dx$$

input `Integrate[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x^2,x]`

output `Integrate[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x^2, x]`

**Rubi [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx$$

↓ 6865

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx$$

input

```
Int[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x^2,x]
```

output

```
$Aborted
```

**Maple [N/A]**

Not integrable

Time = 0.23 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{(x^2 e + d)^{3/2} (a + b \operatorname{arcsech}(cx))}{x^2} dx$$

input

```
int((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^2,x)
```

output

```
int((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^2,x)
```

**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.74

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{(ex^2 + d)^{\frac{3}{2}} (b \operatorname{arsech}(cx) + a)}{x^2} dx$$

input `integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^2,x, algorithm="fricas")`

output `integral((a*e*x^2 + a*d + (b*e*x^2 + b*d)*arcsech(c*x))*sqrt(e*x^2 + d)/x^2, x)`

**Sympy [N/A]**

Not integrable

Time = 20.38 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{(a + b \operatorname{asech}(cx)) (d + ex^2)^{\frac{3}{2}}}{x^2} dx$$

input `integrate((e*x**2+d)**(3/2)*(a+b*asech(c*x))/x**2,x)`

output `Integral((a + b*asech(c*x))*(d + e*x**2)**(3/2)/x**2, x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx = \text{Exception raised: ValueError}$$

input `integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^2,x, algorithm="maxima")`



output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{(ex^2 + d)^{3/2} (b \operatorname{arsech}(cx) + a)}{x^2} dx$$

input

```
integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^2,x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)^(3/2)*(b*arcsech(c*x) + a)/x^2, x)
```

**Mupad [N/A]**

Not integrable

Time = 4.50 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{(ex^2 + d)^{3/2} (a + b \operatorname{acosh}(\frac{1}{cx}))}{x^2} dx$$

input

```
int(((d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))))/x^2,x)
```

output

```
int(((d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))))/x^2, x)
```

**Reduce [N/A]**

Not integrable

Time = 200.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^2} dx = \int \frac{(ex^2 + d)^{\frac{3}{2}} (a \operatorname{sech}(cx) b + a)}{x^2} dx$$

input `int((e*x^2+d)^(3/2)*(a+b*asech(c*x))/x^2,x)`output `int((e*x^2+d)^(3/2)*(a+b*asech(c*x))/x^2,x)`

$$3.146 \quad \int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^4} dx$$

Optimal result	1178
Mathematica [N/A]	1178
Rubi [N/A]	1179
Maple [N/A]	1179
Fricas [N/A]	1180
Sympy [N/A]	1180
Maxima [F(-2)]	1180
Giac [N/A]	1181
Mupad [N/A]	1181
Reduce [N/A]	1182

### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^4} dx = \operatorname{Int}\left(\frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^4}, x\right)$$

output `Defer(Int)((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^4,x)`

### Mathematica [N/A]

Not integrable

Time = 12.66 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^4} dx$$

input `Integrate[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x^4,x]`

output `Integrate[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x^4, x]`

**Rubi [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx$$

↓ 6865

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx$$

input `Int[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x^4,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.41 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{(x^2e + d)^{\frac{3}{2}} (a + b \operatorname{arcsech}(cx))}{x^4} dx$$

input `int((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^4,x)`

output `int((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^4,x)`

**Fricas [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.74

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{(ex^2 + d)^{\frac{3}{2}} (b \operatorname{arsech}(cx) + a)}{x^4} dx$$

input `integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^4,x, algorithm="fricas")`

output `integral((a*e*x^2 + a*d + (b*e*x^2 + b*d)*arcsech(c*x))*sqrt(e*x^2 + d)/x^4, x)`

**Sympy [N/A]**

Not integrable

Time = 23.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{(a + b \operatorname{asech}(cx)) (d + ex^2)^{\frac{3}{2}}}{x^4} dx$$

input `integrate((e*x**2+d)**(3/2)*(a+b*asech(c*x))/x**4,x)`

output `Integral((a + b*asech(c*x))*(d + e*x**2)**(3/2)/x**4, x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx = \text{Exception raised: ValueError}$$

input `integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^4,x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{(ex^2 + d)^{3/2} (b \operatorname{arsech}(cx) + a)}{x^4} dx$$

input

```
integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^4,x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)^(3/2)*(b*arcsech(c*x) + a)/x^4, x)
```

**Mupad [N/A]**

Not integrable

Time = 4.24 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{(ex^2 + d)^{3/2} (a + b \operatorname{acosh}(\frac{1}{cx}))}{x^4} dx$$

input

```
int(((d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))))/x^4,x)
```

output

```
int(((d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))))/x^4, x)
```

**Reduce [N/A]**

Not integrable

Time = 200.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^4} dx = \int \frac{(ex^2 + d)^{\frac{3}{2}} (a \operatorname{sech}(cx) b + a)}{x^4} dx$$

input `int((e*x^2+d)^(3/2)*(a+b*asech(c*x))/x^4,x)`output `int((e*x^2+d)^(3/2)*(a+b*asech(c*x))/x^4,x)`

$$3.147 \quad \int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^6} dx$$

Optimal result . . . . .	1183
Mathematica [C] (verified) . . . . .	1184
Rubi [A] (verified) . . . . .	1185
Maple [F] . . . . .	1190
Fricas [A] (verification not implemented) . . . . .	1190
Sympy [F] . . . . .	1191
Maxima [F(-2)] . . . . .	1191
Giac [F] . . . . .	1191
Mupad [F(-1)] . . . . .	1192
Reduce [F] . . . . .	1192

**Optimal result**

Integrand size = 23, antiderivative size = 409

$$\int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^6} dx = \frac{4b(c^2d+2e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{75x^3}$$

$$+ \frac{b(8c^4d^2+23c^2de+23e^2)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{75dx}$$

$$+ \frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{25x^5} - \frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5}$$

$$+ \frac{bc(8c^4d^2+23c^2de+23e^2)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{d+ex^2}E(\arcsin(cx)|-\frac{e}{c^2d})}{75d\sqrt{1+\frac{ex^2}{d}}}$$

$$- \frac{b(c^2d+e)(8c^4d^2+19c^2de+15e^2)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1+\frac{ex^2}{d}}\operatorname{EllipticF}(\arcsin(cx),-\frac{e}{c^2d})}{75cd\sqrt{d+ex^2}}$$



output

```

4/75*b*(c^2*d+2*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*(e*x
^2+d)^(1/2)/x^3+1/75*b*(8*c^4*d^2+23*c^2*d*e+23*e^2)*(1/(c*x+1))^(1/2)*(c*
x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*(e*x^2+d)^(1/2)/d/x+1/25*b*(1/(c*x+1))^(1/2)
*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*(e*x^2+d)^(3/2)/x^5-1/5*(e*x^2+d)^(5/2)*
(a+b*arcsech(c*x))/d/x^5+1/75*b*c*(8*c^4*d^2+23*c^2*d*e+23*e^2)*(1/(c*x+1)
)^(1/2)*(c*x+1)^(1/2)*(e*x^2+d)^(1/2)*EllipticE(c*x,(-e/c^2/d)^(1/2))/d/(1
+e*x^2/d)^(1/2)-1/75*b*(c^2*d+e)*(8*c^4*d^2+19*c^2*d*e+15*e^2)*(1/(c*x+1))
^(1/2)*(c*x+1)^(1/2)*(1+e*x^2/d)^(1/2)*EllipticF(c*x,(-e/c^2/d)^(1/2))/c/d
/(e*x^2+d)^(1/2)

```

### Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 25.26 (sec) , antiderivative size = 620, normalized size of antiderivative = 1.52

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx = -\frac{15a(d+ex^2)^3}{x^5} + \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(d+ex^2)(23e^2x^4+dex^2(11+23c^2x^2)+d^2(3+4c^2x^2+8c^4x^4))}{x^5}$$

input

```
Integrate[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x^6,x]
```

output

```

((-15*a*(d + e*x^2)^3)/x^5 + (b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(d + e
*x^2)*(23*e^2*x^4 + d*e*x^2*(11 + 23*c^2*x^2) + d^2*(3 + 4*c^2*x^2 + 8*c^4
*x^4)))/x^5 - (15*b*(d + e*x^2)^3*ArcSech[c*x])/x^5 + (b*Sqrt[(1 - c*x)/(1
+ c*x)]*(-(c^2*(8*c^4*d^2 + 23*c^2*d*e + 23*e^2)*(d + e*x^2)) - (I*(c*Sqr
t[d] - I*Sqrt[e])^2*(1 + c*x)*Sqrt[(c*(Sqrt[d] - I*Sqrt[e]*x))/((c*Sqrt[d]
- I*Sqrt[e])*(1 + c*x))]*Sqrt[(c*(Sqrt[d] + I*Sqrt[e]*x))/((c*Sqrt[d] + I
*Sqrt[e])*(1 + c*x))]*((8*c^4*d^2 + 23*c^2*d*e + 23*e^2)*EllipticE[I*ArcSi
nh[Sqrt[((c^2*d + e)*(1 - c*x))/((c*Sqrt[d] + I*Sqrt[e])^2*(1 + c*x))]]], (
c*Sqrt[d] + I*Sqrt[e])^2/(c*Sqrt[d] - I*Sqrt[e])^2 + 2*Sqrt[e]*((8*I)*c^3
*d^(3/2) - 12*c^2*d*Sqrt[e] + (7*I)*c*Sqrt[d]*e - 15*e^(3/2))*EllipticF[I*
ArcSinh[Sqrt[((c^2*d + e)*(1 - c*x))/((c*Sqrt[d] + I*Sqrt[e])^2*(1 + c*x))
]], (c*Sqrt[d] + I*Sqrt[e])^2/(c*Sqrt[d] - I*Sqrt[e])^2)))/Sqrt[-((c*Sqrt
[d] - I*Sqrt[e])*(-1 + c*x))/((c*Sqrt[d] + I*Sqrt[e])*(1 + c*x)))]/c)/(7
5*d*Sqrt[d + e*x^2])

```

**Rubi [A] (verified)**

Time = 0.71 (sec) , antiderivative size = 338, normalized size of antiderivative = 0.83, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.522$ , Rules used = {6855, 27, 376, 442, 445, 25, 27, 399, 323, 321, 330, 327}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^6} dx \\
 & \quad \downarrow \text{6855} \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int -\frac{(ex^2+d)^{5/2}}{5dx^6\sqrt{1-c^2x^2}} dx - \frac{(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{5dx^5} \\
 & \quad \downarrow \text{27} \\
 & -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(ex^2+d)^{5/2}}{x^6\sqrt{1-c^2x^2}} dx}{5d} - \frac{(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{5dx^5} \\
 & \quad \downarrow \text{376} \\
 & -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{5} \int \frac{\sqrt{ex^2+d}(e(dc^2+5e)x^2+4d(dc^2+2e))}{x^4\sqrt{1-c^2x^2}} dx - \frac{d\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{5x^5} \right)}{5d} \\
 & \quad \downarrow \text{442} \\
 & -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{5} \left( \frac{1}{3} \int \frac{e(4d^2c^4+11dec^2+15e^2)x^2+d(8d^2c^4+23dec^2+23e^2)}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx - \frac{4d\sqrt{1-c^2x^2}(c^2d+2e)\sqrt{d+ex^2}}{3x^3} \right) - \frac{d\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{5x^5} \right)}{5d} \\
 & \quad \downarrow \text{445} \\
 & \frac{(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{5dx^5}
 \end{aligned}$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\left(\frac{1}{3}\left(-\frac{\int-\frac{de(4d^2c^4-(8d^2c^4+23dec^2+23e^2)x^2c^2+11dec^2+15e^2)}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx}{d}-\frac{\sqrt{1-c^2x^2}(8c^4d^2+23c^2de+23e^2)\sqrt{d+ex^2}}{x}\right)\right)\right)$$


---


$$\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5}$$

↓ 25

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\left(\frac{1}{3}\left(\frac{\int\frac{de(4d^2c^4-(8d^2c^4+23dec^2+23e^2)x^2c^2+11dec^2+15e^2)}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx}{d}-\frac{\sqrt{1-c^2x^2}(8c^4d^2+23c^2de+23e^2)\sqrt{d+ex^2}}{x}\right)\right)\right)$$


---


$$\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5}$$

↓ 27

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\left(\frac{1}{3}\left(e\int\frac{4d^2c^4-(8d^2c^4+23dec^2+23e^2)x^2c^2+11dec^2+15e^2}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx-\frac{\sqrt{1-c^2x^2}(8c^4d^2+23c^2de+23e^2)\sqrt{d+ex^2}}{x}\right)\right)\right)$$


---


$$\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5}$$

↓ 399

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\left(\frac{1}{3}\left(e\left(\frac{(c^2d+e)(8c^4d^2+19c^2de+15e^2)\int\frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx}{e}-\frac{c^2(8c^4d^2+23c^2de+23e^2)\int\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}dx}{e}\right)\right)\right)\right)$$


---


$$\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5}$$

↓ 323

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\left(\frac{1}{3}\left(e\left(\frac{(c^2d+e)(8c^4d^2+19c^2de+15e^2)\sqrt{\frac{ex^2}{d}+1}\int\frac{1}{\sqrt{1-c^2x^2}\sqrt{\frac{ex^2}{d}+1}}dx}{e\sqrt{d+ex^2}}-\frac{c^2(8c^4d^2+23c^2de+23e^2)\int\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}dx}{e}\right)\right)\right)\right)$$


---


$$\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5}$$

↓ 321

$$\begin{aligned}
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\left(\frac{1}{3}\left(e\left(\frac{(c^2d+e)(8c^4d^2+19c^2de+15e^2)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}}-\frac{c^2(8c^4d^2+23c^2de+23e^2)\int\frac{\sqrt{ex^2}}{\sqrt{1-}}}{e}\right.\right.\right. \\
 & \left.\left.\left.\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5}\right.\right.\right. \\
 & \left.\left.\left.\downarrow 330\right.\right.\right. \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\left(\frac{1}{3}\left(e\left(\frac{(c^2d+e)(8c^4d^2+19c^2de+15e^2)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}}-\frac{c^2(8c^4d^2+23c^2de+23e^2)\sqrt{d+ex^2}}{e\sqrt{\frac{ex^2}{d}+1}}\right.\right.\right. \\
 & \left.\left.\left.\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5}\right.\right.\right. \\
 & \left.\left.\left.\downarrow 327\right.\right.\right. \\
 & \left.\left.\left.\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5}\right.\right.\right. \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{5}\left(\frac{1}{3}\left(e\left(\frac{(c^2d+e)(8c^4d^2+19c^2de+15e^2)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}}-\frac{c(8c^4d^2+23c^2de+23e^2)\sqrt{d+ex^2}E}{e\sqrt{\frac{ex^2}{d}+1}}\right.\right.\right. \\
 & \left.\left.\left.\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5dx^5}\right.\right.\right. \\
 & \left.\left.\left.\downarrow 5d\right.\right.\right.
 \end{aligned}$$

input `Int[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x^6,x]`

output `-1/5*((d + e*x^2)^(5/2)*(a + b*ArcSech[c*x]))/(d*x^5) - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-1/5*(d*Sqrt[1 - c^2*x^2]*(d + e*x^2)^(3/2))/x^5 + ((-4*d*(c^2*d + 2*e)*Sqrt[1 - c^2*x^2]*Sqrt[d + e*x^2])/(3*x^3) + (-(((8*c^4*d^2 + 23*c^2*d*e + 23*e^2)*Sqrt[1 - c^2*x^2]*Sqrt[d + e*x^2])/x) + e*(-((c*(8*c^4*d^2 + 23*c^2*d*e + 23*e^2)*Sqrt[d + e*x^2]*EllipticE[ArcSin[c*x], -e/(c^2*d)]))/(e*Sqrt[1 + (e*x^2)/d])) + ((c^2*d + e)*(8*c^4*d^2 + 19*c^2*d*e + 15*e^2)*Sqrt[1 + (e*x^2)/d]*EllipticF[ArcSin[c*x], -e/(c^2*d)]))/(c*e*Sqrt[d + e*x^2]))/3)/5)/(5*d)`

## Definitions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27  $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 321  $\text{Int}[1/(\text{Sqrt}[(\text{a}_) + (\text{b}_.)*(x_)^2]*\text{Sqrt}[(\text{c}_) + (\text{d}_.)*(x_)^2]), \text{x\_Symbol}] \rightarrow \text{Simp}[(1/(\text{Sqrt}[\text{a}]*\text{Sqrt}[\text{c}]*\text{Rt}[-\text{d}/\text{c}, 2]))*\text{EllipticF}[\text{ArcSin}[\text{Rt}[-\text{d}/\text{c}, 2]*\text{x}], \text{b}*(\text{c}/(\text{a}*\text{d}))], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{NegQ}[\text{d}/\text{c}] \ \&\& \ \text{GtQ}[\text{c}, 0] \ \&\& \ \text{GtQ}[\text{a}, 0] \ \&\& \ \text{!(NegQ}[\text{b}/\text{a}] \ \&\& \ \text{SimplerSqrtQ}[-\text{b}/\text{a}, -\text{d}/\text{c}])$
- rule 323  $\text{Int}[1/(\text{Sqrt}[(\text{a}_) + (\text{b}_.)*(x_)^2]*\text{Sqrt}[(\text{c}_) + (\text{d}_.)*(x_)^2]), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Sqrt}[1 + (\text{d}/\text{c})*\text{x}^2]/\text{Sqrt}[\text{c} + \text{d}*\text{x}^2] \quad \text{Int}[1/(\text{Sqrt}[\text{a} + \text{b}*\text{x}^2]*\text{Sqrt}[1 + (\text{d}/\text{c})*\text{x}^2]), \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{!GtQ}[\text{c}, 0]$
- rule 327  $\text{Int}[\text{Sqrt}[(\text{a}_) + (\text{b}_.)*(x_)^2]/\text{Sqrt}[(\text{c}_) + (\text{d}_.)*(x_)^2], \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{a}]/(\text{Sqrt}[\text{c}]*\text{Rt}[-\text{d}/\text{c}, 2]))*\text{EllipticE}[\text{ArcSin}[\text{Rt}[-\text{d}/\text{c}, 2]*\text{x}], \text{b}*(\text{c}/(\text{a}*\text{d}))], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{NegQ}[\text{d}/\text{c}] \ \&\& \ \text{GtQ}[\text{c}, 0] \ \&\& \ \text{GtQ}[\text{a}, 0]$
- rule 330  $\text{Int}[\text{Sqrt}[(\text{a}_) + (\text{b}_.)*(x_)^2]/\text{Sqrt}[(\text{c}_) + (\text{d}_.)*(x_)^2], \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Sqrt}[\text{a} + \text{b}*\text{x}^2]/\text{Sqrt}[1 + (\text{b}/\text{a})*\text{x}^2] \quad \text{Int}[\text{Sqrt}[1 + (\text{b}/\text{a})*\text{x}^2]/\text{Sqrt}[\text{c} + \text{d}*\text{x}^2], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{NegQ}[\text{d}/\text{c}] \ \&\& \ \text{GtQ}[\text{c}, 0] \ \&\& \ \text{!GtQ}[\text{a}, 0]$
- rule 376  $\text{Int}[(\text{e}_.)*(x_)^{\text{m}_}*(\text{a}_) + (\text{b}_.)*(x_)^2)^{\text{p}_}*(\text{c}_) + (\text{d}_.)*(x_)^2)^{\text{q}_}, \text{x\_Symbol}] \rightarrow \text{Simp}[\text{c}*(\text{e}*\text{x})^{\text{m} + 1}*(\text{a} + \text{b}*\text{x}^2)^{\text{p} + 1}*(\text{c} + \text{d}*\text{x}^2)^{\text{q} - 1}/(\text{a}*\text{e}*(\text{m} + 1))), \text{x}] - \text{Simp}[1/(\text{a}*\text{e}^{2*(\text{m} + 1)}) \quad \text{Int}[(\text{e}*\text{x})^{\text{m} + 2}*(\text{a} + \text{b}*\text{x}^2)^{\text{p}*(\text{c} + \text{d}*\text{x}^2)^{\text{q} - 2}}*\text{Simp}[\text{c}*(\text{b}*\text{c} - \text{a}*\text{d})*(\text{m} + 1) + 2*\text{c}*(\text{b}*\text{c}*(\text{p} + 1) + \text{a}*\text{d}*(\text{q} - 1)) + \text{d}*((\text{b}*\text{c} - \text{a}*\text{d})*(\text{m} + 1) + 2*\text{b}*\text{c}*(\text{p} + \text{q}))*\text{x}^2, \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{p}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*\text{c} - \text{a}*\text{d}, 0] \ \&\& \ \text{GtQ}[\text{q}, 1] \ \&\& \ \text{LtQ}[\text{m}, -1] \ \&\& \ \text{IntBinomialQ}[\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{m}, 2, \text{p}, \text{q}, \text{x}]$

rule 399

```
Int[((e_) + (f_)*(x_)^2)/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] := Simp[f/b Int[Sqrt[a + b*x^2]/Sqrt[c + d*x^2], x], x] + Simp[(b*e - a*f)/b Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && !((PosQ[b/a] && PosQ[d/c]) || (NegQ[b/a] && (PosQ[d/c] || (GtQ[a, 0] && (!GtQ[c, 0] || SimplerSqrtQ[-b/a, -d/c])))))
```

rule 442

```
Int[((g_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2), x_Symbol] := Simp[e*(g*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^q/(a*g*(m + 1))), x] - Simp[1/(a*g^2*(m + 1)) Int[(g*x)^(m + 2)*(a + b*x^2)^p*(c + d*x^2)^(q - 1)*Simp[c*(b*e - a*f)*(m + 1) + e*2*(b*c*(p + 1) + a*d*q) + d*((b*e - a*f)*(m + 1) + b*e*2*(p + q + 1))*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && GtQ[q, 0] && LtQ[m, -1] && !(EqQ[q, 1] && SimplerQ[e + f*x^2, c + d*x^2])
```

rule 445

```
Int[((g_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2), x_Symbol] := Simp[e*(g*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(a*c*g*(m + 1))), x] + Simp[1/(a*c*g^2*(m + 1)) Int[(g*x)^(m + 2)*(a + b*x^2)^p*(c + d*x^2)^q*Simp[a*f*c*(m + 1) - e*(b*c + a*d)*(m + 2 + 1) - e*2*(b*c*p + a*d*q) - b*e*d*(m + 2*(p + q + 2) + 1)*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, p, q}, x] && LtQ[m, -1]
```

rule 6855

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

**Maple [F]**

$$\int \frac{(x^2e + d)^{\frac{3}{2}} (a + b \operatorname{arcsech}(cx))}{x^6} dx$$

input `int((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^6,x)`

output `int((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^6,x)`

**Fricas [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 338, normalized size of antiderivative = 0.83

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx =$$

$$15 (bcde^2x^4 + 2bcd^2ex^2 + bcd^3)\sqrt{ex^2 + d} \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}+1}{cx}\right) + \left(15acde^2x^4 + 30acd^2ex^2 + 15acd^3 - \dots\right)$$

input `integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^6,x, algorithm="fricas")`

output `-1/75*(15*(b*c*d*e^2*x^4 + 2*b*c*d^2*e*x^2 + b*c*d^3)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + (15*a*c*d*e^2*x^4 + 30*a*c*d^2*e*x^2 + 15*a*c*d^3 - (3*b*c^2*d^3*x + (8*b*c^6*d^3 + 23*b*c^4*d^2*e + 23*b*c^2*d*e^2)*x^5 + (4*b*c^4*d^3 + 11*b*c^2*d^2*e)*x^3)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d) - ((8*b*c^8*d^3 + 23*b*c^6*d^2*e + 23*b*c^4*d*e^2)*x^5*elliptic_e(arcsin(c*x), -e/(c^2*d)) - (8*b*c^8*d^3 + (23*b*c^6 + 4*b*c^4)*d^2*e + (23*b*c^4 + 11*b*c^2)*d*e^2 + 15*b*e^3)*x^5*elliptic_f(arcsin(c*x), -e/(c^2*d)))*sqrt(d))/(c*d^2*x^5)`

**Sympy [F]**

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx = \int \frac{(a + b \operatorname{asech}(cx)) (d + ex^2)^{3/2}}{x^6} dx$$

input `integrate((e*x**2+d)**(3/2)*(a+b*asech(c*x))/x**6,x)`

output `Integral((a + b*asech(c*x))*(d + e*x**2)**(3/2)/x**6, x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx = \text{Exception raised: ValueError}$$

input `integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^6,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx = \int \frac{(ex^2 + d)^{3/2} (b \operatorname{arsech}(cx) + a)}{x^6} dx$$

input `integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^6,x, algorithm="giac")`

output `integrate((e*x^2 + d)^(3/2)*(b*arcsech(c*x) + a)/x^6, x)`



**Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx = \int \frac{(ex^2 + d)^{3/2} (a + b \operatorname{acosh}(\frac{1}{cx}))}{x^6} dx$$

input `int(((d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))))/x^6,x)`output `int(((d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))))/x^6, x)`**Reduce [F]**

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^6} dx = \int \frac{(ex^2 + d)^{3/2} (a \operatorname{sech}(cx) b + a)}{x^6} dx$$

input `int((e*x^2+d)^(3/2)*(a+b*asech(c*x))/x^6,x)`output `int((e*x^2+d)^(3/2)*(a+b*asech(c*x))/x^6,x)`

**3.148**  $\int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^8} dx$

Optimal result	1193
Mathematica [C] (verified)	1194
Rubi [A] (verified)	1195
Maple [F]	1200
Fricas [A] (verification not implemented)	1200
Sympy [F(-1)]	1201
Maxima [F(-2)]	1201
Giac [F]	1202
Mupad [F(-1)]	1202
Reduce [F]	1202

**Optimal result**

Integrand size = 23, antiderivative size = 556

$$\int \frac{(d+ex^2)^{3/2} (a+b\operatorname{sech}^{-1}(cx))}{x^8} dx = \frac{b(120c^4d^2 + 159c^2de - 37e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} \sqrt{d+ex^2}}{3675dx^3}$$

$$+ \frac{b(240c^6d^3 + 528c^4d^2e + 193c^2de^2 - 247e^3) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} \sqrt{d+ex^2}}{3675d^2x}$$

$$+ \frac{b(30c^2d + 11e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} (d+ex^2)^{3/2}}{1225dx^5}$$

$$+ \frac{b\sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} (d+ex^2)^{5/2}}{49dx^7}$$

$$- \frac{(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{7dx^7} + \frac{2e(d+ex^2)^{5/2} (a+b\operatorname{sech}^{-1}(cx))}{35d^2x^5}$$

$$+ \frac{bc(240c^6d^3 + 528c^4d^2e + 193c^2de^2 - 247e^3) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{d+ex^2} E(\arcsin(cx) | -\frac{e}{c^2d})}{3675d^2 \sqrt{1+\frac{ex^2}{d}}}$$

$$- \frac{2b(c^2d + e) (120c^6d^3 + 204c^4d^2e + 17c^2de^2 - 105e^3) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1+\frac{ex^2}{d}} \operatorname{EllipticF}(\arcsin(cx), -\frac{e}{c^2d})}{3675cd^2 \sqrt{d+ex^2}}$$

output

```

1/3675*b*(120*c^4*d^2+159*c^2*d*e-37*e^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*
(-c^2*x^2+1)^(1/2)*(e*x^2+d)^(1/2)/d/x^3+1/3675*b*(240*c^6*d^3+528*c^4*d^2
*e+193*c^2*d*e^2-247*e^3)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)
*(e*x^2+d)^(1/2)/d^2/x+1/1225*b*(30*c^2*d+11*e)*(1/(c*x+1))^(1/2)*(c*x+1)
^(1/2)*(-c^2*x^2+1)^(1/2)*(e*x^2+d)^(3/2)/d/x^5+1/49*b*(1/(c*x+1))^(1/2)*
(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*(e*x^2+d)^(5/2)/d/x^7-1/7*(e*x^2+d)^(5/2)
*(a+b*arcsech(c*x))/d/x^7+2/35*e*(e*x^2+d)^(5/2)*(a+b*arcsech(c*x))/d^2/x^
5+1/3675*b*c*(240*c^6*d^3+528*c^4*d^2*e+193*c^2*d*e^2-247*e^3)*(1/(c*x+1))
^(1/2)*(c*x+1)^(1/2)*(e*x^2+d)^(1/2)*EllipticE(c*x,(-e/c^2/d)^(1/2))/d^2/(
1+e*x^2/d)^(1/2)-2/3675*b*(c^2*d+e)*(120*c^6*d^3+204*c^4*d^2*e+17*c^2*d*e^
2-105*e^3)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(1+e*x^2/d)^(1/2)*EllipticF(c*x
,(-e/c^2/d)^(1/2))/c/d^2/(e*x^2+d)^(1/2)

```

### Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 26.74 (sec) , antiderivative size = 728, normalized size of antiderivative = 1.31

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^8} dx = \frac{105a(d+ex^2)^3(-5d+2ex^2)}{x^7} + \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(d+ex^2)(-247e^3x^6+de^2x^4(71+193c^2x^2))}{x^7}$$

input

```
Integrate[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x^8,x]
```

output

```

((105*a*(d + e*x^2)^3*(-5*d + 2*e*x^2))/x^7 + (b*Sqrt[(1 - c*x)/(1 + c*x)]
*(1 + c*x)*(d + e*x^2)*(-247*e^3*x^6 + d*e^2*x^4*(71 + 193*c^2*x^2) + 3*d^
2*e*x^2*(61 + 83*c^2*x^2 + 176*c^4*x^4) + 15*d^3*(5 + 6*c^2*x^2 + 8*c^4*x^
4 + 16*c^6*x^6)))/x^7 + (105*b*(d + e*x^2)^3*(-5*d + 2*e*x^2)*ArcSech[c*x]
)/x^7 + (b*Sqrt[(1 - c*x)/(1 + c*x)]*(-(c^2*(240*c^6*d^3 + 528*c^4*d^2*e +
193*c^2*d*e^2 - 247*e^3)*(d + e*x^2)) - (I*(c*Sqrt[d] - I*Sqrt[e])^2*(1 +
c*x)*Sqrt[(c*(Sqrt[d] - I*Sqrt[e]*x))/((c*Sqrt[d] - I*Sqrt[e])*(1 + c*x))
]*Sqrt[(c*(Sqrt[d] + I*Sqrt[e]*x))/((c*Sqrt[d] + I*Sqrt[e])*(1 + c*x))])*((
240*c^6*d^3 + 528*c^4*d^2*e + 193*c^2*d*e^2 - 247*e^3)*EllipticE[I*ArcSinh
[Sqrt[((c^2*d + e)*(1 - c*x))/((c*Sqrt[d] + I*Sqrt[e])^2*(1 + c*x))]]], (c*
Sqrt[d] + I*Sqrt[e])^2/(c*Sqrt[d] - I*Sqrt[e])^2 + 2*Sqrt[e]*((240*I)*c^5
*d^(5/2) - 360*c^4*d^2*Sqrt[e] + (48*I)*c^3*d^(3/2)*e - 207*c^2*d*e^(3/2)
- (173*I)*c*Sqrt[d]*e^2 + 210*e^(5/2))*EllipticF[I*ArcSinh[Sqrt[((c^2*d +
e)*(1 - c*x))/((c*Sqrt[d] + I*Sqrt[e])^2*(1 + c*x))]]], (c*Sqrt[d] + I*Sqrt
[e])^2/(c*Sqrt[d] - I*Sqrt[e])^2))/Sqrt[-(((c*Sqrt[d] - I*Sqrt[e])*(-1 +
c*x))/((c*Sqrt[d] + I*Sqrt[e])*(1 + c*x)))]))/c/(3675*d^2*Sqrt[d + e*x^2]
)

```

### Rubi [A] (verified)

Time = 0.92 (sec) , antiderivative size = 462, normalized size of antiderivative = 0.83, number of steps used = 13, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.565$ , Rules used = {6855, 27, 442, 442, 442, 445, 25, 27, 399, 323, 321, 330, 327}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^8} dx$$

$$\downarrow 6855$$

$$b \sqrt{\frac{1}{cx + 1}} \sqrt{cx + 1} \int -\frac{(5d - 2ex^2) (ex^2 + d)^{5/2}}{35d^2 x^8 \sqrt{1 - c^2 x^2}} dx + \frac{2e(d + ex^2)^{5/2} (a + b \operatorname{sech}^{-1}(cx))}{35d^2 x^5} - \frac{(d + ex^2)^{5/2} (a + b \operatorname{sech}^{-1}(cx))}{7dx^7}$$

$$\downarrow 27$$

$$\begin{aligned}
 & - \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(5d-2ex^2)(ex^2+d)^{5/2}}{x^8\sqrt{1-c^2x^2}} dx}{35d^2} + \frac{2e(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{35d^2x^5} - \\
 & \qquad \qquad \qquad \frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{7dx^7} \\
 & \qquad \qquad \qquad \downarrow 442 \\
 & - \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{7} \int \frac{(ex^2+d)^{3/2}((5c^2d-14e)ex^2+d(30dc^2+11e))}{x^6\sqrt{1-c^2x^2}} dx - \frac{5d\sqrt{1-c^2x^2}(d+ex^2)^{5/2}}{7x^7} \right)}{35d^2} + \\
 & \qquad \qquad \qquad \frac{2e(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{35d^2x^5} - \frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{7dx^7} \\
 & \qquad \qquad \qquad \downarrow 442 \\
 & - \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{7} \left( \frac{1}{5} \int \frac{\sqrt{ex^2+d}(2e(15d^2c^4+18dec^2-35e^2)x^2+d(120d^2c^4+159dec^2-37e^2))}{x^4\sqrt{1-c^2x^2}} dx - \frac{d\sqrt{1-c^2x^2}(30c^2d+11e)(d+ex^2)}{5x^5} \right) \right)}{35d^2} \\
 & \qquad \qquad \qquad \frac{2e(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{35d^2x^5} - \frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{7dx^7} \\
 & \qquad \qquad \qquad \downarrow 442 \\
 & - \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{7} \left( \frac{1}{5} \left( \frac{1}{3} \int \frac{e(120d^3c^6+249d^2ec^4+71de^2c^2-210e^3)x^2+d(240d^3c^6+528d^2ec^4+193de^2c^2-247e^3)}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx - \frac{d\sqrt{1-c^2x^2}(120d^3c^6+249d^2ec^4+71de^2c^2-210e^3)}{3x^3} \right) \right) \right)}{35d^2} \\
 & \qquad \qquad \qquad \frac{2e(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{35d^2x^5} - \frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{7dx^7} \\
 & \qquad \qquad \qquad \downarrow 445 \\
 & - \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{1}{7} \left( \frac{1}{5} \left( \frac{1}{3} \left( - \int \frac{de(120d^3c^6+249d^2ec^4+71de^2c^2-(240d^3c^6+528d^2ec^4+193de^2c^2-247e^3)x^2c^2-210e^3)}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx - \frac{\sqrt{1-c^2x^2}(120d^3c^6+249d^2ec^4+71de^2c^2-210e^3)}{3d} \right) \right) \right) \right)}{35d^2} \\
 & \qquad \qquad \qquad \frac{2e(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{35d^2x^5} - \frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{7dx^7} \\
 & \qquad \qquad \qquad \downarrow 25
 \end{aligned}$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{1}{3}\left(\int\frac{de(120d^3c^6+249d^2ec^4+71de^2c^2-(240d^3c^6+528d^2ec^4+193de^2c^2-247e^3)x^2c^2-210e^3)}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx-\frac{\sqrt{1-c^2x^2}(240c^6d^3+528c^4d^2e+193c^2de^2-247e^3)}{e}\right)\right)\right)\right)$$

$$\frac{2e(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{35d^2x^5}-\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{7dx^7}$$

↓ 27

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{1}{3}\left(e\int\frac{120d^3c^6+249d^2ec^4+71de^2c^2-(240d^3c^6+528d^2ec^4+193de^2c^2-247e^3)x^2c^2-210e^3}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx-\frac{\sqrt{1-c^2x^2}(240c^6d^3+528c^4d^2e+193c^2de^2-247e^3)}{e}\right)\right)\right)\right)$$

$$\frac{2e(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{35d^2x^5}-\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{7dx^7}$$

↓ 399

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{1}{3}\left(e\left(\frac{2(c^2d+e)(120c^6d^3+204c^4d^2e+17c^2de^2-105e^3)}{e}\int\frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx-\frac{c^2(240c^6d^3+528c^4d^2e+193c^2de^2-247e^3)}{e}\right)\right)\right)\right)$$

$$\frac{2e(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{35d^2x^5}-\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{7dx^7}$$

↓ 323

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{1}{3}\left(e\left(\frac{2(c^2d+e)(120c^6d^3+204c^4d^2e+17c^2de^2-105e^3)\sqrt{\frac{ex^2}{d}+1}}{e\sqrt{d+ex^2}}\int\frac{1}{\sqrt{1-c^2x^2}\sqrt{\frac{ex^2}{d}+1}}dx-\frac{c^2(240c^6d^3+528c^4d^2e+193c^2de^2-247e^3)}{e}\right)\right)\right)\right)$$

$$\frac{2e(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{35d^2x^5}-\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{7dx^7}$$

↓ 321

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{1}{3}\left(e\left(\frac{2(c^2d+e)(120c^6d^3+204c^4d^2e+17c^2de^2-105e^3)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}}-\frac{c^2(240c^6d^3+528c^4d^2e+193c^2de^2-247e^3)}{e}\right)\right)\right)\right)$$

$$\frac{2e(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{35d^2x^5}-\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{7dx^7}$$

↓ 330

$$\begin{aligned}
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{1}{3}\left(e\left(\frac{2(c^2d+e)(120c^6d^3+204c^4d^2e+17c^2de^2-105e^3)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}}-\frac{c^2(240c^6d^3+528c^4d^2e+193c^2de^2-247e^3)\sqrt{d+ex^2}}{x^5}+\frac{(-1/3*(d*(120*c^4*d^2+159*c^2*d*e-37*e^2)*\sqrt{1-c^2*x^2}*\sqrt{d+ex^2})}{x^3}+(-(((240*c^6*d^3+528*c^4*d^2*e+193*c^2*d*e^2-247*e^3)*\sqrt{1-c^2*x^2}*\sqrt{d+ex^2})/x)+e*(-((c*(240*c^6*d^3+528*c^4*d^2*e+193*c^2*d*e^2-247*e^3)*\sqrt{d+ex^2}*\operatorname{EllipticE}[\operatorname{ArcSin}[c*x],-(e/(c^2*d))])/(e*\sqrt{1+(e*x^2)/d}))+2*(c^2*d+e)*(120*c^6*d^3+204*c^4*d^2*e+17*c^2*d*e^2-105*e^3)*\sqrt{1+(e*x^2)/d}*\operatorname{EllipticF}[\operatorname{ArcSin}[c*x],-(e/(c^2*d))])/(c*e*\sqrt{d+ex^2}))/3)/5)/7)\right)/35d^2x^5-\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{7dx^7}\right) \\
 & \quad \quad \quad \downarrow \text{327} \\
 & \frac{2e(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{35d^2x^5}-\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{7dx^7} \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{1}{3}\left(e\left(\frac{2(c^2d+e)(120c^6d^3+204c^4d^2e+17c^2de^2-105e^3)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}}-\frac{c^2(240c^6d^3+528c^4d^2e+193c^2de^2-247e^3)\sqrt{d+ex^2}}{x^5}+\frac{(-1/3*(d*(120*c^4*d^2+159*c^2*d*e-37*e^2)*\sqrt{1-c^2*x^2}*\sqrt{d+ex^2})}{x^3}+(-(((240*c^6*d^3+528*c^4*d^2*e+193*c^2*d*e^2-247*e^3)*\sqrt{1-c^2*x^2}*\sqrt{d+ex^2})/x)+e*(-((c*(240*c^6*d^3+528*c^4*d^2*e+193*c^2*d*e^2-247*e^3)*\sqrt{d+ex^2}*\operatorname{EllipticE}[\operatorname{ArcSin}[c*x],-(e/(c^2*d))])/(e*\sqrt{1+(e*x^2)/d}))+2*(c^2*d+e)*(120*c^6*d^3+204*c^4*d^2*e+17*c^2*d*e^2-105*e^3)*\sqrt{1+(e*x^2)/d}*\operatorname{EllipticF}[\operatorname{ArcSin}[c*x],-(e/(c^2*d))])/(c*e*\sqrt{d+ex^2}))/3)/5)/7)\right)/35d^2x^5-\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{7dx^7}\right)
 \end{aligned}$$

```
input Int[((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/x^8,x]
```

```
output -1/7*((d + e*x^2)^(5/2)*(a + b*ArcSech[c*x]))/(d*x^7) + (2*e*(d + e*x^2)^(5/2)*(a + b*ArcSech[c*x]))/(35*d^2*x^5) - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-5*d*Sqrt[1 - c^2*x^2]*(d + e*x^2)^(5/2))/(7*x^7) + (-1/5*(d*(30*c^2*d + 11*e)*Sqrt[1 - c^2*x^2]*(d + e*x^2)^(3/2))/x^5 + (-1/3*(d*(120*c^4*d^2 + 159*c^2*d*e - 37*e^2)*Sqrt[1 - c^2*x^2]*Sqrt[d + e*x^2])/x^3 + (-(((240*c^6*d^3 + 528*c^4*d^2*e + 193*c^2*d*e^2 - 247*e^3)*Sqrt[1 - c^2*x^2]*Sqrt[d + e*x^2])/x) + e*(-((c*(240*c^6*d^3 + 528*c^4*d^2*e + 193*c^2*d*e^2 - 247*e^3)*Sqrt[d + e*x^2]*EllipticE[ArcSin[c*x], -(e/(c^2*d))])/(e*Sqrt[1 + (e*x^2)/d])) + (2*(c^2*d + e)*(120*c^6*d^3 + 204*c^4*d^2*e + 17*c^2*d*e^2 - 105*e^3)*Sqrt[1 + (e*x^2)/d]*EllipticF[ArcSin[c*x], -(e/(c^2*d))])/(c*e*Sqrt[d + e*x^2])))/3)/5)/7))/(35*d^2)
```

**Defintions of rubi rules used**

```
rule 25 Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

rule 321 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := S  
imp[(1/(Sqrt[a]*Sqrt[c]*Rt[-d/c, 2]))*EllipticF[ArcSin[Rt[-d/c, 2]*x], b*(c  
/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a,  
0] && !(NegQ[b/a] && SimplerSqrtQ[-b/a, -d/c])`

rule 323 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := S  
imp[Sqrt[1 + (d/c)*x^2]/Sqrt[c + d*x^2] Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (  
d/c)*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]`

rule 327 `Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[  
(Sqrt[a]/(Sqrt[c]*Rt[-d/c, 2]))*EllipticE[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d)  
)], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]`

rule 330 `Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[  
Sqrt[a + b*x^2]/Sqrt[1 + (b/a)*x^2] Int[Sqrt[1 + (b/a)*x^2]/Sqrt[c + d*x^  
2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a,  
0]`

rule 399 `Int[((e_) + (f_.)*(x_)^2)/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)  
^2]), x_Symbol] := Simp[f/b Int[Sqrt[a + b*x^2]/Sqrt[c + d*x^2], x], x] +  
Simp[(b*e - a*f)/b Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; Fr  
eeQ[{a, b, c, d, e, f}, x] && !((PosQ[b/a] && PosQ[d/c]) || (NegQ[b/a] &&  
(PosQ[d/c] || (GtQ[a, 0] && (!GtQ[c, 0] || SimplerSqrtQ[-b/a, -d/c]))))`

rule 442 `Int[((g_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_  
.)*((e_) + (f_.)*(x_)^2), x_Symbol] := Simp[e*(g*x)^(m + 1)*(a + b*x^2)^(p  
+ 1)*((c + d*x^2)^q/(a*g*(m + 1))), x] - Simp[1/(a*g^2*(m + 1)) Int[(g*x)  
^(m + 2)*(a + b*x^2)^p*(c + d*x^2)^(q - 1)*Simp[c*(b*e - a*f)*(m + 1) + e*2  
*(b*c*(p + 1) + a*d*q) + d*((b*e - a*f)*(m + 1) + b*e*2*(p + q + 1))*x^2, x  
], x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && GtQ[q, 0] && LtQ[m, -1]  
&& !(EqQ[q, 1] && SimplerQ[e + f*x^2, c + d*x^2])`



rule 445

```
Int[((g._)*(x_))^(m_)*((a_) + (b._)*(x_)^2)^(p_)*((c_) + (d._)*(x_)^2)^(q_
.)*(e_) + (f._)*(x_)^2), x_Symbol] := Simp[e*(g*x)^(m + 1)*(a + b*x^2)^(p
+ 1)*((c + d*x^2)^(q + 1)/(a*c*g*(m + 1))), x] + Simp[1/(a*c*g^2*(m + 1))
Int[(g*x)^(m + 2)*(a + b*x^2)^p*(c + d*x^2)^q*Simp[a*f*c*(m + 1) - e*(b*c
+ a*d)*(m + 2 + 1) - e*2*(b*c*p + a*d*q) - b*e*d*(m + 2*(p + q + 2) + 1)*x^
2, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, p, q}, x] && LtQ[m, -1]
```

rule 6855

```
Int[((a_) + ArcSech[(c._)*(x_)]*(b._))*((f._)*(x_))^(m_)*((d_) + (e._)*(
x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Si
mp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)]
Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x]] /; Fre
eQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] &&
GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2
*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

**Maple [F]**

$$\int \frac{(x^2e + d)^{\frac{3}{2}} (a + b \operatorname{arcsech}(cx))}{x^8} dx$$

input

```
int((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^8,x)
```

output

```
int((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^8,x)
```

**Fricas [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 447, normalized size of antiderivative = 0.80

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^8} dx = \frac{105 (2bcde^3x^6 - bcd^2e^2x^4 - 8bcd^3ex^2 - 5bcd^4)\sqrt{ex^2 + d} \log\left(\frac{cx\sqrt{ex^2 + d}}{d + ex^2}\right) + \dots}{x^8}$$

input

```
integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^8,x, algorithm="fricas")
```

output

```
1/3675*(105*(2*b*c*d*e^3*x^6 - b*c*d^2*e^2*x^4 - 8*b*c*d^3*e*x^2 - 5*b*c*d^4)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + (210*a*c*d*e^3*x^6 - 105*a*c*d^2*e^2*x^4 - 840*a*c*d^3*e*x^2 - 525*a*c*d^4 + (75*b*c^2*d^4*x + (240*b*c^8*d^4 + 528*b*c^6*d^3*e + 193*b*c^4*d^2*e^2 - 247*b*c^2*d*e^3)*x^7 + (120*b*c^6*d^4 + 249*b*c^4*d^3*e + 71*b*c^2*d^2*e^2)*x^5 + 3*(30*b*c^4*d^4 + 61*b*c^2*d^3*e)*x^3)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d) + ((240*b*c^10*d^4 + 528*b*c^8*d^3*e + 193*b*c^6*d^2*e^2 - 247*b*c^4*d*e^3)*x^7*elliptic_e(arcsin(c*x), -e/(c^2*d)) - (240*b*c^10*d^4 + 24*(22*b*c^8 + 5*b*c^6)*d^3*e + (193*b*c^6 + 249*b*c^4)*d^2*e^2 - (247*b*c^4 - 71*b*c^2)*d*e^3 - 210*b*e^4)*x^7*elliptic_f(arcsin(c*x), -e/(c^2*d)))*sqrt(d))/(c*d^3*x^7)
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^8} dx = \text{Timed out}$$

input

```
integrate((e*x**2+d)**(3/2)*(a+b*asech(c*x))/x**8,x)
```

output

Timed out

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^8} dx = \text{Exception raised: ValueError}$$

input

```
integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^8,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e
```

**Giac [F]**

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^8} dx = \int \frac{(ex^2 + d)^{3/2} (b \operatorname{arsech}(cx) + a)}{x^8} dx$$

input `integrate((e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/x^8,x, algorithm="giac")`

output `integrate((e*x^2 + d)^(3/2)*(b*arcsech(c*x) + a)/x^8, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^8} dx = \int \frac{(ex^2 + d)^{3/2} (a + b \operatorname{acosh}(\frac{1}{cx}))}{x^8} dx$$

input `int(((d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))))/x^8,x)`

output `int(((d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))))/x^8, x)`

**Reduce [F]**

$$\int \frac{(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{x^8} dx = \int \frac{(ex^2 + d)^{3/2} (a \operatorname{sech}(cx) b + a)}{x^8} dx$$

input `int((e*x^2+d)^(3/2)*(a+b*asech(c*x))/x^8,x)`

output `int((e*x^2+d)^(3/2)*(a+b*asech(c*x))/x^8,x)`

**3.149**  $\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx$

Optimal result	1203
Mathematica [A] (verified)	1204
Rubi [A] (verified)	1205
Maple [F]	1210
Fricas [A] (verification not implemented)	1210
Sympy [F]	1211
Maxima [F(-2)]	1212
Giac [F]	1212
Mupad [F(-1)]	1212
Reduce [F]	1213

**Optimal result**

Integrand size = 23, antiderivative size = 356

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx$$

$$= \frac{b(19c^2d - 9e) \sqrt{\frac{1}{1+cx}} \sqrt{1 + cx} \sqrt{1 - c^2x^2} \sqrt{d + ex^2}}{120c^4e^2}$$

$$- \frac{b \sqrt{\frac{1}{1+cx}} \sqrt{1 + cx} \sqrt{1 - c^2x^2} (d + ex^2)^{3/2}}{20c^2e^2} + \frac{d^2 \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx))}{e^3}$$

$$- \frac{2d(d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{3e^3} + \frac{(d + ex^2)^{5/2} (a + b \operatorname{sech}^{-1}(cx))}{5e^3}$$

$$- \frac{b(45c^4d^2 - 10c^2de + 9e^2) \sqrt{\frac{1}{1+cx}} \sqrt{1 + cx} \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{120c^5e^{5/2}}$$

$$- \frac{8bd^{5/2} \sqrt{\frac{1}{1+cx}} \sqrt{1 + cx} \operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{15e^3}$$

output

$$\begin{aligned} & 1/120*b*(19*c^2*d-9*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)* \\ & (e*x^2+d)^(1/2)/c^4/e^2-1/20*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1) \\ & )^(1/2)*(e*x^2+d)^(3/2)/c^2/e^2+d^2*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/e^3 \\ & -2/3*d*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/e^3+1/5*(e*x^2+d)^(5/2)*(a+b*arc \\ & sech(c*x))/e^3-1/120*b*(45*c^4*d^2-10*c^2*d*e+9*e^2)*(1/(c*x+1))^(1/2)*(c* \\ & x+1)^(1/2)*arctan(e^(1/2)*(-c^2*x^2+1)^(1/2)/c/(e*x^2+d)^(1/2))/c^5/e^(5/2) \\ & )-8/15*b*d^(5/2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh((e*x^2+d)^(1/2)/d \\ & ^{(1/2)/(-c^2*x^2+1)^(1/2))/e^3 \end{aligned}$$

**Mathematica [A] (verified)**

Time = 22.20 (sec) , antiderivative size = 366, normalized size of antiderivative = 1.03

$$\begin{aligned} & \int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx \\ & = \frac{\sqrt{d + ex^2} \left( 8ac^4(8d^2 - 4dex^2 + 3e^2x^4) - be\sqrt{\frac{1-cx}{1+cx}}(1 + cx)(9e + c^2(-13d + 6ex^2)) + 8bc^4(8d^2 - 4dex^2 + 3e^2x^4) \right)}{120c^4e^3} \\ & + \frac{b\sqrt{\frac{1-cx}{1+cx}}\sqrt{1 - c^2x^2} \left( \sqrt{-c^2}\sqrt{-c^2d - e}\sqrt{e}(45c^4d^2 - 10c^2de + 9e^2) \sqrt{\frac{c^2(d+ex^2)}{c^2d+e}} \arcsin\left(\frac{c\sqrt{e}\sqrt{1-c^2x^2}}{\sqrt{-c^2}\sqrt{-c^2d-e}}\right) + 6 \right)}{120c^7e^3(-1 + cx)\sqrt{d + ex^2}} \end{aligned}$$

input

Integrate[(x^5\*(a + b\*ArcSech[c\*x]))/Sqrt[d + e\*x^2],x]

output

$$\begin{aligned} & (\operatorname{Sqrt}[d + e*x^2]*(8*a*c^4*(8*d^2 - 4*d*e*x^2 + 3*e^2*x^4) - b*e*\operatorname{Sqrt}[(1 - \\ & c*x)/(1 + c*x)]*(1 + c*x)*(9*e + c^2*(-13*d + 6*e*x^2)) + 8*b*c^4*(8*d^2 - \\ & 4*d*e*x^2 + 3*e^2*x^4)*\operatorname{ArcSech}[c*x])/(120*c^4*e^3) + (b*\operatorname{Sqrt}[(1 - c*x)/( \\ & 1 + c*x)]*\operatorname{Sqrt}[1 - c^2*x^2]*(\operatorname{Sqrt}[-c^2]*\operatorname{Sqrt}[-(c^2*d) - e]*\operatorname{Sqrt}[e]*(45*c^4 \\ & *d^2 - 10*c^2*d*e + 9*e^2)*\operatorname{Sqrt}[(c^2*(d + e*x^2))/(c^2*d + e)]*\operatorname{ArcSin}[(c*\operatorname{S} \\ & \operatorname{qrt}[e]*\operatorname{Sqrt}[1 - c^2*x^2])/(\operatorname{Sqrt}[-c^2]*\operatorname{Sqrt}[-(c^2*d) - e])] + 64*c^7*d^(5/2) \\ & )*\operatorname{Sqrt}[-d - e*x^2]*\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[1 - c^2*x^2])/(\operatorname{Sqrt}[-d - e*x^2])]) / \\ & (120*c^7*e^3*(-1 + c*x)*\operatorname{Sqrt}[d + e*x^2]) \end{aligned}$$

**Rubi [A] (verified)**

Time = 1.36 (sec) , antiderivative size = 306, normalized size of antiderivative = 0.86, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.522$ , Rules used = {6855, 27, 7282, 2118, 27, 171, 27, 175, 66, 104, 218, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx \\
 & \quad \downarrow \text{6855} \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{\sqrt{ex^2+d}(3e^2x^4 - 4dex^2 + 8d^2)}{15e^3x\sqrt{1-c^2x^2}} dx + \frac{d^2\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{e^3} + \\
 & \quad \frac{(d+ex^2)^{5/2}(a + b\operatorname{sech}^{-1}(cx))}{5e^3} - \frac{2d(d+ex^2)^{3/2}(a + b\operatorname{sech}^{-1}(cx))}{3e^3} \\
 & \quad \downarrow \text{27} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{\sqrt{ex^2+d}(3e^2x^4 - 4dex^2 + 8d^2)}{x\sqrt{1-c^2x^2}} dx}{15e^3} + \frac{d^2\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{e^3} + \\
 & \quad \frac{(d+ex^2)^{5/2}(a + b\operatorname{sech}^{-1}(cx))}{5e^3} - \frac{2d(d+ex^2)^{3/2}(a + b\operatorname{sech}^{-1}(cx))}{3e^3} \\
 & \quad \downarrow \text{7282} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{\sqrt{ex^2+d}(3e^2x^4 - 4dex^2 + 8d^2)}{x^2\sqrt{1-c^2x^2}} dx^2}{30e^3} + \frac{d^2\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{e^3} + \\
 & \quad \frac{(d+ex^2)^{5/2}(a + b\operatorname{sech}^{-1}(cx))}{5e^3} - \frac{2d(d+ex^2)^{3/2}(a + b\operatorname{sech}^{-1}(cx))}{3e^3} \\
 & \quad \downarrow \text{2118} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{\int -\frac{e\sqrt{ex^2+d}(32c^2d^2 - (19c^2d - 9e)ex^2)}{2x^2\sqrt{1-c^2x^2}} dx^2}{2c^2e} - \frac{3e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2} \right)}{30e^3} + \\
 & \quad \frac{d^2\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{e^3} + \frac{(d+ex^2)^{5/2}(a + b\operatorname{sech}^{-1}(cx))}{5e^3} - \\
 & \quad \frac{2d(d+ex^2)^{3/2}(a + b\operatorname{sech}^{-1}(cx))}{3e^3} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{\int\frac{\sqrt{ex^2+d}(32c^2d^2-(19c^2d-9e)ex^2)}{x^2\sqrt{1-c^2x^2}}dx^2}{4c^2}-\frac{3e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2}\right)$$


---


$$\frac{30e^3}{e^3}\frac{d^2\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3}+\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3}-\frac{2d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3}$$

↓ 171

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{e\sqrt{1-c^2x^2}(19c^2d-9e)\sqrt{d+ex^2}}{c^2}-\frac{\int-\frac{64d^3c^4+e(45d^2c^4-10dec^2+9e^2)x^2}{2x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2}{4c^2}-\frac{3e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2}\right)$$


---


$$\frac{30e^3}{e^3}\frac{d^2\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3}+\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3}-\frac{2d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3}$$

↓ 27

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{\int\frac{64d^3c^4+e(45d^2c^4-10dec^2+9e^2)x^2}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2}{4c^2}+\frac{e\sqrt{1-c^2x^2}(19c^2d-9e)\sqrt{d+ex^2}}{c^2}-\frac{3e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2}\right)$$


---


$$\frac{30e^3}{e^3}\frac{d^2\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3}+\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3}-\frac{2d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3}$$

↓ 175

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{64c^4d^3\int\frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2+e(45c^4d^2-10c^2de+9e^2)\int\frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2}{2c^2}+\frac{e\sqrt{1-c^2x^2}(19c^2d-9e)\sqrt{d+ex^2}}{c^2}-\frac{3e\sqrt{1-c^2x^2}(d+ex^2)^{3/2}}{2c^2}\right)$$


---


$$\frac{30e^3}{e^3}\frac{d^2\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3}+\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3}-\frac{2d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3}$$

↓ 66

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{64c^4d^3\int\frac{1}{x^2\sqrt{1-c^2x^2}}dx^2+2e(45c^4d^2-10c^2de+9e^2)\int\frac{1}{-ex^4-c^2}\frac{d\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}}}{2c^2}+\frac{e\sqrt{1-c^2x^2}(19c^2d-9e)\sqrt{d+ex^2}}{c^2}\right)-\frac{3e\sqrt{1-c^2x^2}}{2}$$

---


$$\frac{d^2\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3}+\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3}-\frac{2d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3}$$

↓ 104

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{128c^4d^3\int\frac{1}{x^4-d}\frac{d\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}+2e(45c^4d^2-10c^2de+9e^2)\int\frac{1}{-ex^4-c^2}\frac{d\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}}}{2c^2}+\frac{e\sqrt{1-c^2x^2}(19c^2d-9e)\sqrt{d+ex^2}}{c^2}\right)-\frac{3e\sqrt{1-c^2x^2}}{2}$$

---


$$\frac{d^2\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3}+\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3}-\frac{2d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3}$$

↓ 218

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{128c^4d^3\int\frac{1}{x^4-d}\frac{d\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}-\frac{2\sqrt{e}(45c^4d^2-10c^2de+9e^2)\arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{2c^2}}{4c^2}+\frac{e\sqrt{1-c^2x^2}(19c^2d-9e)\sqrt{d+ex^2}}{c^2}\right)-\frac{3e\sqrt{1-c^2x^2}}{2}$$

---


$$\frac{d^2\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3}+\frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3}-\frac{2d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3}$$

↓ 220



$$\frac{d^2\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3} + \frac{(d+ex^2)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{5e^3} - \frac{2d(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} + b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\frac{2\sqrt{e}(45c^4d^2-10c^2de+9e^2)\arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c} - 128c^4d^{5/2}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{2c^2} + \frac{e\sqrt{1-c^2x^2}(19c^2d-9e)\sqrt{d+ex^2}}{4c^2} \right) \frac{1}{30e^3}$$

input `Int[(x^5*(a + b*ArcSech[c*x]))/Sqrt[d + e*x^2],x]`

output `(d^2*Sqrt[d + e*x^2]*(a + b*ArcSech[c*x])/e^3 - (2*d*(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/(3*e^3) + ((d + e*x^2)^(5/2)*(a + b*ArcSech[c*x]))/(5*e^3) + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-3*e*Sqrt[1 - c^2*x^2]*(d + e*x^2)^(3/2))/(2*c^2) + (((19*c^2*d - 9*e)*e*Sqrt[1 - c^2*x^2]*Sqrt[d + e*x^2])/c^2 + ((-2*Sqrt[e]*(45*c^4*d^2 - 10*c^2*d*e + 9*e^2)*ArcTan[(Sqrt[e]*Sqrt[1 - c^2*x^2])/(c*Sqrt[d + e*x^2])])/c - 128*c^4*d^(5/2)*ArcTanh[Sqrt[d + e*x^2]/(Sqrt[d]*Sqrt[1 - c^2*x^2])])/(2*c^2))/(4*c^2)))/(30*e^3)`

**Defintions of rubi rules used**

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 104 `Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 171 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[h*(a + b*x)^m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Simp[1/(d*f*(m + n + p + 2)) Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegersQ[2*m, 2*n, 2*p]`

rule 175 `Int[(((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)))/((a_.) + (b_.)*(x_)), x_] := Simp[h/b Int[(c + d*x)^n*(e + f*x)^p, x], x] + Simp[(b*g - a*h)/b Int[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 2118 `Int[(Px_)*((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := With[{q = Expon[Px, x], k = Coeff[Px, x, Expon[Px, x]]}, Simp[k*(a + b*x)^(m + q - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*b^(q - 1)*(m + n + p + q + 1))), x] + Simp[1/(d*f*b^q*(m + n + p + q + 1)) Int[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p*ExpandToSum[d*f*b^q*(m + n + p + q + 1)*Px - d*f*k*(m + n + p + q + 1)*(a + b*x)^q + k*(a + b*x)^(q - 2)*(a^2*d*f*(m + n + p + q + 1) - b*(b*c*e*(m + q - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*(m + q) + n + p) - b*(d*e*(m + q + n) + c*f*(m + q + p)))]*x, x], x] /; NeQ[m + n + p + q + 1, 0] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x]`

rule 6855

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))*((f_.)*(x_.))^(m_.)*((d_.) + (e_.)*(
x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Si
mp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)]
Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x]] /; Fre
eQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] &&
GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2
*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

rule 7282

```
Int[(u_)/(x_), x_Symbol] := With[{lst = PowerVariableExpn[u, 0, x]}, Simp[1
/lst[[2]] Subst[Int[NormalizeIntegrand[Simplify[lst[[1]]/x], x], x], x, (
lst[[3]]*x)^lst[[2]]], x] /; !FalseQ[lst] && NeQ[lst[[2]], 0] /; NonsumQ[
u] && !RationalFunctionQ[u, x]
```

**Maple [F]**

$$\int \frac{x^5(a + b \operatorname{arcsech}(cx))}{\sqrt{x^2e + d}} dx$$

input

```
int(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x)
```

output

```
int(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x)
```

**Fricas [A] (verification not implemented)**

Time = 0.66 (sec) , antiderivative size = 1679, normalized size of antiderivative = 4.72

$$\int \frac{x^5(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \text{Too large to display}$$

input

```
integrate(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="fricas")
```

output

```
[1/480*(64*b*c^5*d^(5/2)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 + 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) - (45*b*c^4*d^2 - 10*b*c^2*d*e + 9*b*e^2)*sqrt(-e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e + 8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^4*e*x^3 + (c^4*d - c^2*e)*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e^2) + 32*(3*b*c^5*e^2*x^4 - 4*b*c^5*d*e*x^2 + 8*b*c^5*d^2)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 4*(24*a*c^5*e^2*x^4 - 32*a*c^5*d*e*x^2 + 64*a*c^5*d^2 - (6*b*c^4*e^2*x^3 - (13*b*c^4*d*e - 9*b*c^2*e^2)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d))/(c^5*e^3), 1/240*(32*b*c^5*d^(5/2)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 + 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) - (45*b*c^4*d^2 - 10*b*c^2*d*e + 9*b*e^2)*sqrt(e)*arctan(1/2*(2*c^2*e*x^3 + (c^2*d - e)*x)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*e^2*x^4 + (c^2*d*e - e^2)*x^2 - d*e) + 16*(3*b*c^5*e^2*x^4 - 4*b*c^5*d*e*x^2 + 8*b*c^5*d^2)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 2*(24*a*c^5*e^2*x^4 - 32*a*c^5*d*e*x^2 + 64*a*c^5*d^2 - (6*b*c^4*e^2*x^3 - (13*b*c^4*d*e - 9*b*c^2*e^2)*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d))/(c^5*e^3), -1/480*(128*b*c^5*sqrt(-d)*d^2*arctan(-1/2*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(-d)*sqrt(-(c^...
```

## Sympy [F]

$$\int \frac{x^5(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{x^5(a + b \operatorname{asech}(cx))}{\sqrt{d + ex^2}} dx$$

input

```
integrate(x**5*(a+b*asech(c*x))/sqrt(d + e*x**2), x)
```

output

```
Integral(x**5*(a + b*asech(c*x))/sqrt(d + e*x**2), x)
```

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \text{Exception raised: ValueError}$$

input `integrate(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^5}{\sqrt{ex^2 + d}} dx$$

input `integrate(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^5/sqrt(e*x^2 + d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{x^5 (a + b \operatorname{acosh}(\frac{1}{cx}))}{\sqrt{ex^2 + d}} dx$$

input `int((x^5*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(1/2),x)`

output `int((x^5*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(1/2), x)`

**Reduce [F]**

$$\int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{x^5(\operatorname{asech}(cx)b + a)}{\sqrt{ex^2 + d}} dx$$

input `int(x^5*(a+b*asech(c*x))/(e*x^2+d)^(1/2),x)`

output `int(x^5*(a+b*asech(c*x))/(e*x^2+d)^(1/2),x)`

$$3.150 \quad \int \frac{x^3 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx$$

Optimal result	1214
Mathematica [A] (verified)	1215
Rubi [A] (verified)	1215
Maple [F]	1219
Fricas [B] (verification not implemented)	1219
Sympy [F]	1220
Maxima [F(-2)]	1221
Giac [F]	1221
Mupad [F(-1)]	1221
Reduce [F]	1222

### Optimal result

Integrand size = 23, antiderivative size = 251

$$\int \frac{x^3 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = -\frac{b \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} \sqrt{d+ex^2}}{6c^2e} - \frac{d \sqrt{d+ex^2} (a + b \operatorname{sech}^{-1}(cx))}{e^2} + \frac{(d+ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{3e^2} + \frac{b(3c^2d - e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{6c^3e^{3/2}} + \frac{2bd^{3/2} \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{3e^2}$$

output

```
-1/6*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*(e*x^2+d)^(1/2)/
c^2/e-d*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/e^2+1/3*(e*x^2+d)^(3/2)*(a+b*ar
csech(c*x))/e^2+1/6*b*(3*c^2*d-e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctan(e
^(1/2)*(-c^2*x^2+1)^(1/2)/c/(e*x^2+d)^(1/2))/c^3/e^(3/2)+2/3*b*d^(3/2)*(1/
(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh((e*x^2+d)^(1/2)/d^(1/2)/(-c^2*x^2+1)^(
1/2))/e^2
```

**Mathematica [A] (verified)**

Time = 21.82 (sec) , antiderivative size = 406, normalized size of antiderivative = 1.62

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx$$

$$= -\frac{\sqrt{d + ex^2} \left( be \sqrt{\frac{1-cx}{1+cx}} (1 + cx) + 2ac^2(2d - ex^2) + 2bc^2(2d - ex^2) \operatorname{sech}^{-1}(cx) \right)}{6c^2e^2}$$

$$- \frac{b \sqrt{\frac{1-cx}{1+cx}} \sqrt{1 - c^2x^2} \left( -3(-c^2)^{3/2} d \sqrt{-c^2d - e} \sqrt{e} \sqrt{\frac{c^2(d+ex^2)}{c^2d+e}} \arcsin \left( \frac{c \sqrt{e} \sqrt{1-c^2x^2}}{\sqrt{-c^2} \sqrt{-c^2d - e}} \right) + \sqrt{-c^2} \sqrt{-c^2d - e} \right)}{6c^5e^2(-1 + cx) \sqrt{d + ex^2}}$$

input `Integrate[(x^3*(a + b*ArcSech[c*x]))/Sqrt[d + e*x^2],x]`

output `-1/6*(Sqrt[d + e*x^2]*(b*e*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x) + 2*a*c^2*(2*d - e*x^2) + 2*b*c^2*(2*d - e*x^2)*ArcSech[c*x]))/(c^2*e^2) - (b*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*(-3*(-c^2)^(3/2)*d*Sqrt[-(c^2*d) - e]*Sqrt[e]*Sqrt[(c^2*(d + e*x^2))/(c^2*d + e)]*ArcSin[(c*Sqrt[e]*Sqrt[1 - c^2*x^2])/(Sqrt[-c^2]*Sqrt[-(c^2*d) - e]]) + Sqrt[-c^2]*Sqrt[-(c^2*d) - e]*e^(3/2)*Sqrt[(c^2*(d + e*x^2))/(c^2*d + e)]*ArcSin[(Sqrt[-c^2]*Sqrt[e]*Sqrt[1 - c^2*x^2])/(c*Sqrt[-(c^2*d) - e]]) + 4*c^5*d^(3/2)*Sqrt[-d - e*x^2]*ArcTan[(Sqrt[d]*Sqrt[1 - c^2*x^2])/Sqrt[-d - e*x^2]]))/(6*c^5*e^2*(-1 + c*x)*Sqrt[d + e*x^2])`

**Rubi [A] (verified)**

Time = 0.51 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.86, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.435$ , Rules used = {6855, 27, 435, 171, 27, 175, 66, 104, 218, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx$$

↓ 6855



$$\begin{aligned}
& b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int -\frac{(2d-ex^2)\sqrt{ex^2+d}}{3e^2x\sqrt{1-c^2x^2}}dx + \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2} - \\
& \quad \frac{d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^2} \\
& \quad \downarrow 27 \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(2d-ex^2)\sqrt{ex^2+d}}{x\sqrt{1-c^2x^2}}dx}{3e^2} + \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2} - \\
& \quad \frac{d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^2} \\
& \quad \downarrow 435 \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(2d-ex^2)\sqrt{ex^2+d}}{x^2\sqrt{1-c^2x^2}}dx^2}{6e^2} + \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2} - \\
& \quad \frac{d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^2} \\
& \quad \downarrow 171 \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} - \frac{\int -\frac{4c^2d^2+(3c^2d-e)ex^2}{2x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2}{c^2} \right)}{6e^2} + \\
& \quad \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2} - \frac{d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^2} \\
& \quad \downarrow 27 \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{4c^2d^2+(3c^2d-e)ex^2}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2}{2c^2} + \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} \right)}{6e^2} + \\
& \quad \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2} - \frac{d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^2} \\
& \quad \downarrow 175 \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{4c^2d^2 \int \frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2 + e(3c^2d-e) \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2}{2c^2} + \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} \right)}{6e^2} + \\
& \quad \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2} - \frac{d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^2} \\
& \quad \downarrow 66
\end{aligned}$$

$$\begin{aligned}
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{4c^2d^2\int\frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2+2e(3c^2d-e)\int\frac{1}{-ex^4-c^2}d\frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}}+\frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2}\right)}{6e^2} \\
 & \quad - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2} - \frac{d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^2} \\
 & \qquad \qquad \qquad \downarrow 104 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{8c^2d^2\int\frac{1}{x^4-d}d\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}+2e(3c^2d-e)\int\frac{1}{-ex^4-c^2}d\frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}}+\frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2}\right)}{6e^2} \\
 & \quad - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2} - \frac{d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^2} \\
 & \qquad \qquad \qquad \downarrow 218 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{8c^2d^2\int\frac{1}{x^4-d}d\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}-\frac{2\sqrt{e}(3c^2d-e)\arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{2c^2}+\frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2}\right)}{6e^2} \\
 & \quad - \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2} - \frac{d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^2} \\
 & \qquad \qquad \qquad \downarrow 220 \\
 & \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^2} - \frac{d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^2} - \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{-\frac{2\sqrt{e}(3c^2d-e)\arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c}-8c^2d^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{2c^2}+\frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2}\right)}{6e^2}
 \end{aligned}$$

input `Int[(x^3*(a + b*ArcSech[c*x]))/Sqrt[d + e*x^2],x]`

output `-((d*Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/e^2) + ((d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]))/(3*e^2) - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((e*Sqrt[1 - c^2*x^2]*Sqrt[d + e*x^2])/c^2 + ((-2*(3*c^2*d - e)*Sqrt[e]*ArcTan[(Sqrt[e]*Sqrt[1 - c^2*x^2])/(c*Sqrt[d + e*x^2])])/c - 8*c^2*d^(3/2)*ArcTanh[Sqrt[d + e*x^2]/(Sqrt[d]*Sqrt[1 - c^2*x^2])])/(2*c^2)))/(6*e^2)`

## Definitions of rubi rules used

- rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 66  $\text{Int}[1/(\text{Sqrt}[(a_*) + (b_*)(x_)]*\text{Sqrt}[(c_*) + (d_*)(x_)]), x\_Symbol] \rightarrow \text{Simp}[2 \text{ Subst}[\text{Int}[1/(b - d*x^2), x], x, \text{Sqrt}[a + b*x]/\text{Sqrt}[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ !\text{GtQ}[c - a*(d/b), 0]$
- rule 104  $\text{Int}[(((a_*) + (b_*)(x_))^{(m_*)}*((c_*) + (d_*)(x_))^{(n_*)})/((e_*) + (f_*)(x_))), x_] \rightarrow \text{With}[\{q = \text{Denominator}[m]\}, \text{Simp}[q \text{ Subst}[\text{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{EqQ}[m + n + 1, 0] \ \&\& \ \text{RationalQ}[n] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{SimplerQ}[a + b*x, c + d*x]$
- rule 171  $\text{Int}[(((a_*) + (b_*)(x_))^{(m_*)}*((c_*) + (d_*)(x_))^{(n_*)}*((e_*) + (f_*)(x_))^{(p_*)}*((g_*) + (h_*)(x_))), x_] \rightarrow \text{Simp}[h*(a + b*x)^m*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)}/(d*f*(m+n+p+2))), x] + \text{Simp}[1/(d*f*(m+n+p+2)) \text{ Int}[(a + b*x)^{(m-1)}*(c + d*x)^n*(e + f*x)^p*\text{Simp}[a*d*f*g*(m+n+p+2) - h*(b*c*e*m + a*(d*e*(n+1) + c*f*(p+1))) + (b*d*f*g*(m+n+p+2) + h*(a*d*f*m - b*(d*e*(m+n+1) + c*f*(m+p+1)))*x, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, h, n, p\}, x] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{NeQ}[m+n+p+2, 0] \ \&\& \ \text{IntegersQ}[2*m, 2*n, 2*p]$
- rule 175  $\text{Int}[(((c_*) + (d_*)(x_))^{(n_*)}*((e_*) + (f_*)(x_))^{(p_*)}*((g_*) + (h_*)(x_))))/((a_*) + (b_*)(x_)), x_] \rightarrow \text{Simp}[h/b \text{ Int}[(c + d*x)^n*(e + f*x)^p, x], x] + \text{Simp}[(b*g - a*h)/b \text{ Int}[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, h, n, p\}, x]$
- rule 218  $\text{Int}[((a_*) + (b_*)(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

rule 220 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 435 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2)^(r_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2) * (a + b*x)^p * (c + d*x)^q * (e + f*x)^r, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, f, p, q, r}, x] && IntegerQ[(m - 1)/2]`

rule 6855 `Int[((a_) + ArcSech[(c_)*(x_)]*(b_))*((f_)*(x_)^(m_))*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))`

## Maple [F]

$$\int \frac{x^3(a + b \operatorname{arcsech}(cx))}{\sqrt{x^2e + d}} dx$$

input `int(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x)`

output `int(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x)`

## Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 330 vs. 2(159) = 318.

Time = 0.33 (sec) , antiderivative size = 1389, normalized size of antiderivative = 5.53

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \text{Too large to display}$$

input `integrate(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="fricas")`

output `[1/24*(4*b*c^3*d^(3/2)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 - 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) + (3*b*c^2*d - b*e)*sqrt(-e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e + 8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^4*e*x^3 + (c^4*d - c^2*e)*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e^2) + 8*(b*c^3*e*x^2 - 2*b*c^3*d)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 4*(2*a*c^3*e*x^2 - b*c^2*e*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 4*a*c^3*d)*sqrt(e*x^2 + d))/(c^3*e^2), 1/12*(2*b*c^3*d^(3/2)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 - 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) + (3*b*c^2*d - b*e)*sqrt(e)*arctan(1/2*(2*c^2*e*x^3 + (c^2*d - e)*x)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*e^2*x^4 + (c^2*d*e - e^2)*x^2 - d*e) + 4*(b*c^3*e*x^2 - 2*b*c^3*d)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 2*(2*a*c^3*e*x^2 - b*c^2*e*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 4*a*c^3*d)*sqrt(e*x^2 + d))/(c^3*e^2), 1/24*(8*b*c^3*sqrt(-d)*d*arctan(-1/2*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(-d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*d*e*x^4 + (c^2*d^2 - d*e)*x^2 - d^2)) + (3*b*c^2*d - b*e)*sqrt(-e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e + 8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^4*e*x^3 + (c^4*d - c^2*e)*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(-(c^2*...`

## Sympy [F]

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{x^3(a + b \operatorname{asech}(cx))}{\sqrt{d + ex^2}} dx$$

input `integrate(x**3*(a+b*asech(c*x))/sqrt(d + e*x**2),x)`

output `Integral(x**3*(a + b*asech(c*x))/sqrt(d + e*x**2), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \text{Exception raised: ValueError}$$

input `integrate(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^3}{\sqrt{ex^2 + d}} dx$$

input `integrate(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^3/sqrt(e*x^2 + d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(\frac{1}{cx}))}{\sqrt{ex^2 + d}} dx$$

input `int((x^3*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(1/2),x)`

output `int((x^3*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(1/2), x)`

**Reduce [F]**

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{x^3(\operatorname{asech}(cx)b + a)}{\sqrt{ex^2 + d}} dx$$

input `int(x^3*(a+b*asech(c*x))/(e*x^2+d)^(1/2),x)`

output `int(x^3*(a+b*asech(c*x))/(e*x^2+d)^(1/2),x)`

**3.151** 
$$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{\sqrt{d+ex^2}} dx$$

Optimal result	1223
Mathematica [A] (verified)	1224
Rubi [A] (verified)	1224
Maple [F]	1227
Fricas [B] (verification not implemented)	1228
Sympy [F]	1229
Maxima [F]	1229
Giac [F]	1229
Mupad [F(-1)]	1230
Reduce [F]	1230

**Optimal result**

Integrand size = 21, antiderivative size = 153

$$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{\sqrt{d+ex^2}} dx = \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e} - \frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c\sqrt{e}} - \frac{b\sqrt{d}\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{e}$$

output

```
(e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/e-b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arc
tan(e^(1/2)*(-c^2*x^2+1)^(1/2)/c/(e*x^2+d)^(1/2))/c/e^(1/2)-b*d^(1/2)*(1/(
c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh((e*x^2+d)^(1/2)/d^(1/2)/(-c^2*x^2+1)^(
1/2))/e
```



**Mathematica [A] (verified)**

Time = 18.93 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.56

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \frac{\sqrt{d + ex^2}(a + b \operatorname{sech}^{-1}(cx))}{e} + \frac{b\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\left(\sqrt{-c^2}\sqrt{-c^2d-e}\sqrt{e}\sqrt{\frac{c^2(d+ex^2)}{c^2d+e}}\arcsin\left(\frac{c\sqrt{e}\sqrt{1-c^2x^2}}{\sqrt{-c^2}\sqrt{-c^2d-e}}\right) + c^3\sqrt{d}\sqrt{-d-ex^2}\arctan\left(\frac{\sqrt{d}\sqrt{-d-ex^2}}{c^3e(-1+cx)\sqrt{d+ex^2}}\right)\right)}{c^3e(-1+cx)\sqrt{d+ex^2}}$$

input `Integrate[(x*(a + b*ArcSech[c*x]))/Sqrt[d + e*x^2],x]`

output  $(\sqrt{d + ex^2}(a + b \operatorname{ArcSech}[cx]))/e + (b \sqrt{(1 - cx)/(1 + cx)} \sqrt{1 - c^2x^2} (\sqrt{-c^2} \sqrt{-c^2d - e} \sqrt{e} \sqrt{(c^2(d + ex^2)/(c^2d + e))} \operatorname{ArcSin}[(c \sqrt{e} \sqrt{1 - c^2x^2})/(\sqrt{-c^2} \sqrt{-c^2d - e})]) + c^3 \sqrt{d} \sqrt{-d - ex^2} \operatorname{ArcTan}[(\sqrt{d} \sqrt{1 - c^2x^2})/(\sqrt{-d - ex^2})])/(c^3 e (-1 + cx) \sqrt{d + ex^2}))$

**Rubi [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.90, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {6853, 2036, 354, 140, 27, 66, 104, 218, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx$$

↓ 6853

$$\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{\sqrt{ex^2+d}}{x\sqrt{1-cx}\sqrt{cx+1}} dx}{e} + \frac{\sqrt{d + ex^2}(a + b \operatorname{sech}^{-1}(cx))}{e}$$

↓ 2036

$$\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{\sqrt{ex^2+d}}{x\sqrt{1-c^2x^2}} dx}{e} + \frac{\sqrt{d + ex^2}(a + b \operatorname{sech}^{-1}(cx))}{e}$$

$$\begin{aligned}
& \downarrow 354 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{\sqrt{ex^2+d}}{x^2\sqrt{1-c^2x^2}} dx^2}{2e} + \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e} \\
& \downarrow 140 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( e \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 + \int \frac{d}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 \right)}{2e} + \\
& \quad \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e} \\
& \downarrow 27 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( e \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 + d \int \frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 \right)}{2e} + \\
& \quad \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e} \\
& \downarrow 66 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( d \int \frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 + 2e \int \frac{1}{-ex^4-c^2} d \frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}} \right)}{2e} + \\
& \quad \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e} \\
& \downarrow 104 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( 2d \int \frac{1}{x^4-d} d \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} + 2e \int \frac{1}{-ex^4-c^2} d \frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}} \right)}{2e} + \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e} \\
& \downarrow 218 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( 2d \int \frac{1}{x^4-d} d \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} - \frac{2\sqrt{e} \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c} \right)}{2e} + \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e} \\
& \downarrow 220 \\
& \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e} + \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{2\sqrt{e} \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c} - 2\sqrt{d} \operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right) \right)}{2e}
\end{aligned}$$

input `Int[(x*(a + b*ArcSech[c*x]))/Sqrt[d + e*x^2],x]`

output `(Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/e + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-2*Sqrt[e]*ArcTan[(Sqrt[e]*Sqrt[1 - c^2*x^2])/(c*Sqrt[d + e*x^2])])/c - 2*Sqrt[d]*ArcTanh[Sqrt[d + e*x^2]/(Sqrt[d]*Sqrt[1 - c^2*x^2])]))/(2*e)`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 104 `Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 140 `Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)), x_] := Simp[b*d^(m + n)*f^p Int[(a + b*x)^(m - 1)/(c + d*x)^m, x], x] + Int[(a + b*x)^(m - 1)*((e + f*x)^p/(c + d*x)^m)*ExpandToSum[(a + b*x)*(c + d*x)^(-p - 1) - (b*d^(-p - 1)*f^p)/(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[m + n + p + 1, 0] && ILtQ[p, 0] && (GtQ[m, 0] || SumSimplerQ[m, -1] || !(GtQ[n, 0] || SumSimplerQ[n, -1]))`

rule 218 `Int[(((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 220 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 354 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]`

rule 2036 `Int[(u_)*((c_) + (d_)*(x_)^(n_))^(q_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_), x_Symbol] := Int[u*(a1*a2 + b1*b2*x^n)^p*(c + d*x^n)^q, x] /; FreeQ[{a1, b1, a2, b2, c, d, n, p, q}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))`

rule 6853 `Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*(x_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcSech[c*x])/(2*e*(p + 1))), x] + Simp[b*(Sqrt[1 + c*x]/(2*e*(p + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x^2)^(p + 1)/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[p, -1]`

## Maple [F]

$$\int \frac{x(a + b \operatorname{arcsech}(cx))}{\sqrt{x^2 e + d}} dx$$

input `int(x*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2), x)`

output `int(x*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2), x)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 257 vs. 2(95) = 190.

Time = 0.23 (sec) , antiderivative size = 1102, normalized size of antiderivative = 7.20

$$\int \frac{x(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \text{Too large to display}$$

input `integrate(x*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="fricas")`

output

```
[1/4*(4*sqrt(e*x^2 + d)*b*c*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + b*c*sqrt(d)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 + 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) + 4*sqrt(e*x^2 + d)*a*c - b*sqrt(-e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e + 8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^4*e*x^3 + (c^4*d - c^2*e)*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e^2))/(c*e), 1/4*(4*sqrt(e*x^2 + d)*b*c*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + b*c*sqrt(d)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 + 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) + 4*sqrt(e*x^2 + d)*a*c - 2*b*sqrt(e)*arctan(1/2*(2*c^2*e*x^3 + (c^2*d - e)*x)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*e^2*x^4 + (c^2*d*e - e^2)*x^2 - d*e))/(c*e), -1/4*(2*b*c*sqrt(-d)*arctan(-1/2*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(-d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*d*e*x^4 + (c^2*d^2 - d*e)*x^2 - d^2)) - 4*sqrt(e*x^2 + d)*b*c*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - 4*sqrt(e*x^2 + d)*a*c + b*sqrt(-e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e + 8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^4*e*x^3 + (c^4*d - c^2*e)*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e^2))/(c*e), -1/2*(b*c*sqrt(-d)*arctan(-1/2*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(-d)*sqrt(-(c^2*x^2 - 1)/(c^2*...
```

**Sympy [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{x(a + b \operatorname{arsech}(cx))}{\sqrt{d + ex^2}} dx$$

input `integrate(x*(a+b*asech(c*x))/(e*x**2+d)**(1/2),x)`

output `Integral(x*(a + b*asech(c*x))/sqrt(d + e*x**2), x)`

**Maxima [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x}{\sqrt{ex^2 + d}} dx$$

input `integrate(x*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="maxima")`

output `b*(sqrt(e*x^2 + d)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)/e - integrate((2*(c^2*e*x^2 - e)*x*log(sqrt(x)) + (c^2*e*x^2*log(c) - e*log(c))*x + (2*(c^2*e*x^2 - e)*x*log(sqrt(x)) + ((e*log(c) + e)*c^2*x^2 + c^2*d - e*log(c))*x)*e^(1/2*log(c*x + 1) + 1/2*log(-c*x + 1)))/((c^2*e*x^2 + (c^2*e*x^2 - e)*e^(1/2*log(c*x + 1) + 1/2*log(-c*x + 1)) - e)*sqrt(e*x^2 + d)), x) + sqrt(e*x^2 + d)*a/e`

**Giac [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x}{\sqrt{ex^2 + d}} dx$$

input `integrate(x*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x/sqrt(e*x^2 + d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{x(a + b \operatorname{acosh}(\frac{1}{cx}))}{\sqrt{ex^2 + d}} dx$$

input `int((x*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(1/2),x)`

output `int((x*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(1/2), x)`

**Reduce [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{x(a \operatorname{sech}(cx) b + a)}{\sqrt{ex^2 + d}} dx$$

input `int(x*(a+b*asech(c*x))/(e*x^2+d)^(1/2),x)`

output `int(x*(a+b*asech(c*x))/(e*x^2+d)^(1/2),x)`

$$3.152 \quad \int \frac{a+b\operatorname{sech}^{-1}(cx)}{x\sqrt{d+ex^2}} dx$$

Optimal result	1231
Mathematica [N/A]	1231
Rubi [N/A]	1232
Maple [N/A]	1232
Fricas [N/A]	1233
Sympy [N/A]	1233
Maxima [F(-2)]	1233
Giac [N/A]	1234
Mupad [N/A]	1234
Reduce [N/A]	1235

### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x\sqrt{d + ex^2}} dx = \operatorname{Int}\left(\frac{a + b\operatorname{sech}^{-1}(cx)}{x\sqrt{d + ex^2}}, x\right)$$

output `Defer(Int)((a+b*arcsech(c*x))/x/(e*x^2+d)^(1/2),x)`

### Mathematica [N/A]

Not integrable

Time = 2.04 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x\sqrt{d + ex^2}} dx = \int \frac{a + b\operatorname{sech}^{-1}(cx)}{x\sqrt{d + ex^2}} dx$$

input `Integrate[(a + b*ArcSech[c*x])/(x*Sqrt[d + e*x^2]),x]`

output `Integrate[(a + b*ArcSech[c*x])/(x*Sqrt[d + e*x^2]), x]`



**Rubi [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x \sqrt{d + ex^2}} dx$$

↓ 6865

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x \sqrt{d + ex^2}} dx$$

input `Int[(a + b*ArcSech[c*x])/(x*Sqrt[d + e*x^2]),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{a + b \operatorname{arcsech}(cx)}{x \sqrt{x^2 e + d}} dx$$

input `int((a+b*arcsech(c*x))/x/(e*x^2+d)^(1/2),x)`

output `int((a+b*arcsech(c*x))/x/(e*x^2+d)^(1/2),x)`

**Fricas [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.35

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x\sqrt{d + ex^2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{\sqrt{ex^2 + d}} dx$$

input `integrate((a+b*arcsech(c*x))/x/(e*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)/(e*x^3 + d*x), x)`

**Sympy [N/A]**

Not integrable

Time = 2.87 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x\sqrt{d + ex^2}} dx = \int \frac{a + b \operatorname{asech}(cx)}{x\sqrt{d + ex^2}} dx$$

input `integrate((a+b*asech(c*x))/x/(e*x**2+d)**(1/2),x)`

output `Integral((a + b*asech(c*x))/(x*sqrt(d + e*x**2)), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x\sqrt{d + ex^2}} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/x/(e*x^2+d)^(1/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x \sqrt{d + ex^2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{\sqrt{ex^2 + d}} dx$$

input

```
integrate((a+b*arcsech(c*x))/x/(e*x^2+d)^(1/2),x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)/(sqrt(e*x^2 + d)*x), x)
```

**Mupad [N/A]**

Not integrable

Time = 4.27 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x \sqrt{d + ex^2}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x \sqrt{ex^2 + d}} dx$$

input

```
int((a + b*acosh(1/(c*x)))/(x*(d + e*x^2)^(1/2)),x)
```

output

```
int((a + b*acosh(1/(c*x)))/(x*(d + e*x^2)^(1/2)), x)
```

**Reduce [N/A]**

Not integrable

Time = 200.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x \sqrt{d + ex^2}} dx = \int \frac{a \operatorname{sech}(cx) b + a}{x \sqrt{ex^2 + d}} dx$$

input `int((a+b*asech(c*x))/x/(e*x^2+d)^(1/2),x)`output `int((a+b*asech(c*x))/x/(e*x^2+d)^(1/2),x)`

$$3.153 \quad \int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^3\sqrt{d+ex^2}} dx$$

Optimal result	1236
Mathematica [N/A]	1236
Rubi [N/A]	1237
Maple [N/A]	1237
Fricas [N/A]	1238
Sympy [N/A]	1238
Maxima [F(-2)]	1238
Giac [N/A]	1239
Mupad [N/A]	1239
Reduce [N/A]	1240

### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^3\sqrt{d + ex^2}} dx = \operatorname{Int}\left(\frac{a + b\operatorname{sech}^{-1}(cx)}{x^3\sqrt{d + ex^2}}, x\right)$$

output `Defer(Int)((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(1/2),x)`

### Mathematica [N/A]

Not integrable

Time = 5.16 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^3\sqrt{d + ex^2}} dx = \int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^3\sqrt{d + ex^2}} dx$$

input `Integrate[(a + b*ArcSech[c*x])/(x^3*Sqrt[d + e*x^2]),x]`

output `Integrate[(a + b*ArcSech[c*x])/(x^3*Sqrt[d + e*x^2]), x]`

**Rubi [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 \sqrt{d + ex^2}} dx$$

↓ 6865

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 \sqrt{d + ex^2}} dx$$

input `Int[(a + b*ArcSech[c*x])/(x^3*Sqrt[d + e*x^2]),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{a + b \operatorname{arcsech}(cx)}{x^3 \sqrt{x^2 e + d}} dx$$

input `int((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(1/2),x)`

output `int((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(1/2),x)`

**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.43

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 \sqrt{d + ex^2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{\sqrt{ex^2 + dx^3}} dx$$

input `integrate((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)/(e*x^5 + d*x^3), x)`

**Sympy [N/A]**

Not integrable

Time = 9.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 \sqrt{d + ex^2}} dx = \int \frac{a + b \operatorname{asech}(cx)}{x^3 \sqrt{d + ex^2}} dx$$

input `integrate((a+b*asech(c*x))/x**3/(e*x**2+d)**(1/2),x)`

output `Integral((a + b*asech(c*x))/(x**3*sqrt(d + e*x**2)), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 \sqrt{d + ex^2}} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(1/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 \sqrt{d + ex^2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{\sqrt{ex^2 + d} x^3} dx$$

input

```
integrate((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(1/2),x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)/(sqrt(e*x^2 + d)*x^3), x)
```

**Mupad [N/A]**

Not integrable

Time = 4.43 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 \sqrt{d + ex^2}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x^3 \sqrt{ex^2 + d}} dx$$

input

```
int((a + b*acosh(1/(c*x)))/(x^3*(d + e*x^2)^(1/2)),x)
```

output

```
int((a + b*acosh(1/(c*x)))/(x^3*(d + e*x^2)^(1/2)), x)
```



**Reduce [N/A]**

Not integrable

Time = 200.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 \sqrt{d + ex^2}} dx = \int \frac{a \operatorname{sech}(cx) b + a}{x^3 \sqrt{ex^2 + d}} dx$$

input `int((a+b*asech(c*x))/x^3/(e*x^2+d)^(1/2),x)`output `int((a+b*asech(c*x))/x^3/(e*x^2+d)^(1/2),x)`

$$3.154 \quad \int \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx$$

Optimal result	1241
Mathematica [N/A]	1241
Rubi [N/A]	1242
Maple [N/A]	1242
Fricas [N/A]	1243
Sympy [N/A]	1243
Maxima [F(-2)]	1243
Giac [N/A]	1244
Mupad [N/A]	1244
Reduce [N/A]	1245

### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \operatorname{Int} \left( \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}}, x \right)$$

output `Defer(Int)(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x)`

### Mathematica [N/A]

Not integrable

Time = 10.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx$$

input `Integrate[(x^2*(a + b*ArcSech[c*x]))/Sqrt[d + e*x^2],x]`

output `Integrate[(x^2*(a + b*ArcSech[c*x]))/Sqrt[d + e*x^2], x]`

**Rubi [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx$$

↓ 6865

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx$$

input `Int[(x^2*(a + b*ArcSech[c*x]))/Sqrt[d + e*x^2],x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.16 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{x^2(a + b \operatorname{arcsech}(cx))}{\sqrt{x^2e + d}} dx$$

input `int(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x)`

output `int(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x)`

**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^2}{\sqrt{ex^2 + d}} dx$$

input `integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral((b*x^2*arcsech(c*x) + a*x^2)/sqrt(e*x^2 + d), x)`

**Sympy [N/A]**

Not integrable

Time = 4.44 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{x^2(a + b \operatorname{asech}(cx))}{\sqrt{d + ex^2}} dx$$

input `integrate(x**2*(a+b*asech(c*x))/(e*x**2+d)**(1/2),x)`

output `Integral(x**2*(a + b*asech(c*x))/sqrt(d + e*x**2), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \text{Exception raised: ValueError}$$

input `integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^2}{\sqrt{ex^2 + d}} dx$$

input

```
integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)*x^2/sqrt(e*x^2 + d), x)
```

**Mupad [N/A]**

Not integrable

Time = 4.42 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(\frac{1}{cx}))}{\sqrt{ex^2 + d}} dx$$

input

```
int((x^2*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(1/2),x)
```

output

```
int((x^2*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(1/2), x)
```

**Reduce [N/A]**

Not integrable

Time = 200.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{x^2(\operatorname{asech}(cx)b + a)}{\sqrt{ex^2 + d}} dx$$

input `int(x^2*(a+b*asech(c*x))/(e*x^2+d)^(1/2),x)`output `int(x^2*(a+b*asech(c*x))/(e*x^2+d)^(1/2),x)`

$$3.155 \quad \int \frac{a+b\operatorname{sech}^{-1}(cx)}{\sqrt{d+ex^2}} dx$$

Optimal result	1246
Mathematica [N/A]	1246
Rubi [N/A]	1247
Maple [N/A]	1247
Fricas [N/A]	1248
Sympy [N/A]	1248
Maxima [F(-2)]	1248
Giac [N/A]	1249
Mupad [N/A]	1249
Reduce [N/A]	1250

### Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{\sqrt{d + ex^2}} dx = \operatorname{Int}\left(\frac{a + b\operatorname{sech}^{-1}(cx)}{\sqrt{d + ex^2}}, x\right)$$

output `Defer(Int)((a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x)`

### Mathematica [N/A]

Not integrable

Time = 0.70 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{\sqrt{d + ex^2}} dx = \int \frac{a + b\operatorname{sech}^{-1}(cx)}{\sqrt{d + ex^2}} dx$$

input `Integrate[(a + b*ArcSech[c*x])/Sqrt[d + e*x^2],x]`

output `Integrate[(a + b*ArcSech[c*x])/Sqrt[d + e*x^2], x]`

**Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex^2}} dx$$

↓ 6865

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex^2}} dx$$

input `Int[(a + b*ArcSech[c*x])/Sqrt[d + e*x^2], x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.16 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{a + b \operatorname{arcsech}(cx)}{\sqrt{x^2 e + d}} dx$$

input `int((a+b*arcsech(c*x))/(e*x^2+d)^(1/2), x)`

output `int((a+b*arcsech(c*x))/(e*x^2+d)^(1/2), x)`



**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex^2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{\sqrt{ex^2 + d}} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral((b*arcsech(c*x) + a)/sqrt(e*x^2 + d), x)`

**Sympy [N/A]**

Not integrable

Time = 1.35 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex^2}} dx = \int \frac{a + b \operatorname{asech}(cx)}{\sqrt{d + ex^2}} dx$$

input `integrate((a+b*asech(c*x))/(e*x**2+d)**(1/2),x)`

output `Integral((a + b*asech(c*x))/sqrt(d + e*x**2), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex^2}} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex^2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{\sqrt{ex^2 + d}} dx$$

input

```
integrate((a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)/sqrt(e*x^2 + d), x)
```

**Mupad [N/A]**

Not integrable

Time = 4.28 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.20

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex^2}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{\sqrt{ex^2 + d}} dx$$

input

```
int((a + b*acosh(1/(c*x)))/(d + e*x^2)^(1/2),x)
```

output

```
int((a + b*acosh(1/(c*x)))/(d + e*x^2)^(1/2), x)
```

**Reduce [N/A]**

Not integrable

Time = 67.93 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{\sqrt{d + ex^2}} dx = \int \frac{a \operatorname{sech}(cx) b + a}{\sqrt{ex^2 + d}} dx$$

input `int((a+b*asech(c*x))/(e*x^2+d)^(1/2),x)`output `int((a+b*asech(c*x))/(e*x^2+d)^(1/2),x)`

**3.156**  $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^2\sqrt{d+ex^2}} dx$

Optimal result	1251
Mathematica [C] (verified)	1252
Rubi [A] (verified)	1252
Maple [F]	1256
Fricas [A] (verification not implemented)	1256
Sympy [F]	1257
Maxima [F(-2)]	1257
Giac [F]	1258
Mupad [F(-1)]	1258
Reduce [F]	1258

**Optimal result**

Integrand size = 23, antiderivative size = 218

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^2\sqrt{d + ex^2}} dx$$

$$= \frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{dx} - \frac{\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{dx}$$

$$+ \frac{bc\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1+\frac{ex^2}{d}}E(\arcsin(cx) | -\frac{e}{c^2d})}{\sqrt{d+ex^2}}$$

$$- \frac{b(c^2d + e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1+\frac{ex^2}{d}}\operatorname{EllipticF}(\arcsin(cx), -\frac{e}{c^2d})}{cd\sqrt{d+ex^2}}$$

output

```
b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*(e*x^2+d)^(1/2)/d/x-(
e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/d/x+b*c*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*
(1+e*x^2/d)^(1/2)*EllipticE(c*x,(-e/c^2/d)^(1/2))/(e*x^2+d)^(1/2)-b*(c^2*d
+e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(1+e*x^2/d)^(1/2)*EllipticF(c*x,(-e/c^
2/d)^(1/2))/c/d/(e*x^2+d)^(1/2)
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 23.81 (sec) , antiderivative size = 501, normalized size of antiderivative = 2.30

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 \sqrt{d + ex^2}} dx =$$

$$a\left(\frac{d}{x} + ex\right) + bc\sqrt{\frac{1-cx}{1+cx}}(d + ex^2) - \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)(d+ex^2)}{x} + \frac{b(d+ex^2)\operatorname{sech}^{-1}(cx)}{x} + \frac{b\sqrt{\frac{1-cx}{1+cx}}\sqrt{\frac{c(\sqrt{d+i\sqrt{ex}})}{(c\sqrt{d+i\sqrt{e}})(1+cx)}}(i\sqrt{d-$$

input

```
Integrate[(a + b*ArcSech[c*x])/(x^2*Sqrt[d + e*x^2]),x]
```

output

```
-((a*(d/x + e*x) + b*c*Sqrt[(1 - c*x)/(1 + c*x)]*(d + e*x^2) - (b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(d + e*x^2))/x + (b*(d + e*x^2)*ArcSech[c*x])/x + (b*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[(c*(Sqrt[d] + I*Sqrt[e]*x))/((c*Sqrt[d] + I*Sqrt[e])*(1 + c*x))]*(I*Sqrt[d] + Sqrt[e]*x)*((c*Sqrt[d] - I*Sqrt[e])*EllipticE[I*ArcSinh[Sqrt[((c^2*d + e)*(1 - c*x))/((c*Sqrt[d] + I*Sqrt[e])^2*(1 + c*x))]]], (c*Sqrt[d] + I*Sqrt[e])^2/(c*Sqrt[d] - I*Sqrt[e])^2 + (2*I)*Sqrt[e]*EllipticF[I*ArcSinh[Sqrt[((c^2*d + e)*(1 - c*x))/((c*Sqrt[d] + I*Sqrt[e])^2*(1 + c*x))]]], (c*Sqrt[d] + I*Sqrt[e])^2/(c*Sqrt[d] - I*Sqrt[e])^2))/Sqrt[-(((c*Sqrt[d] - I*Sqrt[e])*(-1 + c*x))/((c*Sqrt[d] + I*Sqrt[e])*(1 + c*x)))]*Sqrt[(c*(Sqrt[d] - I*Sqrt[e]*x))/((c*Sqrt[d] - I*Sqrt[e])*(1 + c*x))]))/(d*Sqrt[d + e*x^2]))
```

**Rubi [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.85, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.435$ , Rules used = {6855, 25, 27, 377, 27, 326, 323, 321, 330, 327}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 \sqrt{d + ex^2}} dx \\
& \quad \downarrow \text{6855} \\
& b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int -\frac{\sqrt{ex^2+d}}{dx^2 \sqrt{1-c^2x^2}} dx - \frac{\sqrt{d+ex^2}(a + b \operatorname{sech}^{-1}(cx))}{dx} \\
& \quad \downarrow \text{25} \\
& -b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{\sqrt{ex^2+d}}{dx^2 \sqrt{1-c^2x^2}} dx - \frac{\sqrt{d+ex^2}(a + b \operatorname{sech}^{-1}(cx))}{dx} \\
& \quad \downarrow \text{27} \\
& -\frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{\sqrt{ex^2+d}}{x^2 \sqrt{1-c^2x^2}} dx}{d} - \frac{\sqrt{d+ex^2}(a + b \operatorname{sech}^{-1}(cx))}{dx} \\
& \quad \downarrow \text{377} \\
& -\frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \left( \int \frac{e \sqrt{1-c^2x^2}}{\sqrt{ex^2+d}} dx - \frac{\sqrt{1-c^2x^2} \sqrt{d+ex^2}}{x} \right)}{d} - \frac{\sqrt{d+ex^2}(a + b \operatorname{sech}^{-1}(cx))}{dx} \\
& \quad \downarrow \text{27} \\
& -\frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \left( e \int \frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}} dx - \frac{\sqrt{1-c^2x^2} \sqrt{d+ex^2}}{x} \right)}{d} - \frac{\sqrt{d+ex^2}(a + b \operatorname{sech}^{-1}(cx))}{dx} \\
& \quad \downarrow \text{326} \\
& \frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \left( e \left( \frac{(c^2d+e) \int \frac{1}{\sqrt{1-c^2x^2} \sqrt{ex^2+d}} dx}{e} - \frac{c^2 \int \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} dx}{e} \right) - \frac{\sqrt{1-c^2x^2} \sqrt{d+ex^2}}{x} \right)}{d} - \frac{\sqrt{d+ex^2}(a + b \operatorname{sech}^{-1}(cx))}{dx} \\
& \quad \downarrow \text{323} \\
& \frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \left( e \left( \frac{(c^2d+e) \sqrt{\frac{ex^2}{d}+1} \int \frac{1}{\sqrt{1-c^2x^2} \sqrt{\frac{ex^2}{d}+1}} dx}{e \sqrt{d+ex^2}} - \frac{c^2 \int \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} dx}{e} \right) - \frac{\sqrt{1-c^2x^2} \sqrt{d+ex^2}}{x} \right)}{d} - \frac{\sqrt{d+ex^2}(a + b \operatorname{sech}^{-1}(cx))}{dx} \\
& \quad \downarrow \text{321}
\end{aligned}$$

$$\begin{aligned}
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(e\left(\frac{(c^2d+e)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}}-\frac{c^2\int\frac{\sqrt{\frac{ex^2}{d}+1}}{\sqrt{1-c^2x^2}}dx}{e}\right)-\frac{\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{x}\right)}{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))} \\
 & \quad \downarrow \text{330} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(e\left(\frac{(c^2d+e)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}}-\frac{c^2\sqrt{d+ex^2}\int\frac{\sqrt{\frac{ex^2}{d}+1}}{\sqrt{1-c^2x^2}}dx}{e\sqrt{\frac{ex^2}{d}+1}}\right)-\frac{\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{x}\right)}{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))} \\
 & \quad \downarrow \text{327} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(e\left(\frac{(c^2d+e)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}}-\frac{c\sqrt{d+ex^2}E\left(\arcsin(cx)\middle|-\frac{e}{c^2d}\right)}{e\sqrt{\frac{ex^2}{d}+1}}\right)-\frac{\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{x}\right)}{d}
 \end{aligned}$$

input `Int[(a + b*ArcSech[c*x])/(x^2*Sqrt[d + e*x^2]),x]`

output `-((Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/(d*x)) - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-(Sqrt[1 - c^2*x^2]*Sqrt[d + e*x^2])/x) + e*(-((c*Sqrt[d + e*x^2]*EllipticE[ArcSin[c*x], -(e/(c^2*d))])/(e*Sqrt[1 + (e*x^2)/d])) + ((c^2*d + e)*Sqrt[1 + (e*x^2)/d]*EllipticF[ArcSin[c*x], -(e/(c^2*d))])/(c*e*Sqrt[d + e*x^2])))`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 321 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[imp[(1/(Sqrt[a]*Sqrt[c]*Rt[-d/c, 2]))*EllipticF[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-b/a, -d/c])]`
- rule 323 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[Sqrt[1 + (d/c)*x^2]/Sqrt[c + d*x^2] Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d/c)*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]`
- rule 326 `Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[b/d Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Simp[(b*c - a*d)/d Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]`
- rule 327 `Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]/(Sqrt[c]*Rt[-d/c, 2]))*EllipticE[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]`
- rule 330 `Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[Sqrt[a + b*x^2]/Sqrt[1 + (b/a)*x^2] Int[Sqrt[1 + (b/a)*x^2]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]`



rule 377

```
Int[((e._)*(x_))^(m_)*((a_) + (b._)*(x_)^2)^(p_)*((c_) + (d._)*(x_)^2)^(q_)
, x_Symbol] := Simp[(e*x)^(m+1)*(a + b*x^2)^(p+1)*((c + d*x^2)^q/(a*e*(
m+1))), x] - Simp[1/(a*e^2*(m+1)) Int[(e*x)^(m+2)*(a + b*x^2)^p*(c
+ d*x^2)^(q-1)*Simp[b*c*(m+1) + 2*(b*c*(p+1) + a*d*q) + d*(b*(m+1)
+ 2*b*(p+q+1))*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b
*c - a*d, 0] && LtQ[0, q, 1] && LtQ[m, -1] && IntBinomialQ[a, b, c, d, e, m
, 2, p, q, x]
```

rule 6855

```
Int[((a_) + ArcSech[(c._)*(x_)])*(b._))*((f._)*(x_))^(m_)*((d_) + (e._)*(
x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Si
mp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)]
Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; Fre
eQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m-1)/2, 0] &&
GtQ[m+2*p+3, 0])) || (IGtQ[(m+1)/2, 0] && !(ILtQ[p, 0] && GtQ[m+2
*p+3, 0])) || (ILtQ[(m+2*p+1)/2, 0] && !ILtQ[(m-1)/2, 0]))
```

## Maple [F]

$$\int \frac{a + b \operatorname{arcsech}(cx)}{x^2 \sqrt{x^2 e + d}} dx$$

input

```
int((a+b*arcsech(c*x))/x^2/(e*x^2+d)^(1/2),x)
```

output

```
int((a+b*arcsech(c*x))/x^2/(e*x^2+d)^(1/2),x)
```

## Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.71

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 \sqrt{d + ex^2}} dx =$$

$$\frac{\sqrt{ex^2 + d} bcd \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2} + 1}}{cx}\right) - \left(bc^2 dx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - acd\right) \sqrt{ex^2 + d} - (bc^4 dx E(\arcsin(cx) | -\frac{e}{c^2 d})}{cd^2 x}$$

input

```
integrate((a+b*arcsech(c*x))/x^2/(e*x^2+d)^(1/2),x, algorithm="fricas")
```

output

```
-(sqrt(e*x^2 + d)*b*c*d*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)
) - (b*c^2*d*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - a*c*d)*sqrt(e*x^2 + d) - (
b*c^4*d*x*elliptic_e(arcsin(c*x), -e/(c^2*d)) - (b*c^4*d + b*e)*x*elliptic
_f(arcsin(c*x), -e/(c^2*d)))*sqrt(d))/(c*d^2*x)
```

## Sympy [F]

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 \sqrt{d + ex^2}} dx = \int \frac{a + b \operatorname{asech}(cx)}{x^2 \sqrt{d + ex^2}} dx$$

input

```
integrate((a+b*asech(c*x))/x**2/(e*x**2+d)**(1/2),x)
```

output

```
Integral((a + b*asech(c*x))/(x**2*sqrt(d + e*x**2)), x)
```

## Maxima [F(-2)]

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 \sqrt{d + ex^2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((a+b*arcsech(c*x))/x^2/(e*x^2+d)^(1/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 \sqrt{d + ex^2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{\sqrt{ex^2 + d}} dx$$

input `integrate((a+b*arcsech(c*x))/x^2/(e*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/(sqrt(e*x^2 + d)*x^2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 \sqrt{d + ex^2}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x^2 \sqrt{ex^2 + d}} dx$$

input `int((a + b*acosh(1/(c*x)))/(x^2*(d + e*x^2)^(1/2)),x)`

output `int((a + b*acosh(1/(c*x)))/(x^2*(d + e*x^2)^(1/2)), x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 \sqrt{d + ex^2}} dx = \int \frac{a \operatorname{sech}(cx) b + a}{x^2 \sqrt{ex^2 + d}} dx$$

input `int((a+b*asech(c*x))/x^2/(e*x^2+d)^(1/2),x)`

output `int((a+b*asech(c*x))/x^2/(e*x^2+d)^(1/2),x)`

**3.157**  $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^4\sqrt{d+ex^2}} dx$

Optimal result	1259
Mathematica [C] (verified)	1260
Rubi [A] (verified)	1261
Maple [F]	1266
Fricas [A] (verification not implemented)	1266
Sympy [F]	1267
Maxima [F(-2)]	1267
Giac [F]	1267
Mupad [F(-1)]	1268
Reduce [F]	1268

**Optimal result**

Integrand size = 23, antiderivative size = 346

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^4\sqrt{d+ex^2}} dx$$

$$= \frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{9dx^3}$$

$$+ \frac{b(2c^2d - 5e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{9d^2x}$$

$$- \frac{\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{3dx^3} + \frac{2e\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{3d^2x}$$

$$+ \frac{bc(2c^2d - 5e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{d+ex^2}E(\arcsin(cx) | -\frac{e}{c^2d})}{9d^2\sqrt{1+\frac{ex^2}{d}}}$$

$$- \frac{2b(c^2d - 3e)(c^2d + e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1+\frac{ex^2}{d}}\operatorname{EllipticF}(\arcsin(cx), -\frac{e}{c^2d})}{9cd^2\sqrt{d+ex^2}}$$

output

```

1/9*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*(e*x^2+d)^(1/2)/d
/x^3+1/9*b*(2*c^2*d-5*e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)
)*(e*x^2+d)^(1/2)/d^2/x-1/3*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/d/x^3+2/3*e
*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/d^2/x+1/9*b*c*(2*c^2*d-5*e)*(1/(c*x+1)
)^(1/2)*(c*x+1)^(1/2)*(e*x^2+d)^(1/2)*EllipticE(c*x,(-e/c^2/d)^(1/2))/d^2/
(1+e*x^2/d)^(1/2)-2/9*b*(c^2*d-3*e)*(c^2*d+e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1
/2)*(1+e*x^2/d)^(1/2)*EllipticF(c*x,(-e/c^2/d)^(1/2))/c/d^2/(e*x^2+d)^(1/2)
)

```

### Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 24.17 (sec) , antiderivative size = 612, normalized size of antiderivative = 1.77

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4 \sqrt{d + ex^2}} dx$$

$$\frac{bd\sqrt{\frac{1-cx}{1+cx}}(d+ex^2)}{x^3} + \frac{bcd\sqrt{\frac{1-cx}{1+cx}}(d+ex^2)}{x^2} + \frac{b(2c^2d-5e)\sqrt{\frac{1-cx}{1+cx}}(d+ex^2)}{x} - \frac{3a(d-2ex^2)(d+ex^2)}{x^3} - \frac{3b(d-2ex^2)(d+ex^2)\operatorname{sech}^{-1}(cx)}{x^3}$$

=

input

```
Integrate[(a + b*ArcSech[c*x])/(x^4*Sqrt[d + e*x^2]),x]
```

output

```

((b*d*Sqrt[(1 - c*x)/(1 + c*x)]*(d + e*x^2))/x^3 + (b*c*d*Sqrt[(1 - c*x)/(
1 + c*x)]*(d + e*x^2))/x^2 + (b*(2*c^2*d - 5*e)*Sqrt[(1 - c*x)/(1 + c*x)]*
(d + e*x^2))/x - (3*a*(d - 2*e*x^2)*(d + e*x^2))/x^3 - (3*b*(d - 2*e*x^2)*
(d + e*x^2)*ArcSech[c*x])/x^3 - (b*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[(c*(Sqrt
[d] + I*Sqrt[e]*x))/((c*Sqrt[d] + I*Sqrt[e])*(1 + c*x))]*(I*Sqrt[d] + Sqrt
[e]*x)*((2*c^3*d^(3/2) - (2*I)*c^2*d*Sqrt[e] - 5*c*Sqrt[d]*e + (5*I)*e^(3/
2))*EllipticE[I*ArcSinh[Sqrt[((c^2*d + e)*(1 - c*x))/((c*Sqrt[d] + I*Sqrt[
e])^2*(1 + c*x))]]], (c*Sqrt[d] + I*Sqrt[e])^2/(c*Sqrt[d] - I*Sqrt[e])^2 +
2*((2*I)*c^2*d - c*Sqrt[d]*Sqrt[e] - (6*I)*e)*Sqrt[e]*EllipticF[I*ArcSinh
[Sqrt[((c^2*d + e)*(1 - c*x))/((c*Sqrt[d] + I*Sqrt[e])^2*(1 + c*x))]]], (c
Sqrt[d] + I*Sqrt[e])^2/(c*Sqrt[d] - I*Sqrt[e])^2))/((Sqrt[-((c*Sqrt[d] -
I*Sqrt[e])*(-1 + c*x))/((c*Sqrt[d] + I*Sqrt[e])*(1 + c*x))])*Sqrt[(c*(Sqrt
[d] - I*Sqrt[e]*x))/((c*Sqrt[d] - I*Sqrt[e])*(1 + c*x))]))/(9*d^2*Sqrt[d +
e*x^2])

```

## Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 288, normalized size of antiderivative = 0.83, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.478$ , Rules used = {6855, 27, 442, 445, 25, 27, 399, 323, 321, 330, 327}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4 \sqrt{d + ex^2}} dx \\
 & \quad \downarrow \text{6855} \\
 & b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int -\frac{(d-2ex^2)\sqrt{ex^2+d}}{3d^2 x^4 \sqrt{1-c^2x^2}} dx + \frac{2e\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3d^2 x} - \\
 & \quad \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3dx^3} \\
 & \quad \downarrow \text{27} \\
 & -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(d-2ex^2)\sqrt{ex^2+d}}{x^4\sqrt{1-c^2x^2}} dx}{3d^2} + \frac{2e\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3d^2 x} - \\
 & \quad \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3dx^3} \\
 & \quad \downarrow \text{442}
 \end{aligned}$$

$$\begin{aligned}
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\int\frac{(c^2d-6e)ex^2+d(2c^2d-5e)}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx-\frac{d\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{3x^3}\right)}{3d^2} + \\
& \frac{2e\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3d^2x} - \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3dx^3} \\
& \quad \downarrow 445 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\left(-\int\frac{de\left(-\left((2c^2d-5e)x^2c^2\right)+dc^2-6e\right)}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx-\frac{\sqrt{1-c^2x^2}(2c^2d-5e)\sqrt{d+ex^2}}{x}\right)-\frac{d\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{3x^3}\right)}{3d^2} + \\
& \frac{2e\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3d^2x} - \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3dx^3} \\
& \quad \downarrow 25 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\left(\int\frac{de\left(-\left((2c^2d-5e)x^2c^2\right)+dc^2-6e\right)}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx-\frac{\sqrt{1-c^2x^2}(2c^2d-5e)\sqrt{d+ex^2}}{x}\right)-\frac{d\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{3x^3}\right)}{3d^2} + \\
& \frac{2e\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3d^2x} - \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3dx^3} \\
& \quad \downarrow 27 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\left(e\int\frac{-\left((2c^2d-5e)x^2c^2\right)+dc^2-6e}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx-\frac{\sqrt{1-c^2x^2}(2c^2d-5e)\sqrt{d+ex^2}}{x}\right)-\frac{d\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{3x^3}\right)}{3d^2} + \\
& \frac{2e\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3d^2x} - \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3dx^3} \\
& \quad \downarrow 399 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\left(e\left(\frac{2(c^2d-3e)(c^2d+e)\int\frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx}{e}-\frac{c^2(2c^2d-5e)\int\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}dx}{e}\right)-\frac{\sqrt{1-c^2x^2}(2c^2d-5e)\sqrt{d+ex^2}}{x}\right)}{3d^2} + \\
& \frac{2e\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3d^2x} - \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3dx^3} \\
& \quad \downarrow 323
\end{aligned}$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\left(e\left(\frac{2(c^2d-3e)(c^2d+e)\sqrt{\frac{ex^2}{d}+1}\int\frac{1}{\sqrt{1-c^2x^2}\sqrt{\frac{ex^2}{d}+1}}dx-\frac{c^2(2c^2d-5e)\int\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}dx}{e\sqrt{d+ex^2}}\right)-\frac{\sqrt{1-c^2x^2}(2c^2d-5e)}{x}\right)\right)$$

$$\frac{2e\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3d^2x}-\frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3dx^3}$$

321

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\left(e\left(\frac{2(c^2d-3e)(c^2d+e)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)-\frac{c^2(2c^2d-5e)\int\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}dx}{e\sqrt{d+ex^2}}\right)-\frac{\sqrt{1-c^2x^2}(2c^2d-5e)}{x}\right)\right)$$

$$\frac{2e\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3d^2x}-\frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3dx^3}$$

330

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\left(e\left(\frac{2(c^2d-3e)(c^2d+e)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)-\frac{c^2(2c^2d-5e)\sqrt{d+ex^2}\int\frac{\sqrt{\frac{ex^2}{d}+1}}{\sqrt{1-c^2x^2}}dx}{e\sqrt{\frac{ex^2}{d}+1}}\right)-\frac{\sqrt{1-c^2x^2}(2c^2d-5e)}{x}\right)\right)$$

$$\frac{2e\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3d^2x}-\frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3dx^3}$$

327

$$\frac{2e\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3d^2x}-\frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{3dx^3}$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\frac{1}{3}\left(e\left(\frac{2(c^2d-3e)(c^2d+e)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)-\frac{c(2c^2d-5e)\sqrt{d+ex^2}E\left(\arcsin(cx)\middle|-\frac{e}{c^2d}\right)}{e\sqrt{\frac{ex^2}{d}+1}}\right)-\frac{\sqrt{1-c^2x^2}(2c^2d-5e)}{x}\right)\right)$$

3d<sup>2</sup>

input

```
Int[(a + b*ArcSech[c*x])/(x^4*sqrt[d + e*x^2]),x]
```



output

```
-1/3*(Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/(d*x^3) + (2*e*Sqrt[d + e*x^2]
*(a + b*ArcSech[c*x]))/(3*d^2*x) - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(
-1/3*(d*Sqrt[1 - c^2*x^2]*Sqrt[d + e*x^2])/x^3 + (-((2*c^2*d - 5*e)*Sqrt[
1 - c^2*x^2]*Sqrt[d + e*x^2])/x) + e*(-((c*(2*c^2*d - 5*e)*Sqrt[d + e*x^2]
*EllipticE[ArcSin[c*x], -(e/(c^2*d))])/(e*Sqrt[1 + (e*x^2)/d])) + (2*(c^2*
d - 3*e)*(c^2*d + e)*Sqrt[1 + (e*x^2)/d]*EllipticF[ArcSin[c*x], -(e/(c^2*d
))])/(c*e*Sqrt[d + e*x^2])))/3)/(3*d^2)
```

**Defintions of rubi rules used**

rule 25

```
Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 321

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> S
imp[(1/(Sqrt[a]*Sqrt[c]*Rt[-d/c, 2]))*EllipticF[ArcSin[Rt[-d/c, 2]*x], b*(c
/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a,
0] && !(NegQ[b/a] && SimplerSqrtQ[-b/a, -d/c])
```

rule 323

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> S
imp[Sqrt[1 + (d/c)*x^2]/Sqrt[c + d*x^2] Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (
d/c)*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]
```

rule 327

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[
(Sqrt[a]/(Sqrt[c]*Rt[-d/c, 2]))*EllipticE[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d)
)], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]
```

rule 330

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[
Sqrt[a + b*x^2]/Sqrt[1 + (b/a)*x^2] Int[Sqrt[1 + (b/a)*x^2]/Sqrt[c + d*x^
2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a,
0]
```

rule 399

```
Int[((e_) + (f_)*(x_)^2)/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] := Simp[f/b Int[Sqrt[a + b*x^2]/Sqrt[c + d*x^2], x], x] + Simp[(b*e - a*f)/b Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && !((PosQ[b/a] && PosQ[d/c]) || (NegQ[b/a] && (PosQ[d/c] || (GtQ[a, 0] && (!GtQ[c, 0] || SimplerSqrtQ[-b/a, -d/c])))))
```

rule 442

```
Int[((g_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2), x_Symbol] := Simp[e*(g*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^q/(a*g*(m + 1))), x] - Simp[1/(a*g^2*(m + 1)) Int[(g*x)^(m + 2)*(a + b*x^2)^p*(c + d*x^2)^(q - 1)*Simp[c*(b*e - a*f)*(m + 1) + e*2*(b*c*(p + 1) + a*d*q) + d*((b*e - a*f)*(m + 1) + b*e*2*(p + q + 1))*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && GtQ[q, 0] && LtQ[m, -1] && !(EqQ[q, 1] && SimplerQ[e + f*x^2, c + d*x^2])
```

rule 445

```
Int[((g_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2), x_Symbol] := Simp[e*(g*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(a*c*g*(m + 1))), x] + Simp[1/(a*c*g^2*(m + 1)) Int[(g*x)^(m + 2)*(a + b*x^2)^p*(c + d*x^2)^q*Simp[a*f*c*(m + 1) - e*(b*c + a*d)*(m + 2 + 1) - e*2*(b*c*p + a*d*q) - b*e*d*(m + 2*(p + q + 2) + 1))*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, p, q}, x] && LtQ[m, -1]
```

rule 6855

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

**Maple [F]**

$$\int \frac{a + b \operatorname{arcsech}(cx)}{x^4 \sqrt{x^2 e + d}} dx$$

input `int((a+b*arcsech(c*x))/x^4/(e*x^2+d)^(1/2),x)`

output `int((a+b*arcsech(c*x))/x^4/(e*x^2+d)^(1/2),x)`

**Fricas [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 244, normalized size of antiderivative = 0.71

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4 \sqrt{d + ex^2}} dx$$

$$= \frac{3(2bcde x^2 - bcd^2) \sqrt{ex^2 + d} \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2} + 1}}{cx}\right) + (6acde x^2 - 3acd^2 + (bc^2 d^2 x + (2bc^4 d^2 - 5bc^2 de)x^3) \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}}) \sqrt{ex^2 + d} + ((2b^2 c^6 d^2 - 5b^2 c^4 d^2 e) x^3 \operatorname{elliptic}_e(\arcsin(cx), -e/(c^2 d)) - (2b^2 c^6 d^2 - (5b^2 c^4 - b^2 c^2) d^2 e - 6b^2 e^2) x^3 \operatorname{elliptic}_f(\arcsin(cx), -e/(c^2 d))) \sqrt{d}}{c^3 d^3 x^3}$$

input `integrate((a+b*arcsech(c*x))/x^4/(e*x^2+d)^(1/2),x, algorithm="fricas")`

output `1/9*(3*(2*b*c*d*e*x^2 - b*c*d^2)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + (6*a*c*d*e*x^2 - 3*a*c*d^2 + (b*c^2*d^2*x + (2*b*c^4*d^2 - 5*b*c^2*d*e)*x^3)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d) + ((2*b*c^6*d^2 - 5*b*c^4*d*e)*x^3*elliptic_e(arcsin(c*x), -e/(c^2*d)) - (2*b*c^6*d^2 - (5*b*c^4 - b*c^2)*d^2*e - 6*b*e^2)*x^3*elliptic_f(arcsin(c*x), -e/(c^2*d)))*sqrt(d))/(c*d^3*x^3)`

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4 \sqrt{d + ex^2}} dx = \int \frac{a + b \operatorname{arsech}(cx)}{x^4 \sqrt{d + ex^2}} dx$$

input `integrate((a+b*asech(c*x))/x**4/(e*x**2+d)**(1/2),x)`

output `Integral((a + b*asech(c*x))/(x**4*sqrt(d + e*x**2)), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4 \sqrt{d + ex^2}} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/x^4/(e*x^2+d)^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4 \sqrt{d + ex^2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{\sqrt{ex^2 + dx^4}} dx$$

input `integrate((a+b*arcsech(c*x))/x^4/(e*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/(sqrt(e*x^2 + d)*x^4), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4 \sqrt{d + ex^2}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x^4 \sqrt{ex^2 + d}} dx$$

input `int((a + b*acosh(1/(c*x)))/(x^4*(d + e*x^2)^(1/2)),x)`

output `int((a + b*acosh(1/(c*x)))/(x^4*(d + e*x^2)^(1/2)), x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^4 \sqrt{d + ex^2}} dx = \int \frac{a \operatorname{sech}(cx) b + a}{x^4 \sqrt{ex^2 + d}} dx$$

input `int((a+b*asech(c*x))/x^4/(e*x^2+d)^(1/2),x)`

output `int((a+b*asech(c*x))/x^4/(e*x^2+d)^(1/2),x)`

**3.158** 
$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

Optimal result	1269
Mathematica [A] (verified)	1270
Rubi [A] (verified)	1270
Maple [F]	1274
Fricas [B] (verification not implemented)	1275
Sympy [F]	1276
Maxima [F(-2)]	1276
Giac [F]	1276
Mupad [F(-1)]	1277
Reduce [F]	1277

**Optimal result**

Integrand size = 23, antiderivative size = 278

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = -\frac{b \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2} \sqrt{d+ex^2}}{6c^2e^2} - \frac{d^2(a + b \operatorname{sech}^{-1}(cx))}{e^3 \sqrt{d+ex^2}} - \frac{2d\sqrt{d+ex^2}(a + b \operatorname{sech}^{-1}(cx))}{e^3} + \frac{(d+ex^2)^{3/2}(a + b \operatorname{sech}^{-1}(cx))}{3e^3} + \frac{b(9c^2d - e) \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{6c^3e^{5/2}} + \frac{8bd^{3/2} \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{3e^3}$$

output

```
-1/6*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*(e*x^2+d)^(1/2)/
c^2/e^2-d^2*(a+b*arcsech(c*x))/e^3/(e*x^2+d)^(1/2)-2*d*(e*x^2+d)^(1/2)*(a+
b*arcsech(c*x))/e^3+1/3*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x))/e^3+1/6*b*(9*c^
2*d-e)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctan(e^(1/2)*(-c^2*x^2+1)^(1/2)/c
/(e*x^2+d)^(1/2))/c^3/e^(5/2)+8/3*b*d^(3/2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2
)*arctanh((e*x^2+d)^(1/2)/d^(1/2)/(-c^2*x^2+1)^(1/2))/e^3
```

**Mathematica [A] (verified)**

Time = 22.08 (sec) , antiderivative size = 436, normalized size of antiderivative = 1.57

$$\int \frac{x^5(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \frac{-be\sqrt{\frac{1-cx}{1+cx}}(1+cx)(d+ex^2) - 2ac^2(8d^2 + 4dex^2 - e^2x^4) - 2bc^2(8d^2 + 4dex^2 - e^2x^4)}{6c^2e^3\sqrt{d+ex^2}} - \frac{b\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\left(-9(-c^2)^{3/2}d\sqrt{-c^2d-e}\sqrt{e}\sqrt{\frac{c^2(d+ex^2)}{c^2d+e}}\arcsin\left(\frac{c\sqrt{e}\sqrt{1-c^2x^2}}{\sqrt{-c^2}\sqrt{-c^2d-e}}\right) + \sqrt{-c^2}\sqrt{-c^2d-e}e^{3/2}\right)}{6c^5e^3(-1+cx)\sqrt{d+ex^2}}$$

input

```
Integrate[(x^5*(a + b*ArcSech[c*x]))/(d + e*x^2)^(3/2), x]
```

output

```
(-(b*e*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(d + e*x^2)) - 2*a*c^2*(8*d^2 + 4*d*e*x^2 - e^2*x^4) - 2*b*c^2*(8*d^2 + 4*d*e*x^2 - e^2*x^4)*ArcSech[c*x]) / (6*c^2*e^3*Sqrt[d + e*x^2]) - (b*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*(-9*(-c^2)^(3/2)*d*Sqrt[-(c^2*d) - e]*Sqrt[e]*Sqrt[(c^2*(d + e*x^2))/(c^2*d + e)]*ArcSin[(c*Sqrt[e]*Sqrt[1 - c^2*x^2])/(Sqrt[-c^2]*Sqrt[-(c^2*d) - e])] + Sqrt[-c^2]*Sqrt[-(c^2*d) - e]*e^(3/2)*Sqrt[(c^2*(d + e*x^2))/(c^2*d + e)]*ArcSin[(Sqrt[-c^2]*Sqrt[e]*Sqrt[1 - c^2*x^2])/(c*Sqrt[-(c^2*d) - e])] + 16*c^5*d^(3/2)*Sqrt[-d - e*x^2]*ArcTan[(Sqrt[d]*Sqrt[1 - c^2*x^2])/Sqrt[-d - e*x^2]])) / (6*c^5*e^3*(-1 + c*x)*Sqrt[d + e*x^2])
```

**Rubi [A] (verified)**

Time = 1.35 (sec) , antiderivative size = 243, normalized size of antiderivative = 0.87, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.435$ , Rules used = {6855, 27, 7282, 2118, 27, 175, 66, 104, 218, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^5(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

↓ 6855

$$\begin{aligned}
& b\sqrt{\frac{1}{cx+1}\sqrt{cx+1}} \int -\frac{-e^2x^4+4dex^2+8d^2}{3e^3x\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx - \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} - \\
& \frac{2d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3} + \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} \\
& \quad \downarrow 27 \\
& \frac{b\sqrt{\frac{1}{cx+1}\sqrt{cx+1}} \int \frac{-e^2x^4+4dex^2+8d^2}{x\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx}{3e^3} - \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} - \\
& \frac{2d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3} + \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} \\
& \quad \downarrow 7282 \\
& \frac{b\sqrt{\frac{1}{cx+1}\sqrt{cx+1}} \int \frac{-e^2x^4+4dex^2+8d^2}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2}{6e^3} - \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} - \\
& \frac{2d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3} + \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} \\
& \quad \downarrow 2118 \\
& \frac{b\sqrt{\frac{1}{cx+1}\sqrt{cx+1}} \left( \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} - \frac{\int -\frac{e(16c^2d^2+(9c^2d-e)ex^2)}{2x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2}{c^2e} \right)}{6e^3} - \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} - \\
& \frac{2d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3} + \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} \\
& \quad \downarrow 27 \\
& \frac{b\sqrt{\frac{1}{cx+1}\sqrt{cx+1}} \left( \frac{\int \frac{16c^2d^2+(9c^2d-e)ex^2}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2}{2c^2} + \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} \right)}{6e^3} - \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} - \\
& \frac{2d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3} + \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} \\
& \quad \downarrow 175 \\
& \frac{b\sqrt{\frac{1}{cx+1}\sqrt{cx+1}} \left( \frac{16c^2d^2 \int \frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 + e(9c^2d-e) \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2}{2c^2} + \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} \right)}{6e^3} - \\
& \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} - \frac{2d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3} + \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} \\
& \quad \downarrow 66
\end{aligned}$$



$$\begin{aligned}
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{16c^2d^2 \int \frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 + 2e(9c^2d-e) \int \frac{1}{-ex^4-c^2} d \frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}} + \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} \right)}{6e^3} \\
 & \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} - \frac{2d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3} + \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} \\
 & \quad \downarrow 104 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{32c^2d^2 \int \frac{1}{x^4-d} d \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} + 2e(9c^2d-e) \int \frac{1}{-ex^4-c^2} d \frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}} + \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} \right)}{6e^3} \\
 & \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} - \frac{2d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3} + \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} \\
 & \quad \downarrow 218 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{32c^2d^2 \int \frac{1}{x^4-d} d \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} - \frac{2\sqrt{e}(9c^2d-e) \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c} + \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} \right)}{6e^3} \\
 & \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} - \frac{2d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3} + \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} \\
 & \quad \downarrow 220 \\
 & \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} - \frac{2d\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3} + \frac{(d+ex^2)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3e^3} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{2\sqrt{e}(9c^2d-e) \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c} - 32c^2d^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right) + \frac{e\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{c^2} \right)}{6e^3}
 \end{aligned}$$

input `Int[(x^5*(a + b*ArcSech[c*x]))/(d + e*x^2)^(3/2),x]`

output

$$-\left(\frac{d^2(a + b \operatorname{ArcSech}[c*x])}{e^3 \sqrt{d + e*x^2}}\right) - \left(\frac{2*d*\sqrt{d + e*x^2} * (a + b \operatorname{ArcSech}[c*x])}{e^3} + \frac{(d + e*x^2)^{3/2} * (a + b \operatorname{ArcSech}[c*x])}{3*e^3} - \frac{(b*\sqrt{(1 + c*x)^{-1}}*\sqrt{1 + c*x}*((e*\sqrt{1 - c^2*x^2})*\sqrt{d + e*x^2}))/c^2 + ((-2*(9*c^2*d - e)*\sqrt{e}*\operatorname{ArcTan}[(\sqrt{e}*\sqrt{1 - c^2*x^2}))/(\sqrt{d + e*x^2})])}{c} - 32*c^2*d^{3/2}*\operatorname{ArcTanh}[\sqrt{d + e*x^2}/(\sqrt{d}*\sqrt{1 - c^2*x^2})]}{(2*c^2)}\right)/(6*e^3)$$

### Defintions of rubi rules used

rule 27

$$\operatorname{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \operatorname{Simp}[a \operatorname{Int}[Fx, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{MatchQ}[Fx, (b_*)(Gx_)] /; \operatorname{FreeQ}[b, x]$$

rule 66

$$\operatorname{Int}[1/(\sqrt{(a_*) + (b_*)(x_*)}*\sqrt{(c_*) + (d_*)(x_*)}), x\_Symbol] \rightarrow \operatorname{Simp}[2 \operatorname{Subst}[\operatorname{Int}[1/(b - d*x^2), x], x, \sqrt{a + b*x}/\sqrt{c + d*x}], x] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ !\operatorname{GtQ}[c - a*(d/b), 0]$$

rule 104

$$\operatorname{Int}[(((a_*) + (b_*)(x_*)^m)*((c_*) + (d_*)(x_*)^n))/((e_*) + (f_*)(x_*)^q), x_] \rightarrow \operatorname{With}[\{q = \operatorname{Denominator}[m]\}, \operatorname{Simp}[q \operatorname{Subst}[\operatorname{Int}[x^{(q*(m+1) - 1)/(b*e - a*f - (d*e - c*f)*x^q}], x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x] /; \operatorname{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \operatorname{EqQ}[m + n + 1, 0] \ \&\& \ \operatorname{RationalQ}[n] \ \&\& \ \operatorname{LtQ}[-1, m, 0] \ \&\& \ \operatorname{SimplerQ}[a + b*x, c + d*x]$$

rule 175

$$\operatorname{Int}[(((c_*) + (d_*)(x_*)^n)*((e_*) + (f_*)(x_*)^p)*((g_*) + (h_*)(x_*)^q))/((a_*) + (b_*)(x_*)^r), x_] \rightarrow \operatorname{Simp}[h/b \operatorname{Int}[(c + d*x)^n*(e + f*x)^p, x], x] + \operatorname{Simp}[(b*g - a*h)/b \operatorname{Int}[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; \operatorname{FreeQ}[\{a, b, c, d, e, f, g, h, n, p\}, x]$$

rule 218

$$\operatorname{Int}[((a_*) + (b_*)(x_*)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[a/b, 2]/a)*\operatorname{ArcTan}[x/\operatorname{Rt}[a/b, 2]], x] /; \operatorname{FreeQ}[\{a, b\}, x] \ \&\& \ \operatorname{PosQ}[a/b]$$

rule 220

$$\operatorname{Int}[((a_*) + (b_*)(x_*)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(-\operatorname{Rt}[-a, 2]*\operatorname{Rt}[b, 2])^{-1})*\operatorname{ArcTanh}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[-a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \ \&\& \ \operatorname{NegQ}[a/b] \ \&\& \ (\operatorname{LtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$$

rule 2118

```
Int[(Px_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*
(x_))^(p_), x_Symbol] := With[{q = Expon[Px, x], k = Coeff[Px, x, Expo
n[Px, x]]}, Simp[k*(a + b*x)^(m + q - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p +
1)/(d*f*b^(q - 1)*(m + n + p + q + 1))), x] + Simp[1/(d*f*b^q*(m + n + p +
q + 1)) Int[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p*ExpandToSum[d*f*b^q*(m +
n + p + q + 1)*Px - d*f*k*(m + n + p + q + 1)*(a + b*x)^q + k*(a + b*x)^(q
- 2)*(a^2*d*f*(m + n + p + q + 1) - b*(b*c*e*(m + q - 1) + a*(d*e*(n + 1) +
c*f*(p + 1))) + b*(a*d*f*(2*(m + q) + n + p) - b*(d*e*(m + q + n) + c*f*(m
+ q + p))*x), x], x], x] /; NeQ[m + n + p + q + 1, 0] /; FreeQ[{a, b, c,
d, e, f, m, n, p}, x] && PolyQ[Px, x]
```

rule 6855

```
Int[((a_) + ArcSech[(c_)*(x_)]*(b_))*((f_)*(x_))^(m_)*((d_) + (e_)*(
x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Si
mp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)]
Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x]] /; Fre
eQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] &&
GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2
*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

rule 7282

```
Int[(u_)/(x_), x_Symbol] := With[{lst = PowerVariableExpn[u, 0, x]}, Simp[1
/lst[[2]] Subst[Int[NormalizeIntegrand[Simplify[lst[[1]]/x], x], x], x, (
lst[[3]]*x)^lst[[2]]], x] /; !FalseQ[lst] && NeQ[lst[[2]], 0] /; NonsumQ[
u] && !RationalFunctionQ[u, x]
```

## Maple [F]

$$\int \frac{x^5(a + b \operatorname{arcsech}(cx))}{(x^2e + d)^{\frac{3}{2}}} dx$$

input

```
int(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x)
```

output

```
int(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 425 vs.  $2(184) = 368$ .

Time = 0.33 (sec) , antiderivative size = 1771, normalized size of antiderivative = 6.37

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \text{Too large to display}$$

input `integrate(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="fricas")`

output

```
[1/24*((9*b*c^2*d^2 - b*d*e + (9*b*c^2*d*e - b*e^2)*x^2)*sqrt(-e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e + 8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^4*e*x^3 + (c^4*d - c^2*e)*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e^2) + 8*(b*c^3*e^2*x^4 - 4*b*c^3*d*e*x^2 - 8*b*c^3*d^2)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 16*(b*c^3*d*e*x^2 + b*c^3*d^2)*sqrt(d)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 - 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) + 4*(2*a*c^3*e^2*x^4 - 8*a*c^3*d*e*x^2 - 16*a*c^3*d^2 - (b*c^2*e^2*x^3 + b*c^2*d*e*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d)/(c^3*e^4*x^2 + c^3*d*e^3), 1/12*((9*b*c^2*d^2 - b*d*e + (9*b*c^2*d*e - b*e^2)*x^2)*sqrt(e)*arctan(1/2*(2*c^2*e*x^3 + (c^2*d - e)*x)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*e^2*x^4 + (c^2*d*e - e^2)*x^2 - d*e) + 4*(b*c^3*e^2*x^4 - 4*b*c^3*d*e*x^2 - 8*b*c^3*d^2)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 8*(b*c^3*d*e*x^2 + b*c^3*d^2)*sqrt(d)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 - 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) + 2*(2*a*c^3*e^2*x^4 - 8*a*c^3*d*e*x^2 - 16*a*c^3*d^2 - (b*c^2*e^2*x^3 + b*c^2*d*e*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d)/(c^3*e^4*x^2 + c^3*d*e^3), 1/24*(32*(b*c^3*d*e*x^2 + b*c^3*d^2)*sqrt(-d)*arctan(-1/2*((c^3*d ...
```

**Sympy [F]**

$$\int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x^5(a + b \operatorname{arsech}(cx))}{(d + ex^2)^{\frac{3}{2}}} dx$$

input `integrate(x**5*(a+b*asech(c*x))/(e*x**2+d)**(3/2),x)`

output `Integral(x**5*(a + b*asech(c*x))/(d + e*x**2)**(3/2), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^5}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^5/(e*x^2 + d)^(3/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^5(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x^5(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^{3/2}} dx$$

input `int((x^5*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(3/2), x)`

output `int((x^5*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(3/2), x)`

**Reduce [F]**

$$\int \frac{x^5(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x^5(a \operatorname{sech}(cx) b + a)}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `int(x^5*(a+b*asech(c*x))/(e*x^2+d)^(3/2), x)`

output `int(x^5*(a+b*asech(c*x))/(e*x^2+d)^(3/2), x)`

**3.159** 
$$\int \frac{x^3 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

Optimal result	1278
Mathematica [A] (verified)	1278
Rubi [A] (verified)	1279
Maple [F]	1282
Fricas [B] (verification not implemented)	1282
Sympy [F]	1283
Maxima [F(-2)]	1284
Giac [F]	1284
Mupad [F(-1)]	1284
Reduce [F]	1285

**Optimal result**

Integrand size = 23, antiderivative size = 177

$$\int \frac{x^3 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \frac{d(a + b \operatorname{sech}^{-1}(cx))}{e^2 \sqrt{d + ex^2}} + \frac{\sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx))}{e^2} - \frac{b \sqrt{\frac{1}{1+cx}} \sqrt{1 + cx} \arctan\left(\frac{\sqrt{e} \sqrt{1 - c^2 x^2}}{c \sqrt{d + ex^2}}\right)}{ce^{3/2}} - \frac{2b \sqrt{d} \sqrt{\frac{1}{1+cx}} \sqrt{1 + cx} \operatorname{arctanh}\left(\frac{\sqrt{d + ex^2}}{\sqrt{d} \sqrt{1 - c^2 x^2}}\right)}{e^2}$$

output

```
d*(a+b*arcsech(c*x))/e^2/(e*x^2+d)^(1/2)+(e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/e^2-b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctan(e^(1/2)*(-c^2*x^2+1)^(1/2)/c/(e*x^2+d)^(1/2))/c/e^(3/2)-2*b*d^(1/2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh((e*x^2+d)^(1/2)/d^(1/2)/(-c^2*x^2+1)^(1/2))/e^2
```

**Mathematica [A] (verified)**

Time = 21.97 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.41

$$\int \frac{x^3 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \frac{(2d + ex^2) (a + b \operatorname{sech}^{-1}(cx))}{e^2 \sqrt{d + ex^2}} + \frac{b \sqrt{\frac{1-cx}{1+cx}} \sqrt{1 - c^2 x^2} \left( \sqrt{-c^2} \sqrt{-c^2 d - e} \sqrt{e} \sqrt{\frac{c^2 (d + ex^2)}{c^2 d + e}} \arcsin\left(\frac{c \sqrt{e} \sqrt{1 - c^2 x^2}}{\sqrt{-c^2} \sqrt{-c^2 d - e}}\right) + 2c^3 \sqrt{d} \sqrt{-d - ex^2} \arctan\left(\frac{\sqrt{d + ex^2}}{\sqrt{d} \sqrt{1 - c^2 x^2}}\right) \right)}{c^3 e^2 (-1 + cx) \sqrt{d + ex^2}}$$

input `Integrate[(x^3*(a + b*ArcSech[c*x]))/(d + e*x^2)^(3/2),x]`

output `((2*d + e*x^2)*(a + b*ArcSech[c*x]))/(e^2*Sqrt[d + e*x^2]) + (b*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*(Sqrt[-c^2]*Sqrt[-(c^2*d) - e]*Sqrt[e]*Sqrt[(c^2*(d + e*x^2))/(c^2*d + e)]*ArcSin[(c*Sqrt[e]*Sqrt[1 - c^2*x^2])/(Sqrt[-c^2]*Sqrt[-(c^2*d) - e]]) + 2*c^3*Sqrt[d]*Sqrt[-d - e*x^2]*ArcTan[(Sqrt[d]*Sqrt[1 - c^2*x^2])/Sqrt[-d - e*x^2]])/(c^3*e^2*(-1 + c*x)*Sqrt[d + e*x^2])`

### Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.91, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.348$ , Rules used = {6855, 27, 435, 175, 66, 104, 218, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx \\
 & \quad \downarrow \text{6855} \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{ex^2 + 2d}{e^2x\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx + \frac{\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{e^2} + \\
 & \quad \frac{d(a + b\operatorname{sech}^{-1}(cx))}{e^2\sqrt{d+ex^2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{ex^2+2d}{x\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx}{e^2} + \frac{\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{e^2} + \frac{d(a + b\operatorname{sech}^{-1}(cx))}{e^2\sqrt{d+ex^2}} \\
 & \quad \downarrow \text{435} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{ex^2+2d}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2}{2e^2} + \frac{\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{e^2} + \frac{d(a + b\operatorname{sech}^{-1}(cx))}{e^2\sqrt{d+ex^2}} \\
 & \quad \downarrow \text{175}
 \end{aligned}$$



$$\begin{aligned}
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(e\int\frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2+2d\int\frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2\right)}{2e^2} + \\
& \quad \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^2} + \frac{d(a+b\operatorname{sech}^{-1}(cx))}{e^2\sqrt{d+ex^2}} \\
& \quad \downarrow 66 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(2d\int\frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2+2e\int\frac{1}{-ex^4-c^2}d\frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}}\right)}{2e^2} + \\
& \quad \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^2} + \frac{d(a+b\operatorname{sech}^{-1}(cx))}{e^2\sqrt{d+ex^2}} \\
& \quad \downarrow 104 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(4d\int\frac{1}{x^4-d}d\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}+2e\int\frac{1}{-ex^4-c^2}d\frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}}\right)}{2e^2} + \\
& \quad \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^2} + \frac{d(a+b\operatorname{sech}^{-1}(cx))}{e^2\sqrt{d+ex^2}} \\
& \quad \downarrow 218 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(4d\int\frac{1}{x^4-d}d\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}-\frac{2\sqrt{e}\arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c}\right)}{2e^2} + \\
& \quad \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^2} + \frac{d(a+b\operatorname{sech}^{-1}(cx))}{e^2\sqrt{d+ex^2}} \\
& \quad \downarrow 220 \\
& \frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^2} + \frac{d(a+b\operatorname{sech}^{-1}(cx))}{e^2\sqrt{d+ex^2}} + \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{2\sqrt{e}\arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c}-4\sqrt{d}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)\right)}{2e^2}
\end{aligned}$$

input `Int[(x^3*(a + b*ArcSech[c*x]))/(d + e*x^2)^(3/2),x]`

output `(d*(a + b*ArcSech[c*x]))/(e^2*Sqrt[d + e*x^2]) + (Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/e^2 + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-2*Sqrt[e]*ArcTan[(Sqrt[e]*Sqrt[1 - c^2*x^2])/(c*Sqrt[d + e*x^2])])/c - 4*Sqrt[d]*ArcTanh[Sqrt[d + e*x^2]/(Sqrt[d]*Sqrt[1 - c^2*x^2])]))/(2*e^2)`

## Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 66 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`
- rule 104 `Int[(((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 175 `Int[(((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_)*((g_) + (h_)*(x_)))/((a_) + (b_)*(x_)), x_] := Simp[h/b Int[(c + d*x)^n*(e + f*x)^p, x], x] + Simp[(b*g - a*h)/b Int[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 220 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`
- rule 435 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2)^(r_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q*(e + f*x)^r, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, f, p, q, r}, x] && IntegerQ[(m - 1)/2]`

rule 6855

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))*((f_.)*(x_.))^(m_.)*((d_.) + (e_.)*(
x_)^2)^(p_.), x_Symbol] :> With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Si
mp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)]
Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x]] /; Fre
eQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] &&
GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2
*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

**Maple [F]**

$$\int \frac{x^3(a + b \operatorname{arcsech}(cx))}{(x^2e + d)^{\frac{3}{2}}} dx$$

input

```
int(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x)
```

output

```
int(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 310 vs.  $2(117) = 234$ .

Time = 0.25 (sec) , antiderivative size = 1311, normalized size of antiderivative = 7.41

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \text{Too large to display}$$

input

```
integrate(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="fricas")
```

output

```

[-1/4*((b*e*x^2 + b*d)*sqrt(-e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e +
8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^4*e*x^3 + (c^4*d - c^2*e)*x)*sqrt(e*x^2
+ d)*sqrt(-e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e^2) - 4*(b*c*e*x^2 + 2*b*
c*d)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) -
2*(b*c*e*x^2 + b*c*d)*sqrt(d)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c
^2*d^2 - d*e)*x^2 + 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d
)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) - 4*(a*c*e*x^2 + 2*a*c*d)*s
qrt(e*x^2 + d)/(c*e^3*x^2 + c*d*e^2), -1/2*((b*e*x^2 + b*d)*sqrt(e)*arcta
n(1/2*(2*c^2*e*x^3 + (c^2*d - e)*x)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(-(c^2*x^2
- 1)/(c^2*x^2)))/(c^2*e^2*x^4 + (c^2*d*e - e^2)*x^2 - d*e)) - 2*(b*c*e*x^2
+ 2*b*c*d)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(
c*x)) - (b*c*e*x^2 + b*c*d)*sqrt(d)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 -
8*(c^2*d^2 - d*e)*x^2 + 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*s
qrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) - 2*(a*c*e*x^2 + 2*a*c
*d)*sqrt(e*x^2 + d)/(c*e^3*x^2 + c*d*e^2), -1/4*(4*(b*c*e*x^2 + b*c*d)*sq
rt(-d)*arctan(-1/2*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(-d)*
sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*d*e*x^4 + (c^2*d^2 - d*e)*x^2 - d^2))
+ (b*e*x^2 + b*d)*sqrt(-e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e + 8*(c
^4*d*e - c^2*e^2)*x^2 - 4*(2*c^4*e*x^3 + (c^4*d - c^2*e)*x)*sqrt(e*x^2 + d)
*sqrt(-e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e^2) - 4*(b*c*e*x^2 + 2*b*c*...

```

## Sympy [F]

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x^3(a + b \operatorname{asech}(cx))}{(d + ex^2)^{3/2}} dx$$

input

```
integrate(x**3*(a+b*asech(c*x))/(e*x**2+d)**(3/2),x)
```

output

```
Integral(x**3*(a + b*asech(c*x))/(d + e*x**2)**(3/2), x)
```

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^3}{(ex^2 + d)^{3/2}} dx$$

input `integrate(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^3/(e*x^2 + d)^(3/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^{3/2}} dx$$

input `int((x^3*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(3/2),x)`

output `int((x^3*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(3/2), x)`

**Reduce [F]**

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x^3(a \operatorname{sech}(cx) b + a)}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `int(x^3*(a+b*asech(c*x))/(e*x^2+d)^(3/2),x)`

output `int(x^3*(a+b*asech(c*x))/(e*x^2+d)^(3/2),x)`

**3.160** 
$$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{3/2}} dx$$

Optimal result	1286
Mathematica [A] (verified)	1286
Rubi [A] (verified)	1287
Maple [F]	1289
Fricas [B] (verification not implemented)	1289
Sympy [F]	1290
Maxima [F]	1290
Giac [F]	1291
Mupad [F(-1)]	1291
Reduce [F]	1291

**Optimal result**

Integrand size = 21, antiderivative size = 87

$$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{3/2}} dx = -\frac{a+b\operatorname{sech}^{-1}(cx)}{e\sqrt{d+ex^2}} + \frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{\sqrt{de}}$$

output

```
-(a+b*arcsech(c*x))/e/(e*x^2+d)^(1/2)+b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh((e*x^2+d)^(1/2)/d^(1/2)/(-c^2*x^2+1)^(1/2))/d^(1/2)/e
```

**Mathematica [A] (verified)**

Time = 19.88 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.55

$$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{3/2}} dx = -\frac{a+b\operatorname{sech}^{-1}(cx)}{e\sqrt{d+ex^2}} - \frac{b\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\sqrt{-d-ex^2}\arctan\left(\frac{\sqrt{d}\sqrt{1-c^2x^2}}{\sqrt{-d-ex^2}}\right)}{\sqrt{de}(-1+cx)\sqrt{d+ex^2}}$$

input

```
Integrate[(x*(a + b*ArcSech[c*x]))/(d + e*x^2)^(3/2),x]
```

output

$$-\left(\frac{a + b \operatorname{ArcSech}[c x]}{e \sqrt{d + e x^2}}\right) - \left(\frac{b \sqrt{\frac{1 - c x}{1 + c x}} \sqrt{1 - c^2 x^2} \sqrt{-d - e x^2} \operatorname{ArcTan}\left[\frac{\sqrt{d} \sqrt{1 - c^2 x^2}}{\sqrt{-d - e x^2}}\right]}{\sqrt{d} e (-1 + c x) \sqrt{d + e x^2}}\right)$$
**Rubi [A] (verified)**

Time = 0.46 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {6853, 2036, 354, 104, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

↓ 6853

$$-\frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{1}{x \sqrt{1-cx} \sqrt{cx+1} \sqrt{ex^2+d}} dx}{e} - \frac{a + b \operatorname{sech}^{-1}(cx)}{e \sqrt{d + ex^2}}$$

↓ 2036

$$-\frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{1}{x \sqrt{1-c^2x^2} \sqrt{ex^2+d}} dx}{e} - \frac{a + b \operatorname{sech}^{-1}(cx)}{e \sqrt{d + ex^2}}$$

↓ 354

$$-\frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{1}{x^2 \sqrt{1-c^2x^2} \sqrt{ex^2+d}} dx^2}{2e} - \frac{a + b \operatorname{sech}^{-1}(cx)}{e \sqrt{d + ex^2}}$$

↓ 104

$$-\frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \int \frac{1}{x^4-d} d \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}}{e} - \frac{a + b \operatorname{sech}^{-1}(cx)}{e \sqrt{d + ex^2}}$$

↓ 220

$$\frac{b \sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d} \sqrt{1-c^2x^2}}\right)}{\sqrt{de}} - \frac{a + b \operatorname{sech}^{-1}(cx)}{e \sqrt{d + ex^2}}$$



input `Int[(x*(a + b*ArcSech[c*x]))/(d + e*x^2)^(3/2),x]`

output `-((a + b*ArcSech[c*x])/(e*Sqrt[d + e*x^2])) + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*ArcTanh[Sqrt[d + e*x^2]/(Sqrt[d]*Sqrt[1 - c^2*x^2])])/(Sqrt[d]*e)`

### Defintions of rubi rules used

rule 104 `Int[(((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 220 `Int[(((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 354 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]`

rule 2036 `Int[(u_.)*((c_) + (d_.)*(x_)^(n_.))^(q_.)*((a1_) + (b1_.)*(x_)^(non2_.))^(p_.)*((a2_) + (b2_.)*(x_)^(non2_.))^(p_.), x_Symbol] := Int[u*(a1*a2 + b1*b2*x^n)^p*(c + d*x^n)^q, x] /; FreeQ[{a1, b1, a2, b2, c, d, n, p, q}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))`

rule 6853

```
Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))*(x_.)*((d_.) + (e_.)*(x_)^2)^(p_.),
x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcSech[c*x])/(2*e*(p + 1))),
x] + Simp[b*(Sqrt[1 + c*x]/(2*e*(p + 1)))*Sqrt[1/(1 + c*x)] Int[(d + e*x
^2)^(p + 1)/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x] /; FreeQ[{a, b, c, d, e
, p}, x] && NeQ[p, -1]
```

**Maple [F]**

$$\int \frac{x(a + b \operatorname{arcsech}(cx))}{(x^2e + d)^{\frac{3}{2}}} dx$$

input

```
int(x*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x)
```

output

```
int(x*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 183 vs. 2(57) = 114.

Time = 0.15 (sec) , antiderivative size = 379, normalized size of antiderivative = 4.36

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \frac{4\sqrt{ex^2 + d}bd \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}+1}}{cx}\right) + 4\sqrt{ex^2 + d}ad - (bex^2 + bd)\sqrt{d} \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}+1}}{cx}\right) + 2\sqrt{ex^2 + d}bd \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}+1}}{cx}\right) + 2\sqrt{ex^2 + d}ad - (bex^2 + bd)\sqrt{-d} \arctan\left(-\frac{((c^3d-ce)x^3-2cdx)\sqrt{ex^2+d}\sqrt{-d}}{2(c^2dex^4+(c^2d^2-de)x}\right)}{4(de^2x^2 + d^2e)}$$

input

```
integrate(x*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="fricas")
```

output

```
[-1/4*(4*sqrt(e*x^2 + d)*b*d*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 4*sqrt(e*x^2 + d)*a*d - (b*e*x^2 + b*d)*sqrt(d)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 - 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4))/(d*e^2*x^2 + d^2*e), -1/2*(2*sqrt(e*x^2 + d)*b*d*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + 2*sqrt(e*x^2 + d)*a*d - (b*e*x^2 + b*d)*sqrt(-d)*arctan(-1/2*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(-d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*d*e*x^4 + (c^2*d^2 - d*e)*x^2 - d^2)))/(d*e^2*x^2 + d^2*e)]
```

**Sympy [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x(a + b \operatorname{asech}(cx))}{(d + ex^2)^{\frac{3}{2}}} dx$$

input

```
integrate(x*(a+b*asech(c*x))/(e*x**2+d)**(3/2),x)
```

output

```
Integral(x*(a + b*asech(c*x))/(d + e*x**2)**(3/2), x)
```

**Maxima [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input

```
integrate(x*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="maxima")
```

output

```
b*integrate(x*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(e*x^2 + d)^(3/2), x) - a/(sqrt(e*x^2 + d)*e)
```

**Giac [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x/(e*x^2 + d)^(3/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^{3/2}} dx$$

input `int((x*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(3/2),x)`

output `int((x*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(3/2), x)`

**Reduce [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x(a \operatorname{sech}(cx) b + a)}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `int(x*(a+b*asech(c*x))/(e*x^2+d)^(3/2),x)`

output `int(x*(a+b*asech(c*x))/(e*x^2+d)^(3/2),x)`

$$3.161 \quad \int \frac{a+b\operatorname{sech}^{-1}(cx)}{x(d+ex^2)^{3/2}} dx$$

Optimal result	1292
Mathematica [N/A]	1292
Rubi [N/A]	1293
Maple [N/A]	1293
Fricas [N/A]	1294
Sympy [N/A]	1294
Maxima [F(-2)]	1294
Giac [N/A]	1295
Mupad [N/A]	1295
Reduce [N/A]	1296

### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{3/2}} dx = \operatorname{Int}\left(\frac{a + b\operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{3/2}}, x\right)$$

output `Defer(Int)((a+b*arcsech(c*x))/x/(e*x^2+d)^(3/2),x)`

### Mathematica [N/A]

Not integrable

Time = 7.97 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{3/2}} dx = \int \frac{a + b\operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{3/2}} dx$$

input `Integrate[(a + b*ArcSech[c*x])/(x*(d + e*x^2)^(3/2)),x]`

output `Integrate[(a + b*ArcSech[c*x])/(x*(d + e*x^2)^(3/2)), x]`

**Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x (d + ex^2)^{3/2}} dx$$

↓ 6865

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x (d + ex^2)^{3/2}} dx$$

input `Int[(a + b*ArcSech[c*x])/(x*(d + e*x^2)^(3/2)),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.20 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{a + b \operatorname{arcsech}(cx)}{x (x^2 e + d)^{\frac{3}{2}}} dx$$

input `int((a+b*arcsech(c*x))/x/(e*x^2+d)^(3/2),x)`

output `int((a+b*arcsech(c*x))/x/(e*x^2+d)^(3/2),x)`

**Fricas [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.83

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{3/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^{\frac{3}{2}} x} dx$$

input `integrate((a+b*arcsech(c*x))/x/(e*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)/(e^2*x^5 + 2*d*e*x^3 + d^2*x), x)`

**Sympy [N/A]**

Not integrable

Time = 23.17 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{3/2}} dx = \int \frac{a + b \operatorname{asech}(cx)}{x(d + ex^2)^{\frac{3}{2}}} dx$$

input `integrate((a+b*asech(c*x))/x/(e*x**2+d)**(3/2),x)`

output `Integral((a + b*asech(c*x))/(x*(d + e*x**2)**(3/2)), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/x/(e*x^2+d)^(3/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{3/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^{\frac{3}{2}} x} dx$$

input

```
integrate((a+b*arcsech(c*x))/x/(e*x^2+d)^(3/2),x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)/((e*x^2 + d)^(3/2)*x), x)
```

**Mupad [N/A]**

Not integrable

Time = 4.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x(e x^2 + d)^{3/2}} dx$$

input

```
int((a + b*acosh(1/(c*x)))/(x*(d + e*x^2)^(3/2)),x)
```

output

```
int((a + b*acosh(1/(c*x)))/(x*(d + e*x^2)^(3/2)), x)
```



**Reduce [N/A]**

Not integrable

Time = 200.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x (d + ex^2)^{3/2}} dx = \int \frac{a \operatorname{sech}(cx) b + a}{x (ex^2 + d)^{\frac{3}{2}}} dx$$

input `int((a+b*asech(c*x))/x/(e*x^2+d)^(3/2),x)`output `int((a+b*asech(c*x))/x/(e*x^2+d)^(3/2),x)`

$$3.162 \quad \int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^3(d+ex^2)^{3/2}} dx$$

Optimal result	1297
Mathematica [N/A]	1297
Rubi [N/A]	1298
Maple [N/A]	1298
Fricas [N/A]	1299
Sympy [N/A]	1299
Maxima [F(-2)]	1299
Giac [N/A]	1300
Mupad [N/A]	1300
Reduce [N/A]	1301

### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^3(d + ex^2)^{3/2}} dx = \operatorname{Int}\left(\frac{a + b\operatorname{sech}^{-1}(cx)}{x^3(d + ex^2)^{3/2}}, x\right)$$

output `Defer(Int)((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(3/2),x)`

### Mathematica [N/A]

Not integrable

Time = 10.51 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^3(d + ex^2)^{3/2}} dx = \int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^3(d + ex^2)^{3/2}} dx$$

input `Integrate[(a + b*ArcSech[c*x])/(x^3*(d + e*x^2)^(3/2)),x]`

output `Integrate[(a + b*ArcSech[c*x])/(x^3*(d + e*x^2)^(3/2)), x]`

**Rubi [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{3/2}} dx$$

↓ 6865

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{3/2}} dx$$

input `Int[(a + b*ArcSech[c*x])/(x^3*(d + e*x^2)^(3/2)),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.38 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{a + b \operatorname{arcsech}(cx)}{x^3 (x^2 e + d)^{3/2}} dx$$

input `int((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(3/2),x)`

output `int((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(3/2),x)`

**Fricas [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.91

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{3/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^{\frac{3}{2}} x^3} dx$$

input `integrate((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)/(e^2*x^7 + 2*d*e*x^5 + d^2*x^3), x)`

**Sympy [N/A]**

Not integrable

Time = 97.82 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{3/2}} dx = \int \frac{a + b \operatorname{asech}(cx)}{x^3 (d + ex^2)^{\frac{3}{2}}} dx$$

input `integrate((a+b*asech(c*x))/x**3/(e*x**2+d)**(3/2),x)`

output `Integral((a + b*asech(c*x))/(x**3*(d + e*x**2)**(3/2)), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(3/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{3/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^{\frac{3}{2}} x^3} dx$$

input

```
integrate((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(3/2),x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)/((e*x^2 + d)^(3/2)*x^3), x)
```

**Mupad [N/A]**

Not integrable

Time = 4.27 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x^3 (ex^2 + d)^{3/2}} dx$$

input

```
int((a + b*acosh(1/(c*x)))/(x^3*(d + e*x^2)^(3/2)),x)
```

output

```
int((a + b*acosh(1/(c*x)))/(x^3*(d + e*x^2)^(3/2)), x)
```

**Reduce [N/A]**

Not integrable

Time = 200.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{3/2}} dx = \int \frac{a \operatorname{sech}(cx) b + a}{x^3 (ex^2 + d)^{\frac{3}{2}}} dx$$

input `int((a+b*asech(c*x))/x^3/(e*x^2+d)^(3/2),x)`output `int((a+b*asech(c*x))/x^3/(e*x^2+d)^(3/2),x)`

$$3.163 \quad \int \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

Optimal result	1302
Mathematica [N/A]	1302
Rubi [N/A]	1303
Maple [N/A]	1303
Fricas [N/A]	1304
Sympy [N/A]	1304
Maxima [F(-2)]	1304
Giac [N/A]	1305
Mupad [N/A]	1305
Reduce [N/A]	1306

### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \operatorname{Int} \left( \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}}, x \right)$$

output `Defer(Int)(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x)`

### Mathematica [N/A]

Not integrable

Time = 14.14 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

input `Integrate[(x^4*(a + b*ArcSech[c*x]))/(d + e*x^2)^(3/2),x]`

output `Integrate[(x^4*(a + b*ArcSech[c*x]))/(d + e*x^2)^(3/2), x]`

**Rubi [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

↓ 6865

$$\int \frac{x^4(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

input `Int[(x^4*(a + b*ArcSech[c*x]))/(d + e*x^2)^(3/2),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.20 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{x^4(a + b \operatorname{arcsech}(cx))}{(x^2e + d)^{\frac{3}{2}}} dx$$

input `int(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x)`

output `int(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x)`



**Fricas [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.04

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^4}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral((b*x^4*arcsech(c*x) + a*x^4)*sqrt(e*x^2 + d)/(e^2*x^4 + 2*d*e*x^2 + d^2), x)`

**Sympy [N/A]**

Not integrable

Time = 47.52 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x^4(a + b \operatorname{asech}(cx))}{(d + ex^2)^{\frac{3}{2}}} dx$$

input `integrate(x**4*(a+b*asech(c*x))/(e*x**2+d)**(3/2),x)`

output `Integral(x**4*(a + b*asech(c*x))/(d + e*x**2)**(3/2), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{x^4(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^4}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input

```
integrate(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)*x^4/(e*x^2 + d)^(3/2), x)
```

**Mupad [N/A]**

Not integrable

Time = 4.18 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{x^4(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x^4(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^{3/2}} dx$$

input

```
int((x^4*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(3/2),x)
```

output

```
int((x^4*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(3/2), x)
```

**Reduce [N/A]**

Not integrable

Time = 200.04 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x^4(\operatorname{asech}(cx) b + a)}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `int(x^4*(a+b*asech(c*x))/(e*x^2+d)^(3/2),x)`output `int(x^4*(a+b*asech(c*x))/(e*x^2+d)^(3/2),x)`

$$3.164 \quad \int \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

Optimal result	1307
Mathematica [N/A]	1307
Rubi [N/A]	1308
Maple [N/A]	1308
Fricas [N/A]	1309
Sympy [N/A]	1309
Maxima [F(-2)]	1309
Giac [N/A]	1310
Mupad [N/A]	1310
Reduce [N/A]	1311

### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \operatorname{Int} \left( \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}}, x \right)$$

output `Defer(Int)(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x)`

### Mathematica [N/A]

Not integrable

Time = 5.71 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

input `Integrate[(x^2*(a + b*ArcSech[c*x]))/(d + e*x^2)^(3/2),x]`

output `Integrate[(x^2*(a + b*ArcSech[c*x]))/(d + e*x^2)^(3/2), x]`

**Rubi [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

↓ 6865

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

input `Int[(x^2*(a + b*ArcSech[c*x]))/(d + e*x^2)^(3/2),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{x^2(a + b \operatorname{arcsech}(cx))}{(x^2e + d)^{\frac{3}{2}}} dx$$

input `int(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x)`

output `int(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x)`

**Fricas [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.04

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^2}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral((b*x^2*arcsech(c*x) + a*x^2)*sqrt(e*x^2 + d)/(e^2*x^4 + 2*d*e*x^2 + d^2), x)`

**Sympy [N/A]**

Not integrable

Time = 11.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x^2(a + b \operatorname{asech}(cx))}{(d + ex^2)^{\frac{3}{2}}} dx$$

input `integrate(x**2*(a+b*asech(c*x))/(e*x**2+d)**(3/2),x)`

output `Integral(x**2*(a + b*asech(c*x))/(d + e*x**2)**(3/2), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^2}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input

```
integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)*x^2/(e*x^2 + d)^(3/2), x)
```

**Mupad [N/A]**

Not integrable

Time = 3.82 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^{3/2}} dx$$

input

```
int((x^2*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(3/2),x)
```

output

```
int((x^2*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(3/2), x)
```

**Reduce [N/A]**

Not integrable

Time = 200.05 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{x^2(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{x^2(\operatorname{asech}(cx) b + a)}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `int(x^2*(a+b*asech(c*x))/(e*x^2+d)^(3/2),x)`output `int(x^2*(a+b*asech(c*x))/(e*x^2+d)^(3/2),x)`



**3.165**  $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex^2)^{3/2}} dx$

Optimal result	1312
Mathematica [C] (verified)	1313
Rubi [A] (verified)	1313
Maple [F]	1315
Fricas [A] (verification not implemented)	1315
Sympy [F]	1316
Maxima [F]	1316
Giac [F]	1316
Mupad [F(-1)]	1317
Reduce [F]	1317

**Optimal result**

Integrand size = 20, antiderivative size = 92

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{(d + ex^2)^{3/2}} dx = \frac{x(a + b\operatorname{sech}^{-1}(cx))}{d\sqrt{d + ex^2}} + \frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1 + cx}\sqrt{1 + \frac{ex^2}{d}} \operatorname{EllipticF}(\arcsin(cx), -\frac{e}{c^2d})}{cd\sqrt{d + ex^2}}$$

output

```
x*(a+b*arcsech(c*x))/d/(e*x^2+d)^(1/2)+b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(1+e*x^2/d)^(1/2)*EllipticF(c*x,(-e/c^2/d)^(1/2))/c/d/(e*x^2+d)^(1/2)
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 48.59 (sec) , antiderivative size = 334, normalized size of antiderivative = 3.63

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{3/2}} dx = \frac{x(a + b \operatorname{sech}^{-1}(cx))}{d\sqrt{d + ex^2}} + \frac{2ib\sqrt{\frac{1-cx}{1+cx}} \sqrt{\frac{(c\sqrt{d}+i\sqrt{e})(1+cx)}{(c\sqrt{d}-i\sqrt{e})(-1+cx)}}} (-i\sqrt{d} + \sqrt{ex}) \sqrt{-\frac{-1+\frac{i\sqrt{ex}}{\sqrt{d}}+c\left(\frac{i\sqrt{d}}{\sqrt{e}}+x\right)}{1-cx}}} \operatorname{EllipticF}\left(\arcsin\left(\sqrt{\frac{1+\frac{ic\sqrt{d}}{\sqrt{e}}-cx+\frac{i\sqrt{ex}}{\sqrt{d}}}{2-2cx}}}\right)\right)}{d(c\sqrt{d} + i\sqrt{e}) \sqrt{\frac{1+\frac{ic\sqrt{d}}{\sqrt{e}}-cx+\frac{i\sqrt{ex}}{\sqrt{d}}}{1-cx}}} \sqrt{d + ex^2}}$$

input `Integrate[(a + b*ArcSech[c*x])/(d + e*x^2)^(3/2),x]`

output `(x*(a + b*ArcSech[c*x]))/(d*Sqrt[d + e*x^2]) + ((2*I)*b*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[((c*Sqrt[d] + I*Sqrt[e])*(1 + c*x))/((c*Sqrt[d] - I*Sqrt[e])*(-1 + c*x))]*((-I)*Sqrt[d] + Sqrt[e]*x)*Sqrt[-((-1 + (I*Sqrt[e]*x)/Sqrt[d] + c*((I*Sqrt[d])/Sqrt[e] + x))/(1 - c*x))]*EllipticF[ArcSin[Sqrt[(1 + (I*c*Sqrt[d])/Sqrt[e] - c*x + (I*Sqrt[e]*x)/Sqrt[d])/(2 - 2*c*x)]], ((-4*I)*c*Sqrt[d]*Sqrt[e])/(c*Sqrt[d] - I*Sqrt[e]^2)]/(d*(c*Sqrt[d] + I*Sqrt[e])*Sqrt[(1 + (I*c*Sqrt[d])/Sqrt[e] - c*x + (I*Sqrt[e]*x)/Sqrt[d])/(1 - c*x)]*Sqrt[d + e*x^2])`

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {6845, 27, 323, 321}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{3/2}} dx$$

↓ 6845

$$\begin{aligned}
& b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{d\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx + \frac{x(a+b\operatorname{sech}^{-1}(cx))}{d\sqrt{d+ex^2}} \\
& \quad \downarrow 27 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx}{d} + \frac{x(a+b\operatorname{sech}^{-1}(cx))}{d\sqrt{d+ex^2}} \\
& \quad \downarrow 323 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\sqrt{\frac{ex^2}{d}+1} \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{\frac{ex^2}{d}+1}} dx}{d\sqrt{d+ex^2}} + \frac{x(a+b\operatorname{sech}^{-1}(cx))}{d\sqrt{d+ex^2}} \\
& \quad \downarrow 321 \\
& \frac{x(a+b\operatorname{sech}^{-1}(cx))}{d\sqrt{d+ex^2}} + \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\sqrt{\frac{ex^2}{d}+1} \operatorname{EllipticF}(\arcsin(cx), -\frac{e}{c^2d})}{cd\sqrt{d+ex^2}}
\end{aligned}$$

input `Int[(a + b*ArcSech[c*x])/(d + e*x^2)^(3/2),x]`

output `(x*(a + b*ArcSech[c*x]))/(d*Sqrt[d + e*x^2]) + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*Sqrt[1 + (e*x^2)/d]*EllipticF[ArcSin[c*x], -(e/(c^2*d))])/(c*d*Sqrt[d + e*x^2])`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 321 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(1/(Sqrt[a]*Sqrt[c]*Rt[-d/c, 2]))*EllipticF[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-b/a, -d/c])`

rule 323

```
Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] := S
imp[Sqrt[1 + (d/c)*x^2]/Sqrt[c + d*x^2] Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (
d/c)*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]
```

rule 6845

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_))*((d_) + (e_)*(x_)^2)^(p_), x_Sym
bol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u
, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*
Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e}, x] && (
IGtQ[p, 0] || ILtQ[p + 1/2, 0])
```

**Maple [F]**

$$\int \frac{a + b \operatorname{arcsech}(cx)}{(x^2 e + d)^{\frac{3}{2}}} dx$$

input

```
int((a+b*arcsech(c*x))/(e*x^2+d)^(3/2), x)
```

output

```
int((a+b*arcsech(c*x))/(e*x^2+d)^(3/2), x)
```

**Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.16

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{3/2}} dx = \frac{\sqrt{ex^2 + d} b c d x \log\left(\frac{cx \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2} + 1}}{cx}\right) + \sqrt{ex^2 + d} a c d x + (b e x^2 + b d) \sqrt{d} F(\arcsin(cx))}{cd^2 e x^2 + cd^3}$$

input

```
integrate((a+b*arcsech(c*x))/(e*x^2+d)^(3/2), x, algorithm="fricas")
```

output

```
(sqrt(e*x^2 + d)*b*c*d*x*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x
)) + sqrt(e*x^2 + d)*a*c*d*x + (b*e*x^2 + b*d)*sqrt(d)*elliptic_f(arcsin(c
*x), -e/(c^2*d)))/(c*d^2*e*x^2 + c*d^3)
```

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{3/2}} dx = \int \frac{a + b \operatorname{asech}(cx)}{(d + ex^2)^{\frac{3}{2}}} dx$$

input `integrate((a+b*asech(c*x))/(e*x**2+d)**(3/2), x)`

output `Integral((a + b*asech(c*x))/(d + e*x**2)**(3/2), x)`

**Maxima [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{3/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x^2+d)^(3/2), x, algorithm="maxima")`

output `b*integrate(log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(e*x^2 + d)^(3/2), x) + a*x/(sqrt(e*x^2 + d)*d)`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{3/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x^2+d)^(3/2), x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/(e*x^2 + d)^(3/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{(ex^2 + d)^{3/2}} dx$$

input `int((a + b*acosh(1/(c*x)))/(d + e*x^2)^(3/2), x)`output `int((a + b*acosh(1/(c*x)))/(d + e*x^2)^(3/2), x)`**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{3/2}} dx = \int \frac{a \operatorname{sech}(cx) b + a}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `int((a+b*asech(c*x))/(e*x^2+d)^(3/2), x)`output `int((a+b*asech(c*x))/(e*x^2+d)^(3/2), x)`

**3.166**  $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^2(d+ex^2)^{3/2}} dx$

Optimal result	1318
Mathematica [C] (verified)	1319
Rubi [A] (verified)	1319
Maple [F]	1323
Fricas [A] (verification not implemented)	1324
Sympy [F]	1324
Maxima [F(-2)]	1325
Giac [F]	1325
Mupad [F(-1)]	1325
Reduce [F]	1326

**Optimal result**

Integrand size = 23, antiderivative size = 249

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^2(d + ex^2)^{3/2}} dx = \frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{d^2x} + \frac{a + b\operatorname{sech}^{-1}(cx)}{dx\sqrt{d+ex^2}}$$

$$- \frac{2\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{d^2x} + \frac{bc\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{d+ex^2}E(\arcsin(cx) | -\frac{e}{c^2d})}{d^2\sqrt{1+\frac{ex^2}{d}}}$$

$$- \frac{b(c^2d + 2e)\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1+\frac{ex^2}{d}}\operatorname{EllipticF}(\arcsin(cx), -\frac{e}{c^2d})}{cd^2\sqrt{d+ex^2}}$$

output

```
b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)*(e*x^2+d)^(1/2)/d^2/x
+(a+b*arcsech(c*x))/d/x/(e*x^2+d)^(1/2)-2*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x
))/d^2/x+b*c*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(e*x^2+d)^(1/2)*EllipticE(c*x
,(-e/c^2/d)^(1/2))/d^2/(1+e*x^2/d)^(1/2)-b*(c^2*d+2*e)*(1/(c*x+1))^(1/2)*(
c*x+1)^(1/2)*(1+e*x^2/d)^(1/2)*EllipticF(c*x,(-e/c^2/d)^(1/2))/c/d^2/(e*x^
2+d)^(1/2)
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 23.96 (sec) , antiderivative size = 501, normalized size of antiderivative = 2.01

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^{3/2}} dx = \frac{b \sqrt{\frac{1-cx}{1+cx}} (1+cx)(d+ex^2)}{x} - \frac{a(d+2ex^2)}{x} - \frac{b(d+2ex^2) \operatorname{sech}^{-1}(cx)}{x} + \frac{b \sqrt{\frac{1-cx}{1+cx}} \left( -c^2 (d+ex^2) + \frac{(1+cx) \sqrt{\frac{c}{cx}}}{(cx)} \right)}{x}$$

input `Integrate[(a + b*ArcSech[c*x])/(x^2*(d + e*x^2)^(3/2)),x]`

output

```
((b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(d + e*x^2))/x - (a*(d + 2*e*x^2))
/x - (b*(d + 2*e*x^2)*ArcSech[c*x])/x + (b*Sqrt[(1 - c*x)/(1 + c*x)]*(-(c^
2*(d + e*x^2)) + ((1 + c*x)*Sqrt[(c*(Sqrt[d] - I*Sqrt[e]*x))/((c*Sqrt[d] -
I*Sqrt[e])*(1 + c*x))])*Sqrt[(c*(Sqrt[d] + I*Sqrt[e]*x))/((c*Sqrt[d] + I*S
qrt[e])*(1 + c*x))])*((-I)*(c*Sqrt[d] - I*Sqrt[e])^2*EllipticE[I*ArcSinh[Sq
rt[((c^2*d + e)*(1 - c*x))/((c*Sqrt[d] + I*Sqrt[e])^2*(1 + c*x))]]], (c*Sqr
t[d] + I*Sqrt[e])^2/(c*Sqrt[d] - I*Sqrt[e])^2 + 2*(c*Sqrt[d] - (2*I)*Sqrt
[e])*Sqrt[e]*EllipticF[I*ArcSinh[Sqrt[((c^2*d + e)*(1 - c*x))/((c*Sqrt[d]
+ I*Sqrt[e])^2*(1 + c*x))]]], (c*Sqrt[d] + I*Sqrt[e])^2/(c*Sqrt[d] - I*Sqrt
[e])^2))/Sqrt[-((c*Sqrt[d] - I*Sqrt[e])*(-1 + c*x))/((c*Sqrt[d] + I*Sqrt
[e])*(1 + c*x))]))/c)/(d^2*Sqrt[d + e*x^2])
```

**Rubi [A] (verified)**

Time = 0.55 (sec) , antiderivative size = 214, normalized size of antiderivative = 0.86, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.478$ , Rules used = {6855, 25, 27, 445, 25, 27, 399, 323, 321, 330, 327}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^{3/2}} dx$$



$$\begin{aligned}
& \downarrow 6855 \\
& b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int -\frac{2ex^2+d}{d^2x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx - \frac{2ex(a+b\operatorname{sech}^{-1}(cx))}{d^2\sqrt{d+ex^2}} - \frac{a+b\operatorname{sech}^{-1}(cx)}{dx\sqrt{d+ex^2}} \\
& \downarrow 25 \\
& -b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{2ex^2+d}{d^2x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx - \frac{2ex(a+b\operatorname{sech}^{-1}(cx))}{d^2\sqrt{d+ex^2}} - \frac{a+b\operatorname{sech}^{-1}(cx)}{dx\sqrt{d+ex^2}} \\
& \downarrow 27 \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{2ex^2+d}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx}{d^2} - \frac{2ex(a+b\operatorname{sech}^{-1}(cx))}{d^2\sqrt{d+ex^2}} - \frac{a+b\operatorname{sech}^{-1}(cx)}{dx\sqrt{d+ex^2}} \\
& \downarrow 445 \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( -\frac{\int -\frac{de(2-c^2x^2)}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx}{d} - \frac{\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{x} \right)}{d^2} - \frac{2ex(a+b\operatorname{sech}^{-1}(cx))}{d^2\sqrt{d+ex^2}} - \frac{a+b\operatorname{sech}^{-1}(cx)}{dx\sqrt{d+ex^2}} \\
& \downarrow 25 \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{de(2-c^2x^2)}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx}{d} - \frac{\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{x} \right)}{d^2} - \frac{2ex(a+b\operatorname{sech}^{-1}(cx))}{d^2\sqrt{d+ex^2}} - \frac{a+b\operatorname{sech}^{-1}(cx)}{dx\sqrt{d+ex^2}} \\
& \downarrow 27 \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( e \int \frac{2-c^2x^2}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx - \frac{\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{x} \right)}{d^2} - \frac{2ex(a+b\operatorname{sech}^{-1}(cx))}{d^2\sqrt{d+ex^2}} - \frac{a+b\operatorname{sech}^{-1}(cx)}{dx\sqrt{d+ex^2}} \\
& \downarrow 399 \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( e \left( \frac{(c^2d+2e) \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx}{e} - \frac{c^2 \int \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}dx}{e} \right) - \frac{\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{x} \right)}{d^2} - \frac{2ex(a+b\operatorname{sech}^{-1}(cx))}{d^2\sqrt{d+ex^2}} - \frac{a+b\operatorname{sech}^{-1}(cx)}{dx\sqrt{d+ex^2}}
\end{aligned}$$

↓ 323

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(e\left(\frac{(c^2d+2e)\sqrt{\frac{ex^2}{d}+1}\int\frac{1}{\sqrt{1-c^2x^2}\sqrt{\frac{ex^2}{d}+1}}dx}{e\sqrt{d+ex^2}}-\frac{c^2\int\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}dx}{e}\right)-\frac{\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{x}\right)$$

---


$$\frac{2ex(a+b\operatorname{sech}^{-1}(cx))}{d^2\sqrt{d+ex^2}}-\frac{a+b\operatorname{sech}^{-1}(cx)}{dx\sqrt{d+ex^2}}$$

↓ 321

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(e\left(\frac{(c^2d+2e)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}}-\frac{c^2\int\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}dx}{e}\right)-\frac{\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{x}\right)$$

---


$$\frac{2ex(a+b\operatorname{sech}^{-1}(cx))}{d^2\sqrt{d+ex^2}}-\frac{a+b\operatorname{sech}^{-1}(cx)}{dx\sqrt{d+ex^2}}$$

↓ 330

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(e\left(\frac{(c^2d+2e)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}}-\frac{c^2\sqrt{d+ex^2}\int\frac{\sqrt{\frac{ex^2}{d}+1}}{\sqrt{1-c^2x^2}}dx}{e\sqrt{\frac{ex^2}{d}+1}}\right)-\frac{\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{x}\right)$$

---


$$\frac{2ex(a+b\operatorname{sech}^{-1}(cx))}{d^2\sqrt{d+ex^2}}-\frac{a+b\operatorname{sech}^{-1}(cx)}{dx\sqrt{d+ex^2}}$$

↓ 327

$$\frac{2ex(a+b\operatorname{sech}^{-1}(cx))}{d^2\sqrt{d+ex^2}}-\frac{a+b\operatorname{sech}^{-1}(cx)}{dx\sqrt{d+ex^2}}-\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(e\left(\frac{(c^2d+2e)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx),-\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}}-\frac{c\sqrt{d+ex^2}E\left(\arcsin(cx)\middle|-\frac{e}{c^2d}\right)}{e\sqrt{\frac{ex^2}{d}+1}}\right)-\frac{\sqrt{1-c^2x^2}\sqrt{d+ex^2}}{x}\right)}{d^2}$$

input

`Int[(a + b*ArcSech[c*x])/(x^2*(d + e*x^2)^(3/2)),x]`

output

$$-\left(\frac{a + b \operatorname{ArcSech}[c x]}{d x \sqrt{d + e x^2}}\right) - \frac{2 e x (a + b \operatorname{ArcSech}[c x])}{d^2 \sqrt{d + e x^2}} - \frac{b \sqrt{(1 + c x)^{-1}} \sqrt{1 + c x} \left(-\left(\sqrt{1 - c^2 x^2} \sqrt{d + e x^2}\right) / x + e \left(-\left(c \sqrt{d + e x^2} \operatorname{EllipticE}[\operatorname{ArcSin}[c x], -\left(e / (c^2 d)\right)]\right) / \left(e \sqrt{1 + (e x^2) / d}\right)\right) + \left((c^2 d + 2 e) \sqrt{1 + (e x^2) / d} \operatorname{EllipticF}[\operatorname{ArcSin}[c x], -\left(e / (c^2 d)\right)]\right) / \left(c e \sqrt{d + e x^2}\right)\right)}{d^2}$$

### Defintions of rubi rules used

rule 25

$$\operatorname{Int}[-(F x), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F x, x], x]$$

rule 27

$$\operatorname{Int}[(a) (F x), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[a \operatorname{Int}[F x, x], x] /; \operatorname{FreeQ}[a, x] \&\& \operatorname{!MatchQ}[F x, (b) (G x) /; \operatorname{FreeQ}[b, x]]$$

rule 321

$$\operatorname{Int}[1 / (\sqrt{(a) + (b) (x)^2} \sqrt{(c) + (d) (x)^2}), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(1 / (\sqrt{a} \sqrt{c} \operatorname{Rt}[-d/c, 2])) \operatorname{EllipticF}[\operatorname{ArcSin}[\operatorname{Rt}[-d/c, 2] x], b (c / (a d))], x] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \&\& \operatorname{NegQ}[d/c] \&\& \operatorname{GtQ}[c, 0] \&\& \operatorname{GtQ}[a, 0] \&\& \operatorname{!(NegQ}[b/a] \&\& \operatorname{SimplerSqrtQ}[-b/a, -d/c])$$

rule 323

$$\operatorname{Int}[1 / (\sqrt{(a) + (b) (x)^2} \sqrt{(c) + (d) (x)^2}), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\sqrt{1 + (d/c) x^2} / \sqrt{c + d x^2} \operatorname{Int}[1 / (\sqrt{a + b x^2} \sqrt{1 + (d/c) x^2}), x], x] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \&\& \operatorname{!GtQ}[c, 0]$$

rule 327

$$\operatorname{Int}[\sqrt{(a) + (b) (x)^2} / \sqrt{(c) + (d) (x)^2}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(\sqrt{a} / (\sqrt{c} \operatorname{Rt}[-d/c, 2])) \operatorname{EllipticE}[\operatorname{ArcSin}[\operatorname{Rt}[-d/c, 2] x], b (c / (a d))], x] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \&\& \operatorname{NegQ}[d/c] \&\& \operatorname{GtQ}[c, 0] \&\& \operatorname{GtQ}[a, 0]$$

rule 330

$$\operatorname{Int}[\sqrt{(a) + (b) (x)^2} / \sqrt{(c) + (d) (x)^2}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\sqrt{a + b x^2} / \sqrt{1 + (b/a) x^2} \operatorname{Int}[\sqrt{1 + (b/a) x^2} / \sqrt{c + d x^2}, x], x] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \&\& \operatorname{NegQ}[d/c] \&\& \operatorname{GtQ}[c, 0] \&\& \operatorname{!GtQ}[a, 0]$$

rule 399

```
Int[((e_) + (f_)*(x_)^2)/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] := Simp[f/b Int[Sqrt[a + b*x^2]/Sqrt[c + d*x^2], x], x] + Simp[(b*e - a*f)/b Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && !((PosQ[b/a] && PosQ[d/c]) || (NegQ[b/a] && (PosQ[d/c] || (GtQ[a, 0] && (!GtQ[c, 0] || SimplerSqrtQ[-b/a, -d/c])))))
```

rule 445

```
Int[((g_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2), x_Symbol] := Simp[e*(g*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(a*c*g*(m + 1))), x] + Simp[1/(a*c*g^2*(m + 1)) Int[(g*x)^(m + 2)*(a + b*x^2)^p*(c + d*x^2)^q*Simp[a*f*c*(m + 1) - e*(b*c + a*d)*(m + 2 + 1) - e*2*(b*c*p + a*d*q) - b*e*d*(m + 2*(p + q + 2) + 1)*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, p, q}, x] && LtQ[m, -1]
```

rule 6855

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

## Maple [F]

$$\int \frac{a + b \operatorname{arcsech}(cx)}{x^2 (x^2 e + d)^{\frac{3}{2}}} dx$$

input

```
int((a+b*arcsech(c*x))/x^2/(e*x^2+d)^(3/2), x)
```

output

```
int((a+b*arcsech(c*x))/x^2/(e*x^2+d)^(3/2), x)
```

**Fricas [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 238, normalized size of antiderivative = 0.96

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^{3/2}} dx =$$

$$\frac{(2bcdex^2 + bcd^2)\sqrt{ex^2 + d} \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}}+1}{cx}\right) + \left(2acdex^2 + acd^2 - (bc^2dex^3 + bc^2d^2x)\sqrt{-\frac{c^2x^2-1}{c^2x^2}}\right)\sqrt{d}}{\dots}$$

input `integrate((a+b*arcsech(c*x))/x^2/(e*x^2+d)^(3/2),x, algorithm="fricas")`

output `-((2*b*c*d*e*x^2 + b*c*d^2)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + (2*a*c*d*e*x^2 + a*c*d^2 - (b*c^2*d*e*x^3 + b*c^2*d^2*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d) - ((b*c^4*d*e*x^3 + b*c^4*d^2*x)*elliptic_e(arcsin(c*x), -e/(c^2*d)) - ((b*c^4*d*e + 2*b*e^2)*x^3 + (b*c^4*d^2 + 2*b*d*e)*x)*elliptic_f(arcsin(c*x), -e/(c^2*d)))*sqrt(d))/(c*d^3*e*x^3 + c*d^4*x)`

**Sympy [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^{3/2}} dx = \int \frac{a + b \operatorname{asech}(cx)}{x^2 (d + ex^2)^{\frac{3}{2}}} dx$$

input `integrate((a+b*asech(c*x))/(x**2/(e*x**2+d)**(3/2)),x)`

output `Integral((a + b*asech(c*x))/(x**2*(d + e*x**2)**(3/2)), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/x^2/(e*x^2+d)^(3/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation \*may\* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^{3/2}} dx = \int \frac{b \operatorname{ar} \operatorname{sech}(cx) + a}{(ex^2 + d)^{\frac{3}{2}} x^2} dx$$

input `integrate((a+b*arcsech(c*x))/x^2/(e*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/((e*x^2 + d)^(3/2)*x^2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x^2 (ex^2 + d)^{3/2}} dx$$

input `int((a + b*acosh(1/(c*x)))/(x^2*(d + e*x^2)^(3/2)),x)`

output `int((a + b*acosh(1/(c*x)))/(x^2*(d + e*x^2)^(3/2)), x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^2 (d + ex^2)^{3/2}} dx = \int \frac{a \operatorname{sech}(cx) b + a}{x^2 (ex^2 + d)^{\frac{3}{2}}} dx$$

input `int((a+b*asech(c*x))/x^2/(e*x^2+d)^(3/2),x)`

output `int((a+b*asech(c*x))/x^2/(e*x^2+d)^(3/2),x)`

$$3.167 \quad \int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx$$

Optimal result	1327
Mathematica [A] (verified)	1328
Rubi [A] (verified)	1328
Maple [F]	1332
Fricas [B] (verification not implemented)	1333
Sympy [F(-1)]	1334
Maxima [F(-2)]	1334
Giac [F]	1334
Mupad [F(-1)]	1335
Reduce [F]	1335

### Optimal result

Integrand size = 23, antiderivative size = 272

$$\begin{aligned} \int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx &= -\frac{bd \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{3e^2 (c^2d + e) \sqrt{d + ex^2}} \\ &\quad - \frac{d^2 (a + b \operatorname{sech}^{-1}(cx))}{3e^3 (d + ex^2)^{3/2}} + \frac{2d (a + b \operatorname{sech}^{-1}(cx))}{e^3 \sqrt{d + ex^2}} \\ &\quad + \frac{\sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx))}{e^3} - \frac{b \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{ce^{5/2}} \\ &\quad - \frac{8b\sqrt{d} \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{3e^3} \end{aligned}$$

output

```
-1/3*b*d*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/e^2/(c^2*d+e)/
(e*x^2+d)^(1/2)-1/3*d^2*(a+b*arcsech(c*x))/e^3/(e*x^2+d)^(3/2)+2*d*(a+b*ar
csech(c*x))/e^3/(e*x^2+d)^(1/2)+(e*x^2+d)^(1/2)*(a+b*arcsech(c*x))/e^3-b*(
1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctan(e^(1/2)*(-c^2*x^2+1)^(1/2)/c/(e*x^2+
d)^(1/2))/c/e^(5/2)-8/3*b*d^(1/2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh(
(e*x^2+d)^(1/2)/d^(1/2)/(-c^2*x^2+1)^(1/2))/e^3
```



**Mathematica [A] (verified)**

Time = 22.10 (sec) , antiderivative size = 348, normalized size of antiderivative = 1.28

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \frac{-bde \sqrt{\frac{1-cx}{1+cx}} (1+cx) (d+ex^2) + a(c^2d+e) (8d^2+12dex^2+3e^2x^4) + b(c^2d+e) (8d^2+12dex^2+3e^2x^4)}{3e^3(c^2d+e)(d+ex^2)^{3/2}} + \frac{b \sqrt{\frac{1-cx}{1+cx}} \sqrt{1-c^2x^2} \left( 3\sqrt{-c^2} \sqrt{-c^2d-e} \sqrt{e} \sqrt{\frac{c^2(d+ex^2)}{c^2d+e}} \arcsin\left(\frac{c\sqrt{e}\sqrt{1-c^2x^2}}{\sqrt{-c^2}\sqrt{-c^2d-e}}\right) + 8c^3\sqrt{d}\sqrt{-d-ex^2} \arctan\left(\frac{\sqrt{d}\sqrt{1-c^2x^2}}{\sqrt{-d-ex^2}}\right) \right)}{3c^3e^3(-1+cx)\sqrt{d+ex^2}}$$

input

```
Integrate[(x^5*(a + b*ArcSech[c*x]))/(d + e*x^2)^(5/2), x]
```

output

```
(-(b*d*e*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(d + e*x^2)) + a*(c^2*d + e)*(8*d^2 + 12*d*e*x^2 + 3*e^2*x^4) + b*(c^2*d + e)*(8*d^2 + 12*d*e*x^2 + 3*e^2*x^4)*ArcSech[c*x])/(3*e^3*(c^2*d + e)*(d + e*x^2)^(3/2)) + (b*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*(3*Sqrt[-c^2]*Sqrt[-(c^2*d) - e]*Sqrt[e]*Sqrt[(c^2*(d + e*x^2))/(c^2*d + e)]*ArcSin[(c*Sqrt[e]*Sqrt[1 - c^2*x^2])/(Sqrt[-c^2]*Sqrt[-(c^2*d) - e]]) + 8*c^3*Sqrt[d]*Sqrt[-d - e*x^2]*ArcTan[(Sqrt[d]*Sqrt[1 - c^2*x^2])/Sqrt[-d - e*x^2]])/(3*c^3*e^3*(-1 + c*x)*Sqrt[d + e*x^2])
```

**Rubi [A] (verified)**

Time = 1.45 (sec) , antiderivative size = 229, normalized size of antiderivative = 0.84, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.435$ , Rules used = {6855, 27, 7282, 2117, 27, 175, 66, 104, 218, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx$$

↓ 6855

$$\begin{aligned}
& b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{3e^2x^4 + 12dex^2 + 8d^2}{3e^3x\sqrt{1-c^2x^2}(ex^2+d)^{3/2}} dx - \frac{d^2(a + b\operatorname{sech}^{-1}(cx))}{3e^3(d+ex^2)^{3/2}} + \\
& \quad \frac{2d(a + b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} + \frac{\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{e^3} \\
& \quad \downarrow 27 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{3e^2x^4 + 12dex^2 + 8d^2}{x\sqrt{1-c^2x^2}(ex^2+d)^{3/2}} dx}{3e^3} - \frac{d^2(a + b\operatorname{sech}^{-1}(cx))}{3e^3(d+ex^2)^{3/2}} + \frac{2d(a + b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} + \\
& \quad \frac{\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{e^3} \\
& \quad \downarrow 7282 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{3e^2x^4 + 12dex^2 + 8d^2}{x^2\sqrt{1-c^2x^2}(ex^2+d)^{3/2}} dx^2}{6e^3} - \frac{d^2(a + b\operatorname{sech}^{-1}(cx))}{3e^3(d+ex^2)^{3/2}} + \frac{2d(a + b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} + \\
& \quad \frac{\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{e^3} \\
& \quad \downarrow 2117 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{2 \int \frac{d(dc^2+e)(3ex^2+8d)}{2x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2}{d(c^2d+e)} - \frac{2de\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{6e^3} - \frac{d^2(a + b\operatorname{sech}^{-1}(cx))}{3e^3(d+ex^2)^{3/2}} + \\
& \quad \frac{2d(a + b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} + \frac{\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{e^3} \\
& \quad \downarrow 27 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \int \frac{3ex^2+8d}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 - \frac{2de\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{6e^3} - \frac{d^2(a + b\operatorname{sech}^{-1}(cx))}{3e^3(d+ex^2)^{3/2}} + \\
& \quad \frac{2d(a + b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} + \frac{\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{e^3} \\
& \quad \downarrow 175 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( 3e \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 + 8d \int \frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2 - \frac{2de\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{6e^3} - \\
& \quad \frac{d^2(a + b\operatorname{sech}^{-1}(cx))}{3e^3(d+ex^2)^{3/2}} + \frac{2d(a + b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}} + \frac{\sqrt{d+ex^2}(a + b\operatorname{sech}^{-1}(cx))}{e^3} \\
& \quad \downarrow 66
\end{aligned}$$

$$\begin{aligned}
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(8d\int\frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2+6e\int\frac{1}{-ex^4-c^2}d\frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}}-\frac{2de\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}}\right)}{6e^3} \\
& \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{3e^3(d+ex^2)^{3/2}}+\frac{2d(a+b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}}+\frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3} \\
& \quad \downarrow 104 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(6e\int\frac{1}{-ex^4-c^2}d\frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}}+16d\int\frac{1}{x^4-d}d\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}-\frac{2de\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}}\right)}{6e^3} \\
& \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{3e^3(d+ex^2)^{3/2}}+\frac{2d(a+b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}}+\frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3} \\
& \quad \downarrow 218 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(16d\int\frac{1}{x^4-d}d\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}-\frac{6\sqrt{e}\arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c}-\frac{2de\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}}\right)}{6e^3} \\
& \frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{3e^3(d+ex^2)^{3/2}}+\frac{2d(a+b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}}+\frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3} \\
& \quad \downarrow 220 \\
& -\frac{d^2(a+b\operatorname{sech}^{-1}(cx))}{3e^3(d+ex^2)^{3/2}}+\frac{2d(a+b\operatorname{sech}^{-1}(cx))}{e^3\sqrt{d+ex^2}}+\frac{\sqrt{d+ex^2}(a+b\operatorname{sech}^{-1}(cx))}{e^3}+ \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{6\sqrt{e}\arctan\left(\frac{\sqrt{e}\sqrt{1-c^2x^2}}{c\sqrt{d+ex^2}}\right)}{c}-16\sqrt{d}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)-\frac{2de\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}}\right)}{6e^3}
\end{aligned}$$

input `Int[(x^5*(a + b*ArcSech[c*x]))/(d + e*x^2)^(5/2),x]`

output

```

-1/3*(d^2*(a + b*ArcSech[c*x]))/(e^3*(d + e*x^2)^(3/2)) + (2*d*(a + b*ArcSech[c*x]))/(e^3*sqrt[d + e*x^2]) + (sqrt[d + e*x^2]*(a + b*ArcSech[c*x]))/e^3 + (b*sqrt[(1 + c*x)^(-1)]*sqrt[1 + c*x]*((-2*d*e*sqrt[1 - c^2*x^2]))/((c^2*d + e)*sqrt[d + e*x^2]) - (6*sqrt[e]*ArcTan[(sqrt[e]*sqrt[1 - c^2*x^2])]/(c*sqrt[d + e*x^2]))/c - 16*sqrt[d]*ArcTanh[sqrt[d + e*x^2]/(sqrt[d]*sqrt[1 - c^2*x^2])])/((6*e^3)

```

## Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`
- rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`
- rule 104 `Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 175 `Int[(((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)))/((a_.) + (b_.)*(x_)), x_] := Simp[h/b Int[(c + d*x)^n*(e + f*x)^p, x], x] + Simp[(b*g - a*h)/b Int[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 2117

```
Int[(Px_)*((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)
*(x_))^(p_), x_Symbol] := With[{Qx = PolynomialQuotient[Px, a + b*x, x],
R = PolynomialRemainder[Px, a + b*x, x]}, Simp[b*R*(a + b*x)^(m + 1)*(c +
d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Si
mp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*
(e + f*x)^p*ExpandToSum[(m + 1)*(b*c - a*d)*(b*e - a*f)*Qx + a*d*f*R*(m + 1)
) - b*R*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*R*(m + n + p + 3)*x, x]
, x]] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && PolyQ[Px, x] && LtQ[m, -
1] && IntegersQ[2*m, 2*n, 2*p]
```

rule 6855

```
Int[((a_.) + ArcSech[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_)*((d_.) + (e_.)*(
x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Si
mp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)]
Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x]] /; Fre
eQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] &&
GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2
*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

rule 7282

```
Int[(u_)/(x_), x_Symbol] := With[{lst = PowerVariableExpn[u, 0, x]}, Simp[1
/lst[[2]] Subst[Int[NormalizeIntegrand[Simplify[lst[[1]]/x], x], x], x, (
lst[[3]]*x)^lst[[2]]], x] /; !FalseQ[lst] && NeQ[lst[[2]], 0] /; NonsumQ[
u] && !RationalFunctionQ[u, x]
```

## Maple [F]

$$\int \frac{x^5(a + b \operatorname{arcsech}(cx))}{(x^2e + d)^{\frac{5}{2}}} dx$$

input

```
int(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2), x)
```

output

```
int(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2), x)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 586 vs.  $2(180) = 360$ .

Time = 0.35 (sec) , antiderivative size = 2415, normalized size of antiderivative = 8.88

$$\int \frac{x^5(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \text{Too large to display}$$

input `integrate(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="fricas")`

output

```
[ -1/12*(3*(b*c^2*d^3 + (b*c^2*d*e^2 + b*e^3)*x^4 + b*d^2*e + 2*(b*c^2*d^2*e + b*d*e^2)*x^2)*sqrt(-e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e + 8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^4*e*x^3 + (c^4*d - c^2*e)*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + e^2) - 4*(8*b*c^3*d^3 + 8*b*c*d^2*e + 3*(b*c^3*d*e^2 + b*c*e^3)*x^4 + 12*(b*c^3*d^2*e + b*c*d*e^2)*x^2)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - 8*(b*c^3*d^3 + b*c*d^2*e + (b*c^3*d*e^2 + b*c*e^3)*x^4 + 2*(b*c^3*d^2*e + b*c*d*e^2)*x^2)*sqrt(d)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 + 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) - 4*(8*a*c^3*d^3 + 8*a*c*d^2*e + 3*(a*c^3*d*e^2 + a*c*e^3)*x^4 + 12*(a*c^3*d^2*e + a*c*d*e^2)*x^2 - (b*c^2*d*e^2*x^3 + b*c^2*d^2*e*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d))/(c^3*d^3*e^3 + c*d^2*e^4 + (c^3*d*e^5 + c*e^6)*x^4 + 2*(c^3*d^2*e^4 + c*d*e^5)*x^2), -1/6*(3*(b*c^2*d^3 + (b*c^2*d*e^2 + b*e^3)*x^4 + b*d^2*e + 2*(b*c^2*d^2*e + b*d*e^2)*x^2)*sqrt(e)*arctan(1/2*(2*c^2*e*x^3 + (c^2*d - e)*x)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*e^2*x^4 + (c^2*d*e - e^2)*x^2 - d*e) - 2*(8*b*c^3*d^3 + 8*b*c*d^2*e + 3*(b*c^3*d*e^2 + b*c*e^3)*x^4 + 12*(b*c^3*d^2*e + b*c*d*e^2)*x^2)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - 4*(b*c^3*d^3 + b*c*d^2*e + (b*c^3*d*e^2 + b*c*e^3)*x^4 + 2*(b*c^3*d^2*e + b*c*d*e^2)*x^2)*sqrt(d)*log(...
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \text{Timed out}$$

input `integrate(x**5*(a+b*asech(c*x))/(e*x**2+d)**(5/2),x)`

output `Timed out`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**Giac [F]**

$$\int \frac{x^5 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^5}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input `integrate(x^5*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^5/(e*x^2 + d)^(5/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^5(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{x^5(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^{5/2}} dx$$

input `int((x^5*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(5/2), x)`

output `int((x^5*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(5/2), x)`

**Reduce [F]**

$$\int \frac{x^5(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{x^5(a \operatorname{sech}(cx) b + a)}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input `int(x^5*(a+b*asech(c*x))/(e*x^2+d)^(5/2), x)`

output `int(x^5*(a+b*asech(c*x))/(e*x^2+d)^(5/2), x)`



**3.168** 
$$\int \frac{x^3 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx$$

Optimal result	1336
Mathematica [A] (verified)	1337
Rubi [A] (verified)	1337
Maple [F]	1340
Fricas [B] (verification not implemented)	1340
Sympy [F(-1)]	1341
Maxima [F]	1341
Giac [F]	1342
Mupad [F(-1)]	1342
Reduce [F]	1342

**Optimal result**

Integrand size = 23, antiderivative size = 179

$$\int \frac{x^3 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \frac{b \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{3e(c^2d+e)\sqrt{d+ex^2}} + \frac{d(a+b \operatorname{sech}^{-1}(cx))}{3e^2(d+ex^2)^{3/2}}$$

$$- \frac{a+b \operatorname{sech}^{-1}(cx)}{e^2 \sqrt{d+ex^2}} + \frac{2b \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{3\sqrt{de^2}}$$

output

```
1/3*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/e/(c^2*d+e)/(e*x^
2+d)^(1/2)+1/3*d*(a+b*arcsech(c*x))/e^2/(e*x^2+d)^(3/2)-(a+b*arcsech(c*x))
/e^2/(e*x^2+d)^(1/2)+2/3*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh((e*x^2+
d)^(1/2)/d^(1/2)/(-c^2*x^2+1)^(1/2))/d^(1/2)/e^2
```

**Mathematica [A] (verified)**

Time = 0.43 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.22

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \frac{be\sqrt{\frac{1-cx}{1+cx}}(1+cx)(d+ex^2) - a(c^2d+e)(2d+3ex^2) - b(c^2d+e)(2d+3ex^2)}{3e^2(c^2d+e)(d+ex^2)^{3/2}} - \frac{2b\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\sqrt{-d-ex^2} \arctan\left(\frac{\sqrt{d}\sqrt{1-c^2x^2}}{\sqrt{-d-ex^2}}\right)}{3\sqrt{de^2(-1+cx)}\sqrt{d+ex^2}}$$

input

```
Integrate[(x^3*(a + b*ArcSech[c*x]))/(d + e*x^2)^(5/2),x]
```

output

```
(b*e*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(d + e*x^2) - a*(c^2*d + e)*(2*d + 3*e*x^2) - b*(c^2*d + e)*(2*d + 3*e*x^2)*ArcSech[c*x])/(3*e^2*(c^2*d + e)*(d + e*x^2)^(3/2)) - (2*b*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*Sqrt[-d - e*x^2]*ArcTan[(Sqrt[d]*Sqrt[1 - c^2*x^2])/Sqrt[-d - e*x^2]])/(3*Sqrt[d]*e^2*(-1 + c*x)*Sqrt[d + e*x^2])
```

**Rubi [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.88, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$ , Rules used = {6855, 27, 435, 169, 27, 104, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx$$

↓ 6855

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int -\frac{3ex^2 + 2d}{3e^2x\sqrt{1-c^2x^2}(ex^2+d)^{3/2}} dx - \frac{a + b \operatorname{sech}^{-1}(cx)}{e^2\sqrt{d+ex^2}} + \frac{d(a + b \operatorname{sech}^{-1}(cx))}{3e^2(d+ex^2)^{3/2}}$$

↓ 27

$$\begin{aligned}
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\int\frac{3ex^2+2d}{x\sqrt{1-c^2x^2}(ex^2+d)^{3/2}}dx}{3e^2}-\frac{a+b\operatorname{sech}^{-1}(cx)}{e^2\sqrt{d+ex^2}}+\frac{d(a+b\operatorname{sech}^{-1}(cx))}{3e^2(d+ex^2)^{3/2}} \\
& \quad \downarrow 435 \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\int\frac{3ex^2+2d}{x^2\sqrt{1-c^2x^2}(ex^2+d)^{3/2}}dx^2}{6e^2}-\frac{a+b\operatorname{sech}^{-1}(cx)}{e^2\sqrt{d+ex^2}}+\frac{d(a+b\operatorname{sech}^{-1}(cx))}{3e^2(d+ex^2)^{3/2}} \\
& \quad \downarrow 169 \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(2\int\frac{d(dc^2+e)}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2-\frac{2e\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}}\right)}{6e^2}-\frac{a+b\operatorname{sech}^{-1}(cx)}{e^2\sqrt{d+ex^2}}+ \\
& \quad \frac{d(a+b\operatorname{sech}^{-1}(cx))}{3e^2(d+ex^2)^{3/2}} \\
& \quad \downarrow 27 \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(2\int\frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}}dx^2-\frac{2e\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}}\right)}{6e^2}-\frac{a+b\operatorname{sech}^{-1}(cx)}{e^2\sqrt{d+ex^2}}+ \\
& \quad \frac{d(a+b\operatorname{sech}^{-1}(cx))}{3e^2(d+ex^2)^{3/2}} \\
& \quad \downarrow 104 \\
& -\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(4\int\frac{1}{x^4-d}d\frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}-\frac{2e\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}}\right)}{6e^2}-\frac{a+b\operatorname{sech}^{-1}(cx)}{e^2\sqrt{d+ex^2}}+ \\
& \quad \frac{d(a+b\operatorname{sech}^{-1}(cx))}{3e^2(d+ex^2)^{3/2}} \\
& \quad \downarrow 220 \\
& -\frac{a+b\operatorname{sech}^{-1}(cx)}{e^2\sqrt{d+ex^2}}+\frac{d(a+b\operatorname{sech}^{-1}(cx))}{3e^2(d+ex^2)^{3/2}}- \\
& \quad \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(-\frac{4\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{\sqrt{d}}-\frac{2e\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}}\right)}{6e^2}
\end{aligned}$$

input `Int[(x^3*(a + b*ArcSech[c*x]))/(d + e*x^2)^(5/2),x]`

output

```
(d*(a + b*ArcSech[c*x]))/(3*e^2*(d + e*x^2)^(3/2)) - (a + b*ArcSech[c*x])/
(e^2*Sqrt[d + e*x^2]) - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((-2*e*Sqrt[
1 - c^2*x^2]))/((c^2*d + e)*Sqrt[d + e*x^2]) - (4*ArcTanh[Sqrt[d + e*x^2]/(
Sqrt[d]*Sqrt[1 - c^2*x^2])])/Sqrt[d]))/(6*e^2)
```

### Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

rule 104

```
Int[(((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_))/((e_) + (f_)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

rule 169

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_
))^(p_)*((g_) + (h_)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c +
d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + S
imp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n
*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*
h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x],
x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && LtQ[m, -1] && IntegersQ[
2*m, 2*n, 2*p]
```

rule 220

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-
1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] &&
(LtQ[a, 0] || GtQ[b, 0])
```

rule 435

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((
e_) + (f_)*(x_)^2)^(r_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)
*(a + b*x)^p*(c + d*x)^q*(e + f*x)^r, x], x, x^2], x] /; FreeQ[{a, b, c, d,
e, f, p, q, r}, x] && IntegerQ[(m - 1)/2]
```

rule 6855

```

Int[((a_.) + ArcSech[(c_.)*(x_.)]*(b_.))*((f_.)*(x_.))^(m_.)*((d_.) + (e_.)*(
x_)^2)^(p_.), x_Symbol] :> With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Si
mp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)]
Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x]] /; Fre
eQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] &&
GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2
*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))

```

**Maple [F]**

$$\int \frac{x^3(a + b \operatorname{arcsech}(cx))}{(x^2e + d)^{\frac{5}{2}}} dx$$

input

```
int(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x)
```

output

```
int(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 386 vs.  $2(117) = 234$ .

Time = 0.22 (sec) , antiderivative size = 786, normalized size of antiderivative = 4.39

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \text{Too large to display}$$

input

```
integrate(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="fricas")
```

output

```
[-1/6*(2*(2*b*c^2*d^3 + 2*b*d^2*e + 3*(b*c^2*d^2*e + b*d*e^2)*x^2)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (b*c^2*d^3 + (b*c^2*d*e^2 + b*e^3)*x^4 + b*d^2*e + 2*(b*c^2*d^2*e + b*d*e^2)*x^2)*sqrt(d)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 - 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) + 2*(2*a*c^2*d^3 + 2*a*d^2*e + 3*(a*c^2*d^2*e + a*d*e^2)*x^2 - (b*c*d*e^2*x^3 + b*c*d^2*e*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d))/(c^2*d^4*e^2 + d^3*e^3 + (c^2*d^2*e^4 + d*e^5)*x^4 + 2*(c^2*d^3*e^3 + d^2*e^4)*x^2), 1/3*((b*c^2*d^3 + (b*c^2*d*e^2 + b*e^3)*x^4 + b*d^2*e + 2*(b*c^2*d^2*e + b*d*e^2)*x^2)*sqrt(-d)*arctan(-1/2*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(-d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*d*e*x^4 + (c^2*d^2 - d*e)*x^2 - d^2)) - (2*b*c^2*d^3 + 2*b*d^2*e + 3*(b*c^2*d^2*e + b*d*e^2)*x^2)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (2*a*c^2*d^3 + 2*a*d^2*e + 3*(a*c^2*d^2*e + a*d*e^2)*x^2 - (b*c*d*e^2*x^3 + b*c*d^2*e*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d))/(c^2*d^4*e^2 + d^3*e^3 + (c^2*d^2*e^4 + d*e^5)*x^4 + 2*(c^2*d^3*e^3 + d^2*e^4)*x^2)]
```

## Sympy [F(-1)]

Timed out.

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \text{Timed out}$$

input

```
integrate(x**3*(a+b*asech(c*x))/(e*x**2+d)**(5/2),x)
```

output

Timed out

## Maxima [F]

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^3}{(ex^2 + d)^{5/2}} dx$$

input

```
integrate(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="maxima")
```

output  $-1/3*a*(3*x^2/((e*x^2 + d)^{(3/2)}*e) + 2*d/((e*x^2 + d)^{(3/2)}*e^2)) + b*\text{integrate}(x^3*\log(\text{sqrt}(1/(c*x) + 1))*\text{sqrt}(1/(c*x) - 1) + 1/(c*x))/(e*x^2 + d)^{(5/2)}, x)$

### Giac [F]

$$\int \frac{x^3(a + b\text{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{(b \text{arsech}(cx) + a)x^3}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input `integrate(x^3*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^3/(e*x^2 + d)^(5/2), x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{x^3(a + b\text{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{x^3(a + b \text{acosh}(\frac{1}{cx}))}{(ex^2 + d)^{5/2}} dx$$

input `int((x^3*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(5/2),x)`

output `int((x^3*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(5/2), x)`

### Reduce [F]

$$\int \frac{x^3(a + b\text{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{x^3(\text{asech}(cx) b + a)}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input `int(x^3*(a+b*asech(c*x))/(e*x^2+d)^(5/2),x)`

output `int(x^3*(a+b*asech(c*x))/(e*x^2+d)^(5/2),x)`



**3.169** 
$$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{5/2}} dx$$

Optimal result	1344
Mathematica [A] (verified)	1345
Rubi [A] (verified)	1345
Maple [F]	1348
Fricas [B] (verification not implemented)	1348
Sympy [F(-1)]	1349
Maxima [F]	1349
Giac [F]	1349
Mupad [F(-1)]	1350
Reduce [F]	1350

**Optimal result**

Integrand size = 21, antiderivative size = 154

$$\int \frac{x(a+b\operatorname{sech}^{-1}(cx))}{(d+ex^2)^{5/2}} dx = -\frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{3d(c^2d+e)\sqrt{d+ex^2}} - \frac{a+b\operatorname{sech}^{-1}(cx)}{3e(d+ex^2)^{3/2}} + \frac{b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{3d^{3/2}e}$$

output

```
-1/3*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/d/(c^2*d+e)/(e*x^2+d)^(1/2)-1/3*(a+b*arcsech(c*x))/e/(e*x^2+d)^(3/2)+1/3*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*arctanh((e*x^2+d)^(1/2)/d^(1/2)/(-c^2*x^2+1)^(1/2))/d^(3/2)/e
```

**Mathematica [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.32

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \frac{-ad(c^2d + e) - be\sqrt{\frac{1-cx}{1+cx}}(1 + cx)(d + ex^2) - bd(c^2d + e) \operatorname{sech}^{-1}(cx)}{3de(c^2d + e)(d + ex^2)^{3/2}} - \frac{b\sqrt{\frac{1-cx}{1+cx}}\sqrt{1 - c^2x^2}\sqrt{-d - ex^2} \arctan\left(\frac{\sqrt{d}\sqrt{1-c^2x^2}}{\sqrt{-d-ex^2}}\right)}{3d^{3/2}e(-1 + cx)\sqrt{d + ex^2}}$$

input

```
Integrate[(x*(a + b*ArcSech[c*x]))/(d + e*x^2)^(5/2),x]
```

output

```
(-(a*d*(c^2*d + e)) - b*e*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*(d + e*x^2) - b*d*(c^2*d + e)*ArcSech[c*x])/(3*d*e*(c^2*d + e)*(d + e*x^2)^(3/2)) - (b*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*Sqrt[-d - e*x^2]*ArcTan[(Sqrt[d]*Sqrt[1 - c^2*x^2])/Sqrt[-d - e*x^2]])/(3*d^(3/2)*e*(-1 + c*x)*Sqrt[d + e*x^2])
```

**Rubi [A] (verified)**

Time = 0.55 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.88, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {6853, 2036, 354, 107, 104, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx$$

↓ 6853

$$-\frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{x\sqrt{1-cx}\sqrt{cx+1}(ex^2+d)^{3/2}} dx}{3e} - \frac{a + b \operatorname{sech}^{-1}(cx)}{3e(d + ex^2)^{3/2}}$$

↓ 2036

$$\begin{aligned}
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{x\sqrt{1-c^2x^2}(ex^2+d)^{3/2}} dx}{3e} - \frac{a + b\operatorname{sech}^{-1}(cx)}{3e(d+ex^2)^{3/2}} \\
 & \quad \downarrow \text{354} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{1}{x^2\sqrt{1-c^2x^2}(ex^2+d)^{3/2}} dx^2}{6e} - \frac{a + b\operatorname{sech}^{-1}(cx)}{3e(d+ex^2)^{3/2}} \\
 & \quad \downarrow \text{107} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{1}{x^2\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx^2}{d} + \frac{2e\sqrt{1-c^2x^2}}{d(c^2d+e)\sqrt{d+ex^2}} \right)}{6e} - \frac{a + b\operatorname{sech}^{-1}(cx)}{3e(d+ex^2)^{3/2}} \\
 & \quad \downarrow \text{104} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{2 \int \frac{1}{x^4-d} \frac{d\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}}}{d} + \frac{2e\sqrt{1-c^2x^2}}{d(c^2d+e)\sqrt{d+ex^2}} \right)}{6e} - \frac{a + b\operatorname{sech}^{-1}(cx)}{3e(d+ex^2)^{3/2}} \\
 & \quad \downarrow \text{220} \\
 & \frac{a + b\operatorname{sech}^{-1}(cx)}{3e(d+ex^2)^{3/2}} - \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{2e\sqrt{1-c^2x^2}}{d(c^2d+e)\sqrt{d+ex^2}} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{d^{3/2}} \right)}{6e}
 \end{aligned}$$

input `Int[(x*(a + b*ArcSech[c*x]))/(d + e*x^2)^(5/2),x]`

output `-1/3*(a + b*ArcSech[c*x])/(e*(d + e*x^2)^(3/2)) - (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((2*e*Sqrt[1 - c^2*x^2])/(d*(c^2*d + e)*Sqrt[d + e*x^2]) - (2*ArcTanh[Sqrt[d + e*x^2]/(Sqrt[d]*Sqrt[1 - c^2*x^2])])/d^(3/2)))/(6*e)`

## Defintions of rubi rules used

rule 104  $\text{Int}[\frac{((a_.) + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}}{((e_.) + (f_.)*(x_))}, x_] := \text{With}[\{q = \text{Denominator}[m]\}, \text{Simp}[q \text{ Subst}[\text{Int}[x^{(q*(m+1)-1)} / (b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)} / (c + d*x)^{(1/q)}], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{EqQ}[m + n + 1, 0] \&\& \text{RationalQ}[n] \&\& \text{LtQ}[-1, m, 0] \&\& \text{SimplerQ}[a + b*x, c + d*x]$

rule 107  $\text{Int}[\frac{((a_.) + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}*((e_.) + (f_.)*(x_))^{(p_)}}{x}], x_] := \text{Simp}[b*(a + b*x)^{(m+1)}*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)}) / ((m+1)*(b*c - a*d)*(b*e - a*f)), x] + \text{Simp}[(a*d*f*(m+1) + b*c*f*(n+1) + b*d*e*(p+1)) / ((m+1)*(b*c - a*d)*(b*e - a*f)) \text{Int}[(a + b*x)^{(m+1)}*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \&\& \text{EqQ}[\text{Simplify}[m + n + p + 3], 0] \&\& (\text{LtQ}[m, -1] || \text{SumSimplerQ}[m, 1])$

rule 220  $\text{Int}[\frac{((a_.) + (b_.)*(x_)^2)^{-1}}{x\_Symbol}], x_] := \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[b, 2])^{-1})*\text{ArcTanh}[\text{Rt}[b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a/b] \&\& (\text{LtQ}[a, 0] || \text{GtQ}[b, 0])$

rule 354  $\text{Int}[(x_)^{(m_)}*((a_.) + (b_.)*(x_)^2)^{(p_)}*((c_.) + (d_.)*(x_)^2)^{(q_)}], x\_Symbol] := \text{Simp}[1/2 \text{Subst}[\text{Int}[x^{((m-1)/2)}*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; \text{FreeQ}[\{a, b, c, d, p, q\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{IntegerQ}[(m-1)/2]$

rule 2036  $\text{Int}[(u_)*((c_.) + (d_.)*(x_)^{(n_)}))^{(q_)}*((a1_.) + (b1_.)*(x_)^{(non2_)}))^{(p_)}*((a2_.) + (b2_.)*(x_)^{(non2_)}))^{(p_)}], x\_Symbol] := \text{Int}[u*(a1*a2 + b1*b2*x^n)^p*(c + d*x^n)^q, x] /; \text{FreeQ}[\{a1, b1, a2, b2, c, d, n, p, q\}, x] \&\& \text{EqQ}[\text{non2}, n/2] \&\& \text{EqQ}[a2*b1 + a1*b2, 0] \&\& (\text{IntegerQ}[p] || (\text{GtQ}[a1, 0] \&\& \text{GtQ}[a2, 0]))$

rule 6853  $\text{Int}[(a_.) + \text{ArcSech}[(c_.)*(x_)]*(b_.)*(x_)*((d_.) + (e_.)*(x_)^2)^{(p_)}], x\_Symbol] := \text{Simp}[(d + e*x^2)^{(p+1)}*((a + b*\text{ArcSech}[c*x]) / (2*e*(p+1))), x] + \text{Simp}[b*(\text{Sqrt}[1 + c*x] / (2*e*(p+1))) * \text{Sqrt}[1 / (1 + c*x)] \text{Int}[(d + e*x^2)^{(p+1)} / (x*\text{Sqrt}[1 - c*x]*\text{Sqrt}[1 + c*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, p\}, x] \&\& \text{NeQ}[p, -1]$

**Maple [F]**

$$\int \frac{x(a + b \operatorname{arcsech}(cx))}{(x^2e + d)^{\frac{5}{2}}} dx$$

input `int(x*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x)`

output `int(x*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 339 vs. 2(94) = 188.

Time = 0.19 (sec) , antiderivative size = 692, normalized size of antiderivative = 4.49

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \left[ \frac{4(bc^2d^3 + bd^2e)\sqrt{ex^2 + d} \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}+1}}{cx}\right) - (bc^2d^3 + (bc^2de^2 + be^3)x^4}{(d + ex^2)^{5/2}} \right]$$

input `integrate(x*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="fricas")`

output `[-1/12*(4*(b*c^2*d^3 + b*d^2*e)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (b*c^2*d^3 + (b*c^2*d*e^2 + b*e^3)*x^4 + b*d^2*e + 2*(b*c^2*d^2*e + b*d*e^2)*x^2)*sqrt(d)*log(((c^4*d^2 - 6*c^2*d*e + e^2)*x^4 - 8*(c^2*d^2 - d*e)*x^2 - 4*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 8*d^2)/x^4) + 4*(a*c^2*d^3 + a*d^2*e + (b*c*d*e^2*x^3 + b*c*d^2*e*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d)/(c^2*d^5*e + d^4*e^2 + (c^2*d^3*e^3 + d^2*e^4)*x^4 + 2*(c^2*d^4*e^2 + d^3*e^3)*x^2), 1/6*((b*c^2*d^3 + (b*c^2*d*e^2 + b*e^3)*x^4 + b*d^2*e + 2*(b*c^2*d^2*e + b*d*e^2)*x^2)*sqrt(-d)*arctan(-1/2*((c^3*d - c*e)*x^3 - 2*c*d*x)*sqrt(e*x^2 + d)*sqrt(-d)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))/(c^2*d*e*x^4 + (c^2*d^2 - d*e)*x^2 - d^2) - 2*(b*c^2*d^3 + b*d^2*e)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - 2*(a*c^2*d^3 + a*d^2*e + (b*c*d*e^2*x^3 + b*c*d^2*e*x)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d)/(c^2*d^5*e + d^4*e^2 + (c^2*d^3*e^3 + d^2*e^4)*x^4 + 2*(c^2*d^4*e^2 + d^3*e^3)*x^2)]`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \text{Timed out}$$

input `integrate(x*(a+b*asech(c*x))/(e*x**2+d)**(5/2),x)`

output `Timed out`

**Maxima [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input `integrate(x*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="maxima")`

output `b*integrate(x*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(e*x^2 + d)^(5/2), x) - 1/3*a/((e*x^2 + d)^(3/2)*e)`

**Giac [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input `integrate(x*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x/(e*x^2 + d)^(5/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{x(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^{5/2}} dx$$

input `int((x*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(5/2),x)`

output `int((x*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(5/2), x)`

**Reduce [F]**

$$\int \frac{x(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{x(a \operatorname{sech}(cx) b + a)}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input `int(x*(a+b*asech(c*x))/(e*x^2+d)^(5/2),x)`

output `int(x*(a+b*asech(c*x))/(e*x^2+d)^(5/2),x)`

$$3.170 \quad \int \frac{a+b\operatorname{sech}^{-1}(cx)}{x(d+ex^2)^{5/2}} dx$$

Optimal result	1351
Mathematica [N/A]	1351
Rubi [N/A]	1352
Maple [N/A]	1352
Fricas [N/A]	1353
Sympy [F(-1)]	1353
Maxima [F(-2)]	1353
Giac [N/A]	1354
Mupad [N/A]	1354
Reduce [N/A]	1355

### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{5/2}} dx = \operatorname{Int}\left(\frac{a + b\operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{5/2}}, x\right)$$

output `Defer(Int)((a+b*arcsech(c*x))/x/(e*x^2+d)^(5/2),x)`

### Mathematica [N/A]

Not integrable

Time = 13.68 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{5/2}} dx = \int \frac{a + b\operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{5/2}} dx$$

input `Integrate[(a + b*ArcSech[c*x])/(x*(d + e*x^2)^(5/2)),x]`

output `Integrate[(a + b*ArcSech[c*x])/(x*(d + e*x^2)^(5/2)), x]`



**Rubi [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x (d + ex^2)^{5/2}} dx$$

↓ 6865

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x (d + ex^2)^{5/2}} dx$$

input `Int[(a + b*ArcSech[c*x])/(x*(d + e*x^2)^(5/2)),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{a + b \operatorname{arcsech}(cx)}{x (x^2 e + d)^{5/2}} dx$$

input `int((a+b*arcsech(c*x))/x/(e*x^2+d)^(5/2),x)`

output `int((a+b*arcsech(c*x))/x/(e*x^2+d)^(5/2),x)`

**Fricas [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 53, normalized size of antiderivative = 2.30

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{5/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^{\frac{5}{2}} x} dx$$

input `integrate((a+b*arcsech(c*x))/x/(e*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)/(e^3*x^7 + 3*d*e^2*x^5 + 3*d^2*e*x^3 + d^3*x), x)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{5/2}} dx = \text{Timed out}$$

input `integrate((a+b*asech(c*x))/x/(e*x**2+d)**(5/2),x)`

output `Timed out`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/x/(e*x^2+d)^(5/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{5/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^{5/2} x} dx$$

input

```
integrate((a+b*arcsech(c*x))/x/(e*x^2+d)^(5/2),x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)/((e*x^2 + d)^(5/2)*x), x)
```

**Mupad [N/A]**

Not integrable

Time = 4.08 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x(d + ex^2)^{5/2}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x(e x^2 + d)^{5/2}} dx$$

input

```
int((a + b*acosh(1/(c*x)))/(x*(d + e*x^2)^(5/2)),x)
```

output

```
int((a + b*acosh(1/(c*x)))/(x*(d + e*x^2)^(5/2)), x)
```

**Reduce [N/A]**

Not integrable

Time = 200.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x (d + ex^2)^{5/2}} dx = \int \frac{a \operatorname{sech}(cx) b + a}{x (ex^2 + d)^{5/2}} dx$$

input `int((a+b*asech(c*x))/x/(e*x^2+d)^(5/2),x)`output `int((a+b*asech(c*x))/x/(e*x^2+d)^(5/2),x)`

$$3.171 \quad \int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^3(d+ex^2)^{5/2}} dx$$

Optimal result	1356
Mathematica [N/A]	1356
Rubi [N/A]	1357
Maple [N/A]	1357
Fricas [N/A]	1358
Sympy [F(-1)]	1358
Maxima [F(-2)]	1358
Giac [N/A]	1359
Mupad [N/A]	1359
Reduce [N/A]	1360

### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^3(d + ex^2)^{5/2}} dx = \operatorname{Int}\left(\frac{a + b\operatorname{sech}^{-1}(cx)}{x^3(d + ex^2)^{5/2}}, x\right)$$

output `Defer(Int)((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(5/2),x)`

### Mathematica [N/A]

Not integrable

Time = 15.70 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^3(d + ex^2)^{5/2}} dx = \int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^3(d + ex^2)^{5/2}} dx$$

input `Integrate[(a + b*ArcSech[c*x])/(x^3*(d + e*x^2)^(5/2)),x]`

output `Integrate[(a + b*ArcSech[c*x])/(x^3*(d + e*x^2)^(5/2)), x]`

**Rubi [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{5/2}} dx$$

↓ 6865

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{5/2}} dx$$

input `Int[(a + b*ArcSech[c*x])/(x^3*(d + e*x^2)^(5/2)),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.42 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{a + b \operatorname{arcsech}(cx)}{x^3 (x^2 e + d)^{5/2}} dx$$

input `int((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(5/2),x)`

output `int((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(5/2),x)`

**Fricas [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.39

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{5/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^{5/2} x^3} dx$$

input `integrate((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)/(e^3*x^9 + 3*d*e^2*x^7 + 3*d^2*e*x^5 + d^3*x^3), x)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{5/2}} dx = \text{Timed out}$$

input `integrate((a+b*asech(c*x))/x**3/(e*x**2+d)**(5/2),x)`

output `Timed out`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(5/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{5/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^{5/2} x^3} dx$$

input

```
integrate((a+b*arcsech(c*x))/x^3/(e*x^2+d)^(5/2),x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)/((e*x^2 + d)^(5/2)*x^3), x)
```

**Mupad [N/A]**

Not integrable

Time = 4.44 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{5/2}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x^3 (ex^2 + d)^{5/2}} dx$$

input

```
int((a + b*acosh(1/(c*x)))/(x^3*(d + e*x^2)^(5/2)),x)
```

output

```
int((a + b*acosh(1/(c*x)))/(x^3*(d + e*x^2)^(5/2)), x)
```



**Reduce [N/A]**

Not integrable

Time = 200.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^3 (d + ex^2)^{5/2}} dx = \int \frac{a \operatorname{sech}(cx) b + a}{x^3 (ex^2 + d)^{\frac{5}{2}}} dx$$

input `int((a+b*asech(c*x))/x^3/(e*x^2+d)^(5/2),x)`output `int((a+b*asech(c*x))/x^3/(e*x^2+d)^(5/2),x)`

**3.172** 
$$\int \frac{x^6 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx$$

Optimal result	1361
Mathematica [N/A]	1361
Rubi [N/A]	1362
Maple [N/A]	1362
Fricas [N/A]	1363
Sympy [F(-1)]	1363
Maxima [F(-2)]	1363
Giac [N/A]	1364
Mupad [N/A]	1364
Reduce [N/A]	1365

**Optimal result**

Integrand size = 23, antiderivative size = 23

$$\int \frac{x^6 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \operatorname{Int} \left( \frac{x^6 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}}, x \right)$$

output `Defer(Int)(x^6*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x)`

**Mathematica [N/A]**

Not integrable

Time = 15.05 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{x^6 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{x^6 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx$$

input `Integrate[(x^6*(a + b*ArcSech[c*x]))/(d + e*x^2)^(5/2),x]`

output `Integrate[(x^6*(a + b*ArcSech[c*x]))/(d + e*x^2)^(5/2), x]`

**Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^6(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx$$

↓ 6865

$$\int \frac{x^6(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx$$

input `Int[(x^6*(a + b*ArcSech[c*x]))/(d + e*x^2)^(5/2),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{x^6(a + b \operatorname{arcsech}(cx))}{(x^2e + d)^{5/2}} dx$$

input `int(x^6*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x)`

output `int(x^6*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x)`

**Fricas [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.52

$$\int \frac{x^6(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^6}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input `integrate(x^6*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral((b*x^6*arcsech(c*x) + a*x^6)*sqrt(e*x^2 + d)/(e^3*x^6 + 3*d*e^2*x^4 + 3*d^2*e*x^2 + d^3), x)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{x^6(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \text{Timed out}$$

input `integrate(x**6*(a+b*asech(c*x))/(e*x**2+d)**(5/2),x)`

output `Timed out`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^6(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate(x^6*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.14 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{x^6(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^6}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input

```
integrate(x^6*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)*x^6/(e*x^2 + d)^(5/2), x)
```

**Mupad [N/A]**

Not integrable

Time = 4.20 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{x^6(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{x^6(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^{5/2}} dx$$

input

```
int((x^6*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(5/2),x)
```

output

```
int((x^6*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(5/2), x)
```

**Reduce [N/A]**

Not integrable

Time = 200.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{x^6 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{x^6 (a \operatorname{sech}(cx) b + a)}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input `int(x^6*(a+b*asech(c*x))/(e*x^2+d)^(5/2),x)`output `int(x^6*(a+b*asech(c*x))/(e*x^2+d)^(5/2),x)`

$$3.173 \quad \int \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx$$

Optimal result	1366
Mathematica [N/A]	1366
Rubi [N/A]	1367
Maple [N/A]	1367
Fricas [N/A]	1368
Sympy [F(-1)]	1368
Maxima [F(-2)]	1368
Giac [N/A]	1369
Mupad [N/A]	1369
Reduce [N/A]	1370

### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \operatorname{Int} \left( \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}}, x \right)$$

output

```
Defer(Int)(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x)
```

### Mathematica [N/A]

Not integrable

Time = 14.80 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{x^4 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx$$

input

```
Integrate[(x^4*(a + b*ArcSech[c*x]))/(d + e*x^2)^(5/2),x]
```

output

```
Integrate[(x^4*(a + b*ArcSech[c*x]))/(d + e*x^2)^(5/2), x]
```

**Rubi [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx$$

↓ 6865

$$\int \frac{x^4(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx$$

input `Int[(x^4*(a + b*ArcSech[c*x]))/(d + e*x^2)^(5/2),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{x^4(a + b \operatorname{arcsech}(cx))}{(x^2e + d)^{5/2}} dx$$

input `int(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x)`

output `int(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x)`



**Fricas [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.52

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^4}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input `integrate(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral((b*x^4*arcsech(c*x) + a*x^4)*sqrt(e*x^2 + d)/(e^3*x^6 + 3*d*e^2*x^4 + 3*d^2*e*x^2 + d^3), x)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \text{Timed out}$$

input `integrate(x**4*(a+b*asech(c*x))/(e*x**2+d)**(5/2),x)`

output `Timed out`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**Giac [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{x^4(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^4}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input

```
integrate(x^4*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="giac")
```

output

```
integrate((b*arcsech(c*x) + a)*x^4/(e*x^2 + d)^(5/2), x)
```

**Mupad [N/A]**

Not integrable

Time = 4.08 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{x^4(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{x^4(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^{5/2}} dx$$

input

```
int((x^4*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(5/2),x)
```

output

```
int((x^4*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(5/2), x)
```

**Reduce [N/A]**

Not integrable

Time = 200.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{x^4(a + b\operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{x^4(\operatorname{asech}(cx)b + a)}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input `int(x^4*(a+b*asech(c*x))/(e*x^2+d)^(5/2),x)`output `int(x^4*(a+b*asech(c*x))/(e*x^2+d)^(5/2),x)`

**3.174** 
$$\int \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx$$

Optimal result	1371
Mathematica [C] (verified)	1372
Rubi [A] (verified)	1372
Maple [F]	1376
Fricas [B] (verification not implemented)	1376
Sympy [F(-1)]	1377
Maxima [F]	1377
Giac [F]	1377
Mupad [F(-1)]	1378
Reduce [F]	1378

**Optimal result**

Integrand size = 23, antiderivative size = 243

$$\int \frac{x^2 (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = -\frac{bx \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{3d(c^2d + e) \sqrt{d + ex^2}} + \frac{x^3 (a + b \operatorname{sech}^{-1}(cx))}{3d(d + ex^2)^{3/2}} - \frac{bc \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1 + \frac{ex^2}{d}} E(\arcsin(cx) | -\frac{e}{c^2d})}{3e(c^2d + e) \sqrt{d + ex^2}} + \frac{b \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1 + \frac{ex^2}{d}} \operatorname{EllipticF}(\arcsin(cx), -\frac{e}{c^2d})}{3cde \sqrt{d + ex^2}}$$

output

```
-1/3*b*x*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/d/(c^2*d+e)/(e*x^2+d)^(1/2)+1/3*x^3*(a+b*arcsech(c*x))/d/(e*x^2+d)^(3/2)-1/3*b*c*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(1+e*x^2/d)^(1/2)*EllipticE(c*x,(-e/c^2/d)^(1/2))/e/(c^2*d+e)/(e*x^2+d)^(1/2)+1/3*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(1+e*x^2/d)^(1/2)*EllipticF(c*x,(-e/c^2/d)^(1/2))/c/d/e/(e*x^2+d)^(1/2)
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 1.98 (sec) , antiderivative size = 488, normalized size of antiderivative = 2.01

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \frac{ax^3 - \frac{b\sqrt{\frac{1-cx}{1+cx}}(-cd+ex)(d+ex^2)}{e(c^2d+e)} + bx^3 \operatorname{sech}^{-1}(cx) + \frac{b\sqrt{\frac{1-cx}{1+cx}}(1+cx)\sqrt{\frac{c(\sqrt{d+i\sqrt{ex}})}{(c\sqrt{d+i\sqrt{e}})(1+cx)}}}{\sqrt{\frac{d+ex^2}{d+ex^2}}}}{e(c^2d+e)}$$

input `Integrate[(x^2*(a + b*ArcSech[c*x]))/(d + e*x^2)^(5/2),x]`

output

```
(a*x^3 - (b*Sqrt[(1 - c*x)/(1 + c*x)]*(-(c*d) + e*x)*(d + e*x^2))/(e*(c^2*d + e)) + b*x^3*ArcSech[c*x] + (b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*Sqrt[(c*(Sqrt[d] + I*Sqrt[e]*x))/((c*Sqrt[d] + I*Sqrt[e])*(1 + c*x))]*Sqrt[(c*(I*Sqrt[d] + Sqrt[e]*x))/((I*c*Sqrt[d] + Sqrt[e])*(1 + c*x))]*(d + e*x^2)*((I*c*Sqrt[d] + Sqrt[e])*EllipticE[I*ArcSinh[Sqrt[((c^2*d + e)*(1 - c*x)))/((c*Sqrt[d] + I*Sqrt[e])^2*(1 + c*x))]], (c*Sqrt[d] + I*Sqrt[e])^2/(c*Sqrt[d] - I*Sqrt[e])^2] - 2*Sqrt[e]*EllipticF[I*ArcSinh[Sqrt[((c^2*d + e)*(1 - c*x)))/((c*Sqrt[d] + I*Sqrt[e])^2*(1 + c*x))]], (c*Sqrt[d] + I*Sqrt[e])^2/(c*Sqrt[d] - I*Sqrt[e])^2))/(c*(c*Sqrt[d] + I*Sqrt[e])*e*Sqrt[(I*c*Sqrt[d] + Sqrt[e])*(-1 + c*x)]/((-I)*c*Sqrt[d] + Sqrt[e])*(1 + c*x)))/(3*d*(d + e*x^2)^(3/2))
```

**Rubi [A] (verified)**

Time = 0.53 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.84, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.348$ , Rules used = {6855, 27, 373, 326, 323, 321, 330, 327}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx$$

$$\begin{aligned}
 & \downarrow \text{6855} \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^2}{3d\sqrt{1-c^2x^2}(ex^2+d)^{3/2}} dx + \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} \\
 & \downarrow \text{27} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{x^2}{\sqrt{1-c^2x^2}(ex^2+d)^{3/2}} dx}{3d} + \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} \\
 & \downarrow \text{373} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{\sqrt{1-c^2x^2}}{\sqrt{ex^2+d}} dx}{c^2d+e} - \frac{x\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{3d} + \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} \\
 & \downarrow \text{326} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{(c^2d+e) \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx}{e} - \frac{c^2 \int \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} dx}{e} - \frac{x\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{3d} + \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} \\
 & \downarrow \text{323} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{(c^2d+e)\sqrt{\frac{ex^2}{d}+1} \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{\frac{ex^2}{d}+1}} dx}{e\sqrt{d+ex^2}} - \frac{c^2 \int \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} dx}{e} - \frac{x\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{3d} + \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} \\
 & \downarrow \text{321} \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{(c^2d+e)\sqrt{\frac{ex^2}{d}+1} \operatorname{EllipticF}\left(\arcsin(cx), -\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}} - \frac{c^2 \int \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} dx}{e} - \frac{x\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{3d} + \frac{x^3(a+b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} \\
 & \downarrow \text{330}
 \end{aligned}$$

$$\begin{aligned}
& b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{(c^2d+e)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx), -\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}} - \frac{c^2\sqrt{d+ex^2} \int \frac{\sqrt{\frac{ex^2}{d}+1}}{\sqrt{1-c^2x^2}} dx}{e\sqrt{\frac{ex^2}{d}+1}} - \frac{x\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right) \\
& \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} \\
& \quad \downarrow \text{327} \\
& \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} + \\
& b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{(c^2d+e)\sqrt{\frac{ex^2}{d}+1}\operatorname{EllipticF}\left(\arcsin(cx), -\frac{e}{c^2d}\right)}{ce\sqrt{d+ex^2}} - \frac{c\sqrt{d+ex^2}E\left(\arcsin(cx)\middle|-\frac{e}{c^2d}\right)}{e\sqrt{\frac{ex^2}{d}+1}} - \frac{x\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right) \\
& \frac{3d}{3d}
\end{aligned}$$

input `Int[(x^2*(a + b*ArcSech[c*x]))/(d + e*x^2)^(5/2),x]`

output `(x^3*(a + b*ArcSech[c*x]))/(3*d*(d + e*x^2)^(3/2)) + (b*sqrt[(1 + c*x)^(-1)]*sqrt[1 + c*x]*(-(x*sqrt[1 - c^2*x^2])/((c^2*d + e)*sqrt[d + e*x^2])) + (-(c*sqrt[d + e*x^2]*EllipticE[ArcSin[c*x], -(e/(c^2*d))]))/(e*sqrt[1 + (e*x^2)/d])) + ((c^2*d + e)*sqrt[1 + (e*x^2)/d]*EllipticF[ArcSin[c*x], -(e/(c^2*d))])/(c*e*sqrt[d + e*x^2]))/(c^2*d + e))/(3*d)`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 321 `Int[1/(sqrt[(a_) + (b_.)*(x_)^2]*sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(1/(sqrt[a]*sqrt[c]*Rt[-d/c, 2]))*EllipticF[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-b/a, -d/c])`

rule 323 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[  
imp[Sqrt[1 + (d/c)*x^2]/Sqrt[c + d*x^2] Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d/c)*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]`

rule 326 `Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[  
b/d Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Simp[(b*c - a*d)/d Int[  
1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] &&  
PosQ[d/c] && NegQ[b/a]`

rule 327 `Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[  
(Sqrt[a]/(Sqrt[c]*Rt[-d/c, 2]))*EllipticE[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]`

rule 330 `Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[  
Sqrt[a + b*x^2]/Sqrt[1 + (b/a)*x^2] Int[Sqrt[1 + (b/a)*x^2]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]`

rule 373 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[e*(e*x)^(m - 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*(b*c - a*d)*(p + 1))), x] - Simp[e^2/(2*(b*c - a*d)*(p + 1)) Int[(e*x)^(m - 2)*(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[c*(m - 1) + d*(m + 2*p + 2*q + 3)*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[m, 1] && LeQ[m, 3] && IntBinomialQ[a, b, c, d, e, m, 2, p, q, x]`

rule 6855 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))`



**Maple [F]**

$$\int \frac{x^2(a + b \operatorname{arcsech}(cx))}{(x^2e + d)^{\frac{5}{2}}} dx$$

input `int(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2), x)`

output `int(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2), x)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 335 vs.  $2(163) = 326$ .

Time = 0.15 (sec) , antiderivative size = 335, normalized size of antiderivative = 1.38

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \frac{(bc^3d^2e + bcde^2)\sqrt{ex^2 + d}x^3 \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{e^2x^2}+1}}{cx}\right) + \left((ac^3d^2e + acde^2)x^3 - ($$

input `integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2), x, algorithm="fricas")`

output `1/3*((b*c^3*d^2*e + b*c*d*e^2)*sqrt(e*x^2 + d)*x^3*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) + ((a*c^3*d^2*e + a*c*d*e^2)*x^3 - (b*c^2*d*e^2*x^4 + b*c^2*d^2*e*x^2)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d) - ((b*c^4*d*e^2*x^4 + 2*b*c^4*d^2*e*x^2 + b*c^4*d^3)*elliptic_e(arcsin(c*x), -e/(c^2*d)) - (b*c^4*d^3 + (b*c^4*d*e^2 + b*e^3)*x^4 + b*d^2*e + 2*(b*c^4*d^2*e + b*d*e^2)*x^2)*elliptic_f(arcsin(c*x), -e/(c^2*d)))*sqrt(d))/(c^3*d^5*e + c*d^4*e^2 + (c^3*d^3*e^3 + c*d^2*e^4)*x^4 + 2*(c^3*d^4*e^2 + c*d^3*e^3)*x^2)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \text{Timed out}$$

input `integrate(x**2*(a+b*asech(c*x))/(e*x**2+d)**(5/2),x)`

output `Timed out`

**Maxima [F]**

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^2}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input `integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="maxima")`

output `-1/3*a*(x/((e*x^2 + d)^(3/2)*e) - x/(sqrt(e*x^2 + d)*d*e)) + b*integrate(x^2*log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(e*x^2 + d)^(5/2), x)`

**Giac [F]**

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^2}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input `integrate(x^2*(a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^2/(e*x^2 + d)^(5/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^{5/2}} dx$$

input `int((x^2*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(5/2), x)`

output `int((x^2*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(5/2), x)`

**Reduce [F]**

$$\int \frac{x^2(a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{5/2}} dx = \int \frac{x^2(a \operatorname{sech}(cx) b + a)}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input `int(x^2*(a+b*asech(c*x))/(e*x^2+d)^(5/2), x)`

output `int(x^2*(a+b*asech(c*x))/(e*x^2+d)^(5/2), x)`

**3.175**  $\int \frac{a+b\operatorname{sech}^{-1}(cx)}{(d+ex^2)^{5/2}} dx$

Optimal result	1379
Mathematica [C] (verified)	1380
Rubi [A] (verified)	1380
Maple [F]	1384
Fricas [B] (verification not implemented)	1385
Sympy [F(-1)]	1385
Maxima [F]	1386
Giac [F]	1386
Mupad [F(-1)]	1386
Reduce [F]	1387

**Optimal result**

Integrand size = 20, antiderivative size = 266

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{(d + ex^2)^{5/2}} dx = \frac{bcx\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1-c^2x^2}}{3d^2(c^2d+e)\sqrt{d+ex^2}} + \frac{x(a + b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}}$$

$$+ \frac{2x(a + b\operatorname{sech}^{-1}(cx))}{3d^2\sqrt{d+ex^2}} + \frac{bc\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{d+ex^2}E(\arcsin(cx) | -\frac{e}{c^2d})}{3d^2(c^2d+e)\sqrt{1+\frac{ex^2}{d}}}$$

$$+ \frac{2b\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\sqrt{1+\frac{ex^2}{d}}\operatorname{EllipticF}(\arcsin(cx), -\frac{e}{c^2d})}{3cd^2\sqrt{d+ex^2}}$$

output

```
1/3*b*e*x*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/d^2/(c^2*d+e)
/(e*x^2+d)^(1/2)+1/3*x*(a+b*arcsech(c*x))/d/(e*x^2+d)^(3/2)+2/3*x*(a+b*arc
sech(c*x))/d^2/(e*x^2+d)^(1/2)+1/3*b*c*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(e*
x^2+d)^(1/2)*EllipticE(c*x,(-e/c^2/d)^(1/2))/d^2/(c^2*d+e)/(1+e*x^2/d)^(1/
2)+2/3*b*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(1+e*x^2/d)^(1/2)*EllipticF(c*x,(
-e/c^2/d)^(1/2))/c/d^2/(e*x^2+d)^(1/2)
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 24.74 (sec) , antiderivative size = 517, normalized size of antiderivative = 1.94

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{5/2}} dx = \frac{b \sqrt{\frac{1-cx}{1+cx}} (-cd+ex)(d+ex^2)}{c^2 d+e} + ax(3d + 2ex^2) + bx(3d + 2ex^2) \operatorname{sech}^{-1}(cx) - \frac{ib \sqrt{\frac{1-cx}{1+cx}} (1+cx)}{c^2 d+e}$$

input

```
Integrate[(a + b*ArcSech[c*x])/(d + e*x^2)^(5/2), x]
```

output

```
((b*Sqrt[(1 - c*x)/(1 + c*x)]*(-(c*d) + e*x)*(d + e*x^2))/(c^2*d + e) + a*x*(3*d + 2*e*x^2) + b*x*(3*d + 2*e*x^2)*ArcSech[c*x] - (I*b*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*Sqrt[(c*(Sqrt[d] - I*Sqrt[e]*x))/((c*Sqrt[d] - I*Sqrt[e])*(1 + c*x))]*Sqrt[(c*(Sqrt[d] + I*Sqrt[e]*x))/((c*Sqrt[d] + I*Sqrt[e])*(1 + c*x))]*(d + e*x^2)*((c*Sqrt[d] - I*Sqrt[e])*EllipticE[I*ArcSinh[Sqrt[((c^2*d + e)*(1 - c*x))/((c*Sqrt[d] + I*Sqrt[e])^2*(1 + c*x))]]], (c*Sqrt[d] + I*Sqrt[e])^2/(c*Sqrt[d] - I*Sqrt[e])^2] - 2*(3*c*Sqrt[d] + (2*I)*Sqrt[e])*EllipticF[I*ArcSinh[Sqrt[((c^2*d + e)*(1 - c*x))/((c*Sqrt[d] + I*Sqrt[e])^2*(1 + c*x))]]], (c*Sqrt[d] + I*Sqrt[e])^2/(c*Sqrt[d] - I*Sqrt[e])^2))/((c*(c*Sqrt[d] + I*Sqrt[e])*Sqrt[-(((c*Sqrt[d] - I*Sqrt[e])*(-1 + c*x))/((c*Sqrt[d] + I*Sqrt[e])*(1 + c*x))))]/(3*d^2*(d + e*x^2)^(3/2))
```

**Rubi [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 224, normalized size of antiderivative = 0.84, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6845, 27, 402, 25, 27, 399, 323, 321, 330, 327}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{5/2}} dx$$

$$\begin{aligned}
& \downarrow 6845 \\
& b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{2ex^2+3d}{3d^2\sqrt{1-c^2x^2}(ex^2+d)^{3/2}} dx + \frac{2x(a+b\operatorname{sech}^{-1}(cx))}{3d^2\sqrt{d+ex^2}} + \\
& \quad \frac{x(a+b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} \\
& \downarrow 27 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{2ex^2+3d}{\sqrt{1-c^2x^2}(ex^2+d)^{3/2}} dx}{3d^2} + \frac{2x(a+b\operatorname{sech}^{-1}(cx))}{3d^2\sqrt{d+ex^2}} + \frac{x(a+b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} \\
& \downarrow 402 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{ex\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} - \frac{\int -\frac{d(ex^2c^2+3dc^2+2e)}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx}{d(c^2d+e)} \right)}{3d^2} + \frac{2x(a+b\operatorname{sech}^{-1}(cx))}{3d^2\sqrt{d+ex^2}} + \\
& \quad \frac{x(a+b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} \\
& \downarrow 25 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{d(ex^2c^2+3dc^2+2e)}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx}{d(c^2d+e)} + \frac{ex\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{3d^2} + \frac{2x(a+b\operatorname{sech}^{-1}(cx))}{3d^2\sqrt{d+ex^2}} + \\
& \quad \frac{x(a+b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} \\
& \downarrow 27 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{ex^2c^2+3dc^2+2e}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx}{c^2d+e} + \frac{ex\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{3d^2} + \frac{2x(a+b\operatorname{sech}^{-1}(cx))}{3d^2\sqrt{d+ex^2}} + \\
& \quad \frac{x(a+b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} \\
& \downarrow 399 \\
& \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{c^2 \int \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} dx + 2(c^2d+e) \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{ex^2+d}} dx}{c^2d+e} + \frac{ex\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{3d^2} + \\
& \quad \frac{2x(a+b\operatorname{sech}^{-1}(cx))}{3d^2\sqrt{d+ex^2}} + \frac{x(a+b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}}
\end{aligned}$$

$$\begin{aligned}
 & \downarrow 323 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{c^2 \int \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} dx + \frac{2(c^2d+e)\sqrt{\frac{ex^2}{d}+1} \int \frac{1}{\sqrt{1-c^2x^2}\sqrt{\frac{ex^2}{d}+1}} dx}{c^2d+e} + \frac{ex\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{3d^2} + \\
 & \frac{2x(a + b\operatorname{sech}^{-1}(cx))}{3d^2\sqrt{d+ex^2}} + \frac{x(a + b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} \\
 & \downarrow 321 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{c^2 \int \frac{\sqrt{ex^2+d}}{\sqrt{1-c^2x^2}} dx + \frac{2(c^2d+e)\sqrt{\frac{ex^2}{d}+1} \operatorname{EllipticF}\left(\arcsin(cx), -\frac{e}{c^2d}\right)}{c\sqrt{d+ex^2}}}{c^2d+e} + \frac{ex\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{3d^2} + \\
 & \frac{2x(a + b\operatorname{sech}^{-1}(cx))}{3d^2\sqrt{d+ex^2}} + \frac{x(a + b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} \\
 & \downarrow 330 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\frac{c^2\sqrt{d+ex^2} \int \frac{\sqrt{\frac{ex^2}{d}+1}}{\sqrt{1-c^2x^2}} dx + 2(c^2d+e)\sqrt{\frac{ex^2}{d}+1} \operatorname{EllipticF}\left(\arcsin(cx), -\frac{e}{c^2d}\right)}{\sqrt{\frac{ex^2}{d}+1}}}{c^2d+e} + \frac{ex\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{3d^2} + \\
 & \frac{2x(a + b\operatorname{sech}^{-1}(cx))}{3d^2\sqrt{d+ex^2}} + \frac{x(a + b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} \\
 & \downarrow 327 \\
 & \frac{2x(a + b\operatorname{sech}^{-1}(cx))}{3d^2\sqrt{d+ex^2}} + \frac{x(a + b\operatorname{sech}^{-1}(cx))}{3d(d+ex^2)^{3/2}} + \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\frac{2(c^2d+e)\sqrt{\frac{ex^2}{d}+1} \operatorname{EllipticF}\left(\arcsin(cx), -\frac{e}{c^2d}\right)}{c\sqrt{d+ex^2}} + \frac{c\sqrt{d+ex^2} E\left(\arcsin(cx) \middle| -\frac{e}{c^2d}\right)}{\sqrt{\frac{ex^2}{d}+1}}}{c^2d+e} + \frac{ex\sqrt{1-c^2x^2}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{3d^2}
 \end{aligned}$$

input `Int[(a + b*ArcSech[c*x])/(d + e*x^2)^(5/2), x]`

output

```
(x*(a + b*ArcSech[c*x]))/(3*d*(d + e*x^2)^(3/2)) + (2*x*(a + b*ArcSech[c*x
]))/(3*d^2*Sqrt[d + e*x^2]) + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*((e*x*
Sqrt[1 - c^2*x^2])/((c^2*d + e)*Sqrt[d + e*x^2]) + ((c*Sqrt[d + e*x^2]*Ell
ipticE[ArcSin[c*x], -(e/(c^2*d))])/Sqrt[1 + (e*x^2)/d] + (2*(c^2*d + e)*Sq
rt[1 + (e*x^2)/d]*EllipticF[ArcSin[c*x], -(e/(c^2*d))])/(c*Sqrt[d + e*x^2
]))/(c^2*d + e))/(3*d^2)
```

### Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

rule 321

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := S
imp[(1/(Sqrt[a]*Sqrt[c]*Rt[-d/c, 2]))*EllipticF[ArcSin[Rt[-d/c, 2]*x], b*(c
/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a,
0] && !(NegQ[b/a] && SimplerSqrtQ[-b/a, -d/c])
```

rule 323

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := S
imp[Sqrt[1 + (d/c)*x^2]/Sqrt[c + d*x^2] Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (
d/c)*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]
```

rule 327

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[
(Sqrt[a]/(Sqrt[c]*Rt[-d/c, 2]))*EllipticE[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d)
)], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]
```

rule 330

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[
Sqrt[a + b*x^2]/Sqrt[1 + (b/a)*x^2] Int[Sqrt[1 + (b/a)*x^2]/Sqrt[c + d*x^
2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a,
0]
```



rule 399 `Int[((e_) + (f_)*(x_)^2)/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] := Simp[f/b Int[Sqrt[a + b*x^2]/Sqrt[c + d*x^2], x], x] + Simp[(b*e - a*f)/b Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && !((PosQ[b/a] && PosQ[d/c]) || (NegQ[b/a] && (PosQ[d/c] || (GtQ[a, 0] && (!GtQ[c, 0] || SimplerSqrtQ[-b/a, -d/c])))))`

rule 402 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2), x_Symbol] := Simp[(-(b*e - a*f))*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(a^2*(b*c - a*d)*(p + 1))), x] + Simp[1/(a^2*(b*c - a*d)*(p + 1)) Int[(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[c*(b*e - a*f) + e^2*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(2*(p + q + 2) + 1)*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, q}, x] && LtQ[p, -1]`

rule 6845 `Int[((a_) + ArcSech[(c_)*(x_)])*(b_))*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e}, x] && (IGtQ[p, 0] || ILtQ[p + 1/2, 0])`

## Maple **[F]**

$$\int \frac{a + b \operatorname{arcsech}(cx)}{(x^2 e + d)^{\frac{5}{2}}} dx$$

input `int((a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x)`

output `int((a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 397 vs.  $2(182) = 364$ .

Time = 0.13 (sec) , antiderivative size = 397, normalized size of antiderivative = 1.49

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{5/2}} dx = \frac{(2(bc^3d^2e + bcde^2)x^3 + 3(bc^3d^3 + bcd^2e)x)\sqrt{ex^2 + d} \log\left(\frac{cx\sqrt{\frac{-c^2x^2-1}{c^2x^2}+1}}{cx}\right) + (2(a$$

input `integrate((a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="fricas")`

output `1/3*((2*(b*c^3*d^2*e + b*c*d*e^2)*x^3 + 3*(b*c^3*d^3 + b*c*d^2*e)*x)*sqrt(e*x^2 + d)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2))+ 1)/(c*x)) + (2*(a*c^3*d^2*e + a*c*d*e^2)*x^3 + 3*(a*c^3*d^3 + a*c*d^2*e)*x + (b*c^2*d*e^2*x^4 + b*c^2*d^2*e*x^2)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)))*sqrt(e*x^2 + d) + ((b*c^4*d*e^2*x^4 + 2*b*c^4*d^2*e*x^2 + b*c^4*d^3)*elliptic_e(arcsin(c*x), -e/(c^2*d)) - ((b*c^4 - 3*b*c^2)*d*e^2 - 2*b*e^3)*x^4 + (b*c^4 - 3*b*c^2)*d^3 - 2*b*d^2*e + 2*((b*c^4 - 3*b*c^2)*d^2*e - 2*b*d*e^2)*x^2)*elliptic_f(arcsin(c*x), -e/(c^2*d))*sqrt(d))/(c^3*d^6 + c*d^5*e + (c^3*d^4*e^2 + c*d^3*e^3)*x^4 + 2*(c^3*d^5*e + c*d^4*e^2)*x^2)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{5/2}} dx = \text{Timed out}$$

input `integrate((a+b*asech(c*x))/(e*x**2+d)**(5/2),x)`

output `Timed out`

**Maxima [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{5/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^{5/2}} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="maxima")`

output `1/3*a*(2*x/(sqrt(e*x^2 + d)*d^2) + x/((e*x^2 + d)^(3/2)*d)) + b*integrate(log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(e*x^2 + d)^(5/2), x)`

**Giac [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{5/2}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{(ex^2 + d)^{5/2}} dx$$

input `integrate((a+b*arcsech(c*x))/(e*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/(e*x^2 + d)^(5/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{5/2}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{(ex^2 + d)^{5/2}} dx$$

input `int((a + b*acosh(1/(c*x)))/(d + e*x^2)^(5/2),x)`

output `int((a + b*acosh(1/(c*x)))/(d + e*x^2)^(5/2), x)`

**Reduce [F]**

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{(d + ex^2)^{5/2}} dx = \int \frac{a \operatorname{sech}(cx) b + a}{(ex^2 + d)^{5/2}} dx$$

input `int((a+b*asech(c*x))/(e*x^2+d)^(5/2),x)`

output `int((a+b*asech(c*x))/(e*x^2+d)^(5/2),x)`

### 3.176 $\int (fx)^m (d + ex^2)^3 (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	1388
Mathematica [A] (verified)	1389
Rubi [A] (verified)	1390
Maple [F]	1394
Fricas [F]	1395
Sympy [F]	1395
Maxima [F]	1395
Giac [F]	1396
Mupad [F(-1)]	1397
Reduce [F]	1397

#### Optimal result

Integrand size = 23, antiderivative size = 593

$$\int (fx)^m (d + ex^2)^3 (a + b\operatorname{sech}^{-1}(cx)) dx =$$

$$\frac{be\left(e^2(15 + 8m + m^2)^2 + 3c^2de(3 + m)^2(42 + 13m + m^2) + 3c^4d^2(840 + 638m + 179m^2 + 22m^3 + m^4)\right)}{c^6 f(2 + m)(3 + m)(4 + m)(5 + m)(6 + m)(7 + m)}$$

$$- \frac{be^2(e(5 + m)^2 + 3c^2d(42 + 13m + m^2))(fx)^{3+m} \sqrt{\frac{1}{1+cx}} \sqrt{1 + cx} \sqrt{1 - c^2x^2}}{c^4 f^3(4 + m)(5 + m)(6 + m)(7 + m)}$$

$$- \frac{be^3(fx)^{5+m} \sqrt{\frac{1}{1+cx}} \sqrt{1 + cx} \sqrt{1 - c^2x^2}}{c^2 f^5(6 + m)(7 + m)}$$

$$+ \frac{d^3(fx)^{1+m} (a + b\operatorname{sech}^{-1}(cx))}{f(1 + m)} + \frac{3d^2e(fx)^{3+m} (a + b\operatorname{sech}^{-1}(cx))}{f^3(3 + m)}$$

$$+ \frac{3de^2(fx)^{5+m} (a + b\operatorname{sech}^{-1}(cx))}{f^5(5 + m)} + \frac{e^3(fx)^{7+m} (a + b\operatorname{sech}^{-1}(cx))}{f^7(7 + m)}$$

$$+ \frac{b\left(c^6 d^3(2 + m)(4 + m)(6 + m) + \frac{e(1+m)^2(e^2(15+8m+m^2)^2 + 3c^2de(3+m)^2(42+13m+m^2) + 3c^4d^2(840+638m+179m^2+22m^3+m^4))}{(3+m)(5+m)(7+m)}\right)}{c^6 f(1 + m)^2(2 + m)(4 + m)(6 + m)}$$

output

```

-b*e*(e^2*(m^2+8*m+15)^2+3*c^2*d*e*(3+m)^2*(m^2+13*m+42)+3*c^4*d^2*(m^4+22
*m^3+179*m^2+638*m+840))*(f*x)^(1+m)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2
*x^2+1)^(1/2)/c^6/f/(2+m)/(3+m)/(4+m)/(5+m)/(6+m)/(7+m)-b*e^2*(e*(5+m)^2+3
*c^2*d*(m^2+13*m+42))*(f*x)^(3+m)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^
2+1)^(1/2)/c^4/f^3/(4+m)/(5+m)/(6+m)/(7+m)-b*e^3*(f*x)^(5+m)*(1/(c*x+1))^(
1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^2/f^5/(6+m)/(7+m)+d^3*(f*x)^(1+m)*
(a+b*arcsech(c*x))/f/(1+m)+3*d^2*e*(f*x)^(3+m)*(a+b*arcsech(c*x))/f^3/(3+m
)+3*d*e^2*(f*x)^(5+m)*(a+b*arcsech(c*x))/f^5/(5+m)+e^3*(f*x)^(7+m)*(a+b*ar
csech(c*x))/f^7/(7+m)+b*(c^6*d^3*(2+m)*(4+m)*(6+m)+e*(1+m)^2*(e^2*(m^2+8*m
+15)^2+3*c^2*d*e*(3+m)^2*(m^2+13*m+42)+3*c^4*d^2*(m^4+22*m^3+179*m^2+638*m
+840))/(3+m)/(5+m)/(7+m))*(f*x)^(1+m)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*hype
rgeom([1/2, 1/2+1/2*m], [3/2+1/2*m], c^2*x^2)/c^6/f/(1+m)^2/(2+m)/(4+m)/(6+m
)

```

**Mathematica [A] (verified)**

Time = 2.90 (sec) , antiderivative size = 441, normalized size of antiderivative = 0.74

$$\begin{aligned}
& \int (fx)^m (d + ex^2)^3 (a + b\operatorname{sech}^{-1}(cx)) dx \\
&= x(fx)^m \left( \frac{ad^3}{1+m} + \frac{3ad^2ex^2}{3+m} + \frac{3ade^2x^4}{5+m} + \frac{ae^3x^6}{7+m} + \frac{bd^3\operatorname{sech}^{-1}(cx)}{1+m} \right. \\
&\quad + \frac{3bd^2ex^2\operatorname{sech}^{-1}(cx)}{3+m} + \frac{3bde^2x^4\operatorname{sech}^{-1}(cx)}{5+m} + \frac{be^3x^6\operatorname{sech}^{-1}(cx)}{7+m} \\
&\quad - \frac{bd^3\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2x^2\right)}{(1+m)^2(-1+cx)} \\
&\quad - \frac{3bd^2ex^2\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3+m}{2}, \frac{5+m}{2}, c^2x^2\right)}{(3+m)^2(-1+cx)} \\
&\quad - \frac{3bde^2x^4\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5+m}{2}, \frac{7+m}{2}, c^2x^2\right)}{(5+m)^2(-1+cx)} \\
&\quad \left. - \frac{be^3x^6\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7+m}{2}, \frac{9+m}{2}, c^2x^2\right)}{(7+m)^2(-1+cx)} \right)
\end{aligned}$$

input

```
Integrate[(f*x)^m*(d + e*x^2)^3*(a + b*ArcSech[c*x]), x]
```

output

```
x*(f*x)^m*((a*d^3)/(1 + m) + (3*a*d^2*e*x^2)/(3 + m) + (3*a*d*e^2*x^4)/(5 + m) + (a*e^3*x^6)/(7 + m) + (b*d^3*ArcSech[c*x])/(1 + m) + (3*b*d^2*e*x^2*ArcSech[c*x])/(3 + m) + (3*b*d*e^2*x^4*ArcSech[c*x])/(5 + m) + (b*e^3*x^6*ArcSech[c*x])/(7 + m) - (b*d^3*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])/((1 + m)^2*(-1 + c*x)) - (3*b*d^2*e*x^2*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (3 + m)/2, (5 + m)/2, c^2*x^2])/((3 + m)^2*(-1 + c*x)) - (3*b*d*e^2*x^4*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (5 + m)/2, (7 + m)/2, c^2*x^2])/((5 + m)^2*(-1 + c*x)) - (b*e^3*x^6*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (7 + m)/2, (9 + m)/2, c^2*x^2])/((7 + m)^2*(-1 + c*x))
```

**Rubi [A] (verified)**

Time = 2.18 (sec) , antiderivative size = 527, normalized size of antiderivative = 0.89, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$ , Rules used = {6855, 2340, 25, 1590, 25, 363, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2)^3 (fx)^m (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$\downarrow 6855$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(fx)^m \left( \frac{e^3 x^6}{m+7} + \frac{3de^2 x^4}{m+5} + \frac{3d^2 ex^2}{m+3} + \frac{d^3}{m+1} \right)}{\sqrt{1-c^2 x^2}} dx +$$

$$\frac{d^3 (fx)^{m+1} (a + b\operatorname{sech}^{-1}(cx))}{f(m+1)} + \frac{3d^2 e (fx)^{m+3} (a + b\operatorname{sech}^{-1}(cx))}{f^3(m+3)} +$$

$$\frac{3de^2 (fx)^{m+5} (a + b\operatorname{sech}^{-1}(cx))}{f^5(m+5)} + \frac{e^3 (fx)^{m+7} (a + b\operatorname{sech}^{-1}(cx))}{f^7(m+7)}$$

$$\downarrow 2340$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\int -\frac{(fx)^m\left(\frac{e^2(3d(m^2+13m+42)c^2+e(m+5)^2)x^4}{(m+5)(m+7)}+\frac{3c^2d^2e(m+6)x^2}{m+3}+\frac{c^2d^3(m+6)}{m+1}\right)}{\sqrt{1-c^2x^2}}}{c^2(m+6)}dx - \frac{e^3\sqrt{1-c^2x^2}(fx)^{m+1}}{c^2f^5(m+6)(m+7)}\right. \\ \left.+\frac{d^3(fx)^{m+1}(a+b\operatorname{sech}^{-1}(cx))}{f(m+1)}+\frac{3d^2e(fx)^{m+3}(a+b\operatorname{sech}^{-1}(cx))}{f^3(m+3)}+\frac{3de^2(fx)^{m+5}(a+b\operatorname{sech}^{-1}(cx))}{f^5(m+5)}+\frac{e^3(fx)^{m+7}(a+b\operatorname{sech}^{-1}(cx))}{f^7(m+7)}\right)$$

25

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\int \frac{(fx)^m\left(\frac{e^2(3d(m^2+13m+42)c^2+e(m+5)^2)x^4}{(m+5)(m+7)}+\frac{3c^2d^2e(m+6)x^2}{m+3}+\frac{c^2d^3(m+6)}{m+1}\right)}{\sqrt{1-c^2x^2}}}{c^2(m+6)}dx - \frac{e^3\sqrt{1-c^2x^2}(fx)^{m+5}}{c^2f^5(m+6)(m+7)}\right) \\ \left.+\frac{d^3(fx)^{m+1}(a+b\operatorname{sech}^{-1}(cx))}{f(m+1)}+\frac{3d^2e(fx)^{m+3}(a+b\operatorname{sech}^{-1}(cx))}{f^3(m+3)}+\frac{3de^2(fx)^{m+5}(a+b\operatorname{sech}^{-1}(cx))}{f^5(m+5)}+\frac{e^3(fx)^{m+7}(a+b\operatorname{sech}^{-1}(cx))}{f^7(m+7)}\right)$$

1590

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\left(\int -\frac{(fx)^m\left(\frac{d^3(m+4)(m+6)c^4}{m+1}+\frac{e(3d^2(m^4+22m^3+179m^2+638m+840)c^4+3de(m+3)^2(m^2+13m+42)c^2+e^2(m^2+8m+15)^2)}{(m+3)(m+5)(m+7)}\right)}{\sqrt{1-c^2x^2}}}{c^2(m+4)}dx - \frac{e^3\sqrt{1-c^2x^2}(fx)^{m+5}}{c^2f^5(m+6)(m+7)}\right) \\ \left.+\frac{d^3(fx)^{m+1}(a+b\operatorname{sech}^{-1}(cx))}{f(m+1)}+\frac{3d^2e(fx)^{m+3}(a+b\operatorname{sech}^{-1}(cx))}{f^3(m+3)}+\frac{3de^2(fx)^{m+5}(a+b\operatorname{sech}^{-1}(cx))}{f^5(m+5)}+\frac{e^3(fx)^{m+7}(a+b\operatorname{sech}^{-1}(cx))}{f^7(m+7)}\right)$$

25



$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{(fx)^m \left( \frac{d^3(m+4)(m+6)c^4}{m+1} + \frac{e \left( 3d^2(m^4+22m^3+179m^2+638m+840)c^4 + 3de(m+3)^2(m^2+13m+42)c^2 + e^2(m^2+8m+15)^2 \right) x^2}{(m+3)(m+5)(m+7)} \right)}{\frac{\sqrt{1-c^2x^2}}{c^2(m+4)}}}{c^2(m+6)} \right)$$

$$\frac{d^3(fx)^{m+1} (a + b\operatorname{sech}^{-1}(cx))}{f(m+1)} + \frac{3d^2e(fx)^{m+3} (a + b\operatorname{sech}^{-1}(cx))}{f^3(m+3)} +$$

$$\frac{3de^2(fx)^{m+5} (a + b\operatorname{sech}^{-1}(cx))}{f^5(m+5)} + \frac{e^3(fx)^{m+7} (a + b\operatorname{sech}^{-1}(cx))}{f^7(m+7)}$$

↓ 363

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{\left( \frac{c^4d^3(m+4)(m+6)}{m+1} + \frac{e^{(m+1)} \left( 3c^4d^2(m^4+22m^3+179m^2+638m+840) + 3c^2de(m+3)^2(m^2+13m+42) + e^2(m^2+8m+15)^2 \right)}{c^2(m+2)(m+3)(m+5)(m+7)} \right)}{c^2(m+6)}}{c^2(m+6)} \right)$$

$$\frac{d^3(fx)^{m+1} (a + b\operatorname{sech}^{-1}(cx))}{f(m+1)} + \frac{3d^2e(fx)^{m+3} (a + b\operatorname{sech}^{-1}(cx))}{f^3(m+3)} +$$

$$\frac{3de^2(fx)^{m+5} (a + b\operatorname{sech}^{-1}(cx))}{f^5(m+5)} + \frac{e^3(fx)^{m+7} (a + b\operatorname{sech}^{-1}(cx))}{f^7(m+7)}$$

↓ 278

$$\frac{d^3(fx)^{m+1} (a + b\operatorname{sech}^{-1}(cx))}{f(m+1)} + \frac{3d^2e(fx)^{m+3} (a + b\operatorname{sech}^{-1}(cx))}{f^3(m+3)} +$$

$$\frac{3de^2(fx)^{m+5} (a + b\operatorname{sech}^{-1}(cx))}{f^5(m+5)} + \frac{e^3(fx)^{m+7} (a + b\operatorname{sech}^{-1}(cx))}{f^7(m+7)} +$$

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{(fx)^{m+1} \left( \frac{c^4d^3(m+4)(m+6)}{m+1} + \frac{e^{(m+1)} \left( 3c^4d^2(m^4+22m^3+179m^2+638m+840) + 3c^2de(m+3)^2(m^2+13m+42) + e^2(m^2+8m+15)^2 \right)}{c^2(m+2)(m+3)(m+5)(m+7)} \right)}{f(m+1)}}{f(m+1)} \right)$$

input Int[(f\*x)^m\*(d + e\*x^2)^3\*(a + b\*ArcSech[c\*x]),x]

output

```
(d^3*(f*x)^(1 + m)*(a + b*ArcSech[c*x]))/(f*(1 + m)) + (3*d^2*e*(f*x)^(3 +
m)*(a + b*ArcSech[c*x]))/(f^3*(3 + m)) + (3*d*e^2*(f*x)^(5 + m)*(a + b*Ar
cSech[c*x]))/(f^5*(5 + m)) + (e^3*(f*x)^(7 + m)*(a + b*ArcSech[c*x]))/(f^7
*(7 + m)) + b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-((e^3*(f*x)^(5 + m)*Sqr
t[1 - c^2*x^2])/(c^2*f^5*(6 + m)*(7 + m))) + (-((e^2*(e*(5 + m)^2 + 3*c^2*
d*(42 + 13*m + m^2))*(f*x)^(3 + m)*Sqrt[1 - c^2*x^2])/(c^2*f^3*(4 + m)*(5
+ m)*(7 + m))) + (-((e*(e^2*(15 + 8*m + m^2)^2 + 3*c^2*d*e*(3 + m)^2*(42 +
13*m + m^2) + 3*c^4*d^2*(840 + 638*m + 179*m^2 + 22*m^3 + m^4))*(f*x)^(1
+ m)*Sqrt[1 - c^2*x^2])/(c^2*f*(2 + m)*(3 + m)*(5 + m)*(7 + m))) + (((c^4*
d^3*(4 + m)*(6 + m))/(1 + m) + (e*(1 + m)*(e^2*(15 + 8*m + m^2)^2 + 3*c^2*
d*e*(3 + m)^2*(42 + 13*m + m^2) + 3*c^4*d^2*(840 + 638*m + 179*m^2 + 22*m^
3 + m^4)))/(c^2*(2 + m)*(3 + m)*(5 + m)*(7 + m)))*(f*x)^(1 + m)*Hypergeome
tric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])/(f*(1 + m))/(c^2*(4 + m))/(c
^2*(6 + m))
```

### Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 278

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((
c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (
-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0
] || GtQ[a, 0])
```

rule 363

```
Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2), x
_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(b*e*(m + 2*p + 3))),
x] - Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(b*(m + 2*p + 3)) Int[(e*x)^
m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d
, 0] && NeQ[m + 2*p + 3, 0]
```

rule 1590

```
Int[((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (
c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[c^p*(f*x)^(m + 4*p - 1)*((d + e*x^2)^
(q + 1)/(e*f^(4*p - 1)*(m + 4*p + 2*q + 1))), x] + Simp[1/(e*(m + 4*p + 2*q
+ 1)) Int[(f*x)^m*(d + e*x^2)^q*ExpandToSum[e*(m + 4*p + 2*q + 1)*((a +
b*x^2 + c*x^4)^p - c^p*x^(4*p)) - d*c^p*(m + 4*p - 1)*x^(4*p - 2), x], x],
x] /; FreeQ[{a, b, c, d, e, f, m, q}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p,
0] && !IntegerQ[q] && NeQ[m + 4*p + 2*q + 1, 0]
```

rule 2340

```
Int[(Pq)*((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[
{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[f*(c*x)^(m + q - 1
)*((a + b*x^2)^(p + 1)/(b*c^(q - 1)*(m + q + 2*p + 1))), x] + Simp[1/(b*(m
+ q + 2*p + 1)) Int[(c*x)^m*(a + b*x^2)^p*ExpandToSum[b*(m + q + 2*p + 1)
*Pq - b*f*(m + q + 2*p + 1)*x^q - a*f*(m + q - 1)*x^(q - 2), x], x], x] /;
GtQ[q, 1] && NeQ[m + q + 2*p + 1, 0] /; FreeQ[{a, b, c, m, p}, x] && PolyQ
[Pq, x] && (!IGtQ[m, 0] || IGtQ[p + 1/2, -1])
```

rule 6855

```
Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.))*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(
x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Si
mp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)]
Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; Fre
eQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] &&
GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2
*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

## Maple [F]

$$\int (fx)^m (x^2e + d)^3 (a + b \operatorname{arcsech}(cx)) dx$$

input

```
int((f*x)^m*(e*x^2+d)^3*(a+b*arcsech(c*x)),x)
```

output

```
int((f*x)^m*(e*x^2+d)^3*(a+b*arcsech(c*x)),x)
```

**Fricas [F]**

$$\int (fx)^m (d + ex^2)^3 (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^3 (b \operatorname{arsech}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^3*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral((a*e^3*x^6 + 3*a*d*e^2*x^4 + 3*a*d^2*e*x^2 + a*d^3 + (b*e^3*x^6 + 3*b*d*e^2*x^4 + 3*b*d^2*e*x^2 + b*d^3)*arcsech(c*x))*(f*x)^m, x)`

**Sympy [F]**

$$\int (fx)^m (d + ex^2)^3 (a + b \operatorname{sech}^{-1}(cx)) dx = \int (fx)^m (a + b \operatorname{asech}(cx)) (d + ex^2)^3 dx$$

input `integrate((f*x)**m*(e*x**2+d)**3*(a+b*asech(c*x)),x)`

output `Integral((f*x)**m*(a + b*asech(c*x))*(d + e*x**2)**3, x)`

**Maxima [F]**

$$\int (fx)^m (d + ex^2)^3 (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^3 (b \operatorname{arsech}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^3*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output

```

a*e^3*f^m*x^7*x^m/(m + 7) + 3*a*d*e^2*f^m*x^5*x^m/(m + 5) + 3*a*d^2*e*f^m*
x^3*x^m/(m + 3) + (f*x)^(m + 1)*a*d^3/(f*(m + 1)) + ((m^3 + 9*m^2 + 23*m
+ 15)*b*e^3*f^m*x^7*x^m + 3*(m^3 + 11*m^2 + 31*m + 21)*b*d*e^2*f^m*x^5*x^m
+ 3*(m^3 + 13*m^2 + 47*m + 35)*b*d^2*e*f^m*x^3*x^m + (m^3 + 15*m^2 + 71*m
+ 105)*b*d^3*f^m*x*x^m)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1) - ((m^3 + 9
*m^2 + 23*m + 15)*b*e^3*f^m*x^7*x^m + 3*(m^3 + 11*m^2 + 31*m + 21)*b*d*e^2
*f^m*x^5*x^m + 3*(m^3 + 13*m^2 + 47*m + 35)*b*d^2*e*f^m*x^3*x^m + (m^3 + 1
5*m^2 + 71*m + 105)*b*d^3*f^m*x*x^m)*log(x))/(m^4 + 16*m^3 + 86*m^2 + 176*
m + 105) - integrate((b*c^2*e^3*f^m*(m + 7)*x^2*log(c) - (e^3*f^m*(m + 7)*
log(c) - e^3*f^m)*b)*x^6*x^m/(c^2*(m + 7)*x^2 - m - 7), x) - integrate(3*(
b*c^2*d*e^2*f^m*(m + 5)*x^2*log(c) - (d*e^2*f^m*(m + 5)*log(c) - d*e^2*f^m
)*b)*x^4*x^m/(c^2*(m + 5)*x^2 - m - 5), x) - integrate(3*(b*c^2*d^2*e*f^m*
(m + 3)*x^2*log(c) - (d^2*e*f^m*(m + 3)*log(c) - d^2*e*f^m)*b)*x^2*x^m/(c^
2*(m + 3)*x^2 - m - 3), x) - integrate((b*c^2*d^3*f^m*(m + 1)*x^2*log(c) -
(d^3*f^m*(m + 1)*log(c) - d^3*f^m)*b)*x^m/(c^2*(m + 1)*x^2 - m - 1), x) +
integrate(((m^3 + 9*m^2 + 23*m + 15)*b*c^2*e^3*f^m*x^8*x^m + 3*(m^3 + 11*
m^2 + 31*m + 21)*b*c^2*d*e^2*f^m*x^6*x^m + 3*(m^3 + 13*m^2 + 47*m + 35)*b*
c^2*d^2*e*f^m*x^4*x^m + (m^3 + 15*m^2 + 71*m + 105)*b*c^2*d^3*f^m*x^2*x^m)
/((m^4 + 16*m^3 + 86*m^2 + 176*m + 105)*c^2*x^2 - m^4 - 16*m^3 + ((m^4 + 1
6*m^3 + 86*m^2 + 176*m + 105)*c^2*x^2 - m^4 - 16*m^3 - 86*m^2 - 176*m - ...

```

**Giac [F]**

$$\int (fx)^m (d + ex^2)^3 (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^3 (b \operatorname{arsech}(cx) + a)(fx)^m dx$$

input

```
integrate((f*x)^m*(e*x^2+d)^3*(a+b*arcsech(c*x)),x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)^3*(b*arcsech(c*x) + a)*(f*x)^m, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int (fx)^m (d+ex^2)^3 (a+b\operatorname{sech}^{-1}(cx)) dx = \int (fx)^m (ex^2+d)^3 \left( a+b\operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int((f*x)^m*(d + e*x^2)^3*(a + b*acosh(1/(c*x))),x)`

output `int((f*x)^m*(d + e*x^2)^3*(a + b*acosh(1/(c*x))), x)`

**Reduce [F]**

$$\int (fx)^m (d+ex^2)^3 (a+b\operatorname{sech}^{-1}(cx)) dx = \text{Too large to display}$$

input `int((f*x)^m*(e*x^2+d)^3*(a+b*asech(c*x)),x)`

output `(f**m*(x**m*a*d**3*m**3*x + 15*x**m*a*d**3*m**2*x + 71*x**m*a*d**3*m*x + 105*x**m*a*d**3*x + 3*x**m*a*d**2*e*m**3*x**3 + 39*x**m*a*d**2*e*m**2*x**3 + 141*x**m*a*d**2*e*m*x**3 + 105*x**m*a*d**2*e*x**3 + 3*x**m*a*d*e**2*m**3*x**5 + 33*x**m*a*d*e**2*m**2*x**5 + 93*x**m*a*d*e**2*m*x**5 + 63*x**m*a*d*e**2*x**5 + x**m*a*e**3*m**3*x**7 + 9*x**m*a*e**3*m**2*x**7 + 23*x**m*a*e**3*m*x**7 + 15*x**m*a*e**3*x**7 + int(x**m*asech(c*x)*x**6,x)*b*e**3*m**4 + 16*int(x**m*asech(c*x)*x**6,x)*b*e**3*m**3 + 86*int(x**m*asech(c*x)*x**6,x)*b*e**3*m**2 + 176*int(x**m*asech(c*x)*x**6,x)*b*e**3*m + 105*int(x**m*asech(c*x)*x**6,x)*b*e**3 + 3*int(x**m*asech(c*x)*x**4,x)*b*d*e**2*m**4 + 48*int(x**m*asech(c*x)*x**4,x)*b*d*e**2*m**3 + 258*int(x**m*asech(c*x)*x**4,x)*b*d*e**2*m**2 + 528*int(x**m*asech(c*x)*x**4,x)*b*d*e**2*m + 315*int(x**m*asech(c*x)*x**4,x)*b*d*e**2 + 3*int(x**m*asech(c*x)*x**2,x)*b*d**2*e*m**4 + 48*int(x**m*asech(c*x)*x**2,x)*b*d**2*e*m**3 + 258*int(x**m*asech(c*x)*x**2,x)*b*d**2*e*m**2 + 528*int(x**m*asech(c*x)*x**2,x)*b*d**2*e*m + 315*int(x**m*asech(c*x)*x**2,x)*b*d**2*e + int(x**m*asech(c*x),x)*b*d**3*m**4 + 16*int(x**m*asech(c*x),x)*b*d**3*m**3 + 86*int(x**m*asech(c*x),x)*b*d**3*m**2 + 176*int(x**m*asech(c*x),x)*b*d**3*m + 105*int(x**m*asech(c*x),x)*b*d**3))/(m**4 + 16*m**3 + 86*m**2 + 176*m + 105)`

### 3.177 $\int (fx)^m (d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	1398
Mathematica [A] (verified)	1399
Rubi [A] (verified)	1399
Maple [F]	1402
Fricas [F]	1403
Sympy [F]	1403
Maxima [F]	1403
Giac [F]	1404
Mupad [F(-1)]	1404
Reduce [F]	1405

#### Optimal result

Integrand size = 23, antiderivative size = 372

$$\int (fx)^m (d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= -\frac{be(e(3+m)^2 + 2c^2d(20 + 9m + m^2))(fx)^{1+m} \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{c^4 f(2+m)(3+m)(4+m)(5+m)}$$

$$- \frac{be^2(fx)^{3+m} \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{c^2 f^3(4+m)(5+m)} + \frac{d^2(fx)^{1+m} (a + b\operatorname{sech}^{-1}(cx))}{f(1+m)}$$

$$+ \frac{2de(fx)^{3+m} (a + b\operatorname{sech}^{-1}(cx))}{f^3(3+m)} + \frac{e^2(fx)^{5+m} (a + b\operatorname{sech}^{-1}(cx))}{f^5(5+m)}$$

$$+ \frac{b(c^4 d^2(2+m)(3+m)(4+m)(5+m) + e(1+m)^2 (e(3+m)^2 + 2c^2d(20 + 9m + m^2))) (fx)^{1+m} \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{c^4 f(1+m)^2(2+m)(3+m)(4+m)(5+m)}$$

output

```
-b*e*(e*(3+m)^2+2*c^2*d*(m^2+9*m+20))*(f*x)^(1+m)*(1/(c*x+1))^(1/2)*(c*x+1)
)^(1/2)*(-c^2*x^2+1)^(1/2)/c^4/f/(2+m)/(3+m)/(4+m)/(5+m)-b*e^2*(f*x)^(3+m)
*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^2/f^3/(4+m)/(5+m)+d^
2*(f*x)^(1+m)*(a+b*arcsech(c*x))/f/(1+m)+2*d*e*(f*x)^(3+m)*(a+b*arcsech(c*
x))/f^3/(3+m)+e^2*(f*x)^(5+m)*(a+b*arcsech(c*x))/f^5/(5+m)+b*(c^4*d^2*(2+m)
)*(3+m)*(4+m)*(5+m)+e*(1+m)^2*(e*(3+m)^2+2*c^2*d*(m^2+9*m+20))*(f*x)^(1+m)
)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*hypergeom([1/2, 1/2+1/2*m], [3/2+1/2*m], c
^2*x^2)/c^4/f/(1+m)^2/(2+m)/(3+m)/(4+m)/(5+m)
```

**Mathematica [A] (verified)**

Time = 0.49 (sec) , antiderivative size = 322, normalized size of antiderivative = 0.87

$$\int (fx)^m (d + ex^2)^2 (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= x(fx)^m \left( \frac{ad^2}{1+m} + \frac{2adex^2}{3+m} + \frac{ae^2x^4}{5+m} + \frac{bd^2\operatorname{sech}^{-1}(cx)}{1+m} + \frac{2bdex^2\operatorname{sech}^{-1}(cx)}{3+m} \right.$$

$$+ \frac{be^2x^4\operatorname{sech}^{-1}(cx)}{5+m} - \frac{bd^2\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2x^2\right)}{(1+m)^2(-1+cx)}$$

$$- \frac{2bdex^2\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3+m}{2}, \frac{5+m}{2}, c^2x^2\right)}{(3+m)^2(-1+cx)}$$

$$\left. - \frac{be^2x^4\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5+m}{2}, \frac{7+m}{2}, c^2x^2\right)}{(5+m)^2(-1+cx)} \right)$$

input `Integrate[(f*x)^m*(d + e*x^2)^2*(a + b*ArcSech[c*x]),x]`

output `x*(f*x)^m*((a*d^2)/(1+m) + (2*a*d*e*x^2)/(3+m) + (a*e^2*x^4)/(5+m) + (b*d^2*ArcSech[c*x])/(1+m) + (2*b*d*e*x^2*ArcSech[c*x])/(3+m) + (b*e^2*x^4*ArcSech[c*x])/(5+m) - (b*d^2*Sqrt[(1-c*x)/(1+c*x)]*Sqrt[1-c^2*x^2]*Hypergeometric2F1[1/2, (1+m)/2, (3+m)/2, c^2*x^2])/((1+m)^2*(-1+c*x)) - (2*b*d*e*x^2*Sqrt[(1-c*x)/(1+c*x)]*Sqrt[1-c^2*x^2]*Hypergeometric2F1[1/2, (3+m)/2, (5+m)/2, c^2*x^2])/((3+m)^2*(-1+c*x)) - (b*e^2*x^4*Sqrt[(1-c*x)/(1+c*x)]*Sqrt[1-c^2*x^2]*Hypergeometric2F1[1/2, (5+m)/2, (7+m)/2, c^2*x^2])/((5+m)^2*(-1+c*x))`

**Rubi [A] (verified)**

Time = 0.67 (sec) , antiderivative size = 328, normalized size of antiderivative = 0.88, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$ , Rules used = {6855, 27, 1590, 25, 363, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.



$$\int (d + ex^2)^2 (fx)^m (a + b \operatorname{sech}^{-1}(cx)) dx$$

↓ 6855

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(fx)^m (e^2(m+1)(m+3)x^4 + 2de(m+1)(m+5)x^2 + d^2(m+3)(m+5))}{(m^3 + 9m^2 + 23m + 15)\sqrt{1-c^2x^2}} dx +$$

$$\frac{d^2(fx)^{m+1} (a + b \operatorname{sech}^{-1}(cx))}{f(m+1)} + \frac{2de(fx)^{m+3} (a + b \operatorname{sech}^{-1}(cx))}{f^3(m+3)} + \frac{e^2(fx)^{m+5} (a + b \operatorname{sech}^{-1}(cx))}{f^5(m+5)}$$

↓ 27

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(fx)^m (e^2(m+1)(m+3)x^4 + 2de(m+1)(m+5)x^2 + d^2(m+3)(m+5))}{\sqrt{1-c^2x^2}} dx +$$

$$\frac{d^2(fx)^{m+1} (a + b \operatorname{sech}^{-1}(cx))}{f(m+1)} + \frac{m^3 + 9m^2 + 23m + 15}{f^3(m+3)} \frac{2de(fx)^{m+3} (a + b \operatorname{sech}^{-1}(cx))}{f^3(m+3)} + \frac{e^2(fx)^{m+5} (a + b \operatorname{sech}^{-1}(cx))}{f^5(m+5)}$$

↓ 1590

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( - \frac{\int \frac{(fx)^m (c^2(m+3)(m+4)(m+5)d^2 + e(m+1)(2d(m^2+9m+20)c^2 + e(m+3)^2)x^2)}{\sqrt{1-c^2x^2}} dx}{c^2(m+4)} - \frac{e^2(m+1)(m+3)\sqrt{1-c^2x^2}(fx)^{m+3}}{c^2 f^3(m+4)} \right)$$

$$\frac{d^2(fx)^{m+1} (a + b \operatorname{sech}^{-1}(cx))}{f(m+1)} + \frac{m^3 + 9m^2 + 23m + 15}{f^3(m+3)} \frac{2de(fx)^{m+3} (a + b \operatorname{sech}^{-1}(cx))}{f^3(m+3)} + \frac{e^2(fx)^{m+5} (a + b \operatorname{sech}^{-1}(cx))}{f^5(m+5)}$$

↓ 25

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\int \frac{(fx)^m (c^2(m+3)(m+4)(m+5)d^2 + e(m+1)(2d(m^2+9m+20)c^2 + e(m+3)^2)x^2)}{\sqrt{1-c^2x^2}} dx}{c^2(m+4)} - \frac{e^2(m+1)(m+3)\sqrt{1-c^2x^2}(fx)^{m+3}}{c^2 f^3(m+4)} \right) +$$

$$\frac{d^2(fx)^{m+1} (a + b \operatorname{sech}^{-1}(cx))}{f(m+1)} + \frac{m^3 + 9m^2 + 23m + 15}{f^3(m+3)} \frac{2de(fx)^{m+3} (a + b \operatorname{sech}^{-1}(cx))}{f^3(m+3)} + \frac{e^2(fx)^{m+5} (a + b \operatorname{sech}^{-1}(cx))}{f^5(m+5)}$$

↓ 363

$$b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{\left( \frac{c^4 d^2(m+3)(m+4)(m+5) + \frac{e(m+1)^2(2c^2 d(m^2+9m+20) + e(m+3)^2)}{m+2}}{c^2} \right) \int \frac{(fx)^m}{\sqrt{1-c^2x^2}} dx}{c^2(m+4)} - \frac{e(m+1)\sqrt{1-c^2x^2}(fx)^{m+1}(2c^2 d(m^2+9m+20) + e(m+3)^2)}{c^2 f(m+2)} \right) +$$

$$\frac{d^2(fx)^{m+1} (a + b \operatorname{sech}^{-1}(cx))}{f(m+1)} + \frac{2de(fx)^{m+3} (a + b \operatorname{sech}^{-1}(cx))}{f^3(m+3)} + \frac{m^3 + 9m^2 + 23m + 15}{f^5(m+5)} \frac{e^2(fx)^{m+5} (a + b \operatorname{sech}^{-1}(cx))}{f^5(m+5)}$$

$$\begin{aligned}
 & \downarrow 278 \\
 & \frac{d^2(fx)^{m+1} (a + b\operatorname{sech}^{-1}(cx))}{f(m+1)} + \frac{2de(fx)^{m+3} (a + b\operatorname{sech}^{-1}(cx))}{f^3(m+3)} + \\
 & \frac{e^2(fx)^{m+5} (a + b\operatorname{sech}^{-1}(cx))}{f^5(m+5)} + \\
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{(fx)^{m+1} \left( c^4 d^2 (m+3)(m+4)(m+5) + \frac{e(m+1)^2 (2c^2 d(m^2+9m+20) + e(m+3)^2)}{m+2} \right)}{c^2 f(m+1)} \operatorname{Hypergeometric2F1} \left( \frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, c^2 x^2 \right) - \frac{e(m+1)}{c^2(m+4)} \right) \\
 & \hline
 & m^3 + 9m^2 + 23m + 15
 \end{aligned}$$

```
input Int[(f*x)^m*(d + e*x^2)^2*(a + b*ArcSech[c*x]),x]
```

```
output (d^2*(f*x)^(1 + m)*(a + b*ArcSech[c*x]))/(f*(1 + m)) + (2*d*e*(f*x)^(3 + m)*(a + b*ArcSech[c*x]))/(f^3*(3 + m)) + (e^2*(f*x)^(5 + m)*(a + b*ArcSech[c*x]))/(f^5*(5 + m)) + (b*Sqrt[(1 + c*x)^(-1)]*Sqrt[1 + c*x]*(-(e^2*(1 + m)*(3 + m)*(f*x)^(3 + m)*Sqrt[1 - c^2*x^2])/(c^2*f^3*(4 + m)))) + (-(e*(1 + m)*(e*(3 + m)^2 + 2*c^2*d*(20 + 9*m + m^2))*(f*x)^(1 + m)*Sqrt[1 - c^2*x^2])/(c^2*f*(2 + m))) + ((c^4*d^2*(3 + m)*(4 + m)*(5 + m) + (e*(1 + m)^2*(e*(3 + m)^2 + 2*c^2*d*(20 + 9*m + m^2))))/(2 + m))*(f*x)^(1 + m)*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2]/(c^2*f*(1 + m))/(c^2*(4 + m)))/(15 + 23*m + 9*m^2 + m^3)
```

**Defintions of rubi rules used**

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 278 Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])
```

rule 363

```
Int[((e._)*(x._))^(m._)*((a._) + (b._)*(x._)^2)^(p._)*((c._) + (d._)*(x._)^2), x
_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(b*e*(m + 2*p + 3))),
x] - Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(b*(m + 2*p + 3)) Int[(e*x)^
m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d
, 0] && NeQ[m + 2*p + 3, 0]
```

rule 1590

```
Int(((f._)*(x._))^(m._)*((d._) + (e._)*(x._)^2)^(q._)*((a._) + (b._)*(x._)^2 + (
c._)*(x._)^4)^(p._), x_Symbol] := Simp[c^p*(f*x)^(m + 4*p - 1)*((d + e*x^2)^
(q + 1)/(e*f^(4*p - 1)*(m + 4*p + 2*q + 1))), x] + Simp[1/(e*(m + 4*p + 2*q
+ 1)) Int[(f*x)^m*(d + e*x^2)^q*ExpandToSum[e*(m + 4*p + 2*q + 1)*((a +
b*x^2 + c*x^4)^p - c^p*x^(4*p)) - d*c^p*(m + 4*p - 1)*x^(4*p - 2), x], x],
x] /; FreeQ[{a, b, c, d, e, f, m, q}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p,
0] && !IntegerQ[q] && NeQ[m + 4*p + 2*q + 1, 0]
```

rule 6855

```
Int[((a._) + ArcSech[(c._)*(x._)]*(b._))*((f._)*(x._))^(m._)*((d._) + (e._)*(
x._)^2)^(p._), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Si
mp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)]
Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x]] /; Fre
eQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] &&
GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2
*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))
```

## Maple [F]

$$\int (fx)^m (x^2e + d)^2 (a + b \operatorname{arcsech}(cx)) dx$$

input

```
int((f*x)^m*(e*x^2+d)^2*(a+b*arcsech(c*x)),x)
```

output

```
int((f*x)^m*(e*x^2+d)^2*(a+b*arcsech(c*x)),x)
```

**Fricas [F]**

$$\int (fx)^m (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^2 (b \operatorname{arsech}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^2*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral((a*e^2*x^4 + 2*a*d*e*x^2 + a*d^2 + (b*e^2*x^4 + 2*b*d*e*x^2 + b*d^2)*arcsech(c*x))*(f*x)^m, x)`

**Sympy [F]**

$$\int (fx)^m (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \int (fx)^m (a + b \operatorname{asech}(cx)) (d + ex^2)^2 dx$$

input `integrate((f*x)**m*(e*x**2+d)**2*(a+b*asech(c*x)),x)`

output `Integral((f*x)**m*(a + b*asech(c*x))*(d + e*x**2)**2, x)`

**Maxima [F]**

$$\int (fx)^m (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^2 (b \operatorname{arsech}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^2*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output

```

a*e^2*f^m*x^5*x^m/(m + 5) + 2*a*d*e*f^m*x^3*x^m/(m + 3) + (f*x)^(m + 1)*a*
d^2/(f*(m + 1)) + (((m^2 + 4*m + 3)*b*e^2*f^m*x^5*x^m + 2*(m^2 + 6*m + 5)*
b*d*e*f^m*x^3*x^m + (m^2 + 8*m + 15)*b*d^2*f^m*x*x^m)*log(sqrt(c*x + 1)*sq
rt(-c*x + 1) + 1) - ((m^2 + 4*m + 3)*b*e^2*f^m*x^5*x^m + 2*(m^2 + 6*m + 5)
*b*d*e*f^m*x^3*x^m + (m^2 + 8*m + 15)*b*d^2*f^m*x*x^m)*log(x))/(m^3 + 9*m^
2 + 23*m + 15) - integrate((b*c^2*e^2*f^m*(m + 5)*x^2*log(c) - (e^2*f^m*(m
+ 5)*log(c) - e^2*f^m)*b)*x^4*x^m/(c^2*(m + 5)*x^2 - m - 5), x) - integra
te(2*(b*c^2*d*e*f^m*(m + 3)*x^2*log(c) - (d*e*f^m*(m + 3)*log(c) - d*e*f^m
)*b)*x^2*x^m/(c^2*(m + 3)*x^2 - m - 3), x) - integrate((b*c^2*d^2*f^m*(m +
1)*x^2*log(c) - (d^2*f^m*(m + 1)*log(c) - d^2*f^m)*b)*x^m/(c^2*(m + 1)*x^
2 - m - 1), x) + integrate(((m^2 + 4*m + 3)*b*c^2*e^2*f^m*x^6*x^m + 2*(m^2
+ 6*m + 5)*b*c^2*d*e*f^m*x^4*x^m + (m^2 + 8*m + 15)*b*c^2*d^2*f^m*x^2*x^m
)/((m^3 + 9*m^2 + 23*m + 15)*c^2*x^2 - m^3 + ((m^3 + 9*m^2 + 23*m + 15)*c^
2*x^2 - m^3 - 9*m^2 - 23*m - 15)*sqrt(c*x + 1)*sqrt(-c*x + 1) - 9*m^2 - 23
*m - 15), x)

```

**Giac [F]**

$$\int (fx)^m (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^2 (b \operatorname{arsech}(cx) + a)(fx)^m dx$$

input

```
integrate((f*x)^m*(e*x^2+d)^2*(a+b*arcsech(c*x)),x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)^2*(b*arcsech(c*x) + a)*(f*x)^m, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int (fx)^m (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx = \int (fx)^m (ex^2 + d)^2 \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input

```
int((f*x)^m*(d + e*x^2)^2*(a + b*acosh(1/(c*x))),x)
```

output

```
int((f*x)^m*(d + e*x^2)^2*(a + b*acosh(1/(c*x))), x)
```

**Reduce [F]**

$$\int (fx)^m (d + ex^2)^2 (a + b \operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{f^m (x^m a d^2 m^2 x + 8x^m a d^2 m x + 15x^m a d^2 x + 2x^m a d e m^2 x^3 + 12x^m a d e m x^3 + 10x^m a d e x^3 + x^m a e^2 m^2 x^5 + 8x^m a e^2 m x^5 + 3x^m a e^2 x^5 + \int (x^m a \operatorname{sech}^{-1}(cx)) x^4 dx) b e^2 m^3 + 9 \int (x^m a \operatorname{sech}^{-1}(cx)) x^4 dx) b e^2 m^2 + 23 \int (x^m a \operatorname{sech}^{-1}(cx)) x^4 dx) b e^2 m + 15 \int (x^m a \operatorname{sech}^{-1}(cx)) x^4 dx) b e^2 + 2 \int (x^m a \operatorname{sech}^{-1}(cx)) x^2 dx) b d e m^3 + 18 \int (x^m a \operatorname{sech}^{-1}(cx)) x^2 dx) b d e m^2 + 46 \int (x^m a \operatorname{sech}^{-1}(cx)) x^2 dx) b d e m + 30 \int (x^m a \operatorname{sech}^{-1}(cx)) x^2 dx) b d e + \int (x^m a \operatorname{sech}^{-1}(cx)) dx) b d^2 m^3 + 9 \int (x^m a \operatorname{sech}^{-1}(cx)) dx) b d^2 m^2 + 23 \int (x^m a \operatorname{sech}^{-1}(cx)) dx) b d^2 m + 15 \int (x^m a \operatorname{sech}^{-1}(cx)) dx) b d^2)}{(m^3 + 9m^2 + 23m + 15)}$$

input `int((f*x)^m*(e*x^2+d)^2*(a+b*asech(c*x)),x)`

output

```
(f**m*(x**m*a*d**2*m**2*x + 8*x**m*a*d**2*m*x + 15*x**m*a*d**2*x + 2*x**m*a*d*e*m**2*x**3 + 12*x**m*a*d*e*m*x**3 + 10*x**m*a*d*e*x**3 + x**m*a*e**2*m**2*x**5 + 4*x**m*a*e**2*m*x**5 + 3*x**m*a*e**2*x**5 + int(x**m*asech(c*x)*x**4,x)*b*e**2*m**3 + 9*int(x**m*asech(c*x)*x**4,x)*b*e**2*m**2 + 23*int(x**m*asech(c*x)*x**4,x)*b*e**2*m + 15*int(x**m*asech(c*x)*x**4,x)*b*e**2 + 2*int(x**m*asech(c*x)*x**2,x)*b*d*e*m**3 + 18*int(x**m*asech(c*x)*x**2,x)*b*d*e*m**2 + 46*int(x**m*asech(c*x)*x**2,x)*b*d*e*m + 30*int(x**m*asech(c*x)*x**2,x)*b*d*e + int(x**m*asech(c*x),x)*b*d**2*m**3 + 9*int(x**m*asech(c*x),x)*b*d**2*m**2 + 23*int(x**m*asech(c*x),x)*b*d**2*m + 15*int(x**m*asech(c*x),x)*b*d**2))/(m**3 + 9*m**2 + 23*m + 15)
```

### 3.178 $\int (fx)^m (d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	1406
Mathematica [A] (verified)	1407
Rubi [A] (verified)	1407
Maple [F]	1409
Fricas [F]	1410
Sympy [F]	1410
Maxima [F]	1410
Giac [F]	1411
Mupad [F(-1)]	1411
Reduce [F]	1412

#### Optimal result

Integrand size = 21, antiderivative size = 206

$$\int (fx)^m (d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx = -\frac{be(fx)^{1+m} \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \sqrt{1-c^2x^2}}{c^2 f(2+m)(3+m)} + \frac{d(fx)^{1+m} (a + b\operatorname{sech}^{-1}(cx))}{f(1+m)} + \frac{e(fx)^{3+m} (a + b\operatorname{sech}^{-1}(cx))}{f^3(3+m)} + \frac{b(e(1+m)^2 + c^2d(2+m)(3+m)) (fx)^{1+m} \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2x^2\right)}{c^2 f(1+m)^2(2+m)(3+m)}$$

output

```
-b*e*(f*x)^(1+m)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/c^2/f/(2+m)/(3+m)+d*(f*x)^(1+m)*(a+b*arcsech(c*x))/f/(1+m)+e*(f*x)^(3+m)*(a+b*arcsech(c*x))/f^3/(3+m)+b*(e*(1+m)^2+c^2*d*(2+m)*(3+m))*(f*x)^(1+m)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)*hypergeom([1/2, 1/2+1/2*m], [3/2+1/2*m], c^2*x^2)/c^2/f/(1+m)^2/(2+m)/(3+m)
```

**Mathematica [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.92

$$\int (fx)^m (d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= x(fx)^m \left( -\frac{bd\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2x^2\right)}{(1+m)^2(-1+cx)} \right. \\ \left. + \frac{(3+m)(d(3+m)+e(1+m)x^2)(a+b\operatorname{sech}^{-1}(cx))}{1+m} - \frac{be x^2\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^2x^2}\operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3+m}{2}, \frac{5+m}{2}, c^2x^2\right)}{-1+cx} \right) \frac{1}{(3+m)^2}$$

input `Integrate[(f*x)^m*(d + e*x^2)*(a + b*ArcSech[c*x]),x]`

output `x*(f*x)^m*(-((b*d*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])/((1 + m)^2*(-1 + c*x))) + (((3 + m)*(d*(3 + m) + e*(1 + m)*x^2)*(a + b*ArcSech[c*x]))/(1 + m) - (b*e*x^2*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (3 + m)/2, (5 + m)/2, c^2*x^2])/(-1 + c*x))/(3 + m)^2)`

**Rubi [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 184, normalized size of antiderivative = 0.89, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$ , Rules used = {6855, 27, 363, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2) (fx)^m (a + b\operatorname{sech}^{-1}(cx)) dx$$

↓ 6855



$$\begin{aligned}
 & b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(fx)^m (e(m+1)x^2 + d(m+3))}{(m^2 + 4m + 3)\sqrt{1-c^2x^2}} dx + \frac{d(fx)^{m+1} (a + b\operatorname{sech}^{-1}(cx))}{f(m+1)} + \\
 & \quad \frac{e(fx)^{m+3} (a + b\operatorname{sech}^{-1}(cx))}{f^3(m+3)} \\
 & \quad \downarrow 27 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \int \frac{(fx)^m (e(m+1)x^2 + d(m+3))}{\sqrt{1-c^2x^2}} dx}{m^2 + 4m + 3} + \frac{d(fx)^{m+1} (a + b\operatorname{sech}^{-1}(cx))}{f(m+1)} + \\
 & \quad \frac{e(fx)^{m+3} (a + b\operatorname{sech}^{-1}(cx))}{f^3(m+3)} \\
 & \quad \downarrow 363 \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \left( \frac{e(m+1)^2}{c^2(m+2)} + d(m+3) \right) \int \frac{(fx)^m}{\sqrt{1-c^2x^2}} dx - \frac{e(m+1)\sqrt{1-c^2x^2}(fx)^{m+1}}{c^2 f(m+2)} \right)}{m^2 + 4m + 3} + \\
 & \quad \frac{d(fx)^{m+1} (a + b\operatorname{sech}^{-1}(cx))}{f(m+1)} + \frac{e(fx)^{m+3} (a + b\operatorname{sech}^{-1}(cx))}{f^3(m+3)} \\
 & \quad \downarrow 278 \\
 & \frac{d(fx)^{m+1} (a + b\operatorname{sech}^{-1}(cx))}{f(m+1)} + \frac{e(fx)^{m+3} (a + b\operatorname{sech}^{-1}(cx))}{f^3(m+3)} + \\
 & \frac{b\sqrt{\frac{1}{cx+1}}\sqrt{cx+1} \left( \frac{(fx)^{m+1} \left( \frac{e(m+1)^2}{c^2(m+2)} + d(m+3) \right) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, c^2x^2\right)}{f(m+1)} - \frac{e(m+1)\sqrt{1-c^2x^2}(fx)^{m+1}}{c^2 f(m+2)} \right)}{m^2 + 4m + 3}
 \end{aligned}$$

input `Int[(f*x)^m*(d + e*x^2)*(a + b*ArcSech[c*x]),x]`

output `(d*(f*x)^(1 + m)*(a + b*ArcSech[c*x]))/(f*(1 + m)) + (e*(f*x)^(3 + m)*(a + b*ArcSech[c*x]))/(f^3*(3 + m)) + (b*sqrt[(1 + c*x)^(-1)]*sqrt[1 + c*x]*(-((e*(1 + m)*(f*x)^(1 + m)*sqrt[1 - c^2*x^2])/(c^2*f*(2 + m))) + (((e*(1 + m)^2)/(c^2*(2 + m)) + d*(3 + m))*(f*x)^(1 + m)*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])/(f*(1 + m))))/(3 + 4*m + m^2)`

## Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 278 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1))]*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 363 `Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(b*e*(m + 2*p + 3))), x] - Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(b*(m + 2*p + 3)) Int[(e*x)^(m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + 2*p + 3, 0]`

rule 6855 `Int[((a_) + ArcSech[(c_)*(x_)])*(b_))*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcSech[c*x]) u, x] + Simp[b*Sqrt[1 + c*x]*Sqrt[1/(1 + c*x)] Int[SimplifyIntegrand[u/(x*Sqrt[1 - c*x]*Sqrt[1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && ((IGtQ[p, 0] && !(ILtQ[(m - 1)/2, 0] && GtQ[m + 2*p + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(ILtQ[p, 0] && GtQ[m + 2*p + 3, 0])) || (ILtQ[(m + 2*p + 1)/2, 0] && !ILtQ[(m - 1)/2, 0]))`

## Maple [F]

$$\int (fx)^m (x^2e + d) (a + b \operatorname{arcsech}(cx)) dx$$

input `int((f*x)^m*(e*x^2+d)*(a+b*arcsech(c*x)),x)`

output `int((f*x)^m*(e*x^2+d)*(a+b*arcsech(c*x)),x)`

**Fricas [F]**

$$\int (fx)^m (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)(b \operatorname{arsech}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral((a*e*x^2 + a*d + (b*e*x^2 + b*d)*arcsech(c*x))*(f*x)^m, x)`

**Sympy [F]**

$$\int (fx)^m (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx = \int (fx)^m (a + b \operatorname{asech}(cx)) (d + ex^2) dx$$

input `integrate((f*x)**m*(e*x**2+d)*(a+b*asech(c*x)),x)`

output `Integral((f*x)**m*(a + b*asech(c*x))*(d + e*x**2), x)`

**Maxima [F]**

$$\int (fx)^m (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)(b \operatorname{arsech}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output

```
a*e*f^m*x^3*x^m/(m + 3) + (f*x)^(m + 1)*a*d/(f*(m + 1)) + ((b*e*f^m*(m + 1)
)*x^3*x^m + b*d*f^m*(m + 3)*x*x^m)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1) -
(b*e*f^m*(m + 1)*x^3*x^m + b*d*f^m*(m + 3)*x*x^m)*log(x))/(m^2 + 4*m + 3)
- integrate((b*c^2*e*f^m*(m + 3)*x^2*log(c) - (e*f^m*(m + 3)*log(c) - e*f
^m)*b)*x^2*x^m/(c^2*(m + 3)*x^2 - m - 3), x) - integrate((b*c^2*d*f^m*(m +
1)*x^2*log(c) - (d*f^m*(m + 1)*log(c) - d*f^m)*b)*x^m/(c^2*(m + 1)*x^2 -
m - 1), x) + integrate((b*c^2*e*f^m*(m + 1)*x^4*x^m + b*c^2*d*f^m*(m + 3)*
x^2*x^m)/((m^2 + 4*m + 3)*c^2*x^2 + ((m^2 + 4*m + 3)*c^2*x^2 - m^2 - 4*m -
3)*sqrt(c*x + 1)*sqrt(-c*x + 1) - m^2 - 4*m - 3), x)
```

**Giac [F]**

$$\int (fx)^m (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)(b \operatorname{ar} \operatorname{sech}(cx) + a)(fx)^m dx$$

input

```
integrate((f*x)^m*(e*x^2+d)*(a+b*arcsech(c*x)),x, algorithm="giac")
```

output

```
integrate((e*x^2 + d)*(b*arcsech(c*x) + a)*(f*x)^m, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int (fx)^m (d + ex^2) (a + b \operatorname{sech}^{-1}(cx)) dx = \int (fx)^m (ex^2 + d) \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input

```
int((f*x)^m*(d + e*x^2)*(a + b*acosh(1/(c*x))),x)
```

output

```
int((f*x)^m*(d + e*x^2)*(a + b*acosh(1/(c*x))), x)
```

**Reduce [F]**

$$\int (fx)^m (d + ex^2) (a + b\operatorname{sech}^{-1}(cx)) dx$$

$$= \frac{f^m (x^m a d m x + 3x^m a d x + x^m a e m x^3 + x^m a e x^3 + (\int x^m a \operatorname{sech}(cx) x^2 dx) b e m^2 + 4(\int x^m a \operatorname{sech}(cx) x^2 dx))}{m^2 + 4m + 3}$$

input `int((f*x)^m*(e*x^2+d)*(a+b*asech(c*x)),x)`

output `(f**m*(x**m*a*d*m*x + 3*x**m*a*d*x + x**m*a*e*m*x**3 + x**m*a*e*x**3 + int(x**m*asech(c*x)*x**2,x)*b*e*m**2 + 4*int(x**m*asech(c*x)*x**2,x)*b*e*m + 3*int(x**m*asech(c*x)*x**2,x)*b*e + int(x**m*asech(c*x),x)*b*d*m**2 + 4*int(x**m*asech(c*x),x)*b*d*m + 3*int(x**m*asech(c*x),x)*b*d))/(m**2 + 4*m + 3)`

**3.179** 
$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx$$

Optimal result	1413
Mathematica [N/A]	1413
Rubi [N/A]	1414
Maple [N/A]	1414
Fricas [N/A]	1415
Sympy [N/A]	1415
Maxima [N/A]	1415
Giac [N/A]	1416
Mupad [N/A]	1416
Reduce [N/A]	1417

**Optimal result**

Integrand size = 23, antiderivative size = 23

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \operatorname{Int} \left( \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{d + ex^2}, x \right)$$

output

```
Defer(Int)((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d),x)
```

**Mathematica [N/A]**

Not integrable

Time = 2.37 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx$$

input

```
Integrate[((f*x)^m*(a + b*ArcSech[c*x]))/(d + e*x^2),x]
```

output

```
Integrate[((f*x)^m*(a + b*ArcSech[c*x]))/(d + e*x^2), x]
```

**Rubi [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx$$

↓ 6865

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx$$

input `Int[((f*x)^m*(a + b*ArcSech[c*x]))/(d + e*x^2),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 1.08 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{arcsech}(cx))}{x^2 e + d} dx$$

input `int((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d),x)`

output `int((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d),x)`

**Fricas [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)(fx)^m}{ex^2 + d} dx$$

input `integrate((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d),x, algorithm="fricas")`

output `integral((b*arcsech(c*x) + a)*(f*x)^m/(e*x^2 + d), x)`

**Sympy [N/A]**

Not integrable

Time = 10.46 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \int \frac{(fx)^m (a + b \operatorname{asech}(cx))}{d + ex^2} dx$$

input `integrate((f*x)**m*(a+b*asech(c*x))/(e*x**2+d),x)`

output `Integral((f*x)**m*(a + b*asech(c*x))/(d + e*x**2), x)`

**Maxima [N/A]**

Not integrable

Time = 0.14 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)(fx)^m}{ex^2 + d} dx$$

input `integrate((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d),x, algorithm="maxima")`



output `integrate((b*arcsech(c*x) + a)*(f*x)^m/(e*x^2 + d), x)`

### Giac [N/A]

Not integrable

Time = 0.11 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)(fx)^m}{ex^2 + d} dx$$

input `integrate((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*(f*x)^m/(e*x^2 + d), x)`

### Mupad [N/A]

Not integrable

Time = 3.79 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.26

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx = \int \frac{(fx)^m (a + b \operatorname{acosh}(\frac{1}{cx}))}{ex^2 + d} dx$$

input `int(((f*x)^m*(a + b*acosh(1/(c*x))))/(d + e*x^2),x)`

output `int(((f*x)^m*(a + b*acosh(1/(c*x))))/(d + e*x^2), x)`

**Reduce [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.87

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{d + ex^2} dx = f^m \left( \left( \int \frac{x^m}{ex^2 + d} dx \right) a + \left( \int \frac{x^m \operatorname{asech}(cx)}{ex^2 + d} dx \right) b \right)$$

input `int((f*x)^m*(a+b*asech(c*x))/(e*x^2+d),x)`output `f**m*(int(x**m/(d + e*x**2),x)*a + int((x**m*asech(c*x))/(d + e*x**2),x)*b)`

$$3.180 \quad \int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx$$

Optimal result	1418
Mathematica [N/A]	1418
Rubi [N/A]	1419
Maple [N/A]	1419
Fricas [N/A]	1420
Sympy [F(-1)]	1420
Maxima [N/A]	1420
Giac [N/A]	1421
Mupad [N/A]	1421
Reduce [N/A]	1421

### Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \operatorname{Int} \left( \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2}, x \right)$$

output `Defer(Int)((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^2,x)`

### Mathematica [N/A]

Not integrable

Time = 4.94 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx$$

input `Integrate[((f*x)^m*(a + b*ArcSech[c*x]))/(d + e*x^2)^2,x]`

output `Integrate[((f*x)^m*(a + b*ArcSech[c*x]))/(d + e*x^2)^2, x]`

**Rubi [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx$$

↓ 6865

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx$$

input

```
Int[((f*x)^m*(a + b*ArcSech[c*x]))/(d + e*x^2)^2,x]
```

output

```
$Aborted
```

**Maple [N/A]**

Not integrable

Time = 1.25 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{arcsech}(cx))}{(x^2e + d)^2} dx$$

input

```
int((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^2,x)
```

output

```
int((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^2,x)
```

**Fricas [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.57

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)(fx)^m}{(ex^2 + d)^2} dx$$

input `integrate((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*arcsech(c*x) + a)*(f*x)^m/(e^2*x^4 + 2*d*e*x^2 + d^2), x)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \text{Timed out}$$

input `integrate((f*x)**m*(a+b*asech(c*x))/(e*x**2+d)**2,x)`

output `Timed out`

**Maxima [N/A]**

Not integrable

Time = 0.16 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)(fx)^m}{(ex^2 + d)^2} dx$$

input `integrate((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="maxima")`

output `integrate((b*arcsech(c*x) + a)*(f*x)^m/(e*x^2 + d)^2, x)`

**Giac [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)(fx)^m}{(ex^2 + d)^2} dx$$

input `integrate((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*(f*x)^m/(e*x^2 + d)^2, x)`

**Mupad [N/A]**

Not integrable

Time = 3.72 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.26

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = \int \frac{(fx)^m (a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^2} dx$$

input `int(((f*x)^m*(a + b*acosh(1/(c*x))))/(d + e*x^2)^2,x)`

output `int(((f*x)^m*(a + b*acosh(1/(c*x))))/(d + e*x^2)^2, x)`

**Reduce [N/A]**

Not integrable

Time = 0.91 (sec) , antiderivative size = 65, normalized size of antiderivative = 2.83

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^2} dx = f^m \left( \left( \int \frac{x^m}{e^2 x^4 + 2de x^2 + d^2} dx \right) a + \left( \int \frac{x^m \operatorname{asech}(cx)}{e^2 x^4 + 2de x^2 + d^2} dx \right) b \right)$$

input `int((f*x)^m*(a+b*asech(c*x))/(e*x^2+d)^2,x)`

output `f**m*(int(x**m/(d**2 + 2*d*e*x**2 + e**2*x**4),x)*a + int((x**m*asech(c*x))/(d**2 + 2*d*e*x**2 + e**2*x**4),x)*b)`

### 3.181 $\int (fx)^m (d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx$

Optimal result	1423
Mathematica [N/A]	1423
Rubi [N/A]	1424
Maple [N/A]	1424
Fricas [N/A]	1425
Sympy [F(-1)]	1425
Maxima [N/A]	1425
Giac [N/A]	1426
Mupad [N/A]	1426
Reduce [N/A]	1427

#### Optimal result

Integrand size = 25, antiderivative size = 25

$$\int (fx)^m (d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \operatorname{Int}\left((fx)^m (d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)), x\right)$$

output

```
Defer(Int)((f*x)^m*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x)
```

#### Mathematica [N/A]

Not integrable

Time = 0.94 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08

$$\int (fx)^m (d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int (fx)^m (d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx$$

input

```
Integrate[(f*x)^m*(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]),x]
```

output

```
Integrate[(f*x)^m*(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]), x]
```



**Rubi [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2)^{3/2} (fx)^m (a + b \operatorname{sech}^{-1}(cx)) dx$$

↓ 6865

$$\int (d + ex^2)^{3/2} (fx)^m (a + b \operatorname{sech}^{-1}(cx)) dx$$

input `Int[(f*x)^m*(d + e*x^2)^(3/2)*(a + b*ArcSech[c*x]),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.50 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int (fx)^m (x^2e + d)^{\frac{3}{2}} (a + b \operatorname{arcsech}(cx)) dx$$

input `int((f*x)^m*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x)`

output `int((f*x)^m*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x)`

**Fricas [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.68

$$\int (fx)^m (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^{\frac{3}{2}} (b \operatorname{ar} \operatorname{sech}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral((a*e*x^2 + a*d + (b*e*x^2 + b*d)*arcsech(c*x))*sqrt(e*x^2 + d)*(f*x)^m, x)`

**Sympy [F(-1)]**

Timed out.

$$\int (fx)^m (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \text{Timed out}$$

input `integrate((f*x)**m*(e*x**2+d)**(3/2)*(a+b*asech(c*x)),x)`

output `Timed out`

**Maxima [N/A]**

Not integrable

Time = 0.16 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int (fx)^m (d + ex^2)^{3/2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^{\frac{3}{2}} (b \operatorname{ar} \operatorname{sech}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `integrate((e*x^2 + d)^(3/2)*(b*arcsech(c*x) + a)*(f*x)^m, x)`

### Giac [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int (fx)^m (d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int (ex^2 + d)^{\frac{3}{2}} (b \operatorname{ar} \operatorname{sech}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^(3/2)*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate((e*x^2 + d)^(3/2)*(b*arcsech(c*x) + a)*(f*x)^m, x)`

### Mupad [N/A]

Not integrable

Time = 3.68 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.16

$$\int (fx)^m (d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int (fx)^m (ex^2 + d)^{3/2} \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int((f*x)^m*(d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))),x)`

output `int((f*x)^m*(d + e*x^2)^(3/2)*(a + b*acosh(1/(c*x))), x)`

**Reduce [N/A]**

Not integrable

Time = 200.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int (fx)^m (d + ex^2)^{3/2} (a + b\operatorname{sech}^{-1}(cx)) dx = \int (fx)^m (ex^2 + d)^{\frac{3}{2}} (\operatorname{asech}(cx)b + a) dx$$

input `int((f*x)^m*(e*x^2+d)^(3/2)*(a+b*asech(c*x)),x)`output `int((f*x)^m*(e*x^2+d)^(3/2)*(a+b*asech(c*x)),x)`

### 3.182 $\int (fx)^m \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx$

Optimal result	1428
Mathematica [N/A]	1428
Rubi [N/A]	1429
Maple [N/A]	1429
Fricas [N/A]	1430
Sympy [N/A]	1430
Maxima [N/A]	1430
Giac [N/A]	1431
Mupad [N/A]	1431
Reduce [N/A]	1432

#### Optimal result

Integrand size = 25, antiderivative size = 25

$$\int (fx)^m \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \operatorname{Int}\left((fx)^m \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)), x\right)$$

output `Defer(Int)((f*x)^m*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x)`

#### Mathematica [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08

$$\int (fx)^m \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int (fx)^m \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx$$

input `Integrate[(f*x)^m*Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]),x]`

output `Integrate[(f*x)^m*Sqrt[d + e*x^2]*(a + b*ArcSech[c*x]), x]`

**Rubi [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{d + ex^2}(fx)^m (a + b\operatorname{sech}^{-1}(cx)) dx$$

↓ 6865

$$\int \sqrt{d + ex^2}(fx)^m (a + b\operatorname{sech}^{-1}(cx)) dx$$

input `Int[(f*x)^m*sqrt[d + e*x^2]*(a + b*ArcSech[c*x]),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.51 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int (fx)^m \sqrt{x^2e + d}(a + b \operatorname{arcsech}(cx)) dx$$

input `int((f*x)^m*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x)`

output `int((f*x)^m*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x)`

**Fricas [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int (fx)^m \sqrt{d+ex^2} (a+b\operatorname{sech}^{-1}(cx)) dx = \int \sqrt{ex^2+d} (b \operatorname{arsech}(cx) + a) (fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)*(f*x)^m, x)`

**Sympy [N/A]**

Not integrable

Time = 10.97 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int (fx)^m \sqrt{d+ex^2} (a+b\operatorname{sech}^{-1}(cx)) dx = \int (fx)^m (a+b \operatorname{asech}(cx)) \sqrt{d+ex^2} dx$$

input `integrate((f*x)**m*(e*x**2+d)**(1/2)*(a+b*asech(c*x)),x)`

output `Integral((f*x)**m*(a + b*asech(c*x))*sqrt(d + e*x**2), x)`

**Maxima [N/A]**

Not integrable

Time = 0.14 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int (fx)^m \sqrt{d+ex^2} (a+b\operatorname{sech}^{-1}(cx)) dx = \int \sqrt{ex^2+d} (b \operatorname{arsech}(cx) + a) (fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="maxima")`

output `integrate(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)*(f*x)^m, x)`

### Giac [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int (fx)^m \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int \sqrt{ex^2 + d} (b \operatorname{ar} \operatorname{sech}(cx) + a) (fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^(1/2)*(a+b*arcsech(c*x)),x, algorithm="giac")`

output `integrate(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)*(f*x)^m, x)`

### Mupad [N/A]

Not integrable

Time = 3.56 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.16

$$\int (fx)^m \sqrt{d + ex^2} (a + b \operatorname{sech}^{-1}(cx)) dx = \int (fx)^m \sqrt{ex^2 + d} \left( a + b \operatorname{acosh}\left(\frac{1}{cx}\right) \right) dx$$

input `int((f*x)^m*(d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))),x)`

output `int((f*x)^m*(d + e*x^2)^(1/2)*(a + b*acosh(1/(c*x))), x)`



**Reduce [N/A]**

Not integrable

Time = 200.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int (fx)^m \sqrt{d+ex^2} (a+b\operatorname{sech}^{-1}(cx)) dx = \int (fx)^m \sqrt{ex^2+d} (a\operatorname{sech}(cx) b+a) dx$$

input `int((f*x)^m*(e*x^2+d)^(1/2)*(a+b*asech(c*x)),x)`output `int((f*x)^m*(e*x^2+d)^(1/2)*(a+b*asech(c*x)),x)`

$$3.183 \quad \int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx$$

Optimal result	1433
Mathematica [N/A]	1433
Rubi [N/A]	1434
Maple [N/A]	1434
Fricas [N/A]	1435
Sympy [N/A]	1435
Maxima [N/A]	1435
Giac [N/A]	1436
Mupad [N/A]	1436
Reduce [N/A]	1437

### Optimal result

Integrand size = 25, antiderivative size = 25

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \operatorname{Int} \left( \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}}, x \right)$$

output `Defer(Int)((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x)`

### Mathematica [N/A]

Not integrable

Time = 0.97 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx$$

input `Integrate[((f*x)^m*(a + b*ArcSech[c*x]))/Sqrt[d + e*x^2],x]`

output `Integrate[((f*x)^m*(a + b*ArcSech[c*x]))/Sqrt[d + e*x^2], x]`

**Rubi [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx$$

↓ 6865

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx$$

input `Int[((f*x)^m*(a + b*ArcSech[c*x]))/Sqrt[d + e*x^2],x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.52 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{(fx)^m (a + b \operatorname{arcsech}(cx))}{\sqrt{x^2 e + d}} dx$$

input `int((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x)`

output `int((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x)`

**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{(b \operatorname{arsech}(cx) + a)(fx)^m}{\sqrt{ex^2 + d}} dx$$

input `integrate((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral((b*arcsech(c*x) + a)*(f*x)^m/sqrt(e*x^2 + d), x)`

**Sympy [N/A]**

Not integrable

Time = 4.47 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{(fx)^m (a + b \operatorname{asech}(cx))}{\sqrt{d + ex^2}} dx$$

input `integrate((f*x)**m*(a+b*asech(c*x))/(e*x**2+d)**(1/2),x)`

output `Integral((f*x)**m*(a + b*asech(c*x))/sqrt(d + e*x**2), x)`

**Maxima [N/A]**

Not integrable

Time = 0.17 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{(b \operatorname{arsech}(cx) + a)(fx)^m}{\sqrt{ex^2 + d}} dx$$

input `integrate((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="maxima")`

output `integrate((b*arcsech(c*x) + a)*(f*x)^m/sqrt(e*x^2 + d), x)`

### Giac [N/A]

Not integrable

Time = 0.12 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)(fx)^m}{\sqrt{ex^2 + d}} dx$$

input `integrate((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*(f*x)^m/sqrt(e*x^2 + d), x)`

### Mupad [N/A]

Not integrable

Time = 3.60 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.16

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{(fx)^m (a + b \operatorname{acosh}(\frac{1}{cx}))}{\sqrt{ex^2 + d}} dx$$

input `int(((f*x)^m*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(1/2),x)`

output `int(((f*x)^m*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(1/2), x)`

**Reduce [N/A]**

Not integrable

Time = 200.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{d + ex^2}} dx = \int \frac{(fx)^m (a \operatorname{sech}(cx) b + a)}{\sqrt{ex^2 + d}} dx$$

input `int((f*x)^m*(a+b*asech(c*x))/(e*x^2+d)^(1/2),x)`output `int((f*x)^m*(a+b*asech(c*x))/(e*x^2+d)^(1/2),x)`

$$3.184 \quad \int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

Optimal result	1438
Mathematica [N/A]	1438
Rubi [N/A]	1439
Maple [N/A]	1439
Fricas [N/A]	1440
Sympy [N/A]	1440
Maxima [N/A]	1441
Giac [N/A]	1441
Mupad [N/A]	1441
Reduce [N/A]	1442

### Optimal result

Integrand size = 25, antiderivative size = 25

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \operatorname{Int} \left( \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}}, x \right)$$

output `Defer(Int)((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x)`

### Mathematica [N/A]

Not integrable

Time = 1.19 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

input `Integrate[((f*x)^m*(a + b*ArcSech[c*x]))/(d + e*x^2)^(3/2),x]`

output `Integrate[((f*x)^m*(a + b*ArcSech[c*x]))/(d + e*x^2)^(3/2), x]`

**Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

↓ 6865

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx$$

input `Int[((f*x)^m*(a + b*ArcSech[c*x]))/(d + e*x^2)^(3/2),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.53 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{(fx)^m (a + b \operatorname{arcsech}(cx))}{(x^2 e + d)^{\frac{3}{2}}} dx$$

input `int((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x)`

output `int((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x)`



**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.80

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{(b \operatorname{arsech}(cx) + a)(fx)^m}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)*(b*arcsech(c*x) + a)*(f*x)^m/(e^2*x^4 + 2*d*e*x^2 + d^2), x)`

**Sympy [N/A]**

Not integrable

Time = 30.18 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{(fx)^m (a + b \operatorname{asech}(cx))}{(d + ex^2)^{\frac{3}{2}}} dx$$

input `integrate((f*x)**m*(a+b*asech(c*x))/(e*x**2+d)**(3/2),x)`

output `Integral((f*x)**m*(a + b*asech(c*x))/(d + e*x**2)**(3/2), x)`

**Maxima [N/A]**

Not integrable

Time = 0.15 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)(fx)^m}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="maxima")`

output `integrate((b*arcsech(c*x) + a)*(f*x)^m/(e*x^2 + d)^(3/2), x)`

**Giac [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)(fx)^m}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arcsech(c*x))/(e*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*(f*x)^m/(e*x^2 + d)^(3/2), x)`

**Mupad [N/A]**

Not integrable

Time = 3.68 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.16

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{(fx)^m (a + b \operatorname{acosh}(\frac{1}{cx}))}{(ex^2 + d)^{3/2}} dx$$

input `int(((f*x)^m*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(3/2),x)`

output `int(((f*x)^m*(a + b*acosh(1/(c*x))))/(d + e*x^2)^(3/2), x)`

### Reduce [N/A]

Not integrable

Time = 200.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{sech}^{-1}(cx))}{(d + ex^2)^{3/2}} dx = \int \frac{(fx)^m (a \operatorname{sech}(cx) b + a)}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `int((f*x)^m*(a+b*asech(c*x))/(e*x^2+d)^(3/2),x)`

output `int((f*x)^m*(a+b*asech(c*x))/(e*x^2+d)^(3/2),x)`

**3.185**  $\int \frac{x^{11} (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx$

Optimal result	1443
Mathematica [A] (verified)	1444
Rubi [A] (verified)	1444
Maple [F]	1447
Fricas [A] (verification not implemented)	1447
Sympy [F(-1)]	1448
Maxima [F]	1448
Giac [F(-2)]	1449
Mupad [F(-1)]	1449
Reduce [F]	1450

**Optimal result**

Integrand size = 26, antiderivative size = 473

$$\int \frac{x^{11} (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx = -\frac{4b\sqrt{1 - c^2 x^2} \sqrt{1 + c^2 x^2}}{15c^{13} \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} x} + \frac{7b\sqrt{1 - c^2 x^2} (1 + c^2 x^2)^{3/2}}{90c^{13} \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} x}$$

$$- \frac{13b\sqrt{1 - c^2 x^2} (1 + c^2 x^2)^{5/2}}{150c^{13} \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} x}$$

$$+ \frac{3b\sqrt{1 - c^2 x^2} (1 + c^2 x^2)^{7/2}}{70c^{13} \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} x} - \frac{b\sqrt{1 - c^2 x^2} (1 + c^2 x^2)^{9/2}}{90c^{13} \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} x}$$

$$- \frac{\sqrt{1 - c^4 x^4} (a + b \operatorname{sech}^{-1}(cx))}{2c^{12}}$$

$$+ \frac{(1 - c^4 x^4)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{3c^{12}}$$

$$- \frac{(1 - c^4 x^4)^{5/2} (a + b \operatorname{sech}^{-1}(cx))}{10c^{12}}$$

$$+ \frac{4b\sqrt{1 - c^2 x^2} \operatorname{arctanh}(\sqrt{1 + c^2 x^2})}{15c^{13} \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} x}$$

output

$$\begin{aligned} & -4/15*b*(-c^2*x^2+1)^{(1/2)}*(c^2*x^2+1)^{(1/2)}/c^{13}/(-1+1/c/x)^{(1/2)}/(1+1/c/x)^{(1/2)}/x+7/90*b*(-c^2*x^2+1)^{(1/2)}*(c^2*x^2+1)^{(3/2)}/c^{13}/(-1+1/c/x)^{(1/2)}/(1+1/c/x)^{(1/2)}/x-13/150*b*(-c^2*x^2+1)^{(1/2)}*(c^2*x^2+1)^{(5/2)}/c^{13}/(-1+1/c/x)^{(1/2)}/(1+1/c/x)^{(1/2)}/x+3/70*b*(-c^2*x^2+1)^{(1/2)}*(c^2*x^2+1)^{(7/2)}/c^{13}/(-1+1/c/x)^{(1/2)}/(1+1/c/x)^{(1/2)}/x-1/90*b*(-c^2*x^2+1)^{(1/2)}*(c^2*x^2+1)^{(9/2)}/c^{13}/(-1+1/c/x)^{(1/2)}/(1+1/c/x)^{(1/2)}/x-1/2*(-c^4*x^4+1)^{(1/2)}*(a+b*\operatorname{arcsech}(c*x))/c^{12}+1/3*(-c^4*x^4+1)^{(3/2)}*(a+b*\operatorname{arcsech}(c*x))/c^{12}-1/10*(-c^4*x^4+1)^{(5/2)}*(a+b*\operatorname{arcsech}(c*x))/c^{12}+4/15*b*(-c^2*x^2+1)^{(1/2)}*a \operatorname{rctanh}((c^2*x^2+1)^{(1/2)})/c^{13}/(-1+1/c/x)^{(1/2)}/(1+1/c/x)^{(1/2)}/x \end{aligned}$$
**Mathematica [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 213, normalized size of antiderivative = 0.45

$$\int \frac{x^{11}(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx$$

$$= \frac{-105a\sqrt{1 - c^4 x^4}(8 + 4c^4 x^4 + 3c^8 x^8) + \frac{b\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^4 x^4}(768+36c^2 x^2+78c^4 x^4+5c^6 x^6+35c^8 x^8)}{-1+cx} - 105b\sqrt{1 - c^4 x^4}(8 - 3150c^{12}}{3150c^{12}}$$

input

$$\text{Integrate}[(x^{11}*(a + b*\text{ArcSech}[c*x]))/\text{Sqrt}[1 - c^4*x^4], x]$$

output

$$\begin{aligned} & (-105*a*\text{Sqrt}[1 - c^4*x^4]*(8 + 4*c^4*x^4 + 3*c^8*x^8) + (b*\text{Sqrt}[(1 - c*x)/(1 + c*x)]*\text{Sqrt}[1 - c^4*x^4]*(768 + 36*c^2*x^2 + 78*c^4*x^4 + 5*c^6*x^6 + 35*c^8*x^8))/(-1 + c*x) - 105*b*\text{Sqrt}[1 - c^4*x^4]*(8 + 4*c^4*x^4 + 3*c^8*x^8)*\text{ArcSech}[c*x] + 840*b*\text{Log}[x*(1 - c*x)] - 840*b*\text{Log}[1 - c*x - \text{Sqrt}[(1 - c*x)/(1 + c*x)]]*\text{Sqrt}[1 - c^4*x^4])/(3150*c^{12}) \end{aligned}$$
**Rubi [A] (verified)**Time = 0.73 (sec) , antiderivative size = 239, normalized size of antiderivative = 0.51, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {6863, 27, 1388, 2331, 2123, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{11}(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4x^4}} dx$$

↓ 6863

$$\frac{b\sqrt{1 - c^2x^2} \int -\frac{\sqrt{1 - c^4x^4}(3c^8x^8 + 4c^4x^4 + 8)}{30c^{12}x\sqrt{1 - c^2x^2}} dx}{cx\sqrt{\frac{1}{cx} - 1}\sqrt{\frac{1}{cx} + 1}} - \frac{(1 - c^4x^4)^{5/2}(a + b\operatorname{sech}^{-1}(cx))}{10c^{12}} +$$

$$\frac{(1 - c^4x^4)^{3/2}(a + b\operatorname{sech}^{-1}(cx))}{3c^{12}} - \frac{\sqrt{1 - c^4x^4}(a + b\operatorname{sech}^{-1}(cx))}{2c^{12}}$$

↓ 27

$$-\frac{b\sqrt{1 - c^2x^2} \int \frac{\sqrt{1 - c^4x^4}(3c^8x^8 + 4c^4x^4 + 8)}{x\sqrt{1 - c^2x^2}} dx}{30c^{13}x\sqrt{\frac{1}{cx} - 1}\sqrt{\frac{1}{cx} + 1}} - \frac{(1 - c^4x^4)^{5/2}(a + b\operatorname{sech}^{-1}(cx))}{10c^{12}} +$$

$$\frac{(1 - c^4x^4)^{3/2}(a + b\operatorname{sech}^{-1}(cx))}{3c^{12}} - \frac{\sqrt{1 - c^4x^4}(a + b\operatorname{sech}^{-1}(cx))}{2c^{12}}$$

↓ 1388

$$-\frac{b\sqrt{1 - c^2x^2} \int \frac{\sqrt{c^2x^2 + 1}(3c^8x^8 + 4c^4x^4 + 8)}{x} dx}{30c^{13}x\sqrt{\frac{1}{cx} - 1}\sqrt{\frac{1}{cx} + 1}} - \frac{(1 - c^4x^4)^{5/2}(a + b\operatorname{sech}^{-1}(cx))}{10c^{12}} +$$

$$\frac{(1 - c^4x^4)^{3/2}(a + b\operatorname{sech}^{-1}(cx))}{3c^{12}} - \frac{\sqrt{1 - c^4x^4}(a + b\operatorname{sech}^{-1}(cx))}{2c^{12}}$$

↓ 2331

$$-\frac{b\sqrt{1 - c^2x^2} \int \frac{\sqrt{c^2x^2 + 1}(3c^8x^8 + 4c^4x^4 + 8)}{x^2} dx^2}{60c^{13}x\sqrt{\frac{1}{cx} - 1}\sqrt{\frac{1}{cx} + 1}} - \frac{(1 - c^4x^4)^{5/2}(a + b\operatorname{sech}^{-1}(cx))}{10c^{12}} +$$

$$\frac{(1 - c^4x^4)^{3/2}(a + b\operatorname{sech}^{-1}(cx))}{3c^{12}} - \frac{\sqrt{1 - c^4x^4}(a + b\operatorname{sech}^{-1}(cx))}{2c^{12}}$$

↓ 2123

$$-\frac{b\sqrt{1 - c^2x^2} \int \left(3c^2(c^2x^2 + 1)^{7/2} - 9c^2(c^2x^2 + 1)^{5/2} + 13c^2(c^2x^2 + 1)^{3/2} - 7c^2\sqrt{c^2x^2 + 1} + \frac{8\sqrt{c^2x^2 + 1}}{x^2}\right) dx^2}{60c^{13}x\sqrt{\frac{1}{cx} - 1}\sqrt{\frac{1}{cx} + 1}}$$

$$\frac{(1 - c^4x^4)^{5/2}(a + b\operatorname{sech}^{-1}(cx))}{10c^{12}} + \frac{(1 - c^4x^4)^{3/2}(a + b\operatorname{sech}^{-1}(cx))}{3c^{12}} -$$

$$\frac{\sqrt{1 - c^4x^4}(a + b\operatorname{sech}^{-1}(cx))}{2c^{12}}$$

↓ 2009

$$\frac{-\frac{(1-c^4x^4)^{5/2}(a+b\operatorname{sech}^{-1}(cx))}{10c^{12}} + \frac{(1-c^4x^4)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{3c^{12}} - \frac{\sqrt{1-c^4x^4}(a+b\operatorname{sech}^{-1}(cx))}{2c^{12}}}{b\sqrt{1-c^2x^2}\left(-16\operatorname{arctanh}\left(\sqrt{c^2x^2+1}\right) + \frac{2}{3}(c^2x^2+1)^{9/2} - \frac{18}{7}(c^2x^2+1)^{7/2} + \frac{26}{5}(c^2x^2+1)^{5/2} - \frac{14}{3}(c^2x^2+1)^{3/2}\right)}}{60c^{13}x\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}}$$

input `Int[(x^11*(a + b*ArcSech[c*x]))/Sqrt[1 - c^4*x^4], x]`

output `-1/2*(Sqrt[1 - c^4*x^4]*(a + b*ArcSech[c*x]))/c^12 + ((1 - c^4*x^4)^(3/2)*(a + b*ArcSech[c*x]))/(3*c^12) - ((1 - c^4*x^4)^(5/2)*(a + b*ArcSech[c*x]))/(10*c^12) - (b*Sqrt[1 - c^2*x^2]*(16*Sqrt[1 + c^2*x^2] - (14*(1 + c^2*x^2)^(3/2))/3 + (26*(1 + c^2*x^2)^(5/2))/5 - (18*(1 + c^2*x^2)^(7/2))/7 + (2*(1 + c^2*x^2)^(9/2))/3 - 16*ArcTanh[Sqrt[1 + c^2*x^2]]))/(60*c^13*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]*x)`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 1388 `Int[(u_)*((a_) + (c_.)*(x_)^(n2_.))^ (p_.)*((d_) + (e_.)*(x_)^(n_.))^ (q_.), x_Symbol] := Int[u*(d + e*x^n)^(p + q)*(a/d + (c/e)*x^n)^p, x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[c*d^2 + a*e^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[d, 0]))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2123 `Int[(Px_)*((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] := Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && PolyQ[Px, x] && (IntegersQ[m, n] || IGtQ[m, -2])`

rule 2331

```
Int[(Pq_)*(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 S
ubst[Int[x^((m - 1)/2)*SubstFor[x^2, Pq, x]*(a + b*x)^p, x], x, x^2], x] /;
FreeQ[{a, b, p}, x] && PolyQ[Pq, x^2] && IntegerQ[(m - 1)/2]
```

rule 6863

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_)]*(u_), x_Symbol] := With[{v = IntHid
e[u, x]}, Simp[(a + b*ArcSech[c*x]) v, x] + Simp[b*(Sqrt[1 - c^2*x^2]/(c*
x*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) Int[SimplifyIntegrand[v/(x*Sqrt[
1 - c^2*x^2]), x], x], x] /; InverseFunctionFreeQ[v, x] /; FreeQ[{a, b, c}
, x]
```

**Maple [F]**

$$\int \frac{x^{11}(a + b \operatorname{arcsech}(cx))}{\sqrt{-c^4x^4 + 1}} dx$$

input

```
int(x^11*(a+b*arcsech(c*x))/(-c^4*x^4+1)^(1/2),x)
```

output

```
int(x^11*(a+b*arcsech(c*x))/(-c^4*x^4+1)^(1/2),x)
```

**Fricas [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 393, normalized size of antiderivative = 0.83

$$\int \frac{x^{11}(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4x^4}} dx =$$

$$\frac{105(3bc^{10}x^{10} - 3bc^8x^8 + 4bc^6x^6 - 4bc^4x^4 + 8bc^2x^2 - 8b)\sqrt{-c^4x^4 + 1} \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}+1}}{cx}\right) - (35bc^9}{-}$$

input

```
integrate(x^11*(a+b*arcsech(c*x))/(-c^4*x^4+1)^(1/2),x, algorithm="fricas"
)
```



output

```
-1/3150*(105*(3*b*c^10*x^10 - 3*b*c^8*x^8 + 4*b*c^6*x^6 - 4*b*c^4*x^4 + 8*
b*c^2*x^2 - 8*b)*sqrt(-c^4*x^4 + 1)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)
) + 1)/(c*x)) - (35*b*c^9*x^9 + 5*b*c^7*x^7 + 78*b*c^5*x^5 + 36*b*c^3*x^3
+ 768*b*c*x)*sqrt(-c^4*x^4 + 1)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 420*(b*c^
2*x^2 - b)*log((c^2*x^2 + sqrt(-c^4*x^4 + 1)*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*
x^2)) - 1)/(c^2*x^2 - 1)) - 420*(b*c^2*x^2 - b)*log(-(c^2*x^2 - sqrt(-c^4*
x^4 + 1)*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/(c^2*x^2 - 1)) + 105*(3*a
*c^10*x^10 - 3*a*c^8*x^8 + 4*a*c^6*x^6 - 4*a*c^4*x^4 + 8*a*c^2*x^2 - 8*a)*
sqrt(-c^4*x^4 + 1))/(c^14*x^2 - c^12)
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{x^{11}(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx = \text{Timed out}$$

input

```
integrate(x**11*(a+b*asech(c*x))/(-c**4*x**4+1)**(1/2),x)
```

output

Timed out

**Maxima [F]**

$$\int \frac{x^{11}(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx = \int \frac{(b \operatorname{ar} \operatorname{sech}(cx) + a)x^{11}}{\sqrt{-c^4 x^4 + 1}} dx$$

input

```
integrate(x^11*(a+b*arcsech(c*x))/(-c^4*x^4+1)^(1/2),x, algorithm="maxima"
)
```

output

```
-1/30*a*(3*(-c^4*x^4 + 1)^(5/2)/c^12 - 10*(-c^4*x^4 + 1)^(3/2)/c^12 + 15*sqrt(-c^4*x^4 + 1)/c^12) + 1/30*b*((3*c^12*x^12 + c^8*x^8 + 4*c^4*x^4 - 8)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)/(sqrt(c^2*x^2 + 1)*sqrt(c*x + 1)*sqrt(-c*x + 1)*c^12) - 30*integrate(1/30*(30*c^10*x^21*log(c) + 60*c^10*x^21*log(sqrt(x)) + (60*c^10*x^21*log(sqrt(x)) + (3*c^10*x^10*(10*log(c) + 1) + 3*c^8*x^8 + 4*c^6*x^6 + 4*c^4*x^4 + 8*c^2*x^2 + 8)*x^11)*e^(1/2*log(c*x + 1) + 1/2*log(-c*x + 1)))/((c^10*x^10*e^(log(c*x + 1) + log(-c*x + 1)) + c^10*x^10*e^(1/2*log(c*x + 1) + 1/2*log(-c*x + 1)))*sqrt(c^2*x^2 + 1)), x))
```

**Giac [F(-2)]**

Exception generated.

$$\int \frac{x^{11}(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx = \text{Exception raised: TypeError}$$

input

```
integrate(x^11*(a+b*arcsech(c*x))/(-c^4*x^4+1)^(1/2),x, algorithm="giac")
```

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^{11}(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx = \int \frac{x^{11}(a + b \operatorname{acosh}(\frac{1}{cx}))}{\sqrt{1 - c^4 x^4}} dx$$

input

```
int((x^11*(a + b*acosh(1/(c*x))))/(1 - c^4*x^4)^(1/2),x)
```

output

```
int((x^11*(a + b*acosh(1/(c*x))))/(1 - c^4*x^4)^(1/2), x)
```

**Reduce [F]**

$$\int \frac{x^{11}(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx$$

$$= \frac{-3\sqrt{-c^4 x^4 + 1} a c^8 x^8 - 4\sqrt{-c^4 x^4 + 1} a c^4 x^4 - 8\sqrt{-c^4 x^4 + 1} a - 30 \left( \int \frac{\sqrt{-c^4 x^4 + 1} a \operatorname{sech}(cx) x^{11}}{c^4 x^4 - 1} dx \right) b c^{12}}{30 c^{12}}$$

input `int(x^11*(a+b*asech(c*x))/(-c^4*x^4+1)^(1/2),x)`

output `( - 3*sqrt( - c**4*x**4 + 1)*a*c**8*x**8 - 4*sqrt( - c**4*x**4 + 1)*a*c**4*x**4 - 8*sqrt( - c**4*x**4 + 1)*a - 30*int((sqrt( - c**4*x**4 + 1)*asech(c*x)*x**11)/(c**4*x**4 - 1),x)*b*c**12)/(30*c**12)`

**3.186** 
$$\int \frac{x^7 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx$$

Optimal result	1451
Mathematica [A] (verified)	1452
Rubi [A] (warning: unable to verify)	1452
Maple [F]	1455
Fricas [A] (verification not implemented)	1455
Sympy [F]	1456
Maxima [F]	1456
Giac [F(-2)]	1457
Mupad [F(-1)]	1457
Reduce [F]	1457

**Optimal result**

Integrand size = 26, antiderivative size = 316

$$\int \frac{x^7 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx = -\frac{b\sqrt{1 - c^2 x^2} \sqrt{1 + c^2 x^2}}{3c^9 \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}} + \frac{b\sqrt{1 - c^2 x^2} (1 + c^2 x^2)^{3/2}}{18c^9 \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}} - \frac{b\sqrt{1 - c^2 x^2} (1 + c^2 x^2)^{5/2}}{30c^9 \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}} - \frac{\sqrt{1 - c^4 x^4} (a + b \operatorname{sech}^{-1}(cx))}{2c^8} + \frac{(1 - c^4 x^4)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{6c^8} + \frac{b\sqrt{1 - c^2 x^2} \operatorname{arctanh}(\sqrt{1 + c^2 x^2})}{3c^9 \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}}$$

output

```
-1/3*b*(-c^2*x^2+1)^(1/2)*(c^2*x^2+1)^(1/2)/c^9/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)/x+1/18*b*(-c^2*x^2+1)^(1/2)*(c^2*x^2+1)^(3/2)/c^9/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)/x-1/30*b*(-c^2*x^2+1)^(1/2)*(c^2*x^2+1)^(5/2)/c^9/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)/x-1/2*(-c^4*x^4+1)^(1/2)*(a+b*arcsech(c*x))/c^8+1/6*(-c^4*x^4+1)^(3/2)*(a+b*arcsech(c*x))/c^8+1/3*b*(-c^2*x^2+1)^(1/2)*arctanh((c^2*x^2+1)^(1/2))/c^9/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)/x
```

**Mathematica [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.56

$$\int \frac{x^7(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx$$

$$= \frac{-15a\sqrt{1 - c^4 x^4}(2 + c^4 x^4) + \frac{b\sqrt{\frac{1-cx}{1+cx}}\sqrt{1-c^4 x^4}(28+c^2 x^2+3c^4 x^4)}{-1+cx} - 15b\sqrt{1 - c^4 x^4}(2 + c^4 x^4) \operatorname{sech}^{-1}(cx) + 30b \log}{90c^8}$$

input `Integrate[(x^7*(a + b*ArcSech[c*x]))/Sqrt[1 - c^4*x^4],x]`output `(-15*a*Sqrt[1 - c^4*x^4]*(2 + c^4*x^4) + (b*Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^4*x^4]*(28 + c^2*x^2 + 3*c^4*x^4))/(-1 + c*x) - 15*b*Sqrt[1 - c^4*x^4]*(2 + c^4*x^4)*ArcSech[c*x] + 30*b*Log[x*(1 - c*x)] - 30*b*Log[1 - c*x - Sqrt[(1 - c*x)/(1 + c*x)]*Sqrt[1 - c^4*x^4]])/(90*c^8)`**Rubi [A] (warning: unable to verify)**Time = 0.56 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.53, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {6863, 27, 1388, 1579, 517, 25, 1584, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^7(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx$$

$$\downarrow \text{6863}$$

$$\frac{b\sqrt{1 - c^2 x^2} \int -\frac{\sqrt{1 - c^4 x^4}(c^4 x^4 + 2)}{6c^8 x \sqrt{1 - c^2 x^2}} dx}{cx \sqrt{\frac{1}{cx} - 1} \sqrt{\frac{1}{cx} + 1}} + \frac{(1 - c^4 x^4)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{6c^8} -$$

$$\frac{\sqrt{1 - c^4 x^4}(a + b \operatorname{sech}^{-1}(cx))}{2c^8}$$

$$\downarrow \text{27}$$

$$\begin{aligned}
& -\frac{b\sqrt{1-c^2x^2} \int \frac{\sqrt{1-c^4x^4}(c^4x^4+2)}{x\sqrt{1-c^2x^2}} dx}{6c^9x\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}} + \frac{(1-c^4x^4)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{6c^8} - \\
& \quad \frac{\sqrt{1-c^4x^4}(a+b\operatorname{sech}^{-1}(cx))}{2c^8} \\
& \quad \downarrow \text{1388} \\
& -\frac{b\sqrt{1-c^2x^2} \int \frac{\sqrt{c^2x^2+1}(c^4x^4+2)}{x} dx}{6c^9x\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}} + \frac{(1-c^4x^4)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{6c^8} - \\
& \quad \frac{\sqrt{1-c^4x^4}(a+b\operatorname{sech}^{-1}(cx))}{2c^8} \\
& \quad \downarrow \text{1579} \\
& -\frac{b\sqrt{1-c^2x^2} \int \frac{\sqrt{c^2x^2+1}(c^4x^4+2)}{x^2} dx^2}{12c^9x\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}} + \frac{(1-c^4x^4)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{6c^8} - \\
& \quad \frac{\sqrt{1-c^4x^4}(a+b\operatorname{sech}^{-1}(cx))}{2c^8} \\
& \quad \downarrow \text{517} \\
& -\frac{b\sqrt{1-c^2x^2} \int -\frac{x^4(c^4x^8-2c^4x^4+3c^4)}{1-x^4} d\sqrt{c^2x^2+1}}{6c^{13}x\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}} + \frac{(1-c^4x^4)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{6c^8} - \\
& \quad \frac{\sqrt{1-c^4x^4}(a+b\operatorname{sech}^{-1}(cx))}{2c^8} \\
& \quad \downarrow \text{25} \\
& \frac{b\sqrt{1-c^2x^2} \int \frac{x^4(c^4x^8-2c^4x^4+3c^4)}{1-x^4} d\sqrt{c^2x^2+1}}{6c^{13}x\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}} + \frac{(1-c^4x^4)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{6c^8} - \\
& \quad \frac{\sqrt{1-c^4x^4}(a+b\operatorname{sech}^{-1}(cx))}{2c^8} \\
& \quad \downarrow \text{1584} \\
& \frac{b\sqrt{1-c^2x^2} \int \left(-c^4x^8+c^4x^4-2c^4+\frac{2c^4}{1-x^4}\right) d\sqrt{c^2x^2+1}}{6c^{13}x\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}} + \frac{(1-c^4x^4)^{3/2}(a+b\operatorname{sech}^{-1}(cx))}{6c^8} - \\
& \quad \frac{\sqrt{1-c^4x^4}(a+b\operatorname{sech}^{-1}(cx))}{2c^8} \\
& \quad \downarrow \text{2009}
\end{aligned}$$

$$\frac{(1 - c^4 x^4)^{3/2} (a + b \operatorname{sech}^{-1}(cx))}{6c^8} - \frac{\sqrt{1 - c^4 x^4} (a + b \operatorname{sech}^{-1}(cx))}{b\sqrt{1 - c^2 x^2} \left( -2c^4 \operatorname{arctanh}(\sqrt{c^2 x^2 + 1}) + \frac{c^4 x^{10}}{5} - \frac{c^4 x^6}{3} + 2c^4 \sqrt{c^2 x^2 + 1} \right)}$$

$$\frac{6c^{13} x \sqrt{\frac{1}{cx} - 1} \sqrt{\frac{1}{cx} + 1}}{6c^{13} x \sqrt{\frac{1}{cx} - 1} \sqrt{\frac{1}{cx} + 1}}$$

input `Int[(x^7*(a + b*ArcSech[c*x]))/Sqrt[1 - c^4*x^4],x]`

output `-1/2*(Sqrt[1 - c^4*x^4]*(a + b*ArcSech[c*x]))/c^8 + ((1 - c^4*x^4)^(3/2)*(a + b*ArcSech[c*x]))/(6*c^8) - (b*Sqrt[1 - c^2*x^2]*(-1/3*(c^4*x^6) + (c^4*x^10)/5 + 2*c^4*Sqrt[1 + c^2*x^2] - 2*c^4*ArcTanh[Sqrt[1 + c^2*x^2]]))/(6*c^13*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]*x)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 517 `Int[((e_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[2*(e^m/d^(m + 2*p + 1)) Subst[Int[x^(2*n + 1)*(-c + x^2)^m*(b*c^2 + a*d^2 - 2*b*c*x^2 + b*x^4)^p, x], x, Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && ILtQ[m, 0] && IntegerQ[n + 1/2]`

rule 1388 `Int[(u_)*((a_) + (c_)*(x_)^(n2_))^(p_)*((d_) + (e_)*(x_)^(n_))^(q_), x_Symbol] := Int[u*(d + e*x^n)^(p + q)*(a/d + (c/e)*x^n)^p, x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[c*d^2 + a*e^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[d, 0]))`

rule 1579 `Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, c, d, e, p, q}, x] && IntegerQ[(m + 1)/2]`

rule 1584 `Int[((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Int[ExpandIntegrand[(f*x)^m*(d + e*x^2)^q*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, q}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && IGtQ[q, -2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6863 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.)*(u_), x_Symbol] := With[{v = IntHide[u, x]}, Simp[(a + b*ArcSech[c*x]) v, x] + Simp[b*(Sqrt[1 - c^2*x^2]/(c*x*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)])) Int[SimplifyIntegrand[v/(x*Sqrt[1 - c^2*x^2]), x], x], x] /; InverseFunctionFreeQ[v, x] /; FreeQ[{a, b, c}, x]`

## Maple [F]

$$\int \frac{x^7(a + b \operatorname{arcsech}(cx))}{\sqrt{-c^4x^4 + 1}} dx$$

input `int(x^7*(a+b*arcsech(c*x))/(-c^4*x^4+1)^(1/2),x)`

output `int(x^7*(a+b*arcsech(c*x))/(-c^4*x^4+1)^(1/2),x)`

## Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 336, normalized size of antiderivative = 1.06

$$\int \frac{x^7(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4x^4}} dx =$$

$$15(bc^6x^6 - bc^4x^4 + 2bc^2x^2 - 2b)\sqrt{-c^4x^4 + 1} \log\left(\frac{cx\sqrt{-\frac{c^2x^2-1}{c^2x^2}+1}}{cx}\right) - (3bc^5x^5 + bc^3x^3 + 28bcx)\sqrt{-c^4x^4 + 1}$$

input `integrate(x^7*(a+b*arcsech(c*x))/(-c^4*x^4+1)^(1/2),x, algorithm="fricas")`



output

```
-1/90*(15*(b*c^6*x^6 - b*c^4*x^4 + 2*b*c^2*x^2 - 2*b)*sqrt(-c^4*x^4 + 1)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (3*b*c^5*x^5 + b*c^3*x^3 + 28*b*c*x)*sqrt(-c^4*x^4 + 1)*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 15*(b*c^2*x^2 - b)*log((c^2*x^2 + sqrt(-c^4*x^4 + 1)*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/(c^2*x^2 - 1)) - 15*(b*c^2*x^2 - b)*log(-(c^2*x^2 - sqrt(-c^4*x^4 + 1)*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/(c^2*x^2 - 1)) + 15*(a*c^6*x^6 - a*c^4*x^4 + 2*a*c^2*x^2 - 2*a)*sqrt(-c^4*x^4 + 1))/(c^10*x^2 - c^8)
```

**Sympy [F]**

$$\int \frac{x^7(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx = \int \frac{x^7(a + b \operatorname{asech}(cx))}{\sqrt{-(cx - 1)(cx + 1)(c^2 x^2 + 1)}} dx$$

input

```
integrate(x**7*(a+b*asech(c*x))/(-c**4*x**4+1)**(1/2), x)
```

output

```
Integral(x**7*(a + b*asech(c*x))/sqrt(-(c*x - 1)*(c*x + 1)*(c**2*x**2 + 1)), x)
```

**Maxima [F]**

$$\int \frac{x^7(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^7}{\sqrt{-c^4 x^4 + 1}} dx$$

input

```
integrate(x^7*(a+b*arcsech(c*x))/(-c^4*x^4+1)^(1/2), x, algorithm="maxima")
```

output

```
1/6*a*((-c^4*x^4 + 1)^(3/2)/c^8 - 3*sqrt(-c^4*x^4 + 1)/c^8) + 1/6*b*((c^8*x^8 + c^4*x^4 - 2)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)/(sqrt(c^2*x^2 + 1))*sqrt(c*x + 1)*sqrt(-c*x + 1)*c^8) - 6*integrate(1/6*(6*c^6*x^13*log(c) + 12*c^6*x^13*log(sqrt(x)) + (12*c^6*x^13*log(sqrt(x)) + (c^6*x^6*(6*log(c) + 1) + c^4*x^4 + 2*c^2*x^2 + 2)*x^7)*e^(1/2*log(c*x + 1) + 1/2*log(-c*x + 1)))/((c^6*x^6*e^(log(c*x + 1) + log(-c*x + 1)) + c^6*x^6*e^(1/2*log(c*x + 1) + 1/2*log(-c*x + 1)))*sqrt(c^2*x^2 + 1)), x)
```

**Giac [F(-2)]**

Exception generated.

$$\int \frac{x^7(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^7*(a+b*arcsech(c*x))/(-c^4*x^4+1)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^7(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx = \int \frac{x^7(a + b \operatorname{acosh}(\frac{1}{cx}))}{\sqrt{1 - c^4 x^4}} dx$$

input `int((x^7*(a + b*acosh(1/(c*x))))/(1 - c^4*x^4)^(1/2),x)`

output `int((x^7*(a + b*acosh(1/(c*x))))/(1 - c^4*x^4)^(1/2), x)`

**Reduce [F]**

$$\begin{aligned} & \int \frac{x^7(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx \\ &= \frac{-\sqrt{-c^4 x^4 + 1} a c^4 x^4 - 2\sqrt{-c^4 x^4 + 1} a - 6 \left( \int \frac{\sqrt{-c^4 x^4 + 1} \operatorname{asech}(cx) x^7}{c^4 x^4 - 1} dx \right) b c^8}{6c^8} \end{aligned}$$

input `int(x^7*(a+b*asech(c*x))/(-c^4*x^4+1)^(1/2),x)`

output

```
( - sqrt( - c**4*x**4 + 1)*a*c**4*x**4 - 2*sqrt( - c**4*x**4 + 1)*a - 6*in  
t((sqrt( - c**4*x**4 + 1)*asech(c*x)*x**7)/(c**4*x**4 - 1),x)*b*c**8)/(6*c  
**8)
```

**3.187**  $\int \frac{x^3 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx$

Optimal result	1459
Mathematica [A] (verified)	1459
Rubi [A] (verified)	1460
Maple [F]	1462
Fricas [B] (verification not implemented)	1463
Sympy [F]	1463
Maxima [F]	1464
Giac [F]	1464
Mupad [F(-1)]	1464
Reduce [F]	1465

**Optimal result**

Integrand size = 26, antiderivative size = 159

$$\int \frac{x^3 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx = -\frac{b\sqrt{1 - c^2 x^2} \sqrt{1 + c^2 x^2}}{2c^5 \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} x} - \frac{\sqrt{1 - c^4 x^4} (a + b \operatorname{sech}^{-1}(cx))}{2c^4} + \frac{b\sqrt{1 - c^2 x^2} \operatorname{arctanh}(\sqrt{1 + c^2 x^2})}{2c^5 \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}} x}$$

output

```
-1/2*b*(-c^2*x^2+1)^(1/2)*(c^2*x^2+1)^(1/2)/c^5/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)/x-1/2*(-c^4*x^4+1)^(1/2)*(a+b*arcsech(c*x))/c^4+1/2*b*(-c^2*x^2+1)^(1/2)*arctanh((c^2*x^2+1)^(1/2))/c^5/(-1+1/c/x)^(1/2)/(1+1/c/x)^(1/2)/x
```

**Mathematica [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.88

$$\int \frac{x^3 (a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx = \frac{a\sqrt{1 - c^4 x^4} + \frac{b\sqrt{1 - c^4 x^4}}{\sqrt{\frac{1 - cx}{1 + cx}}(1 + cx)} + b\sqrt{1 - c^4 x^4} \operatorname{sech}^{-1}(cx) - b \log(x(1 - cx)) + b \log\left(1 - cx - \sqrt{\frac{1 - cx}{1 + cx}} \sqrt{1 - c^4 x^4}\right)}{2c^4}$$

input `Integrate[(x^3*(a + b*ArcSech[c*x]))/Sqrt[1 - c^4*x^4],x]`

output `-1/2*(a*Sqrt[1 - c^4*x^4] + (b*Sqrt[1 - c^4*x^4])/(Sqrt[(1 - c*x)/(1 + c*x)])*(1 + c*x)) + b*Sqrt[1 - c^4*x^4]*ArcSech[c*x] - b*Log[x*(1 - c*x)] + b*Log[1 - c*x - Sqrt[(1 - c*x)/(1 + c*x)]]*Sqrt[1 - c^4*x^4])/c^4`

### Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.71, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.269$ , Rules used = {6863, 27, 1388, 243, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4x^4}} dx \\
 & \quad \downarrow \text{6863} \\
 & \frac{b\sqrt{1 - c^2x^2} \int -\frac{\sqrt{1 - c^4x^4}}{2c^4x\sqrt{1 - c^2x^2}} dx}{cx\sqrt{\frac{1}{cx} - 1}\sqrt{\frac{1}{cx} + 1}} - \frac{\sqrt{1 - c^4x^4}(a + b\operatorname{sech}^{-1}(cx))}{2c^4} \\
 & \quad \downarrow \text{27} \\
 & \frac{b\sqrt{1 - c^2x^2} \int \frac{\sqrt{1 - c^4x^4}}{x\sqrt{1 - c^2x^2}} dx}{2c^5x\sqrt{\frac{1}{cx} - 1}\sqrt{\frac{1}{cx} + 1}} - \frac{\sqrt{1 - c^4x^4}(a + b\operatorname{sech}^{-1}(cx))}{2c^4} \\
 & \quad \downarrow \text{1388} \\
 & \frac{b\sqrt{1 - c^2x^2} \int \frac{\sqrt{c^2x^2 + 1}}{x} dx}{2c^5x\sqrt{\frac{1}{cx} - 1}\sqrt{\frac{1}{cx} + 1}} - \frac{\sqrt{1 - c^4x^4}(a + b\operatorname{sech}^{-1}(cx))}{2c^4} \\
 & \quad \downarrow \text{243} \\
 & \frac{b\sqrt{1 - c^2x^2} \int \frac{\sqrt{c^2x^2 + 1}}{x^2} dx^2}{4c^5x\sqrt{\frac{1}{cx} - 1}\sqrt{\frac{1}{cx} + 1}} - \frac{\sqrt{1 - c^4x^4}(a + b\operatorname{sech}^{-1}(cx))}{2c^4} \\
 & \quad \downarrow \text{60}
 \end{aligned}$$

$$\begin{aligned}
& \frac{b\sqrt{1-c^2x^2}\left(\int \frac{1}{x^2\sqrt{c^2x^2+1}}dx^2 + 2\sqrt{c^2x^2+1}\right)}{4c^5x\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}} - \frac{\sqrt{1-c^4x^4}(a+b\operatorname{sech}^{-1}(cx))}{2c^4} \\
& \quad \downarrow 73 \\
& \frac{b\sqrt{1-c^2x^2}\left(\frac{2\int \frac{x^4-1}{c^2-\frac{1}{c^2}}d\sqrt{c^2x^2+1}}{c^2-\frac{1}{c^2}} + 2\sqrt{c^2x^2+1}\right)}{4c^5x\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}} - \frac{\sqrt{1-c^4x^4}(a+b\operatorname{sech}^{-1}(cx))}{2c^4} \\
& \quad \downarrow 221 \\
& \frac{\sqrt{1-c^4x^4}(a+b\operatorname{sech}^{-1}(cx))}{2c^4} - \frac{b\sqrt{1-c^2x^2}\left(2\sqrt{c^2x^2+1} - 2\operatorname{arctanh}\left(\sqrt{c^2x^2+1}\right)\right)}{4c^5x\sqrt{\frac{1}{cx}-1}\sqrt{\frac{1}{cx}+1}}
\end{aligned}$$

input `Int[(x^3*(a + b*ArcSech[c*x]))/Sqrt[1 - c^4*x^4],x]`

output `-1/2*(Sqrt[1 - c^4*x^4]*(a + b*ArcSech[c*x]))/c^4 - (b*Sqrt[1 - c^2*x^2]*(2*Sqrt[1 + c^2*x^2] - 2*ArcTanh[Sqrt[1 + c^2*x^2]]))/(4*c^5*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]*x)`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[  
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +  
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt  
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL  
 inearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x  
 /Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[In  
 t[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I  
 ntegerQ[(m - 1)/2]`

rule 1388 `Int[(u_.)*((a_) + (c_.)*(x_)^(n2_.))^(p_.)*((d_) + (e_.)*(x_)^(n_))^(q_.),  
 x_Symbol] := Int[u*(d + e*x^n)^(p + q)*(a/d + (c/e)*x^n)^p, x] /; FreeQ[{a,  
 c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[c*d^2 + a*e^2, 0] && (Integer  
 Q[p] || (GtQ[a, 0] && GtQ[d, 0]))`

rule 6863 `Int[((a_.) + ArcSech[(c_.)*(x_)])*(b_.)*(u_), x_Symbol] := With[{v = IntHid  
 e[u, x]}, Simp[(a + b*ArcSech[c*x]) v, x] + Simp[b*(Sqrt[1 - c^2*x^2]/(c*  
 x*Sqrt[-1 + 1/(c*x)]*Sqrt[1 + 1/(c*x)]) Int[SimplifyIntegrand[v/(x*Sqrt[  
 1 - c^2*x^2]), x], x], x] /; InverseFunctionFreeQ[v, x] /; FreeQ[{a, b, c},  
 x]`

## Maple [F]

$$\int \frac{x^3(a + b \operatorname{arcsech}(cx))}{\sqrt{-c^4x^4 + 1}} dx$$

input `int(x^3*(a+b*arcsech(c*x))/(-c^4*x^4+1)^(1/2),x)`

output `int(x^3*(a+b*arcsech(c*x))/(-c^4*x^4+1)^(1/2),x)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 279 vs.  $2(135) = 270$ .

Time = 0.12 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.75

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx$$

$$= \frac{2\sqrt{-c^4 x^4 + 1} b c x \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} - 2\sqrt{-c^4 x^4 + 1} (b c^2 x^2 - b) \log\left(\frac{c x \sqrt{-\frac{c^2 x^2 - 1}{c^2 x^2}} + 1}{c x}\right) - (b c^2 x^2 - b) \log\left(\frac{c^2 x^2 + \sqrt{-c^4 x^4 + 1}}{c^2 x^2 - 1}\right)}{4(c^6 x^2 - c^4)}$$

input `integrate(x^3*(a+b*arcsech(c*x))/(-c^4*x^4+1)^(1/2),x, algorithm="fricas")`

output `1/4*(2*sqrt(-c^4*x^4 + 1)*b*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 2*sqrt(-c^4*x^4 + 1)*(b*c^2*x^2 - b)*log((c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) + 1)/(c*x)) - (b*c^2*x^2 - b)*log((c^2*x^2 + sqrt(-c^4*x^4 + 1)*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/(c^2*x^2 - 1)) + (b*c^2*x^2 - b)*log(-(c^2*x^2 - sqrt(-c^4*x^4 + 1)*c*x*sqrt(-(c^2*x^2 - 1)/(c^2*x^2)) - 1)/(c^2*x^2 - 1)) - 2*sqrt(-c^4*x^4 + 1)*(a*c^2*x^2 - a))/(c^6*x^2 - c^4)`

**Sympy [F]**

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx = \int \frac{x^3(a + b \operatorname{asech}(cx))}{\sqrt{-(cx - 1)(cx + 1)(c^2 x^2 + 1)}} dx$$

input `integrate(x**3*(a+b*asech(c*x))/(-c**4*x**4+1)**(1/2),x)`

output `Integral(x**3*(a + b*asech(c*x))/sqrt(-(c*x - 1)*(c*x + 1)*(c**2*x**2 + 1)), x)`



**Maxima [F]**

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4x^4}} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^3}{\sqrt{-c^4x^4 + 1}} dx$$

input `integrate(x^3*(a+b*arcsech(c*x))/(-c^4*x^4+1)^(1/2),x, algorithm="maxima")`

output `1/2*b*((c^4*x^4 - 1)*log(sqrt(c*x + 1)*sqrt(-c*x + 1) + 1)/(sqrt(c^2*x^2 + 1)*sqrt(c*x + 1)*sqrt(-c*x + 1)*c^4) - 2*integrate(1/2*(2*c^2*x^5*log(c) + 4*c^2*x^5*log(sqrt(x)) + (4*c^2*x^5*log(sqrt(x)) + (c^2*x^2*(2*log(c) + 1) + 1)*x^3)*e^(1/2*log(c*x + 1) + 1/2*log(-c*x + 1)))/((c^2*x^2*e^(log(c*x + 1) + log(-c*x + 1)) + c^2*x^2*e^(1/2*log(c*x + 1) + 1/2*log(-c*x + 1)))*sqrt(c^2*x^2 + 1)), x) - 1/2*sqrt(-c^4*x^4 + 1)*a/c^4`

**Giac [F]**

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4x^4}} dx = \int \frac{(b \operatorname{arsech}(cx) + a)x^3}{\sqrt{-c^4x^4 + 1}} dx$$

input `integrate(x^3*(a+b*arcsech(c*x))/(-c^4*x^4+1)^(1/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)*x^3/sqrt(-c^4*x^4 + 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + b\operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4x^4}} dx = \int \frac{x^3(a + b \operatorname{acosh}(\frac{1}{cx}))}{\sqrt{1 - c^4x^4}} dx$$

input `int((x^3*(a + b*acosh(1/(c*x))))/(1 - c^4*x^4)^(1/2),x)`

output `int((x^3*(a + b*acosh(1/(c*x))))/(1 - c^4*x^4)^(1/2), x)`

**Reduce [F]**

$$\int \frac{x^3(a + b \operatorname{sech}^{-1}(cx))}{\sqrt{1 - c^4 x^4}} dx = \frac{-\sqrt{-c^4 x^4 + 1} a - 2 \left( \int \frac{\sqrt{-c^4 x^4 + 1} \operatorname{asech}(cx) x^3}{c^4 x^4 - 1} dx \right) b c^4}{2c^4}$$

input `int(x^3*(a+b*asech(c*x))/(-c^4*x^4+1)^(1/2),x)`

output `(-sqrt(-c**4*x**4+1)*a-2*int((sqrt(-c**4*x**4+1)*asech(c*x)*x**3)/(c**4*x**4-1),x)*b*c**4)/(2*c**4)`

$$3.188 \quad \int \frac{a+b\operatorname{sech}^{-1}(cx)}{x\sqrt{1-c^4x^4}} dx$$

Optimal result	1466
Mathematica [N/A]	1466
Rubi [N/A]	1467
Maple [N/A]	1467
Fricas [N/A]	1468
Sympy [N/A]	1468
Maxima [N/A]	1468
Giac [N/A]	1469
Mupad [N/A]	1469
Reduce [N/A]	1470

### Optimal result

Integrand size = 26, antiderivative size = 26

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x\sqrt{1 - c^4x^4}} dx = \operatorname{Int}\left(\frac{a + b\operatorname{sech}^{-1}(cx)}{x\sqrt{1 - c^4x^4}}, x\right)$$

output `Defer(Int)((a+b*arcsech(c*x))/x/(-c^4*x^4+1)^(1/2),x)`

### Mathematica [N/A]

Not integrable

Time = 0.50 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x\sqrt{1 - c^4x^4}} dx = \int \frac{a + b\operatorname{sech}^{-1}(cx)}{x\sqrt{1 - c^4x^4}} dx$$

input `Integrate[(a + b*ArcSech[c*x])/(x*Sqrt[1 - c^4*x^4]),x]`

output `Integrate[(a + b*ArcSech[c*x])/(x*Sqrt[1 - c^4*x^4]), x]`

**Rubi [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x \sqrt{1 - c^4 x^4}} dx$$

↓ 6865

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x \sqrt{1 - c^4 x^4}} dx$$

input `Int[(a + b*ArcSech[c*x])/(x*Sqrt[1 - c^4*x^4]),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \frac{a + b \operatorname{arcsech}(cx)}{x \sqrt{-c^4 x^4 + 1}} dx$$

input `int((a+b*arcsech(c*x))/x/(-c^4*x^4+1)^(1/2),x)`

output `int((a+b*arcsech(c*x))/x/(-c^4*x^4+1)^(1/2),x)`

**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.42

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x\sqrt{1 - c^4x^4}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{\sqrt{-c^4x^4 + 1}x} dx$$

input `integrate((a+b*arcsech(c*x))/x/(-c^4*x^4+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^4*x^4 + 1)*(b*arcsech(c*x) + a)/(c^4*x^5 - x), x)`

**Sympy [N/A]**

Not integrable

Time = 3.09 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.31

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x\sqrt{1 - c^4x^4}} dx = \int \frac{a + b \operatorname{asech}(cx)}{x\sqrt{-(cx - 1)(cx + 1)(c^2x^2 + 1)}} dx$$

input `integrate((a+b*asech(c*x))/x/(-c**4*x**4+1)**(1/2),x)`

output `Integral((a + b*asech(c*x))/(x*sqrt(-(c*x - 1)*(c*x + 1)*(c**2*x**2 + 1))), x)`

**Maxima [N/A]**

Not integrable

Time = 0.57 (sec) , antiderivative size = 100, normalized size of antiderivative = 3.85

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x\sqrt{1 - c^4x^4}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{\sqrt{-c^4x^4 + 1}x} dx$$

input `integrate((a+b*arcsech(c*x))/x/(-c^4*x^4+1)^(1/2),x, algorithm="maxima")`

output `-1/4*a*(log(sqrt(-c^4*x^4 + 1) + 1) - log(sqrt(-c^4*x^4 + 1) - 1)) + b*integrate(log(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(sqrt(-(c^2*x^2 + 1)*(c*x + 1)*(c*x - 1))*x), x)`

### Giac [N/A]

Not integrable

Time = 0.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x \sqrt{1 - c^4 x^4}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{\sqrt{-c^4 x^4 + 1} x} dx$$

input `integrate((a+b*arcsech(c*x))/x/(-c^4*x^4+1)^(1/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/(sqrt(-c^4*x^4 + 1)*x), x)`

### Mupad [N/A]

Not integrable

Time = 4.72 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.15

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x \sqrt{1 - c^4 x^4}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x \sqrt{1 - c^4 x^4}} dx$$

input `int((a + b*acosh(1/(c*x)))/(x*(1 - c^4*x^4)^(1/2)),x)`

output `int((a + b*acosh(1/(c*x)))/(x*(1 - c^4*x^4)^(1/2)), x)`

**Reduce [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.92

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x \sqrt{1 - c^4 x^4}} dx = - \left( \int \frac{\sqrt{-c^4 x^4 + 1} \operatorname{asech}(cx)}{c^4 x^5 - x} dx \right) b + \frac{\log \left( \tan \left( \frac{\operatorname{asin}(c^2 x^2)}{2} \right) \right) a}{2}$$

input

```
int((a+b*asech(c*x))/x/(-c^4*x^4+1)^(1/2),x)
```

output

```
( - 2*int((sqrt( - c**4*x**4 + 1)*asech(c*x))/(c**4*x**5 - x),x)*b + log(tan(asin(c**2*x**2)/2))*a)/2
```

$$3.189 \quad \int \frac{a+b\operatorname{sech}^{-1}(cx)}{x^5\sqrt{1-c^4x^4}} dx$$

Optimal result	1471
Mathematica [N/A]	1471
Rubi [N/A]	1472
Maple [N/A]	1472
Fricas [N/A]	1473
Sympy [N/A]	1473
Maxima [N/A]	1473
Giac [N/A]	1474
Mupad [N/A]	1474
Reduce [N/A]	1475

### Optimal result

Integrand size = 26, antiderivative size = 26

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^5\sqrt{1-c^4x^4}} dx = \operatorname{Int}\left(\frac{a + b\operatorname{sech}^{-1}(cx)}{x^5\sqrt{1-c^4x^4}}, x\right)$$

output `Defer(Int)((a+b*arcsech(c*x))/x^5/(-c^4*x^4+1)^(1/2),x)`

### Mathematica [N/A]

Not integrable

Time = 7.13 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

$$\int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^5\sqrt{1-c^4x^4}} dx = \int \frac{a + b\operatorname{sech}^{-1}(cx)}{x^5\sqrt{1-c^4x^4}} dx$$

input `Integrate[(a + b*ArcSech[c*x])/(x^5*Sqrt[1 - c^4*x^4]),x]`

output `Integrate[(a + b*ArcSech[c*x])/(x^5*Sqrt[1 - c^4*x^4]), x]`



**Rubi [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5 \sqrt{1 - c^4 x^4}} dx$$

↓ 6865

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5 \sqrt{1 - c^4 x^4}} dx$$

input `Int[(a + b*ArcSech[c*x])/(x^5*Sqrt[1 - c^4*x^4]),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \frac{a + b \operatorname{arcsech}(cx)}{x^5 \sqrt{-c^4 x^4 + 1}} dx$$

input `int((a+b*arcsech(c*x))/x^5/(-c^4*x^4+1)^(1/2),x)`

output `int((a+b*arcsech(c*x))/x^5/(-c^4*x^4+1)^(1/2),x)`

**Fricas [N/A]**

Not integrable

Time = 0.11 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.50

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5 \sqrt{1 - c^4 x^4}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{\sqrt{-c^4 x^4 + 1} x^5} dx$$

input `integrate((a+b*arcsech(c*x))/x^5/(-c^4*x^4+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^4*x^4 + 1)*(b*arcsech(c*x) + a)/(c^4*x^9 - x^5), x)`

**Sympy [N/A]**

Not integrable

Time = 28.59 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.38

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5 \sqrt{1 - c^4 x^4}} dx = \int \frac{a + b \operatorname{asech}(cx)}{x^5 \sqrt{-(cx - 1)(cx + 1)(c^2 x^2 + 1)}} dx$$

input `integrate((a+b*asech(c*x))/x**5/(-c**4*x**4+1)**(1/2),x)`

output `Integral((a + b*asech(c*x))/(x**5*sqrt(-(c*x - 1)*(c*x + 1)*(c**2*x**2 + 1))), x)`

**Maxima [N/A]**

Not integrable

Time = 0.63 (sec) , antiderivative size = 124, normalized size of antiderivative = 4.77

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5 \sqrt{1 - c^4 x^4}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{\sqrt{-c^4 x^4 + 1} x^5} dx$$

input `integrate((a+b*arcsech(c*x))/x^5/(-c^4*x^4+1)^(1/2),x, algorithm="maxima")`

output `-1/8*(c^4*log(sqrt(-c^4*x^4 + 1) + 1) - c^4*log(sqrt(-c^4*x^4 + 1) - 1) + 2*sqrt(-c^4*x^4 + 1)/x^4)*a + b*integrate(log(sqrt(1/(c*x) + 1))*sqrt(1/(c*x) - 1) + 1/(c*x))/(sqrt(-c^2*x^2 + 1)*(c*x + 1)*(c*x - 1))*x^5, x)`

### Giac [N/A]

Not integrable

Time = 0.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5 \sqrt{1 - c^4 x^4}} dx = \int \frac{b \operatorname{arsech}(cx) + a}{\sqrt{-c^4 x^4 + 1} x^5} dx$$

input `integrate((a+b*arcsech(c*x))/x^5/(-c^4*x^4+1)^(1/2),x, algorithm="giac")`

output `integrate((b*arcsech(c*x) + a)/(sqrt(-c^4*x^4 + 1)*x^5), x)`

### Mupad [N/A]

Not integrable

Time = 4.23 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.15

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5 \sqrt{1 - c^4 x^4}} dx = \int \frac{a + b \operatorname{acosh}\left(\frac{1}{cx}\right)}{x^5 \sqrt{1 - c^4 x^4}} dx$$

input `int((a + b*acosh(1/(c*x)))/(x^5*(1 - c^4*x^4)^(1/2)),x)`

output `int((a + b*acosh(1/(c*x)))/(x^5*(1 - c^4*x^4)^(1/2)), x)`

**Reduce [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 79, normalized size of antiderivative = 3.04

$$\int \frac{a + b \operatorname{sech}^{-1}(cx)}{x^5 \sqrt{1 - c^4 x^4}} dx$$

$$= \frac{-\sqrt{-c^4 x^4 + 1} a - 4 \left( \int \frac{\sqrt{-c^4 x^4 + 1} \operatorname{asech}(cx)}{c^4 x^9 - x^5} dx \right) b x^4 + \log \left( \tan \left( \frac{\operatorname{asin}(c^2 x^2)}{2} \right) \right) a c^4 x^4}{4x^4}$$

input

```
int((a+b*asech(c*x))/x^5/(-c^4*x^4+1)^(1/2),x)
```

output

```
( - sqrt( - c**4*x**4 + 1)*a - 4*int((sqrt( - c**4*x**4 + 1)*asech(c*x))/(
c**4*x**9 - x**5),x)*b*x**4 + log(tan(asin(c**2*x**2)/2))*a*c**4*x**4)/(4*
x**4)
```

# CHAPTER 4

## APPENDIX

4.1	Listing of Grading functions . . . . .	1476
4.2	Links to plain text integration problems used in this report for each CAS .	1494

### 4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

#### Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal."}
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal."}
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order of result is higher than in optimal."}
  ]
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]==Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]==Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]==Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]==Plus || Head[expn]==Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```





## Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022  add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result    := ExpnType(result);
      ExpnType_optimal   := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#     is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

## Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```



```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

## SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

## 4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file