

Computer Algebra Independent Integration Tests

Summer 2024

7-Inverse-hyperbolic-functions/7.6-Inverse-hyperbolic-
cosecant/348-7.6

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 10:36am

Contents

1	Introduction	3
1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	Performance based on number of rules Rubi used	11
1.5	Performance based on number of steps Rubi used	12
1.6	Solved integrals histogram based on leaf size of result	13
1.7	Solved integrals histogram based on CPU time used	14
1.8	Leaf size vs. CPU time used	15
1.9	list of integrals with no known antiderivative	16
1.10	List of integrals solved by CAS but has no known antiderivative	16
1.11	list of integrals solved by CAS but failed verification	16
1.12	Timing	17
1.13	Verification	17
1.14	Important notes about some of the results	18
1.15	Current tree layout of integration tests	21
1.16	Design of the test system	22
2	detailed summary tables of results	23
2.1	List of integrals sorted by grade for each CAS	24
2.2	Detailed conclusion table per each integral for all CAS systems	28
2.3	Detailed conclusion table specific for Rubi results	36
3	Listing of integrals	38
3.1	$\int x^3 \operatorname{csch}^{-1}(a + bx) dx$	40
3.2	$\int x^2 \operatorname{csch}^{-1}(a + bx) dx$	47
3.3	$\int x \operatorname{csch}^{-1}(a + bx) dx$	53
3.4	$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x} dx$	60
3.5	$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^2} dx$	70
3.6	$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^3} dx$	77

3.7	$\int (e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$	87
3.8	$\int (e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$	97
3.9	$\int (e + fx) (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$	105
3.10	$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$	112
3.11	$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{e + fx} dx$	119
3.12	$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^2} dx$	130
3.13	$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^3} dx$	138
3.14	$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx$	146
3.15	$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx$	152
3.16	$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx$	157
3.17	$\int \operatorname{csch}^{-1}(\sqrt{x}) dx$	162
3.18	$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx$	167
3.19	$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx$	174
3.20	$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx$	180
3.21	$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx$	186
3.22	$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx$	192
3.23	$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx$	197
3.24	$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx$	204
3.25	$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx$	211
3.26	$\int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{a}{b} + dx} dx$	218
3.27	$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx$	225
3.28	$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx$	231
4	Appendix	237
4.1	Listing of Grading functions	237
4.2	Links to plain text integration problems used in this report for each CAS255	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	Performance based on number of rules Rubi used	11
1.5	Performance based on number of steps Rubi used	12
1.6	Solved integrals histogram based on leaf size of result	13
1.7	Solved integrals histogram based on CPU time used	14
1.8	Leaf size vs. CPU time used	15
1.9	list of integrals with no known antiderivative	16
1.10	List of integrals solved by CAS but has no known antiderivative	16
1.11	list of integrals solved by CAS but failed verification	16
1.12	Timing	17
1.13	Verification	17
1.14	Important notes about some of the results	18
1.15	Current tree layout of integration tests	21
1.16	Design of the test system	22

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [28]. This is test number [348].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (28)	0.00 (0)
Mathematica	96.43 (27)	3.57 (1)
Fricas	53.57 (15)	46.43 (13)
Maple	50.00 (14)	50.00 (14)
Maxima	35.71 (10)	64.29 (18)
Mupad	17.86 (5)	82.14 (23)
Reduce	3.57 (1)	96.43 (27)
Sympy	3.57 (1)	96.43 (27)
Giac	0.00 (0)	100.00 (28)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	71.429	0.000	28.571	0.000
Mathematica	53.571	21.429	21.429	3.571
Maple	42.857	7.143	0.000	50.000
Maxima	35.714	0.000	0.000	64.286
Fricas	28.571	25.000	0.000	46.429
Sympy	3.571	0.000	0.000	96.429
Giac	0.000	0.000	0.000	100.000
Mupad	0.000	17.857	0.000	82.143
Reduce	0.000	3.571	0.000	96.429

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	1	100.00	0.00	0.00
Fricas	13	84.62	0.00	15.38
Maple	14	100.00	0.00	0.00
Maxima	18	100.00	0.00	0.00
Mupad	23	0.00	100.00	0.00
Reduce	27	100.00	0.00	0.00
Sympy	27	92.59	7.41	0.00
Giac	28	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.03
Sympy	0.07
Fricas	0.11
Reduce	0.15
Maple	0.23
Rubi	0.63
Mathematica	1.89
Mupad	3.00
Giac	-nan(ind)

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Reduce	13.00	0.81	13.00	0.81
Sympy	14.00	0.88	14.00	0.88
Mupad	29.40	0.75	33.00	0.87
Maxima	56.20	0.87	57.50	1.00
Maple	93.00	1.08	54.50	0.90
Rubi	172.61	1.03	78.00	1.00
Fricas	178.73	2.49	58.00	1.38
Mathematica	564.44	1.79	95.00	1.17
Giac	-nan(ind)	-nan(ind)	nan	nan

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {10, 18, 23, 24, 25, 26, 27, 28}

Mathematica {7, 8, 12, 13}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	24
2.2	Detailed conclusion table per each integral for all CAS systems	28
2.3	Detailed conclusion table specific for Rubi results	36

2.1 List of integrals sorted by grade for each CAS

Rubi	24
Mma	24
Maple	25
Fricas	25
Maxima	25
Giac	26
Mupad	26
Sympy	26
Reduce	27

Rubi

A grade { 1, 2, 3, 5, 6, 7, 8, 9, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 27, 28 }

B grade { }

C grade { 4, 10, 11, 18, 23, 24, 25, 26 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 6, 14, 15, 16, 17, 18, 19, 20, 21, 22, 24, 26 }

B grade { 5, 9, 10, 25, 27, 28 }

C grade { 4, 7, 8, 12, 13, 23 }

F normal fail { 11 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 14, 15, 16, 17, 19, 20, 21, 22, 27 }

B grade { 5, 6 }

C grade { }

F normal fail { 4, 7, 8, 9, 10, 11, 12, 13, 18, 23, 24, 25, 26, 28 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 14, 15, 16, 17, 19, 20, 21, 22 }

B grade { 1, 2, 3, 5, 6, 27, 28 }

C grade { }

F normal fail { 4, 7, 8, 9, 10, 11, 12, 13, 18, 24, 26 }

F(-1) timedout fail { }

F(-2) exception fail { 23, 25 }

Maxima

A grade { 14, 15, 16, 17, 19, 20, 21, 22, 27, 28 }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 18, 23, 24, 25, 26 }

F(-1) timedout fail { }

F(-2) exception fail { }

Giac

A grade { }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 17, 19, 22, 27, 28 }

C grade { }

F normal fail { }

F(-1) timedout fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 18, 20, 21, 23, 24, 25, 26 }

F(-2) exception fail { }

Sympy

A grade { 22 }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 23, 24, 25, 26 }

F(-1) timedout fail { 27, 28 }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 22 }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 23, 24, 25,
26, 27, 28 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	147	150	149	227	0	325	0	0	12	0
N.S.	1	1.02	1.01	1.54	0.00	2.21	0.00	0.00	0.08	0.00
time (sec)	N/A	0.518	0.177	0.323	0.000	0.126	0.000	0.000	0.179	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	110	129	170	0	306	0	0	12	0
N.S.	1	1.00	1.17	1.55	0.00	2.78	0.00	0.00	0.11	0.00
time (sec)	N/A	0.381	0.109	0.241	0.000	0.148	0.000	0.000	0.181	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	74	110	97	0	285	0	0	10	0
N.S.	1	0.99	1.47	1.29	0.00	3.80	0.00	0.00	0.13	0.00
time (sec)	N/A	0.425	0.057	0.240	0.000	0.104	0.000	0.000	0.165	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	162	225	427	0	0	0	0	0	12	0
N.S.	1	1.39	2.64	0.00	0.00	0.00	0.00	0.00	0.07	0.00
time (sec)	N/A	1.015	0.302	0.000	0.000	0.000	0.000	0.000	0.160	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	72	141	127	0	343	0	0	12	0
N.S.	1	1.14	2.24	2.02	0.00	5.44	0.00	0.00	0.19	0.00
time (sec)	N/A	0.424	0.101	0.466	0.000	0.119	0.000	0.000	0.157	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	114	143	220	318	0	461	0	0	12	0
N.S.	1	1.25	1.93	2.79	0.00	4.04	0.00	0.00	0.11	0.00
time (sec)	N/A	0.745	0.297	0.487	0.000	0.129	0.000	0.000	0.172	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	501	476	1429	0	0	0	0	0	197	0
N.S.	1	0.95	2.85	0.00	0.00	0.00	0.00	0.00	0.39	0.00
time (sec)	N/A	1.066	9.980	0.000	0.000	0.000	0.000	0.000	0.357	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	351	336	864	0	0	0	0	0	136	0
N.S.	1	0.96	2.46	0.00	0.00	0.00	0.00	0.00	0.39	0.00
time (sec)	N/A	0.772	7.865	0.000	0.000	0.000	0.000	0.000	0.303	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	194	199	427	0	0	0	0	0	76	0
N.S.	1	1.03	2.20	0.00	0.00	0.00	0.00	0.00	0.39	0.00
time (sec)	N/A	0.588	2.882	0.000	0.000	0.000	0.000	0.000	0.238	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	B	F	F	F	F	F	F	F(-1)
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	85	84	176	0	0	0	0	0	32	0
N.S.	1	0.99	2.07	0.00	0.00	0.00	0.00	0.00	0.38	0.00
time (sec)	N/A	0.470	0.199	0.000	0.000	0.000	0.000	0.000	0.195	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	475	568	0	0	0	0	0	0	59	0
N.S.	1	1.20	0.00	0.00	0.00	0.00	0.00	0.00	0.12	0.00
time (sec)	N/A	2.599	0.000	0.000	0.000	0.000	0.000	0.000	0.226	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	448	461	1874	0	0	0	0	0	157	0
N.S.	1	1.03	4.18	0.00	0.00	0.00	0.00	0.00	0.35	0.00
time (sec)	N/A	1.368	12.514	0.000	0.000	0.000	0.000	0.000	0.226	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	1024	1006	8350	0	0	0	0	0	320	0
N.S.	1	0.98	8.15	0.00	0.00	0.00	0.00	0.00	0.31	0.00
time (sec)	N/A	2.455	13.272	0.000	0.000	0.000	0.000	0.000	0.480	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	114	79	47	40	58	55	0	0	9	0
N.S.	1	0.69	0.41	0.35	0.51	0.48	0.00	0.00	0.08	0.00
time (sec)	N/A	0.232	0.024	0.207	0.023	0.090	0.000	0.000	0.347	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	68	42	35	46	50	0	0	9	0
N.S.	1	0.76	0.47	0.39	0.52	0.56	0.00	0.00	0.10	0.00
time (sec)	N/A	0.215	0.019	0.152	0.028	0.083	0.000	0.000	0.306	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	55	35	28	34	43	0	0	7	0
N.S.	1	0.86	0.55	0.44	0.53	0.67	0.00	0.00	0.11	0.00
time (sec)	N/A	0.205	0.018	0.161	0.030	0.109	0.000	0.000	0.274	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	24	21	18	36	0	0	5	18
N.S.	1	1.00	0.77	0.68	0.58	1.16	0.00	0.00	0.16	0.58
time (sec)	N/A	0.171	2.167	0.159	0.031	0.107	0.000	0.000	0.224	3.822

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	F	F	F	F	F	F	F(-1)
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	46	58	46	0	0	0	0	0	9	0
N.S.	1	1.26	1.00	0.00	0.00	0.00	0.00	0.00	0.20	0.00
time (sec)	N/A	0.463	0.018	0.000	0.000	0.000	0.000	0.000	0.206	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	54	42	46	65	44	0	0	9	33
N.S.	1	0.86	0.67	0.73	1.03	0.70	0.00	0.00	0.14	0.52
time (sec)	N/A	0.206	0.018	0.164	0.030	0.092	0.000	0.000	0.189	3.630

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	77	47	57	92	53	0	0	9	0
N.S.	1	0.86	0.52	0.63	1.02	0.59	0.00	0.00	0.10	0.00
time (sec)	N/A	0.215	0.025	0.160	0.024	0.096	0.000	0.000	0.194	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	98	52	67	116	58	0	0	9	0
N.S.	1	0.85	0.45	0.58	1.01	0.50	0.00	0.00	0.08	0.00
time (sec)	N/A	0.228	0.030	0.158	0.026	0.095	0.000	0.000	0.189	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	17	16	22	14	0	13	14
N.S.	1	1.00	1.12	1.06	1.00	1.38	0.88	0.00	0.81	0.88
time (sec)	N/A	0.191	0.003	0.155	0.027	0.089	0.075	0.000	0.155	0.072

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	C	F	F	F(-2)	F	F	F	F(-1)
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	61	77	64	0	0	0	0	0	12	0
N.S.	1	1.26	1.05	0.00	0.00	0.00	0.00	0.00	0.20	0.00
time (sec)	N/A	0.492	0.058	0.000	0.000	0.000	0.000	0.000	0.176	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	F	F	F	F	F	F	F(-1)
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	54	68	50	0	0	0	0	0	12	0
N.S.	1	1.26	0.93	0.00	0.00	0.00	0.00	0.00	0.22	0.00
time (sec)	N/A	0.497	0.020	0.000	0.000	0.000	0.000	0.000	0.150	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	B	F	F	F(-2)	F	F	F	F(-1)
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	77	97	236	0	0	0	0	0	12	0
N.S.	1	1.26	3.06	0.00	0.00	0.00	0.00	0.00	0.16	0.00
time (sec)	N/A	0.482	0.682	0.000	0.000	0.000	0.000	0.000	0.161	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	F	F	F	F	F	F	F(-1)
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	61	73	53	0	0	0	0	0	21	0
N.S.	1	1.20	0.87	0.00	0.00	0.00	0.00	0.00	0.34	0.00
time (sec)	N/A	0.535	0.048	0.000	0.000	0.000	0.000	0.000	0.163	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	B	F(-1)	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	38	95	52	57	266	0	0	14	42
N.S.	1	0.83	2.07	1.13	1.24	5.78	0.00	0.00	0.30	0.91
time (sec)	N/A	0.302	0.104	0.166	0.034	0.120	0.000	0.000	0.153	3.961

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	A	B	F(-1)	F	F	B
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	46	40	93	0	60	334	0	0	17	40
N.S.	1	0.87	2.02	0.00	1.30	7.26	0.00	0.00	0.37	0.87
time (sec)	N/A	0.323	0.126	0.000	0.023	0.123	0.000	0.000	0.189	3.510

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [3] had the largest ratio of [1.6250000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	9	8	1.02	10	0.800
2	A	6	5	1.00	10	0.500
3	A	14	13	0.99	8	1.625
4	C	17	16	1.39	10	1.600
5	A	9	8	1.14	10	0.800
6	A	17	16	1.25	10	1.600
7	A	6	5	0.95	20	0.250
8	A	6	5	0.96	20	0.250
9	A	6	5	1.03	18	0.278
10	C	9	8	0.99	12	0.667
11	C	16	15	1.20	20	0.750
12	A	6	5	1.03	20	0.250
13	A	6	5	0.98	20	0.250
14	A	4	4	0.69	10	0.400
15	A	4	4	0.76	10	0.400
16	A	4	4	0.86	8	0.500
17	A	2	2	1.00	6	0.333
18	C	11	10	1.26	10	1.000
19	A	6	5	0.86	10	0.500
20	A	7	6	0.86	10	0.600
21	A	8	7	0.85	10	0.700

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	3	3	1.00	4	0.750
23	C	11	10	1.26	10	1.000
24	C	11	10	1.26	10	1.000
25	C	11	10	1.26	10	1.000
26	C	12	11	1.20	19	0.579
27	A	7	6	0.83	12	0.500
28	A	7	6	0.87	14	0.429

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^3 \operatorname{csch}^{-1}(a + bx) dx$	40
3.2	$\int x^2 \operatorname{csch}^{-1}(a + bx) dx$	47
3.3	$\int x \operatorname{csch}^{-1}(a + bx) dx$	53
3.4	$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x} dx$	60
3.5	$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^2} dx$	70
3.6	$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^3} dx$	77
3.7	$\int (e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$	87
3.8	$\int (e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$	97
3.9	$\int (e + fx) (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$	105
3.10	$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$	112
3.11	$\int \frac{(a+b \operatorname{csch}^{-1}(c+dx))^2}{e+fx} dx$	119
3.12	$\int \frac{(a+b \operatorname{csch}^{-1}(c+dx))^2}{(e+fx)^2} dx$	130
3.13	$\int \frac{(a+b \operatorname{csch}^{-1}(c+dx))^2}{(e+fx)^3} dx$	138
3.14	$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx$	146
3.15	$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx$	152
3.16	$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx$	157
3.17	$\int \operatorname{csch}^{-1}(\sqrt{x}) dx$	162
3.18	$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx$	167
3.19	$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx$	174
3.20	$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx$	180
3.21	$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx$	186
3.22	$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx$	192

3.23	$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx$	197
3.24	$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx$	204
3.25	$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx$	211
3.26	$\int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx$	218
3.27	$\int x^3 \operatorname{csch}^{-1}(a+bx^4) dx$	225
3.28	$\int x^{-1+n} \operatorname{csch}^{-1}(a+bx^n) dx$	231

3.1 $\int x^3 \operatorname{csch}^{-1}(a + bx) dx$

Optimal result	40
Mathematica [A] (verified)	41
Rubi [A] (verified)	41
Maple [A] (verified)	44
Fricas [B] (verification not implemented)	44
Sympy [F]	45
Maxima [F]	45
Giac [F]	46
Mupad [F(-1)]	46
Reduce [F]	46

Optimal result

Integrand size = 10, antiderivative size = 147

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx = -\frac{(2 - 17a^2)(a + bx)\sqrt{1 + \frac{1}{(a+bx)^2}}}{12b^4} + \frac{x^2(a + bx)\sqrt{1 + \frac{1}{(a+bx)^2}}}{12b^2}$$

$$- \frac{a(a + bx)^2\sqrt{1 + \frac{1}{(a+bx)^2}}}{3b^4} - \frac{a^4 \operatorname{csch}^{-1}(a + bx)}{4b^4}$$

$$+ \frac{1}{4}x^4 \operatorname{csch}^{-1}(a + bx) + \frac{a(1 - 2a^2) \operatorname{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx)^2}}\right)}{2b^4}$$

output

```
-1/12*(-17*a^2+2)*(b*x+a)*(1+1/(b*x+a)^2)^(1/2)/b^4+1/12*x^2*(b*x+a)*(1+1/
(b*x+a)^2)^(1/2)/b^2-1/3*a*(b*x+a)^2*(1+1/(b*x+a)^2)^(1/2)/b^4-1/4*a^4*arc
csch(b*x+a)/b^4+1/4*x^4*arccsch(b*x+a)+1/2*a*(-2*a^2+1)*arctanh((1+1/(b*x+
a)^2)^(1/2))/b^4
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.01

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx$$

$$= \frac{\sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}}(-2a + 13a^3 - 2bx + 9a^2bx - 3ab^2x^2 + b^3x^3) + 3b^4x^4 \operatorname{csch}^{-1}(a + bx) - 3a^4 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}{12b^4}$$

input

```
Integrate[x^3*ArcCsch[a + b*x],x]
```

output

```
(Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]*(-2*a + 13*a^3 - 2*b*x + 9*a^2*b*x - 3*a*b^2*x^2 + b^3*x^3) + 3*b^4*x^4*ArcCsch[a + b*x] - 3*a^4*ArcSinh[(a + b*x)^(-1)] + 6*a*(1 - 2*a^2)*Log[(a + b*x)*(1 + Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2])])/(12*b^4)
```

Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.02, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {6876, 25, 5992, 3042, 4269, 3042, 4536, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx$$

$$\downarrow 6876$$

$$-\frac{\int b^3 x^3 (a + bx)^2 \sqrt{1 + \frac{1}{(a+bx)^2}} \operatorname{csch}^{-1}(a + bx) d\operatorname{csch}^{-1}(a + bx)}{b^4}$$

$$\downarrow 25$$

$$\frac{\int -b^3 x^3 (a + bx)^2 \sqrt{1 + \frac{1}{(a+bx)^2}} \operatorname{csch}^{-1}(a + bx) d\operatorname{csch}^{-1}(a + bx)}{b^4}$$

$$\downarrow 5992$$

$$\frac{\frac{1}{4} \int b^4 x^4 d\operatorname{csch}^{-1}(a + bx) - \frac{1}{4} b^4 x^4 \operatorname{csch}^{-1}(a + bx)}{b^4}$$

↓ 3042

$$\frac{-\frac{1}{4} b^4 x^4 \operatorname{csch}^{-1}(a + bx) + \frac{1}{4} \int (a - i \operatorname{csc}(i \operatorname{csch}^{-1}(a + bx)))^4 d\operatorname{csch}^{-1}(a + bx)}{b^4}$$

↓ 4269

$$\frac{\frac{1}{4} \left(\frac{1}{3} \int -bx(3a^3 + 8(a + bx)^2 a + (2 - 9a^2)(a + bx)) d\operatorname{csch}^{-1}(a + bx) - \frac{1}{3} b^2 x^2 (a + bx) \sqrt{\frac{1}{(a + bx)^2} + 1} \right) - \frac{1}{4} b^4 x^4 \operatorname{csch}^{-1}(a + bx)}{b^4}$$

↓ 3042

$$\frac{-\frac{1}{4} b^4 x^4 \operatorname{csch}^{-1}(a + bx) + \frac{1}{4} \left(-\frac{1}{3} b^2 x^2 (a + bx) \sqrt{\frac{1}{(a + bx)^2} + 1} + \frac{1}{3} \int (a - i \operatorname{csc}(i \operatorname{csch}^{-1}(a + bx))) (3a^3 - 8 \operatorname{csc}(i \operatorname{csch}^{-1}(a + bx))) d\operatorname{csch}^{-1}(a + bx) \right)}{b^4}$$

↓ 4536

$$\frac{\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int (6a^4 + 12(1 - 2a^2)(a + bx)a - 2(2 - 17a^2)(a + bx)^2) d\operatorname{csch}^{-1}(a + bx) + 4a \sqrt{\frac{1}{(a + bx)^2} + 1} (a + bx)^2 \right) \right)}{b^4}$$

↓ 2009

$$\frac{\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(6a^4 \operatorname{csch}^{-1}(a + bx) - 12(1 - 2a^2) a \operatorname{arctanh} \left(\sqrt{\frac{1}{(a + bx)^2} + 1} \right) + 2(2 - 17a^2)(a + bx) \sqrt{\frac{1}{(a + bx)^2} + 1} \right) + 4a \sqrt{\frac{1}{(a + bx)^2} + 1} (a + bx)^2 \right) \right)}{b^4}$$

input `Int[x^3*ArcCsch[a + b*x],x]`

output `-((-1/4*(b^4*x^4*ArcCsch[a + b*x]) + (-1/3*(b^2*x^2*(a + b*x)*Sqrt[1 + (a + b*x)^(-2)])) + (4*a*(a + b*x)^2*Sqrt[1 + (a + b*x)^(-2)] + (2*(2 - 17*a^2)*(a + b*x)*Sqrt[1 + (a + b*x)^(-2)] + 6*a^4*ArcCsch[a + b*x] - 12*a*(1 - 2*a^2)*ArcTanh[Sqrt[1 + (a + b*x)^(-2)]])/2)/3)/4)/b^4`

Defintions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 2009 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Simp}[\text{IntSum}[\text{u}, \text{x}], \text{x}] \text{ /; SumQ}[\text{u}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ /; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4269 $\text{Int}[(\text{csc}[(\text{c}_.) + (\text{d}_.) * (\text{x}_)] * (\text{b}_.) + (\text{a}_.))^{\text{n}_.}, \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{b}^2) * \text{Cot}[\text{c} + \text{d} * \text{x}] * ((\text{a} + \text{b} * \text{Csc}[\text{c} + \text{d} * \text{x}])^{\text{n} - 2} / (\text{d} * (\text{n} - 1))), \text{x}] + \text{Simp}[1 / (\text{n} - 1) \text{Int}[(\text{a} + \text{b} * \text{Csc}[\text{c} + \text{d} * \text{x}])^{\text{n} - 3} * \text{Simp}[\text{a}^3 * (\text{n} - 1) + (\text{b} * (\text{b}^2 * (\text{n} - 2) + 3 * \text{a}^2 * (\text{n} - 1))) * \text{Csc}[\text{c} + \text{d} * \text{x}] + (\text{a} * \text{b}^2 * (3 * \text{n} - 4)) * \text{Csc}[\text{c} + \text{d} * \text{x}]^2, \text{x}], \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \&\& \text{GtQ}[\text{n}, 2] \&\& \text{IntegerQ}[2 * \text{n}]$
- rule 4536 $\text{Int}[(\text{A}_.) + \text{csc}[(\text{e}_.) + (\text{f}_.) * (\text{x}_)] * (\text{B}_.) + \text{csc}[(\text{e}_.) + (\text{f}_.) * (\text{x}_)]^2 * (\text{C}_.) * (\text{csc}[(\text{e}_.) + (\text{f}_.) * (\text{x}_)] * (\text{b}_.) + (\text{a}_.)), \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{b}) * \text{C} * \text{Csc}[\text{e} + \text{f} * \text{x}] * (\text{Cot}[\text{e} + \text{f} * \text{x}] / (2 * \text{f})), \text{x}] + \text{Simp}[1/2 \quad \text{Int}[\text{Simp}[2 * \text{A} * \text{a} + (2 * \text{B} * \text{a} + \text{b} * (2 * \text{A} + \text{C})) * \text{Csc}[\text{e} + \text{f} * \text{x}] + 2 * (\text{a} * \text{C} + \text{B} * \text{b}) * \text{Csc}[\text{e} + \text{f} * \text{x}]^2, \text{x}], \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{e}, \text{f}, \text{A}, \text{B}, \text{C}\}, \text{x}]$
- rule 5992 $\text{Int}[\text{Coth}[(\text{c}_.) + (\text{d}_.) * (\text{x}_)] * \text{Csch}[(\text{c}_.) + (\text{d}_.) * (\text{x}_)] * (\text{Csch}[(\text{c}_.) + (\text{d}_.) * (\text{x}_)] * (\text{b}_.) + (\text{a}_.))^{\text{n}_.} * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.))^{\text{m}_.}, \text{x_Symbol}] \rightarrow \text{Simp}[(-(\text{e} + \text{f} * \text{x})^{\text{m}}) * ((\text{a} + \text{b} * \text{Csch}[\text{c} + \text{d} * \text{x}])^{\text{n} + 1} / (\text{b} * \text{d} * (\text{n} + 1))), \text{x}] + \text{Simp}[\text{f} * (\text{m} / (\text{b} * \text{d} * (\text{n} + 1))) \quad \text{Int}[(\text{e} + \text{f} * \text{x})^{\text{m} - 1} * (\text{a} + \text{b} * \text{Csch}[\text{c} + \text{d} * \text{x}])^{\text{n} + 1}, \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{n}\}, \text{x}] \&\& \text{IGtQ}[\text{m}, 0] \&\& \text{NeQ}[\text{n}, -1]$
- rule 6876 $\text{Int}[(\text{a}_.) + \text{ArcCsch}[(\text{c}_.) + (\text{d}_.) * (\text{x}_)] * (\text{b}_.)]^{\text{p}_.} * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.))^{\text{m}_.}, \text{x_Symbol}] \rightarrow \text{Simp}[-(\text{d}^{\text{m} + 1})^{-1} \quad \text{Subst}[\text{Int}[(\text{a} + \text{b} * \text{x})^{\text{p}} * \text{Csch}[\text{x}] * \text{Coth}[\text{x}] * (\text{d} * \text{e} - \text{c} * \text{f} + \text{f} * \text{Csch}[\text{x}])^{\text{m}}, \text{x}], \text{x}, \text{ArcCsch}[\text{c} + \text{d} * \text{x}]], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \&\& \text{IGtQ}[\text{p}, 0] \&\& \text{IntegerQ}[\text{m}]$

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.54

method	result
derivativedivides	$\frac{\operatorname{arccsch}(bx+a)a^4}{4} - \operatorname{arccsch}(bx+a)a^3(bx+a) + \frac{3 \operatorname{arccsch}(bx+a)a^2(bx+a)^2}{2} - \operatorname{arccsch}(bx+a)a(bx+a)^3 + \frac{\operatorname{arccsch}(bx+a)(bx+a)^4}{4}$
default	$\frac{\operatorname{arccsch}(bx+a)a^4}{4} - \operatorname{arccsch}(bx+a)a^3(bx+a) + \frac{3 \operatorname{arccsch}(bx+a)a^2(bx+a)^2}{2} - \operatorname{arccsch}(bx+a)a(bx+a)^3 + \frac{\operatorname{arccsch}(bx+a)(bx+a)^4}{4}$
parts	$\frac{x^4 \operatorname{arccsch}(bx+a)}{4} + \frac{\sqrt{b^2x^2+2bxa+a^2+1} \left(x^2\sqrt{b^2x^2+2bxa+a^2+1}b^2\sqrt{b^2}-3a^4 \operatorname{arctanh}\left(\frac{1}{\sqrt{b^2x^2+2bxa+a^2+1}}\right)\sqrt{b^2}- \right)}{\dots}$

input `int(x^3*arccsch(b*x+a),x,method=_RETURNVERBOSE)`output
$$\frac{1}{b^4} \left(\frac{1}{4} \operatorname{arccsch}(bx+a) a^4 - \operatorname{arccsch}(bx+a) a^3 (bx+a) + \frac{3}{2} \operatorname{arccsch}(bx+a) a^2 (bx+a)^2 - \operatorname{arccsch}(bx+a) a (bx+a)^3 + \frac{1}{4} \operatorname{arccsch}(bx+a) (bx+a)^4 \right) - \frac{1}{12} \left((bx+a)^2 + 1 \right)^{1/2} \left(3a^4 \operatorname{arctanh}\left(\frac{1}{\left((bx+a)^2 + 1\right)^{1/2}}\right) + 12a^3 \operatorname{arcsinh}(bx+a) - 18a^2 \left((bx+a)^2 + 1 \right)^{1/2} + 6a (bx+a) \left((bx+a)^2 + 1 \right)^{1/2} - (bx+a)^2 \left((bx+a)^2 + 1 \right)^{1/2} - 6a \operatorname{arcsinh}(bx+a) + 2 \left((bx+a)^2 + 1 \right)^{1/2} \right) / \left(\left((bx+a)^2 + 1 \right) / (bx+a)^2 \right)^{1/2} / (bx+a)$$
Fricas [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 325 vs. $2(127) = 254$.

Time = 0.13 (sec) , antiderivative size = 325, normalized size of antiderivative = 2.21

$$\int x^3 \operatorname{csch}^{-1}(a+bx) dx$$

$$= \frac{3b^4x^4 \log\left(\frac{(bx+a)\sqrt{\frac{b^2x^2+2abx+a^2+1}{b^2x^2+2abx+a^2}}+1}{bx+a}\right) - 3a^4 \log\left(-bx + (bx+a)\sqrt{\frac{b^2x^2+2abx+a^2+1}{b^2x^2+2abx+a^2}} - a + 1\right) + 3a^4 \log\left(-\right)}{\dots}$$

input `integrate(x^3*arccsch(b*x+a),x, algorithm="fricas")`

output
$$\frac{1}{12} \cdot (3b^4x^4 \log\left(\frac{(bx+a)\sqrt{(b^2x^2+2abx+a^2+1)}}{b^2x^2+2abx+a^2}\right) + 1) / (bx+a) - 3a^4 \log(-bx+(bx+a)\sqrt{(b^2x^2+2abx+a^2+1)}) / (b^2x^2+2abx+a^2) - a + 1 + 3a^4 \log(-bx+(bx+a)\sqrt{(b^2x^2+2abx+a^2+1)}) / (b^2x^2+2abx+a^2) - a - 1 + 6(2a^3 - a) \log(-bx+(bx+a)\sqrt{(b^2x^2+2abx+a^2+1)}) / (b^2x^2+2abx+a^2) - a + (b^3x^3 - 3ab^2x^2 + 13a^3 + (9a^2 - 2)bx - 2a) \sqrt{(b^2x^2+2abx+a^2+1)} / (b^2x^2+2abx+a^2) / b^4$$

Sympy [F]

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx = \int x^3 \operatorname{acsch}(a + bx) dx$$

input `integrate(x**3*acsch(b*x+a),x)`

output `Integral(x**3*acsch(a + b*x), x)`

Maxima [F]

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx = \int x^3 \operatorname{arcsch}(bx + a) dx$$

input `integrate(x^3*arccsch(b*x+a),x, algorithm="maxima")`

output
$$-1/2 \cdot (-Ia^3 + Ia) \cdot (\log(I \cdot (b^2x + ab) / b + 1) - \log(-I \cdot (b^2x + ab) / b + 1)) / b^4 + 1/8 \cdot (2b^4x^4 \log(\sqrt{(b^2x^2+2abx+a^2+1)} + 1) + b^2x^2 - 6abx - (a^4 - 6a^2 + 1) \log(b^2x^2+2abx+a^2+1) - 2(b^4x^4 - a^4) \log(bx+a)) / b^4 + \operatorname{integrate}(1/4 \cdot (b^2x^5 + abx^4) / (b^2x^2+2abx+a^2+(b^2x^2+2abx+a^2+1)^{(3/2)}+1), x)$$

Giac [F]

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx = \int x^3 \operatorname{arcsch}(bx + a) dx$$

input `integrate(x^3*arccsch(b*x+a),x, algorithm="giac")`

output `integrate(x^3*arccsch(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx = \int x^3 \operatorname{asinh}\left(\frac{1}{a + bx}\right) dx$$

input `int(x^3*asinh(1/(a + b*x)),x)`

output `int(x^3*asinh(1/(a + b*x)), x)`

Reduce [F]

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx = \int \operatorname{acsch}(bx + a) x^3 dx$$

input `int(x^3*acsch(b*x+a),x)`

output `int(acsch(a + b*x)*x**3,x)`

3.2 $\int x^2 \operatorname{csch}^{-1}(a + bx) dx$

Optimal result	47
Mathematica [A] (verified)	47
Rubi [A] (verified)	48
Maple [A] (verified)	50
Fricas [B] (verification not implemented)	50
Sympy [F]	51
Maxima [F]	51
Giac [F]	52
Mupad [F(-1)]	52
Reduce [F]	52

Optimal result

Integrand size = 10, antiderivative size = 110

$$\int x^2 \operatorname{csch}^{-1}(a + bx) dx = -\frac{5a(a + bx)\sqrt{1 + \frac{1}{(a+bx)^2}}}{6b^3} + \frac{x(a + bx)\sqrt{1 + \frac{1}{(a+bx)^2}}}{6b^2} + \frac{a^3 \operatorname{csch}^{-1}(a + bx)}{3b^3} + \frac{1}{3}x^3 \operatorname{csch}^{-1}(a + bx) - \frac{(1 - 6a^2) \operatorname{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx)^2}}\right)}{6b^3}$$

output `-5/6*a*(b*x+a)*(1+1/(b*x+a)^2)^(1/2)/b^3+1/6*x*(b*x+a)*(1+1/(b*x+a)^2)^(1/2)/b^2+1/3*a^3*arccsch(b*x+a)/b^3+1/3*x^3*arccsch(b*x+a)-1/6*(-6*a^2+1)*arctanh((1+1/(b*x+a)^2)^(1/2))/b^3`

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.17

$$\int x^2 \operatorname{csch}^{-1}(a + bx) dx = \frac{(-5a^2 - 4abx + b^2x^2) \sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}} + 2b^3x^3 \operatorname{csch}^{-1}(a + bx) + 2a^3 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right) + (-1 + 6a^2) \log\left(a + bx\right)}{6b^3}$$

input `Integrate[x^2*ArcCsch[a + b*x],x]`

output
$$\frac{((-5a^2 - 4abx + b^2x^2)\sqrt{(1 + a^2 + 2abx + b^2x^2)/(a + bx)^2} + 2b^3x^3\text{ArcCsch}[a + bx] + 2a^3\text{ArcSinh}[(a + bx)^{-1}] + (-1 + 6a^2)\text{Log}[(a + bx)(1 + \sqrt{(1 + a^2 + 2abx + b^2x^2)/(a + bx)^2})])}{(6b^3)}$$

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6876, 5992, 3042, 4269, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \text{csch}^{-1}(a + bx) dx$$

$$\downarrow 6876$$

$$-\frac{\int b^2 x^2 (a + bx)^2 \sqrt{1 + \frac{1}{(a+bx)^2}} \text{csch}^{-1}(a + bx) d\text{csch}^{-1}(a + bx)}{b^3}$$

$$\downarrow 5992$$

$$-\frac{-\frac{1}{3} \int -b^3 x^3 d\text{csch}^{-1}(a + bx) - \frac{1}{3} b^3 x^3 \text{csch}^{-1}(a + bx)}{b^3}$$

$$\downarrow 3042$$

$$-\frac{-\frac{1}{3} b^3 x^3 \text{csch}^{-1}(a + bx) - \frac{1}{3} \int (a - i \csc(i \text{csch}^{-1}(a + bx)))^3 d\text{csch}^{-1}(a + bx)}{b^3}$$

$$\downarrow 4269$$

$$-\frac{\frac{1}{3} \left(-\frac{1}{2} \int (2a^3 + 5(a + bx)^2 a + (1 - 6a^2)(a + bx)) d\text{csch}^{-1}(a + bx) - \frac{1}{2} bx \sqrt{\frac{1}{(a+bx)^2} + 1} (a + bx) \right) - \frac{1}{3} b^3 x^3 \text{csch}^{-1}(a + bx)}{b^3}$$

$$\downarrow 2009$$

$$\frac{\frac{1}{3}\left(\frac{1}{2}\left(-2a^3\operatorname{csch}^{-1}(a+bx) + (1-6a^2)\operatorname{arctanh}\left(\sqrt{\frac{1}{(a+bx)^2}+1}\right) + 5a(a+bx)\sqrt{\frac{1}{(a+bx)^2}+1}\right) - \frac{1}{2}bx(a+bx)\sqrt{\frac{1}{(a+bx)^2}+1}\right)}{b^3}$$

input `Int[x^2*ArcCsch[a + b*x],x]`

output `-((-1/3*(b^3*x^3*ArcCsch[a + b*x]) + (-1/2*(b*x*(a + b*x)*Sqrt[1 + (a + b*x)^(-2)])) + (5*a*(a + b*x)*Sqrt[1 + (a + b*x)^(-2)] - 2*a^3*ArcCsch[a + b*x] + (1 - 6*a^2)*ArcTanh[Sqrt[1 + (a + b*x)^(-2)]])/2)/3)/b^3)`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :=> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4269 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^ (n_), x_Symbol] :=> Simp[(-b^2)*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n - 2)/(d*(n - 1))), x] + Simp[1/(n - 1) Int[(a + b*Csc[c + d*x])^(n - 3)*Simp[a^3*(n - 1) + (b*(b^2*(n - 2) + 3*a^2*(n - 1))*Csc[c + d*x] + (a*b^2*(3*n - 4))*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 2] && IntegerQ[2*n]`

rule 5992 `Int[Coth[(c_.) + (d_.)*(x_)]*Csch[(c_.) + (d_.)*(x_)]*(Csch[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^ (n_.)*((e_.) + (f_.)*(x_))^ (m_.), x_Symbol] :=> Simp[(-(e + f*x)^m)*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

rule 6876 `Int[((a_.) + ArcCsch[(c_.) + (d_.)*(x_)]*(b_.))^ (p_.)*((e_.) + (f_.)*(x_))^ (m_.), x_Symbol] :=> Simp[-(d^(m + 1))^ (-1) Subst[Int[(a + b*x)^p*Csch[x]*Coth[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.55

method	result
derivativedivides	$-\frac{\operatorname{arccsch}(bx+a)a^3}{3} + \operatorname{arccsch}(bx+a)a^2(bx+a) - \operatorname{arccsch}(bx+a)a(bx+a)^2 + \frac{\operatorname{arccsch}(bx+a)(bx+a)^3}{3} + \frac{\sqrt{(bx+a)^2+1} \left(2a^3 \operatorname{arctanh}\left(\frac{1}{\sqrt{b^2x^2+2bxa+a^2+1}}\right) \sqrt{b^2+6} \ln\left(\frac{b^2x+\sqrt{b^2x^2+2bxa+a^2+1}\sqrt{b^2+6}}{\sqrt{b^2+6}}\right) \right)}{b^3}$
default	$-\frac{\operatorname{arccsch}(bx+a)a^3}{3} + \operatorname{arccsch}(bx+a)a^2(bx+a) - \operatorname{arccsch}(bx+a)a(bx+a)^2 + \frac{\operatorname{arccsch}(bx+a)(bx+a)^3}{3} + \frac{\sqrt{(bx+a)^2+1} \left(2a^3 \operatorname{arctanh}\left(\frac{1}{\sqrt{b^2x^2+2bxa+a^2+1}}\right) \sqrt{b^2+6} \ln\left(\frac{b^2x+\sqrt{b^2x^2+2bxa+a^2+1}\sqrt{b^2+6}}{\sqrt{b^2+6}}\right) \right)}{b^3}$
parts	$\frac{x^3 \operatorname{arccsch}(bx+a)}{3} + \frac{\sqrt{b^2x^2+2bxa+a^2+1} \left(2a^3 \operatorname{arctanh}\left(\frac{1}{\sqrt{b^2x^2+2bxa+a^2+1}}\right) \sqrt{b^2+6} \ln\left(\frac{b^2x+\sqrt{b^2x^2+2bxa+a^2+1}\sqrt{b^2+6}}{\sqrt{b^2+6}}\right) \right)}{6b^3 \sqrt{b^2x^2+2bxa+a^2+1}}$

input `int(x^2*arccsch(b*x+a),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{b^3} * (-1/3 * \operatorname{arccsch}(b*x+a) * a^3 + \operatorname{arccsch}(b*x+a) * a^2 * (b*x+a) - \operatorname{arccsch}(b*x+a) * a * (b*x+a)^2 + 1/3 * \operatorname{arccsch}(b*x+a) * (b*x+a)^3 + 1/6 * ((b*x+a)^2 + 1)^{(1/2)} * (2 * a^3 * \operatorname{arctanh}(1 / ((b*x+a)^2 + 1)^{(1/2)}) + 6 * a^2 * \operatorname{arcsinh}(b*x+a) - 6 * a * ((b*x+a)^2 + 1)^{(1/2)} + (b*x+a) * ((b*x+a)^2 + 1)^{(1/2)} - \operatorname{arcsinh}(b*x+a)) / (((b*x+a)^2 + 1) / (b*x+a)^2)^{(1/2)} / (b*x+a))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 306 vs. 2(94) = 188.

Time = 0.15 (sec) , antiderivative size = 306, normalized size of antiderivative = 2.78

$$\int x^2 \operatorname{csch}^{-1}(a + bx) dx$$

$$= \frac{2b^3x^3 \log\left(\frac{(bx+a)\sqrt{\frac{b^2x^2+2abx+a^2+1}{b^2x^2+2abx+a^2}}+1}{bx+a}\right) + 2a^3 \log\left(-bx + (bx+a)\sqrt{\frac{b^2x^2+2abx+a^2+1}{b^2x^2+2abx+a^2}} - a + 1\right) - 2a^3 \log\left(-\right)}{6b^3 \sqrt{b^2x^2+2abx+a^2+1}}$$

input `integrate(x^2*arccsch(b*x+a),x, algorithm="fricas")`

output

```
1/6*(2*b^3*x^3*log(((b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)) + 2*a^3*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a + 1) - 2*a^3*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a - 1) - (6*a^2 - 1)*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a) + (b^2*x^2 - 4*a*b*x - 5*a^2)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)))/b^3
```

Sympy [F]

$$\int x^2 \operatorname{csch}^{-1}(a + bx) dx = \int x^2 \operatorname{acsch}(a + bx) dx$$

input

```
integrate(x**2*acsch(b*x+a),x)
```

output

```
Integral(x**2*acsch(a + b*x), x)
```

Maxima [F]

$$\int x^2 \operatorname{csch}^{-1}(a + bx) dx = \int x^2 \operatorname{arcsch}(bx + a) dx$$

input

```
integrate(x^2*arccsch(b*x+a),x, algorithm="maxima")
```

output

```
-1/6*(3*I*a^2 - I)*(log(I*(b^2*x + a*b)/b + 1) - log(-I*(b^2*x + a*b)/b + 1))/b^3 + 1/6*(2*b^3*x^3*log(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) + 1) + 2*b*x + (a^3 - 3*a)*log(b^2*x^2 + 2*a*b*x + a^2 + 1) - 2*(b^3*x^3 + a^3)*log(b*x + a))/b^3 + integrate(1/3*(b^2*x^4 + a*b*x^3)/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 + 1)^(3/2) + 1), x)
```

Giac [F]

$$\int x^2 \operatorname{csch}^{-1}(a + bx) dx = \int x^2 \operatorname{arcsch}(bx + a) dx$$

input `integrate(x^2*arccsch(b*x+a),x, algorithm="giac")`

output `integrate(x^2*arccsch(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{csch}^{-1}(a + bx) dx = \int x^2 \operatorname{asinh}\left(\frac{1}{a + bx}\right) dx$$

input `int(x^2*asinh(1/(a + b*x)),x)`

output `int(x^2*asinh(1/(a + b*x)), x)`

Reduce [F]

$$\int x^2 \operatorname{csch}^{-1}(a + bx) dx = \int \operatorname{acsch}(bx + a) x^2 dx$$

input `int(x^2*acsch(b*x+a),x)`

output `int(acsch(a + b*x)*x**2,x)`

3.3 $\int x \operatorname{csch}^{-1}(a + bx) dx$

Optimal result	53
Mathematica [A] (verified)	53
Rubi [A] (verified)	54
Maple [A] (verified)	57
Fricas [B] (verification not implemented)	57
Sympy [F]	58
Maxima [F]	58
Giac [F]	59
Mupad [F(-1)]	59
Reduce [F]	59

Optimal result

Integrand size = 8, antiderivative size = 75

$$\int x \operatorname{csch}^{-1}(a + bx) dx = \frac{(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}}{2b^2} - \frac{a^2 \operatorname{csch}^{-1}(a + bx)}{2b^2} + \frac{1}{2} x^2 \operatorname{csch}^{-1}(a + bx) - \frac{a \operatorname{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx)^2}}\right)}{b^2}$$

output

$1/2*(b*x+a)*(1+1/(b*x+a)^2)^{(1/2)}/b^2-1/2*a^2*\operatorname{arccsch}(b*x+a)/b^2+1/2*x^2*a \operatorname{rccsch}(b*x+a)-a*\operatorname{arctanh}((1+1/(b*x+a)^2)^{(1/2)})/b^2$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.47

$$\int x \operatorname{csch}^{-1}(a + bx) dx = \frac{(a + bx) \sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}} + b^2 x^2 \operatorname{csch}^{-1}(a + bx) - a^2 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right) - 2a \log\left((a + bx) \left(1 + \sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}}\right)\right)}{2b^2}$$

input

`Integrate[x*ArcCsch[a + b*x], x]`

output

```
((a + b*x)*Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2] + b^2*x^2*ArcCs
ch[a + b*x] - a^2*ArcSinh[(a + b*x)^(-1)] - 2*a*Log[(a + b*x)*(1 + Sqrt[(1
+ a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2])])/(2*b^2)
```

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.99, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 1.625$, Rules used = {6876, 25, 5992, 3042, 4260, 25, 26, 3042, 25, 26, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{csch}^{-1}(a + bx) dx$$

$$\downarrow 6876$$

$$\frac{\int bx(a + bx)^2 \sqrt{1 + \frac{1}{(a+bx)^2}} \operatorname{csch}^{-1}(a + bx) d\operatorname{csch}^{-1}(a + bx)}{b^2}$$

$$\downarrow 25$$

$$\frac{\int -bx(a + bx)^2 \sqrt{1 + \frac{1}{(a+bx)^2}} \operatorname{csch}^{-1}(a + bx) d\operatorname{csch}^{-1}(a + bx)}{b^2}$$

$$\downarrow 5992$$

$$\frac{\frac{1}{2} \int b^2 x^2 d\operatorname{csch}^{-1}(a + bx) - \frac{1}{2} b^2 x^2 \operatorname{csch}^{-1}(a + bx)}{b^2}$$

$$\downarrow 3042$$

$$\frac{-\frac{1}{2} b^2 x^2 \operatorname{csch}^{-1}(a + bx) + \frac{1}{2} \int (a - i \csc(i \operatorname{csch}^{-1}(a + bx)))^2 d\operatorname{csch}^{-1}(a + bx)}{b^2}$$

$$\downarrow 4260$$

$$\frac{-\frac{1}{2} b^2 x^2 \operatorname{csch}^{-1}(a + bx) + \frac{1}{2} (-2ia \int -i(a + bx) d\operatorname{csch}^{-1}(a + bx) - \int -(a + bx)^2 d\operatorname{csch}^{-1}(a + bx) + a^2 \operatorname{csch}^{-1}(a + bx))}{b^2}$$

$$\downarrow 25$$

$$\frac{-\frac{1}{2}b^2x^2\operatorname{csch}^{-1}(a+bx) + \frac{1}{2}(-2ia \int -i(a+bx)d\operatorname{csch}^{-1}(a+bx) + \int (a+bx)^2d\operatorname{csch}^{-1}(a+bx) + a^2\operatorname{csch}^{-1}(a+bx))}{b^2}$$

↓ 26

$$\frac{\frac{1}{2}(-2a \int (a+bx)d\operatorname{csch}^{-1}(a+bx) + \int (a+bx)^2d\operatorname{csch}^{-1}(a+bx) + a^2\operatorname{csch}^{-1}(a+bx)) - \frac{1}{2}b^2x^2\operatorname{csch}^{-1}(a+bx)}{b^2}$$

↓ 3042

$$\frac{-\frac{1}{2}b^2x^2\operatorname{csch}^{-1}(a+bx) + \frac{1}{2}(-2a \int i \csc(\operatorname{icsch}^{-1}(a+bx))d\operatorname{csch}^{-1}(a+bx) + \int -\csc(\operatorname{icsch}^{-1}(a+bx))^2d\operatorname{csch}^{-1}(a+bx))}{b^2}$$

↓ 25

$$\frac{-\frac{1}{2}b^2x^2\operatorname{csch}^{-1}(a+bx) + \frac{1}{2}(-2a \int i \csc(\operatorname{icsch}^{-1}(a+bx))d\operatorname{csch}^{-1}(a+bx) - \int \csc(\operatorname{icsch}^{-1}(a+bx))^2d\operatorname{csch}^{-1}(a+bx))}{b^2}$$

↓ 26

$$\frac{-\frac{1}{2}b^2x^2\operatorname{csch}^{-1}(a+bx) + \frac{1}{2}(-2ia \int \csc(\operatorname{icsch}^{-1}(a+bx))d\operatorname{csch}^{-1}(a+bx) - \int \csc(\operatorname{icsch}^{-1}(a+bx))^2d\operatorname{csch}^{-1}(a+bx))}{b^2}$$

↓ 4254

$$\frac{-\frac{1}{2}b^2x^2\operatorname{csch}^{-1}(a+bx) + \frac{1}{2}(-i \int 1d(-i(a+bx)\sqrt{1+\frac{1}{(a+bx)^2}}) - 2ia \int \csc(\operatorname{icsch}^{-1}(a+bx))d\operatorname{csch}^{-1}(a+bx))}{b^2}$$

↓ 24

$$\frac{-\frac{1}{2}b^2x^2\operatorname{csch}^{-1}(a+bx) + \frac{1}{2}(-2ia \int \csc(\operatorname{icsch}^{-1}(a+bx))d\operatorname{csch}^{-1}(a+bx) + a^2\operatorname{csch}^{-1}(a+bx) - (a+bx)\sqrt{\frac{1}{(a+bx)^2}+1})}{b^2}$$

↓ 4257

$$\frac{\frac{1}{2}(a^2\operatorname{csch}^{-1}(a+bx) + 2a\operatorname{arctanh}\left(\sqrt{\frac{1}{(a+bx)^2}+1}\right) - (a+bx)\sqrt{\frac{1}{(a+bx)^2}+1}) - \frac{1}{2}b^2x^2\operatorname{csch}^{-1}(a+bx)}{b^2}$$

input `Int[x*ArcCsch[a + b*x], x]`

output
$$\frac{-((-1/2*(b^2*x^2*ArcCsch[a + b*x]) + (-((a + b*x)*Sqrt[1 + (a + b*x)^{-2}]) + a^2*ArcCsch[a + b*x] + 2*a*ArcTanh[Sqrt[1 + (a + b*x)^{-2}]])/2)/b^2}$$

Defintions of rubi rules used

rule 24
$$\text{Int}[a_, x_Symbol] \text{ :> Simp}[a*x, x] \text{ /; FreeQ}[a, x]$$

rule 25
$$\text{Int}[-(Fx_), x_Symbol] \text{ :> Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$$

rule 26
$$\text{Int}[(\text{Complex}[0, a_])*(Fx_), x_Symbol] \text{ :> Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{ Int}[Fx, x], x] \text{ /; FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$$

rule 3042
$$\text{Int}[u_, x_Symbol] \text{ :> Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$$

rule 4254
$$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_)]^{(n_.)}, x_Symbol] \text{ :> Simp}[-d^{(-1)} \text{ Subst}[\text{Int}[\text{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \text{Cot}[c + d*x]], x] \text{ /; FreeQ}\{c, d\}, x] \ \&\& \ \text{IGtQ}[n/2, 0]$$

rule 4257
$$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_)], x_Symbol] \text{ :> Simp}[-\text{ArcTanh}[\text{Cos}[c + d*x]]/d, x] \text{ /; FreeQ}\{c, d\}, x]$$

rule 4260
$$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.) + (a_))^{2}, x_Symbol] \text{ :> Simp}[a^2*x, x] + (\text{Simp}[2*a*b \text{ Int}[\text{Csc}[c + d*x], x], x] + \text{Simp}[b^2 \text{ Int}[\text{Csc}[c + d*x]^2, x], x]) \text{ /; FreeQ}\{a, b, c, d\}, x]$$

rule 5992
$$\text{Int}[\text{Coth}[(c_.) + (d_.)*(x_)]*\text{Csch}[(c_.) + (d_.)*(x_)]*(\text{Csch}[(c_.) + (d_.)*(x_)]*(b_.) + (a_))^{(n_.)}*((e_.) + (f_.)*(x_))^{(m_.)}, x_Symbol] \text{ :> Simp}[(-(e + f*x)^m)*((a + b*\text{Csch}[c + d*x])^{(n + 1)}/(b*d*(n + 1))), x] + \text{Simp}[f*(m/(b*d*(n + 1))) \text{ Int}[(e + f*x)^{(m - 1)}*(a + b*\text{Csch}[c + d*x])^{(n + 1)}, x], x] \text{ /; FreeQ}\{a, b, c, d, e, f, n\}, x] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{NeQ}[n, -1]$$

rule 6876

```
Int[((a_.) + ArcCsch[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Csch[x]*Coth[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.29

method	result
derivativedivides	$\frac{\frac{\operatorname{arccsch}(bx+a)(bx+a)^2}{2} - \operatorname{arccsch}(bx+a)a(bx+a) - \frac{\sqrt{(bx+a)^2+1} (2a \operatorname{arcsinh}(bx+a) - \sqrt{(bx+a)^2+1})}{2(bx+a)\sqrt{\frac{(bx+a)^2+1}{(bx+a)^2}}}}{b^2}$
default	$\frac{\frac{\operatorname{arccsch}(bx+a)(bx+a)^2}{2} - \operatorname{arccsch}(bx+a)a(bx+a) - \frac{\sqrt{(bx+a)^2+1} (2a \operatorname{arcsinh}(bx+a) - \sqrt{(bx+a)^2+1})}{2(bx+a)\sqrt{\frac{(bx+a)^2+1}{(bx+a)^2}}}}{b^2}$
parts	$\frac{x^2 \operatorname{arccsch}(bx+a)}{2} - \frac{\sqrt{b^2x^2+2bxa+a^2+1} \left(a^2 \operatorname{arctanh}\left(\frac{1}{\sqrt{b^2x^2+2bxa+a^2+1}}\right) \sqrt{b^2+2a} \ln\left(\frac{b^2x+\sqrt{b^2x^2+2bxa+a^2+1} \sqrt{b^2}}{\sqrt{b^2}}\right) \right)}{2b^2 \sqrt{\frac{b^2x^2+2bxa+a^2+1}{(bx+a)^2}} (bx+a) \sqrt{b^2}}$

input

```
int(x*arccsch(b*x+a), x, method=_RETURNVERBOSE)
```

output

```
1/b^2*(1/2*arccsch(b*x+a)*(b*x+a)^2-arccsch(b*x+a)*a*(b*x+a)-1/2*((b*x+a)^2+1)^(1/2)*(2*a*arcsinh(b*x+a)-((b*x+a)^2+1)^(1/2))/(b*x+a)/(((b*x+a)^2+1)/(b*x+a)^2)^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 285 vs. 2(65) = 130.

Time = 0.10 (sec) , antiderivative size = 285, normalized size of antiderivative = 3.80

$$\int x \operatorname{csch}^{-1}(a + bx) dx$$

$$= \frac{b^2x^2 \log\left(\frac{(bx+a)\sqrt{\frac{b^2x^2+2abx+a^2+1}{b^2x^2+2abx+a^2}+1}}{bx+a}\right) - a^2 \log\left(-bx + (bx+a)\sqrt{\frac{b^2x^2+2abx+a^2+1}{b^2x^2+2abx+a^2}} - a + 1\right) + a^2 \log\left(-bx + \dots\right)}{\dots}$$

input `integrate(x*arccsch(b*x+a),x, algorithm="fricas")`

output
$$\frac{1}{2}(b^2x^2 \log\left(\frac{(bx+a)\sqrt{(b^2x^2+2abx+a^2+1)}}{(b^2x^2+2abx+a^2)}+1\right) - a^2 \log\left(\frac{-bx+(bx+a)\sqrt{(b^2x^2+2abx+a^2+1)}}{(b^2x^2+2abx+a^2)}-a+1\right) + a^2 \log\left(\frac{-bx+(bx+a)\sqrt{(b^2x^2+2abx+a^2+1)}}{(b^2x^2+2abx+a^2)}-a-1\right) + 2a \log\left(\frac{-bx+(bx+a)\sqrt{(b^2x^2+2abx+a^2+1)}}{(b^2x^2+2abx+a^2)}-a\right) + (bx+a)\sqrt{(b^2x^2+2abx+a^2+1)}/(b^2x^2+2abx+a^2)))/b^2$$

Sympy [F]

$$\int x \operatorname{csch}^{-1}(a+bx) dx = \int x \operatorname{acsch}(a+bx) dx$$

input `integrate(x*acsch(b*x+a),x)`

output `Integral(x*acsch(a + b*x), x)`

Maxima [F]

$$\int x \operatorname{csch}^{-1}(a+bx) dx = \int x \operatorname{arcsch}(bx+a) dx$$

input `integrate(x*arccsch(b*x+a),x, algorithm="maxima")`

output
$$\frac{1}{2}I*a*(\log(I*(b^2*x + a*b)/b + 1) - \log(-I*(b^2*x + a*b)/b + 1))/b^2 + 1/4*(2*b^2*x^2*\log(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) + 1) - (a^2 - 1)*\log(b^2*x^2 + 2*a*b*x + a^2 + 1) - 2*(b^2*x^2 - a^2)*\log(b*x + a))/b^2 + \operatorname{integrate}(1/2*(b^2*x^3 + a*b*x^2)/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 + 1)^{(3/2)} + 1), x)$$

Giac [F]

$$\int x \operatorname{csch}^{-1}(a + bx) dx = \int x \operatorname{arcsch}(bx + a) dx$$

input `integrate(x*arccsch(b*x+a),x, algorithm="giac")`

output `integrate(x*arccsch(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{csch}^{-1}(a + bx) dx = \int x \operatorname{asinh}\left(\frac{1}{a + bx}\right) dx$$

input `int(x*asinh(1/(a + b*x)),x)`

output `int(x*asinh(1/(a + b*x)), x)`

Reduce [F]

$$\int x \operatorname{csch}^{-1}(a + bx) dx = \int \operatorname{acsch}(bx + a) x dx$$

input `int(x*acsch(b*x+a),x)`

output `int(acsch(a + b*x)*x,x)`

3.4 $\int \frac{\operatorname{csch}^{-1}(a+bx)}{x} dx$

Optimal result	60
Mathematica [C] (verified)	61
Rubi [C] (verified)	62
Maple [F]	67
Fricas [F]	67
Sympy [F]	68
Maxima [F]	68
Giac [F]	68
Mupad [F(-1)]	69
Reduce [F]	69

Optimal result

Integrand size = 10, antiderivative size = 162

$$\begin{aligned} \int \frac{\operatorname{csch}^{-1}(a+bx)}{x} dx &= \operatorname{csch}^{-1}(a+bx) \log \left(1 - \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{1 - \sqrt{1+a^2}} \right) \\ &\quad + \operatorname{csch}^{-1}(a+bx) \log \left(1 - \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{1 + \sqrt{1+a^2}} \right) \\ &\quad - \operatorname{csch}^{-1}(a+bx) \log \left(1 - e^{2\operatorname{csch}^{-1}(a+bx)} \right) \\ &\quad + \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{1 - \sqrt{1+a^2}} \right) + \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{1 + \sqrt{1+a^2}} \right) \\ &\quad - \frac{1}{2} \operatorname{PolyLog} \left(2, e^{2\operatorname{csch}^{-1}(a+bx)} \right) \end{aligned}$$

output

```
arccsch(b*x+a)*ln(1-a*(1/(b*x+a)+(1+1/(b*x+a)^2)^(1/2))/(1-(a^2+1)^(1/2)))
+arccsch(b*x+a)*ln(1-a*(1/(b*x+a)+(1+1/(b*x+a)^2)^(1/2))/(1+(a^2+1)^(1/2)))
)-arccsch(b*x+a)*ln(1-(1/(b*x+a)+(1+1/(b*x+a)^2)^(1/2))^2)+polylog(2,a*(1/
(b*x+a)+(1+1/(b*x+a)^2)^(1/2))/(1-(a^2+1)^(1/2)))+polylog(2,a*(1/(b*x+a)+(
1+1/(b*x+a)^2)^(1/2))/(1+(a^2+1)^(1/2)))-1/2*polylog(2,(1/(b*x+a)+(1+1/(b*
x+a)^2)^(1/2))^2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 427, normalized size of antiderivative = 2.64

$$\begin{aligned}
 & \int \frac{\operatorname{csch}^{-1}(a+bx)}{x} dx \\
 &= \frac{1}{8} \left(\pi^2 - 4i\pi \operatorname{csch}^{-1}(a+bx) - 8\operatorname{csch}^{-1}(a+bx)^2 \right. \\
 &\quad - 32 \arcsin \left(\frac{\sqrt{\frac{-i+a}{a}}}{\sqrt{2}} \right) \arctan \left(\frac{(1-ia) \cot \left(\frac{1}{4}(\pi + 2i \operatorname{csch}^{-1}(a+bx)) \right)}{\sqrt{1+a^2}} \right) \\
 &\quad \quad - 8\operatorname{csch}^{-1}(a+bx) \log \left(1 - e^{-2\operatorname{csch}^{-1}(a+bx)} \right) \\
 &\quad \quad + 4i\pi \log \left(1 - \frac{(-1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) \\
 &\quad \quad + 8\operatorname{csch}^{-1}(a+bx) \log \left(1 - \frac{(-1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) \\
 &\quad \quad + 16i \arcsin \left(\frac{\sqrt{\frac{-i+a}{a}}}{\sqrt{2}} \right) \log \left(1 - \frac{(-1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) \\
 &\quad \quad \quad + 4i\pi \log \left(1 + \frac{(1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) \\
 &\quad \quad \quad + 8\operatorname{csch}^{-1}(a+bx) \log \left(1 + \frac{(1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) \\
 &\quad - 16i \arcsin \left(\frac{\sqrt{\frac{-i+a}{a}}}{\sqrt{2}} \right) \log \left(1 + \frac{(1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) - 4i\pi \log \left(\frac{bx}{a+bx} \right) \\
 &\quad \quad + 4 \operatorname{PolyLog} \left(2, e^{-2\operatorname{csch}^{-1}(a+bx)} \right) + 8 \operatorname{PolyLog} \left(2, \frac{(-1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) \\
 &\quad \quad \quad + 8 \operatorname{PolyLog} \left(2, -\frac{(1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) \left. \right)
 \end{aligned}$$

input `Integrate[ArcCsch[a + b*x]/x,x]`

output

```
(Pi^2 - (4*I)*Pi*ArcCsch[a + b*x] - 8*ArcCsch[a + b*x]^2 - 32*ArcSin[Sqrt[
(-I + a)/a]/Sqrt[2]]*ArcTan[((1 - I*a)*Cot[(Pi + (2*I)*ArcCsch[a + b*x])/4
])/Sqrt[1 + a^2]] - 8*ArcCsch[a + b*x]*Log[1 - E^(-2*ArcCsch[a + b*x])] +
(4*I)*Pi*Log[1 - ((-1 + Sqrt[1 + a^2])*E^ArcCsch[a + b*x])/a] + 8*ArcCsch[
a + b*x]*Log[1 - ((-1 + Sqrt[1 + a^2])*E^ArcCsch[a + b*x])/a] + (16*I)*Arc
Sin[Sqrt[(-I + a)/a]/Sqrt[2]]*Log[1 - ((-1 + Sqrt[1 + a^2])*E^ArcCsch[a +
b*x])/a] + (4*I)*Pi*Log[1 + ((1 + Sqrt[1 + a^2])*E^ArcCsch[a + b*x])/a] +
8*ArcCsch[a + b*x]*Log[1 + ((1 + Sqrt[1 + a^2])*E^ArcCsch[a + b*x])/a] - (
16*I)*ArcSin[Sqrt[(-I + a)/a]/Sqrt[2]]*Log[1 + ((1 + Sqrt[1 + a^2])*E^ArcC
sch[a + b*x])/a] - (4*I)*Pi*Log[(b*x)/(a + b*x)] + 4*PolyLog[2, E^(-2*ArcC
sch[a + b*x])] + 8*PolyLog[2, ((-1 + Sqrt[1 + a^2])*E^ArcCsch[a + b*x])/a]
+ 8*PolyLog[2, -(((1 + Sqrt[1 + a^2])*E^ArcCsch[a + b*x])/a)])/8
```

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.01 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.39, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 1.600$, Rules used = {6876, 25, 6130, 6103, 25, 3042, 26, 4199, 25, 2620, 2715, 2838, 6095, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}^{-1}(a+bx)}{x} dx \\
 & \quad \downarrow \text{6876} \\
 & - \int \frac{(a+bx)^2 \sqrt{1 + \frac{1}{(a+bx)^2}} \operatorname{csch}^{-1}(a+bx)}{bx} d\operatorname{csch}^{-1}(a+bx) \\
 & \quad \downarrow \text{25} \\
 & \int - \frac{(a+bx)^2 \sqrt{\frac{1}{(a+bx)^2} + 1} \operatorname{csch}^{-1}(a+bx)}{bx} d\operatorname{csch}^{-1}(a+bx) \\
 & \quad \downarrow \text{6130}
 \end{aligned}$$

$$\int \frac{(a+bx)\sqrt{\frac{1}{(a+bx)^2} + 1}\operatorname{csch}^{-1}(a+bx)}{\frac{a}{a+bx} - 1} d\operatorname{csch}^{-1}(a+bx)$$

↓ 6103

$$a \int -\frac{\sqrt{1 + \frac{1}{(a+bx)^2}}\operatorname{csch}^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d\operatorname{csch}^{-1}(a+bx) - \int (a+bx)\sqrt{1 + \frac{1}{(a+bx)^2}}\operatorname{csch}^{-1}(a+bx) d\operatorname{csch}^{-1}(a+bx)$$

↓ 25

$$- \int (a+bx)\sqrt{1 + \frac{1}{(a+bx)^2}}\operatorname{csch}^{-1}(a+bx) d\operatorname{csch}^{-1}(a+bx) - a \int \frac{\sqrt{1 + \frac{1}{(a+bx)^2}}\operatorname{csch}^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d\operatorname{csch}^{-1}(a+bx)$$

↓ 3042

$$-a \int \frac{\sqrt{1 + \frac{1}{(a+bx)^2}}\operatorname{csch}^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d\operatorname{csch}^{-1}(a+bx) - \int -i\operatorname{csch}^{-1}(a+bx) \tan\left(i\operatorname{csch}^{-1}(a+bx) + \frac{\pi}{2}\right) d\operatorname{csch}^{-1}(a+bx)$$

↓ 26

$$-a \int \frac{\sqrt{1 + \frac{1}{(a+bx)^2}}\operatorname{csch}^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d\operatorname{csch}^{-1}(a+bx) + i \int \operatorname{csch}^{-1}(a+bx) \tan\left(i\operatorname{csch}^{-1}(a+bx) + \frac{\pi}{2}\right) d\operatorname{csch}^{-1}(a+bx)$$

↓ 4199

$$-a \int \frac{\sqrt{1 + \frac{1}{(a+bx)^2}}\operatorname{csch}^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d\operatorname{csch}^{-1}(a+bx) + i \left(2i \int -\frac{e^{2\operatorname{csch}^{-1}(a+bx)}\operatorname{csch}^{-1}(a+bx)}{1 - e^{2\operatorname{csch}^{-1}(a+bx)}} d\operatorname{csch}^{-1}(a+bx) - \frac{1}{2}i\operatorname{csch}^{-1}(a+bx)^2 \right)$$

↓ 25

$$-a \int \frac{\sqrt{1 + \frac{1}{(a+bx)^2}}\operatorname{csch}^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d\operatorname{csch}^{-1}(a+bx) + i \left(-2i \int \frac{e^{2\operatorname{csch}^{-1}(a+bx)}\operatorname{csch}^{-1}(a+bx)}{1 - e^{2\operatorname{csch}^{-1}(a+bx)}} d\operatorname{csch}^{-1}(a+bx) - \frac{1}{2}i\operatorname{csch}^{-1}(a+bx)^2 \right)$$

$$\begin{aligned} & \downarrow 2620 \\ & -a \int \frac{\sqrt{1 + \frac{1}{(a+bx)^2}} \operatorname{csch}^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d\operatorname{csch}^{-1}(a+bx) + \\ & i \left(-2i \left(\frac{1}{2} \int \log \left(1 - e^{2\operatorname{csch}^{-1}(a+bx)} \right) d\operatorname{csch}^{-1}(a+bx) - \frac{1}{2} \operatorname{csch}^{-1}(a+bx) \log \left(1 - e^{2\operatorname{csch}^{-1}(a+bx)} \right) \right) - \frac{1}{2} i \operatorname{csch}^{-1}(a+bx)^2 \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 2715 \\ & -a \int \frac{\sqrt{1 + \frac{1}{(a+bx)^2}} \operatorname{csch}^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d\operatorname{csch}^{-1}(a+bx) + \\ & i \left(-2i \left(\frac{1}{4} \int e^{-2\operatorname{csch}^{-1}(a+bx)} \log \left(1 - e^{2\operatorname{csch}^{-1}(a+bx)} \right) d e^{2\operatorname{csch}^{-1}(a+bx)} - \frac{1}{2} \operatorname{csch}^{-1}(a+bx) \log \left(1 - e^{2\operatorname{csch}^{-1}(a+bx)} \right) \right) \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 2838 \\ & -a \int \frac{\sqrt{1 + \frac{1}{(a+bx)^2}} \operatorname{csch}^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d\operatorname{csch}^{-1}(a+bx) + \\ & i \left(-2i \left(-\frac{1}{4} \operatorname{PolyLog} \left(2, e^{2\operatorname{csch}^{-1}(a+bx)} \right) - \frac{1}{2} \operatorname{csch}^{-1}(a+bx) \log \left(1 - e^{2\operatorname{csch}^{-1}(a+bx)} \right) \right) - \frac{1}{2} i \operatorname{csch}^{-1}(a+bx)^2 \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 6095 \\ & -a \left(\int \frac{e^{\operatorname{csch}^{-1}(a+bx)} \operatorname{csch}^{-1}(a+bx)}{-e^{\operatorname{csch}^{-1}(a+bx)} a - \sqrt{a^2+1} + 1} d\operatorname{csch}^{-1}(a+bx) + \int \frac{e^{\operatorname{csch}^{-1}(a+bx)} \operatorname{csch}^{-1}(a+bx)}{-e^{\operatorname{csch}^{-1}(a+bx)} a + \sqrt{a^2+1} + 1} d\operatorname{csch}^{-1}(a+bx) + \operatorname{csch}^{-1}(a+bx) \right) \\ & i \left(-2i \left(-\frac{1}{4} \operatorname{PolyLog} \left(2, e^{2\operatorname{csch}^{-1}(a+bx)} \right) - \frac{1}{2} \operatorname{csch}^{-1}(a+bx) \log \left(1 - e^{2\operatorname{csch}^{-1}(a+bx)} \right) \right) - \frac{1}{2} i \operatorname{csch}^{-1}(a+bx)^2 \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 2620 \\ & -a \left(\frac{\int \log \left(1 - \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{1 - \sqrt{a^2+1}} \right) d\operatorname{csch}^{-1}(a+bx)}{a} + \frac{\int \log \left(1 - \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{\sqrt{a^2+1} + 1} \right) d\operatorname{csch}^{-1}(a+bx)}{a} - \operatorname{csch}^{-1}(a+bx) \right) \\ & i \left(-2i \left(-\frac{1}{4} \operatorname{PolyLog} \left(2, e^{2\operatorname{csch}^{-1}(a+bx)} \right) - \frac{1}{2} \operatorname{csch}^{-1}(a+bx) \log \left(1 - e^{2\operatorname{csch}^{-1}(a+bx)} \right) \right) - \frac{1}{2} i \operatorname{csch}^{-1}(a+bx)^2 \right) \end{aligned}$$

$$\downarrow 2715$$

$$\begin{aligned}
& -a \left(\frac{\int e^{-\operatorname{csch}^{-1}(a+bx)} \log \left(1 - \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{1-\sqrt{a^2+1}} \right) de^{\operatorname{csch}^{-1}(a+bx)}}{a} + \frac{\int e^{-\operatorname{csch}^{-1}(a+bx)} \log \left(1 - \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{\sqrt{a^2+1}+1} \right) de^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) \\
& i \left(-2i \left(-\frac{1}{4} \operatorname{PolyLog} \left(2, e^{2\operatorname{csch}^{-1}(a+bx)} \right) - \frac{1}{2} \operatorname{csch}^{-1}(a+bx) \log \left(1 - e^{2\operatorname{csch}^{-1}(a+bx)} \right) \right) - \frac{1}{2} i \operatorname{csch}^{-1}(a+bx)^2 \right) \\
& \quad \downarrow \text{2838} \\
& -a \left(-\frac{\operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{1-\sqrt{a^2+1}} \right)}{a} - \frac{\operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{\sqrt{a^2+1}+1} \right)}{a} - \frac{\operatorname{csch}^{-1}(a+bx) \log \left(1 - \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{1-\sqrt{a^2+1}} \right)}{a} \right) \\
& i \left(-2i \left(-\frac{1}{4} \operatorname{PolyLog} \left(2, e^{2\operatorname{csch}^{-1}(a+bx)} \right) - \frac{1}{2} \operatorname{csch}^{-1}(a+bx) \log \left(1 - e^{2\operatorname{csch}^{-1}(a+bx)} \right) \right) - \frac{1}{2} i \operatorname{csch}^{-1}(a+bx)^2 \right)
\end{aligned}$$

input `Int[ArcCsch[a + b*x]/x, x]`

output `-(a*(ArcCsch[a + b*x]^2/(2*a) - (ArcCsch[a + b*x]*Log[1 - (a*E^ArcCsch[a + b*x])/(1 - Sqrt[1 + a^2])])/a - (ArcCsch[a + b*x]*Log[1 - (a*E^ArcCsch[a + b*x])/(1 + Sqrt[1 + a^2])])/a - PolyLog[2, (a*E^ArcCsch[a + b*x])/(1 - Sqrt[1 + a^2])]/a - PolyLog[2, (a*E^ArcCsch[a + b*x])/(1 + Sqrt[1 + a^2])]/a)) + I*((-1/2*I)*ArcCsch[a + b*x]^2 - (2*I)*(-1/2*(ArcCsch[a + b*x]*Log[1 - E^(2*ArcCsch[a + b*x])]) - PolyLog[2, E^(2*ArcCsch[a + b*x])]/4))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4199 `Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi))]/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 6095 `Int[(Cosh[(c_) + (d_)*(x_)]*((e_) + (f_)*(x_))^(m_))/((a_) + (b_)*Sinh[(c_) + (d_)*(x_)]), x_Symbol] := Simp[-(e + f*x)^(m + 1)/(b*f*(m + 1)), x] + (Int[(e + f*x)^m*(E^(c + d*x)/(a - Rt[a^2 + b^2, 2] + b*E^(c + d*x))), x] + Int[(e + f*x)^m*(E^(c + d*x)/(a + Rt[a^2 + b^2, 2] + b*E^(c + d*x))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[a^2 + b^2, 0]`

rule 6103 `Int[(Coth[(c_) + (d_)*(x_)]^(n_))*((e_) + (f_)*(x_))^(m_))/((a_) + (b_)*Sinh[(c_) + (d_)*(x_)]), x_Symbol] := Simp[1/a Int[(e + f*x)^m*Coth[c + d*x]^n, x], x] - Simp[b/a Int[(e + f*x)^m*Cosh[c + d*x]*(Coth[c + d*x])^(n - 1)/(a + b*Sinh[c + d*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

rule 6130

```
Int[((e_.) + (f_.)*(x_)^(m_.)*(F_)[(c_.) + (d_.)*(x_)]^(n_.)*(G_)[(c_.) +
(d_.)*(x_)]^(p_.))/(Csch[(c_.) + (d_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := I
nt[(e + f*x)^m*Sinh[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Sinh[c + d*x
])), x] /; FreeQ[{a, b, c, d, e, f}, x] && HyperbolicQ[F] && HyperbolicQ[G]
&& IntegerQ[m, n, p]
```

rule 6876

```
Int[((a_.) + ArcCsch[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Simp[-(d^(m + 1))^(−1) Subst[Int[(a + b*x)^p*Csch[x]*C
oth[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Maple [F]

$$\int \frac{\operatorname{arccsch}(bx + a)}{x} dx$$

input

```
int(arccsch(b*x+a)/x,x)
```

output

```
int(arccsch(b*x+a)/x,x)
```

Fricas [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arcsch}(bx + a)}{x} dx$$

input

```
integrate(arccsch(b*x+a)/x,x, algorithm="fricas")
```

output

```
integral(arccsch(b*x + a)/x, x)
```

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{acsch}(a + bx)}{x} dx$$

input `integrate(acsch(b*x+a)/x,x)`

output `Integral(acsch(a + b*x)/x, x)`

Maxima [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arcsch}(bx + a)}{x} dx$$

input `integrate(arccsch(b*x+a)/x,x, algorithm="maxima")`

output `integrate(arccsch(b*x + a)/x, x)`

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arcsch}(bx + a)}{x} dx$$

input `integrate(arccsch(b*x+a)/x,x, algorithm="giac")`

output `integrate(arccsch(b*x + a)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{a+bx}\right)}{x} dx$$

input `int(asinh(1/(a + b*x))/x,x)`output `int(asinh(1/(a + b*x))/x, x)`**Reduce [F]**

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{acsch}(bx + a)}{x} dx$$

input `int(acsch(b*x+a)/x,x)`output `int(acsch(a + b*x)/x,x)`

3.5 $\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^2} dx$

Optimal result	70
Mathematica [B] (verified)	70
Rubi [A] (verified)	71
Maple [B] (verified)	74
Fricas [B] (verification not implemented)	74
Sympy [F]	75
Maxima [F]	75
Giac [F]	76
Mupad [F(-1)]	76
Reduce [F]	76

Optimal result

Integrand size = 10, antiderivative size = 63

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^2} dx = -\frac{b\operatorname{csch}^{-1}(a+bx)}{a} - \frac{\operatorname{csch}^{-1}(a+bx)}{x} + \frac{2b\operatorname{arctanh}\left(\frac{a+\tanh\left(\frac{1}{2}\operatorname{csch}^{-1}(a+bx)\right)}{\sqrt{1+a^2}}\right)}{a\sqrt{1+a^2}}$$

output `-b*arccsch(b*x+a)/a-arccsch(b*x+a)/x+2*b*arctanh((a+tanh(1/2*arccsch(b*x+a)))/(a^2+1)^(1/2))/a/(a^2+1)^(1/2)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 141 vs. 2(63) = 126.

Time = 0.10 (sec) , antiderivative size = 141, normalized size of antiderivative = 2.24

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^2} dx = -\frac{\operatorname{csch}^{-1}(a+bx)}{x} - \frac{b\left(\sqrt{1+a^2}\operatorname{arcsinh}\left(\frac{1}{a+bx}\right) + \log(x) - \log\left(1+a^2+abx+a\sqrt{1+a^2}\sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}} + \sqrt{1+a^2}bx\sqrt{1+a^2}\right)\right)}{a\sqrt{1+a^2}}$$

input `Integrate[ArcCsch[a + b*x]/x^2,x]`

output
$$-(\text{ArcCsch}[a + b*x]/x) - (b(\text{Sqrt}[1 + a^2]*\text{ArcSinh}[(a + b*x)^{-1}] + \text{Log}[x] - \text{Log}[1 + a^2 + a*b*x + a*\text{Sqrt}[1 + a^2]*\text{Sqrt}[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2] + \text{Sqrt}[1 + a^2]*b*x*\text{Sqrt}[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]]))/(a*\text{Sqrt}[1 + a^2])$$

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.14, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {6876, 5992, 3042, 4270, 3042, 3139, 1083, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\text{csch}^{-1}(a + bx)}{x^2} dx \\ & \quad \downarrow 6876 \\ & -b \int \frac{(a + bx)^2 \sqrt{1 + \frac{1}{(a+bx)^2}} \text{csch}^{-1}(a + bx)}{b^2 x^2} d\text{csch}^{-1}(a + bx) \\ & \quad \downarrow 5992 \\ & -b \left(\int -\frac{1}{bx} d\text{csch}^{-1}(a + bx) + \frac{\text{csch}^{-1}(a + bx)}{bx} \right) \\ & \quad \downarrow 3042 \\ & -b \left(\frac{\text{csch}^{-1}(a + bx)}{bx} + \int \frac{1}{a - i \csc(i \text{csch}^{-1}(a + bx))} d\text{csch}^{-1}(a + bx) \right) \\ & \quad \downarrow 4270 \\ & -b \left(-\frac{\int \frac{1}{1 - \frac{a}{a+bx}} d\text{csch}^{-1}(a + bx)}{a} + \frac{\text{csch}^{-1}(a + bx)}{a} + \frac{\text{csch}^{-1}(a + bx)}{bx} \right) \\ & \quad \downarrow 3042 \end{aligned}$$

$$-b \left(\frac{\int \frac{1}{ia \sin(i \operatorname{csch}^{-1}(a+bx)) + 1} d \operatorname{csch}^{-1}(a+bx)}{a} + \frac{\operatorname{csch}^{-1}(a+bx)}{a} + \frac{\operatorname{csch}^{-1}(a+bx)}{bx} \right)$$

↓ 3139

$$-b \left(\frac{2 \int \frac{1}{-\tanh^2\left(\frac{1}{2} \operatorname{csch}^{-1}(a+bx)\right) - 2a \tanh\left(\frac{1}{2} \operatorname{csch}^{-1}(a+bx)\right) + 1} d \tanh\left(\frac{1}{2} \operatorname{csch}^{-1}(a+bx)\right)}{a} + \frac{\operatorname{csch}^{-1}(a+bx)}{a} + \frac{\operatorname{csch}^{-1}(a+bx)}{bx} \right)$$

↓ 1083

$$-b \left(\frac{4 \int \frac{1}{4(a^2+1) - \left(-2a - 2 \tanh\left(\frac{1}{2} \operatorname{csch}^{-1}(a+bx)\right)\right)^2} d\left(-2a - 2 \tanh\left(\frac{1}{2} \operatorname{csch}^{-1}(a+bx)\right)\right)}{a} + \frac{\operatorname{csch}^{-1}(a+bx)}{a} + \frac{\operatorname{csch}^{-1}(a+bx)}{bx} \right)$$

↓ 219

$$-b \left(\frac{2 \operatorname{arctanh}\left(\frac{-2 \tanh\left(\frac{1}{2} \operatorname{csch}^{-1}(a+bx)\right) - 2a}{2\sqrt{a^2+1}}\right)}{a\sqrt{a^2+1}} + \frac{\operatorname{csch}^{-1}(a+bx)}{a} + \frac{\operatorname{csch}^{-1}(a+bx)}{bx} \right)$$

input `Int[ArcCsch[a + b*x]/x^2,x]`

output `-(b*(ArcCsch[a + b*x]/a + ArcCsch[a + b*x]/(b*x) + (2*ArcTanh[(-2*a - 2*
Tanh[ArcCsch[a + b*x]/2])/(2*
Sqrt[1 + a^2])])/(a*Sqrt[1 + a^2])))`

Definitions of rubi rules used

- rule 219 $\text{Int}[(a_ + (b_ \cdot x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[-b, 2])) \cdot \text{ArcTanh}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 1083 $\text{Int}[(a_ + (b_ \cdot x_) + (c_ \cdot x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[-2 \ \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4 \cdot a \cdot c - x^2, x], x], x, b + 2 \cdot c \cdot x], x] /; \text{FreeQ}\{a, b, c\}, x]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3139 $\text{Int}[(a_ + (b_ \cdot \sin[(c_ \cdot x_) + (d_ \cdot x_)])^{-1}, x_Symbol] \rightarrow \text{With}\{e = \text{FreeFactors}[\text{Tan}[(c + d \cdot x)/2], x]\}, \text{Simp}[2 \cdot (e/d) \ \text{Subst}[\text{Int}[1/(a + 2 \cdot b \cdot e \cdot x + a \cdot e^2 \cdot x^2), x], x, \text{Tan}[(c + d \cdot x)/2]/e], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$
- rule 4270 $\text{Int}[(\text{csc}[(c_ \cdot x_) + (d_ \cdot x_)] \cdot (b_ \cdot x_) + (a_))^{-1}, x_Symbol] \rightarrow \text{Simp}[x/a, x] - \text{Simp}[1/a \ \text{Int}[1/(1 + (a/b) \cdot \text{Sin}[c + d \cdot x]), x], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$
- rule 5992 $\text{Int}[\text{Coth}[(c_ \cdot x_) + (d_ \cdot x_)] \cdot \text{Csch}[(c_ \cdot x_) + (d_ \cdot x_)] \cdot (\text{Csch}[(c_ \cdot x_) + (d_ \cdot x_)] \cdot (b_ \cdot x_) + (a_))^{(n_ \cdot x_)} \cdot ((e_ \cdot x_) + (f_ \cdot x_))^{(m_ \cdot x_)}, x_Symbol] \rightarrow \text{Simp}[(-(e + f \cdot x)^m \cdot ((a + b \cdot \text{Csch}[c + d \cdot x])^{(n + 1)} / (b \cdot d \cdot (n + 1))), x] + \text{Simp}[f \cdot (m / (b \cdot d \cdot (n + 1))) \ \text{Int}[(e + f \cdot x)^{(m - 1)} \cdot (a + b \cdot \text{Csch}[c + d \cdot x])^{(n + 1)}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{NeQ}[n, -1]$
- rule 6876 $\text{Int}[(a_ \cdot \text{ArcCsch}[(c_ \cdot x_) + (d_ \cdot x_)] \cdot (b_ \cdot x_))^{(p_ \cdot x_)} \cdot ((e_ \cdot x_) + (f_ \cdot x_))^{(m_ \cdot x_)}, x_Symbol] \rightarrow \text{Simp}[-(d^{(m + 1)})^{-1} \ \text{Subst}[\text{Int}[(a + b \cdot x)^p \cdot \text{Csch}[x] \cdot \text{Coth}[x] \cdot (d \cdot e - c \cdot f + f \cdot \text{Csch}[x])^m, x], x, \text{ArcCsch}[c + d \cdot x]], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{IntegerQ}[m]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 126 vs. $2(57) = 114$.

Time = 0.47 (sec) , antiderivative size = 127, normalized size of antiderivative = 2.02

method	result
derivativedivides	$b \left(-\frac{\operatorname{arccsch}(bx+a)}{bx} - \frac{\sqrt{(bx+a)^2+1} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{(bx+a)^2+1}}\right) \sqrt{a^2+1} - \ln\left(\frac{2\sqrt{a^2+1}\sqrt{(bx+a)^2+1}+2(bx+a)a+2}{bx}\right) \right)}{\sqrt{\frac{(bx+a)^2+1}{(bx+a)^2}} (bx+a)a\sqrt{a^2+1}} \right)$
default	$b \left(-\frac{\operatorname{arccsch}(bx+a)}{bx} - \frac{\sqrt{(bx+a)^2+1} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{(bx+a)^2+1}}\right) \sqrt{a^2+1} - \ln\left(\frac{2\sqrt{a^2+1}\sqrt{(bx+a)^2+1}+2(bx+a)a+2}{bx}\right) \right)}{\sqrt{\frac{(bx+a)^2+1}{(bx+a)^2}} (bx+a)a\sqrt{a^2+1}} \right)$
parts	$-\frac{\operatorname{arccsch}(bx+a)}{x} - \frac{b\sqrt{b^2x^2+2bxa+a^2+1} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{b^2x^2+2bxa+a^2+1}}\right) \sqrt{a^2+1} - \ln\left(\frac{2a^2+2+2bxa+2\sqrt{a^2+1}\sqrt{b^2x^2+2bxa+a^2+1}}{x}\right) \right)}{\sqrt{\frac{b^2x^2+2bxa+a^2+1}{(bx+a)^2}} (bx+a)a\sqrt{a^2+1}}$

input `int(arccsch(b*x+a)/x^2,x,method=_RETURNVERBOSE)`

output `b*(-1/b/x*arccsch(b*x+a)-((b*x+a)^2+1)^(1/2)*(arctanh(1/((b*x+a)^2+1)^(1/2)))*(a^2+1)^(1/2)-ln(2*((a^2+1)^(1/2)*((b*x+a)^2+1)^(1/2)+(b*x+a)*a+1)/b/x))/(((b*x+a)^2+1)/(b*x+a)^2)^(1/2)/(b*x+a)/a/(a^2+1)^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 343 vs. $2(57) = 114$.

Time = 0.12 (sec) , antiderivative size = 343, normalized size of antiderivative = 5.44

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^2} dx =$$

$$(a^2+1)bx \log\left(-bx+(bx+a)\sqrt{\frac{b^2x^2+2abx+a^2+1}{b^2x^2+2abx+a^2}}-a+1\right) - (a^2+1)bx \log\left(-bx+(bx+a)\sqrt{\frac{b^2x^2+2a}{b^2x^2+2}}$$

input `integrate(arccsch(b*x+a)/x^2,x, algorithm="fricas")`

output

```

-((a^2 + 1)*b*x*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a + 1) - (a^2 + 1)*b*x*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a - 1) - sqrt(a^2 + 1)*b*x*log(-(a^2*b*x + a^3 + (a*b*x + a^2 + (a*b*x + a^2)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)*sqrt(a^2 + 1) + (a^3 + (a^2 + 1)*b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) + a)/x) + (a^3 + a)*log(((b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)))/((a^3 + a)*x)

```

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{acsch}(a + bx)}{x^2} dx$$

input

```
integrate(acsch(b*x+a)/x**2,x)
```

output

```
Integral(acsch(a + b*x)/x**2, x)
```

Maxima [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{arcsch}(bx + a)}{x^2} dx$$

input

```
integrate(arccsch(b*x+a)/x^2,x, algorithm="maxima")
```

output

```

-1/2*I*b*(log(I*(b^2*x + a*b)/b + 1) - log(-I*(b^2*x + a*b)/b + 1))/(a^2 + 1) - b*log(x)/(a^3 + a) - 1/2*(a^2*b*x*log(b^2*x^2 + 2*a*b*x + a^2 + 1) - 2*(a^3 + (a^2*b + b)*x + a)*log(b*x + a) + 2*(a^3 + a)*log(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) + 1))/((a^3 + a)*x) - integrate((b^2*x + a*b)/(b^2*x^3 + 2*a*b*x^2 + (a^2 + 1)*x + (b^2*x^3 + 2*a*b*x^2 + (a^2 + 1)*x)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)), x)

```

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{arcsch}(bx + a)}{x^2} dx$$

input `integrate(arccsch(b*x+a)/x^2,x, algorithm="giac")`

output `integrate(arccsch(b*x + a)/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{a+bx}\right)}{x^2} dx$$

input `int(asinh(1/(a + b*x))/x^2,x)`

output `int(asinh(1/(a + b*x))/x^2, x)`

Reduce [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{acsch}(bx + a)}{x^2} dx$$

input `int(acsch(b*x+a)/x^2,x)`

output `int(acsch(a + b*x)/x**2,x)`

3.6 $\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^3} dx$

Optimal result	77
Mathematica [A] (verified)	78
Rubi [A] (verified)	78
Maple [B] (verified)	83
Fricas [B] (verification not implemented)	84
Sympy [F]	84
Maxima [F]	85
Giac [F]	85
Mupad [F(-1)]	85
Reduce [F]	86

Optimal result

Integrand size = 10, antiderivative size = 114

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^3} dx = \frac{b(a+bx)\sqrt{1+\frac{1}{(a+bx)^2}}}{2a(1+a^2)x} + \frac{b^2\operatorname{csch}^{-1}(a+bx)}{2a^2} - \frac{\operatorname{csch}^{-1}(a+bx)}{2x^2} - \frac{(1+2a^2)b^2\operatorname{arctanh}\left(\frac{a+\tanh\left(\frac{1}{2}\operatorname{csch}^{-1}(a+bx)\right)}{\sqrt{1+a^2}}\right)}{a^2(1+a^2)^{3/2}}$$

output

```
1/2*b*(b*x+a)*(1+1/(b*x+a)^2)^(1/2)/a/(a^2+1)/x+1/2*b^2*arccsch(b*x+a)/a^2
-1/2*arccsch(b*x+a)/x^2-(2*a^2+1)*b^2*arctanh((a+tanh(1/2*arccsch(b*x+a)))/
(a^2+1)^(1/2))/a^2/(a^2+1)^(3/2)
```

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.93

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^3} dx = \frac{1}{2} \left(\frac{b(a+bx) \sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}}}{a(1+a^2)x} - \frac{\operatorname{csch}^{-1}(a+bx)}{x^2} \right. \\ \left. + \frac{b^2 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}{a^2} + \frac{(1+2a^2)b^2 \log(x)}{a^2(1+a^2)^{3/2}} \right. \\ \left. - \frac{(1+2a^2)b^2 \log\left(1+a^2+abx+a\sqrt{1+a^2}\sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}} + \sqrt{1+a^2}bx\sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}}\right)}{a^2(1+a^2)^{3/2}} \right)$$

input `Integrate[ArcCsch[a + b*x]/x^3,x]`

output `((b*(a + b*x)*Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2])/(a*(1 + a^2)*x) - ArcCsch[a + b*x]/x^2 + (b^2*ArcSinh[(a + b*x)^(-1)]/a^2 + ((1 + 2*a^2)*b^2*Log[x]))/(a^2*(1 + a^2)^(3/2)) - ((1 + 2*a^2)*b^2*Log[1 + a^2 + a*b*x + a*Sqrt[1 + a^2]*Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2] + Sqrt[1 + a^2]*b*x*Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]))/(a^2*(1 + a^2)^(3/2)))/2`

Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.25, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 1.600$, Rules used = {6876, 25, 5992, 3042, 4272, 25, 3042, 4407, 26, 3042, 26, 4318, 3042, 3139, 1083, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^3} dx$$

↓ 6876

$$\begin{aligned}
& -b^2 \int \frac{(a+bx)^2 \sqrt{1 + \frac{1}{(a+bx)^2}} \operatorname{csch}^{-1}(a+bx)}{b^3 x^3} d\operatorname{csch}^{-1}(a+bx) \\
& \quad \downarrow 25 \\
& b^2 \int -\frac{(a+bx)^2 \sqrt{1 + \frac{1}{(a+bx)^2}} \operatorname{csch}^{-1}(a+bx)}{b^3 x^3} d\operatorname{csch}^{-1}(a+bx) \\
& \quad \downarrow 5992 \\
& -b^2 \left(\frac{\operatorname{csch}^{-1}(a+bx)}{2b^2 x^2} - \frac{1}{2} \int \frac{1}{b^2 x^2} d\operatorname{csch}^{-1}(a+bx) \right) \\
& \quad \downarrow 3042 \\
& -b^2 \left(\frac{\operatorname{csch}^{-1}(a+bx)}{2b^2 x^2} - \frac{1}{2} \int \frac{1}{(a - i \csc(i \operatorname{csch}^{-1}(a+bx)))^2} d\operatorname{csch}^{-1}(a+bx) \right) \\
& \quad \downarrow 4272 \\
& -b^2 \left(\frac{1}{2} \left(\frac{\int \frac{a^2 + (a+bx)a+1}{bx} d\operatorname{csch}^{-1}(a+bx)}{a(a^2+1)} - \frac{(a+bx) \sqrt{\frac{1}{(a+bx)^2} + 1}}{a(a^2+1)bx} \right) + \frac{\operatorname{csch}^{-1}(a+bx)}{2b^2 x^2} \right) \\
& \quad \downarrow 25 \\
& -b^2 \left(\frac{1}{2} \left(-\frac{\int -\frac{a^2 + (a+bx)a+1}{bx} d\operatorname{csch}^{-1}(a+bx)}{a(a^2+1)} - \frac{\sqrt{\frac{1}{(a+bx)^2} + 1}(a+bx)}{a(a^2+1)bx} \right) + \frac{\operatorname{csch}^{-1}(a+bx)}{2b^2 x^2} \right) \\
& \quad \downarrow 3042 \\
& -b^2 \left(\frac{\operatorname{csch}^{-1}(a+bx)}{2b^2 x^2} + \frac{1}{2} \left(-\frac{\sqrt{\frac{1}{(a+bx)^2} + 1}(a+bx)}{a(a^2+1)bx} - \frac{\int \frac{a^2 + i \csc(i \operatorname{csch}^{-1}(a+bx))a+1}{a - i \csc(i \operatorname{csch}^{-1}(a+bx))} d\operatorname{csch}^{-1}(a+bx)}{a(a^2+1)} \right) \right) \\
& \quad \downarrow 4407 \\
& -b^2 \left(\frac{\operatorname{csch}^{-1}(a+bx)}{2b^2 x^2} + \frac{1}{2} \left(-\frac{\sqrt{\frac{1}{(a+bx)^2} + 1}(a+bx)}{a(a^2+1)bx} - \frac{(a^2+1)\operatorname{csch}^{-1}(a+bx)}{a} + \frac{i(2a^2+1) \int \frac{i(a+bx)}{bx} d\operatorname{csch}^{-1}(a+bx)}{a(a^2+1)} \right) \right) \\
& \quad \downarrow 26
\end{aligned}$$

$$-b^2 \left(\frac{1}{2} \left(-\frac{(2a^2+1) \int -\frac{a+bx}{bx} d\operatorname{csch}^{-1}(a+bx) + (a^2+1)\operatorname{csch}^{-1}(a+bx)}{a(a^2+1)} - \frac{\sqrt{\frac{1}{(a+bx)^2} + 1}(a+bx)}{a(a^2+1)bx} \right) + \frac{\operatorname{csch}^{-1}(a+bx)}{2b^2x^2} \right)$$

↓ 3042

$$-b^2 \left(\frac{\operatorname{csch}^{-1}(a+bx)}{2b^2x^2} + \frac{1}{2} \left(-\frac{\sqrt{\frac{1}{(a+bx)^2} + 1}(a+bx)}{a(a^2+1)bx} - \frac{(a^2+1)\operatorname{csch}^{-1}(a+bx)}{a} + \frac{(2a^2+1) \int \frac{i \operatorname{csc}(i\operatorname{csch}^{-1}(a+bx))}{a-i \operatorname{csc}(i\operatorname{csch}^{-1}(a+bx))} d\operatorname{csch}^{-1}}{a(a^2+1)} \right) \right)$$

↓ 26

$$-b^2 \left(\frac{\operatorname{csch}^{-1}(a+bx)}{2b^2x^2} + \frac{1}{2} \left(-\frac{\sqrt{\frac{1}{(a+bx)^2} + 1}(a+bx)}{a(a^2+1)bx} - \frac{(a^2+1)\operatorname{csch}^{-1}(a+bx)}{a} + \frac{i(2a^2+1) \int \frac{\operatorname{csc}(i\operatorname{csch}^{-1}(a+bx))}{a-i \operatorname{csc}(i\operatorname{csch}^{-1}(a+bx))} d\operatorname{csch}^{-1}}{a(a^2+1)} \right) \right)$$

↓ 4318

$$-b^2 \left(\frac{1}{2} \left(-\frac{(a^2+1)\operatorname{csch}^{-1}(a+bx)}{a} - \frac{(2a^2+1) \int \frac{1}{1-\frac{a}{a+bx}} d\operatorname{csch}^{-1}(a+bx)}{a(a^2+1)} - \frac{\sqrt{\frac{1}{(a+bx)^2} + 1}(a+bx)}{a(a^2+1)bx} \right) + \frac{\operatorname{csch}^{-1}(a+bx)}{2b^2x^2} \right)$$

↓ 3042

$$-b^2 \left(\frac{\operatorname{csch}^{-1}(a+bx)}{2b^2x^2} + \frac{1}{2} \left(-\frac{\sqrt{\frac{1}{(a+bx)^2} + 1}(a+bx)}{a(a^2+1)bx} - \frac{(a^2+1)\operatorname{csch}^{-1}(a+bx)}{a} - \frac{(2a^2+1) \int \frac{1}{ia \sin(i\operatorname{csch}^{-1}(a+bx))+1} d\operatorname{csch}^{-1}}{a(a^2+1)} \right) \right)$$

↓ 3139

$$-b^2 \left(\frac{1}{2} \left(-\frac{(a^2+1)\operatorname{csch}^{-1}(a+bx)}{a} - \frac{2(2a^2+1) \int \frac{1}{-\tanh^2(\frac{1}{2}\operatorname{csch}^{-1}(a+bx))-2a \tanh(\frac{1}{2}\operatorname{csch}^{-1}(a+bx))+1} d \tanh(\frac{1}{2}\operatorname{csch}^{-1}(a+bx))}{a(a^2+1)} - \frac{\sqrt{\frac{1}{(a+bx)^2} + 1}(a+bx)}{a(a^2+1)bx} \right) + \frac{\operatorname{csch}^{-1}(a+bx)}{2b^2x^2} \right)$$

↓ 1083

$$-b^2 \left(\frac{1}{2} \left(-\frac{4(2a^2+1) \int \frac{1}{4(a^2+1) - (-2a-2 \tanh(\frac{1}{2} \operatorname{csch}^{-1}(a+bx)))^2} dx (-2a-2 \tanh(\frac{1}{2} \operatorname{csch}^{-1}(a+bx)))}{a(a^2+1)} + \frac{(a^2+1) \operatorname{csch}^{-1}(a+bx)}{a} - \sqrt{\frac{1}{(a+bx)^2} + 1} (a+bx) \right) \right)$$

↓ 219

$$-b^2 \left(\frac{1}{2} \left(-\frac{2(2a^2+1) \operatorname{arctanh}\left(\frac{-2 \tanh(\frac{1}{2} \operatorname{csch}^{-1}(a+bx)) - 2a}{2\sqrt{a^2+1}}\right)}{a\sqrt{a^2+1}} + \frac{(a^2+1) \operatorname{csch}^{-1}(a+bx)}{a} - \frac{\sqrt{\frac{1}{(a+bx)^2} + 1} (a+bx)}{a(a^2+1)bx} \right) \right) + \frac{\operatorname{csch}^{-1}(a+bx)}{2}$$

input `Int[ArcCsch[a + b*x]/x^3,x]`

output `-(b^2*(ArcCsch[a + b*x]/(2*b^2*x^2) + (-(((a + b*x)*Sqrt[1 + (a + b*x)^(-2)]))/(a*(1 + a^2)*b*x)) - (((1 + a^2)*ArcCsch[a + b*x])/a + (2*(1 + 2*a^2)*ArcTanh[(-2*a - 2*Tanh[ArcCsch[a + b*x]/2])/(2*Sqrt[1 + a^2])])/(a*Sqrt[1 + a^2]))/(a*(1 + a^2)))/2)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3139 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 4272 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.) + (a_)^(n_), x_Symbol] := Simp[b^2*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n + 1)/(a*d*(n + 1)*(a^2 - b^2))), x] + Simp[1/(a*(n + 1)*(a^2 - b^2)) Int[(a + b*Csc[c + d*x])^(n + 1)*Simp[(a^2 - b^2)*(n + 1) - a*b*(n + 1)*Csc[c + d*x] + b^2*(n + 2)*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4318 `Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_), x_Symbol] := Simp[1/b Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4407 `Int[(csc[(e_.) + (f_.)*(x_)])*(d_.) + (c_))/(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_), x_Symbol] := Simp[c*(x/a), x] - Simp[(b*c - a*d)/a Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 5992 `Int[Coth[(c_.) + (d_.)*(x_)])*Csch[(c_.) + (d_.)*(x_)])*(Csch[(c_.) + (d_.)*(x_)])*(b_.) + (a_)^(n_.)*((e_.) + (f_.)*(x_)^(m_.), x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

rule 6876

```
Int[((a_.) + ArcCsch[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Csch[x]*Coth[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 317 vs. 2(100) = 200.

Time = 0.49 (sec) , antiderivative size = 318, normalized size of antiderivative = 2.79

method	result
parts	$-\frac{\operatorname{arccsch}(bx+a)}{2x^2} + \frac{b\sqrt{b^2x^2+2bxa+a^2+1} \left((a^2+1)^{\frac{3}{2}} \operatorname{arctanh}\left(\frac{1}{\sqrt{b^2x^2+2bxa+a^2+1}}\right) a^2bx - 2 \ln\left(\frac{2a^2+2+2bxa+2\sqrt{a^2+1}}{\dots}\right) \right)}{\dots}$
derivativedivides	$b^2 \left(-\frac{\operatorname{arccsch}(bx+a)}{2b^2x^2} - \frac{\sqrt{(bx+a)^2+1} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{(bx+a)^2+1}}\right) (a^2+1)^{\frac{3}{2}} a^3 - \operatorname{arctanh}\left(\frac{1}{\sqrt{(bx+a)^2+1}}\right) (a^2+1)^{\frac{3}{2}} a^2 \right)}{\dots} \right)$
default	$b^2 \left(-\frac{\operatorname{arccsch}(bx+a)}{2b^2x^2} - \frac{\sqrt{(bx+a)^2+1} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{(bx+a)^2+1}}\right) (a^2+1)^{\frac{3}{2}} a^3 - \operatorname{arctanh}\left(\frac{1}{\sqrt{(bx+a)^2+1}}\right) (a^2+1)^{\frac{3}{2}} a^2 \right)}{\dots} \right)$

input

```
int(arccsch(b*x+a)/x^3,x,method=_RETURNVERBOSE)
```

output

```
-1/2*arccsch(b*x+a)/x^2+1/2*b*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)*((a^2+1)^(3/2)*arctanh(1/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))*a^2*b*x-2*ln(2*(b*x*a+(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a^2+1)/x)*a^4*b*x+b*arctanh(1/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))*x*(a^2+1)^(3/2)+(a^2+1)^(3/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)*a-3*ln(2*(b*x*a+(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a^2+1)/x)*a^2*b*x-b*ln(2*(b*x*a+(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a^2+1)/x)*x)/((b^2*x^2+2*a*b*x+a^2+1)/(b*x+a)^2)^(1/2)/(b*x+a)/a^2/(a^2+1)^(5/2)/x
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 461 vs. $2(100) = 200$.

Time = 0.13 (sec) , antiderivative size = 461, normalized size of antiderivative = 4.04

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^3} dx$$

$$(2a^2 + 1)\sqrt{a^2 + 1}b^2x^2 \log \left(-\frac{a^2bx + a^3 - (abx + a^2 + (abx + a^2)\sqrt{\frac{b^2x^2 + 2abx + a^2 + 1}{b^2x^2 + 2abx + a^2} + 1})\sqrt{a^2 + 1} + (a^3 + (a^2 + 1)bx + a)\sqrt{\frac{b^2x^2 + 2abx + a^2 + 1}{b^2x^2 + 2abx + a^2}}}{x} \right)$$

input `integrate(arccsch(b*x+a)/x^3,x, algorithm="fricas")`

output

```
1/2*((2*a^2 + 1)*sqrt(a^2 + 1)*b^2*x^2*log(-(a^2*b*x + a^3 - (a*b*x + a^2
+ (a*b*x + a^2)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^
2)) + 1)*sqrt(a^2 + 1) + (a^3 + (a^2 + 1)*b*x + a)*sqrt((b^2*x^2 + 2*a*b*x
+ a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) + a)/x) + (a^4 + 2*a^2 + 1)*b^2*x^2
*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x
+ a^2)) - a + 1) - (a^4 + 2*a^2 + 1)*b^2*x^2*log(-b*x + (b*x + a)*sqrt((
b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a - 1) + (a^3 +
a)*b^2*x^2 - (a^6 + 2*a^4 + a^2)*log(((b*x + a)*sqrt((b^2*x^2 + 2*a*b*x +
a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)) + ((a^3 + a)*b^2*x^2 +
(a^4 + a^2)*b*x)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x +
a^2)))/((a^6 + 2*a^4 + a^2)*x^2)
```

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{acsch}(a + bx)}{x^3} dx$$

input `integrate(acsch(b*x+a)/x**3,x)`

output `Integral(acsch(a + b*x)/x**3, x)`

Maxima [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{arcsch}(bx + a)}{x^3} dx$$

input `integrate(arccsch(b*x+a)/x^3,x, algorithm="maxima")`

output `1/2*I*a*b^2*(log(I*(b^2*x + a*b)/b + 1) - log(-I*(b^2*x + a*b)/b + 1))/(a^4 + 2*a^2 + 1) + 1/2*(3*a^2*b^2 + b^2)*log(x)/(a^6 + 2*a^4 + a^2) + 1/4*((a^4*b^2 - a^2*b^2)*x^2*log(b^2*x^2 + 2*a*b*x + a^2 + 1) + 2*(a^3*b + a*b)*x + 2*(a^6 + 2*a^4 - (a^4*b^2 + 2*a^2*b^2 + b^2)*x^2 + a^2)*log(b*x + a) - 2*(a^6 + 2*a^4 + a^2)*log(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) + 1))/((a^6 + 2*a^4 + a^2)*x^2) - integrate(1/2*(b^2*x + a*b)/(b^2*x^4 + 2*a*b*x^3 + (a^2 + 1)*x^2 + (b^2*x^4 + 2*a*b*x^3 + (a^2 + 1)*x^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)), x)`

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{arcsch}(bx + a)}{x^3} dx$$

input `integrate(arccsch(b*x+a)/x^3,x, algorithm="giac")`

output `integrate(arccsch(b*x + a)/x^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{a+bx}\right)}{x^3} dx$$

input `int(asinh(1/(a + b*x))/x^3,x)`

output `int(asinh(1/(a + b*x))/x^3, x)`

Reduce [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{acsch}(bx + a)}{x^3} dx$$

input `int(acsch(b*x+a)/x^3,x)`

output `int(acsch(a + b*x)/x**3,x)`

3.7 $\int (e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$

Optimal result	88
Mathematica [C] (warning: unable to verify)	89
Rubi [A] (verified)	90
Maple [F]	92
Fricas [F]	93
Sympy [F]	93
Maxima [F]	93
Giac [F]	94
Mupad [F(-1)]	95
Reduce [F]	95

Optimal result

Integrand size = 20, antiderivative size = 501

$$\begin{aligned}
& \int (e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx \\
&= \frac{b^2 f^2 (de - cf)x}{d^3} + \frac{b^2 f^3 (c + dx)^2}{12d^4} - \frac{bf^3 (c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{3d^4} \\
&+ \frac{3bf(de - cf)^2 (c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{d^4} \\
&+ \frac{bf^2 (de - cf)(c + dx)^2 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{d^4} \\
&+ \frac{bf^3 (c + dx)^3 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{6d^4} \\
&- \frac{(de - cf)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4d^4 f} + \frac{(e + fx)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4f} \\
&- \frac{2bf^2 (de - cf) (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&+ \frac{4b(de - cf)^3 (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} - \frac{b^2 f^3 \log(c + dx)}{3d^4} \\
&+ \frac{3b^2 f (de - cf)^2 \log(c + dx)}{d^4} - \frac{b^2 f^2 (de - cf) \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&+ \frac{2b^2 (de - cf)^3 \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&+ \frac{b^2 f^2 (de - cf) \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&- \frac{2b^2 (de - cf)^3 \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4}
\end{aligned}$$

output

```

b^2*f^2*(-c*f+d*e)*x/d^3+1/12*b^2*f^3*(d*x+c)^2/d^4-1/3*b*f^3*(d*x+c)*(1+1
/(d*x+c)^2)^(1/2)*(a+b*arccsch(d*x+c))/d^4+3*b*f*(-c*f+d*e)^2*(d*x+c)*(1+1
/(d*x+c)^2)^(1/2)*(a+b*arccsch(d*x+c))/d^4+b*f^2*(-c*f+d*e)*(d*x+c)^2*(1+1
/(d*x+c)^2)^(1/2)*(a+b*arccsch(d*x+c))/d^4+1/6*b*f^3*(d*x+c)^3*(1+1/(d*x+c
)^2)^(1/2)*(a+b*arccsch(d*x+c))/d^4-1/4*(-c*f+d*e)^4*(a+b*arccsch(d*x+c))^
2/d^4/f+1/4*(f*x+e)^4*(a+b*arccsch(d*x+c))^2/f-2*b*f^2*(-c*f+d*e)*(a+b*arc
csch(d*x+c))*arctanh(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))/d^4+4*b*(-c*f+d*e)^3
*(a+b*arccsch(d*x+c))*arctanh(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))/d^4-1/3*b^2
*f^3*ln(d*x+c)/d^4+3*b^2*f*(-c*f+d*e)^2*ln(d*x+c)/d^4-b^2*f^2*(-c*f+d*e)*p
olylog(2,-1/(d*x+c)-(1+1/(d*x+c)^2)^(1/2))/d^4+2*b^2*(-c*f+d*e)^3*polylog(
2,-1/(d*x+c)-(1+1/(d*x+c)^2)^(1/2))/d^4+b^2*f^2*(-c*f+d*e)*polylog(2,1/(d*
x+c)+(1+1/(d*x+c)^2)^(1/2))/d^4-2*b^2*(-c*f+d*e)^3*polylog(2,1/(d*x+c)+(1+
1/(d*x+c)^2)^(1/2))/d^4

```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 9.98 (sec) , antiderivative size = 1429, normalized size of antiderivative = 2.85

$$\int (e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \text{Too large to display}$$

input

```
Integrate[(e + f*x)^3*(a + b*ArcCsch[c + d*x])^2,x]
```

output

```

a^2*e^3*x + (3*a^2*e^2*f*x^2)/2 + a^2*e*f^2*x^3 + (a^2*f^3*x^4)/4 + (a*b*(
3*x*(4*e^3 + 6*e^2*f*x + 4*e*f^2*x^2 + f^3*x^3)*ArcCsch[c + d*x] + (f*(c +
d*x)*Sqrt[(1 + c^2 + 2*c*d*x + d^2*x^2)/(c + d*x)^2]*((-2 + 13*c^2)*f^2 -
2*c*d*f*(15*e + 2*f*x) + d^2*(18*e^2 + 6*e*f*x + f^2*x^2)) - 3*c*(-4*d^3*
e^3 + 6*c*d^2*e^2*f - 4*c^2*d*e*f^2 + c^3*f^3)*ArcSinh[(c + d*x)^(-1)] + 6
*(2*d^3*e^3 - 6*c*d^2*e^2*f + (-1 + 6*c^2)*d*e*f^2 + c*(1 - 2*c^2)*f^3)*Lo
g[(c + d*x)*(1 + Sqrt[(1 + c^2 + 2*c*d*x + d^2*x^2)/(c + d*x)^2])]/d^4)/
6 - (b^2*e^3*(-(ArcCsch[c + d*x]*((c + d*x)*ArcCsch[c + d*x] - 2*Log[1 - E
^(-ArcCsch[c + d*x])]) + 2*Log[1 + E^(-ArcCsch[c + d*x])])) + 2*PolyLog[2,
-E^(-ArcCsch[c + d*x])]) - 2*PolyLog[2, E^(-ArcCsch[c + d*x])]))/d - (3*b^2
*d*e^2*f*x*((c + d*x)*Sqrt[1 + (c + d*x)^(-2)]*ArcCsch[c + d*x])/d^2 + ((
c + d*x)^2*ArcCsch[c + d*x]^2)/(2*d^2) - (c*ArcCsch[c + d*x]^2*Coth[ArcCsc
h[c + d*x]/2])/(2*d^2) - Log[(c + d*x)^(-1)]/d^2 - ((2*I)*c*(I*ArcCsch[c +
d*x]*(Log[1 - E^(-ArcCsch[c + d*x])]) - Log[1 + E^(-ArcCsch[c + d*x])])) +
I*(PolyLog[2, -E^(-ArcCsch[c + d*x])]) - PolyLog[2, E^(-ArcCsch[c + d*x])])
)/d^2 + (c*ArcCsch[c + d*x]^2*Tanh[ArcCsch[c + d*x]/2])/(2*d^2))/((c + d
*x)*(-1 + c/(c + d*x))) - (b^2*e*f^2*(2*(-2 + 12*c*ArcCsch[c + d*x] + ArcC
sch[c + d*x]^2 - 6*c^2*ArcCsch[c + d*x]^2)*Coth[ArcCsch[c + d*x]/2] + 2*Ar
cCsch[c + d*x]*(-1 + 3*c*ArcCsch[c + d*x])*Csch[ArcCsch[c + d*x]/2]^2 - (A
rcCsch[c + d*x]^2*Csch[ArcCsch[c + d*x]/2]^4)/(2*(c + d*x)) - 48*c*Log[...

```

Rubi [A] (verified)

Time = 1.07 (sec) , antiderivative size = 476, normalized size of antiderivative = 0.95, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6876, 5992, 3042, 4678, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)^3 (a + bcsch^{-1}(c + dx))^2 dx$$

$$\downarrow 6876$$

$$\frac{f(c + dx)^2 \sqrt{1 + \frac{1}{(c+dx)^2}} (de - cf + f(c + dx))^3 (a + bcsch^{-1}(c + dx))^2 dcsch^{-1}(c + dx)}{d^4}$$

$$\downarrow 5992$$

$$\begin{aligned}
 & \frac{b \int (de - cf + f(c + dx))^4 (a + b \operatorname{csch}^{-1}(c + dx)) d \operatorname{csch}^{-1}(c + dx)}{2f} - \frac{(f(c + dx) - cf + de)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4f} \\
 & \qquad \qquad \qquad \downarrow d^4 \qquad \qquad \qquad 3042 \\
 & \frac{(f(c + dx) - cf + de)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4f} + \frac{b \int (a + b \operatorname{csch}^{-1}(c + dx)) (de - cf + if \operatorname{csc}(i \operatorname{csch}^{-1}(c + dx)))^4 d \operatorname{csch}^{-1}(c + dx)}{2f} \\
 & \qquad \qquad \qquad \downarrow d^4 \qquad \qquad \qquad 4678 \\
 & b \int \left(d^4 \left(\frac{cf(-4d^3e^3 + 6cd^2fe^2 - 4c^2df^2e + c^3f^3)}{d^4e^4} + 1 \right) (a + b \operatorname{csch}^{-1}(c + dx)) e^4 + 4d^3f \left(1 - \frac{cf(3d^2e^2 - 3cdf e + c^2f^2)}{d^3e^3} \right) (c + dx) (a + b \operatorname{csch}^{-1}(c + dx)) e \right) \\
 & \qquad \qquad \qquad \downarrow 2009 \\
 & b \left(4f^3(de - cf) \operatorname{arctanh}(e^{\operatorname{csch}^{-1}(c + dx)}) (a + b \operatorname{csch}^{-1}(c + dx)) - 8f(de - cf)^3 \operatorname{arctanh}(e^{\operatorname{csch}^{-1}(c + dx)}) (a + b \operatorname{csch}^{-1}(c + dx)) - 2f^3(c + dx) \right)
 \end{aligned}$$

input `Int[(e + f*x)^3*(a + b*ArcCsch[c + d*x])^2,x]`

output `-((-1/4*((d*e - c*f + f*(c + d*x))^4*(a + b*ArcCsch[c + d*x])^2)/f + (b*(-2*b*f^3*(d*e - c*f)*(c + d*x) - (b*f^4*(c + d*x)^2)/6 + (2*f^4*(c + d*x)*Sqrt[1 + (c + d*x)^(-2)]*(a + b*ArcCsch[c + d*x]))/3 - 6*f^2*(d*e - c*f)^2*(c + d*x)*Sqrt[1 + (c + d*x)^(-2)]*(a + b*ArcCsch[c + d*x]) - 2*f^3*(d*e - c*f)*(c + d*x)^2*Sqrt[1 + (c + d*x)^(-2)]*(a + b*ArcCsch[c + d*x]) - (f^4*(c + d*x)^3*Sqrt[1 + (c + d*x)^(-2)]*(a + b*ArcCsch[c + d*x]))/3 + ((d*e - c*f)^4*(a + b*ArcCsch[c + d*x])^2)/(2*b) + 4*f^3*(d*e - c*f)*(a + b*ArcCsch[c + d*x])*ArcTanh[E^ArcCsch[c + d*x]] - 8*f*(d*e - c*f)^3*(a + b*ArcCsch[c + d*x])*ArcTanh[E^ArcCsch[c + d*x]] - (2*b*f^4*Log[(c + d*x)^(-1)])/3 + 6*b*f^2*(d*e - c*f)^2*Log[(c + d*x)^(-1)] + 2*b*f^3*(d*e - c*f)*PolyLog[2, -E^ArcCsch[c + d*x]] - 4*b*f*(d*e - c*f)^3*PolyLog[2, -E^ArcCsch[c + d*x]] - 2*b*f^3*(d*e - c*f)*PolyLog[2, E^ArcCsch[c + d*x]] + 4*b*f*(d*e - c*f)^3*PolyLog[2, E^ArcCsch[c + d*x]]))/(2*f))/d^4`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4678 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

rule 5992 `Int[Coth[(c_.) + (d_.)*(x_)]*Csch[(c_.) + (d_.)*(x_)]*(Csch[(c_.) + (d_.)*(x_)]*(b_.) + (a_)^(n_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

rule 6876 `Int[((a_.) + ArcCsch[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Csch[x]*Coth[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

Maple **[F]**

$$\int (fx + e)^3 (a + b \operatorname{arccsch}(dx + c))^2 dx$$

input `int((f*x+e)^3*(a+b*arccsch(d*x+c))^2,x)`

output `int((f*x+e)^3*(a+b*arccsch(d*x+c))^2,x)`

Fricas [F]

$$\int (e + fx)^3 (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (fx + e)^3 (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

input `integrate((f*x+e)^3*(a+b*arccsch(d*x+c))^2,x, algorithm="fricas")`

output `integral(a^2*f^3*x^3 + 3*a^2*e*f^2*x^2 + 3*a^2*e^2*f*x + a^2*e^3 + (b^2*f^3*x^3 + 3*b^2*e*f^2*x^2 + 3*b^2*e^2*f*x + b^2*e^3)*arccsch(d*x + c)^2 + 2*(a*b*f^3*x^3 + 3*a*b*e*f^2*x^2 + 3*a*b*e^2*f*x + a*b*e^3)*arccsch(d*x + c), x)`

Sympy [F]

$$\int (e + fx)^3 (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (a + b \operatorname{arcsch}(c + dx))^2 (e + fx)^3 dx$$

input `integrate((f*x+e)**3*(a+b*arcsch(d*x+c))**2,x)`

output `Integral((a + b*arcsch(c + d*x))**2*(e + f*x)**3, x)`

Maxima [F]

$$\int (e + fx)^3 (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (fx + e)^3 (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

input `integrate((f*x+e)^3*(a+b*arccsch(d*x+c))^2,x, algorithm="maxima")`

output

```

1/4*a^2*f^3*x^4 + a^2*e*f^2*x^3 + 3/2*a^2*e^2*f*x^2 + a^2*e^3*x + (2*(d*x
+ c)*arccsch(d*x + c) + log(sqrt(1/(d*x + c)^2 + 1) + 1) - log(sqrt(1/(d*x
+ c)^2 + 1) - 1))*a*b*e^3/d + 1/4*(b^2*f^3*x^4 + 4*b^2*e*f^2*x^3 + 6*b^2*
e^2*f*x^2 + 4*b^2*e^3*x)*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1)^2 - in
tegrate(-1/2*(2*(b^2*d^2*f^3*x^5 + b^2*c^2*e^3 + b^2*e^3 + (3*b^2*d^2*e*f^
2 + 2*b^2*c*d*f^3)*x^4 + (6*b^2*c*d*e*f^2 + b^2*c^2*f^3 + (3*d^2*e^2*f + f
^3)*b^2)*x^3 + (6*b^2*c*d*e^2*f + 3*b^2*c^2*e*f^2 + (d^2*e^3 + 3*e*f^2)*b^
2)*x^2 + (2*b^2*c*d*e^3 + 3*b^2*c^2*e^2*f + 3*b^2*e^2*f)*x)*log(d*x + c)^2
- 4*(a*b*d^2*f^3*x^5 + (3*a*b*d^2*e*f^2 + 2*a*b*c*d*f^3)*x^4 + (6*a*b*c*d
*e*f^2 + a*b*c^2*f^3 + (3*d^2*e^2*f + f^3)*a*b)*x^3 + 3*(2*a*b*c*d*e^2*f +
a*b*c^2*e*f^2 + a*b*e*f^2)*x^2 + 3*(a*b*c^2*e^2*f + a*b*e^2*f)*x)*log(d*x
+ c) + (4*a*b*d^2*f^3*x^5 + 4*(3*a*b*d^2*e*f^2 + 2*a*b*c*d*f^3)*x^4 + 4*(
6*a*b*c*d*e*f^2 + a*b*c^2*f^3 + (3*d^2*e^2*f + f^3)*a*b)*x^3 + 12*(2*a*b*c
*d*e^2*f + a*b*c^2*e*f^2 + a*b*e*f^2)*x^2 + 12*(a*b*c^2*e^2*f + a*b*e^2*f)
*x - 4*(b^2*d^2*f^3*x^5 + b^2*c^2*e^3 + b^2*e^3 + (3*b^2*d^2*e*f^2 + 2*b^2
*c*d*f^3)*x^4 + (6*b^2*c*d*e*f^2 + b^2*c^2*f^3 + (3*d^2*e^2*f + f^3)*b^2)*
x^3 + (6*b^2*c*d*e^2*f + 3*b^2*c^2*e*f^2 + (d^2*e^3 + 3*e*f^2)*b^2)*x^2 +
(2*b^2*c*d*e^3 + 3*b^2*c^2*e^2*f + 3*b^2*e^2*f)*x)*log(d*x + c) + ((4*a*b*
d^2*f^3 - b^2*d^2*f^3)*x^5 + (12*a*b*d^2*e*f^2 - 4*b^2*d^2*e*f^2 + (8*a*b*
d*f^3 - b^2*d*f^3)*c)*x^4 - 2*(3*b^2*d^2*e^2*f - 2*a*b*c^2*f^3 - 2*(3*d...

```

Giac [F]

$$\int (e + fx)^3 (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (fx + e)^3 (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

input

```
integrate((f*x+e)^3*(a+b*arccsch(d*x+c))^2,x, algorithm="giac")
```

output

```
integrate((f*x + e)^3*(b*arccsch(d*x + c) + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int (e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \int (e + fx)^3 \left(a + b \operatorname{asinh}\left(\frac{1}{c + dx}\right) \right)^2 dx$$

input `int((e + f*x)^3*(a + b*asinh(1/(c + d*x)))^2,x)`output `int((e + f*x)^3*(a + b*asinh(1/(c + d*x)))^2, x)`**Reduce [F]**

$$\begin{aligned} \int (e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = & 2 \left(\int \operatorname{acsch}(dx + c) dx \right) ab e^3 \\ & + \left(\int \operatorname{acsch}(dx + c)^2 dx \right) b^2 e^3 \\ & + 2 \left(\int \operatorname{acsch}(dx + c) x^3 dx \right) ab f^3 \\ & + 6 \left(\int \operatorname{acsch}(dx + c) x^2 dx \right) ab e f^2 \\ & + 6 \left(\int \operatorname{acsch}(dx + c) x dx \right) ab e^2 f \\ & + \left(\int \operatorname{acsch}(dx + c)^2 x^3 dx \right) b^2 f^3 \\ & + 3 \left(\int \operatorname{acsch}(dx + c)^2 x^2 dx \right) b^2 e f^2 \\ & + 3 \left(\int \operatorname{acsch}(dx + c)^2 x dx \right) b^2 e^2 f \\ & + a^2 e^3 x + \frac{3a^2 e^2 f x^2}{2} + a^2 e f^2 x^3 + \frac{a^2 f^3 x^4}{4} \end{aligned}$$

input `int((f*x+e)^3*(a+b*acsch(d*x+c))^2,x)`

output

```
(8*int(acsch(c + d*x),x)*a*b*e**3 + 4*int(acsch(c + d*x)**2,x)*b**2*e**3 +
 8*int(acsch(c + d*x)*x**3,x)*a*b*f**3 + 24*int(acsch(c + d*x)*x**2,x)*a*b
*e*f**2 + 24*int(acsch(c + d*x)*x,x)*a*b*e**2*f + 4*int(acsch(c + d*x)**2*
x**3,x)*b**2*f**3 + 12*int(acsch(c + d*x)**2*x**2,x)*b**2*e*f**2 + 12*int(
acsch(c + d*x)**2*x,x)*b**2*e**2*f + 4*a**2*e**3*x + 6*a**2*e**2*f*x**2 +
4*a**2*e*f**2*x**3 + a**2*f**3*x**4)/4
```

3.8 $\int (e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$

Optimal result	97
Mathematica [C] (warning: unable to verify)	98
Rubi [A] (verified)	99
Maple [F]	101
Fricas [F]	102
Sympy [F]	102
Maxima [F]	102
Giac [F]	103
Mupad [F(-1)]	104
Reduce [F]	104

Optimal result

Integrand size = 20, antiderivative size = 351

$$\begin{aligned}
 & \int (e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx \\
 &= \frac{b^2 f^2 x}{3d^2} + \frac{2bf(de - cf)(c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{d^3} \\
 &+ \frac{bf^2(c + dx)^2 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{3d^3} \\
 &- \frac{(de - cf)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2}{3d^3 f} + \frac{(e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2}{3f} \\
 &- \frac{2bf^2(a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{3d^3} \\
 &+ \frac{4b(de - cf)^2 (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^3} \\
 &+ \frac{2b^2 f(de - cf) \log(c + dx)}{d^3} - \frac{b^2 f^2 \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{3d^3} \\
 &+ \frac{2b^2(de - cf)^2 \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^3} \\
 &+ \frac{b^2 f^2 \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{3d^3} - \frac{2b^2(de - cf)^2 \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^3}
 \end{aligned}$$

output

```

1/3*b^2*f^2*x/d^2+2*b*f*(-c*f+d*e)*(d*x+c)*(1+1/(d*x+c)^2)^(1/2)*(a+b*arccsch(d*x+c))/d^3+1/3*b*f^2*(d*x+c)^2*(1+1/(d*x+c)^2)^(1/2)*(a+b*arccsch(d*x+c))/d^3-1/3*(-c*f+d*e)^3*(a+b*arccsch(d*x+c))^2/d^3/f+1/3*(f*x+e)^3*(a+b*arccsch(d*x+c))^2/f-2/3*b*f^2*(a+b*arccsch(d*x+c))*arctanh(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))/d^3+4*b*(-c*f+d*e)^2*(a+b*arccsch(d*x+c))*arctanh(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))/d^3+2*b^2*f*(-c*f+d*e)*ln(d*x+c)/d^3-1/3*b^2*f^2*polylog(2,-1/(d*x+c)-(1+1/(d*x+c)^2)^(1/2))/d^3+2*b^2*(-c*f+d*e)^2*polylog(2,-1/(d*x+c)-(1+1/(d*x+c)^2)^(1/2))/d^3+1/3*b^2*f^2*polylog(2,1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))/d^3-2*b^2*(-c*f+d*e)^2*polylog(2,1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))/d^3

```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 7.87 (sec) , antiderivative size = 864, normalized size of antiderivative = 2.46

$$\int (e + fx)^2 (a + bcsch^{-1}(c + dx))^2 dx = \text{Too large to display}$$

input

```
Integrate[(e + f*x)^2*(a + b*ArcCsch[c + d*x])^2,x]
```

output

```

a^2*e^2*x + a^2*e*f*x^2 + (a^2*f^2*x^3)/3 + (a*b*(2*x*(3*e^2 + 3*e*f*x + f
^2*x^2)*ArcCsch[c + d*x] + (-(f*(c + d*x)*Sqrt[(1 + c^2 + 2*c*d*x + d^2*x^
2)/(c + d*x)^2]*(5*c*f - d*(6*e + f*x))) + 2*c*(3*d^2*e^2 - 3*c*d*e*f + c^
2*f^2)*ArcSinh[(c + d*x)^(-1)] + (6*d^2*e^2 - 12*c*d*e*f + (-1 + 6*c^2)*f^
2)*Log[(c + d*x)*(1 + Sqrt[(1 + c^2 + 2*c*d*x + d^2*x^2)/(c + d*x)^2]))/d
^3))/3 - (b^2*e^2*(-(ArcCsch[c + d*x]*((c + d*x)*ArcCsch[c + d*x] - 2*Log[
1 - E^(-ArcCsch[c + d*x])) + 2*Log[1 + E^(-ArcCsch[c + d*x]))]) + 2*PolyLo
g[2, -E^(-ArcCsch[c + d*x])] - 2*PolyLog[2, E^(-ArcCsch[c + d*x])]))/d - (
2*b^2*d*e*f*x*((c + d*x)*Sqrt[1 + (c + d*x)^(-2)]*ArcCsch[c + d*x])/d^2 +
((c + d*x)^2*ArcCsch[c + d*x]^2)/(2*d^2) - (c*ArcCsch[c + d*x]^2*Coth[Arc
Csch[c + d*x]/2])/(2*d^2) - Log[(c + d*x)^(-1)]/d^2 - ((2*I)*c*(I*ArcCsch[
c + d*x]*(Log[1 - E^(-ArcCsch[c + d*x])] - Log[1 + E^(-ArcCsch[c + d*x])])
+ I*(PolyLog[2, -E^(-ArcCsch[c + d*x])] - PolyLog[2, E^(-ArcCsch[c + d*x]
)])))/d^2 + (c*ArcCsch[c + d*x]^2*Tanh[ArcCsch[c + d*x]/2])/(2*d^2))/((c
+ d*x)*(-1 + c/(c + d*x))) - (b^2*f^2*(2*(-2 + 12*c*ArcCsch[c + d*x] + Arc
Csch[c + d*x]^2 - 6*c^2*ArcCsch[c + d*x]^2)*Coth[ArcCsch[c + d*x]/2] + 2*A
rcCsch[c + d*x]*(-1 + 3*c*ArcCsch[c + d*x])*Csch[ArcCsch[c + d*x]/2]^2 - (
ArcCsch[c + d*x]^2*Csch[ArcCsch[c + d*x]/2]^4)/(2*(c + d*x)) - 48*c*Log[(c
+ d*x)^(-1)] + 8*(-1 + 6*c^2)*(ArcCsch[c + d*x]*(Log[1 - E^(-ArcCsch[c +
d*x])]) - Log[1 + E^(-ArcCsch[c + d*x])]) + PolyLog[2, -E^(-ArcCsch[c + ...

```

Rubi [A] (verified)

Time = 0.77 (sec) , antiderivative size = 336, normalized size of antiderivative = 0.96, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6876, 5992, 3042, 4678, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)^2 (a + bcsch^{-1}(c + dx))^2 dx$$

$$\downarrow 6876$$

$$\frac{\int (c + dx)^2 \sqrt{1 + \frac{1}{(c+dx)^2}} (de - cf + f(c + dx))^2 (a + bcsch^{-1}(c + dx))^2 dcsch^{-1}(c + dx)}{d^3}$$

$$\downarrow 5992$$

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4678 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

rule 5992 `Int[Coth[(c_.) + (d_.)*(x_)]*Csch[(c_.) + (d_.)*(x_)]*(Csch[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

rule 6876 `Int[((a_.) + ArcCsch[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Csch[x]*Coth[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

Maple **[F]**

$$\int (fx + e)^2 (a + b \operatorname{arccsch}(dx + c))^2 dx$$

input `int((f*x+e)^2*(a+b*arccsch(d*x+c))^2,x)`

output `int((f*x+e)^2*(a+b*arccsch(d*x+c))^2,x)`

Fricas [F]

$$\int (e + fx)^2 (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (fx + e)^2 (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

input `integrate((f*x+e)^2*(a+b*arccsch(d*x+c))^2,x, algorithm="fricas")`

output `integral(a^2*f^2*x^2 + 2*a^2*e*f*x + a^2*e^2 + (b^2*f^2*x^2 + 2*b^2*e*f*x + b^2*e^2)*arccsch(d*x + c)^2 + 2*(a*b*f^2*x^2 + 2*a*b*e*f*x + a*b*e^2)*arccsch(d*x + c), x)`

Sympy [F]

$$\int (e + fx)^2 (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (a + b \operatorname{arcsch}(c + dx))^2 (e + fx)^2 dx$$

input `integrate((f*x+e)**2*(a+b*arcsch(d*x+c))**2,x)`

output `Integral((a + b*arcsch(c + d*x))**2*(e + f*x)**2, x)`

Maxima [F]

$$\int (e + fx)^2 (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (fx + e)^2 (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

input `integrate((f*x+e)^2*(a+b*arccsch(d*x+c))^2,x, algorithm="maxima")`

output

```

1/3*a^2*f^2*x^3 + a^2*e*f*x^2 + a^2*e^2*x + (2*(d*x + c)*arccsch(d*x + c)
+ log(sqrt(1/(d*x + c)^2 + 1) + 1) - log(sqrt(1/(d*x + c)^2 + 1) - 1))*a*b
*e^2/d + 1/3*(b^2*f^2*x^3 + 3*b^2*e*f*x^2 + 3*b^2*e^2*x)*log(sqrt(d^2*x^2
+ 2*c*d*x + c^2 + 1) + 1)^2 - integrate(-1/3*(3*(b^2*d^2*f^2*x^4 + b^2*c^2
*e^2 + b^2*e^2 + 2*(b^2*d^2*e*f + b^2*c*d*f^2))*x^3 + (4*b^2*c*d*e*f + b^2*
c^2*f^2 + (d^2*e^2 + f^2)*b^2))*x^2 + 2*(b^2*c*d*e^2 + b^2*c^2*e*f + b^2*e*
f)*x)*log(d*x + c)^2 - 6*(a*b*d^2*f^2*x^4 + 2*(a*b*d^2*e*f + a*b*c*d*f^2)*
x^3 + (4*a*b*c*d*e*f + a*b*c^2*f^2 + a*b*f^2))*x^2 + 2*(a*b*c^2*e*f + a*b*e
*f)*x)*log(d*x + c) + 2*(3*a*b*d^2*f^2*x^4 + 6*(a*b*d^2*e*f + a*b*c*d*f^2)
*x^3 + 3*(4*a*b*c*d*e*f + a*b*c^2*f^2 + a*b*f^2))*x^2 + 6*(a*b*c^2*e*f + a*
b*e*f)*x - 3*(b^2*d^2*f^2*x^4 + b^2*c^2*e^2 + b^2*e^2 + 2*(b^2*d^2*e*f + b
^2*c*d*f^2))*x^3 + (4*b^2*c*d*e*f + b^2*c^2*f^2 + (d^2*e^2 + f^2)*b^2))*x^2
+ 2*(b^2*c*d*e^2 + b^2*c^2*e*f + b^2*e*f)*x)*log(d*x + c) + ((3*a*b*d^2*f^
2 - b^2*d^2*f^2)*x^4 + (6*a*b*d^2*e*f - 3*b^2*d^2*e*f + (6*a*b*d*f^2 - b^2
*d*f^2)*c)*x^3 - 3*(b^2*d^2*e^2 - a*b*c^2*f^2 - a*b*f^2 - (4*a*b*d*e*f - b
^2*d*e*f)*c)*x^2 - 3*(b^2*c*d*e^2 - 2*a*b*c^2*e*f - 2*a*b*e*f)*x - 3*(b^2*
d^2*f^2*x^4 + b^2*c^2*e^2 + b^2*e^2 + 2*(b^2*d^2*e*f + b^2*c*d*f^2))*x^3 +
(4*b^2*c*d*e*f + b^2*c^2*f^2 + (d^2*e^2 + f^2)*b^2))*x^2 + 2*(b^2*c*d*e^2 +
b^2*c^2*e*f + b^2*e*f)*x)*log(d*x + c))*sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1)
)*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1) + 3*sqrt(d^2*x^2 + 2*c*d*x...

```

Giac [F]

$$\int (e + fx)^2 (a + b \operatorname{arcsch}(c + dx))^2 dx = \int (fx + e)^2 (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

input

```
integrate((f*x+e)^2*(a+b*arccsch(d*x+c))^2,x, algorithm="giac")
```

output

```
integrate((f*x + e)^2*(b*arccsch(d*x + c) + a)^2, x)
```


Mupad [F(-1)]

Timed out.

$$\int (e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \int (e + fx)^2 \left(a + b \operatorname{asinh}\left(\frac{1}{c + dx}\right) \right)^2 dx$$

input `int((e + f*x)^2*(a + b*asinh(1/(c + d*x)))^2,x)`output `int((e + f*x)^2*(a + b*asinh(1/(c + d*x)))^2, x)`**Reduce [F]**

$$\begin{aligned} \int (e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = & 2 \left(\int \operatorname{acsch}(dx + c) dx \right) ab e^2 \\ & + \left(\int \operatorname{acsch}(dx + c)^2 dx \right) b^2 e^2 \\ & + 2 \left(\int \operatorname{acsch}(dx + c) x^2 dx \right) ab f^2 \\ & + 4 \left(\int \operatorname{acsch}(dx + c) x dx \right) ab e f \\ & + \left(\int \operatorname{acsch}(dx + c)^2 x^2 dx \right) b^2 f^2 \\ & + 2 \left(\int \operatorname{acsch}(dx + c)^2 x dx \right) b^2 e f \\ & + a^2 e^2 x + a^2 e f x^2 + \frac{a^2 f^2 x^3}{3} \end{aligned}$$

input `int((f*x+e)^2*(a+b*acsch(d*x+c))^2,x)`output `(6*int(acsch(c + d*x),x)*a*b*e**2 + 3*int(acsch(c + d*x)**2,x)*b**2*e**2 + 6*int(acsch(c + d*x)*x**2,x)*a*b*f**2 + 12*int(acsch(c + d*x)*x,x)*a*b*e*f + 3*int(acsch(c + d*x)**2*x**2,x)*b**2*f**2 + 6*int(acsch(c + d*x)**2*x,x)*b**2*e*f + 3*a**2*e**2*x + 3*a**2*e*f*x**2 + a**2*f**2*x**3)/3`

3.9 $\int (e + fx) (a + bcsch^{-1}(c + dx))^2 dx$

Optimal result	105
Mathematica [B] (verified)	106
Rubi [A] (verified)	106
Maple [F]	108
Fricas [F]	109
Sympy [F]	109
Maxima [F]	109
Giac [F]	110
Mupad [F(-1)]	110
Reduce [F]	111

Optimal result

Integrand size = 18, antiderivative size = 194

$$\int (e + fx) (a + bcsch^{-1}(c + dx))^2 dx$$

$$= \frac{bf(c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + bcsch^{-1}(c + dx))}{d^2}$$

$$- \frac{(de - cf)^2 (a + bcsch^{-1}(c + dx))^2}{2d^2 f} + \frac{(e + fx)^2 (a + bcsch^{-1}(c + dx))^2}{2f}$$

$$+ \frac{4b(de - cf) (a + bcsch^{-1}(c + dx)) \operatorname{arctanh}\left(e^{csch^{-1}(c+dx)}\right)}{d^2} + \frac{b^2 f \log(c + dx)}{d^2}$$

$$+ \frac{2b^2(de - cf) \operatorname{PolyLog}\left(2, -e^{csch^{-1}(c+dx)}\right)}{d^2} - \frac{2b^2(de - cf) \operatorname{PolyLog}\left(2, e^{csch^{-1}(c+dx)}\right)}{d^2}$$

output

```
b*f*(d*x+c)*(1+1/(d*x+c)^2)^(1/2)*(a+b*arccsch(d*x+c))/d^2-1/2*(-c*f+d*e)^2*(a+b*arccsch(d*x+c))^2/d^2/f+1/2*(f*x+e)^2*(a+b*arccsch(d*x+c))^2/f+4*b*(-c*f+d*e)*(a+b*arccsch(d*x+c))*arctanh(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))/d^2+b^2*f*ln(d*x+c)/d^2+2*b^2*(-c*f+d*e)*polylog(2,-1/(d*x+c)-(1+1/(d*x+c)^2)^(1/2))/d^2-2*b^2*(-c*f+d*e)*polylog(2,1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))/d^2
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 427 vs. $2(194) = 388$.

Time = 2.88 (sec) , antiderivative size = 427, normalized size of antiderivative = 2.20

$$\int (e + fx) (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$$

$$= \frac{2a^2(de - cf)(c + dx) + a^2 f(c + dx)^2 + 2abf(c + dx) \left(\sqrt{1 + \frac{1}{(c+dx)^2}} + (c + dx) \operatorname{csch}^{-1}(c + dx) \right) + 2b^2 f}{}$$

input

```
Integrate[(e + f*x)*(a + b*ArcCsch[c + d*x])^2,x]
```

output

```
(2*a^2*(d*e - c*f)*(c + d*x) + a^2*f*(c + d*x)^2 + 2*a*b*f*(c + d*x)*(Sqrt
[1 + (c + d*x)^(-2)] + (c + d*x)*ArcCsch[c + d*x]) + 2*b^2*f*((c + d*x)*Sqr
rt[1 + (c + d*x)^(-2)]*ArcCsch[c + d*x] + ((c + d*x)^2*ArcCsch[c + d*x]^2)
/2 - Log[(c + d*x)^(-1)]) + 4*a*b*d*e*((c + d*x)*ArcCsch[c + d*x] + Log[Csch
ch[ArcCsch[c + d*x]/2]/(2*(c + d*x))] - Log[Sinh[ArcCsch[c + d*x]/2]]) - 4
*a*b*c*f*((c + d*x)*ArcCsch[c + d*x] + Log[Csch[ArcCsch[c + d*x]/2]/(2*(c
+ d*x))] - Log[Sinh[ArcCsch[c + d*x]/2]]) + 2*b^2*d*e*(ArcCsch[c + d*x]*((
c + d*x)*ArcCsch[c + d*x] - 2*Log[1 - E^(-ArcCsch[c + d*x])]) + 2*Log[1 + E
^(-ArcCsch[c + d*x])]) - 2*PolyLog[2, -E^(-ArcCsch[c + d*x])] + 2*PolyLog[
2, E^(-ArcCsch[c + d*x])] - 2*b^2*c*f*(ArcCsch[c + d*x]*((c + d*x)*ArcCsch
[c + d*x] - 2*Log[1 - E^(-ArcCsch[c + d*x])] + 2*Log[1 + E^(-ArcCsch[c +
d*x])]) - 2*PolyLog[2, -E^(-ArcCsch[c + d*x])] + 2*PolyLog[2, E^(-ArcCsch[
c + d*x])])))/(2*d^2)
```

Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {6876, 5992, 3042, 4678, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (e + fx) (a + b\operatorname{csch}^{-1}(c + dx))^2 dx \\
 & \quad \downarrow \text{6876} \\
 & \frac{\int (c + dx)^2 \sqrt{1 + \frac{1}{(c+dx)^2}} (de - cf + f(c + dx)) (a + b\operatorname{csch}^{-1}(c + dx))^2 d\operatorname{csch}^{-1}(c + dx)}{d^2} \\
 & \quad \downarrow \text{5992} \\
 & \frac{b \int (de - cf + f(c + dx))^2 (a + b\operatorname{csch}^{-1}(c + dx)) d\operatorname{csch}^{-1}(c + dx)}{f} - \frac{(f(c + dx) - cf + de)^2 (a + b\operatorname{csch}^{-1}(c + dx))^2}{2f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(f(c + dx) - cf + de)^2 (a + b\operatorname{csch}^{-1}(c + dx))^2}{2f} + \frac{b \int (a + b\operatorname{csch}^{-1}(c + dx)) (de - cf + if \operatorname{csc}(i\operatorname{csch}^{-1}(c + dx)))^2 d\operatorname{csch}^{-1}(c + dx)}{d^2} \\
 & \quad \downarrow \text{4678} \\
 & \frac{b \int \left(d^2 \left(\frac{cf(c f - 2de)}{d^2 e^2} + 1 \right) (a + b\operatorname{csch}^{-1}(c + dx)) e^2 + 2df \left(1 - \frac{cf}{de} \right) (c + dx) (a + b\operatorname{csch}^{-1}(c + dx)) e + f^2 (c + dx)^2 (a + b\operatorname{csch}^{-1}(c + dx)) \right) d\operatorname{csch}^{-1}(c + dx)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{b \left(-4f(de - cf) \operatorname{arctanh}(e^{\operatorname{csch}^{-1}(c + dx)}) (a + b\operatorname{csch}^{-1}(c + dx)) + \frac{(de - cf)^2 (a + b\operatorname{csch}^{-1}(c + dx))^2}{2b} - \left(f^2 (c + dx) \sqrt{\frac{1}{(c + dx)^2} + 1} (a + b\operatorname{csch}^{-1}(c + dx)) \right) \right)}{f}
 \end{aligned}$$

input `Int[(e + f*x)*(a + b*ArcCsch[c + d*x])^2,x]`

output `-((-1/2*((d*e - c*f + f*(c + d*x))^2*(a + b*ArcCsch[c + d*x])^2)/f + (b*(-f^2*(c + d*x)*Sqrt[1 + (c + d*x)^(-2)]*(a + b*ArcCsch[c + d*x])) + ((d*e - c*f)^2*(a + b*ArcCsch[c + d*x])^2)/(2*b) - 4*f*(d*e - c*f)*(a + b*ArcCsch[c + d*x])*ArcTanh[E^ArcCsch[c + d*x]] + b*f^2*Log[(c + d*x)^(-1)] - 2*b*f*(d*e - c*f)*PolyLog[2, -E^ArcCsch[c + d*x]] + 2*b*f*(d*e - c*f)*PolyLog[2, E^ArcCsch[c + d*x]]))/f/d^2)`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4678 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

rule 5992 `Int[Coth[(c_.) + (d_.)*(x_)]*Csch[(c_.) + (d_.)*(x_)]*(Csch[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

rule 6876 `Int[((a_.) + ArcCsch[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Csch[x]*Coth[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

Maple **[F]**

$$\int (fx + e)(a + b \operatorname{arccsch}(dx + c))^2 dx$$

input `int((f*x+e)*(a+b*arccsch(d*x+c))^2,x)`

output `int((f*x+e)*(a+b*arccsch(d*x+c))^2,x)`

Fricas [F]

$$\int (e + fx) (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (fx + e)(b \operatorname{arcsch}(dx + c) + a)^2 dx$$

input `integrate((f*x+e)*(a+b*arccsch(d*x+c))^2,x, algorithm="fricas")`

output `integral(a^2*f*x + a^2*e + (b^2*f*x + b^2*e)*arccsch(d*x + c)^2 + 2*(a*b*f*x + a*b*e)*arccsch(d*x + c), x)`

Sympy [F]

$$\int (e + fx) (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (a + b \operatorname{arcsch}(c + dx))^2 (e + fx) dx$$

input `integrate((f*x+e)*(a+b*arcsch(d*x+c))**2,x)`

output `Integral((a + b*arcsch(c + d*x))**2*(e + f*x), x)`

Maxima [F]

$$\int (e + fx) (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (fx + e)(b \operatorname{arcsch}(dx + c) + a)^2 dx$$

input `integrate((f*x+e)*(a+b*arccsch(d*x+c))^2,x, algorithm="maxima")`

output

```

1/2*a^2*f*x^2 + a^2*e*x + (2*(d*x + c)*arccsch(d*x + c) + log(sqrt(1/(d*x
+ c)^2 + 1) + 1) - log(sqrt(1/(d*x + c)^2 + 1) - 1))*a*b*e/d + 1/2*(b^2*f*
x^2 + 2*b^2*e*x)*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1)^2 - integrate(
-((b^2*d^2*f*x^3 + b^2*c^2*e + b^2*e + (b^2*d^2*e + 2*b^2*c*d*f)*x^2 + (2*
b^2*c*d*e + b^2*c^2*f + b^2*f)*x)*log(d*x + c)^2 - 2*(a*b*d^2*f*x^3 + 2*a*
b*c*d*f*x^2 + (a*b*c^2*f + a*b*f)*x)*log(d*x + c) + (2*a*b*d^2*f*x^3 + 4*a
*b*c*d*f*x^2 + 2*(a*b*c^2*f + a*b*f)*x - 2*(b^2*d^2*f*x^3 + b^2*c^2*e + b^
2*e + (b^2*d^2*e + 2*b^2*c*d*f)*x^2 + (2*b^2*c*d*e + b^2*c^2*f + b^2*f)*x)
*log(d*x + c) + sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1)*((2*a*b*d^2*f - b^2*d^2*
f)*x^3 - (2*b^2*d^2*e - (4*a*b*d*f - b^2*d*f)*c)*x^2 - 2*(b^2*c*d*e - a*b*
c^2*f - a*b*f)*x - 2*(b^2*d^2*f*x^3 + b^2*c^2*e + b^2*e + (b^2*d^2*e + 2*b
^2*c*d*f)*x^2 + (2*b^2*c*d*e + b^2*c^2*f + b^2*f)*x)*log(d*x + c)))*log(sq
rt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1) + sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1)*
(b^2*d^2*f*x^3 + b^2*c^2*e + b^2*e + (b^2*d^2*e + 2*b^2*c*d*f)*x^2 + (2*b^
2*c*d*e + b^2*c^2*f + b^2*f)*x)*log(d*x + c)^2 - 2*(a*b*d^2*f*x^3 + 2*a*b*
c*d*f*x^2 + (a*b*c^2*f + a*b*f)*x)*log(d*x + c)))/(d^2*x^2 + 2*c*d*x + c^2
+ (d^2*x^2 + 2*c*d*x + c^2 + 1)^(3/2) + 1), x)

```

Giac [F]

$$\int (e + fx) (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (fx + e) (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

input

```
integrate((f*x+e)*(a+b*arccsch(d*x+c))^2,x, algorithm="giac")
```

output

```
integrate((f*x + e)*(b*arccsch(d*x + c) + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int (e + fx) (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (e + fx) \left(a + b \operatorname{asinh} \left(\frac{1}{c + dx} \right) \right)^2 dx$$

input

```
int((e + f*x)*(a + b*asinh(1/(c + d*x)))^2,x)
```

output `int((e + f*x)*(a + b*asinh(1/(c + d*x)))^2, x)`

Reduce [F]

$$\int (e + fx) (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = 2 \left(\int \operatorname{acsch}(dx + c) dx \right) abe$$

$$+ \left(\int \operatorname{acsch}(dx + c)^2 dx \right) b^2 e$$

$$+ 2 \left(\int \operatorname{acsch}(dx + c) x dx \right) abf$$

$$+ \left(\int \operatorname{acsch}(dx + c)^2 x dx \right) b^2 f + a^2 ex + \frac{a^2 f x^2}{2}$$

input `int((f*x+e)*(a+b*acsch(d*x+c))^2,x)`

output `(4*int(acsch(c + d*x),x)*a*b*e + 2*int(acsch(c + d*x)**2,x)*b**2*e + 4*int(acsch(c + d*x)*x,x)*a*b*f + 2*int(acsch(c + d*x)**2*x,x)*b**2*f + 2*a**2*e*x + a**2*f*x**2)/2`

3.10 $\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$

Optimal result	112
Mathematica [B] (verified)	113
Rubi [C] (warning: unable to verify)	113
Maple [F]	116
Fricas [F]	116
Sympy [F]	116
Maxima [F]	117
Giac [F]	117
Mupad [F(-1)]	117
Reduce [F]	118

Optimal result

Integrand size = 12, antiderivative size = 85

$$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \frac{(c + dx) (a + b \operatorname{csch}^{-1}(c + dx))^2}{d} + \frac{4b(a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c + dx)}\right)}{d} + \frac{2b^2 \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c + dx)}\right)}{d} - \frac{2b^2 \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c + dx)}\right)}{d}$$

output

```
(d*x+c)*(a+b*arccsch(d*x+c))^2/d+4*b*(a+b*arccsch(d*x+c))*arctanh(1/(d*x+c)
)+(1+1/(d*x+c)^2)^(1/2)/d+2*b^2*polylog(2,-1/(d*x+c)-(1+1/(d*x+c)^2)^(1/2)
)/d-2*b^2*polylog(2,1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))/d
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 176 vs. $2(85) = 170$.

Time = 0.20 (sec) , antiderivative size = 176, normalized size of antiderivative = 2.07

$$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$$

$$= \frac{a^2 c + a^2 dx + 2ab(c + dx) \operatorname{csch}^{-1}(c + dx) + b^2 c \operatorname{csch}^{-1}(c + dx)^2 + b^2 dx \operatorname{csch}^{-1}(c + dx)^2 - 2b^2 \operatorname{csch}^{-1}(c + dx)}{d}$$

input `Integrate[(a + b*ArcCsch[c + d*x])^2,x]`

output `(a^2*c + a^2*d*x + 2*a*b*(c + d*x)*ArcCsch[c + d*x] + b^2*c*ArcCsch[c + d*x]^2 + b^2*d*x*ArcCsch[c + d*x]^2 - 2*b^2*ArcCsch[c + d*x]*Log[1 - E^(-ArcCsch[c + d*x])] + 2*b^2*ArcCsch[c + d*x]*Log[1 + E^(-ArcCsch[c + d*x])] + 2*a*b*Log[Cosh[ArcCsch[c + d*x]/2]] - 2*a*b*Log[Sinh[ArcCsch[c + d*x]/2]] - 2*b^2*PolyLog[2, -E^(-ArcCsch[c + d*x])] + 2*b^2*PolyLog[2, E^(-ArcCsch[c + d*x])])/d`

Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.47 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.99, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {6870, 6834, 5975, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$$

$$\downarrow 6870$$

$$\frac{\int (a + b \operatorname{csch}^{-1}(c + dx))^2 d(c + dx)}{d}$$

$$\downarrow 6834$$

$$\begin{aligned}
& \frac{\int (c+dx)^2 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c+dx))^2 d \operatorname{csch}^{-1}(c+dx)}{d} \\
& \quad \downarrow 5975 \\
& \frac{2b \int (c+dx) (a + b \operatorname{csch}^{-1}(c+dx)) d \operatorname{csch}^{-1}(c+dx) - (c+dx) (a + b \operatorname{csch}^{-1}(c+dx))^2}{d} \\
& \quad \downarrow 3042 \\
& \frac{-(c+dx) (a + b \operatorname{csch}^{-1}(c+dx))^2 + 2b \int i (a + b \operatorname{csch}^{-1}(c+dx)) \operatorname{csc}(i \operatorname{csch}^{-1}(c+dx)) d \operatorname{csch}^{-1}(c+dx)}{d} \\
& \quad \downarrow 26 \\
& \frac{-(c+dx) (a + b \operatorname{csch}^{-1}(c+dx))^2 + 2ib \int (a + b \operatorname{csch}^{-1}(c+dx)) \operatorname{csc}(i \operatorname{csch}^{-1}(c+dx)) d \operatorname{csch}^{-1}(c+dx)}{d} \\
& \quad \downarrow 4670 \\
& \frac{-(c+dx) (a + b \operatorname{csch}^{-1}(c+dx))^2 + 2ib \left(ib \int \log(1 - e^{\operatorname{csch}^{-1}(c+dx)}) d \operatorname{csch}^{-1}(c+dx) - ib \int \log(1 + e^{\operatorname{csch}^{-1}(c+dx)}) d \operatorname{csch}^{-1}(c+dx) \right)}{d} \\
& \quad \downarrow 2715 \\
& \frac{-(c+dx) (a + b \operatorname{csch}^{-1}(c+dx))^2 + 2ib \left(-ib \int e^{-\operatorname{csch}^{-1}(c+dx)} \log(1 + e^{\operatorname{csch}^{-1}(c+dx)}) d e^{\operatorname{csch}^{-1}(c+dx)} + ib \int e^{-\operatorname{csch}^{-1}(c+dx)} d e^{\operatorname{csch}^{-1}(c+dx)} \right)}{d} \\
& \quad \downarrow 2838 \\
& \frac{-(c+dx) (a + b \operatorname{csch}^{-1}(c+dx))^2 + 2ib \left(2i \operatorname{arctanh}(e^{\operatorname{csch}^{-1}(c+dx)}) (a + b \operatorname{csch}^{-1}(c+dx)) + ib \operatorname{PolyLog}(2, -c - d \operatorname{csch}^{-1}(c+dx)) \right)}{d}
\end{aligned}$$

input

```
Int[(a + b*ArcCsch[c + d*x])^2,x]
```

output

```
-((-(c + d*x)*(a + b*ArcCsch[c + d*x])^2) + (2*I)*b*((2*I)*(a + b*ArcCsch[c + d*x])*ArcTanh[E^ArcCsch[c + d*x]] - I*b*PolyLog[2, E^ArcCsch[c + d*x]] + I*b*PolyLog[2, -c - d*x]))/d
```

Definitions of rubi rules used

- rule 26 $\text{Int}[(\text{Complex}[0, a_])*(F_{x_}), x_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_{x_}, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 2715 $\text{Int}[\text{Log}[(a_ + (b_)*(F_)^{(e_)*((c_ + (d_)*(x_)))})^{(n_)}], x_Symbol] \rightarrow \text{Simp}[1/(d_*e*n*\text{Log}[F]) \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^{(e*(c + d*x)})^n), x] /; \text{FreeQ}\{F, a, b, c, d, e, n\}, x] \ \&\& \ \text{GtQ}[a, 0]$
- rule 2838 $\text{Int}[\text{Log}[(c_)*((d_ + (e_)*(x_)^{(n_)}))/x], x_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n/n, x] /; \text{FreeQ}\{c, d, e, n\}, x] \ \&\& \ \text{EqQ}[c*d, 1]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4670 $\text{Int}[\text{csc}[(e_ + (\text{Complex}[0, fz_])*(f_)*(x_))*((c_ + (d_)*(x_))^{(m_)}), x_Symbol] \rightarrow \text{Simp}[-2*(c + d*x)^m*(\text{ArcTanh}[E^{((-I)*e + f*fz*x)}/(f*fz*I)], x] + (-\text{Simp}[d*(m/(f*fz*I)) \text{Int}[(c + d*x)^{(m-1)}*\text{Log}[1 - E^{((-I)*e + f*fz*x}], x], x] + \text{Simp}[d*(m/(f*fz*I)) \text{Int}[(c + d*x)^{(m-1)}*\text{Log}[1 + E^{((-I)*e + f*fz*x}], x], x]) /; \text{FreeQ}\{c, d, e, f, fz\}, x] \ \&\& \ \text{IGtQ}[m, 0]$
- rule 5975 $\text{Int}[\text{Coth}[(a_ + (b_)*(x_))^{(p_)}]*\text{Csch}[(a_ + (b_)*(x_))^{(n_)}]*((c_ + (d_)*(x_))^{(m_)}), x_Symbol] \rightarrow \text{Simp}[(-c + d*x)^m*(\text{Csch}[a + b*x]^n/(b*n)), x] + \text{Simp}[d*(m/(b*n)) \text{Int}[(c + d*x)^{(m-1)}*\text{Csch}[a + b*x]^n, x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x] \ \&\& \ \text{EqQ}[p, 1] \ \&\& \ \text{GtQ}[m, 0]$
- rule 6834 $\text{Int}[(a_ + \text{ArcCsch}[(c_)*(x_)]*(b_))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[-c^{(-1)} \text{Subst}[\text{Int}[(a + b*x)^n*\text{Csch}[x]*\text{Coth}[x], x], x, \text{ArcCsch}[c*x]], x] /; \text{FreeQ}\{a, b, c, n\}, x] \ \&\& \ \text{IGtQ}[n, 0]$
- rule 6870 $\text{Int}[(a_ + \text{ArcCsch}[(c_ + (d_)*(x_)]*(b_))^{(p_)}), x_Symbol] \rightarrow \text{Simp}[1/d \text{Subst}[\text{Int}[(a + b*\text{ArcCsch}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{IGtQ}[p, 0]$

Maple [F]

$$\int (a + b \operatorname{arccsch}(dx + c))^2 dx$$

input `int((a+b*arccsch(d*x+c))^2,x)`

output `int((a+b*arccsch(d*x+c))^2,x)`

Fricas [F]

$$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \int (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

input `integrate((a+b*arccsch(d*x+c))^2,x, algorithm="fricas")`

output `integral(b^2*arccsch(d*x + c)^2 + 2*a*b*arccsch(d*x + c) + a^2, x)`

Sympy [F]

$$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \int (a + b \operatorname{acsch}(c + dx))^2 dx$$

input `integrate((a+b*acsch(d*x+c))**2,x)`

output `Integral((a + b*acsch(c + d*x))**2, x)`

Maxima [F]

$$\int (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

input `integrate((a+b*arccsch(d*x+c))^2,x, algorithm="maxima")`

output `(x*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1)^2 - integrate(-((d^2*x^2 + 2*c*d*x + c^2 + 1)^(3/2)*log(d*x + c)^2 + (d^2*x^2 + 2*c*d*x + c^2 + 1)*log(d*x + c)^2 - 2*((d^2*x^2 + 2*c*d*x + c^2 + 1)*log(d*x + c) + sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1)*(d^2*x^2 + c*d*x + (d^2*x^2 + 2*c*d*x + c^2 + 1)*log(d*x + c)))*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1))/(d^2*x^2 + 2*c*d*x + c^2 + (d^2*x^2 + 2*c*d*x + c^2 + 1)^(3/2) + 1), x))*b^2 + a^2*x + (2*(d*x + c)*arccsch(d*x + c) + log(sqrt(1/(d*x + c)^2 + 1) + 1) - log(sqrt(1/(d*x + c)^2 + 1) - 1))*a*b/d`

Giac [F]

$$\int (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

input `integrate((a+b*arccsch(d*x+c))^2,x, algorithm="giac")`

output `integrate((b*arccsch(d*x + c) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int \left(a + b \operatorname{asinh}\left(\frac{1}{c + dx}\right) \right)^2 dx$$

input `int((a + b*asinh(1/(c + d*x)))^2,x)`

output `int((a + b*asinh(1/(c + d*x)))^2, x)`

Reduce [F]

$$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = 2 \left(\int \operatorname{acsch}(dx + c) dx \right) ab + \left(\int \operatorname{acsch}(dx + c)^2 dx \right) b^2 + a^2 x$$

input `int((a+b*acsch(d*x+c))^2,x)`

output `2*int(acsch(c + d*x),x)*a*b + int(acsch(c + d*x)**2,x)*b**2 + a**2*x`

$$3.11 \quad \int \frac{\left(a+b\operatorname{csch}^{-1}(c+dx)\right)^2}{e+fx} dx$$

Optimal result	120
Mathematica [F]	121
Rubi [C] (verified)	121
Maple [F]	127
Fricas [F]	128
Sympy [F]	128
Maxima [F]	128
Giac [F]	129
Mupad [F(-1)]	129
Reduce [F]	129

Optimal result

Integrand size = 20, antiderivative size = 475

$$\begin{aligned}
& \int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{e + fx} dx \\
&= - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2 \log\left(1 - e^{2 \operatorname{csch}^{-1}(c + dx)}\right)}{f} \\
&+ \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2 \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f} \\
&+ \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2 \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f} \\
&- \frac{b(a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, e^{2 \operatorname{csch}^{-1}(c + dx)}\right)}{f} \\
&+ \frac{2b(a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f} \\
&+ \frac{2b(a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f} \\
&+ \frac{b^2 \operatorname{PolyLog}\left(3, e^{2 \operatorname{csch}^{-1}(c + dx)}\right)}{2f} - \frac{2b^2 \operatorname{PolyLog}\left(3, -\frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f} \\
&- \frac{2b^2 \operatorname{PolyLog}\left(3, -\frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f}
\end{aligned}$$

output

```

-(a+b*arccsch(d*x+c))^2*ln(1-(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))^2)/f+(a+b*arccsch(d*x+c))^2*ln(1+(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2)))*(-c*f+d*e)/(f-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2))/f+(a+b*arccsch(d*x+c))^2*ln(1+(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2)))*(-c*f+d*e)/(f+(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2))/f-b*(a+b*arccsch(d*x+c))*polylog(2,(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))^2)/f+2*b*(a+b*arccsch(d*x+c))*polylog(2,-(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2)))*(-c*f+d*e)/(f-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2))/f+2*b*(a+b*arccsch(d*x+c))*polylog(2,-(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2)))*(-c*f+d*e)/(f+(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2))/f+1/2*b^2*polylog(3,(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))^2)/f-2*b^2*polylog(3,-(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2)))*(-c*f+d*e)/(f-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2))/f-2*b^2*polylog(3,-(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2)))*(-c*f+d*e)/(f+(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2))/f

```

Mathematica [F]

$$\int \frac{(a + b \operatorname{arcsch}^{-1}(c + dx))^2}{e + fx} dx = \int \frac{(a + b \operatorname{arcsch}^{-1}(c + dx))^2}{e + fx} dx$$

input

```
Integrate[(a + b*ArcCsch[c + d*x])^2/(e + f*x), x]
```

output

```
Integrate[(a + b*ArcCsch[c + d*x])^2/(e + f*x), x]
```

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 2.60 (sec) , antiderivative size = 568, normalized size of antiderivative = 1.20, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6876, 6130, 6103, 3042, 26, 4199, 25, 2620, 3011, 2720, 6095, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{(a + b\operatorname{csch}^{-1}(c + dx))^2}{e + fx} dx \\
& \quad \downarrow \text{6876} \\
& - \int \frac{(c + dx)^2 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b\operatorname{csch}^{-1}(c + dx))^2}{de - cf + f(c + dx)} d\operatorname{csch}^{-1}(c + dx) \\
& \quad \downarrow \text{6130} \\
& - \int \frac{(c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b\operatorname{csch}^{-1}(c + dx))^2}{f + \frac{de - cf}{c + dx}} d\operatorname{csch}^{-1}(c + dx) \\
& \quad \downarrow \text{6103} \\
& \frac{(de - cf) \int \frac{\sqrt{1 + \frac{1}{(c+dx)^2}} (a + b\operatorname{csch}^{-1}(c + dx))^2}{f + \frac{de - cf}{c + dx}} d\operatorname{csch}^{-1}(c + dx)}{f} \\
& \frac{\int (c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b\operatorname{csch}^{-1}(c + dx))^2 d\operatorname{csch}^{-1}(c + dx)}{f} \\
& \quad \downarrow \text{3042} \\
& \frac{(de - cf) \int \frac{\sqrt{1 + \frac{1}{(c+dx)^2}} (a + b\operatorname{csch}^{-1}(c + dx))^2}{f + \frac{de - cf}{c + dx}} d\operatorname{csch}^{-1}(c + dx)}{f} \\
& \frac{\int -i(a + b\operatorname{csch}^{-1}(c + dx))^2 \tan\left(i\operatorname{csch}^{-1}(c + dx) + \frac{\pi}{2}\right) d\operatorname{csch}^{-1}(c + dx)}{f} \\
& \quad \downarrow \text{26} \\
& \frac{(de - cf) \int \frac{\sqrt{1 + \frac{1}{(c+dx)^2}} (a + b\operatorname{csch}^{-1}(c + dx))^2}{f + \frac{de - cf}{c + dx}} d\operatorname{csch}^{-1}(c + dx)}{f} \\
& \frac{i \int (a + b\operatorname{csch}^{-1}(c + dx))^2 \tan\left(i\operatorname{csch}^{-1}(c + dx) + \frac{\pi}{2}\right) d\operatorname{csch}^{-1}(c + dx)}{f} \\
& \quad \downarrow \text{4199} \\
& \frac{(de - cf) \int \frac{\sqrt{1 + \frac{1}{(c+dx)^2}} (a + b\operatorname{csch}^{-1}(c + dx))^2}{f + \frac{de - cf}{c + dx}} d\operatorname{csch}^{-1}(c + dx)}{f} \\
& \frac{i \left(2i \int -\frac{e^{2\operatorname{csch}^{-1}(c + dx)} (a + b\operatorname{csch}^{-1}(c + dx))^2}{1 - e^{2\operatorname{csch}^{-1}(c + dx)}} d\operatorname{csch}^{-1}(c + dx) - \frac{i(a + b\operatorname{csch}^{-1}(c + dx))^3}{3b} \right)}{f}
\end{aligned}$$

$$\begin{array}{c}
 \downarrow 25 \\
 \frac{(de - cf) \int \frac{\sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c+dx))^2}{f + \frac{de - cf}{c+dx}} d \operatorname{csch}^{-1}(c + dx)}{f} + \\
 i \left(-2i \int \frac{e^{2 \operatorname{csch}^{-1}(c+dx)} (a + b \operatorname{csch}^{-1}(c+dx))^2}{1 - e^{2 \operatorname{csch}^{-1}(c+dx)}} d \operatorname{csch}^{-1}(c + dx) - \frac{i (a + b \operatorname{csch}^{-1}(c+dx))^3}{3b} \right) \\
 \hline
 \downarrow 2620 \\
 \frac{(de - cf) \int \frac{\sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c+dx))^2}{f + \frac{de - cf}{c+dx}} d \operatorname{csch}^{-1}(c + dx)}{f} + \\
 i \left(-2i \left(b \int (a + b \operatorname{csch}^{-1}(c + dx)) \log \left(1 - e^{2 \operatorname{csch}^{-1}(c+dx)} \right) d \operatorname{csch}^{-1}(c + dx) - \frac{1}{2} \log \left(1 - e^{2 \operatorname{csch}^{-1}(c+dx)} \right) (a + b \operatorname{csch}^{-1}(c + dx)) \right) \right) \\
 \hline
 \downarrow 3011 \\
 \frac{(de - cf) \int \frac{\sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c+dx))^2}{f + \frac{de - cf}{c+dx}} d \operatorname{csch}^{-1}(c + dx)}{f} + \\
 i \left(-2i \left(b \left(\frac{1}{2} b \int \operatorname{PolyLog} \left(2, e^{2 \operatorname{csch}^{-1}(c+dx)} \right) d \operatorname{csch}^{-1}(c + dx) - \frac{1}{2} \operatorname{PolyLog} \left(2, e^{2 \operatorname{csch}^{-1}(c+dx)} \right) (a + b \operatorname{csch}^{-1}(c + dx)) \right) \right) \right) \\
 \hline
 \downarrow 2720 \\
 \frac{(de - cf) \int \frac{\sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c+dx))^2}{f + \frac{de - cf}{c+dx}} d \operatorname{csch}^{-1}(c + dx)}{f} + \\
 i \left(-2i \left(b \left(\frac{1}{4} b \int e^{-2 \operatorname{csch}^{-1}(c+dx)} \operatorname{PolyLog} \left(2, e^{2 \operatorname{csch}^{-1}(c+dx)} \right) d e^{2 \operatorname{csch}^{-1}(c+dx)} - \frac{1}{2} \operatorname{PolyLog} \left(2, e^{2 \operatorname{csch}^{-1}(c+dx)} \right) (a + b \operatorname{csch}^{-1}(c + dx)) \right) \right) \right) \\
 \hline
 \downarrow 6095
 \end{array}$$

$$(de - cf) \left(\int \frac{e^{\operatorname{csch}^{-1}(c+dx)} (a+b\operatorname{csch}^{-1}(c+dx))^2}{f+e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)-\sqrt{d^2e^2-2cdf e+c^2f^2+f^2}} d\operatorname{csch}^{-1}(c+dx) + \int \frac{e^{\operatorname{csch}^{-1}(c+dx)} (a+b\operatorname{csch}^{-1}(c+dx))^2}{f+e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)+\sqrt{d^2e^2-2cdf e+c^2f^2+f^2}} d\operatorname{csch}^{-1}(c+dx) \right)$$

$$i \left(-2i \left(b \left(\frac{1}{4} b \int e^{-2\operatorname{csch}^{-1}(c+dx)} \operatorname{PolyLog} \left(2, e^{2\operatorname{csch}^{-1}(c+dx)} \right) de^{2\operatorname{csch}^{-1}(c+dx)} - \frac{1}{2} \operatorname{PolyLog} \left(2, e^{2\operatorname{csch}^{-1}(c+dx)} \right) (a + b\operatorname{csch}^{-1}(c+dx)) \right) \right)$$

f

↓ 2620

$$(de - cf) \left(- \frac{2b \int (a+b\operatorname{csch}^{-1}(c+dx)) \log \left(\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f-\sqrt{d^2e^2-2cdf e+(c^2+1)f^2}} + 1 \right) d\operatorname{csch}^{-1}(c+dx)}{de-cf} - \frac{2b \int (a+b\operatorname{csch}^{-1}(c+dx)) \log \left(\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f+\sqrt{d^2e^2-2cdf e+(c^2+1)f^2}} + 1 \right) d\operatorname{csch}^{-1}(c+dx)}{de-cf} \right)$$

$$i \left(-2i \left(b \left(\frac{1}{4} b \int e^{-2\operatorname{csch}^{-1}(c+dx)} \operatorname{PolyLog} \left(2, e^{2\operatorname{csch}^{-1}(c+dx)} \right) de^{2\operatorname{csch}^{-1}(c+dx)} - \frac{1}{2} \operatorname{PolyLog} \left(2, e^{2\operatorname{csch}^{-1}(c+dx)} \right) (a + b\operatorname{csch}^{-1}(c+dx)) \right) \right)$$

f

↓ 3011

$$(de - cf) \left(- \frac{2b \left(b \int \operatorname{PolyLog} \left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f-\sqrt{d^2e^2-2cdf e+(c^2+1)f^2}} \right) d\operatorname{csch}^{-1}(c+dx) - (a+b\operatorname{csch}^{-1}(c+dx)) \operatorname{PolyLog} \left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f-\sqrt{d^2e^2-2cdf e+(c^2+1)f^2}} \right) \right)}{de-cf} \right)$$

$$i \left(-2i \left(b \left(\frac{1}{4} b \int e^{-2\operatorname{csch}^{-1}(c+dx)} \operatorname{PolyLog} \left(2, e^{2\operatorname{csch}^{-1}(c+dx)} \right) de^{2\operatorname{csch}^{-1}(c+dx)} - \frac{1}{2} \operatorname{PolyLog} \left(2, e^{2\operatorname{csch}^{-1}(c+dx)} \right) (a + b\operatorname{csch}^{-1}(c+dx)) \right) \right)$$

f

↓ 2720

$$(de - cf) \left(- \frac{2b \left(b \int e^{-\operatorname{csch}^{-1}(c+dx)} \operatorname{PolyLog} \left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f-\sqrt{d^2e^2-2cdf e+(c^2+1)f^2}} \right) de^{\operatorname{csch}^{-1}(c+dx)} - (a+b\operatorname{csch}^{-1}(c+dx)) \operatorname{PolyLog} \left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f-\sqrt{d^2e^2-2cdf e+(c^2+1)f^2}} \right) \right)}{de-cf} \right)$$

$$i \left(-2i \left(b \left(\frac{1}{4} b \int e^{-2\operatorname{csch}^{-1}(c+dx)} \operatorname{PolyLog} \left(2, e^{2\operatorname{csch}^{-1}(c+dx)} \right) de^{2\operatorname{csch}^{-1}(c+dx)} - \frac{1}{2} \operatorname{PolyLog} \left(2, e^{2\operatorname{csch}^{-1}(c+dx)} \right) (a + b\operatorname{csch}^{-1}(c+dx)) \right) \right)$$

f

↓ 7143

$$(de - cf) \left(-\frac{2b \left(b \operatorname{PolyLog} \left(3, -\frac{e^{\operatorname{csch}^{-1}(c+dx)(de-cf)}}}{f - \sqrt{d^2 e^2 - 2cdf e + (c^2 + 1)f^2}} \right) - (a + b \operatorname{csch}^{-1}(c+dx)) \operatorname{PolyLog} \left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)(de-cf)}}}{f - \sqrt{d^2 e^2 - 2cdf e + (c^2 + 1)f^2}} \right) \right)}{de - cf} - \dots \right)$$

$$i \left(-2i \left(b \left(\frac{1}{4} b \operatorname{PolyLog} \left(3, e^{2 \operatorname{csch}^{-1}(c+dx)} \right) - \frac{1}{2} \operatorname{PolyLog} \left(2, e^{2 \operatorname{csch}^{-1}(c+dx)} \right) (a + b \operatorname{csch}^{-1}(c + dx)) \right) \right) - \frac{1}{2} \log \left(1 - e^{2 \dots} \right) \right)$$

f

```
input Int[(a + b*ArcCsch[c + d*x])^2/(e + f*x),x]
```

```
output (I*((( -1/3*I)*(a + b*ArcCsch[c + d*x])^3)/b - (2*I)*(-1/2*((a + b*ArcCsch[c + d*x])^2*Log[1 - E^(2*ArcCsch[c + d*x])]) + b*(-1/2*((a + b*ArcCsch[c + d*x])*PolyLog[2, E^(2*ArcCsch[c + d*x])]) + (b*PolyLog[3, E^(2*ArcCsch[c + d*x])]))/4))))/f + ((d*e - c*f)*(-1/3*(a + b*ArcCsch[c + d*x])^3/(b*(d*e - c*f)) + ((a + b*ArcCsch[c + d*x])^2*Log[1 + (E^ArcCsch[c + d*x]*(d*e - c*f))/(f - Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]]))/(d*e - c*f) + ((a + b*ArcCsch[c + d*x])^2*Log[1 + (E^ArcCsch[c + d*x]*(d*e - c*f))/(f + Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]]))/(d*e - c*f) - (2*b*((a + b*ArcCsch[c + d*x])*PolyLog[2, -((E^ArcCsch[c + d*x]*(d*e - c*f))/(f - Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2])]) + b*PolyLog[3, -((E^ArcCsch[c + d*x]*(d*e - c*f))/(f - Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2])])])/(d*e - c*f) - (2*b*((a + b*ArcCsch[c + d*x])*PolyLog[2, -((E^ArcCsch[c + d*x]*(d*e - c*f))/(f + Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2])]) + b*PolyLog[3, -((E^ArcCsch[c + d*x]*(d*e - c*f))/(f + Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2])])])/(d*e - c*f)))/f
```

Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 26 Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)
*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4199

```
Int[(((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_
.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp
[2*I Int[(((c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x
)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && In
tegerQ[4*k] && IGtQ[m, 0]
```

rule 6095

```
Int[(Cosh[(c_) + (d_)*(x_)]*((e_) + (f_)*(x_))^(m_))/((a_) + (b_)*Sin
h[(c_) + (d_)*(x_)]), x_Symbol] := Simp[-(e + f*x)^(m + 1)/(b*f*(m + 1)),
x] + (Int[(e + f*x)^m*(E^(c + d*x)/(a - Rt[a^2 + b^2, 2] + b*E^(c + d*x))
, x] + Int[(e + f*x)^m*(E^(c + d*x)/(a + Rt[a^2 + b^2, 2] + b*E^(c + d*x))
, x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[a^2 + b^2, 0]
```

rule 6103 `Int[(Coth[(c_.) + (d_.)*(x_)]^(n_.)*((e_.) + (f_.)*(x_))^(m_.))/((a_.) + (b_.)*Sinh[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[1/a Int[(e + f*x)^m*Coth[c + d*x]^n, x], x] - Simp[b/a Int[(e + f*x)^m*Cosh[c + d*x]*(Coth[c + d*x]^(n - 1)/(a + b*Sinh[c + d*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

rule 6130 `Int[(((e_.) + (f_.)*(x_))^(m_.)*(F_)[(c_.) + (d_.)*(x_)]^(n_.)*(G_)[(c_.) + (d_.)*(x_)]^(p_.))/(Csch[(c_.) + (d_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Int[(e + f*x)^m*Sinh[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Sinh[c + d*x])), x] /; FreeQ[{a, b, c, d, e, f}, x] && HyperbolicQ[F] && HyperbolicQ[G] && IntegersQ[m, n, p]`

rule 6876 `Int[((a_.) + ArcCsch[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Csch[x]*Coth[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

Maple [F]

$$\int \frac{(a + b \operatorname{arccsch}(dx + c))^2}{fx + e} dx$$

input `int((a+b*arccsch(d*x+c))^2/(f*x+e),x)`

output `int((a+b*arccsch(d*x+c))^2/(f*x+e),x)`

Fricas [F]

$$\int \frac{(a + b \operatorname{arcsch}^{-1}(c + dx))^2}{e + fx} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{fx + e} dx$$

input `integrate((a+b*arccsch(d*x+c))^2/(f*x+e),x, algorithm="fricas")`

output `integral((b^2*arccsch(d*x + c)^2 + 2*a*b*arccsch(d*x + c) + a^2)/(f*x + e), x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arcsch}^{-1}(c + dx))^2}{e + fx} dx = \int \frac{(a + b \operatorname{acsch}(c + dx))^2}{e + fx} dx$$

input `integrate((a+b*acsch(d*x+c))**2/(f*x+e),x)`

output `Integral((a + b*acsch(c + d*x))**2/(e + f*x), x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arcsch}^{-1}(c + dx))^2}{e + fx} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{fx + e} dx$$

input `integrate((a+b*arccsch(d*x+c))^2/(f*x+e),x, algorithm="maxima")`

output `a^2*log(f*x + e)/f + integrate(b^2*log(sqrt(1/(d*x + c)^2 + 1) + 1/(d*x + c))^2/(f*x + e) + 2*a*b*log(sqrt(1/(d*x + c)^2 + 1) + 1/(d*x + c))/(f*x + e), x)`

Giac [F]

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{e + fx} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{fx + e} dx$$

input `integrate((a+b*arccsch(d*x+c))^2/(f*x+e),x, algorithm="giac")`

output `integrate((b*arccsch(d*x + c) + a)^2/(f*x + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{e + fx} dx = \int \frac{(a + b \operatorname{asinh}(\frac{1}{c+dx}))^2}{e + fx} dx$$

input `int((a + b*asinh(1/(c + d*x)))^2/(e + f*x),x)`

output `int((a + b*asinh(1/(c + d*x)))^2/(e + f*x), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{e + fx} dx \\ &= \frac{2 \left(\int \frac{\operatorname{acsch}(dx+c)}{fx+e} dx \right) abf + \left(\int \frac{\operatorname{acsch}(dx+c)^2}{fx+e} dx \right) b^2 f + \log(fx + e) a^2}{f} \end{aligned}$$

input `int((a+b*acsch(d*x+c))^2/(f*x+e),x)`

output `(2*int(acsch(c + d*x)/(e + f*x),x)*a*b*f + int(acsch(c + d*x)**2/(e + f*x),x)*b**2*f + log(e + f*x)*a**2)/f`

3.12
$$\int \frac{(a+b\operatorname{csch}^{-1}(c+dx))^2}{(e+fx)^2} dx$$

Optimal result	130
Mathematica [C] (warning: unable to verify)	131
Rubi [A] (verified)	132
Maple [F]	135
Fricas [F]	135
Sympy [F]	135
Maxima [F]	136
Giac [F]	136
Mupad [F(-1)]	137
Reduce [F]	137

Optimal result

Integrand size = 20, antiderivative size = 448

$$\begin{aligned} & \int \frac{(a + b\operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^2} dx \\ &= \frac{d(a + b\operatorname{csch}^{-1}(c + dx))^2}{f(de - cf)} - \frac{(a + b\operatorname{csch}^{-1}(c + dx))^2}{f(e + fx)} \\ & \quad - \frac{2bd(a + b\operatorname{csch}^{-1}(c + dx)) \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f - \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdef + (1 + c^2)f^2}} \\ & \quad + \frac{2bd(a + b\operatorname{csch}^{-1}(c + dx)) \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f + \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdef + (1 + c^2)f^2}} \\ & \quad - \frac{2b^2d \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f - \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdef + (1 + c^2)f^2}} \\ & \quad + \frac{2b^2d \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f + \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdef + (1 + c^2)f^2}} \end{aligned}$$

output

```
d*(a+b*arccsch(d*x+c))^2/f/(-c*f+d*e)-(a+b*arccsch(d*x+c))^2/f/(f*x+e)-2*b
*d*(a+b*arccsch(d*x+c))*ln(1+(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))*(-c*f+d*e)/
(f-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/(-c*f+d*e)/(d^2*e^2-2*c*d*e*f+(
c^2+1)*f^2)^(1/2)+2*b*d*(a+b*arccsch(d*x+c))*ln(1+(1/(d*x+c)+(1+1/(d*x+c)^
2)^(1/2))*(-c*f+d*e)/(f+(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/(-c*f+d*e)
/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)-2*b^2*d*polylog(2,-(1/(d*x+c)+(1+1/
(d*x+c)^2)^(1/2))*(-c*f+d*e)/(f-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/(-
c*f+d*e)/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)+2*b^2*d*polylog(2,-(1/(d*x+
c)+(1+1/(d*x+c)^2)^(1/2))*(-c*f+d*e)/(f+(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1
/2)))/(-c*f+d*e)/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)
```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 12.51 (sec) , antiderivative size = 1874, normalized size of antiderivative = 4.18

$$\int \frac{(a + b \operatorname{arcsch}(c + dx))^2}{(e + fx)^2} dx = \text{Too large to display}$$

input

```
Integrate[(a + b*ArcCsch[c + d*x])^2/(e + f*x)^2,x]
```

output

```

-(a^2/(f*(e + f*x))) - (2*a*b*(c + d*x)^2*(f + (d*e - c*f)/(c + d*x))^2*(A
rcCsch[c + d*x]/(f + (d*e)/(c + d*x) - (c*f)/(c + d*x)) - (2*ArcTan[(d*e -
c*f - f*Tanh[ArcCsch[c + d*x]/2])/Sqrt[-(d^2*e^2) + 2*c*d*e*f - (1 + c^2)
*f^2]])/Sqrt[-(d^2*e^2) + 2*c*d*e*f - (1 + c^2)*f^2]))/(d*(-(d*e) + c*f)*(
e + f*x)^2) - (b^2*(c + d*x)^2*(f + (d*e - c*f)/(c + d*x))^2*(ArcCsch[c +
d*x]^2/(((d*e) + c*f)*(f + (d*e - c*f)/(c + d*x))) + (2*(((I)*Pi*ArcTanh
[-(d*e) + c*f + f*Tanh[ArcCsch[c + d*x]/2])/Sqrt[f^2 + (d*e - c*f)^2]))/S
qrt[f^2 + (d*e - c*f)^2] - ((2*I)*ArcCos[(I*f)/(-(d*e) + c*f)]*ArcTan[((d*
e - (I + c)*f)*Cot[(Pi + (2*I)*ArcCsch[c + d*x])/4])/Sqrt[-(d^2*e^2) + 2*c
*d*e*f - (1 + c^2)*f^2]] + (Pi - (2*I)*ArcCsch[c + d*x])*ArcTanh[(((I)*d*
e + f + I*c*f)*Tan[(Pi + (2*I)*ArcCsch[c + d*x])/4])/Sqrt[-(d^2*e^2) + 2*c
*d*e*f - (1 + c^2)*f^2]] + (ArcCos[(I*f)/(-(d*e) + c*f)] + 2*ArcTan[((d*e
- (I + c)*f)*Cot[(Pi + (2*I)*ArcCsch[c + d*x])/4])/Sqrt[-(d^2*e^2) + 2*c*d
*e*f - (1 + c^2)*f^2]] - (2*I)*ArcTanh[(((I)*d*e + f + I*c*f)*Tan[(Pi + (
2*I)*ArcCsch[c + d*x])/4])/Sqrt[-(d^2*e^2) + 2*c*d*e*f - (1 + c^2)*f^2]])*
Log[-(((I)*(-1)^(3/4)*Sqrt[-(d^2*e^2) + 2*c*d*e*f - (1 + c^2)*f^2])/(Sqrt[2]*E
^((ArcCsch[c + d*x]/2)*Sqrt[I*(-(d*e) + c*f)]*Sqrt[f + (d*e - c*f)/(c + d*x
)])))] + (ArcCos[(I*f)/(-(d*e) + c*f)] - 2*ArcTan[((d*e - (I + c)*f)*Cot[(P
i + (2*I)*ArcCsch[c + d*x])/4])/Sqrt[-(d^2*e^2) + 2*c*d*e*f - (1 + c^2)*f^
2]] + (2*I)*ArcTanh[(((I)*d*e + f + I*c*f)*Tan[(Pi + (2*I)*ArcCsch[c +...

```

Rubi [A] (verified)

Time = 1.37 (sec) , antiderivative size = 461, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6876, 5992, 3042, 4679, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^2} dx$$

$$\downarrow 6876$$

$$-d \int \frac{(c + dx)^2 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))^2}{(de - cf + f(c + dx))^2} d \operatorname{csch}^{-1}(c + dx)$$

$$\downarrow 5992$$

$$\begin{aligned}
 & -d \left(\frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(f(c + dx) - cf + de)} - \frac{2b \int \frac{a + b \operatorname{csch}^{-1}(c + dx)}{de - cf + f(c + dx)} d \operatorname{csch}^{-1}(c + dx)}{f} \right) \\
 & \quad \downarrow \text{3042} \\
 & -d \left(\frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(f(c + dx) - cf + de)} - \frac{2b \int \frac{a + b \operatorname{csch}^{-1}(c + dx)}{de - cf + i f \operatorname{csc}(i \operatorname{csch}^{-1}(c + dx))} d \operatorname{csch}^{-1}(c + dx)}{f} \right) \\
 & \quad \downarrow \text{4679} \\
 & -d \left(\frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(f(c + dx) - cf + de)} - \frac{2b \int \left(\frac{a + b \operatorname{csch}^{-1}(c + dx)}{de - cf} + \frac{f(a + b \operatorname{csch}^{-1}(c + dx))}{(cf - de) \left(f + \frac{de(1 - \frac{cf}{de})}{c + dx} \right)} \right) d \operatorname{csch}^{-1}(c + dx)}{f} \right) \\
 & \quad \downarrow \text{2009} \\
 & -d \left(\frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(f(c + dx) - cf + de)} - \frac{2b \left(-\frac{f(a + b \operatorname{csch}^{-1}(c + dx)) \log \left(\frac{(de - cf)e^{\operatorname{csch}^{-1}(c + dx)}}{f - \sqrt{(c^2 + 1)f^2 - 2cdef + d^2e^2}} + 1 \right)}{(de - cf)\sqrt{(c^2 + 1)f^2 - 2cdef + d^2e^2}} + \frac{f(a + b \operatorname{csch}^{-1}(c + dx)) \log \left(\frac{(de - cf)e^{\operatorname{csch}^{-1}(c + dx)}}{f - \sqrt{(c^2 + 1)f^2 - 2cdef + d^2e^2}} + 1 \right)}{(de - cf)\sqrt{(c^2 + 1)f^2 - 2cdef + d^2e^2}} \right)}{f} \right)
 \end{aligned}$$

input `Int[(a + b*ArcCsch[c + d*x])^2/(e + f*x)^2,x]`

output

```

-(d*((a + b*ArcCsch[c + d*x])^2/(f*(d*e - c*f + f*(c + d*x))) - (2*b*((a +
b*ArcCsch[c + d*x])^2/(2*b*(d*e - c*f)) - (f*(a + b*ArcCsch[c + d*x])*Log
[1 + (E^ArcCsch[c + d*x]*(d*e - c*f))/(f - Sqrt[d^2*e^2 - 2*c*d*e*f + (1 +
c^2)*f^2])))/(d*e - c*f)*Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]) + (f
*(a + b*ArcCsch[c + d*x])*Log[1 + (E^ArcCsch[c + d*x]*(d*e - c*f))/(f + Sq
rt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2])))/((d*e - c*f)*Sqrt[d^2*e^2 - 2*c
*d*e*f + (1 + c^2)*f^2]) - (b*f*PolyLog[2, -((E^ArcCsch[c + d*x]*(d*e - c*
f))/(f - Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2])))/((d*e - c*f)*Sqrt[d
^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]) + (b*f*PolyLog[2, -((E^ArcCsch[c + d*
x]*(d*e - c*f))/(f + Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2])))/((d*e -
c*f)*Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]))/f))

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4679

```
Int[(csc[(e_) + (f_)*(x_)]*(b_) + (a_)^(n_))*((c_) + (d_)*(x_))^(m_)
, x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Si
n[e + f*x]^n)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGt
Q[m, 0]
```

rule 5992

```
Int[Coth[(c_) + (d_)*(x_)]*Csch[(c_) + (d_)*(x_)]*(Csch[(c_) + (d_)*(
x_)]*(b_) + (a_)^(n_))*((e_) + (f_)*(x_))^(m_), x_Symbol] := Simp[(-(e
+ f*x)^m)*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b
*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

rule 6876

```
Int[((a_) + ArcCsch[(c_) + (d_)*(x_)]*(b_))^(p_))*((e_) + (f_)*(x_))^(
m_), x_Symbol] := Simp[-(d^(m + 1))^( -1) Subst[Int[(a + b*x)^p*Csch[x]*C
oth[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Maple [F]

$$\int \frac{(a + b \operatorname{arccsch}(dx + c))^2}{(fx + e)^2} dx$$

input `int((a+b*arccsch(d*x+c))^2/(f*x+e)^2,x)`

output `int((a+b*arccsch(d*x+c))^2/(f*x+e)^2,x)`

Fricas [F]

$$\int \frac{(a + b \operatorname{bsch}^{-1}(c + dx))^2}{(e + fx)^2} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{(fx + e)^2} dx$$

input `integrate((a+b*arccsch(d*x+c))^2/(f*x+e)^2,x, algorithm="fricas")`

output `integral((b^2*arccsch(d*x + c)^2 + 2*a*b*arccsch(d*x + c) + a^2)/(f^2*x^2 + 2*e*f*x + e^2), x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{bsch}^{-1}(c + dx))^2}{(e + fx)^2} dx = \int \frac{(a + b \operatorname{acsch}(c + dx))^2}{(e + fx)^2} dx$$

input `integrate((a+b*acsch(d*x+c))**2/(f*x+e)**2,x)`

output `Integral((a + b*acsch(c + d*x))**2/(e + f*x)**2, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arcsch}^{-1}(c + dx))^2}{(e + fx)^2} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{(fx + e)^2} dx$$

input `integrate((a+b*arccsch(d*x+c))^2/(f*x+e)^2,x, algorithm="maxima")`

output

```
-b^2*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1)/(f^2*x + e*f) - a^2/(f^2
*x + e*f) - integrate(-((b^2*d^2*f*x^2 + 2*b^2*c*d*f*x + (c^2*f + f)*b^2)*
log(d*x + c)^2 - 2*(a*b*d^2*f*x^2 + 2*a*b*c*d*f*x + (c^2*f + f)*a*b)*log(d
*x + c) + 2*(a*b*d^2*f*x^2 + 2*a*b*c*d*f*x + (c^2*f + f)*a*b - (b^2*d^2*f*
x^2 + 2*b^2*c*d*f*x + (c^2*f + f)*b^2)*log(d*x + c) + (b^2*c*d*e + (c^2*f
+ f)*a*b + (a*b*d^2*f + b^2*d^2*f)*x^2 + (2*a*b*c*d*f + (d^2*e + c*d*f)*b^
2)*x - (b^2*d^2*f*x^2 + 2*b^2*c*d*f*x + (c^2*f + f)*b^2)*log(d*x + c))*sq
rt(d^2*x^2 + 2*c*d*x + c^2 + 1))*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1)
+ sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1)*((b^2*d^2*f*x^2 + 2*b^2*c*d*f*x + (c^
2*f + f)*b^2)*log(d*x + c)^2 - 2*(a*b*d^2*f*x^2 + 2*a*b*c*d*f*x + (c^2*f +
f)*a*b)*log(d*x + c)))/(d^2*f^3*x^4 + c^2*e^2*f + 2*(d^2*e*f^2 + c*d*f^3)
*x^3 + e^2*f + (d^2*e^2*f + 4*c*d*e*f^2 + c^2*f^3 + f^3)*x^2 + 2*(c*d*e^2*
f + c^2*e*f^2 + e*f^2)*x + (d^2*f^3*x^4 + c^2*e^2*f + 2*(d^2*e*f^2 + c*d*f
^3)*x^3 + e^2*f + (d^2*e^2*f + 4*c*d*e*f^2 + c^2*f^3 + f^3)*x^2 + 2*(c*d*e
^2*f + c^2*e*f^2 + e*f^2)*x)*sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1)), x)
```

Giac [F]

$$\int \frac{(a + b \operatorname{arcsch}^{-1}(c + dx))^2}{(e + fx)^2} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{(fx + e)^2} dx$$

input `integrate((a+b*arccsch(d*x+c))^2/(f*x+e)^2,x, algorithm="giac")`

output

```
integrate((b*arccsch(d*x + c) + a)^2/(f*x + e)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^2} dx = \int \frac{(a + b \operatorname{asinh}(\frac{1}{c+dx}))^2}{(e + fx)^2} dx$$

input `int((a + b*asinh(1/(c + d*x)))^2/(e + f*x)^2,x)`output `int((a + b*asinh(1/(c + d*x)))^2/(e + f*x)^2, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^2} dx$$

$$= \frac{2 \left(\int \frac{\operatorname{acsch}(dx+c)}{f^2 x^2 + 2efx + e^2} dx \right) ab e^2 + 2 \left(\int \frac{\operatorname{acsch}(dx+c)}{f^2 x^2 + 2efx + e^2} dx \right) abefx + \left(\int \frac{\operatorname{acsch}(dx+c)^2}{f^2 x^2 + 2efx + e^2} dx \right) b^2 e^2 + \left(\int \frac{\operatorname{acsch}(dx+c)^2}{f^2 x^2 + 2efx + e^2} dx \right) b^2 e^2}{e(fx + e)}$$

input `int((a+b*acsch(d*x+c))^2/(f*x+e)^2,x)`output `(2*int(acsch(c + d*x)/(e**2 + 2*e*f*x + f**2*x**2),x)*a*b*e**2 + 2*int(acsch(c + d*x)/(e**2 + 2*e*f*x + f**2*x**2),x)*a*b*e*f*x + int(acsch(c + d*x)**2/(e**2 + 2*e*f*x + f**2*x**2),x)*b**2*e**2 + int(acsch(c + d*x)**2/(e**2 + 2*e*f*x + f**2*x**2),x)*b**2*e*f*x + a**2*x)/(e*(e + f*x))`

$$3.13 \quad \int \frac{\left(a + b \operatorname{csch}^{-1}(c + dx)\right)^2}{(e + fx)^3} dx$$

Optimal result	138
Mathematica [C] (warning: unable to verify)	139
Rubi [A] (verified)	140
Maple [F]	142
Fricas [F]	143
Sympy [F]	143
Maxima [F]	143
Giac [F]	144
Mupad [F(-1)]	145
Reduce [F]	145

Optimal result

Integrand size = 20, antiderivative size = 1024

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^3} dx = \text{Too large to display}$$

output

```

-b*d^2*f*(1+1/(d*x+c)^2)^(1/2)*(a+b*arccsch(d*x+c))/(-c*f+d*e)/(d^2*e^2-2*
c*d*e*f+(c^2+1)*f^2)/(f+(-c*f+d*e)/(d*x+c))+1/2*d^2*(a+b*arccsch(d*x+c))^2
/f/(-c*f+d*e)^2-1/2*(a+b*arccsch(d*x+c))^2/f/(f*x+e)^2+b*d^2*f^2*(a+b*arcc
sch(d*x+c))*ln(1+(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))*(-c*f+d*e)/(f-(d^2*e^2-
2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/(-c*f+d*e)^2/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2
)^(3/2)-2*b*d^2*(a+b*arccsch(d*x+c))*ln(1+(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2)
))*(-c*f+d*e)/(f-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/(-c*f+d*e)^2/(d^2*
e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)-b*d^2*f^2*(a+b*arccsch(d*x+c))*ln(1+(1/(d
*x+c)+(1+1/(d*x+c)^2)^(1/2))*(-c*f+d*e)/(f+(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)
^(1/2)))/(-c*f+d*e)^2/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(3/2)+2*b*d^2*(a+b*a
rccsch(d*x+c))*ln(1+(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))*(-c*f+d*e)/(f+(d^2*e
^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/(-c*f+d*e)^2/(d^2*e^2-2*c*d*e*f+(c^2+1)*
f^2)^(1/2)+b^2*d^2*f*ln(f+(-c*f+d*e)/(d*x+c))/(-c*f+d*e)^2/(d^2*e^2-2*c*d*
e*f+(c^2+1)*f^2)+b^2*d^2*f^2*polylog(2,-(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))*
(-c*f+d*e)/(f-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/(-c*f+d*e)^2/(d^2*e^
2-2*c*d*e*f+(c^2+1)*f^2)^(3/2)-2*b^2*d^2*polylog(2,-(1/(d*x+c)+(1+1/(d*x+c)
)^2)^(1/2))*(-c*f+d*e)/(f-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/(-c*f+d*
e)^2/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)-b^2*d^2*f^2*polylog(2,-(1/(d*x+
c)+(1+1/(d*x+c)^2)^(1/2))*(-c*f+d*e)/(f+(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1
/2)))/(-c*f+d*e)^2/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(3/2)+2*b^2*d^2*poly...

```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 13.27 (sec) , antiderivative size = 8350, normalized size of antiderivative = 8.15

$$\int \frac{(a + b \operatorname{arcsch}(c + dx))^2}{(e + fx)^3} dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*ArcCsch[c + d*x])^2/(e + f*x)^3,x]
```

output

Result too large to show

Rubi [A] (verified)

Time = 2.45 (sec) , antiderivative size = 1006, normalized size of antiderivative = 0.98, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6876, 5992, 3042, 4679, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^3} dx \\
 & \quad \downarrow \text{6876} \\
 & -d^2 \int \frac{(c + dx)^2 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))^2}{(de - cf + f(c + dx))^3} d \operatorname{csch}^{-1}(c + dx) \\
 & \quad \downarrow \text{5992} \\
 & -d^2 \left(\frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{2f(f(c + dx) - cf + de)^2} - \frac{b \int \frac{a + b \operatorname{csch}^{-1}(c + dx)}{(de - cf + f(c + dx))^2} d \operatorname{csch}^{-1}(c + dx)}{f} \right) \\
 & \quad \downarrow \text{3042} \\
 & -d^2 \left(\frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{2f(f(c + dx) - cf + de)^2} - \frac{b \int \frac{a + b \operatorname{csch}^{-1}(c + dx)}{(de - cf + if \operatorname{csc}(i \operatorname{csch}^{-1}(c + dx)))^2} d \operatorname{csch}^{-1}(c + dx)}{f} \right) \\
 & \quad \downarrow \text{4679} \\
 & -d^2 \left(\frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{2f(f(c + dx) - cf + de)^2} - \frac{b \int \left(\frac{(a + b \operatorname{csch}^{-1}(c + dx)) f^2}{(de - cf)^2 \left(f + \frac{de(1 - \frac{cf}{de})}{c + dx} \right)^2} + \frac{2(a + b \operatorname{csch}^{-1}(c + dx)) f}{(de - cf)^2 \left(-f - \frac{de(1 - \frac{cf}{de})}{c + dx} \right)} + \frac{a + b \operatorname{csch}^{-1}(c + dx)}{(de - cf)^2} \right) dc}{f} \right) \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4679 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Sin[e + f*x]^n)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGtQ[m, 0]`

rule 5992 `Int[Coth[(c_.) + (d_.)*(x_)]*Csch[(c_.) + (d_.)*(x_)]*(Csch[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

rule 6876 `Int[((a_.) + ArcCsch[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[-(d^(m + 1))^(-1) Subst[Int[(a + b*x)^p*Csch[x]*Coth[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

Maple [F]

$$\int \frac{(a + b \operatorname{arccsch}(dx + c))^2}{(fx + e)^3} dx$$

input `int((a+b*arccsch(d*x+c))^2/(f*x+e)^3,x)`

output `int((a+b*arccsch(d*x+c))^2/(f*x+e)^3,x)`

Fricas [F]

$$\int \frac{(a + b \operatorname{arcsch}^{-1}(c + dx))^2}{(e + fx)^3} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{(fx + e)^3} dx$$

input `integrate((a+b*arccsch(d*x+c))^2/(f*x+e)^3,x, algorithm="fricas")`

output `integral((b^2*arccsch(d*x + c)^2 + 2*a*b*arccsch(d*x + c) + a^2)/(f^3*x^3 + 3*e*f^2*x^2 + 3*e^2*f*x + e^3), x)`

Sympy [F]

$$\int \frac{(a + b \operatorname{arcsch}^{-1}(c + dx))^2}{(e + fx)^3} dx = \int \frac{(a + b \operatorname{acsch}(c + dx))^2}{(e + fx)^3} dx$$

input `integrate((a+b*acsch(d*x+c))**2/(f*x+e)**3,x)`

output `Integral((a + b*acsch(c + d*x))**2/(e + f*x)**3, x)`

Maxima [F]

$$\int \frac{(a + b \operatorname{arcsch}^{-1}(c + dx))^2}{(e + fx)^3} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{(fx + e)^3} dx$$

input `integrate((a+b*arccsch(d*x+c))^2/(f*x+e)^3,x, algorithm="maxima")`

output

```

-1/2*b^2*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1)^2/(f^3*x^2 + 2*e*f^2*x
+ e^2*f) - 1/2*a^2/(f^3*x^2 + 2*e*f^2*x + e^2*f) - integrate(-((b^2*d^2*f
*x^2 + 2*b^2*c*d*f*x + (c^2*f + f)*b^2)*log(d*x + c)^2 - 2*(a*b*d^2*f*x^2
+ 2*a*b*c*d*f*x + (c^2*f + f)*a*b)*log(d*x + c) + (2*a*b*d^2*f*x^2 + 4*a*b
*c*d*f*x + 2*(c^2*f + f)*a*b - 2*(b^2*d^2*f*x^2 + 2*b^2*c*d*f*x + (c^2*f +
f)*b^2)*log(d*x + c) + (b^2*c*d*e + 2*(c^2*f + f)*a*b + (2*a*b*d^2*f + b^
2*d^2*f)*x^2 + (4*a*b*c*d*f + (d^2*e + c*d*f)*b^2)*x - 2*(b^2*d^2*f*x^2 +
2*b^2*c*d*f*x + (c^2*f + f)*b^2)*log(d*x + c))*sqrt(d^2*x^2 + 2*c*d*x + c^
2 + 1))*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1) + sqrt(d^2*x^2 + 2*c*d*
x + c^2 + 1)*((b^2*d^2*f*x^2 + 2*b^2*c*d*f*x + (c^2*f + f)*b^2)*log(d*x +
c)^2 - 2*(a*b*d^2*f*x^2 + 2*a*b*c*d*f*x + (c^2*f + f)*a*b)*log(d*x + c)))/
(d^2*f^4*x^5 + c^2*e^3*f + (3*d^2*e*f^3 + 2*c*d*f^4)*x^4 + e^3*f + (3*d^2*
e^2*f^2 + 6*c*d*e*f^3 + c^2*f^4 + f^4)*x^3 + (d^2*e^3*f + 6*c*d*e^2*f^2 +
3*c^2*e*f^3 + 3*e*f^3)*x^2 + (2*c*d*e^3*f + 3*c^2*e^2*f^2 + 3*e^2*f^2)*x +
(d^2*f^4*x^5 + c^2*e^3*f + (3*d^2*e*f^3 + 2*c*d*f^4)*x^4 + e^3*f + (3*d^2
*e^2*f^2 + 6*c*d*e*f^3 + c^2*f^4 + f^4)*x^3 + (d^2*e^3*f + 6*c*d*e^2*f^2 +
3*c^2*e*f^3 + 3*e*f^3)*x^2 + (2*c*d*e^3*f + 3*c^2*e^2*f^2 + 3*e^2*f^2)*x)
*sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1)), x)

```

Giac [F]

$$\int \frac{(a + b \operatorname{arcsch}^{-1}(c + dx))^2}{(e + fx)^3} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{(fx + e)^3} dx$$

input

```
integrate((a+b*arccsch(d*x+c))^2/(f*x+e)^3,x, algorithm="giac")
```

output

```
integrate((b*arccsch(d*x + c) + a)^2/(f*x + e)^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^3} dx = \int \frac{(a + b \operatorname{asinh}(\frac{1}{c+dx}))^2}{(e + fx)^3} dx$$

input `int((a + b*asinh(1/(c + d*x)))^2/(e + f*x)^3,x)`output `int((a + b*asinh(1/(c + d*x)))^2/(e + f*x)^3, x)`**Reduce [F]**

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^3} dx$$

$$= \frac{4 \left(\int \frac{\operatorname{acsch}(dx+c)}{f^3 x^3 + 3e f^2 x^2 + 3e^2 f x + e^3} dx \right) a b e^2 f + 8 \left(\int \frac{\operatorname{acsch}(dx+c)}{f^3 x^3 + 3e f^2 x^2 + 3e^2 f x + e^3} dx \right) a b e f^2 x + 4 \left(\int \frac{\operatorname{acsch}(dx+c)}{f^3 x^3 + 3e f^2 x^2 + 3e^2 f x + e^3} dx \right) a b e^2 f^2 x + 4 \left(\int \frac{\operatorname{acsch}(dx+c)}{f^3 x^3 + 3e f^2 x^2 + 3e^2 f x + e^3} dx \right) a b e^2 f^2 x^2 + 4 \left(\int \frac{\operatorname{acsch}(dx+c)}{f^3 x^3 + 3e f^2 x^2 + 3e^2 f x + e^3} dx \right) a b e^2 f^2 x^3 + 4 \left(\int \frac{\operatorname{acsch}(dx+c)}{f^3 x^3 + 3e f^2 x^2 + 3e^2 f x + e^3} dx \right) a b e^2 f^2 x^4 + \dots$$

input `int((a+b*acsch(d*x+c))^2/(f*x+e)^3,x)`output `(4*int(acsch(c + d*x)/(e**3 + 3*e**2*f*x + 3*e*f**2*x**2 + f**3*x**3),x)*a*b*e**2*f + 8*int(acsch(c + d*x)/(e**3 + 3*e**2*f*x + 3*e*f**2*x**2 + f**3*x**3),x)*a*b*e*f**2*x + 4*int(acsch(c + d*x)/(e**3 + 3*e**2*f*x + 3*e*f**2*x**2 + f**3*x**3),x)*a*b*f**3*x**2 + 2*int(acsch(c + d*x)**2/(e**3 + 3*e**2*f*x + 3*e*f**2*x**2 + f**3*x**3),x)*b**2*e**2*f + 4*int(acsch(c + d*x)**2/(e**3 + 3*e**2*f*x + 3*e*f**2*x**2 + f**3*x**3),x)*b**2*e*f**2*x + 2*int(acsch(c + d*x)**2/(e**3 + 3*e**2*f*x + 3*e*f**2*x**2 + f**3*x**3),x)*b**2*e*f**2*x**2 + 2*int(acsch(c + d*x)**2/(e**3 + 3*e**2*f*x + 3*e*f**2*x**2 + f**3*x**3),x)*b**2*f**3*x**2 - a**2)/(2*f*(e**2 + 2*e*f*x + f**2*x**2))`

3.14 $\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx$

Optimal result	146
Mathematica [A] (verified)	146
Rubi [A] (verified)	147
Maple [A] (verified)	148
Fricas [A] (verification not implemented)	149
Sympy [F]	149
Maxima [A] (verification not implemented)	150
Giac [F]	150
Mupad [F(-1)]	150
Reduce [F]	151

Optimal result

Integrand size = 10, antiderivative size = 114

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = -\frac{\sqrt{-1-x}\sqrt{x}}{4\sqrt{-x}} - \frac{(-1-x)^{3/2}\sqrt{x}}{4\sqrt{-x}} - \frac{3(-1-x)^{5/2}\sqrt{x}}{20\sqrt{-x}} - \frac{(-1-x)^{7/2}\sqrt{x}}{28\sqrt{-x}} + \frac{1}{4}x^4 \operatorname{csch}^{-1}(\sqrt{x})$$

output

```
-1/4*(-1-x)^(1/2)*x^(1/2)/(-x)^(1/2)-1/4*(-1-x)^(3/2)*x^(1/2)/(-x)^(1/2)-3/20*(-1-x)^(5/2)*x^(1/2)/(-x)^(1/2)-1/28*(-1-x)^(7/2)*x^(1/2)/(-x)^(1/2)+1/4*x^4*arccsch(x^(1/2))
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.41

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{140} \sqrt{1 + \frac{1}{x}} \sqrt{x} (-16 + 8x - 6x^2 + 5x^3) + \frac{1}{4} x^4 \operatorname{csch}^{-1}(\sqrt{x})$$

input

```
Integrate[x^3*ArcCsch[Sqrt[x]], x]
```

output $(\text{Sqrt}[1 + x^{(-1)}] * \text{Sqrt}[x] * (-16 + 8*x - 6*x^2 + 5*x^3))/140 + (x^4 * \text{ArcCsch}[\text{Sqrt}[x]])/4$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.69, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {6900, 27, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \text{csch}^{-1}(\sqrt{x}) dx$$

$$\downarrow 6900$$

$$\frac{1}{4}x^4 \text{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \frac{x^3}{2\sqrt{-x-1}} dx}{4\sqrt{-x}}$$

$$\downarrow 27$$

$$\frac{1}{4}x^4 \text{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \frac{x^3}{\sqrt{-x-1}} dx}{8\sqrt{-x}}$$

$$\downarrow 53$$

$$\frac{1}{4}x^4 \text{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \left(-(-x-1)^{5/2} - 3(-x-1)^{3/2} - 3\sqrt{-x-1} - \frac{1}{\sqrt{-x-1}} \right) dx}{8\sqrt{-x}}$$

$$\downarrow 2009$$

$$\frac{1}{4}x^4 \text{csch}^{-1}(\sqrt{x}) - \frac{\left(\frac{2}{7}(-x-1)^{7/2} + \frac{6}{5}(-x-1)^{5/2} + 2(-x-1)^{3/2} + 2\sqrt{-x-1} \right) \sqrt{x}}{8\sqrt{-x}}$$

input $\text{Int}[x^3 * \text{ArcCsch}[\text{Sqrt}[x]], x]$

output $-1/8 * ((2 * \text{Sqrt}[-1 - x] + 2 * (-1 - x)^{(3/2)} + (6 * (-1 - x)^{(5/2)}))/5 + (2 * (-1 - x)^{(7/2)})/7) * \text{Sqrt}[x] / \text{Sqrt}[-x] + (x^4 * \text{ArcCsch}[\text{Sqrt}[x]])/4$

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6900 `Int[((a_.) + ArcCsch[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcCsch[u])/(d*(m + 1))), x] - Simp[b*(u/(d*(m + 1)*Sqrt[-u^2])) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[-1 - u^2]))], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]`

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.35

method	result	size
parts	$\frac{x^4 \operatorname{arccsch}(\sqrt{x})}{4} + \frac{\sqrt{\frac{x+1}{x}} \sqrt{x} (5x^3 - 6x^2 + 8x - 16)}{140}$	40
derivativedivides	$\frac{x^4 \operatorname{arccsch}(\sqrt{x})}{4} + \frac{(x+1)(5x^3 - 6x^2 + 8x - 16)}{140\sqrt{\frac{x+1}{x}} \sqrt{x}}$	43
default	$\frac{x^4 \operatorname{arccsch}(\sqrt{x})}{4} + \frac{(x+1)(5x^3 - 6x^2 + 8x - 16)}{140\sqrt{\frac{x+1}{x}} \sqrt{x}}$	43

input `int(x^3*arccsch(x^(1/2)),x,method=_RETURNVERBOSE)`

output `1/4*x^4*arccsch(x^(1/2))+1/140*((x+1)/x)^(1/2)*x^(1/2)*(5*x^3-6*x^2+8*x-16)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.48

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{4} x^4 \log \left(\frac{x \sqrt{\frac{x+1}{x}} + \sqrt{x}}{x} \right) + \frac{1}{140} (5x^3 - 6x^2 + 8x - 16) \sqrt{x} \sqrt{\frac{x+1}{x}}$$

input `integrate(x^3*arccsch(x^(1/2)),x, algorithm="fricas")`

output `1/4*x^4*log((x*sqrt((x + 1)/x) + sqrt(x))/x) + 1/140*(5*x^3 - 6*x^2 + 8*x - 16)*sqrt(x)*sqrt((x + 1)/x)`

Sympy [F]

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = \int x^3 \operatorname{acsch}(\sqrt{x}) dx$$

input `integrate(x**3*acsch(x**(1/2)),x)`

output `Integral(x**3*acsch(sqrt(x)), x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.51

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{28} x^{\frac{7}{2}} \left(\frac{1}{x} + 1\right)^{\frac{7}{2}} - \frac{3}{20} x^{\frac{5}{2}} \left(\frac{1}{x} + 1\right)^{\frac{5}{2}} + \frac{1}{4} x^4 \operatorname{arcsch}(\sqrt{x}) + \frac{1}{4} x^{\frac{3}{2}} \left(\frac{1}{x} + 1\right)^{\frac{3}{2}} - \frac{1}{4} \sqrt{x} \sqrt{\frac{1}{x} + 1}$$

input `integrate(x^3*arccsch(x^(1/2)),x, algorithm="maxima")`output `1/28*x^(7/2)*(1/x + 1)^(7/2) - 3/20*x^(5/2)*(1/x + 1)^(5/2) + 1/4*x^4*arccsch(sqrt(x)) + 1/4*x^(3/2)*(1/x + 1)^(3/2) - 1/4*sqrt(x)*sqrt(1/x + 1)`**Giac [F]**

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = \int x^3 \operatorname{arcsch}(\sqrt{x}) dx$$

input `integrate(x^3*arccsch(x^(1/2)),x, algorithm="giac")`output `integrate(x^3*arccsch(sqrt(x)), x)`**Mupad [F(-1)]**

Timed out.

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = \int x^3 \operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right) dx$$

input `int(x^3*asinh(1/x^(1/2)),x)`output `int(x^3*asinh(1/x^(1/2)), x)`

Reduce [F]

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = \int \operatorname{acsch}(\sqrt{x}) x^3 dx$$

input `int(x^3*acsch(x^(1/2)),x)`

output `int(acsch(sqrt(x))*x**3,x)`

3.15 $\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx$

Optimal result	152
Mathematica [A] (verified)	152
Rubi [A] (verified)	153
Maple [A] (verified)	154
Fricas [A] (verification not implemented)	155
Sympy [F]	155
Maxima [A] (verification not implemented)	155
Giac [F]	156
Mupad [F(-1)]	156
Reduce [F]	156

Optimal result

Integrand size = 10, antiderivative size = 89

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{\sqrt{-1-x}\sqrt{x}}{3\sqrt{-x}} + \frac{2(-1-x)^{3/2}\sqrt{x}}{9\sqrt{-x}} + \frac{(-1-x)^{5/2}\sqrt{x}}{15\sqrt{-x}} + \frac{1}{3}x^3 \operatorname{csch}^{-1}(\sqrt{x})$$

output

$$\frac{1}{3}(-1-x)^{(1/2)}x^{(1/2)} / (-x)^{(1/2)} + \frac{2}{9}(-1-x)^{(3/2)}x^{(1/2)} / (-x)^{(1/2)} + \frac{1}{15}(-1-x)^{(5/2)}x^{(1/2)} / (-x)^{(1/2)} + \frac{1}{3}x^3 \operatorname{arccsch}(x^{(1/2)})$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.47

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{45} \sqrt{1 + \frac{1}{x}} \sqrt{x} (8 - 4x + 3x^2) + \frac{1}{3} x^3 \operatorname{csch}^{-1}(\sqrt{x})$$

input

```
Integrate[x^2*ArcCsch[Sqrt[x]],x]
```

output

```
(Sqrt[1 + x^(-1)]*Sqrt[x]*(8 - 4*x + 3*x^2))/45 + (x^3*ArcCsch[Sqrt[x]])/3
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.76, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {6900, 27, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \operatorname{csch}^{-1}(\sqrt{x}) \, dx \\
 & \quad \downarrow \text{6900} \\
 & \frac{1}{3} x^3 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \frac{x^2}{2\sqrt{-x-1}} \, dx}{3\sqrt{-x}} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{3} x^3 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \frac{x^2}{\sqrt{-x-1}} \, dx}{6\sqrt{-x}} \\
 & \quad \downarrow \text{53} \\
 & \frac{1}{3} x^3 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \left((-x-1)^{3/2} + 2\sqrt{-x-1} + \frac{1}{\sqrt{-x-1}} \right) \, dx}{6\sqrt{-x}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{3} x^3 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\left(-\frac{2}{5}(-x-1)^{5/2} - \frac{4}{3}(-x-1)^{3/2} - 2\sqrt{-x-1} \right) \sqrt{x}}{6\sqrt{-x}}
 \end{aligned}$$

input `Int [x^2*ArcCsch[Sqrt [x]] , x]`

output `-1/6*((-2*Sqrt[-1 - x] - (4*(-1 - x)^(3/2)))/3 - (2*(-1 - x)^(5/2))/5)*Sqrt [x])/Sqrt [-x] + (x^3*ArcCsch[Sqrt [x]])/3`

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 53 `Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6900 `Int[((a_.) + ArcCsch[u_]*(b_.))*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcCsch[u])/(d*(m + 1))), x] - Simp[b*(u/(d*(m + 1)*Sqrt[-u^2])) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[-1 - u^2]))], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.39

method	result	size
parts	$\frac{x^3 \operatorname{arccsch}(\sqrt{x})}{3} + \frac{\sqrt{\frac{x+1}{x}} \sqrt{x} (3x^2 - 4x + 8)}{45}$	35
derivativedivides	$\frac{x^3 \operatorname{arccsch}(\sqrt{x})}{3} + \frac{(x+1)(3x^2 - 4x + 8)}{45\sqrt{\frac{x+1}{x}} \sqrt{x}}$	38
default	$\frac{x^3 \operatorname{arccsch}(\sqrt{x})}{3} + \frac{(x+1)(3x^2 - 4x + 8)}{45\sqrt{\frac{x+1}{x}} \sqrt{x}}$	38

input `int(x^2*arccsch(x^(1/2)),x,method=_RETURNVERBOSE)`

output `1/3*x^3*arccsch(x^(1/2))+1/45*((x+1)/x)^(1/2)*x^(1/2)*(3*x^2-4*x+8)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.56

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{3} x^3 \log \left(\frac{x \sqrt{\frac{x+1}{x}} + \sqrt{x}}{x} \right) + \frac{1}{45} (3x^2 - 4x + 8) \sqrt{x} \sqrt{\frac{x+1}{x}}$$

input `integrate(x^2*arccsch(x^(1/2)),x, algorithm="fricas")`output `1/3*x^3*log((x*sqrt((x + 1)/x) + sqrt(x))/x) + 1/45*(3*x^2 - 4*x + 8)*sqrt(x)*sqrt((x + 1)/x)`**Sympy [F]**

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx = \int x^2 \operatorname{acsch}(\sqrt{x}) dx$$

input `integrate(x**2*acsch(x**(1/2)),x)`output `Integral(x**2*acsch(sqrt(x)), x)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.52

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{15} x^{\frac{5}{2}} \left(\frac{1}{x} + 1 \right)^{\frac{5}{2}} + \frac{1}{3} x^3 \operatorname{arcsch}(\sqrt{x}) - \frac{2}{9} x^{\frac{3}{2}} \left(\frac{1}{x} + 1 \right)^{\frac{3}{2}} + \frac{1}{3} \sqrt{x} \sqrt{\frac{1}{x} + 1}$$

input `integrate(x^2*arccsch(x^(1/2)),x, algorithm="maxima")`

output $1/15*x^{(5/2)}*(1/x + 1)^{(5/2)} + 1/3*x^3*arccsch(sqrt(x)) - 2/9*x^{(3/2)}*(1/x + 1)^{(3/2)} + 1/3*sqrt(x)*sqrt(1/x + 1)$

Giac [F]

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) \, dx = \int x^2 \operatorname{arcsch}(\sqrt{x}) \, dx$$

input `integrate(x^2*arccsch(x^(1/2)),x, algorithm="giac")`

output `integrate(x^2*arccsch(sqrt(x)), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) \, dx = \int x^2 \operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right) \, dx$$

input `int(x^2*asinh(1/x^(1/2)),x)`

output `int(x^2*asinh(1/x^(1/2)), x)`

Reduce [F]

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) \, dx = \int \operatorname{acsch}(\sqrt{x}) x^2 \, dx$$

input `int(x^2*acsch(x^(1/2)),x)`

output `int(acsch(sqrt(x))*x**2,x)`

3.16 $\int x \operatorname{csch}^{-1}(\sqrt{x}) dx$

Optimal result	157
Mathematica [A] (verified)	157
Rubi [A] (verified)	158
Maple [A] (verified)	159
Fricas [A] (verification not implemented)	160
Sympy [F]	160
Maxima [A] (verification not implemented)	160
Giac [F]	161
Mupad [F(-1)]	161
Reduce [F]	161

Optimal result

Integrand size = 8, antiderivative size = 64

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx = -\frac{\sqrt{-1-x}\sqrt{x}}{2\sqrt{-x}} - \frac{(-1-x)^{3/2}\sqrt{x}}{6\sqrt{-x}} + \frac{1}{2}x^2 \operatorname{csch}^{-1}(\sqrt{x})$$

output

```
-1/2*(-1-x)^(1/2)*x^(1/2)/(-x)^(1/2)-1/6*(-1-x)^(3/2)*x^(1/2)/(-x)^(1/2)+1/2*x^2*arccsch(x^(1/2))
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.55

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{6}\sqrt{1+\frac{1}{x}}(-2+x)\sqrt{x} + \frac{1}{2}x^2 \operatorname{csch}^{-1}(\sqrt{x})$$

input

```
Integrate[x*ArcCsch[Sqrt[x]],x]
```

output

```
(Sqrt[1 + x^(-1)]*(-2 + x)*Sqrt[x])/6 + (x^2*ArcCsch[Sqrt[x]])/2
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.86, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6900, 27, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \operatorname{csch}^{-1}(\sqrt{x}) \, dx \\
 & \quad \downarrow \text{6900} \\
 & \frac{1}{2}x^2 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \frac{x}{2\sqrt{-x-1}} \, dx}{2\sqrt{-x}} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{2}x^2 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \frac{x}{\sqrt{-x-1}} \, dx}{4\sqrt{-x}} \\
 & \quad \downarrow \text{53} \\
 & \frac{1}{2}x^2 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \left(-\sqrt{-x-1} - \frac{1}{\sqrt{-x-1}} \right) \, dx}{4\sqrt{-x}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2}x^2 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\left(\frac{2}{3}(-x-1)^{3/2} + 2\sqrt{-x-1} \right) \sqrt{x}}{4\sqrt{-x}}
 \end{aligned}$$

input `Int [x*ArcCsch[Sqrt [x]] , x]`

output `-1/4*((2*Sqrt [-1 - x] + (2*(-1 - x)^(3/2))/3)*Sqrt [x])/Sqrt [-x] + (x^2*ArcCsch [Sqrt [x]])/2`

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 53 `Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6900 `Int[((a_.) + ArcCsch[u_]*(b_.))*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcCsch[u])/(d*(m + 1))), x] - Simp[b*(u/(d*(m + 1)*Sqrt[-u^2])) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[-1 - u^2]))], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]`

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.44

method	result	size
parts	$\frac{x^2 \operatorname{arccsch}(\sqrt{x})}{2} + \frac{\sqrt{\frac{x+1}{x}} \sqrt{x} (x-2)}{6}$	28
derivativedivides	$\frac{x^2 \operatorname{arccsch}(\sqrt{x})}{2} + \frac{(x+1)(x-2)}{6\sqrt{\frac{x+1}{x}} \sqrt{x}}$	31
default	$\frac{x^2 \operatorname{arccsch}(\sqrt{x})}{2} + \frac{(x+1)(x-2)}{6\sqrt{\frac{x+1}{x}} \sqrt{x}}$	31

input `int(x*arccsch(x^(1/2)),x,method=_RETURNVERBOSE)`

output `1/2*x^2*arccsch(x^(1/2))+1/6*((x+1)/x)^(1/2)*x^(1/2)*(x-2)`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.67

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) \, dx = \frac{1}{2} x^2 \log \left(\frac{x \sqrt{\frac{x+1}{x}} + \sqrt{x}}{x} \right) + \frac{1}{6} (x-2) \sqrt{x} \sqrt{\frac{x+1}{x}}$$

input `integrate(x*arccsch(x^(1/2)),x, algorithm="fricas")`

output `1/2*x^2*log((x*sqrt((x + 1)/x) + sqrt(x))/x) + 1/6*(x - 2)*sqrt(x)*sqrt((x + 1)/x)`

Sympy [F]

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) \, dx = \int x \operatorname{acsch}(\sqrt{x}) \, dx$$

input `integrate(x*acsch(x**(1/2)),x)`

output `Integral(x*acsch(sqrt(x)), x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.53

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) \, dx = \frac{1}{6} x^{\frac{3}{2}} \left(\frac{1}{x} + 1 \right)^{\frac{3}{2}} + \frac{1}{2} x^2 \operatorname{arcsch}(\sqrt{x}) - \frac{1}{2} \sqrt{x} \sqrt{\frac{1}{x} + 1}$$

input `integrate(x*arccsch(x^(1/2)),x, algorithm="maxima")`

output `1/6*x^(3/2)*(1/x + 1)^(3/2) + 1/2*x^2*arccsch(sqrt(x)) - 1/2*sqrt(x)*sqrt(1/x + 1)`

Giac [F]

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx = \int x \operatorname{arcsch}(\sqrt{x}) dx$$

input `integrate(x*arccsch(x^(1/2)),x, algorithm="giac")`

output `integrate(x*arccsch(sqrt(x)), x)`

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx = \int x \operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right) dx$$

input `int(x*asinh(1/x^(1/2)),x)`

output `int(x*asinh(1/x^(1/2)), x)`

Reduce [F]

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx = \int \operatorname{acsch}(\sqrt{x}) x dx$$

input `int(x*acsch(x^(1/2)),x)`

output `int(acsch(sqrt(x))*x,x)`

3.17 $\int \operatorname{csch}^{-1}(\sqrt{x}) dx$

Optimal result	162
Mathematica [A] (verified)	162
Rubi [A] (verified)	163
Maple [A] (verified)	164
Fricas [A] (verification not implemented)	164
Sympy [F]	165
Maxima [A] (verification not implemented)	165
Giac [F]	165
Mupad [B] (verification not implemented)	166
Reduce [F]	166

Optimal result

Integrand size = 6, antiderivative size = 31

$$\int \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{\sqrt{-1-x}\sqrt{x}}{\sqrt{-x}} + x\operatorname{csch}^{-1}(\sqrt{x})$$

output

```
(-1-x)^(1/2)*x^(1/2)/(-x)^(1/2)+x*arccsch(x^(1/2))
```

Mathematica [A] (verified)

Time = 2.17 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.77

$$\int \operatorname{csch}^{-1}(\sqrt{x}) dx = \sqrt{1 + \frac{1}{x}}\sqrt{x} + x\operatorname{csch}^{-1}(\sqrt{x})$$

input

```
Integrate[ArcCsch[Sqrt[x]],x]
```

output

```
Sqrt[1 + x^(-1)]*Sqrt[x] + x*ArcCsch[Sqrt[x]]
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6898, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{csch}^{-1}(\sqrt{x}) \, dx$$

$$\downarrow \text{6898}$$

$$x \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \frac{1}{2\sqrt{-x-1}} \, dx}{\sqrt{-x}}$$

$$\downarrow \text{17}$$

$$\frac{\sqrt{-x-1}\sqrt{x}}{\sqrt{-x}} + x \operatorname{csch}^{-1}(\sqrt{x})$$

input `Int[ArcCsch[Sqrt[x]], x]`

output `(Sqrt[-1 - x]*Sqrt[x])/Sqrt[-x] + x*ArcCsch[Sqrt[x]]`

Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 6898 `Int[ArcCsch[u_], x_Symbol] := Simp[x*ArcCsch[u], x] - Simp[u/Sqrt[-u^2] Int[SimplifyIntegrand[x*(D[u, x]/(u*Sqrt[-1 - u^2])), x], x], x] /; InverseFunctionFreeQ[u, x] && !FunctionOfExponentialQ[u, x]`

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.68

method	result	size
parts	$x \operatorname{arccsch}(\sqrt{x}) + \sqrt{x} \sqrt{\frac{x+1}{x}}$	21
derivativedivides	$x \operatorname{arccsch}(\sqrt{x}) + \frac{x+1}{\sqrt{\frac{x+1}{x}} \sqrt{x}}$	24
default	$x \operatorname{arccsch}(\sqrt{x}) + \frac{x+1}{\sqrt{\frac{x+1}{x}} \sqrt{x}}$	24

input `int(arccsch(x^(1/2)),x,method=_RETURNVERBOSE)`output `x*arccsch(x^(1/2))+x^(1/2)*((x+1)/x)^(1/2)`**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.16

$$\int \operatorname{csch}^{-1}(\sqrt{x}) dx = x \log \left(\frac{x \sqrt{\frac{x+1}{x}} + \sqrt{x}}{x} \right) + \sqrt{x} \sqrt{\frac{x+1}{x}}$$

input `integrate(arccsch(x^(1/2)),x, algorithm="fricas")`output `x*log((x*sqrt((x + 1)/x) + sqrt(x))/x) + sqrt(x)*sqrt((x + 1)/x)`

Sympy [F]

$$\int \operatorname{csch}^{-1}(\sqrt{x}) \, dx = \int \operatorname{acsch}(\sqrt{x}) \, dx$$

input `integrate(acsch(x**(1/2)),x)`

output `Integral(acsch(sqrt(x)), x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.58

$$\int \operatorname{csch}^{-1}(\sqrt{x}) \, dx = x \operatorname{arcsch}(\sqrt{x}) + \sqrt{x} \sqrt{\frac{1}{x} + 1}$$

input `integrate(arccsch(x^(1/2)),x, algorithm="maxima")`

output `x*arccsch(sqrt(x)) + sqrt(x)*sqrt(1/x + 1)`

Giac [F]

$$\int \operatorname{csch}^{-1}(\sqrt{x}) \, dx = \int \operatorname{arcsch}(\sqrt{x}) \, dx$$

input `integrate(arccsch(x^(1/2)),x, algorithm="giac")`

output `integrate(arccsch(sqrt(x)), x)`

Mupad [B] (verification not implemented)

Time = 3.82 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.58

$$\int \operatorname{csch}^{-1}(\sqrt{x}) dx = x \operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right) + \sqrt{x} \sqrt{\frac{1}{x} + 1}$$

input `int(asinh(1/x^(1/2)),x)`

output `x*asinh(1/x^(1/2)) + x^(1/2)*(1/x + 1)^(1/2)`

Reduce [F]

$$\int \operatorname{csch}^{-1}(\sqrt{x}) dx = \int \operatorname{acsch}(\sqrt{x}) dx$$

input `int(acsch(x^(1/2)),x)`

output `int(acsch(sqrt(x)),x)`

3.18 $\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx$

Optimal result	167
Mathematica [A] (verified)	167
Rubi [C] (warning: unable to verify)	168
Maple [F]	171
Fricas [F]	171
Sympy [F]	171
Maxima [F]	172
Giac [F]	172
Mupad [F(-1)]	172
Reduce [F]	173

Optimal result

Integrand size = 10, antiderivative size = 46

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = \operatorname{csch}^{-1}(\sqrt{x})^2 - 2\operatorname{csch}^{-1}(\sqrt{x}) \log\left(1 - e^{2\operatorname{csch}^{-1}(\sqrt{x})}\right) - \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(\sqrt{x})}\right)$$

output

```
arccsch(x^(1/2))^2-2*arccsch(x^(1/2))*ln(1-(1/x^(1/2)+(1+1/x)^(1/2))^2)-polylog(2, (1/x^(1/2)+(1+1/x)^(1/2))^2)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = \operatorname{csch}^{-1}(\sqrt{x}) \left(\operatorname{csch}^{-1}(\sqrt{x}) - 2 \log\left(1 - e^{2\operatorname{csch}^{-1}(\sqrt{x})}\right) \right) - \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(\sqrt{x})}\right)$$

input

```
Integrate[ArcCsch[Sqrt[x]]/x,x]
```


output

```
ArcCsch[Sqrt[x]]*(ArcCsch[Sqrt[x]] - 2*Log[1 - E^(2*ArcCsch[Sqrt[x]])]) -
PolyLog[2, E^(2*ArcCsch[Sqrt[x]])]
```

Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.46 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.26, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {7267, 6836, 6190, 3042, 26, 4199, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx \\
 & \quad \downarrow \text{7267} \\
 & 2 \int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{\sqrt{x}} d\sqrt{x} \\
 & \quad \downarrow \text{6836} \\
 & -2 \int \frac{\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)}{\sqrt{x}} d\frac{1}{\sqrt{x}} \\
 & \quad \downarrow \text{6190} \\
 & -2 \int \sqrt{1 + \frac{1}{x}} \sqrt{x} \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) d\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) \\
 & \quad \downarrow \text{3042} \\
 & -2 \int -i \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) \tan\left(i \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) + \frac{\pi}{2}\right) d\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) \\
 & \quad \downarrow \text{26} \\
 & 2i \int \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) \tan\left(i \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) + \frac{\pi}{2}\right) d\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) \\
 & \quad \downarrow \text{4199}
 \end{aligned}$$

$$2i \left(2i \int -\frac{e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)} \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)}{1 - e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)}} d\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) - \frac{ix}{2} \right)$$

↓ 25

$$2i \left(-2i \int \frac{e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)} \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)}{1 - e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)}} d\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) - \frac{ix}{2} \right)$$

↓ 2620

$$2i \left(-2i \left(\frac{1}{2} \int \log \left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)} \right) d\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) - \frac{1}{2} \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) \log \left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)} \right) \right) - \frac{ix}{2} \right)$$

↓ 2715

$$2i \left(-2i \left(\frac{1}{4} \int e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)} \log \left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)} \right) de^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)} - \frac{1}{2} \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) \log \left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)} \right) \right) - \frac{ix}{2} \right)$$

↓ 2838

$$2i \left(-2i \left(-\frac{1}{4} \operatorname{PolyLog} \left(2, e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)} \right) - \frac{1}{2} \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) \log \left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)} \right) \right) - \frac{ix}{2} \right)$$

input `Int[ArcCsch[Sqrt[x]]/x,x]`

output `(2*I)*((-1/2*I)*x - (2*I)*(-1/2*(ArcSinh[1/Sqrt[x]]*Log[1 - E^(2*ArcSinh[1/Sqrt[x]])]) - PolyLog[2, E^(2*ArcSinh[1/Sqrt[x]])]/4))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 $\text{Int}[(((F_)^{(g_)*((e_)+(f_)*(x_)))^{(n_)*((c_)+(d_)*(x_))^{(m_)}})/((a_)+(b_)*((F_)^{(g_)*((e_)+(f_)*(x_)))^{(n_)}), x_Symbol] \rightarrow \text{Simp} [((c + d*x)^m/(b*f*g*n*\text{Log}[F]))*\text{Log}[1 + b*((F^{(g*(e + f*x)))^n/a}], x] - \text{Simp}[d*(m/(b*f*g*n*\text{Log}[F])) \text{Int}[(c + d*x)^{m-1}*\text{Log}[1 + b*((F^{(g*(e + f*x)))^n/a}], x], x] /; \text{FreeQ}\{F, a, b, c, d, e, f, g, n\}, x\} \&\& \text{IGtQ}[m, 0]$

rule 2715 $\text{Int}[\text{Log}[a_ + (b_)*((F_)^{(e_)*((c_)+(d_)*(x_)))^{(n_)}], x_Symbol] \rightarrow \text{Simp}[1/(d*e*n*\text{Log}[F]) \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^{(e*(c + d*x)))^n}], x] /; \text{FreeQ}\{F, a, b, c, d, e, n\}, x\} \&\& \text{GtQ}[a, 0]$

rule 2838 $\text{Int}[\text{Log}[(c_)*((d_)+(e_)*(x_)^{(n_)})]/(x_), x_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n]/n, x] /; \text{FreeQ}\{c, d, e, n\}, x\} \&\& \text{EqQ}[c*d, 1]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4199 $\text{Int}[((c_)+(d_)*(x_))^{(m_)*\text{tan}[(e_)+\text{Pi}*(k_)+(Complex[0, fz_])*(f_)*(x_)], x_Symbol] \rightarrow \text{Simp}[(-I)*((c + d*x)^{(m+1)}/(d*(m+1))), x] + \text{Simp}[2*I \text{Int}[(c + d*x)^m*(E^{(2*((-I)*e + f*fz*x))}/(1 + E^{(2*((-I)*e + f*fz*x))})/E^{(2*I*k*Pi)}))/E^{(2*I*k*Pi)}, x], x] /; \text{FreeQ}\{c, d, e, f, fz\}, x\} \&\& \text{IntegerQ}[4*k] \&\& \text{IGtQ}[m, 0]$

rule 6190 $\text{Int}[((a_)+\text{ArcSinh}[(c_)*(x_)]*(b_))^{(n_)} / (x_), x_Symbol] \rightarrow \text{Simp}[1/b \text{Subst}[\text{Int}[x^n*\text{Coth}[-a/b + x/b], x], x, a + b*\text{ArcSinh}[c*x], x] /; \text{FreeQ}\{a, b, c\}, x\} \&\& \text{IGtQ}[n, 0]$

rule 6836 $\text{Int}[((a_)+\text{ArcSch}[(c_)*(x_)]*(b_)) / (x_), x_Symbol] \rightarrow -\text{Subst}[\text{Int}[(a + b*\text{ArcSinh}[x/c])/x, x], x, 1/x] /; \text{FreeQ}\{a, b, c\}, x\}$

rule 7267 $\text{Int}[u_, x_Symbol] \rightarrow \text{With}\{[1st = \text{SubstForFractionalPowerOfLinear}[u, x]\}, \text{Simp}[1st[[2]]*1st[[4]] \text{Subst}[\text{Int}[1st[[1]], x], x, 1st[[3]]^{(1/1st[[2]])}], x] /; !\text{FalseQ}[1st] \&\& \text{SubstForFractionalPowerQ}[u, 1st[[3]], x]$

Maple [F]

$$\int \frac{\operatorname{arccsch}(\sqrt{x})}{x} dx$$

input `int(arccsch(x^(1/2))/x,x)`

output `int(arccsch(x^(1/2))/x,x)`

Fricas [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arcsch}(\sqrt{x})}{x} dx$$

input `integrate(arccsch(x^(1/2))/x,x, algorithm="fricas")`

output `integral(arccsch(sqrt(x))/x, x)`

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{acsch}(\sqrt{x})}{x} dx$$

input `integrate(acsch(x**(1/2))/x,x)`

output `Integral(acsch(sqrt(x))/x, x)`

Maxima [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arcsch}(\sqrt{x})}{x} dx$$

input `integrate(arccsch(x^(1/2))/x,x, algorithm="maxima")`

output `integrate(arccsch(sqrt(x))/x, x)`

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arcsch}(\sqrt{x})}{x} dx$$

input `integrate(arccsch(x^(1/2))/x,x, algorithm="giac")`

output `integrate(arccsch(sqrt(x))/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right)}{x} dx$$

input `int(asinh(1/x^(1/2))/x,x)`

output `int(asinh(1/x^(1/2))/x, x)`

Reduce [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{acsch}(\sqrt{x})}{x} dx$$

input `int(acsch(x^(1/2))/x,x)`

output `int(acsch(sqrt(x))/x,x)`

3.19 $\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx$

Optimal result	174
Mathematica [A] (verified)	174
Rubi [A] (verified)	175
Maple [A] (verified)	177
Fricas [A] (verification not implemented)	177
Sympy [F]	178
Maxima [A] (verification not implemented)	178
Giac [F]	178
Mupad [B] (verification not implemented)	179
Reduce [F]	179

Optimal result

Integrand size = 10, antiderivative size = 63

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{-1-x}}{2\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} - \frac{\sqrt{x} \arctan(\sqrt{-1-x})}{2\sqrt{-x}}$$

output

$1/2*(-1-x)^{(1/2)/(-x)^{(1/2)/x^{(1/2)}-\operatorname{arccsch}(x^{(1/2)})/x-1/2*x^{(1/2)}*\arctan((-1-x)^{(1/2))/(-x)^{(1/2)}}$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.67

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{\frac{1+x}{x}}}{2\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} - \frac{1}{2} \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)$$

input

`Integrate[ArcCsch[Sqrt[x]]/x^2,x]`

output

`Sqrt[(1 + x)/x]/(2*Sqrt[x]) - ArcCsch[Sqrt[x]]/x - ArcSinh[1/Sqrt[x]]/2`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.86, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6900, 27, 52, 73, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx \\
 & \quad \downarrow \text{6900} \\
 & \frac{\sqrt{x} \int \frac{1}{2\sqrt{-x-1}x^2} dx}{\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} \\
 & \quad \downarrow \text{27} \\
 & \frac{\sqrt{x} \int \frac{1}{\sqrt{-x-1}x^2} dx}{2\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} \\
 & \quad \downarrow \text{52} \\
 & \frac{\sqrt{x} \left(\frac{\sqrt{-x-1}}{x} - \frac{1}{2} \int \frac{1}{\sqrt{-x-1}x} dx \right)}{2\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} \\
 & \quad \downarrow \text{73} \\
 & \frac{\sqrt{x} \left(\int \frac{1}{x} d\sqrt{-x-1} + \frac{\sqrt{-x-1}}{x} \right)}{2\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} \\
 & \quad \downarrow \text{217} \\
 & \frac{\sqrt{x} \left(\frac{\sqrt{-x-1}}{x} - \arctan(\sqrt{-x-1}) \right)}{2\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x}
 \end{aligned}$$

input `Int[ArcCsch[Sqrt[x]]/x^2,x]`

output `-(ArcCsch[Sqrt[x]]/x) + (Sqrt[x]*(Sqrt[-1 - x]/x - ArcTan[Sqrt[-1 - x]]))/(2*Sqrt[-x])`

Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 6900 `Int[((a_.) + ArcCsch[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcCsch[u])/(d*(m + 1))), x] - Simp[b*(u/(d*(m + 1)*Sqrt[-u^2])) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[-1 - u^2]))], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]`

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.73

method	result	size
derivativedivides	$-\frac{\operatorname{arccsch}(\sqrt{x})}{x} - \frac{\sqrt{x+1} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{x+1}}\right)x - \sqrt{x+1} \right)}{2\sqrt{\frac{x+1}{x}} x^{\frac{3}{2}}}$	46
default	$-\frac{\operatorname{arccsch}(\sqrt{x})}{x} - \frac{\sqrt{x+1} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{x+1}}\right)x - \sqrt{x+1} \right)}{2\sqrt{\frac{x+1}{x}} x^{\frac{3}{2}}}$	46
parts	$-\frac{\operatorname{arccsch}(\sqrt{x})}{x} - \frac{\sqrt{\frac{x+1}{x}} \sqrt{x} (\ln(\sqrt{x+1}+1)x - \ln(\sqrt{x+1}-1)x - 2\sqrt{x+1})}{4\sqrt{x+1} (\sqrt{x+1}+1) (\sqrt{x+1}-1)}$	77

input `int(arccsch(x^(1/2))/x^2,x,method=_RETURNVERBOSE)`

output $-\operatorname{arccsch}(x^{1/2})/x - 1/2*(x+1)^{1/2}*(\operatorname{arctanh}(1/(x+1)^{1/2}))*x - (x+1)^{1/2}) / ((x+1)/x)^{1/2}/x^{3/2}$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.70

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = -\frac{(x+2) \log\left(\frac{x\sqrt{\frac{x+1}{x}} + \sqrt{x}}{x}\right) - \sqrt{x}\sqrt{\frac{x+1}{x}}}{2x}$$

input `integrate(arccsch(x^(1/2))/x^2,x, algorithm="fricas")`

output $-1/2*((x+2)*\log((x*\sqrt{(x+1)/x} + \sqrt{x})/x) - \sqrt{x}*\sqrt{(x+1)/x})/x$

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = \int \frac{\operatorname{acsch}(\sqrt{x})}{x^2} dx$$

input `integrate(acsch(x**(1/2))/x**2,x)`

output `Integral(acsch(sqrt(x))/x**2, x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.03

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{x}\sqrt{\frac{1}{x}+1}}{2\left(x\left(\frac{1}{x}+1\right)-1\right)} - \frac{\operatorname{arcsch}(\sqrt{x})}{x} - \frac{1}{4} \log\left(\sqrt{x}\sqrt{\frac{1}{x}+1}+1\right) + \frac{1}{4} \log\left(\sqrt{x}\sqrt{\frac{1}{x}+1}-1\right)$$

input `integrate(arccsch(x^(1/2))/x^2,x, algorithm="maxima")`

output `1/2*sqrt(x)*sqrt(1/x + 1)/(x*(1/x + 1) - 1) - arccsch(sqrt(x))/x - 1/4*log(sqrt(x)*sqrt(1/x + 1) + 1) + 1/4*log(sqrt(x)*sqrt(1/x + 1) - 1)`

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = \int \frac{\operatorname{arcsch}(\sqrt{x})}{x^2} dx$$

input `integrate(arccsch(x^(1/2))/x^2,x, algorithm="giac")`

output `integrate(arccsch(sqrt(x))/x^2, x)`

Mupad [B] (verification not implemented)

Time = 3.63 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.52

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{\frac{1}{x} + 1}}{2\sqrt{x}} - \frac{2 \operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right) \left(\frac{1}{2\sqrt{x}} + \frac{\sqrt{x}}{4}\right)}{\sqrt{x}}$$

input `int(asinh(1/x^(1/2))/x^2,x)`

output `(1/x + 1)^(1/2)/(2*x^(1/2)) - (2*asinh(1/x^(1/2))*(1/(2*x^(1/2)) + x^(1/2)/4))/x^(1/2)`

Reduce [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = \int \frac{\operatorname{acsch}(\sqrt{x})}{x^2} dx$$

input `int(acsch(x^(1/2))/x^2,x)`

output `int(acsch(sqrt(x))/x**2,x)`

3.20 $\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx$

Optimal result	180
Mathematica [A] (verified)	180
Rubi [A] (verified)	181
Maple [A] (verified)	183
Fricas [A] (verification not implemented)	183
Sympy [F]	184
Maxima [A] (verification not implemented)	184
Giac [F]	184
Mupad [F(-1)]	185
Reduce [F]	185

Optimal result

Integrand size = 10, antiderivative size = 90

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = \frac{\sqrt{-1-x}}{8\sqrt{-x}x^{3/2}} - \frac{3\sqrt{-1-x}}{16\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{2x^2} + \frac{3\sqrt{x} \arctan(\sqrt{-1-x})}{16\sqrt{-x}}$$

output $1/8*(-1-x)^{(1/2)/(-x)^{(1/2)/x^{(3/2)}}-3/16*(-1-x)^{(1/2)/(-x)^{(1/2)/x^{(1/2)}}-1/2*\operatorname{arccsch}(x^{(1/2)})/x^2+3/16*x^{(1/2)}*\arctan((-1-x)^{(1/2)/(-x)^{(1/2)}}$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.52

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = \frac{\sqrt{1+\frac{1}{x}}(2-3x)\sqrt{x} - 8\operatorname{csch}^{-1}(\sqrt{x}) + 3x^2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)}{16x^2}$$

input `Integrate[ArcCsch[Sqrt[x]]/x^3,x]`

output $(\operatorname{Sqrt}[1+x^{(-1)}]*(2-3*x)*\operatorname{Sqrt}[x]-8*\operatorname{ArcCsch}[\operatorname{Sqrt}[x]]+3*x^2*\operatorname{ArcSinh}[1/\operatorname{Sqrt}[x]])/(16*x^2)$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.86, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6900, 27, 52, 52, 73, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx \\
 & \quad \downarrow \text{6900} \\
 & \frac{\sqrt{x} \int \frac{1}{2\sqrt{-x-1}x^3} dx}{2\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{2x^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{\sqrt{x} \int \frac{1}{\sqrt{-x-1}x^3} dx}{4\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{2x^2} \\
 & \quad \downarrow \text{52} \\
 & \frac{\sqrt{x} \left(\frac{\sqrt{-x-1}}{2x^2} - \frac{3}{4} \int \frac{1}{\sqrt{-x-1}x^2} dx \right)}{4\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{2x^2} \\
 & \quad \downarrow \text{52} \\
 & \frac{\sqrt{x} \left(\frac{\sqrt{-x-1}}{2x^2} - \frac{3}{4} \left(\frac{\sqrt{-x-1}}{x} - \frac{1}{2} \int \frac{1}{\sqrt{-x-1}x} dx \right) \right)}{4\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{2x^2} \\
 & \quad \downarrow \text{73} \\
 & \frac{\sqrt{x} \left(\frac{\sqrt{-x-1}}{2x^2} - \frac{3}{4} \left(\int \frac{1}{x} d\sqrt{-x-1} + \frac{\sqrt{-x-1}}{x} \right) \right)}{4\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{2x^2} \\
 & \quad \downarrow \text{217} \\
 & \frac{\sqrt{x} \left(\frac{\sqrt{-x-1}}{2x^2} - \frac{3}{4} \left(\frac{\sqrt{-x-1}}{x} - \arctan(\sqrt{-x-1}) \right) \right)}{4\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{2x^2}
 \end{aligned}$$

input `Int[ArcCsch[Sqrt[x]]/x^3,x]`

output

$$-1/2*\text{ArcSch}[\text{Sqrt}[x]]/x^2 + (\text{Sqrt}[x]*(\text{Sqrt}[-1 - x]/(2*x^2) - (3*(\text{Sqrt}[-1 - x]/x - \text{ArcTan}[\text{Sqrt}[-1 - x]]))/4))/(4*\text{Sqrt}[-x])$$

Defintions of rubi rules used

rule 27

$$\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] \text{ /; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] \text{ /; FreeQ}[b, x]$$

rule 52

$$\text{Int}[(a_.) + (b_.)*(x_)^(m_)*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - \text{Simp}[d*((m + n + 2)/((b*c - a*d)*(m + 1)) \text{ Int}[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] \text{ /; FreeQ}[\{a, b, c, d, n\}, x] \ \&\& \ \text{ILtQ}[m, -1] \ \&\& \ \text{FractionQ}[n] \ \&\& \ \text{LtQ}[n, 0]$$

rule 73

$$\text{Int}[(a_.) + (b_.)*(x_)^(m_)*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \text{ Subst}[\text{Int}[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] \text{ /; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$$

rule 217

$$\text{Int}[(a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^(-1))*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] \text{ /; FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 6900

$$\text{Int}[(a_.) + \text{ArcSch}[u_]*(b_.)*((c_.) + (d_.)*(x_)^(m_)), x_Symbol] \rightarrow \text{Simp}[(c + d*x)^(m + 1)*((a + b*\text{ArcSch}[u])/(d*(m + 1))), x] - \text{Simp}[b*(u/(d*(m + 1)*\text{Sqrt}[-u^2])) \text{ Int}[\text{SimplifyIntegrand}[(c + d*x)^(m + 1)*(D[u, x]/(u*\text{Sqrt}[-1 - u^2])), x], x], x] \text{ /; FreeQ}[\{a, b, c, d, m\}, x] \ \&\& \ \text{NeQ}[m, -1] \ \&\& \ \text{InverseFunctionFreeQ}[u, x] \ \&\& \ !\text{FunctionOfQ}[(c + d*x)^(m + 1), u, x] \ \&\& \ !\text{FunctionOfExponentialQ}[u, x]$$

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.63

method	result	size
derivativedivides	$-\frac{\operatorname{arccsch}(\sqrt{x})}{2x^2} + \frac{\sqrt{x+1} \left(3 \operatorname{arctanh}\left(\frac{1}{\sqrt{x+1}}\right)x^2 - 3\sqrt{x+1}x + 2\sqrt{x+1} \right)}{16\sqrt{\frac{x+1}{x}}x^{\frac{5}{2}}}$	57
default	$-\frac{\operatorname{arccsch}(\sqrt{x})}{2x^2} + \frac{\sqrt{x+1} \left(3 \operatorname{arctanh}\left(\frac{1}{\sqrt{x+1}}\right)x^2 - 3\sqrt{x+1}x + 2\sqrt{x+1} \right)}{16\sqrt{\frac{x+1}{x}}x^{\frac{5}{2}}}$	57
parts	$-\frac{\operatorname{arccsch}(\sqrt{x})}{2x^2} - \frac{\sqrt{\frac{x+1}{x}}\sqrt{x} \left(3 \ln(\sqrt{x+1}-1)x^2 - 3 \ln(\sqrt{x+1}+1)x^2 + 6\sqrt{x+1}x - 4\sqrt{x+1} \right)}{32\sqrt{x+1}(\sqrt{x+1}-1)^2(\sqrt{x+1}+1)^2}$	90

input `int(arccsch(x^(1/2))/x^3,x,method=_RETURNVERBOSE)`

output `-1/2*arccsch(x^(1/2))/x^2+1/16*(x+1)^(1/2)*(3*arctanh(1/(x+1)^(1/2))*x^2-3*(x+1)^(1/2)*x+2*(x+1)^(1/2))/((x+1)/x)^(1/2)/x^(5/2)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.59

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = -\frac{(3x-2)\sqrt{x}\sqrt{\frac{x+1}{x}} - (3x^2-8)\log\left(\frac{x\sqrt{\frac{x+1}{x}}+\sqrt{x}}{x}\right)}{16x^2}$$

input `integrate(arccsch(x^(1/2))/x^3,x, algorithm="fricas")`

output `-1/16*((3*x - 2)*sqrt(x)*sqrt((x + 1)/x) - (3*x^2 - 8)*log((x*sqrt((x + 1)/x) + sqrt(x))/x))/x^2`

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\operatorname{acsch}(\sqrt{x})}{x^3} dx$$

input `integrate(acsch(x**(1/2))/x**3,x)`

output `Integral(acsch(sqrt(x))/x**3, x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.02

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = -\frac{3x^{\frac{3}{2}}\left(\frac{1}{x}+1\right)^{\frac{3}{2}} - 5\sqrt{x}\sqrt{\frac{1}{x}+1}}{16\left(x^2\left(\frac{1}{x}+1\right)^2 - 2x\left(\frac{1}{x}+1\right) + 1\right)} - \frac{\operatorname{arcsch}(\sqrt{x})}{2x^2} \\ + \frac{3}{32} \log\left(\sqrt{x}\sqrt{\frac{1}{x}+1} + 1\right) - \frac{3}{32} \log\left(\sqrt{x}\sqrt{\frac{1}{x}+1} - 1\right)$$

input `integrate(arccsch(x^(1/2))/x^3,x, algorithm="maxima")`

output `-1/16*(3*x^(3/2)*(1/x + 1)^(3/2) - 5*sqrt(x)*sqrt(1/x + 1))/(x^2*(1/x + 1)
^2 - 2*x*(1/x + 1) + 1) - 1/2*arccsch(sqrt(x))/x^2 + 3/32*log(sqrt(x)*sqrt
(1/x + 1) + 1) - 3/32*log(sqrt(x)*sqrt(1/x + 1) - 1)`

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\operatorname{arcsch}(\sqrt{x})}{x^3} dx$$

input `integrate(arccsch(x^(1/2))/x^3,x, algorithm="giac")`

output `integrate(arccsch(sqrt(x))/x^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right)}{x^3} dx$$

input `int(asinh(1/x^(1/2))/x^3,x)`

output `int(asinh(1/x^(1/2))/x^3, x)`

Reduce [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\operatorname{acsch}(\sqrt{x})}{x^3} dx$$

input `int(acsch(x^(1/2))/x^3,x)`

output `int(acsch(sqrt(x))/x**3,x)`

3.21 $\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx$

Optimal result	186
Mathematica [A] (verified)	186
Rubi [A] (verified)	187
Maple [A] (verified)	189
Fricas [A] (verification not implemented)	189
Sympy [F]	190
Maxima [A] (verification not implemented)	190
Giac [F]	191
Mupad [F(-1)]	191
Reduce [F]	191

Optimal result

Integrand size = 10, antiderivative size = 115

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = \frac{\sqrt{-1-x}}{18\sqrt{-x}x^{5/2}} - \frac{5\sqrt{-1-x}}{72\sqrt{-x}x^{3/2}} + \frac{5\sqrt{-1-x}}{48\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3} - \frac{5\sqrt{x} \arctan(\sqrt{-1-x})}{48\sqrt{-x}}$$

output

```
1/18*(-1-x)^(1/2)/(-x)^(1/2)/x^(5/2)-5/72*(-1-x)^(1/2)/(-x)^(1/2)/x^(3/2)+
5/48*(-1-x)^(1/2)/(-x)^(1/2)/x^(1/2)-1/3*arccsch(x^(1/2))/x^3-5/48*x^(1/2)
*arctan((-1-x)^(1/2))/(-x)^(1/2)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.45

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = \frac{\sqrt{1 + \frac{1}{x}}\sqrt{x}(8 - 10x + 15x^2) - 48\operatorname{csch}^{-1}(\sqrt{x}) - 15x^3\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)}{144x^3}$$

input

```
Integrate[ArcCsch[Sqrt[x]]/x^4,x]
```

output

```
(Sqrt[1 + x^(-1)]*Sqrt[x]*(8 - 10*x + 15*x^2) - 48*ArcCsch[Sqrt[x]] - 15*x^3*ArcSinh[1/Sqrt[x]])/(144*x^3)
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.85, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {6900, 27, 52, 52, 52, 73, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx \\
 & \quad \downarrow 6900 \\
 & \frac{\sqrt{x} \int \frac{1}{2\sqrt{-x-1}x^4} dx}{3\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3} \\
 & \quad \downarrow 27 \\
 & \frac{\sqrt{x} \int \frac{1}{\sqrt{-x-1}x^4} dx}{6\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3} \\
 & \quad \downarrow 52 \\
 & \frac{\sqrt{x} \left(\frac{\sqrt{-x-1}}{3x^3} - \frac{5}{6} \int \frac{1}{\sqrt{-x-1}x^3} dx \right)}{6\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3} \\
 & \quad \downarrow 52 \\
 & \frac{\sqrt{x} \left(\frac{\sqrt{-x-1}}{3x^3} - \frac{5}{6} \left(\frac{\sqrt{-x-1}}{2x^2} - \frac{3}{4} \int \frac{1}{\sqrt{-x-1}x^2} dx \right) \right)}{6\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3} \\
 & \quad \downarrow 52 \\
 & \frac{\sqrt{x} \left(\frac{\sqrt{-x-1}}{3x^3} - \frac{5}{6} \left(\frac{\sqrt{-x-1}}{2x^2} - \frac{3}{4} \left(\frac{\sqrt{-x-1}}{x} - \frac{1}{2} \int \frac{1}{\sqrt{-x-1}x} dx \right) \right) \right)}{6\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3} \\
 & \quad \downarrow 73
 \end{aligned}$$

$$\frac{\sqrt{x} \left(\frac{\sqrt{-x-1}}{3x^3} - \frac{5}{6} \left(\frac{\sqrt{-x-1}}{2x^2} - \frac{3}{4} \left(\int \frac{1}{x} d\sqrt{-x-1} + \frac{\sqrt{-x-1}}{x} \right) \right) \right)}{6\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3}$$

↓ 217

$$\frac{\sqrt{x} \left(\frac{\sqrt{-x-1}}{3x^3} - \frac{5}{6} \left(\frac{\sqrt{-x-1}}{2x^2} - \frac{3}{4} \left(\frac{\sqrt{-x-1}}{x} - \arctan(\sqrt{-x-1}) \right) \right) \right)}{6\sqrt{-x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3}$$

input `Int[ArcCsch[Sqrt[x]]/x^4,x]`

output `-1/3*ArcCsch[Sqrt[x]]/x^3 + (Sqrt[x]*(Sqrt[-1 - x]/(3*x^3) - (5*(Sqrt[-1 - x]/(2*x^2) - (3*(Sqrt[-1 - x]/x - ArcTan[Sqrt[-1 - x]]))/4))/6)/(6*Sqrt[-x])`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 52 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 6900

```
Int[((a_.) + ArcCsch[u_]*(b_.))*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] :> Simp[(c + d*x)^(m + 1)*((a + b*ArcCsch[u])/(d*(m + 1))), x] - Simp[b*(u/(d*(m + 1)*Sqrt[-u^2])) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[-1 - u^2]))], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]
```

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.58

method	result	size
derivativedivides	$-\frac{\operatorname{arccsch}(\sqrt{x})}{3x^3} - \frac{\sqrt{x+1} \left(15 \operatorname{arctanh}\left(\frac{1}{\sqrt{x+1}}\right)x^3 - 15\sqrt{x+1}x^2 + 10\sqrt{x+1}x - 8\sqrt{x+1} \right)}{144\sqrt{\frac{x+1}{x}}x^{\frac{7}{2}}}$	67
default	$-\frac{\operatorname{arccsch}(\sqrt{x})}{3x^3} - \frac{\sqrt{x+1} \left(15 \operatorname{arctanh}\left(\frac{1}{\sqrt{x+1}}\right)x^3 - 15\sqrt{x+1}x^2 + 10\sqrt{x+1}x - 8\sqrt{x+1} \right)}{144\sqrt{\frac{x+1}{x}}x^{\frac{7}{2}}}$	67
parts	$-\frac{\operatorname{arccsch}(\sqrt{x})}{3x^3} + \frac{\sqrt{\frac{x+1}{x}}\sqrt{x} \left(15 \ln(\sqrt{x+1}-1)x^3 - 15 \ln(\sqrt{x+1}+1)x^3 + 30\sqrt{x+1}x^2 - 20\sqrt{x+1}x + 16\sqrt{x+1} \right)}{288\sqrt{x+1}(\sqrt{x+1}-1)^3(\sqrt{x+1}+1)^3}$	100

input

```
int(arccsch(x^(1/2))/x^4,x,method=_RETURNVERBOSE)
```

output

```
-1/3*arccsch(x^(1/2))/x^3-1/144*(x+1)^(1/2)*(15*arctanh(1/(x+1)^(1/2))*x^3-15*(x+1)^(1/2)*x^2+10*(x+1)^(1/2)*x-8*(x+1)^(1/2))/((x+1)/x)^(1/2)/x^(7/2)
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.50

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = \frac{(15x^2 - 10x + 8)\sqrt{x}\sqrt{\frac{x+1}{x}} - 3(5x^3 + 16)\log\left(\frac{x\sqrt{\frac{x+1}{x}} + \sqrt{x}}{x}\right)}{144x^3}$$

input

```
integrate(arccsch(x^(1/2))/x^4,x, algorithm="fricas")
```

output $1/144*((15*x^2 - 10*x + 8)*\sqrt{x}*\sqrt{(x + 1)/x} - 3*(5*x^3 + 16)*\log((x*\sqrt{(x + 1)/x} + \sqrt{x})/x))/x^3$

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\operatorname{acsch}(\sqrt{x})}{x^4} dx$$

input `integrate(acsch(x**(1/2))/x**4, x)`

output `Integral(acsch(sqrt(x))/x**4, x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.01

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = \frac{15 x^{\frac{5}{2}} \left(\frac{1}{x} + 1\right)^{\frac{5}{2}} - 40 x^{\frac{3}{2}} \left(\frac{1}{x} + 1\right)^{\frac{3}{2}} + 33 \sqrt{x} \sqrt{\frac{1}{x} + 1}}{144 \left(x^3 \left(\frac{1}{x} + 1\right)^3 - 3 x^2 \left(\frac{1}{x} + 1\right)^2 + 3 x \left(\frac{1}{x} + 1\right) - 1\right)} - \frac{\operatorname{arcsch}(\sqrt{x})}{3 x^3} - \frac{5}{96} \log \left(\sqrt{x} \sqrt{\frac{1}{x} + 1} + 1 \right) + \frac{5}{96} \log \left(\sqrt{x} \sqrt{\frac{1}{x} + 1} - 1 \right)$$

input `integrate(arccsch(x^(1/2))/x^4,x, algorithm="maxima")`

output $1/144*(15*x^(5/2)*(1/x + 1)^(5/2) - 40*x^(3/2)*(1/x + 1)^(3/2) + 33*\sqrt{x}*\sqrt{1/x + 1})/(x^3*(1/x + 1)^3 - 3*x^2*(1/x + 1)^2 + 3*x*(1/x + 1) - 1) - 1/3*\operatorname{arccsch}(\sqrt{x})/x^3 - 5/96*\log(\sqrt{x}*\sqrt{1/x + 1} + 1) + 5/96*\log(\sqrt{x}*\sqrt{1/x + 1} - 1)$

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\operatorname{arcsch}(\sqrt{x})}{x^4} dx$$

input `integrate(arccsch(x^(1/2))/x^4,x, algorithm="giac")`

output `integrate(arccsch(sqrt(x))/x^4, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right)}{x^4} dx$$

input `int(asinh(1/x^(1/2))/x^4,x)`

output `int(asinh(1/x^(1/2))/x^4, x)`

Reduce [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\operatorname{acsch}(\sqrt{x})}{x^4} dx$$

input `int(acsch(x^(1/2))/x^4,x)`

output `int(acsch(sqrt(x))/x**4,x)`

3.22 $\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx$

Optimal result	192
Mathematica [A] (verified)	192
Rubi [A] (verified)	193
Maple [A] (verified)	194
Fricas [A] (verification not implemented)	194
Sympy [A] (verification not implemented)	195
Maxima [A] (verification not implemented)	195
Giac [F]	195
Mupad [B] (verification not implemented)	196
Reduce [B] (verification not implemented)	196

Optimal result

Integrand size = 4, antiderivative size = 16

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = -\sqrt{1+x^2} + x \operatorname{arcsinh}(x)$$

output

```
-(x^2+1)^(1/2)+x*arcsinh(x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = -\sqrt{1+x^2} + x \operatorname{csch}^{-1}\left(\frac{1}{x}\right)$$

input

```
Integrate[ArcCsch[x^(-1)],x]
```

output

```
-Sqrt[1 + x^2] + x*ArcCsch[x^(-1)]
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6882, 6187, 241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx \\ & \quad \downarrow \text{6882} \\ & \int \operatorname{arcsinh}(x) dx \\ & \quad \downarrow \text{6187} \\ & x \operatorname{arcsinh}(x) - \int \frac{x}{\sqrt{x^2 + 1}} dx \\ & \quad \downarrow \text{241} \\ & x \operatorname{arcsinh}(x) - \sqrt{x^2 + 1} \end{aligned}$$

input `Int[ArcCsch[x^(-1)], x]`

output `-Sqrt[1 + x^2] + x*ArcSinh[x]`

Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

rule 6187 `Int[((a_.) + ArcSinh[(c_.)*(x_)])*(b_.))^(n_.), x_Symbol] :> Simp[x*(a + b*ArcSinh[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcSinh[c*x])^(n - 1)/Sqrt[1 + c^2*x^2]), x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

rule 6882

```
Int[ArcCsch[(c_.)/((a_.) + (b_.)*(x_)^(n_.))]^(m_.)*(u_.), x_Symbol] := Int
[u*ArcSinh[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]
```

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

method	result	size
parts	$x \operatorname{arccsch}\left(\frac{1}{x}\right) - \sqrt{x^2 + 1}$	17
derivativedivides	$x \operatorname{arccsch}\left(\frac{1}{x}\right) - \frac{x^2\left(\frac{1}{x^2}+1\right)}{\sqrt{\left(\frac{1}{x^2}+1\right)x^2}}$	29
default	$x \operatorname{arccsch}\left(\frac{1}{x}\right) - \frac{x^2\left(\frac{1}{x^2}+1\right)}{\sqrt{\left(\frac{1}{x^2}+1\right)x^2}}$	29

```
input int(arccsch(1/x), x, method=_RETURNVERBOSE)
```

```
output x*arccsch(1/x)-(x^2+1)^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = x \log\left(x + \sqrt{x^2 + 1}\right) - \sqrt{x^2 + 1}$$

```
input integrate(arccsch(1/x), x, algorithm="fricas")
```

```
output x*log(x + sqrt(x^2 + 1)) - sqrt(x^2 + 1)
```

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = x \operatorname{acsch}\left(\frac{1}{x}\right) - \sqrt{x^2 + 1}$$

input `integrate(acsch(1/x), x)`output `x*acsch(1/x) - sqrt(x**2 + 1)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = x \operatorname{arcsch}\left(\frac{1}{x}\right) - \sqrt{x^2 + 1}$$

input `integrate(arccsch(1/x), x, algorithm="maxima")`output `x*arccsch(1/x) - sqrt(x^2 + 1)`**Giac [F]**

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = \int \operatorname{arcsch}\left(\frac{1}{x}\right) dx$$

input `integrate(arccsch(1/x), x, algorithm="giac")`output `integrate(arccsch(1/x), x)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = x \operatorname{asinh}(x) - \sqrt{x^2 + 1}$$

input `int(asinh(x),x)`

output `x*asinh(x) - (x^2 + 1)^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = \operatorname{asinh}(x) x - \sqrt{x^2 + 1}$$

input `int(acsch(1/x),x)`

output `asinh(x)*x - sqrt(x**2 + 1)`

3.23 $\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx$

Optimal result	197
Mathematica [C] (verified)	197
Rubi [C] (warning: unable to verify)	198
Maple [F]	201
Fricas [F(-2)]	201
Sympy [F]	202
Maxima [F]	202
Giac [F]	202
Mupad [F(-1)]	203
Reduce [F]	203

Optimal result

Integrand size = 10, antiderivative size = 61

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = \frac{\operatorname{csch}^{-1}(ax^n)^2}{2n} - \frac{\operatorname{csch}^{-1}(ax^n) \log\left(1 - e^{2\operatorname{csch}^{-1}(ax^n)}\right)}{n} - \frac{\operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(ax^n)}\right)}{2n}$$

output

$1/2*\operatorname{arccsch}(a*x^n)^2/n - \operatorname{arccsch}(a*x^n)*\ln(1 - (1/a/(x^n) + (1+1/a^2/(x^n)^2)^(1/2))^2)/n - 1/2*\operatorname{polylog}(2, (1/a/(x^n) + (1+1/a^2/(x^n)^2)^(1/2))^2)/n$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.06 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.05

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = -\frac{x^{-n} {}_3F_2\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}; \frac{3}{2}, \frac{3}{2}; -\frac{x^{-2n}}{a^2}\right)}{an} + \left(\operatorname{csch}^{-1}(ax^n) - \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)\right) \log(x)$$

input `Integrate[ArcCsch[a*x^n]/x,x]`

output `-(HypergeometricPFQ[{1/2, 1/2, 1/2}, {3/2, 3/2}, -(1/(a^2*x^(2*n)))]/(a*n*x^n)) + (ArcCsch[a*x^n] - ArcSinh[1/(a*x^n)])*Log[x]`

Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.49 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.26, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {7282, 6836, 6190, 3042, 26, 4199, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx \\
 & \quad \downarrow \text{7282} \\
 & \frac{\int x^{-n} \operatorname{csch}^{-1}(ax^n) dx^n}{n} \\
 & \quad \downarrow \text{6836} \\
 & \frac{\int x^{-n} \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right) dx^{-n}}{n} \\
 & \quad \downarrow \text{6190} \\
 & \frac{\int ax^n \sqrt{\frac{x^{-2n}}{a^2} + 1} \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right) d\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)}{n} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int -i \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right) \tan\left(i \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right) + \frac{\pi}{2}\right) d\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)}{n} \\
 & \quad \downarrow \text{26} \\
 & \frac{i \int \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right) \tan\left(i \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right) + \frac{\pi}{2}\right) d\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)}{n}
 \end{aligned}$$

$$\begin{array}{c}
\downarrow 4199 \\
\frac{i \left(2i \int -\frac{e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)} \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)}{1-e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)}} d\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right) - \frac{1}{2}ix^{2n} \right)}{n} \\
\downarrow 25 \\
\frac{i \left(-2i \int \frac{e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)} \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)}{1-e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)}} d\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right) - \frac{1}{2}ix^{2n} \right)}{n} \\
\downarrow 2620 \\
\frac{i \left(-2i \left(\frac{1}{2} \int \log \left(1 - e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)} \right) d\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right) - \frac{1}{2} \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right) \log \left(1 - e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)} \right) \right) - \frac{1}{2}ix^{2n} \right)}{n} \\
\downarrow 2715 \\
\frac{i \left(-2i \left(\frac{1}{4} \int e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)} \log \left(1 - e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)} \right) de^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)} - \frac{1}{2} \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right) \log \left(1 - e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)} \right) \right) \right)}{n} \\
\downarrow 2838 \\
\frac{i \left(-2i \left(-\frac{1}{4} \operatorname{PolyLog} \left(2, e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)} \right) - \frac{1}{2} \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right) \log \left(1 - e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)} \right) \right) - \frac{1}{2}ix^{2n} \right)}{n}
\end{array}$$

input `Int[ArcCsch[a*x^n]/x,x]`

output `(I*((-1/2*I)*x^(2*n) - (2*I)*(-1/2*(ArcSinh[1/(a*x^n)]*Log[1 - E^(2*ArcSinh[1/(a*x^n)])]) - PolyLog[2, E^(2*ArcSinh[1/(a*x^n)])]/4)))/n`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 26 $\text{Int}[(\text{Complex}[0, \text{a}_])*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], \text{a}]) \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{EqQ}[\text{a}^2, 1]$
- rule 2620 $\text{Int}[(((\text{F}_)^{((\text{g}_)*((\text{e}_.) + (\text{f}_.)*(x_))))^{(n_.)}*((\text{c}_.) + (\text{d}_.)*(x_))^{(m_.)})/((\text{a}_.) + (\text{b}_.)*((\text{F}_)^{((\text{g}_)*((\text{e}_.) + (\text{f}_.)*(x_))))^{(n_.)}), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{d}*x)^m/(\text{b}*f*g*n*\text{Log}[\text{F}]))*\text{Log}[1 + \text{b}*((\text{F}^{(g*(e + f*x))})^n/\text{a})], \text{x}] - \text{Simp}[\text{d}*(m/(\text{b}*f*g*n*\text{Log}[\text{F}])) \quad \text{Int}[(\text{c} + \text{d}*x)^{(m-1)}*\text{Log}[1 + \text{b}*((\text{F}^{(g*(e + f*x))})^n/\text{a})], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{n}\}, \text{x}] \ \&\& \ \text{IGtQ}[\text{m}, 0]$
- rule 2715 $\text{Int}[\text{Log}[(\text{a}_.) + (\text{b}_.)*((\text{F}_)^{((\text{e}_.)*((\text{c}_.) + (\text{d}_.)*(x_))))^{(n_.)}], \text{x_Symbol}] \rightarrow \text{Simp}[1/(\text{d}*e*n*\text{Log}[\text{F}]) \quad \text{Subst}[\text{Int}[\text{Log}[\text{a} + \text{b}*x]/x, \text{x}], \text{x}, (\text{F}^{(e*(c + d*x))})^n], \text{x}] \text{ ; FreeQ}[\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{n}\}, \text{x}] \ \&\& \ \text{GtQ}[\text{a}, 0]$
- rule 2838 $\text{Int}[\text{Log}[(\text{c}_.)*((\text{d}_.) + (\text{e}_.)*(x_))^{(n_.)}]/(x_), \text{x_Symbol}] \rightarrow \text{Simp}[-\text{PolyLog}[2, (\text{-c})*e*x^n]/n, \text{x}] \text{ ; FreeQ}[\{\text{c}, \text{d}, \text{e}, \text{n}\}, \text{x}] \ \&\& \ \text{EqQ}[\text{c}*d, 1]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4199 $\text{Int}[(((\text{c}_.) + (\text{d}_.)*(x_))^{(m_.)}*\text{tan}[(\text{e}_.) + \text{Pi}*(\text{k}_.) + (\text{Complex}[0, \text{fz}_])*(\text{f}_.)*(x_)]), \text{x_Symbol}] \rightarrow \text{Simp}[(-1)*((\text{c} + \text{d}*x)^{(m+1})/(\text{d}*(m+1))), \text{x}] + \text{Simp}[2*I \quad \text{Int}[((\text{c} + \text{d}*x)^m*(\text{E}^{(2*((-1)*e + f*fz*x))}/(1 + \text{E}^{(2*((-1)*e + f*fz*x))})/\text{E}^{(2*I*k*Pi)})))/\text{E}^{(2*I*k*Pi)}, \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{c}, \text{d}, \text{e}, \text{f}, \text{fz}\}, \text{x}] \ \&\& \ \text{IntegerQ}[4*k] \ \&\& \ \text{IGtQ}[\text{m}, 0]$
- rule 6190 $\text{Int}[((\text{a}_.) + \text{ArcSinh}[(\text{c}_.)*(x_)]*(\text{b}_.))^{(n_.)}/(x_), \text{x_Symbol}] \rightarrow \text{Simp}[1/\text{b} \quad \text{Subst}[\text{Int}[\text{x}^n*\text{Coth}[-\text{a}/\text{b} + \text{x}/\text{b}], \text{x}], \text{x}, \text{a} + \text{b}*ArcSinh[\text{c}*x]], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}\}, \text{x}] \ \&\& \ \text{IGtQ}[\text{n}, 0]$

rule 6836 `Int[((a_.) + ArcCsch[(c_.)*(x_.)]*(b_.))/(x_), x_Symbol] := -Subst[Int[(a + b*ArcSinh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]`

rule 7282 `Int[(u_)/(x_), x_Symbol] := With[{lst = PowerVariableExpn[u, 0, x]}, Simp[1/lst[[2]] Subst[Int[NormalizeIntegrand[Simplify[lst[[1]]/x], x], x], x, (lst[[3]]*x)^lst[[2]]], x] /; !FalseQ[lst] && NeQ[lst[[2]], 0] /; NonsumQ[u] && !RationalFunctionQ[u, x]`

Maple **[F]**

$$\int \frac{\operatorname{arccsch}(ax^n)}{x} dx$$

input `int(arccsch(a*x^n)/x,x)`

output `int(arccsch(a*x^n)/x,x)`

Fricas **[F(-2)]**

Exception generated.

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = \text{Exception raised: TypeError}$$

input `integrate(arccsch(a*x^n)/x,x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{acsch}(ax^n)}{x} dx$$

input `integrate(acsch(a*x**n)/x,x)`

output `Integral(acsch(a*x**n)/x, x)`

Maxima [F]

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{arcsch}(ax^n)}{x} dx$$

input `integrate(arccsch(a*x^n)/x,x, algorithm="maxima")`

output `a^2*n*integrate(x^(2*n)*log(x)/(a^2*x*x^(2*n) + (a^2*x*x^(2*n) + x)*sqrt(a^2*x^(2*n) + 1) + x), x) + n*integrate(log(x)/(a^2*x*x^(2*n) + x), x) - log(a)*log(x) - log(x)*log(x^n) + log(x)*log(sqrt(a^2*x^(2*n) + 1) + 1)`

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{arcsch}(ax^n)}{x} dx$$

input `integrate(arccsch(a*x^n)/x,x, algorithm="giac")`

output `integrate(arccsch(a*x^n)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{ax^n}\right)}{x} dx$$

input `int(asinh(1/(a*x^n))/x,x)`output `int(asinh(1/(a*x^n))/x, x)`**Reduce [F]**

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{acsch}(x^na)}{x} dx$$

input `int(acsch(a*x^n)/x,x)`output `int(acsch(x**n*a)/x,x)`

3.24 $\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx$

Optimal result	204
Mathematica [A] (verified)	204
Rubi [C] (warning: unable to verify)	205
Maple [F]	208
Fricas [F]	208
Sympy [F]	208
Maxima [F]	209
Giac [F]	209
Mupad [F(-1)]	209
Reduce [F]	210

Optimal result

Integrand size = 10, antiderivative size = 54

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = \frac{1}{10} \operatorname{csch}^{-1}(ax^5)^2 - \frac{1}{5} \operatorname{csch}^{-1}(ax^5) \log\left(1 - e^{2\operatorname{csch}^{-1}(ax^5)}\right) - \frac{1}{10} \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(ax^5)}\right)$$

output

```
1/10*arccsch(a*x^5)^2-1/5*arccsch(a*x^5)*ln(1-(1/a/x^5+(1+1/a^2/x^10)^(1/2))^-2)-1/10*polylog(2,(1/a/x^5+(1+1/a^2/x^10)^(1/2))^-2)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.93

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = \frac{1}{10} \left(\operatorname{csch}^{-1}(ax^5)^2 - 2\operatorname{csch}^{-1}(ax^5) \log\left(1 - e^{2\operatorname{csch}^{-1}(ax^5)}\right) - \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(ax^5)}\right) \right)$$

input

```
Integrate[ArcCsch[a*x^5]/x,x]
```

output

```
(ArcCsch[a*x^5]^2 - 2*ArcCsch[a*x^5]*Log[1 - E^(2*ArcCsch[a*x^5])] - PolyLog[2, E^(2*ArcCsch[a*x^5])])/10
```

Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.50 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.26, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {7282, 6836, 6190, 3042, 26, 4199, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx$$

$$\downarrow 7282$$

$$\frac{1}{5} \int \frac{\operatorname{csch}^{-1}(ax^5)}{x^5} dx^5$$

$$\downarrow 6836$$

$$-\frac{1}{5} \int \frac{\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)}{x^5} d\frac{1}{x^5}$$

$$\downarrow 6190$$

$$-\frac{1}{5} \int a \sqrt{1 + \frac{1}{x^{10}a^2}} x^5 \operatorname{arcsinh}\left(\frac{1}{ax^5}\right) d\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)$$

$$\downarrow 3042$$

$$-\frac{1}{5} \int -i \operatorname{arcsinh}\left(\frac{1}{ax^5}\right) \tan\left(i \operatorname{arcsinh}\left(\frac{1}{ax^5}\right) + \frac{\pi}{2}\right) d\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)$$

$$\downarrow 26$$

$$\frac{1}{5} i \int \operatorname{arcsinh}\left(\frac{1}{ax^5}\right) \tan\left(i \operatorname{arcsinh}\left(\frac{1}{ax^5}\right) + \frac{\pi}{2}\right) d\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)$$

$$\downarrow 4199$$

$$\frac{1}{5}i \left(2i \int -\frac{e^{2\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)} \operatorname{arcsinh}\left(\frac{1}{ax^5}\right)}{1 - e^{2\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)}} d\operatorname{arcsinh}\left(\frac{1}{ax^5}\right) - \frac{ix^{10}}{2} \right)$$

↓ 25

$$\frac{1}{5}i \left(-2i \int \frac{e^{2\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)} \operatorname{arcsinh}\left(\frac{1}{ax^5}\right)}{1 - e^{2\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)}} d\operatorname{arcsinh}\left(\frac{1}{ax^5}\right) - \frac{ix^{10}}{2} \right)$$

↓ 2620

$$\frac{1}{5}i \left(-2i \left(\frac{1}{2} \int \log \left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)} \right) d\operatorname{arcsinh}\left(\frac{1}{ax^5}\right) - \frac{1}{2} \operatorname{arcsinh}\left(\frac{1}{ax^5}\right) \log \left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)} \right) \right) - \frac{ix^{10}}{2} \right)$$

↓ 2715

$$\frac{1}{5}i \left(-2i \left(\frac{1}{4} \int e^{2\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)} \log \left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)} \right) de^{2\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)} - \frac{1}{2} \operatorname{arcsinh}\left(\frac{1}{ax^5}\right) \log \left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)} \right) \right) \right)$$

↓ 2838

$$\frac{1}{5}i \left(-2i \left(-\frac{1}{4} \operatorname{PolyLog} \left(2, e^{2\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)} \right) - \frac{1}{2} \operatorname{arcsinh}\left(\frac{1}{ax^5}\right) \log \left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{ax^5}\right)} \right) \right) - \frac{ix^{10}}{2} \right)$$

input `Int[ArcCsch[a*x^5]/x,x]`

output `(I/5)*((-1/2*I)*x^10 - (2*I)*(-1/2*(ArcSinh[1/(a*x^5)]*Log[1 - E^(2*ArcSinh[1/(a*x^5)])]) - PolyLog[2, E^(2*ArcSinh[1/(a*x^5)])]/4))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2715

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x
)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

rule 2838

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4199

```
Int[(((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_
.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp
[2*I Int[(((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x
)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && In
tegerQ[4*k] && IGtQ[m, 0]
```

rule 6190

```
Int[(((a_) + ArcSinh[(c_)*(x_)])*(b_))^(n_)/(x_), x_Symbol] := Simp[1/b
Subst[Int[x^n*Coth[-a/b + x/b], x], x, a + b*ArcSinh[c*x], x] /; FreeQ[{a
, b, c}, x] && IGtQ[n, 0]
```

rule 6836

```
Int[(((a_) + ArcSch[(c_)*(x_)])*(b_))/(x_), x_Symbol] := -Subst[Int[(a +
b*ArcSinh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]
```


rule 7282

```
Int[(u_)/(x_), x_Symbol] := With[{lst = PowerVariableExpn[u, 0, x]}, Simp[1
/lst[[2]] Subst[Int[NormalizeIntegrand[Simplify[lst[[1]]/x], x], x], x, (
lst[[3]]*x)^lst[[2]]], x] /; !FalseQ[lst] && NeQ[lst[[2]], 0] /; NonsumQ[
u] && !RationalFunctionQ[u, x]
```

Maple [F]

$$\int \frac{\operatorname{arccsch}(ax^5)}{x} dx$$

input `int(arccsch(a*x^5)/x,x)`output `int(arccsch(a*x^5)/x,x)`**Fricas [F]**

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arcsch}(ax^5)}{x} dx$$

input `integrate(arccsch(a*x^5)/x,x, algorithm="fricas")`output `integral(arccsch(a*x^5)/x, x)`**Sympy [F]**

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{acsch}(ax^5)}{x} dx$$

input `integrate(acsch(a*x**5)/x,x)`output `Integral(acsch(a*x**5)/x, x)`

Maxima [F]

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arcsch}(ax^5)}{x} dx$$

input `integrate(arccsch(a*x^5)/x,x, algorithm="maxima")`

output `5*a^2*integrate(x^9*log(x)/(a^2*x^10 + (a^2*x^10 + 1)^(3/2) + 1), x) - 1/2*log(a^2*x^10 + 1)*log(x) - log(a)*log(x) - 5/2*log(x)^2 + log(x)*log(sqrt(a^2*x^10 + 1) + 1) - 1/20*dilog(-a^2*x^10)`

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arcsch}(ax^5)}{x} dx$$

input `integrate(arccsch(a*x^5)/x,x, algorithm="giac")`

output `integrate(arccsch(a*x^5)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{ax^5}\right)}{x} dx$$

input `int(asinh(1/(a*x^5))/x,x)`

output `int(asinh(1/(a*x^5))/x, x)`

Reduce [F]

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{acsch}(ax^5)}{x} dx$$

input `int(acsch(a*x^5)/x,x)`

output `int(acsch(a*x**5)/x,x)`

3.25 $\int \operatorname{csch}^{-1}(ce^{a+bx}) dx$

Optimal result	211
Mathematica [B] (verified)	211
Rubi [C] (warning: unable to verify)	212
Maple [F]	215
Fricas [F(-2)]	215
Sympy [F]	216
Maxima [F]	216
Giac [F]	216
Mupad [F(-1)]	217
Reduce [F]	217

Optimal result

Integrand size = 10, antiderivative size = 77

$$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx = \frac{\operatorname{csch}^{-1}(ce^{a+bx})^2}{2b} - \frac{\operatorname{csch}^{-1}(ce^{a+bx}) \log(1 - e^{2\operatorname{csch}^{-1}(ce^{a+bx})})}{b} - \frac{\operatorname{PolyLog}(2, e^{2\operatorname{csch}^{-1}(ce^{a+bx})})}{2b}$$

```
output 1/2*arccsch(c*exp(b*x+a))^2/b-arccsch(c*exp(b*x+a))*ln(1-(1/c/(exp(1)^(b*x+a)))+(1+1/c^2/(exp(1)^(b*x+a))^2)^(1/2))^2)/b-1/2*polylog(2,(1/c/(exp(1)^(b*x+a)))+(1+1/c^2/(exp(1)^(b*x+a))^2)^(1/2))^2)/b
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 236 vs. 2(77) = 154.

Time = 0.68 (sec) , antiderivative size = 236, normalized size of antiderivative = 3.06

$$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx = x\operatorname{csch}^{-1}(ce^{a+bx}) + \frac{e^{-a-bx}\sqrt{1+c^2e^{2(a+bx)}}\left(\log^2(-c^2e^{2(a+bx)}) + \operatorname{arctanh}\left(\sqrt{1+c^2e^{2(a+bx)}}\right)(-8bx + 4\log(-c^2e^{2(a+bx)}))\right)}{2}$$

input `Integrate[ArcCsch[c*E^(a + b*x)],x]`

output `x*ArcCsch[c*E^(a + b*x)] + (E^(-a - b*x)*Sqrt[1 + c^2*E^(2*(a + b*x))]*(Log[-(c^2*E^(2*(a + b*x))])^2 + ArcTanh[Sqrt[1 + c^2*E^(2*(a + b*x))]]*(-8*b*x + 4*Log[-(c^2*E^(2*(a + b*x))])) - 4*Log[-(c^2*E^(2*(a + b*x))])*Log[(1 + Sqrt[1 + c^2*E^(2*(a + b*x))])/2] + 2*Log[(1 + Sqrt[1 + c^2*E^(2*(a + b*x))])/2]^2 - 4*PolyLog[2, (1 - Sqrt[1 + c^2*E^(2*(a + b*x))])/2])/(8*b*c*Sqrt[1 + 1/(c^2*E^(2*(a + b*x)))]))`

Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.48 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.26, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {2720, 6836, 6190, 3042, 26, 4199, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \operatorname{csch}^{-1}(ce^{a+bx}) dx \\
 & \quad \downarrow 2720 \\
 & \frac{\int e^{-a-bx} \operatorname{csch}^{-1}(ce^{a+bx}) de^{a+bx}}{b} \\
 & \quad \downarrow 6836 \\
 & -\frac{\int e^{-a-bx} \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right) de^{-a-bx}}{b} \\
 & \quad \downarrow 6190 \\
 & -\frac{\int ce^{a+bx} \sqrt{1 + \frac{e^{-2a-2bx}}{c^2}} \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right) d\operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}{b} \\
 & \quad \downarrow 3042 \\
 & -\frac{\int -i \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right) \tan\left(i \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right) + \frac{\pi}{2}\right) d\operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}{b}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 26 \\
& \frac{i \int \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right) \tan\left(i \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right) + \frac{\pi}{2}\right) d \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}{b} \\
& \downarrow 4199 \\
& \frac{i \left(2i \int -\frac{e^{a+bx+2 \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}}{1-e^{2 \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}} d \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right) - \frac{1}{2} i e^{2a+2bx} \right)}{b} \\
& \downarrow 25 \\
& \frac{i \left(-2i \int \frac{e^{a+bx+2 \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}}{1-e^{2 \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}} d \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right) - \frac{1}{2} i e^{2a+2bx} \right)}{b} \\
& \downarrow 2620 \\
& \frac{i \left(-2i \left(\frac{1}{2} \int \log\left(1 - e^{2 \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}\right) d \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right) - \frac{1}{2} \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 - e^{2 \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}\right) \right) \right) - \frac{1}{2}}{b} \\
& \downarrow 2715 \\
& \frac{i \left(-2i \left(\frac{1}{4} \int e^{2 \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)} \log\left(1 - e^{2 \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}\right) d e^{2 \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)} - \frac{1}{2} \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 - e^{2 \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}\right) \right) \right)}{b} \\
& \downarrow 2838 \\
& \frac{i \left(-2i \left(-\frac{1}{4} \operatorname{PolyLog}\left(2, e^{2 \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}\right) - \frac{1}{2} \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 - e^{2 \operatorname{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}\right) \right) \right) - \frac{1}{2} i e^{2a+2bx}}{b}
\end{aligned}$$

input `Int[ArcCsch[c*E^(a + b*x)],x]`

output `(I*((-1/2*I)*E^(2*a + 2*b*x) - (2*I)*(-1/2*(ArcSinh[E^(-a - b*x)/c]*Log[1 - E^(2*ArcSinh[E^(-a - b*x)/c]]) - PolyLog[2, E^(2*ArcSinh[E^(-a - b*x)/c]])/4)))/b`

Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))* (F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4199 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 6190 `Int[((a_.) + ArcSinh[(c_.)*(x_)*(b_.)]^(n_.))/(x_), x_Symbol] := Simp[1/b Subst[Int[x^n*Coth[-a/b + x/b], x], x, a + b*ArcSinh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

rule 6836 `Int[((a_.) + ArcCsch[(c_.)*(x_)*(b_.)]/(x_), x_Symbol] := -Subst[Int[(a + b*ArcSinh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]`

Maple [F]

$$\int \operatorname{arccsch}(e^{bx+a}c) dx$$

input `int(arccsch(exp(b*x+a)*c),x)`

output `int(arccsch(exp(b*x+a)*c),x)`

Fricas [F(-2)]

Exception generated.

$$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx = \text{Exception raised: TypeError}$$

input `integrate(arccsch(c*exp(b*x+a)),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

Sympy [F]

$$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx = \int \operatorname{acsch}(ce^{a+bx}) dx$$

input `integrate(acsch(c*exp(b*x+a)),x)`

output `Integral(acsch(c*exp(a + b*x)), x)`

Maxima [F]

$$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx = \int \operatorname{arcsch}(ce^{(bx+a)}) dx$$

input `integrate(arccsch(c*exp(b*x+a)),x, algorithm="maxima")`

output `b*c^2*integrate(x*e^(2*b*x + 2*a)/(c^2*e^(2*b*x + 2*a) + (c^2*e^(2*b*x + 2*a) + 1)^(3/2) + 1), x) - 1/2*b*x^2 - (a + log(c))*x + x*log(sqrt(c^2*e^(2*b*x + 2*a) + 1) + 1) - 1/4*(2*b*x*log(c^2*e^(2*b*x + 2*a) + 1) + dilog(-c^2*e^(2*b*x + 2*a)))/b`

Giac [F]

$$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx = \int \operatorname{arcsch}(ce^{(bx+a)}) dx$$

input `integrate(arccsch(c*exp(b*x+a)),x, algorithm="giac")`

output `integrate(arccsch(c*e^(b*x + a)), x)`

Mupad [F(-1)]

Timed out.

$$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx = \int \operatorname{asinh}\left(\frac{e^{-a-bx}}{c}\right) dx$$

input `int(asinh(exp(- a - b*x)/c),x)`output `int(asinh(exp(- a - b*x)/c), x)`**Reduce [F]**

$$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx = \int \operatorname{acsch}(e^{bx+a}c) dx$$

input `int(acsch(c*exp(b*x+a)),x)`output `int(acsch(e**(a + b*x)*c),x)`

3.26 $\int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx$

Optimal result	218
Mathematica [A] (verified)	218
Rubi [C] (warning: unable to verify)	219
Maple [F]	222
Fricas [F]	223
Sympy [F]	223
Maxima [F]	223
Giac [F]	224
Mupad [F(-1)]	224
Reduce [F]	224

Optimal result

Integrand size = 19, antiderivative size = 61

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \frac{\operatorname{csch}^{-1}(a+bx)^2}{2d} - \frac{\operatorname{csch}^{-1}(a+bx) \log\left(1 - e^{2\operatorname{csch}^{-1}(a+bx)}\right)}{d} - \frac{\operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(a+bx)}\right)}{2d}$$

output

```
1/2*arccsch(b*x+a)^2/d-arccsch(b*x+a)*ln(1-(1/(b*x+a)+(1+1/(b*x+a)^2)^(1/2))^2)/d-1/2*polylog(2,(1/(b*x+a)+(1+1/(b*x+a)^2)^(1/2))^2)/d
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.87

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \frac{\operatorname{csch}^{-1}(a+bx)^2 - 2\operatorname{csch}^{-1}(a+bx) \log\left(1 - e^{2\operatorname{csch}^{-1}(a+bx)}\right) - \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(a+bx)}\right)}{2d}$$

input `Integrate[ArcCsch[a + b*x]/((a*d)/b + d*x), x]`

output `(ArcCsch[a + b*x]^2 - 2*ArcCsch[a + b*x]*Log[1 - E^(2*ArcCsch[a + b*x])] - PolyLog[2, E^(2*ArcCsch[a + b*x])])/(2*d)`

Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.53 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.20, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.579$, Rules used = {6874, 27, 6836, 6190, 3042, 26, 4199, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{ad}{b} + dx} dx \\
 & \quad \downarrow \text{6874} \\
 & \int \frac{b \operatorname{csch}^{-1}(a+bx) d(a+bx)}{d(a+bx)} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{\operatorname{csch}^{-1}(a+bx) d(a+bx)}{a+bx} \\
 & \quad \downarrow \text{6836} \\
 & \int (a+bx) \operatorname{arcsinh}\left(\frac{1}{a+bx}\right) d\frac{1}{a+bx} \\
 & \quad \downarrow \text{6190} \\
 & \int (a+bx) \sqrt{1 + \frac{1}{(a+bx)^2}} \operatorname{arcsinh}\left(\frac{1}{a+bx}\right) d\operatorname{arcsinh}\left(\frac{1}{a+bx}\right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{\int -i \operatorname{arcsinh}\left(\frac{1}{a+bx}\right) \tan\left(i \operatorname{arcsinh}\left(\frac{1}{a+bx}\right) + \frac{\pi}{2}\right) d \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}{d} \\
& \quad \downarrow \text{26} \\
& \frac{i \int \operatorname{arcsinh}\left(\frac{1}{a+bx}\right) \tan\left(i \operatorname{arcsinh}\left(\frac{1}{a+bx}\right) + \frac{\pi}{2}\right) d \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}{d} \\
& \quad \downarrow \text{4199} \\
& \frac{i \left(2i \int -\frac{e^{2 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)} \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}{1 - e^{2 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}} d \operatorname{arcsinh}\left(\frac{1}{a+bx}\right) - \frac{i}{2(a+bx)^2} \right)}{d} \\
& \quad \downarrow \text{25} \\
& \frac{i \left(-2i \int \frac{e^{2 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)} \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}{1 - e^{2 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}} d \operatorname{arcsinh}\left(\frac{1}{a+bx}\right) - \frac{i}{2(a+bx)^2} \right)}{d} \\
& \quad \downarrow \text{2620} \\
& \frac{i \left(-2i \left(\frac{1}{2} \int \log\left(1 - e^{2 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}\right) d \operatorname{arcsinh}\left(\frac{1}{a+bx}\right) - \frac{1}{2} \operatorname{arcsinh}\left(\frac{1}{a+bx}\right) \log\left(1 - e^{2 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}\right) \right) - \frac{i}{2(a+bx)^2} \right)}{d} \\
& \quad \downarrow \text{2715} \\
& \frac{i \left(-2i \left(\frac{1}{4} \int (a+bx) \log(-a-bx+1) d e^{2 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)} - \frac{1}{2} \operatorname{arcsinh}\left(\frac{1}{a+bx}\right) \log\left(1 - e^{2 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}\right) \right) - \frac{i}{2(a+bx)^2} \right)}{d} \\
& \quad \downarrow \text{2838} \\
& \frac{i \left(-2i \left(-\frac{1}{4} \operatorname{PolyLog}\left(2, e^{2 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}\right) - \frac{1}{2} \operatorname{arcsinh}\left(\frac{1}{a+bx}\right) \log\left(1 - e^{2 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}\right) \right) - \frac{i}{2(a+bx)^2} \right)}{d}
\end{aligned}$$

input

```
Int[ArcCsch[a + b*x]/((a*d)/b + d*x), x]
```

output $(I*((-1/2*I)/(a + b*x)^2 - (2*I)*(-1/2*(ArcSinh[(a + b*x)^{-1}]*Log[1 - E^{(2*ArcSinh[(a + b*x)^{-1}]])]) - PolyLog[2, E^{(2*ArcSinh[(a + b*x)^{-1}]])})/4))/d$

Defintions of rubi rules used

rule 25 $Int[-(Fx_), x_Symbol] \rightarrow Simp[Identity[-1] Int[Fx, x], x]$

rule 26 $Int[(Complex[0, a_]*)(Fx_), x_Symbol] \rightarrow Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] \&\& EqQ[a^2, 1]$

rule 27 $Int[(a_)*(Fx_), x_Symbol] \rightarrow Simp[a Int[Fx, x], x] /; FreeQ[a, x] \&\& !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]$

rule 2620 $Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^((n_)*((c_) + (d_)*(x_))^(m_)))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^((n_))), x_Symbol] \rightarrow Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] \&\& IGtQ[m, 0]$

rule 2715 $Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^((n_))], x_Symbol] \rightarrow Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] \&\& GtQ[a, 0]$

rule 2838 $Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] \rightarrow Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] \&\& EqQ[c*d, 1]$

rule 3042 $Int[u_, x_Symbol] \rightarrow Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]$

rule 4199 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 6190 `Int[((a_.) + ArcSinh[(c_.)*(x_)]*(b_.))^(n_.)/(x_), x_Symbol] := Simp[1/b Subst[Int[x^n*Coth[-a/b + x/b], x], x, a + b*ArcSinh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

rule 6836 `Int[((a_.) + ArcCsch[(c_.)*(x_)]*(b_.))/(x_), x_Symbol] := -Subst[Int[(a + b*ArcSinh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]`

rule 6874 `Int[((a_.) + ArcCsch[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_)^(m_.), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcCsch[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]`

Maple **[F]**

$$\int \frac{\operatorname{arccsch}(bx + a)}{\frac{ad}{b} + dx} dx$$

input `int(arccsch(b*x+a)/(a*d/b+d*x),x)`

output `int(arccsch(b*x+a)/(a*d/b+d*x),x)`

Fricas [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{arcsch}(bx + a)}{dx + \frac{ad}{b}} dx$$

input `integrate(arccsch(b*x+a)/(a*d/b+d*x),x, algorithm="fricas")`

output `integral(b*arccsch(b*x + a)/(b*d*x + a*d), x)`

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \frac{b \int \frac{\operatorname{acsch}(a+bx)}{a+bx} dx}{d}$$

input `integrate(acsch(b*x+a)/(a*d/b+d*x),x)`

output `b*Integral(acsch(a + b*x)/(a + b*x), x)/d`

Maxima [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{arcsch}(bx + a)}{dx + \frac{ad}{b}} dx$$

input `integrate(arccsch(b*x+a)/(a*d/b+d*x),x, algorithm="maxima")`

output `-1/4*(2*log(b^2*x^2 + 2*a*b*x + a^2 + 1)*log(b*x + a) + dilog(-b^2*x^2 - 2*a*b*x - a^2))/d - 1/2*(log(b*x + a)^2 - 2*log(b*x + a)*log(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) + 1))/d + integrate((b^2*x + a*b)*log(b*x + a)/(b^2*d*x^2 + 2*a*b*d*x + a^2*d + (b^2*d*x^2 + 2*a*b*d*x + a^2*d + d)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) + d), x)`

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{arcsch}(bx + a)}{dx + \frac{ad}{b}} dx$$

input `integrate(arccsch(b*x+a)/(a*d/b+d*x),x, algorithm="giac")`

output `integrate(arccsch(b*x + a)/(d*x + a*d/b), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{a+bx}\right)}{dx + \frac{ad}{b}} dx$$

input `int(asinh(1/(a + b*x)))/(d*x + (a*d)/b),x)`

output `int(asinh(1/(a + b*x)))/(d*x + (a*d)/b), x)`

Reduce [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \frac{\left(\int \frac{\operatorname{acsch}(bx+a)}{bx+a} dx\right) b}{d}$$

input `int(acsch(b*x+a)/(a*d/b+d*x),x)`

output `(int(acsch(a + b*x)/(a + b*x),x)*b)/d`

3.27 $\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx$

Optimal result	225
Mathematica [B] (verified)	225
Rubi [A] (warning: unable to verify)	226
Maple [A] (verified)	228
Fricas [B] (verification not implemented)	228
Sympy [F(-1)]	229
Maxima [A] (verification not implemented)	229
Giac [F]	230
Mupad [B] (verification not implemented)	230
Reduce [F]	230

Optimal result

Integrand size = 12, antiderivative size = 46

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx = \frac{(a + bx^4) \operatorname{csch}^{-1}(a + bx^4)}{4b} + \frac{\operatorname{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx^4)^2}}\right)}{4b}$$

output

```
1/4*(b*x^4+a)*arccsch(b*x^4+a)/b+1/4*arctanh((1+1/(b*x^4+a)^2)^(1/2))/b
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 95 vs. 2(46) = 92.

Time = 0.10 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.07

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx = \frac{(a + bx^4) \operatorname{csch}^{-1}(a + bx^4)}{4b} - \frac{\sqrt{1 + (a + bx^4)^2} \log\left(-a - bx^4 + \sqrt{1 + (a + bx^4)^2}\right)}{4b(a + bx^4) \sqrt{1 + \frac{1}{(a+bx^4)^2}}}$$

input

```
Integrate[x^3*ArcCsch[a + b*x^4],x]
```

output

```
((a + b*x^4)*ArcCsch[a + b*x^4])/(4*b) - (Sqrt[1 + (a + b*x^4)^2]*Log[-a -
b*x^4 + Sqrt[1 + (a + b*x^4)^2]])/(4*b*(a + b*x^4)*Sqrt[1 + (a + b*x^4)^(
-2)])
```

Rubi [A] (warning: unable to verify)

Time = 0.30 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.83, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {7266, 6868, 895, 798, 73, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx$$

$$\downarrow 7266$$

$$\frac{1}{4} \int \operatorname{csch}^{-1}(bx^4 + a) dx^4$$

$$\downarrow 6868$$

$$\frac{1}{4} \left(\int \frac{1}{(bx^4 + a) \sqrt{1 + \frac{1}{(bx^4 + a)^2}}} dx^4 + \frac{(a + bx^4) \operatorname{csch}^{-1}(a + bx^4)}{b} \right)$$

$$\downarrow 895$$

$$\frac{1}{4} \left(\frac{\int \frac{1}{\sqrt{1 + \frac{1}{x^8}} x^4} d(bx^4 + a)}{b} + \frac{(a + bx^4) \operatorname{csch}^{-1}(a + bx^4)}{b} \right)$$

$$\downarrow 798$$

$$\frac{1}{4} \left(\frac{(a + bx^4) \operatorname{csch}^{-1}(a + bx^4)}{b} - \frac{\int \frac{1}{\sqrt{1 + \frac{1}{x^8}} x^4} d\frac{1}{x^8}}{2b} \right)$$

$$\downarrow 73$$

$$\frac{1}{4} \left(\frac{(a + bx^4) \operatorname{csch}^{-1}(a + bx^4)}{b} - \frac{\int \frac{1}{x^8 - 1} d\sqrt{1 + \frac{1}{x^8}}}{b} \right)$$

$$\downarrow 220$$

$$\frac{1}{4} \left(\frac{(a + bx^4) \operatorname{csch}^{-1}(a + bx^4)}{b} + \frac{\operatorname{arctanh}\left(\sqrt{\frac{1}{x^8} + 1}\right)}{b} \right)$$

input `Int[x^3*ArcCsch[a + b*x^4],x]`

output `((a + b*x^4)*ArcCsch[a + b*x^4])/b + ArcTanh[Sqrt[1 + x^(-8)]]/b)/4`

Defintions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-
 1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] &&
 (LtQ[a, 0] || GtQ[b, 0])`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst
 [Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a,
 b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 895 `Int[(u_)^(m_.)*((a_) + (b_.)*(v_)^(n_))^(p_.), x_Symbol] := Simp[u^m/(Coeff
 icient[v, x, 1]*v^m) Subst[Int[x^m*(a + b*x^n)^p, x], x, v], x] /; FreeQ[
 {a, b, m, n, p}, x] && LinearPairQ[u, v, x]`

rule 6868 `Int[ArcCsch[(c_) + (d_.)*(x_)], x_Symbol] := Simp[(c + d*x)*(ArcCsch[c + d*
 x]/d), x] + Int[1/((c + d*x)*Sqrt[1 + 1/(c + d*x)^2]), x] /; FreeQ[{c, d},
 x]`

rule 7266

```
Int[(u_)*(x_)^(m_), x_Symbol] := Simp[1/(m + 1) Subst[Int[SubstFor[x^(m
+ 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && Function
OfQ[x^(m + 1), u, x]
```

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.13

method	result	size
derivativedivides	$\frac{(bx^4+a) \operatorname{arccsch}(bx^4+a) + \ln\left(bx^4+a+(bx^4+a)\sqrt{1+\frac{1}{(bx^4+a)^2}}\right)}{4b}$	52
default	$\frac{(bx^4+a) \operatorname{arccsch}(bx^4+a) + \ln\left(bx^4+a+(bx^4+a)\sqrt{1+\frac{1}{(bx^4+a)^2}}\right)}{4b}$	52

input

```
int(x^3*arccsch(b*x^4+a),x,method=_RETURNVERBOSE)
```

output

```
1/4/b*((b*x^4+a)*arccsch(b*x^4+a)+ln(b*x^4+a+(b*x^4+a)*(1+1/(b*x^4+a)^2)^(
1/2)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 266 vs. 2(40) = 80.

Time = 0.12 (sec) , antiderivative size = 266, normalized size of antiderivative = 5.78

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx$$

$$= \frac{bx^4 \log\left(\frac{(bx^4+a)\sqrt{\frac{b^2x^8+2abx^4+a^2+1}{b^2x^8+2abx^4+a^2}}+1}{bx^4+a}\right) + a \log\left(-bx^4 + (bx^4 + a)\sqrt{\frac{b^2x^8+2abx^4+a^2+1}{b^2x^8+2abx^4+a^2}} - a + 1\right) - a \log(-bx^4 + a)}{4b}$$

input

```
integrate(x^3*arccsch(b*x^4+a),x, algorithm="fricas")
```

output

```
1/4*(b*x^4*log(((b*x^4 + a)*sqrt((b^2*x^8 + 2*a*b*x^4 + a^2 + 1)/(b^2*x^8
+ 2*a*b*x^4 + a^2)) + 1)/(b*x^4 + a)) + a*log(-b*x^4 + (b*x^4 + a)*sqrt((b
^2*x^8 + 2*a*b*x^4 + a^2 + 1)/(b^2*x^8 + 2*a*b*x^4 + a^2)) - a + 1) - a*log
(-b*x^4 + (b*x^4 + a)*sqrt((b^2*x^8 + 2*a*b*x^4 + a^2 + 1)/(b^2*x^8 + 2*a
*b*x^4 + a^2)) - a - 1) - log(-b*x^4 + (b*x^4 + a)*sqrt((b^2*x^8 + 2*a*b*x
^4 + a^2 + 1)/(b^2*x^8 + 2*a*b*x^4 + a^2)) - a))/b
```

Sympy [F(-1)]

Timed out.

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx = \text{Timed out}$$

input

```
integrate(x**3*acsch(b*x**4+a), x)
```

output

Timed out

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.24

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx$$

$$= \frac{2(bx^4 + a) \operatorname{arcsch}(bx^4 + a) + \log\left(\sqrt{\frac{1}{(bx^4+a)^2} + 1} + 1\right) - \log\left(\sqrt{\frac{1}{(bx^4+a)^2} + 1} - 1\right)}{8b}$$

input

```
integrate(x^3*arccsch(b*x^4+a), x, algorithm="maxima")
```

output

```
1/8*(2*(b*x^4 + a)*arccsch(b*x^4 + a) + log(sqrt(1/(b*x^4 + a)^2 + 1) + 1)
- log(sqrt(1/(b*x^4 + a)^2 + 1) - 1))/b
```

Giac [F]

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx = \int x^3 \operatorname{arcsch}(bx^4 + a) dx$$

input `integrate(x^3*arccsch(b*x^4+a),x, algorithm="giac")`

output `integrate(x^3*arccsch(b*x^4 + a), x)`

Mupad [B] (verification not implemented)

Time = 3.96 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.91

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx = \frac{\operatorname{atanh}\left(\sqrt{\frac{1}{(bx^4+a)^2} + 1}\right)}{4b} + \frac{\operatorname{asinh}\left(\frac{1}{bx^4+a}\right) (bx^4 + a)}{4b}$$

input `int(x^3*asinh(1/(a + b*x^4)),x)`

output `atanh((1/(a + b*x^4)^2 + 1)^(1/2))/(4*b) + (asinh(1/(a + b*x^4))*(a + b*x^4))/(4*b)`

Reduce [F]

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx = \int \operatorname{acsch}(bx^4 + a) x^3 dx$$

input `int(x^3*acsch(b*x^4+a),x)`

output `int(acsch(a + b*x**4)*x**3,x)`

3.28 $\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx$

Optimal result	231
Mathematica [B] (verified)	231
Rubi [A] (warning: unable to verify)	232
Maple [F]	234
Fricas [B] (verification not implemented)	234
Sympy [F(-1)]	235
Maxima [A] (verification not implemented)	235
Giac [F]	236
Mupad [B] (verification not implemented)	236
Reduce [F]	236

Optimal result

Integrand size = 14, antiderivative size = 46

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx = \frac{(a + bx^n) \operatorname{csch}^{-1}(a + bx^n)}{bn} + \frac{\operatorname{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx^n)^2}}\right)}{bn}$$

output

```
(a+b*x^n)*arccsch(a+b*x^n)/b/n+arctanh((1+1/(a+b*x^n)^2)^(1/2))/b/n
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 93 vs. 2(46) = 92.

Time = 0.13 (sec) , antiderivative size = 93, normalized size of antiderivative = 2.02

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx = \frac{(a + bx^n)^2 \operatorname{csch}^{-1}(a + bx^n) - \frac{\sqrt{1+(a+bx^n)^2} \log\left(-a-bx^n + \sqrt{1+(a+bx^n)^2}\right)}{\sqrt{1+\frac{1}{(a+bx^n)^2}}}}{bn(a + bx^n)}$$

input

```
Integrate[x^(-1 + n)*ArcCsch[a + b*x^n], x]
```


output

```
((a + b*x^n)^2*ArcCsch[a + b*x^n] - (Sqrt[1 + (a + b*x^n)^2]*Log[-a - b*x^n + Sqrt[1 + (a + b*x^n)^2]])/Sqrt[1 + (a + b*x^n)^(-2)]/(b*n*(a + b*x^n))
```

Rubi [A] (warning: unable to verify)

Time = 0.32 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.87, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {7266, 6868, 895, 798, 73, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{n-1} \operatorname{csch}^{-1}(a + bx^n) dx$$

$$\downarrow \text{7266}$$

$$\int \operatorname{csch}^{-1}(bx^n + a) dx^n$$

$$\downarrow \text{6868}$$

$$\int \frac{1}{(bx^n+a)\sqrt{1+\frac{1}{(bx^n+a)^2}}} dx^n + \frac{(a+bx^n)\operatorname{csch}^{-1}(a+bx^n)}{b}$$

$$\downarrow \text{895}$$

$$\frac{\int \frac{x^{-n}}{\sqrt{x^{-2n+1}}} d(bx^n+a)}{b} + \frac{(a+bx^n)\operatorname{csch}^{-1}(a+bx^n)}{b}$$

$$\downarrow \text{798}$$

$$\frac{(a+bx^n)\operatorname{csch}^{-1}(a+bx^n)}{b} - \frac{\int \frac{x^{-n}}{\sqrt{x^{-2n+1}}} dx^{-2n}}{2b}$$

$$\downarrow \text{73}$$

$$\frac{(a+bx^n)\operatorname{csch}^{-1}(a+bx^n)}{b} - \frac{\int \frac{1}{x^{2n-1}} d\sqrt{x^{-2n+1}}}{b}$$

$$\downarrow \text{220}$$

$$\frac{\frac{(a+bx^n)\operatorname{csch}^{-1}(a+bx^n)}{b} + \frac{\operatorname{arctanh}(\sqrt{x^{-2n}+1})}{b}}{n}$$

input `Int[x^(-1 + n)*ArcCsch[a + b*x^n], x]`

output `((a + b*x^n)*ArcCsch[a + b*x^n])/b + ArcTanh[Sqrt[1 + x^(-2*n)]]/b/n`

Defintions of rubi rules used

rule 73 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]`

rule 220 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-
1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] &&
(LtQ[a, 0] || GtQ[b, 0])`

rule 798 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst
[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a,
b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 895 `Int[(u_)^(m_)*((a_) + (b_)*(v_)^(n_))^(p_), x_Symbol] := Simp[u^m/(Coeff
icient[v, x, 1]*v^m) Subst[Int[x^m*(a + b*x^n)^p, x], x, v], x] /; FreeQ[
{a, b, m, n, p}, x] && LinearPairQ[u, v, x]`

rule 6868 `Int[ArcCsch[(c_) + (d_)*(x_)], x_Symbol] := Simp[(c + d*x)*(ArcCsch[c + d*
x]/d), x] + Int[1/((c + d*x)*Sqrt[1 + 1/(c + d*x)^2]), x] /; FreeQ[{c, d},
x]`

rule 7266

```
Int[(u_)*(x_)^(m_.), x_Symbol] := Simp[1/(m + 1) Subst[Int[SubstFor[x^(m
+ 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && Function
OfQ[x^(m + 1), u, x]
```

Maple [F]

$$\int x^{-1+n} \operatorname{arccsch}(a + b x^n) dx$$

input

```
int(x^(-1+n)*arccsch(a+b*x^n),x)
```

output

```
int(x^(-1+n)*arccsch(a+b*x^n),x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 334 vs. $2(44) = 88$.

Time = 0.12 (sec) , antiderivative size = 334, normalized size of antiderivative = 7.26

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + b x^n) dx$$

$$a \log \left(-b \cosh(n \log(x)) - b \sinh(n \log(x)) - a + \sqrt{\frac{2ab + (a^2 + b^2 + 1) \cosh(n \log(x)) - (a^2 - b^2 + 1) \sinh(n \log(x))}{\cosh(n \log(x)) - \sinh(n \log(x))}} + 1 \right)$$

=

input

```
integrate(x^(-1+n)*arccsch(a+b*x^n),x, algorithm="fricas")
```

output

```
(a*log(-b*cosh(n*log(x)) - b*sinh(n*log(x)) - a + sqrt((2*a*b + (a^2 + b^2 + 1)*cosh(n*log(x)) - (a^2 - b^2 + 1)*sinh(n*log(x))))/(cosh(n*log(x)) - sinh(n*log(x)))) + 1) - a*log(-b*cosh(n*log(x)) - b*sinh(n*log(x)) - a + sqrt((2*a*b + (a^2 + b^2 + 1)*cosh(n*log(x)) - (a^2 - b^2 + 1)*sinh(n*log(x))))/(cosh(n*log(x)) - sinh(n*log(x)))) - 1) + (b*cosh(n*log(x)) + b*sinh(n*log(x)))*log((sqrt((2*a*b + (a^2 + b^2 + 1)*cosh(n*log(x)) - (a^2 - b^2 + 1)*sinh(n*log(x))))/(cosh(n*log(x)) - sinh(n*log(x)))) + 1)/(b*cosh(n*log(x)) + b*sinh(n*log(x)) + a)) - log(-b*cosh(n*log(x)) - b*sinh(n*log(x)) - a + sqrt((2*a*b + (a^2 + b^2 + 1)*cosh(n*log(x)) - (a^2 - b^2 + 1)*sinh(n*log(x))))/(cosh(n*log(x)) - sinh(n*log(x))))))/(b*n)
```

Sympy [F(-1)]

Timed out.

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx = \text{Timed out}$$

input

```
integrate(x**(-1+n)*acsch(a+b*x**n), x)
```

output

Timed out

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.30

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx$$

$$= \frac{2(bx^n + a) \operatorname{arcsch}(bx^n + a) + \log\left(\sqrt{\frac{1}{(bx^n + a)^2} + 1} + 1\right) - \log\left(\sqrt{\frac{1}{(bx^n + a)^2} + 1} - 1\right)}{2bn}$$

input

```
integrate(x^(-1+n)*arccsch(a+b*x^n), x, algorithm="maxima")
```

output

```
1/2*(2*(b*x^n + a)*arccsch(b*x^n + a) + log(sqrt(1/(b*x^n + a)^2 + 1) + 1) - log(sqrt(1/(b*x^n + a)^2 + 1) - 1))/(b*n)
```

Giac [F]

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx = \int x^{n-1} \operatorname{arcsch}(bx^n + a) dx$$

input `integrate(x^(-1+n)*arccsch(a+b*x^n),x, algorithm="giac")`

output `integrate(x^(n - 1)*arccsch(b*x^n + a), x)`

Mupad [B] (verification not implemented)

Time = 3.51 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.87

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx = \frac{\operatorname{atanh}\left(\sqrt{\frac{1}{(a+bx^n)^2} + 1}\right) + \operatorname{asinh}\left(\frac{1}{a+bx^n}\right) (a + bx^n)}{bn}$$

input `int(x^(n - 1)*asinh(1/(a + b*x^n)),x)`

output `(atanh((1/(a + b*x^n)^2 + 1)^(1/2)) + asinh(1/(a + b*x^n))*(a + b*x^n))/(b*n)`

Reduce [F]

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx = \int \frac{x^n \operatorname{acsch}(x^n b + a)}{x} dx$$

input `int(x^(-1+n)*acsch(a+b*x^n),x)`

output `int((x**n*acsch(x**n*b + a))/x,x)`

CHAPTER 4

APPENDIX

4.1	Listing of Grading functions	237
4.2	Links to plain text integration problems used in this report for each CAS .	255

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
          finalresult={"C","Result contains complex when optimal does not."}
        ]
      ,(*ELSE*)(*result does not contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
          ]
        ]
      ,(*ELSE*)(*expnResult>expnOptimal*)
        If[FreeQ[result,Integrate] && FreeQ[result,Int],
          finalresult={"C","Result contains higher order function than in optimal. Order "
          ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]==Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]==Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]==Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]==Plus || Head[expn]==Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```



```

    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 5]],
    If [AppellFunctionQ [Head [expn]],
        Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 6]],
    If [Head [expn] === RootSum,
        Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 7]],
    If [Head [expn] === Integrate || Head [expn] === Int,
        Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 8]],
    9]]]]]]]]]]]

```

```

ElementaryFunctionQ [func_] :=
  MemberQ [{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

```

```

SpecialFunctionQ [func_] :=
  MemberQ [{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

```

```

HypergeometricFunctionQ [func_] :=
  MemberQ [{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```

AppellFunctionQ [func_] :=
  MemberQ [{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022  add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result   := ExpnType(result);
      ExpnType_optimal  := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#     is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```



```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file