


A Solution Manual For

**Advanced Mathematical Methods  
for Scientists and Engineers,  
Bender and Orszag. Springer  
October 29, 1999**



**Nasser M. Abbasi**

October 12, 2023

# Contents

1	Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS. page 136	2
---	--	---

# 1 Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS. page 136

1.1	problem 3.5 . . . . .	3
1.2	problem 3.6 (a) . . . . .	4
1.3	problem 3.6 (b) . . . . .	5
1.4	problem 3.6 (c) . . . . .	6
1.5	problem 3.6 (d) . . . . .	7
1.6	problem 3.24 (a) . . . . .	8
1.7	problem 3.24 (b) . . . . .	9
1.8	problem 3.24 (c) . . . . .	10
1.9	problem 3.24 (d) . . . . .	11
1.10	problem 3.24 (e) . . . . .	12
1.11	problem 3.24 (f) . . . . .	13
1.12	problem 3.24 (g) . . . . .	14
1.13	problem 3.24 (h) . . . . .	15
1.14	problem 3.24 (i) . . . . .	16
1.15	problem 3.25 $v=1/2$ . . . . .	17
1.16	problem 3.25 $v=3/2$ . . . . .	18
1.17	problem 3.25 $v=5/2$ . . . . .	19
1.18	problem 3.26 . . . . .	20
1.19	problem 3.48 (a) . . . . .	21
1.20	problem 3.48 (b) . . . . .	22
1.21	problem 3.48 (c) . . . . .	23
1.22	problem 3.48 (d) . . . . .	24
1.23	problem 3.50 . . . . .	26

## 1.1 problem 3.5

Internal problem ID [4726]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.5.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _exact, _linear, _homogeneous]`

$$(x - 1)(x - 2)y'' + (4x - 6)y' + 2y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = 1]$$

With the expansion point for the power series method at  $x = 0$ .

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

Order:=6;

`dsolve([(x-1)*(x-2)*diff(y(x),x$2)+(4*x-6)*diff(y(x),x)+2*y(x)=0,y(0) = 2, D(y)(0) = 1],y(x),`

$$y(x) = 2 + x + \frac{1}{2}x^2 + \frac{1}{4}x^3 + \frac{1}{8}x^4 + \frac{1}{16}x^5 + O(x^6)$$

### ✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 34

`AsymptoticDSolveValue[{(x-1)*(x-2)*y'[x]+(4*x-6)*y'[x]+2*y[x]==0,{y[0]==2,y'[0]==1}},y[x],{x`

$$y(x) \rightarrow \frac{x^5}{16} + \frac{x^4}{8} + \frac{x^3}{4} + \frac{x^2}{2} + x + 2$$

## 1.2 problem 3.6 (a)

Internal problem ID [4727]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.6 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`

$$y'' - 2y'x + 8y = 0$$

With initial conditions

$$[y(0) = 4, y'(0) = 0]$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;
dsolve([diff(y(x),x$2)-2*x*diff(y(x),x)+8*y(x)=0,y(0) = 4, D(y)(0) = 0],y(x),type='series',x=
```

$$y(x) = 4 - 16x^2 + \frac{16}{3}x^4 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 17

```
AsymptoticDSolveValue[{y'[x]-2*x*y'[x]+8*y[x]==0,{y[0]==4,y'[0]==0}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{16x^4}{3} - 16x^2 + 4$$

### 1.3 problem 3.6 (b)

Internal problem ID [4728]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.6 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`

$$y'' - 2y'x + 8y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 4]$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;
dsolve([diff(y(x),x$2)-2*x*diff(y(x),x)+8*y(x)=0,y(0) = 0, D(y)(0) = 4],y(x),type='series',x=
```

$$y(x) = 4x - 4x^3 + \frac{2}{5}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 19

```
AsymptoticDSolveValue[{y'[x]-2*x*y'[x]+8*y[x]==0,{y[0]==0,y'[0]==4}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{2x^5}{5} - 4x^3 + 4x$$

## 1.4 problem 3.6 (c)

Internal problem ID [4729]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.6 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [\_Gegenbauer]

$$(-x^2 + 1)y'' - 2y'x + 12y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 3]$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
Order:=6;
dsolve([(1-x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+12*y(x)=0,y(0) = 0, D(y)(0) = 3],y(x),type='s
```

$$y(x) = -5x^3 + 3x$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 12

```
AsymptoticDSolveValue[{{(1-x^2)*y'[x]-2*x*y'[x]+12*y[x]==0,{y[0]==0,y'[0]==3}},y[x],{x,0,5}]
```

$$y(x) \rightarrow 3x - 5x^3$$

## 1.5 problem 3.6 (d)

Internal problem ID [4730]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.6 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - (x - 1)y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

With the expansion point for the power series method at  $x = 0$ .

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
Order:=6;
dsolve([diff(y(x),x$2)=(x-1)*y(x),y(0) = 1, D(y)(0) = 0],y(x),type='series',x=0);
```

$$y(x) = 1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{30}x^5 + O(x^6)$$

### ✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 33

```
AsymptoticDSolveValue[{y'[x]==(x-1)*y[x],{y[0]==1,y'[0]==0}},y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{x^5}{30} + \frac{x^4}{24} + \frac{x^3}{6} - \frac{x^2}{2} + 1$$



## 1.6 problem 3.24 (a)

Internal problem ID [4731]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.24 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x(2+x)y'' + 2y'(x+1) - 2y = 0$$

With the expansion point for the power series method at  $x = 0$ .

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 43

```
Order:=6;
dsolve(x*(x+2)*diff(y(x),x$2)+2*(x+1)*diff(y(x),x)-2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (\ln(x) c_2 + c_1) (1 + x + O(x^6)) + \left( -\frac{5}{2}x - \frac{3}{8}x^2 + \frac{1}{12}x^3 - \frac{5}{192}x^4 + \frac{3}{320}x^5 + O(x^6) \right) c_2$$

### ✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 53

```
AsymptoticDSolveValue[x*(x+2)*y'[x]+2*(x+1)*y'[x]-2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{3x^5}{320} - \frac{5x^4}{192} + \frac{x^3}{12} - \frac{3x^2}{8} - \frac{5x}{2} + (x+1) \log(x) \right) + c_1(x+1)$$

## 1.7 problem 3.24 (b)

Internal problem ID [4732]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.24 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [[\_Emden, \_Fowler]]

$$xy'' + y = 0$$

With the expansion point for the power series method at  $x = 0$ .

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 58

```
Order:=6;
dsolve(x*diff(y(x),x$2)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x \left( 1 - \frac{1}{2}x + \frac{1}{12}x^2 - \frac{1}{144}x^3 + \frac{1}{2880}x^4 - \frac{1}{86400}x^5 + O(x^6) \right) \\ + c_2 \left( \ln(x) \left( -x + \frac{1}{2}x^2 - \frac{1}{12}x^3 + \frac{1}{144}x^4 - \frac{1}{2880}x^5 + O(x^6) \right) \right. \\ \left. + \left( 1 - \frac{3}{4}x^2 + \frac{7}{36}x^3 - \frac{35}{1728}x^4 + \frac{101}{86400}x^5 + O(x^6) \right) \right)$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 85

```
AsymptoticDSolveValue[x*y''[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{1}{144}x(x^3 - 12x^2 + 72x - 144) \log(x) + \frac{-47x^4 + 480x^3 - 2160x^2 + 1728x + 1728}{1728} \right) \\ + c_2 \left( \frac{x^5}{2880} - \frac{x^4}{144} + \frac{x^3}{12} - \frac{x^2}{2} + x \right)$$

## 1.8 problem 3.24 (c)

Internal problem ID [4733]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.24 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + (e^x - 1)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 39

```
Order:=6;
dsolve(diff(y(x),x$2)+(exp(x)-1)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{120}x^5\right) y(0) + \left(x - \frac{1}{12}x^4 - \frac{1}{40}x^5\right) D(y)(0) + O(x^6)$$

### ✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 49

```
AsymptoticDSolveValue[y''[x]+(Exp[x]-1)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( -\frac{x^5}{40} - \frac{x^4}{12} + x \right) + c_1 \left( -\frac{x^5}{120} - \frac{x^4}{24} - \frac{x^3}{6} + 1 \right)$$

## 1.9 problem 3.24 (d)

Internal problem ID [4734]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.24 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x(1-x)y'' - 3y'x - y = 0$$

With the expansion point for the power series method at  $x = 0$ .

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 60

```
Order:=6;
dsolve(x*(1-x)*diff(y(x),x$2)-3*x*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1x(1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + O(x^6)) \\ & + (x + 2x^2 + 3x^3 + 4x^4 + 5x^5 + O(x^6)) \ln(x) c_2 \\ & + (1 + 3x + 5x^2 + 7x^3 + 9x^4 + 11x^5 + O(x^6)) c_2 \end{aligned}$$

### ✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 63

```
AsymptoticDSolveValue[x*(1-x)*y''[x]-3*x*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(x^4 + x^3 + x^2 + (4x^3 + 3x^2 + 2x + 1)x \log(x) + x + 1) + c_2(5x^5 + 4x^4 + 3x^3 + 2x^2 + x)$$

## 1.10 problem 3.24 (e)

Internal problem ID [4735]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.24 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F(`

$$2xy'' - y' + x^2y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 28

```
Order:=6;
dsolve(2*x*diff(y(x),x$2)-diff(y(x),x)+x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{\frac{3}{2}} \left( 1 - \frac{1}{27} x^3 + O(x^6) \right) + c_2 \left( 1 - \frac{1}{9} x^3 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 33

```
AsymptoticDSolveValue[2*x*y''[x]-y'[x]+x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( 1 - \frac{x^3}{9} \right) + c_1 \left( 1 - \frac{x^3}{27} \right) x^{3/2}$$

## 1.11 problem 3.24 (f)

Internal problem ID [4736]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.24 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$\sin(x) y'' - 2y' \cos(x) - y \sin(x) = 0$$

With the expansion point for the power series method at  $x = 0$ .

### ✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 32

```
Order:=6;
dsolve(sin(x)*diff(y(x),x$2)-2*cos(x)*diff(y(x),x)-sin(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^3 \left( 1 - \frac{1}{10} x^2 + \frac{1}{280} x^4 + O(x^6) \right) + c_2 \left( 12 - 6x^2 + \frac{1}{2} x^4 + O(x^6) \right)$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 44

```
AsymptoticDSolveValue[Sin[x]*y''[x]-2*Cos[x]*y'[x]-Sin[x]*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{x^4}{24} - \frac{x^2}{2} + 1 \right) + c_2 \left( \frac{x^7}{280} - \frac{x^5}{10} + x^3 \right)$$

## 1.12 problem 3.24 (g)

Internal problem ID [4737]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.24 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [[\_Emden, \_Fowler]]

$$y'' - x^2 y = 0$$

With the expansion point for the power series method at  $x = 0$ .

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;
dsolve(diff(y(x),x$2)-x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{x^4}{12}\right) y(0) + \left(x + \frac{1}{20}x^5\right) D(y)(0) + O(x^6)$$

### ✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y''[x]-x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{x^5}{20} + x \right) + c_1 \left( \frac{x^4}{12} + 1 \right)$$

### 1.13 problem 3.24 (h)

Internal problem ID [4738]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.24 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`, `[_2nd_order, _linear]`, ‘

$$x(2+x)y'' + y'(x+1) - 4y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 38

```
Order:=6;
dsolve(x*(x+2)*diff(y(x),x$2)+(x+1)*diff(y(x),x)-4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1\sqrt{x} \left( 1 + \frac{5}{4}x + \frac{7}{32}x^2 - \frac{3}{128}x^3 + \frac{11}{2048}x^4 - \frac{13}{8192}x^5 + O(x^6) \right) + c_2(1 + 4x + 2x^2 + O(x^6))$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 60

```
AsymptoticDSolveValue[x*(x+2)*y''[x]+(x+1)*y'[x]-4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2(2x^2 + 4x + 1) + c_1\sqrt{x} \left( -\frac{13x^5}{8192} + \frac{11x^4}{2048} - \frac{3x^3}{128} + \frac{7x^2}{32} + \frac{5x}{4} + 1 \right)$$



## 1.14 problem 3.24 (i)

Internal problem ID [4739]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.24 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$xy'' + \left(\frac{1}{2} - x\right)y' - y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```
Order:=6;
dsolve(x*diff(y(x),x$2)+(1/2-x)*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1\sqrt{x} \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + O(x^6)\right) \\ + c_2 \left(1 + 2x + \frac{4}{3}x^2 + \frac{8}{15}x^3 + \frac{16}{105}x^4 + \frac{32}{945}x^5 + O(x^6)\right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 79

```
AsymptoticDSolveValue[x*y''[x]+(1/2-x)*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1\sqrt{x} \left(\frac{x^5}{120} + \frac{x^4}{24} + \frac{x^3}{6} + \frac{x^2}{2} + x + 1\right) + c_2 \left(\frac{32x^5}{945} + \frac{16x^4}{105} + \frac{8x^3}{15} + \frac{4x^2}{3} + 2x + 1\right)$$

## 1.15 problem 3.25 $v=1/2$

Internal problem ID [4740]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.25  $v=1/2$ .

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y' x + \left(x^2 + \frac{1}{4}\right) y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 45

```
Order:=6;
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2+(1/2)^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{-\frac{i}{2}} \left( 1 + \left( -\frac{1}{5} - \frac{i}{10} \right) x^2 + \left( \frac{7}{680} + \frac{3i}{340} \right) x^4 + O(x^6) \right) \\ + c_2 x^{\frac{i}{2}} \left( 1 + \left( -\frac{1}{5} + \frac{i}{10} \right) x^2 + \left( \frac{7}{680} - \frac{3i}{340} \right) x^4 + O(x^6) \right)$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
AsymptoticDSolveValue[x^2*y''+x*y'[x]+(x^2+(1/2)^2)*y[x]==0,y[x],{x,0,5}]
```

Timed out

## 1.16 problem 3.25 $v=3/2$

Internal problem ID [4741]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.25  $v=3/2$ .

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y' x + \left(x^2 + \frac{9}{4}\right) y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 45

```
Order:=6;
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2+(3/2)^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{-\frac{3i}{2}} \left( 1 + \left( -\frac{1}{13} - \frac{3i}{26} \right) x^2 + \left( -\frac{1}{2600} + \frac{9i}{1300} \right) x^4 + O(x^6) \right) \\ + c_2 x^{\frac{3i}{2}} \left( 1 + \left( -\frac{1}{13} + \frac{3i}{26} \right) x^2 + \left( -\frac{1}{2600} - \frac{9i}{1300} \right) x^4 + O(x^6) \right)$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
AsymptoticDSolveValue[x^2*y''+x*y'[x]+(x^2+(3/2)^2)*y[x]==0,y[x],{x,0,5}]
```

Timed out

## 1.17 problem 3.25 $v=5/2$

Internal problem ID [4742]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.25  $v=5/2$ .

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y' x + \left(x^2 + \frac{25}{4}\right) y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

```
Order:=6;
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2+(5/2)^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{-\frac{5i}{2}} \left( 1 + \left( -\frac{1}{29} - \frac{5i}{58} \right) x^2 + \left( -\frac{17}{9512} + \frac{15i}{4756} \right) x^4 + O(x^6) \right) \\ + c_2 x^{\frac{5i}{2}} \left( 1 + \left( -\frac{1}{29} + \frac{5i}{58} \right) x^2 + \left( -\frac{17}{9512} - \frac{15i}{4756} \right) x^4 + O(x^6) \right)$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
AsymptoticDSolveValue[x^2*y''+x*y'[x]+(x^2+(5/2)^2)*y[x]==0,y[x],{x,0,5}]
```

Timed out

## 1.18 problem 3.26

Internal problem ID [4743]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.26.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x - 1)y'' - y'x + y = 0$$

With the expansion point for the power series method at  $x = 0$ .

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;
dsolve((x-1)*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5\right) y(0) + D(y)(0)x + O(x^6)$$

### ✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 41

```
AsymptoticDSolveValue[(x-1)*y''[x]-x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{x^5}{120} + \frac{x^4}{24} + \frac{x^3}{6} + \frac{x^2}{2} + 1 \right) + c_2 x$$

## 1.19 problem 3.48 (a)

Internal problem ID [4744]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.48 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_linear]`

$$y' + xy - \cos(x) = 0$$

With the expansion point for the power series method at  $x = 0$ .

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
Order:=6;
dsolve(diff(y(x),x)+x*y(x)=cos(x),y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4\right) y(0) + x - \frac{x^3}{2} + \frac{13x^5}{120} + O(x^6)$$

### ✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 38

```
AsymptoticDSolveValue[y'[x]+x*y[x]==Cos[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{13x^5}{120} - \frac{x^3}{2} + c_1 \left( \frac{x^4}{8} - \frac{x^2}{2} + 1 \right) + x$$

## 1.20 problem 3.48 (b)

Internal problem ID [4745]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.48 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_linear]

$$y' + xy - \frac{1}{x^3} = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(x),x)+x*y(x)=1/x^3,y(x), singsol=all)
```

$$y(x) = \left( -\frac{e^{\frac{x^2}{2}}}{2x^2} - \frac{\text{Ei}_1\left(-\frac{x^2}{2}\right)}{4} + c_1 \right) e^{-\frac{x^2}{2}}$$

### ✓ Solution by Mathematica

Time used: 0.061 (sec). Leaf size: 38

```
DSolve[y'[x]+x*y[x]==1/x^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4} \left( -\frac{2}{x^2} + e^{-\frac{x^2}{2}} \left( \text{ExpIntegralEi}\left(\frac{x^2}{2}\right) + 4c_1 \right) \right)$$

## 1.21 problem 3.48 (c)

Internal problem ID [4746]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.48 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x^3 y'' + y - \frac{1}{x^4} = 0$$

With the expansion point for the power series method at  $x = 0$ .

✗ Solution by Maple

```
Order:=6;
dsolve(x^3*diff(y(x),x$2)+y(x)=1/x^4,y(x),type='series',x=0);
```

No solution found

✓ Solution by Mathematica

Time used: 0.361 (sec). Leaf size: 800

```
AsymptoticDSolveValue[x^3*y''[x]+y[x]==1/x^4,y[x],{x,0,5}]
```

$$y(x) \rightarrow e^{-\frac{2i}{\sqrt{x}}} x^{3/4} \left( \frac{33424574007825x^5}{281474976710656} - \frac{468131288625ix^{9/2}}{8796093022208} - \frac{14783093325x^4}{549755813888} \right. \\ \left. + \frac{66891825ix^{7/2}}{4294967296} + \frac{2837835x^3}{268435456} - \frac{72765ix^{5/2}}{8388608} - \frac{4725x^2}{524288} + \frac{105ix^{3/2}}{8192} + \frac{15x}{512} - \frac{3i\sqrt{x}}{16} \right. \\ \left. + 1 \right) c_1 + e^{\frac{2i}{\sqrt{x}}} x^{3/4} \left( \frac{33424574007825x^5}{281474976710656} + \frac{468131288625ix^{9/2}}{8796093022208} - \frac{14783093325x^4}{549755813888} - \frac{66891825ix^{7/2}}{4294967296} + \frac{2837835x^3}{268435456} \right.$$



## 1.22 problem 3.48 (d)

Internal problem ID [4747]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.48 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$xy'' - 2y' + y - \cos(x) = 0$$

With the expansion point for the power series method at  $x = 0$ .

**X** Solution by Maple

```
Order:=6;
dsolve(x*diff(y(x),x$2)-2*diff(y(x),x)+y(x)=cos(x),y(x),type='series',x=0);
```

No solution found

✓ Solution by Mathematica

Time used: 0.144 (sec). Leaf size: 312

AsymptoticDSolveValue[x\*y'[x]-2\*y'[x]+y[x]==Cos[x],y[x],{x,0,5}]

$$\begin{aligned}
y(x) \rightarrow & c_1 \left( x^4 \left( \frac{\log(x)}{48} - \frac{5}{192} \right) - \frac{1}{12} x^3 \log(x) + \frac{x^2}{4} + \frac{x}{2} + 1 \right) \\
& + c_2 \left( -\frac{x^5}{806400} + \frac{x^4}{20160} - \frac{x^3}{720} + \frac{x^2}{40} - \frac{x}{4} + 1 \right) x^3 + \left( -\frac{x^5}{806400} + \frac{x^4}{20160} - \frac{x^3}{720} \right. \\
& \quad \left. + \frac{x^2}{40} - \frac{x}{4} + 1 \right) x^3 \left( \frac{x^6 (-20160 \log^2(x) + 141222 \log(x) - 201569)}{3135283200} \right. \\
& \quad + \frac{x^5 (22277 - 114360 \log(x))}{435456000} + \frac{x^4 (69541 - 29064 \log(x))}{34836480} + \frac{x^3 (1860 \log(x) + 193)}{388800} \\
& \quad \left. - \frac{1}{6x^2} + \frac{x^2 (4 \log(x) - 23)}{1152} - \frac{1}{6x} + \frac{1}{36} x (-\log(x) - 2) - \frac{\log(x)}{12} \right) \\
& + \left( \frac{x^6 (5791 - 672 \log(x))}{8709120} - \frac{589x^5}{302400} - \frac{89x^4}{8640} + \frac{19x^3}{360} + \frac{x^2}{24} - \frac{x}{3} \right) \left( x^4 \left( \frac{\log(x)}{48} - \frac{5}{192} \right) \right. \\
& \quad \left. - \frac{1}{12} x^3 \log(x) + \frac{x^2}{4} + \frac{x}{2} + 1 \right)
\end{aligned}$$

## 1.23 problem 3.50

Internal problem ID [4748]

**Book:** Advanced Mathematical Methods for Scientists and Engineers, Bender and Orszag. Springer  
October 29, 1999

**Section:** Chapter 3. APPROXIMATE SOLUTION OF LINEAR DIFFERENTIAL EQUATIONS.  
page 136

**Problem number:** 3.50.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_linear]

$$y' - \frac{y}{x} - \cos(x) = 0$$

With the expansion point for the power series method at  $x = 0$ .

✗ Solution by Maple

```
Order:=6;
dsolve(diff(y(x),x)-y(x)/x=cos(x),y(x),type='series',x=0);
```

No solution found

✓ Solution by Mathematica

Time used: 0.054 (sec). Leaf size: 34

```
AsymptoticDSolveValue[y'[x]-y[x]/x==Cos[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow x \left( -\frac{x^6}{4320} + \frac{x^4}{96} - \frac{x^2}{4} + \log(x) \right) + c_1 x$$