


A Solution Manual For

Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961



Nasser M. Abbasi

October 12, 2023

Contents

1 Exercis 2, page 5

2

1 Exercis 2, page 5

1.1	problem 2(a)	3
1.2	problem 2(b)	4
1.3	problem 2(c)	5
1.4	problem 2(d)	6
1.5	problem 2(e)	7
1.6	problem 2(a)	8
1.7	problem 3(a)	9
1.8	problem 3(b)	10
1.9	problem 3(c)	11
1.10	problem 3(d)	12
1.11	problem 3(e)	13
1.12	problem 3(f)	14
1.13	problem 3(g)	15
1.14	problem 3(h)	16
1.15	problem 3(i)	17

1.1 problem 2(a)

Internal problem ID [1923]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - 2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)=2,y(x), singsol=all)
```

$$y(x) = 2x + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 11

```
DSolve[y'[x]==2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2x + c_1$$

1.2 problem 2(b)

Internal problem ID [1924]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - 2e^{3x} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=2*exp(3*x),y(x), singsol=all)
```

$$y(x) = \frac{2e^{3x}}{3} + c_1$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 17

```
DSolve[y'[x]==2*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2e^{3x}}{3} + c_1$$

1.3 problem 2(c)

Internal problem ID [1925]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - \frac{2}{\sqrt{1-x^2}} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)=2/sqrt(1-x^2),y(x), singsol=all)
```

$$y(x) = 2 \arcsin(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 26

```
DSolve[y'[x]==2/Sqrt[1-x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -4 \cot^{-1} \left(\frac{x+1}{\sqrt{1-x^2}} \right) + c_1$$

1.4 problem 2(d)

Internal problem ID [1926]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(d).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - e^{x^2} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x)=exp(x^2),y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{\pi} \operatorname{erfi}(x)}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 16

```
DSolve[y'[x]==Exp[x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x^2} \operatorname{DawsonF}(x) + c_1$$

1.5 problem 2(e)

Internal problem ID [1927]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(e).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - x e^{x^2} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=x*exp(x^2),y(x), singsol=all)
```

$$y(x) = \frac{e^{x^2}}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 17

```
DSolve[y'[x]==x*Exp[x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{x^2}}{2} + c_1$$

1.6 problem 2(a)

Internal problem ID [1928]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - \arcsin(x) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)=arcsin(x),y(x), singsol=all)
```

$$y(x) = x \arcsin(x) + \sqrt{-x^2 + 1} + c_1$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 23

```
DSolve[y'[x]==ArcSin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \arcsin(x) + \sqrt{1 - x^2} + c_1$$

1.7 problem 3(a)

Internal problem ID [1929]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$-yx + y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=y(x)*x,y(x), singsol=all)
```

$$y(x) = e^{\frac{x^2}{2}} c_1$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 22

```
DSolve[y'[x]==y[x]*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{\frac{x^2}{2}}$$

$$y(x) \rightarrow 0$$

1.8 problem 3(b)

Internal problem ID [1930]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' - x^2 y^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)=y(x)^2*x^2,y(x), singsol=all)
```

$$y(x) = \frac{3}{-x^3 + 3c_1}$$

✓ Solution by Mathematica

Time used: 0.109 (sec). Leaf size: 22

```
DSolve[y'[x]==y[x]^2*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{3}{x^3 + 3c_1}$$

$$y(x) \rightarrow 0$$

1.9 problem 3(c)

Internal problem ID [1931]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' + e^y x = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)=-x*exp(y(x)),y(x), singsol=all)
```

$$y(x) = \ln\left(\frac{2}{x^2 + 2c_1}\right)$$

✓ Solution by Mathematica

Time used: 0.287 (sec). Leaf size: 19

```
DSolve[y'[x]==-x*Exp[y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(2) - \log(x^2 - 2c_1)$$

1.10 problem 3(d)

Internal problem ID [1932]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(d).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' \sin(y) - x^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)*sin(y(x))=x^2,y(x), singsol=all)
```

$$y(x) = \pi - \arccos\left(\frac{x^3}{3} + c_1\right)$$

✓ Solution by Mathematica

Time used: 0.448 (sec). Leaf size: 37

```
DSolve[y'[x]*Sin[y[x]]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\arccos\left(-\frac{x^3}{3} - c_1\right)$$

$$y(x) \rightarrow \arccos\left(-\frac{x^3}{3} - c_1\right)$$

1.11 problem 3(e)

Internal problem ID [1933]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(e).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y'x - \sqrt{1 - y^2} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)*x=sqrt(1-y(x)^2),y(x), singsol=all)
```

$$y(x) = \sin(\ln(x) + c_1)$$

✓ Solution by Mathematica

Time used: 0.198 (sec). Leaf size: 29

```
DSolve[y'[x]*x==Sqrt[1-y[x]^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \cos(\log(x) + c_1)$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 1$$

$$y(x) \rightarrow \text{Interval}[\{-1, 1\}]$$

1.12 problem 3(f)

Internal problem ID [1934]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(f).

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$y'^2 - y^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve((diff(y(x),x))^2-y(x)^2=0,y(x), singsol=all)
```

$$y(x) = c_1 e^x$$

$$y(x) = e^{-x} c_1$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 28

```
DSolve[(y'[x])^2-y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-x}$$

$$y(x) \rightarrow c_1 e^x$$

$$y(x) \rightarrow 0$$

1.13 problem 3(g)

Internal problem ID [1935]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(g).

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$y'^2 - 3y' + 2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)^2-3*diff(y(x),x)+2=0,y(x), singsol=all)
```

$$y(x) = 2x + c_1$$

$$y(x) = c_1 + x$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 19

```
DSolve[(y'[x])^2-3*y'[x]+2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x + c_1$$

$$y(x) \rightarrow 2x + c_1$$

1.14 problem 3(h)

Internal problem ID [1936]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(h).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$(x^2 + 1)y' - 1 = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 8

```
dsolve((1+x^2)*diff(y(x),x)=1,y(x), singsol=all)
```

$$y(x) = \arctan(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 10

```
DSolve[(1+x^2)*y'[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arctan(x) + c_1$$

1.15 problem 3(i)

Internal problem ID [1937]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(i).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$\sin(x)y' - 1 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)*sin(x)=1,y(x), singsol=all)
```

$$y(x) = -\ln(\csc(x) + \cot(x)) + c_1$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 24

```
DSolve[y'[x]*Sin[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log\left(\sin\left(\frac{x}{2}\right)\right) - \log\left(\cos\left(\frac{x}{2}\right)\right) + c_1$$