


A Solution Manual For

**Elementary differential equations.**

**By Earl D. Rainville, Phillip E.  
Bedient. Macmillan Publishing  
Co. NY. 6th edition. 1981.**



**Nasser M. Abbasi**

October 12, 2023

# Contents

1	CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309	2
2	CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314	25
3	CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320	43
4	CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324	81

**1 CHAPTER 16. Nonlinear equations. Section 94.**  
**Factoring the left member. EXERCISES Page 309**

1.1	problem 1	3
1.2	problem 2	4
1.3	problem 3	5
1.4	problem 4	6
1.5	problem 5	7
1.6	problem 6	8
1.7	problem 7	9
1.8	problem 8	10
1.9	problem 9	11
1.10	problem 10	12
1.11	problem 11	13
1.12	problem 12	14
1.13	problem 13	15
1.14	problem 14	16
1.15	problem 15	17
1.16	problem 16	19
1.17	problem 17	21
1.18	problem 18	23
1.19	problem 19	24

## 1.1 problem 1

Internal problem ID [6013]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 1.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type [\_separable]

$$x^2 y'^2 - y^2 = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x)^2-y(x)^2=0,y(x), singsol=all)
```

$$y(x) = c_1 x$$

$$y(x) = \frac{c_1}{x}$$

### ✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 24

```
DSolve[x^2*(y'[x])^2-y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{x}$$

$$y(x) \rightarrow c_1 x$$

$$y(x) \rightarrow 0$$

## 1.2 problem 2

Internal problem ID [6014]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 2.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type [\_quadrature]

$$xy'^2 - (3y + 2x)y' + 6y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x*diff(y(x),x)^2-(2*x+3*y(x))*diff(y(x),x)+6*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1x^3$$

$$y(x) = 2x + c_1$$

### ✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 26

```
DSolve[x*(y'[x])^2-(2*x+3*y[x])*y'[x]+6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x^3$$

$$y(x) \rightarrow 2x + c_1$$

$$y(x) \rightarrow 0$$

### 1.3 problem 3

Internal problem ID [6015]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 3.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type [\_separable]

$$x^2y'^2 - 5xyy' + 6y^2 = 0$$

#### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve(x^2*diff(y(x),x)^2-5*x*y(x)*diff(y(x),x)+6*y(x)^2=0,y(x), singsol=all)
```

$$y(x) = c_1x^3$$

$$y(x) = c_1x^2$$

#### ✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 26

```
DSolve[x^2*(y'[x])^2-5*x*y[x]*y'[x]+6*y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x^2$$

$$y(x) \rightarrow c_1x^3$$

$$y(x) \rightarrow 0$$

## 1.4 problem 4

Internal problem ID [6016]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 4.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type [\_separable]

$$x^2y'^2 + y'x - y^2 - y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(x^2*diff(y(x),x)^2+x*diff(y(x),x)-y(x)^2-y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1x$$

$$y(x) = \frac{-x + c_1}{x}$$

### ✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 31

```
DSolve[x^2*(y'[x])^2+x*y'[x]-y[x]^2-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x$$

$$y(x) \rightarrow -1 + \frac{c_1}{x}$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 0$$

## 1.5 problem 5

Internal problem ID [6017]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 5.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type [\_quadrature]

$$xy'^2 + (1 - x^2y) y' - xy = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(x*diff(y(x),x)^2+(1-x^2*y(x))*diff(y(x),x)-x*y(x)=0,y(x), singsol=all)
```

$$y(x) = -\ln(x) + c_1$$

$$y(x) = e^{\frac{x^2}{2}} c_1$$

### ✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 28

```
DSolve[x*(y'[x])^2+(1-x^2*y[x])*y'[x]-x*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{\frac{x^2}{2}}$$

$$y(x) \rightarrow -\log(x) + c_1$$



## 1.6 problem 6

Internal problem ID [6018]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 6.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type [\_quadrature]

$$y'^2 - (x^2y + 3)y' + 3x^2y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x)^2-(x^2*y(x)+3)*diff(y(x),x)+3*x^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{x^3}{3}}$$

$$y(x) = 3x + c_1$$

### ✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 27

```
DSolve[(y'[x])^2-(x^2*y[x]+3)*y'[x]+3*x^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{\frac{x^3}{3}}$$

$$y(x) \rightarrow 3x + c_1$$

## 1.7 problem 7

Internal problem ID [6019]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 7.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type [\_quadrature]

$$xy'^2 - (xy + 1)y' + y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(x*diff(y(x),x)^2-(1+x*y(x))*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \ln(x) + c_1$$

$$y(x) = e^x c_1$$

### ✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 20

```
DSolve[x*(y'[x])^2-(1+x*y[x])*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x$$

$$y(x) \rightarrow \log(x) + c_1$$

## 1.8 problem 8

Internal problem ID [6020]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 8.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type [\_separable]

$$y'^2 - x^2 y^2 = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x)^2-x^2*y(x)^2=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{x^2}{2}} c_1$$

$$y(x) = e^{-\frac{x^2}{2}} c_1$$

### ✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 38

```
DSolve[(y'[x])^2-x^2*y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-\frac{x^2}{2}}$$

$$y(x) \rightarrow c_1 e^{\frac{x^2}{2}}$$

$$y(x) \rightarrow 0$$

## 1.9 problem 9

Internal problem ID [6021]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 9.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cla`

$$(y + x)^2 y' - y^2 = 0$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 48

```
dsolve((x+y(x))^2*diff(y(x),x)^2=y(x)^2,y(x), singsol=all)
```

$$y(x) = e^{\text{LambertW}(x e^{c_1}) - c_1}$$

$$y(x) = -x - \sqrt{x^2 + 2c_1}$$

$$y(x) = -x + \sqrt{x^2 + 2c_1}$$

### ✓ Solution by Mathematica

Time used: 3.909 (sec). Leaf size: 101

```
DSolve[(x+y[x])^2*(y'[x])^2==y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x - \sqrt{x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -x + \sqrt{x^2 + e^{2c_1}}$$

$$y(x) \rightarrow \frac{x}{W(e^{-c_1}x)}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -\sqrt{x^2} - x$$

$$y(x) \rightarrow \sqrt{x^2} - x$$

## 1.10 problem 10

Internal problem ID [6022]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 10.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type [\_quadrature]

$$yy'^2 + (x - y^2)y' - xy = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve(y(x)*diff(y(x),x)^2+(x-y(x)^2)*diff(y(x),x)-x*y(x)=0,y(x), singsol=all)
```

$$y(x) = \sqrt{-x^2 + c_1}$$

$$y(x) = -\sqrt{-x^2 + c_1}$$

$$y(x) = e^x c_1$$

### ✓ Solution by Mathematica

Time used: 0.133 (sec). Leaf size: 54

```
DSolve[y[x]*(y'[x])^2+(x-y[x]^2)*y'[x]-x*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x$$

$$y(x) \rightarrow -\sqrt{-x^2 + 2c_1}$$

$$y(x) \rightarrow \sqrt{-x^2 + 2c_1}$$

$$y(x) \rightarrow 0$$

## 1.11 problem 11

Internal problem ID [6023]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 11.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type [\_separable]

$$y'^2 - xy(y+x)y' + x^3y^3 = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve(diff(y(x),x)^2-x*y(x)*(x+y(x))*diff(y(x),x)+x^3*y(x)^3=0,y(x), singsol=all)
```

$$y(x) = \frac{2}{-x^2 + 2c_1}$$

$$y(x) = c_1 e^{\frac{x^3}{3}}$$

### ✓ Solution by Mathematica

Time used: 0.138 (sec). Leaf size: 38

```
DSolve[(y'[x])^2-x*y[x]*(x+y[x])*y'[x]+x^3*y[x]^3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{\frac{x^3}{3}}$$

$$y(x) \rightarrow -\frac{2}{x^2 + 2c_1}$$

$$y(x) \rightarrow 0$$

## 1.12 problem 12

Internal problem ID [6024]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 12.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type [\_quadrature]

$$(-y + 4x)y'^2 + 6(-y + x)y' + 2x - 5y = 0$$

### ✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 55

```
dsolve((4*x-y(x))*diff(y(x),x)^2+6*(x-y(x))*diff(y(x),x)+2*x-5*y(x)=0,y(x), singsol=all)
```

$$y(x) = -x + c_1$$

$$y(x) = -\frac{4c_1x - \sqrt{-12c_1x + 1} - 1}{2c_1}$$

$$y(x) = -\frac{4c_1x + \sqrt{-12c_1x + 1} - 1}{2c_1}$$

### ✓ Solution by Mathematica

Time used: 1.115 (sec). Leaf size: 90

```
DSolve[(4*x-y[x])*(y'[x])^2+6*(x-y[x])*y'[x]+2*x-5*y[x]==0,y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow \frac{1}{2} \left( -4x - e^{\frac{c_1}{2}} \sqrt{12x + e^{c_1}} - e^{c_1} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left( -4x + e^{\frac{c_1}{2}} \sqrt{12x + e^{c_1}} - e^{c_1} \right)$$

$$y(x) \rightarrow -x + c_1$$

## 1.13 problem 13

Internal problem ID [6025]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 13.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cla`

$$(-y + x)^2 y' - y^2 = 0$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 45

```
dsolve((x-y(x))^2*diff(y(x),x)^2=y(x)^2,y(x), singsol=all)
```

$$y(x) = x - \sqrt{x^2 - 2c_1}$$

$$y(x) = x + \sqrt{x^2 - 2c_1}$$

$$y(x) = e^{\text{LambertW}(-x e^{-c_1}) + c_1}$$

### ✓ Solution by Mathematica

Time used: 4.364 (sec). Leaf size: 99

```
DSolve[(x-y[x])^2*(y'[x])^2==y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x - \sqrt{x^2 - e^{2c_1}}$$

$$y(x) \rightarrow x + \sqrt{x^2 - e^{2c_1}}$$

$$y(x) \rightarrow e^{W(-e^{-c_1}x) + c_1}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow x - \sqrt{x^2}$$

$$y(x) \rightarrow \sqrt{x^2} + x$$



## 1.14 problem 14

Internal problem ID [6026]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 14.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type [\_quadrature]

$$xyy'^2 + (xy^2 - 1)y' - y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve(x*y(x)*diff(y(x),x)^2+(x*y(x)^2-1)*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = \sqrt{2 \ln(x) + c_1}$$

$$y(x) = -\sqrt{2 \ln(x) + c_1}$$

$$y(x) = e^{-x}c_1$$

### ✓ Solution by Mathematica

Time used: 0.106 (sec). Leaf size: 57

```
DSolve[x*y[x]*(y'[x])^2+(x*y[x]^2-1)*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-x}$$

$$y(x) \rightarrow -\sqrt{2} \sqrt{\log(x) + c_1}$$

$$y(x) \rightarrow \sqrt{2} \sqrt{\log(x) + c_1}$$

$$y(x) \rightarrow 0$$

## 1.15 problem 15

Internal problem ID [6027]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 15.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _dAlembert]`

$$(y^2 + x^2)^2 y' - 4x^2 y^2 = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 301

```
dsolve((x^2+y(x)^2)^2*diff(y(x),x)-4*x^2*y(x)^2,y(x), singsol=all)
```

$$y(x) = -\frac{-1 + \sqrt{4c_1^2 x^2 + 1}}{2c_1}$$

$$y(x) = \frac{1 + \sqrt{4c_1^2 x^2 + 1}}{2c_1}$$

$$y(x) = \frac{\frac{(4+4\sqrt{4c_1^3 x^6+1})^{\frac{1}{3}}}{2} - \frac{2x^2 c_1}{(4+4\sqrt{4c_1^3 x^6+1})^{\frac{1}{3}}}}{\sqrt{c_1}}$$

$$y(x) = \frac{-\frac{(4+4\sqrt{4c_1^3 x^6+1})^{\frac{1}{3}}}{4} + \frac{x^2 c_1}{(4+4\sqrt{4c_1^3 x^6+1})^{\frac{1}{3}}} - \frac{i\sqrt{3} \left( \frac{(4+4\sqrt{4c_1^3 x^6+1})^{\frac{1}{3}}}{2} + \frac{2x^2 c_1}{(4+4\sqrt{4c_1^3 x^6+1})^{\frac{1}{3}}} \right)}{2}}{\sqrt{c_1}}$$

$$y(x) = \frac{-\frac{(4+4\sqrt{4c_1^3 x^6+1})^{\frac{1}{3}}}{4} + \frac{x^2 c_1}{(4+4\sqrt{4c_1^3 x^6+1})^{\frac{1}{3}}} + \frac{i\sqrt{3} \left( \frac{(4+4\sqrt{4c_1^3 x^6+1})^{\frac{1}{3}}}{2} + \frac{2x^2 c_1}{(4+4\sqrt{4c_1^3 x^6+1})^{\frac{1}{3}}} \right)}{2}}{\sqrt{c_1}}$$

✓ Solution by Mathematica

Time used: 16.202 (sec). Leaf size: 306

`DSolve[(x^2+y[x]^2)^2*(y'[x])^2==4*x^2*y[x]^2,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{1}{2} \left( -\sqrt{4x^2 + e^{2c_1}} - e^{c_1} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left( \sqrt{4x^2 + e^{2c_1}} - e^{c_1} \right)$$

$$y(x) \rightarrow \frac{\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}{\sqrt[3]{2}} - \frac{\sqrt[3]{2}x^2}{\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}$$

$$y(x) \rightarrow \frac{2\sqrt[3]{-2x^2} + (-2)^{2/3} (\sqrt{4x^6 + e^{6c_1}} + e^{3c_1})^{2/3}}{2\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}$$

$$y(x) \rightarrow -\frac{2(-1)^{2/3}x^2 + \sqrt[3]{-2}(\sqrt{4x^6 + e^{6c_1}} + e^{3c_1})^{2/3}}{2^{2/3}\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}$$

$$y(x) \rightarrow 0$$

## 1.16 problem 16

Internal problem ID [6028]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 16.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cla`

$$(y + x)^2 y'^2 + (2y^2 + xy - x^2) y' + y(y - x) = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 85

```
dsolve((y(x)+x)^2*diff(y(x),x)^2+(2*y(x)^2+x*y(x)-x^2)*diff(y(x),x)+y(x)*(y(x)-x)=0,y(x), sin
```

$$y(x) = -x - \sqrt{x^2 + 2c_1}$$

$$y(x) = -x + \sqrt{x^2 + 2c_1}$$

$$y(x) = \frac{-c_1 x - \sqrt{2c_1^2 x^2 + 1}}{c_1}$$

$$y(x) = \frac{-c_1 x + \sqrt{2c_1^2 x^2 + 1}}{c_1}$$

✓ Solution by Mathematica

Time used: 0.65 (sec). Leaf size: 172

`DSolve[(y[x]+x)^2*(y'[x])^2+(2*y[x]^2+x*y[x]-x^2)*y'[x]+y[x]*(y[x]-x)==0,y[x],x,IncludeSingularSolutions->True]`

$$y(x) \rightarrow -x - \sqrt{x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -x + \sqrt{x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -x - \sqrt{2x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -x + \sqrt{2x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -\sqrt{x^2} - x$$

$$y(x) \rightarrow \sqrt{x^2} - x$$

$$y(x) \rightarrow -\sqrt{2}\sqrt{x^2} - x$$

$$y(x) \rightarrow \sqrt{2}\sqrt{x^2} - x$$

## 1.17 problem 17

Internal problem ID [6029]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 17.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _Bernoulli]`

$$xy(y^2 + x^2)(y'^2 - 1) - y'(x^4 + x^2y^2 + y^4) = 0$$

✓ Solution by Maple

Time used: 0.594 (sec). Leaf size: 250

```
dsolve(x*y(x)*(x^2+y(x)^2)*(diff(y(x),x)^2-1)=diff(y(x),x)*(x^4+x^2*y(x)^2+y(x)^4),y(x),sing
```

$$y(x) = \frac{\sqrt{x^2 c_1 (c_1 x^2 - \sqrt{c_1^2 x^4 + 1})}}{x (c_1 x^2 - \sqrt{c_1^2 x^4 + 1}) c_1}$$

$$y(x) = \frac{\sqrt{x^2 c_1 (c_1 x^2 + \sqrt{c_1^2 x^4 + 1})}}{x (c_1 x^2 + \sqrt{c_1^2 x^4 + 1}) c_1}$$

$$y(x) = -\frac{\sqrt{x^2 c_1 (c_1 x^2 - \sqrt{c_1^2 x^4 + 1})}}{x (c_1 x^2 - \sqrt{c_1^2 x^4 + 1}) c_1}$$

$$y(x) = -\frac{\sqrt{x^2 c_1 (c_1 x^2 + \sqrt{c_1^2 x^4 + 1})}}{x (c_1 x^2 + \sqrt{c_1^2 x^4 + 1}) c_1}$$

$$y(x) = \sqrt{2 \ln(x) + c_1} x$$

$$y(x) = -\sqrt{2 \ln(x) + c_1} x$$

✓ Solution by Mathematica

Time used: 8.924 (sec). Leaf size: 248

`DSolve[x*y[x]*(x^2+y[x]^2)*((y'[x])^2-1)==y'[x]*(x^4+x^2*y[x]^2+y[x]^4),y[x],x,IncludeSingular`

$$y(x) \rightarrow -\sqrt{-x^2 - \sqrt{x^4 + e^{4c_1}}}$$

$$y(x) \rightarrow \sqrt{-x^2 - \sqrt{x^4 + e^{4c_1}}}$$

$$y(x) \rightarrow -\sqrt{-x^2 + \sqrt{x^4 + e^{4c_1}}}$$

$$y(x) \rightarrow \sqrt{-x^2 + \sqrt{x^4 + e^{4c_1}}}$$

$$y(x) \rightarrow -x\sqrt{2\log(x) + c_1}$$

$$y(x) \rightarrow x\sqrt{2\log(x) + c_1}$$

$$y(x) \rightarrow -\sqrt{-\sqrt{x^4} - x^2}$$

$$y(x) \rightarrow \sqrt{-\sqrt{x^4} - x^2}$$

$$y(x) \rightarrow -\sqrt{\sqrt{x^4} - x^2}$$

$$y(x) \rightarrow \sqrt{\sqrt{x^4} - x^2}$$

## 1.18 problem 18

Internal problem ID [6030]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 18.

**ODE order:** 1.

**ODE degree:** 3.

CAS Maple gives this as type [\_quadrature]

$$xy^3 - (x^2 + x + y)y'^2 + (x^2 + xy + y)y' - xy = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(x*diff(y(x),x)^3-(x^2+x+y(x))*diff(y(x),x)^2+(x^2+x*y(x)+y(x))*diff(y(x),x)-x*y(x)=0,y
```

$$y(x) = c_1x$$

$$y(x) = x + c_1$$

$$y(x) = \frac{x^2}{2} + c_1$$

### ✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 36

```
DSolve[x*(y'[x])^3-(x^2+x+y[x])*(y'[x])^2+(x^2+x*y[x]+y[x])*y'[x]-x*y[x]==0,y[x],x,IncludeSin
```

$$y(x) \rightarrow c_1x$$

$$y(x) \rightarrow x + c_1$$

$$y(x) \rightarrow \frac{x^2}{2} + c_1$$

$$y(x) \rightarrow 0$$



## 1.19 problem 19

Internal problem ID [6031]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

**Problem number:** 19.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type [\_quadrature]

$$xyy'^2 + (y + x)y' + 1 = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 32

```
dsolve(x*y(x)*diff(y(x),x)^2+(x+y(x))*diff(y(x),x)+1=0,y(x), singsol=all)
```

$$y(x) = -\ln(x) + c_1$$

$$y(x) = \sqrt{-2x + c_1}$$

$$y(x) = -\sqrt{-2x + c_1}$$

### ✓ Solution by Mathematica

Time used: 0.063 (sec). Leaf size: 53

```
DSolve[x*y[x]*(y'[x])^2+(x+y[x])*y'[x]+1==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{2}\sqrt{-x + c_1}$$

$$y(x) \rightarrow \sqrt{2}\sqrt{-x + c_1}$$

$$y(x) \rightarrow -\log(x) + c_1$$

**2 CHAPTER 16. Nonlinear equations. Section 97.**  
**The p-discriminant equation. EXERCISES Page**  
**314**

2.1	problem 8 . . . . .	26
2.2	problem 9 . . . . .	27
2.3	problem 10 . . . . .	29
2.4	problem 11 . . . . .	31
2.5	problem 12 . . . . .	32
2.6	problem 13 . . . . .	34
2.7	problem 14 . . . . .	36
2.8	problem 15 . . . . .	38
2.9	problem 16 . . . . .	41

## 2.1 problem 8

Internal problem ID [6032]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

**Problem number:** 8.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _dAlembert]`

$$xy'^2 - 2yy' + 4x = 0$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 29

```
dsolve(x*diff(y(x),x)^2-2*y(x)*diff(y(x),x)+4*x=0,y(x), singsol=all)
```

$$y(x) = -2x$$

$$y(x) = 2x$$

$$y(x) = -\frac{\left(-\frac{x^2}{c_1^2} - 4\right) c_1}{2}$$

### ✓ Solution by Mathematica

Time used: 0.282 (sec). Leaf size: 43

```
DSolve[x*(y'[x])^2-2*y[x]*y'[x]+4*x==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2x \cosh(-\log(x) + c_1)$$

$$y(x) \rightarrow -2x \cosh(\log(x) + c_1)$$

$$y(x) \rightarrow -2x$$

$$y(x) \rightarrow 2x$$

## 2.2 problem 9

Internal problem ID [6033]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

**Problem number:** 9.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[_homogeneous, 'class G'], _rational]`

$$3x^4y'^2 - y'x - y = 0$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 147

```
dsolve(3*x^4*diff(y(x),x)^2-x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{1}{12x^2}$$

$$y(x) = \frac{-c_1(-c_1 + 2ix\sqrt{3}) - c_1^2 - 6x^2}{6x^2c_1^2}$$

$$y(x) = \frac{-c_1(-c_1 - 2ix\sqrt{3}) - c_1^2 - 6x^2}{6x^2c_1^2}$$

$$y(x) = \frac{c_1(c_1 + 2ix\sqrt{3}) - 6x^2 - c_1^2}{6c_1^2x^2}$$

$$y(x) = \frac{c_1(c_1 - 2ix\sqrt{3}) - 6x^2 - c_1^2}{6c_1^2x^2}$$

✓ Solution by Mathematica

Time used: 0.494 (sec). Leaf size: 123

```
DSolve[3*x^4*(y'[x])^2-x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ \frac{x \sqrt{12x^2y(x) + 1} \operatorname{arctanh}(\sqrt{12x^2y(x) + 1})}{\sqrt{12x^4y(x) + x^2}} - \frac{1}{2} \log(y(x)) = c_1, y(x) \right]$$

$$\text{Solve} \left[ \frac{x \sqrt{12x^2y(x) + 1} \operatorname{arctanh}(\sqrt{12x^2y(x) + 1})}{\sqrt{12x^4y(x) + x^2}} - \frac{1}{2} \log(y(x)) = c_1, y(x) \right]$$

$$y(x) \rightarrow 0$$

## 2.3 problem 10

Internal problem ID [6034]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

**Problem number:** 10.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$y'^2 - y'x - y = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 77

```
dsolve(diff(y(x),x)^2-x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$\frac{c_1}{\sqrt{2x - 2\sqrt{x^2 + 4y(x)}}} + \frac{2x}{3} + \frac{\sqrt{x^2 + 4y(x)}}{3} = 0$$

$$\frac{c_1}{\sqrt{2x + 2\sqrt{x^2 + 4y(x)}}} + \frac{2x}{3} - \frac{\sqrt{x^2 + 4y(x)}}{3} = 0$$

✓ Solution by Mathematica

Time used: 60.17 (sec). Leaf size: 965

`DSolve[(y'[x])^2 - x*y'[x] - y[x] == 0, y[x], x, IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{\left(x^2 + \sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}}\right)^2 + 8e^{3c_1}x}{4\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}}}$$

$$y(x) \rightarrow \frac{1}{8} \left( 4x^2 + \frac{(-1 - i\sqrt{3})x(x^3 + 8e^{3c_1})}{\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}}} + i(\sqrt{3} + i)\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}} \right)$$

$$y(x) \rightarrow \frac{1}{8} \left( 4x^2 + \frac{i(\sqrt{3} + i)x(x^3 + 8e^{3c_1})}{\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}}} - (1 + i\sqrt{3})\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}} \right)$$

$$y(x) \rightarrow \frac{2\sqrt[3]{2}x^4 + 2^{2/3}\left(-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}(4x^3 + e^{3c_1})^3 + e^{6c_1}}\right)^{2/3} + 4x^2\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}(4x^3 + e^{3c_1})^3 + e^{6c_1}}}}{8\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}(4x^3 + e^{3c_1})^3 + e^{6c_1}}}}$$

$$y(x) \rightarrow \frac{1}{16} \left( 8x^2 - \frac{4\sqrt[3]{-2}x(x^3 - 2e^{3c_1})}{\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}(4x^3 + e^{3c_1})^3 + e^{6c_1}}}} + 2(-2)^{2/3}\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}(4x^3 + e^{3c_1})^3 + e^{6c_1}}} \right)$$

$$y(x) \rightarrow \frac{x^2}{2} + \frac{(-1)^{2/3}x(x^3 - 2e^{3c_1})}{2 \cdot 2^{2/3}\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}(4x^3 + e^{3c_1})^3 + e^{6c_1}}}} - \frac{1}{4}\sqrt[3]{-\frac{1}{2}\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}(4x^3 + e^{3c_1})^3 + e^{6c_1}}}}$$

## 2.4 problem 11

Internal problem ID [6035]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

**Problem number:** 11.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _Clairaut]`

$$y'^2 - y'x + y = 0$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)^2-x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x^2}{4}$$

$$y(x) = -c_1^2 + c_1x$$

### ✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 25

```
DSolve[(y'[x])^2-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1(x - c_1)$$

$$y(x) \rightarrow \frac{x^2}{4}$$



## 2.5 problem 12

Internal problem ID [6036]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

**Problem number:** 12.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$y'^2 + 4x^5y' - 12x^4y = 0$$

✓ Solution by Maple

Time used: 0.219 (sec). Leaf size: 23

```
dsolve(diff(y(x),x)^2+4*x^5*diff(y(x),x)-12*x^4*y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{x^6}{3}$$

$$y(x) = c_1x^3 + \frac{3}{4}c_1^2$$

✓ Solution by Mathematica

Time used: 1.306 (sec). Leaf size: 217

`DSolve[(y'[x])^2+4*x^5*y'[x]-12*x^4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$\text{Solve} \left[ \frac{1}{6} \left( \log(y(x)) - \frac{x^2 \sqrt{x^6 + 3y(x)} \log(y(x))}{\sqrt{x^4 (x^6 + 3y(x))}} \right) + \frac{x^2 \sqrt{x^6 + 3y(x)} \log(\sqrt{x^6 + 3y(x)} + x^3)}{3\sqrt{x^4 (x^6 + 3y(x))}} = c_1, y(x) \right]$$

$$\text{Solve} \left[ \frac{1}{6} \left( \frac{x^2 \sqrt{x^6 + 3y(x)} \log(y(x))}{\sqrt{x^4 (x^6 + 3y(x))}} + \log(y(x)) \right) - \frac{x^2 \sqrt{x^6 + 3y(x)} \log(\sqrt{x^6 + 3y(x)} + x^3)}{3\sqrt{x^4 (x^6 + 3y(x))}} = c_1, y(x) \right]$$

$$y(x) \rightarrow -\frac{x^6}{3}$$

## 2.6 problem 13

Internal problem ID [6037]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

**Problem number:** 13.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _rational]`

$$4y^3y'^2 - 4y'x + y = 0$$

### ✓ Solution by Maple

Time used: 0.265 (sec). Leaf size: 81

```
dsolve(4*y(x)^3*diff(y(x),x)^2-4*x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \sqrt{-x}$$

$$y(x) = -\sqrt{-x}$$

$$y(x) = \sqrt{x}$$

$$y(x) = -\sqrt{x}$$

$$y(x) = 0$$

$$y(x) = \text{RootOf} \left( -\ln(x) + \int^{-z} -\frac{2(-a^4 + \sqrt{-a^4 + 1} - 1)}{-a(-a^4 - 1)} d_a + c_1 \right) \sqrt{x}$$

✓ Solution by Mathematica

Time used: 0.561 (sec). Leaf size: 282

`DSolve[4*y[x]^3*(y'[x])^2-4*x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow -e^{\frac{c_1}{4}} \sqrt[4]{e^{c_1} - 2ix}$$

$$y(x) \rightarrow -ie^{\frac{c_1}{4}} \sqrt[4]{e^{c_1} - 2ix}$$

$$y(x) \rightarrow ie^{\frac{c_1}{4}} \sqrt[4]{e^{c_1} - 2ix}$$

$$y(x) \rightarrow e^{\frac{c_1}{4}} \sqrt[4]{e^{c_1} - 2ix}$$

$$y(x) \rightarrow -e^{\frac{c_1}{4}} \sqrt[4]{2ix + e^{c_1}}$$

$$y(x) \rightarrow -ie^{\frac{c_1}{4}} \sqrt[4]{2ix + e^{c_1}}$$

$$y(x) \rightarrow ie^{\frac{c_1}{4}} \sqrt[4]{2ix + e^{c_1}}$$

$$y(x) \rightarrow e^{\frac{c_1}{4}} \sqrt[4]{2ix + e^{c_1}}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -\sqrt{x}$$

$$y(x) \rightarrow -i\sqrt{x}$$

$$y(x) \rightarrow i\sqrt{x}$$

$$y(x) \rightarrow \sqrt{x}$$

## 2.7 problem 14

Internal problem ID [6038]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

**Problem number:** 14.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _rational]`

$$4y^3y'^2 + 4y'x + y = 0$$

✓ Solution by Maple

Time used: 0.297 (sec). Leaf size: 287

```
dsolve(4*y(x)^3*diff(y(x),x)^2+4*x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$\int_{-b}^x \frac{-2_a + \sqrt{-y(x)^4 + _a^2}}{2y(x)^4 + 6_a^2} d_a + \int^{y(x)} \left( \frac{_{f^\beta}}{-_{f^4} + \sqrt{-_{f^4} + x^2 x - x^2}} \right. \\ \left. - \left( \int_{-b}^x \left( -\frac{_{f^\beta}}{\sqrt{-_{f^4} + _a^2} (_{f^4} + 3_a^2)} - \frac{2(-2_a + \sqrt{-_{f^4} + _a^2})_{f^\beta}}{(_{f^4} + 3_a^2)^2} \right) d_a \right) \right) d_f \\ + c_1 = 0$$

$$\int_{-b}^x \frac{2_a + \sqrt{-y(x)^4 + _a^2}}{2(y(x)^4 + 3_a^2)} d_a + \int^{y(x)} \left( -\frac{_{f^\beta}}{-_{f^4} + \sqrt{-_{f^4} + x^2 x + x^2}} \right. \\ \left. - \left( \int_{-b}^x \left( \frac{_{f^\beta}}{\sqrt{-_{f^4} + _a^2} (_{f^4} + 3_a^2)} + \frac{2(2_a + \sqrt{-_{f^4} + _a^2})_{f^\beta}}{(_{f^4} + 3_a^2)^2} \right) d_a \right) \right) d_f \\ + c_1 = 0$$

✓ Solution by Mathematica

Time used: 60.319 (sec). Leaf size: 2815

```
DSolve[4*y[x]^3*(y'[x])^2+4*x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

## 2.8 problem 15

Internal problem ID [6039]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

**Problem number:** 15.

**ODE order:** 1.

**ODE degree:** 3.

CAS Maple gives this as type [`_dAlembert`]

$$y'^3 + xy'^2 - y = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 1473

```
dsolve(diff(y(x),x)^3+x*diff(y(x),x)^2-y(x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x)$$

$$= \left( \frac{(-36x^2 - 54x + 108c_1 - 8x^3 + 27 + 6\sqrt{-48c_1x^3 - 216c_1x^2 - 24x^3 + 324c_1^2 - 324c_1x - 108x^2 + 162c_1})}{6} \right.$$

$$+ x \left( \frac{(-36x^2 - 54x + 108c_1 - 8x^3 + 27 + 6\sqrt{-48c_1x^3 - 216c_1x^2 - 24x^3 + 324c_1^2 - 324c_1x - 108x^2 + 162c_1})}{6} \right.$$

$$y(x)$$

$$= \left( - \frac{(-36x^2 - 54x + 108c_1 - 8x^3 + 27 + 6\sqrt{-48c_1x^3 - 216c_1x^2 - 24x^3 + 324c_1^2 - 324c_1x - 108x^2 + 162c_1})}{12} \right.$$

$$+ x \left( - \frac{(-36x^2 - 54x + 108c_1 - 8x^3 + 27 + 6\sqrt{-48c_1x^3 - 216c_1x^2 - 24x^3 + 324c_1^2 - 324c_1x - 108x^2 + 162c_1})}{12} \right.$$

$$y(x)$$

$$= \left( - \frac{(-36x^2 - 54x + 108c_1 - 8x^3 + 27 + 6\sqrt{-48c_1x^3 - 216c_1x^2 - 24x^3 + 324c_1^2 - 324c_1x - 108x^2 + 162c_1})}{12} \right.$$

$$+ x \left( - \frac{(-36x^2 - 54x + 108c_1 - 8x^3 + 27 + 6\sqrt{-48c_1x^3 - 216c_1x^2 - 24x^3 + 324c_1^2 - 324c_1x - 108x^2 + 162c_1})}{12} \right.$$



✓ Solution by Mathematica

Time used: 83.056 (sec). Leaf size: 1410

`DSolve[(y'[x])^3+x*(y'[x])^2-y[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{1}{24} \left( -4x^2 + 2x \left( 6 \right. \right. \\ \left. \left. + \sqrt[3]{-2x(2x(2x+9)+27) + 3 \left( 2\sqrt{6} \sqrt{-((1+2c_1)(x(2x(2x+9)+27) - 27c_1)) + 9 + 36c_1} \right)} + 9 + 36c_1 \right) \right) \\ \left. + 3 \left( 9 \right. \right. \\ \left. \left. + \sqrt[3]{-2x(2x(2x+9)+27) + 3 \left( 2\sqrt{6} \sqrt{-((1+2c_1)(x(2x(2x+9)+27) - 27c_1)) + 9 + 36c_1} \right)} + 9 + 36c_1 \right) \right) \\ + \frac{24c_1(2x+3)^3 - (2x+3)^3 \left( -2x(2x(2x+9)+27) + 3 \left( 2\sqrt{6} \sqrt{-((1+2c_1)(x(2x(2x+9)+27) - 27c_1)) + 9 + 36c_1} \right)} \right)}{24}$$

$$y(x) \rightarrow \frac{1}{6} \left( 2(3-2x)x - 6x \right. \\ \left. - \frac{i(\sqrt{3}-i)x(2x+3)^2}{\sqrt[3]{-2x(2x(2x+9)+27) + 3 \left( 2\sqrt{6} \sqrt{-((1+2c_1)(x(2x(2x+9)+27) - 27c_1)) + 9 + 36c_1} \right)} + \frac{1}{16} \left( -4x \right. \right. \\ \left. \left. - \frac{i(\sqrt{3}-i)(2x+3)^2}{\sqrt[3]{-2x(2x(2x+9)+27) + 3 \left( 2\sqrt{6} \sqrt{-((1+2c_1)(x(2x(2x+9)+27) - 27c_1)) + 9 + 36c_1} \right)} \right)} \right. \\ \left. + i(\sqrt{3}+i) \sqrt[3]{-2x(2x(2x+9)+27) + 3 \left( 2\sqrt{6} \sqrt{-((1+2c_1)(x(2x(2x+9)+27) - 27c_1)) + 9 + 36c_1} \right)} + 9 + 36c_1 \right) \right. \\ \left. + 6 \right)^2 + i(\sqrt{3} \\ + i) x \sqrt[3]{-2x(2x(2x+9)+27) + 3 \left( 2\sqrt{6} \sqrt{-((1+2c_1)(x(2x(2x+9)+27) - 27c_1)) + 9 + 36c_1} \right)} \\ \left. \right)$$

## 2.9 problem 16

Internal problem ID [6040]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

**Problem number:** 16.

**ODE order:** 1.

**ODE degree:** 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$y^4 y'^3 - 6y'x + 2y = 0$$

### ✓ Solution by Maple

Time used: 0.25 (sec). Leaf size: 183

```
dsolve(y(x)^4*diff(y(x),x)^3-6*x*diff(y(x),x)+2*y(x)=0,y(x), singsol=all)
```

$$y(x) = \sqrt{-ix\sqrt{3} - x}$$

$$y(x) = \sqrt{ix\sqrt{3} - x}$$

$$y(x) = -\sqrt{-ix\sqrt{3} - x}$$

$$y(x) = -\sqrt{ix\sqrt{3} - x}$$

$$y(x) = \sqrt{x} \sqrt{2}$$

$$y(x) = -\sqrt{x} \sqrt{2}$$

$$y(x) = 0$$

$$y(x) = \frac{(-4c_1^3 + 24c_1x)^{\frac{1}{3}}}{2}$$

$$y(x) = -\frac{(-4c_1^3 + 24c_1x)^{\frac{1}{3}}}{4} - \frac{i\sqrt{3}(-4c_1^3 + 24c_1x)^{\frac{1}{3}}}{4}$$

$$y(x) = -\frac{(-4c_1^3 + 24c_1x)^{\frac{1}{3}}}{4} + \frac{i\sqrt{3}(-4c_1^3 + 24c_1x)^{\frac{1}{3}}}{4}$$

✓ Solution by Mathematica

Time used: 69.212 (sec). Leaf size: 22649

```
DSolve[y[x]^4*(y'[x])^3-6*x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

### 3 CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

3.1	problem 3	44
3.2	problem 4	46
3.3	problem 5	48
3.4	problem 6	49
3.5	problem 7	50
3.6	problem 8	51
3.7	problem 9	53
3.8	problem 10	55
3.9	problem 11	56
3.10	problem 12	58
3.11	problem 13	59
3.12	problem 14	60
3.13	problem 15	61
3.14	problem 16	62
3.15	problem 17	63
3.16	problem 19	65
3.17	problem 20	67
3.18	problem 21	68
3.19	problem 22	69
3.20	problem 23	70
3.21	problem 24	72
3.22	problem 25	73
3.23	problem 26	75
3.24	problem 27	76
3.25	problem 28	78
3.26	problem 29	80

### 3.1 problem 3

Internal problem ID [6041]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 3.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$y'^2 + x^3 y' - 2yx^2 = 0$$

✓ Solution by Maple

Time used: 0.219 (sec). Leaf size: 23

```
dsolve(diff(y(x),x)^2+x^3*diff(y(x),x)-2*x^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{x^4}{8}$$

$$y(x) = c_1 x^2 + 2c_1^2$$

✓ Solution by Mathematica

Time used: 1.162 (sec). Leaf size: 209

`DSolve[(y'[x])^2+x^3*y'[x]-2*x^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$\text{Solve} \left[ \frac{\sqrt{x^6 + 8x^2y(x)} \log(\sqrt{x^4 + 8y(x)} + x^2)}{2x\sqrt{x^4 + 8y(x)}} + \frac{1}{4} \left( 1 - \frac{\sqrt{x^6 + 8x^2y(x)}}{x\sqrt{x^4 + 8y(x)}} \right) \log(y(x)) = c_1, y(x) \right]$$

$$\text{Solve} \left[ \frac{1}{4} \left( \frac{\sqrt{x^6 + 8x^2y(x)}}{x\sqrt{x^4 + 8y(x)}} + 1 \right) \log(y(x)) - \frac{\sqrt{x^6 + 8x^2y(x)} \log(\sqrt{x^4 + 8y(x)} + x^2)}{2x\sqrt{x^4 + 8y(x)}} = c_1, y(x) \right]$$

$$y(x) \rightarrow -\frac{x^4}{8}$$

## 3.2 problem 4

Internal problem ID [6042]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 4.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$y'^2 + 4x^5y' - 12x^4y = 0$$

✓ Solution by Maple

Time used: 0.219 (sec). Leaf size: 23

```
dsolve(diff(y(x),x)^2+4*x^5*diff(y(x),x)-12*x^4*y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{x^6}{3}$$

$$y(x) = c_1x^3 + \frac{3}{4}c_1^2$$

✓ Solution by Mathematica

Time used: 0.574 (sec). Leaf size: 217

`DSolve[(y'[x])^2+4*x^5*y'[x]-12*x^4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$\text{Solve} \left[ \frac{1}{6} \left( \log(y(x)) - \frac{x^2 \sqrt{x^6 + 3y(x)} \log(y(x))}{\sqrt{x^4 (x^6 + 3y(x))}} \right) + \frac{x^2 \sqrt{x^6 + 3y(x)} \log(\sqrt{x^6 + 3y(x)} + x^3)}{3\sqrt{x^4 (x^6 + 3y(x))}} = c_1, y(x) \right]$$

$$\text{Solve} \left[ \frac{1}{6} \left( \frac{x^2 \sqrt{x^6 + 3y(x)} \log(y(x))}{\sqrt{x^4 (x^6 + 3y(x))}} + \log(y(x)) \right) - \frac{x^2 \sqrt{x^6 + 3y(x)} \log(\sqrt{x^6 + 3y(x)} + x^3)}{3\sqrt{x^4 (x^6 + 3y(x))}} = c_1, y(x) \right]$$

$$y(x) \rightarrow -\frac{x^6}{3}$$



### 3.3 problem 5

Internal problem ID [6043]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 5.

**ODE order:** 1.

**ODE degree:** 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$2xy'^3 - 6yy'^2 + x^4 = 0$$

#### ✓ Solution by Maple

Time used: 0.297 (sec). Leaf size: 56

```
dsolve(2*x*diff(y(x),x)^3-6*y(x)*diff(y(x),x)^2+x^4=0,y(x), singsol=all)
```

$$y(x) = \frac{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x^2}{2}$$

$$y(x) = \frac{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x^2}{2}$$

$$y(x) = \frac{x^2}{2}$$

$$y(x) = \frac{1}{6c_1^2} + \frac{c_1x^3}{3}$$

#### ✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[2*x*(y'[x])^3-6*y[x]*(y'[x])^2+x^4==0,y[x],x,IncludeSingularSolutions -> True]
```

Timed out

### 3.4 problem 6

Internal problem ID [6044]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 6.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _Clairaut]`

$$y'^2 - y'x + y = 0$$

#### ✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)^2-x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x^2}{4}$$

$$y(x) = -c_1^2 + c_1x$$

#### ✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 25

```
DSolve[(y'[x])^2-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1(x - c_1)$$

$$y(x) \rightarrow \frac{x^2}{4}$$

### 3.5 problem 7

Internal problem ID [6045]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 7.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _Clairaut]`

$$y - y'x - ky'^2 = 0$$

#### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 24

```
dsolve(y(x)=diff(y(x),x)*x+k*diff(y(x),x)^2,y(x), singsol=all)
```

$$y(x) = -\frac{x^2}{4k}$$

$$y(x) = c_1^2 k + c_1 x$$

#### ✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 28

```
DSolve[y[x]==y'[x]*x+k*(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1(x + c_1 k)$$

$$y(x) \rightarrow -\frac{x^2}{4k}$$

### 3.6 problem 8

Internal problem ID [6046]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 8.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[_homogeneous, 'class G']`

$$x^8 y'^2 + 3y'x + 9y = 0$$

✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 42

```
dsolve(x^8*diff(y(x),x)^2+3*x*diff(y(x),x)+9*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{1}{4x^6}$$

$$y(x) = \frac{-x^3 + c_1}{x^3 c_1^2}$$

$$y(x) = -\frac{x^3 + c_1}{x^3 c_1^2}$$

✓ Solution by Mathematica

Time used: 0.537 (sec). Leaf size: 130

`DSolve[x^8*(y'[x])^2+3*x*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$\text{Solve} \left[ \frac{x\sqrt{4x^6y(x)-1} \arctan\left(\sqrt{4x^6y(x)-1}\right)}{3\sqrt{x^2-4x^8y(x)}} - \frac{1}{6} \log(y(x)) = c_1, y(x) \right]$$

$$\text{Solve} \left[ \frac{\sqrt{x^2-4x^8y(x)} \arctan\left(\sqrt{4x^6y(x)-1}\right)}{3x\sqrt{4x^6y(x)-1}} - \frac{1}{6} \log(y(x)) = c_1, y(x) \right]$$

$$y(x) \rightarrow 0$$

### 3.7 problem 9

Internal problem ID [6047]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 9.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational]`

$$x^4 y'^2 + 2x^3 y' y - 4 = 0$$

✓ Solution by Maple

Time used: 0.109 (sec). Leaf size: 49

```
dsolve(x^4*diff(y(x),x)^2+2*x^3*y(x)*diff(y(x),x)-4=0,y(x), singsol=all)
```

$$y(x) = -\frac{2i}{x}$$

$$y(x) = \frac{2i}{x}$$

$$y(x) = \frac{2 \sinh(-\ln(x) + c_1)}{x}$$

$$y(x) = -\frac{2 \sinh(-\ln(x) + c_1)}{x}$$

✓ Solution by Mathematica

Time used: 0.642 (sec). Leaf size: 71

```
DSolve[x^4*(y'[x])^2+2*x^3*y[x]*y'[x]-4==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{4e^{c_1}}{x^2} - \frac{e^{-c_1}}{4}$$

$$y(x) \rightarrow \frac{e^{-c_1}}{4} - \frac{4e^{c_1}}{x^2}$$

$$y(x) \rightarrow -\frac{2i}{x}$$

$$y(x) \rightarrow \frac{2i}{x}$$

### 3.8 problem 10

Internal problem ID [6048]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 10.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _dAlembert]`

$$xy'^2 - 2yy' + 4x = 0$$

#### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 29

```
dsolve(x*diff(y(x),x)^2-2*y(x)*diff(y(x),x)+4*x=0,y(x), singsol=all)
```

$$y(x) = -2x$$

$$y(x) = 2x$$

$$y(x) = -\frac{\left(-\frac{x^2}{c_1^2} - 4\right) c_1}{2}$$

#### ✓ Solution by Mathematica

Time used: 0.249 (sec). Leaf size: 43

```
DSolve[x*(y'[x])^2-2*y[x]*y'[x]+4*x==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2x \cosh(-\log(x) + c_1)$$

$$y(x) \rightarrow -2x \cosh(\log(x) + c_1)$$

$$y(x) \rightarrow -2x$$

$$y(x) \rightarrow 2x$$



### 3.9 problem 11

Internal problem ID [6049]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 11.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[_homogeneous, 'class G'], _rational]`

$$3x^4y'^2 - y'x - y = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 147

```
dsolve(3*x^4*diff(y(x),x)^2-x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{1}{12x^2}$$

$$y(x) = \frac{-c_1(-c_1 + 2ix\sqrt{3}) - c_1^2 - 6x^2}{6x^2c_1^2}$$

$$y(x) = \frac{-c_1(-c_1 - 2ix\sqrt{3}) - c_1^2 - 6x^2}{6x^2c_1^2}$$

$$y(x) = \frac{c_1(c_1 + 2ix\sqrt{3}) - 6x^2 - c_1^2}{6c_1^2x^2}$$

$$y(x) = \frac{c_1(c_1 - 2ix\sqrt{3}) - 6x^2 - c_1^2}{6c_1^2x^2}$$

✓ Solution by Mathematica

Time used: 0.473 (sec). Leaf size: 123

```
DSolve[3*x^4*(y'[x])^2-x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ \frac{x \sqrt{12x^2y(x) + 1} \operatorname{arctanh}(\sqrt{12x^2y(x) + 1})}{\sqrt{12x^4y(x) + x^2}} - \frac{1}{2} \log(y(x)) = c_1, y(x) \right]$$

$$\text{Solve} \left[ \frac{x \sqrt{12x^2y(x) + 1} \operatorname{arctanh}(\sqrt{12x^2y(x) + 1})}{\sqrt{12x^4y(x) + x^2}} - \frac{1}{2} \log(y(x)) = c_1, y(x) \right]$$

$$y(x) \rightarrow 0$$

### 3.10 problem 12

Internal problem ID [6050]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 12.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _rational, _dAlembert]`

$$xy'^2 + (x - y)y' + 1 - y = 0$$

#### ✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 56

```
dsolve(x*diff(y(x),x)^2+(x-y(x))*diff(y(x),x)+1-y(x)=0,y(x), singsol=all)
```

$$y(x) = -x - 2\sqrt{x}$$

$$y(x) = -x + 2\sqrt{x}$$

$$y(x) = \frac{(-c_1^2 - c_1)x}{-1 - c_1} - \frac{1}{-1 - c_1}$$

#### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 46

```
DSolve[x*(y'[x])^2+(x-y[x])*y'[x]+1-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x + \frac{1}{1 + c_1}$$

$$y(x) \rightarrow -x - 2\sqrt{x}$$

$$y(x) \rightarrow 2\sqrt{x} - x$$

### 3.11 problem 13

Internal problem ID [6051]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 13.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[_1st_order, _with_linear_symmetries], _rational, _Clairaut]`

$$y'(y'x - y + k) + a = 0$$

#### ✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 41

```
dsolve(diff(y(x),x)*( x*diff(y(x),x)-y(x)+k )+a=0,y(x), singsol=all)
```

$$y(x) = k - 2\sqrt{ax}$$

$$y(x) = k + 2\sqrt{ax}$$

$$y(x) = c_1x + \frac{c_1k + a}{c_1}$$

#### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 58

```
DSolve[y'[x]*( x*y'[x]-y[x]+k )+a==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{a}{c_1} + k + c_1x$$

$$y(x) \rightarrow \text{Indeterminate}$$

$$y(x) \rightarrow k - 2\sqrt{a}\sqrt{x}$$

$$y(x) \rightarrow 2\sqrt{a}\sqrt{x} + k$$

### 3.12 problem 14

Internal problem ID [6052]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 14.

**ODE order:** 1.

**ODE degree:** 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$x^6 y'^3 - 3y'x - 3y = 0$$

#### ✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 32

```
dsolve(x^6*diff(y(x),x)^3-3*x*diff(y(x),x)-3*y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{2}{3x^{\frac{3}{2}}}$$

$$y(x) = \frac{2}{3x^{\frac{3}{2}}}$$

$$y(x) = \frac{c_1^3}{3} - \frac{c_1}{x}$$

#### ✓ Solution by Mathematica

Time used: 134.736 (sec). Leaf size: 24834

```
DSolve[x^6*(y'[x])^3-3*x*y'[x]-3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

### 3.13 problem 15

Internal problem ID [6053]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 15.

**ODE order:** 1.

**ODE degree:** 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$y - x^6 y'^3 + y'x = 0$$

#### ✓ Solution by Maple

Time used: 0.234 (sec). Leaf size: 36

```
dsolve(y(x)=x^6*diff(y(x),x)^3-x*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = -\frac{2\sqrt{3}}{9x^{\frac{3}{2}}}$$

$$y(x) = \frac{2\sqrt{3}}{9x^{\frac{3}{2}}}$$

$$y(x) = c_1^3 - \frac{c_1}{x}$$

#### ✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y[x]==x^6*(y'[x])^3-x*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

Timed out

### 3.14 problem 16

Internal problem ID [6054]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 16.

**ODE order:** 1.

**ODE degree:** 4.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$xy'^4 - 2yy'^3 + 12x^3 = 0$$

#### ✓ Solution by Maple

Time used: 0.297 (sec). Leaf size: 62

```
dsolve(x*diff(y(x),x)^4-2*y(x)*diff(y(x),x)^3+12*x^3=0,y(x), singsol=all)
```

$$y(x) = -\frac{2\sqrt{-6x}x}{3}$$

$$y(x) = \frac{2\sqrt{-6x}x}{3}$$

$$y(x) = -\frac{2\sqrt{6}x^{\frac{3}{2}}}{3}$$

$$y(x) = \frac{2\sqrt{6}x^{\frac{3}{2}}}{3}$$

$$y(x) = 6c_1^3 + \frac{x^2}{2c_1}$$

#### ✓ Solution by Mathematica

Time used: 36.401 (sec). Leaf size: 30947

```
DSolve[x*(y'[x])^4-2*y[x]*(y'[x])^3+12*x^3==0,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

### 3.15 problem 17

Internal problem ID [6055]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 17.

**ODE order:** 1.

**ODE degree:** 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _Clairaut]`

$$xy'^3 - yy'^2 + 1 = 0$$

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 80

```
dsolve(x*diff(y(x),x)^3-y(x)*diff(y(x),x)^2+1=0,y(x), singsol=all)
```

$$y(x) = \frac{3 \cdot 2^{\frac{1}{3}} (x^2)^{\frac{1}{3}}}{2}$$

$$y(x) = -\frac{3 \cdot 2^{\frac{1}{3}} (x^2)^{\frac{1}{3}}}{4} - \frac{3i\sqrt{3} \cdot 2^{\frac{1}{3}} (x^2)^{\frac{1}{3}}}{4}$$

$$y(x) = -\frac{3 \cdot 2^{\frac{1}{3}} (x^2)^{\frac{1}{3}}}{4} + \frac{3i\sqrt{3} \cdot 2^{\frac{1}{3}} (x^2)^{\frac{1}{3}}}{4}$$

$$y(x) = c_1 x + \frac{1}{c_1^2}$$



✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 69

```
DSolve[x*(y'[x])^3-y[x]*(y'[x])^2+1==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 x + \frac{1}{c_1^2}$$

$$y(x) \rightarrow 3 \left( -\frac{1}{2} \right)^{2/3} x^{2/3}$$

$$y(x) \rightarrow \frac{3x^{2/3}}{2^{2/3}}$$

$$y(x) \rightarrow -\frac{3\sqrt[3]{-1}x^{2/3}}{2^{2/3}}$$

### 3.16 problem 19

Internal problem ID [6056]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 19.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$y'^2 - y'x - y = 0$$

✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 77

```
dsolve(diff(y(x),x)^2-x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$\frac{c_1}{\sqrt{2x - 2\sqrt{x^2 + 4y(x)}}} + \frac{2x}{3} + \frac{\sqrt{x^2 + 4y(x)}}{3} = 0$$

$$\frac{c_1}{\sqrt{2x + 2\sqrt{x^2 + 4y(x)}}} + \frac{2x}{3} - \frac{\sqrt{x^2 + 4y(x)}}{3} = 0$$

✓ Solution by Mathematica

Time used: 60.18 (sec). Leaf size: 965

`DSolve[(y'[x])^2 - x*y'[x] - y[x] == 0, y[x], x, IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{\left(x^2 + \sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}}\right)^2 + 8e^{3c_1}x}{4\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}}}$$

$$y(x) \rightarrow \frac{1}{8} \left( 4x^2 + \frac{(-1 - i\sqrt{3})x(x^3 + 8e^{3c_1})}{\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}}} + i(\sqrt{3} + i)\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}} \right)$$

$$y(x) \rightarrow \frac{1}{8} \left( 4x^2 + \frac{i(\sqrt{3} + i)x(x^3 + 8e^{3c_1})}{\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}}} - (1 + i\sqrt{3})\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}} \right)$$

$$y(x) \rightarrow \frac{2\sqrt[3]{2}x^4 + 2^{2/3}\left(-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}(4x^3 + e^{3c_1})^3 + e^{6c_1}}\right)^{2/3} + 4x^2\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}(4x^3 + e^{3c_1})^3 + e^{6c_1}}}}{8\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}(4x^3 + e^{3c_1})^3 + e^{6c_1}}}}$$

$$y(x) \rightarrow \frac{1}{16} \left( 8x^2 - \frac{4\sqrt[3]{-2}x(x^3 - 2e^{3c_1})}{\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}(4x^3 + e^{3c_1})^3 + e^{6c_1}}}} + 2(-2)^{2/3}\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}(4x^3 + e^{3c_1})^3 + e^{6c_1}}} \right)$$

$$y(x) \rightarrow \frac{x^2}{2} + \frac{(-1)^{2/3}x(x^3 - 2e^{3c_1})}{2 \cdot 2^{2/3}\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}(4x^3 + e^{3c_1})^3 + e^{6c_1}}}} - \frac{1}{4}\sqrt[3]{-\frac{1}{2}\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}(4x^3 + e^{3c_1})^3 + e^{6c_1}}}}$$

### 3.17 problem 20

Internal problem ID [6057]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 20.

**ODE order:** 1.

**ODE degree:** 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$2y'^3 + y'x - 2y = 0$$

#### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 79

```
dsolve(2*diff(y(x),x)^3+x*diff(y(x),x)-2*y(x)=0,y(x), singsol=all)
```

$$y(x) = \left( -\frac{c_1}{12} - \frac{\sqrt{c_1^2 + 24x}}{12} \right)^3 + \frac{\left( -\frac{c_1}{12} - \frac{\sqrt{c_1^2 + 24x}}{12} \right) x}{2}$$

$$y(x) = \left( -\frac{c_1}{12} + \frac{\sqrt{c_1^2 + 24x}}{12} \right)^3 + \frac{\left( -\frac{c_1}{12} + \frac{\sqrt{c_1^2 + 24x}}{12} \right) x}{2}$$

#### ✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[2*(y'[x])^3+x*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Timed out

### 3.18 problem 21

Internal problem ID [6058]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 21.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$2y'^2 + y'x - 2y = 0$$

#### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 36

```
dsolve(2*diff(y(x),x)^2+x*diff(y(x),x)-2*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{2 \operatorname{LambertW}\left(\frac{x e^{\frac{c_1}{4}}}{4}\right) - \frac{c_1}{2}} + \frac{e^{\operatorname{LambertW}\left(\frac{x e^{\frac{c_1}{4}}}{4}\right) - \frac{c_1}{4}} x}{2}$$

#### ✓ Solution by Mathematica

Time used: 1.125 (sec). Leaf size: 130

```
DSolve[2*(y'[x])^2+x*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ \frac{\frac{1}{2}x\sqrt{x^2 + 16y(x)} - 8y(x) \log(\sqrt{x^2 + 16y(x)} - x) + \frac{x^2}{2}}{8y(x)} = c_1, y(x) \right]$$

$$\text{Solve} \left[ \frac{\frac{1}{2}x\sqrt{x^2 + 16y(x)} - 8y(x) \log(\sqrt{x^2 + 16y(x)} - x) - \frac{x^2}{2}}{8y(x)} + \log(y(x)) = c_1, y(x) \right]$$

$$y(x) \rightarrow 0$$

### 3.19 problem 22

Internal problem ID [6059]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 22.

**ODE order:** 1.

**ODE degree:** 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$y'^3 + 2y'x - y = 0$$

#### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 173

```
dsolve(diff(y(x),x)^3+2*x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{(-6\sqrt{x^2 + 3c_1} - 6x)^{\frac{3}{2}}}{27} - \frac{2\sqrt{-6\sqrt{x^2 + 3c_1} - 6x}x}{3}$$

$$y(x) = \frac{(-6\sqrt{x^2 + 3c_1} - 6x)^{\frac{3}{2}}}{27} + \frac{2\sqrt{-6\sqrt{x^2 + 3c_1} - 6x}x}{3}$$

$$y(x) = -\frac{(6\sqrt{x^2 + 3c_1} - 6x)^{\frac{3}{2}}}{27} - \frac{2\sqrt{6\sqrt{x^2 + 3c_1} - 6x}x}{3}$$

$$y(x) = \frac{(6\sqrt{x^2 + 3c_1} - 6x)^{\frac{3}{2}}}{27} + \frac{2\sqrt{6\sqrt{x^2 + 3c_1} - 6x}x}{3}$$

#### ✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(y'[x])^3+2*x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Timed out

### 3.20 problem 23

Internal problem ID [6060]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 23.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational, _dAlembert]`

$$4xy'^2 - 3yy' + 3 = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 153

```
dsolve(4*x*diff(y(x),x)^2-3*y(x)*diff(y(x),x)+3=0,y(x), singsol=all)
```

$$y(x) = -\frac{2\sqrt{x(3 + \sqrt{16c_1x + 9})}}{3} - \frac{2x}{\sqrt{x(3 + \sqrt{16c_1x + 9})}}$$

$$y(x) = \frac{2\sqrt{x(3 + \sqrt{16c_1x + 9})}}{3} + \frac{2x}{\sqrt{x(3 + \sqrt{16c_1x + 9})}}$$

$$y(x) = -\frac{2\sqrt{-x(-3 + \sqrt{16c_1x + 9})}}{3} - \frac{2x}{\sqrt{-x(-3 + \sqrt{16c_1x + 9})}}$$

$$y(x) = \frac{2\sqrt{-x(-3 + \sqrt{16c_1x + 9})}}{3} + \frac{2x}{\sqrt{-x(-3 + \sqrt{16c_1x + 9})}}$$

✓ Solution by Mathematica

Time used: 23.354 (sec). Leaf size: 187

`DSolve[4*x*(y'[x])^2-3*y[x]*y'[x]+3==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow -\frac{\sqrt{432x - e^{-\frac{c_1}{2}} (-144x + e^{c_1})^{3/2} + e^{c_1}}}{6\sqrt{3}}$$

$$y(x) \rightarrow \frac{\sqrt{432x - e^{-\frac{c_1}{2}} (-144x + e^{c_1})^{3/2} + e^{c_1}}}{6\sqrt{3}}$$

$$y(x) \rightarrow -\frac{\sqrt{432x + e^{-\frac{c_1}{2}} (-144x + e^{c_1})^{3/2} + e^{c_1}}}{6\sqrt{3}}$$

$$y(x) \rightarrow \frac{\sqrt{432x + e^{-\frac{c_1}{2}} (-144x + e^{c_1})^{3/2} + e^{c_1}}}{6\sqrt{3}}$$



### 3.21 problem 24

Internal problem ID [6061]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 24.

**ODE order:** 1.

**ODE degree:** 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$y'^3 - y'x + 2y = 0$$

#### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 83

```
dsolve(diff(y(x),x)^3-x*diff(y(x),x)+2*y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{\left(\frac{c_1}{6} - \frac{\sqrt{c_1^2 - 12x}}{6}\right)^3}{2} + \frac{\left(\frac{c_1}{6} - \frac{\sqrt{c_1^2 - 12x}}{6}\right)x}{2}$$

$$y(x) = -\frac{\left(\frac{c_1}{6} + \frac{\sqrt{c_1^2 - 12x}}{6}\right)^3}{2} + \frac{\left(\frac{c_1}{6} + \frac{\sqrt{c_1^2 - 12x}}{6}\right)x}{2}$$

#### ✓ Solution by Mathematica

Time used: 29.553 (sec). Leaf size: 10134

```
DSolve[(y'[x])^3-x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

### 3.22 problem 25

Internal problem ID [6062]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 25.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$5y'^2 + 6y'x - 2y = 0$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 85

```
dsolve(5*diff(y(x),x)^2+6*x*diff(y(x),x)-2*y(x)=0,y(x), singsol=all)
```

$$\frac{c_1}{\left(-15x - 5\sqrt{9x^2 + 10y(x)}\right)^{\frac{3}{2}}} + \frac{2x}{5} - \frac{\sqrt{9x^2 + 10y(x)}}{5} = 0$$

$$\frac{c_1}{\left(-15x + 5\sqrt{9x^2 + 10y(x)}\right)^{\frac{3}{2}}} + \frac{2x}{5} + \frac{\sqrt{9x^2 + 10y(x)}}{5} = 0$$

✓ Solution by Mathematica

Time used: 13.202 (sec). Leaf size: 771

`DSolve[5*(y'[x])^2+6*x*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \text{Root}\left[4\#1^5 + 4\#1^4x^2 + \#1^3x^4 + 1000\#1^2e^{5c_1}x + 900\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 25000e^{10c_1}\&, 1\right]$$

$$y(x) \rightarrow \text{Root}\left[4\#1^5 + 4\#1^4x^2 + \#1^3x^4 + 1000\#1^2e^{5c_1}x + 900\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 25000e^{10c_1}\&, 2\right]$$

$$y(x) \rightarrow \text{Root}\left[4\#1^5 + 4\#1^4x^2 + \#1^3x^4 + 1000\#1^2e^{5c_1}x + 900\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 25000e^{10c_1}\&, 3\right]$$

$$y(x) \rightarrow \text{Root}\left[4\#1^5 + 4\#1^4x^2 + \#1^3x^4 + 1000\#1^2e^{5c_1}x + 900\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 25000e^{10c_1}\&, 4\right]$$

$$y(x) \rightarrow \text{Root}\left[4\#1^5 + 4\#1^4x^2 + \#1^3x^4 + 1000\#1^2e^{5c_1}x + 900\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 25000e^{10c_1}\&, 5\right]$$

$$y(x) \rightarrow \text{Root}\left[100000\#1^5 + 100000\#1^4x^2 + 25000\#1^3x^4 - 1000\#1^2e^{5c_1}x - 900\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 1\right]$$

$$y(x) \rightarrow \text{Root}\left[100000\#1^5 + 100000\#1^4x^2 + 25000\#1^3x^4 - 1000\#1^2e^{5c_1}x - 900\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 2\right]$$

$$y(x) \rightarrow \text{Root}\left[100000\#1^5 + 100000\#1^4x^2 + 25000\#1^3x^4 - 1000\#1^2e^{5c_1}x - 900\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 3\right]$$

$$y(x) \rightarrow \text{Root}\left[100000\#1^5 + 100000\#1^4x^2 + 25000\#1^3x^4 - 1000\#1^2e^{5c_1}x - 900\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 4\right]$$

$$y(x) \rightarrow \text{Root}\left[100000\#1^5 + 100000\#1^4x^2 + 25000\#1^3x^4 - 1000\#1^2e^{5c_1}x - 900\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 5\right]$$

$$y(x) \rightarrow 0$$

### 3.23 problem 26

Internal problem ID [6063]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 26.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type [\_rational, \_dAlembert]

$$2xy'^2 + (-y + 2x)y' + 1 - y = 0$$

#### ✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 146

```
dsolve(2*x*diff(y(x),x)^2+(2*x-y(x))*diff(y(x),x)+1-y(x)=0,y(x), singsol=all)
```

$$y(x) = - \left( -2 \left( e^{\text{RootOf}(-e^{3-Z}x+2e^{2-Z}x+c_1e^{-Z}+Ze^{-Z}-xe^{-Z}+1)} - 1 \right)^2 \right. \\ \left. - 2e^{\text{RootOf}(-e^{3-Z}x+2e^{2-Z}x+c_1e^{-Z}+Ze^{-Z}-xe^{-Z}+1)} \right. \\ \left. + 2 \right) e^{-\text{RootOf}(-e^{3-Z}x+2e^{2-Z}x+c_1e^{-Z}+Ze^{-Z}-xe^{-Z}+1)} x \\ + e^{-\text{RootOf}(-e^{3-Z}x+2e^{2-Z}x+c_1e^{-Z}+Ze^{-Z}-xe^{-Z}+1)}$$

#### ✓ Solution by Mathematica

Time used: 1.344 (sec). Leaf size: 49

```
DSolve[2*x*(y'[x])^2+(2*x-y[x])*y'[x]+1-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ \left\{ x = \frac{\frac{1}{K[1]+1} + \log(K[1] + 1)}{K[1]^2} + \frac{c_1}{K[1]^2}, y(x) = 2xK[1] + \frac{1}{K[1] + 1} \right\}, \{y(x), K[1]\} \right]$$

### 3.24 problem 27

Internal problem ID [6064]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 27.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$5y'^2 + 3y'x - y = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 85

```
dsolve(5*diff(y(x),x)^2+3*x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$\frac{c_1}{\left(-30x - 10\sqrt{9x^2 + 20y(x)}\right)^{\frac{3}{2}}} + \frac{2x}{5} - \frac{\sqrt{9x^2 + 20y(x)}}{5} = 0$$

$$\frac{c_1}{\left(-30x + 10\sqrt{9x^2 + 20y(x)}\right)^{\frac{3}{2}}} + \frac{2x}{5} + \frac{\sqrt{9x^2 + 20y(x)}}{5} = 0$$

✓ Solution by Mathematica

Time used: 13.45 (sec). Leaf size: 771

`DSolve[5*(y'[x])^2+3*x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 8\#1^4x^2 + \#1^3x^4 + 4000\#1^2e^{5c_1}x + 1800\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 200000e^{10c_1}\&, 1\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 8\#1^4x^2 + \#1^3x^4 + 4000\#1^2e^{5c_1}x + 1800\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 200000e^{10c_1}\&, 2\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 8\#1^4x^2 + \#1^3x^4 + 4000\#1^2e^{5c_1}x + 1800\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 200000e^{10c_1}\&, 3\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 8\#1^4x^2 + \#1^3x^4 + 4000\#1^2e^{5c_1}x + 1800\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 200000e^{10c_1}\&, 4\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 8\#1^4x^2 + \#1^3x^4 + 4000\#1^2e^{5c_1}x + 1800\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 200000e^{10c_1}\&, 5\right]$$

$$y(x) \rightarrow \text{Root}\left[3200000\#1^5 + 1600000\#1^4x^2 + 200000\#1^3x^4 - 4000\#1^2e^{5c_1}x - 1800\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 1\right]$$

$$y(x) \rightarrow \text{Root}\left[3200000\#1^5 + 1600000\#1^4x^2 + 200000\#1^3x^4 - 4000\#1^2e^{5c_1}x - 1800\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 2\right]$$

$$y(x) \rightarrow \text{Root}\left[3200000\#1^5 + 1600000\#1^4x^2 + 200000\#1^3x^4 - 4000\#1^2e^{5c_1}x - 1800\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 3\right]$$

$$y(x) \rightarrow \text{Root}\left[3200000\#1^5 + 1600000\#1^4x^2 + 200000\#1^3x^4 - 4000\#1^2e^{5c_1}x - 1800\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 4\right]$$

$$y(x) \rightarrow \text{Root}\left[3200000\#1^5 + 1600000\#1^4x^2 + 200000\#1^3x^4 - 4000\#1^2e^{5c_1}x - 1800\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 5\right]$$

$$y(x) \rightarrow 0$$

### 3.25 problem 28

Internal problem ID [6065]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

**Problem number:** 28.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$y'^2 + 3y'x - y = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 85

```
dsolve(diff(y(x),x)^2+3*x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$\frac{c_1}{\left(-6x - 2\sqrt{9x^2 + 4y(x)}\right)^{\frac{3}{2}}} + \frac{2x}{5} - \frac{\sqrt{9x^2 + 4y(x)}}{5} = 0$$

$$\frac{c_1}{\left(-6x + 2\sqrt{9x^2 + 4y(x)}\right)^{\frac{3}{2}}} + \frac{2x}{5} + \frac{\sqrt{9x^2 + 4y(x)}}{5} = 0$$

✓ Solution by Mathematica

Time used: 13.436 (sec). Leaf size: 776

```
DSolve[(y'[x])^2+3*x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 40\#1^4x^2 + 25\#1^3x^4 + 160\#1^2e^{5c_1}x + 360\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 64e^{10c_1}\&, 1\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 40\#1^4x^2 + 25\#1^3x^4 + 160\#1^2e^{5c_1}x + 360\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 64e^{10c_1}\&, 2\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 40\#1^4x^2 + 25\#1^3x^4 + 160\#1^2e^{5c_1}x + 360\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 64e^{10c_1}\&, 3\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 40\#1^4x^2 + 25\#1^3x^4 + 160\#1^2e^{5c_1}x + 360\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 64e^{10c_1}\&, 4\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 40\#1^4x^2 + 25\#1^3x^4 + 160\#1^2e^{5c_1}x + 360\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 64e^{10c_1}\&, 5\right]$$

$$y(x) \rightarrow \text{Root}\left[1024\#1^5 + 2560\#1^4x^2 + 1600\#1^3x^4 - 160\#1^2e^{5c_1}x - 360\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 1\right]$$

$$y(x) \rightarrow \text{Root}\left[1024\#1^5 + 2560\#1^4x^2 + 1600\#1^3x^4 - 160\#1^2e^{5c_1}x - 360\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 2\right]$$

$$y(x) \rightarrow \text{Root}\left[1024\#1^5 + 2560\#1^4x^2 + 1600\#1^3x^4 - 160\#1^2e^{5c_1}x - 360\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 3\right]$$

$$y(x) \rightarrow \text{Root}\left[1024\#1^5 + 2560\#1^4x^2 + 1600\#1^3x^4 - 160\#1^2e^{5c_1}x - 360\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 4\right]$$

$$y(x) \rightarrow \text{Root}\left[1024\#1^5 + 2560\#1^4x^2 + 1600\#1^3x^4 - 160\#1^2e^{5c_1}x - 360\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 5\right]$$

$$y(x) \rightarrow 0$$



### 3.26 problem 29

Internal problem ID [6066]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320


**Problem number:** 29.

**ODE order:** 1.

**ODE degree:** 2.

CAS Maple gives this as type `[_homogeneous, 'class G'], _rational]`

$$y - y'x - x^3y'^2 = 0$$

 Solution by Maple

```
dsolve(y(x)=x*diff(y(x),x)+x^3*diff(y(x),x)^2,y(x), singsol=all)
```

No solution found

 Solution by Mathematica

Time used: 105.583 (sec). Leaf size: 7052

```
DSolve[y[x]==x*y'[x]+x^3*(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

**4 CHAPTER 16. Nonlinear equations. Section 101.  
Independent variable missing. EXERCISES Page  
324**

4.1	problem 1	83
4.2	problem 2	84
4.3	problem 3	85
4.4	problem 4	86
4.5	problem 5	87
4.6	problem 6	88
4.7	problem 7	89
4.8	problem 9	90
4.9	problem 10	91
4.10	problem 11	92
4.11	problem 12	95
4.12	problem 13	96
4.13	problem 14	97
4.14	problem 15	98
4.15	problem 16	99
4.16	problem 17	100
4.17	problem 18	101
4.18	problem 19	102
4.19	problem 20	103
4.20	problem 21	104
4.21	problem 23	105
4.22	problem 24	106
4.23	problem 25	107
4.24	problem 26	108
4.25	problem 27	110
4.26	problem 28	111
4.27	problem 30	112
4.28	problem 31	113
4.29	problem 32	114
4.30	problem 33	115
4.31	problem 34	117
4.32	problem 35	118
4.33	problem 36	119
4.34	problem 37	120
4.35	problem 38	121
4.36	problem 39	122
4.37	problem 40	123

4.38 problem 41	124
4.39 problem 42	126
4.40 problem 43	127

## 4.1 problem 1

Internal problem ID [6067]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 1.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]`

$$y'' - xy'^3 = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 37

```
dsolve(diff(y(x),x$2)=x*(diff(y(x),x))^3,y(x), singsol=all)
```

$$y(x) = \arctan\left(\frac{x}{\sqrt{-x^2 + c_1}}\right) + c_2$$

$$y(x) = -\arctan\left(\frac{x}{\sqrt{-x^2 + c_1}}\right) + c_2$$

### ✓ Solution by Mathematica

Time used: 10.872 (sec). Leaf size: 57

```
DSolve[y'[x]==x*(y'[x])^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \arctan\left(\frac{x}{\sqrt{-x^2 - 2c_1}}\right)$$

$$y(x) \rightarrow \arctan\left(\frac{x}{\sqrt{-x^2 - 2c_1}}\right) + c_2$$

$$y(x) \rightarrow c_2$$

## 4.2 problem 2

Internal problem ID [6068]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 2.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^2 y'' + y'^2 - 2y'x = 0$$

With initial conditions

$$[y(2) = 5, y'(2) = -4]$$

### ✓ Solution by Maple

Time used: 0.171 (sec). Leaf size: 24

```
dsolve([x^2*diff(y(x),x$2)+diff(y(x),x)^2-2*x*diff(y(x),x)=0,y(2) = 5, D(y)(2) = -4],y(x), si
```

$$y(x) = \frac{x^2}{2} + 3x + 9 \ln(x - 3) - 3 - 9i\pi$$

### ✓ Solution by Mathematica

Time used: 0.49 (sec). Leaf size: 26

```
DSolve[{x^2*y'[x]+(y'[x])^2-2*x*y'[x]==0,{y[2]==5,y'[2]==-4}},y[x],x,IncludeSingularSolution
```

$$y(x) \rightarrow \frac{1}{2}x(x + 6) + 9 \log(x - 3) - 9i\pi - 3$$

### 4.3 problem 3

Internal problem ID [6069]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 3.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^2 y'' + y'^2 - 2y'x = 0$$

With initial conditions

$$[y(2) = 5, y'(2) = 2]$$

#### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 11

```
dsolve([x^2*diff(y(x),x$2)+diff(y(x),x)^2-2*x*diff(y(x),x)=0,y(2) = 5, D(y)(2) = 2],y(x), sin
```

$$y(x) = \frac{x^2}{2} + 3$$

#### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 14

```
DSolve[{x^2*y'[x]+(y'[x])^2-2*x*y'[x]==0,{y[2]==5,y'[2]==2}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{1}{2}(x^2 + 6)$$

## 4.4 problem 4

Internal problem ID [6070]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 4.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear], _L`

$$yy'' + y'^2 = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 33

```
dsolve(y(x)*diff(y(x),x$2)+diff(y(x),x)^2=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = \sqrt{2c_1x + 2c_2}$$

$$y(x) = -\sqrt{2c_1x + 2c_2}$$

### ✓ Solution by Mathematica

Time used: 0.067 (sec). Leaf size: 20

```
DSolve[y[x]*y'[x]+(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2\sqrt{2x - c_1}$$

## 4.5 problem 5

Internal problem ID [6071]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 5.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]`,

$$y^2 y'' + y'^3 = 0$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 34

```
dsolve(y(x)^2*diff(y(x),x$2)+diff(y(x),x)^3=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1$$

$$y(x) = e^{-\text{LambertW}(-c_1 e^{-c_2} e^{-x}) - c_2 - x}$$

### ✓ Solution by Mathematica

Time used: 0.086 (sec). Leaf size: 37

```
DSolve[y[x]^2*y'[x]+(y'[x])^3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 \left( 1 + \frac{1}{\text{InverseFunction} \left[ -\frac{1}{\#1} - \log(\#1) + \log(\#1 + 1) \& \right] [-x + c_1]} \right)$$



## 4.6 problem 6

Internal problem ID [6072]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 6.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible,`

$$(1 + y)y'' - y'^2 = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve((y(x)+1)*diff(y(x),x$2)=diff(y(x),x)^2,y(x), singsol=all)
```

$$y(x) = -1$$

$$y(x) = e^{c_1 x} c_2 - 1$$

### ✓ Solution by Mathematica

Time used: 0.088 (sec). Leaf size: 21

```
DSolve[(y[x]+1)*y'[x]==(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -1 + \frac{e^{c_1(x+c_2)}}{c_1}$$

## 4.7 problem 7

Internal problem ID [6073]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 7.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_y_y1]`

$$2ay'' + y'^3 = 0$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 29

```
dsolve(2*a*diff(y(x),x$2)+diff(y(x),x)^3=0,y(x), singsol=all)
```

$$y(x) = 2\sqrt{(x + c_1)a} + c_2$$

$$y(x) = -2\sqrt{(x + c_1)a} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.303 (sec). Leaf size: 51

```
DSolve[2*a*y'[x]+(y'[x])^3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - 2\sqrt{a}\sqrt{x - 2ac_1}$$

$$y(x) \rightarrow 2\sqrt{a}\sqrt{x - 2ac_1} + c_2$$

## 4.8 problem 9

Internal problem ID [6074]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 9.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' - y' - x^5 = 0$$

With initial conditions

$$\left[ y(1) = \frac{1}{2}, y'(1) = 1 \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve([x*dif(y(x),x$2)=dif(y(x),x)+x^5,y(1) = 1/2, D(y)(1) = 1],y(x), singsol=all)
```

$$y(x) = \frac{1}{24}x^6 + \frac{3}{8}x^2 + \frac{1}{12}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 19

```
DSolve[{x*y'[x]==y'[x]+x^5,{y[1]==1/2,y'[1]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{24}(x^6 + 9x^2 + 2)$$

## 4.9 problem 10

Internal problem ID [6075]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 10.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' + y' + x = 0$$

With initial conditions

$$\left[ y(2) = -1, y'(2) = -\frac{1}{2} \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve([x*diff(y(x),x$2)+diff(y(x),x)+x=0,y(2) = -1, D(y)(2) = -1/2],y(x), singsol=all)
```

$$y(x) = -\frac{x^2}{4} + \ln(x) - \ln(2)$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 19

```
DSolve[{x*y''[x]+y'[x]+x==0,{y[2]==-1,y'[2]==-1/2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log\left(\frac{x}{2}\right) - \frac{x^2}{4}$$

## 4.10 problem 11

Internal problem ID [6076]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 11.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]`,

$$y'' - 2yy'^3 = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 386

```
dsolve(diff(y(x),x$2)=2*y(x)*diff(y(x),x)^3,y(x), singsol=all)
```

$$y(x) = c_1$$

$$y(x) = \frac{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}{2} + \frac{2c_1}{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}$$

$$y(x) = -\frac{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}{4c_1} - \frac{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}{2} + \frac{i\sqrt{3} \left( \frac{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}{2} - \frac{2c_1}{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}} \right)}{2}$$

$$y(x) = -\frac{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}{4c_1} - \frac{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}{2} + \frac{i\sqrt{3} \left( \frac{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}{2} - \frac{2c_1}{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}} \right)}{2}$$

✓ Solution by Mathematica

Time used: 0.299 (sec). Leaf size: 346

```
DSolve[y'[x]==2*y[x]*(y'[x])^3,y[x],x,IncludeSingularSolutions -> True]
```

$y(x)$

$$\rightarrow \frac{\sqrt[3]{2}c_1}{\sqrt[3]{\sqrt{9x^2 + 18c_2x + 4c_1^3 + 9c_2^2} + 3x + 3c_2}} - \frac{\sqrt[3]{\sqrt{9x^2 + 18c_2x + 4c_1^3 + 9c_2^2} + 3x + 3c_2}}{\sqrt[3]{2}}$$

$$y(x) \rightarrow \frac{2^{2/3}(1 - i\sqrt{3}) (\sqrt{9x^2 + 18c_2x + 4c_1^3 + 9c_2^2} + 3x + 3c_2)^{2/3} + \sqrt[3]{2}(-2 - 2i\sqrt{3}) c_1}{4\sqrt[3]{\sqrt{9x^2 + 18c_2x + 4c_1^3 + 9c_2^2} + 3x + 3c_2}}$$

$$y(x) \rightarrow \frac{2^{2/3}(1 + i\sqrt{3}) (\sqrt{9x^2 + 18c_2x + 4c_1^3 + 9c_2^2} + 3x + 3c_2)^{2/3} + 2i\sqrt[3]{2}(\sqrt{3} + i) c_1}{4\sqrt[3]{\sqrt{9x^2 + 18c_2x + 4c_1^3 + 9c_2^2} + 3x + 3c_2}}$$

## 4.11 problem 12

Internal problem ID [6077]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 12.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]`,

$$yy'' + y'^3 - y'^2 = 0$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 44

```
dsolve(y(x)*diff(y(x),x$2)+diff(y(x),x)^3-diff(y(x),x)^2=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1$$

$$y(x) = e^{-\frac{c_1 \operatorname{LambertW}\left(\frac{e^{\frac{c_2}{c_1}} e^{\frac{x}{c_1}}}{c_1}\right) - c_2 - x}{c_1}}$$

### ✓ Solution by Mathematica

Time used: 0.108 (sec). Leaf size: 29

```
DSolve[y[x]*y'[x]+(y'[x])^3-(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{c_1 W\left(e^{e^{-c_1}(x+c_2)-c_1}\right)}$$



## 4.12 problem 13

Internal problem ID [6078]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 13.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + \beta^2 y = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+beta^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(\beta x) + c_2 \cos(\beta x)$$

### ✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 20

```
DSolve[y''[x]+\[Beta]^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(\beta x) + c_2 \sin(\beta x)$$

### 4.13 problem 14

Internal problem ID [6079]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 14.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]`,

$$yy'' + y'^3 = 0$$

#### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 27

```
dsolve(y(x)*diff(y(x),x$2)+diff(y(x),x)^3=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1$$

$$y(x) = e^{\text{LambertW}((c_2+x)e^{c_1}e^{-1})-c_1+1}$$

#### ✓ Solution by Mathematica

Time used: 0.093 (sec). Leaf size: 25

```
DSolve[y[x]*y'[x]+(y'[x])^3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{W(e^{-1-c_1}(x+c_2))+1+c_1}$$

## 4.14 problem 15

Internal problem ID [6080]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 15.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' \cos(x) - y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$2)*cos(x)=diff(y(x),x),y(x), singsol=all)
```

$$y(x) = c_1 + (\ln(\sec(x) + \tan(x)) - \ln(\cos(x)))c_2$$

### ✓ Solution by Mathematica

Time used: 0.158 (sec). Leaf size: 25

```
DSolve[y''[x]*Cos[x]==y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \log\left(e^{4\operatorname{arctanh}\left(\tan\left(\frac{x}{2}\right)\right)} + 1\right) + c_2$$

## 4.15 problem 16

Internal problem ID [6081]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 16.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]`

$$y'' - xy'^2 = 0$$

With initial conditions

$$\left[ y(2) = \frac{\pi}{4}, y'(2) = -\frac{1}{4} \right]$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 8

```
dsolve([diff(y(x),x$2)=x*diff(y(x),x)^2,y(2) = 1/4*Pi, D(y)(2) = -1/4],y(x), singsol=all)
```

$$y(x) = \operatorname{arccot}\left(\frac{x}{2}\right)$$

### ✓ Solution by Mathematica

Time used: 1.165 (sec). Leaf size: 19

```
DSolve[{y'[x]==x*(y'[x])^2,{y[2]==1/4*Pi,y'[2]==-1/4}},y[x],x,IncludeSingularSolutions -> Tr
```

$$y(x) \rightarrow \frac{1}{2} \left( \pi - 2 \arctan\left(\frac{x}{2}\right) \right)$$

## 4.16 problem 17

Internal problem ID [6082]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 17.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]`

$$y'' - xy'^2 = 0$$

With initial conditions

$$\left[ y(0) = 1, y'(0) = \frac{1}{2} \right]$$

### ✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 10

```
dsolve([diff(y(x),x$2)=x*diff(y(x),x)^2,y(0) = 1, D(y)(0) = 1/2],y(x), singsol=all)
```

$$y(x) = \operatorname{arctanh}\left(\frac{x}{2}\right) + 1$$

### ✓ Solution by Mathematica

Time used: 0.217 (sec). Leaf size: 13

```
DSolve[{y'[x]==x*(y'[x])^2,{y[0]==1,y'[0]==1/2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \operatorname{arctanh}\left(\frac{x}{2}\right) + 1$$

## 4.17 problem 18

Internal problem ID [6083]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 18.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]`

$$y'' + e^{-2y} = 0$$

With initial conditions

$$[y(3) = 0, y'(3) = 1]$$

### ✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 12

```
dsolve([diff(y(x),x$2)=-exp(-2*y(x)),y(3) = 0, D(y)(3) = 1],y(x), singsol=all)
```

$$y(x) = \frac{\ln((x-2)^2)}{2}$$

### ✓ Solution by Mathematica

Time used: 0.147 (sec). Leaf size: 9

```
DSolve[{y'[x]==-Exp[-2*y[x]],{y[3]==0,y'[3]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(x-2)$$

## 4.18 problem 19

Internal problem ID [6084]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 19.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]`

$$y'' + e^{-2y} = 0$$

With initial conditions

$$[y(3) = 0, y'(3) = -1]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve([diff(y(x),x$2)=-exp(-2*y(x)),y(3) = 0, D(y)(3) = -1],y(x), singsol=all)
```

$$y(x) = \frac{\ln((x-4)^2)}{2}$$

### ✓ Solution by Mathematica

Time used: 0.133 (sec). Leaf size: 11

```
DSolve[{y'[x]==-Exp[-2*y[x]],{y[3]==0,y'[3]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(4 - x)$$

## 4.19 problem 20

Internal problem ID [6085]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 20.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]`

$$2y'' - \sin(2y) = 0$$

With initial conditions

$$\left[ y(0) = \frac{\pi}{2}, y'(0) = 1 \right]$$

### ✓ Solution by Maple

Time used: 134.11 (sec). Leaf size: 1495

```
dsolve([2*diff(y(x),x$2)=sin(2*y(x)),y(0) = 1/2*Pi, D(y)(0) = 1],y(x), singsol=all)
```

Expression too large to display

### ✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{2*y'[x]==Sin[2*y[x]],{y[0]==Pi/2,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

{}



## 4.20 problem 21

Internal problem ID [6086]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 21.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]`

$$2y'' - \sin(2y) = 0$$

With initial conditions

$$\left[ y(0) = -\frac{\pi}{2}, y'(0) = 1 \right]$$

### ✓ Solution by Maple

Time used: 98.39 (sec). Leaf size: 1490

```
dsolve([2*diff(y(x),x$2)=sin(2*y(x)),y(0) = -1/2*Pi, D(y)(0) = 1],y(x), singsol=all)
```

Expression too large to display

### ✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{2*y'[x]==Sin[2*y[x]],{y[0]==-Pi/2,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True
```

```
{}
```

## 4.21 problem 23

Internal problem ID [6087]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 23.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^3 y'' - x^2 y' + x^2 - 3 = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(x^3*diff(y(x),x$2)-x^2*diff(y(x),x)=3-x^2,y(x), singsol=all)
```

$$y(x) = \frac{c_1 x^2}{2} + \frac{1}{x} + x + c_2$$

### ✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 21

```
DSolve[x^3*y'[x]-x^2*y'[x]==3-x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1 x^2}{2} + x + \frac{1}{x} + c_2$$

## 4.22 problem 24

Internal problem ID [6088]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 24.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible,`

$$y'' - y'^2 = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(diff(y(x),x$2)=diff(y(x),x)^2,y(x), singsol=all)
```

$$y(x) = -\ln(-c_1x - c_2)$$

### ✓ Solution by Mathematica

Time used: 0.186 (sec). Leaf size: 15

```
DSolve[y''[x]==(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \log(x + c_1)$$

## 4.23 problem 25

Internal problem ID [6089]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 25.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' - e^x y'^2 = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)=exp(x)*diff(y(x),x)^2,y(x), singsol=all)
```

$$y(x) = \frac{\ln(e^x)}{c_1} - \frac{\ln(e^x - c_1)}{c_1} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.902 (sec). Leaf size: 37

```
DSolve[y''[x]==Exp[x](y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-x + \log(e^x + c_1) + c_1 c_2}{c_1}$$

$$y(x) \rightarrow \text{Indeterminate}$$

$$y(x) \rightarrow c_2$$

## 4.24 problem 26

Internal problem ID [6090]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 26.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]`

$$2y'' - y'^3 \sin(2x) = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 78

```
dsolve(2*dif(y(x),x$2)=dif(y(x),x)^3*sin(2*x),y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{-\left(\sin(x)^2 - \frac{1}{c_1^2}\right) c_1^2 \operatorname{InverseJacobiAM}(x, c_1)}}{\sqrt{-\sin(x)^2 + \frac{1}{c_1^2}}} + c_2$$

$$y(x) = -\frac{\sqrt{-\left(\sin(x)^2 - \frac{1}{c_1^2}\right) c_1^2 \operatorname{InverseJacobiAM}(x, c_1)}}{\sqrt{-\sin(x)^2 + \frac{1}{c_1^2}}} + c_2$$

✓ Solution by Mathematica

Time used: 5.916 (sec). Leaf size: 118

```
DSolve[2*y'[x]==(y'[x])^3*Sin[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \frac{\sqrt{\frac{\cos(2x)+1-4c_1}{1-2c_1}} \operatorname{EllipticF}\left(x, \frac{1}{1-2c_1}\right)}{\sqrt{\cos(2x)+1-4c_1}}$$

$$y(x) \rightarrow \frac{\sqrt{\frac{\cos(2x)+1-4c_1}{1-2c_1}} \operatorname{EllipticF}\left(x, \frac{1}{1-2c_1}\right)}{\sqrt{\cos(2x)+1-4c_1}} + c_2$$

$$y(x) \rightarrow c_2$$

## 4.25 problem 27

Internal problem ID [6091]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 27.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]`

$$x^2 y'' + y'^2 = 0$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 21

```
dsolve(x^2*diff(y(x),x$2)+diff(y(x),x)^2=0,y(x), singsol=all)
```

$$y(x) = \frac{x}{c_1} + \frac{\ln(c_1 x - 1)}{c_1^2} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.533 (sec). Leaf size: 47

```
DSolve[x^2*y'[x]+(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x}{c_1} + \frac{\log(1 + c_1 x)}{c_1^2} + c_2$$

$$y(x) \rightarrow c_2$$

$$y(x) \rightarrow -\frac{x^2}{2} + c_2$$

## 4.26 problem 28

Internal problem ID [6092]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 28.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_xy]`

$$y'' - 1 - y'^2 = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$2)=1+diff(y(x),x)^2,y(x), singsol=all)
```

$$y(x) = -\ln\left(\frac{-c_2 + \tan(x) c_1}{\sec(x)}\right)$$

### ✓ Solution by Mathematica

Time used: 1.769 (sec). Leaf size: 16

```
DSolve[y' '[x]==1+(y' [x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \log(\cos(x + c_1))$$



## 4.27 problem 30

Internal problem ID [6093]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 30.

**ODE order:** 2.

**ODE degree:** 2.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - (1 + y'^2)^{\frac{3}{2}} = 0$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 51

```
dsolve(diff(y(x),x$2)=(1+diff(y(x),x)^2)^(3/2),y(x), singsol=all)
```

$$y(x) = -ix + c_1$$

$$y(x) = ix + c_1$$

$$y(x) = (c_1 + x + 1)(c_1 + x - 1) \sqrt{-\frac{1}{c_1^2 + 2c_1x + x^2 - 1}} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.258 (sec). Leaf size: 53

```
DSolve[y' '[x]==(1+(y' [x])^2)^(3/2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - i\sqrt{(x-1+c_1)(x+1+c_1)}$$

$$y(x) \rightarrow i\sqrt{(x-1+c_1)(x+1+c_1)} + c_2$$

## 4.28 problem 31

Internal problem ID [6094]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 31.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_y_y1]`

$$yy'' - y'^2(1 - y' \sin(y) - yy' \cos(y)) = 0$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 24

```
dsolve(y(x)*diff(y(x),x$2)=diff(y(x),x)^2*(1-diff(y(x),x)*sin(y(x))-y(x)*diff(y(x),x)*cos(y(x))),y(x),x)
```

$$y(x) = c_1$$

$$-\cos(y(x)) + c_1 \ln(y(x)) - x - c_2 = 0$$

### ✓ Solution by Mathematica

Time used: 0.181 (sec). Leaf size: 23

```
DSolve[y[x]*y'[x]==(y'[x])^2*(1-y'[x]*Sin[y[x]]-y[x]*y'[x]*Cos[y[x]]),y[x],x,IncludeSingular
```

$$y(x) \rightarrow \text{InverseFunction}[-\cos(\#1) + c_1 \log(\#1)\&][x + c_2]$$

## 4.29 problem 32

Internal problem ID [6095]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 32.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$(1 + y^2) y'' + y'^3 + y' = 0$$

### ✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 130

```
dsolve((1+y(x)^2)*diff(y(x),x$2)+diff(y(x),x)^3+diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -i$$

$$y(x) = i$$

$$y(x) = c_1$$

$$y(x) = \frac{i(c_1 - 1)}{1 + c_1}$$

$$\frac{-c_1^2 + 2c_1 + c_1^2 c_2 + x c_1^2 - 1 + 4 \operatorname{LambertW}\left(-\frac{i e^{-\frac{c_1 c_2}{4}} e^{-\frac{c_1 x}{4}} e^{\frac{c_1}{4}} e^{-\frac{c_2}{2}} e^{-\frac{x}{2}} e^{-\frac{1}{2}} e^{-\frac{c_2}{4 c_1}} e^{-\frac{x}{4 c_1}} e^{\frac{1}{4 c_1}} (c_1 - 1)}{4 c_1}\right) c_1 + 2 c_1 c_2 + 2 c_1 x + c_2 + x}{e^{4 c_1}}}{1 + c_1}$$

### ✓ Solution by Mathematica

Time used: 0.194 (sec). Leaf size: 42

```
DSolve[(1+y[x]^2)*y'[x]+(y'[x])^3+y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \csc(c_1) \sec(c_1) W\left(\sin(c_1) e^{-((x+c_2) \cos^2(c_1) - \sin^2(c_1))}\right) + \tan(c_1)$$

### 4.30 problem 33

Internal problem ID [6096]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 33.

**ODE order:** 2.

**ODE degree:** 2.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$\left( yy'' + 1 + y'^2 \right)^2 - \left( 1 + y'^2 \right)^3 = 0$$

#### ✓ Solution by Maple

Time used: 0.14 (sec). Leaf size: 131

```
dsolve((y(x)*diff(y(x),x$2)+1+diff(y(x),x)^2)^2=(1+diff(y(x),x)^2)^3,y(x), singsol=all)
```

$$y(x) = -ix + c_1$$

$$y(x) = ix + c_1$$

$$y(x) = 0$$

$$y(x) = -c_1 - \sqrt{c_1^2 - c_2^2 - 2xc_2 - x^2}$$

$$y(x) = -c_1 + \sqrt{c_1^2 - c_2^2 - 2xc_2 - x^2}$$

$$y(x) = c_1 - \sqrt{c_1^2 - c_2^2 - 2xc_2 - x^2}$$

$$y(x) = c_1 + \sqrt{c_1^2 - c_2^2 - 2xc_2 - x^2}$$

✓ Solution by Mathematica

Time used: 6.299 (sec). Leaf size: 121

```
DSolve[(y[x]*y'[x]+1+(y'[x])^2)^2==(1+(y'[x])^2)^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{e^{2c_1} - (x + c_2)^2} - e^{c_1}$$

$$y(x) \rightarrow e^{c_1} - \sqrt{e^{2c_1} - (x + c_2)^2}$$

$$y(x) \rightarrow \sqrt{e^{2c_1} - (x + c_2)^2} - e^{c_1}$$

$$y(x) \rightarrow \sqrt{e^{2c_1} - (x + c_2)^2} + e^{c_1}$$

### 4.31 problem 34

Internal problem ID [6097]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 34.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^2 y'' - y'(2x - y') = 0$$

With initial conditions

$$[y(-1) = 5, y'(-1) = 1]$$

#### ✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 20

```
dsolve([x^2*diff(y(x),x$2)=diff(y(x),x)*(2*x-diff(y(x),x)),y(-1) = 5, D(y)(-1) = 1],y(x), sin
```

$$y(x) = \frac{x^2}{2} - 2x + 4 \ln(x + 2) + \frac{5}{2}$$

#### ✓ Solution by Mathematica

Time used: 0.49 (sec). Leaf size: 22

```
DSolve[{x^2*y''[x]==y'[x]*(2*x-y'[x]),{y[-1]==5,y'[-1]==1}},y[x],x,IncludeSingularSolutions -
```

$$y(x) \rightarrow \frac{1}{2}((x - 4)x + 8 \log(x + 2) + 5)$$

## 4.32 problem 35

Internal problem ID [6098]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 35.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^2 y'' - y'(3x - 2y') = 0$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 22

```
dsolve(x^2*diff(y(x),x$2)=diff(y(x),x)*(3*x-2*diff(y(x),x)),y(x), singsol=all)
```

$$y(x) = \frac{x^2}{2} + \frac{c_1 \ln(x^2 - c_1)}{2} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.327 (sec). Leaf size: 28

```
DSolve[x^2*y''[x]==y'[x]*(3*x-2*y'[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(x^2 - c_1 \log(x^2 + c_1) + 2c_2)$$

### 4.33 problem 36

Internal problem ID [6099]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 36.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_y], _Liouville, [_2nd_order, _reducible,`

$$xy'' - y'(2 - 3y'x) = 0$$

#### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve(x*diff(y(x),x$2)=diff(y(x),x)*(2-3*x*diff(y(x),x)),y(x), singsol=all)
```

$$y(x) = \frac{\ln(c_1 x^3 + 3c_2)}{3}$$

#### ✓ Solution by Mathematica

Time used: 0.255 (sec). Leaf size: 19

```
DSolve[x*y'[x]==y'[x]*(2-3*x*y'[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} \log(x^3 + c_1) + c_2$$



### 4.34 problem 37

Internal problem ID [6100]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 37.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^4 y'' - y'(y' + x^3) = 0$$

With initial conditions

$$[y(1) = 2, y'(1) = 1]$$

#### ✓ Solution by Maple

Time used: 0.172 (sec). Leaf size: 25

```
dsolve([x^4*diff(y(x),x$2)=diff(y(x),x)*(diff(y(x),x)+x^3),y(1) = 2, D(y)(1) = 1],y(x), sings
```

$$y(x) = x^2 - \ln(-x^2 - 1) + 1 + \ln(2) + i\pi$$

#### ✓ Solution by Mathematica

Time used: 0.891 (sec). Leaf size: 20

```
DSolve[{x^4*y''[x]==y'[x]*(y'[x]+x^3)},{y[1]==2,y'[1]==1},y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow x^2 - \log(x^2 + 1) + 1 + \log(2)$$

### 4.35 problem 38

Internal problem ID [6101]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 38.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_xy]`

$$y'' - 2x - (x^2 - y')^2 = 0$$

#### ✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$2)=2*x+(x^2-diff(y(x),x))^2,y(x), singsol=all)
```

$$y(x) = \frac{x^3}{3} - \ln(xc_2 - c_1)$$

#### ✓ Solution by Mathematica

Time used: 0.312 (sec). Leaf size: 24

```
DSolve[y'[x]==2*x+(x^2-y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{3} - \log(-x + c_1) + c_2$$

### 4.36 problem 39

Internal problem ID [6102]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 39.

**ODE order:** 2.

**ODE degree:** 2.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y''^2 - 2y'' + y'^2 - 2xy' + x^2 = 0$$

With initial conditions

$$\left[ y(0) = \frac{1}{2}, y'(0) = 1 \right]$$

#### ✓ Solution by Maple

Time used: 0.313 (sec). Leaf size: 23

```
dsolve([diff(y(x),x$2)^2-2*diff(y(x),x$2)+diff(y(x),x)^2-2*x*diff(y(x),x)+x^2=0,y(0) = 1/2, D
```

$$y(x) = \frac{(x+1)^2}{2}$$

$$y(x) = \frac{x^2}{2} + \sin(x) + \frac{1}{2}$$

#### ✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{(y'[x])^2-2*y'[x]+(y'[x])^2-2*x*y'[x]+x^2==0,{y[0]==1/2,y'[0]==1}},y[x],x,IncludeSi
```

Not solved

### 4.37 problem 40

Internal problem ID [6103]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 40.

**ODE order:** 2.

**ODE degree:** 2.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y''^2 - xy'' + y' = 0$$

#### ✓ Solution by Maple

Time used: 0.11 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)^2-x*diff(y(x),x$2)+diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x^3}{12} + c_1$$

$$y(x) = \frac{1}{2}c_1x^2 - xc_1^2 + c_2$$

#### ✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 21

```
DSolve[(y'[x])^2-x*y''[x]+y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}c_1x(x - 2c_1) + c_2$$

**4.38 problem 41**

Internal problem ID [6104]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillian Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 41.

**ODE order:** 2.

**ODE degree:** 3.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y''^3 - 12y'(xy'' - 2y') = 0$$

✓ Solution by Maple

Time used: 0.344 (sec). Leaf size: 174

```
dsolve(diff(y(x),x$2)^3=12*diff(y(x),x)*(x*diff(y(x),x$2)-2*diff(y(x),x)),y(x), singsol=all)
```

$$y(x) = \frac{x^4}{9} + c_1$$

$$y(x) = c_1$$

$$y(x) = \int \text{RootOf} \left( -6 \ln(x) - \left( \int -z \frac{3-f \sqrt{\frac{1}{-f(9-f-4)}} 2^{\frac{1}{3}} \left( \left( 3 \sqrt{\frac{1}{-f(9-f-4)}} -f + 1 \right)^2 (9-f-4)^4 \right)^{\frac{1}{3}} - 2 2^{\frac{2}{3}} \left( \left( 3 \sqrt{\frac{1}{-f(9-f-4)}} -f + 1 \right) \right)}{-f(9-f-4)} + 6c_1 \right) x^3 dx + c_2 \right)$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(y'[x])^3==12*y'[x]*(x*y'[x]-2*y'[x]),y[x],x,IncludeSingularSolutions -> True]
```

Not solved

### 4.39 problem 42

Internal problem ID [6105]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 42.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]`

$$3yy'y'' - y'^3 + 1 = 0$$

#### ✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 87

```
dsolve(3*y(x)*diff(y(x),x)*diff(y(x),x$2)=diff(y(x),x)^3-1,y(x), singsol=all)
```

$$\frac{3(c_1y(x) + 1)^{\frac{2}{3}}}{2c_1} - x - c_2 = 0$$

$$\frac{3(c_1y(x) + 1)^{\frac{2}{3}}}{c_1(-1 + i\sqrt{3})} - x - c_2 = 0$$

$$-\frac{3(c_1y(x) + 1)^{\frac{2}{3}}}{c_1(1 + i\sqrt{3})} - x - c_2 = 0$$

#### ✓ Solution by Mathematica

Time used: 0.174 (sec). Leaf size: 126

```
DSolve[3*y[x]*y'[x]*y''[x]==(y'[x])^3-1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{9}e^{-3c_1} \left( -9 + 2\sqrt{6}(e^{3c_1}(x + c_2))^{3/2} \right)$$

$$y(x) \rightarrow \frac{1}{9}e^{-3c_1} \left( -9 + 2\sqrt{6}(-\sqrt[3]{-1}e^{3c_1}(x + c_2))^{3/2} \right)$$

$$y(x) \rightarrow \frac{1}{9}e^{-3c_1} \left( -9 + 2\sqrt{6}((-1)^{2/3}e^{3c_1}(x + c_2))^{3/2} \right)$$

## 4.40 problem 43

Internal problem ID [6106]

**Book:** Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

**Section:** CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

**Problem number:** 43.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$4yy'^2y'' - y'^4 - 3 = 0$$

✓ Solution by Maple

Time used: 0.14 (sec). Leaf size: 91

```
dsolve(4*y(x)*diff(y(x),x)^2*diff(y(x),x$2)=diff(y(x),x)^4+3,y(x), singsol=all)
```

$$-\frac{4(c_1y(x) - 3)^{\frac{3}{4}}}{3c_1} - x - c_2 = 0$$

$$\frac{4(c_1y(x) - 3)^{\frac{3}{4}}}{3c_1} - x - c_2 = 0$$

$$-\frac{4i(c_1y(x) - 3)^{\frac{3}{4}}}{3c_1} - x - c_2 = 0$$

$$\frac{4i(c_1y(x) - 3)^{\frac{3}{4}}}{3c_1} - x - c_2 = 0$$



✓ Solution by Mathematica

Time used: 0.241 (sec). Leaf size: 156

`DSolve[4*y[x]*(y'[x])^2*y''[x]==(y'[x])^4+3,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{3}{8}e^{-4c_1} \left( 8 + \sqrt[3]{6}(-e^{4c_1}(x + c_2))^{4/3} \right)$$

$$y(x) \rightarrow \frac{3}{8}e^{-4c_1} \left( 8 + \sqrt[3]{6}(-ie^{4c_1}(x + c_2))^{4/3} \right)$$

$$y(x) \rightarrow \frac{3}{8}e^{-4c_1} \left( 8 + \sqrt[3]{6}(ie^{4c_1}(x + c_2))^{4/3} \right)$$

$$y(x) \rightarrow \frac{3}{8}e^{-4c_1} \left( 8 + \sqrt[3]{6}(e^{4c_1}(x + c_2))^{4/3} \right)$$