

A Solution Manual For

First order enumerated odes

Nasser M. Abbasi

October 12, 2023

Contents

1 section 1

2

1 section 1

1.1	problem 1	4
1.2	problem 2	5
1.3	problem 3	6
1.4	problem 4	7
1.5	problem 5	8
1.6	problem 6	9
1.7	problem 7	10
1.8	problem 8	11
1.9	problem 9	12
1.10	problem 10	13
1.11	problem 11	14
1.12	problem 12	15
1.13	problem 13	16
1.14	problem 14	17
1.15	problem 15	18
1.16	problem 16	19
1.17	problem 17	20
1.18	problem 18	21
1.19	problem 19	22
1.20	problem 20	23
1.21	problem 21	24
1.22	problem 22	25
1.23	problem 23	26
1.24	problem 24	27
1.25	problem 25	28
1.26	problem 26	29
1.27	problem 27	30
1.28	problem 28	31
1.29	problem 29	32
1.30	problem 30	33
1.31	problem 31	34
1.32	problem 32	35
1.33	problem 33	36
1.34	problem 34	37
1.35	problem 35	38
1.36	problem 36	39
1.37	problem 37	40
1.38	problem 38	41
1.39	problem 39	42
1.40	problem 40	43

1.41 problem 41	44
1.42 problem 42	45
1.43 problem 43	46
1.44 problem 44	47
1.45 problem 45	48
1.46 problem 46	49
1.47 problem 47	50
1.48 problem 48	51
1.49 problem 49	52
1.50 problem 50	53
1.51 problem 51	54
1.52 problem 52	55
1.53 problem 53	56
1.54 problem 54	57
1.55 problem 55	60
1.56 problem 56	61
1.57 problem 57	62
1.58 problem 58	63
1.59 problem 59	65
1.60 problem 60	67
1.61 problem 61	68
1.62 problem 62	69
1.63 problem 63	70
1.64 problem 64	71
1.65 problem 65	72
1.66 problem 66	73
1.67 problem 67	74
1.68 problem 68	75
1.69 problem 69	76
1.70 problem 70	77

1.1 problem 1

Internal problem ID [6563]

Book: First order enumerated odes

Section: section 1

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve(diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 7

```
DSolve[y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.2 problem 2

Internal problem ID [6564]

Book: First order enumerated odes

Section: section 1

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - a = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)=a,y(x), singsol=all)
```

$$y(x) = ax + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 11

```
DSolve[y'[x]==a,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow ax + c_1$$

1.3 problem 3

Internal problem ID [6565]

Book: First order enumerated odes

Section: section 1

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - x = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(diff(y(x),x)=x,y(x), singsol=all)
```

$$y(x) = \frac{x^2}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 15

```
DSolve[y'[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2}{2} + c_1$$

1.4 problem 4

Internal problem ID [6566]

Book: First order enumerated odes

Section: section 1

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - 1 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 7

```
dsolve(diff(y(x),x)=1,y(x), singsol=all)
```

$$y(x) = c_1 + x$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 9

```
DSolve[y'[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x + c_1$$

1.5 problem 5

Internal problem ID [6567]

Book: First order enumerated odes

Section: section 1

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - ax = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=a*x,y(x), singsol=all)
```

$$y(x) = \frac{ax^2}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 16

```
DSolve[y'[x]==a*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{ax^2}{2} + c_1$$

1.6 problem 6

Internal problem ID [6568]

Book: First order enumerated odes

Section: section 1

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' - axy = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x)=a*x*y(x),y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{ax^2}{2}}$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 23

```
DSolve[y'[x]==a*x*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{\frac{ax^2}{2}}$$

$$y(x) \rightarrow 0$$

1.7 problem 7

Internal problem ID [6569]

Book: First order enumerated odes

Section: section 1

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' - ax - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)=a*x+y(x),y(x), singsol=all)
```

$$y(x) = -ax - a + c_1 e^x$$

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 18

```
DSolve[y'[x]==a*x+y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -a(x + 1) + c_1 e^x$$

1.8 problem 8

Internal problem ID [6570]

Book: First order enumerated odes

Section: section 1

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear, 'class A']`

$$y' - ax - by = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x)=a*x+b*y(x),y(x), singsol=all)
```

$$y(x) = -\frac{ax}{b} - \frac{a}{b^2} + e^{xb}c_1$$

✓ Solution by Mathematica

Time used: 0.053 (sec). Leaf size: 25

```
DSolve[y'[x]==a*x+b*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{abx + a}{b^2} + c_1e^{bx}$$

1.9 problem 9

Internal problem ID [6571]

Book: First order enumerated odes

Section: section 1

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve(diff(y(x),x)=y(x),y(x), singsol=all)
```

$$y(x) = c_1 e^x$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 16

```
DSolve[y'[x]==y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x$$

$$y(x) \rightarrow 0$$

1.10 problem 10

Internal problem ID [6572]

Book: First order enumerated odes

Section: section 1

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - by = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)=b*y(x),y(x), singsol=all)
```

$$y(x) = e^{xb}c_1$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 18

```
DSolve[y'[x]==b*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{bx}$$

$$y(x) \rightarrow 0$$

1.11 problem 11

Internal problem ID [6573]

Book: First order enumerated odes

Section: section 1

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_Riccati, _special]]`

$$y' - ax - by^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 59

```
dsolve(diff(y(x),x)=a*x+b*y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{(ba)^{\frac{1}{3}} \left(\text{AiryAi} \left(1, -(ba)^{\frac{1}{3}} x \right) c_1 + \text{AiryBi} \left(1, -(ba)^{\frac{1}{3}} x \right) \right)}{b \left(c_1 \text{AiryAi} \left(-(ba)^{\frac{1}{3}} x \right) + \text{AiryBi} \left(-(ba)^{\frac{1}{3}} x \right) \right)}$$

✓ Solution by Mathematica

Time used: 0.164 (sec). Leaf size: 165

```
DSolve[y'[x]==a*x+b*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt{a}\sqrt{x} \left(-\text{BesselJ} \left(-\frac{2}{3}, \frac{2}{3}\sqrt{a}\sqrt{bx^{3/2}} \right) + c_1 \text{BesselJ} \left(\frac{2}{3}, \frac{2}{3}\sqrt{a}\sqrt{bx^{3/2}} \right) \right)}{\sqrt{b} \left(\text{BesselJ} \left(\frac{1}{3}, \frac{2}{3}\sqrt{a}\sqrt{bx^{3/2}} \right) + c_1 \text{BesselJ} \left(-\frac{1}{3}, \frac{2}{3}\sqrt{a}\sqrt{bx^{3/2}} \right) \right)}$$

$$y(x) \rightarrow \frac{ax^2 {}_0\tilde{F}_1 \left(; \frac{5}{3}; -\frac{1}{9}abx^3 \right)}{3 {}_0\tilde{F}_1 \left(; \frac{2}{3}; -\frac{1}{9}abx^3 \right)}$$

1.12 problem 12

Internal problem ID [6574]

Book: First order enumerated odes

Section: section 1

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y'c = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve(c*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 7

```
DSolve[c*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.13 problem 13

Internal problem ID [6575]

Book: First order enumerated odes

Section: section 1

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y'c - a = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(c*diff(y(x),x)=a,y(x), singsol=all)
```

$$y(x) = \frac{ax}{c} + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 14

```
DSolve[c*y'[x]==a,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{ax}{c} + c_1$$

1.14 problem 14

Internal problem ID [6576]

Book: First order enumerated odes

Section: section 1

Problem number: 14.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y'c - ax = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(c*diff(y(x),x)=a*x,y(x), singsol=all)
```

$$y(x) = \frac{ax^2}{2c} + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 19

```
DSolve[c*y'[x]==a*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{ax^2}{2c} + c_1$$

1.15 problem 15

Internal problem ID [6577]

Book: First order enumerated odes

Section: section 1

Problem number: 15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear, 'class A']`

$$y'c - ax - y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(c*diff(y(x),x)=a*x+y(x),y(x), singsol=all)
```

$$y(x) = -ac - ax + e^{\frac{x}{c}}c_1$$

✓ Solution by Mathematica

Time used: 0.051 (sec). Leaf size: 22

```
DSolve[c*y'[x]==a*x+y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -a(c + x) + c_1 e^{\frac{x}{c}}$$

1.16 problem 16

Internal problem ID [6578]

Book: First order enumerated odes

Section: section 1

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y'c - ax - by = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(c*diff(y(x),x)=a*x+b*y(x),y(x), singsol=all)
```

$$y(x) = -\frac{ax}{b} - \frac{ac}{b^2} + e^{\frac{bx}{c}} c_1$$

✓ Solution by Mathematica

Time used: 0.057 (sec). Leaf size: 28

```
DSolve[c*y'[x]==a*x+b*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{a(bx + c)}{b^2} + c_1 e^{\frac{bx}{c}}$$

1.17 problem 17

Internal problem ID [6579]

Book: First order enumerated odes

Section: section 1

Problem number: 17.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y'c - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(c*diff(y(x),x)=y(x),y(x), singsol=all)
```

$$y(x) = e^{\frac{x}{c}} c_1$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 20

```
DSolve[c*y'[x]==y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{\frac{x}{c}}$$

$$y(x) \rightarrow 0$$

1.18 problem 18

Internal problem ID [6580]

Book: First order enumerated odes

Section: section 1

Problem number: 18.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y'c - by = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(c*diff(y(x),x)=b*y(x),y(x), singsol=all)
```

$$y(x) = e^{\frac{xb}{c}} c_1$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 21

```
DSolve[c*y'[x]==b*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{\frac{bx}{c}}$$

$$y(x) \rightarrow 0$$

1.19 problem 19

Internal problem ID [6581]

Book: First order enumerated odes

Section: section 1

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_Riccati, _special]]`

$$y'c - ax - by^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 75

```
dsolve(c*diff(y(x),x)=a*x+b*y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{\left(\frac{ba}{c^2}\right)^{\frac{1}{3}} \left(\text{AiryAi}\left(1, -\left(\frac{ba}{c^2}\right)^{\frac{1}{3}} x\right) c_1 + \text{AiryBi}\left(1, -\left(\frac{ba}{c^2}\right)^{\frac{1}{3}} x\right) \right) c}{b \left(c_1 \text{AiryAi}\left(-\left(\frac{ba}{c^2}\right)^{\frac{1}{3}} x\right) + \text{AiryBi}\left(-\left(\frac{ba}{c^2}\right)^{\frac{1}{3}} x\right) \right)}$$

✓ Solution by Mathematica

Time used: 0.217 (sec). Leaf size: 214

```
DSolve[c*y'[x]==a*x+b*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt{x} \sqrt{\frac{a}{c}} \left(-\text{BesselJ}\left(-\frac{2}{3}, \frac{2}{3} \sqrt{\frac{a}{c}} \sqrt{\frac{b}{c}} x^{3/2}\right) + c_1 \text{BesselJ}\left(\frac{2}{3}, \frac{2}{3} \sqrt{\frac{a}{c}} \sqrt{\frac{b}{c}} x^{3/2}\right) \right)}{\sqrt{\frac{b}{c}} \left(\text{BesselJ}\left(\frac{1}{3}, \frac{2}{3} \sqrt{\frac{a}{c}} \sqrt{\frac{b}{c}} x^{3/2}\right) + c_1 \text{BesselJ}\left(-\frac{1}{3}, \frac{2}{3} \sqrt{\frac{a}{c}} \sqrt{\frac{b}{c}} x^{3/2}\right) \right)}$$

$$y(x) \rightarrow \frac{ax^2 {}_0\tilde{F}_1\left(\frac{5}{3}; -\frac{abx^3}{9c^2}\right)}{3c {}_0\tilde{F}_1\left(\frac{2}{3}; -\frac{abx^3}{9c^2}\right)}$$

1.20 problem 20

Internal problem ID [6582]

Book: First order enumerated odes

Section: section 1

Problem number: 20.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_Riccati, _special]]`

$$y'c - \frac{ax + by^2}{r} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 91

```
dsolve(c*diff(y(x),x)=(a*x+b*y(x)^2)/r,y(x), singsol=all)
```

$$y(x) = \frac{\left(\frac{ba}{r^2c^2}\right)^{\frac{1}{3}} \left(\text{AiryAi}\left(1, -\left(\frac{ba}{r^2c^2}\right)^{\frac{1}{3}}x\right) c_1 + \text{AiryBi}\left(1, -\left(\frac{ba}{r^2c^2}\right)^{\frac{1}{3}}x\right) \right) rc}{b \left(c_1 \text{AiryAi}\left(-\left(\frac{ba}{r^2c^2}\right)^{\frac{1}{3}}x\right) + \text{AiryBi}\left(-\left(\frac{ba}{r^2c^2}\right)^{\frac{1}{3}}x\right) \right)}$$

✓ Solution by Mathematica

Time used: 0.27 (sec). Leaf size: 253

```
DSolve[c*y'[x]==(a*x+b*y[x]^2)/r,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt{x}\sqrt{\frac{a}{cr}} \left(-\text{BesselJ}\left(-\frac{2}{3}, \frac{2}{3}\sqrt{\frac{a}{cr}}\sqrt{\frac{b}{cr}}x^{3/2}\right) + c_1 \text{BesselJ}\left(\frac{2}{3}, \frac{2}{3}\sqrt{\frac{a}{cr}}\sqrt{\frac{b}{cr}}x^{3/2}\right) \right)}{\sqrt{\frac{b}{cr}} \left(\text{BesselJ}\left(\frac{1}{3}, \frac{2}{3}\sqrt{\frac{a}{cr}}\sqrt{\frac{b}{cr}}x^{3/2}\right) + c_1 \text{BesselJ}\left(-\frac{1}{3}, \frac{2}{3}\sqrt{\frac{a}{cr}}\sqrt{\frac{b}{cr}}x^{3/2}\right) \right)}$$

$$y(x) \rightarrow \frac{ax^2 {}_0\tilde{F}_1\left(\frac{5}{3}; -\frac{abx^3}{9c^2r^2}\right)}{3cr {}_0\tilde{F}_1\left(\frac{2}{3}; -\frac{abx^3}{9c^2r^2}\right)}$$

1.21 problem 21

Internal problem ID [6583]

Book: First order enumerated odes

Section: section 1

Problem number: 21.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [rational, Riccati]

$$y'c - \frac{ax + by^2}{rx} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 94

```
dsolve(c*diff(y(x),x)=(a*x+b*y(x)^2)/(r*x),y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{\frac{xba}{r^2c^2}} cr \left(\text{BesselY} \left(1, 2\sqrt{\frac{xba}{r^2c^2}} \right) c_1 + \text{BesselJ} \left(1, 2\sqrt{\frac{xba}{r^2c^2}} \right) \right)}{b \left(c_1 \text{BesselY} \left(0, 2\sqrt{\frac{xba}{r^2c^2}} \right) + \text{BesselJ} \left(0, 2\sqrt{\frac{xba}{r^2c^2}} \right) \right)}$$

✓ Solution by Mathematica

Time used: 0.281 (sec). Leaf size: 160

```
DSolve[c*y'[x]==(a*x+b*y[x]^2)/(r*x),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\frac{2\sqrt{acr}\sqrt{x}Y_1\left(\frac{2\sqrt{a}\sqrt{b}\sqrt{x}}{cr}\right)}{\sqrt{b}} + ac_1x {}_0\tilde{F}_1\left(; 2; -\frac{abx}{c^2r^2}\right)}{2crY_0\left(\frac{2\sqrt{a}\sqrt{b}\sqrt{x}}{cr}\right) + cc_1r {}_0\tilde{F}_1\left(; 1; -\frac{abx}{c^2r^2}\right)}$$

$$y(x) \rightarrow \frac{ax {}_0\tilde{F}_1\left(; 2; -\frac{abx}{c^2r^2}\right)}{cr {}_0\tilde{F}_1\left(; 1; -\frac{abx}{c^2r^2}\right)}$$

1.22 problem 22

Internal problem ID [6584]

Book: First order enumerated odes

Section: section 1

Problem number: 22.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [rational, Riccati]

$$y'c - \frac{ax + by^2}{rx^2} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 106

```
dsolve(c*diff(y(x),x)=(a*x+b*y(x)^2)/(r*x^2),y(x), singsol=all)
```

$$y(x) = \frac{a \left(\text{BesselY} \left(0, 2\sqrt{\frac{ba}{c^2 r^2 x}} \right) c_1 + \text{BesselJ} \left(0, 2\sqrt{\frac{ba}{c^2 r^2 x}} \right) \right)}{cr \sqrt{\frac{ba}{c^2 r^2 x}} \left(c_1 \text{BesselY} \left(1, 2\sqrt{\frac{ba}{c^2 r^2 x}} \right) + \text{BesselJ} \left(1, 2\sqrt{\frac{ba}{c^2 r^2 x}} \right) \right)}$$

✓ Solution by Mathematica

Time used: 0.302 (sec). Leaf size: 179

```
DSolve[c*y'[x]==(a*x+b*y[x]^2)/(r*x^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt{acr}x \left(c_1 {}_0\tilde{F}_1 \left(; 1; -\frac{ab}{c^2 r^2 x} \right) + 2iY_0 \left(\frac{2\sqrt{a}\sqrt{b}\sqrt{\frac{1}{x}}}{cr} \right) \right)}{\sqrt{abc}c_1 {}_0\tilde{F}_1 \left(; 2; -\frac{ab}{c^2 r^2 x} \right) + \frac{2i\sqrt{bcr}Y_1 \left(\frac{2\sqrt{a}\sqrt{b}\sqrt{\frac{1}{x}}}{cr} \right)}{\sqrt{\frac{1}{x}}}}$$

$$y(x) \rightarrow \frac{crx {}_0\tilde{F}_1 \left(; 1; -\frac{ab}{c^2 r^2 x} \right)}{b {}_0\tilde{F}_1 \left(; 2; -\frac{ab}{c^2 r^2 x} \right)}$$

1.23 problem 23

Internal problem ID [6585]

Book: First order enumerated odes

Section: section 1

Problem number: 23.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_rational, _Bernoulli]

$$y'c - \frac{ax + by^2}{y} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 69

```
dsolve(c*dif(y(x),x)=(a*x+b*y(x)^2)/y(x),y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{4e^{\frac{2xb}{c}}c_1b^2 - 4axb - 2ac}}{2b}$$

$$y(x) = \frac{\sqrt{4e^{\frac{2xb}{c}}c_1b^2 - 4axb - 2ac}}{2b}$$

✓ Solution by Mathematica

Time used: 5.139 (sec). Leaf size: 85

```
DSolve[c*y'[x]==(a*x+b*y[x]^2)/y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{i\sqrt{abx + \frac{ac}{2} + b^2c_1\left(-e^{\frac{2bx}{c}}\right)}}{b}$$

$$y(x) \rightarrow \frac{i\sqrt{abx + \frac{ac}{2} + b^2c_1\left(-e^{\frac{2bx}{c}}\right)}}{b}$$

1.24 problem 24

Internal problem ID [6586]

Book: First order enumerated odes

Section: section 1

Problem number: 24.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$a \sin(x) y x y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(a*sin(x)*y(x)*x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 12

```
DSolve[a*Sin[x]*y[x]*x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow c_1$$

1.25 problem 25

Internal problem ID [6587]

Book: First order enumerated odes

Section: section 1

Problem number: 25.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$f(x) \sin(x) y x y' \pi = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(f(x)*sin(x)*y(x)*x*diff(y(x),x)*Pi=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 12

```
DSolve[f(x)*Sin[x]*y[x]*x*y'[x]*Pi==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow c_1$$

1.26 problem 26

Internal problem ID [6588]

Book: First order enumerated odes

Section: section 1

Problem number: 26.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' - \sin(x) - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)=sin(x)+y(x),y(x), singsol=all)
```

$$y(x) = -\frac{\cos(x)}{2} - \frac{\sin(x)}{2} + c_1 e^x$$

✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 24

```
DSolve[y'[x]==Sin[x]+y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sin(x)}{2} - \frac{\cos(x)}{2} + c_1 e^x$$

1.27 problem 27

Internal problem ID [6589]

Book: First order enumerated odes

Section: section 1

Problem number: 27.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [Riccati]

$$y' - \sin(x) - y^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
dsolve(diff(y(x),x)=sin(x)+y(x)^2,y(x), singsol=all)
```

$$y(x) = -\frac{c_1 \text{MathieuSPrime}\left(0, -2, -\frac{\pi}{4} + \frac{x}{2}\right) + \text{MathieuCPrime}\left(0, -2, -\frac{\pi}{4} + \frac{x}{2}\right)}{2\left(c_1 \text{MathieuS}\left(0, -2, -\frac{\pi}{4} + \frac{x}{2}\right) + \text{MathieuC}\left(0, -2, -\frac{\pi}{4} + \frac{x}{2}\right)\right)}$$

✓ Solution by Mathematica

Time used: 0.181 (sec). Leaf size: 105

```
DSolve[y'[x]==Sin[x]+y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-\text{MathieuSPrime}\left[0, -2, \frac{1}{4}(\pi - 2x)\right] + c_1 \text{MathieuCPrime}\left[0, -2, \frac{1}{4}(\pi - 2x)\right]}{2\left(\text{MathieuS}\left[0, -2, \frac{1}{4}(2x - \pi)\right] + c_1 \text{MathieuC}\left[0, -2, \frac{1}{4}(\pi - 2x)\right]\right)}$$

$$y(x) \rightarrow \frac{\text{MathieuCPrime}\left[0, -2, \frac{1}{4}(\pi - 2x)\right]}{2\text{MathieuC}\left[0, -2, \frac{1}{4}(\pi - 2x)\right]}$$

1.28 problem 28

Internal problem ID [6590]

Book: First order enumerated odes

Section: section 1

Problem number: 28.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' - \cos(x) - \frac{y}{x} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)=cos(x)+y(x)/x,y(x), singsol=all)
```

$$y(x) = (\text{Ci}(x) + c_1) x$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 12

```
DSolve[y'[x]==Cos[x]+y[x]/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(\text{CosIntegral}(x) + c_1)$$

1.29 problem 29

Internal problem ID [6591]

Book: First order enumerated odes

Section: section 1

Problem number: 29.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [Riccati]

$$y' - \cos(x) - \frac{y^2}{x} = 0$$

X Solution by Maple

```
dsolve(diff(y(x),x)=cos(x)+y(x)^2/x,y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y'[x]==Cos[x]+y[x]^2/x,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

1.30 problem 30

Internal problem ID [6592]

Book: First order enumerated odes

Section: section 1

Problem number: 30.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_Riccati]

$$y' - x - y - by^2 = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 105

```
dsolve(diff(y(x),x)=x+y(x)+b*y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{2b^{\frac{1}{3}} \text{AiryAi}\left(1, -\frac{4xb-1}{4b^{\frac{2}{3}}}\right) c_1 + 2 \text{AiryBi}\left(1, -\frac{4xb-1}{4b^{\frac{2}{3}}}\right) b^{\frac{1}{3}} - \text{AiryAi}\left(-\frac{4xb-1}{4b^{\frac{2}{3}}}\right) c_1 - \text{AiryBi}\left(-\frac{4xb-1}{4b^{\frac{2}{3}}}\right)}{2b \left(\text{AiryAi}\left(-\frac{4xb-1}{4b^{\frac{2}{3}}}\right) c_1 + \text{AiryBi}\left(-\frac{4xb-1}{4b^{\frac{2}{3}}}\right) \right)}$$

✓ Solution by Mathematica

Time used: 0.175 (sec). Leaf size: 195

```
DSolve[y'[x]==x+y[x]+b*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\text{AiryBi}\left(\frac{\frac{1}{4}-bx}{(-b)^{2/3}}\right) + c_1 \text{AiryAi}\left(\frac{\frac{1}{4}-bx}{(-b)^{2/3}}\right) + 2\sqrt[3]{-b} \left(\text{AiryBiPrime}\left(\frac{\frac{1}{4}-bx}{(-b)^{2/3}}\right) + c_1 \text{AiryAiPrime}\left(\frac{\frac{1}{4}-bx}{(-b)^{2/3}}\right) \right)}{2b \left(\text{AiryBi}\left(\frac{\frac{1}{4}-bx}{(-b)^{2/3}}\right) + c_1 \text{AiryAi}\left(\frac{\frac{1}{4}-bx}{(-b)^{2/3}}\right) \right)}$$

$$y(x) \rightarrow -\frac{2\sqrt[3]{-b} \text{AiryAiPrime}\left(\frac{\frac{1}{4}-bx}{(-b)^{2/3}}\right)}{\text{AiryAi}\left(\frac{\frac{1}{4}-bx}{(-b)^{2/3}}\right)} + 1$$

1.31 problem 31

Internal problem ID [6593]

Book: First order enumerated odes

Section: section 1

Problem number: 31.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y'x = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve(x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 7

```
DSolve[x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.32 problem 32

Internal problem ID [6594]

Book: First order enumerated odes

Section: section 1

Problem number: 32.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$5y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve(5*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 7

```
DSolve[5*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.33 problem 33

Internal problem ID [6595]

Book: First order enumerated odes

Section: section 1

Problem number: 33.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$ey' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve(exp(1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 7

```
DSolve[Exp[1]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.34 problem 34

Internal problem ID [6596]

Book: First order enumerated odes

Section: section 1

Problem number: 34.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$\pi y' = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 5

```
dsolve(Pi*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 7

```
DSolve[Pi*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.35 problem 35

Internal problem ID [6597]

Book: First order enumerated odes

Section: section 1

Problem number: 35.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' \sin(x) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve(sin(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 7

```
DSolve[Sin[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.36 problem 36

Internal problem ID [6598]

Book: First order enumerated odes

Section: section 1

Problem number: 36.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$f(x)y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve(f(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 7

```
DSolve[f[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.37 problem 37

Internal problem ID [6599]

Book: First order enumerated odes

Section: section 1

Problem number: 37.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y'x - 1 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve(x*diff(y(x),x)=1,y(x), singsol=all)
```

$$y(x) = \ln(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 10

```
DSolve[x*y'[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(x) + c_1$$

1.38 problem 38

Internal problem ID [6600]

Book: First order enumerated odes

Section: section 1

Problem number: 38.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y'x - \sin(x) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve(x*diff(y(x),x)=sin(x),y(x), singsol=all)
```

$$y(x) = \text{Si}(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 10

```
DSolve[x*y'[x]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{Si}(x) + c_1$$

1.39 problem 39

Internal problem ID [6601]

Book: First order enumerated odes

Section: section 1

Problem number: 39.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y'(-1 + x) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve((x-1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 7

```
DSolve[(x-1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.40 problem 40

Internal problem ID [6602]

Book: First order enumerated odes

Section: section 1

Problem number: 40.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$yy' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve(y(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = -c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 12

```
DSolve[y[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow c_1$$

1.41 problem 41

Internal problem ID [6603]

Book: First order enumerated odes

Section: section 1

Problem number: 41.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$yxy' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(x*y(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 12

```
DSolve[x*y[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow c_1$$

1.42 problem 42

Internal problem ID [6604]

Book: First order enumerated odes

Section: section 1

Problem number: 42.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$xy \sin(x) y' = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 9

```
dsolve(x*y(x)*sin(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 12

```
DSolve[x*y[x]*Sin[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow c_1$$

1.43 problem 43

Internal problem ID [6605]

Book: First order enumerated odes

Section: section 1

Problem number: 43.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$\pi y \sin(x) y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(Pi*y(x)*sin(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 12

```
DSolve[Pi*y[x]*Sin[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow c_1$$

1.44 problem 44

Internal problem ID [6606]

Book: First order enumerated odes

Section: section 1

Problem number: 44.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$x \sin(x) y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve(x*sin(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 7

```
DSolve[x*Sin[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.45 problem 45

Internal problem ID [6607]

Book: First order enumerated odes

Section: section 1

Problem number: 45.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$x \sin(x) y'^2 = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 5

```
dsolve(x*sin(x)*diff(y(x),x)^2=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 7

```
DSolve[x*Sin[x]*y'[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.46 problem 46

Internal problem ID [6608]

Book: First order enumerated odes

Section: section 1

Problem number: 46.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$yy'^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(y(x)*diff(y(x),x)^2=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 12

```
DSolve[y[x]*(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow c_1$$

1.47 problem 47

Internal problem ID [6609]

Book: First order enumerated odes

Section: section 1

Problem number: 47.

ODE order: 1.

ODE degree: 0.

CAS Maple gives this as type [_quadrature]

$$y'^n = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve(diff(y(x),x)^n=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 15

```
DSolve[(y'[x])^n==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0^{\frac{1}{n}} x + c_1$$

1.48 problem 48

Internal problem ID [6610]

Book: First order enumerated odes

Section: section 1

Problem number: 48.

ODE order: 1.

ODE degree: 0.

CAS Maple gives this as type [_quadrature]

$$xy^n = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve(x*diff(y(x),x)^n=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 15

```
DSolve[x*(y'[x])^n==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0^{\frac{1}{n}}x + c_1$$

1.49 problem 49

Internal problem ID [6611]

Book: First order enumerated odes

Section: section 1

Problem number: 49.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$y'^2 - x = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)^2=x,y(x), singsol=all)
```

$$y(x) = \frac{2x^{\frac{3}{2}}}{3} + c_1$$

$$y(x) = -\frac{2x^{\frac{3}{2}}}{3} + c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 33

```
DSolve[(y'[x])^2==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2x^{3/2}}{3} + c_1$$

$$y(x) \rightarrow \frac{2x^{3/2}}{3} + c_1$$

1.50 problem 50

Internal problem ID [6612]

Book: First order enumerated odes

Section: section 1

Problem number: 50.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$y'^2 - x - y = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 24

```
dsolve(diff(y(x),x)^2=x+y(x),y(x), singsol=all)
```

$$y(x) = \left(-\text{LambertW}\left(-c_1 e^{-\frac{x}{2}-1}\right) - 1\right)^2 - x$$

✓ Solution by Mathematica

Time used: 17.832 (sec). Leaf size: 98

```
DSolve[(y'[x])^2==x+y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow W\left(-e^{-\frac{x}{2}-1-\frac{c_1}{2}}\right) \left(2 + W\left(-e^{-\frac{x}{2}-1-\frac{c_1}{2}}\right)\right) - x + 1$$

$$y(x) \rightarrow W\left(e^{\frac{1}{2}(-x-2+c_1)}\right) \left(2 + W\left(e^{\frac{1}{2}(-x-2+c_1)}\right)\right) - x + 1$$

$$y(x) \rightarrow 1 - x$$

1.51 problem 51

Internal problem ID [6613]

Book: First order enumerated odes

Section: section 1

Problem number: 51.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _dAlembert]`

$$y'^2 - \frac{y}{x} = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 39

```
dsolve(diff(y(x),x)^2=y(x)/x,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = \frac{(x + \sqrt{xc_1})^2}{x}$$

$$y(x) = \frac{(-x + \sqrt{xc_1})^2}{x}$$

✓ Solution by Mathematica

Time used: 0.046 (sec). Leaf size: 46

```
DSolve[(y'[x])^2==y[x]/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}(-2\sqrt{x} + c_1)^2$$

$$y(x) \rightarrow \frac{1}{4}(2\sqrt{x} + c_1)^2$$

$$y(x) \rightarrow 0$$

1.52 problem 52

Internal problem ID [6614]

Book: First order enumerated odes

Section: section 1

Problem number: 52.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_separable]

$$y'^2 - \frac{y^2}{x} = 0$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 27

```
dsolve(diff(y(x),x)^2=y(x)^2/x,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1 e^{-2\sqrt{x}}$$

$$y(x) = c_1 e^{2\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.064 (sec). Leaf size: 38

```
DSolve[(y'[x])^2==y[x]^2/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-2\sqrt{x}}$$

$$y(x) \rightarrow c_1 e^{2\sqrt{x}}$$

$$y(x) \rightarrow 0$$

1.53 problem 53

Internal problem ID [6615]

Book: First order enumerated odes

Section: section 1

Problem number: 53.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_homogeneous, 'class G']]`

$$y'^2 - \frac{y^3}{x} = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 29

```
dsolve(diff(y(x),x)^2=y(x)^3/x,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = \text{WeierstrassP}\left(\frac{x2^{\frac{2}{3}}}{\sqrt{x2^{\frac{2}{3}}}} + c_1, 0, 0\right) 2^{\frac{2}{3}}$$

✓ Solution by Mathematica

Time used: 0.069 (sec). Leaf size: 42

```
DSolve[(y'[x])^2==y[x]^3/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{4}{(-2\sqrt{x} + c_1)^2}$$

$$y(x) \rightarrow \frac{4}{(2\sqrt{x} + c_1)^2}$$

$$y(x) \rightarrow 0$$

1.54 problem 54

Internal problem ID [6616]

Book: First order enumerated odes

Section: section 1

Problem number: 54.

ODE order: 1.

ODE degree: 3.

CAS Maple gives this as type `[_homogeneous, 'class G'], _rational]`

$$y'^3 - \frac{y^2}{x} = 0$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 453

```
dsolve(diff(y(x),x)^3=y(x)^2/x,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = -\frac{3x^{\frac{4}{3}}c_1}{8} + \frac{3x^{\frac{2}{3}}c_1^2}{8} - \frac{c_1^3}{8} + \frac{x^2}{8}$$

$$y(x) = -\frac{3\left(-\frac{x^{\frac{2}{3}}}{2} - \frac{i\sqrt{3}x^{\frac{2}{3}}}{2}\right)^2 c_1}{8} + \frac{3\left(-\frac{x^{\frac{2}{3}}}{2} - \frac{i\sqrt{3}x^{\frac{2}{3}}}{2}\right) c_1^2}{8} - \frac{c_1^3}{8} + \frac{x^2}{8}$$

$$y(x) = -\frac{3\left(-\frac{x^{\frac{2}{3}}}{2} + \frac{i\sqrt{3}x^{\frac{2}{3}}}{2}\right)^2 c_1}{8} + \frac{3\left(-\frac{x^{\frac{2}{3}}}{2} + \frac{i\sqrt{3}x^{\frac{2}{3}}}{2}\right) c_1^2}{8} - \frac{c_1^3}{8} + \frac{x^2}{8}$$

$$y(x) = \frac{3\left(2x^{\frac{2}{3}} + c_1\right)^2 c_1}{64} - \frac{3c_1^2\left(2x^{\frac{2}{3}} + c_1\right)}{64} + \frac{c_1^3}{64} + \frac{x^2}{8}$$

$$y(x) = \frac{3\left(-x^{\frac{2}{3}} + c_1 - i\sqrt{3}x^{\frac{2}{3}}\right)^2 c_1}{64} - \frac{3c_1^2\left(-x^{\frac{2}{3}} + c_1 - i\sqrt{3}x^{\frac{2}{3}}\right)}{64} + \frac{c_1^3}{64} + \frac{x^2}{8}$$

$$y(x) = \frac{3\left(-x^{\frac{2}{3}} + c_1 + i\sqrt{3}x^{\frac{2}{3}}\right)^2 c_1}{64} - \frac{3c_1^2\left(-x^{\frac{2}{3}} + c_1 + i\sqrt{3}x^{\frac{2}{3}}\right)}{64} + \frac{c_1^3}{64} + \frac{x^2}{8}$$

$$y(x) = -\frac{3\left(2x^{\frac{2}{3}} - c_1\right)^2 c_1}{64} - \frac{3c_1^2\left(2x^{\frac{2}{3}} - c_1\right)}{64} - \frac{c_1^3}{64} + \frac{x^2}{8}$$

$$y(x) = -\frac{3\left(-x^{\frac{2}{3}} - c_1 - i\sqrt{3}x^{\frac{2}{3}}\right)^2 c_1}{64} - \frac{3c_1^2\left(-x^{\frac{2}{3}} - c_1 - i\sqrt{3}x^{\frac{2}{3}}\right)}{64} - \frac{c_1^3}{64} + \frac{x^2}{8}$$

$$y(x) = -\frac{3\left(-x^{\frac{2}{3}} - c_1 + i\sqrt{3}x^{\frac{2}{3}}\right)^2 c_1}{64} - \frac{3c_1^2\left(-x^{\frac{2}{3}} - c_1 + i\sqrt{3}x^{\frac{2}{3}}\right)}{64} - \frac{c_1^3}{64} + \frac{x^2}{8}$$

✓ Solution by Mathematica

Time used: 0.083 (sec). Leaf size: 152

```
DSolve[(y'[x])^3==y[x]^2/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{216} (3x^{2/3} + 2c_1)^3$$

$$y(x) \rightarrow \frac{1}{216} \left(18i(\sqrt{3} + i) c_1^2 x^{2/3} - 27i(\sqrt{3} - i) c_1 x^{4/3} + 27x^2 + 8c_1^3 \right)$$

$$y(x) \rightarrow \frac{1}{216} \left(-18i(\sqrt{3} - i) c_1^2 x^{2/3} + 27i(\sqrt{3} + i) c_1 x^{4/3} + 27x^2 + 8c_1^3 \right)$$

$$y(x) \rightarrow 0$$

1.55 problem 55

Internal problem ID [6617]

Book: First order enumerated odes

Section: section 1

Problem number: 55.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [[_homogeneous, 'class G']]

$$y'^2 - \frac{1}{yx} = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 43

```
dsolve(diff(y(x),x)^2=1/(y(x)*x),y(x), singsol=all)
```

$$\frac{(xy(x))^{\frac{3}{2}}}{x^{\frac{3}{2}}} - 3\sqrt{x} - c_1 = 0$$

$$\frac{(xy(x))^{\frac{3}{2}}}{x^{\frac{3}{2}}} + 3\sqrt{x} - c_1 = 0$$

✓ Solution by Mathematica

Time used: 3.51 (sec). Leaf size: 53

```
DSolve[(y'[x])^2==1/(y[x]*x),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left(\frac{3}{2}\right)^{2/3} (-2\sqrt{x} + c_1)^{2/3}$$

$$y(x) \rightarrow \left(\frac{3}{2}\right)^{2/3} (2\sqrt{x} + c_1)^{2/3}$$

1.56 problem 56

Internal problem ID [6618]

Book: First order enumerated odes

Section: section 1

Problem number: 56.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [[_homogeneous, 'class G']]

$$y'^2 - \frac{1}{xy^3} = 0$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 43

```
dsolve(diff(y(x),x)^2=1/(x*y(x)^3),y(x), singsol=all)
```

$$\frac{(xy(x))^{\frac{5}{2}}}{x^{\frac{5}{2}}} - 5\sqrt{x} - c_1 = 0$$

$$\frac{(xy(x))^{\frac{5}{2}}}{x^{\frac{5}{2}}} + 5\sqrt{x} - c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.104 (sec). Leaf size: 53

```
DSolve[(y'[x])^2==1/(x*y[x]^3),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left(\frac{5}{2}\right)^{2/5} (-2\sqrt{x} + c_1)^{2/5}$$

$$y(x) \rightarrow \left(\frac{5}{2}\right)^{2/5} (2\sqrt{x} + c_1)^{2/5}$$

1.57 problem 57

Internal problem ID [6619]

Book: First order enumerated odes

Section: section 1

Problem number: 57.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_separable]

$$y'^2 - \frac{1}{x^2 y^3} = 0$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 29

```
dsolve(diff(y(x),x)^2=1/(x^2*y(x)^3),y(x), singsol=all)
```

$$\ln(x) - \frac{2y(x)^{\frac{5}{2}}}{5} - c_1 = 0$$

$$\ln(x) + \frac{2y(x)^{\frac{5}{2}}}{5} - c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.127 (sec). Leaf size: 45

```
DSolve[(y'[x])^2==1/(x^2*y[x]^3),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left(\frac{5}{2}\right)^{2/5} (-\log(x) + c_1)^{2/5}$$

$$y(x) \rightarrow \left(\frac{5}{2}\right)^{2/5} (\log(x) + c_1)^{2/5}$$

1.58 problem 58

Internal problem ID [6620]

Book: First order enumerated odes

Section: section 1

Problem number: 58.

ODE order: 1.

ODE degree: 4.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational]`

$$y'^4 - \frac{1}{xy^3} = 0$$

✓ Solution by Maple

Time used: 0.109 (sec). Leaf size: 99

```
dsolve(diff(y(x),x)^4=1/(x*y(x)^3),y(x), singsol=all)
```

$$\frac{3(y(x)x^3)^{\frac{7}{4}}}{x^{\frac{21}{4}}} - 7x^{\frac{3}{4}} - c_1 = 0$$

$$\frac{3i(y(x)x^3)^{\frac{7}{4}}}{x^{\frac{21}{4}}} - 7x^{\frac{3}{4}} - c_1 = 0$$

$$\frac{3i(y(x)x^3)^{\frac{7}{4}}}{x^{\frac{21}{4}}} + 7x^{\frac{3}{4}} - c_1 = 0$$

$$\frac{3(y(x)x^3)^{\frac{7}{4}}}{x^{\frac{21}{4}}} + 7x^{\frac{3}{4}} - c_1 = 0$$

✓ Solution by Mathematica

Time used: 7.391 (sec). Leaf size: 129

```
DSolve[(y'[x])^4==1/(x*y[x]^3),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\left(-\frac{28x^{3/4}}{3} + 7c_1\right)^{4/7}}{2\sqrt[7]{2}}$$

$$y(x) \rightarrow \frac{\left(7c_1 - \frac{28ix^{3/4}}{3}\right)^{4/7}}{2\sqrt[7]{2}}$$

$$y(x) \rightarrow \frac{\left(\frac{28ix^{3/4}}{3} + 7c_1\right)^{4/7}}{2\sqrt[7]{2}}$$

$$y(x) \rightarrow \frac{\left(\frac{28x^{3/4}}{3} + 7c_1\right)^{4/7}}{2\sqrt[7]{2}}$$

1.59 problem 59

Internal problem ID [6621]

Book: First order enumerated odes

Section: section 1

Problem number: 59.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_separable]

$$y'^2 - \frac{1}{x^3 y^4} = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 185

```
dsolve(diff(y(x),x)^2=1/(x^3*y(x)^4),y(x), singsol=all)
```

$$y(x) = \left(\frac{c_1 \sqrt{x} - 6}{\sqrt{x}} \right)^{\frac{1}{3}}$$

$$y(x) = -\frac{\left(\frac{c_1 \sqrt{x} - 6}{\sqrt{x}} \right)^{\frac{1}{3}}}{2} - \frac{i\sqrt{3} \left(\frac{c_1 \sqrt{x} - 6}{\sqrt{x}} \right)^{\frac{1}{3}}}{2}$$

$$y(x) = -\frac{\left(\frac{c_1 \sqrt{x} - 6}{\sqrt{x}} \right)^{\frac{1}{3}}}{2} + \frac{i\sqrt{3} \left(\frac{c_1 \sqrt{x} - 6}{\sqrt{x}} \right)^{\frac{1}{3}}}{2}$$

$$y(x) = \left(\frac{c_1 \sqrt{x} + 6}{\sqrt{x}} \right)^{\frac{1}{3}}$$

$$y(x) = -\frac{\left(\frac{c_1 \sqrt{x} + 6}{\sqrt{x}} \right)^{\frac{1}{3}}}{2} - \frac{i\sqrt{3} \left(\frac{c_1 \sqrt{x} + 6}{\sqrt{x}} \right)^{\frac{1}{3}}}{2}$$

$$y(x) = -\frac{\left(\frac{c_1 \sqrt{x} + 6}{\sqrt{x}} \right)^{\frac{1}{3}}}{2} + \frac{i\sqrt{3} \left(\frac{c_1 \sqrt{x} + 6}{\sqrt{x}} \right)^{\frac{1}{3}}}{2}$$

✓ Solution by Mathematica

Time used: 3.614 (sec). Leaf size: 157

`DSolve[(y'[x])^2==1/(x^3*y[x]^4),y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow -\sqrt[3]{-3} \sqrt[3]{-\frac{2}{\sqrt{x}}} + c_1$$

$$y(x) \rightarrow \sqrt[3]{3} \sqrt[3]{-\frac{2}{\sqrt{x}}} + c_1$$

$$y(x) \rightarrow (-1)^{2/3} \sqrt[3]{3} \sqrt[3]{-\frac{2}{\sqrt{x}}} + c_1$$

$$y(x) \rightarrow -\sqrt[3]{-3} \sqrt[3]{\frac{2}{\sqrt{x}}} + c_1$$

$$y(x) \rightarrow \sqrt[3]{3} \sqrt[3]{\frac{2}{\sqrt{x}}} + c_1$$

$$y(x) \rightarrow (-1)^{2/3} \sqrt[3]{3} \sqrt[3]{\frac{2}{\sqrt{x}}} + c_1$$

1.60 problem 60

Internal problem ID [6622]

Book: First order enumerated odes

Section: section 1

Problem number: 60.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$y' - \sqrt{1 + 6x + y} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 57

```
dsolve(diff(y(x),x)=(1+6*x+y(x))^(1/2),y(x), singsol=all)
```

$$\begin{aligned} x - 2\sqrt{1 + 6x + y(x)} - 6 \ln \left(-6 + \sqrt{1 + 6x + y(x)} \right) \\ + 6 \ln \left(6 + \sqrt{1 + 6x + y(x)} \right) + 6 \ln (-35 + y(x) + 6x) - c_1 = 0 \end{aligned}$$

✓ Solution by Mathematica

Time used: 13.621 (sec). Leaf size: 108

```
DSolve[y'[x]==(1+6*x+y[x])^(1/2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 36W \left(-\frac{1}{6} e^{\frac{1}{72}(-6x-73+6c_1)} \right) \left(2 + W \left(-\frac{1}{6} e^{\frac{1}{72}(-6x-73+6c_1)} \right) \right) - 6x + 35$$

$$y(x) \rightarrow 35 - 6x$$

$$y(x) \rightarrow 36W \left(-\frac{1}{6} e^{\frac{1}{72}(-6x-73)} \right) \left(W \left(-\frac{1}{6} e^{\frac{1}{72}(-6x-73)} \right) + 2 \right) - 6x + 35$$

1.61 problem 61

Internal problem ID [6623]

Book: First order enumerated odes

Section: section 1

Problem number: 61.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$y' - (1 + 6x + y)^{\frac{1}{3}} = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 79

```
dsolve(diff(y(x),x)=(1+6*x+y(x))^(1/3),y(x), singsol=all)
```

$$x - \frac{3(1 + 6x + y(x))^{\frac{2}{3}}}{2} + 36 \ln \left((1 + 6x + y(x))^{\frac{2}{3}} - 6(1 + 6x + y(x))^{\frac{1}{3}} + 36 \right) - 72 \ln \left(6 + (1 + 6x + y(x))^{\frac{1}{3}} \right) - 36 \ln (217 + y(x) + 6x) + 18(1 + 6x + y(x))^{\frac{1}{3}} - c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.245 (sec). Leaf size: 66

```
DSolve[y'[x]==(1+6*x+y[x])^(1/3),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{1}{6} \left(y(x) - 9(y(x) + 6x + 1)^{2/3} + 108 \sqrt[3]{y(x) + 6x + 1} - 648 \log \left(\sqrt[3]{y(x) + 6x + 1} + 6 \right) + 6x + 1 \right) - \frac{y(x)}{6} = c_1, y(x) \right]$$

1.62 problem 62

Internal problem ID [6624]

Book: First order enumerated odes

Section: section 1

Problem number: 62.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$y' - (1 + 6x + y)^{\frac{1}{4}} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 109

```
dsolve(diff(y(x),x)=(1+6*x+y(x))^(1/4),y(x), singsol=all)
```

$$\begin{aligned} & x + 216 \ln(1295 - y(x) - 6x) + 12\sqrt{1 + 6x + y(x)} - 216 \ln\left(\sqrt{1 + 6x + y(x)} + 36\right) \\ & + 216 \ln\left(\sqrt{1 + 6x + y(x)} - 36\right) - 144(1 + 6x + y(x))^{\frac{1}{4}} - 432 \ln\left((1 + 6x + y(x))^{\frac{1}{4}} - 6\right) \\ & + 432 \ln\left(6 + (1 + 6x + y(x))^{\frac{1}{4}}\right) - \frac{4(1 + 6x + y(x))^{\frac{3}{4}}}{3} - c_1 = 0 \end{aligned}$$

✓ Solution by Mathematica

Time used: 2.504 (sec). Leaf size: 79

```
DSolve[y'[x]==(1+6*x+y[x])^(1/4),y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned} & \text{Solve} \left[\frac{1}{6} \left(y(x) - 8(y(x) + 6x + 1)^{3/4} + 72\sqrt{y(x) + 6x + 1} - 864\sqrt[4]{y(x) + 6x + 1} \right. \right. \\ & \left. \left. + 5184 \log\left(\sqrt[4]{y(x) + 6x + 1} + 6\right) + 6x + 1\right) - \frac{y(x)}{6} = c_1, y(x) \right] \end{aligned}$$

1.63 problem 63

Internal problem ID [6625]

Book: First order enumerated odes

Section: section 1

Problem number: 63.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$y' - (a + bx + y)^4 = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 49

```
dsolve(diff(y(x),x)=(a+b*x+y(x))^4,y(x), singsol=all)
```

$$y(x) = -xb + \text{RootOf} \left(-x + \int^{-z} \frac{1}{-a^4 + 4_a^3a + 6_a^2a^2 + 4_a a^3 + a^4 + b} d_a + c_1 \right)$$

✓ Solution by Mathematica

Time used: 0.431 (sec). Leaf size: 163

```
DSolve[y'[x]==(a+b*x+y[x])^4,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{2\sqrt{2} \arctan \left(1 - \frac{\sqrt{2}(a+bx+y(x))}{\sqrt[4]{b}} \right) - 2\sqrt{2} \arctan \left(\frac{\sqrt{2}(a+bx+y(x))}{\sqrt[4]{b}} + 1 \right) + \sqrt{2} \log \left((a + bx + y(x))^2 - \sqrt{2} \right)}{8b^{3/4}} \right]$$

1.64 problem 64

Internal problem ID [6626]

Book: First order enumerated odes

Section: section 1

Problem number: 64.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$y' - (\pi + x + 7y)^{\frac{7}{2}} = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 74

```
dsolve(diff(y(x),x)=(Pi+x+7*y(x))^(7/2),y(x), singsol=all)
```

$$y(x) = -\frac{x}{7} + \text{RootOf}\left(-x + 7\left(\int^{-z} \frac{1}{7\pi^3\sqrt{\pi+7_a} + 147\pi^2_a\sqrt{\pi+7_a} + 1029\pi_a^2\sqrt{\pi+7_a} + 2401_a^3\sqrt{\pi+7_a} + 1} d_a + c_1\right)\right)$$

✓ Solution by Mathematica

Time used: 30.501 (sec). Leaf size: 43

```
DSolve[y'[x]==(Pi+x+7*y[x])^(7/2),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[-(7y(x) + x + \pi) \left(\text{Hypergeometric2F1}\left(\frac{2}{7}, 1, \frac{9}{7}, -7(x + 7y(x) + \pi)^{7/2}\right) - 1\right) - 7y(x) = c_1, y(x)\right]$$

1.65 problem 65

Internal problem ID [6627]

Book: First order enumerated odes

Section: section 1

Problem number: 65.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$y' - (a + xb + yc)^6 = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 94

```
dsolve(diff(y(x),x)=(a+b*x+c*y(x))^6,y(x), singsol=all)
```

$$y(x) = \frac{\text{RootOf}\left(\left(\int^{-Z} \frac{1}{c^7 - a^6 + 6a^5ac^6 + 15a^4a^2c^5 + 20a^3a^3c^4 + 15a^2a^4c^3 + 6aa^5c^2 + a^6c + b} d_a\right) c - x + c_1\right) c - xb}{c}$$

✓ Solution by Mathematica

Time used: 1.762 (sec). Leaf size: 274

```
DSolve[y'[x]==(a+b*x+c*y[x])^6,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[\frac{-4\sqrt[6]{b} \arctan\left(\frac{\sqrt[6]{C(a+bx+cy(x))}}{\sqrt[6]{b}}\right) + 2\sqrt[6]{b} \arctan\left(\sqrt{3} - \frac{2\sqrt[6]{C(a+bx+cy(x))}}{\sqrt[6]{b}}\right) - 2\sqrt[6]{b} \arctan\left(\frac{2\sqrt[6]{C(a+bx+cy(x))}}{\sqrt[6]{b}}\right)}{-\frac{cy(x)}{b} = c_1, y(x)}\right]$$

1.66 problem 66

Internal problem ID [6628]

Book: First order enumerated odes

Section: section 1

Problem number: 66.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' - e^{x+y} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
dsolve(diff(y(x),x)=exp(x+y(x)),y(x), singsol=all)
```

$$y(x) = \ln\left(-\frac{1}{e^x + c_1}\right)$$

✓ Solution by Mathematica

Time used: 0.772 (sec). Leaf size: 18

```
DSolve[y'[x]==Exp[x+y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\log(-e^x - c_1)$$

1.67 problem 67

Internal problem ID [6629]

Book: First order enumerated odes

Section: section 1

Problem number: 67.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [[_homogeneous, 'class C'], _dAlembert]

$$y' - 10 - e^{x+y} = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)=10+exp(x+y(x)),y(x), singsol=all)
```

$$y(x) = -x + \ln\left(\frac{11}{e^{-11x}c_1 - 1}\right)$$

✓ Solution by Mathematica

Time used: 3.171 (sec). Leaf size: 42

```
DSolve[y'[x]==10+Exp[x+y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log\left(-\frac{11e^{10x+11c_1}}{-1 + e^{11(x+c_1)}}\right)$$

$$y(x) \rightarrow \log(-11e^{-x})$$

1.68 problem 68

Internal problem ID [6630]

Book: First order enumerated odes

Section: section 1

Problem number: 68.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x),G(x)]']]`

$$y' - 10e^{x+y} - x^2 = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve(diff(y(x),x)=10*exp(x+y(x))+x^2,y(x), singsol=all)
```

$$y(x) = \frac{x^3}{3} - \ln\left(-c_1 - 10\left(\int e^x e^{\frac{x^3}{3}} dx\right)\right)$$

✓ Solution by Mathematica

Time used: 0.403 (sec). Leaf size: 115

```
DSolve[y'[x]==10*Exp[x+y[x]]+x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[\int_1^{y(x)} -\frac{1}{10}e^{-K[2]}\left(10e^{K[2]}\int_1^x -\frac{1}{10}e^{\frac{K[1]^3}{3}-K[2]}K[1]^2 dK[1] + e^{\frac{x^3}{3}}\right) dK[2] + \int_1^x \left(\frac{1}{10}e^{\frac{K[1]^3}{3}-y(x)}K[1]^2 + e^{\frac{K[1]^3}{3}+K[1]}\right) dK[1] = c_1, y(x)\right]$$

1.69 problem 69

Internal problem ID [6631]

Book: First order enumerated odes

Section: section 1

Problem number: 69.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x),G(x)]']]`

$$y' - x e^{x+y} - \sin(x) = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve(diff(y(x),x)=x*exp(x+y(x))+sin(x),y(x), singsol=all)
```

$$y(x) = -\cos(x) - \ln\left(-c_1 - \left(\int x e^x e^{-\cos(x)} dx\right)\right)$$

✓ Solution by Mathematica

Time used: 3.673 (sec). Leaf size: 100

```
DSolve[y'[x]==x*Exp[x+y[x]]+Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[\int_1^x \left(-e^{K[1]-\cos(K[1])} K[1] - e^{-\cos(K[1])-y(x)} \sin(K[1])\right) dK[1] + \int_1^{y(x)} -e^{-\cos(x)-K[2]} \left(e^{\cos(x)+K[2]} \int_1^x e^{-\cos(K[1]-K[2]} \sin(K[1]) dK[1] - 1\right) dK[2] = c_1, y(x)\right]$$

1.70 problem 70

Internal problem ID [6632]

Book: First order enumerated odes

Section: section 1

Problem number: 70.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x),G(x)]']]`

$$y' - 5e^{x^2+20y} - \sin(x) = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 30

```
dsolve(diff(y(x),x)=5*exp(x^2+20*y(x))+sin(x),y(x), singsol=all)
```

$$y(x) = -\cos(x) - \frac{\ln\left(-20c_1 - 100\left(\int e^{x^2} e^{-20\cos(x)} dx\right)\right)}{20}$$

✓ Solution by Mathematica

Time used: 9.745 (sec). Leaf size: 140

```
DSolve[y'[x]==5*Exp[x^2+20*y[x]]+Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[\int_1^x -\frac{1}{100}e^{-20\cos(K[1])-20y(x)}\left(\sin(K[1]) + 5e^{K[1]^2+20y(x)}\right) dK[1] + \int_1^{y(x)} -\frac{1}{100}e^{-20\cos(x)-20K[2]}\left(100e^{20\cos(x)+20K[2]}\int_1^x\left(\frac{1}{5}e^{-20\cos(K[1])-20K[2]}\left(\sin(K[1]) + 5e^{K[1]^2+20K[2]}\right) - e^{K[1]^2-20\cos(x)} - 1\right) dK[2] = c_1, y(x)\right]$$