


A Solution Manual For

**Mathematical methods for
physics and engineering, Riley,
Hobson, Bence, second edition,
2002**



Nasser M. Abbasi

October 12, 2023

Contents

1	Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490	2
2	Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523	33
3	Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550	53

1 Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

1.1	problem Problem 14.2 (a)	3
1.2	problem Problem 14.2 (b)	4
1.3	problem Problem 14.2 (c)	5
1.4	problem Problem 14.3 (a)	6
1.5	problem Problem 14.3 (b)	8
1.6	problem Problem 14.3 (c)	9
1.7	problem Problem 14.5 (a)	10
1.8	problem Problem 14.5 (b)	11
1.9	problem Problem 14.5 (c)	12
1.10	problem Problem 14.6	14
1.11	problem Problem 14.11	15
1.12	problem Problem 14.14	16
1.13	problem Problem 14.15	17
1.14	problem Problem 14.16	18
1.15	problem Problem 14.17	19
1.16	problem Problem 14.23 (a)	20
1.17	problem Problem 14.23 (b)	21
1.18	problem Problem 14.24 (a)	22
1.19	problem Problem 14.24 (b)	23
1.20	problem Problem 14.24 (c)	24
1.21	problem Problem 14.24 (d)	25
1.22	problem Problem 14.26	26
1.23	problem Problem 14.28	27
1.24	problem Problem 14.29	29
1.25	problem Problem 14.30 (a)	30
1.26	problem Problem 14.30 (b)	31
1.27	problem Problem 14.31	32

1.1 problem Problem 14.2 (a)

Internal problem ID [1977]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.2 (a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' - xy^3 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x)-x*y(x)^3=0,y(x), singsol=all)
```

$$y(x) = \frac{1}{\sqrt{-x^2 + c_1}}$$

$$y(x) = -\frac{1}{\sqrt{-x^2 + c_1}}$$

✓ Solution by Mathematica

Time used: 0.153 (sec). Leaf size: 44

```
DSolve[y'[x]-x*y[x]^3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\sqrt{-x^2 - 2c_1}}$$

$$y(x) \rightarrow \frac{1}{\sqrt{-x^2 - 2c_1}}$$

$$y(x) \rightarrow 0$$

1.2 problem Problem 14.2 (b)

Internal problem ID [1978]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.2 (b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_separable]`

$$\frac{y'}{\tan(x)} - \frac{y}{x^2 + 1} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)/tan(x)-y(x)/(1+x^2)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\int \frac{\tan(x)}{x^2+1} dx}$$

✓ Solution by Mathematica

Time used: 9.792 (sec). Leaf size: 34

```
DSolve[y'[x]/Tan[x]-y[x]/(1+x^2)==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \exp\left(\int_1^x \frac{\tan(K[1])}{K[1]^2 + 1} dK[1]\right)$$

$$y(x) \rightarrow 0$$

1.3 problem Problem 14.2 (c)

Internal problem ID [1979]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.2 (c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y'x^2 + xy^2 - 4y^2 = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve(x^2*diff(y(x),x)+x*y(x)^2=4*y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{x}{4 + x \ln(x) + c_1 x}$$

✓ Solution by Mathematica

Time used: 0.138 (sec). Leaf size: 24

```
DSolve[y'[x]+x*y[x]^2==4*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2}{(x-8)x - 2c_1}$$

$$y(x) \rightarrow 0$$

1.4 problem Problem 14.3 (a)

Internal problem ID [1980]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.3 (a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_exact, _rational, [_1st_order, ‘_with_symmetry_[F(x)*G(y),0]’

$$y(2x^2y^2 + 1)y' + x(y^4 + 1) = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 119

```
dsolve(y(x)*(2*x^2*y(x)^2+1)*diff(y(x),x)+x*(y(x)^4+1)=0,y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{-2 - 2\sqrt{-4x^4 - 8c_1x^2 + 1}}}{2x}$$

$$y(x) = \frac{\sqrt{-2 - 2\sqrt{-4x^4 - 8c_1x^2 + 1}}}{2x}$$

$$y(x) = -\frac{\sqrt{2} \sqrt{-1 + \sqrt{-4x^4 - 8c_1x^2 + 1}}}{2x}$$

$$y(x) = \frac{\sqrt{2} \sqrt{-1 + \sqrt{-4x^4 - 8c_1x^2 + 1}}}{2x}$$

✓ Solution by Mathematica

Time used: 11.805 (sec). Leaf size: 197

`DSolve[y[x]*(2*x^2*y[x]^2+1)*y'[x]+x*(y[x]^4+1)==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow -\frac{\sqrt{-\frac{1+\sqrt{-4x^4+8c_1x^2+1}}{x^2}}}{\sqrt{2}}$$

$$y(x) \rightarrow \frac{\sqrt{-\frac{1+\sqrt{-4x^4+8c_1x^2+1}}{x^2}}}{\sqrt{2}}$$

$$y(x) \rightarrow -\frac{\sqrt{\frac{-1+\sqrt{-4x^4+8c_1x^2+1}}{x^2}}}{\sqrt{2}}$$

$$y(x) \rightarrow \frac{\sqrt{\frac{-1+\sqrt{-4x^4+8c_1x^2+1}}{x^2}}}{\sqrt{2}}$$

$$y(x) \rightarrow -\sqrt[4]{-1}$$

$$y(x) \rightarrow \sqrt[4]{-1}$$

$$y(x) \rightarrow -(-1)^{3/4}$$

$$y(x) \rightarrow (-1)^{3/4}$$

1.5 problem Problem 14.3 (b)

Internal problem ID [1981]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.3 (b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$2y'x + 3x + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(2*x*diff(y(x),x)+3*x+y(x)=0,y(x), singsol=all)
```

$$y(x) = -x + \frac{c_1}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 17

```
DSolve[2*x*y'[x]+3*x+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x + \frac{c_1}{\sqrt{x}}$$

1.6 problem Problem 14.3 (c)

Internal problem ID [1982]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.3 (c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x)*G(y),0]'], [_Abel, '2nd typ`

$$(\cos(x)^2 + y \sin(2x)) y' + y^2 = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve((cos(x)^2+y(x)*sin(2*x))*diff(y(x),x)+y(x)^2=0,y(x), singsol=all)
```

$$c_1 + y(x)^2 \tan(x) + y(x) = 0$$

✓ Solution by Mathematica

Time used: 1.666 (sec). Leaf size: 80

```
DSolve[(Cos[x]^2+y[x]*Sin[2*x])*y'[x]+y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2} \cot(x) \left(1 + \sqrt{\sec^2(x) \sqrt{\cos(x)(\cos(x) + 4c_1 \sin(x))}} \right)$$

$$y(x) \rightarrow \frac{1}{2} \cot(x) \left(-1 + \sqrt{\sec^2(x) \sqrt{\cos(x)(\cos(x) + 4c_1 \sin(x))}} \right)$$

$$y(x) \rightarrow 0$$

1.7 problem Problem 14.5 (a)

Internal problem ID [1983]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.5 (a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear]`

$$(1 - x^2) y' + 4yx - (1 - x^2)^{\frac{3}{2}} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve((1-x^2)*diff(y(x),x)+2*x*y(x)+2*x*y(x)=(1-x^2)^(3/2),y(x), singsol=all)
```

$$y(x) = (x^4 - 2x^2 + 1) c_1 - \sqrt{-x^2 + 1} x (x^2 - 1)$$

✓ Solution by Mathematica

Time used: 0.114 (sec). Leaf size: 29

```
DSolve[(1-x^2)*y'[x]+2*x*y[x]+2*x*y[x]==(1-x^2)^(3/2),y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow (x^2 - 1)^2 \left(\frac{x}{\sqrt{1 - x^2}} + c_1 \right)$$

1.8 problem Problem 14.5 (b)

Internal problem ID [1984]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.5 (b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear]`

$$y' - \cot(x)y + \frac{1}{\sin(x)} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(diff(y(x),x)-y(x)*cot(x)+1/sin(x)=0,y(x), singsol=all)
```

$$y(x) = (\cot(x) + c_1) \sin(x)$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 13

```
DSolve[y'[x]-y[x]*Cot[x]+1/Sin[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \cos(x) + c_1 \sin(x)$$

1.9 problem Problem 14.5 (c)

Internal problem ID [1985]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.5 (c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational]`

$$(y^3 + x)y' - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 260

```
dsolve((x+y(x)^3)*diff(y(x),x)=y(x),y(x), singsol=all)
```

$$y(x) = \frac{\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{1}{3}}}{3} - \frac{2c_1}{\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{1}{3}}}$$

$$y(x) = -\frac{\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{1}{3}}}{6} + \frac{c_1}{\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{1}{3}}}$$

$$- \frac{i\sqrt{3} \left(\frac{\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{1}{3}}}{3} + \frac{2c_1}{\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{1}{3}}} \right)}{2}$$

$$y(x) = -\frac{\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{1}{3}}}{6} + \frac{c_1}{\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{1}{3}}}$$

$$+ \frac{i\sqrt{3} \left(\frac{\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{1}{3}}}{3} + \frac{2c_1}{\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{1}{3}}} \right)}{2}$$

✓ Solution by Mathematica

Time used: 1.662 (sec). Leaf size: 227

```
DSolve[(x+y[x]^3)*y'[x]==y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2 \cdot 3^{2/3} c_1 - \sqrt[3]{3} (-9x + \sqrt{81x^2 + 24c_1^3})^{2/3}}{3 \sqrt[3]{-9x + \sqrt{81x^2 + 24c_1^3}}}$$

$$y(x) \rightarrow \frac{-(-1)^{2/3} (-9x + \sqrt{81x^2 + 24c_1^3})^{2/3} - 2 \sqrt[3]{-3} c_1}{3^{2/3} \sqrt[3]{-9x + \sqrt{81x^2 + 24c_1^3}}}$$

$$y(x) \rightarrow \frac{2 \sqrt[3]{-3} (-9x + \sqrt{81x^2 + 24c_1^3})^{2/3} + 4(-3)^{2/3} c_1}{6 \sqrt[3]{-9x + \sqrt{81x^2 + 24c_1^3}}}$$

$$y(x) \rightarrow 0$$

1.10 problem Problem 14.6

Internal problem ID [1986]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, _Bernoulli]`

$$y' + \frac{2x^2 + y^2 + x}{yx} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

```
dsolve(diff(y(x),x) = - (2*x^2+y(x)^2+x)/(x*y(x)),y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{-9x^4 - 6x^3 + 9c_1}}{3x}$$

$$y(x) = \frac{\sqrt{-9x^4 - 6x^3 + 9c_1}}{3x}$$

✓ Solution by Mathematica

Time used: 0.244 (sec). Leaf size: 56

```
DSolve[y'[x] == - (2*x^2+y[x]^2+x)/(x*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{-\frac{1}{3}x^3(3x+2) + c_1}}{x}$$

$$y(x) \rightarrow \frac{\sqrt{-\frac{1}{3}x^3(3x+2) + c_1}}{x}$$

1.11 problem Problem 14.11

Internal problem ID [1987]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cla`

$$(y - x)y' + 2x + 3y = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 28

```
dsolve((y(x)-x)*diff(y(x),x)+2*x+3*y(x)=0,y(x), singsol=all)
```

$$y(x) = \tan \left(\text{RootOf} \left(-4_Z + \ln \left(\frac{1}{\cos(_Z)^2} \right) + 2 \ln(x) + 2c_1 \right) \right) x - x$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 45

```
DSolve[(y[x]-x)*y'[x]+2*x+3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{1}{2} \log \left(\frac{y(x)^2}{x^2} + \frac{2y(x)}{x} + 2 \right) - 2 \arctan \left(\frac{y(x)}{x} + 1 \right) = -\log(x) + c_1, y(x) \right]$$

1.12 problem Problem 14.14

Internal problem ID [1988]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.14.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class C', [_Abel, '2nd type', 'class C'], _dA`

$$y' - \frac{1}{x + 2y + 1} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 21

```
dsolve(diff(y(x),x) = 1/(x+2*y(x)+1),y(x), singsol=all)
```

$$y(x) = -\text{LambertW}\left(-\frac{c_1 e^{-\frac{x}{2}-\frac{3}{2}}}{2}\right) - \frac{x}{2} - \frac{3}{2}$$

✓ Solution by Mathematica

Time used: 60.041 (sec). Leaf size: 34

```
DSolve[y'[x] == 1/(x+2*y[x]+1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(-2W\left(-\frac{1}{2}c_1 e^{-\frac{x}{2}-\frac{3}{2}}\right) - x - 3 \right)$$

1.13 problem Problem 14.15

Internal problem ID [1989]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cla`

$$y' + \frac{x + y}{3x + 3y - 4} = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 29

```
dsolve(diff(y(x),x) = - (x+y(x))/(3*x+3*y(x)-4),y(x), singsol=all)
```

$$y(x) = e^{-\text{LambertW}\left(\frac{3e^x e^{-3} e^{-c_1}}{2}\right) + x - 3 - c_1} + 2 - x$$

✓ Solution by Mathematica

Time used: 3.532 (sec). Leaf size: 33

```
DSolve[y'[x] == - (x+y[x])/(3*x+3*y[x]-4),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2}{3}W(-e^{x-1+c_1}) - x + 2$$

$$y(x) \rightarrow 2 - x$$

1.14 problem Problem 14.16

Internal problem ID [1990]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' - \tan(x) \cos(y) (\cos(y) + \sin(y)) = 0$$

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 18

```
dsolve(diff(y(x),x) = tan(x)*cos(y(x))*( cos(y(x)) + sin(y(x)) ),y(x), singsol=all)
```

$$y(x) = -\arctan\left(\frac{\cos(x) - c_1}{\cos(x)}\right)$$

✓ Solution by Mathematica

Time used: 60.531 (sec). Leaf size: 143

```
DSolve[y'[x]==Tan[x]*Cos[y[x]]*( Cos[y[x]] + Sin[y[x]] ),y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow -\sec^{-1}\left(\sec(x)\left(-\sqrt{\cos(2x) - 2e^{\frac{c_1}{2}}\cos(x) + 1 + e^{c_1}}\right)\right)$$

$$y(x) \rightarrow \sec^{-1}\left(\sec(x)\left(-\sqrt{\cos(2x) - 2e^{\frac{c_1}{2}}\cos(x) + 1 + e^{c_1}}\right)\right)$$

$$y(x) \rightarrow -\sec^{-1}\left(\sec(x)\sqrt{\cos(2x) - 2e^{\frac{c_1}{2}}\cos(x) + 1 + e^{c_1}}\right)$$

$$y(x) \rightarrow \sec^{-1}\left(\sec(x)\sqrt{\cos(2x) - 2e^{\frac{c_1}{2}}\cos(x) + 1 + e^{c_1}}\right)$$

1.15 problem Problem 14.17

Internal problem ID [1991]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.17.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _exact, _rational, [_Abel, '2nd typ`

$$x(-2x^2y + 1)y' + y - 3x^2y^2 = 0$$

With initial conditions

$$\left[y(1) = \frac{1}{2} \right]$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 35

```
dsolve([x*(1-2*x^2*y(x))*diff(y(x),x) +y(x) = 3*x^2*y(x)^2,y(1) = 1/2],y(x), singsol=all)
```

$$y(x) = \frac{1 - \sqrt{1-x}}{2x^2}$$

$$y(x) = \frac{1 + \sqrt{1-x}}{2x^2}$$

✓ Solution by Mathematica

Time used: 0.571 (sec). Leaf size: 50

```
DSolve[{x*(1-2*x^2*y[x])*y'[x] +y[x] == 3*x^2*y[x]^2,y[1]==1/2},y[x],x,IncludeSingularSolutio
```

$$y(x) \rightarrow \frac{1}{2 \left(\sqrt{-((x-1)x^2)} + x \right)}$$

$$y(x) \rightarrow \frac{\sqrt{-((x-1)x^2)} + x}{2x^3}$$

1.16 problem Problem 14.23 (a)

Internal problem ID [1992]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.23 (a) .

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear]`

$$y' + \frac{xy}{a^2 + x^2} - x = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 26

```
dsolve(diff(y(x),x)+ (x*y(x))/(a^2+x^2)=x,y(x), singsol=all)
```

$$y(x) = \frac{a^2}{3} + \frac{x^2}{3} + \frac{c_1}{\sqrt{a^2 + x^2}}$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 31

```
DSolve[y'[x]+ (x*y[x])/(a^2+x^2)==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}(a^2 + x^2) + \frac{c_1}{\sqrt{a^2 + x^2}}$$

1.17 problem Problem 14.23 (b)

Internal problem ID [1993]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.23 (b) .

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' - \frac{4y^2}{x^2} + y^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)= 4*y(x)^2/x^2 - y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{x}{c_1x + x^2 + 4}$$

✓ Solution by Mathematica

Time used: 0.15 (sec). Leaf size: 24

```
DSolve[y'[x]== 4*y[x]^2/x^2 - y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x}{x^2 - c_1x + 4}$$

$$y(x) \rightarrow 0$$

1.18 problem Problem 14.24 (a)

Internal problem ID [1994]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.24 (a) .

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear]`

$$y' - \frac{y}{x} - 1 = 0$$

With initial conditions

$$[y(1) = -1]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve([diff(y(x),x)-y(x)/x=1,y(1) = -1],y(x), singsol=all)
```

$$y(x) = (-1 + \ln(x))x$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 11

```
DSolve[{y'[x]-y[x]/x==1,y[1]==-1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(\log(x) - 1)$$

1.19 problem Problem 14.24 (b)

Internal problem ID [1995]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.24 (b) .

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear]`

$$y' - y \tan(x) - 1 = 0$$

With initial conditions

$$\left[y\left(\frac{\pi}{4}\right) = 3 \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
dsolve([diff(y(x),x)-y(x)*tan(x)=1,y(1/4*Pi) = 3],y(x), singsol=all)
```

$$y(x) = \tan(x) + \sec(x) \sqrt{2}$$

✓ Solution by Mathematica

Time used: 0.046 (sec). Leaf size: 16

```
DSolve[{y'[x]-y[x]*Tan[x]==1,y[Pi/4]==3},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (\sin(x) + \sqrt{2}) \sec(x)$$

1.20 problem Problem 14.24 (c)

Internal problem ID [1996]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.24 (c) .

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Riccati]`

$$y' - \frac{y^2}{x^2} - \frac{1}{4} = 0$$

With initial conditions

$$[y(1) = 1]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 17

```
dsolve([diff(y(x),x)-y(x)^2/x^2=1/4,y(1) = 1],y(x), singsol=all)
```

$$y(x) = \frac{x(\ln(x) - 4)}{2\ln(x) - 4}$$

✓ Solution by Mathematica

Time used: 0.129 (sec). Leaf size: 20

```
DSolve[{y'[x]-y[x]^2/x^2==1/4,y[1]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x(\log(x) - 4)}{2(\log(x) - 2)}$$

1.21 problem Problem 14.24 (d)

Internal problem ID [1997]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.24 (d) .

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Riccati]`

$$y' - \frac{y^2}{x^2} - \frac{1}{4} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve(diff(y(x),x)-y(x)^2/x^2=1/4,y(x), singsol=all)
```

$$y(x) = \frac{x(\ln(x) + c_1 - 2)}{2\ln(x) + 2c_1}$$

✓ Solution by Mathematica

Time used: 0.095 (sec). Leaf size: 31

```
DSolve[y'[x]-y[x]^2/x^2==1/4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \left(\frac{1}{2} - \frac{1}{\log(x) + 4c_1} \right)$$

$$y(x) \rightarrow \frac{x}{2}$$

1.22 problem Problem 14.26

Internal problem ID [1998]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.26.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear]`

$$\sin(x) y' + 2 \cos(x) y - 1 = 0$$

With initial conditions

$$\left[y\left(\frac{\pi}{2}\right) = 1 \right]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 10

```
dsolve([sin(x)*diff(y(x),x)+2*y(x)*cos(x)=1,y(1/2*Pi) = 1],y(x), singsol=all)
```

$$y(x) = \frac{1}{\cos(x) + 1}$$

✓ Solution by Mathematica

Time used: 0.099 (sec). Leaf size: 11

```
DSolve[{Sin[x]*y'[x]+2*y[x]*Cos[x]==1,y[Pi/2]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{\cos(x) + 1}$$

1.23 problem Problem 14.28

Internal problem ID [1999]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.28.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cla`

$$(5x + y - 7)y' - 3 - 3x - 3y = 0$$

✓ Solution by Maple

Time used: 0.563 (sec). Leaf size: 327

```
dsolve((5*x+y(x)-7)*diff(y(x),x)=3*(x+y(x)+1),y(x), singsol=all)
```

$$y(x) = -3$$

$$+ \frac{144(-2+x) \left(-\frac{\left(1-216(-2+x)^2 c_1 + 12\sqrt{324(-2+x)^4 c_1^2 - 3(-2+x)^2 c_1}\right)^{\frac{1}{3}}}{24} - \frac{1}{24\left(1-216(-2+x)^2 c_1 + 12\sqrt{324(-2+x)^4 c_1^2 - 3(-2+x)^2 c_1}\right)} \right)}{-6 \left(1-216(-2+x)^2 c_1 + 12\sqrt{324(-2+x)^4 c_1^2 - 3(-2+x)^2 c_1}\right)^{\frac{1}{3}} - \frac{6}{\left(1-216(-2+x)^2 c_1 + 12\sqrt{324(-2+x)^4 c_1^2 - 3(-2+x)^2 c_1}\right)}}$$

✓ Solution by Mathematica

Time used: 60.176 (sec). Leaf size: 629

`DSolve[(5*x+y[x]-7)*y'[x]==3*(x+y[x]+1),y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow -5x$$

$$+ \frac{6(x-2)}{\sqrt[3]{-e^{\frac{3c_1}{4}}(x-2)^4 + 2e^{\frac{3c_1}{8}}(x-2)^2 + \sqrt{e^{\frac{3c_1}{8}}(x-2)^2 \left(-1 + e^{\frac{3c_1}{8}}(x-2)^2\right)^3 - 1}} + \frac{\sqrt[3]{-e^{\frac{3c_1}{4}}(x-2)^4 + 2e^{\frac{3c_1}{8}}(x-2)^2 + \sqrt{e^{\frac{3c_1}{8}}(x-2)^2 \left(-1 + e^{\frac{3c_1}{8}}(x-2)^2\right)^3 - 1}}}{1}$$

+ 7

$$y(x) \rightarrow -5x$$

$$+ \frac{12(x-2)}{\sqrt[3]{-e^{\frac{3c_1}{4}}(x-2)^4 + 2e^{\frac{3c_1}{8}}(x-2)^2 + \sqrt{e^{\frac{3c_1}{8}}(x-2)^2 \left(-1 + e^{\frac{3c_1}{8}}(x-2)^2\right)^3 - 1}} + \frac{i(\sqrt{3}+i)\sqrt[3]{-e^{\frac{3c_1}{4}}(x-2)^4 + 2e^{\frac{3c_1}{8}}(x-2)^2 + \sqrt{e^{\frac{3c_1}{8}}(x-2)^2 \left(-1 + e^{\frac{3c_1}{8}}(x-2)^2\right)^3 - 1}}{1+i\sqrt{3}}$$

+ 7

$$y(x) \rightarrow -5x$$

$$+ \frac{12(x-2)}{\sqrt[3]{-e^{\frac{3c_1}{4}}(x-2)^4 + 2e^{\frac{3c_1}{8}}(x-2)^2 + \sqrt{e^{\frac{3c_1}{8}}(x-2)^2 \left(-1 + e^{\frac{3c_1}{8}}(x-2)^2\right)^3 - 1}} + \frac{(-1-i\sqrt{3})\sqrt[3]{-e^{\frac{3c_1}{4}}(x-2)^4 + 2e^{\frac{3c_1}{8}}(x-2)^2 + \sqrt{e^{\frac{3c_1}{8}}(x-2)^2 \left(-1 + e^{\frac{3c_1}{8}}(x-2)^2\right)^3 - 1}}{1-i\sqrt{3}}$$

+ 7

1.24 problem Problem 14.29

Internal problem ID [2000]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.29.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, _Bernoulli]`

$$y'x + y - \frac{y^2}{x^{\frac{3}{2}}} = 0$$

With initial conditions

$$[y(1) = 1]$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 18

```
dsolve([x*diff(y(x),x)+y(x)-y(x)^2/x^(3/2)=0,y(1) = 1],y(x), singsol=all)
```

$$y(x) = \frac{5x^{\frac{3}{2}}}{3x^{\frac{5}{2}} + 2}$$

✓ Solution by Mathematica

Time used: 0.16 (sec). Leaf size: 23

```
DSolve[{x*y'[x]+y[x]-y[x]^2/x^(3/2)==0,y[1]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{5x^{3/2}}{3x^{5/2} + 2}$$

1.25 problem Problem 14.30 (a)

Internal problem ID [2001]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.30 (a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x)*G(y),0]']]`

$$(2 \sin(y) - x) y' - \tan(y) = 0$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve([(2*sin(y(x))-x)*diff(y(x),x)=tan(y(x)),y(0) = 0],y(x), singsol=all)
```

$$y(x) = 0$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 6

```
DSolve[{(2*Sin[y[x]]-x)*y'[x]==Tan[y[x]],y[0]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

1.26 problem Problem 14.30 (b)

Internal problem ID [2002]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.30 (b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x)*G(y),0]']]`

$$(2 \sin(y) - x) y' - \tan(y) = 0$$

With initial conditions

$$\left[y(0) = \frac{\pi}{2} \right]$$

✓ Solution by Maple

Time used: 6.969 (sec). Leaf size: 18

```
dsolve([(2*sin(y(x))-x)*diff(y(x),x)=tan(y(x)),y(0) = 1/2*Pi],y(x), singsol=all)
```

$$y(x) = \arcsin\left(\frac{x}{2} + \frac{\sqrt{x^2 + 4}}{2}\right)$$

✓ Solution by Mathematica

Time used: 17.52 (sec). Leaf size: 67

```
DSolve[{(2*Sin[y[x]]-x)*y'[x]==Tan[y[x]],y[0]==Pi/2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \cot^{-1}\left(\sqrt{\frac{x^2}{2} - \frac{1}{2}\sqrt{x^4 + 4x^2}}\right)$$

$$y(x) \rightarrow \cot^{-1}\left(\frac{\sqrt{x^2 + \sqrt{x^2(x^2 + 4)}}}{\sqrt{2}}\right)$$

1.27 problem Problem 14.31

Internal problem ID [2003]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 14, First order ordinary differential equations. 14.4 Exercises, page 490

Problem number: Problem 14.31.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible,`

$$y'' + y'^2 + y' = 0$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 18

```
dsolve([diff(y(x),x$2)+ (diff(y(x),x))^2+diff(y(x),x)=0,y(0) = 0],y(x), singsol=all)
```

$$y(x) = \ln(e^x c_2 - c_2 + 1) - x$$

✓ Solution by Mathematica

Time used: 0.335 (sec). Leaf size: 54

```
DSolve[{y'[x]+(y'[x])^2+y'[x]==0,y[0]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(-e^x) - \log(e^x) - i\pi$$

$$y(x) \rightarrow -\log(e^x) + \log(-e^x + e^{c_1}) - \log(-1 + e^{c_1})$$

2 Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

2.1	problem Problem 15.1	34
2.2	problem Problem 15.2(a)	35
2.3	problem Problem 15.2(b)	36
2.4	problem Problem 15.4	37
2.5	problem Problem 15.5(a)	38
2.6	problem Problem 15.5(b)	39
2.7	problem Problem 15.7	40
2.8	problem Problem 15.9(a)	41
2.9	problem Problem 15.9(b)	42
2.10	problem Problem 15.21	45
2.11	problem Problem 15.22	46
2.12	problem Problem 15.23	47
2.13	problem Problem 15.24(a)	48
2.14	problem Problem 15.24(b)	49
2.15	problem Problem 15.33	50
2.16	problem Problem 15.34	51
2.17	problem Problem 15.35	52

2.1 problem Problem 15.1

Internal problem ID [2004]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x'' + \omega_0^2 x - a \cos(\omega t) = 0$$

With initial conditions

$$[x(0) = 0, x'(0) = 0]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 28

```
dsolve([diff(x(t),t$2)+ (omega__0)^2*x(t)=a*cos(omega*t),x(0) = 0, D(x)(0) = 0],x(t), singsol
```

$$x(t) = \frac{a(\cos(\omega_0 t) - \cos(\omega t))}{\omega^2 - \omega_0^2}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 33

```
DSolve[{x''[t]+(Subscript[\[Omega],0])^2*x[t]==a*Cos[\[Omega]*t],{x[0]==0,x'[0]==0}},x[t],t,I
```

$$x(t) \rightarrow \frac{a(\cos(t\omega_0) - \cos(t\omega))}{\omega^2 - \omega_0^2}$$

2.2 problem Problem 15.2(a)

Internal problem ID [2005]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.2(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$f'' + 2f' + 5f = 0$$

With initial conditions

$$[f(0) = 1, f'(0) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve([diff(f(t),t$2)+2*diff(f(t),t)+5*f(t)=0,f(0) = 1, D(f)(0) = 0],f(t), singsol=all)
```

$$f(t) = \frac{e^{-t}(\sin(2t) + 2\cos(2t))}{2}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 25

```
DSolve[{f''[t]+2*f'[t]+5*f[t]==0,{f[0]==1,f'[0]==0}},f[t],t,IncludeSingularSolutions -> True]
```

$$f(t) \rightarrow \frac{1}{2}e^{-t}(\sin(2t) + 2\cos(2t))$$

2.3 problem Problem 15.2(b)

Internal problem ID [2006]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.2(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$f'' + 2f' + 5f - e^{-t} \cos(3t) = 0$$

With initial conditions

$$[f(0) = 0, f'(0) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 25

```
dsolve([diff(f(t),t$2)+2*diff(f(t),t)+5*f(t)=exp(-t)*cos(3*t),f(0) = 0, D(f)(0) = 0],f(t), si
```

$$f(t) = -\frac{(4 \cos(t)^3 - 2 \cos(t)^2 - 3 \cos(t) + 1) e^{-t}}{5}$$

✓ Solution by Mathematica

Time used: 0.075 (sec). Leaf size: 25

```
DSolve[{f''[t]+2*f'[t]+5*f[t]==Exp[-t]*Cos[3*t],{f[0]==0,f'[0]==0}},f[t],t,IncludeSingularSol
```

$$f(t) \rightarrow \frac{1}{5}e^{-t}(\cos(2t) - \cos(3t))$$

2.4 problem Problem 15.4

Internal problem ID [2007]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$f'' + 6f' + 9f - e^{-t} = 0$$

With initial conditions

$$[f(0) = 0, f'(0) = \lambda]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 26

```
dsolve([diff(f(t),t$2)+6*diff(f(t),t)+9*f(t)=exp(-t),f(0) = 0, D(f)(0) = lambda],f(t), singso
```

$$f(t) = \frac{(-1 + (4\lambda - 2)t)e^{-3t}}{4} + \frac{e^{-t}}{4}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 28

```
DSolve[{f''[t]+6*f'[t]+9*f[t]==Exp[-t],{f[0]==0,f'[0]==\[Lambda]}},f[t],t,IncludeSingularSolu
```

$$f(t) \rightarrow \frac{1}{4}e^{-3t}((4\lambda - 2)t + e^{2t} - 1)$$

2.5 problem Problem 15.5(a)

Internal problem ID [2008]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.5(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$f'' + 8f' + 12f - 12e^{-4t} = 0$$

With initial conditions

$$[f(0) = 0, f'(0) = 0]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 23

```
dsolve([diff(f(t),t$2)+8*diff(f(t),t)+12*f(t)=12*exp(-4*t),f(0) = 0, D(f)(0) = 0],f(t), sings
```

$$f(t) = \frac{3e^{-6t}}{2} + \frac{3e^{-2t}}{2} - 3e^{-4t}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 16

```
DSolve[{f''[t]+8*f'[t]+12*f[t]==12*Exp[-4*t]},{f[0]==0,f'[0]==0},f[t],t,IncludeSingularSoluti
```

$$f(t) \rightarrow 6e^{-4t} \sinh^2(t)$$

2.6 problem Problem 15.5(b)

Internal problem ID [2009]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.5(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$f'' + 8f' + 12f - 12e^{-4t} = 0$$

With initial conditions

$$[f(0) = 0, f'(0) = -2]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 21

```
dsolve([diff(f(t),t$2)+8*diff(f(t),t)+12*f(t)=12*exp(-4*t),f(0) = 0, D(f)(0) = -2],f(t), sing
```

$$f(t) = 2e^{-6t} + e^{-2t} - 3e^{-4t}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 25

```
DSolve[{f'[t]+8*f'[t]+12*f[t]==12*Exp[-4*t]},{f[0]==0,f'[0]==-2}],f[t],t,IncludeSingularSolut
```

$$f(t) \rightarrow e^{-6t}(-3e^{2t} + e^{4t} + 2)$$

2.7 problem Problem 15.7

Internal problem ID [2010]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 2y' + y - 4e^{-x} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)+y(x)=4*exp(-x),y(x), singsol=all)
```

$$y(x) = e^{-x}c_2 + xe^{-x}c_1 + 2x^2e^{-x}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 22

```
DSolve[y''[x]+2*y'[x]+y[x]==4*Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(x(2x + c_2) + c_1)$$

2.8 problem Problem 15.9(a)

Internal problem ID [2011]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.9(a).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$y''' - 12y' + 16y - 32x + 8 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$3)-12*diff(y(x),x)+16*y(x)=32*x-8,y(x), singsol=all)
```

$$y(x) = 1 + 2x + c_1e^{-4x} + c_2e^{2x} + c_3e^{2x}x$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 31

```
DSolve[y'''[x]-12*y'[x]+16*y[x]==32*x-8,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2x + c_1e^{-4x} + e^{2x}(c_3x + c_2) + 1$$

2.9 problem Problem 15.9(b)

Internal problem ID [2012]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.9(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`, `[_2nd_order, _reducible]`

$$-\frac{y'^2}{y^2} + \frac{y''}{y} + \frac{2a \coth(2ax) y'}{y} - 2a^2 = 0$$

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 407

`dsolve(diff(1/y(x)*diff(y(x),x),x)+(2*a*coth(2*a*x))*(1/y(x)*diff(y(x),x))=2*a^2,y(x), sings`

$$y(x) = \frac{2^{\frac{3}{4}} e^{-ax} e^{-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(e^{4ax}+1)e^{-2ax}}{2}\right)}{c_1 a}} e^{-\frac{c_2}{c_1}} \left((e^{8ax} - 2e^{4ax} + 1) e^{\frac{3\sqrt{2} \operatorname{arctanh}\left(\frac{(e^{4ax}+1)e^{-2ax}}{2}\right)}{c_1 a}} \right)^{\frac{1}{4}}}{2}$$

$$y(x) = \frac{2^{\frac{3}{4}} e^{-ax} e^{-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(e^{4ax}+1)e^{-2ax}}{2}\right)}{c_1 a}} e^{-\frac{c_2}{c_1}} \left((e^{8ax} - 2e^{4ax} + 1) e^{\frac{3\sqrt{2} \operatorname{arctanh}\left(\frac{(e^{4ax}+1)e^{-2ax}}{2}\right)}{c_1 a}} \right)^{\frac{1}{4}}}{2}$$

$$y(x) = \frac{i 2^{\frac{3}{4}} e^{-ax} e^{-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(e^{4ax}+1)e^{-2ax}}{2}\right)}{c_1 a}} e^{-\frac{c_2}{c_1}} \left((e^{8ax} - 2e^{4ax} + 1) e^{\frac{3\sqrt{2} \operatorname{arctanh}\left(\frac{(e^{4ax}+1)e^{-2ax}}{2}\right)}{c_1 a}} \right)^{\frac{1}{4}}}{2}$$

$$y(x) = \frac{i 2^{\frac{3}{4}} e^{-ax} e^{-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(e^{4ax}+1)e^{-2ax}}{2}\right)}{c_1 a}} e^{-\frac{c_2}{c_1}} \left((e^{8ax} - 2e^{4ax} + 1) e^{\frac{3\sqrt{2} \operatorname{arctanh}\left(\frac{(e^{4ax}+1)e^{-2ax}}{2}\right)}{c_1 a}} \right)^{\frac{1}{4}}}{2}$$

✓ Solution by Mathematica

Time used: 0.487 (sec). Leaf size: 287

```
DSolve[D[1/y[x]*y'[x],x]+(2*a*Coth[1/y[x]*y'[x]])==2*a^2,y[x],x,IncludeSingularSolutions ->T
```

$y(x)$

$$\rightarrow c_2 \exp \left(-\text{PolyLog} \left(2, \frac{(a+1) \exp \left(-2 \text{InverseFunction} \left[\frac{-((a+1) \log(1-\tanh(\#1)))+(a-1) \log(\tanh(\#1)+1)+2 \log(1-a \tanh(\#1))}{2(a^2-1)} \right]}{a-1} \right) \right) \right)$$

2.10 problem Problem 15.21

Internal problem ID [2013]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - y'x + y - x = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=x,y(x), singsol=all)
```

$$y(x) = c_2 x + \ln(x) c_1 x + \frac{\ln(x)^2 x}{2}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 25

```
DSolve[x^2*y''[x]-x*y'[x]+y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}x(\log^2(x) + 2c_2 \log(x) + 2c_1)$$

2.11 problem Problem 15.22

Internal problem ID [2014]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$(x + 1)^2 y'' + 3(x + 1) y' + y - x^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 50

```
dsolve((x+1)^2*diff(y(x),x$2)+3*(x+1)*diff(y(x),x)+y(x)=x^2,y(x), singsol=all)
```

$$y(x) = \frac{\ln(x+1)c_1}{x+1} + \frac{c_2}{x+1} - \frac{-2x^3 + 3x^2 + 6\ln(x+1) - 6x}{18(x+1)}$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 42

```
DSolve[(x+1)^2*y''[x]+3*(x+1)*y'[x]+y[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x(x(2x-3)+6)+6(-1+3c_2)\log(x+1)+18c_1}{18(x+1)}$$

2.12 problem Problem 15.23

Internal problem ID [2015]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`, `[_2nd_order, _linear]`, ‘

$$(-2 + x)y'' + 3y' + \frac{4y}{x^2} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve((x-2)*diff(y(x),x$2)+3*diff(y(x),x)+4*y(x)/x^2=0,y(x), singsol=all)
```

$$y(x) = \frac{(3x - 4)c_1}{x(-2 + x)^2} + \frac{x^2c_2}{(-2 + x)^2}$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 44

```
DSolve[(x-2)*y''[x]+3*y'[x]+4*y[x]/x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{6c_1x^3 + c_2(3x - 4)}{6\sqrt{2 - x}(x - 2)^{3/2}x}$$

2.13 problem Problem 15.24(a)

Internal problem ID [2016]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.24(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - y - x^n = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 81

```
dsolve(diff(y(x),x$2)-y(x)=x^n,y(x), singsol=all)
```

$$y(x) = e^{-x}c_2 + c_1e^x + \frac{e^{-x} \left(-((n\Gamma(n, -x) - \Gamma(n + 1))(-x)^{-n} + e^x)x^n(n + 1) + e^{\frac{3x}{2}}x^{\frac{n}{2}} \text{WhittakerM}\left(\frac{n}{2}, \frac{n}{2} + \frac{1}{2}, x\right) \right)}{2n + 2}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 50

```
DSolve[y''[x]-y[x]==x^n,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}e^{-x}(x^{n+1} \text{ExpIntegralE}(-n, -x) + e^{2x}(-\Gamma(n + 1, x) + 2c_1) + 2c_2)$$

2.14 problem Problem 15.24(b)

Internal problem ID [2017]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.24(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 2y' + y - 2e^x x = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x)=2*x*exp(x),y(x), singsol=all)
```

$$y(x) = e^x c_2 + x e^x c_1 + \frac{e^x x^3}{3}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 25

```
DSolve[y''[x]-2*y'[x]+y[x]==2*x*Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}e^x(x^3 + 3c_2x + 3c_1)$$

2.15 problem Problem 15.33

Internal problem ID [2018]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.33.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _exact, _nonlinear]]`

Solve

$$2yy''' + 2(y + 3y')y'' + 2(y')^2 - \sin(x) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 97

```
dsolve(2*y(x)*diff(y(x),x$3)+2*(y(x)+3*diff(y(x),x))*diff(y(x),x$2)+2*(diff(y(x),x))^2=sin(x))
```

$$y(x) = -\frac{e^{-x}\sqrt{2}\sqrt{e^x(-4xe^xc_1 + \cos(x)e^x - \sin(x)e^x + 4c_1e^x - 4c_3e^x + 4c_2)}}{2}$$

$$y(x) = \frac{e^{-x}\sqrt{2}\sqrt{e^x(-4xe^xc_1 + \cos(x)e^x - \sin(x)e^x + 4c_1e^x - 4c_3e^x + 4c_2)}}{2}$$

✓ Solution by Mathematica

Time used: 0.456 (sec). Leaf size: 84

```
DSolve[2*y[x]*y'''[x]+2*(y[x]+3*y'[x])*y''[x]+2*(y'[x])^2==Sin[x],y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow -\frac{\sqrt{-\sin(x) + \cos(x) + 2c_1(x-1) + 2c_3e^{-x} - 4c_2}}{\sqrt{2}}$$

$$y(x) \rightarrow \frac{\sqrt{-\sin(x) + \cos(x) + 2c_1(x-1) + 2c_3e^{-x} - 4c_2}}{\sqrt{2}}$$

2.16 problem Problem 15.34

Internal problem ID [2019]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.34.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_y]]`

$$xy''' + 2y'' - Ax = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(x*dif(y(x),x$3)+2*dif(y(x),x$2)=A*x,y(x), singsol=all)
```

$$y(x) = \frac{Ax^3}{18} - \ln(x)c_1 + c_2x + c_3$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 26

```
DSolve[x*y'''[x]+2*y''[x]==A*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{Ax^3}{18} + c_3x - c_1 \log(x) + c_2$$

2.17 problem Problem 15.35

Internal problem ID [2020]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 15, Higher order ordinary differential equations. 15.4 Exercises, page 523

Problem number: Problem 15.35.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y'x + (4x^2 + 6)y - e^{-x^2} \sin(2x) = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 42

```
dsolve(diff(y(x),x$2)+4*x*diff(y(x),x)+(4*x^2+6)*y(x)=exp(-x^2)*sin(2*x),y(x), singsol=all)
```

$$y(x) = e^{-x^2} \cos(2x) c_2 + e^{-x^2} \sin(2x) c_1 - \frac{e^{-x^2} x \cos(2x)}{4}$$

✓ Solution by Mathematica

Time used: 0.059 (sec). Leaf size: 52

```
DSolve[y''[x]+4*x*y'[x]+(4*x^2+6)*y[x]==Exp[-x^2]*Sin[2*x],y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow \frac{1}{32} e^{-x(x+2i)} (-4x - e^{4ix} (4x + i + 8ic_2) + i + 32c_1)$$

3 Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550

3.1	problem Problem 16.1	54
3.2	problem Problem 16.2	55
3.3	problem Problem 16.3	56
3.4	problem Problem 16.4	57
3.5	problem Problem 16.6	58
3.6	problem Problem 16.8	59
3.7	problem Problem 16.9	60
3.8	problem Problem 16.10	61
3.9	problem Problem 16.11	63
3.10	problem Problem 16.12 (a)	64
3.11	problem Problem 16.12 (b)	65
3.12	problem Problem 16.13	66
3.13	problem Problem 16.14	67
3.14	problem Problem 16.15	69

3.1 problem Problem 16.1

Internal problem ID [2021]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550

Problem number: Problem 16.1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [_Gegenbauer]

$$(-z^2 + 1)y'' - 3zy' + \lambda y = 0$$

With the expansion point for the power series method at $z = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 63

Order:=6;

```
dsolve((1-z^2)*diff(y(z),z$2)-3*z*diff(y(z),z)+lambda*y(z)=0,y(z),type='series',z=0);
```

$$y(z) = \left(1 - \frac{\lambda z^2}{2} + \frac{\lambda(\lambda - 8)z^4}{24}\right) y(0) + \left(z - \frac{(\lambda - 3)z^3}{6} + \frac{(\lambda - 3)(\lambda - 15)z^5}{120}\right) D(y)(0) + O(z^6)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 80

```
AsymptoticDSolveValue[(1-z^2)*y'[z]-3*z*y'[z]+[Lambda]*y[z]==0,y[z],{z,0,5}]
```

$$y(z) \rightarrow c_2 \left(\frac{\lambda^2 z^5}{120} - \frac{3\lambda z^5}{20} + \frac{3z^5}{8} - \frac{\lambda z^3}{6} + \frac{z^3}{2} + z \right) + c_1 \left(\frac{\lambda^2 z^4}{24} - \frac{\lambda z^4}{3} - \frac{\lambda z^2}{2} + 1 \right)$$

3.2 problem Problem 16.2

Internal problem ID [2022]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550

Problem number: Problem 16.2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4zy'' + 2(1 - z)y' - y = 0$$

With the expansion point for the power series method at $z = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 44

```
Order:=6;
dsolve(4*z*diff(y(z),z$2)+2*(1-z)*diff(y(z),z)-y(z)=0,y(z),type='series',z=0);
```

$$y(z) = c_1 \sqrt{z} \left(1 + \frac{1}{3}z + \frac{1}{15}z^2 + \frac{1}{105}z^3 + \frac{1}{945}z^4 + \frac{1}{10395}z^5 + O(z^6) \right) \\ + c_2 \left(1 + \frac{1}{2}z + \frac{1}{8}z^2 + \frac{1}{48}z^3 + \frac{1}{384}z^4 + \frac{1}{3840}z^5 + O(z^6) \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 85

```
AsymptoticDSolveValue[4*z*y''[z]+2*(1-z)*y'[z]-y[z]==0,y[z],{z,0,5}]
```

$$y(z) \rightarrow c_1 \sqrt{z} \left(\frac{z^5}{10395} + \frac{z^4}{945} + \frac{z^3}{105} + \frac{z^2}{15} + \frac{z}{3} + 1 \right) + c_2 \left(\frac{z^5}{3840} + \frac{z^4}{384} + \frac{z^3}{48} + \frac{z^2}{8} + \frac{z}{2} + 1 \right)$$

3.3 problem Problem 16.3

Internal problem ID [2023]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550

Problem number: Problem 16.3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F(`

$$zy'' - 2y' + 9z^5y = 0$$

With the expansion point for the power series method at $z = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 28

```
Order:=7;
dsolve(z*dif(y(z),z$2)-2*dif(y(z),z)+9*z^5*y(z)=0,y(z),type='series',z=0);
```

$$y(z) = c_1 z^3 \left(1 - \frac{1}{6} z^6 + O(z^7) \right) + c_2 (12 - 6z^6 + O(z^7))$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 12

```
AsymptoticDSolveValue[z*y''[z]-2*y'[z]+9*z^5*y[z]==0,y[z],{z,0,6}]
```

$$y(z) \rightarrow c_2 z^3 + c_1$$

3.4 problem Problem 16.4

Internal problem ID [2024]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550

Problem number: Problem 16.4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$f'' + 2(z-1)f' + 4f = 0$$

With the expansion point for the power series method at $z = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 52

```
Order:=6;
dsolve(diff(f(z),z$2)+2*(z-1)*diff(f(z),z)+4*f(z)=0,f(z),type='series',z=0);
```

$$f(z) = \left(1 - 2z^2 - \frac{4}{3}z^3 + \frac{2}{3}z^4 + \frac{14}{15}z^5\right) f(0) + \left(z + z^2 - \frac{1}{3}z^3 - \frac{5}{6}z^4 - \frac{1}{6}z^5\right) D(f)(0) + O(z^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 127

```
AsymptoticDSolveValue[f''[z]+2*(z-a)*f'[z]+4*f[z]==0,f[z],{z,0,5}]
```

$$f(z) \rightarrow c_1 \left(-\frac{4}{15}a^3z^5 - \frac{2a^2z^4}{3} + \frac{6az^5}{5} - \frac{4az^3}{3} + \frac{4z^4}{3} - 2z^2 + 1 \right) \\ + c_2 \left(\frac{2a^4z^5}{15} + \frac{a^3z^4}{3} - \frac{4a^2z^5}{5} + \frac{2a^2z^3}{3} - \frac{7az^4}{6} + az^2 + \frac{z^5}{2} - z^3 + z \right)$$

3.5 problem Problem 16.6

Internal problem ID [2025]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550

Problem number: Problem 16.6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$z^2 y'' - \frac{3zy'}{2} + (z+1)y = 0$$

With the expansion point for the power series method at $z = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 47

Order:=6;

```
dsolve(z^2*diff(y(z),z$2)-3/2*z*diff(y(z),z)+(1+z)*y(z)=0,y(z),type='series',z=0);
```

$$y(z) = c_1 \sqrt{z} \left(1 + 2z - 2z^2 + \frac{4}{9}z^3 - \frac{2}{45}z^4 + \frac{4}{1575}z^5 + O(z^6) \right) \\ + c_2 z^2 \left(1 - \frac{2}{5}z + \frac{2}{35}z^2 - \frac{4}{945}z^3 + \frac{2}{10395}z^4 - \frac{4}{675675}z^5 + O(z^6) \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 84

```
AsymptoticDSolveValue[z^2*y''[z]-3/2*z*y'[z]+(1+z)*y[z]==0,y[z],{z,0,5}]
```

$$y(z) \rightarrow c_1 \left(-\frac{4z^5}{675675} + \frac{2z^4}{10395} - \frac{4z^3}{945} + \frac{2z^2}{35} - \frac{2z}{5} + 1 \right) z^2 \\ + c_2 \left(\frac{4z^5}{1575} - \frac{2z^4}{45} + \frac{4z^3}{9} - 2z^2 + 2z + 1 \right) \sqrt{z}$$

3.6 problem Problem 16.8

Internal problem ID [2026]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550

Problem number: Problem 16.8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_Lienard]`

$$zy'' - 2y' + yz = 0$$

With the expansion point for the power series method at $z = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

```
Order:=6;
dsolve(z*dif(y(z),z$2)-2*dif(y(z),z)+z*y(z)=0,y(z),type='series',z=0);
```

$$y(z) = c_1 z^3 \left(1 - \frac{1}{10} z^2 + \frac{1}{280} z^4 + O(z^6) \right) + c_2 \left(12 + 6z^2 - \frac{3}{2} z^4 + O(z^6) \right)$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 44

```
AsymptoticDSolveValue[z*y'[z]-2*y'[z]+z*y[z]==0,y[z],{z,0,5}]
```

$$y(z) \rightarrow c_1 \left(-\frac{z^4}{8} + \frac{z^2}{2} + 1 \right) + c_2 \left(\frac{z^7}{280} - \frac{z^5}{10} + z^3 \right)$$

3.7 problem Problem 16.9

Internal problem ID [2027]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550

Problem number: Problem 16.9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$y'' - 2zy' - 2y = 0$$

With the expansion point for the power series method at $z = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
Order:=6;
dsolve(diff(y(z),z$2)-2*z*diff(y(z),z)-2*y(z)=0,y(z),type='series',z=0);
```

$$y(z) = \left(1 + z^2 + \frac{1}{2}z^4\right) y(0) + \left(z + \frac{2}{3}z^3 + \frac{4}{15}z^5\right) D(y)(0) + O(z^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 38

```
AsymptoticDSolveValue[y'[z]-2*z*y'[z]-2*y[z]==0,y[z],{z,0,5}]
```

$$y(z) \rightarrow c_2 \left(\frac{4z^5}{15} + \frac{2z^3}{3} + z \right) + c_1 \left(\frac{z^4}{2} + z^2 + 1 \right)$$

3.8 problem Problem 16.10

Internal problem ID [2028]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550

Problem number: Problem 16.10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [_Jacobi]

$$z(1-z)y'' + (1-z)y' + \lambda y = 0$$

With the expansion point for the power series method at $z = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 261

Order:=6;

dsolve(z*(1-z)*diff(y(z),z\$2)+(1-z)*diff(y(z),z)+lambda*y(z)=0,y(z),type='series',z=0);

$$\begin{aligned}
 y(z) = & \left(2\lambda z + \left(\frac{1}{4}\lambda - \frac{3}{4}\lambda^2 \right) z^2 + \left(-\frac{37}{108}\lambda^2 + \frac{2}{27}\lambda + \frac{11}{108}\lambda^3 \right) z^3 \right. \\
 & \left. + \left(\frac{139}{1728}\lambda^3 - \frac{649}{3456}\lambda^2 + \frac{1}{32}\lambda - \frac{25}{3456}\lambda^4 \right) z^4 \right. \\
 & \left. + \left(-\frac{13}{1600}\lambda^4 + \frac{8467}{144000}\lambda^3 - \frac{2527}{21600}\lambda^2 + \frac{2}{125}\lambda + \frac{137}{432000}\lambda^5 \right) z^5 + O(z^6) \right) c_2 \\
 & + \left(1 - \lambda z + \frac{1}{4}(-1 + \lambda)\lambda z^2 - \frac{1}{36}\lambda(\lambda^2 - 5\lambda + 4)z^3 + \frac{1}{576}\lambda(\lambda^3 - 14\lambda^2 + 49\lambda - 36)z^4 \right. \\
 & \left. - \frac{1}{14400}\lambda(-1 + \lambda)(\lambda - 4)(\lambda - 16)(\lambda - 9)z^5 + O(z^6) \right) (c_2 \ln(z) + c_1)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 940

AsymptoticDSolveValue[z*(1-z)*y''[z]+(1-z)*y'[z]+\[Lambda]*y[z]==0,y[z],{z,0,5}]

$$\begin{aligned}
y(z) \rightarrow & \left(\frac{1}{25} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \frac{1}{9} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) \lambda \right. \right. \\
& \quad \left. \left. - \frac{1}{16} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \frac{1}{9} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) \lambda - \lambda \right) \lambda - \lambda \right) z^5 \right. \\
& + \frac{1}{16} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \frac{1}{9} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) \lambda - \lambda \right) z^4 + \frac{1}{9} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) z^3 \\
& + \frac{1}{4}(\lambda^2 - \lambda) z^2 - \lambda z + 1 \Big) c_1 + c_2 \left(-\frac{2}{125} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \frac{1}{9} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) \lambda \right. \right. \\
& \quad \left. \left. - \frac{1}{16} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \frac{1}{9} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) \lambda - \lambda \right) \lambda - \lambda \right) z^5 \right. \\
& \quad \left. + \frac{1}{25} \left(\frac{\lambda^3}{2} - 2\lambda^2 + \frac{1}{4}(\lambda^2 - \lambda) \lambda + \frac{2}{27} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) \lambda \right. \right. \\
& \quad \left. \left. - \frac{1}{9} \left(\frac{\lambda^3}{2} - 2\lambda^2 + \frac{1}{4}(\lambda^2 - \lambda) \lambda \right) \lambda + \frac{1}{32} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \frac{1}{9} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) \lambda - \lambda \right) \lambda \right. \right. \\
& \quad \left. \left. - \frac{1}{16} \left(\frac{\lambda^3}{2} - 2\lambda^2 + \frac{1}{4}(\lambda^2 - \lambda) \lambda + \frac{2}{27} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) \lambda - \frac{1}{9} \left(\frac{\lambda^3}{2} - 2\lambda^2 + \frac{1}{4}(\lambda^2 - \lambda) \lambda \right) \lambda \right) \lambda \right) z^5 \right. \\
& \quad \left. - \frac{1}{32} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \frac{1}{9} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) \lambda - \lambda \right) z^4 + \frac{1}{16} \left(\frac{\lambda^3}{2} - 2\lambda^2 \right. \right. \\
& \quad \left. \left. + \frac{1}{4}(\lambda^2 - \lambda) \lambda + \frac{2}{27} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) \lambda - \frac{1}{9} \left(\frac{\lambda^3}{2} - 2\lambda^2 + \frac{1}{4}(\lambda^2 - \lambda) \lambda \right) \lambda \right) z^4 \right. \\
& \quad \left. - \frac{2}{27} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) z^3 + \frac{1}{9} \left(\frac{\lambda^3}{2} - 2\lambda^2 + \frac{1}{4}(\lambda^2 - \lambda) \lambda \right) z^3 - \frac{\lambda^2 z^2}{2} - \frac{1}{4}(\lambda^2 - \lambda) z^2 + 2\lambda z \right. \\
& \quad \left. + \left(\frac{1}{25} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \frac{1}{9} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) \lambda - \frac{1}{16} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \frac{1}{9} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) \lambda - \lambda \right) \lambda \right. \right. \right. \\
& \quad \quad \left. \left. + \frac{1}{16} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \frac{1}{9} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) \lambda - \lambda \right) z^4 \right. \right. \\
& \quad \left. \left. + \frac{1}{9} \left(\lambda^2 - \frac{1}{4}(\lambda^2 - \lambda) \lambda - \lambda \right) z^3 + \frac{1}{4}(\lambda^2 - \lambda) z^2 - \lambda z + 1 \right) \log(z) \right)
\end{aligned}$$

3.9 problem Problem 16.11

Internal problem ID [2029]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550

Problem number: Problem 16.11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$zy'' + (2z - 3)y' + \frac{4y}{z} = 0$$

With the expansion point for the power series method at $z = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 69

```
Order:=6;
dsolve(z*diff(y(z),z$2)+(2*z-3)*diff(y(z),z)+4/z*y(z)=0,y(z),type='series',z=0);
```

$$y(z) = z^2 \left((c_2 \ln(z) + c_1) \left(1 - 4z + 6z^2 - \frac{16}{3}z^3 + \frac{10}{3}z^4 - \frac{8}{5}z^5 + O(z^6) \right) + \left(6z - 13z^2 + \frac{124}{9}z^3 - \frac{173}{18}z^4 + \frac{374}{75}z^5 + O(z^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 116

```
AsymptoticDSolveValue[z*y''[z]+(2*z-3)*y'[z]+4/z*y[z]==0,y[z],{z,0,5}]
```

$$y(z) \rightarrow c_1 \left(-\frac{8z^5}{5} + \frac{10z^4}{3} - \frac{16z^3}{3} + 6z^2 - 4z + 1 \right) z^2 + c_2 \left(\left(\frac{374z^5}{75} - \frac{173z^4}{18} + \frac{124z^3}{9} - 13z^2 + 6z \right) z^2 + \left(-\frac{8z^5}{5} + \frac{10z^4}{3} - \frac{16z^3}{3} + 6z^2 - 4z + 1 \right) z^2 \log(z) \right)$$

3.10 problem Problem 16.12 (a)

Internal problem ID [2030]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550

Problem number: Problem 16.12 (a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(z^2 + 5z + 6)y'' + 2y = 0$$

With the expansion point for the power series method at $z = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 49

```
Order:=6;
dsolve((z^2+5*z+6)*diff(y(z),z$2)+2*y(z)=0,y(z),type='series',z=0);
```

$$y(z) = \left(1 - \frac{1}{6}z^2 + \frac{5}{108}z^3 - \frac{13}{1296}z^4 + \frac{5}{2592}z^5\right) y(0) \\ + \left(z - \frac{1}{18}z^3 + \frac{5}{216}z^4 - \frac{17}{2160}z^5\right) D(y)(0) + O(z^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 63

```
AsymptoticDSolveValue[(z^2+5*z+6)*y''[z]+2*y[z]==0,y[z],{z,0,5}]
```

$$y(z) \rightarrow c_2 \left(-\frac{17z^5}{2160} + \frac{5z^4}{216} - \frac{z^3}{18} + z \right) + c_1 \left(\frac{5z^5}{2592} - \frac{13z^4}{1296} + \frac{5z^3}{108} - \frac{z^2}{6} + 1 \right)$$

3.11 problem Problem 16.12 (b)

Internal problem ID [2031]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550

Problem number: Problem 16.12 (b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$(z^2 + 5z + 7)y'' + 2y = 0$$

With the expansion point for the power series method at $z = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

```
Order:=6;
dsolve((z^2+5*z+7)*diff(y(z),z$2)+2*y(z)=0,y(z),type='series',z=0);
```

$$y(z) = \left(1 - \frac{1}{7}z^2 + \frac{5}{147}z^3 - \frac{11}{2058}z^4 + \frac{5}{14406}z^5\right) y(0) \\ + \left(z - \frac{1}{21}z^3 + \frac{5}{294}z^4 - \frac{47}{10290}z^5\right) D(y)(0) + O(z^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 63

```
AsymptoticDSolveValue[(z^2+5*z+7)*y'[z]+2*y[z]==0,y[z],{z,0,5}]
```

$$y(z) \rightarrow c_2 \left(-\frac{47z^5}{10290} + \frac{5z^4}{294} - \frac{z^3}{21} + z \right) + c_1 \left(\frac{5z^5}{14406} - \frac{11z^4}{2058} + \frac{5z^3}{147} - \frac{z^2}{7} + 1 \right)$$

3.12 problem Problem 16.13

Internal problem ID [2032]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550

Problem number: Problem 16.13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y'' + \frac{y}{z^3} = 0$$

With the expansion point for the power series method at $z = 0$.

X Solution by Maple

```
Order:=6;
dsolve(diff(y(z),z$2)+1/z^3*y(z)=0,y(z),type='series',z=0);
```

No solution found

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 222

```
AsymptoticDSolveValue[y''[z]+1/z^3*y[z]==0,y[z],{z,0,5}]
```

$$y(z) \rightarrow c_1 e^{-\frac{2i}{\sqrt{z}} z^{3/4}} \left(-\frac{468131288625iz^{9/2}}{8796093022208} + \frac{66891825iz^{7/2}}{4294967296} - \frac{72765iz^{5/2}}{8388608} + \frac{105iz^{3/2}}{8192} \right. \\ \left. + \frac{33424574007825z^5}{281474976710656} - \frac{14783093325z^4}{549755813888} + \frac{2837835z^3}{268435456} - \frac{4725z^2}{524288} + \frac{15z}{512} - \frac{3i\sqrt{z}}{16} \right. \\ \left. + 1 \right) + c_2 e^{\frac{2i}{\sqrt{z}} z^{3/4}} \left(\frac{468131288625iz^{9/2}}{8796093022208} - \frac{66891825iz^{7/2}}{4294967296} + \frac{72765iz^{5/2}}{8388608} - \frac{105iz^{3/2}}{8192} + \frac{33424574007825z^5}{281474976710656} - \frac{14783093325z^4}{549755813888} + \frac{2837835z^3}{268435456} - \frac{4725z^2}{524288} + \frac{15z}{512} - \frac{3i\sqrt{z}}{16} \right)$$

3.13 problem Problem 16.14

Internal problem ID [2033]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550

Problem number: Problem 16.14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [_Laguerre]

$$zy'' + (1 - z)y' + \lambda y = 0$$

With the expansion point for the power series method at $z = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 309

```
Order:=6;
dsolve(z*dif(y(z),z$2)+(1-z)*dif(y(z),z)+lambda*y(z)=0,y(z),type='series',z=0);
```

$$\begin{aligned}
 y(z) = & \left((2\lambda + 1)z + \left(\frac{1}{4}\lambda + \frac{1}{4} - \frac{3}{4}\lambda^2 \right) z^2 + \left(-\frac{2}{9}\lambda^2 + \frac{1}{27}\lambda + \frac{1}{18} + \frac{11}{108}\lambda^3 \right) z^3 \right. \\
 & \left. + \left(\frac{7}{192}\lambda^3 - \frac{167}{3456}\lambda^2 + \frac{1}{192}\lambda + \frac{1}{96} - \frac{25}{3456}\lambda^4 \right) z^4 \right. \\
 & \left. + \left(\frac{1}{1500}\lambda - \frac{37}{4320}\lambda^2 + \frac{719}{86400}\lambda^3 - \frac{61}{21600}\lambda^4 + \frac{137}{432000}\lambda^5 + \frac{1}{600} \right) z^5 + O(z^6) \right) c_2 \\
 & + \left(1 - \lambda z + \frac{1}{4}(-1 + \lambda)\lambda z^2 - \frac{1}{36}(\lambda - 2)(-1 + \lambda)\lambda z^3 + \frac{1}{576}(\lambda - 3)(\lambda - 2)(-1 + \lambda)\lambda z^4 \right. \\
 & \left. - \frac{1}{14400}(\lambda - 4)(\lambda - 3)(\lambda - 2)(-1 + \lambda)\lambda z^5 + O(z^6) \right) (c_2 \ln(z) + c_1)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 415

AsymptoticDSolveValue[z*y''[z]+(1-z)*y'[z]+\[Lambda]*y[z]==0,y[z],{z,0,5}]

$$\begin{aligned}
y(z) \rightarrow & c_1 \left(-\frac{(\lambda-4)(\lambda-3)(\lambda-2)(\lambda-1)\lambda z^5}{14400} + \frac{1}{576}(\lambda-3)(\lambda-2)(\lambda-1)\lambda z^4 \right. \\
& \left. - \frac{1}{36}(\lambda-2)(\lambda-1)\lambda z^3 + \frac{1}{4}(\lambda-1)\lambda z^2 - \lambda z + 1 \right) + c_2 \left(\frac{(\lambda-4)(\lambda-3)(\lambda-2)(\lambda-1)z^5}{14400} \right. \\
& + \frac{(\lambda-4)(\lambda-3)(\lambda-2)\lambda z^5}{14400} + \frac{(\lambda-4)(\lambda-3)(\lambda-1)\lambda z^5}{14400} + \frac{(\lambda-4)(\lambda-2)(\lambda-1)\lambda z^5}{14400} \\
& \left. + \frac{137(\lambda-4)(\lambda-3)(\lambda-2)(\lambda-1)\lambda z^5}{432000} + \frac{(\lambda-3)(\lambda-2)(\lambda-1)\lambda z^5}{14400} \right. \\
& - \frac{1}{576}(\lambda-3)(\lambda-2)(\lambda-1)z^4 - \frac{1}{576}(\lambda-3)(\lambda-2)\lambda z^4 - \frac{1}{576}(\lambda-3)(\lambda-1)\lambda z^4 \\
& - \frac{25(\lambda-3)(\lambda-2)(\lambda-1)\lambda z^4}{3456} - \frac{1}{576}(\lambda-2)(\lambda-1)\lambda z^4 + \frac{1}{36}(\lambda-2)(\lambda-1)z^3 \\
& + \frac{1}{36}(\lambda-2)\lambda z^3 + \frac{11}{108}(\lambda-2)(\lambda-1)\lambda z^3 + \frac{1}{36}(\lambda-1)\lambda z^3 - \frac{1}{4}(\lambda-1)z^2 - \frac{3}{4}(\lambda-1)\lambda z^2 \\
& - \frac{\lambda z^2}{4} + \left(-\frac{(\lambda-4)(\lambda-3)(\lambda-2)(\lambda-1)\lambda z^5}{14400} + \frac{1}{576}(\lambda-3)(\lambda-2)(\lambda-1)\lambda z^4 \right. \\
& \left. - \frac{1}{36}(\lambda-2)(\lambda-1)\lambda z^3 + \frac{1}{4}(\lambda-1)\lambda z^2 - \lambda z + 1 \right) \log(z) + 2\lambda z + z
\end{aligned}$$

3.14 problem Problem 16.15

Internal problem ID [2034]

Book: Mathematical methods for physics and engineering, Riley, Hobson, Bence, second edition, 2002

Section: Chapter 16, Series solutions of ODEs. Section 16.6 Exercises, page 550

Problem number: Problem 16.15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [_Gegenbauer, [_2nd_order, _linear, ‘_with_symmetry_[0,F(x)]’]]

$$(-z^2 + 1)y'' - zy' + m^2y = 0$$

With the expansion point for the power series method at $z = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 71

```
Order:=6;
dsolve((1-z^2)*diff(y(z),z$2)-z*diff(y(z),z)+m^2*y(z)=0,y(z),type='series',z=0);
```

$$y(z) = \left(1 - \frac{m^2 z^2}{2} + \frac{m^2(m^2 - 4)z^4}{24}\right) y(0) + \left(z - \frac{(m^2 - 1)z^3}{6} + \frac{(m^4 - 10m^2 + 9)z^5}{120}\right) D(y)(0) + O(z^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 88

```
AsymptoticDSolveValue[(1-z^2)*y'[z]-z*y'[z]+m^2*y[z]==0,y[z],{z,0,5}]
```

$$y(z) \rightarrow c_2 \left(\frac{m^4 z^5}{120} - \frac{m^2 z^5}{12} - \frac{m^2 z^3}{6} + \frac{3z^5}{40} + \frac{z^3}{6} + z \right) + c_1 \left(\frac{m^4 z^4}{24} - \frac{m^2 z^4}{6} - \frac{m^2 z^2}{2} + 1 \right)$$