


A Solution Manual For

**Differential Equations by Shepley
L. Ross. Third edition. John
Willey. New Delhi. 2004.**



Nasser M. Abbasi

October 12, 2023

Contents

1	Chapter 1, Differential equations and their solutions. Exercises page 13	2
2	Chapter 1, section 1.3. Exercises page 22	18
3	Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37	30
4	Chapter 2, section 2.2 (Separable equations). Exercises page 47	49

1 Chapter 1, Differential equations and their solutions. Exercises page 13

1.1	problem 1(a)	3
1.2	problem 1(b)	4
1.3	problem 1(c)	5
1.4	problem 1(d)	6
1.5	problem 2(a)	7
1.6	problem 2(b)	8
1.7	problem 3(a)	9
1.8	problem 3(b)	10
1.9	problem 4(a)	11
1.10	problem 4(b)	12
1.11	problem 5(a)	13
1.12	problem 5(b)	14
1.13	problem 6(a)	15
1.14	problem 6(b)	16
1.15	problem 7(a)	17

1.1 problem 1(a)

Internal problem ID [10552]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, Differential equations and their solutions. Exercises page 13

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + y - x - 1 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)+y(x)=1+x,y(x), singsol=all)
```

$$y(x) = x + e^{-x}c_1$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 15

```
DSolve[y'[x]+y[x]==1+x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x + c_1 e^{-x}$$

1.2 problem 1(b)

Internal problem ID [10553]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, Differential equations and their solutions. Exercises page 13

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 7y' + 12y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-7*diff(y(x),x)+12*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{3x}c_1 + c_2e^{4x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 20

```
DSolve[y''[x]-7*y'[x]+12*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{3x}(c_2e^x + c_1)$$

1.3 problem 1(c)

Internal problem ID [10554]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, Differential equations and their solutions. Exercises page 13

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 3y' + 2y - 4x^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)+2*y(x)=4*x^2,y(x), singsol=all)
```

$$y(x) = e^{2x}c_1 + c_2e^x + 2x^2 + 6x + 7$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 26

```
DSolve[y''[x]-3*y'[x]+2*y[x]==4*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2x(x + 3) + e^x(c_2e^x + c_1) + 7$$

1.4 problem 1(d)

Internal problem ID [10555]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, Differential equations and their solutions. Exercises page 13

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x^2 + 1)y'' + 4y'x + 2y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve((1+x^2)*diff(y(x),x$2)+4*x*diff(y(x),x)+2*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{xc_1 + c_2}{x^2 + 1}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 20

```
DSolve[(1+x^2)*y''[x]+4*x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2x + c_1}{x^2 + 1}$$

1.5 problem 2(a)

Internal problem ID [10556]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, Differential equations and their solutions. Exercises page 13

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _exact, _rational, _Bernoulli]`

$$2yxy' + x^2 + y^2 = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 49

```
dsolve(2*x*y(x)*diff(y(x),x)+x^2+y(x)^2=0,y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{3} \sqrt{x(-x^3 + 3c_1)}}{3x}$$

$$y(x) = \frac{\sqrt{3} \sqrt{x(-x^3 + 3c_1)}}{3x}$$

✓ Solution by Mathematica

Time used: 0.24 (sec). Leaf size: 60

```
DSolve[2*x*y[x]*y'[x]+x^2+y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{-x^3 + 3c_1}}{\sqrt{3}\sqrt{x}}$$

$$y(x) \rightarrow \frac{\sqrt{-x^3 + 3c_1}}{\sqrt{3}\sqrt{x}}$$

1.6 problem 2(b)

Internal problem ID [10557]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, Differential equations and their solutions. Exercises page 13

Problem number: 2(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational, _Bernoulli]`

$$y'x + y - x^3y^3 = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 30

```
dsolve(x*diff(y(x),x)+y(x)=x^3*y(x)^3,y(x), singsol=all)
```

$$y(x) = \frac{1}{\sqrt{-2x + c_1} x}$$

$$y(x) = -\frac{1}{\sqrt{-2x + c_1} x}$$

✓ Solution by Mathematica

Time used: 0.369 (sec). Leaf size: 44

```
DSolve[x*y'[x]+y[x]==x^3*y[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\sqrt{x^2(-2x + c_1)}}$$

$$y(x) \rightarrow \frac{1}{\sqrt{x^2(-2x + c_1)}}$$

$$y(x) \rightarrow 0$$

1.7 problem 3(a)

Internal problem ID [10558]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, Differential equations and their solutions. Exercises page 13

Problem number: 3(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 3y - 3x^2e^{-3x} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)+3*y(x)=3*x^2*exp(-3*x),y(x), singsol=all)
```

$$y(x) = (x^3 + c_1) e^{-3x}$$

✓ Solution by Mathematica

Time used: 0.061 (sec). Leaf size: 17

```
DSolve[y'[x]+3*y[x]==3*x^2*Exp[-3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x}(x^3 + c_1)$$

1.8 problem 3(b)

Internal problem ID [10559]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, Differential equations and their solutions. Exercises page 13

Problem number: 3(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' + 4xy - 8x = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)+4*x*y(x)=8*x,y(x), singsol=all)
```

$$y(x) = 2 + e^{-2x^2} c_1$$

✓ Solution by Mathematica

Time used: 0.045 (sec). Leaf size: 22

```
DSolve[y'[x]+4*x*y[x]==8*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 + c_1 e^{-2x^2}$$

$$y(x) \rightarrow 2$$

1.9 problem 4(a)

Internal problem ID [10560]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, Differential equations and their solutions. Exercises page 13

Problem number: 4(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' - 8y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)-8*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{4x} + c_2 e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 22

```
DSolve[y''[x]-2*y'[x]-8*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(c_2 e^{6x} + c_1)$$

1.10 problem 4(b)

Internal problem ID [10561]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, Differential equations and their solutions. Exercises page 13

Problem number: 4(b).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 2y'' - 4y' + 8y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$3)-2*diff(y(x),x$2)-4*diff(y(x),x)+8*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-2x}c_1 + c_2e^{2x} + c_3e^{2x}x$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 27

```
DSolve[y'''[x]-2*y''[x]-4*y'[x]+8*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(e^{4x}(c_3x + c_2) + c_1)$$

1.11 problem 5(a)

Internal problem ID [10562]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, Differential equations and their solutions. Exercises page 13

Problem number: 5(a).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 3y'' - 4y' + 12y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$3)-3*diff(y(x),x$2)-4*diff(y(x),x)+12*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{3x}c_1 + c_2e^{-2x} + c_3e^{2x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 29

```
DSolve[y'''[x]-3*y''[x]-4*y'[x]+12*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(e^{4x}(c_3e^x + c_2) + c_1)$$

1.12 problem 5(b)

Internal problem ID [10563]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, Differential equations and their solutions. Exercises page 13

Problem number: 5(b).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _fully, _exact, _linear]]`

$$x^3 y''' + 2x^2 y'' - 10y'x - 8y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(x^3*diff(y(x),x$3)+2*x^2*diff(y(x),x$2)-10*x*diff(y(x),x)-8*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1}{x} + c_2 x^4 + \frac{c_3}{x^2}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 22

```
DSolve[x^3*y'''[x]+2*x^2*y''[x]-10*x*y'[x]-8*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_3 x^6 + c_2 x + c_1}{x^2}$$

1.13 problem 6(a)

Internal problem ID [10564]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, Differential equations and their solutions. Exercises page 13

Problem number: 6(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 2y - 6e^x - 4xe^{-2x} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x)+2*y(x)=6*exp(x)+4*x*exp(-2*x),y(x), singsol=all)
```

$$y(x) = (2x^2 + 2e^{3x} + c_1) e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.082 (sec). Leaf size: 25

```
DSolve[y'[x]+2*y[x]==6*Exp[x]+4*x*Exp[-2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2e^x + e^{-2x}(2x^2 + c_1)$$

1.14 problem 6(b)

Internal problem ID [10565]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, Differential equations and their solutions. Exercises page 13

Problem number: 6(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y' + 4y + 8 \sin(2x) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+4*y(x)=-8*sin(2*x),y(x), singsol=all)
```

$$y(x) = c_2 e^{2x} + x e^{2x} c_1 - \cos(2x)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 25

```
DSolve[y''[x]-4*y'[x]+4*y[x]==-8*Sin[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\cos(2x) + e^{2x}(c_2 x + c_1)$$

1.15 problem 7(a)

Internal problem ID [10566]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, Differential equations and their solutions. Exercises page 13

Problem number: 7(a).

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$y'^2 - 4y = 0$$

✓ Solution by Maple

Time used: 0.171 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)^2-4*y(x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1^2 - 2xc_1 + x^2$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 38

```
DSolve[(y'[x])^2-4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}(-2x + c_1)^2$$

$$y(x) \rightarrow \frac{1}{4}(2x + c_1)^2$$

$$y(x) \rightarrow 0$$

2 Chapter 1, section 1.3. Exercises page 22

2.1	problem 1	19
2.2	problem 2(a)	20
2.3	problem 2(b)	21
2.4	problem 3(a)	22
2.5	problem 4(a)	23
2.6	problem 4(b)	24
2.7	problem 4(c)	25
2.8	problem 5	26
2.9	problem 6(a)	27
2.10	problem 6(b)	28
2.11	problem 8	29

2.1 problem 1

Internal problem ID [10567]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, section 1.3. Exercises page 22

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y' - 6y = 0$$

With initial conditions

$$[y(0) = 6, y'(0) = 2]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)+diff(y(x),x)-6*y(x)=0,y(0) = 6, D(y)(0) = 2],y(x), singsol=all)
```

$$y(x) = (4e^{5x} + 2)e^{-3x}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 20

```
DSolve[{y'[x]+y'[x]-6*y[x]==0,{y[0]==6,y'[0]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x}(4e^{5x} + 2)$$

2.2 problem 2(a)

Internal problem ID [10568]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, section 1.3. Exercises page 22

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + y - 2e^{-x}x = 0$$

With initial conditions

$$[y(0) = 2]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve([diff(y(x),x)+y(x)=2*x*exp(-x),y(0) = 2],y(x), singsol=all)
```

$$y(x) = (x^2 + 2) e^{-x}$$

✓ Solution by Mathematica

Time used: 0.056 (sec). Leaf size: 16

```
DSolve[{y'[x]+y[x]==2*x*Exp[-x],{y[0]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(x^2 + 2)$$

2.3 problem 2(b)

Internal problem ID [10569]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, section 1.3. Exercises page 22

Problem number: 2(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + y - 2e^{-x}x = 0$$

With initial conditions

$$[y(-1) = e + 3]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 17

```
dsolve([diff(y(x),x)+y(x)=2*x*exp(-x),y(-1) = exp(1)+3],y(x), singsol=all)
```

$$y(x) = (x^2 + 3e^{-1})e^{-x}$$

✓ Solution by Mathematica

Time used: 0.057 (sec). Leaf size: 20

```
DSolve[{y'[x]+y[x]==2*x*Exp[-x],{y[-1]==Exp[1]+3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x-1}(ex^2 + 3)$$

2.4 problem 3(a)

Internal problem ID [10570]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, section 1.3. Exercises page 22

Problem number: 3(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y' - 12y = 0$$

With initial conditions

$$[y(0) = 5, y'(0) = 6]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)-diff(y(x),x)-12*y(x)=0,y(0) = 5, D(y)(0) = 6],y(x), singsol=all)
```

$$y(x) = (3e^{7x} + 2)e^{-3x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 20

```
DSolve[{y'[x]-y'[x]-12*y[x]==0,{y[0]==5,y'[0]==6}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x}(3e^{7x} + 2)$$

2.5 problem 4(a)

Internal problem ID [10571]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, section 1.3. Exercises page 22

Problem number: 4(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y = 0$$

With initial conditions

$$\left[y(0) = 0, y'\left(\frac{\pi}{2}\right) = 1 \right]$$

✗ Solution by Maple

```
dsolve([diff(y(x),x$2)+y(x)=0,y(0) = 0, D(y)(1/2*Pi) = 1],y(x), singsol=all)
```

No solution found

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y'[x]+y[x]==0,{y[0]==0,y'[Pi/2]==1}},y[x],x,IncludeSingularSolutions -> True]
```

{}

2.6 problem 4(b)

Internal problem ID [10572]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, section 1.3. Exercises page 22

Problem number: 4(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y = 0$$

With initial conditions

$$\left[y(0) = 0, y'\left(\frac{\pi}{2}\right) = -1 \right]$$

✗ Solution by Maple

```
dsolve([diff(y(x),x$2)+y(x)=0,y(0) = 0, D(y)(1/2*Pi) = -1],y(x), singsol=all)
```

No solution found

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y'[x]+y[x]==0,{y[0]==0,y'[Pi/2]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

{}

2.7 problem 4(c)

Internal problem ID [10573]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, section 1.3. Exercises page 22

Problem number: 4(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y = 0$$

With initial conditions

$$[y(0) = 0, y'(\pi) = 1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 8

```
dsolve([diff(y(x),x$2)+y(x)=0,y(0) = 0, D(y)(Pi) = 1],y(x), singsol=all)
```

$$y(x) = -\sin(x)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 9

```
DSolve[{y'[x]+y[x]==0,{y[0]==0,y'[Pi]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sin(x)$$

2.8 problem 5

Internal problem ID [10574]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, section 1.3. Exercises page 22

Problem number: 5.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$x^3 y''' - 3x^2 y'' + 6y'x - 6y = 0$$

With initial conditions

$$[y(2) = 0, y'(2) = 2, y''(2) = 6]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve([x^3*diff(y(x),x$3)-3*x^2*diff(y(x),x$2)+6*x*diff(y(x),x)-6*y(x)=0,y(2) = 0, D(y)(2) =
```

$$y(x) = x^3 - 3x^2 + 2x$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 13

```
DSolve[{x^3*y'''[x]-3*x^2*y''[x]+6*x*y'[x]-6*y[x]==0,{y[2]==0,y'[2]==2,y''[2]==6}},y[x],x,Inc
```

$$y(x) \rightarrow (x - 2)(x - 1)x$$

2.9 problem 6(a)

Internal problem ID [10575]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, section 1.3. Exercises page 22

Problem number: 6(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' - x^2 \sin(y) = 0$$

With initial conditions

$$[y(1) = -2]$$

✓ Solution by Maple

Time used: 2.719 (sec). Leaf size: 97

```
dsolve([diff(y(x),x)=x^2*sin(y(x)),y(1) = -2],y(x), singsol=all)
```

$$y(x) = \arctan \left(\frac{2 \sin(2) e^{\frac{(x-1)(x^2+x+1)}{3}}}{(-1 + \cos(2)) e^{\frac{2(x-1)(x^2+x+1)}{3}} - \cos(2) - 1}, \frac{(-\cos(2) + 1) e^{\frac{2(x-1)(x^2+x+1)}{3}} - \cos(2) - 1}{(-1 + \cos(2)) e^{\frac{2(x-1)(x^2+x+1)}{3}} - \cos(2) - 1} \right)$$

✓ Solution by Mathematica

Time used: 6.739 (sec). Leaf size: 24

```
DSolve[{y'[x]==x^2*Sin[y[x]],{y[1]==-2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2 \cot^{-1} \left(e^{\frac{1}{3} - \frac{x^3}{3}} \cot(1) \right)$$

2.10 problem 6(b)

Internal problem ID [10576]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, section 1.3. Exercises page 22

Problem number: 6(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' - \frac{y^2}{x-2} = 0$$

With initial conditions

$$[y(1) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve([diff(y(x),x)=y(x)^2/(x-2),y(1) = 0],y(x), singsol=all)
```

$$y(x) = 0$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 6

```
DSolve[{y'[x]==y[x]^2/(x-2),{y[1]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

2.11 problem 8

Internal problem ID [10577]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 1, section 1.3. Exercises page 22

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - y^{\frac{1}{3}} = 0$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve([diff(y(x),x)=y(x)^(1/3),y(0) = 0],y(x), singsol=all)
```

$$y(x) = 0$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 21

```
DSolve[{y'[x]==y[x]^(1/3),{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2}{3} \sqrt{\frac{2}{3}} x^{3/2}$$

3 Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

3.1	problem 1	31
3.2	problem 2	32
3.3	problem 3	33
3.4	problem 4	34
3.5	problem 5	35
3.6	problem 7	36
3.7	problem 8	37
3.8	problem 9	38
3.9	problem 10	39
3.10	problem 11	40
3.11	problem 12	41
3.12	problem 13	42
3.13	problem 14	43
3.14	problem 15	44
3.15	problem 16	45
3.16	problem 21	46
3.17	problem 22	47
3.18	problem 24	48

3.1 problem 1

Internal problem ID [10578]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _exact, _rational, [_Abel, '2nd typ`

$$3x + 2y + (y + 2x)y' = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 49

```
dsolve((3*x+2*y(x))+(2*x+y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-2xc_1 - \sqrt{c_1^2x^2 + 1}}{c_1}$$

$$y(x) = \frac{-2xc_1 + \sqrt{c_1^2x^2 + 1}}{c_1}$$

✓ Solution by Mathematica

Time used: 0.538 (sec). Leaf size: 79

```
DSolve[(3*x+2*y[x])+(2*x+y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2x - \sqrt{x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -2x + \sqrt{x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -\sqrt{x^2} - 2x$$

$$y(x) \rightarrow \sqrt{x^2} - 2x$$

3.2 problem 2

Internal problem ID [10579]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_exact, _rational, [_1st_order, ‘_with_symmetry_[F(x)*G(y),0]’

$$y^2 + 3 + (2xy - 4)y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 37

```
dsolve((y(x)^2+3)+(2*x*y(x)-4)*diff(y(x),x)=0,y(x), singsol=all)
```

$$c_1 + \frac{1}{(i\sqrt{3} - y(x))(i(xy(x) - 4)\sqrt{3} - 3x)} = 0$$

✓ Solution by Mathematica

Time used: 0.392 (sec). Leaf size: 77

```
DSolve[(y[x]^2+3)+(2*x*y[x]-4)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2 - \sqrt{4 + x(-3x + c_1)}}{x}$$

$$y(x) \rightarrow \frac{2 + \sqrt{4 + x(-3x + c_1)}}{x}$$

$$y(x) \rightarrow -i\sqrt{3}$$

$$y(x) \rightarrow i\sqrt{3}$$

3.3 problem 3

Internal problem ID [10580]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_exact, _rational, [_1st_order, '_with_symmetry_[F(x),G(x)]]]`,

$$2xy + 1 + (x^2 + 4y)y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
dsolve((2*x*y(x)+1)+(x^2+4*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{x^2}{4} - \frac{\sqrt{x^4 - 8c_1 - 8x}}{4}$$

$$y(x) = -\frac{x^2}{4} + \frac{\sqrt{x^4 - 8c_1 - 8x}}{4}$$

✓ Solution by Mathematica

Time used: 0.133 (sec). Leaf size: 61

```
DSolve[(2*x*y[x]+1)+(x^2+4*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4} \left(-x^2 - \sqrt{x^4 - 8x + 16c_1} \right)$$

$$y(x) \rightarrow \frac{1}{4} \left(-x^2 + \sqrt{x^4 - 8x + 16c_1} \right)$$

3.4 problem 4

Internal problem ID [10581]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, [_Abel, '2nd type', 'class A']]`

$$3x^2y + 2 - (x^3 + y)y' = 0$$

✗ Solution by Maple

```
dsolve((3*x^2*y(x)+2)-(x^3+y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

No solution found

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(3*x^2+2)-(x^3+y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

3.5 problem 5

Internal problem ID [10582]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_exact, _rational, [_Abel, '2nd type', 'class B']]

$$6xy + 2y^2 - 5 + (3x^2 + 4xy - 6)y' = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 67

```
dsolve((6*x*y(x)+2*y(x)^2-5)+(3*x^2+4*x*y(x)-6)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-3x^2 + 6 + \sqrt{9x^4 - 8xc_1 + 4x^2 + 36}}{4x}$$

$$y(x) = -\frac{3x^2 + \sqrt{9x^4 - 8xc_1 + 4x^2 + 36} - 6}{4x}$$

✓ Solution by Mathematica

Time used: 0.471 (sec). Leaf size: 79

```
DSolve[(6*x*y[x]+2*y[x]^2-5)+(3*x^2+4*x*y[x]-6)*y'[x]==0,y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow -\frac{3x^2 + \sqrt{9x^4 + 4x^2 + 16c_1x + 36} - 6}{4x}$$

$$y(x) \rightarrow \frac{-3x^2 + \sqrt{9x^4 + 4x^2 + 16c_1x + 36} + 6}{4x}$$

3.6 problem 7

Internal problem ID [10583]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_exact, [_Abel, '2nd type', 'class A']]

$$y \sec(x)^2 + \sec(x) \tan(x) + (\tan(x) + 2y) y' = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 63

```
dsolve((y(x)*sec(x)^2+sec(x)*tan(x))+(tan(x)+2*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{\sin(x) - \sqrt{-4 \cos(x)^2 c_1 + \sin(x)^2 - 4 \cos(x)}}{2 \cos(x)}$$

$$y(x) = -\frac{\sin(x) + \sqrt{-4 \cos(x)^2 c_1 + \sin(x)^2 - 4 \cos(x)}}{2 \cos(x)}$$

✓ Solution by Mathematica

Time used: 1.148 (sec). Leaf size: 96

```
DSolve[(y[x]*Sec[x]^2+Sec[x]*Tan[x])+(Tan[x]+2*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{1}{4} \left(-2 \tan(x) - \sqrt{\sec^2(x) \sqrt{-16 \cos(x) + (-2 + 8c_1) \cos(2x) + 2 + 8c_1}} \right)$$

$$y(x) \rightarrow \frac{1}{4} \left(-2 \tan(x) + \sqrt{\sec^2(x) \sqrt{-16 \cos(x) + (-2 + 8c_1) \cos(2x) + 2 + 8c_1}} \right)$$

3.7 problem 8

Internal problem ID [10584]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, [_1st_order, ' _with_symmetry_[F(x)*G(y),0]']]`

$$\frac{x}{y^2} + x + \left(\frac{x^2}{y^3} + y \right) y' = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 47

```
dsolve((x/y(x)^2+x)+(x^2/y(x)^3+y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$\frac{y(x)^2 x^2}{2y(x)^2 + 2} + \frac{y(x)^2}{2} - \frac{1}{2(y(x)^2 + 1)} - \ln(y(x)^2 + 1) + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.284 (sec). Leaf size: 55

```
DSolve[(x/y[x]^2+x)+(x^2/y[x]^3+y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{x^2 y(x)^2}{2(y(x)^2 + 1)} + \frac{y(x)^2}{2} - \frac{1}{2(y(x)^2 + 1)} - \log(y(x)^2 + 1) = c_1, y(x) \right]$$

3.8 problem 9

Internal problem ID [10585]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$\frac{(2s-1)s'}{t} + \frac{s-s^2}{t^2} = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 31

```
dsolve((2*s(t)-1)/t*diff(s(t),t)+(s(t)-s(t)^2)/t^2=0,s(t), singsol=all)
```

$$s(t) = \frac{1}{2} - \frac{\sqrt{4c_1t+1}}{2}$$

$$s(t) = \frac{1}{2} + \frac{\sqrt{4c_1t+1}}{2}$$

✓ Solution by Mathematica

Time used: 0.407 (sec). Leaf size: 59

```
DSolve[(2*s[t]-1)/t*s'[t]+(s[t]-s[t]^2)/t^2==0,s[t],t,IncludeSingularSolutions -> True]
```

$$s(t) \rightarrow \frac{1}{2}(1 - \sqrt{1 - 4e^{c_1t}})$$

$$s(t) \rightarrow \frac{1}{2}(1 + \sqrt{1 - 4e^{c_1t}})$$

$$s(t) \rightarrow 0$$

$$s(t) \rightarrow 1$$

3.9 problem 10

Internal problem ID [10586]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational]`

$$\frac{2y^{\frac{3}{2}} + 1}{x^{\frac{1}{3}}} + (3\sqrt{x}\sqrt{y} - 1)y' = 0$$

X Solution by Maple

```
dsolve((2*y(x)^(3/2)+1)/x^(1/3)+(3*x^(1/2)*y(x)^(1/2)-1)*diff(y(x),x)=0,y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(2*y[x]^(3/2)+1)/x^(1/3)+(3*x^(1/2)*y[x]^(1/2)-1)*y'[x]==0,y[x],x,IncludeSingularSolut
```

Timed out

3.10 problem 11

Internal problem ID [10587]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_exact, _rational, [_1st_order, '_with_symmetry_[F(x),G(x)]'],

$$2xy - 3 + (x^2 + 4y) y' = 0$$

With initial conditions

$$[y(1) = 2]$$

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 22

```
dsolve([(2*x*y(x)-3)+(x^2+4*y(x))*diff(y(x),x)=0,y(1) = 2],y(x), singsol=all)
```

$$y(x) = -\frac{x^2}{4} + \frac{\sqrt{x^4 + 24x + 56}}{4}$$

✓ Solution by Mathematica

Time used: 0.137 (sec). Leaf size: 27

```
DSolve[{(2*x*y[x]-3)+(x^2+4*y[x])*y'[x]==0,{y[1]==2}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{1}{4} \left(\sqrt{x^4 + 24x + 56} - x^2 \right)$$

3.11 problem 12

Internal problem ID [10588]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [exact, rational]

$$3y^2x^2 - y^3 + 2x + (2x^3y - 3y^2x + 1)y' = 0$$

With initial conditions

$$[y(-2) = 1]$$

✗ Solution by Maple

```
dsolve([(3*x^2*y(x)^2-y(x)^3+2*x)+(2*x^3*y(x)-3*x*y(x)^2+1)*diff(y(x),x)=0,y(-2) = 1],y(x), s
```

No solution found

✓ Solution by Mathematica

Time used: 56.288 (sec). Leaf size: 250

```
DSolve[{(3*x^2*y[x]^2-y[x]^3+2*x)+(2*x^3*y[x]-3*x*y[x]^2+1)*y'[x]==0,{y[-2]==1}},y[x],x,Inclu
```

$y(x)$

$$\rightarrow \frac{2^{\frac{3}{2}}(1 - i\sqrt{3})x^6 + 4^{\frac{3}{2}}\sqrt{-2x^9 - 36x^4 - 27x^2 + 3\sqrt{3}\sqrt{x^3}(4x^{10} + 4x^8 + 44x^5 + 72x^3 + 27x - 4)}x^3 + (1 + i\sqrt{3})x^6}{12x^3\sqrt{-2x^9 - 36x^4 - 27x^2 + 3\sqrt{3}\sqrt{x^3}}}$$

3.12 problem 13

Internal problem ID [10589]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_exact, [_Abel, '2nd type', 'class B']]

$$2y \sin(x) \cos(x) + y^2 \sin(x) + (\sin(x)^2 - 2y \cos(x)) y' = 0$$

With initial conditions

$$[y(0) = 3]$$

✓ Solution by Maple

Time used: 0.359 (sec). Leaf size: 24

```
dsolve([(2*y(x)*sin(x)*cos(x)+y(x)^2*sin(x))+(sin(x)^2-2*y(x)*cos(x))*diff(y(x),x)]=0,y(0) = 3
```

$$y(x) = \frac{\left(\sin(x)^2 + \sqrt{\sin(x)^4 + 36 \cos(x)}\right) \sec(x)}{2}$$

✓ Solution by Mathematica

Time used: 1.267 (sec). Leaf size: 32

```
DSolve[{(2*y[x]*Sin[x]*Cos[x]+y[x]^2*SIn[x])+(Sin[x]^2-2*y[x]*Cos[x])*y'[x]==0,{y[0]==3}},y[x]
```

$$y(x) \rightarrow \frac{1}{2} \sec(x) \left(-\cos^2(x) + \sqrt{\sin^4(x) + 36 \cos(x)} + 1 \right)$$

3.13 problem 14

Internal problem ID [10590]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 14.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_exact, [_Abel, '2nd type', 'class B']]`

$$e^x y + 2e^x + y^2 + (e^x + 2xy)y' = 0$$

With initial conditions

$$[y(0) = 6]$$

✓ Solution by Maple

Time used: 0.437 (sec). Leaf size: 29

```
dsolve([(y(x)*exp(x)+2*exp(x)+y(x)^2)+(exp(x)+2*x*y(x))*diff(y(x),x)=0,y(0) = 6],y(x), singso
```

$$y(x) = \frac{-e^x + \sqrt{e^{2x} - 8xe^x + 32x}}{2x}$$

✓ Solution by Mathematica

Time used: 31.602 (sec). Leaf size: 37

```
DSolve[{(y[x]*Exp[x]+2*Exp[x]+y[x]^2)+(Exp[x]+2*x*y[x])*y'[x]==0,{y[0]==6}},y[x],x,IncludeSin
```

$$y(x) \rightarrow \frac{\sqrt{-8e^x x + 32x + e^{2x}} - e^x}{2x}$$

3.14 problem 15

Internal problem ID [10591]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_exact, _rational, [_1st_order, ‘_with_symmetry_[F(x)*G(y),0]’

$$\frac{3-y}{x^2} + \frac{(y^2-2x)y'}{y^2x} = 0$$

With initial conditions

$$[y(-1) = 2]$$

✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 21

```
dsolve([(3-y(x))/x^2+((y(x)^2-2*x)/(x*y(x)^2))*diff(y(x),x)=0,y(-1) = 2],y(x), singsol=all)
```

$$y(x) = x + \frac{3}{2} + \frac{\sqrt{4x^2 + 4x + 9}}{2}$$

✓ Solution by Mathematica

Time used: 1.216 (sec). Leaf size: 26

```
DSolve[{(3-y[x])/x^2+( (y[x]^2-2*x)/(x*y[x]^2) )*y'[x]==0,{y[-1]==2}},y[x],x,IncludeSingularS
```

$$y(x) \rightarrow \frac{1}{2} \left(2x + \sqrt{4x(x+1) + 9} + 3 \right)$$

3.15 problem 16

Internal problem ID [10592]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _exact, _rational]`

$$\frac{1 + 8xy^{\frac{2}{3}}}{x^{\frac{2}{3}}y^{\frac{1}{3}}} + \frac{(2x^{\frac{4}{3}}y^{\frac{2}{3}} - x^{\frac{1}{3}})y'}{y^{\frac{4}{3}}} = 0$$

With initial conditions

$$[y(1) = 8]$$

✓ Solution by Maple

Time used: 0.25 (sec). Leaf size: 55

```
dsolve([(1+8*x*y(x)^(2/3))/(x^(2/3)*y(x)^(1/3))+((2*x^(4/3)*y(x)^(2/3)-x^(1/3))/(y(x)^(4/3)))]
```

$$y(x) = \text{RootOf}\left(64_Z^7x^4 + 96_Z^5x^3 - 729_Z^4 + 48x^2_Z + 8x_Z^{\frac{1}{3}}\right)$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{(1+8*x*y[x]^(2/3))/(x^(2/3)*y[x]^(1/3))+((2*x^(4/3)*y[x]^(2/3)-x^(1/3))/(y[x]^(4/3)))]
```

{}

3.16 problem 21

Internal problem ID [10593]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 21.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational, _Bernoulli]`

$$4x + 3y^2 + 2yxy' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 38

```
dsolve((4*x+3*y(x)^2)+(2*x*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{x(-x^4 + c_1)}}{x^2}$$

$$y(x) = -\frac{\sqrt{x(-x^4 + c_1)}}{x^2}$$

✓ Solution by Mathematica

Time used: 0.228 (sec). Leaf size: 46

```
DSolve[(4*x+3*y[x]^2)+(2*x*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{-x^4 + c_1}}{x^{3/2}}$$

$$y(x) \rightarrow \frac{\sqrt{-x^4 + c_1}}{x^{3/2}}$$

3.17 problem 22

Internal problem ID [10594]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 22.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$y^2 + 2xy - x^2y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve((y(x)^2+2*x*y(x))-x^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x^2}{-x + c_1}$$

✓ Solution by Mathematica

Time used: 0.144 (sec). Leaf size: 22

```
DSolve[(y[x]^2+2*x*y[x])-x^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2}{-x + c_1}$$

$$y(x) \rightarrow 0$$

3.18 problem 24

Internal problem ID [10595]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.1 (Exact differential equations and integrating factors). Exercises page 37

Problem number: 24.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _rational]`

$$y + x(x^2 + y^2)^2 + (y(x^2 + y^2)^2 - x)y' = 0$$

✓ Solution by Maple

Time used: 0.218 (sec). Leaf size: 39

```
dsolve((y(x)+x*(x^2+y(x)^2)^2)+(y(x)*(x^2+y(x)^2)^2-x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\tan(\text{RootOf}(-\tan(_Z)^4 x^4 - 2\tan(_Z)^2 x^4 - x^4 + 4c_1 - 4_Z)) x$$

✓ Solution by Mathematica

Time used: 0.097 (sec). Leaf size: 40

```
DSolve[(y[x]+x*(x^2+y[x]^2)^2)+(y[x]*(x^2+y[x]^2)^2-x)*y'[x]==0,y[x],x,IncludeSingularSolutions->True]
```

$$\text{Solve}\left[\arctan\left(\frac{x}{y(x)}\right) + \frac{x^4}{4} + \frac{1}{2}x^2y(x)^2 + \frac{y(x)^4}{4} = c_1, y(x)\right]$$

4 Chapter 2, section 2.2 (Separable equations).

Exercises page 47

4.1	problem 1	50
4.2	problem 2	51
4.3	problem 3	52
4.4	problem 4	53
4.5	problem 5	54
4.6	problem 6	55
4.7	problem 7	56
4.8	problem 8	58
4.9	problem 9	59
4.10	problem 10	60
4.11	problem 11	61
4.12	problem 12	62
4.13	problem 13	64
4.14	problem 14	65
4.15	problem 15	66
4.16	problem 16	67
4.17	problem 17	68
4.18	problem 18	69
4.19	problem 19	70
4.20	problem 20	71
4.21	problem 22(a)	72
4.22	problem 22(b)	73
4.23	problem 23(a)	74
4.24	problem 23(b)	76

4.1 problem 1

Internal problem ID [10596]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$4xy + (x^2 + 1)y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve((4*x*y(x))+(x^2+1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1}{(x^2 + 1)^2}$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 20

```
DSolve[(4*x*y[x])+(x^2+1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{(x^2 + 1)^2}$$

$$y(x) \rightarrow 0$$

4.2 problem 2

Internal problem ID [10597]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$xy + 2x + y + 2 + (x^2 + 2x)y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve((x*y(x)+2*x+y(x)+2)+(x^2+2*x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -2 + \frac{c_1}{\sqrt{x(x+2)}}$$

✓ Solution by Mathematica

Time used: 0.095 (sec). Leaf size: 27

```
DSolve[(x*y[x]+2*x+y[x]+2)+(x^2+2*x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2 + \frac{c_1}{\sqrt{x}\sqrt{x+2}}$$

$$y(x) \rightarrow -2$$

4.3 problem 3

Internal problem ID [10598]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$2r(s^2 + 1) + (r^4 + 1)s' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(2*r*(s(r)^2+1)+(r^4+1)*diff(s(r),r)=0,s(r), singsol=all)
```

$$s(r) = -\tan(\arctan(r^2) + 2c_1)$$

✓ Solution by Mathematica

Time used: 0.316 (sec). Leaf size: 31

```
DSolve[2*r*(s[r]^2+1)+(r^4+1)*s'[r]==0,s[r],r,IncludeSingularSolutions -> True]
```

$$s(r) \rightarrow -\tan(\arctan(r^2) - c_1)$$

$$s(r) \rightarrow -i$$

$$s(r) \rightarrow i$$

4.4 problem 4

Internal problem ID [10599]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$\csc(y) + y' \sec(x) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(csc(y(x))+sec(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \arccos(\sin(x) + c_1)$$

✓ Solution by Mathematica

Time used: 0.443 (sec). Leaf size: 27

```
DSolve[Csc[y[x]]+Sec[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\arccos(\sin(x) - c_1)$$

$$y(x) \rightarrow \arccos(\sin(x) - c_1)$$

4.5 problem 5

Internal problem ID [10600]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$\tan(\theta) + 2r\theta' = 0$$

✓ Solution by Maple

Time used: 0.469 (sec). Leaf size: 21

```
dsolve(tan(theta(r))+2*r*diff(theta(r),r)=0,theta(r), singsol=all)
```

$$\theta(r) = \arcsin\left(\frac{1}{\sqrt{c_1 r}}\right)$$

$$\theta(r) = -\arcsin\left(\frac{1}{\sqrt{c_1 r}}\right)$$

✓ Solution by Mathematica

Time used: 8.404 (sec). Leaf size: 21

```
DSolve[Tan[theta[r]]+2*r*theta'[r]==0,theta[r],r,IncludeSingularSolutions -> True]
```

$$\theta(r) \rightarrow \arcsin\left(\frac{e^{c_1}}{\sqrt{r}}\right)$$

$$\theta(r) \rightarrow 0$$

4.6 problem 6

Internal problem ID [10601]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$(e^v + 1) \cos(u) + e^v(1 + \sin(u)) v' = 0$$

✓ Solution by Maple

Time used: 0.922 (sec). Leaf size: 28

```
dsolve((exp(v(u))+1)*cos(u) + exp(v(u))*(1+sin(u))*diff(v(u),u)=0,v(u), singsol=all)
```

$$v(u) = -\ln\left(-\frac{1 + \sin(u)}{-1 + \sin(u) e^{c_1} + e^{c_1}}\right) - c_1$$

✓ Solution by Mathematica

Time used: 4.67 (sec). Leaf size: 28

```
DSolve[(Exp[v[u]]+1)*Cos[u] + Exp[v[u]]*(1+Sin[u])*v'[u]==0,v[u],u,IncludeSingularSolutions -
```

$$v(u) \rightarrow \log\left(-1 + \frac{e^{c_1}}{\sin(u) + 1}\right)$$

$$v(u) \rightarrow i\pi$$

4.7 problem 7

Internal problem ID [10602]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$(x + 4)(y^2 + 1) + y(x^2 + 3x + 2)y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 122

```
dsolve((x+4)*(y(x)^2+1) + y(x)*(x^2+3*x+2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{-x^6 + c_1x^4 - 6x^5 + 8c_1x^3 + 24x^2c_1 + 100x^3 + 32xc_1 + 345x^2 + 16c_1 + 474x + 239}}{(x + 1)^3}$$

$$y(x) = -\frac{\sqrt{-x^6 + c_1x^4 - 6x^5 + 8c_1x^3 + 24x^2c_1 + 100x^3 + 32xc_1 + 345x^2 + 16c_1 + 474x + 239}}{(x + 1)^3}$$

✓ Solution by Mathematica

Time used: 4.966 (sec). Leaf size: 126

```
DSolve[(x+4)*(y[x]^2+1) + y[x]*(x^2+3*x+2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{-(x+1)^6 + e^{2c_1}(x+2)^4}}{(x+1)^3}$$

$$y(x) \rightarrow \frac{\sqrt{-(x+1)^6 + e^{2c_1}(x+2)^4}}{(x+1)^3}$$

$$y(x) \rightarrow -i$$

$$y(x) \rightarrow i$$

$$y(x) \rightarrow \frac{(x+1)^3}{\sqrt{-(x+1)^6}}$$

$$y(x) \rightarrow \frac{\sqrt{-(x+1)^6}}{(x+1)^3}$$

4.8 problem 8

Internal problem ID [10603]

Book: Differential Equations by Shepley L. Ross. Third edition. John Willey. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear]`

$$x + y - y'x = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve((x+y(x))- x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = (\ln(x) + c_1)x$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 12

```
DSolve[(x+y[x])- x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(\log(x) + c_1)$$

4.9 problem 9

Internal problem ID [10604]

Book: Differential Equations by Shepley L. Ross. Third edition. John Willey. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cla`

$$2xy + 3y^2 - (2xy + x^2)y' = 0$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 35

```
dsolve((2*x*y(x)+3*y(x)^2)- (2*x*y(x)+x^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \left(-\frac{1}{2} - \frac{\sqrt{4xc_1 + 1}}{2}\right)x$$

$$y(x) = \left(-\frac{1}{2} + \frac{\sqrt{4xc_1 + 1}}{2}\right)x$$

✓ Solution by Mathematica

Time used: 0.406 (sec). Leaf size: 61

```
DSolve[(2*x*y[x]+3*y[x]^2)- (2*x*y[x]+x^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2}x \left(1 + \sqrt{1 + 4e^{c_1}x}\right)$$

$$y(x) \rightarrow \frac{1}{2}x \left(-1 + \sqrt{1 + 4e^{c_1}x}\right)$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -x$$

4.10 problem 10

Internal problem ID [10605]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _dAlembert]`

$$v^3 + (u^3 - uv^2) v' = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 21

```
dsolve(v(u)^3+ (u^3-u*v(u)^2)*diff(v(u),u)=0,v(u), singsol=all)
```

$$v(u) = e^{-\frac{\text{LambertW}\left(-\frac{e^{-2c_1}}{u^2}\right)}{2} - c_1}$$

✓ Solution by Mathematica

Time used: 5.391 (sec). Leaf size: 56

```
DSolve[v[u]^3+ (u^3-u*v[u]^2)*v'[u]==0,v[u],u,IncludeSingularSolutions -> True]
```

$$v(u) \rightarrow -iu \sqrt{W\left(-\frac{e^{-2c_1}}{u^2}\right)}$$

$$v(u) \rightarrow iu \sqrt{W\left(-\frac{e^{-2c_1}}{u^2}\right)}$$

$$v(u) \rightarrow 0$$

4.11 problem 11

Internal problem ID [10606]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _dAlembert]`

$$x \tan\left(\frac{y}{x}\right) + y - y'x = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 10

```
dsolve((x*tan(y(x)/x)+y(x))- x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \arcsin(xc_1)x$$

✓ Solution by Mathematica

Time used: 5.033 (sec). Leaf size: 19

```
DSolve[(x*Tan[y[x]/x]+y[x])- x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \arcsin(e^{c_1}x)$$

$$y(x) \rightarrow 0$$

4.12 problem 12

Internal problem ID [10607]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _exact, _rational, _dAlembert]`

$$(2s^2 + 2st + t^2) s' + s^2 + 2st - t^2 = 0$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 421

`dsolve((2*s(t)^2+2*s(t)*t+t^2)*diff(s(t),t)+(s(t)^2+2*s(t)*t-t^2)=0,s(t), singsol=all)`

$$s(t) = \frac{\left(4t^3c_1^3+2+\sqrt{17c_1^6t^6+16t^3c_1^3+4}\right)^{\frac{1}{3}}}{2} - \frac{t^2c_1^2}{2\left(4t^3c_1^3+2+\sqrt{17c_1^6t^6+16t^3c_1^3+4}\right)^{\frac{1}{3}}} - \frac{c_1t}{2}$$

$s(t)$

$$= \frac{-\left(4t^3c_1^3+2+\sqrt{17c_1^6t^6+16t^3c_1^3+4}\right)^{\frac{1}{3}}}{4} + \frac{t^2c_1^2}{4\left(4t^3c_1^3+2+\sqrt{17c_1^6t^6+16t^3c_1^3+4}\right)^{\frac{1}{3}}} - \frac{c_1t}{2} - \frac{i\sqrt{3}\left(\frac{\left(4t^3c_1^3+2+\sqrt{17c_1^6t^6+16t^3c_1^3+4}\right)^{\frac{1}{3}}}{2} + \frac{1}{2\left(4t^3c_1^3+2+\sqrt{17c_1^6t^6+16t^3c_1^3+4}\right)^{\frac{1}{3}}}\right)}{2}$$

$s(t)$

$$= \frac{-\left(4t^3c_1^3+2+\sqrt{17c_1^6t^6+16t^3c_1^3+4}\right)^{\frac{1}{3}}}{4} + \frac{t^2c_1^2}{4\left(4t^3c_1^3+2+\sqrt{17c_1^6t^6+16t^3c_1^3+4}\right)^{\frac{1}{3}}} - \frac{c_1t}{2} + \frac{i\sqrt{3}\left(\frac{\left(4t^3c_1^3+2+\sqrt{17c_1^6t^6+16t^3c_1^3+4}\right)^{\frac{1}{3}}}{2} + \frac{1}{2\left(4t^3c_1^3+2+\sqrt{17c_1^6t^6+16t^3c_1^3+4}\right)^{\frac{1}{3}}}\right)}{2}$$

✓ Solution by Mathematica

Time used: 43.63 (sec). Leaf size: 613

`DSolve[(2*s[t]^2+2*s[t]*t+t^2)*s'[t]+(s[t]^2+2*s[t]*t-t^2)==0,s[t],t,IncludeSingularSolutions`

$$s(t) \rightarrow \frac{1}{2} \left(\sqrt[3]{4t^3 + \sqrt{17t^6 + 16e^{3c_1}t^3 + 4e^{6c_1}} + 2e^{3c_1}} - \frac{t^2}{\sqrt[3]{4t^3 + \sqrt{17t^6 + 16e^{3c_1}t^3 + 4e^{6c_1}} + 2e^{3c_1}}} - t \right)$$

$$s(t) \rightarrow \frac{1}{8} \left(2i(\sqrt{3} + i) \sqrt[3]{4t^3 + \sqrt{17t^6 + 16e^{3c_1}t^3 + 4e^{6c_1}} + 2e^{3c_1}} + \frac{(2 + 2i\sqrt{3})t^2}{\sqrt[3]{4t^3 + \sqrt{17t^6 + 16e^{3c_1}t^3 + 4e^{6c_1}} + 2e^{3c_1}}} - 4t \right)$$

$$s(t) \rightarrow \frac{1}{8} \left((-2 - 2i\sqrt{3}) \sqrt[3]{4t^3 + \sqrt{17t^6 + 16e^{3c_1}t^3 + 4e^{6c_1}} + 2e^{3c_1}} + \frac{(2 - 2i\sqrt{3})t^2}{\sqrt[3]{4t^3 + \sqrt{17t^6 + 16e^{3c_1}t^3 + 4e^{6c_1}} + 2e^{3c_1}}} - 4t \right)$$

$$s(t) \rightarrow \frac{1}{2} \left(\sqrt[3]{\sqrt{17}\sqrt{t^6} + 4t^3} - \frac{t^2}{\sqrt[3]{\sqrt{17}\sqrt{t^6} + 4t^3}} - t \right)$$

$$s(t) \rightarrow \frac{1}{4} \left((-1 - i\sqrt{3}) \sqrt[3]{\sqrt{17}\sqrt{t^6} + 4t^3} + \frac{(1 - i\sqrt{3})t^2}{\sqrt[3]{\sqrt{17}\sqrt{t^6} + 4t^3}} - 2t \right)$$

$$s(t) \rightarrow \frac{1}{4} \left(i(\sqrt{3} + i) \sqrt[3]{\sqrt{17}\sqrt{t^6} + 4t^3} + \frac{(1 + i\sqrt{3})t^2}{\sqrt[3]{\sqrt{17}\sqrt{t^6} + 4t^3}} - 2t \right)$$

4.13 problem 13

Internal problem ID [10608]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A'], _dAlembert]`

$$x^3 + y^2 \sqrt{y^2 + x^2} - xy \sqrt{y^2 + x^2} y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve((x^3+y(x)^2*sqrt(x^2+y(x)^2))-x*y(x)*sqrt(x^2+y(x)^2)*diff(y(x),x)=0,y(x), singsol=all
```

$$-\frac{(y(x)^2 + x^2)^{\frac{3}{2}}}{x^3} + 3 \ln(x) - c_1 = 0$$

✓ Solution by Mathematica

Time used: 19.618 (sec). Leaf size: 265

```
DSolve[(x^3+y[x]^2*Sqrt[x^2+y[x]^2])-x*y[x]*Sqrt[x^2+y[x]^2]*y'[x]==0,y[x],x,IncludeSingularS
```

$$y(x) \rightarrow -\sqrt{-x^2 - \frac{1}{2} \sqrt[6]{3} (\sqrt{3} + 3i) \sqrt[3]{x^6 (\log(x) + c_1)^2}}$$

$$y(x) \rightarrow \sqrt{-x^2 - \frac{1}{2} \sqrt[6]{3} (\sqrt{3} + 3i) \sqrt[3]{x^6 (\log(x) + c_1)^2}}$$

$$y(x) \rightarrow -\sqrt{-x^2 - \frac{1}{2} \sqrt[6]{3} (\sqrt{3} - 3i) \sqrt[3]{x^6 (\log(x) + c_1)^2}}$$

$$y(x) \rightarrow \sqrt{-x^2 - \frac{1}{2} \sqrt[6]{3} (\sqrt{3} - 3i) \sqrt[3]{x^6 (\log(x) + c_1)^2}}$$

$$y(x) \rightarrow -\sqrt{-x^2 + 3^{2/3} \sqrt[3]{x^6 (\log(x) + c_1)^2}}$$

$$y(x) \rightarrow \sqrt{-x^2 + 3^{2/3} \sqrt[3]{x^6 (\log(x) + c_1)^2}}$$

4.14 problem 14

Internal problem ID [10609]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 14.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$\sqrt{x+y} + \sqrt{x-y} + (\sqrt{x-y} - \sqrt{x+y}) y' = 0$$

✓ Solution by Maple

Time used: 3.781 (sec). Leaf size: 32

```
dsolve((sqrt(x+y(x))+sqrt(x-y(x)))+(sqrt(x-y(x))-sqrt(x+y(x)))*diff(y(x),x)=0,y(x), singsol=a
```

$$\ln(x) + \ln\left(\frac{y(x)}{x}\right) - \operatorname{arctanh}\left(\frac{1}{\sqrt{-\frac{y(x)^2}{x^2} + 1}}\right) - c_1 = 0$$

✓ Solution by Mathematica

Time used: 1.717 (sec). Leaf size: 66

```
DSolve[(Sqrt[x+y[x]]+Sqrt[x-y[x]])+(Sqrt[x-y[x]]-Sqrt[x+y[x]])*y'[x]==0,y[x],x,IncludeSingular
```

$$y(x) \rightarrow -\frac{1}{4}e^{\frac{c_1}{2}}\sqrt{e^{c_1} - 8ix}$$

$$y(x) \rightarrow \frac{1}{4}e^{\frac{c_1}{2}}\sqrt{e^{c_1} - 8ix}$$

$$y(x) \rightarrow 0$$

4.15 problem 15

Internal problem ID [10610]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y + 2 + y(x + 4)y' = 0$$

With initial conditions

$$[y(-3) = -1]$$

✓ Solution by Maple

Time used: 0.219 (sec). Leaf size: 18

```
dsolve([(y(x)+2)+(y(x)*(x+4))*diff(y(x),x)=0,y(-3) = -1],y(x), singsol=all)
```

$$y(x) = -2 \operatorname{LambertW}\left(-\frac{\sqrt{x+4}e^{-\frac{1}{2}}}{2}\right) - 2$$

✓ Solution by Mathematica

Time used: 9.319 (sec). Leaf size: 26

```
DSolve[{(y[x]+2)+(y[x]*(x+4))*y'[x]==0,{y[-3]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2\left(W\left(-\frac{\sqrt{x+4}}{2\sqrt{e}}\right) + 1\right)$$

4.16 problem 16

Internal problem ID [10611]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$8 \cos(y)^2 + \csc(x)^2 y' = 0$$

With initial conditions

$$\left[y\left(\frac{\pi}{12}\right) = \frac{\pi}{4} \right]$$

✓ Solution by Maple

Time used: 0.234 (sec). Leaf size: 20

```
dsolve([(8*cos(y(x))^2)+csc(x)^2*diff(y(x),x)=0,y(1/12*Pi) = 1/4*Pi],y(x), singsol=all)
```

$$y(x) = -\arctan\left(-\frac{\pi}{3} + 4x - 2\sin(2x)\right)$$

✓ Solution by Mathematica

Time used: 0.752 (sec). Leaf size: 21

```
DSolve[{(8*Cos[y[x]]^2)+Csc[x]^2*y'[x]==0,{y[Pi/12]==Pi/4}},y[x],x,IncludeSingularSolutions -
```

$$y(x) \rightarrow \arctan\left(-4x + 2\sin(2x) + \frac{\pi}{3}\right)$$

4.17 problem 17

Internal problem ID [10612]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 17.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$(3x + 8)(y^2 + 4) - 4y(x^2 + 5x + 6)y' = 0$$

With initial conditions

$$[y(1) = 2]$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 21

```
dsolve([(3*x+8)*(y(x)^2+4)-4*y(x)*(x^2+5*x+6)*diff(y(x),x)=0,y(1) = 2],y(x), singsol=all)
```

$$y(x) = \frac{2\sqrt{-9 + (3x + 6)\sqrt{3 + x}}}{3}$$

✓ Solution by Mathematica

Time used: 4.372 (sec). Leaf size: 36

```
DSolve[{(3*x+8)*(y[x]^2+4)-4*y[x]*(x^2+5*x+6)*y'[x]==0,{y[1]==2}},y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow \frac{2\sqrt{\sqrt{x+3}x + 2\sqrt{x+3} - 3}}{\sqrt{3}}$$

4.18 problem 18

Internal problem ID [10613]

Book: Differential Equations by Shepley L. Ross. Third edition. John Willey. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 18.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$x^2 + 3y^2 - 2yxy' = 0$$

With initial conditions

$$[y(2) = 6]$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 13

```
dsolve([(x^2+3*y(x)^2)-2*x*y(x)*diff(y(x),x)=0,y(2) = 6],y(x), singsol=all)
```

$$y(x) = \sqrt{5x - 1} x$$

✓ Solution by Mathematica

Time used: 0.275 (sec). Leaf size: 16

```
DSolve[{(x^2+3*y[x]^2)-2*x*y[x]*y'[x]==0,{y[2]==6}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x\sqrt{5x - 1}$$

4.19 problem 19

Internal problem ID [10614]

Book: Differential Equations by Shepley L. Ross. Third edition. John Willey. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cla`

$$2x - 5y + (4x - y)y' = 0$$

With initial conditions

$$[y(1) = 4]$$

✓ Solution by Maple

Time used: 0.297 (sec). Leaf size: 35

```
dsolve([(2*x-5*y(x))+(4*x-y(x))*diff(y(x),x)=0,y(1) = 4],y(x), singsol=all)
```

$$y(x) = 6 - 2x - 6\sqrt{1-x}$$

$$y(x) = 6 - 2x + 6\sqrt{1-x}$$

✓ Solution by Mathematica

Time used: 1.302 (sec). Leaf size: 41

```
DSolve[{(2*x-5*y[x])+(4*x-y[x])*y'[x]==0,{y[1]==4}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2x - 6i\sqrt{x-1} + 6$$

$$y(x) \rightarrow -2x + 6i\sqrt{x-1} + 6$$

4.20 problem 20

Internal problem ID [10615]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 20.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cla`

$$3x^2 + 9xy + 5y^2 - (6x^2 + 4xy) y' = 0$$

With initial conditions

$$[y(2) = -6]$$

✓ Solution by Maple

Time used: 0.329 (sec). Leaf size: 21

```
dsolve([(3*x^2+9*x*y(x)+5*y(x)^2)-(6*x^2+4*x*y(x))*diff(y(x),x)=0,y(2) = -6],y(x), singsol=al
```

$$y(x) = -\frac{\left(3 + \sqrt{-3 + 6\sqrt{x}\sqrt{2}}\right)x}{2}$$

✓ Solution by Mathematica

Time used: 21.827 (sec). Leaf size: 30

```
DSolve[{(3*x^2+9*x*y[x]+5*y[x]^2)-(6*x^2+4*x*y[x])*y'[x]==0,{y[2]==-6}},y[x],x,IncludeSingula
```

$$y(x) \rightarrow -\frac{1}{2}\left(\sqrt{6\sqrt{2}\sqrt{x}-3}+3\right)x$$

4.21 problem 22(a)

Internal problem ID [10616]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 22(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _exact, _rational, [_Abel, '2nd typ`

$$x + 2y + (2x - y)y' = 0$$

✓ Solution by Maple

Time used: 0.093 (sec). Leaf size: 51

```
dsolve((x+2*y(x))+(2*x-y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{2xc_1 - \sqrt{5c_1^2x^2 + 1}}{c_1}$$

$$y(x) = \frac{2xc_1 + \sqrt{5c_1^2x^2 + 1}}{c_1}$$

✓ Solution by Mathematica

Time used: 0.458 (sec). Leaf size: 94

```
DSolve[(x+2*y[x])+(2*x-y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2x - \sqrt{5x^2 + e^{2c_1}}$$

$$y(x) \rightarrow 2x + \sqrt{5x^2 + e^{2c_1}}$$

$$y(x) \rightarrow 2x - \sqrt{5}\sqrt{x^2}$$

$$y(x) \rightarrow \sqrt{5}\sqrt{x^2} + 2x$$

4.22 problem 22(b)

Internal problem ID [10617]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 22(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _exact, _rational, [_Abel, '2nd typ`

$$3x - y - (x + y)y' = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 51

```
dsolve((3*x-y(x))-(x+y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-xc_1 - \sqrt{4c_1^2x^2 + 1}}{c_1}$$

$$y(x) = \frac{-xc_1 + \sqrt{4c_1^2x^2 + 1}}{c_1}$$

✓ Solution by Mathematica

Time used: 0.445 (sec). Leaf size: 85

```
DSolve[(3*x-y[x])-(x+y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x - \sqrt{4x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -x + \sqrt{4x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -2\sqrt{x^2} - x$$

$$y(x) \rightarrow 2\sqrt{x^2} - x$$

4.23 problem 23(a)

Internal problem ID [10618]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 23(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _exact, _rational, _dAlembert]`

$$x^2 + 2y^2 + (4xy - y^2) y' = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 441

```
dsolve((x^2+2*y(x)^2)+(4*x*y(x)-y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\frac{\left(4+68x^3c_1^3+4\sqrt{33c_1^6x^6+34x^3c_1^3+1}\right)^{\frac{1}{3}}}{2} + \frac{8c_1^2x^2}{\left(4+68x^3c_1^3+4\sqrt{33c_1^6x^6+34x^3c_1^3+1}\right)^{\frac{1}{3}}} + 2xc_1}{c_1}$$

$y(x)$

$$= \frac{\frac{\left(4+68x^3c_1^3+4\sqrt{33c_1^6x^6+34x^3c_1^3+1}\right)^{\frac{1}{3}}}{4} - \frac{4c_1^2x^2}{\left(4+68x^3c_1^3+4\sqrt{33c_1^6x^6+34x^3c_1^3+1}\right)^{\frac{1}{3}}} + 2xc_1 - \frac{i\sqrt{3}\left(\frac{\left(4+68x^3c_1^3+4\sqrt{33c_1^6x^6+34x^3c_1^3+1}\right)^{\frac{1}{3}}}{2}\right)}{2}}{c_1}$$

$y(x)$

$$= \frac{\frac{\left(4+68x^3c_1^3+4\sqrt{33c_1^6x^6+34x^3c_1^3+1}\right)^{\frac{1}{3}}}{4} - \frac{4c_1^2x^2}{\left(4+68x^3c_1^3+4\sqrt{33c_1^6x^6+34x^3c_1^3+1}\right)^{\frac{1}{3}}} + 2xc_1 + \frac{i\sqrt{3}\left(\frac{\left(4+68x^3c_1^3+4\sqrt{33c_1^6x^6+34x^3c_1^3+1}\right)^{\frac{1}{3}}}{2}\right)}{2}}{c_1}$$

✓ Solution by Mathematica

Time used: 31.117 (sec). Leaf size: 564

`DSolve[(x^2+2*y[x]^2)+(4*x*y[x]-y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{\sqrt[3]{17x^3 + \sqrt{33x^6 + 34e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}}{\sqrt[3]{2}} + \frac{4\sqrt[3]{2}x^2}{\sqrt[3]{17x^3 + \sqrt{33x^6 + 34e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}} + 2x$$

$$y(x) \rightarrow \frac{1}{2} \left((-2)^{2/3} \sqrt[3]{17x^3 + \sqrt{33x^6 + 34e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}} - \frac{8\sqrt[3]{-2}x^2}{\sqrt[3]{17x^3 + \sqrt{33x^6 + 34e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}} + 4x \right)$$

$$y(x) \rightarrow x \left(2 + \frac{4(-1)^{2/3}\sqrt[3]{2}x}{\sqrt[3]{17x^3 + \sqrt{33x^6 + 34e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}} \right) - \sqrt[3]{-\frac{1}{2}} \sqrt[3]{17x^3 + \sqrt{33x^6 + 34e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{\sqrt{33}\sqrt{x^6} + 17x^3}}{\sqrt[3]{2}} + \frac{4\sqrt[3]{2}x^2}{\sqrt[3]{\sqrt{33}\sqrt{x^6} + 17x^3}} + 2x$$

$$y(x) \rightarrow -\sqrt[3]{-\frac{1}{2}} \sqrt[3]{\sqrt{33}\sqrt{x^6} + 17x^3} + \frac{4(-1)^{2/3}\sqrt[3]{2}x^2}{\sqrt[3]{\sqrt{33}\sqrt{x^6} + 17x^3}} + 2x$$

$$y(x) \rightarrow \frac{1}{2} \left((-2)^{2/3} \sqrt[3]{\sqrt{33}\sqrt{x^6} + 17x^3} - \frac{8\sqrt[3]{-2}x^2}{\sqrt[3]{\sqrt{33}\sqrt{x^6} + 17x^3}} + 4x \right)$$

4.24 problem 23(b)

Internal problem ID [10619]

Book: Differential Equations by Shepley L. Ross. Third edition. John Wiley. New Delhi. 2004.

Section: Chapter 2, section 2.2 (Separable equations). Exercises page 47

Problem number: 23(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _exact, _rational, [_Abel, '2nd typ`

$$2x^2 + 2xy + y^2 + (2xy + x^2) y' = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 73

```
dsolve((2*x^2+2*x*y(x)+y(x)^2)+(x^2+2*x*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{3c_1^2 x^2 - \sqrt{-15x^4 c_1^4 + 12x c_1}}{6x c_1^2}$$

$$y(x) = -\frac{3c_1^2 x^2 + \sqrt{-15x^4 c_1^4 + 12x c_1}}{6x c_1^2}$$

✓ Solution by Mathematica

Time used: 0.789 (sec). Leaf size: 150

`DSolve[(2*x^2+2*x*y[x]+y[x]^2)+(x^2+2*x*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{1}{6} \left(-3x - \frac{\sqrt{3}\sqrt{-5x^3 + 4e^{3c_1}}}{\sqrt{x}} \right)$$

$$y(x) \rightarrow \frac{1}{6} \left(-3x + \frac{\sqrt{3}\sqrt{-5x^3 + 4e^{3c_1}}}{\sqrt{x}} \right)$$

$$y(x) \rightarrow \frac{1}{6} x \left(\frac{\sqrt{15}x^{3/2}}{\sqrt{-x^3}} - 3 \right)$$

$$y(x) \rightarrow \frac{\sqrt{\frac{5}{3}}\sqrt{-x^3}}{2\sqrt{x}} - \frac{x}{2}$$