


A Solution Manual For

**AN INTRODUCTION TO  
ORDINARY DIFFERENTIAL  
EQUATIONS** by **JAMES C.  
ROBINSON**. Cambridge  
University Press 2004



**Nasser M. Abbasi**

March 3, 2024

# Contents

1	Chapter 5, Trivial differential equations. Exercises page 33	2
2	Chapter 7, Scalar autonomous ODEs. Exercises page 56	13
3	Chapter 8, Separable equations. Exercises page 72	20
4	Chapter 9, First order linear equations and the integrating factor. Exercises page 86	33
5	Chapter 10, Two tricks for nonlinear equations. Exercises page 97	43
6	Chapter 12, Homogeneous second order linear equations. Exercises page 118	54
7	Chapter 14, Inhomogeneous second order linear equations. Exercises page 140	70
8	Chapter 15, Resonance. Exercises page 148	84
9	Chapter 16, Higher order linear equations with constant coefficients. Exercises page 153	87
10	Chapter 17, Reduction of order. Exercises page 162	92
11	Chapter 18, The variation of constants formula. Exercises page 168	99
12	Chapter 19, CauchyEuler equations. Exercises page 174	106
13	Chapter 20, Series solutions of second order linear equations. Exercises page 195	118
14	Chapter 26, Explicit solutions of coupled linear systems. Exercises page 257	130
15	Chapter 28, Distinct real eigenvalues. Exercises page 282	138
16	Chapter 29, Complex eigenvalues. Exercises page 292	144
17	Chapter 30, A repeated real eigenvalue. Exercises page 299	150

# 1 Chapter 5, Trivial differential equations.

## Exercises page 33

1.1	problem 5.1 (i)	3
1.2	problem 5.1 (ii)	4
1.3	problem 5.1 (iii)	5
1.4	problem 5.1 (iv)	6
1.5	problem 5.1 (v)	7
1.6	problem 5.4 (i)	8
1.7	problem 5.4 (ii)	9
1.8	problem 5.4 (iii)	10
1.9	problem 5.4 (iv)	11
1.10	problem 5.4 (v)	12

## 1.1 problem 5.1 (i)

Internal problem ID [11648]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 5, Trivial differential equations. Exercises page 33

**Problem number:** 5.1 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$x' = \cos(t) + \sin(t)$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve(diff(x(t),t)=sin(t)+cos(t),x(t), singsol=all)
```

$$x(t) = -\cos(t) + \sin(t) + c_1$$

### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 14

```
DSolve[x'[t]==Sin[t]+Cos[t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \sin(t) - \cos(t) + c_1$$

## 1.2 problem 5.1 (ii)

Internal problem ID [11649]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 5, Trivial differential equations. Exercises page 33

**Problem number:** 5.1 (ii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = \frac{1}{x^2 - 1}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)=1/(x^2-1),y(x), singsol=all)
```

$$y(x) = -\operatorname{arctanh}(x) + c_1$$

### ✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 26

```
DSolve[y'[x]==1/(x^2-1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(\log(1-x) - \log(x+1)) + 2c_1$$

### 1.3 problem 5.1 (iii)

Internal problem ID [11650]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 5, Trivial differential equations. Exercises page 33

**Problem number:** 5.1 (iii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_quadrature`]

$$u' = 4t \ln(t)$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(u(t),t)=4*t*ln(t),u(t), singsol=all)
```

$$u(t) = 2t^2 \ln(t) - t^2 + c_1$$

#### ✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 20

```
DSolve[u'[t]==4*t*Log[t],u[t],t,IncludeSingularSolutions -> True]
```

$$u(t) \rightarrow -t^2 + 2t^2 \log(t) + c_1$$

## 1.4 problem 5.1 (iv)

Internal problem ID [11651]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 5, Trivial differential equations. Exercises page 33

**Problem number:** 5.1 (iv).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$z' = x e^{-2x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(z(x),x)=x*exp(-2*x),z(x), singsol=all)
```

$$z(x) = -\frac{(2x + 1)e^{-2x}}{4} + c_1$$

### ✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 22

```
DSolve[z'[x]==x*Exp[-2*x],z[x],x,IncludeSingularSolutions -> True]
```

$$z(x) \rightarrow -\frac{1}{4}e^{-2x}(2x + 1) + c_1$$



## 1.5 problem 5.1 (v)

Internal problem ID [11652]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 5, Trivial differential equations. Exercises page 33

**Problem number:** 5.1 (v).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$T' = e^{-t} \sin(2t)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(T(t),t)=exp(-t)*sin(2*t),T(t), singsol=all)
```

$$T(t) = -\frac{2e^{-t} \cos(2t)}{5} - \frac{e^{-t} \sin(2t)}{5} + c_1$$

### ✓ Solution by Mathematica

Time used: 0.055 (sec). Leaf size: 28

```
DSolve[T'[t]==Exp[-t]*Sin[2*t],T[t],t,IncludeSingularSolutions -> True]
```

$$T(t) \rightarrow -\frac{1}{5}e^{-t}(\sin(2t) + 2 \cos(2t)) + c_1$$

## 1.6 problem 5.4 (i)

Internal problem ID [11653]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 5, Trivial differential equations. Exercises page 33

**Problem number:** 5.4 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_quadrature`]

$$x' = \sec(t)^2$$

With initial conditions

$$\left[ x\left(\frac{\pi}{4}\right) = 0 \right]$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 8

```
dsolve([diff(x(t),t)=sec(t)^2,x(1/4*Pi) = 0],x(t), singsol=all)
```

$$x(t) = \tan(t) - 1$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 9

```
DSolve[{x'[t]==Sec[t]^2,{x[Pi/4]==0}},x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \tan(t) - 1$$

## 1.7 problem 5.4 (ii)

Internal problem ID [11654]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 5, Trivial differential equations. Exercises page 33

**Problem number:** 5.4 (ii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' = x - \frac{1}{3}x^3$$

With initial conditions

$$[y(-1) = 1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve([diff(y(x),x)=x-1/3*x^3,y(-1) = 1],y(x), singsol=all)
```

$$y(x) = -\frac{(x^2 - 3)^2}{12} + \frac{4}{3}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 21

```
DSolve[{y'[x]==x-1/3*x^3,{y[-1]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{12}(-x^4 + 6x^2 + 7)$$

## 1.8 problem 5.4 (iii)

Internal problem ID [11655]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 5, Trivial differential equations. Exercises page 33

**Problem number:** 5.4 (iii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_quadrature`]

$$x' = 2 \sin(t)^2$$

With initial conditions

$$\left[ x\left(\frac{\pi}{4}\right) = \frac{\pi}{4} \right]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([diff(x(t),t)=2*sin(t)^2,x(1/4*Pi) = 1/4*Pi],x(t), singsol=all)
```

$$x(t) = t + \frac{1}{2} - \frac{\sin(2t)}{2}$$

### ✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 16

```
DSolve[{x'[t]==2*Sin[t]^2,{x[Pi/4]==Pi/4}},x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow t - \sin(t) \cos(t) + \frac{1}{2}$$

## 1.9 problem 5.4 (iv)

Internal problem ID [11656]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 5, Trivial differential equations. Exercises page 33

**Problem number:** 5.4 (iv).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_quadrature`]

$$xV' = x^2 + 1$$

With initial conditions

$$[V(1) = 1]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([x*diff(V(x),x)=1+x^2,V(1) = 1],V(x), singsol=all)
```

$$V(x) = \frac{x^2}{2} + \ln(x) + \frac{1}{2}$$

### ✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 18

```
DSolve[{x*V'[x]==1+x^2,{V[1]==1}},V[x],x,IncludeSingularSolutions -> True]
```

$$V(x) \rightarrow \frac{1}{2}(x^2 + 2 \log(x) + 1)$$

## 1.10 problem 5.4 (v)

Internal problem ID [11657]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 5, Trivial differential equations. Exercises page 33

**Problem number:** 5.4 (v).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$3e^{3t}x + e^{3t}x' = e^{-t}$$

With initial conditions

$$[x(0) = 3]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve([diff(x(t)*exp(3*t),t)=exp(-t),x(0) = 3],x(t), singsol=all)
```

$$x(t) = -(e^{-t} - 4)e^{-3t}$$

### ✓ Solution by Mathematica

Time used: 0.092 (sec). Leaf size: 18

```
DSolve[{D[x[t]*Exp[3*t],t]==Exp[-t],{x[0]==3}},x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow e^{-4t}(4e^t - 1)$$

## 2 Chapter 7, Scalar autonomous ODEs. Exercises

page 56

2.1	problem 7.1 (i)	14
2.2	problem 7.1 (ii)	15
2.3	problem 7.1 (iii)	16
2.4	problem 7.1 (iv)	17
2.5	problem 7.1 (v)	19

## 2.1 problem 7.1 (i)

Internal problem ID [11658]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 7, Scalar autonomous ODEs. Exercises page 56

**Problem number:** 7.1 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$x' + x = 1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(x(t),t)=-x(t)+1,x(t), singsol=all)
```

$$x(t) = 1 + e^{-t}c_1$$

### ✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 20

```
DSolve[x'[t]==-x[t]+1,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow 1 + c_1 e^{-t}$$

$$x(t) \rightarrow 1$$



## 2.2 problem 7.1 (ii)

Internal problem ID [11659]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 7, Scalar autonomous ODEs. Exercises page 56

**Problem number:** 7.1 (ii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$x' - x(-x + 2) = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(x(t),t)=x(t)*(2-x(t)),x(t), singsol=all)
```

$$x(t) = \frac{2}{1 + 2e^{-2t}c_1}$$

### ✓ Solution by Mathematica

Time used: 0.503 (sec). Leaf size: 36

```
DSolve[x'[t]==x[t]*(2-x[t]),x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{2e^{2t}}{e^{2t} + e^{2c_1}}$$

$$x(t) \rightarrow 0$$

$$x(t) \rightarrow 2$$

## 2.3 problem 7.1 (iii)

Internal problem ID [11660]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 7, Scalar autonomous ODEs. Exercises page 56

**Problem number:** 7.1 (iii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$x' - (x + 1)(-x + 2) \sin(x) = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 26

```
dsolve(diff(x(t),t)=(1+x(t))*(2-x(t))*sin(x(t)),x(t), singsol=all)
```

$$t + \int^{x(t)} \frac{1}{(\_a + 1)(\_a - 2) \sin(\_a)} d\_a + c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 15.593 (sec). Leaf size: 52

```
DSolve[x'[t]==(1+x[t])*(2-x[t])*Sin[x[t]],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \text{InverseFunction} \left[ \int_1^{\#1} \frac{\csc(K[1])}{(K[1] - 2)(K[1] + 1)} dK[1] \& \right] [-t + c_1]$$

$$x(t) \rightarrow -1$$

$$x(t) \rightarrow 0$$

$$x(t) \rightarrow 2$$

## 2.4 problem 7.1 (iv)

Internal problem ID [11661]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 7, Scalar autonomous ODEs. Exercises page 56

**Problem number:** 7.1 (iv).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$x' + x(1 - x)(-x + 2) = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 22

```
dsolve(diff(x(t),t)=-x(t)*(1-x(t))*(2-x(t)),x(t), singsol=all)
```

$$x(t) = \frac{e^t c_1}{\sqrt{-1 + e^{2t} c_1^2}} + 1$$

✓ Solution by Mathematica

Time used: 19.885 (sec). Leaf size: 159

```
DSolve[x'[t]==-x[t]*(1-x[t])*(2-x[t]),x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{e^{2t} - \sqrt{e^{4t} + e^{2(t+c_1)}} + e^{2c_1}}{e^{2t} + e^{2c_1}}$$

$$x(t) \rightarrow \frac{e^{2t} + \sqrt{e^{4t} + e^{2(t+c_1)}} + e^{2c_1}}{e^{2t} + e^{2c_1}}$$

$$x(t) \rightarrow 0$$

$$x(t) \rightarrow 1$$

$$x(t) \rightarrow 2$$

$$x(t) \rightarrow 1 - e^{-2t} \sqrt{e^{4t}}$$

$$x(t) \rightarrow e^{-2t} \sqrt{e^{4t}} + 1$$

## 2.5 problem 7.1 (v)

Internal problem ID [11662]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 7, Scalar autonomous ODEs. Exercises page 56

**Problem number:** 7.1 (v).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$x' - x^2 + x^4 = 0$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 47

```
dsolve(diff(x(t),t)=x(t)^2-x(t)^4,x(t), singsol=all)
```

$$x(t) = e^{\text{RootOf}(\ln(e^{-Z}-2)e^{-Z}+2c_1e^{-Z}-Z e^{-Z}+2t e^{-Z}-\ln(e^{-Z}-2)-2c_1+_Z-2t+2) - 1}$$

### ✓ Solution by Mathematica

Time used: 0.414 (sec). Leaf size: 53

```
DSolve[x'[t]==x[t]^2-x[t]^4,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \text{InverseFunction} \left[ \frac{1}{\#1} + \frac{1}{2} \log(1 - \#1) - \frac{1}{2} \log(\#1 + 1) \& \right] [-t + c_1]$$

$$x(t) \rightarrow -1$$

$$x(t) \rightarrow 0$$

$$x(t) \rightarrow 1$$

### 3 Chapter 8, Separable equations. Exercises page 72

3.1	problem 8.1 (i)	21
3.2	problem 8.1 (ii)	22
3.3	problem 8.1 (iii)	23
3.4	problem 8.1 (iv)	24
3.5	problem 8.1 (v)	25
3.6	problem 8.2	26
3.7	problem 8.3	27
3.8	problem 8.4	28
3.9	problem 8.5	29
3.10	problem 8.6	30
3.11	problem 8.7	31
3.12	problem 8.8	32

### 3.1 problem 8.1 (i)

Internal problem ID [11663]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 8, Separable equations. Exercises page 72

**Problem number:** 8.1 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$x' - t^3(1 - x) = 0$$

With initial conditions

$$[x(0) = 3]$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve([diff(x(t),t)=t^3*(1-x(t)),x(0) = 3],x(t), singsol=all)
```

$$x(t) = 1 + 2e^{-\frac{t^4}{4}}$$

#### ✓ Solution by Mathematica

Time used: 0.074 (sec). Leaf size: 18

```
DSolve[{x'[t]==t^3*(1-x[t]),{x[0]==3}},x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow 2e^{-\frac{t^4}{4}} + 1$$

### 3.2 problem 8.1 (ii)

Internal problem ID [11664]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 8, Separable equations. Exercises page 72

**Problem number:** 8.1 (ii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_separable`]

$$y' - (y^2 + 1) \tan(x) = 0$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.14 (sec). Leaf size: 12

```
dsolve([diff(y(x),x)=(1+y(x)^2)*tan(x),y(0) = 1],y(x), singsol=all)
```

$$y(x) = \cot\left(\frac{\pi}{4} + \ln(\cos(x))\right)$$

✓ Solution by Mathematica

Time used: 0.472 (sec). Leaf size: 15

```
DSolve[{y'[x]==(1+y[x]^2)*Tan[x],{y[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \cot\left(\log(\cos(x)) + \frac{\pi}{4}\right)$$



### 3.3 problem 8.1 (iii)

Internal problem ID [11665]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 8, Separable equations. Exercises page 72

**Problem number:** 8.1 (iii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$x' - t^2x = 0$$

#### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 12

```
dsolve(diff(x(t),t)=t^2*x(t),x(t), singsol=all)
```

$$x(t) = c_1 e^{\frac{t^3}{3}}$$

#### ✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 22

```
DSolve[x'[t]==t^2*x[t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow c_1 e^{\frac{t^3}{3}}$$

$$x(t) \rightarrow 0$$

### 3.4 problem 8.1 (iv)

Internal problem ID [11666]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 8, Separable equations. Exercises page 72

**Problem number:** 8.1 (iv).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$x' + x^2 = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(x(t),t)=-x(t)^2,x(t), singsol=all)
```

$$x(t) = \frac{1}{t + c_1}$$

#### ✓ Solution by Mathematica

Time used: 0.169 (sec). Leaf size: 18

```
DSolve[x'[t]==-x[t]^2,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{1}{t - c_1}$$

$$x(t) \rightarrow 0$$

### 3.5 problem 8.1 (v)

Internal problem ID [11667]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 8, Separable equations. Exercises page 72

**Problem number:** 8.1 (v).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y' - y^2 e^{-t^2} = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(t),t)=exp(-t^2)*y(t)^2,y(t), singsol=all)
```

$$y(t) = -\frac{2}{\sqrt{\pi} \operatorname{erf}(t) - 2c_1}$$

#### ✓ Solution by Mathematica

Time used: 0.347 (sec). Leaf size: 27

```
DSolve[y'[t]==Exp[-t^2]*y[t]^2,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -\frac{2}{\sqrt{\pi} \operatorname{erf}(t) + 2c_1}$$

$$y(t) \rightarrow 0$$

### 3.6 problem 8.2

Internal problem ID [11668]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 8, Separable equations. Exercises page 72

**Problem number:** 8.2 .

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$x' + px = q$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(x(t),t)+p*x(t)=q,x(t), singsol=all)
```

$$x(t) = \frac{q}{p} + e^{-pt}c_1$$

#### ✓ Solution by Mathematica

Time used: 0.063 (sec). Leaf size: 29

```
DSolve[x'[t]+p*x[t]==q,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{q}{p} + c_1 e^{-pt}$$

$$x(t) \rightarrow \frac{q}{p}$$

### 3.7 problem 8.3

Internal problem ID [11669]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 8, Separable equations. Exercises page 72

**Problem number:** 8.3.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_separable`]

$$xy' - ky = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(x*diff(y(x),x)=k*y(x),y(x), singsol=all)
```

$$y(x) = c_1 x^k$$

#### ✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 16

```
DSolve[x*y'[x]==k*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 x^k$$

$$y(x) \rightarrow 0$$

### 3.8 problem 8.4

Internal problem ID [11670]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 8, Separable equations. Exercises page 72

**Problem number:** 8.4.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$i' - p(t)i = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(diff(i(t),t)=p(t)*i(t),i(t), singsol=all)
```

$$i(t) = c_1 e^{\int p(t) dt}$$

#### ✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 25

```
DSolve[i'[t]==p[t]*i[t],i[t],t,IncludeSingularSolutions -> True]
```

$$i(t) \rightarrow c_1 \exp\left(\int_1^t p(K[1])dK[1]\right)$$

$$i(t) \rightarrow 0$$

### 3.9 problem 8.5

Internal problem ID [11671]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 8, Separable equations. Exercises page 72

**Problem number:** 8.5.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_quadrature]`

$$x' - \lambda x = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(x(t),t)=lambda*x(t),x(t), singsol=all)
```

$$x(t) = c_1 e^{\lambda t}$$

#### ✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 18

```
DSolve[x'[t]==\[Lambda]*x[t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow c_1 e^{\lambda t}$$

$$x(t) \rightarrow 0$$

### 3.10 problem 8.6

Internal problem ID [11672]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 8, Separable equations. Exercises page 72

**Problem number:** 8.6.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$mv' - kv^2 = -mg$$

#### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 29

```
dsolve(m*diff(v(t),t)=-m*g+k*v(t)^2,v(t), singsol=all)
```

$$v(t) = -\frac{\tanh\left(\frac{\sqrt{mgk}(t+c_1)}{m}\right)\sqrt{mgk}}{k}$$

#### ✓ Solution by Mathematica

Time used: 14.167 (sec). Leaf size: 87

```
DSolve[m*v'[t]==-m*g+k*v[t]^2,v[t],t,IncludeSingularSolutions -> True]
```

$$v(t) \rightarrow \frac{\sqrt{g}\sqrt{m}\tanh\left(\frac{\sqrt{g}\sqrt{k}(-t+c_1m)}{\sqrt{m}}\right)}{\sqrt{k}}$$

$$v(t) \rightarrow -\frac{\sqrt{g}\sqrt{m}}{\sqrt{k}}$$

$$v(t) \rightarrow \frac{\sqrt{g}\sqrt{m}}{\sqrt{k}}$$



### 3.11 problem 8.7

Internal problem ID [11673]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 8, Separable equations. Exercises page 72

**Problem number:** 8.7.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$x' - kx + x^2 = 0$$

With initial conditions

$$[x(0) = x_0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 22

```
dsolve([diff(x(t),t)=k*x(t)-x(t)^2,x(0) = x_0],x(t), singsol=all)
```

$$x(t) = \frac{kx_0}{(-x_0 + k)e^{-kt} + x_0}$$

✓ Solution by Mathematica

Time used: 1.052 (sec). Leaf size: 26

```
DSolve[{x'[t]==k*x[t]-x[t]^2,{x[0]==x0}},x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{kx_0e^{kt}}{x_0(e^{kt} - 1) + k}$$

### 3.12 problem 8.8

Internal problem ID [11674]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 8, Separable equations. Exercises page 72

**Problem number:** 8.8.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_quadrature]`

$$x' + x(k^2 + x^2) = 0$$

With initial conditions

$$[x(0) = x_0]$$

**✗** Solution by Maple

```
dsolve([diff(x(t),t)=-x(t)*(k^2+x(t)^2),x(0) = x__0],x(t), singsol=all)
```

No solution found

**✓** Solution by Mathematica

Time used: 1.848 (sec). Leaf size: 62

```
DSolve[{x'[t]==-x[t]*(k^2+x[t]^2)},{x[0]==x0}],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow -\frac{k}{\sqrt{e^{2k^2t} \left( \frac{k^2}{x_0^2} + 1 \right) - 1}}$$

$$x(t) \rightarrow \frac{k}{\sqrt{e^{2k^2t} \left( \frac{k^2}{x_0^2} + 1 \right) - 1}}$$

## 4 Chapter 9, First order linear equations and the integrating factor. Exercises page 86

4.1	problem 9.1 (i)	34
4.2	problem 9.1 (ii)	35
4.3	problem 9.1 (iii)	36
4.4	problem 9.1 (iv)	37
4.5	problem 9.1 (v)	38
4.6	problem 9.1 (vi)	39
4.7	problem 9.1 (vii)	40
4.8	problem 9.1 (viii)	41
4.9	problem 9.4	42

## 4.1 problem 9.1 (i)

Internal problem ID [11675]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 9, First order linear equations and the integrating factor. Exercises page 86

**Problem number:** 9.1 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y' + \frac{y}{x} = x^2$$

With initial conditions

$$[y(0) = y_0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 9

```
dsolve([diff(y(x),x)+y(x)/x=x^2,y(0) = y__0],y(x), singsol=all)
```

$$y(x) = \frac{x^3}{4}$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y'[x]+y[x]/x==x^2,{y[0]==y0}},y[x],x,IncludeSingularSolutions -> True]
```

Not solved

## 4.2 problem 9.1 (ii)

Internal problem ID [11676]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 9, First order linear equations and the integrating factor. Exercises page 86

**Problem number:** 9.1 (ii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$tx + x' = 4t$$

With initial conditions

$$[x(0) = 2]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([diff(x(t),t)+t*x(t)=4*t,x(0) = 2],x(t), singsol=all)
```

$$x(t) = 4 - 2e^{-\frac{t^2}{2}}$$

### ✓ Solution by Mathematica

Time used: 0.066 (sec). Leaf size: 18

```
DSolve[{x'[t]+t*x[t]==4*t,{x[0]==2}},x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow 4 - 2e^{-\frac{t^2}{2}}$$

### 4.3 problem 9.1 (iii)

Internal problem ID [11677]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 9, First order linear equations and the integrating factor. Exercises page 86

**Problem number:** 9.1 (iii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$z' - z \tan(y) = \sin(y)$$

#### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve(diff(z(y),y)=z(y)*tan(y)+sin(y),z(y), singsol=all)
```

$$z(y) = \frac{-\frac{\cos(2y)}{4} + c_1}{\cos(y)}$$

#### ✓ Solution by Mathematica

Time used: 0.075 (sec). Leaf size: 17

```
DSolve[z'[y]==z[y]*Tan[y]+Sin[y],z[y],y,IncludeSingularSolutions -> True]
```

$$z(y) \rightarrow -\frac{\cos(y)}{2} + c_1 \sec(y)$$

## 4.4 problem 9.1 (iv)

Internal problem ID [11678]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 9, First order linear equations and the integrating factor. Exercises page 86

**Problem number:** 9.1 (iv).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y' + e^{-x}y = 1$$

With initial conditions

$$[y(0) = e]$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 23

```
dsolve([diff(y(x),x)+exp(-x)*y(x)=1,y(0) = exp(1)],y(x), singsol=all)
```

$$y(x) = -(-\text{Ei}_1(e^{-x}) - 1 + \text{Ei}_1(1)) e^{e^{-x}}$$

✓ Solution by Mathematica

Time used: 0.104 (sec). Leaf size: 27

```
DSolve[{y'[x]+Exp[-x]*y[x]==1,{y[0]==Exp[1]}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{e^{-x}} (-\text{ExpIntegralEi}(-e^{-x}) + \text{ExpIntegralEi}(-1) + 1)$$

## 4.5 problem 9.1 (v)

Internal problem ID [11679]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 9, First order linear equations and the integrating factor. Exercises page 86

**Problem number:** 9.1 (v).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$x' + x \tanh(t) = 3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(x(t),t)+x(t)*tanh(t)=3,x(t), singsol=all)
```

$$x(t) = \frac{3 \sinh(t) + c_1}{\cosh(t)}$$

### ✓ Solution by Mathematica

Time used: 0.094 (sec). Leaf size: 15

```
DSolve[x'[t]+x[t]*Tanh[t]==3,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \operatorname{sech}(t)(3 \sinh(t) + c_1)$$



## 4.6 problem 9.1 (vi)

Internal problem ID [11680]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 9, First order linear equations and the integrating factor. Exercises page 86

**Problem number:** 9.1 (vi).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y' + 2y \cot(x) = 5$$

With initial conditions

$$\left[ y\left(\frac{\pi}{2}\right) = 1 \right]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve([diff(y(x),x)+2*y(x)*cot(x)=5,y(1/2*Pi) = 1],y(x), singsol=all)
```

$$y(x) = \frac{-10x + 5 \sin(2x) - 4 + 5\pi}{-2 + 2 \cos(2x)}$$

✓ Solution by Mathematica

Time used: 0.082 (sec). Leaf size: 27

```
DSolve[{y'[x]+2*y[x]*Cot[x]==5,{y[Pi/2]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}(10x - 5 \sin(2x) - 5\pi + 4) \csc^2(x)$$

## 4.7 problem 9.1 (vii)

Internal problem ID [11681]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 9, First order linear equations and the integrating factor. Exercises page 86

**Problem number:** 9.1 (vii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$x' + 5x = t$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(x(t),t)+5*x(t)=t,x(t), singsol=all)
```

$$x(t) = \frac{t}{5} - \frac{1}{25} + e^{-5t}c_1$$

### ✓ Solution by Mathematica

Time used: 0.071 (sec). Leaf size: 22

```
DSolve[x'[t]+5*x[t]==t,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{t}{5} + c_1 e^{-5t} - \frac{1}{25}$$

## 4.8 problem 9.1 (viii)

Internal problem ID [11682]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 9, First order linear equations and the integrating factor. Exercises page 86

**Problem number:** 9.1 (viii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$x' + \left(a + \frac{1}{t}\right)x = b$$

With initial conditions

$$[x(1) = x_0]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 38

```
dsolve([diff(x(t),t)+(a+1/t)*x(t)=b,x(1) = x__0],x(t), singsol=all)
```

$$x(t) = \frac{(x_0 a^2 - ba + b) e^{-a(t-1)} + b(at - 1)}{a^2 t}$$

✓ Solution by Mathematica

Time used: 0.101 (sec). Leaf size: 48

```
DSolve[{x'[t]+(a+1/t)*x[t]==b,{x[1]==x0}],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{e^{-at}(e^a a^2 x_0 + be^{at}(at - 1) - (a - 1)e^a b)}{a^2 t}$$

## 4.9 problem 9.4

Internal problem ID [11683]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 9, First order linear equations and the integrating factor. Exercises page 86

**Problem number:** 9.4.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$T' + k(T - \mu - a \cos(\omega(t - \phi))) = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 62

```
dsolve(diff(T(t),t)=-k*(T(t)- (mu+a*cos( omega*(t-phi))))),T(t), singsol=all)
```

$$T(t) = e^{-kt} c_1 - \frac{\sin(\omega(-t + \phi)) ak\omega - \cos(\omega(-t + \phi)) a k^2 - k^2\mu - \mu\omega^2}{k^2 + \omega^2}$$

### ✓ Solution by Mathematica

Time used: 0.511 (sec). Leaf size: 60

```
DSolve[T'[t]==-k*(T[t]- (mu+a*Cos[ omega*(t-phi)])),T[t],t,IncludeSingularSolutions -> True]
```

$$T(t) \rightarrow -\frac{ak\omega \sin(\omega(\phi - t))}{k^2 + \omega^2} + \frac{ak^2 \cos(\omega(\phi - t))}{k^2 + \omega^2} + c_1 e^{-kt} + \mu$$

## 5 Chapter 10, Two tricks for nonlinear equations.

### Exercises page 97

5.1	problem 10.1 (i)	44
5.2	problem 10.1 (ii)	45
5.3	problem 10.1 (iii)	46
5.4	problem 10.1 (iv)	47
5.5	problem 10.2	48
5.6	problem 10.3 (i)	49
5.7	problem 10.3 (ii)	50
5.8	problem 10.4 (i)	51
5.9	problem 10.4 (ii)	52
5.10	problem 10.5	53

## 5.1 problem 10.1 (i)

Internal problem ID [11684]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 10, Two tricks for nonlinear equations. Exercises page 97

**Problem number:** 10.1 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_exact, [_1st_order, ‘_with_symmetry_[F(x),G(x)]’], [_Abel, ‘`

$$2yx + (x^2 + 2y) y' = \sec(x)^2$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 49

```
dsolve((2*x*y(x)- sec(x)^2)+(x^2+2*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{x^2}{2} - \frac{\sqrt{x^4 + 4 \tan(x) - 4c_1}}{2}$$

$$y(x) = -\frac{x^2}{2} + \frac{\sqrt{x^4 + 4 \tan(x) - 4c_1}}{2}$$

### ✓ Solution by Mathematica

Time used: 26.886 (sec). Leaf size: 90

```
DSolve[(2*x*y[x]- Sec[x]^2)+(x^2+2*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left( -x^2 - \sqrt{\sec^2(x) \sqrt{\cos^2(x) (x^4 + 4 \tan(x) + 4c_1)}} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left( -x^2 + \sqrt{\sec^2(x) \sqrt{\cos^2(x) (x^4 + 4 \tan(x) + 4c_1)}} \right)$$

## 5.2 problem 10.1 (ii)

Internal problem ID [11685]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 10, Two tricks for nonlinear equations. Exercises page 97

**Problem number:** 10.1 (ii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y e^x + y x e^x + (x e^x + 2) y' = -1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve((1+exp(x))*y(x)+x*exp(x)*y(x)+(x*exp(x)+2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-x + c_1}{x e^x + 2}$$

### ✓ Solution by Mathematica

Time used: 0.142 (sec). Leaf size: 21

```
DSolve[(1+Exp[x])*y[x]+x*Exp[x]*y[x)+(x*Exp[x]+2)*y'[x]==0,y[x],x,IncludeSingularSolutions -
```

$$y(x) \rightarrow \frac{-x + c_1}{e^x x + 2}$$

### 5.3 problem 10.1 (iii)

Internal problem ID [11686]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 10, Two tricks for nonlinear equations. Exercises page 97

**Problem number:** 10.1 (iii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [exact]

$$(x \cos(y) + \cos(x)) y' + \sin(y) - \sin(x) y = 0$$

#### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 15

```
dsolve((x*cos(y(x))+cos(x))*diff(y(x),x)+sin(y(x))-y(x)*sin(x)=0,y(x), singsol=all)
```

$$\cos(x) y(x) + x \sin(y(x)) + c_1 = 0$$

#### ✓ Solution by Mathematica

Time used: 0.254 (sec). Leaf size: 17

```
DSolve[(x*Cos[y[x]]+Cos[x])*y'[x]+Sin[y[x]]-y[x]*Sin[x]==0,y[x],x,IncludeSingularSolutions -
```

$$\text{Solve}[x \sin(y(x)) + y(x) \cos(x) = c_1, y(x)]$$



## 5.4 problem 10.1 (iv)

Internal problem ID [11687]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 10, Two tricks for nonlinear equations. Exercises page 97

**Problem number:** 10.1 (iv).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_exact]

$$e^x \sin(y) + y + (e^x \cos(y) + x + e^y) y' = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve(exp(x)*sin(y(x))+y(x)+(exp(x)*cos(y(x))+x+exp(y(x)))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x)x + e^x \sin(y(x)) + e^{y(x)} + c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 0.637 (sec). Leaf size: 22

```
DSolve[Exp[x]*Sin[y[x]]+y[x]+(Exp[x]*Cos[y[x]]+x+Exp[y[x]])*y'[x]==0,y[x],x,IncludeSingularSolutions->True]
```

$$\text{Solve}[e^{y(x)} + xy(x) + e^x \sin(y(x)) = c_1, y(x)]$$

## 5.5 problem 10.2

Internal problem ID [11688]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 10, Two tricks for nonlinear equations. Exercises page 97

**Problem number:** 10.2.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x),G(x)]']]`

$$e^{-y} \sec(x) - e^{-y} y' = -2 \cos(x)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 88

```
dsolve(exp(-y(x))*sec(x)+2*cos(x)-exp(-y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \ln \left( \frac{\tan\left(\frac{x}{2}\right)^3 + \tan\left(\frac{x}{2}\right)^2 + \tan\left(\frac{x}{2}\right) + 1}{\tan\left(\frac{x}{2}\right)^3 c_1 + 2 \tan\left(\frac{x}{2}\right)^3 x - \tan\left(\frac{x}{2}\right)^2 c_1 - 2 \tan\left(\frac{x}{2}\right)^2 x + \tan\left(\frac{x}{2}\right) c_1 + 2 \tan\left(\frac{x}{2}\right) x - c_1 - 2x - 4 \tan\left(\frac{x}{2}\right)}$$

✓ Solution by Mathematica

Time used: 2.559 (sec). Leaf size: 33

```
DSolve[Exp[-y[x]]*Sec[x]+2*Cos[x]-Exp[-y[x]]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log \left( \frac{e^{2 \arctanh\left(\tan\left(\frac{x}{2}\right)\right)}}{2(-x + \cos(x) - 2c_1)} \right)$$

## 5.6 problem 10.3 (i)

Internal problem ID [11689]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 10, Two tricks for nonlinear equations. Exercises page 97

**Problem number:** 10.3 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$2yy' = -V'(x)$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve(diff(V(x),x)+2*y(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \sqrt{-V(x) + c_1}$$

$$y(x) = -\sqrt{-V(x) + c_1}$$

### ✓ Solution by Mathematica

Time used: 0.126 (sec). Leaf size: 37

```
DSolve[V'[x]+2*y[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{-V(x) + 2c_1}$$

$$y(x) \rightarrow \sqrt{-V(x) + 2c_1}$$

## 5.7 problem 10.3 (ii)

Internal problem ID [11690]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 10, Two tricks for nonlinear equations. Exercises page 97

**Problem number:** 10.3 (ii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$\left(\frac{1}{y} - a\right) y' = -\frac{2}{x} + b$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 21

```
dsolve((1/y(x)-a)*diff(y(x),x)+2/x-b=0,y(x), singsol=all)
```

$$y(x) = -\frac{\text{LambertW}\left(-\frac{ae^{bx}c_1}{x^2}\right)}{a}$$

### ✓ Solution by Mathematica

Time used: 6.296 (sec). Leaf size: 32

```
DSolve[(1/y[x]-a)*y'[x]+2/x-b==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{W\left(-\frac{ae^{bx}c_1}{x^2}\right)}{a}$$

$$y(x) \rightarrow 0$$

## 5.8 problem 10.4 (i)

Internal problem ID [11691]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 10, Two tricks for nonlinear equations. Exercises page 97

**Problem number:** 10.4 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _Riccati]`

$$yx + y^2 - x^2y' = -x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(x*y(x)+y(x)^2+x^2-x^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \tan(\ln(x) + c_1) x$$

### ✓ Solution by Mathematica

Time used: 0.314 (sec). Leaf size: 13

```
DSolve[x*y[x]+y[x]^2+x^2-x^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \tan(\log(x) + c_1)$$

## 5.9 problem 10.4 (ii)

Internal problem ID [11692]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 10, Two tricks for nonlinear equations. Exercises page 97

**Problem number:** 10.4 (ii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _dAlembert]`

$$x' - \frac{x^2 + t\sqrt{x^2 + t^2}}{tx} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
dsolve(diff(x(t),t)=(x(t)^2+t*sqrt(t^2+x(t)^2))/(t*x(t)),x(t), singsol=all)
```

$$-\frac{\sqrt{t^2 + x(t)^2}}{t} + \ln(t) - c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.512 (sec). Leaf size: 54

```
DSolve[x'[t]==(x[t]^2+t*Sqrt[t^2+x[t]^2))/(t*x[t]),x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow -t\sqrt{\log^2(t) + 2c_1 \log(t) - 1 + c_1^2}$$

$$x(t) \rightarrow t\sqrt{\log^2(t) + 2c_1 \log(t) - 1 + c_1^2}$$

## 5.10 problem 10.5

Internal problem ID [11693]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 10, Two tricks for nonlinear equations. Exercises page 97

**Problem number:** 10.5.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$x' - kx + x^2 = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(x(t),t)=k*x(t)-x(t)^2,x(t), singsol=all)
```

$$x(t) = \frac{k}{1 + e^{-kt}c_1k}$$

### ✓ Solution by Mathematica

Time used: 0.963 (sec). Leaf size: 37

```
DSolve[x'[t]==k*x[t]-x[t]^2,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{ke^{k(t+c_1)}}{-1 + e^{k(t+c_1)}}$$

$$x(t) \rightarrow 0$$

$$x(t) \rightarrow k$$

## 6 Chapter 12, Homogeneous second order linear equations. Exercises page 118

6.1	problem 12.1 (i)	55
6.2	problem 12.1 (ii)	56
6.3	problem 12.1 (iii)	57
6.4	problem 12.1 (iv)	58
6.5	problem 12.1 (v)	59
6.6	problem 12.1 (vi)	60
6.7	problem 12.1 (vii)	61
6.8	problem 12.1 (viii)	62
6.9	problem 12.1 (ix)	63
6.10	problem 12.1 (x)	64
6.11	problem 12.1 (xi)	65
6.12	problem 12.1 (xii)	66
6.13	problem 12.1 (xiii)	67
6.14	problem 12.1 (xiv)	68
6.15	problem 12.1 (xv)	69



## 6.1 problem 12.1 (i)

Internal problem ID [11694]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 12, Homogeneous second order linear equations. Exercises page 118

**Problem number:** 12.1 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$x'' - 3x' + 2x = 0$$

With initial conditions

$$[x(0) = 2, x'(0) = 6]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 15

```
dsolve([diff(x(t),t$2)-3*diff(x(t),t)+2*x(t)=0,x(0) = 2, D(x)(0) = 6],x(t), singsol=all)
```

$$x(t) = -2e^t + 4e^{2t}$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 17

```
DSolve[{x'[t]-3*x'[t]+2*x[t]==0,{x[0]==2,x'[0]==6}},x[t],t,IncludeSingularSolutions -> True
```

$$x(t) \rightarrow 2e^t(2e^t - 1)$$

## 6.2 problem 12.1 (ii)

Internal problem ID [11695]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 12, Homogeneous second order linear equations. Exercises page 118

**Problem number:** 12.1 (ii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y' + 4y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 3]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 11

```
dsolve([diff(y(x),x$2)-4*diff(y(x),x)+4*y(x)=0,y(0) = 0, D(y)(0) = 3],y(x), singsol=all)
```

$$y(x) = 3e^{2x}x$$

### ✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 13

```
DSolve[{y'[x]-4*y'[x]+4*y[x]==0,{y[0]==0,y'[0]==3}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow 3e^{2x}x$$

### 6.3 problem 12.1 (iii)

Internal problem ID [11696]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 12, Homogeneous second order linear equations. Exercises page 118

**Problem number:** 12.1 (iii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$z'' - 4z' + 13z = 0$$

With initial conditions

$$[z(0) = 7, z'(0) = 42]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve([diff(z(t),t$2)-4*diff(z(t),t)+13*z(t)=0,z(0) = 7, D(z)(0) = 42],z(t), singsol=all)
```

$$z(t) = \frac{7e^{2t}(4\sin(3t) + 3\cos(3t))}{3}$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 27

```
DSolve[{z''[t]-4*z'[t]+13*z[t]==0,{z[0]==7,z'[0]==42}},z[t],t,IncludeSingularSolutions -> True]
```

$$z(t) \rightarrow \frac{7}{3}e^{2t}(4\sin(3t) + 3\cos(3t))$$

## 6.4 problem 12.1 (iv)

Internal problem ID [11697]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 12, Homogeneous second order linear equations. Exercises page 118

**Problem number:** 12.1 (iv).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y' - 6y = 0$$

With initial conditions

$$[y(0) = -1, y'(0) = 8]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 15

```
dsolve([diff(y(t),t$2)+diff(y(t),t)-6*y(t)=0,y(0) = -1, D(y)(0) = 8],y(t), singsol=all)
```

$$y(t) = (e^{5t} - 2)e^{-3t}$$

### ✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 18

```
DSolve[{y'[t]+y'[t]-6*y[t]==0,{y[0]==-1,y'[0]==8}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{-3t}(e^{5t} - 2)$$

## 6.5 problem 12.1 (v)

Internal problem ID [11698]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 12, Homogeneous second order linear equations. Exercises page 118

**Problem number:** 12.1 (v).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y' = 0$$

With initial conditions

$$[y(0) = 13, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 5

```
dsolve([diff(y(t),t$2)-4*diff(y(t),t)=0,y(0) = 13, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = 13$$

### ✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 6

```
DSolve[{y'[t]-4*y'[t]==0,{y[0]==13,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 13$$

## 6.6 problem 12.1 (vi)

Internal problem ID [11699]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 12, Homogeneous second order linear equations. Exercises page 118

**Problem number:** 12.1 (vi).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$\theta'' + 4\theta = 0$$

With initial conditions

$$[\theta(0) = 0, \theta'(0) = 10]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve([diff(theta(t),t$2)+4*theta(t)=0,theta(0) = 0, D(theta)(0) = 10],theta(t), singsol=all)
```

$$\theta(t) = 5 \sin(2t)$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 11

```
DSolve[{Theta''[t]+4*Theta[t]==0,{Theta[0]==0,Theta'[0]==10}},Theta[t],t,IncludeSingularSolutions->All]
```

$$\theta(t) \rightarrow 5 \sin(2t)$$

## 6.7 problem 12.1 (vii)

Internal problem ID [11700]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 12, Homogeneous second order linear equations. Exercises page 118

**Problem number:** 12.1 (vii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2y' + 10y = 0$$

With initial conditions

$$[y(0) = 3, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve([diff(y(t),t$2)+2*diff(y(t),t)+10*y(t)=0,y(0) = 3, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = e^{-t}(3 \cos(3t) + \sin(3t))$$

### ✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 22

```
DSolve[{y'[t]+2*y'[t]+10*y[t]==0,{y[0]==3,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{-t}(\sin(3t) + 3 \cos(3t))$$

## 6.8 problem 12.1 (viii)

Internal problem ID [11701]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 12, Homogeneous second order linear equations. Exercises page 118

**Problem number:** 12.1 (viii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$2z'' + 7z' - 4z = 0$$

With initial conditions

$$[z(0) = 0, z'(0) = 9]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve([2*diff(z(t),t$2)+7*diff(z(t),t)-4*z(t)=0,z(0) = 0, D(z)(0) = 9],z(t), singsol=all)
```

$$z(t) = 2\left(e^{\frac{9t}{2}} - 1\right)e^{-4t}$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 49

```
DSolve[{z'[t]+7*z'[t]-4*z[t]==0,{z[0]==3,z'[0]==9}},z[t],t,IncludeSingularSolutions -> True
```

$$z(t) \rightarrow \frac{3}{10}e^{-\frac{1}{2}(7+\sqrt{65})t} \left( (5 + \sqrt{65})e^{\sqrt{65}t} + 5 - \sqrt{65} \right)$$



## 6.9 problem 12.1 (ix)

Internal problem ID [11702]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 12, Homogeneous second order linear equations. Exercises page 118

**Problem number:** 12.1 (ix).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2y' + y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = -1]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve([diff(y(t),t$2)+2*diff(y(t),t)+y(t)=0,y(0) = 0, D(y)(0) = -1],y(t), singsol=all)
```

$$y(t) = -te^{-t}$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 13

```
DSolve[{y'[t]+2*y'[t]+y[t]==0,{y[0]==0,y'[0]==-1}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -e^{-t}t$$

## 6.10 problem 12.1 (x)

Internal problem ID [11703]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 12, Homogeneous second order linear equations. Exercises page 118

**Problem number:** 12.1 (x).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$x'' + 6x' + 10x = 0$$

With initial conditions

$$[x(0) = 3, x'(0) = 1]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve([diff(x(t),t$2)+6*diff(x(t),t)+10*x(t)=0,x(0) = 3, D(x)(0) = 1],x(t), singsol=all)
```

$$x(t) = e^{-3t}(3 \cos(t) + 10 \sin(t))$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 20

```
DSolve[{x''[t]+6*x'[t]+10*x[t]==0,{x[0]==3,x'[0]==1}},x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow e^{-3t}(10 \sin(t) + 3 \cos(t))$$

## 6.11 problem 12.1 (xi)

Internal problem ID [11704]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 12, Homogeneous second order linear equations. Exercises page 118

**Problem number:** 12.1 (xi).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4x'' - 20x' + 21x = 0$$

With initial conditions

$$[x(0) = -4, x'(0) = -12]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve([4*diff(x(t),t$2)-20*diff(x(t),t)+21*x(t)=0,x(0) = -4, D(x)(0) = -12],x(t), singsol=a
```

$$x(t) = -3e^{\frac{7t}{2}} - e^{\frac{3t}{2}}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 23

```
DSolve[{4*x'[t]-20*x'[t]+21*x[t]==0,{x[0]==-4,x'[0]==-12}},x[t],t,IncludeSingularSolutions
```

$$x(t) \rightarrow -e^{3t/2}(3e^{2t} + 1)$$

## 6.12 problem 12.1 (xii)

Internal problem ID [11705]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 12, Homogeneous second order linear equations. Exercises page 118

**Problem number:** 12.1 (xii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y' - 2y = 0$$

With initial conditions

$$[y(0) = 4, y'(0) = -4]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve([diff(y(t),t$2)+diff(y(t),t)-2*y(t)=0,y(0) = 4, D(y)(0) = -4],y(t), singsol=all)
```

$$y(t) = \frac{4(e^{3t} + 2)e^{-2t}}{3}$$

### ✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 21

```
DSolve[{y''[t]+y'[t]-2*y[t]==0,{y[0]==4,y'[0]==-4}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{4}{3}e^{-2t}(e^{3t} + 2)$$

### 6.13 problem 12.1 (xiii)

Internal problem ID [11706]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 12, Homogeneous second order linear equations. Exercises page 118

**Problem number:** 12.1 (xiii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y = 0$$

With initial conditions

$$[y(0) = 10, y'(0) = 0]$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve([diff(y(t),t$2)-4*y(t)=0,y(0) = 10, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = 5e^{-2t} + 5e^{2t}$$

#### ✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 19

```
DSolve[{y'[t]-4*y[t]==0,{y[0]==10,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 5e^{-2t}(e^{4t} + 1)$$

## 6.14 problem 12.1 (xiv)

Internal problem ID [11707]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 12, Homogeneous second order linear equations. Exercises page 118

**Problem number:** 12.1 (xiv).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 4y' + 4y = 0$$

With initial conditions

$$[y(0) = 27, y'(0) = -54]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 10

```
dsolve([diff(y(t),t$2)+4*diff(y(t),t)+4*y(t)=0,y(0) = 27, D(y)(0) = -54],y(t), singsol=all)
```

$$y(t) = 27e^{-2t}$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 12

```
DSolve[{y'[t]+4*y'[t]+4*y[t]==0,{y[0]==27,y'[0]==-54}},y[t],t,IncludeSingularSolutions -> T
```

$$y(t) \rightarrow 27e^{-2t}$$

## 6.15 problem 12.1 (xv)

Internal problem ID [11708]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 12, Homogeneous second order linear equations. Exercises page 118

**Problem number:** 12.1 (xv).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + \omega^2 y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve([diff(y(t),t$2)+omega^2*y(t)=0,y(0) = 0, D(y)(0) = 1],y(t), singsol=all)
```

$$y(t) = \frac{\sin(t\omega)}{\omega}$$

### ✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 13

```
DSolve[{y''[t]+w^2*y[t]==0,{y[0]==0,y'[0]==1}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{\sin(tw)}{w}$$

## 7 Chapter 14, Inhomogeneous second order linear equations. Exercises page 140

7.1	problem 14.1 (i)	71
7.2	problem 14.1 (ii)	72
7.3	problem 14.1 (iii)	73
7.4	problem 14.1 (iv)	74
7.5	problem 14.1 (v)	75
7.6	problem 14.1 (vi)	76
7.7	problem 14.1 (vii)	77
7.8	problem 14.1 (viii)	78
7.9	problem 14.1 (ix)	79
7.10	problem 14.1 (x)	80
7.11	problem 14.1 (xi)	81
7.12	problem 14.2	82
7.13	problem 14.3	83



## 7.1 problem 14.1 (i)

Internal problem ID [11709]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 14, Inhomogeneous second order linear equations. Exercises page 140

**Problem number:** 14.1 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x'' - 4x = t^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(x(t),t$2)-4*x(t)=t^2,x(t), singsol=all)
```

$$x(t) = c_1 e^{2t} + c_2 e^{-2t} - \frac{t^2}{4} - \frac{1}{8}$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 32

```
DSolve[x''[t]-4*x[t]==t^2,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow -\frac{t^2}{4} + c_1 e^{2t} + c_2 e^{-2t} - \frac{1}{8}$$

## 7.2 problem 14.1 (ii)

Internal problem ID [11710]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 14, Inhomogeneous second order linear equations. Exercises page 140

**Problem number:** 14.1 (ii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x'' - 4x' = t^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(x(t),t$2)-4*diff(x(t),t)=t^2,x(t), singsol=all)
```

$$x(t) = -\frac{t^2}{16} - \frac{t^3}{12} + \frac{c_1 e^{4t}}{4} - \frac{t}{32} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.096 (sec). Leaf size: 36

```
DSolve[x''[t]-4*x'[t]==t^2,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{1}{96}(-8t^3 - 6t^2 - 3t + 24c_1 e^{4t} + 96c_2)$$

### 7.3 problem 14.1 (iii)

Internal problem ID [11711]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 14, Inhomogeneous second order linear equations. Exercises page 140

**Problem number:** 14.1 (iii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x'' + x' - 2x = 3e^{-t}$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(x(t),t$2)+diff(x(t),t)-2*x(t)=3*exp(-t),x(t), singsol=all)
```

$$x(t) = c_1 e^t + c_2 e^{-2t} - \frac{3e^{-t}}{2}$$

#### ✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 29

```
DSolve[x''[t]+x'[t]-2*x[t]==3*Exp[-t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow -\frac{3e^{-t}}{2} + c_1 e^{-2t} + c_2 e^t$$

## 7.4 problem 14.1 (iv)

Internal problem ID [11712]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 14, Inhomogeneous second order linear equations. Exercises page 140

**Problem number:** 14.1 (iv).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x'' + x' - 2x = e^t$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(x(t),t$2)+diff(x(t),t)-2*x(t)=exp(t),x(t), singsol=all)
```

$$x(t) = c_1 e^t + c_2 e^{-2t} + \frac{t e^t}{3}$$

### ✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 29

```
DSolve[x''[t]+x'[t]-2*x[t]==Exp[t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow c_1 e^{-2t} + e^t \left( \frac{t}{3} - \frac{1}{9} + c_2 \right)$$

## 7.5 problem 14.1 (v)

Internal problem ID [11713]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 14, Inhomogeneous second order linear equations. Exercises page 140

**Problem number:** 14.1 (v).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x'' + 2x' + x = e^{-t}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(x(t),t$2)+2*diff(x(t),t)+x(t)=exp(-t),x(t), singsol=all)
```

$$x(t) = c_1 t e^{-t} + \frac{e^{-t} t^2}{2} + c_2 e^{-t}$$

### ✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 27

```
DSolve[x''[t]+2*x'[t]+x[t]==Exp[-t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{1}{2} e^{-t} (t^2 + 2c_2 t + 2c_1)$$

## 7.6 problem 14.1 (vi)

Internal problem ID [11714]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 14, Inhomogeneous second order linear equations. Exercises page 140

**Problem number:** 14.1 (vi).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x'' + \omega^2 x = \sin(\alpha t)$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve(diff(x(t),t$2)+omega^2*x(t)=sin(alpha*t),x(t), singsol=all)
```

$$x(t) = \sin(t\omega) c_2 + \cos(t\omega) c_1 - \frac{\sin(\alpha t)}{\alpha^2 - \omega^2}$$

### ✓ Solution by Mathematica

Time used: 0.359 (sec). Leaf size: 56

```
DSolve[x''[t]+w^2*x[t]==Sin[a*t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{-(c_1(a^2 - w^2) \cos(tw)) + c_2(w^2 - a^2) \sin(tw) + \sin(at)}{(w - a)(a + w)}$$

## 7.7 problem 14.1 (vii)

Internal problem ID [11715]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 14, Inhomogeneous second order linear equations. Exercises page 140

**Problem number:** 14.1 (vii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x'' + \omega^2 x = \sin(\omega t)$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```
dsolve(diff(x(t),t$2)+omega^2*x(t)=sin(omega*t),x(t), singsol=all)
```

$$x(t) = \sin(t\omega) c_2 + \cos(t\omega) c_1 + \frac{\sin(t\omega) - \cos(t\omega)\omega t}{2\omega^2}$$

### ✓ Solution by Mathematica

Time used: 0.081 (sec). Leaf size: 29

```
DSolve[x''[t]+w^2*x[t]==Sin[w*t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \left(-\frac{t}{2w} + c_1\right) \cos(tw) + c_2 \sin(tw)$$

## 7.8 problem 14.1 (viii)

Internal problem ID [11716]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 14, Inhomogeneous second order linear equations. Exercises page 140

**Problem number:** 14.1 (viii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x'' + 2x' + 10x = e^{-t}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(x(t),t$2)+2*diff(x(t),t)+10*x(t)=exp(-t),x(t), singsol=all)
```

$$x(t) = e^{-t} \sin(3t) c_2 + e^{-t} \cos(3t) c_1 + \frac{e^{-t}}{9}$$

### ✓ Solution by Mathematica

Time used: 0.086 (sec). Leaf size: 32

```
DSolve[x''[t]+2*x'[t]+10*x[t]==Exp[-t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{1}{9}e^{-t}(9c_2 \cos(3t) + 9c_1 \sin(3t) + 1)$$



## 7.9 problem 14.1 (ix)

Internal problem ID [11717]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 14, Inhomogeneous second order linear equations. Exercises page 140

**Problem number:** 14.1 (ix).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x'' + 2x' + 10x = e^{-t} \cos(3t)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 43

```
dsolve(diff(x(t),t$2)+2*diff(x(t),t)+10*x(t)=exp(-t)*cos(3*t),x(t), singsol=all)
```

$$x(t) = e^{-t} \sin(3t) c_2 + e^{-t} \cos(3t) c_1 + \frac{e^{-t}(\cos(3t) + 3t \sin(3t))}{18}$$

### ✓ Solution by Mathematica

Time used: 0.084 (sec). Leaf size: 38

```
DSolve[x''[t]+2*x'[t]+10*x[t]==Exp[-t]*Cos[3*t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{1}{36} e^{-t} ((1 + 36c_2) \cos(3t) + 6(t + 6c_1) \sin(3t))$$

## 7.10 problem 14.1 (x)

Internal problem ID [11718]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 14, Inhomogeneous second order linear equations. Exercises page 140

**Problem number:** 14.1 (x).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x'' + 6x' + 10x = e^{-2t} \cos(t)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve(diff(x(t),t$2)+6*diff(x(t),t)+10*x(t)=exp(-2*t)*cos(t),x(t), singsol=all)
```

$$x(t) = \sin(t) e^{-3t} c_2 + \cos(t) e^{-3t} c_1 + \frac{e^{-2t}(\cos(t) + 2 \sin(t))}{5}$$

### ✓ Solution by Mathematica

Time used: 0.087 (sec). Leaf size: 33

```
DSolve[x''[t]+6*x'[t]+10*x[t]==Exp[-3*t]*Cos[t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{1}{2} e^{-3t} ((1 + 2c_2) \cos(t) + (t + 2c_1) \sin(t))$$

## 7.11 problem 14.1 (xi)

Internal problem ID [11719]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 14, Inhomogeneous second order linear equations. Exercises page 140

**Problem number:** 14.1 (xi).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x'' + 4x' + 4x = e^{2t}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(x(t),t$2)+4*diff(x(t),t)+4*x(t)=exp(2*t),x(t), singsol=all)
```

$$x(t) = \frac{e^{2t}}{16} + c_1 t e^{-2t} + c_2 e^{-2t}$$

### ✓ Solution by Mathematica

Time used: 0.064 (sec). Leaf size: 28

```
DSolve[x''[t]+4*x'[t]+4*x[t]==Exp[2*t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{e^{2t}}{16} + e^{-2t}(c_2 t + c_1)$$

## 7.12 problem 14.2

Internal problem ID [11720]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 14, Inhomogeneous second order linear equations. Exercises page 140

**Problem number:** 14.2.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x'' + x' - 2x = 12e^{-t} - 6e^t$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(diff(x(t),t$2)+diff(x(t),t)-2*x(t)=12*exp(-t)-6*exp(t),x(t), singsol=all)
```

$$x(t) = c_2 e^{-2t} + c_1 e^t - 6e^{-t} - 2te^t + \frac{2e^t}{3}$$

### ✓ Solution by Mathematica

Time used: 0.093 (sec). Leaf size: 34

```
DSolve[x''[t]+x'[t]-2*x[t]==12*Exp[-t]-6*Exp[t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow e^{-2t} \left( -6e^t + e^{3t} \left( -2t + \frac{2}{3} + c_2 \right) + c_1 \right)$$

## 7.13 problem 14.3

Internal problem ID [11721]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 14, Inhomogeneous second order linear equations. Exercises page 140

**Problem number:** 14.3.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x'' + 4x = 289t e^t \sin(2t)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 48

```
dsolve(diff(x(t),t$2)+4*x(t)=289*t*exp(t)*sin(2*t),x(t), singsol=all)
```

$$x(t) = c_2 \sin(2t) + c_1 \cos(2t) - e^t(68 \cos(2t)t - 17t \sin(2t) - 76 \cos(2t) + 2 \sin(2t))$$

### ✓ Solution by Mathematica

Time used: 0.05 (sec). Leaf size: 40

```
DSolve[x''[t]+4*x[t]==289*t*Exp[t]*Sin[2*t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow (e^t(76 - 68t) + c_1) \cos(2t) + (e^t(17t - 2) + c_2) \sin(2t)$$

## 8 Chapter 15, Resonance. Exercises page 148

8.1	problem 15.1	85
8.2	problem 15.3	86

## 8.1 problem 15.1

Internal problem ID [11722]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 15, Resonance. Exercises page 148

**Problem number:** 15.1.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x'' + \omega^2 x = \cos(\alpha t)$$

With initial conditions

$$[x(0) = 0, x'(0) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve([diff(x(t),t$2)+omega^2*x(t)=cos(alpha*t),x(0) = 0, D(x)(0) = 0],x(t), singsol=all)
```

$$x(t) = \frac{\cos(t\omega) - \cos(\alpha t)}{\alpha^2 - \omega^2}$$

✓ Solution by Mathematica

Time used: 0.25 (sec). Leaf size: 28

```
DSolve[{x''[t]+w^2*x[t]==Cos[a*t],{x[0]==0,x'[0]==0}},x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{\cos(tw) - \cos(at)}{a^2 - w^2}$$

## 8.2 problem 15.3

Internal problem ID [11723]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 15, Resonance. Exercises page 148

**Problem number:** 15.3.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x'' + \omega^2 x = \cos(\omega t)$$

With initial conditions

$$[x(0) = 0, x'(0) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([diff(x(t),t$2)+omega^2*x(t)=cos(omega*t),x(0) = 0, D(x)(0) = 0],x(t), singsol=all)
```

$$x(t) = \frac{\sin(t\omega) t}{2\omega}$$

✓ Solution by Mathematica

Time used: 0.099 (sec). Leaf size: 17

```
DSolve[{x''[t]+w^2*x[t]==Cos[w*t],{x[0]==0,x'[0]==0}},x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{t \sin(tw)}{2w}$$



## 9 Chapter 16, Higher order linear equations with constant coefficients. Exercises page 153

9.1	problem 16.1 (i) . . . . .	88
9.2	problem 16.1 (ii) . . . . .	89
9.3	problem 16.1 (iii) . . . . .	90
9.4	problem 16.1 (iv) . . . . .	91

## 9.1 problem 16.1 (i)

Internal problem ID [11724]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 16, Higher order linear equations with constant coefficients. Exercises page 153

**Problem number:** 16.1 (i).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$x''' - 6x'' + 11x' - 6x = e^{-t}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(x(t),t$3)-6*diff(x(t),t$2)+11*diff(x(t),t)-6*x(t)=exp(-t),x(t), singsol=all)
```

$$x(t) = c_1 e^t + e^{2t} c_2 + c_3 e^{3t} - \frac{e^{-t}}{24}$$

### ✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 37

```
DSolve[x'''[t]-6*x''[t]+11*x'[t]-6*x[t]==Exp[-t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow -\frac{e^{-t}}{24} + c_1 e^t + c_2 e^{2t} + c_3 e^{3t}$$

## 9.2 problem 16.1 (ii)

Internal problem ID [11725]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 16, Higher order linear equations with constant coefficients. Exercises page 153

**Problem number:** 16.1 (ii).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - 3y'' + 2y = \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 59

```
dsolve(diff(y(x),x$3)-3*diff(y(x),x$2)+2*y(x)=sin(x),y(x), singsol=all)
```

$$y(x) = \frac{-15 \sin(x) - 3 \cos(x)}{6(5 + 2\sqrt{3})(-5 + 2\sqrt{3})} + e^x c_1 + e^{(1+\sqrt{3})x} c_2 + e^{-(\sqrt{3}-1)x} c_3$$

### ✓ Solution by Mathematica

Time used: 0.206 (sec). Leaf size: 49

```
DSolve[y'''[x]-3*y''[x]+2*y[x]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{26} \left( 5 \sin(x) + \cos(x) + 26e^x \left( c_1 e^{-\sqrt{3}x} + c_2 e^{\sqrt{3}x} + c_3 \right) \right)$$

### 9.3 problem 16.1 (iii)

Internal problem ID [11726]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 16, Higher order linear equations with constant coefficients. Exercises page 153

**Problem number:** 16.1 (iii).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$x'''' - 4x''' + 8x'' - 8x' + 4x = \sin(t)$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 39

```
dsolve(diff(x(t),t$4)-4*diff(x(t),t$3)+8*diff(x(t),t$2)-8*diff(x(t),t)+4*x(t)=sin(t),x(t), s
```

$$x(t) = -\frac{3 \sin(t)}{25} + \frac{4 \cos(t)}{25} + c_1 e^t \cos(t) + c_2 e^t \sin(t) + c_3 e^t \cos(t) t + c_4 e^t \sin(t) t$$

#### ✓ Solution by Mathematica

Time used: 0.258 (sec). Leaf size: 42

```
DSolve[x''''[t]-4*x'''[t]+8*x''[t]-8*x'[t]+4*x[t]==Sin[t],x[t],t,IncludeSingularSolutions ->
```

$$x(t) \rightarrow \left( \frac{4}{25} + e^t(c_4 t + c_3) \right) \cos(t) + \left( -\frac{3}{25} + e^t(c_2 t + c_1) \right) \sin(t)$$

## 9.4 problem 16.1 (iv)

Internal problem ID [11727]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 16, Higher order linear equations with constant coefficients. Exercises page 153

**Problem number:** 16.1 (iv).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _with_linear_symmetries]]`

$$x'''' - 5x'' + 4x = e^t$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

```
dsolve(diff(x(t),t$4)-5*diff(x(t),t$2)+4*x(t)=exp(t),x(t), singsol=all)
```

$$x(t) = -\frac{te^t}{6} + c_1e^t + c_2e^{-2t} + c_3e^{-t} + e^{2t}c_4$$

### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 45

```
DSolve[x''''[t]-5*x''[t]+4*x[t]==Exp[t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow e^{-2t} \left( c_2e^t + e^{3t} \left( -\frac{t}{6} - \frac{1}{36} + c_3 \right) + c_4e^{4t} + c_1 \right)$$

**10 Chapter 17, Reduction of order. Exercises page  
162**

10.1	problem 17.1	93
10.2	problem 17.2	94
10.3	problem 17.3	95
10.4	problem 17.4	96
10.5	problem 17.5	97
10.6	problem 17.6	98

## 10.1 problem 17.1

Internal problem ID [11728]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 17, Reduction of order. Exercises page 162

**Problem number:** 17.1.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$t^2 y'' - (t^2 + 2t) y' + (t + 2) y = 0$$

Given that one solution of the ode is

$$y_1 = t$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
dsolve([t^2*dif(y(t),t$2)-(t^2+2*t)*dif(y(t),t)+(t+2)*y(t)=0,t],y(t), singsol=all)
```

$$y(t) = c_1 t + c_2 t e^t$$

### ✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 16

```
DSolve[t^2*y'[t]-(t^2+2*t)*y'[t]+(t+2)*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow t(c_2 e^t + c_1)$$

## 10.2 problem 17.2

Internal problem ID [11729]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 17, Reduction of order. Exercises page 162

**Problem number:** 17.2.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x - 1)y'' - xy' + y = 0$$

Given that one solution of the ode is

$$y_1 = e^x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve([(x-1)*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,exp(x)],y(x), singsol=all)
```

$$y(x) = c_1x + c_2e^x$$

### ✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 17

```
DSolve[(x-1)*y''[x]-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1e^x - c_2x$$



### 10.3 problem 17.3

Internal problem ID [11730]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 17, Reduction of order. Exercises page 162

**Problem number:** 17.3.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(\cos(t)t - \sin(t))x'' - x't \sin(t) - x \sin(t) = 0$$

Given that one solution of the ode is

$$x_1 = t$$

#### **X** Solution by Maple

```
dsolve([(t*cos(t)-sin(t))*diff(x(t),t$2)-diff(x(t),t)*t*sin(t)-x(t)*sin(t)=0,t],x(t), singso
```

No solution found

#### **X** Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(t*Cos[t]-Sin[t])*x''[t]-x'[t]*t*Sin[t]-x[t]*Sin[t]==0,x[t],t,IncludeSingularSolution
```

Not solved

## 10.4 problem 17.4

Internal problem ID [11731]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 17, Reduction of order. Exercises page 162

**Problem number:** 17.4.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(-t^2 + t)x'' + (-t^2 + 2)x' + (2 - t)x = 0$$

Given that one solution of the ode is

$$x_1 = e^{-t}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve([(t-t^2)*diff(x(t),t$2)+(2-t^2)*diff(x(t),t)+(2-t)*x(t)=0,exp(-t)],x(t), singsol=all)
```

$$x(t) = \frac{c_1}{t} + c_2 e^{-t}$$

### ✓ Solution by Mathematica

Time used: 0.104 (sec). Leaf size: 42

```
DSolve[(t-t^2)*x''[t]+(2-t^2)*x'[t]+(2-t)*x[t]==0,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{e^{-t}\sqrt{1-t}(c_1 e^t - c_2 t)}{\sqrt{t-1}t}$$

## 10.5 problem 17.5

Internal problem ID [11732]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 17, Reduction of order. Exercises page 162

**Problem number:** 17.5.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [Hermite]

$$y'' - xy' + y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 38

```
dsolve([diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,x],y(x), singsol=all)
```

$$y(x) = c_1x + c_2 \left( i\sqrt{2} \sqrt{\pi} e^{\frac{x^2}{2}} - \pi \operatorname{erf} \left( \frac{i\sqrt{2}x}{2} \right) x \right)$$

### ✓ Solution by Mathematica

Time used: 0.091 (sec). Leaf size: 61

```
DSolve[y''[x]-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{\frac{\pi}{2}}c_2\sqrt{x^2}\operatorname{erfi}\left(\frac{\sqrt{x^2}}{\sqrt{2}}\right) + c_2e^{\frac{x^2}{2}} + \sqrt{2}c_1x$$

## 10.6 problem 17.6

Internal problem ID [11733]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 17, Reduction of order. Exercises page 162

**Problem number:** 17.6.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$\tan(t)x'' - 3x' + (\tan(t) + 3\cot(t))x = 0$$

Given that one solution of the ode is

$$x_1 = \sin(t)$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 15

```
dsolve([tan(t)*diff(x(t),t$2)-3*diff(x(t),t)+(tan(t)+3*cot(t))*x(t)=0,sin(t)],x(t), singsol=
```

$$x(t) = c_1 \sin(t) + c_2 \sin(t) \cos(t)$$

### ✓ Solution by Mathematica

Time used: 0.374 (sec). Leaf size: 24

```
DSolve[Tan[t]*x''[t]-3*x'[t]+(Tan[t]+3*Cot[t])*x[t]==0,x[t],t,IncludeSingularSolutions -> Tr
```

$$x(t) \rightarrow \sqrt{-\sin^2(t)}(c_2 \cos(t) + c_1)$$

## 11 Chapter 18, The variation of constants formula.

### Exercises page 168

11.1	problem 18.1 (i)	100
11.2	problem 18.1 (ii)	101
11.3	problem 18.1 (iii)	102
11.4	problem 18.1 (iv)	103
11.5	problem 18.1 (v)	104
11.6	problem 18.1 (vi)	105

## 11.1 problem 18.1 (i)

Internal problem ID [11734]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 18, The variation of constants formula. Exercises page 168

**Problem number:** 18.1 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - y' - 6y = e^x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-6*y(x)=exp(x),y(x), singsol=all)
```

$$y(x) = c_1 e^{3x} + c_2 e^{-2x} - \frac{e^x}{6}$$

### ✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 29

```
DSolve[y''[x]-y'[x]-6*y[x]==Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{e^x}{6} + c_1 e^{-2x} + c_2 e^{3x}$$

## 11.2 problem 18.1 (ii)

Internal problem ID [11735]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 18, The variation of constants formula. Exercises page 168

**Problem number:** 18.1 (ii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x'' - x = \frac{1}{t}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(x(t),t$2)-x(t)=1/t,x(t), singsol=all)
```

$$x(t) = c_1 e^t + c_2 e^{-t} - \frac{\text{Ei}_1(t) e^t}{2} + \frac{\text{Ei}_1(-t) e^{-t}}{2}$$

### ✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 42

```
DSolve[x''[t]-x[t]==1/t,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{1}{2} e^{-t} (e^{2t} \text{ExpIntegralEi}(-t) - \text{ExpIntegralEi}(t) + 2(c_1 e^{2t} + c_2))$$

### 11.3 problem 18.1 (iii)

Internal problem ID [11736]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 18, The variation of constants formula. Exercises page 168

**Problem number:** 18.1 (iii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y = \cot(2x)$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$2)+4*y(x)=cot(2*x),y(x), singsol=all)
```

$$y(x) = c_1 \cos(2x) + c_2 \sin(2x) + \frac{\sin(2x) \ln(\csc(2x) - \cot(2x))}{4}$$

#### ✓ Solution by Mathematica

Time used: 0.188 (sec). Leaf size: 34

```
DSolve[y''[x]+4*y[x]==Cot[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(2x) + \frac{1}{4} \sin(2x)(\log(\sin(x)) - \log(\cos(x)) + 4c_2)$$



## 11.4 problem 18.1 (iv)

Internal problem ID [11737]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 18, The variation of constants formula. Exercises page 168

**Problem number:** 18.1 (iv).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$t^2 x'' - 2x = t^3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(t^2*diff(x(t),t$2)-2*x(t)=t^3,x(t), singsol=all)
```

$$x(t) = \frac{t^3}{4} + \frac{c_1}{t} + c_2 t^2$$

### ✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 25

```
DSolve[t^2*x''[t]-2*x[t]==t^3,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{t^3}{4} + c_2 t^2 + \frac{c_1}{t}$$

## 11.5 problem 18.1 (v)

Internal problem ID [11738]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 18, The variation of constants formula. Exercises page 168

**Problem number:** 18.1 (v).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x'' - 4x' = \tan(t)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(x(t),t$2)-4*diff(x(t),t)=tan(t),x(t), singsol=all)
```

$$x(t) = \int \left( \int \tan(t) e^{-4t} dt + c_1 \right) e^{4t} dt + c_2$$

### ✓ Solution by Mathematica

Time used: 60.232 (sec). Leaf size: 82

```
DSolve[x''[t]-4*x'[t]==Tan[t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \int_1^t \left( e^{4K[1]} c_1 + \frac{1}{20} \left( -5i \operatorname{Hypergeometric2F1}(2i, 1, 1 + 2i, -e^{2iK[1]}) - (2 - 4i) e^{2iK[1]} \operatorname{Hypergeometric2F1}(1, 1 + 2i, 2 + 2i, -e^{2iK[1]}) \right) \right) dK[1] + c_2$$

## 11.6 problem 18.1 (vi)

Internal problem ID [11739]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 18, The variation of constants formula. Exercises page 168

**Problem number:** 18.1 (vi).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$(\tan(x)^2 - 1)y'' - 4\tan(x)^3y' + 2y\sec(x)^4 = (\tan(x)^2 - 1)(1 - 2\sin(x)^2)$$

Given that one solution of the ode is

$$y_1 = \sec(x)^2$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 29

```
dsolve([(tan(x)^2-1)*diff(y(x),x^2)-4*tan(x)^3*diff(y(x),x)+2*y(x)*sec(x)^4=(tan(x)^2-1)*(1-
```

$$y(x) = \sec(x)^2 c_2 + \sec(x) \sin(x) c_1 - \frac{\cos(x)^2}{4} + \frac{x \tan(x)}{2} + \frac{1}{2}$$

### ✓ Solution by Mathematica

Time used: 0.764 (sec). Leaf size: 66

```
DSolve[(Tan[x]^2-1)*y''[x]-4*Tan[x]^3*y'[x]+2*y[x]*Sec[x]^4==(Tan[x]^2-1)*(1-2*Sin[x]^2),y[x]
```

$$y(x) \rightarrow \sqrt{\sin^2(x)} \sec(x) \arctan\left(\frac{\cos(x)}{1 - \sqrt{\sin^2(x)}}\right) - \frac{1}{4} \cos^2(x) + c_1 \sec^2(x) + c_2 \sqrt{\sin^2(x)} \sec(x) + \frac{1}{2}$$

## 12 Chapter 19, CauchyEuler equations. Exercises

page 174

12.1	problem 19.1 (i)	107
12.2	problem 19.1 (ii)	108
12.3	problem 19.1 (iii)	109
12.4	problem 19.1 (iv)	110
12.5	problem 19.1 (v)	111
12.6	problem 19.1 (vi)	112
12.7	problem 19.1 (vii)	113
12.8	problem 19.1 (viii)	114
12.9	problem 19.1 (ix)	115
12.10	problem 19.1 (x)	116
12.11	problem 19.2	117

## 12.1 problem 19.1 (i)

Internal problem ID [11740]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 19, CauchyEuler equations. Exercises page 174

**Problem number:** 19.1 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$x^2y'' - 4xy' + 6y = 0$$

With initial conditions

$$[y(1) = 0, y'(1) = 1]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve([x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+6*y(x)=0,y(1) = 0, D(y)(1) = 1],y(x), singsol=all)
```

$$y(x) = x^2(x - 1)$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 12

```
DSolve[{x^2*y''[x]-4*x*y'[x]+6*y[x]==0,{y[1]==0,y'[1]==1}},y[x],x,IncludeSingularSolutions -
```

$$y(x) \rightarrow (x - 1)x^2$$

## 12.2 problem 19.1 (ii)

Internal problem ID [11741]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 19, CauchyEuler equations. Exercises page 174

**Problem number:** 19.1 (ii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$4x^2y'' + y = 0$$

With initial conditions

$$[y(1) = 1, y'(1) = 0]$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 13

```
dsolve([4*x^2*diff(y(x),x$2)+y(x)=0,y(1) = 1, D(y)(1) = 0],y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{x}(-2 + \ln(x))}{2}$$

### ✓ Solution by Mathematica

Time used: 0.046 (sec). Leaf size: 47

```
DSolve[{x^2*y'[x]+y[x]==0,{y[1]==1,y'[1]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{3}\sqrt{x}\left(\sqrt{3}\sin\left(\frac{1}{2}\sqrt{3}\log(x)\right) - 3\cos\left(\frac{1}{2}\sqrt{3}\log(x)\right)\right)$$

## 12.3 problem 19.1 (iii)

Internal problem ID [11742]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 19, CauchyEuler equations. Exercises page 174

**Problem number:** 19.1 (iii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$t^2 x'' - 5x't + 10x = 0$$

With initial conditions

$$[x(1) = 2, x'(1) = 1]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 19

```
dsolve([t^2*diff(x(t),t$2)-5*t*diff(x(t),t)+10*x(t)=0,x(1) = 2, D(x)(1) = 1],x(t), singsol=a
```

$$x(t) = t^3(-5 \sin(\ln(t)) + 2 \cos(\ln(t)))$$

### ✓ Solution by Mathematica

Time used: 0.197 (sec). Leaf size: 256

```
DSolve[{t^2*x''[t]-5*t*x'[t]+10*x[t]==0,{x[1]==2,x'[1]==1}},x[t],t,IncludeSingularSolutions -
```

$$x(t) \rightarrow \frac{2\sqrt{t}((\text{BesselI}(-1 - i\sqrt{39}, 2\sqrt{5}) + \text{BesselI}(1 - i\sqrt{39}, 2\sqrt{5})) \text{BesselI}(i\sqrt{39}, 2\sqrt{5}\sqrt{t}) - (\text{BesselI}(-1 + i\sqrt{39}, 2\sqrt{5}) + \text{BesselI}(1 + i\sqrt{39}, 2\sqrt{5})) \text{BesselI}(-i\sqrt{39}, 2\sqrt{5}\sqrt{t}))}{\text{BesselI}(i\sqrt{39}, 2\sqrt{5})(\text{BesselI}(-1 - i\sqrt{39}, 2\sqrt{5}) + \text{BesselI}(1 - i\sqrt{39}, 2\sqrt{5})) - \text{BesselI}(-i\sqrt{39}, 2\sqrt{5})(\text{BesselI}(-1 + i\sqrt{39}, 2\sqrt{5}) + \text{BesselI}(1 + i\sqrt{39}, 2\sqrt{5}))}$$

## 12.4 problem 19.1 (iv)

Internal problem ID [11743]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 19, CauchyEuler equations. Exercises page 174

**Problem number:** 19.1 (iv).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$t^2 x'' + x't - x = 0$$

With initial conditions

$$[x(1) = 1, x'(1) = 1]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 5

```
dsolve([t^2*diff(x(t),t$2)+t*diff(x(t),t)-x(t)=0,x(1) = 1, D(x)(1) = 1],x(t), singsol=all)
```

$$x(t) = t$$

### ✓ Solution by Mathematica

Time used: 0.119 (sec). Leaf size: 172

```
DSolve[{t^2*x''[t]+t*x'[t]-x[t]==0,{x[1]==1,x'[1]==1}},x[t],t,IncludeSingularSolutions -> True]
```

$x(t)$

$$\rightarrow \frac{\sqrt{t}((\text{BesselJ}(\sqrt{5}, 2) - \text{BesselJ}(-1 + \sqrt{5}, 2) + \text{BesselJ}(1 + \sqrt{5}, 2)) \text{BesselJ}(-\sqrt{5}, 2\sqrt{t}) - (\text{BesselJ}(\sqrt{5}, 2) - \text{BesselJ}(-1 - \sqrt{5}, 2) + \text{BesselJ}(1 - \sqrt{5}, 2)) \text{BesselJ}(\sqrt{5}, 2\sqrt{t}))}{\text{BesselJ}(\sqrt{5}, 2) (\text{BesselJ}(-1 - \sqrt{5}, 2) - \text{BesselJ}(1 - \sqrt{5}, 2)) + \text{BesselJ}(-\sqrt{5}, 2) (\text{BesselJ}(-1 + \sqrt{5}, 2) - \text{BesselJ}(1 + \sqrt{5}, 2))}$$



## 12.5 problem 19.1 (v)

Internal problem ID [11744]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 19, CauchyEuler equations. Exercises page 174

**Problem number:** 19.1 (v).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 z'' + 3xz' + 4z = 0$$

With initial conditions

$$[z(1) = 0, z'(1) = 5]$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 19

```
dsolve([x^2*diff(z(x),x$2)+3*x*diff(z(x),x)+4*z(x)=0,z(1) = 0, D(z)(1) = 5],z(x), singsol=al
```

$$z(x) = \frac{5\sqrt{3} \sin(\sqrt{3} \ln(x))}{3x}$$

### ✓ Solution by Mathematica

Time used: 0.145 (sec). Leaf size: 220

```
DSolve[{x^2*z''[x]+3*x*z'[x]+4*z[x]==0,{z[1]==0,z'[1]==5}},z[x],x,IncludeSingularSolutions ->
```

$$z(x) \rightarrow \frac{10\sqrt{x}(\text{BesselJ}(i\sqrt{15}, 2\sqrt{3}) \text{BesselJ}(-i\sqrt{15}, 2\sqrt{3}\sqrt{x}) - \text{BesselJ}(-i\sqrt{15}, 2\sqrt{3}\sqrt{x}))}{\sqrt{3}(\text{BesselJ}(i\sqrt{15}, 2\sqrt{3})(\text{BesselJ}(-1 - i\sqrt{15}, 2\sqrt{3}) - \text{BesselJ}(1 - i\sqrt{15}, 2\sqrt{3})) + \text{BesselJ}(-i\sqrt{15}, 2\sqrt{3}))}$$

## 12.6 problem 19.1 (vi)

Internal problem ID [11745]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 19, CauchyEuler equations. Exercises page 174

**Problem number:** 19.1 (vi).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x^2 y'' - xy' - 3y = 0$$

With initial conditions

$$[y(1) = 1, y'(1) = -1]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 7

```
dsolve([x^2*diff(y(x),x$2)-x*diff(y(x),x)-3*y(x)=0,y(1) = 1, D(y)(1) = -1],y(x), singsol=all
```

$$y(x) = \frac{1}{x}$$

### ✓ Solution by Mathematica

Time used: 0.138 (sec). Leaf size: 169

```
DSolve[{x^2*y''[x]-x*y'[x]-3*y[x]==0,{y[1]==1,y'[1]==-1}},y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow \frac{\sqrt{x}((3 \text{BesselI}(-\sqrt{13}, 2) + \text{BesselI}(-1 - \sqrt{13}, 2) + \text{BesselI}(1 - \sqrt{13}, 2)) \text{BesselI}(\sqrt{13}, 2\sqrt{x}) - (3 \text{BesselI}(\sqrt{13}, 2) (\text{BesselI}(-1 - \sqrt{13}, 2) + \text{BesselI}(1 - \sqrt{13}, 2)) - \text{BesselI}(\sqrt{13}, 2) \text{BesselI}(\sqrt{13}, 2\sqrt{x})))}{\text{BesselI}(\sqrt{13}, 2) (\text{BesselI}(-1 - \sqrt{13}, 2) + \text{BesselI}(1 - \sqrt{13}, 2)) - \text{BesselI}(\sqrt{13}, 2) \text{BesselI}(\sqrt{13}, 2)}$$

## 12.7 problem 19.1 (vii)

Internal problem ID [11746]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 19, CauchyEuler equations. Exercises page 174

**Problem number:** 19.1 (vii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$4t^2 x'' + 8x't + 5x = 0$$

With initial conditions

$$[x(1) = 2, x'(1) = 0]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve([4*t^2*diff(x(t),t$2)+8*t*diff(x(t),t)+5*x(t)=0,x(1) = 2, D(x)(1) = 0],x(t), singsol=
```

$$x(t) = \frac{\sin(\ln(t)) + 2 \cos(\ln(t))}{\sqrt{t}}$$

### ✓ Solution by Mathematica

Time used: 0.101 (sec). Leaf size: 232

```
DSolve[{4*t^2*x'[t]+8*t*x[t]+5*x[t]==0,{x[1]==2,x'[1]==0}},x[t],t,IncludeSingularSolutions
```

$$x(t) \rightarrow \frac{\sqrt{t}((2 \text{BesselJ}(-1 + 2i, 2\sqrt{2}) + \sqrt{2} \text{BesselJ}(2i, 2\sqrt{2}) - 2 \text{BesselJ}(1 + 2i, 2\sqrt{2})) \text{BesselJ}(-2i, 2\sqrt{2}\sqrt{t}) - \text{BesselJ}(-1 + 2i, 2\sqrt{2}) \text{BesselJ}(-2i, 2\sqrt{2}) - \text{BesselJ}(-1 - 2i, 2\sqrt{2}) \text{BesselJ}(2i, 2\sqrt{2}\sqrt{t}))}{\text{BesselJ}(-1 + 2i, 2\sqrt{2}) \text{BesselJ}(-2i, 2\sqrt{2}) - \text{BesselJ}(-1 - 2i, 2\sqrt{2}) \text{BesselJ}(2i, 2\sqrt{2})}$$

## 12.8 problem 19.1 (viii)

Internal problem ID [11747]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 19, CauchyEuler equations. Exercises page 174

**Problem number:** 19.1 (viii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2y'' - 5xy' + 5y = 0$$

With initial conditions

$$[y(1) = -2, y'(1) = 1]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([x^2*diff(y(x),x$2)-5*x*diff(y(x),x)+5*y(x)=0,y(1) = -2, D(y)(1) = 1],y(x), singsol=a
```

$$y(x) = \frac{3}{4}x^5 - \frac{11}{4}x$$

### ✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 17

```
DSolve[{x^2*y''[x]-5*x*y'[x]+5*y[x]==0,{y[1]==-2,y'[1]==1}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{1}{4}x(3x^4 - 11)$$

## 12.9 problem 19.1 (ix)

Internal problem ID [11748]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 19, CauchyEuler equations. Exercises page 174

**Problem number:** 19.1 (ix).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$3x^2z'' + 5xz' - z = 0$$

With initial conditions

$$[z(1) = 2, z'(1) = -1]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([3*x^2*diff(z(x),x$2)+5*x*diff(z(x),x)-z(x)=0,z(1) = 2, D(z)(1) = -1],z(x), singsol=a
```

$$z(x) = \frac{3x^{\frac{4}{3}} + 5}{4x}$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 21

```
DSolve[{3*x^2*z'[x]+5*x*z'[x]-z[x]==0,{z[1]==2,z'[1]==-1}},z[x],x,IncludeSingularSolutions
```

$$z(x) \rightarrow \frac{3x^{4/3} + 5}{4x}$$

## 12.10 problem 19.1 (x)

Internal problem ID [11749]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 19, CauchyEuler equations. Exercises page 174

**Problem number:** 19.1 (x).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$t^2 x'' + 3x't + 13x = 0$$

With initial conditions

$$[x(1) = -1, x'(1) = 2]$$

### ✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 32

```
dsolve([t^2*diff(x(t),t$2)+3*t*diff(x(t),t)+13*x(t)=0,x(1) = -1, D(x)(1) = 2],x(t), singsol=
```

$$x(t) = \frac{\sqrt{3} \sin(2\sqrt{3} \ln(t)) - 6 \cos(2\sqrt{3} \ln(t))}{6t}$$

### ✓ Solution by Mathematica

Time used: 0.051 (sec). Leaf size: 41

```
DSolve[{t^2*x''[t]+3*t*x'[t]+13*x[t]==0,{x[1]==-1,x'[1]==2}},x[t],t,IncludeSingularSolutions
```

$$x(t) \rightarrow \frac{\sqrt{3} \sin(2\sqrt{3} \log(t)) - 6 \cos(2\sqrt{3} \log(t))}{6t}$$

## 12.11 problem 19.2

Internal problem ID [11750]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 19, CauchyEuler equations. Exercises page 174

**Problem number:** 19.2.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$ay'' + (b - a)y' + cy = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 67

```
dsolve(a*diff(y(z),z$2)+(b-a)*diff(y(z),z)+c*y(z)=0,y(z), singsol=all)
```

$$y(z) = c_1 e^{\frac{(-b+a+\sqrt{a^2-2ba-4ca+b^2})z}{2a}} + c_2 e^{-\frac{(b-a+\sqrt{a^2-2ba-4ca+b^2})z}{2a}}$$

### ✓ Solution by Mathematica

Time used: 0.073 (sec). Leaf size: 72

```
DSolve[a*y''[z]+(b-a)*y'[z]+c*y[z]==0,y[z],z,IncludeSingularSolutions -> True]
```

$$y(z) \rightarrow \left( c_2 e^{\frac{z\sqrt{a^2-2a(b+2c)+b^2}}{a}} + c_1 \right) \exp\left( -\frac{z\left(\sqrt{a^2-2a(b+2c)+b^2}-a+b\right)}{2a} \right)$$

## 13 Chapter 20, Series solutions of second order linear equations. Exercises page 195

13.1	problem 20.1	119
13.2	problem 20.2 (i)	120
13.3	problem 20.2 (ii)	121
13.4	problem 20.2 (iii)	122
13.5	problem 20.2 (iv) ( $k=-2$ )	123
13.6	problem 20.2 (iv) ( $k=2$ )	124
13.7	problem 20.3	125
13.8	problem 20.4	126
13.9	problem 20.5	127
13.10	problem 20.7	128



## 13.1 problem 20.1

Internal problem ID [11751]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 20, Series solutions of second order linear equations. Exercises page 195

**Problem number:** 20.1.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [Gegenbauer]

$$(-x^2 + 1)y'' - 2xy' + n(n+1)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 101

```
Order:=6;  
dsolve((1-x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+n*(n+1)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{n(n+1)x^2}{2} + \frac{n(n^3 + 2n^2 - 5n - 6)x^4}{24}\right) y(0) \\ + \left(x - \frac{(n^2 + n - 2)x^3}{6} + \frac{(n^4 + 2n^3 - 13n^2 - 14n + 24)x^5}{120}\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 120

```
AsymptoticDSolveValue[(1-x^2)*y''[x]-2*x*y'[x]+n*(n+1)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{1}{120} (n^2 + n)^2 x^5 + \frac{7}{60} (-n^2 - n) x^5 + \frac{1}{6} (-n^2 - n) x^3 + \frac{x^5}{5} + \frac{x^3}{3} + x \right) \\ + c_1 \left( \frac{1}{24} (n^2 + n)^2 x^4 + \frac{1}{4} (-n^2 - n) x^4 + \frac{1}{2} (-n^2 - n) x^2 + 1 \right)$$

## 13.2 problem 20.2 (i)

Internal problem ID [11752]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 20, Series solutions of second order linear equations. Exercises page 195

**Problem number:** 20.2 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [Hermite]

$$y'' - xy' + y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4\right) y(0) + xD(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 27

```
AsymptoticDSolveValue[y''[x]-x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^4}{24} - \frac{x^2}{2} + 1\right) + c_2 x$$

### 13.3 problem 20.2 (ii)

Internal problem ID [11753]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 20, Series solutions of second order linear equations. Exercises page 195

**Problem number:** 20.2 (ii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$(x^2 + 1)y'' + y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 34

```
Order:=6;  
dsolve((1+x^2)*diff(y(x),x$2)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4\right)y(0) + \left(x - \frac{1}{6}x^3 + \frac{7}{120}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[(1+x^2)*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{7x^5}{120} - \frac{x^3}{6} + x \right) + c_1 \left( \frac{x^4}{8} - \frac{x^2}{2} + 1 \right)$$

### 13.4 problem 20.2 (iii)

Internal problem ID [11754]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 20, Series solutions of second order linear equations. Exercises page 195

**Problem number:** 20.2 (iii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$2xy'' + y' - 2y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

```
Order:=6;  
dsolve(2*x*diff(y(x),x$2)+diff(y(x),x)-2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1\sqrt{x} \left( 1 + \frac{2}{3}x + \frac{2}{15}x^2 + \frac{4}{315}x^3 + \frac{2}{2835}x^4 + \frac{4}{155925}x^5 + O(x^6) \right) \\ + c_2 \left( 1 + 2x + \frac{2}{3}x^2 + \frac{4}{45}x^3 + \frac{2}{315}x^4 + \frac{4}{14175}x^5 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 83

```
AsymptoticDSolveValue[2*x*y'[x]+y'[x]-2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1\sqrt{x} \left( \frac{4x^5}{155925} + \frac{2x^4}{2835} + \frac{4x^3}{315} + \frac{2x^2}{15} + \frac{2x}{3} + 1 \right) \\ + c_2 \left( \frac{4x^5}{14175} + \frac{2x^4}{315} + \frac{4x^3}{45} + \frac{2x^2}{3} + 2x + 1 \right)$$

### 13.5 problem 20.2 (iv) (k=-2)

Internal problem ID [11755]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 20, Series solutions of second order linear equations. Exercises page 195

**Problem number:** 20.2 (iv) (k=-2).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 2xy' - 4y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
Order:=6;  
dsolve(diff(y(x),x$2)-2*x*diff(y(x),x)-4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + 2x^2 + \frac{4}{3}x^4\right) y(0) + \left(x + x^3 + \frac{1}{2}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 36

```
AsymptoticDSolveValue[y'[x]-2*x*y'[x]-4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{x^5}{2} + x^3 + x \right) + c_1 \left( \frac{4x^4}{3} + 2x^2 + 1 \right)$$

## 13.6 problem 20.2 (iv) (k=2)

Internal problem ID [11756]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 20, Series solutions of second order linear equations. Exercises page 195

**Problem number:** 20.2 (iv) (k=2).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 2xy' + 4y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
Order:=6;  
dsolve(diff(y(x),x$2)-2*x*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (-2x^2 + 1)y(0) + \left(x - \frac{1}{3}x^3 - \frac{1}{30}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 33

```
AsymptoticDSolveValue[y''[x]-2*x*y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(1 - 2x^2) + c_2\left(-\frac{x^5}{30} - \frac{x^3}{3} + x\right)$$

## 13.7 problem 20.3

Internal problem ID [11757]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 20, Series solutions of second order linear equations. Exercises page 195

**Problem number:** 20.3.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x(1-x)y'' - 3xy' - y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 60

```
Order:=6;  
dsolve(x*(1-x)*diff(y(x),x$2)-3*x*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1x(1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + O(x^6)) \\ & + (x + 2x^2 + 3x^3 + 4x^4 + 5x^5 + O(x^6)) \ln(x) c_2 \\ & + (1 + 3x + 5x^2 + 7x^3 + 9x^4 + 11x^5 + O(x^6)) c_2 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.059 (sec). Leaf size: 63

```
AsymptoticDSolveValue[x*(1-x)*y''[x]-3*x*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$\begin{aligned} y(x) \rightarrow & c_1(x^4 + x^3 + x^2 + (4x^3 + 3x^2 + 2x + 1)x \log(x) + x + 1) \\ & + c_2(5x^5 + 4x^4 + 3x^3 + 2x^2 + x) \end{aligned}$$

## 13.8 problem 20.4

Internal problem ID [11758]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 20, Series solutions of second order linear equations. Exercises page 195

**Problem number:** 20.4.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + xy' - yx^2 = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 41

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) \left( 1 + \frac{1}{4}x^2 + \frac{1}{64}x^4 + O(x^6) \right) + \left( -\frac{1}{4}x^2 - \frac{3}{128}x^4 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 60

```
AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]-x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{x^4}{64} + \frac{x^2}{4} + 1 \right) + c_2 \left( -\frac{3x^4}{128} - \frac{x^2}{4} + \left( \frac{x^4}{64} + \frac{x^2}{4} + 1 \right) \log(x) \right)$$



## 13.9 problem 20.5

Internal problem ID [11759]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 20, Series solutions of second order linear equations. Exercises page 195

**Problem number:** 20.5.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [Bessel]

$$x^2 y'' + x y' + (x^2 - 1) y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-1)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 x^2 \left(1 - \frac{1}{8} x^2 + \frac{1}{192} x^4 + O(x^6)\right) + c_2 \left(\ln(x) \left(x^2 - \frac{1}{8} x^4 + O(x^6)\right) + \left(-2 + \frac{3}{32} x^4 + O(x^6)\right)\right)}{x}$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 58

```
AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]+(x^2-1)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{x^5}{192} - \frac{x^3}{8} + x \right) + c_1 \left( \frac{1}{16} x (x^2 - 8) \log(x) - \frac{5x^4 - 16x^2 - 64}{64x} \right)$$

## 13.10 problem 20.7

Internal problem ID [11760]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 20, Series solutions of second order linear equations. Exercises page 195

**Problem number:** 20.7.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [Bessel]

$$x^2 y'' + xy' + (-n^2 + x^2)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 77

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-n^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{-n} \left( 1 + \frac{1}{4n-4} x^2 + \frac{1}{32} \frac{1}{(n-2)(n-1)} x^4 + O(x^6) \right) \\ + c_2 x^n \left( 1 - \frac{1}{4n+4} x^2 + \frac{1}{32} \frac{1}{(n+2)(n+1)} x^4 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 160

```
AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]+(x^2-n^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{x^4}{(-n^2 - n + (1 - n)(2 - n) + 2)(-n^2 - n + (3 - n)(4 - n) + 4)} - \frac{x^2}{-n^2 - n + (1 - n)(2 - n) + 2} + 1 \right) x^{-n} + c_1 \left( \frac{x^4}{(-n^2 + n + (n + 1)(n + 2) + 2)(-n^2 + n + (n + 3)(n + 4) + 4)} - \frac{x^2}{-n^2 + n + (n + 1)(n + 2) + 2} + 1 \right) x^n$$

## 14 Chapter 26, Explicit solutions of coupled linear systems. Exercises page 257

14.1	problem 26.1 (i)	131
14.2	problem 26.1 (ii)	132
14.3	problem 26.1 (iii)	133
14.4	problem 26.1 (iv)	134
14.5	problem 26.1 (v)	135
14.6	problem 26.1 (vi)	136
14.7	problem 26.1 (vii)	137

## 14.1 problem 26.1 (i)

Internal problem ID [11761]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 26, Explicit solutions of coupled linear systems. Exercises page 257

**Problem number:** 26.1 (i).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 4x(t) - y(t)$$

$$y'(t) = 2x(t) + y(t) + t^2$$

With initial conditions

$$[x(0) = 0, y(0) = 1]$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 52

```
dsolve([diff(x(t),t) = 4*x(t)-y(t), diff(y(t),t) = 2*x(t)+y(t)+t^2, x(0) = 0, y(0) = 1],[x(t),y(t)])
```

$$x(t) = -\frac{t^2}{6} + \frac{5e^{2t}}{4} - \frac{29e^{3t}}{27} - \frac{5t}{18} - \frac{19}{108}$$

$$y(t) = \frac{5e^{2t}}{2} - \frac{29e^{3t}}{27} - \frac{2t^2}{3} - \frac{7t}{9} - \frac{23}{54}$$

✓ Solution by Mathematica

Time used: 0.128 (sec). Leaf size: 64

```
DSolve[{x'[t]==4*x[t]-y[t],y'[t]==2*x[t]+y[t]+t^2},{x[0]==0,y[0]==1},{x[t],y[t]},t,IncludeSingularSolutions->True]
```

$$x(t) \rightarrow \frac{1}{108}(-18t^2 - 30t + 135e^{2t} - 116e^{3t} - 19)$$

$$y(t) \rightarrow \frac{1}{54}(-36t^2 - 42t + 135e^{2t} - 58e^{3t} - 23)$$

## 14.2 problem 26.1 (ii)

Internal problem ID [11762]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 26, Explicit solutions of coupled linear systems. Exercises page 257

**Problem number:** 26.1 (ii).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= x(t) - 4y(t) + 2 \cos(t)^2 - 1 \\y'(t) &= x(t) + y(t)\end{aligned}$$

With initial conditions

$$[x(0) = 1, y(0) = 1]$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 66

```
dsolve([diff(x(t),t) = x(t)-4*y(t)+cos(2*t), diff(y(t),t) = x(t)+y(t), x(0) = 1, y(0) = 1], [
```

$$x(t) = \frac{26 e^t \cos(2t)}{17} - \frac{32 e^t \sin(2t)}{17} - \frac{9 \cos(2t)}{17} + \frac{2 \sin(2t)}{17}$$

$$y(t) = \frac{13 e^t \sin(2t)}{17} + \frac{16 e^t \cos(2t)}{17} - \frac{4 \sin(2t)}{17} + \frac{\cos(2t)}{17}$$

✓ Solution by Mathematica

Time used: 0.167 (sec). Leaf size: 90

```
DSolve[{x'[t]==x[t]-4*y[t]+cos(2*t), y'[t]==x[t]+y[t]},{x[0]==1,y[0]==1},{x[t],y[t]},t,Includ
```

$$x(t) \rightarrow \frac{1}{25} (2(3 - 5t) \cos - e^t (6 \cos - 25) \cos(2t) + 2e^t (4 \cos - 25) \sin(2t))$$

$$y(t) \rightarrow \frac{1}{50} (4(5t + 2) \cos - 2e^t (4 \cos - 25) \cos(2t) - e^t (6 \cos - 25) \sin(2t))$$

### 14.3 problem 26.1 (iii)

Internal problem ID [11763]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 26, Explicit solutions of coupled linear systems. Exercises page 257

**Problem number:** 26.1 (iii).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= 2x(t) + 2y(t) \\y'(t) &= 6x(t) + 3y(t) + e^t\end{aligned}$$

With initial conditions

$$[x(0) = 0, y(0) = 1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 42

```
dsolve([diff(x(t),t) = 2*x(t)+2*y(t), diff(y(t),t) = 6*x(t)+3*y(t)+exp(t), x(0) = 0, y(0) = 1])
```

$$x(t) = \frac{12 e^{6t}}{35} - \frac{e^{-t}}{7} - \frac{e^t}{5}$$

$$y(t) = \frac{24 e^{6t}}{35} + \frac{3 e^{-t}}{14} + \frac{e^t}{10}$$

✓ Solution by Mathematica

Time used: 0.101 (sec). Leaf size: 58

```
DSolve[{x'[t]==2*x[t]+2*y[t], y'[t]==6*x[t]+3*y[t]+Exp[t]}, {x[0]==0, y[0]==1}, {x[t], y[t]}, t, Integrate -> True]
```

$$x(t) \rightarrow \frac{1}{35} e^{-t} (-7e^{2t} + 12e^{7t} - 5)$$

$$y(t) \rightarrow \frac{1}{70} e^{-t} (7e^{2t} + 48e^{7t} + 15)$$

## 14.4 problem 26.1 (iv)

Internal problem ID [11764]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 26, Explicit solutions of coupled linear systems. Exercises page 257

**Problem number:** 26.1 (iv).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= 5x(t) - 4y(t) + e^{3t} \\y'(t) &= x(t) + y(t)\end{aligned}$$

With initial conditions

$$[x(0) = 1, y(0) = -1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve([diff(x(t),t) = 5*x(t)-4*y(t)+exp(3*t), diff(y(t),t) = x(t)+y(t), x(0) = 1, y(0) = -1
```

$$x(t) = e^{3t}(t^2 + 7t + 1)$$

$$y(t) = \frac{e^{3t}(t^2 + 6t - 2)}{2}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 39

```
DSolve[{x'[t]==5*x[t]-4*y[t]+Exp[3*t], y'[t]==x[t]+y[t]}, {x[0]==1, y[0]==-1}, {x[t], y[t]}, t, Inc
```

$$x(t) \rightarrow e^{3t}(t^2 + 7t + 1)$$

$$y(t) \rightarrow \frac{1}{2}e^{3t}(t^2 + 6t - 2)$$



## 14.5 problem 26.1 (v)

Internal problem ID [11765]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 26, Explicit solutions of coupled linear systems. Exercises page 257

**Problem number:** 26.1 (v).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= 2x(t) + 5y(t) \\y'(t) &= -2x(t) + 4 \cos(t)^3 - 3 \cos(t)\end{aligned}$$

With initial conditions

$$[x(0) = 2, y(0) = -1]$$

✓ Solution by Maple

Time used: 0.093 (sec). Leaf size: 66

```
dsolve([diff(x(t),t) = 2*x(t)+5*y(t), diff(y(t),t) = -2*x(t)+cos(3*t), x(0) = 2, y(0) = -1],
```

$$x(t) = -\frac{16 e^t \sin(3t)}{111} + \frac{69 e^t \cos(3t)}{37} - \frac{30 \sin(3t)}{37} + \frac{5 \cos(3t)}{37}$$

$$y(t) = -\frac{121 e^t \sin(3t)}{111} - \frac{17 e^t \cos(3t)}{37} - \frac{20 \cos(3t)}{37} + \frac{9 \sin(3t)}{37}$$

✓ Solution by Mathematica

Time used: 0.363 (sec). Leaf size: 70

```
DSolve[{x'[t]==2*x[t]+5*y[t],y'[t]==-2*x[t]+Cos[3*t]},{x[0]==2,y[0]==-1},{x[t],y[t]},t,Inclu
```

$$x(t) \rightarrow \frac{1}{111} (3(69e^t + 5) \cos(3t) - 2(8e^t + 45) \sin(3t))$$

$$y(t) \rightarrow \frac{1}{111} (-(121e^t - 27) \sin(3t) - 3(17e^t + 20) \cos(3t))$$

## 14.6 problem 26.1 (vi)

Internal problem ID [11766]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 26, Explicit solutions of coupled linear systems. Exercises page 257

**Problem number:** 26.1 (vi).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= x(t) + y(t) + e^{-t} \\y'(t) &= 4x(t) - 2y(t) + e^{2t}\end{aligned}$$

With initial conditions

$$[x(0) = 1, y(0) = -1]$$

✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 60

```
dsolve([diff(x(t),t) = x(t)+y(t)+exp(-t), diff(y(t),t) = 4*x(t)-2*y(t)+exp(2*t)], x(0) = 1, y(0) = -1)
```

$$x(t) = \frac{62 e^{2t}}{75} + \frac{e^{2t}t}{5} + \frac{17 e^{-3t}}{50} - \frac{e^{-t}}{6}$$

$$y(t) = \frac{77 e^{2t}}{75} - \frac{34 e^{-3t}}{25} + \frac{e^{2t}t}{5} - \frac{2 e^{-t}}{3}$$

✓ Solution by Mathematica

Time used: 0.69 (sec). Leaf size: 67

```
DSolve[{x'[t]==x[t]+y[t]+Exp[-t], y'[t]==4*x[t]-2*y[t]+Exp[2*t]}, {x[0]==1, y[0]==-1}, {x[t], y[t]}
```

$$x(t) \rightarrow \frac{1}{150} e^{-3t} (2e^{5t} (15t + 62) - 25e^{2t} + 51)$$

$$y(t) \rightarrow \frac{1}{75} e^{-3t} (e^{5t} (15t + 77) - 50e^{2t} - 102)$$

## 14.7 problem 26.1 (vii)

Internal problem ID [11767]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 26, Explicit solutions of coupled linear systems. Exercises page 257

**Problem number:** 26.1 (vii).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 8x(t) + 14y(t)$$

$$y'(t) = 7x(t) + y(t)$$

With initial conditions

$$[x(0) = 1, y(0) = 1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve([diff(x(t),t) = 8*x(t)+14*y(t), diff(y(t),t) = 7*x(t)+y(t), x(0) = 1, y(0) = 1],[x(t)
```

$$x(t) = \frac{4e^{15t}}{3} - \frac{e^{-6t}}{3}$$

$$y(t) = \frac{2e^{15t}}{3} + \frac{e^{-6t}}{3}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 44

```
DSolve[{x'[t]==8*x[t]+14*y[t],y'[t]==7*x[t]+y[t]},{x[0]==1,y[0]==1},{x[t],y[t]},t,IncludeSin
```

$$x(t) \rightarrow \frac{1}{3}e^{-6t}(4e^{21t} - 1)$$

$$y(t) \rightarrow \frac{1}{3}e^{-6t}(2e^{21t} + 1)$$

**15 Chapter 28, Distinct real eigenvalues. Exercises**  
**page 282**

15.1	problem 28.2 (i)	139
15.2	problem 28.2 (ii)	140
15.3	problem 28.2 (iii)	141
15.4	problem 28.2 (iv)	142
15.5	problem 28.6 (iii)	143

## 15.1 problem 28.2 (i)

Internal problem ID [11777]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 28, Distinct real eigenvalues. Exercises page 282

**Problem number:** 28.2 (i).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 8x(t) + 14y(t)$$

$$y'(t) = 7x(t) + y(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 36

```
dsolve([diff(x(t),t)=8*x(t)+14*y(t),diff(y(t),t)=7*x(t)+y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = 2c_1e^{15t} - c_2e^{-6t}$$

$$y(t) = c_1e^{15t} + c_2e^{-6t}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 71

```
DSolve[{x'[t]==8*x[t]+14*y[t],y'[t]==7*x[t]+y[t]},{x[t],y[t]},t,IncludeSingularSolutions ->
```

$$x(t) \rightarrow \frac{1}{3}e^{-6t}(c_1(2e^{21t} + 1) + 2c_2(e^{21t} - 1))$$

$$y(t) \rightarrow \frac{1}{3}e^{-6t}(c_1(e^{21t} - 1) + c_2(e^{21t} + 2))$$

## 15.2 problem 28.2 (ii)

Internal problem ID [11778]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 28, Distinct real eigenvalues. Exercises page 282

**Problem number:** 28.2 (ii).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= 2x(t) \\ y'(t) &= -5x(t) - 3y(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve([diff(x(t),t)=2*x(t),diff(y(t),t)=-5*x(t)-3*y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = -c_2 e^{2t}$$

$$y(t) = c_1 e^{-3t} + c_2 e^{2t}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 36

```
DSolve[{x'[t]==2*x[t],y'[t]==-5*x[t]-3*y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow c_1 e^{2t}$$

$$y(t) \rightarrow e^{-3t}(c_1(-e^{5t}) + c_1 + c_2)$$

### 15.3 problem 28.2 (iii)

Internal problem ID [11779]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 28, Distinct real eigenvalues. Exercises page 282

**Problem number:** 28.2 (iii).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 11x(t) - 2y(t)$$

$$y'(t) = 3x(t) + 4y(t)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 36

```
dsolve([diff(x(t),t)=11*x(t)-2*y(t),diff(y(t),t)=3*x(t)+4*y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = 2c_1e^{10t} + \frac{c_2e^{5t}}{3}$$

$$y(t) = c_1e^{10t} + c_2e^{5t}$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 95

```
DSolve[{x'[t]==2*x[t]-2*y[t],y'[t]==3*x[t]+4*y[t]},{x[t],y[t]},t,IncludeSingularSolutions ->
```

$$x(t) \rightarrow \frac{1}{5}e^{3t} \left( 5c_1 \cos(\sqrt{5}t) - \sqrt{5}(c_1 + 2c_2) \sin(\sqrt{5}t) \right)$$

$$y(t) \rightarrow \frac{1}{5}e^{3t} \left( 5c_2 \cos(\sqrt{5}t) + \sqrt{5}(3c_1 + c_2) \sin(\sqrt{5}t) \right)$$

## 15.4 problem 28.2 (iv)

Internal problem ID [11780]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 28, Distinct real eigenvalues. Exercises page 282

**Problem number:** 28.2 (iv).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= x(t) + 20y(t) \\y'(t) &= 40x(t) - 19y(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```
dsolve([diff(x(t),t)=x(t)+20*y(t),diff(y(t),t)=40*x(t)-19*y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = c_1 e^{21t} - \frac{c_2 e^{-39t}}{2}$$

$$y(t) = c_1 e^{21t} + c_2 e^{-39t}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 71

```
DSolve[{x'[t]==x[t]+20*y[t],y'[t]==40*x[t]-19*y[t]},{x[t],y[t]},t,IncludeSingularSolutions -
```

$$x(t) \rightarrow \frac{1}{3} e^{-39t} (c_1 (2e^{60t} + 1) + c_2 (e^{60t} - 1))$$

$$y(t) \rightarrow \frac{1}{3} e^{-39t} (2c_1 (e^{60t} - 1) + c_2 (e^{60t} + 2))$$



## 15.5 problem 28.6 (iii)

Internal problem ID [11781]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 28, Distinct real eigenvalues. Exercises page 282

**Problem number:** 28.6 (iii).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = -2x(t) + 2y(t)$$

$$y'(t) = x(t) - y(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve([diff(x(t),t)=-2*x(t)+2*y(t),diff(y(t),t)=x(t)-y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = -2c_2e^{-3t} + c_1$$

$$y(t) = c_1 + c_2e^{-3t}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 71

```
DSolve[{x'[t]==-2*x[t]+2*y[t],y'[t]==x[t]-y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{1}{3}e^{-3t}(c_1(e^{3t} + 2) + 2c_2(e^{3t} - 1))$$

$$y(t) \rightarrow \frac{1}{3}e^{-3t}(c_1(e^{3t} - 1) + c_2(2e^{3t} + 1))$$

**16 Chapter 29, Complex eigenvalues. Exercises**  
**page 292**

16.1	problem 29.3 (i)	145
16.2	problem 29.3 (ii)	147
16.3	problem 29.3 (iii)	148
16.4	problem 29.3 (iv)	149

## 16.1 problem 29.3 (i)

Internal problem ID [11782]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 29, Complex eigenvalues. Exercises page 292

**Problem number:** 29.3 (i).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= -y(t) \\ y'(t) &= x(t) - y(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 84

```
dsolve([diff(x(t),t)=-y(t),diff(y(t),t)=x(t)-y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = -\frac{e^{-\frac{t}{2}} \left( \sin\left(\frac{\sqrt{3}t}{2}\right) \sqrt{3} c_2 - \cos\left(\frac{\sqrt{3}t}{2}\right) \sqrt{3} c_1 - \sin\left(\frac{\sqrt{3}t}{2}\right) c_1 - \cos\left(\frac{\sqrt{3}t}{2}\right) c_2 \right)}{2}$$

$$y(t) = e^{-\frac{t}{2}} \left( \sin\left(\frac{\sqrt{3}t}{2}\right) c_1 + \cos\left(\frac{\sqrt{3}t}{2}\right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 112

```
DSolve[{x'[t]==-y[t],y'[t]==x[t]-y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{1}{3}e^{-t/2} \left( 3c_1 \cos\left(\frac{\sqrt{3}t}{2}\right) + \sqrt{3}(c_1 - 2c_2) \sin\left(\frac{\sqrt{3}t}{2}\right) \right)$$

$$y(t) \rightarrow \frac{1}{3}e^{-t/2} \left( 3c_2 \cos\left(\frac{\sqrt{3}t}{2}\right) + \sqrt{3}(2c_1 - c_2) \sin\left(\frac{\sqrt{3}t}{2}\right) \right)$$

## 16.2 problem 29.3 (ii)

Internal problem ID [11783]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 29, Complex eigenvalues. Exercises page 292

**Problem number:** 29.3 (ii).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = -2x(t) + 3y(t)$$

$$y'(t) = -6x(t) + 4y(t)$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 54

```
dsolve([diff(x(t),t)=-2*x(t)+3*y(t),diff(y(t),t)=-6*x(t)+4*y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = \frac{e^t(\sin(3t)c_1 + \sin(3t)c_2 - \cos(3t)c_1 + \cos(3t)c_2)}{2}$$

$$y(t) = e^t(\sin(3t)c_1 + \cos(3t)c_2)$$

### ✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 56

```
DSolve[{x'[t]==-2*x[t]+3*y[t],y'[t]==-6*x[t]+4*y[t]},{x[t],y[t]},t,IncludeSingularSolutions
```

$$x(t) \rightarrow e^t(c_1 \cos(3t) + (c_2 - c_1) \sin(3t))$$

$$y(t) \rightarrow e^t(c_2 \cos(3t) + (c_2 - 2c_1) \sin(3t))$$

## 16.3 problem 29.3 (iii)

Internal problem ID [11784]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 29, Complex eigenvalues. Exercises page 292

**Problem number:** 29.3 (iii).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = -11x(t) - 2y(t)$$

$$y'(t) = 13x(t) - 9y(t)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 59

```
dsolve([diff(x(t),t)=-11*x(t)-2*y(t),diff(y(t),t)=13*x(t)-9*y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = -\frac{e^{-10t}(\sin(5t)c_1 + 5\sin(5t)c_2 - 5\cos(5t)c_1 + \cos(5t)c_2)}{13}$$

$$y(t) = e^{-10t}(\sin(5t)c_1 + \cos(5t)c_2)$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 69

```
DSolve[{x'[t]==-11*x[t]-2*y[t],y'[t]==13*x[t]-9*y[t]},{x[t],y[t]},t,IncludeSingularSolutions
```

$$x(t) \rightarrow \frac{1}{5}e^{-10t}(5c_1 \cos(5t) - (c_1 + 2c_2) \sin(5t))$$

$$y(t) \rightarrow \frac{1}{5}e^{-10t}(5c_2 \cos(5t) + (13c_1 + c_2) \sin(5t))$$

## 16.4 problem 29.3 (iv)

Internal problem ID [11785]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 29, Complex eigenvalues. Exercises page 292

**Problem number:** 29.3 (iv).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 7x(t) - 5y(t)$$

$$y'(t) = 10x(t) - 3y(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 58

```
dsolve([diff(x(t),t)=7*x(t)-5*y(t),diff(y(t),t)=10*x(t)-3*y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = \frac{e^{2t}(\sin(5t)c_1 - \sin(5t)c_2 + \cos(5t)c_1 + \cos(5t)c_2)}{2}$$

$$y(t) = e^{2t}(\sin(5t)c_1 + \cos(5t)c_2)$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 62

```
DSolve[{x'[t]==7*x[t]-5*y[t],y'[t]==10*x[t]-3*y[t]},{x[t],y[t]},t,IncludeSingularSolutions -
```

$$x(t) \rightarrow e^{2t}(c_1 \cos(5t) + (c_1 - c_2) \sin(5t))$$

$$y(t) \rightarrow e^{2t}(c_2 \cos(5t) + (2c_1 - c_2) \sin(5t))$$

## 17 Chapter 30, A repeated real eigenvalue.

### Exercises page 299

17.1	problem 30.1 (i)	151
17.2	problem 30.1 (ii)	152
17.3	problem 30.1 (iii)	153
17.4	problem 30.1 (iv)	154
17.5	problem 30.1 (v)	156
17.6	problem 30.5 (iii)	157



## 17.1 problem 30.1 (i)

Internal problem ID [11786]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 30, A repeated real eigenvalue. Exercises page 299

**Problem number:** 30.1 (i).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 5x(t) - 4y(t)$$

$$y'(t) = x(t) + y(t)$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

```
dsolve([diff(x(t),t)=5*x(t)-4*y(t),diff(y(t),t)=x(t)+y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = e^{3t}(2c_2t + 2c_1 + c_2)$$

$$y(t) = e^{3t}(c_2t + c_1)$$

### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 45

```
DSolve[{x'[t]==5*x[t]-4*y[t],y'[t]==x[t]+y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow e^{3t}(2c_1t - 4c_2t + c_1)$$

$$y(t) \rightarrow e^{3t}((c_1 - 2c_2)t + c_2)$$

## 17.2 problem 30.1 (ii)

Internal problem ID [11787]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 30, A repeated real eigenvalue. Exercises page 299

**Problem number:** 30.1 (ii).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = -6x(t) + 2y(t)$$

$$y'(t) = -2x(t) - 2y(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve([diff(x(t),t)=-6*x(t)+2*y(t),diff(y(t),t)=-2*x(t)-2*y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = \frac{e^{-4t}(2c_2t + 2c_1 - c_2)}{2}$$

$$y(t) = e^{-4t}(c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 46

```
DSolve[{x'[t]==-6*x[t]+2*y[t],y'[t]==-2*x[t]-2*y[t]},{x[t],y[t]},t,IncludeSingularSolutions
```

$$x(t) \rightarrow e^{-4t}(-2c_1t + 2c_2t + c_1)$$

$$y(t) \rightarrow e^{-4t}(-2c_1t + 2c_2t + c_2)$$

### 17.3 problem 30.1 (iii)

Internal problem ID [11788]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 30, A repeated real eigenvalue. Exercises page 299

**Problem number:** 30.1 (iii).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = -3x(t) - y(t)$$

$$y'(t) = x(t) - 5y(t)$$

#### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 29

```
dsolve([diff(x(t),t)=-3*x(t)-y(t),diff(y(t),t)=x(t)-5*y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = e^{-4t}(c_2t + c_1 + c_2)$$

$$y(t) = e^{-4t}(c_2t + c_1)$$

#### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 44

```
DSolve[{x'[t]==-3*x[t]-y[t],y'[t]==x[t]-5*y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow e^{-4t}(c_1(t+1) - c_2t)$$

$$y(t) \rightarrow e^{-4t}((c_1 - c_2)t + c_2)$$

## 17.4 problem 30.1 (iv)

Internal problem ID [11789]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 30, A repeated real eigenvalue. Exercises page 299

**Problem number:** 30.1 (iv).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 13x(t)$$

$$y'(t) = 13y(t)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve([diff(x(t),t)=13*x(t),diff(y(t),t)=13*y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = c_1 e^{13t}$$

$$y(t) = c_2 e^{13t}$$

✓ Solution by Mathematica

Time used: 0.065 (sec). Leaf size: 65

```
DSolve[{x'[t]==13*x[t],y'[t]==13*y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow c_1 e^{13t}$$

$$y(t) \rightarrow c_2 e^{13t}$$

$$x(t) \rightarrow c_1 e^{13t}$$

$$y(t) \rightarrow 0$$

$$x(t) \rightarrow 0$$

$$y(t) \rightarrow c_2 e^{13t}$$

$$x(t) \rightarrow 0$$

$$y(t) \rightarrow 0$$

## 17.5 problem 30.1 (v)

Internal problem ID [11790]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 30, A repeated real eigenvalue. Exercises page 299

**Problem number:** 30.1 (v).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 7x(t) - 4y(t)$$

$$y'(t) = x(t) + 3y(t)$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 32

```
dsolve([diff(x(t),t)=7*x(t)-4*y(t),diff(y(t),t)=x(t)+3*y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = e^{5t}(2c_2t + 2c_1 + c_2)$$

$$y(t) = e^{5t}(c_2t + c_1)$$

### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 45

```
DSolve[{x'[t]==7*x[t]-4*y[t],y'[t]==x[t]+3*y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> T
```

$$x(t) \rightarrow e^{5t}(2c_1t - 4c_2t + c_1)$$

$$y(t) \rightarrow e^{5t}((c_1 - 2c_2)t + c_2)$$

## 17.6 problem 30.5 (iii)

Internal problem ID [11791]

**Book:** AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS by JAMES C. ROBINSON. Cambridge University Press 2004

**Section:** Chapter 30, A repeated real eigenvalue. Exercises page 299

**Problem number:** 30.5 (iii).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = -x(t) + y(t)$$

$$y'(t) = -x(t) + y(t)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve([diff(x(t),t)=-x(t)+y(t),diff(y(t),t)=-x(t)+y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = c_1 t - c_1 + c_2$$

$$y(t) = c_1 t + c_2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 32

```
DSolve[{x'[t]==-x[t]+y[t],y'[t]==-x[t]+y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow c_1(-t) + c_2 t + c_1$$

$$y(t) \rightarrow (c_2 - c_1)t + c_2$$