


A Solution Manual For

A treatise on ordinary and
partial differential equations by
William Woolsey Johnson. 1913



Nasser M. Abbasi

March 3, 2024

Contents

1	Chapter 1, Nature and meaning of a differential equation between two variables. page 12	2
2	Chapter 2, Equations of the first order and degree. page 20	11
3	Chapter VII, Solutions in series. Examples XIV. page 177	24
4	Chapter VII, Solutions in series. Examples XV. page 194	43
5	Chapter VII, Solutions in series. Examples XVI. page 220	60
6	Chapter IX, Special forms of differential equations. Examples XVII. page 247	67

1 Chapter 1, Nature and meaning of a differential equation between two variables. page 12

1.1	problem 1	3
1.2	problem 2	4
1.3	problem 3	5
1.4	problem 4	7
1.5	problem 5	8
1.6	problem 9	9

1.1 problem 1

Internal problem ID [4681]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter 1, Nature and meaning of a differential equation between two variables. page 12

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' + y \tan(x) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve(diff(y(x),x)+y(x)*tan(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \cos(x)$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 15

```
DSolve[y'[x]+y[x]*Tan[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(x)$$

$$y(x) \rightarrow 0$$

1.2 problem 2

Internal problem ID [4682]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter 1, Nature and meaning of a differential equation between two variables. page 12

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$x^2y'' - 2y'x + 2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_2x^2 + c_1x$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 14

```
DSolve[x^2*y'[x]-2*x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(c_2x + c_1)$$

1.3 problem 3

Internal problem ID [4683]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter 1, Nature and meaning of a differential equation between two variables. page 12

Problem number: 3.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _dAlembert]`

$$yy'^2 + 2y'x - y = 0$$

✓ Solution by Maple

Time used: 0.172 (sec). Leaf size: 75

```
dsolve(y(x)*diff(y(x),x)^2+2*x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = -ix$$

$$y(x) = ix$$

$$y(x) = 0$$

$$y(x) = \sqrt{c_1^2 - 2c_1x}$$

$$y(x) = \sqrt{c_1^2 + 2c_1x}$$

$$y(x) = -\sqrt{c_1^2 - 2c_1x}$$

$$y(x) = -\sqrt{c_1^2 + 2c_1x}$$

✓ Solution by Mathematica

Time used: 0.462 (sec). Leaf size: 126

```
DSolve[y[x]*(y'[x])^2+2*x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -e^{\frac{c_1}{2}} \sqrt{-2x + e^{c_1}}$$

$$y(x) \rightarrow e^{\frac{c_1}{2}} \sqrt{-2x + e^{c_1}}$$

$$y(x) \rightarrow -e^{\frac{c_1}{2}} \sqrt{2x + e^{c_1}}$$

$$y(x) \rightarrow e^{\frac{c_1}{2}} \sqrt{2x + e^{c_1}}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -ix$$

$$y(x) \rightarrow ix$$

1.4 problem 4

Internal problem ID [4684]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter 1, Nature and meaning of a differential equation between two variables. page 12

Problem number: 4.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$y'^2(-x^2 + 1) = -1$$

✓ Solution by Maple

Time used: 0.172 (sec). Leaf size: 33

```
dsolve(diff(y(x),x)^2*(1-x^2)+1=0,y(x), singsol=all)
```

$$y(x) = \ln(x + \sqrt{x^2 - 1}) + c_1$$

$$y(x) = -\ln(x + \sqrt{x^2 - 1}) + c_1$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 89

```
DSolve[y'[x]^2*(1-x^2)+1==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(\log \left(1 - \frac{x}{\sqrt{x^2 - 1}} \right) - \log \left(\frac{x}{\sqrt{x^2 - 1}} + 1 \right) + 2c_1 \right)$$

$$y(x) \rightarrow \frac{1}{2} \left(-\log \left(1 - \frac{x}{\sqrt{x^2 - 1}} \right) + \log \left(\frac{x}{\sqrt{x^2 - 1}} + 1 \right) + 2c_1 \right)$$

1.5 problem 5

Internal problem ID [4685]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter 1, Nature and meaning of a differential equation between two variables. page 12

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' - ya = e^{xa}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=exp(a*x)+a*y(x),y(x), singsol=all)
```

$$y(x) = (x + c_1) e^{ax}$$

✓ Solution by Mathematica

Time used: 0.044 (sec). Leaf size: 15

```
DSolve[y'[x]==Exp[a*x]+a*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{ax}(x + c_1)$$

1.6 problem 9

Internal problem ID [4686]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter 1, Nature and meaning of a differential equation between two variables. page 12

Problem number: 9.

ODE order: 2.

ODE degree: 2.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$(y'^2 + 1)^3 - a^2 y''^2 = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 96

```
dsolve((diff(y(x),x)^2+1)^3=a^2*(diff(y(x),x$2))^2,y(x), singsol=all)
```

$$y(x) = -ix + c_1$$

$$y(x) = ix + c_1$$

$$y(x) = -\frac{(a+x+c_1)(a-x-c_1)}{\sqrt{a^2-c_1^2-2c_1x-x^2}} + c_2$$

$$y(x) = \frac{(a+x+c_1)(a-x-c_1)}{\sqrt{a^2-c_1^2-2c_1x-x^2}} + c_2$$

✓ Solution by Mathematica

Time used: 0.658 (sec). Leaf size: 141

```
DSolve[(y'[x]^2+1)^3==a^2*(y''[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - i\sqrt{a^2(-1 + c_1^2) - 2ac_1x + x^2}$$

$$y(x) \rightarrow i\sqrt{a^2(-1 + c_1^2) - 2ac_1x + x^2} + c_2$$

$$y(x) \rightarrow c_2 - i\sqrt{a^2(-1 + c_1^2) + 2ac_1x + x^2}$$

$$y(x) \rightarrow i\sqrt{a^2(-1 + c_1^2) + 2ac_1x + x^2} + c_2$$

2 Chapter 2, Equations of the first order and degree. page 20

2.1	problem 1	12
2.2	problem 2	13
2.3	problem 3	14
2.4	problem 4	15
2.5	problem 5	17
2.6	problem 6	20
2.7	problem 7	21
2.8	problem 8	22
2.9	problem 9	23

2.1 problem 1

Internal problem ID [4687]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter 2, Equations of the first order and degree. page 20

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(1+x)y + (1-y)xy' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve((1+x)*y(x)+(1-y(x))*x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\text{LambertW}\left(-\frac{e^{-x}}{c_1 x}\right)$$

✓ Solution by Mathematica

Time used: 3.094 (sec). Leaf size: 28

```
DSolve[(1+x)*y[x]+(1-y[x])*x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -W\left(-\frac{e^{-x-c_1}}{x}\right)$$

$$y(x) \rightarrow 0$$

2.2 problem 2

Internal problem ID [4688]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter 2, Equations of the first order and degree. page 20

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - y^2ax = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x)=a*y(x)^2*x,y(x), singsol=all)
```

$$y(x) = \frac{2}{-ax^2 + 2c_1}$$

✓ Solution by Mathematica

Time used: 0.127 (sec). Leaf size: 24

```
DSolve[y'[x]==a*y[x]^2*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2}{ax^2 + 2c_1}$$

$$y(x) \rightarrow 0$$

2.3 problem 3

Internal problem ID [4689]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter 2, Equations of the first order and degree. page 20

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y^2 + xy^2 + (x^2 - yx^2) y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve((y(x)^2+x*y(x)^2)+(x^2-y(x)*x^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{x \ln(x) + \text{LambertW}\left(-\frac{e^{-c_1 + \frac{1}{x}}}{x}\right) x + c_1 x - 1}{x}}$$

✓ Solution by Mathematica

Time used: 5.302 (sec). Leaf size: 30

```
DSolve[(y[x]^2+x*y[x]^2)+(x^2-y[x]*x^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{W\left(-\frac{e^{\frac{1}{x}-c_1}}{x}\right)}$$

$$y(x) \rightarrow 0$$

2.4 problem 4

Internal problem ID [4690]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter 2, Equations of the first order and degree. page 20

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$xy(x^2 + 1)y' - y^2 = 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
dsolve(x*y(x)*(1+x^2)*diff(y(x),x)=1+y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{(x^2 + 1)(c_1 x^2 - 1)}}{x^2 + 1}$$

$$y(x) = -\frac{\sqrt{(x^2 + 1)(c_1 x^2 - 1)}}{x^2 + 1}$$

✓ Solution by Mathematica

Time used: 1.206 (sec). Leaf size: 131

```
DSolve[x*y[x]*(1+x^2)*y'[x]==1+y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{-1 + (-1 + e^{2c_1}) x^2}}{\sqrt{x^2 + 1}}$$

$$y(x) \rightarrow \frac{\sqrt{-1 + (-1 + e^{2c_1}) x^2}}{\sqrt{x^2 + 1}}$$

$$y(x) \rightarrow -i$$

$$y(x) \rightarrow i$$

$$y(x) \rightarrow -\frac{\sqrt{-x^2 - 1}}{\sqrt{x^2 + 1}}$$

$$y(x) \rightarrow \frac{\sqrt{-x^2 - 1}}{\sqrt{x^2 + 1}}$$

2.5 problem 5

Internal problem ID [4691]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson.
1913

Section: Chapter 2, Equations of the first order and degree. page 20

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$\frac{x}{1+y} - \frac{yy'}{1+x} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 720

`dsolve(x/(1+y(x))=y(x)/(1+x)*diff(y(x),x),y(x), singsol=all)`

$y(x)$

$$= \frac{\left(-1 + 4x^3 + 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}{2} + \frac{1}{2 \left(-1 + 4x^3 + 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}} - \frac{1}{2}$$

$y(x) =$

$$- \frac{\left(-1 + 4x^3 + 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}{4} - \frac{1}{4 \left(-1 + 4x^3 + 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}} - \frac{1}{2} + i\sqrt{3} \left(\frac{\left(-1 + 4x^3 + 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}{2} - \frac{1}{2 \left(-1 + 4x^3 + 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}} \right)$$

$y(x) =$

$$- \frac{\left(-1 + 4x^3 + 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}{4} - \frac{1}{4 \left(-1 + 4x^3 + 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}} - \frac{1}{2} + i\sqrt{3} \left(\frac{\left(-1 + 4x^3 + 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}{2} - \frac{1}{2 \left(-1 + 4x^3 + 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}} \right) + \frac{1}{2}$$

✓ Solution by Mathematica

Time used: 4.125 (sec). Leaf size: 346

`DSolve[x/(1+y[x])==y[x]/(1+x)*y'[x],y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{1}{2} \left(\sqrt[3]{4x^3 + 6x^2 + \sqrt{-1 + (4x^3 + 6x^2 - 1 + 12c_1)^2 - 1 + 12c_1}} \right. \\ \left. + \frac{1}{\sqrt[3]{4x^3 + 6x^2 + \sqrt{-1 + (4x^3 + 6x^2 - 1 + 12c_1)^2 - 1 + 12c_1}} - 1} \right)$$

$$y(x) \rightarrow \frac{1}{8} \left(2i(\sqrt{3} + i) \sqrt[3]{4x^3 + 6x^2 + \sqrt{-1 + (4x^3 + 6x^2 - 1 + 12c_1)^2 - 1 + 12c_1}} \right. \\ \left. + \frac{-2 - 2i\sqrt{3}}{\sqrt[3]{4x^3 + 6x^2 + \sqrt{-1 + (4x^3 + 6x^2 - 1 + 12c_1)^2 - 1 + 12c_1}} - 4} \right)$$

$$y(x) \rightarrow \frac{1}{8} \left(-2(1 + i\sqrt{3}) \sqrt[3]{4x^3 + 6x^2 + \sqrt{-1 + (4x^3 + 6x^2 - 1 + 12c_1)^2 - 1 + 12c_1}} \right. \\ \left. + \frac{2i(\sqrt{3} + i)}{\sqrt[3]{4x^3 + 6x^2 + \sqrt{-1 + (4x^3 + 6x^2 - 1 + 12c_1)^2 - 1 + 12c_1}} - 4} \right)$$

2.6 problem 6

Internal problem ID [4692]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter 2, Equations of the first order and degree. page 20

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' + y^2 b^2 = a^2$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 40

```
dsolve(diff(y(x),x)+b^2*y(x)^2=a^2,y(x), singsol=all)
```

$$y(x) = -\frac{a(e^{-2abc_1-2xba} + 1)}{b(e^{-2abc_1-2xba} - 1)}$$

✓ Solution by Mathematica

Time used: 3.208 (sec). Leaf size: 37

```
DSolve[y'[x]+b^2*y[x]^2==a^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{a \tanh(ab(x + c_1))}{b}$$

$$y(x) \rightarrow -\frac{a}{b}$$

$$y(x) \rightarrow \frac{a}{b}$$

2.7 problem 7

Internal problem ID [4693]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter 2, Equations of the first order and degree. page 20

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{y^2 + 1}{x^2 + 1} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)=(y(x)^2+1)/(x^2+1),y(x), singsol=all)
```

$$y(x) = \tan(\arctan(x) + c_1)$$

✓ Solution by Mathematica

Time used: 0.237 (sec). Leaf size: 25

```
DSolve[y'[x]==(y[x]^2+1)/(x^2+1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan(\arctan(x) + c_1)$$

$$y(x) \rightarrow -i$$

$$y(x) \rightarrow i$$

2.8 problem 8

Internal problem ID [4694]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter 2, Equations of the first order and degree. page 20

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$\sin(x) \cos(y) - \cos(x) \sin(y) y' = 0$$

✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 11

```
dsolve(sin(x)*cos(y(x))=cos(x)*sin(y(x))*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = \arccos\left(\frac{\cos(x)}{c_1}\right)$$

✓ Solution by Mathematica

Time used: 5.183 (sec). Leaf size: 47

```
DSolve[Sin[x]*Cos[y[x]]==Cos[x]*Sin[y[x]]*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\arccos\left(\frac{1}{2}c_1 \cos(x)\right)$$

$$y(x) \rightarrow \arccos\left(\frac{1}{2}c_1 \cos(x)\right)$$

$$y(x) \rightarrow -\frac{\pi}{2}$$

$$y(x) \rightarrow \frac{\pi}{2}$$

2.9 problem 9

Internal problem ID [4695]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter 2, Equations of the first order and degree. page 20

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$axy' + 2y - xyy' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 40

```
dsolve(a*x*diff(y(x),x)+2*y(x)=x*y(x)*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = e^{-\frac{a \operatorname{LambertW}\left(-x^{-\frac{2}{a}} e^{-\frac{2c_1}{a}}\right) + 2 \ln(x) + 2c_1}{a}}$$

✓ Solution by Mathematica

Time used: 60.019 (sec). Leaf size: 29

```
DSolve[a*x*y'[x]+2*y[x]==x*y[x]*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -aW\left(-\frac{e^{\frac{c_1}{a}} x^{-2/a}}{a}\right)$$

3 Chapter VII, Solutions in series. Examples XIV. page 177

3.1	problem 1	25
3.2	problem 2	27
3.3	problem 3	28
3.4	problem 4	29
3.5	problem 5	30
3.6	problem 7	31
3.7	problem 8	32
3.8	problem 9	34
3.9	problem 10	35
3.10	problem 11	36
3.11	problem 13	37
3.12	problem 14	38
3.13	problem 15	40
3.14	problem 16	41
3.15	problem 18	42

3.1 problem 1

Internal problem ID [4696]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XIV. page 177

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$xy'' + (x+n)y' + (n+1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 248

```
Order:=6;  
dsolve(x*diff(y(x),x$2)+(x+n)*diff(y(x),x)+(n+1)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{-n+1} \left(1 + 2 \frac{1}{-2+n} x + 3 \frac{1}{(-3+n)(-2+n)} x^2 + 4 \frac{1}{(n-4)(-3+n)(-2+n)} x^3 \right. \\ \left. + 5 \frac{1}{(n-5)(n-4)(-3+n)(-2+n)} x^4 \right. \\ \left. + 6 \frac{1}{(n-6)(n-5)(n-4)(-3+n)(-2+n)} x^5 + O(x^6) \right) \\ + c_2 \left(1 + \frac{-n-1}{n} x + \frac{1}{2} \frac{n+2}{n} x^2 - \frac{1}{6} \frac{n+3}{n} x^3 + \frac{1}{24} \frac{n+4}{n} x^4 - \frac{1}{120} \frac{n+5}{n} x^5 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 519

AsymptoticDSolveValue[x*y''[x]+(x+n)*y'[x]+(n+1)*y[x]==0,y[x],{x,0,5}]

$$\begin{aligned}
 & y(x) \\
 \rightarrow & c_2 \left(\frac{(-n-1)(n+2)(n+3)(n+4)(n+5)x^5}{n(2n+2)(3n+6)(4n+12)(5n+20)} - \frac{(-n-1)(n+2)(n+3)(n+4)x^4}{n(2n+2)(3n+6)(4n+12)} \right. \\
 & \quad \left. + \frac{(-n-1)(n+2)(n+3)x^3}{n(2n+2)(3n+6)} - \frac{(-n-1)(n+2)x^2}{n(2n+2)} + \frac{(-n-1)x}{n} + 1 \right) \\
 & + c_1 \left(- \frac{720x^5}{((1-n)(2-n)+n(2-n))((2-n)(3-n)+n(3-n))((3-n)(4-n)+n(4-n))((4-n)(5-n))} \right. \\
 & \quad \left. + \frac{120x^4}{((1-n)(2-n)+n(2-n))((2-n)(3-n)+n(3-n))((3-n)(4-n)+n(4-n))((4-n)(5-n))} \right. \\
 & \quad \left. + \frac{24x^3}{((1-n)(2-n)+n(2-n))((2-n)(3-n)+n(3-n))((3-n)(4-n)+n(4-n))} \right. \\
 & \quad \left. - \frac{6x^2}{((1-n)(2-n)+n(2-n))((2-n)(3-n)+n(3-n))((3-n)(4-n)+n(4-n))} \right. \\
 & \quad \left. + \frac{2x}{((1-n)(2-n)+n(2-n))((2-n)(3-n)+n(3-n))} - \frac{2x}{(1-n)(2-n)+n(2-n)} + 1 \right) x^{1-n}
 \end{aligned}$$

3.2 problem 2

Internal problem ID [4697]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XIV. page 177

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y'' + yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)+x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{x^3}{6}\right) y(0) + \left(x - \frac{1}{12}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y''[x]+x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{x^4}{12}\right) + c_1 \left(1 - \frac{x^3}{6}\right)$$

3.3 problem 3

Internal problem ID [4698]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XIV. page 177

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$2x^2y'' - y'x + (-x^2 + 1)y = x^2$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

```
Order:=6;
dsolve(2*x^2*diff(y(x),x$2)-x*diff(y(x),x)+(1-x^2)*y(x)=x^2,y(x),type='series',x=0);
```

$$y(x) = c_1\sqrt{x} \left(1 + \frac{1}{6}x^2 + \frac{1}{168}x^4 + O(x^6) \right) + c_2x \left(1 + \frac{1}{10}x^2 + \frac{1}{360}x^4 + O(x^6) \right) + x^2 \left(\frac{1}{3} + \frac{1}{63}x^2 + O(x^4) \right)$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 160

```
AsymptoticDSolveValue[2*x^2*y''[x]-x*y'[x]+(1-x^2)*y[x]==x^2,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) + c_1\sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + \sqrt{x} \left(-\frac{x^{11/2}}{1980} - \frac{x^{7/2}}{35} - \frac{2x^{3/2}}{3} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + x \left(\frac{x^5}{840} + \frac{x^3}{18} + x \right) \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right)$$

3.4 problem 4

Internal problem ID [4699]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XIV. page 177

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$xy'' + 2y' + a^3x^2y = 2$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 50

```
Order:=6;  
dsolve(x*diff(y(x),x$2)+2*diff(y(x),x)+a^3*x^2*y(x)=2,y(x),type='series',x=0);
```

$$y(x) = c_1 \left(1 - \frac{1}{12} a^3 x^3 + O(x^6) \right) + \frac{c_2 \left(1 - \frac{1}{6} a^3 x^3 + O(x^6) \right)}{x} + x \left(1 - \frac{1}{20} a^3 x^3 + O(x^5) \right)$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 136

```
AsymptoticDSolveValue[x*y''[x]+2*y'[x]+a^3*x^2*y[x]==2,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{a^6 x^6}{504} - \frac{a^3 x^3}{12} + 1 \right) + \frac{c_2 \left(\frac{a^6 x^6}{180} - \frac{a^3 x^3}{6} + 1 \right)}{x} \\ + \left(2x - \frac{a^3 x^4}{12} \right) \left(\frac{a^6 x^6}{504} - \frac{a^3 x^3}{12} + 1 \right) + \frac{\left(\frac{a^3 x^5}{30} - x^2 \right) \left(\frac{a^6 x^6}{180} - \frac{a^3 x^3}{6} + 1 \right)}{x}$$

3.5 problem 5

Internal problem ID [4700]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XIV. page 177

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + ax^2y = 1 + x$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
Order:=6;  
dsolve(diff(y(x),x$2)+a*x^2*y(x)=1+x,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{ax^4}{12}\right)y(0) + \left(x - \frac{1}{20}ax^5\right)D(y)(0) + \frac{x^2}{2} + \frac{x^3}{6} + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 44

```
AsymptoticDSolveValue[y''[x]+a*x^2*y[x]==1+x,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{ax^5}{20}\right) + c_1 \left(1 - \frac{ax^4}{12}\right) + \frac{x^3}{6} + \frac{x^2}{2}$$

3.6 problem 7

Internal problem ID [4701]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XIV. page 177

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^4 y'' + y'x + y = 0$$

With the expansion point for the power series method at $x = 0$.

 Solution by Maple

```
Order:=6;  
dsolve(x^4*diff(y(x),x$2)+x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

No solution found

 Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 49

```
AsymptoticDSolveValue[x^4*y''[x]+x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_1(1-x^2)}{x} + c_2 e^{\frac{1}{2x^2}} (420x^6 + 45x^4 + 6x^2 + 1) x^4$$

3.7 problem 8

Internal problem ID [4702]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XIV. page 177

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + (2x^2 + x) y' - 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 45

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)+(x+2*x^2)*diff(y(x),x)-4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^2 \left(1 - \frac{4}{5}x + \frac{2}{5}x^2 - \frac{16}{105}x^3 + \frac{1}{21}x^4 - \frac{4}{315}x^5 + O(x^6) \right) + \frac{c_2(-144 + 192x - 96x^2 + 32x^4 - \frac{128}{5}x^5 + O(x^6))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 208

```
AsymptoticDSolveValue[x^2*y''[x]+(x+2*x^2)*y'[x]-4*y[x]==2,y[x],{x,0,5}]
```

$$\begin{aligned}
 y(x) \rightarrow & \frac{c_1 \left(\frac{2x^2}{3} - \frac{4x}{3} + 1 \right)}{x^2} + c_2 \left(-\frac{4x^5}{315} + \frac{x^4}{21} - \frac{16x^3}{105} + \frac{2x^2}{5} - \frac{4x}{5} + 1 \right) x^2 \\
 & + \left(-\frac{4x^5}{315} + \frac{x^4}{21} - \frac{16x^3}{105} + \frac{2x^2}{5} - \frac{4x}{5} + 1 \right) \left(\frac{7x^6}{2430} + \frac{19x^5}{2025} + \frac{5x^4}{216} + \frac{2x^3}{45} + \frac{x^2}{18} \right. \\
 & \left. - \frac{1}{4x^2} - \frac{1}{3x} \right) x^2 + \frac{\left(\frac{2x^2}{3} - \frac{4x}{3} + 1 \right) \left(-\frac{x^6}{84} - \frac{4x^5}{105} - \frac{x^4}{10} - \frac{x^3}{5} - \frac{x^2}{4} \right)}{x^2}
 \end{aligned}$$

3.8 problem 9

Internal problem ID [4703]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XIV. page 177

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(-x^2 + x)y'' + 3y' + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 36

```
Order:=6;  
dsolve((x-x^2)*diff(y(x),x$2)+3*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 \left(1 - \frac{2}{3}x + \frac{1}{6}x^2 + O(x^6) \right) + \frac{c_2(-2 + 8x - 12x^2 + 8x^3 - 2x^4 + O(x^6))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.05 (sec). Leaf size: 40

```
AsymptoticDSolveValue[(x-x^2)*y'[x]+3*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(x^2 + \frac{1}{x^2} - 4x - \frac{4}{x} + 6 \right) + c_2 \left(\frac{x^2}{6} - \frac{2x}{3} + 1 \right)$$

3.9 problem 10

Internal problem ID [4704]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XIV. page 177

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`

$$(4x^3 - 14x^2 - 2x)y'' - (6x^2 - 7x + 1)y' + (6x - 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
Order:=6;
```

```
dsolve((4*x^3-14*x^2-2*x)*diff(y(x),x$2)-(6*x^2-7*x+1)*diff(y(x),x)+(6*x-1)*y(x)=0,y(x),type
```

$$y(x) = c_1\sqrt{x}(1 + 2x + O(x^6)) + c_2(1 - x + O(x^6))$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 25

```
AsymptoticDSolveValue[(4*x^3-14*x^2-2*x)*y''[x]-(6*x^2-7*x+1)*y'[x]+(6*x-1)*y[x]==0,y[x],{x,
```

$$y(x) \rightarrow c_1\sqrt{x}(2x + 1) + c_2(1 - x)$$

3.10 problem 11

Internal problem ID [4705]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XIV. page 177

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + x^2 y' + (x - 2)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 43

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)+x^2*diff(y(x),x)+(x-2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^2 \left(1 - \frac{3}{4}x + \frac{3}{10}x^2 - \frac{1}{12}x^3 + \frac{1}{56}x^4 - \frac{1}{320}x^5 + O(x^6) \right) \\ + \frac{c_2 (12 - 2x^3 + \frac{3}{2}x^4 - \frac{3}{5}x^5 + O(x^6))}{x}$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 60

```
AsymptoticDSolveValue[x^2*y''[x]+x^2*y'[x]+(x-2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^3}{8} - \frac{x^2}{6} + \frac{1}{x} \right) + c_2 \left(\frac{x^6}{56} - \frac{x^5}{12} + \frac{3x^4}{10} - \frac{3x^3}{4} + x^2 \right)$$

3.11 problem 13

Internal problem ID [4706]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XIV. page 177

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - x^2 y' + (x - 2)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 47

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)-x^2*diff(y(x),x)+(x-2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^2 \left(1 + \frac{1}{4}x + \frac{1}{20}x^2 + \frac{1}{120}x^3 + \frac{1}{840}x^4 + \frac{1}{6720}x^5 + O(x^6) \right) \\ + \frac{c_2 (12 + 12x + 6x^2 + 2x^3 + \frac{1}{2}x^4 + \frac{1}{10}x^5 + O(x^6))}{x}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 66

```
AsymptoticDSolveValue[x^2*y''[x]-x^2*y'[x]+(x-2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^3}{24} + \frac{x^2}{6} + \frac{x}{2} + \frac{1}{x} + 1 \right) + c_2 \left(\frac{x^6}{840} + \frac{x^5}{120} + \frac{x^4}{20} + \frac{x^3}{4} + x^2 \right)$$

3.12 problem 14

Internal problem ID [4707]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XIV. page 177

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(1 - 4x)y'' + ((-n + 1)x - (6 - 4n)x^2)y' + n(-n + 1)xy = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 471

```
Order:=6;
```

```
dsolve(x^2*(1-4*x)*diff(y(x),x$2)+((-1-n)*x-(6-4*n)*x^2)*diff(y(x),x)+n*(1-n)*x*y(x)=0,y(x),t
```

$$y(x) = x^n c_1 \left(1 + nx + \frac{1}{2}n(n+3)x^2 + \frac{1}{6}(n+5)(n+4)nx^3 + \frac{1}{24}n(n+5)(n+7)(n+6)x^4 + \frac{1}{120}(n+9)(n+8)(n+7)(n+6)nx^5 + O(x^6) \right) + \left(1 - nx + \frac{1}{2}n(-3+n)x^2 - \frac{1}{6}(n-4)(n-5)nx^3 + \frac{1}{24}n(n-5)(n-6)(n-7)x^4 - \frac{1}{120}(n-6)(n-7)(n-8)(n-9)nx^5 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 2114

AsymptoticDSolveValue[x^2*(1-4*x)*y'[x]+((1-n)*x-(6-4*n)*x^2)*y'[x]+n*(1-n)*x*y[x]==0,y[x],

$$\begin{aligned}
 & y(x) \\
 & \left(\left(512n - 256(n - n^2) - \frac{(n^2+n)(64(n-n^2)-128(n+1))}{(1-n)(n+1)+n(n+1)} - \frac{(16(n-n^2)-32(n+2)) \left(8n-4(n-n^2) - \frac{(n^2+n)(-n^2+n-2(n+1))}{(1-n)(n+1)+n(n+1)} \right)}{(1-n)(n+2)+(n+1)(n+2)} \right) \right. \\
 & \left. + \left(128n - 64(n - n^2) - \frac{(n^2+n)(16(n-n^2)-32(n+1))}{(1-n)(n+1)+n(n+1)} - \frac{(4(n-n^2)-8(n+2)) \left(8n-4(n-n^2) - \frac{(n^2+n)(-n^2+n-2(n+1))}{(1-n)(n+1)+n(n+1)} \right)}{(1-n)(n+2)+(n+1)(n+2)} \right) \right. \\
 & \left. + \left(32n - 16(n - n^2) - \frac{(n^2+n)(4(n-n^2)-8(n+1))}{(1-n)(n+1)+n(n+1)} - \frac{(-n^2+n-2(n+2)) \left(8n-4(n-n^2) - \frac{(n^2+n)(-n^2+n-2(n+1))}{(1-n)(n+1)+n(n+1)} \right)}{(1-n)(n+2)+(n+1)(n+2)} \right) x^3 \right) \\
 & \frac{(1-n)(n+4) + (1-n)(n+3) + (n+2)(n+3)}{39}
 \end{aligned}$$

3.13 problem 15

Internal problem ID [4708]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XIV. page 177

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + (x^2 + x) y' + (x - 9) y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 41

```
Order:=6;
dsolve(x^2*diff(y(x),x$2)+(x+x^2)*diff(y(x),x)+(x-9)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^3 \left(1 - \frac{4}{7}x + \frac{5}{28}x^2 - \frac{5}{126}x^3 + \frac{1}{144}x^4 - \frac{1}{990}x^5 + O(x^6) \right) + \frac{c_2(-86400 + 34560x - 4320x^2 + O(x^6))}{x^3}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 60

```
AsymptoticDSolveValue[x^2*y'[x]+(x+x^2)*y'[x]+(x-9)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{x^3} - \frac{2}{5x^2} + \frac{1}{20x} \right) + c_2 \left(\frac{x^7}{144} - \frac{5x^6}{126} + \frac{5x^5}{28} - \frac{4x^4}{7} + x^3 \right)$$

3.14 problem 16

Internal problem ID [4709]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XIV. page 177

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries], [_2nd_order, _linear,`

$$(a^2 + x^2) y'' + y'x - yn^2 = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 72

```
Order:=6;
dsolve((a^2+x^2)*diff(y(x),x$2)+x*diff(y(x),x)-n^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{n^2 x^2}{2a^2} + \frac{n^2(n^2 - 4)x^4}{24a^4}\right) y(0) + \left(x + \frac{(n^2 - 1)x^3}{6a^2} + \frac{(n^4 - 10n^2 + 9)x^5}{120a^4}\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 112

```
AsymptoticDSolveValue[(a^2+x^2)*y'[x]+x*y'[x]-n^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{n^4 x^5}{120a^4} - \frac{n^2 x^5}{12a^4} + \frac{3x^5}{40a^4} + \frac{n^2 x^3}{6a^2} - \frac{x^3}{6a^2} + x \right) + c_1 \left(\frac{n^4 x^4}{24a^4} - \frac{n^2 x^4}{6a^4} + \frac{n^2 x^2}{2a^2} + 1 \right)$$

3.15 problem 18

Internal problem ID [4710]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XIV. page 177

Problem number: 18.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [_Gegenbauer, [_2nd_order, _linear, ‘_with_symmetry_[0,F(x)]’]

$$(-x^2 + 1)y'' - y'x + a^2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 71

```
Order:=6;  
dsolve((1-x^2)*diff(y(x),x$2)-x*diff(y(x),x)+a^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{a^2 x^2}{2} + \frac{a^2(a^2 - 4)x^4}{24}\right) y(0) + \left(x - \frac{(a^2 - 1)x^3}{6} + \frac{(a^4 - 10a^2 + 9)x^5}{120}\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 88

```
AsymptoticDSolveValue[(1-x^2)*y''[x]-x*y'[x]+a^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{a^4 x^5}{120} - \frac{a^2 x^5}{12} - \frac{a^2 x^3}{6} + \frac{3x^5}{40} + \frac{x^3}{6} + x \right) + c_1 \left(\frac{a^4 x^4}{24} - \frac{a^2 x^4}{6} - \frac{a^2 x^2}{2} + 1 \right)$$

4 Chapter VII, Solutions in series. Examples XV.

page 194

4.1	problem 1	44
4.2	problem 2	45
4.3	problem 3	46
4.4	problem 4	47
4.5	problem 5	48
4.6	problem 6	49
4.7	problem 7	51
4.8	problem 8	52
4.9	problem 9	54
4.10	problem 10	55
4.11	problem 11	56
4.12	problem 12	57
4.13	problem 13	58
4.14	problem 14	59

4.1 problem 1

Internal problem ID [4711]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XV. page 194

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$xy'' + y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 59

```
Order:=6;  
dsolve(x*diff(y(x),x$2)+diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 - x + \frac{1}{4}x^2 - \frac{1}{36}x^3 + \frac{1}{576}x^4 - \frac{1}{14400}x^5 + O(x^6) \right) \\ + \left(2x - \frac{3}{4}x^2 + \frac{11}{108}x^3 - \frac{25}{3456}x^4 + \frac{137}{432000}x^5 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 111

```
AsymptoticDSolveValue[x*y''[x]+y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^5}{14400} + \frac{x^4}{576} - \frac{x^3}{36} + \frac{x^2}{4} - x + 1 \right) + c_2 \left(\frac{137x^5}{432000} - \frac{25x^4}{3456} + \frac{11x^3}{108} - \frac{3x^2}{4} \right. \\ \left. + \left(-\frac{x^5}{14400} + \frac{x^4}{576} - \frac{x^3}{36} + \frac{x^2}{4} - x + 1 \right) \log(x) + 2x \right)$$

4.2 problem 2

Internal problem ID [4712]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XV. page 194

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$xy'' + y' + pxy = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 59

```
Order:=6;  
dsolve(x*diff(y(x),x$2)+diff(y(x),x)+p*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 - \frac{1}{4} p x^2 + \frac{1}{64} p^2 x^4 + O(x^6) \right) + \left(\frac{p}{4} x^2 - \frac{3}{128} p^2 x^4 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 72

```
AsymptoticDSolveValue[x*y''[x]+y'[x]+p*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{p^2 x^4}{64} - \frac{p x^2}{4} + 1 \right) + c_2 \left(-\frac{3}{128} p^2 x^4 + \left(\frac{p^2 x^4}{64} - \frac{p x^2}{4} + 1 \right) \log(x) + \frac{p x^2}{4} \right)$$

4.3 problem 3

Internal problem ID [4713]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XV. page 194

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$xy'' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 58

```
Order:=6;  
dsolve(x*diff(y(x),x$2)+y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1 x \left(1 - \frac{1}{2}x + \frac{1}{12}x^2 - \frac{1}{144}x^3 + \frac{1}{2880}x^4 - \frac{1}{86400}x^5 + O(x^6) \right) \\ & + c_2 \left(\ln(x) \left(-x + \frac{1}{2}x^2 - \frac{1}{12}x^3 + \frac{1}{144}x^4 - \frac{1}{2880}x^5 + O(x^6) \right) \right. \\ & \left. + \left(1 - \frac{3}{4}x^2 + \frac{7}{36}x^3 - \frac{35}{1728}x^4 + \frac{101}{86400}x^5 + O(x^6) \right) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 85

```
AsymptoticDSolveValue[x*y''[x]+y[x]==0,y[x],{x,0,5}]
```

$$\begin{aligned} y(x) \rightarrow & c_1 \left(\frac{1}{144}x(x^3 - 12x^2 + 72x - 144) \log(x) \right. \\ & \left. + \frac{-47x^4 + 480x^3 - 2160x^2 + 1728x + 1728}{1728} \right) + c_2 \left(\frac{x^5}{2880} - \frac{x^4}{144} + \frac{x^3}{12} - \frac{x^2}{2} + x \right) \end{aligned}$$

4.4 problem 4

Internal problem ID [4714]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XV. page 194

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^3 y'' - (2x - 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

 Solution by Maple

```
Order:=6;  
dsolve(x^3*diff(y(x),x$2)-(2*x-1)*y(x)=0,y(x),type='series',x=0);
```

No solution found

 Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 222

```
AsymptoticDSolveValue[x^3*y'[x]-(2*x-1)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 e^{-\frac{2i}{\sqrt{x}} x^{3/4}} \left(-\frac{1159525191825ix^{9/2}}{8796093022208} + \frac{218243025ix^{7/2}}{4294967296} - \frac{405405ix^{5/2}}{8388608} + \frac{3465ix^{3/2}}{8192} \right. \\ \left. + \frac{75369137468625x^5}{281474976710656} - \frac{41247931725x^4}{549755813888} + \frac{11486475x^3}{268435456} - \frac{45045x^2}{524288} - \frac{945x}{512} - \frac{35i\sqrt{x}}{16} \right. \\ \left. + 1 \right) + c_2 e^{\frac{2i}{\sqrt{x}} x^{3/4}} \left(\frac{1159525191825ix^{9/2}}{8796093022208} - \frac{218243025ix^{7/2}}{4294967296} + \frac{405405ix^{5/2}}{8388608} - \frac{3465ix^{3/2}}{8192} + \frac{75369137468625}{281474976710656} \right)$$

4.5 problem 5

Internal problem ID [4715]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XV. page 194

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + x(1+x)y' + (3x-1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 65

```
Order:=6;
dsolve(x^2*diff(y(x),x$2)+x*(x+1)*diff(y(x),x)+(3*x-1)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 x^2 \left(1 - \frac{4}{3}x + \frac{5}{6}x^2 - \frac{1}{3}x^3 + \frac{7}{72}x^4 - \frac{1}{45}x^5 + O(x^6)\right) + c_2 (\ln(x) (6x^2 - 8x^3 + 5x^4 - 2x^5 + O(x^6)) + (-2x^2 + 2x - 1) \ln(x))}{x}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 85

```
AsymptoticDSolveValue[x^2*y''[x]+x*(x+1)*y'[x]+(3*x-1)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{13x^4 - 12x^3 - 4x^2 + 8x + 4}{4x} - \frac{1}{2}x(5x^2 - 8x + 6) \log(x) \right) + c_2 \left(\frac{7x^5}{72} - \frac{x^4}{3} + \frac{5x^3}{6} - \frac{4x^2}{3} + x \right)$$

4.6 problem 6

Internal problem ID [4716]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XV. page 194

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(-x^2 + x)y'' - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 58

```
Order:=6;  
dsolve((x-x^2)*diff(y(x),x$2)-y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1 x \left(1 + \frac{1}{2}x + \frac{1}{4}x^2 + \frac{7}{48}x^3 + \frac{91}{960}x^4 + \frac{637}{9600}x^5 + O(x^6) \right) \\ & + c_2 \left(\ln(x) \left(x + \frac{1}{2}x^2 + \frac{1}{4}x^3 + \frac{7}{48}x^4 + \frac{91}{960}x^5 + O(x^6) \right) \right. \\ & \left. + \left(1 - \frac{1}{4}x^2 - \frac{1}{12}x^3 - \frac{17}{576}x^4 - \frac{311}{28800}x^5 + O(x^6) \right) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.044 (sec). Leaf size: 87

```
AsymptoticDSolveValue[(x-x^2)*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{48} x (7x^3 + 12x^2 + 24x + 48) \log(x) + \frac{1}{576} (-185x^4 - 336x^3 - 720x^2 - 1152x + 576) \right) + c_2 \left(\frac{91x^5}{960} + \frac{7x^4}{48} + \frac{x^3}{4} + \frac{x^2}{2} + x \right)$$

4.7 problem 7

Internal problem ID [4717]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XV. page 194

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_elliptic, _class_I]]`

$$x(-x^2 + 1)y'' + (-3x^2 + 1)y' - yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 41

```
Order:=6;  
dsolve(x*(1-x^2)*diff(y(x),x$2)+(1-3*x^2)*diff(y(x),x)-x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 + \frac{1}{4}x^2 + \frac{9}{64}x^4 + O(x^6) \right) + \left(\frac{1}{4}x^2 + \frac{21}{128}x^4 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 60

```
AsymptoticDSolveValue[x*(1-x^2)*y'[x]+(1-3*x^2)*y'[x]-x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{9x^4}{64} + \frac{x^2}{4} + 1 \right) + c_2 \left(\frac{21x^4}{128} + \frac{x^2}{4} + \left(\frac{9x^4}{64} + \frac{x^2}{4} + 1 \right) \log(x) \right)$$

4.8 problem 8

Internal problem ID [4718]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XV. page 194

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [[_Emden, _Fowler]]

$$y'' + \frac{ay}{x^{\frac{3}{2}}} = 0$$

With the expansion point for the power series method at $x = 0$.

X Solution by Maple

```
Order:=6;  
dsolve(diff(y(x),x$2)+a/x^(3/2)*y(x)=0,y(x),type='series',x=0);
```

No solution found

✓ Solution by Mathematica

Time used: 0.244 (sec). Leaf size: 576

AsymptoticDSolveValue[y''[x]+a/x^(3/2)*y[x]==0,y[x],{x,0,5}]

$$\begin{aligned}
 y(x) \rightarrow & \\
 & - \frac{16x^5(126a^{10}c_2 \log(x) - 252\pi a^{10}c_1 + 504\gamma a^{10}c_2 - 1423a^{10}c_2 + 252a^{10}c_2 \log(a) + 504a^{10}c_2 \log(2))}{281302875\pi} \\
 & + \frac{32x^{9/2}(1260a^9c_2 \log(x) - 2520\pi a^9c_1 + 5040\gamma a^9c_2 - 13663a^9c_2 + 2520a^9c_2 \log(a) + 5040a^9c_2 \log(2))}{281302875\pi} \\
 & - \frac{8x^4(140a^8c_2 \log(x) - 280\pi a^8c_1 + 560\gamma a^8c_2 - 1447a^8c_2 + 280a^8c_2 \log(a) + 560a^8c_2 \log(2))}{496125\pi} \\
 & + \frac{128x^{7/2}(105a^7c_2 \log(x) - 210\pi a^7c_1 + 420\gamma a^7c_2 - 1024a^7c_2 + 210a^7c_2 \log(a) + 420a^7c_2 \log(2))}{496125\pi} \\
 & - \frac{32x^3(15a^6c_2 \log(x) - 30\pi a^6c_1 + 60\gamma a^6c_2 - 136a^6c_2 + 30a^6c_2 \log(a) + 60a^6c_2 \log(2))}{2025\pi} \\
 & + \frac{32x^{5/2}(30a^5c_2 \log(x) - 60\pi a^5c_1 + 120\gamma a^5c_2 - 247a^5c_2 + 60a^5c_2 \log(a) + 120a^5c_2 \log(2))}{675\pi} \\
 & - \frac{8x^2(6a^4c_2 \log(x) - 12\pi a^4c_1 + 24\gamma a^4c_2 - 43a^4c_2 + 12a^4c_2 \log(a) + 24a^4c_2 \log(2))}{9\pi} \\
 & + \frac{32x^{3/2}(3a^3c_2 \log(x) - 6\pi a^3c_1 + 12\gamma a^3c_2 - 17a^3c_2 + 6a^3c_2 \log(a) + 12a^3c_2 \log(2))}{9\pi} \\
 & - \frac{8x(a^2c_2 \log(x) - 2\pi a^2c_1 + 4\gamma a^2c_2 - 3a^2c_2 + 2a^2c_2 \log(a) + 4a^2c_2 \log(2))}{\pi} \\
 & + \frac{8ac_2\sqrt{x}}{\pi} + \frac{2c_2}{\pi}
 \end{aligned}$$

4.9 problem 9

Internal problem ID [4719]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XV. page 194

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - (x^2 + 4x) y' + 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 61

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)-(x^2+4*x)*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = x \left(c_1 x^3 \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + O(x^6) \right) \right. \\ \left. + c_2 (\ln(x) (6x^3 + 6x^4 + 3x^5 + O(x^6)) \right. \\ \left. + (12 - 6x + 6x^2 + 11x^3 + 5x^4 + x^5 + O(x^6))) \right)$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 74

```
AsymptoticDSolveValue[x^2*y'[x]-(x^2+4*x)*y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{2}(x+1)x^4 \log(x) + \frac{1}{4}(x^4 + 3x^3 + 2x^2 - 2x + 4)x \right) + c_2 \left(\frac{x^8}{24} + \frac{x^7}{6} + \frac{x^6}{2} + x^5 + x^4 \right)$$

4.10 problem 10

Internal problem ID [4720]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XV. page 194

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_elliptic, _class_II]]`

$$x(-x^2 + 1)y'' + (-x^2 + 1)y' + yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 41

```
Order:=6;  
dsolve(x*(1-x^2)*diff(y(x),x$2)+(1-x^2)*diff(y(x),x)+x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 - \frac{1}{4}x^2 - \frac{3}{64}x^4 + O(x^6)\right) + \left(\frac{1}{4}x^2 + \frac{1}{128}x^4 + O(x^6)\right) c_2$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 60

```
AsymptoticDSolveValue[x*(1-x^2)*y''[x]+(1-x^2)*y'[x]+x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{3x^4}{64} - \frac{x^2}{4} + 1\right) + c_2 \left(\frac{x^4}{128} + \frac{x^2}{4} + \left(-\frac{3x^4}{64} - \frac{x^2}{4} + 1\right) \log(x)\right)$$

4.11 problem 11

Internal problem ID [4721]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XV. page 194

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Jacobi]

$$4x(1-x)y'' - 4y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 60

```
Order:=6;
```

```
dsolve(4*x*(1-x)*diff(y(x),x$2)-4*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^2 \left(1 + \frac{3}{4}x + \frac{75}{128}x^2 + \frac{245}{512}x^3 + \frac{6615}{16384}x^4 + \frac{22869}{65536}x^5 + O(x^6) \right) \\ + c_2 \left(\ln(x) \left(\frac{1}{16}x^2 + \frac{3}{64}x^3 + \frac{75}{2048}x^4 + \frac{245}{8192}x^5 + O(x^6) \right) \right. \\ \left. + \left(-2 + \frac{1}{2}x + \frac{1}{2}x^2 + \frac{3}{8}x^3 + \frac{2415}{8192}x^4 + \frac{23779}{98304}x^5 + O(x^6) \right) \right)$$

✓ Solution by Mathematica

Time used: 0.05 (sec). Leaf size: 86

```
AsymptoticDSolveValue[4*x*(1-x)*y'[x]-4*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{135x^4 + 192x^3 + 256x^2 - 4096x + 16384}{16384} - \frac{x^2(75x^2 + 96x + 128) \log(x)}{4096} \right) \\ + c_2 \left(\frac{6615x^6}{16384} + \frac{245x^5}{512} + \frac{75x^4}{128} + \frac{3x^3}{4} + x^2 \right)$$

4.12 problem 12

Internal problem ID [4722]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XV. page 194

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x^3 y'' + y = x^{\frac{3}{2}}$$

With the expansion point for the power series method at $x = 0$.

✗ Solution by Maple

```
Order:=6;
dsolve(x^3*diff(y(x),x$2)+y(x)=x^(3/2),y(x),type='series',x=0);
```

No solution found

✓ Solution by Mathematica

Time used: 0.251 (sec). Leaf size: 688

```
AsymptoticDSolveValue[x^3*y'[x]+y[x]==x^(3/2),y[x],{x,0,5}]
```

$y(x)$

$$\begin{aligned}
 & e^{\frac{2i}{\sqrt{x}}} x^{3/4} \left(\frac{468131288625ix^{9/2}}{8796093022208} - \frac{66891825ix^{7/2}}{4294967296} + \frac{72765ix^{5/2}}{8388608} - \frac{105ix^{3/2}}{8192} + \frac{33424574007825x^5}{281474976710656} - \frac{14783093325x^4}{549755813888} + \frac{2837835x^3}{268435456} \right) \\
 & + e^{-\frac{2i}{\sqrt{x}}} x^{3/4} \left(-\frac{468131288625ix^{9/2}}{8796093022208} + \frac{66891825ix^{7/2}}{4294967296} - \frac{72765ix^{5/2}}{8388608} + \frac{105ix^{3/2}}{8192} + \frac{33424574007825x^5}{281474976710656} - \frac{14783093325x^4}{549755813888} + \frac{2837835x^3}{268435456} \right) \\
 & + c_1 e^{-\frac{2i}{\sqrt{x}}} x^{3/4} \left(-\frac{468131288625ix^{9/2}}{8796093022208} + \frac{66891825ix^{7/2}}{4294967296} - \frac{72765ix^{5/2}}{8388608} + \frac{105ix^{3/2}}{8192} + \frac{33424574007825x^5}{281474976710656} - \frac{14783093325x^4}{549755813888} + \frac{2837835x^3}{268435456} \right)
 \end{aligned}$$

4.13 problem 13

Internal problem ID [4723]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XV. page 194

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$2x^2y'' - (3x + 2)y' + \frac{(2x - 1)y}{x} = \sqrt{x}$$

With the expansion point for the power series method at $x = 0$.

X Solution by Maple

Order:=6;

`dsolve(2*x^2*diff(y(x),x$2)-(3*x+2)*diff(y(x),x)+(2*x-1)/x*y(x)=x^(1/2),y(x),type='series',x`

No solution found

✓ Solution by Mathematica

Time used: 0.158 (sec). Leaf size: 222

`AsymptoticDSolveValue[2*x^2*y''[x]-(3*x+2)*y'[x]+(2*x-1)/x*y[x]==x^(1/2),y[x],{x,0,5}]`

$$y(x) \rightarrow \frac{1}{256}e^{-1/x} \left(-\frac{405405x^5}{16} + \frac{45045x^4}{16} - \frac{693x^3}{2} + \frac{189x^2}{4} - 7x + 1 \right) x^4 \left(\frac{2e^{\frac{1}{x}}(15663375x^7 + 20072325x^6 + 10329540x^5 + 4131816x^4 + 2754544x^3 + 5509088x^2 - 64x - 11018112\sqrt{\pi}\operatorname{erfi}\left(\frac{1}{\sqrt{x}}\right))}{x^{3/2}} \right) + c_2 e^{-1/x} \left(-\frac{405405x^5}{16} + \frac{45045x^4}{16} - \frac{693x^3}{2} + \frac{189x^2}{4} - 7x + 1 \right) x^4 + \frac{\left(\frac{5x}{2} + 1\right) \left(-\frac{15015x^6}{64} + \frac{693x^5}{20} - \frac{189x^4}{32} + 7 \right)}{\sqrt{x}}$$

4.14 problem 14

Internal problem ID [4724]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XV. page 194

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$(-x^2 + x)y'' + 3y' + 2y = 3x^2$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 50

```
Order:=6;  
dsolve((x-x^2)*diff(y(x),x$2)+3*diff(y(x),x)+2*y(x)=3*x^2,y(x),type='series',x=0);
```

$$y(x) = c_1 \left(1 - \frac{2}{3}x + \frac{1}{6}x^2 + O(x^6) \right) + \frac{c_2(-2 + 8x - 12x^2 + 8x^3 - 2x^4 + O(x^6))}{x^2} \\ + x^3 \left(\frac{1}{5} + \frac{1}{30}x + \frac{1}{105}x^2 + O(x^3) \right)$$

✓ Solution by Mathematica

Time used: 0.049 (sec). Leaf size: 91

```
AsymptoticDSolveValue[(x-x^2)*y''[x]+3*y'[x]+2*y[x]==3*x^2,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^2}{6} - \frac{2x}{3} + 1 \right) + \frac{c_2(1 - 4x)}{x^2} + \frac{(1 - 4x) \left(-\frac{5x^6}{6} - \frac{3x^5}{10} \right)}{x^2} \\ + \left(\frac{x^2}{6} - \frac{2x}{3} + 1 \right) \left(-5x^6 - \frac{9x^5}{5} + \frac{x^3}{2} \right)$$

5 Chapter VII, Solutions in series. Examples XVI.
page 220

5.1	problem 5	61
5.2	problem 6	62
5.3	problem 8	63
5.4	problem 9	64
5.5	problem 10	65
5.6	problem 11	66

5.1 problem 5

Internal problem ID [4725]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XVI. page 220

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Jacobi]

$$x(1-x)y'' + \left(\frac{3}{2} - 2x\right)y' - \frac{y}{4} = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 34

```
Order:=6;
```

```
dsolve(x*(1-x)*diff(y(x),x$2)+(3/2-2*x)*diff(y(x),x)-1/4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_2 \left(1 + \frac{1}{6}x + \frac{3}{40}x^2 + \frac{5}{112}x^3 + \frac{35}{1152}x^4 + \frac{63}{2816}x^5 + O(x^6)\right) \sqrt{x} + c_1(1 + O(x^6))}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 50

```
AsymptoticDSolveValue[x*(1-x)*y''[x]+(3/2-2*x)*y'[x]-1/4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{63x^5}{2816} + \frac{35x^4}{1152} + \frac{5x^3}{112} + \frac{3x^2}{40} + \frac{x}{6} + 1 \right) + \frac{c_2}{\sqrt{x}}$$

5.2 problem 6

Internal problem ID [4726]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XVI. page 220

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x(1-x)y'' + y'x - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 42

```
Order:=6;  
dsolve(2*x*(1-x)*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \ln(x) \left(\frac{1}{2}x + O(x^6) \right) c_2 + c_1(1 + O(x^6)) x \\ + \left(1 - \frac{1}{2}x + \frac{1}{8}x^2 + \frac{1}{32}x^3 + \frac{5}{384}x^4 + \frac{7}{1024}x^5 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.051 (sec). Leaf size: 43

```
AsymptoticDSolveValue[2*x*(1-x)*y'[x]+x*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{384}(5x^4 + 12x^3 + 48x^2 - 768x + 384) + \frac{1}{2}x \log(x) \right) + c_2x$$

5.3 problem 8

Internal problem ID [4727]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XVI. page 220

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Jacobi]

$$2x(1-x)y'' + (1-11x)y' - 10y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

```
Order:=6;  
dsolve(2*x*(1-x)*diff(y(x),x$2)+(1-11*x)*diff(y(x),x)-10*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1\sqrt{x}(1 + 5x + 14x^2 + 30x^3 + 55x^4 + 91x^5 + O(x^6)) \\ + c_2(1 + 10x + 35x^2 + 84x^3 + 165x^4 + 286x^5 + O(x^6))$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 65

```
AsymptoticDSolveValue[2*x*(1-x)*y'[x]+(1-11*x)*y'[x]-10*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1\sqrt{x}(91x^5 + 55x^4 + 30x^3 + 14x^2 + 5x + 1) \\ + c_2(286x^5 + 165x^4 + 84x^3 + 35x^2 + 10x + 1)$$

5.4 problem 9

Internal problem ID [4728]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XVI. page 220

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Jacobi]

$$x(1-x)y'' + \frac{(-2x+1)y'}{3} + \frac{20y}{9} = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 36

```
Order:=6;
```

```
dsolve(x*(1-x)*diff(y(x),x$2)+1/3*(1-2*x)*diff(y(x),x)+20/9*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{\frac{2}{3}} \left(1 - \frac{6}{5}x + O(x^6) \right) + c_2 \left(1 - \frac{20}{3}x + \frac{35}{9}x^2 + \frac{50}{81}x^3 + \frac{65}{243}x^4 + \frac{112}{729}x^5 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 57

```
AsymptoticDSolveValue[x*(1-x)*y''[x]+1/3*(1-2*x)*y'[x]+20/9*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(1 - \frac{6x}{5} \right) x^{2/3} + c_2 \left(\frac{112x^5}{729} + \frac{65x^4}{243} + \frac{50x^3}{81} + \frac{35x^2}{9} - \frac{20x}{3} + 1 \right)$$

5.5 problem 10

Internal problem ID [4729]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XVI. page 220

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$2x(1-x)y'' + y' + 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 38

```
Order:=6;  
dsolve(2*x*(1-x)*diff(y(x),x$2)+diff(y(x),x)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1\sqrt{x} \left(1 - \frac{3}{2}x + \frac{3}{8}x^2 + \frac{1}{16}x^3 + \frac{3}{128}x^4 + \frac{3}{256}x^5 + O(x^6) \right) \\ + c_2 \left(1 - 4x + \frac{8}{3}x^2 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 62

```
AsymptoticDSolveValue[2*x*(1-x)*y'[x]+y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{8x^2}{3} - 4x + 1 \right) + c_1\sqrt{x} \left(\frac{3x^5}{256} + \frac{3x^4}{128} + \frac{x^3}{16} + \frac{3x^2}{8} - \frac{3x}{2} + 1 \right)$$

5.6 problem 11

Internal problem ID [4730]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter VII, Solutions in series. Examples XVI. page 220

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4y'' + \frac{3(-x^2 + 2)y}{(-x^2 + 1)^2} = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(4*diff(y(x),x$2)+3*(2-x^2)/(1-x^2)^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{3}{4}x^2 - \frac{3}{32}x^4\right) y(0) + \left(x - \frac{1}{4}x^3 - \frac{3}{32}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[4*y''[x]+3*(2-x^2)/(1-x^2)^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(-\frac{3x^5}{32} - \frac{x^3}{4} + x \right) + c_1 \left(-\frac{3x^4}{32} - \frac{3x^2}{4} + 1 \right)$$

6 Chapter IX, Special forms of differential equations. Examples XVII. page 247

6.1	problem 1	68
6.2	problem 2	69
6.3	problem 3	70
6.4	problem 4	71
6.5	problem 5	72
6.6	problem 6	73
6.7	problem 7	74
6.8	problem 8	75
6.9	problem 9	76
6.10	problem 10	77
6.11	problem 11	78
6.12	problem 12	79
6.13	problem 15	80
6.14	problem 18	81
6.15	problem 19	82
6.16	problem 20	83
6.17	problem 21	84

6.1 problem 1

Internal problem ID [4731]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, _Riccati]`

$$y' + y^2 = \frac{a^2}{x^4}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 38

```
dsolve(diff(y(x),x)+y(x)^2=a^2/x^4,y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{-a^2} \tan\left(\frac{\sqrt{-a^2}(c_1x-1)}{x}\right) - x}{x^2}$$

✓ Solution by Mathematica

Time used: 0.384 (sec). Leaf size: 71

```
DSolve[y'[x]+y[x]^2==a^2/x^4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-2a^2c_1e^{\frac{2a}{x}} + 2ac_1xe^{\frac{2a}{x}} + a + x}{x^2 \left(1 + 2ac_1e^{\frac{2a}{x}}\right)}$$

$$y(x) \rightarrow \frac{x - a}{x^2}$$

6.2 problem 2

Internal problem ID [4732]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$u'' - \frac{a^2 u}{x^{\frac{2}{3}}} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 43

```
dsolve(diff(u(x), x$2) - a^2*x^(-2/3)*u(x)=0, u(x), singsol=all)
```

$$u(x) = c_1 \sqrt{x} \operatorname{BesselJ}\left(\frac{3}{4}, \frac{3\sqrt{-a^2} x^{\frac{2}{3}}}{2}\right) + c_2 \sqrt{x} \operatorname{BesselY}\left(\frac{3}{4}, \frac{3\sqrt{-a^2} x^{\frac{2}{3}}}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.05 (sec). Leaf size: 79

```
DSolve[u''[x] - a^2*x^(-2/3)*u[x]==0, u[x], x, IncludeSingularSolutions -> True]
```

$$u(x) \rightarrow \frac{3^{3/4} a^{3/4} \sqrt{x} \left(16c_1 \operatorname{Gamma}\left(\frac{5}{4}\right) \operatorname{BesselI}\left(-\frac{3}{4}, \frac{3}{2} a x^{2/3}\right) + 3(-1)^{3/4} c_2 \operatorname{Gamma}\left(\frac{3}{4}\right) \operatorname{BesselI}\left(\frac{3}{4}, \frac{3}{2} a x^{2/3}\right) \right)}{8\sqrt{2}}$$

6.3 problem 3

Internal problem ID [4733]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$u'' - \frac{2u'}{x} - a^2u = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(diff(u(x),x$2)-2/x*diff(u(x),x)-a^2*u(x)=0,u(x), singsol=all)
```

$$u(x) = c_1 e^{ax}(ax - 1) + c_2 e^{-ax}(ax + 1)$$

✓ Solution by Mathematica

Time used: 0.13 (sec). Leaf size: 68

```
DSolve[u''[x]-2/x*u'[x]-a^2*u[x]==0,u[x],x,IncludeSingularSolutions -> True]
```

$$u(x) \rightarrow \frac{\sqrt{\frac{2}{\pi}}\sqrt{x}((iac_2x + c_1) \sinh(ax) - (ac_1x + ic_2) \cosh(ax))}{a\sqrt{-iax}}$$

6.4 problem 4

Internal problem ID [4734]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$u'' + \frac{2u'}{x} - a^2u = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(diff(u(x),x$2)+2/x*diff(u(x),x)-a^2*u(x)=0,u(x), singsol=all)
```

$$u(x) = \frac{c_1 \sinh(ax)}{x} + \frac{c_2 \cosh(ax)}{x}$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 35

```
DSolve[u''[x]+2/x*u'[x]-a^2*u[x]==0,u[x],x,IncludeSingularSolutions -> True]
```

$$u(x) \rightarrow \frac{2ac_1e^{-ax} + c_2e^{ax}}{2ax}$$

6.5 problem 5

Internal problem ID [4735]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$u'' + \frac{2u'}{x} + a^2u = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(u(x),x$2)+2/x*diff(u(x),x)+a^2*u(x)=0,u(x), singsol=all)
```

$$u(x) = \frac{c_1 \sin(ax)}{x} + \frac{c_2 \cos(ax)}{x}$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 42

```
DSolve[u''[x]+2/x*u'[x]+a^2*u[x]==0,u[x],x,IncludeSingularSolutions -> True]
```

$$u(x) \rightarrow \frac{e^{-iax} \left(2c_1 - \frac{ic_2 e^{2iax}}{a} \right)}{2x}$$

6.6 problem 6

Internal problem ID [4736]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$u'' + \frac{4u'}{x} - a^2u = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve(diff(u(x),x$2)+4/x*diff(u(x),x)-a^2*u(x)=0,u(x), singsol=all)
```

$$u(x) = \frac{c_1 e^{ax}(ax - 1)}{x^3} + \frac{c_2 e^{-ax}(ax + 1)}{x^3}$$

✓ Solution by Mathematica

Time used: 0.091 (sec). Leaf size: 68

```
DSolve[u''[x]+4/x*u'[x]-a^2*u[x]==0,u[x],x,IncludeSingularSolutions -> True]
```

$$u(x) \rightarrow \frac{\sqrt{\frac{2}{\pi}}((iac_2x + c_1) \sinh(ax) - (ac_1x + ic_2) \cosh(ax))}{ax^{5/2}\sqrt{-iax}}$$

6.7 problem 7

Internal problem ID [4737]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$u'' + \frac{4u'}{x} + a^2u = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 41

```
dsolve(diff(u(x),x$2)+4/x*diff(u(x),x)+a^2*u(x)=0,u(x), singsol=all)
```

$$u(x) = \frac{c_1(\cos(ax)ax - \sin(ax))}{x^3} + \frac{c_2(\cos(ax) + \sin(ax)ax)}{x^3}$$

✓ Solution by Mathematica

Time used: 0.094 (sec). Leaf size: 57

```
DSolve[u''[x]+4/x*u'[x]+a^2*u[x]==0,u[x],x,IncludeSingularSolutions -> True]
```

$$u(x) \rightarrow -\frac{\sqrt{\frac{2}{\pi}}((ac_1x + c_2)\cos(ax) + (ac_2x - c_1)\sin(ax))}{x^{3/2}(ax)^{3/2}}$$

6.8 problem 8

Internal problem ID [4738]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - a^2y - \frac{6y}{x^2} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 50

```
dsolve(diff(y(x),x$2)-a^2*y(x)=6*y(x)/x^2,y(x), singsol=all)
```

$$y(x) = \frac{c_1 e^{ax}(a^2 x^2 - 3ax + 3)}{x^2} + \frac{c_2 e^{-ax}(a^2 x^2 + 3ax + 3)}{x^2}$$

✓ Solution by Mathematica

Time used: 0.138 (sec). Leaf size: 90

```
DSolve[y''[x]-a^2*y[x]==6*y[x]/x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt{\frac{2}{\pi}}((a^2 c_2 x^2 - 3iac_1 x + 3c_2) \cosh(ax) + i(c_1(a^2 x^2 + 3) + 3iac_2 x) \sinh(ax))}{a^2 x^{3/2} \sqrt{-iax}}$$

6.9 problem 9

Internal problem ID [4739]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + yn^2 - \frac{6y}{x^2} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 61

```
dsolve(diff(y(x),x$2)+n^2*y(x)=6*y(x)/x^2,y(x), singsol=all)
```

$$y(x) = \frac{c_1((n^2x^2 - 3) \cos(nx) - 3 \sin(nx) nx)}{x^2} + \frac{c_2(3 \cos(nx) nx + (n^2x^2 - 3) \sin(nx))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.143 (sec). Leaf size: 79

```
DSolve[y''[x]+n^2*y[x]==6*y[x]/x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{\frac{2}{\pi}}\sqrt{x}((c_2(-n^2)x^2 + 3c_1nx + 3c_2) \cos(nx) + (c_1(n^2x^2 - 3) + 3c_2nx) \sin(nx))}{(nx)^{5/2}}$$

6.10 problem 10

Internal problem ID [4740]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y' x - \left(x^2 + \frac{1}{4}\right) y = 0$$

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 19

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-(x^2+1/4)*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 \sinh(x)}{\sqrt{x}} + \frac{c_2 \cosh(x)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 32

```
DSolve[x^2*y'[x]+x*y'[x]-(x^2+1/4)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{-x}(c_2 e^{2x} + 2c_1)}{2\sqrt{x}}$$

6.11 problem 11

Internal problem ID [4741]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y' x + \frac{(-9a^2 + 4x^2)y}{4a^2} = 0$$

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 45

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(4*x^2-9*a^2)/(4*a^2)*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 e^{\frac{ix}{a}} (ix - a)}{x^{\frac{3}{2}}} + \frac{c_2 e^{-\frac{ix}{a}} (ix + a)}{x^{\frac{3}{2}}}$$

✓ Solution by Mathematica

Time used: 0.101 (sec). Leaf size: 62

```
DSolve[x^2*y''[x]+x*y'[x]+(4*x^2-9*a^2)/(4*a^2)*y[x]==0,y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow -\frac{\sqrt{\frac{2}{\pi}} \left((ac_2 + c_1 x) \cos\left(\frac{x}{a}\right) + (c_2 x - ac_1) \sin\left(\frac{x}{a}\right) \right)}{x \sqrt{\frac{x}{a}}}$$

6.12 problem 12

Internal problem ID [4742]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y' x + \left(x^2 - \frac{25}{4}\right) y = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 45

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-25/4)*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 e^{ix} (x^2 + 3ix - 3)}{x^{\frac{5}{2}}} + \frac{c_2 e^{-ix} (-x^2 + 3ix + 3)}{x^{\frac{5}{2}}}$$

✓ Solution by Mathematica

Time used: 0.111 (sec). Leaf size: 59

```
DSolve[x^2*y''[x]+x*y'[x]+(x^2-25/4)*y[x]==0,y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow -\frac{\sqrt{\frac{2}{\pi}}((-c_2 x^2 + 3c_1 x + 3c_2) \cos(x) + (c_1(x^2 - 3) + 3c_2 x) \sin(x))}{x^{5/2}}$$

6.13 problem 15

Internal problem ID [4743]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + qy' - \frac{2y}{x^2} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$2)+q*diff(y(x),x)=2*y(x)/x^2,y(x), singsol=all)
```

$$y(x) = \frac{c_1(qx - 2)}{x} + \frac{c_2 e^{-qx}(qx + 2)}{x}$$

✓ Solution by Mathematica

Time used: 0.064 (sec). Leaf size: 80

```
DSolve[y''[x]+q*y'[x]==2*y[x]/x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{qx^{3/2}e^{-\frac{qx}{2}}(2(ic_2qx + 2c_1)\sinh\left(\frac{qx}{2}\right) - 2(c_1qx + 2ic_2)\cosh\left(\frac{qx}{2}\right))}{\sqrt{\pi}(-iqx)^{5/2}}$$

6.14 problem 18

Internal problem ID [4744]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 18.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + e^{2x}y - yn^2 = 0$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+exp(2*x)*y(x)=n^2*y(x),y(x), singsol=all)
```

$$y(x) = c_1 \text{BesselJ}(n, e^x) + c_2 \text{BesselY}(n, e^x)$$

✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 46

```
DSolve[y''[x]+Exp[2*x]*y[x]==n^2*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \text{Gamma}(1 - n) \text{BesselJ}\left(-n, \sqrt{e^{2x}}\right) + c_2 \text{Gamma}(n + 1) \text{BesselJ}\left(n, \sqrt{e^{2x}}\right)$$

6.15 problem 19

Internal problem ID [4745]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y'' + \frac{y}{4x} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)+y(x)/(4*x)=0,y(x), singsol=all)
```

$$y(x) = c_1\sqrt{x} \text{ BesselJ}(1, \sqrt{x}) + c_2\sqrt{x} \text{ BesselY}(1, \sqrt{x})$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 38

```
DSolve[y''[x]+y[x]/(4*x)==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}\sqrt{x}(c_1 \text{ BesselJ}(1, \sqrt{x}) + 2ic_2 \text{ BesselY}(1, \sqrt{x}))$$

6.16 problem 20

Internal problem ID [4746]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y''x + y' + y = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 23

```
dsolve(x*diff(y(x),x$2)+diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \text{BesselJ}(0, 2\sqrt{x}) + c_2 \text{BesselY}(0, 2\sqrt{x})$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 31

```
DSolve[x*y''[x]+y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \text{BesselJ}(0, 2\sqrt{x}) + 2c_2 \text{BesselY}(0, 2\sqrt{x})$$

6.17 problem 21

Internal problem ID [4747]

Book: A treatise on ordinary and partial differential equations by William Woolsey Johnson. 1913

Section: Chapter IX, Special forms of differential equations. Examples XVII. page 247

Problem number: 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y''x + 3y' + 4yx^3 = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(x*diff(y(x),x$2)+3*diff(y(x),x)+4*x^3*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 \sin(x^2)}{x^2} + \frac{c_2 \cos(x^2)}{x^2}$$

✓ Solution by Mathematica

Time used: 0.08 (sec). Leaf size: 41

```
DSolve[x*y''[x]+3*y'[x]+4*x^3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{4c_1 e^{-ix^2} - ic_2 e^{ix^2}}{4x^2}$$