


A Solution Manual For

**An introduction to Ordinary
Differential Equations. Earl A.
Coddington. Dover. NY 1961**



Nasser M. Abbasi

March 3, 2024

Contents

1	Chapter 1.3 Introduction– Linear equations of First Order. Page 38	3
2	Chapter 1.6 Introduction– Linear equations of First Order. Page 41	15
3	Chapter 1. Introduction– Linear equations of First Order. Page 45	26
4	Chapter 2. Linear equations with constant coefficients. Page 52	38
5	Chapter 2. Linear equations with constant coefficients. Page 59	52
6	Chapter 2. Linear equations with constant coefficients. Page 69	57
7	Chapter 2. Linear equations with constant coefficients. Page 74	69
8	Chapter 2. Linear equations with constant coefficients. Page 79	78
9	Chapter 2. Linear equations with constant coefficients. Page 83	81
10	Chapter 2. Linear equations with constant coefficients. Page 89	92
11	Chapter 2. Linear equations with constant coefficients. Page 93	99
12	Chapter 3. Linear equations with variable coefficients. Page 108	109
13	Chapter 3. Linear equations with variable coefficients. Page 121	113
14	Chapter 3. Linear equations with variable coefficients. Page 124	121
15	Chapter 3. Linear equations with variable coefficients. Page 130	125
16	Chapter 4. Linear equations with Regular Singular Points. Page 149	138
17	Chapter 4. Linear equations with Regular Singular Points. Page 154	148
18	Chapter 4. Linear equations with Regular Singular Points. Page 159	164
19	Chapter 4. Linear equations with Regular Singular Points. Page 166	169
20	Chapter 4. Linear equations with Regular Singular Points. Page 182	181
21	Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190	183

22	Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198	202
23	Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238	219
24	Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 250	231
25	Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 254	237

1 Chapter 1.3 Introduction– Linear equations of First Order. Page 38

1.1	problem 1 (a)	4
1.2	problem 1 (b)	5
1.3	problem 1 (d)	6
1.4	problem 2 (a)	7
1.5	problem 2 (b)	8
1.6	problem 2 (c)	9
1.7	problem 2 (f)	10
1.8	problem 2 (h)	11
1.9	problem 3(a)	12
1.10	problem 4(a)	13
1.11	problem 5(a)	14

1.1 problem 1 (a)

Internal problem ID [5912]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 1 (a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' = e^{3x} + \sin(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)=exp(3*x)+sin(x),y(x), singsol=all)
```

$$y(x) = \frac{e^{3x}}{3} - \cos(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 21

```
DSolve[y'[x]==Exp[3*x]+Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{3x}}{3} - \cos(x) + c_1$$

1.2 problem 1 (b)

Internal problem ID [5913]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 1 (b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$y'' = x + 2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)=2+x,y(x), singsol=all)
```

$$y(x) = \frac{1}{6}x^3 + x^2 + c_1x + c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 22

```
DSolve[y''[x]==2+x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{6} + x^2 + c_2x + c_1$$

1.3 problem 1 (d)

Internal problem ID [5914]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 1 (d).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _quadrature]]`

$$y''' = x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$3)=x^2,y(x), singsol=all)
```

$$y(x) = \frac{1}{60}x^5 + \frac{1}{2}c_1x^2 + xc_2 + c_3$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 25

```
DSolve[y'''[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^5}{60} + c_3x^2 + c_2x + c_1$$

1.4 problem 2 (a)

Internal problem ID [5915]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 2 (a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' + \cos(x)y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve(diff(y(x),x)+cos(x)*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-\sin(x)}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 19

```
DSolve[y'[x]+Cos[x]*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-\sin(x)}$$

$$y(x) \rightarrow 0$$

1.5 problem 2 (b)

Internal problem ID [5916]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 2 (b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + \cos(x)y = \cos(x)\sin(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)+cos(x)*y(x)=sin(x)*cos(x),y(x), singsol=all)
```

$$y(x) = \sin(x) - 1 + c_1 e^{-\sin(x)}$$

✓ Solution by Mathematica

Time used: 0.049 (sec). Leaf size: 18

```
DSolve[y'[x]+Cos[x]*y[x]==Sin[x]*Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sin(x) + c_1 e^{-\sin(x)} - 1$$

1.6 problem 2 (c)

Internal problem ID [5917]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 2 (c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x$2)-y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-x}c_1 + e^xc_2$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 20

```
DSolve[y''[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1e^x + c_2e^{-x}$$

1.7 problem 2 (f)

Internal problem ID [5918]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 2 (f).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 4y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(2x) + c_2 \cos(2x)$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 20

```
DSolve[y''[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(2x) + c_2 \sin(2x)$$

1.8 problem 2 (h)

Internal problem ID [5919]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 2 (h).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + k^2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+k^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(kx) + c_2 \cos(kx)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 20

```
DSolve[y''[x]+k^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(kx) + c_2 \sin(kx)$$

1.9 problem 3(a)

Internal problem ID [5920]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 3(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y' + 5y = 2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)+5*y(x)=2,y(x), singsol=all)
```

$$y(x) = \frac{2}{5} + e^{-5x}c_1$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 24

```
DSolve[y'[x]+5*y[x]==2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2}{5} + c_1 e^{-5x}$$

$$y(x) \rightarrow \frac{2}{5}$$

1.10 problem 4(a)

Internal problem ID [5921]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 4(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$y'' = 3x + 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x$2)=3*x+1,y(x), singsol=all)
```

$$y(x) = \frac{1}{2}x^3 + \frac{1}{2}x^2 + c_1x + c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 25

```
DSolve[y''[x]==3*x+1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(x^3 + x^2 + 2c_2x + 2c_1)$$

1.11 problem 5(a)

Internal problem ID [5922]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 5(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - yk = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)=k*y(x),y(x), singsol=all)
```

$$y(x) = c_1 e^{kx}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 18

```
DSolve[y'[x]==k*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{kx}$$

$$y(x) \rightarrow 0$$

2 Chapter 1.6 Introduction– Linear equations of First Order. Page 41

2.1	problem 1(a)	16
2.2	problem 1(b)	17
2.3	problem 1(c)	18
2.4	problem 1(d)	19
2.5	problem 1(e)	20
2.6	problem 2	21
2.7	problem 3	22
2.8	problem 4	23
2.9	problem 5	24
2.10	problem 7	25

2.1 problem 1(a)

Internal problem ID [5923]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y' - 2y = 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)-2*y(x)=1,y(x), singsol=all)
```

$$y(x) = -\frac{1}{2} + e^{2x}c_1$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 24

```
DSolve[y'[x]-2*y[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2} + c_1 e^{2x}$$

$$y(x) \rightarrow -\frac{1}{2}$$

2.2 problem 1(b)

Internal problem ID [5924]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 1(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + y = e^x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)+y(x)=exp(x),y(x), singsol=all)
```

$$y(x) = \frac{e^x}{2} + e^{-x}c_1$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 21

```
DSolve[y'[x]+y[x]==Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^x}{2} + c_1 e^{-x}$$

2.3 problem 1(c)

Internal problem ID [5925]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 1(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' - 2y = x^2 + x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x)-2*y(x)=x^2+x,y(x), singsol=all)
```

$$y(x) = -\frac{x^2}{2} - x - \frac{1}{2} + e^{2x}c_1$$

✓ Solution by Mathematica

Time used: 0.073 (sec). Leaf size: 23

```
DSolve[y'[x]-2*y[x]==x^2+x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2}(x+1)^2 + c_1 e^{2x}$$

2.4 problem 1(d)

Internal problem ID [5926]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 1(d).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y + 3y' = 2e^{-x}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve(3*diff(y(x),x)+y(x)=2*exp(-x),y(x), singsol=all)
```

$$y(x) = -e^{-x} + e^{-\frac{x}{3}}c_1$$

✓ Solution by Mathematica

Time used: 0.051 (sec). Leaf size: 23

```
DSolve[3*y'[x]+y[x]==2*Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(-1 + c_1 e^{2x/3})$$

2.5 problem 1(e)

Internal problem ID [5927]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 1(e).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 3y = e^{ix}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)+3*y(x)=exp(I*x),y(x), singsol=all)
```

$$y(x) = \left(\left(\frac{3}{10} - \frac{i}{10} \right) e^{(3+i)x} + c_1 \right) e^{-3x}$$

✓ Solution by Mathematica

Time used: 0.054 (sec). Leaf size: 29

```
DSolve[y'[x]+3*y[x]==Exp[I*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left(\frac{3}{10} - \frac{i}{10} \right) e^{ix} + c_1 e^{-3x}$$

2.6 problem 2

Internal problem ID [5928]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + iy = x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)+I*y(x)=x,y(x), singsol=all)
```

$$y(x) = -ix + 1 + e^{-ix}c_1$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 22

```
DSolve[y'[x]+I*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -ix + c_1e^{-ix} + 1$$

2.7 problem 3

Internal problem ID [5929]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$Ly' + Ry = E$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(L*diff(y(x),x)+R*y(x)=E,y(x), singsol=all)
```

$$y(x) = \frac{E}{R} + e^{-\frac{Rx}{L}} c_1$$

✓ Solution by Mathematica

Time used: 0.056 (sec). Leaf size: 23

```
DSolve[L*y'[x]+R*y[x]==E0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{E0 - E0e^{-\frac{Rx}{L}}}{R}$$

2.8 problem 4

Internal problem ID [5930]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$Ly' + Ry = E \sin(\omega x)$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 47

```
dsolve([L*diff(y(x),x)+R*y(x)=E*sin(omega*x),y(0) = 0],y(x), singsol=all)
```

$$y(x) = -\frac{E \left(L \cos(\omega x) \omega - e^{-\frac{Rx}{L}} L \omega - \sin(\omega x) R \right)}{\omega^2 L^2 + R^2}$$

✓ Solution by Mathematica

Time used: 0.115 (sec). Leaf size: 47

```
DSolve[{L*y'[x]+R*y[x]==E0*Sin[\[Omega]*x],{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{E0 \left(L \omega e^{-\frac{Rx}{L}} - L \omega \cos(x\omega) + R \sin(x\omega) \right)}{L^2 \omega^2 + R^2}$$

2.9 problem 5

Internal problem ID [5931]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$Ly' + Ry = E e^{i\omega x}$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 38

```
dsolve([L*diff(y(x),x)+R*y(x)=E*exp(I*omega*x),y(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{E \left(e^{\frac{x(iL\omega + R)}{L}} - 1 \right) e^{-\frac{Rx}{L}}}{iL\omega + R}$$

✓ Solution by Mathematica

Time used: 0.101 (sec). Leaf size: 43

```
DSolve[{L*y'[x]+R*y[x]==E0*Exp[I*\[Omega]*x],{y[0]==0}},y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow \frac{E0 e^{-\frac{Rx}{L}} \left(-1 + e^{\frac{x(R+iL\omega)}{L}} \right)}{R + iL\omega}$$

2.10 problem 7

Internal problem ID [5932]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + ya = b(x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)+a*y(x)=b(x),y(x), singsol=all)
```

$$y(x) = \left(\int b(x) e^{ax} dx + c_1 \right) e^{-ax}$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 32

```
DSolve[y'[x]+a*y[x]==b[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-ax} \left(\int_1^x e^{aK[1]} b(K[1]) dK[1] + c_1 \right)$$

3 Chapter 1. Introduction– Linear equations of First Order. Page 45

3.1	problem 1(a)	27
3.2	problem 1(b)	28
3.3	problem 1(c)	29
3.4	problem 1(d)	30
3.5	problem 1(e)	31
3.6	problem 2	32
3.7	problem 3	33
3.8	problem 8	34
3.9	problem 14(a)	35
3.10	problem 14(b)	36
3.11	problem 14(b)	37

3.1 problem 1(a)

Internal problem ID [5933]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$2yx + y' = x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)+2*x*y(x)=x,y(x), singsol=all)
```

$$y(x) = \frac{1}{2} + c_1 e^{-x^2}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 26

```
DSolve[y'[x]+2*x*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} + c_1 e^{-x^2}$$

$$y(x) \rightarrow \frac{1}{2}$$

3.2 problem 1(b)

Internal problem ID [5934]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 1(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y'x + y = 3x^3 - 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(x*diff(y(x),x)+y(x)=3*x^3-1,y(x), singsol=all)
```

$$y(x) = \frac{\frac{3}{4}x^4 - x + c_1}{x}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 20

```
DSolve[x*y'[x]+y[x]==3*x^3-1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3x^3}{4} + \frac{c_1}{x} - 1$$

3.3 problem 1(c)

Internal problem ID [5935]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 1(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' + e^x y = 3 e^x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve(diff(y(x),x)+exp(x)*y(x)=3*exp(x),y(x), singsol=all)
```

$$y(x) = 3 + e^{-e^x} c_1$$

✓ Solution by Mathematica

Time used: 0.057 (sec). Leaf size: 22

```
DSolve[y'[x]+Exp[x]*y[x]==3*Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3 + c_1 e^{-e^x}$$

$$y(x) \rightarrow 3$$

3.4 problem 1(d)

Internal problem ID [5936]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 1(d).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' - y \tan(x) = e^{\sin(x)}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)-tan(x)*y(x)=exp(sin(x)),y(x), singsol=all)
```

$$y(x) = \frac{e^{\sin(x)} + c_1}{\cos(x)}$$

✓ Solution by Mathematica

Time used: 0.149 (sec). Leaf size: 15

```
DSolve[y'[x]-Tan[x]*y[x]==Exp[Sin[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sec(x) (e^{\sin(x)} + c_1)$$

3.5 problem 1(e)

Internal problem ID [5937]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 1(e).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$2yx + y' = x e^{-x^2}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 18

```
dsolve(diff(y(x),x)+2*x*y(x)=x*exp(-x^2),y(x), singsol=all)
```

$$y(x) = \left(\frac{x^2}{2} + c_1 \right) e^{-x^2}$$

✓ Solution by Mathematica

Time used: 0.057 (sec). Leaf size: 24

```
DSolve[y'[x]+2*x*y[x]==x*Exp[-x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^{-x^2} (x^2 + 2c_1)$$

3.6 problem 2

Internal problem ID [5938]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + \cos(x)y = e^{-\sin(x)}$$

With initial conditions

$$[y(\pi) = \pi]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve([diff(y(x),x)+cos(x)*y(x)=exp(-sin(x)),y(Pi) = Pi],y(x), singsol=all)
```

$$y(x) = e^{-\sin(x)}x$$

✓ Solution by Mathematica

Time used: 0.138 (sec). Leaf size: 13

```
DSolve[{y'[x]+Cos[x]*y[x]==Exp[-Sin[x]],{y[Pi]==Pi}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow xe^{-\sin(x)}$$

3.7 problem 3

Internal problem ID [5939]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$x^2y' + 2yx = 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(x^2*diff(y(x),x)+2*x*y(x)=1,y(x), singsol=all)
```

$$y(x) = \frac{x + c_1}{x^2}$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 13

```
DSolve[x^2*y'[x]+2*x*y[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x + c_1}{x^2}$$

3.8 problem 8

Internal problem ID [5940]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 2y = b(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x)+2*y(x)=b(x),y(x), singsol=all)
```

$$y(x) = \left(\int b(x) e^{2x} dx + c_1 \right) e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.045 (sec). Leaf size: 31

```
DSolve[y'[x]+2*y[x]==b[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} \left(\int_1^x e^{2K[1]} b(K[1]) dK[1] + c_1 \right)$$

3.9 problem 14(a)

Internal problem ID [5941]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 14(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - y = 1$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 8

```
dsolve([diff(y(x),x)=1+y(x),y(0) = 0],y(x), singsol=all)
```

$$y(x) = e^x - 1$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 10

```
DSolve[{y'[x]==1+y[x],{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x - 1$$

3.10 problem 14(b)

Internal problem ID [5942]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 14(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$-y^2 + y' = 1$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 6

```
dsolve([diff(y(x),x)=1+y(x)^2,y(0) = 0],y(x), singsol=all)
```

$$y(x) = \tan(x)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 7

```
DSolve[{y'[x]==1+y[x]^2,{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan(x)$$

3.11 problem 14(b)

Internal problem ID [5943]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 14(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$-y^2 + y' = 1$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 6

```
dsolve([diff(y(x),x)=1+y(x)^2,y(0) = 0],y(x), singsol=all)
```

$$y(x) = \tan(x)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 7

```
DSolve[{y'[x]==1+y[x]^2,{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan(x)$$

4 Chapter 2. Linear equations with constant coefficients. Page 52

4.1	problem 1(a)	39
4.2	problem 1(b)	40
4.3	problem 1(c)	41
4.4	problem 1(d)	42
4.5	problem 1(e)	43
4.6	problem 1(f)	44
4.7	problem 1(g)	45
4.8	problem 2(a)	46
4.9	problem 2(b)	47
4.10	problem 3(a)	48
4.11	problem 3(b)	49
4.12	problem 3(c)	50
4.13	problem 3(d)	51

4.1 problem 1(a)

Internal problem ID [5944]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-4*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{2x}c_1 + c_2e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 22

```
DSolve[y''[x]-4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(c_1e^{4x} + c_2)$$

4.2 problem 1(b)

Internal problem ID [5945]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$3y'' + 2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(3*diff(y(x),x$2)+2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin\left(\frac{\sqrt{6}x}{3}\right) + c_2 \cos\left(\frac{\sqrt{6}x}{3}\right)$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 32

```
DSolve[3*y''[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos\left(\sqrt{\frac{2}{3}}x\right) + c_2 \sin\left(\sqrt{\frac{2}{3}}x\right)$$

4.3 problem 1(c)

Internal problem ID [5946]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 16y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+16*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(4x) + c_2 \cos(4x)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 20

```
DSolve[y''[x]+16*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(4x) + c_2 \sin(4x)$$

4.4 problem 1(d)

Internal problem ID [5947]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$y'' = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 9

```
dsolve(diff(y(x),x$2)=0,y(x), singsol=all)
```

$$y(x) = c_1x + c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 12

```
DSolve[y''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2x + c_1$$

4.5 problem 1(e)

Internal problem ID [5948]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2iy' + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$2)+2*I*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-ix} \sin(\sqrt{2}x) + c_2 e^{-ix} \cos(\sqrt{2}x)$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 38

```
DSolve[y''[x]+2*I*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-i(1+\sqrt{2})x} (c_2 e^{2i\sqrt{2}x} + c_1)$$

4.6 problem 1(f)

Internal problem ID [5949]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y' + 5y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+5*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(x) e^{2x} + c_2 \cos(x) e^{2x}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 22

```
DSolve[y''[x]-4*y'[x]+5*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(c_2 \cos(x) + c_1 \sin(x))$$

4.7 problem 1(g)

Internal problem ID [5950]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 1(g).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + (-1 + 3i)y' - 3iy = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x$2)+(3*I-1)*diff(y(x),x)-3*I*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-3ix} + e^x c_2$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 22

```
DSolve[y''[x]+(3*I-1)*y'[x]-3*I*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-3ix} + c_2 e^x$$

4.8 problem 2(a)

Internal problem ID [5951]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 2(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y' - 6y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 18

```
dsolve([diff(y(x),x$2)+diff(y(x),x)-6*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{(3e^{5x} + 2)e^{-3x}}{5}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 23

```
DSolve[{y''[x]+y'[x]-6*y[x]==0,{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{5}e^{-3x}(3e^{5x} + 2)$$

4.9 problem 2(b)

Internal problem ID [5952]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 2(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y' - 6y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve([diff(y(x),x$2)+diff(y(x),x)-6*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{(e^{5x} - 1)e^{-3x}}{5}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 21

```
DSolve[{y''[x]+y'[x]-6*y[x]==0,{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{5}e^{-3x}(e^{5x} - 1)$$

4.10 problem 3(a)

Internal problem ID [5953]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 3(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y = 0$$

With initial conditions

$$\left[y(0) = 1, y\left(\frac{\pi}{2}\right) = 2 \right]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve([diff(y(x),x$2)+y(x)=0,y(0) = 1, y(1/2*Pi) = 2],y(x), singsol=all)
```

$$y(x) = 2 \sin(x) + \cos(x)$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 12

```
DSolve[{y'[x]+y[x]==0,{y[0]==1,y[Pi/2]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 \sin(x) + \cos(x)$$

4.11 problem 3(b)

Internal problem ID [5954]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 3(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y = 0$$

With initial conditions

$$[y(0) = 0, y(\pi) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 8

```
dsolve([diff(y(x),x$2)+y(x)=0,y(0) = 0, y(Pi) = 0],y(x), singsol=all)
```

$$y(x) = c_1 \sin(x)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 10

```
DSolve[{y''[x]+y[x]==0,{y[0]==0,y[Pi]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \sin(x)$$

4.12 problem 3(c)

Internal problem ID [5955]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 3(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y = 0$$

With initial conditions

$$\left[y(0) = 0, y'\left(\frac{\pi}{2}\right) = 0 \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 8

```
dsolve([diff(y(x),x$2)+y(x)=0,y(0) = 0, D(y)(1/2*Pi) = 0],y(x), singsol=all)
```

$$y(x) = c_1 \sin(x)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 10

```
DSolve[{y'[x]+y[x]==0,{y[0]==0,y'[Pi/2]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \sin(x)$$

4.13 problem 3(d)

Internal problem ID [5956]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 3(d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y = 0$$

With initial conditions

$$\left[y(0) = 0, y\left(\frac{\pi}{2}\right) = 0 \right]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve([diff(y(x),x$2)+y(x)=0,y(0) = 0, y(1/2*Pi) = 0],y(x), singsol=all)
```

$$y(x) = 0$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 6

```
DSolve[{y''[x]+y[x]==0,{y[0]==0,y[Pi/2]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

5 Chapter 2. Linear equations with constant coefficients. Page 59

5.1	problem 1(a)	53
5.2	problem 1(b)	54
5.3	problem 1(c)	55
5.4	problem 1(d)	56

5.1 problem 1(a)

Internal problem ID [5957]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 59

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' - 3y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)-2*diff(y(x),x)-3*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{e^{3x}}{4} - \frac{e^{-x}}{4}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 21

```
DSolve[{y'[x]-2*y'[x]-3*y[x]==0,{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{1}{4}e^{-x}(e^{4x} - 1)$$

5.2 problem 1(b)

Internal problem ID [5958]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 59

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + (1 + 4i)y' + y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 5

```
dsolve([diff(y(x),x$2)+(4*I+1)*diff(y(x),x)+y(x)=0,y(0) = 0, D(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = 0$$

✓ Solution by Mathematica

Time used: 0.054 (sec). Leaf size: 6

```
DSolve[{y'[x]+(4*I+1)*y'[x]+y[x]==0,{y[0]==0,y'[0]==0}},y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow 0$$

5.3 problem 1(c)

Internal problem ID [5959]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 59

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + (-1 + 3i)y' - 3iy = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve([diff(y(x),x$2)+(3*I-1)*diff(y(x),x)-3*I*y(x)=0,y(0) = 2, D(y)(0) = 0],y(x), singsol=
```

$$y(x) = \left(\frac{1}{5} - \frac{3i}{5}\right) e^{-3ix} + \left(\frac{9}{5} + \frac{3i}{5}\right) e^x$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 31

```
DSolve[{y'[x]+(3*I-1)*y'[x]-3*I*y[x]==0,{y[0]==2,y'[0]==0}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{1}{5} e^{-3ix} ((9 + 3i)e^{(1+3i)x} + (1 - 3i))$$

5.4 problem 1(d)

Internal problem ID [5960]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 59

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 10y = 0$$

With initial conditions

$$[y(0) = \pi, y'(0) = \pi^2]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 27

```
dsolve([diff(y(x),x$2)+10*y(x)=0,y(0) = Pi, D(y)(0) = Pi^2],y(x), singsol=all)
```

$$y(x) = \frac{\pi(\pi\sqrt{10} \sin(\sqrt{10}x) + 10 \cos(\sqrt{10}x))}{10}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 33

```
DSolve[{y'[x]+10*y[x]==0,{y[0]==Pi,y'[0]==Pi^2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\pi^2 \sin(\sqrt{10}x)}{\sqrt{10}} + \pi \cos(\sqrt{10}x)$$

6 Chapter 2. Linear equations with constant coefficients. Page 69

6.1	problem 1(a)	58
6.2	problem 1(b)	59
6.3	problem 1(c)	60
6.4	problem 1(d)	61
6.5	problem 1(e)	62
6.6	problem 1(f)	63
6.7	problem 1(g)	64
6.8	problem 1(h)	65
6.9	problem 1(i)	66
6.10	problem 1(j)	67
6.11	problem 4(c)	68

6.1 problem 1(a)

Internal problem ID [5961]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y = \cos(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)+4*y(x)=cos(x),y(x), singsol=all)
```

$$y(x) = \sin(2x) c_2 + \cos(2x) c_1 + \frac{\cos(x)}{3}$$

✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 26

```
DSolve[y''[x]+4*y[x]==Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\cos(x)}{3} + c_1 \cos(2x) + c_2 \sin(2x)$$

6.2 problem 1(b)

Internal problem ID [5962]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = \sin(3x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)+9*y(x)=sin(3*x),y(x), singsol=all)
```

$$y(x) = \sin(3x) c_2 + \cos(3x) c_1 - \frac{\cos(3x) x}{6}$$

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 33

```
DSolve[y''[x]+9*y[x]==Sin[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left(-\frac{x}{6} + c_1\right) \cos(3x) + \frac{1}{36}(1 + 36c_2) \sin(3x)$$

6.3 problem 1(c)

Internal problem ID [5963]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \tan(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)+y(x)=tan(x),y(x), singsol=all)
```

$$y(x) = \sin(x) c_2 + \cos(x) c_1 - \cos(x) \ln(\sec(x) + \tan(x))$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 23

```
DSolve[y''[x]+y[x]==Tan[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \cos(x)(-\operatorname{arctanh}(\sin(x))) + c_1 \cos(x) + c_2 \sin(x)$$

6.4 problem 1(d)

Internal problem ID [5964]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 2iy' + y = x$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$2)+2*I*diff(y(x),x)+y(x)=x,y(x), singsol=all)
```

$$y(x) = e^{-ix} \sin(\sqrt{2}x) c_2 + e^{-ix} \cos(\sqrt{2}x) c_1 - 2i + x$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 44

```
DSolve[y''[x]+2*I*y'[x]+y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x + c_1 e^{-i(1+\sqrt{2})x} + c_2 e^{i(\sqrt{2}-1)x} - 2i$$

6.5 problem 1(e)

Internal problem ID [5965]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y' + 5y = 3e^{-x} + 2x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+5*y(x)=3*exp(-x)+2*x^2,y(x), singsol=all)
```

$$y(x) = \sin(x) e^{2x} c_2 + \cos(x) e^{2x} c_1 + \frac{3e^{-x}}{10} + \frac{2x^2}{5} + \frac{16x}{25} + \frac{44}{125}$$

✓ Solution by Mathematica

Time used: 0.316 (sec). Leaf size: 47

```
DSolve[y''[x]-4*y'[x]+5*y[x]==3*Exp[-x]+2*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{250}(100x^2 + 160x + 75e^{-x} + 88) + c_2 e^{2x} \cos(x) + c_1 e^{2x} \sin(x)$$

6.6 problem 1(f)

Internal problem ID [5966]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 7y' + 6y = \sin(x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)-7*diff(y(x),x)+6*y(x)=sin(x),y(x), singsol=all)
```

$$y(x) = c_2 e^{6x} + e^x c_1 + \frac{7 \cos(x)}{74} + \frac{5 \sin(x)}{74}$$

✓ Solution by Mathematica

Time used: 0.063 (sec). Leaf size: 32

```
DSolve[y''[x]-7*y'[x]+6*y[x]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{5 \sin(x)}{74} + \frac{7 \cos(x)}{74} + c_1 e^x + c_2 e^{6x}$$

6.7 problem 1(g)

Internal problem ID [5967]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(g).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = 2 \sin(x) \sin(2x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)+y(x)=2*sin(x)*sin(2*x),y(x), singsol=all)
```

$$y(x) = \sin(x) c_2 + \cos(x) c_1 + \frac{\sin(x) (-\cos(x) \sin(x) + x)}{2}$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 33

```
DSolve[y''[x]+y[x]==2*Sin[x]*Sin[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{8}(\cos(3x) + (-1 + 8c_1) \cos(x) + 4(x + 2c_2) \sin(x))$$

6.8 problem 1(h)

Internal problem ID [5968]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(h).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \sec(x)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)+y(x)=sec(x),y(x), singsol=all)
```

$$y(x) = \sin(x) c_2 + \cos(x) c_1 - \ln(\sec(x)) \cos(x) + \sin(x) x$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 22

```
DSolve[y''[x]+y[x]==Sec[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x + c_2) \sin(x) + \cos(x)(\log(\cos(x)) + c_1)$$

6.9 problem 1(i)

Internal problem ID [5969]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(i).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4y'' - y = e^x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(4*diff(y(x),x$2)-y(x)=exp(x),y(x), singsol=all)
```

$$y(x) = e^{-\frac{x}{2}}c_2 + e^{\frac{x}{2}}c_1 + \frac{e^x}{3}$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 33

```
DSolve[4*y''[x]-y[x]==Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^x}{3} + c_1 e^{x/2} + c_2 e^{-x/2}$$

6.10 problem 1(j)

Internal problem ID [5970]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(j).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$6y'' + 5y' - 6y = x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(6*diff(y(x),x$2)+5*diff(y(x),x)-6*y(x)=x,y(x), singsol=all)
```

$$y(x) = e^{-\frac{3x}{2}} c_2 + e^{\frac{2x}{3}} c_1 - \frac{x}{6} - \frac{5}{36}$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 34

```
DSolve[6*y''[x]+5*y'[x]-6*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x}{6} + c_1 e^{2x/3} + c_2 e^{-3x/2} - \frac{5}{36}$$

6.11 problem 4(c)

Internal problem ID [5971]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 4(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + \omega^2 y = A \cos(\omega x)$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 18

```
dsolve([diff(y(x),x$2)+omega^2*y(x)=A*cos(omega*x),y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{\sin(\omega x)(Ax + 2)}{2\omega}$$

✓ Solution by Mathematica

Time used: 0.058 (sec). Leaf size: 21

```
DSolve[{y''[x]+\[Omega]^2*y[x]==A*Cos[\[Omega]*x],{y[0]==0,y'[0]==1}},y[x],x,IncludeSingular
```

$$y(x) \rightarrow \frac{(Ax + 2) \sin(x\omega)}{2\omega}$$

7 Chapter 2. Linear equations with constant coefficients. Page 74

7.1	problem 4(a)	70
7.2	problem 4(b)	71
7.3	problem 4(c)	72
7.4	problem 4(d)	73
7.5	problem 4(f)	74
7.6	problem 4(g)	75
7.7	problem 4(h)	76
7.8	problem 4(i)	77

7.1 problem 4(a)

Internal problem ID [5972]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(a).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 8y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$3)-8*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{2x}c_1 + c_2e^{-x} \sin(\sqrt{3}x) + c_3e^{-x} \cos(\sqrt{3}x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 42

```
DSolve[y'''[x]-8*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} \left(c_1 e^{3x} + c_2 \cos(\sqrt{3}x) + c_3 \sin(\sqrt{3}x) \right)$$

7.2 problem 4(b)

Internal problem ID [5973]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(b).

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 16y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 65

```
dsolve(diff(y(x),x$4)+16*y(x)=0,y(x), singsol=all)
```

$$y(x) = -c_1 e^{-\sqrt{2}x} \sin(\sqrt{2}x) - c_2 e^{\sqrt{2}x} \sin(\sqrt{2}x) \\ + c_3 e^{-\sqrt{2}x} \cos(\sqrt{2}x) + c_4 e^{\sqrt{2}x} \cos(\sqrt{2}x)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 67

```
DSolve[y''''[x]+16*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\sqrt{2}x} \left((c_1 e^{2\sqrt{2}x} + c_2) \cos(\sqrt{2}x) + (c_4 e^{2\sqrt{2}x} + c_3) \sin(\sqrt{2}x) \right)$$

7.3 problem 4(c)

Internal problem ID [5974]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(c).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 5y'' + 6y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$3)-5*diff(y(x),x$2)+6*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1 + e^{2x}c_2 + c_3e^{3x}$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 30

```
DSolve[y'''[x]-5*y''[x]+6*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}c_1e^{2x} + \frac{1}{3}c_2e^{3x} + c_3$$

7.4 problem 4(d)

Internal problem ID [5975]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(d).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - iy'' + 4y' - 4iy = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$3)-I*diff(y(x),x$2)+4*diff(y(x),x)-4*I*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{2ix} + c_2 e^{ix} + c_3 e^{-2ix}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 36

```
DSolve[y'''[x]-I*y''[x]+4*y'[x]-4*I*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2ix} (c_2 e^{4ix} + c_3 e^{3ix} + c_1)$$

7.5 problem 4(f)

Internal problem ID [5976]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(f).

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 5y'' + 4y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$4)+5*diff(y(x),x$2)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(2x) + c_2 \cos(2x) + c_3 \sin(x) + c_4 \cos(x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
DSolve[y''''[x]+5*y''[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(2x) + c_4 \sin(x) + \cos(x)(2c_2 \sin(x) + c_3)$$

7.6 problem 4(g)

Internal problem ID [5977]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(g).

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 16y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$4)-16*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{2x}c_1 + c_2e^{-2x} + c_3 \sin(2x) + c_4 \cos(2x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 36

```
DSolve[y''''[x]-16*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1e^{2x} + c_3e^{-2x} + c_2 \cos(2x) + c_4 \sin(2x)$$

7.7 problem 4(h)

Internal problem ID [5978]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(h).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 3y' - 2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$3)-3*diff(y(x),x)-2*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{2x}c_1 + c_2e^{-x} + c_3e^{-x}x$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 26

```
DSolve[y'''[x]-3*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(c_2x + c_3e^{3x} + c_1)$$

7.8 problem 4(i)

Internal problem ID [5979]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(i).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 3iy'' - 3y' + iy = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$3)-3*I*diff(y(x),x$2)-3*diff(y(x),x)+I*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{ix} + c_2 e^{ix} x + c_3 e^{ix} x^2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 25

```
DSolve[y'''[x]-3*I*y''[x]-3*y'[x]+I*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{ix}(x(c_3 x + c_2) + c_1)$$

8 Chapter 2. Linear equations with constant coefficients. Page 79

8.1	problem 1(c)	79
8.2	problem 2(c)	80

8.1 problem 1(c)

Internal problem ID [5980]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 79

Problem number: 1(c).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 4y' = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1, y''(0) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$3)-4*diff(y(x),x)=0,y(0) = 0, D(y)(0) = 1, (D@@2)(y)(0) = 0],y(x), sings
```

$$y(x) = \frac{e^{2x}}{4} - \frac{e^{-2x}}{4}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 69

```
DSolve[{y'''[x]-4*y[x]==0,{y[0]==0,y'[0]==1,y''[0]==0}},y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow \frac{e^{-\frac{x}{\sqrt[3]{2}}} \left(e^{\frac{3x}{\sqrt[3]{2}}} + \sqrt{3} \sin \left(\frac{\sqrt{3}x}{\sqrt[3]{2}} \right) - \cos \left(\frac{\sqrt{3}x}{\sqrt[3]{2}} \right) \right)}{3 \cdot 2^{2/3}}$$

8.2 problem 2(c)

Internal problem ID [5981]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 79

Problem number: 2(c).

ODE order: 5.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y^{(5)} - y'''' - y' + y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0, y''(0) = 0, y'''(0) = 0, y''''(0) = 0]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 28

```
dsolve([diff(y(x),x$5)-diff(y(x),x$4)-diff(y(x),x)+y(x)=0,y(0) = 1, D(y)(0) = 0, D@@2)(y)(0)
```

$$y(x) = \frac{e^{-x}}{8} + \frac{(-2x + 5)e^x}{8} + \frac{\cos(x)}{4} - \frac{\sin(x)}{4}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 34

```
DSolve[{y'''''[x]-y''''[x]-y'[x]+y[x]==0,{y[0]==1,y'[0]==0,y''[0]==0,y''''[0]==0,y'''''[0]==0}
```

$$y(x) \rightarrow \frac{1}{8}(-2e^x x + e^{-x} + 5e^x - 2\sin(x) + 2\cos(x))$$

9 Chapter 2. Linear equations with constant coefficients. Page 83

9.1	problem 1(a)	82
9.2	problem 1(b)	83
9.3	problem 1(c)	84
9.4	problem 1(d)	85
9.5	problem 1(e)	87
9.6	problem 2	88
9.7	problem 3(a)	89
9.8	problem 3(b)	90
9.9	problem 5(b)	91

9.1 problem 1(a)

Internal problem ID [5982]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x$2)+y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(x) + c_2 \cos(x)$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 16

```
DSolve[y''[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(x) + c_2 \sin(x)$$

9.2 problem 1(b)

Internal problem ID [5983]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x$2)-y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-x}c_1 + e^xc_2$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 20

```
DSolve[y''[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1e^x + c_2e^{-x}$$

9.3 problem 1(c)

Internal problem ID [5984]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 1(c).

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$4)-y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-x}c_1 + e^x c_2 + c_3 \sin(x) + c_4 \cos(x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
DSolve[y''''[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x + c_3 e^{-x} + c_2 \cos(x) + c_4 \sin(x)$$

9.4 problem 1(d)

Internal problem ID [5985]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 1(d).

ODE order: 5.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y^{(5)} + 2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 369

```
dsolve(diff(y(x),x$5)+2*y(x)=0,y(x), singsol=all)
```

$$\begin{aligned}
 & y(x) \\
 = & c_1 e^{\left(\frac{2^{\frac{1}{5}} \cos\left(\frac{\pi}{5}\right) \sqrt{5}}{4} - \frac{\cos\left(\frac{\pi}{5}\right) 2^{\frac{1}{5}}}{4} + \frac{2^{\frac{7}{10}} \sqrt{5+\sqrt{5}} \sin\left(\frac{\pi}{5}\right)}{4} - \frac{i 2^{\frac{7}{10}} \sqrt{5+\sqrt{5}} \cos\left(\frac{\pi}{5}\right)}{4} + \frac{i 2^{\frac{1}{5}} \sin\left(\frac{\pi}{5}\right) \sqrt{5}}{4} - \frac{i \sin\left(\frac{\pi}{5}\right) 2^{\frac{1}{5}}}{4} \right) x} \\
 & + c_2 e^{\left(-\frac{2^{\frac{1}{5}} \cos\left(\frac{\pi}{5}\right) \sqrt{5}}{4} - \frac{\cos\left(\frac{\pi}{5}\right) 2^{\frac{1}{5}}}{4} + \frac{2^{\frac{7}{10}} \sqrt{5-\sqrt{5}} \sin\left(\frac{\pi}{5}\right)}{4} - \frac{i 2^{\frac{7}{10}} \sqrt{5-\sqrt{5}} \cos\left(\frac{\pi}{5}\right)}{4} - \frac{i 2^{\frac{1}{5}} \sin\left(\frac{\pi}{5}\right) \sqrt{5}}{4} - \frac{i \sin\left(\frac{\pi}{5}\right) 2^{\frac{1}{5}}}{4} \right) x} \\
 & + c_3 e^{\left(-\frac{2^{\frac{1}{5}} \cos\left(\frac{\pi}{5}\right) \sqrt{5}}{4} - \frac{\cos\left(\frac{\pi}{5}\right) 2^{\frac{1}{5}}}{4} - \frac{2^{\frac{7}{10}} \sqrt{5-\sqrt{5}} \sin\left(\frac{\pi}{5}\right)}{4} + \frac{i 2^{\frac{7}{10}} \sqrt{5-\sqrt{5}} \cos\left(\frac{\pi}{5}\right)}{4} - \frac{i 2^{\frac{1}{5}} \sin\left(\frac{\pi}{5}\right) \sqrt{5}}{4} - \frac{i \sin\left(\frac{\pi}{5}\right) 2^{\frac{1}{5}}}{4} \right) x} \\
 & + c_4 e^{\left(\frac{2^{\frac{1}{5}} \cos\left(\frac{\pi}{5}\right) \sqrt{5}}{4} - \frac{\cos\left(\frac{\pi}{5}\right) 2^{\frac{1}{5}}}{4} - \frac{2^{\frac{7}{10}} \sqrt{5+\sqrt{5}} \sin\left(\frac{\pi}{5}\right)}{4} + \frac{i 2^{\frac{7}{10}} \sqrt{5+\sqrt{5}} \cos\left(\frac{\pi}{5}\right)}{4} + \frac{i 2^{\frac{1}{5}} \sin\left(\frac{\pi}{5}\right) \sqrt{5}}{4} - \frac{i \sin\left(\frac{\pi}{5}\right) 2^{\frac{1}{5}}}{4} \right) x} \\
 & + c_5 e^{\left(\cos\left(\frac{\pi}{5}\right) 2^{\frac{1}{5}} + i \sin\left(\frac{\pi}{5}\right) 2^{\frac{1}{5}} \right) x}
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 180

```
DSolve[y'''''[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\frac{(\sqrt{5}-1)x}{2 \cdot 2^{4/5}}} \left(c_5 e^{\frac{(\sqrt{5}-5)x}{2 \cdot 2^{4/5}}} \right. \\ \left. + c_3 e^{\frac{\sqrt{5}x}{2^{4/5}}} \cos\left(\frac{\sqrt{5-\sqrt{5}x}}{2 \cdot 2^{3/10}}\right) + c_4 \cos\left(\frac{\sqrt{5+\sqrt{5}x}}{2 \cdot 2^{3/10}}\right) + c_2 e^{\frac{\sqrt{5}x}{2^{4/5}}} \sin\left(\frac{\sqrt{5-\sqrt{5}x}}{2 \cdot 2^{3/10}}\right) + c_1 \sin\left(\frac{\sqrt{5+\sqrt{5}x}}{2 \cdot 2^{3/10}}\right) \right)$$

9.5 problem 1(e)

Internal problem ID [5986]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 1(e).

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 5y'' + 4y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$4)-5*diff(y(x),x$2)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{2x}c_1 + c_2e^{-2x} + c_3e^{-x} + c_4e^x$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 35

```
DSolve[y''''[x]-5*y''[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(c_2e^x + e^{3x}(c_4e^x + c_3) + c_1)$$

9.6 problem 2

Internal problem ID [5987]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 2.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' + y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1, y''(0) = 0]$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 39

```
dsolve([diff(y(x),x$3)+y(x)=0,y(0) = 0, D(y)(0) = 1, (D@@2)(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{\left(\sqrt{3} e^{\frac{3x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + e^{\frac{3x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) - 1\right) e^{-x}}{3}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 59

```
DSolve[{y'''[x]+y[x]==0,{y[0]==0,y'[0]==1,y''[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} e^{-x} \left(\sqrt{3} e^{3x/2} \sin\left(\frac{\sqrt{3}x}{2}\right) + e^{3x/2} \cos\left(\frac{\sqrt{3}x}{2}\right) - 1 \right)$$

9.7 problem 3(a)

Internal problem ID [5988]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 3(a).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - iy'' + y' - iy = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$3)-I*diff(y(x),x$2)+diff(y(x),x)-I*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-ix}c_1 + c_2e^{ix} + c_3e^{ix}x$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 31

```
DSolve[y'''[x]-I*y''[x]+y'[x]-I*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-ix}(e^{2ix}(c_3x + c_2) + c_1)$$

9.8 problem 3(b)

Internal problem ID [5989]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 3(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2iy' - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$2)-2*I*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{ix} + c_2 e^{ix} x$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 20

```
DSolve[y''[x]-2*I*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{ix}(c_2 x + c_1)$$

9.9 problem 5(b)

Internal problem ID [5990]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 5(b).

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - k^4 y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 0, y(1) = 0, y'(1) = 0]$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 5

```
dsolve([diff(y(x),x$4)-k^4*y(x)=0,y(0) = 0, D(y)(0) = 0, y(1) = 0, D(y)(1) = 0],y(x), singsol
```

$$y(x) = 0$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 6

```
DSolve[{y''''[x]-k^4*y[x]==0,{y[0]==0,y[1]==0,y'[0]==0,y'[1]==0}},y[x],x,IncludeSingularSolu
```

$$y(x) \rightarrow 0$$

10 Chapter 2. Linear equations with constant coefficients. Page 89

10.1	problem 1(a)	93
10.2	problem 1(b)	94
10.3	problem 1(c)	95
10.4	problem 1(d)	96
10.5	problem 1(e)	97
10.6	problem 1(f)	98

10.1 problem 1(a)

Internal problem ID [5991]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 89

Problem number: 1(a).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$y''' - y = x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 38

```
dsolve(diff(y(x),x$3)-y(x)=x,y(x), singsol=all)
```

$$y(x) = -x + e^x c_1 + c_2 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_3 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 57

```
DSolve[y'''[x]-y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x + c_1 e^x + c_2 e^{-x/2} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_3 e^{-x/2} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

10.2 problem 1(b)

Internal problem ID [5992]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 89

Problem number: 1(b).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$y''' - 8y = e^{ix}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 44

```
dsolve(diff(y(x),x$3)-8*y(x)=exp(I*x),y(x), singsol=all)
```

$$y(x) = \left(-\frac{8}{65} + \frac{i}{65}\right) e^{ix} + e^{2x} c_1 + c_2 e^{-x} \cos(\sqrt{3}x) + c_3 e^{-x} \sin(\sqrt{3}x)$$

✓ Solution by Mathematica

Time used: 0.472 (sec). Leaf size: 59

```
DSolve[y'''[x]-8*y[x]==Exp[I*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{65} e^{-x} \left(-(8-i)e^{(1+i)x} + 65c_1 e^{3x} + 65c_2 \cos(\sqrt{3}x) + 65c_3 \sin(\sqrt{3}x) \right)$$

10.3 problem 1(c)

Internal problem ID [5993]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 89

Problem number: 1(c).

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' + 16y = \cos(x)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 85

```
dsolve(diff(y(x),x$4)+16*y(x)=cos(x),y(x), singsol=all)
```

$$y(x) = -\frac{\cos(x)}{(5+2\sqrt{2})(-5+2\sqrt{2})} + c_1 e^{\sqrt{2}x} \cos(\sqrt{2}x) \\ + c_2 e^{\sqrt{2}x} \sin(\sqrt{2}x) + c_3 e^{-\sqrt{2}x} \cos(\sqrt{2}x) + c_4 e^{-\sqrt{2}x} \sin(\sqrt{2}x)$$

✓ Solution by Mathematica

Time used: 0.762 (sec). Leaf size: 74

```
DSolve[y''''[x]+16*y[x]==Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\cos(x)}{17} + e^{-\sqrt{2}x} \left((c_1 e^{2\sqrt{2}x} + c_2) \cos(\sqrt{2}x) + (c_4 e^{2\sqrt{2}x} + c_3) \sin(\sqrt{2}x) \right)$$

10.4 problem 1(d)

Internal problem ID [5994]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 89

Problem number: 1(d).

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _with_linear_symmetries]]`

$$y'''' - 4y''' + 6y'' - 4y' + y = e^x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$4)-4*diff(y(x),x$3)+6*diff(y(x),x$2)-4*diff(y(x),x)+y(x)=exp(x),y(x), sin
```

$$y(x) = \frac{e^x x^4}{24} + e^x c_1 + c_2 e^x x + c_3 e^x x^2 + c_4 e^x x^3$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 39

```
DSolve[y''''[x]-4*y'''[x]+6*y''[x]-4*y'[x]+y[x]==Exp[x],y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow \frac{1}{24} e^x (x^4 + 24c_4 x^3 + 24c_3 x^2 + 24c_2 x + 24c_1)$$

10.5 problem 1(e)

Internal problem ID [5995]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 89

Problem number: 1(e).

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - y = \cos(x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

```
dsolve(diff(y(x),x$4)-y(x)=cos(x),y(x), singsol=all)
```

$$y(x) = -\frac{\cos(x)}{4} - \frac{\sin(x)x}{4} + \cos(x)c_1 + e^x c_2 + c_3 \sin(x) + c_4 e^{-x}$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 40

```
DSolve[y''''[x]-y[x]==Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x + c_3 e^{-x} + \left(-\frac{1}{2} + c_2\right) \cos(x) + \left(-\frac{x}{4} + c_4\right) \sin(x)$$

10.6 problem 1(f)

Internal problem ID [5996]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 89

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 2iy' - y = e^{ix} - 2e^{-ix}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 44

```
dsolve(diff(y(x),x$2)-2*I*diff(y(x),x)-y(x)=exp(I*x)-2*exp(-I*x),y(x), singsol=all)
```

$$y(x) = c_2 e^{ix} + e^{ix} c_1 x + \frac{(x^2 + 2ix + 2) \cos(x)}{2} + \frac{x(ix - 2) \sin(x)}{2}$$

✓ Solution by Mathematica

Time used: 0.177 (sec). Leaf size: 39

```
DSolve[y''[x]-2*I*y'[x]-y[x]==Exp[I*x]-2*Exp[-I*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^{-ix} (1 + e^{2ix} (x^2 + 2c_2 x + 2c_1))$$

11 Chapter 2. Linear equations with constant coefficients. Page 93

11.1	problem 1(a)	100
11.2	problem 1(b)	101
11.3	problem 1(c)	102
11.4	problem 1(d)	103
11.5	problem 1(e)	104
11.6	problem 1(f)	105
11.7	problem 1(g)	106
11.8	problem 1(h)	107
11.9	problem 1(i)	108

11.1 problem 1(a)

Internal problem ID [5997]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y = \cos(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)+4*y(x)=cos(x),y(x), singsol=all)
```

$$y(x) = \sin(2x) c_2 + \cos(2x) c_1 + \frac{\cos(x)}{3}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 26

```
DSolve[y''[x]+4*y[x]==Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\cos(x)}{3} + c_1 \cos(2x) + c_2 \sin(2x)$$

11.2 problem 1(b)

Internal problem ID [5998]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y = \sin(2x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)+4*y(x)=sin(2*x),y(x), singsol=all)
```

$$y(x) = \sin(2x) c_2 + \cos(2x) c_1 - \frac{x \cos(2x)}{4}$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 33

```
DSolve[y''[x]+4*y[x]==Sin[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left(-\frac{x}{4} + c_1\right) \cos(2x) + \frac{1}{8}(1 + 16c_2) \sin(x) \cos(x)$$

11.3 problem 1(c)

Internal problem ID [5999]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y = 3e^{2x} + 4e^{-x}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$2)-4*y(x)=3*exp(2*x)+4*exp(-x),y(x), singsol=all)
```

$$y(x) = e^{2x}c_2 + e^{-2x}c_1 + \frac{3(-1 + 4x)e^{2x}}{16} - \frac{4e^{-x}}{3}$$

✓ Solution by Mathematica

Time used: 0.345 (sec). Leaf size: 86

```
DSolve[y''[x]-4*y[x]==3*exp[2*x]+4*Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} \left(e^{4x} \int_1^x \frac{1}{4} e^{-3K[1]} (3e^{K[1]} \exp(2K[1]) + 4) dK[1] + \int_1^x -\frac{1}{4} e^{K[2]} (3e^{K[2]} \exp(2K[2]) + 4) dK[2] + c_1 e^{4x} + c_2 \right)$$

11.4 problem 1(d)

Internal problem ID [6000]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - y' - 2y = x^2 + \cos(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-2*y(x)=x^2+cos(x),y(x), singsol=all)
```

$$y(x) = e^{2x}c_2 + e^{-x}c_1 - \frac{x^2}{2} - \frac{3\cos(x)}{10} - \frac{\sin(x)}{10} + \frac{x}{2} - \frac{3}{4}$$

✓ Solution by Mathematica

Time used: 0.142 (sec). Leaf size: 44

```
DSolve[y''[x]-y'[x]-2*y[x]==x^2+Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{20}(-10x^2 + 10x - 2\sin(x) - 6\cos(x) - 15) + c_1e^{-x} + c_2e^{2x}$$

11.5 problem 1(e)

Internal problem ID [6001]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = x^2 e^{3x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$2)+9*y(x)=x^2*exp(3*x),y(x), singsol=all)
```

$$y(x) = \sin(3x) c_2 + \cos(3x) c_1 + \frac{(3x - 1)^2 e^{3x}}{162}$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 36

```
DSolve[y''[x]+9*y[x]==x^2*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{162} e^{3x} (1 - 3x)^2 + c_1 \cos(3x) + c_2 \sin(3x)$$

11.6 problem 1(f)

Internal problem ID [6002]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = x e^x \cos(2x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 37

```
dsolve(diff(y(x),x$2)+y(x)=x*exp(x)*cos(2*x),y(x), singsol=all)
```

$$y(x) = \sin(x) c_2 + \cos(x) c_1 + \frac{(-5x + 11) e^x \cos(2x)}{50} + \frac{e^x (x - \frac{1}{5}) \sin(2x)}{5}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 45

```
DSolve[y''[x]+y[x]==x*Exp[x]*Cos[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{50} e^x (2(1 - 5x) \sin(2x) + (5x - 11) \cos(2x)) + c_1 \cos(x) + c_2 \sin(x)$$

11.7 problem 1(g)

Internal problem ID [6003]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(g).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + iy' + 2y = 2 \cosh(2x) + e^{-2x}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$2)+I*diff(y(x),x)+2*y(x)=2*cosh(2*x)+exp(-2*x),y(x), singsol=all)
```

$$y(x) = c_2 e^{ix} + e^{-2ix} c_1 + \left(\frac{3}{10} + \frac{i}{10}\right) e^{-2x} + \left(\frac{3}{20} - \frac{i}{20}\right) e^{2x}$$

✓ Solution by Mathematica

Time used: 0.157 (sec). Leaf size: 48

```
DSolve[y''[x]+I*y'[x]+2*y[x]==2*Cosh[2*x]+Exp[-2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{20} e^{-2x} ((3-i)e^{4x} + (6+2i)) + c_1 e^{-2ix} + c_2 e^{ix}$$

11.8 problem 1(h)

Internal problem ID [6004]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(h).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _quadrature]]`

$$y''' = x^2 + e^{-x} \sin(x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 36

```
dsolve(diff(y(x),x$3)=x^2+exp(-x)*sin(x),y(x), singsol=all)
```

$$y(x) = \frac{x^5}{60} + \frac{c_1 x^2}{2} - \frac{\cos(x) e^{-x}}{4} + \frac{\sin(x) e^{-x}}{4} + x c_2 + c_3$$

✓ Solution by Mathematica

Time used: 0.114 (sec). Leaf size: 47

```
DSolve[y'''[x]==x^2+Exp[-x]*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^5}{60} + c_3 x^2 + \frac{1}{4} e^{-x} \sin(x) - \frac{1}{4} e^{-x} \cos(x) + c_2 x + c_1$$

11.9 problem 1(i)

Internal problem ID [6005]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(i).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' + 3y'' + 3y' + y = e^{-x}x^2$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 36

```
dsolve(diff(y(x),x$3)+3*diff(y(x),x$2)+3*diff(y(x),x)+y(x)=x^2*exp(-x),y(x), singsol=all)
```

$$y(x) = \frac{x^5 e^{-x}}{60} + e^{-x}c_1 + c_2 e^{-x}x + c_3 e^{-x}x^2$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 34

```
DSolve[y'''[x]+3*y''[x]+3*y'[x]+y[x]==x^2*Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{60} e^{-x} (x^5 + 60c_3 x^2 + 60c_2 x + 60c_1)$$

12 Chapter 3. Linear equations with variable coefficients. Page 108

12.1	problem 1(c.1)	110
12.2	problem 1(c.2)	111
12.3	problem 2	112

12.1 problem 1(c.1)

Internal problem ID [6006]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 108

Problem number: 1(c.1).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$y'' + \frac{y'}{x} - \frac{y}{x^2} = 0$$

With initial conditions

$$[y(1) = 1, y'(1) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve([diff(y(x),x$2)+1/x*diff(y(x),x)-1/x^2*y(x)=0,y(1) = 1, D(y)(1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{1}{2x} + \frac{x}{2}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 17

```
DSolve[{y'[x]+1/x*y'[x]-1/x^2*y[x]==0,{y[1]==1,y'[1]==0}},y[x],x,IncludeSingularSolutions -
```

$$y(x) \rightarrow \frac{x^2 + 1}{2x}$$

12.2 problem 1(c.2)

Internal problem ID [6007]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 108

Problem number: 1(c.2).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$y'' + \frac{y'}{x} - \frac{y}{x^2} = 0$$

With initial conditions

$$[y(1) = 0, y'(1) = 1]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
dsolve([diff(y(x),x$2)+1/x*diff(y(x),x)-1/x^2*y(x)=0,y(1) = 0, D(y)(1) = 1],y(x), singsol=all)
```

$$y(x) = -\frac{1}{2x} + \frac{x}{2}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 17

```
DSolve[{y'[x]+1/x*y'[x]-1/x^2*y[x]==0,{y[1]==0,y'[1]==1}},y[x],x,IncludeSingularSolutions -
```

$$y(x) \rightarrow \frac{x^2 - 1}{2x}$$

12.3 problem 2

Internal problem ID [6008]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 108

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(3x - 1)^2 y'' + (9x - 3) y' - 9y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve((3*x-1)^2*diff(y(x),x$2)+(9*x-3)*diff(y(x),x)-9*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1}{x - \frac{1}{3}} + \left(x - \frac{1}{3}\right) c_2$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 39

```
DSolve[(3*x-1)^2*y''[x]+(9*x-3)*y'[x]-9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1(-9x^2 + 6x - 2) - 3ic_2x(3x - 2)}{6x - 2}$$

13 Chapter 3. Linear equations with variable coefficients. Page 121

13.1	problem 1(a)	114
13.2	problem 1(b)	115
13.3	problem 1(c)	116
13.4	problem 1(d)	117
13.5	problem 1(e)	118
13.6	problem 1(f)	119
13.7	problem 2	120

13.1 problem 1(a)

Internal problem ID [6009]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 121

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2y'' - 7y'x + 15y = 0$$

Given that one solution of the ode is

$$y_1 = x^3$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 15

```
dsolve([x^2*diff(y(x),x$2)-7*x*diff(y(x),x)+15*y(x)=0,x^3],y(x), singsol=all)
```

$$y(x) = c_2x^5 + c_1x^3$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 18

```
DSolve[x^2*y'[x]-7*x*y'[x]+15*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^3(c_2x^2 + c_1)$$

13.2 problem 1(b)

Internal problem ID [6010]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 121

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - y'x + y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,x],y(x), singsol=all)
```

$$y(x) = c_1 x + c_2 x \ln(x)$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 15

```
DSolve[x^2*y''[x]-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(c_2 \log(x) + c_1)$$

13.3 problem 1(c)

Internal problem ID [6011]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 121

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 4y'x + (4x^2 - 2)y = 0$$

Given that one solution of the ode is

$$y_1 = e^{x^2}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve([diff(y(x),x$2)-4*x*diff(y(x),x)+(4*x^2-2)*y(x)=0,exp(x^2)],y(x), singsol=all)
```

$$y(x) = c_1 e^{x^2} + c_2 x e^{x^2}$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 18

```
DSolve[y''[x]-4*x*y'[x]+(4*x^2-2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x^2} (c_2 x + c_1)$$

13.4 problem 1(d)

Internal problem ID [6012]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 121

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Laguerre]

$$xy'' - y'(1+x) + y = 0$$

Given that one solution of the ode is

$$y_1 = e^x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve([x*diff(y(x),x$2)-(x+1)*diff(y(x),x)+y(x)=0,exp(x)],y(x), singsol=all)
```

$$y(x) = c_1(x+1) + e^x c_2$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 19

```
DSolve[x*y''[x]-(x+1)*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x - c_2(x+1)$$

13.5 problem 1(e)

Internal problem ID [6013]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 121

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Gegenbauer]

$$(-x^2 + 1)y'' - 2y'x + 2y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 26

```
dsolve([(1-x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=0,x],y(x), singsol=all)
```

$$y(x) = c_1x + c_2\left(\frac{\ln(x-1)x}{2} - \frac{\ln(x+1)x}{2} + 1\right)$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 33

```
DSolve[(1-x^2)*y'[x]-2*x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x - \frac{1}{2}c_2(x \log(1-x) - x \log(x+1) + 2)$$

13.6 problem 1(f)

Internal problem ID [6014]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 121

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 2y'x + 2y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve([diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=0,x],y(x), singsol=all)
```

$$y(x) = c_1x + c_2\left(\sqrt{\pi} \operatorname{erfi}(x)x - e^{x^2}\right)$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 43

```
DSolve[y''[x]-2*x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{\pi}c_2\sqrt{x^2}\operatorname{erfi}\left(\sqrt{x^2}\right) + c_2e^{x^2} + 2c_1x$$

13.7 problem 2

Internal problem ID [6015]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 121

Problem number: 2.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$x^3 y''' - 3x^2 y'' + 6y'x - 6y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve([x^3*diff(y(x),x$3)-3*x^2*diff(y(x),x$2)+6*x*diff(y(x),x)-6*y(x)=0,x],y(x), singsol=a
```

$$y(x) = c_2 x^3 + c_1 x^2 + c_3 x$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 19

```
DSolve[x^3*y'''[x]-3*x^2*y''[x]+6*x*y'[x]-6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(x(c_3 x + c_2) + c_1)$$

14 Chapter 3. Linear equations with variable coefficients. Page 124

14.1 problem 1	122
14.2 problem 2	123
14.3 problem 3	124

14.1 problem 1

Internal problem ID [6016]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 124

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x^2 y'' - 2y = 0$$

Given that one solution of the ode is

$$y_1 = x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([x^2*diff(y(x),x$2)-2*y(x)=0,x^2],y(x), singsol=all)
```

$$y(x) = c_1 x^2 + \frac{c_2}{x}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 18

```
DSolve[x^2*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 x^3 + c_1}{x}$$

14.2 problem 2

Internal problem ID [6017]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 124

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - y'x + y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,x],y(x), singsol=all)
```

$$y(x) = c_1 x + c_2 x \ln(x)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 15

```
DSolve[x^2*y''[x]-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(c_2 \log(x) + c_1)$$

14.3 problem 3

Internal problem ID [6018]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 124

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + 4y'x + y(x^2 + 2) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(x^2*diff(y(x),x$2)+4*x*diff(y(x),x)+(2+x^2)*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 \sin(x)}{x^2} + \frac{c_2 \cos(x)}{x^2}$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 37

```
DSolve[x^2*y'[x]+4*x*y'[x]+(2+x^2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2c_1 e^{-ix} - ic_2 e^{ix}}{2x^2}$$

15 Chapter 3. Linear equations with variable coefficients. Page 130

15.1	problem 1(a)	126
15.2	problem 1(b)	127
15.3	problem 1(c)	128
15.4	problem 1(d)	129
15.5	problem 1(e)	130
15.6	problem 2	131
15.7	problem 3	132
15.8	problem 4	133
15.9	problem 5	134
15.10	problem 6	135
15.11	problem 7	136
15.12	problem 8	137

15.1 problem 1(a)

Internal problem ID [6019]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [_Hermite]

$$y'' - y'x + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4\right) y(0) + D(y)(0)x + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 27

```
AsymptoticDSolveValue[y' '[x]-x*y' [x]+y [x]==0,y [x] ,{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^4}{24} - \frac{x^2}{2} + 1\right) + c_2 x$$

15.2 problem 1(b)

Internal problem ID [6020]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 3x^2y' - yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)+3*x^2*diff(y(x),x)-x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{x^3}{6}\right) y(0) + \left(x - \frac{1}{6}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y'[x]+3*x^2*y'[x]-x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{x^4}{6}\right) + c_1 \left(\frac{x^3}{6} + 1\right)$$

15.3 problem 1(c)

Internal problem ID [6021]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y'' - yx^2 = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)-x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{x^4}{12}\right) y(0) + \left(x + \frac{1}{20}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y''[x]-x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{20} + x\right) + c_1 \left(\frac{x^4}{12} + 1\right)$$

15.4 problem 1(d)

Internal problem ID [6022]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + y'x^3 + yx^2 = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)+x^3*diff(y(x),x)+x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{x^4}{12}\right) y(0) + \left(x - \frac{1}{10}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y'[x]+x^3*y'[x]+x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{x^5}{10}\right) + c_1 \left(1 - \frac{x^4}{12}\right)$$

15.5 problem 1(e)

Internal problem ID [6023]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4\right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[y''[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{120} - \frac{x^3}{6} + x \right) + c_1 \left(\frac{x^4}{24} - \frac{x^2}{2} + 1 \right)$$

15.6 problem 2

Internal problem ID [6024]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + (x - 1)^2 y' - (x - 1)y = 0$$

With initial conditions

$$[y(1) = 1, y'(1) = 0]$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
Order:=6;
dsolve([diff(y(x),x$2)+(x-1)^2*diff(y(x),x)-(x-1)*y(x)=0,y(1) = 1, D(y)(1) = 0],y(x),type='s
```

$$y(x) = 1 + \frac{1}{6}(x - 1)^3 + O((x - 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 14

```
AsymptoticDSolveValue[{y''[x]+(x-1)^2*y'[x]-(x-1)*y[x]==0,{y[1]==1,y'[1]==0}},y[x],{x,1,5}]
```

$$y(x) \rightarrow \frac{1}{6}(x - 1)^3 + 1$$

15.7 problem 3

Internal problem ID [6025]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$(x^2 + 1)y'' + y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;  
dsolve([(1+x^2)*diff(y(x),x$2)+y(x)=0,y(0) = 0, D(y)(0) = 1],y(x),type='series',x=0);
```

$$y(x) = x - \frac{1}{6}x^3 + \frac{7}{120}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 19

```
AsymptoticDSolveValue[{(1+x^2)*y'[x]+y[x]==0,{y[0]==0,y'[0]==1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{7x^5}{120} - \frac{x^3}{6} + x$$

15.8 problem 4

Internal problem ID [6026]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + e^x y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
Order:=6;  
dsolve([diff(y(x),x$2)+exp(x)*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x),type='series',x=0);
```

$$y(x) = 1 - \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{40}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 56

```
AsymptoticDSolveValue[{y'[x]+Exp[x]*y[x]==0,{}},y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(-\frac{x^5}{60} - \frac{x^4}{12} - \frac{x^3}{6} + x \right) + c_1 \left(\frac{x^5}{40} - \frac{x^3}{6} - \frac{x^2}{2} + 1 \right)$$

15.9 problem 5

Internal problem ID [6027]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 5.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$y''' - xy = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0, y''(0) = 0]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 14

```
dsolve([diff(y(x),x$3)-x*y(x)=0,y(0) = 1, D(y)(0) = 0, (D@@2)(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = \text{hypergeom} \left(\left[\right], \left[\frac{1}{2}, \frac{3}{4} \right], \frac{x^4}{64} \right)$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 21

```
DSolve[{y'''[x]-x*y[x]==0,{y[0]==1,y'[0]==0,y''[0]==0}},y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow {}_0F_2 \left(; \frac{1}{2}, \frac{3}{4}; \frac{x^4}{64} \right)$$

15.10 problem 6

Internal problem ID [6028]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Gegenbauer]

$$(-x^2 + 1)y'' - 2y'x + \alpha(\alpha + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 101

```
Order:=6;
```

```
dsolve((1-x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+alpha*(alpha+1)*y(x)=0,y(x),type='series',x=0
```

$$y(x) = \left(1 - \frac{\alpha(\alpha + 1)x^2}{2} + \frac{\alpha(\alpha^3 + 2\alpha^2 - 5\alpha - 6)x^4}{24}\right) y(0) \\ + \left(x - \frac{(\alpha^2 + \alpha - 2)x^3}{6} + \frac{(\alpha^4 + 2\alpha^3 - 13\alpha^2 - 14\alpha + 24)x^5}{120}\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 127

```
AsymptoticDSolveValue[(1-x^2)*y''[x]-2*x*y'[x]+\[Alpha]*(\[Alpha]+1)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{1}{60} (-\alpha^2 - \alpha) x^5 - \frac{1}{120} (-\alpha^2 - \alpha) (\alpha^2 + \alpha) x^5 - \frac{1}{10} (\alpha^2 + \alpha) x^5 + \frac{x^5}{5} \right. \\ \left. - \frac{1}{6} (\alpha^2 + \alpha) x^3 + \frac{x^3}{3} + x \right) + c_1 \left(\frac{1}{24} (\alpha^2 + \alpha)^2 x^4 - \frac{1}{4} (\alpha^2 + \alpha) x^4 - \frac{1}{2} (\alpha^2 + \alpha) x^2 + 1 \right)$$

15.11 problem 7

Internal problem ID [6029]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [_Gegenbauer, [_2nd_order, _linear, ‘_with_symmetry_[0,F(x)]’]

$$(-x^2 + 1)y'' - y'x + \alpha^2 y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 33

```
dsolve((1-x^2)*diff(y(x),x$2)-x*diff(y(x),x)+alpha^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \left(x + \sqrt{x^2 - 1}\right)^{-\alpha} + c_2 \left(x + \sqrt{x^2 - 1}\right)^{\alpha}$$

✓ Solution by Mathematica

Time used: 0.089 (sec). Leaf size: 91

```
DSolve[(1-x^2)*y'[x]-x*y'[x]+\[Alpha]^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cosh \left(\frac{1}{2} \alpha \left(\log \left(1 - \frac{x}{\sqrt{x^2 - 1}} \right) - \log \left(\frac{x}{\sqrt{x^2 - 1}} + 1 \right) \right) \right) - i c_2 \sinh \left(\frac{1}{2} \alpha \left(\log \left(1 - \frac{x}{\sqrt{x^2 - 1}} \right) - \log \left(\frac{x}{\sqrt{x^2 - 1}} + 1 \right) \right) \right)$$

15.12 problem 8

Internal problem ID [6030]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 2y'x + 2\alpha y = 0$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$2)-2*x*diff(y(x),x)+2*alpha*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 x \operatorname{KummerM}\left(\frac{1}{2} - \frac{\alpha}{2}, \frac{3}{2}, x^2\right) + c_2 x \operatorname{KummerU}\left(\frac{1}{2} - \frac{\alpha}{2}, \frac{3}{2}, x^2\right)$$

✓ Solution by Mathematica

Time used: 0.062 (sec). Leaf size: 91

```
DSolve[(1-x^2)*y'[x]-x*y'[x]+\[Alpha]^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cosh\left(\frac{1}{2}\alpha\left(\log\left(1 - \frac{x}{\sqrt{x^2-1}}\right) - \log\left(\frac{x}{\sqrt{x^2-1}} + 1\right)\right)\right) - ic_2 \sinh\left(\frac{1}{2}\alpha\left(\log\left(1 - \frac{x}{\sqrt{x^2-1}}\right) - \log\left(\frac{x}{\sqrt{x^2-1}} + 1\right)\right)\right)$$

16 Chapter 4. Linear equations with Regular Singular Points. Page 149

16.1	problem 1(a)	139
16.2	problem 1(b)	140
16.3	problem 1(c)	141
16.4	problem 1(d)	142
16.5	problem 1(e)	143
16.6	problem 2(a)	144
16.7	problem 2(b)	145
16.8	problem 2(c)	146
16.9	problem 2(d)	147

16.1 problem 1(a)

Internal problem ID [6031]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$x^2y'' + 2y'x - 6y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)+2*x*diff(y(x),x)-6*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1x^2 + \frac{c_2}{x^3}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]+2*x*y'[x]-6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2x^5 + c_1}{x^3}$$

16.2 problem 1(b)

Internal problem ID [6032]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$2x^2y'' + y'x - y = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
dsolve(2*x^2*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1}{\sqrt{x}} + xc_2$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 18

```
DSolve[2*x^2*y''[x]+x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{\sqrt{x}} + c_2x$$

16.3 problem 1(c)

Internal problem ID [6033]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$x^2y'' + y'x - 4y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-4*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1x^2 + \frac{c_2}{x^2}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]+x*y'[x]-4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2x^4 + c_1}{x^2}$$

16.4 problem 1(d)

Internal problem ID [6034]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 5y'x + 9y = x^2$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 20

```
dsolve(x^2*diff(y(x),x$2)-5*x*diff(y(x),x)+9*y(x)=x^2,y(x), singsol=all)
```

$$y(x) = c_2 x^3 + x^3 \ln(x) c_1 + x^2$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 22

```
DSolve[x^2*y''[x]-5*x*y'[x]+9*y[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2(c_1 x + 3c_2 x \log(x) + 1)$$

16.5 problem 1(e)

Internal problem ID [6035]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 1(e).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _exact, _linear, _homogeneous]]`

$$x^3 y''' + 2x^2 y'' - y'x + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(x^3*diff(y(x),x$3)+2*x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1}{x} + xc_2 + c_3x \ln(x)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 22

```
DSolve[x^3*y'''[x]+2*x^2*y''[x]-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{x} + c_2x + c_3x \log(x)$$

16.6 problem 2(a)

Internal problem ID [6036]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 2(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y' x + 4y = 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+4*y(x)=1,y(x), singsol=all)
```

$$y(x) = \sin(2 \ln(x)) c_2 + \cos(2 \ln(x)) c_1 + \frac{1}{4}$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 25

```
DSolve[x^2*y''[x]+x*y'[x]+4*y[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(2 \log(x)) + c_2 \sin(2 \log(x)) + \frac{1}{4}$$

16.7 problem 2(b)

Internal problem ID [6037]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 2(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - 3y'x + 5y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(x^2*diff(y(x),x$2)-3*x*diff(y(x),x)+5*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(\ln(x)) x^2 + c_2 \cos(\ln(x)) x^2$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 22

```
DSolve[x^2*y''[x]-3*x*y'[x]+5*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2(c_2 \cos(\log(x)) + c_1 \sin(\log(x)))$$

16.8 problem 2(c)

Internal problem ID [6038]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 2(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' + (-2 - i) x y' + 3i y = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve(x^2*diff(y(x),x$2)-(2+I)*x*diff(y(x),x)+3*I*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 x^3 + c_2 x^i$$

✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 20

```
DSolve[x^2*y''[x]-(2+I)*x*y'[x]+3*I*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 x^i + c_2 x^3$$

16.9 problem 2(d)

Internal problem ID [6039]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 2(d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2y'' + y'x - 4\pi y = x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-4*Pi*y(x)=x,y(x), singsol=all)
```

$$y(x) = x^{-2\sqrt{\pi}}c_2 + x^{2\sqrt{\pi}}c_1 - \frac{x}{4\pi - 1}$$

✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 39

```
DSolve[x^2*y''[x]+x*y'[x]-4*Pi*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2x^{2\sqrt{\pi}} + c_1x^{-2\sqrt{\pi}} + \frac{x}{1 - 4\pi}$$

17 Chapter 4. Linear equations with Regular Singular Points. Page 154

17.1 problem 1(a)	149
17.2 problem 1(b)	150
17.3 problem 1(c)	152
17.4 problem 1(d)	153
17.5 problem 1(e)	155
17.6 problem 1(f)	156
17.7 problem 1(g)	158
17.8 problem 2(b)	160
17.9 problem 2(c)	161
17.10problem 2(d)	162

17.1 problem 1(a)

Internal problem ID [6040]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + (x^2 + x) y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 53

```
Order:=8;
dsolve(x^2*diff(y(x),x$2)+(x+x^2)*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x \left(1 - \frac{1}{3}x + \frac{1}{12}x^2 - \frac{1}{60}x^3 + \frac{1}{360}x^4 - \frac{1}{2520}x^5 + \frac{1}{20160}x^6 - \frac{1}{181440}x^7 + O(x^8) \right) \\ + \frac{c_2 \left(-2 + 2x - x^2 + \frac{1}{3}x^3 - \frac{1}{12}x^4 + \frac{1}{60}x^5 - \frac{1}{360}x^6 + \frac{1}{2520}x^7 + O(x^8) \right)}{x}$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 92

```
AsymptoticDSolveValue[x^2*y'[x]+(x+x^2)*y'[x]-y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^5}{720} - \frac{x^4}{120} + \frac{x^3}{24} - \frac{x^2}{6} + \frac{x}{2} + \frac{1}{x} - 1 \right) \\ + c_2 \left(\frac{x^7}{20160} - \frac{x^6}{2520} + \frac{x^5}{360} - \frac{x^4}{60} + \frac{x^3}{12} - \frac{x^2}{3} + x \right)$$

17.2 problem 1(b)

Internal problem ID [6041]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$3x^2y'' + y'x^6 + 2yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 70

```
Order:=8;
dsolve(3*x^2*diff(y(x),x$2)+x^6*diff(y(x),x)+2*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x \left(1 - \frac{1}{3}x + \frac{1}{27}x^2 - \frac{1}{486}x^3 + \frac{1}{14580}x^4 - \frac{7291}{656100}x^5 + \frac{225991}{41334300}x^6 - \frac{2522341}{3472081200}x^7 + O(x^8) \right) + c_2 \left(\ln(x) \left(-\frac{2}{3}x + \frac{2}{9}x^2 - \frac{2}{81}x^3 + \frac{1}{729}x^4 - \frac{1}{21870}x^5 + \frac{7291}{984150}x^6 - \frac{225991}{62001450}x^7 + O(x^8) \right) + \left(1 - \frac{1}{3}x^2 + \frac{14}{243}x^3 - \frac{35}{8748}x^4 + \frac{101}{656100}x^5 + \frac{69199}{14762250}x^6 + \frac{19882543}{4340101500}x^7 + O(x^8) \right) \right)$$

✓ Solution by Mathematica

Time used: 0.044 (sec). Leaf size: 121

```
AsymptoticDSolveValue[3*x^2*y'[x]+x^6*y'[x]+2*x*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{x(7291x^5 - 45x^4 + 1350x^3 - 24300x^2 + 218700x - 656100) \log(x)}{984150} + \frac{-80332x^6 + 5895x^5 - 158625x^4 + 2430000x^3 - 16402500x^2 + 19683000x + 29524500}{29524500} \right) + c_2 \left(\frac{225991x^7}{41334300} - \frac{7291x^6}{656100} + \frac{x^5}{14580} - \frac{x^4}{486} + \frac{x^3}{27} - \frac{x^2}{3} + x \right)$$

17.3 problem 1(c)

Internal problem ID [6042]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 5y' + 3yx^2 = 0$$

With the expansion point for the power series method at $x = 0$.

X Solution by Maple

```
Order:=8;  
dsolve(x^2*diff(y(x),x$2)-5*diff(y(x),x)+3*x^2*y(x)=0,y(x),type='series',x=0);
```

No solution found

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 106

```
AsymptoticDSolveValue[x^2*y''[x]-5*y'[x]+3*x^2*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{339x^7}{8750} + \frac{49x^6}{1250} + \frac{18x^5}{625} + \frac{3x^4}{50} + \frac{x^3}{5} + 1 \right) + c_2 e^{-5/x} \left(-\frac{302083x^7}{218750} + \frac{5243x^6}{6250} - \frac{357x^5}{625} + \frac{113x^4}{250} - \frac{49x^3}{125} + \frac{6x^2}{25} - \frac{2x}{5} + 1 \right) x^2$$

17.4 problem 1(d)

Internal problem ID [6043]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y''x + 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 70

```
Order:=8;  
dsolve(x*diff(y(x),x$2)+4*y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1 x \left(1 - 2x + \frac{4}{3}x^2 - \frac{4}{9}x^3 + \frac{4}{45}x^4 - \frac{8}{675}x^5 + \frac{16}{14175}x^6 - \frac{8}{99225}x^7 + O(x^8) \right) \\ & + c_2 \left(\ln(x) \left((-4)x + 8x^2 - \frac{16}{3}x^3 + \frac{16}{9}x^4 - \frac{16}{45}x^5 + \frac{32}{675}x^6 - \frac{64}{14175}x^7 + O(x^8) \right) \right. \\ & \left. + \left(1 - 12x^2 + \frac{112}{9}x^3 - \frac{140}{27}x^4 + \frac{808}{675}x^5 - \frac{1792}{10125}x^6 + \frac{9056}{496125}x^7 + O(x^8) \right) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 119

```
AsymptoticDSolveValue[x*y''[x]+4*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{4}{675} x (8x^5 - 60x^4 + 300x^3 - 900x^2 + 1350x - 675) \log(x) \right. \\ \left. + \frac{-2272x^6 + 15720x^5 - 70500x^4 + 180000x^3 - 202500x^2 + 40500x + 10125}{10125} \right) \\ + c_2 \left(\frac{16x^7}{14175} - \frac{8x^6}{675} + \frac{4x^5}{45} - \frac{4x^4}{9} + \frac{4x^3}{3} - 2x^2 + x \right)$$

17.5 problem 1(e)

Internal problem ID [6044]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Gegenbauer]

$$(-x^2 + 1)y'' - 2y'x + 2y = 0$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 49

```
Order:=8;  
dsolve((1-x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=1);
```

$$y(x) = \left(-\frac{5}{2}(x-1) - \frac{3}{8}(x-1)^2 + \frac{1}{12}(x-1)^3 - \frac{5}{192}(x-1)^4 + \frac{3}{320}(x-1)^5 - \frac{7}{1920}(x-1)^6 + \frac{1}{672}(x-1)^7 + O((x-1)^8) \right) c_2 + (1 + (x-1) + O((x-1)^8)) (\ln(x-1) c_2 + c_1)$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 86

```
AsymptoticDSolveValue[(1-x^2)*y''[x]-2*x*y'[x]+2*y[x]==0,y[x],{x,1,7}]
```

$$y(x) \rightarrow c_1 x + c_2 \left(\frac{1}{672}(x-1)^7 - \frac{7(x-1)^6}{1920} + \frac{3}{320}(x-1)^5 - \frac{5}{192}(x-1)^4 + \frac{1}{12}(x-1)^3 - \frac{3}{8}(x-1)^2 - 2(x-1) + \frac{1-x}{2} + x \log(x-1) \right)$$

17.6 problem 1(f)

Internal problem ID [6045]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + x - 2)^2 y'' + 3(x + 2)y' + (x - 1)y = 0$$

With the expansion point for the power series method at $x = -2$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 57

```
Order:=8;
```

```
dsolve((x^2+x-2)^2*diff(y(x),x$2)+3*(x+2)*diff(y(x),x)+(x-1)*y(x)=0,y(x),type='series',x=-2)
```

$y(x)$

$$= c_1 \left(1 - \frac{5}{9}(x + 2) + \frac{23}{324}(x + 2)^2 + \frac{271}{43740}(x + 2)^3 + \frac{10517}{12597120}(x + 2)^4 + \frac{778801}{6235574400}(x + 2)^5 + \frac{16965493}{942818849280}(x + 2)^6 + \dots \right)$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 148

AsymptoticDSolveValue[(x^2+x-2)^2*y'[x]+3*(x+2)*y'[x]+(x-1)*y[x]==0,y[x],{x,-2,7}]

$$y(x) \rightarrow c_1(x+2) \left(-\frac{52991201(x+2)^7}{11727918720000} - \frac{5797423(x+2)^6}{290405606400} - \frac{709507(x+2)^5}{8066822400} \right. \\ \left. - \frac{11093(x+2)^4}{28304640} - \frac{53(x+2)^3}{29484} - \frac{11(x+2)^2}{1260} + \frac{1}{21}(-x-2) + 1 \right) \\ + \frac{c_2 \left(\frac{899971067(x+2)^7}{458981357990400} + \frac{16965493(x+2)^6}{942818849280} + \frac{778801(x+2)^5}{6235574400} + \frac{10517(x+2)^4}{12597120} + \frac{271(x+2)^3}{43740} + \frac{23}{324}(x+2)^2 - \frac{5(x+2)}{9} + 1 \right)}{\sqrt[3]{x+2}}$$

17.7 problem 1(g)

Internal problem ID [6046]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 1(g).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y' \sin(x) + \cos(x) y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.14 (sec). Leaf size: 53

```
Order:=8;  
dsolve(x^2*diff(y(x),x$2)+sin(x)*diff(y(x),x)+cos(x)*y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1 x^{-i} \left(1 + \left(\frac{1}{12} + \frac{i}{24} \right) x^2 + \left(\frac{29}{28800} + \frac{67i}{28800} \right) x^4 \right. \\ & \left. + \left(-\frac{893}{14515200} - \frac{17i}{4838400} \right) x^6 + O(x^8) \right) + c_2 x^i \left(1 + \left(\frac{1}{12} - \frac{i}{24} \right) x^2 \right. \\ & \left. + \left(\frac{29}{28800} - \frac{67i}{28800} \right) x^4 + \left(-\frac{893}{14515200} + \frac{17i}{4838400} \right) x^6 + O(x^8) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 112

```
AsymptoticDSolveValue[x^2*y''[x]+Sin[x]*y'[x]+Cos[x]*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 x^{-i} \left(\left(-\frac{26459}{59222016000} - \frac{12449i}{7402752000} \right) x^8 - \left(\frac{893}{14515200} + \frac{17i}{4838400} \right) x^6 \right. \\ \left. + \left(\frac{29}{28800} + \frac{67i}{28800} \right) x^4 + \left(\frac{1}{12} + \frac{i}{24} \right) x^2 + 1 \right) \\ + c_2 x^i \left(\left(-\frac{26459}{59222016000} + \frac{12449i}{7402752000} \right) x^8 - \left(\frac{893}{14515200} - \frac{17i}{4838400} \right) x^6 \right. \\ \left. + \left(\frac{29}{28800} - \frac{67i}{28800} \right) x^4 + \left(\frac{1}{12} - \frac{i}{24} \right) x^2 + 1 \right)$$

17.8 problem 2(b)

Internal problem ID [6047]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 2(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y' x + \left(-\frac{1}{4} + x^2\right) y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 39

```
Order:=8;
```

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-1/4)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 x \left(1 - \frac{1}{6}x^2 + \frac{1}{120}x^4 - \frac{1}{5040}x^6 + O(x^8)\right) + c_2 \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{720}x^6 + O(x^8)\right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 76

```
AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]+(x^2-1/4)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^{11/2}}{720} + \frac{x^{7/2}}{24} - \frac{x^{3/2}}{2} + \frac{1}{\sqrt{x}}\right) + c_2 \left(-\frac{x^{13/2}}{5040} + \frac{x^{9/2}}{120} - \frac{x^{5/2}}{6} + \sqrt{x}\right)$$

17.9 problem 2(c)

Internal problem ID [6048]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 2(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' + (4x^4 - 5x)y' + y(x^2 + 2) = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 51

```
Order:=8;
```

```
dsolve(4*x^2*diff(y(x),x$2)+(4*x^4-5*x)*diff(y(x),x)+(x^2+2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{\frac{1}{4}} \left(1 - \frac{1}{2}x^2 - \frac{1}{15}x^3 + \frac{1}{72}x^4 + \frac{137}{1950}x^5 + \frac{307}{36720}x^6 - \frac{7169}{3439800}x^7 + O(x^8) \right) + c_2 x^2 \left(1 - \frac{1}{30}x^2 - \frac{8}{57}x^3 + \frac{1}{2760}x^4 + \frac{64}{12825}x^5 + \frac{147181}{9753840}x^6 - \frac{4037}{72268875}x^7 + O(x^8) \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 106

```
AsymptoticDSolveValue[4*x^2*y'[x]+(4*x^4-5*x)*y'[x]+(x^2+2)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{4037x^7}{72268875} + \frac{147181x^6}{9753840} + \frac{64x^5}{12825} + \frac{x^4}{2760} - \frac{8x^3}{57} - \frac{x^2}{30} + 1 \right) x^2 + c_2 \left(-\frac{7169x^7}{3439800} + \frac{307x^6}{36720} + \frac{137x^5}{1950} + \frac{x^4}{72} - \frac{x^3}{15} - \frac{x^2}{2} + 1 \right) \sqrt[4]{x}$$

17.10 problem 2(d)

Internal problem ID [6049]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 2(d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + (-3x^2 + x) y' + e^x y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 85

```
Order:=8;
dsolve(x^2*diff(y(x),x$2)+(x-3*x^2)*diff(y(x),x)+exp(x)*y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1 x^{-i} \left(1 + (1-i)x + \left(\frac{7}{16} - \frac{13i}{16} \right) x^2 + \left(\frac{7}{39} - \frac{395i}{936} \right) x^3 + \left(\frac{2117}{29952} - \frac{5197i}{29952} \right) x^4 \right. \\ & + \left(\frac{5521}{217152} - \frac{642043i}{10857600} \right) x^5 + \left(\frac{782461}{97718400} - \frac{8813057i}{521164800} \right) x^6 \\ & \left. + \left(\frac{1238071931}{580056422400} - \frac{3271304833i}{812078991360} \right) x^7 + O(x^8) \right) \\ & + c_2 x^i \left(1 + (1+i)x + \left(\frac{7}{16} + \frac{13i}{16} \right) x^2 + \left(\frac{7}{39} + \frac{395i}{936} \right) x^3 + \left(\frac{2117}{29952} + \frac{5197i}{29952} \right) x^4 \right. \\ & + \left(\frac{5521}{217152} + \frac{642043i}{10857600} \right) x^5 + \left(\frac{782461}{97718400} + \frac{8813057i}{521164800} \right) x^6 \\ & \left. + \left(\frac{1238071931}{580056422400} + \frac{3271304833i}{812078991360} \right) x^7 + O(x^8) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 122

```
AsymptoticDSolveValue[x^2*y'[x]+(x-3*x^2)*y'[x]+Exp[x]*y[x]==0,y[x],{x,0,7}]
```

$$\begin{aligned} y(x) \rightarrow & \left(\frac{1}{97718400} + \frac{11i}{1563494400} \right) c_1 x^i \left((1302761 + 756800i)x^6 \right. \\ & + (4384656 + 2763936i)x^5 + (12605400 + 8289000i)x^4 \\ & + (31161600 + 19814400i)x^3 + (66096000 + 33955200i)x^2 \\ & \left. + (111974400 + 20736000i)x + (66355200 - 45619200i) \right) \\ & - \left(\frac{11}{1563494400} + \frac{i}{97718400} \right) c_2 x^{-i} \left((756800 + 1302761i)x^6 \right. \\ & + (2763936 + 4384656i)x^5 + (8289000 + 12605400i)x^4 \\ & + (19814400 + 31161600i)x^3 + (33955200 + 66096000i)x^2 \\ & \left. + (20736000 + 111974400i)x - (45619200 - 66355200i) \right) \end{aligned}$$

18 Chapter 4. Linear equations with Regular Singular Points. Page 159

18.1 problem 1(a)	165
18.2 problem 1(b)	166
18.3 problem 2	167

18.1 problem 1(a)

Internal problem ID [6050]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 159

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$3x^2y'' + 5y'x + 3yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 52

```
Order:=8;
dsolve(3*x^2*diff(y(x),x$2)+5*x*diff(y(x),x)+3*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_2 \left(1 - \frac{3}{5}x + \frac{9}{80}x^2 - \frac{9}{880}x^3 + \frac{27}{49280}x^4 - \frac{81}{4188800}x^5 + \frac{81}{167552000}x^6 - \frac{243}{26975872000}x^7 + O(x^8) \right) x^{\frac{2}{3}} + c_1 (1 - 3x + \dots)}{x^{\frac{2}{3}}}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 111

```
AsymptoticDSolveValue[3*x^2*y''[x]+5*x*y'[x]+3*x*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{243x^7}{26975872000} + \frac{81x^6}{167552000} - \frac{81x^5}{4188800} + \frac{27x^4}{49280} - \frac{9x^3}{880} + \frac{9x^2}{80} - \frac{3x}{5} + 1 \right) + \frac{c_2 \left(-\frac{243x^7}{619673600} + \frac{81x^6}{4659200} - \frac{81x^5}{145600} + \frac{27x^4}{2240} - \frac{9x^3}{56} + \frac{9x^2}{8} - 3x + 1 \right)}{x^{2/3}}$$

18.2 problem 1(b)

Internal problem ID [6051]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 159

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Lienard]

$$x^2 y'' + y' x + y x^2 = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 47

```
Order:=8;  
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (\ln(x) c_2 + c_1) \left(1 - \frac{1}{4}x^2 + \frac{1}{64}x^4 - \frac{1}{2304}x^6 + O(x^8) \right) \\ + \left(\frac{1}{4}x^2 - \frac{3}{128}x^4 + \frac{11}{13824}x^6 + O(x^8) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 81

```
AsymptoticDSolveValue[x^2*y'[x]+x*y'[x]+x^2*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^6}{2304} + \frac{x^4}{64} - \frac{x^2}{4} + 1 \right) \\ + c_2 \left(\frac{11x^6}{13824} - \frac{3x^4}{128} + \frac{x^2}{4} + \left(-\frac{x^6}{2304} + \frac{x^4}{64} - \frac{x^2}{4} + 1 \right) \log(x) \right)$$

18.3 problem 2

Internal problem ID [6052]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 159

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' + y' x e^x + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 85

```
Order:=8;
dsolve(x^2*diff(y(x),x$2)+x*exp(x)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1 x^{-i} \left(1 + \left(-\frac{2}{5} + \frac{i}{5} \right) x + \left(\frac{3}{80} + \frac{i}{80} \right) x^2 + \left(\frac{67}{9360} - \frac{9i}{1040} \right) x^3 \right. \\ & \left. + \left(-\frac{103}{149760} - \frac{229i}{149760} \right) x^4 + \left(-\frac{2831}{7238400} + \frac{607i}{4343040} \right) x^5 \right. \\ & \left. + \left(-\frac{59077}{1563494400} + \frac{26063i}{260582400} \right) x^6 + \left(\frac{22952047}{2030197478400} + \frac{8634893i}{580056422400} \right) x^7 \right. \\ & \left. + O(x^8) \right) + c_2 x^i \left(1 + \left(-\frac{2}{5} - \frac{i}{5} \right) x + \left(\frac{3}{80} - \frac{i}{80} \right) x^2 + \left(\frac{67}{9360} + \frac{9i}{1040} \right) x^3 \right. \\ & \left. + \left(-\frac{103}{149760} + \frac{229i}{149760} \right) x^4 + \left(-\frac{2831}{7238400} - \frac{607i}{4343040} \right) x^5 \right. \\ & \left. + \left(-\frac{59077}{1563494400} - \frac{26063i}{260582400} \right) x^6 + \left(\frac{22952047}{2030197478400} - \frac{8634893i}{580056422400} \right) x^7 \right. \\ & \left. + O(x^8) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 122

```
AsymptoticDSolveValue[x^2*y''[x]+x*Exp[x]*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$\begin{aligned} y(x) \rightarrow & \left(\frac{11}{1563494400} + \frac{i}{97718400} \right) c_2 x^{-i} \left((4913 + 7070i)x^6 - (8568 - 32328i)x^5 \right. \\ & - (132840 + 24120i)x^4 - (247680 + 869760i)x^3 + (2540160 - 1918080i)x^2 \\ & \left. - (4976640 - 35665920i)x + (45619200 - 66355200i) \right) \\ & - \left(\frac{1}{97718400} + \frac{11i}{1563494400} \right) c_1 x^i \left((7070 + 4913i)x^6 + (32328 - 8568i)x^5 \right. \\ & - (24120 + 132840i)x^4 - (869760 + 247680i)x^3 - (1918080 - 2540160i)x^2 \\ & \left. + (35665920 - 4976640i)x - (66355200 - 45619200i) \right) \end{aligned}$$

19 Chapter 4. Linear equations with Regular Singular Points. Page 166

19.1	problem 1(i)	170
19.2	problem 1(ii)	171
19.3	problem 1(iii)	173
19.4	problem 3(a)	174
19.5	problem 3(b)	176
19.6	problem 3(c)	177
19.7	problem 3(d)	178
19.8	problem 3(e)	179
19.9	problem 3(f)	180

19.1 problem 1(i)

Internal problem ID [6053]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 1(i).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2y'' + (x^2 + 5x)y' + (x^2 - 2)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 55

```
Order:=8;
```

```
dsolve(2*x^2*diff(y(x),x$2)+(5*x+x^2)*diff(y(x),x)+(x^2-2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_2 x^{\frac{5}{2}} \left(1 - \frac{1}{14}x - \frac{25}{504}x^2 + \frac{197}{33264}x^3 + \frac{1921}{3459456}x^4 - \frac{11653}{103783680}x^5 + \frac{12923}{21171870720}x^6 + \frac{917285}{1126343522304}x^7 + O(x^8) \right) + c_1}{x^2}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 116

```
AsymptoticDSolveValue[2*x^2*y''[x]+(5*x+x^2)*y'[x]+(x^2-2)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{917285x^7}{1126343522304} + \frac{12923x^6}{21171870720} - \frac{11653x^5}{103783680} + \frac{1921x^4}{3459456} + \frac{197x^3}{33264} - \frac{25x^2}{504} - \frac{x}{14} + 1 \right) + \frac{c_2 \left(-\frac{4x^7}{35721} + \frac{101x^6}{45360} - \frac{x^5}{540} - \frac{19x^4}{216} + \frac{2x^3}{9} + \frac{5x^2}{6} - \frac{2x}{3} + 1 \right)}{x^2}$$

19.2 problem 1(ii)

Internal problem ID [6054]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 1(ii).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' - 4y'xe^x + 3\cos(x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 81

```
Order:=8;
```

```
dsolve(4*x^2*diff(y(x),x$2)-4*x*exp(x)*diff(y(x),x)+3*cos(x)*y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & \sqrt{x} \left(x \left(1 + \frac{3}{4}x + \frac{1}{2}x^2 + \frac{103}{384}x^3 + \frac{669}{5120}x^4 + \frac{54731}{921600}x^5 + \frac{123443}{4838400}x^6 \right. \right. \\ & \left. \left. + \frac{30273113}{2890137600}x^7 + O(x^8) \right) c_1 \right. \\ & + c_2 \left(\ln(x) \left(\frac{1}{2}x + \frac{3}{8}x^2 + \frac{1}{4}x^3 + \frac{103}{768}x^4 + \frac{669}{10240}x^5 + \frac{54731}{1843200}x^6 + \frac{123443}{9676800}x^7 + O(x^8) \right) \right. \\ & \left. \left. + \left(1 + x + \frac{3}{4}x^2 + \frac{59}{144}x^3 + \frac{5701}{27648}x^4 + \frac{17519}{184320}x^5 + \frac{6852157}{165888000}x^6 + \frac{417496453}{24385536000}x^7 + O(x^8) \right) \right) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.146 (sec). Leaf size: 146

```
AsymptoticDSolveValue[4*x^2*y'[x]-4*x*Exp[x]*y'[x]+3*Cos[x]*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(\frac{123443x^{15/2}}{4838400} + \frac{54731x^{13/2}}{921600} + \frac{669x^{11/2}}{5120} + \frac{103x^{9/2}}{384} + \frac{x^{7/2}}{2} + \frac{3x^{5/2}}{4} + x^{3/2} \right) + c_1 \left(\frac{(54731x^5 + 120420x^4 + 247200x^3 + 460800x^2 + 691200x + 921600) x^{3/2} \log(x)}{1843200} + \frac{(192636x^5 + 120420x^4 + 247200x^3 + 460800x^2 + 691200x + 921600) x^{3/2}}{1843200} \right)$$

19.3 problem 1(iii)

Internal problem ID [6055]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 1(iii).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(-x^2 + 1)x^2y'' + 3(x^2 + x)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 81

```
Order:=8;
```

```
dsolve((1-x^2)*x^2*diff(y(x),x$2)+3*(x+x^2)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{(\ln(x)c_2 + c_1) \left(1 + 3x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{16}x^4 - \frac{43}{1200}x^5 + \frac{161}{7200}x^6 - \frac{1837}{117600}x^7 + O(x^8)\right) + \left((-9)x - \frac{7}{2}x^2 + \frac{7}{9}x^3\right)}{x}$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 84

```
AsymptoticDSolveValue[(1-x^2)*y''[x]+3*(x+x^2)*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(\frac{53x^7}{630} + \frac{5x^6}{24} + \frac{2x^5}{15} - \frac{x^4}{4} - \frac{2x^3}{3} + x \right) + c_1 \left(-\frac{19x^7}{420} - \frac{x^6}{144} + \frac{3x^5}{20} + \frac{5x^4}{24} - \frac{x^2}{2} + 1 \right)$$

19.4 problem 3(a)

Internal problem ID [6056]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 3(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + 3y'x + (1+x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 81

```
Order:=8;  
dsolve(x^2*diff(y(x),x$2)+3*x*diff(y(x),x)+(1+x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{(\ln(x) c_2 + c_1) \left(1 - x + \frac{1}{4}x^2 - \frac{1}{36}x^3 + \frac{1}{576}x^4 - \frac{1}{14400}x^5 + \frac{1}{518400}x^6 - \frac{1}{25401600}x^7 + O(x^8)\right) + \left(2x - \frac{3}{4}x^2 + \dots\right)}{x}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 164

```
AsymptoticDSolveValue[x^2*y''[x]+3*x*y'[x]+(1+x)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow \frac{c_1 \left(-\frac{x^7}{25401600} + \frac{x^6}{518400} - \frac{x^5}{14400} + \frac{x^4}{576} - \frac{x^3}{36} + \frac{x^2}{4} - x + 1 \right)}{x} + c_2 \left(\frac{\frac{121x^7}{592704000} - \frac{49x^6}{5184000} + \frac{137x^5}{432000} - \frac{25x^4}{3456} + \frac{11x^3}{108} - \frac{3x^2}{4} + 2x}{x} + \frac{\left(-\frac{x^7}{25401600} + \frac{x^6}{518400} - \frac{x^5}{14400} + \frac{x^4}{576} - \frac{x^3}{36} + \frac{x^2}{4} - x + 1 \right) \log(x)}{x} \right)$$

19.5 problem 3(b)

Internal problem ID [6057]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 3(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2y'' + 2x^2y' - 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 53

```
Order:=8;  
dsolve(x^2*diff(y(x),x$2)+2*x^2*diff(y(x),x)-2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1x^2\left(1 - x + \frac{3}{5}x^2 - \frac{4}{15}x^3 + \frac{2}{21}x^4 - \frac{1}{35}x^5 + \frac{1}{135}x^6 - \frac{8}{4725}x^7 + O(x^8)\right) \\ + \frac{c_2(12 - 12x + 8x^3 - 8x^4 + \frac{24}{5}x^5 - \frac{32}{15}x^6 + \frac{16}{21}x^7 + O(x^8))}{x}$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 87

```
AsymptoticDSolveValue[x^2*y''[x]+2*x^2*y'[x]-2*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1\left(-\frac{8x^5}{45} + \frac{2x^4}{5} - \frac{2x^3}{3} + \frac{2x^2}{3} + \frac{1}{x} - 1\right) + c_2\left(\frac{x^8}{135} - \frac{x^7}{35} + \frac{2x^6}{21} - \frac{4x^5}{15} + \frac{3x^4}{5} - x^3 + x^2\right)$$

19.6 problem 3(c)

Internal problem ID [6058]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 3(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + 5y'x + (-x^3 + 3)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
Order:=8;  
dsolve(x^2*diff(y(x),x$2)+5*x*diff(y(x),x)+(3-x^3)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 \left(1 + \frac{1}{15}x^3 + \frac{1}{720}x^6 + O(x^8)\right)}{x} + \frac{c_2 \left(-2 - \frac{2}{3}x^3 - \frac{1}{36}x^6 + O(x^8)\right)}{x^3}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 40

```
AsymptoticDSolveValue[x^2*y'[x]+5*x*y'[x]+(3-3*x^3)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^3}{8} + \frac{1}{x^3} + 1 \right) + c_2 \left(\frac{x^5}{80} + \frac{x^2}{5} + \frac{1}{x} \right)$$

19.7 problem 3(d)

Internal problem ID [6059]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 3(d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 2x(1+x)y' + 2(1+x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 53

```
Order:=8;  
dsolve(x^2*diff(y(x),x$2)-2*x*(x+1)*diff(y(x),x)+2*(x+1)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^2 \left(1 + x + \frac{2}{3}x^2 + \frac{1}{3}x^3 + \frac{2}{15}x^4 + \frac{2}{45}x^5 + \frac{4}{315}x^6 + \frac{1}{315}x^7 + O(x^8) \right) \\ + c_2 x \left(1 + 2x + 2x^2 + \frac{4}{3}x^3 + \frac{2}{3}x^4 + \frac{4}{15}x^5 + \frac{4}{45}x^6 + \frac{8}{315}x^7 + O(x^8) \right)$$

✓ Solution by Mathematica

Time used: 0.086 (sec). Leaf size: 92

```
AsymptoticDSolveValue[x^2*y'[x]-2*x*(x+1)*y'[x]+2*(1+x)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{4x^7}{45} + \frac{4x^6}{15} + \frac{2x^5}{3} + \frac{4x^4}{3} + 2x^3 + 2x^2 + x \right) \\ + c_2 \left(\frac{4x^8}{315} + \frac{2x^7}{45} + \frac{2x^6}{15} + \frac{x^5}{3} + \frac{2x^4}{3} + x^3 + x^2 \right)$$

19.8 problem 3(e)

Internal problem ID [6060]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 3(e).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Bessel]

$$x^2 y'' + y'x + y(x^2 - 1) = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 53

```
Order:=8;
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-1)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 x^2 \left(1 - \frac{1}{8}x^2 + \frac{1}{192}x^4 - \frac{1}{9216}x^6 + O(x^8)\right) + c_2 (\ln(x) \left(x^2 - \frac{1}{8}x^4 + \frac{1}{192}x^6 + O(x^8)\right) + \left(-2 + \frac{3}{32}x^4 - \frac{7}{1152}x^6\right))}{x}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 75

```
AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]+(x^2-1)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(-\frac{x^7}{9216} + \frac{x^5}{192} - \frac{x^3}{8} + x \right) + c_1 \left(\frac{5x^6 - 90x^4 + 288x^2 + 1152}{1152x} - \frac{1}{384}x(x^4 - 24x^2 + 192) \log(x) \right)$$

19.9 problem 3(f)

Internal problem ID [6061]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 3(f).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`, `[_2nd_order, _linear,`

$$x^2 y'' - 2x^2 y' + (-2 + 4x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 55

```
Order:=8;
dsolve(x^2*diff(y(x),x$2)-2*x^2*diff(y(x),x)+(4*x-2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^2 (1 + O(x^8)) + \frac{c_2 (\ln(x) ((-48)x^3 + O(x^8)) + (12 + 36x + 72x^2 + 88x^3 - 24x^4 - \frac{24}{5}x^5 - \frac{16}{15}x^6 - \frac{8}{35}x^7 + O(x^8)))}{x}$$

✓ Solution by Mathematica

Time used: 0.1 (sec). Leaf size: 58

```
AsymptoticDSolveValue[x^2*y''[x]-2*x^2*y'[x]+(4*x-2)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 x^2 + c_1 \left(-4x^2 \log(x) - \frac{4x^6 + 18x^5 + 90x^4 - 390x^3 - 270x^2 - 135x - 45}{45x} \right)$$

**20 Chapter 4. Linear equations with Regular
Singular Points. Page 182**

20.1 problem 4 182

20.1 problem 4

Internal problem ID [6062]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 182

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Gegenbauer]

$$(-x^2 + 1)y'' - 2y'x + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
Order:=8;  
dsolve((1-x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x^2 - \frac{1}{3}x^4 - \frac{1}{5}x^6\right)y(0) + D(y)(0)x + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 32

```
AsymptoticDSolveValue[(1-x^2)*y'[x]-2*x*y'[x]+2*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^6}{5} - \frac{x^4}{3} - x^2 + 1\right) + c_2 x$$

21 Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

21.1 problem 1(a)	184
21.2 problem 1(b)	185
21.3 problem 1(c)	186
21.4 problem 1(d)	189
21.5 problem 1(e)	190
21.6 problem 2(a)	191
21.7 problem 3(a)	192
21.8 problem 3(b)	193
21.9 problem 4(a)	194
21.10 problem 4(b)	195
21.11 problem 4(c)	196
21.12 problem 4(d)	197
21.13 problem 5(a)	198
21.14 problem 5(b)	199
21.15 problem 5(c)	200
21.16 problem 6(b)	201

21.1 problem 1(a)

Internal problem ID [6063]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - yx^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=x^2*y(x),y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{x^3}{3}}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 22

```
DSolve[y'[x]==x^2*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{\frac{x^3}{3}}$$

$$y(x) \rightarrow 0$$

21.2 problem 1(b)

Internal problem ID [6064]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 1(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$yy' = x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(y(x)*diff(y(x),x)=x,y(x), singsol=all)
```

$$y(x) = \sqrt{x^2 + c_1}$$

$$y(x) = -\sqrt{x^2 + c_1}$$

✓ Solution by Mathematica

Time used: 0.081 (sec). Leaf size: 35

```
DSolve[y[x]*y'[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{x^2 + 2c_1}$$

$$y(x) \rightarrow \sqrt{x^2 + 2c_1}$$

21.3 problem 1(c)

Internal problem ID [6065]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 1(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$y' - \frac{x^2 + x}{y - y^2} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 720

```
dsolve(diff(y(x),x)=(x+x^2)/(y(x)-y(x)^2),y(x), singsol=all)
```

$y(x)$

$$= \frac{\left(1 - 4x^3 - 6x^2 - 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}{2} + \frac{1}{2\left(1 - 4x^3 - 6x^2 - 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}} + \frac{1}{2}$$

$y(x) =$

$$-\frac{\left(1 - 4x^3 - 6x^2 - 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}{4} - \frac{1}{4\left(1 - 4x^3 - 6x^2 - 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}} + \frac{1}{2} + \frac{i\sqrt{3}\left(\frac{\left(1 - 4x^3 - 6x^2 - 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}{2} - \frac{1}{2\left(1 - 4x^3 - 6x^2 - 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}\right)}{2}$$

$y(x) =$

$$-\frac{\left(1 - 4x^3 - 6x^2 - 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}{4} - \frac{1}{4\left(1 - 4x^3 - 6x^2 - 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}} + \frac{1}{2} + \frac{i\sqrt{3}\left(\frac{\left(1 - 4x^3 - 6x^2 - 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}{2} - \frac{1}{2\left(1 - 4x^3 - 6x^2 - 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}\right)}{2}$$

✓ Solution by Mathematica

Time used: 4.147 (sec). Leaf size: 346

`DSolve[y'[x]==(x+x^2)/(y[x]-y[x]^2),y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{1}{2} \left(\sqrt[3]{-4x^3 - 6x^2 + \sqrt{-1 + (-4x^3 - 6x^2 + 1 + 12c_1)^2 + 1 + 12c_1}} + \frac{1}{\sqrt[3]{-4x^3 - 6x^2 + \sqrt{-1 + (-4x^3 - 6x^2 + 1 + 12c_1)^2 + 1 + 12c_1}} + 1} \right)$$

$$y(x) \rightarrow \frac{1}{8} \left(2i(\sqrt{3} + i) \sqrt[3]{-4x^3 - 6x^2 + \sqrt{-1 + (-4x^3 - 6x^2 + 1 + 12c_1)^2 + 1 + 12c_1}} + \frac{-2 - 2i\sqrt{3}}{\sqrt[3]{-4x^3 - 6x^2 + \sqrt{-1 + (-4x^3 - 6x^2 + 1 + 12c_1)^2 + 1 + 12c_1}} + 4} \right)$$

$$y(x) \rightarrow \frac{1}{8} \left(-2(1 + i\sqrt{3}) \sqrt[3]{-4x^3 - 6x^2 + \sqrt{-1 + (-4x^3 - 6x^2 + 1 + 12c_1)^2 + 1 + 12c_1}} + \frac{2i(\sqrt{3} + i)}{\sqrt[3]{-4x^3 - 6x^2 + \sqrt{-1 + (-4x^3 - 6x^2 + 1 + 12c_1)^2 + 1 + 12c_1}} + 4} \right)$$

21.4 problem 1(d)

Internal problem ID [6066]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 1(d).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{e^{x-y}}{1 + e^x} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=exp(x-y(x))/(1+exp(x)),y(x), singsol=all)
```

$$y(x) = \ln(\ln(e^x + 1) + c_1)$$

✓ Solution by Mathematica

Time used: 0.465 (sec). Leaf size: 15

```
DSolve[y'[x]==Exp[x-y[x]]/(1+Exp[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(\log(e^x + 1) + c_1)$$

21.5 problem 1(e)

Internal problem ID [6067]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 1(e).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - x^2 y^2 = -4x^2$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 28

```
dsolve(diff(y(x),x)=x^2*y(x)^2-4*x^2,y(x), singsol=all)
```

$$y(x) = -\frac{2\left(e^{\frac{4x^3}{3}}c_1 + 1\right)}{-1 + e^{\frac{4x^3}{3}}c_1}$$

✓ Solution by Mathematica

Time used: 0.258 (sec). Leaf size: 52

```
DSolve[y'[x]==x^2*y[x]^2-4*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2 - 2e^{\frac{4x^3}{3} + 4c_1}}{1 + e^{\frac{4x^3}{3} + 4c_1}}$$

$$y(x) \rightarrow -2$$

$$y(x) \rightarrow 2$$

21.6 problem 2(a)

Internal problem ID [6068]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$-y^2 + y' = 0$$

With initial conditions

$$[y(x_0) = y_0]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 18

```
dsolve([diff(y(x),x)=y(x)^2,y(x__0) = y__0],y(x), singsol=all)
```

$$y(x) = -\frac{y_0}{-1 + (x - x_0) y_0}$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 16

```
DSolve[{y'[x]==x^2*y[x],{y[x0]==y0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow y_0 e^{x^2(x-x_0)}$$

21.7 problem 3(a)

Internal problem ID [6069]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 3(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y' - 2\sqrt{y} = 0$$

With initial conditions

$$[y(x_0) = y_0]$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 28

```
dsolve([diff(y(x),x)=2*sqrt(y(x)),y(x_0) = y_0],y(x), singsol=all)
```

$$y(x) = (2x - 2x_0)\sqrt{y_0} + x^2 - 2xx_0 + x_0^2 + y_0$$

✓ Solution by Mathematica

Time used: 0.108 (sec). Leaf size: 33

```
DSolve[{y'[x]==2*Sqrt[y[x]],{y[x0]==y0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x - x_0 + \sqrt{y_0})^2$$

$$y(x) \rightarrow (-x + x_0 + \sqrt{y_0})^2$$

21.8 problem 3(b)

Internal problem ID [6070]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 3(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_quadrature]`

$$y' - 2\sqrt{y} = 0$$

With initial conditions

$$[y(x_0) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve([diff(y(x),x)=2*sqrt(y(x)),y(x__0) = 0],y(x), singsol=all)
```

$$y(x) = 0$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 6

```
DSolve[{y'[x]==2*Sqrt[y[x]],{y[x0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

21.9 problem 4(a)

Internal problem ID [6071]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 4(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{x+y}{x-y} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
dsolve(diff(y(x),x)=(x+y(x))/(x-y(x)),y(x), singsol=all)
```

$$y(x) = \tan \left(\text{RootOf} \left(-2_Z + \ln \left(\frac{1}{\cos(_Z)^2} \right) + 2 \ln(x) + 2c_1 \right) \right) x$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 36

```
DSolve[y'[x]==(x+y[x])/(x-y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{1}{2} \log \left(\frac{y(x)^2}{x^2} + 1 \right) - \arctan \left(\frac{y(x)}{x} \right) = -\log(x) + c_1, y(x) \right]$$

21.10 problem 4(b)

Internal problem ID [6072]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 4(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{y^2}{x^2 + yx} = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 20

```
dsolve(diff(y(x),x)=y(x)^2/(x*y(x)+x^2),y(x), singsol=all)
```

$$y(x) = e^{-\text{LambertW}\left(\frac{e^{-c_1}}{x}\right) - c_1}$$

✓ Solution by Mathematica

Time used: 2.317 (sec). Leaf size: 21

```
DSolve[y'[x]==y[x]^2/(x*y[x]+x^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow xW\left(\frac{e^{c_1}}{x}\right)$$

$$y(x) \rightarrow 0$$

21.11 problem 4(c)

Internal problem ID [6073]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 4(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Riccati]`

$$y' - \frac{x^2 + yx + y^2}{x^2} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve(diff(y(x),x)=(x^2+x*y(x)+y(x)^2)/x^2,y(x), singsol=all)
```

$$y(x) = \tan(\ln(x) + c_1)x$$

✓ Solution by Mathematica

Time used: 0.188 (sec). Leaf size: 13

```
DSolve[y'[x]==(x^2+x*y[x]+y[x]^2)/x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \tan(\log(x) + c_1)$$

21.12 problem 4(d)

Internal problem ID [6074]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 4(d).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _dAlembert]`

$$y' - \frac{y + x e^{-\frac{2y}{x}}}{x} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)=(y(x)+x*exp(-2*y(x)/x))/x,y(x), singsol=all)
```

$$y(x) = \frac{\ln(2 \ln(x) + 2c_1) x}{2}$$

✓ Solution by Mathematica

Time used: 0.412 (sec). Leaf size: 18

```
DSolve[y'[x]==(y[x]+x*Exp[-2*y[x]/x])/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} x \log(2(\log(x) + c_1))$$

21.13 problem 5(a)

Internal problem ID [6075]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 5(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{x - y + 2}{y + x - 1} = 0$$

✓ Solution by Maple

Time used: 0.203 (sec). Leaf size: 35

```
dsolve(diff(y(x),x)=(x-y(x)+2)/(x+y(x)-1),y(x), singsol=all)
```

$$y(x) = \frac{3}{2} - \frac{(2x + 1)c_1 + \sqrt{2(2x + 1)^2 c_1^2 + 1}}{2c_1}$$

✓ Solution by Mathematica

Time used: 0.154 (sec). Leaf size: 53

```
DSolve[y'[x]==(x-y[x]+2)/(x+y[x]-1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{2x^2 + 2x + 1 + c_1} - x + 1$$

$$y(x) \rightarrow \sqrt{2x^2 + 2x + 1 + c_1} - x + 1$$

21.14 problem 5(b)

Internal problem ID [6076]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 5(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{2x + 3y + 1}{x - 2y - 1} = 0$$

✓ Solution by Maple

Time used: 0.219 (sec). Leaf size: 59

```
dsolve(diff(y(x),x)=(2*x+3*y(x)+1)/(x-2*y(x)-1),y(x), singsol=all)
```

$$y(x) = -\frac{5}{14} - \frac{x}{2} + \frac{\sqrt{3}(7x-1) \tan\left(\text{RootOf}\left(\sqrt{3} \ln\left(\frac{3(7x-1)^2}{4} + \frac{3(7x-1)^2 \tan(-Z)^2}{4}\right) + 2\sqrt{3}c_1 - 4_Z\right)\right)}{14}$$

✓ Solution by Mathematica

Time used: 0.12 (sec). Leaf size: 85

```
DSolve[y'[x]==(2*x+3*y[x]+1)/(x-2*y[x]-1),y[x],x,IncludeSingularSolutions->True]
```

$$\text{Solve}\left[32\sqrt{3} \arctan\left(\frac{4y(x) + 5x + 1}{\sqrt{3}(-2y(x) + x - 1)}\right) = 3\left(8 \log\left(\frac{4(7x^2 + 7y(x)^2 + (7x + 5)y(x) + x + 1)}{(1 - 7x)^2}\right) + 16 \log(7x - 1) + 7c_1\right), y(x)\right]$$

21.15 problem 5(c)

Internal problem ID [6077]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 5(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{y + x + 1}{2x + 2y - 1} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve(diff(y(x),x)=(x+y(x)+1)/(2*x+2*y(x)-1),y(x), singsol=all)
```

$$y(x) = e^{-\text{LambertW}(-2e^{-3x}e^{3c_1})-3x+3c_1} - x$$

✓ Solution by Mathematica

Time used: 4.2 (sec). Leaf size: 32

```
DSolve[y'[x]==(x+y[x]+1)/(2*x+2*y[x]-1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x - \frac{1}{2}W(-e^{-3x-1+c_1})$$

$$y(x) \rightarrow -x$$

21.16 problem 6(b)

Internal problem ID [6078]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 6(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _rational, _Riccati]`

$$y' - \frac{(y+x-1)^2}{2(x+2)^2} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)=1/2*((x+y(x)-1)/(x+2))^2,y(x), singsol=all)
```

$$y(x) = 3 + \tan\left(\frac{\ln(x+2)}{2} + \frac{c_1}{2}\right)(x+2)$$

✓ Solution by Mathematica

Time used: 0.411 (sec). Leaf size: 99

```
DSolve[y'[x]==1/2*((x+y[x]-1)/(x+2))^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2^i(x+2)^i x + (2+3i)2^i(x+2)^i - 2ic_1 x - (6+4i)c_1}{i2^i(x+2)^i - 2c_1}$$

$$y(x) \rightarrow ix + (3+2i)$$

$$y(x) \rightarrow ix + (3+2i)$$

22 Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

22.1	problem 1(a)	203
22.2	problem 1(b)	205
22.3	problem 1(c)	206
22.4	problem 1(d)	207
22.5	problem 1(e)	209
22.6	problem 1(f)	210
22.7	problem 1(g)	211
22.8	problem 1(h)	212
22.9	problem 2(a)	213
22.10	problem 2(b)	215
22.11	problem 2(c)	216
22.12	problem 2(d)	218

22.1 problem 1(a)

Internal problem ID [6079]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _exact, _rational, _dAlembert]`

$$2yx + (3y^2 + x^2)y' = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 257

```
dsolve(2*x*y(x)+(x^2+3*y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\frac{\left(108+12\sqrt{12c_1^3x^6+81}\right)^{\frac{1}{3}}}{6} - \frac{2x^2c_1}{\left(108+12\sqrt{12c_1^3x^6+81}\right)^{\frac{1}{3}}}}{\sqrt{c_1}}$$

$y(x)$

$$= \frac{-\frac{\left(108+12\sqrt{12c_1^3x^6+81}\right)^{\frac{1}{3}}}{12} + \frac{x^2c_1}{\left(108+12\sqrt{12c_1^3x^6+81}\right)^{\frac{1}{3}}} - \frac{i\sqrt{3}\left(\frac{\left(108+12\sqrt{12c_1^3x^6+81}\right)^{\frac{1}{3}}}{6} + \frac{2x^2c_1}{\left(108+12\sqrt{12c_1^3x^6+81}\right)^{\frac{1}{3}}}\right)}{2}}{\sqrt{c_1}}$$

$y(x)$

$$= \frac{-\frac{\left(108+12\sqrt{12c_1^3x^6+81}\right)^{\frac{1}{3}}}{12} + \frac{x^2c_1}{\left(108+12\sqrt{12c_1^3x^6+81}\right)^{\frac{1}{3}}} + \frac{i\sqrt{3}\left(\frac{\left(108+12\sqrt{12c_1^3x^6+81}\right)^{\frac{1}{3}}}{6} + \frac{2x^2c_1}{\left(108+12\sqrt{12c_1^3x^6+81}\right)^{\frac{1}{3}}}\right)}{2}}{\sqrt{c_1}}$$

✓ Solution by Mathematica

Time used: 27.686 (sec). Leaf size: 442

```
DSolve[2*x*y[x]+(x^2+3*y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-2\sqrt[3]{3}x^2 + \sqrt[3]{2}(\sqrt{12x^6 + 81e^{2c_1}} + 9e^{c_1})^{2/3}}{6^{2/3}\sqrt[3]{\sqrt{12x^6 + 81e^{2c_1}} + 9e^{c_1}}}$$

$$y(x) \rightarrow \frac{i2^{2/3}\sqrt[3]{3}(\sqrt{3} + i)(\sqrt{12x^6 + 81e^{2c_1}} + 9e^{c_1})^{2/3} + 2\sqrt[3]{2}\sqrt[6]{3}(\sqrt{3} + 3i)x^2}{12\sqrt[3]{\sqrt{12x^6 + 81e^{2c_1}} + 9e^{c_1}}}$$

$$y(x) \rightarrow \frac{2^{2/3}\sqrt[3]{3}(-1 - i\sqrt{3})(\sqrt{12x^6 + 81e^{2c_1}} + 9e^{c_1})^{2/3} + 2\sqrt[3]{2}\sqrt[6]{3}(\sqrt{3} - 3i)x^2}{12\sqrt[3]{\sqrt{12x^6 + 81e^{2c_1}} + 9e^{c_1}}}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow \frac{\sqrt[3]{x^6} - x^2}{\sqrt{3}\sqrt[6]{x^6}}$$

$$y(x) \rightarrow \frac{(\sqrt{3} - 3i)x^2 - (\sqrt{3} + 3i)\sqrt[3]{x^6}}{6\sqrt[6]{x^6}}$$

$$y(x) \rightarrow \frac{(\sqrt{3} + 3i)x^2 - (\sqrt{3} - 3i)\sqrt[3]{x^6}}{6\sqrt[6]{x^6}}$$

22.2 problem 1(b)

Internal problem ID [6080]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$yx + (x + y)y' = -x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve((x^2+x*y(x))+(x+y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -x$$

$$y(x) = -\frac{x^2}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.132 (sec). Leaf size: 53

```
DSolve[(x^2+y[x])+(x+y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x - \sqrt{-\frac{2x^3}{3} + x^2 + c_1}$$

$$y(x) \rightarrow -x + \sqrt{-\frac{2x^3}{3} + x^2 + c_1}$$

22.3 problem 1(c)

Internal problem ID [6081]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$e^y(1+y)y' = -e^x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve(exp(x)+(exp(y(x))*(y(x)+1))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \text{LambertW}(-c_1 - e^x)$$

✓ Solution by Mathematica

Time used: 60.161 (sec). Leaf size: 14

```
DSolve[Exp[x]+(Exp[y[x]]*(y[x]+1))*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow W(-e^x + c_1)$$

22.4 problem 1(d)

Internal problem ID [6082]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(d).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$\cos(x) \cos(y)^2 - \sin(x) \sin(2y) y' = 0$$

✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 25

```
dsolve(cos(x)*cos(y(x))^2-sin(x)*sin(2*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \arccos\left(\frac{1}{\sqrt{c_1 \sin(x)}}\right)$$

$$y(x) = \pi - \arccos\left(\frac{1}{\sqrt{c_1 \sin(x)}}\right)$$

✓ Solution by Mathematica

Time used: 6.536 (sec). Leaf size: 73

```
DSolve[Cos[x]*Cos[y[x]]^2-Sin[x]*Sin[2*y[x]]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\pi}{2}$$

$$y(x) \rightarrow \frac{\pi}{2}$$

$$y(x) \rightarrow -\arccos\left(-\frac{c_1}{4\sqrt{\sin(x)}}\right)$$

$$y(x) \rightarrow \arccos\left(-\frac{c_1}{4\sqrt{\sin(x)}}\right)$$

$$y(x) \rightarrow -\frac{\pi}{2}$$

$$y(x) \rightarrow \frac{\pi}{2}$$

22.5 problem 1(e)

Internal problem ID [6083]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(e).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y^3 x^2 - y^2 x^3 y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(x^2*y(x)^3-x^3*y(x)^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1 x$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 19

```
DSolve[x^2*y[x]^3-x^3*y[x]^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow c_1 x$$

$$y(x) \rightarrow 0$$

22.6 problem 1(f)

Internal problem ID [6084]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(f).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _exact, _rational, [_Abel, '2nd ty`

$$y + (x - y)y' = -x$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 49

```
dsolve((x+y(x))+(x-y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 x - \sqrt{2c_1^2 x^2 + 1}}{c_1}$$

$$y(x) = \frac{c_1 x + \sqrt{2c_1^2 x^2 + 1}}{c_1}$$

✓ Solution by Mathematica

Time used: 0.449 (sec). Leaf size: 86

```
DSolve[(x+y[x])+(x-y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x - \sqrt{2x^2 + e^{2c_1}}$$

$$y(x) \rightarrow x + \sqrt{2x^2 + e^{2c_1}}$$

$$y(x) \rightarrow x - \sqrt{2}\sqrt{x^2}$$

$$y(x) \rightarrow \sqrt{2}\sqrt{x^2} + x$$

22.7 problem 1(g)

Internal problem ID [6085]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(g).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_exact]

$$2e^{2x}y + 2x \cos(y) + (e^{2x} - x^2 \sin(y))y' = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 19

```
dsolve((2*y(x)*exp(2*x)+2*x*cos(y(x)))+(exp(2*x)-x^2*sin(y(x)))*diff(y(x),x)=0,y(x), singular
```

$$\cos(y(x))x^2 + y(x)e^{2x} + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.414 (sec). Leaf size: 30

```
DSolve[(2*y[x]*Exp[2*x]+2*x*Cos[y[x]])+(Exp[2*x]-x^2*Sin[y[x]])*y'[x]==0,y[x],x,IncludeSingu
```

$$\text{Solve} \left[2 \left(\frac{1}{2} x^2 \cos(y(x)) + \frac{1}{2} e^{2x} y(x) \right) = c_1, y(x) \right]$$

22.8 problem 1(h)

Internal problem ID [6086]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(h).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y + y'x = -3 \ln(x) x^2 - x^2$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve((3*x^2*ln(x)+x^2+y(x))+x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-x^3 \ln(x) + c_1}{x}$$

✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 19

```
DSolve[(3*x^2*Log[x]+x^2+y[x])+x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-x^3 \log(x) + c_1}{x}$$

22.9 problem 2(a)

Internal problem ID [6087]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$2y^3 + 3xy^2y' = -2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 99

```
dsolve((2*y(x)^3+2)+(3*x*y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{((-x^2 + c_1)x)^{\frac{1}{3}}}{x}$$

$$y(x) = -\frac{((-x^2 + c_1)x)^{\frac{1}{3}}}{2x} - \frac{i\sqrt{3}((-x^2 + c_1)x)^{\frac{1}{3}}}{2x}$$

$$y(x) = -\frac{((-x^2 + c_1)x)^{\frac{1}{3}}}{2x} + \frac{i\sqrt{3}((-x^2 + c_1)x)^{\frac{1}{3}}}{2x}$$

✓ Solution by Mathematica

Time used: 0.281 (sec). Leaf size: 215

```
DSolve[(3*y[x]^3+2)+(3*x*y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt[3]{-\frac{1}{3}\sqrt[3]{-2x^3 + e^{9c_1}}}}{x}$$

$$y(x) \rightarrow \frac{\sqrt[3]{-2x^3 + e^{9c_1}}}{\sqrt[3]{3}x}$$

$$y(x) \rightarrow \frac{(-1)^{2/3}\sqrt[3]{-2x^3 + e^{9c_1}}}{\sqrt[3]{3}x}$$

$$y(x) \rightarrow \sqrt[3]{-\frac{2}{3}}$$

$$y(x) \rightarrow -\sqrt[3]{\frac{2}{3}}$$

$$y(x) \rightarrow -(-1)^{2/3}\sqrt[3]{\frac{2}{3}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{-\frac{2}{3}x^2}}{(-x^3)^{2/3}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{\frac{2}{3}\sqrt[3]{-x^3}}}{x}$$

$$y(x) \rightarrow \frac{(-1)^{2/3}\sqrt[3]{\frac{2}{3}\sqrt[3]{-x^3}}}{x}$$

22.10 problem 2(b)

Internal problem ID [6088]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 2(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$\cos(x) \cos(y) - 2y' \sin(y) \sin(x) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(cos(x)*cos(y(x))-2*sin(x)*sin(y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \arccos\left(\frac{1}{\sqrt{c_1 \sin(x)}}\right)$$

$$y(x) = \pi - \arccos\left(\frac{1}{\sqrt{c_1 \sin(x)}}\right)$$

✓ Solution by Mathematica

Time used: 0.491 (sec). Leaf size: 43

```
DSolve[Cos[x]*cos[y[x]]-(2*Sin[x]*Sin[y[x]])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{InverseFunction}\left[\int_1^{\#1} \frac{\sin(K[1])}{\cos(K[1])} dK[1] \&\amp; \left[\frac{1}{2} \log(\sin(x)) + c_1\right]\right]$$

$$y(x) \rightarrow \cos^{(-1)}(0)$$

22.11 problem 2(c)

Internal problem ID [6089]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 2(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [[_homogeneous, 'class G'], _rational, [_Abel, '2nd type', 'cl

$$5y^2x^3 + 2y + (3yx^4 + 2x)y' = 0$$

✓ Solution by Maple

Time used: 0.328 (sec). Leaf size: 347

`dsolve((5*x^3*y(x)^2+2*y(x))+(3*x^4*y(x)+2*x)*diff(y(x),x)=0,y(x), singsol=all)`

$$y(x) = \frac{\left(\frac{6 \left((108x^2 + 12\sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{1}{3}}}{c_1} + \frac{72c_1}{\left((108x^2 + 12\sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{1}{3}}} \right)^2}{1296} - 1$$

$$y(x) = \frac{\left(-\frac{3 \left((108x^2 + 12\sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{1}{3}}}{c_1} - \frac{36c_1}{\left((108x^2 + 12\sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{1}{3}}} - 18i\sqrt{3} \left(\frac{\left((108x^2 + 12\sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{1}{3}}}{6c_1} - \frac{2c_1}{\left((108x^2 + 12\sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{1}{3}}} \right) \right)^2}{1296} - 1$$

$$y(x) = \frac{\left(-\frac{3 \left((108x^2 + 12\sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{1}{3}}}{c_1} - \frac{36c_1}{\left((108x^2 + 12\sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{1}{3}}} + 18i\sqrt{3} \left(\frac{\left((108x^2 + 12\sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{1}{3}}}{6c_1} - \frac{2c_1}{\left((108x^2 + 12\sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{1}{3}}} \right) \right)^2}{1296} - 1$$

✓ Solution by Mathematica

Time used: 49.208 (sec). Leaf size: 400

`DSolve[(5*x^3*y[x]^2+2*y[x])+(3*x^4*y[x]+2*x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> Tr`

$$y(x) \rightarrow \frac{-2x^2 + \frac{2x^4}{\sqrt[3]{\frac{27c_1x^{10}}{2} - x^6 + \frac{3}{2}\sqrt{3}\sqrt{c_1x^{16}(-4 + 27c_1x^4)}}}} + 2^{2/3}\sqrt[3]{27c_1x^{10} - 2x^6 + 3\sqrt{3}\sqrt{c_1x^{16}(-4 + 27c_1x^4)}}}{6x^5}$$

$$y(x) \rightarrow \frac{-4x^2 - \frac{2(1+i\sqrt{3})x^4}{\sqrt[3]{\frac{27c_1x^{10}}{2} - x^6 + \frac{3}{2}\sqrt{3}\sqrt{c_1x^{16}(-4 + 27c_1x^4)}}}} + i2^{2/3}(\sqrt{3} + i)\sqrt[3]{27c_1x^{10} - 2x^6 + 3\sqrt{3}\sqrt{c_1x^{16}(-4 + 27c_1x^4)}}}{12x^5}$$

$$y(x) \rightarrow \frac{4x^2 - \frac{2i(\sqrt{3}+i)x^4}{\sqrt[3]{\frac{27c_1x^{10}}{2} - x^6 + \frac{3}{2}\sqrt{3}\sqrt{c_1x^{16}(-4 + 27c_1x^4)}}}} + 2^{2/3}(1 + i\sqrt{3})\sqrt[3]{27c_1x^{10} - 2x^6 + 3\sqrt{3}\sqrt{c_1x^{16}(-4 + 27c_1x^4)}}}{12x^5}$$

22.12 problem 2(d)

Internal problem ID [6090]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 2(d).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$e^y + e^y x + x e^y y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve((exp(y(x))+x*exp(y(x)))+(x*exp(y(x)))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -x - \ln(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 15

```
DSolve[(Exp[y[x]]+x*Exp[y[x]])+(x*Exp[y[x]])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x - \log(x) + c_1$$

23 Chapter 6. Existence and uniqueness of solutions to systems and nth order equations.

Page 238

23.1 problem 1(a)	220
23.2 problem 1(b)	221
23.3 problem 1(c)	222
23.4 problem 1(d)	224
23.5 problem 1(e)	225
23.6 problem 1(f)	226
23.7 problem 2	227
23.8 problem 3	228
23.9 problem 5(b)	229
23.10problem 5(c)	230

23.1 problem 1(a)

Internal problem ID [6091]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y' = 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x$2)+diff(y(x),x)=1,y(x), singsol=all)
```

$$y(x) = -e^{-x}c_1 + x + c_2$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 18

```
DSolve[y''[x]+y'[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x - c_1 e^{-x} + c_2$$

23.2 problem 1(b)

Internal problem ID [6092]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' + y'e^x = e^x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x$2)+exp(x)*diff(y(x),x)=exp(x),y(x), singsol=all)
```

$$y(x) = -c_1 \operatorname{Ei}_1(e^x) + x + c_2$$

✓ Solution by Mathematica

Time used: 0.081 (sec). Leaf size: 18

```
DSolve[y''[x]+Exp[x]*y'[x]==Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \operatorname{ExpIntegralEi}(-e^x) + x + c_2$$

23.3 problem 1(c)

Internal problem ID [6093]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible]`

$$yy'' + 4y'^2 = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 158

```
dsolve(y(x)*diff(y(x),x$2)+4*diff(y(x),x)^2=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = (5c_1x + 5c_2)^{\frac{1}{5}}$$

$$y(x) = \left(-\frac{\sqrt{5}}{4} - \frac{1}{4} - \frac{i\sqrt{2}\sqrt{5-\sqrt{5}}}{4} \right) (5c_1x + 5c_2)^{\frac{1}{5}}$$

$$y(x) = \left(-\frac{\sqrt{5}}{4} - \frac{1}{4} + \frac{i\sqrt{2}\sqrt{5-\sqrt{5}}}{4} \right) (5c_1x + 5c_2)^{\frac{1}{5}}$$

$$y(x) = \left(\frac{\sqrt{5}}{4} - \frac{1}{4} - \frac{i\sqrt{2}\sqrt{5+\sqrt{5}}}{4} \right) (5c_1x + 5c_2)^{\frac{1}{5}}$$

$$y(x) = \left(\frac{\sqrt{5}}{4} - \frac{1}{4} + \frac{i\sqrt{2}\sqrt{5+\sqrt{5}}}{4} \right) (5c_1x + 5c_2)^{\frac{1}{5}}$$

✓ Solution by Mathematica

Time used: 0.178 (sec). Leaf size: 20

```
DSolve[y[x]*y'[x]+4*(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 \sqrt[5]{5x - c_1}$$

23.4 problem 1(d)

Internal problem ID [6094]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + k^2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+k^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(kx) + c_2 \cos(kx)$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 20

```
DSolve[y''[x]+k^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(kx) + c_2 \sin(kx)$$

23.5 problem 1(e)

Internal problem ID [6095]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear],`

$$y'' - yy' = 0$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)=y(x)*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = \frac{\tan\left(\frac{(c_2+x)\sqrt{2}}{2c_1}\right)\sqrt{2}}{c_1}$$

✓ Solution by Mathematica

Time used: 16.739 (sec). Leaf size: 34

```
DSolve[y''[x]==y[x]*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt{2}\sqrt{c_1} \tan\left(\frac{\sqrt{c_1}(x+c_2)}{\sqrt{2}}\right)$$

23.6 problem 1(f)

Internal problem ID [6096]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' - 2y' = x^3$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x*diff(y(x),x$2)-2*diff(y(x),x)=x^3,y(x), singsol=all)
```

$$y(x) = \frac{1}{4}x^4 + \frac{1}{3}c_1x^3 + c_2$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 24

```
DSolve[x*y''[x]-2*y'[x]==x^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^4}{4} + \frac{c_1x^3}{3} + c_2$$

23.7 problem 2

Internal problem ID [6097]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_xy]]`

$$y'' - y'^2 = 1$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 7

```
dsolve([diff(y(x),x$2)=1+diff(y(x),x)^2,y(0) = 0, D(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = \ln(\sec(x))$$

✓ Solution by Mathematica

Time used: 2.581 (sec). Leaf size: 27

```
DSolve[{y'[x]==1+(y'[x])^2,{y[0]==0,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\log(-\cos(x)) + i\pi$$

$$y(x) \rightarrow -\log(\cos(x))$$

23.8 problem 3

Internal problem ID [6098]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_poly_y`

$$y'' + \frac{1}{2y'^2} = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = -1]$$

✓ Solution by Maple

Time used: 0.438 (sec). Leaf size: 26

```
dsolve([diff(y(x),x$2)=-1/(2*diff(y(x),x)^2),y(0) = 1, D(y)(0) = -1],y(x), singsol=all)
```

$$y(x) = \frac{3(x + \frac{2}{3})(-12x - 8)^{\frac{1}{3}}(-1 + i\sqrt{3})}{16} + \frac{3}{2}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 27

```
DSolve[{y'[x]==-1/(2*(y'[x])^2),{y[0]==1,y'[0]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{8}(12 - (-2)^{2/3}(-3x - 2)^{4/3})$$

23.9 problem 5(b)

Internal problem ID [6099]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 5(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$y'' + \sin(y) = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = \beta]$$

✓ Solution by Maple

Time used: 0.812 (sec). Leaf size: 53

```
dsolve([diff(y(x),x$2)+sin(y(x))=0,y(0) = 0, D(y)(0) = beta],y(x), singsol=all)
```

$$y(x) = \text{RootOf} \left(- \left(\int_0^{-z} \frac{1}{\sqrt{2 \cos(_a) + \beta^2 - 2}} d_a \right) + x \right)$$

$$y(x) = \text{RootOf} \left(\int_0^{-z} \frac{1}{\sqrt{2 \cos(_a) + \beta^2 - 2}} d_a + x \right)$$

✓ Solution by Mathematica

Time used: 0.621 (sec). Leaf size: 19

```
DSolve[{y'[x]+Sin[y[x]]==0,{y[0]==0,y'[0]==\[Beta]}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 \text{JacobiAmplitude} \left(\frac{x\beta}{2}, \frac{4}{\beta^2} \right)$$

23.10 problem 5(c)

Internal problem ID [6100]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 5(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$y'' + \sin(y) = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 2]$$

✓ Solution by Maple

Time used: 0.156 (sec). Leaf size: 23

```
dsolve([diff(y(x),x$2)+sin(y(x))=0,y(0) = 0, D(y)(0) = 2],y(x), singsol=all)
```

$$y(x) = \text{RootOf} \left(- \left(\int_0^{-z} \frac{1}{\sqrt{2 \cos(_a) + 2}} d_a \right) + x \right)$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y'[x]+Sin[y[x]]==0,{y[0]==0,y'[0]==2}},y[x],x,IncludeSingularSolutions -> True]
```

{}

24 Chapter 6. Existence and uniqueness of solutions to systems and nth order equations.

Page 250

24.1 problem 3	232
24.2 problem 4	233
24.3 problem 5	235

24.1 problem 3

Internal problem ID [6101]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 250

Problem number: 3.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(x) &= y_1(x) \\y_2'(x) &= y_1(x) + y_2(x)\end{aligned}$$

With initial conditions

$$[y_1(0) = 1, y_2(0) = 2]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 16

```
dsolve([diff(y__1(x),x) = y__1(x), diff(y__2(x),x) = y__1(x)+y__2(x), y__1(0) = 1, y__2(0) = 2])
```

$$y_1(x) = e^x$$

$$y_2(x) = e^x(x + 2)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 18

```
DSolve[{y1'[x]==y1[x],y2'[x]==y1[x]+y2[x]},{y1[0]==1,y2[0]==2},{y1[x],y2[x]},x,IncludeSingularSolutions->True]
```

$$y1(x) \rightarrow e^x$$

$$y2(x) \rightarrow e^x(x + 2)$$

24.2 problem 4

Internal problem ID [6102]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 250

Problem number: 4.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(x) &= y_2(x) \\ y_2'(x) &= 6y_1(x) + y_2(x)\end{aligned}$$

With initial conditions

$$[y_1(0) = 1, y_2(0) = -1]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 34

```
dsolve([diff(y__1(x),x) = y__2(x), diff(y__2(x),x) = 6*y__1(x)+y__2(x), y__1(0) = 1, y__2(0)
```

$$y_1(x) = \frac{e^{3x}}{5} + \frac{4e^{-2x}}{5}$$

$$y_2(x) = \frac{3e^{3x}}{5} - \frac{8e^{-2x}}{5}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 42

```
DSolve[{y1'[x]==y2[x],y2'[x]==6*y1[x]+y2[x]},{y1[0]==1,y2[0]==-1},{y1[x],y2[x]},x,IncludeSin
```

$$y1(x) \rightarrow \frac{1}{5}e^{-2x}(e^{5x} + 4)$$

$$y2(x) \rightarrow \frac{1}{5}e^{-2x}(3e^{5x} - 8)$$

24.3 problem 5

Internal problem ID [6103]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 250

Problem number: 5.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(x) &= y_1(x) + y_2(x) \\y_2'(x) &= y_1(x) + y_2(x) + e^{3x}\end{aligned}$$

With initial conditions

$$[y_1(0) = 0, y_2(0) = 0]$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 36

```
dsolve([diff(y__1(x),x) = y__1(x)+y__2(x), diff(y__2(x),x) = y__1(x)+y__2(x)+exp(3*x), y__1(0)=0, y__2(0)=0])
```

$$y_1(x) = -\frac{e^{2x}}{2} + \frac{e^{3x}}{3} + \frac{1}{6}$$

$$y_2(x) = -\frac{e^{2x}}{2} + \frac{2e^{3x}}{3} - \frac{1}{6}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 46

```
DSolve[{y1'[x]==y1[x]+y2[x],y2'[x]==y1[x]+y2[x]+Exp[3*x]},{y1[0]==0,y2[0]==0},{y1[x],y2[x]},
```

$$y1(x) \rightarrow \frac{1}{6}(e^x - 1)^2(2e^x + 1)$$

$$y2(x) \rightarrow \frac{1}{6}(-3e^{2x} + 4e^{3x} - 1)$$

25 Chapter 6. Existence and uniqueness of solutions to systems and nth order equations.

Page 254

25.1 problem 2 238

25.1 problem 2

Internal problem ID [6104]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 254

Problem number: 2.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(x) = 3y_1(x) + xy_3(x)$$

$$y_2'(x) = y_2(x) + x^3y_3(x)$$

$$y_3'(x) = 2xy_2(x) - y_2(x) + e^x y_3(x)$$

X Solution by Maple

```
dsolve([diff(y__1(x),x)=3*y__1(x)+x*y__3(x),diff(y__2(x),x)=y__2(x)+x^3*y__3(x),diff(y__3(x)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y1'[x]==3*y1[x]+x*y3[x],y2'[x]==y2[x]+x^3*y3[x],y3'[x]==2*x*y1[x]-y2[x]+Exp[x]*y3[x]
```

Not solved