


A Solution Manual For

**Elementary Differential
Equations, Martin, Reissner,
2nd ed, 1961**



Nasser M. Abbasi

March 3, 2024

Contents

1 Exercis 2, page 5

2

1 Exercis 2, page 5

1.1	problem 2(a)	3
1.2	problem 2(b)	4
1.3	problem 2(c)	5
1.4	problem 2(d)	6
1.5	problem 2(e)	7
1.6	problem 2(a)	8
1.7	problem 3(a)	9
1.8	problem 3(b)	10
1.9	problem 3(c)	11
1.10	problem 3(d)	12
1.11	problem 3(e)	13
1.12	problem 3(f)	14
1.13	problem 3(g)	15
1.14	problem 3(h)	16
1.15	problem 3(i)	17

1.1 problem 2(a)

Internal problem ID [2432]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' = 2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)=2,y(x), singsol=all)
```

$$y(x) = 2x + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 11

```
DSolve[y'[x]==2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2x + c_1$$

1.2 problem 2(b)

Internal problem ID [2433]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' = 2e^{3x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=2*exp(3*x),y(x), singsol=all)
```

$$y(x) = \frac{2e^{3x}}{3} + c_1$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 17

```
DSolve[y'[x]==2*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2e^{3x}}{3} + c_1$$

1.3 problem 2(c)

Internal problem ID [2434]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' = \frac{2}{\sqrt{1-x^2}}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)=2/sqrt(1-x^2),y(x), singsol=all)
```

$$y(x) = 2 \arcsin(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 28

```
DSolve[y'[x]==2/Sqrt[1-x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -4 \arctan\left(\frac{\sqrt{1-x^2}}{x+1}\right) + c_1$$

1.4 problem 2(d)

Internal problem ID [2435]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(d).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y' = e^{x^2}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x)=exp(x^2),y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{\pi} \operatorname{erfi}(x)}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 19

```
DSolve[y'[x]==Exp[x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}\sqrt{\pi}\operatorname{erfi}(x) + c_1$$

1.5 problem 2(e)

Internal problem ID [2436]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(e).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' = x e^{x^2}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=x*exp(x^2),y(x), singsol=all)
```

$$y(x) = \frac{e^{x^2}}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 17

```
DSolve[y'[x]==x*Exp[x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{x^2}}{2} + c_1$$

1.6 problem 2(a)

Internal problem ID [2437]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' = \arcsin(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)=arcsin(x),y(x), singsol=all)
```

$$y(x) = x \arcsin(x) + \sqrt{-x^2 + 1} + c_1$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 23

```
DSolve[y'[x]==ArcSin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \arcsin(x) + \sqrt{1 - x^2} + c_1$$

1.7 problem 3(a)

Internal problem ID [2438]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' - yx = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=y(x)*x,y(x), singsol=all)
```

$$y(x) = e^{\frac{x^2}{2}} c_1$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 22

```
DSolve[y'[x]==y[x]*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{\frac{x^2}{2}}$$

$$y(x) \rightarrow 0$$

1.8 problem 3(b)

Internal problem ID [2439]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - y^2x^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)=y(x)^2*x^2,y(x), singsol=all)
```

$$y(x) = \frac{3}{-x^3 + 3c_1}$$

✓ Solution by Mathematica

Time used: 0.107 (sec). Leaf size: 22

```
DSolve[y'[x]==y[x]^2*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{3}{x^3 + 3c_1}$$

$$y(x) \rightarrow 0$$

1.9 problem 3(c)

Internal problem ID [2440]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_separable]`

$$y' + x e^y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)=-x*exp(y(x)),y(x), singsol=all)
```

$$y(x) = \ln\left(\frac{2}{x^2 + 2c_1}\right)$$

✓ Solution by Mathematica

Time used: 0.307 (sec). Leaf size: 19

```
DSolve[y'[x]==-x*Exp[y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(2) - \log(x^2 - 2c_1)$$

1.10 problem 3(d)

Internal problem ID [2441]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(d).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' \sin(y) = x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)*sin(y(x))=x^2,y(x), singsol=all)
```

$$y(x) = \pi - \arccos\left(\frac{x^3}{3} + c_1\right)$$

✓ Solution by Mathematica

Time used: 0.509 (sec). Leaf size: 37

```
DSolve[y'[x]*Sin[y[x]]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\arccos\left(-\frac{x^3}{3} - c_1\right)$$

$$y(x) \rightarrow \arccos\left(-\frac{x^3}{3} - c_1\right)$$

1.11 problem 3(e)

Internal problem ID [2442]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(e).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$xy' - \sqrt{1 - y^2} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)*x=sqrt(1-y(x)^2),y(x), singsol=all)
```

$$y(x) = \sin(\ln(x) + c_1)$$

✓ Solution by Mathematica

Time used: 0.217 (sec). Leaf size: 29

```
DSolve[y'[x]*x==Sqrt[1-y[x]^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \cos(\log(x) + c_1)$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 1$$

$$y(x) \rightarrow \text{Interval}[\{-1, 1\}]$$

1.12 problem 3(f)

Internal problem ID [2443]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(f).

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$y'^2 - y^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve((diff(y(x),x))^2-y(x)^2=0,y(x), singsol=all)
```

$$y(x) = c_1 e^x$$

$$y(x) = e^{-x} c_1$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 28

```
DSolve[(y'[x])^2-y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-x}$$

$$y(x) \rightarrow c_1 e^x$$

$$y(x) \rightarrow 0$$

1.13 problem 3(g)

Internal problem ID [2444]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(g).

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$y'^2 - 3y' = -2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)^2-3*diff(y(x),x)+2=0,y(x), singsol=all)
```

$$y(x) = 2x + c_1$$

$$y(x) = c_1 + x$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 19

```
DSolve[(y'[x])^2-3*y'[x]+2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x + c_1$$

$$y(x) \rightarrow 2x + c_1$$

1.14 problem 3(h)

Internal problem ID [2445]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(h).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$(x^2 + 1) y' = 1$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 8

```
dsolve((1+x^2)*diff(y(x),x)=1,y(x), singsol=all)
```

$$y(x) = \arctan(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 10

```
DSolve[(1+x^2)*y'[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arctan(x) + c_1$$

1.15 problem 3(i)

Internal problem ID [2446]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(i).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' \sin(x) = 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)*sin(x)=1,y(x), singsol=all)
```

$$y(x) = -\ln(\csc(x) + \cot(x)) + c_1$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 13

```
DSolve[y'[x]*Sin[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\operatorname{arctanh}(\cos(x)) + c_1$$