


A Solution Manual For

**Elementary differential
equations. By Earl D. Rainville,
Phillip E. Bedient. Macmillian
Publishing Co. NY. 6th edition.
1981.**



Nasser M. Abbasi

March 3, 2024

Contents

1	CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309	2
2	CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314	29
3	CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320	47
4	CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324	86

**1 CHAPTER 16. Nonlinear equations. Section 94.
Factoring the left member. EXERCISES Page
309**

1.1	problem 1	3
1.2	problem 2	4
1.3	problem 3	5
1.4	problem 4	6
1.5	problem 5	7
1.6	problem 6	8
1.7	problem 7	9
1.8	problem 8	10
1.9	problem 9	11
1.10	problem 10	13
1.11	problem 11	14
1.12	problem 12	15
1.13	problem 13	16
1.14	problem 14	18
1.15	problem 15	19
1.16	problem 16	22
1.17	problem 17	24
1.18	problem 18	27
1.19	problem 19	28

1.1 problem 1

Internal problem ID [6767]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 1.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [separable]

$$x^2 y'^2 - y^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x)^2-y(x)^2=0,y(x), singsol=all)
```

$$y(x) = c_1 x$$

$$y(x) = \frac{c_1}{x}$$

✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 24

```
DSolve[x^2*(y'[x])^2-y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{x}$$

$$y(x) \rightarrow c_1 x$$

$$y(x) \rightarrow 0$$

1.2 problem 2

Internal problem ID [6768]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 2.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$xy'^2 - (2x + 3y)y' + 6y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x*diff(y(x),x)^2-(2*x+3*y(x))*diff(y(x),x)+6*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1x^3$$

$$y(x) = 2x + c_1$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 26

```
DSolve[x*(y'[x])^2-(2*x+3*y[x])*y'[x]+6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x^3$$

$$y(x) \rightarrow 2x + c_1$$

$$y(x) \rightarrow 0$$

1.3 problem 3

Internal problem ID [6769]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 3.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_separable]

$$x^2y'^2 - 5xyy' + 6y^2 = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve(x^2*diff(y(x),x)^2-5*x*y(x)*diff(y(x),x)+6*y(x)^2=0,y(x), singsol=all)
```

$$y(x) = c_1x^3$$

$$y(x) = c_1x^2$$

✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 26

```
DSolve[x^2*(y'[x])^2-5*x*y[x]*y'[x]+6*y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x^2$$

$$y(x) \rightarrow c_1x^3$$

$$y(x) \rightarrow 0$$

1.4 problem 4

Internal problem ID [6770]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 4.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_separable]

$$x^2 y'^2 + y'x - y^2 - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(x^2*diff(y(x),x)^2+x*diff(y(x),x)-y(x)^2-y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 x$$

$$y(x) = \frac{-x + c_1}{x}$$

✓ Solution by Mathematica

Time used: 0.045 (sec). Leaf size: 31

```
DSolve[x^2*(y'[x])^2+x*y'[x]-y[x]^2-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 x$$

$$y(x) \rightarrow -1 + \frac{c_1}{x}$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 0$$

1.5 problem 5

Internal problem ID [6771]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 5.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$xy'^2 + (1 - x^2y)y' - yx = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(x*diff(y(x),x)^2+(1-x^2*y(x))*diff(y(x),x)-x*y(x)=0,y(x), singsol=all)
```

$$y(x) = -\ln(x) + c_1$$

$$y(x) = e^{\frac{x^2}{2}} c_1$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 28

```
DSolve[x*(y'[x])^2+(1-x^2*y[x])*y'[x]-x*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{\frac{x^2}{2}}$$

$$y(x) \rightarrow -\log(x) + c_1$$

1.6 problem 6

Internal problem ID [6772]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 6.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$y'^2 - (x^2y + 3)y' + 3x^2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x)^2-(x^2*y(x)+3)*diff(y(x),x)+3*x^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{x^3}{3}}$$

$$y(x) = 3x + c_1$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 27

```
DSolve[(y'[x])^2-(x^2*y[x]+3)*y'[x]+3*x^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{\frac{x^3}{3}}$$

$$y(x) \rightarrow 3x + c_1$$

1.7 problem 7

Internal problem ID [6773]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 7.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$xy'^2 - (1 + yx)y' + y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(x*diff(y(x),x)^2-(1+x*y(x))*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \ln(x) + c_1$$

$$y(x) = e^x c_1$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 20

```
DSolve[x*(y'[x])^2-(1+x*y[x])*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x$$

$$y(x) \rightarrow \log(x) + c_1$$

1.8 problem 8

Internal problem ID [6774]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 8.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [separable]

$$y'^2 - x^2 y^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x)^2-x^2*y(x)^2=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{x^2}{2}} c_1$$

$$y(x) = e^{-\frac{x^2}{2}} c_1$$

✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 38

```
DSolve[(y'[x])^2-x^2*y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-\frac{x^2}{2}}$$

$$y(x) \rightarrow c_1 e^{\frac{x^2}{2}}$$

$$y(x) \rightarrow 0$$

1.9 problem 9

Internal problem ID [6775]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 9.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$(x + y)^2 y'^2 - y^2 = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 48

```
dsolve((x+y(x))^2*diff(y(x),x)^2=y(x)^2,y(x), singsol=all)
```

$$y(x) = e^{\text{LambertW}(x e^{c_1}) - c_1}$$

$$y(x) = -x - \sqrt{x^2 + 2c_1}$$

$$y(x) = -x + \sqrt{x^2 + 2c_1}$$

✓ Solution by Mathematica

Time used: 4.023 (sec). Leaf size: 101

```
DSolve[(x+y[x])^2*(y'[x])^2==y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x - \sqrt{x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -x + \sqrt{x^2 + e^{2c_1}}$$

$$y(x) \rightarrow \frac{x}{W(e^{-c_1}x)}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -\sqrt{x^2} - x$$

$$y(x) \rightarrow \sqrt{x^2} - x$$

1.10 problem 10

Internal problem ID [6776]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 10.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$yy'^2 + (x - y^2)y' - yx = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve(y(x)*diff(y(x),x)^2+(x-y(x)^2)*diff(y(x),x)-x*y(x)=0,y(x), singsol=all)
```

$$y(x) = \sqrt{-x^2 + c_1}$$

$$y(x) = -\sqrt{-x^2 + c_1}$$

$$y(x) = e^x c_1$$

✓ Solution by Mathematica

Time used: 0.134 (sec). Leaf size: 54

```
DSolve[y[x]*(y'[x])^2+(x-y[x]^2)*y'[x]-x*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x$$

$$y(x) \rightarrow -\sqrt{-x^2 + 2c_1}$$

$$y(x) \rightarrow \sqrt{-x^2 + 2c_1}$$

$$y(x) \rightarrow 0$$

1.11 problem 11

Internal problem ID [6777]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 11.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_separable]

$$y'^2 - xy(x+y)y' + x^3y^3 = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve(diff(y(x),x)^2-x*y(x)*(x+y(x))*diff(y(x),x)+x^3*y(x)^3=0,y(x), singsol=all)
```

$$y(x) = \frac{2}{-x^2 + 2c_1}$$

$$y(x) = c_1 e^{\frac{x^3}{3}}$$

✓ Solution by Mathematica

Time used: 0.122 (sec). Leaf size: 38

```
DSolve[(y'[x])^2-x*y[x]*(x+y[x])*y'[x]+x^3*y[x]^3==0,y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow c_1 e^{\frac{x^3}{3}}$$

$$y(x) \rightarrow -\frac{2}{x^2 + 2c_1}$$

$$y(x) \rightarrow 0$$

1.12 problem 12

Internal problem ID [6778]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 12.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$(4x - y)y'^2 + 6(x - y)y' - 5y = -2x$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 55

```
dsolve((4*x-y(x))*diff(y(x),x)^2+6*(x-y(x))*diff(y(x),x)+2*x-5*y(x)=0,y(x), singsol=all)
```

$$y(x) = -x + c_1$$

$$y(x) = -\frac{4c_1x - \sqrt{-12c_1x + 1} - 1}{2c_1}$$

$$y(x) = -\frac{4c_1x + \sqrt{-12c_1x + 1} - 1}{2c_1}$$

✓ Solution by Mathematica

Time used: 1.077 (sec). Leaf size: 90

```
DSolve[(4*x-y[x])*(y'[x])^2+6*(x-y[x])*y'[x]+2*x-5*y[x]==0,y[x],x,IncludeSingularSolutions -
```

$$y(x) \rightarrow \frac{1}{2} \left(-4x - e^{\frac{c_1}{2}} \sqrt{12x + e^{c_1}} - e^{c_1} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left(-4x + e^{\frac{c_1}{2}} \sqrt{12x + e^{c_1}} - e^{c_1} \right)$$

$$y(x) \rightarrow -x + c_1$$

1.13 problem 13

Internal problem ID [6779]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 13.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$(x - y)^2 y'^2 - y^2 = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 45

```
dsolve((x-y(x))^2*diff(y(x),x)^2=y(x)^2,y(x), singsol=all)
```

$$y(x) = x - \sqrt{x^2 - 2c_1}$$

$$y(x) = x + \sqrt{x^2 - 2c_1}$$

$$y(x) = e^{\text{LambertW}(-x e^{-c_1}) + c_1}$$

✓ Solution by Mathematica

Time used: 4.446 (sec). Leaf size: 99

```
DSolve[(x-y[x])^2*(y'[x])^2==y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x - \sqrt{x^2 - e^{2c_1}}$$

$$y(x) \rightarrow x + \sqrt{x^2 - e^{2c_1}}$$

$$y(x) \rightarrow -\frac{x}{W(-e^{-c_1}x)}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow x - \sqrt{x^2}$$

$$y(x) \rightarrow \sqrt{x^2} + x$$

1.14 problem 14

Internal problem ID [6780]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 14.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$yy'^2x + (xy^2 - 1)y' - y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve(x*y(x)*diff(y(x),x)^2+(x*y(x)^2-1)*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = \sqrt{2 \ln(x) + c_1}$$

$$y(x) = -\sqrt{2 \ln(x) + c_1}$$

$$y(x) = e^{-x}c_1$$

✓ Solution by Mathematica

Time used: 0.108 (sec). Leaf size: 57

```
DSolve[x*y[x]*(y'[x])^2+(x*y[x]^2-1)*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-x}$$

$$y(x) \rightarrow -\sqrt{2} \sqrt{\log(x) + c_1}$$

$$y(x) \rightarrow \sqrt{2} \sqrt{\log(x) + c_1}$$

$$y(x) \rightarrow 0$$

1.15 problem 15

Internal problem ID [6781]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 15.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _dAlembert]`

$$(x^2 + y^2)^2 y' - 4x^2 y^2 = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 301

`dsolve((x^2+y(x)^2)^2*diff(y(x),x)^2=4*x^2*y(x)^2,y(x), singsol=all)`

$$y(x) = -\frac{-1 + \sqrt{4c_1^2x^2 + 1}}{2c_1}$$

$$y(x) = \frac{1 + \sqrt{4c_1^2x^2 + 1}}{2c_1}$$

$$y(x) = \frac{\frac{\left(4+4\sqrt{4c_1^3x^6+1}\right)^{\frac{1}{3}}}{2} - \frac{2x^2c_1}{\left(4+4\sqrt{4c_1^3x^6+1}\right)^{\frac{1}{3}}}}{\sqrt{c_1}}$$

$$y(x) = \frac{-\frac{\left(4+4\sqrt{4c_1^3x^6+1}\right)^{\frac{1}{3}}}{4} + \frac{x^2c_1}{\left(4+4\sqrt{4c_1^3x^6+1}\right)^{\frac{1}{3}}} - \frac{i\sqrt{3}\left(\frac{\left(4+4\sqrt{4c_1^3x^6+1}\right)^{\frac{1}{3}}}{2} + \frac{2x^2c_1}{\left(4+4\sqrt{4c_1^3x^6+1}\right)^{\frac{1}{3}}}\right)}{2}}{\sqrt{c_1}}$$

$$y(x) = \frac{-\frac{\left(4+4\sqrt{4c_1^3x^6+1}\right)^{\frac{1}{3}}}{4} + \frac{x^2c_1}{\left(4+4\sqrt{4c_1^3x^6+1}\right)^{\frac{1}{3}}} + \frac{i\sqrt{3}\left(\frac{\left(4+4\sqrt{4c_1^3x^6+1}\right)^{\frac{1}{3}}}{2} + \frac{2x^2c_1}{\left(4+4\sqrt{4c_1^3x^6+1}\right)^{\frac{1}{3}}}\right)}{2}}{\sqrt{c_1}}$$

✓ Solution by Mathematica

Time used: 15.845 (sec). Leaf size: 345

`DSolve[(x^2+y[x]^2)^2*(y'[x])^2==4*x^2*y[x]^2,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{1}{2} \left(-\sqrt{4x^2 + e^{2c_1}} - e^{c_1} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left(\sqrt{4x^2 + e^{2c_1}} - e^{c_1} \right)$$

$$y(x) \rightarrow \frac{\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}{\sqrt[3]{2}} - \frac{\sqrt[3]{2}x^2}{\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}$$

$$y(x) \rightarrow \frac{i2^{2/3}(\sqrt{3} + i) (\sqrt{4x^6 + e^{6c_1}} + e^{3c_1})^{2/3} + \sqrt[3]{2}(2 + 2i\sqrt{3}) x^2}{4\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}$$

$$y(x) \rightarrow \frac{(1 - i\sqrt{3}) x^2}{2^{2/3} \sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}} - \frac{(1 + i\sqrt{3}) \sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}{2\sqrt[3]{2}}$$

$$y(x) \rightarrow 0$$

1.16 problem 16

Internal problem ID [6782]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 16.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$(x + y)^2 y'^2 + (2y^2 + yx - x^2) y' + y(-x + y) = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 85

```
dsolve((y(x)+x)^2*diff(y(x),x)^2+(2*y(x)^2+x*y(x)-x^2)*diff(y(x),x)+y(x)*(y(x)-x)=0,y(x), si
```

$$y(x) = -x - \sqrt{x^2 + 2c_1}$$

$$y(x) = -x + \sqrt{x^2 + 2c_1}$$

$$y(x) = \frac{-c_1x - \sqrt{2c_1^2x^2 + 1}}{c_1}$$

$$y(x) = \frac{-c_1x + \sqrt{2c_1^2x^2 + 1}}{c_1}$$

✓ Solution by Mathematica

Time used: 0.492 (sec). Leaf size: 172

```
DSolve[(y[x]+x)^2*(y'[x])^2+(2*y[x]^2+x*y[x]-x^2)*y'[x]+y[x]*(y[x]-x)==0,y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow -x - \sqrt{x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -x + \sqrt{x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -x - \sqrt{2x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -x + \sqrt{2x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -\sqrt{x^2} - x$$

$$y(x) \rightarrow \sqrt{x^2} - x$$

$$y(x) \rightarrow -\sqrt{2}\sqrt{x^2} - x$$

$$y(x) \rightarrow \sqrt{2}\sqrt{x^2} - x$$

1.17 problem 17

Internal problem ID [6783]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 17.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$xy(x^2 + y^2)(y'^2 - 1) - y'(x^4 + x^2y^2 + y^4) = 0$$

✓ Solution by Maple

Time used: 0.594 (sec). Leaf size: 250

```
dsolve(x*y(x)*(x^2+y(x)^2)*(diff(y(x),x)^2-1)=diff(y(x),x)*(x^4+x^2*y(x)^2+y(x)^4),y(x), sin
```

$$y(x) = \frac{\sqrt{x^2 c_1 (c_1 x^2 - \sqrt{c_1^2 x^4 + 1})}}{x (c_1 x^2 - \sqrt{c_1^2 x^4 + 1}) c_1}$$

$$y(x) = \frac{\sqrt{x^2 c_1 (c_1 x^2 + \sqrt{c_1^2 x^4 + 1})}}{x (c_1 x^2 + \sqrt{c_1^2 x^4 + 1}) c_1}$$

$$y(x) = -\frac{\sqrt{x^2 c_1 (c_1 x^2 - \sqrt{c_1^2 x^4 + 1})}}{x (c_1 x^2 - \sqrt{c_1^2 x^4 + 1}) c_1}$$

$$y(x) = -\frac{\sqrt{x^2 c_1 (c_1 x^2 + \sqrt{c_1^2 x^4 + 1})}}{x (c_1 x^2 + \sqrt{c_1^2 x^4 + 1}) c_1}$$

$$y(x) = \sqrt{2 \ln(x) + c_1} x$$

$$y(x) = -\sqrt{2 \ln(x) + c_1} x$$

✓ Solution by Mathematica

Time used: 9.298 (sec). Leaf size: 248

```
DSolve[x*y[x]*(x^2+y[x]^2)*((y'[x])^2-1)==y'[x]*(x^4+x^2*y[x]^2+y[x]^4),y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow -\sqrt{-x^2 - \sqrt{x^4 + e^{4c_1}}}$$

$$y(x) \rightarrow \sqrt{-x^2 - \sqrt{x^4 + e^{4c_1}}}$$

$$y(x) \rightarrow -\sqrt{-x^2 + \sqrt{x^4 + e^{4c_1}}}$$

$$y(x) \rightarrow \sqrt{-x^2 + \sqrt{x^4 + e^{4c_1}}}$$

$$y(x) \rightarrow -x\sqrt{2\log(x) + c_1}$$

$$y(x) \rightarrow x\sqrt{2\log(x) + c_1}$$

$$y(x) \rightarrow -\sqrt{-\sqrt{x^4} - x^2}$$

$$y(x) \rightarrow \sqrt{-\sqrt{x^4} - x^2}$$

$$y(x) \rightarrow -\sqrt{\sqrt{x^4} - x^2}$$

$$y(x) \rightarrow \sqrt{\sqrt{x^4} - x^2}$$

1.18 problem 18

Internal problem ID [6784]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 18.

ODE order: 1.

ODE degree: 3.

CAS Maple gives this as type `[_quadrature]`

$$xy'^3 - (x^2 + x + y)y'^2 + (x^2 + yx + y)y' - yx = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(x*diff(y(x),x)^3-(x^2+x+y(x))*diff(y(x),x)^2+(x^2+x*y(x)+y(x))*diff(y(x),x)-x*y(x)=0,
```

$$y(x) = c_1x$$

$$y(x) = x + c_1$$

$$y(x) = \frac{x^2}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.044 (sec). Leaf size: 36

```
DSolve[x*(y'[x])^3-(x^2+x+y[x])*(y'[x])^2+(x^2+x*y[x]+y[x])*y'[x]-x*y[x]==0,y[x],x,IncludeSi
```

$$y(x) \rightarrow c_1x$$

$$y(x) \rightarrow x + c_1$$

$$y(x) \rightarrow \frac{x^2}{2} + c_1$$

$$y(x) \rightarrow 0$$

1.19 problem 19

Internal problem ID [6785]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 94. Factoring the left member. EXERCISES Page 309

Problem number: 19.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type [_quadrature]

$$yy'^2x + (x + y)y' = -1$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 32

```
dsolve(x*y(x)*diff(y(x),x)^2+(x+y(x))*diff(y(x),x)+1=0,y(x), singsol=all)
```

$$y(x) = -\ln(x) + c_1$$

$$y(x) = \sqrt{-2x + c_1}$$

$$y(x) = -\sqrt{-2x + c_1}$$

✓ Solution by Mathematica

Time used: 0.062 (sec). Leaf size: 53

```
DSolve[x*y[x]*(y'[x])^2+(x+y[x])*y'[x]+1==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{2}\sqrt{-x + c_1}$$

$$y(x) \rightarrow \sqrt{2}\sqrt{-x + c_1}$$

$$y(x) \rightarrow -\log(x) + c_1$$

**2 CHAPTER 16. Nonlinear equations. Section 97.
The p-discriminant equation. EXERCISES Page
314**

2.1	problem 8	30
2.2	problem 9	31
2.3	problem 10	33
2.4	problem 11	35
2.5	problem 12	36
2.6	problem 13	38
2.7	problem 14	40
2.8	problem 15	42
2.9	problem 16	45

2.1 problem 8

Internal problem ID [6786]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

Problem number: 8.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _dAlembert]`

$$xy'^2 - 2y'y = -4x$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 29

```
dsolve(x*diff(y(x),x)^2-2*y(x)*diff(y(x),x)+4*x=0,y(x), singsol=all)
```

$$y(x) = -2x$$

$$y(x) = 2x$$

$$y(x) = -\frac{\left(-\frac{x^2}{c_1^2} - 4\right) c_1}{2}$$

✓ Solution by Mathematica

Time used: 0.294 (sec). Leaf size: 43

```
DSolve[x*(y'[x])^2-2*y[x]*y'[x]+4*x==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2x \cosh(-\log(x) + c_1)$$

$$y(x) \rightarrow -2x \cosh(\log(x) + c_1)$$

$$y(x) \rightarrow -2x$$

$$y(x) \rightarrow 2x$$

2.2 problem 9

Internal problem ID [6787]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

Problem number: 9.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational]`

$$3x^4y'^2 - y'x - y = 0$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 147

```
dsolve(3*x^4*diff(y(x),x)^2-x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{1}{12x^2}$$

$$y(x) = \frac{-c_1(-c_1 + 2ix\sqrt{3}) - c_1^2 - 6x^2}{6x^2c_1^2}$$

$$y(x) = \frac{-c_1(-c_1 - 2ix\sqrt{3}) - c_1^2 - 6x^2}{6x^2c_1^2}$$

$$y(x) = \frac{c_1(c_1 + 2ix\sqrt{3}) - 6x^2 - c_1^2}{6c_1^2x^2}$$

$$y(x) = \frac{c_1(c_1 - 2ix\sqrt{3}) - 6x^2 - c_1^2}{6c_1^2x^2}$$

✓ Solution by Mathematica

Time used: 0.518 (sec). Leaf size: 123

```
DSolve[3*x^4*(y'[x])^2-x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[-\frac{x\sqrt{12x^2y(x)+1}\operatorname{arctanh}\left(\sqrt{12x^2y(x)+1}\right)}{\sqrt{12x^4y(x)+x^2}} - \frac{1}{2}\log(y(x)) = c_1, y(x) \right]$$

$$\text{Solve} \left[\frac{x\sqrt{12x^2y(x)+1}\operatorname{arctanh}\left(\sqrt{12x^2y(x)+1}\right)}{\sqrt{12x^4y(x)+x^2}} - \frac{1}{2}\log(y(x)) = c_1, y(x) \right]$$

$$y(x) \rightarrow 0$$

2.3 problem 10

Internal problem ID [6788]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

Problem number: 10.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$y'^2 - y'x - y = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 77

```
dsolve(diff(y(x),x)^2-x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$\frac{c_1}{\sqrt{2x - 2\sqrt{x^2 + 4y(x)}}} + \frac{2x}{3} + \frac{\sqrt{x^2 + 4y(x)}}{3} = 0$$
$$\frac{c_1}{\sqrt{2x + 2\sqrt{x^2 + 4y(x)}}} + \frac{2x}{3} - \frac{\sqrt{x^2 + 4y(x)}}{3} = 0$$

✓ Solution by Mathematica

Time used: 60.178 (sec). Leaf size: 1003

`DSolve[(y'[x])^2 - x*y'[x] - y[x] == 0, y[x], x, IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{\left(x^2 + \sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}\right)^2 + 8e^{3c_1}x}{4\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}}$$

$$y(x) \rightarrow \frac{1}{8} \left(4x^2 - \frac{i(\sqrt{3} - i)x(x^3 + 8e^{3c_1})}{\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}} + i(\sqrt{3} + i)\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}} \right)$$

$$y(x) \rightarrow \frac{1}{8} \left(4x^2 + \frac{i(\sqrt{3} + i)x(x^3 + 8e^{3c_1})}{\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}} - (1 + i\sqrt{3})\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}} \right)$$

$$y(x) \rightarrow \frac{2\sqrt[3]{2}x^4 + 2^{2/3}\left(-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}}(4x^3 + e^{3c_1})^3 + e^{6c_1}\right)^{2/3} + 4x^2\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}}(4x^3 + e^{3c_1})^3 + e^{6c_1}}}{8\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}}(4x^3 + e^{3c_1})^3 + e^{6c_1}}}$$

$$y(x) \rightarrow \frac{1}{16} \left(8x^2 + \frac{2\sqrt[3]{2}(1 + i\sqrt{3})x(-x^3 + 2e^{3c_1})}{\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}}(4x^3 + e^{3c_1})^3 + e^{6c_1}}} + i2^{2/3}(\sqrt{3} + i)\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}}(4x^3 + e^{3c_1})^3 + e^{6c_1}} \right)$$

$$y(x) \rightarrow \frac{1}{16} \left(8x^2 + \frac{2i\sqrt[3]{2}(\sqrt{3} + i)x(x^3 - 2e^{3c_1})}{\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}}(4x^3 + e^{3c_1})^3 + e^{6c_1}}} - 2^{2/3}(1 + i\sqrt{3})\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}}(4x^3 + e^{3c_1})^3 + e^{6c_1}} \right)$$

2.4 problem 11

Internal problem ID [6789]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

Problem number: 11.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[_1st_order, _with_linear_symmetries], _Clairaut]`

$$y'^2 - y'x + y = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)^2-x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x^2}{4}$$

$$y(x) = -c_1^2 + c_1x$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 25

```
DSolve[(y'[x])^2-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1(x - c_1)$$

$$y(x) \rightarrow \frac{x^2}{4}$$

2.5 problem 12

Internal problem ID [6790]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

Problem number: 12.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$y'^2 + 4x^5y' - 12x^4y = 0$$

✓ Solution by Maple

Time used: 0.219 (sec). Leaf size: 23

```
dsolve(diff(y(x),x)^2+4*x^5*diff(y(x),x)-12*x^4*y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{x^6}{3}$$

$$y(x) = c_1x^3 + \frac{3}{4}c_1^2$$

✓ Solution by Mathematica

Time used: 1.361 (sec). Leaf size: 217

`DSolve[(y'[x])^2+4*x^5*y'[x]-12*x^4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$\text{Solve} \left[\frac{1}{6} \left(\log(y(x)) - \frac{x^2 \sqrt{x^6 + 3y(x)} \log(y(x))}{\sqrt{x^4 (x^6 + 3y(x))}} \right) + \frac{x^2 \sqrt{x^6 + 3y(x)} \log(\sqrt{x^6 + 3y(x)} + x^3)}{3\sqrt{x^4 (x^6 + 3y(x))}} = c_1, y(x) \right]$$

$$\text{Solve} \left[\frac{1}{6} \left(\frac{x^2 \sqrt{x^6 + 3y(x)} \log(y(x))}{\sqrt{x^4 (x^6 + 3y(x))}} + \log(y(x)) \right) - \frac{x^2 \sqrt{x^6 + 3y(x)} \log(\sqrt{x^6 + 3y(x)} + x^3)}{3\sqrt{x^4 (x^6 + 3y(x))}} = c_1, y(x) \right]$$

$$y(x) \rightarrow -\frac{x^6}{3}$$

2.6 problem 13

Internal problem ID [6791]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

Problem number: 13.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _rational]`

$$4y^3y'^2 - 4y'x + y = 0$$

✓ Solution by Maple

Time used: 0.265 (sec). Leaf size: 81

```
dsolve(4*y(x)^3*diff(y(x),x)^2-4*x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \sqrt{-x}$$

$$y(x) = -\sqrt{-x}$$

$$y(x) = \sqrt{x}$$

$$y(x) = -\sqrt{x}$$

$$y(x) = 0$$

$$y(x) = \text{RootOf} \left(-\ln(x) + \int^{-z} -\frac{2(-a^4 + \sqrt{-a^4 + 1} - 1)}{-a(-a^4 - 1)} da + c_1 \right) \sqrt{x}$$

✓ Solution by Mathematica

Time used: 0.587 (sec). Leaf size: 282

```
DSolve[4*y[x]^3*(y'[x])^2-4*x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -e^{\frac{c_1}{4}} \sqrt[4]{e^{c_1} - 2ix}$$

$$y(x) \rightarrow -ie^{\frac{c_1}{4}} \sqrt[4]{e^{c_1} - 2ix}$$

$$y(x) \rightarrow ie^{\frac{c_1}{4}} \sqrt[4]{e^{c_1} - 2ix}$$

$$y(x) \rightarrow e^{\frac{c_1}{4}} \sqrt[4]{e^{c_1} - 2ix}$$

$$y(x) \rightarrow -e^{\frac{c_1}{4}} \sqrt[4]{2ix + e^{c_1}}$$

$$y(x) \rightarrow -ie^{\frac{c_1}{4}} \sqrt[4]{2ix + e^{c_1}}$$

$$y(x) \rightarrow ie^{\frac{c_1}{4}} \sqrt[4]{2ix + e^{c_1}}$$

$$y(x) \rightarrow e^{\frac{c_1}{4}} \sqrt[4]{2ix + e^{c_1}}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -\sqrt{x}$$

$$y(x) \rightarrow -i\sqrt{x}$$

$$y(x) \rightarrow i\sqrt{x}$$

$$y(x) \rightarrow \sqrt{x}$$

2.7 problem 14

Internal problem ID [6792]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

Problem number: 14.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _rational]`

$$4y^3y'^2 + 4y'x + y = 0$$

✓ Solution by Maple

Time used: 0.297 (sec). Leaf size: 287

```
dsolve(4*y(x)^3*diff(y(x),x)^2+4*x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$\int_{-b}^x \frac{-2a + \sqrt{-y(x)^4 + a^2}}{2y(x)^4 + 6a^2} da + \int^{y(x)} \left(\frac{-f^\beta}{-f^4 + \sqrt{-f^4 + x^2} x - x^2} - \left(\int_{-b}^x \left(-\frac{-f^\beta}{\sqrt{-f^4 + a^2} (f^4 + 3a^2)} - \frac{2(-2a + \sqrt{-f^4 + a^2}) - f^\beta}{(f^4 + 3a^2)^2} \right) da \right) \right) df + c_1 = 0$$

$$\int_{-b}^x \frac{-2a + \sqrt{-y(x)^4 + a^2}}{2(y(x)^4 + 3a^2)} da + \int^{y(x)} \left(-\frac{-f^\beta}{-f^4 + \sqrt{-f^4 + x^2} x + x^2} - \left(\int_{-b}^x \left(\frac{-f^\beta}{\sqrt{-f^4 + a^2} (f^4 + 3a^2)} + \frac{2(2a + \sqrt{-f^4 + a^2}) - f^\beta}{(f^4 + 3a^2)^2} \right) da \right) \right) df + c_1 = 0$$

✓ Solution by Mathematica

Time used: 60.284 (sec). Leaf size: 2815

```
DSolve[4*y[x]^3*(y'[x])^2+4*x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

2.8 problem 15

Internal problem ID [6793]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

Problem number: 15.

ODE order: 1.

ODE degree: 3.

CAS Maple gives this as type [`dAlembert`]

$$y'^3 + xy'^2 - y = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 1473

```
dsolve(diff(y(x),x)^3+x*diff(y(x),x)^2-y(x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x)$$

$$= \left(\frac{\left(-36x^2 - 54x + 108c_1 - 8x^3 + 27 + 6\sqrt{-48c_1x^3 - 216c_1x^2 - 24x^3 + 324c_1^2 - 324c_1x - 108x^2 + 162}\right)}{6}$$

$$+ x \left(\frac{\left(-36x^2 - 54x + 108c_1 - 8x^3 + 27 + 6\sqrt{-48c_1x^3 - 216c_1x^2 - 24x^3 + 324c_1^2 - 324c_1x - 108x^2 + 162}\right)}{6}$$

$$y(x)$$

$$= \left(- \frac{\left(-36x^2 - 54x + 108c_1 - 8x^3 + 27 + 6\sqrt{-48c_1x^3 - 216c_1x^2 - 24x^3 + 324c_1^2 - 324c_1x - 108x^2 + 162}\right)}{12}$$

$$+ x \left(- \frac{\left(-36x^2 - 54x + 108c_1 - 8x^3 + 27 + 6\sqrt{-48c_1x^3 - 216c_1x^2 - 24x^3 + 324c_1^2 - 324c_1x - 108x^2 + 162}\right)}{12}$$

$$y(x)$$

$$= \left(- \frac{\left(-36x^2 - 54x + 108c_1 - 8x^3 + 27 + 6\sqrt{-48c_1x^3 - 216c_1x^2 - 24x^3 + 324c_1^2 - 324c_1x - 108x^2 + 162}\right)}{12}$$

$$+ x \left(- \frac{\left(-36x^2 - 54x + 108c_1 - 8x^3 + 27 + 6\sqrt{-48c_1x^3 - 216c_1x^2 - 24x^3 + 324c_1^2 - 324c_1x - 108x^2 + 162}\right)}{12}$$

✓ Solution by Mathematica

Time used: 84.497 (sec). Leaf size: 1516

`DSolve[(y'[x])^3+x*(y'[x])^2-y[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{-16x^4 + 8 \left(\sqrt[3]{-8x^3 - 36x^2 - 54x + 108c_1 + 6\sqrt{6}\sqrt{-((4x^3 + 18x^2 + 27x - 27c_1)(2c_1 + 1))} + 27 - 12 \right)}{\dots}$$

$$y(x) \rightarrow \frac{1}{6} \left(\frac{i(\sqrt{3} - i)x(2x + 3)^2}{\sqrt[3]{-8x^3 - 36x^2 + 6\sqrt{6}\sqrt{-((1 + 2c_1)(4x^3 + 18x^2 + 27x - 27c_1))} - 54x + 27 + 108c_1}} \right. \\ \left. + \frac{1}{16} \left(\frac{i(\sqrt{3} - i)(2x + 3)^2}{\sqrt[3]{-8x^3 - 36x^2 + 6\sqrt{6}\sqrt{-((1 + 2c_1)(4x^3 + 18x^2 + 27x - 27c_1))} - 54x + 27 + 108c_1}} \right. \right. \\ \left. \left. + i(\sqrt{3} + i) \sqrt[3]{-8x^3 - 36x^2 + 6\sqrt{6}\sqrt{-((1 + 2c_1)(4x^3 + 18x^2 + 27x - 27c_1))} - 54x + 27 + 108c_1} \right. \right. \\ \left. \left. - 4x + 6 \right)^2 + i(\sqrt{3} + i) x \sqrt[3]{-8x^3 - 36x^2 + 6\sqrt{6}\sqrt{-((1 + 2c_1)(4x^3 + 18x^2 + 27x - 27c_1))} - 54x + 27 + 108c_1} \right. \\ \left. + 2(3 - 2x)x - 6x + 6c_1 \right)$$

$$y(x) \rightarrow \frac{1}{6} \left(\frac{i(\sqrt{3} + i)x(2x + 3)^2}{\sqrt[3]{-8x^3 - 36x^2 + 6\sqrt{6}\sqrt{-((1 + 2c_1)(4x^3 + 18x^2 + 27x - 27c_1))} - 54x + 27 + 108c_1}} \right. \\ \left. + \frac{1}{16} \left(\frac{(1 - i\sqrt{3})(2x + 3)^2}{\sqrt[3]{-8x^3 - 36x^2 + 6\sqrt{6}\sqrt{-((1 + 2c_1)(4x^3 + 18x^2 + 27x - 27c_1))} - 54x + 27 + 108c_1}} \right. \right. \\ \left. \left. + (1 + i\sqrt{3}) \sqrt[3]{-8x^3 - 36x^2 + 6\sqrt{6}\sqrt{-((1 + 2c_1)(4x^3 + 18x^2 + 27x - 27c_1))} - 54x + 27 + 108c_1} \right. \right. \\ \left. \left. + 4x - 6 \right)^2 - (1 + i\sqrt{3}) x \sqrt[3]{-8x^3 - 36x^2 + 6\sqrt{6}\sqrt{-((1 + 2c_1)(4x^3 + 18x^2 + 27x - 27c_1))} - 54x + 27 + 108c_1} \right. \\ \left. + 2(3 - 2x)x - 6x + 6c_1 \right)$$

2.9 problem 16

Internal problem ID [6794]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 97. The p-discriminant equation. EXERCISES Page 314

Problem number: 16.

ODE order: 1.

ODE degree: 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$y^4 y' - 6y'x + 2y = 0$$

✓ Solution by Maple

Time used: 0.25 (sec). Leaf size: 183

```
dsolve(y(x)^4*diff(y(x),x)^3-6*x*diff(y(x),x)+2*y(x)=0,y(x), singsol=all)
```

$$y(x) = \sqrt{-ix\sqrt{3} - x}$$

$$y(x) = \sqrt{ix\sqrt{3} - x}$$

$$y(x) = -\sqrt{-ix\sqrt{3} - x}$$

$$y(x) = -\sqrt{ix\sqrt{3} - x}$$

$$y(x) = \sqrt{x} \sqrt{2}$$

$$y(x) = -\sqrt{x} \sqrt{2}$$

$$y(x) = 0$$

$$y(x) = \frac{(-4c_1^3 + 24c_1x)^{\frac{1}{3}}}{2}$$

$$y(x) = -\frac{(-4c_1^3 + 24c_1x)^{\frac{1}{3}}}{4} - \frac{i\sqrt{3}(-4c_1^3 + 24c_1x)^{\frac{1}{3}}}{4}$$

$$y(x) = -\frac{(-4c_1^3 + 24c_1x)^{\frac{1}{3}}}{4} + \frac{i\sqrt{3}(-4c_1^3 + 24c_1x)^{\frac{1}{3}}}{4}$$

✓ Solution by Mathematica

Time used: 70.054 (sec). Leaf size: 22649

```
DSolve[y[x]^4*(y'[x])^3-6*x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

**3 CHAPTER 16. Nonlinear equations. Section 99.
Clairaut's equation. EXERCISES Page 320**

3.1	problem 3	48
3.2	problem 4	50
3.3	problem 5	52
3.4	problem 6	53
3.5	problem 7	54
3.6	problem 8	55
3.7	problem 9	57
3.8	problem 10	59
3.9	problem 11	60
3.10	problem 12	62
3.11	problem 13	63
3.12	problem 14	64
3.13	problem 15	65
3.14	problem 16	66
3.15	problem 17	68
3.16	problem 19	70
3.17	problem 20	72
3.18	problem 21	73
3.19	problem 22	74
3.20	problem 23	75
3.21	problem 24	77
3.22	problem 25	78
3.23	problem 26	80
3.24	problem 27	81
3.25	problem 28	83
3.26	problem 29	85

3.1 problem 3

Internal problem ID [6795]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 3.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$y'^2 + x^3 y' - 2x^2 y = 0$$

✓ Solution by Maple

Time used: 0.219 (sec). Leaf size: 23

```
dsolve(diff(y(x),x)^2+x^3*diff(y(x),x)-2*x^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{x^4}{8}$$

$$y(x) = c_1 x^2 + 2c_1^2$$

✓ Solution by Mathematica

Time used: 1.255 (sec). Leaf size: 209

```
DSolve[(y'[x])^2+x^3*y'[x]-2*x^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{\sqrt{x^6 + 8x^2y(x)} \log(\sqrt{x^4 + 8y(x)} + x^2)}{2x\sqrt{x^4 + 8y(x)}} + \frac{1}{4} \left(1 - \frac{\sqrt{x^6 + 8x^2y(x)}}{x\sqrt{x^4 + 8y(x)}} \right) \log(y(x)) = c_1, y(x) \right]$$
$$\text{Solve} \left[\frac{1}{4} \left(\frac{\sqrt{x^6 + 8x^2y(x)}}{x\sqrt{x^4 + 8y(x)}} + 1 \right) \log(y(x)) - \frac{\sqrt{x^6 + 8x^2y(x)} \log(\sqrt{x^4 + 8y(x)} + x^2)}{2x\sqrt{x^4 + 8y(x)}} = c_1, y(x) \right]$$
$$y(x) \rightarrow -\frac{x^4}{8}$$

3.2 problem 4

Internal problem ID [6796]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 4.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$y'^2 + 4x^5y' - 12x^4y = 0$$

✓ Solution by Maple

Time used: 0.219 (sec). Leaf size: 23

```
dsolve(diff(y(x),x)^2+4*x^5*diff(y(x),x)-12*x^4*y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{x^6}{3}$$

$$y(x) = c_1x^3 + \frac{3}{4}c_1^2$$

✓ Solution by Mathematica

Time used: 0.603 (sec). Leaf size: 217

```
DSolve[(y'[x])^2+4*x^5*y'[x]-12*x^4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{1}{6} \left(\log(y(x)) - \frac{x^2 \sqrt{x^6 + 3y(x)} \log(y(x))}{\sqrt{x^4 (x^6 + 3y(x))}} \right) + \frac{x^2 \sqrt{x^6 + 3y(x)} \log(\sqrt{x^6 + 3y(x)} + x^3)}{3\sqrt{x^4 (x^6 + 3y(x))}} = c_1, y(x) \right]$$

$$\text{Solve} \left[\frac{1}{6} \left(\frac{x^2 \sqrt{x^6 + 3y(x)} \log(y(x))}{\sqrt{x^4 (x^6 + 3y(x))}} + \log(y(x)) \right) - \frac{x^2 \sqrt{x^6 + 3y(x)} \log(\sqrt{x^6 + 3y(x)} + x^3)}{3\sqrt{x^4 (x^6 + 3y(x))}} = c_1, y(x) \right]$$

$$y(x) \rightarrow -\frac{x^6}{3}$$

3.3 problem 5

Internal problem ID [6797]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 5.

ODE order: 1.

ODE degree: 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$2xy'^3 - 6yy'^2 = -x^4$$

✓ Solution by Maple

Time used: 0.297 (sec). Leaf size: 56

```
dsolve(2*x*diff(y(x),x)^3-6*y(x)*diff(y(x),x)^2+x^4=0,y(x), singsol=all)
```

$$y(x) = \frac{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x^2}{2}$$

$$y(x) = \frac{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x^2}{2}$$

$$y(x) = \frac{x^2}{2}$$

$$y(x) = \frac{1}{6c_1^2} + \frac{c_1x^3}{3}$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[2*x*(y'[x])^3-6*y[x]*(y'[x])^2+x^4==0,y[x],x,IncludeSingularSolutions -> True]
```

Timed out

3.4 problem 6

Internal problem ID [6798]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES

Page 320

Problem number: 6.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[_1st_order, _with_linear_symmetries], _Clairaut]`

$$y'^2 - y'x + y = 0$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)^2-x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x^2}{4}$$

$$y(x) = -c_1^2 + c_1x$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 25

```
DSolve[(y'[x])^2-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1(x - c_1)$$

$$y(x) \rightarrow \frac{x^2}{4}$$

3.5 problem 7

Internal problem ID [6799]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 7.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[_1st_order, _with_linear_symmetries], _Clairaut]`

$$y - y'x - ky'^2 = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 24

```
dsolve(y(x)=diff(y(x),x)*x+k*diff(y(x),x)^2,y(x), singsol=all)
```

$$y(x) = -\frac{x^2}{4k}$$

$$y(x) = c_1^2 k + c_1 x$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 28

```
DSolve[y[x]==y'[x]*x+k*(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1(x + c_1 k)$$

$$y(x) \rightarrow -\frac{x^2}{4k}$$

3.6 problem 8

Internal problem ID [6800]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 8.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_homogeneous, 'class G']]`

$$x^8 y'^2 + 3y'x + 9y = 0$$

✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 42

```
dsolve(x^8*diff(y(x),x)^2+3*x*diff(y(x),x)+9*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{1}{4x^6}$$

$$y(x) = \frac{-x^3 + c_1}{x^3 c_1^2}$$

$$y(x) = -\frac{x^3 + c_1}{x^3 c_1^2}$$

✓ Solution by Mathematica

Time used: 0.583 (sec). Leaf size: 130

```
DSolve[x^8*(y'[x])^2+3*x*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{x\sqrt{4x^6y(x)-1} \arctan\left(\sqrt{4x^6y(x)-1}\right)}{3\sqrt{x^2-4x^8y(x)}} - \frac{1}{6} \log(y(x)) = c_1, y(x) \right]$$

$$\text{Solve} \left[\frac{\sqrt{x^2-4x^8y(x)} \arctan\left(\sqrt{4x^6y(x)-1}\right)}{3x\sqrt{4x^6y(x)-1}} - \frac{1}{6} \log(y(x)) = c_1, y(x) \right]$$

$$y(x) \rightarrow 0$$

3.7 problem 9

Internal problem ID [6801]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 9.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational]`

$$x^4 y'^2 + 2y' y x^3 = 4$$

✓ Solution by Maple

Time used: 0.109 (sec). Leaf size: 49

```
dsolve(x^4*diff(y(x),x)^2+2*x^3*y(x)*diff(y(x),x)-4=0,y(x), singsol=all)
```

$$y(x) = -\frac{2i}{x}$$

$$y(x) = \frac{2i}{x}$$

$$y(x) = \frac{2 \sinh(-\ln(x) + c_1)}{x}$$

$$y(x) = -\frac{2 \sinh(-\ln(x) + c_1)}{x}$$

✓ Solution by Mathematica

Time used: 0.688 (sec). Leaf size: 71

```
DSolve[x^4*(y'[x])^2+2*x^3*y[x]*y'[x]-4==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{4e^{c_1}}{x^2} - \frac{e^{-c_1}}{4}$$

$$y(x) \rightarrow \frac{e^{-c_1}}{4} - \frac{4e^{c_1}}{x^2}$$

$$y(x) \rightarrow -\frac{2i}{x}$$

$$y(x) \rightarrow \frac{2i}{x}$$

3.8 problem 10

Internal problem ID [6802]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 10.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _dAlembert]`

$$xy'^2 - 2y'y = -4x$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 29

```
dsolve(x*diff(y(x),x)^2-2*y(x)*diff(y(x),x)+4*x=0,y(x), singsol=all)
```

$$y(x) = -2x$$

$$y(x) = 2x$$

$$y(x) = -\frac{\left(-\frac{x^2}{c_1^2} - 4\right) c_1}{2}$$

✓ Solution by Mathematica

Time used: 0.202 (sec). Leaf size: 43

```
DSolve[x*(y'[x])^2-2*y[x]*y'[x]+4*x==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2x \cosh(-\log(x) + c_1)$$

$$y(x) \rightarrow -2x \cosh(\log(x) + c_1)$$

$$y(x) \rightarrow -2x$$

$$y(x) \rightarrow 2x$$

3.9 problem 11

Internal problem ID [6803]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 11.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational]`

$$3x^4y'^2 - y'x - y = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 147

```
dsolve(3*x^4*diff(y(x),x)^2-x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{1}{12x^2}$$

$$y(x) = \frac{-c_1(-c_1 + 2ix\sqrt{3}) - c_1^2 - 6x^2}{6x^2c_1^2}$$

$$y(x) = \frac{-c_1(-c_1 - 2ix\sqrt{3}) - c_1^2 - 6x^2}{6x^2c_1^2}$$

$$y(x) = \frac{c_1(c_1 + 2ix\sqrt{3}) - 6x^2 - c_1^2}{6c_1^2x^2}$$

$$y(x) = \frac{c_1(c_1 - 2ix\sqrt{3}) - 6x^2 - c_1^2}{6c_1^2x^2}$$

✓ Solution by Mathematica

Time used: 0.512 (sec). Leaf size: 123

```
DSolve[3*x^4*(y'[x])^2-x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[-\frac{x\sqrt{12x^2y(x)+1}\operatorname{arctanh}\left(\sqrt{12x^2y(x)+1}\right)}{\sqrt{12x^4y(x)+x^2}} - \frac{1}{2}\log(y(x)) = c_1, y(x) \right]$$

$$\text{Solve} \left[\frac{x\sqrt{12x^2y(x)+1}\operatorname{arctanh}\left(\sqrt{12x^2y(x)+1}\right)}{\sqrt{12x^4y(x)+x^2}} - \frac{1}{2}\log(y(x)) = c_1, y(x) \right]$$

$$y(x) \rightarrow 0$$

3.10 problem 12

Internal problem ID [6804]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 12.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _rational, _dAlembert]`

$$xy'^2 + (x - y)y' - y = -1$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 56

```
dsolve(x*diff(y(x),x)^2+(x-y(x))*diff(y(x),x)+1-y(x)=0,y(x), singsol=all)
```

$$y(x) = -x - 2\sqrt{x}$$

$$y(x) = -x + 2\sqrt{x}$$

$$y(x) = \frac{(-c_1^2 - c_1)x}{-1 - c_1} - \frac{1}{-1 - c_1}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 46

```
DSolve[x*(y'[x])^2+(x-y[x])*y'[x]+1-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x + \frac{1}{1 + c_1}$$

$$y(x) \rightarrow -x - 2\sqrt{x}$$

$$y(x) \rightarrow 2\sqrt{x} - x$$

3.11 problem 13

Internal problem ID [6805]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 13.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _rational, _Clairaut]`

$$y'(y'x - y + k) = -a$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 41

```
dsolve(diff(y(x),x)*( x*diff(y(x),x)-y(x)+k )+a=0,y(x), singsol=all)
```

$$y(x) = k - 2\sqrt{ax}$$

$$y(x) = k + 2\sqrt{ax}$$

$$y(x) = c_1x + \frac{c_1k + a}{c_1}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 58

```
DSolve[y'[x]*( x*y'[x]-y[x]+k )+a==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{a}{c_1} + k + c_1x$$

$$y(x) \rightarrow \text{Indeterminate}$$

$$y(x) \rightarrow k - 2\sqrt{a}\sqrt{x}$$

$$y(x) \rightarrow 2\sqrt{a}\sqrt{x} + k$$

3.12 problem 14

Internal problem ID [6806]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 14.

ODE order: 1.

ODE degree: 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$x^6 y'^3 - 3y'x - 3y = 0$$

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 32

```
dsolve(x^6*diff(y(x),x)^3-3*x*diff(y(x),x)-3*y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{2}{3x^{\frac{3}{2}}}$$

$$y(x) = \frac{2}{3x^{\frac{3}{2}}}$$

$$y(x) = \frac{c_1^3}{3} - \frac{c_1}{x}$$

✓ Solution by Mathematica

Time used: 136.42 (sec). Leaf size: 24834

```
DSolve[x^6*(y'[x])^3-3*x*y'[x]-3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

3.13 problem 15

Internal problem ID [6807]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 15.

ODE order: 1.

ODE degree: 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$y - x^6 y'^3 + y'x = 0$$

✓ Solution by Maple

Time used: 0.234 (sec). Leaf size: 36

```
dsolve(y(x)=x^6*diff(y(x),x)^3-x*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = -\frac{2\sqrt{3}}{9x^{\frac{3}{2}}}$$

$$y(x) = \frac{2\sqrt{3}}{9x^{\frac{3}{2}}}$$

$$y(x) = c_1^3 - \frac{c_1}{x}$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y[x]==x^6*(y'[x])^3-x*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

Timed out

3.14 problem 16

Internal problem ID [6808]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 16.

ODE order: 1.

ODE degree: 4.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$xy'^4 - 2yy'^3 = -12x^3$$

✓ Solution by Maple

Time used: 0.297 (sec). Leaf size: 62

```
dsolve(x*diff(y(x),x)^4-2*y(x)*diff(y(x),x)^3+12*x^3=0,y(x), singsol=all)
```

$$y(x) = -\frac{2\sqrt{-6x}x}{3}$$

$$y(x) = \frac{2\sqrt{-6x}x}{3}$$

$$y(x) = -\frac{2\sqrt{6}x^{\frac{3}{2}}}{3}$$

$$y(x) = \frac{2\sqrt{6}x^{\frac{3}{2}}}{3}$$

$$y(x) = 6c_1^3 + \frac{x^2}{2c_1}$$

✓ Solution by Mathematica

Time used: 37.824 (sec). Leaf size: 30947

```
DSolve[x*(y'[x])^4-2*y[x]*(y'[x])^3+12*x^3==0,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

3.15 problem 17

Internal problem ID [6809]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 17.

ODE order: 1.

ODE degree: 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _Clairaut]`

$$xy'^3 - yy'^2 = -1$$

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 80

```
dsolve(x*diff(y(x),x)^3-y(x)*diff(y(x),x)^2+1=0,y(x), singsol=all)
```

$$y(x) = \frac{3 \cdot 2^{\frac{1}{3}} (x^2)^{\frac{1}{3}}}{2}$$

$$y(x) = -\frac{3 \cdot 2^{\frac{1}{3}} (x^2)^{\frac{1}{3}}}{4} - \frac{3i\sqrt{3} \cdot 2^{\frac{1}{3}} (x^2)^{\frac{1}{3}}}{4}$$

$$y(x) = -\frac{3 \cdot 2^{\frac{1}{3}} (x^2)^{\frac{1}{3}}}{4} + \frac{3i\sqrt{3} \cdot 2^{\frac{1}{3}} (x^2)^{\frac{1}{3}}}{4}$$

$$y(x) = c_1 x + \frac{1}{c_1^2}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 69

```
DSolve[x*(y'[x])^3-y[x]*(y'[x])^2+1==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 x + \frac{1}{c_1^2}$$

$$y(x) \rightarrow 3 \left(-\frac{1}{2} \right)^{2/3} x^{2/3}$$

$$y(x) \rightarrow \frac{3x^{2/3}}{2^{2/3}}$$

$$y(x) \rightarrow -\frac{3\sqrt[3]{-1}x^{2/3}}{2^{2/3}}$$

3.16 problem 19

Internal problem ID [6810]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES

Page 320

Problem number: 19.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$y'^2 - y'x - y = 0$$

✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 77

```
dsolve(diff(y(x),x)^2-x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$\frac{c_1}{\sqrt{2x - 2\sqrt{x^2 + 4y(x)}}} + \frac{2x}{3} + \frac{\sqrt{x^2 + 4y(x)}}{3} = 0$$
$$\frac{c_1}{\sqrt{2x + 2\sqrt{x^2 + 4y(x)}}} + \frac{2x}{3} - \frac{\sqrt{x^2 + 4y(x)}}{3} = 0$$

✓ Solution by Mathematica

Time used: 60.129 (sec). Leaf size: 1003

`DSolve[(y'[x])^2 - x*y'[x] - y[x] == 0, y[x], x, IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{\left(x^2 + \sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}\right)^2 + 8e^{3c_1}x}{4\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}}$$

$$y(x) \rightarrow \frac{1}{8} \left(4x^2 - \frac{i(\sqrt{3} - i)x(x^3 + 8e^{3c_1})}{\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}} + i(\sqrt{3} + i)\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}} \right)$$

$$y(x) \rightarrow \frac{1}{8} \left(4x^2 + \frac{i(\sqrt{3} + i)x(x^3 + 8e^{3c_1})}{\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}}} - (1 + i\sqrt{3})\sqrt[3]{-x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}}(-x^3 + e^{3c_1})^3 + 8e^{6c_1}} \right)$$

$$y(x) \rightarrow \frac{2\sqrt[3]{2}x^4 + 2^{2/3}\left(-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}}(4x^3 + e^{3c_1})^3 + e^{6c_1}\right)^{2/3} + 4x^2\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}}(4x^3 + e^{3c_1})^3 + e^{6c_1}}}{8\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}}(4x^3 + e^{3c_1})^3 + e^{6c_1}}}$$

$$y(x) \rightarrow \frac{1}{16} \left(8x^2 + \frac{2\sqrt[3]{2}(1 + i\sqrt{3})x(-x^3 + 2e^{3c_1})}{\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}}(4x^3 + e^{3c_1})^3 + e^{6c_1}}} + i2^{2/3}(\sqrt{3} + i)\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}}(4x^3 + e^{3c_1})^3 + e^{6c_1}} \right)$$

$$y(x) \rightarrow \frac{1}{16} \left(8x^2 + \frac{2i\sqrt[3]{2}(\sqrt{3} + i)x(x^3 - 2e^{3c_1})}{\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}}(4x^3 + e^{3c_1})^3 + e^{6c_1}}} - 2^{2/3}(1 + i\sqrt{3})\sqrt[3]{-2x^6 - 10e^{3c_1}x^3 + \sqrt{e^{3c_1}}(4x^3 + e^{3c_1})^3 + e^{6c_1}} \right)$$

3.17 problem 20

Internal problem ID [6811]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 20.

ODE order: 1.

ODE degree: 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$2y'^3 + y'x - 2y = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 79

```
dsolve(2*diff(y(x),x)^3+x*diff(y(x),x)-2*y(x)=0,y(x), singsol=all)
```

$$y(x) = \left(-\frac{c_1}{12} - \frac{\sqrt{c_1^2 + 24x}}{12} \right)^3 + \frac{\left(-\frac{c_1}{12} - \frac{\sqrt{c_1^2 + 24x}}{12} \right) x}{2}$$

$$y(x) = \left(-\frac{c_1}{12} + \frac{\sqrt{c_1^2 + 24x}}{12} \right)^3 + \frac{\left(-\frac{c_1}{12} + \frac{\sqrt{c_1^2 + 24x}}{12} \right) x}{2}$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[2*(y'[x])^3+x*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Timed out

3.18 problem 21

Internal problem ID [6812]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

Problem number: 21.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$2y'^2 + y'x - 2y = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 36

```
dsolve(2*diff(y(x),x)^2+x*diff(y(x),x)-2*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{2 \operatorname{LambertW}\left(\frac{x e^{\frac{c_1}{4}}}{4}\right) - \frac{c_1}{2}} + \frac{e^{\operatorname{LambertW}\left(\frac{x e^{\frac{c_1}{4}}}{4}\right) - \frac{c_1}{4}} x}{2}$$

✓ Solution by Mathematica

Time used: 1.194 (sec). Leaf size: 130

```
DSolve[2*(y'[x])^2+x*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[-\frac{\frac{1}{2}x\sqrt{x^2+16y(x)} - 8y(x)\log\left(\sqrt{x^2+16y(x)} - x\right) + \frac{x^2}{2}}{8y(x)} = c_1, y(x) \right]$$

$$\text{Solve} \left[\frac{\frac{1}{2}x\sqrt{x^2+16y(x)} - 8y(x)\log\left(\sqrt{x^2+16y(x)} - x\right) - \frac{x^2}{2}}{8y(x)} + \log(y(x)) = c_1, y(x) \right]$$

$y(x) \rightarrow 0$

3.19 problem 22

Internal problem ID [6813]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 22.

ODE order: 1.

ODE degree: 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$y'^3 + 2y'x - y = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 173

```
dsolve(diff(y(x),x)^3+2*x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{(-6\sqrt{x^2 + 3c_1} - 6x)^{\frac{3}{2}}}{27} - \frac{2\sqrt{-6\sqrt{x^2 + 3c_1} - 6x}x}{3}$$

$$y(x) = \frac{(-6\sqrt{x^2 + 3c_1} - 6x)^{\frac{3}{2}}}{27} + \frac{2\sqrt{-6\sqrt{x^2 + 3c_1} - 6x}x}{3}$$

$$y(x) = -\frac{(6\sqrt{x^2 + 3c_1} - 6x)^{\frac{3}{2}}}{27} - \frac{2\sqrt{6\sqrt{x^2 + 3c_1} - 6x}x}{3}$$

$$y(x) = \frac{(6\sqrt{x^2 + 3c_1} - 6x)^{\frac{3}{2}}}{27} + \frac{2\sqrt{6\sqrt{x^2 + 3c_1} - 6x}x}{3}$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(y'[x])^3+2*x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Timed out

3.20 problem 23

Internal problem ID [6814]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 23.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational, _dAlembert]`

$$4xy'^2 - 3y'y = -3$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 153

```
dsolve(4*x*diff(y(x),x)^2-3*y(x)*diff(y(x),x)+3=0,y(x), singsol=all)
```

$$y(x) = -\frac{2\sqrt{x(3 + \sqrt{16c_1x + 9})}}{3} - \frac{2x}{\sqrt{x(3 + \sqrt{16c_1x + 9})}}$$

$$y(x) = \frac{2\sqrt{x(3 + \sqrt{16c_1x + 9})}}{3} + \frac{2x}{\sqrt{x(3 + \sqrt{16c_1x + 9})}}$$

$$y(x) = -\frac{2\sqrt{-x(-3 + \sqrt{16c_1x + 9})}}{3} - \frac{2x}{\sqrt{-x(-3 + \sqrt{16c_1x + 9})}}$$

$$y(x) = \frac{2\sqrt{-x(-3 + \sqrt{16c_1x + 9})}}{3} + \frac{2x}{\sqrt{-x(-3 + \sqrt{16c_1x + 9})}}$$

✓ Solution by Mathematica

Time used: 23.695 (sec). Leaf size: 187

```
DSolve[4*x*(y'[x])^2-3*y[x]*y'[x]+3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{432x - e^{-\frac{c_1}{2}} (-144x + e^{c_1})^{3/2} + e^{c_1}}}{6\sqrt{3}}$$

$$y(x) \rightarrow \frac{\sqrt{432x - e^{-\frac{c_1}{2}} (-144x + e^{c_1})^{3/2} + e^{c_1}}}{6\sqrt{3}}$$

$$y(x) \rightarrow -\frac{\sqrt{432x + e^{-\frac{c_1}{2}} (-144x + e^{c_1})^{3/2} + e^{c_1}}}{6\sqrt{3}}$$

$$y(x) \rightarrow \frac{\sqrt{432x + e^{-\frac{c_1}{2}} (-144x + e^{c_1})^{3/2} + e^{c_1}}}{6\sqrt{3}}$$

3.21 problem 24

Internal problem ID [6815]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 24.

ODE order: 1.

ODE degree: 3.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$y'^3 - y'x + 2y = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 83

```
dsolve(diff(y(x),x)^3-x*diff(y(x),x)+2*y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{\left(\frac{c_1}{6} - \frac{\sqrt{c_1^2 - 12x}}{6}\right)^3}{2} + \frac{\left(\frac{c_1}{6} - \frac{\sqrt{c_1^2 - 12x}}{6}\right)x}{2}$$
$$y(x) = -\frac{\left(\frac{c_1}{6} + \frac{\sqrt{c_1^2 - 12x}}{6}\right)^3}{2} + \frac{\left(\frac{c_1}{6} + \frac{\sqrt{c_1^2 - 12x}}{6}\right)x}{2}$$

✓ Solution by Mathematica

Time used: 29.375 (sec). Leaf size: 10134

```
DSolve[(y'[x])^3-x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

3.22 problem 25

Internal problem ID [6816]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320

Problem number: 25.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$5y'^2 + 6y'x - 2y = 0$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 85

```
dsolve(5*diff(y(x),x)^2+6*x*diff(y(x),x)-2*y(x)=0,y(x), singsol=all)
```

$$\frac{c_1}{\left(-15x - 5\sqrt{9x^2 + 10y(x)}\right)^{\frac{3}{2}}} + \frac{2x}{5} - \frac{\sqrt{9x^2 + 10y(x)}}{5} = 0$$

$$\frac{c_1}{\left(-15x + 5\sqrt{9x^2 + 10y(x)}\right)^{\frac{3}{2}}} + \frac{2x}{5} + \frac{\sqrt{9x^2 + 10y(x)}}{5} = 0$$

✓ Solution by Mathematica

Time used: 14.31 (sec). Leaf size: 771

```
DSolve[5*(y'[x])^2+6*x*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{Root}\left[4\#1^5 + 4\#1^4x^2 + \#1^3x^4 + 1000\#1^2e^{5c_1}x + 900\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 25000e^{10c_1}\&, 1\right]$$

$$y(x) \rightarrow \text{Root}\left[4\#1^5 + 4\#1^4x^2 + \#1^3x^4 + 1000\#1^2e^{5c_1}x + 900\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 25000e^{10c_1}\&, 2\right]$$

$$y(x) \rightarrow \text{Root}\left[4\#1^5 + 4\#1^4x^2 + \#1^3x^4 + 1000\#1^2e^{5c_1}x + 900\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 25000e^{10c_1}\&, 3\right]$$

$$y(x) \rightarrow \text{Root}\left[4\#1^5 + 4\#1^4x^2 + \#1^3x^4 + 1000\#1^2e^{5c_1}x + 900\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 25000e^{10c_1}\&, 4\right]$$

$$y(x) \rightarrow \text{Root}\left[4\#1^5 + 4\#1^4x^2 + \#1^3x^4 + 1000\#1^2e^{5c_1}x + 900\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 25000e^{10c_1}\&, 5\right]$$

$$y(x) \rightarrow \text{Root}\left[100000\#1^5 + 100000\#1^4x^2 + 25000\#1^3x^4 - 1000\#1^2e^{5c_1}x - 900\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 1\right]$$

$$y(x) \rightarrow \text{Root}\left[100000\#1^5 + 100000\#1^4x^2 + 25000\#1^3x^4 - 1000\#1^2e^{5c_1}x - 900\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 2\right]$$

$$y(x) \rightarrow \text{Root}\left[100000\#1^5 + 100000\#1^4x^2 + 25000\#1^3x^4 - 1000\#1^2e^{5c_1}x - 900\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 3\right]$$

$$y(x) \rightarrow \text{Root}\left[100000\#1^5 + 100000\#1^4x^2 + 25000\#1^3x^4 - 1000\#1^2e^{5c_1}x - 900\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 4\right]$$

$$y(x) \rightarrow \text{Root}\left[100000\#1^5 + 100000\#1^4x^2 + 25000\#1^3x^4 - 1000\#1^2e^{5c_1}x - 900\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 5\right]$$

$$y(x) \rightarrow 0$$

3.23 problem 26

Internal problem ID [6817]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES Page 320

Problem number: 26.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[_rational, _dAlembert]`

$$2xy'^2 + (2x - y)y' - y = -1$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 146

```
dsolve(2*x*diff(y(x),x)^2+(2*x-y(x))*diff(y(x),x)+1-y(x)=0,y(x), singsol=all)
```

$$y(x) = - \left(-2 \left(e^{\text{RootOf}(-e^3 - Zx + 2e^2 - Zx + c_1 e^{-Z} + Ze^{-Z} - x e^{-Z} + 1)} - 1 \right)^2 - 2 e^{\text{RootOf}(-e^3 - Zx + 2e^2 - Zx + c_1 e^{-Z} + Ze^{-Z} - x e^{-Z} + 1)} + 2 \right) e^{-\text{RootOf}(-e^3 - Zx + 2e^2 - Zx + c_1 e^{-Z} + Ze^{-Z} - x e^{-Z} + 1)} x + e^{-\text{RootOf}(-e^3 - Zx + 2e^2 - Zx + c_1 e^{-Z} + Ze^{-Z} - x e^{-Z} + 1)}$$

✓ Solution by Mathematica

Time used: 1.438 (sec). Leaf size: 49

```
DSolve[2*x*(y'[x])^2+(2*x-y[x])*y'[x]+1-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\left\{ x = \frac{\frac{1}{K[1]+1} + \log(K[1] + 1)}{K[1]^2} + \frac{c_1}{K[1]^2}, y(x) = 2xK[1] + \frac{1}{K[1] + 1} \right\}, \{y(x), K[1]\} \right]$$

3.24 problem 27

Internal problem ID [6818]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES

Page 320

Problem number: 27.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$5y'^2 + 3y'x - y = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 85

```
dsolve(5*diff(y(x),x)^2+3*x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$\frac{c_1}{\left(-30x - 10\sqrt{9x^2 + 20y(x)}\right)^{\frac{3}{2}}} + \frac{2x}{5} - \frac{\sqrt{9x^2 + 20y(x)}}{5} = 0$$

$$\frac{c_1}{\left(-30x + 10\sqrt{9x^2 + 20y(x)}\right)^{\frac{3}{2}}} + \frac{2x}{5} + \frac{\sqrt{9x^2 + 20y(x)}}{5} = 0$$

✓ Solution by Mathematica

Time used: 14.529 (sec). Leaf size: 771

```
DSolve[5*(y'[x])^2+3*x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 8\#1^4x^2 + \#1^3x^4 + 4000\#1^2e^{5c_1}x + 1800\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 200000e^{10c_1}\&, 1\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 8\#1^4x^2 + \#1^3x^4 + 4000\#1^2e^{5c_1}x + 1800\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 200000e^{10c_1}\&, 2\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 8\#1^4x^2 + \#1^3x^4 + 4000\#1^2e^{5c_1}x + 1800\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 200000e^{10c_1}\&, 3\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 8\#1^4x^2 + \#1^3x^4 + 4000\#1^2e^{5c_1}x + 1800\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 200000e^{10c_1}\&, 4\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 8\#1^4x^2 + \#1^3x^4 + 4000\#1^2e^{5c_1}x + 1800\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 200000e^{10c_1}\&, 5\right]$$

$$y(x) \rightarrow \text{Root}\left[3200000\#1^5 + 1600000\#1^4x^2 + 200000\#1^3x^4 - 4000\#1^2e^{5c_1}x - 1800\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 1\right]$$

$$y(x) \rightarrow \text{Root}\left[3200000\#1^5 + 1600000\#1^4x^2 + 200000\#1^3x^4 - 4000\#1^2e^{5c_1}x - 1800\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 2\right]$$

$$y(x) \rightarrow \text{Root}\left[3200000\#1^5 + 1600000\#1^4x^2 + 200000\#1^3x^4 - 4000\#1^2e^{5c_1}x - 1800\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 3\right]$$

$$y(x) \rightarrow \text{Root}\left[3200000\#1^5 + 1600000\#1^4x^2 + 200000\#1^3x^4 - 4000\#1^2e^{5c_1}x - 1800\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 4\right]$$

$$y(x) \rightarrow \text{Root}\left[3200000\#1^5 + 1600000\#1^4x^2 + 200000\#1^3x^4 - 4000\#1^2e^{5c_1}x - 1800\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 5\right]$$

$$y(x) \rightarrow 0$$

3.25 problem 28

Internal problem ID [6819]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES

Page 320

Problem number: 28.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _dAlembert]`

$$y'^2 + 3y'x - y = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 85

```
dsolve(diff(y(x),x)^2+3*x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$\frac{c_1}{\left(-6x - 2\sqrt{9x^2 + 4y(x)}\right)^{\frac{3}{2}}} + \frac{2x}{5} - \frac{\sqrt{9x^2 + 4y(x)}}{5} = 0$$

$$\frac{c_1}{\left(-6x + 2\sqrt{9x^2 + 4y(x)}\right)^{\frac{3}{2}}} + \frac{2x}{5} + \frac{\sqrt{9x^2 + 4y(x)}}{5} = 0$$

✓ Solution by Mathematica

Time used: 14.495 (sec). Leaf size: 776

```
DSolve[(y'[x])^2+3*x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 40\#1^4x^2 + 25\#1^3x^4 + 160\#1^2e^{5c_1}x + 360\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 64e^{10c_1}\&, 1\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 40\#1^4x^2 + 25\#1^3x^4 + 160\#1^2e^{5c_1}x + 360\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 64e^{10c_1}\&, 2\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 40\#1^4x^2 + 25\#1^3x^4 + 160\#1^2e^{5c_1}x + 360\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 64e^{10c_1}\&, 3\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 40\#1^4x^2 + 25\#1^3x^4 + 160\#1^2e^{5c_1}x + 360\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 64e^{10c_1}\&, 4\right]$$

$$y(x) \rightarrow \text{Root}\left[16\#1^5 + 40\#1^4x^2 + 25\#1^3x^4 + 160\#1^2e^{5c_1}x + 360\#1e^{5c_1}x^3 + 216e^{5c_1}x^5 - 64e^{10c_1}\&, 5\right]$$

$$y(x) \rightarrow \text{Root}\left[1024\#1^5 + 2560\#1^4x^2 + 1600\#1^3x^4 - 160\#1^2e^{5c_1}x - 360\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 1\right]$$

$$y(x) \rightarrow \text{Root}\left[1024\#1^5 + 2560\#1^4x^2 + 1600\#1^3x^4 - 160\#1^2e^{5c_1}x - 360\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 2\right]$$

$$y(x) \rightarrow \text{Root}\left[1024\#1^5 + 2560\#1^4x^2 + 1600\#1^3x^4 - 160\#1^2e^{5c_1}x - 360\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 3\right]$$

$$y(x) \rightarrow \text{Root}\left[1024\#1^5 + 2560\#1^4x^2 + 1600\#1^3x^4 - 160\#1^2e^{5c_1}x - 360\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 4\right]$$

$$y(x) \rightarrow \text{Root}\left[1024\#1^5 + 2560\#1^4x^2 + 1600\#1^3x^4 - 160\#1^2e^{5c_1}x - 360\#1e^{5c_1}x^3 - 216e^{5c_1}x^5 - e^{10c_1}\&, 5\right]$$

$$y(x) \rightarrow 0$$

3.26 problem 29

Internal problem ID [6820]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 99. Clairaut's equation. EXERCISES
Page 320


Problem number: 29.

ODE order: 1.

ODE degree: 2.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational]`

$$y - y'x - x^3y'^2 = 0$$

 Solution by Maple

```
dsolve(y(x)=x*diff(y(x),x)+x^3*diff(y(x),x)^2,y(x), singsol=all)
```

No solution found

 Solution by Mathematica

Time used: 105.562 (sec). Leaf size: 7052

```
DSolve[y[x]==x*y'[x]+x^3*(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

4 CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES

Page 324

4.1	problem 1	88
4.2	problem 2	89
4.3	problem 3	90
4.4	problem 4	91
4.5	problem 5	92
4.6	problem 6	93
4.7	problem 7	94
4.8	problem 9	95
4.9	problem 10	96
4.10	problem 11	97
4.11	problem 12	100
4.12	problem 13	101
4.13	problem 14	102
4.14	problem 15	103
4.15	problem 16	104
4.16	problem 17	105
4.17	problem 18	106
4.18	problem 19	107
4.19	problem 20	108
4.20	problem 21	109
4.21	problem 23	110
4.22	problem 24	111
4.23	problem 25	112
4.24	problem 26	113
4.25	problem 27	115
4.26	problem 28	116
4.27	problem 30	117
4.28	problem 31	118
4.29	problem 32	119
4.30	problem 33	121
4.31	problem 34	123
4.32	problem 35	124
4.33	problem 36	125
4.34	problem 37	126

4.35	problem 38	127
4.36	problem 39	128
4.37	problem 40	129
4.38	problem 41	130
4.39	problem 42	132
4.40	problem 43	134

4.1 problem 1

Internal problem ID [6821]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing.
EXERCISES Page 324

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]`

$$y'' - xy'^3 = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 37

```
dsolve(diff(y(x),x$2)=x*(diff(y(x),x))^3,y(x), singsol=all)
```

$$y(x) = \arctan\left(\frac{x}{\sqrt{-x^2 + c_1}}\right) + c_2$$

$$y(x) = -\arctan\left(\frac{x}{\sqrt{-x^2 + c_1}}\right) + c_2$$

✓ Solution by Mathematica

Time used: 10.922 (sec). Leaf size: 57

```
DSolve[y''[x]==x*(y'[x])^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \arctan\left(\frac{x}{\sqrt{-x^2 - 2c_1}}\right)$$

$$y(x) \rightarrow \arctan\left(\frac{x}{\sqrt{-x^2 - 2c_1}}\right) + c_2$$

$$y(x) \rightarrow c_2$$

4.2 problem 2

Internal problem ID [6822]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing.
EXERCISES Page 324

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^2 y'' + y'^2 - 2y'x = 0$$

With initial conditions

$$[y(2) = 5, y'(2) = -4]$$

✓ Solution by Maple

Time used: 0.171 (sec). Leaf size: 24

```
dsolve([x^2*diff(y(x),x$2)+diff(y(x),x)^2-2*x*diff(y(x),x)=0,y(2) = 5, D(y)(2) = -4],y(x), s
```

$$y(x) = \frac{x^2}{2} + 3x + 9 \ln(x - 3) - 3 - 9i\pi$$

✓ Solution by Mathematica

Time used: 0.478 (sec). Leaf size: 28

```
DSolve[{x^2*y'[x]+(y'[x])^2-2*x*y'[x]==0,{y[2]==5,y'[2]==-4}},y[x],x,IncludeSingularSolutio
```

$$y(x) \rightarrow \frac{x^2}{2} + 3x + 9 \log(x - 3) - 9i\pi - 3$$

4.3 problem 3

Internal problem ID [6823]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^2 y'' + y'^2 - 2y'x = 0$$

With initial conditions

$$[y(2) = 5, y'(2) = 2]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 11

```
dsolve([x^2*diff(y(x),x$2)+diff(y(x),x)^2-2*x*diff(y(x),x)=0,y(2) = 5, D(y)(2) = 2],y(x), si
```

$$y(x) = \frac{x^2}{2} + 3$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 14

```
DSolve[{x^2*y'[x]+(y'[x])^2-2*x*y'[x]==0,{y[2]==5,y'[2]==2}},y[x],x,IncludeSingularSolution
```

$$y(x) \rightarrow \frac{1}{2}(x^2 + 6)$$

4.4 problem 4

Internal problem ID [6824]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear],`

$$yy'' + y'^2 = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 33

```
dsolve(y(x)*diff(y(x),x$2)+diff(y(x),x)^2=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = \sqrt{2c_1x + 2c_2}$$

$$y(x) = -\sqrt{2c_1x + 2c_2}$$

✓ Solution by Mathematica

Time used: 0.172 (sec). Leaf size: 20

```
DSolve[y[x]*y'[x]+(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2\sqrt{2x - c_1}$$

4.5 problem 5

Internal problem ID [6825]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1],`

$$y^2 y'' + y'^3 = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 34

```
dsolve(y(x)^2*diff(y(x),x$2)+diff(y(x),x)^3=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1$$

$$y(x) = e^{-\text{LambertW}(-c_1 e^{-c_2 e^{-x}}) - c_2 - x}$$

✓ Solution by Mathematica

Time used: 0.609 (sec). Leaf size: 37

```
DSolve[y[x]^2*y'[x]+(y'[x])^3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 \left(1 + \frac{1}{\text{InverseFunction} \left[-\frac{1}{\#1} - \log(\#1) + \log(\#1 + 1) \right] [-x + c_1]} \right)$$

4.6 problem 6

Internal problem ID [6826]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible]`

$$(y + 1)y'' - y'^2 = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve((y(x)+1)*diff(y(x),x$2)=diff(y(x),x)^2,y(x), singsol=all)
```

$$y(x) = -1$$

$$y(x) = e^{c_1 x} c_2 - 1$$

✓ Solution by Mathematica

Time used: 1.193 (sec). Leaf size: 26

```
DSolve[(y[x]+1)*y'[x]==(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -1 + \frac{e^{c_1(x+c_2)}}{c_1}$$

$$y(x) \rightarrow \text{Indeterminate}$$

4.7 problem 7

Internal problem ID [6827]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_y_y1]]`

$$2ay'' + y'^3 = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 29

```
dsolve(2*a*diff(y(x),x$2)+diff(y(x),x)^3=0,y(x), singsol=all)
```

$$y(x) = 2\sqrt{(x + c_1)a} + c_2$$

$$y(x) = -2\sqrt{(x + c_1)a} + c_2$$

✓ Solution by Mathematica

Time used: 0.33 (sec). Leaf size: 51

```
DSolve[2*a*y''[x]+(y'[x])^3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - 2\sqrt{a}\sqrt{x - 2ac_1}$$

$$y(x) \rightarrow 2\sqrt{a}\sqrt{x - 2ac_1} + c_2$$

4.8 problem 9

Internal problem ID [6828]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' - y' = x^5$$

With initial conditions

$$\left[y(1) = \frac{1}{2}, y'(1) = 1 \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve([x*diff(y(x),x$2)=diff(y(x),x)+x^5,y(1) = 1/2, D(y)(1) = 1],y(x), singsol=all)
```

$$y(x) = \frac{1}{24}x^6 + \frac{3}{8}x^2 + \frac{1}{12}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 19

```
DSolve[{x*y''[x]==y'[x]+x^5,{y[1]==1/2,y'[1]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{24}(x^6 + 9x^2 + 2)$$

4.9 problem 10

Internal problem ID [6829]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing.
EXERCISES Page 324

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' + y' = -x$$

With initial conditions

$$\left[y(2) = -1, y'(2) = -\frac{1}{2} \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve([x*diff(y(x),x$2)+diff(y(x),x)+x=0,y(2) = -1, D(y)(2) = -1/2],y(x), singsol=all)
```

$$y(x) = -\frac{x^2}{4} + \ln(x) - \ln(2)$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 19

```
DSolve[{x*y''[x]+y'[x]+x==0,{y[2]==-1,y'[2]==-1/2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log\left(\frac{x}{2}\right) - \frac{x^2}{4}$$

4.10 problem 11

Internal problem ID [6830]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing.
EXERCISES Page 324

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1],`

$$y'' - 2yy'^3 = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 386

```
dsolve(diff(y(x),x$2)=2*y(x)*diff(y(x),x)^3,y(x), singsol=all)
```

$$y(x) = c_1$$

$$y(x) = \frac{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}{2} + \frac{2c_1}{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}$$

$$y(x)$$

$$= -\frac{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}{4c_1} - \frac{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}{2} + \frac{i\sqrt{3}\left(\frac{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}{2} - \frac{2c_1}{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}\right)}{2}$$

$$y(x)$$

$$= -\frac{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}{4c_1} - \frac{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}{2} + \frac{i\sqrt{3}\left(\frac{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}{2} - \frac{2c_1}{\left(-12c_2 - 12x + 4\sqrt{-4c_1^3 + 9c_2^2 + 18xc_2 + 9x^2}\right)^{\frac{1}{3}}}\right)}{2}$$

✓ Solution by Mathematica

Time used: 7.768 (sec). Leaf size: 351

```
DSolve[y''[x]==2*y[x]*(y'[x])^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt[3]{2}c_1}{\sqrt[3]{\sqrt{9x^2 + 18c_2x + 4c_1^3 + 9c_2^2} + 3x + 3c_2}} - \frac{\sqrt[3]{\sqrt{9x^2 + 18c_2x + 4c_1^3 + 9c_2^2} + 3x + 3c_2}}{\sqrt[3]{2}}$$

$$y(x) \rightarrow \frac{2^{2/3}(1 - i\sqrt{3}) (\sqrt{9x^2 + 18c_2x + 4c_1^3 + 9c_2^2} + 3x + 3c_2)^{2/3} + \sqrt[3]{2}(-2 - 2i\sqrt{3}) c_1}{4\sqrt[3]{\sqrt{9x^2 + 18c_2x + 4c_1^3 + 9c_2^2} + 3x + 3c_2}}$$

$$y(x) \rightarrow \frac{2^{2/3}(1 + i\sqrt{3}) (\sqrt{9x^2 + 18c_2x + 4c_1^3 + 9c_2^2} + 3x + 3c_2)^{2/3} + 2i\sqrt[3]{2}(\sqrt{3} + i) c_1}{4\sqrt[3]{\sqrt{9x^2 + 18c_2x + 4c_1^3 + 9c_2^2} + 3x + 3c_2}}$$

$$y(x) \rightarrow 0$$

4.11 problem 12

Internal problem ID [6831]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1],`

$$yy'' + y'^3 - y'^2 = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 44

```
dsolve(y(x)*diff(y(x),x$2)+diff(y(x),x)^3-diff(y(x),x)^2=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1$$

$$y(x) = e^{-\frac{c_1 \operatorname{LambertW}\left(\frac{e^{\frac{c_2}{c_1}} e^{\frac{x}{c_1}}}{c_1}\right) - c_2 - x}{c_1}}$$

✓ Solution by Mathematica

Time used: 22.067 (sec). Leaf size: 32

```
DSolve[y[x]*y'[x]+(y'[x])^3-(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{c_1 W\left(e^{e^{-c_1}(x - e^{c_1} c_1 + c_2)}\right)}$$

4.12 problem 13

Internal problem ID [6832]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + \beta^2 y = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+beta^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(\beta x) + c_2 \cos(\beta x)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 20

```
DSolve[y''[x]+\[Beta]^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(\beta x) + c_2 \sin(\beta x)$$

4.13 problem 14

Internal problem ID [6833]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1],`

$$yy'' + y'^3 = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 27

```
dsolve(y(x)*diff(y(x),x$2)+diff(y(x),x)^3=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1$$

$$y(x) = e^{\text{LambertW}((c_2+x)e^{c_1}e^{-1})-c_1+1}$$

✓ Solution by Mathematica

Time used: 60.095 (sec). Leaf size: 26

```
DSolve[y[x]*y'[x]+(y'[x])^3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x + c_2}{W(e^{-1-c_1}(x + c_2))}$$

4.14 problem 15

Internal problem ID [6834]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' \cos(x) - y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$2)*cos(x)=diff(y(x),x),y(x), singsol=all)
```

$$y(x) = c_1 + (\ln(\sec(x) + \tan(x)) - \ln(\cos(x))) c_2$$

✓ Solution by Mathematica

Time used: 0.181 (sec). Leaf size: 25

```
DSolve[y''[x]*Cos[x]==y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \log\left(e^{4\operatorname{arctanh}\left(\tan\left(\frac{x}{2}\right)\right)} + 1\right) + c_2$$

4.15 problem 16

Internal problem ID [6835]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing.
EXERCISES Page 324

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]`

$$y'' - xy'^2 = 0$$

With initial conditions

$$\left[y(2) = \frac{\pi}{4}, y'(2) = -\frac{1}{4} \right]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 8

```
dsolve([diff(y(x),x$2)=x*diff(y(x),x)^2,y(2) = 1/4*Pi, D(y)(2) = -1/4],y(x), singsol=all)
```

$$y(x) = \operatorname{arccot}\left(\frac{x}{2}\right)$$

✓ Solution by Mathematica

Time used: 1.241 (sec). Leaf size: 19

```
DSolve[{y'[x]==x*(y'[x])^2,{y[2]==1/4*Pi,y'[2]==-1/4}},y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow \frac{1}{2} \left(\pi - 2 \arctan\left(\frac{x}{2}\right) \right)$$

4.16 problem 17

Internal problem ID [6836]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing.
EXERCISES Page 324

Problem number: 17.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]`

$$y'' - xy'^2 = 0$$

With initial conditions

$$\left[y(0) = 1, y'(0) = \frac{1}{2} \right]$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 10

```
dsolve([diff(y(x),x$2)=x*diff(y(x),x)^2,y(0) = 1, D(y)(0) = 1/2],y(x), singsol=all)
```

$$y(x) = \operatorname{arctanh}\left(\frac{x}{2}\right) + 1$$

✓ Solution by Mathematica

Time used: 0.229 (sec). Leaf size: 13

```
DSolve[{y'[x]==x*(y'[x])^2,{y[0]==1,y'[0]==1/2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \operatorname{arctanh}\left(\frac{x}{2}\right) + 1$$

4.17 problem 18

Internal problem ID [6837]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing.
EXERCISES Page 324

Problem number: 18.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$y'' + e^{-2y} = 0$$

With initial conditions

$$[y(3) = 0, y'(3) = 1]$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 12

```
dsolve([diff(y(x),x$2)=-exp(-2*y(x)),y(3) = 0, D(y)(3) = 1],y(x), singsol=all)
```

$$y(x) = \frac{\ln((x-2)^2)}{2}$$

✓ Solution by Mathematica

Time used: 0.157 (sec). Leaf size: 9

```
DSolve[{y'[x]==-Exp[-2*y[x]],{y[3]==0,y'[3]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(x-2)$$

4.18 problem 19

Internal problem ID [6838]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing.
EXERCISES Page 324

Problem number: 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$y'' + e^{-2y} = 0$$

With initial conditions

$$[y(3) = 0, y'(3) = -1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve([diff(y(x),x$2)=-exp(-2*y(x)),y(3) = 0, D(y)(3) = -1],y(x), singsol=all)
```

$$y(x) = \frac{\ln((x-4)^2)}{2}$$

✓ Solution by Mathematica

Time used: 0.146 (sec). Leaf size: 11

```
DSolve[{y'[x]==-Exp[-2*y[x]],{y[3]==0,y'[3]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(4 - x)$$

4.19 problem 20

Internal problem ID [6839]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing.

EXERCISES Page 324

Problem number: 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$2y'' - \sin(2y) = 0$$

With initial conditions

$$\left[y(0) = \frac{\pi}{2}, y'(0) = 1 \right]$$

✓ Solution by Maple

Time used: 134.11 (sec). Leaf size: 1495

```
dsolve([2*diff(y(x),x$2)=sin(2*y(x)),y(0) = 1/2*Pi, D(y)(0) = 1],y(x), singsol=all)
```

Expression too large to display

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{2*y'[x]==Sin[2*y[x]],{y[0]==Pi/2,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True
```

```
{}
```

4.20 problem 21

Internal problem ID [6840]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing.
EXERCISES Page 324

Problem number: 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$2y'' - \sin(2y) = 0$$

With initial conditions

$$\left[y(0) = -\frac{\pi}{2}, y'(0) = 1 \right]$$

✓ Solution by Maple

Time used: 98.39 (sec). Leaf size: 1490

```
dsolve([2*diff(y(x),x$2)=sin(2*y(x)),y(0) = -1/2*Pi, D(y)(0) = 1],y(x), singsol=all)
```

Expression too large to display

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{2*y'[x]==Sin[2*y[x]],{y[0]==-Pi/2,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

{}

4.21 problem 23

Internal problem ID [6841]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^3 y'' - x^2 y' = -x^2 + 3$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(x^3*diff(y(x),x$2)-x^2*diff(y(x),x)=3-x^2,y(x), singsol=all)
```

$$y(x) = \frac{c_1 x^2}{2} + \frac{1}{x} + x + c_2$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 21

```
DSolve[x^3*y'[x]-x^2*y'[x]==3-x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1 x^2}{2} + x + \frac{1}{x} + c_2$$

4.22 problem 24

Internal problem ID [6842]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing.
EXERCISES Page 324

Problem number: 24.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible]`

$$y'' - y'^2 = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(diff(y(x),x$2)=diff(y(x),x)^2,y(x), singsol=all)
```

$$y(x) = -\ln(-c_1x - c_2)$$

✓ Solution by Mathematica

Time used: 0.197 (sec). Leaf size: 15

```
DSolve[y''[x]==(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \log(x + c_1)$$

4.23 problem 25

Internal problem ID [6843]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 25.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' - e^x y'^2 = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)=exp(x)*diff(y(x),x)^2,y(x), singsol=all)
```

$$y(x) = \frac{\ln(e^x)}{c_1} - \frac{\ln(e^x - c_1)}{c_1} + c_2$$

✓ Solution by Mathematica

Time used: 0.985 (sec). Leaf size: 37

```
DSolve[y''[x]==Exp[x](y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-x + \log(e^x + c_1) + c_1 c_2}{c_1}$$

$$y(x) \rightarrow \text{Indeterminate}$$

$$y(x) \rightarrow c_2$$

4.24 problem 26

Internal problem ID [6844]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 26.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]`

$$2y'' - y'^3 \sin(2x) = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 78

```
dsolve(2*diff(y(x),x$2)=diff(y(x),x)^3*sin(2*x),y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{-\left(\sin(x)^2 - \frac{1}{c_1^2}\right) c_1^2 \operatorname{InverseJacobiAM}(x, c_1)}}{\sqrt{-\sin(x)^2 + \frac{1}{c_1^2}}} + c_2$$
$$y(x) = -\frac{\sqrt{-\left(\sin(x)^2 - \frac{1}{c_1^2}\right) c_1^2 \operatorname{InverseJacobiAM}(x, c_1)}}{\sqrt{-\sin(x)^2 + \frac{1}{c_1^2}}} + c_2$$

✓ Solution by Mathematica

Time used: 6.102 (sec). Leaf size: 120

```
DSolve[2*y''[x]==(y'[x])^3*Sin[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \frac{\sqrt{-\frac{\cos(2x)+1-4c_1}{-1+2c_1}} \operatorname{EllipticF}\left(x, \frac{1}{1-2c_1}\right)}{\sqrt{\cos(2x)+1-4c_1}}$$

$$y(x) \rightarrow \frac{\sqrt{-\frac{\cos(2x)+1-4c_1}{-1+2c_1}} \operatorname{EllipticF}\left(x, \frac{1}{1-2c_1}\right)}{\sqrt{\cos(2x)+1-4c_1}} + c_2$$

$$y(x) \rightarrow c_2$$

4.25 problem 27

Internal problem ID [6845]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 27.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]`

$$x^2y'' + y'^2 = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 21

```
dsolve(x^2*diff(y(x),x$2)+diff(y(x),x)^2=0,y(x), singsol=all)
```

$$y(x) = \frac{x}{c_1} + \frac{\ln(c_1x - 1)}{c_1^2} + c_2$$

✓ Solution by Mathematica

Time used: 0.57 (sec). Leaf size: 47

```
DSolve[x^2*y'[x]+(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x}{c_1} + \frac{\log(1 + c_1x)}{c_1^2} + c_2$$

$$y(x) \rightarrow c_2$$

$$y(x) \rightarrow -\frac{x^2}{2} + c_2$$

4.26 problem 28

Internal problem ID [6846]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing.
EXERCISES Page 324

Problem number: 28.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_xy]]`

$$y'' - y'^2 = 1$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$2)=1+diff(y(x),x)^2,y(x), singsol=all)
```

$$y(x) = -\ln\left(\frac{-c_2 + \tan(x) c_1}{\sec(x)}\right)$$

✓ Solution by Mathematica

Time used: 1.97 (sec). Leaf size: 16

```
DSolve[y''[x]==1+(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \log(\cos(x + c_1))$$

4.27 problem 30

Internal problem ID [6847]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 30.

ODE order: 2.

ODE degree: 2.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - (1 + y'^2)^{\frac{3}{2}} = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 51

```
dsolve(diff(y(x),x$2)=(1+diff(y(x),x)^2)^(3/2),y(x), singsol=all)
```

$$y(x) = -ix + c_1$$

$$y(x) = ix + c_1$$

$$y(x) = (c_1 + x + 1)(c_1 + x - 1) \sqrt{-\frac{1}{c_1^2 + 2c_1x + x^2 - 1}} + c_2$$

✓ Solution by Mathematica

Time used: 0.269 (sec). Leaf size: 59

```
DSolve[y''[x]==(1+(y'[x])^2)^(3/2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - i\sqrt{x^2 + 2c_1x - 1 + c_1^2}$$

$$y(x) \rightarrow i\sqrt{x^2 + 2c_1x - 1 + c_1^2} + c_2$$

4.28 problem 31

Internal problem ID [6848]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 31.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_y_y1]]`

$$yy'' - y'^2(1 - y' \sin(y) - yy' \cos(y)) = 0$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 24

```
dsolve(y(x)*diff(y(x),x$2)=diff(y(x),x)^2*(1-diff(y(x),x)*sin(y(x))-y(x)*diff(y(x),x)*cos(y(x))),y(x),x)
```

$$y(x) = c_1$$
$$- \cos(y(x)) + c_1 \ln(y(x)) - x - c_2 = 0$$

✓ Solution by Mathematica

Time used: 0.489 (sec). Leaf size: 69

```
DSolve[y[x]*y'[x]==(y'[x])^2*(1-y'[x]*Sin[y[x]]-y[x]*y'[x]*Cos[y[x]]),y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \text{InverseFunction}[-\cos(\#1) + c_1 \log(\#1)\&][x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction}[-\cos(\#1) - c_1 \log(\#1)\&][x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction}[-\cos(\#1) + c_1 \log(\#1)\&][x + c_2]$$

4.29 problem 32

Internal problem ID [6849]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 32.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$(1 + y^2) y'' + y'^3 + y' = 0$$

✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 130

```
dsolve((1+y(x)^2)*diff(y(x),x$2)+diff(y(x),x)^3+diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -i$$

$$y(x) = i$$

$$y(x) = c_1$$

$$y(x) = \frac{i(c_1 - 1)}{1 + c_1}$$

$$e^{-\frac{-c_1^2 + 2c_1 + c_1^2 c_2 + x c_1^2 - 1 + 4 \operatorname{LambertW}\left(-\frac{i e^{-\frac{c_1 c_2}{4}} e^{-\frac{c_1 x}{4}} e^{\frac{c_1}{4}} e^{-\frac{c_2}{2}} e^{-\frac{x}{2}} e^{-\frac{1}{2}} e^{-\frac{c_2}{4 c_1}} e^{-\frac{x}{4 c_1}} e^{\frac{1}{4 c_1}} (c_1 - 1)}{4 c_1}\right) c_1 + 2 c_1 c_2 + 2 c_1 x + c_2 + x}{4 c_1}}}{1 + c_1}$$

✓ Solution by Mathematica

Time used: 57.998 (sec). Leaf size: 56

```
DSolve[(1+y[x]^2)*y'[x]+(y'[x])^3+y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \csc(c_1) \sec(c_1) W\left(\sin(c_1) e^{-((x+c_2) \cos^2(c_1)) - \sin^2(c_1)}\right) + \tan(c_1)$$

$$y(x) \rightarrow e^{-x-c_2}$$

4.30 problem 33

Internal problem ID [6850]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 33.

ODE order: 2.

ODE degree: 2.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$\left(yy'' + 1 + y'^2 \right)^2 - \left(1 + y'^2 \right)^3 = 0$$

✓ Solution by Maple

Time used: 0.14 (sec). Leaf size: 131

```
dsolve((y(x)*diff(y(x),x$2)+1+diff(y(x),x)^2)^2=(1+diff(y(x),x)^2)^3,y(x), singsol=all)
```

$$y(x) = -ix + c_1$$

$$y(x) = ix + c_1$$

$$y(x) = 0$$

$$y(x) = -c_1 - \sqrt{c_1^2 - c_2^2 - 2xc_2 - x^2}$$

$$y(x) = -c_1 + \sqrt{c_1^2 - c_2^2 - 2xc_2 - x^2}$$

$$y(x) = c_1 - \sqrt{c_1^2 - c_2^2 - 2xc_2 - x^2}$$

$$y(x) = c_1 + \sqrt{c_1^2 - c_2^2 - 2xc_2 - x^2}$$

✓ Solution by Mathematica

Time used: 45.659 (sec). Leaf size: 155

```
DSolve[(y[x]*y'[x]+1+(y'[x])^2)^2==(1+(y'[x])^2)^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{e^{2c_1} - (x + c_2)^2} - e^{c_1}$$

$$y(x) \rightarrow e^{c_1} - \sqrt{e^{2c_1} - (x + c_2)^2}$$

$$y(x) \rightarrow \sqrt{e^{2c_1} - (x + c_2)^2} - e^{c_1}$$

$$y(x) \rightarrow \sqrt{e^{2c_1} - (x + c_2)^2} + e^{c_1}$$

$$y(x) \rightarrow -\sqrt{-(x + c_2)^2}$$

$$y(x) \rightarrow \sqrt{-(x + c_2)^2}$$

4.31 problem 34

Internal problem ID [6851]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 34.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^2 y'' - y'(2x - y') = 0$$

With initial conditions

$$[y(-1) = 5, y'(-1) = 1]$$

✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 20

```
dsolve([x^2*diff(y(x),x$2)=diff(y(x),x)*(2*x-diff(y(x),x)),y(-1) = 5, D(y)(-1) = 1],y(x), si
```

$$y(x) = \frac{x^2}{2} - 2x + 4 \ln(x + 2) + \frac{5}{2}$$

✓ Solution by Mathematica

Time used: 0.52 (sec). Leaf size: 23

```
DSolve[{x^2*y''[x]==y'[x]*(2*x-y'[x]),{y[-1]==5,y'[-1]==1}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{1}{2}(x^2 - 4x + 8 \log(x + 2) + 5)$$

4.32 problem 35

Internal problem ID [6852]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 35.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^2 y'' - y'(3x - 2y') = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 22

```
dsolve(x^2*diff(y(x),x$2)=diff(y(x),x)*(3*x-2*diff(y(x),x)),y(x), singsol=all)
```

$$y(x) = \frac{x^2}{2} + \frac{c_1 \ln(x^2 - c_1)}{2} + c_2$$

✓ Solution by Mathematica

Time used: 0.333 (sec). Leaf size: 28

```
DSolve[x^2*y'[x]==y'[x]*(3*x-2*y'[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(x^2 - c_1 \log(x^2 + c_1) + 2c_2)$$

4.33 problem 36

Internal problem ID [6853]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 36.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _missing_y], _Liouville, [_2nd_order, _reducible]`

$$xy'' - y'(2 - 3y'/x) = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve(x*diff(y(x),x$2)=diff(y(x),x)*(2-3*x*diff(y(x),x)),y(x), singsol=all)
```

$$y(x) = \frac{\ln(c_1 x^3 + 3c_2)}{3}$$

✓ Solution by Mathematica

Time used: 0.267 (sec). Leaf size: 19

```
DSolve[x*y'[x]==y'[x]*(2-3*x*y'[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} \log(x^3 + c_1) + c_2$$

4.34 problem 37

Internal problem ID [6854]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 37.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^4 y'' - y'(y' + x^3) = 0$$

With initial conditions

$$[y(1) = 2, y'(1) = 1]$$

✓ Solution by Maple

Time used: 0.172 (sec). Leaf size: 25

```
dsolve([x^4*diff(y(x),x$2)=diff(y(x),x)*(diff(y(x),x)+x^3),y(1) = 2, D(y)(1) = 1],y(x), sing
```

$$y(x) = x^2 - \ln(-x^2 - 1) + 1 + \ln(2) + i\pi$$

✓ Solution by Mathematica

Time used: 0.929 (sec). Leaf size: 20

```
DSolve[{x^4*y'[x]==y'[x]*(y'[x]+x^3)},{y[1]==2,y'[1]==1}],y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow x^2 - \log(x^2 + 1) + 1 + \log(2)$$

4.35 problem 38

Internal problem ID [6855]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 38.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_xy]]`

$$y'' - (x^2 - y')^2 = 2x$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$2)=2*x+(x^2-diff(y(x),x))^2,y(x), singsol=all)
```

$$y(x) = \frac{x^3}{3} - \ln(xc_2 - c_1)$$

✓ Solution by Mathematica

Time used: 0.298 (sec). Leaf size: 24

```
DSolve[y''[x]==2*x+(x^2-y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{3} - \log(-x + c_1) + c_2$$

4.36 problem 39

Internal problem ID [6856]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing.
EXERCISES Page 324

Problem number: 39.

ODE order: 2.

ODE degree: 2.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y''^2 - 2y'' + y'^2 - 2y'x = -x^2$$

With initial conditions

$$\left[y(0) = \frac{1}{2}, y'(0) = 1 \right]$$

✓ Solution by Maple

Time used: 0.313 (sec). Leaf size: 23

```
dsolve([diff(y(x),x$2)^2-2*diff(y(x),x$2)+diff(y(x),x)^2-2*x*diff(y(x),x)+x^2=0,y(0) = 1/2,
```

$$y(x) = \frac{(x+1)^2}{2}$$

$$y(x) = \frac{x^2}{2} + \sin(x) + \frac{1}{2}$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{(y'[x])^2-2*y'[x]+(y[x])^2-2*x*y'[x]+x^2==0,{y[0]==1/2,y'[0]==1}},y[x],x,IncludeS
```

Not solved

4.37 problem 40

Internal problem ID [6857]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 40.

ODE order: 2.

ODE degree: 2.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y''^2 - xy'' + y' = 0$$

✓ Solution by Maple

Time used: 0.11 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)^2-x*diff(y(x),x$2)+diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x^3}{12} + c_1$$

$$y(x) = \frac{1}{2}c_1x^2 - xc_1^2 + c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 24

```
DSolve[(y'[x])^2-x*y'[x]+y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1x^2}{2} - c_1^2x + c_2$$

4.38 problem 41

Internal problem ID [6858]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing.
EXERCISES Page 324

Problem number: 41.

ODE order: 2.

ODE degree: 3.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y''^3 - 12y'(xy'' - 2y') = 0$$

✓ Solution by Maple

Time used: 0.344 (sec). Leaf size: 174

```
dsolve(diff(y(x),x$2)^3=12*diff(y(x),x)*(x*diff(y(x),x$2)-2*diff(y(x),x)),y(x), singsol=all)
```

$$y(x) = \frac{x^4}{9} + c_1$$

$$y(x) = c_1$$

$$y(x) = \int \text{RootOf} \left(-6 \ln(x) \right. \\ \left. - \left(\int -z \frac{3-f \sqrt{\frac{1}{-f(9-f-4)}} 2^{\frac{1}{3}} \left(\left(3 \sqrt{\frac{1}{-f(9-f-4)}} -f+1 \right)^2 (9-f-4)^4 \right)^{\frac{1}{3}} - 2 2^{\frac{2}{3}} \left(\left(3 \sqrt{\frac{1}{-f(9-f-4)}} -f+1 \right) \right)}{-f(9-f-4)} \right. \right. \\ \left. \left. + 6c_1 \right) x^3 dx + c_2 \right.$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(y'[x])^3==12*y'[x]*(x*y'[x]-2*y'[x]),y[x],x,IncludeSingularSolutions -> True]
```

Not solved

4.39 problem 42

Internal problem ID [6859]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 42.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$3yy'y'' - y'^3 = -1$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 87

```
dsolve(3*y(x)*diff(y(x),x)*diff(y(x),x$2)=diff(y(x),x)^3-1,y(x), singsol=all)
```

$$\begin{aligned}\frac{3(c_1y(x) + 1)^{\frac{2}{3}}}{2c_1} - x - c_2 &= 0 \\ \frac{3(c_1y(x) + 1)^{\frac{2}{3}}}{c_1(-1 + i\sqrt{3})} - x - c_2 &= 0 \\ -\frac{3(c_1y(x) + 1)^{\frac{2}{3}}}{c_1(1 + i\sqrt{3})} - x - c_2 &= 0\end{aligned}$$

✓ Solution by Mathematica

Time used: 45.036 (sec). Leaf size: 126

```
DSolve[3*y[x]*y'[x]*y''[x]==(y'[x])^3-1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{9}e^{-3c_1} \left(-9 + 2\sqrt{6} \left(e^{3c_1} (x + c_2) \right)^{3/2} \right)$$

$$y(x) \rightarrow \frac{1}{9}e^{-3c_1} \left(-9 + 2\sqrt{6} \left(-\sqrt[3]{-1} e^{3c_1} (x + c_2) \right)^{3/2} \right)$$

$$y(x) \rightarrow \frac{1}{9}e^{-3c_1} \left(-9 + 2\sqrt{6} \left((-1)^{2/3} e^{3c_1} (x + c_2) \right)^{3/2} \right)$$

4.40 problem 43

Internal problem ID [6860]

Book: Elementary differential equations. By Earl D. Rainville, Phillip E. Bedient. Macmillan Publishing Co. NY. 6th edition. 1981.

Section: CHAPTER 16. Nonlinear equations. Section 101. Independent variable missing. EXERCISES Page 324

Problem number: 43.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$4yy'^2y'' - y'^4 = 3$$

✓ Solution by Maple

Time used: 0.14 (sec). Leaf size: 91

```
dsolve(4*y(x)*diff(y(x),x)^2*diff(y(x),x$2)=diff(y(x),x)^4+3,y(x), singsol=all)
```

$$-\frac{4(c_1y(x) - 3)^{\frac{3}{4}}}{3c_1} - x - c_2 = 0$$

$$\frac{4(c_1y(x) - 3)^{\frac{3}{4}}}{3c_1} - x - c_2 = 0$$

$$-\frac{4i(c_1y(x) - 3)^{\frac{3}{4}}}{3c_1} - x - c_2 = 0$$

$$\frac{4i(c_1y(x) - 3)^{\frac{3}{4}}}{3c_1} - x - c_2 = 0$$

✓ Solution by Mathematica

Time used: 60.242 (sec). Leaf size: 156

```
DSolve[4*y[x]*(y'[x])^2*y''[x]==(y'[x])^4+3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3}{8}e^{-4c_1} \left(8 + \sqrt[3]{6}(-e^{4c_1}(x + c_2))^{4/3} \right)$$

$$y(x) \rightarrow \frac{3}{8}e^{-4c_1} \left(8 + \sqrt[3]{6}(-ie^{4c_1}(x + c_2))^{4/3} \right)$$

$$y(x) \rightarrow \frac{3}{8}e^{-4c_1} \left(8 + \sqrt[3]{6}(ie^{4c_1}(x + c_2))^{4/3} \right)$$

$$y(x) \rightarrow \frac{3}{8}e^{-4c_1} \left(8 + \sqrt[3]{6}(e^{4c_1}(x + c_2))^{4/3} \right)$$