


A Solution Manual For

**Elementary differential
equations and boundary value
problems, 10th ed., Boyce and
DiPrima**



Nasser M. Abbasi

March 3, 2024

Contents

1	Section 2.1. Page 40	3
2	Section 2.2. Page 48	35
3	Section 2.4. Page 76	78
4	Section 2.5. Page 88	97
5	Section 2.6. Page 100	111
6	Miscellaneous problems, end of chapter 2. Page 133	146
7	Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144	185
8	Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164	210
9	Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172	241
10	Chapter 3, Second order linear equations, section 3.6, Variation of Parameters. page 190	272
11	Chapter 3, Second order linear equations, 3.7 Mechanical and Electrical Vibrations. page 203	295
12	Chapter 3, Second order linear equations, 3.7 Forced Vibrations. page 217	298
13	Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263	303
14	Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269	326
15	Chapter 7.5, Homogeneous Linear Systems with Constant Coefficients. page 407	345
16	Chapter 7.6, Complex Eigenvalues. page 417	348

17 Chapter 7.8, Repeated Eigenvalues. page 436	368
18 Chapter 7.9, Nonhomogeneous Linear Systems. page 447	386
19 Chapter 9.1, The Phase Plane: Linear Systems. page 505	403
20 Chapter 9.2, Autonomous Systems and Stability. page 517	420

1 Section 2.1. Page 40

1.1	problem 1	4
1.2	problem 2	5
1.3	problem 3	6
1.4	problem 4	7
1.5	problem 5	8
1.6	problem 6	9
1.7	problem 7	10
1.8	problem 8	11
1.9	problem 9	12
1.10	problem 10	13
1.11	problem 11	14
1.12	problem 12	15
1.13	problem 13	16
1.14	problem 14	17
1.15	problem 15	18
1.16	problem 16	19
1.17	problem 17	20
1.18	problem 18	21
1.19	problem 19	22
1.20	problem 20	23
1.21	problem 21	24
1.22	problem 22	25
1.23	problem 23	26
1.24	problem 24	27
1.25	problem 25	28
1.26	problem 26	29
1.27	problem 27	30
1.28	problem 28	31
1.29	problem 29	32
1.30	problem 30	33
1.31	problem 31	34

1.1 problem 1

Internal problem ID [448]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$3y + y' = e^{-2t} + t$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(3*y(t)+diff(y(t),t) = exp(-2*t)+t,y(t), singsol=all)
```

$$y(t) = \frac{t}{3} - \frac{1}{9} + e^{-2t} + c_1 e^{-3t}$$

✓ Solution by Mathematica

Time used: 0.086 (sec). Leaf size: 27

```
DSolve[3*y[t]+y'[t] == Exp[-2*t]+t,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{t}{3} + e^{-2t} + c_1 e^{-3t} - \frac{1}{9}$$

1.2 problem 2

Internal problem ID [449]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$-2y + y' = e^{2t}t^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(-2*y(t)+diff(y(t),t) = exp(2*t)*t^2,y(t), singsol=all)
```

$$y(t) = \left(\frac{t^3}{3} + c_1\right) e^{2t}$$

✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 22

```
DSolve[-2*y[t]+y'[t]== Exp[2*t]*t^2,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{3}e^{2t}(t^3 + 3c_1)$$

1.3 problem 3

Internal problem ID [450]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y + y' = 1 + te^{-t}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(y(t)+diff(y(t),t) = 1+t/exp(t),y(t), singsol=all)
```

$$y(t) = \left(\frac{t^2}{2} + e^t + c_1 \right) e^{-t}$$

✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 27

```
DSolve[y[t]+y'[t] == 1+t/Exp[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{2}e^{-t}(t^2 + 2e^t + 2c_1)$$

1.4 problem 4

Internal problem ID [451]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$\frac{y}{t} + y' = 3 \cos(2t)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(y(t)/t+diff(y(t),t) = 3*cos(2*t),y(t), singsol=all)
```

$$y(t) = \frac{\frac{3 \cos(2t)}{4} + \frac{3 \sin(2t)t}{2} + C_1}{t}$$

✓ Solution by Mathematica

Time used: 0.073 (sec). Leaf size: 30

```
DSolve[y[t]/t+y'[t] == 3*Cos[2*t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{6t \sin(2t) + 3 \cos(2t) + 4c_1}{4t}$$

1.5 problem 5

Internal problem ID [452]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$-2y + y' = 3e^t$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(-2*y(t)+diff(y(t),t) = 3*exp(t),y(t), singsol=all)
```

$$y(t) = -3e^t + c_1e^{2t}$$

✓ Solution by Mathematica

Time used: 0.046 (sec). Leaf size: 17

```
DSolve[-2*y[t]+y'[t] == 3*Exp[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^t(-3 + c_1e^t)$$

1.6 problem 6

Internal problem ID [453]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$2y + ty' = \sin(t)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(2*y(t)+t*diff(y(t),t) = sin(t),y(t), singsol=all)
```

$$y(t) = \frac{\sin(t) - \cos(t)t + c_1}{t^2}$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 19

```
DSolve[2*y[t]+t*y'[t]== Sin[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{\sin(t) - t \cos(t) + c_1}{t^2}$$

1.7 problem 7

Internal problem ID [454]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$2yt + y' = 2t e^{-t^2}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(2*t*y(t)+diff(y(t),t) = 2*t/exp(t^2),y(t), singsol=all)
```

$$y(t) = (t^2 + c_1) e^{-t^2}$$

✓ Solution by Mathematica

Time used: 0.055 (sec). Leaf size: 19

```
DSolve[2*t*y[t]+y'[t] == 2*t/Exp[t^2],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{-t^2} (t^2 + c_1)$$

1.8 problem 8

Internal problem ID [455]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$4yt + (t^2 + 1)y' = \frac{1}{(t^2 + 1)^2}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(4*t*y(t)+(t^2+1)*diff(y(t),t) = 1/(t^2+1)^2,y(t), singsol=all)
```

$$y(t) = \frac{\arctan(t) + c_1}{(t^2 + 1)^2}$$

✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 18

```
DSolve[4*t*y[t]+(t^2+1)*y'[t] == 1/(t^2+1)^2,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{\arctan(t) + c_1}{(t^2 + 1)^2}$$

1.9 problem 9

Internal problem ID [456]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y + 2y' = 3t$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(y(t)+2*diff(y(t),t) = 3*t,y(t), singsol=all)
```

$$y(t) = 3t - 6 + e^{-\frac{t}{2}}c_1$$

✓ Solution by Mathematica

Time used: 0.045 (sec). Leaf size: 20

```
DSolve[y[t]+2*y'[t] == 3*t,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 3t + c_1 e^{-t/2} - 6$$

1.10 problem 10

Internal problem ID [457]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$ty' - y = t^2 e^{-t}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve(-y(t)+t*diff(y(t),t) = t^2/exp(t),y(t), singsol=all)
```

$$y(t) = (-e^{-t} + c_1) t$$

✓ Solution by Mathematica

Time used: 0.053 (sec). Leaf size: 17

```
DSolve[-y[t]+t*y'[t] == t^2/Exp[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow t(-e^{-t} + c_1)$$

1.11 problem 11

Internal problem ID [458]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y + y' = 5 \sin(2t)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(y(t)+diff(y(t),t) = 5*sin(2*t),y(t), singsol=all)
```

$$y(t) = -2 \cos(2t) + \sin(2t) + e^{-t}c_1$$

✓ Solution by Mathematica

Time used: 0.087 (sec). Leaf size: 24

```
DSolve[y[t]+y'[t] == 5*Sin[2*t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \sin(2t) - 2 \cos(2t) + c_1 e^{-t}$$

1.12 problem 12

Internal problem ID [459]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y + 2y' = 3t^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(y(t)+2*diff(y(t),t) = 3*t^2,y(t), singsol=all)
```

$$y(t) = 3t^2 - 12t + 24 + e^{-\frac{t}{2}}c_1$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 25

```
DSolve[y[t]+2*y'[t] == 3*t^2,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 3t^2 - 12t + c_1 e^{-t/2} + 24$$

1.13 problem 13

Internal problem ID [460]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$-y + y' = 2e^{2t}$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve([-y(t)+diff(y(t),t) = 2*exp(2*t)*t,y(0) = 1],y(t), singsol=all)
```

$$y(t) = (2t - 2)e^{2t} + 3e^t$$

✓ Solution by Mathematica

Time used: 0.053 (sec). Leaf size: 19

```
DSolve[{-y[t]+y'[t] == 2*Exp[2*t]*t,y[0]==1},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^t(2e^t(t - 1) + 3)$$

1.14 problem 14

Internal problem ID [461]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 14.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear, 'class A']`

$$2y + y' = te^{-2t}$$

With initial conditions

$$[y(1) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([2*y(t)+diff(y(t),t) = t/exp(2*t),y(1) = 0],y(t), singsol=all)
```

$$y(t) = \frac{(t^2 - 1)e^{-2t}}{2}$$

✓ Solution by Mathematica

Time used: 0.054 (sec). Leaf size: 19

```
DSolve[{2*y[t]+y'[t] == t/Exp[2*t],y[1]==0},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{2}e^{-2t}(t^2 - 1)$$

1.15 problem 15

Internal problem ID [462]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$2y + ty' = t^2 - t + 1$$

With initial conditions

$$\left[y(1) = \frac{1}{2} \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 19

```
dsolve([2*y(t)+t*diff(y(t),t) = t^2-t+1,y(1) = 1/2],y(t), singsol=all)
```

$$y(t) = \frac{t^2}{4} - \frac{t}{3} + \frac{1}{2} + \frac{1}{12t^2}$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 22

```
DSolve[{2*y[t]+t*y'[t] == t^2-t+1,y[1]==1/2},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{12} \left(3t^2 + \frac{1}{t^2} - 4t + 6 \right)$$

1.16 problem 16

Internal problem ID [463]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + \frac{2y}{t} = \frac{\cos(t)}{t^2}$$

With initial conditions

$$[y(\pi) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 10

```
dsolve([2*y(t)/t+diff(y(t),t) = cos(t)/t^2,y(Pi) = 0],y(t), singsol=all)
```

$$y(t) = \frac{\sin(t)}{t^2}$$

✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 11

```
DSolve[{2*y[t]/t+y'[t] == Cos[t]/t^2,y[Pi]==0},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{\sin(t)}{t^2}$$

1.17 problem 17

Internal problem ID [464]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 17.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$-2y + y' = e^{2t}$$

With initial conditions

$$[y(0) = 2]$$

✓ Solution by Maple

Time used: 0.328 (sec). Leaf size: 12

```
dsolve([-2*y(t)+diff(y(t),t) = exp(2*t),y(0) = 2],y(t), singsol=all)
```

$$y(t) = (2 + t)e^{2t}$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 14

```
DSolve[{-2*y[t]+y'[t] == Exp[2*t],y[0]==2},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{2t}(t + 2)$$

1.18 problem 18

Internal problem ID [465]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 18.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$2y + ty' = \sin(t)$$

With initial conditions

$$\left[y\left(\frac{\pi}{2}\right) = 1 \right]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 22

```
dsolve([2*y(t)+t*diff(y(t),t) = sin(t),y(1/2*Pi) = 1],y(t), singsol=all)
```

$$y(t) = \frac{\sin(t) - \cos(t)t + \frac{\pi^2}{4} - 1}{t^2}$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 26

```
DSolve[{2*y[t]+t*y'[t] == Sin[t],y[Pi/2]==1},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{4 \sin(t) - 4t \cos(t) + \pi^2 - 4}{4t^2}$$

1.19 problem 19

Internal problem ID [466]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$4yt^2 + y/t^3 = e^{-t}$$

With initial conditions

$$[y(-1) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve([4*t^2*y(t)+t^3*diff(y(t),t) = exp(-t),y(-1) = 0],y(t), singsol=all)
```

$$y(t) = -\frac{(t+1)e^{-t}}{t^4}$$

✓ Solution by Mathematica

Time used: 0.067 (sec). Leaf size: 18

```
DSolve[{4*t^2*y[t]+t^3*y'[t] == Exp[-t],y[-1]==0},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -\frac{e^{-t}(t+1)}{t^4}$$

1.20 problem 20

Internal problem ID [467]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 20.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$(t + 1)y + ty' = t$$

With initial conditions

$$[y(\ln(2)) = 1]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve([(1+t)*y(t)+t*diff(y(t),t) = t,y(ln(2)) = 1],y(t), singsol=all)
```

$$y(t) = \frac{t - 1 + 2e^{-t}}{t}$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 23

```
DSolve[{(1+t)*y[t]+t*y'[t]== t,y[Log[2]]==1},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{e^{-t}(e^t(t - 1) + 2)}{t}$$

1.21 problem 21

Internal problem ID [468]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 21.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$-\frac{y}{2} + y' = 2 \cos(t)$$

With initial conditions

$$[y(0) = a]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve([-1/2*y(t)+diff(y(t),t) = 2*cos(t),y(0) = a],y(t), singsol=all)
```

$$y(t) = -\frac{4 \cos(t)}{5} + \frac{8 \sin(t)}{5} + e^{\frac{t}{2}} a + \frac{4 e^{\frac{t}{2}}}{5}$$

✓ Solution by Mathematica

Time used: 0.054 (sec). Leaf size: 31

```
DSolve[{-1/2*y[t]+y'[t] == 2*Cos[t],y[0]==a},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{5}((5a + 4)e^{t/2} + 8 \sin(t) - 4 \cos(t))$$

1.22 problem 22

Internal problem ID [469]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 22.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$-y + 2y' = e^{\frac{t}{3}}$$

With initial conditions

$$[y(0) = a]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve([-y(t)+2*diff(y(t),t) = exp(1/3*t),y(0) = a],y(t), singsol=all)
```

$$y(t) = \left(-3 + (a + 3)e^{\frac{t}{6}}\right)e^{\frac{t}{3}}$$

✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 26

```
DSolve[{-y[t]+2*y'[t] == Exp[1/3*t],y[0]==a},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{t/3}((a + 3)e^{t/6} - 3)$$

1.23 problem 23

Internal problem ID [470]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 23.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$-2y + 3y' = e^{-\frac{\pi t}{2}}$$

With initial conditions

$$[y(0) = a]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 35

```
dsolve([-2*y(t)+3*diff(y(t),t) = exp(-1/2*Pi*t),y(0) = a],y(t), singsol=all)
```

$$y(t) = \frac{\left(3\pi a - 2e^{t\left(-\frac{\pi}{2} - \frac{2}{3}\right)} + 4a + 2\right)e^{\frac{2t}{3}}}{3\pi + 4}$$

✓ Solution by Mathematica

Time used: 0.083 (sec). Leaf size: 43

```
DSolve[{-2*y[t]+3*y'[t] == Exp[-1/2*Pi*t],y[0]==a},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{e^{2t/3} \left((4 + 3\pi)a - 2e^{-\frac{1}{6}(4+3\pi)t} + 2 \right)}{4 + 3\pi}$$

1.24 problem 24

Internal problem ID [471]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 24.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$(t + 1)y + ty' = 2te^{-t}$$

With initial conditions

$$[y(1) = a]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve([(1+t)*y(t)+t*diff(y(t),t) = 2*t/exp(t),y(1) = a],y(t), singsol=all)
```

$$y(t) = \frac{(t^2 + ae - 1)e^{-t}}{t}$$

✓ Solution by Mathematica

Time used: 0.065 (sec). Leaf size: 22

```
DSolve[{(1+t)*y[t]+t*y'[t] == 2*t/Exp[t],y[1]==a},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{e^{-t}(ea + t^2 - 1)}{t}$$

1.25 problem 25

Internal problem ID [472]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 25.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$2y + ty' = \frac{\sin(t)}{t}$$

With initial conditions

$$\left[y\left(-\frac{\pi}{2}\right) = a \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 19

```
dsolve([2*y(t)+t*diff(y(t),t) = sin(t)/t,y(-1/2*Pi) = a],y(t), singsol=all)
```

$$y(t) = \frac{-\cos(t) + \frac{a\pi^2}{4}}{t^2}$$

✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 22

```
DSolve[{2*y[t]+t*y'[t] == Sin[t]/t,y[-Pi/2]==a},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{\pi^2 a - 4 \cos(t)}{4t^2}$$

1.26 problem 26

Internal problem ID [473]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 26.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$\cos(t)y + \sin(t)y' = e^t$$

With initial conditions

$$[y(1) = a]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 18

```
dsolve([cos(t)*y(t)+sin(t)*diff(y(t),t) = exp(t),y(1) = a],y(t), singsol=all)
```

$$y(t) = \csc(t) (e^t + a \sin(1) - e)$$

✓ Solution by Mathematica

Time used: 0.062 (sec). Leaf size: 19

```
DSolve[{Cos[t]*y[t]+Sin[t]*y'[t] == Exp[t],y[1]==a},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \csc(t) (a \sin(1) + e^t - e)$$

1.27 problem 27

Internal problem ID [474]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 27.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + \frac{y}{2} = 2 \cos(t)$$

With initial conditions

$$[y(0) = -1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve([1/2*y(t)+diff(y(t),t) = 2*cos(t),y(0) = -1],y(t), singsol=all)
```

$$y(t) = \frac{4 \cos(t)}{5} + \frac{8 \sin(t)}{5} - \frac{9 e^{-\frac{t}{2}}}{5}$$

✓ Solution by Mathematica

Time used: 0.049 (sec). Leaf size: 27

```
DSolve[{1/2*y[t]+y'[t] == 2*Cos[t],y[0]==-1},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{5}(-9e^{-t/2} + 8 \sin(t) + 4 \cos(t))$$

1.28 problem 28

Internal problem ID [475]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 28.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$\frac{2y}{3} + y' = -\frac{t}{2} + 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(2/3*y(t)+diff(y(t),t) = 1-1/2*t,y(t), singsol=all)
```

$$y(t) = -\frac{3t}{4} + \frac{21}{8} + e^{-\frac{2t}{3}} c_1$$

✓ Solution by Mathematica

Time used: 0.061 (sec). Leaf size: 24

```
DSolve[2/3*y[t]+y'[t] == 1-1/2*t,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -\frac{3t}{4} + c_1 e^{-2t/3} + \frac{21}{8}$$

1.29 problem 29

Internal problem ID [476]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 29.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$\frac{y}{4} + y' = 3 + 2 \cos(2t)$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
dsolve([1/4*y(t)+diff(y(t),t) = 3+2*cos(2*t),y(0) = 0],y(t), singsol=all)
```

$$y(t) = 12 + \frac{8 \cos(2t)}{65} + \frac{64 \sin(2t)}{65} - \frac{788 e^{-t/4}}{65}$$

✓ Solution by Mathematica

Time used: 0.158 (sec). Leaf size: 32

```
DSolve[{1/4*y[t]+y'[t] == 3+2*Cos[2*t],y[0]==0},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{4}{65}(-197e^{-t/4} + 16 \sin(2t) + 2 \cos(2t) + 195)$$

1.30 problem 30

Internal problem ID [477]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 30.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$-y + y' = 1 + 3 \sin(t)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(-y(t)+diff(y(t),t) = 1+3*sin(t),y(t), singsol=all)
```

$$y(t) = -1 - \frac{3 \cos(t)}{2} - \frac{3 \sin(t)}{2} + c_1 e^t$$

✓ Solution by Mathematica

Time used: 0.072 (sec). Leaf size: 25

```
DSolve[-y[t]+y'[t] == 1+3*Sin[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -\frac{3 \sin(t)}{2} - \frac{3 \cos(t)}{2} + c_1 e^t - 1$$

1.31 problem 31

Internal problem ID [478]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.1. Page 40

Problem number: 31.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear, 'class A']`

$$-\frac{3y}{2} + y' = 2e^t + 3t$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(-3/2*y(t)+diff(y(t),t) = 2*exp(t)+3*t,y(t), singsol=all)
```

$$y(t) = -2t - \frac{4}{3} - 4e^t + e^{\frac{3t}{2}}c_1$$

✓ Solution by Mathematica

Time used: 0.1 (sec). Leaf size: 27

```
DSolve[-3/2*y[t]+y'[t] == 2*Exp[t]+3*t,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -2t - 4e^t + c_1e^{3t/2} - \frac{4}{3}$$

2 Section 2.2. Page 48

2.1	problem 1	36
2.2	problem 2	37
2.3	problem 3	38
2.4	problem 4	39
2.5	problem 5	40
2.6	problem 6	41
2.7	problem 7	42
2.8	problem 8	43
2.9	problem 9	46
2.10	problem 10	47
2.11	problem 11	48
2.12	problem 12	49
2.13	problem 13	50
2.14	problem 14	51
2.15	problem 15	52
2.16	problem 16	53
2.17	problem 17	54
2.18	problem 18	55
2.19	problem 19	56
2.20	problem 20	57
2.21	problem 21	58
2.22	problem 22	59
2.23	problem 23	60
2.24	problem 24	61
2.25	problem 25	62
2.26	problem 26	63
2.27	problem 27	64
2.28	problem 28	65
2.29	problem 29	66
2.30	problem 31	67
2.31	problem 32	68
2.32	problem 33	69
2.33	problem 34	71
2.34	problem 35	74
2.35	problem 36	75
2.36	problem 37	76
2.37	problem 38	77

2.1 problem 1

Internal problem ID [479]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{x^2}{y} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(x),x) = x^2/y(x),y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{6x^3 + 9c_1}}{3}$$

$$y(x) = \frac{\sqrt{6x^3 + 9c_1}}{3}$$

✓ Solution by Mathematica

Time used: 0.093 (sec). Leaf size: 50

```
DSolve[y'[x] == x^2/y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{\frac{2}{3}}\sqrt{x^3 + 3c_1}$$

$$y(x) \rightarrow \sqrt{\frac{2}{3}}\sqrt{x^3 + 3c_1}$$

2.2 problem 2

Internal problem ID [480]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{x^2}{(x^3 + 1)y} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
dsolve(diff(y(x),x) = x^2/(x^3+1)/y(x),y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{6 \ln(x^3 + 1) + 9c_1}}{3}$$

$$y(x) = \frac{\sqrt{6 \ln(x^3 + 1) + 9c_1}}{3}$$

✓ Solution by Mathematica

Time used: 0.12 (sec). Leaf size: 56

```
DSolve[y'[x] == x^2/(x^3+1)/y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{\frac{2}{3}} \sqrt{\log(x^3 + 1) + 3c_1}$$

$$y(x) \rightarrow \sqrt{\frac{2}{3}} \sqrt{\log(x^3 + 1) + 3c_1}$$

2.3 problem 3

Internal problem ID [481]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$\sin(x)y^2 + y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve(sin(x)*y(x)^2+diff(y(x),x) = 0,y(x), singsol=all)
```

$$y(x) = -\frac{1}{\cos(x) - c_1}$$

✓ Solution by Mathematica

Time used: 0.133 (sec). Leaf size: 19

```
DSolve[Sin[x]*y[x]^2+y'[x]== 0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\cos(x) + c_1}$$

$$y(x) \rightarrow 0$$

2.4 problem 4

Internal problem ID [482]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{3x^2 - 1}{3 + 2y} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 45

```
dsolve(diff(y(x),x) = (3*x^2-1)/(3+2*y(x)),y(x), singsol=all)
```

$$y(x) = -\frac{3}{2} - \frac{\sqrt{4x^3 + 4c_1 - 4x + 9}}{2}$$

$$y(x) = -\frac{3}{2} + \frac{\sqrt{4x^3 + 4c_1 - 4x + 9}}{2}$$

✓ Solution by Mathematica

Time used: 0.121 (sec). Leaf size: 59

```
DSolve[y'[x] == (3*x^2-1)/(3+2*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(-3 - \sqrt{4x^3 - 4x + 9 + 4c_1} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left(-3 + \sqrt{4x^3 - 4x + 9 + 4c_1} \right)$$

2.5 problem 5

Internal problem ID [483]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \cos(x)^2 \cos(2y)^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x) = cos(x)^2*cos(2*y(x))^2,y(x), singsol=all)
```

$$y(x) = \frac{\arctan\left(x + 2c_1 + \frac{\sin(2x)}{2}\right)}{2}$$

✓ Solution by Mathematica

Time used: 1.312 (sec). Leaf size: 63

```
DSolve[y'[x] == Cos[x]^2*Cos[2*y[x]]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \arctan\left(x + \sin(x) \cos(x) + \frac{c_1}{4}\right)$$

$$y(x) \rightarrow \frac{1}{2} \arctan\left(x + \sin(x) \cos(x) + \frac{c_1}{4}\right)$$

$$y(x) \rightarrow -\frac{\pi}{4}$$

$$y(x) \rightarrow \frac{\pi}{4}$$

2.6 problem 6

Internal problem ID [484]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y'x - \sqrt{1 - y^2} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(x*diff(y(x),x) = (1-y(x)^2)^(1/2),y(x), singsol=all)
```

$$y(x) = \sin(\ln(x) + c_1)$$

✓ Solution by Mathematica

Time used: 0.206 (sec). Leaf size: 29

```
DSolve[x*y'[x] == (1-y[x]^2)^(1/2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \cos(\log(x) + c_1)$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 1$$

$$y(x) \rightarrow \text{Interval}[\{-1, 1\}]$$

2.7 problem 7

Internal problem ID [485]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type ['y=_G(x,y)']

$$y' - \frac{-e^{-x} + x}{e^y + x} = 0$$

X Solution by Maple

```
dsolve(diff(y(x),x) = (-exp(-x)+x)/(exp(y(x))+x),y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y'[x] == (-Exp[-x]+x)/(Exp[y[x]]+x),y[x],x,IncludeSingularSolutions -> True]
```

Not solved

2.8 problem 8

Internal problem ID [486]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$y' - \frac{x^2}{1 + y^2} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 353

```
dsolve(diff(y(x),x) = x^2/(1+y(x)^2),y(x), singsol=all)
```

$$\begin{aligned}
 y(x) &= \frac{\left(4x^3 + 12c_1 + 4\sqrt{x^6 + 6c_1x^3 + 9c_1^2 + 4}\right)^{\frac{1}{3}}}{2} \\
 &\quad - \frac{\left(4x^3 + 12c_1 + 4\sqrt{x^6 + 6c_1x^3 + 9c_1^2 + 4}\right)^{\frac{1}{3}}}{2} \\
 y(x) &= -\frac{\left(4x^3 + 12c_1 + 4\sqrt{x^6 + 6c_1x^3 + 9c_1^2 + 4}\right)^{\frac{1}{3}}}{4} \\
 &\quad + \frac{\left(4x^3 + 12c_1 + 4\sqrt{x^6 + 6c_1x^3 + 9c_1^2 + 4}\right)^{\frac{1}{3}}}{1} \\
 &\quad - \frac{i\sqrt{3} \left(\frac{\left(4x^3 + 12c_1 + 4\sqrt{x^6 + 6c_1x^3 + 9c_1^2 + 4}\right)^{\frac{1}{3}}}{2} + \frac{2}{\left(4x^3 + 12c_1 + 4\sqrt{x^6 + 6c_1x^3 + 9c_1^2 + 4}\right)^{\frac{1}{3}}} \right)}{2} \\
 y(x) &= -\frac{\left(4x^3 + 12c_1 + 4\sqrt{x^6 + 6c_1x^3 + 9c_1^2 + 4}\right)^{\frac{1}{3}}}{4} \\
 &\quad + \frac{\left(4x^3 + 12c_1 + 4\sqrt{x^6 + 6c_1x^3 + 9c_1^2 + 4}\right)^{\frac{1}{3}}}{1} \\
 &\quad + \frac{i\sqrt{3} \left(\frac{\left(4x^3 + 12c_1 + 4\sqrt{x^6 + 6c_1x^3 + 9c_1^2 + 4}\right)^{\frac{1}{3}}}{2} + \frac{2}{\left(4x^3 + 12c_1 + 4\sqrt{x^6 + 6c_1x^3 + 9c_1^2 + 4}\right)^{\frac{1}{3}}} \right)}{2}
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 2.246 (sec). Leaf size: 307

`DSolve[y'[x]== x^2/(1+y[x]^2),y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{-2 + \sqrt[3]{2}(x^3 + \sqrt{x^6 + 6c_1x^3 + 4 + 9c_1^2 + 3c_1})^{2/3}}{2^{2/3}\sqrt[3]{x^3 + \sqrt{x^6 + 6c_1x^3 + 4 + 9c_1^2 + 3c_1}}}$$

$$y(x) \rightarrow \frac{i(\sqrt{3} + i)\sqrt[3]{x^3 + \sqrt{x^6 + 6c_1x^3 + 4 + 9c_1^2 + 3c_1}}}{2\sqrt[3]{2}} + \frac{1 + i\sqrt{3}}{2^{2/3}\sqrt[3]{x^3 + \sqrt{x^6 + 6c_1x^3 + 4 + 9c_1^2 + 3c_1}}}$$

$$y(x) \rightarrow \frac{1 - i\sqrt{3}}{2^{2/3}\sqrt[3]{x^3 + \sqrt{x^6 + 6c_1x^3 + 4 + 9c_1^2 + 3c_1}}} - \frac{(1 + i\sqrt{3})\sqrt[3]{x^3 + \sqrt{x^6 + 6c_1x^3 + 4 + 9c_1^2 + 3c_1}}}{2\sqrt[3]{2}}$$

2.9 problem 9

Internal problem ID [487]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - (1 - 2x)y^2 = 0$$

With initial conditions

$$\left[y(0) = -\frac{1}{6} \right]$$

✓ Solution by Maple

Time used: 0.109 (sec). Leaf size: 14

```
dsolve([diff(y(x),x) = (1-2*x)*y(x)^2,y(0) = -1/6],y(x), singsol=all)
```

$$y(x) = \frac{1}{x^2 - x - 6}$$

✓ Solution by Mathematica

Time used: 0.131 (sec). Leaf size: 15

```
DSolve[{y'[x] == (1-2*x)*y[x]^2,y[0]==-1/6},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{x^2 - x - 6}$$

2.10 problem 10

Internal problem ID [488]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{1 - 2x}{y} = 0$$

With initial conditions

$$[y(1) = -2]$$

✓ Solution by Maple

Time used: 0.359 (sec). Leaf size: 18

```
dsolve([diff(y(x),x) = (1-2*x)/y(x),y(1) = -2],y(x), singsol=all)
```

$$y(x) = -\sqrt{-2x^2 + 2x + 4}$$

✓ Solution by Mathematica

Time used: 0.094 (sec). Leaf size: 24

```
DSolve[{y'[x] == (1-2*x)/y[x],y[1]==-2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{2}\sqrt{-x^2 + x + 2}$$

2.11 problem 11

Internal problem ID [489]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y'e^{-x}y = -x$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 17

```
dsolve([x+y(x)*diff(y(x),x)/exp(x) = 0,y(0) = 1],y(x), singsol=all)
```

$$y(x) = \sqrt{-1 - 2x e^x + 2 e^x}$$

✓ Solution by Mathematica

Time used: 1.763 (sec). Leaf size: 19

```
DSolve[{x+y[x]*y'[x]/Exp[x] == 0,y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt{-2e^x(x-1) - 1}$$

2.12 problem 12

Internal problem ID [490]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$r' - \frac{r^2}{x} = 0$$

With initial conditions

$$[r(1) = 2]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 14

```
dsolve([diff(r(x),x) = r(x)^2/x,r(1) = 2],r(x), singsol=all)
```

$$r(x) = -\frac{2}{2 \ln(x) - 1}$$

✓ Solution by Mathematica

Time used: 0.128 (sec). Leaf size: 15

```
DSolve[{r'[x] == r[x]^2/x,r[1]==2},r[x],x,IncludeSingularSolutions -> True]
```

$$r(x) \rightarrow \frac{2}{1 - 2 \log(x)}$$

2.13 problem 13

Internal problem ID [491]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{2x}{y + x^2y} = 0$$

With initial conditions

$$[y(0) = -2]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 18

```
dsolve([diff(y(x),x) = 2*x/(y(x)+x^2*y(x)),y(0) = -2],y(x), singsol=all)
```

$$y(x) = -\sqrt{2 \ln(x^2 + 1) + 4}$$

✓ Solution by Mathematica

Time used: 0.091 (sec). Leaf size: 24

```
DSolve[{y'[x] == 2*x/(y[x]+x^2*y[x]),y[0]==-2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{2} \sqrt{\log(x^2 + 1) + 2}$$

2.14 problem 14

Internal problem ID [492]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 14.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{xy^2}{\sqrt{x^2 + 1}} = 0$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 17

```
dsolve([diff(y(x),x) = x*y(x)^2/(x^2+1)^(1/2),y(0) = 1],y(x), singsol=all)
```

$$y(x) = -\frac{1}{\sqrt{x^2 + 1} - 2}$$

✓ Solution by Mathematica

Time used: 0.183 (sec). Leaf size: 20

```
DSolve[{y'[x] == x*y[x]^2/(x^2+1)^(1/2),y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2 - \sqrt{x^2 + 1}}$$

2.15 problem 15

Internal problem ID [493]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{2x}{1+2y} = 0$$

With initial conditions

$$[y(2) = 0]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 17

```
dsolve([diff(y(x),x) = 2*x/(1+2*y(x)),y(2) = 0],y(x), singsol=all)
```

$$y(x) = -\frac{1}{2} + \frac{\sqrt{4x^2 - 15}}{2}$$

✓ Solution by Mathematica

Time used: 0.115 (sec). Leaf size: 22

```
DSolve[{y'[x] == 2*x/(1+2*y[x]),y[2]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(\sqrt{4x^2 - 15} - 1 \right)$$

2.16 problem 16

Internal problem ID [494]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{x(x^2 + 1)}{4y^3} = 0$$

With initial conditions

$$\left[y(0) = -\frac{\sqrt{2}}{2} \right]$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 15

```
dsolve([diff(y(x),x) = 1/4*x*(x^2+1)/y(x)^3,y(0) = -1/2*2^(1/2)],y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{2x^2 + 2}}{2}$$

✓ Solution by Mathematica

Time used: 0.23 (sec). Leaf size: 23

```
DSolve[{y'[x] == 1/4*x*(x^2+1)/y[x]^3,y[0]==-(1/Sqrt[2])},y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow -\frac{\sqrt[4]{(x^2 + 1)^2}}{\sqrt{2}}$$

2.17 problem 17

Internal problem ID [495]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 17.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{-e^x + 3x^2}{-5 + 2y} = 0$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.266 (sec). Leaf size: 21

```
dsolve([diff(y(x),x) = (-exp(x)+3*x^2)/(-5+2*y(x)),y(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{5}{2} - \frac{\sqrt{13 + 4x^3 - 4e^x}}{2}$$

✓ Solution by Mathematica

Time used: 0.891 (sec). Leaf size: 29

```
DSolve[{y'[x] == (-Exp[x]+3*x^2)/(-5+2*y[x]),y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(5 - \sqrt{4x^3 - 4e^x + 13} \right)$$

2.18 problem 18

Internal problem ID [496]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 18.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{-e^x + e^{-x}}{3 + 4y} = 0$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.391 (sec). Leaf size: 29

```
dsolve([diff(y(x),x) = (exp(-x)-exp(x))/(3+4*y(x)),y(0) = 1],y(x), singsol=all)
```

$$y(x) = -\frac{3}{4} + \frac{\sqrt{e^x(-8e^{2x} + 65e^x - 8)}e^{-x}}{4}$$

✓ Solution by Mathematica

Time used: 1.347 (sec). Leaf size: 29

```
DSolve[{y'[x] == (Exp[-x]-Exp[x])/(3+4*y[x]),y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4} \left(\sqrt{-8e^{-x} - 8e^x + 65} - 3 \right)$$

2.19 problem 19

Internal problem ID [497]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$\cos(3y)y' = -\sin(2x)$$

With initial conditions

$$\left[y\left(\frac{\pi}{2}\right) = 0 \right]$$

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 15

```
dsolve([sin(2*x)+cos(3*y(x))*diff(y(x),x) = 0,y(1/2*Pi) = 0],y(x), singsol=all)
```

$$y(x) = \frac{\arcsin\left(\frac{3}{2} + \frac{3\cos(2x)}{2}\right)}{3}$$

✓ Solution by Mathematica

Time used: 0.614 (sec). Leaf size: 16

```
DSolve[{Sin[2*x]+Cos[3*y[x]]*y'[x] == 0,y[Pi/2]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} \arcsin(3 \cos^2(x))$$

2.20 problem 20

Internal problem ID [498]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 20.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$\sqrt{-x^2 + 1} y^2 y' = \arcsin(x)$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.157 (sec). Leaf size: 16

```
dsolve([(-x^2+1)^(1/2)*y(x)^2*diff(y(x),x) = arcsin(x),y(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{(8 + 12 \arcsin(x)^2)^{\frac{1}{3}}}{2}$$

✓ Solution by Mathematica

Time used: 0.527 (sec). Leaf size: 19

```
DSolve[{-x^2+1)^(1/2)*y[x]^2*y'[x] == ArcSin[x],y[0]==1},y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow \sqrt[3]{\frac{3 \arcsin(x)^2}{2} + 1}$$

2.21 problem 21

Internal problem ID [499]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 21.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{3x^2 + 1}{-6y + 3y^2} = 0$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.172 (sec). Leaf size: 109

```
dsolve([diff(y(x),x) = (3*x^2+1)/(-6*y(x)+3*y(x)^2),y(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{(1 + i\sqrt{3}) (4x^3 + 4x + 4\sqrt{x^6 + 2x^4 + x^2 - 4})^{\frac{2}{3}} - 4i\sqrt{3} - 4(4x^3 + 4x + 4\sqrt{x^6 + 2x^4 + x^2 - 4})^{\frac{1}{3}} + 4}{4 (4x^3 + 4x + 4\sqrt{x^6 + 2x^4 + x^2 - 4})^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 4.019 (sec). Leaf size: 158

```
DSolve[{y'[x] == (3*x^2+1)/(-6*y[x]+3*y[x]^2),y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-i2^{2/3}\sqrt{3}(x^3 + \sqrt{x^6 + 2x^4 + x^2 - 4} + x)^{2/3} - 2^{2/3}(x^3 + \sqrt{x^6 + 2x^4 + x^2 - 4} + x)^{2/3} + 4\sqrt[3]{x^3 + \sqrt{x^6 + 2x^4 + x^2 - 4}}}{4\sqrt[3]{x^3 + \sqrt{x^6 + 2x^4 + x^2 - 4} + x}}$$

2.22 problem 22

Internal problem ID [500]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 22.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{3x^2}{-4 + 3y^2} = 0$$

With initial conditions

$$[y(1) = 0]$$

✓ Solution by Maple

Time used: 0.469 (sec). Leaf size: 73

```
dsolve([diff(y(x),x) = 3*x^2/(-4+3*y(x)^2),y(1) = 0],y(x), singsol=all)
```

$$y(x) = -\frac{(1 + i\sqrt{3}) (-108 + 108x^3 + 12\sqrt{81x^6 - 162x^3 - 687})^{\frac{2}{3}} - 48i\sqrt{3} + 48}{12 (-108 + 108x^3 + 12\sqrt{81x^6 - 162x^3 - 687})^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 9.526 (sec). Leaf size: 137

```
DSolve[{y'[x]== 3*x^2/(-4+3*y[x]^2),y[1]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-i\sqrt{3}2^{2/3}(9x^3 + \sqrt{81x^6 - 162x^3 - 687} - 9)^{2/3} - \sqrt[3]{2}\sqrt[3]{3}(9x^3 + \sqrt{81x^6 - 162x^3 - 687} - 9)^{2/3} - 8\sqrt{3}}{2 \cdot 2^{2/3}3^{5/6} \sqrt[3]{9x^3 + \sqrt{81x^6 - 162x^3 - 687} - 9}}$$

2.23 problem 23

Internal problem ID [501]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 23.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - 2y^2 - xy^2 = 0$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.11 (sec). Leaf size: 16

```
dsolve([diff(y(x),x) = 2*y(x)^2+x*y(x)^2,y(0) = 1],y(x), singsol=all)
```

$$y(x) = -\frac{2}{x^2 + 4x - 2}$$

✓ Solution by Mathematica

Time used: 0.127 (sec). Leaf size: 17

```
DSolve[{y'[x] == 2*y[x]^2+x*y[x]^2,y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2}{x^2 + 4x - 2}$$

2.24 problem 24

Internal problem ID [502]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 24.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{2 - e^x}{3 + 2y} = 0$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.25 (sec). Leaf size: 19

```
dsolve([diff(y(x),x) = (2-exp(x))/(3+2*y(x)),y(0) = 0],y(x), singsol=all)
```

$$y(x) = -\frac{3}{2} + \frac{\sqrt{13 - 4e^x + 8x}}{2}$$

✓ Solution by Mathematica

Time used: 0.737 (sec). Leaf size: 25

```
DSolve[{y'[x] == (2-Exp[x])/(3+2*y[x]),y[0]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(\sqrt{8x - 4e^x + 13} - 3)$$

2.25 problem 25

Internal problem ID [503]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 25.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{2 \cos(2x)}{3 + 2y} = 0$$

With initial conditions

$$[y(0) = -1]$$

✓ Solution by Maple

Time used: 0.39 (sec). Leaf size: 18

```
dsolve([diff(y(x),x) = 2*cos(2*x)/(3+2*y(x)),y(0) = -1],y(x), singsol=all)
```

$$y(x) = -\frac{3}{2} + \frac{\sqrt{1 + 4 \sin(2x)}}{2}$$

✓ Solution by Mathematica

Time used: 0.175 (sec). Leaf size: 23

```
DSolve[{y'[x] == 2*Cos[2*x]/(3+2*y[x]),y[0]==-1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(\sqrt{4 \sin(2x) + 1} - 3 \right)$$

2.26 problem 26

Internal problem ID [504]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 26.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - 2(x + 1)(1 + y^2) = 0$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 12

```
dsolve([diff(y(x),x) = 2*(1+x)*(1+y(x)^2),y(0) = 0],y(x), singsol=all)
```

$$y(x) = \tan(x^2 + 2x)$$

✓ Solution by Mathematica

Time used: 0.199 (sec). Leaf size: 11

```
DSolve[{y'[x] == 2*(1+x)*(1+y[x]^2),y[0]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan(x(x + 2))$$

2.27 problem 27

Internal problem ID [505]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 27.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{t(4-y)y}{3} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(t),t) = 1/3*t*(4-y(t))*y(t),y(t), singsol=all)
```

$$y(t) = \frac{4}{1 + 4e^{-\frac{2t^2}{3}}c_1}$$

✓ Solution by Mathematica

Time used: 0.248 (sec). Leaf size: 44

```
DSolve[y'[t]== 1/3*t*(4-y[t])*y[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{4e^{\frac{2t^2}{3}}}{e^{\frac{2t^2}{3}} + e^{4c_1}}$$

$$y(t) \rightarrow 0$$

$$y(t) \rightarrow 4$$

2.28 problem 28

Internal problem ID [506]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 28.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{ty(4-y)}{t+1} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 55

```
dsolve(diff(y(t),t) = t*y(t)*(4-y(t))/(1+t),y(t), singsol=all)
```

$$y(t) = \frac{4}{1 + 4e^{-4t}c_1t^4 + 16e^{-4t}c_1t^3 + 24e^{-4t}c_1t^2 + 16e^{-4t}c_1t + 4e^{-4t}c_1}$$

✓ Solution by Mathematica

Time used: 1.337 (sec). Leaf size: 42

```
DSolve[y'[t] == t*y[t]*(4-y[t])/(1+t),y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{4e^{4t}}{e^{4t} + e^{4c_1}(t+1)^4}$$

$$y(t) \rightarrow 0$$

$$y(t) \rightarrow 4$$

2.29 problem 29

Internal problem ID [507]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 29.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - \frac{b + ay}{d + cy} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 203

```
dsolve(diff(y(x),x) = (b+a*y(x))/(d+c*y(x)),y(x), singsol=all)
```

$$y(x) = \frac{c_1 a^2 + x a^2 - \left(-\text{LambertW} \left(-\frac{c e^{\frac{c_1 a^2}{ad-bc} + \frac{x a^2}{ad-bc} + \frac{bc}{ad-bc}}}{-ad+bc} \right) + \frac{c_1 a^2 + x a^2 + bc}{ad-bc} \right) ad + \left(-\text{LambertW} \left(-\frac{c e^{\frac{c_1 a^2}{ad-bc} + \frac{x a^2}{ad-bc} + \frac{bc}{ad-bc}}}{-ad+bc} \right) + \frac{c_1 a^2 + x a^2 + bc}{ad-bc} \right) ad}{ac}$$

✓ Solution by Mathematica

Time used: 16.166 (sec). Leaf size: 83

```
DSolve[y'[x] == (b+a*y[x])/(d+c*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-bc + (ad - bc)W \left(-\frac{c \left(e^{-1 - \frac{a^2(x+c_1)}{bc}} \right)^{\frac{bc}{bc-ad}}}{bc-ad} \right)}{ac}$$

$$y(x) \rightarrow -\frac{b}{a}$$

2.30 problem 31

Internal problem ID [508]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 31.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _Riccati]`

$$y' - \frac{x^2 + yx + y^2}{x^2} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 11

```
dsolve(diff(y(x),x) = (x^2+x*y(x)+y(x)^2)/x^2,y(x), singsol=all)
```

$$y(x) = \tan(\ln(x) + c_1)x$$

✓ Solution by Mathematica

Time used: 0.199 (sec). Leaf size: 13

```
DSolve[y'[x] == (x^2+x*y[x]+y[x]^2)/x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \tan(\log(x) + c_1)$$

2.31 problem 32

Internal problem ID [509]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 32.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$y' - \frac{x^2 + 3y^2}{2xy} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x) = (x^2+3*y(x)^2)/(2*x*y(x)),y(x), singsol=all)
```

$$y(x) = \sqrt{c_1 x - 1} x$$

$$y(x) = -\sqrt{c_1 x - 1} x$$

✓ Solution by Mathematica

Time used: 0.251 (sec). Leaf size: 34

```
DSolve[y'[x] == (x^2+3*y[x]^2)/(2*x*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x\sqrt{-1 + c_1 x}$$

$$y(x) \rightarrow x\sqrt{-1 + c_1 x}$$

2.32 problem 33

Internal problem ID [510]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 33.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{4y - 3x}{2x - y} = 0$$

✓ Solution by Maple

Time used: 0.235 (sec). Leaf size: 28

```
dsolve(diff(y(x),x) = (4*y(x)-3*x)/(2*x-y(x)),y(x), singsol=all)
```

$$y(x) = \text{RootOf} \left(_Z^{20} c_1 x^4 - _Z^4 + 4 \right)^4 x - 3x$$

✓ Solution by Mathematica

Time used: 3.328 (sec). Leaf size: 336

```
DSolve[y'[x] == (4*y[x]-3*x)/(2*x-y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{Root}\left[\#1^5 + 15\#1^4x + 90\#1^3x^2 + 270\#1^2x^3 + \#1(405x^4 - e^{4c_1}) + 243x^5 + e^{4c_1}x\&, 1\right]$$

$$y(x) \rightarrow \text{Root}\left[\#1^5 + 15\#1^4x + 90\#1^3x^2 + 270\#1^2x^3 + \#1(405x^4 - e^{4c_1}) + 243x^5 + e^{4c_1}x\&, 2\right]$$

$$y(x) \rightarrow \text{Root}\left[\#1^5 + 15\#1^4x + 90\#1^3x^2 + 270\#1^2x^3 + \#1(405x^4 - e^{4c_1}) + 243x^5 + e^{4c_1}x\&, 3\right]$$

$$y(x) \rightarrow \text{Root}\left[\#1^5 + 15\#1^4x + 90\#1^3x^2 + 270\#1^2x^3 + \#1(405x^4 - e^{4c_1}) + 243x^5 + e^{4c_1}x\&, 4\right]$$

$$y(x) \rightarrow \text{Root}\left[\#1^5 + 15\#1^4x + 90\#1^3x^2 + 270\#1^2x^3 + \#1(405x^4 - e^{4c_1}) + 243x^5 + e^{4c_1}x\&, 5\right]$$

2.33 problem 34

Internal problem ID [511]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 34.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$y' + \frac{4x + 3y}{y + 2x} = 0$$

✓ Solution by Maple

Time used: 0.203 (sec). Leaf size: 1282

`dsolve(diff(y(x),x) = - (4*x+3*y(x))/(2*x+y(x)),y(x), singsol=all)`

$$y(x) = \frac{\left(4\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}-\frac{16x^3c_1}{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}\right)^2}{64c_1x^2} - x^3$$

$$y(x) = \frac{\left(-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)^6\left(4\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}-\frac{16x^3c_1}{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}\right)^2}{64c_1x^2} - x^3$$

$$y(x) = \frac{\left(-\frac{1}{2}+\frac{i\sqrt{3}}{2}\right)^6\left(4\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}-\frac{16x^3c_1}{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}\right)^2}{64c_1x^2} - x^3$$

$$y(x) = \frac{\left(-2\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}+\frac{8x^3c_1}{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}-4i\sqrt{3}\left(\frac{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}{2}+\frac{2x^3c_1}{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}\right)\right)^2}{64c_1x^2} - x^3$$

$$y(x) = \frac{\left(-2\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}+\frac{8x^3c_1}{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}+4i\sqrt{3}\left(\frac{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}{2}+\frac{2x^3c_1}{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}\right)\right)^2}{64c_1x^2} - x^3$$

$$y(x) = \frac{\left(-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)^6\left(-2\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}+\frac{8x^3c_1}{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}-4i\sqrt{3}\left(\frac{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}{2}+\frac{2x^3c_1}{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}\right)\right)^2}{64c_1x^2}$$

$$y(x) = \frac{\left(-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)^6\left(-2\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}+\frac{8x^3c_1}{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}+4i\sqrt{3}\left(\frac{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}{2}+\frac{2x^3c_1}{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}\right)\right)^2}{64c_1x^2}$$

$$y(x) = \frac{72}{64c_1x^2}\left(-\frac{1}{2}+\frac{i\sqrt{3}}{2}\right)^6\left(-2\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}+\frac{8x^3c_1}{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}-4i\sqrt{3}\left(\frac{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}{2}+\frac{2x^3c_1}{\left(4c_1x^3+4\sqrt{4c_1^3x^9+c_1^2x^6}\right)^{\frac{1}{3}}}\right)\right)^2$$

✓ Solution by Mathematica

Time used: 20.375 (sec). Leaf size: 484

`DSolve[y'[x] == - (4*x+3*y[x])/(2*x+y[x]), y[x], x, IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{\sqrt[3]{2x^3 + \sqrt{4e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}}{\sqrt[3]{2}} + \frac{\sqrt[3]{2}x^2}{\sqrt[3]{2x^3 + \sqrt{4e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}} - 3x$$

$$y(x) \rightarrow \frac{i(\sqrt{3} + i) \sqrt[3]{2x^3 + \sqrt{4e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}}{2\sqrt[3]{2}} - \frac{(1 + i\sqrt{3})x^2}{2^{2/3} \sqrt[3]{2x^3 + \sqrt{4e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}} - 3x$$

$$y(x) \rightarrow -\frac{(1 + i\sqrt{3}) \sqrt[3]{2x^3 + \sqrt{4e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}}{2\sqrt[3]{2}} + \frac{i(\sqrt{3} + i)x^2}{2^{2/3} \sqrt[3]{2x^3 + \sqrt{4e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}} - 3x$$

$$y(x) \rightarrow \sqrt[3]{x^3} + \frac{(x^3)^{2/3}}{x} - 3x$$

$$y(x) \rightarrow \frac{1}{2} \left(i(\sqrt{3} + i) \sqrt[3]{x^3} + \frac{(-1 - i\sqrt{3})(x^3)^{2/3}}{x} - 6x \right)$$

$$y(x) \rightarrow \frac{1}{2} \left((-1 - i\sqrt{3}) \sqrt[3]{x^3} + \frac{i(\sqrt{3} + i)(x^3)^{2/3}}{x} - 6x \right)$$

2.34 problem 35

Internal problem ID [512]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 35.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{x + 3y}{x - y} = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 21

```
dsolve(diff(y(x),x) = (x+3*y(x))/(x-y(x)),y(x), singsol=all)
```

$$y(x) = -\frac{x(\text{LambertW}(-2c_1x) + 2)}{\text{LambertW}(-2c_1x)}$$

✓ Solution by Mathematica

Time used: 0.113 (sec). Leaf size: 33

```
DSolve[y'[x] == (x+3*y[x])/(x-y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{2}{\frac{y(x)}{x} + 1} + \log \left(\frac{y(x)}{x} + 1 \right) = -\log(x) + c_1, y(x) \right]$$

2.35 problem 36

Internal problem ID [513]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 36.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _Riccati]`

$$3yx + y^2 - y'x^2 = -x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve((x^2+3*x*y(x)+y(x)^2)-x^2* diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{x(\ln(x) + c_1 + 1)}{\ln(x) + c_1}$$

✓ Solution by Mathematica

Time used: 0.141 (sec). Leaf size: 28

```
DSolve[(x^2+3*x*y[x]+y[x]^2)-x^2* y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x(\log(x) + 1 + c_1)}{\log(x) + c_1}$$

$$y(x) \rightarrow -x$$

2.36 problem 37

Internal problem ID [514]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 37.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$y' - \frac{x^2 - 3y^2}{2yx} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

```
dsolve(diff(y(x),x) = (x^2-3*y(x)^2)/(2*x*y(x)),y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{5} \sqrt{x(x^5 + 5c_1)}}{5x^2}$$

$$y(x) = \frac{\sqrt{5} \sqrt{x(x^5 + 5c_1)}}{5x^2}$$

✓ Solution by Mathematica

Time used: 0.211 (sec). Leaf size: 50

```
DSolve[y'[x] == (x^2-3*y[x]^2)/(2*x*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{\frac{x^5}{5} + c_1}}{x^{3/2}}$$

$$y(x) \rightarrow \frac{\sqrt{\frac{x^5}{5} + c_1}}{x^{3/2}}$$

2.37 problem 38

Internal problem ID [515]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.2. Page 48

Problem number: 38.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$y' - \frac{3y^2 - x^2}{2yx} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 26

```
dsolve(diff(y(x),x) = (3*y(x)^2-x^2)/(2*x*y(x)),y(x), singsol=all)
```

$$y(x) = \sqrt{c_1x + 1}x$$

$$y(x) = -\sqrt{c_1x + 1}x$$

✓ Solution by Mathematica

Time used: 0.209 (sec). Leaf size: 34

```
DSolve[y'[x] == (3*y[x]^2-x^2)/(2*x*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x\sqrt{1 + c_1x}$$

$$y(x) \rightarrow x\sqrt{1 + c_1x}$$

3 Section 2.4. Page 76

3.1	problem 1	79
3.2	problem 2	80
3.3	problem 3	81
3.4	problem 4	82
3.5	problem 5	83
3.6	problem 6	84
3.7	problem 11	85
3.8	problem 12	88
3.9	problem 13	89
3.10	problem 14	90
3.11	problem 15	91
3.12	problem 16	92
3.13	problem 17	93
3.14	problem 18	94
3.15	problem 19	95
3.16	problem 20	96

3.1 problem 1

Internal problem ID [516]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$\ln(t)y + (t - 3)y' = 2t$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 56

```
dsolve(ln(t)*y(t)+(-3+t)*diff(y(t),t) = 2*t,y(t), singsol=all)
```

$$y(t) = 3^{-\ln(-t+3)} \left(\int -2t(-t+3)^{-1+\ln(3)} e^{-\ln(3)^2} e^{-\operatorname{dilog}(\frac{t}{3})} dt + c_1 \right) e^{\ln(3)^2} e^{\operatorname{dilog}(\frac{t}{3})}$$

✓ Solution by Mathematica

Time used: 0.188 (sec). Leaf size: 69

```
DSolve[Log[t]*y[t]+(-3+t)*y'[t] == 2*t,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{\operatorname{PolyLog}(2,1-\frac{t}{3})-\log(3)\log(t-3)} \left(\int_1^t \frac{2e^{\log(3)\log(K[1]-3)-\operatorname{PolyLog}(2,1-\frac{K[1]}{3})} K[1]}{K[1]-3} dK[1] + c_1 \right)$$

3.2 problem 2

Internal problem ID [517]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y + (t - 4)ty' = 0$$

With initial conditions

$$[y(2) = 1]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 19

```
dsolve([y(t)+(-4+t)*t*diff(y(t),t) = 0,y(2) = 1],y(t), singsol=all)
```

$$y(t) = \frac{\left(\frac{1}{2} + \frac{i}{2}\right) \sqrt{2} t^{\frac{1}{4}}}{(-4 + t)^{\frac{1}{4}}}$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 20

```
DSolve[{y[t]+(-4+t)*t*y'[t] == 0,y[2]==1},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{\sqrt[4]{t}}{\sqrt[4]{4-t}}$$

3.3 problem 3

Internal problem ID [518]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$\tan(t)y + y' = \sin(t)$$

With initial conditions

$$[y(\pi) = 0]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve([tan(t)*y(t)+diff(y(t),t) = sin(t),y(Pi) = 0],y(t), singsol=all)
```

$$y(t) = (-\ln(\cos(t)) + i\pi) \cos(t)$$

✓ Solution by Mathematica

Time used: 0.073 (sec). Leaf size: 20

```
DSolve[{Tan[t]*y[t]+y'[t] == Sin[t],y[Pi]==0},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow i \cos(t)(\pi + i \log(\cos(t)))$$

3.4 problem 4

Internal problem ID [519]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$2yt + (-t^2 + 4)y' = 3t^2$$

With initial conditions

$$[y(-3) = 1]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 42

```
dsolve([2*t*y(t)+(-t^2+4)*diff(y(t),t) = 3*t^2,y(-3) = 1],y(t), singsol=all)
```

$$y(t) = \frac{3t}{2} + \frac{3 \ln(2+t)t^2}{8} - \frac{3 \ln(2+t)}{2} - \frac{3 \ln(t-2)t^2}{8} \\ + \frac{3 \ln(t-2)}{2} + \frac{11t^2}{10} - \frac{22}{5} + \frac{3 \ln(5)t^2}{8} - \frac{3 \ln(5)}{2}$$

✓ Solution by Mathematica

Time used: 0.049 (sec). Leaf size: 67

```
DSolve[{2*t*y[t]+(-t^2+4)*y'[t] == 3*t^2,y[-3]==1},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{40}(-15i\pi t^2 + 44t^2 + 15t^2 \log(5) - 15(t^2 - 4) \log(2 - t) + 15(t^2 - 4) \log(t + 2) \\ + 60t + 60i\pi - 176 - 60 \log(5))$$

3.5 problem 5

Internal problem ID [520]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$2yt + (-t^2 + 4)y' = 3t^2$$

With initial conditions

$$[y(1) = -3]$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 46

```
dsolve([2*t*y(t)+(-t^2+4)*diff(y(t),t) = 3*t^2,y(1) = -3],y(t), singsol=all)
```

$$y(t) = -6 + \frac{3(t^2 - 4) \ln(2 + t)}{8} + \frac{3i\pi t^2}{8} - \frac{3 \ln(3) t^2}{8} \\ - \frac{3 \ln(t - 2) t^2}{8} - \frac{3i\pi}{2} + \frac{3t^2}{2} + \frac{3t}{2} + \frac{3 \ln(3)}{2} + \frac{3 \ln(t - 2)}{2}$$

✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 52

```
DSolve[{2*t*y[t]+(-t^2+4)*y'[t] == 3*t^2,y[1]==-3},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -\frac{3}{8}(-4t^2 + t^2 \log(3) + (t^2 - 4) \log(2 - t) - (t^2 - 4) \log(t + 2) - 4t + 16 - 4 \log(3))$$

3.6 problem 6

Internal problem ID [521]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear]`

$$y + \ln(t) y' = \cot(t)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
dsolve(y(t)+ln(t)*diff(y(t),t) = cot(t),y(t), singsol=all)
```

$$y(t) = \left(\int \frac{\cot(t) e^{-\text{Ei}_1(-\ln(t))}}{\ln(t)} dt + c_1 \right) e^{\text{Ei}_1(-\ln(t))}$$

✓ Solution by Mathematica

Time used: 0.142 (sec). Leaf size: 36

```
DSolve[y[t]+Log[t]*y'[t] == Cot[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{-\text{LogIntegral}(t)} \left(\int_1^t \frac{e^{\text{LogIntegral}(K[1])} \cot(K[1])}{\log(K[1])} dK[1] + c_1 \right)$$

3.7 problem 11

Internal problem ID [522]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$y' - \frac{t^2 + 1}{3y - y^2} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 660

`dsolve(diff(y(t),t) = (t^2+1)/(3*y(t)-y(t)^2),y(t), singsol=all)`

$y(t)$

$$= \frac{\left(27 - 4t^3 - 12c_1 - 12t + 2\sqrt{4t^6 + 24c_1t^3 + 24t^4 - 54t^3 + 36c_1^2 + 72c_1t + 36t^2 - 162c_1 - 162t}\right)^{\frac{1}{3}}}{2} + \frac{2\left(27 - 4t^3 - 12c_1 - 12t + 2\sqrt{4t^6 + 24c_1t^3 + 24t^4 - 54t^3 + 36c_1^2 + 72c_1t + 36t^2 - 162c_1 - 162t}\right)^{\frac{1}{3}}}{9} + \frac{3}{2}$$

$y(t) =$

$$-\frac{\left(27 - 4t^3 - 12c_1 - 12t + 2\sqrt{4t^6 + 24c_1t^3 + 24t^4 - 54t^3 + 36c_1^2 + 72c_1t + 36t^2 - 162c_1 - 162t}\right)^{\frac{1}{3}}}{4} - \frac{4\left(27 - 4t^3 - 12c_1 - 12t + 2\sqrt{4t^6 + 24c_1t^3 + 24t^4 - 54t^3 + 36c_1^2 + 72c_1t + 36t^2 - 162c_1 - 162t}\right)^{\frac{1}{3}}}{9} + \frac{3}{2} + \frac{i\sqrt{3}\left(\frac{\left(27 - 4t^3 - 12c_1 - 12t + 2\sqrt{4t^6 + 24c_1t^3 + 24t^4 - 54t^3 + 36c_1^2 + 72c_1t + 36t^2 - 162c_1 - 162t}\right)^{\frac{1}{3}}}{2} - \frac{2\left(27 - 4t^3 - 12c_1 - 12t + 2\sqrt{4t^6 + 24c_1t^3 + 24t^4 - 54t^3 + 36c_1^2 + 72c_1t + 36t^2 - 162c_1 - 162t}\right)^{\frac{1}{3}}}{2}\right)}{2}$$

$y(t) =$

$$-\frac{\left(27 - 4t^3 - 12c_1 - 12t + 2\sqrt{4t^6 + 24c_1t^3 + 24t^4 - 54t^3 + 36c_1^2 + 72c_1t + 36t^2 - 162c_1 - 162t}\right)^{\frac{1}{3}}}{4} - \frac{4\left(27 - 4t^3 - 12c_1 - 12t + 2\sqrt{4t^6 + 24c_1t^3 + 24t^4 - 54t^3 + 36c_1^2 + 72c_1t + 36t^2 - 162c_1 - 162t}\right)^{\frac{1}{3}}}{9} + \frac{3}{2} + \frac{i\sqrt{3}\left(\frac{\left(27 - 4t^3 - 12c_1 - 12t + 2\sqrt{4t^6 + 24c_1t^3 + 24t^4 - 54t^3 + 36c_1^2 + 72c_1t + 36t^2 - 162c_1 - 162t}\right)^{\frac{1}{3}}}{2} - \frac{2\left(27 - 4t^3 - 12c_1 - 12t + 2\sqrt{4t^6 + 24c_1t^3 + 24t^4 - 54t^3 + 36c_1^2 + 72c_1t + 36t^2 - 162c_1 - 162t}\right)^{\frac{1}{3}}}{2}\right)}{2}$$

✓ Solution by Mathematica

Time used: 3.185 (sec). Leaf size: 343

`DSolve[y'[t] == (t^2+1)/(3*y[t]-y[t]^2),y[t],t,IncludeSingularSolutions -> True]`

$$y(t) \rightarrow \frac{1}{2} \left(\sqrt[3]{-4t^3 + \sqrt{-729 + (4t^3 + 12t - 3(9 + 4c_1))^2} - 12t + 27 + 12c_1} + \frac{9}{\sqrt[3]{-4t^3 + \sqrt{-729 + (4t^3 + 12t - 3(9 + 4c_1))^2} - 12t + 27 + 12c_1}} + 3 \right)$$

$$y(t) \rightarrow \frac{1}{4} \left(i(\sqrt{3} + i) \sqrt[3]{-4t^3 + \sqrt{-729 + (4t^3 + 12t - 3(9 + 4c_1))^2} - 12t + 27 + 12c_1} - \frac{9(1 + i\sqrt{3})}{\sqrt[3]{-4t^3 + \sqrt{-729 + (4t^3 + 12t - 3(9 + 4c_1))^2} - 12t + 27 + 12c_1}} + 6 \right)$$

$$y(t) \rightarrow \frac{1}{4} \left(- \left((1 + i\sqrt{3}) \sqrt[3]{-4t^3 + \sqrt{-729 + (4t^3 + 12t - 3(9 + 4c_1))^2} - 12t + 27 + 12c_1} + \frac{9i(\sqrt{3} + i)}{\sqrt[3]{-4t^3 + \sqrt{-729 + (4t^3 + 12t - 3(9 + 4c_1))^2} - 12t + 27 + 12c_1}} + 6 \right) \right)$$

3.8 problem 12

Internal problem ID [523]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{\cot(t)y}{1+y} = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 9

```
dsolve(diff(y(t),t) = cot(t)*y(t)/(1+y(t)),y(t), singsol=all)
```

$$y(t) = \text{LambertW}(c_1 \sin(t))$$

✓ Solution by Mathematica

Time used: 1.602 (sec). Leaf size: 18

```
DSolve[y'[t] == Cot[t]*y[t]/(1+y[t]),y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow W(e^{c_1} \sin(t))$$

$$y(t) \rightarrow 0$$

3.9 problem 13

Internal problem ID [524]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' + \frac{4t}{y} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve(diff(y(t),t) = -4*t/y(t),y(t), singsol=all)
```

$$y(t) = \sqrt{-4t^2 + c_1}$$

$$y(t) = -\sqrt{-4t^2 + c_1}$$

✓ Solution by Mathematica

Time used: 0.087 (sec). Leaf size: 46

```
DSolve[y'[t]== -4*t/y[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -\sqrt{2}\sqrt{-2t^2 + c_1}$$

$$y(t) \rightarrow \sqrt{2}\sqrt{-2t^2 + c_1}$$

3.10 problem 14

Internal problem ID [525]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 14.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - 2ty^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(t),t) = 2*t*y(t)^2,y(t), singsol=all)
```

$$y(t) = \frac{1}{-t^2 + c_1}$$

✓ Solution by Mathematica

Time used: 0.111 (sec). Leaf size: 20

```
DSolve[y'[t] == 2*t*y[t]^2,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -\frac{1}{t^2 + c_1}$$

$$y(t) \rightarrow 0$$

3.11 problem 15

Internal problem ID [526]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y^3 + y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(y(t)^3+diff(y(t),t) = 0,y(t), singsol=all)
```

$$y(t) = \frac{1}{\sqrt{2t + c_1}}$$
$$y(t) = -\frac{1}{\sqrt{2t + c_1}}$$

✓ Solution by Mathematica

Time used: 0.124 (sec). Leaf size: 40

```
DSolve[y[t]^3+y'[t] == 0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -\frac{1}{\sqrt{2t - 2c_1}}$$
$$y(t) \rightarrow \frac{1}{\sqrt{2t - 2c_1}}$$
$$y(t) \rightarrow 0$$

3.12 problem 16

Internal problem ID [527]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{t^2}{(t^3 + 1)y} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
dsolve(diff(y(t),t) = t^2/(t^3+1)/y(t),y(t), singsol=all)
```

$$y(t) = -\frac{\sqrt{6 \ln(t^3 + 1) + 9c_1}}{3}$$

$$y(t) = \frac{\sqrt{6 \ln(t^3 + 1) + 9c_1}}{3}$$

✓ Solution by Mathematica

Time used: 0.123 (sec). Leaf size: 56

```
DSolve[y'[t] == t^2/(t^3+1)/y[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -\sqrt{\frac{2}{3}} \sqrt{\log(t^3 + 1) + 3c_1}$$

$$y(t) \rightarrow \sqrt{\frac{2}{3}} \sqrt{\log(t^3 + 1) + 3c_1}$$

3.13 problem 17

Internal problem ID [528]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 17.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - t(3 - y)y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve(diff(y(t),t) = t*(3-y(t))*y(t),y(t), singsol=all)
```

$$y(t) = \frac{3}{1 + 3e^{-\frac{3t^2}{2}} c_1}$$

✓ Solution by Mathematica

Time used: 0.266 (sec). Leaf size: 44

```
DSolve[y'[t] == t*(3-y[t])*y[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{3e^{\frac{3t^2}{2}}}{e^{\frac{3t^2}{2}} + e^{3c_1}}$$

$$y(t) \rightarrow 0$$

$$y(t) \rightarrow 3$$

3.14 problem 18

Internal problem ID [529]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 18.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_Bernoulli]

$$y' - y(3 - yt) = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 20

```
dsolve(diff(y(t),t) = y(t)*(3-t*y(t)),y(t), singsol=all)
```

$$y(t) = \frac{9}{-1 + 9c_1e^{-3t} + 3t}$$

✓ Solution by Mathematica

Time used: 0.118 (sec). Leaf size: 35

```
DSolve[y'[t] == y[t]*(3-t*y[t]),y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{9e^{3t}}{e^{3t}(3t - 1) + 9c_1}$$

$$y(t) \rightarrow 0$$

3.15 problem 19

Internal problem ID [530]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [Bernoulli]

$$y' + y(3 - yt) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(t),t) = -y(t)*(3-t*y(t)),y(t), singsol=all)
```

$$y(t) = \frac{9}{1 + 9c_1e^{3t} + 3t}$$

✓ Solution by Mathematica

Time used: 0.126 (sec). Leaf size: 28

```
DSolve[y'[t] == -y[t]*(3-t*y[t]),y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{9}{3t + 9c_1e^{3t} + 1}$$

$$y(t) \rightarrow 0$$

3.16 problem 20

Internal problem ID [531]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.4. Page 76

Problem number: 20.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [Riccati]

$$y' + y^2 = t - 1$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 31

```
dsolve(diff(y(t),t) = t-1-y(t)^2,y(t), singsol=all)
```

$$y(t) = \frac{\text{AiryAi}(1, t - 1) c_1 + \text{AiryBi}(1, t - 1)}{\text{AiryAi}(t - 1) c_1 + \text{AiryBi}(t - 1)}$$

✓ Solution by Mathematica

Time used: 0.123 (sec). Leaf size: 47

```
DSolve[y'[t] == t-1-y[t]^2,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{\text{AiryBiPrime}(t - 1) + c_1 \text{AiryAiPrime}(t - 1)}{\text{AiryBi}(t - 1) + c_1 \text{AiryAi}(t - 1)}$$

$$y(t) \rightarrow \frac{\text{AiryAiPrime}(t - 1)}{\text{AiryAi}(t - 1)}$$

4 Section 2.5. Page 88

4.1	problem 1	98
4.2	problem 3	99
4.3	problem 4	101
4.4	problem 5	102
4.5	problem 6	103
4.6	problem 7	104
4.7	problem 9	105
4.8	problem 10	106
4.9	problem 11	108
4.10	problem 12	109
4.11	problem 13	110

4.1 problem 1

Internal problem ID [532]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.5. Page 88

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y' - ay - by^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x) = a*y(x)+b*y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{a}{e^{-ax}c_1a - b}$$

✓ Solution by Mathematica

Time used: 0.888 (sec). Leaf size: 45

```
DSolve[y'[x]== a*y[x]+b*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{ae^{a(x+c_1)}}{-1 + be^{a(x+c_1)}}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -\frac{a}{b}$$

4.2 problem 3

Internal problem ID [533]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.5. Page 88

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - y(-2 + y)(y - 1) = 0$$

✓ Solution by Maple

Time used: 0.203 (sec). Leaf size: 73

```
dsolve(diff(y(t),t) = y(t)*(-2+y(t))*(-1+y(t)),y(t), singsol=all)
```

$$y(t) = -\frac{e^{2t}c_1}{\left(-\frac{1}{\sqrt{-c_1e^{2t}+1}} - 1\right)(c_1e^{2t} - 1)}$$

$$y(t) = -\frac{e^{2t}c_1}{\left(\frac{1}{\sqrt{-c_1e^{2t}+1}} - 1\right)(c_1e^{2t} - 1)}$$

✓ Solution by Mathematica

Time used: 11.055 (sec). Leaf size: 100

```
DSolve[y'[t] == y[t]*(-2+y[t])*(-1+y[t]),y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{-\sqrt{1 + e^{2(t+c_1)}} + e^{2(t+c_1)} + 1}{1 + e^{2(t+c_1)}}$$

$$y(t) \rightarrow \frac{\sqrt{1 + e^{2(t+c_1)}} + e^{2(t+c_1)} + 1}{1 + e^{2(t+c_1)}}$$

$$y(t) \rightarrow 0$$

$$y(t) \rightarrow 1$$

$$y(t) \rightarrow 2$$

4.3 problem 4

Internal problem ID [534]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.5. Page 88

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - e^y = -1$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 15

```
dsolve(diff(y(t),t) = -1+exp(y(t)),y(t), singsol=all)
```

$$y(t) = \ln\left(-\frac{1}{c_1 e^t - 1}\right)$$

✓ Solution by Mathematica

Time used: 0.817 (sec). Leaf size: 28

```
DSolve[y'[t]== -1+Exp[y[t]],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \log\left(\frac{1}{2}\left(1 - \tanh\left(\frac{t + c_1}{2}\right)\right)\right)$$

$$y(t) \rightarrow 0$$

4.4 problem 5

Internal problem ID [535]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.5. Page 88

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - e^{-y} = -1$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 18

```
dsolve(diff(y(t),t) = -1+exp(-y(t)),y(t), singsol=all)
```

$$y(t) = -t + \ln(e^{t+c_1} - 1) - c_1$$

✓ Solution by Mathematica

Time used: 0.853 (sec). Leaf size: 21

```
DSolve[y'[t] == -1+Exp[-y[t]],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \log(1 + e^{-t+c_1})$$

$$y(t) \rightarrow 0$$

4.5 problem 6

Internal problem ID [536]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.5. Page 88

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y' + \frac{2 \arctan(y)}{1 + y^2} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(t),t) = -2*arctan(y(t))/(1+y(t)^2),y(t), singsol=all)
```

$$t + \int^{y(t)} \frac{-a^2 + 1}{2 \arctan(a)} da + c_1 = 0$$

✓ Solution by Mathematica

Time used: 1.013 (sec). Leaf size: 38

```
DSolve[y'[t] == -2*ArcTan[y[t]]/(1+y[t]^2),y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} \frac{K[1]^2 + 1}{\arctan(K[1])} dK[1] \& \right] [-2t + c_1]$$

$$y(t) \rightarrow 0$$

4.6 problem 7

Internal problem ID [537]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.5. Page 88

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' + k(y - 1)^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(t),t) = -k*(-1+y(t))^2,y(t), singsol=all)
```

$$y(t) = \frac{c_1 k + t k + 1}{k(t + c_1)}$$

✓ Solution by Mathematica

Time used: 0.139 (sec). Leaf size: 30

```
DSolve[y'[t]== -k*(-1+y[t])^2,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{kt + 1 - c_1}{kt - c_1}$$

$$y(t) \rightarrow 1$$

4.7 problem 9

Internal problem ID [538]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.5. Page 88

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y' - y^2(y^2 - 1) = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 47

```
dsolve(diff(y(t),t) = y(t)^2*(y(t)^2-1),y(t), singsol=all)
```

$$y(t) = e^{\text{RootOf}(-\ln(e^{-Z}-2)e^{-Z}+2c_1e^{-Z}+Ze^{-Z}+2te^{-Z}+\ln(e^{-Z}-2)-2c_1-Z-2t-2)} - 1$$

✓ Solution by Mathematica

Time used: 0.246 (sec). Leaf size: 51

```
DSolve[y'[t] == y[t]^2*(y[t]^2-1),y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \text{InverseFunction}\left[\frac{1}{\#1} + \frac{1}{2}\log(1 - \#1) - \frac{1}{2}\log(\#1 + 1)\&\right][t + c_1]$$

$$y(t) \rightarrow -1$$

$$y(t) \rightarrow 0$$

$$y(t) \rightarrow 1$$

4.8 problem 10

Internal problem ID [539]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.5. Page 88

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y' - y(1 - y^2) = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 29

```
dsolve(diff(y(t),t) = y(t)*(1-y(t)^2),y(t), singsol=all)
```

$$y(t) = \frac{1}{\sqrt{c_1 e^{-2t} + 1}}$$

$$y(t) = -\frac{1}{\sqrt{c_1 e^{-2t} + 1}}$$

✓ Solution by Mathematica

Time used: 0.676 (sec). Leaf size: 100

```
DSolve[y'[t]== y[t]*(1-y[t]^2),y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -\frac{e^t}{\sqrt{e^{2t} + e^{2c_1}}}$$

$$y(t) \rightarrow \frac{e^t}{\sqrt{e^{2t} + e^{2c_1}}}$$

$$y(t) \rightarrow -1$$

$$y(t) \rightarrow 0$$

$$y(t) \rightarrow 1$$

$$y(t) \rightarrow -\frac{e^t}{\sqrt{e^{2t}}}$$

$$y(t) \rightarrow \frac{e^t}{\sqrt{e^{2t}}}$$

4.9 problem 11

Internal problem ID [540]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.5. Page 88

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y' + b\sqrt{y} - ay = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(t),t) = -b*y(t)^(1/2)+a*y(t),y(t), singsol=all)
```

$$-\frac{b}{a} - e^{\frac{at}{2}} c_1 + \sqrt{y(t)} = 0$$

✓ Solution by Mathematica

Time used: 0.844 (sec). Leaf size: 55

```
DSolve[y'[t] == -b*y[t]^(1/2)+a*y[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{e^{-ac_1} \left(e^{\frac{at}{2}} - be^{\frac{ac_1}{2}} \right)^2}{a^2}$$

$$y(t) \rightarrow 0$$

$$y(t) \rightarrow \frac{b^2}{a^2}$$

4.10 problem 12

Internal problem ID [541]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.5. Page 88

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y' - y^2(4 - y^2) = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 49

```
dsolve(diff(y(t),t) = y(t)^2*(4-y(t)^2),y(t), singsol=all)
```

$$y(t) = e^{\text{RootOf}(\ln(e^{-Z}-4)e^{-Z}+16c_1e^{-Z}-Ze^{-Z}+16t e^{-Z}-2\ln(e^{-Z}-4)-32c_1+2_Z-32t+4) - 2}$$

✓ Solution by Mathematica

Time used: 0.247 (sec). Leaf size: 57

```
DSolve[y'[t] == y[t]^2*(4-y[t]^2),y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \text{InverseFunction}\left[\frac{1}{4\#1} + \frac{1}{16} \log(2 - \#1) - \frac{1}{16} \log(\#1 + 2)\&\right] [-t + c_1]$$

$$y(t) \rightarrow -2$$

$$y(t) \rightarrow 0$$

$$y(t) \rightarrow 2$$

4.11 problem 13

Internal problem ID [542]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.5. Page 88

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y' - (1 - y)^2 y^2 = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 66

```
dsolve(diff(y(t),t) = (1-y(t))^2*y(t)^2,y(t), singsol=all)
```

$$y(t) = e^{\text{RootOf}(-2 \ln(e^{-Z}+1)e^{2-Z}+c_1e^{2-Z}+2_Ze^{2-Z}+te^{2-Z}-2 \ln(e^{-Z}+1)e^{-Z}+c_1e^{-Z}+2_Ze^{-Z}+te^{-Z}+2e^{-Z}+1)} + 1$$

✓ Solution by Mathematica

Time used: 0.365 (sec). Leaf size: 50

```
DSolve[y'[t] == (1-y[t])^2*y[t]^2,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \text{InverseFunction} \left[-\frac{1}{\#1-1} - \frac{1}{\#1} - 2 \log(1-\#1) + 2 \log(\#1) \& \right] [t + c_1]$$

$$y(t) \rightarrow 0$$

$$y(t) \rightarrow 1$$

5 Section 2.6. Page 100

5.1	problem 1	112
5.2	problem 2	113
5.3	problem 3	114
5.4	problem 4	117
5.5	problem 5	118
5.6	problem 6	120
5.7	problem 7	121
5.8	problem 8	122
5.9	problem 9	123
5.10	problem 10	124
5.11	problem 11	125
5.12	problem 12	126
5.13	problem 13	127
5.14	problem 14	128
5.15	problem 19	129
5.16	problem 21	130
5.17	problem 22	131
5.18	problem 25	132
5.19	problem 26	135
5.20	problem 27	136
5.21	problem 28	137
5.22	problem 29	138
5.23	problem 30	139
5.24	problem 30	141
5.25	problem 32	144

5.1 problem 1

Internal problem ID [543]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(-2 + 2y)y' = -3 - 2x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 43

```
dsolve(3+2*x+(-2+2*y(x))*diff(y(x),x) = 0,y(x), singsol=all)
```

$$y(x) = 1 - \sqrt{-x^2 - c_1 - 3x + 1}$$

$$y(x) = 1 + \sqrt{-x^2 - c_1 - 3x + 1}$$

✓ Solution by Mathematica

Time used: 0.115 (sec). Leaf size: 51

```
DSolve[3+2*x+(-2+2*y[x])*y'[x] == 0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 1 - \sqrt{-x^2 - 3x + 1 + 2c_1}$$

$$y(x) \rightarrow 1 + \sqrt{-x^2 - 3x + 1 + 2c_1}$$

5.2 problem 2

Internal problem ID [544]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$4y + (2x - 2y)y' = -2x$$

✓ Solution by Maple

Time used: 0.203 (sec). Leaf size: 56

```
dsolve(2*x+4*y(x)+(2*x-2*y(x))*diff(y(x),x) = 0,y(x), singsol=all)
```

$$-\frac{\ln\left(-\frac{x^2+3xy(x)-y(x)^2}{x^2}\right)}{2} - \frac{\sqrt{13} \operatorname{arctanh}\left(\frac{(-2y(x)+3x)\sqrt{13}}{13x}\right)}{13} - \ln(x) - c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.065 (sec). Leaf size: 63

```
DSolve[2*x+4*y[x]+(2*x-2*y[x])*y'[x]== 0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[\frac{1}{26}\left(\left(13 + \sqrt{13}\right)\log\left(-\frac{2y(x)}{x} + \sqrt{13} + 3\right) - \left(\sqrt{13} - 13\right)\log\left(\frac{2y(x)}{x} + \sqrt{13} - 3\right)\right) = -\log(x) + c_1, y(x)\right]$$

5.3 problem 3

Internal problem ID [545]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_exact`, `_rational`]

$$-2yx + (3 - x^2 + 6y^2) y' = -3x^2 - 2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 582

`dsolve(2+3*x^2-2*x*y(x)+(3-x^2+6*y(x)^2)*diff(y(x),x) = 0,y(x), singsol=all)`

$$\begin{aligned}
 y(x) &= \frac{\left(-54x^3 - 54c_1 - 108x + 6\sqrt{75x^6 + 162c_1x^3 + 378x^4 + 81c_1^2 + 324c_1x + 162x^2 + 162}\right)^{\frac{1}{3}}}{6\left(-\frac{x^2}{6} + \frac{1}{2}\right)} \\
 &- \frac{\left(-54x^3 - 54c_1 - 108x + 6\sqrt{75x^6 + 162c_1x^3 + 378x^4 + 81c_1^2 + 324c_1x + 162x^2 + 162}\right)^{\frac{1}{3}}}{6\left(-\frac{x^2}{6} + \frac{1}{2}\right)} \\
 y(x) &= \frac{\left(-54x^3 - 54c_1 - 108x + 6\sqrt{75x^6 + 162c_1x^3 + 378x^4 + 81c_1^2 + 324c_1x + 162x^2 + 162}\right)^{\frac{1}{3}}}{12\left(-\frac{x^2}{2} + \frac{3}{2}\right)} \\
 &+ \frac{\left(-54x^3 - 54c_1 - 108x + 6\sqrt{75x^6 + 162c_1x^3 + 378x^4 + 81c_1^2 + 324c_1x + 162x^2 + 162}\right)^{\frac{1}{3}}}{12\left(-\frac{x^2}{2} + \frac{3}{2}\right)} \\
 &- \frac{i\sqrt{3}\left(\frac{\left(-54x^3 - 54c_1 - 108x + 6\sqrt{75x^6 + 162c_1x^3 + 378x^4 + 81c_1^2 + 324c_1x + 162x^2 + 162}\right)^{\frac{1}{3}}}{6} + \frac{-x^2 + 3}{2\left(-54x^3 - 54c_1 - 108x + 6\sqrt{75x^6 + 162c_1x^3 + 378x^4 + 81c_1^2 + 324c_1x + 162x^2 + 162}\right)^{\frac{1}{3}}}\right)}{2} \\
 y(x) &= \frac{\left(-54x^3 - 54c_1 - 108x + 6\sqrt{75x^6 + 162c_1x^3 + 378x^4 + 81c_1^2 + 324c_1x + 162x^2 + 162}\right)^{\frac{1}{3}}}{12\left(-\frac{x^2}{2} + \frac{3}{2}\right)} \\
 &+ \frac{\left(-54x^3 - 54c_1 - 108x + 6\sqrt{75x^6 + 162c_1x^3 + 378x^4 + 81c_1^2 + 324c_1x + 162x^2 + 162}\right)^{\frac{1}{3}}}{12\left(-\frac{x^2}{2} + \frac{3}{2}\right)} \\
 &- \frac{i\sqrt{3}\left(\frac{\left(-54x^3 - 54c_1 - 108x + 6\sqrt{75x^6 + 162c_1x^3 + 378x^4 + 81c_1^2 + 324c_1x + 162x^2 + 162}\right)^{\frac{1}{3}}}{6} + \frac{-x^2 + 3}{2\left(-54x^3 - 54c_1 - 108x + 6\sqrt{75x^6 + 162c_1x^3 + 378x^4 + 81c_1^2 + 324c_1x + 162x^2 + 162}\right)^{\frac{1}{3}}}\right)}{2} \\
 &+ \frac{\left(-54x^3 - 54c_1 - 108x + 6\sqrt{75x^6 + 162c_1x^3 + 378x^4 + 81c_1^2 + 324c_1x + 162x^2 + 162}\right)^{\frac{1}{3}}}{2}
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 8.724 (sec). Leaf size: 421

`DSolve[2+3*x^2-2*x*y[x]+(3-x^2+6*y[x]^2)*y'[x] == 0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow -\frac{x^2 - 3}{\frac{\sqrt[3]{6}\sqrt[3]{9x^3 + \sqrt{3}\sqrt{-2(x^2 - 3)^3 + 27(x^3 + 2x + c_1)^2 + 18x + 9c_1}}}{6^{2/3}}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{6}(1 + i\sqrt{3})(x^2 - 3) + (1 - i\sqrt{3})\left(9x^3 + \sqrt{3}\sqrt{-2(x^2 - 3)^3 + 27(x^3 + 2x + c_1)^2 + 18x + 9c_1}\right)^{2/3}}{2 \cdot 6^{2/3}\sqrt[3]{9x^3 + \sqrt{3}\sqrt{-2(x^2 - 3)^3 + 27(x^3 + 2x + c_1)^2 + 18x + 9c_1}}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{6}(1 - i\sqrt{3})(x^2 - 3) + (1 + i\sqrt{3})\left(9x^3 + \sqrt{3}\sqrt{-2(x^2 - 3)^3 + 27(x^3 + 2x + c_1)^2 + 18x + 9c_1}\right)^{2/3}}{2 \cdot 6^{2/3}\sqrt[3]{9x^3 + \sqrt{3}\sqrt{-2(x^2 - 3)^3 + 27(x^3 + 2x + c_1)^2 + 18x + 9c_1}}}$$

5.4 problem 4

Internal problem ID [546]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$2y + 2xy^2 + (2x + 2x^2y) y' = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 31

```
dsolve(2*y(x)+2*x*y(x)^2+(2*x+2*x^2*y(x))*diff(y(x),x) = 0,y(x), singsol=all)
```

$$y(x) = -\frac{1}{x}$$

$$y(x) = \frac{-1 - c_1}{x}$$

$$y(x) = \frac{c_1 - 1}{x}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 29

```
DSolve[2*y[x]+2*x*y[x]^2+(2*x+2*x^2*y[x])*y'[x] == 0,y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow -\frac{1}{x}$$

$$y(x) \rightarrow \frac{c_1}{x}$$

$$y(x) \rightarrow -\frac{1}{x}$$

5.5 problem 5

Internal problem ID [547]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{-ax - by}{bx + cy} = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 83

```
dsolve(diff(y(x),x) = (-a*x-b*y(x))/(b*x+c*y(x)),y(x), singsol=all)
```

$$y(x) = -\frac{bxc_1 - \sqrt{-ac c_1^2 x^2 + b^2 c_1^2 x^2 + c}}{cc_1}$$

$$y(x) = -\frac{bxc_1 + \sqrt{-ac c_1^2 x^2 + b^2 c_1^2 x^2 + c}}{cc_1}$$

✓ Solution by Mathematica

Time used: 17.783 (sec). Leaf size: 139

```
DSolve[y'[x]== (-a*x-b*y[x])/(b*x+c*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{bx + \sqrt{-acx^2 + b^2x^2 + ce^{2c_1}}}{c}$$

$$y(x) \rightarrow \frac{-bx + \sqrt{b^2x^2 + c(-ax^2 + e^{2c_1})}}{c}$$

$$y(x) \rightarrow -\frac{\sqrt{x^2(b^2 - ac)} + bx}{c}$$

$$y(x) \rightarrow \frac{\sqrt{x^2(b^2 - ac)} - bx}{c}$$

5.6 problem 6

Internal problem ID [548]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{-ax + by}{bx - cy} = 0$$

✓ Solution by Maple

Time used: 0.609 (sec). Leaf size: 52

```
dsolve(diff(y(x),x) = (-a*x+b*y(x))/(b*x-c*y(x)),y(x), singsol=all)
```

$$y(x) = \text{RootOf} \left(c_Z^2 - a - e^{\text{RootOf} \left(\tanh \left(\frac{\sqrt{ac} (2c_1 + _Z + 2 \ln(x))}{2b} \right)^2 a - a - e^{-Z} \right)} \right) x$$

✓ Solution by Mathematica

Time used: 0.072 (sec). Leaf size: 58

```
DSolve[y'[x] == (-a*x+b*y[x])/(b*x-c*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[-\frac{\text{barctanh} \left(\frac{\sqrt{cy(x)}}{\sqrt{ax}} \right)}{\sqrt{a}\sqrt{c}} - \frac{1}{2} \log \left(\frac{cy(x)^2}{x^2} - a \right) = \log(x) + c_1, y(x) \right]$$

5.7 problem 7

Internal problem ID [549]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [exact]

$$e^x \sin(y) - 2 \sin(x) y + (2 \cos(x) + e^x \cos(y)) y' = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 17

```
dsolve(exp(x)*sin(y(x))-2*sin(x)*y(x)+(2*cos(x)+exp(x)*cos(y(x)))*diff(y(x),x) = 0,y(x), sin
```

$$e^x \sin(y(x)) + 2 \cos(x) y(x) + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.282 (sec). Leaf size: 20

```
DSolve[Exp[x]*Sin[y[x]]-2*SIn[x]*y[x]+(2*Cos[x]+Exp[x]*Cos[y[x]])*y'[x] == 0,y[x],x,IncludeS
```

$$\text{Solve}[e^x \sin(y(x)) + 2y(x) \cos(x) = c_1, y(x)]$$

5.8 problem 8

Internal problem ID [550]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type ['x=_G(y,y')']

$$e^x \sin(y) + 3y - (3x - e^x \sin(y))y' = 0$$

X Solution by Maple

```
dsolve(exp(x)*sin(y(x))+3*y(x)-(3*x-exp(x)*sin(y(x)))*diff(y(x),x) = 0,y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[Exp[x]*Sin[y[x]]+3*y[x]-(3*x-Exp[x]*Sin[y[x]])*y'[x] == 0,y[x],x,IncludeSingularSolut
```

Not solved

5.9 problem 9

Internal problem ID [551]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_exact]

$$-2e^{yx} \sin(2x) + e^{yx} \cos(2x) y + (-3 + e^{yx} x \cos(2x)) y' = -2x$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 40

```
dsolve(2*x-2*exp(x*y(x))*sin(2*x)+exp(x*y(x))*cos(2*x)*y(x)+(-3+exp(x*y(x))*x*cos(2*x))*diff
```

$$y(x) = -\frac{-x^3 - c_1 x + 3 \operatorname{LambertW}\left(-\frac{x \cos(2x) e^{\frac{x^3}{3}} e^{\frac{c_1 x}{3}}}{3}\right)}{3x}$$

✓ Solution by Mathematica

Time used: 5.197 (sec). Leaf size: 48

```
DSolve[2*x-2*Exp[x*y[x]]*Sin[2*x]+Exp[x*y[x]]*Cos[2*x]*y[x]+(-3+Exp[x*y[x]]*x*cos[2*x])*y'[x]
```

$$y(x) \rightarrow \frac{-3W\left(-\frac{1}{3}x e^{\frac{1}{3}x(x^2-c_1)} \cos(2x)\right) + x^3 - c_1 x}{3x}$$

5.10 problem 10

Internal problem ID [552]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$\frac{y}{x} + (\ln(x) - 2)y' = -6x$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 18

```
dsolve((y(x)/x+6*x)+(ln(x)-2)*diff(y(x),x) = 0,y(x), singsol=all)
```

$$y(x) = \frac{-3x^2 + c_1}{\ln(x) - 2}$$

✓ Solution by Mathematica

Time used: 0.079 (sec). Leaf size: 20

```
DSolve[(y[x]/x+6*x)+(Log[x]-2)*y'[x] == 0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-3x^2 + c_1}{\log(x) - 2}$$

5.11 problem 11

Internal problem ID [553]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_Abel, '2nd type', 'class B']]`

$$yx + (\ln(x)y + yx)y' = -x \ln(x)$$

X Solution by Maple

```
dsolve((x*ln(x)+x*y(x))+(y(x)*ln(x)+x*y(x))*diff(y(x),x) = 0,y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(x*Log[x]+x*y[x])+(y[x]*Log[x]+x*y[x])*y'[x] == 0,y[x],x,IncludeSingularSolutions ->
```

Not solved

5.12 problem 12

Internal problem ID [554]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$\frac{x}{(x^2 + y^2)^{\frac{3}{2}}} + \frac{yy'}{(x^2 + y^2)^{\frac{3}{2}}} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(x/(x^2+y(x)^2)^(3/2)+y(x)*diff(y(x),x)/(x^2+y(x)^2)^(3/2) = 0,y(x), singsol=all)
```

$$y(x) = \sqrt{-x^2 + c_1}$$

$$y(x) = -\sqrt{-x^2 + c_1}$$

✓ Solution by Mathematica

Time used: 0.09 (sec). Leaf size: 39

```
DSolve[x/(x^2+y[x]^2)^(3/2)+y[x]*y'[x]/(x^2+y[x]^2)^(3/2) == 0,y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow -\sqrt{-x^2 + 2c_1}$$

$$y(x) \rightarrow \sqrt{-x^2 + 2c_1}$$

5.13 problem 13

Internal problem ID [555]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _exact, _rational, [_Abel, '2nd ty`

$$-y + (-x + 2y)y' = -2x$$

With initial conditions

$$[y(1) = 3]$$

✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 19

```
dsolve([2*x-y(x)+(-x+2*y(x))*diff(y(x),x) = 0,y(1) = 3],y(x), singsol=all)
```

$$y(x) = \frac{x}{2} + \frac{\sqrt{-3x^2 + 28}}{2}$$

✓ Solution by Mathematica

Time used: 0.456 (sec). Leaf size: 22

```
DSolve[{2*x-y[x]+(-x+2*y[x])*y'[x] == 0,y[1]==3},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(\sqrt{28 - 3x^2} + x \right)$$

5.14 problem 14

Internal problem ID [556]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 14.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_exact, _rational, [_1st_order, ' _with_symmetry_[F(x),G(x)]]'`

$$y + (x - 4y)y' = -9x^2 + 1$$

With initial conditions

$$[y(1) = 0]$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 25

```
dsolve([-1+9*x^2+y(x)+(x-4*y(x))*diff(y(x),x) = 0,y(1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{x}{4} - \frac{\sqrt{24x^3 + x^2 - 8x - 16}}{4}$$

✓ Solution by Mathematica

Time used: 0.148 (sec). Leaf size: 34

```
DSolve[{-1+9*x^2+y[x]+(x-4*y[x])*y'[x] == 0,y[1]==0},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{1}{4} \left(x + i\sqrt{-24x^3 - x^2 + 8x + 16} \right)$$

5.15 problem 19

Internal problem ID [557]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y^3 x^2 + x(1 + y^2) y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve(x^2*y(x)^3+x*(1+y(x)^2)*diff(y(x),x) = 0,y(x), singsol=all)
```

$$y(x) = e^{\frac{\text{LambertW}\left(\frac{e^{x^2+2c_1}}{2}\right) - \frac{x^2}{2} - c_1}{2}}$$

✓ Solution by Mathematica

Time used: 4.095 (sec). Leaf size: 46

```
DSolve[x^2*y[x]^3+x*(1+y[x]^2)*y'[x] == 0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\sqrt{W(e^{x^2-2c_1})}}$$

$$y(x) \rightarrow \frac{1}{\sqrt{W(e^{x^2-2c_1})}}$$

$$y(x) \rightarrow 0$$

5.16 problem 21

Internal problem ID [558]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 21.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x)*G(y),0]']]`

$$y + (2x - e^y y) y' = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 27

```
dsolve(y(x)+(2*x-exp(y(x))*y(x))*diff(y(x),x) = 0,y(x), singsol=all)
```

$$x - \frac{(y(x)^2 - 2y(x) + 2) e^{y(x)} + c_1}{y(x)^2} = 0$$

✓ Solution by Mathematica

Time used: 0.225 (sec). Leaf size: 32

```
DSolve[y[x]+(2*x-Exp[y[x]]*y[x])*y'[x] == 0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[x = \frac{e^{y(x)}(y(x)^2 - 2y(x) + 2)}{y(x)^2} + \frac{c_1}{y(x)^2}, y(x) \right]$$

5.17 problem 22

Internal problem ID [559]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 22.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(2 + x) \sin(y) + x \cos(y) y' = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve((2+x)*sin(y(x))+x*cos(y(x))*diff(y(x),x) = 0,y(x), singsol=all)
```

$$y(x) = \arcsin\left(\frac{e^{-x}}{c_1 x^2}\right)$$

✓ Solution by Mathematica

Time used: 51.022 (sec). Leaf size: 23

```
DSolve[(2+x)*Sin[y[x]]+x*Cos[y[x]]*y'[x] == 0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arcsin\left(\frac{e^{-x+c_1}}{x^2}\right)$$

$$y(x) \rightarrow 0$$

5.18 problem 25

Internal problem ID [560]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 25.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class D'], _rational]`

$$2yx + 3x^2y + y^3 + (x^2 + y^2)y' = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 420

`dsolve(2*x*y(x)+3*x^2*y(x)+y(x)^3+(x^2+y(x)^2)*diff(y(x),x) = 0,y(x), singsol=all)`

$$y(x) = \frac{e^{-3x} \left(\left(4 + 4\sqrt{4x^6 e^{6x} c_1^2 + 1} \right) e^{6x} c_1^2 \right)^{\frac{1}{3}}}{2c_1} - \frac{2x^2 e^{3x} c_1}{\left(\left(4 + 4\sqrt{4x^6 e^{6x} c_1^2 + 1} \right) e^{6x} c_1^2 \right)^{\frac{1}{3}}}$$

$$y(x) = -\frac{e^{-3x} \left(\left(4 + 4\sqrt{4x^6 e^{6x} c_1^2 + 1} \right) e^{6x} c_1^2 \right)^{\frac{1}{3}}}{4c_1} + \frac{x^2 e^{3x} c_1}{\left(\left(4 + 4\sqrt{4x^6 e^{6x} c_1^2 + 1} \right) e^{6x} c_1^2 \right)^{\frac{1}{3}}}$$

$$- \frac{i\sqrt{3} \left(\frac{e^{-3x} \left(\left(4 + 4\sqrt{4x^6 e^{6x} c_1^2 + 1} \right) e^{6x} c_1^2 \right)^{\frac{1}{3}}}{2c_1} + \frac{2x^2 e^{3x} c_1}{\left(\left(4 + 4\sqrt{4x^6 e^{6x} c_1^2 + 1} \right) e^{6x} c_1^2 \right)^{\frac{1}{3}}} \right)}{2}$$

$$y(x) = -\frac{e^{-3x} \left(\left(4 + 4\sqrt{4x^6 e^{6x} c_1^2 + 1} \right) e^{6x} c_1^2 \right)^{\frac{1}{3}}}{4c_1} + \frac{x^2 e^{3x} c_1}{\left(\left(4 + 4\sqrt{4x^6 e^{6x} c_1^2 + 1} \right) e^{6x} c_1^2 \right)^{\frac{1}{3}}}$$

$$+ \frac{i\sqrt{3} \left(\frac{e^{-3x} \left(\left(4 + 4\sqrt{4x^6 e^{6x} c_1^2 + 1} \right) e^{6x} c_1^2 \right)^{\frac{1}{3}}}{2c_1} + \frac{2x^2 e^{3x} c_1}{\left(\left(4 + 4\sqrt{4x^6 e^{6x} c_1^2 + 1} \right) e^{6x} c_1^2 \right)^{\frac{1}{3}}} \right)}{2}$$

✓ Solution by Mathematica

Time used: 60.305 (sec). Leaf size: 383

`DSolve[2*x*y[x]+3*x^2*y[x]+y[x]^3+(x^2+y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions->`

$$y(x) \rightarrow \frac{e^{-3x} \left(-2e^{6x} x^2 + \sqrt[3]{2} \left(\sqrt{4e^{18x} x^6 + e^{6(2x+c_1)}} + e^{6x+3c_1} \right)^{2/3} \right)}{2^{2/3} \sqrt[3]{\sqrt{4e^{18x} x^6 + e^{6(2x+c_1)}} + e^{6x+3c_1}}}$$

$$y(x) \rightarrow \frac{i(\sqrt{3} + i) e^{-3x} \sqrt[3]{\sqrt{4e^{18x} x^6 + e^{6(2x+c_1)}} + e^{6x+3c_1}}}{2\sqrt[3]{2}} + \frac{(1 + i\sqrt{3}) e^{3x} x^2}{2^{2/3} \sqrt[3]{\sqrt{4e^{18x} x^6 + e^{6(2x+c_1)}} + e^{6x+3c_1}}}$$

$$y(x) \rightarrow \frac{(1 - i\sqrt{3}) e^{3x} x^2}{2^{2/3} \sqrt[3]{\sqrt{4e^{18x} x^6 + e^{6(2x+c_1)}} + e^{6x+3c_1}}} - \frac{(1 + i\sqrt{3}) e^{-3x} \sqrt[3]{\sqrt{4e^{18x} x^6 + e^{6(2x+c_1)}} + e^{6x+3c_1}}}{2\sqrt[3]{2}}$$

5.19 problem 26

Internal problem ID [561]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 26.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' - y = e^{2x} - 1$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve(diff(y(x),x) = -1+exp(2*x)+y(x),y(x), singsol=all)
```

$$y(x) = e^{2x} + 1 + e^x c_1$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 18

```
DSolve[y'[x] == -1+Exp[2*x]+y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x} + c_1 e^x + 1$$

5.20 problem 27

Internal problem ID [562]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 27.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x)*G(y),0]']]`

$$\left(-\sin(y) + \frac{x}{y}\right) y' = -1$$

✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 23

```
dsolve(1+(-sin(y(x))+x/y(x))*diff(y(x),x) = 0,y(x), singsol=all)
```

$$x - \frac{-y(x) \cos(y(x)) + \sin(y(x)) + c_1}{y(x)} = 0$$

✓ Solution by Mathematica

Time used: 0.143 (sec). Leaf size: 29

```
DSolve[1+(-Sin[y[x]]+x/y[x])*y'[x] == 0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[x = \frac{\sin(y(x)) - y(x) \cos(y(x))}{y(x)} + \frac{c_1}{y(x)}, y(x)\right]$$

5.21 problem 28

Internal problem ID [563]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 28.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, _with_exponential_symmetries]]`

$$y + (-e^{-2y} + 2yx) y' = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 24

```
dsolve(y(x)+(-exp(-2*y(x))+2*x*y(x))*diff(y(x),x) = 0,y(x), singsol=all)
```

$$y(x) = e^{\text{RootOf}(c_1 e^{-2e^{-Z}} + Z e^{-2e^{-Z}} - x)}$$

✓ Solution by Mathematica

Time used: 0.243 (sec). Leaf size: 25

```
DSolve[y[x]+(-Exp[-2*y[x]]+2*x*y[x])*y'[x] == 0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}[x = e^{-2y(x)} \log(y(x)) + c_1 e^{-2y(x)}, y(x)]$$

5.22 problem 29

Internal problem ID [564]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 29.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x)*G(y),0]']]`

$$(e^x \cot(y) + 2 \csc(y) y) y' = -e^x$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 15

```
dsolve(exp(x)+(exp(x)*cot(y(x))+2*csc(y(x))*y(x))*diff(y(x),x) = 0,y(x), singsol=all)
```

$$e^x \sin(y(x)) + y(x)^2 + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.312 (sec). Leaf size: 18

```
DSolve[Exp[x]+(Exp[x]*Cot[y[x]]+2*Csc[y[x]]*y[x])*y'[x] == 0,y[x],x,IncludeSingularSolutions
```

$$\text{Solve}[y(x)^2 + e^x \sin(y(x)) = c_1, y(x)]$$

5.23 problem 30

Internal problem ID [565]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 30.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [rational]

$$\frac{4x^3}{y^2} + \frac{3}{y} + \left(\frac{3x}{y^2} + 4y \right) y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve(4*x^3/y(x)^2+3/y(x)+(3*x/y(x)^2+4*y(x))*diff(y(x),x) = 0,y(x), singsol=all)
```

$$x^4 + y(x)^4 + 3xy(x) + c_1 = 0$$

✓ Solution by Mathematica

Time used: 60.148 (sec). Leaf size: 1181

`DSolve[4*x^3/y[x]^2+3/y[x]+(3*x/y[x]^2+4*y[x])*y'[x]== 0,y[x],x,IncludeSingularSolutions ->`

$y(x) \rightarrow$

$$-\frac{1}{2} \sqrt{\frac{6x}{\sqrt{\frac{4\sqrt[3]{2}(x^4-c_1)}{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}} + \frac{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}}}{3\sqrt[3]{2}}}} - \sqrt[3]{243x^2}}$$

$$-\frac{1}{2} \sqrt{\frac{4\sqrt[3]{2}(x^4 - c_1)}{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}} + \frac{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}}}{3\sqrt[3]{2}}}}$$

$y(x)$

$$\rightarrow \frac{1}{2} \sqrt{\frac{6x}{\sqrt{\frac{4\sqrt[3]{2}(x^4-c_1)}{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}} + \frac{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}}}{3\sqrt[3]{2}}}} - \sqrt[3]{243x^2}}$$

$$-\frac{1}{2} \sqrt{\frac{4\sqrt[3]{2}(x^4 - c_1)}{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}} + \frac{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}}}{3\sqrt[3]{2}}}}$$

$y(x)$

$$\rightarrow \frac{1}{2} \sqrt{\frac{4\sqrt[3]{2}(x^4 - c_1)}{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}} + \frac{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}}}{3\sqrt[3]{2}}}}$$

$$-\frac{1}{2} \sqrt{\frac{6x}{\sqrt{\frac{4\sqrt[3]{2}(x^4-c_1)}{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}} + \frac{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}}}{3\sqrt[3]{2}}}} - \sqrt[3]{243x^2}}$$

$y(x)$

$$\rightarrow \frac{1}{2} \sqrt{\frac{6x}{\sqrt{\frac{4\sqrt[3]{2}(x^4-c_1)}{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}} + \frac{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}}}{3\sqrt[3]{2}}}} - \sqrt[3]{243x^2}}$$

$$\frac{1}{2} \sqrt{\frac{4\sqrt[3]{2}(x^4 - c_1)}{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}} + \frac{\sqrt[3]{243x^2 + \sqrt{59049x^4 - 6912(x^4 - c_1)^3}}}{3\sqrt[3]{2}}}}$$

5.24 problem 30

Internal problem ID [566]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 30.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational]`

$$\frac{6}{y} + \left(\frac{x^2}{y} + \frac{3y}{x} \right) y' = -3x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 430

`dsolve(3*x+6/y(x)+(x^2/y(x)+3*y(x)/x)*diff(y(x),x) = 0,y(x), singsol=all)`

$$\begin{aligned}
 y(x) &= \frac{\left(-324x^2 - 108c_1 + 12\sqrt{12x^9 + 729x^4 + 486c_1x^2 + 81c_1^2}\right)^{\frac{1}{3}}}{6} \\
 &\quad - \frac{\left(-324x^2 - 108c_1 + 12\sqrt{12x^9 + 729x^4 + 486c_1x^2 + 81c_1^2}\right)^{\frac{1}{3}}}{2x^3} \\
 y(x) &= -\frac{\left(-324x^2 - 108c_1 + 12\sqrt{12x^9 + 729x^4 + 486c_1x^2 + 81c_1^2}\right)^{\frac{1}{3}}}{12} \\
 &\quad + \frac{\left(-324x^2 - 108c_1 + 12\sqrt{12x^9 + 729x^4 + 486c_1x^2 + 81c_1^2}\right)^{\frac{1}{3}}}{x^3} \\
 &\quad + \frac{\left(-324x^2 - 108c_1 + 12\sqrt{12x^9 + 729x^4 + 486c_1x^2 + 81c_1^2}\right)^{\frac{1}{3}}}{2} \\
 &\quad + \frac{i\sqrt{3} \left(\frac{\left(-324x^2 - 108c_1 + 12\sqrt{12x^9 + 729x^4 + 486c_1x^2 + 81c_1^2}\right)^{\frac{1}{3}}}{6} + \frac{2x^3}{\left(-324x^2 - 108c_1 + 12\sqrt{12x^9 + 729x^4 + 486c_1x^2 + 81c_1^2}\right)^{\frac{1}{3}}} \right)}{2} \\
 y(x) &= -\frac{\left(-324x^2 - 108c_1 + 12\sqrt{12x^9 + 729x^4 + 486c_1x^2 + 81c_1^2}\right)^{\frac{1}{3}}}{12} \\
 &\quad + \frac{\left(-324x^2 - 108c_1 + 12\sqrt{12x^9 + 729x^4 + 486c_1x^2 + 81c_1^2}\right)^{\frac{1}{3}}}{x^3} \\
 &\quad + \frac{i\sqrt{3} \left(\frac{\left(-324x^2 - 108c_1 + 12\sqrt{12x^9 + 729x^4 + 486c_1x^2 + 81c_1^2}\right)^{\frac{1}{3}}}{6} + \frac{2x^3}{\left(-324x^2 - 108c_1 + 12\sqrt{12x^9 + 729x^4 + 486c_1x^2 + 81c_1^2}\right)^{\frac{1}{3}}} \right)}{2}
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 4.558 (sec). Leaf size: 331

`DSolve[3*x+6/y[x]+(x^2/y[x]+3*y[x]/x)*y'[x] == 0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{\sqrt[3]{-81x^2 + \sqrt{108x^9 + 729(-3x^2 + c_1)^2 + 27c_1}}}{3\sqrt[3]{2}\sqrt[3]{2x^3}} - \frac{\sqrt[3]{-81x^2 + \sqrt{108x^9 + 729(-3x^2 + c_1)^2 + 27c_1}}}{\sqrt[3]{2x^3}}$$

$$y(x) \rightarrow \frac{(-1 + i\sqrt{3})\sqrt[3]{-81x^2 + \sqrt{108x^9 + 729(-3x^2 + c_1)^2 + 27c_1}}}{6\sqrt[3]{2}} + \frac{(1 + i\sqrt{3})x^3}{2^{2/3}\sqrt[3]{-81x^2 + \sqrt{108x^9 + 729(-3x^2 + c_1)^2 + 27c_1}}}$$

$$y(x) \rightarrow \frac{(1 - i\sqrt{3})x^3}{2^{2/3}\sqrt[3]{-81x^2 + \sqrt{108x^9 + 729(-3x^2 + c_1)^2 + 27c_1}}} - \frac{(1 + i\sqrt{3})\sqrt[3]{-81x^2 + \sqrt{108x^9 + 729(-3x^2 + c_1)^2 + 27c_1}}}{6\sqrt[3]{2}}$$

5.25 problem 32

Internal problem ID [567]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Section 2.6. Page 100

Problem number: 32.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$3yx + y^2 + (x^2 + yx)y' = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 59

```
dsolve(3*x*y(x)+y(x)^2+(x^2+x*y(x))*diff(y(x),x) = 0,y(x), singsol=all)
```

$$y(x) = \frac{-c_1x^2 - \sqrt{c_1^2x^4 + 1}}{c_1x}$$

$$y(x) = \frac{-c_1x^2 + \sqrt{c_1^2x^4 + 1}}{c_1x}$$

✓ Solution by Mathematica

Time used: 0.661 (sec). Leaf size: 93

```
DSolve[3*x*y[x]+y[x]^2+(x^2+x*y[x])*y'[x]== 0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x^2 + \sqrt{x^4 + e^{2c_1}}}{x}$$

$$y(x) \rightarrow -x + \frac{\sqrt{x^4 + e^{2c_1}}}{x}$$

$$y(x) \rightarrow -\frac{\sqrt{x^4 + x^2}}{x}$$

$$y(x) \rightarrow \frac{\sqrt{x^4}}{x} - x$$

6 Miscellaneous problems, end of chapter 2. Page 133

6.1	problem 1	147
6.2	problem 2	148
6.3	problem 3	149
6.4	problem 4	150
6.5	problem 5	151
6.6	problem 6	152
6.7	problem 7	153
6.8	problem 8	156
6.9	problem 9	157
6.10	problem 10	158
6.11	problem 11	159
6.12	problem 12	160
6.13	problem 13	161
6.14	problem 14	162
6.15	problem 15	163
6.16	problem 16	164
6.17	problem 17	165
6.18	problem 18	166
6.19	problem 19	167
6.20	problem 20	170
6.21	problem 21	171
6.22	problem 22	174
6.23	problem 23	175
6.24	problem 24	176
6.25	problem 25	177
6.26	problem 26	178
6.27	problem 27	179
6.28	problem 28	180
6.29	problem 29	181
6.30	problem 30	182
6.31	problem 31	183

6.1 problem 1

Internal problem ID [568]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' - \frac{x^3 - 2y}{x} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x) = (x^3-2*y(x))/x,y(x), singsol=all)
```

$$y(x) = \frac{\frac{x^5}{5} + c_1}{x^2}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 19

```
DSolve[y'[x]== (x^3-2*y[x])/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{5} + \frac{c_1}{x^2}$$

6.2 problem 2

Internal problem ID [569]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{\cos(x) + 1}{2 - \sin(y)} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x) = (1+cos(x))/(2-sin(y(x))),y(x), singsol=all)
```

$$x + \sin(x) - 2y(x) - \cos(y(x)) + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.372 (sec). Leaf size: 27

```
DSolve[y'[x] == (1+Cos[x])/(2-Sin[y[x]]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{InverseFunction}[-2\#1 - \cos(\#1)\&][-x - \sin(x) + c_1]$$

6.3 problem 3

Internal problem ID [570]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [rational]

$$y' - \frac{y + 2x}{3 - x + 3y^2} = 0$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.109 (sec). Leaf size: 75

```
dsolve([diff(y(x),x) = (2*x+y(x))/(3-x+3*y(x)^2),y(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{(108x^2 + 12\sqrt{81x^4 - 12x^3 + 108x^2 - 324x + 324})^{\frac{2}{3}} + 12x - 36}{6(108x^2 + 12\sqrt{81x^4 - 12x^3 + 108x^2 - 324x + 324})^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 5.408 (sec). Leaf size: 114

```
DSolve[{y'[x] == (2*x+y[x])/(3-x+3*y[x]^2),y[0]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt[3]{2}(\sqrt{3}\sqrt{27x^4 - 4x^3 + 36x^2 - 108x + 108} - 9x^2)^{2/3} + 2\sqrt[3]{3}x - 6\sqrt[3]{3}}{6^{2/3}\sqrt[3]{\sqrt{3}\sqrt{27x^4 - 4x^3 + 36x^2 - 108x + 108} - 9x^2}}$$

6.4 problem 4

Internal problem ID [571]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - y + 2yx = -6x + 3$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x) = 3-6*x+y(x)-2*x*y(x),y(x), singsol=all)
```

$$y(x) = -3 + e^{-x(x-1)}c_1$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 24

```
DSolve[y'[x] == 3-6*x+y[x]-2*x*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -3 + c_1 e^{x-x^2}$$

$$y(x) \rightarrow -3$$

6.5 problem 5

Internal problem ID [572]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, [_Abel, '2nd type', 'class B']]`

$$y' - \frac{-1 - 2yx - y^2}{x^2 + 2yx} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 57

```
dsolve(diff(y(x),x) = (-1-2*x*y(x)-y(x)^2)/(x^2+2*x*y(x)),y(x), singsol=all)
```

$$y(x) = \frac{-x^2 + \sqrt{x^4 - 4c_1x - 4x^2}}{2x}$$

$$y(x) = -\frac{x^2 + \sqrt{x^4 - 4c_1x - 4x^2}}{2x}$$

✓ Solution by Mathematica

Time used: 0.502 (sec). Leaf size: 67

```
DSolve[y'[x] == (-1-2*x*y[x]-y[x]^2)/(x^2+2*x*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x^2 + \sqrt{x(x^3 - 4x + 4c_1)}}{2x}$$

$$y(x) \rightarrow \frac{-x^2 + \sqrt{x(x^3 - 4x + 4c_1)}}{2x}$$

6.6 problem 6

Internal problem ID [573]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$yx + y'x + y = 1$$

With initial conditions

$$[y(1) = 0]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 18

```
dsolve([x*y(x)+x*diff(y(x),x) = 1-y(x),y(1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{1 - e^{1-x}}{x}$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 20

```
DSolve[{x*y[x]+x*y'[x] == 1-y[x],y[1]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1 - e^{1-x}}{x}$$

6.7 problem 7

Internal problem ID [574]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [**_separable**]

$$y' - \frac{4x^3 + 1}{y(2 + 3y)} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 660

```
dsolve(diff(y(x),x) = (4*x^3+1)/(y(x)*(2+3*y(x))),y(x), singsol=all)
```

$$y(x) = \frac{\left(-8 + 108x^4 + 108c_1 + 108x + 12\sqrt{81x^8 + 162x^4c_1 + 162x^5 - 12x^4 + 81c_1^2 + 162c_1x + 81x^2 - 12c_1 - 12x}\right)}{6} + \frac{3\left(-8 + 108x^4 + 108c_1 + 108x + 12\sqrt{81x^8 + 162x^4c_1 + 162x^5 - 12x^4 + 81c_1^2 + 162c_1x + 81x^2 - 12c_1 - 12x}\right)}{2} - \frac{1}{3}$$

$$y(x) = \frac{\left(-8 + 108x^4 + 108c_1 + 108x + 12\sqrt{81x^8 + 162x^4c_1 + 162x^5 - 12x^4 + 81c_1^2 + 162c_1x + 81x^2 - 12c_1 - 12x}\right)}{12} - \frac{3\left(-8 + 108x^4 + 108c_1 + 108x + 12\sqrt{81x^8 + 162x^4c_1 + 162x^5 - 12x^4 + 81c_1^2 + 162c_1x + 81x^2 - 12c_1 - 12x}\right)}{1} - \frac{1}{3} + i\sqrt{3} \left(\frac{\left(-8 + 108x^4 + 108c_1 + 108x + 12\sqrt{81x^8 + 162x^4c_1 + 162x^5 - 12x^4 + 81c_1^2 + 162c_1x + 81x^2 - 12c_1 - 12x}\right)^{\frac{1}{3}}}{6} - \frac{3\left(-8 + 108x^4 + 108c_1 + 108x + 12\sqrt{81x^8 + 162x^4c_1 + 162x^5 - 12x^4 + 81c_1^2 + 162c_1x + 81x^2 - 12c_1 - 12x}\right)^{\frac{1}{3}}}{2} \right)$$

$$y(x) = \frac{\left(-8 + 108x^4 + 108c_1 + 108x + 12\sqrt{81x^8 + 162x^4c_1 + 162x^5 - 12x^4 + 81c_1^2 + 162c_1x + 81x^2 - 12c_1 - 12x}\right)}{12} - \frac{3\left(-8 + 108x^4 + 108c_1 + 108x + 12\sqrt{81x^8 + 162x^4c_1 + 162x^5 - 12x^4 + 81c_1^2 + 162c_1x + 81x^2 - 12c_1 - 12x}\right)}{1} - \frac{1}{3} + i\sqrt{3} \left(\frac{\left(-8 + 108x^4 + 108c_1 + 108x + 12\sqrt{81x^8 + 162x^4c_1 + 162x^5 - 12x^4 + 81c_1^2 + 162c_1x + 81x^2 - 12c_1 - 12x}\right)^{\frac{1}{3}}}{6} - \frac{3\left(-8 + 108x^4 + 108c_1 + 108x + 12\sqrt{81x^8 + 162x^4c_1 + 162x^5 - 12x^4 + 81c_1^2 + 162c_1x + 81x^2 - 12c_1 - 12x}\right)^{\frac{1}{3}}}{2} \right) + \frac{3\left(-8 + 108x^4 + 108c_1 + 108x + 12\sqrt{81x^8 + 162x^4c_1 + 162x^5 - 12x^4 + 81c_1^2 + 162c_1x + 81x^2 - 12c_1 - 12x}\right)}{2}$$

✓ Solution by Mathematica

Time used: 4.502 (sec). Leaf size: 356

`DSolve[y'[x]== (4*x^3+1)/(y[x]*(2+3*y[x])),y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{1}{6} \left(2^{2/3} \sqrt[3]{27x^4 + \sqrt{-4 + (27x^4 + 27x - 2 + 27c_1)^2} + 27x - 2 + 27c_1} \right. \\ \left. + \frac{2^3 \sqrt{2}}{\sqrt[3]{27x^4 + \sqrt{-4 + (27x^4 + 27x - 2 + 27c_1)^2} + 27x - 2 + 27c_1}} - 2 \right)$$

$$y(x) \rightarrow \frac{1}{12} \left(i 2^{2/3} (\sqrt{3} + i) \sqrt[3]{27x^4 + \sqrt{-4 + (27x^4 + 27x - 2 + 27c_1)^2} + 27x - 2 + 27c_1} \right. \\ \left. - \frac{2^3 \sqrt{2} (1 + i \sqrt{3})}{\sqrt[3]{27x^4 + \sqrt{-4 + (27x^4 + 27x - 2 + 27c_1)^2} + 27x - 2 + 27c_1}} - 4 \right)$$

$$y(x) \rightarrow \frac{1}{12} \left(-2^{2/3} (1 + i \sqrt{3}) \sqrt[3]{27x^4 + \sqrt{-4 + (27x^4 + 27x - 2 + 27c_1)^2} + 27x - 2 + 27c_1} \right. \\ \left. + \frac{2i \sqrt[3]{2} (\sqrt{3} + i)}{\sqrt[3]{27x^4 + \sqrt{-4 + (27x^4 + 27x - 2 + 27c_1)^2} + 27x - 2 + 27c_1}} - 4 \right)$$

6.8 problem 8

Internal problem ID [575]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$2y + y'x = \frac{\sin(x)}{x}$$

With initial conditions

$$[y(2) = 1]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve([2*y(x)+x*diff(y(x),x) = sin(x)/x,y(2) = 1],y(x), singsol=all)
```

$$y(x) = \frac{-\cos(x) + 4 + \cos(2)}{x^2}$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 17

```
DSolve[{2*y[x]+x*y'[x] == Sin[x]/x,y[2]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-\cos(x) + 4 + \cos(2)}{x^2}$$

6.9 problem 9

Internal problem ID [576]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, [_1st_order, ‘_with_symmetry_[F(x),G(x)]’], [_Abel]`

$$y' - \frac{-1 - 2yx}{x^2 + 2y} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
dsolve(diff(y(x),x) = (-1-2*x*y(x))/(x^2+2*y(x)),y(x), singsol=all)
```

$$y(x) = -\frac{x^2}{2} - \frac{\sqrt{x^4 - 4c_1 - 4x}}{2}$$

$$y(x) = -\frac{x^2}{2} + \frac{\sqrt{x^4 - 4c_1 - 4x}}{2}$$

✓ Solution by Mathematica

Time used: 0.15 (sec). Leaf size: 61

```
DSolve[y'[x]== (-1-2*x*y[x])/(x^2+2*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(-x^2 - \sqrt{x^4 - 4x + 4c_1} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left(-x^2 + \sqrt{x^4 - 4x + 4c_1} \right)$$

6.10 problem 10

Internal problem ID [577]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$\frac{yy'}{y-2} = -\frac{-x^2+x+1}{x^2}$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 26

```
dsolve((-x^2+x+1)/x^2+y(x)*diff(y(x),x)/(-2+y(x)) = 0,y(x), singsol=all)
```

$$y(x) = 2 \operatorname{LambertW}\left(\frac{c_1 e^{\frac{x}{2}-1+\frac{1}{2x}}}{2\sqrt{x}}\right) + 2$$

✓ Solution by Mathematica

Time used: 60.036 (sec). Leaf size: 63

```
DSolve[(-x^2+x+1)/x^2+y[x]*y'[x]/(-2+y[x]) == 0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 \left(1 + W\left(-\frac{1}{2} \sqrt{\frac{e^{x+\frac{1}{x}-2+c_1}}{x}}\right) \right)$$

$$y(x) \rightarrow 2 \left(1 + W\left(\frac{1}{2} \sqrt{\frac{e^{x+\frac{1}{x}-2+c_1}}{x}}\right) \right)$$

6.11 problem 11

Internal problem ID [578]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [exact]

$$y + (e^y + x)y' = -x^2$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 38

```
dsolve(x^2+y(x)+(exp(y(x))+x)*diff(y(x),x) = 0,y(x), singsol=all)
```

$$y(x) = -\text{LambertW}\left(\frac{e^{-\frac{x^2}{3}} e^{-\frac{c_1}{x}}}{x}\right) - \frac{x^3 + 3c_1}{3x}$$

✓ Solution by Mathematica

Time used: 3.877 (sec). Leaf size: 42

```
DSolve[x^2+y[x]+(Exp[y[x]]+x)*y'[x] == 0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -W\left(\frac{e^{-\frac{x^2}{3} + \frac{c_1}{x}}}{x}\right) - \frac{x^2}{3} + \frac{c_1}{x}$$

6.12 problem 12

Internal problem ID [579]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + y = \frac{1}{1 + e^x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(y(x)+diff(y(x),x) = 1/(1+exp(x)),y(x), singsol=all)
```

$$y(x) = (\ln(1 + e^x) + c_1) e^{-x}$$

✓ Solution by Mathematica

Time used: 0.067 (sec). Leaf size: 20

```
DSolve[y[x]+y'[x] == 1/(1+Exp[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(\log(e^x + 1) + c_1)$$

6.13 problem 13

Internal problem ID [580]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - y^2 - 2xy^2 = 1 + 2x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve(diff(y(x),x) = 1+2*x+y(x)^2+2*x*y(x)^2,y(x), singsol=all)
```

$$y(x) = \tan(x^2 + c_1 + x)$$

✓ Solution by Mathematica

Time used: 0.178 (sec). Leaf size: 13

```
DSolve[y'[x] == 1+2*x+y[x]^2+2*x*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan(x^2 + x + c_1)$$

6.14 problem 14

Internal problem ID [581]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 14.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _exact, _rational, [_Abel, '2nd ty`

$$y + (x + 2y)y' = -x$$

With initial conditions

$$[y(2) = 3]$$

✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 19

```
dsolve([x+y(x)+(x+2*y(x))*diff(y(x),x) = 0,y(2) = 3],y(x), singsol=all)
```

$$y(x) = -\frac{x}{2} + \frac{\sqrt{-x^2 + 68}}{2}$$

✓ Solution by Mathematica

Time used: 0.458 (sec). Leaf size: 24

```
DSolve[{x+y[x]+(x+2*y[x])*y'[x] == 0,y[2]==3},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(\sqrt{68 - x^2} - x)$$

6.15 problem 15

Internal problem ID [582]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(1 + e^x)y' - y + e^xy = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve((1+exp(x))*diff(y(x),x) = y(x)-exp(x)*y(x),y(x), singsol=all)
```

$$y(x) = \frac{c_1 e^x}{(1 + e^x)^2}$$

✓ Solution by Mathematica

Time used: 0.077 (sec). Leaf size: 23

```
DSolve[(1+Exp[x])*y'[x]== y[x]-Exp[x]*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1 e^x}{(e^x + 1)^2}$$

$$y(x) \rightarrow 0$$

6.16 problem 16

Internal problem ID [583]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [NONE]

$$y' - \frac{-e^{2y} \cos(x) + \cos(y) e^{-x}}{2e^{2y} \sin(x) - \sin(y) e^{-x}} = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 21

```
dsolve(diff(y(x), x) = (-exp(2*y(x))*cos(x)+cos(y(x))/exp(x))/(2*exp(2*y(x))*sin(x)-sin(y(x))))
```

$$e^{2y(x)} \sin(x) + \cos(y(x)) e^{-x} + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.473 (sec). Leaf size: 25

```
DSolve[y' [x] == (-Exp[2*y[x]]*Cos[x]+Cos[y[x]]/Exp[x])/(2*Exp[2*y[x]]*Sin[x]-Sin[y[x]]/Exp[x])
```

$$\text{Solve}[e^{2y(x)} \sin(x) + e^{-x} \cos(y(x)) = c_1, y(x)]$$

6.17 problem 17

Internal problem ID [584]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 17.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' - 3y = e^{2x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x) = exp(2*x)+3*y(x),y(x), singsol=all)
```

$$y(x) = (-e^{-x} + c_1) e^{3x}$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 19

```
DSolve[y'[x]== Exp[2*x]+3*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(-1 + c_1 e^x)$$

6.18 problem 18

Internal problem ID [585]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 18.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$2y + y' = e^{-x^2-2x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(2*y(x)+diff(y(x),x) = exp(-x^2-2*x),y(x), singsol=all)
```

$$y(x) = \left(\frac{\sqrt{\pi} \operatorname{erf}(x)}{2} + c_1 \right) e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.09 (sec). Leaf size: 27

```
DSolve[2*y[x]+y'[x] == Exp[-x^2-2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}e^{-2x}(\sqrt{\pi}\operatorname{erf}(x) + 2c_1)$$

6.19 problem 19

Internal problem ID [586]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational]`

$$y' - \frac{3x^2 - 2y - y^3}{2x + 3xy^2} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 507

`dsolve(diff(y(x),x) = (3*x^2-2*y(x)-y(x)^3)/(2*x+3*x*y(x)^2),y(x), singsol=all)`

$$\begin{aligned}
 y(x) &= \frac{\left(\left(108x^3 + 12\sqrt{3} \sqrt{27x^6 - 54c_1x^3 + 27c_1^2 + 32x^2} - 108c_1\right) x^2\right)^{\frac{1}{3}}}{6x} \\
 &\quad - \frac{\left(\left(108x^3 + 12\sqrt{3} \sqrt{27x^6 - 54c_1x^3 + 27c_1^2 + 32x^2} - 108c_1\right) x^2\right)^{\frac{1}{3}}}{4x} \\
 y(x) &= -\frac{\left(\left(108x^3 + 12\sqrt{3} \sqrt{27x^6 - 54c_1x^3 + 27c_1^2 + 32x^2} - 108c_1\right) x^2\right)^{\frac{1}{3}}}{2x} \\
 &\quad + \frac{\left(\left(108x^3 + 12\sqrt{3} \sqrt{27x^6 - 54c_1x^3 + 27c_1^2 + 32x^2} - 108c_1\right) x^2\right)^{\frac{1}{3}}}{12x} \\
 &\quad - \frac{i\sqrt{3} \left(\frac{\left(\left(108x^3 + 12\sqrt{3} \sqrt{27x^6 - 54c_1x^3 + 27c_1^2 + 32x^2} - 108c_1\right) x^2\right)^{\frac{1}{3}}}{6x} + \frac{4x}{\left(\left(108x^3 + 12\sqrt{3} \sqrt{27x^6 - 54c_1x^3 + 27c_1^2 + 32x^2} - 108c_1\right) x^2\right)^{\frac{1}{3}}} \right)}{2} \\
 y(x) &= -\frac{\left(\left(108x^3 + 12\sqrt{3} \sqrt{27x^6 - 54c_1x^3 + 27c_1^2 + 32x^2} - 108c_1\right) x^2\right)^{\frac{1}{3}}}{2x} \\
 &\quad + \frac{\left(\left(108x^3 + 12\sqrt{3} \sqrt{27x^6 - 54c_1x^3 + 27c_1^2 + 32x^2} - 108c_1\right) x^2\right)^{\frac{1}{3}}}{12x} \\
 &\quad + \frac{i\sqrt{3} \left(\frac{\left(\left(108x^3 + 12\sqrt{3} \sqrt{27x^6 - 54c_1x^3 + 27c_1^2 + 32x^2} - 108c_1\right) x^2\right)^{\frac{1}{3}}}{6x} + \frac{4x}{\left(\left(108x^3 + 12\sqrt{3} \sqrt{27x^6 - 54c_1x^3 + 27c_1^2 + 32x^2} - 108c_1\right) x^2\right)^{\frac{1}{3}}} \right)}{2}
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 32.075 (sec). Leaf size: 358

`DSolve[y'[x] == (3*x^2-2*y[x]-y[x]^3)/(2*x+3*x*y[x]^2),y[x],x,IncludeSingularSolutions -> Tr`

$$y(x) \rightarrow \frac{\sqrt[3]{27x^5 + 27c_1x^2 + \sqrt{864x^6 + 729x^4(x^3 + c_1)^2}}}{\frac{3\sqrt[3]{2}x}{2\sqrt[3]{2}x}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{2}(1 + i\sqrt{3})x}{\sqrt[3]{27x^5 + 27c_1x^2 + \sqrt{864x^6 + 729x^4(x^3 + c_1)^2}}}$$

$$- \frac{(1 - i\sqrt{3})\sqrt[3]{27x^5 + 27c_1x^2 + \sqrt{864x^6 + 729x^4(x^3 + c_1)^2}}}{6\sqrt[3]{2}x}$$

$$y(x) \rightarrow \frac{\sqrt[3]{2}(1 - i\sqrt{3})x}{\sqrt[3]{27x^5 + 27c_1x^2 + \sqrt{864x^6 + 729x^4(x^3 + c_1)^2}}}$$

$$- \frac{(1 + i\sqrt{3})\sqrt[3]{27x^5 + 27c_1x^2 + \sqrt{864x^6 + 729x^4(x^3 + c_1)^2}}}{6\sqrt[3]{2}x}$$

6.20 problem 20

Internal problem ID [587]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 20.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - e^{x+y} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x) = exp(x+y(x)),y(x), singsol=all)
```

$$y(x) = \ln\left(-\frac{1}{e^x + c_1}\right)$$

✓ Solution by Mathematica

Time used: 0.752 (sec). Leaf size: 18

```
DSolve[y'[x] == Exp[x+y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\log(-e^x - c_1)$$

6.21 problem 21

Internal problem ID [588]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 21.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational]`

$$\frac{-4 + 6yx + 2y^2}{3x^2 + 4yx + 3y^2} + y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 561

`dsolve((-4+6*x*y(x)+2*y(x)^2)/(3*x^2+4*x*y(x)+3*y(x)^2)+diff(y(x),x)=0,y(x), singsol=all)`

$y(x)$

$$= \frac{\left(152x^3 - 108c_1 + 432x + 12\sqrt{216x^6 - 228c_1x^3 + 912x^4 + 81c_1^2 - 648c_1x + 1296x^2}\right)^{\frac{1}{3}}}{\frac{6}{10x^2}}$$

$$- \frac{3 \left(152x^3 - 108c_1 + 432x + 12\sqrt{216x^6 - 228c_1x^3 + 912x^4 + 81c_1^2 - 648c_1x + 1296x^2}\right)^{\frac{1}{3}}}{\frac{2x}{3}}$$

$y(x) =$

$$\frac{\left(152x^3 - 108c_1 + 432x + 12\sqrt{216x^6 - 228c_1x^3 + 912x^4 + 81c_1^2 - 648c_1x + 1296x^2}\right)^{\frac{1}{3}}}{\frac{12}{5x^2}}$$

$$+ \frac{3 \left(152x^3 - 108c_1 + 432x + 12\sqrt{216x^6 - 228c_1x^3 + 912x^4 + 81c_1^2 - 648c_1x + 1296x^2}\right)^{\frac{1}{3}}}{\frac{2x}{3}}$$

$$- \frac{i\sqrt{3} \left(\frac{\left(152x^3 - 108c_1 + 432x + 12\sqrt{216x^6 - 228c_1x^3 + 912x^4 + 81c_1^2 - 648c_1x + 1296x^2}\right)^{\frac{1}{3}}}{6} + \frac{10x^2}{3 \left(152x^3 - 108c_1 + 432x + 12\sqrt{216x^6 - 228c_1x^3 + 912x^4 + 81c_1^2 - 648c_1x + 1296x^2}\right)^{\frac{1}{3}}} \right)}{2}$$

$y(x) =$

$$\frac{\left(152x^3 - 108c_1 + 432x + 12\sqrt{216x^6 - 228c_1x^3 + 912x^4 + 81c_1^2 - 648c_1x + 1296x^2}\right)^{\frac{1}{3}}}{\frac{12}{5x^2}}$$

$$+ \frac{3 \left(152x^3 - 108c_1 + 432x + 12\sqrt{216x^6 - 228c_1x^3 + 912x^4 + 81c_1^2 - 648c_1x + 1296x^2}\right)^{\frac{1}{3}}}{\frac{2x}{3}}$$

$$+ \frac{i\sqrt{3} \left(\frac{\left(152x^3 - 108c_1 + 432x + 12\sqrt{216x^6 - 228c_1x^3 + 912x^4 + 81c_1^2 - 648c_1x + 1296x^2}\right)^{\frac{1}{3}}}{6} + \frac{10x^2}{3 \left(152x^3 - 108c_1 + 432x + 12\sqrt{216x^6 - 228c_1x^3 + 912x^4 + 81c_1^2 - 648c_1x + 1296x^2}\right)^{\frac{1}{3}}} \right)}{2}$$

✓ Solution by Mathematica

Time used: 4.748 (sec). Leaf size: 383

`DSolve[(-4+6*x*y[x]+2*y[x]^2)/(3*x^2+4*x*y[x]+3*y[x]^2)+y'[x]==0,y[x],x,IncludeSingularSolut`

$$y(x) \rightarrow \frac{1}{6} \left(2^{2/3} \sqrt[3]{38x^3 + \sqrt{500x^6 + (38x^3 + 108x + 27c_1)^2} + 108x + 27c_1} - \frac{10\sqrt[3]{2}x^2}{\sqrt[3]{38x^3 + \sqrt{500x^6 + (38x^3 + 108x + 27c_1)^2} + 108x + 27c_1}} - 4x \right)$$

$$y(x) \rightarrow \frac{1}{12} \left(i2^{2/3}(\sqrt{3} + i) \sqrt[3]{38x^3 + \sqrt{500x^6 + (38x^3 + 108x + 27c_1)^2} + 108x + 27c_1} + \frac{10\sqrt[3]{2}(1 + i\sqrt{3})x^2}{\sqrt[3]{38x^3 + \sqrt{500x^6 + (38x^3 + 108x + 27c_1)^2} + 108x + 27c_1}} - 8x \right)$$

$$y(x) \rightarrow \frac{1}{12} \left(-2^{2/3}(1 + i\sqrt{3}) \sqrt[3]{38x^3 + \sqrt{500x^6 + (38x^3 + 108x + 27c_1)^2} + 108x + 27c_1} + \frac{10\sqrt[3]{2}(1 - i\sqrt{3})x^2}{\sqrt[3]{38x^3 + \sqrt{500x^6 + (38x^3 + 108x + 27c_1)^2} + 108x + 27c_1}} - 8x \right)$$

6.22 problem 22

Internal problem ID [589]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 22.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{x^2 - 1}{1 + y^2} = 0$$

With initial conditions

$$[y(-1) = 1]$$

✓ Solution by Maple

Time used: 0.172 (sec). Leaf size: 87

```
dsolve([diff(y(x),x) = (x^2-1)/(1+y(x)^2),y(-1) = 1],y(x), singsol=all)
```

$$y(x) = \frac{(8 + 4x^3 - 12x + 4\sqrt{x^6 - 6x^4 + 4x^3 + 9x^2 - 12x + 8})^{\frac{2}{3}} - 4}{2(8 + 4x^3 - 12x + 4\sqrt{x^6 - 6x^4 + 4x^3 + 9x^2 - 12x + 8})^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 2.95 (sec). Leaf size: 97

```
DSolve[{y'[x]== (x^2-1)/(1+y[x]^2),y[-1]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt[3]{2}(x^3 + \sqrt{x^6 - 6x^4 + 4x^3 + 9x^2 - 12x + 8} - 3x + 2)^{2/3} - 2}{2^{2/3} \sqrt[3]{x^3 + \sqrt{x^6 - 6x^4 + 4x^3 + 9x^2 - 12x + 8} - 3x + 2}}$$

6.23 problem 23

Internal problem ID [590]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 23.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$(t + 1)y + ty' = e^{2t}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 20

```
dsolve((1+t)*y(t)+t*diff(y(t),t) = exp(2*t),y(t), singsol=all)
```

$$y(t) = \frac{\left(\frac{e^{3t}}{3} + c_1\right) e^{-t}}{t}$$

✓ Solution by Mathematica

Time used: 0.053 (sec). Leaf size: 27

```
DSolve[(1+t)*y[t]+t*y'[t] == Exp[2*t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{e^{2t} + 3c_1 e^{-t}}{3t}$$

6.24 problem 24

Internal problem ID [591]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 24.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$2 \cos(x) \sin(x) \sin(y) + \cos(y) \sin(x)^2 y' = 0$$

✓ Solution by Maple

Time used: 0.109 (sec). Leaf size: 18

```
dsolve(2*cos(x)*sin(x)*sin(y(x))+cos(y(x))*sin(x)^2*diff(y(x),x) = 0,y(x), singsol=all)
```

$$y(x) = -\arcsin\left(\frac{2c_1}{-1 + \cos(2x)}\right)$$

✓ Solution by Mathematica

Time used: 5.176 (sec). Leaf size: 21

```
DSolve[2*Cos[x]*Sin[x]*Sin[y[x]]+Cos[y[x]]*Sin[x]^2*y'[x] == 0,y[x],x,IncludeSingularSolutio
```

$$y(x) \rightarrow \arcsin\left(\frac{1}{2}c_1 \csc^2(x)\right)$$

$$y(x) \rightarrow 0$$

6.25 problem 25

Internal problem ID [592]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 25.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_exact, _rational]`

$$\frac{2x}{y} - \frac{y}{x^2 + y^2} + \left(-\frac{x^2}{y^2} + \frac{x}{x^2 + y^2} \right) y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve(2*x/y(x)-y(x)/(x^2+y(x)^2)+(-x^2/y(x)^2+x/(x^2+y(x)^2))*diff(y(x),x) = 0,y(x), singso
```

$$y(x) = \frac{x}{\tan(\text{RootOf}(-_Z + x \tan(_Z) + c_1))}$$

✓ Solution by Mathematica

Time used: 0.255 (sec). Leaf size: 23

```
DSolve[2*x/y[x]-y[x]/(x^2+y[x]^2)+(-x^2/y[x]^2+x/(x^2+y[x]^2))*y'[x] == 0,y[x],x,IncludeSing
```

$$\text{Solve} \left[\arctan \left(\frac{x}{y(x)} \right) - \frac{x^2}{y(x)} = c_1, y(x) \right]$$

6.26 problem 26

Internal problem ID [593]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 26.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A'], _dAlembert]`

$$y'x - e^{\frac{y}{x}}x - y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(x*diff(y(x),x) = exp(y(x)/x)*x+y(x),y(x), singsol=all)
```

$$y(x) = \ln\left(-\frac{1}{\ln(x) + c_1}\right)x$$

✓ Solution by Mathematica

Time used: 0.316 (sec). Leaf size: 18

```
DSolve[x*y'[x] == Exp[y[x]/x]*x+y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x \log(-\log(x) - c_1)$$

6.27 problem 27

Internal problem ID [594]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 27.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, [_1st_order, ‘_with_symmetry_[F(x)*G(y),0]’]]`

$$y' - \frac{x}{x^2 + y + y^3} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 40

```
dsolve(diff(y(x),x) = x/(x^2+y(x)+y(x)^3),y(x), singsol=all)
```

$$c_1 - e^{-2y(x)}x^2 - \frac{(4y(x)^3 + 6y(x)^2 + 10y(x) + 5)e^{-2y(x)}}{4} = 0$$

✓ Solution by Mathematica

Time used: 0.173 (sec). Leaf size: 48

```
DSolve[y'[x] == x/(x^2+y[x]+y[x]^3),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[-\frac{1}{2}x^2e^{-2y(x)} - \frac{1}{8}e^{-2y(x)}(4y(x)^3 + 6y(x)^2 + 10y(x) + 5) = c_1, y(x)\right]$$

6.28 problem 28

Internal problem ID [595]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 28.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$2y + ty' = -3t$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(3*t+2*y(t) = -t*diff(y(t),t),y(t), singsol=all)
```

$$y(t) = -t + \frac{c_1}{t^2}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 15

```
DSolve[3*t+2*y[t] == -t*y'[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -t + \frac{c_1}{t^2}$$

6.29 problem 29

Internal problem ID [596]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 29.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{x+y}{x-y} = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 24

```
dsolve(diff(y(x),x) = (x+y(x))/(x-y(x)),y(x), singsol=all)
```

$$y(x) = \tan \left(\text{RootOf} \left(-2_Z + \ln \left(\frac{1}{\cos(_Z)^2} \right) + 2 \ln(x) + 2c_1 \right) \right) x$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 36

```
DSolve[y'[x] == (x+y[x])/(x-y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{1}{2} \log \left(\frac{y(x)^2}{x^2} + 1 \right) - \arctan \left(\frac{y(x)}{x} \right) = -\log(x) + c_1, y(x) \right]$$

6.30 problem 30

Internal problem ID [597]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 30.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$2yx + 3y^2 - (x^2 + 2yx) y' = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```
dsolve(2*x*y(x)+3*y(x)^2-(x^2+2*x*y(x))*diff(y(x),x) = 0,y(x), singsol=all)
```

$$y(x) = \left(-\frac{1}{2} - \frac{\sqrt{4c_1x + 1}}{2} \right) x$$

$$y(x) = \left(-\frac{1}{2} + \frac{\sqrt{4c_1x + 1}}{2} \right) x$$

✓ Solution by Mathematica

Time used: 0.408 (sec). Leaf size: 61

```
DSolve[2*x*y[x]+3*y[x]^2-(x^2+2*x*y[x])*y'[x] == 0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2}x \left(1 + \sqrt{1 + 4e^{c_1x}} \right)$$

$$y(x) \rightarrow \frac{1}{2}x \left(-1 + \sqrt{1 + 4e^{c_1x}} \right)$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -x$$

6.31 problem 31

Internal problem ID [598]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Miscellaneous problems, end of chapter 2. Page 133

Problem number: 31.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{-3x^2y - y^2}{2x^3 + 3yx} = 0$$

With initial conditions

$$[y(1) = -2]$$

✓ Solution by Maple

Time used: 1.281 (sec). Leaf size: 111

```
dsolve([diff(y(x),x) = (-3*x^2*y(x)-y(x)^2)/(2*x^3+3*x*y(x)),y(1) = -2],y(x), singsol=all)
```

$$y(x) = \frac{(i\sqrt{3} - 1) \left(-(x^7 - 6\sqrt{3}\sqrt{x^7 + 27} + 54) x^2 \right)^{\frac{2}{3}} - x^3 \left(ix^3\sqrt{3} + x^3 + 2 \left(-(x^7 - 6\sqrt{3}\sqrt{x^7 + 27} + 54) x^2 \right)^{\frac{1}{3}} \right)}{6 \left(-(x^7 - 6\sqrt{3}\sqrt{x^7 + 27} + 54) x^2 \right)^{\frac{1}{3}} x}$$

✓ Solution by Mathematica

Time used: 40.923 (sec). Leaf size: 136

```
DSolve[{y'[x]== (-3*x^2*y[x]-y[x]^2)/(2*x^3+3*x*y[x]),y[1]==-2},y[x],x,IncludeSingularSoluti
```

$$y(x) \rightarrow \frac{i \left((\sqrt{3} + i) x^3 - (\sqrt{3} - i) x^3 + (\sqrt{3} + i) \sqrt[3]{-x^9 - 54x^2 + 6\sqrt{3}\sqrt{x^4(x^7 + 27)}} - \frac{(\sqrt{3} - i)x^6}{\sqrt[3]{-x^9 - 54x^2 + 6\sqrt{3}\sqrt{x^4(x^7 + 27)}}} \right)}{6x}$$

7 Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

7.1	problem 1	186
7.2	problem 2	187
7.3	problem 3	188
7.4	problem 4	189
7.5	problem 5	190
7.6	problem 6	191
7.7	problem 7	192
7.8	problem 8	193
7.9	problem 9	194
7.10	problem 10	195
7.11	problem 11	196
7.12	problem 12	197
7.13	problem 13	198
7.14	problem 14	199
7.15	problem 15	200
7.16	problem 16	201
7.17	problem 19	202
7.18	problem 20	203
7.19	problem 21	204
7.20	problem 22	205
7.21	problem 23	206
7.22	problem 24	207
7.23	problem 25	208
7.24	problem 26	209

7.1 problem 1

Internal problem ID [599]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2y' - 3y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x$2) +2*diff(y(x),x)-3*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-3x} + e^x c_2$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 20

```
DSolve[y''[x]+2*y'[x]-3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-3x} + c_2 e^x$$

7.2 problem 2

Internal problem ID [600]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 3y' + 2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2) +3*diff(y(x),x)+2*y(x) = 0,y(x), singsol=all)
```

$$y(x) = e^{-x}c_1 + c_2e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 20

```
DSolve[y''[x]+3*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(c_2e^x + c_1)$$

7.3 problem 3

Internal problem ID [601]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$6y'' - y' - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(6*diff(y(x),x$2) -diff(y(x),x)-y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{x}{3}} + c_2 e^{\frac{x}{2}}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 26

```
DSolve[6*y''[x]-y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/3} (c_2 e^{5x/6} + c_1)$$

7.4 problem 4

Internal problem ID [602]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$2y'' - 3y' + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(2*diff(y(x),x$2) -3*diff(y(x),x)+y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{x}{2}} + e^x c_2$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 35

```
DSolve[y''[x]-3*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\frac{1}{2}(\sqrt{5}-3)x} (c_2 e^{\sqrt{5}x} + c_1)$$

7.5 problem 5

Internal problem ID [603]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 5y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve(diff(y(x),x$2) +5*diff(y(x),x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 + c_2 e^{-5x}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 19

```
DSolve[y''[x]+5*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \frac{1}{5}c_1 e^{-5x}$$

7.6 problem 6

Internal problem ID [604]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4y'' - 9y = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve(4*diff(y(x),x$2) -9*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{3x}{2}} + c_2 e^{\frac{3x}{2}}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 24

```
DSolve[4*y''[x]-9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x/2}(c_1 e^{3x} + c_2)$$

7.7 problem 7

Internal problem ID [605]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 9y' + 9y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2) -9*diff(y(x),x)+9*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{3(3+\sqrt{5})x}{2}} + c_2 e^{-\frac{3(\sqrt{5}-3)x}{2}}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 36

```
DSolve[y''[x]-9*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\frac{3}{2}(\sqrt{5}-3)x} (c_2 e^{3\sqrt{5}x} + c_1)$$

7.8 problem 8

Internal problem ID [606]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' - 2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2) -2*diff(y(x),x)-2*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{(1+\sqrt{3})x} + c_2 e^{-(\sqrt{3}-1)x}$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 34

```
DSolve[y''[x]-2*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x-\sqrt{3}x} \left(c_2 e^{2\sqrt{3}x} + c_1 \right)$$

7.9 problem 9

Internal problem ID [607]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y' - 2y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve([diff(y(x),x$2) +diff(y(x),x)-2*y(x) = 0,y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{(e^{3x} - 1)e^{-2x}}{3}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 21

```
DSolve[{y''[x]+y'[x]-2*y[x]==0,{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}e^{-2x}(e^{3x} - 1)$$

7.10 problem 10

Internal problem ID [608]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 4y' + 3y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = -1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2) +4*diff(y(x),x)+3*y(x) = 0,y(0) = 2, D(y)(0) = -1],y(x), singsol=all)
```

$$y(x) = \frac{5e^{-x}}{2} - \frac{e^{-3x}}{2}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 23

```
DSolve[{y''[x]+4*y'[x]+3*y[x]==0,{y[0]==2,y'[0]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}e^{-3x}(5e^{2x} - 1)$$

7.11 problem 11

Internal problem ID [609]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$6y'' - 5y' + y = 0$$

With initial conditions

$$[y(0) = 4, y'(0) = 0]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve([6*diff(y(x),x$2) -5*diff(y(x),x)+y(x) = 0,y(0) = 4, D(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = -8e^{\frac{x}{2}} + 12e^{\frac{x}{3}}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 48

```
DSolve[{6*y''[x]-5*y'[x]+2*y[x]==0,{y[0]==4,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{4}{23}e^{5x/12} \left(23 \cos\left(\frac{\sqrt{23}x}{12}\right) - 5\sqrt{23} \sin\left(\frac{\sqrt{23}x}{12}\right) \right)$$

7.12 problem 12

Internal problem ID [610]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 3y' = 0$$

With initial conditions

$$[y(0) = -2, y'(0) = 3]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 12

```
dsolve([diff(y(x),x$2) +3*diff(y(x),x) = 0,y(0) = -2, D(y)(0) = 3],y(x), singsol=all)
```

$$y(x) = -1 - e^{-3x}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 14

```
DSolve[{y'[x]+3*y'[x]==0,{y[0]==-2,y'[0]==3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -e^{-3x} - 1$$

7.13 problem 13

Internal problem ID [611]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 5y' + 3y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

✓ Solution by Maple

Time used: 0.422 (sec). Leaf size: 39

```
dsolve([diff(y(x),x$2) +5*diff(y(x),x)+3*y(x) = 0,y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{(5\sqrt{13} + 13) e^{\frac{(-5+\sqrt{13})x}{2}}}{26} + \frac{(-5\sqrt{13} + 13) e^{-\frac{(5+\sqrt{13})x}{2}}}{26}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 51

```
DSolve[{y'[x]+5*y'[x]+3*y[x]==0,{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{1}{26} e^{-\frac{1}{2}(5+\sqrt{13})x} \left((13 + 5\sqrt{13}) e^{\sqrt{13}x} + 13 - 5\sqrt{13} \right)$$

7.14 problem 14

Internal problem ID [612]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$2y'' + y' - 4y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 30

```
dsolve([2*diff(y(x),x$2) +diff(y(x),x)-4*y(x) = 0,y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = -\frac{2\left(-e^{\frac{(-1+\sqrt{33})x}{4}} + e^{\frac{(1+\sqrt{33})x}{4}}\right)\sqrt{33}}{33}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 40

```
DSolve[{2*y''[x]+y'[x]-4*y[x]==0,{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{2e^{-\frac{1}{4}(1+\sqrt{33})x}\left(e^{\frac{\sqrt{33}x}{2}} - 1\right)}{\sqrt{33}}$$

7.15 problem 15

Internal problem ID [613]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 8y' - 9y = 0$$

With initial conditions

$$[y(1) = 1, y'(1) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 19

```
dsolve([diff(y(x),x$2) +8*diff(y(x),x)-9*y(x) = 0,y(1) = 1, D(y)(1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{e^{9-9x}}{10} + \frac{9e^{x-1}}{10}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 26

```
DSolve[{y''[x]+8*y'[x]-9*y[x]==0,{y[1]==1,y'[1]==0}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{1}{10}e^{9-9x} + \frac{9e^{x-1}}{10}$$

7.16 problem 16

Internal problem ID [614]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4y'' - y = 0$$

With initial conditions

$$[y(-2) = 1, y'(-2) = -1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve([4*diff(y(x),x$2) -y(x) = 0,y(-2) = 1, D(y)(-2) = -1],y(x), singsol=all)
```

$$y(x) = -\frac{e^{1+\frac{x}{2}}}{2} + \frac{3e^{-1-\frac{x}{2}}}{2}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 25

```
DSolve[{4*y'[x]-y[x]==0,{y[-2]==1,y'[-2]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2}e^{-\frac{x}{2}-1}(e^{x+2} - 3)$$

7.17 problem 19

Internal problem ID [615]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y = 0$$

With initial conditions

$$\left[y(0) = \frac{5}{4}, y'(0) = -\frac{3}{4} \right]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([diff(y(x),x$2) -y(x) = 0,y(0) = 5/4, D(y)(0) = -3/4],y(x), singsol=all)
```

$$y(x) = e^{-x} + \frac{e^x}{4}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 18

```
DSolve[{y'[x]-y[x]==0,{y[0]==5/4,y'[0]==-3/4}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} + \frac{e^x}{4}$$

7.18 problem 20

Internal problem ID [616]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$2y'' - 3y' + y = 0$$

With initial conditions

$$\left[y(0) = 2, y'(0) = \frac{1}{2} \right]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve([2*diff(y(x),x$2) -3*diff(y(x),x)+y(x) = 0,y(0) = 2, D(y)(0) = 1/2],y(x), singsol=all
```

$$y(x) = 3e^{\frac{x}{2}} - e^x$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 20

```
DSolve[{2*y'[x]-3*y'[x]+y[x]==0,{y[0]==2,y'[0]==1/2}},y[x],x,IncludeSingularSolutions -> Tr
```

$$y(x) \rightarrow 3e^{x/2} - e^x$$

7.19 problem 21

Internal problem ID [617]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y' - 2y = 0$$

With initial conditions

$$[y(0) = \alpha, y'(0) = 2]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve([diff(y(x),x$2) -diff(y(x),x)-2*y(x) = 0,y(0) = alpha, D(y)(0) = 2],y(x), singsol=all
```

$$y(x) = \frac{(2\alpha - 2)e^{-x}}{3} + \frac{e^{2x}(\alpha + 2)}{3}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 29

```
DSolve[{y''[x]-y'[x]-2*y[x]==0,{y[0]==\[Alpha],y'[0]==2}},y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow \frac{1}{3}e^{-x}(2(\alpha - 1) + (\alpha + 2)e^{3x})$$

7.20 problem 22

Internal problem ID [618]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4y'' - y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = \beta]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 22

```
dsolve([4*diff(y(x),x$2) -y(x) = 0,y(0) = 2, D(y)(0) = beta],y(x), singsol=all)
```

$$y(x) = (1 + \beta)e^{\frac{x}{2}} - (\beta - 1)e^{-\frac{x}{2}}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 25

```
DSolve[{4*y'[x]-y[x]==0,{y[0]==2,y'[0]==[Beta]}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/2}(-\beta + (\beta + 1)e^x + 1)$$

7.21 problem 23

Internal problem ID [619]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - (2\alpha - 1)y' + \alpha(\alpha - 1)y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x$2) -(2*alpha-1)*diff(y(x),x)+alpha*(alpha-1)*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\alpha x} + c_2 e^{(\alpha-1)x}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 24

```
DSolve[y''[x]-(2*[Alpha]-1)*y'[x]+\[Alpha]*(\[Alpha]-1)*y[x]==0,y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow c_1 e^{(\alpha-1)x} + c_2 e^{\alpha x}$$

7.22 problem 24

Internal problem ID [620]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 24.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + (3 - \alpha)y' - 2(\alpha - 1)y = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 19

```
dsolve(diff(y(x),x$2) +(3-alpha)*diff(y(x),x)-2*(alpha-1)*y(x) = 0,y(x), singsol=all)
```

$$y(x) = e^{-2x}c_1 + c_2e^{(\alpha-1)x}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 24

```
DSolve[y''[x]+(3-\[Alpha])*y'[x]-2*(\[Alpha]-1)*y[x]==0,y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow e^{-2x}(c_1e^{\alpha x+x} + c_2)$$

7.23 problem 25

Internal problem ID [621]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 25.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$2y'' + 3y' - 2y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = -\beta]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve([2*diff(y(x),x$2) +3*diff(y(x),x)-2*y(x) = 0,y(0) = 1, D(y)(0) = -beta],y(x), singsol
```

$$y(x) = -\frac{\left(2e^{\frac{5x}{2}}\beta - 4e^{\frac{5x}{2}} - 2\beta - 1\right)e^{-2x}}{5}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 67

```
DSolve[{y'[x]+3*y'[x]-2*y[x]==0,{y[0]==1,y'[0]==-\[Beta]}],y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{1}{34}e^{-\frac{1}{2}(3+\sqrt{17})x} \left(2\sqrt{17}\beta + (-2\sqrt{17}\beta + 3\sqrt{17} + 17)e^{\sqrt{17}x} - 3\sqrt{17} + 17\right)$$

7.24 problem 26

Internal problem ID [622]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.1 Homogeneous Equations with Constant Coefficients, page 144

Problem number: 26.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 5y' + 6y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = \beta]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 23

```
dsolve([diff(y(x),x$2) +5*diff(y(x),x)+6*y(x) = 0,y(0) = 2, D(y)(0) = beta],y(x), singsol=all)
```

$$y(x) = e^{-2x}(6 + \beta) + (-\beta - 4)e^{-3x}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 23

```
DSolve[{y'[x]+5*y'[x]+6*y[x]==0,{y[0]==2,y'[0]==\[Beta]}},y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow e^{-3x}(-\beta + (\beta + 6)e^x - 4)$$

8 Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

8.1	problem 7	211
8.2	problem 8	212
8.3	problem 9	213
8.4	problem 10	214
8.5	problem 11	215
8.6	problem 12	216
8.7	problem 13	217
8.8	problem 14	218
8.9	problem 15	219
8.10	problem 16	220
8.11	problem 17	221
8.12	problem 18	222
8.13	problem 19	223
8.14	problem 20	224
8.15	problem 21	225
8.16	problem 22	226
8.17	problem 23	227
8.18	problem 24	228
8.19	problem 25	229
8.20	problem 26	230
8.21	problem 35	231
8.22	problem 36	232
8.23	problem 37	233
8.24	problem 38	234
8.25	problem 39	235
8.26	problem 40	236
8.27	problem 41	237
8.28	problem 42	238
8.29	problem 44	239
8.30	problem 46	240

8.1 problem 7

Internal problem ID [623]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' + 2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2) -2*diff(y(x),x)+2*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(x) e^x + c_2 \cos(x) e^x$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 20

```
DSolve[y''[x]-2*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(c_2 \cos(x) + c_1 \sin(x))$$

8.2 problem 8

Internal problem ID [624]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' + 6y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2) -2*diff(y(x),x)+6*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^x \sin(\sqrt{5}x) + c_2 e^x \cos(\sqrt{5}x)$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 32

```
DSolve[y''[x]-2*y'[x]+6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x \left(c_2 \cos(\sqrt{5}x) + c_1 \sin(\sqrt{5}x) \right)$$

8.3 problem 9

Internal problem ID [625]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2y' - 8y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2) +2*diff(y(x),x)-8*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-4x} + c_2 e^{2x}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 22

```
DSolve[y''[x]+2*y'[x]-8*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-4x} (c_2 e^{6x} + c_1)$$

8.4 problem 10

Internal problem ID [626]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2y' + 2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2) +2*diff(y(x),x)+2*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-x} \sin(x) + c_2 e^{-x} \cos(x)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 22

```
DSolve[y''[x]+2*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(c_2 \cos(x) + c_1 \sin(x))$$

8.5 problem 11

Internal problem ID [627]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 6y' + 13y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2) +6*diff(y(x),x)+13*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-3x} \sin(2x) + c_2 e^{-3x} \cos(2x)$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 26

```
DSolve[y''[x]+6*y'[x]+13*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x}(c_2 \cos(2x) + c_1 \sin(2x))$$

8.6 problem 12

Internal problem ID [628]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4y'' + 9y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(4*diff(y(x),x$2) +9*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 \sin\left(\frac{3x}{2}\right) + c_2 \cos\left(\frac{3x}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 20

```
DSolve[y''[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(3x) + c_2 \sin(3x)$$

8.7 problem 13

Internal problem ID [629]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2y' + \frac{5y}{4} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2) +2*diff(y(x),x)+125/100*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-x} \sin\left(\frac{x}{2}\right) + c_2 e^{-x} \cos\left(\frac{x}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 30

```
DSolve[y''[x]+2*y'[x]+125/100*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} \left(c_2 \cos\left(\frac{x}{2}\right) + c_1 \sin\left(\frac{x}{2}\right) \right)$$

8.8 problem 14

Internal problem ID [630]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$9y'' + 9y' - 4y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(9*diff(y(x),x$2) +9*diff(y(x),x)-4*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{4x}{3}} + c_2 e^{\frac{x}{3}}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 26

```
DSolve[9*y''[x]+9*y'[x]-4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-4x/3}(c_2 e^{5x/3} + c_1)$$

8.9 problem 15

Internal problem ID [631]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y' + \frac{5y}{4} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2) +diff(y(x),x)+125/100*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{x}{2}} \sin(x) + c_2 e^{-\frac{x}{2}} \cos(x)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 24

```
DSolve[y''[x]+y'[x]+125/100*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/2}(c_2 \cos(x) + c_1 \sin(x))$$

8.10 problem 16

Internal problem ID [632]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 4y' + \frac{25y}{4} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)+ 4*diff(y(x),x)+625/100*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-2x} \sin\left(\frac{3x}{2}\right) + c_2 e^{-2x} \cos\left(\frac{3x}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 30

```
DSolve[y''[x]+4*y'[x]+625/100*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} \left(c_2 \cos\left(\frac{3x}{2}\right) + c_1 \sin\left(\frac{3x}{2}\right) \right)$$

8.11 problem 17

Internal problem ID [633]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 17.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 4y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 10

```
dsolve([diff(y(x),x$2)+ 4*y(x) = 0,y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{\sin(2x)}{2}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 10

```
DSolve[{y'[x]+4*y[x]==0,{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sin(x) \cos(x)$$

8.12 problem 18

Internal problem ID [634]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 18.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 4y' + 5y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve([diff(y(x),x$2)+ 4*diff(y(x),x)+5*y(x) = 0,y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = e^{-2x}(2 \sin(x) + \cos(x))$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 18

```
DSolve[{y'[x]+4*y'[x]+5*y[x]==0,{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow e^{-2x}(2 \sin(x) + \cos(x))$$

8.13 problem 19

Internal problem ID [635]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' + 5y = 0$$

With initial conditions

$$\left[y\left(\frac{\pi}{2}\right) = 0, y'\left(\frac{\pi}{2}\right) = 2 \right]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 16

```
dsolve([diff(y(x),x$2)- 2*diff(y(x),x)+5*y(x) = 0,y(1/2*Pi) = 0, D(y)(1/2*Pi) = 2],y(x), sin
```

$$y(x) = -\sin(2x)e^{-\frac{\pi}{2}+x}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 20

```
DSolve[{y''[x]-2*y'[x]+5*y[x]==0,{y[Pi/2]==0,y'[Pi/2]==2}},y[x],x,IncludeSingularSolutions -
```

$$y(x) \rightarrow -e^{x-\frac{\pi}{2}} \sin(2x)$$

8.14 problem 20

Internal problem ID [636]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y = 0$$

With initial conditions

$$\left[y\left(\frac{\pi}{3}\right) = 2, y'\left(\frac{\pi}{3}\right) = -4 \right]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 23

```
dsolve([diff(y(x),x$2)+y(x) = 0,y(1/3*Pi) = 2, D(y)(1/3*Pi) = -4],y(x), singsol=all)
```

$$y(x) = (\sin(x) + 2 \cos(x)) \sqrt{3} + \cos(x) - 2 \sin(x)$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 28

```
DSolve[{y'[x]+y[x]==0,{y[Pi/3]==2,y'[Pi/3]==-4}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (\sqrt{3} - 2) \sin(x) + (1 + 2\sqrt{3}) \cos(x)$$

8.15 problem 21

Internal problem ID [637]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y' + \frac{5y}{4} = 0$$

With initial conditions

$$[y(0) = 3, y'(0) = 1]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 19

```
dsolve([diff(y(x),x$2)+ diff(y(x),x)+125/100*y(x) = 0,y(0) = 3, D(y)(0) = 1],y(x), singsol=a
```

$$y(x) = \frac{e^{-\frac{x}{2}}(5 \sin(x) + 6 \cos(x))}{2}$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 25

```
DSolve[{y'[x]+y'[x]+125/100*y[x]==0,{y[0]==3,y'[0]==1}},y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow \frac{1}{2}e^{-x/2}(5 \sin(x) + 6 \cos(x))$$

8.16 problem 22

Internal problem ID [638]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2y' + 2y = 0$$

With initial conditions

$$\left[y\left(\frac{\pi}{4}\right) = 2, y'\left(\frac{\pi}{4}\right) = -2 \right]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 21

```
dsolve([diff(y(x),x$2)+ 2*diff(y(x),x)+2*y(x) = 0,y(1/4*Pi) = 2, D(y)(1/4*Pi) = -2],y(x), si
```

$$y(x) = \sqrt{2} e^{-x+\frac{\pi}{4}} (\cos(x) + \sin(x))$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 27

```
DSolve[{y'[x]+2*y'[x]+2*y[x]==0,{y[Pi/4]==2,y'[Pi/4]==-2}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \sqrt{2} e^{\frac{\pi}{4}-x} (\sin(x) + \cos(x))$$

8.17 problem 23

Internal problem ID [639]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$u'' - u' + 2u = 0$$

With initial conditions

$$[u(0) = 2, u'(0) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 31

```
dsolve([diff(u(x),x$2)- diff(u(x),x)+2*u(x) = 0,u(0) = 2, D(u)(0) = 0],u(x), singsol=all)
```

$$u(x) = -\frac{2e^{\frac{x}{2}}\left(\sqrt{7}\sin\left(\frac{\sqrt{7}x}{2}\right) - 7\cos\left(\frac{\sqrt{7}x}{2}\right)\right)}{7}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 19

```
DSolve[{u'[x]+4*u'[x]+5*u[x]==0,{u[0]==2,u'[0]==0}},u[x],x,IncludeSingularSolutions -> True
```

$$u(x) \rightarrow 2e^{-2x}(2\sin(x) + \cos(x))$$

8.18 problem 24

Internal problem ID [640]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 24.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$5u'' + 2u' + 7u = 0$$

With initial conditions

$$[u(0) = 2, u'(0) = 1]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 32

```
dsolve([5*diff(u(x),x$2)+ 2*diff(u(x),x)+7*u(x) = 0,u(0) = 2, D(u)(0) = 1],u(x), singsol=all
```

$$u(x) = \frac{e^{-\frac{x}{5}} \left(7\sqrt{34} \sin\left(\frac{\sqrt{34}x}{5}\right) + 68 \cos\left(\frac{\sqrt{34}x}{5}\right) \right)}{34}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 48

```
DSolve[{5*u''[x]+2*u'[x]+7*u[x]==0,{u[0]==2,u'[0]==1}},u[x],x,IncludeSingularSolutions -> Tr
```

$$u(x) \rightarrow \frac{1}{34} e^{-x/5} \left(7\sqrt{34} \sin\left(\frac{\sqrt{34}x}{5}\right) + 68 \cos\left(\frac{\sqrt{34}x}{5}\right) \right)$$

8.19 problem 25

Internal problem ID [641]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 25.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2y' + 6y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = \alpha]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 32

```
dsolve([diff(y(x),x$2)+ 2*diff(y(x),x)+6*y(x) = 0,y(0) = 2, D(y)(0) = alpha],y(x), singsol=a
```

$$y(x) = \frac{(\sqrt{5}(\alpha + 2) \sin(\sqrt{5}x) + 10 \cos(\sqrt{5}x)) e^{-x}}{5}$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 42

```
DSolve[{y''[x]+2*y'[x]+6*y[x]==0,{y[0]==2,y'[0]==\[Alpha]}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{1}{5}e^{-x} \left(\sqrt{5}(\alpha + 2) \sin(\sqrt{5}x) + 10 \cos(\sqrt{5}x) \right)$$

8.20 problem 26

Internal problem ID [642]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 26.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2ay' + (a^2 + 1)y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)+ 2*a*diff(y(x),x)+(a^2+1)*y(x) = 0,y(0) = 1, D(y)(0) = 0],y(x), sings
```

$$y(x) = e^{-ax}(a \sin(x) + \cos(x))$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 94

```
DSolve[{y'[x]+2*a*y'[x]+(a^2+1)*y[x]==0,{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{e^{-((\sqrt{a^2-a-1}+a)x)} \left(a \left(e^{2\sqrt{a^2-a-1}x} - 1 \right) + \sqrt{a^2-a-1} \left(e^{2\sqrt{a^2-a-1}x} + 1 \right) \right)}{2\sqrt{a^2-a-1}}$$

8.21 problem 35

Internal problem ID [643]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 35.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$t^2 y'' + t y' + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(t^2*diff(y(t),t$2)+ t*diff(y(t),t)+y(t) = 0,y(t), singsol=all)
```

$$y(t) = c_1 \sin(\ln(t)) + c_2 \cos(\ln(t))$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 18

```
DSolve[t^2*y'[t]+t*y'[t]+y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow c_1 \cos(\log(t)) + c_2 \sin(\log(t))$$

8.22 problem 36

Internal problem ID [644]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 36.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$t^2 y'' + 4ty' + 2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(t^2*diff(y(t),t$2)+ 4*t*diff(y(t),t)+2*y(t) = 0,y(t), singsol=all)
```

$$y(t) = \frac{c_1}{t} + \frac{c_2}{t^2}$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 34

```
DSolve[t^2*y''[t]+4*t*y'[t]+y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow t^{-\frac{3}{2}-\frac{\sqrt{5}}{2}} \left(c_2 t^{\sqrt{5}} + c_1 \right)$$

8.23 problem 37

Internal problem ID [645]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 37.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$t^2 y'' + 3ty' + \frac{5y}{4} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(t^2*diff(y(t),t$2)+ 3*t*diff(y(t),t)+125/100*y(t) = 0,y(t), singsol=all)
```

$$y(t) = \frac{c_1 \sin\left(\frac{\ln(t)}{2}\right)}{t} + \frac{c_2 \cos\left(\frac{\ln(t)}{2}\right)}{t}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 30

```
DSolve[t^2*y''[t]+3*t*y'[t]+125/100*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{c_2 \cos\left(\frac{\log(t)}{2}\right) + c_1 \sin\left(\frac{\log(t)}{2}\right)}{t}$$

8.24 problem 38

Internal problem ID [646]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 38.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$t^2 y'' - 4ty' - 6y = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 15

```
dsolve(t^2*diff(y(t),t$2)- 4*t*diff(y(t),t)-6*y(t) = 0,y(t), singsol=all)
```

$$y(t) = \frac{c_1}{t} + c_2 t^6$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 18

```
DSolve[t^2*y''[t]-4*t*y'[t]-6*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{c_2 t^7 + c_1}{t}$$

8.25 problem 39

Internal problem ID [647]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 39.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$t^2y'' - 4ty' + 6y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(t^2*diff(y(t),t$2)-4*t*diff(y(t),t)+6*y(t) = 0,y(t), singsol=all)
```

$$y(t) = c_1t^3 + c_2t^2$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 16

```
DSolve[t^2*y'[t]-4*t*y'[t]+6*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow t^2(c_2t + c_1)$$

8.26 problem 40

Internal problem ID [648]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 40.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$t^2 y'' - t y' + 5y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(t^2*diff(y(t),t$2)- t*diff(y(t),t)+5*y(t) = 0,y(t), singsol=all)
```

$$y(t) = c_1 t \sin(2 \ln(t)) + c_2 t \cos(2 \ln(t))$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 24

```
DSolve[t^2*y''[t]-t*y'[t]+5*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow t(c_2 \cos(2 \log(t)) + c_1 \sin(2 \log(t)))$$

8.27 problem 41

Internal problem ID [649]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 41.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$t^2 y'' + 3ty' - 3y = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
dsolve(t^2*diff(y(t),t$2)+ 3*t*diff(y(t),t)-3*y(t) = 0,y(t), singsol=all)
```

$$y(t) = \frac{c_1}{t^3} + c_2 t$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 16

```
DSolve[t^2*y''[t]+3*t*y'[t]-3*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{c_1}{t^3} + c_2 t$$

8.28 problem 42

Internal problem ID [650]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 42.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$t^2 y'' + 7ty' + 10y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(t^2*diff(y(t),t$2)+ 7*t*diff(y(t),t)+10*y(t) = 0,y(t), singsol=all)
```

$$y(t) = \frac{c_1 \sin(\ln(t))}{t^3} + \frac{c_2 \cos(\ln(t))}{t^3}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 22

```
DSolve[t^2*y''[t]+7*t*y'[t]+10*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{c_2 \cos(\log(t)) + c_1 \sin(\log(t))}{t^3}$$

8.29 problem 44

Internal problem ID [651]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 44.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`, `[_2nd_order, _linear,`

$$y'' + ty' + e^{-t^2}y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 65

```
dsolve(diff(y(t),t$2)+ t*diff(y(t),t)+exp(-t^2)*y(t) = 0,y(t), singsol=all)
```

$$y(t) = c_1 \sin \left(\frac{\sqrt{2} e^{\frac{t^2}{2}} \sqrt{\pi} \operatorname{erf} \left(\frac{t\sqrt{2}}{2} \right)}{2\sqrt{e^{t^2}}} \right) + c_2 \cos \left(\frac{\sqrt{2} e^{\frac{t^2}{2}} \sqrt{\pi} \operatorname{erf} \left(\frac{t\sqrt{2}}{2} \right)}{2\sqrt{e^{t^2}}} \right)$$

✓ Solution by Mathematica

Time used: 0.059 (sec). Leaf size: 102

```
DSolve[y''[t]+t*y'[t]+exp(-t^2)*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{-\frac{1}{4}(\sqrt{4\exp+1}+1)t^2} \left(c_1 \operatorname{HermiteH} \left(-\frac{1}{2} - \frac{1}{2\sqrt{4\exp+1}}, \frac{\sqrt{4\exp+1}t}{\sqrt{2}} \right) + c_2 \operatorname{Hypergeometric1F1} \left(\frac{1}{4} \left(1 + \frac{1}{\sqrt{4\exp+1}} \right), \frac{1}{2}, \frac{1}{2} \sqrt{4\exp+1}t^2 \right) \right)$$

8.30 problem 46

Internal problem ID [652]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.3 Complex Roots of the Characteristic Equation , page 164

Problem number: 46.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$ty'' + (t^2 - 1)y' + yt^3 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
dsolve(t*diff(y(t),t$2)+(t^2-1)*diff(y(t),t)+t^3*y(t) = 0,y(t), singsol=all)
```

$$y(t) = c_1 e^{-\frac{t^2}{4}} \cos\left(\frac{t^2 \sqrt{3}}{4}\right) + c_2 e^{-\frac{t^2}{4}} \sin\left(\frac{t^2 \sqrt{3}}{4}\right)$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 48

```
DSolve[t*y''[t]+(t^2-1)*y'[t]+t^3*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{-\frac{t^2}{4}} \left(c_2 \cos\left(\frac{\sqrt{3}t^2}{4}\right) + c_1 \sin\left(\frac{\sqrt{3}t^2}{4}\right) \right)$$

9 Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

9.1	problem 1	242
9.2	problem 2	243
9.3	problem 3	244
9.4	problem 4	245
9.5	problem 5	246
9.6	problem 6	247
9.7	problem 7	248
9.8	problem 8	249
9.9	problem 9	250
9.10	problem 10	251
9.11	problem 11	252
9.12	problem 12	253
9.13	problem 13	254
9.14	problem 14	255
9.15	problem 15	256
9.16	problem 16	257
9.17	problem 23	258
9.18	problem 24	259
9.19	problem 25	260
9.20	problem 26	261
9.21	problem 27	262
9.22	problem 28	263
9.23	problem 29	264
9.24	problem 30	265
9.25	problem 40	266
9.26	problem 41	267
9.27	problem 42	268
9.28	problem 43	269
9.29	problem 44	270
9.30	problem 45	271

9.1 problem 1

Internal problem ID [653]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x) = 0,y(x), singsol=all)
```

$$y(x) = e^x c_1 + c_2 e^x x$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 16

```
DSolve[y''[x]-2*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(c_2 x + c_1)$$

9.2 problem 2

Internal problem ID [654]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$9y'' + 6y' + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(9*diff(y(x),x$2)+6*diff(y(x),x)+y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{x}{3}} + c_2 e^{-\frac{x}{3}} x$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 20

```
DSolve[9*y''[x]+6*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/3}(c_2 x + c_1)$$

9.3 problem 3

Internal problem ID [655]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4y'' - 4y' - 3y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(4*diff(y(x),x$2)-4*diff(y(x),x)-3*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{3x}{2}} + c_2 e^{-\frac{x}{2}}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 24

```
DSolve[4*y''[x]-4*y'[x]-3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/2}(c_2 e^{2x} + c_1)$$

9.4 problem 4

Internal problem ID [656]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4y'' + 12y' + 9y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(4*diff(y(x),x$2)+12*diff(y(x),x)+9*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{3x}{2}} + c_2 x e^{-\frac{3x}{2}}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 20

```
DSolve[4*y''[x]+12*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x/2}(c_2 x + c_1)$$

9.5 problem 5

Internal problem ID [657]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' + 10y = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+10*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^x \sin(3x) + c_2 e^x \cos(3x)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 24

```
DSolve[y''[x]-2*y'[x]+10*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(c_2 \cos(3x) + c_1 \sin(3x))$$

9.6 problem 6

Internal problem ID [658]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 6y' + 9y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)-6*diff(y(x),x)+9*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{3x} + c_2 e^{3x} x$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 18

```
DSolve[y''[x]-6*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{3x}(c_2 x + c_1)$$

9.7 problem 7

Internal problem ID [659]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4y'' + 17y' + 4y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(4*diff(y(x),x$2)+17*diff(y(x),x)+4*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-4x} + c_2 e^{-\frac{x}{4}}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 24

```
DSolve[4*y''[x]+17*y'[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-4x} (c_1 e^{15x/4} + c_2)$$

9.8 problem 8

Internal problem ID [660]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$16y'' + 24y' + 9y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(16*diff(y(x),x$2)+24*diff(y(x),x)+9*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{3x}{4}} + c_2 e^{-\frac{3x}{4}} x$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 20

```
DSolve[16*y'[x]+24*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x/4}(c_2 x + c_1)$$

9.9 problem 9

Internal problem ID [661]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$25y'' - 20y' + 4y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(25*diff(y(x),x$2)-20*diff(y(x),x)+4*y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{2x}{5}} + c_2 e^{\frac{2x}{5}} x$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 20

```
DSolve[25*y'[x]-20*y'[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x/5}(c_2 x + c_1)$$

9.10 problem 10

Internal problem ID [662]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$2y'' + 2y' + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(2*diff(y(x),x$2)+2*diff(y(x),x)+y(x) = 0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{x}{2}} \sin\left(\frac{x}{2}\right) + c_2 e^{-\frac{x}{2}} \cos\left(\frac{x}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 32

```
DSolve[2*y''[x]+2*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/2} \left(c_2 \cos\left(\frac{x}{2}\right) + c_1 \sin\left(\frac{x}{2}\right) \right)$$

9.11 problem 11

Internal problem ID [663]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$9y'' - 12y' + 4y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = -1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve([9*dif(y(t),t$2)-12*dif(y(t),t)+4*y(t) = 0,y(0) = 2, D(y)(0) = -1],y(t), singsol=all)
```

$$y(t) = -\frac{e^{\frac{2t}{3}}(-6 + 7t)}{3}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 15

```
DSolve[{9*y'[t]-12*y'[t]+4*y[t]==0,{y[0]==2,y'[0]==-1}},y[t],t,IncludeSingularSolutions ->
```

$$y(t) \rightarrow -e^{2t/3}t$$

9.12 problem 12

Internal problem ID [664]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 6y' + 9y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 2]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve([diff(y(t),t$2)-6*diff(y(t),t)+9*y(t) = 0,y(0) = 0, D(y)(0) = 2],y(t), singsol=all)
```

$$y(t) = 2e^{3t}t$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 13

```
DSolve[{y'[t]-6*y'[t]+9*y[t]==0,{y[0]==0,y'[0]==2}},y[t],t,IncludeSingularSolutions -> True
```

$$y(t) \rightarrow 2e^{3t}t$$

9.13 problem 13

Internal problem ID [665]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$9y'' + 6y' + 82y = 0$$

With initial conditions

$$[y(0) = -1, y'(0) = 2]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 23

```
dsolve([9*dif(y(t),t$2)+6*dif(y(t),t)+82*y(t) = 0,y(0) = -1, D(y)(0) = 2],y(t), singsol=all)
```

$$y(t) = \frac{e^{-\frac{t}{3}}(5 \sin(3t) - 9 \cos(3t))}{9}$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 29

```
DSolve[{9*y'[t]+6*y'[t]+82*y[t]==0,{y[0]==-1,y'[0]==2}},y[t],t,IncludeSingularSolutions ->
```

$$y(t) \rightarrow \frac{1}{9}e^{-t/3}(5 \sin(3t) - 9 \cos(3t))$$

9.14 problem 14

Internal problem ID [666]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 4y' + 4y = 0$$

With initial conditions

$$[y(-1) = 2, y'(-1) = 1]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 16

```
dsolve([diff(y(x),x$2)+4*diff(y(x),x)+4*y(x) = 0,y(-1) = 2, D(y)(-1) = 1],y(x), singsol=all)
```

$$y(x) = e^{-2x-2}(5x + 7)$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 18

```
DSolve[{y'[x]+4*y'[x]+4*y[x]==0,{y[-1]==2,y'[-1]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2(x+1)}(5x + 7)$$

9.15 problem 15

Internal problem ID [667]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4y'' + 12y' + 9y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = -4]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve([4*dif(y(t),t$2)+12*dif(y(t),t)+9*y(t) = 0,y(0) = 1, D(y)(0) = -4],y(t), singsol=all)
```

$$y(t) = -\frac{e^{-\frac{3t}{2}}(-2 + 5t)}{2}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 21

```
DSolve[{4*y'[t]+12*y'[t]+9*y[t]==0,{y[0]==1,y'[0]==-4}},y[t],t,IncludeSingularSolutions ->
```

$$y(t) \rightarrow \frac{1}{2}e^{-3t/2}(2 - 5t)$$

9.16 problem 16

Internal problem ID [668]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y' + \frac{y}{4} = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = b]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve([diff(y(t),t$2)-diff(y(t),t)+25/100*y(t) = 0,y(0) = 2, D(y)(0) = b],y(t), singsol=all
```

$$y(t) = e^{\frac{t}{2}}(2 + t(b - 1))$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 20

```
DSolve[{y'[t]-y'[t]+25/100*y[t]==0,{y[0]==2,y'[0]==b}},y[t],t,IncludeSingularSolutions -> T
```

$$y(t) \rightarrow e^{t/2}((b - 1)t + 2)$$

9.17 problem 23

Internal problem ID [669]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$t^2 y'' - 4ty' + 6y = 0$$

Given that one solution of the ode is

$$y_1 = t^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([t^2*diff(y(t),t$2)-4*t*diff(y(t),t)+6*y(t)=0,t^2],y(t), singsol=all)
```

$$y(t) = c_1 t^3 + c_2 t^2$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 16

```
DSolve[t^2*y''[t]-4*t*y'[t]+6*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow t^2(c_2 t + c_1)$$

9.18 problem 24

Internal problem ID [670]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 24.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$t^2 y'' + 2ty' - 2y = 0$$

Given that one solution of the ode is

$$y_1 = t$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([t^2*diff(y(t),t$2)+2*t*diff(y(t),t)-2*y(t)=0,t],y(t), singsol=all)
```

$$y(t) = \frac{c_1}{t^2} + c_2 t$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 16

```
DSolve[t^2*y''[t]+2*t*y'[t]-2*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{c_1}{t^2} + c_2 t$$

9.19 problem 25

Internal problem ID [671]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 25.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$t^2 y'' + 3ty' + y = 0$$

Given that one solution of the ode is

$$y_1 = \frac{1}{t}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve([t^2*diff(y(t),t$2)+3*t*diff(y(t),t)+y(t)=0,1/t],y(t), singsol=all)
```

$$y(t) = \frac{c_1}{t} + \frac{c_2 \ln(t)}{t}$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 17

```
DSolve[t^2*y'[t]+3*t*y'[t]+y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{c_2 \log(t) + c_1}{t}$$

9.20 problem 26

Internal problem ID [672]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 26.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$t^2 y'' - t(2+t)y' + (2+t)y = 0$$

Given that one solution of the ode is

$$y_1 = t$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([t^2*diff(y(t),t$2)-t*(t+2)*diff(y(t),t)+(t+2)*y(t)=0,t],y(t), singsol=all)
```

$$y(t) = c_1 t + c_2 t e^t$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 16

```
DSolve[t^2*y'[t]-t*(t+2)*y'[t]+(t+2)*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow t(c_2 e^t + c_1)$$

9.21 problem 27

Internal problem ID [673]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 27.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$xy'' - y' + 4x^3y = 0$$

Given that one solution of the ode is

$$y_1 = \sin(x^2)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve([x*diff(y(x),x$2)-diff(y(x),x)+4*x^3*y(x)=0,sin(x^2)],y(x), singsol=all)
```

$$y(x) = c_1 \sin(x^2) + c_2 \cos(x^2)$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 20

```
DSolve[x*y''[x]-y'[x]+4*x^3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(x^2) + c_2 \sin(x^2)$$

9.22 problem 28

Internal problem ID [674]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 28.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x - 1)y'' - y'x + y = 0$$

Given that one solution of the ode is

$$y_1 = e^x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve([(x-1)*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,exp(x)],y(x), singsol=all)
```

$$y(x) = c_1x + e^xc_2$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 17

```
DSolve[(x-1)*y'[x]-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1e^x - c_2x$$

9.23 problem 29

Internal problem ID [675]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 29.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - \left(x - \frac{3}{16}\right) y = 0$$

Given that one solution of the ode is

$$y_1 = x^{\frac{1}{4}} e^{2\sqrt{x}}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve([x^2*diff(y(x),x$2)-(x-1875/10000)*y(x)=0,x^(1/4)*exp(2*sqrt(x))],y(x), singsol=all)
```

$$y(x) = c_1 x^{\frac{1}{4}} \sinh(2\sqrt{x}) + c_2 x^{\frac{1}{4}} \cosh(2\sqrt{x})$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 41

```
DSolve[x^2*y''[x]-(x-1875/10000)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^{-2\sqrt{x}} \sqrt[4]{x} (2c_1 e^{4\sqrt{x}} - c_2)$$

9.24 problem 30

Internal problem ID [676]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 30.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y' x + \left(x^2 - \frac{1}{4}\right) y = 0$$

Given that one solution of the ode is

$$y_1 = \frac{\sin(x)}{\sqrt{x}}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve([x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-25/100)*y(x)=0,x^(-1/2)*sin(x)],y(x), singsol
```

$$y(x) = \frac{c_1 \sin(x)}{\sqrt{x}} + \frac{c_2 \cos(x)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 39

```
DSolve[x^2*y'[x]+x*y'[x]+(x^2-25/100)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{-ix}(2c_1 - ic_2 e^{2ix})}{2\sqrt{x}}$$

9.25 problem 40

Internal problem ID [677]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 40.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$t^2 y'' - 3ty' + 4y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(t^2*diff(y(t),t$2)-3*t*diff(y(t),t)+4*y(t)=0,y(t), singsol=all)
```

$$y(t) = c_1 t^2 + c_2 t^2 \ln(t)$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 18

```
DSolve[t^2*y'[t]-3*t*y'[t]+4*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow t^2(2c_2 \log(t) + c_1)$$

9.26 problem 41

Internal problem ID [678]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 41.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$t^2 y'' + 2ty' + \frac{y}{4} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(t^2*diff(y(t),t$2)+2*t*diff(y(t),t)+25/100*y(t)=0,y(t), singsol=all)
```

$$y(t) = \frac{c_1}{\sqrt{t}} + \frac{c_2 \ln(t)}{\sqrt{t}}$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 24

```
DSolve[t^2*y''[t]+2*t*y'[t]+25/100*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{c_2 \log(t) + 2c_1}{2\sqrt{t}}$$

9.27 problem 42

Internal problem ID [679]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 42.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$2t^2y'' - 5ty' + 5y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(2*t^2*diff(y(t),t$2)-5*t*diff(y(t),t)+5*y(t)=0,y(t), singsol=all)
```

$$y(t) = c_1t^{\frac{5}{2}} + c_2t$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 18

```
DSolve[2*t^2*y''[t]-5*t*y'[t]+5*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow t(c_2t^{3/2} + c_1)$$

9.28 problem 43

Internal problem ID [680]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 43.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$t^2 y'' + 3ty' + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(t^2*diff(y(t),t$2)+3*t*diff(y(t),t)+y(t)=0,y(t), singsol=all)
```

$$y(t) = \frac{c_1}{t} + \frac{c_2 \ln(t)}{t}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 17

```
DSolve[t^2*y''[t]+3*t*y'[t]+y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{c_2 \log(t) + c_1}{t}$$

9.29 problem 44

Internal problem ID [681]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 44.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$4t^2y'' - 8ty' + 9y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(4*t^2*diff(y(t),t$2)-8*t*diff(y(t),t)+9*y(t)=0,y(t), singsol=all)
```

$$y(t) = c_1t^{\frac{3}{2}} + c_2t^{\frac{3}{2}} \ln(t)$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 25

```
DSolve[4*t^2*y''[t]-8*t*y'[t]+9*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{2}t^{3/2}(3c_2 \log(t) + 2c_1)$$

9.30 problem 45

Internal problem ID [682]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.4 Repeated roots, reduction of order , page 172

Problem number: 45.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$t^2 y'' + 5ty' + 13y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(t^2*diff(y(t),t$2)+5*t*diff(y(t),t)+13*y(t)=0,y(t), singsol=all)
```

$$y(t) = \frac{c_1 \sin(3 \ln(t))}{t^2} + \frac{c_2 \cos(3 \ln(t))}{t^2}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 26

```
DSolve[t^2*y''[t]+5*t*y'[t]+13*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{c_2 \cos(3 \log(t)) + c_1 \sin(3 \log(t))}{t^2}$$

**10 Chapter 3, Second order linear equations,
section 3.6, Variation of Parameters. page 190**

10.1 problem 1	273
10.2 problem 2	274
10.3 problem 3	275
10.4 problem 4	276
10.5 problem 5	277
10.6 problem 6	278
10.7 problem 7	279
10.8 problem 8	280
10.9 problem 9	281
10.10problem 10	282
10.11problem 11	283
10.12problem 12	284
10.13problem 13	285
10.14problem 14	286
10.15problem 15	287
10.16problem 16	288
10.17problem 17	289
10.18problem 20	290
10.19problem 29	291
10.20problem 30	292
10.21problem 31	293
10.22problem 32	294

10.1 problem 1

Internal problem ID [683]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters.
page 190

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 5y' + 6y = 2e^t$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve(diff(y(t),t$2)-5*diff(y(t),t)+6*y(t) = 2*exp(t),y(t), singsol=all)
```

$$y(t) = c_2e^{3t} + c_1e^{2t} + e^t$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 25

```
DSolve[y''[t]-5*y'[t]+6*y[t]== 2*Exp[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^t(c_1e^t + c_2e^{2t} + 1)$$

10.2 problem 2

Internal problem ID [684]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters. page 190

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - y' - 2y = 2e^{-t}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(t),t$2)-diff(y(t),t)-2*y(t) = 2*exp(-t),y(t), singsol=all)
```

$$y(t) = c_2 e^{-t} + c_1 e^{2t} - \frac{2t e^{-t}}{3}$$

✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 32

```
DSolve[y''[t]-y'[t]-2*y[t] == 2*Exp[-t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{9} e^{-t} (-6t + 9c_2 e^{3t} - 2 + 9c_1)$$

10.3 problem 3

Internal problem ID [685]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters.
page 190

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 2y' + y = 3e^{-t}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve(diff(y(t),t$2)+2*diff(y(t),t)+y(t) = 3*exp(-t),y(t), singsol=all)
```

$$y(t) = c_2 e^{-t} + t e^{-t} c_1 + \frac{3 e^{-t} t^2}{2}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 29

```
DSolve[y''[t]+2*y'[t]+y[t] == 3*Exp[-t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{2} e^{-t} (3t^2 + 2c_2 t + 2c_1)$$

10.4 problem 4

Internal problem ID [686]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters.
page 190

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4y'' - 4y' + y = 16e^{\frac{t}{2}}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve(4*diff(y(t),t$2)-4*diff(y(t),t)+y(t) = 16*exp(t/2),y(t), singsol=all)
```

$$y(t) = c_2 e^{\frac{t}{2}} + t e^{\frac{t}{2}} c_1 + 2t^2 e^{\frac{t}{2}}$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 25

```
DSolve[4*y''[t]-4*y'[t]+y[t]== 16*Exp[t/2],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{t/2}(2t^2 + c_2 t + c_1)$$

10.5 problem 5

Internal problem ID [687]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters.
page 190

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \tan(t)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 23

```
dsolve(diff(y(t),t$2)+y(t) = tan(t),y(t), singsol=all)
```

$$y(t) = c_2 \sin(t) + \cos(t) c_1 - \cos(t) \ln(\sec(t) + \tan(t))$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 23

```
DSolve[y''[t]+y[t] == Tan[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \cos(t)(-\arctanh(\sin(t))) + c_1 \cos(t) + c_2 \sin(t)$$

10.6 problem 6

Internal problem ID [688]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters.
page 190

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = 9 \sec(3t)^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(t),t$2)+9*y(t) = 9*sec(3*t)^2,y(t), singsol=all)
```

$$y(t) = c_2 \sin(3t) + c_1 \cos(3t) + \ln(\sec(3t) + \tan(3t)) \sin(3t) - 1$$

✓ Solution by Mathematica

Time used: 0.117 (sec). Leaf size: 31

```
DSolve[y''[t]+9*y[t] == 9*Sec[3*t]^2,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow c_1 \cos(3t) + \sin(3t) \coth^{-1}(\sin(3t)) + c_2 \sin(3t) - 1$$

10.7 problem 7

Internal problem ID [689]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters.
page 190

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y' + 4y = \frac{e^{-2t}}{t^2}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve(diff(y(t),t$2)+4*diff(y(t),t)+4*y(t) = t^(-2)*exp(-2*t),y(t), singsol=all)
```

$$y(t) = e^{-2t}c_2 + e^{-2t}tc_1 - (\ln(t) + 1)e^{-2t}$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 23

```
DSolve[y''[t]+4*y'[t]+4*y[t] == t^(-2)*Exp[-2*t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{-2t}(-\log(t) + c_2t - 1 + c_1)$$

10.8 problem 8

Internal problem ID [690]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters.
page 190

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y = 3 \csc(2t)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```
dsolve(diff(y(t),t$2)+4*y(t) = 3*csc(2*t),y(t), singsol=all)
```

$$y(t) = c_2 \sin(2t) + c_1 \cos(2t) - \frac{3 \ln(\csc(2t)) \sin(2t)}{4} - \frac{3 \cos(2t) t}{2}$$

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 39

```
DSolve[y''[t]+4*y[t] ==3*Csc[2*t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \left(-\frac{3t}{2} + c_1\right) \cos(2t) + \frac{1}{4} \sin(2t)(3 \log(\sin(2t)) + 4c_2)$$

10.9 problem 9

Internal problem ID [691]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters. page 190

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = 2 \sec\left(\frac{t}{2}\right)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 43

```
dsolve(diff(y(t),t$2)+y(t) = 2*sec(t/2),y(t), singsol=all)
```

$$y(t) = c_2 \sin(t) + \cos(t) c_1 + \cos\left(\frac{t}{2}\right) \left(-8 \left(\ln(2) + \ln\left(\csc(t) \sin\left(\frac{t}{2}\right) \left(\sin\left(\frac{t}{2}\right) + 1\right)\right)\right) \sin\left(\frac{t}{2}\right) + 8\right)$$

✓ Solution by Mathematica

Time used: 0.114 (sec). Leaf size: 35

```
DSolve[y''[t]+y[t]== 2*Sec[t/2],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -4 \sin(t) \operatorname{arctanh}\left(\sin\left(\frac{t}{2}\right)\right) + 8 \cos\left(\frac{t}{2}\right) + c_1 \cos(t) + c_2 \sin(t)$$

10.10 problem 10

Internal problem ID [692]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters. page 190

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 2y' + y = \frac{e^t}{t^2 + 1}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(diff(y(t),t$2)-2*diff(y(t),t)+y(t) = exp(t)/(1+t^2),y(t), singsol=all)
```

$$y(t) = c_2 e^t + t e^t c_1 + e^t \left(-\frac{\ln(t^2 + 1)}{2} + t \arctan(t) \right)$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 35

```
DSolve[y''[t]-2*y'[t]+y[t] == Exp[t]/(1+t^2),y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{2} e^t (2t \arctan(t) - \log(t^2 + 1) + 2(c_2 t + c_1))$$

10.11 problem 11

Internal problem ID [693]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters.
page 190

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 5y' + 6y = g(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 46

```
dsolve(diff(y(t),t$2)-5*diff(y(t),t)+6*y(t) = g(t),y(t), singsol=all)
```

$$y(t) = c_2 e^{3t} + c_1 e^{2t} + \left(\int g(t) e^{-3t} dt \right) e^{3t} - \left(\int g(t) e^{-2t} dt \right) e^{2t}$$

✓ Solution by Mathematica

Time used: 0.064 (sec). Leaf size: 59

```
DSolve[y''[t]-5*y'[t]+6*y[t] == g[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{2t} \left(\int_1^t -e^{-2K[1]} g(K[1]) dK[1] + e^t \int_1^t e^{-3K[2]} g(K[2]) dK[2] + c_2 e^t + c_1 \right)$$

10.12 problem 12

Internal problem ID [694]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters. page 190

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y = g(t)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
dsolve(diff(y(t),t$2)+4*y(t) = g(t),y(t), singsol=all)
```

$$y(t) = c_2 \sin(2t) + c_1 \cos(2t) + \frac{(\int \cos(2t) g(t) dt) \sin(2t)}{2} - \frac{(\int \sin(2t) g(t) dt) \cos(2t)}{2}$$

✓ Solution by Mathematica

Time used: 0.093 (sec). Leaf size: 67

```
DSolve[y''[t]+4*y[t] == g[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \cos(2t) \int_1^t -\cos(K[1])g(K[1]) \sin(K[1])dK[1] \\ + \sin(2t) \int_1^t \frac{1}{2} \cos(2K[2])g(K[2])dK[2] + c_1 \cos(2t) + c_2 \sin(2t)$$

10.13 problem 13

Internal problem ID [695]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters. page 190

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$t^2 y'' - 2y = 3t^2 - 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(t^2*diff(y(t),t$2)-2*y(t) = 3*t^2-1,y(t), singsol=all)
```

$$y(t) = c_2 t^2 + t^2 \ln(t) + \frac{1}{2} + \frac{c_1}{t}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 31

```
DSolve[t^2*y'[t]-2*y[t] == 3*t^2-1,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow t^2 \log(t) + \left(-\frac{1}{3} + c_2\right) t^2 + \frac{c_1}{t} + \frac{1}{2}$$

10.14 problem 14

Internal problem ID [696]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters.
page 190

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$t^2 y'' - t(2+t)y' + (2+t)y = 2t^3$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 18

```
dsolve(t^2*diff(y(t),t$2)-t*(t+2)*diff(y(t),t)+(t+2)*y(t) = 2*t^3,y(t), singsol=all)
```

$$y(t) = c_2 t + t e^t c_1 - 2t^2$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 20

```
DSolve[t^2*y'[t]-t*(t+2)*y'[t]+(t+2)*y[t] == 2*t^3,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow t(-2t + c_2 e^t - 2 + c_1)$$

10.15 problem 15

Internal problem ID [697]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters.
page 190

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$ty'' - (t + 1)y' + y = e^{2t}t^2$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(t*diff(y(t),t$2)-(1+t)*diff(y(t),t)+y(t) = t^2*exp(2*t),y(t), singsol=all)
```

$$y(t) = (t + 1)c_2 + c_1e^t + \frac{(t - 1)e^{2t}}{2}$$

✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 31

```
DSolve[t*y''[t]-(1+t)*y'[t]+y[t] == t^2*Exp[2*t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{2}e^{2t}(t - 1) + c_1e^t - c_2(t + 1)$$

10.16 problem 16

Internal problem ID [698]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters.
page 190

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(1-t)y'' + ty' - y = 2(t-1)^2 e^{-t}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve((1-t)*diff(y(t),t$2)+t*diff(y(t),t)-y(t) = 2*(t-1)^2*exp(-t),y(t), singsol=all)
```

$$y(t) = c_2 t + c_1 e^t - \frac{(2t-1)e^{-t}}{2}$$

✓ Solution by Mathematica

Time used: 0.07 (sec). Leaf size: 30

```
DSolve[(1-t)*y''[t]+t*y'[t]-y[t] == 2*(t-1)^2*Exp[-t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{-t} \left(\frac{1}{2} - t \right) + c_1 e^t - c_2 t$$

10.17 problem 17

Internal problem ID [699]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters.
page 190

Problem number: 17.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x^2 y'' - 3y'x + 4y = \ln(x) x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(x^2*diff(y(x),x$2)-3*x*diff(y(x),x)+4*y(x) = x^2*ln(x),y(x), singsol=all)
```

$$y(x) = x^2 c_2 + \ln(x) c_1 x^2 + \frac{\ln(x)^3 x^2}{6}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 27

```
DSolve[x^2*y''[x]-3*x*y'[x]+4*y[x] == x^2*Log[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{6} x^2 (\log^3(x) + 12c_2 \log(x) + 6c_1)$$

10.18 problem 20

Internal problem ID [700]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters. page 190

Problem number: 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x^2 y'' + y'x + \left(x^2 - \frac{1}{4}\right) y = g(x)$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 51

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-25/100)*y(x) = g(x),y(x), singsol=all)
```

$$y(x) = \frac{\sin(x) c_2}{\sqrt{x}} + \frac{\cos(x) c_1}{\sqrt{x}} + \frac{\left(\int \frac{\cos(x)g(x)}{x^{\frac{3}{2}}} dx\right) \sin(x) - \left(\int \frac{\sin(x)g(x)}{x^{\frac{3}{2}}} dx\right) \cos(x)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.203 (sec). Leaf size: 107

```
DSolve[x^2*y'[x]+x*y'[x]+(x^2-25/100)*y[x] == g[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{-ix} \left(2 \int_1^x \frac{ie^{iK[1]}g(K[1])}{2K[1]^{3/2}} dK[1] - ie^{2ix} \int_1^x \frac{e^{-iK[2]}g(K[2])}{K[2]^{3/2}} dK[2] - ic_2 e^{2ix} + 2c_1 \right)}{2\sqrt{x}}$$

10.19 problem 29

Internal problem ID [701]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters.
page 190

Problem number: 29.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$t^2 y'' - 2ty' + 2y = 4t^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(t^2*diff(y(t),t$2)-2*t*diff(y(t),t)+2*y(t) = 4*t^2,y(t), singsol=all)
```

$$y(t) = c_2 t^2 + c_1 t + 4t^2(-1 + \ln(t))$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 21

```
DSolve[t^2*y'[t]-2*t*y'[t]+2*y[t] ==4*t^2,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow t(4t \log(t) + (-4 + c_2)t + c_1)$$

10.20 problem 30

Internal problem ID [702]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters. page 190

Problem number: 30.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$t^2 y'' + 7ty' + 5y = t$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(t^2*diff(y(t),t$2)+7*t*diff(y(t),t)+5*y(t) = t,y(t), singsol=all)
```

$$y(t) = \frac{c_2 + \frac{2\left(c_1 + \frac{t^2}{2}\right)^3}{3} - c_1 \left(c_1 + \frac{t^2}{2}\right)^2}{t^5}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 23

```
DSolve[t^2*y''[t]+7*t*y'[t]+5*y[t] ==t,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{c_1}{t^5} + \frac{t}{12} + \frac{c_2}{t}$$

10.21 problem 31

Internal problem ID [703]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters.
page 190

Problem number: 31.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$ty'' - (t + 1)y' + y = e^{2t}t^2$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(t*diff(y(t),t$2)-(1+t)*diff(y(t),t)+y(t) = t^2*exp(2*t),y(t), singsol=all)
```

$$y(t) = (t + 1)c_2 + c_1e^t + \frac{(t - 1)e^{2t}}{2}$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 31

```
DSolve[t*y''[t]-(1+t)*y'[t]+y[t] ==t^2*Exp[2*t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{2}e^{2t}(t - 1) + c_1e^t - c_2(t + 1)$$

10.22 problem 32

Internal problem ID [704]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, section 3.6, Variation of Parameters.
page 190

Problem number: 32.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(1-t)y'' + ty' - y = 2(t-1)e^{-t}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 46

```
dsolve((1-t)*diff(y(t),t$2)+t*diff(y(t),t)-y(t) = 2*(t-1)*exp(-t),y(t), singsol=all)
```

$$y(t) = c_2 t + c_1 e^t - 2 \left(t e^{t-1} \text{Ei}_1(t-1) - e^{2t-2} \text{Ei}_1(2t-2) - \frac{1}{2} \right) e^{-t}$$

✓ Solution by Mathematica

Time used: 0.187 (sec). Leaf size: 47

```
DSolve[(1-t)*y'[t]+t*y''[t]-y[t] ==2*(t-1)*Exp[-t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -2e^{t-2} \text{ExpIntegralEi}(2-2t) + \frac{2t \text{ExpIntegralEi}(1-t)}{e} + e^{-t} + c_1 e^t - c_2 t$$

**11 Chapter 3, Second order linear equations, 3.7
Mechanical and Electrical Vibrations. page 203**

11.1 problem 28	296
11.2 problem 29	297

11.1 problem 28

Internal problem ID [705]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.7 Mechanical and Electrical Vibrations. page 203

Problem number: 28.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$u'' + 2u = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(u(t),t$2)+2*u(t) = 0,u(t), singsol=all)
```

$$u(t) = c_1 \sin(t\sqrt{2}) + c_2 \cos(t\sqrt{2})$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 28

```
DSolve[u''[t]+2*u[t] ==0,u[t],t,IncludeSingularSolutions -> True]
```

$$u(t) \rightarrow c_1 \cos(\sqrt{2}t) + c_2 \sin(\sqrt{2}t)$$

11.2 problem 29

Internal problem ID [706]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.7 Mechanical and Electrical Vibrations. page 203

Problem number: 29.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$u'' + \frac{u'}{4} + 2u = 0$$

With initial conditions

$$[u(0) = 0, u'(0) = 2]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 20

```
dsolve([diff(u(t),t$2)+1/4*diff(u(t),t)+2*u(t) = 0,u(0) = 0, D(u)(0) = 2],u(t), singsol=all)
```

$$u(t) = \frac{16\sqrt{127}e^{-\frac{t}{8}} \sin\left(\frac{\sqrt{127}t}{8}\right)}{127}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 30

```
DSolve[{u''[t]+1/4*u'[t]+2*u[t] ==0,{u[0]==0,u'[0]==2}},u[t],t,IncludeSingularSolutions -> T
```

$$u(t) \rightarrow \frac{16e^{-t/8} \sin\left(\frac{\sqrt{127}t}{8}\right)}{\sqrt{127}}$$

**12 Chapter 3, Second order linear equations, 3.7
Forced Vibrations. page 217**

12.1 problem 21	299
12.2 problem 22	300
12.3 problem 23	301
12.4 problem 24	302

12.1 problem 21

Internal problem ID [707]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.7 Forced Vibrations. page 217

Problem number: 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$u'' + \frac{u'}{8} + 4u = 3 \cos\left(\frac{t}{4}\right)$$

With initial conditions

$$[u(0) = 2, u'(0) = 0]$$

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 46

```
dsolve([diff(u(t),t$2)+125/1000*diff(u(t),t)+4*u(t) = 3*cos(t/4),u(0) = 2, D(u)(0) = 0],u(t))
```

$$u(t) = \frac{19274 e^{-\frac{t}{16}} \sqrt{1023} \sin\left(\frac{\sqrt{1023}t}{16}\right)}{16242171} + \frac{19658 e^{-\frac{t}{16}} \cos\left(\frac{\sqrt{1023}t}{16}\right)}{15877} + \frac{96 \sin\left(\frac{t}{4}\right)}{15877} + \frac{12096 \cos\left(\frac{t}{4}\right)}{15877}$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 71

```
DSolve[{u''[t]+125/1000*u'[t]+4*u[t] ==3*Cos[t/4],{u[0]==0,u'[0]==0}},u[t],t,IncludeSingular
```

$$u(t) \rightarrow \frac{32\left(1023 \sin\left(\frac{t}{4}\right) - 130\sqrt{1023}e^{-t/16} \sin\left(\frac{\sqrt{1023}t}{16}\right) + 128898 \cos\left(\frac{t}{4}\right) - 128898e^{-t/16} \cos\left(\frac{\sqrt{1023}t}{16}\right)\right)}{5414057}$$

12.2 problem 22

Internal problem ID [708]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.7 Forced Vibrations. page 217

Problem number: 22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$u'' + \frac{u'}{8} + 4u = 3 \cos(2t)$$

With initial conditions

$$[u(0) = 2, u'(0) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 40

```
dsolve([diff(u(t),t$2)+125/1000*diff(u(t),t)+4*u(t) = 3*cos(2*t),u(0) = 2, D(u)(0) = 0],u(t))
```

$$u(t) = -\frac{382 e^{-\frac{t}{16}} \sqrt{1023} \sin\left(\frac{\sqrt{1023}t}{16}\right)}{1023} + 2 e^{-\frac{t}{16}} \cos\left(\frac{\sqrt{1023}t}{16}\right) + 12 \sin(2t)$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 39

```
DSolve[{u''[t]+125/1000*u'[t]+4*u[t] ==3*Cos[2*t]},{u[0]==0,u'[0]==0},u[t],t,IncludeSingular
```

$$u(t) \rightarrow 12 \sin(2t) - 128 \sqrt{\frac{3}{341}} e^{-t/16} \sin\left(\frac{\sqrt{1023}t}{16}\right)$$

12.3 problem 23

Internal problem ID [709]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.7 Forced Vibrations. page 217

Problem number: 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$u'' + \frac{u'}{8} + 4u = 3 \cos(6t)$$

With initial conditions

$$[u(0) = 2, u'(0) = 0]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 46

```
dsolve([diff(u(t),t$2)+125/1000*diff(u(t),t)+4*u(t) = 3*cos(6*t),u(0) = 2, D(u)(0) = 0],u(t))
```

$$u(t) = \frac{2806 e^{-\frac{t}{16}} \sqrt{1023} \sin\left(\frac{\sqrt{1023} t}{16}\right)}{1524549} + \frac{34322 e^{-\frac{t}{16}} \cos\left(\frac{\sqrt{1023} t}{16}\right)}{16393} + \frac{36 \sin(6t)}{16393} - \frac{1536 \cos(6t)}{16393}$$

✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 74

```
DSolve[{u''[t]+125/1000*u'[t]+4*u[t] ==3*Cos[6*t],{u[0]==0,u'[0]==0}},u[t],t,IncludeSingular
```

$$u(t) \rightarrow \frac{4e^{-t/16} \left(-3069e^{t/16} \sin(6t) + 160\sqrt{1023} \sin\left(\frac{\sqrt{1023}t}{16}\right) + 130944e^{t/16} \cos(6t) - 130944 \cos\left(\frac{\sqrt{1023}t}{16}\right) \right)}{5590013}$$

12.4 problem 24

Internal problem ID [710]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 3, Second order linear equations, 3.7 Forced Vibrations. page 217

Problem number: 24.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [NONE]

$$u'' + u' + \frac{u^3}{5} = \cos(t)$$

With initial conditions

$$[u(0) = 2, u'(0) = 0]$$

X Solution by Maple

```
dsolve([diff(u(t),t$2)+diff(u(t),t)+1/5*u(t)^3 = cos(t),u(0) = 2, D(u)(0) = 0],u(t), singsol
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{u''[t]+u'[t]+1/5*u[t]^3 ==3*Cos[t],{u[0]==0,u'[0]==0}},u[t],t,IncludeSingularSolutio
```

Not solved

13 Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

13.1 problem 1	304
13.2 problem 2	305
13.3 problem 4	306
13.4 problem 5	307
13.5 problem 6	308
13.6 problem 7	309
13.7 problem 9	310
13.8 problem 10	311
13.9 problem 11	312
13.10problem 12	313
13.11problem 13	314
13.12problem 15	315
13.13problem 16	316
13.14problem 17	317
13.15problem 18	318
13.16problem 21	319
13.17problem 23	320
13.18problem 24	321
13.19problem 25	322
13.20problem 26	323
13.21problem 27	324
13.22problem 28	325

13.1 problem 1

Internal problem ID [711]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 + \frac{1}{24}x^4\right) y(0) + \left(x + \frac{1}{6}x^3 + \frac{1}{120}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[y''[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{120} + \frac{x^3}{6} + x \right) + c_1 \left(\frac{x^4}{24} + \frac{x^2}{2} + 1 \right)$$

13.2 problem 2

Internal problem ID [712]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$y'' - y'x - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)-x*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 + \frac{1}{8}x^4\right)y(0) + \left(x + \frac{1}{3}x^3 + \frac{1}{15}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[y''[x]-x*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{15} + \frac{x^3}{3} + x \right) + c_1 \left(\frac{x^4}{8} + \frac{x^2}{2} + 1 \right)$$

13.3 problem 4

Internal problem ID [713]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y'' + k^2 x^2 y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
Order:=6;  
dsolve(diff(y(x),x$2)+k^2*x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{k^2 x^4}{12}\right) y(0) + \left(x - \frac{1}{20} k^2 x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 34

```
AsymptoticDSolveValue[y''[x]+k^2*x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{k^2 x^5}{20}\right) + c_1 \left(1 - \frac{k^2 x^4}{12}\right)$$

13.4 problem 5

Internal problem ID [714]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$(1 - x)y'' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 49

```
Order:=6;  
dsolve((1-x)*diff(y(x),x$2)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 - \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{60}x^5\right) y(0) + \left(x - \frac{1}{6}x^3 - \frac{1}{12}x^4 - \frac{1}{24}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 63

```
AsymptoticDSolveValue[(1-x)*y''[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(-\frac{x^5}{24} - \frac{x^4}{12} - \frac{x^3}{6} + x \right) + c_1 \left(-\frac{x^5}{60} - \frac{x^4}{24} - \frac{x^3}{6} - \frac{x^2}{2} + 1 \right)$$

13.5 problem 6

Internal problem ID [715]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 2)y'' - y'x + 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve((2+x^2)*diff(y(x),x$2)-x*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x^2 + \frac{1}{6}x^4\right)y(0) + \left(x - \frac{1}{4}x^3 + \frac{7}{160}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 40

```
AsymptoticDSolveValue[(2+x^2)*y'[x]-x*y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{7x^5}{160} - \frac{x^3}{4} + x \right) + c_1 \left(\frac{x^4}{6} - x^2 + 1 \right)$$

13.6 problem 7

Internal problem ID [716]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + y'x + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)+x*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x^2 + \frac{1}{3}x^4\right)y(0) + \left(x - \frac{1}{2}x^3 + \frac{1}{8}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 40

```
AsymptoticDSolveValue[y''[x]+x*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{8} - \frac{x^3}{2} + x \right) + c_1 \left(\frac{x^4}{3} - x^2 + 1 \right)$$

13.7 problem 9

Internal problem ID [717]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 1)y'' - 4y'x + 6y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
Order:=6;  
dsolve((1+x^2)*diff(y(x),x$2)-4*x*diff(y(x),x)+6*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = y(0) + D(y)(0)x - 3y(0)x^2 - \frac{D(y)(0)x^3}{3}$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 26

```
AsymptoticDSolveValue[(1+x^2)*y'[x]-4*x*y'[x]+6*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{x^3}{3} \right) + c_1(1 - 3x^2)$$

13.8 problem 10

Internal problem ID [718]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(-x^2 + 4)y'' + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
Order:=6;  
dsolve((4-x^2)*diff(y(x),x$2)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(-\frac{x^2}{4} + 1\right) y(0) + \left(x - \frac{1}{12}x^3 - \frac{1}{240}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 35

```
AsymptoticDSolveValue[(4-x^2)*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(1 - \frac{x^2}{4}\right) + c_2 \left(-\frac{x^5}{240} - \frac{x^3}{12} + x\right)$$

13.9 problem 11

Internal problem ID [719]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(-x^2 + 3)y'' - 3y'x - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve((3-x^2)*diff(y(x),x$2)-3*x*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{6}x^2 + \frac{1}{24}x^4\right)y(0) + \left(x + \frac{2}{9}x^3 + \frac{8}{135}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 70

```
AsymptoticDSolveValue[(3-x^2)*y''[x]-3*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{13x^5}{1080} + \frac{x^4}{36} + \frac{x^3}{18} + \frac{x^2}{6} + 1 \right) + c_2 \left(\frac{49x^5}{1080} + \frac{7x^4}{72} + \frac{2x^3}{9} + \frac{x^2}{2} + x \right)$$

13.10 problem 12

Internal problem ID [720]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(1-x)y'' + y'x - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
Order:=6;  
dsolve((1-x)*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5\right)y(0) + D(y)(0)x + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 41

```
AsymptoticDSolveValue[(1-x)*y''[x]+x*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^5}{120} + \frac{x^4}{24} + \frac{x^3}{6} + \frac{x^2}{2} + 1 \right) + c_2 x$$

13.11 problem 13

Internal problem ID [721]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2y'' + y'x + 3y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(2*diff(y(x),x$2)+x*diff(y(x),x)+3*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{3}{4}x^2 + \frac{5}{32}x^4\right) y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{20}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[2*y''[x]+x*y'[x]+3*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{20} - \frac{x^3}{3} + x \right) + c_1 \left(\frac{5x^4}{32} - \frac{3x^2}{4} + 1 \right)$$

13.12 problem 15

Internal problem ID [722]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$y'' - y'x - y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = 1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([diff(y(x),x$2)-x*diff(y(x),x)-y(x)=0,y(0) = 2, D(y)(0) = 1],y(x),type='series',x=0);
```

$$y(x) = 2 + x + x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4 + \frac{1}{15}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 30

```
AsymptoticDSolveValue[{y'[x]-x*y'[x]-y[x]==0,{y[0]==2,y'[0]==1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{x^5}{15} + \frac{x^4}{4} + \frac{x^3}{3} + x^2 + x + 2$$

13.13 problem 16

Internal problem ID [723]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 2)y'' - y'x + 4y = 0$$

With initial conditions

$$[y(0) = -1, y'(0) = 3]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;  
dsolve([(2+x^2)*diff(y(x),x$2)-x*diff(y(x),x)+4*y(x)=0,y(0) = -1, D(y)(0) = 3],y(x),type='se
```

$$y(x) = -1 + 3x + x^2 - \frac{3}{4}x^3 - \frac{1}{6}x^4 + \frac{21}{160}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 32

```
AsymptoticDSolveValue[{(2+x^2)*y''[x]-x*y'[x]+4*y[x]==0,{y[0]==-1,y'[0]==3}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{21x^5}{160} - \frac{x^4}{6} - \frac{3x^3}{4} + x^2 + 3x - 1$$

13.14 problem 17

Internal problem ID [724]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 17.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + y'x + 2y = 0$$

With initial conditions

$$[y(0) = 4, y'(0) = -1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;  
dsolve([diff(y(x),x$2)+x*diff(y(x),x)+2*y(x)=0,y(0) = 4, D(y)(0) = -1],y(x),type='series',x=
```

$$y(x) = 4 - x - 4x^2 + \frac{1}{2}x^3 + \frac{4}{3}x^4 - \frac{1}{8}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 34

```
AsymptoticDSolveValue[{y'[x]+x*y'[x]+2*y[x]==0,{y[0]==4,y'[0]==-1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{x^5}{8} + \frac{4x^4}{3} + \frac{x^3}{2} - 4x^2 - x + 4$$

13.15 problem 18

Internal problem ID [725]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 18.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(1-x)y'' + y'x - y = 0$$

With initial conditions

$$[y(0) = -3, y'(0) = 2]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

Order:=6;

```
dsolve([(1-x)*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=0,y(0) = -3, D(y)(0) = 2],y(x),type='series
```

$$y(x) = -3 + 2x - \frac{3}{2}x^2 - \frac{1}{2}x^3 - \frac{1}{8}x^4 - \frac{1}{40}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 36

```
AsymptoticDSolveValue[{(1-x)*y''[x]+x*y'[x]-y[x]==0,{y[0]==-3,y'[0]==2}},y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{x^5}{40} - \frac{x^4}{8} - \frac{x^3}{2} - \frac{3x^2}{2} + 2x - 3$$

13.16 problem 21

Internal problem ID [726]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 2y'x + \lambda y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 63

```
Order:=6;  
dsolve(diff(y(x),x$2)-2*x*diff(y(x),x)+lambda*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{\lambda x^2}{2} + \frac{\lambda(\lambda - 4)x^4}{24}\right) y(0) + \left(x - \frac{(\lambda - 2)x^3}{6} + \frac{(\lambda - 2)(-6 + \lambda)x^5}{120}\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 80

```
AsymptoticDSolveValue[y''[x]-2*x*y'[x]+\[Lambda]*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{\lambda^2 x^5}{120} - \frac{\lambda x^5}{15} + \frac{x^5}{10} - \frac{\lambda x^3}{6} + \frac{x^3}{3} + x \right) + c_1 \left(\frac{\lambda^2 x^4}{24} - \frac{\lambda x^4}{6} - \frac{\lambda x^2}{2} + 1 \right)$$

13.17 problem 23

Internal problem ID [727]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$y'' - y'x - y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;
```

```
dsolve([diff(y(x),x$2)-x*diff(y(x),x)-y(x)=0,y(0) = 1, D(y)(0) = 0],y(x),type='series',x=0);
```

$$y(x) = 1 + \frac{1}{2}x^2 + \frac{1}{8}x^4 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 19

```
AsymptoticDSolveValue[{y'[x]-x*y'[x]-y[x]==0,{y[0]==1,y'[0]==0}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{x^4}{8} + \frac{x^2}{2} + 1$$

13.18 problem 24

Internal problem ID [728]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 24.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 2)y'' - y'x + 4y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;  
dsolve([(2+x^2)*diff(y(x),x$2)-x*diff(y(x),x)+4*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x),type='series')
```

$$y(x) = 1 - x^2 + \frac{1}{6}x^4 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 17

```
AsymptoticDSolveValue[{(2+x^2)*y''[x]-x*y'[x]+4*y[x]==0,{y[0]==1,y'[0]==0}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{x^4}{6} - x^2 + 1$$

13.19 problem 25

Internal problem ID [729]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 25.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + y'x + 2y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;  
dsolve([diff(y(x),x$2)+x*diff(y(x),x)+2*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x),type='series',x=0)
```

$$y(x) = x - \frac{1}{2}x^3 + \frac{1}{8}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 19

```
AsymptoticDSolveValue[{y'[x]+x*y'[x]+2*y[x]==0,{y[0]==0,y'[0]==1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{x^5}{8} - \frac{x^3}{2} + x$$

13.20 problem 26

Internal problem ID [730]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 26.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(-x^2 + 4)y'' + y'x + 2y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;  
dsolve([(4-x^2)*diff(y(x),x$2)+x*diff(y(x),x)+2*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x),type='series')
```

$$y(x) = x - \frac{1}{8}x^3 - \frac{1}{640}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 19

```
AsymptoticDSolveValue[{(4-x^2)*y''[x]+x*y'[x]+2*y[x]==0,{y[0]==0,y'[0]==1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{x^5}{640} - \frac{x^3}{8} + x$$

13.21 problem 27

Internal problem ID [731]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 27.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [[_Emden, _Fowler]]

$$y'' + x^2y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
Order:=6;  
dsolve([diff(y(x),x$2)+x^2*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x),type='series',x=0);
```

$$y(x) = 1 - \frac{1}{12}x^4 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 12

```
AsymptoticDSolveValue[{y'[x]+x^2*y[x]==0,{y[0]==1,y'[0]==0}},y[x],{x,0,5}]
```

$$y(x) \rightarrow 1 - \frac{x^4}{12}$$

13.22 problem 28

Internal problem ID [732]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.2, Series Solutions Near an Ordinary Point, Part I. page 263

Problem number: 28.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`, `[_2nd_order, _linear,`

$$(1 - x)y'' + y'x - 2y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
Order:=6;
```

```
dsolve([(1-x)*diff(y(x),x$2)+x*diff(y(x),x)-2*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x),type='series')
```

$$y(x) = x + \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{1}{24}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 26

```
AsymptoticDSolveValue[{{(1-x)*y''[x]+x*y'[x]-2*y[x]==0,{y[0]==0,y'[0]==1}},y[x]},{x,0,5}]
```

$$y(x) \rightarrow \frac{x^5}{24} + \frac{x^4}{12} + \frac{x^3}{6} + x$$

14 Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

14.1	problem 1	327
14.2	problem 2	328
14.3	problem 3	329
14.4	problem 4	330
14.5	problem 5. case $x_0 = 0$	331
14.6	problem 5. case $x_0 = 4$	332
14.7	problem 6. case $x_0 = 0$	333
14.8	problem 6. case $x_0 = 4$ only	334
14.9	problem 6. case $x_0 = -4$	335
14.10	problem 7. case $x_0 = 0$	336
14.11	problem 7. case $x_0 = 2$	337
14.12	problem 8	339
14.13	problem 10	340
14.14	problem 16	341
14.15	problem 17	342
14.16	problem 19	343
14.17	problem 22	344

14.1 problem 1

Internal problem ID [733]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$y'' + y'x + y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;
```

```
dsolve([diff(y(x),x$2)+x*diff(y(x),x)+y(x)=0,y(0) = 1, D(y)(0) = 0],y(x),type='series',x=0);
```

$$y(x) = 1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 19

```
AsymptoticDSolveValue[{y'[x]+x*y'[x]+y[x]==0,{y[0]==1,y'[0]==0}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{x^4}{8} - \frac{x^2}{2} + 1$$

14.2 problem 2

Internal problem ID [734]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$y'' + \sin(x)y' + \cos(x)y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

Order:=6;

```
dsolve([diff(y(x),x$2)+sin(x)*diff(y(x),x)+cos(x)*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x),type='s
```

$$y(x) = x - \frac{1}{3}x^3 + \frac{1}{10}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 19

```
AsymptoticDSolveValue[{y'[x]+Sin[x]*y'[x]+Cos[x]*y[x]==0,{y[0]==0,y'[0]==1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{x^5}{10} - \frac{x^3}{3} + x$$

14.3 problem 3

Internal problem ID [735]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + (x + 1) y' + 3 \ln(x) y = 0$$

With initial conditions

$$[y(1) = 2, y'(1) = 0]$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

`Order:=6;`

`dsolve([x^2*diff(y(x),x$2)+(1+x)*diff(y(x),x)+3*ln(x)*y(x)=0,y(1) = 2, D(y)(1) = 0],y(x),typ`

$$y(x) = 2 - (x - 1)^3 + \frac{7}{4}(x - 1)^4 - \frac{49}{20}(x - 1)^5 + O((x - 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 30

`AsymptoticDSolveValue[{x^2*y''[x]+(1+x)*y'[x]+3*Log[x]*y[x]==0,{y[1]==2,y'[1]==0}},y[x],{x,1`

$$y(x) \rightarrow -\frac{49}{20}(x - 1)^5 + \frac{7}{4}(x - 1)^4 - (x - 1)^3 + 2$$

14.4 problem 4

Internal problem ID [736]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + y'x^2 + \sin(x)y = 0$$

With initial conditions

$$[y(0) = a_0, y'(0) = a_1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;
```

```
dsolve([diff(y(x),x$2)+x^2*diff(y(x),x)+sin(x)*y(x)=0,y(0) = a__0, D(y)(0) = a__1],y(x),type
```

$$y(x) = a_0 + a_1x - \frac{1}{6}a_0x^3 - \frac{1}{6}a_1x^4 + \frac{1}{120}a_0x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 32

```
AsymptoticDSolveValue[{y'[x]+x^2*y'[x]+Sin[x]*y[x]==0,{y[0]==a0,y'[0]==a1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{a_0x^5}{120} - \frac{a_0x^3}{6} + a_0 - \frac{a_1x^4}{6} + a_1x$$

14.5 problem 5. case $x_0 = 0$

Internal problem ID [737]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 5. case $x_0 = 0$.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 4y' + 6yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 47

```
Order:=6;  
dsolve(diff(y(x),x$2)+4*diff(y(x),x)+6*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x^3 + x^4 - \frac{4}{5}x^5\right) y(0) + \left(x - 2x^2 + \frac{8}{3}x^3 - \frac{19}{6}x^4 + \frac{47}{15}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 55

```
AsymptoticDSolveValue[y''[x]+4*y'[x]+6*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{4x^5}{5} + x^4 - x^3 + 1 \right) + c_2 \left(\frac{47x^5}{15} - \frac{19x^4}{6} + \frac{8x^3}{3} - 2x^2 + x \right)$$

14.6 problem 5. case $x_0 = 4$

Internal problem ID [738]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 5. case $x_0 = 4$.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 4y' + 6yx = 0$$

With the expansion point for the power series method at $x = 4$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 54

```
Order:=6;  
dsolve(diff(y(x),x$2)+4*diff(y(x),x)+6*x*y(x)=0,y(x),type='series',x=4);
```

$$y(x) = \left(1 - 12(x-4)^2 + 15(x-4)^3 + 9(x-4)^4 - \frac{108(x-4)^5}{5} \right) y(4) \\ + \left(x - 4 - 2(x-4)^2 - \frac{4(x-4)^3}{3} + \frac{29(x-4)^4}{6} - \frac{5(x-4)^5}{3} \right) D(y)(4) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 79

```
AsymptoticDSolveValue[y''[x]+4*y'[x]+6*x*y[x]==0,y[x],{x,4,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{108}{5}(x-4)^5 + 9(x-4)^4 + 15(x-4)^3 - 12(x-4)^2 + 1 \right) \\ + c_2 \left(-\frac{5}{3}(x-4)^5 + \frac{29}{6}(x-4)^4 - \frac{4}{3}(x-4)^3 - 2(x-4)^2 + x - 4 \right)$$

14.7 problem 6. case $x_0 = 0$

Internal problem ID [739]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 6. case $x_0 = 0$.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 - 2x - 3)y'' + y'x + 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 49

```
Order:=6;  
dsolve((x^2-2*x-3)*diff(y(x),x$2)+x*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{2}{3}x^2 - \frac{4}{27}x^3 + \frac{16}{81}x^4 - \frac{1}{9}x^5\right) y(0) \\ + \left(x + \frac{5}{18}x^3 - \frac{5}{54}x^4 + \frac{7}{72}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 63

```
AsymptoticDSolveValue[(x^2-2*x-3)*y'[x]+x*y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{7x^5}{72} - \frac{5x^4}{54} + \frac{5x^3}{18} + x \right) + c_1 \left(-\frac{x^5}{9} + \frac{16x^4}{81} - \frac{4x^3}{27} + \frac{2x^2}{3} + 1 \right)$$

14.8 problem 6. case $x_0 = 4$ only

Internal problem ID [740]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 6. case $x_0 = 4$ only.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 - 2x - 3)y'' + y'x + 4y = 0$$

With the expansion point for the power series method at $x = 4$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
Order:=6;
```

```
dsolve((x^2-2*x-3)*diff(y(x),x$2)+x*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=4);
```

$$y(x) = \left(1 - \frac{2(x-4)^2}{5} + \frac{4(x-4)^3}{15} - \frac{4(x-4)^4}{25} + \frac{199(x-4)^5}{1875}\right) y(4) \\ + \left(x - 4 - \frac{2(x-4)^2}{5} + \frac{(x-4)^3}{10} - \frac{2(x-4)^4}{75} + \frac{157(x-4)^5}{15000}\right) D(y)(4) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 87

```
AsymptoticDSolveValue[(x^2-2*x-3)*y'[x]+x*y'[x]+4*y[x]==0,y[x],{x,4,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{199(x-4)^5}{1875} - \frac{4}{25}(x-4)^4 + \frac{4}{15}(x-4)^3 - \frac{2}{5}(x-4)^2 + 1 \right) \\ + c_2 \left(\frac{157(x-4)^5}{15000} - \frac{2}{75}(x-4)^4 + \frac{1}{10}(x-4)^3 - \frac{2}{5}(x-4)^2 + x - 4 \right)$$

14.9 problem 6. case $x_0 = -4$

Internal problem ID [741]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 6. case $x_0 = -4$.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 - 2x - 3)y'' + y'x + 4y = 0$$

With the expansion point for the power series method at $x = -4$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 54

```
Order:=6;
```

```
dsolve((x^2-2*x-3)*diff(y(x),x$2)+x*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=-4);
```

$$y(x) = \left(1 - \frac{2(x+4)^2}{21} - \frac{4(x+4)^3}{189} - \frac{4(x+4)^4}{1323} - \frac{(x+4)^5}{3087}\right) y(-4) \\ + \left(x+4 + \frac{2(x+4)^2}{21} - \frac{(x+4)^3}{54} - \frac{11(x+4)^4}{1323} - \frac{157(x+4)^5}{74088}\right) D(y)(-4) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 87

```
AsymptoticDSolveValue[(x^2-2*x-3)*y'[x]+x*y'[x]+4*y[x]==0,y[x],{x,-4,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{(x+4)^5}{3087} - \frac{4(x+4)^4}{1323} - \frac{4}{189}(x+4)^3 - \frac{2}{21}(x+4)^2 + 1 \right) \\ + c_2 \left(-\frac{157(x+4)^5}{74088} - \frac{11(x+4)^4}{1323} - \frac{1}{54}(x+4)^3 + \frac{2}{21}(x+4)^2 + x+4 \right)$$

14.10 problem 7. case $x_0 = 0$

Internal problem ID [742]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 7. case $x_0 = 0$.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^3 + 1)y'' + 4y'x + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
Order:=6;  
dsolve((1+x^3)*diff(y(x),x$2)+4*x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{3}{8}x^4 + \frac{1}{20}x^5\right)y(0) + \left(x - \frac{5}{6}x^3 + \frac{13}{24}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 49

```
AsymptoticDSolveValue[(1+x^3)*y''[x]+4*x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{13x^5}{24} - \frac{5x^3}{6} + x \right) + c_1 \left(\frac{x^5}{20} + \frac{3x^4}{8} - \frac{x^2}{2} + 1 \right)$$

14.11 problem 7. case $x_0 = 2$

Internal problem ID [743]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 7. case $x_0 = 2$.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^3 + 1)y'' + 4y'x + y = 0$$

With the expansion point for the power series method at $x = 2$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
Order:=6;  
dsolve((1+x^3)*diff(y(x),x$2)+4*x*diff(y(x),x)+y(x)=0,y(x),type='series',x=2);
```

$$y(x) = \left(1 - \frac{(-2+x)^2}{18} + \frac{10(-2+x)^3}{243} - \frac{451(-2+x)^4}{17496} + \frac{1151(-2+x)^5}{78732}\right) y(2) \\ + \left(-2+x - \frac{4(-2+x)^2}{9} + \frac{115(-2+x)^3}{486} - \frac{271(-2+x)^4}{2187} + \frac{9713(-2+x)^5}{157464}\right) D(y)(2) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 87

```
AsymptoticDSolveValue[(1+x^3)*y''[x]+4*x*y'[x]+y[x]==0,y[x],{x,2,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1151(x-2)^5}{78732} - \frac{451(x-2)^4}{17496} + \frac{10}{243}(x-2)^3 - \frac{1}{18}(x-2)^2 + 1 \right) \\ + c_2 \left(\frac{9713(x-2)^5}{157464} - \frac{271(x-2)^4}{2187} + \frac{115}{486}(x-2)^3 - \frac{4}{9}(x-2)^2 + x - 2 \right)$$

14.12 problem 8

Internal problem ID [744]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$xy'' + y = 0$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

```
Order:=6;  
dsolve(x*diff(y(x),x$2)+y(x)=0,y(x),type='series',x=1);
```

$$y(x) = \left(1 - \frac{(x-1)^2}{2} + \frac{(x-1)^3}{6} - \frac{(x-1)^4}{24} + \frac{(x-1)^5}{60}\right) y(1) \\ + \left(x - 1 - \frac{(x-1)^3}{6} + \frac{(x-1)^4}{12} - \frac{(x-1)^5}{24}\right) D(y)(1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 78

```
AsymptoticDSolveValue[x*y''[x]+y[x]==0,y[x],{x,1,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{60}(x-1)^5 - \frac{1}{24}(x-1)^4 + \frac{1}{6}(x-1)^3 - \frac{1}{2}(x-1)^2 + 1 \right) \\ + c_2 \left(-\frac{1}{24}(x-1)^5 + \frac{1}{12}(x-1)^4 - \frac{1}{6}(x-1)^3 + x - 1 \right)$$

14.13 problem 10

Internal problem ID [745]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [_Gegenbauer, [_2nd_order, _linear, '_with_symmetry_[0,F(x)]']

$$(-x^2 + 1)y'' - y'x + \alpha^2 y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 71

```
Order:=6;  
dsolve((1-x^2)*diff(y(x),x$2)-x*diff(y(x),x)+alpha^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{\alpha^2 x^2}{2} + \frac{\alpha^2(\alpha^2 - 4)x^4}{24}\right) y(0) + \left(x - \frac{(\alpha^2 - 1)x^3}{6} + \frac{(\alpha^4 - 10\alpha^2 + 9)x^5}{120}\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 88

```
AsymptoticDSolveValue[(1-x^2)*y''[x]-x*y'[x]+\[Alpha]^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{\alpha^4 x^5}{120} - \frac{\alpha^2 x^5}{12} + \frac{3x^5}{40} - \frac{\alpha^2 x^3}{6} + \frac{x^3}{6} + x \right) + c_1 \left(\frac{\alpha^4 x^4}{24} - \frac{\alpha^2 x^4}{6} - \frac{\alpha^2 x^2}{2} + 1 \right)$$

14.14 problem 16

Internal problem ID [746]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5\right)y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 37

```
AsymptoticDSolveValue[y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^5}{120} + \frac{x^4}{24} + \frac{x^3}{6} + \frac{x^2}{2} + x + 1 \right)$$

14.15 problem 17

Internal problem ID [747]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 17.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$-yx + y' = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
Order:=6;  
dsolve(diff(y(x),x)-x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 + \frac{1}{8}x^4\right)y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 22

```
AsymptoticDSolveValue[y'[x]-x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^4}{8} + \frac{x^2}{2} + 1 \right)$$

14.16 problem 19

Internal problem ID [748]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(1 - x)y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
Order:=6;  
dsolve((1-x)*diff(y(x),x)=y(x),y(x),type='series',x=0);
```

$$y(x) = (x^5 + x^4 + x^3 + x^2 + x + 1)y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 21

```
AsymptoticDSolveValue[(1-x)*y'[x]==y[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(x^5 + x^4 + x^3 + x^2 + x + 1)$$

14.17 problem 22

Internal problem ID [749]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 5.3, Series Solutions Near an Ordinary Point, Part II. page 269

Problem number: 22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Gegenbauer]

$$(-x^2 + 1)y'' - 2y'x + \alpha(1 + \alpha)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 101

```
Order:=6;
```

```
dsolve((1-x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+alpha*(alpha+1)*y(x)=0,y(x),type='series',x=0
```

$$y(x) = \left(1 - \frac{\alpha(1 + \alpha)x^2}{2} + \frac{\alpha(\alpha^3 + 2\alpha^2 - 5\alpha - 6)x^4}{24}\right) y(0) \\ + \left(x - \frac{(\alpha^2 + \alpha - 2)x^3}{6} + \frac{(\alpha^4 + 2\alpha^3 - 13\alpha^2 - 14\alpha + 24)x^5}{120}\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 127

```
AsymptoticDSolveValue[(1-x^2)*y''[x]-2*x*y'[x]+\[Alpha]*(\[Alpha]+1)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{1}{60} (-\alpha^2 - \alpha) x^5 - \frac{1}{120} (-\alpha^2 - \alpha) (\alpha^2 + \alpha) x^5 - \frac{1}{10} (\alpha^2 + \alpha) x^5 + \frac{x^5}{5} \right. \\ \left. - \frac{1}{6} (\alpha^2 + \alpha) x^3 + \frac{x^3}{3} + x \right) + c_1 \left(\frac{1}{24} (\alpha^2 + \alpha)^2 x^4 - \frac{1}{4} (\alpha^2 + \alpha) x^4 - \frac{1}{2} (\alpha^2 + \alpha) x^2 + 1 \right)$$

**15 Chapter 7.5, Homogeneous Linear Systems
with Constant Coefficients. page 407**

15.1 problem 30 346

15.1 problem 30

Internal problem ID [750]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.5, Homogeneous Linear Systems with Constant Coefficients. page 407

Problem number: 30.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= -\frac{x_1(t)}{10} + \frac{3x_2(t)}{40} \\x_2'(t) &= \frac{x_1(t)}{10} - \frac{x_2(t)}{5}\end{aligned}$$

With initial conditions

$$[x_1(0) = -17, x_2(0) = -21]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 34

```
dsolve([diff(x__1(t),t) = -1/10*x__1(t)+3/40*x__2(t), diff(x__2(t),t) = 1/10*x__1(t)-1/5*x__2(t)], [x__1(t), x__2(t)])
```

$$x_1(t) = \frac{29 e^{-\frac{t}{4}}}{8} - \frac{165 e^{-\frac{t}{20}}}{8}$$

$$x_2(t) = -\frac{29 e^{-\frac{t}{4}}}{4} - \frac{55 e^{-\frac{t}{20}}}{4}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 52

```
DSolve[{x1'[t]==-1/10*x1[t]+3/40*x2[t],x2'[t]==1/10*x1[t]-1/5*x2[t]},{x1[0]==-17,x2[0]==-21}
```

$$x1(t) \rightarrow \frac{1}{8}e^{-t/4}(29 - 165e^{t/5})$$

$$x2(t) \rightarrow -\frac{1}{4}e^{-t/4}(55e^{t/5} + 29)$$

16 Chapter 7.6, Complex Eigenvalues. page 417

16.1 problem 1	349
16.2 problem 2	350
16.3 problem 3	351
16.4 problem 4	352
16.5 problem 5	353
16.6 problem 6	354
16.7 problem 7	355
16.8 problem 8	357
16.9 problem 9	359
16.10problem 10	360
16.11problem 11	361
16.12problem 12	362
16.13problem 23	363
16.14problem 24	365
16.15problem 25	367

16.1 problem 1

Internal problem ID [751]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.6, Complex Eigenvalues. page 417

Problem number: 1.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 3x_1(t) - 2x_2(t)$$

$$x_2'(t) = 4x_1(t) - x_2(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 54

```
dsolve([diff(x__1(t),t)=3*x__1(t)-2*x__2(t),diff(x__2(t),t)=4*x__1(t)-1*x__2(t)],[x__1(t), x
```

$$x_1(t) = \frac{e^t(c_1 \sin(2t) - c_2 \sin(2t) + c_1 \cos(2t) + c_2 \cos(2t))}{2}$$

$$x_2(t) = e^t(c_1 \sin(2t) + c_2 \cos(2t))$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 58

```
DSolve[{x1'[t]==3*x1[t]-2*x2[t],x2'[t]==4*x1[t]-1*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x_1(t) \rightarrow e^t(c_1 \cos(2t) + (c_1 - c_2) \sin(2t))$$

$$x_2(t) \rightarrow e^t(c_2 \cos(2t) + (2c_1 - c_2) \sin(2t))$$

16.2 problem 2

Internal problem ID [752]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.6, Complex Eigenvalues. page 417

Problem number: 2.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -x_1(t) - 4x_2(t)$$

$$x_2'(t) = x_1(t) - x_2(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 46

```
dsolve([diff(x__1(t),t)=-1*x__1(t)-4*x__2(t),diff(x__2(t),t)=1*x__1(t)-1*x__2(t)],[x__1(t),
```

$$x_1(t) = -2e^{-t}(c_2 \sin(2t) - c_1 \cos(2t))$$

$$x_2(t) = e^{-t}(c_1 \sin(2t) + c_2 \cos(2t))$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 55

```
DSolve[{x1'[t]==-1*x1[t]-4*x2[t],x2'[t]==1*x1[t]-1*x2[t]},{x1[t],x2[t]},t,IncludeSingularSol
```

$$x1(t) \rightarrow e^{-t}(c_1 \cos(2t) - 2c_2 \sin(2t))$$

$$x2(t) \rightarrow \frac{1}{2}e^{-t}(2c_2 \cos(2t) + c_1 \sin(2t))$$

16.3 problem 3

Internal problem ID [753]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.6, Complex Eigenvalues. page 417

Problem number: 3.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 2x_1(t) - 5x_2(t)$$

$$x_2'(t) = x_1(t) - 2x_2(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 37

```
dsolve([diff(x__1(t),t)=2*x__1(t)-5*x__2(t),diff(x__2(t),t)=1*x__1(t)-2*x__2(t)],[x__1(t), x
```

$$x_1(t) = \cos(t) c_1 - c_2 \sin(t) + 2c_1 \sin(t) + 2c_2 \cos(t)$$

$$x_2(t) = c_1 \sin(t) + c_2 \cos(t)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 41

```
DSolve[{x1'[t]==2*x1[t]-5*x2[t],x2'[t]==1*x1[t]-2*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x1(t) \rightarrow c_1(2 \sin(t) + \cos(t)) - 5c_2 \sin(t)$$

$$x2(t) \rightarrow c_2 \cos(t) + (c_1 - 2c_2) \sin(t)$$

16.4 problem 4

Internal problem ID [754]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.6, Complex Eigenvalues. page 417

Problem number: 4.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 2x_1(t) - \frac{5x_2(t)}{2} \\x_2'(t) &= \frac{9x_1(t)}{5} - x_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 58

```
dsolve([diff(x__1(t),t)=2*x__1(t)-5/2*x__2(t),diff(x__2(t),t)=9/5*x__1(t)-1*x__2(t)], [x__1(t)
```

$$x_1(t) = \frac{5e^{\frac{t}{2}} \left(\sin\left(\frac{3t}{2}\right) c_1 - \sin\left(\frac{3t}{2}\right) c_2 + \cos\left(\frac{3t}{2}\right) c_1 + \cos\left(\frac{3t}{2}\right) c_2 \right)}{6}$$

$$x_2(t) = e^{\frac{t}{2}} \left(\sin\left(\frac{3t}{2}\right) c_1 + \cos\left(\frac{3t}{2}\right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 84

```
DSolve[{x1'[t]==2*x1[t]-5/2*x2[t],x2'[t]==9/5*x1[t]-1*x2[t]},{x1[t],x2[t]},t,IncludeSingular
```

$$x1(t) \rightarrow \frac{1}{3}e^{t/2} \left(3c_1 \cos\left(\frac{3t}{2}\right) + (3c_1 - 5c_2) \sin\left(\frac{3t}{2}\right) \right)$$

$$x2(t) \rightarrow \frac{1}{5}e^{t/2} \left(5c_2 \cos\left(\frac{3t}{2}\right) + (6c_1 - 5c_2) \sin\left(\frac{3t}{2}\right) \right)$$

16.5 problem 5

Internal problem ID [755]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.6, Complex Eigenvalues. page 417

Problem number: 5.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= x_1(t) - x_2(t) \\x_2'(t) &= 5x_1(t) - 3x_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 48

```
dsolve([diff(x__1(t),t)=1*x__1(t)-1*x__2(t),diff(x__2(t),t)=5*x__1(t)-3*x__2(t)],[x__1(t), x
```

$$x_1(t) = \frac{e^{-t}(\cos(t)c_1 - c_2 \sin(t) + 2c_1 \sin(t) + 2c_2 \cos(t))}{5}$$

$$x_2(t) = e^{-t}(c_1 \sin(t) + c_2 \cos(t))$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 56

```
DSolve[{x1'[t]==1*x1[t]-1*x2[t],x2'[t]==5*x1[t]-3*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x1(t) \rightarrow e^{-t}(c_1 \cos(t) + (2c_1 - c_2) \sin(t))$$

$$x2(t) \rightarrow e^{-t}(c_2 \cos(t) + (5c_1 - 2c_2) \sin(t))$$

16.6 problem 6

Internal problem ID [756]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.6, Complex Eigenvalues. page 417

Problem number: 6.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= x_1(t) + 2x_2(t) \\x_2'(t) &= -5x_1(t) - x_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 50

```
dsolve([diff(x__1(t),t)=1*x__1(t)+2*x__2(t),diff(x__2(t),t)=-5*x__1(t)-1*x__2(t)], [x__1(t),
```

$$x_1(t) = -\frac{3c_1 \cos(3t)}{5} + \frac{3c_2 \sin(3t)}{5} - \frac{c_1 \sin(3t)}{5} - \frac{c_2 \cos(3t)}{5}$$

$$x_2(t) = c_1 \sin(3t) + c_2 \cos(3t)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 54

```
DSolve[{x1'[t]==1*x1[t]+2*x2[t],x2'[t]==-5*x1[t]-1*x2[t]},{x1[t],x2[t]},t,IncludeSingularSol
```

$$x_1(t) \rightarrow c_1 \cos(3t) + \frac{1}{3}(c_1 + 2c_2) \sin(3t)$$

$$x_2(t) \rightarrow c_2 \cos(3t) - \frac{1}{3}(5c_1 + c_2) \sin(3t)$$

16.7 problem 7

Internal problem ID [757]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.6, Complex Eigenvalues. page 417

Problem number: 7.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t)$$

$$x_2'(t) = 2x_1(t) + x_2(t) - 2x_3(t)$$

$$x_3'(t) = 3x_1(t) + 2x_2(t) + x_3(t)$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 54

```
dsolve([diff(x__1(t),t)=1*x__1(t)+0*x__2(t)+0*x__3(t),diff(x__2(t),t)=2*x__1(t)+1*x__2(t)-2*
```

$$x_1(t) = c_1 e^t$$

$$x_2(t) = -\frac{e^t(2c_3 \sin(2t) - 2c_2 \cos(2t) + 3c_1)}{2}$$

$$x_3(t) = e^t(c_2 \sin(2t) + c_3 \cos(2t) + c_1)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 95

```
DSolve[{x1'[t]==1*x1[t]+0*x2[t]+0*x3[t],x2'[t]==2*x1[t]+1*x2[t]-2*x3[t],x3'[t]==3*x1[t]+2*x2[t]}
```

$$x1(t) \rightarrow c_1 e^t$$

$$x2(t) \rightarrow \frac{1}{2} e^t ((3c_1 + 2c_2) \cos(2t) + 2(c_1 - c_3) \sin(2t) - 3c_1)$$

$$x3(t) \rightarrow \frac{1}{2} e^t (-2(c_1 - c_3) \cos(2t) + (3c_1 + 2c_2) \sin(2t) + 2c_1)$$

16.8 problem 8

Internal problem ID [758]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.6, Complex Eigenvalues. page 417

Problem number: 8.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -3x_1(t) + 2x_3(t)$$

$$x_2'(t) = x_1(t) - x_2(t)$$

$$x_3'(t) = -2x_1(t) - x_2(t)$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 174

```
dsolve([diff(x__1(t),t)=-3*x__1(t)+0*x__2(t)+2*x__3(t),diff(x__2(t),t)=1*x__1(t)-1*x__2(t)-0
```

$$x_1(t) = 2c_1e^{-2t} + \frac{2c_2e^{-t} \sin(t\sqrt{2})}{3} - \frac{c_2e^{-t}\sqrt{2} \cos(t\sqrt{2})}{3} \\ + \frac{2c_3e^{-t} \cos(t\sqrt{2})}{3} + \frac{c_3e^{-t}\sqrt{2} \sin(t\sqrt{2})}{3}$$

$$x_2(t) = -2c_1e^{-2t} - \frac{c_2e^{-t} \sin(t\sqrt{2})}{3} - \frac{c_2e^{-t}\sqrt{2} \cos(t\sqrt{2})}{3} \\ - \frac{c_3e^{-t} \cos(t\sqrt{2})}{3} + \frac{c_3e^{-t}\sqrt{2} \sin(t\sqrt{2})}{3}$$

$$x_3(t) = c_1e^{-2t} + c_2e^{-t} \sin(t\sqrt{2}) + c_3e^{-t} \cos(t\sqrt{2})$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 235

```
DSolve[{x1'[t]==-3*x1[t]+0*x2[t]+2*x3[t],x2'[t]==1*x1[t]-1*x2[t]-0*x3[t],x3'[t]==-2*x1[t]-1*x2[t]+0*x3[t]},x1,x2,x3,t]
```

$$x1(t) \rightarrow \frac{1}{3}e^{-2t} \left((c_1 + 2(c_2 + c_3))e^t \cos(\sqrt{2}t) - \sqrt{2}(2c_1 + c_2 - 2c_3)e^t \sin(\sqrt{2}t) + 2(c_1 - c_2 - c_3) \right)$$

$$x2(t) \rightarrow \frac{1}{6}e^{-2t} \left(2(2c_1 + c_2 - 2c_3)e^t \cos(\sqrt{2}t) + \sqrt{2}(c_1 + 2(c_2 + c_3))e^t \sin(\sqrt{2}t) + 4(-c_1 + c_2 + c_3) \right)$$

$$x3(t) \rightarrow \frac{1}{6}e^{-2t} \left(-2(c_1 - c_2 - 4c_3)e^t \cos(\sqrt{2}t) - \sqrt{2}(5c_1 + 4c_2 - 2c_3)e^t \sin(\sqrt{2}t) + 2(c_1 - c_2 - c_3) \right)$$

16.9 problem 9

Internal problem ID [759]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.6, Complex Eigenvalues. page 417

Problem number: 9.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t) - 5x_2(t)$$

$$x_2'(t) = x_1(t) - 3x_2(t)$$

With initial conditions

$$[x_1(0) = 1, x_2(0) = 1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

```
dsolve([diff(x__1(t),t) = x__1(t)-5*x__2(t), diff(x__2(t),t) = x__1(t)-3*x__2(t), x__1(0) =
```

$$x_1(t) = e^{-t}(\cos(t) - 3 \sin(t))$$

$$x_2(t) = e^{-t}(-\sin(t) + \cos(t))$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 34

```
DSolve[{x1'[t]==1*x1[t]-5*x2[t],x2'[t]==1*x1[t]-3*x2[t]},{x1[0]==1,x2[0]==1},{x1[t],x2[t]},t
```

$$x1(t) \rightarrow e^{-t}(\cos(t) - 3 \sin(t))$$

$$x2(t) \rightarrow e^{-t}(\cos(t) - \sin(t))$$

16.10 problem 10

Internal problem ID [760]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.6, Complex Eigenvalues. page 417

Problem number: 10.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -3x_1(t) + 2x_2(t)$$

$$x_2'(t) = -x_1(t) - x_2(t)$$

With initial conditions

$$[x_1(0) = 1, x_2(0) = -2]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 37

```
dsolve([diff(x__1(t),t) = -3*x__1(t)+2*x__2(t), diff(x__2(t),t) = -x__1(t)-x__2(t), x__1(0)
```

$$x_1(t) = -e^{-2t}(-\cos(t) + 5\sin(t))$$

$$x_2(t) = e^{-2t}(-3\sin(t) - 2\cos(t))$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 27

```
DSolve[{x1'[t]==-3*x1[t]+2*x2[t],x2'[t]==-1*x1[t]-1*x2[t]},{x1[0]==1,x2[0]==-1},{x1[t],x2[t]}
```

$$x1(t) \rightarrow e^{-2t}(\sin(t) + \cos(t))$$

$$x2(t) \rightarrow e^{-2t} \cos(t)$$

16.11 problem 11

Internal problem ID [761]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.6, Complex Eigenvalues. page 417

Problem number: 11.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= \frac{3x_1(t)}{4} - 2x_2(t) \\x_2'(t) &= x_1(t) - \frac{5x_2(t)}{4}\end{aligned}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 45

```
dsolve([diff(x__1(t),t)=3/4*x__1(t)-2*x__2(t),diff(x__2(t),t)=1*x__1(t)-5/4*x__2(t)], [x__1(t)
```

$$x_1(t) = e^{-\frac{t}{4}}(c_1 \cos(t) + c_2 \cos(t) + c_1 \sin(t) - c_2 \sin(t))$$

$$x_2(t) = e^{-\frac{t}{4}}(c_1 \sin(t) + c_2 \cos(t))$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 56

```
DSolve[{x1'[t]==3/4*x1[t]-2*x2[t],x2'[t]==1*x1[t]-5/4*x2[t]},{x1[t],x2[t]},t,IncludeSingular
```

$$x1(t) \rightarrow e^{-t/4}(c_1 \cos(t) + (c_1 - 2c_2) \sin(t))$$

$$x2(t) \rightarrow e^{-t/4}(c_2 \cos(t) + (c_1 - c_2) \sin(t))$$

16.12 problem 12

Internal problem ID [762]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.6, Complex Eigenvalues. page 417

Problem number: 12.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= -\frac{4x_1(t)}{5} + 2x_2(t) \\x_2'(t) &= -x_1(t) + \frac{6x_2(t)}{5}\end{aligned}$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 48

```
dsolve([diff(x__1(t),t)=-4/5*x__1(t)+2*x__2(t),diff(x__2(t),t)=-1*x__1(t)+6/5*x__2(t)], [x__1
```

$$x_1(t) = -e^{\frac{t}{5}}(\cos(t) c_1 - c_2 \cos(t) - c_1 \sin(t) - c_2 \sin(t))$$

$$x_2(t) = e^{\frac{t}{5}}(c_1 \sin(t) + c_2 \cos(t))$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 56

```
DSolve[{x1'[t]==-4/5*x1[t]+2*x2[t],x2'[t]==-1*x1[t]+6/5*x2[t]},{x1[t],x2[t]},t,IncludeSingular
```

$$x_1(t) \rightarrow e^{t/5}(c_1 \cos(t) - (c_1 - 2c_2) \sin(t))$$

$$x_2(t) \rightarrow e^{t/5}(c_2(\sin(t) + \cos(t)) - c_1 \sin(t))$$

16.13 problem 23

Internal problem ID [763]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.6, Complex Eigenvalues. page 417

Problem number: 23.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -\frac{x_1(t)}{4} + x_2(t)$$

$$x_2'(t) = -x_1(t) - \frac{x_2(t)}{4}$$

$$x_3'(t) = -\frac{x_3(t)}{4}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 47

```
dsolve([diff(x__1(t),t)=-1/4*x__1(t)+1*x__2(t)+0*x__3(t),diff(x__2(t),t)=-1*x__1(t)-1/4*x__2(t),diff(x__3(t),t)=-1/4*x__3(t)),x__1(t),x__2(t),x__3(t))
```

$$x_1(t) = -e^{-\frac{t}{4}}(\cos(t) c_1 - c_2 \sin(t))$$

$$x_2(t) = e^{-\frac{t}{4}}(c_1 \sin(t) + c_2 \cos(t))$$

$$x_3(t) = c_3 e^{-\frac{t}{4}}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 110

```
DSolve[{x1'[t]==-1/4*x1[t]+1*x2[t]+0*x3[t],x2'[t]==-1*x1[t]-1/4*x2[t]+0*x3[t],x3'[t]==0*x1[t]
```

$$x1(t) \rightarrow e^{-t/4}(c_1 \cos(t) + c_2 \sin(t))$$

$$x2(t) \rightarrow e^{-t/4}(c_2 \cos(t) - c_1 \sin(t))$$

$$x3(t) \rightarrow c_3 e^{-t/4}$$

$$x1(t) \rightarrow e^{-t/4}(c_1 \cos(t) + c_2 \sin(t))$$

$$x2(t) \rightarrow e^{-t/4}(c_2 \cos(t) - c_1 \sin(t))$$

$$x3(t) \rightarrow 0$$

16.14 problem 24

Internal problem ID [764]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.6, Complex Eigenvalues. page 417

Problem number: 24.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -\frac{x_1(t)}{4} + x_2(t)$$

$$x_2'(t) = -x_1(t) - \frac{x_2(t)}{4}$$

$$x_3'(t) = \frac{x_3(t)}{10}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 47

```
dsolve([diff(x__1(t),t)=-1/4*x__1(t)+1*x__2(t)+0*x__3(t),diff(x__2(t),t)=-1*x__1(t)-1/4*x__2(t),diff(x__3(t),t)=x__3(t)/10],x__1(t),x__2(t),x__3(t))
```

$$x_1(t) = -e^{-\frac{t}{4}}(\cos(t) c_1 - c_2 \sin(t))$$

$$x_2(t) = e^{-\frac{t}{4}}(c_1 \sin(t) + c_2 \cos(t))$$

$$x_3(t) = c_3 e^{\frac{t}{10}}$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 110

```
DSolve[{x1'[t]==-1/4*x1[t]+1*x2[t]+0*x3[t],x2'[t]==-1*x1[t]-1/4*x2[t]+0*x3[t],x3'[t]==0*x1[t]
```

$$x1(t) \rightarrow e^{-t/4}(c_1 \cos(t) + c_2 \sin(t))$$

$$x2(t) \rightarrow e^{-t/4}(c_2 \cos(t) - c_1 \sin(t))$$

$$x3(t) \rightarrow c_3 e^{t/10}$$

$$x1(t) \rightarrow e^{-t/4}(c_1 \cos(t) + c_2 \sin(t))$$

$$x2(t) \rightarrow e^{-t/4}(c_2 \cos(t) - c_1 \sin(t))$$

$$x3(t) \rightarrow 0$$

16.15 problem 25

Internal problem ID [765]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.6, Complex Eigenvalues. page 417

Problem number: 25.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= -\frac{x_1(t)}{2} - \frac{x_2(t)}{8} \\x_2'(t) &= 2x_1(t) - \frac{x_2(t)}{2}\end{aligned}$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 46

```
dsolve([diff(x__1(t),t)=-1/2*x__1(t)-1/8*x__2(t),diff(x__2(t),t)=2*x__1(t)-1/2*x__2(t)], [x__
```

$$x_1(t) = \frac{e^{-\frac{t}{2}} \left(\cos\left(\frac{t}{2}\right) c_1 - \sin\left(\frac{t}{2}\right) c_2 \right)}{4}$$

$$x_2(t) = e^{-\frac{t}{2}} \left(c_2 \cos\left(\frac{t}{2}\right) + c_1 \sin\left(\frac{t}{2}\right) \right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 68

```
DSolve[{x1'[t]==-1/2*x1[t]-1/8*x2[t],x2'[t]==2*x1[t]-1/2*x2[t]},{x1[t],x2[t]},t,IncludeSingu
```

$$x1(t) \rightarrow \frac{1}{4} e^{-t/2} \left(4c_1 \cos\left(\frac{t}{2}\right) - c_2 \sin\left(\frac{t}{2}\right) \right)$$

$$x2(t) \rightarrow e^{-t/2} \left(c_2 \cos\left(\frac{t}{2}\right) + 4c_1 \sin\left(\frac{t}{2}\right) \right)$$

17 Chapter 7.8, Repeated Eigenvalues. page 436

17.1 problem 1	369
17.2 problem 2	370
17.3 problem 3	371
17.4 problem 4	372
17.5 problem 5	373
17.6 problem 6	375
17.7 problem 7	377
17.8 problem 8	378
17.9 problem 9	379
17.10problem 10	381
17.11problem 11	382
17.12problem 12	384

17.1 problem 1

Internal problem ID [766]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.8, Repeated Eigenvalues. page 436

Problem number: 1.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 3x_1(t) - 4x_2(t)$$

$$x_2'(t) = x_1(t) - x_2(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve([diff(x__1(t),t)=3*x__1(t)-4*x__2(t),diff(x__2(t),t)=1*x__1(t)-1*x__2(t)],[x__1(t), x
```

$$x_1(t) = e^t(2c_2t + 2c_1 + c_2)$$

$$x_2(t) = e^t(c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 41

```
DSolve[{x1'[t]==3*x1[t]-4*x2[t],x2'[t]==1*x1[t]-1*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x1(t) \rightarrow e^t(2c_1t - 4c_2t + c_1)$$

$$x2(t) \rightarrow e^t((c_1 - 2c_2)t + c_2)$$

17.2 problem 2

Internal problem ID [767]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.8, Repeated Eigenvalues. page 436

Problem number: 2.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 4x_1(t) - 2x_2(t)$$

$$x_2'(t) = 8x_1(t) - 4x_2(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 24

```
dsolve([diff(x__1(t),t)=4*x__1(t)-2*x__2(t),diff(x__2(t),t)=8*x__1(t)-4*x__2(t)],[x__1(t), x
```

$$x_1(t) = \frac{1}{8}c_1 + \frac{1}{2}c_1t + \frac{1}{2}c_2$$

$$x_2(t) = c_1t + c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 34

```
DSolve[{x1'[t]==4*x1[t]-2*x2[t],x2'[t]==8*x1[t]-4*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x1(t) \rightarrow 4c_1t - 2c_2t + c_1$$

$$x2(t) \rightarrow 8c_1t - 4c_2t + c_2$$

17.3 problem 3

Internal problem ID [768]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.8, Repeated Eigenvalues. page 436

Problem number: 3.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= -\frac{3x_1(t)}{2} + x_2(t) \\x_2'(t) &= -\frac{x_1(t)}{4} - \frac{x_2(t)}{2}\end{aligned}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

```
dsolve([diff(x__1(t),t)=-3/2*x__1(t)+1*x__2(t),diff(x__2(t),t)=-1/4*x__1(t)-1/2*x__2(t)], [x__
```

$$x_1(t) = 2e^{-t}(c_2t + c_1 - 2c_2)$$

$$x_2(t) = e^{-t}(c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 54

```
DSolve[{x1'[t]==-3/2*x1[t]+1*x2[t],x2'[t]==-1/4*x1[t]-1/2*x2[t]},{x1[t],x2[t]},t,IncludeSing
```

$$x1(t) \rightarrow \frac{1}{2}e^{-t}(2c_2t - c_1(t - 2))$$

$$x2(t) \rightarrow \frac{1}{4}e^{-t}(c_1(-t) + 2c_2t + 4c_2)$$

17.4 problem 4

Internal problem ID [769]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.8, Repeated Eigenvalues. page 436

Problem number: 4.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= -3x_1(t) + \frac{5x_2(t)}{2} \\x_2'(t) &= -\frac{5x_1(t)}{2} + 2x_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve([diff(x__1(t),t)=-3*x__1(t)+5/2*x__2(t),diff(x__2(t),t)=-5/2*x__1(t)+2*x__2(t)], [x__1
```

$$x_1(t) = \frac{e^{-\frac{t}{2}}(5c_2t + 5c_1 - 2c_2)}{5}$$

$$x_2(t) = e^{-\frac{t}{2}}(c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 59

```
DSolve[{x1'[t]==-3*x1[t]+5/2*x2[t],x2'[t]==-5/2*x1[t]+2*x2[t]},{x1[t],x2[t]},t,IncludeSingular
```

$$x_1(t) \rightarrow \frac{1}{2}e^{-t/2}(c_1(2 - 5t) + 5c_2t)$$

$$x_2(t) \rightarrow \frac{1}{2}e^{-t/2}(-5c_1t + 5c_2t + 2c_2)$$

17.5 problem 5

Internal problem ID [770]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.8, Repeated Eigenvalues. page 436

Problem number: 5.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t) + x_2(t) + x_3(t)$$

$$x_2'(t) = 2x_1(t) + x_2(t) - x_3(t)$$

$$x_3'(t) = -x_2(t) + x_3(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 76

```
dsolve([diff(x__1(t),t)=1*x__1(t)+1*x__2(t)+1*x__3(t),diff(x__2(t),t)=2*x__1(t)+1*x__2(t)-1*
```

$$x_1(t) = -\frac{3e^{-t}c_1}{2} - c_3e^{2t}$$

$$x_2(t) = 2e^{-t}c_1 - c_2e^{2t} - e^{2t}c_3t - c_3e^{2t}$$

$$x_3(t) = e^{-t}c_1 + c_2e^{2t} + e^{2t}c_3t$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 164

```
DSolve[{x1'[t]==1*x1[t]+1*x2[t]+1*x3[t],x2'[t]==2*x1[t]+1*x2[t]-1*x3[t],x3'[t]==0*x1[t]-1*x2[t]}
```

$$x1(t) \rightarrow \frac{1}{3}e^{-t}(c_1(2e^{3t} + 1) + (c_2 + c_3)(e^{3t} - 1))$$

$$x2(t) \rightarrow \frac{1}{9}e^{-t}(c_1(e^{3t}(6t + 4) - 4) + c_2(e^{3t}(3t + 5) + 4) + c_3(e^{3t}(3t - 4) + 4))$$

$$x3(t) \rightarrow \frac{1}{9}e^{-t}(c_1(e^{3t}(2 - 6t) - 2) + c_2(2 - e^{3t}(3t + 2)) - c_3(e^{3t}(3t - 7) - 2))$$

17.6 problem 6

Internal problem ID [771]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.8, Repeated Eigenvalues. page 436

Problem number: 6.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_2(t) + x_3(t)$$

$$x_2'(t) = x_1(t) + x_3(t)$$

$$x_3'(t) = x_1(t) + x_2(t)$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 64

```
dsolve([diff(x__1(t),t)=0*x__1(t)+1*x__2(t)+1*x__3(t),diff(x__2(t),t)=1*x__1(t)+0*x__2(t)+1*
```

$$x_1(t) = c_2 e^{2t} - 2c_3 e^{-t} - e^{-t} c_1$$

$$x_2(t) = c_2 e^{2t} + c_3 e^{-t} + e^{-t} c_1$$

$$x_3(t) = c_2 e^{2t} + c_3 e^{-t}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 124

```
DSolve[{x1'[t]==0*x1[t]+1*x2[t]+1*x3[t],x2'[t]==1*x1[t]+0*x2[t]+1*x3[t],x3'[t]==1*x1[t]+1*x2[t]}
```

$$x1(t) \rightarrow \frac{1}{3}e^{-t}(c_1(e^{3t} + 2) + (c_2 + c_3)(e^{3t} - 1))$$

$$x2(t) \rightarrow \frac{1}{3}e^{-t}(c_1(e^{3t} - 1) + c_2(e^{3t} + 2) + c_3(e^{3t} - 1))$$

$$x3(t) \rightarrow \frac{1}{3}e^{-t}(c_1(e^{3t} - 1) + c_2(e^{3t} - 1) + c_3(e^{3t} + 2))$$

17.7 problem 7

Internal problem ID [772]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.8, Repeated Eigenvalues. page 436

Problem number: 7.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= x_1(t) - 4x_2(t) \\x_2'(t) &= 4x_1(t) - 7x_2(t)\end{aligned}$$

With initial conditions

$$[x_1(0) = 3, x_2(0) = 2]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 29

```
dsolve([diff(x__1(t),t) = x__1(t)-4*x__2(t), diff(x__2(t),t) = 4*x__1(t)-7*x__2(t), x__1(0)
```

$$x_1(t) = \frac{e^{-3t}(16t + 12)}{4}$$

$$x_2(t) = e^{-3t}(4t + 2)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 34

```
DSolve[{x1'[t]==1*x1[t]-4*x2[t],x2'[t]==1*x1[t]-4*x2[t]},{x1[0]==3,x2[0]==2},{x1[t],x2[t]},t
```

$$x1(t) \rightarrow \frac{5e^{-3t}}{3} + \frac{4}{3}$$

$$x2(t) \rightarrow \frac{5e^{-3t}}{3} + \frac{1}{3}$$

17.8 problem 8

Internal problem ID [773]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.8, Repeated Eigenvalues. page 436

Problem number: 8.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= -\frac{5x_1(t)}{2} + \frac{3x_2(t)}{2} \\x_2'(t) &= -\frac{3x_1(t)}{2} + \frac{x_2(t)}{2}\end{aligned}$$

With initial conditions

$$[x_1(0) = 3, x_2(0) = -1]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 29

```
dsolve([diff(x__1(t),t) = -5/2*x__1(t)+3/2*x__2(t), diff(x__2(t),t) = -3/2*x__1(t)+1/2*x__2(t)], [x__1(0)=3, x__2(0)=-1])
```

$$x_1(t) = \frac{e^{-t}(-18t + 9)}{3}$$

$$x_2(t) = e^{-t}(-6t - 1)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 31

```
DSolve[{x1'[t]==-5/2*x1[t]+3/2*x2[t], x2'[t]==-3/2*x1[t]+1/2*x2[t]}, {x1[0]==3, x2[0]==-1}, {x1[t], x2[t]}
```

$$x1(t) \rightarrow e^{-t}(3 - 6t)$$

$$x2(t) \rightarrow -e^{-t}(6t + 1)$$

17.9 problem 9

Internal problem ID [774]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.8, Repeated Eigenvalues. page 436

Problem number: 9.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 2x_1(t) + \frac{3x_2(t)}{2} \\x_2'(t) &= -\frac{3x_1(t)}{2} - x_2(t)\end{aligned}$$

With initial conditions

$$[x_1(0) = 3, x_2(0) = -2]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 29

```
dsolve([diff(x__1(t),t) = 2*x__1(t)+3/2*x__2(t), diff(x__2(t),t) = -3/2*x__1(t)-x__2(t), x__
```

$$x_1(t) = -\frac{e^{\frac{t}{2}}\left(-\frac{9t}{2} - 9\right)}{3}$$

$$x_2(t) = e^{\frac{t}{2}}\left(-\frac{3t}{2} - 2\right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 38

```
DSolve[{x1'[t]==2*x1[t]+3/2*x2[t],x2'[t]==-3/2*x1[t]-1*x2[t]},{x1[0]==3,x2[0]==-2},{x1[t],x2[t]}
```

$$x_1(t) \rightarrow \frac{3}{2}e^{t/2}(t+2)$$

$$x_2(t) \rightarrow -\frac{1}{2}e^{t/2}(3t+4)$$

17.10 problem 10

Internal problem ID [775]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.8, Repeated Eigenvalues. page 436

Problem number: 10.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 3x_1(t) + 9x_2(t)$$

$$x_2'(t) = -x_1(t) - 3x_2(t)$$

With initial conditions

$$[x_1(0) = 2, x_2(0) = 4]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 18

```
dsolve([diff(x__1(t),t) = 3*x__1(t)+9*x__2(t), diff(x__2(t),t) = -x__1(t)-3*x__2(t), x__1(0)
```

$$x_1(t) = 42t + 2$$

$$x_2(t) = -14t + 4$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 18

```
DSolve[{x1'[t]==3*x1[t]+9*x2[t],x2'[t]==-1*x1[t]-3*x2[t]},{x1[0]==2,x2[0]==4},{x1[t],x2[t]},
```

$$x1(t) \rightarrow 42t + 2$$

$$x2(t) \rightarrow 4 - 14t$$

17.11 problem 11

Internal problem ID [776]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.8, Repeated Eigenvalues. page 436

Problem number: 11.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t)$$

$$x_2'(t) = -4x_1(t) + x_2(t)$$

$$x_3'(t) = 3x_1(t) + 6x_2(t) + 2x_3(t)$$

With initial conditions

$$[x_1(0) = -1, x_2(0) = 2, x_3(0) = -30]$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 40

```
dsolve([diff(x__1(t),t) = x__1(t), diff(x__2(t),t) = -4*x__1(t)+x__2(t), diff(x__3(t),t) = 3
```

$$x_1(t) = -e^t$$

$$x_2(t) = -\frac{e^t(-192t - 96)}{48}$$

$$x_3(t) = 3e^{2t} - 33e^t - 24te^t$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 39

```
DSolve[{x1'[t]==1*x1[t]+0*x2[t]+0*x3[t],x2'[t]==-4*x1[t]+1*x2[t]+0*x3[t],x3'[t]==3*x1[t]+6*x
```

$$x1(t) \rightarrow -e^t$$

$$x2(t) \rightarrow 2e^t(2t + 1)$$

$$x3(t) \rightarrow 3e^t(-8t + e^t - 11)$$

17.12 problem 12

Internal problem ID [777]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.8, Repeated Eigenvalues. page 436

Problem number: 12.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -\frac{5x_1(t)}{2} + x_2(t) + x_3(t)$$

$$x_2'(t) = x_1(t) - \frac{5x_2(t)}{2} + x_3(t)$$

$$x_3'(t) = x_1(t) + x_2(t) - \frac{5x_3(t)}{2}$$

With initial conditions

$$[x_1(0) = 2, x_2(0) = 3, x_3(0) = -1]$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 50

```
dsolve([diff(x__1(t),t) = -5/2*x__1(t)+x__2(t)+x__3(t), diff(x__2(t),t) = x__1(t)-5/2*x__2(t)
```

$$x_1(t) = \frac{2e^{-\frac{7t}{2}}}{3} + \frac{4e^{-\frac{t}{2}}}{3}$$

$$x_2(t) = \frac{5e^{-\frac{7t}{2}}}{3} + \frac{4e^{-\frac{t}{2}}}{3}$$

$$x_3(t) = -\frac{7e^{-\frac{7t}{2}}}{3} + \frac{4e^{-\frac{t}{2}}}{3}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 71

```
DSolve[{x1'[t]==-5/2*x1[t]+1*x2[t]+1*x3[t],x2'[t]==1*x1[t]-5/2*x2[t]+1*x3[t],x3'[t]==1*x1[t]
```

$$x1(t) \rightarrow \frac{2}{3}e^{-7t/2}(2e^{3t} + 1)$$

$$x2(t) \rightarrow \frac{1}{3}e^{-7t/2}(4e^{3t} + 5)$$

$$x3(t) \rightarrow \frac{1}{3}e^{-7t/2}(4e^{3t} - 7)$$

18 Chapter 7.9, Nonhomogeneous Linear Systems.
page 447

18.1 problem 1	387
18.2 problem 2	388
18.3 problem 3	390
18.4 problem 4	391
18.5 problem 5	392
18.6 problem 6	393
18.7 problem 7	394
18.8 problem 8	395
18.9 problem 9	396
18.10problem 10	397
18.11problem 11	398
18.12problem 12	399
18.13problem 13	400
18.14problem 18	401

18.1 problem 1

Internal problem ID [778]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.9, Nonhomogeneous Linear Systems. page 447

Problem number: 1.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 2x_1(t) - x_2(t) + e^t \\x_2'(t) &= 3x_1(t) - 2x_2(t) + t\end{aligned}$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 54

```
dsolve([diff(x__1(t),t)=2*x__1(t)-1*x__2(t)+exp(t),diff(x__2(t),t)=3*x__1(t)-2*x__2(t)+t],[x
```

$$x_1(t) = \frac{c_2 e^{-t}}{3} + c_1 e^t + \frac{3t e^t}{2} - \frac{e^t}{4} + t$$

$$x_2(t) = c_2 e^{-t} + c_1 e^t + \frac{3t e^t}{2} - \frac{3e^t}{4} + 2t - 1$$

✓ Solution by Mathematica

Time used: 0.123 (sec). Leaf size: 97

```
DSolve[{x1'[t]==2*x1[t]-1*x2[t]+Exp[t],x2'[t]==3*x1[t]-2*x2[t]+t},{x1[t],x2[t]},t,IncludeSin
```

$$x_1(t) \rightarrow \frac{1}{4} e^{-t} (4e^t t + e^{2t} (6t - 1 + 6c_1 - 2c_2) - 2c_1 + 2c_2)$$

$$x_2(t) \rightarrow \frac{1}{4} e^{-t} (e^t (8t - 4) + e^{2t} (6t - 3 + 6c_1 - 2c_2) - 6c_1 + 6c_2)$$

18.2 problem 2

Internal problem ID [779]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.9, Nonhomogeneous Linear Systems. page 447

Problem number: 2.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= x_1(t) + \sqrt{3}x_2(t) + e^t \\x_2'(t) &= \sqrt{3}x_1(t) - x_2(t) + \sqrt{3}e^{-t}\end{aligned}$$

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 67

```
dsolve([diff(x__1(t),t)=1*x__1(t)+sqrt(3)*x__2(t)+exp(t),diff(x__2(t),t)=sqrt(3)*x__1(t)-1*x__2(t)+sqrt(3)*exp(-t)],t)
```

$$x_1(t) = e^{2t}\sqrt{3}c_2 - \frac{e^{-2t}\sqrt{3}c_1}{3} - \frac{2e^t}{3} - e^{-t}$$

$$x_2(t) = c_2e^{2t} + c_1e^{-2t} + \frac{2\sqrt{3}e^{-t}}{3} - \frac{e^t\sqrt{3}}{3}$$

✓ Solution by Mathematica

Time used: 2.501 (sec). Leaf size: 313

`DSolve[{x1'[t]==1*x1[t]+Sqrt[4]*x2[t]+Exp[t],x2'[t]==Sqrt[3]*x1[t]-1*x2[t]+Sqrt[3]*Exp[-t]},`

$$x1(t) \rightarrow \frac{1}{6} \left(-6e^{-t} - \frac{2(6 + \sqrt{3}) e^t}{1 + 2\sqrt{3}} + \frac{(3(\sqrt{1 + 2\sqrt{3}} - 1) c_1 - 6c_2) e^{-\sqrt{1+2\sqrt{3}}t}}{\sqrt{1 + 2\sqrt{3}}} \right. \\ \left. + \frac{3((1 + \sqrt{1 + 2\sqrt{3}}) c_1 + 2c_2) e^{\sqrt{1+2\sqrt{3}}t}}{\sqrt{1 + 2\sqrt{3}}} \right)$$

$$x2(t) \rightarrow \frac{1}{4} \left(4e^{-t} - 2e^t + \frac{2((6 + \sqrt{3}) c_1 + (1 + 2\sqrt{3})(\sqrt{1 + 2\sqrt{3}} - 1) c_2) e^{\sqrt{1+2\sqrt{3}}t}}{(1 + 2\sqrt{3})^{3/2}} \right. \\ \left. + \frac{(2(1 + 2\sqrt{3})(1 + \sqrt{1 + 2\sqrt{3}}) c_2 - 2(6 + \sqrt{3}) c_1) e^{-\sqrt{1+2\sqrt{3}}t}}{(1 + 2\sqrt{3})^{3/2}} \right)$$

18.3 problem 3

Internal problem ID [780]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.9, Nonhomogeneous Linear Systems. page 447

Problem number: 3.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 2x_1(t) - 5x_2(t) - \cos(t) \\x_2'(t) &= x_1(t) - 2x_2(t) + \sin(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 59

```
dsolve([diff(x__1(t),t)=2*x__1(t)-5*x__2(t)-cos(t),diff(x__2(t),t)=1*x__1(t)-2*x__2(t)+sin(t)
```

$$x_1(t) = c_2 \cos(t) - c_1 \sin(t) - \sin(t)t + 2c_2 \sin(t) + 2 \cos(t) c_1 - 3 \sin(t) + 2 \cos(t)t$$

$$x_2(t) = c_2 \sin(t) + \cos(t) c_1 - \sin(t) + \cos(t)t$$

✓ Solution by Mathematica

Time used: 0.069 (sec). Leaf size: 61

```
DSolve[{x1'[t]==2*x1[t]-5*x2[t]-Cos[t],x2'[t]==1*x1[t]-2*x2[t]+Sin[t]},{x1[t],x2[t]},t,Inclu
```

$$x_1(t) \rightarrow \left(2t - \frac{1}{2} + c_1\right) \cos(t) - (t - 1 - 2c_1 + 5c_2) \sin(t)$$

$$x_2(t) \rightarrow (t - 1 + c_2) \cos(t) + \frac{1}{2}(1 + 2c_1 - 4c_2) \sin(t)$$

18.4 problem 4

Internal problem ID [781]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.9, Nonhomogeneous Linear Systems. page 447

Problem number: 4.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= x_1(t) + x_2(t) + e^{-2t} \\x_2'(t) &= 4x_1(t) - 2x_2(t) - 2e^t\end{aligned}$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 45

```
dsolve([diff(x__1(t),t)=1*x__1(t)+1*x__2(t)+exp(-2*t),diff(x__2(t),t)=4*x__1(t)-2*x__2(t)-2*
```

$$x_1(t) = c_2 e^{2t} - \frac{c_1 e^{-3t}}{4} + \frac{e^t}{2}$$

$$x_2(t) = c_2 e^{2t} + c_1 e^{-3t} - e^{-2t}$$

✓ Solution by Mathematica

Time used: 0.585 (sec). Leaf size: 84

```
DSolve[{x1'[t]==1*x1[t]+1*x2[t]+Exp[-2*t],x2'[t]==4*x1[t]-2*x2[t]-2*Exp[t]},{x1[t],x2[t]},t,
```

$$x_1(t) \rightarrow \frac{e^t}{2} + \frac{1}{5}(c_1 - c_2)e^{-3t} + \frac{1}{5}(4c_1 + c_2)e^{2t}$$

$$x_2(t) \rightarrow \frac{1}{5}e^{-3t}(-5e^t + (4c_1 + c_2)e^{5t} - 4c_1 + 4c_2)$$

18.5 problem 5

Internal problem ID [782]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.9, Nonhomogeneous Linear Systems. page 447

Problem number: 5.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 4x_1(t) - 2x_2(t) + \frac{1}{t^3} \\x_2'(t) &= 8x_1(t) - 4x_2(t) - \frac{1}{t^2}\end{aligned}$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 47

```
dsolve([diff(x__1(t),t)=4*x__1(t)-2*x__2(t)+1/(t^3),diff(x__2(t),t)=8*x__1(t)-4*x__2(t)-1/(t
```

$$x_1(t) = \frac{c_1 t}{2} - 2 \ln(t) + \frac{c_1}{8} + \frac{c_2}{2} + \frac{2}{t} - \frac{1}{2t^2}$$

$$x_2(t) = \frac{5}{t} - 4 \ln(t) + c_1 t + c_2$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 61

```
DSolve[{x1'[t]==4*x1[t]-2*x2[t]+1/(t^3),x2'[t]==8*x1[t]-4*x2[t]-1/(t^2)},{x1[t],x2[t]},t,Inc
```

$$x_1(t) \rightarrow -\frac{1}{2t^2} + \frac{2}{t} - 2 \log(t) + 4c_1 t - 2c_2 t - 2 + c_1$$

$$x_2(t) \rightarrow \frac{5}{t} - 4 \log(t) + 8c_1 t - 4c_2 t - 4 + c_2$$

18.6 problem 6

Internal problem ID [783]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.9, Nonhomogeneous Linear Systems. page 447

Problem number: 6.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= -4x_1(t) + 2x_2(t) + \frac{1}{t} \\x_2'(t) &= 2x_1(t) - x_2(t) + \frac{2}{t} + 4\end{aligned}$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 45

```
dsolve([diff(x__1(t),t)=-4*x__1(t)+2*x__2(t)+1/t,diff(x__2(t),t)=2*x__1(t)-1*x__2(t)+2/t+4],
```

$$x_1(t) = \frac{2e^{-5t}c_1}{5} + \ln(-5t) + \frac{c_2}{2} + \frac{8t}{5} - \frac{2}{5}$$

$$x_2(t) = 2 \ln(-5t) - \frac{e^{-5t}c_1}{5} + \frac{16t}{5} + c_2$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 86

```
DSolve[{x1'[t]==-4*x1[t]+2*x2[t]+1/t,x2'[t]==2*x1[t]-1*x2[t]+2/t+4},{x1[t],x2[t]},t,IncludeS
```

$$x_1(t) \rightarrow \frac{1}{25} (40t + 25 \log(t) + 20c_1 e^{-5t} - 10c_2 e^{-5t} - 8 + 5c_1 + 10c_2)$$

$$x_2(t) \rightarrow \frac{1}{25} (80t + 50 \log(t) - 10c_1 e^{-5t} + 5c_2 e^{-5t} + 4 + 10c_1 + 20c_2)$$

18.7 problem 7

Internal problem ID [784]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.9, Nonhomogeneous Linear Systems. page 447

Problem number: 7.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= x_1(t) + x_2(t) + 2e^t \\x_2'(t) &= 4x_1(t) + x_2(t) - e^t\end{aligned}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 44

```
dsolve([diff(x__1(t),t)=1*x__1(t)+1*x__2(t)+2*exp(t),diff(x__2(t),t)=4*x__1(t)+1*x__2(t)-exp
```

$$x_1(t) = -\frac{c_2 e^{-t}}{2} + \frac{c_1 e^{3t}}{2} + \frac{e^t}{4}$$

$$x_2(t) = c_2 e^{-t} + c_1 e^{3t} - 2e^t$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 80

```
DSolve[{x1'[t]==1*x1[t]+1*x2[t]+2*Exp[t],x2'[t]==4*x1[t]+1*x2[t]-Exp[t]},{x1[t],x2[t]},t,Inc
```

$$x_1(t) \rightarrow \frac{1}{4} e^{-t} (e^{2t} + (2c_1 + c_2) e^{4t} + 2c_1 - c_2)$$

$$x_2(t) \rightarrow \frac{1}{2} e^{-t} (-4e^{2t} + (2c_1 + c_2) e^{4t} - 2c_1 + c_2)$$

18.8 problem 8

Internal problem ID [785]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.9, Nonhomogeneous Linear Systems. page 447

Problem number: 8.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 2x_1(t) - x_2(t) + e^t \\x_2'(t) &= 3x_1(t) - 2x_2(t) - e^t\end{aligned}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 43

```
dsolve([diff(x__1(t),t)=2*x__1(t)-1*x__2(t)+exp(t),diff(x__2(t),t)=3*x__1(t)-2*x__2(t)-exp(t)
```

$$x_1(t) = \frac{c_2 e^{-t}}{3} + c_1 e^t + e^t + 2t e^t$$

$$x_2(t) = c_2 e^{-t} + c_1 e^t + 2t e^t$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 80

```
DSolve[{x1'[t]==2*x1[t]-1*x2[t]+Exp[t],x2'[t]==3*x1[t]-2*x2[t]-Exp[t]},{x1[t],x2[t]},t,Inclu
```

$$x_1(t) \rightarrow \frac{1}{2} e^{-t} (e^{2t} (4t - 1 + 3c_1 - c_2) - c_1 + c_2)$$

$$x_2(t) \rightarrow \frac{1}{2} e^{-t} (e^{2t} (4t - 3 + 3c_1 - c_2) - 3c_1 + 3c_2)$$

18.9 problem 9

Internal problem ID [786]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.9, Nonhomogeneous Linear Systems. page 447

Problem number: 9.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= -\frac{5x_1(t)}{4} + \frac{3x_2(t)}{4} + 2t \\x_2'(t) &= \frac{3x_1(t)}{4} - \frac{5x_2(t)}{4} + e^t\end{aligned}$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 51

```
dsolve([diff(x__1(t),t)=-5/4*x__1(t)+3/4*x__2(t)+2*t,diff(x__2(t),t)=3/4*x__1(t)-5/4*x__2(t)
```

$$x_1(t) = c_2 e^{-\frac{t}{2}} - c_1 e^{-2t} - \frac{17}{4} + \frac{e^t}{6} + \frac{5t}{2}$$

$$x_2(t) = c_2 e^{-\frac{t}{2}} + c_1 e^{-2t} + \frac{3t}{2} + \frac{e^t}{2} - \frac{15}{4}$$

✓ Solution by Mathematica

Time used: 0.349 (sec). Leaf size: 101

```
DSolve[{x1'[t]==-5/4*x1[t]+3/4*x2[t]+2*t,x2'[t]==3/4*x1[t]-5/4*x2[t]+Exp[t]},{x1[t],x2[t]},t
```

$$x_1(t) \rightarrow \frac{1}{12}(30t + 2e^t + 6(c_1 - c_2)e^{-2t} + 6(c_1 + c_2)e^{-t/2} - 51)$$

$$x_2(t) \rightarrow \frac{1}{4}e^{-2t}(3e^{2t}(2t - 5) + 2e^{3t} + 2(c_1 + c_2)e^{3t/2} - 2c_1 + 2c_2)$$

18.10 problem 10

Internal problem ID [787]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.9, Nonhomogeneous Linear Systems. page 447

Problem number: 10.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= -3x_1(t) + \sqrt{2}x_2(t) + e^{-t} \\x_2'(t) &= \sqrt{2}x_1(t) - 2x_2(t) - e^{-t}\end{aligned}$$

✓ Solution by Maple

Time used: 0.109 (sec). Leaf size: 91

```
dsolve([diff(x__1(t),t)=-3*x__1(t)+sqrt(2)*x__2(t)+exp(-t),diff(x__2(t),t)=sqrt(2)*x__1(t)-2
```

$$x_1(t) = \frac{te^{-t}}{3} + \frac{e^{-t}}{3} + \frac{e^{-t}\sqrt{2}c_1}{2} - \frac{te^{-t}\sqrt{2}}{3} - \sqrt{2}e^{-4t}c_2 + \frac{\sqrt{2}e^{-t}}{6}$$

$$x_2(t) = e^{-4t}c_2 + e^{-t}c_1 + \frac{te^{-t}\sqrt{2}}{3} - \frac{2te^{-t}}{3}$$

✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 128

```
DSolve[{x1'[t]==-3*x1[t]+Sqrt[2]*x2[t]+Exp[-t],x2'[t]==Sqrt[2]*x1[t]-2*x2[t]-Exp[-t]},{x1[t]
```

$$x_1(t) \rightarrow \frac{1}{9}e^{-4t} \left(e^{3t} \left(-3(\sqrt{2}-1)t + \sqrt{2} + 2 + 3c_1 + 3\sqrt{2}c_2 \right) + 6c_1 - 3\sqrt{2}c_2 \right)$$

$$x_2(t) \rightarrow \frac{1}{9}e^{-4t} \left(e^{3t} \left(3(\sqrt{2}-2)t - \sqrt{2} - 1 + 3\sqrt{2}c_1 + 6c_2 \right) - 3\sqrt{2}c_1 + 3c_2 \right)$$

18.11 problem 11

Internal problem ID [788]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.9, Nonhomogeneous Linear Systems. page 447

Problem number: 11.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 2x_1(t) - 5x_2(t) \\x_2'(t) &= x_1(t) - 2x_2(t) + \cos(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.109 (sec). Leaf size: 60

```
dsolve([diff(x__1(t),t)=2*x__1(t)-5*x__2(t)+0,diff(x__2(t),t)=1*x__1(t)-2*x__2(t)+cos(t)], [x
```

$$x_1(t) = 2 \cos(t) c_1 + c_2 \cos(t) - c_1 \sin(t) + 2c_2 \sin(t) - \frac{5 \sin(t) t}{2} - \frac{5 \cos(t)}{2}$$

$$x_2(t) = c_2 \sin(t) + \cos(t) c_1 + \frac{\cos(t) t}{2} - \cos(t) - \sin(t) t$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 60

```
DSolve[{x1'[t]==2*x1[t]-5*x2[t]+0,x2'[t]==1*x1[t]-2*x2[t]-Cos[t]},{x1[t],x2[t]},t,IncludeSin
```

$$x_1(t) \rightarrow \left(\frac{5}{2} + c_1\right) \cos(t) + \frac{1}{2}(5t + 4c_1 - 10c_2) \sin(t)$$

$$x_2(t) \rightarrow \left(-\frac{t}{2} + 1 + c_2\right) \cos(t) + (t + c_1 - 2c_2) \sin(t)$$

18.12 problem 12

Internal problem ID [789]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.9, Nonhomogeneous Linear Systems. page 447

Problem number: 12.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 2x_1(t) - 5x_2(t) + \csc(t)$$

$$x_2'(t) = x_1(t) - 2x_2(t) + \sec(t)$$

✓ Solution by Maple

Time used: 0.39 (sec). Leaf size: 113

```
dsolve([diff(x__1(t),t)=2*x__1(t)-5*x__2(t)+csc(t),diff(x__2(t),t)=1*x__1(t)-2*x__2(t)+sec(t)
```

$$\begin{aligned}x_1(t) = & -5 \ln(\cos(t)) \cos(t) + \cos(t) \ln(\sin(t)) + 2 \cos(t) c_1 \\ & + c_2 \cos(t) - 2 \cos(t) t + 2 \sin(t) \ln(\sin(t)) - c_1 \sin(t) \\ & + 2c_2 \sin(t) - 4 \sin(t) t - 2 \sin(t) - \sec(t) + \frac{\sin(t)^2}{\cos(t)}\end{aligned}$$

$$\begin{aligned}x_2(t) = & -2 \ln(\cos(t)) \cos(t) + \cos(t) c_1 - \ln(\cos(t)) \sin(t) \\ & + \sin(t) \ln(\sin(t)) + c_2 \sin(t) - 2 \sin(t) t - \sin(t)\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 79

```
DSolve[{x1'[t]==2*x1[t]-5*x2[t]+Csc[t],x2'[t]==1*x1[t]-2*x2[t]+Sec[t]},{x1[t],x2[t]},t,Inclu
```

$$\begin{aligned}x1(t) \rightarrow & \sin(t)(-4t + 2 \log(\sin(t)) + 2c_1 - 5c_2) \\ & + \cos(t)(-2t + \log(\sin(t)) - 5 \log(\cos(t)) + c_1)\end{aligned}$$

$$x2(t) \rightarrow \cos(t)(-2 \log(\cos(t)) + c_2) + \sin(t)(-2t + \log(\sin(t)) - \log(\cos(t)) + c_1 - 2c_2)$$

18.13 problem 13

Internal problem ID [790]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.9, Nonhomogeneous Linear Systems. page 447

Problem number: 13.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= -\frac{x_1(t)}{2} - \frac{x_2(t)}{8} + \frac{e^{-\frac{t}{2}}}{2} \\x_2'(t) &= 2x_1(t) - \frac{x_2(t)}{2}\end{aligned}$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 47

```
dsolve([diff(x__1(t),t)=-1/2*x__1(t)-1/8*x__2(t)+1/2*exp(-t/2),diff(x__2(t),t)=2*x__1(t)-1/2
```

$$x_1(t) = \frac{e^{-\frac{t}{2}} \left(c_2 \cos\left(\frac{t}{2}\right) - c_1 \sin\left(\frac{t}{2}\right) \right)}{4}$$

$$x_2(t) = e^{-\frac{t}{2}} \left(\cos\left(\frac{t}{2}\right) c_1 + \sin\left(\frac{t}{2}\right) c_2 + 4 \right)$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 69

```
DSolve[{x1'[t]==-1/2*x1[t]-1/8*x2[t]+1/2*Exp[-t/2],x2'[t]==2*x1[t]-1/2*x2[t]+0},{x1[t],x2[t]
```

$$x_1(t) \rightarrow \frac{1}{4} e^{-t/2} \left(4c_1 \cos\left(\frac{t}{2}\right) - c_2 \sin\left(\frac{t}{2}\right) \right)$$

$$x_2(t) \rightarrow e^{-t/2} \left(c_2 \cos\left(\frac{t}{2}\right) + 4c_1 \sin\left(\frac{t}{2}\right) + 4 \right)$$

18.14 problem 18

Internal problem ID [791]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 7.9, Nonhomogeneous Linear Systems. page 447

Problem number: 18.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= -2x_1(t) + x_2(t) + 2e^{-t} \\x_2'(t) &= x_1(t) - 2x_2(t) + 3t\end{aligned}$$

With initial conditions

$$[x_1(0) = \alpha_1, x_2(0) = \alpha_2]$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 93

```
dsolve([diff(x__1(t),t) = -2*x__1(t)+x__2(t)+2*exp(-t), diff(x__2(t),t) = x__1(t)-2*x__2(t)+
```

$$x_1(t) = \left(\frac{3}{2} + \frac{\alpha_2}{2} + \frac{\alpha_1}{2}\right) e^{-t} - \left(\frac{2}{3} + \frac{\alpha_2}{2} - \frac{\alpha_1}{2}\right) e^{-3t} + \frac{e^{-t}}{2} + t e^{-t} - \frac{4}{3} + t$$

$$x_2(t) = \left(\frac{3}{2} + \frac{\alpha_2}{2} + \frac{\alpha_1}{2}\right) e^{-t} + \left(\frac{2}{3} + \frac{\alpha_2}{2} - \frac{\alpha_1}{2}\right) e^{-3t} + t e^{-t} + 2t - \frac{5}{3} - \frac{e^{-t}}{2}$$

✓ Solution by Mathematica

Time used: 0.061 (sec). Leaf size: 122

```
DSolve[{x1'[t]==-2*x1[t]+1*x2[t]+2*Exp[-t],x2'[t]==1*x1[t]-2*x2[t]+3*t},{x1[0]==a1,x2[0]==a2}
```

$$x1(t) \rightarrow \frac{1}{6}e^{-3t}(3a1(e^{2t} + 1) + 3a2(e^{2t} - 1) + 12e^{2t} - 8e^{3t} + 6e^{2t}t + 6e^{3t}t - 4)$$

$$x2(t) \rightarrow \frac{1}{6}e^{-3t}(3a1(e^{2t} - 1) + 3a2(e^{2t} + 1) + 6e^{2t}(t + 1) + 2e^{3t}(6t - 5) + 4)$$

19 Chapter 9.1, The Phase Plane: Linear Systems.
page 505

19.1 problem 1	404
19.2 problem 2	405
19.3 problem 3	406
19.4 problem 4	407
19.5 problem 5	408
19.6 problem 6	409
19.7 problem 7	410
19.8 problem 8	411
19.9 problem 9	412
19.10problem 10	413
19.11problem 11	414
19.12problem 12	416
19.13problem 13	417
19.14problem 14	418
19.15problem 15	419

19.1 problem 1

Internal problem ID [792]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.1, The Phase Plane: Linear Systems. page 505

Problem number: 1.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 3x_1(t) - 2x_2(t)$$

$$x_2'(t) = 2x_1(t) - 2x_2(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 36

```
dsolve([diff(x__1(t),t)=3*x__1(t)-2*x__2(t),diff(x__2(t),t)=2*x__1(t)-2*x__2(t)],[x__1(t), x
```

$$x_1(t) = \frac{e^{-t}c_1}{2} + 2c_2e^{2t}$$

$$x_2(t) = e^{-t}c_1 + c_2e^{2t}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 73

```
DSolve[{x1'[t]==3*x1[t]-2*x2[t],x2'[t]==2*x1[t]-2*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x_1(t) \rightarrow \frac{1}{3}e^{-t}(c_1(4e^{3t} - 1) - 2c_2(e^{3t} - 1))$$

$$x_2(t) \rightarrow \frac{1}{3}e^{-t}(2c_1(e^{3t} - 1) - c_2(e^{3t} - 4))$$

19.2 problem 2

Internal problem ID [793]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.1, The Phase Plane: Linear Systems. page 505

Problem number: 2.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 5x_1(t) - x_2(t)$$

$$x_2'(t) = 3x_1(t) + x_2(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 35

```
dsolve([diff(x__1(t),t)=5*x__1(t)-1*x__2(t),diff(x__2(t),t)=3*x__1(t)+1*x__2(t)],[x__1(t), x
```

$$x_1(t) = c_1 e^{4t} + \frac{c_2 e^{2t}}{3}$$

$$x_2(t) = c_1 e^{4t} + c_2 e^{2t}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 73

```
DSolve[{x1'[t]==5*x1[t]-1*x2[t],x2'[t]==3*x1[t]+1*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x_1(t) \rightarrow \frac{1}{2}e^{2t}(c_1(3e^{2t} - 1) - c_2(e^{2t} - 1))$$

$$x_2(t) \rightarrow \frac{1}{2}e^{2t}(3c_1(e^{2t} - 1) - c_2(e^{2t} - 3))$$

19.3 problem 3

Internal problem ID [794]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.1, The Phase Plane: Linear Systems. page 505

Problem number: 3.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 2x_1(t) - x_2(t) \\x_2'(t) &= 3x_1(t) - 2x_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 31

```
dsolve([diff(x__1(t),t)=2*x__1(t)-1*x__2(t),diff(x__2(t),t)=3*x__1(t)-2*x__2(t)],[x__1(t), x
```

$$x_1(t) = \frac{e^{-t}c_1}{3} + c_2e^t$$

$$x_2(t) = e^{-t}c_1 + c_2e^t$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 73

```
DSolve[{x1'[t]==2*x1[t]-1*x2[t],x2'[t]==3*x1[t]-2*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x_1(t) \rightarrow \frac{1}{2}e^{-t}(c_1(3e^{2t} - 1) - c_2(e^{2t} - 1))$$

$$x_2(t) \rightarrow \frac{1}{2}e^{-t}(3c_1(e^{2t} - 1) - c_2(e^{2t} - 3))$$

19.4 problem 4

Internal problem ID [795]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.1, The Phase Plane: Linear Systems. page 505

Problem number: 4.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= x_1(t) - 4x_2(t) \\x_2'(t) &= 4x_1(t) - 7x_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 33

```
dsolve([diff(x__1(t),t)=1*x__1(t)-4*x__2(t),diff(x__2(t),t)=4*x__1(t)-7*x__2(t)],[x__1(t), x
```

$$x_1(t) = \frac{e^{-3t}(4c_2t + 4c_1 + c_2)}{4}$$

$$x_2(t) = e^{-3t}(c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 46

```
DSolve[{x1'[t]==1*x1[t]-4*x2[t],x2'[t]==4*x1[t]-7*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x1(t) \rightarrow e^{-3t}(4c_1t - 4c_2t + c_1)$$

$$x2(t) \rightarrow e^{-3t}(4(c_1 - c_2)t + c_2)$$

19.5 problem 5

Internal problem ID [796]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.1, The Phase Plane: Linear Systems. page 505

Problem number: 5.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t) - 5x_2(t)$$

$$x_2'(t) = x_1(t) - 3x_2(t)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
dsolve([diff(x__1(t),t)=1*x__1(t)-5*x__2(t),diff(x__2(t),t)=1*x__1(t)-3*x__2(t)],[x__1(t), x
```

$$x_1(t) = e^{-t}(\cos(t) c_1 - c_2 \sin(t) + 2c_1 \sin(t) + 2c_2 \cos(t))$$

$$x_2(t) = e^{-t}(c_1 \sin(t) + c_2 \cos(t))$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 54

```
DSolve[{x1'[t]==1*x1[t]-5*x2[t],x2'[t]==1*x1[t]-3*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x1(t) \rightarrow e^{-t}(c_1 \cos(t) + (2c_1 - 5c_2) \sin(t))$$

$$x2(t) \rightarrow e^{-t}(c_2 \cos(t) + (c_1 - 2c_2) \sin(t))$$

19.6 problem 6

Internal problem ID [797]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.1, The Phase Plane: Linear Systems. page 505

Problem number: 6.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 2x_1(t) - 5x_2(t)$$

$$x_2'(t) = x_1(t) - 2x_2(t)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 37

```
dsolve([diff(x__1(t),t)=2*x__1(t)-5*x__2(t),diff(x__2(t),t)=1*x__1(t)-2*x__2(t)],[x__1(t), x
```

$$x_1(t) = \cos(t) c_1 - c_2 \sin(t) + 2c_1 \sin(t) + 2c_2 \cos(t)$$

$$x_2(t) = c_1 \sin(t) + c_2 \cos(t)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 41

```
DSolve[{x1'[t]==2*x1[t]-5*x2[t],x2'[t]==1*x1[t]-2*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x1(t) \rightarrow c_1(2 \sin(t) + \cos(t)) - 5c_2 \sin(t)$$

$$x2(t) \rightarrow c_2 \cos(t) + (c_1 - 2c_2) \sin(t)$$

19.7 problem 7

Internal problem ID [798]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.1, The Phase Plane: Linear Systems. page 505

Problem number: 7.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 3x_1(t) - 2x_2(t)$$

$$x_2'(t) = 4x_1(t) - x_2(t)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
dsolve([diff(x__1(t),t)=3*x__1(t)-2*x__2(t),diff(x__2(t),t)=4*x__1(t)-1*x__2(t)],[x__1(t), x
```

$$x_1(t) = \frac{e^t(c_1 \sin(2t) - c_2 \sin(2t) + c_1 \cos(2t) + c_2 \cos(2t))}{2}$$

$$x_2(t) = e^t(c_1 \sin(2t) + c_2 \cos(2t))$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 58

```
DSolve[{x1'[t]==3*x1[t]-2*x2[t],x2'[t]==4*x1[t]-1*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x_1(t) \rightarrow e^t(c_1 \cos(2t) + (c_1 - c_2) \sin(2t))$$

$$x_2(t) \rightarrow e^t(c_2 \cos(2t) + (2c_1 - c_2) \sin(2t))$$

19.8 problem 8

Internal problem ID [799]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.1, The Phase Plane: Linear Systems. page 505

Problem number: 8.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= -x_1(t) - x_2(t) \\x_2'(t) &= -\frac{5x_2(t)}{2}\end{aligned}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 28

```
dsolve([diff(x__1(t),t)=-1*x__1(t)-1*x__2(t),diff(x__2(t),t)=0*x__1(t)-25/10*x__2(t)], [x__1(t),x__2(t)])
```

$$x_1(t) = \frac{2c_2 e^{-\frac{5t}{2}}}{3} + e^{-t} c_1$$

$$x_2(t) = c_2 e^{-\frac{5t}{2}}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 47

```
DSolve[{x1'[t]==-1*x1[t]-1*x2[t],x2'[t]==0*x1[t]-25/10*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolutions->True]
```

$$x_1(t) \rightarrow \left(c_1 - \frac{2c_2}{3} \right) e^{-t} + \frac{2}{3} c_2 e^{-5t/2}$$

$$x_2(t) \rightarrow c_2 e^{-5t/2}$$

19.9 problem 9

Internal problem ID [800]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.1, The Phase Plane: Linear Systems. page 505

Problem number: 9.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 3x_1(t) - 4x_2(t)$$

$$x_2'(t) = x_1(t) - x_2(t)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 28

```
dsolve([diff(x__1(t),t)=3*x__1(t)-4*x__2(t),diff(x__2(t),t)=1*x__1(t)-1*x__2(t)],[x__1(t), x
```

$$x_1(t) = e^t(2c_2t + 2c_1 + c_2)$$

$$x_2(t) = e^t(c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 41

```
DSolve[{x1'[t]==3*x1[t]-4*x2[t],x2'[t]==1*x1[t]-1*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x1(t) \rightarrow e^t(2c_1t - 4c_2t + c_1)$$

$$x2(t) \rightarrow e^t((c_1 - 2c_2)t + c_2)$$

19.10 problem 10

Internal problem ID [801]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.1, The Phase Plane: Linear Systems. page 505

Problem number: 10.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= x_1(t) + 2x_2(t) \\x_2'(t) &= -5x_1(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 84

```
dsolve([diff(x__1(t),t)=1*x__1(t)+2*x__2(t),diff(x__2(t),t)=-5*x__1(t)-0*x__2(t)],[x__1(t),
```

$$x_1(t) = \frac{e^{\frac{t}{2}} \left(\sin\left(\frac{\sqrt{39}t}{2}\right) \sqrt{39} c_2 - \cos\left(\frac{\sqrt{39}t}{2}\right) \sqrt{39} c_1 - \sin\left(\frac{\sqrt{39}t}{2}\right) c_1 - \cos\left(\frac{\sqrt{39}t}{2}\right) c_2 \right)}{10}$$

$$x_2(t) = e^{\frac{t}{2}} \left(\sin\left(\frac{\sqrt{39}t}{2}\right) c_1 + \cos\left(\frac{\sqrt{39}t}{2}\right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 54

```
DSolve[{x1'[t]==1*x1[t]+2*x2[t],x2'[t]==-5*x1[t]-1*x2[t]},{x1[t],x2[t]},t,IncludeSingularSol
```

$$x_1(t) \rightarrow c_1 \cos(3t) + \frac{1}{3}(c_1 + 2c_2) \sin(3t)$$

$$x_2(t) \rightarrow c_2 \cos(3t) - \frac{1}{3}(5c_1 + c_2) \sin(3t)$$

19.11 problem 11

Internal problem ID [802]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.1, The Phase Plane: Linear Systems. page 505

Problem number: 11.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -x_1(t)$$

$$x_2'(t) = -x_2(t)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 20

```
dsolve([diff(x__1(t),t)=-1*x__1(t)-0*x__2(t),diff(x__2(t),t)=0*x__1(t)-1*x__2(t)],[x__1(t),
```

$$x_1(t) = e^{-t}c_1$$

$$x_2(t) = c_2e^{-t}$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 65

```
DSolve[{x1'[t]==-1*x1[t]-0*x2[t],x2'[t]==0*x1[t]-1*x2[t]},{x1[t],x2[t]},t,IncludeSingularSol
```

$$x1(t) \rightarrow c_1 e^{-t}$$

$$x2(t) \rightarrow c_2 e^{-t}$$

$$x1(t) \rightarrow c_1 e^{-t}$$

$$x2(t) \rightarrow 0$$

$$x1(t) \rightarrow 0$$

$$x2(t) \rightarrow c_2 e^{-t}$$

$$x1(t) \rightarrow 0$$

$$x2(t) \rightarrow 0$$

19.12 problem 12

Internal problem ID [803]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.1, The Phase Plane: Linear Systems. page 505

Problem number: 12.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 2x_1(t) - \frac{5x_2(t)}{2} \\x_2'(t) &= \frac{9x_1(t)}{5} - x_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 58

```
dsolve([diff(x__1(t),t)=2*x__1(t)-5/2*x__2(t),diff(x__2(t),t)=9/5*x__1(t)-1*x__2(t)], [x__1(t)
```

$$x_1(t) = \frac{5e^{\frac{t}{2}} \left(\sin\left(\frac{3t}{2}\right) c_1 - \sin\left(\frac{3t}{2}\right) c_2 + \cos\left(\frac{3t}{2}\right) c_1 + \cos\left(\frac{3t}{2}\right) c_2 \right)}{6}$$

$$x_2(t) = e^{\frac{t}{2}} \left(\sin\left(\frac{3t}{2}\right) c_1 + \cos\left(\frac{3t}{2}\right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 84

```
DSolve[{x1'[t]==2*x1[t]-5/2*x2[t],x2'[t]==9/5*x1[t]-1*x2[t]},{x1[t],x2[t]},t,IncludeSingular
```

$$x1(t) \rightarrow \frac{1}{3}e^{t/2} \left(3c_1 \cos\left(\frac{3t}{2}\right) + (3c_1 - 5c_2) \sin\left(\frac{3t}{2}\right) \right)$$

$$x2(t) \rightarrow \frac{1}{5}e^{t/2} \left(5c_2 \cos\left(\frac{3t}{2}\right) + (6c_1 - 5c_2) \sin\left(\frac{3t}{2}\right) \right)$$

19.13 problem 13

Internal problem ID [804]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.1, The Phase Plane: Linear Systems. page 505

Problem number: 13.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= x_1(t) + x_2(t) - 2 \\x_2'(t) &= x_1(t) - x_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 70

```
dsolve([diff(x__1(t),t)=1*x__1(t)+1*x__2(t)-2,diff(x__2(t),t)=1*x__1(t)-1*x__2(t)],[x__1(t),
```

$$x_1(t) = \sqrt{2} e^{t\sqrt{2}} c_2 - \sqrt{2} e^{-t\sqrt{2}} c_1 + e^{t\sqrt{2}} c_2 + e^{-t\sqrt{2}} c_1 + 1$$

$$x_2(t) = e^{t\sqrt{2}} c_2 + e^{-t\sqrt{2}} c_1 + 1$$

✓ Solution by Mathematica

Time used: 0.097 (sec). Leaf size: 160

```
DSolve[{x1'[t]==1*x1[t]+1*x2[t]-2,x2'[t]==1*x1[t]-1*x2[t]},{x1[t],x2[t]},t,IncludeSingularSo
```

$$x_1(t) \rightarrow \frac{1}{4} e^{-\sqrt{2}t} \left(4e^{\sqrt{2}t} + \left((2 + \sqrt{2}) c_1 + \sqrt{2} c_2 \right) e^{2\sqrt{2}t} - \left((\sqrt{2} - 2) c_1 \right) - \sqrt{2} c_2 \right)$$

$$x_2(t) \rightarrow \frac{1}{4} e^{-\sqrt{2}t} \left(4e^{\sqrt{2}t} + \left(\sqrt{2} c_1 - (\sqrt{2} - 2) c_2 \right) e^{2\sqrt{2}t} - \sqrt{2} c_1 + (2 + \sqrt{2}) c_2 \right)$$

19.14 problem 14

Internal problem ID [805]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.1, The Phase Plane: Linear Systems. page 505

Problem number: 14.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= -2x_1(t) + x_2(t) - 2 \\x_2'(t) &= x_1(t) - 2x_2(t) + 1\end{aligned}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 36

```
dsolve([diff(x__1(t),t)=-2*x__1(t)+1*x__2(t)-2,diff(x__2(t),t)=1*x__1(t)-2*x__2(t)+1],[x__1(t),x__2(t)])
```

$$x_1(t) = e^{-t}c_1 - c_2e^{-3t} - 1$$

$$x_2(t) = e^{-t}c_1 + c_2e^{-3t}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 72

```
DSolve[{x1'[t]==-2*x1[t]+1*x2[t]-2,x2'[t]==1*x1[t]-2*x2[t]+1},{x1[t],x2[t]},t,IncludeSingularSolutions->True]
```

$$x_1(t) \rightarrow \frac{1}{2}e^{-3t}(-2e^{3t} + (c_1 + c_2)e^{2t} + c_1 - c_2)$$

$$x_2(t) \rightarrow \frac{1}{2}e^{-3t}(c_1(e^{2t} - 1) + c_2(e^{2t} + 1))$$

19.15 problem 15

Internal problem ID [806]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.1, The Phase Plane: Linear Systems. page 505

Problem number: 15.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= -x_1(t) - x_2(t) - 1 \\x_2'(t) &= 2x_1(t) - x_2(t) + 5\end{aligned}$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 61

```
dsolve([diff(x__1(t),t)=-1*x__1(t)-1*x__2(t)-1,diff(x__2(t),t)=2*x__1(t)-1*x__2(t)+5],[x__1(t)
```

$$x_1(t) = -2 - \frac{\sqrt{2}e^{-t}(c_1 \sin(t\sqrt{2}) - c_2 \cos(t\sqrt{2}))}{2}$$

$$x_2(t) = 1 + e^{-t}(c_2 \sin(t\sqrt{2}) + \cos(t\sqrt{2})c_1)$$

✓ Solution by Mathematica

Time used: 0.414 (sec). Leaf size: 85

```
DSolve[{x1'[t]==-1*x1[t]-1*x2[t]-1,x2'[t]==2*x1[t]-1*x2[t]+5},{x1[t],x2[t]},t,IncludeSingular
```

$$x1(t) \rightarrow c_1 e^{-t} \cos(\sqrt{2}t) - \frac{c_2 e^{-t} \sin(\sqrt{2}t)}{\sqrt{2}} - 2$$

$$x2(t) \rightarrow e^{-t}(e^t + c_2 \cos(\sqrt{2}t) + \sqrt{2}c_1 \sin(\sqrt{2}t))$$

20 Chapter 9.2, Autonomous Systems and Stability. page 517

20.1 problem 1	421
20.2 problem 2 part 1	422
20.3 problem 2 part 2	423
20.4 problem 3 part 1	424
20.5 problem 3 part 2	425

20.1 problem 1

Internal problem ID [807]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.2, Autonomous Systems and Stability. page 517

Problem number: 1.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x'(t) &= -x(t) \\ y'(t) &= -2y(t)\end{aligned}$$

With initial conditions

$$[x(0) = 4, y(0) = 2]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 20

```
dsolve([diff(x(t),t) = -x(t), diff(y(t),t) = -2*y(t), x(0) = 4, y(0) = 2],[x(t), y(t)], sing
```

$$x(t) = 4e^{-t}$$

$$y(t) = 2e^{-2t}$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 22

```
DSolve[{x'[t]==-1*x[t]+0*y[t],y'[t]==-2*y[t]},{x[0]==4,y[0]==2},{x[t],y[t]},t,IncludeSingula
```

$$x(t) \rightarrow 4e^{-t}$$

$$y(t) \rightarrow 2e^{-2t}$$

20.2 problem 2 part 1

Internal problem ID [808]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.2, Autonomous Systems and Stability. page 517

Problem number: 2 part 1.

ODE order: 1.

ODE degree: 1.

Solve

$$x'(t) = -x(t)$$

$$y'(t) = 2y(t)$$

With initial conditions

$$[x(0) = 4, y(0) = 2]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve([diff(x(t),t) = -x(t), diff(y(t),t) = 2*y(t), x(0) = 4, y(0) = 2],[x(t), y(t)], sings
```

$$x(t) = 4e^{-t}$$

$$y(t) = 2e^{2t}$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 22

```
DSolve[{x'[t]==-1*x[t]+0*y[t],y'[t]==0*x[t]+2*y[t]},{x[0]==4,y[0]==2},{x[t],y[t]},t,IncludeS
```

$$x(t) \rightarrow 4e^{-t}$$

$$y(t) \rightarrow 2e^{2t}$$

20.3 problem 2 part 2

Internal problem ID [809]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.2, Autonomous Systems and Stability. page 517

Problem number: 2 part 2.

ODE order: 1.

ODE degree: 1.

Solve

$$x'(t) = -x(t)$$

$$y'(t) = 2y(t)$$

With initial conditions

$$[x(0) = 4, y(0) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve([diff(x(t),t) = -x(t), diff(y(t),t) = 2*y(t), x(0) = 4, y(0) = 0],[x(t), y(t)], sings
```

$$x(t) = 4e^{-t}$$

$$y(t) = 0$$

✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 16

```
DSolve[{x'[t]==-1*x[t]+0*y[t],y'[t]==0*x[t]+2*y[t]},{x[0]==4,y[0]==0},{x[t],y[t]},t,IncludeS
```

$$x(t) \rightarrow 4e^{-t}$$

$$y(t) \rightarrow 0$$

20.4 problem 3 part 1

Internal problem ID [810]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.2, Autonomous Systems and Stability. page 517

Problem number: 3 part 1.

ODE order: 1.

ODE degree: 1.

Solve

$$x'(t) = -y(t)$$

$$y'(t) = x(t)$$

With initial conditions

$$[x(0) = 4, y(0) = 0]$$

✓ Solution by Maple

Time used: 36.859 (sec). Leaf size: 16

```
dsolve([diff(x(t),t) = -y(t), diff(y(t),t) = x(t), x(0) = 4, y(0) = 0],[x(t), y(t)], singsol
```

$$x(t) = 4 \cos(t)$$

$$y(t) = 4 \sin(t)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 16

```
DSolve[{x'[t]==-0*x[t]-1*y[t],y'[t]==1*x[t]+0*y[t]},{x[0]==4,y[0]==0},{x[t],y[t]},t,IncludeS
```

$$x(t) \rightarrow 4 \cos(t)$$

$$y(t) \rightarrow 4 \sin(t)$$

20.5 problem 3 part 2

Internal problem ID [811]

Book: Elementary differential equations and boundary value problems, 10th ed., Boyce and DiPrima

Section: Chapter 9.2, Autonomous Systems and Stability. page 517

Problem number: 3 part 2.

ODE order: 1.

ODE degree: 1.

Solve

$$x'(t) = -y(t)$$

$$y'(t) = x(t)$$

With initial conditions

$$[x(0) = 0, y(0) = 4]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve([diff(x(t),t) = -y(t), diff(y(t),t) = x(t), x(0) = 0, y(0) = 4],[x(t), y(t)], singsol
```

$$x(t) = -4 \sin(t)$$

$$y(t) = 4 \cos(t)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 16

```
DSolve[{x'[t]==-0*x[t]-1*y[t],y'[t]==1*x[t]+0*y[t]},{x[0]==0,y[0]==4},{x[t],y[t]},t,IncludeS
```

$$x(t) \rightarrow -4 \sin(t)$$

$$y(t) \rightarrow 4 \cos(t)$$