


A Solution Manual For

**Engineering Mathematics. By
K. A. Stroud. 5th edition.
Industrial press Inc. NY. 2001**



Nasser M. Abbasi

March 3, 2024

Contents

1	Program 24. First order differential equations. Test exercise 24. page 1067	2
2	Program 24. First order differential equations. Further problems 24. page 1068	16
3	Program 25. Second order differential equations. Test Exercise 25. page 1093	77
4	Program 25. Second order differential equations. Further problems 25. page 1094	86

1 Program 24. First order differential equations.

Test exercise 24. page 1067

1.1	problem 1	3
1.2	problem 2	4
1.3	problem 3	5
1.4	problem 4	6
1.5	problem 5	7
1.6	problem 6	8
1.7	problem 7	9
1.8	problem 8	11
1.9	problem 9	12
1.10	problem 10	13
1.11	problem 11	14
1.12	problem 12	15

1.1 problem 1

Internal problem ID [5075]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Test exercise 24. page 1067

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y'x = x^2 + 2x - 3$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(x*diff(y(x),x)=x^2+2*x-3,y(x), singsol=all)
```

$$y(x) = \frac{x^2}{2} + 2x - 3 \ln(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 22

```
DSolve[x*y'[x]==x^2+2*x-3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2}{2} + 2x - 3 \log(x) + c_1$$

1.2 problem 2

Internal problem ID [5076]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Test exercise 24. page 1067

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(1+x)^2 y' - y^2 = 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve((1+x)^2*diff(y(x),x)=1+y(x)^2,y(x), singsol=all)
```

$$y(x) = \tan\left(\frac{c_1 x + c_1 - 1}{x + 1}\right)$$

✓ Solution by Mathematica

Time used: 0.264 (sec). Leaf size: 32

```
DSolve[(1+x)^2*y'[x]==1+y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\tan\left(\frac{1}{x+1} - c_1\right)$$

$$y(x) \rightarrow -i$$

$$y(x) \rightarrow i$$

1.3 problem 3

Internal problem ID [5077]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Test exercise 24. page 1067

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 2y = e^{3x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)+2*y(x)=exp(3*x),y(x), singsol=all)
```

$$y(x) = \left(\frac{e^{5x}}{5} + c_1 \right) e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 23

```
DSolve[y'[x]+2*y[x]==Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{3x}}{5} + c_1 e^{-2x}$$

1.4 problem 4

Internal problem ID [5078]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Test exercise 24. page 1067

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y'x - y = x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(x*diff(y(x),x)-y(x)=x^2,y(x), singsol=all)
```

$$y(x) = (x + c_1)x$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 11

```
DSolve[x*y'[x]-y[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(x + c_1)$$

1.5 problem 5

Internal problem ID [5079]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Test exercise 24. page 1067

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$x^2 y' = x^3 \sin(3x) + 4$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(x^2*diff(y(x),x)=x^3*sin(3*x)+4,y(x), singsol=all)
```

$$y(x) = \frac{\sin(3x)}{9} - \frac{x \cos(3x)}{3} - \frac{4}{x} + c_1$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 30

```
DSolve[x^2*y'[x]==x^3*Sin[3*x]+4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{4}{x} + \frac{1}{9} \sin(3x) - \frac{1}{3} x \cos(3x) + c_1$$

1.6 problem 6

Internal problem ID [5080]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Test exercise 24. page 1067

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$x \cos(y) y' - \sin(y) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve(x*cos(y(x))*diff(y(x),x)-sin(y(x))=0,y(x), singsol=all)
```

$$y(x) = \arcsin(c_1 x)$$

✓ Solution by Mathematica

Time used: 9.024 (sec). Leaf size: 17

```
DSolve[x*Cos[y[x]]*y'[x]-Sin[y[x]]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arcsin(e^{c_1} x)$$

$$y(x) \rightarrow 0$$

1.7 problem 7

Internal problem ID [5081]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Test exercise 24. page 1067

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _dAlembert]`

$$(x^3 + xy^2)y' - 2y^3 = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 47

```
dsolve((x^3+x*y(x)^2)*diff(y(x),x)=2*y(x)^3,y(x), singsol=all)
```

$$y(x) = \left(\frac{c_1 x}{2} - \frac{\sqrt{c_1^2 x^2 + 4}}{2} \right) x$$

$$y(x) = \left(\frac{c_1 x}{2} + \frac{\sqrt{c_1^2 x^2 + 4}}{2} \right) x$$

✓ Solution by Mathematica

Time used: 1.2 (sec). Leaf size: 83

```
DSolve[(x^3+x*y[x]^2)*y'[x]==2*y[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2}x\left(\sqrt{4 + e^{2c_1}x^2} + e^{c_1}x\right)$$

$$y(x) \rightarrow \frac{1}{2}x\left(\sqrt{4 + e^{2c_1}x^2} - e^{c_1}x\right)$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -x$$

$$y(x) \rightarrow x$$

1.8 problem 8

Internal problem ID [5082]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Test exercise 24. page 1067

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$(x^2 - 1)y' + 2yx = x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve((x^2-1)*diff(y(x),x)+2*x*y(x)=x,y(x), singsol=all)
```

$$y(x) = \frac{\frac{x^2}{2} + c_1}{(x-1)(x+1)}$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 31

```
DSolve[(x^2-1)*y'[x]+2*x*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2 + 2c_1}{2(x^2 - 1)}$$

$$y(x) \rightarrow \frac{1}{2}$$

1.9 problem 9

Internal problem ID [5083]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Test exercise 24. page 1067

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + y \tanh(x) = 2 \sinh(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)+y(x)*tanh(x)=2*sinh(x),y(x), singsol=all)
```

$$y(x) = \frac{\frac{\cosh(2x)}{2} + c_1}{\cosh(x)}$$

✓ Solution by Mathematica

Time used: 0.098 (sec). Leaf size: 20

```
DSolve[y'[x]+y[x]*Tanh[x]==2*Sinh[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \operatorname{sech}(x) (\cosh(2x) + 2c_1)$$

1.10 problem 10

Internal problem ID [5084]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Test exercise 24. page 1067

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y'x - 2y = x^3 \cos(x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve(x*diff(y(x),x)-2*y(x)=x^3*cos(x),y(x), singsol=all)
```

$$y(x) = (\sin(x) + c_1) x^2$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 14

```
DSolve[x*y'[x]-2*y[x]==x^3*Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2(\sin(x) + c_1)$$

1.11 problem 11

Internal problem ID [5085]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Test exercise 24. page 1067

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational, _Bernoulli]`

$$y' + \frac{y}{x} - y^3 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x),x)+y(x)/x=y(x)^3,y(x), singsol=all)
```

$$y(x) = \frac{1}{\sqrt{c_1 x^2 + 2x}}$$

$$y(x) = -\frac{1}{\sqrt{c_1 x^2 + 2x}}$$

✓ Solution by Mathematica

Time used: 0.375 (sec). Leaf size: 40

```
DSolve[y'[x]+y[x]/x==y[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\sqrt{x(2+c_1x)}}$$

$$y(x) \rightarrow \frac{1}{\sqrt{x(2+c_1x)}}$$

$$y(x) \rightarrow 0$$

1.12 problem 12

Internal problem ID [5086]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Test exercise 24. page 1067

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational, _Bernoulli]`

$$y'x + 3y - x^2y^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x*diff(y(x),x)+3*y(x)=x^2*y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{1}{x^2(c_1x + 1)}$$

✓ Solution by Mathematica

Time used: 0.137 (sec). Leaf size: 22

```
DSolve[x*y'[x]+3*y[x]==x^2*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{x^2 + c_1x^3}$$

$$y(x) \rightarrow 0$$

2 Program 24. First order differential equations. Further problems 24. page 1068

2.1	problem 1	18
2.2	problem 2	19
2.3	problem 3	20
2.4	problem 4	22
2.5	problem 5	23
2.6	problem 6	24
2.7	problem 7	26
2.8	problem 8	27
2.9	problem 9	29
2.10	problem 10	31
2.11	problem 11	33
2.12	problem 12	34
2.13	problem 13	35
2.14	problem 14	36
2.15	problem 15	37
2.16	problem 16	38
2.17	problem 17	39
2.18	problem 18	40
2.19	problem 19	41
2.20	problem 20	42
2.21	problem 21	43
2.22	problem 22	44
2.23	problem 23	46
2.24	problem 24	47
2.25	problem 25	48
2.26	problem 26	49
2.27	problem 27	50
2.28	problem 28	52
2.29	problem 29	53
2.30	problem 30	54
2.31	problem 31	55
2.32	problem 32	56
2.33	problem 33	59
2.34	problem 34	60
2.35	problem 35	61
2.36	problem 36	62

2.37	problem 37	63
2.38	problem 38	64
2.39	problem 39	65
2.40	problem 40	66
2.41	problem 41	67
2.42	problem 42	68
2.43	problem 43	69
2.44	problem 44	70
2.45	problem 45	71
2.46	problem 46	72
2.47	problem 47	73
2.48	problem 48	75
2.49	problem 49	76

2.1 problem 1

Internal problem ID [5087]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$x(y - 3)y' - 4y = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 25

```
dsolve(x*(y(x)-3)*diff(y(x),x)=4*y(x),y(x), singsol=all)
```

$$y(x) = e^{-\text{LambertW}\left(-\frac{e^{-\frac{4c_1}{3}}}{3x^{\frac{4}{3}}}\right) - \frac{4\ln(x)}{3} - \frac{4c_1}{3}}$$

✓ Solution by Mathematica

Time used: 13.068 (sec). Leaf size: 94

```
DSolve[x*(y[x]-3)*y'[x]==4*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -3W\left(\frac{1}{3}\sqrt[3]{-\frac{e^{-c_1}}{x^4}}\right)$$

$$y(x) \rightarrow -3W\left(-\frac{1}{3}\sqrt[3]{-1}\sqrt[3]{-\frac{e^{-c_1}}{x^4}}\right)$$

$$y(x) \rightarrow -3W\left(\frac{1}{3}(-1)^{2/3}\sqrt[3]{-\frac{e^{-c_1}}{x^4}}\right)$$

$$y(x) \rightarrow 0$$

2.2 problem 2

Internal problem ID [5088]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(x^3 + 1)y' - yx^2 = 0$$

With initial conditions

$$[y(1) = 2]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve([(1+x^3)*diff(y(x),x)=x^2*y(x),y(1) = 2],y(x), singsol=all)
```

$$y(x) = 2^{\frac{2}{3}}(x^3 + 1)^{\frac{1}{3}}$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 20

```
DSolve[{(1+x^3)*y'[x]==x^2*y[x],{y[1]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2^{2/3}\sqrt[3]{x^3 + 1}$$

2.3 problem 3

Internal problem ID [5089]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(1 + y)^2 y' = -x^3$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 89

```
dsolve(x^3+(y(x)+1)^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{(-6x^4 - 24c_1)^{\frac{1}{3}}}{2} - 1$$

$$y(x) = -\frac{(-6x^4 - 24c_1)^{\frac{1}{3}}}{4} - \frac{i\sqrt{3}(-6x^4 - 24c_1)^{\frac{1}{3}}}{4} - 1$$

$$y(x) = -\frac{(-6x^4 - 24c_1)^{\frac{1}{3}}}{4} + \frac{i\sqrt{3}(-6x^4 - 24c_1)^{\frac{1}{3}}}{4} - 1$$

✓ Solution by Mathematica

Time used: 0.483 (sec). Leaf size: 110

```
DSolve[x^3+(y[x]+1)^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -1 + \frac{\sqrt[3]{-3x^4 + 4 + 12c_1}}{2^{2/3}}$$

$$y(x) \rightarrow -1 + \frac{i(\sqrt{3} + i) \sqrt[3]{-3x^4 + 4 + 12c_1}}{2^{2/3}}$$

$$y(x) \rightarrow -1 - \frac{(1 + i\sqrt{3}) \sqrt[3]{-3x^4 + 4 + 12c_1}}{2^{2/3}}$$

2.4 problem 4

Internal problem ID [5090]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$\cos(y) + (1 + e^{-x}) \sin(y) y' = 0$$

With initial conditions

$$y(0) = \frac{\pi}{4}$$

✓ Solution by Maple

Time used: 0.156 (sec). Leaf size: 14

```
dsolve([cos(y(x))+(1+exp(-x))*sin(y(x))*diff(y(x),x)=0,y(0) = 1/4*Pi],y(x), singsol=all)
```

$$y(x) = \arccos\left(\frac{\sqrt{2}(e^x + 1)}{4}\right)$$

✓ Solution by Mathematica

Time used: 50.086 (sec). Leaf size: 20

```
DSolve[{Cos[y[x]]+(1+Exp[-x])*Sin[y[x]]*y'[x]==0,{y[0]==Pi/4}},y[x],x,IncludeSingularSolutio
```

$$y(x) \rightarrow \arccos\left(\frac{e^x + 1}{2\sqrt{2}}\right)$$

2.5 problem 5

Internal problem ID [5091]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$x^2(1+y) + y^2(x-1)y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve(x^2*(y(x)+1)+y(x)^2*(x-1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$\frac{x^2}{2} + x + \ln(x-1) + \frac{y(x)^2}{2} - y(x) + \ln(y(x)+1) + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.42 (sec). Leaf size: 56

```
DSolve[x^2*(y[x]+1)+y[x]^2*(x-1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{1}{2}(\#1+1)^2 - 2(\#1+1) + \log(\#1+1) \& \right] \left[-\frac{x^2}{2} - x - \log(x-1) + \frac{3}{2} + c_1 \right]$$

$$y(x) \rightarrow -1$$

2.6 problem 6

Internal problem ID [5092]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [[_homogeneous, 'class A'], _exact, _rational, [_Abel, '2nd ty

$$(2y - x)y' - y = 2x$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 53

```
dsolve((2*y(x)-x)*diff(y(x),x)=2*x+y(x),y(x), singsol=all)
```

$$y(x) = \frac{\frac{c_1 x}{2} - \frac{\sqrt{5c_1^2 x^2 + 4}}{2}}{c_1}$$

$$y(x) = \frac{\frac{c_1 x}{2} + \frac{\sqrt{5c_1^2 x^2 + 4}}{2}}{c_1}$$

✓ Solution by Mathematica

Time used: 0.454 (sec). Leaf size: 102

```
DSolve[(2*y[x]-x)*y'[x]==2*x+y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(x - \sqrt{5x^2 - 4e^{c_1}} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left(x + \sqrt{5x^2 - 4e^{c_1}} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left(x - \sqrt{5}\sqrt{x^2} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left(\sqrt{5}\sqrt{x^2} + x \right)$$

2.7 problem 7

Internal problem ID [5093]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$yx + y^2 + (x^2 - yx) y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve((x*y(x)+y(x)^2)+(x^2-x*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{e^{-\text{LambertW}\left(-\frac{e^{-2c_1}}{x^2}\right)-2c_1}}{x}$$

✓ Solution by Mathematica

Time used: 2.801 (sec). Leaf size: 25

```
DSolve[(x*y[x]+y[x]^2)+(x^2-x*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -xW\left(-\frac{e^{-c_1}}{x^2}\right)$$

$$y(x) \rightarrow 0$$

2.8 problem 8

Internal problem ID [5094]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _Bernoulli]`

$$y^3 - 3xy^2y' = -x^3$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 90

```
dsolve((x^3+y(x)^3)=3*x*y(x)^2*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = \frac{(4x^3 + 8c_1x)^{\frac{1}{3}}}{2}$$

$$y(x) = -\frac{(4x^3 + 8c_1x)^{\frac{1}{3}}}{4} - \frac{i\sqrt{3}(4x^3 + 8c_1x)^{\frac{1}{3}}}{4}$$

$$y(x) = -\frac{(4x^3 + 8c_1x)^{\frac{1}{3}}}{4} + \frac{i\sqrt{3}(4x^3 + 8c_1x)^{\frac{1}{3}}}{4}$$

✓ Solution by Mathematica

Time used: 0.21 (sec). Leaf size: 90

```
DSolve[(x^3+y[x]^3)==3*x*y[x]^2*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt[3]{-\frac{1}{2}\sqrt[3]{x}\sqrt[3]{x^2+2c_1}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{x}\sqrt[3]{x^2+2c_1}}{\sqrt[3]{2}}$$

$$y(x) \rightarrow \frac{(-1)^{2/3}\sqrt[3]{x}\sqrt[3]{x^2+2c_1}}{\sqrt[3]{2}}$$

2.9 problem 9

Internal problem ID [5095]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$y + (4y + 3x)y' = 3x$$

✓ Solution by Maple

Time used: 0.25 (sec). Leaf size: 278

```
dsolve(y(x)-3*x+(4*y(x)+3*x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{3x^8 c_1 \text{RootOf}(_Z^{64} c_1 x^8 + 12_Z^{56} c_1 x^8 + 48_Z^{48} c_1 x^8 + 64_Z^{40} c_1 x^8 - 1)^{56} + 24x^8 c_1 \text{RootOf}(_Z^{64} c_1 x^8 - 1)}{2c_1 x^7 \text{RootOf}(_Z^{64} c_1 x^8 + 12_Z^{56} c_1 x^8 + 48_Z^{48} c_1 x^8 + 64_Z^{40} c_1 x^8 - 1)^{40} \left(\text{RootOf}(_Z^{64} c_1 x^8 - 1) \right)}$$

✓ Solution by Mathematica

Time used: 5.296 (sec). Leaf size: 673

```
DSolve[y[x]-3*x+(4*y[x]+3*x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{Root}\left[256\#1^8 + 512\#1^7x - 512\#1^6x^2 - 896\#1^5x^3 + 800\#1^4x^4 + 352\#1^3x^5 - 576\#1^2x^6 + 216\#1x^7 - 27x^8 + e^{8c_1}\&, 1\right]$$

$$y(x) \rightarrow \text{Root}\left[256\#1^8 + 512\#1^7x - 512\#1^6x^2 - 896\#1^5x^3 + 800\#1^4x^4 + 352\#1^3x^5 - 576\#1^2x^6 + 216\#1x^7 - 27x^8 + e^{8c_1}\&, 2\right]$$

$$y(x) \rightarrow \text{Root}\left[256\#1^8 + 512\#1^7x - 512\#1^6x^2 - 896\#1^5x^3 + 800\#1^4x^4 + 352\#1^3x^5 - 576\#1^2x^6 + 216\#1x^7 - 27x^8 + e^{8c_1}\&, 3\right]$$

$$y(x) \rightarrow \text{Root}\left[256\#1^8 + 512\#1^7x - 512\#1^6x^2 - 896\#1^5x^3 + 800\#1^4x^4 + 352\#1^3x^5 - 576\#1^2x^6 + 216\#1x^7 - 27x^8 + e^{8c_1}\&, 4\right]$$

$$y(x) \rightarrow \text{Root}\left[256\#1^8 + 512\#1^7x - 512\#1^6x^2 - 896\#1^5x^3 + 800\#1^4x^4 + 352\#1^3x^5 - 576\#1^2x^6 + 216\#1x^7 - 27x^8 + e^{8c_1}\&, 5\right]$$

$$y(x) \rightarrow \text{Root}\left[256\#1^8 + 512\#1^7x - 512\#1^6x^2 - 896\#1^5x^3 + 800\#1^4x^4 + 352\#1^3x^5 - 576\#1^2x^6 + 216\#1x^7 - 27x^8 + e^{8c_1}\&, 6\right]$$

$$y(x) \rightarrow \text{Root}\left[256\#1^8 + 512\#1^7x - 512\#1^6x^2 - 896\#1^5x^3 + 800\#1^4x^4 + 352\#1^3x^5 - 576\#1^2x^6 + 216\#1x^7 - 27x^8 + e^{8c_1}\&, 7\right]$$

$$y(x) \rightarrow \text{Root}\left[256\#1^8 + 512\#1^7x - 512\#1^6x^2 - 896\#1^5x^3 + 800\#1^4x^4 + 352\#1^3x^5 - 576\#1^2x^6 + 216\#1x^7 - 27x^8 + e^{8c_1}\&, 8\right]$$

2.10 problem 10

Internal problem ID [5096]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _dAlembert]`

$$(x^3 + 3xy^2) y' - y^3 - 3yx^2 = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 23

```
dsolve((x^3+3*x*y(x)^2)*diff(y(x),x)=y(x)^3+3*x^2*y(x),y(x), singsol=all)
```

$$y(x) = \text{RootOf}(_Z^4 c_1 x - c_1 x - _Z)^2 x$$

✓ Solution by Mathematica

Time used: 60.142 (sec). Leaf size: 1659

`DSolve[(x^3+3*x*y[x]^2)*y'[x]==y[x]^3+3*x^2*y[x],y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{1}{6} \left(-\sqrt{3} \sqrt{4x^2 + \frac{16\sqrt[3]{2}x^4}{\sqrt[3]{128x^6 + 27e^{2c_1}x^2 + 3\sqrt{768e^{2c_1}x^8 + 81e^{4c_1}x^4}}}} + \frac{\sqrt[3]{128x^6 + 27e^{2c_1}x^2 + 3\sqrt{768e^{2c_1}x^8 + 81e^{4c_1}x^4}}}{\sqrt[3]{2}} \right)$$

$$-3 \sqrt{\frac{8x^2}{3} - \frac{16\sqrt[3]{2}x^4}{3\sqrt[3]{128x^6 + 27e^{2c_1}x^2 + 3\sqrt{768e^{2c_1}x^8 + 81e^{4c_1}x^4}}}} - \sqrt{4x^2 + \frac{16\sqrt[3]{2}x^4}{\sqrt[3]{128x^6 + 27e^{2c_1}x^2 + 3\sqrt{768e^{2c_1}x^8 + 81e^{4c_1}x^4}}}}$$

$$y(x) \rightarrow \frac{1}{6} \left(3 \sqrt{\frac{8x^2}{3} - \frac{16\sqrt[3]{2}x^4}{3\sqrt[3]{128x^6 + 27e^{2c_1}x^2 + 3\sqrt{768e^{2c_1}x^8 + 81e^{4c_1}x^4}}}} - \sqrt{4x^2 + \frac{16\sqrt[3]{2}x^4}{\sqrt[3]{128x^6 + 27e^{2c_1}x^2 + 3\sqrt{768e^{2c_1}x^8 + 81e^{4c_1}x^4}}}} \right)$$

$$-\sqrt{3} \sqrt{4x^2 + \frac{16\sqrt[3]{2}x^4}{\sqrt[3]{128x^6 + 27e^{2c_1}x^2 + 3\sqrt{768e^{2c_1}x^8 + 81e^{4c_1}x^4}}}} + \frac{\sqrt[3]{128x^6 + 27e^{2c_1}x^2 + 3\sqrt{768e^{2c_1}x^8 + 81e^{4c_1}x^4}}}{\sqrt[3]{2}}$$

$$y(x) \rightarrow \frac{1}{6} \left(\sqrt{3} \sqrt{4x^2 + \frac{16\sqrt[3]{2}x^4}{\sqrt[3]{128x^6 + 27e^{2c_1}x^2 + 3\sqrt{768e^{2c_1}x^8 + 81e^{4c_1}x^4}}}} + \frac{\sqrt[3]{128x^6 + 27e^{2c_1}x^2 + 3\sqrt{768e^{2c_1}x^8 + 81e^{4c_1}x^4}}}{\sqrt[3]{2}} \right)$$

2.11 problem 11

Internal problem ID [5097]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y'x - y = x^3 + 3x^2 - 2x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(x*diff(y(x),x)-y(x)=x^3+3*x^2-2*x,y(x), singsol=all)
```

$$y(x) = \left(\frac{x^2}{2} + 3x - 2 \ln(x) + c_1 \right) x$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 24

```
DSolve[x*y'[x]-y[x]==x^3+3*x^2-2*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \left(\frac{x^2}{2} + 3x - 2 \log(x) + c_1 \right)$$

2.12 problem 12

Internal problem ID [5098]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + y \tan(x) = \sin(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)+y(x)*tan(x)=sin(x),y(x), singsol=all)
```

$$y(x) = (-\ln(\cos(x)) + c_1) \cos(x)$$

✓ Solution by Mathematica

Time used: 0.058 (sec). Leaf size: 16

```
DSolve[y'[x]+y[x]*Tan[x]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \cos(x)(-\log(\cos(x)) + c_1)$$

2.13 problem 13

Internal problem ID [5099]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y'x - y = x^3 \cos(x)$$

With initial conditions

$$[y(\pi) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([x*diff(y(x),x)-y(x)=x^3*cos(x),y(Pi) = 0],y(x), singsol=all)
```

$$y(x) = (\cos(x) + \sin(x)x + 1)x$$

✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 15

```
DSolve[{x*y'[x]-y[x]==x^3*Cos[x],{y[Pi]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(x \sin(x) + \cos(x) + 1)$$

2.14 problem 14

Internal problem ID [5100]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 14.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(x^2 + 1) y' + 3yx = 5x$$

With initial conditions

$$[y(1) = 2]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve([(1+x^2)*diff(y(x),x)+3*x*y(x)=5*x,y(1) = 2],y(x), singsol=all)
```

$$y(x) = \frac{5}{3} + \frac{2\sqrt{2}}{3(x^2 + 1)^{\frac{3}{2}}}$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 27

```
DSolve[{(1+x^2)*y'[x]+3*x*y[x]==5*x,{y[1]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2\sqrt{2}}{3(x^2 + 1)^{3/2}} + \frac{5}{3}$$

2.15 problem 15

Internal problem ID [5101]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + y \cot(x) = 5e^{\cos(x)}$$

With initial conditions

$$\left[y\left(\frac{\pi}{2}\right) = -4 \right]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([diff(y(x),x)+y(x)*cot(x)=5*exp(cos(x)),y(1/2*Pi) = -4],y(x), singsol=all)
```

$$y(x) = -5e^{\cos(x)} \csc(x) + \csc(x)$$

✓ Solution by Mathematica

Time used: 0.1 (sec). Leaf size: 16

```
DSolve[{y'[x]+y[x]*Cot[x]==5*Exp[Cos[x]],{y[Pi/2]==-4}},y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow (1 - 5e^{\cos(x)}) \csc(x)$$

2.16 problem 16

Internal problem ID [5102]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$(3x + 3y - 4)y' + y = -x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve((3*x+3*y(x)-4)*diff(y(x),x)=-x+y(x)),y(x), singsol=all)
```

$$y(x) = e^{-\text{LambertW}\left(\frac{3e^x e^{-3} e^{-c_1}}{2}\right) + x - 3 - c_1} + 2 - x$$

✓ Solution by Mathematica

Time used: 3.675 (sec). Leaf size: 33

```
DSolve[(3*x+3*y[x]-4)*y'[x]==-(x+y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2}{3}W(-e^{x-1+c_1}) - x + 2$$

$$y(x) \rightarrow 2 - x$$

2.17 problem 17

Internal problem ID [5103]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 17.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, [_1st_order, ‘_with_symmetry_[F(x)*G(y),0]’], [_Ab`

$$-xy^2 - (x + yx^2)y' = -x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 56

```
dsolve((x-x*y(x)^2)=(x+x^2*y(x))*diff(y(x),x),y(x), singsol=all)
```

$$x + \frac{\sqrt{(y(x)-1)(y(x)+1)} \ln\left(y(x) + \sqrt{y(x)^2 - 1}\right)}{(y(x)-1)(y(x)+1)} - \frac{c_1}{\sqrt{y(x)-1}\sqrt{y(x)+1}} = 0$$

✓ Solution by Mathematica

Time used: 0.127 (sec). Leaf size: 55

```
DSolve[(x-x*y[x]^2)==(x+x^2*y[x])*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[x = -\frac{2 \arctan\left(\frac{\sqrt{1-y(x)^2}}{y(x)+1}\right)}{\sqrt{1-y(x)^2}} + \frac{c_1}{\sqrt{1-y(x)^2}}, y(x)\right]$$

2.18 problem 18

Internal problem ID [5104]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 18.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _rational, [_Abel, '2nd type', 'cl`

$$-y + (4y + x - 1)y' = 1 - x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve((x-y(x)-1)+(4*y(x)+x-1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{\tan\left(\text{RootOf}\left(\ln\left(\frac{1}{\cos(_Z)^2}\right) - _Z + 2\ln(x-1) + 2c_1\right)\right)(x-1)}{2}$$

✓ Solution by Mathematica

Time used: 0.059 (sec). Leaf size: 58

```
DSolve[(x-y[x]-1)+(4*y[x]+x-1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[2\arctan\left(\frac{2y(x)-2x+2}{4y(x)+x-1}\right) + 2\log\left(\frac{4}{5}\left(\frac{4y(x)^2}{(x-1)^2} + 1\right)\right) + 4\log(x-1) + 5c_1 = 0, y(x)\right]$$

2.19 problem 19

Internal problem ID [5105]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$3y + (7y - 3x + 3)y' = 7x - 7$$

✓ Solution by Maple

Time used: 0.187 (sec). Leaf size: 706

```
dsolve((3*y(x)-7*x+7)+(7*y(x)-3*x+3)*diff(y(x),x)=0,y(x), singsol=all)
```

Expression too large to display

✓ Solution by Mathematica

Time used: 60.706 (sec). Leaf size: 7785

```
DSolve[(3*y[x]-7*x+7)+(7*y[x]-3*x+3)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

2.20 problem 20

Internal problem ID [5106]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 20.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational]`

$$y(1 + yx) + x(1 + yx + x^2y^2) y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 38

```
dsolve(y(x)*(x*y(x)+1)+x*(1+x*y(x)+x^2*y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{e^{\text{RootOf}(-2\ln(x)e^{2-Z}+2c_1e^{2-Z}+2_Ze^{2-Z}-2e^{-Z}-1)}}}{x}$$

✓ Solution by Mathematica

Time used: 0.11 (sec). Leaf size: 30

```
DSolve[y[x]*(x*y[x]+1)+x*(1+x*y[x]+x^2*y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> T
```

$$\text{Solve} \left[\frac{-\frac{1}{2x^2} - \frac{y(x)}{x}}{y(x)^2} + \log(y(x)) = c_1, y(x) \right]$$

2.21 problem 21

Internal problem ID [5107]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 21.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_Bernoulli]

$$y' + y - xy^3 = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 39

```
dsolve(diff(y(x),x)+y(x)=x*y(x)^3,y(x), singsol=all)
```

$$y(x) = -\frac{2}{\sqrt{2 + 4e^{2x}c_1 + 4x}}$$

$$y(x) = \frac{2}{\sqrt{2 + 4e^{2x}c_1 + 4x}}$$

✓ Solution by Mathematica

Time used: 2.704 (sec). Leaf size: 50

```
DSolve[y'[x]+y[x]==x*y[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\sqrt{x + c_1 e^{2x} + \frac{1}{2}}}$$

$$y(x) \rightarrow \frac{1}{\sqrt{x + c_1 e^{2x} + \frac{1}{2}}}$$

$$y(x) \rightarrow 0$$

2.22 problem 22

Internal problem ID [5108]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 22.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _Bernoulli]`

$$y' + y - y^4 e^x = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 194

```
dsolve(diff(y(x),x)+y(x)=y(x)^4*exp(x),y(x), singsol=all)
```

$$y(x) = \frac{2^{\frac{1}{3}} \left((2c_1 e^{3x} + 3e^x)^2 \right)^{\frac{1}{3}}}{2c_1 e^{3x} + 3e^x}$$
$$y(x) = -\frac{2^{\frac{1}{3}} \left((2c_1 e^{3x} + 3e^x)^2 \right)^{\frac{1}{3}}}{2(2c_1 e^{3x} + 3e^x)} - \frac{i\sqrt{3} 2^{\frac{1}{3}} \left((2c_1 e^{3x} + 3e^x)^2 \right)^{\frac{1}{3}}}{2(2c_1 e^{3x} + 3e^x)}$$
$$y(x) = -\frac{2^{\frac{1}{3}} \left((2c_1 e^{3x} + 3e^x)^2 \right)^{\frac{1}{3}}}{2(2c_1 e^{3x} + 3e^x)} + \frac{i\sqrt{3} 2^{\frac{1}{3}} \left((2c_1 e^{3x} + 3e^x)^2 \right)^{\frac{1}{3}}}{4c_1 e^{3x} + 6e^x}$$

✓ Solution by Mathematica

Time used: 4.751 (sec). Leaf size: 90

```
DSolve[y'[x]+y[x]==y[x]^4*Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt[3]{-2}}{\sqrt[3]{e^x(3+2c_1e^{2x})}}$$

$$y(x) \rightarrow \frac{1}{\sqrt[3]{\frac{3e^x}{2} + c_1e^{3x}}}$$

$$y(x) \rightarrow \frac{(-1)^{2/3}}{\sqrt[3]{\frac{3e^x}{2} + c_1e^{3x}}}$$

$$y(x) \rightarrow 0$$

2.23 problem 23

Internal problem ID [5109]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 23.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_Bernoulli]

$$2y' + y - y^3(x-1) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(2*diff(y(x),x)+y(x)=y(x)^3*(x-1),y(x), singsol=all)
```

$$y(x) = \frac{1}{\sqrt{e^x c_1 + x}}$$
$$y(x) = -\frac{1}{\sqrt{e^x c_1 + x}}$$

✓ Solution by Mathematica

Time used: 2.721 (sec). Leaf size: 40

```
DSolve[2*y'[x]+y[x]==y[x]^3*(x-1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\sqrt{x + c_1 e^x}}$$
$$y(x) \rightarrow \frac{1}{\sqrt{x + c_1 e^x}}$$
$$y(x) \rightarrow 0$$

2.24 problem 24

Internal problem ID [5110]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 24.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_Bernoulli]

$$y' - 2y \tan(x) - y^2 \tan(x)^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x)-2*y(x)*tan(x)=y(x)^2*tan(x)^2,y(x), singsol=all)
```

$$y(x) = -\frac{3}{\cos(x)^2 (\tan(x)^3 - 3c_1)}$$

✓ Solution by Mathematica

Time used: 0.519 (sec). Leaf size: 31

```
DSolve[y'[x]-2*y[x]*Tan[x]==y[x]^2*Tan[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3}{-\sin^2(x) \tan(x) + 3c_1 \cos^2(x)}$$
$$y(x) \rightarrow 0$$

2.25 problem 25

Internal problem ID [5111]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 25.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [Bernoulli]

$$y' + y \tan(x) - y^3 \sec(x)^4 = 0$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 92

```
dsolve(diff(y(x),x)+y(x)*tan(x)=y(x)^3*sec(x)^4,y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{\cos(x)(c_1 \cos(x) - 2 \sin(x))(\sin(x)^4 + 2 \cos(x)^2 - 1)}}{\cos(x)(c_1 \cos(x) - 2 \sin(x))}$$

$$y(x) = -\frac{\sqrt{\cos(x)(c_1 \cos(x) - 2 \sin(x))(\sin(x)^4 + 2 \cos(x)^2 - 1)}}{\cos(x)(c_1 \cos(x) - 2 \sin(x))}$$

✓ Solution by Mathematica

Time used: 4.061 (sec). Leaf size: 48

```
DSolve[y'[x]+y[x]*Tan[x]==y[x]^3*Sec[x]^4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\sqrt{\sec^2(x)(-2 \tan(x) + c_1)}}$$

$$y(x) \rightarrow \frac{1}{\sqrt{\sec^2(x)(-2 \tan(x) + c_1)}}$$

$$y(x) \rightarrow 0$$

2.26 problem 26

Internal problem ID [5112]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 26.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$(-x^2 + 1) y' - yx = 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 48

```
dsolve((1-x^2)*diff(y(x),x)=1+x*y(x),y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{(x-1)(x+1)} \ln(x + \sqrt{x^2-1})}{(x-1)(x+1)} + \frac{c_1}{\sqrt{x-1}\sqrt{x+1}}$$

✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 54

```
DSolve[(1-x^2)*y'[x]==1+x*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\log\left(1 - \frac{x}{\sqrt{x^2-1}}\right) - \log\left(\frac{x}{\sqrt{x^2-1}} + 1\right) + 2c_1}{2\sqrt{x^2-1}}$$

2.27 problem 27

Internal problem ID [5113]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 27.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$xyy' - (1+x)\sqrt{-1+y} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(x*y(x)*diff(y(x),x)-(1+x)*sqrt(y(x)-1)=0,y(x), singsol=all)
```

$$x + \ln(x) - \frac{2\sqrt{y(x)-1}(y(x)+2)}{3} + c_1 = 0$$

✓ Solution by Mathematica

Time used: 5.614 (sec). Leaf size: 582

`DSolve[x*y[x]*y'[x]-(1+x)*Sqrt[y[x]-1]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{1}{2} \sqrt[3]{9x^2 + 3\sqrt{(x + \log(x) + c_1)^2 (9x^2 + 9\log^2(x) + 18c_1x + 18(x + c_1)\log(x) + 16 + 9c_1^2) + 9\log^2(x)}} + \frac{\sqrt[3]{9x^2 + 3\sqrt{(x + \log(x) + c_1)^2 (9x^2 + 9\log^2(x) + 18c_1x + 18(x + c_1)\log(x) + 16 + 9c_1^2) + 9\log^2(x)}}}{-1}$$

$$y(x) \rightarrow \frac{1}{4}i(\sqrt{3} + i) \sqrt[3]{9x^2 + 3\sqrt{(x + \log(x) + c_1)^2 (9x^2 + 9\log^2(x) + 18c_1x + 18(x + c_1)\log(x) + 16 + 9c_1^2) + 9\log^2(x)}} + \frac{\sqrt[3]{9x^2 + 3\sqrt{(x + \log(x) + c_1)^2 (9x^2 + 9\log^2(x) + 18c_1x + 18(x + c_1)\log(x) + 16 + 9c_1^2) + 9\log^2(x)}}}{-1}$$

$$y(x) \rightarrow -\frac{1}{4}i(\sqrt{3} - i) \sqrt[3]{9x^2 + 3\sqrt{(x + \log(x) + c_1)^2 (9x^2 + 9\log^2(x) + 18c_1x + 18(x + c_1)\log(x) + 16 + 9c_1^2) + 9\log^2(x)}} + \frac{\sqrt[3]{9x^2 + 3\sqrt{(x + \log(x) + c_1)^2 (9x^2 + 9\log^2(x) + 18c_1x + 18(x + c_1)\log(x) + 16 + 9c_1^2) + 9\log^2(x)}}}{-1}$$

$$y(x) \rightarrow 1$$

2.28 problem 28

Internal problem ID [5114]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 28.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _dAlembert]`

$$-2yx + 5y^2 - (x^2 + 2yx + y^2) y' = -x^2$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 35

```
dsolve((x^2-2*x*y(x)+5*y(x)^2)=(x^2+2*x*y(x)+y(x)^2)*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = e^{\text{RootOf}(\ln(x)e^{2-Z} + c_1e^{2-Z} + Ze^{2-Z} - 4e^{-Z} - 2)} x + x$$

✓ Solution by Mathematica

Time used: 0.343 (sec). Leaf size: 41

```
DSolve[(x^2-2*x*y[x]+5*y[x]^2)==(x^2+2*x*y[x]+y[x]^2)*y'[x],y[x],x,IncludeSingularSolutions
```

$$\text{Solve} \left[\frac{2 - \frac{4y(x)}{x}}{\left(\frac{y(x)}{x} - 1\right)^2} + \log\left(\frac{y(x)}{x} - 1\right) = -\log(x) + c_1, y(x) \right]$$

2.29 problem 29

Internal problem ID [5115]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 29.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [Bernoulli]

$$y' - y \cot(x) - y^2 \sec(x)^2 = 0$$

With initial conditions

$$\left[y\left(\frac{\pi}{4}\right) = -1 \right]$$

✓ Solution by Maple

Time used: 0.469 (sec). Leaf size: 18

```
dsolve([diff(y(x),x)-y(x)*cot(x)=y(x)^2*sec(x)^2,y(1/4*Pi) = -1],y(x), singsol=all)
```

$$y(x) = \frac{2 \sin(x)}{\sqrt{2} - 2 \sec(x)}$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y'[x]-y[x]*Cot[x]==y[x]^2*Sec[x]^2,{y[Pi/2]==-1}],y[x],x,IncludeSingularSolutions ->
```

```
{}
```

2.30 problem 30

Internal problem ID [5116]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 30.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y + (x^2 - 4x)y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(y(x)+(x^2-4*x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 x^{\frac{1}{4}}}{(x-4)^{\frac{1}{4}}}$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 27

```
DSolve[y[x]+(x^2-4*x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1 \sqrt[4]{x}}{\sqrt[4]{4-x}}$$

$$y(x) \rightarrow 0$$

2.31 problem 31

Internal problem ID [5117]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 31.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' - y \tan(x) = \cos(x) - 2 \sin(x) x$$

With initial conditions

$$\left[y\left(\frac{\pi}{6}\right) = 0 \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 22

```
dsolve([diff(y(x),x)-y(x)*tan(x)=cos(x)-2*x*sin(x),y(1/6*Pi) = 0],y(x), singsol=all)
```

$$y(x) = \frac{(4x \cos(2x) - \pi + 4x) \sec(x)}{8}$$

✓ Solution by Mathematica

Time used: 0.073 (sec). Leaf size: 25

```
DSolve[{y'[x]-y[x]*Tan[x]==Cos[x]-2*x*Sin[x],{y[Pi/6]==0}},y[x],x,IncludeSingularSolutions -
```

$$y(x) \rightarrow \frac{1}{8}(4x + 4x \cos(2x) - \pi) \sec(x)$$

2.32 problem 32

Internal problem ID [5118]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 32.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{2yx + y^2}{x^2 + 2yx} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 380

`dsolve(diff(y(x),x)=(2*x*y(x)+y(x)^2)/(x^2+2*x*y(x)),y(x), singsol=all)`

$$y(x) = \frac{12^{\frac{1}{3}} \left(x \left(\sqrt{3} \sqrt{\frac{x(27c_1x-4)}{c_1}} + 9x \right) c_1^2 \right)^{\frac{1}{3}}}{6c_1} + \frac{x12^{\frac{2}{3}}}{6 \left(x \left(\sqrt{3} \sqrt{\frac{x(27c_1x-4)}{c_1}} + 9x \right) c_1^2 \right)^{\frac{1}{3}}} + x$$

$$y(x) = -\frac{12^{\frac{1}{3}} \left(x \left(\sqrt{3} \sqrt{\frac{x(27c_1x-4)}{c_1}} + 9x \right) c_1^2 \right)^{\frac{1}{3}}}{12c_1} - \frac{x12^{\frac{2}{3}}}{12 \left(x \left(\sqrt{3} \sqrt{\frac{x(27c_1x-4)}{c_1}} + 9x \right) c_1^2 \right)^{\frac{1}{3}}} + x - \frac{i\sqrt{3} \left(\frac{12^{\frac{1}{3}} \left(x \left(\sqrt{3} \sqrt{\frac{x(27c_1x-4)}{c_1}} + 9x \right) c_1^2 \right)^{\frac{1}{3}}}{6c_1} - \frac{x12^{\frac{2}{3}}}{6 \left(x \left(\sqrt{3} \sqrt{\frac{x(27c_1x-4)}{c_1}} + 9x \right) c_1^2 \right)^{\frac{1}{3}}} \right)}{2}$$

$$y(x) = -\frac{12^{\frac{1}{3}} \left(x \left(\sqrt{3} \sqrt{\frac{x(27c_1x-4)}{c_1}} + 9x \right) c_1^2 \right)^{\frac{1}{3}}}{12c_1} - \frac{x12^{\frac{2}{3}}}{12 \left(x \left(\sqrt{3} \sqrt{\frac{x(27c_1x-4)}{c_1}} + 9x \right) c_1^2 \right)^{\frac{1}{3}}} + x + \frac{i\sqrt{3} \left(\frac{12^{\frac{1}{3}} \left(x \left(\sqrt{3} \sqrt{\frac{x(27c_1x-4)}{c_1}} + 9x \right) c_1^2 \right)^{\frac{1}{3}}}{6c_1} - \frac{x12^{\frac{2}{3}}}{6 \left(x \left(\sqrt{3} \sqrt{\frac{x(27c_1x-4)}{c_1}} + 9x \right) c_1^2 \right)^{\frac{1}{3}}} \right)}{2}$$

✓ Solution by Mathematica

Time used: 56.42 (sec). Leaf size: 404

`DSolve[y'[x]==(2*x*y[x]+y[x]^2)/(x^2+2*x*y[x]),y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow -\frac{\sqrt[3]{\frac{2}{3}}e^{c_1}x}{\sqrt[3]{\sqrt{3}\sqrt{e^{2c_1}x^3(27x+4e^{c_1})}-9e^{c_1}x^2}}} + \frac{\sqrt[3]{\sqrt{3}\sqrt{e^{2c_1}x^3(27x+4e^{c_1})}-9e^{c_1}x^2}}}{\sqrt[3]{23^{2/3}}} + x$$

$$y(x) \rightarrow \frac{(1+i\sqrt{3})e^{c_1}x}{2^{2/3}\sqrt[3]{3\sqrt{3}\sqrt{e^{2c_1}x^3(27x+4e^{c_1})}-27e^{c_1}x^2}}} + \frac{i(\sqrt{3}+i)\sqrt[3]{\sqrt{3}\sqrt{e^{2c_1}x^3(27x+4e^{c_1})}-9e^{c_1}x^2}}}{2\sqrt[3]{23^{2/3}}} + x$$

$$y(x) \rightarrow \frac{(1-i\sqrt{3})e^{c_1}x}{2^{2/3}\sqrt[3]{3\sqrt{3}\sqrt{e^{2c_1}x^3(27x+4e^{c_1})}-27e^{c_1}x^2}}} - \frac{(1+i\sqrt{3})\sqrt[3]{\sqrt{3}\sqrt{e^{2c_1}x^3(27x+4e^{c_1})}-9e^{c_1}x^2}}}{2\sqrt[3]{23^{2/3}}} + x$$

2.33 problem 33

Internal problem ID [5119]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 33.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(x^2 + 1) y' - x(1 + y) = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve((1+x^2)*diff(y(x),x)=x*(1+y(x)),y(x), singsol=all)
```

$$y(x) = c_1 \sqrt{x^2 + 1} - 1$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 24

```
DSolve[(1+x^2)*y'[x]==x*(1+y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -1 + c_1 \sqrt{x^2 + 1}$$

$$y(x) \rightarrow -1$$

2.34 problem 34

Internal problem ID [5120]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 34.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y'x + 2y = 3x - 1$$

With initial conditions

$$[y(2) = 1]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 12

```
dsolve([x*diff(y(x),x)+2*y(x)=3*x-1,y(2) = 1],y(x), singsol=all)
```

$$y(x) = x - \frac{1}{2} - \frac{2}{x^2}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 15

```
DSolve[{x*y'[x]+2*y[x]==3*x-1,{y[2]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2}{x^2} + x - \frac{1}{2}$$

2.35 problem 35

Internal problem ID [5121]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 35.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$x^2y' - y^2 + xyy' = 0$$

With initial conditions

$$[y(1) = 1]$$

✓ Solution by Maple

Time used: 0.187 (sec). Leaf size: 13

```
dsolve([x^2*diff(y(x),x)=y(x)^2-x*y(x)*diff(y(x),x),y(1) = 1],y(x), singsol=all)
```

$$y(x) = \text{LambertW}\left(\frac{e}{x}\right) x$$

✓ Solution by Mathematica

Time used: 2.335 (sec). Leaf size: 13

```
DSolve[{x^2*y'[x]==y[x]^2-x*y[x]*y'[x],{y[1]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow xW\left(\frac{e}{x}\right)$$

2.36 problem 36

Internal problem ID [5122]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 36.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - e^{3x-2y} = 0$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 15

```
dsolve([diff(y(x),x)=exp(3*x-2*y(x)),y(0) = 0],y(x), singsol=all)
```

$$y(x) = -\frac{\ln(3)}{2} + \frac{\ln(1 + 2e^{3x})}{2}$$

✓ Solution by Mathematica

Time used: 0.881 (sec). Leaf size: 23

```
DSolve[{y'[x]==Exp[3*x-2*y[x]],{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \log \left(\frac{1}{3} (2e^{3x} + 1) \right)$$

2.37 problem 37

Internal problem ID [5123]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 37.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + \frac{y}{x} = \sin(2x)$$

With initial conditions

$$\left[y\left(\frac{\pi}{4}\right) = 2 \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 25

```
dsolve([diff(y(x),x)+1/x*y(x)=sin(2*x),y(1/4*Pi) = 2],y(x), singsol=all)
```

$$y(x) = \frac{-2x \cos(2x) + 2\pi + \sin(2x) - 1}{4x}$$

✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 28

```
DSolve[{y'[x]+1/x*y[x]==Sin[2*x],{y[Pi/4]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sin(2x) - 2x \cos(2x) + 2\pi - 1}{4x}$$

2.38 problem 38

Internal problem ID [5124]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 38.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$y^2 + x^2 y' - x y y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(y(x)^2+x^2*diff(y(x),x)=x*y(x)*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = e^{-\text{LambertW}\left(-\frac{e^{-c_1}}{x}\right) - c_1}$$

✓ Solution by Mathematica

Time used: 2.23 (sec). Leaf size: 25

```
DSolve[y[x]^2+x^2*y'[x]==x*y[x]*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -xW\left(-\frac{e^{-c_1}}{x}\right)$$

$$y(x) \rightarrow 0$$

2.39 problem 39

Internal problem ID [5125]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 39.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _exact, _rational, _Bernoulli]`

$$2xyy' + y^2 = x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 45

```
dsolve(2*x*y(x)*diff(y(x),x)=x^2-y(x)^2,y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{3} \sqrt{x(x^3 + 3c_1)}}{3x}$$

$$y(x) = \frac{\sqrt{3} \sqrt{x(x^3 + 3c_1)}}{3x}$$

✓ Solution by Mathematica

Time used: 0.2 (sec). Leaf size: 56

```
DSolve[2*x*y[x]*y'[x]==x^2-y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{x^3 + 3c_1}}{\sqrt{3}\sqrt{x}}$$

$$y(x) \rightarrow \frac{\sqrt{x^3 + 3c_1}}{\sqrt{3}\sqrt{x}}$$

2.40 problem 40

Internal problem ID [5126]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 40.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{x - 2y + 1}{2x - 4y} = 0$$

With initial conditions

$$[y(1) = 1]$$

✓ Solution by Maple

Time used: 0.109 (sec). Leaf size: 17

```
dsolve([diff(y(x),x)=(x-2*y(x)+1)/(2*x-4*y(x)),y(1) = 1],y(x), singsol=all)
```

$$y(x) = \frac{x}{2} + \frac{\sqrt{-2x+3}}{2}$$

✓ Solution by Mathematica

Time used: 0.115 (sec). Leaf size: 24

```
DSolve[{y'[x]==(x-2*y[x]+1)/(2*x-4*y[x]),{y[1]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(x - i\sqrt{2x-3})$$

2.41 problem 41

Internal problem ID [5127]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 41.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$(-x^3 + 1)y' + yx^2 = x^2(-x^3 + 1)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve((1-x^3)*diff(y(x),x)+x^2*y(x)=x^2*(1-x^3),y(x), singsol=all)
```

$$y(x) = \frac{x^3}{2} - \frac{1}{2} + (x^3 - 1)^{\frac{1}{3}} c_1$$

✓ Solution by Mathematica

Time used: 0.054 (sec). Leaf size: 27

```
DSolve[(1-x^3)*y'[x]+x^2*y[x]==x^2*(1-x^3),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(x^3 + 2c_1 \sqrt[3]{x^3 - 1} - 1 \right)$$

2.42 problem 42

Internal problem ID [5128]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 42.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + \frac{y}{x} = \sin(x)$$

With initial conditions

$$\left[y\left(\frac{\pi}{2}\right) = 0 \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve([diff(y(x),x)+y(x)/x=sin(x),y(1/2*Pi) = 0],y(x), singsol=all)
```

$$y(x) = \frac{\sin(x) - \cos(x)x - 1}{x}$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 18

```
DSolve[{y'[x]+y[x]/x==Sin[x],{y[Pi/2]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sin(x) - x \cos(x) - 1}{x}$$

2.43 problem 43

Internal problem ID [5129]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 43.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' + xy^2 = -x$$

With initial conditions

$$[y(1) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve([diff(y(x),x)+x*x*y(x)^2=0,y(1) = 0],y(x), singsol=all)
```

$$y(x) = -\tan\left(\frac{x^2}{2} - \frac{1}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.215 (sec). Leaf size: 17

```
DSolve[{y'[x]+x*x*y[x]^2==0,{y[1]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan\left(\frac{1}{2}(1 - x^2)\right)$$

2.44 problem 44

Internal problem ID [5130]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 44.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + \left(\frac{1}{x} - \frac{2x}{-x^2 + 1} \right) y = \frac{1}{-x^2 + 1}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x)+(1/x-(2*x)/(1-x^2))*y(x)=1/(1-x^2),y(x), singsol=all)
```

$$y(x) = \frac{-\frac{x^2}{2} + c_1}{x(x^2 - 1)}$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 25

```
DSolve[y'[x]+(1/x-(2*x)/(1-x^2))*y[x]==1/(1-x^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2 + 2c_1}{2x - 2x^3}$$

2.45 problem 45

Internal problem ID [5131]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 45.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$(x^2 + 1) y' + yx = (x^2 + 1)^{\frac{3}{2}}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve((1+x^2)*diff(y(x),x)+x*y(x)=(1+x^2)^(3/2),y(x), singsol=all)
```

$$y(x) = \frac{\frac{1}{3}x^3 + x + c_1}{\sqrt{x^2 + 1}}$$

✓ Solution by Mathematica

Time used: 0.065 (sec). Leaf size: 29

```
DSolve[(1+x^2)*y'[x]+x*y[x]==(1+x^2)^(3/2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3 + 3x + 3c_1}{3\sqrt{x^2 + 1}}$$

2.46 problem 46

Internal problem ID [5132]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 46.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$x(y^2 + 1) - y(x^2 + 1)y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(x*(1+y(x)^2)-y(x)*(1+x^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \sqrt{c_1 x^2 + c_1 - 1}$$

$$y(x) = -\sqrt{c_1 x^2 + c_1 - 1}$$

✓ Solution by Mathematica

Time used: 0.499 (sec). Leaf size: 61

```
DSolve[x*(1+y[x]^2)-y[x]*(1+x^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{-1 + e^{2c_1} (x^2 + 1)}$$

$$y(x) \rightarrow \sqrt{-1 + e^{2c_1} (x^2 + 1)}$$

$$y(x) \rightarrow -i$$

$$y(x) \rightarrow i$$

2.47 problem 47

Internal problem ID [5133]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 47.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$\frac{r \tan(\theta) r'}{a^2 - r^2} = 1$$

With initial conditions

$$\left[r\left(\frac{\pi}{4}\right) = 0 \right]$$

✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 37

```
dsolve([r(theta)*tan(theta)/(a^2-r(theta)^2)*diff(r(theta),theta)=1,r(1/4*Pi) = 0],r(theta),
```

$$r(\theta) = -\frac{a\sqrt{-4\cos(\theta)^2 + 2\csc(\theta)}}{2}$$

$$r(\theta) = \frac{a\sqrt{-4\cos(\theta)^2 + 2\csc(\theta)}}{2}$$

✓ Solution by Mathematica

Time used: 0.149 (sec). Leaf size: 51

```
DSolve[{r[\[Theta]]*Tan[\[Theta]]/(a^2-r[\[Theta]]^2)*r'[\[Theta]]==1,{r[Pi/4]==0}},r[\[Theta]]
```

$$r(\theta) \rightarrow -\sqrt{\frac{a^2 \cos(2\theta)}{\cos(2\theta) - 1}}$$

$$r(\theta) \rightarrow \sqrt{\frac{a^2 \cos(2\theta)}{\cos(2\theta) - 1}}$$

2.48 problem 48

Internal problem ID [5134]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 48.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + y \cot(x) = \cos(x)$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 8

```
dsolve([diff(y(x),x)+y(x)*cot(x)=cos(x),y(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{\sin(x)}{2}$$

✓ Solution by Mathematica

Time used: 0.103 (sec). Leaf size: 11

```
DSolve[{y'[x]+y[x]*Cot[x]==Cos[x],{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sin(x)}{2}$$

2.49 problem 49

Internal problem ID [5135]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 24. First order differential equations. Further problems 24. page 1068

Problem number: 49.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational, _Bernoulli]`

$$y' + \frac{y}{x} - xy^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)+y(x)/x=x*y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{1}{(-x + c_1)x}$$

✓ Solution by Mathematica

Time used: 0.135 (sec). Leaf size: 23

```
DSolve[y'[x]+y[x]/x==x*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{x^2 - c_1 x}$$

$$y(x) \rightarrow 0$$

3 Program 25. Second order differential equations.

Test Excercise 25. page 1093

3.1	problem 1	78
3.2	problem 2	79
3.3	problem 3	80
3.4	problem 4	81
3.5	problem 5	82
3.6	problem 6	83
3.7	problem 7	84
3.8	problem 8	85

3.1 problem 1

Internal problem ID [5136]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Test Exercise 25. page 1093

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y' - 2y = 8$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-2*y(x)=8,y(x), singsol=all)
```

$$y(x) = e^{2x}c_2 + e^{-x}c_1 - 4$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 23

```
DSolve[y''[x]-y'[x]-2*y[x]==8,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1e^{-x} + c_2e^{2x} - 4$$

3.2 problem 2

Internal problem ID [5137]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Test Exercise 25. page 1093

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 4y = 10e^{3x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)-4*y(x)=10*exp(3*x),y(x), singsol=all)
```

$$y(x) = e^{2x}c_2 + e^{-2x}c_1 + 2e^{3x}$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 29

```
DSolve[y''[x]-4*y[x]==10*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(2e^{5x} + c_1e^{4x} + c_2)$$

3.3 problem 3

Internal problem ID [5138]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Test Exercise 25. page 1093

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 2y' + y = e^{-2x}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)+y(x)=exp(-2*x),y(x), singsol=all)
```

$$y(x) = c_2 e^{-x} + e^{-x} x c_1 + e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 24

```
DSolve[y''[x]+2*y'[x]+y[x]==Exp[-2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(1 + e^x(c_2 x + c_1))$$

3.4 problem 4

Internal problem ID [5139]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Test Exercise 25. page 1093

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 25y = 5x^2 + x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)+25*y(x)=5*x^2+x,y(x), singsol=all)
```

$$y(x) = \sin(5x) c_2 + \cos(5x) c_1 + \frac{x^2}{5} + \frac{x}{25} - \frac{2}{125}$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 34

```
DSolve[y''[x]+25*y[x]==5*x^2+x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{125}(25x^2 + 5x - 2) + c_1 \cos(5x) + c_2 \sin(5x)$$

3.5 problem 5

Internal problem ID [5140]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Test Exercise 25. page 1093

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 2y' + y = 4 \sin(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x)=4*sin(x),y(x), singsol=all)
```

$$y(x) = e^x c_2 + e^x x c_1 + 2 \cos(x)$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 21

```
DSolve[y''[x]-2*y'[x]+y[x]==4*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 \cos(x) + e^x(c_2 x + c_1)$$

3.6 problem 6

Internal problem ID [5141]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Test Exercise 25. page 1093

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 4y' + 5y = 2e^{-2x}$$

With initial conditions

$$[y(0) = 1, y'(0) = -2]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve([diff(y(x),x$2)+4*diff(y(x),x)+5*y(x)=2*exp(-2*x),y(0) = 1, D(y)(0) = -2],y(x), sings
```

$$y(x) = -e^{-2x}(\cos(x) - 2)$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 16

```
DSolve[{y''[x]+4*y'[x]+5*y[x]==2*Exp[-2*x]},{y[0]==1,y'[0]==-2},y[x],x,IncludeSingularSoluti
```

$$y(x) \rightarrow -e^{-2x}(\cos(x) - 2)$$

3.7 problem 7

Internal problem ID [5142]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Test Exercise 25. page 1093

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$3y'' - 2y' - y = 2x - 3$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(3*diff(y(x),x$2)-2*diff(y(x),x)-y(x)=2*x-3,y(x), singsol=all)
```

$$y(x) = e^{-\frac{x}{3}}c_2 + e^x c_1 - 2x + 7$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 26

```
DSolve[3*y''[x]-2*y'[x]-y[x]==2*x-3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2x + c_1 e^{-x/3} + c_2 e^x + 7$$

3.8 problem 8

Internal problem ID [5143]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Test Exercise 25. page 1093

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 6y' + 8y = 8e^{4x}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)-6*diff(y(x),x)+8*y(x)=8*exp(4*x),y(x), singsol=all)
```

$$y(x) = \left(\frac{e^{2x}(8x + c_1 - 4)}{2} + c_2 \right) e^{2x}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 27

```
DSolve[y''[x]-6*y'[x]+8*y[x]==8*Exp[4*x],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow c_1 e^{2x} + e^{4x}(4x - 2 + c_2)$$

4 Program 25. Second order differential equations. Further problems 25. page 1094

4.1	problem 1	87
4.2	problem 2	88
4.3	problem 3	89
4.4	problem 4	90
4.5	problem 5	91
4.6	problem 6	92
4.7	problem 7	93
4.8	problem 8	94
4.9	problem 9	95
4.10	problem 10	96
4.11	problem 12	97
4.12	problem 13	98
4.13	problem 14	99
4.14	problem 15	100
4.15	problem 16	101
4.16	problem 17	102
4.17	problem 18	103
4.18	problem 19	104
4.19	problem 20	105

4.1 problem 1

Internal problem ID [5144]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2y'' - 7y' - 4y = e^{3x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(2*diff(y(x),x$2)-7*diff(y(x),x)-4*y(x)=exp(3*x),y(x), singsol=all)
```

$$y(x) = c_2 e^{4x} + e^{-\frac{x}{2}} c_1 - \frac{e^{3x}}{7}$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 33

```
DSolve[2*y''[x]-7*y'[x]-4*y[x]==Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{e^{3x}}{7} + c_1 e^{-x/2} + c_2 e^{4x}$$

4.2 problem 2

Internal problem ID [5145]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 6y' + 9y = 54x + 18$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$2)-6*diff(y(x),x)+9*y(x)=54*x+18,y(x), singsol=all)
```

$$y(x) = e^{3x}c_2 + e^{3x}xc_1 + 6x + 6$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 27

```
DSolve[y''[x]-6*y'[x]+9*y[x]==54*x+18,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{3x} + x(6 + c_2 e^{3x}) + 6$$

4.3 problem 3

Internal problem ID [5146]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 5y' + 6y = 100 \sin(4x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$2)-5*diff(y(x),x)+6*y(x)=100*sin(4*x),y(x), singsol=all)
```

$$y(x) = e^{2x}c_2 + c_1e^{3x} - 2 \sin(4x) + 4 \cos(4x)$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 33

```
DSolve[y''[x]-5*y'[x]+6*y[x]==100*Sin[4*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2 \sin(4x) + 4 \cos(4x) + e^{2x}(c_2e^x + c_1)$$

4.4 problem 4

Internal problem ID [5147]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 2y' + y = 4 \sinh(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 38

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)+y(x)=4*sinh(x),y(x), singsol=all)
```

$$y(x) = c_2 e^{-x} + e^{-x} x c_1 + \frac{(-2x^2 + 2x + 1)e^{-x}}{2} + \frac{e^x}{2}$$

✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 31

```
DSolve[y''[x]+2*y'[x]+y[x]==4*Sinh[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^x}{2} + e^{-x}(-x^2 + c_2 x + c_1)$$

4.5 problem 5

Internal problem ID [5148]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y' - 2y = 2 \cosh(2x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

```
dsolve(diff(y(x),x$2)+diff(y(x),x)-2*y(x)=2*cosh(2*x),y(x), singsol=all)
```

$$y(x) = e^x c_2 + e^{-2x} c_1 + \frac{(-12x - 7) e^{-2x}}{36} + \frac{e^{2x}}{4}$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 39

```
DSolve[y''[x]+y'[x]-2*y[x]==2*Cosh[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{36} e^{-2x} (-12x + 9e^{4x} + 36c_2 e^{3x} - 4 + 36c_1)$$

4.6 problem 6

Internal problem ID [5149]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - y' + 10y = 20 - e^{2x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 38

```
dsolve(diff(y(x),x$2)-diff(y(x),x)+10*y(x)=20-exp(2*x),y(x), singsol=all)
```

$$y(x) = e^{\frac{x}{2}} \sin\left(\frac{\sqrt{39}x}{2}\right) c_2 + e^{\frac{x}{2}} \cos\left(\frac{\sqrt{39}x}{2}\right) c_1 + 2 - \frac{e^{2x}}{12}$$

✓ Solution by Mathematica

Time used: 1.291 (sec). Leaf size: 58

```
DSolve[y''[x]-y'[x]+10*y[x]==20-Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{e^{2x}}{12} + c_2 e^{x/2} \cos\left(\frac{\sqrt{39}x}{2}\right) + c_1 e^{x/2} \sin\left(\frac{\sqrt{39}x}{2}\right) + 2$$

4.7 problem 7

Internal problem ID [5150]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y' + 4y = 2 \cos(x)^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)+4*y(x)=2*cos(x)^2,y(x), singsol=all)
```

$$y(x) = c_2 e^{-2x} + e^{-2x} x c_1 + \frac{1}{4} + \frac{\sin(2x)}{8}$$

✓ Solution by Mathematica

Time used: 0.132 (sec). Leaf size: 29

```
DSolve[y''[x]+4*y'[x]+4*y[x]==2*Cos[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{8}(\sin(2x) + 8e^{-2x}(c_2 x + c_1) + 2)$$

4.8 problem 8

Internal problem ID [5151]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 4y' + 3y = x + e^{2x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+3*y(x)=x+exp(2*x),y(x), singsol=all)
```

$$y(x) = e^x c_2 + c_1 e^{3x} - e^{2x} + \frac{x}{3} + \frac{4}{9}$$

✓ Solution by Mathematica

Time used: 0.129 (sec). Leaf size: 35

```
DSolve[y''[x]-4*y'[x]+3*y[x]==x+Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x}{3} - e^{2x} + c_1 e^x + c_2 e^{3x} + \frac{4}{9}$$

4.9 problem 9

Internal problem ID [5152]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 2y' + 3y = x^2 - 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+3*y(x)=x^2-1,y(x), singsol=all)
```

$$y(x) = e^x \sin(\sqrt{2}x) c_2 + e^x \cos(\sqrt{2}x) c_1 + \frac{x^2}{3} + \frac{4x}{9} - \frac{7}{27}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 48

```
DSolve[y''[x]-2*y'[x]+3*y[x]==x^2-1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{27}(9x^2 + 12x - 7) + c_2 e^x \cos(\sqrt{2}x) + c_1 e^x \sin(\sqrt{2}x)$$

4.10 problem 10

Internal problem ID [5153]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 9y = e^{3x} + \sin(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
dsolve(diff(y(x),x$2)-9*y(x)=exp(3*x)+sin(x),y(x), singsol=all)
```

$$y(x) = e^{-3x}c_2 + c_1e^{3x} + \frac{(-1 + 6x)e^{3x}}{36} - \frac{\sin(x)}{10}$$

✓ Solution by Mathematica

Time used: 0.126 (sec). Leaf size: 37

```
DSolve[y''[x]-9*y[x]==Exp[3*x]+Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sin(x)}{10} + e^{3x}\left(\frac{x}{6} - \frac{1}{36} + c_1\right) + c_2e^{-3x}$$

4.11 problem 12

Internal problem ID [5154]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x'' + 4x' + 3x = e^{-3t}$$

With initial conditions

$$\left[x(0) = \frac{1}{2}, x'(0) = -2 \right]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve([diff(x(t),t$2)+4*diff(x(t),t)+3*x(t)=exp(-3*t),x(0) = 1/2, D(x)(0) = -2],x(t), sings
```

$$x(t) = -\frac{e^{-3t}(t-1)}{2}$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 17

```
DSolve[{x''[t]+4*x'[t]+3*x[t]==Exp[-3*t],{x[0]==1/2,x'[0]==-2}},x[t],t,IncludeSingularSoluti
```

$$x(t) \rightarrow -\frac{1}{2}e^{-3t}(t-1)$$

4.12 problem 13

Internal problem ID [5155]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y' + 5y = 6 \sin(t)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(t),t$2)+4*diff(y(t),t)+5*y(t)=6*sin(t),y(t), singsol=all)
```

$$y(t) = e^{-2t} \sin(t) c_2 + e^{-2t} \cos(t) c_1 - \frac{3 \cos(t)}{4} + \frac{3 \sin(t)}{4}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 36

```
DSolve[y''[t]+4*y'[t]+5*y[t]==6*Sin[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \left(-\frac{3}{4} + c_2 e^{-2t}\right) \cos(t) + \left(\frac{3}{4} + c_1 e^{-2t}\right) \sin(t)$$

4.13 problem 14

Internal problem ID [5156]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x'' - 3x' + 2x = \sin(t)$$

With initial conditions

$$[x(0) = 0, x'(0) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve([diff(x(t),t$2)-3*diff(x(t),t)+2*x(t)=sin(t),x(0) = 0, D(x)(0) = 0],x(t), singsol=all
```

$$x(t) = \frac{e^{2t}}{5} + \frac{3 \cos(t)}{10} + \frac{\sin(t)}{10} - \frac{e^t}{2}$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 27

```
DSolve[{x''[t]-3*x'[t]+2*x[t]==Sin[t],{x[0]==0,x'[0]==0}},x[t],t,IncludeSingularSolutions ->
```

$$x(t) \rightarrow \frac{1}{10}(e^t(2e^t - 5) + \sin(t) + 3 \cos(t))$$

4.14 problem 15

Internal problem ID [5157]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 3y' + 2y = 3 \sin(x)$$

With initial conditions

$$\left[y(0) = -\frac{9}{10}, y'(0) = -\frac{7}{10} \right]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve([diff(y(x),x$2)+3*diff(y(x),x)+2*y(x)=3*sin(x),y(0) = -9/10, D(y)(0) = -7/10],y(x), s
```

$$y(x) = e^{-2x} - \frac{9 \cos(x)}{10} + \frac{3 \sin(x)}{10} - e^{-x}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 30

```
DSolve[{y'[x]+3*y'[x]+2*y[x]==3*Sin[x],{y[0]==-9/10,y'[0]==-7/10}},y[x],x,IncludeSingularSo
```

$$y(x) \rightarrow -e^{-2x}(e^x - 1) + \frac{3 \sin(x)}{10} - \frac{9 \cos(x)}{10}$$

4.15 problem 16

Internal problem ID [5158]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 6y' + 10y = 50x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)+6*diff(y(x),x)+10*y(x)=50*x,y(x), singsol=all)
```

$$y(x) = e^{-3x} \sin(x) c_2 + e^{-3x} \cos(x) c_1 + 5x - 3$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 30

```
DSolve[y''[x]+6*y'[x]+10*y[x]==50*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 5x + c_2 e^{-3x} \cos(x) + c_1 e^{-3x} \sin(x) - 3$$

4.16 problem 17

Internal problem ID [5159]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 17.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x'' + 2x' + 2x = 85 \sin(3t)$$

With initial conditions

$$[x(0) = 0, x'(0) = -20]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 33

```
dsolve([diff(x(t),t$2)+2*diff(x(t),t)+2*x(t)=85*sin(3*t),x(0) = 0, D(x)(0) = -20],x(t), sing
```

$$x(t) = (7 \sin(t) + 6 \cos(t)) e^{-t} - 6 \cos(3t) - 7 \sin(3t)$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 36

```
DSolve[{x'[t]+2*x'[t]+2*x[t]==85*Sin[3*t]},{x[0]==0,x'[0]==-20}],x[t],t,IncludeSingularSolut
```

$$x(t) \rightarrow 7e^{-t} \sin(t) - 7 \sin(3t) + 6e^{-t} \cos(t) - 6 \cos(3t)$$

4.17 problem 18

Internal problem ID [5160]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 18.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y = 3 \sin(x)$$

With initial conditions

$$\left[y(0) = 0, y'\left(\frac{\pi}{2}\right) = 1 \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
dsolve([diff(y(x),x$2)=3*sin(x)-4*y(x),y(0) = 0, D(y)(1/2*Pi) = 1],y(x), singsol=all)
```

$$y(x) = -\frac{\sin(2x)}{2} + \sin(x)$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 13

```
DSolve[{y''[x]==3*Sin[x]-4*y[x],{y[0]==0,y'[Pi/2]==1}},y[x],x,IncludeSingularSolutions -> Tr
```

$$y(x) \rightarrow -(\sin(x)(\cos(x) - 1))$$

4.18 problem 19

Internal problem ID [5161]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$\frac{x''}{2} + 48x = 0$$

With initial conditions

$$\left[x(0) = \frac{1}{6}, x'(0) = 0 \right]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve([1/2*diff(x(t),t$2)=-48*x(t),x(0) = 1/6, D(x)(0) = 0],x(t), singsol=all)
```

$$x(t) = \frac{\cos(4\sqrt{6}t)}{6}$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 18

```
DSolve[{1/2*x''[t]==-48*x[t],{x[0]==1/6,x'[0]==0}},x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{1}{6} \cos(4\sqrt{6}t)$$

4.19 problem 20

Internal problem ID [5162]

Book: Engineering Mathematics. By K. A. Stroud. 5th edition. Industrial press Inc. NY. 2001

Section: Program 25. Second order differential equations. Further problems 25. page 1094

Problem number: 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x'' + 5x' + 6x = \cos(t)$$

With initial conditions

$$\left[x(0) = \frac{1}{10}, x'(0) = 0 \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 25

```
dsolve([diff(x(t),t$2)+5*diff(x(t),t)+6*x(t)=cos(t),x(0) = 1/10, D(x)(0) = 0],x(t), singsol=
```

$$x(t) = \frac{e^{-3t}}{10} - \frac{e^{-2t}}{10} + \frac{\cos(t)}{10} + \frac{\sin(t)}{10}$$

✓ Solution by Mathematica

Time used: 0.059 (sec). Leaf size: 26

```
DSolve[{x''[t]+5*x'[t]+6*x[t]==Cos[t],{x[0]==1/10,x'[0]==0}},x[t],t,IncludeSingularSolutions
```

$$x(t) \rightarrow \frac{1}{10}(e^{-3t} - e^{-2t} + \sin(t) + \cos(t))$$