


A Solution Manual For

**Notes on Diffy Qs. Differential
Equations for Engineers. By by
Jiri Lebl, 2013.**



Nasser M. Abbasi

March 3, 2024

Contents

1	Chapter 7. POWER SERIES METHODS. 7.2.1 Exercises. page 290	2
2	Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300	15

1 Chapter 7. POWER SERIES METHODS. 7.2.1 Exercises. page 290

1.1	problem 7.2.1	3
1.2	problem 7.2.2	4
1.3	problem 7.2.3	5
1.4	problem 7.2.4	6
1.5	problem 7.2.5	7
1.6	problem 7.2.6	8
1.7	problem 7.2.7	9
1.8	problem 7.2.8 part(a)	10
1.9	problem 7.2.8 part(b)	11
1.10	problem 7.2.101	12
1.11	problem 7.2.102	13
1.12	problem 7.2.103	14

1.1 problem 7.2.1

Internal problem ID [5503]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.2.1 Exercises. page 290

Problem number: 7.2.1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y = 0$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)+y(x)=0,y(x),type='series',x=1);
```

$$y(x) = \left(1 - \frac{(x-1)^2}{2} + \frac{(x-1)^4}{24}\right) y(1) + \left(x-1 - \frac{(x-1)^3}{6} + \frac{(x-1)^5}{120}\right) D(y)(1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 51

```
AsymptoticDSolveValue[y''[x]+y[x]==0,y[x],{x,1,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{24}(x-1)^4 - \frac{1}{2}(x-1)^2 + 1 \right) + c_2 \left(\frac{1}{120}(x-1)^5 - \frac{1}{6}(x-1)^3 + x-1 \right)$$

1.2 problem 7.2.2

Internal problem ID [5504]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.2.1 Exercises. page 290

Problem number: 7.2.2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y'' + 4yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)+4*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{2x^3}{3}\right) y(0) + \left(x - \frac{1}{3}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y''[x]+4*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{x^4}{3}\right) + c_1 \left(1 - \frac{2x^3}{3}\right)$$

1.3 problem 7.2.3

Internal problem ID [5505]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.2.1 Exercises. page 290

Problem number: 7.2.3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y'' - yx = 0$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

```
Order:=6;  
dsolve(diff(y(x),x$2)-x*y(x)=0,y(x),type='series',x=1);
```

$$y(x) = \left(1 + \frac{(x-1)^2}{2} + \frac{(x-1)^3}{6} + \frac{(x-1)^4}{24} + \frac{(x-1)^5}{30}\right) y(1) \\ + \left(x-1 + \frac{(x-1)^3}{6} + \frac{(x-1)^4}{12} + \frac{(x-1)^5}{120}\right) D(y)(1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 78

```
AsymptoticDSolveValue[y''[x]-x*y[x]==0,y[x],{x,1,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{30}(x-1)^5 + \frac{1}{24}(x-1)^4 + \frac{1}{6}(x-1)^3 + \frac{1}{2}(x-1)^2 + 1 \right) \\ + c_2 \left(\frac{1}{120}(x-1)^5 + \frac{1}{12}(x-1)^4 + \frac{1}{6}(x-1)^3 + x-1 \right)$$

1.4 problem 7.2.4

Internal problem ID [5506]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.2.1 Exercises. page 290

Problem number: 7.2.4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_Emden, _Fowler]`

$$y'' + yx^2 = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)+x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{x^4}{12}\right) y(0) + \left(x - \frac{1}{20}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y'[x]+x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{x^5}{20}\right) + c_1 \left(1 - \frac{x^4}{12}\right)$$

1.5 problem 7.2.5

Internal problem ID [5507]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.2.1 Exercises. page 290

Problem number: 7.2.5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
Order:=6;  
dsolve(diff(y(x),x)-x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 + \frac{1}{8}x^4\right) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 22

```
AsymptoticDSolveValue[y'[x]-x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^4}{8} + \frac{x^2}{2} + 1 \right)$$

1.6 problem 7.2.6

Internal problem ID [5508]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.2.1 Exercises. page 290

Problem number: 7.2.6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [_Gegenbauer, [_2nd_order, _linear, '_with_symmetry_[0,F(x)]']

$$(-x^2 + 1)y'' - y'x + p^2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 71

```
Order:=6;
```

```
dsolve((1-x^2)*diff(y(x),x$2)-x*diff(y(x),x)+p^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{p^2 x^2}{2} + \frac{p^2(p^2 - 4)x^4}{24}\right) y(0) \\ + \left(x - \frac{(p^2 - 1)x^3}{6} + \frac{(p^4 - 10p^2 + 9)x^5}{120}\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 88

```
AsymptoticDSolveValue[(1-x^2)*y''[x]-x*y'[x]+p^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{p^4 x^5}{120} - \frac{p^2 x^5}{12} - \frac{p^2 x^3}{6} + \frac{3x^5}{40} + \frac{x^3}{6} + x \right) + c_1 \left(\frac{p^4 x^4}{24} - \frac{p^2 x^4}{6} - \frac{p^2 x^2}{2} + 1 \right)$$

1.7 problem 7.2.7

Internal problem ID [5509]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.2.1 Exercises. page 290

Problem number: 7.2.7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 1) y'' - 2y'x + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
Order:=6;  
dsolve((1+x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = y(0) + D(y)(0)x - x^2y(0)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 18

```
AsymptoticDSolveValue[(1+x^2)*y''[x]-2*x*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(1 - x^2) + c_2x$$

1.8 problem 7.2.8 part(a)

Internal problem ID [5510]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.2.1 Exercises. page 290

Problem number: 7.2.8 part(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$(x^2 + 1)y'' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve((x^2+1)*diff(y(x),x$2)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4\right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{7}{120}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[(x^2+1)*y''[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{7x^5}{120} - \frac{x^3}{6} + x \right) + c_1 \left(\frac{x^4}{8} - \frac{x^2}{2} + 1 \right)$$

1.9 problem 7.2.8 part(b)

Internal problem ID [5511]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.2.1 Exercises. page 290

Problem number: 7.2.8 part(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$xy'' + y = 0$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

```
Order:=6;  
dsolve(x*diff(y(x),x$2)+y(x)=0,y(x),type='series',x=1);
```

$$y(x) = \left(1 - \frac{(x-1)^2}{2} + \frac{(x-1)^3}{6} - \frac{(x-1)^4}{24} + \frac{(x-1)^5}{60}\right) y(1) \\ + \left(x - 1 - \frac{(x-1)^3}{6} + \frac{(x-1)^4}{12} - \frac{(x-1)^5}{24}\right) D(y)(1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 78

```
AsymptoticDSolveValue[x*y''[x]+y[x]==0,y[x],{x,1,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{60}(x-1)^5 - \frac{1}{24}(x-1)^4 + \frac{1}{6}(x-1)^3 - \frac{1}{2}(x-1)^2 + 1 \right) \\ + c_2 \left(-\frac{1}{24}(x-1)^5 + \frac{1}{12}(x-1)^4 - \frac{1}{6}(x-1)^3 + x - 1 \right)$$

1.10 problem 7.2.101

Internal problem ID [5512]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.2.1 Exercises. page 290

Problem number: 7.2.101.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y'' + 2yx^3 = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
Order:=6;  
dsolve(diff(y(x),x$2)+2*x^3*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{x^5}{10}\right) y(0) + D(y)(0)x + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 20

```
AsymptoticDSolveValue[y'[x]+2*x^3*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(1 - \frac{x^5}{10}\right) + c_2 x$$

1.11 problem 7.2.102

Internal problem ID [5513]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.2.1 Exercises. page 290

Problem number: 7.2.102.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - yx = \frac{1}{1-x}$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
Order:=6;  
dsolve([diff(y(x),x$2)-x*y(x)=1/(1-x),y(0) = 0, D(y)(0) = 0],y(x),type='series',x=0);
```

$$y(x) = \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{3}{40}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 56

```
AsymptoticDSolveValue[{y'[x]-x*y[x]==1/(1-x),{}} ,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{3x^5}{40} + \frac{x^4}{12} + c_2 \left(\frac{x^4}{12} + x \right) + \frac{x^3}{6} + c_1 \left(\frac{x^3}{6} + 1 \right) + \frac{x^2}{2}$$

1.12 problem 7.2.103

Internal problem ID [5514]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.2.1 Exercises. page 290

Problem number: 7.2.103.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \sqrt{x} \left(x^{-\frac{\sqrt{5}}{2}} c_1 + x^{\frac{\sqrt{5}}{2}} c_2 \right) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 38

```
AsymptoticDSolveValue[x^2*y''[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x^{\frac{1}{2}(1+\sqrt{5})} + c_2 x^{\frac{1}{2}(1-\sqrt{5})}$$

2 Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

2.1	problem 7.3.3	16
2.2	problem 7.3.4	18
2.3	problem 7.3.5	19
2.4	problem 7.3.6	20
2.5	problem 7.3.7	21
2.6	problem 7.3.8 (a)	22
2.7	problem 7.3.8 (b)	24
2.8	problem 7.3.8 (c)	25
2.9	problem 7.3.8 (d)	27
2.10	problem 7.3.8 (e)	28
2.11	problem 7.3.101 (a)	29
2.12	problem 7.3.101 (b)	30
2.13	problem 7.3.101 (c)	31
2.14	problem 7.3.101 (d)	32
2.15	problem 7.3.101 (e)	33
2.16	problem 7.3.102	34
2.17	problem 7.3.103	35
2.18	problem 7.3.104 (d)	36

2.1 problem 7.3.3

Internal problem ID [5515]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2y'' + y'x + (1+x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 69

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(1+x)*y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1 x^{-i} \left(1 + \left(-\frac{1}{5} - \frac{2i}{5} \right) x + \left(-\frac{1}{40} + \frac{3i}{40} \right) x^2 + \left(\frac{3}{520} - \frac{7i}{1560} \right) x^3 \right. \\ & \left. + \left(-\frac{1}{2496} + \frac{i}{12480} \right) x^4 + \left(\frac{9}{603200} + \frac{i}{361920} \right) x^5 + O(x^6) \right) \\ & + c_2 x^i \left(1 + \left(-\frac{1}{5} + \frac{2i}{5} \right) x + \left(-\frac{1}{40} - \frac{3i}{40} \right) x^2 + \left(\frac{3}{520} + \frac{7i}{1560} \right) x^3 \right. \\ & \left. + \left(-\frac{1}{2496} - \frac{i}{12480} \right) x^4 + \left(\frac{9}{603200} - \frac{i}{361920} \right) x^5 + O(x^6) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 90

```
AsymptoticDSolveValue[x^2*y'[x]+x*y'[x]+(1+x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \left(\frac{1}{12480} + \frac{i}{2496} \right) c_2 x^{-i} (ix^4 - (8 + 16i)x^3 + (168 + 96i)x^2 - (1056 - 288i)x + (480 - 2400i)) - \left(\frac{1}{2496} + \frac{i}{12480} \right) c_1 x^i (x^4 - (16 + 8i)x^3 + (96 + 168i)x^2 + (288 - 1056i)x - (2400 - 480i))$$

2.2 problem 7.3.4

Internal problem ID [5516]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \sqrt{x} \left(x^{-\frac{\sqrt{5}}{2}} c_1 + x^{\frac{\sqrt{5}}{2}} c_2 \right) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 38

```
AsymptoticDSolveValue[x^2*y''[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x^{\frac{1}{2}(1+\sqrt{5})} + c_2 x^{\frac{1}{2}(1-\sqrt{5})}$$

2.3 problem 7.3.5

Internal problem ID [5517]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y'' + \frac{y'}{x} - yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
Order:=6;  
dsolve(diff(y(x),x$2)+1/x*diff(y(x),x)-x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (\ln(x) c_2 + c_1) \left(1 + \frac{1}{9}x^3 + O(x^6) \right) + \left(-\frac{2}{27}x^3 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 39

```
AsymptoticDSolveValue[y''[x]+1/x*y'[x]-x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^3}{9} + 1 \right) + c_2 \left(\left(\frac{x^3}{9} + 1 \right) \log(x) - \frac{2x^3}{27} \right)$$

2.4 problem 7.3.6

Internal problem ID [5518]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$2xy'' + y' - yx^2 = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 28

```
Order:=6;  
dsolve(2*x*diff(y(x),x$2)+diff(y(x),x)-x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1\sqrt{x} \left(1 + \frac{1}{21}x^3 + O(x^6)\right) + c_2 \left(1 + \frac{1}{15}x^3 + O(x^6)\right)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 33

```
AsymptoticDSolveValue[2*x*y''[x]+y'[x]-x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1\sqrt{x} \left(\frac{x^3}{21} + 1\right) + c_2 \left(\frac{x^3}{15} + 1\right)$$

2.5 problem 7.3.7

Internal problem ID [5519]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2y'' - y'x - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 37

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)-x*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = x\left(x^{-\sqrt{2}}c_1 + x^{\sqrt{2}}c_2\right) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
AsymptoticDSolveValue[x^2*y''[x]-x*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1x^{1+\sqrt{2}} + c_2x^{1-\sqrt{2}}$$

2.6 problem 7.3.8 (a)

Internal problem ID [5520]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.8 (a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 1)y'' + yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 58

```
Order:=6;  
dsolve(x^2*(1+x^2)*diff(y(x),x$2)+x*y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1 x \left(1 - \frac{1}{2}x + \frac{1}{12}x^2 + \frac{11}{144}x^3 - \frac{83}{2880}x^4 - \frac{2557}{86400}x^5 + O(x^6) \right) \\ & + c_2 \left(\ln(x) \left(-x + \frac{1}{2}x^2 - \frac{1}{12}x^3 - \frac{11}{144}x^4 + \frac{83}{2880}x^5 + O(x^6) \right) \right. \\ & \left. + \left(1 - \frac{3}{4}x^2 + \frac{13}{36}x^3 + \frac{25}{1728}x^4 - \frac{8743}{86400}x^5 + O(x^6) \right) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 87

```
AsymptoticDSolveValue[x^2*(1+x^2)*y'[x]+x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{157x^4 + 768x^3 - 2160x^2 + 1728x + 1728}{1728} - \frac{1}{144}x(11x^3 + 12x^2 - 72x + 144) \log(x) \right) + c_2 \left(-\frac{83x^5}{2880} + \frac{11x^4}{144} + \frac{x^3}{12} - \frac{x^2}{2} + x \right)$$

2.7 problem 7.3.8 (b)

Internal problem ID [5521]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.8 (b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2y'' + y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

X Solution by Maple

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)+diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

No solution found

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 84

```
AsymptoticDSolveValue[x^2*y'[x]+y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 e^{\frac{1}{x}} \left(\frac{59241x^5}{40} + \frac{1911x^4}{8} + \frac{91x^3}{2} + \frac{21x^2}{2} + 3x + 1 \right) x^2 \\ + c_1 \left(-\frac{91x^5}{40} + \frac{7x^4}{8} - \frac{x^3}{2} + \frac{x^2}{2} - x + 1 \right)$$

2.8 problem 7.3.8 (c)

Internal problem ID [5522]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.8 (c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$xy'' + y'x^3 + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 58

```
Order:=6;  
dsolve(x*dif(y(x),x$2)+x^3*dif(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1 x \left(1 - \frac{1}{2}x + \frac{1}{12}x^2 - \frac{13}{144}x^3 + \frac{157}{2880}x^4 - \frac{877}{86400}x^5 + O(x^6) \right) \\ & + c_2 \left(\ln(x) \left(-x + \frac{1}{2}x^2 - \frac{1}{12}x^3 + \frac{13}{144}x^4 - \frac{157}{2880}x^5 + O(x^6) \right) \right. \\ & \quad \left. + \left(1 - \frac{3}{4}x^2 + \frac{7}{36}x^3 + \frac{25}{1728}x^4 + \frac{6377}{86400}x^5 + O(x^6) \right) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 87

```
AsymptoticDSolveValue[x*y'[x]+x^3*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{144} x (13x^3 - 12x^2 + 72x - 144) \log(x) + \frac{-131x^4 + 480x^3 - 2160x^2 + 1728x + 1728}{1728} \right) + c_2 \left(\frac{157x^5}{2880} - \frac{13x^4}{144} + \frac{x^3}{12} - \frac{x^2}{2} + x \right)$$

2.9 problem 7.3.8 (d)

Internal problem ID [5523]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.8 (d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$xy'' + y'x - e^x y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 54

```
Order:=6;  
dsolve(x*dif(y(x),x$2)+x*dif(y(x),x)-exp(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x \left(1 + \frac{1}{6}x^2 + \frac{1}{72}x^3 + \frac{7}{480}x^4 + \frac{29}{10800}x^5 + O(x^6) \right) \\ + c_2 \left(\ln(x) \left(x + \frac{1}{6}x^3 + \frac{1}{72}x^4 + \frac{7}{480}x^5 + O(x^6) \right) \right. \\ \left. + \left(1 - x - \frac{2}{9}x^3 - \frac{11}{864}x^4 - \frac{109}{4800}x^5 + O(x^6) \right) \right)$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 70

```
AsymptoticDSolveValue[x*y''[x]+x*y'[x]-Exp[x]*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{7x^5}{480} + \frac{x^4}{72} + \frac{x^3}{6} + x \right) \\ + c_1 \left(\frac{1}{864} (-23x^4 - 336x^3 - 1728x + 864) + \frac{1}{72} x (x^3 + 12x^2 + 72) \log(x) \right)$$

2.10 problem 7.3.8 (e)

Internal problem ID [5524]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.8 (e).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$x^2 y'' + x^2 y' + y x^2 = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)+x^2*diff(y(x),x)+x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{120}x^5\right) y(0) + \left(x - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{120}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 56

```
AsymptoticDSolveValue[x^2*y''[x]+x^2*y'[x]+x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(-\frac{x^5}{120} + \frac{x^4}{24} - \frac{x^2}{2} + x \right) + c_1 \left(-\frac{x^5}{120} + \frac{x^3}{6} - \frac{x^2}{2} + 1 \right)$$

2.11 problem 7.3.101 (a)

Internal problem ID [5525]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.101 (a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(\frac{1}{24}x^4 - \frac{1}{2}x^2 + 1 \right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5 \right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[y''[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{120} - \frac{x^3}{6} + x \right) + c_1 \left(\frac{x^4}{24} - \frac{x^2}{2} + 1 \right)$$

2.12 problem 7.3.101 (b)

Internal problem ID [5526]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.101 (b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^3 y'' + (1+x)y = 0$$

With the expansion point for the power series method at $x = 0$.

X Solution by Maple

```
Order:=6;
dsolve(x^3*diff(y(x),x$2)+(1+x)*y(x)=0,y(x),type='series',x=0);
```

No solution found

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 222

```
AsymptoticDSolveValue[x^3*y'[x]+(1+x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 e^{-\frac{2i}{\sqrt{x}} x^{3/4}} \left(\frac{520667425699057ix^{9/2}}{131941395333120} - \frac{21896102683ix^{7/2}}{21474836480} + \frac{19100991ix^{5/2}}{41943040} - \frac{3367ix^{3/2}}{8192} - \frac{194208949785748261x^5}{21110623253299200} + \frac{5189376335871x^4}{2748779069440} - \frac{846810601x^3}{1342177280} + \frac{205387x^2}{524288} - \frac{273x}{512} + \frac{13i\sqrt{x}}{16} \right) + c_2 e^{\frac{2i}{\sqrt{x}} x^{3/4}} \left(-\frac{520667425699057ix^{9/2}}{131941395333120} + \frac{21896102683ix^{7/2}}{21474836480} - \frac{19100991ix^{5/2}}{41943040} + \frac{3367ix^{3/2}}{8192} - \frac{1942089}{21110623253299200} \right)$$

2.13 problem 7.3.101 (c)

Internal problem ID [5527]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.101 (c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$xy'' + x^5y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 58

```
Order:=6;
dsolve(x*dif(y(x),x$2)+x^5*dif(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 \left(1 - \frac{1}{2}x + \frac{1}{12}x^2 - \frac{1}{144}x^3 + \frac{1}{2880}x^4 - \frac{2881}{86400}x^5 + O(x^6) \right) \\ + c_2 \left(\ln(x) \left(-x + \frac{1}{2}x^2 - \frac{1}{12}x^3 + \frac{1}{144}x^4 - \frac{1}{2880}x^5 + O(x^6) \right) \right. \\ \left. + \left(1 - \frac{3}{4}x^2 + \frac{7}{36}x^3 - \frac{35}{1728}x^4 + \frac{101}{86400}x^5 + O(x^6) \right) \right)$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 85

```
AsymptoticDSolveValue[x*y''[x]+x^5*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{144}x(x^3 - 12x^2 + 72x - 144) \log(x) \right. \\ \left. + \frac{-47x^4 + 480x^3 - 2160x^2 + 1728x + 1728}{1728} \right) + c_2 \left(\frac{x^5}{2880} - \frac{x^4}{144} + \frac{x^3}{12} - \frac{x^2}{2} + x \right)$$

2.14 problem 7.3.101 (d)

Internal problem ID [5528]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.101 (d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$\sin(x)y'' - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.219 (sec). Leaf size: 58

```
Order:=6;  
dsolve(sin(x)*diff(y(x),x$2)-y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1x \left(1 + \frac{1}{2}x + \frac{1}{12}x^2 + \frac{1}{48}x^3 + \frac{1}{192}x^4 + \frac{37}{28800}x^5 + O(x^6) \right) \\ & + c_2 \left(\ln(x) \left(x + \frac{1}{2}x^2 + \frac{1}{12}x^3 + \frac{1}{48}x^4 + \frac{1}{192}x^5 + O(x^6) \right) \right. \\ & \left. + \left(1 - \frac{3}{4}x^2 - \frac{1}{6}x^3 - \frac{5}{192}x^4 - \frac{257}{28800}x^5 + O(x^6) \right) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 85

```
AsymptoticDSolveValue[Sin[x]*y''[x]-y[x]==0,y[x],{x,0,5}]
```

$$\begin{aligned} y(x) \rightarrow & c_1 \left(\frac{1}{48}x(x^3 + 4x^2 + 24x + 48) \log(x) + \frac{1}{64}(-3x^4 - 16x^3 - 80x^2 - 64x + 64) \right) \\ & + c_2 \left(\frac{x^5}{192} + \frac{x^4}{48} + \frac{x^3}{12} + \frac{x^2}{2} + x \right) \end{aligned}$$

2.15 problem 7.3.101 (e)

Internal problem ID [5529]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.101 (e).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$\cos(x)y'' - \sin(x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 29

```
Order:=6;  
dsolve(cos(x)*diff(y(x),x$2)-sin(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{6}x^3 + \frac{1}{60}x^5\right) y(0) + \left(x + \frac{1}{12}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 35

```
AsymptoticDSolveValue[Cos[x]*y''[x]-Sin[x]*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^4}{12} + x \right) + c_1 \left(\frac{x^5}{60} + \frac{x^3}{6} + 1 \right)$$

2.16 problem 7.3.102

Internal problem ID [5530]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.102.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \sqrt{x} \left(x^{-\frac{\sqrt{5}}{2}} c_1 + x^{\frac{\sqrt{5}}{2}} c_2 \right) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 38

```
AsymptoticDSolveValue[x^2*y''[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x^{\frac{1}{2}(1+\sqrt{5})} + c_2 x^{\frac{1}{2}(1-\sqrt{5})}$$

2.17 problem 7.3.103

Internal problem ID [5531]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.103.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + \left(x - \frac{3}{4}\right) y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 65

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)+(x-3/4)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 x^2 \left(1 - \frac{1}{3}x + \frac{1}{24}x^2 - \frac{1}{360}x^3 + \frac{1}{8640}x^4 - \frac{1}{302400}x^5 + O(x^6)\right) + c_2 (\ln(x) \left(x^2 - \frac{1}{3}x^3 + \frac{1}{24}x^4 - \frac{1}{360}x^5 + O(x^6)\right))}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 101

```
AsymptoticDSolveValue[x^2*y''[x]+(x-3/4)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^{11/2}}{8640} - \frac{x^{9/2}}{360} + \frac{x^{7/2}}{24} - \frac{x^{5/2}}{3} + x^{3/2} \right) + c_1 \left(\frac{31x^4 - 176x^3 + 144x^2 + 576x + 576}{576\sqrt{x}} - \frac{1}{48}x^{3/2}(x^2 - 8x + 24) \log(x) \right)$$

2.18 problem 7.3.104 (d)

Internal problem ID [5532]

Book: Notes on Diffy Qs. Differential Equations for Engineers. By by Jiri Lebl, 2013.

Section: Chapter 7. POWER SERIES METHODS. 7.3.2 The method of Frobenius. Exercises. page 300

Problem number: 7.3.104 (d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - y' x + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = x(\ln(x) c_2 + c_1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 14

```
AsymptoticDSolveValue[x^2*y''[x]-x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x + c_2 x \log(x)$$