


A Solution Manual For

**Ordinary Differential Equations,**  
**By Tenenbaum and Pollard.**  
**Dover, NY 1963**



**Nasser M. Abbasi**

March 3, 2024

# Contents

1	Chapter 2. Special types of differential equations of the first kind. Lesson 7	2
2	Chapter 2. Special types of differential equations of the first kind. Lesson 8	18
3	Chapter 2. Special types of differential equations of the first kind. Lesson 9	34
4	Chapter 2. Special types of differential equations of the first kind. Lesson 10	54
5	Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations	101
6	Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods	129
7	Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients	194
8	Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients	229
9	Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters	255
10	Chapter 8. Special second order equations. Lesson 35. Independent variable $x$ absent	276

# 1 Chapter 2. Special types of differential equations of the first kind. Lesson 7

1.1	problem First order with homogeneous Coefficients. Exercise 7.2, page 61 .	3
1.2	problem First order with homogeneous Coefficients. Exercise 7.3, page 61 .	5
1.3	problem First order with homogeneous Coefficients. Exercise 7.4, page 61 .	6
1.4	problem First order with homogeneous Coefficients. Exercise 7.5, page 61 .	7
1.5	problem First order with homogeneous Coefficients. Exercise 7.6, page 61 .	8
1.6	problem First order with homogeneous Coefficients. Exercise 7.7, page 61 .	9
1.7	problem First order with homogeneous Coefficients. Exercise 7.8, page 61 .	10
1.8	problem First order with homogeneous Coefficients. Exercise 7.9, page 61 .	11
1.9	problem First order with homogeneous Coefficients. Exercise 7.10, page 61	12
1.10	problem First order with homogeneous Coefficients. Exercise 7.11, page 61	13
1.11	problem First order with homogeneous Coefficients. Exercise 7.12, page 61	14
1.12	problem First order with homogeneous Coefficients. Exercise 7.13, page 61	15
1.13	problem First order with homogeneous Coefficients. Exercise 7.14, page 61	16
1.14	problem First order with homogeneous Coefficients. Exercise 7.15, page 61	17

## 1.1 problem First order with homogeneous Coefficients. Exercise 7.2, page 61

Internal problem ID [4427]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 7

**Problem number:** First order with homogeneous Coefficients. Exercise 7.2, page 61.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _exact, _rational, _dAlembert]`

$$2yx + (x^2 + y^2) y' = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 257

```
dsolve(2*x*y(x)+(x^2+y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\frac{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}{2} - \frac{2x^2c_1}{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}}{\sqrt{c_1}}$$

$$y(x) = \frac{-\frac{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}{4} + \frac{x^2c_1}{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}} - \frac{i\sqrt{3}\left(\frac{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}{2} + \frac{2x^2c_1}{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}\right)}{2}}{\sqrt{c_1}}$$

$$y(x) = \frac{-\frac{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}{4} + \frac{x^2c_1}{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}} + \frac{i\sqrt{3}\left(\frac{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}{2} + \frac{2x^2c_1}{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}\right)}{2}}{\sqrt{c_1}}$$

✓ Solution by Mathematica

Time used: 15.191 (sec). Leaf size: 401

`DSolve[2*x*y[x]+(x^2+y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}{\sqrt[3]{2}} - \frac{\sqrt[3]{2}x^2}{\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}$$

$$y(x) \rightarrow \frac{i2^{2/3}(\sqrt{3} + i)(\sqrt{4x^6 + e^{6c_1}} + e^{3c_1})^{2/3} + \sqrt[3]{2}(2 + 2i\sqrt{3})x^2}{4\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}$$

$$y(x) \rightarrow \frac{(1 - i\sqrt{3})x^2}{2^{2/3}\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}} - \frac{(1 + i\sqrt{3})\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}{2\sqrt[3]{2}}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow \frac{1}{2}\sqrt[6]{x^6} \left( \frac{(1 - i\sqrt{3})(x^6)^{2/3}}{x^4} - i\sqrt{3} - 1 \right)$$

$$y(x) \rightarrow \frac{1}{2}\sqrt[6]{x^6} \left( \frac{(1 + i\sqrt{3})(x^6)^{2/3}}{x^4} + i\sqrt{3} - 1 \right)$$

$$y(x) \rightarrow \sqrt[6]{x^6} - \frac{(x^6)^{5/6}}{x^4}$$

## 1.2 problem First order with homogeneous Coefficients. Exercise 7.3, page 61

Internal problem ID [4428]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 7

**Problem number:** First order with homogeneous Coefficients. Exercise 7.3, page 61.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _dAlembert]`

$$\left( x + \sqrt{y^2 - yx} \right) y' - y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve((x+sqrt(y(x)^2-x*y(x)))*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$\ln(y(x)) + \frac{2\sqrt{y(x)(y(x)-x)}}{y(x)} - c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 0.291 (sec). Leaf size: 43

```
DSolve[(x+Sqrt[y[x]^2-x*y[x]])*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ \frac{2\sqrt{\frac{y(x)}{x} - 1}}{\sqrt{\frac{y(x)}{x}}} + \log\left(\frac{y(x)}{x}\right) = -\log(x) + c_1, y(x) \right]$$

### 1.3 problem First order with homogeneous Coefficients. Exercise 7.4, page 61

Internal problem ID [4429]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 7

**Problem number:** First order with homogeneous Coefficients. Exercise 7.4, page 61.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$y - (-y + x)y' = -x$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
dsolve((x+y(x))-(-x-y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \tan \left( \text{RootOf} \left( -2\_Z + \ln \left( \frac{1}{\cos(\_Z)^2} \right) + 2 \ln(x) + 2c_1 \right) \right) x$$

#### ✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 36

```
DSolve[(x+y[x])-(-x-y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ \frac{1}{2} \log \left( \frac{y(x)^2}{x^2} + 1 \right) - \arctan \left( \frac{y(x)}{x} \right) = -\log(x) + c_1, y(x) \right]$$

## 1.4 problem First order with homogeneous Coefficients. Exercise 7.5, page 61

Internal problem ID [4430]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 7

**Problem number:** First order with homogeneous Coefficients. Exercise 7.5, page 61.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _dAlembert]`

$$xy' - y - x \sin\left(\frac{y}{x}\right) = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

```
dsolve(x*diff(y(x),x)-y(x)-x*sin(y(x)/x)=0,y(x), singsol=all)
```

$$y(x) = \arctan\left(\frac{2xc_1}{c_1^2x^2 + 1}, -\frac{c_1^2x^2 - 1}{c_1^2x^2 + 1}\right) x$$

### ✓ Solution by Mathematica

Time used: 0.325 (sec). Leaf size: 52

```
DSolve[x*y'[x]-y[x]-x*Sin[y[x]/x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x \arccos(-\tanh(\log(x) + c_1))$$

$$y(x) \rightarrow x \arccos(-\tanh(\log(x) + c_1))$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -\pi x$$

$$y(x) \rightarrow \pi x$$



## 1.5 problem First order with homogeneous Coefficients. Exercise 7.6, page 61

Internal problem ID [4431]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 7

**Problem number:** First order with homogeneous Coefficients. Exercise 7.6, page 61.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _dAlembert]`

$$2x^2y + y^3 + (y^2x - 2x^3)y' = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve((2*x^2*y(x)+y(x)^3)+(x*y(x)^2-2*x^3)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \sqrt{-\frac{2}{\text{LambertW}(-2c_1x^4)}} x$$

### ✓ Solution by Mathematica

Time used: 5.64 (sec). Leaf size: 66

```
DSolve[(2*x^2*y[x]+y[x]^3)+(x*y[x]^2-2*x^3)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow -\frac{i\sqrt{2}x}{\sqrt{W(-2e^{-2c_1x^4})}}$$

$$y(x) \rightarrow \frac{i\sqrt{2}x}{\sqrt{W(-2e^{-2c_1x^4})}}$$

$$y(x) \rightarrow 0$$

**1.6 problem First order with homogeneous Coefficients.  
Exercise 7.7, page 61**

Internal problem ID [4432]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 7

**Problem number:** First order with homogeneous Coefficients. Exercise 7.7, page 61.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _dAlembert]`

$$y^2 + (x\sqrt{y^2 - x^2} - yx) y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve(y(x)^2+(x*sqrt(y(x)^2-x^2)-x*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$\frac{\sqrt{y(x)^2 - x^2}}{xy(x)} + \frac{1}{x} - c_1 = 0$$

✓ Solution by Mathematica

Time used: 2.247 (sec). Leaf size: 111

```
DSolve[y[x]^2+(x*Sqrt[y[x]^2-x^2]-x*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ \begin{aligned} & -\frac{\sqrt{\frac{y(x)^2}{x^2} - 1} \left( \log \left( \sqrt{\frac{y(x)}{x} + 1} - 1 \right) + \log \left( \sqrt{\frac{y(x)}{x} + 1} + 1 \right) \right)}{\sqrt{\frac{y(x)}{x} - 1} \sqrt{\frac{y(x)}{x} + 1}} \\ & - 2 \log \left( \sqrt{\frac{y(x)}{x} - 1} - \sqrt{\frac{y(x)}{x} + 1} \right) = \log(x) + c_1, y(x) \end{aligned} \right]$$

## 1.7 problem First order with homogeneous Coefficients. Exercise 7.8, page 61

Internal problem ID [4433]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 7

**Problem number:** First order with homogeneous Coefficients. Exercise 7.8, page 61.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _dAlembert]`

$$\frac{y \cos\left(\frac{y}{x}\right)}{x} - \left(\frac{x \sin\left(\frac{y}{x}\right)}{y} + \cos\left(\frac{y}{x}\right)\right) y' = 0$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 15

```
dsolve(y(x)/x*cos(y(x)/x)-(x/y(x)*sin(y(x)/x)+cos(y(x)/x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \text{RootOf}(\_Z x c_1 \sin(\_Z) - 1) x$$

### ✓ Solution by Mathematica

Time used: 0.247 (sec). Leaf size: 27

```
DSolve[y[x]/x*Cos[y[x]/x]-(x/y[x]*Sin[y[x]/x]+Cos[y[x]/x])*y'[x]==0,y[x],x,IncludeSingularSo
```

$$\text{Solve}\left[\log\left(\frac{y(x)}{x}\right) + \log\left(\sin\left(\frac{y(x)}{x}\right)\right) = -\log(x) + c_1, y(x)\right]$$

## 1.8 problem First order with homogeneous Coefficients. Exercise 7.9, page 61

Internal problem ID [4434]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 7

**Problem number:** First order with homogeneous Coefficients. Exercise 7.9, page 61.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _dAlembert]`

$$y + x \ln\left(\frac{y}{x}\right) y' - 2xy' = 0$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 18

```
dsolve(y(x)+x*ln(y(x)/x)*diff(y(x),x)-2*x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = e^{-\text{LambertW}(-exc_1)+1}x$$

### ✓ Solution by Mathematica

Time used: 5.502 (sec). Leaf size: 35

```
DSolve[y[x]+x*Log[y[x]/x]*y'[x]-2*x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -e^{c_1} W(-e^{1-c_1} x)$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow ex$$

## 1.9 problem First order with homogeneous Coefficients. Exercise 7.10, page 61

Internal problem ID [4435]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 7

**Problem number:** First order with homogeneous Coefficients. Exercise 7.10, page 61.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _dAlembert]`

$$2y e^{\frac{x}{y}} + \left(y - 2x e^{\frac{x}{y}}\right) y' = 0$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 23

```
dsolve(2*y(x)*exp(x/y(x))+(y(x)-2*x*exp(x/y(x)))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x}{\text{RootOf}\left(-\frac{Z e^{-2e^{-Z}}}{c_1} - x\right)}$$

### ✓ Solution by Mathematica

Time used: 0.247 (sec). Leaf size: 29

```
DSolve[2*y[x]*Exp[x/y[x]]+(y[x]-2*x*Exp[x/y[x]])*y'[x]==0,y[x],x,IncludeSingularSolutions ->
```

$$\text{Solve}\left[-2e^{\frac{x}{y(x)}} - \log\left(\frac{y(x)}{x}\right) = \log(x) + c_1, y(x)\right]$$

**1.10 problem First order with homogeneous Coefficients.  
Exercise 7.11, page 61**

Internal problem ID [4436]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 7

**Problem number:** First order with homogeneous Coefficients. Exercise 7.11, page 61.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _dAlembert]`

$$x e^{\frac{y}{x}} - y \sin\left(\frac{y}{x}\right) + x \sin\left(\frac{y}{x}\right) y' = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 63

```
dsolve((x*exp(y(x)/x)-y(x)*sin(y(x)/x))+x*sin(y(x)/x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \text{RootOf}\left(e^{2-Z}\left(4 \ln(x)^2 e^{2-Z} + 8 \ln(x) e^{2-Z} c_1 + 4 c_1^2 e^{2-Z} - 4 \ln(x) \sin(\_Z) e^{-Z} - 4 \sin(\_Z) e^{-Z} c_1 + 2 \sin(\_Z)^2 - 1\right) x\right)$$

✓ Solution by Mathematica

Time used: 0.328 (sec). Leaf size: 39

```
DSolve[(x*Exp[y[x]/x]-y[x]*Sin[y[x]/x])+x*SIn[y[x]/x]*y'[x]==0,y[x],x,IncludeSingularSolutio
```

$$\text{Solve}\left[-\frac{1}{2} e^{-\frac{y(x)}{x}} \left(\sin\left(\frac{y(x)}{x}\right) + \cos\left(\frac{y(x)}{x}\right)\right) = -\log(x) + c_1, y(x)\right]$$

## 1.11 problem First order with homogeneous Coefficients. Exercise 7.12, page 61

Internal problem ID [4437]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 7

**Problem number:** First order with homogeneous Coefficients. Exercise 7.12, page 61.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$y^2 - 2yy'x = -x^2$$

With initial conditions

$$[y(-1) = 0]$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 23

```
dsolve([(x^2+y(x)^2)=2*x*y(x)*diff(y(x),x),y(-1) = 0],y(x), singsol=all)
```

$$y(x) = \sqrt{x(x+1)}$$

$$y(x) = -\sqrt{x(x+1)}$$

### ✓ Solution by Mathematica

Time used: 0.19 (sec). Leaf size: 36

```
DSolve[{(x^2+y[x]^2)==2*x*y[x]*y'[x],y[-1]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{x}\sqrt{x+1}$$

$$y(x) \rightarrow \sqrt{x}\sqrt{x+1}$$

## 1.12 problem First order with homogeneous Coefficients. Exercise 7.13, page 61

Internal problem ID [4438]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 7

**Problem number:** First order with homogeneous Coefficients. Exercise 7.13, page 61.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _dAlembert]`

$$x e^{\frac{y}{x}} + y - xy' = 0$$

With initial conditions

$$[y(1) = 0]$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 15

```
dsolve([(x*exp(y(x)/x)+y(x))=x*diff(y(x),x),y(1) = 0],y(x), singsol=all)
```

$$y(x) = \ln\left(-\frac{1}{\ln(x) - 1}\right)x$$

### ✓ Solution by Mathematica

Time used: 0.316 (sec). Leaf size: 15

```
DSolve[{(x*Exp[y[x]/x]+y[x])==x*y'[x],y[1]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x \log(1 - \log(x))$$



### 1.13 problem First order with homogeneous Coefficients. Exercise 7.14, page 61

Internal problem ID [4439]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 7

**Problem number:** First order with homogeneous Coefficients. Exercise 7.14, page 61.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _dAlembert]`

$$y' - \frac{y}{x} + \csc\left(\frac{y}{x}\right) = 0$$

With initial conditions

$$[y(1) = 0]$$

#### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 16

```
dsolve([diff(y(x),x)-y(x)/x+csc(y(x)/x)=0,y(1) = 0],y(x), singsol=all)
```

$$y(x) = x(1 - 2 \arccos(\ln(x) + 1))$$

#### ✓ Solution by Mathematica

Time used: 0.394 (sec). Leaf size: 24

```
DSolve[{y'[x]-y[x]/x+Csc[y[x]/x]==0,y[1]==0},y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow -x \arccos(\log(x) + 1)$$

$$y(x) \rightarrow x \arccos(\log(x) + 1)$$

## 1.14 problem First order with homogeneous Coefficients. Exercise 7.15, page 61

Internal problem ID [4440]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 7

**Problem number:** First order with homogeneous Coefficients. Exercise 7.15, page 61.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$yx - y^2 - x^2y' = 0$$

With initial conditions

$$[y(1) = 1]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 12

```
dsolve([(x*y(x)-y(x)^2)-x^2*diff(y(x),x)=0,y(1) = 1],y(x), singsol=all)
```

$$y(x) = \frac{x}{\ln(x) + 1}$$

### ✓ Solution by Mathematica

Time used: 0.139 (sec). Leaf size: 13

```
DSolve[{(x*y[x]-y[x]^2)-x^2*y'[x]==0,y[1]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x}{\log(x) + 1}$$

## **2 Chapter 2. Special types of differential equations of the first kind. Lesson 8**

- 2.1 problem Differential equations with Linear Coefficients. Exercise 8.1, page 69 19
- 2.2 problem Differential equations with Linear Coefficients. Exercise 8.2, page 69 20
- 2.3 problem Differential equations with Linear Coefficients. Exercise 8.3, page 69 21
- 2.4 problem Differential equations with Linear Coefficients. Exercise 8.4, page 69 22
- 2.5 problem Differential equations with Linear Coefficients. Exercise 8.5, page 69 23
- 2.6 problem Differential equations with Linear Coefficients. Exercise 8.6, page 69 24
- 2.7 problem Differential equations with Linear Coefficients. Exercise 8.7, page 69 25
- 2.8 problem Differential equations with Linear Coefficients. Exercise 8.8, page 69 26
- 2.9 problem Differential equations with Linear Coefficients. Exercise 8.9, page 69 27
- 2.10 problem Differential equations with Linear Coefficients. Exercise 8.10, page 69 28
- 2.11 problem Differential equations with Linear Coefficients. Exercise 8.11, page 69 29
- 2.12 problem Differential equations with Linear Coefficients. Exercise 8.12, page 69 30
- 2.13 problem Differential equations with Linear Coefficients. Exercise 8.13, page 69 31
- 2.14 problem Differential equations with Linear Coefficients. Exercise 8.14, page 69 33

## 2.1 problem Differential equations with Linear Coefficients. Exercise 8.1, page 69

Internal problem ID [4441]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 8

**Problem number:** Differential equations with Linear Coefficients. Exercise 8.1, page 69.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$2y - (2x - 4y)y' = -x + 4$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 31

```
dsolve((x+2*y(x)-4)-(2*x-4*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 1 - \frac{\tan\left(\text{RootOf}\left(2\_Z + \ln\left(\frac{1}{\cos(\_Z)^2}\right) + 2\ln(x-2) + 2c_1\right)\right)(x-2)}{2}$$

### ✓ Solution by Mathematica

Time used: 0.057 (sec). Leaf size: 63

```
DSolve[(x+2*y[x]-4)-(2*x-4*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[2\arctan\left(\frac{-2y(x)-x+4}{x-2y(x)}\right) + \log\left(\frac{x^2+4y(x)^2-8y(x)-4x+8}{2(x-2)^2}\right) + 2\log(x-2) + c_1 = 0, y(x)\right]$$

## 2.2 problem Differential equations with Linear Coefficients. Exercise 8.2, page 69

Internal problem ID [4442]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 8

**Problem number:** Differential equations with Linear Coefficients. Exercise 8.2, page 69.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$2y - (3x + 2y - 1)y' = -1 - 3x$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 21

```
dsolve((3*x+2*y(x)+1)-(3*x+2*y(x)-1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{3x}{2} - \frac{2 \operatorname{LambertW}\left(-\frac{e^{\frac{1}{4}} e^{-\frac{25x}{4}} c_1}{4}\right)}{5} + \frac{1}{10}$$

### ✓ Solution by Mathematica

Time used: 4.816 (sec). Leaf size: 43

```
DSolve[(3*x+2*y[x]+1)-(3*x+2*y[x]-1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{10} \left( -4W\left(-e^{-\frac{25x}{4}-1+c_1}\right) - 15x + 1 \right)$$

$$y(x) \rightarrow \frac{1}{10} - \frac{3x}{2}$$

## 2.3 problem Differential equations with Linear Coefficients. Exercise 8.3, page 69

Internal problem ID [4443]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 8

**Problem number:** Differential equations with Linear Coefficients. Exercise 8.3, page 69.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y + (2x + 2y + 2)y' = -x - 1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve((x+y(x)+1)+(2*x+2*y(x)+2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -x - 1$$

$$y(x) = -\frac{x}{2} + c_1$$

### ✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 22

```
DSolve[(x+y[x]+1)+(2*x+2*y[x]+2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x - 1$$

$$y(x) \rightarrow -\frac{x}{2} + c_1$$

## 2.4 problem Differential equations with Linear Coefficients. Exercise 8.4, page 69

Internal problem ID [4444]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 8

**Problem number:** Differential equations with Linear Coefficients. Exercise 8.4, page 69.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$y + (2x + 2y - 3)y' = 1 - x$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 29

```
dsolve((x+y(x)-1)+(2*x+2*y(x)-3)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = e^{-\text{LambertW}(2e^xe^{-4}e^{-c_1})+x-4-c_1} + 2 - x$$

### ✓ Solution by Mathematica

Time used: 4.725 (sec). Leaf size: 33

```
DSolve[(x+y[x]-1)+(2*x+2*y[x]-3)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(W(-e^{x-1+c_1}) - 2x + 4)$$

$$y(x) \rightarrow 2 - x$$

## 2.5 problem Differential equations with Linear Coefficients. Exercise 8.5, page 69

Internal problem ID [4445]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 8

**Problem number:** Differential equations with Linear Coefficients. Exercise 8.5, page 69.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$y - (x - y - 1)y' = 1 - x$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 29

```
dsolve((x+y(x)-1)-(x-y(x)-1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\tan\left(\text{RootOf}\left(2_Z + \ln\left(\frac{1}{\cos(_Z)^2}\right) + 2\ln(x-1) + 2c_1\right)\right)(x-1)$$

### ✓ Solution by Mathematica

Time used: 0.057 (sec). Leaf size: 48

```
DSolve[(x+y[x]-1)-(x-y[x]-1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[2 \arctan\left(\frac{y(x) + x - 1}{-y(x) + x - 1}\right) = \log\left(\frac{1}{2}\left(\frac{y(x)^2}{(x-1)^2} + 1\right)\right) + 2\log(x-1) + c_1, y(x)\right]$$



## 2.6 problem Differential equations with Linear Coefficients. Exercise 8.6, page 69

Internal problem ID [4446]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 8

**Problem number:** Differential equations with Linear Coefficients. Exercise 8.6, page 69.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$y + (2x + 2y - 1)y' = -x$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve((x+y(x))+(2*x+2*y(x)-1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = e^{-\text{LambertW}(2e^x e^{-2} e^{-c_1}) + x - 2 - c_1} - x + 1$$

### ✓ Solution by Mathematica

Time used: 1.056 (sec). Leaf size: 33

```
DSolve[(x+y[x])+(2*x+2*y[x]-1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(W(-e^{x-1+c_1}) - 2x + 2)$$

$$y(x) \rightarrow 1 - x$$

## 2.7 problem Differential equations with Linear Coefficients. Exercise 8.7, page 69

Internal problem ID [4447]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 8

**Problem number:** Differential equations with Linear Coefficients. Exercise 8.7, page 69.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$7y + (1 + 2x)y' = 3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve((7*y(x)-3)+(2*x+1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{3}{7} + \frac{c_1}{(1 + 2x)^{\frac{7}{2}}}$$

### ✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 28

```
DSolve[(7*y[x]-3)+(2*x+1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3}{7} + \frac{c_1}{(2x + 1)^{7/2}}$$

$$y(x) \rightarrow \frac{3}{7}$$

## 2.8 problem Differential equations with Linear Coefficients. Exercise 8.8, page 69

Internal problem ID [4448]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 8

**Problem number:** Differential equations with Linear Coefficients. Exercise 8.8, page 69.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$2y + (3x + 6y + 3)y' = -x$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```
dsolve((x+2*y(x))+(3*x+6*y(x)+3)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{e^{-\text{LambertW}\left(-\frac{e^{-\frac{x}{6}} e^{-\frac{3}{2}} e^{\frac{c_1}{6}}}{2}\right) - \frac{x}{6} - \frac{3}{2} + \frac{c_1}{6}}}{2} - \frac{3}{2} - \frac{x}{2}$$

### ✓ Solution by Mathematica

Time used: 4.834 (sec). Leaf size: 43

```
DSolve[(x+2*y[x])+(3*x+6*y[x]+3)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(-2W(-e^{-\frac{x}{6}-1+c_1}) - x - 3)$$

$$y(x) \rightarrow \frac{1}{2}(-x - 3)$$

## 2.9 problem Differential equations with Linear Coefficients. Exercise 8.9, page 69

Internal problem ID [4449]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 8

**Problem number:** Differential equations with Linear Coefficients. Exercise 8.9, page 69.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$2y + (y - 1)y' = -x$$

### ✓ Solution by Maple

Time used: 0.234 (sec). Leaf size: 27

```
dsolve((x+2*y(x))+(y(x)-1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 1 - \frac{(2+x)(\text{LambertW}(c_1(2+x)) + 1)}{\text{LambertW}(c_1(2+x))}$$

### ✓ Solution by Mathematica

Time used: 1.178 (sec). Leaf size: 143

```
DSolve[(x+2*y[x])+(y[x]-1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ \frac{(-2)^{2/3} \left( - \left( (x+1) \log \left( - \frac{3(-2)^{2/3}(x+2)}{y(x)-1} \right) \right) + x \log \left( \frac{3(-2)^{2/3}(y(x)+x+1)}{y(x)-1} \right) + \log \left( \frac{3(-2)^{2/3}(y(x)+x+1)}{y(x)-1} \right) \right)}{9(y(x)+x+1)} \right]$$

## 2.10 problem Differential equations with Linear Coefficients. Exercise 8.10, page 69

Internal problem ID [4450]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 8

**Problem number:** Differential equations with Linear Coefficients. Exercise 8.10, page 69.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _exact, _rational, [_Abel, '2nd ty`

$$-2y - (2x + 7y - 1)y' = -4 - 3x$$

### ✓ Solution by Maple

Time used: 0.531 (sec). Leaf size: 38

```
dsolve((3*x-2*y(x)+4)-(2*x+7*y(x)-1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{11}{25} - \frac{\frac{2(25x+26)c_1}{7} + \frac{\sqrt{25(25x+26)^2c_1^2+7}}{7}}{25c_1}$$

### ✓ Solution by Mathematica

Time used: 0.126 (sec). Leaf size: 65

```
DSolve[(3*x-2*y[x]+4)-(2*x+7*y[x]-1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{7} \left( -\sqrt{25x^2 + 52x + 1 + 49c_1} - 2x + 1 \right)$$

$$y(x) \rightarrow \frac{1}{7} \left( \sqrt{25x^2 + 52x + 1 + 49c_1} - 2x + 1 \right)$$

## 2.11 problem Differential equations with Linear Coefficients. Exercise 8.11, page 69

Internal problem ID [4451]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 8

**Problem number:** Differential equations with Linear Coefficients. Exercise 8.11, page 69.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$y + (3x + 3y - 4)y' = -x$$

With initial conditions

$$[y(1) = 0]$$

✓ Solution by Maple

Time used: 0.172 (sec). Leaf size: 19

```
dsolve([(x+y(x))+(3*x+3*y(x)-4)*diff(y(x),x)=0,y(1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{2 \operatorname{LambertW}\left(-1, -\frac{3e^{x-\frac{5}{2}}}{2}\right)}{3} + 2 - x$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{(x+y[x])+(3*x+3*y[x]-4)*y'[x]==0,y[1]==0},y[x],x,IncludeSingularSolutions -> True]
```

{}

## 2.12 problem Differential equations with Linear Coefficients. Exercise 8.12, page 69

Internal problem ID [4452]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 8

**Problem number:** Differential equations with Linear Coefficients. Exercise 8.12, page 69.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$2y - (-1 + x + 2y)y' = -3x - 3$$

✓ Solution by Maple

Time used: 0.422 (sec). Leaf size: 46

```
dsolve((3*x+2*y(x)+3)-(x+2*y(x)-1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{9}{2} - \frac{\text{RootOf}((2x+4)^5 c_1 Z^{25} - 5(2x+4)^5 c_1 Z^{20} - 2)^5 (2x+4)}{4} + \frac{3x}{2}$$

✓ Solution by Mathematica

Time used: 60.094 (sec). Leaf size: 3081

```
DSolve[(3*x+2*y[x]+3)-(x+2*y[x]-1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

## 2.13 problem Differential equations with Linear Coefficients. Exercise 8.13, page 69

Internal problem ID [4453]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 8

**Problem number:** Differential equations with Linear Coefficients. Exercise 8.13, page 69.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _rational, [_Abel, '2nd type', 'cl`

$$y + (2x + y + 3)y' = -7$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.156 (sec). Leaf size: 87

```
dsolve([(y(x)+7)+(2*x+y(x)+3)*diff(y(x),x)=0,y(0) = 1],y(x), singsol=all)
```

$$y(x) = \left(-x^3 + 6x^2 - 12x + 72 + 8\sqrt{-2x^3 + 12x^2 - 24x + 80}\right)^{\frac{1}{3}} + \frac{(x-2)^2}{\left(-x^3 + 6x^2 - 12x + 72 + 8\sqrt{-2x^3 + 12x^2 - 24x + 80}\right)^{\frac{1}{3}}} - x - 5$$



✓ Solution by Mathematica

Time used: 6.783 (sec). Leaf size: 198

```
DSolve[{(y[x]+7)+(2*x+y[x]+3)*y'[x]==0,y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) = \frac{x^2 - \left( \sqrt[3]{-x^3 + 6x^2 + 8\sqrt{2}\sqrt{-x^3 + 6x^2 - 12x + 40}} - 12x + 72 + 4 \right) x + (-x^3 + 6x^2 + 8\sqrt{2}\sqrt{-x^3 + 6x^2 - 12x + 40})}{\sqrt[3]{-x^3 + 6x^2 + 8\sqrt{2}\sqrt{-x^3 + 6x^2 - 12x + 40}}}$$

## 2.14 problem Differential equations with Linear Coefficients. Exercise 8.14, page 69

Internal problem ID [4454]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 8

**Problem number:** Differential equations with Linear Coefficients. Exercise 8.14, page 69.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$y - (x - y - 4)y' = -2 - x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve((x+y(x)+2)-(x-y(x)-4)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -3 - \tan \left( \text{RootOf} \left( 2\_Z + \ln \left( \frac{1}{\cos(\_Z)^2} \right) + 2 \ln(x - 1) + 2c_1 \right) \right) (x - 1)$$

### ✓ Solution by Mathematica

Time used: 0.056 (sec). Leaf size: 58

```
DSolve[(x+y[x]+2)-(x-y[x]-4)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ 2 \arctan \left( \frac{y(x) + x + 2}{y(x) - x + 4} \right) + \log \left( \frac{x^2 + y(x)^2 + 6y(x) - 2x + 10}{2(x - 1)^2} \right) + 2 \log(x - 1) + c_1 = 0, y(x) \right]$$

### **3 Chapter 2. Special types of differential equations of the first kind. Lesson 9**

3.1	problem Exact Differential equations. Exercise 9.4, page 79 . . . . .	35
3.2	problem Exact Differential equations. Exercise 9.5, page 79 . . . . .	38
3.3	problem Exact Differential equations. Exercise 9.6, page 79 . . . . .	39
3.4	problem Exact Differential equations. Exercise 9.7, page 79 . . . . .	41
3.5	problem Exact Differential equations. Exercise 9.8, page 79 . . . . .	42
3.6	problem Exact Differential equations. Exercise 9.9, page 79 . . . . .	43
3.7	problem Exact Differential equations. Exercise 9.10, page 79 . . . . .	44
3.8	problem Exact Differential equations. Exercise 9.11, page 79 . . . . .	45
3.9	problem Exact Differential equations. Exercise 9.12, page 79 . . . . .	46
3.10	problem Exact Differential equations. Exercise 9.13, page 79 . . . . .	48
3.11	problem Exact Differential equations. Exercise 9.15, page 79 . . . . .	51
3.12	problem Exact Differential equations. Exercise 9.16, page 79 . . . . .	52
3.13	problem Exact Differential equations. Exercise 9.17, page 79 . . . . .	53

### 3.1 problem Exact Differential equations. Exercise 9.4, page 79

Internal problem ID [4455]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 9

**Problem number:** Exact Differential equations. Exercise 9.4, page 79.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_exact`, `_rational`]

$$3x^2y + 8y^2x + (x^3 + 8x^2y + 12y^2) y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 597

`dsolve((3*x^2*y(x)+8*x*y(x)^2)+(x^3+8*x^2*y(x)+12*y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)`

$$\begin{aligned}
 y(x) &= \frac{\left(9x^5 - 27c_1 - 8x^6 + 3\sqrt{-3x^{10} + 3x^9 + 48c_1x^6 - 54c_1x^5 + 81c_1^2}\right)^{\frac{1}{3}}}{6\left(\frac{1}{12}x^3 - \frac{1}{9}x^4\right)} - \frac{x^2}{3} \\
 &\quad - \frac{\left(9x^5 - 27c_1 - 8x^6 + 3\sqrt{-3x^{10} + 3x^9 + 48c_1x^6 - 54c_1x^5 + 81c_1^2}\right)^{\frac{1}{3}}}{\left(9x^5 - 27c_1 - 8x^6 + 3\sqrt{-3x^{10} + 3x^9 + 48c_1x^6 - 54c_1x^5 + 81c_1^2}\right)^{\frac{1}{3}}} \\
 y(x) &= -\frac{\left(9x^5 - 27c_1 - 8x^6 + 3\sqrt{-3x^{10} + 3x^9 + 48c_1x^6 - 54c_1x^5 + 81c_1^2}\right)^{\frac{1}{3}}}{\frac{12}{\frac{1}{4}x^3 - \frac{1}{3}x^4}} - \frac{x^2}{3} \\
 &\quad + \frac{\left(9x^5 - 27c_1 - 8x^6 + 3\sqrt{-3x^{10} + 3x^9 + 48c_1x^6 - 54c_1x^5 + 81c_1^2}\right)^{\frac{1}{3}}}{\left(9x^5 - 27c_1 - 8x^6 + 3\sqrt{-3x^{10} + 3x^9 + 48c_1x^6 - 54c_1x^5 + 81c_1^2}\right)^{\frac{1}{3}}} \\
 &\quad - \frac{i\sqrt{3}\left(\frac{\left(9x^5 - 27c_1 - 8x^6 + 3\sqrt{-3x^{10} + 3x^9 + 48c_1x^6 - 54c_1x^5 + 81c_1^2}\right)^{\frac{1}{3}}}{6} + \frac{\frac{1}{2}x^3 - \frac{2}{3}x^4}{\left(9x^5 - 27c_1 - 8x^6 + 3\sqrt{-3x^{10} + 3x^9 + 48c_1x^6 - 54c_1x^5 + 81c_1^2}\right)^{\frac{1}{3}}}\right)}{2} \\
 y(x) &= -\frac{\left(9x^5 - 27c_1 - 8x^6 + 3\sqrt{-3x^{10} + 3x^9 + 48c_1x^6 - 54c_1x^5 + 81c_1^2}\right)^{\frac{1}{3}}}{\frac{12}{\frac{1}{4}x^3 - \frac{1}{3}x^4}} - \frac{x^2}{3} \\
 &\quad + \frac{\left(9x^5 - 27c_1 - 8x^6 + 3\sqrt{-3x^{10} + 3x^9 + 48c_1x^6 - 54c_1x^5 + 81c_1^2}\right)^{\frac{1}{3}}}{\left(9x^5 - 27c_1 - 8x^6 + 3\sqrt{-3x^{10} + 3x^9 + 48c_1x^6 - 54c_1x^5 + 81c_1^2}\right)^{\frac{1}{3}}} \\
 &\quad + \frac{i\sqrt{3}\left(\frac{\left(9x^5 - 27c_1 - 8x^6 + 3\sqrt{-3x^{10} + 3x^9 + 48c_1x^6 - 54c_1x^5 + 81c_1^2}\right)^{\frac{1}{3}}}{6} + \frac{\frac{1}{2}x^3 - \frac{2}{3}x^4}{\left(9x^5 - 27c_1 - 8x^6 + 3\sqrt{-3x^{10} + 3x^9 + 48c_1x^6 - 54c_1x^5 + 81c_1^2}\right)^{\frac{1}{3}}}\right)}{2}
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 1.703 (sec). Leaf size: 474

`DSolve[(3*x^2*y[x]+8*x*y[x]^2)+(x^3+8*x^2*y[x]+12*y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSol`

$$y(x) \rightarrow \frac{1}{6} \left( -2x^2 + \sqrt[3]{-8x^6 + 9x^5 + 3\sqrt{3}\sqrt{-x^{10} + x^9 - 16c_1x^6 + 18c_1x^5 + 27c_1^2 + 27c_1}} \right. \\ \left. + \frac{(4x - 3)x^3}{\sqrt[3]{-8x^6 + 9x^5 + 3\sqrt{3}\sqrt{-x^{10} + x^9 - 16c_1x^6 + 18c_1x^5 + 27c_1^2 + 27c_1}}} \right)$$

$$y(x) \rightarrow \frac{1}{48} \left( -16x^2 + 4i(\sqrt{3} \right. \\ \left. + i) \sqrt[3]{-8x^6 + 9x^5 + 3\sqrt{3}\sqrt{-x^{10} + x^9 - 16c_1x^6 + 18c_1x^5 + 27c_1^2 + 27c_1}} \right. \\ \left. - \frac{4i(\sqrt{3} - i)(4x - 3)x^3}{\sqrt[3]{-8x^6 + 9x^5 + 3\sqrt{3}\sqrt{-x^{10} + x^9 - 16c_1x^6 + 18c_1x^5 + 27c_1^2 + 27c_1}}} \right)$$

$$y(x) \rightarrow \frac{1}{48} \left( -16x^2 - 4(1 \right. \\ \left. + i\sqrt{3}) \sqrt[3]{-8x^6 + 9x^5 + 3\sqrt{3}\sqrt{-x^{10} + x^9 - 16c_1x^6 + 18c_1x^5 + 27c_1^2 + 27c_1}} \right. \\ \left. + \frac{4i(\sqrt{3} + i)(4x - 3)x^3}{\sqrt[3]{-8x^6 + 9x^5 + 3\sqrt{3}\sqrt{-x^{10} + x^9 - 16c_1x^6 + 18c_1x^5 + 27c_1^2 + 27c_1}}} \right)$$

### 3.2 problem Exact Differential equations. Exercise 9.5, page 79

Internal problem ID [4456]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 9

**Problem number:** Exact Differential equations. Exercise 9.5, page 79.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class D', _exact, _rational, [_Abel, '2nd ty`

$$\frac{1 + 2yx}{y} + \frac{(y - x)y'}{y^2} = 0$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve((2*x*y(x)+1)/y(x)+(y(x)-x)/y(x)^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{x}{\text{LambertW}(-e^{x^2}c_1x)}$$

#### ✓ Solution by Mathematica

Time used: 5.208 (sec). Leaf size: 29

```
DSolve[(2*x*y[x]+1)/y[x]+(y[x]-x)/y[x]^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x}{W(x(-e^{x^2-c_1}))}$$

$$y(x) \rightarrow 0$$

### 3.3 problem Exact Differential equations. Exercise 9.6, page 79

Internal problem ID [4457]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 9

**Problem number:** Exact Differential equations. Exercise 9.6, page 79.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _exact, _rational, _dAlembert]`

$$2yx + (x^2 + y^2) y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 257

```
dsolve(2*x*y(x)+(x^2+y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\frac{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}{2} - \frac{2x^2c_1}{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}}{\sqrt{c_1}}$$

$$y(x) = \frac{-\frac{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}{4} + \frac{x^2c_1}{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}} - \frac{i\sqrt{3}\left(\frac{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}{2} + \frac{2x^2c_1}{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}\right)}{2}}{\sqrt{c_1}}$$

$$y(x) = \frac{-\frac{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}{4} + \frac{x^2c_1}{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}} + \frac{i\sqrt{3}\left(\frac{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}{2} + \frac{2x^2c_1}{(4+4\sqrt{4x^6c_1^3+1})^{\frac{1}{3}}}\right)}{2}}{\sqrt{c_1}}$$



✓ Solution by Mathematica

Time used: 15.514 (sec). Leaf size: 401

`DSolve[2*x*y[x]+(x^2+y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}{\sqrt[3]{2}} - \frac{\sqrt[3]{2}x^2}{\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}$$

$$y(x) \rightarrow \frac{i2^{2/3}(\sqrt{3} + i)(\sqrt{4x^6 + e^{6c_1}} + e^{3c_1})^{2/3} + \sqrt[3]{2}(2 + 2i\sqrt{3})x^2}{4\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}$$

$$y(x) \rightarrow \frac{(1 - i\sqrt{3})x^2}{2^{2/3}\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}} - \frac{(1 + i\sqrt{3})\sqrt[3]{\sqrt{4x^6 + e^{6c_1}} + e^{3c_1}}}{2\sqrt[3]{2}}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow \frac{1}{2}\sqrt[6]{x^6} \left( \frac{(1 - i\sqrt{3})(x^6)^{2/3}}{x^4} - i\sqrt{3} - 1 \right)$$

$$y(x) \rightarrow \frac{1}{2}\sqrt[6]{x^6} \left( \frac{(1 + i\sqrt{3})(x^6)^{2/3}}{x^4} + i\sqrt{3} - 1 \right)$$

$$y(x) \rightarrow \sqrt[6]{x^6} - \frac{(x^6)^{5/6}}{x^4}$$

### 3.4 problem Exact Differential equations. Exercise 9.7, page 79

Internal problem ID [4458]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 9

**Problem number:** Exact Differential equations. Exercise 9.7, page 79.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [exact]

$$e^x \sin(y) + e^{-y} - (x e^{-y} - e^x \cos(y)) y' = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve((exp(x)*sin(y(x))+exp(-y(x)))-(x*exp(-y(x))-exp(x)*cos(y(x)))*diff(y(x),x)=0,y(x), si
```

$$e^x \sin(y(x)) + x e^{-y(x)} + c_1 = 0$$

#### ✓ Solution by Mathematica

Time used: 0.389 (sec). Leaf size: 24

```
DSolve[(Exp[x]*Sin[y[x]]+Exp[-y[x]])-(x*Exp[-y[x]]-Exp[x]*Cos[y[x]])*y'[x]==0,y[x],x,Include
```

$$\text{Solve}[x(-e^{-y(x)}) - e^x \sin(y(x)) = c_1, y(x)]$$

### 3.5 problem Exact Differential equations. Exercise 9.8, page 79

Internal problem ID [4459]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 9

**Problem number:** Exact Differential equations. Exercise 9.8, page 79.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_exact, [\_1st\_order, ‘\_with\_symmetry\_[F(x)\*G(y),0]’]]

$$\cos(y) - (x \sin(y) - y^2) y' = 0$$

#### ✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 20

```
dsolve(cos(y(x))-(x*sin(y(x))-y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$x - \frac{-\frac{y(x)^3}{3} + c_1}{\cos(y(x))} = 0$$

#### ✓ Solution by Mathematica

Time used: 0.124 (sec). Leaf size: 23

```
DSolve[Cos[y[x]]-(x*Sin[y[x]]-y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ x = -\frac{1}{3}y(x)^3 \sec(y(x)) + c_1 \sec(y(x)), y(x) \right]$$

### 3.6 problem Exact Differential equations. Exercise 9.9, page 79

Internal problem ID [4460]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 9

**Problem number:** Exact Differential equations. Exercise 9.9, page 79.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [exact]

$$-2yx + e^y + (y - x^2 + x e^y) y' = -x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve((x-2*x*y(x)+exp(y(x)))+(y(x)-x^2+x*exp(y(x)))*diff(y(x),x)=0,y(x), singsol=all)
```

$$-y(x)x^2 + xe^{y(x)} + \frac{x^2}{2} + \frac{y(x)^2}{2} + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.315 (sec). Leaf size: 35

```
DSolve[(x-2*x*y[x]+Exp[y[x]])+(y[x]-x^2+x*Exp[y[x]])*y'[x]==0,y[x],x,IncludeSingularSolution
```

$$\text{Solve}\left[x^2(-y(x)) + \frac{x^2}{2} + xe^{y(x)} + \frac{y(x)^2}{2} = c_1, y(x)\right]$$

### 3.7 problem Exact Differential equations. Exercise 9.10, page 79

Internal problem ID [4461]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 9

**Problem number:** Exact Differential equations. Exercise 9.10, page 79.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [exact]

$$y^2 - (-2yx + e^y)y' = -x^2 + x$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve((x^2-x+y(x)^2)-(exp(y(x))-2*x*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$\frac{x^3}{3} + y(x)^2 x - \frac{x^2}{2} - e^{y(x)} + c_1 = 0$$

#### ✓ Solution by Mathematica

Time used: 0.198 (sec). Leaf size: 32

```
DSolve[(x^2-x+y[x]^2)-(Exp[y[x]]-2*x*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[-\frac{x^3}{3} + \frac{x^2}{2} - xy(x)^2 + e^{y(x)} = c_1, y(x)\right]$$

### 3.8 problem Exact Differential equations. Exercise 9.11, page 79

Internal problem ID [4462]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 9

**Problem number:** Exact Differential equations. Exercise 9.11, page 79.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [exact]

$$y \cos(x) + (2y + \sin(x) - \sin(y)) y' = -2x$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve((2*x+y(x)*cos(x))+(2*y(x)+sin(x)-sin(y(x)))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) \sin(x) + x^2 + y(x)^2 + \cos(y(x)) + c_1 = 0$$

#### ✓ Solution by Mathematica

Time used: 0.198 (sec). Leaf size: 22

```
DSolve[(2*x+y[x]*Cos[x])+(2*y[x]+Sin[x]-Sin[y[x]])*y'[x]==0,y[x],x,IncludeSingularSolutions
```

$$\text{Solve}[x^2 + y(x)^2 + y(x) \sin(x) + \cos(y(x)) = c_1, y(x)]$$

### 3.9 problem Exact Differential equations. Exercise 9.12, page 79

Internal problem ID [4463]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 9

**Problem number:** Exact Differential equations. Exercise 9.12, page 79.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _exact, _dAlembert]`

$$x\sqrt{x^2 + y^2} - \frac{x^2 y y'}{y - \sqrt{x^2 + y^2}} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(x*sqrt(x^2+y(x)^2)-(x^2*y(x))/(y(x)- sqrt(x^2+y(x)^2))*diff(y(x),x)=0,y(x), singsol=a
```

$$c_1 + (x^2 + y(x)^2)^{\frac{3}{2}} + y(x)^3 = 0$$

✓ Solution by Mathematica

Time used: 60.259 (sec). Leaf size: 2125

`DSolve[x*Sqrt[x^2+y[x]^2]-(x^2*y[x])/(y[x]-Sqrt[x^2+y[x]^2])*y'[x]==0,y[x],x,IncludeSingularities->True]`

$y(x) \rightarrow$

$$x^2 \sqrt{\frac{e^{6c_1}}{x^4} - 6x^2 + \frac{3(5x^6 - 4e^{6c_1})}{\sqrt[3]{-11x^{12} + 14e^{6c_1}x^6 + 2\sqrt{(-x^6 + e^{6c_1})(x^6 + e^{6c_1})^3} - 2e^{12c_1}}} + \frac{3\sqrt[3]{-11x^{12} + 14e^{6c_1}x^6 + 2\sqrt{(-x^6 + e^{6c_1})(x^6 + e^{6c_1})^3} - 2e^{12c_1}}}{\sqrt[3]{-11x^{12} + 14e^{6c_1}x^6 + 2\sqrt{(-x^6 + e^{6c_1})(x^6 + e^{6c_1})^3} - 2e^{12c_1}}}}$$


---

$y(x)$

$$x^2 \left( - \sqrt{\frac{e^{6c_1}}{x^4} - 6x^2 + \frac{3(5x^6 - 4e^{6c_1})}{\sqrt[3]{-11x^{12} + 14e^{6c_1}x^6 + 2\sqrt{(-x^6 + e^{6c_1})(x^6 + e^{6c_1})^3} - 2e^{12c_1}}} + \frac{3\sqrt[3]{-11x^{12} + 14e^{6c_1}x^6 + 2\sqrt{(-x^6 + e^{6c_1})(x^6 + e^{6c_1})^3} - 2e^{12c_1}}}{\sqrt[3]{-11x^{12} + 14e^{6c_1}x^6 + 2\sqrt{(-x^6 + e^{6c_1})(x^6 + e^{6c_1})^3} - 2e^{12c_1}}}} \right)$$


---

$\rightarrow$

$y(x)$

$$x^2 \sqrt{\frac{e^{6c_1}}{x^4} - 6x^2 + \frac{3(5x^6 - 4e^{6c_1})}{\sqrt[3]{-11x^{12} + 14e^{6c_1}x^6 + 2\sqrt{(-x^6 + e^{6c_1})(x^6 + e^{6c_1})^3} - 2e^{12c_1}}} + \frac{3\sqrt[3]{-11x^{12} + 14e^{6c_1}x^6 + 2\sqrt{(-x^6 + e^{6c_1})(x^6 + e^{6c_1})^3} - 2e^{12c_1}}}{\sqrt[3]{-11x^{12} + 14e^{6c_1}x^6 + 2\sqrt{(-x^6 + e^{6c_1})(x^6 + e^{6c_1})^3} - 2e^{12c_1}}}}$$


---

$\rightarrow$

$y(x)$

$$x^2 \sqrt{\frac{e^{6c_1}}{x^4} - 6x^2 + \frac{3(5x^6 - 4e^{6c_1})}{\sqrt[3]{-11x^{12} + 14e^{6c_1}x^6 + 2\sqrt{(-x^6 + e^{6c_1})(x^6 + e^{6c_1})^3} - 2e^{12c_1}}} + \frac{3\sqrt[3]{-11x^{12} + 14e^{6c_1}x^6 + 2\sqrt{(-x^6 + e^{6c_1})(x^6 + e^{6c_1})^3} - 2e^{12c_1}}}{\sqrt[3]{-11x^{12} + 14e^{6c_1}x^6 + 2\sqrt{(-x^6 + e^{6c_1})(x^6 + e^{6c_1})^3} - 2e^{12c_1}}}}$$


---

$\rightarrow$



### 3.10 problem Exact Differential equations. Exercise 9.13, page 79

Internal problem ID [4464]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 9

**Problem number:** Exact Differential equations. Exercise 9.13, page 79.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_exact`]

$$y^3 - (y^2 + 1 - 3y^2x) y' = -4x^3 + \sin(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 1162

`dsolve((4*x^3-sin(x)+y(x)^3)-(y(x)^2+1-3*x*y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)`

$$y(x) = \frac{\left( \left( -12x^4 + 4\sqrt{\frac{27x^9 - 9x^8 + 54\cos(x)x^5 + 54c_1x^5 - 18\cos(x)x^4 - 18c_1x^4 + 27x\cos(x)^2 + 54c_1x\cos(x) + 27c_1^2x - 9\cos(x)^2 - 18c_1\cos(x) - 9c_1^2}{3x-1}} \right) \right)}{6x - 2} + \frac{6x - 2}{2}$$

$$+ \frac{\left( \left( -12x^4 + 4\sqrt{\frac{27x^9 - 9x^8 + 54\cos(x)x^5 + 54c_1x^5 - 18\cos(x)x^4 - 18c_1x^4 + 27x\cos(x)^2 + 54c_1x\cos(x) + 27c_1^2x - 9\cos(x)^2 - 18c_1\cos(x) - 9c_1^2}{3x-1}} \right) \right)}{6x - 2}$$

$$y(x) = \frac{\left( \left( -12x^4 + 4\sqrt{\frac{27x^9 - 9x^8 + 54\cos(x)x^5 + 54c_1x^5 - 18\cos(x)x^4 - 18c_1x^4 + 27x\cos(x)^2 + 54c_1x\cos(x) + 27c_1^2x - 9\cos(x)^2 - 18c_1\cos(x) - 9c_1^2}{3x-1}} \right) \right)}{4(3x - 1)} - \frac{4(3x - 1)}{1}$$

$$+ i\sqrt{3} \left( \frac{\left( \left( -12x^4 + 4\sqrt{\frac{27x^9 - 9x^8 + 54\cos(x)x^5 + 54c_1x^5 - 18\cos(x)x^4 - 18c_1x^4 + 27x\cos(x)^2 + 54c_1x\cos(x) + 27c_1^2x - 9\cos(x)^2 - 18c_1\cos(x) - 9c_1^2 - 4}{3x-1}} \right) \right)}{6x - 2} \right)$$

$$y(x) = \frac{\left( \left( -12x^4 + 4\sqrt{\frac{27x^9 - 9x^8 + 54\cos(x)x^5 + 54c_1x^5 - 18\cos(x)x^4 - 18c_1x^4 + 27x\cos(x)^2 + 54c_1x\cos(x) + 27c_1^2x - 9\cos(x)^2 - 18c_1\cos(x) - 9c_1^2}{3x-1}} \right) \right)}{4(3x - 1)} - \frac{4(3x - 1)}{1}$$

$$+ i\sqrt{3} \left( \frac{\left( \left( -12x^4 + 4\sqrt{\frac{27x^9 - 9x^8 + 54\cos(x)x^5 + 54c_1x^5 - 18\cos(x)x^4 - 18c_1x^4 + 27x\cos(x)^2 + 54c_1x\cos(x) + 27c_1^2x - 9\cos(x)^2 - 18c_1\cos(x) - 9c_1^2 - 4}{3x-1}} \right) \right)}{6x - 2} \right)$$

✓ Solution by Mathematica

Time used: 60.207 (sec). Leaf size: 682

`DSolve[(4*x^3-Sin[x]+y[x]^3)-(y[x]^2+1-3*x*y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions`

$$y(x) \rightarrow \frac{\sqrt[3]{2} \left( -27x^6 + 18x^5 - 3x^4 + \frac{1}{27} \sqrt{4(9-27x)^3 + 6561(1-3x)^4(x^4 + \cos(x) - c_1)^2} - 27x^2 \cos(x) + 27c_1 \right)}{2^{2/3}(3x-1) \sqrt[3]{-27x^6 + 18x^5 - 3x^4 + \frac{1}{27} \sqrt{4(9-27x)^3 + 6561(1-3x)^4(x^4 + \cos(x) - c_1)^2} - 27x^2 \cos(x) + 27c_1}}$$

$$y(x) \rightarrow \frac{9i \sqrt[3]{2} (\sqrt{3} + i) \left( -27x^6 + 18x^5 - 3x^4 + \frac{1}{27} \sqrt{4(9-27x)^3 + 6561(1-3x)^4(x^4 + \cos(x) - c_1)^2} - 27x^2 \cos(x) + 27c_1 \right)}{18 \cdot 2^{2/3} (3x-1) \sqrt[3]{-27x^6 + 18x^5 - 3x^4 + \frac{1}{27} \sqrt{4(9-27x)^3 + 6561(1-3x)^4(x^4 + \cos(x) - c_1)^2} - 27x^2 \cos(x) + 27c_1}}$$

$$y(x) \rightarrow \frac{i(\sqrt{3} + i)}{2^{2/3} \sqrt[3]{-27x^6 + 18x^5 - 3x^4 + \frac{1}{27} \sqrt{4(9-27x)^3 + 6561(1-3x)^4(x^4 + \cos(x) - c_1)^2} - 27x^2 \cos(x) + 27c_1} \cdot \frac{(1 + i\sqrt{3}) \sqrt[3]{-54x^6 + 36x^5 - 6x^4 + \frac{2}{27} \sqrt{4(9-27x)^3 + 6561(1-3x)^4(x^4 + \cos(x) - c_1)^2} - 54x^2 \cos(x) + 54c_1}}{2 \cdot 2^{2/3} (3x-1)}$$

### 3.11 problem Exact Differential equations. Exercise 9.15, page 79

Internal problem ID [4465]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 9

**Problem number:** Exact Differential equations. Exercise 9.15, page 79.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_exact`, `_Bernoulli`]

$$e^x (y^3 + y^3 x + 1) + 3y^2 (x e^x - 6) y' = 0$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 38

```
dsolve([exp(x)*(y(x)^3+x*y(x)^3+1)+3*y(x)^2*(x*exp(x)-6)*diff(y(x),x)=0,y(0) = 1],y(x), sing
```

$$y(x) = \frac{(-1 + i\sqrt{3}) (-(e^x + 5) (e^x x - 6)^2)^{\frac{1}{3}}}{2 e^x x - 12}$$

✓ Solution by Mathematica

Time used: 1.114 (sec). Leaf size: 28

```
DSolve[{Exp[x]*(y[x]^3+x*y[x]^3+1)+3*y[x]^2*(x*Exp[x]-6)*y'[x]==0,y[0]==1},y[x],x,IncludeSin
```

$$y(x) \rightarrow \frac{\sqrt[3]{-e^x - 5}}{\sqrt[3]{e^x x - 6}}$$

### 3.12 problem Exact Differential equations. Exercise 9.16, page 79

Internal problem ID [4466]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 9

**Problem number:** Exact Differential equations. Exercise 9.16, page 79.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$\sin(x) \cos(y) + \cos(x) \sin(y) y' = 0$$

With initial conditions

$$\left[ y\left(\frac{\pi}{4}\right) = \frac{\pi}{4} \right]$$

✓ Solution by Maple

Time used: 0.328 (sec). Leaf size: 11

```
dsolve([sin(x)*cos(y(x))+cos(x)*sin(y(x))*diff(y(x),x)=0,y(1/4*Pi) = 1/4*Pi],y(x), singsol=a
```

$$y(x) = \arccos\left(\frac{\sec(x)}{2}\right)$$

✓ Solution by Mathematica

Time used: 6.111 (sec). Leaf size: 12

```
DSolve[{Sin[x]*Cos[y[x]]+Cos[x]*Sin[y[x]]*y'[x]==0,y[Pi/4]==Pi/4},y[x],x,IncludeSingularSolu
```

$$y(x) \rightarrow \arccos\left(\frac{\sec(x)}{2}\right)$$

### 3.13 problem Exact Differential equations. Exercise 9.17, page 79

Internal problem ID [4467]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 9

**Problem number:** Exact Differential equations. Exercise 9.17, page 79.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [exact]

$$y^2 e^{y^2 x} + (2xy e^{y^2 x} - 3y^2) y' = -4x^3$$

With initial conditions

$$[y(1) = 0]$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 23

```
dsolve([(y(x)^2*exp(x*y(x)^2)+4*x^3)+(2*x*y(x)*exp(x*y(x)^2)-3*y(x)^2)*diff(y(x),x)=0,y(1)=0],y(x))
```

$$y(x) = \text{RootOf} \left( -e^{-Z^2 x} - x^4 + \_Z^3 + 2 \right)$$

✓ Solution by Mathematica

Time used: 0.332 (sec). Leaf size: 23

```
DSolve[{(y[x]^2*Exp[x*y[x]^2]+4*x^3)+(2*x*y[x]*Exp[x*y[x]^2]-3*y[x]^2)*y'[x]==0,y[1]==0},y[x]]
```

$$\text{Solve} \left[ x^4 + e^{xy(x)^2} - y(x)^3 = 2, y(x) \right]$$

## 4 Chapter 2. Special types of differential equations of the first kind. Lesson 10

4.1	problem Recognizable Exact Differential equations. Integrating factors. Example 10.51, page 90 . . . . .	56
4.2	problem Recognizable Exact Differential equations. Integrating factors. Example 10.52, page 90 . . . . .	57
4.3	problem Recognizable Exact Differential equations. Integrating factors. Example 10.661, page 90 . . . . .	58
4.4	problem Recognizable Exact Differential equations. Integrating factors. Example 10.701, page 90 . . . . .	59
4.5	problem Recognizable Exact Differential equations. Integrating factors. Example 10.741, page 90 . . . . .	60
4.6	problem Recognizable Exact Differential equations. Integrating factors. Example 10.781, page 90 . . . . .	62
4.7	problem Recognizable Exact Differential equations. Integrating factors. Example 10.81, page 90 . . . . .	63
4.8	problem Recognizable Exact Differential equations. Integrating factors. Example 10.83, page 90 . . . . .	64
4.9	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.1, page 90 . . . . .	66
4.10	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.2, page 90 . . . . .	69
4.11	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.3, page 90 . . . . .	71
4.12	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.4, page 90 . . . . .	72
4.13	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.5, page 90 . . . . .	73
4.14	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.6, page 90 . . . . .	74
4.15	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.7, page 90 . . . . .	75
4.16	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.8, page 90 . . . . .	77
4.17	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.9, page 90 . . . . .	80
4.18	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.10, page 90 . . . . .	81

4.19	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.11, page 90 . . . . .	82
4.20	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.12, page 90 . . . . .	83
4.21	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.13, page 90 . . . . .	85
4.22	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.14, page 90 . . . . .	88
4.23	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.15, page 90 . . . . .	91
4.24	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.16, page 90 . . . . .	92
4.25	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.17, page 90 . . . . .	94
4.26	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.18, page 90 . . . . .	95
4.27	problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.19, page 90 . . . . .	98



## 4.1 problem Recognizable Exact Differential equations. Integrating factors. Example 10.51, page 90

Internal problem ID [4468]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Example 10.51, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y^2 + y - xy' = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
dsolve((y(x)^2+y(x))-x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x}{c_1 - x}$$

### ✓ Solution by Mathematica

Time used: 0.274 (sec). Leaf size: 32

```
DSolve[(y[x]^2+y[x])-x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{c_1 x}}{1 - e^{c_1 x}}$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 0$$

## 4.2 problem Recognizable Exact Differential equations. Integrating factors. Example 10.52, page 90

Internal problem ID [4469]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Example 10.52, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_separable`]

$$y \sec(x) + y' \sin(x) = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve((y(x)*sec(x))+sin(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1}{\tan(x)}$$

### ✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 15

```
DSolve[(y[x]*Sec[x])+Sin[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cot(x)$$

$$y(x) \rightarrow 0$$

### 4.3 problem Recognizable Exact Differential equations. Integrating factors. Example 10.661, page 90

Internal problem ID [4470]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Example 10.661, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type ['y=\_G(x,y)']

$$-\sin(y) + \cos(y) y' = -e^x$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve((exp(x)-sin(y(x)))+cos(y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\arcsin((x + c_1) e^x)$$

#### ✓ Solution by Mathematica

Time used: 11.754 (sec). Leaf size: 16

```
DSolve[(Exp[x]-Sin[y[x]])+Cos[y[x]]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\arcsin(e^x(x + c_1))$$

#### 4.4 problem Recognizable Exact Differential equations. Integrating factors. Example 10.701, page 90

Internal problem ID [4471]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Example 10.701, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$yx + (x^2 + 1)y' = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve((x*y(x))+(1+x^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1}{\sqrt{x^2 + 1}}$$

#### ✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 22

```
DSolve[(x*y[x])+(1+x^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{\sqrt{x^2 + 1}}$$

$$y(x) \rightarrow 0$$

## 4.5 problem Recognizable Exact Differential equations. Integrating factors. Example 10.741, page 90

Internal problem ID [4472]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Example 10.741, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_rational, [_Abel, '2nd type', 'class C']]`

$$y^3 + y^2x + y + (x^3 + x^2y + x) y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 118

```
dsolve((y(x)^3+x*y(x)^2+y(x))+(x^3+x^2*y(x)+x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x^4 + 2x^2 + 1}{x \left( \sqrt{\frac{c_1 x^4 + c_1 x^2 - 1}{x^2(x^2+1)}} (x^2 + 1)^{\frac{3}{2}} - x^2 - 1 \right)}$$
$$y(x) = -\frac{x^4 + 2x^2 + 1}{x \left( x^2 + \sqrt{\frac{c_1 x^4 + c_1 x^2 - 1}{x^2(x^2+1)}} (x^2 + 1)^{\frac{3}{2}} + 1 \right)}$$

✓ Solution by Mathematica

Time used: 3.726 (sec). Leaf size: 114

```
DSolve[(y[x]^3+x*y[x]^2+y[x])+(x^3+x^2*y[x]+x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow -\frac{\sqrt{\frac{1}{x^3}x(x^2+1)}}{\sqrt{\frac{1}{x^3}x^2 - \sqrt{c_1x^3 - \frac{1}{x} + c_1x}}}$$

$$y(x) \rightarrow -\frac{\sqrt{\frac{1}{x^3}x(x^2+1)}}{\sqrt{\frac{1}{x^3}x^2 + \sqrt{c_1x^3 - \frac{1}{x} + c_1x}}}$$

$$y(x) \rightarrow 0$$

## 4.6 problem Recognizable Exact Differential equations. Integrating factors. Example 10.781, page 90

Internal problem ID [4473]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Example 10.781, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$3y - xy' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve((3*y(x))-(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1x^3$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 16

```
DSolve[(3*y[x])-(x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x^3$$

$$y(x) \rightarrow 0$$

## 4.7 problem Recognizable Exact Differential equations. Integrating factors. Example 10.81, page 90

Internal problem ID [4474]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Example 10.81, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_separable`]

$$y - 3xy' = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 9

```
dsolve((y(x))-(3*x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1 x^{\frac{1}{3}}$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 18

```
DSolve[(y[x])-(3*x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \sqrt[3]{x}$$

$$y(x) \rightarrow 0$$



## 4.8 problem Recognizable Exact Differential equations. Integrating factors. Example 10.83, page 90

Internal problem ID [4475]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Example 10.83, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational]`

$$y(2y^3x^2 + 3) + x(y^3x^2 - 1)y' = 0$$

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 39

```
dsolve((y(x)*(2*x^2*y(x)^3+3))+x*(x^2*y(x)^3-1))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{e^{-\frac{11c_1}{3}} x^3}{\text{RootOf}(11 e^{11c_1} \_Z^{15} - e^{11c_1} \_Z^{11} + 4x^{11})^5}$$

✓ Solution by Mathematica

Time used: 10.635 (sec). Leaf size: 1081

`DSolve[(y[x]*(2*x^2*y[x]^3+3))+(x*(x^2*y[x]^3-1))*y'[x]==0,y[x],x,IncludeSingularSolutions`

$$y(x) \rightarrow \text{Root} \left[ 1024\#1^{15}x^{22} + 14080\#1^{12}x^{20} + 77440\#1^9x^{18} + 212960\#1^6x^{16} - \#1^4e^{\frac{44c_1}{3}} \right. \\ \left. + 292820\#1^3x^{14} + 161051x^{12} \&, 1 \right]$$

$$y(x) \rightarrow \text{Root} \left[ 1024\#1^{15}x^{22} + 14080\#1^{12}x^{20} + 77440\#1^9x^{18} + 212960\#1^6x^{16} - \#1^4e^{\frac{44c_1}{3}} \right. \\ \left. + 292820\#1^3x^{14} + 161051x^{12} \&, 2 \right]$$

$$y(x) \rightarrow \text{Root} \left[ 1024\#1^{15}x^{22} + 14080\#1^{12}x^{20} + 77440\#1^9x^{18} + 212960\#1^6x^{16} - \#1^4e^{\frac{44c_1}{3}} \right. \\ \left. + 292820\#1^3x^{14} + 161051x^{12} \&, 3 \right]$$

$$y(x) \rightarrow \text{Root} \left[ 1024\#1^{15}x^{22} + 14080\#1^{12}x^{20} + 77440\#1^9x^{18} + 212960\#1^6x^{16} - \#1^4e^{\frac{44c_1}{3}} \right. \\ \left. + 292820\#1^3x^{14} + 161051x^{12} \&, 4 \right]$$

$$y(x) \rightarrow \text{Root} \left[ 1024\#1^{15}x^{22} + 14080\#1^{12}x^{20} + 77440\#1^9x^{18} + 212960\#1^6x^{16} - \#1^4e^{\frac{44c_1}{3}} \right. \\ \left. + 292820\#1^3x^{14} + 161051x^{12} \&, 5 \right]$$

$$y(x) \rightarrow \text{Root} \left[ 1024\#1^{15}x^{22} + 14080\#1^{12}x^{20} + 77440\#1^9x^{18} + 212960\#1^6x^{16} - \#1^4e^{\frac{44c_1}{3}} \right. \\ \left. + 292820\#1^3x^{14} + 161051x^{12} \&, 6 \right]$$

$$y(x) \rightarrow \text{Root} \left[ 1024\#1^{15}x^{22} + 14080\#1^{12}x^{20} + 77440\#1^9x^{18} + 212960\#1^6x^{16} - \#1^4e^{\frac{44c_1}{3}} \right. \\ \left. + 292820\#1^3x^{14} + 161051x^{12} \&, 7 \right]$$

$$y(x) \rightarrow \text{Root} \left[ 1024\#1^{15}x^{22} + 14080\#1^{12}x^{20} + 77440\#1^9x^{18} + 212960\#1^6x^{16} - \#1^4e^{\frac{44c_1}{3}} \right. \\ \left. + 292820\#1^3x^{14} + 161051x^{12} \&, 8 \right]$$

$$y(x) \rightarrow \text{Root} \left[ 1024\#1^{15}x^{22} + 14080\#1^{12}x^{20} + 77440\#1^9x^{18} + 212960\#1^6x^{16} - \#1^4e^{\frac{44c_1}{3}} \right. \\ \left. + 292820\#1^3x^{14} + 161051x^{12} \&, 9 \right]$$

$$y(x) \rightarrow \text{Root} \left[ 1024\#1^{15}x^{22} + 14080\#1^{12}x^{20} + 77440\#1^9x^{18} + 212960\#1^6x^{16} - \#1^4e^{\frac{44c_1}{3}} \right. \\ \left. + 292820\#1^3x^{14} + 161051x^{12} \&, 10 \right]$$

$$y(x) \rightarrow \text{Root} \left[ 1024\#1^{15}x^{22} + 14080\#1^{12}x^{20} + 77440\#1^9x^{18} + 212960\#1^6x^{16} - \#1^4e^{\frac{44c_1}{3}} \right. \\ \left. + 292820\#1^3x^{14} + 161051x^{12} \&, 11 \right]$$

## 4.9 problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.1, page 90

Internal problem ID [4476]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.1, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _exact, _rational, _dAlembert]`

$$2yx + (x^2 + y^2) y' = -x^2$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 417

`dsolve((2*x*y(x)+x^2)+(x^2+y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)`

$$y(x) = \frac{\left(4-4x^3c_1^{\frac{3}{2}}+4\sqrt{5x^6c_1^3-2x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}{2} - \frac{2x^2c_1}{\left(4-4x^3c_1^{\frac{3}{2}}+4\sqrt{5x^6c_1^3-2x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}$$

$y(x)$

$$= \frac{\left(4-4x^3c_1^{\frac{3}{2}}+4\sqrt{5x^6c_1^3-2x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}{4} + \frac{x^2c_1}{\left(4-4x^3c_1^{\frac{3}{2}}+4\sqrt{5x^6c_1^3-2x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}} - \frac{i\sqrt{3}\left(\frac{\left(4-4x^3c_1^{\frac{3}{2}}+4\sqrt{5x^6c_1^3-2x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}{2} + \frac{1}{\left(4-4x^3c_1^{\frac{3}{2}}+4\sqrt{5x^6c_1^3-2x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}\right)}{2}$$

$y(x)$

$$= \frac{\left(4-4x^3c_1^{\frac{3}{2}}+4\sqrt{5x^6c_1^3-2x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}{4} + \frac{x^2c_1}{\left(4-4x^3c_1^{\frac{3}{2}}+4\sqrt{5x^6c_1^3-2x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}} + \frac{i\sqrt{3}\left(\frac{\left(4-4x^3c_1^{\frac{3}{2}}+4\sqrt{5x^6c_1^3-2x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}{2} + \frac{1}{\left(4-4x^3c_1^{\frac{3}{2}}+4\sqrt{5x^6c_1^3-2x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}\right)}{2}$$

✓ Solution by Mathematica

Time used: 23.867 (sec). Leaf size: 597

`DSolve[(2*x*y[x]+x^2)+(x^2+y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{\sqrt[3]{-x^3 + \sqrt{5x^6 - 2e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}}{\sqrt[3]{2}} - \frac{\sqrt[3]{2}x^2}{\sqrt[3]{-x^3 + \sqrt{5x^6 - 2e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{2}(2 + 2i\sqrt{3})x^2 + i2^{2/3}(\sqrt{3} + i)(-x^3 + \sqrt{5x^6 - 2e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1})^{2/3}}{4\sqrt[3]{-x^3 + \sqrt{5x^6 - 2e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}}$$

$$y(x) \rightarrow \frac{(1 - i\sqrt{3})x^2}{2^{2/3}\sqrt[3]{-x^3 + \sqrt{5x^6 - 2e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}} - \frac{(1 + i\sqrt{3})\sqrt[3]{-x^3 + \sqrt{5x^6 - 2e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}}{2\sqrt[3]{2}}$$

$$y(x) \rightarrow \frac{2\sqrt[3]{-2x^2} + (-2)^{2/3}(\sqrt{5}\sqrt{x^6} - x^3)^{2/3}}{2\sqrt[3]{\sqrt{5}\sqrt{x^6} - x^3}}$$

$$y(x) \rightarrow \frac{(2\sqrt{5}\sqrt{x^6} - 2x^3)^{2/3} - 2\sqrt[3]{2}x^2}{2\sqrt[3]{\sqrt{5}\sqrt{x^6} - x^3}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{2}(2 - 2i\sqrt{3})x^2 + (-1 - i\sqrt{3})(2\sqrt{5}\sqrt{x^6} - 2x^3)^{2/3}}{4\sqrt[3]{\sqrt{5}\sqrt{x^6} - x^3}}$$

## 4.10 problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.2, page 90

Internal problem ID [4477]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.2, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [exact]

$$y \cos(x) + (y^3 + \sin(x)) y' = -x^2$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 21

```
dsolve((x^2+y(x)*cos(x))+(y(x)^3+sin(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$\frac{x^3}{3} + y(x) \sin(x) + \frac{y(x)^4}{4} + c_1 = 0$$

✓ Solution by Mathematica

Time used: 60.198 (sec). Leaf size: 1119

`DSolve[(x^2+y[x]*Cos[x])+(y[x]^3+Sin[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{\sqrt{\frac{4x^3 + (27\sin^2(x) + \sqrt{729\sin^4(x) - 64(x^3 - 3c_1)^3})^{2/3} - 12c_1}{3\sqrt[3]{27\sin^2(x) + \sqrt{729\sin^4(x) - 64(x^3 - 3c_1)^3}}}}}{\sqrt{6}}$$

$$-\frac{1}{2} \sqrt{\frac{8(x^3 - 3c_1)}{3\sqrt[3]{27\sin^2(x) + \sqrt{729\sin^4(x) - 64(x^3 - 3c_1)^3}}} - \frac{2}{3}\sqrt[3]{27\sin^2(x) + \sqrt{729\sin^4(x) - 64(x^3 - 3c_1)^3}}}$$

$$y(x) \rightarrow \frac{\sqrt{\frac{4x^3 + (27\sin^2(x) + \sqrt{729\sin^4(x) - 64(x^3 - 3c_1)^3})^{2/3} - 12c_1}{3\sqrt[3]{27\sin^2(x) + \sqrt{729\sin^4(x) - 64(x^3 - 3c_1)^3}}}}}{\sqrt{6}}$$

$$+\frac{1}{2} \sqrt{\frac{8(x^3 - 3c_1)}{3\sqrt[3]{27\sin^2(x) + \sqrt{729\sin^4(x) - 64(x^3 - 3c_1)^3}}} - \frac{2}{3}\sqrt[3]{27\sin^2(x) + \sqrt{729\sin^4(x) - 64(x^3 - 3c_1)^3}}}$$

$$y(x) \rightarrow -\frac{\sqrt{\frac{4x^3 + (27\sin^2(x) + \sqrt{729\sin^4(x) - 64(x^3 - 3c_1)^3})^{2/3} - 12c_1}{3\sqrt[3]{27\sin^2(x) + \sqrt{729\sin^4(x) - 64(x^3 - 3c_1)^3}}}}}{\sqrt{6}}$$

$$-\frac{1}{2} \sqrt{\frac{8(x^3 - 3c_1)}{3\sqrt[3]{27\sin^2(x) + \sqrt{729\sin^4(x) - 64(x^3 - 3c_1)^3}}} - \frac{2}{3}\sqrt[3]{27\sin^2(x) + \sqrt{729\sin^4(x) - 64(x^3 - 3c_1)^3}}}$$

$$y(x) \rightarrow \frac{1}{2} \sqrt{\frac{8(x^3 - 3c_1)}{3\sqrt[3]{27\sin^2(x) + \sqrt{729\sin^4(x) - 64(x^3 - 3c_1)^3}}} - \frac{2}{3}\sqrt[3]{27\sin^2(x) + \sqrt{729\sin^4(x) - 64(x^3 - 3c_1)^3}}}$$

## 4.11 problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.3, page 90

Internal problem ID [4478]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.3, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_rational, _Bernoulli]`

$$yy'x + y^2 = -x^2 - x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

```
dsolve((x^2+y(x)^2+x)+(x*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{-18x^4 - 24x^3 + 36c_1}}{6x}$$

$$y(x) = \frac{\sqrt{-18x^4 - 24x^3 + 36c_1}}{6x}$$

### ✓ Solution by Mathematica

Time used: 0.242 (sec). Leaf size: 60

```
DSolve[(x^2+y[x]^2+x)+(x*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{-\frac{x^4}{2} - \frac{2x^3}{3} + c_1}}{x}$$

$$y(x) \rightarrow \frac{\sqrt{-\frac{x^4}{2} - \frac{2x^3}{3} + c_1}}{x}$$



## 4.12 problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.4, page 90

Internal problem ID [4479]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.4, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_exact]`

$$-2yx + e^y + (y - x^2 + x e^y) y' = -x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve((x-2*x*y(x)+exp(y(x)))+(y(x)-x^2+x*exp(y(x)))*diff(y(x),x)=0,y(x), singsol=all)
```

$$-y(x)x^2 + x e^{y(x)} + \frac{x^2}{2} + \frac{y(x)^2}{2} + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.316 (sec). Leaf size: 35

```
DSolve[(x-2*x*y[x]+Exp[y[x]])+(y[x]-x^2+x*Exp[y[x]])*y'[x]==0,y[x],x,IncludeSingularSolution
```

$$\text{Solve}\left[x^2(-y(x)) + \frac{x^2}{2} + x e^{y(x)} + \frac{y(x)^2}{2} = c_1, y(x)\right]$$

### 4.13 problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.5, page 90

Internal problem ID [4480]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.5, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_exact]`

$$e^x \sin(y) + e^{-y} - (x e^{-y} - e^x \cos(y)) y' = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve((exp(x)*sin(y(x))+exp(-y(x)))-(x*exp(-y(x))-exp(x)*cos(y(x)))*diff(y(x),x)=0,y(x), si
```

$$e^x \sin(y(x)) + x e^{-y(x)} + c_1 = 0$$

#### ✓ Solution by Mathematica

Time used: 0.377 (sec). Leaf size: 24

```
DSolve[(Exp[x]*Sin[y[x]]+Exp[-y[x]])-(x*Exp[-y[x]]-Exp[x]*Cos[y[x]])*y'[x]==0,y[x],x,Include
```

$$\text{Solve}[x(-e^{-y(x)}) - e^x \sin(y(x)) = c_1, y(x)]$$

## 4.14 problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.6, page 90

Internal problem ID [4481]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.6, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _rational]`

$$-y^2 - y - (x^2 - y^2 - x) y' = -x^2$$

### ✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 28

```
dsolve((x^2-y(x)^2-y(x))-(x^2-y(x)^2-x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$-2y(x) + \ln(x + y(x)) - \ln(y(x) - x) + 2x - c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 0.242 (sec). Leaf size: 32

```
DSolve[(x^2-y[x]^2-y[x])-(x^2-y[x]^2-x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ -\frac{e^{2x-2y(x)}(y(x) + x)}{2(x - y(x))} = c_1, y(x) \right]$$

#### 4.15 problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.7, page 90

Internal problem ID [4482]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.7, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_rational]`

$$y^2 x^4 - y + (y^4 x^2 - x) y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve((x^4*y(x)^2-y(x))+(x^2*y(x)^4-x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$-\frac{x^3}{3} - \frac{1}{xy(x)} - \frac{y(x)^3}{3} + c_1 = 0$$

✓ Solution by Mathematica

Time used: 60.131 (sec). Leaf size: 1507

`DSolve[(x^4*y[x]^2-y[x])+(x^2*y[x]^4-x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{1}{4} \left( \sqrt{2} \sqrt{\frac{8\sqrt[3]{2}x + 2^{2/3} \left( x^9 - 6c_1x^6 + 9c_1^2x^3 + \sqrt{x^2(-256x + (x^4 - 3c_1x)^4) \right)^{2/3}}{x^3 \sqrt{x^9 - 6c_1x^6 + 9c_1^2x^3 + \sqrt{x^2(-256x + (x^4 - 3c_1x)^4)}}}} \right. \\ \left. - 2 \sqrt{\frac{\sqrt[3]{x(x^4 - 3c_1x)^2 + \sqrt{x^2(-256x + (x^4 - 3c_1x)^4)}}}{\sqrt[3]{2}x} - \frac{2\sqrt{2}(x^3 - 3c_1)}{\sqrt{x^3 \sqrt{x^9 - 6c_1x^6 + 9c_1^2x^3 + \sqrt{x^2(-256x + (x^4 - 3c_1x)^4)}}}} \right)$$

$$y(x) \rightarrow \frac{1}{4} \left( \sqrt{2} \sqrt{\frac{8\sqrt[3]{2}x + 2^{2/3} \left( x^9 - 6c_1x^6 + 9c_1^2x^3 + \sqrt{x^2(-256x + (x^4 - 3c_1x)^4) \right)^{2/3}}{x^3 \sqrt{x^9 - 6c_1x^6 + 9c_1^2x^3 + \sqrt{x^2(-256x + (x^4 - 3c_1x)^4)}}}} \right. \\ \left. + 2 \sqrt{\frac{\sqrt[3]{x(x^4 - 3c_1x)^2 + \sqrt{x^2(-256x + (x^4 - 3c_1x)^4)}}}{\sqrt[3]{2}x} - \frac{2\sqrt{2}(x^3 - 3c_1)}{\sqrt{x^3 \sqrt{x^9 - 6c_1x^6 + 9c_1^2x^3 + \sqrt{x^2(-256x + (x^4 - 3c_1x)^4)}}}} \right)$$

$$y(x) \rightarrow \frac{1}{4} \left( -\sqrt{2} \sqrt{\frac{8\sqrt[3]{2}x + 2^{2/3} \left( x^9 - 6c_1x^6 + 9c_1^2x^3 + \sqrt{x^2(-256x + (x^4 - 3c_1x)^4) \right)^{2/3}}{x^3 \sqrt{x^9 - 6c_1x^6 + 9c_1^2x^3 + \sqrt{x^2(-256x + (x^4 - 3c_1x)^4)}}}} \right. \\ \left. - 2 \sqrt{\frac{\sqrt[3]{x(x^4 - 3c_1x)^2 + \sqrt{x^2(-256x + (x^4 - 3c_1x)^4)}}}{\sqrt[3]{2}x} + \frac{2\sqrt{2}(x^3 - 3c_1)}{\sqrt{x^3 \sqrt{x^9 - 6c_1x^6 + 9c_1^2x^3 + \sqrt{x^2(-256x + (x^4 - 3c_1x)^4)}}}} \right)$$

#### 4.16 problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.8, page 90

Internal problem ID [4483]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.8, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational]`

$$y(2x + y^3) - x(2x - y^3) y' = 0$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 420

`dsolve((y(x)*(2*x+y(x)^3))-(x*(2*x-y(x)^3))*diff(y(x),x)=0,y(x), singsol=all)`

$$y(x) = \frac{\left(-108x^4 + 8c_1^3 + 12\sqrt{81x^4 - 12c_1^3 x^2}\right)^{\frac{1}{3}}}{6x} + \frac{2c_1^2}{3x \left(-108x^4 + 8c_1^3 + 12\sqrt{81x^4 - 12c_1^3 x^2}\right)^{\frac{1}{3}}} + \frac{c_1}{3x}$$

$$y(x) = -\frac{\left(-108x^4 + 8c_1^3 + 12\sqrt{81x^4 - 12c_1^3 x^2}\right)^{\frac{1}{3}}}{12x} - \frac{c_1^2}{3x \left(-108x^4 + 8c_1^3 + 12\sqrt{81x^4 - 12c_1^3 x^2}\right)^{\frac{1}{3}}} + \frac{c_1}{3x} - \frac{i\sqrt{3} \left( \frac{\left(-108x^4 + 8c_1^3 + 12\sqrt{81x^4 - 12c_1^3 x^2}\right)^{\frac{1}{3}}}{6x} - \frac{2c_1^2}{3x \left(-108x^4 + 8c_1^3 + 12\sqrt{81x^4 - 12c_1^3 x^2}\right)^{\frac{1}{3}}} \right)}{2}$$

$$y(x) = -\frac{\left(-108x^4 + 8c_1^3 + 12\sqrt{81x^4 - 12c_1^3 x^2}\right)^{\frac{1}{3}}}{12x} - \frac{c_1^2}{3x \left(-108x^4 + 8c_1^3 + 12\sqrt{81x^4 - 12c_1^3 x^2}\right)^{\frac{1}{3}}} + \frac{c_1}{3x} + \frac{i\sqrt{3} \left( \frac{\left(-108x^4 + 8c_1^3 + 12\sqrt{81x^4 - 12c_1^3 x^2}\right)^{\frac{1}{3}}}{6x} - \frac{2c_1^2}{3x \left(-108x^4 + 8c_1^3 + 12\sqrt{81x^4 - 12c_1^3 x^2}\right)^{\frac{1}{3}}} \right)}{2}$$

✓ Solution by Mathematica

Time used: 11.386 (sec). Leaf size: 371

`DSolve[(y[x]*(2*x+y[x]^3))-(x*(2*x-y[x]^3))*y'[x]==0,y[x],x,IncludeSingularSolutions -> True`

$y(x) \rightarrow$

$$\frac{\frac{2^{\frac{3}{2}}\sqrt{2}c_1^2}{\sqrt[3]{27x^4 + 3\sqrt{81x^8 + 12c_1^3x^4 + 2c_1^3}}} + 2^{2/3}\sqrt[3]{27x^4 + 3\sqrt{81x^8 + 12c_1^3x^4 + 2c_1^3}} + 2c_1}{6x}$$

$y(x)$

$$\rightarrow \frac{\frac{2^{\frac{3}{2}}\sqrt{2}(1+i\sqrt{3})c_1^2}{\sqrt[3]{27x^4 + 3\sqrt{81x^8 + 12c_1^3x^4 + 2c_1^3}}} + 2^{2/3}(1-i\sqrt{3})\sqrt[3]{27x^4 + 3\sqrt{81x^8 + 12c_1^3x^4 + 2c_1^3}} - 4c_1}{12x}$$

$y(x)$

$$\rightarrow \frac{\frac{2^{\frac{3}{2}}\sqrt{2}(1-i\sqrt{3})c_1^2}{\sqrt[3]{27x^4 + 3\sqrt{81x^8 + 12c_1^3x^4 + 2c_1^3}}} + 2^{2/3}(1+i\sqrt{3})\sqrt[3]{27x^4 + 3\sqrt{81x^8 + 12c_1^3x^4 + 2c_1^3}} - 4c_1}{12x}$$

$y(x) \rightarrow 0$



## 4.17 problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.9, page 90

Internal problem ID [4484]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.9, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [exact]

$$\arctan(yx) + \frac{yx - 2y^2x}{1 + x^2y^2} + \frac{(x^2 - 2x^2y)y'}{1 + x^2y^2} = 0$$

### ✓ Solution by Maple

Time used: 0.093 (sec). Leaf size: 24

```
dsolve((arctan(x*y(x))+(x*y(x)-2*x*y(x)^2)/(1+x^2*y(x)^2))+((x^2-2*x^2*y(x))/(1+x^2*y(x)^2))
```

$$y(x) = \frac{\tan(\text{RootOf}(\_Zx - \ln(\tan(\_Z)^2 + 1) + c_1))}{x}$$

### ✓ Solution by Mathematica

Time used: 0.173 (sec). Leaf size: 26

```
DSolve[(ArcTan[x*y[x]]+(x*y[x]-2*x*y[x]^2)/(1+x^2*y[x]^2))+((x^2-2*x^2*y[x])/(1+x^2*y[x]^2))
```

$$\text{Solve}[\log(x^2y(x)^2 + 1) - x \arctan(xy(x)) = c_1, y(x)]$$

## 4.18 problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.10, page 90

Internal problem ID [4485]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.10, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type ['y=\_G(x,y)']

$$(y e^y - x e^x) y' = -e^x(x + 1)$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve((exp(x)*(x+1))+(y(x)*exp(y(x))-x*exp(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$x e^{-y(x)+x} + \frac{y(x)^2}{2} + c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 0.291 (sec). Leaf size: 26

```
DSolve[(Exp[x]*(x+1))+(y[x]*Exp[y[x]]-x*Exp[x])*y'[x]==0,y[x],x,IncludeSingularSolutions ->
```

$$\text{Solve} \left[ -\frac{1}{2}y(x)^2 - x e^{x-y(x)} = c_1, y(x) \right]$$

## 4.19 problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.11, page 90

Internal problem ID [4486]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.11, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class D', _exact, _rational, [_Abel, '2nd ty`

$$\frac{yx + 1}{y} + \frac{(-x + 2y)y'}{y^2} = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 20

```
dsolve(((x*y(x)+1)/y(x))+((2*y(x)-x)/y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{x}{2 \operatorname{LambertW}\left(-\frac{e^{\frac{x^2}{4}} c_1 x}{2}\right)}$$

### ✓ Solution by Mathematica

Time used: 3.618 (sec). Leaf size: 37

```
DSolve[((x*y[x]+1)/y[x])+((2*y[x]-x)/y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x}{2W\left(-\frac{1}{2}xe^{\frac{1}{4}(x^2-2c_1)}\right)}$$

$$y(x) \rightarrow 0$$

## 4.20 problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.12, page 90

Internal problem ID [4487]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.12, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$y^2 - 3yx + (yx - x^2) y' = 2x^2$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 59

```
dsolve((y(x)^2-3*x*y(x)-2*x^2)+(x*y(x)-x^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 x^2 - \sqrt{2c_1^2 x^4 + 1}}{c_1 x}$$

$$y(x) = \frac{c_1 x^2 + \sqrt{2c_1^2 x^4 + 1}}{c_1 x}$$

✓ Solution by Mathematica

Time used: 0.657 (sec). Leaf size: 99

```
DSolve[(y[x]^2-3*x*y[x]-2*x^2)+(x*y[x]-x^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow x - \frac{\sqrt{2x^4 + e^{2c_1}}}{x}$$

$$y(x) \rightarrow x + \frac{\sqrt{2x^4 + e^{2c_1}}}{x}$$

$$y(x) \rightarrow x - \frac{\sqrt{2}\sqrt{x^4}}{x}$$

$$y(x) \rightarrow \frac{\sqrt{2}\sqrt{x^4}}{x} + x$$

**4.21 problem Recognizable Exact Differential equations.  
Integrating factors. Exercise 10.13, page 90**

Internal problem ID [4488]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.13, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_rational, [_Abel, '2nd type', 'class B']]`

$$(1 + 2x + y)y - x(-1 + x + 2y)y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 493

`dsolve((y(x)*(y(x)+2*x+1))-(x*(2*y(x)+x-1))*diff(y(x),x)=0,y(x), singsol=all)`

$$\begin{aligned}
 y(x) &= \frac{3 \cdot 5^{\frac{1}{3}} \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2 - 160c_1x + 80c_1 - x}{c_1}} + 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}}{40c_1} \\
 &\quad + \frac{3x5^{\frac{2}{3}}}{40 \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2 - 160c_1x + 80c_1 - x}{c_1}} + 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}} + x - 1 \\
 y(x) &= - \frac{3 \cdot 5^{\frac{1}{3}} \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2 - 160c_1x + 80c_1 - x}{c_1}} + 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}}{80c_1} \\
 &\quad - \frac{3x5^{\frac{2}{3}}}{80 \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2 - 160c_1x + 80c_1 - x}{c_1}} + 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}} + x - 1 \\
 &\quad - \frac{i\sqrt{3} \left( \frac{3 \cdot 5^{\frac{1}{3}} \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2 - 160c_1x + 80c_1 - x}{c_1}} + 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}}{40c_1} - \frac{3x5^{\frac{2}{3}}}{40 \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2 - 160c_1x + 80c_1 - x}{c_1}} + 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}} \right)}{2} \\
 y(x) &= - \frac{3 \cdot 5^{\frac{1}{3}} \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2 - 160c_1x + 80c_1 - x}{c_1}} + 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}}{80c_1} \\
 &\quad - \frac{3x5^{\frac{2}{3}}}{80 \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2 - 160c_1x + 80c_1 - x}{c_1}} + 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}} + x - 1 \\
 &\quad + \frac{i\sqrt{3} \left( \frac{3 \cdot 5^{\frac{1}{3}} \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2 - 160c_1x + 80c_1 - x}{c_1}} + 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}}{40c_1} - \frac{3x5^{\frac{2}{3}}}{40 \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2 - 160c_1x + 80c_1 - x}{c_1}} + 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}} \right)}{2}
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 41.715 (sec). Leaf size: 463

DSolve[(y[x]\*(y[x]+2\*x+1))-(x\*(2\*y[x]+x-1))\*y'[x]==0,y[x],x,IncludeSingularSolutions -> True

$$y(x) \rightarrow -\frac{\sqrt[3]{2}x}{\sqrt[3]{-27c_1^2x^2 + \sqrt{108c_1^3x^3 + (27c_1^2x - 27c_1^2x^2)^2} + 27c_1^2x}} + \frac{\sqrt[3]{-27c_1^2x^2 + \sqrt{108c_1^3x^3 + (27c_1^2x - 27c_1^2x^2)^2} + 27c_1^2x}}{3\sqrt[3]{2}c_1} + x - 1$$

$$y(x) \rightarrow \frac{(1 + i\sqrt{3})x}{2^{2/3}\sqrt[3]{-27c_1^2x^2 + \sqrt{108c_1^3x^3 + (27c_1^2x - 27c_1^2x^2)^2} + 27c_1^2x}} - \frac{(1 - i\sqrt{3})\sqrt[3]{-27c_1^2x^2 + \sqrt{108c_1^3x^3 + (27c_1^2x - 27c_1^2x^2)^2} + 27c_1^2x}}{6\sqrt[3]{2}c_1} + x - 1$$

$$y(x) \rightarrow \frac{(1 - i\sqrt{3})x}{2^{2/3}\sqrt[3]{-27c_1^2x^2 + \sqrt{108c_1^3x^3 + (27c_1^2x - 27c_1^2x^2)^2} + 27c_1^2x}} - \frac{(1 + i\sqrt{3})\sqrt[3]{-27c_1^2x^2 + \sqrt{108c_1^3x^3 + (27c_1^2x - 27c_1^2x^2)^2} + 27c_1^2x}}{6\sqrt[3]{2}c_1} + x - 1$$

$y(x) \rightarrow$  Indeterminate

$y(x) \rightarrow x - 1$



**4.22 problem Recognizable Exact Differential equations.  
Integrating factors. Exercise 10.14, page 90**

Internal problem ID [4489]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.14, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_rational, [_Abel, '2nd type', 'class B']]`

$$y(2x - y - 1) + x(-1 - x + 2y)y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 499

`dsolve((y(x)*(2*x-y(x)-1))+(x*(2*y(x)-x-1))*diff(y(x),x)=0,y(x), singsol=all)`

$$\begin{aligned}
 y(x) &= \frac{3 \cdot 5^{\frac{1}{3}} \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2+160c_1x+80c_1-x}{c_1}} - 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}}{40c_1} \\
 &\quad + \frac{3x5^{\frac{2}{3}}}{40 \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2+160c_1x+80c_1-x}{c_1}} - 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}} - x - 1 \\
 y(x) &= - \frac{3 \cdot 5^{\frac{1}{3}} \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2+160c_1x+80c_1-x}{c_1}} - 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}}{80c_1} \\
 &\quad - \frac{3x5^{\frac{2}{3}}}{80 \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2+160c_1x+80c_1-x}{c_1}} - 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}} - x - 1 \\
 &\quad - \frac{i\sqrt{3} \left( \frac{3 \cdot 5^{\frac{1}{3}} \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2+160c_1x+80c_1-x}{c_1}} - 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}}{40c_1} - \frac{3x5^{\frac{2}{3}}}{40 \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2+160c_1x+80c_1-x}{c_1}} - 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}} \right)}{2} \\
 y(x) &= - \frac{3 \cdot 5^{\frac{1}{3}} \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2+160c_1x+80c_1-x}{c_1}} - 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}}{80c_1} \\
 &\quad - \frac{3x5^{\frac{2}{3}}}{80 \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2+160c_1x+80c_1-x}{c_1}} - 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}} - x - 1 \\
 &\quad + \frac{i\sqrt{3} \left( \frac{3 \cdot 5^{\frac{1}{3}} \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2+160c_1x+80c_1-x}{c_1}} - 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}}{40c_1} - \frac{3x5^{\frac{2}{3}}}{40 \left( x \left( \sqrt{5} \sqrt{\frac{80c_1x^2+160c_1x+80c_1-x}{c_1}} - 20x - 20 \right) c_1^2 \right)^{\frac{1}{3}}} \right)}{2}
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 40.285 (sec). Leaf size: 471

`DSolve[(y[x]*(2*x-y[x]-1))+(x*(2*y[x]-x-1))*y'[x]==0,y[x],x,IncludeSingularSolutions -> True`

$$y(x) \rightarrow -\frac{\sqrt[3]{2}x}{\sqrt[3]{27c_1^2x^2 + \sqrt{(27c_1^2x^2 + 27c_1^2x)^2 - 108c_1^3x^3} + 27c_1^2x}} - \frac{\sqrt[3]{27c_1^2x^2 + \sqrt{(27c_1^2x^2 + 27c_1^2x)^2 - 108c_1^3x^3} + 27c_1^2x}}{3\sqrt[3]{2}c_1} - x - 1$$

$$y(x) \rightarrow \frac{(1 + i\sqrt{3})x}{2^{2/3}\sqrt[3]{27c_1^2x^2 + \sqrt{(27c_1^2x^2 + 27c_1^2x)^2 - 108c_1^3x^3} + 27c_1^2x}} + \frac{(1 - i\sqrt{3})\sqrt[3]{27c_1^2x^2 + \sqrt{(27c_1^2x^2 + 27c_1^2x)^2 - 108c_1^3x^3} + 27c_1^2x}}{6\sqrt[3]{2}c_1} - x - 1$$

$$y(x) \rightarrow \frac{(1 - i\sqrt{3})x}{2^{2/3}\sqrt[3]{27c_1^2x^2 + \sqrt{(27c_1^2x^2 + 27c_1^2x)^2 - 108c_1^3x^3} + 27c_1^2x}} + \frac{(1 + i\sqrt{3})\sqrt[3]{27c_1^2x^2 + \sqrt{(27c_1^2x^2 + 27c_1^2x)^2 - 108c_1^3x^3} + 27c_1^2x}}{6\sqrt[3]{2}c_1} - x - 1$$

$y(x) \rightarrow$  Indeterminate

$y(x) \rightarrow -x - 1$

## 4.23 problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.15, page 90

Internal problem ID [4490]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.15, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class G'], _exact, _rational, [_Abel, '2nd ty`

$$y^2 + 12x^2y + (2yx + 4x^3)y' = 0$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 50

```
dsolve((y(x)^2+12*x^2*y(x))+(2*x*y(x)+4*x^3)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-2x^3 + \sqrt{4x^6 + c_1x}}{x}$$

$$y(x) = -\frac{2x^3 + \sqrt{4x^6 + c_1x}}{x}$$

### ✓ Solution by Mathematica

Time used: 0.431 (sec). Leaf size: 58

```
DSolve[(y[x]^2+12*x^2*y[x])+(2*x*y[x]+4*x^3)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2x^3 + \sqrt{x(4x^5 + c_1)}}{x}$$

$$y(x) \rightarrow \frac{-2x^3 + \sqrt{x(4x^5 + c_1)}}{x}$$

## 4.24 problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.16, page 90

Internal problem ID [4491]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.16, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$3(x + y)^2 + x(2x + 3y)y' = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 63

```
dsolve((3*(y(x)+x)^2)+(x*(3*y(x)+2*x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-\frac{2c_1x^2}{3} - \frac{\sqrt{-2c_1^2x^4+6}}{6}}{c_1x}$$

$$y(x) = \frac{-\frac{2c_1x^2}{3} + \frac{\sqrt{-2c_1^2x^4+6}}{6}}{c_1x}$$

✓ Solution by Mathematica

Time used: 1.741 (sec). Leaf size: 135

```
DSolve[(3*(y[x]+x)^2)+(x*(3*y[x]+2*x))*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{4x^2 + \sqrt{-2x^4 + 6e^{4c_1}}}{6x}$$

$$y(x) \rightarrow \frac{-4x^2 + \sqrt{-2x^4 + 6e^{4c_1}}}{6x}$$

$$y(x) \rightarrow -\frac{\sqrt{2}\sqrt{-x^4} + 4x^2}{6x}$$

$$y(x) \rightarrow \frac{\sqrt{2}\sqrt{-x^4} - 4x^2}{6x}$$

## 4.25 problem Recognizable Exact Differential equations. Integrating factors. Exercise 10.17, page 90

Internal problem ID [4492]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.17, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_rational]`

$$y - (x + x^2 + y^2) y' = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 30

```
dsolve((y(x))-(y(x)^2+x^2+x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$c_1 + \frac{e^{-2iy(x)}(ix + y(x))}{2iy(x) + 2x} = 0$$

✓ Solution by Mathematica

Time used: 0.102 (sec). Leaf size: 18

```
DSolve[(y[x])-(y[x]^2+x^2+x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ y(x) - \arctan \left( \frac{x}{y(x)} \right) = c_1, y(x) \right]$$

**4.26 problem Recognizable Exact Differential equations.  
Integrating factors. Exercise 10.18, page 90**

Internal problem ID [4493]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.18, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_exact, _rational, [_1st_order, ' _with_symmetry_[F(x)*G(y),0]`

$$2yx + (a + x^2 + y^2) y' = 0$$



✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 470

`dsolve((2*x*y(x))+(x^2+y(x)^2+a)*diff(y(x),x)=0,y(x), singsol=all)`

$$\begin{aligned}
 y(x) &= \frac{\left(-12c_1 + 4\sqrt{4x^6 + 12ax^4 + 12a^2x^2 + 4a^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2(x^2 + a)} \\
 &\quad - \frac{\left(-12c_1 + 4\sqrt{4x^6 + 12ax^4 + 12a^2x^2 + 4a^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2(x^2 + a)} \\
 y(x) &= -\frac{\left(-12c_1 + 4\sqrt{4x^6 + 12ax^4 + 12a^2x^2 + 4a^3 + 9c_1^2}\right)^{\frac{1}{3}}}{4(x^2 + a)} \\
 &\quad + \frac{\left(-12c_1 + 4\sqrt{4x^6 + 12ax^4 + 12a^2x^2 + 4a^3 + 9c_1^2}\right)^{\frac{1}{3}}}{4(x^2 + a)} \\
 &\quad - \frac{i\sqrt{3} \left( \frac{\left(-12c_1 + 4\sqrt{4x^6 + 12ax^4 + 12a^2x^2 + 4a^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2} + \frac{2x^2 + 2a}{\left(-12c_1 + 4\sqrt{4x^6 + 12ax^4 + 12a^2x^2 + 4a^3 + 9c_1^2}\right)^{\frac{1}{3}}} \right)}{2} \\
 y(x) &= -\frac{\left(-12c_1 + 4\sqrt{4x^6 + 12ax^4 + 12a^2x^2 + 4a^3 + 9c_1^2}\right)^{\frac{1}{3}}}{4(x^2 + a)} \\
 &\quad + \frac{\left(-12c_1 + 4\sqrt{4x^6 + 12ax^4 + 12a^2x^2 + 4a^3 + 9c_1^2}\right)^{\frac{1}{3}}}{4(x^2 + a)} \\
 &\quad + \frac{i\sqrt{3} \left( \frac{\left(-12c_1 + 4\sqrt{4x^6 + 12ax^4 + 12a^2x^2 + 4a^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2} + \frac{2x^2 + 2a}{\left(-12c_1 + 4\sqrt{4x^6 + 12ax^4 + 12a^2x^2 + 4a^3 + 9c_1^2}\right)^{\frac{1}{3}}} \right)}{2}
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 4.319 (sec). Leaf size: 299

```
DSolve[(2*x*y[x])+(x^2+y[x]^2+a)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt[3]{2} \left( \sqrt{4(a+x^2)^3 + 9c_1^2 + 3c_1} \right)^{2/3} - 2a - 2x^2}{2^{2/3} \sqrt[3]{\sqrt{4(a+x^2)^3 + 9c_1^2 + 3c_1}}}$$

$$y(x) \rightarrow \frac{(1+i\sqrt{3})(a+x^2)}{2^{2/3} \sqrt[3]{\sqrt{4(a+x^2)^3 + 9c_1^2 + 3c_1}}} + \frac{i(\sqrt{3}+i) \sqrt[3]{\sqrt{4(a+x^2)^3 + 9c_1^2 + 3c_1}}}{2\sqrt[3]{2}}$$

$$y(x) \rightarrow \frac{(1-i\sqrt{3})(a+x^2)}{2^{2/3} \sqrt[3]{\sqrt{4(a+x^2)^3 + 9c_1^2 + 3c_1}}} - \frac{i(\sqrt{3}-i) \sqrt[3]{\sqrt{4(a+x^2)^3 + 9c_1^2 + 3c_1}}}{2\sqrt[3]{2}}$$

$$y(x) \rightarrow 0$$

**4.27 problem Recognizable Exact Differential equations.  
Integrating factors. Exercise 10.19, page 90**

Internal problem ID [4494]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 10

**Problem number:** Recognizable Exact Differential equations. Integrating factors. Exercise 10.19, page 90.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_exact, _rational]`

$$2yx + (a + x^2 + y^2) y' = -x^2 - b$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 810

```
dsolve((2*x*y(x)+x^2+b)+(y(x)^2+x^2+a)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\left(-4x^3 - 12xb - 12c_1 + 4\sqrt{5x^6 + 12ax^4 + 6bx^4 + 12a^2x^2 + 9b^2x^2 + 6c_1x^3 + 4a^3 + 18bc_1x + 9c_1^2}\right)^{\frac{1}{3}}}{2(x^2 + a)}$$

$$y(x) = \frac{\left(-4x^3 - 12xb - 12c_1 + 4\sqrt{5x^6 + 12ax^4 + 6bx^4 + 12a^2x^2 + 9b^2x^2 + 6c_1x^3 + 4a^3 + 18bc_1x + 9c_1^2}\right)^{\frac{1}{3}}}{x^2 + a} + \frac{\left(-4x^3 - 12xb - 12c_1 + 4\sqrt{5x^6 + 12ax^4 + 6bx^4 + 12a^2x^2 + 9b^2x^2 + 6c_1x^3 + 4a^3 + 18bc_1x + 9c_1^2}\right)^{\frac{1}{3}}}{2} + \frac{i\sqrt{3} \left(\frac{\left(-4x^3 - 12xb - 12c_1 + 4\sqrt{5x^6 + 12ax^4 + 6bx^4 + 12a^2x^2 + 9b^2x^2 + 6c_1x^3 + 4a^3 + 18bc_1x + 9c_1^2}\right)^{\frac{1}{3}}}{2}\right)}{\left(-4x^3 - 12xb - 12c_1 + 4\sqrt{5x^6 + 12ax^4 + 6bx^4 + 12a^2x^2 + 9b^2x^2 + 6c_1x^3 + 4a^3 + 18bc_1x + 9c_1^2}\right)^{\frac{1}{3}}}$$

$$y(x) = \frac{\left(-4x^3 - 12xb - 12c_1 + 4\sqrt{5x^6 + 12ax^4 + 6bx^4 + 12a^2x^2 + 9b^2x^2 + 6c_1x^3 + 4a^3 + 18bc_1x + 9c_1^2}\right)^{\frac{1}{3}}}{x^2 + a} + \frac{\left(-4x^3 - 12xb - 12c_1 + 4\sqrt{5x^6 + 12ax^4 + 6bx^4 + 12a^2x^2 + 9b^2x^2 + 6c_1x^3 + 4a^3 + 18bc_1x + 9c_1^2}\right)^{\frac{1}{3}}}{2} + \frac{i\sqrt{3} \left(\frac{\left(-4x^3 - 12xb - 12c_1 + 4\sqrt{5x^6 + 12ax^4 + 6bx^4 + 12a^2x^2 + 9b^2x^2 + 6c_1x^3 + 4a^3 + 18bc_1x + 9c_1^2}\right)^{\frac{1}{3}}}{2}\right)}{\left(-4x^3 - 12xb - 12c_1 + 4\sqrt{5x^6 + 12ax^4 + 6bx^4 + 12a^2x^2 + 9b^2x^2 + 6c_1x^3 + 4a^3 + 18bc_1x + 9c_1^2}\right)^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 6.558 (sec). Leaf size: 396

`DSolve[(2*x*y[x]+x^2+b)+(y[x]^2+x^2+a)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{\sqrt[3]{2} \left( \sqrt{4(a+x^2)^3 + (3bx+x^3-3c_1)^2} - 3bx - x^3 + 3c_1 \right)^{2/3} - 2a - 2x^2}{2^{2/3} \sqrt[3]{\sqrt{4(a+x^2)^3 + (3bx+x^3-3c_1)^2} - 3bx - x^3 + 3c_1}}$$

$$y(x) \rightarrow \frac{(1+i\sqrt{3})(a+x^2)}{2^{2/3} \sqrt[3]{\sqrt{4(a+x^2)^3 + (3bx+x^3-3c_1)^2} - 3bx - x^3 + 3c_1}} + \frac{i(\sqrt{3}+i) \sqrt[3]{\sqrt{4(a+x^2)^3 + (3bx+x^3-3c_1)^2} - 3bx - x^3 + 3c_1}}{2\sqrt[3]{2}}$$

$$y(x) \rightarrow \frac{(1-i\sqrt{3})(a+x^2)}{2^{2/3} \sqrt[3]{\sqrt{4(a+x^2)^3 + (3bx+x^3-3c_1)^2} - 3bx - x^3 + 3c_1}} - \frac{i(\sqrt{3}-i) \sqrt[3]{\sqrt{4(a+x^2)^3 + (3bx+x^3-3c_1)^2} - 3bx - x^3 + 3c_1}}{2\sqrt[3]{2}}$$

## 5 Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

5.1	problem Exercise 11.1, page 97 . . . . .	102
5.2	problem Exercise 11.2, page 97 . . . . .	103
5.3	problem Exercise 11.3, page 97 . . . . .	104
5.4	problem Exercise 11.4, page 97 . . . . .	105
5.5	problem Exercise 11.5, page 97 . . . . .	106
5.6	problem Exercise 11.6, page 97 . . . . .	107
5.7	problem Exercise 11.7, page 97 . . . . .	108
5.8	problem Exercise 11.8, page 97 . . . . .	109
5.9	problem Exercise 11.9, page 97 . . . . .	110
5.10	problem Exercise 11.11, page 97 . . . . .	111
5.11	problem Exercise 11.12, page 97 . . . . .	112
5.12	problem Exercise 11.11, page 97 . . . . .	113
5.13	problem Exercise 11.14, page 97 . . . . .	114
5.14	problem Exercise 11.15, page 97 . . . . .	115
5.15	problem Exercise 11.16, page 97 . . . . .	116
5.16	problem Exercise 11.17, page 97 . . . . .	117
5.17	problem Exercise 11.18, page 97 . . . . .	118
5.18	problem Exercise 11.19, page 97 . . . . .	119
5.19	problem Exercise 11.20, page 97 . . . . .	120
5.20	problem Exercise 11.21, page 97 . . . . .	121
5.21	problem Exercise 11.22, page 97 . . . . .	122
5.22	problem Exercise 11.23, page 97 . . . . .	123
5.23	problem Exercise 11.24, page 97 . . . . .	124
5.24	problem Exercise 11.26, page 97 . . . . .	125
5.25	problem Exercise 11.27, page 97 . . . . .	126
5.26	problem Exercise 11.28, page 97 . . . . .	127
5.27	problem Exercise 11.29, page 97 . . . . .	128

## 5.1 problem Exercise 11.1, page 97

Internal problem ID [4495]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.1, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$xy' + y = x^3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x*diff(y(x),x)+y(x)=x^3,y(x), singsol=all)
```

$$y(x) = \frac{\frac{x^4}{4} + c_1}{x}$$

### ✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 19

```
DSolve[x*y'[x]+y[x]==x^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{4} + \frac{c_1}{x}$$

## 5.2 problem Exercise 11.2, page 97

Internal problem ID [4496]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.2, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' + ay = b$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)+a*y(x)=b,y(x), singsol=all)
```

$$y(x) = \frac{b}{a} + e^{-ax}c_1$$

### ✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 29

```
DSolve[y'[x]+a*y[x]==b,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{b}{a} + c_1 e^{-ax}$$

$$y(x) \rightarrow \frac{b}{a}$$



### 5.3 problem Exercise 11.3, page 97

Internal problem ID [4497]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.3, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_Bernoulli]

$$xy' + y - y^2 \ln(x) = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(x*diff(y(x),x)+y(x)=y(x)^2*ln(x),y(x), singsol=all)
```

$$y(x) = \frac{1}{1 + c_1 x + \ln(x)}$$

#### ✓ Solution by Mathematica

Time used: 0.143 (sec). Leaf size: 20

```
DSolve[x*y'[x]+y[x]==y[x]^2*Log[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{\log(x) + c_1 x + 1}$$

$$y(x) \rightarrow 0$$

## 5.4 problem Exercise 11.4, page 97

Internal problem ID [4498]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.4, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$x' + 2yx = e^{-y^2}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(x(y),y)+2*y*x(y)=exp(-y^2),x(y), singsol=all)
```

$$x(y) = (y + c_1) e^{-y^2}$$

✓ Solution by Mathematica

Time used: 0.054 (sec). Leaf size: 17

```
DSolve[x'[y]+2*y*x[y]==Exp[-y^2],x[y],y,IncludeSingularSolutions -> True]
```

$$x(y) \rightarrow e^{-y^2}(y + c_1)$$

## 5.5 problem Exercise 11.5, page 97

Internal problem ID [4499]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.5, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$r' - (r + e^{-\theta}) \tan(\theta) = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(r(theta),theta)=(r(theta)+exp(-theta))*tan(theta),r(theta), singsol=all)
```

$$r(\theta) = \frac{c_1}{\cos(\theta)} - \frac{e^{-\theta}(\cos(\theta) + \sin(\theta))}{2 \cos(\theta)}$$

### ✓ Solution by Mathematica

Time used: 0.093 (sec). Leaf size: 24

```
DSolve[r' [\Theta]==(r [\Theta]+Exp[-\Theta])*Tan [\Theta],r [\Theta], \Theta, Include
```

$$r(\theta) \rightarrow -\frac{1}{2}e^{-\theta}(\tan(\theta) + 1) + c_1 \sec(\theta)$$

## 5.6 problem Exercise 11.6, page 97

Internal problem ID [4500]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.6, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y' - \frac{2xy}{x^2 + 1} = 1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)-(2*x*y(x))/(x^2+1)=1,y(x), singsol=all)
```

$$y(x) = (\arctan(x) + c_1)(x^2 + 1)$$

### ✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 16

```
DSolve[y'[x]-2*x*y[x]/(x^2+1)==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x^2 + 1)(\arctan(x) + c_1)$$

## 5.7 problem Exercise 11.7, page 97

Internal problem ID [4501]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.7, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [Bernoulli]

$$y' + y - y^3 x = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
dsolve(diff(y(x),x)+y(x)=x*y(x)^3,y(x), singsol=all)
```

$$y(x) = -\frac{2}{\sqrt{2 + 4c_1 e^{2x} + 4x}}$$

$$y(x) = \frac{2}{\sqrt{2 + 4c_1 e^{2x} + 4x}}$$

### ✓ Solution by Mathematica

Time used: 2.606 (sec). Leaf size: 50

```
DSolve[y'[x]+y[x]==x*y[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\sqrt{x + c_1 e^{2x} + \frac{1}{2}}}$$

$$y(x) \rightarrow \frac{1}{\sqrt{x + c_1 e^{2x} + \frac{1}{2}}}$$

$$y(x) \rightarrow 0$$

## 5.8 problem Exercise 11.8, page 97

Internal problem ID [4502]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.8, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_rational, \_Bernoulli]

$$(-x^3 + 1)y' - 2(x + 1)y - y^{\frac{5}{2}} = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 45

```
dsolve((1-x^3)*diff(y(x),x)-2*(1+x)*y(x)=y(x)^(5/2),y(x), singsol=all)
```

$$-\frac{c_1}{\frac{x^2}{(x-1)^2} + \frac{x}{(x-1)^2} + \frac{1}{(x-1)^2}} + \frac{1}{y(x)^{\frac{3}{2}}} + \frac{3}{4(x^2 + x + 1)} = 0$$

### ✓ Solution by Mathematica

Time used: 3.024 (sec). Leaf size: 41

```
DSolve[(1-x^3)*y'[x]-2*(1+x)*y[x]==y[x]^(5/2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2\sqrt[3]{2}}{\left(\frac{-3+4c_1(x-1)^2}{x^2+x+1}\right)^{2/3}}$$

$$y(x) \rightarrow 0$$

## 5.9 problem Exercise 11.9, page 97

Internal problem ID [4503]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.9, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$\tan(\theta) r' - r = \tan(\theta)^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(tan(theta)*diff(r(theta),theta)-r(theta)=tan(theta)^2,r(theta), singsol=all)
```

$$r(\theta) = (\ln(\sec(\theta) + \tan(\theta)) + c_1) \sin(\theta)$$

### ✓ Solution by Mathematica

Time used: 0.051 (sec). Leaf size: 14

```
DSolve[Tan[\[Theta]]*r'[\[Theta]]-r[\[Theta]]==Tan[\[Theta]]^2,r[\[Theta]],\[Theta],IncludeS
```

$$r(\theta) \rightarrow \sin(\theta) (\coth^{-1}(\sin(\theta)) + c_1)$$

## 5.10 problem Exercise 11.11, page 97

Internal problem ID [4504]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.11, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 2y = 3e^{-2x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)+2*y(x)=3*exp(-2*x),y(x), singsol=all)
```

$$y(x) = (3x + c_1)e^{-2x}$$

### ✓ Solution by Mathematica

Time used: 0.051 (sec). Leaf size: 17

```
DSolve[y'[x]+2*y[x]==3*Exp[-2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(3x + c_1)$$



## 5.11 problem Exercise 11.12, page 97

Internal problem ID [4505]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.12, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 2y = \frac{3e^{-2x}}{4}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)+2*y(x)=3/4*exp(-2*x),y(x), singsol=all)
```

$$y(x) = \left( \frac{3x}{4} + c_1 \right) e^{-2x}$$

### ✓ Solution by Mathematica

Time used: 0.05 (sec). Leaf size: 22

```
DSolve[y'[x]+2*y[x]==3/4*Exp[-2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}e^{-2x}(3x + 4c_1)$$

## 5.12 problem Exercise 11.11, page 97

Internal problem ID [4506]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.11, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 2y = \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)+2*y(x)=sin(x),y(x), singsol=all)
```

$$y(x) = -\frac{\cos(x)}{5} + \frac{2 \sin(x)}{5} + c_1 e^{-2x}$$

### ✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 26

```
DSolve[y'[x]+2*y[x]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2 \sin(x)}{5} - \frac{\cos(x)}{5} + c_1 e^{-2x}$$

### 5.13 problem Exercise 11.14, page 97

Internal problem ID [4507]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.14, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y' + y \cos(x) = e^{2x}$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)+y(x)*cos(x)=exp(2*x),y(x), singsol=all)
```

$$y(x) = \left( \int e^{2x+\sin(x)} dx + c_1 \right) e^{-\sin(x)}$$

#### ✓ Solution by Mathematica

Time used: 0.735 (sec). Leaf size: 32

```
DSolve[y'[x]+y[x]*Cos[x]==Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\sin(x)} \left( \int_1^x e^{2K[1]+\sin(K[1])} dK[1] + c_1 \right)$$

## 5.14 problem Exercise 11.15, page 97

Internal problem ID [4508]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.15, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y' + y \cos(x) = \frac{\sin(2x)}{2}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)+y(x)*cos(x)=1/2*sin(2*x),y(x), singsol=all)
```

$$y(x) = \sin(x) - 1 + e^{-\sin(x)} c_1$$

### ✓ Solution by Mathematica

Time used: 0.05 (sec). Leaf size: 18

```
DSolve[y'[x]+y[x]*Cos[x]==1/2*Sin[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sin(x) + c_1 e^{-\sin(x)} - 1$$

## 5.15 problem Exercise 11.16, page 97

Internal problem ID [4509]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.16, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$xy' + y = x \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x*diff(y(x),x)+y(x)=x*sin(x),y(x), singsol=all)
```

$$y(x) = \frac{-x \cos(x) + \sin(x) + c_1}{x}$$

### ✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 19

```
DSolve[x*y'[x]+y[x]==x*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sin(x) - x \cos(x) + c_1}{x}$$

## 5.16 problem Exercise 11.17, page 97

Internal problem ID [4510]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.17, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$xy' - y = \sin(x)x^2$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve(x*diff(y(x),x)-y(x)=x^2*sin(x),y(x), singsol=all)
```

$$y(x) = (-\cos(x) + c_1)x$$

### ✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 14

```
DSolve[x*y'[x]-y[x]==x^2*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(-\cos(x) + c_1)$$

## 5.17 problem Exercise 11.18, page 97

Internal problem ID [4511]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.18, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class D', _rational, _Bernoulli]`

$$xy' + y^2x - y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(x*diff(y(x),x)+x*y(x)^2-y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{2x}{x^2 + 2c_1}$$

### ✓ Solution by Mathematica

Time used: 0.149 (sec). Leaf size: 23

```
DSolve[x*y'[x]+x*y[x]^2-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2x}{x^2 + 2c_1}$$

$$y(x) \rightarrow 0$$

## 5.18 problem Exercise 11.19, page 97

Internal problem ID [4512]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.19, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_Bernoulli]

$$xy' - y(2 \ln(x)y - 1) = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x*diff(y(x),x)-y(x)*(2*y(x)*ln(x)-1)=0,y(x), singsol=all)
```

$$y(x) = \frac{1}{2 + c_1x + 2 \ln(x)}$$

### ✓ Solution by Mathematica

Time used: 0.14 (sec). Leaf size: 22

```
DSolve[x*y'[x]-y[x]*(2*y[x]*Log[x]-1)==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2 \log(x) + c_1x + 2}$$

$$y(x) \rightarrow 0$$



## 5.19 problem Exercise 11.20, page 97

Internal problem ID [4513]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.20, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class D'], _rational, _Bernoulli]`

$$x^2(-1+x)y' - y^2 - x(x-2)y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(x^2*(x-1)*diff(y(x),x)-y(x)^2-x*(x-2)*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x^2}{c_1x - c_1 + 1}$$

### ✓ Solution by Mathematica

Time used: 0.191 (sec). Leaf size: 25

```
DSolve[x^2*(x-1)*y'[x]-y[x]^2-x*(x-2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2}{c_1(-x) + 1 + c_1}$$

$$y(x) \rightarrow 0$$

## 5.20 problem Exercise 11.21, page 97

Internal problem ID [4514]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.21, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' - y = e^x$$

With initial conditions

$$[y(0) = 1]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 10

```
dsolve([diff(y(x),x)-y(x)=exp(x),y(0) = 1],y(x), singsol=all)
```

$$y(x) = e^x(x + 1)$$

### ✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 12

```
DSolve[{y'[x]-y[x]==Exp[x],{y[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(x + 1)$$

## 5.21 problem Exercise 11.22, page 97

Internal problem ID [4515]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.22, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y' + \frac{y}{x} - \frac{y^2}{x} = 0$$

With initial conditions

$$[y(-1) = 1]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve([diff(y(x),x)+y(x)/x=y(x)^2/x,y(-1) = 1],y(x), singsol=all)
```

$$y(x) = 1$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 6

```
DSolve[{y'[x]+y[x]/x==y[x]^2/x,{y[-1]==1}],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 1$$

## 5.22 problem Exercise 11.23, page 97

Internal problem ID [4516]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.23, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_Bernoulli]

$$2 \cos(x) y' - \sin(x) y + y^3 = 0$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.36 (sec). Leaf size: 33

```
dsolve([2*cos(x)*diff(y(x),x)=y(x)*sin(x)-y(x)^3,y(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{(2 \cos(x)^2 - 1) (\cos(x) - \sin(x))}}{2 \cos(x)^2 - 1}$$

✓ Solution by Mathematica

Time used: 0.369 (sec). Leaf size: 14

```
DSolve[{2*Cos[x]*y'[x]==y[x]*Sin[x]-y[x]^3,{y[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{\sqrt{\sin(x) + \cos(x)}}$$

## 5.23 problem Exercise 11.24, page 97

Internal problem ID [4517]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.24, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x)*G(y),0]']]`

$$(x - \cos(y))y' + \tan(y) = 0$$

With initial conditions

$$y(1) = \frac{\pi}{6}$$

✓ Solution by Maple

Time used: 1.235 (sec). Leaf size: 29

```
dsolve([(x-cos(y(x)))*diff(y(x),x)+tan(y(x))=0,y(1) = 1/6*Pi],y(x), singsol=all)
```

$$y(x) = \text{RootOf}\left(24x \sin(\_Z) + 3\sqrt{3} - 6 \sin(2\_Z) + 2\pi - 12\_Z - 12\right)$$

✓ Solution by Mathematica

Time used: 0.216 (sec). Leaf size: 45

```
DSolve[{(x-Cos[y[x]])*y'[x]+Tan[y[x]]==0,{y[1]==Pi/6}},y[x],x,IncludeSingularSolutions -> Tr
```

$$\text{Solve}\left[x = \frac{1}{24}\left(12 - 3\sqrt{3} - 2\pi\right) \csc(y(x)) + \left(\frac{y(x)}{2} + \frac{1}{4} \sin(2y(x))\right) \csc(y(x)), y(x)\right]$$

## 5.24 problem Exercise 11.26, page 97

Internal problem ID [4518]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.26, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_rational, _Riccati]`

$$y' - \frac{2y}{x} + \frac{y^2}{x} = x^3$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)=x^3+2/x*y(x)-1/x*y(x)^2,y(x), singsol=all)
```

$$y(x) = i \tan\left(-\frac{ix^2}{2} + c_1\right) x^2$$

### ✓ Solution by Mathematica

Time used: 0.162 (sec). Leaf size: 75

```
DSolve[y'[x]==x^3+2/x*y[x]-1/x*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2 \left( i \cosh\left(\frac{x^2}{2}\right) + c_1 \sinh\left(\frac{x^2}{2}\right) \right)}{i \sinh\left(\frac{x^2}{2}\right) + c_1 \cosh\left(\frac{x^2}{2}\right)}$$

$$y(x) \rightarrow x^2 \tanh\left(\frac{x^2}{2}\right)$$

## 5.25 problem Exercise 11.27, page 97

Internal problem ID [4519]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.27, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [Riccati]

$$y' + y^2 \sin(x) = 2 \tan(x) \sec(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 41

```
dsolve(diff(y(x),x)=2*tan(x)*sec(x)-y(x)^2*sin(x),y(x), singsol=all)
```

$$y(x) = \frac{\sec(x) \tan(x)}{\sin(x) (c_1 \cos(x)^2 + \sec(x))} - \frac{2c_1 \cos(x)}{c_1 \cos(x)^2 + \sec(x)}$$

### ✓ Solution by Mathematica

Time used: 0.88 (sec). Leaf size: 32

```
DSolve[y'[x]==2*Tan[x]*Sec[x]-y[x]^2*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sec(x) (-2 \cos^3(x) + c_1)}{\cos^3(x) + c_1}$$

$$y(x) \rightarrow \sec(x)$$

## 5.26 problem Exercise 11.28, page 97

Internal problem ID [4520]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.28, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, _Riccati]`

$$y' + \frac{y}{x} + y^2 = \frac{1}{x^2}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)=1/x^2-y(x)/x-y(x)^2,y(x), singsol=all)
```

$$y(x) = -\frac{\tanh(-\ln(x) + c_1)}{x}$$

### ✓ Solution by Mathematica

Time used: 1.192 (sec). Leaf size: 62

```
DSolve[y'[x]==1/x^2-y[x]/x-y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{i \tan(c_1 - i \log(x))}{x}$$
$$y(x) \rightarrow -\frac{-x^2 + e^{2i \text{Interval}\{0,\pi\}}}{x^3 + x e^{2i \text{Interval}\{0,\pi\}}}$$



## 5.27 problem Exercise 11.29, page 97

Internal problem ID [4521]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 11, Bernoulli Equations

**Problem number:** Exercise 11.29, page 97.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _Riccati]`

$$y' - \frac{y}{x} + \frac{y^2}{x^2} = 1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(diff(y(x),x)=1+y(x)/x-y(x)^2/x^2,y(x), singsol=all)
```

$$y(x) = \tanh(\ln(x) + c_1) x$$

### ✓ Solution by Mathematica

Time used: 0.539 (sec). Leaf size: 43

```
DSolve[y'[x]==1+y[x]/x-y[x]^2/x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x(x^2 - e^{2c_1})}{x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -x$$

$$y(x) \rightarrow x$$

## 6 Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

6.1	problem Exercise 12.1, page 103 . . . . .	131
6.2	problem Exercise 12.2, page 103 . . . . .	132
6.3	problem Exercise 12.3, page 103 . . . . .	133
6.4	problem Exercise 12.4, page 103 . . . . .	135
6.5	problem Exercise 12.5, page 103 . . . . .	136
6.6	problem Exercise 12.6, page 103 . . . . .	137
6.7	problem Exercise 12.7, page 103 . . . . .	138
6.8	problem Exercise 12.8, page 103 . . . . .	139
6.9	problem Exercise 12.9, page 103 . . . . .	140
6.10	problem Exercise 12.10, page 103 . . . . .	141
6.11	problem Exercise 12.11, page 103 . . . . .	142
6.12	problem Exercise 12.12, page 103 . . . . .	143
6.13	problem Exercise 12.13, page 103 . . . . .	144
6.14	problem Exercise 12.14, page 103 . . . . .	145
6.15	problem Exercise 12.15, page 103 . . . . .	146
6.16	problem Exercise 12.16, page 103 . . . . .	147
6.17	problem Exercise 12.17, page 103 . . . . .	148
6.18	problem Exercise 12.18, page 103 . . . . .	149
6.19	problem Exercise 12.19, page 103 . . . . .	150
6.20	problem Exercise 12.20, page 103 . . . . .	151
6.21	problem Exercise 12.21, page 103 . . . . .	152
6.22	problem Exercise 12.22, page 103 . . . . .	153
6.23	problem Exercise 12.23, page 103 . . . . .	154
6.24	problem Exercise 12.24, page 103 . . . . .	155
6.25	problem Exercise 12.25, page 103 . . . . .	156
6.26	problem Exercise 12.26, page 103 . . . . .	157
6.27	problem Exercise 12.27, page 103 . . . . .	158
6.28	problem Exercise 12.28, page 103 . . . . .	159
6.29	problem Exercise 12.29, page 103 . . . . .	160
6.30	problem Exercise 12.30, page 103 . . . . .	162
6.31	problem Exercise 12.31, page 103 . . . . .	163
6.32	problem Exercise 12.32, page 103 . . . . .	164
6.33	problem Exercise 12.33, page 103 . . . . .	165
6.34	problem Exercise 12.34, page 103 . . . . .	166

6.35	problem Exercise 12.35, page 103 . . . . .	167
6.36	problem Exercise 12.36, page 103 . . . . .	168
6.37	problem Exercise 12.37, page 103 . . . . .	171
6.38	problem Exercise 12.38, page 103 . . . . .	172
6.39	problem Exercise 12.39, page 103 . . . . .	173
6.40	problem Exercise 12.40, page 103 . . . . .	174
6.41	problem Exercise 12.41, page 103 . . . . .	176
6.42	problem Exercise 12.42, page 103 . . . . .	177
6.43	problem Exercise 12.43, page 103 . . . . .	178
6.44	problem Exercise 12.44, page 103 . . . . .	181
6.45	problem Exercise 12.45, page 103 . . . . .	182
6.46	problem Exercise 12.46, page 103 . . . . .	185
6.47	problem Exercise 12.47, page 103 . . . . .	187
6.48	problem Exercise 12.48, page 103 . . . . .	190
6.49	problem Exercise 12.49, page 103 . . . . .	191
6.50	problem Exercise 12.50, page 103 . . . . .	193

## 6.1 problem Exercise 12.1, page 103

Internal problem ID [4522]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.1, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_Bernoulli]

$$2yy'x + (x + 1)y^2 = e^x$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 59

```
dsolve(2*x*y(x)*diff(y(x),x)+(1+x)*y(x)^2=exp(x),y(x), singsol=all)
```

$$y(x) = -\frac{e^{-x}\sqrt{2}\sqrt{e^x x(e^{2x} + 2c_1)}}{2x}$$

$$y(x) = \frac{e^{-x}\sqrt{2}\sqrt{e^x x(e^{2x} + 2c_1)}}{2x}$$

### ✓ Solution by Mathematica

Time used: 7.324 (sec). Leaf size: 66

```
DSolve[2*x*y[x]*y'[x]+(1+x)*y[x]^2==Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{e^x + 2c_1 e^{-x}}}{\sqrt{2}\sqrt{x}}$$

$$y(x) \rightarrow \frac{\sqrt{e^x + 2c_1 e^{-x}}}{\sqrt{2}\sqrt{x}}$$

## 6.2 problem Exercise 12.2, page 103

Internal problem ID [4523]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.2, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type ['y=\_G(x,y)']

$$\cos(y) y' + \sin(y) = x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(cos(y(x))*diff(y(x),x)+sin(y(x))=x^2,y(x), singsol=all)
```

$$y(x) = \arcsin\left(\left(e^x x^2 - 2e^x x + 2e^x - c_1\right) e^{-x}\right)$$

### ✓ Solution by Mathematica

Time used: 14.047 (sec). Leaf size: 23

```
DSolve[Cos[y[x]]*y'[x]+Sin[y[x]]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arcsin\left(x^2 - 2x - 2c_1 e^{-x} + 2\right)$$

### 6.3 problem Exercise 12.3, page 103

Internal problem ID [4524]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.3, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$(x + 1)y' - y - (x + 1)\sqrt{1 + y} = 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 160

```
dsolve((x+1)*diff(y(x),x)-(y(x)+1)=(x+1)*sqrt(y(x)+1),y(x), singsol=all)
```

$$\begin{aligned} & \frac{\sqrt{y(x)+1}x}{(-x^2-2x+y(x))(\sqrt{y(x)+1}-1-x)} \\ & + \frac{2x}{(-x^2-2x+y(x))(\sqrt{y(x)+1}-1-x)} \\ & + \frac{x^2}{(-x^2-2x+y(x))(\sqrt{y(x)+1}-1-x)} \\ & + \frac{\sqrt{y(x)+1}}{(-x^2-2x+y(x))(\sqrt{y(x)+1}-1-x)} \\ & + \frac{1}{(-x^2-2x+y(x))(\sqrt{y(x)+1}-1-x)} - c_1 = 0 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.244 (sec). Leaf size: 60

```
DSolve[(x+1)*y'[x]-(y[x]+1)==(x+1)*Sqrt[y[x]+1],y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ \frac{2\sqrt{y(x)+1} \arctan\left(\frac{x+1}{\sqrt{-y(x)-1}}\right)}{\sqrt{-y(x)-1}} + \log(y(x) - (x+1)^2 + 1) - \log(x+1) = c_1, y(x) \right]$$

## 6.4 problem Exercise 12.4, page 103

Internal problem ID [4525]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.4, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$e^y(y' + 1) = e^x$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 16

```
dsolve(exp(y(x))*(diff(y(x),x)+1)=exp(x),y(x), singsol=all)
```

$$y(x) = x + \ln\left(\frac{c_1 e^{-2x}}{2} + \frac{1}{2}\right)$$

### ✓ Solution by Mathematica

Time used: 1.32 (sec). Leaf size: 22

```
DSolve[Exp[y[x]]*(y'[x]+1)==Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x + \log\left(\frac{e^{2x}}{2} + c_1\right)$$



## 6.5 problem Exercise 12.5, page 103

Internal problem ID [4526]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.5, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y' \sin(y) + \sin(x) \cos(y) = \sin(x)$$

### ✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)*sin(y(x))+sin(x)*cos(y(x))=sin(x),y(x), singsol=all)
```

$$y(x) = \arccos(e^{-\cos(x)}c_1 + 1)$$

### ✓ Solution by Mathematica

Time used: 0.792 (sec). Leaf size: 81

```
DSolve[y'[x]*Sin[y[x]]+Sin[x]*Cos[y[x]]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

$$\text{Solve} \left[ 2 \cos(x) \tan\left(\frac{y(x)}{2}\right) e^{\arctanh(\cos(y(x)))} - \sqrt{\sin^2(y(x))} \csc\left(\frac{y(x)}{2}\right) \sec\left(\frac{y(x)}{2}\right) \left(\log\left(\sec^2\left(\frac{y(x)}{2}\right)\right)\right) - 2 \log\left(\tan\left(\frac{y(x)}{2}\right)\right) = c_1, y(x) \right]$$

$$y(x) \rightarrow 0$$

## 6.6 problem Exercise 12.6, page 103

Internal problem ID [4527]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.6, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$(-y + x)^2 y' = 4$$

### ✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 27

```
dsolve((x-y(x))^2*diff(y(x),x)=4,y(x), singsol=all)
```

$$y(x) - \ln(y(x) - x + 2) + \ln(y(x) - x - 2) - c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 0.202 (sec). Leaf size: 36

```
DSolve[(x-y[x])^2*y'[x]==4,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ y(x) - 4 \left( \frac{1}{4} \log(y(x) - x + 2) - \frac{1}{4} \log(-y(x) + x + 2) \right) = c_1, y(x) \right]$$

## 6.7 problem Exercise 12.7, page 103

Internal problem ID [4528]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.7, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _dAlembert]`

$$xy' - y - \sqrt{x^2 + y^2} = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve(x*diff(y(x),x)-y(x)=sqrt(x^2+y(x)^2),y(x), singsol=all)
```

$$\frac{y(x)}{x^2} + \frac{\sqrt{x^2 + y(x)^2}}{x^2} - c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 0.337 (sec). Leaf size: 27

```
DSolve[x*y'[x]-y[x]==Sqrt[x^2+y[x]^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}e^{-c_1}(-1 + e^{2c_1}x^2)$$

## 6.8 problem Exercise 12.8, page 103

Internal problem ID [4529]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.8, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _exact, _rational, [_Abel, '2nd ty`

$$(3x + 2y + 1)y' + 3y = -4x - 2$$

### ✓ Solution by Maple

Time used: 0.312 (sec). Leaf size: 33

```
dsolve((3*x+2*y(x)+1)*diff(y(x),x)+(4*x+3*y(x)+2)=0,y(x), singsol=all)
```

$$y(x) = -2 - \frac{\frac{3c_1(x-1)}{2} + \frac{\sqrt{(x-1)^2 c_1^2 + 4}}{2}}{c_1}$$

### ✓ Solution by Mathematica

Time used: 0.123 (sec). Leaf size: 61

```
DSolve[(3*x+2*y[x]+1)*y'[x]+(4*x+3*y[x]+2)==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left( -\sqrt{x^2 - 2x + 1 + 4c_1} - 3x - 1 \right)$$

$$y(x) \rightarrow \frac{1}{2} \left( \sqrt{x^2 - 2x + 1 + 4c_1} - 3x - 1 \right)$$

## 6.9 problem Exercise 12.9, page 103

Internal problem ID [4530]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.9, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _dAlembert]`

$$(x^2 - y^2) y' - 2yx = 0$$

### ✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 45

```
dsolve((x^2-y(x)^2)*diff(y(x),x)=2*x*y(x),y(x), singsol=all)
```

$$y(x) = -\frac{-1 + \sqrt{-4c_1^2x^2 + 1}}{2c_1}$$

$$y(x) = \frac{1 + \sqrt{-4c_1^2x^2 + 1}}{2c_1}$$

### ✓ Solution by Mathematica

Time used: 0.982 (sec). Leaf size: 66

```
DSolve[(x^2-y[x]^2)*y'[x]==2*x*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left( e^{c_1} - \sqrt{-4x^2 + e^{2c_1}} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left( \sqrt{-4x^2 + e^{2c_1}} + e^{c_1} \right)$$

$$y(x) \rightarrow 0$$

## 6.10 problem Exercise 12.10, page 103

Internal problem ID [4531]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.10, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$y + (1 + e^{2x}y^2)y' = 0$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 18

```
dsolve(y(x)+(1+y(x)^2*exp(2*x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{e^{-x}}{\sqrt{\text{LambertW}(c_1 e^{-2x})}}$$

### ✓ Solution by Mathematica

Time used: 3.33 (sec). Leaf size: 57

```
DSolve[y[x]+(1+y[x]^2*Exp[2*x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{e^{-x}}{\sqrt{W(e^{-2x+2c_1})}}$$

$$y(x) \rightarrow \frac{e^{-x}}{\sqrt{W(e^{-2x+2c_1})}}$$

$$y(x) \rightarrow 0$$

## 6.11 problem Exercise 12.11, page 103

Internal problem ID [4532]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.11, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, _Bernoulli]`

$$x^2y + y^2 + y'x^3 = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve((x^2*y(x)+y(x)^2)+x^3*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{3x^2}{3c_1x^3 - 1}$$

### ✓ Solution by Mathematica

Time used: 0.143 (sec). Leaf size: 26

```
DSolve[(x^2*y[x]+y[x]^2)+x^3*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3x^2}{-1 + 3c_1x^3}$$

$$y(x) \rightarrow 0$$

## 6.12 problem Exercise 12.12, page 103

Internal problem ID [4533]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.12, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_exact]

$$y^2 e^{y^2 x} + (2xy e^{y^2 x} - 3y^2) y' = -4x^3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve((y(x)^2*exp(x*y(x)^2)+4*x^3)+(2*x*y(x)*exp(x*y(x)^2)-3*y(x)^2)*diff(y(x),x)=0,y(x), s
```

$$e^{y(x)^2 x} + x^4 - y(x)^3 + c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 0.279 (sec). Leaf size: 24

```
DSolve[(y[x]^2*Exp[x*y[x]^2]+4*x^3)+(2*x*y[x]*Exp[x*y[x]^2]-3*y[x]^2)*y'[x]==0,y[x],x,Includ
```

$$\text{Solve}\left[x^4 + e^{xy(x)^2} - y(x)^3 = c_1, y(x)\right]$$



## 6.13 problem Exercise 12.13, page 103

Internal problem ID [4534]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.13, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries]]`

$$y' - (x^2 + 2y - 1)^{\frac{2}{3}} = -x$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)=(x^2+2*y(x)-1)^(2/3)-x,y(x), singsol=all)
```

$$x - \frac{3(x^2 + 2y(x) - 1)^{\frac{1}{3}}}{2} - c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 0.214 (sec). Leaf size: 40

```
DSolve[y'[x]==(x^2+2*y[x]-1)^(2/3)-x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{54}(8x^3 - 3(9 + 8c_1)x^2 + 24c_1^2x + 27 - 8c_1^3)$$

## 6.14 problem Exercise 12.14, page 103

Internal problem ID [4535]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.14, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_Bernoulli]

$$xy' + y - x^2(e^x + 1)y^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(x*diff(y(x),x)+y(x)=x^2*(1+exp(x))*y(x)^2,y(x), singsol=all)
```

$$y(x) = -\frac{1}{(x + e^x - c_1)x}$$

✓ Solution by Mathematica

Time used: 0.249 (sec). Leaf size: 55

```
DSolve[x*y'[x]+y[x]==x^2*(1+exp[x])*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{-x \int_1^x (\exp(K[1]) + 1) dK[1] + c_1 x}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -\frac{1}{x \int_1^x (\exp(K[1]) + 1) dK[1]}$$

## 6.15 problem Exercise 12.15, page 103

Internal problem ID [4536]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.15, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$2y - xy \ln(x) - 2x \ln(x) y' = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve((2*y(x)-x*y(x)*ln(x))-2*x*ln(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{x}{2}} \ln(x)$$

### ✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 22

```
DSolve[(2*y[x]-x*y[x]*Log[x])-2*x*Log[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-x/2} \log(x)$$

$$y(x) \rightarrow 0$$

## 6.16 problem Exercise 12.16, page 103

Internal problem ID [4537]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.16, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + ay = k e^{bx}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x)+a*y(x)=k*exp(b*x),y(x), singsol=all)
```

$$y(x) = \left( \frac{k e^{x(a+b)}}{a+b} + c_1 \right) e^{-ax}$$

### ✓ Solution by Mathematica

Time used: 0.072 (sec). Leaf size: 33

```
DSolve[y'[x]+a*y[x]==k*Exp[b*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{-ax} (k e^{x(a+b)} + c_1 (a+b))}{a+b}$$

## 6.17 problem Exercise 12.17, page 103

Internal problem ID [4538]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.17, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _Riccati]`

$$y' - (x + y)^2 = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)=(x+y(x))^2,y(x), singsol=all)
```

$$y(x) = -x - \tan(c_1 - x)$$

### ✓ Solution by Mathematica

Time used: 0.472 (sec). Leaf size: 14

```
DSolve[y'[x]==(x+y[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x + \tan(x + c_1)$$

## 6.18 problem Exercise 12.18, page 103

Internal problem ID [4539]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.18, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_Bernoulli]

$$y' + 8y^3x^3 + 2yx = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 43

```
dsolve(diff(y(x),x)+8*x^3*y(x)^3+2*x*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{1}{\sqrt{e^{2x^2}c_1 - 4x^2 - 2}}$$
$$y(x) = -\frac{1}{\sqrt{e^{2x^2}c_1 - 4x^2 - 2}}$$

### ✓ Solution by Mathematica

Time used: 7.034 (sec). Leaf size: 58

```
DSolve[y'[x]+8*x^3*y[x]^3+2*x*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\sqrt{-4x^2 + c_1e^{2x^2} - 2}}$$
$$y(x) \rightarrow \frac{1}{\sqrt{-4x^2 + c_1e^{2x^2} - 2}}$$
$$y(x) \rightarrow 0$$

## 6.19 problem Exercise 12.19, page 103

Internal problem ID [4540]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.19, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [NONE]

$$(xy\sqrt{x^2 - y^2} + x)y' - y + x^2\sqrt{x^2 - y^2} = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 34

```
dsolve((x*y(x)*sqrt(x^2-y(x)^2)+x)*diff(y(x),x)=y(x)-x^2*sqrt(x^2-y(x)^2),y(x), singsol=all)
```

$$\frac{y(x)^2}{2} + \arctan\left(\frac{y(x)}{\sqrt{x^2 - y(x)^2}}\right) + \frac{x^2}{2} - c_1 = 0$$

✓ Solution by Mathematica

Time used: 1.772 (sec). Leaf size: 44

```
DSolve[(x*y[x]*Sqrt[x^2-y[x]^2]+x)*y'[x]==y[x]-x^2*Sqrt[x^2-y[x]^2],y[x],x,IncludeSingularSo
```

$$\text{Solve}\left[-\arctan\left(\frac{\sqrt{x^2 - y(x)^2}}{y(x)}\right) + \frac{x^2}{2} + \frac{y(x)^2}{2} = c_1, y(x)\right]$$

## 6.20 problem Exercise 12.20, page 103

Internal problem ID [4541]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.20, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + ay = b \sin(kx)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 38

```
dsolve(diff(y(x),x)+a*y(x)=b*sin(k*x),y(x), singsol=all)
```

$$y(x) = e^{-ax} c_1 - \frac{b(k \cos(kx) - \sin(kx) a)}{a^2 + k^2}$$

### ✓ Solution by Mathematica

Time used: 0.083 (sec). Leaf size: 40

```
DSolve[y'[x]+a*y[x]==b*Sin[k*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{b(a \sin(kx) - k \cos(kx))}{a^2 + k^2} + c_1 e^{-ax}$$



## 6.21 problem Exercise 12.21, page 103

Internal problem ID [4542]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.21, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$xy' - y^2 = -1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(x*diff(y(x),x)-y(x)^2+1=0,y(x), singsol=all)
```

$$y(x) = -\tanh(\ln(x) + c_1)$$

### ✓ Solution by Mathematica

Time used: 0.486 (sec). Leaf size: 43

```
DSolve[x*y'[x]-y[x]^2+1==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1 - e^{2c_1} x^2}{1 + e^{2c_1} x^2}$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 1$$

## 6.22 problem Exercise 12.22, page 103

Internal problem ID [4543]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.22, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x)*G(y),0]']]`

$$(y^2 + \sin(x) a) y' = \cos(x)$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 41

```
dsolve((y(x)^2+a*sin(x))*diff(y(x),x)=cos(x),y(x), singsol=all)
```

$$-e^{-ay(x)} \sin(x) - \frac{(a^2 y(x)^2 + 2ay(x) + 2) e^{-ay(x)}}{a^3} + c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 0.194 (sec). Leaf size: 45

```
DSolve[(y[x]^2+a*Sin[x])*y'[x]==Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[\sin(x) (-e^{-ay(x)}) - \frac{e^{-ay(x)}(a^2 y(x)^2 + 2ay(x) + 2)}{a^3} = c_1, y(x)\right]$$

## 6.23 problem Exercise 12.23, page 103

Internal problem ID [4544]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.23, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _dAlembert]`

$$xy' - x e^{\frac{y}{x}} - y = x$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(x*diff(y(x),x)=x*exp(y(x)/x)+x+y(x),y(x), singsol=all)
```

$$y(x) = \left( \ln \left( -\frac{x}{x e^{c_1} - 1} \right) + c_1 \right) x$$

### ✓ Solution by Mathematica

Time used: 4.512 (sec). Leaf size: 38

```
DSolve[x*y'[x]==x*Exp[y[x]/x]+x+y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \log \left( \frac{1}{2} \left( -1 + \tanh \left( \frac{1}{2} (-\log(x) - c_1) \right) \right) \right)$$

$$y(x) \rightarrow i\pi x$$

## 6.24 problem Exercise 12.24, page 103

Internal problem ID [4545]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.24, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y' + y \cos(x) = e^{-\sin(x)}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x)+y(x)*cos(x)=exp(-sin(x)),y(x), singsol=all)
```

$$y(x) = (x + c_1) e^{-\sin(x)}$$

### ✓ Solution by Mathematica

Time used: 0.123 (sec). Leaf size: 16

```
DSolve[y'[x]+y[x]*Cos[x]==Exp[-Sin[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x + c_1) e^{-\sin(x)}$$

## 6.25 problem Exercise 12.25, page 103

Internal problem ID [4546]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.25, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class G']]`

$$xy' - y(\ln(yx) - 1) = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve(x*diff(y(x),x)-y(x)*(ln(x*y(x))-1)=0,y(x), singsol=all)
```

$$y(x) = \frac{e^{\frac{x}{c_1}}}{x}$$

### ✓ Solution by Mathematica

Time used: 0.186 (sec). Leaf size: 24

```
DSolve[x*y'[x]-y[x]*(Log[x*y[x]]-1)==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{e^{c_1}x}}{x}$$

$$y(x) \rightarrow \frac{1}{x}$$

## 6.26 problem Exercise 12.26, page 103

Internal problem ID [4547]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.26, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class D', _rational, _Bernoulli]`

$$x^3y' - y^2 - x^2y = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 15

```
dsolve(x^3*diff(y(x),x)-y(x)^2-x^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x^2}{c_1x + 1}$$

### ✓ Solution by Mathematica

Time used: 0.129 (sec). Leaf size: 22

```
DSolve[x^3*y'[x]-y[x]^2-x^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2}{1 + c_1x}$$

$$y(x) \rightarrow 0$$

## 6.27 problem Exercise 12.27, page 103

Internal problem ID [4548]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.27, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$xy' + ay = -bx^n$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(x*diff(y(x),x)+a*y(x)+b*x^n=0,y(x), singsol=all)
```

$$y(x) = -\frac{bx^n}{a+n} + x^{-a}c_1$$

### ✓ Solution by Mathematica

Time used: 0.056 (sec). Leaf size: 25

```
DSolve[x*y'[x]+a*y[x]+b*x^n==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{bx^n}{a+n} + c_1x^{-a}$$

## 6.28 problem Exercise 12.28, page 103

Internal problem ID [4549]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.28, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _dAlembert]`

$$xy' - x \sin\left(\frac{y}{x}\right) - y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```
dsolve(x*diff(y(x),x)-x*sin(y(x)/x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = \arctan\left(\frac{2xc_1}{c_1^2x^2 + 1}, -\frac{c_1^2x^2 - 1}{c_1^2x^2 + 1}\right) x$$

### ✓ Solution by Mathematica

Time used: 0.321 (sec). Leaf size: 52

```
DSolve[x*y'[x]-x*Sin[y[x]/x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x \arccos(-\tanh(\log(x) + c_1))$$

$$y(x) \rightarrow x \arccos(-\tanh(\log(x) + c_1))$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -\pi x$$

$$y(x) \rightarrow \pi x$$



## 6.29 problem Exercise 12.29, page 103

Internal problem ID [4550]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.29, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$y^2 - 3yx + (yx - x^2) y' = 2x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 59

```
dsolve((x*y(x)-x^2)*diff(y(x),x)+y(x)^2-3*x*y(x)-2*x^2=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 x^2 - \sqrt{2c_1^2 x^4 + 1}}{c_1 x}$$

$$y(x) = \frac{c_1 x^2 + \sqrt{2c_1^2 x^4 + 1}}{c_1 x}$$

✓ Solution by Mathematica

Time used: 0.625 (sec). Leaf size: 99

```
DSolve[(x*y[x]-x^2)*y'[x]+y[x]^2-3*x*y[x]-2*x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x - \frac{\sqrt{2x^4 + e^{2c_1}}}{x}$$

$$y(x) \rightarrow x + \frac{\sqrt{2x^4 + e^{2c_1}}}{x}$$

$$y(x) \rightarrow x - \frac{\sqrt{2}\sqrt{x^4}}{x}$$

$$y(x) \rightarrow \frac{\sqrt{2}\sqrt{x^4}}{x} + x$$

## 6.30 problem Exercise 12.30, page 103

Internal problem ID [4551]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.30, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_exact, _rational, [_Abel, '2nd type', 'class B']]`

$$(3 + 6yx + x^2)y' + 3y^2 + 2yx = -2x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 71

```
dsolve((6*x*y(x)+x^2+3)*diff(y(x),x)+3*y(x)^2+2*x*y(x)+2*x=0,y(x), singsol=all)
```

$$y(x) = \frac{-x^2 - 3 + \sqrt{x^4 - 12x^3 - 12c_1x + 6x^2 + 9}}{6x}$$

$$y(x) = -\frac{x^2 + \sqrt{x^4 - 12x^3 - 12c_1x + 6x^2 + 9} + 3}{6x}$$

### ✓ Solution by Mathematica

Time used: 0.477 (sec). Leaf size: 83

```
DSolve[(6*x*y[x]+x^2+3)*y'[x]+3*y[x]^2+2*x*y[x]+2*x==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x^2 + \sqrt{x^4 - 12x^3 + 6x^2 + 36c_1x + 9} + 3}{6x}$$

$$y(x) \rightarrow -\frac{x^2 - \sqrt{x^4 - 12x^3 + 6x^2 + 36c_1x + 9} + 3}{6x}$$

## 6.31 problem Exercise 12.31, page 103

Internal problem ID [4552]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.31, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _Riccati]`

$$x^2y' + y^2 + yx = -x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(x^2*diff(y(x),x)+y(x)^2+x*y(x)+x^2=0,y(x), singsol=all)
```

$$y(x) = -\frac{x(\ln(x) + c_1 - 1)}{\ln(x) + c_1}$$

### ✓ Solution by Mathematica

Time used: 0.139 (sec). Leaf size: 31

```
DSolve[x^2*y'[x]+y[x]^2+x*y[x]+x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x(\log(x) - 1 - c_1)}{-\log(x) + c_1}$$

$$y(x) \rightarrow -x$$

## 6.32 problem Exercise 12.32, page 103

Internal problem ID [4553]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.32, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$(x^2 - 1)y' + 2yx = \cos(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve((x^2-1)*diff(y(x),x)+2*x*y(x)-cos(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\sin(x) + c_1}{(x-1)(x+1)}$$

### ✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 18

```
DSolve[(x^2-1)*y'[x]+2*x*y[x]-Cos[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sin(x) + c_1}{x^2 - 1}$$

### 6.33 problem Exercise 12.33, page 103

Internal problem ID [4554]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.33, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_exact, \_rational, [\_Abel, '2nd type', 'class B']]

$$(x^2y - 1)y' + y^2x = 1$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 50

```
dsolve((x^2*y(x)-1)*diff(y(x),x)+x*y(x)^2-1=0,y(x), singsol=all)
```

$$y(x) = \frac{1 + \sqrt{-2c_1x^2 + 2x^3 + 1}}{x^2}$$
$$y(x) = -\frac{-1 + \sqrt{-2c_1x^2 + 2x^3 + 1}}{x^2}$$

#### ✓ Solution by Mathematica

Time used: 0.505 (sec). Leaf size: 57

```
DSolve[(x^2*y[x]-1)*y'[x]+x*y[x]^2-1==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1 - \sqrt{2x^3 + c_1x^2 + 1}}{x^2}$$
$$y(x) \rightarrow \frac{1 + \sqrt{2x^3 + c_1x^2 + 1}}{x^2}$$

## 6.34 problem Exercise 12.34, page 103

Internal problem ID [4555]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.34, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$(x^2 - 1) y' + yx - 3y^2x = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve((x^2-1)*diff(y(x),x)+x*y(x)-3*x*y(x)^2=0,y(x), singsol=all)
```

$$y(x) = \frac{1}{3 + \sqrt{x-1}\sqrt{x+1}c_1}$$

✓ Solution by Mathematica

Time used: 2.214 (sec). Leaf size: 35

```
DSolve[(x^2-1)*y'[x]+x*y[x]-3*x*y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3 + e^{c_1}\sqrt{x^2-1}}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow \frac{1}{3}$$

## 6.35 problem Exercise 12.35, page 103

Internal problem ID [4556]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.35, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$(x^2 - 1)y' - 2xy \ln(y) = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve((x^2-1)*diff(y(x),x)-2*x*y(x)*ln(y(x))=0,y(x), singsol=all)
```

$$y(x) = e^{c_1(x+1)(x-1)}$$

### ✓ Solution by Mathematica

Time used: 0.223 (sec). Leaf size: 22

```
DSolve[(x^2-1)*y'[x]-2*x*y[x]*Log[y[x]]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{e^{c_1}(x^2-1)}$$

$$y(x) \rightarrow 1$$



## 6.36 problem Exercise 12.36, page 103

Internal problem ID [4557]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.36, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_exact`, `_rational`]

$$(1 + x^2 + y^2) y' + 2yx = -x^2 - 3$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 570

`dsolve((x^2+y(x)^2+1)*diff(y(x),x)+2*x*y(x)+x^2+3=0,y(x), singsol=all)`

$$\begin{aligned}
 y(x) &= \frac{\left(-4x^3 - 12c_1 - 36x + 4\sqrt{5x^6 + 6c_1x^3 + 30x^4 + 9c_1^2 + 54c_1x + 93x^2 + 4}\right)^{\frac{1}{3}}}{2(x^2 + 1)} \\
 &\quad - \frac{\left(-4x^3 - 12c_1 - 36x + 4\sqrt{5x^6 + 6c_1x^3 + 30x^4 + 9c_1^2 + 54c_1x + 93x^2 + 4}\right)^{\frac{1}{3}}}{2(x^2 + 1)} \\
 y(x) &= -\frac{\left(-4x^3 - 12c_1 - 36x + 4\sqrt{5x^6 + 6c_1x^3 + 30x^4 + 9c_1^2 + 54c_1x + 93x^2 + 4}\right)^{\frac{1}{3}}}{x^2 + 1} \\
 &\quad + \frac{\left(-4x^3 - 12c_1 - 36x + 4\sqrt{5x^6 + 6c_1x^3 + 30x^4 + 9c_1^2 + 54c_1x + 93x^2 + 4}\right)^{\frac{1}{3}}}{x^2 + 1} \\
 &\quad - \frac{i\sqrt{3} \left( \frac{\left(-4x^3 - 12c_1 - 36x + 4\sqrt{5x^6 + 6c_1x^3 + 30x^4 + 9c_1^2 + 54c_1x + 93x^2 + 4}\right)^{\frac{1}{3}}}{2} + \frac{2x^2 + 2}{\left(-4x^3 - 12c_1 - 36x + 4\sqrt{5x^6 + 6c_1x^3 + 30x^4 + 9c_1^2 + 54c_1x + 93x^2 + 4}\right)^{\frac{1}{3}}} \right)}{2} \\
 y(x) &= -\frac{\left(-4x^3 - 12c_1 - 36x + 4\sqrt{5x^6 + 6c_1x^3 + 30x^4 + 9c_1^2 + 54c_1x + 93x^2 + 4}\right)^{\frac{1}{3}}}{x^2 + 1} \\
 &\quad + \frac{\left(-4x^3 - 12c_1 - 36x + 4\sqrt{5x^6 + 6c_1x^3 + 30x^4 + 9c_1^2 + 54c_1x + 93x^2 + 4}\right)^{\frac{1}{3}}}{x^2 + 1} \\
 &\quad - \frac{i\sqrt{3} \left( \frac{\left(-4x^3 - 12c_1 - 36x + 4\sqrt{5x^6 + 6c_1x^3 + 30x^4 + 9c_1^2 + 54c_1x + 93x^2 + 4}\right)^{\frac{1}{3}}}{2} + \frac{2x^2 + 2}{\left(-4x^3 - 12c_1 - 36x + 4\sqrt{5x^6 + 6c_1x^3 + 30x^4 + 9c_1^2 + 54c_1x + 93x^2 + 4}\right)^{\frac{1}{3}}} \right)}{2} \\
 &\quad + \frac{\left(-4x^3 - 12c_1 - 36x + 4\sqrt{5x^6 + 6c_1x^3 + 30x^4 + 9c_1^2 + 54c_1x + 93x^2 + 4}\right)^{\frac{1}{3}}}{x^2 + 1}
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 5.385 (sec). Leaf size: 411

`DSolve[(x^2+y[x]^2+1)*y'[x]+2*x*y[x]+x^2+3==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{\sqrt[3]{-27x^3 + \sqrt{4(9x^2 + 9)^3 + 729(x^3 + 9x - 3c_1)^2 - 243x + 81c_1}}}{3\sqrt[3]{2}} - \frac{3\sqrt[3]{2}(x^2 + 1)}{\sqrt[3]{-27x^3 + \sqrt{4(9x^2 + 9)^3 + 729(x^3 + 9x - 3c_1)^2 - 243x + 81c_1}}}$$

$$y(x) \rightarrow \frac{3(1 + i\sqrt{3})(x^2 + 1)}{2^{2/3}\sqrt[3]{-27x^3 + \sqrt{4(9x^2 + 9)^3 + 729(x^3 + 9x - 3c_1)^2 - 243x + 81c_1}}} + \frac{(-1 + i\sqrt{3})\sqrt[3]{-27x^3 + \sqrt{4(9x^2 + 9)^3 + 729(x^3 + 9x - 3c_1)^2 - 243x + 81c_1}}}{6\sqrt[3]{2}}$$

$$y(x) \rightarrow \frac{3(1 - i\sqrt{3})(x^2 + 1)}{2^{2/3}\sqrt[3]{-27x^3 + \sqrt{4(9x^2 + 9)^3 + 729(x^3 + 9x - 3c_1)^2 - 243x + 81c_1}}} - \frac{(1 + i\sqrt{3})\sqrt[3]{-27x^3 + \sqrt{4(9x^2 + 9)^3 + 729(x^3 + 9x - 3c_1)^2 - 243x + 81c_1}}}{6\sqrt[3]{2}}$$

## 6.37 problem Exercise 12.37, page 103

Internal problem ID [4558]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.37, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$\cos(x)y' + y = -(\sin(x) + 1)\cos(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x)*cos(x)+y(x)+(1+sin(x))*cos(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-2 \ln(\sec(x) + \tan(x)) + 2 \ln(\cos(x)) + \sin(x) + c_1}{\sec(x) + \tan(x)}$$

✓ Solution by Mathematica

Time used: 0.671 (sec). Leaf size: 40

```
DSolve[y'[x]*Cos[x]+y[x]+(1+Sin[x])*Cos[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2\operatorname{arctanh}\left(\tan\left(\frac{x}{2}\right)\right)} \left( \sin(x) + 4 \log\left(\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)\right) + c_1 \right)$$

## 6.38 problem Exercise 12.38, page 103

Internal problem ID [4559]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.38, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _exact, _rational, [_Abel, '2nd ty`

$$(2yx + 4x^3)y' + y^2 + 12x^2y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 50

```
dsolve((2*x*y(x)+4*x^3)*diff(y(x),x)+y(x)^2+12*x^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-2x^3 + \sqrt{4x^6 + c_1x}}{x}$$

$$y(x) = -\frac{2x^3 + \sqrt{4x^6 + c_1x}}{x}$$

### ✓ Solution by Mathematica

Time used: 0.441 (sec). Leaf size: 58

```
DSolve[(2*x*y[x]+4*x^3)*y'[x]+y[x]^2+12*x^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2x^3 + \sqrt{x(4x^5 + c_1)}}{x}$$

$$y(x) \rightarrow \frac{-2x^3 + \sqrt{x(4x^5 + c_1)}}{x}$$

## 6.39 problem Exercise 12.39, page 103

Internal problem ID [4560]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.39, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_rational, [_1st_order, ‘_with_symmetry_[F(x)*G(y),0]’], [_Ab`

$$(x^2 - y) y' = -x$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 23

```
dsolve((x^2-y(x))*diff(y(x),x)+x=0,y(x), singsol=all)
```

$$y(x) = x^2 + \frac{\text{LambertW}\left(4c_1 e^{-2x^2-1}\right)}{2} + \frac{1}{2}$$

### ✓ Solution by Mathematica

Time used: 5.105 (sec). Leaf size: 40

```
DSolve[(x^2-y[x])*y'[x]+x==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2 + \frac{1}{2} \left( 1 + W\left(-e^{-2x^2-1+c_1}\right) \right)$$

$$y(x) \rightarrow x^2 + \frac{1}{2}$$

## 6.40 problem Exercise 12.40, page 103

Internal problem ID [4561]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.40, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational, [_Abel, '2nd type', 'cl`

$$(x^2 - y) y' - 4yx = 0$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 53

```
dsolve((x^2-y(x))*diff(y(x),x)-4*x*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 \left( c_1 - \sqrt{c_1^2 - 4x^2} \right)}{2} - x^2$$

$$y(x) = \frac{c_1 \left( c_1 + \sqrt{c_1^2 - 4x^2} \right)}{2} - x^2$$

✓ Solution by Mathematica

Time used: 2.441 (sec). Leaf size: 246

`DSolve[(x^2-y[x])*y'[x]-4*x*y[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow x^2 \left( 1 + \frac{2 - 2i}{\frac{i\sqrt{2}}{\sqrt{x^2 \cosh\left(\frac{2c_1}{9}\right) + x^2 \sinh\left(\frac{2c_1}{9}\right) - i}} - (1 - i)} \right)$$

$$y(x) \rightarrow x^2 \left( 1 + \frac{2 - 2i}{(-1 + i) - \frac{i\sqrt{2}}{\sqrt{x^2 \cosh\left(\frac{2c_1}{9}\right) + x^2 \sinh\left(\frac{2c_1}{9}\right) - i}}} \right)$$

$$y(x) \rightarrow x^2 \left( 1 + \frac{2 - 2i}{(-1 + i) - \frac{\sqrt{2}}{\sqrt{x^2 \cosh\left(\frac{2c_1}{9}\right) + x^2 \sinh\left(\frac{2c_1}{9}\right) + i}}} \right)$$

$$y(x) \rightarrow x^2 \left( 1 + \frac{2 - 2i}{\frac{\sqrt{2}}{\sqrt{x^2 \cosh\left(\frac{2c_1}{9}\right) + x^2 \sinh\left(\frac{2c_1}{9}\right) + i}} - (1 - i)} \right)$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -x^2$$



## 6.41 problem Exercise 12.41, page 103

Internal problem ID [4562]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.41, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _Bernoulli]`

$$yy'x + y^2 = -x^2$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 39

```
dsolve(x*y(x)*diff(y(x),x)+x^2+y(x)^2=0,y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{-2x^4 + 4c_1}}{2x}$$

$$y(x) = \frac{\sqrt{-2x^4 + 4c_1}}{2x}$$

### ✓ Solution by Mathematica

Time used: 0.211 (sec). Leaf size: 46

```
DSolve[x*y[x]*y'[x]+x^2+y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{-\frac{x^4}{2} + c_1}}{x}$$

$$y(x) \rightarrow \frac{\sqrt{-\frac{x^4}{2} + c_1}}{x}$$

## 6.42 problem Exercise 12.42, page 103

Internal problem ID [4563]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.42, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$2yy'x - y^2 = -3x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(2*x*y(x)*diff(y(x),x)+3*x^2-y(x)^2=0,y(x), singsol=all)
```

$$y(x) = \sqrt{c_1x - 3x^2}$$

$$y(x) = -\sqrt{c_1x - 3x^2}$$

### ✓ Solution by Mathematica

Time used: 0.306 (sec). Leaf size: 35

```
DSolve[2*x*y[x]*y'[x]+3*x^2-y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{x(-3x + c_1)}$$

$$y(x) \rightarrow \sqrt{x(-3x + c_1)}$$

## 6.43 problem Exercise 12.43, page 103

Internal problem ID [4564]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.43, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A'], _rational, _dAlembert]`

$$(2y^3x - x^4)y' + 2yx^3 - y^4 = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 447

`dsolve((2*x*y(x)^3-x^4)*diff(y(x),x)+2*x^3*y(x)-y(x)^4=0,y(x), singsol=all)`

$$\begin{aligned}
 y(x) &= \frac{12^{\frac{1}{3}} \left( x \left( -9c_1x^2 + \sqrt{3} \sqrt{\frac{x(27c_1^3x^3-4)}{c_1}} \right) c_1^2 \right)^{\frac{1}{3}}}{6c_1} \\
 &+ \frac{x12^{\frac{2}{3}}}{12 \left( x \left( -9c_1x^2 + \sqrt{3} \sqrt{\frac{x(27c_1^3x^3-4)}{c_1}} \right) c_1^2 \right)^{\frac{1}{3}}} \\
 y(x) &= - \frac{12^{\frac{1}{3}} \left( x \left( -9c_1x^2 + \sqrt{3} \sqrt{\frac{x(27c_1^3x^3-4)}{c_1}} \right) c_1^2 \right)^{\frac{1}{3}}}{12c_1} \\
 &- \frac{x12^{\frac{2}{3}}}{12 \left( x \left( -9c_1x^2 + \sqrt{3} \sqrt{\frac{x(27c_1^3x^3-4)}{c_1}} \right) c_1^2 \right)^{\frac{1}{3}}} \\
 &- \frac{i\sqrt{3} \left( \frac{12^{\frac{1}{3}} \left( x \left( -9c_1x^2 + \sqrt{3} \sqrt{\frac{x(27c_1^3x^3-4)}{c_1}} \right) c_1^2 \right)^{\frac{1}{3}}}{6c_1} - \frac{x12^{\frac{2}{3}}}{6 \left( x \left( -9c_1x^2 + \sqrt{3} \sqrt{\frac{x(27c_1^3x^3-4)}{c_1}} \right) c_1^2 \right)^{\frac{1}{3}}} \right)}{2} \\
 y(x) &= - \frac{12^{\frac{1}{3}} \left( x \left( -9c_1x^2 + \sqrt{3} \sqrt{\frac{x(27c_1^3x^3-4)}{c_1}} \right) c_1^2 \right)^{\frac{1}{3}}}{12c_1} \\
 &- \frac{x12^{\frac{2}{3}}}{12 \left( x \left( -9c_1x^2 + \sqrt{3} \sqrt{\frac{x(27c_1^3x^3-4)}{c_1}} \right) c_1^2 \right)^{\frac{1}{3}}} \\
 &+ \frac{i\sqrt{3} \left( \frac{12^{\frac{1}{3}} \left( x \left( -9c_1x^2 + \sqrt{3} \sqrt{\frac{x(27c_1^3x^3-4)}{c_1}} \right) c_1^2 \right)^{\frac{1}{3}}}{6c_1} - \frac{x12^{\frac{2}{3}}}{6 \left( x \left( -9c_1x^2 + \sqrt{3} \sqrt{\frac{x(27c_1^3x^3-4)}{c_1}} \right) c_1^2 \right)^{\frac{1}{3}}} \right)}{2}
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 60.224 (sec). Leaf size: 331

```
DSolve[(2*x*y[x]^3-x^4)*y'[x]+2*x^3*y[x]-y[x]^4==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt[3]{2}(-9x^3 + \sqrt{81x^6 - 12e^{3c_1}x^3})^{2/3} + 2\sqrt[3]{3}e^{c_1}x}{6^{2/3}\sqrt[3]{-9x^3 + \sqrt{81x^6 - 12e^{3c_1}x^3}}}$$

$$y(x) \rightarrow \frac{i\sqrt[3]{2}\sqrt[6]{3}(\sqrt{3} + i)(-9x^3 + \sqrt{81x^6 - 12e^{3c_1}x^3})^{2/3} - 2(\sqrt{3} + 3i)e^{c_1}x}{2 \cdot 2^{2/3}3^{5/6}\sqrt[3]{-9x^3 + \sqrt{81x^6 - 12e^{3c_1}x^3}}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{2}\sqrt[6]{3}(-1 - i\sqrt{3})(-9x^3 + \sqrt{81x^6 - 12e^{3c_1}x^3})^{2/3} - 2(\sqrt{3} - 3i)e^{c_1}x}{2 \cdot 2^{2/3}3^{5/6}\sqrt[3]{-9x^3 + \sqrt{81x^6 - 12e^{3c_1}x^3}}}$$

## 6.44 problem Exercise 12.44, page 103

Internal problem ID [4565]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.44, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational]`

$$(-1 + yx)^2 xy' + (1 + x^2y^2) y = 0$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 34

```
dsolve((x*y(x)-1)^2*x*diff(y(x),x)+(x^2*y(x)^2+1)*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{e^{\text{RootOf}(-2e^{-Z}\ln(x)-e^{2-Z}+2e^{-Z}c_1+2_Ze^{-Z}+1)}}{x}$$

✓ Solution by Mathematica

Time used: 0.108 (sec). Leaf size: 25

```
DSolve[(x*y[x]-1)^2*x*y'[x]+(x^2*y[x]^2+1)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[xy(x) - \frac{1}{xy(x)} - 2\log(y(x)) = c_1, y(x)\right]$$

## 6.45 problem Exercise 12.45, page 103

Internal problem ID [4566]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.45, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _exact, _rational, _dAlembert]`

$$(x^2 + y^2) y' + 2x(y + 2x) = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 417

`dsolve((x^2+y(x)^2)*diff(y(x),x)+2*x*(2*x+y(x))=0,y(x), singsol=all)`

$$y(x) = \frac{\left(4-16x^3c_1^{\frac{3}{2}}+4\sqrt{20x^6c_1^3-8x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}{2} - \frac{2x^2c_1}{\left(4-16x^3c_1^{\frac{3}{2}}+4\sqrt{20x^6c_1^3-8x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}$$

$$y(x) = -\frac{\left(4-16x^3c_1^{\frac{3}{2}}+4\sqrt{20x^6c_1^3-8x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}{4} + \frac{x^2c_1}{\left(4-16x^3c_1^{\frac{3}{2}}+4\sqrt{20x^6c_1^3-8x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}} - \frac{i\sqrt{3}\left(\frac{\left(4-16x^3c_1^{\frac{3}{2}}+4\sqrt{20x^6c_1^3-8x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}{2} + \frac{\left(4-16x^3c_1^{\frac{3}{2}}+4\sqrt{20x^6c_1^3-8x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}{2}\right)}{\sqrt{c_1}}$$

$$y(x) = -\frac{\left(4-16x^3c_1^{\frac{3}{2}}+4\sqrt{20x^6c_1^3-8x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}{4} + \frac{x^2c_1}{\left(4-16x^3c_1^{\frac{3}{2}}+4\sqrt{20x^6c_1^3-8x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}} + \frac{i\sqrt{3}\left(\frac{\left(4-16x^3c_1^{\frac{3}{2}}+4\sqrt{20x^6c_1^3-8x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}{2} + \frac{\left(4-16x^3c_1^{\frac{3}{2}}+4\sqrt{20x^6c_1^3-8x^3c_1^{\frac{3}{2}}+1}\right)^{\frac{1}{3}}}{2}\right)}{\sqrt{c_1}}$$



✓ Solution by Mathematica

Time used: 18.874 (sec). Leaf size: 593

DSolve[(x^2+y[x]^2)\*y'[x]+2\*x\*(2\*x+y[x])==0,y[x],x,IncludeSingularSolutions -> True]

$$y(x) \rightarrow \frac{\sqrt[3]{-4x^3 + \sqrt{20x^6 - 8e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}}{\sqrt[3]{2}} - \frac{\sqrt[3]{2}x^2}{\sqrt[3]{-4x^3 + \sqrt{20x^6 - 8e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{2}(2 + 2i\sqrt{3})x^2 + i2^{2/3}(\sqrt{3} + i)(-4x^3 + \sqrt{20x^6 - 8e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1})^{2/3}}{4\sqrt[3]{-4x^3 + \sqrt{20x^6 - 8e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}}$$

$$y(x) \rightarrow \frac{(1 - i\sqrt{3})x^2}{2^{2/3}\sqrt[3]{-4x^3 + \sqrt{20x^6 - 8e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}} - \frac{(1 + i\sqrt{3})\sqrt[3]{-4x^3 + \sqrt{20x^6 - 8e^{3c_1}x^3 + e^{6c_1}} + e^{3c_1}}}{2\sqrt[3]{2}}$$

$$y(x) \rightarrow \sqrt[3]{\sqrt{5}\sqrt{x^6} - 2x^3} - \frac{x^2}{\sqrt[3]{\sqrt{5}\sqrt{x^6} - 2x^3}}$$

$$y(x) \rightarrow \frac{(1 - i\sqrt{3})x^2 + (-1 - i\sqrt{3})(\sqrt{5}\sqrt{x^6} - 2x^3)^{2/3}}{2\sqrt[3]{\sqrt{5}\sqrt{x^6} - 2x^3}}$$

$$y(x) \rightarrow \frac{(1 + i\sqrt{3})x^2 + i(\sqrt{3} + i)(\sqrt{5}\sqrt{x^6} - 2x^3)^{2/3}}{2\sqrt[3]{\sqrt{5}\sqrt{x^6} - 2x^3}}$$

## 6.46 problem Exercise 12.46, page 103

Internal problem ID [4567]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.46, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _exact, _rational, _Bernoulli]`

$$3xy^2y' + y^3 = 2x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 99

```
dsolve(3*x*y(x)^2*diff(y(x),x)+y(x)^3-2*x=0,y(x), singsol=all)
```

$$y(x) = \frac{((x^2 + c_1) x^2)^{\frac{1}{3}}}{x}$$

$$y(x) = -\frac{((x^2 + c_1) x^2)^{\frac{1}{3}}}{2x} - \frac{i\sqrt{3}((x^2 + c_1) x^2)^{\frac{1}{3}}}{2x}$$

$$y(x) = -\frac{((x^2 + c_1) x^2)^{\frac{1}{3}}}{2x} + \frac{i\sqrt{3}((x^2 + c_1) x^2)^{\frac{1}{3}}}{2x}$$

✓ Solution by Mathematica

Time used: 0.224 (sec). Leaf size: 72

```
DSolve[3*x*y[x]^2*y'[x]+y[x]^3-2*x==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt[3]{x^2 + c_1}}{\sqrt[3]{x}}$$

$$y(x) \rightarrow -\frac{\sqrt[3]{-1}\sqrt[3]{x^2 + c_1}}{\sqrt[3]{x}}$$

$$y(x) \rightarrow \frac{(-1)^{2/3}\sqrt[3]{x^2 + c_1}}{\sqrt[3]{x}}$$

## 6.47 problem Exercise 12.47, page 103

Internal problem ID [4568]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.47, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A'], _rational, _dAlembert]`

$$2y^3y' + y^2x = x^3$$

✓ Solution by Maple

Time used: 0.453 (sec). Leaf size: 711

`dsolve(2*y(x)^3*diff(y(x),x)+x*y(x)^2-x^3=0,y(x), singsol=all)`

$$y(x) = -\frac{\sqrt{2\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}+\frac{2x^4c_1^2}{\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}}-2c_1x^2}}{2\sqrt{c_1}}$$

$$y(x) = \frac{\sqrt{2\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}+\frac{2x^4c_1^2}{\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}}-2c_1x^2}}{2\sqrt{c_1}}$$

$$y(x) = \frac{\sqrt{-\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}-\frac{x^4c_1^2}{\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}}-2c_1x^2-2i\sqrt{3}\left(\frac{\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}}{2}-\frac{1}{2\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)}\right)}}{2\sqrt{c_1}}$$

$$y(x) = \frac{\sqrt{-\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}-\frac{x^4c_1^2}{\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}}-2c_1x^2-2i\sqrt{3}\left(\frac{\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}}{2}-\frac{1}{2\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)}\right)}}{2\sqrt{c_1}}$$

$$y(x) = \frac{\sqrt{-\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}-\frac{x^4c_1^2}{\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}}-2c_1x^2+2i\sqrt{3}\left(\frac{\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}}{2}-\frac{1}{2\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)}\right)}}{2\sqrt{c_1}}$$

$$y(x) = \frac{\sqrt{-\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}-\frac{x^4c_1^2}{\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}}-2c_1x^2+2i\sqrt{3}\left(\frac{\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)^{\frac{1}{3}}}{2}-\frac{1}{2\left(2+x^6c_1^3+2\sqrt{x^6c_1^3+1}\right)}\right)}}{2\sqrt{c_1}}$$

✓ Solution by Mathematica

Time used: 60.13 (sec). Leaf size: 714

`DSolve[2*y[x]^3*y'[x]+x*y[x]^2-x^3==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow -\frac{\sqrt{\sqrt[3]{x^6 + 2\sqrt{e^{24c_1} - e^{12c_1}x^6} - 2e^{12c_1}} - x^2 + \frac{x^4}{\sqrt[3]{x^6 + 2\sqrt{e^{24c_1} - e^{12c_1}x^6} - 2e^{12c_1}}}}}{\sqrt{2}}$$

$$y(x) \rightarrow \frac{\sqrt{\sqrt[3]{x^6 + 2\sqrt{e^{24c_1} - e^{12c_1}x^6} - 2e^{12c_1}} - x^2 + \frac{x^4}{\sqrt[3]{x^6 + 2\sqrt{e^{24c_1} - e^{12c_1}x^6} - 2e^{12c_1}}}}}{\sqrt{2}}$$

$$y(x) \rightarrow -\frac{1}{2}\sqrt{\left(-1 - i\sqrt{3}\right)\sqrt[3]{x^6 + 2\sqrt{e^{24c_1} - e^{12c_1}x^6} - 2e^{12c_1}} - 2x^2 + \frac{i(\sqrt{3} + i)x^4}{\sqrt[3]{x^6 + 2\sqrt{e^{24c_1} - e^{12c_1}x^6} - 2e^{12c_1}}}}$$

$$y(x) \rightarrow \frac{1}{2}\sqrt{\left(-1 - i\sqrt{3}\right)\sqrt[3]{x^6 + 2\sqrt{e^{24c_1} - e^{12c_1}x^6} - 2e^{12c_1}} - 2x^2 + \frac{i(\sqrt{3} + i)x^4}{\sqrt[3]{x^6 + 2\sqrt{e^{24c_1} - e^{12c_1}x^6} - 2e^{12c_1}}}}$$

$$y(x) \rightarrow -\frac{1}{2}\sqrt{i(\sqrt{3} + i)\sqrt[3]{x^6 + 2\sqrt{e^{24c_1} - e^{12c_1}x^6} - 2e^{12c_1}} - 2x^2 + \frac{(-1 - i\sqrt{3})x^4}{\sqrt[3]{x^6 + 2\sqrt{e^{24c_1} - e^{12c_1}x^6} - 2e^{12c_1}}}}$$

$$y(x) \rightarrow \frac{1}{2}\sqrt{i(\sqrt{3} + i)\sqrt[3]{x^6 + 2\sqrt{e^{24c_1} - e^{12c_1}x^6} - 2e^{12c_1}} - 2x^2 + \frac{(-1 - i\sqrt{3})x^4}{\sqrt[3]{x^6 + 2\sqrt{e^{24c_1} - e^{12c_1}x^6} - 2e^{12c_1}}}}$$

## 6.48 problem Exercise 12.48, page 103

Internal problem ID [4569]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.48, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_rational]`

$$(2y^3x + yx + x^2)y' - yx + y^2 = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve((2*x*y(x)^3+x*y(x)+x^2)*diff(y(x),x)-x*y(x)+y(x)^2=0,y(x), singsol=all)
```

$$y(x) = e^{\text{RootOf}(-e^{3-Z} - e^{-Z} \ln(x) + e^{-Z} c_1 - Z e^{-Z} + x)}$$

✓ Solution by Mathematica

Time used: 0.225 (sec). Leaf size: 23

```
DSolve[(2*x*y[x]^3+x*y[x]+x^2)*y'[x]-x*y[x]+y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ y(x)^2 - \frac{x}{y(x)} + \log(y(x)) + \log(x) = c_1, y(x) \right]$$

## 6.49 problem Exercise 12.49, page 103

Internal problem ID [4570]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.49, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$(2y^3 + y) y' = 2x^3 + x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 113

```
dsolve((2*y(x)^3+y(x))*diff(y(x),x)-2*x^3-x=0,y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{-2 - 2\sqrt{4x^4 + 4x^2 + 8c_1 + 1}}}{2}$$

$$y(x) = \frac{\sqrt{-2 - 2\sqrt{4x^4 + 4x^2 + 8c_1 + 1}}}{2}$$

$$y(x) = -\frac{\sqrt{-2 + 2\sqrt{4x^4 + 4x^2 + 8c_1 + 1}}}{2}$$

$$y(x) = \frac{\sqrt{-2 + 2\sqrt{4x^4 + 4x^2 + 8c_1 + 1}}}{2}$$



✓ Solution by Mathematica

Time used: 2.313 (sec). Leaf size: 151

```
DSolve[(2*y[x]^3+y[x])*y'[x]-2*x^3-x==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{-1 - \sqrt{4x^4 + 4x^2 + 1 + 8c_1}}}{\sqrt{2}}$$

$$y(x) \rightarrow \frac{\sqrt{-1 - \sqrt{4x^4 + 4x^2 + 1 + 8c_1}}}{\sqrt{2}}$$

$$y(x) \rightarrow -\frac{\sqrt{-1 + \sqrt{4x^4 + 4x^2 + 1 + 8c_1}}}{\sqrt{2}}$$

$$y(x) \rightarrow \frac{\sqrt{-1 + \sqrt{4x^4 + 4x^2 + 1 + 8c_1}}}{\sqrt{2}}$$

## 6.50 problem Exercise 12.50, page 103

Internal problem ID [4571]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 2. Special types of differential equations of the first kind. Lesson 12, Miscellaneous Methods

**Problem number:** Exercise 12.50, page 103.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y' - e^{-y+x} = -e^x$$

### ✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 20

```
dsolve(diff(y(x),x)-exp(x-y(x))+exp(x)=0,y(x), singsol=all)
```

$$y(x) = -e^x + \ln(-1 + e^{e^x+c_1}) - c_1$$

### ✓ Solution by Mathematica

Time used: 2.135 (sec). Leaf size: 23

```
DSolve[y'[x]-Exp[x-y[x]]+Exp[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(1 + e^{-e^x+c_1})$$

$$y(x) \rightarrow 0$$

## 7 Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

7.1	problem Exercise 20.1, page 220 . . . . .	195
7.2	problem Exercise 20.2, page 220 . . . . .	196
7.3	problem Exercise 20.3, page 220 . . . . .	197
7.4	problem Exercise 20.5, page 220 . . . . .	198
7.5	problem Exercise 20.6, page 220 . . . . .	199
7.6	problem Exercise 20.7, page 220 . . . . .	200
7.7	problem Exercise 20.8, page 220 . . . . .	201
7.8	problem Exercise 20.9, page 220 . . . . .	202
7.9	problem Exercise 20.10, page 220 . . . . .	203
7.10	problem Exercise 20.11, page 220 . . . . .	204
7.11	problem Exercise 20.12, page 220 . . . . .	205
7.12	problem Exercise 20.13, page 220 . . . . .	206
7.13	problem Exercise 20.14, page 220 . . . . .	207
7.14	problem Exercise 20.15, page 220 . . . . .	208
7.15	problem Exercise 20.16, page 220 . . . . .	209
7.16	problem Exercise 20.17, page 220 . . . . .	210
7.17	problem Exercise 20.18, page 220 . . . . .	211
7.18	problem Exercise 20.19, page 220 . . . . .	212
7.19	problem Exercise 20.20, page 220 . . . . .	213
7.20	problem Exercise 20.21, page 220 . . . . .	214
7.21	problem Exercise 20.22, page 220 . . . . .	215
7.22	problem Exercise 20.23, page 220 . . . . .	216
7.23	problem Exercise 20.24, page 220 . . . . .	217
7.24	problem Exercise 20.25, page 220 . . . . .	218
7.25	problem Exercise 20.26, page 220 . . . . .	219
7.26	problem Exercise 20.27, page 220 . . . . .	220
7.27	problem Exercise 20.28, page 220 . . . . .	221
7.28	problem Exercise 20.29, page 220 . . . . .	222
7.29	problem Exercise 20.30, page 220 . . . . .	223
7.30	problem Exercise 20, problem 31, page 220 . . . . .	224
7.31	problem Exercise 20, problem 32, page 220 . . . . .	225
7.32	problem Exercise 20, problem 33, page 220 . . . . .	226
7.33	problem Exercise 20, problem 34, page 220 . . . . .	227
7.34	problem Exercise 20, problem 35, page 220 . . . . .	228

## 7.1 problem Exercise 20.1, page 220

Internal problem ID [4572]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.1, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1 + c_2 e^{-2x}$$

### ✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 19

```
DSolve[y''[x]+2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \frac{1}{2}c_1 e^{-2x}$$

## 7.2 problem Exercise 20.2, page 220

Internal problem ID [4573]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.2, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 3y' + 2y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)+2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^x + c_2 e^{2x}$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 18

```
DSolve[y''[x]-3*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(c_2 e^x + c_1)$$

### 7.3 problem Exercise 20.3, page 220

Internal problem ID [4574]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.3, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x$2)-y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1e^x + c_2e^{-x}$$

#### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 20

```
DSolve[y''[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1e^x + c_2e^{-x}$$

## 7.4 problem Exercise 20.5, page 220

Internal problem ID [4575]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.5, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$6y'' - 11y' + 4y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(6*diff(y(x),x$2)-11*diff(y(x),x)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{4x}{3}} + c_2 e^{\frac{x}{2}}$$

### ✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 35

```
DSolve[y''[x]-11*y'[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\frac{1}{2}(\sqrt{105}-11)x} \left( c_2 e^{\sqrt{105}x} + c_1 \right)$$

## 7.5 problem Exercise 20.6, page 220

Internal problem ID [4576]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.6, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2y' - y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{(\sqrt{2}-1)x} + c_2 e^{-(1+\sqrt{2})x}$$

### ✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 34

```
DSolve[y''[x]+2*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-((1+\sqrt{2})x)} \left( c_2 e^{2\sqrt{2}x} + c_1 \right)$$



## 7.6 problem Exercise 20.7, page 220

Internal problem ID [4577]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.7, page 220.

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' + y'' - 10y' - 6y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
dsolve(diff(y(x),x$3)+diff(y(x),x$2)-10*diff(y(x),x)-6*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{3x}c_1 + c_2e^{(-2+\sqrt{2})x} + c_3e^{-(2+\sqrt{2})x}$$

### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 43

```
DSolve[y'''[x]+y''[x]-10*y'[x]-6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1e^{-((2+\sqrt{2})x)} + c_2e^{(\sqrt{2}-2)x} + c_3e^{3x}$$

## 7.7 problem Exercise 20.8, page 220

Internal problem ID [4578]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.8, page 220.

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - y''' - 4y'' + 4y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$4)-diff(y(x),x$3)-4*diff(y(x),x$2)+4*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1 + e^x c_2 + c_3 e^{-2x} + c_4 e^{2x}$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 36

```
DSolve[y''''[x]-y'''[x]-4*y''[x]+4*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2}c_1 e^{-2x} + c_2 e^x + \frac{1}{2}c_3 e^{2x} + c_4$$

## 7.8 problem Exercise 20.9, page 220

Internal problem ID [4579]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.9, page 220.

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 4y''' + y'' - 4y' - 2y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
dsolve(diff(y(x),x$4)+4*diff(y(x),x$3)+diff(y(x),x$2)-4*diff(y(x),x)-2*y(x)=0,y(x), singsol=
```

$$y(x) = c_1 e^x + c_2 e^{-x} + c_3 e^{(-2+\sqrt{2})x} + c_4 e^{-(2+\sqrt{2})x}$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 49

```
DSolve[y''''[x]+4*y'''[x]+y''[x]-4*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-((2+\sqrt{2})x)} + c_2 e^{(\sqrt{2}-2)x} + c_3 e^{-x} + c_4 e^x$$

## 7.9 problem Exercise 20.10, page 220

Internal problem ID [4580]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.10, page 220.

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - a^2y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 38

```
dsolve(diff(y(x),x$4)-a^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\sqrt{a}x} + c_2 e^{-\sqrt{a}x} + c_3 \sin(\sqrt{a}x) + c_4 \cos(\sqrt{a}x)$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 53

```
DSolve[y''''[x]-a^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 e^{-\sqrt{a}x} + c_4 e^{\sqrt{a}x} + c_1 \cos(\sqrt{a}x) + c_3 \sin(\sqrt{a}x)$$

## 7.10 problem Exercise 20.11, page 220

Internal problem ID [4581]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.11, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2ky' - 2y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$2)-2*k*diff(y(x),x)-2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{(k+\sqrt{k^2+2})x} + c_2 e^{(k-\sqrt{k^2+2})x}$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 44

```
DSolve[y''[x]-2*k*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{(k-\sqrt{k^2+2})x} + c_2 e^{(\sqrt{k^2+2}+k)x}$$

## 7.11 problem Exercise 20.12, page 220

Internal problem ID [4582]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.12, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 4ky' - 12k^2y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x$2)+4*k*diff(y(x),x)-12*k^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1e^{-6kx} + c_2e^{2kx}$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 24

```
DSolve[y''[x]+4*k*y'[x]-12*k^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-6kx}(c_2e^{8kx} + c_1)$$

## 7.12 problem Exercise 20.13, page 220

Internal problem ID [4583]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.13, page 220.

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _quadrature]]`

$$y'''' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$4)=0,y(x), singsol=all)
```

$$y(x) = \frac{1}{6}c_1x^3 + \frac{1}{2}c_2x^2 + c_3x + c_4$$

### ✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 22

```
DSolve[y''''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(x(c_4x + c_3) + c_2) + c_1$$

## 7.13 problem Exercise 20.14, page 220

Internal problem ID [4584]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.14, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 4y' + 4y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-2x} + c_2 e^{-2x} x$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 18

```
DSolve[y''[x]+4*y'[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(c_2 x + c_1)$$



## 7.14 problem Exercise 20.15, page 220

Internal problem ID [4585]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.15, page 220.

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$3y''' + 5y'' + y' - y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(3*diff(y(x),x$3)+5*diff(y(x),x$2)+diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{x}{3}} + c_2 e^{-x} + c_3 e^{-x} x$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 28

```
DSolve[3*y'''[x]+5*y''[x]+y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} (c_1 e^{4x/3} + c_3 x + c_2)$$

## 7.15 problem Exercise 20.16, page 220

Internal problem ID [4586]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.16, page 220.

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 6y'' + 12y' - 8y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$3)-6*diff(y(x),x$2)+12*diff(y(x),x)-8*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{2x} + c_2 e^{2x} x + c_3 e^{2x} x^2$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 23

```
DSolve[y'''[x]-6*y''[x]+12*y'[x]-8*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(x(c_3 x + c_2) + c_1)$$

## 7.16 problem Exercise 20.17, page 220

Internal problem ID [4587]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.17, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2ay' + ya^2 = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)-2*a*diff(y(x),x)+a^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{ax} + c_2 e^{ax} x$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 18

```
DSolve[y''[x]-2*a*y'[x]+a^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{ax}(c_2 x + c_1)$$

## 7.17 problem Exercise 20.18, page 220

Internal problem ID [4588]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.18, page 220.

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 3y''' = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$4)+3*diff(y(x),x$3)=0,y(x), singsol=all)
```

$$y(x) = c_1 + c_2x + c_3x^2 + c_4e^{-3x}$$

### ✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 28

```
DSolve[y''''[x]+3*y'''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{27}c_1e^{-3x} + x(c_4x + c_3) + c_2$$

## 7.18 problem Exercise 20.19, page 220

Internal problem ID [4589]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.19, page 220.

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 2y'' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$4)-2*diff(y(x),x$2)=0,y(x), singsol=all)
```

$$y(x) = c_1 + c_2x + c_3e^{\sqrt{2}x} + c_4e^{-\sqrt{2}x}$$

### ✓ Solution by Mathematica

Time used: 0.073 (sec). Leaf size: 42

```
DSolve[y''''[x]-2*y''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}e^{-\sqrt{2}x} \left( c_1 e^{2\sqrt{2}x} + c_2 \right) + c_4x + c_3$$

## 7.19 problem Exercise 20.20, page 220

Internal problem ID [4590]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.20, page 220.

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 2y''' - 11y'' - 12y' + 36y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$4)+2*diff(y(x),x$3)-11*diff(y(x),x$2)-12*diff(y(x),x)+36*y(x)=0,y(x), sin
```

$$y(x) = c_1 e^{-3x} + c_2 e^{-3x} x + c_3 e^{2x} + c_4 e^{2x} x$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 35

```
DSolve[y''''[x]+2*y'''[x]-11*y''[x]-12*y'[x]+36*y[x]==0,y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow e^{-3x} (c_3 e^{5x} + x(c_4 e^{5x} + c_2) + c_1)$$

## 7.20 problem Exercise 20.21, page 220

Internal problem ID [4591]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.21, page 220.

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$36y'''' - 37y'' + 4y' + 5y = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 29

```
dsolve(36*diff(y(x),x$4)-37*diff(y(x),x$2)+4*diff(y(x),x)+5*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-x} + c_2 e^{\frac{x}{2}} + c_3 e^{-\frac{x}{3}} + c_4 e^{\frac{5x}{6}}$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 44

```
DSolve[36*y''''[x]-37*y''[x]+4*y'[x]+5*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} (c_1 e^{11x/6} + c_2 e^{2x/3} + c_3 e^{3x/2} + c_4)$$

## 7.21 problem Exercise 20.22, page 220

Internal problem ID [4592]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.22, page 220.

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 8y'' + 36y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 48

```
dsolve(diff(y(x),x$4)-8*diff(y(x),x$2)+36*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\sqrt{5}x} \sin(x) - c_2 e^{-\sqrt{5}x} \sin(x) + c_3 e^{\sqrt{5}x} \cos(x) + c_4 e^{-\sqrt{5}x} \cos(x)$$

### ✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 142

```
DSolve[y''''[x]-8*y''[x]+36*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\sqrt{6}x \cos\left(\frac{1}{2} \arctan\left(\frac{\sqrt{5}}{2}\right)\right)} \left( \left( c_3 e^{2\sqrt{6}x \cos\left(\frac{1}{2} \arctan\left(\frac{\sqrt{5}}{2}\right)\right)} + c_2 \right) \cos\left(\sqrt{6}x \sin\left(\frac{1}{2} \arctan\left(\frac{\sqrt{5}}{2}\right)\right)\right) \right. \\ \left. + \sin\left(\sqrt{6}x \sin\left(\frac{1}{2} \arctan\left(\frac{\sqrt{5}}{2}\right)\right)\right) \left( c_1 e^{2\sqrt{6}x \cos\left(\frac{1}{2} \arctan\left(\frac{\sqrt{5}}{2}\right)\right)} + c_4 \right) \right)$$



## 7.22 problem Exercise 20.23, page 220

Internal problem ID [4593]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.23, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' + 5y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+5*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^x \sin(2x) + c_2 e^x \cos(2x)$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 24

```
DSolve[y''[x]-2*y'[x]+5*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x (c_2 \cos(2x) + c_1 \sin(2x))$$

## 7.23 problem Exercise 20.24, page 220

Internal problem ID [4594]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.24, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y' + y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$2)-diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + c_2 e^{\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

### ✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 42

```
DSolve[y''[x]-y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x/2} \left( c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right)$$

## 7.24 problem Exercise 20.25, page 220

Internal problem ID [4595]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.25, page 220.

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 5y'' + 6y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 37

```
dsolve(diff(y(x),x$4)+5*diff(y(x),x$2)+6*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(\sqrt{2}x) + c_2 \cos(\sqrt{2}x) + c_3 \sin(\sqrt{3}x) + c_4 \cos(\sqrt{3}x)$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 50

```
DSolve[y''''[x]+5*y''[x]+6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3 \cos(\sqrt{2}x) + c_1 \cos(\sqrt{3}x) + c_4 \sin(\sqrt{2}x) + c_2 \sin(\sqrt{3}x)$$

## 7.25 problem Exercise 20.26, page 220

Internal problem ID [4596]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.26, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y' + 20y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+20*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{2x} \sin(4x) + c_2 e^{2x} \cos(4x)$$

### ✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 26

```
DSolve[y''[x]-4*y'[x]+20*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(c_2 \cos(4x) + c_1 \sin(4x))$$

## 7.26 problem Exercise 20.27, page 220

Internal problem ID [4597]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.27, page 220.

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 4y'' + 4y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
dsolve(diff(y(x),x$4)+4*diff(y(x),x$2)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(\sqrt{2}x) + c_2 \cos(\sqrt{2}x) + c_3 \sin(\sqrt{2}x)x + c_4 \cos(\sqrt{2}x)x$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 38

```
DSolve[y''''[x]+4*y''[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (c_2x + c_1) \cos(\sqrt{2}x) + (c_4x + c_3) \sin(\sqrt{2}x)$$

## 7.27 problem Exercise 20.28, page 220

Internal problem ID [4598]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.28, page 220.

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' + 8y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$3)+8*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-2x} + c_2 e^x \sin(\sqrt{3}x) + c_3 e^x \cos(\sqrt{3}x)$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 42

```
DSolve[y'''[x]+8*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-2x} + c_3 e^x \cos(\sqrt{3}x) + c_2 e^x \sin(\sqrt{3}x)$$

## 7.28 problem Exercise 20.29, page 220

Internal problem ID [4599]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.29, page 220.

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 4y'' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$4)+4*diff(y(x),x$2)=0,y(x), singsol=all)
```

$$y(x) = c_1 + c_2x + c_3 \sin(2x) + c_4 \cos(2x)$$

### ✓ Solution by Mathematica

Time used: 0.118 (sec). Leaf size: 32

```
DSolve[y''''[x]+4*y''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_4x - \frac{1}{4}c_1 \cos(2x) - \frac{1}{4}c_2 \sin(2x) + c_3$$

## 7.29 problem Exercise 20.30, page 220

Internal problem ID [4600]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20.30, page 220.

**ODE order:** 5.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y^{(5)} + 2y''' + y' = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$5)+2*diff(y(x),x$3)+diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1 + c_2 \sin(x) + c_3 \cos(x) + c_4 \sin(x)x + c_5 \cos(x)x$$

### ✓ Solution by Mathematica

Time used: 0.05 (sec). Leaf size: 35

```
DSolve[y'''''[x]+2*y'''[x]+y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (-c_4x + c_2 - c_3) \cos(x) + (c_2x + c_1 + c_4) \sin(x) + c_5$$



### 7.30 problem Exercise 20, problem 31, page 220

Internal problem ID [4601]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20, problem 31, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$y'' = 0$$

With initial conditions

$$[y(1) = 2, y'(1) = -1]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve([diff(y(x),x$2)=0,y(1) = 2, D(y)(1) = -1],y(x), singsol=all)
```

$$y(x) = 3 - x$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 10

```
DSolve[{y''[x]==0,{y[1]==2,y'[1]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3 - x$$

### 7.31 problem Exercise 20, problem 32, page 220

Internal problem ID [4602]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20, problem 32, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 4y' + 4y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 1]$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([diff(y(x),x$2)+4*diff(y(x),x)+4*y(x)=0,y(0) = 1, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = e^{-2x}(3x + 1)$$

#### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 16

```
DSolve[{y'[x]+4*y'[x]+4*y[x]==0,{y[0]==1,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow e^{-2x}(3x + 1)$$

## 7.32 problem Exercise 20, problem 33, page 220

Internal problem ID [4603]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20, problem 33, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' + 5y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = 1]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve([diff(y(x),x$2)-2*diff(y(x),x)+5*y(x)=0,y(0) = 2, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{e^x(-\sin(2x) + 4\cos(2x))}{2}$$

### ✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 25

```
DSolve[{y'[x]-2*y'[x]+5*y[x]==0,{y[0]==2,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{1}{2}e^x(4\cos(2x) - \sin(2x))$$

### 7.33 problem Exercise 20, problem 34, page 220

Internal problem ID [4604]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20, problem 34, page 220.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y' + 20y = 0$$

With initial conditions

$$\left[ y\left(\frac{\pi}{2}\right) = 1, y'\left(\frac{\pi}{2}\right) = 1 \right]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve([diff(y(x),x$2)-4*diff(y(x),x)+20*y(x)=0,y(1/2*Pi) = 1, D(y)(1/2*Pi) = 1],y(x), sings
```

$$y(x) = \frac{(-\sin(4x) + 4\cos(4x))e^{2x-\pi}}{4}$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 31

```
DSolve[{y'[x]-4*y'[x]+20*y[x]==0,{y[Pi/2]==1,y'[Pi/2]==1}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{1}{4}e^{2x-\pi}(4\cos(4x) - \sin(4x))$$

## 7.34 problem Exercise 20, problem 35, page 220

Internal problem ID [4605]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 20. Constant coefficients

**Problem number:** Exercise 20, problem 35, page 220.

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$3y''' + 5y'' + y' - y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1, y''(0) = -1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve([3*diff(y(x),x$3)+5*diff(y(x),x$2)+diff(y(x),x)-y(x)=0,y(0) = 0, D(y)(0) = 1, (D@@2)(
```

$$y(x) = \frac{\left(9e^{\frac{4x}{3}} + 4x - 9\right)e^{-x}}{16}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 28

```
DSolve[{3*y'''[x]+5*y''[x]+y'[x]-y[x]==0,{y[0]==0,y'[0]==1,y''[0]==-1}},y[x],x,IncludeSingular
```

$$y(x) \rightarrow \frac{1}{16}e^{-x}(4x + 9e^{4x/3} - 9)$$

## 8 Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

8.1	problem Exercise 21.3, page 231 . . . . .	230
8.2	problem Exercise 21.4, page 231 . . . . .	231
8.3	problem Exercise 21.5, page 231 . . . . .	232
8.4	problem Exercise 21.6, page 231 . . . . .	233
8.5	problem Exercise 21.7, page 231 . . . . .	234
8.6	problem Exercise 21.8, page 231 . . . . .	235
8.7	problem Exercise 21.9, page 231 . . . . .	236
8.8	problem Exercise 21.10, page 231 . . . . .	237
8.9	problem Exercise 21.11, page 231 . . . . .	238
8.10	problem Exercise 21.13, page 231 . . . . .	239
8.11	problem Exercise 21.14, page 231 . . . . .	240
8.12	problem Exercise 21.15, page 231 . . . . .	241
8.13	problem Exercise 21.16, page 231 . . . . .	242
8.14	problem Exercise 21.17, page 231 . . . . .	243
8.15	problem Exercise 21.19, page 231 . . . . .	244
8.16	problem Exercise 21.20, page 231 . . . . .	245
8.17	problem Exercise 21.21, page 231 . . . . .	246
8.18	problem Exercise 21.22, page 231 . . . . .	247
8.19	problem Exercise 21.24, page 231 . . . . .	248
8.20	problem Exercise 21.27, page 231 . . . . .	249
8.21	problem Exercise 21.28, page 231 . . . . .	250
8.22	problem Exercise 21.29, page 231 . . . . .	251
8.23	problem Exercise 21.31, page 231 . . . . .	252
8.24	problem Exercise 21.32, page 231 . . . . .	253
8.25	problem Exercise 21.33, page 231 . . . . .	254

## 8.1 problem Exercise 21.3, page 231

Internal problem ID [4606]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.3, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 3y' + 2y = 4$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x$2)+3*diff(y(x),x)+2*y(x)=4,y(x), singsol=all)
```

$$y(x) = -c_1 e^{-2x} + c_2 e^{-x} + 2$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 23

```
DSolve[y''[x]+3*y'[x]+2*y[x]==4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-2x} + c_2 e^{-x} + 2$$

## 8.2 problem Exercise 21.4, page 231

Internal problem ID [4607]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.4, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 3y' + 2y = 12e^x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$2)+3*diff(y(x),x)+2*y(x)=12*exp(x),y(x), singsol=all)
```

$$y(x) = -c_1e^{-2x} + 2e^x + c_2e^{-x}$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 27

```
DSolve[y''[x]+3*y'[x]+2*y[x]==12*Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(2e^{3x} + c_2e^x + c_1)$$



### 8.3 problem Exercise 21.5, page 231

Internal problem ID [4608]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.5, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 3y' + 2y = e^{ix}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$2)+3*diff(y(x),x)+2*y(x)=exp(I*x),y(x), singsol=all)
```

$$y(x) = \left( \left( \frac{1}{10} - \frac{3i}{10} \right) e^{ix+x} - c_1 e^{-x} + c_2 \right) e^{-x}$$

✓ Solution by Mathematica

Time used: 0.046 (sec). Leaf size: 37

```
DSolve[y''[x]+3*y'[x]+2*y[x]==Exp[I*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left( \frac{1}{10} - \frac{3i}{10} \right) e^{ix} + c_1 e^{-2x} + c_2 e^{-x}$$

## 8.4 problem Exercise 21.6, page 231

Internal problem ID [4609]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.6, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 3y' + 2y = \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)+3*diff(y(x),x)+2*y(x)=sin(x),y(x), singsol=all)
```

$$y(x) = -c_1 e^{-2x} - \frac{3 \cos(x)}{10} + \frac{\sin(x)}{10} + c_2 e^{-x}$$

### ✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 32

```
DSolve[y''[x]+3*y'[x]+2*y[x]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{10}(\sin(x) - 3 \cos(x) + 10e^{-2x}(c_2 e^x + c_1))$$

## 8.5 problem Exercise 21.7, page 231

Internal problem ID [4610]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.7, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 3y' + 2y = \cos(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)+3*diff(y(x),x)+2*y(x)=cos(x),y(x), singsol=all)
```

$$y(x) = -c_1 e^{-2x} + \frac{\cos(x)}{10} + \frac{3 \sin(x)}{10} + c_2 e^{-x}$$

### ✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 32

```
DSolve[y''[x]+3*y'[x]+2*y[x]==Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{10} (3 \sin(x) + \cos(x) + 10e^{-2x}(c_2 e^x + c_1))$$

## 8.6 problem Exercise 21.8, page 231

Internal problem ID [4611]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.8, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 3y' + 2y = 8 + 6e^x + 2\sin(x)$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$2)+3*diff(y(x),x)+2*y(x)=8+6*exp(x)+2*sin(x),y(x), singsol=all)
```

$$y(x) = -c_1 e^{-2x} + 4 + e^x - \frac{3 \cos(x)}{5} + \frac{\sin(x)}{5} + c_2 e^{-x}$$

### ✓ Solution by Mathematica

Time used: 0.165 (sec). Leaf size: 38

```
DSolve[y''[x]+3*y'[x]+2*y[x]==8+6*Exp[x]+2*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x + \frac{\sin(x)}{5} - \frac{3 \cos(x)}{5} + c_1 e^{-2x} + c_2 e^{-x} + 4$$

## 8.7 problem Exercise 21.9, page 231

Internal problem ID [4612]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.9, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + y' + y = x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 37

```
dsolve(diff(y(x),x$2)+diff(y(x),x)+y(x)=x^2,y(x), singsol=all)
```

$$y(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) c_2 + e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + x^2 - 2x$$

### ✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 54

```
DSolve[y''[x]+y'[x]+y[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/2} \left( e^{x/2} (x-2)x + c_2 \cos\left(\frac{\sqrt{3}x}{2}\right) + c_1 \sin\left(\frac{\sqrt{3}x}{2}\right) \right)$$

## 8.8 problem Exercise 21.10, page 231

Internal problem ID [4613]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.10, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 2y' - 8y = 9x e^x + 10 e^{-x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)-8*y(x)=9*x*exp(x)+10*exp(-x),y(x), singsol=all)
```

$$y(x) = e^{4x} c_2 + c_1 e^{-2x} - e^x x - 2 e^{-x}$$

### ✓ Solution by Mathematica

Time used: 0.145 (sec). Leaf size: 35

```
DSolve[y''[x]-2*y'[x]-8*y[x]==9*x*Exp[x]+10*Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} (-e^{3x} x - 2e^x + c_2 e^{6x} + c_1)$$

## 8.9 problem Exercise 21.11, page 231

Internal problem ID [4614]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.11, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' - 3y' = 2e^{2x} \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)=2*exp(2*x)*sin(x),y(x), singsol=all)
```

$$y(x) = \frac{e^{3x} c_1}{3} - \frac{e^{2x} \cos(x)}{5} - \frac{3e^{2x} \sin(x)}{5} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.245 (sec). Leaf size: 33

```
DSolve[y''[x]-3*y'[x]==2*Exp[2*x]*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{15} e^{2x} (-9 \sin(x) - 3 \cos(x) + 5c_1 e^x) + c_2$$

## 8.10 problem Exercise 21.13, page 231

Internal problem ID [4615]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.13, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' + y' = x^2 + 2x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)+diff(y(x),x)=x^2+2*x,y(x), singsol=all)
```

$$y(x) = \frac{x^3}{3} - c_1 e^{-x} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 24

```
DSolve[y''[x]+y'[x]==x^2+2*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{3} - c_1 e^{-x} + c_2$$



## 8.11 problem Exercise 21.14, page 231

Internal problem ID [4616]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.14, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' + y' = x + \sin(2x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(x),x$2)+diff(y(x),x)=x+sin(2*x),y(x), singsol=all)
```

$$y(x) = \frac{x^2}{2} - c_1 e^{-x} - \frac{\sin(2x)}{5} - \frac{\cos(2x)}{10} - x + c_2$$

### ✓ Solution by Mathematica

Time used: 0.359 (sec). Leaf size: 43

```
DSolve[y''[x]+y'[x]==x+Sin[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2}{2} - x - \frac{1}{5} \sin(2x) - \frac{1}{10} \cos(2x) - c_1 e^{-x} + c_2$$

## 8.12 problem Exercise 21.15, page 231

Internal problem ID [4617]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.15, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = 4x \sin(x)$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)+y(x)=4*x*sin(x),y(x), singsol=all)
```

$$y(x) = c_2 \sin(x) + c_1 \cos(x) - x(x \cos(x) - \sin(x))$$

### ✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 27

```
DSolve[y''[x]+y[x]==4*x*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left(-x^2 + \frac{1}{2} + c_1\right) \cos(x) + (x + c_2) \sin(x)$$

### 8.13 problem Exercise 21.16, page 231

Internal problem ID [4618]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.16, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y = x \sin(2x)$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(x),x$2)+4*y(x)=x*sin(2*x),y(x), singsol=all)
```

$$y(x) = \sin(2x)c_2 + c_1 \cos(2x) + \frac{\sin(2x)x}{16} - \frac{x^2 \cos(2x)}{8}$$

#### ✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 38

```
DSolve[y''[x]+4*y[x]==x*Sin[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{64}((-8x^2 + 1 + 64c_1) \cos(2x) + 4(x + 16c_2) \sin(2x))$$

## 8.14 problem Exercise 21.17, page 231

Internal problem ID [4619]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.17, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 2y' + y = x^2e^{-x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)+y(x)=x^2*exp(-x),y(x), singsol=all)
```

$$y(x) = c_2e^{-x} + e^{-x}c_1x + \frac{x^4e^{-x}}{12}$$

### ✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 27

```
DSolve[y''[x]+2*y'[x]+y[x]==x^2*Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{12}e^{-x}(x^4 + 12c_2x + 12c_1)$$

## 8.15 problem Exercise 21.19, page 231

Internal problem ID [4620]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.19, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 3y' + 2y = e^{-2x} + x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 40

```
dsolve(diff(y(x),x$2)+3*diff(y(x),x)+2*y(x)=exp(-2*x)+x^2,y(x), singsol=all)
```

$$y(x) = -c_1 e^{-2x} - \frac{3x}{2} + \frac{7}{4} - x e^{-2x} - e^{-2x} + \frac{x^2}{2} + c_2 e^{-x}$$

### ✓ Solution by Mathematica

Time used: 0.078 (sec). Leaf size: 41

```
DSolve[y''[x]+3*y'[x]+2*y[x]==Exp[-2*x]+x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}(2x^2 - 6x + 7) + e^{-2x}(-x - 1 + c_1) + c_2 e^{-x}$$

## 8.16 problem Exercise 21.20, page 231

Internal problem ID [4621]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.20, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 3y' + 2y = x e^{-x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)+2*y(x)=x*exp(-x),y(x), singsol=all)
```

$$y(x) = \left( c_1 e^x + \frac{5 e^{-2x}}{36} + \frac{x e^{-2x}}{6} + c_2 \right) e^x$$

### ✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 34

```
DSolve[y''[x]-3*y'[x]+2*y[x]==x*Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{36} e^{-x} (6x + 5) + c_1 e^x + c_2 e^{2x}$$

## 8.17 problem Exercise 21.21, page 231

Internal problem ID [4622]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.21, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + y' - 6y = x + e^{2x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
dsolve(diff(y(x),x$2)+diff(y(x),x)-6*y(x)=x+exp(2*x),y(x), singsol=all)
```

$$y(x) = e^{-3x}c_2 + c_1e^{2x} - \frac{1}{36} + \frac{(-1 + 5x)e^{2x}}{25} - \frac{x}{6}$$

### ✓ Solution by Mathematica

Time used: 0.085 (sec). Leaf size: 40

```
DSolve[y''[x]+y'[x]-6*y[x]==x+Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{36}(-6x - 1) + c_1e^{-3x} + e^{2x}\left(\frac{x}{5} - \frac{1}{25} + c_2\right)$$

## 8.18 problem Exercise 21.22, page 231

Internal problem ID [4623]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.22, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \sin(x) + e^{-x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)+y(x)=sin(x)+exp(-x),y(x), singsol=all)
```

$$y(x) = c_2 \sin(x) + c_1 \cos(x) + \frac{e^{-x}}{2} - \frac{x \cos(x)}{2}$$

### ✓ Solution by Mathematica

Time used: 0.337 (sec). Leaf size: 36

```
DSolve[y''[x]+y[x]==Sin[x]+Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}(2e^{-x} + \sin(x) - 2x \cos(x) + 4c_1 \cos(x) + 4c_2 \sin(x))$$



## 8.19 problem Exercise 21.24, page 231

Internal problem ID [4624]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.24, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \sin(x)^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$2)+y(x)=sin(x)^2,y(x), singsol=all)
```

$$y(x) = c_2 \sin(x) + c_1 \cos(x) + \frac{1}{2} + \frac{\cos(2x)}{6}$$

### ✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 27

```
DSolve[y''[x]+y[x]==Sin[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{6}(\cos(2x) + 6c_1 \cos(x) + 6c_2 \sin(x) + 3)$$

## 8.20 problem Exercise 21.27, page 231

Internal problem ID [4625]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.27, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \sin(x) \sin(2x)$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)+y(x)=sin(2*x)*sin(x),y(x), singsol=all)
```

$$y(x) = c_2 \sin(x) + c_1 \cos(x) + \frac{\sin(x) (-\cos(x) \sin(x) + x)}{4}$$

### ✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 33

```
DSolve[y''[x]+y[x]==Sin[2*x]*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{16}(\cos(3x) + (-1 + 16c_1) \cos(x) + 4(x + 4c_2) \sin(x))$$

## 8.21 problem Exercise 21.28, page 231

Internal problem ID [4626]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.28, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 5y' - 6y = e^{3x}$$

With initial conditions

$$[y(0) = 2, y'(0) = 1]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve([diff(y(x),x$2)-5*diff(y(x),x)-6*y(x)=exp(3*x),y(0) = 2, D(y)(0) = 1],y(x), singsol=a
```

$$y(x) = \frac{45 e^{-x}}{28} + \frac{10 e^{6x}}{21} - \frac{e^{3x}}{12}$$

### ✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 30

```
DSolve[{y'[x]-5*y'[x]-6*y[x]==Exp[3*x],{y[0]==2,y'[0]==1}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{1}{84} e^{-x} (-7e^{4x} + 40e^{7x} + 135)$$

## 8.22 problem Exercise 21.29, page 231

Internal problem ID [4627]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.29, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - y' - 2y = 5 \sin(x)$$

With initial conditions

$$[y(0) = 1, y'(0) = -1]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve([diff(y(x),x$2)-diff(y(x),x)-2*y(x)=5*sin(x),y(0) = 1, D(y)(0) = -1],y(x), singsol=all
```

$$y(x) = \frac{e^{-x}}{6} + \frac{e^{2x}}{3} + \frac{\cos(x)}{2} - \frac{3 \sin(x)}{2}$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 30

```
DSolve[{y''[x]-y'[x]-2*y[x]==5*Sin[x],{y[0]==1,y'[0]==-1}},y[x],x,IncludeSingularSolutions -
```

$$y(x) \rightarrow \frac{1}{6}(e^{-x} + 2e^{2x} - 9 \sin(x) + 3 \cos(x))$$

## 8.23 problem Exercise 21.31, page 231

Internal problem ID [4628]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.31, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = 8 \cos(x)$$

With initial conditions

$$\left[ y\left(\frac{\pi}{2}\right) = -1, y'\left(\frac{\pi}{2}\right) = 1 \right]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)+9*y(x)=8*cos(x),y(1/2*Pi) = -1, D(y)(1/2*Pi) = 1],y(x), singsol=all)
```

$$y(x) = \sin(3x) + \frac{2 \cos(3x)}{3} + \cos(x)$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 20

```
DSolve[{y''[x]+9*y[x]==8*Cos[x],{y[Pi/2]==-1,y'[Pi/2]==1}},y[x],x,IncludeSingularSolutions -
```

$$y(x) \rightarrow \sin(3x) + \cos(x) + \frac{2}{3} \cos(3x)$$

## 8.24 problem Exercise 21.32, page 231

Internal problem ID [4629]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.32, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 5y' + 6y = e^x(2x - 3)$$

With initial conditions

$$[y(0) = 1, y'(0) = 3]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve([diff(y(x),x$2)-5*diff(y(x),x)+6*y(x)=exp(x)*(2*x-3),y(0) = 1, D(y)(0) = 3],y(x), sin
```

$$y(x) = e^{2x} + e^x x$$

### ✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 35

```
DSolve[{y''[x]-5*y'[x]-6*y[x]==Exp[x]*(2*x-3)},{y[0]==1,y'[0]==3}],y[x],x,IncludeSingularSolu
```

$$y(x) \rightarrow \frac{1}{175}e^{-x}(-7e^{2x}(5x - 9) + 87e^{7x} + 25)$$

## 8.25 problem Exercise 21.33, page 231

Internal problem ID [4630]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 21. Undetermined Coefficients

**Problem number:** Exercise 21.33, page 231.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 3y' + 2y = e^{-x}$$

With initial conditions

$$[y(0) = 1, y'(0) = -1]$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 21

```
dsolve([diff(y(x),x$2)-3*diff(y(x),x)+2*y(x)=exp(-x),y(0) = 1, D(y)(0) = -1],y(x), singsol=a
```

$$y(x) = -\frac{5e^{2x}}{3} + \frac{5e^x}{2} + \frac{e^{-x}}{6}$$

### ✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 31

```
DSolve[{y'[x]-3*y'[x]+2*y[x]==Exp[-x],{y[0]==1,y'[0]==-1}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{e^{-x}}{6} + \frac{5e^x}{2} - \frac{5e^{2x}}{3}$$

## 9 Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

9.1	problem Exercise 22.1, page 240 . . . . .	256
9.2	problem Exercise 22.2, page 240 . . . . .	257
9.3	problem Exercise 22.3, page 240 . . . . .	258
9.4	problem Exercise 22.4, page 240 . . . . .	259
9.5	problem Exercise 22.5, page 240 . . . . .	260
9.6	problem Exercise 22.6, page 240 . . . . .	261
9.7	problem Exercise 22.7, page 240 . . . . .	262
9.8	problem Exercise 22.8, page 240 . . . . .	263
9.9	problem Exercise 22.9, page 240 . . . . .	264
9.10	problem Exercise 22.10, page 240 . . . . .	265
9.11	problem Exercise 22.11, page 240 . . . . .	266
9.12	problem Exercise 22.12, page 240 . . . . .	267
9.13	problem Exercise 22.13, page 240 . . . . .	268
9.14	problem Exercise 22.14, page 240 . . . . .	269
9.15	problem Exercise 22.15, page 240 . . . . .	270
9.16	problem Exercise 22, problem 16, page 240 . . . . .	271
9.17	problem Exercise 22, problem 17, page 240 . . . . .	272
9.18	problem Exercise 22, problem 18, page 240 . . . . .	273
9.19	problem Exercise 22, problem 19, page 240 . . . . .	274
9.20	problem Exercise 22, problem 20, page 240 . . . . .	275



## 9.1 problem Exercise 22.1, page 240

Internal problem ID [4631]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22.1, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \sec(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)+y(x)=sec(x),y(x), singsol=all)
```

$$y(x) = c_2 \sin(x) + c_1 \cos(x) + x \sin(x) - \ln(\sec(x)) \cos(x)$$

### ✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 22

```
DSolve[y''[x]+y[x]==Sec[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x + c_2) \sin(x) + \cos(x)(\log(\cos(x)) + c_1)$$

## 9.2 problem Exercise 22.2, page 240

Internal problem ID [4632]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22.2, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \cot(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)+y(x)=cot(x),y(x), singsol=all)
```

$$y(x) = c_2 \sin(x) + c_1 \cos(x) + \sin(x) \ln(\csc(x) - \cot(x))$$

### ✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 33

```
DSolve[y''[x]+y[x]==Cot[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(x) + \sin(x) \left( \log\left(\sin\left(\frac{x}{2}\right)\right) - \log\left(\cos\left(\frac{x}{2}\right)\right) \right) + c_2$$

### 9.3 problem Exercise 22.3, page 240

Internal problem ID [4633]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22.3, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \sec(x)^2$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)+y(x)=sec(x)^2,y(x), singsol=all)
```

$$y(x) = c_2 \sin(x) + c_1 \cos(x) + \ln(\sec(x) + \tan(x)) \sin(x) - 1$$

#### ✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 28

```
DSolve[y''[x]+y[x]==Sec[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 \sin(x) \operatorname{arctanh}\left(\tan\left(\frac{x}{2}\right)\right) + c_1 \cos(x) + c_2 \sin(x) - 1$$

## 9.4 problem Exercise 22.4, page 240

Internal problem ID [4634]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22.4, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - y = \sin(x)^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$2)-y(x)=sin(x)^2,y(x), singsol=all)
```

$$y(x) = e^x c_2 + c_1 e^{-x} + \frac{\cos(x)^2}{5} - \frac{3}{5}$$

### ✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 30

```
DSolve[y''[x]-y[x]==Sin[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{10}(\cos(2x) - 5) + c_1 e^x + c_2 e^{-x}$$

## 9.5 problem Exercise 22.5, page 240

Internal problem ID [4635]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22.5, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \sin(x)^2$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$2)+y(x)=sin(x)^2,y(x), singsol=all)
```

$$y(x) = c_2 \sin(x) + c_1 \cos(x) + \frac{1}{2} + \frac{\cos(2x)}{6}$$

### ✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 27

```
DSolve[y''[x]+y[x]==Sin[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{6}(\cos(2x) + 6c_1 \cos(x) + 6c_2 \sin(x) + 3)$$

## 9.6 problem Exercise 22.6, page 240

Internal problem ID [4636]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22.6, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 3y' + 2y = 12e^x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$2)+3*diff(y(x),x)+2*y(x)=12*exp(x),y(x), singsol=all)
```

$$y(x) = -c_1e^{-2x} + 2e^x + c_2e^{-x}$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 27

```
DSolve[y''[x]+3*y'[x]+2*y[x]==12*Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(2e^{3x} + c_2e^x + c_1)$$

## 9.7 problem Exercise 22.7, page 240

Internal problem ID [4637]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22.7, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 2y' + y = x^2e^{-x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)+y(x)=x^2*exp(-x),y(x), singsol=all)
```

$$y(x) = c_2e^{-x} + e^{-x}c_1x + \frac{x^4e^{-x}}{12}$$

### ✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 27

```
DSolve[y''[x]+2*y'[x]+y[x]==x^2*Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{12}e^{-x}(x^4 + 12c_2x + 12c_1)$$

## 9.8 problem Exercise 22.8, page 240

Internal problem ID [4638]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22.8, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = 4x \sin(x)$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)+y(x)=4*x*sin(x),y(x), singsol=all)
```

$$y(x) = c_2 \sin(x) + c_1 \cos(x) - x(x \cos(x) - \sin(x))$$

### ✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 27

```
DSolve[y''[x]+y[x]==4*x*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left(-x^2 + \frac{1}{2} + c_1\right) \cos(x) + (x + c_2) \sin(x)$$



## 9.9 problem Exercise 22.9, page 240

Internal problem ID [4639]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22.9, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 2y' + y = e^{-x} \ln(x)$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 33

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)+y(x)=exp(-x)*ln(x),y(x), singsol=all)
```

$$y(x) = c_2 e^{-x} + e^{-x} c_1 x + \frac{x^2(2 \ln(x) - 3) e^{-x}}{4}$$

### ✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 36

```
DSolve[y''[x]+2*y'[x]+y[x]==Exp[-x]*Log[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4} e^{-x} (-3x^2 + 2x^2 \log(x) + 4c_2 x + 4c_1)$$

## 9.10 problem Exercise 22.10, page 240

Internal problem ID [4640]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22.10, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \csc(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)+y(x)=csc(x),y(x), singsol=all)
```

$$y(x) = c_2 \sin(x) + c_1 \cos(x) - \ln(\csc(x)) \sin(x) - x \cos(x)$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 24

```
DSolve[y''[x]+y[x]==Csc[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (-x + c_1) \cos(x) + \sin(x)(\log(\sin(x)) + c_2)$$

## 9.11 problem Exercise 22.11, page 240

Internal problem ID [4641]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22.11, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \tan(x)^2$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)+y(x)=tan(x)^2,y(x), singsol=all)
```

$$y(x) = c_2 \sin(x) + c_1 \cos(x) - 2 + \ln(\sec(x) + \tan(x)) \sin(x)$$

### ✓ Solution by Mathematica

Time used: 0.109 (sec). Leaf size: 23

```
DSolve[y''[x]+y[x]==Tan[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sin(x) \operatorname{arctanh}(\sin(x)) + c_1 \cos(x) + c_2 \sin(x) - 2$$

## 9.12 problem Exercise 22.12, page 240

Internal problem ID [4642]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22.12, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 2y' + y = \frac{e^{-x}}{x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)+y(x)=exp(-x)/x,y(x), singsol=all)
```

$$y(x) = c_2 e^{-x} + e^{-x} c_1 x + x(\ln(x) - 1) e^{-x}$$

### ✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 24

```
DSolve[y''[x]+2*y'[x]+y[x]==Exp[-x]/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(x \log(x) + (-1 + c_2)x + c_1)$$

## 9.13 problem Exercise 22.13, page 240

Internal problem ID [4643]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22.13, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \csc(x) \sec(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$2)+y(x)=sec(x)*csc(x),y(x), singsol=all)
```

$$y(x) = c_2 \sin(x) + c_1 \cos(x) + \sin(x) \ln(\csc(x) - \cot(x)) - \ln(\sec(x) + \tan(x)) \cos(x)$$

### ✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 30

```
DSolve[y''[x]+y[x]==Sec[x]*Csc[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sin(x) \operatorname{arctanh}(\cos(x)) + c_1 \cos(x) + c_2 \sin(x) + \cos(x) \left(-\operatorname{coth}^{-1}(\sin(x))\right)$$

## 9.14 problem Exercise 22.14, page 240

Internal problem ID [4644]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22.14, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 2y' + y = e^x \ln(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x)=exp(x)*ln(x),y(x), singsol=all)
```

$$y(x) = e^x c_2 + e^x c_1 x + \frac{e^x x^2 (2 \ln(x) - 3)}{4}$$

### ✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 34

```
DSolve[y''[x]-2*y'[x]+y[x]==Exp[x]*Log[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4} e^x (-3x^2 + 2x^2 \log(x) + 4c_2 x + 4c_1)$$

## 9.15 problem Exercise 22.15, page 240

Internal problem ID [4645]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22.15, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 3y' + 2y = \cos(e^{-x})$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)+2*y(x)=cos(exp(-x)),y(x), singsol=all)
```

$$y(x) = (c_1 e^x - e^x - e^x \cos(e^{-x}) + c_2) e^x$$

### ✓ Solution by Mathematica

Time used: 0.082 (sec). Leaf size: 29

```
DSolve[y''[x]-3*y'[x]+2*y[x]==Cos[Exp[-x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(-e^x \cos(e^{-x}) + c_2 e^x + c_1)$$

## 9.16 problem Exercise 22, problem 16, page 240

Internal problem ID [4646]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22, problem 16, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2y'' - xy' + y = x$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 20

```
dsolve(x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=x,y(x), singsol=all)
```

$$y(x) = c_2x + x \ln(x) c_1 + \frac{\ln(x)^2 x}{2}$$

### ✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 25

```
DSolve[x^2*y''[x]-x*y'[x]+y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}x(\log^2(x) + 2c_2 \log(x) + 2c_1)$$



## 9.17 problem Exercise 22, problem 17, page 240

Internal problem ID [4647]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22, problem 17, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$y'' - \frac{2y'}{x} + \frac{2y}{x^2} = x \ln(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)-2/x*diff(y(x),x)+2/x^2*y(x)=x*ln(x),y(x), singsol=all)
```

$$y(x) = c_1x + c_2x^2 + \frac{x^3(2 \ln(x) - 3)}{4}$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 32

```
DSolve[y''[x]-2/x*y'[x]+2/x^2*y[x]==x*Log[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}x(-3x^2 + 2x^2 \log(x) + 4c_2x + 4c_1)$$

## 9.18 problem Exercise 22, problem 18, page 240

Internal problem ID [4648]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22, problem 18, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2y'' + xy' - 4y = x^3$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-4*y(x)=x^3,y(x), singsol=all)
```

$$y(x) = \frac{c_2}{x^2} + c_1x^2 + \frac{x^3}{5}$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 25

```
DSolve[x^2*y''[x]+x*y'[x]-4*y[x]==x^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{5} + c_2x^2 + \frac{c_1}{x^2}$$

## 9.19 problem Exercise 22, problem 19, page 240

Internal problem ID [4649]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22, problem 19, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$x^2y'' + xy' - y = x^2e^{-x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=x^2*exp(-x),y(x), singsol=all)
```

$$y(x) = \frac{c_1}{x} + c_2x + \frac{e^{-x}(x+1)}{x}$$

### ✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 27

```
DSolve[x^2*y''[x]+x*y'[x]-y[x]==x^2*Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2x^2 + e^{-x}(x+1) + c_1}{x}$$

## 9.20 problem Exercise 22, problem 20, page 240

Internal problem ID [4650]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 4. Higher order linear differential equations. Lesson 22. Variation of Parameters

**Problem number:** Exercise 22, problem 20, page 240.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$2x^2y'' + 3xy' - y = \frac{1}{x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(2*x^2*diff(y(x),x$2)+3*x*diff(y(x),x)-y(x)=1/x,y(x), singsol=all)
```

$$y(x) = \frac{c_1}{x} + c_2\sqrt{x} - \frac{3\ln(x) + 2}{9x}$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 31

```
DSolve[2*x^2*y''[x]+3*x*y'[x]-y[x]==1/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{9c_2x^{3/2} - 3\log(x) - 2 + 9c_1}{9x}$$

## 10 Chapter 8. Special second order equations.

### Lesson 35. Independent variable $x$ absent

10.1	problem Exercise 35.1, page 504 . . . . .	277
10.2	problem Exercise 35.2, page 504 . . . . .	278
10.3	problem Exercise 35.3, page 504 . . . . .	279
10.4	problem Exercise 35.4, page 504 . . . . .	280
10.5	problem Exercise 35.5, page 504 . . . . .	281
10.6	problem Exercise 35.6, page 504 . . . . .	282
10.7	problem Exercise 35.7, page 504 . . . . .	284
10.8	problem Exercise 35.8, page 504 . . . . .	285
10.9	problem Exercise 35.9, page 504 . . . . .	286
10.10	problem Exercise 35.10, page 504 . . . . .	287
10.11	problem Exercise 35.11, page 504 . . . . .	288
10.12	problem Exercise 35.12, page 504 . . . . .	290
10.13	problem Exercise 35.13, page 504 . . . . .	291
10.14	problem Exercise 35.14, page 504 . . . . .	292
10.15	problem Exercise 35.15, page 504 . . . . .	293
10.16	problem Exercise 35.16, page 504 . . . . .	294
10.17	problem Exercise 35.17, page 504 . . . . .	295
10.18	problem Exercise 35.18, page 504 . . . . .	296
10.19	problem Exercise 35.19, page 504 . . . . .	297
10.20	problem Exercise 35.20, page 504 . . . . .	298
10.21	problem Exercise 35.21, page 504 . . . . .	299
10.22	problem Exercise 35.23(a), page 504 . . . . .	300
10.23	problem Exercise 35.23(b), page 504 . . . . .	301
10.24	problem Exercise 35.23(c), page 504 . . . . .	302

## 10.1 problem Exercise 35.1, page 504

Internal problem ID [4651]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.1, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear], _`

$$y'' - 2yy' = 0$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 16

```
dsolve(diff(y(x),x$2)=2*y(x)*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = \frac{\tan\left(\frac{c_2+x}{c_1}\right)}{c_1}$$

### ✓ Solution by Mathematica

Time used: 9.872 (sec). Leaf size: 24

```
DSolve[y''[x]==2*y[x]*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt{c_1} \tan(\sqrt{c_1}(x + c_2))$$

## 10.2 problem Exercise 35.2, page 504

Internal problem ID [4652]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.2, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$y^3 y'' = k$$

### ✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 70

```
dsolve(y(x)^3*diff(y(x),x$2)=k,y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{c_1 (c_1^2 c_2^2 + 2c_1^2 c_2 x + c_1^2 x^2 + k)}}{c_1}$$

$$y(x) = -\frac{\sqrt{c_1 (c_1^2 c_2^2 + 2c_1^2 c_2 x + c_1^2 x^2 + k)}}{c_1}$$

### ✓ Solution by Mathematica

Time used: 2.878 (sec). Leaf size: 63

```
DSolve[y[x]^3*y'[x]==k,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{k + c_1^2(x + c_2)^2}}{\sqrt{c_1}}$$

$$y(x) \rightarrow \frac{\sqrt{k + c_1^2(x + c_2)^2}}{\sqrt{c_1}}$$

$$y(x) \rightarrow \text{Indeterminate}$$

### 10.3 problem Exercise 35.3, page 504

Internal problem ID [4653]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable  $x$  absent

**Problem number:** Exercise 35.3, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$yy'' - y'^2 = -1$$

✓ Solution by Maple

Time used: 0.14 (sec). Leaf size: 79

```
dsolve(y(x)*diff(y(x),x$2)=(diff(y(x),x))^2-1,y(x), singsol=all)
```

$$y(x) = \frac{c_1 \left( e^{-\frac{2c_2}{c_1}} e^{-\frac{2x}{c_1}} - 1 \right) e^{\frac{c_2}{c_1}} e^{\frac{x}{c_1}}}{2}$$

$$y(x) = \frac{c_1 \left( e^{\frac{2c_2}{c_1}} e^{\frac{2x}{c_1}} - 1 \right) e^{-\frac{c_2}{c_1}} e^{-\frac{x}{c_1}}}{2}$$

✓ Solution by Mathematica

Time used: 60.201 (sec). Leaf size: 85

```
DSolve[y[x]*y'[x]==(y'[x])^2-1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{ie^{-c_1} \tanh(e^{c_1}(x+c_2))}{\sqrt{-\operatorname{sech}^2(e^{c_1}(x+c_2))}}$$

$$y(x) \rightarrow \frac{ie^{-c_1} \tanh(e^{c_1}(x+c_2))}{\sqrt{-\operatorname{sech}^2(e^{c_1}(x+c_2))}}$$



## 10.4 problem Exercise 35.4, page 504

Internal problem ID [4654]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.4, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^2 y'' + xy' = 1$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve(x^2*diff(y(x),x$2)+x*(diff(y(x),x))=1,y(x), singsol=all)
```

$$y(x) = \frac{\ln(x)^2}{2} + c_1 \ln(x) + c_2$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 21

```
DSolve[x^2*y''[x]+x*y'[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\log^2(x)}{2} + c_1 \log(x) + c_2$$

## 10.5 problem Exercise 35.5, page 504

Internal problem ID [4655]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.5, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' - y' = x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x*diff(y(x),x$2)-diff(y(x),x)=x^2,y(x), singsol=all)
```

$$y(x) = \frac{1}{3}x^3 + \frac{1}{2}c_1x^2 + c_2$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 24

```
DSolve[x*y''[x]-y'[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{3} + \frac{c_1x^2}{2} + c_2$$

## 10.6 problem Exercise 35.6, page 504

Internal problem ID [4656]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.6, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible]`

$$(1 + y)y'' - 3y'^2 = 0$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 59

```
dsolve((y(x)+1)*diff(y(x),x$2)=3*(diff(y(x),x))^2,y(x), singsol=all)
```

$$y(x) = -1$$

$$y(x) = -\frac{\sqrt{-2c_1x - 2c_2} - 1}{\sqrt{-2c_1x - 2c_2}}$$

$$y(x) = -\frac{\sqrt{-2c_1x - 2c_2} + 1}{\sqrt{-2c_1x - 2c_2}}$$

✓ Solution by Mathematica

Time used: 1.485 (sec). Leaf size: 107

```
DSolve[(y[x]+1)*y'[x]==3*(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2c_1x + \sqrt{2}\sqrt{-c_1(x+c_2)} + 2c_2c_1}{2c_1(x+c_2)}$$

$$y(x) \rightarrow \frac{-2c_1x + \sqrt{2}\sqrt{-c_1(x+c_2)} - 2c_2c_1}{2c_1(x+c_2)}$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow \text{Indeterminate}$$

## 10.7 problem Exercise 35.7, page 504

Internal problem ID [4657]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.7, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$r'' + \frac{k}{r^2} = 0$$

### ✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 369

```
dsolve(diff(r(t),t$2)=-k/(r(t)^2),r(t), singsol=all)
```

$$r(t) = c_1 \left( c_1^2 k^2 - 2k c_1 e^{\text{RootOf}\left(\text{csgn}\left(\frac{1}{c_1}\right) c_1^4 k^2 + 2\_Z c_1^3 k e^{-Z} - \text{csgn}\left(\frac{1}{c_1}\right) e^{2-Z} c_1^2 - 2 \text{csgn}\left(\frac{1}{c_1}\right) e^{-Z} c_2 - 2 \text{csgn}\left(\frac{1}{c_1}\right) e^{-Z} t\right)} + e^{2 \text{RootOf}\left(\text{csgn}\left(\frac{1}{c_1}\right) e^{2-Z} c_1^2 - 2 \text{csgn}\left(\frac{1}{c_1}\right) e^{-Z} c_2 - 2 \text{csgn}\left(\frac{1}{c_1}\right) e^{-Z} t\right)} \right)$$

$$r(t) = c_1 \left( c_1^2 k^2 - 2k c_1 e^{\text{RootOf}\left(\text{csgn}\left(\frac{1}{c_1}\right) c_1^4 k^2 + 2\_Z c_1^3 k e^{-Z} - \text{csgn}\left(\frac{1}{c_1}\right) e^{2-Z} c_1^2 + 2 \text{csgn}\left(\frac{1}{c_1}\right) e^{-Z} c_2 + 2 \text{csgn}\left(\frac{1}{c_1}\right) e^{-Z} t\right)} + e^{2 \text{RootOf}\left(\text{csgn}\left(\frac{1}{c_1}\right) e^{2-Z} c_1^2 + 2 \text{csgn}\left(\frac{1}{c_1}\right) e^{-Z} c_2 + 2 \text{csgn}\left(\frac{1}{c_1}\right) e^{-Z} t\right)} \right)$$

### ✓ Solution by Mathematica

Time used: 0.169 (sec). Leaf size: 65

```
DSolve[r''[t]==-k/(r[t]^2),r[t],t,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ \left( \frac{r(t) \sqrt{\frac{2k}{r(t)} + c_1}}{c_1} - \frac{2k \arctanh\left(\frac{\sqrt{\frac{2k}{r(t)} + c_1}}{\sqrt{c_1}}\right)}{c_1^{3/2}} \right)^2 = (t + c_2)^2, r(t) \right]$$

## 10.8 problem Exercise 35.8, page 504

Internal problem ID [4658]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.8, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]`

$$y'' - \frac{3y^2k}{2} = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 15

```
dsolve(diff(y(x),x$2)=3/2*k*y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{4 \operatorname{WeierstrassP}(x + c_1, 0, c_2)}{k}$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y''[x]==3/2*(k*y[x]^2),y[x],x,IncludeSingularSolutions -> True]
```

Not solved

## 10.9 problem Exercise 35.9, page 504

Internal problem ID [4659]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.9, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$y'' - 2ky^3 = 0$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$2)=2*k*y(x)^3,y(x), singsol=all)
```

$$y(x) = c_2 \operatorname{JacobiSN}\left(\left(\sqrt{-k}x + c_1\right) c_2, i\right)$$

✓ Solution by Mathematica

Time used: 61.304 (sec). Leaf size: 115

```
DSolve[y''[x]==2*k*y[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\operatorname{isn}\left(\left(-1\right)^{3/4}\sqrt{\sqrt{k}\sqrt{c_1}(x+c_2)^2}-1\right)}{\sqrt{\frac{i\sqrt{k}}{\sqrt{c_1}}}}$$

$$y(x) \rightarrow \frac{\operatorname{isn}\left(\left(-1\right)^{3/4}\sqrt{\sqrt{k}\sqrt{c_1}(x+c_2)^2}-1\right)}{\sqrt{\frac{i\sqrt{k}}{\sqrt{c_1}}}}$$

## 10.10 problem Exercise 35.10, page 504

Internal problem ID [4660]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.10, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear], [`

$$yy'' + y'^2 - y' = 0$$

### ✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 37

```
dsolve(y(x)*diff(y(x),x$2)+(diff(y(x),x))^2-diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = -c_1 \left( \text{LambertW} \left( -\frac{e^{-1} e^{-\frac{c_2}{c_1}} e^{-\frac{x}{c_1}}}{c_1} \right) + 1 \right)$$

### ✓ Solution by Mathematica

Time used: 60.084 (sec). Leaf size: 32

```
DSolve[y[x]*y'[x]+(y'[x])^2-y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -c_1 \left( 1 + W \left( -\frac{e^{-\frac{x+c_1+c_2}{c_1}}}{c_1} \right) \right)$$



## 10.11 problem Exercise 35.11, page 504

Internal problem ID [4661]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.11, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$r'' - \frac{h^2}{r^3} + \frac{k}{r^2} = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 441

```
dsolve(diff(r(t),t$2)= h^2/r(t)^3-k/r(t)^2,r(t), singsol=all)
```

$$r(t) = \frac{c_1 \left( c_1^2 k^2 - 2k c_1 e^{\text{RootOf}\left(\text{csgn}\left(\frac{1}{c_1}\right) c_1^4 k^2 + 2_Z c_1^3 k e^{-Z} - \text{csgn}\left(\frac{1}{c_1}\right) e^{2-Z} c_1^2 + \text{csgn}\left(\frac{1}{c_1}\right) c_1^2 h^2 - 2 \text{csgn}\left(\frac{1}{c_1}\right) e^{-Z} c_2 - 2 \text{csgn}\left(\frac{1}{c_1}\right) e^{-Z} t\right)} + e^{2-Z} \right)}{\dots}$$

$$r(t) = \frac{c_1 \left( c_1^2 k^2 - 2k c_1 e^{\text{RootOf}\left(\text{csgn}\left(\frac{1}{c_1}\right) c_1^4 k^2 + 2_Z c_1^3 k e^{-Z} - \text{csgn}\left(\frac{1}{c_1}\right) e^{2-Z} c_1^2 + \text{csgn}\left(\frac{1}{c_1}\right) c_1^2 h^2 + 2 \text{csgn}\left(\frac{1}{c_1}\right) e^{-Z} c_2 + 2 \text{csgn}\left(\frac{1}{c_1}\right) e^{-Z} t\right)} + e^{2-Z} \right)}{\dots}$$

✓ Solution by Mathematica

Time used: 1.099 (sec). Leaf size: 130

```
DSolve[r''[t]==h^2/r[t]^3-k/r[t]^2,r[t],t,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ \frac{\left( \sqrt{c_1}(-h^2 + r(t)(2k + c_1 r(t))) - k\sqrt{-h^2 + r(t)(2k + c_1 r(t))} \operatorname{arctanh} \left( \frac{k + c_1 r(t)}{\sqrt{c_1} \sqrt{-h^2 + r(t)(2k + c_1 r(t))}} \right) \right)^2}{c_1^3 r(t)^2 \left( -\frac{h^2}{r(t)^2} + \frac{2k}{r(t)} + c_1 \right)} + c_2)^2, r(t) \right]$$

## 10.12 problem Exercise 35.12, page 504

Internal problem ID [4662]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.12, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]`,

$$yy'' + y'^3 - y'^2 = 0$$

### ✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 44

```
dsolve(y(x)*diff(y(x),x$2)+(diff(y(x),x))^3-diff(y(x),x)^2=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1$$

$$y(x) = e^{-\frac{c_1 \operatorname{LambertW}\left(\frac{e^{\frac{c_2}{c_1}} e^{\frac{x}{c_1}}}{c_1}\right) - c_2 - x}{c_1}}$$

### ✓ Solution by Mathematica

Time used: 22.229 (sec). Leaf size: 32

```
DSolve[y[x]*y'[x]+(y'[x])^3-(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{c_1 W\left(e^{e^{-c_1}(x - e^{c_1}c_1 + c_2)}\right)}$$

## 10.13 problem Exercise 35.13, page 504

Internal problem ID [4663]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.13, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible]`

$$yy'' - 3y'^2 = 0$$

### ✓ Solution by Maple

Time used: 0.11 (sec). Leaf size: 33

```
dsolve(y(x)*diff(y(x),x$2)-3*(diff(y(x),x))^2=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = \frac{1}{\sqrt{-2c_1x - 2c_2}}$$

$$y(x) = -\frac{1}{\sqrt{-2c_1x - 2c_2}}$$

### ✓ Solution by Mathematica

Time used: 0.106 (sec). Leaf size: 14

```
DSolve[y[x]*y'[x]-(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 e^{c_1 x}$$

## 10.14 problem Exercise 35.14, page 504

Internal problem ID [4664]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.14, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]`

$$(x^2 + 1)y'' + y'^2 = -1$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 29

```
dsolve((1+x^2)*diff(y(x),x$2)+(diff(y(x),x))^2+1=0,y(x), singsol=all)
```

$$y(x) = \frac{x}{c_1} - \frac{(-c_1^2 - 1) \ln(c_1 x - 1)}{c_1^2} + c_2$$

### ✓ Solution by Mathematica

Time used: 7.091 (sec). Leaf size: 33

```
DSolve[(1+x^2)*y'[x]+(y'[x])^2+1==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x \cot(c_1) + \csc^2(c_1) \log(-x \sin(c_1) - \cos(c_1)) + c_2$$

## 10.15 problem Exercise 35.15, page 504

Internal problem ID [4665]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable  $x$  absent

**Problem number:** Exercise 35.15, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$(x^2 + 1)y'' + 2x(y' + 1) = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve((1+x^2)*diff(y(x),x$2)+2*x*(diff(y(x),x)+1)=0,y(x), singsol=all)
```

$$y(x) = -x + (c_1 + 1) \arctan(x) + c_2$$

### ✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 18

```
DSolve[(1+x^2)*y'[x]+2*x*(y'[x]+1)==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (1 + c_1) \arctan(x) - x + c_2$$

## 10.16 problem Exercise 35.16, page 504

Internal problem ID [4666]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.16, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible]`

$$(1 + y)y'' - 3y'^2 = 0$$

With initial conditions

$$\left[ y(1) = 0, y'(1) = -\frac{1}{2} \right]$$

✓ Solution by Maple

Time used: 0.375 (sec). Leaf size: 15

```
dsolve([(y(x)+1)*diff(y(x),x$2)=3*(diff(y(x),x))^2,y(1) = 0, D(y)(1) = -1/2],y(x), singsol=a
```

$$y(x) = \frac{-x + \sqrt{x}}{x}$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{(y[x]+1)*y'[x]==3*(y'[x])^3,{y[1]==0,y'[0]==-1/2}},y[x],x,IncludeSingularSolutions
```

{}

## 10.17 problem Exercise 35.17, page 504

Internal problem ID [4667]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.17, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear], [`

$$y'' - y'e^y = 0$$

With initial conditions

$$[y(3) = 0, y'(3) = 1]$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 12

```
dsolve([diff(y(x),x$2)=diff(y(x),x)*exp(y(x)),y(3) = 0, D(y)(3) = 1],y(x), singsol=all)
```

$$y(x) = -\ln(-x + 4)$$

### ✓ Solution by Mathematica

Time used: 7.673 (sec). Leaf size: 13

```
DSolve[{y'[x]==y'[x]*Exp[y[x]],{y[3]==0,y'[3]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\log(4 - x)$$



## 10.18 problem Exercise 35.18, page 504

Internal problem ID [4668]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.18, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear], _`

$$y'' - 2yy' = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 2]$$

✓ Solution by Maple

Time used: 0.109 (sec). Leaf size: 10

```
dsolve([diff(y(x),x$2)=2*y(x)*diff(y(x),x),y(0) = 1, D(y)(0) = 2],y(x), singsol=all)
```

$$y(x) = \tan\left(x + \frac{\pi}{4}\right)$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y'[x]==2*y[x]*y'[x],{y[0]==1,y'[0]==2}},y[x],x,IncludeSingularSolutions -> True]
```

{}

## 10.19 problem Exercise 35.19, page 504

Internal problem ID [4669]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.19, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$2y'' - e^y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 15

```
dsolve([2*diff(y(x),x$2)=exp(y(x)),y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = 2 \ln(2) + \ln\left(\frac{1}{(x-2)^2}\right)$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 15

```
DSolve[{2*y'[x]==Exp[y[x]],{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2 \log\left(1 - \frac{x}{2}\right)$$

## 10.20 problem Exercise 35.20, page 504

Internal problem ID [4670]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.20, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^2 y'' + y' x = 1$$

With initial conditions

$$[y(1) = 1, y'(1) = 2]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve([x^2*diff(y(x),x$2)+x*diff(y(x),x)=1,y(1) = 1, D(y)(1) = 2],y(x), singsol=all)
```

$$y(x) = \frac{\ln(x)^2}{2} + 2 \ln(x) + 1$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 19

```
DSolve[{x^2*y'[x]+x*y'[x]==1,{y[1]==1,y'[1]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(\log^2(x) + 4 \log(x) + 2)$$

## 10.21 problem Exercise 35.21, page 504

Internal problem ID [4671]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.21, page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' - y' = x^2$$

With initial conditions

$$[y(1) = 0, y'(1) = -1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve([x*diff(y(x),x$2)-diff(y(x),x)=x^2,y(1) = 0, D(y)(1) = -1],y(x), singsol=all)
```

$$y(x) = \frac{1}{3}x^3 - x^2 + \frac{2}{3}$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 19

```
DSolve[{x*y'[x]-y'[x]==x^2,{y[1]==0,y'[1]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}(x^3 - 3x^2 + 2)$$

## 10.22 problem Exercise 35.23(a), page 504

Internal problem ID [4672]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.23(a), page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [\_Liouville, [\_2nd\_order, \_with\_linear\_symmetries], [\_2nd\_order]

$$xyy'' - 2xy'^2 + yy' = 0$$

### ✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 18

```
dsolve(x*y(x)*diff(y(x),x$2)-2*x*(diff(y(x),x))^2+y(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = -\frac{1}{c_1 \ln(x) + c_2}$$

### ✓ Solution by Mathematica

Time used: 0.243 (sec). Leaf size: 22

```
DSolve[x*y[x]*y'[x]-2*x*(y'[x])^2+y[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2}{-\log(x) + c_1}$$

$$y(x) \rightarrow 0$$

## 10.23 problem Exercise 35.23(b), page 504

Internal problem ID [4673]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.23(b), page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _exact, _nonlinear], _Liouville, [_2nd_order, _w`

$$xyy'' + xy'^2 - yy' = 0$$

### ✓ Solution by Maple

Time used: 0.11 (sec). Leaf size: 35

```
dsolve(x*y(x)*diff(y(x),x$2)+x*(diff(y(x),x))^2-y(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = \sqrt{c_1x^2 + 2c_2}$$

$$y(x) = -\sqrt{c_1x^2 + 2c_2}$$

### ✓ Solution by Mathematica

Time used: 0.241 (sec). Leaf size: 18

```
DSolve[x*y[x]*y'[x]+x*(y'[x])^2-y[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2\sqrt{x^2 + c_1}$$

## 10.24 problem Exercise 35.23(c), page 504

Internal problem ID [4674]

**Book:** Ordinary Differential Equations, By Tenenbaum and Pollard. Dover, NY 1963

**Section:** Chapter 8. Special second order equations. Lesson 35. Independent variable x absent

**Problem number:** Exercise 35.23(c), page 504.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries], [_2nd_order, _reducibl`

$$xyy'' - 2xy'^2 + (1 + y)y' = 0$$

### ✓ Solution by Maple

Time used: 0.204 (sec). Leaf size: 22

```
dsolve(x*y(x)*diff(y(x),x$2)-2*x*(diff(y(x),x))^2+(1+y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1 \tanh\left(\frac{\ln(x) - c_2}{2c_1}\right)$$

### ✓ Solution by Mathematica

Time used: 20.549 (sec). Leaf size: 52

```
DSolve[x*y[x]*y'[x]-2*x*(y'[x])^2+(1+y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{\tan\left(\frac{\sqrt{c_1}(\log(x)-c_2)}{\sqrt{2}}\right)}{\sqrt{2}\sqrt{c_1}}$$

$$y(x) \rightarrow \frac{1}{2}(\log(x) - c_2)$$