

A Solution Manual For

# Ordinary Differential Equations, Robert H. Martin, 1983



**Nasser M. Abbasi**

March 3, 2024

# Contents

1	Problem 1.1-2, page 6	2
2	Problem 1.1-3, page 6	10
3	Problem 1.1-4, page 7	17
4	Problem 1.1-5, page 7	20
5	Problem 1.1-6, page 7	22
6	Problem 1.2-1, page 12	27
7	Problem 1.2-2, page 12	37
8	Problem 1.2-3, page 12	44

# 1 Problem 1.1-2, page 6

1.1	problem 1.1-2 (a)	3
1.2	problem 1.1-2 (b)	4
1.3	problem 1.1-2 (c)	5
1.4	problem 1.1-2 (d)	6
1.5	problem 1.1-2 (e)	7
1.6	problem 1.1-2 (f)	8
1.7	problem 1.1-2 (g)	9

## 1.1 problem 1.1-2 (a)

Internal problem ID [2447]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-2, page 6

**Problem number:** 1.1-2 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = t^2 + 3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(t),t)=t^2+3,y(t), singsol=all)
```

$$y(t) = \frac{1}{3}t^3 + 3t + c_1$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 18

```
DSolve[y'[t]==t^2+3,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{t^3}{3} + 3t + c_1$$

## 1.2 problem 1.1-2 (b)

Internal problem ID [2448]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-2, page 6

**Problem number:** 1.1-2 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = t e^{2t}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(t),t)=t*exp(2*t),y(t), singsol=all)
```

$$y(t) = \frac{(2t - 1)e^{2t}}{4} + c_1$$

### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 22

```
DSolve[y'[t]==t*Exp[2*t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{4}e^{2t}(2t - 1) + c_1$$

### 1.3 problem 1.1-2 (c)

Internal problem ID [2449]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-2, page 6

**Problem number:** 1.1-2 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = \sin(3t)$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(t),t)=sin(3*t),y(t), singsol=all)
```

$$y(t) = -\frac{\cos(3t)}{3} + c_1$$

#### ✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 16

```
DSolve[y'[t]==Sin[3*t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -\frac{1}{3} \cos(3t) + c_1$$

## 1.4 problem 1.1-2 (d)

Internal problem ID [2450]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-2, page 6

**Problem number:** 1.1-2 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = \sin(t)^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(t),t)=sin(t)^2,y(t), singsol=all)
```

$$y(t) = \frac{t}{2} + c_1 - \frac{\sin(2t)}{4}$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 21

```
DSolve[y'[t]==Sin[t]^2,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{t}{2} - \frac{1}{4} \sin(2t) + c_1$$

## 1.5 problem 1.1-2 (e)

Internal problem ID [2451]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-2, page 6

**Problem number:** 1.1-2 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = \frac{t}{t^2 + 4}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(t),t)=t/(t^2+4),y(t), singsol=all)
```

$$y(t) = \frac{\ln(t^2 + 4)}{2} + c_1$$

### ✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 18

```
DSolve[y'[t]==t/(t^2+4),y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{2} \log(t^2 + 4) + c_1$$



## 1.6 problem 1.1-2 (f)

Internal problem ID [2452]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-2, page 6

**Problem number:** 1.1-2 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' = \ln(t)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(t),t)=ln(t),y(t), singsol=all)
```

$$y(t) = t \ln(t) - t + c_1$$

### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 15

```
DSolve[y'[t]==Log[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -t + t \log(t) + c_1$$

## 1.7 problem 1.1-2 (g)

Internal problem ID [2453]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-2, page 6

**Problem number:** 1.1-2 (g).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = \frac{t}{\sqrt{t} + 1}$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 27

```
dsolve(diff(y(t),t)=t/(sqrt(t)+1),y(t), singsol=all)
```

$$y(t) = \frac{2t^{\frac{3}{2}}}{3} - t + 2\sqrt{t} - 2\ln(\sqrt{t} + 1) + c_1$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 25

```
DSolve[y'[t]==1/(1+Sqrt[t]),y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 2\sqrt{t} - 2\log(\sqrt{t} + 1) + c_1$$

## **2 Problem 1.1-3, page 6**

2.1	problem 1.1-3 (a) . . . . .	11
2.2	problem 1.1-3 (b) . . . . .	12
2.3	problem 1.1-3 (c) . . . . .	13
2.4	problem 1.1-3 (d) . . . . .	14
2.5	problem 1.1-3 (e) . . . . .	15
2.6	problem 1.1-3 (f) . . . . .	16

## 2.1 problem 1.1-3 (a)

Internal problem ID [2454]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-3, page 6

**Problem number:** 1.1-3 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - 2y = -4$$

With initial conditions

$$[y(0) = 5]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 12

```
dsolve([diff(y(t),t)=2*y(t)-4,y(0) = 5],y(t), singsol=all)
```

$$y(t) = 2 + 3e^{2t}$$

### ✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 14

```
DSolve[{y'[t]==2*y[t]-4,y[0]==5},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 3e^{2t} + 2$$

## 2.2 problem 1.1-3 (b)

Internal problem ID [2455]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-3, page 6

**Problem number:** 1.1-3 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' + y^3 = 0$$

With initial conditions

$$[y(1) = 3]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 13

```
dsolve([diff(y(t),t)=-y(t)^3,y(1) = 3],y(t), singsol=all)
```

$$y(t) = \frac{3}{\sqrt{18t - 17}}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 16

```
DSolve[{y'[t]==-y[t]^3,y[1]==3},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{3}{\sqrt{18t - 17}}$$

## 2.3 problem 1.1-3 (c)

Internal problem ID [2456]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-3, page 6

**Problem number:** 1.1-3 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y' - \frac{e^t}{y} = 0$$

With initial conditions

$$[y(\ln(2)) = -8]$$

### ✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 14

```
dsolve([diff(y(t),t)=exp(t)/y(t),y(ln(2)) = -8],y(t), singsol=all)
```

$$y(t) = -\sqrt{2e^t + 60}$$

### ✓ Solution by Mathematica

Time used: 0.594 (sec). Leaf size: 21

```
DSolve[{y'[t]==Exp[t]/y[t],y[Log[2]]==-8},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -\sqrt{2}\sqrt{e^t + 30}$$

## 2.4 problem 1.1-3 (d)

Internal problem ID [2457]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-3, page 6

**Problem number:** 1.1-3 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = t e^{2t}$$

With initial conditions

$$[y(1) = 5]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 21

```
dsolve([diff(y(t),t)=t*exp(2*t),y(1) = 5],y(t), singsol=all)
```

$$y(t) = \frac{(2t - 1) e^{2t}}{4} + 5 - \frac{e^2}{4}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 27

```
DSolve[{y'[t]==t*Exp[2*t],y[1]==5},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{4}(e^{2t}(2t - 1) - e^2 + 20)$$

## 2.5 problem 1.1-3 (e)

Internal problem ID [2458]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-3, page 6

**Problem number:** 1.1-3 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = \sin(t)^2$$

With initial conditions

$$\left[ y\left(\frac{\pi}{6}\right) = 3 \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 23

```
dsolve([diff(y(t),t)=sin(t)^2,y(1/6*Pi) = 3],y(t), singsol=all)
```

$$y(t) = \frac{t}{2} + 3 - \frac{\pi}{12} + \frac{\sqrt{3}}{8} - \frac{\sin(2t)}{4}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 31

```
DSolve[{y'[t]==Sin[t]^2,y[Pi/6]==3},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{24} \left( 3(4t + \sqrt{3} + 24) - 6\sin(2t) - 2\pi \right)$$



## 2.6 problem 1.1-3 (f)

Internal problem ID [2459]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-3, page 6

**Problem number:** 1.1-3 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_quadrature`]

$$y' = 8e^{4t} + t$$

With initial conditions

$$[y(0) = 12]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([diff(y(t),t)=8*exp(4*t)+t,y(0) = 12],y(t), singsol=all)
```

$$y(t) = \frac{t^2}{2} + 2e^{4t} + 10$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 21

```
DSolve[{y'[t]==8*Exp[4*t]+t,y[0]==12},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{2}(t^2 + 4e^{4t} + 20)$$

### **3 Problem 1.1-4, page 7**

3.1	problem 1.1-4 (a) . . . . .	18
3.2	problem 1.1-4 (b) . . . . .	19

### 3.1 problem 1.1-4 (a)

Internal problem ID [2460]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-4, page 7

**Problem number:** 1.1-4 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - \frac{y}{t} = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 7

```
dsolve(diff(y(t),t)=y(t)/t,y(t), singsol=all)
```

$$y(t) = tc_1$$

#### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 14

```
DSolve[y'[t]==y[t]/t,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow c_1 t$$

$$y(t) \rightarrow 0$$

### 3.2 problem 1.1-4 (b)

Internal problem ID [2461]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-4, page 7

**Problem number:** 1.1-4 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' + \frac{t}{y} = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(t),t)=-t/y(t),y(t), singsol=all)
```

$$y(t) = \sqrt{-t^2 + c_1}$$

$$y(t) = -\sqrt{-t^2 + c_1}$$

#### ✓ Solution by Mathematica

Time used: 0.105 (sec). Leaf size: 39

```
DSolve[y'[t]==-t/y[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -\sqrt{-t^2 + 2c_1}$$

$$y(t) \rightarrow \sqrt{-t^2 + 2c_1}$$

## 4 Problem 1.1-5, page 7

4.1	problem 1.1-5 . . . . .	21
-----	-------------------------	----

## 4.1 problem 1.1-5

Internal problem ID [2462]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-5, page 7

**Problem number:** 1.1-5.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - y^2 + y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve(diff(y(t),t)=y(t)^2-y(t),y(t), singsol=all)
```

$$y(t) = \frac{1}{1 + c_1 e^t}$$

### ✓ Solution by Mathematica

Time used: 0.242 (sec). Leaf size: 25

```
DSolve[y'[t]==y[t]^2-y[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{1 + e^{t+c_1}}$$

$$y(t) \rightarrow 0$$

$$y(t) \rightarrow 1$$

## **5 Problem 1.1-6, page 7**

5.1	problem 1.1-6 (a) . . . . .	23
5.2	problem 1.1-6 (b) . . . . .	24
5.3	problem 1.1-6 (c) . . . . .	25
5.4	problem 1.1-6 (d) . . . . .	26

## 5.1 problem 1.1-6 (a)

Internal problem ID [2463]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-6, page 7

**Problem number:** 1.1-6 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - y = -1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(t),t)=y(t)-1,y(t), singsol=all)
```

$$y(t) = 1 + c_1 e^t$$

### ✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 18

```
DSolve[y'[t]==y[t]-1,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 1 + c_1 e^t$$

$$y(t) \rightarrow 1$$



## 5.2 problem 1.1-6 (b)

Internal problem ID [2464]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-6, page 7

**Problem number:** 1.1-6 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' + y = 1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(t),t)=1-y(t),y(t), singsol=all)
```

$$y(t) = 1 + e^{-t}c_1$$

### ✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 20

```
DSolve[y'[t]==1-y[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 1 + c_1 e^{-t}$$

$$y(t) \rightarrow 1$$

### 5.3 problem 1.1-6 (c)

Internal problem ID [2465]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-6, page 7

**Problem number:** 1.1-6 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - y^3 + y^2 = 0$$

#### ✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 16

```
dsolve(diff(y(t),t)=y(t)^3-y(t)^2,y(t), singsol=all)
```

$$y(t) = \frac{1}{\text{LambertW}(-c_1 e^{t-1}) + 1}$$

#### ✓ Solution by Mathematica

Time used: 0.227 (sec). Leaf size: 38

```
DSolve[y'[t]==y[t]^3-y[t]^2,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \text{InverseFunction} \left[ \frac{1}{\#1} + \log(1 - \#1) - \log(\#1) \& \right] [t + c_1]$$

$$y(t) \rightarrow 0$$

$$y(t) \rightarrow 1$$

## 5.4 problem 1.1-6 (d)

Internal problem ID [2466]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.1-6, page 7

**Problem number:** 1.1-6 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' + y^2 = 1$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 8

```
dsolve(diff(y(t),t)=1-y(t)^2,y(t), singsol=all)
```

$$y(t) = \tanh(t + c_1)$$

### ✓ Solution by Mathematica

Time used: 0.713 (sec). Leaf size: 44

```
DSolve[y'[t]==1-y[t]^2,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{e^{2t} - e^{2c_1}}{e^{2t} + e^{2c_1}}$$

$$y(t) \rightarrow -1$$

$$y(t) \rightarrow 1$$

## 6 Problem 1.2-1, page 12

6.1	problem 1.2-1 (a)	28
6.2	problem 1.2-1 (b)	29
6.3	problem 1.2-1 (c)	30
6.4	problem 1.2-1 (d)	31
6.5	problem 1.2-1 (e)	32
6.6	problem 1.2-1 (f)	33
6.7	problem 1.2-1 (g)	34
6.8	problem 1.2-1 (h)	35
6.9	problem 1.2-1 (i)	36

## 6.1 problem 1.2-1 (a)

Internal problem ID [2467]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-1, page 12

**Problem number:** 1.2-1 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y' - y(t^2 + 1) = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(t),t)=(t^2+1)*y(t),y(t), singsol=all)
```

$$y(t) = c_1 e^{\frac{t(t^2+3)}{3}}$$

### ✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 24

```
DSolve[y'[t]==(t^2+1)*y[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow c_1 e^{\frac{t^3}{3}+t}$$

$$y(t) \rightarrow 0$$

## 6.2 problem 1.2-1 (b)

Internal problem ID [2468]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-1, page 12

**Problem number:** 1.2-1 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' + y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(t),t)=-y(t),y(t), singsol=all)
```

$$y(t) = e^{-t}c_1$$

### ✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 18

```
DSolve[y'[t]==-y[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow c_1 e^{-t}$$

$$y(t) \rightarrow 0$$

### 6.3 problem 1.2-1 (c)

Internal problem ID [2469]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-1, page 12

**Problem number:** 1.2-1 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_linear, 'class A']`

$$y' - 2y = e^{-3t}$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve(diff(y(t),t)=2*y(t)+exp(-3*t),y(t), singsol=all)
```

$$y(t) = \left( -\frac{e^{-5t}}{5} + c_1 \right) e^{2t}$$

#### ✓ Solution by Mathematica

Time used: 0.065 (sec). Leaf size: 23

```
DSolve[y'[t]==2*y[t]+Exp[-3*t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -\frac{e^{-3t}}{5} + c_1 e^{2t}$$

## 6.4 problem 1.2-1 (d)

Internal problem ID [2470]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-1, page 12

**Problem number:** 1.2-1 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_linear, 'class A']`

$$y' - 2y = e^{2t}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(t),t)=2*y(t)+exp(2*t),y(t), singsol=all)
```

$$y(t) = (t + c_1)e^{2t}$$

### ✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 15

```
DSolve[y'[t]==2*y[t]+Exp[2*t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{2t}(t + c_1)$$



## 6.5 problem 1.2-1 (e)

Internal problem ID [2471]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-1, page 12

**Problem number:** 1.2-1 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + y = t$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(t),t)=-y(t)+t,y(t), singsol=all)
```

$$y(t) = t - 1 + e^{-t}c_1$$

### ✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 16

```
DSolve[y'[t]==-y[t]+t,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow t + c_1 e^{-t} - 1$$

## 6.6 problem 1.2-1 (f)

Internal problem ID [2472]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-1, page 12

**Problem number:** 1.2-1 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_linear]`

$$ty' + 2y = \sin(t)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(t*diff(y(t),t)+2*y(t)=sin(t),y(t), singsol=all)
```

$$y(t) = \frac{\sin(t) - \cos(t)t + c_1}{t^2}$$

### ✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 19

```
DSolve[t*y'[t]+2*y[t]==Sin[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{\sin(t) - t \cos(t) + c_1}{t^2}$$

## 6.7 problem 1.2-1 (g)

Internal problem ID [2473]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-1, page 12

**Problem number:** 1.2-1 (g).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_linear]`

$$y' - y \tan(t) = \sec(t)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(t),t)=y(t)*tan(t)+sec(t),y(t), singsol=all)
```

$$y(t) = \frac{t + c_1}{\cos(t)}$$

### ✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 12

```
DSolve[y'[t]==y[t]*Tan[t]+Sec[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow (t + c_1) \sec(t)$$

## 6.8 problem 1.2-1 (h)

Internal problem ID [2474]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-1, page 12

**Problem number:** 1.2-1 (h).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_linear]`

$$y' - \frac{2ty}{t^2 + 1} = t + 1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(t),t)=2*t/(t^2+1)*y(t)+t+1,y(t), singsol=all)
```

$$y(t) = \left( \frac{\ln(t^2 + 1)}{2} + \arctan(t) + c_1 \right) (t^2 + 1)$$

### ✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 26

```
DSolve[y'[t]==2*t/(t^2+1)*y[t]+t+1,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow (t^2 + 1) \left( \arctan(t) + \frac{1}{2} \log(t^2 + 1) + c_1 \right)$$

## 6.9 problem 1.2-1 (i)

Internal problem ID [2475]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-1, page 12

**Problem number:** 1.2-1 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_linear]`

$$y' - y \tan(t) = \sec(t)^3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(t),t)=y(t)*tan(t)+sec(t)^3,y(t), singsol=all)
```

$$y(t) = \frac{\tan(t) + c_1}{\cos(t)}$$

### ✓ Solution by Mathematica

Time used: 0.057 (sec). Leaf size: 13

```
DSolve[y'[t]==y[t]*Tan[t]+Sec[t]^3,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \sec(t)(\tan(t) + c_1)$$

## 7 Problem 1.2-2, page 12

7.1	problem 1.2-2 (a)	38
7.2	problem 1.2-2 (b)	39
7.3	problem 1.2-2 (c)	40
7.4	problem 1.2-2 (d)	41
7.5	problem 1.2-2 (e)	42
7.6	problem 1.2-2 (f)	43

## 7.1 problem 1.2-2 (a)

Internal problem ID [2476]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-2, page 12

**Problem number:** 1.2-2 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - y = 0$$

With initial conditions

$$[y(0) = 2]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 8

```
dsolve([diff(y(t),t)=y(t),y(0) = 2],y(t), singsol=all)
```

$$y(t) = 2e^t$$

### ✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 10

```
DSolve[{y'[t]==y[t],y[0]==2},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 2e^t$$

## 7.2 problem 1.2-2 (b)

Internal problem ID [2477]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-2, page 12

**Problem number:** 1.2-2 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - 2y = 0$$

With initial conditions

$$[y(\ln(3)) = 3]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve([diff(y(t),t)=2*y(t),y(ln(3)) = 3],y(t), singsol=all)
```

$$y(t) = \frac{e^{2t}}{3}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 14

```
DSolve[{y'[t]==2*y[t],y[Log[3]]==3},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{e^{2t}}{3}$$



### 7.3 problem 1.2-2 (c)

Internal problem ID [2478]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-2, page 12

**Problem number:** 1.2-2 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_linear]`

$$ty' - y = t^3$$

With initial conditions

$$[y(1) = -2]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve([t*diff(y(t),t)=y(t)+t^3,y(1) = -2],y(t), singsol=all)
```

$$y(t) = \frac{(t^2 - 5)t}{2}$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 27

```
DSolve[{y'[t]==y[t]+t^3,y[1]==-2},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -t^3 - 3t^2 - 6t + 14e^{t-1} - 6$$

## 7.4 problem 1.2-2 (d)

Internal problem ID [2479]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-2, page 12

**Problem number:** 1.2-2 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_linear]`

$$y' + y \tan(t) = \sec(t)$$

With initial conditions

$$[y(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 6

```
dsolve([diff(y(t),t)=-tan(t)*y(t)+sec(t),y(0) = 0],y(t), singsol=all)
```

$$y(t) = \sin(t)$$

### ✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 7

```
DSolve[{y'[t]==-Tan[t]*y[t]+Sec[t],y[0]==0},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \sin(t)$$

## 7.5 problem 1.2-2 (e)

Internal problem ID [2480]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-2, page 12

**Problem number:** 1.2-2 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_separable`]

$$y' - \frac{2y}{t+1} = 0$$

With initial conditions

$$[y(0) = 6]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve([diff(y(t),t)=2/(1+t)*y(t),y(0) = 6],y(t), singsol=all)
```

$$y(t) = 6(t+1)^2$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 12

```
DSolve[{y'[t]==2/(1+t)*y[t],y[0]==6},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 6(t+1)^2$$

## 7.6 problem 1.2-2 (f)

Internal problem ID [2481]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-2, page 12

**Problem number:** 1.2-2 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_linear]`

$$ty' + y = t^3$$

With initial conditions

$$[y(1) = 2]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve([t*diff(y(t),t)=-y(t)+t^3,y(1) = 2],y(t), singsol=all)
```

$$y(t) = \frac{t^4 + 7}{4t}$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 27

```
DSolve[{y'[t]==-y[t]+t^3,y[1]==2},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow t^3 - 3t^2 + 6t + 4e^{1-t} - 6$$

## 8 Problem 1.2-3, page 12

8.1	problem 1.2-3 (a) . . . . .	45
8.2	problem 1.2-3 (b) . . . . .	46
8.3	problem 1.2-3 (c) . . . . .	47
8.4	problem 1.2-3 (d) . . . . .	48

## 8.1 problem 1.2-3 (a)

Internal problem ID [2482]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-3, page 12

**Problem number:** 1.2-3 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' + 4 \tan(2t) y = \tan(2t)$$

With initial conditions

$$\left[ y\left(\frac{\pi}{8}\right) = 2 \right]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve([diff(y(t),t)+4*tan(2*t)*y(t)=tan(2*t),y(1/8*Pi) = 2],y(t), singsol=all)
```

$$y(t) = \frac{7 \cos(2t)^2}{2} + \frac{1}{4}$$

### ✓ Solution by Mathematica

Time used: 0.098 (sec). Leaf size: 15

```
DSolve[{y'[t]+4*Tan[2*t]*y[t]==Tan[2*t],y[Pi/8]==2},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{7}{4} \cos(4t) + 2$$

## 8.2 problem 1.2-3 (b)

Internal problem ID [2483]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-3, page 12

**Problem number:** 1.2-3 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_linear]`

$$t \ln(t) y' + y = t \ln(t)$$

With initial conditions

$$[y(e) = 1]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve([t*ln(t)*diff(y(t),t)=t*ln(t)-y(t),y(exp(1)) = 1],y(t), singsol=all)
```

$$y(t) = \frac{t \ln(t) - t + 1}{\ln(t)}$$

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 19

```
DSolve[{t*Log[t]*y'[t]==t*Log[t]-y[t],y[Exp[1]]==1},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{-t + t \log(t) + 1}{\log(t)}$$

### 8.3 problem 1.2-3 (c)

Internal problem ID [2484]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-3, page 12

**Problem number:** 1.2-3 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y' - \frac{2y}{-t^2 + 1} = 3$$

With initial conditions

$$\left[ y\left(\frac{1}{2}\right) = 1 \right]$$

#### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 34

```
dsolve([diff(y(t),t)=2/(1-t^2)*y(t)+3,y(1/2) = 1],y(t), singsol=all)
```

$$y(t) = \frac{(t+1)(18t - 36 \ln(t+1) - 11 + 36 \ln(3) - 36 \ln(2))}{6t - 6}$$

#### ✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 34

```
DSolve[{y'[t]==2/(1-t^2)*y[t]+3,y[1/2]==1},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{(t+1)(18t - 36 \log(t+1) - 11 + 36 \log(\frac{3}{2}))}{6(t-1)}$$



## 8.4 problem 1.2-3 (d)

Internal problem ID [2485]

**Book:** Ordinary Differential Equations, Robert H. Martin, 1983

**Section:** Problem 1.2-3, page 12

**Problem number:** 1.2-3 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y' + \cot(t)y = 6 \cos(t)^2$$

With initial conditions

$$\left[ y\left(\frac{\pi}{4}\right) = 3 \right]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 18

```
dsolve([diff(y(t),t)=-cot(t)*y(t)+6*cos(t)^2,y(1/4*Pi) = 3],y(t), singsol=all)
```

$$y(t) = -2 \csc(t) \left( \cos(t)^3 - \sqrt{2} \right)$$

✓ Solution by Mathematica

Time used: 0.06 (sec). Leaf size: 23

```
DSolve[{y'[t]==-Cot[t]*y[t]+6*Cos[t]^2,y[Pi/4]==3},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 2\sqrt{2} \csc(t) - 2 \cos^2(t) \cot(t)$$