


A Solution Manual For

# Ordinary Differential Equations.

An introduction to the  
fundamentals. Kenneth B.  
Howell. second edition. CRC  
Press. FL, USA. 2020



**Nasser M. Abbasi**

March 3, 2024

# Contents

1	Chapter 2. Integration and differential equations. Additional exercises. page 32	3
2	Chapter 3. Some basics about First order equations. Additional exercises. page 63	52
3	Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90	66
4	Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103	120
5	Chapter 6. Simplifying through simplification. Additional exercises. page 114	151
6	Chapter 7. The exact form and general integrating factors. Additional exercises. page 141	184
7	Chapter 8. Review exercises for part of part II. page 143	214
8	Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259	271
9	Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277	328
10	Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294	364
11	Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334	384
12	Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369	424
13	Chapter 20. Euler equations. Additional exercises page 382	451
14	Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391	484

15 Chapter 22. Method of undetermined coefficients. Additional exercises page 412	506
16 Chapter 24. Variation of parameters. Additional exercises page 444	590
17 Chapter 25. Review exercises for part III. page 447	614
18 Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496	667
19 Chapter 28. The inverse Laplace transform. Additional Exercises. page 509	682
20 Chapter 29. Convolution. Additional Exercises. page 523	694
21 Chapter 30. Piecewise-defined functions and periodic functions. Additional Exercises. page 553	705
22 Chapter 31. Delta Functions. Additional Exercises. page 572	717
23 Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641	737
24 Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678	772
25 Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715	808
26 Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739	849
27 Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786	870
28 Chapter 39. Critical points, Direction fields and trajectories. Additional Exercises. page 815	891

# 1 Chapter 2. Integration and differential equations.

## Additional exercises. page 32

1.1	problem 2.2 (a)	5
1.2	problem 2.2 (b)	6
1.3	problem 2.2 (c)	7
1.4	problem 2.2 (d)	8
1.5	problem 2.2 (e)	9
1.6	problem 2.2 (f)	10
1.7	problem 2.2 (g)	11
1.8	problem 2.2 (h)	12
1.9	problem 2.2 (i)	13
1.10	problem 2.2 (j)	15
1.11	problem 2.3 (a)	16
1.12	problem 2.3 (b)	17
1.13	problem 2.3 (c)	18
1.14	problem 2.3 (d)	19
1.15	problem 2.3 (e)	20
1.16	problem 2.3 (f)	21
1.17	problem 2.3 (g)	22
1.18	problem 2.3 (h)	23
1.19	problem 2.3 (i)	24
1.20	problem 2.3 (j)	25
1.21	problem 2.3 (k)	26
1.22	problem 2.3 (L)	27
1.23	problem 2.4 (a)	28
1.24	problem 2.4 (b)	29
1.25	problem 2.4 (c)	30
1.26	problem 2.4 (d)	31
1.27	problem 2.4 (e)	32
1.28	problem 2.4 (f)	33
1.29	problem 2.4 (g)	34
1.30	problem 2.5 (a)	35
1.31	problem 2.5 (b i)	36
1.32	problem 2.5 (b ii)	37
1.33	problem 2.6 (a)	38
1.34	problem 2.6 (b i)	39
1.35	problem 2.6 (b ii)	40

1.36	problem 2.6 (b iii)	41
1.37	problem 2.7 a	42
1.38	problem 2.7 b	43
1.39	problem 2.7 c	44
1.40	problem 2.7 d	45
1.41	problem 2.7 e	46
1.42	problem 2.7 f	47
1.43	problem 2.9 a	48
1.44	problem 2.9 b	49
1.45	problem 2.9 c	50

## 1.1 problem 2.2 (a)

Internal problem ID [12923]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.2 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' = 3 - \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(diff(y(x),x)=3-sin(x),y(x), singsol=all)
```

$$y = 3x + \cos(x) + c_1$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 13

```
DSolve[y'[x]==3-Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3x + \cos(x) + c_1$$

## 1.2 problem 2.2 (b)

Internal problem ID [12924]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.2 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' + \sin(y) = 3$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 26

```
dsolve(diff(y(x),x)=3-sin(y(x)),y(x), singsol=all)
```

$$y = 2 \arctan \left( \frac{(4 \tan((x + c_1) \sqrt{2}) + \sqrt{2}) \sqrt{2}}{6} \right)$$

### ✓ Solution by Mathematica

Time used: 5.504 (sec). Leaf size: 83

```
DSolve[y'[x]==3-Sin[y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2 \arctan \left( \frac{1}{3} \left( -1 - 2\sqrt{2} \tan \left( \sqrt{2}(x - c_1) \right) \right) \right)$$

$$y(x) \rightarrow 2 \arctan \left( \frac{1}{3} \left( 1 + 2\sqrt{2} \tan \left( \sqrt{2}(x - c_1) \right) \right) \right)$$

$$y(x) \rightarrow \arcsin(3)$$

$$y(x) \rightarrow \text{Interval}[\{-\pi, \pi\}]$$



### 1.3 problem 2.2 (c)

Internal problem ID [12925]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.2 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 4y = e^{2x}$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)+4*y(x)=exp(2*x),y(x), singsol=all)
```

$$y = \left( \frac{e^{6x}}{6} + c_1 \right) e^{-4x}$$

#### ✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 23

```
DSolve[y'[x]+4*y[x]==Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{2x}}{6} + c_1 e^{-4x}$$

## 1.4 problem 2.2 (d)

Internal problem ID [12926]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.2 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y'x = \arcsin(x^2)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 109

```
dsolve(x*diff(y(x),x)=arcsin(x^2),y(x), singsol=all)
```

$$y = -\frac{i \arcsin(x^2)^2}{4} + \frac{\arcsin(x^2) \ln(1 + ix^2 + \sqrt{-x^4 + 1})}{2} - \frac{i \operatorname{polylog}(2, -ix^2 - \sqrt{-x^4 + 1})}{2} + \frac{\arcsin(x^2) \ln(1 - ix^2 - \sqrt{-x^4 + 1})}{2} - \frac{i \operatorname{polylog}(2, ix^2 + \sqrt{-x^4 + 1})}{2} + c_1$$

### ✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 56

```
DSolve[x*y'[x]==ArcSin[x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{4}i \left( \arcsin(x^2)^2 + \operatorname{PolyLog}\left(2, e^{2i \arcsin(x^2)}\right) \right) + \frac{1}{2} \arcsin(x^2) \log\left(1 - e^{2i \arcsin(x^2)}\right) + c_1$$

## 1.5 problem 2.2 (e)

Internal problem ID [12927]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.2 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$yy' = 2x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(y(x)*diff(y(x),x)=2*x,y(x), singsol=all)
```

$$y = \sqrt{2x^2 + c_1}$$

$$y = -\sqrt{2x^2 + c_1}$$

### ✓ Solution by Mathematica

Time used: 0.075 (sec). Leaf size: 42

```
DSolve[y[x]*y'[x]==2*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{2}\sqrt{x^2 + c_1}$$

$$y(x) \rightarrow \sqrt{2}\sqrt{x^2 + c_1}$$

## 1.6 problem 2.2 (f)

Internal problem ID [12928]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.2 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$y'' = \frac{1+x}{-1+x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)=(x+1)/(x-1),y(x), singsol=all)
```

$$y = \frac{x^2}{2} + 2 \ln(-1+x)(-1+x) + 2 - 2x + c_1x + c_2$$

### ✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 30

```
DSolve[y''[x]==(x+1)/(x-1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2}{2} + 2(x-1) \log(x-1) + (-2+c_2)x + c_1$$

## 1.7 problem 2.2 (g)

Internal problem ID [12929]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.2 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$x^2 y'' = 1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(x^2*diff(y(x),x$2)=1,y(x), singsol=all)
```

$$y = -\ln(x) + c_1 x + c_2$$

### ✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 16

```
DSolve[x^2*y''[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\log(x) + c_2 x + c_1$$

## 1.8 problem 2.2 (h)

Internal problem ID [12930]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.2 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _with_linear_symmetries]]`

$$y^2 y'' = 8x^2$$

### **X** Solution by Maple

```
dsolve(y(x)^2*diff(y(x),x$2)=8*x^2,y(x), singsol=all)
```

No solution found

### **X** Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y[x]^2*y'[x]==8*x^2,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

## 1.9 problem 2.2 (i)

Internal problem ID [12931]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.2 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 3y' + 8y = e^{-x^2}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 121

```
dsolve(diff(y(x),x$2)+3*diff(y(x),x)+8*y(x)=exp(-x^2),y(x), singsol=all)
```

$$y = \sin\left(\frac{\sqrt{23}x}{2}\right) e^{-\frac{3x}{2}} c_2 + e^{-\frac{3x}{2}} \cos\left(\frac{\sqrt{23}x}{2}\right) c_1$$
$$+ \frac{\sqrt{23} e^{-\frac{3x}{2} - \frac{7}{8}} \sqrt{\pi} \left( e^{\frac{3i\sqrt{23}}{8}} \left( i \cos\left(\frac{\sqrt{23}x}{2}\right) + \sin\left(\frac{\sqrt{23}x}{2}\right) \right) \operatorname{erf}\left(x - \frac{3}{4} - \frac{i\sqrt{23}}{4}\right) - e^{-\frac{3i\sqrt{23}}{8}} \left( i \cos\left(\frac{\sqrt{23}x}{2}\right) - \sin\left(\frac{\sqrt{23}x}{2}\right) \right) \right)}{46}$$

✓ Solution by Mathematica

Time used: 0.519 (sec). Leaf size: 205

```
DSolve[y''[x]+3*y'[x]+8*y[x]==Exp[-x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned} y(x) \rightarrow & \frac{1}{46} e^{\frac{1}{8}(-12x-3i\sqrt{23}-7)} \left( -i e^{\frac{3i\sqrt{23}}{4}} \sqrt{23} \pi \operatorname{erf} \left( -x + \frac{i\sqrt{23}}{4} + \frac{3}{4} \right) \left( \cos \left( \frac{\sqrt{23}x}{2} \right) \right. \right. \\ & \left. \left. - i \sin \left( \frac{\sqrt{23}x}{2} \right) \right) \right. \\ & \left. + i \sqrt{23} \pi \operatorname{erf} \left( \frac{1}{4}(-4x - i\sqrt{23} + 3) \right) \left( \cos \left( \frac{\sqrt{23}x}{2} \right) + i \sin \left( \frac{\sqrt{23}x}{2} \right) \right) \right. \\ & \left. + 46 e^{\frac{7}{8} + \frac{3i\sqrt{23}}{8}} \left( c_2 \cos \left( \frac{\sqrt{23}x}{2} \right) + c_1 \sin \left( \frac{\sqrt{23}x}{2} \right) \right) \right) \end{aligned}$$



## 1.10 problem 2.2 (j)

Internal problem ID [12932]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.2 (j).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^2 y'' + 3y'x = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(x^2*diff(y(x),x$2)+3*x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = c_1 + \frac{c_2}{x^2}$$

### ✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 17

```
DSolve[x^2*y''[x]+3*x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \frac{c_1}{2x^2}$$

## 1.11 problem 2.3 (a)

Internal problem ID [12933]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.3 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' = 4x^3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)=4*x^3,y(x), singsol=all)
```

$$y = x^4 + c_1$$

### ✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 11

```
DSolve[y'[x]==4*x^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^4 + c_1$$

## 1.12 problem 2.3 (b)

Internal problem ID [12934]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.3 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' = 20 e^{-4x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=20*exp(-4*x),y(x), singsol=all)
```

$$y = -5e^{-4x} + c_1$$

### ✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 15

```
DSolve[y'[x]==20*Exp[-4*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -5e^{-4x} + c_1$$

### 1.13 problem 2.3 (c)

Internal problem ID [12935]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.3 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y'x = -\sqrt{x} + 2$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x*diff(y(x),x)+sqrt(x)=2,y(x), singsol=all)
```

$$y = -2\sqrt{x} + 2 \ln(x) + c_1$$

#### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 19

```
DSolve[x*y'[x]+Sqrt[x]==2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2\sqrt{x} + 2 \log(x) + c_1$$

## 1.14 problem 2.3 (d)

Internal problem ID [12936]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.3 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$\sqrt{x+4}y' = 1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(sqrt(x+4)*diff(y(x),x)=1,y(x), singsol=all)
```

$$y = 2\sqrt{x+4} + c_1$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 17

```
DSolve[Sqrt[x+4]*y'[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2\sqrt{x+4} + c_1$$

## 1.15 problem 2.3 (e)

Internal problem ID [12937]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.3 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' = x \cos(x^2)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=x*cos(x^2),y(x), singsol=all)
```

$$y = \frac{\sin(x^2)}{2} + c_1$$

### ✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 16

```
DSolve[y'[x]==x*Cos[x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sin(x^2)}{2} + c_1$$

## 1.16 problem 2.3 (f)

Internal problem ID [12938]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.3 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_quadrature`]

$$y' = x \cos(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=x*cos(x),y(x), singsol=all)
```

$$y = \cos(x) + x \sin(x) + c_1$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 14

```
DSolve[y'[x]==x*Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \sin(x) + \cos(x) + c_1$$

## 1.17 problem 2.3 (g)

Internal problem ID [12939]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.3 (g).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$-(x^2 - 9) y' = -x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(x=(x^2-9)*diff(y(x),x),y(x), singsol=all)
```

$$y = \frac{\ln(x^2 - 9)}{2} + c_1$$

### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 18

```
DSolve[x==(x^2-9)*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \log(x^2 - 9) + c_1$$



## 1.18 problem 2.3 (h)

Internal problem ID [12940]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.3 (h).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$-(x^2 - 9)y' = -1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(1=(x^2-9)*diff(y(x),x),y(x), singsol=all)
```

$$y = \frac{\ln(x-3)}{6} - \frac{\ln(x+3)}{6} + c_1$$

### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 26

```
DSolve[1==(x^2-9)*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{6}(\log(3-x) - \log(x+3) + 6c_1)$$

## 1.19 problem 2.3 (i)

Internal problem ID [12941]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.3 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$9y' = x^2 - 1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(1=x^2-9*diff(y(x),x),y(x), singsol=all)
```

$$y = \frac{1}{27}x^3 - \frac{1}{9}x + c_1$$

### ✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 20

```
DSolve[1==x^2-9*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{27} - \frac{x}{9} + c_1$$

## 1.20 problem 2.3 (j)

Internal problem ID [12942]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.3 (j).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$y'' = \sin(2x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x$2)=sin(2*x),y(x), singsol=all)
```

$$y = -\frac{\sin(2x)}{4} + c_1x + c_2$$

### ✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 20

```
DSolve[y''[x]==Sin[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{4}\sin(2x) + c_2x + c_1$$

## 1.21 problem 2.3 (k)

Internal problem ID [12943]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.3 (k).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$y'' = x + 3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x$2)-3=x,y(x), singsol=all)
```

$$y = \frac{1}{6}x^3 + \frac{3}{2}x^2 + c_1x + c_2$$

### ✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 26

```
DSolve[y''[x]-3==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{6} + \frac{3x^2}{2} + c_2x + c_1$$

## 1.22 problem 2.3 (L)

Internal problem ID [12944]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.3 (L).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _quadrature]]`

$$y'''' = 1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$4)=1,y(x), singsol=all)
```

$$y = \frac{1}{24}x^4 + \frac{1}{6}c_1x^3 + \frac{1}{2}c_2x^2 + c_3x + c_4$$

### ✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 31

```
DSolve[y''''[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^4}{24} + c_4x^3 + c_3x^2 + c_2x + c_1$$

## 1.23 problem 2.4 (a)

Internal problem ID [12945]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.4 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = 40x e^{2x}$$

With initial conditions

$$[y(0) = 4]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve([diff(y(x),x)=4*x*10*exp(2*x),y(0) = 4],y(x), singsol=all)
```

$$y = 14 + (20x - 10) e^{2x}$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 21

```
DSolve[{y'[x]==4*x*10*Exp[2*x],{y[0]==4}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2(5e^{2x}(2x - 1) + 7)$$

## 1.24 problem 2.4 (b)

Internal problem ID [12946]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.4 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_quadrature`]

$$(x + 6)^{\frac{1}{3}} y' = 1$$

With initial conditions

$$[y(2) = 10]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve([(x+6)^(1/3)*diff(y(x),x)=1,y(2) = 10],y(x), singsol=all)
```

$$y = \frac{3(x + 6)^{\frac{2}{3}}}{2} + 4$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 18

```
DSolve[{(x+6)^(1/3)*y'[x]==1,{y[2]==10}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3}{2}(x + 6)^{2/3} + 4$$

## 1.25 problem 2.4 (c)

Internal problem ID [12947]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.4 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = \frac{-1 + x}{1 + x}$$

With initial conditions

$$[y(0) = 8]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([diff(y(x),x)=(x-1)/(x+1),y(0) = 8],y(x), singsol=all)
```

$$y = x - 2 \ln(1 + x) + 8$$

### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 14

```
DSolve[{y'[x]==(x-1)/(x+1)},{y[0]==8}],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x - 2 \log(x + 1) + 8$$



## 1.26 problem 2.4 (d)

Internal problem ID [12948]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.4 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y'x = \sqrt{x} - 2$$

With initial conditions

$$[y(1) = 6]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve([x*diff(y(x),x)+2=sqrt(x),y(1) = 6],y(x), singsol=all)
```

$$y = 2\sqrt{x} - 2 \ln(x) + 4$$

### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 18

```
DSolve[{x*y'[x]+2==Sqrt[x],{y[1]==6}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2(\sqrt{x} - \log(x) + 2)$$

## 1.27 problem 2.4 (e)

Internal problem ID [12949]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.4 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_quadrature]`

$$\cos(x) y' = \sin(x)$$

With initial conditions

$$[y(0) = 3]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 11

```
dsolve([cos(x)*diff(y(x),x)-sin(x)=0,y(0) = 3],y(x), singsol=all)
```

$$y = -\ln(\cos(x)) + 3$$

### ✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 12

```
DSolve[{Cos[x]*y'[x]-Sin[x]==0,{y[0]==3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3 - \log(\cos(x))$$

## 1.28 problem 2.4 (f)

Internal problem ID [12950]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.4 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_quadrature]`

$$(x^2 + 1) y' = 1$$

With initial conditions

$$[y(0) = 3]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve([(x^2+1)*diff(y(x),x)=1,y(0) = 3],y(x), singsol=all)
```

$$y = \arctan(x) + 3$$

### ✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 9

```
DSolve[{(x^2+1)*y'[x]==1,{y[0]==3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arctan(x) + 3$$

## 1.29 problem 2.4 (g)

Internal problem ID [12951]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.4 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$y''x = \sqrt{x} - 2$$

With initial conditions

$$[y(1) = 8, y'(1) = 6]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve([x*diff(y(x),x$2)+2=sqrt(x),y(1) = 8, D(y)(1) = 6],y(x), singsol=all)
```

$$y = \frac{4x^{\frac{3}{2}}}{3} - 2 \ln(x)x + 6x + \frac{2}{3}$$

### ✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 26

```
DSolve[{x*y''[x]+2==Sqrt[x],{y[1]==8,y'[1]==6}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2}{3}(2x^{3/2} + 9x - 3x \log(x) + 1)$$

### 1.30 problem 2.5 (a)

Internal problem ID [12952]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.5 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = \sin\left(\frac{x}{2}\right)$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=sin(x/2),y(x), singsol=all)
```

$$y = -2 \cos\left(\frac{x}{2}\right) + c_1$$

#### ✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 16

```
DSolve[y'[x]==Sin[x/2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2 \cos\left(\frac{x}{2}\right) + c_1$$

### 1.31 problem 2.5 (b i)

Internal problem ID [12953]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.5 (b i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_quadrature`]

$$y' = \sin\left(\frac{x}{2}\right)$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve([diff(y(x),x)=sin(x/2),y(0) = 0],y(x), singsol=all)
```

$$y = -2 \cos\left(\frac{x}{2}\right) + 2$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 15

```
DSolve[{y'[x]==Sin[x/2]},{y[0]==0}],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 4 \sin^2\left(\frac{x}{4}\right)$$

## 1.32 problem 2.5 (b ii)

Internal problem ID [12954]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.5 (b ii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_quadrature`]

$$y' = \sin\left(\frac{x}{2}\right)$$

With initial conditions

$$[y(0) = 3]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve([diff(y(x),x)=sin(x/2),y(0) = 3],y(x), singsol=all)
```

$$y = -2 \cos\left(\frac{x}{2}\right) + 5$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 15

```
DSolve[{y'[x]==Sin[x/2]},{y[0]==3}],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 5 - 2 \cos\left(\frac{x}{2}\right)$$

### 1.33 problem 2.6 (a)

Internal problem ID [12955]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.6 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = 3\sqrt{x+3}$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x)=3*sqrt(x+3),y(x), singsol=all)
```

$$y = 2(x+3)^{\frac{3}{2}} + c_1$$

#### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 17

```
DSolve[y'[x]==3*Sqrt[x+3],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2(x+3)^{3/2} + c_1$$



### 1.34 problem 2.6 (b i)

Internal problem ID [12956]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.6 (b i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_quadrature]`

$$y' = 3\sqrt{x+3}$$

With initial conditions

$$[y(1) = 16]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve([diff(y(x),x)=3*sqrt(x+3),y(1) = 16],y(x), singsol=all)
```

$$y = 2(x+3)^{\frac{3}{2}}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 14

```
DSolve[{y'[x]==3*Sqrt[x+3],{y[1]==16}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2(x+3)^{3/2}$$

### 1.35 problem 2.6 (b ii)

Internal problem ID [12957]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.6 (b ii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' = 3\sqrt{x+3}$$

With initial conditions

$$[y(1) = 20]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
dsolve([diff(y(x),x)=3*sqrt(x+3),y(1) = 20],y(x), singsol=all)
```

$$y = 4 + (2x + 6)\sqrt{x+3}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 16

```
DSolve[{y'[x]==3*Sqrt[x+3]},{y[1]==20}],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2((x+3)^{3/2} + 2)$$

### 1.36 problem 2.6 (b iii)

Internal problem ID [12958]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.6 (b iii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' = 3\sqrt{x+3}$$

With initial conditions

$$[y(1) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([diff(y(x),x)=3*sqrt(x+3),y(1) = 0],y(x), singsol=all)
```

$$y = -16 + (2x + 6)\sqrt{x + 3}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 16

```
DSolve[{y'[x]==3*Sqrt[x+3]},{y[1]==0}],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2((x + 3)^{3/2} - 8)$$

## 1.37 problem 2.7 a

Internal problem ID [12959]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.7 a.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_quadrature`]

$$y' = x e^{-x^2}$$

With initial conditions

$$[y(0) = 3]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([diff(y(x),x)=x*exp(-x^2),y(0) = 3],y(x), singsol=all)
```

$$y = -\frac{e^{-x^2}}{2} + \frac{7}{2}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 20

```
DSolve[{y'[x]==x*Exp[-x^2],{y[0]==3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{7}{2} - \frac{e^{-x^2}}{2}$$

## 1.38 problem 2.7 b

Internal problem ID [12960]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.7 b.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' = \frac{x}{\sqrt{x^2 + 5}}$$

With initial conditions

$$[y(2) = 7]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve([diff(y(x),x)=x/sqrt(x^2+5),y(2) = 7],y(x), singsol=all)
```

$$y = \sqrt{x^2 + 5} + 4$$

### ✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 16

```
DSolve[{y'[x]==x/Sqrt[x^2+5]},{y[2]==7}],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt{x^2 + 5} + 4$$

### 1.39 problem 2.7 c

Internal problem ID [12961]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.7 c.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' = \frac{1}{x^2 + 1}$$

With initial conditions

$$[y(1) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve([diff(y(x),x)=1/(x^2+1),y(1) = 0],y(x), singsol=all)
```

$$y = \arctan(x) - \frac{\pi}{4}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 13

```
DSolve[{y'[x]==1/(x^2+1)},{y[1]==0}],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arctan(x) - \frac{\pi}{4}$$

## 1.40 problem 2.7 d

Internal problem ID [12962]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.7 d.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = e^{-9x^2}$$

With initial conditions

$$[y(0) = 1]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([diff(y(x),x)=exp(-9*x^2),y(0) = 1],y(x), singsol=all)
```

$$y = \frac{\sqrt{\pi} \operatorname{erf}(3x)}{6} + 1$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 20

```
DSolve[{y'[x]==Exp[-9*x^2],{y[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{6}\sqrt{\pi}\operatorname{erf}(3x) + 1$$

## 1.41 problem 2.7 e

Internal problem ID [12963]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.7 e.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_quadrature]`

$$y'x = \sin(x)$$

With initial conditions

$$[y(0) = 4]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve([x*diff(y(x),x)=sin(x),y(0) = 4],y(x), singsol=all)
```

$$y = \text{Si}(x) + 4$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 9

```
DSolve[{x*y'[x]==Sin[x],{y[0]==4}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{Si}(x) + 4$$



## 1.42 problem 2.7 f

Internal problem ID [12964]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.7 f.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y'x = \sin(x^2)$$

With initial conditions

$$[y(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve([x*diff(y(x),x)=sin(x^2),y(0) = 0],y(x), singsol=all)
```

$$y = \frac{\text{Si}(x^2)}{2}$$

### ✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 13

```
DSolve[{x*y'[x]==Sin[x^2]},{y[0]==0}],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\text{Si}(x^2)}{2}$$

## 1.43 problem 2.9 a

Internal problem ID [12965]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.9 a.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = \begin{cases} 0 & x < 0 \\ 1 & 0 \leq x \end{cases}$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([diff(y(x),x)=piecewise(x<0,0,x>=0,1),y(0) = 0],y(x), singsol=all)
```

$$y = \begin{cases} 0 & x < 0 \\ x & 0 \leq x \end{cases}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 9

```
DSolve[{y'[x]==UnitStep[x],{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x\theta(x)$$

## 1.44 problem 2.9 b

Internal problem ID [12966]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.9 b.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = \begin{cases} 0 & x < 1 \\ 1 & 1 \leq x \end{cases}$$

With initial conditions

$$[y(0) = 2]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve([diff(y(x),x)=piecewise(x<1,0,x>=1,1),y(0) = 2],y(x), singsol=all)
```

$$y = \begin{cases} 2 & x < 1 \\ 1 + x & 1 \leq x \end{cases}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 15

```
DSolve[{y'[x]==Piecewise[{{0,x<1},{1,x>=1}}],{y[0]==2}],y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow \begin{cases} 2 & x \leq 1 \\ x + 1 & \text{True} \end{cases}$$

## 1.45 problem 2.9 c

Internal problem ID [12967]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 2. Integration and differential equations. Additional exercises. page 32

**Problem number:** 2.9 c.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = \begin{cases} 0 & x < 1 \\ 1 & 1 \leq x < 2 \\ 0 & 2 \leq x \end{cases}$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve([diff(y(x),x)=piecewise(x<1,0,1<=x and x<2,1,2<=x,0),y(0) = 0],y(x), singsol=all)
```

$$y = \begin{cases} 0 & x < 1 \\ -1 + x & x < 2 \\ 1 & 2 \leq x \end{cases}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 23

```
DSolve[{y'[x]==Piecewise[{{0,x<1},{1,1<=x<2},{0,2<=x}}],{y[0]==0}],y[x],x,IncludeSingularSol
```

$$y(x) \rightarrow \begin{cases} 0 & x \leq 1 \\ x - 1 & 1 < x \leq 2 \\ 1 & \text{True} \end{cases}$$

## 2 Chapter 3. Some basics about First order equations. Additional exercises. page 63

2.1	problem 3.4 a . . . . .	53
2.2	problem 3.4 b . . . . .	54
2.3	problem 3.4 c . . . . .	55
2.4	problem 3.4 d . . . . .	57
2.5	problem 3.4 e . . . . .	58
2.6	problem 3.4 f . . . . .	59
2.7	problem 3.4 g . . . . .	61
2.8	problem 3.4 h . . . . .	62
2.9	problem 3.4 i . . . . .	63
2.10	problem 3.4 j . . . . .	64
2.11	problem 3.6 . . . . .	65

## 2.1 problem 3.4 a

Internal problem ID [12968]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 3. Some basics about First order equations. Additional exercises. page 63

**Problem number:** 3.4 a.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y' + 3yx = 6x$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)+3*x*y(x)=6*x,y(x), singsol=all)
```

$$y = 2 + c_1 e^{-\frac{3x^2}{2}}$$

### ✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 24

```
DSolve[y'[x]+3*x*y[x]==6*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 + c_1 e^{-\frac{3x^2}{2}}$$

$$y(x) \rightarrow 2$$

## 2.2 problem 3.4 b

Internal problem ID [12969]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 3. Some basics about First order equations. Additional exercises. page 63

**Problem number:** 3.4 b.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [ $y = G(x, y')$ ]

$$\sin(x + y) - yy' = 0$$

**X** Solution by Maple

```
dsolve(sin(x+y(x))-y(x)*diff(y(x),x)=0,y(x), singsol=all)
```

No solution found

**X** Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[Sin[x+y[x]]-y[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Not solved



## 2.3 problem 3.4 c

Internal problem ID [12970]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 3. Some basics about First order equations. Additional exercises. page 63

**Problem number:** 3.4 c.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' - y^3 = 8$$

### ✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 58

```
dsolve(diff(y(x),x)-y(x)^3=8,y(x), singsol=all)
```

$$y = 1 + \sqrt{3} \tan \left( \text{RootOf} \left( -\sqrt{3} \ln \left( \frac{1}{3 \tan(\_Z)^2 + 3} \right) - 2\sqrt{3} \ln \left( 3 + \sqrt{3} \tan(\_Z) \right) + 24\sqrt{3} c_1 + 24\sqrt{3} x - 6\_Z \right) \right)$$

✓ Solution by Mathematica

Time used: 0.206 (sec). Leaf size: 83

```
DSolve[y'[x]-y[x]^3==8,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{InverseFunction} \left[ -\frac{1}{24} \log(\#1^2 - 2\#1 + 4) + \frac{\arctan\left(\frac{\#1-1}{\sqrt{3}}\right)}{4\sqrt{3}} + \frac{1}{12} \log(\#1 + 2) \& \right] [x + c_1]$$

$$y(x) \rightarrow -2$$

$$y(x) \rightarrow 2\sqrt[3]{-1}$$

$$y(x) \rightarrow -2(-1)^{2/3}$$

## 2.4 problem 3.4 d

Internal problem ID [12971]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 3. Some basics about First order equations. Additional exercises. page 63

**Problem number:** 3.4 d.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y'x^2 + xy^2 = x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(x^2*diff(y(x),x)+x*y(x)^2=x,y(x), singsol=all)
```

$$y = \tanh(\ln(x) + c_1)$$

### ✓ Solution by Mathematica

Time used: 1.054 (sec). Leaf size: 40

```
DSolve[x^2*y'[x]+x*y[x]^2==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2 - e^{2c_1}}{x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 1$$

## 2.5 problem 3.4 e

Internal problem ID [12972]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 3. Some basics about First order equations. Additional exercises. page 63

**Problem number:** 3.4 e.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Riccati, _special]]`

$$y' - y^2 = x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x),x)-y(x)^2=x,y(x), singsol=all)
```

$$y = \frac{c_1 \text{AiryAi}(1, -x) + \text{AiryBi}(1, -x)}{c_1 \text{AiryAi}(-x) + \text{AiryBi}(-x)}$$

### ✓ Solution by Mathematica

Time used: 0.114 (sec). Leaf size: 195

```
DSolve[y'[x]-y[x]^2==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^{3/2} \left( -2 \text{BesselJ} \left( -\frac{2}{3}, \frac{2x^{3/2}}{3} \right) + c_1 \left( \text{BesselJ} \left( \frac{2}{3}, \frac{2x^{3/2}}{3} \right) - \text{BesselJ} \left( -\frac{4}{3}, \frac{2x^{3/2}}{3} \right) \right) \right) - c_1 \text{BesselJ} \left( -\frac{1}{3}, \frac{2x^{3/2}}{3} \right)}{2x \left( \text{BesselJ} \left( \frac{1}{3}, \frac{2x^{3/2}}{3} \right) + c_1 \text{BesselJ} \left( -\frac{1}{3}, \frac{2x^{3/2}}{3} \right) \right)}$$
$$y(x) \rightarrow -\frac{x^{3/2} \text{BesselJ} \left( -\frac{4}{3}, \frac{2x^{3/2}}{3} \right) - x^{3/2} \text{BesselJ} \left( \frac{2}{3}, \frac{2x^{3/2}}{3} \right) + \text{BesselJ} \left( -\frac{1}{3}, \frac{2x^{3/2}}{3} \right)}{2x \text{BesselJ} \left( -\frac{1}{3}, \frac{2x^{3/2}}{3} \right)}$$

## 2.6 problem 3.4 f

Internal problem ID [12973]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 3. Some basics about First order equations. Additional exercises. page 63

**Problem number:** 3.4 f.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y^3 - 25y + y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(y(x)^3-25*y(x)+diff(y(x),x)=0,y(x), singsol=all)
```

$$y = -\frac{5}{\sqrt{25 e^{-50x} c_1 + 1}}$$

$$y = \frac{5}{\sqrt{25 e^{-50x} c_1 + 1}}$$

✓ Solution by Mathematica

Time used: 0.68 (sec). Leaf size: 110

```
DSolve[y[x]^3-25*y[x]+y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{5e^{25x}}{\sqrt{e^{50x} + e^{50c_1}}}$$

$$y(x) \rightarrow \frac{5e^{25x}}{\sqrt{e^{50x} + e^{50c_1}}}$$

$$y(x) \rightarrow -5$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow 5$$

$$y(x) \rightarrow -\frac{5e^{25x}}{\sqrt{e^{50x}}}$$

$$y(x) \rightarrow \frac{5e^{25x}}{\sqrt{e^{50x}}}$$

## 2.7 problem 3.4 g

Internal problem ID [12974]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 3. Some basics about First order equations. Additional exercises. page 63

**Problem number:** 3.4 g.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$(x - 2)y' - y = 3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve((x-2)*diff(y(x),x)=y(x)+3,y(x), singsol=all)
```

$$y = -3 + (x - 2)c_1$$

### ✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 18

```
DSolve[(x-2)*y'[x]==y[x]+3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -3 + c_1(x - 2)$$

$$y(x) \rightarrow -3$$

## 2.8 problem 3.4 h

Internal problem ID [12975]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 3. Some basics about First order equations. Additional exercises. page 63

**Problem number:** 3.4 h.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$(y - 2)y' = x - 3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
dsolve((y(x)-2)*diff(y(x),x)=x-3,y(x), singsol=all)
```

$$y = 2 - \sqrt{x^2 + 2c_1 - 6x + 4}$$

$$y = 2 + \sqrt{x^2 + 2c_1 - 6x + 4}$$

### ✓ Solution by Mathematica

Time used: 0.112 (sec). Leaf size: 47

```
DSolve[(y[x]-2)*y'[x]==x-3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 - \sqrt{x^2 - 6x + 4 + 2c_1}$$

$$y(x) \rightarrow 2 + \sqrt{x^2 - 6x + 4 + 2c_1}$$



## 2.9 problem 3.4 i

Internal problem ID [12976]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 3. Some basics about First order equations. Additional exercises. page 63

**Problem number:** 3.4 i.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' + 2y - y^2 = -2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x)+2*y(x)-y(x)^2=-2,y(x), singsol=all)
```

$$y = \frac{(\sqrt{3} - 3 \tanh((x + c_1) \sqrt{3})) \sqrt{3}}{3}$$

### ✓ Solution by Mathematica

Time used: 0.394 (sec). Leaf size: 76

```
DSolve[y'[x]+2*y[x]-y[x]^2== -2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-(\sqrt{3} - 1) e^{2\sqrt{3}(x+c_1)} + 1 + \sqrt{3}}{1 + e^{2\sqrt{3}(x+c_1)}}$$

$$y(x) \rightarrow 1 - \sqrt{3}$$

$$y(x) \rightarrow 1 + \sqrt{3}$$

## 2.10 problem 3.4 j

Internal problem ID [12977]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 3. Some basics about First order equations. Additional exercises. page 63

**Problem number:** 3.4 j.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [Riccati]

$$y' + (8 - x)y - y^2 = -8x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
dsolve(diff(y(x),x)+(8-x)*y(x)-y(x)^2=-8*x,y(x), singsol=all)
```

$$y = 8 + \frac{e^{\frac{1}{2}x^2+8x}}{c_1 + \frac{i\sqrt{\pi}e^{-32\sqrt{2}}\operatorname{erf}\left(\frac{i\sqrt{2}x+4i\sqrt{2}}{2}\right)}{2}}$$

### ✓ Solution by Mathematica

Time used: 0.176 (sec). Leaf size: 54

```
DSolve[y'[x]+(8-x)*y[x]-y[x]^2==-8*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 8 + \frac{2e^{\frac{1}{2}(x+8)^2}}{-\sqrt{2\pi}\operatorname{erfi}\left(\frac{x+8}{\sqrt{2}}\right) + 2e^{32}c_1}$$
$$y(x) \rightarrow 8$$

## 2.11 problem 3.6

Internal problem ID [12978]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 3. Some basics about First order equations. Additional exercises. page 63

**Problem number:** 3.6.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' - 2\sqrt{y} = 0$$

With initial conditions

$$[y(1) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve([diff(y(x),x)=2*sqrt(y(x)),y(1) = 0],y(x), singsol=all)
```

$$y = 0$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 6

```
DSolve[{y'[x]==2*Sqrt[y[x]],{y[1]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

### 3 Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

3.1	problem 4.3 (a)	68
3.2	problem 4.3 (b)	69
3.3	problem 4.3 (c)	70
3.4	problem 4.3 (d)	71
3.5	problem 4.3 (e)	72
3.6	problem 4.3 (f)	73
3.7	problem 4.3 (g)	74
3.8	problem 4.3 (h)	75
3.9	problem 4.3 (i)	76
3.10	problem 4.3 (j)	78
3.11	problem 4.4 (a)	79
3.12	problem 4.4 (b)	80
3.13	problem 4.4 (c)	81
3.14	problem 4.4 (d)	82
3.15	problem 4.4 (e)	83
3.16	problem 4.4 (f)	84
3.17	problem 4.5 (a)	85
3.18	problem 4.5 (b)	86
3.19	problem 4.5 (c)	87
3.20	problem 4.5 (d)	88
3.21	problem 4.6 (a)	89
3.22	problem 4.6 (b)	90
3.23	problem 4.6 (c)	91
3.24	problem 4.6 (d)	92
3.25	problem 4.6 (e)	93
3.26	problem 4.6 (f)	94
3.27	problem 4.7 (a)	95
3.28	problem 4.7 (b)	96
3.29	problem 4.7 (c)	97
3.30	problem 4.7 (d)	98
3.31	problem 4.7 (e)	99
3.32	problem 4.7 (f)	100
3.33	problem 4.7 (g)	103
3.34	problem 4.7 (h)	104
3.35	problem 4.7 (i)	105

3.36	problem 4.7 (j)	106
3.37	problem 4.7 (k)	107
3.38	problem 4.7 (L)	108
3.39	problem 4.7 (m)	110
3.40	problem 4.7 (n)	111
3.41	problem 4.7 (o)	112
3.42	problem 4.8 (a)	113
3.43	problem 4.8 (b)	114
3.44	problem 4.8 (c)	115
3.45	problem 4.8 (d)	116
3.46	problem 4.8 (e)	117
3.47	problem 4.8 (f)	118
3.48	problem 4.8 (g)	119

### 3.1 problem 4.3 (a)

Internal problem ID [12979]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.3 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - 3y^2 + y^2 \sin(x) = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)=3*y(x)^2-y(x)^2*sin(x),y(x), singsol=all)
```

$$y = -\frac{1}{\cos(x) - c_1 + 3x}$$

#### ✓ Solution by Mathematica

Time used: 0.156 (sec). Leaf size: 22

```
DSolve[y'[x]==3*y[x]^2-y[x]^2*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{3x + \cos(x) + c_1}$$

$$y(x) \rightarrow 0$$

## 3.2 problem 4.3 (b)

Internal problem ID [12980]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.3 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y' + \sin(x)y = 3x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x)=3*x-y(x)*sin(x),y(x), singsol=all)
```

$$y = \left( \int 3e^{-\cos(x)} x dx + c_1 \right) e^{\cos(x)}$$

### ✓ Solution by Mathematica

Time used: 0.785 (sec). Leaf size: 31

```
DSolve[y'[x]==3*x-y[x]*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{\cos(x)} \left( \int_1^x 3e^{-\cos(K[1])} K[1] dK[1] + c_1 \right)$$

### 3.3 problem 4.3 (c)

Internal problem ID [12981]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.3 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_rational, _Riccati]`

$$y'x - (x - y)^2 = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 92

```
dsolve(x*diff(y(x),x)=(x-y(x))^2,y(x), singsol=all)
```

$$y = \frac{c_1 \sqrt{x} \text{BesselK}(1, 2\sqrt{x})}{\text{BesselK}(0, 2\sqrt{x}) c_1 + \text{BesselI}(0, 2\sqrt{x})} + \frac{(\sqrt{x} \text{BesselK}(0, 2\sqrt{x}) c_1 + \text{BesselI}(0, 2\sqrt{x}) \sqrt{x} - \text{BesselI}(1, 2\sqrt{x})) \sqrt{x}}{\text{BesselK}(0, 2\sqrt{x}) c_1 + \text{BesselI}(0, 2\sqrt{x})}$$

#### ✓ Solution by Mathematica

Time used: 0.251 (sec). Leaf size: 121

```
DSolve[x*y'[x]==(x-y[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x K_0(2\sqrt{x}) + \sqrt{x} K_1(2\sqrt{x}) + c_1 x \text{BesselI}(0, 2\sqrt{x}) - c_1 \sqrt{x} \text{BesselI}(1, 2\sqrt{x})}{K_0(2\sqrt{x}) + c_1 \text{BesselI}(0, 2\sqrt{x})}$$

$$y(x) \rightarrow x - \frac{\sqrt{x} \text{BesselI}(1, 2\sqrt{x})}{\text{BesselI}(0, 2\sqrt{x})}$$



### 3.4 problem 4.3 (d)

Internal problem ID [12982]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.3 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = \sqrt{x^2 + 1}$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(diff(y(x),x)=sqrt(1+x^2),y(x), singsol=all)
```

$$y = \frac{x\sqrt{x^2 + 1}}{2} + \frac{\operatorname{arcsinh}(x)}{2} + c_1$$

#### ✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 40

```
DSolve[y'[x]==Sqrt[1+x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}\sqrt{x^2 + 1}x - \frac{1}{2}\log(\sqrt{x^2 + 1} - x) + c_1$$

### 3.5 problem 4.3 (e)

Internal problem ID [12983]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.3 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' + 4y = 8$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)+4*y(x)=8,y(x), singsol=all)
```

$$y = 2 + c_1 e^{-4x}$$

#### ✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 20

```
DSolve[y'[x]+4*y[x]==8,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 + c_1 e^{-4x}$$

$$y(x) \rightarrow 2$$

### 3.6 problem 4.3 (f)

Internal problem ID [12984]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.3 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' + yx = 4x$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)+x*y(x)=4*x,y(x), singsol=all)
```

$$y = 4 + c_1 e^{-\frac{x^2}{2}}$$

#### ✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 24

```
DSolve[y'[x]+x*y[x]==4*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 4 + c_1 e^{-\frac{x^2}{2}}$$

$$y(x) \rightarrow 4$$

### 3.7 problem 4.3 (g)

Internal problem ID [12985]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.3 (g).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 4y = x^2$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x)+4*y(x)=x^2,y(x), singsol=all)
```

$$y = \frac{x^2}{4} - \frac{x}{8} + \frac{1}{32} + c_1 e^{-4x}$$

#### ✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 28

```
DSolve[y'[x]+4*y[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{32}(8x^2 - 4x + 1) + c_1 e^{-4x}$$

### 3.8 problem 4.3 (h)

Internal problem ID [12986]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.3 (h).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - yx + 2y = -3x + 6$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)=x*y(x)-3*x-2*y(x)+6,y(x), singsol=all)
```

$$y = 3 + c_1 e^{\frac{x(x-4)}{2}}$$

#### ✓ Solution by Mathematica

Time used: 0.079 (sec). Leaf size: 25

```
DSolve[y'[x]==x*y[x]-3*x-2*y[x]+6,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3 + c_1 e^{\frac{1}{2}(x-4)x}$$

$$y(x) \rightarrow 3$$

### 3.9 problem 4.3 (i)

Internal problem ID [12987]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.3 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$y' - \sin(x + y) = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 25

```
dsolve(diff(y(x),x)=sin(x+y(x)),y(x), singsol=all)
```

$$y = -x - 2 \arctan\left(\frac{c_1 - x - 2}{-x + c_1}\right)$$

✓ Solution by Mathematica

Time used: 35.052 (sec). Leaf size: 541

```
DSolve[y'[x]==Sin[x+y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2 \arccos \left( \frac{(x+c_1) \sin\left(\frac{x}{2}\right) - (x-2+c_1) \cos\left(\frac{x}{2}\right)}{\sqrt{2}\sqrt{x^2+2(-1+c_1)x+2+c_1^2-2c_1}} \right)$$

$$y(x) \rightarrow 2 \arccos \left( \frac{(x+c_1) \sin\left(\frac{x}{2}\right) - (x-2+c_1) \cos\left(\frac{x}{2}\right)}{\sqrt{2}\sqrt{x^2+2(-1+c_1)x+2+c_1^2-2c_1}} \right)$$

$$y(x) \rightarrow -2 \arccos \left( \frac{(x-2+c_1) \cos\left(\frac{x}{2}\right) - (x+c_1) \sin\left(\frac{x}{2}\right)}{\sqrt{2}\sqrt{x^2+2(-1+c_1)x+2+c_1^2-2c_1}} \right)$$

$$y(x) \rightarrow 2 \arccos \left( \frac{(x-2+c_1) \cos\left(\frac{x}{2}\right) - (x+c_1) \sin\left(\frac{x}{2}\right)}{\sqrt{2}\sqrt{x^2+2(-1+c_1)x+2+c_1^2-2c_1}} \right)$$

$$y(x) \rightarrow -2 \arccos \left( \frac{\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)}{\sqrt{2}} \right)$$

$$y(x) \rightarrow 2 \arccos \left( \frac{\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)}{\sqrt{2}} \right)$$

$$y(x) \rightarrow -2 \arccos \left( \frac{\sin\left(\frac{x}{2}\right) - \cos\left(\frac{x}{2}\right)}{\sqrt{2}} \right)$$

$$y(x) \rightarrow 2 \arccos \left( \frac{\sin\left(\frac{x}{2}\right) - \cos\left(\frac{x}{2}\right)}{\sqrt{2}} \right)$$

$$y(x) \rightarrow -2 \arccos \left( \frac{(x-2) \cos\left(\frac{x}{2}\right) - x \sin\left(\frac{x}{2}\right)}{\sqrt{2}\sqrt{x^2-2x+2}} \right)$$

$$y(x) \rightarrow 2 \arccos \left( \frac{(x-2) \cos\left(\frac{x}{2}\right) - x \sin\left(\frac{x}{2}\right)}{\sqrt{2}\sqrt{x^2-2x+2}} \right)$$

$$y(x) \rightarrow -2 \arccos \left( \frac{x \sin\left(\frac{x}{2}\right) - (x-2) \cos\left(\frac{x}{2}\right)}{\sqrt{2}\sqrt{x^2-2x+2}} \right)$$

$$y(x) \rightarrow 2 \arccos \left( \frac{x \sin\left(\frac{x}{2}\right) - (x-2) \cos\left(\frac{x}{2}\right)}{\sqrt{2}\sqrt{x^2-2x+2}} \right)$$

### 3.10 problem 4.3 (j)

Internal problem ID [12988]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.3 (j).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$yy' - e^{x-3y^2} = 0$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 39

```
dsolve(y(x)*diff(y(x),x)=exp(x-3*y(x)^2),y(x), singsol=all)
```

$$y = -\frac{\sqrt{3} \sqrt{\ln(6e^x + 6c_1)}}{3}$$

$$y = \frac{\sqrt{3} \sqrt{\ln(6e^x + 6c_1)}}{3}$$

#### ✓ Solution by Mathematica

Time used: 3.781 (sec). Leaf size: 48

```
DSolve[y[x]*y'[x]==Exp[x-3*y[x]^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{\log(6(e^x + c_1))}}{\sqrt{3}}$$

$$y(x) \rightarrow \frac{\sqrt{\log(6(e^x + c_1))}}{\sqrt{3}}$$



### 3.11 problem 4.4 (a)

Internal problem ID [12989]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.4 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - \frac{x}{y} = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x)=x/y(x),y(x), singsol=all)
```

$$y = \sqrt{x^2 + c_1}$$
$$y = -\sqrt{x^2 + c_1}$$

#### ✓ Solution by Mathematica

Time used: 0.088 (sec). Leaf size: 35

```
DSolve[y'[x]==x/y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{x^2 + 2c_1}$$
$$y(x) \rightarrow \sqrt{x^2 + 2c_1}$$

### 3.12 problem 4.4 (b)

Internal problem ID [12990]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.4 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - y^2 = 9$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)=y(x)^2+9,y(x), singsol=all)
```

$$y = 3 \tan(3x + 3c_1)$$

#### ✓ Solution by Mathematica

Time used: 0.199 (sec). Leaf size: 28

```
DSolve[y'[x]==y[x]^2+9,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3 \tan(3(x + c_1))$$

$$y(x) \rightarrow -3i$$

$$y(x) \rightarrow 3i$$

### 3.13 problem 4.4 (c)

Internal problem ID [12991]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.4 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y'xy - y^2 = 9$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(x*y(x)*diff(y(x),x)=y(x)^2+9,y(x), singsol=all)
```

$$y = \sqrt{c_1 x^2 - 9}$$

$$y = -\sqrt{c_1 x^2 - 9}$$

#### ✓ Solution by Mathematica

Time used: 0.401 (sec). Leaf size: 57

```
DSolve[x*y[x]*y'[x]==y[x]^2+9,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{-9 + e^{2c_1} x^2}$$

$$y(x) \rightarrow \sqrt{-9 + e^{2c_1} x^2}$$

$$y(x) \rightarrow -3i$$

$$y(x) \rightarrow 3i$$

### 3.14 problem 4.4 (d)

Internal problem ID [12992]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.4 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - \frac{y^2 + 1}{x^2 + 1} = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)=(y(x)^2+1)/(x^2+1),y(x), singsol=all)
```

$$y = \tan(\arctan(x) + c_1)$$

#### ✓ Solution by Mathematica

Time used: 0.225 (sec). Leaf size: 25

```
DSolve[y'[x]==(y[x]^2+1)/(x^2+1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan(\arctan(x) + c_1)$$

$$y(x) \rightarrow -i$$

$$y(x) \rightarrow i$$

### 3.15 problem 4.4 (e)

Internal problem ID [12993]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.4 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$\cos(y) y' = \sin(x)$$

#### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 11

```
dsolve(cos(y(x))*diff(y(x),x)=sin(x),y(x), singsol=all)
```

$$y = \arcsin(-\cos(x) + c_1)$$

#### ✓ Solution by Mathematica

Time used: 0.311 (sec). Leaf size: 13

```
DSolve[Cos[y[x]]*y'[x]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arcsin(-\cos(x) + c_1)$$

### 3.16 problem 4.4 (f)

Internal problem ID [12994]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.4 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y' - e^{2x-3y} = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)=exp(2*x-3*y(x)),y(x), singsol=all)
```

$$y = \frac{\ln\left(\frac{3e^{2x}}{2} + 3c_1\right)}{3}$$

#### ✓ Solution by Mathematica

Time used: 0.855 (sec). Leaf size: 24

```
DSolve[y'[x]==Exp[2*x-3*y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} \log\left(\frac{3}{2}(e^{2x} + 2c_1)\right)$$

### 3.17 problem 4.5 (a)

Internal problem ID [12995]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.5 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - \frac{x}{y} = 0$$

With initial conditions

$$[y(1) = 3]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 11

```
dsolve([diff(y(x),x)=x/y(x),y(1) = 3],y(x), singsol=all)
```

$$y = \sqrt{x^2 + 8}$$

✓ Solution by Mathematica

Time used: 0.11 (sec). Leaf size: 14

```
DSolve[{y'[x]==x/y[x],{y[1]==3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt{x^2 + 8}$$

### 3.18 problem 4.5 (b)

Internal problem ID [12996]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.5 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - 2yx + y = 2x - 1$$

With initial conditions

$$[y(0) = 2]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([diff(y(x),x)=2*x-1+2*x*y(x)-y(x),y(0) = 2],y(x), singsol=all)
```

$$y = -1 + 3e^{x(-1+x)}$$

✓ Solution by Mathematica

Time used: 0.053 (sec). Leaf size: 16

```
DSolve[{y'[x]==2*x-1+2*x*y[x]-y[x],{y[0]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3e^{(x-1)x} - 1$$



### 3.19 problem 4.5 (c)

Internal problem ID [12997]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.5 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$yy' - xy^2 = x$$

With initial conditions

$$[y(0) = -2]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve([y(x)*diff(y(x),x)=x*y(x)^2+x,y(0) = -2],y(x), singsol=all)
```

$$y = -\sqrt{5e^{x^2} - 1}$$

✓ Solution by Mathematica

Time used: 7.0 (sec). Leaf size: 20

```
DSolve[{y[x]*y'[x]==x*y[x]^2+x,{y[0]==-2}],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{5e^{x^2} - 1}$$

### 3.20 problem 4.5 (d)

Internal problem ID [12998]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.5 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x),G(x)*y+H(x)]']]`

$$yy' - 3\sqrt{xy^2 + 9x} = 0$$

With initial conditions

$$[y(1) = 4]$$

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 17

```
dsolve([y(x)*diff(y(x),x)=3*sqrt(x*y(x)^2+9*x),y(1) = 4],y(x), singsol=all)
```

$$y = 2\sqrt{x^{\frac{3}{2}}(x^{\frac{3}{2}} + 3)}$$

✓ Solution by Mathematica

Time used: 0.294 (sec). Leaf size: 44

```
DSolve[{y[x]*y'[x]==3*Sqrt[x*y[x]^2+9*x],{y[1]==4}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2\sqrt{3x^{3/2} + x^3}$$

$$y(x) \rightarrow 2\sqrt{-7x^{3/2} + x^3 + 10}$$

### 3.21 problem 4.6 (a)

Internal problem ID [12999]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.6 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - yx = -4x$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)=x*y(x)-4*x,y(x), singsol=all)
```

$$y = 4 + c_1 e^{\frac{x^2}{2}}$$

#### ✓ Solution by Mathematica

Time used: 0.062 (sec). Leaf size: 24

```
DSolve[y' [x]==x*y[x]-4*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 4 + c_1 e^{\frac{x^2}{2}}$$

$$y(x) \rightarrow 4$$

### 3.22 problem 4.6 (b)

Internal problem ID [13000]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.6 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - 4y = 2$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)-4*y(x)=2,y(x), singsol=all)
```

$$y = -\frac{1}{2} + e^{4x}c_1$$

#### ✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 24

```
DSolve[y'[x]-4*y[x]==2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2} + c_1 e^{4x}$$

$$y(x) \rightarrow -\frac{1}{2}$$

### 3.23 problem 4.6 (c)

Internal problem ID [13001]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.6 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$yy' - xy^2 = -9x$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(y(x)*diff(y(x),x)=x*y(x)^2-9*x,y(x), singsol=all)
```

$$y = \sqrt{9 + c_1 e^{x^2}}$$

$$y = -\sqrt{9 + c_1 e^{x^2}}$$

#### ✓ Solution by Mathematica

Time used: 1.856 (sec). Leaf size: 53

```
DSolve[y[x]*y'[x]==x*y[x]^2-9*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{9 + e^{x^2+2c_1}}$$

$$y(x) \rightarrow \sqrt{9 + e^{x^2+2c_1}}$$

$$y(x) \rightarrow -3$$

$$y(x) \rightarrow 3$$

### 3.24 problem 4.6 (d)

Internal problem ID [13002]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.6 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - \sin(y) = 0$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 46

```
dsolve(diff(y(x),x)=sin(y(x)),y(x), singsol=all)
```

$$y = \arctan\left(\frac{2e^x c_1}{e^{2x} c_1^2 + 1}, -\frac{e^{2x} c_1^2 - 1}{e^{2x} c_1^2 + 1}\right)$$

#### ✓ Solution by Mathematica

Time used: 0.293 (sec). Leaf size: 44

```
DSolve[y'[x]==Sin[y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\arccos(-\tanh(x + c_1))$$

$$y(x) \rightarrow \arccos(-\tanh(x + c_1))$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -\pi$$

$$y(x) \rightarrow \pi$$

### 3.25 problem 4.6 (e)

Internal problem ID [13003]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.6 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - e^{y^2+x} = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)=exp(x+y(x)^2),y(x), singsol=all)
```

$$e^x - \frac{\operatorname{erf}(y)\sqrt{\pi}}{2} + c_1 = 0$$

#### ✓ Solution by Mathematica

Time used: 0.439 (sec). Leaf size: 19

```
DSolve[y'[x]==Exp[x+y[x]^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \operatorname{erf}^{-1}\left(\frac{2(e^x + c_1)}{\sqrt{\pi}}\right)$$

### 3.26 problem 4.6 (f)

Internal problem ID [13004]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.6 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - 200y + 2y^2 = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)=200*y(x)-2*y(x)^2,y(x), singsol=all)
```

$$y = \frac{100}{1 + 100e^{-200x}c_1}$$

#### ✓ Solution by Mathematica

Time used: 0.285 (sec). Leaf size: 36

```
DSolve[y'[x]==200*y[x]-2*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{100e^{200x}}{e^{200x} + e^{100c_1}}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow 100$$



### 3.27 problem 4.7 (a)

Internal problem ID [13005]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.7 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - xy = -4x$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)=x*y(x)-4*x,y(x), singsol=all)
```

$$y = 4 + c_1 e^{\frac{x^2}{2}}$$

#### ✓ Solution by Mathematica

Time used: 0.111 (sec). Leaf size: 24

```
DSolve[y'[x]==x*y[x]-4*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 4 + c_1 e^{\frac{x^2}{2}}$$

$$y(x) \rightarrow 4$$

### 3.28 problem 4.7 (b)

Internal problem ID [13006]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.7 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - xy + 2y = -3x + 6$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)=x*y(x)-3*x-2*y(x)+6,y(x), singsol=all)
```

$$y = 3 + e^{\frac{x(x-4)}{2}} c_1$$

#### ✓ Solution by Mathematica

Time used: 0.077 (sec). Leaf size: 25

```
DSolve[y'[x]==x*y[x]-3*x-2*y[x]+6,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3 + c_1 e^{\frac{1}{2}(x-4)x}$$

$$y(x) \rightarrow 3$$

### 3.29 problem 4.7 (c)

Internal problem ID [13007]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.7 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - 3y^2 + y^2 \sin(x) = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)=3*y(x)^2-y(x)^2*sin(x),y(x), singsol=all)
```

$$y = -\frac{1}{\cos(x) - c_1 + 3x}$$

#### ✓ Solution by Mathematica

Time used: 0.173 (sec). Leaf size: 22

```
DSolve[y'[x]==3*y[x]^2-y[x]^2*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{3x + \cos(x) + c_1}$$

$$y(x) \rightarrow 0$$

### 3.30 problem 4.7 (d)

Internal problem ID [13008]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.7 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - \tan(y) = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)=tan(y(x)),y(x), singsol=all)
```

$$y = \arcsin(c_1 e^x)$$

#### ✓ Solution by Mathematica

Time used: 40.257 (sec). Leaf size: 17

```
DSolve[y'[x]==Tan[y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arcsin(e^{x+c_1})$$

$$y(x) \rightarrow 0$$

### 3.31 problem 4.7 (e)

Internal problem ID [13009]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.7 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - \frac{y}{x} = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 7

```
dsolve(diff(y(x),x)=y(x)/x,y(x), singsol=all)
```

$$y = c_1x$$

#### ✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 14

```
DSolve[y'[x]==y[x]/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x$$

$$y(x) \rightarrow 0$$

### 3.32 problem 4.7 (f)

Internal problem ID [13010]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.7 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_separable`]

$$y' - \frac{6x^2 + 4}{3y^2 - 4y} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 660

```
dsolve(diff(y(x),x)=(6*x^2+4)/(3*y(x)^2-4*y(x)),y(x), singsol=all)
```

$$\begin{aligned}
 & y \\
 & = \frac{\left(8 + 27x^3 + 27c_1 + 54x + 3\sqrt{81x^6 + 162c_1x^3 + 324x^4 + 48x^3 + 81c_1^2 + 324c_1x + 324x^2 + 48c_1 + 96x}\right)}{3} \\
 & + \frac{4}{3} \left(8 + 27x^3 + 27c_1 + 54x + 3\sqrt{81x^6 + 162c_1x^3 + 324x^4 + 48x^3 + 81c_1^2 + 324c_1x + 324x^2 + 48c_1 + 96x}\right) \\
 & + \frac{2}{3} \\
 & y = \\
 & - \frac{\left(8 + 27x^3 + 27c_1 + 54x + 3\sqrt{81x^6 + 162c_1x^3 + 324x^4 + 48x^3 + 81c_1^2 + 324c_1x + 324x^2 + 48c_1 + 96x}\right)}{6} \\
 & - \frac{2}{3} \\
 & 3 \left(8 + 27x^3 + 27c_1 + 54x + 3\sqrt{81x^6 + 162c_1x^3 + 324x^4 + 48x^3 + 81c_1^2 + 324c_1x + 324x^2 + 48c_1 + 96x}\right) \\
 & + \frac{2}{3} \\
 & i\sqrt{3} \left( \frac{\left(8 + 27x^3 + 27c_1 + 54x + 3\sqrt{81x^6 + 162c_1x^3 + 324x^4 + 48x^3 + 81c_1^2 + 324c_1x + 324x^2 + 48c_1 + 96x}\right)^{\frac{1}{3}}}{3} - \frac{3 \left(8 + 27x^3 + 27c_1 + 54x + 3\sqrt{81x^6 + 162c_1x^3 + 324x^4 + 48x^3 + 81c_1^2 + 324c_1x + 324x^2 + 48c_1 + 96x}\right)^{\frac{1}{3}}}{2} \right) \\
 & y = \\
 & - \frac{\left(8 + 27x^3 + 27c_1 + 54x + 3\sqrt{81x^6 + 162c_1x^3 + 324x^4 + 48x^3 + 81c_1^2 + 324c_1x + 324x^2 + 48c_1 + 96x}\right)}{6} \\
 & - \frac{2}{3} \\
 & 3 \left(8 + 27x^3 + 27c_1 + 54x + 3\sqrt{81x^6 + 162c_1x^3 + 324x^4 + 48x^3 + 81c_1^2 + 324c_1x + 324x^2 + 48c_1 + 96x}\right) \\
 & + \frac{2}{3} \\
 & i\sqrt{3} \left( \frac{\left(8 + 27x^3 + 27c_1 + 54x + 3\sqrt{81x^6 + 162c_1x^3 + 324x^4 + 48x^3 + 81c_1^2 + 324c_1x + 324x^2 + 48c_1 + 96x}\right)^{\frac{1}{3}}}{3} - \frac{3 \left(8 + 27x^3 + 27c_1 + 54x + 3\sqrt{81x^6 + 162c_1x^3 + 324x^4 + 48x^3 + 81c_1^2 + 324c_1x + 324x^2 + 48c_1 + 96x}\right)^{\frac{1}{3}}}{2} \right) \\
 & + \frac{2}{3}
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 2.967 (sec). Leaf size: 356

`DSolve[y'[x]==(6*x^2+4)/(3*y[x]^2-4*y[x]),y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{1}{6} \left( 2^{2/3} \sqrt[3]{54x^3 + \sqrt{-256 + (54x^3 + 108x + 16 + 27c_1)^2} + 108x + 16 + 27c_1} + \frac{8\sqrt[3]{2}}{\sqrt[3]{54x^3 + \sqrt{-256 + (54x^3 + 108x + 16 + 27c_1)^2} + 108x + 16 + 27c_1}} + 4 \right)$$

$$y(x) \rightarrow \frac{1}{12} \left( i2^{2/3} (\sqrt{3} + i) \sqrt[3]{54x^3 + \sqrt{-256 + (54x^3 + 108x + 16 + 27c_1)^2} + 108x + 16 + 27c_1} - \frac{8\sqrt[3]{2}(1 + i\sqrt{3})}{\sqrt[3]{54x^3 + \sqrt{-256 + (54x^3 + 108x + 16 + 27c_1)^2} + 108x + 16 + 27c_1}} + 8 \right)$$

$$y(x) \rightarrow \frac{1}{12} \left( -2^{2/3} (1 + i\sqrt{3}) \sqrt[3]{54x^3 + \sqrt{-256 + (54x^3 + 108x + 16 + 27c_1)^2} + 108x + 16 + 27c_1} + \frac{8i\sqrt[3]{2}(\sqrt{3} + i)}{\sqrt[3]{54x^3 + \sqrt{-256 + (54x^3 + 108x + 16 + 27c_1)^2} + 108x + 16 + 27c_1}} + 8 \right)$$



### 3.33 problem 4.7 (g)

Internal problem ID [13011]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.7 (g).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$(x^2 + 1) y' - y^2 = 1$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve((x^2+1)*diff(y(x),x)=y(x)^2+1,y(x), singsol=all)
```

$$y = \tan(\arctan(x) + c_1)$$

#### ✓ Solution by Mathematica

Time used: 0.208 (sec). Leaf size: 25

```
DSolve[(x^2+1)*y'[x]==y[x]^2+1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan(\arctan(x) + c_1)$$

$$y(x) \rightarrow -i$$

$$y(x) \rightarrow i$$

### 3.34 problem 4.7 (h)

Internal problem ID [13012]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.7 (h).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$(y^2 - 1)y' - 4y^2x = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 59

```
dsolve((y(x)^2-1)*diff(y(x),x)=4*x*y(x)^2,y(x), singsol=all)
```

$$y = x^2 + 2c_1 - \sqrt{x^4 + 4c_1x^2 + 4c_1^2 - 1}$$

$$y = x^2 + 2c_1 + \sqrt{x^4 + 4c_1x^2 + 4c_1^2 - 1}$$

#### ✓ Solution by Mathematica

Time used: 0.273 (sec). Leaf size: 84

```
DSolve[(y[x]^2-1)*y'[x]==4*x*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left( 2x^2 - \sqrt{4x^4 + 4c_1x^2 - 4 + c_1^2} + c_1 \right)$$

$$y(x) \rightarrow \frac{1}{2} \left( 2x^2 + \sqrt{4x^4 + 4c_1x^2 - 4 + c_1^2} + c_1 \right)$$

$$y(x) \rightarrow 0$$

### 3.35 problem 4.7 (i)

Internal problem ID [13013]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.7 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - e^{-y} = 0$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 8

```
dsolve(diff(y(x),x)=exp(-y(x)),y(x), singsol=all)
```

$$y = \ln(x + c_1)$$

#### ✓ Solution by Mathematica

Time used: 0.196 (sec). Leaf size: 10

```
DSolve[y'[x]==Exp[-y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(x + c_1)$$

### 3.36 problem 4.7 (j)

Internal problem ID [13014]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.7 (j).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - e^{-y} = 1$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(diff(y(x),x)=exp(-y(x))+1,y(x), singsol=all)
```

$$y = \ln(c_1 e^x - 1)$$

#### ✓ Solution by Mathematica

Time used: 1.163 (sec). Leaf size: 32

```
DSolve[y'[x]==Exp[-y[x]]+1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(-1 + e^{x+c_1})$$

$$y(x) \rightarrow -i\pi$$

$$y(x) \rightarrow i\pi$$

### 3.37 problem 4.7 (k)

Internal problem ID [13015]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.7 (k).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - 3xy^3 = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x)=3*x*y(x)^3,y(x), singsol=all)
```

$$y = \frac{1}{\sqrt{-3x^2 + c_1}}$$
$$y = -\frac{1}{\sqrt{-3x^2 + c_1}}$$

#### ✓ Solution by Mathematica

Time used: 0.158 (sec). Leaf size: 44

```
DSolve[y'[x]==3*x*y[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\sqrt{-3x^2 - 2c_1}}$$

$$y(x) \rightarrow \frac{1}{\sqrt{-3x^2 - 2c_1}}$$

$$y(x) \rightarrow 0$$

### 3.38 problem 4.7 (L)

Internal problem ID [13016]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.7 (L).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - \frac{2 + \sqrt{x}}{2 + \sqrt{y}} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(diff(y(x),x)=(2+sqrt(x))/(2+sqrt(y(x))),y(x), singsol=all)
```

$$\frac{2x^{\frac{3}{2}}}{3} + 2x - 2y - \frac{2y^{\frac{3}{2}}}{3} + c_1 = 0$$

✓ Solution by Mathematica

Time used: 3.607 (sec). Leaf size: 1162

`DSolve[y'[x]==(2+Sqrt[x])/(2+Sqrt[y[x]]),y[x],x,IncludeSingularSolutions -> True]`

$y(x)$

$$\rightarrow \frac{-8x^{3/2} + \left(24x^{5/2} + 12(-6 + c_1)x^{3/2} + \sqrt{(2x^{3/2} + 6x - 8 + 3c_1)(2x^{3/2} + 6x + 3c_1)^3} + 4x^3 + 36x^2 + 3\right)}{2\sqrt[3]{24x^{5/2} + 12(-6 + c_1)x^{3/2} + \sqrt{(2x^{3/2} + 6x - 8 + 3c_1)(2x^{3/2} + 6x + 3c_1)^3} + 4x^3 + 36x^2 + 3}}$$

$y(x)$

$$\rightarrow \frac{(8 + 8i\sqrt{3})x^{3/2} + i\sqrt{3}\left(24x^{5/2} + 12(-6 + c_1)x^{3/2} + \sqrt{(2x^{3/2} + 6x - 8 + 3c_1)(2x^{3/2} + 6x + 3c_1)^3} + 4x^3 + 36x^2 + 3\right)}{2\sqrt[3]{24x^{5/2} + 12(-6 + c_1)x^{3/2} + \sqrt{(2x^{3/2} + 6x - 8 + 3c_1)(2x^{3/2} + 6x + 3c_1)^3} + 4x^3 + 36x^2 + 3}}$$

$y(x)$

$$\rightarrow \frac{(8 - 8i\sqrt{3})x^{3/2} - i\sqrt{3}\left(24x^{5/2} + 12(-6 + c_1)x^{3/2} + \sqrt{(2x^{3/2} + 6x - 8 + 3c_1)(2x^{3/2} + 6x + 3c_1)^3} + 4x^3 + 36x^2 + 3\right)}{2\sqrt[3]{24x^{5/2} + 12(-6 + c_1)x^{3/2} + \sqrt{(2x^{3/2} + 6x - 8 + 3c_1)(2x^{3/2} + 6x + 3c_1)^3} + 4x^3 + 36x^2 + 3}}$$

### 3.39 problem 4.7 (m)

Internal problem ID [13017]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.7 (m).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - 3y^2x^2 = -3x^2$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)-3*x^2*y(x)^2=-3*x^2,y(x), singsol=all)
```

$$y = -\tanh(x^3 + 3c_1)$$

#### ✓ Solution by Mathematica

Time used: 0.45 (sec). Leaf size: 44

```
DSolve[y'[x]-3*x^2*y[x]^2==-3*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1 - e^{2(x^3+c_1)}}{1 + e^{2(x^3+c_1)}}$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 1$$



### 3.40 problem 4.7 (n)

Internal problem ID [13018]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.7 (n).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - 3y^2x^2 = 3x^2$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)-3*x^2*y(x)^2=3*x^2,y(x), singsol=all)
```

$$y = \tan(x^3 + 3c_1)$$

#### ✓ Solution by Mathematica

Time used: 0.152 (sec). Leaf size: 26

```
DSolve[y'[x]-3*x^2*y[x]^2==3*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan(x^3 + c_1)$$

$$y(x) \rightarrow -i$$

$$y(x) \rightarrow i$$

### 3.41 problem 4.7 (o)

Internal problem ID [13019]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.7 (o).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - 200y + 2y^2 = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)=200*y(x)-2*y(x)^2,y(x), singsol=all)
```

$$y = \frac{100}{1 + 100e^{-200x}c_1}$$

#### ✓ Solution by Mathematica

Time used: 0.258 (sec). Leaf size: 36

```
DSolve[y'[x]==200*y[x]-2*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{100e^{200x}}{e^{200x} + e^{100c_1}}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow 100$$

### 3.42 problem 4.8 (a)

Internal problem ID [13020]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.8 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - 2y = -10$$

With initial conditions

$$[y(0) = 8]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve([diff(y(x),x)-2*y(x)=-10,y(0) = 8],y(x), singsol=all)
```

$$y = 3e^{2x} + 5$$

✓ Solution by Mathematica

Time used: 0.055 (sec). Leaf size: 14

```
DSolve[{y'[x]-2*y[x]==-10,{y[0]==8}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3e^{2x} + 5$$

### 3.43 problem 4.8 (b)

Internal problem ID [13021]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.8 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$yy' = \sin(x)$$

With initial conditions

$$[y(0) = -4]$$

✓ Solution by Maple

Time used: 0.14 (sec). Leaf size: 14

```
dsolve([y(x)*diff(y(x),x)=sin(x),y(0) = -4],y(x), singsol=all)
```

$$y = -\sqrt{-2 \cos(x) + 18}$$

✓ Solution by Mathematica

Time used: 0.11 (sec). Leaf size: 22

```
DSolve[{y[x]*y'[x]==Sin[x],{y[0]==-4}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{2}\sqrt{9 - \cos(x)}$$

### 3.44 problem 4.8 (c)

Internal problem ID [13022]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.8 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - 2xy + y = 2x - 1$$

With initial conditions

$$[y(0) = -1]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve([diff(y(x),x)=2*x-1+2*x*y(x)-y(x),y(0) = -1],y(x), singsol=all)
```

$$y = -1$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 6

```
DSolve[{y'[x]==2*x-1+2*x*y[x]-y[x],{y[0]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -1$$

### 3.45 problem 4.8 (d)

Internal problem ID [13023]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.8 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y'x - y^2 + y = 0$$

With initial conditions

$$[y(2) = 1]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve([x*diff(y(x),x)=y(x)^2-y(x),y(2) = 1],y(x), singsol=all)
```

$$y = 1$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 6

```
DSolve[{x*y'[x]==y[x]^2-y[x],{y[2]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 1$$

### 3.46 problem 4.8 (e)

Internal problem ID [13024]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.8 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y'x - y^2 + y = 0$$

With initial conditions

$$[y(1) = 2]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 11

```
dsolve([x*diff(y(x),x)=y(x)^2-y(x),y(1) = 2],y(x), singsol=all)
```

$$y = -\frac{2}{x-2}$$

✓ Solution by Mathematica

Time used: 0.229 (sec). Leaf size: 12

```
DSolve[{x*y'[x]==y[x]^2-y[x],{y[1]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2}{x-2}$$

### 3.47 problem 4.8 (f)

Internal problem ID [13025]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.8 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - \frac{y^2 - 1}{xy} = 0$$

With initial conditions

$$[y(1) = -2]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 15

```
dsolve([diff(y(x),x)=(y(x)^2-1)/(x*y(x)),y(1) = -2],y(x), singsol=all)
```

$$y = -\sqrt{3x^2 + 1}$$

✓ Solution by Mathematica

Time used: 0.248 (sec). Leaf size: 18

```
DSolve[{y'[x]==(y[x]^2-1)/(x*y[x]),{y[1]==-2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{3x^2 + 1}$$



### 3.48 problem 4.8 (g)

Internal problem ID [13026]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 4. SEPARABLE FIRST ORDER EQUATIONS. Additional exercises. page 90

**Problem number:** 4.8 (g).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$(y^2 - 1) y' - 4xy = 0$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 25

```
dsolve([(y(x)^2-1)*diff(y(x),x)=4*x*y(x),y(0) = 1],y(x), singsol=all)
```

$$y = \frac{e^{-2x^2 - \frac{1}{2}}}{\sqrt{-\frac{e^{-4x^2 - 1}}{\text{LambertW}(-e^{-4x^2 - 1})}}}$$

✓ Solution by Mathematica

Time used: 4.197 (sec). Leaf size: 25

```
DSolve[{(y[x]^2-1)*y'[x]==4*x*y[x],{y[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -i\sqrt{W(-e^{-4x^2-1})}$$

## 4 Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

4.1	problem 5.1 (a) . . . . .	121
4.2	problem 5.1 (b) . . . . .	122
4.3	problem 5.1 (c) . . . . .	123
4.4	problem 5.1 (d) . . . . .	124
4.5	problem 5.1 (e) . . . . .	125
4.6	problem 5.1 (f) . . . . .	126
4.7	problem 5.1 (g) . . . . .	127
4.8	problem 5.1 (h) . . . . .	128
4.9	problem 5.1 (i) . . . . .	129
4.10	problem 5.1 (j) . . . . .	131
4.11	problem 5.2 (a) . . . . .	132
4.12	problem 5.2 (b) . . . . .	133
4.13	problem 5.2 (c) . . . . .	134
4.14	problem 5.2 (d) . . . . .	135
4.15	problem 5.2 (e) . . . . .	136
4.16	problem 5.2 (f) . . . . .	137
4.17	problem 5.2 (g) . . . . .	138
4.18	problem 5.2 (h) . . . . .	139
4.19	problem 5.2 (i) . . . . .	140
4.20	problem 5.2 (j) . . . . .	141
4.21	problem 5.3 (a) . . . . .	142
4.22	problem 5.3 (b) . . . . .	143
4.23	problem 5.3 (c) . . . . .	144
4.24	problem 5.3 (d) . . . . .	145
4.25	problem 5.3 (e) . . . . .	146
4.26	problem 5.3 (f) . . . . .	147
4.27	problem 5.4 (a) . . . . .	148
4.28	problem 5.4 (b) . . . . .	149
4.29	problem 5.4 (c) . . . . .	150

## 4.1 problem 5.1 (a)

Internal problem ID [13027]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.1 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y'x^2 + 3x^2y = \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 73

```
dsolve(x^2*diff(y(x),x)+3*x^2*y(x)=sin(x),y(x), singsol=all)
```

$$y = \left( \frac{ie^{(3+i)x}}{2x} - \frac{\expIntegral_1((-3-i)x)}{2} + \frac{3i \expIntegral_1((-3-i)x)}{2} - \frac{ie^{(3-i)x}}{2x} - \frac{\expIntegral_1((-3+i)x)}{2} - \frac{3i \expIntegral_1((-3+i)x)}{2} + c_1 \right) e^{-3x}$$

### ✓ Solution by Mathematica

Time used: 0.104 (sec). Leaf size: 68

```
DSolve[x^2*y'[x]+3*x^2*y[x]==Sin[x],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{2}e^{-3x} \left( (1+3i) \text{ExpIntegralEi}((3-i)x) + \frac{(1-3i)x \text{ExpIntegralEi}((3+i)x) - ie^{(3-i)x} + ie^{(3+i)x} + 2c_1x}{x} \right)$$

## 4.2 problem 5.1 (b)

Internal problem ID [13028]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.1 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [ $y = G(x, y')$ ]

$$y^2 y' + 3x^2 y = \sin(x)$$

**X** Solution by Maple

```
dsolve(y(x)^2*diff(y(x),x)+3*x^2*y(x)=sin(x),y(x), singsol=all)
```

No solution found

**X** Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y[x]^2*y'[x]+3*x^2*y[x]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

Not solved

### 4.3 problem 5.1 (c)

Internal problem ID [13029]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.1 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [Riccati]

$$y' - y^2x = \sqrt{x}$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 45

```
dsolve(diff(y(x),x)-x*y(x)^2=sqrt(x),y(x), singsol=all)
```

$$y = -\frac{\text{BesselY}\left(-\frac{3}{7}, \frac{4x^{7/4}}{7}\right) c_1 + \text{BesselJ}\left(-\frac{3}{7}, \frac{4x^{7/4}}{7}\right)}{x^{1/4} \left(\text{BesselY}\left(\frac{4}{7}, \frac{4x^{7/4}}{7}\right) c_1 + \text{BesselJ}\left(\frac{4}{7}, \frac{4x^{7/4}}{7}\right)\right)}$$

#### ✓ Solution by Mathematica

Time used: 0.242 (sec). Leaf size: 273

```
DSolve[y'[x]-x*y[x]^2==Sqrt[x],y[x],x,IncludeSingularSolutions -> True]
```

$y(x) \rightarrow$

$$\frac{x^{7/4} \Gamma\left(\frac{11}{7}\right) \text{BesselJ}\left(-\frac{3}{7}, \frac{4x^{7/4}}{7}\right) - x^{7/4} \Gamma\left(\frac{11}{7}\right) \text{BesselJ}\left(\frac{11}{7}, \frac{4x^{7/4}}{7}\right) + 2 \Gamma\left(\frac{11}{7}\right) \text{BesselJ}\left(\frac{11}{7}, \frac{4x^{7/4}}{7}\right)}{2x^2 \left(\Gamma\left(\frac{11}{7}\right) \text{BesselJ}\left(\frac{11}{7}, \frac{4x^{7/4}}{7}\right) - \Gamma\left(\frac{11}{7}\right) \text{BesselJ}\left(-\frac{3}{7}, \frac{4x^{7/4}}{7}\right)\right)}$$

$$y(x) \rightarrow -\frac{x^{7/4} \text{BesselJ}\left(-\frac{11}{7}, \frac{4x^{7/4}}{7}\right) - x^{7/4} \text{BesselJ}\left(\frac{3}{7}, \frac{4x^{7/4}}{7}\right) + 2 \text{BesselJ}\left(-\frac{4}{7}, \frac{4x^{7/4}}{7}\right)}{2x^2 \text{BesselJ}\left(-\frac{4}{7}, \frac{4x^{7/4}}{7}\right)}$$

## 4.4 problem 5.1 (d)

Internal problem ID [13030]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.1 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [Riccati]

$$y' - (xy + 3y)^2 = 1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 55

```
dsolve(diff(y(x),x)=1+(x*y(x)+3*y(x))^2,y(x), singsol=all)
```

$$y = -\frac{\text{BesselY}\left(-\frac{1}{4}, \frac{(x+3)^2}{2}\right) c_1 + \text{BesselJ}\left(-\frac{1}{4}, \frac{(x+3)^2}{2}\right)}{\left(\text{BesselY}\left(\frac{3}{4}, \frac{(x+3)^2}{2}\right) c_1 + \text{BesselJ}\left(\frac{3}{4}, \frac{(x+3)^2}{2}\right)\right)} (x + 3)$$

### ✓ Solution by Mathematica

Time used: 0.45 (sec). Leaf size: 351

```
DSolve[y'[x]==1+(x*y[x]+3*y[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$y(x) \rightarrow$

$$\frac{((x + 3)^3)^{2/3} \text{Gamma}\left(\frac{7}{4}\right) \text{BesselJ}\left(-\frac{1}{4}, \frac{1}{2}((x + 3)^3)^{2/3}\right) + 3 \text{Gamma}\left(\frac{7}{4}\right) \text{BesselJ}\left(\frac{3}{4}, \frac{1}{2}((x + 3)^3)^{2/3}\right)}{2(x + 3)^3 \text{BesselJ}\left(-\frac{3}{4}, \frac{1}{2}((x + 3)^3)^{2/3}\right)}$$

$y(x)$

$$\rightarrow \frac{-((x + 3)^3)^{2/3} \text{BesselJ}\left(-\frac{7}{4}, \frac{1}{2}((x + 3)^3)^{2/3}\right) - 3 \text{BesselJ}\left(-\frac{3}{4}, \frac{1}{2}((x + 3)^3)^{2/3}\right) + ((x + 3)^3)^{2/3} \text{BesselJ}\left(\frac{3}{4}, \frac{1}{2}((x + 3)^3)^{2/3}\right)}{2(x + 3)^3 \text{BesselJ}\left(-\frac{3}{4}, \frac{1}{2}((x + 3)^3)^{2/3}\right)}$$

## 4.5 problem 5.1 (e)

Internal problem ID [13031]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.1 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_linear]`

$$y' - xy - 3y = 1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 37

```
dsolve(diff(y(x),x)=1+x*y(x)+3*y(x),y(x), singsol=all)
```

$$y = \left( \frac{\sqrt{\pi} e^{\frac{9}{2}} \sqrt{2} \operatorname{erf}\left(\frac{\sqrt{2}x}{2} + \frac{3\sqrt{2}}{2}\right)}{2} + c_1 \right) e^{\frac{x(x+6)}{2}}$$

### ✓ Solution by Mathematica

Time used: 0.054 (sec). Leaf size: 47

```
DSolve[y'[x]==1+x*y[x]+3*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^{\frac{1}{2}x(x+6)} \left( e^{9/2} \sqrt{2\pi} \operatorname{erf}\left(\frac{x+3}{\sqrt{2}}\right) + 2c_1 \right)$$

## 4.6 problem 5.1 (f)

Internal problem ID [13032]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.1 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - 4y = 8$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=4*y(x)+8,y(x), singsol=all)
```

$$y = -2 + c_1 e^{4x}$$

### ✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 20

```
DSolve[y'[x]==4*y[x]+8,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2 + c_1 e^{4x}$$

$$y(x) \rightarrow -2$$



## 4.7 problem 5.1 (g)

Internal problem ID [13033]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.1 (g).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' = e^{2x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)-exp(2*x)=0,y(x), singsol=all)
```

$$y = \frac{e^{2x}}{2} + c_1$$

### ✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 17

```
DSolve[y'[x]-Exp[2*x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{2x}}{2} + c_1$$

## 4.8 problem 5.1 (h)

Internal problem ID [13034]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.1 (h).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y' - \sin(x)y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(diff(y(x),x)=y(x)*sin(x),y(x), singsol=all)
```

$$y = c_1 e^{-\cos(x)}$$

### ✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 19

```
DSolve[y'[x]==y[x]*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-\cos(x)}$$

$$y(x) \rightarrow 0$$

## 4.9 problem 5.1 (i)

Internal problem ID [13035]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.1 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' + 4y - y^3 = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 33

```
dsolve(diff(y(x),x)+4*y(x)=y(x)^3,y(x), singsol=all)
```

$$y = -\frac{2}{\sqrt{4e^{8x}c_1 + 1}}$$

$$y = \frac{2}{\sqrt{4e^{8x}c_1 + 1}}$$

✓ Solution by Mathematica

Time used: 0.728 (sec). Leaf size: 56

```
DSolve[y'[x]+4*y[x]==y[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2}{\sqrt{1 + e^{8(x+c_1)}}}$$

$$y(x) \rightarrow \frac{2}{\sqrt{1 + e^{8(x+c_1)}}}$$

$$y(x) \rightarrow -2$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow 2$$

## 4.10 problem 5.1 (j)

Internal problem ID [13036]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.1 (j).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y'x - 827y = -\cos(x^2)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(x*diff(y(x),x)+cos(x^2)=827*y(x),y(x), singsol=all)
```

$$y = \left( \int -\frac{\cos(x^2)}{x^{828}} dx + c_1 \right) x^{827}$$

### ✓ Solution by Mathematica

Time used: 6.501 (sec). Leaf size: 2119

```
DSolve[x*y'[x]+Cos[x^2]==827*y[x],y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

## 4.11 problem 5.2 (a)

Internal problem ID [13037]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.2 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' + 2y = 6$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)+2*y(x)=6,y(x), singsol=all)
```

$$y = 3 + c_1 e^{-2x}$$

### ✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 20

```
DSolve[y'[x]+2*y[x]==6,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3 + c_1 e^{-2x}$$

$$y(x) \rightarrow 3$$

## 4.12 problem 5.2 (b)

Internal problem ID [13038]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.2 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 2y = 20e^{3x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)+2*y(x)=20*exp(3*x),y(x), singsol=all)
```

$$y = (4e^{5x} + c_1)e^{-2x}$$

### ✓ Solution by Mathematica

Time used: 0.051 (sec). Leaf size: 21

```
DSolve[y'[x]+2*y[x]==20*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(4e^{5x} + c_1)$$

### 4.13 problem 5.2 (c)

Internal problem ID [13039]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.2 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' - 4y = 16x$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)=4*y(x)+16*x,y(x), singsol=all)
```

$$y = -4x - 1 + c_1 e^{4x}$$

#### ✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 18

```
DSolve[y'[x]==4*y[x]+16*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -4x + c_1 e^{4x} - 1$$



## 4.14 problem 5.2 (d)

Internal problem ID [13040]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.2 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y' - 2xy = x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)-2*x*y(x)=x,y(x), singsol=all)
```

$$y = -\frac{1}{2} + c_1 e^{x^2}$$

### ✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 24

```
DSolve[y'[x]-2*x*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2} + c_1 e^{x^2}$$

$$y(x) \rightarrow -\frac{1}{2}$$

## 4.15 problem 5.2 (e)

Internal problem ID [13041]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.2 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y'x + 3y = 10x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x*diff(y(x),x)+3*y(x)-10*x^2=0,y(x), singsol=all)
```

$$y = \frac{2x^5 + c_1}{x^3}$$

### ✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 17

```
DSolve[x*y'[x]+3*y[x]-10*x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2x^5 + c_1}{x^3}$$

## 4.16 problem 5.2 (f)

Internal problem ID [13042]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.2 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y'x^2 + 2xy = \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(x^2*diff(y(x),x)+2*x*y(x)=sin(x),y(x), singsol=all)
```

$$y = \frac{-\cos(x) + c_1}{x^2}$$

### ✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 16

```
DSolve[x^2*y'[x]+2*x*y[x]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-\cos(x) + c_1}{x^2}$$

## 4.17 problem 5.2 (g)

Internal problem ID [13043]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.2 (g).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y'x - 3y = \sqrt{x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x*diff(y(x),x)=sqrt(x)+3*y(x),y(x), singsol=all)
```

$$y = \left( -\frac{2}{5x^{\frac{5}{2}}} + c_1 \right) x^3$$

### ✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 21

```
DSolve[x*y'[x]==Sqrt[x]+3*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2\sqrt{x}}{5} + c_1x^3$$

## 4.18 problem 5.2 (h)

Internal problem ID [13044]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.2 (h).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$\cos(x) y' + \sin(x) y = \cos(x)^2$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 10

```
dsolve(cos(x)*diff(y(x),x)+sin(x)*y(x)=cos(x)^2,y(x), singsol=all)
```

$$y = (x + c_1) \cos(x)$$

### ✓ Solution by Mathematica

Time used: 0.058 (sec). Leaf size: 12

```
DSolve[Cos[x]*y'[x]+Sin[x]*y[x]==Cos[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x + c_1) \cos(x)$$

## 4.19 problem 5.2 (i)

Internal problem ID [13045]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.2 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y'x + (5x + 2)y = \frac{20}{x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(x*diff(y(x),x)+(5*x+2)*y(x)=20/x,y(x), singsol=all)
```

$$y = \frac{4}{x^2} + \frac{c_1 e^{-5x}}{x^2}$$

### ✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 12

```
DSolve[Cos[x]*y'[x]+Sin[x]*y[x]==Cos[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x + c_1) \cos(x)$$

## 4.20 problem 5.2 (j)

Internal problem ID [13046]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.2 (j).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$2\sqrt{x}y' + y = 2xe^{-\sqrt{x}}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(2*sqrt(x)*diff(y(x),x)+y(x)=2*x*exp(-sqrt(x)),y(x), singsol=all)
```

$$y = \left( \frac{2x^{\frac{3}{2}}}{3} + c_1 \right) e^{-\sqrt{x}}$$

### ✓ Solution by Mathematica

Time used: 0.074 (sec). Leaf size: 30

```
DSolve[2*Sqrt[x]*y'[x]+y[x]==2*x*Exp[-Sqrt[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}e^{-\sqrt{x}}(2x^{3/2} + 3c_1)$$

## 4.21 problem 5.3 (a)

Internal problem ID [13047]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.3 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_quadrature`]

$$y' - 3y = 6$$

With initial conditions

$$[y(0) = 5]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve([diff(y(x),x)-3*y(x)=6,y(0) = 5],y(x), singsol=all)
```

$$y = 7e^{3x} - 2$$

### ✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 14

```
DSolve[{y'[x]-3*y[x]==6,{y[0]==5}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 7e^{3x} - 2$$



## 4.22 problem 5.3 (b)

Internal problem ID [13048]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.3 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_quadrature`]

$$y' - 3y = 6$$

With initial conditions

$$[y(0) = -2]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve([diff(y(x),x)-3*y(x)=6,y(0) = -2],y(x), singsol=all)
```

$$y = -2$$

### ✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 6

```
DSolve[{y'[x]-3*y[x]==6,{y[0]==-2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2$$

## 4.23 problem 5.3 (c)

Internal problem ID [13049]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.3 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 5y = e^{-3x}$$

With initial conditions

$$[y(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve([diff(y(x),x)+5*y(x)=exp(-3*x),y(0) = 0],y(x), singsol=all)
```

$$y = \frac{(e^{2x} - 1)e^{-5x}}{2}$$

### ✓ Solution by Mathematica

Time used: 0.051 (sec). Leaf size: 21

```
DSolve[{y'[x]+5*y[x]==Exp[-3*x],{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}e^{-5x}(e^{2x} - 1)$$

## 4.24 problem 5.3 (d)

Internal problem ID [13050]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.3 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y'x + 3y = 20x^2$$

With initial conditions

$$[y(1) = 10]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([x*diff(y(x),x)+3*y(x)=20*x^2,y(1) = 10],y(x), singsol=all)
```

$$y = \frac{4x^5 + 6}{x^3}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 16

```
DSolve[{x*y'[x]+3*y[x]==20*x^2,{y[1]==10}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{4x^5 + 6}{x^3}$$

## 4.25 problem 5.3 (e)

Internal problem ID [13051]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.3 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_linear]`

$$y'x - y = x^2 \cos(x)$$

With initial conditions

$$\left[ y\left(\frac{\pi}{2}\right) = 0 \right]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve([x*diff(y(x),x)=y(x)+x^2*cos(x),y(1/2*Pi) = 0],y(x), singsol=all)
```

$$y = x(\sin(x) - 1)$$

✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 11

```
DSolve[{x*y'[x]==y[x]+x^2*Cos[x],{y[Pi/2]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(\sin(x) - 1)$$

## 4.26 problem 5.3 (f)

Internal problem ID [13052]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.3 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$(x^2 + 1)y' - x(3 + 3x^2 - y) = 0$$

With initial conditions

$$[y(2) = 8]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve([(1+x^2)*diff(y(x),x)=x*(3+3*x^2-y(x)),y(2) = 8],y(x), singsol=all)
```

$$y = x^2 + 1 + \frac{3\sqrt{5}}{\sqrt{x^2 + 1}}$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 26

```
DSolve[{(1+x^2)*y'[x]==x*(3+3*x^2-y[x]),{y[2]==8}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2 + \frac{3\sqrt{5}}{\sqrt{x^2 + 1}} + 1$$

## 4.27 problem 5.4 (a)

Internal problem ID [13053]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.4 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y' + 6xy = \sin(x)$$

With initial conditions

$$[y(0) = 4]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 77

```
dsolve([diff(y(x),x)+6*x*y(x)=sin(x),y(0) = 4],y(x), singsol=all)
```

$$y = 4e^{-3x^2} - \frac{\sqrt{3}\sqrt{\pi}e^{\frac{1}{12}-3x^2}\operatorname{erf}\left(\frac{\sqrt{3}}{6}\right)}{6} - \frac{\sqrt{3}\sqrt{\pi}e^{\frac{1}{12}-3x^2}\operatorname{erf}\left(\frac{(6ix-1)\sqrt{3}}{6}\right)}{12} + \frac{\sqrt{3}\sqrt{\pi}e^{\frac{1}{12}-3x^2}\operatorname{erf}\left(\frac{\sqrt{3}(6ix+1)}{6}\right)}{12}$$

✓ Solution by Mathematica

Time used: 0.072 (sec). Leaf size: 105

```
DSolve[{y'[x]+6*x*y[x]==Sin[x],{y[0]==4}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{12}e^{-3x^2} \left( \sqrt[12]{e}\sqrt{3\pi}\operatorname{erf}\left(\frac{1+6ix}{2\sqrt{3}}\right) - 2\sqrt[12]{e}\sqrt{3\pi}\operatorname{erf}\left(\frac{1}{2\sqrt{3}}\right) - i\sqrt[12]{e}\sqrt{3\pi}\operatorname{erfi}\left(\frac{6x+i}{2\sqrt{3}}\right) + 48 \right)$$

## 4.28 problem 5.4 (b)

Internal problem ID [13054]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.4 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y'x^2 + xy = \sqrt{x} \sin(x)$$

With initial conditions

$$[y(2) = 5]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 42

```
dsolve([x^2*diff(y(x),x)+x*y(x)=sqrt(x)*sin(x),y(2) = 5],y(x), singsol=all)
```

$$y = \frac{\sqrt{\pi} \sqrt{2} \operatorname{FresnelS}\left(\frac{\sqrt{2}\sqrt{x}}{\sqrt{\pi}}\right) - \sqrt{\pi} \sqrt{2} \operatorname{FresnelS}\left(\frac{2}{\sqrt{\pi}}\right) + 10}{x}$$

✓ Solution by Mathematica

Time used: 0.093 (sec). Leaf size: 185

```
DSolve[{x^2*y'[x]+x*y[x]==Sqrt[x]*Sin[x],{y[2]==5}},y[x],x,IncludeSingularSolutions -> True]
```

$y(x)$

$$\rightarrow \frac{2i\sqrt{\pi}x\operatorname{erf}(\sqrt{ix}) - (1+i)\sqrt{2\pi}\operatorname{erf}(1+i)\sqrt{ix}\sqrt{x} - 2i\sqrt{\pi}x\operatorname{erfi}(\sqrt{ix}) + (1+i)\sqrt{2\pi}\operatorname{erfi}(1+i)\sqrt{ix}\sqrt{x} - 2}{4\sqrt{ix}x^{3/2}}$$

## 4.29 problem 5.4 (c)

Internal problem ID [13055]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 5. LINEAR FIRST ORDER EQUATIONS. Additional exercises. page 103

**Problem number:** 5.4 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y'x - y = x^2 e^{-x^2}$$

With initial conditions

$$[y(3) = 8]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve([x*diff(y(x),x)-y(x)=x^2*exp(-x^2),y(3) = 8],y(x), singsol=all)
```

$$y = -\frac{\left(-\frac{16}{3} + (\operatorname{erf}(3) - \operatorname{erf}(x))\sqrt{\pi}\right)x}{2}$$

### ✓ Solution by Mathematica

Time used: 0.079 (sec). Leaf size: 30

```
DSolve[{x*y'[x]-y[x]==x^2*Exp[-x^2],{y[3]==8}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{6}x(3\sqrt{\pi}\operatorname{erf}(x) - 3\sqrt{\pi}\operatorname{erf}(3) + 16)$$



## 5 Chapter 6. Simplifying through simplifiction.

### Additional exercises. page 114

5.1	problem 6.1 (a)	152
5.2	problem 6.1 (b)	153
5.3	problem 6.1 (c)	154
5.4	problem 6.2	156
5.5	problem 6.3 (a)	157
5.6	problem 6.3 (b)	158
5.7	problem 6.3 (c)	159
5.8	problem 6.4	161
5.9	problem 6.5 (a)	162
5.10	problem 6.5 (b)	164
5.11	problem 6.5 (c)	165
5.12	problem 6.6	166
5.13	problem 6.7 (a)	167
5.14	problem 6.7 (b)	168
5.15	problem 6.7 (c)	169
5.16	problem 6.7 (d)	170
5.17	problem 6.7 (e)	171
5.18	problem 6.7 (f)	172
5.19	problem 6.7 (g)	173
5.20	problem 6.7 (h)	174
5.21	problem 6.7 (i)	175
5.22	problem 6.7 (j)	176
5.23	problem 6.7 (k)	177
5.24	problem 6.7 (L)	178
5.25	problem 6.7 (m)	180
5.26	problem 6.7 (n)	181
5.27	problem 6.7 (o)	182
5.28	problem 6.7 (p)	183

## 5.1 problem 6.1 (a)

Internal problem ID [13056]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.1 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$y' - \frac{1}{(3x + 3y + 2)^2} = 0$$

### ✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 24

```
dsolve(diff(y(x),x)=1/(3*x+3*y(x)+2)^2,y(x), singsol=all)
```

$$y = -c_1 + \frac{\text{RootOf}(-\_Z + 3c_1 - 3x - 2 + \tan(\_Z))}{3}$$

### ✓ Solution by Mathematica

Time used: 0.162 (sec). Leaf size: 23

```
DSolve[y'[x]==1/(3*x+3*y[x]+2)^2,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[y(x) - \frac{1}{3} \arctan(3y(x) + 3x + 2) = c_1, y(x)\right]$$

## 5.2 problem 6.1 (b)

Internal problem ID [13057]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.1 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{(3x - 2y)^2 + 1}{3x - 2y} = \frac{3}{2}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 39

```
dsolve(diff(y(x),x)=( (3*x-2*y(x))^2+1 )/(3*x-2*y(x))+3/2,y(x), singsol=all)
```

$$y = \frac{3x}{2} - \frac{\sqrt{e^{-4x}c_1 - 1}}{2}$$

$$y = \frac{3x}{2} + \frac{\sqrt{e^{-4x}c_1 - 1}}{2}$$

### ✓ Solution by Mathematica

Time used: 11.283 (sec). Leaf size: 78

```
DSolve[y'[x]==( (3*x-2*y[x])^2+1 )/(3*x-2*y[x])+3/2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left( 3x - \frac{\sqrt{e^{4x} - 4c_1}}{\sqrt{-e^{4x}}} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left( 3x + \frac{\sqrt{e^{4x} - 4c_1}}{\sqrt{-e^{4x}}} \right)$$

### 5.3 problem 6.1 (c)

Internal problem ID [13058]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.1 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _exact, _dAlembert]`

$$\cos(-4y + 8x - 3)y' - 2\cos(-4y + 8x - 3) = 2$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve(cos(4*y(x)-8*x+3)*diff(y(x),x)=2+2*cos(4*y(x)-8*x+3),y(x), singsol=all)
```

$$y = 2x - \frac{3}{4} - \frac{\arcsin(-8x + 8c_1)}{4}$$

✓ Solution by Mathematica

Time used: 64.647 (sec). Leaf size: 1165

`DSolve[Cos[4*y[x]-8*x+3]*y'[x]==2+2*Cos[4*y[x]-8*x+3],y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow -\arccos\left(-\frac{1}{2}\sqrt{2-\sqrt{-2\sqrt{-((64x^2+64c_1x-1+16c_1^2)\cos^2(3-8x))+8(2x+c_1)\sin(3-8x)+2}}}\right)$$

$$y(x) \rightarrow \arccos\left(-\frac{1}{2}\sqrt{2-\sqrt{-2\sqrt{-((64x^2+64c_1x-1+16c_1^2)\cos^2(3-8x))+8(2x+c_1)\sin(3-8x)+2}}}\right)$$

$$y(x) \rightarrow -\arccos\left(\frac{1}{2}\sqrt{2-\sqrt{-2\sqrt{-((64x^2+64c_1x-1+16c_1^2)\cos^2(3-8x))+8(2x+c_1)\sin(3-8x)+2}}}\right)$$

$$y(x) \rightarrow \arccos\left(\frac{1}{2}\sqrt{2-\sqrt{-2\sqrt{-((64x^2+64c_1x-1+16c_1^2)\cos^2(3-8x))+8(2x+c_1)\sin(3-8x)+2}}}\right)$$

$$y(x) \rightarrow -\arccos\left(-\frac{1}{2}\sqrt{2+\sqrt{-2\sqrt{-((64x^2+64c_1x-1+16c_1^2)\cos^2(3-8x))+8(2x+c_1)\sin(3-8x)+2}}}\right)$$

$$y(x) \rightarrow \arccos\left(-\frac{1}{2}\sqrt{2+\sqrt{-2\sqrt{-((64x^2+64c_1x-1+16c_1^2)\cos^2(3-8x))+8(2x+c_1)\sin(3-8x)+2}}}\right)$$

$$y(x) \rightarrow -\arccos\left(\frac{1}{2}\sqrt{2+\sqrt{-2\sqrt{-((64x^2+64c_1x-1+16c_1^2)\cos^2(3-8x))+8(2x+c_1)\sin(3-8x)+2}}}\right)$$

$$y(x) \rightarrow \arccos\left(\frac{1}{2}\sqrt{2+\sqrt{-2\sqrt{-((64x^2+64c_1x-1+16c_1^2)\cos^2(3-8x))+8(2x+c_1)\sin(3-8x)+2}}}\right)$$

$$y(x) \rightarrow -\arccos\left(-\frac{1}{2}\sqrt{2-\sqrt{2}\sqrt{\sqrt{-((64x^2+64c_1x-1+16c_1^2)\cos^2(3-8x))+4(2x+c_1)\sin(3-8x)+1}}}\right)$$

$$y(x) \rightarrow \arccos\left(-\frac{1}{2}\sqrt{2-\sqrt{2}\sqrt{\sqrt{-((64x^2+64c_1x-1+16c_1^2)\cos^2(3-8x))+4(2x+c_1)\sin(3-8x)+1}}}\right)$$

## 5.4 problem 6.2

Internal problem ID [13059]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.2.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _Riccati]`

$$y' - (y - x)^2 = 1$$

With initial conditions

$$\left[ y(0) = \frac{1}{4} \right]$$

### ✓ Solution by Maple

Time used: 0.14 (sec). Leaf size: 18

```
dsolve([diff(y(x),x)=1+(y(x)-x)^2,y(0) = 1/4],y(x), singsol=all)
```

$$y = \frac{x^2 - 4x - 1}{x - 4}$$

### ✓ Solution by Mathematica

Time used: 0.134 (sec). Leaf size: 19

```
DSolve[{y'[x]==1+(y[x]-x)^2,{y[0]==1/4}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2 - 4x - 1}{x - 4}$$

## 5.5 problem 6.3 (a)

Internal problem ID [13060]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.3 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$y'x^2 - xy - y^2 = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x)-x*y(x)=y(x)^2,y(x), singsol=all)
```

$$y = -\frac{x}{\ln(x) - c_1}$$

### ✓ Solution by Mathematica

Time used: 0.125 (sec). Leaf size: 21

```
DSolve[x^2*y'[x]-x*y[x]==y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x}{-\log(x) + c_1}$$

$$y(x) \rightarrow 0$$

## 5.6 problem 6.3 (b)

Internal problem ID [13061]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.3 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$y' - \frac{y}{x} - \frac{x}{y} = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(diff(y(x),x)=y(x)/x+x/y(x),y(x), singsol=all)
```

$$y = \sqrt{2 \ln(x) + c_1} x$$

$$y = -\sqrt{2 \ln(x) + c_1} x$$

### ✓ Solution by Mathematica

Time used: 0.16 (sec). Leaf size: 36

```
DSolve[y'[x]==y[x]/x+x/y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x\sqrt{2\log(x) + c_1}$$

$$y(x) \rightarrow x\sqrt{2\log(x) + c_1}$$



## 5.7 problem 6.3 (c)

Internal problem ID [13062]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.3 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _dAlembert]`

$$\cos\left(\frac{y}{x}\right)\left(y' - \frac{y}{x}\right) - \sin\left(\frac{y}{x}\right) = 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(cos(y(x)/x)*(diff(y(x),x)-y(x)/x)=1+sin(y(x)/x),y(x), singsol=all)
```

$$y = \arcsin(c_1 x - 1) x$$

✓ Solution by Mathematica

Time used: 60.351 (sec). Leaf size: 185

```
DSolve[Cos[y[x]/x]*(y'[x]-y[x]/x)==1+Sin[y[x]/x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\pi x}{2}$$

$$y(x) \rightarrow \frac{3\pi x}{2}$$

$$y(x) \rightarrow -2x \arccos\left(\frac{1}{2}\left(e^{\frac{c_1}{2}}\sqrt{x} - \sqrt{2 - e^{c_1}x}\right)\right)$$

$$y(x) \rightarrow 2x \arccos\left(\frac{1}{2}\left(e^{\frac{c_1}{2}}\sqrt{x} - \sqrt{2 - e^{c_1}x}\right)\right)$$

$$y(x) \rightarrow -2x \arccos\left(\frac{1}{2}\left(e^{\frac{c_1}{2}}\sqrt{x} + \sqrt{2 - e^{c_1}x}\right)\right)$$

$$y(x) \rightarrow 2x \arccos\left(\frac{1}{2}\left(e^{\frac{c_1}{2}}\sqrt{x} + \sqrt{2 - e^{c_1}x}\right)\right)$$

## 5.8 problem 6.4

Internal problem ID [13063]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.4.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{x-y}{x+y} = 0$$

With initial conditions

$$[y(0) = 3]$$

✓ Solution by Maple

Time used: 0.547 (sec). Leaf size: 19

```
dsolve([diff(y(x),x)=(x-y(x))/(x+y(x)),y(0) = 3],y(x), singsol=all)
```

$$y = -x + \sqrt{2x^2 + 9}$$

✓ Solution by Mathematica

Time used: 0.434 (sec). Leaf size: 20

```
DSolve[{y'[x]==(x-y[x])/(x+y[x]),{y[0]==3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt{2x^2 + 9} - x$$

## 5.9 problem 6.5 (a)

Internal problem ID [13064]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.5 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' + 3y - 3y^3 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x)+3*y(x)=3*y(x)^3,y(x), singsol=all)
```

$$y = \frac{1}{\sqrt{e^{6x}c_1 + 1}}$$
$$y = -\frac{1}{\sqrt{e^{6x}c_1 + 1}}$$

✓ Solution by Mathematica

Time used: 0.675 (sec). Leaf size: 58

```
DSolve[y'[x]+3*y[x]==3*y[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\sqrt{1 + e^{6x+2c_1}}}$$

$$y(x) \rightarrow \frac{1}{\sqrt{1 + e^{6x+2c_1}}}$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow 1$$

## 5.10 problem 6.5 (b)

Internal problem ID [13065]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.5 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$y' - \frac{3y}{x} - \frac{y^2}{x^2} = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 20

```
dsolve(diff(y(x),x)-3*y(x)/x=(y(x)/x)^2,y(x), singsol=all)
```

$$y = \frac{2x^3}{-x^2 + 2c_1}$$

### ✓ Solution by Mathematica

Time used: 0.141 (sec). Leaf size: 25

```
DSolve[y'[x]-3*y[x]/x==(y[x]/x)^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2x^3}{x^2 - 2c_1}$$

$$y(x) \rightarrow 0$$

## 5.11 problem 6.5 (c)

Internal problem ID [13066]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.5 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_Bernoulli]

$$y' + 3 \cot(x) y - 6 \cos(x) y^{\frac{2}{3}} = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)+3*cot(x)*y(x)=6*cos(x)*y(x)^(2/3),y(x), singsol=all)
```

$$-\sin(x) - \frac{c_1}{\sin(x)} + y^{\frac{1}{3}} = 0$$

### ✓ Solution by Mathematica

Time used: 0.305 (sec). Leaf size: 24

```
DSolve[y'[x]+3*Cot[x]*y[x]==6*Cos[x]*y[x]^(2/3),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{8} \csc^3(x) (\cos(2x) - 2c_1)^3$$

## 5.12 problem 6.6

Internal problem ID [13067]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.6.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class D', _rational, _Bernoulli]`

$$y' - \frac{y}{x} - \frac{1}{y} = 0$$

With initial conditions

$$[y(1) = 3]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 13

```
dsolve([diff(y(x),x)-1/x*y(x)=1/y(x),y(1) = 3],y(x), singsol=all)
```

$$y = \sqrt{x(11x - 2)}$$

✓ Solution by Mathematica

Time used: 0.251 (sec). Leaf size: 20

```
DSolve[{y'[x]-1/x*y[x]==1/y[x],{y[1]==3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt{x}\sqrt{11x - 2}$$



## 5.13 problem 6.7 (a)

Internal problem ID [13068]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$y' - \frac{y}{x} - \frac{x^2}{y^2} = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 74

```
dsolve(diff(y(x),x)=y(x)/x+(x/y(x))^2,y(x), singsol=all)
```

$$y = (3 \ln(x) + c_1)^{\frac{1}{3}} x$$
$$y = \left( -\frac{(3 \ln(x) + c_1)^{\frac{1}{3}}}{2} - \frac{i\sqrt{3}(3 \ln(x) + c_1)^{\frac{1}{3}}}{2} \right) x$$
$$y = \left( -\frac{(3 \ln(x) + c_1)^{\frac{1}{3}}}{2} + \frac{i\sqrt{3}(3 \ln(x) + c_1)^{\frac{1}{3}}}{2} \right) x$$

### ✓ Solution by Mathematica

Time used: 0.174 (sec). Leaf size: 63

```
DSolve[y'[x]==y[x]/x+(x/y[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \sqrt[3]{3 \log(x) + c_1}$$
$$y(x) \rightarrow -\sqrt[3]{-1} x \sqrt[3]{3 \log(x) + c_1}$$
$$y(x) \rightarrow (-1)^{2/3} x \sqrt[3]{3 \log(x) + c_1}$$

## 5.14 problem 6.7 (b)

Internal problem ID [13069]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$3y' - \sqrt{2x + 3y + 4} = -2$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(3*diff(y(x),x)=-2+sqrt(2*x+3*y(x)+4),y(x), singsol=all)
```

$$x - 2\sqrt{2x + 3y + 4} - c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 0.329 (sec). Leaf size: 51

```
DSolve[3*y'[x]==-2+Sqrt[2*x+3*y[x]+4],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{48}(4(x^2 - 6x - 15) - 4e^{c_1}(x + 1) + e^{2c_1})$$

$$y(x) \rightarrow \frac{1}{12}(x^2 - 6x - 15)$$

## 5.15 problem 6.7 (c)

Internal problem ID [13070]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, _Bernoulli]`

$$3y' + \frac{2y}{x} - 4\sqrt{y} = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(3*diff(y(x),x)+2/x*y(x)=4*sqrt(y(x)),y(x), singsol=all)
```

$$\sqrt{y} - \frac{x}{2} - \frac{c_1}{x^{\frac{1}{3}}} = 0$$

### ✓ Solution by Mathematica

Time used: 0.158 (sec). Leaf size: 26

```
DSolve[3*y'[x]+2/x*y[x]==4*Sqrt[y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{(x^{4/3} + 2c_1)^2}{4x^{2/3}}$$

## 5.16 problem 6.7 (d)

Internal problem ID [13071]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$y' - \frac{1}{\sin(4x - y)} = 4$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)=4+1/sin(4*x-y(x)),y(x), singsol=all)
```

$$y = 4x - \pi + \arccos(-x + c_1)$$

### ✓ Solution by Mathematica

Time used: 0.624 (sec). Leaf size: 33

```
DSolve[y'[x]==4+1/Sin[4*x-y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 4x - \arccos(x - c_1)$$

$$y(x) \rightarrow 4x + \arccos(x - c_1)$$

## 5.17 problem 6.7 (e)

Internal problem ID [13072]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class C'], [_Abel, '2nd type', 'class C'], _d`

$$(y - x)y' = 1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve((y(x)-x)*diff(y(x),x)=1,y(x), singsol=all)
```

$$y = \text{LambertW}(-c_1 e^{-x-1}) + x + 1$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 20

```
DSolve[(y[x]-x)*y'[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow W(c_1(-e^{-x-1})) + x + 1$$

## 5.18 problem 6.7 (f)

Internal problem ID [13073]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$(x + y)y' - y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve((y(x)+x)*diff(y(x),x)=y(x),y(x), singsol=all)
```

$$y = e^{\text{LambertW}(e^{c_1 x}) - c_1}$$

### ✓ Solution by Mathematica

Time used: 3.407 (sec). Leaf size: 23

```
DSolve[(y[x]+x)*y'[x]==y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x}{W(e^{-c_1 x})}$$

$$y(x) \rightarrow 0$$

## 5.19 problem 6.7 (g)

Internal problem ID [13074]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (g).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$(2xy + 2x^2) y' - 2xy - 2y^2 = x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve((2*x*y(x)+2*x^2)*diff(y(x),x)=x^2+2*x*y(x)+2*y(x)^2,y(x), singsol=all)
```

$$y = \left(-1 - \sqrt{1 + \ln(x) + c_1}\right) x$$

$$y = \left(-1 + \sqrt{1 + \ln(x) + c_1}\right) x$$

### ✓ Solution by Mathematica

Time used: 0.243 (sec). Leaf size: 42

```
DSolve[(2*x*y[x]+2*x^2)*y'[x]==x^2+2*x*y[x]+2*y[x]^2,y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow -x \left(1 + \sqrt{\log(x) + 1 + 2c_1}\right)$$

$$y(x) \rightarrow x \left(-1 + \sqrt{\log(x) + 1 + 2c_1}\right)$$

## 5.20 problem 6.7 (h)

Internal problem ID [13075]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (h).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational, _Bernoulli]`

$$y' + \frac{y}{x} - y^3 x^2 = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(diff(y(x),x)+1/x*y(x)=x^2*y(x)^3,y(x), singsol=all)
```

$$y = \frac{1}{\sqrt{c_1 - 2x x}}$$
$$y = -\frac{1}{\sqrt{c_1 - 2x x}}$$

### ✓ Solution by Mathematica

Time used: 0.336 (sec). Leaf size: 44

```
DSolve[y'[x]+1/x*y[x]==x^2*y[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\sqrt{x^2(-2x + c_1)}}$$
$$y(x) \rightarrow \frac{1}{\sqrt{x^2(-2x + c_1)}}$$
$$y(x) \rightarrow 0$$



## 5.21 problem 6.7 (i)

Internal problem ID [13076]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$y' - 2\sqrt{2x + y - 3} = -2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)=2*sqrt(2*x+y(x)-3)-2,y(x), singsol=all)
```

$$x - \sqrt{2x + y - 3} - c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 0.256 (sec). Leaf size: 38

```
DSolve[y'[x]==2*Sqrt[2*x+y[x]-3]-2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2 - e^{c_1}(x + 1) + 4 + \frac{e^{2c_1}}{4}$$

$$y(x) \rightarrow x^2 + 4$$

## 5.22 problem 6.7 (j)

Internal problem ID [13077]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (j).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$y' - 2\sqrt{2x + y - 3} = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 57

```
dsolve(diff(y(x),x)=2*sqrt(2*x+y(x)-3),y(x), singsol=all)
```

$$x - \sqrt{2x + y - 3} - \frac{\ln(\sqrt{2x + y - 3} - 1)}{2} + \frac{\ln(\sqrt{2x + y - 3} + 1)}{2} + \frac{\ln(-4 + 2x + y)}{2} - c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 8.176 (sec). Leaf size: 51

```
DSolve[y'[x]==2*Sqrt[2*x+y[x]-3],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow W\left(-e^{-x+\frac{1}{2}+c_1}\right)^2 + 2W\left(-e^{-x+\frac{1}{2}+c_1}\right) - 2x + 4$$
$$y(x) \rightarrow 4 - 2x$$

## 5.23 problem 6.7 (k)

Internal problem ID [13078]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (k).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _dAlembert]`

$$y'x - y - \sqrt{xy + x^2} = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve(x*diff(y(x),x)-y(x)=sqrt(x*y(x)+x^2),y(x), singsol=all)
```

$$-\frac{x+y}{\sqrt{x(x+y)}} + \frac{\ln(x)}{2} - c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 0.194 (sec). Leaf size: 26

```
DSolve[x*y'[x]-y[x]==Sqrt[x*y[x]+x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}x(\log^2(x) + 2c_1 \log(x) - 4 + c_1^2)$$

## 5.24 problem 6.7 (L)

Internal problem ID [13079]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (L).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _Bernoulli]`

$$y' + 3y - \frac{28e^{2x}}{y^3} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 78

```
dsolve(diff(y(x),x)+3*y(x)=28*exp(2*x)*1/(y(x)^3),y(x), singsol=all)
```

$$y = (8e^{14x} + c_1)^{\frac{1}{4}} e^{-3x}$$

$$y = -(8e^{14x} + c_1)^{\frac{1}{4}} e^{-3x}$$

$$y = -i(8e^{14x} + c_1)^{\frac{1}{4}} e^{-3x}$$

$$y = i(8e^{14x} + c_1)^{\frac{1}{4}} e^{-3x}$$

✓ Solution by Mathematica

Time used: 0.741 (sec). Leaf size: 104

```
DSolve[y'[x]+3*y[x]==28*Exp[2*x]*1/(y[x]^3),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -e^{-3x} \sqrt[4]{8e^{14x} + c_1}$$

$$y(x) \rightarrow -ie^{-3x} \sqrt[4]{8e^{14x} + c_1}$$

$$y(x) \rightarrow ie^{-3x} \sqrt[4]{8e^{14x} + c_1}$$

$$y(x) \rightarrow e^{-3x} \sqrt[4]{8e^{14x} + c_1}$$

## 5.25 problem 6.7 (m)

Internal problem ID [13080]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (m).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _Riccati]`

$$y' - (x - y + 3)^2 = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve(diff(y(x),x)=(x-y(x)+3)^2,y(x), singsol=all)
```

$$y = \frac{x e^{2x} c_1 + 2 e^{2x} c_1 - x - 4}{e^{2x} c_1 - 1}$$

### ✓ Solution by Mathematica

Time used: 0.163 (sec). Leaf size: 29

```
DSolve[y'[x]==(x-y[x]+3)^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x + \frac{1}{\frac{1}{2} + c_1 e^{2x}} + 2$$
$$y(x) \rightarrow x + 2$$

## 5.26 problem 6.7 (n)

Internal problem ID [13081]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (n).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _Clairaut]`

$$y' - 2\sqrt{y + x^2} = -2x$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve(diff(y(x),x)+2*x=2*sqrt(y(x)+x^2),y(x), singsol=all)
```

$$x - \sqrt{y + x^2} - c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 0.736 (sec). Leaf size: 35

```
DSolve[y'[x]+2*x==2*Sqrt[y[x]+x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (1 + e^{c_1})(2x + 1 + e^{c_1})$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow 2x + 1$$

## 5.27 problem 6.7 (o)

Internal problem ID [13082]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (o).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [ $y = G(x, y')$ ]

$$\cos(y) y' + \sin(y) = e^{-x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(cos(y(x))*diff(y(x),x)=exp(-x)-sin(y(x)),y(x), singsol=all)
```

$$y = -\arcsin((-x + c_1)e^{-x})$$

### ✓ Solution by Mathematica

Time used: 11.73 (sec). Leaf size: 16

```
DSolve[Cos[y[x]]*y'[x]==Exp[-x]-Sin[y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arcsin(e^{-x}(x + c_1))$$



## 5.28 problem 6.7 (p)

Internal problem ID [13083]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 6. Simplifying through simplification. Additional exercises. page 114

**Problem number:** 6.7 (p).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, _Riccati]`

$$y' - x \left( 1 + \frac{2y}{x^2} + \frac{y^2}{x^4} \right) = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)=x*(1+2*y(x)/x^2+y(x)^2/x^4),y(x), singsol=all)
```

$$y = -\tan(-\ln(x) + c_1) x^2$$

### ✓ Solution by Mathematica

Time used: 0.187 (sec). Leaf size: 15

```
DSolve[y'[x]==x*(1+2*y[x]/x^2+y[x]^2/x^4),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2 \tan(\log(x) + c_1)$$

## 6 Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

6.1	problem 7.2 . . . . .	185
6.2	problem 7.2 (c) . . . . .	186
6.3	problem 7.4 (a) . . . . .	187
6.4	problem 7.4 (b) . . . . .	189
6.5	problem 7.4 (c) . . . . .	191
6.6	problem 7.4 (d) . . . . .	192
6.7	problem 7.4 (e) . . . . .	193
6.8	problem 7.4 (f) . . . . .	194
6.9	problem 7.4 (g) . . . . .	195
6.10	problem 7.4 (h) . . . . .	196
6.11	problem 7.5 (a) . . . . .	197
6.12	problem 7.5 (b) . . . . .	199
6.13	problem 7.5 (c) . . . . .	201
6.14	problem 7.5 (d) . . . . .	203
6.15	problem 7.5 (e) . . . . .	205
6.16	problem 7.5 (f) . . . . .	207
6.17	problem 7.5 (g) . . . . .	208
6.18	problem 7.5 (h) . . . . .	210
6.19	problem 7.5 (i) . . . . .	212

## 6.1 problem 7.2

Internal problem ID [13084]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.2.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, _Bernoulli]`

$$y' - \frac{1}{y} + \frac{y}{2x} = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve(diff(y(x),x)=1/y(x)-y(x)/(2*x),y(x), singsol=all)
```

$$y = \frac{\sqrt{x(x^2 + c_1)}}{x}$$
$$y = -\frac{\sqrt{x(x^2 + c_1)}}{x}$$

### ✓ Solution by Mathematica

Time used: 0.185 (sec). Leaf size: 42

```
DSolve[y'[x]==1/y[x]-y[x]/(2*x),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{x^2 + c_1}}{\sqrt{x}}$$
$$y(x) \rightarrow \frac{\sqrt{x^2 + c_1}}{\sqrt{x}}$$

## 6.2 problem 7.2 (c)

Internal problem ID [13085]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.2 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _exact, _rational, _Bernoulli]`

$$e^{y^2x-x^2}(y^2 - 2x) + 2e^{y^2x-x^2}xyy' = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 40

```
dsolve(exp(x*y(x)^2-x^2)*(y(x)^2-2*x)+exp(x*y(x)^2-x^2)*2*x*y(x)*diff(y(x),x)=0,y(x), singso
```

$$y = \frac{\sqrt{-x(-x^2 + c_1)}}{x}$$
$$y = -\frac{\sqrt{-x(-x^2 + c_1)}}{x}$$

### ✓ Solution by Mathematica

Time used: 0.12 (sec). Leaf size: 42

```
DSolve[Exp[x*y[x]^2-x^2]*(y[x]^2-2*x)+Exp[x*y[x]^2-x^2]*2*x*y[x]*y'[x]==0,y[x],x,IncludeSing
```

$$y(x) \rightarrow -\frac{\sqrt{x^2 + c_1}}{\sqrt{x}}$$
$$y(x) \rightarrow \frac{\sqrt{x^2 + c_1}}{\sqrt{x}}$$

### 6.3 problem 7.4 (a)

Internal problem ID [13086]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.4 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _exact, _rational, [_Abel, '2nd ty`

$$2xy + y^2 + (2xy + x^2)y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 68

```
dsolve(2*x*y(x)+y(x)^2+(2*x*y(x)+x^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = -\frac{c_1^2 x^2 + \sqrt{c_1^4 x^4 + 4c_1 x}}{2x c_1^2}$$
$$y = \frac{-c_1^2 x^2 + \sqrt{c_1^4 x^4 + 4c_1 x}}{2c_1^2 x}$$

✓ Solution by Mathematica

Time used: 0.579 (sec). Leaf size: 118

```
DSolve[2*x*y[x]+y[x]^2+(2*x*y[x]+x^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left( -x - \frac{\sqrt{x^3 + 4e^{c_1}}}{\sqrt{x}} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left( -x + \frac{\sqrt{x^3 + 4e^{c_1}}}{\sqrt{x}} \right)$$

$$y(x) \rightarrow -\frac{x^{3/2} + \sqrt{x^3}}{2\sqrt{x}}$$

$$y(x) \rightarrow \frac{\sqrt{x^3}}{2\sqrt{x}} - \frac{x}{2}$$

## 6.4 problem 7.4 (b)

Internal problem ID [13087]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.4 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _exact, _rational, _Bernoulli]`

$$2xy^3 + 3y^2y'x^2 = -4x^3$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 99

```
dsolve(2*x*y(x)^3+4*x^3+3*x^2*y(x)^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \frac{((-x^4 + c_1)x)^{\frac{1}{3}}}{x}$$

$$y = -\frac{((-x^4 + c_1)x)^{\frac{1}{3}}}{2x} - \frac{i\sqrt{3}((-x^4 + c_1)x)^{\frac{1}{3}}}{2x}$$

$$y = -\frac{((-x^4 + c_1)x)^{\frac{1}{3}}}{2x} + \frac{i\sqrt{3}((-x^4 + c_1)x)^{\frac{1}{3}}}{2x}$$

✓ Solution by Mathematica

Time used: 0.195 (sec). Leaf size: 78

```
DSolve[2*x*y[x]^3+4*x^3+3*x^2*y[x]^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt[3]{-x^4 + c_1}}{x^{2/3}}$$

$$y(x) \rightarrow -\frac{\sqrt[3]{-1}\sqrt[3]{-x^4 + c_1}}{x^{2/3}}$$

$$y(x) \rightarrow \frac{(-1)^{2/3}\sqrt[3]{-x^4 + c_1}}{x^{2/3}}$$



## 6.5 problem 7.4 (c)

Internal problem ID [13088]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.4 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$3y^2y' = -2 + 2x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 78

```
dsolve(2-2*x+3*y(x)^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = (x^2 + c_1 - 2x)^{\frac{1}{3}}$$
$$y = -\frac{(x^2 + c_1 - 2x)^{\frac{1}{3}}}{2} - \frac{i\sqrt{3}(x^2 + c_1 - 2x)^{\frac{1}{3}}}{2}$$
$$y = -\frac{(x^2 + c_1 - 2x)^{\frac{1}{3}}}{2} + \frac{i\sqrt{3}(x^2 + c_1 - 2x)^{\frac{1}{3}}}{2}$$

### ✓ Solution by Mathematica

Time used: 0.179 (sec). Leaf size: 71

```
DSolve[2-2*x+3*y[x]^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt[3]{x^2 - 2x + 3c_1}$$
$$y(x) \rightarrow -\sqrt[3]{-1} \sqrt[3]{x^2 - 2x + 3c_1}$$
$$y(x) \rightarrow (-1)^{2/3} \sqrt[3]{x^2 - 2x + 3c_1}$$

## 6.6 problem 7.4 (d)

Internal problem ID [13089]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.4 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_exact`, `_rational`, `_Bernoulli`]

$$3y^2x^2 + (2x^3y + 6y)y' = -1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 50

```
dsolve(1+3*x^2*y(x)^2+(2*x^3*y(x)+6*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \frac{\sqrt{(x^3 + 3)(-x + c_1)}}{x^3 + 3}$$
$$y = -\frac{\sqrt{(x^3 + 3)(-x + c_1)}}{x^3 + 3}$$

### ✓ Solution by Mathematica

Time used: 0.212 (sec). Leaf size: 50

```
DSolve[1+3*x^2*y[x]^2+(2*x^3*y[x]+6*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{-x + c_1}}{\sqrt{x^3 + 3}}$$
$$y(x) \rightarrow \frac{\sqrt{-x + c_1}}{\sqrt{x^3 + 3}}$$

## 6.7 problem 7.4 (e)

Internal problem ID [13090]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.4 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _exact, _rational, _dAlembert]`

$$4x^3y + (x^4 - y^4)y' = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(4*x^3*y(x)+(x^4-y(x)^4)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \frac{\text{RootOf}(-5\_Zc_1^4x^4 + \_Z^5 - 1)}{c_1}$$

### ✓ Solution by Mathematica

Time used: 1.472 (sec). Leaf size: 131

```
DSolve[4*x^3*y[x]+(x^4-y[x]^4)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{Root}[\#1^5 - 5\#1x^4 + e^{5c_1}\&, 1]$$

$$y(x) \rightarrow \text{Root}[\#1^5 - 5\#1x^4 + e^{5c_1}\&, 2]$$

$$y(x) \rightarrow \text{Root}[\#1^5 - 5\#1x^4 + e^{5c_1}\&, 3]$$

$$y(x) \rightarrow \text{Root}[\#1^5 - 5\#1x^4 + e^{5c_1}\&, 4]$$

$$y(x) \rightarrow \text{Root}[\#1^5 - 5\#1x^4 + e^{5c_1}\&, 5]$$

$$y(x) \rightarrow 0$$

## 6.8 problem 7.4 (f)

Internal problem ID [13091]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.4 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class G', _exact]`

$$\ln(xy) = -1 - \frac{xy'}{y}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve(1+ln(x*y(x))+x/y(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \frac{e^{\frac{c_1}{x}}}{x}$$

### ✓ Solution by Mathematica

Time used: 0.171 (sec). Leaf size: 17

```
DSolve[1+Log[x*y[x]]+x/y[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{\frac{c_1}{x}}}{x}$$

## 6.9 problem 7.4 (g)

Internal problem ID [13092]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.4 (g).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$e^y + y'e^y x = -1$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 22

```
dsolve(1+exp(y(x))+x*exp(y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = -\ln\left(-\frac{x}{-1 + e^{c_1}x}\right) - c_1$$

### ✓ Solution by Mathematica

Time used: 0.745 (sec). Leaf size: 25

```
DSolve[1+Exp[y[x]]+x*Exp[y[x]]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log\left(-1 + \frac{e^{c_1}}{x}\right)$$

$$y(x) \rightarrow i\pi$$

## 6.10 problem 7.4 (h)

Internal problem ID [13093]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.4 (h).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_1st_order, _with_exponential_symmetries], _exact]`

$$e^y + (e^y x + 1) y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(exp(y(x))+(x*exp(y(x))+1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = -\text{LambertW}(e^{c_1} x) + c_1$$

### ✓ Solution by Mathematica

Time used: 4.529 (sec). Leaf size: 17

```
DSolve[Exp[y[x]]+(x*Exp[y[x]]+1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 - W(e^{c_1} x)$$

## 6.11 problem 7.5 (a)

Internal problem ID [13094]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.5 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y^4 + y^3 y' x = -1$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 70

```
dsolve(1+y(x)^4+x*y(x)^3*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \frac{(-x^4 + c_1)^{\frac{1}{4}}}{x}$$

$$y = -\frac{(-x^4 + c_1)^{\frac{1}{4}}}{x}$$

$$y = -\frac{i(-x^4 + c_1)^{\frac{1}{4}}}{x}$$

$$y = \frac{i(-x^4 + c_1)^{\frac{1}{4}}}{x}$$

✓ Solution by Mathematica

Time used: 0.295 (sec). Leaf size: 218

```
DSolve[1+y[x]^4+x*y[x]^3*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt[4]{-x^4 + e^{4c_1}}}{x}$$

$$y(x) \rightarrow -\frac{i\sqrt[4]{-x^4 + e^{4c_1}}}{x}$$

$$y(x) \rightarrow \frac{i\sqrt[4]{-x^4 + e^{4c_1}}}{x}$$

$$y(x) \rightarrow \frac{\sqrt[4]{-x^4 + e^{4c_1}}}{x}$$

$$y(x) \rightarrow -\sqrt[4]{-1}$$

$$y(x) \rightarrow \sqrt[4]{-1}$$

$$y(x) \rightarrow -(-1)^{3/4}$$

$$y(x) \rightarrow (-1)^{3/4}$$

$$y(x) \rightarrow \frac{ix^3}{(-x^4)^{3/4}}$$

$$y(x) \rightarrow \frac{x^3}{(-x^4)^{3/4}}$$

$$y(x) \rightarrow \frac{i\sqrt[4]{-x^4}}{x}$$

$$y(x) \rightarrow \frac{\sqrt[4]{-x^4}}{x}$$



## 6.12 problem 7.5 (b)

Internal problem ID [13095]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.5 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational]`

$$y + (y^4 - 3x)y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve(y(x)+(y(x)^4-3*x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$x - (-y + c_1)y^3 = 0$$

✓ Solution by Mathematica

Time used: 43.447 (sec). Leaf size: 1270

`DSolve[y[x]+(y[x]^4-3*x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow -\frac{1}{2} \sqrt{\frac{\sqrt[3]{\sqrt{-768x^3 + 81c_1^4x^2 + 9c_1^2x}}}{\sqrt[3]{23^{2/3}}} + \frac{4\sqrt[3]{\frac{2}{3}x}}{\sqrt[3]{\sqrt{-768x^3 + 81c_1^4x^2 + 9c_1^2x}}} + \frac{c_1^2}{4}}$$

$$-\frac{1}{2} \frac{c_1^3}{\sqrt[3]{\sqrt{-768x^3 + 81c_1^4x^2 + 9c_1^2x}}} - \frac{\sqrt[3]{\sqrt{-768x^3 + 81c_1^4x^2 + 9c_1^2x}}}{\sqrt[3]{23^{2/3}}} + \frac{c_1}{4}$$

$$y(x) \rightarrow -\frac{1}{2} \sqrt{\frac{\sqrt[3]{\sqrt{-768x^3 + 81c_1^4x^2 + 9c_1^2x}}}{\sqrt[3]{23^{2/3}}} + \frac{4\sqrt[3]{\frac{2}{3}x}}{\sqrt[3]{\sqrt{-768x^3 + 81c_1^4x^2 + 9c_1^2x}}} + \frac{c_1^2}{4}}$$

$$+\frac{1}{2} \frac{c_1^3}{\sqrt[3]{\sqrt{-768x^3 + 81c_1^4x^2 + 9c_1^2x}}} - \frac{\sqrt[3]{\sqrt{-768x^3 + 81c_1^4x^2 + 9c_1^2x}}}{\sqrt[3]{23^{2/3}}} + \frac{c_1}{4}$$

$$y(x) \rightarrow \frac{1}{2} \sqrt{\frac{\sqrt[3]{\sqrt{-768x^3 + 81c_1^4x^2 + 9c_1^2x}}}{\sqrt[3]{23^{2/3}}} + \frac{4\sqrt[3]{\frac{2}{3}x}}{\sqrt[3]{\sqrt{-768x^3 + 81c_1^4x^2 + 9c_1^2x}}} + \frac{c_1^2}{4}}$$

$$-\frac{1}{2} \frac{c_1^3}{\sqrt[3]{\sqrt{-768x^3 + 81c_1^4x^2 + 9c_1^2x}}} - \frac{\sqrt[3]{\sqrt{-768x^3 + 81c_1^4x^2 + 9c_1^2x}}}{\sqrt[3]{23^{2/3}}} + \frac{c_1}{4}$$

$$1 \frac{\sqrt[3]{\sqrt{-768x^3 + 81c_1^4x^2 + 9c_1^2x}}}{\sqrt[3]{23^{2/3}}} + \frac{4\sqrt[3]{\frac{2}{3}x}}{\sqrt[3]{\sqrt{-768x^3 + 81c_1^4x^2 + 9c_1^2x}}} + \frac{c_1^2}{4}$$

## 6.13 problem 7.5 (c)

Internal problem ID [13096]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.5 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, [_Abel, '2nd type', 'cl`

$$\frac{2y}{x} + (4x^2y - 3)y' = 0$$

✓ Solution by Maple

Time used: 1.953 (sec). Leaf size: 28

```
dsolve(2*y(x)/x+(4*x^2*y(x)-3)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \frac{\text{RootOf}(-Z^{32}c_1 - Z^{24}c_1 - x^8)^8}{x^2}$$

✓ Solution by Mathematica

Time used: 60.256 (sec). Leaf size: 1985

`DSolve[2*y[x]/x+(4*x^2*y[x]-3)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{1}{4x^2}$$

$$-\frac{1}{2} \sqrt{\frac{1}{4x^4} + \frac{4\sqrt[3]{\frac{2}{3}}e^{-8c_1x^2}}{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}\sqrt{e^{24c_1}x^4(-256x^8 + 27e^{8c_1})}}} + \frac{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}\sqrt{e^{24c_1}x^4(-256x^8 + 27e^{8c_1})}}}{\sqrt[3]{23^{2/3}x^2}}}$$

$$-\frac{1}{2} \sqrt{\frac{1}{2x^4} - \frac{4\sqrt[3]{\frac{2}{3}}e^{-8c_1x^2}}{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}\sqrt{e^{24c_1}x^4(-256x^8 + 27e^{8c_1})}}} - \frac{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}\sqrt{e^{24c_1}x^4(-256x^8 + 27e^{8c_1})}}}{\sqrt[3]{23^{2/3}x^2}}}$$

$$y(x) \rightarrow \frac{1}{4x^2}$$

$$-\frac{1}{2} \sqrt{\frac{1}{4x^4} + \frac{4\sqrt[3]{\frac{2}{3}}e^{-8c_1x^2}}{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}\sqrt{e^{24c_1}x^4(-256x^8 + 27e^{8c_1})}}} + \frac{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}\sqrt{e^{24c_1}x^4(-256x^8 + 27e^{8c_1})}}}{\sqrt[3]{23^{2/3}x^2}}}$$

$$+\frac{1}{2} \sqrt{\frac{1}{2x^4} - \frac{4\sqrt[3]{\frac{2}{3}}e^{-8c_1x^2}}{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}\sqrt{e^{24c_1}x^4(-256x^8 + 27e^{8c_1})}}} - \frac{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}\sqrt{e^{24c_1}x^4(-256x^8 + 27e^{8c_1})}}}{\sqrt[3]{23^{2/3}x^2}}}$$

$$y(x) \rightarrow \frac{1}{4x^2}$$

$$+\frac{1}{2} \sqrt{\frac{1}{4x^4} + \frac{4\sqrt[3]{\frac{2}{3}}e^{-8c_1x^2}}{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}\sqrt{e^{24c_1}x^4(-256x^8 + 27e^{8c_1})}}} + \frac{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}\sqrt{e^{24c_1}x^4(-256x^8 + 27e^{8c_1})}}}{\sqrt[3]{23^{2/3}x^2}}}$$

$$-\frac{1}{2} \sqrt{\frac{1}{2x^4} - \frac{4\sqrt[3]{\frac{2}{3}}e^{-8c_1x^2}}{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}\sqrt{e^{24c_1}x^4(-256x^8 + 27e^{8c_1})}}} - \frac{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}\sqrt{e^{24c_1}x^4(-256x^8 + 27e^{8c_1})}}}{\sqrt[3]{23^{2/3}x^2}}}$$

$$y(x) \rightarrow \frac{1}{4x^2}$$

## 6.14 problem 7.5 (d)

Internal problem ID [13097]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.5 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x)*G(y),0]']]`

$$(1 - x \tan(y)) y' = -1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 115

```
dsolve(1+(1-x*tan(y(x)))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \arctan \left( -\frac{x(c_1x + \sqrt{-c_1^2 + x^2 + 1})}{x^2 + 1} + c_1, \frac{c_1x + \sqrt{-c_1^2 + x^2 + 1}}{x^2 + 1} \right)$$

$$y = \arctan \left( \frac{x(-c_1x + \sqrt{-c_1^2 + x^2 + 1})}{x^2 + 1} + c_1, -\frac{-c_1x + \sqrt{-c_1^2 + x^2 + 1}}{x^2 + 1} \right)$$

✓ Solution by Mathematica

Time used: 0.108 (sec). Leaf size: 145

```
DSolve[1+(1-x*Tan[y[x]])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sec^{-1}\left(\frac{c_1x - \sqrt{x^2 + 1 - c_1^2}}{-1 + c_1^2}\right)$$

$$y(x) \rightarrow \sec^{-1}\left(\frac{c_1x - \sqrt{x^2 + 1 - c_1^2}}{-1 + c_1^2}\right)$$

$$y(x) \rightarrow -\sec^{-1}\left(\frac{\sqrt{x^2 + 1 - c_1^2} + c_1x}{-1 + c_1^2}\right)$$

$$y(x) \rightarrow \sec^{-1}\left(\frac{\sqrt{x^2 + 1 - c_1^2} + c_1x}{-1 + c_1^2}\right)$$

## 6.15 problem 7.5 (e)

Internal problem ID [13098]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.5 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$3y + 3y^2 + (2x + 4xy)y' = 0$$

✓ Solution by Maple

Time used: 0.515 (sec). Leaf size: 137

```
dsolve(3*y(x)+3*y(x)^2+(2*x+4*x*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \frac{-\frac{c_1 x}{2} - \frac{\sqrt{c_1^2 x^2 - 4\sqrt{c_1 x}}}{2}}{c_1 x}$$

$$y = \frac{-\frac{c_1 x}{2} + \frac{\sqrt{c_1^2 x^2 - 4\sqrt{c_1 x}}}{2}}{c_1 x}$$

$$y = \frac{-\frac{c_1 x}{2} - \frac{\sqrt{c_1^2 x^2 + 4\sqrt{c_1 x}}}{2}}{c_1 x}$$

$$y = \frac{-\frac{c_1 x}{2} + \frac{\sqrt{c_1^2 x^2 + 4\sqrt{c_1 x}}}{2}}{c_1 x}$$

✓ Solution by Mathematica

Time used: 6.61 (sec). Leaf size: 67

```
DSolve[3*y[x]+3*y[x]^2+(2*x+4*x*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left( -1 - \sqrt{1 + \frac{4e^{c_1}}{x^{3/2}}} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left( -1 + \sqrt{1 + \frac{4e^{c_1}}{x^{3/2}}} \right)$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 0$$



## 6.16 problem 7.5 (f)

Internal problem ID [13099]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.5 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$2x(y + 1) - y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(2*x*(y(x)+1)-diff(y(x),x)=0,y(x), singsol=all)
```

$$y = -1 + c_1 e^{x^2}$$

### ✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 20

```
DSolve[2*x*(y[x]+1)-y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -1 + c_1 e^{x^2}$$

$$y(x) \rightarrow -1$$

## 6.17 problem 7.5 (g)

Internal problem ID [13100]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.5 (g).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, [_Abel, '2nd type', 'cl`

$$2y^3 + (4y^3x^3 - 3y^2x) y' = 0$$

✓ Solution by Maple

Time used: 1.062 (sec). Leaf size: 32

```
dsolve(2*y(x)^3+(4*x^3*y(x)^3-3*x*y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = 0$$

$$y = \frac{\text{RootOf}(\_Z^{32}c_1 - \_Z^{24}c_1 - x^8)^8}{x^2}$$

✓ Solution by Mathematica

Time used: 60.187 (sec). Leaf size: 1990

`DSolve[2*y[x]^3+(4*x^3*y[x]^3-3*x*y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow \frac{1}{4x^2}$$

$$-\frac{1}{2} \sqrt{\frac{1}{4x^4} + \frac{4\sqrt[3]{\frac{2}{3}}e^{-8c_1x^2}}{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}}\sqrt{e^{24c_1x^4}(-256x^8 + 27e^{8c_1})}}} + \frac{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}}\sqrt{e^{24c_1x^4}(-256x^8 + 27e^{8c_1})}}{\sqrt[3]{23^{2/3}x^2}}$$

$$-\frac{1}{2} \sqrt{\frac{1}{2x^4} - \frac{4\sqrt[3]{\frac{2}{3}}e^{-8c_1x^2}}{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}}\sqrt{e^{24c_1x^4}(-256x^8 + 27e^{8c_1})}}} - \frac{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}}\sqrt{e^{24c_1x^4}(-256x^8 + 27e^{8c_1})}}{\sqrt[3]{23^{2/3}x^2}}$$

$$y(x) \rightarrow \frac{1}{4x^2}$$

$$-\frac{1}{2} \sqrt{\frac{1}{4x^4} + \frac{4\sqrt[3]{\frac{2}{3}}e^{-8c_1x^2}}{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}}\sqrt{e^{24c_1x^4}(-256x^8 + 27e^{8c_1})}}} + \frac{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}}\sqrt{e^{24c_1x^4}(-256x^8 + 27e^{8c_1})}}{\sqrt[3]{23^{2/3}x^2}}$$

$$+\frac{1}{2} \sqrt{\frac{1}{2x^4} - \frac{4\sqrt[3]{\frac{2}{3}}e^{-8c_1x^2}}{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}}\sqrt{e^{24c_1x^4}(-256x^8 + 27e^{8c_1})}}} - \frac{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}}\sqrt{e^{24c_1x^4}(-256x^8 + 27e^{8c_1})}}{\sqrt[3]{23^{2/3}x^2}}$$

$$y(x) \rightarrow \frac{1}{4x^2}$$

$$+\frac{1}{2} \sqrt{\frac{1}{4x^4} + \frac{4\sqrt[3]{\frac{2}{3}}e^{-8c_1x^2}}{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}}\sqrt{e^{24c_1x^4}(-256x^8 + 27e^{8c_1})}}} + \frac{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}}\sqrt{e^{24c_1x^4}(-256x^8 + 27e^{8c_1})}}{\sqrt[3]{23^{2/3}x^2}}$$

$$-\frac{1}{2} \sqrt{\frac{1}{2x^4} - \frac{4\sqrt[3]{\frac{2}{3}}e^{-8c_1x^2}}{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}}\sqrt{e^{24c_1x^4}(-256x^8 + 27e^{8c_1})}}} - \frac{\sqrt[3]{9e^{-8c_1x^2} + \sqrt{3}e^{-24c_1}}\sqrt{e^{24c_1x^4}(-256x^8 + 27e^{8c_1})}}{\sqrt[3]{23^{2/3}x^2}}$$

## 6.18 problem 7.5 (h)

Internal problem ID [13101]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.5 (h).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, [_Abel, '2nd type', 'cl`

$$4xy + (3x^2 + 5y) y' = 0$$

✓ Solution by Maple

Time used: 0.593 (sec). Leaf size: 29

```
dsolve(4*x*y(x)+(3*x^2+5*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \text{RootOf}(x^5\_Z^{25} + x^5\_Z^{15} - c_1)^{10} x^2$$

✓ Solution by Mathematica

Time used: 60.064 (sec). Leaf size: 1121

`DSolve[4*x*y[x]+(3*x^2+5*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow -\frac{3x^2}{5}$$

$$+\frac{1}{5\text{Root}[\#1^{10}(11664x^{20} + 11664e^{60c_1}) - 9720\#1^8x^{16} + 1080\#1^7x^{14} + 3105\#1^6x^{12} - 666\#1^5x^{10} - 42]}$$

$$y(x) \rightarrow -\frac{3x^2}{5}$$

$$+\frac{1}{5\text{Root}[\#1^{10}(11664x^{20} + 11664e^{60c_1}) - 9720\#1^8x^{16} + 1080\#1^7x^{14} + 3105\#1^6x^{12} - 666\#1^5x^{10} - 42]}$$

$$y(x) \rightarrow -\frac{3x^2}{5}$$

$$+\frac{1}{5\text{Root}[\#1^{10}(11664x^{20} + 11664e^{60c_1}) - 9720\#1^8x^{16} + 1080\#1^7x^{14} + 3105\#1^6x^{12} - 666\#1^5x^{10} - 42]}$$

$$y(x) \rightarrow -\frac{3x^2}{5}$$

$$+\frac{1}{5\text{Root}[\#1^{10}(11664x^{20} + 11664e^{60c_1}) - 9720\#1^8x^{16} + 1080\#1^7x^{14} + 3105\#1^6x^{12} - 666\#1^5x^{10} - 42]}$$

$$y(x) \rightarrow -\frac{3x^2}{5}$$

$$+\frac{1}{5\text{Root}[\#1^{10}(11664x^{20} + 11664e^{60c_1}) - 9720\#1^8x^{16} + 1080\#1^7x^{14} + 3105\#1^6x^{12} - 666\#1^5x^{10} - 42]}$$

$$y(x) \rightarrow -\frac{3x^2}{5}$$

$$+\frac{1}{5\text{Root}[\#1^{10}(11664x^{20} + 11664e^{60c_1}) - 9720\#1^8x^{16} + 1080\#1^7x^{14} + 3105\#1^6x^{12} - 666\#1^5x^{10} - 42]}$$

$$y(x) \rightarrow -\frac{3x^2}{5}$$

$$+\frac{1}{5\text{Root}[\#1^{10}(11664x^{20} + 11664e^{60c_1}) - 9720\#1^8x^{16} + 1080\#1^7x^{14} + 3105\#1^6x^{12} - 666\#1^5x^{10} - 42]}$$

$$y(x) \rightarrow -\frac{3x^2}{5}$$

$$+\frac{1}{5\text{Root}[\#1^{10}(11664x^{20} + 11664e^{60c_1}) - 9720\#1^8x^{16} + 1080\#1^7x^{14} + 3105\#1^6x^{12} - 666\#1^5x^{10} - 42]}$$

$$y(x) \rightarrow -\frac{3x^2}{5}$$

$$+\frac{1}{5\text{Root}[\#1^{10}(11664x^{20} + 11664e^{60c_1}) - 9720\#1^8x^{16} + 1080\#1^7x^{14} + 3105\#1^6x^{12} - 666\#1^5x^{10} - 42]}$$

## 6.19 problem 7.5 (i)

Internal problem ID [13102]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 7. The exact form and general integrating factors. Additional exercises. page 141

**Problem number:** 7.5 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational]`

$$12y^2x^2 + \left(7x^3y + \frac{x}{y}\right)y' = -6$$

✓ Solution by Maple

Time used: 0.609 (sec). Leaf size: 59

```
dsolve(6+12*x^2*y(x)^2+(7*x^3*y(x)+x/y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \text{RootOf} \left( x^{10} \_Z^{35} - x^{10} \_Z^{30} - \frac{1}{c_1^2} \right)^{15} x^4 \left( \text{RootOf} \left( x^{10} \_Z^{35} - x^{10} \_Z^{30} - \frac{1}{c_1^2} \right)^5 - 1 \right) c_1$$

✓ Solution by Mathematica

Time used: 3.003 (sec). Leaf size: 330

```
DSolve[6+12*x^2*y[x]^2+(7*x^3*y[x]+x/y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{Root} \left[ -\#1^7 - \frac{3\#1^5}{x^2} - \frac{3\#1^3}{x^4} - \frac{\#1}{x^6} + \frac{e^{c_1}}{x^{12}} \&, 1 \right]$$

$$y(x) \rightarrow \text{Root} \left[ -\#1^7 - \frac{3\#1^5}{x^2} - \frac{3\#1^3}{x^4} - \frac{\#1}{x^6} + \frac{e^{c_1}}{x^{12}} \&, 2 \right]$$

$$y(x) \rightarrow \text{Root} \left[ -\#1^7 - \frac{3\#1^5}{x^2} - \frac{3\#1^3}{x^4} - \frac{\#1}{x^6} + \frac{e^{c_1}}{x^{12}} \&, 3 \right]$$

$$y(x) \rightarrow \text{Root} \left[ -\#1^7 - \frac{3\#1^5}{x^2} - \frac{3\#1^3}{x^4} - \frac{\#1}{x^6} + \frac{e^{c_1}}{x^{12}} \&, 4 \right]$$

$$y(x) \rightarrow \text{Root} \left[ -\#1^7 - \frac{3\#1^5}{x^2} - \frac{3\#1^3}{x^4} - \frac{\#1}{x^6} + \frac{e^{c_1}}{x^{12}} \&, 5 \right]$$

$$y(x) \rightarrow \text{Root} \left[ -\#1^7 - \frac{3\#1^5}{x^2} - \frac{3\#1^3}{x^4} - \frac{\#1}{x^6} + \frac{e^{c_1}}{x^{12}} \&, 6 \right]$$

$$y(x) \rightarrow \text{Root} \left[ -\#1^7 - \frac{3\#1^5}{x^2} - \frac{3\#1^3}{x^4} - \frac{\#1}{x^6} + \frac{e^{c_1}}{x^{12}} \&, 7 \right]$$

## 7 Chapter 8. Review exercises for part of part II. page 143

7.1	problem 1	216
7.2	problem 2	217
7.3	problem 3	218
7.4	problem 4	219
7.5	problem 5	220
7.6	problem 6	221
7.7	problem 7	222
7.8	problem 8	223
7.9	problem 9	224
7.10	problem 10	225
7.11	problem 11	227
7.12	problem 12	228
7.13	problem 13	229
7.14	problem 14	230
7.15	problem 15	231
7.16	problem 16	233
7.17	problem 17	234
7.18	problem 18	235
7.19	problem 19	236
7.20	problem 20	237
7.21	problem 21	238
7.22	problem 22	239
7.23	problem 23	240
7.24	problem 24	241
7.25	problem 25	242
7.26	problem 26	243
7.27	problem 27	244
7.28	problem 28	245
7.29	problem 29	246
7.30	problem 30	247
7.31	problem 31	248
7.32	problem 32	249
7.33	problem 33	250
7.34	problem 34	252
7.35	problem 35	253
7.36	problem 36	254



7.37	problem 37	256
7.38	problem 38	257
7.39	problem 39	258
7.40	problem 40	259
7.41	problem 41	260
7.42	problem 42	261
7.43	problem 43	262
7.44	problem 44	263
7.45	problem 45	264
7.46	problem 46	265
7.47	problem 47	267
7.48	problem 48	268
7.49	problem 49	269
7.50	problem 50	270

## 7.1 problem 1

Internal problem ID [13103]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 1.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y'x - 2y = -6x^3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(x*diff(y(x),x)=2*y(x)-6*x^3,y(x), singsol=all)
```

$$y = (-6x + c_1) x^2$$

### ✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 15

```
DSolve[x*y'[x]==2*y[x]-6*x^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2(-6x + c_1)$$

## 7.2 problem 2

Internal problem ID [13104]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 2.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y'x - 2y^2 + 6y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(x*diff(y(x),x)=2*y(x)^2-6*y(x),y(x), singsol=all)
```

$$y = \frac{3}{3c_1x^6 + 1}$$

### ✓ Solution by Mathematica

Time used: 1.886 (sec). Leaf size: 31

```
DSolve[x*y'[x]==2*y[x]^2-6*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3}{1 + e^{3c_1x^6}}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow 3$$

### 7.3 problem 3

Internal problem ID [13105]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 3.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$4y^2 - y^2x^2 + y' = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(4*y(x)^2-x^2*y(x)^2+diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \frac{3}{-x^3 + 3c_1 + 12x}$$

#### ✓ Solution by Mathematica

Time used: 0.125 (sec). Leaf size: 25

```
DSolve[4*y[x]^2-x^2*y[x]^2+y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{3}{x^3 - 12x + 3c_1}$$

$$y(x) \rightarrow 0$$

## 7.4 problem 4

Internal problem ID [13106]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 4.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$y' - \sqrt{x+y} = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 42

```
dsolve(diff(y(x),x)=sqrt(x+y(x)),y(x), singsol=all)
```

$$x - 2\sqrt{x+y} - \ln(\sqrt{x+y} - 1) + \ln(\sqrt{x+y} + 1) + \ln(x+y-1) - c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 8.647 (sec). Leaf size: 59

```
DSolve[y'[x]==Sqrt[x+y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow W\left(-e^{-\frac{x}{2}-1-\frac{c_1}{2}}\right)^2 + 2W\left(-e^{-\frac{x}{2}-1-\frac{c_1}{2}}\right) - x + 1$$

$$y(x) \rightarrow 1 - x$$

## 7.5 problem 5

Internal problem ID [13107]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 5.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y'x^2 = \sqrt{x} + 3$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(x^2*diff(y(x),x)-sqrt(x)=3,y(x), singsol=all)
```

$$y = -\frac{2}{\sqrt{x}} - \frac{3}{x} + c_1$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 20

```
DSolve[x^2*y'[x]-Sqrt[x]==3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2}{\sqrt{x}} - \frac{3}{x} + c_1$$

## 7.6 problem 6

Internal problem ID [13108]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 6.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _dAlembert]`

$$xyy' - y^2 - \sqrt{x^4 + y^2x^2} = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(x*y(x)*diff(y(x),x)-y(x)^2=sqrt(x^4+x^2*y(x)^2),y(x), singsol=all)
```

$$-\frac{x^2 + y^2}{\sqrt{x^4 + y^2x^2}} + \ln(x) - c_1 = 0$$

### ✓ Solution by Mathematica

Time used: 0.278 (sec). Leaf size: 54

```
DSolve[x*y[x]*y'[x]-y[x]^2==Sqrt[x^4+x^2*y[x]^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x\sqrt{\log^2(x) + 2c_1 \log(x) - 1 + c_1^2}$$

$$y(x) \rightarrow x\sqrt{\log^2(x) + 2c_1 \log(x) - 1 + c_1^2}$$

## 7.7 problem 7

Internal problem ID [13109]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 7.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class C'], _Riccati]`

$$y' + 2xy - y^2 = x^2$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve(diff(y(x),x)=y(x)^2-2*x*y(x)+x^2,y(x), singsol=all)
```

$$y = \frac{x e^{2x} c_1 - e^{2x} c_1 - x - 1}{e^{2x} c_1 - 1}$$

### ✓ Solution by Mathematica

Time used: 0.128 (sec). Leaf size: 29

```
DSolve[y'[x]==y[x]^2-2*x*y[x]+x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x + \frac{1}{\frac{1}{2} + c_1 e^{2x}} - 1$$

$$y(x) \rightarrow x - 1$$



## 7.8 problem 8

Internal problem ID [13110]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 8.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$4xy + y'x^2 = 6$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(4*x*y(x)-6+x^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \frac{2x^3 + c_1}{x^4}$$

### ✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 17

```
DSolve[4*x*y[x]-6+x^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2x^3 + c_1}{x^4}$$

## 7.9 problem 9

Internal problem ID [13111]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 9.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _exact, _rational, _Bernoulli]`

$$y^2x + x^2yy' = 6$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(x*y(x)^2-6+x^2*y(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \frac{\sqrt{12x + c_1}}{x}$$

$$y = -\frac{\sqrt{12x + c_1}}{x}$$

### ✓ Solution by Mathematica

Time used: 0.19 (sec). Leaf size: 38

```
DSolve[x*y[x]^2-6+x^2*y[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{12x + c_1}}{x}$$

$$y(x) \rightarrow \frac{\sqrt{12x + c_1}}{x}$$

## 7.10 problem 10

Internal problem ID [13112]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 10.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _Bernoulli]`

$$y^3 + y'xy^2 = -x^3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 96

```
dsolve(x^3+y(x)^3+x*y(x)^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \frac{(-4x^6 + 8c_1)^{\frac{1}{3}}}{2x}$$

$$y = \frac{-\frac{(-4x^6+8c_1)^{\frac{1}{3}}}{4} - \frac{i\sqrt{3}(-4x^6+8c_1)^{\frac{1}{3}}}{4}}{x}$$

$$y = \frac{-\frac{(-4x^6+8c_1)^{\frac{1}{3}}}{4} + \frac{i\sqrt{3}(-4x^6+8c_1)^{\frac{1}{3}}}{4}}{x}$$

✓ Solution by Mathematica

Time used: 0.232 (sec). Leaf size: 80

```
DSolve[x^3+y[x]^3+x*y[x]^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt[3]{-\frac{1}{2}}\sqrt[3]{-x^6+2c_1}}{x}$$

$$y(x) \rightarrow \frac{\sqrt[3]{-\frac{x^6}{2}+c_1}}{x}$$

$$y(x) \rightarrow \frac{(-1)^{2/3}\sqrt[3]{-\frac{x^6}{2}+c_1}}{x}$$

## 7.11 problem 11

Internal problem ID [13113]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 11.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y'x + 3y = x^3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(3*y(x)-x^3+x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \frac{\frac{x^6}{6} + c_1}{x^3}$$

### ✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 19

```
DSolve[3*y[x]-x^3+x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{6} + \frac{c_1}{x^3}$$

## 7.12 problem 12

Internal problem ID [13114]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 12.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_exact, \_rational, \_Bernoulli]

$$2y^2x + (2x^2y + 2y)y' = -1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 50

```
dsolve(1+2*x*y(x)^2+(2*x^2*y(x)+2*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \frac{\sqrt{(x^2 + 1)(-x + c_1)}}{x^2 + 1}$$
$$y = -\frac{\sqrt{(x^2 + 1)(-x + c_1)}}{x^2 + 1}$$

### ✓ Solution by Mathematica

Time used: 0.212 (sec). Leaf size: 50

```
DSolve[1+2*x*y[x]^2+(2*x^2*y[x]+2*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{-x + c_1}}{\sqrt{x^2 + 1}}$$
$$y(x) \rightarrow \frac{\sqrt{-x + c_1}}{\sqrt{x^2 + 1}}$$

## 7.13 problem 13

Internal problem ID [13115]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 13.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class D'], _rational, _Bernoulli]`

$$3xy^3 - y + y'x = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve(3*x*y(x)^3-y(x)+x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = \frac{x}{\sqrt{2x^3 + c_1}}$$

$$y = -\frac{x}{\sqrt{2x^3 + c_1}}$$

### ✓ Solution by Mathematica

Time used: 0.182 (sec). Leaf size: 43

```
DSolve[3*x*y[x]^3-y[x]+x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x}{\sqrt{2x^3 + c_1}}$$

$$y(x) \rightarrow \frac{x}{\sqrt{2x^3 + c_1}}$$

$$y(x) \rightarrow 0$$

## 7.14 problem 14

Internal problem ID [13116]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 14.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$-2xy + (x^2 + 1)y' = -2x^2 - 2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(2+2*x^2-2*x*y(x)+(x^2+1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = (x^2 + 1)(c_1 - 2 \arctan(x))$$

### ✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 18

```
DSolve[2+2*x^2-2*x*y[x]+(x^2+1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x^2 + 1)(-2 \arctan(x) + c_1)$$



## 7.15 problem 15

Internal problem ID [13117]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 15.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$(y^2 - 4)y' - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve((y(x)^2-4)*diff(y(x),x)=y(x),y(x), singsol=all)
```

$$y = e^{-\frac{\text{LambertW}\left(-e^{-\frac{x}{2}-\frac{c_1}{2}}\right)}{2} - \frac{x}{4} - \frac{c_1}{4}}$$

✓ Solution by Mathematica

Time used: 32.653 (sec). Leaf size: 246

```
DSolve[(y[x]^2-4)*y'[x]==y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2i\sqrt{W\left(-\frac{1}{4}\sqrt[4]{e^{-2(x+c_1)}}\right)}$$

$$y(x) \rightarrow 2i\sqrt{W\left(-\frac{1}{4}\sqrt[4]{e^{-2(x+c_1)}}\right)}$$

$$y(x) \rightarrow -2i\sqrt{W\left(-\frac{1}{4}i\sqrt[4]{e^{-2(x+c_1)}}\right)}$$

$$y(x) \rightarrow 2i\sqrt{W\left(-\frac{1}{4}i\sqrt[4]{e^{-2(x+c_1)}}\right)}$$

$$y(x) \rightarrow -2i\sqrt{W\left(\frac{1}{4}i\sqrt[4]{e^{-2(x+c_1)}}\right)}$$

$$y(x) \rightarrow 2i\sqrt{W\left(\frac{1}{4}i\sqrt[4]{e^{-2(x+c_1)}}\right)}$$

$$y(x) \rightarrow -2i\sqrt{W\left(\frac{1}{4}\sqrt[4]{e^{-2(x+c_1)}}\right)}$$

$$y(x) \rightarrow 2i\sqrt{W\left(\frac{1}{4}\sqrt[4]{e^{-2(x+c_1)}}\right)}$$

$$y(x) \rightarrow 0$$

## 7.16 problem 16

Internal problem ID [13118]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 16.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$(x^2 - 4) y' = x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve((x^2-4)*diff(y(x),x)=x,y(x), singsol=all)
```

$$y = \frac{\ln(x^2 - 4)}{2} + c_1$$

### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 18

```
DSolve[(x^2-4)*y'[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \log(x^2 - 4) + c_1$$

## 7.17 problem 17

Internal problem ID [13119]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 17.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y' - \frac{1}{xy - 3x} = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
dsolve(diff(y(x),x)=1/(x*y(x)-3*x),y(x), singsol=all)
```

$$y = 3 - \sqrt{9 + 2 \ln(x) + 2c_1}$$

$$y = 3 + \sqrt{9 + 2 \ln(x) + 2c_1}$$

### ✓ Solution by Mathematica

Time used: 0.099 (sec). Leaf size: 43

```
DSolve[y'[x]==1/(x*y[x]-3*x),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3 - \sqrt{2 \log(x) + 9 + 2c_1}$$

$$y(x) \rightarrow 3 + \sqrt{2 \log(x) + 9 + 2c_1}$$

## 7.18 problem 18

Internal problem ID [13120]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 18.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _rational, _Bernoulli]`

$$y' - \frac{3y}{x+1} + y^2 = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 42

```
dsolve(diff(y(x),x)=3*y(x)/(1+x)-y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{4x^3 + 12x^2 + 12x + 4}{x^4 + 4x^3 + 6x^2 + 4c_1 + 4x + 1}$$

### ✓ Solution by Mathematica

Time used: 0.258 (sec). Leaf size: 41

```
DSolve[y'[x]==3*y[x]/(1+x)-y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{4(x+1)^3}{x^4 + 4x^3 + 6x^2 + 4x + 1 + 4c_1}$$

$$y(x) \rightarrow 0$$

## 7.19 problem 19

Internal problem ID [13121]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 19.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_exact, [_1st_order, ‘_with_symmetry_[F(x)*G(y),0]’]]`

$$\sin(y) + (x + y) \cos(y) y' = 0$$

### ✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 26

```
dsolve(sin(y(x))+(x+y(x))*cos(y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$x - \frac{-\cos(y(x)) - \sin(y(x))y(x) + c_1}{\sin(y(x))} = 0$$

### ✓ Solution by Mathematica

Time used: 0.193 (sec). Leaf size: 29

```
DSolve[Sin[y[x]]+(x+y[x])*Cos[y[x]]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}[x = \csc(y(x))(-y(x) \sin(y(x)) - \cos(y(x))) + c_1 \csc(y(x)), y(x)]$$

## 7.20 problem 20

Internal problem ID [13122]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 20.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$\sin(y) + (x + 1) \cos(y) y' = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve(sin(y(x))+(1+x)*cos(y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \arcsin\left(\frac{1}{c_1(1+x)}\right)$$

### ✓ Solution by Mathematica

Time used: 19.016 (sec). Leaf size: 21

```
DSolve[Sin[y[x]]+(1+x)*Cos[y[x]]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arcsin\left(\frac{e^{c_1}}{x+1}\right)$$

$$y(x) \rightarrow 0$$

## 7.21 problem 21

Internal problem ID [13123]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 21.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$2 \cos(x) y' = -\sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(sin(x)+2*cos(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\ln(\cos(x))}{2} + c_1$$

### ✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 15

```
DSolve[Sin[x]+2*Cos[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \log(\cos(x)) + c_1$$



## 7.22 problem 22

Internal problem ID [13124]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 22.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$xyy' - 2y^2 = 2x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(x*y(x)*diff(y(x),x)=2*(x^2+y(x)^2),y(x), singsol=all)
```

$$y(x) = \sqrt{c_1x^2 - 2}x$$

$$y(x) = -\sqrt{c_1x^2 - 2}x$$

### ✓ Solution by Mathematica

Time used: 0.42 (sec). Leaf size: 38

```
DSolve[x*y[x]*y'[x]==2*(x^2+y[x]^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x\sqrt{-2 + c_1x^2}$$

$$y(x) \rightarrow x\sqrt{-2 + c_1x^2}$$

## 7.23 problem 23

Internal problem ID [13125]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 23.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{x + 2y}{x + 2y + 3} = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)=(x+2*y(x))/(x+2*y(x)+3),y(x), singsol=all)
```

$$y(x) = -\frac{x}{2} + \text{LambertW}\left(\frac{e^{\frac{3x}{2}} e^{\frac{1}{2}c_1}}{2}\right) - \frac{1}{2}$$

### ✓ Solution by Mathematica

Time used: 3.703 (sec). Leaf size: 41

```
DSolve[y'[x]==(x+2*y[x])/(x+2*y[x]+3),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow W\left(-e^{\frac{3x}{2}-1+c_1}\right) - \frac{x}{2} - \frac{1}{2}$$

$$y(x) \rightarrow \frac{1}{2}(-x - 1)$$

## 7.24 problem 24

Internal problem ID [13126]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 24.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{x + 2y}{2x - y} = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
dsolve(diff(y(x),x)=(x+2*y(x))/(2*x-y(x)),y(x), singsol=all)
```

$$y(x) = \tan \left( \text{RootOf} \left( -4\_Z + \ln \left( \frac{1}{\cos(\_Z)^2} \right) + 2 \ln(x) + 2c_1 \right) \right) x$$

### ✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 36

```
DSolve[y'[x]==(x+2*y[x])/(2*x-y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[ \frac{1}{2} \log \left( \frac{y(x)^2}{x^2} + 1 \right) - 2 \arctan \left( \frac{y(x)}{x} \right) = -\log(x) + c_1, y(x) \right]$$

## 7.25 problem 25

Internal problem ID [13127]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 25.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _dAlembert]`

$$y' - \frac{y}{x} - \tan\left(\frac{y}{x}\right) = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)=y(x)/x+tan(y(x)/x),y(x), singsol=all)
```

$$y(x) = \arcsin(c_1 x) x$$

### ✓ Solution by Mathematica

Time used: 4.007 (sec). Leaf size: 19

```
DSolve[y'[x]==y[x]/x+Tan[y[x]/x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \arcsin(e^{c_1} x)$$

$$y(x) \rightarrow 0$$

## 7.26 problem 26

Internal problem ID [13128]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 26.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y' - y^2x - 3y^2 = x + 3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)=x*y(x)^2+3*y(x)^2+x+3,y(x), singsol=all)
```

$$y(x) = \tan\left(\frac{1}{2}x^2 + c_1 + 3x\right)$$

### ✓ Solution by Mathematica

Time used: 0.206 (sec). Leaf size: 19

```
DSolve[y'[x]==x*y[x]^2+3*y[x]^2+x+3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan\left(\frac{x^2}{2} + 3x + c_1\right)$$

## 7.27 problem 27

Internal problem ID [13129]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 27.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_homogeneous, 'class C'], [_Abel, '2nd type', 'class C'], _d`

$$-(x + 2y)y' = -1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(1-(x+2*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\text{LambertW}\left(-\frac{c_1 e^{-\frac{x}{2}-1}}{2}\right) - \frac{x}{2} - 1$$

### ✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 30

```
DSolve[1-(x+2*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -W\left(-\frac{1}{2}c_1 e^{-\frac{x}{2}-1}\right) - \frac{x}{2} - 1$$

## 7.28 problem 28

Internal problem ID [13130]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 28.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_exact, [_1st_order, ‘_with_symmetry_[F(x)*G(y),0]’]]`

$$\ln(y) + \left(\frac{x}{y} + 3\right) y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(ln(y(x))+(x/y(x)+3)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = e^{-\frac{x \operatorname{LambertW}\left(\frac{3e^{\frac{c_1}{x}}}{x}\right) - c_1}{x}}$$

### ✓ Solution by Mathematica

Time used: 0.925 (sec). Leaf size: 29

```
DSolve[Log[y[x]]+(x/y[x]+3)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} x W\left(\frac{3e^{\frac{c_1}{x}}}{x}\right)$$

$$y(x) \rightarrow 1$$

## 7.29 problem 29

Internal problem ID [13131]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 29.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y^2 - y' = -1$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve(y(x)^2+1-diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \tan(x + c_1)$$

### ✓ Solution by Mathematica

Time used: 0.113 (sec). Leaf size: 24

```
DSolve[y[x]^2+1-y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan(x + c_1)$$

$$y(x) \rightarrow -i$$

$$y(x) \rightarrow i$$



## 7.30 problem 30

Internal problem ID [13132]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 30.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' - 3y = 12e^{2x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)-3*y(x)=12*exp(2*x),y(x), singsol=all)
```

$$y(x) = (-12e^{-x} + c_1) e^{3x}$$

### ✓ Solution by Mathematica

Time used: 0.05 (sec). Leaf size: 19

```
DSolve[y'[x]-3*y[x]==12*Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(-12 + c_1 e^x)$$

## 7.31 problem 31

Internal problem ID [13133]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 31.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$xyy' - yx - y^2 = x^2$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve(x*y(x)*diff(y(x),x)=x^2+x*y(x)+y(x)^2,y(x), singsol=all)
```

$$y(x) = e^{-\text{LambertW}\left(-\frac{e^{-c_1}e^{-1}}{x}\right)-c_1-1} - x$$

### ✓ Solution by Mathematica

Time used: 4.224 (sec). Leaf size: 31

```
DSolve[x*y[x]*y'[x]==x^2+x*y[x]+y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x \left( 1 + W \left( -\frac{e^{-1-c_1}}{x} \right) \right)$$

$$y(x) \rightarrow -x$$

## 7.32 problem 32

Internal problem ID [13134]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 32.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$(x + 2)y' = x^3$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve((x+2)*diff(y(x),x)-x^3=0,y(x), singsol=all)
```

$$y(x) = \frac{x^3}{3} - x^2 + 4x - 8 \ln(x + 2) + c_1$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 32

```
DSolve[(x+2)*y'[x]-x^3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{3} - x^2 + 4x - 8 \log(x + 2) + \frac{44}{3} + c_1$$

### 7.33 problem 33

Internal problem ID [13135]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 33.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class D'], _rational, _Bernoulli]`

$$xy^3y' - y^4 = -x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 73

```
dsolve(x*y(x)^3*diff(y(x),x)=y(x)^4-x^2,y(x), singsol=all)
```

$$y(x) = (c_1x^4 + 2x^2)^{\frac{1}{4}}$$

$$y(x) = -(c_1x^4 + 2x^2)^{\frac{1}{4}}$$

$$y(x) = -i(c_1x^4 + 2x^2)^{\frac{1}{4}}$$

$$y(x) = i(c_1x^4 + 2x^2)^{\frac{1}{4}}$$

✓ Solution by Mathematica

Time used: 0.341 (sec). Leaf size: 96

```
DSolve[x*y[x]^3*y'[x]==y[x]^4-x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{x} \sqrt[4]{2 + c_1 x^2}$$

$$y(x) \rightarrow -i\sqrt{x} \sqrt[4]{2 + c_1 x^2}$$

$$y(x) \rightarrow i\sqrt{x} \sqrt[4]{2 + c_1 x^2}$$

$$y(x) \rightarrow \sqrt{x} \sqrt[4]{2 + c_1 x^2}$$

## 7.34 problem 34

Internal problem ID [13136]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 34.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _Bernoulli]`

$$y' - 4y + \frac{16e^{4x}}{y^2} = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 103

```
dsolve(diff(y(x),x)=4*y(x)-16*exp(4*x)/y(x)^2,y(x), singsol=all)
```

$$y(x) = (e^{12x}c_1 + 6e^{4x})^{\frac{1}{3}}$$

$$y(x) = -\frac{(e^{12x}c_1 + 6e^{4x})^{\frac{1}{3}}}{2} - \frac{i\sqrt{3}(e^{12x}c_1 + 6e^{4x})^{\frac{1}{3}}}{2}$$

$$y(x) = -\frac{(e^{12x}c_1 + 6e^{4x})^{\frac{1}{3}}}{2} + \frac{i\sqrt{3}(e^{12x}c_1 + 6e^{4x})^{\frac{1}{3}}}{2}$$

### ✓ Solution by Mathematica

Time used: 3.622 (sec). Leaf size: 90

```
DSolve[y'[x]==4*y[x]-16*Exp[4*x]/y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{4x/3} \sqrt[3]{6 + c_1 e^{8x}}$$

$$y(x) \rightarrow -\sqrt[3]{-1} e^{4x/3} \sqrt[3]{6 + c_1 e^{8x}}$$

$$y(x) \rightarrow (-1)^{2/3} e^{4x/3} \sqrt[3]{6 + c_1 e^{8x}}$$

## 7.35 problem 35

Internal problem ID [13137]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 35.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$2y + (x + 1)y' = 6x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve((2*y(x)-6*x)+(x+1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{2x^3 + 3x^2 + c_1}{(1+x)^2}$$

### ✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 24

```
DSolve[(2*y[x]-6*x)+(x+1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2x^3 + 3x^2 + c_1}{(x+1)^2}$$

## 7.36 problem 36

Internal problem ID [13138]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 36.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _exact, _rational]`

$$y^2x + (yx^2 + 10y^4)y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(x*y(x)^2+(x^2*y(x)+10*y(x)^4)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$\frac{x^2y(x)^2}{2} + 2y(x)^5 + c_1 = 0$$



✓ Solution by Mathematica

Time used: 3.953 (sec). Leaf size: 141

```
DSolve[x*y[x]^2+(x^2*y[x]+10*y[x]^4)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow \text{Root}[4\#1^5 + \#1^2x^2 - 2c_1\&, 1]$$

$$y(x) \rightarrow \text{Root}[4\#1^5 + \#1^2x^2 - 2c_1\&, 2]$$

$$y(x) \rightarrow \text{Root}[4\#1^5 + \#1^2x^2 - 2c_1\&, 3]$$

$$y(x) \rightarrow \text{Root}[4\#1^5 + \#1^2x^2 - 2c_1\&, 4]$$

$$y(x) \rightarrow \text{Root}[4\#1^5 + \#1^2x^2 - 2c_1\&, 5]$$

$$y(x) \rightarrow 0$$

## 7.37 problem 37

Internal problem ID [13139]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 37.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [Bernoulli]

$$yy' - y^2x = 6xe^{4x^2}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 43

```
dsolve(y(x)*diff(y(x),x)-x*y(x)^2=6*x*exp(4*x^2),y(x), singsol=all)
```

$$y(x) = \sqrt{e^{x^2}c_1 + 2e^{4x^2}}$$

$$y(x) = -\sqrt{e^{x^2}c_1 + 2e^{4x^2}}$$

### ✓ Solution by Mathematica

Time used: 1.953 (sec). Leaf size: 62

```
DSolve[y[x]*y'[x]-x*y[x]^2==6*x*Exp[4*x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -e^{\frac{x^2}{2}} \sqrt{2e^{3x^2} + c_1}$$

$$y(x) \rightarrow e^{\frac{x^2}{2}} \sqrt{2e^{3x^2} + c_1}$$

## 7.38 problem 38

Internal problem ID [13140]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 38.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _exact, _rational, _dAlembert]`

$$(y - x + 3)^2 (y' - 1) = 1$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 80

```
dsolve((y(x)-x+3)^2*(diff(y(x),x)-1)=1,y(x), singsol=all)
```

$$y(x) = (-3c_1 + 3x)^{\frac{1}{3}} + x - 3$$

$$y(x) = -\frac{(-3c_1 + 3x)^{\frac{1}{3}}}{2} - \frac{i\sqrt{3}(-3c_1 + 3x)^{\frac{1}{3}}}{2} + x - 3$$

$$y(x) = -\frac{(-3c_1 + 3x)^{\frac{1}{3}}}{2} + \frac{i\sqrt{3}(-3c_1 + 3x)^{\frac{1}{3}}}{2} + x - 3$$

### ✓ Solution by Mathematica

Time used: 0.421 (sec). Leaf size: 95

```
DSolve[(y[x]-x+3)^2*(y'[x]-1)==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x + \sqrt[3]{3}\sqrt[3]{x+9+c_1} - 3$$

$$y(x) \rightarrow x + \frac{1}{2}i\sqrt[3]{3}(\sqrt{3}+i)\sqrt[3]{x+9+c_1} - 3$$

$$y(x) \rightarrow x - \frac{1}{2}\sqrt[3]{3}(1+i\sqrt{3})\sqrt[3]{x+9+c_1} - 3$$

## 7.39 problem 39

Internal problem ID [13141]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 39.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_exact, [_1st_order, ‘_with_symmetry_[F(x),G(x)*y+H(x)]’]]`

$$y e^{yx} + x e^{yx} y' = -x$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve(x+y(x)*exp(x*y(x))+x*exp(x*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\ln\left(-\frac{x^2}{2} - c_1\right)}{x}$$

### ✓ Solution by Mathematica

Time used: 0.412 (sec). Leaf size: 20

```
DSolve[x+y[x]*Exp[x*y[x]]+x*Exp[x*y[x]]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\log\left(-\frac{x^2}{2} + c_1\right)}{x}$$

## 7.40 problem 40

Internal problem ID [13142]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 40.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y^2 - y^2 \cos(x) + y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(y(x)^2-y(x)^2*cos(x)+diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{1}{\sin(x) - c_1 - x}$$

### ✓ Solution by Mathematica

Time used: 0.123 (sec). Leaf size: 22

```
DSolve[y[x]^2-y[x]^2*Cos[x]+y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{x - \sin(x) - c_1}$$

$$y(x) \rightarrow 0$$

## 7.41 problem 41

Internal problem ID [13143]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 41.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 2y = \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)+2*y(x)=sin(x),y(x), singsol=all)
```

$$y(x) = -\frac{\cos(x)}{5} + \frac{2 \sin(x)}{5} + e^{-2x} c_1$$

### ✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 26

```
DSolve[y'[x]+2*y[x]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2 \sin(x)}{5} - \frac{\cos(x)}{5} + c_1 e^{-2x}$$

## 7.42 problem 42

Internal problem ID [13144]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 42.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = -2x + \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)+2*x=sin(x),y(x), singsol=all)
```

$$y(x) = -x^2 - \cos(x) + c_1$$

### ✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 17

```
DSolve[y'[x]+2*x==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x^2 - \cos(x) + c_1$$

## 7.43 problem 43

Internal problem ID [13145]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 43.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y' - y^3 + y^3 \cos(x) = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x),x)=y(x)^3-y(x)^3*cos(x),y(x), singsol=all)
```

$$y(x) = \frac{1}{\sqrt{c_1 - 2x + 2 \sin(x)}}$$
$$y(x) = -\frac{1}{\sqrt{c_1 - 2x + 2 \sin(x)}}$$

### ✓ Solution by Mathematica

Time used: 0.219 (sec). Leaf size: 55

```
DSolve[y'[x]==y[x]^3-y[x]^3*Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\sqrt{2}\sqrt{-x + \sin(x) - c_1}}$$
$$y(x) \rightarrow \frac{1}{\sqrt{2}\sqrt{-x + \sin(x) - c_1}}$$
$$y(x) \rightarrow 0$$



## 7.44 problem 44

Internal problem ID [13146]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 44.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_exact, [\_1st\_order, ['\_with\_symmetry\_[F(x),G(x)\*y+H(x)]]]]

$$y^2 e^{y^2 x} + 2xy e^{y^2 x} y' = 2x$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 40

```
dsolve(y(x)^2*exp(x*y(x)^2)-2*x+2*x*y(x)*exp(x*y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{x \ln(x^2 - c_1)}}{x}$$

$$y(x) = -\frac{\sqrt{x \ln(x^2 - c_1)}}{x}$$

### ✓ Solution by Mathematica

Time used: 1.468 (sec). Leaf size: 44

```
DSolve[y[x]^2*Exp[x*y[x]^2]-2*x+2*x*y[x]*Exp[x*y[x]^2]*y'[x]==0,y[x],x,IncludeSingularSoluti
```

$$y(x) \rightarrow -\frac{\sqrt{\log(x^2 + c_1)}}{\sqrt{x}}$$

$$y(x) \rightarrow \frac{\sqrt{\log(x^2 + c_1)}}{\sqrt{x}}$$

## 7.45 problem 45

Internal problem ID [13147]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 45.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y' - e^{4x+3y} = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)=exp(4*x+3*y(x)),y(x), singsol=all)
```

$$y(x) = -\frac{\ln\left(-\frac{3e^{4x}}{4} - 3c_1\right)}{3}$$

### ✓ Solution by Mathematica

Time used: 0.884 (sec). Leaf size: 24

```
DSolve[y'[x]==Exp[4*x+3*y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{3} \log\left(-\frac{3}{4}(e^{4x} + 4c_1)\right)$$

## 7.46 problem 46

Internal problem ID [13148]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 46.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _dAlembert]`

$$y' - \tan(6x + 3y + 1) = -2$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 185

```
dsolve(diff(y(x),x)=tan(6*x+3*y(x)+1)-2,y(x), singsol=all)
```

$$y(x) = \frac{\arctan\left(c_1 e^{3x}, e^{6x} c_1^2 \sqrt{\frac{e^{-6x}\left(\frac{e^{-6x}}{c_1^2} - 1\right)}{c_1^2}}\right)}{3} - 2x - \frac{1}{3}$$

$$y(x) = \frac{\arctan\left(c_1 e^{3x}, -e^{6x} c_1^2 \sqrt{\frac{e^{-6x}\left(\frac{e^{-6x}}{c_1^2} - 1\right)}{c_1^2}}\right)}{3} - 2x - \frac{1}{3}$$

$$y(x) = \frac{\arctan\left(-c_1 e^{3x}, e^{6x} c_1^2 \sqrt{\frac{e^{-6x}\left(\frac{e^{-6x}}{c_1^2} - 1\right)}{c_1^2}}\right)}{3} - 2x - \frac{1}{3}$$

$$y(x) = \frac{\arctan\left(-c_1 e^{3x}, -e^{6x} c_1^2 \sqrt{\frac{e^{-6x}\left(\frac{e^{-6x}}{c_1^2} - 1\right)}{c_1^2}}\right)}{3} - 2x - \frac{1}{3}$$

✓ Solution by Mathematica

Time used: 60.483 (sec). Leaf size: 25

```
DSolve[y'[x]==Tan[6*x+3*y[x]+1]-2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}(\arcsin(e^{3x-3c_1}) - 6x - 1)$$

## 7.47 problem 47

Internal problem ID [13149]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 47.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$y' - e^{4x+3y} = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)=exp(4*x+3*y(x)),y(x), singsol=all)
```

$$y(x) = -\frac{\ln\left(-\frac{3e^{4x}}{4} - 3c_1\right)}{3}$$

### ✓ Solution by Mathematica

Time used: 0.872 (sec). Leaf size: 24

```
DSolve[y'[x]==Exp[4*x+3*y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{3} \log\left(-\frac{3}{4}(e^{4x} + 4c_1)\right)$$

## 7.48 problem 48

Internal problem ID [13150]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 48.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y' - x(6y + e^{x^2}) = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)=x*(6*y(x)+exp(x^2)),y(x), singsol=all)
```

$$y(x) = -\frac{e^{x^2}}{4} + e^{3x^2} c_1$$

### ✓ Solution by Mathematica

Time used: 0.071 (sec). Leaf size: 25

```
DSolve[y'[x]==x*(6*y[x]+Exp[x^2]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{e^{x^2}}{4} + c_1 e^{3x^2}$$

## 7.49 problem 49

Internal problem ID [13151]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 49.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[_exact, _rational, [_1st_order, ' _with_symmetry_[F(x)*G(y),0]`

$$x(1 - 2y) + (y - x^2) y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 45

```
dsolve(x*(1-2*y(x))+(y(x)-x^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = x^2 - \sqrt{x^4 - x^2 - 2c_1}$$

$$y(x) = x^2 + \sqrt{x^4 - x^2 - 2c_1}$$

### ✓ Solution by Mathematica

Time used: 0.147 (sec). Leaf size: 66

```
DSolve[x*(1-2*y[x])+(y[x]-x^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2 - i\sqrt{-x^4 + x^2 - c_1}$$

$$y(x) \rightarrow x^2 + i\sqrt{-x^4 + x^2 - c_1}$$

$$y(x) \rightarrow \frac{1}{2}$$

## 7.50 problem 50

Internal problem ID [13152]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 8. Review exercises for part of part II. page 143

**Problem number:** 50.

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$y'x^2 + 3yx = 6e^{-x^2}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(x^2*diff(y(x),x)+3*x*y(x)=6*exp(-x^2),y(x), singsol=all)
```

$$y(x) = \frac{-3e^{-x^2} + c_1}{x^3}$$

✓ Solution by Mathematica

Time used: 0.079 (sec). Leaf size: 21

```
DSolve[x^2*y'[x]+3*x*y[x]==6*Exp[-x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-3e^{-x^2} + c_1}{x^3}$$



## 8 Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

8.1	problem 13.1 (a)	273
8.2	problem 13.1 (b)	274
8.3	problem 13.1 (c)	275
8.4	problem 13.1 (d)	276
8.5	problem 13.1 (e)	277
8.6	problem 13.1 (f)	278
8.7	problem 13.2 (a)	279
8.8	problem 13.2 (b)	280
8.9	problem 13.2 (c)	281
8.10	problem 13.2 (d)	282
8.11	problem 13.2 (e)	283
8.12	problem 13.2 (f)	284
8.13	problem 13.2 (g)	285
8.14	problem 13.2 (h)	286
8.15	problem 13.2 (i)	287
8.16	problem 13.3 (a)	288
8.17	problem 13.3 (b)	289
8.18	problem 13.3 (c)	290
8.19	problem 13.3 (d)	291
8.20	problem 13.4 (a)	292
8.21	problem 13.4 (b)	293
8.22	problem 13.4 (c)	294
8.23	problem 13.4 (d)	295
8.24	problem 13.4 (e)	296
8.25	problem 13.4 (f)	297
8.26	problem 13.5 (a)	300
8.27	problem 13.5 (c)	301
8.28	problem 13.5 (d)	302
8.29	problem 13.5 (e)	303
8.30	problem 13.5 (f)	304
8.31	problem 13.5 (g)	305
8.32	problem 13.5 (h)	306
8.33	problem 13.5 (i)	307
8.34	problem 13.5 (j)	308

8.35	problem 13.6 (a)	309
8.36	problem 13.6 (b)	310
8.37	problem 13.6 (c)	311
8.38	problem 13.6 (d)	312
8.39	problem 13.6 (e)	313
8.40	problem 13.6 (f)	314
8.41	problem 13.6 (g)	315
8.42	problem 13.6 (h)	316
8.43	problem 13.7 (c)	317
8.44	problem 13.7 (d)	318
8.45	problem 13.7 (e)	319
8.46	problem 13.8 (i)	320
8.47	problem 13.8 (ii)	321
8.48	problem 13.8 (iii)	322
8.49	problem 13.8 (iv)	323
8.50	problem 13.9 (i)	324
8.51	problem 13.9 (ii)	325
8.52	problem 13.9 (iii)	326
8.53	problem 13.9 (iv)	327

## 8.1 problem 13.1 (a)

Internal problem ID [13153]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.1 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' + 4y' = 18x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x*diff(y(x),x$2)+4*diff(y(x),x)=18*x^2,y(x), singsol=all)
```

$$y = x^3 - \frac{c_1}{3x^3} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 20

```
DSolve[x*y''[x]+4*y'[x]==18*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^3 - \frac{c_1}{3x^3} + c_2$$

## 8.2 problem 13.1 (b)

Internal problem ID [13154]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.1 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' - 2y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(x*diff(y(x),x$2)=2*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = c_2x^3 + c_1$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 19

```
DSolve[y''[x]==2*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}c_1e^{2x} + c_2$$

### 8.3 problem 13.1 (c)

Internal problem ID [13155]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.1 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y' = 0$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 10

```
dsolve(diff(y(x),x$2)=diff(y(x),x),y(x), singsol=all)
```

$$y(x) = c_1 + c_2 e^x$$

#### ✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 14

```
DSolve[y''[x]==y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x + c_2$$

## 8.4 problem 13.1 (d)

Internal problem ID [13156]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.1 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' + 2y' = 8e^{2x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)=8*exp(2*x),y(x), singsol=all)
```

$$y(x) = -\frac{e^{-2x}c_1}{2} + e^{2x} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.059 (sec). Leaf size: 24

```
DSolve[y''[x]+2*y'[x]==8*Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x} - \frac{1}{2}c_1e^{-2x} + c_2$$

## 8.5 problem 13.1 (e)

Internal problem ID [13157]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.1 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' - y' + 2y'x^2 = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(x*diff(y(x),x$2)=diff(y(x),x)-2*x^2*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = c_1 + e^{-x^2} c_2$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 21

```
DSolve[x*y''[x]==y'[x]-2*x^2*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \frac{1}{2}c_1 e^{-x^2}$$

## 8.6 problem 13.1 (f)

Internal problem ID [13158]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.1 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$(x^2 + 1) y'' + 2xy' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve((x^2+1)*diff(y(x),x$2)+2*x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1 + \arctan(x) c_2$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 13

```
DSolve[(x^2+1)*y'[x]+2*x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \arctan(x) + c_2$$



## 8.7 problem 13.2 (a)

Internal problem ID [13159]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.2 (a).

**ODE order:** 2.

**ODE degree:** 2.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]`

$$y'' - 4x\sqrt{y'} = 0$$

### ✓ Solution by Maple

Time used: 0.172 (sec). Leaf size: 51

```
dsolve(diff(y(x),x$2)=4*x*sqrt(diff(y(x),x)),y(x), singsol=all)
```

$$y(x) = c_1$$

$$y(x) = \frac{x^5}{5} - \frac{2x^3}{3c_1} + \frac{x}{c_1^2} + c_2$$

$$y(x) = \frac{x^5}{5} + \frac{2x^3}{3c_1} + \frac{x}{c_1^2} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.095 (sec). Leaf size: 33

```
DSolve[y''[x]==4*x*Sqrt[y'[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^5}{5} + \frac{c_1 x^3}{3} + \frac{c_1^2 x}{4} + c_2$$

## 8.8 problem 13.2 (b)

Internal problem ID [13160]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.2 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear], [`

$$y'y'' = 1$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 33

```
dsolve(diff(y(x),x)*diff(y(x),x$2)=1,y(x), singsol=all)
```

$$y(x) = \frac{(2c_1 + 2x)^{\frac{3}{2}}}{3} + c_2$$

$$y(x) = -\frac{(2c_1 + 2x)^{\frac{3}{2}}}{3} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 49

```
DSolve[y'[x]*y''[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \frac{2}{3}\sqrt{2}(x + c_1)^{3/2}$$

$$y(x) \rightarrow \frac{2}{3}\sqrt{2}(x + c_1)^{3/2} + c_2$$

## 8.9 problem 13.2 (c)

Internal problem ID [13161]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.2 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear],`

$$yy'' + y'^2 = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 33

```
dsolve(y(x)*diff(y(x),x$2)=-((diff(y(x),x)^2),y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = \sqrt{2c_1x + 2c_2}$$

$$y(x) = -\sqrt{2c_1x + 2c_2}$$

### ✓ Solution by Mathematica

Time used: 0.181 (sec). Leaf size: 20

```
DSolve[y[x]*y'[x]==-(y'[x]^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2\sqrt{2x - c_1}$$

## 8.10 problem 13.2 (d)

Internal problem ID [13162]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.2 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]`

$$xy'' - y'^2 + y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x*diff(y(x),x$2)=diff(y(x),x)^2-diff(y(x),x),y(x), singsol=all)
```

$$y(x) = -\frac{\ln(c_1x - 1)}{c_1} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.301 (sec). Leaf size: 38

```
DSolve[x*y'[x]==(y'[x])^2-y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-c_1} \log(1 + e^{c_1}x) + c_2$$

$$y(x) \rightarrow c_2$$

$$y(x) \rightarrow x + c_2$$

## 8.11 problem 13.2 (e)

Internal problem ID [13163]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.2 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' - y'^2 = 6x^5$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 63

```
dsolve(x*diff(y(x),x$2)-diff(y(x),x)^2=6*x^5,y(x), singsol=all)
```

$$y(x) = \int \frac{\sqrt{6} x^{\frac{5}{2}} \left( \text{BesselY} \left( 1, \frac{2x^{\frac{5}{2}}\sqrt{6}}{5} \right) c_1 + \text{BesselJ} \left( 1, \frac{2x^{\frac{5}{2}}\sqrt{6}}{5} \right) \right)}{c_1 \text{BesselY} \left( 0, \frac{2x^{\frac{5}{2}}\sqrt{6}}{5} \right) + \text{BesselJ} \left( 0, \frac{2x^{\frac{5}{2}}\sqrt{6}}{5} \right)} dx + c_2$$

### ✓ Solution by Mathematica

Time used: 60.384 (sec). Leaf size: 109

```
DSolve[x*y''[x]-y'[x]^2==6*x^5,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \int_1^x \frac{\sqrt{6} (2 \text{BesselY} (1, \frac{2}{5}\sqrt{6}K[1]^{5/2}) + \text{BesselJ} (1, \frac{2}{5}\sqrt{6}K[1]^{5/2}) c_1) K[1]^{5/2}}{2 \text{BesselY} (0, \frac{2}{5}\sqrt{6}K[1]^{5/2}) + \text{BesselJ} (0, \frac{2}{5}\sqrt{6}K[1]^{5/2}) c_1} dK[1] + c_2$$

## 8.12 problem 13.2 (f)

Internal problem ID [13164]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.2 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1],`

$$yy'' - y'^2 - y' = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 23

```
dsolve(y(x)*diff(y(x),x$2)-(diff(y(x),x)^2)=diff(y(x),x),y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = \frac{e^{c_2 c_1} e^{c_1 x} + 1}{c_1}$$

### ✓ Solution by Mathematica

Time used: 1.7 (sec). Leaf size: 26

```
DSolve[y[x]*y'[x]-(y'[x]^2)==y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1 + e^{c_1(x+c_2)}}{c_1}$$

$$y(x) \rightarrow \text{Indeterminate}$$

## 8.13 problem 13.2 (g)

Internal problem ID [13165]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.2 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' = -6$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x$2)=2*diff(y(x),x)-6,y(x), singsol=all)
```

$$y(x) = \frac{e^{2x}c_1}{2} + 3x + c_2$$

### ✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 22

```
DSolve[y''[x]==2*y'[x]-6,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3x + \frac{1}{2}c_1e^{2x} + c_2$$

## 8.14 problem 13.2 (h)

Internal problem ID [13166]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.2 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible]`

$$(y - 3)y'' - 2y'^2 = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 25

```
dsolve((y(x)-3)*diff(y(x),x$2)=2*diff(y(x),x)^2,y(x), singsol=all)
```

$$y(x) = 3$$

$$y(x) = \frac{3c_1x + 3c_2 - 1}{c_1x + c_2}$$

### ✓ Solution by Mathematica

Time used: 0.183 (sec). Leaf size: 44

```
DSolve[(y[x]-3)*y'[x]==2*y'[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3c_1x - 1 + 3c_2c_1}{c_1(x + c_2)}$$

$$y(x) \rightarrow 3$$

$$y(x) \rightarrow \text{Indeterminate}$$

$$y(x) \rightarrow 3$$



## 8.15 problem 13.2 (i)

Internal problem ID [13167]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.2 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' + 4y' = 9e^{-3x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)=9*exp(-3*x),y(x), singsol=all)
```

$$y(x) = -\frac{e^{-4x}c_1}{4} - 3e^{-3x} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.074 (sec). Leaf size: 26

```
DSolve[y''[x]+4*y'[x]==9*Exp[-3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -3e^{-3x} - \frac{1}{4}c_1e^{-4x} + c_2$$

## 8.16 problem 13.3 (a)

Internal problem ID [13168]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.3 (a).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - y'' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x$3)=diff(y(x),x$2),y(x), singsol=all)
```

$$y(x) = c_1 + c_2x + c_3e^x$$

### ✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 18

```
DSolve[y'''[x]==y''[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1e^x + c_3x + c_2$$

## 8.17 problem 13.3 (b)

Internal problem ID [13169]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.3 (b).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_y]]`

$$xy''' + 2y'' = 6x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(x*diff(y(x),x$3)+2*diff(y(x),x$2)=6*x,y(x), singsol=all)
```

$$y(x) = \frac{x^3}{3} - c_1 \ln(x) + c_2 x + c_3$$

### ✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 25

```
DSolve[x*y'''[x]+2*y''[x]==6*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{3} + c_3 x - c_1 \log(x) + c_2$$

## 8.18 problem 13.3 (c)

Internal problem ID [13170]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.3 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_3rd_order, _missing_x], [_3rd_order, _missing_y], [_3rd_order,`

✓ Solution by Maple

Time used: 0.203 (sec). Leaf size: 36

```
dsolve(diff(y(x),x$3)=2*sqrt(diff(y(x),x$2)),y(x), singsol=all)
```

$$y(x) = c_1x + c_2$$

$$y(x) = \frac{1}{12}x^4 + \frac{1}{3}c_1x^3 + \frac{1}{2}c_1^2x^2 + c_2x + c_3$$

✓ Solution by Mathematica

Time used: 0.13 (sec). Leaf size: 39

```
DSolve[y'''[x]==2*Sqrt[y''[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^4}{12} + \frac{c_1x^3}{6} + \frac{c_1^2x^2}{8} + c_3x + c_2$$

## 8.19 problem 13.3 (d)

Internal problem ID [13171]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.3 (d).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 2y''' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$4)=-2*diff(y(x),x$3),y(x), singsol=all)
```

$$y(x) = c_1 + c_2x + c_3x^2 + c_4e^{-2x}$$

### ✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 28

```
DSolve[y''''[x]==-2*y'''[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{8}c_1e^{-2x} + x(c_4x + c_3) + c_2$$

## 8.20 problem 13.4 (a)

Internal problem ID [13172]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.4 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible]`

$$yy'' - y'^2 = 0$$

With initial conditions

$$[y(0) = 5, y'(0) = 15]$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 10

```
dsolve([y(x)*diff(y(x),x$2)=diff(y(x),x)^2,y(0) = 5, D(y)(0) = 15],y(x), singsol=all)
```

$$y(x) = 5e^{3x}$$

### ✓ Solution by Mathematica

Time used: 0.129 (sec). Leaf size: 12

```
DSolve[{y[x]*y'[x]==y'[x]^2,{y[0]==5,y'[0]==15}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 5e^{3x}$$

## 8.21 problem 13.4 (b)

Internal problem ID [13173]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.4 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible`

$$3yy'' - 2y'^2 = 0$$

With initial conditions

$$[y(0) = 8, y'(0) = 6]$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 11

```
dsolve([3*y(x)*diff(y(x),x$2)=2*diff(y(x),x)^2,y(0) = 8, D(y)(0) = 6],y(x), singsol=all)
```

$$y(x) = \frac{(x+4)^3}{8}$$

### ✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 14

```
DSolve[{3*y[x]*y'[x]==2*y'[x]^2,{y[0]==8,y'[0]==6}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{1}{8}(x+4)^3$$

## 8.22 problem 13.4 (c)

Internal problem ID [13174]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.4 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear], _`

$$\sin(y) y'' + \cos(y) y'^2 = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(sin(y(x))*diff(y(x),x$2)+cos(y(x))*diff(y(x),x)^2=0,y(x), singsol=all)
```

$$y(x) = \pi - \arccos(c_1 x + c_2)$$

### ✓ Solution by Mathematica

Time used: 11.859 (sec). Leaf size: 29

```
DSolve[Sin[y[x]]*y'[x]+Cos[y[x]]*y'[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\arccos(-c_1(x + c_2))$$

$$y(x) \rightarrow \arccos(-c_1(x + c_2))$$



## 8.23 problem 13.4 (d)

Internal problem ID [13175]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.4 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x$2)=diff(y(x),x),y(x), singsol=all)
```

$$y(x) = c_1 + c_2 e^x$$

### ✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 14

```
DSolve[y''[x]==y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x + c_2$$

## 8.24 problem 13.4 (e)

Internal problem ID [13176]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.4 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear],`

$$yy'' + y'^2 - 2yy' = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 37

```
dsolve(diff(y(x),x)^2+y(x)*diff(y(x),x$2)=2*y(x)*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = \sqrt{e^{2x}c_1 + 2c_2}$$

$$y(x) = -\sqrt{e^{2x}c_1 + 2c_2}$$

### ✓ Solution by Mathematica

Time used: 0.788 (sec). Leaf size: 38

```
DSolve[y'[x]^2+y[x]*y''[x]==2*y[x]*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2\sqrt{e^{2x} + e^{c_1}}$$

$$y(x) \rightarrow c_2\sqrt{e^{2x}}$$

## 8.25 problem 13.4 (f)

Internal problem ID [13177]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.4 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear], _`

$$y^2 y'' + y'' + 2y y'^2 = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 393

`dsolve(y(x)^2*diff(y(x),x$2)+diff(y(x),x$2)+2*y(x)*diff(y(x),x)^2=0,y(x), singsol=all)`

$$y(x) = -i$$

$$y(x) = i$$

$$y(x) = \frac{\left(12c_1x + 12c_2 + 4\sqrt{9c_1^2x^2 + 18c_1c_2x + 9c_2^2 + 4}\right)^{\frac{1}{3}}}{2} - \frac{\left(12c_1x + 12c_2 + 4\sqrt{9c_1^2x^2 + 18c_1c_2x + 9c_2^2 + 4}\right)^{\frac{1}{3}}}{2}$$

$$y(x) = -\frac{\left(12c_1x + 12c_2 + 4\sqrt{9c_1^2x^2 + 18c_1c_2x + 9c_2^2 + 4}\right)^{\frac{1}{3}}}{4} + \frac{\left(12c_1x + 12c_2 + 4\sqrt{9c_1^2x^2 + 18c_1c_2x + 9c_2^2 + 4}\right)^{\frac{1}{3}}}{1} - \frac{i\sqrt{3} \left( \frac{\left(12c_1x + 12c_2 + 4\sqrt{9c_1^2x^2 + 18c_1c_2x + 9c_2^2 + 4}\right)^{\frac{1}{3}}}{2} + \frac{2}{\left(12c_1x + 12c_2 + 4\sqrt{9c_1^2x^2 + 18c_1c_2x + 9c_2^2 + 4}\right)^{\frac{1}{3}}} \right)}{2}$$

$$y(x) = -\frac{\left(12c_1x + 12c_2 + 4\sqrt{9c_1^2x^2 + 18c_1c_2x + 9c_2^2 + 4}\right)^{\frac{1}{3}}}{4} + \frac{\left(12c_1x + 12c_2 + 4\sqrt{9c_1^2x^2 + 18c_1c_2x + 9c_2^2 + 4}\right)^{\frac{1}{3}}}{1} + \frac{i\sqrt{3} \left( \frac{\left(12c_1x + 12c_2 + 4\sqrt{9c_1^2x^2 + 18c_1c_2x + 9c_2^2 + 4}\right)^{\frac{1}{3}}}{2} + \frac{2}{\left(12c_1x + 12c_2 + 4\sqrt{9c_1^2x^2 + 18c_1c_2x + 9c_2^2 + 4}\right)^{\frac{1}{3}}} \right)}{2}$$

✓ Solution by Mathematica

Time used: 54.871 (sec). Leaf size: 307

`DSolve[y[x]^2*y'[x]+y''[x]+2*y[x]*y'[x]^2==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{-2 + \sqrt[3]{2} \left( 3c_1x + \sqrt{4 + 9c_1^2(x + c_2)^2} + 3c_2c_1 \right)^{2/3}}{2^{2/3} \sqrt[3]{3c_1x + \sqrt{4 + 9c_1^2(x + c_2)^2} + 3c_2c_1}}$$

$$y(x) \rightarrow \frac{i(\sqrt{3} + i) \sqrt[3]{3c_1x + \sqrt{4 + 9c_1^2(x + c_2)^2} + 3c_2c_1}}{2\sqrt[3]{2}} + \frac{1 + i\sqrt{3}}{2^{2/3} \sqrt[3]{3c_1x + \sqrt{4 + 9c_1^2(x + c_2)^2} + 3c_2c_1}}$$

$$y(x) \rightarrow \frac{1 - i\sqrt{3}}{2^{2/3} \sqrt[3]{3c_1x + \sqrt{4 + 9c_1^2(x + c_2)^2} + 3c_2c_1}} - \frac{i(\sqrt{3} - i) \sqrt[3]{3c_1x + \sqrt{4 + 9c_1^2(x + c_2)^2} + 3c_2c_1}}{2\sqrt[3]{2}}$$

## 8.26 problem 13.5 (a)

Internal problem ID [13178]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.5 (a).

**ODE order:** 2.

**ODE degree:** 2.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]`

$$y'' - 4x\sqrt{y'} = 0$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 51

```
dsolve(diff(y(x),x$2)=4*x*sqrt(diff(y(x),x)),y(x), singsol=all)
```

$$y(x) = c_1$$

$$y(x) = \frac{x^5}{5} - \frac{2x^3}{3c_1} + \frac{x}{c_1^2} + c_2$$

$$y(x) = \frac{x^5}{5} + \frac{2x^3}{3c_1} + \frac{x}{c_1^2} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.071 (sec). Leaf size: 33

```
DSolve[y''[x]==4*x*Sqrt[y'[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^5}{5} + \frac{c_1 x^3}{3} + \frac{c_1^2 x}{4} + c_2$$

## 8.27 problem 13.5 (c)

Internal problem ID [13179]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.5 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear], [`

$$y'y'' = 1$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 33

```
dsolve(diff(y(x),x)*diff(y(x),x$2)=1,y(x), singsol=all)
```

$$y(x) = \frac{(2c_1 + 2x)^{\frac{3}{2}}}{3} + c_2$$

$$y(x) = -\frac{(2c_1 + 2x)^{\frac{3}{2}}}{3} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 49

```
DSolve[y'[x]*y''[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \frac{2}{3}\sqrt{2}(x + c_1)^{3/2}$$

$$y(x) \rightarrow \frac{2}{3}\sqrt{2}(x + c_1)^{3/2} + c_2$$

## 8.28 problem 13.5 (d)

Internal problem ID [13180]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.5 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]`

$$xy'' - y'^2 + y' = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve(x*diff(y(x),x$2)=diff(y(x),x)^2-diff(y(x),x),y(x), singsol=all)
```

$$y(x) = -\frac{\ln(c_1x - 1)}{c_1} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.338 (sec). Leaf size: 38

```
DSolve[x*y''[x]==y'[x]^2-y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-c_1} \log(1 + e^{c_1}x) + c_2$$

$$y(x) \rightarrow c_2$$

$$y(x) \rightarrow x + c_2$$



## 8.29 problem 13.5 (e)

Internal problem ID [13181]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.5 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' - y' = 6x^5$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x*diff(y(x),x$2)-diff(y(x),x)=6*x^5,y(x), singsol=all)
```

$$y(x) = \frac{1}{4}x^6 + \frac{1}{2}c_1x^2 + c_2$$

### ✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 24

```
DSolve[x*y''[x]-y'[x]==6*x^5,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}(x^6 + 2c_1x^2 + 4c_2)$$

## 8.30 problem 13.5 (f)

Internal problem ID [13182]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.5 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1],`

$$yy'' - y'^2 - y' = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 23

```
dsolve(y(x)*diff(y(x),x$2)-diff(y(x),x)^2=diff(y(x),x),y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = \frac{e^{c_2 c_1} e^{c_1 x} + 1}{c_1}$$

### ✓ Solution by Mathematica

Time used: 1.761 (sec). Leaf size: 26

```
DSolve[y[x]*y'[x]-y'[x]^2==y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1 + e^{c_1(x+c_2)}}{c_1}$$

$$y(x) \rightarrow \text{Indeterminate}$$

### 8.31 problem 13.5 (g)

Internal problem ID [13183]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.5 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible]`

$$yy'' - 2y'^2 = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(y(x)*diff(y(x),x$2)=2*diff(y(x),x)^2,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = -\frac{1}{c_1x + c_2}$$

#### ✓ Solution by Mathematica

Time used: 0.14 (sec). Leaf size: 19

```
DSolve[y[x]*y'[x]==2*y'[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2}{x + c_1}$$

$$y(x) \rightarrow 0$$

## 8.32 problem 13.5 (h)

Internal problem ID [13184]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.5 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible]`

$$(y - 3)y'' - y'^2 = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve((y(x)-3)*diff(y(x),x$2)=diff(y(x),x)^2,y(x), singsol=all)
```

$$y(x) = 3$$

$$y(x) = e^{c_1 x} c_2 + 3$$

### ✓ Solution by Mathematica

Time used: 0.599 (sec). Leaf size: 16

```
DSolve[(y[x]-3)*y'[x]==y'[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3 + e^{c_1(x+c_2)}$$

### 8.33 problem 13.5 (i)

Internal problem ID [13185]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.5 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' + 4y' = 9e^{-3x}$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)=9*exp(-3*x),y(x), singsol=all)
```

$$y(x) = -\frac{e^{-4x}c_1}{4} - 3e^{-3x} + c_2$$

#### ✓ Solution by Mathematica

Time used: 0.058 (sec). Leaf size: 26

```
DSolve[y''[x]+4*y'[x]==9*Exp[-3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -3e^{-3x} - \frac{1}{4}c_1e^{-4x} + c_2$$

### 8.34 problem 13.5 (j)

Internal problem ID [13186]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.5 (j).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible]`

$$y'' - y'(y' - 2) = 0$$

#### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)=diff(y(x),x)*(diff(y(x),x)-2),y(x), singsol=all)
```

$$y(x) = -\ln\left(\frac{e^{-2x}c_1}{2} - c_2\right)$$

#### ✓ Solution by Mathematica

Time used: 60.076 (sec). Leaf size: 23

```
DSolve[y''[x]==y'[x]*(y'[x]-2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - 2\operatorname{arctanh}(1 + 2e^{2(x+c_1)})$$

## 8.35 problem 13.6 (a)

Internal problem ID [13187]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.6 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' + 4y' = 18x^2$$

With initial conditions

$$[y(1) = 8, y'(1) = -3]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 14

```
dsolve([x*diff(y(x),x$2)+4*diff(y(x),x)=18*x^2,y(1) = 8, D(y)(1) = -3],y(x), singsol=all)
```

$$y(x) = x^3 + \frac{2}{x^3} + 5$$

### ✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 15

```
DSolve[{x*y'[x]+4*y'[x]==18*x^2,{y[1]==8,y'[1]==-3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^3 + \frac{2}{x^3} + 5$$

## 8.36 problem 13.6 (b)

Internal problem ID [13188]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.6 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' - 2y' = 0$$

With initial conditions

$$[y(-1) = 4, y'(-1) = 12]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve([x*diff(y(x),x$2)=2*diff(y(x),x),y(-1) = 4, D(y)(-1) = 12],y(x), singsol=all)
```

$$y(x) = 4x^3 + 8$$

### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 12

```
DSolve[{x*y'[x]==2*y'[x],{y[-1]==4,y'[-1]==12}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 4(x^3 + 2)$$



## 8.37 problem 13.6 (c)

Internal problem ID [13189]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.6 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y' = 0$$

With initial conditions

$$[y(0) = 8, y'(0) = 5]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 10

```
dsolve([diff(y(x),x$2)=diff(y(x),x),y(0) = 8, D(y)(0) = 5],y(x), singsol=all)
```

$$y(x) = 3 + 5e^x$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 12

```
DSolve[{y'[x]==y'[x],{y[0]==8,y'[0]==5}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 5e^x + 3$$

## 8.38 problem 13.6 (d)

Internal problem ID [13190]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.6 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' + 2y' = 8e^{2x}$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([diff(y(x),x$2)+2*diff(y(x),x)=8*exp(2*x),y(0) = 0, D(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = e^{-2x} + e^{2x} - 2$$

### ✓ Solution by Mathematica

Time used: 0.054 (sec). Leaf size: 17

```
DSolve[{y''[x]+2*y'[x]==8*Exp[2*x],{y[0]==0,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} + e^{2x} - 2$$

## 8.39 problem 13.6 (e)

Internal problem ID [13191]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.6 (e).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - y'' = 0$$

With initial conditions

$$[y(0) = 10, y'(0) = 5, y''(0) = 2]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve([diff(y(x),x$3)=diff(y(x),x$2),y(0) = 10, D(y)(0) = 5, (D@@2)(y)(0) = 2],y(x), singso
```

$$y(x) = 8 + 3x + 2e^x$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 15

```
DSolve[{y'''[x]==y''[x],{y[0]==10,y'[0]==5,y''[0]==2}},y[x],x,IncludeSingularSolutions -> Tr
```

$$y(x) \rightarrow 3x + 2e^x + 8$$

## 8.40 problem 13.6 (f)

Internal problem ID [13192]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.6 (f).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_y]]`

$$xy''' + 2y'' = 6x$$

With initial conditions

$$[y(1) = 2, y'(1) = 1, y''(1) = 4]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve([x*diff(y(x),x$3)+2*diff(y(x),x$2)=6*x,y(1) = 2, D(y)(1) = 1, (D@@2)(y)(1) = 4],y(x),
```

$$y(x) = \frac{x^3}{3} - 2 \ln(x) + 2x - \frac{1}{3}$$

### ✓ Solution by Mathematica

Time used: 0.045 (sec). Leaf size: 21

```
DSolve[{x*y'''[x]+2*y''[x]==6*x,{y[1]==2,y'[1]==1,y''[1]==4}},y[x],x,IncludeSingularSolution
```

$$y(x) \rightarrow \frac{1}{3}(x^3 + 6x - 6 \log(x) - 1)$$

## 8.41 problem 13.6 (g)

Internal problem ID [13193]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.6 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' + 2y' = 6$$

With initial conditions

$$[y(1) = 4, y'(1) = 5]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([x*diff(y(x),x$2)+2*diff(y(x),x)=6,y(1) = 4, D(y)(1) = 5],y(x), singsol=all)
```

$$y(x) = -\frac{2}{x} + 3x + 3$$

### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 15

```
DSolve[{x*y'[x]+2*y'[x]==6,{y[1]==4,y'[1]==5}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3x - \frac{2}{x} + 3$$

## 8.42 problem 13.6 (h)

Internal problem ID [13194]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.6 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1],`

$$2xy'y'' - y'^2 = -1$$

With initial conditions

$$[y(1) = 0, y'(1) = \sqrt{3}]$$

✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 19

```
dsolve([2*x*diff(y(x),x)*diff(y(x),x$2)=diff(y(x),x)^2-1,y(1) = 0, D(y)(1) = 3^(1/2)],y(x),
```

$$y(x) = \frac{(2x + 1)^{\frac{3}{2}}}{3} - \sqrt{3}$$

✓ Solution by Mathematica

Time used: 0.121 (sec). Leaf size: 26

```
DSolve[{2*x*y'[x]*y''[x]==y'[x]^2-1,{y[1]==0,y'[1]==Sqrt[3]}],y[x],x,IncludeSingularSolution
```

$$y(x) \rightarrow \frac{1}{3} \left( (2x + 1)^{3/2} - 3\sqrt{3} \right)$$

## 8.43 problem 13.7 (c)

Internal problem ID [13195]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.7 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible]`

$$3yy'' - 2y'^2 = 0$$

With initial conditions

$$[y(1) = 1, y'(1) = 9]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 11

```
dsolve([3*y(x)*diff(y(x),x$2)=2*diff(y(x),x)^2,y(1) = 1, D(y)(1) = 9],y(x), singsol=all)
```

$$y(x) = (3x - 2)^3$$

### ✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 12

```
DSolve[{3*y[x]*y'[x]==2*y'[x]^2,{y[1]==1,y'[1]==9}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow (3x - 2)^3$$

## 8.44 problem 13.7 (d)

Internal problem ID [13196]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.7 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible]`

$$yy'' + 2y'^2 - 3yy' = 0$$

With initial conditions

$$\left[ y(0) = 2, y'(0) = \frac{3}{4} \right]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 14

```
dsolve([y(x)*diff(y(x),x$2)+2*diff(y(x),x)^2=3*y(x)*diff(y(x),x),y(0) = 2, D(y)(0) = 3/4],y(x))
```

$$y(x) = (3e^{3x} + 5)^{\frac{1}{3}}$$

✓ Solution by Mathematica

Time used: 1.151 (sec). Leaf size: 18

```
DSolve[{y[x]*y'[x]+2*y'[x]^2==3*y[x]*y'[x],{y[0]==2,y'[0]==3/4}},y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \sqrt[3]{3e^{3x} + 5}$$



## 8.45 problem 13.7 (e)

Internal problem ID [13197]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.7 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear]`, [

$$y'' + y'e^{-y} = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 2]$$

### ✓ Solution by Maple

Time used: 0.109 (sec). Leaf size: 11

```
dsolve([diff(y(x),x$2)=-diff(y(x),x)*exp(-y(x)),y(0) = 0, D(y)(0) = 2],y(x), singsol=all)
```

$$y(x) = \ln(2e^x - 1)$$

### ✓ Solution by Mathematica

Time used: 5.75 (sec). Leaf size: 13

```
DSolve[{y'[x]==-y'[x]*Exp[-y[x]],{y[0]==0,y'[0]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(2e^x - 1)$$

## 8.46 problem 13.8 (i)

Internal problem ID [13198]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.8 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]`

$$y'' + 2xy'^2 = 0$$

With initial conditions

$$[y(0) = 3, y'(0) = 4]$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 12

```
dsolve([diff(y(x),x$2)=-2*x*diff(y(x),x)^2,y(0) = 3, D(y)(0) = 4],y(x), singsol=all)
```

$$y(x) = 2 \arctan(2x) + 3$$

### ✓ Solution by Mathematica

Time used: 0.981 (sec). Leaf size: 13

```
DSolve[{y'[x]==-2*x*y'[x]^2,{y[0]==3,y'[0]==4}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 \arctan(2x) + 3$$

## 8.47 problem 13.8 (ii)

Internal problem ID [13199]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.8 (ii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]`

$$y'' + 2xy'^2 = 0$$

With initial conditions

$$[y(0) = 3, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve([diff(y(x),x$2)=-2*x*diff(y(x),x)^2,y(0) = 3, D(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = 3$$

### ✓ Solution by Mathematica

Time used: 0.949 (sec). Leaf size: 6

```
DSolve[{y'[x]==-2*x*y'[x]^2,{y[0]==3,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3$$

## 8.48 problem 13.8 (iii)

Internal problem ID [13200]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.8 (iii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]`

$$y'' + 2xy'^2 = 0$$

With initial conditions

$$[y(1) = 0, y'(1) = 1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve([diff(y(x),x$2)=-2*x*diff(y(x),x)^2,y(1) = 0, D(y)(1) = 1],y(x), singsol=all)
```

$$y(x) = \frac{x-1}{x}$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y'[x]==-2*x*y'[x]^2,{y[1]==0,y'[1]==1}},y[x],x,IncludeSingularSolutions -> True]
```

{}

## 8.49 problem 13.8 (iv)

Internal problem ID [13201]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.8 (iv).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]`

$$y'' + 2xy'^2 = 0$$

With initial conditions

$$\left[ y(1) = -\frac{1}{4}, y'(1) = 5 \right]$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 29

```
dsolve([diff(y(x),x$2)=-2*x*diff(y(x),x)^2,y(1) = -1/4, D(y)(1) = 5],y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{5} \operatorname{arctanh}\left(\frac{\sqrt{5}x}{2}\right)}{2} + \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{\sqrt{5}}{2}\right)}{2} - \frac{1}{4}$$

### ✓ Solution by Mathematica

Time used: 0.185 (sec). Leaf size: 46

```
DSolve[{y'[x]==-2*x*y'[x]^2,{y[1]==-1/4,y'[1]==5}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4} \left( -2\sqrt{5} \operatorname{arctanh}\left(\frac{\sqrt{5}x}{2}\right) + 2\sqrt{5} \operatorname{arctanh}\left(\frac{\sqrt{5}}{2}\right) - 1 \right)$$

## 8.50 problem 13.9 (i)

Internal problem ID [13202]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.9 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear],`

$$y'' - 2yy' = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

### ✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 6

```
dsolve([diff(y(x),x$2)=2*y(x)*diff(y(x),x),y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = \tan(x)$$

### ✓ Solution by Mathematica

Time used: 10.217 (sec). Leaf size: 7

```
DSolve[{y'[x]==2*y[x]*y'[x],{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan(x)$$

## 8.51 problem 13.9 (ii)

Internal problem ID [13203]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.9 (ii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear],`

$$y'' - 2yy' = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve([diff(y(x),x$2)=2*y(x)*diff(y(x),x),y(0) = 1, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = -\frac{1}{x-1}$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y'[x]==2*y[x]*y'[x],{y[0]==1,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

{}

## 8.52 problem 13.9 (iii)

Internal problem ID [13204]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.9 (iii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear],`

$$y'' - 2yy' = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve([diff(y(x),x$2)=2*y(x)*diff(y(x),x),y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = 1$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y'[x]==2*y[x]*y'[x],{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

```
{}
```



## 8.53 problem 13.9 (iv)

Internal problem ID [13205]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 13. Higher order equations: Extending first order concepts. Additional exercises page 259

**Problem number:** 13.9 (iv).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear],`

$$y'' - 2yy' = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = -1]$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 8

```
dsolve([diff(y(x),x$2)=2*y(x)*diff(y(x),x),y(0) = 0, D(y)(0) = -1],y(x), singsol=all)
```

$$y(x) = -\tanh(x)$$

### ✓ Solution by Mathematica

Time used: 0.684 (sec). Leaf size: 9

```
DSolve[{y'[x]==2*y[x]*y'[x],{y[0]==0,y'[0]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\tanh(x)$$

## 9 Chapter 14. Higher order equations and the reduction of order method. Additional exercises

page 277

9.1	problem 14.1 (a)	329
9.2	problem 14.1 (b)	331
9.3	problem 14.1 (c)	332
9.4	problem 14.1 (d)	333
9.5	problem 14.1 (e)	334
9.6	problem 14.1 (f)	335
9.7	problem 14.1 (g)	336
9.8	problem 14.1 (h)	337
9.9	problem 14.1 (i)	338
9.10	problem 14.1 (j)	339
9.11	problem 14.2 (a)	340
9.12	problem 14.2 (b)	341
9.13	problem 14.2 (c)	342
9.14	problem 14.2 (d)	343
9.15	problem 14.2 (e)	344
9.16	problem 14.2 (f)	345
9.17	problem 14.2 (g)	346
9.18	problem 14.2 (h)	347
9.19	problem 14.2 (i)	348
9.20	problem 14.2 (j)	349
9.21	problem 14.2 (k)	350
9.22	problem 14.2 (L)	351
9.23	problem 14.2 (m)	352
9.24	problem 14.2 (n)	353
9.25	problem 14.3 (a)	354
9.26	problem 14.3 (b)	355
9.27	problem 14.3 (c)	356
9.28	problem 14.3 (d)	357
9.29	problem 14.3 (e)	358
9.30	problem 14.3 (f)	359
9.31	problem 14.5 (a)	360
9.32	problem 14.5 (b)	361
9.33	problem 14.5 (c)	362
9.34	problem 14.5 (d)	363

## 9.1 problem 14.1 (a)

Internal problem ID [13206]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.1 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + x^2 y' - 4y = x^3$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 189

```
dsolve(diff(y(x), x$2)+x^2*diff(y(x), x)-4*y(x)=x^3,y(x), singsol=all)
```

$$\begin{aligned} y(x) = & e^{-\frac{x^3}{3}} \operatorname{HeunT}\left(-4\sqrt[3]{3}, -3, 0, \frac{\sqrt[3]{3}x}{3}\right) c_2 \\ & + e^{-\frac{x^3}{3}} \operatorname{HeunT}\left(-4\sqrt[3]{3}, -3, 0, \frac{\sqrt[3]{3}x}{3}\right) \left( \int \frac{e^{\frac{x^3}{3}}}{\operatorname{HeunT}\left(-4\sqrt[3]{3}, -3, 0, \frac{\sqrt[3]{3}x}{3}\right)^2} dx \right) c_1 \\ & + \operatorname{HeunT}\left(-4\sqrt[3]{3}, \right. \\ & \left. -3, 0, \frac{\sqrt[3]{3}x}{3}\right) \left( \left( \int \operatorname{HeunT}\left(-4\sqrt[3]{3}, -3, 0, \frac{\sqrt[3]{3}x}{3}\right) x^3 dx \right) \left( \int \frac{e^{\frac{x^3}{3}}}{\operatorname{HeunT}\left(-4\sqrt[3]{3}, -3, 0, \frac{\sqrt[3]{3}x}{3}\right)^2} dx \right) \right. \\ & \left. - \left( \int \operatorname{HeunT}\left(-4\sqrt[3]{3}, -3, 0, \frac{\sqrt[3]{3}x}{3}\right) \left( \int \frac{e^{\frac{x^3}{3}}}{\operatorname{HeunT}\left(-4\sqrt[3]{3}, -3, 0, \frac{\sqrt[3]{3}x}{3}\right)^2} dx \right) x^3 dx \right) \right) e^{-\frac{x^3}{3}} \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.589 (sec). Leaf size: 194

`DSolve[y''[x]+x^2*y'[x]-4*y[x]==x^3,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow e^{-\frac{x^3}{3}} \text{HeunT}[4, -2, 0, 0, -1, x] \left( \int_1^x \frac{e^{\frac{K[2]^3}{3}} \text{HeunT}[4, 0, 0, 0, 1, K[2]] K[2]^3}{\text{HeunT}[4, -2, 0, 0, -1, K[2]] \text{HeunTPrime}[4, 0, 0, 0, 1, K[2]] + \text{HeunT}[4, 0, 0, 0, 1, K[2]] (\text{HeunT}[4, -2, 0, 0, -1, K[2]] + c_2)} dx \right) + \text{HeunT}[4, 0, 0, 0, 1, x] \left( \int_1^x \frac{\text{HeunT}[4, -2, 0, 0, -1, K[1]] \text{HeunTPrime}[4, 0, 0, 0, 1, K[1]] + \text{HeunT}[4, 0, 0, 0, 1, K[1]] (\text{HeunT}[4, -2, 0, 0, -1, K[1]] + c_1)}{\text{HeunT}[4, -2, 0, 0, -1, K[1]] \text{HeunTPrime}[4, 0, 0, 0, 1, K[1]] + \text{HeunT}[4, 0, 0, 0, 1, K[1]] (\text{HeunT}[4, -2, 0, 0, -1, K[1]] + c_1)} dx \right)$$

## 9.2 problem 14.1 (b)

Internal problem ID [13207]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.1 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + x^2 y' - 4y = 0$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 74

```
dsolve(diff(y(x), x$2)+x^2*diff(y(x), x)-4*y(x)=0, y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{x^3}{3}} \operatorname{HeunT}\left(-4 3^{\frac{2}{3}}, -3, 0, \frac{3^{\frac{2}{3}} x}{3}\right) + c_2 e^{-\frac{x^3}{3}} \operatorname{HeunT}\left(-4 3^{\frac{2}{3}}, -3, 0, \frac{3^{\frac{2}{3}} x}{3}\right) \left(\int \frac{e^{\frac{x^3}{3}}}{\operatorname{HeunT}\left(-4 3^{\frac{2}{3}}, -3, 0, \frac{3^{\frac{2}{3}} x}{3}\right)^2} dx\right)$$

### ✓ Solution by Mathematica

Time used: 0.061 (sec). Leaf size: 35

```
DSolve[y''[x]+x^2*y'[x]-4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 e^{-\frac{x^3}{3}} \operatorname{HeunT}[4, -2, 0, 0, -1, x] + c_1 \operatorname{HeunT}[4, 0, 0, 0, 1, x]$$

### 9.3 problem 14.1 (c)

Internal problem ID [13208]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.1 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + x^2 y' - 4y = 0$$

#### ✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 74

```
dsolve(diff(y(x), x$2)+x^2*diff(y(x), x)=4*y(x), y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{x^3}{3}} \operatorname{HeunT}\left(-4 3^{\frac{2}{3}}, -3, 0, \frac{3^{\frac{2}{3}} x}{3}\right) + c_2 e^{-\frac{x^3}{3}} \operatorname{HeunT}\left(-4 3^{\frac{2}{3}}, -3, 0, \frac{3^{\frac{2}{3}} x}{3}\right) \left(\int \frac{e^{\frac{x^3}{3}}}{\operatorname{HeunT}\left(-4 3^{\frac{2}{3}}, -3, 0, \frac{3^{\frac{2}{3}} x}{3}\right)^2} dx\right)$$

#### ✓ Solution by Mathematica

Time used: 0.06 (sec). Leaf size: 35

```
DSolve[y''[x]+x^2*y'[x]==4*y[x], y[x], x, IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 e^{-\frac{x^3}{3}} \operatorname{HeunT}[4, -2, 0, 0, -1, x] + c_1 \operatorname{HeunT}[4, 0, 0, 0, 1, x]$$

## 9.4 problem 14.1 (d)

Internal problem ID [13209]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.1 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [NONE]

$$y'' + y'x^2 + 4y - y^3 = 0$$

**X** Solution by Maple

```
dsolve(diff(y(x),x$2)+x^2*diff(y(x),x)+4*y(x)=y(x)^3,y(x), singsol=all)
```

No solution found

**X** Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y''[x]+x^2*y'[x]+4*y[x]==y[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

## 9.5 problem 14.1 (e)

Internal problem ID [13210]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.1 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [linear]

$$xy' + 3y = e^{2x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(x*diff(y(x),x)+3*y(x)=exp(2*x),y(x), singsol=all)
```

$$y(x) = \frac{\frac{(2x^2-2x+1)e^{2x}}{4} + c_1}{x^3}$$

### ✓ Solution by Mathematica

Time used: 0.069 (sec). Leaf size: 33

```
DSolve[x*y'[x]+3*y[x]==Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{2x}(2x^2 - 2x + 1) + 4c_1}{4x^3}$$



## 9.6 problem 14.1 (f)

Internal problem ID [13211]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.1 (f).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' + y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 37

```
dsolve(diff(y(x),x$3)+y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-x} + c_2 e^{\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + c_3 e^{\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

### ✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 56

```
DSolve[y'''[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} \left( c_3 e^{3x/2} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 e^{3x/2} \sin\left(\frac{\sqrt{3}x}{2}\right) + c_1 \right)$$

## 9.7 problem 14.1 (g)

Internal problem ID [13212]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.1 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1],`

$$(y + 1)y'' - y'^3 = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 31

```
dsolve((y(x)+1)*diff(y(x),x$2)=diff(y(x),x)^3,y(x), singsol=all)
```

$$y(x) = -1$$

$$y(x) = c_1$$

$$y(x) = e^{\text{LambertW}(-(c_1+c_2+x)e^{-c_1}e^{-1})+c_1+1} - 1$$

### ✓ Solution by Mathematica

Time used: 0.438 (sec). Leaf size: 93

```
DSolve[(y[x]+1)*y'[x]==y'[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{InverseFunction}[\#1 - (\#1 + 1) \log(\#1 + 1) + \#1(-c_1)\&][x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction}[\#1 - (\#1 + 1) \log(\#1 + 1) + \#1(-(-c_1))\&][x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction}[\#1 - (\#1 + 1) \log(\#1 + 1) + \#1(-c_1)\&][x + c_2]$$

## 9.8 problem 14.1 (h)

Internal problem ID [13213]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.1 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 2y' + 5y = 30e^{3x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)=2*diff(y(x),x)-5*y(x)+30*exp(3*x),y(x), singsol=all)
```

$$y(x) = e^x \sin(2x) c_2 + e^x \cos(2x) c_1 + \frac{15e^{3x}}{4}$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 35

```
DSolve[y''[x]==2*y'[x]-5*y[x]+30*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{15e^{3x}}{4} + c_2 e^x \cos(2x) + c_1 e^x \sin(2x)$$

## 9.9 problem 14.1 (i)

Internal problem ID [13214]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.1 (i).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 6y'' + 3y' - 83y = 25$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 94

```
dsolve(diff(y(x),x$4)+6*diff(y(x),x$2)+3*diff(y(x),x)-83*y(x)-25=0,y(x), singsol=all)
```

$$y(x) = -\frac{25}{83} + c_1 e^{\text{RootOf}(-Z^4+6_Z^2+3_Z-83,\text{index}=1)x} + c_2 e^{\text{RootOf}(-Z^4+6_Z^2+3_Z-83,\text{index}=2)x} \\ + c_3 e^{\text{RootOf}(-Z^4+6_Z^2+3_Z-83,\text{index}=3)x} + c_4 e^{\text{RootOf}(-Z^4+6_Z^2+3_Z-83,\text{index}=4)x}$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 117

```
DSolve[y''''[x]+6*y''[x]+3*y'[x]-83*y[x]-25==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3 \exp(x\text{Root}[\#1^4 + 6\#1^2 + 3\#1 - 83\&, 3]) \\ + c_4 \exp(x\text{Root}[\#1^4 + 6\#1^2 + 3\#1 - 83\&, 4]) \\ + c_2 \exp(x\text{Root}[\#1^4 + 6\#1^2 + 3\#1 - 83\&, 2]) \\ + c_1 \exp(x\text{Root}[\#1^4 + 6\#1^2 + 3\#1 - 83\&, 1]) - \frac{25}{83}$$

## 9.10 problem 14.1 (j)

Internal problem ID [13215]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.1 (j).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_3rd_order, _missing_x], [_3rd_order, _with_linear_symmetrie`

**X** Solution by Maple

```
dsolve(y(x)*diff(y(x),x$3)+6*diff(y(x),x$2)+3*diff(y(x),x)=y(x),y(x), singsol=all)
```

No solution found

**X** Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y[x]*y'''[x]+6*y''[x]+3*y'[x]==y[x],y[x],x,IncludeSingularSolutions -> True]
```

Not solved

## 9.11 problem 14.2 (a)

Internal problem ID [13216]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.2 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 5y' + 6y = 0$$

Given that one solution of the ode is

$$y_1 = e^{2x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)-5*diff(y(x),x)+6*y(x)=0,exp(2*x)],y(x), singsol=all)
```

$$y(x) = e^{2x}c_1 + c_2e^{3x}$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 20

```
DSolve[y''[x]-5*y'[x]+6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(c_2e^x + c_1)$$

## 9.12 problem 14.2 (b)

Internal problem ID [13217]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.2 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 10y' + 25y = 0$$

Given that one solution of the ode is

$$y_1 = e^{5x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve([diff(y(x),x$2)-10*diff(y(x),x)+25*y(x)=0,exp(5*x)],y(x), singsol=all)
```

$$y(x) = c_1 e^{5x} + c_2 e^{5x} x$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 18

```
DSolve[y''[x]-10*y'[x]+25*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{5x}(c_2 x + c_1)$$

## 9.13 problem 14.2 (c)

Internal problem ID [13218]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.2 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2y'' - 6xy' + 12y = 0$$

Given that one solution of the ode is

$$y_1 = x^3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([x^2*diff(y(x),x$2)-6*x*diff(y(x),x)+12*y(x)=0,x^3],y(x), singsol=all)
```

$$y(x) = c_1x^4 + c_2x^3$$

### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 16

```
DSolve[x^2*y''[x]-6*x*y'[x]+12*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^3(c_2x + c_1)$$



## 9.14 problem 14.2 (d)

Internal problem ID [13219]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.2 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$2x^2y'' - xy' + y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([2*x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,x],y(x), singsol=all)
```

$$y(x) = c_1\sqrt{x} + c_2x$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 18

```
DSolve[2*x^2*y''[x]-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1\sqrt{x} + c_2x$$

## 9.15 problem 14.2 (e)

Internal problem ID [13220]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.2 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$4x^2y'' + y = 0$$

Given that one solution of the ode is

$$y_1 = \sqrt{x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve([4*x^2*diff(y(x),x$2)+y(x)=0,sqrt(x)],y(x), singsol=all)
```

$$y(x) = c_1\sqrt{x} + c_2\sqrt{x} \ln(x)$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 24

```
DSolve[4*x^2*y''[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}\sqrt{x}(c_2 \log(x) + 2c_1)$$

## 9.16 problem 14.2 (f)

Internal problem ID [13221]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.2 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - \left(4 + \frac{2}{x}\right) y' + \left(4 + \frac{4}{x}\right) y = 0$$

Given that one solution of the ode is

$$y_1 = e^{2x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve([diff(y(x),x$2)-(4+2/x)*diff(y(x),x)+(4+4/x)*y(x)=0,exp(2*x)],y(x), singsol=all)
```

$$y(x) = e^{2x} c_1 + c_2 e^{2x} x^3$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 25

```
DSolve[y''[x]-(4+2/x)*y'[x]+(4+4/x)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} e^{2x} (c_2 x^3 + 3c_1)$$

## 9.17 problem 14.2 (g)

Internal problem ID [13222]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.2 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x + 1)y'' + xy' - y = 0$$

Given that one solution of the ode is

$$y_1 = e^{-x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([(x+1)*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=0,exp(-x)],y(x), singsol=all)
```

$$y(x) = c_1x + c_2e^{-x}$$

### ✓ Solution by Mathematica

Time used: 0.113 (sec). Leaf size: 31

```
DSolve[(x+1)*y'[x]+x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2ec_2x + c_1 \cosh(x) - c_1 \sinh(x)}{\sqrt{2e}}$$

## 9.18 problem 14.2 (h)

Internal problem ID [13223]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.2 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$y'' - \frac{y'}{x} - 4yx^2 = 0$$

Given that one solution of the ode is

$$y_1 = e^{-x^2}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)-1/x*diff(y(x),x)-4*x^2*y(x)=0,exp(-x^2)],y(x), singsol=all)
```

$$y(x) = c_1 \sinh(x^2) + c_2 \cosh(x^2)$$

### ✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 23

```
DSolve[y''[x]-1/x*y'[x]-4*x^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cosh(x^2) + ic_2 \sinh(x^2)$$

## 9.19 problem 14.2 (i)

Internal problem ID [13224]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.2 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y = 0$$

Given that one solution of the ode is

$$y_1 = \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([diff(y(x),x$2)+y(x)=0,sin(x)],y(x), singsol=all)
```

$$y(x) = c_1 \sin(x) + c_2 \cos(x)$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 16

```
DSolve[y''[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(x) + c_2 \sin(x)$$

## 9.20 problem 14.2 (j)

Internal problem ID [13225]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.2 (j).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$xy'' + (2x + 2)y' + 2y = 0$$

Given that one solution of the ode is

$$y_1 = \frac{1}{x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve([x*diff(y(x),x$2)+(2+2*x)*diff(y(x),x)+2*y(x)=0,1/x],y(x), singsol=all)
```

$$y(x) = \frac{c_1}{x} + \frac{c_2 e^{-2x}}{x}$$

### ✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 24

```
DSolve[x*y''[x]+(2+2*x)*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2c_1 e^{-2x} + c_2}{2x}$$

## 9.21 problem 14.2 (k)

Internal problem ID [13226]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.2 (k).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$\sin(x)^2 y'' - 2 \cos(x) \sin(x) y' + (1 + \cos(x)^2) y = 0$$

Given that one solution of the ode is

$$y_1 = \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([sin(x)^2*diff(y(x),x$2)-2*cos(x)*sin(x)*diff(y(x),x)+(1+cos(x)^2)*y(x)=0,sin(x)],y(x))
```

$$y(x) = c_1 \sin(x) + c_2 \sin(x) x$$

### ✓ Solution by Mathematica

Time used: 0.079 (sec). Leaf size: 15

```
DSolve[Sin[x]^2*y''[x]-2*Cos[x]*Sin[x]*y'[x]+(1+Cos[x]^2)*y[x]==0,y[x],x,IncludeSingularSolu
```

$$y(x) \rightarrow (c_2 x + c_1) \sin(x)$$



## 9.22 problem 14.2 (L)

Internal problem ID [13227]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.2 (L).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 2xy' + (x^2 + 2)y = 0$$

Given that one solution of the ode is

$$y_1 = \sin(x) x$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve([x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+(x^2+2)*y(x)=0,x*sin(x)],y(x), singsol=all)
```

$$y(x) = c_1 \sin(x) x + c_2 \cos(x) x$$

### ✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 33

```
DSolve[x^2*y''[x]-2*x*y'[x]+(x^2+2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-ix} x - \frac{1}{2} i c_2 e^{ix} x$$

## 9.23 problem 14.2 (m)

Internal problem ID [13228]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.2 (m).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$x^2 y'' + x y' + y = 0$$

Given that one solution of the ode is

$$y_1 = \frac{x}{2} - \frac{1}{2x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([x^2*diff(y(x),x$2)+x*diff(y(x),x)+y(x)=0,sinh(ln(x))],y(x), singsol=all)
```

$$y(x) = c_1 \sin(\ln(x)) + c_2 \cos(\ln(x))$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]+x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(\log(x)) + c_2 \sin(\log(x))$$

## 9.24 problem 14.2 (n)

Internal problem ID [13229]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.2 (n).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + xy' + \left(x^2 - \frac{1}{4}\right) y = 0$$

Given that one solution of the ode is

$$y_1 = \frac{\cos(x)}{\sqrt{x}}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve([x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-1/4)*y(x)=0,1/sqrt(x)*cos(x)],y(x), singsol=a
```

$$y(x) = \frac{c_1 \sin(x)}{\sqrt{x}} + \frac{c_2 \cos(x)}{\sqrt{x}}$$

### ✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 39

```
DSolve[x^2*y'[x]+x*y'[x]+(x^2-1/4)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{-ix}(2c_1 - ic_2 e^{2ix})}{2\sqrt{x}}$$

## 9.25 problem 14.3 (a)

Internal problem ID [13230]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.3 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 4y' + 3y = 9e^{2x}$$

Given that one solution of the ode is

$$y_1 = e^{3x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve([diff(y(x),x$2)-4*diff(y(x),x)+3*y(x)=9*exp(2*x),exp(3*x)],y(x), singsol=all)
```

$$y = e^x c_2 + e^{3x} c_1 - 9e^{2x}$$

### ✓ Solution by Mathematica

Time used: 0.069 (sec). Leaf size: 25

```
DSolve[y''[x]-4*y'[x]+3*y[x]==9*Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(-9e^x + c_2e^{2x} + c_1)$$

## 9.26 problem 14.3 (b)

Internal problem ID [13231]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.3 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 6y' + 8y = e^{4x}$$

Given that one solution of the ode is

$$y_1 = e^{2x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve([diff(y(x),x$2)-6*diff(y(x),x)+8*y(x)=exp(4*x),exp(2*x)],y(x), singsol=all)
```

$$y = \left( \frac{e^{2x}(2x + 2c_1 - 1)}{4} + c_2 \right) e^{2x}$$

### ✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 31

```
DSolve[y''[x]-6*y'[x]+8*y[x]==Exp[4*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{2x} + e^{4x} \left( \frac{x}{2} - \frac{1}{4} + c_2 \right)$$

## 9.27 problem 14.3 (c)

Internal problem ID [13232]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.3 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$x^2y'' + y'x - y = \sqrt{x}$$

Given that one solution of the ode is

$$y_1 = x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve([x^2*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=sqrt(x),x],y(x), singsol=all)
```

$$y = c_2x - \frac{4\sqrt{x}}{3} + \frac{c_1}{x}$$

### ✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 25

```
DSolve[x^2*y''[x]+x*y'[x]-y[x]==Sqrt[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{4\sqrt{x}}{3} + \frac{c_1}{x} + c_2x$$

## 9.28 problem 14.3 (d)

Internal problem ID [13233]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.3 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 20y = 27x^5$$

Given that one solution of the ode is

$$y_1 = x^5$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve([x^2*diff(y(x),x$2)-20*y(x)=27*x^5,x^5],y(x), singsol=all)
```

$$y = \frac{c_2}{x^4} + c_1 x^5 + x^5 \left( -\frac{1}{3} + 3 \ln(x) \right)$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 29

```
DSolve[x^2*y'[x]-20*y[x]==27*x^5,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3x^5 \log(x) + \left( -\frac{1}{3} + c_2 \right) x^5 + \frac{c_1}{x^4}$$

## 9.29 problem 14.3 (e)

Internal problem ID [13234]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.3 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$xy'' + (2 + 2x)y' + 2y = 8e^{2x}$$

Given that one solution of the ode is

$$y_1 = \frac{1}{x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve([x*diff(y(x),x$2)+(2+2*x)*diff(y(x),x)+2*y(x)=8*exp(2*x),1/x],y(x), singsol=all)
```

$$y = \frac{e^{-2x}c_2}{x} + \frac{c_1}{x} + \frac{e^{-2x}e^{4x}}{x}$$

### ✓ Solution by Mathematica

Time used: 0.066 (sec). Leaf size: 31

```
DSolve[x*y''[x]+(2+2*x)*y'[x]+2*y[x]==8*Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2e^{2x} + 2c_1e^{-2x} + c_2}{2x}$$



### 9.30 problem 14.3 (f)

Internal problem ID [13235]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.3 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x + 1)y'' + y'x - y = (x + 1)^2$$

Given that one solution of the ode is

$$y_1 = e^{-x}$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve([(x+1)*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=(1+x)^2,exp(-x)],y(x), singsol=all)
```

$$y = c_2x + e^{-x}c_1 + x^2 + 1$$

#### ✓ Solution by Mathematica

Time used: 0.156 (sec). Leaf size: 41

```
DSolve[(x+1)*y''[x]+x*y'[x]-y[x]==(1+x)^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2 + \left(-1 + \sqrt{2}ec_2\right)x + \frac{c_1e^{-x-\frac{1}{2}}}{\sqrt{2}} + 1$$

## 9.31 problem 14.5 (a)

Internal problem ID [13236]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.5 (a).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 9y'' + 27y' - 27y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve([diff(y(x),x$3)-9*diff(y(x),x$2)+27*diff(y(x),x)-27*y(x)=0,exp(3*x)],y(x), singsol=all
```

$$y = e^{3x}c_1 + x e^{3x}c_2 + x^2 e^{3x}c_3$$

### ✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 23

```
DSolve[y'''[x]-9*y''[x]+27*y'[x]-27*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{3x}(x(c_3x + c_2) + c_1)$$

## 9.32 problem 14.5 (b)

Internal problem ID [13237]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.5 (b).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - 9y'' + 27y' - 27y = e^{3x} \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve([diff(y(x),x$3)-9*diff(y(x),x$2)+27*diff(y(x),x)-27*y(x)=exp(3*x)*sin(x),exp(3*x)],y(x))
```

$$y = x^2 e^{3x} c_3 + x e^{3x} c_2 + e^{3x} \cos(x) + e^{3x} c_1$$

### ✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 25

```
DSolve[y'''[x]-9*y''[x]+27*y'[x]-27*y[x]==Exp[3*x]*Sin[x],y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow e^{3x}(\cos(x) + x(c_3 x + c_2) + c_1)$$

### 9.33 problem 14.5 (c)

Internal problem ID [13238]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.5 (c).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 8y'''' + 24y'' - 32y' + 16y = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
dsolve([diff(y(x),x$4)-8*diff(y(x),x$3)+24*diff(y(x),x$2)-32*diff(y(x),x)+16*y(x)=0,exp(2*x)
```

$$y = e^{2x}c_1 + xe^{2x}c_2 + x^2e^{2x}c_3 + x^3e^{2x}c_4$$

#### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 28

```
DSolve[y''''[x]-8*y''''[x]+24*y''[x]-32*y'[x]+16*y[x]==0,y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow e^{2x}(x(x(c_4x + c_3) + c_2) + c_1)$$

### 9.34 problem 14.5 (d)

Internal problem ID [13239]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 14. Higher order equations and the reduction of order method. Additional exercises page 277

**Problem number:** 14.5 (d).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$x^3y''' - 4y'' + 10y' - 12y = 0$$

**X** Solution by Maple

```
dsolve([x^3*diff(y(x),x$3)-4*diff(y(x),x$2)+10*diff(y(x),x)-12*y(x)=0,x^2],y(x), singsol=all
```

No solution found

**X** Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[x^3*y'''[x]-4*y''[x]+10*y'[x]-12*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

## 10 Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

10.1 problem 15.2 (a) . . . . .	365
10.2 problem 15.2 (b) . . . . .	366
10.3 problem 15.2 (c) . . . . .	367
10.4 problem 15.2 (d) . . . . .	368
10.5 problem 15.2 (e) . . . . .	369
10.6 problem 15.2 (f) . . . . .	370
10.7 problem 15.2 (g) . . . . .	371
10.8 problem 15.2 (h) . . . . .	372
10.9 problem 15.2 (i) . . . . .	373
10.10 problem 15.3 . . . . .	374
10.11 problem 15.4 . . . . .	375
10.12 problem 15.5 (a) . . . . .	376
10.13 problem 15.5 (c) . . . . .	377
10.14 problem 15.6 (a) . . . . .	378
10.15 problem 15.6 (b) . . . . .	379
10.16 problem 15.6 (c) . . . . .	380
10.17 problem 15.6 (d) . . . . .	381
10.18 problem 15.7 (a) . . . . .	382
10.19 problem 15.7 (b) . . . . .	383

## 10.1 problem 15.2 (a)

Internal problem ID [13240]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.2 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 4y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = 6]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)+4*y(x)=0,y(0) = 2, D(y)(0) = 6],y(x), singsol=all)
```

$$y = 3 \sin(2x) + 2 \cos(2x)$$

### ✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 18

```
DSolve[{y'[x]+4*y[x]==0,{y[0]==2,y'[0]==6}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3 \sin(2x) + 2 \cos(2x)$$

## 10.2 problem 15.2 (b)

Internal problem ID [13241]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.2 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 12]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)-4*y(x)=0,y(0) = 0, D(y)(0) = 12],y(x), singsol=all)
```

$$y = 3e^{2x} - 3e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 19

```
DSolve[{y'[x]-4*y[x]==0,{y[0]==0,y'[0]==12}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3e^{-2x}(e^{4x} - 1)$$



### 10.3 problem 15.2 (c)

Internal problem ID [13242]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.2 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y' - 6y = 0$$

With initial conditions

$$[y(0) = 8, y'(0) = -9]$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)+diff(y(x),x)-6*y(x)=0,y(0) = 8, D(y)(0) = -9],y(x), singsol=all)
```

$$y = (3e^{5x} + 5)e^{-3x}$$

#### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 20

```
DSolve[{y''[x]+y'[x]-6*y[x]==0,{y[0]==8,y'[0]==-9}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x}(3e^{5x} + 5)$$

## 10.4 problem 15.2 (d)

Internal problem ID [13243]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.2 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y' + 4y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 6]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 14

```
dsolve([diff(y(x),x$2)-4*diff(y(x),x)+4*y(x)=0,y(0) = 1, D(y)(0) = 6],y(x), singsol=all)
```

$$y = e^{2x}(1 + 4x)$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 16

```
DSolve[{y'[x]-4*y'[x]+4*y[x]==0,{y[0]==1,y'[0]==6}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow e^{2x}(4x + 1)$$

## 10.5 problem 15.2 (e)

Internal problem ID [13244]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.2 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$x^2y'' - 4y'x + 6y = 0$$

With initial conditions

$$[y(1) = 0, y'(1) = 4]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+6*y(x)=0,y(1) = 0, D(y)(1) = 4],y(x), singsol=al
```

$$y = 4x^3 - 4x^2$$

### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 13

```
DSolve[{x^2*y'[x]-4*x*y'[x]+6*y[x]==0,{y[1]==0,y'[1]==4}},y[x],x,IncludeSingularSolutions -
```

$$y(x) \rightarrow 4(x - 1)x^2$$

## 10.6 problem 15.2 (f)

Internal problem ID [13245]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.2 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$4x^2y'' + 4y'x - y = 0$$

With initial conditions

$$[y(1) = 8, y'(1) = 1]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve([4*x^2*diff(y(x),x$2)+4*x*diff(y(x),x)-y(x)=0,y(1) = 8, D(y)(1) = 1],y(x), singsol=all)
```

$$y = \frac{5x + 3}{\sqrt{x}}$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 16

```
DSolve[{4*x^2*y''[x]+4*x*y'[x]-y[x]==0,{y[1]==8,y'[1]==1}},y[x],x,IncludeSingularSolutions->
```

$$y(x) \rightarrow \frac{5x + 3}{\sqrt{x}}$$

## 10.7 problem 15.2 (g)

Internal problem ID [13246]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.2 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - y' x + y = 0$$

With initial conditions

$$[y(1) = 5, y'(1) = 3]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve([x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,y(1) = 5, D(y)(1) = 3],y(x), singsol=all)
```

$$y = x(5 - 2 \ln(x))$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 13

```
DSolve[{x^2*y'[x]-x*y'[x]+y[x]==0,{y[1]==5,y'[1]==3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(5 - 2 \log(x))$$

## 10.8 problem 15.2 (h)

Internal problem ID [13247]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.2 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$xy'' - y' + 4x^3y = 0$$

With initial conditions

$$[y(\sqrt{\pi}) = 3, y'(\sqrt{\pi}) = 4]$$

### ✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 20

```
dsolve([x*diff(y(x),x$2)-diff(y(x),x)+4*x^3*y(x)=0,y(Pi^(1/2)) = 3, D(y)(Pi^(1/2)) = 4],y(x)
```

$$y = \frac{-3 \cos(x^2) \sqrt{\pi} - 2 \sin(x^2)}{\sqrt{\pi}}$$

### ✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 23

```
DSolve[{x*y''[x]-y'[x]+4*x^3*y[x]==0,{y[Sqrt[Pi]]==3,y'[Sqrt[Pi]]==4}},y[x],x,IncludeSingular
```

$$y(x) \rightarrow -\frac{2 \sin(x^2)}{\sqrt{\pi}} - 3 \cos(x^2)$$

## 10.9 problem 15.2 (i)

Internal problem ID [13248]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.2 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`, `[_2nd_order, _linear,`

$$(x + 1)^2 y'' - 2(x + 1) y' + 2y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 4]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve([(x+1)^2*diff(y(x),x$2)-2*(x+1)*diff(y(x),x)+2*y(x)=0,y(0) = 0, D(y)(0) = 4],y(x), si
```

$$y = 4x^2 + 4x$$

### ✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 11

```
DSolve[{(x+1)^2*y''[x]-2*(x+1)*y'[x]+2*y[x]==0,{y[0]==0,y'[0]==4}},y[x],x,IncludeSingularSol
```

$$y(x) \rightarrow 4x(x + 1)$$

## 10.10 problem 15.3

Internal problem ID [13249]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.3.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$x^2y'' - 4y'x + 6y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = -4]$$

**X** Solution by Maple

```
dsolve([x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+6*y(x)=0,y(0) = 0, D(y)(0) = -4],y(x), singsol=a
```

No solution found

**X** Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{x^2*y'[x]-4*x*y'[x]+6*y[x]==0,{y[0]==0,y'[0]==-4}},y[x],x,IncludeSingularSolutions
```

```
{}
```



## 10.11 problem 15.4

Internal problem ID [13250]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.4.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$xy'' - y' + 4x^3y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 4]$$

**X** Solution by Maple

```
dsolve([x*diff(y(x),x$2)-diff(y(x),x)+4*x^3*y(x)=0,y(0) = 1, D(y)(0) = 4],y(x), singsol=all)
```

No solution found

**X** Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{x*y'[x]-y'[x]+4*x^3*y[x]==0,{y[0]==1,y'[0]==4}},y[x],x,IncludeSingularSolutions ->
```

```
{}
```

## 10.12 problem 15.5 (a)

Internal problem ID [13251]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.5 (a).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' + 4y' = 0$$

With initial conditions

$$[y(0) = 3, y'(0) = 8, y''(0) = 4]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve([diff(y(x),x$3)+4*diff(y(x),x)=0,y(0) = 3, D(y)(0) = 8, (D@@2)(y)(0) = 4],y(x), sings
```

$$y = 4 + 4 \sin(2x) - \cos(2x)$$

### ✓ Solution by Mathematica

Time used: 0.111 (sec). Leaf size: 19

```
DSolve[{y'''[x]+4*y'[x]==0,{y[0]==3,y'[0]==8,y''[0]==4}},y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow 4 \sin(2x) - \cos(2x) + 4$$

## 10.13 problem 15.5 (c)

Internal problem ID [13252]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.5 (c).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 4, y''(0) = 0, y'''(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$4)-y(x)=0,y(0) = 0, D(y)(0) = 4, (D@@2)(y)(0) = 0, (D@@3)(y)(0) = 0],y(x)
```

$$y = e^x - e^{-x} + 2 \sin(x)$$

### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 20

```
DSolve[{y''''[x]-y[x]==0,{y[0]==0,y'[0]==4,y''[0]==0,y'''[0]==0}},y[x],x,IncludeSingularSolu
```

$$y(x) \rightarrow -e^{-x} + e^x + 2 \sin(x)$$

## 10.14 problem 15.6 (a)

Internal problem ID [13253]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.6 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)-4*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y = \frac{e^{2x}}{2} + \frac{e^{-2x}}{2}$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 21

```
DSolve[{y''[x]-4*y[x]==0,{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}e^{-2x}(e^{4x} + 1)$$

## 10.15 problem 15.6 (b)

Internal problem ID [13254]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.6 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2y' - 3y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve([diff(y(x),x$2)+2*diff(y(x),x)-3*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y = \frac{(e^{4x} - 1)e^{-3x}}{4}$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 21

```
DSolve[{y'[x]+2*y'[x]-3*y[x]==0,{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{1}{4}e^{-3x}(e^{4x} - 1)$$

## 10.16 problem 15.6 (c)

Internal problem ID [13255]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.6 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 10y' + 9y = 0$$

With initial conditions

$$[y(0) = 8, y'(0) = -24]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve([diff(y(x),x$2)-10*diff(y(x),x)+9*y(x)=0,y(0) = 8, D(y)(0) = -24],y(x), singsol=all)
```

$$y = 12e^x - 4e^{9x}$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 17

```
DSolve[{y'[x]-10*y'[x]+9*y[x]==0,{y[0]==8,y'[0]==-24}},y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow -4e^x(e^{8x} - 3)$$

## 10.17 problem 15.6 (d)

Internal problem ID [13256]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.6 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 5y' = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 5

```
dsolve([diff(y(x),x$2)+5*diff(y(x),x)=0,y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y = 1$$

### ✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 6

```
DSolve[{y''[x]+5*y'[x]==0,{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 1$$

## 10.18 problem 15.7 (a)

Internal problem ID [13257]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.7 (a).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 9y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$3)-9*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = c_1 + c_2 e^{-3x} + c_3 e^{3x}$$

### ✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 30

```
DSolve[y'''[x]-9*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}c_1 e^{3x} - \frac{1}{3}c_2 e^{-3x} + c_3$$



## 10.19 problem 15.7 (b)

Internal problem ID [13258]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 15. General solutions to Homogeneous linear differential equations. Additional exercises page 294

**Problem number:** 15.7 (b).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 10y'' + 9y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$4)-10*diff(y(x),x$2)+9*y(x)=0,y(x), singsol=all)
```

$$y = c_1e^x + c_2e^{-3x} + c_3e^{-x} + c_4e^{3x}$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 36

```
DSolve[y''''[x]-10*y''[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1e^{-3x} + c_2e^{-x} + c_3e^x + c_4e^{3x}$$

# 11 Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

11.1 problem 17.1 (a)	386
11.2 problem 17.1 (b)	387
11.3 problem 17.1 (c)	388
11.4 problem 17.1 (d)	389
11.5 problem 17.1 (e)	390
11.6 problem 17.1 (f)	391
11.7 problem 17.2 (a)	392
11.8 problem 17.2 (b)	393
11.9 problem 17.2 (c)	394
11.10 problem 17.2 (d)	395
11.11 problem 17.2 (e)	396
11.12 problem 17.2 (f)	397
11.13 problem 17.3 (a)	398
11.14 problem 17.3 (b)	399
11.15 problem 17.3 (c)	400
11.16 problem 17.3 (d)	401
11.17 problem 17.3 (e)	402
11.18 problem 17.3 (f)	403
11.19 problem 17.4 (a)	404
11.20 problem 17.4 (b)	405
11.21 problem 17.4 (c)	406
11.22 problem 17.4 (d)	407
11.23 problem 17.4 (e)	408
11.24 problem 17.4 (f)	409
11.25 problem 17.5 (a)	410
11.26 problem 17.5 (b)	411
11.27 problem 17.5 (c)	412
11.28 problem 17.5 (d)	413
11.29 problem 17.5 (e)	414
11.30 problem 17.5 (f)	415
11.31 problem 17.6 (a)	416
11.32 problem 17.6 (b)	417
11.33 problem 17.6 (c)	418
11.34 problem 17.6 (d)	419

11.35problem 17.6 (e)	420
11.36problem 17.6 (f)	421
11.37problem 17.7 (a)	422
11.38problem 17.7 (b)	423

## 11.1 problem 17.1 (a)

Internal problem ID [13259]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.1 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 7y' + 10y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-7*diff(y(x),x)+10*y(x)=0,y(x), singsol=all)
```

$$y = e^{2x}c_1 + c_2e^{5x}$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 22

```
DSolve[y''[x]-7*y'[x]+10*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(c_2e^{3x} + c_1)$$

## 11.2 problem 17.1 (b)

Internal problem ID [13260]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.1 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2y' - 24y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)-24*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-6x} + c_2 e^{4x}$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 22

```
DSolve[y''[x]+2*y'[x]-24*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-6x} + c_2 e^{4x}$$

### 11.3 problem 17.1 (c)

Internal problem ID [13261]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.1 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 25y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-25*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-5x} + c_2 e^{5x}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 22

```
DSolve[y''[x]-25*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{5x} + c_2 e^{-5x}$$

## 11.4 problem 17.1 (d)

Internal problem ID [13262]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.1 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 3y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x$2)+3*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = c_1 + c_2 e^{-3x}$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 19

```
DSolve[y''[x]+3*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \frac{1}{3}c_1 e^{-3x}$$

## 11.5 problem 17.1 (e)

Internal problem ID [13263]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.1 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4y'' - y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(4*diff(y(x),x$2)-y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-\frac{x}{2}} + c_2 e^{\frac{x}{2}}$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 22

```
DSolve[4*y''[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/2}(c_1 e^x + c_2)$$



## 11.6 problem 17.1 (f)

Internal problem ID [13264]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.1 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$3y'' + 7y' - 6y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(3*diff(y(x),x$2)+7*diff(y(x),x)-6*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-3x} + c_2 e^{\frac{2x}{3}}$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 24

```
DSolve[3*y''[x]+7*y'[x]-6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{2x/3} + c_2 e^{-3x}$$

## 11.7 problem 17.2 (a)

Internal problem ID [13265]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.2 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 8y' + 15y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)-8*diff(y(x),x)+15*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y = -\frac{3e^{5x}}{2} + \frac{5e^{3x}}{2}$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 23

```
DSolve[{y'[x]-8*y'[x]+15*y[x]==0,{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}e^{3x}(5 - 3e^{2x})$$

## 11.8 problem 17.2 (b)

Internal problem ID [13266]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.2 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 8y' + 15y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)-8*diff(y(x),x)+15*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y = \frac{e^{5x}}{2} - \frac{e^{3x}}{2}$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 21

```
DSolve[{y'[x]-8*y'[x]+15*y[x]==0,{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}e^{3x}(e^{2x} - 1)$$

## 11.9 problem 17.2 (c)

Internal problem ID [13267]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.2 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 8y' + 15y = 0$$

With initial conditions

$$[y(0) = 5, y'(0) = 19]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)-8*diff(y(x),x)+15*y(x)=0,y(0) = 5, D(y)(0) = 19],y(x), singsol=all)
```

$$y = 2e^{5x} + 3e^{3x}$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 20

```
DSolve[{y'[x]-8*y'[x]+15*y[x]==0,{y[0]==5,y'[0]==19}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{3x}(2e^{2x} + 3)$$

## 11.10 problem 17.2 (d)

Internal problem ID [13268]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.2 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 9y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)-9*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y = \frac{e^{-3x}}{2} + \frac{e^{3x}}{2}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 21

```
DSolve[{y''[x]-9*y[x]==0,{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}e^{-3x}(e^{6x} + 1)$$

## 11.11 problem 17.2 (e)

Internal problem ID [13269]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.2 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 9y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)-9*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y = -\frac{e^{-3x}}{6} + \frac{e^{3x}}{6}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 21

```
DSolve[{y'[x]-9*y[x]==0,{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{6}e^{-3x}(e^{6x} - 1)$$

## 11.12 problem 17.2 (f)

Internal problem ID [13270]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.2 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 9y = 0$$

With initial conditions

$$[y(0) = 3, y'(0) = -3]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([diff(y(x),x$2)-9*y(x)=0,y(0) = 3, D(y)(0) = -3],y(x), singsol=all)
```

$$y = 2e^{-3x} + e^{3x}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 18

```
DSolve[{y'[x]-9*y[x]==0,{y[0]==3,y'[0]==-3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x}(e^{6x} + 2)$$

## 11.13 problem 17.3 (a)

Internal problem ID [13271]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.3 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 10y' + 25y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)-10*diff(y(x),x)+25*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{5x} + c_2 e^{5x} x$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 18

```
DSolve[y''[x]-10*y'[x]+25*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{5x}(c_2 x + c_1)$$



## 11.14 problem 17.3 (b)

Internal problem ID [13272]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.3 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2y' + y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y = e^{-x}c_1 + c_2e^{-x}x$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 18

```
DSolve[y''[x]+2*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(c_2x + c_1)$$

## 11.15 problem 17.3 (c)

Internal problem ID [13273]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.3 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4y'' - 4y' + y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve(4*difff(y(x),x$2)-4*difff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{\frac{x}{2}} + c_2 e^{\frac{x}{2}} x$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 20

```
DSolve[4*y''[x]-4*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x/2}(c_2 x + c_1)$$

## 11.16 problem 17.3 (d)

Internal problem ID [13274]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.3 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$25y'' - 10y' + y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(25*diff(y(x),x$2)-10*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{\frac{x}{5}} + c_2 e^{\frac{x}{5}} x$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 20

```
DSolve[25*y'[x]-10*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x/5}(c_2 x + c_1)$$

## 11.17 problem 17.3 (e)

Internal problem ID [13275]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.3 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$16y'' - 24y' + 9y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve(16*diff(y(x),x$2)-24*diff(y(x),x)+9*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{\frac{3x}{4}} + c_2 e^{\frac{3x}{4}} x$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 20

```
DSolve[16*y''[x]-24*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{3x/4}(c_2 x + c_1)$$

## 11.18 problem 17.3 (f)

Internal problem ID [13276]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.3 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$9y'' + 12y' + 4y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(9*diff(y(x),x$2)+12*diff(y(x),x)+4*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-\frac{2x}{3}} + c_2 e^{-\frac{2x}{3}} x$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 20

```
DSolve[9*y''[x]+12*y'[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x/3}(c_2 x + c_1)$$

## 11.19 problem 17.4 (a)

Internal problem ID [13277]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.4 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 8y' + 16y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([diff(y(x),x$2)-8*diff(y(x),x)+16*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y = e^{4x}(1 - 4x)$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 16

```
DSolve[{y'[x]-8*y'[x]+16*y[x]==0,{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{4x}(1 - 4x)$$

## 11.20 problem 17.4 (b)

Internal problem ID [13278]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.4 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 8y' + 16y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 10

```
dsolve([diff(y(x),x$2)-8*diff(y(x),x)+16*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y = e^{4x}x$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 12

```
DSolve[{y'[x]-8*y'[x]+16*y[x]==0,{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{4x}x$$

## 11.21 problem 17.4 (c)

Internal problem ID [13279]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.4 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 8y' + 16y = 0$$

With initial conditions

$$[y(0) = 3, y'(0) = 14]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([diff(y(x),x$2)-8*diff(y(x),x)+16*y(x)=0,y(0) = 3, D(y)(0) = 14],y(x), singsol=all)
```

$$y = e^{4x}(3 + 2x)$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 16

```
DSolve[{y''[x]-8*y'[x]+16*y[x]==0,{y[0]==3,y'[0]==14}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{4x}(2x + 3)$$



## 11.22 problem 17.4 (d)

Internal problem ID [13280]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.4 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4y'' + 4y' + y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([4*dif(y(x),x$2)+4*dif(y(x),x)+y(x)=0,y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y = \frac{e^{-\frac{x}{2}}(2+x)}{2}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 19

```
DSolve[{4*y'[x]+4*y'[x]+y[x]==0,{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{1}{2}e^{-x/2}(x+2)$$

## 11.23 problem 17.4 (e)

Internal problem ID [13281]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.4 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4y'' + 4y' + y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve([4*dif(y(x),x$2)+4*dif(y(x),x)+y(x)=0,y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y = e^{-\frac{x}{2}}$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 14

```
DSolve[{4*y'[x]+4*y'[x]+y[x]==0,{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow e^{-x/2}$$

## 11.24 problem 17.4 (f)

Internal problem ID [13282]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.4 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4y'' + 4y' + y = 0$$

With initial conditions

$$[y(0) = 6, y'(0) = -5]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve([4*dif(y(x),x$2)+4*dif(y(x),x)+y(x)=0,y(0) = 6, D(y)(0) = -5],y(x), singsol=all)
```

$$y = -2e^{-\frac{x}{2}}(-3 + x)$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 17

```
DSolve[{4*y'[x]+4*y'[x]+y[x]==0,{y[0]==6,y'[0]==-5}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2e^{-x/2}(x - 3)$$

## 11.25 problem 17.5 (a)

Internal problem ID [13283]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.5 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 25y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+25*y(x)=0,y(x), singsol=all)
```

$$y = c_1 \sin(5x) + c_2 \cos(5x)$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 20

```
DSolve[y''[x]+25*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(5x) + c_2 \sin(5x)$$

## 11.26 problem 17.5 (b)

Internal problem ID [13284]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.5 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 2y' + 5y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)+5*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-x} \sin(2x) + c_2 e^{-x} \cos(2x)$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 26

```
DSolve[y''[x]+2*y'[x]+5*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(c_2 \cos(2x) + c_1 \sin(2x))$$

## 11.27 problem 17.5 (c)

Internal problem ID [13285]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.5 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' + 5y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+5*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^x \sin(2x) + c_2 e^x \cos(2x)$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 24

```
DSolve[y''[x]-2*y'[x]+5*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(c_2 \cos(2x) + c_1 \sin(2x))$$

## 11.28 problem 17.5 (d)

Internal problem ID [13286]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.5 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y' + 29y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+29*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{2x} \sin(5x) + c_2 e^{2x} \cos(5x)$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 26

```
DSolve[y''[x]-4*y'[x]+29*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(c_2 \cos(5x) + c_1 \sin(5x))$$

## 11.29 problem 17.5 (e)

Internal problem ID [13287]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.5 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$9y'' + 18y' + 10y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(9*diff(y(x),x$2)+18*diff(y(x),x)+10*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-x} \sin\left(\frac{x}{3}\right) + c_2 e^{-x} \cos\left(\frac{x}{3}\right)$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 30

```
DSolve[9*y''[x]+18*y'[x]+10*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} \left( c_2 \cos\left(\frac{x}{3}\right) + c_1 \sin\left(\frac{x}{3}\right) \right)$$



## 11.30 problem 17.5 (f)

Internal problem ID [13288]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.5 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4y'' + y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(4*diff(y(x),x$2)+y(x)=0,y(x), singsol=all)
```

$$y = c_1 \sin\left(\frac{x}{2}\right) + c_2 \cos\left(\frac{x}{2}\right)$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 24

```
DSolve[4*y''[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos\left(\frac{x}{2}\right) + c_2 \sin\left(\frac{x}{2}\right)$$

## 11.31 problem 17.6 (a)

Internal problem ID [13289]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.6 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 16y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve([diff(y(x),x$2)+16*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y = \cos(4x)$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 9

```
DSolve[{y'[x]+16*y[x]==0,{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \cos(4x)$$

## 11.32 problem 17.6 (b)

Internal problem ID [13290]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.6 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 16y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve([diff(y(x),x$2)+16*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y = \frac{\sin(4x)}{4}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 13

```
DSolve[{y'[x]+16*y[x]==0,{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4} \sin(4x)$$

### 11.33 problem 17.6 (c)

Internal problem ID [13291]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.6 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 16y = 0$$

With initial conditions

$$[y(0) = 4, y'(0) = 12]$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)+16*y(x)=0,y(0) = 4, D(y)(0) = 12],y(x), singsol=all)
```

$$y = 3 \sin(4x) + 4 \cos(4x)$$

#### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 18

```
DSolve[{y'[x]+16*y[x]==0,{y[0]==4,y'[0]==12}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3 \sin(4x) + 4 \cos(4x)$$

## 11.34 problem 17.6 (d)

Internal problem ID [13292]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.6 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y' + 13y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
dsolve([diff(y(x),x$2)-4*diff(y(x),x)+13*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y = -\frac{e^{2x}(2 \sin(3x) - 3 \cos(3x))}{3}$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 27

```
DSolve[{y'[x]-4*y'[x]+13*y[x]==0,{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}e^{2x}(3 \cos(3x) - 2 \sin(3x))$$

## 11.35 problem 17.6 (e)

Internal problem ID [13293]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.6 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y' + 13y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve([diff(y(x),x$2)-4*diff(y(x),x)+13*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y = \frac{e^{2x} \sin(3x)}{3}$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 18

```
DSolve[{y''[x]-4*y'[x]+13*y[x]==0,{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}e^{2x} \sin(3x)$$

## 11.36 problem 17.6 (f)

Internal problem ID [13294]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.6 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y' + 13y = 0$$

With initial conditions

$$[y(0) = 5, y'(0) = 31]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve([diff(y(x),x$2)-4*diff(y(x),x)+13*y(x)=0,y(0) = 5, D(y)(0) = 31],y(x), singsol=all)
```

$$y = e^{2x}(7 \sin(3x) + 5 \cos(3x))$$

### ✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 24

```
DSolve[{y'[x]-4*y'[x]+13*y[x]==0,{y[0]==5,y'[0]==31}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(7 \sin(3x) + 5 \cos(3x))$$

## 11.37 problem 17.7 (a)

Internal problem ID [13295]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.7 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y' + \left(\frac{1}{4} + 4\pi^2\right)y = 0$$

With initial conditions

$$\left[ y(0) = 1, y'(0) = \frac{1}{2} \right]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([diff(y(x),x$2)-diff(y(x),x)+(1/4+4*Pi^2)*y(x)=0,y(0) = 1, D(y)(0) = 1/2],y(x), sings
```

$$y = e^{\frac{x}{2}} \cos(2\pi x)$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 18

```
DSolve[{y''[x]-y'[x]+(1/4+4*Pi^2)*y[x]==0,{y[0]==1,y'[0]==1/2}},y[x],x,IncludeSingularSoluti
```

$$y(x) \rightarrow e^{x/2} \cos(2\pi x)$$



## 11.38 problem 17.7 (b)

Internal problem ID [13296]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 17. Second order Homogeneous equations with constant coefficients. Additional exercises page 334

**Problem number:** 17.7 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y' + \left(\frac{1}{4} + 4\pi^2\right)y = 0$$

With initial conditions

$$\left[ y(0) = 1, y'(0) = -\frac{1}{2} \right]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve([diff(y(x),x$2)-diff(y(x),x)+(1/4+4*Pi^2)*y(x)=0,y(0) = 1, D(y)(0) = -1/2],y(x), sing
```

$$y = \frac{e^{\frac{x}{2}}(2\pi \cos(2\pi x) - \sin(2\pi x))}{2\pi}$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 35

```
DSolve[{y''[x]-y'[x]+(1/4+4*Pi^2)*y[x]==0,{y[0]==1,y'[0]==-1/2}},y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow \frac{e^{x/2}(2\pi \cos(2\pi x) - \sin(2\pi x))}{2\pi}$$

## 12 Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

12.1 problem 19.1 (a)	425
12.2 problem 19.1 (b)	426
12.3 problem 19.1 (c)	427
12.4 problem 19.1 (d)	428
12.5 problem 19.1 (e)	429
12.6 problem 19.1 (f)	430
12.7 problem 19.2 (a)	431
12.8 problem 19.2 (b)	432
12.9 problem 19.2 (c)	433
12.10 problem 19.2 (d)	434
12.11 problem 19.2 (e)	435
12.12 problem 19.2 (f)	436
12.13 problem 19.3 (a)	437
12.14 problem 19.3 (b)	438
12.15 problem 19.3 (c)	439
12.16 problem 19.3 (d)	440
12.17 problem 19.4 (a)	441
12.18 problem 19.4 (b)	442
12.19 problem 19.4 (c)	443
12.20 problem 19.4 (d)	444
12.21 problem 19.4 (e)	445
12.22 problem 19.4 (f)	446
12.23 problem 19.4 (g)	447
12.24 problem 19.4 (h)	448
12.25 problem 19.4 (i)	449
12.26 problem 19.4 (j)	450

## 12.1 problem 19.1 (a)

Internal problem ID [13297]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.1 (a).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 4y''' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$4)-4*diff(y(x),x$3)=0,y(x), singsol=all)
```

$$y = c_1 + c_2x + c_3x^2 + c_4e^{4x}$$

### ✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 28

```
DSolve[y''''[x]-4*y'''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{64}c_1e^{4x} + x(c_4x + c_3) + c_2$$

## 12.2 problem 19.1 (b)

Internal problem ID [13298]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.1 (b).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 4y'' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$4)+4*diff(y(x),x$2)=0,y(x), singsol=all)
```

$$y = c_1 + c_2x + c_3 \sin(2x) + c_4 \cos(2x)$$

### ✓ Solution by Mathematica

Time used: 0.077 (sec). Leaf size: 32

```
DSolve[y''''[x]+4*y''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_4x - \frac{1}{4}c_1 \cos(2x) - \frac{1}{4}c_2 \sin(2x) + c_3$$

## 12.3 problem 19.1 (c)

Internal problem ID [13299]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.1 (c).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 34y'' + 225y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$4)-34*diff(y(x),x$2)+225*y(x)=0,y(x), singsol=all)
```

$$y = c_1e^{-5x} + c_2e^{-3x} + c_3e^{5x} + c_4e^{3x}$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 39

```
DSolve[y''''[x]-34*y''[x]+225*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-5x} (e^{2x} (c_3e^{6x} + c_4e^{8x} + c_2) + c_1)$$

## 12.4 problem 19.1 (d)

Internal problem ID [13300]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.1 (d).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 81y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$4)-81*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-3x} + c_2 e^{3x} + c_3 \sin(3x) + c_4 \cos(3x)$$

### ✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 36

```
DSolve[y''''[x]-81*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{3x} + c_3 e^{-3x} + c_2 \cos(3x) + c_4 \sin(3x)$$

## 12.5 problem 19.1 (e)

Internal problem ID [13301]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.1 (e).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 18y'' + 81y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$4)-18*diff(y(x),x$2)+81*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-3x} + c_2 e^{-3x} x + c_3 e^{3x} + c_4 e^{3x} x$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 35

```
DSolve[y''''[x]-18*y''[x]+81*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x} (c_3 e^{6x} + x(c_4 e^{6x} + c_2)) + c_1$$

## 12.6 problem 19.1 (f)

Internal problem ID [13302]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients.  
Additional exercises page 369

**Problem number:** 19.1 (f).

**ODE order:** 5.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y^{(5)} + 18y''' + 81y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
dsolve(diff(y(x),x$5)+18*diff(y(x),x$3)+81*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = c_1 + c_2 \sin(3x) + c_3 \cos(3x) + c_4 \sin(3x)x + c_5 \cos(3x)x$$

### ✓ Solution by Mathematica

Time used: 0.11 (sec). Leaf size: 48

```
DSolve[y'''''[x]+18*y''''[x]+81*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{9}((c_2 - 3(c_4x + c_3)) \cos(3x) + (3c_2x + 3c_1 + c_4) \sin(3x) + 9c_5)$$



## 12.7 problem 19.2 (a)

Internal problem ID [13303]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.2 (a).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - y'' + y' - y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$3)-diff(y(x),x$2)+diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^x + c_2 \sin(x) + c_3 \cos(x)$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 22

```
DSolve[y'''[x]-y''[x]+y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3 e^x + c_1 \cos(x) + c_2 \sin(x)$$

## 12.8 problem 19.2 (b)

Internal problem ID [13304]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.2 (b).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 6y'' + 11y' - 6y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$3)-6*diff(y(x),x$2)+11*diff(y(x),x)-6*y(x)=0,y(x), singsol=all)
```

$$y = e^{2x}c_1 + e^x c_2 + c_3 e^{3x}$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 25

```
DSolve[y'''[x]-6*y''[x]+11*y'[x]-6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(e^x(c_3 e^x + c_2) + c_1)$$

## 12.9 problem 19.2 (c)

Internal problem ID [13305]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients.

Additional exercises page 369

**Problem number:** 19.2 (c).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 8y'' + 37y' - 50y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$3)-8*diff(y(x),x$2)+37*diff(y(x),x)-50*y(x)=0,y(x), singsol=all)
```

$$y = e^{2x}c_1 + c_2e^{3x} \sin(4x) + c_3e^{3x} \cos(4x)$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 34

```
DSolve[y'''[x]-8*y''[x]+37*y'[x]-50*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(c_2e^x \cos(4x) + c_1e^x \sin(4x) + c_3)$$

## 12.10 problem 19.2 (d)

Internal problem ID [13306]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients.

Additional exercises page 369

**Problem number:** 19.2 (d).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 9y'' + 31y' - 39y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$3)-9*diff(y(x),x$2)+31*diff(y(x),x)-39*y(x)=0,y(x), singsol=all)
```

$$y = e^{3x}c_1 + c_2e^{3x} \sin(2x) + c_3e^{3x} \cos(2x)$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 28

```
DSolve[y'''[x]-9*y''[x]+31*y'[x]-39*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{3x}(c_2 \cos(2x) + c_1 \sin(2x) + c_3)$$

## 12.11 problem 19.2 (e)

Internal problem ID [13307]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.2 (e).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + y''' + 2y'' + 4y' - 8y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$4)+diff(y(x),x$3)+2*diff(y(x),x$2)+4*diff(y(x),x)-8*y(x)=0,y(x), singsol=
```

$$y = c_1 e^{-2x} + e^x c_2 + c_3 \sin(2x) + c_4 \cos(2x)$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 34

```
DSolve[y''''[x]+y'''[x]+2*y''[x]+4*y'[x]-8*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3 e^{-2x} + c_4 e^x + c_1 \cos(2x) + c_2 \sin(2x)$$

## 12.12 problem 19.2 (f)

Internal problem ID [13308]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.2 (f).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 2y''' + 10y'' + 18y' + 9y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$4)+2*diff(y(x),x$3)+10*diff(y(x),x$2)+18*diff(y(x),x)+9*y(x)=0,y(x),sing
```

$$y = e^{-x}c_1 + c_2e^{-x}x + c_3 \sin(3x) + c_4 \cos(3x)$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 38

```
DSolve[y''''[x]+2*y'''[x]+10*y''[x]+18*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> Tr
```

$$y(x) \rightarrow e^{-x}(c_4x + c_1e^x \cos(3x) + c_2e^x \sin(3x) + c_3)$$

## 12.13 problem 19.3 (a)

Internal problem ID [13309]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.3 (a).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' + 4y' = 0$$

With initial conditions

$$[y(0) = 4, y'(0) = 6, y''(0) = 8]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve([diff(y(x),x$3)+4*diff(y(x),x)=0,y(0) = 4, D(y)(0) = 6, (D@@2)(y)(0) = 8],y(x), sings
```

$$y = 6 + 3 \sin(2x) - 2 \cos(2x)$$

### ✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 19

```
DSolve[{y'''[x]+4*y'[x]==0,{y[0]==4,y'[0]==6,y''[0]==8}},y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow 3 \sin(2x) - 2 \cos(2x) + 6$$

## 12.14 problem 19.3 (b)

Internal problem ID [13310]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.3 (b).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 6y'' + 12y' - 8y = 0$$

With initial conditions

$$[y(0) = 5, y'(0) = 13, y''(0) = 86]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 19

```
dsolve([diff(y(x),x$3)-6*diff(y(x),x$2)+12*diff(y(x),x)-8*y(x)=0,y(0) = 5, D(y)(0) = 13, D
```

$$y = e^{2x}(27x^2 + 3x + 5)$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 21

```
DSolve[{y'''[x]-6*y''[x]+12*y'[x]-8*y[x]==0,{y[0]==5,y'[0]==13,y''[0]==86}},y[x],x,IncludeSi
```

$$y(x) \rightarrow e^{2x}(27x^2 + 3x + 5)$$



## 12.15 problem 19.3 (c)

Internal problem ID [13311]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients.  
Additional exercises page 369

**Problem number:** 19.3 (c).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 26y'' + 25y = 0$$

With initial conditions

$$[y(0) = 6, y'(0) = -28, y''(0) = -102, y'''(0) = 622]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve([diff(y(x),x$4)+26*diff(y(x),x$2)+25*y(x)=0,y(0) = 6, D(y)(0) = -28, D@@2)(y)(0) = -
```

$$y = -\frac{13 \sin(x)}{4} + 2 \cos(x) - \frac{99 \sin(5x)}{20} + 4 \cos(5x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
DSolve[{y''''[x]+26*y''[x]+25*y[x]==0,{y[0]==6,y'[0]==-28,y''[0]==-102,y'''[0]==622}},y[x],x
```

$$y(x) \rightarrow -\frac{13 \sin(x)}{4} - \frac{99}{20} \sin(5x) + 2 \cos(x) + 4 \cos(5x)$$

## 12.16 problem 19.3 (d)

Internal problem ID [13312]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.3 (d).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + y''' + 9y'' + 9y' = 0$$

With initial conditions

$$[y(0) = 10, y'(0) = 0, y''(0) = 6, y'''(0) = -60]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve([diff(y(x),x$4)+diff(y(x),x$3)+9*diff(y(x),x$2)+9*diff(y(x),x)=0,y(0)=10, D(y)(0)=
```

$$y = 4 + 6e^{-x} + 2\sin(3x)$$

### ✓ Solution by Mathematica

Time used: 0.055 (sec). Leaf size: 20

```
DSolve[{y''''[x]+y'''[x]+9*y''[x]+9*y'[x]==0,{y[0]==10,y'[0]==0,y''[0]==6,y'''[0]==-60}},y[x]
```

$$y(x) \rightarrow 6e^{-x} + 2\sin(3x) + 4$$

## 12.17 problem 19.4 (a)

Internal problem ID [13313]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.4 (a).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 8y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
dsolve(diff(y(x), x$3)-8*y(x)=0,y(x), singsol=all)
```

$$y = e^{2x}c_1 + c_2e^{-x} \sin(\sqrt{3}x) + c_3e^{-x} \cos(\sqrt{3}x)$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 42

```
DSolve[y'''[x]-8*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} \left( c_1 e^{3x} + c_2 \cos(\sqrt{3}x) + c_3 \sin(\sqrt{3}x) \right)$$

## 12.18 problem 19.4 (b)

Internal problem ID [13314]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.4 (b).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' + 216y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 37

```
dsolve(diff(y(x),x$3)+216*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-6x} + c_2 e^{3x} \sin(3\sqrt{3}x) + c_3 e^{3x} \cos(3\sqrt{3}x)$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 48

```
DSolve[y'''[x]+216*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-6x} \left( c_3 e^{9x} \cos(3\sqrt{3}x) + c_2 e^{9x} \sin(3\sqrt{3}x) + c_1 \right)$$

## 12.19 problem 19.4 (c)

Internal problem ID [13315]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.4 (c).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 3y'' - 4y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$4)-3*diff(y(x),x$2)-4*y(x)=0,y(x), singsol=all)
```

$$y = e^{2x}c_1 + c_2e^{-2x} + c_3 \sin(x) + c_4 \cos(x)$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 32

```
DSolve[y''''[x]-3*y''[x]-4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3e^{-2x} + c_4e^{2x} + c_1 \cos(x) + c_2 \sin(x)$$

## 12.20 problem 19.4 (d)

Internal problem ID [13316]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.4 (d).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 13y'' + 36y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$4)+13*diff(y(x),x$2)+36*y(x)=0,y(x), singsol=all)
```

$$y = c_1 \sin(2x) + c_2 \cos(2x) + c_3 \sin(3x) + c_4 \cos(3x)$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 34

```
DSolve[y''''[x]+13*y''[x]+36*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3 \cos(2x) + c_1 \cos(3x) + c_4 \sin(2x) + c_2 \sin(3x)$$

## 12.21 problem 19.4 (e)

Internal problem ID [13317]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.4 (e).

**ODE order:** 6.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y^{(6)} - 3y'''' + 3y'' - y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 43

```
dsolve(diff(y(x),x$6)-3*diff(y(x),x$4)+3*diff(y(x),x$2)-y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^x + c_2 e^x x + c_3 e^x x^2 + c_4 e^{-x} + c_5 e^{-x} x + c_6 e^{-x} x^2$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 50

```
DSolve[y''''''[x]-3*y''''[x]+3*y''[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(x^2(c_6 e^{2x} + c_3) + x(c_5 e^{2x} + c_2) + c_4 e^{2x} + c_1)$$

## 12.22 problem 19.4 (f)

Internal problem ID [13318]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients.

Additional exercises page 369

**Problem number:** 19.4 (f).

**ODE order:** 6.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y^{(6)} - 2y''' + y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 68

```
dsolve(diff(y(x),x$6)-2*diff(y(x),x$3)+y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^x + c_2 e^x x + c_3 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + c_4 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \\ + c_5 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) x + c_6 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) x$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 67

```
DSolve[y''''''[x]-2*y'''[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/2} \left( e^{3x/2} (c_6 x + c_5) + (c_4 x + c_3) \cos\left(\frac{\sqrt{3}x}{2}\right) + (c_2 x + c_1) \sin\left(\frac{\sqrt{3}x}{2}\right) \right)$$



## 12.23 problem 19.4 (g)

Internal problem ID [13319]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.4 (g).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$16y'''' - y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve(16*diff(y(x),x$4)-y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{\frac{x}{2}} + c_2 e^{-\frac{x}{2}} + c_3 \sin\left(\frac{x}{2}\right) + c_4 \cos\left(\frac{x}{2}\right)$$

### ✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 41

```
DSolve[16*y''''[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/2}(c_1 e^x + c_3) + c_2 \cos\left(\frac{x}{2}\right) + c_4 \sin\left(\frac{x}{2}\right)$$

## 12.24 problem 19.4 (h)

Internal problem ID [13320]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.4 (h).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$4y'''' + 15y'' - 4y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(4*diff(y(x),x$4)+15*diff(y(x),x$2)-4*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{\frac{x}{2}} + c_2 e^{-\frac{x}{2}} + c_3 \sin(2x) + c_4 \cos(2x)$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 37

```
DSolve[4*y''''[x]+15*y''[x]-4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/2}(c_4 e^x + c_3) + c_1 \cos(2x) + c_2 \sin(2x)$$

## 12.25 problem 19.4 (i)

Internal problem ID [13321]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients. Additional exercises page 369

**Problem number:** 19.4 (i).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 4y''' + 16y' - 16y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(x),x$4)-4*diff(y(x),x$3)+16*diff(y(x),x)-16*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-2x} + c_2 e^{2x} + c_3 e^{2x} x + c_4 e^{2x} x^2$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 32

```
DSolve[y''''[x]-4*y'''[x]+16*y'[x]-16*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} (e^{4x} (x(c_4 x + c_3) + c_2) + c_1)$$

## 12.26 problem 19.4 (j)

Internal problem ID [13322]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 19. Arbitrary Homogeneous linear equations with constant coefficients.

Additional exercises page 369

**Problem number:** 19.4 (j).

**ODE order:** 6.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y^{(6)} + 16y''' + 64y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 60

```
dsolve(diff(y(x),x$6)+16*diff(y(x),x$3)+64*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-2x} + c_2 e^{-2x} x + c_3 e^x \sin(\sqrt{3} x) + c_4 e^x \cos(\sqrt{3} x) \\ + c_5 e^x \sin(\sqrt{3} x) x + c_6 e^x \cos(\sqrt{3} x) x$$

### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 60

```
DSolve[y''''''[x]+16*y'''[x]+64*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} (c_6 x + e^{3x} (c_4 x + c_3) \cos(\sqrt{3} x) + e^{3x} (c_2 x + c_1) \sin(\sqrt{3} x) + c_5)$$

## 13 Chapter 20. Euler equations. Additional exercises page 382

13.1 problem 20.1 (a)	452
13.2 problem 20.1 (b)	453
13.3 problem 20.1 (c)	454
13.4 problem 20.1 (d)	455
13.5 problem 20.1 (e)	456
13.6 problem 20.1 (f)	457
13.7 problem 20.1 (g)	458
13.8 problem 20.1 (h)	459
13.9 problem 20.1 (i)	460
13.10 problem 20.1 (j)	461
13.11 problem 20.1 (k)	462
13.12 problem 20.1 (L)	463
13.13 problem 20.1 (m)	464
13.14 problem 20.1 (n)	465
13.15 problem 20.1 (o)	466
13.16 problem 20.1 (p)	467
13.17 problem 20.1 (q)	468
13.18 problem 20.1 (r)	469
13.19 problem 20.2 (a)	470
13.20 problem 20.2 (b)	471
13.21 problem 20.2 (c)	472
13.22 problem 20.2 (d)	473
13.23 problem 20.2 (e)	474
13.24 problem 20.2 (f)	475
13.25 problem 20.4 (a)	476
13.26 problem 20.4 (b)	477
13.27 problem 20.4 (c)	478
13.28 problem 20.4 (d)	479
13.29 problem 20.4 (e)	480
13.30 problem 20.4 (f)	481
13.31 problem 20.4 (g)	482
13.32 problem 20.4 (h)	483

## 13.1 problem 20.1 (a)

Internal problem ID [13323]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$x^2y'' - 5y'x + 8y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)-5*x*diff(y(x),x)+8*y(x)=0,y(x), singsol=all)
```

$$y = c_2x^4 + c_1x^2$$

### ✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]-5*x*y'[x]+8*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2(c_2x^2 + c_1)$$

## 13.2 problem 20.1 (b)

Internal problem ID [13324]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x^2 y'' - 2y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)-2*y(x)=0,y(x), singsol=all)
```

$$y = c_1 x^2 + \frac{c_2}{x}$$

### ✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 x^3 + c_1}{x}$$

### 13.3 problem 20.1 (c)

Internal problem ID [13325]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^2 y'' - 2y'x = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(x^2*diff(y(x),x$2)-2*x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = c_2 x^3 + c_1$$

#### ✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 17

```
DSolve[x^2*y''[x]-2*x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1 x^3}{3} + c_2$$



## 13.4 problem 20.1 (d)

Internal problem ID [13326]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$2x^2y'' - y'x + y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(2*x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y = c_1x + c_2\sqrt{x}$$

### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 18

```
DSolve[2*x^2*y''[x]-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1\sqrt{x} + c_2x$$

## 13.5 problem 20.1 (e)

Internal problem ID [13327]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - 5y'x + 9y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x^2*diff(y(x),x$2)-5*x*diff(y(x),x)+9*y(x)=0,y(x), singsol=all)
```

$$y = c_1 x^3 + \ln(x) x^3 c_2$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]-5*x*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^3(3c_2 \log(x) + c_1)$$

## 13.6 problem 20.1 (f)

Internal problem ID [13328]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' + 5y'x + 4y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve(x^2*diff(y(x),x$2)+5*x*diff(y(x),x)+4*y(x)=0,y(x), singsol=all)
```

$$y = \frac{c_1}{x^2} + \frac{c_2 \ln(x)}{x^2}$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]+5*x*y'[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2c_2 \log(x) + c_1}{x^2}$$

## 13.7 problem 20.1 (g)

Internal problem ID [13329]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$4x^2y'' + y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(4*x^2*diff(y(x),x$2)+y(x)=0,y(x), singsol=all)
```

$$y = c_1\sqrt{x} + c_2\sqrt{x} \ln(x)$$

### ✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 24

```
DSolve[4*x^2*y''[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}\sqrt{x}(c_2 \log(x) + 2c_1)$$

## 13.8 problem 20.1 (h)

Internal problem ID [13330]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - 19y'x + 100y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x^2*diff(y(x),x$2)-19*x*diff(y(x),x)+100*y(x)=0,y(x), singsol=all)
```

$$y = c_1 x^{10} + c_2 x^{10} \ln(x)$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]-19*x*y'[x]+100*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^{10}(10c_2 \log(x) + c_1)$$

### 13.9 problem 20.1 (i)

Internal problem ID [13331]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - 5y'x + 29y = 0$$

#### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 31

```
dsolve(x^2*diff(y(x),x$2)-5*x*diff(y(x),x)+29*y(x)=0,y(x), singsol=all)
```

$$y = c_1 x^3 \sin\left(2\sqrt{5} \ln(x)\right) + c_2 x^3 \cos\left(2\sqrt{5} \ln(x)\right)$$

#### ✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 36

```
DSolve[x^2*y'[x]-5*x*y'[x]+29*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^3 \left( c_2 \cos\left(2\sqrt{5} \log(x)\right) + c_1 \sin\left(2\sqrt{5} \log(x)\right) \right)$$

## 13.10 problem 20.1 (j)

Internal problem ID [13332]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (j).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - y'x + 10y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(x^2*diff(y(x),x$2)-x*diff(y(x),x)+10*y(x)=0,y(x), singsol=all)
```

$$y = c_1 x \sin(3 \ln(x)) + c_2 x \cos(3 \ln(x))$$

### ✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 24

```
DSolve[x^2*y'[x]-x*y'[x]+10*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(c_2 \cos(3 \log(x)) + c_1 \sin(3 \log(x)))$$

## 13.11 problem 20.1 (k)

Internal problem ID [13333]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (k).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' + 5y'x + 29y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(x^2*diff(y(x),x$2)+5*x*diff(y(x),x)+29*y(x)=0,y(x), singsol=all)
```

$$y = \frac{c_1 \sin(5 \ln(x))}{x^2} + \frac{c_2 \cos(5 \ln(x))}{x^2}$$

### ✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 26

```
DSolve[x^2*y''[x]+5*x*y'[x]+29*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 \cos(5 \log(x)) + c_1 \sin(5 \log(x))}{x^2}$$



## 13.12 problem 20.1 (L)

Internal problem ID [13334]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (L).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$x^2 y'' + y' x + y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y = c_1 \sin(\ln(x)) + c_2 \cos(\ln(x))$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]+x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(\log(x)) + c_2 \sin(\log(x))$$

### 13.13 problem 20.1 (m)

Internal problem ID [13335]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (m).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$2x^2y'' + 5y'x + y = 0$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(2*x^2*diff(y(x),x$2)+5*x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y = \frac{c_1}{x} + \frac{c_2}{\sqrt{x}}$$

#### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 20

```
DSolve[2*x^2*y''[x]+5*x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2\sqrt{x} + c_1}{x}$$

### 13.14 problem 20.1 (n)

Internal problem ID [13336]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (n).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_Emden, _Fowler]`

$$4x^2y'' + 37y = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(4*x^2*diff(y(x),x$2)+37*y(x)=0,y(x), singsol=all)
```

$$y = c_1\sqrt{x} \sin(3 \ln(x)) + c_2\sqrt{x} \cos(3 \ln(x))$$

#### ✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 28

```
DSolve[4*x^2*y''[x]+37*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt{x}(c_2 \cos(3 \log(x)) + c_1 \sin(3 \log(x)))$$

## 13.15 problem 20.1 (o)

Internal problem ID [13337]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (o).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$x^2 y'' + y' x = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = c_2 \ln(x) + c_1$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 13

```
DSolve[x^2*y'[x]+x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \log(x) + c_2$$

## 13.16 problem 20.1 (p)

Internal problem ID [13338]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (p).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$x^2y'' + y'x - 25y = 0$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-25*y(x)=0,y(x), singsol=all)
```

$$y = \frac{c_1}{x^5} + c_2x^5$$

### ✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]+x*y'[x]-25*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2x^5 + \frac{c_1}{x^5}$$

### 13.17 problem 20.1 (q)

Internal problem ID [13339]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (q).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$4x^2y'' + 8y'x + 5y = 0$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(4*x^2*diff(y(x),x$2)+8*x*diff(y(x),x)+5*y(x)=0,y(x), singsol=all)
```

$$y = \frac{c_1 \sin(\ln(x))}{\sqrt{x}} + \frac{c_2 \cos(\ln(x))}{\sqrt{x}}$$

#### ✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 24

```
DSolve[4*x^2*y''[x]+8*x*y'[x]+5*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 \cos(\log(x)) + c_1 \sin(\log(x))}{\sqrt{x}}$$

### 13.18 problem 20.1 (r)

Internal problem ID [13340]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.1 (r).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$3x^2y'' - 7y'x + 3y = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(3*x^2*diff(y(x),x$2)-7*x*diff(y(x),x)+3*y(x)=0,y(x), singsol=all)
```

$$y = c_1x^{\frac{1}{3}} + c_2x^3$$

#### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 20

```
DSolve[3*x^2*y'[x]-7*x*y'[x]+3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2x^3 + c_1\sqrt[3]{x}$$

### 13.19 problem 20.2 (a)

Internal problem ID [13341]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.2 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$x^2y'' - 2y'x - 10y = 0$$

With initial conditions

$$[y(1) = 5, y'(1) = 4]$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([x^2*diff(y(x),x$2)-2*x*diff(y(x),x)-10*y(x)=0,y(1) = 5, D(y)(1) = 4],y(x), singsol=a
```

$$y = \frac{3}{x^2} + 2x^5$$

#### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 16

```
DSolve[{x^2*y''[x]-2*x*y'[x]-10*y[x]==0,{y[1]==5,y'[1]==4}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{2x^7 + 3}{x^2}$$



## 13.20 problem 20.2 (b)

Internal problem ID [13342]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.2 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [[\_Emden, \_Fowler], [\_2nd\_order, \_linear, ‘\_with\_symmetry\_[0,F

$$4x^2y'' + 4y'x - y = 0$$

With initial conditions

$$[y(4) = 0, y'(4) = 2]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve([4*x^2*diff(y(x),x$2)+4*x*diff(y(x),x)-y(x)=0,y(4) = 0, D(y)(4) = 2],y(x), singsol=all
```

$$y = \frac{4x - 16}{\sqrt{x}}$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 15

```
DSolve[{4*x^2*y''[x]+4*x*y'[x]-y[x]==0,{y[4]==0,y'[4]==2}},y[x],x,IncludeSingularSolutions -
```

$$y(x) \rightarrow \frac{4(x - 4)}{\sqrt{x}}$$

### 13.21 problem 20.2 (c)

Internal problem ID [13343]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.2 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - 11y'x + 36y = 0$$

With initial conditions

$$\left[ y(1) = \frac{1}{2}, y'(1) = 2 \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 14

```
dsolve([x^2*diff(y(x),x$2)-11*x*diff(y(x),x)+36*y(x)=0,y(1) = 1/2, D(y)(1) = 2],y(x), singsol
```

$$y = x^6 \left( \frac{1}{2} - \ln(x) \right)$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 18

```
DSolve[{x^2*y'[x]-11*x*y'[x]+36*y[x]==0,{y[1]==1/2,y'[1]==2}},y[x],x,IncludeSingularSolutio
```

$$y(x) \rightarrow \frac{1}{2}x^6(1 - 2\log(x))$$

## 13.22 problem 20.2 (d)

Internal problem ID [13344]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.2 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - y' x + y = 0$$

With initial conditions

$$[y(1) = 3, y'(1) = 0]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,y(1) = 3, D(y)(1) = 0],y(x), singsol=all)
```

$$y = 3x - 3x \ln(x)$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 12

```
DSolve[{x^2*y'[x]-x*y[x]+y[x]==0,{y[1]==3,y'[1]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -3x(\log(x) - 1)$$

### 13.23 problem 20.2 (e)

Internal problem ID [13345]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.2 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - y' x + 2y = 0$$

With initial conditions

$$[y(1) = 3, y'(1) = 0]$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve([x^2*diff(y(x),x$2)-x*diff(y(x),x)+2*y(x)=0,y(1) = 3, D(y)(1) = 0],y(x), singsol=all)
```

$$y = -3x(\sin(\ln(x)) - \cos(\ln(x)))$$

#### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 17

```
DSolve[{x^2*y'[x]-x*y[x]+2*y[x]==0,{y[1]==3,y'[1]==0}},y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow 3x(\cos(\log(x)) - \sin(\log(x)))$$

## 13.24 problem 20.2 (f)

Internal problem ID [13346]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.2 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - 3y'x + 13y = 0$$

With initial conditions

$$[y(1) = 9, y'(1) = 3]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve([x^2*diff(y(x),x$2)-3*x*diff(y(x),x)+13*y(x)=0,y(1) = 9, D(y)(1) = 3],y(x), singsol=a
```

$$y = x^2(-5 \sin(3 \ln(x)) + 9 \cos(3 \ln(x)))$$

### ✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 24

```
DSolve[{x^2*y'[x]-3*x*y'[x]+13*y[x]==0,{y[1]==9,y'[1]==3}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow x^2(9 \cos(3 \log(x)) - 5 \sin(3 \log(x)))$$

## 13.25 problem 20.4 (a)

Internal problem ID [13347]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.4 (a).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$x^3 y''' + 2x^2 y'' - 4y'x + 4y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(x^3*diff(y(x),x$3)+2*x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+4*y(x)=0,y(x), singsol=all)
```

$$y = c_1 x^2 + \frac{c_2}{x^2} + c_3 x$$

### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 22

```
DSolve[x^3*y'''[x]+2*x^2*y''[x]-4*x*y'[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3 x^2 + \frac{c_1}{x^2} + c_2 x$$

## 13.26 problem 20.4 (b)

Internal problem ID [13348]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.4 (b).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$x^3 y''' + 2x^2 y'' + y' x - y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(x^3*diff(y(x),x$3)+2*x^2*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y = c_1 x + c_2 \sin(\ln(x)) + c_3 \cos(\ln(x))$$

### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 22

```
DSolve[x^3*y'''[x]+2*x^2*y''[x]+x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3 x + c_1 \cos(\log(x)) + c_2 \sin(\log(x))$$

### 13.27 problem 20.4 (c)

Internal problem ID [13349]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.4 (c).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$x^3 y''' - 5x^2 y'' + 14y'x - 18y = 0$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 22

```
dsolve(x^3*diff(y(x),x$3)-5*x^2*diff(y(x),x$2)+14*x*diff(y(x),x)-18*y(x)=0,y(x), singsol=all
```

$$y = c_1 x^2 + c_2 x^3 + c_3 x^3 \ln(x)$$

#### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 22

```
DSolve[x^3*y'''[x]-5*x^2*y''[x]+14*x*y'[x]-18*y[x]==0,y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow x^2(c_2 x + c_3 x \log(x) + c_1)$$



## 13.28 problem 20.4 (d)

Internal problem ID [13350]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.4 (d).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$x^3 y''' - 3x^2 y'' + 7y'x - 8y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 26

```
dsolve(x^3*diff(y(x),x$3)-3*x^2*diff(y(x),x$2)+7*x*diff(y(x),x)-8*y(x)=0,y(x), singsol=all)
```

$$y = c_1 x^2 + \ln(x) x^2 c_2 + \ln(x)^2 x^2 c_3$$

### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 24

```
DSolve[x^3*y'''[x]-3*x^2*y''[x]+7*x*y'[x]-8*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2 (c_3 \log^2(x) + c_2 \log(x) + c_1)$$

### 13.29 problem 20.4 (e)

Internal problem ID [13351]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.4 (e).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _with_linear_symmetries]]`

$$x^4 y'''' + 6x^3 y'''' + 15x^2 y'' + 9y'x + 16y = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 37

```
dsolve(x^4*diff(y(x),x$4)+6*x^3*diff(y(x),x$3)+15*x^2*diff(y(x),x$2)+9*x*diff(y(x),x)+16*y(x))
```

$$y = c_1 \sin(2 \ln(x)) + c_2 \cos(2 \ln(x)) + c_3 \sin(2 \ln(x)) \ln(x) + c_4 \cos(2 \ln(x)) \ln(x)$$

#### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 34

```
DSolve[x^4*y''''[x]+6*x^3*y''''[x]+15*x^2*y''[x]+9*x*y'[x]+16*y[x]==0,y[x],x,IncludeSingularS
```

$$y(x) \rightarrow (c_2 \log(x) + c_1) \cos(2 \log(x)) + (c_4 \log(x) + c_3) \sin(2 \log(x))$$

### 13.30 problem 20.4 (f)

Internal problem ID [13352]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.4 (f).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _exact, _linear, _homogeneous]]`

$$x^4 y'''' + 6x^3 y''' - 3x^2 y'' - 9y'x + 9y = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(x^4*diff(y(x),x$4)+6*x^3*diff(y(x),x$3)-3*x^2*diff(y(x),x$2)-9*x*diff(y(x),x)+9*y(x)=
```

$$y = c_1 x + \frac{c_2}{x^3} + \frac{c_3}{x} + c_4 x^3$$

#### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 28

```
DSolve[x^4*y''''[x]+6*x^3*y'''[x]-3*x^2*y''[x]-9*x*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSol
```

$$y(x) \rightarrow c_4 x^3 + \frac{c_1}{x^3} + c_3 x + \frac{c_2}{x}$$

### 13.31 problem 20.4 (g)

Internal problem ID [13353]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.4 (g).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _with_linear_symmetries]]`

$$x^4 y'''' + 2x^3 y''' + x^2 y'' - y'x + y = 0$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(x^4*diff(y(x),x$4)+2*x^3*diff(y(x),x$3)+x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,y(x)
```

$$y = c_1 x + c_2 x \ln(x) + c_3 x \ln(x)^2 + c_4 x \ln(x)^3$$

#### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 29

```
DSolve[x^4*y''''[x]+2*x^3*y'''[x]+x^2*y''[x]-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow x(c_4 \log^3(x) + c_3 \log^2(x) + c_2 \log(x) + c_1)$$

### 13.32 problem 20.4 (h)

Internal problem ID [13354]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 20. Euler equations. Additional exercises page 382

**Problem number:** 20.4 (h).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[_high_order, _exact, _linear, _homogeneous]`

$$x^4 y'''' + 6x^3 y''' + 7x^2 y'' + y'x - y = 0$$

#### ✓ Solution by Maple

Time used: 0.11 (sec). Leaf size: 23

```
dsolve(x^4*diff(y(x),x$4)+6*x^3*diff(y(x),x$3)+7*x^2*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=0,y(x))
```

$$y = c_1 x + \frac{c_2}{x} + c_3 \sin(\ln(x)) + c_4 \cos(\ln(x))$$

#### ✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 28

```
DSolve[x^4*y''''[x]+6*x^3*y'''[x]+7*x^2*y''[x]+x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutio
```

$$y(x) \rightarrow c_1 x + \frac{c_3}{x} + c_2 \cos(\log(x)) + c_4 \sin(\log(x))$$

## 14 Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

14.1 problem 21.5 (i) . . . . .	485
14.2 problem 21.5 (ii) . . . . .	486
14.3 problem 21.6 (i) . . . . .	487
14.4 problem 21.6 (ii) . . . . .	488
14.5 problem 21.7 . . . . .	489
14.6 problem 21.8 . . . . .	490
14.7 problem 21.9 . . . . .	491
14.8 problem 21.10 . . . . .	492
14.9 problem 21.11 . . . . .	493
14.10 problem 21.12 . . . . .	494
14.11 problem 21.13 (a) . . . . .	495
14.12 problem 21.13 (b) . . . . .	496
14.13 problem 21.13 (c) . . . . .	497
14.14 problem 21.13 (d) . . . . .	498
14.15 problem 21.14 (a) . . . . .	499
14.16 problem 21.14 (b) . . . . .	500
14.17 problem 21.14 (c) . . . . .	501
14.18 problem 21.15 (i) . . . . .	502
14.19 problem 21.15 (ii) . . . . .	503
14.20 problem 21.15 (iii) . . . . .	504
14.21 problem 21.15 (c) . . . . .	505

## 14.1 problem 21.5 (i)

Internal problem ID [13355]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.5 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 4y = 24e^{2x}$$

With initial conditions

$$[y(0) = 6, y'(0) = 6]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)+4*y(x)=24*exp(2*x),y(0) = 6, D(y)(0) = 6],y(x), singsol=all)
```

$$y = 3 \cos(2x) + 3e^{2x}$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 17

```
DSolve[{y''[x]+4*y[x]==24*Exp[2*x],{y[0]==6,y'[0]==6}},y[x],x,IncludeSingularSolutions -> Tr
```

$$y(x) \rightarrow 3(e^{2x} + \cos(2x))$$

## 14.2 problem 21.5 (ii)

Internal problem ID [13356]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.5 (ii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 4y = 24e^{2x}$$

With initial conditions

$$[y(0) = -2, y'(0) = 2]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve([diff(y(x),x$2)+4*y(x)=24*exp(2*x),y(0) = -2, D(y)(0) = 2],y(x), singsol=all)
```

$$y = -2 \sin(2x) - 5 \cos(2x) + 3e^{2x}$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 25

```
DSolve[{y''[x]+4*y[x]==24*Exp[2*x],{y[0]==-2,y'[0]==2}},y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow 3e^{2x} - 2 \sin(2x) - 5 \cos(2x)$$



### 14.3 problem 21.6 (i)

Internal problem ID [13357]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.6 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 2y' - 8y = 8x^2 - 3$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

#### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 25

```
dsolve([diff(y(x),x$2)+2*diff(y(x),x)-8*y(x)=8*x^2-3,y(0) = 0, D(y)(0) = 0],y(x), singsol=all)
```

$$y = \frac{(-12x^2 - 6x)e^{-4x}e^{4x}}{12} + \frac{(e^{6x} - 1)e^{-4x}}{12}$$

#### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 34

```
DSolve[{y'[x]+2*y'[x]-8*y[x]==8*x^2-3,{y[0]==0,y'[0]==0}},y[x],x,IncludeSingularSolutions -
```

$$y(x) \rightarrow \frac{1}{12}e^{-4x}(-6e^{4x}x(2x+1) + e^{6x} - 1)$$

## 14.4 problem 21.6 (ii)

Internal problem ID [13358]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.6 (ii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 2y' - 8y = 8x^2 - 3$$

With initial conditions

$$[y(0) = 1, y'(0) = -3]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve([diff(y(x),x$2)+2*diff(y(x),x)-8*y(x)=8*x^2-3,y(0) = 1, D(y)(0) = -3],y(x), singsol=a
```

$$y = \frac{(-4x^2 - 2x)e^{-4x}e^{4x}}{4} + \frac{(e^{6x} + 3)e^{-4x}}{4}$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 34

```
DSolve[{y'[x]+2*y'[x]-8*y[x]==8*x^2-3,{y[0]==1,y'[0]==-3}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{1}{4}e^{-4x}(-2e^{4x}x(2x+1) + e^{6x} + 3)$$

## 14.5 problem 21.7

Internal problem ID [13359]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.7.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 9y = 36$$

With initial conditions

$$[y(0) = 8, y'(0) = 6]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 18

```
dsolve([diff(y(x),x$2)-9*y(x)=36,y(0) = 8, D(y)(0) = 6],y(x), singsol=all)
```

$$y = 5e^{-3x} + 7e^{3x} - 4$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 21

```
DSolve[{y'[x]-9*y[x]==36,{y[0]==8,y'[0]==6}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 5e^{-3x} + 7e^{3x} - 4$$

## 14.6 problem 21.8

Internal problem ID [13360]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.8.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 3y' - 10y = -6e^{4x}$$

With initial conditions

$$[y(0) = 6, y'(0) = 8]$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 21

```
dsolve([diff(y(x),x$2)-3*diff(y(x),x)-10*y(x)=-6*exp(4*x),y(0) = 6, D(y)(0) = 8],y(x), sings
```

$$y = (2e^{7x} + e^{6x} + 3)e^{-2x}$$

### ✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 25

```
DSolve[{y'[x]-3*y'[x]-10*y[x]==-6*Exp[4*x],{y[0]==6,y'[0]==8}},y[x],x,IncludeSingularSoluti
```

$$y(x) \rightarrow e^{-2x}(e^{6x} + 2e^{7x} + 3)$$

## 14.7 problem 21.9

Internal problem ID [13361]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.9.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 3y' - 10y = 7e^{5x}$$

With initial conditions

$$[y(0) = 12, y'(0) = -2]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve([diff(y(x),x$2)-3*diff(y(x),x)-10*y(x)=7*exp(5*x),y(0) = 12, D(y)(0) = -2],y(x), sing
```

$$y = (9 + (x + 3)e^{7x})e^{-2x}$$

### ✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 22

```
DSolve[{y'[x]-3*y'[x]-10*y[x]==7*Exp[5*x],{y[0]==12,y'[0]==-2}},y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow e^{-2x}(e^{7x}(x + 3) + 9)$$

## 14.8 problem 21.10

Internal problem ID [13362]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.10.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 6y' + 9y = 169 \sin(2x)$$

With initial conditions

$$[y(0) = -10, y'(0) = 9]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve([diff(y(x),x$2)+6*diff(y(x),x)+9*y(x)=169*sin(2*x),y(0) = -10, D(y)(0) = 9],y(x), sin
```

$$y = (5x + 2)e^{-3x} - 12 \cos(2x) + 5 \sin(2x)$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 34

```
DSolve[{y'[x]+6*y'[x]+9*y[x]==169*Sin[2*x],{y[0]==-10,y'[0]==9}},y[x],x,IncludeSingularSolu
```

$$y(x) \rightarrow e^{-3x}(5x + 5e^{3x} \sin(2x) + 2) - 12 \cos(2x)$$

## 14.9 problem 21.11

Internal problem ID [13363]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.11.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2y'' - 4y'x + 6y = 10x + 12$$

With initial conditions

$$[y(1) = 6, y'(1) = 8]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve([x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+6*y(x)=10*x+12,y(1) = 6, D(y)(1) = 8],y(x), sing
```

$$y = 5x^3 - 6x^2 + 5x + 2$$

### ✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 20

```
DSolve[{x^2*y'[x]-4*x*y'[x]+6*y[x]==10*x+12,{y[1]==6,y'[1]==8}},y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow 5x^3 - 6x^2 + 5x + 2$$

## 14.10 problem 21.12

Internal problem ID [13364]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.12.

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + y'' = 1$$

With initial conditions

$$[y(0) = 4, y'(0) = 3, y''(0) = 0, y'''(0) = 2]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve([diff(y(x),x$4)+diff(y(x),x$2)=1,y(0) = 4, D(y)(0) = 3, (D@@2)(y)(0) = 0, (D@@3)(y)(0) = 2],y(x),x)
```

$$y = \frac{x^2}{2} + \cos(x) - 2 \sin(x) + 5x + 3$$

### ✓ Solution by Mathematica

Time used: 0.099 (sec). Leaf size: 23

```
DSolve[{y''''[x]+y''[x]==1,{y[0]==4,y'[0]==3,y''[0]==0,y'''[0]==2}},y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{x^2}{2} + 5x - 2 \sin(x) + \cos(x) + 3$$



## 14.11 problem 21.13 (a)

Internal problem ID [13365]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.13 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 3y' - 10y = e^{4x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)-10*y(x)=exp(4*x),y(x), singsol=all)
```

$$y = c_2 e^{-2x} + c_1 e^{5x} - \frac{e^{4x}}{6}$$

### ✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 31

```
DSolve[y''[x]-3*y'[x]-10*y[x]==Exp[4*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{e^{4x}}{6} + c_1 e^{-2x} + c_2 e^{5x}$$

## 14.12 problem 21.13 (b)

Internal problem ID [13366]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.13 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 3y' - 10y = e^{5x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)-10*y(x)=exp(5*x),y(x), singsol=all)
```

$$y = c_2 e^{-2x} + c_1 e^{5x} + \frac{x e^{5x}}{7}$$

### ✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 31

```
DSolve[y''[x]-3*y'[x]-10*y[x]==Exp[5*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-2x} + e^{5x} \left( \frac{x}{7} - \frac{1}{49} + c_2 \right)$$

### 14.13 problem 21.13 (c)

Internal problem ID [13367]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.13 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 3y' - 10y = -18e^{4x} + 14e^{5x}$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)-10*y(x)=-18*exp(4*x)+14*exp(5*x),y(x), singsol=all)
```

$$y = c_2 e^{-2x} + c_1 e^{5x} + 2x e^{5x} - \frac{2e^{5x}}{7} + 3e^{4x}$$

#### ✓ Solution by Mathematica

Time used: 0.057 (sec). Leaf size: 36

```
DSolve[y''[x]-3*y'[x]-10*y[x]==-18*Exp[4*x]+14*Exp[5*x],y[x],x,IncludeSingularSolutions->T
```

$$y(x) \rightarrow 3e^{4x} + c_1 e^{-2x} + e^{5x} \left( 2x - \frac{2}{7} + c_2 \right)$$

## 14.14 problem 21.13 (d)

Internal problem ID [13368]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.13 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 3y' - 10y = 35e^{5x} + 12e^{4x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)-10*y(x)=35*exp(5*x)+12*exp(4*x),y(x), singsol=all)
```

$$y = c_2 e^{-2x} + c_1 e^{5x} + 5x e^{5x} - \frac{5 e^{5x}}{7} - 2 e^{4x}$$

### ✓ Solution by Mathematica

Time used: 0.054 (sec). Leaf size: 36

```
DSolve[y''[x]-3*y'[x]-10*y[x]==35*Exp[5*x]+12*Exp[4*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2e^{4x} + c_1 e^{-2x} + e^{5x} \left( 5x - \frac{5}{7} + c_2 \right)$$

## 14.15 problem 21.14 (a)

Internal problem ID [13369]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.14 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2y'' - 4y'x + 6y = 1$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve(x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+6*y(x)=1,y(x), singsol=all)
```

$$y = c_2x^2 + c_1x^3 + \frac{1}{6}$$

### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 21

```
DSolve[x^2*y'[x]-4*x*y'[x]+6*y[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2x^3 + c_1x^2 + \frac{1}{6}$$

## 14.16 problem 21.14 (b)

Internal problem ID [13370]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.14 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2y'' - 4y'x + 6y = x$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve(x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+6*y(x)=x,y(x), singsol=all)
```

$$y = c_2x^2 + c_1x^3 + \frac{1}{2}x$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 23

```
DSolve[x^2*y''[x]-4*x*y'[x]+6*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2x^3 + c_1x^2 + \frac{x}{2}$$

## 14.17 problem 21.14 (c)

Internal problem ID [13371]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.14 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2y'' - 4y'x + 6y = 22x + 24$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 19

```
dsolve(x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+6*y(x)=22*x+24,y(x), singsol=all)
```

$$y = c_1x^3 + c_2x^2 + 11x + 4$$

### ✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 22

```
DSolve[x^2*y''[x]-4*x*y'[x]+6*y[x]==22*x+24,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2x^3 + c_1x^2 + 11x + 4$$

## 14.18 problem 21.15 (i)

Internal problem ID [13372]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.15 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 7y'x + 15y = x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(x^2*diff(y(x),x$2)-7*x*diff(y(x),x)+15*y(x)=x^2,y(x), singsol=all)
```

$$y = c_2 x^5 + c_1 x^3 + \frac{1}{3} x^2$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 25

```
DSolve[x^2*y'[x]-7*x*y'[x]+15*y[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 x^5 + c_1 x^3 + \frac{x^2}{3}$$



## 14.19 problem 21.15 (ii)

Internal problem ID [13373]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.15 (ii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2y'' - 7y'x + 15y = x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(x^2*diff(y(x),x$2)-7*x*diff(y(x),x)+15*y(x)=x,y(x), singsol=all)
```

$$y = c_2x^5 + c_1x^3 + \frac{1}{8}x$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 23

```
DSolve[x^2*y''[x]-7*x*y'[x]+15*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2x^5 + c_1x^3 + \frac{x}{8}$$

## 14.20 problem 21.15 (iii)

Internal problem ID [13374]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.15 (iii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2y'' - 7y'x + 15y = 1$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve(x^2*diff(y(x),x$2)-7*x*diff(y(x),x)+15*y(x)=1,y(x), singsol=all)
```

$$y = c_2x^5 + c_1x^3 + \frac{1}{15}$$

### ✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 21

```
DSolve[x^2*y''[x]-7*x*y'[x]+15*y[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2x^5 + c_1x^3 + \frac{1}{15}$$

## 14.21 problem 21.15 (c)

Internal problem ID [13375]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 21. Nonhomogeneous equations in general. Additional exercises page 391

**Problem number:** 21.15 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 7y'x + 15y = 4x^2 + 2x + 3$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
dsolve(x^2*diff(y(x),x$2)-7*x*diff(y(x),x)+15*y(x)=4*x^2+2*x+3,y(x), singsol=all)
```

$$y = c_2 x^5 + c_1 x^3 + \frac{1}{5} + \frac{4}{3} x^2 + \frac{1}{4} x$$

### ✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 33

```
DSolve[x^2*y''[x]-7*x*y'[x]+15*y[x]==4*x^2+2*x+3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 x^5 + c_1 x^3 + \frac{4x^2}{3} + \frac{x}{4} + \frac{1}{5}$$

## 15 Chapter 22. Method of undetermined coefficients. Additional exercises page 412

15.1 problem 22.1 (a)	509
15.2 problem 22.1 (b)	510
15.3 problem 22.1 (c)	511
15.4 problem 22.1 (d)	512
15.5 problem 22.2	513
15.6 problem 22.3 (a)	514
15.7 problem 22.3 (b)	515
15.8 problem 22.3 (c)	516
15.9 problem 22.3 (d)	517
15.10 problem 22.4	518
15.11 problem 22.5 (a)	519
15.12 problem 22.5 (b)	520
15.13 problem 22.5 (c)	521
15.14 problem 22.5 (d)	522
15.15 problem 22.6	523
15.16 problem 22.7 (a)	524
15.17 problem 22.7 (b)	525
15.18 problem 22.7 (c)	526
15.19 problem 22.7 (d)	527
15.20 problem 22.7 (e)	528
15.21 problem 22.7 (f)	529
15.22 problem 22.8	530
15.23 problem 22.9 (a)	531
15.24 problem 22.9 (b)	532
15.25 problem 22.9 (c)	533
15.26 problem 22.9 (d)	534
15.27 problem 22.9 (e)	535
15.28 problem 22.10 (a)	536
15.29 problem 22.10 (b)	537
15.30 problem 22.10 (c)	538
15.31 problem 22.10 (d)	539
15.32 problem 22.10 (e)	540
15.33 problem 22.10 (f)	541
15.34 problem 22.10 (g)	542
15.35 problem 22.10 (h)	543

15.36problem 22.10 (i)	544
15.37problem 22.10 (j)	545
15.38problem 22.10 (k)	546
15.39problem 22.10 (L)	547
15.40problem 22.10 (m)	548
15.41problem 22.10 (n)	549
15.42problem 22.11 (a)	550
15.43problem 22.11 (b)	551
15.44problem 22.11 (c)	552
15.45problem 22.11 (d)	553
15.46problem 22.11 (e)	554
15.47problem 22.11 (f)	555
15.48problem 22.11 (g)	556
15.49problem 22.11 (h)	557
15.50problem 22.11 (i)	558
15.51problem 22.11 (j)	559
15.52problem 22.11 (k)	560
15.53problem 22.11 (L)	561
15.54problem 22.11 (m)	562
15.55problem 22.11 (n)	563
15.56problem 22.12 (a)	564
15.57problem 22.12 (b)	565
15.58problem 22.12 (c)	566
15.59problem 22.12 (d)	567
15.60problem 22.12 (e)	568
15.61problem 22.12 (f)	569
15.62problem 22.12 (g)	570
15.63problem 22.13 (a)	571
15.64problem 22.13 (b)	572
15.65problem 22.13 (c)	573
15.66problem 22.13 (d)	574
15.67problem 22.13 (e)	575
15.68problem 22.13 (f)	576
15.69problem 22.13 (g)	577
15.70problem 22.14 (a)	578
15.71problem 22.14 (b)	579
15.72problem 22.14 (c)	580
15.73problem 22.14 (d)	581
15.74problem 22.15 (a)	582

15.75problem 22.15 (b) . . . . .	583
15.76problem 22.15 (c) . . . . .	584
15.77problem 22.15 (d) . . . . .	585
15.78problem 22.15 (e) . . . . .	586
15.79problem 22.15 (f) . . . . .	587
15.80problem 22.15 (g) . . . . .	588
15.81problem 22.15 (h) . . . . .	589

## 15.1 problem 22.1 (a)

Internal problem ID [13376]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.1 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 9y = 52 e^{2x}$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)+9*y(x)=52*exp(2*x),y(x), singsol=all)
```

$$y = c_2 \sin(3x) + \cos(3x) c_1 + 4 e^{2x}$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 27

```
DSolve[y''[x]+9*y[x]==52*Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 4e^{2x} + c_1 \cos(3x) + c_2 \sin(3x)$$

## 15.2 problem 22.1 (b)

Internal problem ID [13377]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.1 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 6y' + 9y = 27e^{6x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)-6*diff(y(x),x)+9*y(x)=27*exp(6*x),y(x), singsol=all)
```

$$y = c_2 e^{3x} + x e^{3x} c_1 + 3 e^{6x}$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 25

```
DSolve[y''[x]-6*y'[x]+9*y[x]==27*Exp[6*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{3x} (3e^{3x} + c_2 x + c_1)$$



## 15.3 problem 22.1 (c)

Internal problem ID [13378]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.1 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 4y' - 5y = 30e^{-4x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)-5*y(x)=30*exp(-4*x),y(x), singsol=all)
```

$$y = e^{-5x}c_2 + c_1e^x - 6e^{-4x}$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 27

```
DSolve[y''[x]+4*y'[x]-5*y[x]==30*Exp[-4*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-5x}(-6e^x + c_2e^{6x} + c_1)$$

## 15.4 problem 22.1 (d)

Internal problem ID [13379]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.1 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' + 3y' = e^{\frac{x}{2}}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x$2)+3*diff(y(x),x)=exp(x/2),y(x), singsol=all)
```

$$y = -\frac{c_1 e^{-3x}}{3} + \frac{4 e^{\frac{x}{2}}}{7} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.078 (sec). Leaf size: 30

```
DSolve[y''[x]+3*y'[x]==30*Exp[x/2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{120e^{x/2}}{7} - \frac{1}{3}c_1 e^{-3x} + c_2$$

## 15.5 problem 22.2

Internal problem ID [13380]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.2.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 3y' - 10y = -5e^{3x}$$

With initial conditions

$$[y(0) = 5, y'(0) = 3]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 22

```
dsolve([diff(y(x),x$2)-3*diff(y(x),x)-10*y(x)=-5*exp(3*x),y(0) = 5, D(y)(0) = 3],y(x), singularSolutions=false)
```

$$y = \frac{(3e^{7x} + e^{5x} + 6)e^{-2x}}{2}$$

### ✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 28

```
DSolve[{y'[x]-3*y'[x]-10*y[x]==-5*Exp[3*x],{y[0]==5,y'[0]==3}},y[x],x,IncludeSingularSolutions->False]
```

$$y(x) \rightarrow \frac{1}{2}e^{-2x}(e^{5x} + 3e^{7x} + 6)$$

## 15.6 problem 22.3 (a)

Internal problem ID [13381]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.3 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = 10 \cos(2x) + 15 \sin(2x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$2)+9*y(x)=10*cos(2*x)+15*sin(2*x),y(x), singsol=all)
```

$$y = c_2 \sin(3x) + \cos(3x) c_1 + 3 \sin(2x) + 2 \cos(2x)$$

### ✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 32

```
DSolve[y''[x]+9*y[x]==10*Cos[2*x]+15*Sin[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3 \sin(2x) + 2 \cos(2x) + c_1 \cos(3x) + c_2 \sin(3x)$$

## 15.7 problem 22.3 (b)

Internal problem ID [13382]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.3 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 6y' + 9y = 25 \sin(6x)$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$2)-6*diff(y(x),x)+9*y(x)=25*sin(6*x),y(x), singsol=all)
```

$$y = c_2 e^{3x} + x e^{3x} c_1 + \frac{4 \cos(6x)}{9} - \frac{\sin(6x)}{3}$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 35

```
DSolve[y''[x]-6*y'[x]+9*y[x]==25*Sin[6*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{3} \sin(6x) + \frac{4}{9} \cos(6x) + e^{3x}(c_2 x + c_1)$$

## 15.8 problem 22.3 (c)

Internal problem ID [13383]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.3 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' + 3y' = 26 \cos\left(\frac{x}{3}\right) - 12 \sin\left(\frac{x}{3}\right)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)+3*diff(y(x),x)=26*cos(x/3)-12*sin(x/3),y(x), singsol=all)
```

$$y = -\frac{c_1 e^{-3x}}{3} + 27 \sin\left(\frac{x}{3}\right) + 9 \cos\left(\frac{x}{3}\right) + c_2$$

### ✓ Solution by Mathematica

Time used: 0.29 (sec). Leaf size: 35

```
DSolve[y''[x]+3*y'[x]==26*Cos[x/3]-12*Sin[x/3],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 27 \sin\left(\frac{x}{3}\right) + 9 \cos\left(\frac{x}{3}\right) - \frac{1}{3} c_1 e^{-3x} + c_2$$

## 15.9 problem 22.3 (d)

Internal problem ID [13384]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.3 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y' - 5y = \cos(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)-5*y(x)=cos(x),y(x), singsol=all)
```

$$y = e^{-5x}c_2 + c_1e^x - \frac{3 \cos(x)}{26} + \frac{\sin(x)}{13}$$

### ✓ Solution by Mathematica

Time used: 0.066 (sec). Leaf size: 32

```
DSolve[y''[x]+4*y'[x]-5*y[x]==Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sin(x)}{13} - \frac{3 \cos(x)}{26} + c_1e^{-5x} + c_2e^x$$

## 15.10 problem 22.4

Internal problem ID [13385]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.4.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 3y' - 10y = -4 \cos(x) + 7 \sin(x)$$

With initial conditions

$$[y(0) = 8, y'(0) = -5]$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve([diff(y(x),x$2)-3*diff(y(x),x)-10*y(x)=-4*cos(x)+7*sin(x),y(0) = 8, D(y)(0) = -5],y(x)
```

$$y = \frac{((\cos(x) - \sin(x)) e^{2x} + 3 e^{7x} + 12) e^{-2x}}{2}$$

### ✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 30

```
DSolve[{y'[x]-3*y'[x]-10*y[x]==-4*Cos[x]+7*Sin[x]},{y[0]==8,y'[0]==-5}],y[x],x,IncludeSingular
```

$$y(x) \rightarrow \frac{1}{2}(3e^{-2x}(e^{7x} + 4) - \sin(x) + \cos(x))$$



## 15.11 problem 22.5 (a)

Internal problem ID [13386]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.5 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 3y' - 10y = -200$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)-10*y(x)=-200,y(x), singsol=all)
```

$$y = c_2 e^{-2x} + c_1 e^{5x} + 20$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 23

```
DSolve[y''[x]-3*y'[x]-10*y[x]==-200,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-2x} + c_2 e^{5x} + 20$$

## 15.12 problem 22.5 (b)

Internal problem ID [13387]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.5 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y' - 5y = x^3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)-5*y(x)=x^3,y(x), singsol=all)
```

$$y = e^{-5x}c_2 + c_1e^x - \frac{x^3}{5} - \frac{12x^2}{25} - \frac{126x}{125} - \frac{624}{625}$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 39

```
DSolve[y''[x]+4*y'[x]-5*y[x]==x^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{625}(-125x^3 - 300x^2 - 630x - 624) + c_1e^{-5x} + c_2e^x$$

## 15.13 problem 22.5 (c)

Internal problem ID [13388]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.5 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 6y' + 9y = 18x^2 + 3x + 4$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)-6*diff(y(x),x)+9*y(x)=18*x^2+3*x+4,y(x), singsol=all)
```

$$y = c_2 e^{3x} + x e^{3x} c_1 + 2x^2 + 3x + 2$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 32

```
DSolve[y''[x]-6*y'[x]+9*y[x]==18*x^2+3*x+4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2x^2 + x(3 + c_2 e^{3x}) + c_1 e^{3x} + 2$$

## 15.14 problem 22.5 (d)

Internal problem ID [13389]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.5 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = 9x^4 - 9$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)+9*y(x)=9*x^4-9,y(x), singsol=all)
```

$$y = c_2 \sin(3x) + \cos(3x) c_1 + x^4 - \frac{4x^2}{3} - \frac{19}{27}$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 33

```
DSolve[y''[x]+9*y[x]==9*x^4-9,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^4 - \frac{4x^2}{3} + c_1 \cos(3x) + c_2 \sin(3x) - \frac{19}{27}$$

## 15.15 problem 22.6

Internal problem ID [13390]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.6.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = x^3$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve([diff(y(x),x$2)+9*y(x)=x^3,y(0) = 0, D(y)(0) = 0],y(x), singsol=all)
```

$$y = \frac{2 \sin(3x)}{81} + \frac{x^3}{9} - \frac{2x}{27}$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 24

```
DSolve[{y'[x]+9*y[x]==x^3,{y[0]==0,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{81}(9x^3 - 6x + 2 \sin(3x))$$

## 15.16 problem 22.7 (a)

Internal problem ID [13391]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.7 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = 25x \cos(2x)$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$2)+9*y(x)=25*x*cos(2*x),y(x), singsol=all)
```

$$y = c_2 \sin(3x) + \cos(3x) c_1 + 5x \cos(2x) + 4 \sin(2x)$$

### ✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 33

```
DSolve[y''[x]+9*y[x]==25*x*Cos[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 4 \sin(2x) + 5x \cos(2x) + c_1 \cos(3x) + c_2 \sin(3x)$$

## 15.17 problem 22.7 (b)

Internal problem ID [13392]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.7 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 6y' + 9y = \sin(x) e^{2x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)-6*diff(y(x),x)+9*y(x)=exp(2*x)*sin(x),y(x), singsol=all)
```

$$y = c_2 e^{3x} + x e^{3x} c_1 + \frac{e^{2x} \cos(x)}{2}$$

### ✓ Solution by Mathematica

Time used: 0.085 (sec). Leaf size: 29

```
DSolve[y''[x]-6*y'[x]+9*y[x]==Exp[2*x]*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^{2x} (\cos(x) + 2e^x (c_2 x + c_1))$$

## 15.18 problem 22.7 (c)

Internal problem ID [13393]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.7 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = 54x^2e^{3x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)+9*y(x)=54*x^2*exp(3*x),y(x), singsol=all)
```

$$y = c_2 \sin(3x) + \cos(3x) c_1 + 3 \left(x - \frac{1}{3}\right)^2 e^{3x}$$

### ✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 36

```
DSolve[y''[x]+9*y[x]==54*x^2*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}e^{3x}(1 - 3x)^2 + c_1 \cos(3x) + c_2 \sin(3x)$$



## 15.19 problem 22.7 (d)

Internal problem ID [13394]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.7 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$y'' = 6x e^x \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)=6*x*exp(x)*sin(x),y(x), singsol=all)
```

$$y = -3x e^x \cos(x) + 3 e^x \cos(x) + 3 e^x \sin(x) + c_1 x + c_2$$

### ✓ Solution by Mathematica

Time used: 0.104 (sec). Leaf size: 27

```
DSolve[y''[x]==6*x*Exp[x]*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 x + 3e^x(\sin(x) - x \cos(x) + \cos(x)) + c_1$$

## 15.20 problem 22.7 (e)

Internal problem ID [13395]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.7 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 2y' + y = (-6x - 8) \cos(2x) + (8x - 11) \sin(2x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x)=(-6*x-8)*cos(2*x)+(8*x-11)*sin(2*x),y(x), singsol=
```

$$y = e^x c_2 + x e^x c_1 + \sin(2x) + 2x \cos(2x)$$

### ✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 28

```
DSolve[y''[x]-2*y'[x]+y[x]==(-6*x-8)*Cos[2*x]+(8*x-11)*Sin[2*x],y[x],x,IncludeSingularSoluti
```

$$y(x) \rightarrow \sin(2x) + 2x \cos(2x) + e^x(c_2 x + c_1)$$

## 15.21 problem 22.7 (f)

Internal problem ID [13396]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.7 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 2y' + y = (12x - 4)e^{-5x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x)=(12*x-4)*exp(-5*x),y(x), singsol=all)
```

$$y = e^x c_2 + x e^x c_1 + \frac{e^{-5x} x}{3}$$

### ✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 27

```
DSolve[y''[x]-2*y'[x]+y[x]==(12*x-4)*Exp[-5*x],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{3}e^{-5x}x + e^x(c_2x + c_1)$$

## 15.22 problem 22.8

Internal problem ID [13397]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.8.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = 39e^{2x}x$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 26

```
dsolve([diff(y(x),x$2)+9*y(x)=39*x*exp(2*x),y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y = 3xe^{2x} - \frac{5 \sin(3x)}{13} + \frac{25 \cos(3x)}{13} - \frac{12e^{2x}}{13}$$

### ✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 34

```
DSolve[{y'[x]+9*y[x]==39*x*Exp[2*x],{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow \frac{1}{13}(3e^{2x}(13x - 4) - 5 \sin(3x) + 25 \cos(3x))$$

## 15.23 problem 22.9 (a)

Internal problem ID [13398]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.9 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 3y' - 10y = -3e^{-2x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)-10*y(x)=-3*exp(-2*x),y(x), singsol=all)
```

$$y = c_2 e^{-2x} + c_1 e^{5x} + \frac{3x e^{-2x}}{7}$$

### ✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 32

```
DSolve[y''[x]-3*y'[x]-10*y[x]==-3*Exp[-2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{49} e^{-2x} (21x + 49c_2 e^{7x} + 3 + 49c_1)$$

## 15.24 problem 22.9 (b)

Internal problem ID [13399]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.9 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 4y' = 20$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)=20,y(x), singsol=all)
```

$$y = -\frac{e^{-4x}c_1}{4} + 5x + c_2$$

### ✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 22

```
DSolve[y''[x]+4*y'[x]==20,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 5x - \frac{1}{4}c_1e^{-4x} + c_2$$

## 15.25 problem 22.9 (c)

Internal problem ID [13400]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.9 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' + 4y' = x^2$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)=x^2,y(x), singsol=all)
```

$$y = -\frac{x^2}{16} + \frac{x^3}{12} - \frac{e^{-4x}c_1}{4} + \frac{x}{32} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.074 (sec). Leaf size: 36

```
DSolve[y''[x]+4*y'[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{96}(8x^3 - 6x^2 + 3x - 24c_1e^{-4x} + 96c_2)$$

## 15.26 problem 22.9 (d)

Internal problem ID [13401]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.9 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = 3 \sin(3x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)+9*y(x)=3*sin(3*x),y(x), singsol=all)
```

$$y = c_2 \sin(3x) + \cos(3x) c_1 - \frac{\cos(3x) x}{2}$$

### ✓ Solution by Mathematica

Time used: 0.108 (sec). Leaf size: 33

```
DSolve[y''[x]+9*y[x]==3*Sin[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left(-\frac{x}{2} + c_1\right) \cos(3x) + \frac{1}{12}(1 + 12c_2) \sin(3x)$$



## 15.27 problem 22.9 (e)

Internal problem ID [13402]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.9 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 6y' + 9y = 10e^{3x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)-6*diff(y(x),x)+9*y(x)=10*exp(3*x),y(x), singsol=all)
```

$$y = c_2 e^{3x} + x e^{3x} c_1 + 5x^2 e^{3x}$$

### ✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 23

```
DSolve[y''[x]-6*y'[x]+9*y[x]==10*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{3x} (5x^2 + c_2 x + c_1)$$

## 15.28 problem 22.10 (a)

Internal problem ID [13403]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.10 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _linear, _nonhomogeneous]`

$$y'' - 3y' - 10y = (72x^2 - 1)e^{2x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)-10*y(x)=(72*x^2-1)*exp(2*x),y(x), singsol=all)
```

$$y = c_2e^{-2x} + c_1e^{5x} + e^{2x}(-6x^2 - x - 1)$$

### ✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 37

```
DSolve[y''[x]-3*y'[x]-10*y[x]==(72*x^2-1)*Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -e^{2x}(6x^2 + x + 1) + c_1e^{-2x} + c_2e^{5x}$$

## 15.29 problem 22.10 (b)

Internal problem ID [13404]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.10 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 3y' - 10y = 4x e^{6x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)-10*y(x)=4*x*exp(6*x),y(x), singsol=all)
```

$$y = c_2 e^{-2x} + c_1 e^{5x} + \frac{(8x - 9) e^{6x}}{16}$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 36

```
DSolve[y''[x]-3*y'[x]-10*y[x]==4*x*Exp[6*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{16} e^{6x} (8x - 9) + c_1 e^{-2x} + c_2 e^{5x}$$

## 15.30 problem 22.10 (c)

Internal problem ID [13405]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.10 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 10y' + 25y = 6e^{5x}$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)-10*diff(y(x),x)+25*y(x)=6*exp(5*x),y(x), singsol=all)
```

$$y = c_2 e^{5x} + x e^{5x} c_1 + 3x^2 e^{5x}$$

### ✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 23

```
DSolve[y''[x]-10*y'[x]+25*y[x]==6*Exp[5*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{5x} (3x^2 + c_2 x + c_1)$$

### 15.31 problem 22.10 (d)

Internal problem ID [13406]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.10 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 10y' + 25y = 6e^{-5x}$$

#### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)-10*diff(y(x),x)+25*y(x)=6*exp(-5*x),y(x), singsol=all)
```

$$y = c_2 e^{5x} + x e^{5x} c_1 + \frac{3e^{-5x}}{50}$$

#### ✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 28

```
DSolve[y''[x]-10*y'[x]+25*y[x]==6*Exp[-5*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3e^{-5x}}{50} + e^{5x}(c_2x + c_1)$$

## 15.32 problem 22.10 (e)

Internal problem ID [13407]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.10 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y' + 5y = 24 \sin(3x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)+5*y(x)=24*sin(3*x),y(x), singsol=all)
```

$$y = e^{-2x} \sin(x) c_2 + e^{-2x} \cos(x) c_1 - \frac{3 \sin(3x)}{5} - \frac{9 \cos(3x)}{5}$$

### ✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 41

```
DSolve[y''[x]+4*y'[x]+5*y[x]==24*Sin[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{3}{5}(\sin(3x) + 3 \cos(3x)) + c_2 e^{-2x} \cos(x) + c_1 e^{-2x} \sin(x)$$

### 15.33 problem 22.10 (f)

Internal problem ID [13408]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.10 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 4y' + 5y = 8e^{-3x}$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)+5*y(x)=8*exp(-3*x),y(x), singsol=all)
```

$$y = e^{-2x} \sin(x) c_2 + e^{-2x} \cos(x) c_1 + 4e^{-3x}$$

#### ✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 29

```
DSolve[y''[x]+4*y'[x]+5*y[x]==8*Exp[-3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x}(c_2 e^x \cos(x) + c_1 e^x \sin(x) + 4)$$

## 15.34 problem 22.10 (g)

Internal problem ID [13409]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.10 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y' + 5y = \sin(x) e^{2x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+5*y(x)=exp(2*x)*sin(x),y(x), singsol=all)
```

$$y = e^{2x} \sin(x) c_2 + e^{2x} \cos(x) c_1 + \frac{e^{2x}(\sin(x) - \cos(x)x)}{2}$$

### ✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 30

```
DSolve[y''[x]-4*y'[x]+5*y[x]==Exp[2*x]*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2}e^{2x}((x - 2c_2) \cos(x) - 2c_1 \sin(x))$$



## 15.35 problem 22.10 (h)

Internal problem ID [13410]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.10 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y' + 5y = e^{-x} \sin(x)$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 36

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+5*y(x)=exp(-x)*sin(x),y(x), singsol=all)
```

$$y = e^{2x} \sin(x) c_2 + e^{2x} \cos(x) c_1 + \frac{e^{-x}(3 \sin(x) + 2 \cos(x))}{39}$$

### ✓ Solution by Mathematica

Time used: 0.16 (sec). Leaf size: 44

```
DSolve[y''[x]-4*y'[x]+5*y[x]==Exp[-x]*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{78} e^{-x} ((4 + 78c_2 e^{3x}) \cos(x) + 6(1 + 13c_1 e^{3x}) \sin(x))$$

## 15.36 problem 22.10 (i)

Internal problem ID [13411]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.10 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y' + 5y = 100$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+5*y(x)=100,y(x), singsol=all)
```

$$y = e^{2x} \sin(x) c_2 + e^{2x} \cos(x) c_1 + 20$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 27

```
DSolve[y''[x]-4*y'[x]+5*y[x]==100,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 e^{2x} \cos(x) + c_1 e^{2x} \sin(x) + 20$$

## 15.37 problem 22.10 (j)

Internal problem ID [13412]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.10 (j).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 4y' + 5y = e^{-x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+5*y(x)=exp(-x),y(x), singsol=all)
```

$$y = e^{2x} \sin(x) c_2 + e^{2x} \cos(x) c_1 + \frac{e^{-x}}{10}$$

### ✓ Solution by Mathematica

Time used: 0.067 (sec). Leaf size: 35

```
DSolve[y''[x]-4*y'[x]+5*y[x]==Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{-x}}{10} + c_2 e^{2x} \cos(x) + c_1 e^{2x} \sin(x)$$

## 15.38 problem 22.10 (k)

Internal problem ID [13413]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.10 (k).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 4y' + 5y = 10x^2 + 4x + 8$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+5*y(x)=10*x^2+4*x+8,y(x), singsol=all)
```

$$y = e^{2x} \sin(x) c_2 + e^{2x} \cos(x) c_1 + 2x^2 + 4x + 4$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 35

```
DSolve[y''[x]-4*y'[x]+5*y[x]==10*x^2+4*x+8,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2x^2 + 4x + c_2 e^{2x} \cos(x) + c_1 e^{2x} \sin(x) + 4$$

## 15.39 problem 22.10 (L)

Internal problem ID [13414]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.10 (L).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = e^{2x} \sin(x)$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$2)+9*y(x)=exp(2*x)*sin(x),y(x), singsol=all)
```

$$y = c_2 \sin(3x) + c_1 \cos(3x) - \frac{(\cos(x) - 3 \sin(x)) e^{2x}}{40}$$

### ✓ Solution by Mathematica

Time used: 0.255 (sec). Leaf size: 42

```
DSolve[y''[x]+9*y[x]==Exp[2*x]*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3}{40} e^{2x} \sin(x) - \frac{1}{40} e^{2x} \cos(x) + c_1 \cos(3x) + c_2 \sin(3x)$$

## 15.40 problem 22.10 (m)

Internal problem ID [13415]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.10 (m).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = 6 \cos(x) - 3 \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)+y(x)=6*cos(x)-3*sin(x),y(x), singsol=all)
```

$$y = c_2 \sin(x) + c_1 \cos(x) + 3 \cos(x) + 3x \sin(x) + \frac{3x \cos(x)}{2}$$

### ✓ Solution by Mathematica

Time used: 0.096 (sec). Leaf size: 27

```
DSolve[y''[x]+y[x]==6*Cos[x]-3*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left( \frac{3x}{2} + 3 + c_1 \right) \cos(x) + (3x + c_2) \sin(x)$$

## 15.41 problem 22.10 (n)

Internal problem ID [13416]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.10 (n).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = 6 \cos(2x) - 3 \sin(2x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)+y(x)=6*cos(2*x)-3*sin(2*x),y(x), singsol=all)
```

$$y = c_1 \cos(x) + c_2 \sin(x) - 2 \cos(2x) + \sin(2x)$$

### ✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 26

```
DSolve[y''[x]+y[x]==6*Cos[2*x]-3*Sin[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sin(2x) - 2 \cos(2x) + c_1 \cos(x) + c_2 \sin(x)$$

## 15.42 problem 22.11 (a)

Internal problem ID [13417]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.11 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y' + 5y = x^3 e^{-x} \sin(x)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 61

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+5*y(x)=x^3*exp(-x)*sin(x),y(x), singsol=all)
```

$$y = e^{2x} \sin(x) c_2 + e^{2x} \cos(x) c_1 + \frac{2 e^{-x} \left( \left( x^3 + \frac{31}{13} x^2 + \frac{1220}{507} x + \frac{19100}{19773} \right) \cos(x) + \frac{3 \left( x^3 + \frac{18}{13} x^2 + \frac{138}{169} x + \frac{360}{2197} \right) \sin(x)}{2} \right)}{39}$$

✓ Solution by Mathematica

Time used: 0.051 (sec). Leaf size: 70

```
DSolve[y''[x]-4*y'[x]+5*y[x]==x^3*Exp[-x]*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{-x} \left( (39546x^3 + 94302x^2 + 95160x + 771147c_2 e^{3x} + 38200) \cos(x) + 27(2197x^3 + 3042x^2 + 1794x + 28) \sin(x) \right)}{771147}$$



## 15.43 problem 22.11 (b)

Internal problem ID [13418]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.11 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y' + 5y = x^3 e^{2x} \sin(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+5*y(x)=x^3*exp(2*x)*sin(x),y(x), singsol=all)
```

$$y = e^{2x} \sin(x) c_2 + e^{2x} \cos(x) c_1 - \frac{e^{2x}((x^3 - 3x) \cos(x) + (-2x^2 + 3) \sin(x)) x}{8}$$

### ✓ Solution by Mathematica

Time used: 0.116 (sec). Leaf size: 51

```
DSolve[y''[x]-4*y'[x]+5*y[x]==x^3*Exp[2*x]*Sin[x],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{16} e^{2x} (2(2x^3 - 3x + 8c_1) \sin(x) + (-2x^4 + 6x^2 - 3 + 16c_2) \cos(x))$$

## 15.44 problem 22.11 (c)

Internal problem ID [13419]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.11 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 5y' + 6y = x^2e^{-7x} + 2e^{-7x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 33

```
dsolve(diff(y(x),x$2)-5*diff(y(x),x)+6*y(x)=x^2*exp(-7*x)+2*exp(-7*x),y(x), singsol=all)
```

$$y = c_2e^{2x} + c_1e^{3x} + \frac{(4050x^2 + 1710x + 8371)e^{-7x}}{364500}$$

### ✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 41

```
DSolve[y''[x]-5*y'[x]+6*y[x]==x^2*Exp[-7*x]+2*Exp[-7*x],y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow \frac{e^{-7x}(4050x^2 + 1710x + 8371)}{364500} + c_1e^{2x} + c_2e^{3x}$$

## 15.45 problem 22.11 (d)

Internal problem ID [13420]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.11 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 5y' + 6y = x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)-5*diff(y(x),x)+6*y(x)=x^2,y(x), singsol=all)
```

$$y = c_2 e^{2x} + c_1 e^{3x} + \frac{x^2}{6} + \frac{5x}{18} + \frac{19}{108}$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 37

```
DSolve[y''[x]-5*y'[x]+6*y[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2}{6} + \frac{5x}{18} + c_1 e^{2x} + c_2 e^{3x} + \frac{19}{108}$$

## 15.46 problem 22.11 (e)

Internal problem ID [13421]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.11 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 5y' + 6y = 4e^{-8x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)-5*diff(y(x),x)+6*y(x)=4*exp(-8*x),y(x), singsol=all)
```

$$y = c_2 e^{2x} + c_1 e^{3x} + \frac{2e^{-8x}}{55}$$

### ✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 31

```
DSolve[y''[x]-5*y'[x]+6*y[x]==4*Exp[-8*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2e^{-8x}}{55} + c_1 e^{2x} + c_2 e^{3x}$$

## 15.47 problem 22.11 (f)

Internal problem ID [13422]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.11 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 5y' + 6y = 4e^{3x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)-5*diff(y(x),x)+6*y(x)=4*exp(3*x),y(x), singsol=all)
```

$$y = c_2 e^{2x} + c_1 e^{3x} + 4x e^{3x}$$

### ✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 25

```
DSolve[y''[x]-5*y'[x]+6*y[x]==4*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(e^x(4x - 4 + c_2) + c_1)$$

## 15.48 problem 22.11 (g)

Internal problem ID [13423]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.11 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 5y' + 6y = x^2 e^{3x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

```
dsolve(diff(y(x),x$2)-5*diff(y(x),x)+6*y(x)=x^2*exp(3*x),y(x), singsol=all)
```

$$y = c_2 e^{2x} + c_1 e^{3x} + \frac{x(x^2 - 3x + 6) e^{3x}}{3}$$

### ✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 40

```
DSolve[y''[x]-5*y'[x]+6*y[x]==x^2*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} e^{2x} (e^x (x^3 - 3x^2 + 6x - 6 + 3c_2) + 3c_1)$$

## 15.49 problem 22.11 (h)

Internal problem ID [13424]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.11 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 5y' + 6y = x^2 \cos(2x)$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 49

```
dsolve(diff(y(x),x$2)-5*diff(y(x),x)+6*y(x)=x^2*cos(2*x),y(x), singsol=all)
```

$$y = c_2 e^{2x} + c_1 e^{3x} + \frac{(676x^2 - 2080x - 1909) \cos(2x)}{35152} + \frac{(-3380x^2 - 3796x - 725) \sin(2x)}{35152}$$

### ✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 58

```
DSolve[y''[x]-5*y'[x]+6*y[x]==x^2*Cos[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{(676x^2 - 2080x - 1909) \cos(2x) - (3380x^2 + 3796x + 725) \sin(2x)}{35152} + c_1 e^{2x} + c_2 e^{3x}$$

## 15.50 problem 22.11 (i)

Internal problem ID [13425]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.11 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 5y' + 6y = x^2 e^{3x} \sin(2x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 51

```
dsolve(diff(y(x),x$2)-5*diff(y(x),x)+6*y(x)=x^2*exp(3*x)*sin(2*x),y(x), singsol=all)
```

$$y = c_2 e^{2x} + c_1 e^{3x} - \frac{e^{3x} \left( \left( x^2 + \frac{16}{5}x - \frac{109}{50} \right) \cos(2x) + 2 \sin(2x) \left( x^2 - \frac{13}{10}x - \frac{22}{25} \right) \right)}{10}$$

### ✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 63

```
DSolve[y''[x]-5*y'[x]+6*y[x]==x^2*Exp[3*x]*Sin[2*x],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow -\frac{1}{500} e^{3x} (2(50x^2 - 65x - 44) \sin(2x) + (50x^2 + 160x - 109) \cos(2x)) + c_1 e^{2x} + c_2 e^{3x}$$



## 15.51 problem 22.11 (j)

Internal problem ID [13426]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.11 (j).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y' + 20y = e^{4x} \sin(2x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 42

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+20*y(x)=exp(4*x)*sin(2*x),y(x), singsol=all)
```

$$y = e^{2x} \sin(4x) c_2 + e^{2x} \cos(4x) c_1 - \frac{(\cos(2x) - 2 \sin(2x)) e^{4x}}{40}$$

### ✓ Solution by Mathematica

Time used: 0.224 (sec). Leaf size: 54

```
DSolve[y''[x]-4*y'[x]+20*y[x]==Exp[4*x]*Sin[2*x],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow -\frac{1}{40} e^{2x} (e^{2x} \cos(2x) - 2(e^{2x} \sin(2x) + 20c_2 \cos(4x) + 20c_1 \sin(4x)))$$

## 15.52 problem 22.11 (k)

Internal problem ID [13427]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.11 (k).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y' + 20y = e^{2x} \sin(4x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+20*y(x)=exp(2*x)*sin(4*x),y(x), singsol=all)
```

$$y = e^{2x} \sin(4x) c_2 + e^{2x} \cos(4x) c_1 - \frac{e^{2x} \cos(4x) x}{8}$$

### ✓ Solution by Mathematica

Time used: 0.116 (sec). Leaf size: 38

```
DSolve[y''[x]-4*y'[x]+20*y[x]==Exp[2*x]*Sin[4*x],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{64} e^{2x} ((1 + 64c_1) \sin(4x) - 8(x - 8c_2) \cos(4x))$$

## 15.53 problem 22.11 (L)

Internal problem ID [13428]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.11 (L).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y' + 20y = x^3 \sin(4x)$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 65

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+20*y(x)=x^3*sin(4*x),y(x), singsol=all)
```

$$y = e^{2x} \sin(4x) c_2 + e^{2x} \cos(4x) c_1 + \frac{(9826x^3 + 16473x^2 + 15810x + 7815) \cos(4x)}{167042} + \frac{(x^3 + \frac{3}{17}x^2 - \frac{39}{578}x - \frac{45}{4913}) \sin(4x)}{68}$$

### ✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 76

```
DSolve[y''[x]-4*y'[x]+20*y[x]==x^3*Sin[4*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{(9826x^3 + 1734x^2 - 663x - 90) \sin(4x) + 4(9826x^3 + 16473x^2 + 15810x + 7815) \cos(4x)}{668168} + c_2 e^{2x} \cos(4x) + c_1 e^{2x} \sin(4x)$$

## 15.54 problem 22.11 (m)

Internal problem ID [13429]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.11 (m).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 10y' + 25y = 3x^2e^{5x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)-10*diff(y(x),x)+25*y(x)=3*x^2*exp(5*x),y(x), singsol=all)
```

$$y = c_2e^{5x} + xe^{5x}c_1 + \frac{x^4e^{5x}}{4}$$

### ✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 27

```
DSolve[y''[x]-10*y'[x]+25*y[x]==3*x^2*Exp[5*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}e^{5x}(x^4 + 4c_2x + 4c_1)$$

## 15.55 problem 22.11 (n)

Internal problem ID [13430]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.11 (n).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 10y' + 25y = 3x^4$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 37

```
dsolve(diff(y(x),x$2)-10*diff(y(x),x)+25*y(x)=3*x^4,y(x), singsol=all)
```

$$y = c_2 e^{5x} + x e^{5x} c_1 + \frac{3x^4}{25} + \frac{24x^3}{125} + \frac{108x^2}{625} + \frac{288x}{3125} + \frac{72}{3125}$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 47

```
DSolve[y''[x]-10*y'[x]+25*y[x]==3*x^4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3(125x^4 + 200x^3 + 180x^2 + 96x + 24)}{3125} + c_1 e^{5x} + c_2 e^{5x} x$$

## 15.56 problem 22.12 (a)

Internal problem ID [13431]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.12 (a).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_y]]`

$$y'''' - 4y''' = 12e^{-2x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$4)-4*diff(y(x),x$3)=12*exp(-2*x),y(x), singsol=all)
```

$$y = \frac{c_1 e^{4x}}{64} + \frac{c_2 x^2}{2} + \frac{e^{-2x}}{4} + c_3 x + c_4$$

### ✓ Solution by Mathematica

Time used: 0.152 (sec). Leaf size: 37

```
DSolve[y''''[x]-4*y'''[x]==12*Exp[-2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{-2x}}{4} + \frac{1}{64}c_1 e^{4x} + x(c_4 x + c_3) + c_2$$

## 15.57 problem 22.12 (b)

Internal problem ID [13432]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.12 (b).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_y]]`

$$y'''' - 4y''' = 10 \sin(2x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$4)-4*diff(y(x),x$3)=10*sin(2*x),y(x), singsol=all)
```

$$y = \frac{c_1 e^{4x}}{64} + \frac{c_2 x^2}{2} + \frac{\sin(2x)}{8} - \frac{\cos(2x)}{4} + c_3 x + c_4$$

### ✓ Solution by Mathematica

Time used: 0.383 (sec). Leaf size: 46

```
DSolve[y''''[x]-4*y'''[x]==10*Sin[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_4 x^2 + c_3 x + \frac{1}{64}(16x + 8 \sin(2x) - 16 \cos(2x) + c_1 e^{4x}) + c_2$$

## 15.58 problem 22.12 (c)

Internal problem ID [13433]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.12 (c).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_y]]`

$$y'''' - 4y''' = 32e^{4x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$4)-4*diff(y(x),x$3)=32*exp(4*x),y(x), singsol=all)
```

$$y = \frac{c_1 e^{4x}}{64} + \frac{e^{4x} x}{2} - \frac{3e^{4x}}{8} + \frac{c_2 x^2}{2} + c_3 x + c_4$$

### ✓ Solution by Mathematica

Time used: 0.137 (sec). Leaf size: 33

```
DSolve[y''''[x]-4*y'''[x]==32*Exp[4*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{64} e^{4x} (32x - 24 + c_1) + x(c_4 x + c_3) + c_2$$



## 15.59 problem 22.12 (d)

Internal problem ID [13434]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.12 (d).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_y]]`

$$y'''' - 4y''' = 32x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
dsolve(diff(y(x),x$4)-4*diff(y(x),x$3)=32*x,y(x), singsol=all)
```

$$y = -\frac{x^4}{3} - \frac{x^3}{3} + \frac{c_1 e^{4x}}{64} + \frac{c_2 x^2}{2} + c_3 x + c_4$$

### ✓ Solution by Mathematica

Time used: 0.079 (sec). Leaf size: 43

```
DSolve[y''''[x]-4*y'''[x]==32*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x^4}{3} - \frac{x^3}{3} + c_4 x^2 + c_3 x + \frac{1}{64} c_1 e^{4x} + c_2$$

## 15.60 problem 22.12 (e)

Internal problem ID [13435]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.12 (e).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$y''' - y'' + y' - y = x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$3)-diff(y(x),x$2)+diff(y(x),x)-y(x)=x^2,y(x), singsol=all)
```

$$y = -x^2 - 2x + c_1 \cos(x) + c_2 e^x + c_3 \sin(x)$$

### ✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 30

```
DSolve[y'''[x]-y''[x]+y'[x]-y[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x^2 - 2x + c_3 e^x + c_1 \cos(x) + c_2 \sin(x)$$

## 15.61 problem 22.12 (f)

Internal problem ID [13436]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.12 (f).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - y'' + y' - y = 30 \cos(2x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$3)-diff(y(x),x$2)+diff(y(x),x)-y(x)=30*cos(2*x),y(x), singsol=all)
```

$$y = 2 \cos(2x) - 4 \sin(2x) + c_1 \cos(x) + c_2 e^x + c_3 \sin(x)$$

### ✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 34

```
DSolve[y'''[x]-y''[x]+y'[x]-y[x]==30*Cos[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -4 \sin(2x) + 2 \cos(2x) + c_3 e^x + c_1 \cos(x) + c_2 \sin(x)$$

## 15.62 problem 22.12 (g)

Internal problem ID [13437]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.12 (g).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$y''' - y'' + y' - y = 6e^x$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$3)-diff(y(x),x$2)+diff(y(x),x)-y(x)=6*exp(x),y(x), singsol=all)
```

$$y = 3e^x x + c_1 \cos(x) + c_2 e^x + c_3 \sin(x)$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 27

```
DSolve[y'''[x]-y''[x]+y'[x]-y[x]==6*Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(3x - 3 + c_3) + c_1 \cos(x) + c_2 \sin(x)$$

## 15.63 problem 22.13 (a)

Internal problem ID [13438]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.13 (a).

**ODE order:** 5.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_y]]`

$$y^{(5)} + 18y''' + 81y' = x^2 e^{3x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 74

```
dsolve(diff(y(x),x$5)+18*diff(y(x),x$3)+81*diff(y(x),x)=x^2*exp(3*x),y(x), singsol=all)
```

$$y = -\frac{x e^{3x}}{486} + \frac{5 e^{3x}}{4374} + \frac{x^2 e^{3x}}{972} + \frac{c_1 \sin(3x)}{3} - \frac{c_2 \cos(3x)}{3} \\ + c_3 \left( \frac{\cos(3x)}{9} + \frac{x \sin(3x)}{3} \right) + c_4 \left( \frac{\sin(3x)}{9} - \frac{x \cos(3x)}{3} \right) + c_5$$

### ✓ Solution by Mathematica

Time used: 0.497 (sec). Leaf size: 67

```
DSolve[y'''''[x]+18*y'''[x]+81*y'[x]==x^2*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{3x}(9x^2 - 18x + 10)}{8748} + \frac{1}{9}(c_2 - 3(c_4x + c_3)) \cos(3x) + \frac{1}{9}(3c_2x + 3c_1 + c_4) \sin(3x) + c_5$$

## 15.64 problem 22.13 (b)

Internal problem ID [13439]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.13 (b).

**ODE order:** 5.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_y]]`

$$y^{(5)} + 18y''' + 81y' = x^2 \sin(3x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 92

```
dsolve(diff(y(x),x$5)+18*diff(y(x),x$3)+81*diff(y(x),x)=x^2*sin(3*x),y(x), singsol=all)
```

$$y = -\frac{11x^2 \cos(3x)}{3888} + \frac{19 \cos(3x)}{23328} + \frac{13x \sin(3x)}{5832} - \frac{\sin(3x) x^3}{486} + \frac{c_1 \sin(3x)}{3} - \frac{c_2 \cos(3x)}{3} \\ + \frac{\cos(3x) x^4}{1296} + c_3 \left( \frac{\cos(3x)}{9} + \frac{x \sin(3x)}{3} \right) + c_4 \left( \frac{\sin(3x)}{9} - \frac{x \cos(3x)}{3} \right) + c_5$$

### ✓ Solution by Mathematica

Time used: 0.411 (sec). Leaf size: 72

```
DSolve[y'''''[x]+18*y'''[x]+81*y'[x]==x^2*Sin[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{4(-12x^3 + (13 + 1944c_2)x + 648(3c_1 + c_4)) \sin(3x) + (18x^4 - 66x^2 - 7776c_4x + 19 + 2592c_2 - 7776c_3}{23328} + c_5$$

## 15.65 problem 22.13 (c)

Internal problem ID [13440]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.13 (c).

**ODE order:** 5.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_y]]`

$$y^{(5)} + 18y''' + 81y' = x^2 e^{3x} \sin(3x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 118

```
dsolve(diff(y(x),x$5)+18*diff(y(x),x$3)+81*diff(y(x),x)=x^2*exp(3*x)*sin(3*x),y(x), singsol=
```

$$y = c_5 - \frac{31x e^{3x} \cos(3x)}{91125} + \frac{4x e^{3x} \sin(3x)}{3375} + \frac{139 e^{3x} \cos(3x)}{303750} - \frac{1693 e^{3x} \sin(3x)}{2733750} + \frac{c_3 x \sin(3x)}{3} - \frac{c_4 x \cos(3x)}{3} - \frac{x^2 e^{3x} \cos(3x)}{12150} - \frac{7x^2 e^{3x} \sin(3x)}{12150} + \frac{c_1 \sin(3x)}{3} - \frac{c_2 \cos(3x)}{3} + \frac{\cos(3x) c_3}{9} + \frac{c_4 \sin(3x)}{9}$$

### ✓ Solution by Mathematica

Time used: 0.792 (sec). Leaf size: 94

```
DSolve[y'''''[x]+18*y'''[x]+81*y'[x]==x^2*Exp[3*x]*Sin[3*x],y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow -\frac{e^{3x}(1575x^2 - 3240x + 1693) \sin(3x)}{2733750} - \frac{e^{3x}(75x^2 + 310x - 417) \cos(3x)}{911250} + \frac{1}{9}(c_2 - 3(c_4x + c_3)) \cos(3x) + \frac{1}{9}(3c_2x + 3c_1 + c_4) \sin(3x) + c_5$$

## 15.66 problem 22.13 (d)

Internal problem ID [13441]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.13 (d).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - y'' + y' - y = 30 \cos(2x) x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 37

```
dsolve(diff(y(x),x$3)-diff(y(x),x$2)+diff(y(x),x)-y(x)=30*x*cos(2*x),y(x), singsol=all)
```

$$y = \left(-\frac{98}{15} + 2x\right) \cos(2x) + \left(-4x - \frac{64}{15}\right) \sin(2x) + c_1 \cos(x) + c_2 e^x + c_3 \sin(x)$$

### ✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 49

```
DSolve[y'''[x]-y''[x]+y'[x]-y[x]==30*x*Cos[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -4x \sin(2x) - \frac{64}{15} \sin(2x) + \left(2x - \frac{98}{15}\right) \cos(2x) + c_3 e^x + c_1 \cos(x) + c_2 \sin(x)$$



## 15.67 problem 22.13 (e)

Internal problem ID [13442]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.13 (e).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - y'' + y' - y = 3 \cos(x) x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 42

```
dsolve(diff(y(x),x$3)-diff(y(x),x$2)+diff(y(x),x)-y(x)=3*x*cos(x),y(x), singsol=all)
```

$$y = \left(-\frac{3}{8}x^2 - \frac{9}{8}x\right) \cos(x) + \left(\frac{9}{8} + \frac{3}{8}x - \frac{3}{8}x^2\right) \sin(x) + c_1 \cos(x) + c_2 e^x + c_3 \sin(x)$$

### ✓ Solution by Mathematica

Time used: 0.046 (sec). Leaf size: 52

```
DSolve[y''''[x]-y'''[x]+y''[x]-y[x]==3*x*Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{16} \left( -(6x^2 + 18x + 3 - 16c_1) \cos(x) + (-6x^2 + 6x + 15 + 16c_2) \sin(x) + 16c_3 e^x \right)$$

## 15.68 problem 22.13 (f)

Internal problem ID [13443]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.13 (f).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - y'' + y' - y = 3 \cos(x) x e^x$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
dsolve(diff(y(x),x$3)-diff(y(x),x$2)+diff(y(x),x)-y(x)=3*x*exp(x)*cos(x),y(x), singsol=all)
```

$$y = -\frac{3 e^x (10x - 19) \cos(x)}{25} + \frac{3 e^x (5x + 8) \sin(x)}{25} + c_1 \cos(x) + c_2 e^x + c_3 \sin(x)$$

### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 49

```
DSolve[y'''[x]-y''[x]+y'[x]-y[x]==3*x*Exp[x]*Cos[x],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow c_3 e^x + \left( e^x \left( \frac{57}{25} - \frac{6x}{5} \right) + c_1 \right) \cos(x) + \left( \frac{3}{25} e^x (5x + 8) + c_2 \right) \sin(x)$$

## 15.69 problem 22.13 (g)

Internal problem ID [13444]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.13 (g).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - y'' + y' - y = 5x^5 e^{2x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 48

```
dsolve(diff(y(x),x$3)-diff(y(x),x$2)+diff(y(x),x)-y(x)=5*x^5*exp(2*x),y(x), singsol=all)
```

$$y = \frac{(625x^5 - 5625x^4 + 28000x^3 - 91200x^2 + 187320x - 188376) e^{2x}}{625} + c_1 \cos(x) + c_2 e^x + c_3 \sin(x)$$

### ✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 58

```
DSolve[y''''[x]-y'''[x]+y''[x]-y[x]==5*x^5*Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x \left( e^x \left( x^5 - 9x^4 + \frac{224x^3}{5} - \frac{3648x^2}{25} + \frac{37464x}{125} - \frac{188376}{625} \right) + c_3 \right) + c_1 \cos(x) + c_2 \sin(x)$$

## 15.70 problem 22.14 (a)

Internal problem ID [13445]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.14 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 6y' + 9y = 27e^{6x} + 25 \sin(6x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
dsolve(diff(y(x),x$2)-6*diff(y(x),x)+9*y(x)=27*exp(6*x)+25*sin(6*x),y(x), singsol=all)
```

$$y = c_2 e^{3x} + x e^{3x} c_1 + 3 e^{6x} + \frac{4 \cos(6x)}{9} - \frac{\sin(6x)}{3}$$

### ✓ Solution by Mathematica

Time used: 0.409 (sec). Leaf size: 42

```
DSolve[y''[x]-6*y'[x]+9*y[x]==27*Exp[6*x]+25*Sin[6*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{3} \sin(6x) + \frac{4}{9} \cos(6x) + e^{3x} (3e^{3x} + c_2 x + c_1)$$

## 15.71 problem 22.14 (b)

Internal problem ID [13446]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.14 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = 25x \cos(2x) + 3 \sin(3x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 43

```
dsolve(diff(y(x),x$2)+9*y(x)=25*x*cos(2*x)+3*sin(3*x),y(x), singsol=all)
```

$$y = c_2 \sin(3x) + c_1 \cos(3x) + 4 \sin(2x) + 5x \cos(2x) + \frac{\sin(3x)}{12} - \frac{x \cos(3x)}{2}$$

### ✓ Solution by Mathematica

Time used: 0.274 (sec). Leaf size: 39

```
DSolve[y''[x]+9*y[x]==25*x*Cos[2*x]+3*Sin[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 4 \sin(2x) + 5x \cos(2x) + \left(-\frac{x}{2} + c_1\right) \cos(3x) + c_2 \sin(3x)$$

## 15.72 problem 22.14 (c)

Internal problem ID [13447]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.14 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y' + 5y = 5 \sin(x)^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+5*y(x)=5*sin(x)^2,y(x), singsol=all)
```

$$y = e^{2x} \sin(x) c_2 + e^{2x} \cos(x) c_1 - \frac{\cos(2x)}{26} + \frac{4 \sin(2x)}{13} + \frac{1}{2}$$

### ✓ Solution by Mathematica

Time used: 0.168 (sec). Leaf size: 45

```
DSolve[y''[x]-4*y'[x]+5*y[x]==5*Sin[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{4}{13} \sin(2x) - \frac{1}{26} \cos(2x) + c_2 e^{2x} \cos(x) + c_1 e^{2x} \sin(x) + \frac{1}{2}$$

## 15.73 problem 22.14 (d)

Internal problem ID [13448]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.14 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y' + 5y = 20 \sinh(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+5*y(x)=20*sinh(x),y(x), singsol=all)
```

$$y = e^{2x} \sin(x) c_2 + e^{2x} \cos(x) c_1 - e^{-x} + 5 e^x$$

### ✓ Solution by Mathematica

Time used: 0.105 (sec). Leaf size: 38

```
DSolve[y''[x]-4*y'[x]+5*y[x]==20*Sinh[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -e^{-x} + 5e^x + c_2 e^{2x} \cos(x) + c_1 e^{2x} \sin(x)$$

## 15.74 problem 22.15 (a)

Internal problem ID [13449]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.15 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 5y'x + 8y = \frac{5}{x^3}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(x^2*diff(y(x),x$2)-5*x*diff(y(x),x)+8*y(x)=5/x^3,y(x), singsol=all)
```

$$y = c_2 x^4 + c_1 x^2 + \frac{1}{7x^3}$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 25

```
DSolve[x^2*y'[x]-5*x*y'[x]+8*y[x]==5/x^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 x^4 + \frac{1}{7x^3} + c_1 x^2$$



## 15.75 problem 22.15 (b)

Internal problem ID [13450]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.15 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2y'' - y'x + y = \frac{50}{x^3}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(2*x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=50/x^3,y(x), singsol=all)
```

$$y = c_2\sqrt{x} + c_1x + \frac{25}{14x^3}$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 25

```
DSolve[2*x^2*y''[x]-x*y'[x]+y[x]==50/x^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{25}{14x^3} + c_1\sqrt{x} + c_2x$$

## 15.76 problem 22.15 (c)

Internal problem ID [13451]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.15 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$2x^2y'' + 5y'x + y = 85 \cos(2 \ln(x))$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
dsolve(2*x^2*diff(y(x),x$2)+5*x*diff(y(x),x)+y(x)=85*cos(2*ln(x)),y(x), singsol=all)
```

$$y = \frac{c_2 + \int \frac{c_1 + 17 \cos(2 \ln(x))x + 34x \sin(2 \ln(x))}{2x^{\frac{3}{2}}} dx}{\sqrt{x}}$$

### ✓ Solution by Mathematica

Time used: 0.061 (sec). Leaf size: 34

```
DSolve[2*x^2*y'[x]+5*x*y'[x]+y[x]==85*Cos[2*Log[x]],y[x],x,IncludeSingularSolutions->True
```

$$y(x) \rightarrow \frac{c_1}{x} + \frac{c_2}{\sqrt{x}} + 6 \sin(2 \log(x)) - 7 \cos(2 \log(x))$$

## 15.77 problem 22.15 (d)

Internal problem ID [13452]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.15 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$x^2 y'' - 2y = 15 \cos(3 \ln(x)) - 10 \sin(3 \ln(x))$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
dsolve(x^2*diff(y(x),x$2)-2*y(x)=15*cos(3*ln(x))-10*sin(3*ln(x)),y(x), singsol=all)
```

$$y = \sin\left(\frac{3 \ln(x)}{2}\right) \cos\left(\frac{3 \ln(x)}{2}\right) - 3 \cos\left(\frac{3 \ln(x)}{2}\right)^2 + \frac{3}{2} + \frac{c_1}{x} + c_2 x^2$$

### ✓ Solution by Mathematica

Time used: 0.154 (sec). Leaf size: 35

```
DSolve[x^2*y''[x]-2*y[x]==15*Cos[3*Log[x]]-10*Sin[3*Log[x]],y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow c_2 x^2 + \frac{c_1}{x} + \frac{1}{2}(\sin(3 \log(x)) - 3 \cos(3 \log(x)))$$

## 15.78 problem 22.15 (e)

Internal problem ID [13453]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.15 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$3x^2y'' - 7y'x + 3y = 4x^3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(3*x^2*diff(y(x),x$2)-7*x*diff(y(x),x)+3*y(x)=4*x^3,y(x), singsol=all)
```

$$y = c_2x^3 + c_1x^{\frac{1}{3}} + \frac{x^3(-3 + 8 \ln(x))}{16}$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 33

```
DSolve[3*x^2*y'[x]-7*x*y'[x]+3*y[x]==4*x^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}x^3 \log(x) + \left(-\frac{3}{16} + c_2\right)x^3 + c_1\sqrt[3]{x}$$

## 15.79 problem 22.15 (f)

Internal problem ID [13454]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.15 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$2x^2y'' + 5y'x + y = \frac{10}{x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(2*x^2*diff(y(x),x$2)+5*x*diff(y(x),x)+y(x)=10/x,y(x), singsol=all)
```

$$y = \frac{c_1}{x} + \frac{c_2}{\sqrt{x}} - \frac{10(\ln(x) + 2)}{x}$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 25

```
DSolve[2*x^2*y''[x]+5*x*y'[x]+y[x]==10/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-10 \log(x) + c_2 \sqrt{x} - 20 + c_1}{x}$$

## 15.80 problem 22.15 (g)

Internal problem ID [13455]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.15 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 5y'x + 9y = 6x^3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(x^2*diff(y(x),x$2)-5*x*diff(y(x),x)+9*y(x)=6*x^3,y(x), singsol=all)
```

$$y = c_2 x^3 + \ln(x) x^3 c_1 + 3 \ln(x)^2 x^3$$

### ✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 24

```
DSolve[x^2*y'[x]-5*x*y'[x]+9*y[x]==6*x^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^3 (3 \log^2(x) + 3c_2 \log(x) + c_1)$$

## 15.81 problem 22.15 (h)

Internal problem ID [13456]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 22. Method of undetermined coefficients. Additional exercises page 412

**Problem number:** 22.15 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x^2 y'' + 5y'x + 4y = 64 \ln(x) x^2$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 28

```
dsolve(x^2*diff(y(x),x$2)+5*x*diff(y(x),x)+4*y(x)=64*x^2*ln(x),y(x), singsol=all)
```

$$y = \frac{c_2}{x^2} + \frac{\ln(x) c_1}{x^2} + 2x^2(2 \ln(x) - 1)$$

### ✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 29

```
DSolve[x^2*y''[x]+5*x*y'[x]+4*y[x]==64*x^2*Log[x],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{-2x^4 + 2(2x^4 + c_2) \log(x) + c_1}{x^2}$$

## 16 Chapter 24. Variation of parameters.

### Additional exercises page 444

16.1 problem 24.1 (a)	591
16.2 problem 24.1 (b)	592
16.3 problem 24.1 (c)	593
16.4 problem 24.1 (d)	594
16.5 problem 24.1 (e)	595
16.6 problem 24.1 (f)	596
16.7 problem 24.1 (g)	597
16.8 problem 24.1 (h)	598
16.9 problem 24.1 (i)	599
16.10 problem 24.1 (j)	600
16.11 problem 24.1 (k)	601
16.12 problem 24.1 (L)	602
16.13 problem 24.1 (m)	603
16.14 problem 24.1 (n)	604
16.15 problem 24.2 (a)	605
16.16 problem 24.2 (b)	606
16.17 problem 24.3 (a)	607
16.18 problem 24.3 (b)	608
16.19 problem 24.4 (a)	609
16.20 problem 24.4 (b)	610
16.21 problem 24.4 (c)	612
16.22 problem 24.4 (d)	613



## 16.1 problem 24.1 (a)

Internal problem ID [13457]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.1 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 2y'x + 2y = 3\sqrt{x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve(x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=3*sqrt(x),y(x), singsol=all)
```

$$y = c_1 x^2 + c_2 x + 4\sqrt{x}$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 23

```
DSolve[x^2*y''[x]-2*x*y'[x]+2*y[x]==3*Sqrt[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 x^2 + 4\sqrt{x} + c_1 x$$

## 16.2 problem 24.1 (b)

Internal problem ID [13458]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.1 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \cot(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)+y(x)=cot(x),y(x), singsol=all)
```

$$y = c_2 \sin(x) + c_1 \cos(x) + \sin(x) \ln(\csc(x) - \cot(x))$$

### ✓ Solution by Mathematica

Time used: 0.055 (sec). Leaf size: 33

```
DSolve[y''[x]+y[x]==Cot[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(x) + \sin(x) \left( \log\left(\sin\left(\frac{x}{2}\right)\right) - \log\left(\cos\left(\frac{x}{2}\right)\right) \right) + c_2$$

## 16.3 problem 24.1 (c)

Internal problem ID [13459]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.1 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y = \csc(2x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$2)+4*y(x)=csc(2*x),y(x), singsol=all)
```

$$y = \sin(2x) c_2 + \cos(2x) c_1 - \frac{\ln(\csc(2x)) \sin(2x)}{4} - \frac{x \cos(2x)}{2}$$

### ✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 37

```
DSolve[y''[x]+4*y[x]==Csc[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left(-\frac{x}{2} + c_1\right) \cos(2x) + \frac{1}{4} \sin(2x) (\log(\sin(2x)) + 4c_2)$$

## 16.4 problem 24.1 (d)

Internal problem ID [13460]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.1 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 7y' + 10y = 6e^{3x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)-7*diff(y(x),x)+10*y(x)=6*exp(3*x),y(x), singsol=all)
```

$$y = c_1 e^{5x} + c_2 e^{2x} - 3e^{3x}$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 27

```
DSolve[y''[x]-7*y'[x]+10*y[x]==6*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(-3e^x + c_2 e^{3x} + c_1)$$

## 16.5 problem 24.1 (e)

Internal problem ID [13461]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.1 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y' + 4y = (24x^2 + 2)e^{2x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+4*y(x)=(24*x^2+2)*exp(2*x),y(x), singsol=all)
```

$$y = c_2 e^{2x} + c_1 x e^{2x} + e^{2x}(2x^4 + x^2)$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 26

```
DSolve[y''[x]-4*y'[x]+4*y[x]==(24*x^2+2)*Exp[2*x],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow e^{2x}(2x^4 + x^2 + c_2 x + c_1)$$

## 16.6 problem 24.1 (f)

Internal problem ID [13462]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.1 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y' + 4y = \frac{e^{-2x}}{x^2 + 1}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)+4*y(x)=exp(-2*x)/(1+x^2),y(x), singsol=all)
```

$$y = c_2 e^{-2x} + c_1 x e^{-2x} - \frac{e^{-2x}(-2x \arctan(x) + \ln(x^2 + 1))}{2}$$

### ✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 37

```
DSolve[y''[x]+4*y'[x]+4*y[x]==Exp[-2*x]/(1+x^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^{-2x} (2x \arctan(x) - \log(x^2 + 1) + 2(c_2 x + c_1))$$

## 16.7 problem 24.1 (g)

Internal problem ID [13463]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.1 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$x^2 y'' + y' x - y = \sqrt{x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=sqrt(x),y(x), singsol=all)
```

$$y = c_2 x - \frac{4\sqrt{x}}{3} + \frac{c_1}{x}$$

### ✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 25

```
DSolve[x^2*y''[x]+x*y'[x]-y[x]==Sqrt[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{4\sqrt{x}}{3} + \frac{c_1}{x} + c_2 x$$

## 16.8 problem 24.1 (h)

Internal problem ID [13464]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.1 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y' x - 9y = 12x^3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-9*y(x)=12*x^3,y(x), singsol=all)
```

$$y = x^3 \left( -\frac{1}{3} + 2 \ln(x) \right) + \frac{c_1}{x^3} + c_2 x^3$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 29

```
DSolve[x^2*y''[x]+x*y'[x]-9*y[x]==12*x^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2x^3 \log(x) + \left( -\frac{1}{3} + c_2 \right) x^3 + \frac{c_1}{x^3}$$



## 16.9 problem 24.1 (i)

Internal problem ID [13465]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.1 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 3y'x + 4y = x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(x^2*diff(y(x),x$2)-3*x*diff(y(x),x)+4*y(x)=x^2,y(x), singsol=all)
```

$$y = c_2 x^2 + \ln(x) x^2 c_1 + \frac{\ln(x)^2 x^2}{2}$$

### ✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 27

```
DSolve[x^2*y'[x]-3*x*y'[x]+4*y[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}x^2(\log^2(x) + 4c_2 \log(x) + 2c_1)$$

## 16.10 problem 24.1 (j)

Internal problem ID [13466]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.1 (j).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + 5y'x + 4y = \ln(x)$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 22

```
dsolve(x^2*diff(y(x),x$2)+5*x*diff(y(x),x)+4*y(x)=ln(x),y(x), singsol=all)
```

$$y = \frac{\ln(x)}{4} - \frac{1}{4} + \frac{\ln(x) c_1}{x^2} + \frac{c_2}{x^2}$$

### ✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 29

```
DSolve[x^2*y''[x]+5*x*y'[x]+4*y[x]==Log[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{x^2} + \left( \frac{1}{4} + \frac{2c_2}{x^2} \right) \log(x) - \frac{1}{4}$$

## 16.11 problem 24.1 (k)

Internal problem ID [13467]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.1 (k).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$x^2 y'' - 2y = \frac{1}{x-2}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 50

```
dsolve(x^2*diff(y(x),x$2)-2*y(x)=1/(x-2),y(x), singsol=all)
```

$$y = \frac{c_1}{x} + c_2 x^2 - \frac{\ln(x) x^3 - \ln(x-2) x^3 - 2x^2 + 8 \ln(x-2) - 2x}{24x}$$

### ✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 57

```
DSolve[x^2*y''[x]-2*y[x]==1/(x-2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3 \log(2-x) - x^3 \log(x) + 24c_2 x^3 + 2x^2 + 2x - 8 \log(6-3x) + 24c_1}{24x}$$

## 16.12 problem 24.1 (L)

Internal problem ID [13468]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.1 (L).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$xy'' - y' - 4x^3y = x^3e^{x^2}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(x*diff(y(x),x$2)-diff(y(x),x)-4*x^3*y(x)=x^3*exp(x^2),y(x), singsol=all)
```

$$y = \sinh(x^2) c_2 + c_1 \cosh(x^2) + \frac{x^2 e^{x^2}}{8}$$

### ✓ Solution by Mathematica

Time used: 0.069 (sec). Leaf size: 47

```
DSolve[x*y''[x]-y'[x]-4*x^3*y[x]==x^3*Exp[x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{16} \left( (2x^2 - 1 + 16c_1) \cosh(x^2) + \sinh(x^2) \left( \log(e^{2x^2}) - 1 + 16ic_2 \right) \right)$$

## 16.13 problem 24.1 (m)

Internal problem ID [13469]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.1 (m).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$xy'' + (2x + 2)y' + 2y = 8e^{2x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(x*diff(y(x),x$2)+(2+2*x)*diff(y(x),x)+2*y(x)=8*exp(2*x),y(x), singsol=all)
```

$$y = \frac{e^{-2x}c_2}{x} + \frac{c_1}{x} + \frac{e^{-2x}e^{4x}}{x}$$

### ✓ Solution by Mathematica

Time used: 0.068 (sec). Leaf size: 31

```
DSolve[x*y''[x]+(2+2*x)*y'[x]+2*y[x]==8*Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2e^{2x} + 2c_1e^{-2x} + c_2}{2x}$$

## 16.14 problem 24.1 (n)

Internal problem ID [13470]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.1 (n).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x + 1)y'' + y'x - y = (x + 1)^2$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 18

```
dsolve((x+1)*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=(x+1)^2,y(x), singsol=all)
```

$$y = c_2x + c_1e^{-x} + x^2 + 1$$

### ✓ Solution by Mathematica

Time used: 0.163 (sec). Leaf size: 41

```
DSolve[(x+1)*y'[x]+x*y'[x]-y[x]==(x+1)^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2 + \left(-1 + \sqrt{2}ec_2\right)x + \frac{c_1e^{-x-\frac{1}{2}}}{\sqrt{2}} + 1$$

## 16.15 problem 24.2 (a)

Internal problem ID [13471]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.2 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$x^2 y'' - 2y'x - 4y = \frac{10}{x}$$

With initial conditions

$$[y(1) = 3, y'(1) = -15]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve([x^2*diff(y(x),x$2)-2*x*diff(y(x),x)-4*y(x)=10/x,y(1) = 3, D(y)(1) = -15],y(x), sings
```

$$y = \frac{-2x^5 - 2 \ln(x) + 5}{x}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 20

```
DSolve[{x^2*y''[x]-2*x*y'[x]-4*y[x]==10/x,{y[1]==3,y'[1]==-15}},y[x],x,IncludeSingularSoluti
```

$$y(x) \rightarrow \frac{-2x^5 - 2 \log(x) + 5}{x}$$

## 16.16 problem 24.2 (b)

Internal problem ID [13472]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.2 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - y' - 6y = 12e^{2x}$$

With initial conditions

$$[y(0) = 0, y'(0) = 8]$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve([diff(y(x),x$2)-diff(y(x),x)-6*y(x)=12*exp(2*x),y(0) = 0, D(y)(0) = 8],y(x), singsol=
```

$$y = (4e^{5x} - 3e^{4x} - 1)e^{-2x}$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 27

```
DSolve[{y'[x]-y'[x]-6*y[x]==12*Exp[2*x],{y[0]==0,y'[0]==8}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow e^{-2x}(-3e^{4x} + 4e^{5x} - 1)$$



## 16.17 problem 24.3 (a)

Internal problem ID [13473]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.3 (a).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_y]]`

$$y''' - 4y' = 30e^{3x}$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$3)-4*diff(y(x),x)=30*exp(3*x),y(x), singsol=all)
```

$$y = -\frac{c_1 e^{-2x}}{2} + \frac{c_2 e^{2x}}{2} + 2e^{3x} + c_3$$

### ✓ Solution by Mathematica

Time used: 0.046 (sec). Leaf size: 37

```
DSolve[y'''[x]-4*y'[x]==30*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2e^{3x} + \frac{1}{2}c_1 e^{2x} - \frac{1}{2}c_2 e^{-2x} + c_3$$

## 16.18 problem 24.3 (b)

Internal problem ID [13474]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.3 (b).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$x^3 y''' - 3x^2 y'' + 6y'x - 6y = x^3$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve(x^3*diff(y(x),x$3)-3*x^2*diff(y(x),x$2)+6*x*diff(y(x),x)-6*y(x)=x^3,y(x), singsol=all
```

$$y = -\frac{3x^3}{4} + \frac{\ln(x)x^3}{2} + c_3x^3 + c_2x^2 + c_1x$$

### ✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 34

```
DSolve[x^3*y'''[x]-3*x^2*y''[x]+6*x*y'[x]-6*y[x]==x^3,y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{1}{2}x^3 \log(x) + x \left( \left( -\frac{3}{4} + c_3 \right) x^2 + c_2x + c_1 \right)$$

## 16.19 problem 24.4 (a)

Internal problem ID [13475]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.4 (a).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$x^3 y''' - 3x^2 y'' + 6y'x - 6y = e^{-x^2}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 73

```
dsolve(x^3*diff(y(x),x$3)-3*x^2*diff(y(x),x$2)+6*x*diff(y(x),x)-6*y(x)=exp(-x^2),y(x), singularSolutions)
```

$$y = \frac{\left(2\sqrt{\pi} \operatorname{erf}(x) e^{x^2} x^3 - 3\sqrt{\pi} \operatorname{erf}(x) e^{x^2} x - 3 \operatorname{ExpIntegral}_1(x^2) e^{x^2} x^2 + 2x^2 - 1\right) e^{-x^2}}{6} + c_3 x^3 + c_2 x^2 + c_1 x$$

### ✓ Solution by Mathematica

Time used: 0.068 (sec). Leaf size: 77

```
DSolve[x^3*y'''[x]-3*x^2*y''[x]+6*x*y'[x]-6*y[x]==Exp[-x^2],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{6} \left( \sqrt{\pi} (2x^2 - 3) x \operatorname{erf}(x) + 3x^2 \operatorname{ExpIntegralEi}(-x^2) + 6c_3 x^3 + 2e^{-x^2} x^2 - e^{-x^2} + 6c_2 x^2 + 6c_1 x \right)$$

## 16.20 problem 24.4 (b)

Internal problem ID [13476]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.4 (b).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - y'' + y' - y = \tan(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 72

```
dsolve(diff(y(x),x$3)-diff(y(x),x$2)+diff(y(x),x)-y(x)=tan(x),y(x), singsol=all)
```

$$y = \frac{\cos(x)^2}{2} + \frac{\cos(x) \ln(\sec(x) + \tan(x))}{2} - \left( \int -\frac{(\cos(x)^2 + \sin(x)^2) \tan(x) e^{-x}}{2} dx \right) e^x + \frac{\sin(x)^2}{2} - \frac{\sin(x) \ln(\sec(x) + \tan(x))}{2} + c_1 \cos(x) + c_2 e^x + c_3 \sin(x)$$

✓ Solution by Mathematica

Time used: 0.06 (sec). Leaf size: 106

```
DSolve[y''''[x]-y'''[x]+y''[x]-y[x]==Tan[x],y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned}y(x) \rightarrow & -\frac{1}{2} \sin(x) \operatorname{arctanh}(\sin(x)) + \frac{1}{2} \cos(x) \operatorname{arctanh}(\sin(x)) \\ & - \frac{1}{2} i \operatorname{Hypergeometric2F1}\left(\frac{i}{2}, 1, 1 + \frac{i}{2}, -e^{2ix}\right) \\ & - \left(\frac{1}{5} - \frac{i}{10}\right) e^{2ix} \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{i}{2}, 2 + \frac{i}{2}, -e^{2ix}\right) \\ & + c_3 e^x + c_1 \cos(x) + c_2 \sin(x) + \frac{1}{2}\end{aligned}$$

## 16.21 problem 24.4 (c)

Internal problem ID [13477]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.4 (c).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 81y = \sinh(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 46

```
dsolve(diff(y(x),x$4)-81*y(x)=sinh(x),y(x), singsol=all)
```

$$y = \frac{e^{-4x}(-e^{5x} + e^{3x})}{160} + c_1 \cos(3x) + c_2 e^{-3x} + c_3 e^{3x} + c_4 \sin(3x)$$

### ✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 52

```
DSolve[y''''[x]-81*y[x]==Sinh[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{-x}}{160} - \frac{e^x}{160} + c_1 e^{3x} + c_3 e^{-3x} + c_2 \cos(3x) + c_4 \sin(3x)$$

## 16.22 problem 24.4 (d)

Internal problem ID [13478]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 24. Variation of parameters. Additional exercises page 444

**Problem number:** 24.4 (d).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _exact, _linear, _nonhomogeneous]]`

$$x^4 y'''' + 6x^3 y''' - 3x^2 y'' - 9y'x + 9y = 12x \sin(x^2)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 72

```
dsolve(x^4*diff(y(x),x$4)+6*x^3*diff(y(x),x$3)-3*x^2*diff(y(x),x$2)-9*x*diff(y(x),x)+9*y(x)=
```

$$y = -\frac{-2 \operatorname{Ci}(x^2) x^6 + 2 \sin(x^2) x^4 + 6 \operatorname{Si}(x^2) x^4 + 4x^2 \cos(x^2) - c_1 x^2 + 2 \sin(x^2)}{16x^3} + c_2 x + \frac{c_3}{x^3} + c_4 x^3$$

### ✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 79

```
DSolve[x^4*y''''[x]+6*x^3*y'''[x]-3*x^2*y''[x]-9*x*y'[x]+9*y[x]==12*x*Sin[x^2],y[x],x,Includ
```

$$y(x) \rightarrow \frac{x^6 \operatorname{CosIntegral}(x^2) - 3x^4 \operatorname{Si}(x^2) + 8c_4 x^6 + 8c_3 x^4 - \sin(x^2) - 2x^2 \cos(x^2) + 8c_2 x^2 - x^4 \sin(x^2) + 8c_1}{8x^3}$$

## 17 Chapter 25. Review exercises for part III. page 447

17.1 problem 1	616
17.2 problem 2	617
17.3 problem 3	618
17.4 problem 4	619
17.5 problem 5	620
17.6 problem 6	621
17.7 problem 7	622
17.8 problem 8	623
17.9 problem 9	624
17.10problem 10	625
17.11problem 11	626
17.12problem 12	627
17.13problem 13	628
17.14problem 14	629
17.15problem 15	630
17.16problem 16	631
17.17problem 17	632
17.18problem 18	633
17.19problem 19	634
17.20problem 20	635
17.21problem 21	636
17.22problem 22	637
17.23problem 23	638
17.24problem 24	639
17.25problem 25	640
17.26problem 26	641
17.27problem 27	642
17.28problem 28	643
17.29problem 29	644
17.30problem 30	645
17.31problem 31	646
17.32problem 32	647
17.33problem 33	648
17.34problem 34	649
17.35problem 35	650
17.36problem 36	651



17.37problem 37	652
17.38problem 38	653
17.39problem 39	654
17.40problem 40	655
17.41problem 41	656
17.42problem 42	657
17.43problem 43	659
17.44problem 44	660
17.45problem 45	661
17.46problem 46	662
17.47problem 47	663
17.48problem 48	664
17.49problem 49	665
17.50problem 50	666

## 17.1 problem 1

Internal problem ID [13479]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 1.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 36y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+36*y(x)=0,y(x), singsol=all)
```

$$y = c_1 \sin(6x) + c_2 \cos(6x)$$

### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 20

```
DSolve[y''[x]+36*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(6x) + c_2 \sin(6x)$$

## 17.2 problem 2

Internal problem ID [13480]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 2.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 12y' + 36y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)-12*diff(y(x),x)+36*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{6x} + c_2 x e^{6x}$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 18

```
DSolve[y''[x]-12*y'[x]+36*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{6x}(c_2 x + c_1)$$

## 17.3 problem 3

Internal problem ID [13481]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 3.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$x^2y'' + y'x - 9y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-9*y(x)=0,y(x), singsol=all)
```

$$y = c_1x^3 + \frac{c_2}{x^3}$$

### ✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]+x*y'[x]-9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2x^6 + c_1}{x^3}$$

## 17.4 problem 4

Internal problem ID [13482]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 4.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 36y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-36*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-6x} + c_2 e^{6x}$$

### ✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 22

```
DSolve[y''[x]-36*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{6x} + c_2 e^{-6x}$$

## 17.5 problem 5

Internal problem ID [13483]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 5.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 9y' + 14y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-9*diff(y(x),x)+14*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{7x} + c_2 e^{2x}$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 22

```
DSolve[y''[x]-9*y'[x]+14*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(c_2 e^{5x} + c_1)$$

## 17.6 problem 6

Internal problem ID [13484]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 6.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - 7y'x + 16y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x^2*diff(y(x),x$2)-7*x*diff(y(x),x)+16*y(x)=0,y(x), singsol=all)
```

$$y = c_1 x^4 + c_2 x^4 \ln(x)$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]-7*x*y'[x]+16*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^4(4c_2 \log(x) + c_1)$$

## 17.7 problem 7

Internal problem ID [13485]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 7.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$2xy'' + y' = \sqrt{x}$$

### ✓ Solution by Maple

Time used: 0.172 (sec). Leaf size: 16

```
dsolve(2*x*diff(y(x),x$2)+diff(y(x),x)=sqrt(x),y(x), singsol=all)
```

$$y = \frac{\sqrt{x}(x + 6c_1)}{3} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 23

```
DSolve[2*x*y''[x]+y'[x]==Sqrt[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}\sqrt{x}(x + 6c_1) + c_2$$



## 17.8 problem 8

Internal problem ID [13486]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 8.

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 8y'' + 16y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$4)-8*diff(y(x),x$2)+16*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-2x} + c_2 x e^{-2x} + e^{2x} c_3 + x e^{2x} c_4$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 35

```
DSolve[y''''[x]-8*y''[x]+16*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} (c_3 e^{4x} + x(c_4 e^{4x} + c_2) + c_1)$$

## 17.9 problem 9

Internal problem ID [13487]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 9.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 6y' + 9y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)+6*diff(y(x),x)+9*y(x)=0,y(x), singsol=all)
```

$$y = e^{-3x}c_1 + c_2x e^{-3x}$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 18

```
DSolve[y''[x]+6*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x}(c_2x + c_1)$$

## 17.10 problem 10

Internal problem ID [13488]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 10.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 3y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)+3*y(x)=0,y(x), singsol=all)
```

$$y = c_1 \sin(\sqrt{3}x) + c_2 \cos(\sqrt{3}x)$$

### ✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 28

```
DSolve[y''[x]+3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(\sqrt{3}x) + c_2 \sin(\sqrt{3}x)$$

## 17.11 problem 11

Internal problem ID [13489]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 11.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2y'' + 7y'x + 9y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x^2*diff(y(x),x$2)+7*x*diff(y(x),x)+9*y(x)=0,y(x), singsol=all)
```

$$y = \frac{c_1}{x^3} + \frac{c_2 \ln(x)}{x^3}$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]+7*x*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3c_2 \log(x) + c_1}{x^3}$$

## 17.12 problem 12

Internal problem ID [13490]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 12.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' + \frac{5y}{2} = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(x^2*diff(y(x),x$2)+5/2*y(x)=0,y(x), singsol=all)
```

$$y = c_1 \sqrt{x} \sin\left(\frac{3 \ln(x)}{2}\right) + c_2 \sqrt{x} \cos\left(\frac{3 \ln(x)}{2}\right)$$

### ✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 32

```
DSolve[x^2*y''[x]+5/2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt{x} \left( c_2 \cos\left(\frac{3 \log(x)}{2}\right) + c_1 \sin\left(\frac{3 \log(x)}{2}\right) \right)$$

## 17.13 problem 13

Internal problem ID [13491]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 13.

**ODE order:** 5.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y^{(5)} - 6y'''' + 13y''' = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$5)-6*diff(y(x),x$4)+13*diff(y(x),x$3)=0,y(x), singsol=all)
```

$$y = c_1 + c_2x + c_3x^2 + c_4e^{3x} \sin(2x) + c_5 \cos(2x) e^{3x}$$

### ✓ Solution by Mathematica

Time used: 2.104 (sec). Leaf size: 56

```
DSolve[y'''''[x]-6*y''''[x]+13*y'''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_5x^2 + c_4x - \frac{e^{3x}((46c_1 + 9c_2) \cos(2x) + (9c_1 - 46c_2) \sin(2x))}{2197} + c_3$$

## 17.14 problem 14

Internal problem ID [13492]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 14.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - 6y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)-6*y(x)=0,y(x), singsol=all)
```

$$y = c_1 x^3 + \frac{c_2}{x^2}$$

### ✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]-6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 x^5 + c_1}{x^2}$$

## 17.15 problem 15

Internal problem ID [13493]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 15.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 6y' + 25y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)-6*diff(y(x),x)+25*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{3x} \sin(4x) + c_2 e^{3x} \cos(4x)$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 26

```
DSolve[y''[x]-6*y'[x]+25*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{3x}(c_2 \cos(4x) + c_1 \sin(4x))$$



## 17.16 problem 16

Internal problem ID [13494]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 16.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible]`

$$y'' - y'^2 = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x$2)=diff(y(x),x)^2,y(x), singsol=all)
```

$$y = -\ln(-c_1x - c_2)$$

### ✓ Solution by Mathematica

Time used: 0.217 (sec). Leaf size: 15

```
DSolve[y''[x]==y'[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \log(x + c_1)$$

## 17.17 problem 17

Internal problem ID [13495]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 17.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$x^2 y'' + y' x + 9y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+9*y(x)=0,y(x), singsol=all)
```

$$y = c_1 \sin(3 \ln(x)) + c_2 \cos(3 \ln(x))$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 22

```
DSolve[x^2*y'[x]+x*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(3 \log(x)) + c_2 \sin(3 \log(x))$$

## 17.18 problem 18

Internal problem ID [13496]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 18.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 8y' + 25y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)-8*diff(y(x),x)+25*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{4x} \sin(3x) + c_2 e^{4x} \cos(3x)$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 26

```
DSolve[y''[x]-8*y'[x]+25*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{4x}(c_2 \cos(3x) + c_1 \sin(3x))$$

## 17.19 problem 19

Internal problem ID [13497]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 19.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2y'' + 2y'x - 30y = 0$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)+2*x*diff(y(x),x)-30*y(x)=0,y(x), singsol=all)
```

$$y = c_1x^5 + \frac{c_2}{x^6}$$

### ✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]+2*x*y'[x]-30*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{x^6} + c_2x^5$$

## 17.20 problem 20

Internal problem ID [13498]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 20.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y' - 30y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+diff(y(x),x)-30*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{5x} + c_2 e^{-6x}$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 22

```
DSolve[y''[x]+y'[x]-30*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-6x} + c_2 e^{5x}$$

## 17.21 problem 21

Internal problem ID [13499]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 21.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$16y'' - 8y' + y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(16*diff(y(x),x$2)-8*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{\frac{x}{4}} + c_2 x e^{\frac{x}{4}}$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 20

```
DSolve[16*y''[x]-8*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x/4}(c_2 x + c_1)$$

## 17.22 problem 22

Internal problem ID [13500]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 22.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$4x^2y'' + 8y'x + y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(4*x^2*diff(y(x),x$2)+8*x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y = \frac{c_1}{\sqrt{x}} + \frac{c_2 \ln(x)}{\sqrt{x}}$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 24

```
DSolve[4*x^2*y''[x]+8*x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 \log(x) + 2c_1}{2\sqrt{x}}$$

## 17.23 problem 23

Internal problem ID [13501]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 23.

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 6y'' + 12y' = 8$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 67

```
dsolve(diff(y(x),x$3)-6*diff(y(x),x$2)+12*diff(y(x),x)=8,y(x), singsol=all)
```

$$y = \frac{e^{3x} \cos(\sqrt{3}x) c_1}{4} + \frac{c_1 \sqrt{3} e^{3x} \sin(\sqrt{3}x)}{12} - \frac{c_2 \sqrt{3} e^{3x} \cos(\sqrt{3}x)}{12} + \frac{e^{3x} \sin(\sqrt{3}x) c_2}{4} + \frac{2x}{3} + c_3$$

### ✓ Solution by Mathematica

Time used: 0.312 (sec). Leaf size: 71

```
DSolve[y'''[x]-6*y''[x]+12*y'[x]==8,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{12} \left( 8x - \left( \sqrt{3}c_1 - 3c_2 \right) e^{3x} \cos(\sqrt{3}x) + \left( 3c_1 + \sqrt{3}c_2 \right) e^{3x} \sin(\sqrt{3}x) \right) + c_3$$



## 17.24 problem 24

Internal problem ID [13502]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 24.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$2x^2y'' - 3y'x + 2y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(2*x^2*diff(y(x),x$2)-3*x*diff(y(x),x)+2*y(x)=0,y(x), singsol=all)
```

$$y = c_1x^2 + c_2\sqrt{x}$$

### ✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 20

```
DSolve[2*x^2*y''[x]-3*x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2x^2 + c_1\sqrt{x}$$

## 17.25 problem 25

Internal problem ID [13503]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 25.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$9x^2y'' + 3y'x + y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(9*x^2*diff(y(x),x$2)+3*x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y = c_1x^{\frac{1}{3}} + c_2x^{\frac{1}{3}} \ln(x)$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 24

```
DSolve[9*x^2*y''[x]+3*x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}\sqrt[3]{x}(c_2 \log(x) + 3c_1)$$

## 17.26 problem 26

Internal problem ID [13504]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 26.

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 16y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$4)-16*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-2x} + c_2 e^{2x} + c_3 \sin(2x) + c_4 \cos(2x)$$

### ✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 36

```
DSolve[y''''[x]-16*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{2x} + c_3 e^{-2x} + c_2 \cos(2x) + c_4 \sin(2x)$$

## 17.27 problem 27

Internal problem ID [13505]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 27.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$2y'' - 7y' = -3$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(2*diff(y(x),x$2)-7*diff(y(x),x)+3=0,y(x), singsol=all)
```

$$y = \frac{2c_1 e^{\frac{7x}{2}}}{7} + \frac{3x}{7} + c_2$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 26

```
DSolve[2*y''[x]-7*y'[x]+3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3x}{7} + \frac{2}{7}c_1 e^{7x/2} + c_2$$

## 17.28 problem 28

Internal problem ID [13506]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 28.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 20y' + 100y = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)+20*diff(y(x),x)+100*y(x)=0,y(x), singsol=all)
```

$$y = c_1 e^{-10x} + c_2 x e^{-10x}$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 18

```
DSolve[y''[x]+20*y'[x]+100*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-10x}(c_2 x + c_1)$$

## 17.29 problem 29

Internal problem ID [13507]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 29.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' - 3y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(x*diff(y(x),x$2)=3*diff(y(x),x),y(x), singsol=all)
```

$$y = c_2x^4 + c_1$$

### ✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 17

```
DSolve[x*y''[x]==3*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1x^4}{4} + c_2$$

## 17.30 problem 30

Internal problem ID [13508]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 30.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 5y' = 0$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x$2)-5*diff(y(x),x)=0,y(x), singsol=all)
```

$$y = c_1 + c_2 e^{5x}$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 19

```
DSolve[y''[x]-5*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{5}c_1 e^{5x} + c_2$$

## 17.31 problem 31

Internal problem ID [13509]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 31.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 9y' + 14y = 98x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)-9*diff(y(x),x)+14*y(x)=98*x^2,y(x), singsol=all)
```

$$y = c_2 e^{7x} + c_1 e^{2x} + 7x^2 + 9x + \frac{67}{14}$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 33

```
DSolve[y''[x]-9*y'[x]+14*y[x]==98*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 7x^2 + 9x + c_1 e^{2x} + c_2 e^{7x} + \frac{67}{14}$$



## 17.32 problem 32

Internal problem ID [13510]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 32.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 12y' + 36y = 25 \sin(3x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$2)-12*diff(y(x),x)+36*y(x)=25*sin(3*x),y(x), singsol=all)
```

$$y(x) = e^{6x}c_2 + e^{6x}xc_1 + \frac{4 \cos(3x)}{9} + \frac{\sin(3x)}{3}$$

### ✓ Solution by Mathematica

Time used: 0.092 (sec). Leaf size: 35

```
DSolve[y''[x]-12*y'[x]+36*y[x]==25*Sin[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} \sin(3x) + \frac{4}{9} \cos(3x) + e^{6x}(c_2x + c_1)$$

### 17.33 problem 33

Internal problem ID [13511]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 33.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 9y' + 14y = 576x^2e^{-x}$$

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(x),x$2)-9*diff(y(x),x)+14*y(x)=576*x^2*exp(-x),y(x), singsol=all)
```

$$y(x) = e^{7x}c_2 + e^{2x}c_1 + \frac{(288x^2 + 264x + 97)e^{-x}}{12}$$

#### ✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 39

```
DSolve[y''[x]-9*y'[x]+14*y[x]==576*x^2*Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} \left( 24x^2 + 22x + c_1e^{3x} + c_2e^{8x} + \frac{97}{12} \right)$$

## 17.34 problem 34

Internal problem ID [13512]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 34.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 12y' + 36y = 81e^{3x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)-12*diff(y(x),x)+36*y(x)=81*exp(3*x),y(x), singsol=all)
```

$$y(x) = e^{6x}c_2 + e^{6x}xc_1 + 9e^{3x}$$

### ✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 26

```
DSolve[y''[x]-12*y'[x]+36*y[x]==81*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{3x}(9 + e^{3x}(c_2x + c_1))$$

## 17.35 problem 35

Internal problem ID [13513]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 35.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + x y' - 9y = 3\sqrt{x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-9*y(x)=3*sqrt(x),y(x), singsol=all)
```

$$y(x) = \frac{c_2}{x^3} + c_1 x^3 - \frac{12\sqrt{x}}{35}$$

### ✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 27

```
DSolve[x^2*y''[x]+x*y'[x]-9*y[x]==3*Sqrt[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 x^3 + \frac{c_1}{x^3} - \frac{12\sqrt{x}}{35}$$

## 17.36 problem 36

Internal problem ID [13514]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 36.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 12y' + 36y = 3x e^{6x} - 2e^{6x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$2)-12*diff(y(x),x)+36*y(x)=3*x*exp(6*x)-2*exp(6*x),y(x), singsol=all)
```

$$y(x) = e^{6x}c_2 + e^{6x}xc_1 + \frac{e^{6x}x(9x^2 - 18x + 8)}{18}$$

### ✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 32

```
DSolve[y''[x]-12*y'[x]+36*y[x]==3*x*Exp[6*x]-2*Exp[6*x],y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow \frac{1}{2}e^{6x}(x^3 - 2x^2 + 2c_2x + 2c_1)$$

## 17.37 problem 37

Internal problem ID [13515]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 37.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 36y = 6 \sec(6x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$2)+36*y(x)=6*sec(6*x),y(x), singsol=all)
```

$$y(x) = \sin(6x) c_2 + \cos(6x) c_1 + x \sin(6x) - \frac{\ln(\sec(6x)) \cos(6x)}{6}$$

### ✓ Solution by Mathematica

Time used: 0.045 (sec). Leaf size: 32

```
DSolve[y''[x]+36*y[x]==6*Sec[6*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x + c_2) \sin(6x) + \cos(6x) \left( \frac{1}{6} \log(\cos(6x)) + c_1 \right)$$

## 17.38 problem 38

Internal problem ID [13516]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 38.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + 2xy' - 6y = 18 \ln(x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(x^2*diff(y(x),x$2)+2*x*diff(y(x),x)-6*y(x)=18*ln(x),y(x), singsol=all)
```

$$y(x) = \frac{c_2}{x^3} + c_1 x^2 - 3 \ln(x) - \frac{1}{2}$$

### ✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 25

```
DSolve[x^2*y''[x]+2*x*y'[x]-6*y[x]==18*Log[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{x^3} + c_2 x^2 - 3 \log(x) - \frac{1}{2}$$

## 17.39 problem 39

Internal problem ID [13517]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 39.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 6y' + 9y = 10e^{-3x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)+6*diff(y(x),x)+9*y(x)=10*exp(-3*x),y(x), singsol=all)
```

$$y(x) = e^{-3x}c_2 + e^{-3x}xc_1 + 5x^2e^{-3x}$$

### ✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 23

```
DSolve[y''[x]+6*y'[x]+9*y[x]==10*Exp[-3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x}(5x^2 + c_2x + c_1)$$



## 17.40 problem 40

Internal problem ID [13518]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 40.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2y'' - xy' - 2y = 10x^2$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(2*x^2*diff(y(x),x$2)-x*diff(y(x),x)-2*y(x)=10*x^2,y(x), singsol=all)
```

$$y(x) = c_2x^2 + \frac{c_1}{\sqrt{x}} + \frac{2x^2(-2 + 5 \ln(x))}{5}$$

### ✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 31

```
DSolve[2*x^2*y''[x]-x*y'[x]-2*y[x]==10*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2x^2 \log(x) + \left(-\frac{4}{5} + c_2\right)x^2 + \frac{c_1}{\sqrt{x}}$$

## 17.41 problem 41

Internal problem ID [13519]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 41.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 6y' + 9y = 2 \cos(2x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$2)+6*diff(y(x),x)+9*y(x)=2*cos(2*x),y(x), singsol=all)
```

$$y(x) = e^{-3x}c_2 + e^{-3x}xc_1 + \frac{10 \cos(2x)}{169} + \frac{24 \sin(2x)}{169}$$

### ✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 35

```
DSolve[y''[x]+6*y'[x]+9*y[x]==2*Cos[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{24}{169} \sin(2x) + \frac{10}{169} \cos(2x) + e^{-3x}(c_2x + c_1)$$

## 17.42 problem 42

Internal problem ID [13520]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 42.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$xy'' - y' + 3xy'^3 = 0$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 42

```
dsolve(x*diff(y(x),x$2)-diff(y(x),x)=-3*x* diff(y(x),x)^3,y(x), singsol=all)
```

$$y(x) = \int \frac{x}{\sqrt{2x^3 - c_1}} dx + c_2$$

$$y(x) = \int -\frac{x}{\sqrt{2x^3 - c_1}} dx + c_2$$

✓ Solution by Mathematica

Time used: 1.949 (sec). Leaf size: 195

```
DSolve[x*y''[x]-y'[x]==-3*x*(y'[x])^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \frac{x^2 \sqrt{1 + \frac{2x^3}{c_1}} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, -\frac{2x^3}{c_1}\right)}{2\sqrt{2x^3 + c_1}}$$

$$y(x) \rightarrow \frac{x^2 \sqrt{1 + \frac{2x^3}{c_1}} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, -\frac{2x^3}{c_1}\right)}{2\sqrt{2x^3 + c_1}} + c_2$$

$$y(x) \rightarrow c_2$$

$$y(x) \rightarrow -\frac{3\sqrt{x^3} \operatorname{Gamma}\left(\frac{5}{3}\right)}{\sqrt{2}x \operatorname{Gamma}\left(\frac{2}{3}\right)} + c_2$$

$$y(x) \rightarrow \frac{3\sqrt{x^3} \operatorname{Gamma}\left(\frac{5}{3}\right)}{\sqrt{2}x \operatorname{Gamma}\left(\frac{2}{3}\right)} + c_2$$

## 17.43 problem 43

Internal problem ID [13521]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 43.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + 3xy' + 2y = 6$$

### ✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 22

```
dsolve(x^2*diff(y(x),x$2)+3*x*diff(y(x),x)+2*y(x)=6,y(x), singsol=all)
```

$$y(x) = \frac{\sin(\ln(x)) c_2}{x} + \frac{\cos(\ln(x)) c_1}{x} + 3$$

### ✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 25

```
DSolve[x^2*y''[x]+3*x*y'[x]+2*y[x]==6,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3x + c_2 \cos(\log(x)) + c_1 \sin(\log(x))}{x}$$

## 17.44 problem 44

Internal problem ID [13522]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 44.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$x^2 y'' + xy' - y = \frac{1}{x^2 + 1}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=1/(1+x^2),y(x), singsol=all)
```

$$y(x) = \frac{c_1}{x} + c_2 x - \frac{\arctan(x) x^2 + \arctan(x) + x}{2x}$$

### ✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 33

```
DSolve[x^2*y'[x]+x*y'[x]-y[x]==1/(1+x^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x^2 \arctan(x) + \arctan(x) - 2c_2 x^2 + x - 2c_1}{2x}$$

## 17.45 problem 45

Internal problem ID [13523]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 45.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$4y'' - 12y' + 9y = x e^{\frac{3x}{2}}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve(4*diff(y(x),x$2)-12*diff(y(x),x)+9*y(x)=x*exp(3*x/2),y(x), singsol=all)
```

$$y(x) = e^{\frac{3x}{2}} c_2 + x e^{\frac{3x}{2}} c_1 + \frac{x^3 e^{\frac{3x}{2}}}{24}$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 29

```
DSolve[4*y'[x]-12*y'[x]+9*y[x]==x*Exp[3*x/2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{24} e^{3x/2} (x^3 + 24c_2 x + 24c_1)$$

## 17.46 problem 46

Internal problem ID [13524]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 46.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$3y'' + 8y' - 3y = 123x \sin(3x)$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
dsolve(3*diff(y(x),x$2)+8*diff(y(x),x)-3*y(x)=123*x*sin(3*x),y(x), singsol=all)
```

$$y(x) = e^{-3x}c_2 + e^{\frac{x}{3}}c_1 + \frac{(-492x - 241) \cos(3x)}{246} + \frac{(-615x + 324) \sin(3x)}{246}$$

### ✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 50

```
DSolve[3*y''[x]+8*y'[x]-3*y[x]==123*x*Sin[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left(\frac{54}{41} - \frac{5x}{2}\right) \sin(3x) + \left(-2x - \frac{241}{246}\right) \cos(3x) + c_1 e^{x/3} + c_2 e^{-3x}$$



## 17.47 problem 47

Internal problem ID [13525]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 47.

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$y''' + 8y = e^{-2x}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 38

```
dsolve(diff(y(x),x$3)+8*y(x)=exp(-2*x),y(x), singsol=all)
```

$$y(x) = \frac{x e^{-2x}}{12} + c_1 e^{-2x} + c_2 e^x \cos(\sqrt{3}x) + c_3 e^x \sin(\sqrt{3}x)$$

### ✓ Solution by Mathematica

Time used: 0.337 (sec). Leaf size: 57

```
DSolve[y'''[x]+8*y[x]==Exp[-2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{24} e^{-2x} \left( 2x + 24c_3 e^{3x} \cos(\sqrt{3}x) + 24c_2 e^{3x} \sin(\sqrt{3}x) + 1 + 24c_1 \right)$$

## 17.48 problem 48

Internal problem ID [13526]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 48.

**ODE order:** 6.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _with_linear_symmetries]]`

$$y^{(6)} - 64y = e^{-2x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 68

```
dsolve(diff(y(x),x$6)-64*y(x)=exp(-2*x),y(x), singsol=all)
```

$$y(x) = -\frac{x e^{-2x}}{192} + c_1 e^{-2x} + c_2 e^{2x} + c_3 e^x \cos(\sqrt{3}x) + c_4 e^x \sin(\sqrt{3}x) + c_5 e^{-x} \cos(\sqrt{3}x) + c_6 e^{-x} \sin(\sqrt{3}x)$$

### ✓ Solution by Mathematica

Time used: 0.738 (sec). Leaf size: 80

```
DSolve[y''''''[x]-64*y[x]==Exp[-2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{768} e^{-2x} \left( -4x + 768c_1 e^{4x} + 768e^x (c_2 e^{2x} + c_3) \cos(\sqrt{3}x) + 768e^x (c_6 e^{2x} + c_5) \sin(\sqrt{3}x) - 5 + 768c_4 \right)$$

## 17.49 problem 49

Internal problem ID [13527]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 49.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$x^2 y'' + 3xy' + y = \frac{1}{(x+1)^2}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(x^2*diff(y(x),x$2)+3*x*diff(y(x),x)+y(x)=1/(1+x)^2,y(x), singsol=all)
```

$$y(x) = \frac{\ln(x) c_1}{x} + \frac{c_2}{x} + \frac{\ln(1+x) - \ln(x)}{x}$$

### ✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 23

```
DSolve[x^2*y'[x]+3*x*y'[x]+y[x]==1/(1+x)^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\log(x+1) + (-1 + c_2) \log(x) + c_1}{x}$$

## 17.50 problem 50

Internal problem ID [13528]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 25. Review exercises for part III. page 447

**Problem number:** 50.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$x^2 y'' + 3xy' + y = \frac{1}{x}$$

### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(x^2*diff(y(x),x$2)+3*x*diff(y(x),x)+y(x)=1/x,y(x), singsol=all)
```

$$y(x) = \frac{c_2}{x} + \frac{\ln(x) c_1}{x} + \frac{\ln(x)^2}{2x}$$

### ✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 27

```
DSolve[x^2*y''[x]+3*x*y'[x]+y[x]==1/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\log^2(x) + 2c_2 \log(x) + 2c_1}{2x}$$

## 18 Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496

18.1 problem 27.1 (a)	668
18.2 problem 27.1 (b)	669
18.3 problem 27.1 (c)	670
18.4 problem 27.1 (d)	671
18.5 problem 27.1 (e)	672
18.6 problem 27.1 (f)	673
18.7 problem 27.1 (g)	674
18.8 problem 27.1 (h)	675
18.9 problem 27.1 (i)	676
18.10 problem 27.1 (j)	677
18.11 problem 27.1 (k)	678
18.12 problem 27.1 (L)	679
18.13 problem 27.1 (m)	680
18.14 problem 27.4	681

## 18.1 problem 27.1 (a)

Internal problem ID [13529]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496

**Problem number:** 27.1 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' + 4y = 0$$

With initial conditions

$$[y(0) = 3]$$

### ✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 10

```
dsolve([diff(y(t),t)+4*y(t)=0,y(0) = 3],y(t), singsol=all)
```

$$y(t) = 3e^{-4t}$$

### ✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 12

```
DSolve[{y'[t]+4*y[t]==0,{y[0]==3}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 3e^{-4t}$$

## 18.2 problem 27.1 (b)

Internal problem ID [13530]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496

**Problem number:** 27.1 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' - 2y = t^3$$

With initial conditions

$$[y(0) = 4]$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 25

```
dsolve([diff(y(t),t)-2*y(t)=t^3,y(0) = 4],y(t), singsol=all)
```

$$y(t) = -\frac{3t^2}{4} - \frac{t^3}{2} - \frac{3t}{4} + \frac{35e^{2t}}{8} - \frac{3}{8}$$

✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 31

```
DSolve[{y'[t]+4*y[t]==t^3,{y[0]==4}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{128}(32t^3 - 24t^2 + 12t + 515e^{-4t} - 3)$$

### 18.3 problem 27.1 (c)

Internal problem ID [13531]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496

**Problem number:** 27.1 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 3y = \text{Heaviside}(t - 4)$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 20

```
dsolve([diff(y(t),t)+3*y(t)=Heaviside(t-4),y(0) = 0],y(t), singsol=all)
```

$$y(t) = -\frac{\text{Heaviside}(t - 4)(-1 + e^{-3t+12})}{3}$$

✓ Solution by Mathematica

Time used: 0.065 (sec). Leaf size: 27

```
DSolve[{y'[t]+3*y[t]==UnitStep[t-4],{y[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \begin{cases} \frac{1}{3} - \frac{1}{3}e^{12-3t} & t > 4 \\ 0 & \text{True} \end{cases}$$



## 18.4 problem 27.1 (d)

Internal problem ID [13532]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496

**Problem number:** 27.1 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y = t^3$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 25

```
dsolve([diff(y(t),t$2)-4*y(t)=t^3,y(0) = 1, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = \frac{19e^{2t}}{32} + \frac{13e^{-2t}}{32} - \frac{t^3}{4} - \frac{3t}{8}$$

### ✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 34

```
DSolve[{y'[t]-4*y[t]==t^3,{y[0]==1,y'[0]==3}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{32}(-4t(2t^2 + 3) - 11e^{-2t} + 43e^{2t})$$

## 18.5 problem 27.1 (e)

Internal problem ID [13533]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496

**Problem number:** 27.1 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 4y = 20e^{4t}$$

With initial conditions

$$[y(0) = 3, y'(0) = 12]$$

### ✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 21

```
dsolve([diff(y(t),t$2)+4*y(t)=20*exp(4*t),y(0) = 3, D(y)(0) = 12],y(t), singsol=all)
```

$$y(t) = 2 \cos(2t) + 4 \sin(2t) + e^{4t}$$

### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 23

```
DSolve[{y''[t]+4*y[t]==20*Exp[4*t],{y[0]==3,y'[0]==12}},y[t],t,IncludeSingularSolutions -> T
```

$$y(t) \rightarrow e^{4t} + 4 \sin(2t) + 2 \cos(2t)$$

## 18.6 problem 27.1 (f)

Internal problem ID [13534]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496

**Problem number:** 27.1 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y = \sin(2t)$$

With initial conditions

$$[y(0) = 3, y'(0) = 5]$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 20

```
dsolve([diff(y(t),t$2)+4*y(t)=sin(2*t),y(0) = 3, D(y)(0) = 5],y(t), singsol=all)
```

$$y(t) = \frac{21 \sin(2t)}{8} - \frac{\cos(2t)(-12 + t)}{4}$$

### ✓ Solution by Mathematica

Time used: 0.099 (sec). Leaf size: 26

```
DSolve[{y'[t]+4*y[t]==Sin[2*t],{y[0]==3,y'[0]==5}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{21}{8} \sin(2t) + \left(3 - \frac{t}{4}\right) \cos(2t)$$

## 18.7 problem 27.1 (g)

Internal problem ID [13535]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496

**Problem number:** 27.1 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y = 3 \operatorname{Heaviside}(t - 2)$$

With initial conditions

$$[y(0) = 0, y'(0) = 5]$$

### ✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 23

```
dsolve([diff(y(t),t$2)+4*y(t)=3*Heaviside(t-2),y(0) = 0, D(y)(0) = 5],y(t), singsol=all)
```

$$y(t) = \frac{3 \operatorname{Heaviside}(t - 2) \sin(t - 2)^2}{2} + \frac{5 \sin(2t)}{2}$$

### ✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 37

```
DSolve[{y''[t]+4*y[t]==UnitStep[t-2],{y[0]==0,y'[0]==5}},y[t],t,IncludeSingularSolutions ->
```

$$y(t) \rightarrow \begin{cases} 5 \cos(t) \sin(t) & t \leq 2 \\ \frac{1}{4}(-\cos(4 - 2t) + 10 \sin(2t) + 1) & \text{True} \end{cases}$$

## 18.8 problem 27.1 (h)

Internal problem ID [13536]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496

**Problem number:** 27.1 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 5y' + 6y = e^{4t}$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 23

```
dsolve([diff(y(t),t$2)+5*diff(y(t),t)+6*y(t)=exp(4*t),y(0) = 1, D(y)(0) = 0],y(t), singsol=a
```

$$y(t) = \frac{(e^{7t} + 119e^t - 78)e^{-3t}}{42}$$

### ✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 26

```
DSolve[{y'[t]+5*y'[t]+6*y[t]==Exp[4*t],{y[0]==1,y'[0]==0}},y[t],t,IncludeSingularSolutions
```

$$y(t) \rightarrow \frac{1}{42}e^{-3t}(119e^t + e^{7t} - 78)$$

## 18.9 problem 27.1 (i)

Internal problem ID [13537]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496

**Problem number:** 27.1 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 5y' + 6y = t^2 e^{4t}$$

With initial conditions

$$[y(0) = 0, y'(0) = 2]$$

### ✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 27

```
dsolve([diff(y(t),t$2)-5*diff(y(t),t)+6*y(t)=t^2*exp(4*t),y(0) = 0, D(y)(0) = 2],y(t), sings
```

$$y(t) = -\frac{7e^{2t}}{4} + \frac{(2t^2 - 6t + 7)e^{4t}}{4}$$

### ✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 32

```
DSolve[{y'[t]-5*y'[t]+6*y[t]==t^2*Exp[4*t],{y[0]==0,y'[0]==2}},y[t],t,IncludeSingularSoluti
```

$$y(t) \rightarrow \frac{1}{4}e^{2t}(e^{2t}(2t^2 - 6t + 7) - 7)$$

## 18.10 problem 27.1 (j)

Internal problem ID [13538]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496

**Problem number:** 27.1 (j).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 5y' + 6y = 7$$

With initial conditions

$$[y(0) = 2, y'(0) = 4]$$

### ✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 18

```
dsolve([diff(y(t),t$2)-5*diff(y(t),t)+6*y(t)=7,y(0) = 2, D(y)(0) = 4],y(t), singsol=all)
```

$$y(t) = \frac{7}{6} - \frac{3e^{2t}}{2} + \frac{7e^{3t}}{3}$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 25

```
DSolve[{y'[t]-5*y'[t]+6*y[t]==7,{y[0]==2,y'[0]==4}},y[t],t,IncludeSingularSolutions -> True
```

$$y(t) \rightarrow \frac{1}{6}(-9e^{2t} + 14e^{3t} + 7)$$

## 18.11 problem 27.1 (k)

Internal problem ID [13539]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496

**Problem number:** 27.1 (k).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y' + 13y = e^{2t} \sin(3t)$$

With initial conditions

$$[y(0) = 4, y'(0) = 3]$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 26

```
dsolve([diff(y(t),t$2)-4*diff(y(t),t)+13*y(t)=exp(2*t)*sin(3*t),y(0) = 4, D(y)(0) = 3],y(t),
```

$$y(t) = -\frac{e^{2t}(-24 + t) \cos(3t)}{6} - \frac{29 e^{2t} \sin(3t)}{18}$$

### ✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 61

```
DSolve[{y'[t]-4*y'[t]+13*y[t]==Exp(2*t)*Sin[3*t],{y[0]==4,y'[0]==3}},y[t],t,IncludeSingular
```

$$y(t) \rightarrow \frac{1}{600} \left( ((3\text{Exp} - 1000)e^{2t} + 3\text{Exp}(10t + 1)) \sin(3t) \right. \\ \left. + 6((400 - 9\text{Exp})e^{2t} + 3\text{Exp}(5t + 3)) \cos(3t) \right)$$



## 18.12 problem 27.1 (L)

Internal problem ID [13540]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496

**Problem number:** 27.1 (L).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y' + 13y = 4t + 2e^{2t} \sin(3t)$$

With initial conditions

$$[y(0) = 4, y'(0) = 3]$$

### ✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 47

```
dsolve([diff(y(t),t$2)+4*diff(y(t),t)+13*y(t)=4*t+2*exp(2*t)*sin(3*t),y(0) = 4, D(y)(0) = 3])
```

$$y(t) = -\frac{16}{169} + \frac{2 \cosh(2t) (346 \cos(3t) + 313 \sin(3t))}{169} + \frac{(-1423 \cos(3t) - 1226 \sin(3t)) \sinh(2t)}{338} + \frac{4t}{13}$$

### ✓ Solution by Mathematica

Time used: 1.331 (sec). Leaf size: 55

```
DSolve[{y''[t]+4*y'[t]+13*y[t]==4*t+2*Exp[2*t]*Sin[3*t],{y[0]==4,y'[0]==3}},y[t],t,IncludeS
```

$$y(t) \rightarrow \frac{1}{676} e^{-2t} (16e^{2t} (13t - 4) + (26e^{4t} + 2478) \sin(3t) + (2807 - 39e^{4t}) \cos(3t))$$

## 18.13 problem 27.1 (m)

Internal problem ID [13541]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496

**Problem number:** 27.1 (m).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$y''' - 27y = e^{-3t}$$

With initial conditions

$$[y(0) = 2, y'(0) = 3, y''(0) = 4]$$

### ✓ Solution by Maple

Time used: 0.093 (sec). Leaf size: 44

```
dsolve([diff(y(t),t$3)-27*y(t)=exp(-3*t),y(0) = 2, D(y)(0) = 3, (D@@2)(y)(0) = 4],y(t), sing
```

$$y(t) = \frac{14\sqrt{3}e^{-\frac{3t}{2}}\sin\left(\frac{3\sqrt{3}t}{2}\right)}{81} + \frac{70e^{-\frac{3t}{2}}\cos\left(\frac{3\sqrt{3}t}{2}\right)}{81} + \frac{92\cosh(3t)}{81} + \frac{95\sinh(3t)}{81}$$

### ✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y'[t]-27*y[t]==Exp[-3*t],{y[0]==4,y'[0]==3,y''[0]==4}},y[t],t,IncludeSingularSoluti
```

{}

## 18.14 problem 27.4

Internal problem ID [13542]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 27. Differentiation and the Laplace transform. Additional Exercises. page 496

**Problem number:** 27.4.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [Lienard]

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 7

```
dsolve([t*diff(y(t),t$2)+diff(y(t),t)+t*y(t)=0,y(0) = 1, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = \text{BesselJ}(0, t)$$

✓ Solution by Mathematica

Time used: 0.108 (sec). Leaf size: 8

```
DSolve[{t*y'[t]+y'[t]+t*y[t]==0,{y[0]==1,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True
```

$$y(t) \rightarrow \text{BesselJ}(0, t)$$

**19 Chapter 28. The inverse Laplace transform.  
Additional Exercises. page 509**

19.1 problem 28.6 (a)	683
19.2 problem 28.6 (b)	684
19.3 problem 28.6 (c)	685
19.4 problem 28.8 (a)	686
19.5 problem 28.8 (b)	687
19.6 problem 28.8 (c)	688
19.7 problem 28.8 (d)	689
19.8 problem 28.9 (a)	690
19.9 problem 28.9 (b)	691
19.10 problem 28.9 (c)	692
19.11 problem 28.9 (d)	693

## 19.1 problem 28.6 (a)

Internal problem ID [13543]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 28. The inverse Laplace transform. Additional Exercises. page 509

**Problem number:** 28.6 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 9y = 0$$

With initial conditions

$$[y(0) = 4, y'(0) = 9]$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 17

```
dsolve([diff(y(t),t$2)-9*y(t)=0,y(0) = 4, D(y)(0) = 9],y(t), singsol=all)
```

$$y(t) = \frac{7e^{3t}}{2} + \frac{e^{-3t}}{2}$$

### ✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 23

```
DSolve[{y'[t]-9*y[t]==0,{y[0]==4,y'[0]==9}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{2}e^{-3t}(7e^{6t} + 1)$$

## 19.2 problem 28.6 (b)

Internal problem ID [13544]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 28. The inverse Laplace transform. Additional Exercises. page 509

**Problem number:** 28.6 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = 27t^3$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.187 (sec). Leaf size: 19

```
dsolve([diff(y(t),t$2)+9*y(t)=27*t^3,y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = 3t^3 - 2t + \frac{2 \sin(3t)}{3}$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 22

```
DSolve[{y''[t]+9*y[t]==27*t^3,{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 3t^3 - 2t + \frac{2}{3} \sin(3t)$$

### 19.3 problem 28.6 (c)

Internal problem ID [13545]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 28. The inverse Laplace transform. Additional Exercises. page 509

**Problem number:** 28.6 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 8y' + 7y = 165e^{4t}$$

With initial conditions

$$[y(0) = 8, y'(0) = 1]$$

#### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 21

```
dsolve([diff(y(t),t$2)+8*diff(y(t),t)+7*y(t)=165*exp(4*t),y(0) = 8, D(y)(0) = 1],y(t), sings
```

$$y(t) = (3e^{11t} + 4e^{6t} + 1)e^{-7t}$$

#### ✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 25

```
DSolve[{y'[t]+8*y'[t]+7*y[t]==165*Exp[4*t],{y[0]==8,y'[0]==1}},y[t],t,IncludeSingularSoluti
```

$$y(t) \rightarrow e^{-7t} + 4e^{-t} + 3e^{4t}$$

## 19.4 problem 28.8 (a)

Internal problem ID [13546]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 28. The inverse Laplace transform. Additional Exercises. page 509

**Problem number:** 28.8 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 8y' + 17y = 0$$

With initial conditions

$$[y(0) = 3, y'(0) = 12]$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 12

```
dsolve([diff(y(t),t$2)-8*diff(y(t),t)+17*y(t)=0,y(0) = 3, D(y)(0) = 12],y(t), singsol=all)
```

$$y(t) = 3e^{4t} \cos(t)$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 14

```
DSolve[{y'[t]-8*y'[t]+17*y[t]==0,{y[0]==3,y'[0]==12}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 3e^{4t} \cos(t)$$



## 19.5 problem 28.8 (b)

Internal problem ID [13547]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 28. The inverse Laplace transform. Additional Exercises. page 509

**Problem number:** 28.8 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 6y' + 9y = e^{3t}t^2$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 13

```
dsolve([diff(y(t),t$2)-6*diff(y(t),t)+9*y(t)=exp(3*t)*t^2,y(0) = 0, D(y)(0) = 0],y(t), sings
```

$$y(t) = \frac{t^4 e^{3t}}{12}$$

### ✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 17

```
DSolve[{y'[t]-6*y'[t]+9*y[t]==Exp[3*t]*t^2,{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSoluti
```

$$y(t) \rightarrow \frac{1}{12} e^{3t} t^4$$

## 19.6 problem 28.8 (c)

Internal problem ID [13548]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 28. The inverse Laplace transform. Additional Exercises. page 509

**Problem number:** 28.8 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 6y' + 13y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = 8]$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 22

```
dsolve([diff(y(t),t$2)+6*diff(y(t),t)+13*y(t)=0,y(0) = 2, D(y)(0) = 8],y(t), singsol=all)
```

$$y(t) = e^{-3t}(2 \cos(2t) + 7 \sin(2t))$$

### ✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 24

```
DSolve[{y'[t]+6*y'[t]+13*y[t]==0,{y[0]==2,y'[0]==8}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{-3t}(7 \sin(2t) + 2 \cos(2t))$$

## 19.7 problem 28.8 (d)

Internal problem ID [13549]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 28. The inverse Laplace transform. Additional Exercises. page 509

**Problem number:** 28.8 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 8y' + 17y = 0$$

With initial conditions

$$[y(0) = 3, y'(0) = -12]$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 12

```
dsolve([diff(y(t),t$2)+8*diff(y(t),t)+17*y(t)=0,y(0) = 3, D(y)(0) = -12],y(t), singsol=all)
```

$$y(t) = 3e^{-4t} \cos(t)$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 14

```
DSolve[{y'[t]+8*y'[t]+17*y[t]==0,{y[0]==3,y'[0]==-12}},y[t],t,IncludeSingularSolutions -> T
```

$$y(t) \rightarrow 3e^{-4t} \cos(t)$$

## 19.8 problem 28.9 (a)

Internal problem ID [13550]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 28. The inverse Laplace transform. Additional Exercises. page 509

**Problem number:** 28.9 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$y'' = e^t \sin(t)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 15

```
dsolve([diff(y(t),t$2)=exp(t)*sin(t),y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = \frac{1}{2} - \frac{e^t \cos(t)}{2} + \frac{t}{2}$$

### ✓ Solution by Mathematica

Time used: 0.056 (sec). Leaf size: 19

```
DSolve[{y'[t]==Exp[t]*Sin[t],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{2}(t - e^t \cos(t) + 1)$$

## 19.9 problem 28.9 (b)

Internal problem ID [13551]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 28. The inverse Laplace transform. Additional Exercises. page 509

**Problem number:** 28.9 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 4y' + 40y = 122e^{-3t}$$

With initial conditions

$$[y(0) = 0, y'(0) = 8]$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 29

```
dsolve([diff(y(t),t$2)-4*diff(y(t),t)+40*y(t)=122*exp(-3*t),y(0) = 0, D(y)(0) = 8],y(t), sin
```

$$y(t) = -2 \left( -1 + \left( \cos(6t) - \frac{3 \sin(6t)}{2} \right) e^{5t} \right) e^{-3t}$$

### ✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 35

```
DSolve[{y'[t]-4*y'[t]+40*y[t]==122*Exp[-3*t],{y[0]==0,y'[0]==8}},y[t],t,IncludeSingularSolu
```

$$y(t) \rightarrow e^{-3t} (3e^{5t} \sin(6t) - 2e^{5t} \cos(6t) + 2)$$

## 19.10 problem 28.9 (c)

Internal problem ID [13552]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 28. The inverse Laplace transform. Additional Exercises. page 509

**Problem number:** 28.9 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 9y = 24e^{-3t}$$

With initial conditions

$$[y(0) = 6, y'(0) = 2]$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 22

```
dsolve([diff(y(t),t$2)-9*y(t)=24*exp(-3*t),y(0) = 6, D(y)(0) = 2],y(t), singsol=all)
```

$$y(t) = (-4t + 2)e^{-3t} + 4e^{3t}$$

### ✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 23

```
DSolve[{y''[t]-9*y[t]==24*Exp[-3*t],{y[0]==6,y'[0]==2}},y[t],t,IncludeSingularSolutions -> T
```

$$y(t) \rightarrow e^{-3t}(-4t + 4e^{6t} + 2)$$

## 19.11 problem 28.9 (d)

Internal problem ID [13553]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 28. The inverse Laplace transform. Additional Exercises. page 509

**Problem number:** 28.9 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 4y' + 13y = e^{2t} \sin(3t)$$

With initial conditions

$$[y(0) = 4, y'(0) = 3]$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 26

```
dsolve([diff(y(t),t$2)-4*diff(y(t),t)+13*y(t)=exp(2*t)*sin(3*t),y(0) = 4, D(y)(0) = 3],y(t),
```

$$y(t) = -\frac{e^{2t}(-24 + t) \cos(3t)}{6} - \frac{29 e^{2t} \sin(3t)}{18}$$

### ✓ Solution by Mathematica

Time used: 0.13 (sec). Leaf size: 30

```
DSolve[{y'[t]-4*y'[t]+13*y[t]==Exp[2*t]*Sin[3*t],{y[0]==4,y'[0]==3}},y[t],t,IncludeSingular
```

$$y(t) \rightarrow -\frac{1}{18}e^{2t}(29 \sin(3t) + 3(t - 24) \cos(3t))$$

**20 Chapter 29. Convolution. Additional Exercises.**  
**page 523**

20.1	problem 29.6 (a)	695
20.2	problem 29.6 (b)	696
20.3	problem 29.6 (c)	697
20.4	problem 29.6 (d)	698
20.5	problem 29.6 (e)	699
20.6	problem 29.7 (a)	700
20.7	problem 29.7 (b)	701
20.8	problem 29.7 (c)	702
20.9	problem 29.7 (d)	703
20.10	problem 29.7 (e)	704



## 20.1 problem 29.6 (a)

Internal problem ID [13554]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 29. Convolution. Additional Exercises. page 523

**Problem number:** 29.6 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 4y = 1$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 12

```
dsolve([diff(y(t),t$2)+4*y(t)=1,y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = \frac{1}{4} - \frac{\cos(2t)}{4}$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 13

```
DSolve[{y'[t]+4*y[t]==1,{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{\sin^2(t)}{2}$$

## 20.2 problem 29.6 (b)

Internal problem ID [13555]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 29. Convolution. Additional Exercises. page 523

**Problem number:** 29.6 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 4y = t$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 14

```
dsolve([diff(y(t),t$2)+4*y(t)=t,y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = -\frac{\sin(2t)}{8} + \frac{t}{4}$$

### ✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 17

```
DSolve[{y'[t]+4*y[t]==t,{y[0]==0,y'[0]==0}],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{4}(t - \sin(t) \cos(t))$$

## 20.3 problem 29.6 (c)

Internal problem ID [13556]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 29. Convolution. Additional Exercises. page 523

**Problem number:** 29.6 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 4y = e^{3t}$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.093 (sec). Leaf size: 24

```
dsolve([diff(y(t),t$2)+4*y(t)=exp(3*t),y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = -\frac{2 \cos(t)^2}{13} - \frac{3 \sin(t) \cos(t)}{13} + \frac{e^{3t}}{13} + \frac{1}{13}$$

### ✓ Solution by Mathematica

Time used: 0.1 (sec). Leaf size: 29

```
DSolve[{y'[t]+4*y[t]==Exp[3*t],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{26} (2e^{3t} - 3 \sin(2t) - 2 \cos(2t))$$

## 20.4 problem 29.6 (d)

Internal problem ID [13557]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 29. Convolution. Additional Exercises. page 523

**Problem number:** 29.6 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y = \sin(2t)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 18

```
dsolve([diff(y(t),t$2)+4*y(t)=sin(2*t),y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = \frac{\sin(2t)}{8} - \frac{\cos(2t)t}{4}$$

### ✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 21

```
DSolve[{y''[t]+4*y[t]==Sin[2*t],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{8}(\sin(2t) - 2t \cos(2t))$$

## 20.5 problem 29.6 (e)

Internal problem ID [13558]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 29. Convolution. Additional Exercises. page 523

**Problem number:** 29.6 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y = \sin(t)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 15

```
dsolve([diff(y(t),t$2)+4*y(t)=sin(t),y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = -\frac{\sin(2t)}{6} + \frac{\sin(t)}{3}$$

### ✓ Solution by Mathematica

Time used: 0.072 (sec). Leaf size: 15

```
DSolve[{y''[t]+4*y[t]==Sin[t],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -\frac{1}{3} \sin(t)(\cos(t) - 1)$$

## 20.6 problem 29.7 (a)

Internal problem ID [13559]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 29. Convolution. Additional Exercises. page 523

**Problem number:** 29.7 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 6y' + 9y = 1$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 17

```
dsolve([diff(y(t),t$2)-6*diff(y(t),t)+9*y(t)=1,y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = \frac{1}{9} + \frac{e^{3t}(3t - 1)}{9}$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 22

```
DSolve[{y''[t]-6*y'[t]+9*y[t]==1,{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True
```

$$y(t) \rightarrow \frac{1}{9}(e^{3t}(3t - 1) + 1)$$

## 20.7 problem 29.7 (b)

Internal problem ID [13560]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 29. Convolution. Additional Exercises. page 523

**Problem number:** 29.7 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 6y' + 9y = t$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 24

```
dsolve([diff(y(t),t$2)-6*diff(y(t),t)+9*y(t)=t,y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = \frac{2 \left( t \cosh \left( \frac{3t}{2} \right) - \frac{2 \sinh \left( \frac{3t}{2} \right)}{3} \right) e^{\frac{3t}{2}}}{9}$$

### ✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 25

```
DSolve[{y'[t]-6*y'[t]+9*y[t]==t,{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True
```

$$y(t) \rightarrow \frac{1}{27} (3t + e^{3t}(3t - 2) + 2)$$

## 20.8 problem 29.7 (c)

Internal problem ID [13561]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 29. Convolution. Additional Exercises. page 523

**Problem number:** 29.7 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 6y' + 9y = e^{3t}$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 13

```
dsolve([diff(y(t),t$2)-6*diff(y(t),t)+9*y(t)=exp(3*t),y(0) = 0, D(y)(0) = 0],y(t), singsol=a
```

$$y(t) = \frac{e^{3t}t^2}{2}$$

### ✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 17

```
DSolve[{y'[t]-6*y'[t]+9*y[t]==Exp[3*t],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions
```

$$y(t) \rightarrow \frac{1}{2}e^{3t}t^2$$



## 20.9 problem 29.7 (d)

Internal problem ID [13562]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 29. Convolution. Additional Exercises. page 523

**Problem number:** 29.7 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 6y' + 9y = e^{-3t}$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 23

```
dsolve([diff(y(t),t$2)-6*diff(y(t),t)+9*y(t)=exp(-3*t),y(0) = 0, D(y)(0) = 0],y(t), singsol=
```

$$y(t) = \frac{t \cosh(3t)}{6} + \frac{\sinh(3t)(3t - 1)}{18}$$

### ✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 27

```
DSolve[{y''[t]-6*y'[t]+9*y[t]==Exp[-3*t],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions
```

$$y(t) \rightarrow \frac{1}{36}e^{-3t}(e^{6t}(6t - 1) + 1)$$

## 20.10 problem 29.7 (e)

Internal problem ID [13563]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 29. Convolution. Additional Exercises. page 523

**Problem number:** 29.7 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 6y' + 9y = e^t$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 20

```
dsolve([diff(y(t),t$2)-6*diff(y(t),t)+9*y(t)=exp(t),y(0) = 0, D(y)(0) = 0],y(t), singsol=all
```

$$y(t) = \frac{e^t}{4} + \frac{e^{3t}(2t - 1)}{4}$$

### ✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 24

```
DSolve[{y''[t]-6*y'[t]+9*y[t]==Exp[t],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions ->
```

$$y(t) \rightarrow \frac{1}{4}(e^{3t}(2t - 1) + e^t)$$

**21 Chapter 30. Piecewise-defined functions and  
periodic functions. Additional Exercises. page  
553**

21.1	problem 30.6 (a)	706
21.2	problem 30.6 (b)	707
21.3	problem 30.6 (c)	708
21.4	problem 30.6 (d)	709
21.5	problem 30.6 (e)	710
21.6	problem 30.10 (a)	711
21.7	problem 30.10 (b)	713
21.8	problem 30.10 (c)	715

## 21.1 problem 30.6 (a)

Internal problem ID [13564]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 30. Piecewise-defined functions and periodic functions. Additional Exercises. page 553

**Problem number:** 30.6 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = \text{Heaviside}(t - 3)$$

With initial conditions

$$[y(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 12

```
dsolve([diff(y(t),t)=Heaviside(t-3),y(0) = 0],y(t), singsol=all)
```

$$y(t) = \text{Heaviside}(t - 3)(t - 3)$$

### ✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 15

```
DSolve[{y'[t]==UnitStep[t-3],{y[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \begin{cases} t - 3 & t > 3 \\ 0 & \text{True} \end{cases}$$

## 21.2 problem 30.6 (b)

Internal problem ID [13565]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 30. Piecewise-defined functions and periodic functions. Additional Exercises. page 553

**Problem number:** 30.6 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = \text{Heaviside}(t - 3)$$

With initial conditions

$$[y(0) = 4]$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 14

```
dsolve([diff(y(t),t)=Heaviside(t-3),y(0) = 4],y(t), singsol=all)
```

$$y(t) = \text{Heaviside}(t - 3)(t - 3) + 4$$

### ✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 15

```
DSolve[{y'[t]==UnitStep[t-3],{y[0]==4}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \begin{cases} 4 & t \leq 3 \\ t + 1 & \text{True} \end{cases}$$

## 21.3 problem 30.6 (c)

Internal problem ID [13566]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 30. Piecewise-defined functions and periodic functions. Additional Exercises. page 553

**Problem number:** 30.6 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$y'' = \text{Heaviside}(t - 2)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 15

```
dsolve([diff(y(t),t$2)=Heaviside(t-2),y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = \frac{\text{Heaviside}(t - 2)(t - 2)^2}{2}$$

### ✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 21

```
DSolve[{y''[t]==UnitStep[t-2],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \begin{cases} \frac{1}{2}(t - 2)^2 & t > 2 \\ 0 & \text{True} \end{cases}$$

## 21.4 problem 30.6 (d)

Internal problem ID [13567]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 30. Piecewise-defined functions and periodic functions. Additional Exercises. page 553

**Problem number:** 30.6 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$y'' = \text{Heaviside}(t - 2)$$

With initial conditions

$$[y(0) = 4, y'(0) = 6]$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 20

```
dsolve([diff(y(t),t$2)=Heaviside(t-2),y(0) = 4, D(y)(0) = 6],y(t), singsol=all)
```

$$y(t) = 4 + \frac{\text{Heaviside}(t - 2) (t - 2)^2}{2} + 6t$$

### ✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 28

```
DSolve[{y''[t]==UnitStep[t-2],{y[0]==4,y'[0]==6}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \begin{cases} 6t + 4 & t \leq 2 \\ \frac{t^2}{2} + 4t + 6 & \text{True} \end{cases}$$

## 21.5 problem 30.6 (e)

Internal problem ID [13568]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 30. Piecewise-defined functions and periodic functions. Additional Exercises. page 553

**Problem number:** 30.6 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = \text{Heaviside}(t - 10)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 18

```
dsolve([diff(y(t),t$2)+9*y(t)=Heaviside(t-10),y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = \frac{2 \text{Heaviside}(t - 10) \sin\left(\frac{3t}{2} - 15\right)^2}{9}$$

### ✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 26

```
DSolve[{y''[t]+9*y[t]==UnitStep[t-10],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions ->
```

$$y(t) \rightarrow \begin{cases} \frac{2}{9} \sin^2\left(15 - \frac{3t}{2}\right) & t > 10 \\ 0 & \text{True} \end{cases}$$



## 21.6 problem 30.10 (a)

Internal problem ID [13569]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 30. Piecewise-defined functions and periodic functions. Additional Exercises. page 553

**Problem number:** 30.10 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' = \begin{cases} 0 & t < 1 \\ 1 & 1 < t < 3 \\ 0 & 3 < t \end{cases}$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 19

```
dsolve([diff(y(t),t)=piecewise(t<1,0,1<t and t<3,1,t>3,0),y(0) = 0],y(t), singsol=all)
```

$$y(t) = \begin{cases} 0 & t < 1 \\ t - 1 & 1 < t < 3 \\ 2 & 3 \leq t \end{cases}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 23

```
DSolve[{y'[t]==Piecewise[{ {0,t<1},{1,1<t<3},{0,t>3}},{y[0]==0}],y[t],t,IncludeSingularSolu
```

$$y(t) \rightarrow \begin{cases} 0 & t \leq 1 \\ t - 1 & 1 < t \leq 3 \\ 2 & \text{True} \end{cases}$$

## 21.7 problem 30.10 (b)

Internal problem ID [13570]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 30. Piecewise-defined functions and periodic functions. Additional Exercises. page 553

**Problem number:** 30.10 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$y'' = \begin{cases} 0 & t < 1 \\ 1 & 1 < t < 3 \\ 0 & 3 < t \end{cases}$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 37

```
dsolve([diff(y(t),t$2)=piecewise(t<1,0,1<t and t<3,1,t>3,0),y(0) = 0, D(y)(0) = 0],y(t), sin
```

$$y(t) = \begin{cases} 0 & t < 1 \\ \frac{(t-1)^2}{2} & 1 < t < 3 \\ 2t - 4 & 3 \leq t \end{cases}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 33

```
DSolve[{y''[t]==Piecewise[{ {0,t<1},{1,1<t<3},{0,t>3}},{y[0]==0,y'[0]==0}],y[t],t,IncludeSi
```

$$y(t) \rightarrow \begin{cases} 0 & t \leq 1 \\ \frac{1}{2}(t-1)^2 & 1 < t \leq 3 \\ 2(t-2) & \text{True} \end{cases}$$

## 21.8 problem 30.10 (c)

Internal problem ID [13571]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 30. Piecewise-defined functions and periodic functions. Additional Exercises. page 553

**Problem number:** 30.10 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = \begin{cases} 0 & t < 1 \\ 1 & 1 < t < 3 \\ 0 & 3 < t \end{cases}$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

✓ Solution by Maple

Time used: 0.328 (sec). Leaf size: 46

```
dsolve([diff(y(t),t$2)+9*y(t)=piecewise(t<1,0,1<t and t<3,1,t>3,0),y(0) = 0, D(y)(0) = 0],y(t))
```

$$y(t) = \frac{2 \left( \begin{cases} 0 & t < 1 \\ \sin \left( \frac{3t}{2} - \frac{3}{2} \right)^2 & t < 3 \\ -\cos \left( \frac{3t}{2} - \frac{3}{2} \right)^2 + \cos \left( \frac{3t}{2} - \frac{9}{2} \right)^2 & 3 \leq t \end{cases} \right)}{9}$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 47

```
DSolve[{y''[t]+9*y[t]==Piecewise[{0,t<1},{1,1<t<3},{0,t>3}],{y[0]==0,y'[0]==0}},y[t],t,In
```

$$y(t) \rightarrow \begin{cases} \frac{2}{9} \sin^2\left(\frac{3}{2} - \frac{3t}{2}\right) & 1 < t \leq 3 \\ -\frac{2}{9} \sin(3) \sin(6 - 3t) & t > 3 \end{cases}$$

## 22 Chapter 31. Delta Functions. Additional Exercises. page 572

22.1 problem 31.6 (a)	718
22.2 problem 31.6 (b)	719
22.3 problem 31.6 (c)	720
22.4 problem 31.6 (d)	721
22.5 problem 31.6 (e)	722
22.6 problem 31.6 (f)	723
22.7 problem 31.6 (g)	724
22.8 problem 31.7 (a)	725
22.9 problem 31.7 (b)	726
22.10 problem 31.7 (c)	727
22.11 problem 31.7 (d)	728
22.12 problem 31.7 (e)	729
22.13 problem 31.7 (f)	730
22.14 problem 31.7 (g)	731
22.15 problem 31.7 (h)	732
22.16 problem 31.7 (i)	733
22.17 problem 31.7 (j)	734
22.18 problem 31.7 (k)	735
22.19 problem 31.7 (L)	736

## 22.1 problem 31.6 (a)

Internal problem ID [13572]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.6 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' = 3\delta(t - 2)$$

With initial conditions

$$[y(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 10

```
dsolve([diff(y(t),t)=3*Dirac(t-2),y(0) = 0],y(t), singsol=all)
```

$$y(t) = 3 \text{Heaviside}(t - 2)$$

### ✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 11

```
DSolve[{y'[t]==3*DiracDelta[t-2],{y[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 3\theta(t - 2)$$



## 22.2 problem 31.6 (b)

Internal problem ID [13573]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.6 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [quadrature]

$$y' = \delta(t - 2) - \delta(t - 4)$$

With initial conditions

$$[y(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 15

```
dsolve([diff(y(t),t)=Dirac(t-2)-Dirac(t-4),y(0) = 0],y(t), singsol=all)
```

$$y(t) = -\text{Heaviside}(t - 4) + \text{Heaviside}(t - 2)$$

### ✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 16

```
DSolve[{y'[t]==DiracDelta[t-2]-DiracDelta[t-4],{y[0]==0}},y[t],t,IncludeSingularSolutions ->
```

$$y(t) \rightarrow \theta(t - 2) - \theta(t - 4)$$

## 22.3 problem 31.6 (c)

Internal problem ID [13574]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.6 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$y'' = \delta(t - 3)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 12

```
dsolve([diff(y(t),t$2)=Dirac(t-3),y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = \text{Heaviside}(t - 3)(t - 3)$$

### ✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 13

```
DSolve[{y'[t]==DiracDelta[t-3],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow (t - 3)\theta(t - 3)$$

## 22.4 problem 31.6 (d)

Internal problem ID [13575]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.6 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _quadrature]]`

$$y'' = \delta(-1 + t) - \delta(t - 4)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 22

```
dsolve([diff(y(t),t$2)=Dirac(t-1)-Dirac(t-4),y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = (-t + 4) \text{Heaviside}(t - 4) + \text{Heaviside}(t - 1)(t - 1)$$

### ✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 23

```
DSolve[{y'[t]==DiracDelta[t-1]-DiracDelta[t-4],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions->True]
```

$$y(t) \rightarrow (t - 1)\theta(t - 1) - (t - 4)\theta(t - 4)$$

## 22.5 problem 31.6 (e)

Internal problem ID [13576]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.6 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 2y = 4\delta(-1 + t)$$

With initial conditions

$$[y(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 16

```
dsolve([diff(y(t),t)+2*y(t)=4*Dirac(t-1),y(0) = 0],y(t), singsol=all)
```

$$y(t) = 4 \operatorname{Heaviside}(t - 1) e^{2-2t}$$

### ✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 18

```
DSolve[{y'[t]+2*y[t]==4*DiracDelta[t-1],{y[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 4e^{2-2t}\theta(t - 1)$$

## 22.6 problem 31.6 (f)

Internal problem ID [13577]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.6 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \delta(t) + \delta(t - \pi)$$

### ✓ Solution by Maple

Time used: 0.187 (sec). Leaf size: 22

```
dsolve(diff(y(t),t$2)+y(t)=Dirac(t)+Dirac(t-Pi),y(t), singsol=all)
```

$$y = \cos(t) y(0) + \sin(t) (\text{Heaviside}(-t + \pi) + y'(0))$$

### ✓ Solution by Mathematica

Time used: 0.118 (sec). Leaf size: 92

```
DSolve[y''[t]+2*y[t]==DiracDelta[t]+DiracDelta[t-Pi],y[t],t,IncludeSingularSolutions -> True
```

$$y(t) \rightarrow -\frac{\theta(t - \pi) \sin(\sqrt{2}(\pi - t))}{\sqrt{2}} + \frac{\theta(t) \sin(\sqrt{2}t)}{\sqrt{2}} - \frac{\cos(\sqrt{2}\pi) \sin(\sqrt{2}t)}{\sqrt{2}} + c_1 \cos(\sqrt{2}t) + c_2 \sin(\sqrt{2}t)$$

## 22.7 problem 31.6 (g)

Internal problem ID [13578]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.6 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = -2\delta\left(t - \frac{\pi}{2}\right)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 14

```
dsolve([diff(y(t),t$2)+y(t)=-2*Dirac(t-Pi/2),y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = 2 \operatorname{Heaviside}\left(t - \frac{\pi}{2}\right) \cos(t)$$

### ✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 17

```
DSolve[{y''[t]+y[t]==-2*DiracDelta[t-Pi/2],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions->True]
```

$$y(t) \rightarrow 2\theta(2t - \pi) \cos(t)$$

## 22.8 problem 31.7 (a)

Internal problem ID [13579]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.7 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 3y = \delta(t - 2)$$

With initial conditions

$$[y(0) = 2]$$

### ✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 22

```
dsolve([diff(y(t),t)+3*y(t)=Dirac(t-2),y(0) = 2],y(t), singsol=all)
```

$$y(t) = \text{Heaviside}(t - 2) e^{6-3t} + 2 e^{-3t}$$

### ✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 21

```
DSolve[{y'[t]+3*y[t]==DiracDelta[t-2],{y[0]==2}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{-3t}(e^6 \theta(t - 2) + 2)$$

## 22.9 problem 31.7 (b)

Internal problem ID [13580]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.7 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' + 3y' = \delta(t)$$

### ✓ Solution by Maple

Time used: 0.157 (sec). Leaf size: 22

```
dsolve(diff(y(t),t$2)+3*diff(y(t),t)=Dirac(t),y(t), singsol=all)
```

$$y = \frac{1}{3} + \frac{y'(0)}{3} + y(0) - \frac{e^{-3t}(1 + y'(0))}{3}$$

### ✓ Solution by Mathematica

Time used: 0.138 (sec). Leaf size: 27

```
DSolve[y''[t]+3*y'[t]==DiracDelta[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{3}(\theta(t) - e^{-3t}(\theta(t) + c_1)) + c_2$$



## 22.10 problem 31.7 (c)

Internal problem ID [13581]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.7 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$y'' + 3y' = \delta(-1 + t)$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 28

```
dsolve([diff(y(t),t$2)+3*diff(y(t),t)=Dirac(t-1),y(0) = 0, D(y)(0) = 1],y(t), singsol=all)
```

$$y(t) = -\frac{\text{Heaviside}(t-1)e^{-3t+3}}{3} + \frac{\text{Heaviside}(t-1)}{3} + \frac{1}{3} - \frac{e^{-3t}}{3}$$

### ✓ Solution by Mathematica

Time used: 0.14 (sec). Leaf size: 37

```
DSolve[{y''[t]+3*y'[t]==DiracDelta[t-1],{y[0]==0,y'[0]==1}},y[t],t,IncludeSingularSolutions
```

$$y(t) \rightarrow \frac{1}{3}e^{-3t}((e^{3t} - e^3)\theta(t-1) + e^{3t} - 1)$$

## 22.11 problem 31.7 (d)

Internal problem ID [13582]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.7 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 16y = \delta(t - 2)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 16

```
dsolve([diff(y(t),t$2)+16*y(t)=Dirac(t-2),y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = \frac{\text{Heaviside}(t - 2) \sin(-8 + 4t)}{4}$$

### ✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 19

```
DSolve[{y''[t]+16*y[t]==DiracDelta[t-2],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions
```

$$y(t) \rightarrow -\frac{1}{4}\theta(t - 2) \sin(8 - 4t)$$

## 22.12 problem 31.7 (e)

Internal problem ID [13583]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.7 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 16y = \delta(t - 10)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 16

```
dsolve([diff(y(t),t$2)-16*y(t)=Dirac(t-10),y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = \frac{\text{Heaviside}(t - 10) \sinh(-40 + 4t)}{4}$$

### ✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 31

```
DSolve[{y''[t]-16*y[t]==DiracDelta[t-10],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions
```

$$y(t) \rightarrow \frac{1}{8} e^{-4(t+10)} (e^{8t} - e^{80}) \theta(t - 10)$$

## 22.13 problem 31.7 (f)

Internal problem ID [13584]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.7 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + y = \delta(t)$$

With initial conditions

$$[y(0) = 0, y'(0) = -1]$$

### ✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 5

```
dsolve([diff(y(t),t$2)+y(t)=Dirac(t),y(0) = 0, D(y)(0) = -1],y(t), singsol=all)
```

$$y(t) = 0$$

### ✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 16

```
DSolve[{y'[t]+y[t]==DiracDelta[t],{y[0]==0,y'[0]==-1}},y[t],t,IncludeSingularSolutions -> T
```

$$y(t) \rightarrow (\theta(t) - \theta(0) - 1) \sin(t)$$

## 22.14 problem 31.7 (g)

Internal problem ID [13585]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.7 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y' - 12y = \delta(t)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 14

```
dsolve([diff(y(t),t$2)+4*diff(y(t),t)-12*y(t)=Dirac(t),y(0) = 0, D(y)(0) = 0],y(t), singsol=
```

$$y(t) = \frac{e^{-2t} \sinh(4t)}{4}$$

### ✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 28

```
DSolve[{y''[t]+4*y'[t]-12*y[t]==DiracDelta[t],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolu
```

$$y(t) \rightarrow -\frac{1}{8}e^{-6t}(e^{8t} - 1)(\theta(0) - \theta(t))$$

## 22.15 problem 31.7 (h)

Internal problem ID [13586]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.7 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y' - 12y = \delta(t - 3)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 22

```
dsolve([diff(y(t),t$2)+4*diff(y(t),t)-12*y(t)=Dirac(t-3),y(0) = 0, D(y)(0) = 0],y(t), singso
```

$$y(t) = \frac{\text{Heaviside}(t - 3) e^{-2t+6} \sinh(4t - 12)}{4}$$

### ✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 31

```
DSolve[{y''[t]+4*y'[t]-12*y[t]==DiracDelta[t-3],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSo
```

$$y(t) \rightarrow \frac{1}{8} e^{-6(t+1)} (e^{8t} - e^{24}) \theta(t - 3)$$

## 22.16 problem 31.7 (i)

Internal problem ID [13587]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.7 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 6y' + 9y = \delta(t - 4)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 18

```
dsolve([diff(y(t),t$2)+6*diff(y(t),t)+9*y(t)=Dirac(t-4),y(0) = 0, D(y)(0) = 0],y(t), singsol
```

$$y(t) = (t - 4)e^{-3t+12} \text{Heaviside}(t - 4)$$

### ✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 20

```
DSolve[{y'[t]+6*y'[t]+9*y[t]==DiracDelta[t-4],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSol
```

$$y(t) \rightarrow e^{12-3t}(t - 4)\theta(t - 4)$$

## 22.17 problem 31.7 (j)

Internal problem ID [13588]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.7 (j).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 12y' + 45y = \delta(t)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 14

```
dsolve([diff(y(t),t$2)-12*diff(y(t),t)+45*y(t)=Dirac(t),y(0) = 0, D(y)(0) = 0],y(t), singsol
```

$$y(t) = \frac{e^{6t} \sin(3t)}{3}$$

### ✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 25

```
DSolve[{y'[t]-12*y'[t]+45*y[t]==DiracDelta[t],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSol
```

$$y(t) \rightarrow -\frac{1}{3}e^{6t}(\theta(0) - \theta(t)) \sin(3t)$$



## 22.18 problem 31.7 (k)

Internal problem ID [13589]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.7 (k).

**ODE order:** 3.

**ODE degree:** 1.

CAS Maple gives this as type `[[_3rd_order, _missing_y]]`

$$y''' + 9y' = \delta(-1 + t)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0, y''(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 18

```
dsolve([diff(y(t),t$3)+9*diff(y(t),t)=Dirac(t-1),y(0) = 0, D(y)(0) = 0, (D@@2)(y)(0) = 0],y(t))
```

$$y(t) = \frac{2 \operatorname{Heaviside}(t - 1) \sin\left(\frac{3t}{2} - \frac{3}{2}\right)^2}{9}$$

### ✓ Solution by Mathematica

Time used: 0.174 (sec). Leaf size: 21

```
DSolve[{y'''[t]+9*y'[t]==DiracDelta[t-1],{y[0]==0,y'[0]==0,y''[0]==0}},y[t],t,IncludeSingularSolutions->True]
```

$$y(t) \rightarrow -\frac{1}{9}\theta(t - 1)(\cos(3 - 3t) - 1)$$

## 22.19 problem 31.7 (L)

Internal problem ID [13590]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 31. Delta Functions. Additional Exercises. page 572

**Problem number:** 31.7 (L).

**ODE order:** 4.

**ODE degree:** 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 16y = \delta(t)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0, y''(0) = 0, y'''(0) = 0]$$

### ✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 17

```
dsolve([diff(y(t),t$4)-16*y(t)=Dirac(t),y(0) = 0, D(y)(0) = 0, (D@@2)(y)(0) = 0, (D@@3)(y)(0) = 0],y(t),t,Incl
```

$$y(t) = -\frac{\sin(2t)}{16} + \frac{\sinh(2t)}{16}$$

### ✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 39

```
DSolve[{y''''[t]-16*y[t]==DiracDelta[t],{y[0]==0,y'[0]==0,y''[0]==0,y'''[0]==0}},y[t],t,Incl
```

$$y(t) \rightarrow -\frac{1}{32}e^{-2t}(\theta(0) - \theta(t))(e^{4t} - 2e^{2t}\sin(2t) - 1)$$

## 23 Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises.

page 641

23.1 problem 33.3 (a)	738
23.2 problem 33.3 (b)	739
23.3 problem 33.3 (c)	740
23.4 problem 33.3 (d)	741
23.5 problem 33.3 (e)	742
23.6 problem 33.3 (f)	743
23.7 problem 33.3 (g)	744
23.8 problem 33.3 (h)	745
23.9 problem 33.3 (i)	746
23.10 problem 33.3 (j)	747
23.11 problem 33.3 (k)	748
23.12 problem 33.3 (L)	749
23.13 problem 33.5 (a)	750
23.14 problem 33.5 (b)	751
23.15 problem 33.5 (c)	752
23.16 problem 33.5 (d)	753
23.17 problem 33.5 (e)	754
23.18 problem 33.5 (f)	755
23.19 problem 33.5 (g)	756
23.20 problem 33.5 (h)	757
23.21 problem 33.5 (i)	758
23.22 problem 33.5 (j)	759
23.23 problem 33.5 (k)	760
23.24 problem 33.5 (L)	761
23.25 problem 33.9	762
23.26 problem 33.10	763
23.27 problem 33.11 (a)	764
23.28 problem 33.11 (b)	765
23.29 problem 33.11 (c)	766
23.30 problem 33.11 (d)	767
23.31 problem 33.11 (e)	768
23.32 problem 33.11 (f)	769
23.33 problem 33.11 (g)	770
23.34 problem 33.11 (h)	771

## 23.1 problem 33.3 (a)

Internal problem ID [13591]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.3 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' - 2y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
Order:=6;  
dsolve(diff(y(x),x)-2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + 2x + 2x^2 + \frac{4}{3}x^3 + \frac{2}{3}x^4 + \frac{4}{15}x^5\right) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 37

```
AsymptoticDSolveValue[y'[x]-2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{4x^5}{15} + \frac{2x^4}{3} + \frac{4x^3}{3} + 2x^2 + 2x + 1 \right)$$

## 23.2 problem 33.3 (b)

Internal problem ID [13592]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.3 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_separable`]

$$y' - 2yx = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
Order:=6;  
dsolve(diff(y(x),x)-2*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + x^2 + \frac{1}{2}x^4\right) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 18

```
AsymptoticDSolveValue[y'[x]-2*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^4}{2} + x^2 + 1\right)$$

## 23.3 problem 33.3 (c)

Internal problem ID [13593]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.3 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' + \frac{2y}{2x-1} = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
Order:=6;  
dsolve(diff(y(x),x)+2/(2*x-1)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (32x^5 + 16x^4 + 8x^3 + 4x^2 + 2x + 1) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 31

```
AsymptoticDSolveValue[y'[x]+2/(2*x-1)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(32x^5 + 16x^4 + 8x^3 + 4x^2 + 2x + 1)$$

## 23.4 problem 33.3 (d)

Internal problem ID [13594]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.3 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$(x - 3)y' - 2y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
Order:=6;  
dsolve((x-3)*diff(y(x),x)-2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{y(0)(x-3)^2}{9}$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 20

```
AsymptoticDSolveValue[(x-3)*y'[x]-2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{x^2}{9} - \frac{2x}{3} + 1 \right)$$

## 23.5 problem 33.3 (e)

Internal problem ID [13595]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.3 (e).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$(x^2 + 1) y' - 2yx = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
Order:=6;  
dsolve((1+x^2)*diff(y(x),x)-2*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = y(0) (x^2 + 1)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 11

```
AsymptoticDSolveValue[(1+x^2)*y'[x]-2*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 (x^2 + 1)$$



## 23.6 problem 33.3 (f)

Internal problem ID [13596]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.3 (f).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' + \frac{y}{x-1} = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
Order:=6;  
dsolve(diff(y(x),x)+1/(x-1)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (x^5 + x^4 + x^3 + x^2 + x + 1) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 21

```
AsymptoticDSolveValue[y'[x]+1/(x-1)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(x^5 + x^4 + x^3 + x^2 + x + 1)$$

## 23.7 problem 33.3 (g)

Internal problem ID [13597]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.3 (g).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' + \frac{y}{x-1} = 0$$

With the expansion point for the power series method at  $x = 3$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
Order:=6;  
dsolve(diff(y(x),x)+1/(x-1)*y(x)=0,y(x),type='series',x=3);
```

$$y(x) = \left( \frac{5}{2} - \frac{x}{2} + \frac{(x-3)^2}{4} - \frac{(x-3)^3}{8} + \frac{(x-3)^4}{16} - \frac{(x-3)^5}{32} \right) y(3) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 53

```
AsymptoticDSolveValue[y'[x]+1/(x-1)*y[x]==0,y[x],{x,3,5}]
```

$$y(x) \rightarrow c_1 \left( -\frac{1}{32}(x-3)^5 + \frac{1}{16}(x-3)^4 - \frac{1}{8}(x-3)^3 + \frac{1}{4}(x-3)^2 + \frac{3-x}{2} + 1 \right)$$

## 23.8 problem 33.3 (h)

Internal problem ID [13598]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.3 (h).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$(1-x)y' - 2y = 0$$

With the expansion point for the power series method at  $x = 5$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
Order:=6;  
dsolve((1-x)*diff(y(x),x)-2*y(x)=0,y(x),type='series',x=5);
```

$$y(x) = \left( \frac{7}{2} - \frac{x}{2} + \frac{3(x-5)^2}{16} - \frac{(x-5)^3}{16} + \frac{5(x-5)^4}{256} - \frac{3(x-5)^5}{512} \right) y(5) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 53

```
AsymptoticDSolveValue[(1-x)*y'[x]-2*y[x]==0,y[x],{x,5,5}]
```

$$y(x) \rightarrow c_1 \left( -\frac{3}{512}(x-5)^5 + \frac{5}{256}(x-5)^4 - \frac{1}{16}(x-5)^3 + \frac{3}{16}(x-5)^2 + \frac{5-x}{2} + 1 \right)$$

## 23.9 problem 33.3 (i)

Internal problem ID [13599]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.3 (i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [separable]

$$(-x^3 + 2)y' - 3yx^2 = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
Order:=6;  
dsolve((2-x^3)*diff(y(x),x)-3*x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{x^3}{2}\right)y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 15

```
AsymptoticDSolveValue[(2-x^3)*y'[x]-3*x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^3}{2} + 1\right)$$

## 23.10 problem 33.3 (j)

Internal problem ID [13600]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.3 (j).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$(-x^3 + 2)y' + 3yx^2 = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;  
dsolve((2-x^3)*diff(y(x),x)+3*x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = y(0) \left(1 - \frac{x^3}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 15

```
AsymptoticDSolveValue[(2-x^3)*y'[x]+3*x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(1 - \frac{x^3}{2}\right)$$

## 23.11 problem 33.3 (k)

Internal problem ID [13601]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.3 (k).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$(1+x)y' - yx = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
Order:=6;  
dsolve((1+x)*diff(y(x),x)-x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 - \frac{1}{3}x^3 + \frac{3}{8}x^4 - \frac{11}{30}x^5\right) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 36

```
AsymptoticDSolveValue[(1+x)*y'[x]-x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( -\frac{11x^5}{30} + \frac{3x^4}{8} - \frac{x^3}{3} + \frac{x^2}{2} + 1 \right)$$

## 23.12 problem 33.3 (L)

Internal problem ID [13602]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.3 (L).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$(1+x)y' + (1-x)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
Order:=6;  
dsolve((1+x)*diff(y(x),x)+(1-x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x + \frac{3}{2}x^2 - \frac{11}{6}x^3 + \frac{53}{24}x^4 - \frac{103}{40}x^5\right) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 39

```
AsymptoticDSolveValue[(1+x)*y'[x]+(1-x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( -\frac{103x^5}{40} + \frac{53x^4}{24} - \frac{11x^3}{6} + \frac{3x^2}{2} - x + 1 \right)$$

### 23.13 problem 33.5 (a)

Internal problem ID [13603]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.5 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x^2 + 1)y'' - 2y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
Order:=6;  
dsolve((1+x^2)*diff(y(x),x$2)-2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = y(0)(x^2 + 1) + \left(x + \frac{1}{3}x^3 - \frac{1}{15}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 31

```
AsymptoticDSolveValue[(1+x^2)*y''[x]-2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(x^2 + 1) + c_2\left(-\frac{x^5}{15} + \frac{x^3}{3} + x\right)$$



## 23.14 problem 33.5 (b)

Internal problem ID [13604]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.5 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$y'' + y'x + y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)+x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4\right) y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[y'[x]+x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{x^5}{15} - \frac{x^3}{3} + x \right) + c_1 \left( \frac{x^4}{8} - \frac{x^2}{2} + 1 \right)$$

## 23.15 problem 33.5 (c)

Internal problem ID [13605]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.5 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$(x^2 + 4)y'' + 2y'x = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve((4+x^2)*diff(y(x),x$2)+2*x*diff(y(x),x)=0,y(x),type='series',x=0);
```

$$y(x) = y(0) + \left(x - \frac{1}{12}x^3 + \frac{1}{80}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 25

```
AsymptoticDSolveValue[(4+x^2)*y''[x]+2*x*y'[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{x^5}{80} - \frac{x^3}{12} + x \right) + c_1$$

## 23.16 problem 33.5 (d)

Internal problem ID [13606]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.5 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y'' - 3yx^2 = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)-3*x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{x^4}{4}\right) y(0) + \left(x + \frac{3}{20}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y''[x]-3*x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{3x^5}{20} + x \right) + c_1 \left( \frac{x^4}{4} + 1 \right)$$

## 23.17 problem 33.5 (e)

Internal problem ID [13607]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.5 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(-x^2 + 4)y'' - 5y'x - 3y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve((4-x^2)*diff(y(x),x$2)-5*x*diff(y(x),x)-3*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{3}{8}x^2 + \frac{15}{128}x^4\right)y(0) + \left(x + \frac{1}{3}x^3 + \frac{1}{10}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[(4-x^2)*y''[x]-5*x*y'[x]-3*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{x^5}{10} + \frac{x^3}{3} + x \right) + c_1 \left( \frac{15x^4}{128} + \frac{3x^2}{8} + 1 \right)$$

## 23.18 problem 33.5 (f)

Internal problem ID [13608]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.5 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [\_Gegenbauer, [\_2nd\_order, \_linear, ‘\_with\_symmetry\_[0,F(x)]’]

$$(-x^2 + 1)y'' - y'x + 4y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
Order:=6;  
dsolve((1-x^2)*diff(y(x),x$2)-x*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (-2x^2 + 1)y(0) + \left(x - \frac{1}{2}x^3 - \frac{1}{8}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 33

```
AsymptoticDSolveValue[(1-x^2)*y''[x]-x*y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(1 - 2x^2) + c_2\left(-\frac{x^5}{8} - \frac{x^3}{2} + x\right)$$

## 23.19 problem 33.5 (g)

Internal problem ID [13609]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.5 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 2y'x + 6y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
Order:=6;  
dsolve(diff(y(x),x$2)-2*x*diff(y(x),x)+6*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - 3x^2 + \frac{1}{2}x^4\right) y(0) + \left(-\frac{2}{3}x^3 + x\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 33

```
AsymptoticDSolveValue[y''[x]-2*x*y'[x]+6*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{2x^3}{3}\right) + c_1 \left(\frac{x^4}{2} - 3x^2 + 1\right)$$

## 23.20 problem 33.5 (h)

Internal problem ID [13610]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.5 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x^2 - 6x)y'' + 4(x - 3)y' + 2y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

Order:=6;

```
dsolve((x^2-6*x)*diff(y(x),x$2)+4*(x-3)*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 \left( 1 + \frac{1}{6}x + \frac{1}{36}x^2 + \frac{1}{216}x^3 + \frac{1}{1296}x^4 + \frac{1}{7776}x^5 + O(x^6) \right) \\ + \frac{c_2 \left( 1 + \frac{1}{3}x + \frac{1}{18}x^2 + \frac{1}{108}x^3 + \frac{1}{648}x^4 + \frac{1}{3888}x^5 + O(x^6) \right)}{x}$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 64

```
AsymptoticDSolveValue[(x^2-6*x)*y'[x]+4*(x-3)*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{x^3}{1296} + \frac{x^2}{216} + \frac{x}{36} + \frac{1}{x} + \frac{1}{6} \right) + c_2 \left( \frac{x^4}{1296} + \frac{x^3}{216} + \frac{x^2}{36} + \frac{x}{6} + 1 \right)$$

## 23.21 problem 33.5 (i)

Internal problem ID [13611]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.5 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + (x + 2)y' + 2y = 0$$

With the expansion point for the power series method at  $x = -2$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)+(x+2)*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=-2);
```

$$y(x) = \left(1 - (x + 2)^2 + \frac{(x + 2)^4}{3}\right) y(-2) \\ + \left(x + 2 - \frac{(x + 2)^3}{2} + \frac{(x + 2)^5}{8}\right) D(y)(-2) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 49

```
AsymptoticDSolveValue[y'[x]+(x+2)*y'[x]+2*y[x]==0,y[x],{x,-2,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{1}{3}(x + 2)^4 - (x + 2)^2 + 1 \right) + c_2 \left( \frac{1}{8}(x + 2)^5 - \frac{1}{2}(x + 2)^3 + x + 2 \right)$$



## 23.22 problem 33.5 (j)

Internal problem ID [13612]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.5 (j).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x^2 - 2x + 2)y'' + (1 - x)y' - 3y = 0$$

With the expansion point for the power series method at  $x = 1$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

Order:=6;

```
dsolve((x^2-2*x+2)*diff(y(x),x$2)+(1-x)*diff(y(x),x)-3*y(x)=0,y(x),type='series',x=1);
```

$$y(x) = \left(1 + \frac{3(x-1)^2}{2} + \frac{3(x-1)^4}{8}\right) y(1) + \left(x-1 + \frac{2(x-1)^3}{3}\right) D(y)(1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[(x^2-2*x+2)*y'[x]+(1-x)*y'[x]-3*y[x]==0,y[x],{x,1,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{3}{8}(x-1)^4 + \frac{3}{2}(x-1)^2 + 1 \right) + c_2 \left( \frac{2}{3}(x-1)^3 + x-1 \right)$$

## 23.23 problem 33.5 (k)

Internal problem ID [13613]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.5 (k).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 2y' - yx = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
Order:=6;  
dsolve(diff(y(x),x$2)-2*diff(y(x),x)-x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{1}{30}x^5\right) y(0) + \left(x + x^2 + \frac{2}{3}x^3 + \frac{5}{12}x^4 + \frac{13}{60}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 59

```
AsymptoticDSolveValue[y''[x]-2*y'[x]-x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{x^5}{30} + \frac{x^4}{12} + \frac{x^3}{6} + 1 \right) + c_2 \left( \frac{13x^5}{60} + \frac{5x^4}{12} + \frac{2x^3}{3} + x^2 + x \right)$$

## 23.24 problem 33.5 (L)

Internal problem ID [13614]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.5 (L).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - y'x - 2yx = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
Order:=6;  
dsolve(diff(y(x),x$2)-x*diff(y(x),x)-2*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{3}x^3 + \frac{1}{20}x^5\right) y(0) + \left(x + \frac{1}{6}x^3 + \frac{1}{6}x^4 + \frac{1}{40}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 49

```
AsymptoticDSolveValue[y''[x]-x*y'[x]-2*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{x^5}{20} + \frac{x^3}{3} + 1 \right) + c_2 \left( \frac{x^5}{40} + \frac{x^4}{6} + \frac{x^3}{6} + x \right)$$

## 23.25 problem 33.9

Internal problem ID [13615]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.9.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [\_Gegenbauer, [\_2nd\_order, \_linear, ‘\_with\_symmetry\_[0,F(x)]’]

$$(-x^2 + 1)y'' - y'x + \lambda y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 63

```
Order:=6;  
dsolve((1-x^2)*diff(y(x),x$2)-x*diff(y(x),x)+lambda*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{\lambda x^2}{2} + \frac{\lambda(\lambda - 4)x^4}{24}\right) y(0) + \left(x - \frac{(\lambda - 1)x^3}{6} + \frac{(\lambda - 1)(\lambda - 9)x^5}{120}\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 80

```
AsymptoticDSolveValue[(1-x^2)*y'[x]-x*y'[x]+\[Lambda]*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{\lambda^2 x^5}{120} - \frac{\lambda x^5}{12} + \frac{3x^5}{40} - \frac{\lambda x^3}{6} + \frac{x^3}{6} + x \right) + c_1 \left( \frac{\lambda^2 x^4}{24} - \frac{\lambda x^4}{6} - \frac{\lambda x^2}{2} + 1 \right)$$

## 23.26 problem 33.10

Internal problem ID [13616]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.10.

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [Gegenbauer]

$$(-x^2 + 1)y'' - 2y'x + \lambda y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 63

```
Order:=6;
dsolve((1-x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+lambda*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{\lambda x^2}{2} + \frac{\lambda(\lambda - 6)x^4}{24}\right) y(0) + \left(x - \frac{(\lambda - 2)x^3}{6} + \frac{(\lambda - 2)(-12 + \lambda)x^5}{120}\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 80

```
AsymptoticDSolveValue[(1-x^2)*y''[x]-2*x*y'[x]+[Lambda]*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{\lambda^2 x^5}{120} - \frac{7\lambda x^5}{60} + \frac{x^5}{5} - \frac{\lambda x^3}{6} + \frac{x^3}{3} + x \right) + c_1 \left( \frac{\lambda^2 x^4}{24} - \frac{\lambda x^4}{4} - \frac{\lambda x^2}{2} + 1 \right)$$

## 23.27 problem 33.11 (a)

Internal problem ID [13617]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.11 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 4y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
Order:=5;  
dsolve(diff(y(x),x$2)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - 2x^2 + \frac{2}{3}x^4\right) y(0) + \left(-\frac{2}{3}x^3 + x\right) D(y)(0) + O(x^5)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 33

```
AsymptoticDSolveValue[y'[x]+4*y[x]==0,y[x],{x,0,4}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{2x^3}{3}\right) + c_1 \left(\frac{2x^4}{3} - 2x^2 + 1\right)$$

## 23.28 problem 33.11 (b)

Internal problem ID [13618]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.11 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y'' - yx^2 = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
Order:=5;  
dsolve(diff(y(x),x$2)-x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{x^4}{12}\right) y(0) + D(y)(0) x + O(x^5)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 20

```
AsymptoticDSolveValue[y'[x]-x^2*y[x]==0,y[x],{x,0,4}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^4}{12} + 1\right) + c_2 x$$

## 23.29 problem 33.11 (c)

Internal problem ID [13619]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.11 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + e^{2x}y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
Order:=4;  
dsolve(diff(y(x),x$2)+exp(2*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 - \frac{1}{3}x^3\right) y(0) + \left(x - \frac{1}{6}x^3\right) D(y)(0) + O(x^4)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 35

```
AsymptoticDSolveValue[y''[x]+Exp[2*x]*y[x]==0,y[x],{x,0,3}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{x^3}{6}\right) + c_1 \left(-\frac{x^3}{3} - \frac{x^2}{2} + 1\right)$$



### 23.30 problem 33.11 (d)

Internal problem ID [13620]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.11 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$\sin(x)y'' - y = 0$$

With the expansion point for the power series method at  $x = \frac{\pi}{2}$ .

#### ✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
Order:=4;  
dsolve(sin(x)*diff(y(x),x$2)-y(x)=0,y(x),type='series',x=Pi/2);
```

$$y(x) = \left(1 + \frac{(x - \frac{\pi}{2})^2}{2}\right) y\left(\frac{\pi}{2}\right) + \left(x - \frac{\pi}{2} + \frac{(x - \frac{\pi}{2})^3}{6}\right) D(y)\left(\frac{\pi}{2}\right) + O(x^4)$$

#### ✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 45

```
AsymptoticDSolveValue[Sin[x]*y''[x]-y[x]==0,y[x],{x,Pi/2,3}]
```

$$y(x) \rightarrow c_1 \left( \frac{1}{2} \left( x - \frac{\pi}{2} \right)^2 + 1 \right) + c_2 \left( \frac{1}{6} \left( x - \frac{\pi}{2} \right)^3 + x - \frac{\pi}{2} \right)$$

### 23.31 problem 33.11 (e)

Internal problem ID [13621]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.11 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + yx = \sin(x)$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
Order:=5;  
dsolve(diff(y(x),x$2)+x*y(x)=sin(x),y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{x^3}{6}\right) y(0) + \left(x - \frac{1}{12}x^4\right) D(y)(0) + \frac{x^3}{6} + O(x^5)$$

✓ Solution by Mathematica

Time used: 0.045 (sec). Leaf size: 35

```
AsymptoticDSolveValue[y''[x]+x*y[x]==Sin[x],y[x],{x,0,4}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{x^4}{12}\right) + \frac{x^3}{6} + c_1 \left(1 - \frac{x^3}{6}\right)$$

## 23.32 problem 33.11 (f)

Internal problem ID [13622]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.11 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - \sin(x)y' - yx = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
Order:=5;  
dsolve(diff(y(x),x$2)-sin(x)*diff(y(x),x)-x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{x^3}{6}\right) y(0) + \left(x + \frac{1}{6}x^3 + \frac{1}{12}x^4\right) D(y)(0) + O(x^5)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 35

```
AsymptoticDSolveValue[y''[x]-Sin[x]*y'[x]-x*y[x]==0,y[x],{x,0,4}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^3}{6} + 1\right) + c_2 \left(\frac{x^4}{12} + \frac{x^3}{6} + x\right)$$

### 23.33 problem 33.11 (g)

Internal problem ID [13623]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.11 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]`

$$y'' - y^2 = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 42

```
Order:=5;  
dsolve(diff(y(x),x$2)-y(x)^2=0,y(x),type='series',x=0);
```

$$y(x) = \frac{x^4 y(0)^3}{12} + \frac{y(0)^2 x^2}{2} + \left(1 + \frac{D(y)(0) x^3}{3}\right) y(0) + D(y)(0) x + \frac{D(y)(0)^2 x^4}{12} + O(x^5)$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 48

```
AsymptoticDSolveValue[y''[x]-y[x]^2==0,y[x],{x,0,4}]
```

$$y(x) \rightarrow \frac{1}{12}(c_1^3 + c_2^2) x^4 + \frac{1}{3} c_1 c_2 x^3 + \frac{c_1^2 x^2}{2} + c_2 x + c_1$$

## 23.34 problem 33.11 (h)

Internal problem ID [13624]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 33. Power series solutions I: Basic computational methods. Additional Exercises. page 641

**Problem number:** 33.11 (h).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_quadrature]

$$y' + \cos(y) = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 67

```
Order:=5;  
dsolve(diff(y(x),x)+cos(y(x))=0,y(x),type='series',x=0);
```

$$y(x) = y(0) - \cos(y(0))x - \frac{\sin(y(0))\cos(y(0))x^2}{2} + \left( \frac{\cos(y(0))^3}{3} - \frac{\cos(y(0))}{6} \right) x^3 \\ + \frac{\sin(y(0))\left(\cos(y(0))^3 - \frac{\cos(y(0))}{6}\right)x^4}{4} + O(x^5)$$

✓ Solution by Mathematica

Time used: 0.074 (sec). Leaf size: 76

```
AsymptoticDSolveValue[y'[x]+Cos[y[x]]==0,y[x],{x,0,4}]
```

$$y(x) \rightarrow \frac{1}{24}x^4(5\sin(c_1)\cos^3(c_1) - \sin^3(c_1)\cos(c_1)) \\ + \frac{1}{6}x^3(\cos^3(c_1) - \sin^2(c_1)\cos(c_1)) - \frac{1}{2}x^2\sin(c_1)\cos(c_1) - x\cos(c_1) + c_1$$

**24 Chapter 34. Power series solutions II:  
Generalization and theory. Additional  
Exercises. page 678**

24.1 problem 34.5 (a)	773
24.2 problem 34.5 (b)	774
24.3 problem 34.5 (c)	775
24.4 problem 34.5 (d)	778
24.5 problem 34.5 (e)	780
24.6 problem 34.5 (f)	783
24.7 problem 34.5 (g)	784
24.8 problem 34.5 (h)	786
24.9 problem 34.5 (i)	787
24.10 problem 34.5 (j)	788
24.11 problem 34.6 (a)	789
24.12 problem 34.6 (b)	790
24.13 problem 34.6 (c)	791
24.14 problem 34.6 (d)	792
24.15 problem 34.7 (a)	793
24.16 problem 34.7 (b)	795
24.17 problem 34.7 (c)	796
24.18 problem 34.7 (d)	797
24.19 problem 34.7 (e)	798
24.20 problem 34.7 (f)	799
24.21 problem 34.8 b(i)	800
24.22 problem 34.8 b(ii)	801
24.23 problem 34.8 b(iii)	802
24.24 problem 34.8 b(iv)	803
24.25 problem 34.9 b(i)	804
24.26 problem 34.9 b(ii)	805
24.27 problem 34.9 b(iii)	806
24.28 problem 34.9 b(iv)	807

## 24.1 problem 34.5 (a)

Internal problem ID [13625]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.5 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_separable`]

$$y' - e^x y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
Order:=6;  
dsolve(diff(y(x),x)-exp(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + x + x^2 + \frac{5}{6}x^3 + \frac{5}{8}x^4 + \frac{13}{30}x^5\right) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 33

```
AsymptoticDSolveValue[y'[x]-Exp[x]*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{13x^5}{30} + \frac{5x^4}{8} + \frac{5x^3}{6} + x^2 + x + 1 \right)$$

## 24.2 problem 34.5 (b)

Internal problem ID [13626]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.5 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_separable`]

$$y' - \tan(x)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
Order:=6;  
dsolve(diff(y(x),x)-tan(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 + \frac{5}{24}x^4\right) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 22

```
AsymptoticDSolveValue[y'[x]-Tan[x]*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{5x^4}{24} + \frac{x^2}{2} + 1 \right)$$



## 24.3 problem 34.5 (c)

Internal problem ID [13627]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.5 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$\sin(x)y'' + y'x^2 - e^xy = 0$$

With the expansion point for the power series method at  $x = 2$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 509

```
Order:=6;
dsolve(sin(x)*diff(y(x),x$2)+x^2*diff(y(x),x)-exp(x)*y(x)=0,y(x),type='series',x=2);
```

$$\begin{aligned}
 y(x) = & \left( 1 + \frac{e^2 \csc(2) (x-2)^2}{2} - \frac{e^2 (\cos(2) - \sin(2) + 4) \csc(2)^2 (x-2)^3}{6} \right. \\
 & - \frac{\left( (\cos(2) + 6) e^2 - \frac{e^4}{2} \right) \sin(2) + (-6 \cos(2) - 9) e^2}{12} \csc(2)^3 (x-2)^4 \\
 & + \frac{(2 e^4 \sin(2)^2 + (\cos(2))^2 e^2 + (28 e^2 - 2 e^4) \cos(2) + 50 e^2 - 4 e^4) \sin(2) + \cos(2)^3 e^2 - 3 \cos(2)^2 e^2}{60} \\
 & + \left( x - 2 - 2 \csc(2) (x-2)^2 - \frac{\csc(2)^2 (-e^2 \sin(2) - 4 \cos(2) + 4 \sin(2) - 16) (x-2)^3}{6} \right. \\
 & - \frac{(-\sin(2))^2 e^2 + ((\cos(2) + 4) e^2 - 4 \cos(2) - 24) \sin(2) + \cos(2)^2 + 24 \cos(2) + 35}{12} \csc(2)^3 (x-2)^4 \\
 & + \frac{\left( (-3 e^2 \cos(2) - 18 e^2 + \frac{e^4}{2}) \sin(2)^2 + (-6 \cos(2))^2 + (18 e^2 - 112) \cos(2) + 27 e^2 - 198 \right) \sin(2)}{60} \\
 & \left. + O(x^6) \right)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 953

AsymptoticDSolveValue[Sin[x]\*y''[x]+x^2\*y'[x]-Exp[x]\*y[x]==0,y[x],{x,2,5}]

$$\begin{aligned}
 y(x) \rightarrow & c_2 \left( -\frac{1}{60} (6 \csc(2) - 13 \cot(2) \csc(2) + 12 \cot^2(2) \csc(2) - 12 \cot^3(2) \csc(2)) (x-2)^5 \right. \\
 & - \frac{1}{20} (-e^2 \csc(2) + e^2 \cot(2) \csc(2) - e^2 \cot^2(2) \csc(2)) (x-2)^5 \\
 & + \frac{4}{15} \csc(2) (3 \csc(2) - 4 \cot(2) \csc(2) + 4 \cot^2(2) \csc(2)) (x-2)^5 \\
 & + \frac{1}{6} \csc(2) (-e^2 \csc(2) + e^2 \cot(2) \csc(2)) (x-2)^5 \\
 & - \frac{1}{40} (4 \csc(2) - 4 \cot(2) \csc(2)) (-4 \csc(2) + 4 \cot(2) \csc(2)) (x-2)^5 \\
 & + \frac{2}{5} \csc^2(2) (-4 \csc(2) + 4 \cot(2) \csc(2)) (x-2)^5 \\
 & + \frac{1}{40} e^2 \csc(2) (-4 \csc(2) + 4 \cot(2) \csc(2)) (x-2)^5 \\
 & - \frac{2}{5} \csc^2(2) (4 \csc(2) - 4 \cot(2) \csc(2)) (x-2)^5 \\
 & - \frac{1}{120} e^2 \csc(2) (4 \csc(2) - 4 \cot(2) \csc(2)) (x-2)^5 + \frac{32}{15} \csc^4(2) (x-2)^5 \\
 & + \frac{2}{5} e^2 \csc^3(2) (x-2)^5 + \frac{1}{120} e^4 \csc^2(2) (x-2)^5 \\
 & - \frac{1}{12} (3 \csc(2) - 4 \cot(2) \csc(2) + 4 \cot^2(2) \csc(2)) (x-2)^4 \\
 & - \frac{1}{12} (-e^2 \csc(2) + e^2 \cot(2) \csc(2)) (x-2)^4 \\
 & + \frac{1}{2} \csc(2) (4 \csc(2) - 4 \cot(2) \csc(2)) (x-2)^4 - \frac{8}{3} \csc^3(2) (x-2)^4 \\
 & - \frac{1}{3} e^2 \csc^2(2) (x-2)^4 - \frac{1}{6} (4 \csc(2) - 4 \cot(2) \csc(2)) (x-2)^3 + \frac{8}{3} \csc^2(2) (x-2)^3 \\
 & + \frac{1}{6} e^2 \csc(2) (x-2)^3 - 2 \csc(2) (x-2)^2 + x-2 \Big) + c_1 \left( -\frac{1}{60} (-2e^2 \csc(2) \right. \\
 & + 4e^2 \cot(2) \csc(2) - 3e^2 \cot^2(2) \csc(2) + 3e^2 \cot^3(2) \csc(2)) (x-2)^5 \\
 & + \frac{1}{15} \csc(2) (-e^2 \csc(2) + e^2 \cot(2) \csc(2) - e^2 \cot^2(2) \csc(2)) (x-2)^5 \\
 & - \frac{1}{20} e^2 \csc(2) (3 \csc(2) - 4 \cot(2) \csc(2) + 4 \cot^2(2) \csc(2)) (x-2)^5 \\
 & - \frac{1}{40} (-4 \csc(2) + 4 \cot(2) \csc(2)) (-e^2 \csc(2) + e^2 \cot(2) \csc(2)) (x-2)^5 \\
 & - \frac{2}{15} \csc^2(2) (-e^2 \csc(2) + e^2 \cot(2) \csc(2)) (x-2)^5 \\
 & - \frac{1}{30} e^2 \csc(2) (-e^2 \csc(2) + e^2 \cot(2) \csc(2)) (x-2)^5 \\
 & - \frac{1}{10} e^2 \csc^2(2) (-4 \csc(2) + 4 \cot(2) \csc(2)) (x-2)^5
 \end{aligned}$$

## 24.4 problem 34.5 (d)

Internal problem ID [13628]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.5 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$\sinh(x)y'' + y'x^2 - e^xy = 0$$

With the expansion point for the power series method at  $x = 2$ .

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 278

**Order:=6;**

`dsolve(sinh(x)*diff(y(x),x$2)+x^2*diff(y(x),x)-exp(x)*y(x)=0,y(x),type='series',x=2);`

$$\begin{aligned}
 y(x) = & \left( 1 + \frac{e^4(x-2)^2}{e^4-1} - \frac{2e^2(e^2+4e^4)(x-2)^3}{3(e^4-1)^2} + \frac{(e^2+12e^4+\frac{33e^6}{2}+\frac{e^{10}}{2})e^2(x-2)^4}{3(e^4-1)^3} \right. \\
 & \left. - \frac{2(e^2+\frac{53e^4}{2}+98e^6+79e^8+3e^{10}+\frac{5e^{12}}{2})e^2(x-2)^5}{15(e^4-1)^4} \right) y(2) \\
 & + \left( x-2 - \frac{4e^2(x-2)^2}{e^4-1} - \frac{2e^2(-\frac{31e^2}{2}-\frac{e^6}{2}-4)(x-2)^3}{3(e^4-1)^2} \right. \\
 & \left. + \frac{(-47e^2-65e^4-e^6-\frac{7e^8}{2}-\frac{7}{2})e^2(x-2)^4}{3(e^4-1)^3} \right. \\
 & \left. - \frac{2(-\frac{205e^2}{2}-\frac{1537e^4}{4}-\frac{1249e^6}{4}-\frac{85e^8}{4}-17e^{10}+\frac{e^{12}}{4}-\frac{e^{14}}{4}-\frac{11}{4})e^2(x-2)^5}{15(e^4-1)^4} \right) D(y)(2) \\
 & + O(x^6)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 931

AsymptoticDSolveValue[Sinh[x]\*y'[x]+x^2\*y'[x]-Exp[x]\*y[x]==0,y[x],{x,2,5}]

$$\begin{aligned}
 y(x) \rightarrow & c_2 \left( -\frac{1}{60} (-6\text{csch}(2) + 7\text{coth}(2)\text{csch}(2) + 12\text{coth}^2(2)\text{csch}(2) \right. \\
 & \qquad \qquad \qquad \left. - 12\text{coth}^3(2)\text{csch}(2)) (x-2)^5 \right. \\
 & \qquad \qquad \qquad - \frac{1}{20} (e^2\text{coth}(2)\text{csch}(2) - e^2\text{coth}^2(2)\text{csch}(2)) (x-2)^5 \\
 & \qquad \qquad \qquad + \frac{4}{15}\text{csch}(2) (-\text{csch}(2) - 4\text{coth}(2)\text{csch}(2) + 4\text{coth}^2(2)\text{csch}(2)) (x-2)^5 \\
 & \qquad \qquad \qquad + \frac{1}{6}\text{csch}(2) (-e^2\text{csch}(2) + e^2\text{coth}(2)\text{csch}(2)) (x-2)^5 \\
 & \qquad \qquad \qquad - \frac{1}{40} (4\text{csch}(2) - 4\text{coth}(2)\text{csch}(2)) (-4\text{csch}(2) + 4\text{coth}(2)\text{csch}(2)) (x-2)^5 \\
 & \qquad \qquad \qquad + \frac{2}{5}\text{csch}^2(2) (-4\text{csch}(2) + 4\text{coth}(2)\text{csch}(2)) (x-2)^5 \\
 & \qquad \qquad \qquad + \frac{1}{40} e^2\text{csch}(2) (-4\text{csch}(2) + 4\text{coth}(2)\text{csch}(2)) (x-2)^5 \\
 & \qquad \qquad \qquad - \frac{2}{5}\text{csch}^2(2) (4\text{csch}(2) - 4\text{coth}(2)\text{csch}(2)) (x-2)^5 \\
 & \qquad \qquad \qquad - \frac{1}{120} e^2\text{csch}(2) (4\text{csch}(2) - 4\text{coth}(2)\text{csch}(2)) (x-2)^5 + \frac{32}{15}\text{csch}^4(2) (x-2)^5 \\
 & \qquad \qquad \qquad + \frac{2}{5} e^2\text{csch}^3(2) (x-2)^5 + \frac{1}{120} e^4\text{csch}^2(2) (x-2)^5 \\
 & \qquad \qquad \qquad - \frac{1}{12} (-\text{csch}(2) - 4\text{coth}(2)\text{csch}(2) + 4\text{coth}^2(2)\text{csch}(2)) (x-2)^4 \\
 & \qquad \qquad \qquad - \frac{1}{12} (-e^2\text{csch}(2) + e^2\text{coth}(2)\text{csch}(2)) (x-2)^4 \\
 & \qquad \qquad \qquad + \frac{1}{2}\text{csch}(2) (4\text{csch}(2) - 4\text{coth}(2)\text{csch}(2)) (x-2)^4 - \frac{8}{3}\text{csch}^3(2) (x-2)^4 \\
 & \qquad \qquad \qquad - \frac{1}{3} e^2\text{csch}^2(2) (x-2)^4 - \frac{1}{6} (4\text{csch}(2) - 4\text{coth}(2)\text{csch}(2)) (x-2)^3 \\
 & \qquad \qquad \qquad + \frac{8}{3}\text{csch}^2(2) (x-2)^3 + \frac{1}{6} e^2\text{csch}(2) (x-2)^3 - 2\text{csch}(2) (x-2)^2 + x - 2 \Big) \\
 & + c_1 \left( -\frac{1}{60} (e^2\text{csch}(2) - e^2\text{coth}(2)\text{csch}(2) - 3e^2\text{coth}^2(2)\text{csch}(2) \right. \\
 & \qquad \qquad \qquad \left. + 3e^2\text{coth}^3(2)\text{csch}(2)) (x-2)^5 \right. \\
 & \qquad \qquad \qquad + \frac{1}{15}\text{csch}(2) (e^2\text{coth}(2)\text{csch}(2) - e^2\text{coth}^2(2)\text{csch}(2)) (x-2)^5 \\
 & \qquad \qquad \qquad - \frac{1}{20} e^2\text{csch}(2) (-\text{csch}(2) - 4\text{coth}(2)\text{csch}(2) + 4\text{coth}^2(2)\text{csch}(2)) (x-2)^5 \\
 & \qquad \qquad \qquad - \frac{1}{40} (-4\text{csch}(2) + 4\text{coth}(2)\text{csch}(2)) (-e^2\text{csch}(2) + e^2\text{coth}(2)\text{csch}(2)) (x-2)^5 \\
 & \qquad \qquad \qquad - \frac{2}{15}\text{csch}^2(2) (-e^2\text{csch}(2) + e^2\text{coth}(2)\text{csch}(2)) (x-2)^5
 \end{aligned}$$

## 24.5 problem 34.5 (e)

Internal problem ID [13629]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.5 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$\sinh(x)y'' + y'x^2 - \sin(x)y = 0$$

With the expansion point for the power series method at  $x = 2$ .

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 710

Order:=6;

dsolve(sinh(x)\*diff(y(x),x\$2)+x^2\*diff(y(x),x)-sin(x)\*y(x)=0,y(x),type='series',x=2);

$$\begin{aligned}
 y(x) = & \left( 1 + \frac{\sin(2) e^2 (x-2)^2}{e^4 - 1} + \frac{((-8e^2 - e^4 - 1) \sin(2) + \cos(2) (e^4 - 1)) e^2 (x-2)^3}{3(e^4 - 1)^2} \right. \\
 & + \frac{2 \left( \frac{(-e^2 + e^6) \sin(2)^2}{4} + (5e^2 + 9e^4 + e^6) \sin(2) + \cos(2) \left( e^2 - e^6 - \frac{e^8}{4} + \frac{1}{4} \right) \right) e^2 (x-2)^4}{3(e^4 - 1)^3} \\
 & + \frac{2 \left( \left( (e^2 - 2e^6 + e^{10}) \sin(2) - 8e^2 - \frac{41e^4}{4} + 6e^6 + \frac{41e^8}{4} + 2e^{10} + \frac{e^{12}}{4} - \frac{1}{4} \right) \cos(2) + \sin(2) \left( (e^2 + 4e^4 \right. \right.}{15(e^4 - 1)^4} \\
 & \left. \left. + \left( x - 2 - \frac{4e^2(x-2)^2}{e^4 - 1} + \frac{((e^4 - 1) \sin(2) + 32e^2 + 8) e^2 (x-2)^3}{3(e^4 - 1)^2} \right. \right. \right. \\
 & \left. \left. + \frac{2 \left( -\frac{7}{4} + \left( 2e^2 - 2e^6 - \frac{e^8}{4} + \frac{1}{4} \right) \sin(2) + \frac{\left( \frac{1}{2} - e^4 + \frac{e^8}{2} \right) \cos(2)}{2} - 24e^2 - \frac{69e^4}{2} + \frac{e^8}{4} \right) e^2 (x-2)^4}{3(e^4 - 1)^3} \right. \right. \\
 & \left. \left. + \frac{2 \left( \left( -\frac{e^{10}}{4} + \frac{e^6}{2} - \frac{e^2}{4} \right) \cos(2) \right)^2 + \left( -\frac{3}{4} - 5e^2 - \frac{3e^{12}}{4} - 5e^{10} + 10e^6 + \frac{3e^4}{4} + \frac{3e^8}{4} \right) \cos(2) + (5e^{10} - 13e^2}{15(e^4 - 1)} \right. \right. \\
 & \left. \left. + O(x^6) \right)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 934

AsymptoticDSolveValue[Sinh[x]\*y'[x]+x^2\*y'[x]-Sin[x]\*y[x]==0,y[x],{x,2,5}]

$y(x)$

$$\begin{aligned}
 \rightarrow c_2 & \left( -\frac{1}{20} (\cos(2) \coth(2) \operatorname{csch}(2) + \operatorname{csch}(2) \sin(2) - \coth^2(2) \operatorname{csch}(2) \sin(2)) (x-2)^5 \right. \\
 & + \frac{1}{6} \operatorname{csch}(2) (-\cos(2) \operatorname{csch}(2) + \coth(2) \operatorname{csch}(2) \sin(2)) (x-2)^5 \\
 & + \frac{1}{120} \operatorname{csch}^2(2) \sin^2(2) (x-2)^5 \\
 & + \frac{1}{40} \operatorname{csch}(2) (-4 \operatorname{csch}(2) + 4 \coth(2) \operatorname{csch}(2)) \sin(2) (x-2)^5 \\
 & - \frac{1}{120} \operatorname{csch}(2) (4 \operatorname{csch}(2) - 4 \coth(2) \operatorname{csch}(2)) \sin(2) (x-2)^5 + \frac{2}{5} \operatorname{csch}^3(2) \sin(2) (x-2)^5 \\
 & - \frac{1}{60} (-6 \operatorname{csch}(2) + 7 \coth(2) \operatorname{csch}(2) + 12 \coth^2(2) \operatorname{csch}(2) - 12 \coth^3(2) \operatorname{csch}(2)) (x-2)^5 \\
 & + \frac{1}{15} \operatorname{csch}(2) (-\operatorname{csch}(2) - 4 \coth(2) \operatorname{csch}(2) + 4 \coth^2(2) \operatorname{csch}(2)) (x-2)^5 \\
 & - \frac{1}{5} \operatorname{csch}(2) (\operatorname{csch}(2) + 4 \coth(2) \operatorname{csch}(2) - 4 \coth^2(2) \operatorname{csch}(2)) (x-2)^5 \\
 & - \frac{1}{40} (4 \operatorname{csch}(2) - 4 \coth(2) \operatorname{csch}(2)) (-4 \operatorname{csch}(2) + 4 \coth(2) \operatorname{csch}(2)) (x-2)^5 \\
 & + \frac{2}{3} \operatorname{csch}^2(2) (-4 \operatorname{csch}(2) + 4 \coth(2) \operatorname{csch}(2)) (x-2)^5 \\
 & - \frac{2}{15} \operatorname{csch}^2(2) (4 \operatorname{csch}(2) - 4 \coth(2) \operatorname{csch}(2)) (x-2)^5 + \frac{32}{15} \operatorname{csch}^4(2) (x-2)^5 \\
 & - \frac{1}{12} (-\cos(2) \operatorname{csch}(2) + \coth(2) \operatorname{csch}(2) \sin(2)) (x-2)^4 - \frac{1}{3} \operatorname{csch}^2(2) \sin(2) (x-2)^4 \\
 & - \frac{1}{12} (-\operatorname{csch}(2) - 4 \coth(2) \operatorname{csch}(2) + 4 \coth^2(2) \operatorname{csch}(2)) (x-2)^4 \\
 & - \frac{1}{3} \operatorname{csch}(2) (-4 \operatorname{csch}(2) + 4 \coth(2) \operatorname{csch}(2)) (x-2)^4 \\
 & + \frac{1}{6} \operatorname{csch}(2) (4 \operatorname{csch}(2) - 4 \coth(2) \operatorname{csch}(2)) (x-2)^4 - \frac{8}{3} \operatorname{csch}^3(2) (x-2)^4 \\
 & + \frac{1}{6} \operatorname{csch}(2) \sin(2) (x-2)^3 - \frac{1}{6} (4 \operatorname{csch}(2) - 4 \coth(2) \operatorname{csch}(2)) (x-2)^3 \\
 & + \frac{8}{3} \operatorname{csch}^2(2) (x-2)^3 - 2 \operatorname{csch}(2) (x-2)^2 + x-2 \Big) \\
 & + c_1 \left( -\frac{1}{60} (2 \cos(2) \operatorname{csch}(2) - 3 \cos(2) \coth^2(2) \operatorname{csch}(2) - 4 \coth(2) \operatorname{csch}(2) \sin(2)) \right. \\
 & \qquad \qquad \qquad \left. + 3 \coth^3(2) \operatorname{csch}(2) \sin(2)) (x-2)^5 \right. \\
 & + \frac{1}{15} \operatorname{csch}(2) (\cos(2) \coth(2) \operatorname{csch}(2) + \operatorname{csch}(2) \sin(2) - \coth^2(2) \operatorname{csch}(2) \sin(2)) (x-2)^5 \\
 & \qquad \qquad \qquad \left. - \frac{1}{30} \operatorname{csch}(2) \sin(2) (-\cos(2) \operatorname{csch}(2) + \coth(2) \operatorname{csch}(2) \sin(2)) (x-2)^5 \right)
 \end{aligned}$$



## 24.6 problem 34.5 (f)

Internal problem ID [13630]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.5 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$e^{3x}y'' + \sin(x)y' + \frac{2y}{x^2 + 4} = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

```
Order:=6;
```

```
dsolve(exp(3*x)*diff(y(x),x$2)+sin(x)*diff(y(x),x)+2/(x^2+4)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{4}x^2 + \frac{1}{4}x^3 - \frac{1}{8}x^4 - \frac{7}{160}x^5\right) y(0) + \left(x - \frac{1}{4}x^3 + \frac{3}{8}x^4 - \frac{67}{240}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 63

```
AsymptoticDSolveValue[Exp[3*x]*y''[x]+Sin[x]*y'[x]+2/(x^2+4)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( -\frac{67x^5}{240} + \frac{3x^4}{8} - \frac{x^3}{4} + x \right) + c_1 \left( -\frac{7x^5}{160} - \frac{x^4}{8} + \frac{x^3}{4} - \frac{x^2}{4} + 1 \right)$$

## 24.7 problem 34.5 (g)

Internal problem ID [13631]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.5 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + \frac{(e^x + 1)y}{1 - e^x} = 0$$

With the expansion point for the power series method at  $x = 3$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 167

```
Order:=6;  
dsolve(diff(y(x),x$2)+(1+exp(x))/(1-exp(x))*y(x)=0,y(x),type='series',x=3);
```

$$y(x) = \left( 1 + \frac{(1 + e^3)(x - 3)^2}{-2 + 2e^3} - \frac{e^3(x - 3)^3}{3(-1 + e^3)^2} + \frac{(e^9 + 3e^6 + e^3 - 1)(x - 3)^4}{24(-1 + e^3)^3} \right. \\ \left. + \frac{(-10e^9 - 8e^6 + 6e^3)(x - 3)^5}{120(-1 + e^3)^4} \right) y(3) + \left( x - 3 + \frac{(e^6 - 1)(x - 3)^3}{6(-1 + e^3)^2} \right. \\ \left. + \frac{(-e^6 + e^3)(x - 3)^4}{6(-1 + e^3)^3} + \frac{(e^{12} + 6e^9 - 2e^6 - 6e^3 + 1)(x - 3)^5}{120(-1 + e^3)^4} \right) D(y)(3) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 275

AsymptoticDSolveValue[y''[x]+(1+Exp[x])/(1-Exp[x])\*y[x]==0,y[x],{x,3,5}]

$$\begin{aligned}
 y(x) \rightarrow c_1 & \left( -\frac{(e^3 + 4e^6 + e^9)(x-3)^5}{60(e^3-1)^4} - \frac{e^3(1+e^3)(x-3)^5}{60(e^3-1)^3} + \frac{e^3(-1-e^3)(x-3)^5}{20(e^3-1)^3} \right. \\
 & - \frac{(-e^3-e^6)(x-3)^4}{12(e^3-1)^3} + \frac{(-1-e^3)^2(x-3)^4}{24(e^3-1)^2} - \frac{e^3(x-3)^3}{3(e^3-1)^2} + \frac{(1+e^3)(x-3)^2}{2(e^3-1)} \\
 & \left. + 1 \right) + c_2 \left( -\frac{(-e^3-e^6)(x-3)^5}{20(e^3-1)^3} + \frac{(1+e^3)^2(x-3)^5}{120(e^3-1)^2} - \frac{e^3(x-3)^4}{6(e^3-1)^2} \right. \\
 & \left. + \frac{(1+e^3)(x-3)^3}{6(e^3-1)} + x-3 \right)
 \end{aligned}$$

## 24.8 problem 34.5 (h)

Internal problem ID [13632]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.5 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 - 4)y'' + (x^2 + x - 6)y = 0$$

With the expansion point for the power series method at  $x = 2$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

```
Order:=6;  
dsolve((x^2-4)*diff(y(x),x$2)+(x^2+x-6)*y(x)=0,y(x),type='series',x=2);
```

$$y(x) = \left(1 - \frac{5(x-2)^2}{8} + \frac{(x-2)^3}{96} + \frac{49(x-2)^4}{768} - \frac{37(x-2)^5}{15360}\right) y(2) \\ + \left(x - 2 - \frac{5(x-2)^3}{24} + \frac{(x-2)^4}{192} + \frac{47(x-2)^5}{3840}\right) D(y)(2) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 78

```
AsymptoticDSolveValue[(x^2-4)*y'[x]+(x^2+x-6)*y[x]==0,y[x],{x,2,5}]
```

$$y(x) \rightarrow c_1 \left( -\frac{37(x-2)^5}{15360} + \frac{49}{768}(x-2)^4 + \frac{1}{96}(x-2)^3 - \frac{5}{8}(x-2)^2 + 1 \right) \\ + c_2 \left( \frac{47(x-2)^5}{3840} + \frac{1}{192}(x-2)^4 - \frac{5}{24}(x-2)^3 + x - 2 \right)$$

## 24.9 problem 34.5 (i)

Internal problem ID [13633]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.5 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$xy'' + (1 - e^x)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 49

```
Order:=6;  
dsolve(x*diff(y(x),x$2)+(1-exp(x))*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 + \frac{1}{12}x^3 + \frac{1}{18}x^4 + \frac{3}{160}x^5\right) y(0) \\ + \left(x + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{60}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 63

```
AsymptoticDSolveValue[x*y''[x]+(1-Exp[x])*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{x^5}{60} + \frac{x^4}{24} + \frac{x^3}{6} + x \right) + c_1 \left( \frac{3x^5}{160} + \frac{x^4}{18} + \frac{x^3}{12} + \frac{x^2}{2} + 1 \right)$$

## 24.10 problem 34.5 (j)

Internal problem ID [13634]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.5 (j).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$\sin(\pi x^2) y'' + y x^2 = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 46

```
Order:=6;  
dsolve(sin(Pi*x^2)*diff(y(x),x$2)+x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{x^2}{2\pi} + \frac{x^4}{24\pi^2}\right) y(0) + \left(x - \frac{x^3}{6\pi} + \frac{x^5}{120\pi^2}\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 54

```
AsymptoticDSolveValue[Sin[Pi*x^2]*y'[x]+x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{x^5}{120\pi^2} - \frac{x^3}{6\pi} + x \right) + c_1 \left( \frac{x^4}{24\pi^2} - \frac{x^2}{2\pi} + 1 \right)$$

## 24.11 problem 34.6 (a)

Internal problem ID [13635]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.6 (a).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - e^x y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
Order:=6;  
dsolve(diff(y(x),x)-exp(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + x + x^2 + \frac{5}{6}x^3 + \frac{5}{8}x^4 + \frac{13}{30}x^5\right) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 33

```
AsymptoticDSolveValue[y'[x]-Exp[x]*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{13x^5}{30} + \frac{5x^4}{8} + \frac{5x^3}{6} + x^2 + x + 1 \right)$$

## 24.12 problem 34.6 (b)

Internal problem ID [13636]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.6 (b).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' + e^{2x}y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
Order:=6;  
dsolve(diff(y(x),x)+exp(2*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x - \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{3}{8}x^4 + \frac{23}{120}x^5\right) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 39

```
AsymptoticDSolveValue[y'[x]+Exp[2*x]*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{23x^5}{120} + \frac{3x^4}{8} + \frac{x^3}{6} - \frac{x^2}{2} - x + 1 \right)$$



## 24.13 problem 34.6 (c)

Internal problem ID [13637]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.6 (c).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_separable`]

$$y' + \cos(x)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
Order:=6;  
dsolve(diff(y(x),x)+cos(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{8}x^4 + \frac{1}{15}x^5\right) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 32

```
AsymptoticDSolveValue[y'[x]+Cos[x]*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{x^5}{15} - \frac{x^4}{8} + \frac{x^2}{2} - x + 1 \right)$$

## 24.14 problem 34.6 (d)

Internal problem ID [13638]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.6 (d).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_separable`]

$$y' + \ln(x)y = 0$$

With the expansion point for the power series method at  $x = 1$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
Order:=6;  
dsolve(diff(y(x),x)+ln(x)*y(x)=0,y(x),type='series',x=1);
```

$$y(x) = \left( 1 - \frac{(-1+x)^2}{2} + \frac{(-1+x)^3}{6} + \frac{(-1+x)^4}{24} - \frac{(-1+x)^5}{30} \right) y(1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 44

```
AsymptoticDSolveValue[y'[x]+Log[x]*y[x]==0,y[x],{x,1,5}]
```

$$y(x) \rightarrow c_1 \left( -\frac{1}{30}(x-1)^5 + \frac{1}{24}(x-1)^4 + \frac{1}{6}(x-1)^3 - \frac{1}{2}(x-1)^2 + 1 \right)$$

## 24.15 problem 34.7 (a)

Internal problem ID [13639]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.7 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - e^x y = 0$$

With the expansion point for the power series method at  $x = 1$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 87

```
Order:=6;  
dsolve(diff(y(x),x$2)-exp(x)*y(x)=0,y(x),type='series',x=1);
```

$$y(x) = \left( 1 + \frac{e(-1+x)^2}{2} + \frac{e(-1+x)^3}{6} + \left( \frac{e^2}{24} + \frac{e}{24} \right) (-1+x)^4 + \left( \frac{e^2}{30} + \frac{e}{120} \right) (-1+x)^5 \right) y(1) + \left( -1+x + \frac{e(-1+x)^3}{6} + \frac{e(-1+x)^4}{12} + \frac{(e^2+3e)(-1+x)^5}{120} \right) D(y)(1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 121

```
AsymptoticDSolveValue[y''[x]-Exp[x]*y[x]==0,y[x],{x,1,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{1}{30} e^2 (x-1)^5 + \frac{1}{120} e (x-1)^5 + \frac{1}{24} e^2 (x-1)^4 + \frac{1}{24} e (x-1)^4 + \frac{1}{6} e (x-1)^3 + \frac{1}{2} e (x-1)^2 + 1 \right) + c_2 \left( \frac{1}{120} e^2 (x-1)^5 + \frac{1}{40} e (x-1)^5 + \frac{1}{12} e (x-1)^4 + \frac{1}{6} e (x-1)^3 + x - 1 \right)$$

## 24.16 problem 34.7 (b)

Internal problem ID [13640]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.7 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 3y'/x - e^x y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

```
Order:=6;  
dsolve(diff(y(x),x$2)+3*x*diff(y(x),x)-exp(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{6}x^4 - \frac{1}{30}x^5\right) y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{12}x^4 + \frac{19}{120}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 63

```
AsymptoticDSolveValue[y''[x]+3*x*y'[x]-Exp[x]*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{19x^5}{120} + \frac{x^4}{12} - \frac{x^3}{3} + x \right) + c_1 \left( -\frac{x^5}{30} - \frac{x^4}{6} + \frac{x^3}{6} + \frac{x^2}{2} + 1 \right)$$

## 24.17 problem 34.7 (c)

Internal problem ID [13641]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.7 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$xy'' - 3y'x + \sin(x)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=4;  
dsolve(x*diff(y(x),x$2)-3*x*diff(y(x),x)+sin(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 - \frac{1}{2}x^3\right) y(0) + \left(x + \frac{3}{2}x^2 + \frac{4}{3}x^3\right) D(y)(0) + O(x^4)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[x*y''[x]-3*x*y'[x]+Sin[x]*y[x]==0,y[x],{x,0,3}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^3}{2} - \frac{x^2}{2} + 1\right) + c_2 \left(\frac{4x^3}{3} + \frac{3x^2}{2} + x\right)$$

## 24.18 problem 34.7 (d)

Internal problem ID [13642]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.7 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [\_Titchmarsh]

$$y'' + \ln(x)y = 0$$

With the expansion point for the power series method at  $x = 1$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
Order:=6;  
dsolve(diff(y(x),x$2)+ln(x)*y(x)=0,y(x),type='series',x=1);
```

$$y(x) = \left(1 - \frac{(-1+x)^3}{6} + \frac{(-1+x)^4}{24} - \frac{(-1+x)^5}{60}\right) y(1) \\ + \left(-1+x - \frac{(-1+x)^4}{12} + \frac{(-1+x)^5}{40}\right) D(y)(1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 60

```
AsymptoticDSolveValue[y''[x]+Log[x]*y[x]==0,y[x],{x,1,5}]
```

$$y(x) \rightarrow c_1 \left( -\frac{1}{60}(x-1)^5 + \frac{1}{24}(x-1)^4 - \frac{1}{6}(x-1)^3 + 1 \right) \\ + c_2 \left( \frac{1}{40}(x-1)^5 - \frac{1}{12}(x-1)^4 + x - 1 \right)$$

## 24.19 problem 34.7 (e)

Internal problem ID [13643]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.7 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$\sqrt{x}y'' + y = 0$$

With the expansion point for the power series method at  $x = 1$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

```
Order:=6;  
dsolve(sqrt(x)*diff(y(x),x$2)+y(x)=0,y(x),type='series',x=1);
```

$$y(x) = \left(1 - \frac{(-1+x)^2}{2} + \frac{(-1+x)^3}{12} + \frac{(-1+x)^4}{96} - \frac{(-1+x)^5}{960}\right) y(1) \\ + \left(-1+x - \frac{(-1+x)^3}{6} + \frac{(-1+x)^4}{24} - \frac{(-1+x)^5}{96}\right) D(y)(1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 78

```
AsymptoticDSolveValue[Sqrt[x]*y''[x]+y[x]==0,y[x],{x,1,5}]
```

$$y(x) \rightarrow c_1 \left( -\frac{1}{960}(x-1)^5 + \frac{1}{96}(x-1)^4 + \frac{1}{12}(x-1)^3 - \frac{1}{2}(x-1)^2 + 1 \right) \\ + c_2 \left( -\frac{1}{96}(x-1)^5 + \frac{1}{24}(x-1)^4 - \frac{1}{6}(x-1)^3 + x - 1 \right)$$



## 24.20 problem 34.7 (f)

Internal problem ID [13644]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.7 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$y'' + (6x^2 + 2x + 1)y' + (2 + 12x)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
Order:=6;
dsolve(diff(y(x),x$2)+(1+2*x+6*x^2)*diff(y(x),x)+(2+12*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x^2 - \frac{5}{3}x^3 + \frac{11}{12}x^4 + \frac{101}{60}x^5\right)y(0) \\ + \left(x - \frac{1}{2}x^2 - \frac{1}{2}x^3 - \frac{9}{8}x^4 + \frac{41}{40}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 68

```
AsymptoticDSolveValue[y'[x]+(1+2*x+6*x^2)*y'[x]+(2+12*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{101x^5}{60} + \frac{11x^4}{12} - \frac{5x^3}{3} - x^2 + 1 \right) + c_2 \left( \frac{41x^5}{40} - \frac{9x^4}{8} - \frac{x^3}{2} - \frac{x^2}{2} + x \right)$$

## 24.21 problem 34.8 b(i)

Internal problem ID [13645]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.8 b(i).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' - e^x y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 52

```
Order:=10;  
dsolve(diff(y(x),x)-exp(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + x + x^2 + \frac{5}{6}x^3 + \frac{5}{8}x^4 + \frac{13}{30}x^5 + \frac{203}{720}x^6 + \frac{877}{5040}x^7 + \frac{23}{224}x^8 + \frac{1007}{17280}x^9\right) y(0) + O(x^{10})$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 61

```
AsymptoticDSolveValue[y'[x]-Exp[x]*y[x]==0,y[x],{x,0,9}]
```

$$y(x) \rightarrow c_1 \left( \frac{1007x^9}{17280} + \frac{23x^8}{224} + \frac{877x^7}{5040} + \frac{203x^6}{720} + \frac{13x^5}{30} + \frac{5x^4}{8} + \frac{5x^3}{6} + x^2 + x + 1 \right)$$

## 24.22 problem 34.8 b(ii)

Internal problem ID [13646]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.8 b(ii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' + \sqrt{x^2 + 1} y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 46

```
Order:=8;  
dsolve(diff(y(x),x)+sqrt(1+x^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{3}x^3 + \frac{5}{24}x^4 - \frac{1}{15}x^5 + \frac{13}{720}x^6 - \frac{11}{630}x^7\right) y(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 53

```
AsymptoticDSolveValue[y'[x]+Sqrt[1+x^2]*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left( -\frac{11x^7}{630} + \frac{13x^6}{720} - \frac{x^5}{15} + \frac{5x^4}{24} - \frac{x^3}{3} + \frac{x^2}{2} - x + 1 \right)$$

## 24.23 problem 34.8 b(iii)

Internal problem ID [13647]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.8 b(iii).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [`_separable`]

$$\cos(x) y' + y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 46

```
Order:=8;  
dsolve(cos(x)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{3}x^3 + \frac{5}{24}x^4 - \frac{2}{15}x^5 + \frac{61}{720}x^6 - \frac{17}{315}x^7\right) y(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 53

```
AsymptoticDSolveValue[Cos[x]*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left( -\frac{17x^7}{315} + \frac{61x^6}{720} - \frac{2x^5}{15} + \frac{5x^4}{24} - \frac{x^3}{3} + \frac{x^2}{2} - x + 1 \right)$$

## 24.24 problem 34.8 b(iv)

Internal problem ID [13648]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.8 b(iv).

**ODE order:** 1.

**ODE degree:** 1.

CAS Maple gives this as type [\_separable]

$$y' + \sqrt{2x^2 + 1}y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 46

```
Order:=8;  
dsolve(diff(y(x),x)+sqrt(1+2*x^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{2}x^3 + \frac{3}{8}x^4 - \frac{3}{40}x^5 + \frac{1}{80}x^6 - \frac{51}{560}x^7\right)y(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 53

```
AsymptoticDSolveValue[y'[x]+Sqrt[1+2*x^2]*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left( -\frac{51x^7}{560} + \frac{x^6}{80} - \frac{3x^5}{40} + \frac{3x^4}{8} - \frac{x^3}{2} + \frac{x^2}{2} - x + 1 \right)$$

## 24.25 problem 34.9 b(i)

Internal problem ID [13649]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.9 b(i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - e^x y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 69

```
Order:=8;  
dsolve(diff(y(x),x$2)-exp(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{1}{24}x^5 + \frac{13}{720}x^6 + \frac{1}{140}x^7\right) y(0) \\ + \left(x + \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{1}{30}x^5 + \frac{1}{72}x^6 + \frac{29}{5040}x^7\right) D(y)(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 91

```
AsymptoticDSolveValue[y''[x]-Exp[x]*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left( \frac{29x^7}{5040} + \frac{x^6}{72} + \frac{x^5}{30} + \frac{x^4}{12} + \frac{x^3}{6} + x \right) + c_1 \left( \frac{x^7}{140} + \frac{13x^6}{720} + \frac{x^5}{24} + \frac{x^4}{12} + \frac{x^3}{6} + \frac{x^2}{2} + 1 \right)$$

## 24.26 problem 34.9 b(ii)

Internal problem ID [13650]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.9 b(ii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + \cos(x)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 54

```
Order:=10;  
dsolve(diff(y(x),x$2)+cos(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{12}x^4 - \frac{1}{80}x^6 + \frac{11}{8064}x^8\right) y(0) \\ + \left(x - \frac{1}{6}x^3 + \frac{1}{30}x^5 - \frac{19}{5040}x^7 + \frac{29}{72576}x^9\right) D(y)(0) + O(x^{10})$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 70

```
AsymptoticDSolveValue[y'[x]+Cos[x]*y[x]==0,y[x],{x,0,9}]
```

$$y(x) \rightarrow c_2 \left( \frac{29x^9}{72576} - \frac{19x^7}{5040} + \frac{x^5}{30} - \frac{x^3}{6} + x \right) + c_1 \left( \frac{11x^8}{8064} - \frac{x^6}{80} + \frac{x^4}{12} - \frac{x^2}{2} + 1 \right)$$

## 24.27 problem 34.9 b(iii)

Internal problem ID [13651]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.9 b(iii).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$y'' + \sin(x)y' + \cos(x)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
Order:=7;  
dsolve(diff(y(x),x$2)+sin(x)*diff(y(x),x)+cos(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^4 - \frac{31}{720}x^6\right)y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{10}x^5\right)D(y)(0) + O(x^7)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 49

```
AsymptoticDSolveValue[y''[x]+Sin[x]*y'[x]+Cos[x]*y[x]==0,y[x],{x,0,6}]
```

$$y(x) \rightarrow c_2 \left( \frac{x^5}{10} - \frac{x^3}{3} + x \right) + c_1 \left( -\frac{31x^6}{720} + \frac{x^4}{6} - \frac{x^2}{2} + 1 \right)$$



## 24.28 problem 34.9 b(iv)

Internal problem ID [13652]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 34. Power series solutions II: Generalization and theory. Additional Exercises. page 678

**Problem number:** 34.9 b(iv).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$\sqrt{x} y'' + y' + yx = 0$$

With the expansion point for the power series method at  $x = 1$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```
Order:=5;
dsolve(sqrt(x)*diff(y(x),x$2)+diff(y(x),x)+x*y(x)=0,y(x),type='series',x=1);
```

$$y(x) = \left(1 - \frac{(-1+x)^2}{2} + \frac{(-1+x)^3}{12} - \frac{(-1+x)^4}{96}\right) y(1) \\ + \left(-1+x - \frac{(-1+x)^2}{2} + \frac{(-1+x)^3}{12} - \frac{3(-1+x)^4}{32}\right) D(y)(1) + O(x^5)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 69

```
AsymptoticDSolveValue[Sqrt[x]*y''[x]+y'[x]+x*y[x]==0,y[x],{x,1,4}]
```

$$y(x) \rightarrow c_1 \left( -\frac{1}{96}(x-1)^4 + \frac{1}{12}(x-1)^3 - \frac{1}{2}(x-1)^2 + 1 \right) \\ + c_2 \left( -\frac{3}{32}(x-1)^4 + \frac{1}{12}(x-1)^3 - \frac{1}{2}(x-1)^2 + x - 1 \right)$$

**25 Chapter 35. Modified Power series solutions  
and basic method of Frobenius. Additional  
Exercises. page 715**

25.1 problem 35.2 (a)	809
25.2 problem 35.2 (b)	810
25.3 problem 35.2 (c)	811
25.4 problem 35.2 (d)	812
25.5 problem 35.2 (e)	813
25.6 problem 35.2 (f)	814
25.7 problem 35.3 (a)	815
25.8 problem 35.3 (b)	817
25.9 problem 35.3 (c)	818
25.10 problem 35.3 (d)	820
25.11 problem 35.3 (e)	822
25.12 problem 35.3 (f)	824
25.13 problem 35.3 (g)	825
25.14 problem 35.3 (h)	826
25.15 problem 35.4 (a)	827
25.16 problem 35.4 (b)	828
25.17 problem 35.4 (c)	830
25.18 problem 35.4 (d)	831
25.19 problem 35.4 (e)	832
25.20 problem 35.4 (f)	833
25.21 problem 35.4 (g)	834
25.22 problem 35.4 (h)	835
25.23 problem 35.4 (i)	836
25.24 problem 35.4 (j)	838
25.25 problem 35.4 (k)	839
25.26 problem 35.4 (L)	840
25.27 problem 35.4 (m)	841
25.28 problem 35.4 (n)	842
25.29 problem 35.5 (a)	843
25.30 problem 35.5 (b)	845
25.31 problem 35.5 (c)	846
25.32 problem 35.5 (d)	848

## 25.1 problem 35.2 (a)

Internal problem ID [13653]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.2 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`, `[_2nd_order, _linear,`

$$(x - 3)^2 y'' - 2(x - 3) y' + 2y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve((x-3)^2*dif(y(x),x$2)-2*(x-3)*dif(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{(-x^2 + 9) y(0)}{9} - \frac{x D(y)(0) (x - 3)}{3}$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[(x-3)^2*y''[x]-2*(x-3)*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(1 - \frac{x^2}{9}\right) + c_2 \left(x - \frac{x^2}{3}\right)$$

## 25.2 problem 35.2 (b)

Internal problem ID [13654]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.2 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$2x^2y'' + 5y'x + y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
Order:=6;  
dsolve(2*x^2*diff(y(x),x$2)+5*x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1\sqrt{x} + c_2x}{x^{\frac{3}{2}}} + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 18

```
AsymptoticDSolveValue[2*x^2*y''[x]+5*x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_1}{\sqrt{x}} + \frac{c_2}{x}$$

## 25.3 problem 35.2 (c)

Internal problem ID [13655]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.2 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x-1)^2 y'' - 5(x-1)y' + 9y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
Order:=6;
dsolve((x-1)^2*diff(y(x),x$2)-5*(x-1)*diff(y(x),x)+9*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{9}{2}x^2 + \frac{9}{2}x^3 - \frac{3}{4}x^4 - \frac{3}{20}x^5\right) y(0) \\ + \left(x - \frac{5}{2}x^2 + \frac{11}{6}x^3 - \frac{1}{4}x^4 - \frac{1}{20}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 70

```
AsymptoticDSolveValue[(x-1)^2*y''[x]-5*(x-1)*y'[x]+9*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( -\frac{3x^5}{20} - \frac{3x^4}{4} + \frac{9x^3}{2} - \frac{9x^2}{2} + 1 \right) + c_2 \left( -\frac{x^5}{20} - \frac{x^4}{4} + \frac{11x^3}{6} - \frac{5x^2}{2} + x \right)$$

## 25.4 problem 35.2 (d)

Internal problem ID [13656]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.2 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$(x + 2)^2 y'' + (x + 2) y' = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 34

```
Order:=6;  
dsolve((x+2)^2*dif(y(x),x$2)+(x+2)*dif(y(x),x)=0,y(x),type='series',x=0);
```

$$y(x) = y(0) + \left( x - \frac{1}{4}x^2 + \frac{1}{12}x^3 - \frac{1}{32}x^4 + \frac{1}{80}x^5 \right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 39

```
AsymptoticDSolveValue[(x+2)^2*y''[x]+(x+2)*y'[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{x^5}{80} - \frac{x^4}{32} + \frac{x^3}{12} - \frac{x^2}{4} + x \right) + c_1$$

## 25.5 problem 35.2 (e)

Internal problem ID [13657]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.2 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$3(x-2)^2 y'' - 4(x-5)y' + 2y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 54

```
Order:=6;
```

```
dsolve(3*(x-2)^2*diff(y(x),x$2)-4*(x-5)*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{12}x^2 + \frac{1}{54}x^3 + \frac{1}{648}x^4 - \frac{1}{4860}x^5\right) y(0) \\ + \left(x - \frac{5}{6}x^2 + \frac{23}{108}x^3 + \frac{23}{1296}x^4 - \frac{23}{9720}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 70

```
AsymptoticDSolveValue[3*(x-2)^2*y'[x]-4*(x-5)*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( -\frac{x^5}{4860} + \frac{x^4}{648} + \frac{x^3}{54} - \frac{x^2}{12} + 1 \right) + c_2 \left( -\frac{23x^5}{9720} + \frac{23x^4}{1296} + \frac{23x^3}{108} - \frac{5x^2}{6} + x \right)$$

## 25.6 problem 35.2 (f)

Internal problem ID [13658]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.2 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`, `[_2nd_order, _linear,`

$$(x - 5)^2 y'' + (x - 5) y' + 4y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
Order:=6;
dsolve((x-5)^2*diff(y(x),x$2)+(x-5)*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{2}{25}x^2 - \frac{2}{125}x^3 - \frac{7}{3750}x^4 - \frac{1}{9375}x^5\right) y(0) \\ + \left(x + \frac{1}{10}x^2 - \frac{1}{75}x^3 - \frac{3}{500}x^4 - \frac{1}{750}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 70

```
AsymptoticDSolveValue[(x-5)^2*y''[x]+(x-5)*y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( -\frac{x^5}{9375} - \frac{7x^4}{3750} - \frac{2x^3}{125} - \frac{2x^2}{25} + 1 \right) + c_2 \left( -\frac{x^5}{750} - \frac{3x^4}{500} - \frac{x^3}{75} + \frac{x^2}{10} + x \right)$$



## 25.7 problem 35.3 (a)

Internal problem ID [13659]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.3 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + \frac{xy'}{x-2} + \frac{2y}{x+2} = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 1283

Order:=6;

dsolve(x^2\*diff(y(x),x\$2)+x/(x-2)\*diff(y(x),x)+2/(x+2)\*y(x)=0,y(x),type='series',x=0);

$$\begin{aligned}
 y(x) = & x^{\frac{3}{4}} \left( c_2 x^{\frac{i\sqrt{7}}{4}} \left( 1 + \frac{i\sqrt{7} + 11}{8i\sqrt{7} + 16} x + \frac{1}{64} \frac{7i\sqrt{7} + 45}{(2 + i\sqrt{7})(i\sqrt{7} + 4)} x^2 \right. \right. \\
 & + \frac{1}{256} \frac{223i\sqrt{7} - 43}{(2 + i\sqrt{7})(i\sqrt{7} + 4)(i\sqrt{7} + 6)} x^3 \\
 & + \frac{1}{4096} \frac{7577i\sqrt{7} + 979}{(2 + i\sqrt{7})(i\sqrt{7} + 4)(i\sqrt{7} + 6)(i\sqrt{7} + 8)} x^4 \\
 & \left. \left. + \frac{1}{81920} \frac{553875i\sqrt{7} - 1249007}{(2 + i\sqrt{7})(i\sqrt{7} + 4)(i\sqrt{7} + 6)(i\sqrt{7} + 8)(i\sqrt{7} + 10)} x^5 + O(x^6) \right) \right. \\
 & + c_1 x^{-\frac{i\sqrt{7}}{4}} \left( 1 + \frac{i\sqrt{7} - 11}{8i\sqrt{7} - 16} x - \frac{1}{64} \frac{7i\sqrt{7} - 45}{(i\sqrt{7} - 2)(i\sqrt{7} - 4)} x^2 \right. \\
 & + \frac{1}{256} \frac{223i\sqrt{7} + 43}{(i\sqrt{7} - 2)(i\sqrt{7} - 4)(i\sqrt{7} - 6)} x^3 \\
 & - \frac{1}{4096} \frac{7577i\sqrt{7} - 979}{(i\sqrt{7} - 2)(i\sqrt{7} - 4)(i\sqrt{7} - 6)(i\sqrt{7} - 8)} x^4 \\
 & \left. \left. + \frac{1}{81920} \frac{553875i\sqrt{7} + 1249007}{(i\sqrt{7} - 2)(i\sqrt{7} - 4)(i\sqrt{7} - 6)(i\sqrt{7} - 8)(i\sqrt{7} - 10)} x^5 + O(x^6) \right) \right)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 6290

AsymptoticDSolveValue[x^2\*y'[x]+x/(x-2)\*y'[x]+2/(x+2)\*y[x]==0,y[x],{x,0,5}]

Too large to display

## 25.8 problem 35.3 (b)

Internal problem ID [13660]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.3 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^3 y'' + y' x^2 + y = 0$$

With the expansion point for the power series method at  $x = 2$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
Order:=6;
dsolve(x^3*diff(y(x),x$2)+x^2*diff(y(x),x)+y(x)=0,y(x),type='series',x=2);
```

$$y(x) = \left(1 - \frac{(x-2)^2}{16} + \frac{(x-2)^3}{24} - \frac{35(x-2)^4}{1536} + \frac{89(x-2)^5}{7680}\right) y(2) \\ + \left(x-2 - \frac{(x-2)^2}{4} + \frac{(x-2)^3}{16} - \frac{(x-2)^4}{96} - \frac{19(x-2)^5}{7680}\right) D(y)(2) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 87

```
AsymptoticDSolveValue[x^3*y''[x]+x^2*y'[x]+y[x]==0,y[x],{x,2,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{89(x-2)^5}{7680} - \frac{35(x-2)^4}{1536} + \frac{1}{24}(x-2)^3 - \frac{1}{16}(x-2)^2 + 1 \right) \\ + c_2 \left( -\frac{19(x-2)^5}{7680} - \frac{1}{96}(x-2)^4 + \frac{1}{16}(x-2)^3 - \frac{1}{4}(x-2)^2 + x-2 \right)$$

## 25.9 problem 35.3 (c)

Internal problem ID [13661]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.3 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(-x^4 + x^3)y'' + (3x - 1)y' + 827y = 0$$

With the expansion point for the power series method at  $x = 1$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 62

Order:=6;

```
dsolve((x^3-x^4)*diff(y(x),x$2)+(3*x-1)*diff(y(x),x)+827*y(x)=0,y(x),type='series',x=1);
```

$$\begin{aligned} y(x) = & c_1(-1+x)^3 \left( 1 + \frac{409}{2}(-1+x) + \frac{328391}{20}(-1+x)^2 + \frac{128327201}{180}(-1+x)^3 \right. \\ & \left. + \frac{19341852779}{1008}(-1+x)^4 + \frac{6949904889503}{20160}(-1+x)^5 + O((-1+x)^6) \right) \\ & + c_2 \left( \ln(-1+x) \left( 567661070(-1+x)^3 + 116086688815(-1+x)^4 \right. \right. \\ & \left. \left. + \frac{18641478643837}{2}(-1+x)^5 + O((-1+x)^6) \right) \right. \\ & \left. + \left( 12 - 4962(-1+x) + 2059230(-1+x)^2 - 6162812(-1+x)^3 \right. \right. \\ & \left. \left. - \frac{592298912511}{4}(-1+x)^4 - \frac{744988601770307}{40}(-1+x)^5 + O((-1+x)^6) \right) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.067 (sec). Leaf size: 105

```
AsymptoticDSolveValue[(x^3-x^4)*y'[x]+(3*x-1)*y'[x]+827*y[x]==0,y[x],{x,1,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{19341852779(x-1)^7}{1008} + \frac{128327201}{180}(x-1)^6 + \frac{328391}{20}(x-1)^5 + \frac{409}{2}(x-1)^4 + (x-1)^3 \right) + c_1 \left( \frac{1}{144}(-2226119942329(x-1)^4 - 2270644232(x-1)^3 + 24710760(x-1)^2 - 59544(x-1) + 144) + \frac{283830535}{12}(409(x-1) + 2)(x-1)^3 \log(x-1) \right)$$

## 25.10 problem 35.3 (d)

Internal problem ID [13662]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.3 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + \frac{y'}{x-3} + \frac{y}{x-4} = 0$$

With the expansion point for the power series method at  $x = 3$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 55

```
Order:=6;  
dsolve(diff(y(x),x$2)+1/(x-3)*diff(y(x),x)+1/(x-4)*y(x)=0,y(x),type='series',x=3);
```

$$y(x) = (\ln(x-3) c_2 + c_1) \left( 1 + \frac{1}{4}(x-3)^2 + \frac{1}{9}(x-3)^3 + \frac{5}{64}(x-3)^4 + \frac{49}{900}(x-3)^5 + O((x-3)^6) \right) + \left( -\frac{1}{4}(x-3)^2 - \frac{2}{27}(x-3)^3 - \frac{7}{128}(x-3)^4 - \frac{469}{13500}(x-3)^5 + O((x-3)^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 128

```
AsymptoticDSolveValue[y''[x]+1/(x-3)*y'[x]+1/(x-4)*y[x]==0,y[x],{x,3,5}]
```

$$\begin{aligned} y(x) \rightarrow & c_1 \left( \frac{49}{900}(x-3)^5 + \frac{5}{64}(x-3)^4 + \frac{1}{9}(x-3)^3 + \frac{1}{4}(x-3)^2 + 1 \right) \\ & + c_2 \left( -\frac{469(x-3)^5}{13500} - \frac{7}{128}(x-3)^4 - \frac{2}{27}(x-3)^3 - \frac{1}{4}(x-3)^2 \right. \\ & \left. + \left( \frac{49}{900}(x-3)^5 + \frac{5}{64}(x-3)^4 + \frac{1}{9}(x-3)^3 + \frac{1}{4}(x-3)^2 + 1 \right) \log(x-3) \right) \end{aligned}$$

## 25.11 problem 35.3 (e)

Internal problem ID [13663]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.3 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + \frac{y'}{(x-3)^2} + \frac{y}{(x-4)^2} = 0$$

With the expansion point for the power series method at  $x = 4$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 1199

**Order:=6;**

`dsolve(diff(y(x),x$2)+1/(x-3)^2*diff(y(x),x)+1/(x-4)^2*y(x)=0,y(x),type='series',x=4);`

$$\begin{aligned}
 y(x) = & \sqrt{x-4} \left( c_2(x-4)^{\frac{i\sqrt{3}}{2}} \left( 1 - \frac{1}{2}(x-4) + \frac{5i\sqrt{3}+7}{8i\sqrt{3}+16}(x-4)^2 \right. \right. \\
 & - \frac{1}{12} \frac{5+36i\sqrt{3}}{(i\sqrt{3}+3)(i\sqrt{3}+2)}(x-4)^3 + \frac{1}{96} \frac{1313i\sqrt{3}-865}{(i\sqrt{3}+4)(i\sqrt{3}+3)(i\sqrt{3}+2)}(x-4)^4 \\
 & \left. \left. - \frac{1}{240} \frac{-23995+15978i\sqrt{3}}{(i\sqrt{3}+5)(i\sqrt{3}+4)(i\sqrt{3}+3)(i\sqrt{3}+2)}(x-4)^5 + O((x-4)^6) \right) \right. \\
 & \left. + c_1(x-4)^{-\frac{i\sqrt{3}}{2}} \left( 1 - \frac{1}{2}(x-4) + \frac{5i\sqrt{3}-7}{8i\sqrt{3}-16}(x-4)^2 \right. \right. \\
 & + \frac{1}{12} \frac{-5+36i\sqrt{3}}{(i\sqrt{3}-3)(i\sqrt{3}-2)}(x-4)^3 + \frac{1}{96} \frac{1313i\sqrt{3}+865}{(i\sqrt{3}-4)(i\sqrt{3}-3)(i\sqrt{3}-2)}(x-4)^4 \\
 & \left. \left. + \frac{1}{240} \frac{23995+15978i\sqrt{3}}{(i\sqrt{3}-5)(i\sqrt{3}-4)(i\sqrt{3}-3)(i\sqrt{3}-2)}(x-4)^5 + O((x-4)^6) \right) \right)
 \end{aligned}$$



✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 2225

```
AsymptoticDSolveValue[y''[x]+1/(x-3)^2*y'[x]+1/(x-4)^2*y[x]==0,y[x],{x,4,5}]
```

Too large to display

## 25.12 problem 35.3 (f)

Internal problem ID [13664]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.3 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + \left(\frac{1}{x} - \frac{1}{3}\right) y' + \left(\frac{1}{x} - \frac{1}{4}\right) y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 59

**Order:=6;**

```
dsolve(diff(y(x), x$2)+(1/x-1/3)*diff(y(x), x)+(1/x-1/4)*y(x)=0, y(x), type='series', x=0);
```

$$y(x) = (\ln(x) c_2 + c_1) \left(1 - x + \frac{11}{48}x^2 - \frac{47}{1296}x^3 + \frac{11}{3072}x^4 - \frac{653}{2073600}x^5 + O(x^6)\right) \\ + \left(\frac{7}{3}x - \frac{101}{144}x^2 + \frac{10}{81}x^3 - \frac{6721}{497664}x^4 + \frac{229213}{186624000}x^5 + O(x^6)\right) c_2$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 113

```
AsymptoticDSolveValue[y''[x]+(1/x-1/3)*y'[x]+(1/x-1/4)*y[x]==0, y[x], {x, 0, 5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{653x^5}{2073600} + \frac{11x^4}{3072} - \frac{47x^3}{1296} + \frac{11x^2}{48} - x + 1\right) + c_2 \left(\frac{229213x^5}{186624000} - \frac{6721x^4}{497664} + \frac{10x^3}{81} - \frac{101x^2}{144} + \left(-\frac{653x^5}{2073600} + \frac{11x^4}{3072} - \frac{47x^3}{1296} + \frac{11x^2}{48} - x + 1\right) \log(x) + \frac{7x}{3}\right)$$

## 25.13 problem 35.3 (g)

Internal problem ID [13665]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.3 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(4x^2 - 1)y'' + \left(4 - \frac{2}{x}\right)y' + \frac{(-x^2 + 1)y}{x^2 + 1} = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 62

```
Order:=6;
```

```
dsolve((4*x^2-1)*diff(y(x),x$2)+(4-2/x)*diff(y(x),x)+(1-x^2)/(1+x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{c_1 \left(1 + \frac{1}{6}x^2 + \frac{1}{9}x^3 + \frac{1}{24}x^4 + \frac{31}{270}x^5 + O(x^6)\right) x + \left((-4)x - \frac{2}{3}x^3 - \frac{4}{9}x^4 - \frac{1}{6}x^5 + O(x^6)\right) \ln(x) c_2 + (1 + 4x)}{x}$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 79

```
AsymptoticDSolveValue[(4*x^2-1)*y'[x]+(4-2/x)*y'[x]+(1-x^2)/(1+x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{x^4}{24} + \frac{x^3}{9} + \frac{x^2}{6} + 1 \right) + c_1 \left( \frac{229x^4 + 480x^3 - 756x^2 + 1728x + 216}{216x} - \frac{2}{9}(2x^3 + 3x^2 + 18) \log(x) \right)$$

## 25.14 problem 35.3 (h)

Internal problem ID [13666]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.3 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$(x^2 + 4)^2 y'' + y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve((4+x^2)^2*diff(y(x),x$2)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{32}x^2 + \frac{17}{6144}x^4\right) y(0) + \left(x - \frac{1}{96}x^3 + \frac{49}{30720}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[(4+x^2)^2*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{49x^5}{30720} - \frac{x^3}{96} + x \right) + c_1 \left( \frac{17x^4}{6144} - \frac{x^2}{32} + 1 \right)$$

## 25.15 problem 35.4 (a)

Internal problem ID [13667]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.4 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 2y'x + (x^2 + 2)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+(x^2+2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^2 \left( 1 - \frac{1}{6} x^2 + \frac{1}{120} x^4 + O(x^6) \right) + c_2 x \left( 1 - \frac{1}{2} x^2 + \frac{1}{24} x^4 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 44

```
AsymptoticDSolveValue[x^2*y''[x]-2*x*y'[x]+(x^2+2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{x^5}{24} - \frac{x^3}{2} + x \right) + c_2 \left( \frac{x^6}{120} - \frac{x^4}{6} + x^2 \right)$$

## 25.16 problem 35.4 (b)

Internal problem ID [13668]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.4 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' + (-4x + 1)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 69

```
Order:=6;
```

```
dsolve(4*x^2*diff(y(x),x$2)+(1-4*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \sqrt{x} \left( (\ln(x) c_2 + c_1) \left( 1 + x + \frac{1}{4}x^2 + \frac{1}{36}x^3 + \frac{1}{576}x^4 + \frac{1}{14400}x^5 + O(x^6) \right) \right. \\ \left. + \left( (-2)x - \frac{3}{4}x^2 - \frac{11}{108}x^3 - \frac{25}{3456}x^4 - \frac{137}{432000}x^5 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 124

```
AsymptoticDSolveValue[4*x^2*y''[x]+(1-4*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left( \frac{x^5}{14400} + \frac{x^4}{576} + \frac{x^3}{36} + \frac{x^2}{4} + x + 1 \right) \\ + c_2 \left( \sqrt{x} \left( -\frac{137x^5}{432000} - \frac{25x^4}{3456} - \frac{11x^3}{108} - \frac{3x^2}{4} - 2x \right) \right. \\ \left. + \sqrt{x} \left( \frac{x^5}{14400} + \frac{x^4}{576} + \frac{x^3}{36} + \frac{x^2}{4} + x + 1 \right) \log(x) \right)$$

## 25.17 problem 35.4 (c)

Internal problem ID [13669]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.4 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y' x + (4x - 4) y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 61

```
Order:=6;
```

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(4*x-4)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 x^4 \left(1 - \frac{4}{5}x + \frac{4}{15}x^2 - \frac{16}{315}x^3 + \frac{2}{315}x^4 - \frac{8}{14175}x^5 + O(x^6)\right) + c_2 (\ln(x) (256x^4 - \frac{1024}{5}x^5 + O(x^6)) + (-14x^4 - \frac{1024}{5}x^5 + O(x^6)))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 79

```
AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]+(4*x-4)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{4x^4 + 16x^3 + 12x^2 + 12x + 9}{9x^2} - \frac{16}{9}x^2 \log(x) \right) + c_2 \left( \frac{2x^6}{315} - \frac{16x^5}{315} + \frac{4x^4}{15} - \frac{4x^3}{5} + x^2 \right)$$



## 25.18 problem 35.4 (d)

Internal problem ID [13670]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.4 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(-9x^4 + x^2)y'' - 6y'x + 10y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
Order:=6;  
dsolve((x^2-9*x^4)*diff(y(x),x$2)-6*x*diff(y(x),x)+10*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1x^5(1 + 18x^2 + 243x^4 + O(x^6)) + c_2x^2(12 - 108x^2 - 2916x^4 + O(x^6))$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 38

```
AsymptoticDSolveValue[(x^2-9*x^4)*y'[x]-6*x*y'[x]+10*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2(243x^9 + 18x^7 + x^5) + c_1(-243x^6 - 9x^4 + x^2)$$

## 25.19 problem 35.4 (e)

Internal problem ID [13671]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.4 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - y'x + \frac{y}{1-x} = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 47

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)-x*diff(y(x),x)+1/(1-x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left( (\ln(x) c_2 + c_1) (1 - x + O(x^6)) \right) + \left( 2x - \frac{1}{2}x^2 - \frac{1}{6}x^3 - \frac{1}{12}x^4 - \frac{1}{20}x^5 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 60

```
AsymptoticDSolveValue[x^2*y''[x]-x*y'[x]+1/(1-x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( x \left( -\frac{x^5}{20} - \frac{x^4}{12} - \frac{x^3}{6} - \frac{x^2}{2} + 2x \right) + (1-x)x \log(x) \right) + c_1(1-x)x$$

## 25.20 problem 35.4 (f)

Internal problem ID [13672]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.4 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [Lienard]

$$y'' + \frac{y'}{x} + y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 41

```
Order:=6;  
dsolve(diff(y(x),x$2)+1/x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (\ln(x) c_2 + c_1) \left( 1 - \frac{1}{4}x^2 + \frac{1}{64}x^4 + O(x^6) \right) + \left( \frac{1}{4}x^2 - \frac{3}{128}x^4 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 60

```
AsymptoticDSolveValue[y''[x]+1/x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{x^4}{64} - \frac{x^2}{4} + 1 \right) + c_2 \left( -\frac{3x^4}{128} + \frac{x^2}{4} + \left( \frac{x^4}{64} - \frac{x^2}{4} + 1 \right) \log(x) \right)$$

## 25.21 problem 35.4 (g)

Internal problem ID [13673]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.4 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [Bessel]

$$y'' + \frac{y'}{x} + \left(1 - \frac{1}{x^2}\right)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
Order:=6;  
dsolve(diff(y(x),x$2)+1/x*diff(y(x),x)+(1-1/x^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 x^2 \left(1 - \frac{1}{8}x^2 + \frac{1}{192}x^4 + O(x^6)\right) + c_2 \left(\ln(x) \left(x^2 - \frac{1}{8}x^4 + O(x^6)\right) + \left(-2 + \frac{3}{32}x^4 + O(x^6)\right)\right)}{x}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 58

```
AsymptoticDSolveValue[y'[x]+1/x*y'[x]+(1-1/x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{x^5}{192} - \frac{x^3}{8} + x \right) + c_1 \left( \frac{1}{16} x (x^2 - 8) \log(x) - \frac{5x^4 - 16x^2 - 64}{64x} \right)$$

## 25.22 problem 35.4 (h)

Internal problem ID [13674]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.4 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2y'' + (-2x^3 + 5x)y' + (-x^2 + 1)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

Order:=6;

```
dsolve(2*x^2*diff(y(x),x$2)+(5*x-2*x^3)*diff(y(x),x)+(1-x^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 \left(1 - \frac{1}{6}x^2 - \frac{1}{56}x^4 + O(x^6)\right) \sqrt{x} + c_2(1 + O(x^6))x}{x^{\frac{3}{2}}}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 34

```
AsymptoticDSolveValue[2*x^2*y''[x]+(5*x-2*x^3)*y'[x]+(1-x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_2 \left(-\frac{x^4}{56} - \frac{x^2}{6} + 1\right)}{x} + \frac{c_1}{\sqrt{x}}$$

## 25.23 problem 35.4 (i)

Internal problem ID [13675]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.4 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - (2x^2 + 5x) y' + (9 + 4x) y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 69

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)-(5*x+2*x^2)*diff(y(x),x)+(9+4*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left( (\ln(x) c_2 + c_1) \left( 1 + 2x + 2x^2 + \frac{4}{3}x^3 + \frac{2}{3}x^4 + \frac{4}{15}x^5 + O(x^6) \right) \right. \\ \left. + \left( (-2)x - 3x^2 - \frac{22}{9}x^3 - \frac{25}{18}x^4 - \frac{137}{225}x^5 + O(x^6) \right) c_2 \right) x^3$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 116

```
AsymptoticDSolveValue[x^2*y''[x]-(5*x+2*x^2)*y'[x]+(9+4*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{4x^5}{15} + \frac{2x^4}{3} + \frac{4x^3}{3} + 2x^2 + 2x + 1 \right) x^3 \\ + c_2 \left( \left( -\frac{137x^5}{225} - \frac{25x^4}{18} - \frac{22x^3}{9} - 3x^2 - 2x \right) x^3 \right. \\ \left. + \left( \frac{4x^5}{15} + \frac{2x^4}{3} + \frac{4x^3}{3} + 2x^2 + 2x + 1 \right) x^3 \log(x) \right)$$

## 25.24 problem 35.4 (j)

Internal problem ID [13676]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.4 (j).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(-3x^3 + 3x^2)y'' - (5x^2 + 4x)y' + 2y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
Order:=6;
```

```
dsolve((3*x^2-3*x^3)*diff(y(x),x$2)-(4*x+5*x^2)*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0
```

$$y(x) = c_1 x^{\frac{1}{3}} \left( 1 - \frac{1}{2}x - 2x^2 - \frac{7}{2}x^3 - 5x^4 - \frac{13}{2}x^5 + O(x^6) \right) \\ + c_2 x^2 (1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + O(x^6))$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 74

```
AsymptoticDSolveValue[(3*x^2-3*x^3)*y'[x]-(4*x+5*x^2)*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 (6x^5 + 5x^4 + 4x^3 + 3x^2 + 2x + 1) x^2 + c_2 \left( -\frac{13x^5}{2} - 5x^4 - \frac{7x^3}{2} - 2x^2 - \frac{x}{2} + 1 \right) \sqrt[3]{x}$$



## 25.25 problem 35.4 (k)

Internal problem ID [13677]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.4 (k).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [Laguerre]

$$x^2 y'' - (x^2 + x) y' + 4yx = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 52

```
Order:=6;  
dsolve(x^2*dif(y(x),x$2)-(x+x^2)*dif(y(x),x)+4*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^2 \left( 1 - \frac{2}{3}x + \frac{1}{12}x^2 + O(x^6) \right) + (12x^2 - 8x^3 + x^4 + O(x^6)) \ln(x) c_2 \\ + \left( -2 - 8x - 7x^2 + \frac{58}{3}x^3 - \frac{25}{6}x^4 + \frac{1}{15}x^5 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 70

```
AsymptoticDSolveValue[x^2*y'[x]-(x+x^2)*y'[x]+4*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{x^4}{12} - \frac{2x^3}{3} + x^2 \right) \\ + c_1 \left( \frac{1}{6}(14x^4 - 70x^3 + 39x^2 + 24x + 6) - \frac{1}{2}x^2(x^2 - 8x + 12) \log(x) \right)$$

## 25.26 problem 35.4 (L)

Internal problem ID [13678]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.4 (L).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' + 8y'x^2 + y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 67

```
Order:=6;  
dsolve(4*x^2*diff(y(x),x$2)+8*x^2*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \sqrt{x} \left( (\ln(x) c_2 + c_1) \left( 1 - x + \frac{3}{4}x^2 - \frac{5}{12}x^3 + \frac{35}{192}x^4 - \frac{21}{320}x^5 + O(x^6) \right) \right. \\ \left. + \left( -\frac{1}{4}x^2 + \frac{1}{4}x^3 - \frac{19}{128}x^4 + \frac{25}{384}x^5 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 125

```
AsymptoticDSolveValue[4*x^2*y''[x]+8*x^2*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left( -\frac{21x^5}{320} + \frac{35x^4}{192} - \frac{5x^3}{12} + \frac{3x^2}{4} - x + 1 \right) + c_2 \left( \sqrt{x} \left( \frac{25x^5}{384} - \frac{19x^4}{128} + \frac{x^3}{4} - \frac{x^2}{4} \right) \right. \\ \left. + \sqrt{x} \left( -\frac{21x^5}{320} + \frac{35x^4}{192} - \frac{5x^3}{12} + \frac{3x^2}{4} - x + 1 \right) \log(x) \right)$$

## 25.27 problem 35.4 (m)

Internal problem ID [13679]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.4 (m).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + (-x^4 + x) y' + 3yx^3 = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)+(x-x^4)*diff(y(x),x)+3*x^3*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (\ln(x) c_2 + c_1) \left(1 - \frac{1}{3}x^3 + O(x^6)\right) + \left(\frac{1}{3}x^3 + O(x^6)\right) c_2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 39

```
AsymptoticDSolveValue[x^2*y''[x]+(x-x^4)*y'[x]+3*x^3*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(1 - \frac{x^3}{3}\right) + c_2 \left(\frac{x^3}{3} + \left(1 - \frac{x^3}{3}\right) \log(x)\right)$$

## 25.28 problem 35.4 (n)

Internal problem ID [13680]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.4 (n).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(9x^3 + 9x^2)y'' + (27x^2 + 9x)y' + (8x - 1)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 47

Order:=6;

```
dsolve((9*x^2+9*x^3)*diff(y(x),x$2)+(9*x+27*x^2)*diff(y(x),x)+(8*x-1)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{(-x^5 + x^4 - x^3 + x^2 - x + 1)(x^{\frac{2}{3}}c_2 + c_1)}{x^{\frac{1}{3}}} + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 62

```
AsymptoticDSolveValue[(9*x^2+9*x^3)*y'[x]+(9*x+27*x^2)*y'[x]+(8*x-1)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt[3]{x}(-x^5 + x^4 - x^3 + x^2 - x + 1) + \frac{c_2(-x^5 + x^4 - x^3 + x^2 - x + 1)}{\sqrt[3]{x}}$$

## 25.29 problem 35.5 (a)

Internal problem ID [13681]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.5 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x - 3)y'' + (x - 3)y' + y = 0$$

With the expansion point for the power series method at  $x = 3$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 62

```
Order:=6;
```

```
dsolve((x-3)*diff(y(x),x$2)+(x-3)*diff(y(x),x)+y(x)=0,y(x),type='series',x=3);
```

$$\begin{aligned} y(x) = & c_1(x - 3) \left( 1 - (x - 3) + \frac{1}{2}(x - 3)^2 - \frac{1}{6}(x - 3)^3 + \frac{1}{24}(x - 3)^4 - \frac{1}{120}(x - 3)^5 \right. \\ & \left. + O((x - 3)^6) \right) + c_2 \left( \ln(x - 3) \left( -(x - 3) + (x - 3)^2 - \frac{1}{2}(x - 3)^3 + \frac{1}{6}(x - 3)^4 \right. \right. \\ & \left. \left. - \frac{1}{24}(x - 3)^5 + O((x - 3)^6) \right) \right) \\ & + \left( 1 - (x - 3) + \frac{1}{4}(x - 3)^3 - \frac{5}{36}(x - 3)^4 + \frac{13}{288}(x - 3)^5 + O((x - 3)^6) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 105

```
AsymptoticDSolveValue[(x-3)*y''[x]+(x-3)*y'[x]+y[x]==0,y[x],{x,3,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{1}{24}(x-3)^5 - \frac{1}{6}(x-3)^4 + \frac{1}{2}(x-3)^3 - (x-3)^2 + x - 3 \right) \\ + c_1 \left( \frac{1}{36}(-11(x-3)^4 + 27(x-3)^3 - 36(x-3)^2 + 36) \right. \\ \left. + \frac{1}{6}((x-3)^3 - 3(x-3)^2 + 6(x-3) - 6)(x-3) \log(x-3) \right)$$

## 25.30 problem 35.5 (b)

Internal problem ID [13682]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.5 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + \frac{2y'}{x+2} + y = 0$$

With the expansion point for the power series method at  $x = -2$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)+2/(x+2)*diff(y(x),x)+y(x)=0,y(x),type='series',x=-2);
```

$$y(x) = c_1 \left( 1 - \frac{1}{6}(x+2)^2 + \frac{1}{120}(x+2)^4 + O((x+2)^6) \right) \\ + \frac{c_2 \left( 1 - \frac{1}{2}(x+2)^2 + \frac{1}{24}(x+2)^4 + O((x+2)^6) \right)}{x+2}$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 54

```
AsymptoticDSolveValue[y''[x]+2/(x+2)*y'[x]+y[x]==0,y[x],{x,-2,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{1}{24}(x+2)^3 + \frac{1}{2}(-x-2) + \frac{1}{x+2} \right) + c_2 \left( \frac{1}{120}(x+2)^4 - \frac{1}{6}(x+2)^2 + 1 \right)$$

## 25.31 problem 35.5 (c)

Internal problem ID [13683]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.5 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4y'' + \frac{(4x - 3)y}{(x - 1)^2} = 0$$

With the expansion point for the power series method at  $x = 1$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 77

```
Order:=6;  
dsolve(4*diff(y(x),x$2)+(4*x-3)/(x-1)^2*y(x)=0,y(x),type='series',x=1);
```

$$y(x) = \sqrt{-1+x} \left( (\ln(-1+x) c_2 + c_1) \left( 1 - (-1+x) + \frac{1}{4}(-1+x)^2 - \frac{1}{36}(-1+x)^3 + \frac{1}{576}(-1+x)^4 - \frac{1}{14400}(-1+x)^5 + O((-1+x)^6) \right) + \left( 2(-1+x) - \frac{3}{4}(-1+x)^2 + \frac{11}{108}(-1+x)^3 - \frac{25}{3456}(-1+x)^4 + \frac{137}{432000}(-1+x)^5 + O((-1+x)^6) \right) c_2 \right)$$



✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 162

```
AsymptoticDSolveValue[4*y''[x]+(4*x-3)/(x-1)^2*y[x]==0,y[x],{x,1,5}]
```

$$\begin{aligned} y(x) \rightarrow & c_1 \left( -\frac{(x-1)^5}{14400} + \frac{1}{576}(x-1)^4 - \frac{1}{36}(x-1)^3 + \frac{1}{4}(x-1)^2 - x + 2 \right) \sqrt{x-1} \\ & + c_2 \left( \sqrt{x-1} \left( \frac{137(x-1)^5}{432000} - \frac{25(x-1)^4}{3456} + \frac{11}{108}(x-1)^3 - \frac{3}{4}(x-1)^2 + 2(x-1) \right) \right. \\ & \left. + \left( -\frac{(x-1)^5}{14400} + \frac{1}{576}(x-1)^4 - \frac{1}{36}(x-1)^3 + \frac{1}{4}(x-1)^2 - x + 2 \right) \sqrt{x-1} \log(x-1) \right) \end{aligned}$$

## 25.32 problem 35.5 (d)

Internal problem ID [13684]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 35. Modified Power series solutions and basic method of Frobenius. Additional Exercises. page 715

**Problem number:** 35.5 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x - 3)^2 y'' + (x^2 - 3x) y' - 3y = 0$$

With the expansion point for the power series method at  $x = 3$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 49

Order:=6;

```
dsolve((x-3)^2*diff(y(x),x$2)+(x^2-3*x)*diff(y(x),x)-3*y(x)=0,y(x),type='series',x=3);
```

$$y(x) = \frac{c_1(x-3)^4 \left(1 - \frac{1}{5}(x-3) + \frac{1}{30}(x-3)^2 - \frac{1}{210}(x-3)^3 + \frac{1}{1680}(x-3)^4 - \frac{1}{15120}(x-3)^5 + O((x-3)^6)\right) + c_2(x-3)^3}{(x-3)^3}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 81

```
AsymptoticDSolveValue[(x-3)^2*y''[x]+(x^2-3*x)*y'[x]-3*y[x]==0,y[x],{x,3,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{x-3}{24} + \frac{1}{2(x-3)} - \frac{1}{(x-3)^2} + \frac{1}{(x-3)^3} - \frac{1}{6} \right) + c_2 \left( \frac{(x-3)^5}{1680} - \frac{1}{210}(x-3)^4 + \frac{1}{30}(x-3)^3 - \frac{1}{5}(x-3)^2 + x-3 \right)$$

**26 Chapter 36. The big theorem on the the  
Frobenius method. Additional Exercises. page  
739**

26.1 problem 36.2 (a)	850
26.2 problem 36.2 (b)	851
26.3 problem 36.2 (c)	852
26.4 problem 36.2 (d)	853
26.5 problem 36.2 (e)	854
26.6 problem 36.2 (f)	856
26.7 problem 36.2 (g)	857
26.8 problem 36.2 (h)	858
26.9 problem 36.2 (i)	859
26.10 problem 36.2 (j)	860
26.11 problem 36.2 (k)	861
26.12 problem 36.2 (L)	863
26.13 problem 36.6 (a)	865
26.14 problem 36.6 (b)	867
26.15 problem 36.6 (c)	868
26.16 problem 36.6 (d)	869

## 26.1 problem 36.2 (a)

Internal problem ID [13685]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.2 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 2y'x + (-x^2 + 2)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+(2-x^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^2 \left( 1 + \frac{1}{6} x^2 + \frac{1}{120} x^4 + O(x^6) \right) + c_2 x \left( 1 + \frac{1}{2} x^2 + \frac{1}{24} x^4 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 44

```
AsymptoticDSolveValue[x^2*y''[x]-2*x*y'[x]+(2-x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{x^5}{24} + \frac{x^3}{2} + x \right) + c_2 \left( \frac{x^6}{120} + \frac{x^4}{6} + x^2 \right)$$

## 26.2 problem 36.2 (b)

Internal problem ID [13686]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.2 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 2y'x^2 + (x^2 - 2)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 47

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)-2*x^2*diff(y(x),x)+(x^2-2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^2 \left( 1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + O(x^6) \right) + \frac{c_2 (12 + 12x + 6x^2 + 4x^3 + \frac{5}{2}x^4 + \frac{11}{10}x^5 + O(x^6))}{x}$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 62

```
AsymptoticDSolveValue[x^2*y''[x]-2*x^2*y'[x]+(x^2-2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{5x^3}{24} + \frac{x^2}{3} + \frac{x}{2} + \frac{1}{x} + 1 \right) + c_2 \left( \frac{x^6}{24} + \frac{x^5}{6} + \frac{x^4}{2} + x^3 + x^2 \right)$$

## 26.3 problem 36.2 (c)

Internal problem ID [13687]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.2 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [Lienard]

$$y'' + \frac{y'}{x} + y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 41

```
Order:=6;  
dsolve(diff(y(x),x$2)+1/x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (\ln(x) c_2 + c_1) \left( 1 - \frac{1}{4}x^2 + \frac{1}{64}x^4 + O(x^6) \right) + \left( \frac{1}{4}x^2 - \frac{3}{128}x^4 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 60

```
AsymptoticDSolveValue[y''[x]+1/x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{x^4}{64} - \frac{x^2}{4} + 1 \right) + c_2 \left( -\frac{3x^4}{128} + \frac{x^2}{4} + \left( \frac{x^4}{64} - \frac{x^2}{4} + 1 \right) \log(x) \right)$$

## 26.4 problem 36.2 (d)

Internal problem ID [13688]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.2 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(-x^2 + 2)y'' + (4x^2 + 5x)y' + (x^2 + 1)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 47

Order:=6;

`dsolve(x^2*(2-x^2)*diff(y(x),x$2)+(5*x+4*x^2)*diff(y(x),x)+(1+x^2)*y(x)=0,y(x),type='series'`

$y(x)$

$$= \frac{c_1 \left( 1 + 4x + \frac{1}{6}x^2 - \frac{14}{45}x^3 + \frac{209}{2520}x^4 - \frac{823}{28350}x^5 + O(x^6) \right) \sqrt{x} + c_2 \left( 1 + \frac{2}{3}x - \frac{19}{120}x^2 + \frac{1}{180}x^3 - \frac{23}{51840}x^4 + \frac{557}{142560}x^5 \right)}{x^{\frac{3}{2}}}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 86

`AsymptoticDSolveValue[x^2*(2-x^2)*y'[x]+(5*x+4*x^2)*y'[x]+(1+x^2)*y[x]==0,y[x],{x,0,5}]`

$$y(x) \rightarrow \frac{c_1 \left( \frac{557x^5}{1425600} - \frac{23x^4}{51840} + \frac{x^3}{180} - \frac{19x^2}{120} + \frac{2x}{3} + 1 \right)}{\sqrt{x}} + \frac{c_2 \left( -\frac{823x^5}{28350} + \frac{209x^4}{2520} - \frac{14x^3}{45} + \frac{x^2}{6} + 4x + 1 \right)}{x}$$

## 26.5 problem 36.2 (e)

Internal problem ID [13689]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.2 (e).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - (2x^2 + 5x) y' + 9y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 69

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)-(5*x+2*x^2)*diff(y(x),x)+9*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left( (\ln(x) c_2 + c_1) \left( 1 + 6x + 12x^2 + \frac{40}{3}x^3 + 10x^4 + \frac{28}{5}x^5 + O(x^6) \right) \right. \\ \left. + \left( (-10)x - 29x^2 - \frac{346}{9}x^3 - \frac{193}{6}x^4 - \frac{1459}{75}x^5 + O(x^6) \right) c_2 \right) x^3$$



✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 112

```
AsymptoticDSolveValue[x^2*y''[x]-(5*x+2*x^2)*y'[x]+9*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{28x^5}{5} + 10x^4 + \frac{40x^3}{3} + 12x^2 + 6x + 1 \right) x^3 \\ + c_2 \left( \left( -\frac{1459x^5}{75} - \frac{193x^4}{6} - \frac{346x^3}{9} - 29x^2 - 10x \right) x^3 \right. \\ \left. + \left( \frac{28x^5}{5} + 10x^4 + \frac{40x^3}{3} + 12x^2 + 6x + 1 \right) x^3 \log(x) \right)$$

## 26.6 problem 36.2 (f)

Internal problem ID [13690]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.2 (f).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(2x + 1)y'' + y'x + (4x^3 - 4)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 63

```
Order:=6;
```

```
dsolve(x^2*(1+2*x)*diff(y(x),x$2)+x*diff(y(x),x)+(4*x^3-4)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 x^4 \left(1 - \frac{4}{5}x + \frac{4}{5}x^2 - \frac{116}{105}x^3 + \frac{311}{210}x^4 - \frac{358}{175}x^5 + O(x^6)\right) + c_2 (\ln(x) (576x^4 - \frac{2304}{5}x^5 + O(x^6)) + (-144 - \dots)}{x^2}$$

✓ Solution by Mathematica

Time used: 0.044 (sec). Leaf size: 77

```
AsymptoticDSolveValue[x^2*(1+2*x)*y'[x]+x*y'[x]+(4*x^3-4)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{19x^4 + 4x^3 + 12x^2 + 12x + 3}{3x^2} - 4x^2 \log(x) \right) + c_2 \left( \frac{311x^6}{210} - \frac{116x^5}{105} + \frac{4x^4}{5} - \frac{4x^3}{5} + x^2 \right)$$

## 26.7 problem 36.2 (g)

Internal problem ID [13691]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.2 (g).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' + 8y'x + (-4x + 1)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 69

```
Order:=6;
dsolve(4*x^2*diff(y(x),x$2)+8*x*diff(y(x),x)+(1-4*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{(\ln(x) c_2 + c_1) \left(1 + x + \frac{1}{4}x^2 + \frac{1}{36}x^3 + \frac{1}{576}x^4 + \frac{1}{14400}x^5 + O(x^6)\right) + \left((-2)x - \frac{3}{4}x^2 - \frac{11}{108}x^3 - \frac{25}{3456}x^4 - \frac{1}{43200}x^5\right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 124

```
AsymptoticDSolveValue[4*x^2*y'[x]+8*x*y'[x]+(1-4*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_1 \left( \frac{x^5}{14400} + \frac{x^4}{576} + \frac{x^3}{36} + \frac{x^2}{4} + x + 1 \right)}{\sqrt{x}} + c_2 \left( \frac{-\frac{137x^5}{432000} - \frac{25x^4}{3456} - \frac{11x^3}{108} - \frac{3x^2}{4} - 2x}{\sqrt{x}} + \frac{\left( \frac{x^5}{14400} + \frac{x^4}{576} + \frac{x^3}{36} + \frac{x^2}{4} + x + 1 \right) \log(x)}{\sqrt{x}} \right)$$

## 26.8 problem 36.2 (h)

Internal problem ID [13692]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.2 (h).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y'x - (2x + 1)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 63

```
Order:=6;
```

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-(1+2*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 x^2 \left(1 + \frac{2}{3}x + \frac{1}{6}x^2 + \frac{1}{45}x^3 + \frac{1}{540}x^4 + \frac{1}{9450}x^5 + O(x^6)\right) + c_2 (\ln(x) (4x^2 + \frac{8}{3}x^3 + \frac{2}{3}x^4 + \frac{4}{45}x^5 + O(x^6)) + x}{x}$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 83

```
AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]-(1+2*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{31x^4 + 88x^3 + 36x^2 - 72x + 36}{36x} - \frac{1}{3}x(x^2 + 4x + 6) \log(x) \right) + c_2 \left( \frac{x^5}{540} + \frac{x^4}{45} + \frac{x^3}{6} + \frac{2x^2}{3} + x \right)$$

## 26.9 problem 36.2 (i)

Internal problem ID [13693]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.2 (i).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y''x + 4y' + \frac{12y}{(x+2)^2} = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 42

```
Order:=6;
```

```
dsolve(x*diff(y(x),x$2)+4*diff(y(x),x)+12/(x+2)^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 \left( 1 - \frac{3}{4}x + \frac{21}{40}x^2 - \frac{27}{80}x^3 + \frac{33}{160}x^4 - \frac{39}{320}x^5 + O(x^6) \right) \\ + \frac{c_2(12 + 18x + 9x^2 + \frac{9}{8}x^4 - \frac{63}{80}x^5 + O(x^6))}{x^3}$$

✓ Solution by Mathematica

Time used: 0.065 (sec). Leaf size: 59

```
AsymptoticDSolveValue[x*y'[x]+4*y'[x]+12/(x+2)^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{1}{x^3} + \frac{3}{2x^2} + \frac{3}{4x} + \frac{1}{8} \right) + c_2 \left( \frac{33x^4}{160} - \frac{27x^3}{80} + \frac{21x^2}{40} - \frac{3x}{4} + 1 \right)$$

## 26.10 problem 36.2 (j)

Internal problem ID [13694]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.2 (j).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y''x + 4y' + \frac{12y}{(x+2)^2} = 0$$

With the expansion point for the power series method at  $x = -2$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
Order:=6;
dsolve(x*diff(y(x),x$2)+4*diff(y(x),x)+12/(x+2)^2*y(x)=0,y(x),type='series',x=-2);
```

$$y(x) = \frac{c_1(x+2)^5 \left(1 + \frac{3}{2}(x+2) + \frac{3}{2}(x+2)^2 + \frac{5}{4}(x+2)^3 + \frac{15}{16}(x+2)^4 + \frac{21}{32}(x+2)^5 + O((x+2)^6)\right) + c_2(2880)}{(x+2)^2}$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 70

```
AsymptoticDSolveValue[x*y''[x]+4*y'[x]+12/(x+2)^2*y[x]==0,y[x],{x,-2,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{1}{4(x+2)} + \frac{1}{(x+2)^2} + \frac{1}{24} \right) + c_2 \left( \frac{15}{16}(x+2)^7 + \frac{5}{4}(x+2)^6 + \frac{3}{2}(x+2)^5 + \frac{3}{2}(x+2)^4 + (x+2)^3 \right)$$

## 26.11 problem 36.2 (k)

Internal problem ID [13695]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.2 (k).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x - 3)y'' + (x - 3)y' + y = 0$$

With the expansion point for the power series method at  $x = 3$ .

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 62

```
Order:=6;
```

```
dsolve((x-3)*diff(y(x),x$2)+(x-3)*diff(y(x),x)+y(x)=0,y(x),type='series',x=3);
```

$$\begin{aligned} y(x) = & c_1(x - 3) \left( 1 - (x - 3) + \frac{1}{2}(x - 3)^2 - \frac{1}{6}(x - 3)^3 + \frac{1}{24}(x - 3)^4 - \frac{1}{120}(x - 3)^5 \right. \\ & \left. + O((x - 3)^6) \right) + c_2 \left( \ln(x - 3) \left( -(x - 3) + (x - 3)^2 - \frac{1}{2}(x - 3)^3 + \frac{1}{6}(x - 3)^4 \right. \right. \\ & \left. \left. - \frac{1}{24}(x - 3)^5 + O((x - 3)^6) \right) \right) \\ & + \left( 1 - (x - 3) + \frac{1}{4}(x - 3)^3 - \frac{5}{36}(x - 3)^4 + \frac{13}{288}(x - 3)^5 + O((x - 3)^6) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 105

```
AsymptoticDSolveValue[(x-3)*y''[x]+(x-3)*y'[x]+y[x]==0,y[x],{x,3,5}]
```

$$\begin{aligned} y(x) \rightarrow & c_2 \left( \frac{1}{24}(x-3)^5 - \frac{1}{6}(x-3)^4 + \frac{1}{2}(x-3)^3 - (x-3)^2 + x - 3 \right) \\ & + c_1 \left( \frac{1}{36}(-11(x-3)^4 + 27(x-3)^3 - 36(x-3)^2 + 36) \right. \\ & \left. + \frac{1}{6}((x-3)^3 - 3(x-3)^2 + 6(x-3) - 6)(x-3) \log(x-3) \right) \end{aligned}$$



## 26.12 problem 36.2 (L)

Internal problem ID [13696]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.2 (L).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [\_Gegenbauer, [\_2nd\_order, \_linear, ‘\_with\_symmetry\_[0,F(x)]’]

$$(-x^2 + 1)y'' - y'x + 3y = 0$$

With the expansion point for the power series method at  $x = 1$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 46

```
Order:=6;  
dsolve((1-x^2)*diff(y(x),x$2)-x*diff(y(x),x)+3*y(x)=0,y(x),type='series',x=1);
```

$$y(x) = c_1 \sqrt{-1+x} \left( 1 + \frac{11}{12}(-1+x) + \frac{11}{160}(-1+x)^2 - \frac{143}{13440}(-1+x)^3 + \frac{5291}{1935360}(-1+x)^4 - \frac{11063}{12902400}(-1+x)^5 + O((-1+x)^6) \right) + c_2 \left( 1 + 3(-1+x) + (-1+x)^2 - \frac{1}{15}(-1+x)^3 + \frac{1}{70}(-1+x)^4 - \frac{13}{3150}(-1+x)^5 + O((-1+x)^6) \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 101

```
AsymptoticDSolveValue[(1-x^2)*y'[x]-x*y'[x]+3*y[x]==0,y[x],{x,1,5}]
```

$$y(x) \rightarrow c_1 \left( -\frac{11063(x-1)^5}{12902400} + \frac{5291(x-1)^4}{1935360} - \frac{143(x-1)^3}{13440} + \frac{11}{160}(x-1)^2 + \frac{11(x-1)}{12} + 1 \right) \sqrt{x-1} \\ + c_2 \left( -\frac{13(x-1)^5}{3150} + \frac{1}{70}(x-1)^4 - \frac{1}{15}(x-1)^3 + (x-1)^2 + 3(x-1) + 1 \right)$$

## 26.13 problem 36.6 (a)

Internal problem ID [13697]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.6 (a).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' + (-4x + 1)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 69

```
Order:=6;
```

```
dsolve(4*x^2*diff(y(x),x$2)+(1-4*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \sqrt{x} \left( (\ln(x) c_2 + c_1) \left( 1 + x + \frac{1}{4}x^2 + \frac{1}{36}x^3 + \frac{1}{576}x^4 + \frac{1}{14400}x^5 + O(x^6) \right) \right. \\ \left. + \left( (-2)x - \frac{3}{4}x^2 - \frac{11}{108}x^3 - \frac{25}{3456}x^4 - \frac{137}{432000}x^5 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 124

```
AsymptoticDSolveValue[4*x^2*y''[x]+(1-4*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left( \frac{x^5}{14400} + \frac{x^4}{576} + \frac{x^3}{36} + \frac{x^2}{4} + x + 1 \right) \\ + c_2 \left( \sqrt{x} \left( -\frac{137x^5}{432000} - \frac{25x^4}{3456} - \frac{11x^3}{108} - \frac{3x^2}{4} - 2x \right) \right. \\ \left. + \sqrt{x} \left( \frac{x^5}{14400} + \frac{x^4}{576} + \frac{x^3}{36} + \frac{x^2}{4} + x + 1 \right) \log(x) \right)$$

## 26.14 problem 36.6 (b)

Internal problem ID [13698]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.6 (b).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [Lienard]

$$y'' + \frac{y'}{x} + y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 41

```
Order:=6;  
dsolve(diff(y(x),x$2)+1/x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (\ln(x) c_2 + c_1) \left( 1 - \frac{1}{4}x^2 + \frac{1}{64}x^4 + O(x^6) \right) + \left( \frac{1}{4}x^2 - \frac{3}{128}x^4 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 60

```
AsymptoticDSolveValue[y''[x]+1/x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{x^4}{64} - \frac{x^2}{4} + 1 \right) + c_2 \left( -\frac{3x^4}{128} + \frac{x^2}{4} + \left( \frac{x^4}{64} - \frac{x^2}{4} + 1 \right) \log(x) \right)$$

## 26.15 problem 36.6 (c)

Internal problem ID [13699]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.6 (c).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type [Laguerre]

$$x^2 y'' - (x^2 + x) y' + 4yx = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 52

Order:=6;

```
dsolve(x^2*dif(y(x),x$2)-(x+x^2)*dif(y(x),x)+4*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^2 \left( 1 - \frac{2}{3}x + \frac{1}{12}x^2 + O(x^6) \right) + (12x^2 - 8x^3 + x^4 + O(x^6)) \ln(x) c_2 \\ + \left( -2 - 8x - 7x^2 + \frac{58}{3}x^3 - \frac{25}{6}x^4 + \frac{1}{15}x^5 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 70

```
AsymptoticDSolveValue[x^2*y'[x]-(x+x^2)*y'[x]+4*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left( \frac{x^4}{12} - \frac{2x^3}{3} + x^2 \right) \\ + c_1 \left( \frac{1}{6}(14x^4 - 70x^3 + 39x^2 + 24x + 6) - \frac{1}{2}x^2(x^2 - 8x + 12) \log(x) \right)$$

## 26.16 problem 36.6 (d)

Internal problem ID [13700]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 36. The big theorem on the the Frobenius method. Additional Exercises. page 739

**Problem number:** 36.6 (d).

**ODE order:** 2.

**ODE degree:** 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y' x + (4x - 4)y = 0$$

With the expansion point for the power series method at  $x = 0$ .

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 61

```
Order:=6;
```

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(4*x-4)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 x^4 \left(1 - \frac{4}{5}x + \frac{4}{15}x^2 - \frac{16}{315}x^3 + \frac{2}{315}x^4 - \frac{8}{14175}x^5 + O(x^6)\right) + c_2 (\ln(x) (256x^4 - \frac{1024}{5}x^5 + O(x^6))) + (-14}{x^2}$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 79

```
AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]+(4*x-4)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left( \frac{4x^4 + 16x^3 + 12x^2 + 12x + 9}{9x^2} - \frac{16}{9}x^2 \log(x) \right) + c_2 \left( \frac{2x^6}{315} - \frac{16x^5}{315} + \frac{4x^4}{15} - \frac{4x^3}{5} + x^2 \right)$$

**27 Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786**

27.1	problem 38.1	871
27.2	problem 38.2	872
27.3	problem 38.3	873
27.4	problem 38.4	874
27.5	problem 38.5	875
27.6	problem 38.6	876
27.7	problem 38.10 (a)	877
27.8	problem 38.10 (b)	878
27.9	problem 38.10 (c)	879
27.10	problem 38.10 (d)	880
27.11	problem 38.10 (e)	881
27.12	problem 38.10 (f)	882
27.13	problem 38.10 (g)	883
27.14	problem 38.10 (h)	884
27.15	problem 38.10 (i)	885
27.16	problem 38.10 (j)	886
27.17	problem 38.10 (k)	887
27.18	problem 38.10 (L)	888
27.19	problem 38.11	890



## 27.1 problem 38.1

Internal problem ID [13701]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.1.

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= 2y(t) \\ y'(t) &= 1 - 2x(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 36

```
dsolve([diff(x(t),t)=2*y(t),diff(y(t),t)=1-2*x(t)],[x(t), y(t)], singsol=all)
```

$$\begin{aligned}x(t) &= -c_1 \cos(2t) + c_2 \sin(2t) + \frac{1}{2} \\ y(t) &= c_1 \sin(2t) + c_2 \cos(2t)\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 42

```
DSolve[{x'[t]==2*y[t],y'[t]==1-2*x[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$\begin{aligned}x(t) &\rightarrow c_1 \cos(2t) + c_2 \sin(2t) + \frac{1}{2} \\ y(t) &\rightarrow c_2 \cos(2t) - c_1 \sin(2t)\end{aligned}$$

## 27.2 problem 38.2

Internal problem ID [13702]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.2.

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 4x(t) - 3y(t)$$

$$y'(t) = 6x(t) - 7y(t)$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 36

```
dsolve([diff(x(t),t)=4*x(t)-3*y(t),diff(y(t),t)=6*x(t)-7*y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = \frac{3c_1 e^{2t}}{2} + \frac{c_2 e^{-5t}}{3}$$

$$y(t) = c_1 e^{2t} + c_2 e^{-5t}$$

### ✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 74

```
DSolve[{x'[t]==4*x[t]-3*y[t],y'[t]==6*x[t]-7*y[t]},{x[t],y[t]},t,IncludeSingularSolutions ->
```

$$x(t) \rightarrow \frac{1}{7}e^{-5t}(c_1(9e^{7t} - 2) - 3c_2(e^{7t} - 1))$$

$$y(t) \rightarrow \frac{1}{7}e^{-5t}(6c_1(e^{7t} - 1) + c_2(9 - 2e^{7t}))$$

## 27.3 problem 38.3

Internal problem ID [13703]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.3.

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= \frac{15y(t)}{t} - \frac{2x(t)}{t} \\y'(t) &= \frac{x(t)}{t}\end{aligned}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 34

```
dsolve([t*diff(x(t),t)+2*x(t)=15*y(t),t*diff(y(t),t)=x(t)],[x(t), y(t)], singsol=all)
```

$$\begin{aligned}x(t) &= -\frac{-3c_2t^8 + 5c_1}{t^5} \\y(t) &= \frac{c_2t^8 + c_1}{t^5}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 36

```
DSolve[{t*x'[t]+2*x[t]==15*y[t],t*y'[t]==x[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$\begin{aligned}x(t) &\rightarrow 3c_2t^3 - \frac{5c_1}{t^5} \\y(t) &\rightarrow \frac{c_2t^8 + c_1}{t^5}\end{aligned}$$

## 27.4 problem 38.4

Internal problem ID [13704]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.4.

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= x(t) + 2y(t) \\y'(t) &= 5x(t) - 2y(t)\end{aligned}$$

With initial conditions

$$[x(0) = 7, y(0) = -7]$$

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 34

```
dsolve([diff(x(t),t) = x(t)+2*y(t), diff(y(t),t) = 5*x(t)-2*y(t), x(0) = 7, y(0) = -7],[x(t)
```

$$\begin{aligned}x(t) &= 4e^{-4t} + 3e^{3t} \\y(t) &= -10e^{-4t} + 3e^{3t}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 44

```
DSolve[{x'[t]==x[t]+2*y[t],y'[t]==5*x[t]-2*y[t]},{x[0]==8,y[0]==-7},{x[t],y[t]},t,IncludeSin
```

$$\begin{aligned}x(t) &\rightarrow \frac{2}{7}e^{-4t}(13e^{7t} + 15) \\y(t) &\rightarrow \frac{1}{7}e^{-4t}(26e^{7t} - 75)\end{aligned}$$

## 27.5 problem 38.5

Internal problem ID [13705]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.5.

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 5x(t) + 4y(t)$$

$$y'(t) = 8x(t) + y(t)$$

With initial conditions

$$[x(0) = 0, y(0) = 9]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 34

```
dsolve([diff(x(t),t) = 5*x(t)+4*y(t), diff(y(t),t) = 8*x(t)+y(t), x(0) = 0, y(0) = 9],[x(t),
```

$$x(t) = 3e^{9t} - 3e^{-3t}$$

$$y(t) = 3e^{9t} + 6e^{-3t}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 36

```
DSolve[{x'[t]==5*x[t]+4*y[t],y'[t]==8*x[t]+y[t]},{x[0]==0,y[0]==9},{x[t],y[t]},t,IncludeSing
```

$$x(t) \rightarrow 3e^{-3t}(e^{12t} - 1)$$

$$y(t) \rightarrow 3e^{-3t}(e^{12t} + 2)$$

## 27.6 problem 38.6

Internal problem ID [13706]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.6.

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 4x(t) + 2y(t)$$

$$y'(t) = 3x(t) - y(t)$$

With initial conditions

$$[x(0) = 0, y(0) = -21]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 34

```
dsolve([diff(x(t),t) = 4*x(t)+2*y(t), diff(y(t),t) = 3*x(t)-y(t), x(0) = 0, y(0) = -21],[x(t)
```

$$x(t) = -6e^{5t} + 6e^{-2t}$$

$$y(t) = -3e^{5t} - 18e^{-2t}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 36

```
DSolve[{x'[t]==4*x[t]+2*y[t],y'[t]==3*x[t]-y[t]},{x[0]==0,y[0]==-21},{x[t],y[t]},t,IncludeSi
```

$$x(t) \rightarrow -6e^{-2t}(e^{7t} - 1)$$

$$y(t) \rightarrow -3e^{-2t}(e^{7t} + 6)$$

## 27.7 problem 38.10 (a)

Internal problem ID [13707]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.10 (a).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= x(t) + 2y(t) \\y'(t) &= 5x(t) - 2y(t)\end{aligned}$$

With initial conditions

$$[x(0) = 1, y(0) = 15]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 34

```
dsolve([diff(x(t),t) = x(t)+2*y(t), diff(y(t),t) = 5*x(t)-2*y(t), x(0) = 1, y(0) = 15],[x(t)
```

$$\begin{aligned}x(t) &= -4e^{-4t} + 5e^{3t} \\y(t) &= 10e^{-4t} + 5e^{3t}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 37

```
DSolve[{x'[t]==x[t]+2*y[t],y'[t]==5*x[t]-2*y[t]},{x[0]==1,y[0]==15},{x[t],y[t]},t,IncludeSin
```

$$\begin{aligned}x(t) &\rightarrow e^{-4t}(5e^{7t} - 4) \\y(t) &\rightarrow 5e^{-4t}(e^{7t} + 2)\end{aligned}$$

## 27.8 problem 38.10 (b)

Internal problem ID [13708]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.10 (b).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 2y(t)$$

$$y'(t) = 2x(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve([diff(x(t),t)=2*y(t),diff(y(t),t)=2*x(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = c_1 e^{2t} - c_2 e^{-2t}$$

$$y(t) = c_1 e^{2t} + c_2 e^{-2t}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 68

```
DSolve[{x'[t]==2*y[t],y'[t]==2*x[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{1}{2} e^{-2t} (c_1 (e^{4t} + 1) + c_2 (e^{4t} - 1))$$

$$y(t) \rightarrow \frac{1}{2} e^{-2t} (c_1 (e^{4t} - 1) + c_2 (e^{4t} + 1))$$



## 27.9 problem 38.10 (c)

Internal problem ID [13709]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.10 (c).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= 2y(t) \\ y'(t) &= -2x(t)\end{aligned}$$

### ✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve([diff(x(t),t)=2*y(t),diff(y(t),t)=-2*x(t)],[x(t), y(t)], singsol=all)
```

$$\begin{aligned}x(t) &= -c_1 \cos(2t) + c_2 \sin(2t) \\ y(t) &= c_1 \sin(2t) + c_2 \cos(2t)\end{aligned}$$

### ✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 39

```
DSolve[{x'[t]==2*y[t],y'[t]==-2*x[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$\begin{aligned}x(t) &\rightarrow c_1 \cos(2t) + c_2 \sin(2t) \\ y(t) &\rightarrow c_2 \cos(2t) - c_1 \sin(2t)\end{aligned}$$

## 27.10 problem 38.10 (d)

Internal problem ID [13710]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.10 (d).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = -2y(t)$$

$$y'(t) = 8x(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 36

```
dsolve([diff(x(t),t)=-2*y(t),diff(y(t),t)=8*x(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = \frac{c_1 \cos(4t)}{2} - \frac{c_2 \sin(4t)}{2}$$

$$y(t) = c_1 \sin(4t) + c_2 \cos(4t)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 42

```
DSolve[{x'[t]==-2*y[t],y'[t]==8*x[t]},{x[t],y[t]},t,IncludeSingularSolutions->True]
```

$$x(t) \rightarrow c_1 \cos(4t) - \frac{1}{2}c_2 \sin(4t)$$

$$y(t) \rightarrow c_2 \cos(4t) + 2c_1 \sin(4t)$$

## 27.11 problem 38.10 (e)

Internal problem ID [13711]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.10 (e).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 4x(t) - 13y(t)$$

$$y'(t) = x(t)$$

With initial conditions

$$[x(0) = 2, y(0) = 1]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 35

```
dsolve([diff(x(t),t) = 4*x(t)-13*y(t), diff(y(t),t) = x(t), x(0) = 2, y(0) = 1],[x(t), y(t)])
```

$$x(t) = e^{2t}(-3 \sin(3t) + 2 \cos(3t))$$

$$y(t) = e^{2t} \cos(3t)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 37

```
DSolve[{x'[t]==4*x[t]-13*y[t],y'[t]==x[t]},{x[0]==2,y[0]==1},{x[t],y[t]},t,IncludeSingularSolutions->True]
```

$$x(t) \rightarrow e^{2t}(2 \cos(3t) - 3 \sin(3t))$$

$$y(t) \rightarrow e^{2t} \cos(3t)$$

## 27.12 problem 38.10 (f)

Internal problem ID [13712]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.10 (f).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= 3x(t) + 2y(t) \\y'(t) &= -2x(t) + 3y(t)\end{aligned}$$

With initial conditions

$$[x(0) = a_1, y(0) = a_2]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 48

```
dsolve([diff(x(t),t) = 3*x(t)+2*y(t), diff(y(t),t) = -2*x(t)+3*y(t), x(0) = a_1, y(0) = a_2])
```

$$\begin{aligned}x(t) &= -e^{3t}(-a_1 \cos(2t) - a_2 \sin(2t)) \\y(t) &= e^{3t}(-a_1 \sin(2t) + a_2 \cos(2t))\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 47

```
DSolve[{x'[t]==3*x[t]+2*y[t],y'[t]==-2*x[t]+3*y[t]},{x[0]==a1,y[0]==a2},{x[t],y[t]},t,IncludeSolutions->True]
```

$$\begin{aligned}x(t) &\rightarrow e^{3t}(a_1 \cos(2t) + a_2 \sin(2t)) \\y(t) &\rightarrow e^{3t}(a_2 \cos(2t) - a_1 \sin(2t))\end{aligned}$$

## 27.13 problem 38.10 (g)

Internal problem ID [13713]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.10 (g).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= 8x(t) + 2y(t) - 17 \\y'(t) &= 4x(t) + y(t) - 13\end{aligned}$$

With initial conditions

$$[x(0) = 0, y(0) = 0]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 28

```
dsolve([diff(x(t),t) = 8*x(t)+2*y(t)-17, diff(y(t),t) = 4*x(t)+y(t)-13, x(0) = 0, y(0) = 0],
```

$$\begin{aligned}x(t) &= -2e^{9t} + 2 + t \\y(t) &= -e^{9t} - 4t + 1\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 30

```
DSolve[{x'[t]==8*x[t]+2*y[t]-17,y'[t]==4*x[t]+y[t]-13},{x[0]==0,y[0]==0},{x[t],y[t]},t,Inclu
```

$$\begin{aligned}x(t) &\rightarrow t - 2e^{9t} + 2 \\y(t) &\rightarrow -4t - e^{9t} + 1\end{aligned}$$

## 27.14 problem 38.10 (h)

Internal problem ID [13714]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.10 (h).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= 8x(t) + 2y(t) + 7e^{2t} \\y'(t) &= 4x(t) + y(t) - 7e^{2t}\end{aligned}$$

With initial conditions

$$[x(0) = -1, y(0) = 1]$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 24

```
dsolve([diff(x(t),t) = 8*x(t)+2*y(t)+7*exp(2*t), diff(y(t),t) = 4*x(t)+y(t)-7*exp(2*t), x(0)
```

$$\begin{aligned}x(t) &= -\frac{3}{2} + \frac{e^{2t}}{2} \\y(t) &= 6 - 5e^{2t}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 28

```
DSolve[{x'[t]==8*x[t]+2*y[t]+7*Exp[2*t], y'[t]==4*x[t]+y[t]-7*Exp[2*t]}, {x[0]==-1, y[0]==1}, {x
```

$$\begin{aligned}x(t) &\rightarrow \frac{1}{2}(e^{2t} - 3) \\y(t) &\rightarrow 6 - 5e^{2t}\end{aligned}$$

## 27.15 problem 38.10 (i)

Internal problem ID [13715]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.10 (i).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= 4x(t) + 3y(t) - 6e^{3t} \\y'(t) &= x(t) + 6y(t) + 2e^{3t}\end{aligned}$$

With initial conditions

$$[x(0) = 4, y(0) = 0]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 44

```
dsolve([diff(x(t),t) = 4*x(t)+3*y(t)-6*exp(3*t), diff(y(t),t) = x(t)+6*y(t)+2*exp(3*t), x(0)
```

$$\begin{aligned}x(t) &= 3e^{3t} + e^{7t} - 6te^{3t} \\y(t) &= -e^{3t} + e^{7t} + 2te^{3t}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 50

```
DSolve[{x'[t]==4*x[t]+3*y[t]+6*Exp[3*t], y'[t]==x[t]+6*y[t]+2*Exp[3*t]}, {x[0]==4, y[0]==0}, {x
```

$$\begin{aligned}x(t) &\rightarrow \frac{1}{4}e^{3t}(12t + 7e^{4t} + 9) \\y(t) &\rightarrow \frac{1}{4}e^{3t}(-4t + 7e^{4t} - 7)\end{aligned}$$

## 27.16 problem 38.10 (j)

Internal problem ID [13716]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.10 (j).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= -y(t) \\ y'(t) &= 4x(t) + 24t\end{aligned}$$

With initial conditions

$$[x(0) = 0, y(0) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 26

```
dsolve([diff(x(t),t) = -y(t), diff(y(t),t) = 4*x(t)+24*t, x(0) = 0, y(0) = 0], [x(t), y(t)],
```

$$\begin{aligned}x(t) &= 3 \sin(2t) - 6t \\ y(t) &= 6 - 6 \cos(2t)\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 24

```
DSolve[{x'[t]==-y[t],y'[t]==4*x[t]+24*t},{x[0]==0,y[0]==0},{x[t],y[t]},t,IncludeSingularSolu
```

$$\begin{aligned}x(t) &\rightarrow 3 \sin(2t) - 6t \\ y(t) &\rightarrow 12 \sin^2(t)\end{aligned}$$



## 27.17 problem 38.10 (k)

Internal problem ID [13717]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.10 (k).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 4x(t) - 13y(t)$$

$$y'(t) = x(t) + 152 \cos(t)^4 - 152 \cos(t)^2 + 19 - 104 \sin(t) \cos(t)^3 + 52 \sin(t) \cos(t)$$

With initial conditions

$$[x(0) = 13, y(0) = 0]$$

✓ Solution by Maple

Time used: 0.219 (sec). Leaf size: 27

```
dsolve([diff(x(t),t) = 4*x(t)-13*y(t), diff(y(t),t) = x(t)+19*cos(4*t)-13*sin(4*t), x(0) = 13, y(0) = 0])
```

$$x(t) = 13 \cos(4t) + 13 \sin(4t)$$

$$y(t) = 8 \sin(4t)$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 25

```
DSolve[{x'[t]==4*x[t]-13*y[t],y'[t]==x[t]+19*Cos[4*t]-13*Sin[4*t]},{x[0]==13,y[0]==0},{x[t],y[t]}
```

$$x(t) \rightarrow 13(\sin(4t) + \cos(4t))$$

$$y(t) \rightarrow 8 \sin(4t)$$

## 27.18 problem 38.10 (L)

Internal problem ID [13718]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.10 (L).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 4x(t) + 3y(t) + 5 \operatorname{Heaviside}(t - 2)$$

$$y'(t) = x(t) + 6y(t) + 17 \operatorname{Heaviside}(t - 2)$$

With initial conditions

$$[x(0) = 0, y(0) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 67

```
dsolve([diff(x(t),t) = 4*x(t)+3*y(t)+5*Heaviside(t-2), diff(y(t),t) = x(t)+6*y(t)+17*Heaviside(t-2)], [x(t), y(t)])
```

$$x(t) = \operatorname{Heaviside}(t - 2) - 3 \operatorname{Heaviside}(t - 2) e^{3t-6} + 2 \operatorname{Heaviside}(t - 2) e^{-14+7t}$$

$$y(t) = -3 \operatorname{Heaviside}(t - 2) + \operatorname{Heaviside}(t - 2) e^{3t-6} + 2 \operatorname{Heaviside}(t - 2) e^{-14+7t}$$

✓ Solution by Mathematica

Time used: 0.11 (sec). Leaf size: 60

```
DSolve[{x'[t]==4*x[t]+3*y[t]+5*UnitStep[t-2],y'[t]==x[t]+6*y[t]+17*UnitStep[t-2]},{x[0]==0,y
```

$$x(t) \rightarrow \begin{cases} 1 + 2e^{7(t-2)} - 3e^{3t-6} & t \geq 2 \\ 0 & \text{True} \end{cases}$$

$$y(t) \rightarrow \begin{cases} -3 + 2e^{7(t-2)} + e^{3t-6} & t \geq 2 \\ 0 & \text{True} \end{cases}$$

## 27.19 problem 38.11

Internal problem ID [13719]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 38. Systems of differential equations. A starting point. Additional Exercises. page 786

**Problem number:** 38.11.

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 5x(t) + 4y(t)$$

$$y'(t) = 8x(t) + y(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve([diff(x(t),t)=5*x(t)+4*y(t),diff(y(t),t)=8*x(t)+y(t)],[x(t), y(t)], singsol=all)
```

$$x(t) = c_1 e^{9t} - \frac{c_2 e^{-3t}}{2}$$

$$y(t) = c_1 e^{9t} + c_2 e^{-3t}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 71

```
DSolve[{x'[t]==5*x[t]+4*y[t],y'[t]==8*x[t]+y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> T
```

$$x(t) \rightarrow \frac{1}{3} e^{-3t} (c_1 (2e^{12t} + 1) + c_2 (e^{12t} - 1))$$

$$y(t) \rightarrow \frac{1}{3} e^{-3t} (2c_1 (e^{12t} - 1) + c_2 (e^{12t} + 2))$$

**28 Chapter 39. Critical points, Direction fields  
and trajectories. Additional Exercises. page 815**

28.1	problem 39.1 (a)	892
28.2	problem 39.1 (b)	893
28.3	problem 39.1 (c)	895
28.4	problem 39.1 (d)	896
28.5	problem 39.2	897

## 28.1 problem 39.1 (a)

Internal problem ID [13720]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 39. Critical points, Direction fields and trajectories. Additional Exercises. page 815

**Problem number:** 39.1 (a).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= 2x(t) - 5y(t) \\y'(t) &= 3x(t) - 7y(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 86

```
dsolve([diff(x(t),t)=2*x(t)-5*y(t),diff(y(t),t)=3*x(t)-7*y(t)],[x(t), y(t)], singsol=all)
```

$$\begin{aligned}x(t) &= \frac{c_1 e^{\frac{(-5+\sqrt{21})t}{2}} \sqrt{21}}{6} - \frac{c_2 e^{-\frac{(5+\sqrt{21})t}{2}} \sqrt{21}}{6} + \frac{3c_1 e^{\frac{(-5+\sqrt{21})t}{2}}}{2} + \frac{3c_2 e^{-\frac{(5+\sqrt{21})t}{2}}}{2} \\y(t) &= c_1 e^{\frac{(-5+\sqrt{21})t}{2}} + c_2 e^{-\frac{(5+\sqrt{21})t}{2}}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 150

```
DSolve[{x'[t]==2*x[t]-5*y[t],y'[t]==3*x[t]-7*y[t]},{x[t],y[t]},t,IncludeSingularSolutions ->
```

$$\begin{aligned}x(t) &\rightarrow \frac{1}{42} e^{-\frac{1}{2}(5+\sqrt{21})t} \left( 3c_1 \left( (7+3\sqrt{21}) e^{\sqrt{21}t} + 7-3\sqrt{21} \right) - 10\sqrt{21}c_2 \left( e^{\sqrt{21}t} - 1 \right) \right) \\y(t) &\rightarrow \frac{1}{14} e^{-\frac{1}{2}(5+\sqrt{21})t} \left( 2\sqrt{21}c_1 \left( e^{\sqrt{21}t} - 1 \right) - c_2 \left( (3\sqrt{21}-7) e^{\sqrt{21}t} - 7-3\sqrt{21} \right) \right)\end{aligned}$$

## 28.2 problem 39.1 (b)

Internal problem ID [13721]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 39. Critical points, Direction fields and trajectories. Additional Exercises. page 815

**Problem number:** 39.1 (b).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = 2x(t) - 5y(t) + 4$$

$$y'(t) = 3x(t) - 7y(t) + 5$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 88

```
dsolve([diff(x(t),t)=2*x(t)-5*y(t)+4,diff(y(t),t)=3*x(t)-7*y(t)+5],[x(t), y(t)], singsol=all
```

$$x(t) = \frac{e^{\frac{(-5+\sqrt{21})t}{2}} c_2 \sqrt{21}}{6} - \frac{e^{-\frac{(5+\sqrt{21})t}{2}} c_1 \sqrt{21}}{6} + \frac{3e^{\frac{(-5+\sqrt{21})t}{2}} c_2}{2} + \frac{3e^{-\frac{(5+\sqrt{21})t}{2}} c_1}{2} + 3$$

$$y(t) = e^{\frac{(-5+\sqrt{21})t}{2}} c_2 + e^{-\frac{(5+\sqrt{21})t}{2}} c_1 + 2$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 185

```
DSolve[{x'[t]==2*x[t]-5*y[t]+4,y'[t]==3*x[t]-7*y[t]+5},{x[t],y[t]},t,IncludeSingularSolution
```

$$x(t) \rightarrow \frac{1}{42} e^{-\frac{1}{2}(5+\sqrt{21})t} \left( 126 e^{\frac{1}{2}(5+\sqrt{21})t} + \left( 3(7+3\sqrt{21})c_1 - 10\sqrt{21}c_2 \right) e^{\sqrt{21}t} \right. \\ \left. + (21 - 9\sqrt{21})c_1 + 10\sqrt{21}c_2 \right)$$
$$y(t) \rightarrow \frac{1}{14} e^{-\frac{1}{2}(5+\sqrt{21})t} \left( 28 e^{\frac{1}{2}(5+\sqrt{21})t} + \left( 2\sqrt{21}c_1 + (7 - 3\sqrt{21})c_2 \right) e^{\sqrt{21}t} - 2\sqrt{21}c_1 \right. \\ \left. + (7 + 3\sqrt{21})c_2 \right)$$



## 28.3 problem 39.1 (c)

Internal problem ID [13722]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 39. Critical points, Direction fields and trajectories. Additional Exercises. page 815

**Problem number:** 39.1 (c).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= 3x(t) + y(t) \\y'(t) &= 6x(t) + 2y(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 27

```
dsolve([diff(x(t),t)=3*x(t)+y(t),diff(y(t),t)=6*x(t)+2*y(t)],[x(t), y(t)], singsol=all)
```

$$\begin{aligned}x(t) &= \frac{c_2 e^{5t}}{2} - \frac{c_1}{3} \\y(t) &= c_1 + c_2 e^{5t}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 63

```
DSolve[{x'[t]==3*x[t]+y[t],y'[t]==6*x[t]+2*y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> T
```

$$\begin{aligned}x(t) &\rightarrow \frac{1}{5}(c_1(3e^{5t} + 2) + c_2(e^{5t} - 1)) \\y(t) &\rightarrow \frac{1}{5}(6c_1(e^{5t} - 1) + c_2(2e^{5t} + 3))\end{aligned}$$

## 28.4 problem 39.1 (d)

Internal problem ID [13723]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 39. Critical points, Direction fields and trajectories. Additional Exercises. page 815

**Problem number:** 39.1 (d).

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$x'(t) = x(t)y(t) - 6y(t)$$

$$y'(t) = x(t) - y(t) - 5$$

**X** Solution by Maple

```
dsolve([diff(x(t),t)=x(t)*y(t)-6*y(t),diff(y(t),t)=x(t)-y(t)-5],[x(t), y(t)], singsol=all)
```

No solution found

**X** Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{x'[t]==x[t]*y[t]-6*y[t],y'[t]==x[t]-y[t]-5},{x[t],y[t]},t,IncludeSingularSolutions -
```

Not solved

## 28.5 problem 39.2

Internal problem ID [13724]

**Book:** Ordinary Differential Equations. An introduction to the fundamentals. Kenneth B. Howell. second edition. CRC Press. FL, USA. 2020

**Section:** Chapter 39. Critical points, Direction fields and trajectories. Additional Exercises. page 815

**Problem number:** 39.2.

**ODE order:** 1.

**ODE degree:** 1.

Solve

$$\begin{aligned}x'(t) &= -x(t) + 2y(t) \\y'(t) &= 2x(t) - y(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 31

```
dsolve([diff(x(t),t)=-x(t)+2*y(t),diff(y(t),t)=2*x(t)-y(t)],[x(t), y(t)], singsol=all)
```

$$\begin{aligned}x(t) &= c_1 e^t - c_2 e^{-3t} \\y(t) &= c_1 e^t + c_2 e^{-3t}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 68

```
DSolve[{x'[t]==-x[t]+2*y[t],y'[t]==2*x[t]-y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> Tr
```

$$\begin{aligned}x(t) &\rightarrow \frac{1}{2}e^{-3t}(c_1(e^{4t} + 1) + c_2(e^{4t} - 1)) \\y(t) &\rightarrow \frac{1}{2}e^{-3t}(c_1(e^{4t} - 1) + c_2(e^{4t} + 1))\end{aligned}$$