


A Solution Manual For

**Advanced Mathematica, Book2, Perkin
and Perkin, 1992**



Nasser M. Abbasi

May 16, 2024

Contents

1 Chapter 11.3, page 316

2

1 Chapter 11.3, page 316

1.1	problem 1	3
1.2	problem 2	4
1.3	problem 3	5
1.4	problem 4	6
1.5	problem 5	7
1.6	problem 6	8
1.7	problem 7	9
1.8	problem 8	10
1.9	problem 9	11
1.10	problem 10	12
1.11	problem 11	13
1.12	problem 12	14
1.13	problem 13	15
1.14	problem 14	16
1.15	problem 15	17
1.16	problem 16	18
1.17	problem 17	19
1.18	problem 18	20
1.19	problem 19	21
1.20	problem 20	22
1.21	problem 21	23
1.22	problem 22	24
1.23	problem 23	25
1.24	problem 24	26
1.25	problem 25	27
1.26	problem 26	28
1.27	problem 27	29
1.28	problem 28	30

1.1 problem 1

Internal problem ID [3052]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$3y^2y' = 2x - 1$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 60

```
dsolve(3*y(x)^2*diff(y(x),x)=2*x-1,y(x), singsol=all)
```

$$y(x) = (x^2 + c_1 - x)^{\frac{1}{3}}$$
$$y(x) = -\frac{(x^2 + c_1 - x)^{\frac{1}{3}} (1 + i\sqrt{3})}{2}$$
$$y(x) = \frac{(x^2 + c_1 - x)^{\frac{1}{3}} (i\sqrt{3} - 1)}{2}$$

✓ Solution by Mathematica

Time used: 0.257 (sec). Leaf size: 71

```
DSolve[3*y[x]^2*y'[x]==2*x-1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt[3]{x^2 - x + 3c_1}$$
$$y(x) \rightarrow -\sqrt[3]{-1} \sqrt[3]{x^2 - x + 3c_1}$$
$$y(x) \rightarrow (-1)^{2/3} \sqrt[3]{x^2 - x + 3c_1}$$

1.2 problem 2

Internal problem ID [3053]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - 6y^2x = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x)=6*x*y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{1}{-3x^2 + c_1}$$

✓ Solution by Mathematica

Time used: 0.12 (sec). Leaf size: 22

```
DSolve[y'[x]==6*x*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{3x^2 + c_1}$$
$$y(x) \rightarrow 0$$

1.3 problem 3

Internal problem ID [3054]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$y' - e^y \sin(x) = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
dsolve(diff(y(x),x)=exp(y(x))*sin(x),y(x), singsol=all)
```

$$y(x) = -\ln(\cos(x) - c_1)$$

✓ Solution by Mathematica

Time used: 0.332 (sec). Leaf size: 15

```
DSolve[y'[x]==Exp[y[x]]*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\log(\cos(x) - c_1)$$

1.4 problem 4

Internal problem ID [3055]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$y' - e^{-y+x} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)=exp(x-y(x)),y(x), singsol=all)
```

$$y(x) = \ln(e^x + c_1)$$

✓ Solution by Mathematica

Time used: 0.769 (sec). Leaf size: 12

```
DSolve[y'[x]==Exp[x-y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(e^x + c_1)$$

1.5 problem 5

Internal problem ID [3056]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$y' - x \sec(y) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=x*sec(y(x)),y(x), singsol=all)
```

$$y(x) = \arcsin\left(\frac{x^2}{2} + c_1\right)$$

✓ Solution by Mathematica

Time used: 0.436 (sec). Leaf size: 31

```
DSolve[y'[x]==x*Sec[y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arcsin\left(\frac{x^2}{2} + c_1\right)$$

$$y(x) \rightarrow \arcsin\left(\frac{x^2}{2} + c_1\right)$$

1.6 problem 6

Internal problem ID [3057]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - 3 \cos(y)^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=3*cos(y(x))^2,y(x), singsol=all)
```

$$y(x) = \arctan(3x + 3c_1)$$

✓ Solution by Mathematica

Time used: 0.387 (sec). Leaf size: 32

```
DSolve[y'[x]==3*Cos[y[x]]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arctan(3x + 2c_1)$$

$$y(x) \rightarrow -\frac{\pi}{2}$$

$$y(x) \rightarrow \frac{\pi}{2}$$

1.7 problem 7

Internal problem ID [3058]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$xy' - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 7

```
dsolve(x*diff(y(x),x)=y(x),y(x), singsol=all)
```

$$y(x) = c_1x$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 14

```
DSolve[x*y'[x]==y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x$$

$$y(x) \rightarrow 0$$

1.8 problem 8

Internal problem ID [3059]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$(1 - x)y' - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve((1-x)*diff(y(x),x)=y(x),y(x), singsol=all)
```

$$y(x) = \frac{c_1}{x - 1}$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 20

```
DSolve[(1-x)*y'[x]==y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{1 - x}$$
$$y(x) \rightarrow 0$$

1.9 problem 9

Internal problem ID [3060]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{4xy}{x^2 + 1} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x)=(4*x*y(x))/(x^2+1),y(x), singsol=all)
```

$$y(x) = c_1(x^2 + 1)^2$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 20

```
DSolve[y'[x]==(4*x*y[x])/(x^2+1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1(x^2 + 1)^2$$

$$y(x) \rightarrow 0$$

1.10 problem 10

Internal problem ID [3061]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{2y}{x^2 - 1} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)=(2*y(x))/(x^2-1),y(x), singsol=all)
```

$$y(x) = -\frac{(x-1)c_1}{x+1}$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 22

```
DSolve[y'[x]==(2*y[x])/(x^2-1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{c_1(x-1)}{x+1}$$
$$y(x) \rightarrow 0$$

1.11 problem 11

Internal problem ID [3062]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$y'x^2 - y^2 = 0$$

With initial conditions

$$[y(1) = -1]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 14

```
dsolve([x^2*diff(y(x),x)-y(x)^2=0,y(1) = -1],y(x), singsol=all)
```

$$y(x) = -\frac{x}{2x - 1}$$

✓ Solution by Mathematica

Time used: 0.118 (sec). Leaf size: 14

```
DSolve[{x^2*y'[x]-y[x]^2==0,y[1]==-1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x}{1 - 2x}$$

1.12 problem 12

Internal problem ID [3063]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$y' + 2yx = 0$$

With initial conditions

$$[y(0) = 5]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 12

```
dsolve([diff(y(x),x)+2*x*y(x)=0,y(0) = 5],y(x), singsol=all)
```

$$y(x) = 5e^{-x^2}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 14

```
DSolve[{y'[x]+2*x*y[x]==0,y[0]==5},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 5e^{-x^2}$$

1.13 problem 13

Internal problem ID [3064]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$\cot(x) y' - y = 0$$

With initial conditions

$$[y(0) = 2]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve([cot(x)*diff(y(x),x)=y(x),y(0) = 2],y(x), singsol=all)
```

$$y(x) = 2 \sec(x)$$

✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 9

```
DSolve[{Cot[x]*y'[x]==y[x],y[0]==2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 \sec(x)$$

1.14 problem 14

Internal problem ID [3065]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 14.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' - x e^{-2y} = 0$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 12

```
dsolve([diff(y(x),x)=x*exp(-2*y(x)),y(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{\ln(x^2 + 1)}{2}$$

✓ Solution by Mathematica

Time used: 0.333 (sec). Leaf size: 15

```
DSolve[{y'[x]==x*Exp[-2*y[x]],y[0]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \log(x^2 + 1)$$

1.15 problem 15

Internal problem ID [3066]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$y' - 2yx = 2x$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve([diff(y(x),x)-2*x*y(x)=2*x,y(0) = 1],y(x), singsol=all)
```

$$y(x) = 2e^{x^2} - 1$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 14

```
DSolve[{y'[x]-2*x*y[x]==2*x,y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2e^{x^2} - 1$$

1.16 problem 16

Internal problem ID [3067]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$xy' - yx - y = 0$$

With initial conditions

$$[y(1) = 1]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve([x*diff(y(x),x)=x*y(x)+y(x),y(1) = 1],y(x), singsol=all)
```

$$y(x) = x e^{x-1}$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 12

```
DSolve[{x*y'[x]==x*y[x]+y[x],y[1]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x-1}x$$

1.17 problem 17

Internal problem ID [3068]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 17.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$(x^3 + 1) y' = 3 \tan(x) x^2$$

With initial conditions

$$\left[y(0) = \frac{\pi}{2} \right]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 36

```
dsolve([(1+x^3)*diff(y(x),x)=3*x^2*tan(x),y(0) = 1/2*Pi],y(x), singsol=all)
```

$$y(x) = 3 \left(\int_0^x \frac{\tan(z) z^2}{(z+1)(z^2 - z + 1)} dz \right) + \frac{\pi}{2}$$

✓ Solution by Mathematica

Time used: 8.597 (sec). Leaf size: 35

```
DSolve[{(1+x^3)*y'[x]==3*x^2*Tan[x],y[0]==Pi/2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \int_0^x \frac{3K[1]^2 \tan(K[1])}{K[1]^3 + 1} dK[1] + \frac{\pi}{2}$$

1.18 problem 18

Internal problem ID [3069]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 18.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$x \cos(y) y' - \sin(y) = 1$$

With initial conditions

$$[y(1) = 0]$$

✓ Solution by Maple

Time used: 0.188 (sec). Leaf size: 8

```
dsolve([x*cos(y(x))*diff(y(x),x)=1+sin(y(x)),y(1) = 0],y(x), singsol=all)
```

$$y(x) = \arcsin(x - 1)$$

✓ Solution by Mathematica

Time used: 37.067 (sec). Leaf size: 53

```
DSolve[{x*Cos[y[x]]*y'[x]==1+Sin[y[x]],y[1]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2 \arccos\left(\frac{1}{2}(\sqrt{2-x} + \sqrt{x})\right)$$

$$y(x) \rightarrow 2 \arccos\left(\frac{1}{2}(\sqrt{2-x} + \sqrt{x})\right)$$

1.19 problem 19

Internal problem ID [3070]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$xy' - 2y(y - 1) = 0$$

With initial conditions

$$\left[y\left(\frac{1}{2}\right) = 2 \right]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 15

```
dsolve([x*diff(y(x),x)=2*y(x)*(y(x)-1),y(1/2) = 2],y(x), singsol=all)
```

$$y(x) = -\frac{1}{2x^2 - 1}$$

✓ Solution by Mathematica

Time used: 0.339 (sec). Leaf size: 14

```
DSolve[{x*y'[x]==2*y[x]*(y[x]-1),y[1/2]==2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{1 - 2x^2}$$

1.20 problem 20

Internal problem ID [3071]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 20.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$2xy' + y^2 = 1$$

With initial conditions

$$[y(1) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 13

```
dsolve([2*x*diff(y(x),x)=1-y(x)^2,y(1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{x - 1}{x + 1}$$

✓ Solution by Mathematica

Time used: 0.514 (sec). Leaf size: 14

```
DSolve[{2*x*y'[x]==1-y[x]^2,y[1]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x - 1}{x + 1}$$

1.21 problem 21

Internal problem ID [3072]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 21.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$(1 - x)y' - yx = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve((1-x)*diff(y(x),x)=x*y(x),y(x), singsol=all)
```

$$y(x) = \frac{c_1 e^{-x}}{x - 1}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 23

```
DSolve[(1-x)*y'[x]==x*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1 e^{-x}}{x - 1}$$
$$y(x) \rightarrow 0$$

1.22 problem 22

Internal problem ID [3073]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 22.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(x^2 - 1) y' - (x^2 + 1) y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve((x^2-1)*diff(y(x),x)=(x^2+1)*y(x),y(x), singsol=all)
```

$$y(x) = \frac{e^x(x-1)c_1}{x+1}$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 25

```
DSolve[(x^2-1)*y'[x]==(x^2+1)*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{c_1 e^x (x-1)}{x+1}$$
$$y(x) \rightarrow 0$$

1.23 problem 23

Internal problem ID [3074]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 23.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - e^x(y^2 + 1) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)=exp(x)*(y(x)^2+1),y(x), singsol=all)
```

$$y(x) = \tan(e^x + c_1)$$

✓ Solution by Mathematica

Time used: 0.274 (sec). Leaf size: 26

```
DSolve[y'[x]==Exp[x]*(y[x]^2+1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan(e^x + c_1)$$

$$y(x) \rightarrow -i$$

$$y(x) \rightarrow i$$

1.24 problem 24

Internal problem ID [3075]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 24.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y'e^y - 2xe^y = -2x$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 19

```
dsolve(exp(y(x))*diff(y(x),x)+2*x=2*x*exp(y(x)),y(x), singsol=all)
```

$$y(x) = -\ln\left(-\frac{1}{e^{x^2}c_1 - 1}\right)$$

✓ Solution by Mathematica

Time used: 2.015 (sec). Leaf size: 21

```
DSolve[Exp[y[x]]*y'[x]+2*x==2*x*Exp[y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log\left(1 + e^{x^2+c_1}\right)$$

$$y(x) \rightarrow 0$$

1.25 problem 25

Internal problem ID [3076]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 25.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$e^{2x}yy' = -2x$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.109 (sec). Leaf size: 16

```
dsolve([exp(2*x)*y(x)*diff(y(x),x)+2*x=0,y(0) = 1],y(x), singsol=all)
```

$$y(x) = \sqrt{(2x + 1)e^{-2x}}$$

✓ Solution by Mathematica

Time used: 1.816 (sec). Leaf size: 20

```
DSolve[{Exp[2*x]*y[x]*y'[x]+2*x==0,y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt{e^{-2x}(2x + 1)}$$

1.26 problem 26

Internal problem ID [3077]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 26.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$xyy' - \sqrt{y^2 - 9} = 0$$

With initial conditions

$$[y(e^4) = 5]$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 12

```
dsolve([x*y(x)*diff(y(x),x)=sqrt(y(x)^2-9),y(exp(4)) = 5],y(x), singsol=all)
```

$$y(x) = \sqrt{9 + \ln(x)^2}$$

✓ Solution by Mathematica

Time used: 0.256 (sec). Leaf size: 33

```
DSolve[{x*y[x]*y'[x]==Sqrt[y[x]^2-9],y[Exp[4]]==5},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt{\log^2(x) + 9}$$

$$y(x) \rightarrow \sqrt{\log^2(x) - 16 \log(x) + 73}$$

1.27 problem 27

Internal problem ID [3078]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 27.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _exact, _rational, [_Abel, '2nd ty`

$$(y + x - 1)y' + y = x + 1$$

✓ Solution by Maple

Time used: 0.391 (sec). Leaf size: 28

```
dsolve((x+y(x)-1)*diff(y(x),x)=(x-y(x)+1),y(x), singsol=all)
```

$$y(x) = \frac{-c_1 x - \sqrt{2c_1^2 x^2 + 1} + c_1}{c_1}$$

✓ Solution by Mathematica

Time used: 0.163 (sec). Leaf size: 47

```
DSolve[(x+y[x]-1)*y'[x]==(x-y[x]+1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{2x^2 + 1 + c_1} - x + 1$$

$$y(x) \rightarrow \sqrt{2x^2 + 1 + c_1} - x + 1$$

1.28 problem 28

Internal problem ID [3079]

Book: Advanced Mathematica, Book2, Perkin and Perkin, 1992

Section: Chapter 11.3, page 316

Problem number: 28.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$xyy' + y^2 = 2x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(x*y(x)*diff(y(x),x)=2*x^2-y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{x^4 + c_1}}{x}$$
$$y(x) = -\frac{\sqrt{x^4 + c_1}}{x}$$

✓ Solution by Mathematica

Time used: 0.206 (sec). Leaf size: 38

```
DSolve[x*y[x]*y'[x]==2*x^2-y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{x^4 + c_1}}{x}$$
$$y(x) \rightarrow \frac{\sqrt{x^4 + c_1}}{x}$$